MBL/WHOI
LIBRARY

*Gift
of
Arthur L.
Humes*

MBL/WHOI

0 0301 0065372 1

T. S. Stephens
Arthur G. Himes

Binding .55

SMITHSONIAN INSTITUTION.
UNITED STATES NATIONAL MUSEUM.

42
430.7
U6
556

SYNOPSIS OF THE NAIADES, OR PEARLY
FRESH-WATER MUSSELS.

BY

CHARLES TORREY SIMPSON

Aid, Division of Mollusks.

From the Proceedings of the United States National Museum, Vol. XXII, pages 501-1044
(with Plate XVIII).

[No. 1205.]

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1900.

TO THE MEMORY OF

DOCTOR ISAAC LEA,

WHO, GIVING THE BEST YEARS OF HIS LIFE TO THE STUDY OF THE
NAIADES, CLEARLY RECORDED AND SUMPTUOUSLY ILLUSTRATED
HIS RESEARCHES UPON THEM, AND GAVE TO THE UNITED
STATES NATIONAL MUSEUM THE UNRIVALED COL-
LECTION THE STUDY OF WHICH HAS MADE
POSSIBLE THE PREPARATION OF THE
PRESENT WORK; AND TO

MRS. FRANCES LEA CHAMBERLAIN

AND THE

REVEREND DOCTOR LEANDER T. CHAMBERLAIN,

THROUGH WHOSE GENEROSITY THE PREEMINENCE OF THE COLLEC-
TION HAS BEEN MAINTAINED AND THE LITERATURE
GATHERED FOR MONOGRAPHIC STUDY,

THIS WORK IS RESPECTFULLY DEDICATED.

TABLE OF CONTENTS.

	Page.
Introduction	501
Sketch of different classifications	501
Anatomy of the Naiades	503
Shell characters	504
Geographical provinces	504
Map of the Naiad provinces	505
Number of genera and species	505
Beak sculpture	507
Past history of the Naiades	507
Mutelidæ	512
The work of Rafinesque	512
The work of Lamarek	513
The so-called "New School"	513
Synopsis of the genera of Naiades	514
Synopsis of the Naiades	516
Family Unionidæ	516
Subfamily Unioninæ	516
Heterogenæ	516
Genus Truncilla	516
Micromya	524
Lampsilis	526
Pseudospatha	577
Hyriopsis	578
Chamberlainia	582
Cristaria	583
Lepidodesma	586
Pilsbryoconcha	587
Medionidus	588
Nephronaias	591
Glebula	597
Obovaria	599
Plagiola	603
Digeneæ	607
Tritogonia	607
Mesogenæ	609
Cyprogenia	609
Obliquaria	610
Ptychogenæ	612
Ptychobranthus	612
Eschatigenæ	614

Synopsis of the Naiades—Continued.

Page.

Family Unionidæ—Continued.

Subfamily Unioninæ—Continued.

Eschatigenæ—Continued.

Genus *Dromus*..... 614*Diagenæ*..... 615*Strophitus*..... 615*Homogenæ*..... 620*Anodonta*..... 620*Colletopterum*..... 649*Gabillotia*..... 649*Leguminaia*..... 651*Lastena*..... 654*Solenaia*..... 655*Gonidea*..... 657*Anodontoides*..... 658*Pegias*..... 660*Arcidens*..... 661*Symphynota*..... 662*Alasmidonta*..... 666*Hemilastena*..... 673*Margaritana*..... 674*Unio*..... 679*Pleurobema*..... 745*Tetragenæ*..... 765*Quadrula*..... 765*Schistodesmus*..... 803*Gibbosula*..... 804*Cuneopsis*..... 804Subfamily *Hyrianæ*..... 806*Rosanorhamphus*..... 806Genus *Nodularia*..... 806*Harmandia*..... 827*Grandidieria*..... 827*Physunio*..... 830*Dalliella*..... 832*Pseudodon*..... 834*Parreysia*..... 840*Ptychorhynchus*..... 850*Virgus*..... 851*Ctenodesma*..... 852*Rectidens*..... 853*Lamellidens*..... 854*Trapezoidens*..... 858*Arconaia*..... 859*Pseudavicula*..... 860*Arcidopsis*..... 861*Lamphorhamphus*..... 862*Tetraplodon*..... 863*Castalina*..... 865*Castaliella*..... 866*Callonaia*..... 867*Hyria*..... 868*Prisodon*..... 869*Diplodon*..... 872

Synopsis of the Naiades—Continued.	Page.
Family Mutelidæ	895
Genus Spatha	895
Mutela	902
Chelidonopsis	906
Brazzæa	906
Pleiodon	907
Monocondylæa	908
Iheringella	913
Fossula	914
Leila	914
Glabaris	916
Mycetopoda	933
Bibliography	936
Index	999

ERRATA.

Page 509, line 21, *for* varicosa *read* cicatricosa.

Page 516, before Heterogenæ *insert* Exobranchiæ.

Page 548, line 36, *Unio striatus* Sowerby should be preceded by a mark of interrogation.

Page 559, line 6 from the bottom, *for* L. prevostianus *read* U. prevostianus.

Page 591, line 3, *for* Lampsilis *read* Medionidus.

Page 648, *omit* 8th and 9th lines from the bottom.

Page 719, *Unio striatulus* Lea and U. amabilis Lea should be placed in the Argenteus group of Pleurobema.

Page 760, last line, *for* bigbyana *read* bigbyensis.

Page 766, line 6 from bottom, *for* multiplicata *read* heros.

Page 802, *omit* line 8.

Page 806, line 19, *for* æquitoria *read* æquatoria.

Page 823, lines 10 and 11, *for* æquitoria *read* æquatoria.

Page 825, *omit* lines 8 and 9.

Page 851, line 27, *for* incertus *read* murinus.

Page 862, *omit* line 31.

Page 894, *omit* line 28.

Page 906, 13th and following lines, *for* Chelidoneura *read* Chelidonura.

Page 933, *omit* lines 16 and 17.

SYNOPSIS OF THE NAIADES, OR PEARLY FRESH-WATER MUSSELS.

By CHARLES TORREY SIMPSON.

Aid, Division of Mollusks.

In 1836 Dr. Isaac Lea published a Synopsis of the Family of Naiades, which was followed by a second edition in 1838, a third in 1852, and a fourth in 1870. In this work Dr. Lea did not attempt to make any systematic classification, but divided the species into two great genera, *Margarona* and *Platiris*, and subdivided these genera into subgenera, and finally into smaller groups founded on the presence or absence of a dorsal wing, and on the form and sculpture of the shell. This arrangement was regarded by Dr. Lea as merely a temporary one, to be used until a knowledge of the soft parts could be obtained, on which to base a permanent classification.

Several other attempts at a classification of the Naiades have been made, and among these is that of Swainson in 1840, who believed that the whole family formed a circular group, that by beginning at a certain point in it the species or genera would be found closely related to others, and these to others still, and that by following up the chain of relationships one would be brought back to the place of starting. My studies have not borne out Swainson's conclusions, but I have adopted two of his divisions of the Unionidæ, the subfamilies Unioninæ and Hyrianae. His subfamilies Anodontinæ and Alasmodontinæ have no value, and the subfamily Iridinæ belongs in the family Mutelidæ.

In 1847 Gray gave a classification of the Naiades in his List of Recent Genera, in which he divided them into three families: Unionidæ, with the genera *Anodonta*, *Margaritana*, *Monocondylæa*, *Unio*, *Barbala*, *Lamproscapha*, *Anodonta*, *Byssanodonta*, and *Glabaris*; Mutelidæ with *Mutela*, *Leila*, *Pleiodon*, *Paryodon* and *Prisodon*, and Mycetopodidæ, with the genus *Mycetopus*.

Troschel, in 1847, proposed a classification in which the genera were founded on anatomical characters, and though for lack of knowledge

he was led into several serious errors, his work has proved an important aid in classification.

H. and A. Adams followed with a classification in 1857, in which they recognized the families Unionidæ and Mutelidæ, dividing the former into two subfamilies, Unioninæ and Mycetopiinæ, and giving a list of the then known species.

Agassiz, in 1852, published a classification of the North American Unionidæ, based more especially on the anatomy, but much of his work was hasty and careless, and as a result his groups are, for the most part, heterogeneous assemblages of unrelated species.

In 1893, von Ihering published his *Najaden* von San Paulo, in which he showed that while certain of the Naiades began their existence on hatching from the egg as a *glochidium*, with a bivalve shell capable of containing the soft parts, others developed into a *lasidium*, with three segments, the middle one alone having a single shell. He showed that the presence or absence of siphons by which the Mutelidæ had been separated from the Unionidæ was not a constant character, even in individual species, and placed the genera which he believed to have glochidia in the family Unionidæ, and those with lasidia in the Mutelidæ. I consider this the most important discovery that has yet been made in the study of the Naiades.

In 1896 the writer published a classification of the Naiades, based largely on his studies of shell characters. In that it was shown that throughout one great group all the genera had shells with schizodont hinge teeth or vestiges of them, and in the rest, a smaller group, they all had taxodont teeth, or vestiges of them. As all the genera whose shells have schizodont teeth were believed by von Ihering to possess a *glochidium*, and those with taxodont teeth have a *lasidium*, it seemed as though there was a natural division of the *Naiades* into two families Unionidæ and Mutelidæ.

Some years ago the Rev. and Mrs. L. T. Chamberlain, son-in-law and daughter of Dr. Lea, suggested to Mr. Dall that a new edition of the Synopsis, or if necessary a new work, should be prepared, and for this purpose they generously offered to contribute such sums as might be needed for the purchase of material to add to the great collection left by Dr. Lea to the U. S. National Museum at his death, and for literature that might be required in preparing the work. As the writer had made a special study of the Naiades, the task of preparing the new Synopsis fell to him. Since then Mrs. Chamberlain has been called away by death, greatly regretted, but her husband has continued to cheerfully furnish all the needed funds for carrying on the work to completion.

While the labor of preparing the synopsis has been arduous, and while it has been impossible to obtain material needed in many important groups, or all the necessary literature, I feel that my opportunities for study have been exceptionally good. I have had constant access to the great collection of Dr. Lea, the finest, no doubt, ever

made of the Unionidæ, and containing more types than any other in existence. In addition to this the U. S. National Museum possesses a large collection, including many types, and for the most part carefully determined by that able student Dr. James Lewis, as well as the entire Jeffreys collection. A great amount of valuable material has been lately added to it, including the Morelet collection of Naiades, a set of Burmese shells from Fea, a series of New Zealand forms from Suter, a large number of species from von Ihering, from South and Central America and Mexico, the entire series of types of Mr. S. H. and B. H. Wright, and types from many other sources. The writer has personally examined the collections of Say, Conrad, and Rafinesque, in the Academy of Natural Sciences in Philadelphia, as well as the excellent alcoholic series which was the basis of all of Dr. Lea's studies. Through the courtesy of the officers of the New York State Museum of Natural History the entire collection of Naiades of Dr. Gould was loaned to the U. S. National Museum, in order that he might study it. Besides this he has examined the fine collections of Mr. Bryant Walker, Mrs. George Andrews, Prof. A. G. Wetherby, and a great amount of material belonging to Mr. William A. Marsh, Mr. B. H. Wright, and many others. He has made critical examinations of the soft parts of more than four hundred species, American and foreign.

Aside from the careful work of Lea, Troschel, and Pelseneer, little has been done in the way of studying the anatomy of the Naiades. The soft parts of a good many foreign species have been examined, and descriptions published which do not describe. Authors have gone into details of the color of the organs and of insignificant characters, but have paid no attention to really important points.

In my paper on The Classification and Geographical Distribution of the Pearly Fresh-water Mussels,¹ I placed a great variety of forms under the generic name *Unio*. Since that time additional knowledge, gained largely from a study of the soft parts, has led me to the belief that it would be best to dismember this genus somewhat as the old group *Helix* has been dismembered by Pilsbry. Ordinarily the soft parts of most of the Unionidæ show but slight differential characters, but at the time when the ovules pass down into the gills a most remarkable change in those organs generally takes place. In the *Anodonta edentula* of Say, short, horizontal ovisacs are developed, which run directly across the animal, and which at maturity break through the outer walls of the outer gills and pass with their young entire into the water. In the forms typified by *Unio anodontoides* the young are contained only in very distinct vertical or oblique ovisacs in the hinder part of the outer gills; in *Unio crassidens*, *pictorum*, and the like, the embryos fill the entire outer gills, forming thick, smooth pads; in *Unio metacervus*, *trigonus*, *multiplicatus*, and allied forms, they occupy all four of the branchiæ throughout. In *Unio phascolus* the smooth outer

¹Proc. U. S. Nat. Mus., XVIII, 1896, pp. 295-343.

gills begin to be crimped as they are being filled with embryos, until when full, they become a series of marvelous folds. In *U. irroratus* several ovisacs in the center of the outer gills grow out to a great length, become filled with young, and are closely coiled. In *U. cornutus* a few central ovisacs develop so as to project below in a long, straight flap. In all the South American and Australian Unios, so far as is known, the inner gills alone, as a rule, are filled with young, and this is probably the case with the species of the Ethiopian region and most of those of southeastern Asia. After the young have passed out into the water the gills of all the species change back into their ordinary condition, and when not gravid there is great similarity in those of most of the species formerly classed as Unios.

These peculiar evanescent characters, assumed when the gills are transformed into marsupia, seem to be quite constant, and I believe they can be used as a basis for the foundation of genera. When these are once discovered and understood it will be found, on careful examination, that there are minor shell characters that correspond with those of the marsupia, and which help us in placing certain forms of whose anatomy we know but little. Thus the principal shell characters of such species as *Unio trigonus*, *metanerrus*, and *plicatus* are much like those of *U. gibbosus*, *crassidens*, and *buckleyi*. But in the former set all four gills will be found to be filled with young in the gravid female, while in the latter only the outer ones are full. And the former have generally shorter, solidier, more inflated shells and deep beak cavities, while the latter have longer, lighter shells, which have shallow beak cavities. In *Anodonta edentula* the ovisacs are short and run across the gills; in the very similar looking *A. ferussaciana* the outer gills are filled with oblique ovisacs, and the inner have (at least in some cases) more or less ovules. The beak sculpture of these two species is very different.

From the earliest period in which the Naiades have been studied to the present time it has been claimed by some that the sexes were separate, by others that the animals were hermaphroditic. Recent careful studies by such men as Sterki, Taylor, Kelly, and others appear to demonstrate that in the more highly organized Unionidæ (those which have two forms of shells, and have the ovisacs in the hinder part of the outer gills) the sexes are always separate. In the more simply organized Unionidæ (those with but one form of shell and with the embryos occupying the entire gill) the sexes may or may not be separate.

Further study has shown me that the provinces established in the paper I have quoted hold good as there laid down. There is evidently a very close relationship between many of the unionoid forms of southeastern Asia and tropical Africa. So close is this relation that the two regions might be united if it were not for the fact that a large

MAP SHOWING THE DISTRIBUTION

PEARLY FRESH-WATER MUSSELS

number of forms are found in each of them that are not at all closely related to any of those of the other.

These regions may be tabulated as follows:

AREAS OF THE NAIAD REGIONS.¹

	{ Europe.
	{ Northern and Western Asia.
Palaearctic	{ North Africa to the Desert.
	{ Pacific drainage of North America.
Ethiopian	{ Africa, south of the Sahara.
Oriental	{ Asia, south of the Himalayas.
	{ East Indies to the Solomon Islands.
	{ Australia.
Australian	{ Tasmania.
	{ New Zealand. Part of New Guinea.
Neotropical	{ South America.
	{ Central America.
Central American	{ Mexico, east of the Cordillera.
	{ Cuba.
	{ Entire Mississippi Valley and the Gulf drainage from west Florida
	{ to the Rio Grande.
Mississippian	{ Mackenzie River system.
	{ Red River of the North.
	{ Great Lakes.
Atlantic	{ Lower St. Lawrence and rivers of eastern Canada.
	{ Atlantic drainage of the United States.

I have hesitated long before placing a considerable number of the species of Southeastern Asia in and near North American genera. I had hoped to be able to examine the soft parts of a number of these Oriental forms, but have been disappointed, and as nothing is known of the anatomy of most of them I have been compelled to classify by shell characters alone. It is certain that the beak sculpture of a large number of Naiades of this area is essentially concentric, and *not* zigzag radial, as it is in most of the forms of the Oriental region. The shell characters of the ponderous Chinese *Uniones* are certainly much like those of *Quadrula* in the form, the teeth, the beak sculpture, and especially the deep, compressed beak cavities. A great number of fossil forms from the Tertiary strata of Asia and Eastern Europe seem to show the closest relationship to the American *Quadrulas*. *Hyriopsis*, *Cristaria*, *Chamberlainia*, and *Pilsbryoconcha* seem to be related by shell characters to the alate forms of *Lampsilis*, and even the strange *Pseudospatha* of Africa would appear to belong here.

I recognize about one thousand species and 82 varieties of Unionidæ, having reduced to the synonymy a great many names that most authors have believed to stand for valid species. Of these, 533 species and 55 varieties belong in North America and 101 in South America. The list contains 117 species of Mutelidæ and 11 varieties, and of these 80

¹ For map of Naiad Regions see plate.

species are South American. The following list shows the number of species in each genus and the distribution of the genera:

UNIONIDE.

19 Truncilla, United States.
3 varieties.	6 Margaritana, Circumboreal.
2 Micromya, United States.
8 varieties.	145 Unio, North America, Palearctic.
13 varieties.
128 Lampsilis, North America.
4 Pseudospatha, Tropical Africa.	72 Pleurobema, United States.
1 variety.
10 Hyriopsis, Eastern Asia.
3 Chamberlainia, Southeastern Asia.	99 Quadrula, North America, Eastern Asia.
12 varieties.
10 Cristaria, Eastern Asia.
2 Lepidodesma, Eastern Asia.	2 Schistodesma, Eastern Asia.
4 Pilsbryoconcha, Southeastern Asia.
6 Medionidus, United States.	1 Gibbosula, Eastern Asia.
22 Nephronaias, Mexico, Central America, Cuba.	5 Cnecopsis, Eastern Asia.
1 Glebula, United States.	90 Nodularia, Eastern Asia, Asia Minor, Tropical Africa.
6 varieties.
9 Obovaria, United States.
1 variety.	1 Harmandia, Southeastern Asia.
8 Plagiola, North America.	8 Physunio, Southeastern Asia.
2 Tritogonia, United States.
1 variety.	7 Dalliella, Southeastern Asia.
2 Cyprogenia, United States.
1 variety.	11 Grandidieria, Tropical Africa.
1 Obliquaria, United States.	23 Pseudodon, Eastern Asia.
1 variety.
6 Ptychobrancheus, United States.	38 Parreysia, Southeastern Asia, Tropical Africa.
8 varieties.
8 Strophitus, North America.
1 variety.	5 Ptychorhynchus, Eastern Asia.
1 variety.
55 Anodonta, North America, Eastern Asia, Palearctic.
8 varieties.	3 Virgus, Southeastern Asia.
4 Colleopterum, Southeastern Europe.	2 Ctenodesma, Southeastern Asia.
3 Gabillotia, Asia Minor.
1 variety.	5 Rectidens, Southeastern Asia.
7 Leguminaia, Southern Europe, Asia Minor.
1 variety.	12 Lamellidens, Southeastern Asia, Tropical Africa.
5 varieties.
1 Lastena, United States.	5 Trapezoidens, Southeastern Asia.
1 variety.
9 Solenaia, Eastern Asia.	3 Arconaia, Eastern Asia.
1 Gonidea, United States.	1 Pseudavicula, Tropical Africa.
1 Anodontoides, North America.
2 varieties.	1 Arcidopsis, Southeastern Asia.
1 Pegias, United States.	9 Tetraplodon, South America.
1 variety.
1 Arcidens, United States.	5 Castalina, South America.
7 Symphynota, North America.
1 variety.	1 Callonaia, South America.
13 Alasmidonta, North America.
1 Hemilastena, United States.	4 Hyria, South America.
	8 Prisodon, South America.
	94 Diplodon, South America, Australasia, South Africa.
6 varieties.

MUTELIDE.

27 Spatha, Tropical Africa.
3 varieties.	2 Chelidonopsis, Tropical Africa.
9 Mutela, Tropical Africa.	5 Brazzaea, Tropical Africa.
	4 Pleiodon, Tropical Africa.

10 Monocondylaea, South America.

4 Iheringella, South America.

2 Fossula, South America.

3 Leila, South America.

53 Glabaris, South America.

8 varieties.

8 Mycetopoda, South America.

Total: 61 genera of Unionidæ; 11 genera of Mutelidæ.

I am inclined to believe with von Ihering that the primitive beak sculpture of the Unionidæ was radial, and in two species of Unios from what are believed to be Triassic or Permian strata of the Staked Plains of Texas,¹ which are probably the oldest forms known, the beaks clearly show *strictly radial sculpture*. Four other species from the same lot are not in condition to exhibit this character.

Now I take this to belong to the simplest, earliest, and most lowly organized form of unionoid life. I believe that the earlier Unios had the young contained in the inner branchiæ alone, that there has been a gradual development from these primitive forms with simple, dull-colored, smooth shells, those of the male and female being alike, with radially sculptured beaks, the *Endobranchs*, up to the highest forms of to-day, with concentric, doubly looped beak sculpture, with highly painted shells, in which those of the male and female are very different, with the young contained in distinctly marked ovisacs in the hinder part of the outer gills alone, the *Exobranchs*.

The data for following these developments and the migrations of the *Naiades* are meager so far as fossil material is concerned. But, fortunately, while among the higher orders of life genera and even families appear, develop, grow old, and become extinct in a single geological age, the Unionidæ have held on unbroken from the Triassic or probably an earlier geological age until now, and while there has been slow progress in the development of higher characters *the primitive forms have not died out*. I know of no important type of the family among the fossil species that may not be found somewhere to-day among the living ones. They seem to have migrated to a certain region, made a slight advance over the characters of their predecessors, and to have continued down with but little change until to-day. When a new migration was made the same thing was enacted again.

If the Unionidæ originated in North America during the Triassic or some earlier period we may suppose that some members of the family migrated into South America during that or at a later period. All the species of that family in South America have radial beak sculpture (except *Callonaia* and *Prisodon*, in which the beaks seem to be smooth), and the young are contained in the inner gills alone, so far as we know. In some cases this sculpture is strictly radial; more often we find the central or all the bars curving a little toward each other below, and one or two of the middle pairs coalescing, the first move toward concentric beak sculpture. By an old, now partly submerged land bridge in the Antarctic region it is probable that a migration took place from South

¹ Proc. U. S. Nat. Mus., XVIII, 1896, pp. 381-385.

America to New Zealand and Australia, and slight changes in the way of progress are shown there. The beak sculpture is still radial, but the bars are decidedly curved toward each other below; they become slightly broken or nodulous, and each set is generally removed from the other a little, so that a space in the middle of the young shell is smooth. The young are contained in the inner gills, though Suter reports a few in the outer gills of one or two species and in all these Southern Hemisphere forms the shell has a dull color, and is nearly or quite rayless. Another migration took place to Southeastern Asia, and from there there was another to tropical Africa, possibly from Southern India over an old but now lost landway. In the forms of this region the beak sculpture becomes irregularly and variously zigzag radial, the surface is nodulous in many cases, and the shell is often bright colored and painted. Some of the recent species, such as *Unio crispisulcatus*, show nearly strict radial sculpture all over the shells: others which I have placed in the genus *Lamellidens* have almost such sculpture as is seen in the beaks of the Australian forms. In others the sculpture becomes slightly nodulous. Reasoning from analogy, and the few gravid specimens I have examined of these forms with the zigzagged beak sculpture, I presume that they all carry the young in the inner gills, though it is quite probable that some of them may have them in all four gills, and this would be a step in an upward direction. Some of these Asiatic and African forms with zigzag sculpture are quite short, inflated, and solid, and, in general, show characters strongly allied to those of our heavy Mississippi Valley forms (the latter of which I place in *Quadrula*), the various members of the genus *Parreysia*, for example.

It is quite probable that from such forms the genus *Quadrula* sprung; that it developed in Asiatic streams, where it still seems to be represented. In the Tertiary strata of eastern Europe and in parts of Asia this genus seems to be abundantly represented. It is a little difficult to say from some of the living Asiatic forms whether the beak sculpture should be called sharply, doubly looped, or zigzag radial, and the same thing may almost be said of some of our North American forms. There probably existed at that period an old land way across from northeastern Asia to northwestern North America, and one which lasted a long time, or it may have been submerged and then reappeared again, for through long ages this has apparently been a highway for migrating Unionidæ. I think it not unlikely that the immediate progenitors of the magnificent and diversified series of Uniones found fossil in the Laramie beds came over from Asia among the earlier migrations; for it is a significant fact that among the Uniones of the Laramie formation in the United States we have a number of species which in general form closely resemble these recent Asiatic *Parreysias*, and that their beak sculpture is decidedly zigzag radial, just as it is to-day in these oriental forms. Associated with these Laramie species are others in

which the beak sculpture is somewhat concentric and of a peculiar pattern seldom found among living Uniones, the ridges being nearly straight below and sharply pointed behind in a chevron-shaped pattern.

I take it that the next step of progress was the development of the genus *Pleurobema*, or forms which are nearly allied to it; though the strange, smooth, elongated, wedge-shaped *Unceopsis*, with its remarkably deep beak cavities and often distorted shells, and the curious *Gibbosula* seem like *Quadrulas* making an effort to shape their shells something like those of *Unio*. In *Pleurobema* the shell is solid and rather triangular, the beaks are high and typically well forward, the beak sculpture is broken, but most nearly concentric, the beak cavities are shallow, and the young occupy only the outer branchiae. There are great numbers of species I should refer to this or closely allied genera in the oriental tertiaries, and many of them are knobbed. To-day the genus is confined to the Mississippi Valley and the northern drainage of the Gulf of Mexico, and all the species are smooth except *P. asopus* and *P. varicosa*, which I place in the genus with some hesitation.

This genus has an immense development in the waters of the Tennessee drainage, becoming less abundant as we advance northward, until in the Ohio River and its tributaries only three species are found, *Pleurobema clava* Lamarek, *P. varicosa* Lea, and *P. asopus* Green. In the Tennessee and Cumberland systems all the species except the two last mentioned belong to the group typified by *P. clava*. In the Alabama River drainage and a few adjacent streams emptying into the Gulf there is a great development of this genus, but none of the species belong to the *clava* group. Now, if the genus migrated into the Tennessee system from the northwest, and it seems quite probable that it did, its presence in the Alabama River system would seem to prove the theory entertained by some geologists that the Tennessee formerly emptied into the Gulf, flowing down into what is now the Coosa River and into the Alabama. Additional proof of this is found in the fact that quite a number of the common Mississippi Valley species, such as *Unio tuberculatus* Barnes, *U. rectus* Lamarek, *U. trigonus* Lea, *U. anodontoides* Lea, *U. ebenus* Lea, and others are met with in the Tennessee, the Coosa, and the Alabama River systems, and *Unio conradiensis* and *U. cumberlandianus*, Tennessee River forms, occur in the Alabama system. That this southern outflow of the Tennessee ceased a long time ago is shown by the fact that in the Alabama drainage all the northern species of Uniones have a peculiar aspect, and several entire groups of *Pleurobema* have developed, which are distinct from anything found in the Tennessee.

It is probable that the true Unios and their allies were next developed, with simple oval to elongated shells, and moderate beaks showing quite a variety of sculpture. It is generally concentric, being inclined to follow the growth lines, but sometimes the ridges are broken, and again they are somewhat doubly looped. The young occupy the outer

gills only, forming a pad-like marsupium. The genus is now found throughout a large part of the Palearctic region and North America generally, excepting the Pacific drainage. The *Unios* of what seems to be the *Gibbosus* group are abundant in the Tertiary of eastern Europe. *Unio davilai* from Roumania is almost exactly like *U. gibbosus*, a recent species of the Mississippi Valley, and there were species no doubt closely allied in the Tertiary of the western United States. There were forms in Eastern Europe which seemed to connect *Unio* and *Pleurobema*, some of which were like *P. clava* and formed a connecting link with *Quadrula*. There were *Quadrulas* showing relationship with our *Q. trigona*, *rubiginosa*, *pustulosa*, *pyramidata*, and other forms. *Psilunio craviorensis* looks much like our *Pleurobema cesopus* and there are other striking examples of old forms resembling recent ones.

Now, in *Quadrula*, *Unio*, *Alasmidonta*, and some of the other allied genera we not unfrequently see species or specimens which have the ordinary concentric or doubly looped beak sculpture, and in addition to this a number of fine, regularly radiating ridges in front or behind, or sometimes at each end of the ordinary sculpture, an atavistic development, probably. In some groups this character is quite constant; in others it is only occasionally seen. I have never noticed it in any of the higher genera in which the male and female shells are different.

Up to this point the male and female shells had been essentially alike, and the ovisacs were not distinctly outlined, but formed smooth pads, the marsupium occupying practically the whole gill. In the American waters probably, a great change now began to take place in some of the Unionidae. The ovisacs of many forms became distinctly marked out by a deep sulcus; a constriction developed in some instances around near the base of the marsupium, inside and out, so that the lower end of the ovisacs resembled little bulbs, and these were often filled with pigment of a different color from the rest of the ovisac. *Ptychobranchnus*, with its folded marsupium, may have developed from *Unio*; *Cyprogenia* and *Obliquaria*, with their narrow central marsupia, and *Dromus*, in which it is hung all around the outer gills in short, distinct ovisacs, would seem to have descended from *Quadrula* by their shell characters. In all of these the shells of males and females are essentially alike. In *Medionidus* there is sometimes quite a well-marked swelling at the central or post-basal part of the female shell which is absent in the male, but sometimes it is impossible to separate the shells of the two sexes. The marsupium consists of irregular ovisacs just behind or almost at the center of the outer gills. In the strange *Tritogonia* the female shell differs remarkably from that of the male, having developed a broad, posterior wing, filled with a flap of the mantle, while that of the male is swollen, and truncate behind. Of the marsupia of this strange form I know nothing. *Obovaria*, with short, solid, full shells, sometimes having a slight post-basal inflation in those of the female, and *Plagiola*, with triangular shells, often showing the distinction between male and female, recall to some extent *Quadrula* in shell characters, but have the marsupia in the hinder part of the outer gills alone. The ovisacs are distinct, and there is a sulcus below, and this

is the highest type of a branchial uterus. *Lampsilis* is a step farther on, with oval to elongated shells, generally highly colored, with, as a rule, a decided post-basal swelling of that of the female, without (in most cases) a strong post ridge, and the beak sculpture consists of rather numerous, delicate bars, arranged in two distinct (anterior and posterior) loops.

Throughout the entire Mississippi Valley, a portion of the Atlantic drainage, in most of the streams flowing into the Gulf of Mexico, in eastern Mexico, and Central America there is found an extensive group of Uniones for which I have used the name *Proptera* of Rafinesque. Nearly all the shells are more or less dorsally winged, the beak sculpture is feeble, the epidermis is dull, the teeth are often imperfect, and they have a row of dorsal scars running obliquely downward and forward. The nacre generally is some shade of purple, and though the female shell may show a decided post inflation, it may be wanting. The ovisacs are usually fine and numerous, and are placed in the posterior ends of the outer gills. I have given this group subgeneric rank under *Lampsilis*. Now, it would seem probable that North America was the field in which these remarkable developments in the character of the Unionidæ had taken place, for I know of no fossil species elsewhere showing any of these higher characters. If, as I have elsewhere suggested, there is a relationship between *Proptera* and the *Cristarias*, *Hyriopsis*, *Pilsbryconcha*, *Chamberlainia*, and *Pseudospatha* of the Old World, then I should think it likely they, or their progenitors, had migrated thence from North America some time during the early or middle tertiaries. It is possible that the *Cristarias*, *Hyriopsis*, and the like, or closely related but now extinct groups, may have originated in oriental waters, and that the *Propteras*, and the typical *Lampsilis*, may be their offspring.

Truncella marks the highest development of Naiad life, and may be taken to be the latest. Its shells are smooth and generally highly painted, the beak sculpture is fine and doubly looped, the hinge teeth are well developed. The post-basal area is very distinctly marked out and developed in the female. In most cases the shell of this part in the female is thin, of a different texture from the rest, often radially ridged, and decidedly toothed on its edge. There is usually a surprising difference in the shape of the shells of male and female aside from this, so much so that one would never suppose that they belonged to the same species, unless he traced the growth back to the young shells, which are quite alike in the different sexes. The marsupium is like a large kidney, very full, and totally different in appearance when gravid from the rest of the gill. It appears to be protected by a great flap which grows out from the mantle covering it, which is here double. The genus exhibits a great variety of form, so much so that several good subgeneric groups seem well marked out, and we must believe it has been in existence for some time.

Much of the foregoing may be mere conjecture; much is undoubtedly founded on fact. I believe that the living forms of the Unionidæ show a gradual development from the simplest, lowest, and earliest

types up to the highest, most recent, and most complicated, and stand as a sort of index to the progress of the family in the past.

I have not been able to study the Mutelidæ as carefully as I have the Unionidæ. All the species are foreign, and while I have seen the soft parts of a few South American forms, I have never had the opportunity to examine the anatomy of a single African species. I have not been able, from what knowledge I have obtained, to discover any considerable anatomical or conchological differences between the Mutelid genera of Africa and South America which might be used to separate the family into subfamilies.

I have treated the families of the Unionidæ and Mutelidæ together in this work because they have both been classed as Naiades. But the remarkable differences in the embryos, that of the former family being a glochidium with a bivalve shell inclosing the soft parts, and that of the latter a lasidium divided into three segments with a single shell on its middle section, and the shells of the one family having schizodont teeth, while those of the other have taxodont teeth, seem to show that the two great groups are not very nearly related.

In studying the Naiades I have been greatly perplexed on account of the unsatisfactory and confused condition of much of the literature. Early in this century Rafinesque collected the Unionidæ extensively in Kentucky and published a large number of genera, minor groups, and species.

It is probable that there will always be differences of opinion with regard to his work, just as there have been in the past. His figures are more like those made by children, or the caricatures drawn by aboriginal tribes, than the creations of an intelligent naturalist, and the descriptions are too brief in many cases to give any clear idea of the species. The work in the continuation of his monograph is even worse than that in the body of the paper, and tribes, genera, and subgenera are mingled in bewildering confusion, and all these are placed under two great genera. It is impossible in many cases to tell what his meaning is. I have carefully gone over his so-called types in the Academy of Natural Sciences in Philadelphia, making notes and sketches, and stating the names by which they are commonly known. Four times, separated by long intervals, I have studied his original descriptions, with specimens of the same species as these so-called types, in an attempt to determine his species, each time writing down my conclusions, and I believe that quite a number of his species will have to stand. There can be no doubt whatever that many of his so-called types are not types at all, as they do not fit the descriptions by any stretch of imagination.

In applying generic, subgeneric, and sectional names I have used those which have been previously applied wherever it has been possible to do so by the process of elimination, in order that no claim of priority might be urged for them.

A most unfortunate dispute arose among our earlier conchologists in regard to priority of names, one in which a great deal of ill feeling was displayed. I have endeavored wherever it was possible to ascertain the exact date of publication of these disputed names and to credit the species to the earliest described. In some cases it has been

impossible to tell which of two names appeared first, and in such cases I have used that which seemed to be best known.

Many of the species of Lamarck stand on about the same foundation as those of Rafinesque, having only a brief description and no figures. Dr. Lea on at least two occasions went over the types of most of Lamarck's species and has published his conclusions in the Observations, and I am obliged to abide by his decisions, never having seen the types myself.

One of the most perplexing problems has been the work done by European conchologists, and especially by the so-called new school of France. Previous to 1870, the date of issuance of Lea's last edition of the synopsis, thousands of names had been applied to the few species of Europe. But this work was conservative and reasonable compared with that of the new school since then. I have devoted much time and study to this fauna and its literature. To me it seems that there are not more than eighteen or twenty species of Unionidae found in Europe, judging by the same standards I have applied to species elsewhere. Nearly all the authors seem to be more or less at sea as to certain forms of this area, and the reasons for this are probably their want of striking characters and their extreme variability. *Unio pictorum*, *tumidus*, *elongatulus*, and *platyrhynchoideus* have been often taken for each other. *Margaritana margaritifera* has repeatedly been mistaken for the very similar looking *M. crassa*, and each of these has been hopelessly confused with *Unio batavus*.

I have endeavored to consider names applied before 1870 in my synonymy. I have made no effort to do this with those applied by authors to the forms of Europe since that time, as I do not believe that any new species of *Unio*, *Margaritana*, *Anodonta*, or *Leguminaia* have been found there in the last thirty years. The genus *Colletopterum* (1881) is a doubtful one and is very likely only an mutation of the excessively variable *Anodonta cygnea*. In 1892 Arnould Locard, one of the great lights of the new school, stated that there were 208 species of Unios and 250 Anodontas in France alone.¹

Life is too short and valuable to be wasted in any attempt at deciphering such nonsense, and I have not even cumbered the pages of this work with a list of these new species. Those interested can find them in the works of Westerlund and Kobelt.

In cases where the new school has worked on the fauna of Africa or Asia, I have done the best I could to straighten out the synonymy.

I have not attempted to make any analytical key to genera and other groups, because I do not believe that it is possible to construct a key that will be of any real service to the student. At least four-fifths of the shells of the Naiades have the beaks so eroded that it is impossible to form any idea of what their sculpture is like, and the soft parts are inaccessible to the average student and collector. These are both vital characters that must be used in classification. The general arrangement of the groups down to genera is shown in the following table:

¹ Ann. Soc. Ag. Hist. Nat. Lyon, 1892, p. 55.

SYNOPSIS OF THE GENERA OF NAIADES.

Family Unionidae. Hinge with schizodont teeth; embryo a <i>glochidium</i> .	Subfamily Unioninae. Beak sculpture concentric; embryos in outer or all four of the gills. (<i>Exobranchia</i> .)	Ovisacs distinctly marked out by a sulcus.	Male and female shells different.	(Heterogenæ.) Female shell inflated at base in front of posterior ridge; marsupium in hinder part of outer gills.	{ <i>Truncilla</i> Rafinesque. <i>Micromya</i> Agassiz. <i>Lampsilis</i> Rafinesque. <i>Pseudospatha</i> Simpson. <i>Hyriopsis</i> Conrad. <i>Chamberlainia</i> Simpson. <i>Cristaria</i> Schumacher. <i>Lepidodesma</i> Simpson. <i>Pilsbryconcha</i> Simpson. <i>Medionidus</i> Simpson. <i>Nephronaias</i> Crosse and Fischer. <i>Glebula</i> Conrad. <i>Obovaria</i> Rafinesque. <i>Plagiola</i> Rafinesque.
				(Digenæ.) Female shell inflated behind, above posterior ridge.	{ <i>Tritogonia</i> Agassiz.
				(Mesogenæ.) Marsupium a few ovisacs in the center of outer gills.	{ <i>Cyprogenia</i> Agassiz. <i>Obliquaria</i> Rafinesque.
				(Ptychogenæ.) Marsupium occupying the whole outer gills in a series of folds.	{ <i>Ptychobranchus</i> Simpson.
				(Eschatigenæ.) Marsupium occupying the outer border of outer gills.	{ <i>Dromus</i> Simpson.
				(Diagenæ.) Ovisacs filling the outer gills, running crosswise.	{ <i>Strophitus</i> Rafinesque.
		Ovisacs not separated by a sulcus.	Male and female shells alike.	(Homogenæ.) Marsupium filling the entire outer gills and forming smooth pads.	{ <i>Anodonta</i> Bruguiere. <i>Colletopterum</i> Bourguignat. <i>Gabillotia</i> Servain. <i>Leguminaia</i> Conrad. <i>Lastena</i> Rafinesque. <i>Solenia</i> Conrad. <i>Gonidea</i> Conrad. <i>Anodontoides</i> Simpson. <i>Pegias</i> Simpson. <i>Arcidens</i> Simpson. <i>Symphynota</i> Lea. <i>Alasmidonta</i> Say. <i>Hemilastena</i> Agassiz. <i>Margaritana</i> Schumacher. <i>Unio</i> Retzius. <i>Pleurobema</i> Rafinesque.
				(Tetragenæ.) Marsupium filling all four gills, forming smooth pads; beak cavities deep.	{ <i>Quadrula</i> Rafinesque. <i>Schistodesma</i> Simpson. <i>Gibbosula</i> Simpson. <i>Cuneopsis</i> Simpson.

Family Unionidae. Hinge with schizodont teeth; embryo a <i>glochidium</i> .	Subfamily Hyrianae. Beak sculpture radial; male and female shells alike; marsupium filling the inner gills. (<i>Endobranchia</i> .)	(Rosanorhamphus.) Beak sculpture generally zigzag-radial; epidermis often bright.	<i>Nodularia</i> Conrad. <i>Harmandia</i> Rochebrune. <i>Grandidieria</i> Bourguignat. <i>Physunio</i> Simpson. <i>Dallia</i> Simpson. <i>Pseudodon</i> Gould. <i>Parreysia</i> Conrad. <i>Ptychorhynchus</i> Simpson. <i>Virgus</i> Simpson. <i>Ctenodesma</i> Simpson. <i>Rectidens</i> Simpson. <i>Lamellidens</i> Simpson. <i>Trapezoides</i> Simpson. <i>Arconaia</i> Conrad. <i>Pseudavicula</i> Simpson. <i>Arcidopsis</i> Simpson.
Family Mutelidae.		(Lamphorhamphus.) Beak sculpture radial, often curved; shell dull colored.	<i>Tetraplodon</i> Spix. <i>Castalina</i> von Ihering. <i>Castaliella</i> Simpson. <i>Callonaia</i> Simpson. <i>Hyria</i> Lamarek. <i>Prisodon</i> Schumacher. <i>Diplodon</i> Spix. <i>Spatha</i> Lea. <i>Mutela</i> Scopoli. <i>Chelidonopsis</i> Ancey. <i>Brazza</i> Bourguignat. <i>Pleiodon</i> Conrad. <i>Monocondylaea</i> d'Orbigny. <i>Iheringella</i> Pilsbry. <i>Fossula</i> Lea. <i>Glabaris</i> Gray. <i>Leila</i> Gray. <i>Mycetopoda</i> d'Orbigny.

In the preparation of this work I have received uniform courtesy and kindness from students and collectors generally. I am under especial obligations to Mr. Edgar A. Smith, of the British Museum, for examining and reporting on the original manuscript of Solander, and to Mr. H. A. Pilsbry, of the Academy of Natural Sciences of Philadelphia; Mr. A. G. Wetherby, of Magnetic City, North Carolina; Dr. H. von Ihering, Museo Paulista, San Paulo, Brazil; and Mr. H. Suter, Christ Church, New Zealand, for repeated assistance and advice. Mr. L. E. Daniels, of La Porte, Indiana; Mrs. George Andrews, of Knoxville, Tennessee; Hon. J. D. Mitchell, of Victoria, Texas; Mr. Berlin H. Wright, of Penn Yan, New York; Mr. Bryant Walker, of Detroit, Michigan; Dr. W. S. Strode, of Lewistown, Illinois; Mr. H. M. Kelly, of Mount Vernon, Iowa; Mr. William Moss, of Ashton-under-Lyne, England; and many others have contributed valuable material for study.

SYSTEMATIC ARRANGEMENT OF THE NAIADES.

Family UNIONIDÆ.

Shell nacreous, with a thick epidermis; beaks usually sculptured, often showing the remains of the nuclear shell; ligament opisthodontic; hinge with or without teeth, though with vestiges of them in every genus; when present schizodont and arranged as pseudocardinals and laterals; pallial line usually simple; prismatic border ordinarily narrow.

Animal with labial palpi almost always wider than long; anal opening usually separated from the superanal; embryo a glochidium, the soft parts being inclosed in a bivalve shell, and borne in the inner or outer or all four leaves of the branchiæ.

Subfamily UNIONINÆ Swainson, 1840. (Exobranchiæ.)

Shell having essentially concentric beak sculpture.

Animal with labial palpi somewhat drawn out, projecting posteriorly; embryos borne in the outer or in all four gills.

HETEROGENÆ.

Male and female shells different, the latter inflated in the post-basal region; beak sculpture usually doubly looped; embryos contained in ovisacs separated by a sulcus and occupying the hinder part of the outer gills.

Genus TRUNCILLA Rafinesque, 1819.

(Type, *Truncilla triqueter* Rafinesque.)

Truncilla RAFINESQUE, J. de Phys., Chimie, Hist. Nat., LXXXVIII, 1819, p. 427.

Shell rounded or oval, solid, inflated, generally smooth and rayed, with delicate beak sculpture which has a tendency to be doubly looped, that of the female having a very decided inflation in the post-basal region, which is thinner than the rest of the shell, of different texture, often toothed, and usually radiately sculptured; laterals double in each valve, the inner in the right valve smaller. Animal generally having the inner gills united to the abdominal sac; female with a heavy flap of the mantle which fills the post-basal swelling of the shell, and which has an inner ridge inside at some distance above the edge; marsupium very distinct, occupying the swollen part of the shell.

Subgenus TRUNCILLA Rafinesque, 1819.

Shell covered with broken rays, somewhat triangular, and without a wide, radiate, posterior furrow.

(Group of *Truncilla triquetra*.)

Shell greatly inflated, sharply truncate posteriorly; inflation of the female shell at extreme post-basal point.

† TRUNCILLA TRIQUETRA Rafinesque.

* *Truncilla triquetra* RAFINESQUE, Ann. Gen. Sci. Phys. Brux., XIII, 1820, p. 300, pl. LXXXI, figs. 1-4.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 15, pl. II, fig. 1.—* AGASSIZ, Arch. für. Naturg., I, 1852, p. 44.

Unio triquetra SHORT and EATON, Transylvania Jl., 1831, p. 79.—* SAY, Am. Conch., VI, 1834.—* CONRAD, New F.W. Shells, 1834, p. 72.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* SAY, Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* PÆTEL, Conch. Sam., III, 1890, p. 170.

* *Unio triangularis* BARNES, Am. Jl. Sci., VI, 1823, p. 272, pl. XIII, fig. 17.—* HILDRETH, Am. Jl. Sci., X, 1828, p. 287, fig. —* SAY, Am. Conch., No. 1, 1830, pl. IV.—SHORT and EATON, Transylvania Jl., 1831, p. 78.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 554; 3d ed., II, 1839, p. 674.—* REEVE, Conch. Syst., I, 1841, p. 118, pl. LXXXIX, fig. 9.—* HANLEY, Test. Moll., 1842, p. 183; *Biv. Shells, 1843, p. 183.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 10, pl. I, figs. 4, 4b; *Man., II, 1859, p. 137, fig. 662.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXVII, fig. 340.—* CALKINS, Pr. Ottawa Acad. Nat. Sci., 1874, p. 45.

* *Mya triangularis* EATON, Zool. Text Book, 1826, p. 221.

* *Margarita (Unio) triangularis* LEA, Syn., 1836, p. 18; 1838, p. 16.

* *Margaron (Unio) triangularis* LEA, Syn., 1852, p. 23; 1870, p. 36.

* *Unio cuneatus* SWAINSON, Phil. Mag., 1823, p. 112.

* *Unio formosus* LEA, Trans. Am. Phil. Soc., IV, 1834, p. 111, pl. XVI, fig. 41; *Obs. I, 1834, p. 121, pl. XVI, fig. 41.—* CHENU, Ill. Conch., 1858, pl. VIII, figs. 12, 12a, 12b; *Man., 1859, II, p. 138, fig. 664.

Ohio River drainage; western New York to southern Michigan; Iowa; eastern Nebraska to Indian Territory.

(Group of *Truncilla brevidens*.)

Shell somewhat quadrate, not sharply truncate behind; post-basal swelling of female in front of post-basal point and rounded below.

† TRUNCILLA BREVIDENS Lea.

Unio brevidens LEA, Trans. Am. Phil. Soc., IV, 1834, p. 75, pl. IV, fig. 6; *Obs. I, 1834, p. 85, pl. VI, fig. 6.—* HANLEY, Test. Moll., 1842, p. 198; *Biv. Shells, 1843, p. 198, pl. XX, fig. 41.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* CHENU, Ill. Conch., 1858, pl. VIII, figs. 6, 6a, 6b.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

* *Margarita (Unio) brevidens* LEA, Syn., 1836, p. 29; 1838, p. 21.

* *Margaron (Unio) brevidens* LEA, Syn., 1852, p. 32; 1870, p. 50.

†A † placed before a species indicates that the type, or what are believed to be authentic specimens have been examined by the writer.

*An * in front of a reference indicates that it has been verified.

- * *Unio interruptus* CONRAD, New F. W. Shells, 1834, p. 69.—* SAY, Am. Conch., VI, 1834.—* CONRAD, Monog., X, 1838, p. 88, pl. XLVIII; * Pr. Ac. N. Sci., Phila., VI, 1853, p. 250.—* KUSTER, Conch. Cab. Unio, 1861, p. 182, pl. LVII, fig. 3; pl. LX, fig. 2.—* REEVE, Conch. Icon., XVI, 1864, pl. XIV, fig. 56.
 * *Truncilla interrupta* AGASSIZ, Arch. für Naturg., I, 1852, p. 44.

Tennessee drainage.

† TRUNCILLA PENITA Conrad.

- * *Unio penitus* CONRAD, New F. W. Shells, 1834, pp. 33, 70, pl. v, fig. 1.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen., 1836, p. 203.—* HANLEY, Test. Moll., 1842, p. 183; * Biv. Shells, 1843, p. 183.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 18, pl. III, fig. 9.—* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 254.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* REEVE, Conch. Icon., XVI, 1864, pl. XIV, fig. 55.—* PÆTEL, Conch. Sam., III, 1890, p. 162.
 * *Margarita (Unio) penitus* LEA, Syn., 1836, p. 19; 1838, p. 16.
 * *Margaron (Unio) penitus* LEA, Syn., 1852, p. 24; 1870, p. 36.
 * *Unio compactus* LEA, Proc. Ac. N. Sci. Phila., III, 1859, p. 154; * Jl. Ac. N. Sci. Phila., IV, 1859, p. 218, pl. XXVIII, fig. 98; * Obs. VII, 1859, p. 36, pl. XXVIII, fig. 98.
 * *Margaron (Unio) compactus* LEA, Syn., 1870, p. 36.

Alabama and Tombigbee river drainage.

† TRUNCILLA MODICELLA Lea.

- * *Unio modicellus* LEA, Pr. Ac. Phila., III, 1859, p. 171; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 347, pl. LVII, fig. 172; * Obs., VIII, 1860, p. 29, pl. LVII, fig. 172.—* B. H. WRIGHT, Check List, 1888.
 * *Margaron (Unio) modicellus*¹ LEA, Syn., 1870, p. 41.

Connasauga and Chattanooga rivers, northwest Georgia.

The following species² are unfigured and indeterminate:

Truncilla (Unio) perplexus Rafinesque.

Truncilla granulatus Rafinesque.

Unio (Truncilla) metaplata Rafinesque.

† TRUNCILLA LENIOR Lea.

- * *Unio lenis* LEA, Pr. Am. Phil. Soc., I, 1840, p. 286.³
 * *Unio lenior* LEA, Tr. Am. Phil. Soc., VIII, 1843, p. 204, pl. XII, fig. 18; * Obs., III, 1842, p. 42, pl. XII, fig. 18.—* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 251.—* CHENU, Ill. Conch., 1858, pl. XXVII, figs. 4, 4a, 4b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* REEVE, Conch. Icon., XVI, 1864, pl. XVII, fig. 75.—* PÆTEL, Conch. Sam., III, 1890, p. 157.
 * *Margaron (Unio) lenior* LEA, Syn., 1852, p. 39; 1870, p. 62.

Stone River, Tennessee; near Woodville, Alabama.

¹Four specimens of this species are in the U. S. National Museum collection, from the Connasauga River, Georgia, three of which are males, with a form something like that of *T. penita*; the fourth is a young female, which distinctly shows a radiately ridged post-basal swelling.

²These are in continuation of Monography, 1831, p. 4.

³Lea changed his name *lenis* to *lenior*, probably because Conrad had previously used the former for a species of *Unio*, which is no doubt a young *ventricosus*.

†TRUNCILLA METASTRIATA Conrad.

* *Unio metastriatus* CONRAD, Mon., XII, 1840, p. 104, pl. LVII, fig. 2.

Black Warrior River and Woodville, Alabama.

(Group of *Truncilla arcaiformis*.)

Shell greatly inflated, with a sharp posterior ridge, and two lesser ridges above it; inflation of the female flattened on its base.

†TRUNCILLA ARCÆFORMIS Lea.

* *Unio arcaiformis* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 116, pl. XVII, fig. 44; *Obs., I, 1834, p. 126, pl. XVII, fig. 44.—*ANTON, Verz. der Conch., 1839, p. 14.¹—DESHAYES, Tr. Element, Conch., 1839, p. 18, pl. XXIX, fig. 6.—*HANLEY, Test. Moll., 1842, p. 182; *Biv. Shells, 1843, p. 182, pl. XXII, fig. 40.—*CATLOW and REEVE, Conch. Nom., 1845, p. 55.—*DESHAYES, Traité Element., II, 1853, pl. XXIV, fig. 6.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU, Ill. Conch., 1858, pl. XII, figs. 3, 3a, 3b.—*REEVE, Conch. Icon., XVI, 1864, pl. XIV, fig. 57.

**Margarita (Unio) arcaiformis* LEA, Syn., 1836, p. 18; 1838, p. 16.

Margaron (Unio) arcaiformis LEA, Syn., 1852, p. 23; 1870, p. 36.

**Truncilla arcaiformis* AGASSIZ, Arch. für Naturg., I, 1852, p. 44.

Unio nexus SAY, Transylvania Jl., IV, 1831, p. 527; *SAY, Am. Conch., Pt. 6, 1834, pl. LI.²—*CONRAD, New F. W. Shells, 1834, p. 70.—*FERUSSAC, Guer. Mag., 1835, p. 28.—*CONRAD, Monog., VIII, 1837, p. 68, pl. XXXVIII, fig. 1.—*KUSTER, Conch. Cab. Unio, 1854, p. 69, pl. XVII, fig. 2.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—*PÆTEL, Conch. Sam., III, 1890, p. 60.

Tennessee and Cumberland river systems.

Subgenus SCALENARIA (Rafinesque, 1820) Agassiz.

(Type, *Unio sulcatus* Lea.)

Male shell having a wide, radiating, shallow depression in front of the posterior ridge; that of the female having a small, rounded, well-defined, radial post-basal swelling. Animal with marsupium occupying the greater part of the outer branchiæ; mantle enlarged below; branchial opening, with few papillæ.

(Group of *Truncilla sulcata*.)

Shell elliptical, beaks high and well forward, male shell not radiately striated posteriorly.

¹This may be *T. triquetra*.

²According to Binney, *Unio nexus* was first published in December, 1831, (Bib. N. Am. Conch., I, p. 266), and Seudder states that the fourth volume of the Transactions, containing Lea's description of *Unio arcaiformis*, was published at the latter end of 1831, and acknowledged by correspondents as received that year (Bib. of publications of Lea, 1885, p. 3). As I am not able to say which has precedence, I prefer retaining the well-known name of Lea rather than that of Say, which is less known.

†*TRUNCILLA SULCATA* Lea.

* *Unio sulcatus* LEA, Tr. Am. Phil. Soc., III, 1830, p. 430, pl. VIII, fig. 2.—*SAY, Am. Conch., No. 1, 1830, pl. v.—SHORT and EATON, Transylvania Jl., 1831, p. 75.—*LEA Obs., I, 1834, p. 44, pl. VIII, fig. 12.—*HANLEY, Test. Moll., 1842, p. 188; *Biv. Shells, 1843, p. 188.—*CHENU, Bib. Conch., 1st ser., III, 1845, p. 11, pl. I, figs. 5, 5a.—*CATLOW and REEVE, Conch. Nom., 1845, p. 64.—*CHENU, Ill. Conch., 1858, pl. XIII, figs. 3, 3a, 3b.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—*REEVE, Conch. Icon., XVI, 1864, pl. XIV, fig. 60.—*PETEL, Conch. Sam., III, 1890, p. 169.

**Margarita (Unio) sulcatus* LEA, Syn., 1836, p. 22; 1838, p. 18.

**Margaron (Unio) sulcatus* LEA, Syn., 1852, p. 26; 1870, p. 40.

**Unio sulcata* DESHAYES, An. sans Vert, 2d ed., VI, 1835, p. 548; 3d ed., II, 1839, p. 672.

Unio ridibundus SAY, New Harm. Diss., II, 1829, p. 308; *Am. Conch., I, 1830, pl. v.—SHORT and EATON, Transylvania Jl., 1831, p. 75.—*CONRAD, New F. W. Shells, 1834, p. 71.—*CHENU, Bib. Conch., 1st ser., III, 1845, p. 11, pl. I, fig. 6.—*KUSTER, Conch. Cab. Unio, 1862, p. 276, pl. XCIII, fig. 2.—*REEVE, Conch. Icon., XVI, 1864, pl. xv, fig. 63.

**Unio obliquatus* SAY, Am. Conch., VI, 1834.—*CONRAD, New F. W. Shells, 1834, p. 70.—*KUSTER, Conch. Cab., 1862, p. 274, pl. XCII, fig. 3.—*AGASSIZ, Arch. für Naturg., I, 1852, p. 43.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.

**Unio gibbosus* var. *perobliquus* CONRAD, Monog., VI, 1836, p. 50, pl. XXVII, fig. 2. *Unio perobliquus* CONRAD, Cover of Monog., VIII, 1837.

**Unio pectitus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255; Jl. Ac. N. Sci. Phila., 1854, p. 297, pl. XXVII, fig. 4.

TRUNCILLA SULCATA var. *DELICATA* Simpson.¹

Ohio River drainage; the variety in the Detroit River and southern Michigan.

(Group of *Truncilla haysiana*.)

Shell shining, round-oval; posterior part of both male and female distinctly radially sculptured.

†*TRUNCILLA HAYSIANA* Lea.

**Unio haysianus* LEA, Tr. Am. Phil. Soc., V, 1833, p. 35, pl. III, fig. 7; *Obs., I, 1834, p. 147, pl. III, fig. 7.—*HANLEY, Test. Moll., 1842, p. 188; *Biv. Shells, 1843, p. 188.—*CATLOW and REEVE, Conch. Nom., 1845, p. 59.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—*KUSTER, Conch. Cab. Unio, 1856, p. 209, pl. LXIX, fig. 4.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*REEVE, Conch. Icon., XVI, 1864, pl. xv, fig. 62.—*PETEL, Conch. Sam., III, 1890, p. 154.

**Margarita (Unio) haysianus* LEA, Syn. 1836, p. 22; 1838, p. 18.

**Margaron (Unio) haysianus* LEA, Syn. 1852, p. 26; 1870, p. 41.

**Scalenaria haysiana* AGASSIZ, Arch. für Naturg., I, 1852, p. 48.

**Unio sowerbyanus* LEA, Tr. Am. Phil. Soc., V, 1839, p. 68, pl. x, fig. 28;² *Obs. I, 1834, p. 180, pl. x, fig. 28.—*FERUSSAC, Guer. Mag., 1835, p. 29.—*CONRAD,

¹ Mr. Bryant Walker has sent me specimens of a form which I refer to this species that are smaller and in every way more delicate. I propose the above varietal name for it.

² Male of *T. haysiana*. *Unio haysianus* and *U. sowerbyanus* were published in the same paper, the former having precedence.

Monog., VIII, 1837, p. 66, pl. XXXVII, fig. 1.—* HANLEY, Test. Moll., 1842, p. 185; *Biv. Shells, 1843, p. 185.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* KUSTER, Conch. Cab. Unio, 1852, p. 62, pl. XIV, fig. 3.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* REEVE, Conch. Icon., XVI, 1864, pl. XIV, fig. 58.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

**Margarita (Unio) sowerbyanus* LEA, Syn. 1836, p. 20; 1838, p. 17.

**Margaron (Unio) sowerbyanus* LEA, Syn. 1852, p. 25; 1870, p. 38.

Tennessee and Cumberland river systems.

†*TRUNCILLA OTHCALOOGENSIS* Lea.

**Unio othcaloogensis* LEA, Proc. Acad. N. Sci. Phila., IX, 1857, p. 32; ¹ Jl. Ac. N. Sci. Phila., IV, 1858, p. 74, pl. XIV, fig. 54; Obs. VI, 1858, p. 75, pl. XIV, fig. 54.

**Margaron (Unio) othcaloogensis* LEA, Syn. 1870, p. 38.

Othcalooga Creek, Gordon County, Georgia.

Subgenus *DYSNOMIA* Agassiz. 1852.

(Type, *Unio foliatus* Hildreth.)

Shell of the male with a posterior and central radiating ridge, with a wide, flattened space between; that of the female with a greatly produced inflation, which is but little behind the center of the base, it being a continuation of the central ridge. Animal with mantle beautifully maculate on its border; female animal unknown.

†*TRUNCILLA FOLIATA* Hildreth.

**Unio foliatus* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 284, fig. 16.—* CONRAD, New F. W. Shells, 1834, p. 69.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* HANLEY, Test. Moll., 1842, p. 176; *Biv. Shells, 1843, p. 176.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Man., 1859, II, p. 143, fig. 705.—* HARPER, L. and F. W. Shells of Cinti., p. 4.

**Margarita (Unio) foliatus* LEA, Syn. 1836, p. 13; 1838, p. 14.

**Margaron (Unio) foliatus* LEA, Syn. 1852, p. 20; 1870, p. 30.

**Unio flexuosus* CONRAD, Monog. I, 1835, p. 8, pl. IV, fig. 2.—* KUSTER, Conch. Cab. Unio, 1852, pp. 46, 211, pl. IX, fig. 2; LXX, fig. 1.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—* REEVE, Conch. Icon., XVI, 1864, pl. VI, fig. 22; XIII, fig. 53.

**Dysnomia flexuosa* AGASSIZ, Arch. für Naturg. I, 1852, p. 43.

Ohio River drainage.

†*TRUNCILLA STEWARDSONI* Lea.

**Unio stewardsoni* LEA, Tr. Am. Phil. Soc., X, 1852, p. 278, pl. XXIII, fig. 36; * Obs. V, 1852, p. 34, pl. XXIII, fig. 36.—* CONRAD, Pr. Acad. Nat. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* REEVE, Conch. Icon., XVI, 1864, pl. XV, fig. 66.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

**Margaron (Unio) stewardsoni* LEA, Syn. 1852, p. 20; 1870, p. 38.

Tennessee River.

¹ I have seen only the type of this species, apparently a young female, which is a reddish buff-colored, shining shell, somewhat more elongated than *T. haysiana*, but which probably groups with it.

Subgenus *PILEA* Simpson, 1900.

(Type, *Unio personatus* Say.)

Male shell with a wide, shallow, radiating depression in front of the posterior ridge, that of the female with a rounded, foliaceous swelling at the posterior base. Animal with post basal flap of mantle of female very heavy; ovisacs not extending to the top of the marsupium.

(Group of *Truncilla personata*.)

Shell inflated, rather solid, nearly as high as long; female not having a central depression, post basal swelling small.

† *TRUNCILLA PERSONATA* Say.

Unio personatus SAY, New Harm. Diss. II, No. 20, 1829, p. 309.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* CONRAD, Monog. V, 1836, p. 47, pl. XXIV.—* HANLEY, Test. Moll., 1842, p. 202; * Biv. Shells, 1843, p. 232.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* KUSTER, Conch. Cab. Unio, 1852, p. 48, pl. X, fig. 1.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* REEVE, Conch. Icon., XVI, 1864, pl. XV, fig. 64.—* PETEL, Conch. Sam., III, 1890, p. 116.

* *Margarita (Unio) personatus* LEA, Syn. 1836, p. 33; 1838, p. 23.

* *Margaron (Unio) personatus* LEA, Syn. 1852, p. 35; 1870, p. 38.

* *Scalenaria personata* AGASSIZ, Arch. für Naturg. I, 1852, p. 43.

* *Unio pileus* LEA, Trans. Am. Phil. Soc., IV, 1834, p. 119, pl. XVIII, fig. 47.—* CONRAD, New F. W. Shells, 1834, p. 71.—* LEA, Obs. I, 1834, p. 129, pl. XVIII, fig. 47.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* HANLEY, Test. Moll., 1842, p. 185; * Biv. Shells, 1843, p. 185.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* CHENU, Ill. Conch., 1858, pl. XV, figs. 2, 2a, 2b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* REEVE, Conch. Icon., XVI, 1864, pl. XV, fig. 64.—* PETEL, Conch. Sam., III, 1890, p. 163.

* *Margarita (Unio) pileus* LEA, Syn. 1836, p. 20; 1838, p. 17.

* *Margaron (Unio) pileus* LEA, Syn. 1852, p. 24.

* *Unio capillaris* LEA, Tran. Am. Phil. Soc., V, 1834, p. 29, pl. II, fig. 2; * Obs. I, 1834, p. 141, pl. II, fig. 2.—* FERUSSAC, Guer. Mag., 1835, p. 29.

Ohio River drainage. Rare.

(Group of *Truncilla perplexa*.)

Shell of the male with median and posterior radiating ridges, both of which are usually somewhat nodose; postbasal expansion of the female rounded, large, thin, placed far back.

† *TRUNCILLA PERPLEXA* Lea.¹

* *Unio perplexus* LEA, Trans. Am. Phil. Soc., IV, 1831, p. 112, pl. XVII, fig. 42; * Obs. I, 1834, p. 122, pl. XVII, fig. 42.—* HANLEY, Test. Moll., 1842, p. 181; * Biv. Shells, 1843, p. 181, pl. XXII, fig. 39.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* CHENU, Ill. Conch., 1858, pl. XVIII, figs. 5, 5a, 5b.

* *Margarita (Unio) perplexus* LEA, Syn. 1836, p. 17; 1838, p. 16.

¹ Lea's species was published, according to Scudder (Bibliography of Isaac Lea, p. 3), in the latter part of 1831. Rafinesque published a species, "*Truncilla, Unio perplexus*," in the Continuation of a Monograph in October, 1831, which I can not identify. I do not know which name appeared first, but I think it best to retain the well-known and properly described name of Dr. Lea.

- **Margaron (Unio) perplexus* LEA, Syn., 1852, p. 23; 1870, p. 34.
 **Unio gibbosus* FERUSSAC, Gucr. Mag., 1835, p. 27.—CONRAD, Monog. VI, 1836, p. 50, pl. XXVII, fig. 1.—REEVE, Conch. Icon., XVI, 1864, pl. XI, fig. 41, 41a, 41b.
 **Dysnomia gibbosa* AGASSIZ, Arch. für Nat., I, 1852, p. 43.
 **Unio gibbosus*, Raf. v. *perplexus* PÆTEL, Conch. Sam., III, 1890, p. 153.
 **Unio torulosus* CONRAD, New F. W. Shells, 1834, p. 72; *Proc. Acad. N. Sci. Phila., VI, 1853, p. 259.
 **Unio obliquus* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 153, pl. LVIII, figs. 3, 4.

†**TRUNCILLA PERPLEXA** var. **RANGIANA** Lea.

- **Unio rangianus* LEA, Trans. Am. Phil. Soc., VI, 1839, p. 95, pl. XVIII, fig. 56;¹
 *Obs., II, 1838, p. 95, pl. XVIII, fig. 56.—*TROSCHER, Arch. für Naturg., V, 1839, p. 237.—*HANLEY, Test. Moll., 1842, p. 187; *Biv. Shells, 1843, p. 187, pl. XXI, fig. 48.—*CATLOW and REEVE, Conch. Nom., 1845, p. 63.—*CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 256.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU., Ill. Conch., 1858, pl. XXIV, figs. 5, 5a, 5b.—*KUSTER, Conch. Cab. Unio, 1861, p. 257, pl. LXXXVII, fig. 1; LXXXIX, fig. 1.—*REEVE, Conch. Icon., XVI, 1864, pl. XIV, fig. 54.—*PÆTEL, Conch. Sam., III, 1890, p. 164.
 **Margarita (Unio) rangianus* LEA, Syn., 1838, p. 18.
 **Margaron (Unio) rangianus* LEA, Syn., 1852, p. 26; 1870, p. 40.
 **Unio gubernaculum* REEVE, Conch. Icon., XVI, 1865, pl. XXVIII, fig. 146.—PÆTEL, Conch. Sam., III, 1890, p. 154.

†**TRUNCILLA PERPLEXA** var. **CINCINNATIENSIS** Lea.

- **Unio cincinnatiensis* LEA, Proc. Am. Phil. Soc., I, 1840, p. 285; *Tr. Am. Phil. Soc., VIII, 1842, p. 194, pl. VIII, fig. 4; *Obs., III, 1842, p. 32, pl. VIII, fig. 4.—*CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 246.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—*CHENU., Ill. Conch., 1858, pl. XXX, figs. 3, 3a, 3b.—*PÆTEL, Conch. Sam., III, 1890, p. 148.
 **Margaron (Unio) cincinnatiensis* LEA, Syn., 1852, p. 22; 1870, p. 33.
 **Unio phillipsii* REEVE, Conch. Icon., XVI, 1864, pl. IV, fig. 15.

Ohio River drainage; southern Michigan? The variety *rangianus* to south Michigan.

†**TRUNCILLA SAMPSONII** Lea.

- **Unio sampsonii* LEA, Pr. Ac. N. Sci. Phila., VI, 1861, p. 392; *Jl. Ac. N. Sci. Phila., V, 1862, p. 192, pl. XXV, fig. 261; *Obs., IX, 1863, p. 14, pl. XXV, fig. 261; Syn., 1870, p. 40.

Wabash River; Tennessee?

†**TRUNCILLA PROPINQUA** Lea.

- **Unio propinquus* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 83; *Jl. Ac. N. Sci. Phila., V, 1862, p. 63, pl. v, fig. 212; *Obs., VIII, 1862, p. 67, pl. v, fig. 212.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIX, fig. 417.—*PÆTEL, Conch. Sam., III, 1890, p. 164.
 **Margaron (Unio.) propinquus* LEA, Syn., 1870, p. 34.

Tennessee and Cumberland river drainage.

¹Dr. Lea claims that the animal of this form is lighter colored than that of *perplexus*, and quotes Mr. Clark, of Cincinnati, who states that this is known there as the White Mouth. The shells seem to absolutely blend together, and I have observed a wide range of color in the animals of a number of species of the genus.

† TRUNCILLA BIEMARGINATA Lea.

- * *Unio biemarginatus* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 83; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 47, pl. xvi, fig. 45; *Obs., XI, 1867, p. 51, pl. xvi, fig. 45.
 *Margarou (*Unio*) *biemarginatus* Lea, Syn., 1870, p. 38.

Tennessee River drainage.

† TRUNCILLA CAPSÆFORMIS Lea.

- * *Unio capsæformis* LEA, Tr. Am. Phil. Soc., VI, 1834, p. 31, pl. II, fig. 4; *Obs., I, p. 143, pl. II, fig. 4.—*CONRAD, Monég., VIII, 1837, p. 72, pl. XL, fig. 2.—*FERUS-SAC, Guer. Mag., 1839, p. 29.—*HANLEY, Test. Moll., 1842, p. 191, *Biv. Shells, 1843, p. 19, pl. XXI, fig. 5.—*CATLOW and REEVE, Conch. Nom., 1845, p. 57.—*KUSTER, Conch. Cab. Unio, 1852, p. 42, pl. VIII, fig. 5.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 246.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*REEVE, Conch. Icon., XVI, 1864, pl. XVII, figs. 79, 79a, 79b.—*PÆTEL, Conch. Sam., III, 190, p. 147.
 *Margarita (*Unio*) *capsæformis* LEA, Syn., 1836, p. 24; 1838, p. 19.
 *Margarou (*Unio*) *capsæformis* LEA, Syn., 1852, p. 27; 1870, p. 42.

Tennessee River drainage.

† TRUNCILLA FLORENTINA Lea.

- * *Unio florentinus* LEA, Pr. Ac. N. Sci. Phila., VII, 1857, p. 83; *Jl. Ac. N. Sci. Phila., V, 1862, p. 64, pl. v, fig. 213; *Obs., VIII, 1862, p. 68, pl. v, fig. 213.—*B. H. WRIGHT, Check List, 1888.
 *Margarou (*Unio*) *florentinus* LEA, Syn., 1870, p. 42.
 * *Unio turgidulus* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 40; *Jl. Ac. N. Sci. Phila., V, 1862, p. 62, pl. v, fig. 211; *Obs., VIII, 1862, p. 66, pl. v, fig. 211.—*REEVE, Conch. Icon., XVI, 1864, pl. XVII, fig. 80.—*PÆTEL, Conch. Sam., III, 1890, p. 170.
 *Margarou (*Unio*) *turgidulus* LEA, Syn., 1870, p. 42.
 * *Unio nux* KUSTER, Conch. Cab. Unio, 1861, p. 218, pl. LXXIII, fig. 2.
 * *Unio sacculus* REEVE, Conch. Icon., XVI, 1864, pl. xv, fig. 67.—*ANTHONY, Am. Jl. Conch., I, 1865, p. 157, pl. XII, fig. 3.
 * *Unio saccatus* KUSTER, Conch. Cab., 1861, p. 263, pl. LXXXIX, fig. 2.—*PÆTEL, Conch. Sam., III, 1890, p. 166.

Tennessee River drainage; Cumberland River.

† TRUNCILLA DEVIATA Reeve.

- * *Unio deviatu*¹ REEVE, Conch. Icon., XVI, 1864, pl. xv, fig. 61.—*ANTHONY, Am. Jl. Conch., I, 1865, p. 156, pl. XII, fig. 2.—*B. H. WRIGHT, Check List, 1888.—PÆTEL, Conch. Sam., III, 1890, p. 150.
 *Margarou (*Unio*) *deviatus* LEA, Syn., 1870, p. 42.

Cumberland and Tennessee rivers.

Genus MICROMYA (Agassiz, 1852) Simpson.

(Type, *Unio fabalis* Lea.)

Micromya AGASSIZ, Arch. für Nat., 1852, p. 57.

Shell triangular oval, solid, dark, feebly rayed with undulating lines; beak sculpture almost wanting, consisting of a few feeble, doubly-looped

¹Reeve credits this name to Anthony's manuscript in Museum Cuming.

ridges; hinge teeth heavy, laterals club-shaped and truncated posteriorly; post-basal swelling of the female distinct and often rather abrupt, sometimes somewhat irregularly radially ridged, the shell of this part being rather thin.

Animal with mantle fringed below, maculate on the border, which is greatly thickened at the post-basal part in the female and developed into a flap, with a distinct, toothed ridge inside; marsupium occupying the posterior part of the outer branchiæ in numerous distinctly marked ovisacs; inner gills free from abdominal sac in part.

(Group of *Micromya fabalis*.)

Shell small, with faint undulate rays, inflated area of female scarcely radiately striated.

† MICROMYA FABALIS Lea.

* *Unio fabalis*,¹ LEA, Tr. Am. Phil. Soc., IV, 1831, p. 86, pl. x, fig. 6; *Obs., I, 1834, p. 96, pl. x, fig. 16.—*FERUSSAC, Guer. Mag., 1835, p. 27.—*HANLEY, Test. Moll., 1842, p. 196; *Biv. Shells, 1843, p. 196.—*CATLOW and REEVE, Conch. Nom., 1845, p. 59; *H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*CHENU, Ill. Conch., 1858, pl. VIII, figs. 1, 1a, 1b.—*SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVI, fig. 196.

* *Margarita (Unio) fabalis* LEA, Syn., 1836, p. 28; 1838, p. 20.

* *Margaron (Unio) fabalis* LEA, Syn., 1852, p. 31; 1870, p. 49.

Unio capillus SAY, Transylvania Journ., IV, 1831, p. 528.

* *Unio lapillus* SAY, Am. Conch., V, 1832, pl. XLI; VI, 1834, No. 49.—*CONRAD, New, F. W. Shells, 1834, p. 70.; *Monog., VI, 1836, p. 54, pl. XXIX, fig. 12.—*CHENU, Bib. Conch., 1st ser. III, 1845, p. 52, pl. XIV, figs. 1, 1a, 1b.—*KUSTER, Conch. Cab., 1848, p. 53, pl. XI, fig. 3.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—*B. H. WRIGHT, Check List, 1888.—*PETEL, Conch. Sam., III, 1890, p. 156.

Ohio River drainage; Rouge River, near Detroit, Michigan; New York.

(Group of *Micromya cælata*.)

Shell triangular ovate, with a low, rounded posterior ridge; male and female shells distinctly wrinkled behind.

† MICROMYA CÆLATA Conrad.

* *Unio cælatus* CONRAD, Am. Jl. Sci. and Arts, XXV, 1834, p. 338, pl. I, fig. 1; New, F. W. Shells, 1834, p. 29, pl. III, fig. 4, p. 68.—*FERUSSAC, Guer. Mag., 1835, p. 29.—*MÖLLER, Syn. Nov. Gen., 1836, p. 199.—*HANLEY, Test. Moll., 1842, p. 175; *Biv. Shells, 1843, p. 175.—*CHENU, Bib. Conch., 1st ser. III, 1845, p. 16, pl. III, fig. 3.—*CATLOW and REEVE, Conch. Nom., 1845, p. 56.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—*REEVE, Conch.

¹ According to Scudder the volume of the Transactions containing this species appeared in the latter part of 1831, and was acknowledged by correspondents as received that year. Say published his *U. capillus* in the Transylvania Journal for December, 1831, and in the American Conchology changed it to *lapillus*. It is probable that Lea's name was published first.

Icon., XVI, 164, pl. II, fig. 7.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 147.

* *Margarita (Unio) calatus* LEA, Syn., 1836, p. 12; 1838, p. 14.

* *Margaron (Unio) calatus* LEA, Syn., 1852, p. 20; 1870, p. 20.

Tennessee River drainage; cited from Michigan by Sager and Miles, certainly in error. They might have mistaken for this a rather high specimen of *fabalis*.

Genus LAMPSILIS Rafinesque, 1820.

(Type, *Unio oratus* Say.)

Lampsilis RAFINESQUE, Ann. Gen. Sci. Phys. Brux., 1820, p. 298.

Shell oval to elliptical, smooth or slightly concentrically sculptured, usually without a posterior ridge; epidermis generally smooth and shining, often brilliantly rayed; beak sculpture, for the most part, consisting of fine, parallel ridges, which show a tendency to fall into an anterior and posterior loop; hinge with one or two pseudocardinals and one lateral in the right valve, and two pseudocardinals and two laterals in the left; female shell having a moderate and gradual inflation in the post-ventral region opposite the marsupium. Animal with the inner gills usually attached nearly or quite their entire length to the abdominal sac; marsupium occupying the hinder part of the outer gills; ovisacs distinct, separated by sulci, rounded below, having a fold near their bases, the whole projecting below the inner gills; mantle edge double and thickened, often swollen behind into a sort of flap in the female.

Section LAMPSILIS Rafinesque.

(Type, *Unio oratus* Say.)

Shell inflated, rather thin, shining, sometimes having a posterior ridge; beak sculpture coarse, consisting of a few, more or less, parallel ridges, which scarcely fall into loops. Animal having the mantle of the female usually toothed and thickened on the post-basal portion, which develops into a large, curious flap when the gills are filled with ova.

† LAMPSILIS VENTRICOSUS Barnes.

* *Unio ventricosus* BARNES, Am. Jl. Sci., VI, 1823, p. 267, pl. XIII, fig. 14 (outline).—

* SAY, Am. Conch., No. IV, 1832, pl. XXXII.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* C. B. ADAMS, Thompson's History of Vermont, 1842, p. 167; F. W. and L. S. of Vermont, 1842, p. 17.—* HANLEY, Test. Moll., 1842, p. 189; * Biv. Shells, 1843, p. 189, pl. XXIV, fig. 8.—* DE KAY, Zool. of N. Y., Pt. 6, 1843, p. 190.—* CHENU, Bib. Conch., 1st ser. III, 1845, p. 45, pl. XII, figs. 1, 2.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, pp. 492, 495.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLIII, fig. 235.—* B. H. WRIGHT, Check List, 1888.

* *Mya ventricosus* EATON, Zool. Text-Book, 1826, p. 221.

* *Margarita (Unio) ventricosus* LEA, Syn., 1836, p. 23; 1838, p. 18.

* *Margaron (Unio) ventricosus* LEA, Syn., 1852, p. 26; 1870, p. 41.

* *Lampsilis ventricosus* BAKER, Moll. Chicago, Pt. 1, 1898, p. 94, pl. XII, figs. 3-5.—

* SMITH, Bull. U. S. Fish Com., 1899, p. 291, pl. LXXXIII.

- * *Lampsilis ventricosa* STIMPSON, Shells of N. Eng., 1851, p. 14.
- * *Unio radiatus* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 286.
- Unio occidens* LEA, Tr. Am. Phil. Soc., III, 1829, p. 435, pl. x, fig. 16.—SHORT and EATON, Transylvania Jl., 1831, p. 78.—* LEA, Obs., I, 1834, p. 49, pl. x, fig. 16.—* HANLEY, Test. Moll., 1842, p. 189; * Biv. Shells, 1843, p. 189.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* KUSTER, Conch. Cab. Unio, 1856, p. 170, pl. LI, figs. 1, 2.—* CHENU, Ill. Conch., 1858, pl. XII, figs. 5, 5a, 5b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CALKINS, Pr. Ottawa Acad. Sci., 1874, p. 44.—* LATCHFORD, Tr. Ottawa Field. Nat. Club, 1882, p. 51.—* B. H. WRIGHT, Check List, 1888.
- * *Margarita (Unio) occidens* LEA, Syn., 1836, p. 23; 1838, p. 18.
- * *Margaron (Unio) occidens* LEA, Syn., 1852, p. 26; 1870, p. 41.¹
- * *Unio suboratus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 118, pl. XVIII, fig. 46;² * Obs., I, 1834, p. 128, pl. XVIII, fig. 46.—* HANLEY, Test. Moll., 1842, p. 184; * Biv. Shells, 1843, p. 184, pl. XXII, fig. 43.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* CHENU, Ill. Conch., 1858, pl. XII, figs. 6, 6a, 6b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXI, fig. 163; 1868, pl. LXXXV, fig. 456.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.
- * *Margarita (Unio) suboratus* LEA, Syn., 1836, p. 19; 1838, p. 17.
- * *Margaron (Unio) suboratus* LEA, Syn., 1852, p. 24; 1870, p. 37.
- * *Unio cardium* CONRAD, New F. W. Shells, 1834, p. 68.³—SAY, Am. Conch., VI, 1834.—CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 246.—KUSTER, Conch. Cab. Unio, 1856, p. 169, pl. L, figs. 1-4.
- * *Unio cardium* Raf. v. *occidens* PÆTEL, Conch. Sam., III, 1890, p. 174.
- * *Unio cardium* Raf. v. *ventricosus* PÆTEL, Conch. Sam., III, 1890, p. 147.
- * *Unio fasciolus*, FERUSSAC, Guer. Mag. 1835, p. 26.
- * *Unio ovata*, DESHAYES, An. sans. Vert., 3d ed. 1839, p. 669.
- * *Unio oratus* KUSTER, Conch. Cab. Unio, 1852, p. 55, pl. XII, fig. 1.
- * *Unio lenis* CONRAD, Monog., XII, 1840, p. 106, pl. LVIII, fig. 2.⁴
- * *Unio canadensis* LEA, Proc. Acad. N. Sci. Phila., I, 1857, p. 85;⁵ * Jl. Ac. Nat. Sci. Phila., IV, 1860, p. 268, pl. XLIV, fig. 148; * Obs., VII, 1860, p. 86, pl. XLIV, fig. 148.—B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) canadensis* LEA, Syn., 1870, p. 37.
- * *Unio dolabraformis* SOWERBY, Conch. Icon., XVI, 1867, pl. LIX, p. 298.
- * *Unio latissimus* SOWERBY?, Conch. Icon., XVI, 1868, pl. LXVI, fig. 337.

† **LAMPSILIS VENTRICOSUS** var. **SATUR** Lea.

- * *Unio satur* LEA, Pr. Am. Phil. Soc., V., 1852, p. 252;⁶ Tr. Am. Phil. Soc., X, 1852, p. 205, pl. XVII, fig. 19; * Obs., V, 1852, p. 21, pl. XVII, fig. 19.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857,

¹ Lea says this is so close to *U. ventricosus* that it would be well to unite them. Foot note in Synopsis, 1870, p. 41.

² This is, I am sure, a fine male *U. ventricosus* Barnes. Dr. Lea is inclined to this opinion in a footnote on p. 118 of the Transactions.

³ Conrad refers this species to Rafinesque. From the latter author's figure and description of *Lampsilis cardium* it is evident that it is a member of the *ventricosus* group, but whether it is the *U. ventricosus* of Barnes, the *U. capax* of Green, or some other species, I am unable to determine. Conrad and some other authors believe it to be Barnes's species.

⁴ The type in the Philadelphia Academy is a young *U. ventricosus*.

⁵ The type is a diseased individual, and is probably a compressed male *ventricosus*.

⁶ I believe this to be merely a rather delicate, dark-colored variety of *ventricosus*, which inhabits the southwestern part of the Mississippi drainage.

p. 496.—*CHENU, Man., 1859, II, p. 138, fig. 666.—*SOWERBY, Conch. Icon., XVI, 1868, pl. XCII, fig. 501.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 166.

**Margaron (Unio) satur* LEA, Syn., 1852, p. 24; 1870, p. 37.

Entire Mississippi drainage; St. Lawrence system; Nelson River and tributaries; the variety *satur* in the Southwest to the Sabine River, Texas.

† LAMPSILIS EXCAVATUS Lea.

**Unio excavatus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 32; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 71, pl. XIII, fig. 52; *Obs., VI, 1858, p. 71, pl. XIII, fig. 52.—*CHENU, Man., 1859, II, p. 138, fig. 665.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXX, VII, p. 403.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 152.

**Margaron (Unio) excavatus* LEA, Syn., 1870, p. 37.

*? *Unio ovatus* SAY var. *ornatus* CONRAD, Monog., I, 1835, p. 4.¹

**Unio ornatus* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXI, fig. 162.—*B. H. WRIGHT, Check List, 1888.

Tombigbee and Alabama river drainage.

† LAMPSILIS BINOMINATUS Simpson.²

**Unio lineatus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 287; *Tr. Am. Phil. Soc., VIII, 1842, p. 206, pl. XII, fig. 20; *Obs., III, 1842, p. 44, pl. XII, fig. 20.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—*CHENU, Ill. Conch., 1858, pl. XXVIII, figs. 1, 1a, 1b.—*KUSTER, Conch. Cab. Unio, 1862, p. 278, pl. XCIV, fig. 1.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXI, fig. 309.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 157.

**Margaron (Unio) lineatus* LEA, Syn., 1852, p. 27; 1870, p. 41.

Chattahoochee and Flint rivers, Georgia.

† LAMPSILIS CARIOSUS Say.

Unio cariosus SAY, Nich. Encyc., II, 1816, pl. III, fig. 2.³—SHORT and EATON Transylvania Jl. 1831, p. 70.—*SAY, Am. Conch., VI, 1834.—*CONRAD, New F. W. Shells, 1834, p. 68.—*FERUSSAC, Guer. Mag., 1835, p. 26.—*CONRAD, Monog. IV, 1836, p. 40, pl. XIX.—*GOULD, Inv. of Mass., 1841, p. 111, fig. 72.—*BINNEY, Inv. of Mass., 1870, p. 172, fig. 475.—*REEVE, Conch. Syst., I, 1841, p. 119, pl. LXXXIX, fig. 10.—*HANLEY, Test. Moll., 1842, p. 190.—*DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 193, pl. XXI, figs. 243, 244.⁴—*HANLEY, Biv. Shells, 1843, p. 190, pl. XX, fig. 22.—*CATLOW and REEVE, Conch. Nom., 1845, p. 57.—*KUSTER, Conch. Cab. Unio, 1852, p. 24, pl. I, figs. 2, 3.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1855, p. 246.—*DEWEY, Ninth Rep. N. Y. State Cab., 1856, p. 35.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*SOWERBY,

¹ It is hard to tell just what this is, as Conrad does not figure or describe it. He states that while it is probably *excavatus* it is not certainly so, and he does not think it wise to displace Lea's properly described and figured species for it.

² Lea's name was preoccupied by Valenciennes (Encyc. Meth., II, 1827, p. 151, pl. CCXLVIII, fig. 5) for what is probably *U. radiatus*. I change it to *binominatus*.

³ The first edition of Nicholson's Encyclopedia is dated 1816, fide W. G. Binney, Bib. N. Am. Conch., I, 1863, p. 25, and by Lea, Synopsis, 4th ed., pp. 154, 159.

⁴ The figures are very probably taken from *U. ventricosus*.

Conch. Icon., XVI, 1867, pl. LVIII, fig. 294.—* HARTMAN and MICHENER, Conch. Cest., 1874, p. 38, fig. 183.—* LATCHFORD, Tr. Ottawa Field Nat. Club, 1882, p. 51.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 147.—* SIMPSON, Nautilus VIII, 1895, p. 122, 2 figures.¹

* *Margarita (Unio) cariosus* LEA, Syn., 1836, p. 23; 1838, p. 18.

* *Margaron (Unio) cariosus* LEA, Syn., 1852, p. 27; 1870, p. 42.

* *Unio cariosa* LAMARCK, An. sans Vert., VI, 1819, p. 80.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 545; 3d ed., II, 1839, p. 671.

* *Mya cariosa* EATON, Zool. Text Book, 1826, p. 218.

* *Lampsilis cariosa* STIMPSON, Shells of N. Eng., 1851, p. 14.

* *Unio oratus* var. b. LAMARCK, An. sans Vert., VI, 1819, p. 75.

* *Unio ovata* VALENCIENNES, Rec. Obs. Zool. Anat., II, 1833, p. 226, pl. L, figs. 1, 1a, 1b, 1c.

* *Unio viridis* FERUSSAC, Guer. Mag., 1835, p. 27.

* *Unio oratus* CONRAD, Ann. and Mag. Nat. Hist., IV, 1849, p. 301; * Pr. Ac. N. Sci. Phila., IV, 1849, p. 153; * Jl. Ac. N. Sci. Phila., 1850, p. 276, pl. XXXVII; fig. 6; Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.

Atlantic drainage from Georgia to the lower St. Lawrence.²

† LAMPSILIS ALTILIS Conrad.

* *Unio altilis* CONRAD, New F. W. Shells, 1834, p. 43, pl. II, fig. 1, and p. 68.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen., 1836, p. 207.—* HANLEY, Test. Moll., 1842, p. 190; Biv. Shells, 1843, p. 190.—* CATLOW and REEVE, Conch. Nom., 1845, p. 55.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 21, pl. I, fig. 1.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 244.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.

* *Margarita (Unio) altilis* LEA, Syn., 1836, p. 24; 1838, p. 19.

* *Margaron (Unio) altilis* LEA., Syn., 1852, p. 27; 1870, p. 42.

Alabama River drainage; Little Red River, Clinton, Arkansas?

† LAMPSILIS DOLABRÆFORMIS Lea.

* *Unio dolabræformis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 103, pl. XXIV, fig. 113; * Obs., II, 1838, p. 103, pl. XXIV, fig. 113.—* TROSCHIEL, Arch. für Naturg., V, 1839, II, p. 237.—* HANLEY, Test. Moll., 1842, p. 189; * Biv. Shells., 1843, p. 189, pl. XXI, fig. 47.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* CHENU, Ill. Conch., 1858, pl. XXIII, figs. 6, 6a, 6b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Conch. Cab., 1861, p. 170, pl. LI, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

* *Margarita (Unio) dolabræformis* LEA, Syn., 1838, p. 18.

* *Margaron (Unio) dolabræformis* LEA, Syn., 1852, p. 27; 1870, p. 41.

Savannah, Ogeechee, and Altamaha rivers, Georgia.

† LAMPSILIS CAPAX Green.

* *Unio capax* GREEN, Cab. Nat. Hist., II, 1832, p. 290.—* CONRAD, New F. W. Shells, 1834, p. 68.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* CONRAD, Monog., IX, 1837, p. 75, pl. XLII.—* KUSTER, Conch. Cab. Unio, 1852, pp. 21, 65, pl. XV, fig. 3.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A.

¹ By an error the figures of this species and *U. ochraceus* were changed.

² The specimens referred to this species that are often reported from the Mississippi drainage are no doubt *L. ventricosus*.

- ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1866, pl. LII, fig. 274.—* CALKINS, Pr. Ottawa Ac. Nat. Sci., 1874, p. 41.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 147.
- * *Margaron (Unio) capax* LEA, Syn., 1852, p. 27; 1870, p. 42.
- * *Lampsilis capax* SMITH, Bull. U. S. Fish Com., 1899, p. 291, pl. LXXXIV.
- * *Symphynota globosa* LEA, Tr. Am. Phil. Soc., V, 1832, p. 41, pl. IV, fig. 12; * Obs., I, 1834, p. 153, pl. IV, fig. 12.—* FERUSSAC, Guer. Mag., 1835, p. 25.
- * *Margarita (Unio) globosa* LEA, Syn., 1836, p. 23.
- * *Unio globosus* HANLEY, Test. Moll., 1842, p. 188; * Biv. Shells, 1843, p. 189.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.
- * *Margarita (Unio) globosus* LEA, Syn., 1838, p. 18.

Lower Ohio River drainage; southwest to St. Francis River, Arkansas; north to eastern Iowa; Elkhorn and Blue rivers, Nebraska? (Aughey). Abundant locally.

† LAMPSILIS OVATUS Say.¹

- Unio ovatus* SAY, Nich. Encyc., II, 1816, pl. II, fig. 7.—* LAMARCK, An. sans Vert., VI, 1819, p. 75.—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 287.—SHORT and EATON Trans. Jl., 1831, p. 78.—* SAY, Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 70; * Monog. I, 1835, p. 4, pl. II.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* HANLEY, Test. Moll., 1842, p. 184; * Biv. Shells, 1843, p. 184.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* KUSTER, Conch. Cab. Unio, 1852, p. 22, pl. III, fig. 2.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXI, fig. 164.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 162.
- * *Margarita (Unio) ovatus* LEA, Syn., 1836, p. 19; 1838, p. 17.
- * *Margaron (Unio) ovatus*, LEA, Syn., 1852, p. 24; 1870, p. 37.
- * *Unio ovata* LAMARCK, An. sans Vert., VI, 1819, p. 75.—* STARK, Nat. Hist., II, 1828, p. 90.—* VALENCIENNES, Coq. Marines, Biv., I, 1833, pl. 1, figs. 1, 1a, 1b, 1c.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 538; 3d ed., II, 1839, p. 669.
- * *Lampsilis ovata* RAFINESQUE, Ann. Gen. Sci. Phys. Brux., V, 1820, p. 298.
- * *Mya ovata* EATON, Zool. Text Book, 1826, p. 218.
- * *Eglia ovata* SWAINSON, Treat. on, Mal., 1810, p. 266, fig. 49.

Ohio River drainage.

† LAMPSILIS OCHRACEUS Say.

- Unio ochraceus* SAY, Nich. Encyc., 1816, pl. II, fig. 8.—* CONRAD, New F. W. Shells, 1834, p. 70.—* FERUSSAC, Guer. Mag., 1835, p. 25.—* CONRAD, Monog., IV, 1836, p. 37, pl. XVIII, fig. 2.—* GOULD, Inv. Mass., 1841, p. 112, fig. 74; 1870, p. 173, fig. 476.—* HANLEY, Test. Moll., 1842, p. 190; * Biv. Shells, 1843, p. 190, pl. XX, fig. 48?—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 193, pl. LXIX, figs. 237, 238.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* KUSTER, Conch. Cab. Unio, 1856, p. 163, pl. XLVII, fig. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIII, fig. 317.—* HARTMAN and MICHENER, Conch. Cest., 1874, p. 39, fig. 184.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 161.—* SIMPSON, Nautilus, VIII, 1895, p. 122, fig.²
- * *Margaron (Unio) ochraceus* LEA, Syn., 1852, p. 27; 1870, p. 42.
- * *Mya ochracea* EATON, Zool. Text Book, 1826, p. 218.

¹ The name *ovatus* was used for *Unio tumidus* Retz., by Lister, Donovan, and others, but as it was called a *Mya* and the generic name *Unio* was never used with it, Say's name can stand for this species.

² The figure given for this species, by an error of the printer, is a female *cariosus*.

**Symphynota ochracea* LEA, Tr. Am. Phil. Soc., III, 1830, p. 455; *Obs. I, 1834, p. 69.

**Margarita (Unio) ochracea* LEA, Syn., 1836, p. 23; 1838, p. 18.

**Lampsilis ochracea* STIMPSON, Shells of N. Eng., 1851, p. 14.

**Unio crocatus* LEA, Pr. Am. Phil. Soc., II, 1841, p. 31; *Tr. Am. Phil. Soc., VIII, 1842, p. 238, pl. xxii, fig. 52; *Obs., III, 1842, p. 76, pl. xxii, fig. 52.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—*CHENU, Ill. Conch., 1858, pl. xxxiii, figs. 1, 1a, 1b.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 149.

**Margaron (Unio) crocatus* LEA, Syn., 1852, p. 27; 1870, p. 42.

**Unio rosaceus* CONRAD, Pr. Acad. N. Sci., Phila., IV, 1849, p. 153; *Ann. and Mag. Nat. Hist., IV, 1849, p. 303; *Jl. Acad. N. Sci., Phila., I, 1850, p. 275, pl. xxxvii, fig. 5.

**Unio troostensis* SOWERBY, Conch. Icon., XVI, 1866, pl. xxxviii, fig. 210.

**Unio affinis* SOWERBY? Conch. Icon., XVI, 1868, pl. lxiii, fig. 318.

Atlantic drainage, from New England to the Ogeechee River, Georgia.

† LAMPSILIS SPLENDIDUS Lea.

**Unio splendidus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 70, pl. xix, fig. 61; *Obs., II, 1838, p. 70, pl. xix, fig. 61.—*TROSCHER, Arch. für Naturg., V, 1839, Pt. 2, p. 237.—*HANLEY, Test. Moll., 1842, p. 190; *Biv. Shells, 1843, p. 190.—*CATLOW and REEVE, Conch. Nom., 1845, p. 64.—*KUSTER, Conch. Cab. Unio, 1852, p. 55, pl. xii, fig. 2.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU, Ill. Conch., 1858, pl. xvii, figs. 7, 7a, 7b.—*SOWERBY, Conch. Icon., XVI, 1866, pl. xxxi, fig. 161.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 168.

**Margarita (Unio) splendidus* LEA, Syn., 1836, p. 23; 1838, p. 18.

**Margaron (Unio) splendidus* LEA, Syn., 1852, p. 27; 1870, p. 42.

**Unio regularis* SOWERBY, Conch. Icon., XVI, 1868, pl. xxxiv, fig. 181.

Altamaha and Ogeechee rivers, Georgia.

LAMPSILIS DELUMBIS Conrad.¹

**Unio delumbis* CONRAD, New F. W. Shells, 1834, p. 35, pl. v, fig. 3; p. 69.—*FERUSSAC, Guerin Mag., 1835, p. 29.—*MÖLLER, Syn. Nov. Gen., 1836, p. 204.—*CHENU, Bib. Conch., 1st ser., III, 1845, p. 18, pl. ii, fig. 10.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1857, p. 248.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 150.

Small streams near Cooper River, South Carolina.

† LAMPSILIS PEROVALIS Conrad.

**Unio perovalis* CONRAD, New F. W. Shells, 1834, p. 43, pl. ii, fig. 2, p. 71.—*FERUSSAC, Guerin Mag., 1835, p. 29.—*MÖLLER, Syn. Nov. Gen., 1836, p. 207.—*HANLEY, Test. Moll., 1842, p. 191; *Biv. Shells, 1843, p. 191.—*CATLOW and REEVE, Conch. Nom., 1845, p. 62.—*CHENU, Bib. Conch., 1st ser., III, 1845, p. 21, pl. i, fig. 2.—*CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 254.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*KUSTER, Conch. Cab. Unio, 1861, p. 257, pl. lxxxvii, fig. 2.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 163.

**Margarita (Unio) perovalis* LEA, Syn., 1836, p. 24; 1838, p. 19.

**Margaron (Unio) perovalis* LEA, Syn., 1852, p. 27; 1870, p. 42.

Alabama and Black Warrior rivers.

¹This may be a valid species, but it looks from the figure like an elongated *ochraceus*.

† LAMPASILIS PERPASTUS Lea.

- * *Unio perpastus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 60; *Jl. Ac. N. Sci. Phila., V, 1862, p. 69, pl. VII, fig. 219; *Obs., VIII, 1862, p. 73, pl. VII, fig. 219.—
* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) perpastus* LEA, Syn., 1870, p. 43.

Swamp Creek, Georgia; Coosa River, Alabama.

† LAMPASILIS CLARKIANUS Lea.

- * *Unio clarkianus* LEA, Pr. Am. Phil. Soc., V, 1852, p. 251; *Tr. Am. Phil. Soc., X, 1852, p. 273, pl. XXI, fig. 30; *Obs., V, 1852, p. 29, pl. XXI, fig. 30.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 246.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PATEL, Conch. Sam., III, 1890, p. 148.

* *Margaron (Unio) clarkianus* LEA, Syn., 1852, p. 27; 1870, p. 42.

- *† *Unio spillmanii* LEA, Pr. Ac. N. Sci. Phila., XIII, 1861, p. 39; *Jl. Acad. N. Sci. Phila., V, 1862, p. 98, pl. XV, fig. 246; *Obs., VIII, 1862, p. 102, pl. XV, fig. 246.—
* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) spillmanii* LEA, Syn., 1870, p. 42.

Williamsport, Tennessee; near Columbus, Mississippi; Cahawba and Black Warrior rivers, Alabama.

† LAMPASILIS GERHARDTII Lea.

- * *Unio gerhardtii* LEA, Pr. Acad. N. Sci. Phila., VI, 1862, p. 168; *Jl. Ac. N. Sci. Phila., V, 1862, p. 208, pl. XXXI, fig. 277; *Obs., IX, 1863, p. 30, pl. XXXI, fig. 277.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) gerhardtii* LEA, Syn., 1870, p. 35.

Chattanooga (river?), Georgia; Shorter, Alabama.

† LAMPASILIS MULTIRADIATUS Lea.

- * *Unio multiradiatus* LEA, Tr. Am. Phil. Soc., III, 1829, p. 434, pl. IX, fig. 15; *Obs. I, 1834, p. 48, pl. IX, fig. 15.—* CONRAD, N. F. W. Shells, 1834, p. 70.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 549; 3d ed., II, 1839, p. 673.—* HANLEY, Test. Moll., 1842, p. 190; *Biv. Shells, 1843, p. 190, pl. XXI, fig. 10.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. XII, figs. 2, 2a, 2b.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXX, fig. 506a.—* B. H. WRIGHT, Check List, 1888.—
* PATEL, Conch. Sam., III, 1890, p. 160.

* *Margarita (Unio) multiradiatus* LEA, Syn., 1836, p. 24; 1838, p. 19.

* *Margaron (Unio) multiradiatus* LEA, Syn., 1852, p. 31; 1870, p. 42.

* *Unio fasciolus* SAY, Am. Conch., VI, 1834.—* CONRAD, Monography, III, 1836, p. 26, pl. XI, fig. 2.—* KUSTER, Conch. Cab. Unio, 1852, p. 32, pl. v, fig. 4.

*† *Unio perradiatus* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 40; *Jl. Ac. N. Sci. Phila., V, 1862, p. 66, pl. VI, fig. 215; Obs., VIII, 1862, p. 70, pl. VI, fig. 215.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) perradiatus* LEA, Syn., 1870, p. 37.

* *Unio alt'lis*, REEVE, Conch. Icon., XVI, 1865, pl. XXIII, fig. 109.

* *Unio pervalis* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVIII, fig. 209.

Entire Ohio River drainage; Southern Michigan; New York? (Marshall).

† LAMPSILIS DOLIARIS Lea.

Unio doliaris LEA, Pr. Ac. N. Sci. Phila., XVII, 1865, p. 88; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 260, pl. XXXII, fig. 75; *Obs., XII, 1869, p. 20, pl. XXXII, fig. 75.—

*B. H. WRIGHT, Check List, 1888.

*Margaron (*Unio*) *doliaris* LEA, Syn., 1870, p. 42.

Alabama and Tombigbee drainage.

† LAMPSILIS BREVICULUS Call.

**Unio breviculus* CALL, Pr. U. S. Nat. Mus., X, 1887, p. 499, pl. XXVIII; *Tr. Ac. Sci. St. Louis, VII, 1895, p. 6, pl. XVII.

† LAMPSILIS BREVICULUS var. BRITTSI Simpson.

**Lampsilis brittsi* Simpson, Pr. Ac. N. Sci. Phila., 1900, p. 76, pl. v, figs. 1, 2.¹

White and Current rivers, Arkansas; Texas County, Missouri.

† LAMPSILIS BIANGULATUS Lea.²

**Unio biangulatus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 288; *Tr. Am. Phil. Soc., VIII, 1843, p. 197, pl. IX, fig. 8; *Obs., III, 1842, p. 35, pl. IX, fig. 8.—*CONRAD, Pr. Ac. N. Sci. Phila., 1853, p. 245.—*CHENU, Ill. Conch., 1858, pl. XXX, figs. 7, 7a, 7b.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—*KUSTER, Conch. Cab. Unio, 1861, p. 189, pl. LX, fig. 1; pl. LXI, fig. 1.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXX, fig. 421.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 146.

*Margaron (*Unio*) *biangulatus* LEA Syn., 1852, p. 38, 1870, p. 61.

Tennessee drainage.

LAMPSILIS SUBVEXA Conrad.³

**Anodonta subvexa* CONRAD, Am. Jl. Sci., XXV, 1834, p. 341, pl. I, fig. 1; *New F. W. Shells, 1834, p. 73.—*FERUSSAC, Guer. Mag. 1835, p. 25.—*MÖLLER, Syn. Nov. Gen., 1836, p. 194.—*CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 264.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 185.

¹At the time I published the *L. brittsi* I believed it to be a valid species. Since then I have seen additional material which seems to be a connecting link between it and *L. breviculus*. I can not be certain as to the relationships of this form, which seems on the one hand to have characters belonging to the typical section of *Lampsilis* and on the other to be related to *L. spatulatus* and *L. pleasi*.

²The systematic position of this species is a little uncertain. In the form of the shell and the teeth it seems to belong in the *Ventricosus* group, but the texture and broken rays apparently ally it to the *Ligamentinus* group. The soft parts agree best with those of members of the former assemblage, so I place it here.

³I do not know what this is, nor where it belongs, but it is certainly not a true *Anodonta*. The shell, according to Conrad's figure, is rayed throughout, and looks very much like a young *U. ventricosus*. He says that it has a callus resembling an incipient tooth; that it inhabits the Black Warrior River, and that it is very rare. It is just possible that it is related to some of the so-called Margaritanas of the Southern States, such as *M. spillmani*, but I think it more likely some young form of the *Ventricosus* group in which the hinge has been injured. I place it here with the greatest hesitation.

**Margarita (Anodonta) subrexa* LEA, Syn., 1836, p. 52; 1838, p. 31.

**Anodon subrexa* CATLOW and REEVE, Conch. Nom., 1845, p. 68.

**Margaron (Anodonta) subrexa* LEA, Syn., 1852, p. 50; 1870, p. 81.

Black Warrior River, Alabama.

Section EURYNIA Rafinesque, 1820.

(Type, *Unio recta* Lamarck.)

Shell oval to oblong; beak sculpture delicate, doubly looped; mantle double edged and often toothed below; that of the female sometimes developed into a thickened flap at the post base.¹

(Group of *Lampsilis luteolus*.)

Shell shining, generally rayed; beak sculpture often broken, with the posterior loop open behind; pseudocardinals rather small, compressed, smooth inside, and often curved slightly upward.

† LAMPSILIS LUTEOLUS Lamarck.

**Unio luteola* LAMARCK, An. sans Vert., VI, 1819, p. 79.—*DESHAYES, An. sans Vert., 2d ed. VI, 1835, p. 544; 3d ed. II, 1839, p. 671.

**Unio luteolus* HANLEY, Test. Moll., 1842, p. 192.—*DE KAY, Zool. of New York, Pt. 5, 1843, p. 190, pl. xx, fig. 241.—*HANLEY, Biv. Shells, 1843, p. 192, pl. xxiii, fig. 16.—*CATLOW and REEVE, Conch. Nom., 1845, p. 60.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*SOWERBY, Conch. Icon., XVI, 1867, pl. lviii, fig. 293, 293a, 293b.—*LATCHFORD, Tr. Ottawa F. Nat. Club, 1882, p. 51.—*B. H. WRIGHT, Check List, 1888.—*PETEL, Conch. Sam., III, 1890, p. 158.—*CALL, Tr. Acad. Sci. St. Louis, VII, 1895, p. 25, pl. iii.

**Margarita (Unio) luteolus* LEA, Syn., 1836, p. 25; 1838, p. 19.

**Margaron (Unio) luteolus* LEA, Syn., 1852, p. 28; 1870, p. 44.

**Lampsilis luteolus* BAKER, Moll. Chicago, Pt. 1, 1898, p. 103, pls. xi, xxxvii, fig. 12.

**Unio inflatus* BARNES, Am. Jl. Sci., VI, 1823, p. 266.—*CONRAD, New F.W. Shells, 1834, p. 69.

**Mya inflata* EATON, Zool. Text-Book, 1826, p. 221.

**Unio siliquoides* BARNES, Am. Jl. Sci., VI, 1823, p. 269, pl. xiii, fig. 150 (outline).—*FERUSSAC, Guer. Mag., 1835, p. 27.—*CONRAD, Monog., II, 1836, p. 22, pl. x, fig. 1.—*C. B. ADAMS, Thompson's Hist. of Vermont, 1842, p. 167; F. W. and L. S. of Vermont, p. 17, 1842.

**Unio siliquoides* KUSTER, Conch. Cab. Unio, 1852, p. 30, pl. v, fig. 2.

**Mya siliquoides*, EATON, Zool. Text-Book, 1826, p. 221.

**Lampsilis siliquoides* STIMPSON, Shells of New Eng., 1851, p. 14.

**Unio childreni* HANLEY, Biv. Shells, 1843, p. 193, pl. xxiii, fig. 57.

**Unio hyadianus* KUSTER, part, Conch. Cab. Unio, 1861, p. 201, pl. lxvii, fig. 1.²

**Unio distans* ANTHONY, Am. Jl. Conch., I, 1865, p. 156, pl. xiii, fig. 2.—*B. H. WRIGHT, Check List, 1888.—*PETEL, Conch. Sam., III, 1890, p. 151.

**Unio multiradiatus* SOWERBY, Conch. Icon., XVI, 1868, pl. lxi, fig. 306.³

**Unio affinis* SOWERBY, Conch. Icon., XVI, 1868, pl. lxi, fig. 307.

¹ Lea has figured a female animal of *L. radiatus* in Observations II, pl. xv, which has a curious, winglike projection on the mantle at its posterior base.

² There are two figures; the first is *luteolus*, the second is *hyadianus*.

³ Changed in errata to *luteolus*. The figure 306a, pl. lxx, is a genuine *multiradiatus*.

† LAMPSILIS LUTEOLUS var. ROSACEUS De Kay.

- * *Unio rosaceus* DE KAY, Zool. of New York, V, 1843, p. 192, pl. XXXIX, figs. 355, 356; pl. XL, fig. 357.¹—CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 257.—B. H. WRIGHT, Check List, 1888.

¹ Entire Mississippi drainage, southwest to the Brazos River, Texas; St. Lawrence drainage; entire Dominion of Canada east of the Rocky Mountains; the variety *rosaceus* in the St. Lawrence area.

† LAMPSILIS SUPERIORENSIS Marsh.²

- * *Unio superiorenensis* MARSH, Naut., X, 1897, p. 103, pl. I, figs. 1, 2, 5.

Upper Great Lakes region.

† LAMPSILIS BOREALIS A. F. Gray.³

- * *Unio borealis* A. F. GRAY, Tr. Ottawa Field Nat. Club, 1882, p. 53, pl., 3 figures.—* B. H. WRIGHT, Check List, 1888.

Lower St. Lawrence drainage, to Lake Michigan; Lake of the Woods.

† LAMPSILIS RADIATUS Gmelin.

- * *Mya radiata* GMELIN, Syst. Nat., 13th ed., 1792, p. 3220.—* WOOD, Gen. Conch., I, 1815, p. 109.—* DILLWYN, Cat. I, 1817, p. 51.—* WOOD, Index Test. 2d ed., 1825, p. 12, pl. II, fig. 2.—* EATON, Zool. Text-Book, 1826, p. 220.—* HANLEY, Index Test., 3d ed., 1856, p. 16, pl. II, fig. 32.

- * *Unio radiata* LAMARCK, An. sans. Vert., VI, 1819, p. 73.—* DESHAYES, Hist. Nat. des Vers., II, 1830, p. 581; * An. sans. Vert., 2d ed., VI, 1835, p. 535; 3d ed., 1839, p. 668.—* HANLEY, Test. Moll., 1842, p. 212.

- * *Lampsilis radiata* STIMPSON, N. Eng. Shells, 1851, p. 13.

- * *Unio radiatus* SPENGLER, Skriv. Nat. Selsk., V, 1792, p. 3; III, 1793, p. 62.—* CONRAD, New F. W. Shells, 1834, p. 71.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* CONRAD, Monog., II, 1836, p. 24, pl. X, fig. 2.—* SOWERBY, Rich. Fann. Boreali Am., III, 1836, p. 316.—* GOULD, Inv. of Mass., 1841, p. 110, fig. 73; 1870, p. 170, fig. 474.—* HANLEY, Test. Moll., 1842, p. 193; * Biv. Shells, 1843, p. 193.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 256.—* KUSTER, Conch. Cab. Unio., 1852, p. 29, pl. V, fig. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* REEVE, Conch. Icon., XVI, 1865, p. 27, pl. XXVII, fig. 139.—* HARTMAN and MICHE-
NER, Conch. Cest., 1874, p. 87, fig. 182.—* LATCHFORD, Tr. Ottawa F. Nat. Club, 1882, p. 50.—* B. H. WRIGHT, Check List, 1888.—* H. CARPENTER, Naut., IV, 1890, p. 22.—* PÆTEL, Conch. Sam., III, 1890, p. 165.

- * *Margarita (Unio) radiatus* LEA, Syn., 1836, p. 25; 1838, p. 19.

- * *Margaron (Unio) radiatus* LEA, Syn., 1852, p. 29; 1870, p. 44.

- * *Unio pictorum tenuis* India, etc.—CHEMNITZ, Conch. Cab., 1788, p. 23, pl. II, fig. 7.

¹ The type of De Kay's variety is a delicate shell with peculiarly pallid brown epidermis, slightly rayed, and having a rose-colored nacre. This form is often destitute of the fine tinting of the nacre, and sometimes becomes quite dark, even bronzy or brownish black. It is common in the lakes of the St. Lawrence drainage.

² This and *L. borealis* A. F. Gray are doubtful species which seem to stand between *L. luteolus* and *radiatus*. This form is nearer *luteolus*, while Gray's species approaches more nearly to *radiatus*.

³ In Latchford's Notes on Ottawa Unionidæ.

- * *Unio virginiana* LAMARCK, An. sans. Vert., VI, 1819, p. 79.—* DELESSERT, Rec. Coq. Lam., 1841, pl. XI, fig.
- * *Unio lineata* VALENCIENNES, Enc. Meth., II, 1827, p. 151, pl. CCXLVIII, fig. 5.
- * *Unio elongata* GOODRICH, Ill. Nat. Hist., II, 1829, p. 523, fig.
- * *Unio tenebrosus* CONRAD, New F. W. Shells, 1834, p. 42, pl. VII, fig. 1; p. 72.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen. 1836, p. 207.—* CHENU, Bib. Conch., 1st ser., III, 1845, pl. II, fig. 5.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 169.
- * *Unio melinus* CONRAD, Monog. XI, 1838, p. 101, pl. LV, fig. 1.—SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVII, fig. 467.
- * *Unio mellinus* PÆTEL, Conch. Sam., III, 1890, p. 159.
- * *Unio boydianus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 286; * Tr. Am. Phil. Soc., VIII, 1843, p. 216, pl. XVI, fig. 32; * Obs., III, 1842, p. 54, pl. XVI, fig. 22.—* DE KAY, Zool. of New York, Pt. 5, 1843, p. 189.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* CHENU, Ill. Conch., 1858, pl. XXXII, figs. 2, 2a, 2b.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 146.
- * *Margaron (Unio) boydianus* LEA, Syn., 1852, p. 38; 1870, p. 62.
- * *Unio crassus* KUSTER, Conch. Cab. Unio., 1854, p. 113, pl. XXXI, fig. 5.
- * *Mya oblongata*, WOOD, Index Test., 1856, p. 199, pl. I; Supplement, fig. 2.
- * *Unio obliquiradiatus* REEVE, Conch. Icon., XVI, 1865, pl. XXIX, fig. 151.—* PÆTEL, Conch. Sam., III, 1890, p. 161.

† LAMPSILIS RADIATUS var. CONSPICUUS Lea.

- * *Unio conspicuus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 156; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 34, pl. XI, fig. 31; * Obs., XIII, 1874, p. 38, pl. XI, fig. 31.—* B. H. WRIGHT, Check List, 1888.

St. Lawrence drainage; Manitoba; Atlantic region south to North Carolina (var. *conspicuus* in North Carolina and southern Virginia).

Gould¹ reports *L. radiatus* from the north shore of Lake Superior, and there is a shell in the U. S. National Museum from the northwest boundary of Wisconsin which is probably this species.

† LAMPSILIS HYDIANUS Lea.

- * *Unio hydianus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 14, pl. VI, fig. 14; * Obs., II, 1838, p. 14, pl. VI, fig. 14.—* TROSCHEL, Arch. für Naturg., VII, 1839, p. 234.—* HANLEY, Test. Moll., 1842, p. 192; Biv. Shells, 1843, p. 192, pl. XXIII, fig. 6.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* KUSTER, Conch. Cab. Unio., 1861, p. 201, pl. LXVII, fig. 2.—* CHENU, Ill. Conch., 1858, pl. XVII, fig. 3, 3a, 3b; pl. XXIV, figs. 7, 7a, 7b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVII, fig. 203.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
- * *Margarita (Unio) hydianus* LEA, Syn., 1836, p. 25; 1838, p. 19.
- * *Margaron (Unio) hydianus* LEA, Syn., 1852, p. 28; 1870, p. 44.
- * *Unio placitus* LEA, Pr. Am. Phil. Soc., V, 1852, p. 252; * Tr. Am. Phil. Soc., 1852, p. 279, pl. XXIII, fig. 38; * Obs., V, 1852, p. 35, pl. XXIII, fig. 38.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1838, p. 255.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Conch. Cab. Unio., 1861, p. 262, pl. LXXXVIII, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., 1890, p. 163.
- * *Margaron (Unio) placitus* LEA, Syn., 1852, p. 29; 1870, p. 46.

Eastern half of Texas; Indian Territory; Arkansas; east to Alabama.

¹ Agassiz, Lake Superior, p. 245.

² Merely a young *hydianus*.

† *LAMPSILIS APPROXIMUS* Lea.¹

* *Unio approximus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; * Tr. Am. Phil. Soc., X, 1848, p. 74, pl. v, fig. 13; * Obs., IV, 1848, p. 48, pl. v, fig. 13.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.

* *Margaron (Unio) approximus* LEA, Syn., 1852, p. 28; 1870, p. 44.

Southern Louisiana; southern Alabama.

† *LAMPSILIS AFFINIS* Lea.²

* *Unio affinis* LEA, Pr. Am. Phil. Soc., V, 1852, p. 251; * Tr. Am. Phil. Soc., 1855, p. 271, pl. XIX, fig. 26; * Obs., V, 1852, p. 27, pl. XIX, fig. 26.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 244.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.

* *Margaron (Unio) affinis* LEA, Syn., 1852, p. 27; 1870, p. 42.

Louisiana.

LAMPSILIS CONTRARIUS Conrad.³

* *Unio contrarius* CONRAD, Jl. Ac. Nat. Sci. Phila. I, 1850, p. 276, pl. XXXVII, fig. 7. Ogeechee River, Georgia. (Locality doubtful.)

† *LAMPSILIS CLAIBORNENSIS* Lea.

Unio claubornensis LEA, Tr. Am. Phil. Soc., VI, 1838, p. 105, pl. XXIV, fig. 115; * Obs., II, 1838, p. 105, pl. XXIV, fig. 115.—* TROSCHER, Arch. für. Naturg., V, 1839, pl. II, p. 237.—* HANLEY, Test. Moll., 1842, p. 192, Biv. Shells, 1843, p. 192, pl. XXI, fig. 26.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 246.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* CHENU, Ill. Conch., 1858, pl. XXIII, figs. 3, 3a, 3b.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXX, fig. 357.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Margarita (Unio) claubornensis* LEA, Syn., 1838, p. 19.

* *Margaron (Unio) claubornensis* LEA, Syn., 1852, p. 28; 1870, p. 44.

† * *Unio obtusus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 287; * Tr. Am. Phil. Soc., VIII, 1843, p. 201, pl. XI, fig. 13; * Obs., III, 1842, p. 39, pl. XI, fig. 13.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* CHENU, Ill. Conch. 1858, pl. XXXI, figs. 2, 2a, 2b.—* REEVE, Conch. Icon., XVI, 1865, pl. XXVI, fig. 129.—* PÆTEL, Conch. Sam., III, 1890, p. 161.

* *Margaron (Unio) obtusus* LEA, Syn., 1852, p. 39; 1870, p. 62.

* *Unio pallascens* LEA,⁴ Pr. Am. Phil. Soc., IV, 1845, p. 164; * Tr. Am. Phil. Soc., X, 1848, p. 79, pl. VII, fig. 20; * Obs., IV, 1848, p. 79, pl. VII, fig. 20.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* H. and A. ADAMS, Gen. Rec. Moll.,

¹ Probably only a thin, slightly rayed or rayless variety of *L. hydianus*.

² I doubt if this is distinct from *Approximus*.

³ Dr. Lea places this in the synonymy of *Unio lecontianus* Lea. I have never seen an authentic shell, but the figure given by Conrad in the Journal seems to me more like *Lampsilis obtusus*, and I am inclined to think it is a small male specimen of this species having reversed laterals.

⁴ There is only a single specimen in the Lea collection, the type, an old, rather thin, eroded, large shell, in bad condition, but which, I feel sure, is *claubornensis*.

II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

Margaron (Unio) pallescens LEA, Syn., 1852, p. 27; 1870, p. 43.

Gulf drainage of the United States, from the Flint to the Pearl River.

† **LAMPSILIS PORPHYREUS** Lea.¹

* *Unio porphyreus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 60; * JI. Ac. N. Sci. Phila., V, 1862, p. 80, pl. x, fig. 228; * Obs., VIII, 1862, p. 84, pl. x, fig. 228.—B. H. WRIGHT, Check List, 1888.

Margaron (Unio) porphyreus LEA, Syn., 1870, p. 41.

Cahawba and Coosa rivers, Alabama.

† **LAMPSILIS STRAMINEUS** Conrad.

* *Unio stramineus* CONRAD, Am. JI. Sci., XXV, 1834, p. 339, pl. I, fig. 6; * New F. W. Shells, 1834, p. 71.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen., 1836, p. 198.—* CONRAD, Monog., X, 1838, p. 91, pl. L, fig. 1.—* HANLEY, Test. Moll., 1842, p. 209; * Biv. Shells, 1843, p. 209, pl. XXIII, fig. 45.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIX, fig. 217, 1868, pl. LXXX, fig. 422.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

* *Margarita (Unio) stramineus* LEA, Syn., 1836, p. 39; 1838, p. 25.

* *Margaron (Unio) stramineus* LEA, Syn., 1852, p. 28; 1870, p. 35.

Southern Alabama, southern Mississippi.

† **LAMPSILIS REEVIANUS** Lea.

* *Unio reevianus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 272, pl. xx, fig. 28; * Obs., V, 1852, p. 28, pl. xx, fig. 28.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 256.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Conch. Cab. Unio., 1861, p. 247, pl. LXXXIII, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 165.

* *Margaron (Unio) reevianus* LEA, Syn., 1852, p. 28; 1870, p. 44.

Louisiana; Texas; Arkansas.

† **LAMPSILIS POWELLII** Lea.

* *Unio powellii* LEA, Pr. Am. Phil. Soc., V, 1852, p. 252; * Tr. Am. Phil. Soc., X, 1853, p. 270, pl. XIX, fig. 25; * Obs., V, 1852, p. 26, pl. XIX, fig. 25.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* MUSGRAVE, Phot. Conch., 1863, pl. II, fig. 8.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXX, fig. 359.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.

* *Margaron (Unio) powellii* LEA, Syn., 1852, p. 28; 1870, p. 44.

Salina and Clinton, Arkansas; Guadalupe River, Texas; Spring River, Kansas (Cragin).

¹ I am a little uncertain as to where this species should be placed, as I have only seen the two specimens in the Lea collection, which are very poor and distorted. The nacre is purple.

(Group of *Lampsilis ligamentinus*.)

Shell solid, evenly elliptical, not greatly inflated, with a yellowish green epidermis, with broad, usually broken, dark-green rays; beaks not prominent, sculpture fine, restricted in area, pseudocardinals rather small, stumpy; nacre silvery or rarely pinkish. The female shell is not greatly swollen in the post basal region.

† LAMPSILIS LIGAMENTINUS Lamarck.¹

- Unio crassus* SAY, Nich. Encyc., II, 1816, pl. I, fig. 8.—* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 293.—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 278, fig. 1.
 —* FERUSSAC, Guer. Mag., 1835, p. 27.—* HANLEY, Test. Moll., 1842, p. 192;
 * Biv. Shells, 1843, p. 192.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.
 —* SOWERBY, Conch. Icon., XVI, 1866, pl. XL, fig. 220.—* PÆTEL, Conch. Sam., III, 1890, p. 149.
 * *Margarita (Unio) crassus* LEA, Syn., 1836, p. 24; 1838, p. 19.
 * *Mya crassa* EATON, Zool. Text-Book, 1826, p. 217.
 * *Unio ligamentina* LAMARCK, An. sans Vert., VI, 1819, p. 72.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* DESHAYES, An. sans Vert., 2d. ed., VI, 1835, p. 533; 3d ed., II, 1839, p. 668.
 * *Unio ligamentinus* KUSTER, Conch. Cab. Unio, 1852, p. 23, pl. III, fig. 3.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CALKINS, Pr. Ottawa Ac. Sci., 1874, p. 43.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 157.—* CALL, Tr. Ac. Sci. St. Louis, VII, p. 22, pl. XXI.
 * *Margaron (Unio) ligamentinus* LEA, Syn., 1852, p. 28; 1870, p. 43.
 * *Lampsilis ligamentinus* BAKER, Moll. Chicago, Pt. 1, 1898, p. 108, pl. XVI.—* SMITH, Bull. U. S. Fish Com., 1899, p. 291, pl. LXXIX.
 * *Unio ellipticus* BARNES, Am. Jl. Sci., VI, 1823, p. 259, pl. XIII, fig. 19 (outline).—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 278.
 * *Mya elliptica* EATON, Zool. Text-Book, 1826, p. 219.
 * *Unio carinatus* BARNES, Am. Jl. Sci., VI, 1823, p. 259, pl. XI, fig. 10.
 * *Mya carinata* EATON, Zool. Text-Book, 1826, p. 220.
 * *Unio ellipsarius* SAY, Am. Conch., VI, 1834.
 * *Unio fasciatus* SAY, Am. Conch., VI, 1834.—* CONRAD, Monog., 1835, p. 3, pl. I.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* POTIEZ and MICHAUD, Gall. Moll., 1844, 153, pl. LVIII, fig. 5.—* KUSTER, Conch. Cab., 1852, p. 31, pl. v, fig. 3.
 * *Mya gravis* WOOD, Ind. Test. (Hanley) 1856, p. 199, pl. I, fig. 6.
 * *Unio gravis* PÆTEL, Conch. Sam., III, 1890, p. 154.
 * *Unio luteolus* SOWERBY, Conch. Icon., XVI, 1867, pl. LVIII, fig. 293a.
 * *Unio delodontus* SOWERBY, Conch. Icon., XVI, 1867, pl. LVII, fig. 288.
 * *Unio crassidens* SOWERBY, Conch. Icon., XVI, 1868, pl. LXII, fig. 312.
 * *Unio pictus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXII, fig. 313.
 * *Unio venustus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIV, fig. 326.
 * *Unio upsoni* MARSII, Conch., Ex. 1, 1887, p. 51.²—B. H. WRIGHT, Check List, 1888.

¹ Say's name *crassus* applied to this before that of Lamarck was used by Retzius previously for a European *Unio*.

² The description was read before the Mercer County, Illinois, Scientific and Historical Society March 2, 1880. Mr. Marsh had copies of the description published and thinks that the species was described on the records. It was not figured.

† *LAMPSILIS LIGAMENTINUS* var. *GIBBUS* Simpson.¹

* *Unio crassus* SOWERBY, Conch. Icon., XVI, 1868, pl. xcv, fig. 520.

Mississippi River drainage generally; southern Michigan; western New York; Manitoba; Ontario; the variety from the Ohio River southward.

† *LAMPSILIS ORBICULATUS* Hildreth.

* *Unio orbiculatus* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 284.—* HANLEY, Test. Moll., 1842, p. 192; Biv. Shells, 1843, p. 192, pl. XXI, fig. 2.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 254.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Conch. Cab., 1861, p. 216, pl. LXXI, figs. 3, 4.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLIV, fig. 239.—* CALKINS, Pr. Ottawa Ac. Sci., 1874, p. 44.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Margarita (Unio) orbiculatus* LEA, Syn., 1836, p. 25; 1838, p. 19.

* *Margaron (Unio) orbiculatus* LEA, Syn., 1852, p. 28; 1870, p. 43.

* *Unio abruptus* SAY, Am. Conch., No. II, 1831, pl. XVII.—* CONRAD, New F. W. Shells, 1834, p. 67.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 24, pl. VI, figs. 1, 1a, 1b.—* DESHAYES, An. sans Vert., 2d ed., VI, 1335, p. 555; * 3d ed., II, 1839, p. 674.

* *Unio cyclops* FERUSSAC, Guer. Mag., 1835, p. 28.

* *Unio crassus* CONRAD, Monog., IV, 1836, p. 34, pl. XVI.

Ohio and Cumberland rivers; west to the Mississippi River; north to Minnesota. Some specimens can hardly be separated from *L. higginsii*.

† *LAMPSILIS HIGGINSII* Lea.²

* *Unio higginsii* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 84; * Jl. Ac. N. Sci. Phila., V, 1862, p. 188, pl. XXIV, fig. 258; * Obs., IX, 1863, p. 10, pl. XXIV, fig. 258.—

* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXII, fig. 431.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 155.

* *Margaron (Unio) higginsii* LEA, Syn., 1870, p. 41.

Ohio River, west to Iowa, and southwest to Kansas.

† *LAMPSILIS PINGUIS* Lea.³

Unio pinguis LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 84; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 78, pl. xv, fig. 58; * Obs., VI, 1858, p. 78, pl. xv, fig. 58.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) pinguis* LEA, Syn., 1870, p. 44.

St. Peters River, Minnesota. Doubtful.

¹ A peculiarly humped form of *ligamentinus*, common throughout the southern range of the species, may bear the above name.

² A somewhat puzzling species, of which male specimens often approach closely to *Obovaria ellipsis* in form, but the pseudo cardinals are always stumpy and never longitudinal, as they generally are in that genus. It is probably an aberrant form, closely related to *L. orbiculatus*. The females are more inflated in the postbasal region than in any other member of the group.

³ I have seen only one specimen of this, the type, which is an exceedingly puzzling form. The hinge line is injured and curves in strongly in front of the beaks, where there is a large, irregular lunule, and the pseudocardinals are compressed. It may be a young, diseased *orbiculatus*.

† LAMPSILIS TÆNIATUS Conrad.

- * *Unio tæniatus* CONRAD, New F. W. Shells, 1834, pp. 26, 72, pl. IV, fig. 2.¹—*FERUS-SAC, Guer. Mag., 1835, p. 29.—MÜLLER, Syn. Nov. Gen., 1836, p. 201.—*HANLEY, Test. Moll., 1842, p. 191; *Biv. Shells, 1843, p. 191.—*CHENU, Bib. Conch., 1st ser., III, 1845, p. 15.—*CATLOW and REEVE, Conch. Nom., 1845, p. 64.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—*KUSTER, Conch. Cab. Unio, 1862, p. 271, pl. XCI, fig. 4.—*B. H. WRIGHT, Check List, 1888.
- * *Margarita (Unio) tæniatus* LEA, Syn., 1836, p. 24; 1838, p. 19.
- * *Unio tæniatus* PÆTEL, Conch. Sam., III, 1890, p. 169.
- * *Unio menkianus* LEA, Tr. Am. Phil. Soc., V, 1836, p. 76, pl. XIX, fig. 59; *Obs., II, 1838, p. 76, pl. XIX, fig. 59.—*TROSCHER, Arch. für Naturg., V, 1839, Pt. 2, p. 237.—*HANLEY, Test. Moll., 1842, p. 191; *Biv. Shells, 1843, p. 191, pl. XXIII, fig. 4.—*CATLOW and REEVE, Conch. Nom., 1845, p. 61.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—*CHENU, Ill. Conch., 1858, pl. XIX, figs. 4, 4a, 4b.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—*KUSTER, Conch. Cab. Unio, 1861, p. 245, pl. LXXXII, fig. 3.—*REEVE, Conch. Icon., XVI, pl. XXVIII, fig. 140.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 159.
- * *Margarita (Unio) menkianus* LEA, Syn., 1838, p. 19.
- * *Margaron (Unio) menkianus* LEA, Syn., 1852, p. 27; 1870, p. 43.
- * *Unio pulcher* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 6, pl. III, fig. 6; *Obs., II, 1838, p. 6, pl. III, fig. 6.—*TROSCHER, Arch. für Naturg., V, 1839, Pt. 2, p. 234.—*HANLEY, Test. Moll., 1842, p. 193; *Biv. Shells, 1843, p. 193, pl. XXIII, fig. 7.—*CATLOW and REEVE, Conch. Nom., 1845, p. 63.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*CHENU, Ill. Conch., 1858, pl. XVII, figs. 2, 2a, 2b.—*KUSTER, Conch. Cab. Unio, 1861, p. 193, pl. LXI, fig. 5.—*P. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 164.
- * *Margarita (Unio) pulcher* LEA, Syn., 1836, p. 25; 1838, p. 19.
- * *Margaron (Unio) pulcher* LEA, Syn., 1852, p. 28; 1870, p. 44.
- * *Unio interruptus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 15, pl. VI, fig. 15; *Obs., II, 1838, p. 15, pl. VI, fig. 15.—*HANLEY, Test. Moll., 1842, p. 191; *Biv. Shells, 1843, p. 191, pl. XXI, fig. 28.—*CATLOW and REEVE, Conch. Nom., 1845, p. 60.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*CHENU, Ill. Conch., 1858, pl. XVII, figs. 4, 4a, 4b.—*B. H. WRIGHT, Check List, 1888.
- * *Margarita (Unio) interruptus* LEA, Syn., 1836, p. 24; 1838, p. 19.
- * *Margaron (Unio) interruptus* LEA, Syn., 1852, p. 27; 1870, p. 43.
- * *Unio latiradiatus* CONRAD, Monog., XI, 1838, p. 96, pl. LIII; Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—*KUSTER, Conch. Cab. Unio, 1861, p. 208, pl. LXIX, fig. 3.—PÆTEL, Conch. Sam., III, 1890, p. 151.
- * *Unio tennesseensis* LEA,² Pr. Am. Phil. Soc., I, 1840, p. 288; *Tr. Am. Phil. Soc., VIII, 1843, p. 199, pl. x, fig. 11;³ *Obs., III, 1842, p. 37, pl. x, fig. 11.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU, Ill. Conch., 1858, pl. XXVII, figs. 1, 1a, 1b.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 169.
- * *Margaron (Unio) tennesseensis* LEA, Syn., 1852, p. 27; 1870, p. 43.
- * *Unio perdir* REEVE, Conch. Icon., XVI, 1864, pl. XVIII, fig. 82.

Tennessee and Cumberland river systems.

¹ Lea places this species in the synonymy of his *Unio interruptus*, in the Synopsis. The latter was not published until 1838.

² The type is an old female *tæniatus*, very solid, and but slightly rayed.

³ Lea states in the above citation that this species has radial beak sculpture. The beaks of his type are slightly eroded, having the appearance of somewhat rayed ridges, and I have seen the same thing in the eroded beaks of *Unio rotundatus* Lamarck, but it is not true beak sculpture. Many Unios when decomposing exhibit throughout their shells a somewhat radial structure, and that is what is shown in this case.

† LAMPSILIS PICTUS Lea.

- * *Unio pictus* LEA, Tr. Am. Phil. Soc., V., 1834, p. 73, pl. XI, fig. 36; * Obs., I, 1834, p. 185, pl. XI, fig. 52.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 191; * Biv. Shells, 1843, p. 191, pl. XXIII, fig. 35.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Conch. Cab. Unio, 1861, p. 249, pl. LXXXIII, fig. 5.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 163.
- * *Margarita (Unio) pictus* LEA, Syn., 1836, p. 24; 1838, p. 19.
- * *Margaron (Unio) pictus* LEA, Syn., 1852, p. 27; 1870, p. 43.
- † * *Unio lindsleyi* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 306; * Jl. Ac. N. Sci. Phila., 1860, p. 351, pl. LVIII, fig. 176; * Obs., VIII, 1860, p. 33, pl. LVIII, fig. 176.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLII, fig. 233b.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 157.
- * *Margaron (Unio) lindsleyi* LEA, Syn., 1870, p. 43.
- * *Unio camelopardalis* SOWERBY, Conch. Icon., XVI, 1866, pl. XLII, fig. 233a.

Harpeth River, Tennessee.

† LAMPSILIS CAMELOPARDILIS Lea.¹

- * *Unio camelopardilis* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 92; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 355, pl. LIX, fig. 180; * Obs., VIII, 1860, p. 37, pl. LIX, fig. 180.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 147.
- * *Margaron (Unio) camelopardilis*, LEA, Syn., 1870, p. 53.
- † * *Unio fucatus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 92; Jl. Ac. N. Sci. Phila., IV, 1860, p. 353, pl. LIX, fig. 178; * Obs., VIII, 1860, p. 35, pl. LIX, fig. 178.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) fucatus*, LEA, Syn., 1870, p. 43.

Tennessee and Cumberland river systems.

† LAMPSILIS PUNCTATUS Lea.

- * *Unio punctatus* LEA, Pr. Ac. N. Sci. Phila., IX, 1865, p. 89; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 261, pl. XXXII, fig. 76; * Obs., XII, 1869, p. 21, pl. XXXII, fig. 76.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) punctatus* LEA, Syn., 1870, p. 43.

Tennessee River drainage.

† LAMPSILIS PERDIX Lea.

- * *Unio perdix* LEA, Tr. Am. Phil. Soc., V, 1827, p. 72, pl. XI, fig. 31; * Obs., I, 1834, p. 184, pl. XI, fig. 31.—* FERUSSAC Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 189; * Biv. Shells, 1843, p. 188, pl. XX, fig. 20.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* H. and A. ADAMS, Gen. Rec. Moll. II, 1857, p. 494.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIX, fig. 82.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 163.
- * *Margarita (Unio) perdix* LEA, Syn., 1836, p. 23; 1838, p. 18.
- * *Margaron (Unio) perdix* LEA, Syn., 1852, p. 26; 1870, p. 41.
- * *Unio pectorosus* CONRAD, New F. W. Shells, 1834, p. 37, pl. VI, fig. 1, p. 71.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* CONRAD, Monog. III, 1836, p. 25, pl. XI, fig. 1.—* MÜLLER, Syn. Nov. Gen., 1836, p. 205.—* HANLEY, Test. Moll., 1842, p. 189; * Biv. Shells, 1843, p. 189, pl. XXI, fig. 22.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 19, pl. II, fig. 8.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* KUSTER, Conch. Cab. Unio, 1852, p. 26, pl. IV, fig. 2.—* CONRAD,

¹ It is quite probable that this and the preceding species are merely varieties of the same thing.

Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* B. H. WRIGHT, Check List, 1888.—

* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Margarita (Unio) pectorosus* LEA, Syn., 1836, p. 23, 1838, p. 18.

Tennessee and Cumberland river systems.

† *LAMPSILIS BRACTEATUS* Gould.

* *Unio bracteatus* GOULD, Pr. Bost. Soc. N. Hist., V, 1855, p. 228; * *Otia* Conch., 1862, p. 217.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) bracteatus* LEA, Syn., 1870, p. 43.

Llanos, Guadalupe, and Colorado rivers, Texas.

† *LAMPSILIS VENUSTUS* Lea.

* *Unio venustus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 4, pl. II, fig. 4; * *Obs.*, II 1838, p. 4, pl. II, fig. 4.—* TROSCHER, Arch. für. Naturg., V, 1839, II, p. 233.—

* HANLEY, Test. Moll., 1842, p. 192; * *Biv. Shells*, 1843, p. 192, pl. XXIII, fig. 5.—

* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* CHENU, Ill. Conch., 1858, pl. XXII, figs. 1, 1a, 1b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 171.—* CALL, Tr. Acad. Sci. St. Louis, 1895, p. 57.

* *Margarita (Unio) venustus* LEA, Syn., 1836, p. 24; 1838, p. 19.

* *Margaron (Unio) venustus* LEA, Syn., 1852, p. 27; 1870, p. 43.

Potosi, Missouri.

(Group of *Lampsilis anodontoides*.)

Shell elongated, inflated, moderately solid, sharply pointed behind; that of the female much produced at the posterior base, behind which it is diagonally truncate; epidermis very bright, smooth and shining, but feebly rayed; beaks not prominent, marked with very delicate regular sculpture, the posterior loop often open behind; pseudocardinals smooth, generally compressed; nacre shining, white or purplish. Animal with the marsupium very distinct and well rounded below; inner edge of the mantle usually beautifully and evenly toothed behind.

† *LAMPSILIS ANODONTOIDES* Lea.

† * *Unio anodontoides* LEA, Tr. Am. Phil. Soc., IV, 1834, p. 81, pl. VIII, fig. 11;

* *Obs.*, I, 1834, p. 91, pl. VIII, fig. 11.—* HANLEY, Test. Moll., 1842, p. 204.—

* CATLOW and REEVE, Conch. Nom., 1845, p. 55.—* CHENU, Ill. Conch., 1858, pl. XIV, figs. 3, 3a, 3b.—* REEVE, Conch. Icon., XVI, 1865, pl. XIX, fig. 87.—* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Proc. U. S. Nat. Mus., XV, 1892, p. 430, pl. LXXI, fig. 7; LXXII, figs. 1, 2, 4.

* *Margarita (Unio) anodontoides* LEA, Syn., 1836, p. 35; 1838, p. 24.

* *Margaron (Unio) anodontoides* LEA, Syn., 1852, p. 36; 1870, p. 58.

* *Lampsilis anodontoides* BAKER, Moll. Chicago, Pt. 1, 1898, p. 100, pl. x, figs. 1, 2.—* SMITH, Bull. U. S. Fish Com. 1899, p. 290, pl. LXXVIII.

* ? *Unio teres* SAY, Am. Conch., VI, 1834.¹—* CONRAD, New F. W. Shells, 1834, p. 72; Monog. VI, 1836, p. 52, pl. XXVIII.—* FERUSSAC, Guer. Mag., 1835, p. 27.—

* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.—* KUSTER, Conch. Cab. Unio, 1854, p. 68, pl. XVII, figs. 1, 4.

¹ Said to be of Rafinesque, 1820. I can not be positive whether Rafinesque's description is of a young shell of this species or the *fallaciosus*. He gives its length as 3 inches. I therefore use the name applied by Lea, his shell being properly described and figured.

† *LAMPSILIS ANODONTOIDES* var. *FLORIDENSIS* Lea.

- * *Unio floridensis* LEA,¹ Tr. Am. Phil. Soc., X, 1852, p. 274, pl. XXI, fig. 31; * Obs., V, 1852, p. 30, pl. XXI, fig. 31.—* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 249.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 152.
* Margaron (*Unio floridensis* LEA, Syn., 1852, p. 39; 1870, p. 62.

Entire Mississippi River drainage except (probably) the upper Missouri. All the Gulf drainage from the Withlacoochee River, Florida, to the Rio Grande, and into Mexico.

† *LAMPSILIS FALLACIOSUS* (Smith) Simpson.²

- * *Unio anodontoides* REEVE, Conch. Icon., XVI, 1865, pl. XXI, fig. 97.—* CALKINS, Pr. Ottawa Acad., 1874, p. 41.
* *Lampsilis anodontoides* BAKER, Moll. Chicago, Pt. 1, 1898, pl. x, fig. 3.³
* *Unio oricus* SOWERBY, Conch. Icon., 1868, pl. LXIII, fig. 314.
* *Lampsilis fallaciosus* SMITH, Bull. U. S. Fish Com., 1899, p. 291, pl. LXXIX.—* SIMPSON, Pr. Ac. N. Sci., Phila., 1900, p. 74, pl. II, fig. 5.

Upper Mississippi drainage; south to the Cumberland River, Tennessee, and to Arkansas; Red River of the North?

† *LAMPSILIS VIRESCENS* Lea.⁴

- * *Unio virescens* LEA, Pr. Ac. N. Sci., Phila., II, 1858, p. 40; * JI. Ac. N. Sci., Phila., IV, 1860, p. 341, pl. LV, fig. 166; * Obs., VIII, 1860, p. 23, pl. LV, fig. 166.—* B. H. WRIGHT, Check List, 1888.
* Margaron (*Unio virescens* LEA, Syn., 1870, p. 42.

Tennessee River in northern Alabama.

* † *LAMPSILIS AUSTRALIS* Simpson.

- * *Lampsilis australis* SIMPSON,⁵ Pr. Ac. N. Sci., Phila., 1900, p. 75, pl. II, fig. 2.

Little Patsaliga Creek, southeastern Alabama.

† *LAMPSILIS RECTUS* Lamarck.

- * *Unio recta* LAMARCK, An. sans Vert., VI, 1819, p. 74.—* VALENCIENNES, Rec. Obs. Zool., II, 1833, p. 234, pl. LIV, fig. 1.—* DESHAYES, An. sans Vert., 2d ed., VII, 1835, p. 537; 3d ed., II, 1839, p. 669.

¹ *Lampsilis anodontoides* becomes thinner and more delicate in the waters of the eastern Gulf drainage, and in Florida assumes the form called by Lea *Unio floridensis*. The type of the latter is not in the Lea collection.

² Figured and the name (supplied by the writer) given by Smith in Bull. U. S. Fish Commission, but not described.

³ As this species has generally been confounded with the very closely allied *L. anodontoides* of Lea, it is a little difficult to be certain in all cases which form authors have referred to.

⁴ A species which seems to combine some of the characters of *L. anodontoides* and *cariosus*, but which probably should be placed here.

⁵ This species seems to stand partly between *L. rectus* and *L. anodontoides*. In the character of color both of naere and epidermis the former is quite widely separated from the latter, though specimens of *rectus* from Michigan and other northern localities sometimes have a tawny epidermis. In *L. australis* the naere is silvery, and the color of the epidermis approaches to some extent that of *anodontoides*.

- * *Unio (Ligumia) recta* SWAINSON, Treat. on Mal., 1840, pp. 267, 274, fig. LV.
Unio rectus SHORT and EATON, Transylvania JI., 1831, p. 77.—* CONRAD, New F. W. Shells, 1834, p. 71.—* SAY, Am. Conch., VI, 1834.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* CONRAD, Monog., III, 1836, p. 33, pl. XV.—* HANLEY, Test. Moll., 1842, p. 204.—* C. B. ADAMS, Thompson's Hist. of Vt., 1842, p. 167; * F. W. and L. Sh. of Vermont, 1842, p. 17.—* HANLEY, Biv. Shells, 1843, p. 204.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 195.—* POTIEZ and MICHAUD, Gall. Moll., 1844, II, p. 149, pl. LVII, figs. 3, 4.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* KUSTER, Conch. Cab. Unio, 1852, p. 35, pl. VI, fig. 1; VII, fig. 1.—* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 256.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* CHENU, Man., 1859, II, p. 139, fig. 675.—REEVE, Conch. Icon., XVI, 1865, pl. XIX, fig. 86.—* CALKINS, Pr. Ottawa Ac., 1874, p. 44.—* LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 50.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 165.—* CALL, Tr. Ac. Sci., St. Louis, VII, 1895, p. 43, pl. VII.
 * *Margarita (Unio) rectus* LEA, Syn., 1836, p. 34; 1838, p. 23.
 * *Margaron (Unio) rectus* LEA, Syn., 1852, p. 35; 1870, p. 57.
 * *Lampsilis rectus* SMITH, Bull. U. S. Fish Com., 1899, p. 290, pl. LXXVIII.
 † * *Unio latissima* RAFINESQUE, Ann. Gen. Sci. Phys. Brux., Pt. 13, V, 1820, p. 297, pl. LXXX, figs. 14, 15.
 * ? *Unio praelongus* BARNES,¹ Am. JI. of Sci., VI, 1823, p. 261, pl. XIII, fig. 11.—
 * HILDRETH, Am. JI. of Sci., XIV, 1828, p. 286, fig. 18.
 * *Mya praelonga* EATON, Zool. Text-Book, 1826, p. 220.—WOOD, Ind. Test (Hanley), 1856, p. 200, pl. I, Supp., fig. 11.
 * *Eurynea praelonga* STIMPSON, Shells of N. Eng., 1851, p. 13.
 † * *Unio argeri* CONRAD, Monog., VI, 1836, p. 53, pl. XXIX, fig. 1.
 * ? *Unio arquatus*² CONRAD, JI. Ac. Nat. Sci. Phila., 1854, p. 297, pl. XXVI, fig. 8.—B. H. WRIGHT, Check List, 1888.

Entire Mississippi drainage; Alabama River drainage; Red River of the North; St. Lawrence system. DeCamp's *Unio anodontoides*, reported from Michigan, is a tawny colored variety of this species according to Walker.

(Group of *Lampsilis nasutus*.)

Shell elongated, thin, compressed, with a distinct posterior ridge, and drawn out to a long, pointed beak behind; epidermis rather dull, olive green, and feebly rayed; beaks low, sculptured with fine, close-set ridges, which are slightly looped in front and are parallel with the axis of the shell behind; hinge teeth delicate and compressed; nacre bluish; female shell not greatly produced in the post-basal region. Animal with the marsupium often occupying the greater part of the length of the outer gills posteriorly; inner gills free or united to the abdominal sac.

† LAMPSILIS NASUTUS Say.

Unio nasutus SAY, Nich. Encyc., 1st ed., 1816, pl. IV, fig. 1.—* SWAINSON, Zool. Ill., 1st ser., I, 1821, pl. LVII.—* CONRAD, New F. W. Shells, 1834, p. 70.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* CONRAD, Monog., IV, 1836, p. 38, pl. XVIII, fig. 1,

¹Barnes only gives an outline of his shell, which appears to me more like his *U. gibbosus* than *rectus*. However in deference to the opinions of Hildreth, Lea, and others, I place it in the synonymy of *L. rectus*. Hildreth's description answers fairly well to *rectus*, and it was submitted to Barnes for his inspection.

²Lea believes this to be a distorted *rectus*. I am not quite certain of it.

- part.¹—* GOULD, Inv. of Mass., 1841, p. 109, fig. 71.—* BINNEY, Gould's Inv., 1870, p. 169, fig. 473.—* HANLEY, Test. Moll., 1842, p. 206.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 191, pl. xx, fig. 239.—* HANLEY, Biv. Shells, 1843, p. 206.—CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* KUSTER, Conch. Cab. Unio, 1861, p. 223, pl. LXXV, fig. 4.—* REEVE, Conch. Icon., XVI, 1865, pl. XXI, fig. 94.—* HARTMAN and MICHENER, Conch. Cest., 1874, p. 70, fig. 186.—* B. H. WRIGHT, Check List, 1888.—* H. CARPENTER, Naut., III, 1889, p. 94.—* PÆTEL, Conch. Sam., III, 1890, p. 160.
- * *Margarita (Unio) nasutus* LEA, Syn., 1836, p. 37; 1838, p. 24.
- * *Margaron (Unio) nasutus* LEA, Syn., 1852, p. 37; 1870, p. 60.
- * *Unio nasuta* LAMARCK, An. sans Vert., VI, 1819, p. 75.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 538; 3d ed., II, 1839, p. 669.—* HANLEY, Test. Moll., 1842, p. 208; * Biv. Shells, 1843, p. 208.
- * *Mya nasuta* EATON, Zool. Text-Book, 1826, p. 218.—* WOOD, Index Test. (Hanley), 1856, p. 199, pl. 1, Supp., fig. 4.
- * *Eurynea nasuta* AGASSIZ, Shells of New Eng., 1851, p. 13; Arch. für Naturg., I, 1852, p. 45.
- * *Unio rostrata* VALENCIENNES, Rec. Obs. Zool., II, 1833, p. 233, pl. LIII, fig. 3.
- * *Unio vaughanianus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXI, fig. 308.

St. Lawrence drainage; streams flowing into the Atlantic south to North Carolina.

(Group of *Lampsilis subrostratus*.)

Shell elliptical to elongate, distinctly pointed behind, the point well above the middle of the height, that of the female conspicuously swollen at the posterior base and obliquely truncated behind; epidermis varying from olive to brown, generally more or less rayed and slightly roughened; beaks not prominent, sculptured with fine, delicate, parallel bars, which are somewhat looped in front and generally descend obliquely behind; pseudocardinals usually compressed; nacre bluish to purple. Animal with inner gills often more or less free from the abdominal sac; posterior base of mantle of female generally toothed.

† *LAMPSILIS SUBROSTRATUS* Say.

- Unio subrostratus* SAY, N. Harmony Diss., Jan. 15, 1831; *Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 72.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* L. W. SAY, Terr. and Fluv. Shells, 1840, p. 7.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—KUSTER, Conch. Cab., 1861, p. 203, pl. LXVIII.—* LEWIS, Pr. Ac. N. Sci. Phila., VIII, 1878, p. 273.—* CALL, Am. Nat., XIII, p. 392.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch., Sam., III, 1890, p. 168.
- * *Unio nashrillianus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 100, pl. XIV, fig. 43; *Obs., I, 1834, p. 212, pl. XIV, fig. 43.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 193; Biv. Shells, 1843, p. 193, pl. XXIII, fig. 31.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—* KUSTER, Conch. Cab. Unio, 1861, p. 224, pl. LXXVI, figs. 1, 2.—SOWERBY, Conch. Icon., XVI, 1866, pl. XXX, fig. 158.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 160.

¹ Conrad gives three figures under fig. 1. The first and second are *U. fisherianus*; the third is a female *nasutus*.

- **Margarita (Unio) nashvillianus* LEA, Syn., 1836, p. 26; 1838, p. 20.
 **Margaron (Unio) nashvillianus* LEA, Syn., 1852, p. 29.
 **Margaron (Unio) nashvilliensis* LEA, Syn., 1870, p. 45.
 †**Unio mississippiensis* CONRAD, Jl. Ac. N. Sci. Phila., I, 1850, p. 277, pl. xxxviii, fig. 11.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*KUSTER, Conch. Cab. Unio, 1861, p. 245, pl. lxxxii, fig. 3.—*REEVE, Conch. Icon., XVI, 1865, pl. xix, fig. 85.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 159.
 **Margaron (Unio) mississippiensis* LEA, Syn., 1852, p. 29; 1870, p. 60.
 **Unio rutersvillensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 155; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 355, pl. ix, fig. 181; *Obs., VIII, 1860, p. 38, pl. lx, fig. 181.—*B. H. WRIGHT, Check List, 1888.
 **Margaron (Unio) rutersvillensis* LEA, Syn., 1870, p. 43.
 †**Unio topekaensis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 144; *Jl. Ac. N. Sci. Phila., VI, 1869, p. 313, pl. xlix, fig. 126; *Obs., XII, 1869, p. 73, pl. xlix, fig. 126.—*B. H. WRIGHT, Check List, 1888.
 **Margaron (Unio) topekaensis* LEA, Syn., 1870, p. 43.
 **Unio cocoduensis* REEVE, Conch. Icon., XVI, 1865, pl. 24, fig. 117.

Entire Mississippi drainage north to about latitude 41°. Eastern half of Texas.

†***LAMPSILIS LIENOSUS** Conrad.

- **Unio lienosus* CONRAD, Am. Jl. Sci., XXV, 1834, p. 339, pl. i, fig. 4; *New F. W. Shells, 1834, p. 70.—*FERUSSAC, Guer. Mag., 1835, p. 25.—*CONRAD, Monog. VII, 1836, p. 60, pl. xxxiv, fig. 2.—*MÜLLER, Syn. Nov. Gen., 1836, p. 197.—*HANLEY, Test. Moll., 1842, p. 194; *Biv. Shells, 1843, p. 194, pl. xxi, fig. 32.—*CATLOW and REEVE, Conch. Nom., 1845, p. 60.—*KUSTER, Conch. Cab. Unio, 1852, p. 67, pl. xvi, fig. 3.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*SOWERBY, Conch. Icon., XVI, 1866, pl. xxxii, fig. 166; 1868, pl. lxxxv, fig. 388.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 157.
 **Margarita (Unio) lienosus* LEA, Syn., 1836, p. 26; 1838, p. 20.
 **Margaron (Unio) lienosus* LEA, Syn., 1852, p. 29; 1870, p. 45.
 **Unio saxeus* CONRAD, Monog., XII, 1840, p. 109, pl. lx, fig. 1; Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 166.
 **Margaron (Unio) saxeus* LEA, Syn., 1852, p. 27; 1870, p. 42.
 **Unio caliginosus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 165; *Tr. Am. Phil. Soc., X, 1848, p. 79, pl. vii, fig. 21; *Obs., IV, 1848, p. 53, pl. vii, fig. 21.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 246.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*KUSTER, Conch. Cab. Unio, 1861, p. 180, pl. lvi, fig. 7.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 147.
 **Margaron (Unio) caliginosus* LEA, Syn., 1852, p. 29; 1870, p. 45.
 **Unio fontanus* CONRAD,¹ Am. Jl. Sci., II, 1866, p. 279, pl. xv, fig. 13.—*B. H. WRIGHT, Check List, 1888.

LAMPSILIS LIENOSUS var. **UNICOSTATUS** B. H. Wright.

- **Unio unicostatus* B. H. WRIGHT, Naut., XIII, 1899, p. 69.

Lower Mississippi River drainage north to the lower Ohio; east to southwest Georgia.

¹The characters are poor in the indifferent type in the Academy of Natural Sciences in Philadelphia, but I believe that it is a *Lampsilis lienosus*.

† LAMPSILIS CONCESTATOR Lea.

- * *Unio concestator* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 31; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 66, pl. XII, fig. 48; *Obs., VI, 1858, p. 66, pl. XII, fig. 48.—
 *KUSTER, Conch. Cab. Unio, 1861, p. 178, pl. LVI, fig. 3.—*REEVE, Conch. Icon., XVI, 1865, pl. XXVI, fig. 128.—*B. H. WRIGHT, Check List, 1888.—
 *PÆTEL, Conch. Sam., III, 1890, p. 148.—*SIMPSON, Pr. U. S. Nat. Mus., XVI, 1892, p. 416, pl. LVIII, figs. 2-4.

**Margaron (Unio) concestator* LEA, Syn., 1870, p. 45.

- †* *Unio intercedens* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 32; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 77, pl. XV, fig. 57; *Obs., VI, 1858, p. 77, pl. XV, fig. 57.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 155.

**Margaron (Unio) intercedens* LEA, Syn., 1870, p. 45.

- †* *Unio fallax* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 32; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 79, pl. XV, fig. 59; *Obs., VI, 1858, p. 79, pl. XV, fig. 59.—*KUSTER, Conch. Cab. Unio, 1861, p. 206, pl. LXVIII, fig. 7.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXX, fig. 418.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 152.

**Margaron (Unio) fallax* LEA Syn., 1870, p. 45.

- †* *Unio contiguus* LEA,¹ Proc. Ac. N. Sci. Phila., VI, 1861, p. 392; *Jl. Ac. N. Sci. Phila., V, 1862, p. 199, pl. XXVIII, fig. 268; Obs., IX, 1863, p. 21, pl. XXVIII, fig. 268.—B. H. WRIGHT, Check List, 1888.—PÆTEL, Conch. Sam., III, 1890, p. 148.

**Margaron (Unio) contiguus* LEA, Syn., 1870, p. 45.

- **Unio bi-caclatus* REEVE, Conch. Icon., XVI, 1865, pl. XXVI, fig. 130.—*PÆTEL, Conch. Sam., III, 1890, p. 146.

- †* *Unio genuinus* LEA, Pr. Ac. N. Sci. Phila., XX, 1868, p. 161; *Jl. Ac. N. Sci. Phil., VI, 1868, p. 305, pl. XLVI, fig. 117; *Obs., XII, 1869, p. 64, pl. XLVI, fig. 117.—
 *B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) genuinus* LEA, Syn., 1870, p. 45.

North Carolina. in streams flowing into the sea, to Louisiana, and possibly to Texas. Some specimens are very close to *lienosus*.

LAMPSILIS PROPRIUS Lea.²

- * *Unio proprius* LEA, Pr. Ac. N. Sci. Phila., IX, 1865, p. 89; *Jl. Ac. N. Sci. Phila., VI, 1869, p. 256, pl. XXXI, fig. 70; *Obs., XII, 1869, p. 16, pl. XXXI, fig. 70.—
 *B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) proprius* LEA, Syn., 1870, p. 45.

- **Unio striatus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVIII, fig. 407.

Walker County, Georgia; Clinch River, Virginia.

LAMPSILIS PUNICEUS Haldeman.

- * *Unio puniceus* HALDEMAN,³ Jl. Ac. N. Sci. Phila., VIII, 1842, p. 201.—CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) puniceus* LEA, Syn., 1870, p. 46.

Holston River, Tennessee.

¹ A large rather thin *L. concestator*, I think.

² Possibly distinct, but more likely a mere light-colored variety of *L. Vanuxemensis*.

³ I believe that this species has never been figured. From the description I should judge that it belonged in this group, and it may be synonymous with some one of its species.

† LAMPSILIS VANUXEMENSIS Lea.

* *Unio vanuxemensis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 36, pl. XI, fig. 31; * Obs., II, 1838, p. 36, pl. XI, fig. 31.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 436.—* HANLEY, Test. Moll., 1842, p. 193; * Biv. Shells, 1843, p. 193, pl. XXII, fig. 56.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* CHENU, Ill. Conch., 1858, pl. XIX, figs. 3, 3a, 3b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* KUSTER, Conch. Cab. Unio, 1862, p. 272, pl. XCI, fig. 6.—* PÆTEL, Conch. Sam., III, 1890, p. 171.

* *Margarita (Unio) vanuxemensis* LEA, Syn., 1836, p. 26; 1838, p. 19.

* *Margaron (Unio) vanuxemensis* LEA, Syn., 1852, p. 29.

* *Unio vanuxemensis* CONRAD, Pr. Ac. N. Sci. Phila., 1853, p. 259.

* *Margaron (Unio) vanuxemii* LEA, Syn., 1870, p. 46.

* *Unio vanuxemii* B. H. WRIGHT, Check List, 1888.

† * *Unio nitens* LEA, Pr. Am. Phil. Soc., I, 1840, p. 288; * Tr. Am. Phil. Soc., VIII, 1843, p. 205, pl. XII, fig. 19; * Obs., III, 1842, p. 43, pl. XII, fig. 19.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.—* KUSTER, Conch. Cab., 1856, p. 163, pl. XLVII, fig. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* CHENU, Ill. Conch., 1858, pl. XXVIII, figs. 5, 5a, 5b.—* REEVE, Conch. Icon. XVI, 1865, pl. XXVIII, fig. 144.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 161.

* *Margaron (Unio) nitens* LEA, Syn., 1852, p. 29; 1870, p. 45.

† * *Unio umbrans* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 104; * Jl. Ac. N. Sci. Phila., V, 1858, p. 72, pl. XIII, fig. 53; * Obs., VI, 1858, pl. XIII, fig. 53.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIV, fig. 179.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 171.

* *Margaron (Unio) umbrans* LEA, Syn., 1870, p. 45.

† * *Unio tenebricus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 171; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 83, pl. XVII, fig. 63; * Obs., VI, 1858, p. 83, pl. XVI, fig. 63.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) tenebricus* LEA, Syn., 1870, p. 45.

† * *Unio pybasii* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 40; * Jl. Ac. N. Sci. Phila., V, 1862, p. 67, pl. VI, fig. 216; * Obs., VIII, 1862, p. 71, pl. VI, fig. 216.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.

* *Margaron (Unio) pybasii* LEA, Syn., 1870, p. 45.

† * *Unio fabaceus* LEA,¹ Proc. Ac. N. Sci. Phila., V, 1861, p. 38; * Jl. Ac. N. Sci. Phila., V, 1862, p. 90, pl. XIII, fig. 238; * Obs., VIII, 1862, p. 94, pl. XIII, fig. 238.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) fabaceus* LEA, Syn., 1870, p. 46.

† * *Unio copei* LEA,² Pr. Ac. N. Sci. Phila., XII, 1868, p. 144; * Jl. Ac. N. Sci. Phila., VI, 1869, p. 307, pl. XLVII, fig. 120; * Obs., XII, 1869, p. 67, pl. XLVII, fig. 120.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) copei* LEA, Syn., 1870, p. 45.

† * *Unio dispansus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 191; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 19, pl. VI, fig. 16; * Obs., XIII, 1874, p. 23, pl. VI, fig. 16.—* B. H. WRIGHT, Check List, 1888.

Cumberland and Tennessee river systems; headwaters of the Coosa.

† LAMPSILIS OBSCURUS Lea.

* *Unio obscurus* LEA, Tr. Am. Phil. Soc., VI, 1839, p. 7, pl. III, fig. 7, * Obs., II, 1838, p. 7, pl. III, fig. 7.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 234.—* HANLEY, Test. Moll., 1842, p. 194; * Biv. Shells, 1843, p. 194, pl. XXII, fig. 58.—

¹ A small, rather short form of *vanuxemensis*.

² The type is rather thin and high, but is no doubt a mere variation of the above species.

- * CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* CHENU, Ill. Conch., 1858, pl. XVII, figs. 1, 1a, 1b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Conch. Cab. Unio, 1861, p. 220, pl. LXXIV, fig. 3.—* PÆTEL, Conch. Sam., III, 1890, p. 161.
- * *Margarita (Unio) obscurus* LEA, Syn., 1836, p. 26; 1838, p. 20.
- * *Margaron (Unio) obscurus* LEA, Syn., 1852, p. 29; 1870, p. 45.
- * *Margarita (Unio) zeiglerianus* LEA, Syn., 1836, p. 26; 1838, p. 20.
- * *Unio zeiglerianus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 32, pl. x, fig. 27; * Obs., II, 1838, p. 32, pl. x, fig. 27.—* TROSCHER, Arch. für Naturg., V, 1839, Pt. 2, p. 235.—* HANLEY, Test. Moll., 1842, p. 193, * Biv. Shells, 1843, p. 193, pl. XXII, fig. 57.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XXIV, figs. 2, 2a, 2b.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVI, fig. 19.—* B. H. WRIGHT, Check List, 1888.—PÆTEL, Conch. Sam., III, 1890, p. 172.
- * *Margaron (Unio) zeiglerianus* LEA, Syn., 1852, p. 29; 1870, p. 45.¹

Tennessee and Cumberland river systems; Lower Ohio and its tributaries.

† LAMPSILIS PRATTII Lea.

- * *Unio prattii* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 166; * Jl. Ac. N. Sci. Phila., IV, 1859, p. 206, pl. XXIV, figs. 88, 88a; * Obs., VII, 1859, p. 24, pl. XXIV, fig. 88.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) prattii* LEA, Syn., 1870, p. 45.

Chattahoochee River, Georgia.

† LAMPSILIS VAUGHANIANUS Lea.

- * *Margarita (Unio) vauhanianus* LEA, Syn., 1836, p. 39; 1838, p. 25.
- * *Unio vauhanianus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 5, pl. III, fig. 5; * Obs., II, 1838, p. 5, pl. III, fig. 5.—* TROSCHER, Arch. für Naturg., V, 1839, Pt. 2, p. 234.—* HANLEY, Test. Moll., 1842, p. 208; * Biv. Shells, 1843, p. 208, pl. XXII, fig. 48.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.—* CHENU, Ill. Conch., 1858, pl. XXI, figs. 11a, 11b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 171.
- * *Margaron (Unio) vauhanianus* LEA, Syn., 1852, p. 38; 1870, p. 61.

Near Camden, South Carolina.

† LAMPSILIS OGEECHEENSIS Conrad.

- * *Unio ogeecheensis* CONRAD, Ann. and Mag. N. Hist., IV, 1849, p. 300; Jl. Ac. N. Sci. Phila., I, 1850, p. 275, pl. XXXVII, figs. 3, 4; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.
- * *Unio prevostianus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 269, pl. XIX, fig. 24; * Obs., V, 1852, p. 25, pl. XIX, fig. 24.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* MUSGRAVE, Phot. Conch., 1863, pl. ii, fig. 10.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.
- * *Margaron (Unio) prevostianus* LEA, Syn., 1852, p. 29; 1870, p. 44.
- * *Unio proximus* LEA, Pr. Am. Phil. Soc., V, 1852, p. 252; * Tr. Am. Phil. Soc., X, 1852,

¹ I believe that the *L. obscurus* is only the fully adult form of *U. zeiglerianus*, which when quite young shows very delicate and beautiful rays.

² In this case, owing to delay in the publication of the transactions, the name appeared in the Synopsis first, and under the generic name *Margarita*.

p. 271, pl. XX, fig. 27; *Obs., V, 1852, p. 27, pl. XX, fig. 27.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 164.

**Margaron (Unio) proximus* LEA, Syn., 1852, p. 29; 1870, p. 45.

Unio tenerus SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 416, pl. LVIII, figs. 5 and 8.¹

Neuse River, North Carolina; south to north Florida; west to the Etowah River, Georgia; Duck River, Tennessee.

† **LAMPSILIS CONSTRICTUS** Conrad.

**Unio lienosus* var. *constrictus* CONRAD, Monog. X, 1838, p. 91, pl. XLIX, fig. 4.

**Unio constrictus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—*H. and A. ADAMS, Gen. Rec. Moll. II, 1857, p. 493.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVII, fig. 464.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 148.

**Margaron (Unio) constrictus* LEA, Syn., 1852, p. 29; 1870, p. 46.

**Unio genthii* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 85; *Jl. Ac. N. Sci. Phila., 1862, p. 57, pl. II, fig. 204; *Obs. VIII, 1862, p. 61, pl. II, fig. 204.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 153.

**Margaron (Unio) genthii* LEA, Syn., 1870, p. 41.

James River, Virginia; south to South Carolina; Floyd County, northwest Georgia; northern Alabama; Tennessee.

† **LAMPSILIS APICINUS** Lea.

**Unio apicinus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 32; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 76, pl. XIV, fig. 56; *Obs., VI, 1858, p. 76, pl. XIV, fig. 56.—*B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) apicinus* LEA, Syn., 1870, p. 44.

Othcalooga Creek, Gordon County, Georgia.

† **LAMPSILIS OCCIDENTALIS** Conrad.

**Unio occidentalis* CONRAD, Monog., VII, 1836, p. 64, pl. XXXVI, fig. 1; *Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 161.

**Margaron (Unio) occidentalis* LEA, Syn., 1852, p. 33; 1870, p. 53.

Current and Little Red rivers, Arkansas.

† **LAMPSILIS NIGERRIMUS** Lea.

**Unio nigerrimus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 268, pl. XVIII, fig. 23; *Obs., V, 1852, p. 24, pl. XVIII, fig. 23.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 161.

**Margaron (Unio) nigerrimus* LEA, Syn., 1852, p. 31; 1870, p. 49.

**Unio fuligo* REEVE, Conch. Icon., XVI, 1856, pl. XXX, fig. 159.—*PÆTEL, Conch. Sam., III, 1890, p. 153.

Alexandria, Louisiana, to eastern Texas; Indiana?

(Group of *Lampsilis iris*.)

Shell rather small, elliptical, moderately solid in front, thinner behind; epidermis covered with dark green, broken rays, beak sculp-

¹ As of Ravenel, who, however, never described the species.

ture consisting of irregular ridges and corrugations that have a tendency to fall into two loops, which are sometimes slightly nodulous below; nacre bluish, becoming beautifully iridescent posteriorly. Animal having the inner branchiæ free or united to the abdominal sac; mantle border often maculated and generally, in the female, having papillæ below the marsupium; marsupium having a strong, deep sulcus parallel with and near to the base.

† *LAMPSILIS IRIS* Lea.

* *Unio iris* LEA, Tr. Am. Phil. Soc., III, 1830, p. 439, pl. IX, fig. 18; *Obs., I, 1834, p. 53, pl. XI, fig. 18.—*CONRAD, New F. W. Shells, 1834, p. 69.—*FERUSSAC, Gner. Mag., 1835, p. 26.—*DESHAYES, An. sans Vert, 2d ed., VI, 1835, p. 550; 3d. ed., II, 1839, p. 673.—*HANLEY, Test. Moll. 1842, p. 206; *Biv. Shells, 1843, p. 206, pl. XXI, fig. 37.—*CATLOW and REEVE, Conch. Nom., 1845, p. 60.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—*CHENU, Ill. Conch., 1858, pl. VIII, figs. 7, 7a, 7b.—*REEVE, Conch. Icon., XVI, 1865, pl. XXIX, fig. 148.—*CALKINS, Pr. Ottawa Ac. Sci., 1874, p. 42.—*B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) iris* LEA, Syn., 1836, p. 37; 1838, p. 25.

* *Margaron (Unio) iris* LEA, Syn., 1852, p. 38; 1870, p. 60.¹

* *Lampsilis iris* BAKER, Moll. Chicago, Pt. 1, 1898, p. 105, pl. XIII, fig. 1; XIV, fig. 2.

* *Unio creperus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 33, pl. X, fig. 28; *Obs., II, 1838, p. 33, pl. X, fig. 28.—*TROSCHER, Arch. für Naturg., V, 1839, Pt. 2, p. 235.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—*CHENU, Ill. Conch., 1858, pl. XXIII, figs. 5, 5a, 5b.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) creperus* LEA, Syn. 1836, p. 28; 1838, p. 20.

* *Margaron (Unio) creperus* LEA, Syn. 1852, p. 31; 1870, p. 48.

* *Unio cresserus* HANLEY, Test. Moll., 1842, p. 196.—*CATLOW and REEVE, Conch. Icon., 1845, p. 58.

† * *Unio novi-eboraci* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 104, pl. XXIV, fig. 114; *Obs., II, 1838, p. 104, pl. XXIV, fig. 114.—*TROSCHER, Arch. für Naturg., V, 1839, Pt. 2, p. 237.—*HANLEY, Test. Moll., 1842, p. 190; *Biv. Shells, 1843, p. 190.—*DE KAY, Zool. of N. Y., Pt. 5, 1843, pl. XX, fig. 240.—*CATLOW and REEVE, Conch. Nom., 1845, p. 61.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—*CHENU, Ill. Conch., 1858, pl. XIX, figs. 5, 5a, 5b.—*KUSTER, Conch. Cab. Unio. 1861, p. 221, pl. LXXIV, fig. 4.—*SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVIII, fig. 206.—*CALKINS, Pr. Ottawa Ac. Sci. 1874, p. 43.—*B. H. WRIGHT, Check List, 1888.—*PÄTEL, Conch. Sam. III, 1890, p. 161.

* *Margarita (Unio) novi-eboraci* LEA Syn., 1838, p. 19.

* *Margaron (Unio) novi-eboraci* LEA, Syn., 1852, p. 27; 1870, p. 60.

* *Unio radiatus* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 189, pl. XVII, fig. 236.

* *Unio subrostratus* KUSTER, Conch. Cab. Unio. 1861, p. 203, pl. LXVII, fig. 3.—*REEVE, Conch. Icon., XVI, 1864, pl. XVII, fig. 78.

St. Lawrence drainage; west to southern Michigan; Ohio River system; Illinois and Wisconsin.²

¹ Dr. Lea states in the Synopsis in the above reference that Say's *U. subrostratus* is identical with his *iris*. He is certainly mistaken.

² Lea places *Unio opalinus* Anthony (Am. Jl. Conch. II, 1866, p. 146, pl. VII, fig. 2) in the synonymy of the above species. It is an oval, compressed, greenish-yellow shell according to the figure and description, and seems to have been injured anteriorly. I can not tell what it is. It is credited by Anthony to Michigan.

† LAMPSILIS FATUUS Lea.¹

* *Unio fatuus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 287; *Tr. Am. Phil. Soc., VIII, 1842, p. 201, pl. XI, fig. 14; *Obs., III, 1842, p. 39, pl. XI, fig. 14.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—*CHENU, Ill. Conch., 1858, pl. XXXI, figs. 4, 4a, 4b.—*KUSTER, Conch. Cab. Unio, 1862, p. 287, pl. xcvi, fig. 5.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 152.

* *Margaron (Unio) fatuus* LEA, Syn., 1852, p. 38; 1870, p. 61.

* *Unio dactylus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 287; *Tr. Am. Phil. Soc., VIII, 1842, p. 196, pl. IX, fig. 7; *Obs., III, 1842, p. 34, pl. IX, fig. 7.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—*CHENU, Ill. Conch., 1858, pl. XXX, figs. 5, 5a, 5b.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—*B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) dactylus* LEA, Syn., 1852, p. 36; 1870, p. 57.

Tennessee River system. One shell from Beaver River, Pennsylvania seems to be this.

LAMPSILIS PLANICOSTATUS Lea.

Unio planicostatus LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 92; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 354, pl. LIX, fig. 179; *Obs., VIII, 1860, p. 36, pl. LIX, fig. 179.—*B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) planicostatus* LEA, Syn., 1870, p. 43.

Warren County, Kentucky; Clinch River, Tennessee; Tuscumbia, Alabama.

† LAMPSILIS NEBULOSUS Conrad.²

* *Unionebulosus* CONRAD, New F.W. Shells, 1834, p. 28, pl. III, fig. 7; p. 70.—*FERUS-SAC, Guer. Mag., 1835, p. 29.—*MÖLLER, Syn. Nov. Gen., 1836, p. 202.—*CHENU, Bib. Conch., 1st. ser., III, 1845, p. 16, pl. I, fig. 4.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 160.

* *Margarita Unio) cumberlandianus* LEA, Syn., 1836, p. 27; 1838, p. 20.

* *Margaron (Unio) cumberlandianus* LEA, Syn., 1852, p. 30.

†* *Unio cumberlandicus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 25, pl. VII, fig. 19; *Obs., II, 1838, p. 25, pl. VII, fig. 19.—*TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 235.—*HANLEY, Test. Moll., 1842, p. 191; *Biv. Shells, 1843, p. 195, pl. XXII, fig. 59.—*CATLOW and REEVE, Conch. Nom., 1845, p. 58.—*CONRAD, Pr. Ac. N.

¹It is doubtful whether this is more than a variety of the preceding. In general, it has a rather more cylindrical shell, less rayed, and more decidedly and suddenly thickened in front than *iris*, but there are intermediate forms.

²After carefully comparing again and again large series of specimens, I confess that I can not possibly separate the large number of so-called species I have united under this, the oldest name. It may seem a little strange that a species should be found in the Ohio River area and Gulf drainage, but recent research has proven that a large number of our common Upper Mississippi Valley and Ohio River forms are found in the Coosa, Alabama, Black Warrior, and adjoining streams. There is some variation in the brightness of the painting of a number of these so-called species, and in the form of different specimens, but no more than is found in many other abundant, widely-distributed forms, and the figure on pl. III of the New Fresh-water Shells fairly well represents the general manifestation of the males of this species.

- Sci. Phila., VI, 1853, p. 247.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* CHENU, Ill. Conch., 1858, pl. XXIV, figs. 1, 1a, 1b.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVI, fig. 197.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 149.
- * *Margaron (Unio) cumberlandicus* LEA, Syn., 1870, p. 48.
- † * *Unio notatus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 28, pl. VIII, fig. 22; * Obs., II, 1838, p. 28, pl. VIII, fig. 22.—* TROSCHEL, Arch. für Nat., V, 1839, Pt. 2, p. 235.—* HANLEY, Test. Moll., 1842, p. 193; * Biv. Shells, 1843, p. 193, pl. XXIII, fig. 9.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. XXI, figs. 3, 3a, 3b.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 161.
- * *Margarita (Unio) notatus* LEA, Syn., 1836, p. 26; 1838, p. 19.
- * *Margaron (Unio) notatus* LEA, Syn., 1852, p. 29; 1870, p. 45.
- † * *Unio glaber* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 34, pl. X, fig. 29; * Obs., II, 1838, p. 34, pl. X, fig. 29.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 236.—* HANLEY, Test. Moll., 1842, p. 196; * Biv. Moll., 1843, p. 196, pl. XXIII, fig. 2.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. XXIV, figs. 3, 3a, 3b.—* SOWERBY, Conch. Icon., XVI, 1866, pl. LXXXV, fig. 452.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 153.
- * *Margarita (Unio) glaber* LEA, Syn., 1836, p. 28; 1838, p. 20.
- * *Margaron (Unio) glaber* LEA, Syn., 1852, p. 31; 1870, p. 48.
- * *Unio radians* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 32; * Jl. Ac. N. Sci. Phila., IV, 1859, p. 201, pl. XXIII, fig. 84; * Obs., VII, 1859, p. 19, pl. XXIII, fig. 84.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) radians* LEA, Syn., 1870, p. 46.
- † * *Unio jonesii* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 171; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 339, pl. LIV, fig. 164; * Obs., VIII, 1860, p. 21, pl. LIV, fig. 164.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch., Sam., III, 1890, p. 156.
- * *Margaron (Unio) jonesii* LEA, Syn., 1870, p. 48.
- † * *Unio discrepans* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 92; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 340, pl. LV, fig. 165; * Obs., VIII, 1860, p. 22, pl. LV, fig. 165.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIV, fig. 176.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 151.
- * *Margaron (Unio) discrepans* LEA, Syn., 1870, p. 48.
- † * *Unio scitulus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 93; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 342, pl. LV, fig. 167; * Obs., VIII, 1860, p. 24, pl. LV, fig. 167.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) scitulus* LEA, Syn., 1870, p. 45.
- † * *Unio linguaformis* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 305; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 345, pl. LVI, fig. 170; * Obs., VIII, 1860, p. 27, pl. LVI, fig. 170.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) linguaformis* LEA, Syn., 1870, p. 48.
- † * *Unio perpictus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 306; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 350, pl. LVIII, fig. 175; * Obs., VIII, 1860, p. 32, pl. LVIII, fig. 175; * B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) perpictus* LEA, Syn., 1870, p. 44.
- † * *Unio difficilis* LEA,¹ Pr. Ac. N. Sci. Phila., XII, 1868, p. 144; * Jl. Ac. N. Sci. Phila., VI, 1869, p. 311, pl. XLIX, fig. 124; * Obs., XII, 1869, p. 71, pl. XLIX, fig. 124.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) difficilis* LEA, Syn., 1870, p. 46.

¹Two shells are in the Lea collection; one adult and badly worn, the other young, I believe it to be only a form of *nebulosus*.

† ** Unio sparus* LEA, Pr. Ac. N. Sci. Phila., XX, 1868, p. 143; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 306, pl. XLVII, fig. 119; *Obs., XII, p. 66, pl. XLVII, fig. 119.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) sparus* LEA, Syn., 1870, p. 45.

Cumberland and Tennessee rivers systems, Green River, Kentucky; Tombigbee and Alabama rivers drainage; Columbus, Georgia; Wolfs-ville, North Carolina.

† LAMPSILIS MÜHLFELDIANUS Lea.¹

* *Unio mühlfeldianus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 41, pl. XII, fig. 36; *Obs., II, 1838, p. 41, pl. XII, fig. 36.—* TROSCHER, Arch. für Naturg., V, 1839, Pt. 2, p. 236.—* HANLEY, Test. Moll., 1842, p. 195; *Biv. Shells, 1843, p. 196, pl. XXII, fig. 60.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—* CHENU, Ill. Conch., 1858, pl. XVII, figs. 5, 5a, 5b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVIII, fig. 211.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 160.

* *Margarita (Unio) mühlfeldianus* LEA, Syn., 1836, p. 27; 1838, p. 20.

* *Margaron (Unio) mühlfeldianus* LEA, Syn., 1852, p. 30; 1870, p. 48.

Cumberland River; Watauga River near Johnson City, Tennessee.

† LAMPSILIS AMÆNUS Lea.²

* *Unio amœnus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 286; *Tr. Am. Phil. Soc., VIII, 1842, p. 200, pl. x, fig. 12; *Obs., III, 1842, p. 200, pl. x, fig. 12.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 244.—* HANLEY, Biv. Shells, 1856, p. 386, pl. XXIV, fig. 9.—* CHENU, Ill. Conch., 1858, pl. XXXIII, figs. 5, 5a, 5b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIX, fig. 416.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.

* *Margaron (Unio) amœnus* LEA, Syn., 1852, p. 29; 1870, p. 45.

Holston River, Tennessee.

† LAMPSILIS TENER Lea.³

* *Unio tener* LEA, Pr. Am. Phil. Soc., I, 1840, p. 286; *Tr. Am. Phil. Soc., VIII, 1840, p. 198, pl. x, fig. 10; *Obs., III, 1842, p. 36, pl. x, fig. 10.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XXXI, figs. 1, 1a, 1b.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 169.

* *Margaron (Unio) tener* LEA, Syn., 1852, p. 28; 1870, p. 44.

† ** Unio regularis* LEA, Pr. Am. Phil. Soc., II, 1841, p. 82; *Tr. Am. Phil. Soc., VIII, 1842, p. 243, pl. XXV, fig. 59; *Obs., III, 1842, p. 81, pl. XXV, fig. 59.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 256.—* CHENU, Ill. Conch., 1858, pl. XXXIII, fig. 3, 3a, 3b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* SOWERBY,

¹ Lea has only a single specimen in his collection. I consider it a species of doubtful validity.

² Doubtfully valid. More regularly elliptical than *nebulosus*, and the dorsal and basal lines are more nearly parallel.

³ I have only seen the type, a miserably eroded, broken shell, part of which is missing, but I have no doubt that it is the same thing as *regularis*.

Conch. Icon., XVI, 1868, pl. LXVIII, fig. 351.—* B. H. WRIGHT, Check List, 1888.—PÆTEL, Conch. Sam., III, 1890, p. 161.

* *Margaron (Unio) regularis* LEA, Syn., 1852, p. 29; 1870, p. 45.

Big Pigeon and French Broad rivers, Tennessee; near Bowling Green, Kentucky.

† LAMPSILIS SIMUS LEA.

* *Margarita (Unio) simus* LEA, Syn., 1836, p. 29; 1838, p. 21.

* *Unio simus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 26, pl. VIII, fig. 20; * Obs., II, 1838, p. 26, pl. VIII, fig. 20.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 235.—* HANLEY, Test. Moll., 1842, p. 197; * Biv. Shells, 1843, p. 197.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494; * CHENU, Ill. Conch., 1858, pl. XXI, figs. 4, 4a, 4b.—* KUSTER, Conch. Cab. Unio, 1861, p. 248, pl. LXXXIII, fig. 4, p. 262, pl. LXXXVIII, fig. 3.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 167.¹

* *Margaron (Unio) simus* LEA, Syn., 1852, p. 31; 1870, p. 50.

* *Unio notatus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.

* *Unio spatulatus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXV, fig. 328.

* ? *Unio proximus* KUSTER, Conch. Cab. Unio, 1861, p. 248, pl. LXXXIII, fig. 4.

Cumberland and Tennessee river systems; Othcalooga Creek, north-west Georgia.

† LAMPSILIS PLANCUS LEA.²

* *Unio planicus* LEA, Pr. Ac. N. Sci., Phila., IV, 1860, p. 307; * Jl. Ac. N. Sci. Phila., V, 1862, p. 81, pl. x, fig. 229; * Obs., VIII, 1862, p. 85, pl. x, fig. 229.—

* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) planicus* LEA, Syn., 1870, p. 50.

Coosa River at Wetumpka, Alabama.

(Group of *Lampsilis subangulatus*.)

Shell elliptical, inflated, subsolid, drawn to a point behind; that of the female only slightly swollen just behind the central base; bars of the beak sculpture somewhat coarse, feebly double looped; epidermis very smooth, shining, brightly rayed; teeth rather smooth, subcompressed; nacre bluish or purplish. Animal having the marsupium very large; ovisacs numerous; gills large, nearly semicircular.

LAMPSILIS SUBANGULATUS LEA.

* *Unio subangulatus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 287; * Tr. Am. Phil. Soc., VIII, 1842, p. 29, pl. XIII, fig. 23; * Obs., III, 1842, p. 47, pl. XIII, fig. 23.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XXIX, figs. 2, 2a, 2b.—* KUSTER, Conch. Cab. Unio, 1862, p. 278, pl. xciv, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.—* SIMPSON, Pr. U. S. Nat. Mus., XVI, 1892, p. 415, pl. LVIII, fig. 1.

* *Margaron (Unio) subangulatus* LEA, Syn., 1852, p. 29; 1870, p. 45.

* *Unio fasciolus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.

Appalachicola River system, Georgia and Florida.

¹ Spelled *sinus*, but no doubt intended for *simus*.

² Doubtfully distinct from *simus*. I have seen only the type.

† LAMPSILIS KIRKLANDIANUS S. H. Wright.¹

* *Unio kirklandianus* S. H. WRIGHT, Nautilus, X, 1897, p. 136.

* *Lampsilis kirklandianus* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 76, pl. 1, fig. 7.

Ocklocknee river, Leon County, Florida.

(Group of *Lampsilis ellipsiformis*.)

Shell elliptical, subinflated, pointed behind, that of the female but slightly swollen at the post base; behind this inflation there is a slight sinus; beak sculpture doubly looped, the hinder loop sometimes open behind; epidermis dull, rayed with wavy lines, generally arranged in bands; nacre white.

Animal with the marsupium colored below; gills large, inner wholly united to or only free from the abdominal sack a short distance.

† LAMPSILIS ELLIPSIFORMIS Conrad.

* *Unio ellipsiformis* CONRAD, Monog., VIII, 1836, p. 60, pl. xxxiv, fig. 1.—* B. H. WRIGHT, Check List, 1888.

* *Unio spatulatus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; * Tr. Am. Phil. Soc., X, 1848, p. 80, pl. viii, fig. 22; * Obs., IV, 1848, p. 54, pl. viii, fig. 22; Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* KUSTER, Conch. Cab. Unio, 1861, p. 261, pl. lxxxvii, fig. 7.—* CALKINS, Pr. Ottawa, Ac. Sci., 1874, p. 45.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL Conch. Sam., III, 1890, p. 168.

* *Margarona (Unio) spatulatus* LEA.

* *Lampsilis spatulatus* BAKER, Moll. Chicago, Pt. 1, 1898, p. 106, pl. x, fig. 5; XIII, fig. 2.

Upper Mississippi Valley generally; south to about latitude 38°; western New York; southern Michigan; southern Canada; Red River of the North.

† LAMPSILIS PLEASII Marsh.

* *Unio pleasii* MARSH, the Observer (a newspaper), II, May, 1891; Nautilus, V, 1891, p. 2.

Little Red River, Arkansas; Morrisville, Polk County, Missouri.

† LAMPSILIS ARKANSASENSIS Lea.

* *Unio arkansasensis* LEA, Pr. Ac. N. Sci. Phila., VI, 1862, p. 169; * Jl. Ac. N. Sci. Phila., V, 1862, p. 206, pl. xxx, fig. 275; * Obs., IX, 1863, p. 28, pl. xxx, fig. 275.—

* B. H. WRIGHT, Check List, 1888.

Hot Springs, Arkansas; Saline River near Benton, Arkansas. (Call.)

† LAMPSILIS OZARKENSIS Call.

* *Unio ozarkensis* CALL, Pr. U. S. Nat. Mus., X, 1887, p. 498, pl. xxvii.—* B. H. WRIGHT, Check List, 1888.—* CALL, Tr. Ac. Sci., St. Louis, VII, 1895, p. 33, pl. xviii.

Jack's Ford of Current River, Missouri; White River, Arkansas.

¹ A lovely, brilliant little species, nearly allied to *subangulatus*, but probably distinct.

(Group of *Lampsilis trabalis*.)

Shell solid, that of the male sinuate at post base, and sometimes produced posteriorly, female shell wider, only slightly sinuous; beak sculpture rather coarse, feebly doubly looped; epidermis dark, with wavy, capillary rays; hinge heavy; nacre bluish-white or purple; animal unknown.¹

† *LAMPSILIS TRABALIS* Conrad.

* *Unio trabalis* CONRAD, New F. W. Shells, May 3, 1834, p. 27, pl. III, fig. 5; p. 72.—FERUSSAC, Guer. Mag., 1835, p. 29.—*MOLLER, Syn. Nov. Gen., 1836, p. 201.—*CONRAD, Monog., XII, 1840, p. 110, pl. LX, fig. 2.—*CHENU, Bib. Conch., 1st ser., III, 1845, p. 15, pl. II, fig. 3.—*CONRAD, Pr. Ac. N.Sci. Phila., VI, 1853, p. 259.—*B. H. WRIGHT, Check List, 1888.

† *Unio troostensis* LEA, Tr. Am. Phil. Soc., V, 1834 (August or September), p. 71, pl. X, fig. 30; *Obs., I, 1834, p. 183, pl. X, fig. 30.—*FERUSSAC, Guer. Mag., 1835, p. 29.—*HANLEY, Test. Moll., 1842, p. 186; Biv. Shells, 1843, p. 186, pl. XXIII, fig. 24.—CATLOW and REEVE, Conch. Nom., 1845, p. 64.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1858, p. 496.—*KUSTER Conch. Cab. Unio, 1861, p. 193, pl. LXI, fig. 4.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVIII, fig. 406.—*PETEL, Conch. Sam., III, 1890, p. 170.

**Margarita (Unio) troostensis* LEA, Syn., 1836, p. 21; 1838, p. 18.

**Margaron (Unio) troostensis* LEA, Syn., 1852, p. 25.

**Margaron (Unio) troostii* LEA, Syn., 1870, p. 39.

**Unio troostii* B. H. WRIGHT, Check List, 1888.

**Unio ranuxemensis* SOWERBY, Conch. Icon., 1866, pl. XXXIX, fig. 216.

Kentucky and streams of Tennessee; Clinch River, Virginia.

† *LAMPSILIS PERPURPUREUS* Lea.

**Unio perpurpureus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 41; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 46, pl. XVI, fig. 44; *Obs., XI, 1867, p. 50, pl. XVI, fig. 44.—

*B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) perpurpureus* LEA, Syn., 1870, p. 48.

**Unio troostensis* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIX, fig. 415.

Tennessee River, drainage, extending into Virginia in the Clinch.

(Group of *Lampsilis modioliformis*.)

Shell elliptical, thin, quite evenly rounded before and behind, somewhat inflated; beak sculpture consisting of a few parallel bars, looped in front, and generally open; epidermis with beautiful, rather broad, generally unbroken, sometimes slightly wavy rays; nacre brilliantly iridescent posteriorly; teeth compressed. The female shell is greatly

¹ Certain male shells of *L. trabalis* differ so remarkably from those of the females and from any other known *Naiades* that it might be supposed that they belonged to an isolated group. Other male shells, however, differ but little from the females, and in *L. perpurpureus*, a species so close to *L. trabalis* that it is often difficult to separate them, the male shells differ but slightly from those of the females. Through this former species the group seems very closely related to that of *ellipsiformis*, and so on to the assemblage typified by *L. iris*.

expanded posteriorly, the outline sometimes being almost arcuate on the middle base, and sweeping around in a regular curve to the ligament. Animal with mantle border often spotted, and in the female furnished with fine, well-developed papillæ on the post-ventral region; branchial opening large, with many strong papillæ; marsupium projecting greatly below the rest of the gills, rounded below; ovisacs large and distinct.

† *LAMPASILIS MODIOLIFORMIS* Lea.

* *Unio modioliformis* LEA, Tr. Am. Phil. Soc., V, 1834, p. 97, pl. XIII, fig. 40; * Obs., I, 1834, p. 209, pl. XIII, fig. 40.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 209; * Biv. Shells, 1843, p. 209, pl. XXIII, fig. 37.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* CHENU, Man., 1859, II, p. 139, fig. 678.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 159.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 414, pl. LVI, figs. 2, 3, 6.

* *Margarita (Unio) modioliformis* LEA, Syn., 1836, p. 39; 1838, p. 25.

* *Margaron (Unio) modioliformis* LEA, Syn., 1852, p. 39; 1870, p. 44.

* *Unio tenerus* RAVENEL, Cat., 1834, p. 58.¹—* HANLEY, Test. Moll., 1842, p. 209; * Biv. Shells, 1843, p. 209.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.

* *Margarita (Unio) tenerus* LEA, Syn., 1836, p. 39; 1838, p. 25.

* *Margaron (Unio) tenerus* LEA, Syn., 1852, p. 39; 1870, p. 44.

* *Unio delumbis* KUSTER, Conch. Cab., 1861, p. 207, pl. LXIX, fig. 1.

Santee Canal, South Carolina, south to north Florida; probably west to Mississippi.

† *LAMPASILIS GRACILIOR* Lea.

* *Unio gracilior* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 262; * JI. Ac. N. Sci. Phila., IV, p. 56, pl. VIII, fig. 38; * Obs., VI, p. 56, pl. VIII, fig. 38.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) gracilior* LEA, Syn., 1870, p. 45.

† * *Unio obfuscus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 172; * JI. Ac. N. Sci. Phila., IV, 1859, p. 197, pl. XXII, fig. 80; * Obs., VII, 1859, p. 15, pl. XXII, fig. 80.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) obfuscus* LEA, Syn., 1870, p. 45.

Near Macon, Georgia.

† *LAMPASILIS VIBEX* Conrad.

* *Unio vibex* Conrad, New L. and F. W. Shells, 1834, p. 31, pl. IV, fig. 3, p. 72.²—FERUSSAC, Guerin Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen., 1836, p. 203.—

* HANLEY, Test. Moll., 1842, p. 198; * Biv. Shells, 1843, p. 195.—* CHENU,

¹ I am not aware that any description of Ravenel's species has ever been published. In his catalogue, he gives it this name, and refers to his manuscripts only. Specimens that he gave Dr. Lea, which are credited to him from the Santee canal in pencil on the shells, are undoubtedly *modioliformis*. Other specimens from another locality, which Lea has put with these, and has called *Unio tenerus* Ravenel, are undoubtedly *L. prevostianus* Lea. It was these latter specimens which caused me to be mistaken in the identity of *Unio tenerus* in my paper in Proceedings of the United States National Museum, XV, p. 416.

² One lot in the Academy of Sciences is the same as the *exiguus* of Lea, another is the same as Lea's *nashvillianus*. Conrad's figure, which is not very good, agrees fairly well with the former, but is certainly not the latter.

Bib. Conch., 1st ser., III, 1845, p. 17, pl. III, fig. 6.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* B. H. WRIGHT, Check List, 1888.—* P.ETEL, Conch. Sam., III, 1890, p. 171.

**Margarita (Unio) vibex* LEA, Syn., 1836, p. 27; 1838, p. 20.

**Margaron (Unio) vibex* LEA, Syn., 1852, p. 30; 1870, p. 48.

† **Unio exiguus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 287.—* Tr. Am. Phil. Soc., VIII, 1842, p. 191, pl. VII, fig. 1; * Obs., III, 1842, p. 29, pl. VII, fig. 1.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill., Conch., 1858, pl. XXX, figs. 1, 1a, 1b.¹—* SOWERBY, Conch. Icon., XVI, 1866, p. 38, fig. 208.—* B. H. WRIGHT, Check List, 1888.—* P.ETEL, Conch. Sam., III, 1890, p. 152.

**Margaron (Unio) exiguus* LEA, Syn., 1852, p. 27; 1870, p. 43.

† **Unio stagnalis* CONRAD, Ann. and Mag. N. Hist., IV, 1849, p. 300; * Pr. Ac. N. Sci. Phila., IV, 1849, p. 153; * Jl. Ac. N. Sci. Phila., I, 1850, p. 275, pl. XXXVII, fig. 2; Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* B. H. WRIGHT, Check List, 1888.—* P.ETEL, Conch. Sam., III, 1890, p. 168.

**Margaron (Unio) stagnalis* LEA, Syn., 1852, p. 27; 1870, p. 42.

† **Unio rutilans* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 262; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 59, pl. IX, fig. 41; * Obs., VI, 1858, p. 59, pl. IX, fig. 41.—* KUSTER, Conch. Cab. Unio, 1861, p. 258, pl. LXXXVII, fig. 3.—* B. H. WRIGHT, Check List, 1888.—* P.ETEL, Conch. Sam., III, 1890, p. 166.

**Margaron (Unio) rutilans* LEA, Syn., 1870, p. 45.

† **Unio subellipsidis* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 262; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 62, pl. X, fig. 44; * Obs., VI, 1858, p. 62, pl. X, fig. 44.—* B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) subellipsidis* LEA, Syn., 1870, p. 45.

**Unio prevostianus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIII, fig. 315.

**Unio subangulatus* SOWERBY,³ Conch. Icon., XVI, 1868, pl. LXV, fig. 327.

† LAMPSILIS VIBEX var. NIGRINUS Lea.⁴

**Unio nigrinus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 284, pl. XXIV, fig. 44; * Obs., V, 1852, p. 40, pl. XXIV, fig. 44.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* B. H. WRIGHT, Check List, 1888.—* P.ETEL, Conch. Sam., III, 1890, p. 161.

**Margaron (Unio) nigrinus* LEA, Syn., 1852, p. 39; 1870, p. 62.

† **Unio floridensis* S. H. WRIGHT, Conch. Exchange, II, 1888, p. 105.

† **Unio averillii* B. H. WRIGHT, Pr. Ac. N. Sci. Phila., XXIII, 1888, p. 115, pl. III, fig. 2.—B. H. WRIGHT, Check List, 1888.

Ogeechee River, Georgia, west to Jackson, Mississippi; the variety in Florida, south to the Hillsborough River.

¹ Poor figures, not all accurate.

² This seems to be equivalent to a male *subellipsidis* of Lea, of a little more solid, evenly oval or elliptical form than the type. I do not think it worthy of a varietal name, as there is every possible gradation between these shells. I formerly placed *exiguus*, *rutilans*, and *subellipsidis* in the synonymy of *modioliformis* (Proc. U.S. Nat. Mus., XV, 1892, p. 414), and in a young state they seem to be exactly alike, as a general thing; but the examination of a large amount of additional material since then has led me to believe that the latter may perhaps be distinct.

³ Probably a young specimen, rather wide at the posterior end.

⁴ A smaller, shorter, fragile form, generally having a rather dark epidermis, and purplish nacre. It is a southern variety, and gradually merges into the type in the Chattahoochee region.

† *LAMPASILIS DISPAR* Lea.

* *Unio dispar* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 305; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 327, pl. LI, fig. 153; *Obs., VIII, 1860, p. 9, pl. LI, fig. 153.—*B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) dispar* LEA, Syn., 1870, p. 45.

Southwest Georgia.

(Group of *Lampsilis amygdalum*.)

Shell rather small, obovate, inflated, epidermis varying from smooth and shining to somewhat cloth-like, ashy green to blackish, but always showing green tints when seen through transmitted light, indistinctly rayed; beaks rather high, sculptured with fine, parallel bars, arranged in a double loop, that in front being large and rounded, that behind small and rather pointed below; hinge teeth compressed; nacre iridescent behind. The greatest height of the shell is just behind the center; its greatest diameter is just in front of it, or at a point just behind the beaks; the posterior end is often pointed and somewhat raised. Animal with the marsupium large, reaching far below the inner gills, and having a black border; inner gills united to the abdominal sac throughout; anal opening smooth or only slightly crenulate.

† *LAMPASILIS AMYGDALUM* Lea.

Unio amygdalum LEA, Desc. of 12 sp. Uniones, Aug. 19, 1843; * Tr. Am. Phil. Soc., IX, 1846, p. 275, pl. XXXIX, fig. 1; * Obs., IV, 1848, p. 33, pl. XXXIX, fig. 1.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 244.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 426, pl. LXVII, fig. 3.—* CALL, Pr. Ind. Ac. Sci., 1896, p. 115.

* *Margaron (Unio) amygdalum* LEA, Syn., 1852, p. 39; 1870, p. 62.

† * *Unio lepidus* GOULD,¹ Pr. Bost. Soc. N. Hist., VI, 1856, p. 15; Otia Conch., 1862, p. 222.—B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 426, pl. LXVIII, fig. 1; LXIX, fig. 3.

* *Margaron (Unio) lepidus* LEA, Syn., 1870, p. 39.

Florida.

† *LAMPASILIS SUDUS* Lea.²

* *Unio concavus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 260, pl. xv, fig. 11; * Obs., V, 1852, p. 16, pl. xv, fig. 11.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1868, pl. xcii, fig. 504.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Margaron (Unio) concavus* LEA, Syn., 1852, p. 29; 1870, p. 46.

† * *Unio sudus* LEA, Pr. Ac. N. Sci. Phila., IX, p. 170; * Jl. Ac. N. Sci. Phila., IV, 1859, p. 194, pl. xxi, fig. 77; * Obs., VII, 1895, p. 12, pl. xxi, fig. 77.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) sudus* LEA, Syn., 1870, p. 46.

Abbeville, South Carolina.

¹ Having carefully compared the type of this, a large specimen, with *amygdalum*, I have no doubt that both are the same species.

² The name *concavus* applied by Lea was, I believe, used previously by Zelebor for what seems to be *U. tumidus*.

† LAMPSILIS VESICULARIS Lea.¹

* *Unio vesicularis* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 156; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 37, pl. XII, fig. 34; *Obs., XIII, 1874, p. 41, pl. XII, fig. 34.—* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 425, pl. LXVII, fig. 4.

Florida; Flint River, Georgia?

† LAMPSILIS VILLOSUS B. H. Wright.

* *Unio villosus* B. H. WRIGHT, Naut., XII, 1898, p. 32.

Lampsilis villosus SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 77, pl. I, fig. 1.

Suwanee and Escambia rivers, Florida.

† LAMPSILIS TROSSULUS Lea.²

Unio trossulus LEA, Desc. 12 sp. Uniones, 1843; *Tr. Am. Phil. Soc., IX, 1846, p. 278, pl. XL, fig. 6; *Obs., IV, 1848, p. 36, pl. XL, fig. 6.—* B. H. WRIGHT, Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 170.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 427, pl. LXVIII, fig. 3.

* *Margaron (Unio) trossulus* LEA, Syn., 1852, p. 25; 1870, p. 39.

Lake Monroe, Florida.

† LAMPSILIS PELLUCIDUS Lea.

* *Unio pellucidus* LEA, Pr. Am. Phil. Sci., IV, 1845, p. 163.—* Tr. Am. Phil. Soc., X, 1845, p. 70, pl. II, fig. 6; Obs., IV, 1848, p. 44, pl. II, fig. 6.—CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—B. H. WRIGHT, Check List, 1888.—PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Margaron (Unio) pellucidus* LEA, Syn., 1852, p. 39; 1870, p. 62.

Flint and Chattahoochee rivers, Georgia.

† LAMPSILIS MINOR Lea.

Unio minor LEA, Desc. 12 sp. Uniones, 1843; *Tr. Am. Phil. Soc., IX, 1846, p. 276, pl. XXXIX, fig. 3; *Obs., IV, 1848, p. 34, pl. XXXIX, fig. 3.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 425, pl. LXVII, fig. 2.

* *Margaron (Unio) minor* LEA, Syn., 1852, p. 31; 1870, p. 49.

* *Unio stearnsii* B. H. WRIGHT, Check List, 1888.

Florida and southwestern Georgia.

† LAMPSILIS PAPYRACEUS Gould.

* *Unio papyraceus* GOULD, Pr. Bost. Soc. N. Hist., II, 1845, p. 53.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857,

¹ Dr. Lea has only two specimens of this shell, both in very bad condition—dead and eroded. I have been much inclined to believe that it was a form of *L. amygdalum*, but recently a specimen in better condition submitted for my inspection by Mr. B. H. Wright would seem to show that the two are distinct.

² A doubtful species. The type is the only specimen I have ever seen which I can refer with any certainty to this species. It is a small, rather solid shell, and may be an unusually heavy *L. amygdalum*.

p. 492.—*GOULD, *Otia. Conch.*, 1862, p. 197.—B. H. WRIGHT, Check List, 1888.—
*PÆTEL, *Conch. Sam.*, III, 1890, p. 162.—*SIMPSON, *Pr. U. S. Nat. Mus.*, XV,
1892, p. 427, pl. LXVIII, fig. 2.

Margaron (Unio) papyraceus LEA, *Syn.* 1852, p. 38; 1870, p. 62.

Everglades, Florida.

† *LAMPSILIS SINGLEYANUS* Marsh.¹

**Unio singleyanus* MARSH, *Joliet Weekly News* (a newspaper), May 1, 1891; **Nautilus*, V, 1891, p. 29.—*SIMPSON, *Pr. U. S. Nat. Mus.*, XV, 1892, p. 426, pl. LXVIII, figs. 4, 5.

Putnam and Sumpter counties, Florida.

Subgenus *CARUNCULINA* Simpson in Baker, 1898.²

(Type, *Unio texasensis* Lea.)

Shell small, inflated, obovate, rather solid, covered with a thick, dark, often cloth-like epidermis, which is rayless or only feebly rayed; beak sculpture consisting of rather strong, concentric ridges, which form, as a general thing, only a single, rounded loop in front, and are strongly curved upward behind.³ Pseudocardinals compressed, smooth on the inside, generally reflexed upward, somewhat torn on the edges. Shell quite commonly pointed posteriorly, that of the female truncated obliquely on post base. Animal with the marsupium consisting of a few large ovisacs (8 to 13); inner gills wholly or in part free from the abdominal sac; female often having a well-developed caruncle on the mantle below the branchial opening.

† *LAMPSILIS TEXASENSIS* Lea.

**Unio parvus* CONRAD, *Monog.*, II, 1836, p. 20, pl. IX, fig. 1.—*HANLEY, *Biv. Shells*, 1843, p. 196, pl. XXII, fig. 3.—*SOWERBY, *Conch. Icon.*, XVI, 1866, pl. XXXV, fig. 186.

**Unio texasensis* LEA, *Pr. Ac. N. Sci. Phila.*, VI, 1857, p. 84; **Jl. Ac. N. Sci. Phila.*, IV, 1860, p. 359, pl. LXI, fig. 184; **Obs.*, VIII, 1860, p. 41, pl. LXI, fig. 184.—
*SOWERBY, *Conch. Icon.*, XVI, 1866, pl. XI, fig. 218.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, *Conch. Sam.*, III, 1890, p. 161.—*CALL, *Pr. Ind. Acad. Sci.*, 1896, p. III, pl. v, figs. 38-40.

**Margaron (Unio) texasensis* LEA, *Syn.*, 1870, p. 49.

† **Unio bairdianus* LEA,⁴ *Pr. Ac. N. Sci. Phila.*, I, 1857, p. 102; **Jl. Ac. N. Sci. Phila.*, IV, 1860, p. 361, pl. LXI, fig. 186; **Obs.*, VIII, 1860, p. 43, pl. LXI, fig. 186.—
*B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) bairdianus* LEA, *Syn.*, 1870, p. 49.

¹ This may belong to the *parrus* group. The beaks of specimens I have examined were badly eroded.

² Erroneously spelled *Corunculina*.

³ Occasionally there is a vestige of a posterior loop, and I have seen it perfectly developed, especially in specimens of *L. haleianus*. The beak sculpture of this group is remarkably like that of the *Tetralasmus* group of *Unio*, though the two assemblages are not at all closely related.

⁴ A rather delicate variety, but I hardly think worthy of a varietal name. The species varies from being quite slender to short, wide, and inflated.

† * *Unio bealei* LEA, Pr. Ac. N. Sci. Phila., VI, 1862, p. 169; * Jl. Ac. N. Sci. Phila., V, 1862, p. 204, pl. xxx, fig. 273; * Obs., IX, 1863, p. 26, pl. xxx, fig. 273.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) bealei* LEA, Syn. 1870, p. 49.

† **LAMPASILIS TEXASENSIS** var. **COMPRESSUS** Simpson.¹

Texas, north to Kansas; Missouri; southern Illinois and Indiana; south through Tennessee, Alabama, Mississippi, and Louisiana; the variety in southwest Texas.

† **LAMPASILIS MEARNSI** Simpson.

* *Lampsilis mearnsi* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 75, pl. I, fig. 4.

Vicinity of Fort Clarke, Kinney County, Texas.

† **LAMPASILIS PARVUS** Barnes.

* *Unio parvus* BARNES, Am. Jl. Sci., VI, 1823, pl. XIII, fig. 18 (outline).²—SHORT and EATON, Transylvania Journal, 1831, p. 78.—* CONRAD, New F. W. Shells, 1834, p. 70.—* SAY, Am. Conch., VI, 1834.—FERUSSAC, Guer. Mag., 1835, p. 26.—* HANLEY, Test. Moll., 1842, p. 196.—* PHILIPPI, Abbild, I, 1845, p. 19, pl. I, fig. 4.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* KUSTER, Conch. Cab. Unio, 1852, p. 33, pl. v, fig. 5.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 25.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Margarita (Unio) parvus* LEA, Syn., 1836, p. 28; 1838, p. 21.

* *Margaron (Unio) parvus* LEA, Syn., 1852, p. 31; 1870, p. 49.

* *Lampsilis parvus* BAKER, Moll. Chicago, Pt. 1, 1898, p. 109, pl. XIII, fig. 3.

* *Mya parva* EATON, Zool. Text-Book, 1826, p. 222.

Mississippi River drainage generally; southwest to central Texas; western New York; southern Canada; and Michigan doubtfully.

† **LAMPASILIS HALEIANUS** Lea.³

* *Unio haleianus* LEA, Pr. Am. Phil. Soc., II, 1842, p. 224; * Tr. Am. Phil. Soc., VIII, 1842, p. 247, pl. XXVII, fig. 63; * Obs., III, 1842, p. 85, pl. XXVII, fig. 63.—CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen., Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. XXVI, figs. 6, 6a, 6b.—* ? REEVE, Conch. Icon., XVI, 1865, pl. XXIV, fig. 116.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 154.

* *Margaron (Unio) haleianus* LEA, Syn., 1852, p. 31; 1870, p. 49.

Alexandria, Louisiana; eastern Texas.

¹ A remarkably compressed elevated form from southwest Texas may bear this name.

² Barnes's outline shows a small shell, *evenly rounded posteriorly*. I believe that this character of being rounded behind is constant, and is the only one by which it always differs from its near ally, *L. texasensis*, which is more sharp behind and usually obliquely truncate at the post-base. The two species have been continually confounded, and even Dr. Lea has placed in his collection under the name *parvus* a number of specimens which I am sure are his *texasensis*. The former is generally the smaller of the two, though not always, as in northern Illinois it sometimes reaches a length of 2 inches. *L. texasensis* undoubtedly ranges north into northern Missouri, southern Illinois and Indiana, and the true *parvus* is occasionally found well down into Texas.

³ This is often confounded with *L. texasensis*. It is a larger, thinner-shelled form, and is not so decidedly truncate at the post-basal region.

† *LAMPSILIS GLANS* Lea.

- * *Unio glans* LEA, Tr. Am. Phil. Soc., 1834, p. 82, pl. VIII, fig. 12; * Obs., I, 1834, p. 92, pl. VIII, fig. 12.—* CONRAD, New F. W. Shells, 1834, p. 69.—* FERUSSAC, Guerin Mag., 1835, p. 26.—* CONRAD, Monog., II, 1836, p. 21, pl. IX, fig. 2.—* HANLEY, Test. Moll., 1842, p. 196; Biv. Shells, 1843, p. 196, pl. XXII, fig. 33.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* KUSTER, Conch. Cab. Unio, 1852, p. 37, pl. VI, fig. 3.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—* CHENU, III. Conch., 1858, pl. VIII, figs. 9, 9a, 9b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVI, fig. 190.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 154.
- * *Margarita (Unio) glans* LEA, Syn., 1836, p. 28; 1838, p. 21.
- * *Margaron (Unio) glans* LEA, Syn., 1852, p. 31; 1870, p. 49.

Ohio River drainage; Warsaw, Indiana; (probably St. Lawrence drainage) southern Michigan; White River, Carroll County, Arkansas (Call).

† *LAMPSILIS GERMANUS* Lea.

- * *Unio germanus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 40; * JI. Ac. N. Sci. Phila., VI, 1866, p. 49, pl. XIX, fig. 54; * Obs., XI, 1867, p. 53, pl. XIX, fig. 54.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) germanus* LEA, Syn., 1870, p. 49.
- † * *Unio granulatus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 60; * JI. Ac. N. Sci. Phila., VI, 1866, p. 48, pl. XVI, fig. 46; * Obs., XI, 1867, p. 52, pl. XVI, fig. 46.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) granulatus* LEA, Syn., 1870, p. 49.

Coosa River, and Big Prairie Creek, Alabama.

† *LAMPSILIS CYLINDRELLUS* Lea.

- * *Unio cylindrellus* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 144; * JI. Ac. N. Sci. Phila., VIII, 1869, p. 308, pl. XLVIII, fig. 121.—* Obs., XII, 1869, p. 68, pl. XLVIII, fig. 121.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) cylindrellus* LEA, Syn., 1870, p. 49.

Tennessee River drainage.

† *LAMPSILIS MÆSTUS* Lea.¹

- * *Unio mæstus* LEA, Pr. Am. Phil. Soc., II, 1841, p. 82; * Tr. Am. Phil. Soc., VIII, 1842, p. 244, pl. XXVI, fig. 60; * Obs., III, 1842, p. 82, pl. XXVI, fig. 60.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—* CHENU, III. Conch., 1858, pl. XXVIII, figs. 3, 3a, 3b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Conch. Cab. Unio, 1861, p. 226, pl. LXXVI, fig. 5.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 159.
- * *Margaron (Unio) mæstus* LEA, Syn., 1852, p. 31; 1870, p. 49.

French Broad River, Tennessee.

† *LAMPSILIS CROMWELLII* Lea.

- * *Unio Cromwellii* LEA, Pr. Ac. N. Sci. Phila., IX, 1865, p. 89; JI. Ac. N. Sci. Phila., VI, 1869, p. 258, pl. XXXI, fig. 73; Obs., XII, 1869, p. 19, pl. XXXI, fig. 73.—B. H. WRIGHT, Check List, 1888.

¹The only shells I have seen of this are two in the Lea collection, having quite solid, stumpy teeth. They are old, dead, and a little worn, and in such a condition that it is hard to tell much about them. The species may be a large *cylindrellus*, but I can not be sure that they are.

**Margaron (Unio) cromwellii* LEA, Syn., 1870, p. 49.

†*Unio marginis* LEA, Pr. Ac. N. Sci. Phila., 1865, p. 89; *Jl. Ac. N. Sci. Phila., VI, 1869, p. 225, pl. XXXI, fig. 69; *Obs., XII, 1869, p. 15, pl. XXXI, fig. 69.—* B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) marginis* LEA, Syn., 1870, p. 49.

Western Georgia and southeastern Alabama.

† LAMPSILIS CORVUNCULUS Lea.

**Unio corvunculus* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 144; *Jl. Ac. N. Sci. Phila., VI, 1869, p. 314, pl. I, fig. 127; *Obs., XII, 1869, p. 74, pl. I, fig. 127.—* B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) corvunculus* LEA, Syn., 1870, p. 49.

Western Georgia; Village Creek, Jefferson County, Alabama; Lake Ashby, Florida.

† LAMPSILIS PAULUS Lea.

**Unio paulus* LEA, Pr. Am. Phil. Soc., 1840, p. 287; *Tr. Am. Phil. Soc., VIII, 1842, p. 213, pl. XV, fig. 29; *Obs., III, 1842, p. 51, pl. XV, fig. 29.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. XXVII, figs. 5, 5a, 5b.—* KUSTER, Conch. Cab. Unio, 1861, p. 260, pl. LXXXVII, fig. 6.—* B. H. WRIGHT, Check List, 1888.—* PÄTEL, Conch. Sam., III, 1890, p. 162.

**Margaron (Unio) paulus* LEA, Syn., 1852, p. 31; 1870, p. 49.

†**Unio corvinus* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 144; *Jl. Ac. N. Sci. Phila., VI, 1869, p. 310, pl. XLVIII, fig. 123; *Obs., XII, 1869, p. 70, pl. XLVIII, fig. 123.—* B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) corvinus* LEA, Syn., 1870, p. 49.

Flint and Chattahoochee rivers, Georgia.

LAMPSILIS PULLUS Conrad.¹

**Unio pullus* CONRAD, Monog., XI, 1838, p. 100, pl. LIV, fig. 2; *Proc. Acad. Nat. Sci. Phila., VI, 1853, p. 255.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Conch. Cab. Unio, 1862, p. 270, pl. XCI, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÄTEL, Conch. Sam., III, 1890, p. 164.

**Margaron (Unio) pullus* LEA, Syn., 1852, p. 31; 1870, p. 49.

Warm Springs, North Carolina.

Subgenus PROPTERA Rafinesque, 1819.

(Type, *Unio alatus* Say.)

Shell usually large, gaping at the anterior base and edge of dorsal slope, winged along the dorsal region when young and often when adult; beak sculpture feeble, consisting, when developed, of an anterior and posterior loop, the former often wanting; the latter sometimes becomes slightly nodulous; epidermis generally brown, often cloth-like when fresh, rayless or feebly rayed; teeth rather compressed, pseudocardinals

¹ Conrad's figure is a poor one, and I can not be positive just where the species should be placed, but incline to think it a member of the *Texasensis* group. Lea has a shell which he calls the *pullus* of Conrad, which came from Ravenel, labeled "S. Carolina," but which, I think, is not this, but some species of the *subrostratus* group. It is quite probable that *pullus* is the same as Lea's *corvunculus*.

frequently imperfect or nearly wanting; laterals remote; anterior muscle scars often complicated; dorsal scars consisting of a row of from four to thirty distinct, often deep impressions, running from the cavity of the beak obliquely downward anteriorly; nacre purplish. Animal with large branchiæ; marsupium consisting of numerous, generally fine ovisacs, which are often somewhat radial; mantle thickened and distinctly double on the border, often papillose behind where the outer fold develops into a thickened flap; branchial opening with irregular teeth; anal opening smooth or only slightly crenulate.

(Group of *Lampsilis alatus*.)

Shell obovate, strongly winged, subsolid, with a slight posterior and superposterior ridge; epidermis thick, dark, showing irregular growth lines; hinge teeth generally well developed; nacre dark purple, female shell greatly developed in post-basal region. Animal with very numerous and fine ovisacs; marsupium large; inner gills united to abdominal sac throughout.

† LAMPSILIS ALATUS Say.

Unio alatus SAY, Nich. Encyc., II, 1816, pl. iv, fig. 2.—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 285, fig. 17.—* SAY, Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 67; Monog., VII, 1836, p. 57, pl. XXXI.—* SOWERBY, Conch. Man., 1839, fig. 147.—* ANTON, Verz. der Conch., 1839, p. 14.—* REEVE, Conch. Syst., I, 1841, p. 118, pl. LXXXIX, fig. 8.—* C. B. ADAMS, Thompson's History of Vermont, 1842, p. 166, fig.; * L. and F. W. Shells of Vermont, 1842, p. 16, fig.—* HANLEY, Test. Moll., 1842, p. 174; * Biv. Shells, 1843, p. 174.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 195.—* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 146, pl. LVI, fig. 2.—* CATLOW and REEVE, Conch. Nom., 1845, p. 55.—* KUSTER, Conch. Cab., 1852, p. 15, pl. i, fig. 1.—* DEWEY, Ninth Rep. N. Y. Cab. Nat. Hist., 1856, p. 32.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* CHENU, Man., 1859, II, p. 143, fig. 708.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLIV, fig. 242.—* CALKINS, Pr. Ottawa Ac. N. Sci., 1874, p. 41.—* LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 52.—* B. H. WRIGHT, Check List, 1888.—* PATEL, Conch. Sam., III, 1890, p. 144.

* *Margarita (Unio) alatus* LEA, Syn., 1836, p. 11; 1835, p. 13.

* *Margaron (Unio) alatus* LEA, Syn., 1852, p. 19; 1870, p. 28.

* *Lampsilis alatus* BAKER, Moll. Chicago, Pt. 1, 1898, p. 97, pl. XVIII.

* *Unio alata* LAMARCK, An. sans Vert., VI, 1819, p. 76.—* DESHAYES, Encyc. Meth., II, 1830, p. 583; * An. sans Vert., 2d ed., VI, 1835, p. 539; 3d ed., II, 1839, p. 669.

* *Mya alata*, EATON, Zool. Text-Book, 1826, p. 219.—* WOOD, Index Test. (Rev.), 1856, p. 199, pl. i, supp. fig. 3.

* *Symphynota alata* LEA, Tr. Am. Phil. Soc., III, 1830, p. 448; * Obs., 1834, I, p. 62.—SHORT and EATON, Transylvania Jl., 1831, p. 80.—* FERUSSAC, Gner. Mag., 1835, p. 25.

* *Lymnadia alata* SWAINSON, Treat. on Mal., 1840, p. 265, fig. 48; * Exotic Conch., 2d ed., 1841, p. 38, pl. vii.

* *Mysca alata* SWAINSON, Exotic Conch., 2d ed., 1841, p. 28, pl. vii.

* *Metaptera alata* STIMPSON, Shells of N. Eng., 1851, p. 14.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 260.

*? *Metaptera metaptera* RAFINESQUE, Ann. Gen. Sci., Brux, V, 1820, p. 300, pl. LXXX, figs. 20-22.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 15, pl. i, figs. 20-22.

† *LAMPSILIS ALATUS* var. *POULSONI* Conrad.

* *Unio poulsoni* CONRAD, New F. W. Shells, 1834, p. 25, pl. I, p. 71.—* MÜLLER, Syn. Nov. Gen., 1836, p. 200.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 15, pl. I, fig. 7.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.

* *Symphynota poulsoni* FERUSSAC, Gner. Mag., 1835, p. 25.

* *Unio inflatus* CONRAD, Monog., VII, 1836, p. 57, pl. XXXII.¹

Entire Mississippi drainage as far south as Arkansas; St. Lawrence drainage; Red River of the North; the variety in the Alabama and Tombigbee systems.

† *LAMPSILIS COLORADOENSIS* Lea.

* *Unio coloradoensis* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 103; *Jl. Ac. N. Sci. Phila., III, 1858, p. 314, pl. XXXI, fig. 29; *Obs., VI, 1857, p. 34, pl. XXXI, fig. 29.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) coloradoensis* LEA, Syn., 1870, p. 41.

Eastern Texas.

† *LAMPSILIS ROVIROSAI* Pilsbry.

* *Unio (Lampsilis) rovirosai* PILSBRY, Nantilus, XIII, 1900, p. 140.

Laguna de Atasta, near San Juan Bautista, Mexico.

† *LAMPSILIS PURPURATUS* Lamarck.

* *Unio purpurata* LAMARCK, An. sans Vert., VI, 1819, p. 71.—* STARK, Nat. Hist., II, 1828, p. 90.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 533; 3d ed., II, 1839, p. 667.

* *Unio purpuratus* LEA, Obs., I, 1834, p. 199.—* FERUSSAC, Gner. Mag., 1835, p. 26.—* HANLEY, Test. Moll., 1842, p. 208; *Biv. Shells, 1843, p. 208, pl. XXII, fig. 5.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—* KUSTER, Conch. Cab. Unio, 1861, p. 185, pl. LIX, fig. 1.—* REEVE, Conch. Icon., XVI, 1865, pl. XXIV, fig. 115.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.—* PILSBRY, Nantilus, V, 1891, p. 76.—* CALL, Tr. Ac. Sci. St. Louis, VII, No. 1, 1895, p. 38, pl. v.

* *Margarita (Unio) purpuratus* LEA, 1836, p. 39; 1838, p. 25.

* *Margaron (Unio) purpuratus* LEA, Syn., 1852, p. 38; 1870, p. 61.

† *Unio ater* LEA, Tr. Am. Phil. Soc., III, 1830, p. 426, pl. VII, fig. 9; *Obs., I, 1834, p. 40, pl. VII, fig. 9.—* CONRAD, New F. W. Shells, 1834, p. 67.—* CHENU, Ill. Conch., 1858, pl. x, figs. 1a, 1b.

Unio atra DESHAYES, Enc. Meth., II, 1830, p. 582.

* *Unio lugubris* SAY, Am. Conch., V, 1832, pl. XLIII; VI, 1834.

* *Unio poulsoni* SOWERBY, Conch. Icon., XVI, 1866, pl. LI, fig. 270.

* *Unio dolosus* LEA,² Pr. Ac. N. Sci. Phila., IV, 1860, p. 307; *Jl. Ac. N. Sci. Phila., V, 1862, p. 75, pl. IX, fig. 224; *Obs., VIII, 1862, p. 79, pl. IX, fig. 224.—* Sow-

¹ The variety *poulsoni* is narrower anteriorly and more drawn at the post-basal point than the type, and this form is common in the streams that drain into the Gulf east of the Mississippi, but it seems to connect with typical *alatus* in Tennessee.

² A miserable, worn, young specimen of what I am quite sure is *L. purpuratus*. Lea states that the embryos of this are pouch-shaped, while those of *purpuratus* are wedge-shaped, but the form of embryos in a given species is often not constant.

ERBY, Couch. Icon., XVI, 1866, pl. XLI, fig. 228.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

* *Margaron (Unio) dolosus* LEA, Syn., 1870, p. 61.

Eastern Texas, north to Kansas, through southern Missouri, western Tennessee, to the Alabama River drainage.

† *LAMPSILIS PERMISCENS* Lea.¹

* *Unio permiscens* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 112; *Jl. Ac. N. Sci. Phila., V, 1862, p. 102, pl. XVII, fig. 251; *Obs., VIII, 1862, p. 106, pl. XVII, fig. 251.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) permiscens* LEA, Syn., 1870, p. 61.

Tombigbee River, Columbus, Mississippi.

† *LAMPSILIS GOULDII* Lea.²

* *Unio gouldii* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 165; *Tr. Am. Phil. Soc., X, 1848, p. 76, pl. VI, fig. 16; *Obs., IV, 1848, p. 50, pl. VI, fig. 16.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 153.

* *Margaron (Unio) gouldii* LEA, Syn., 1852, p. 29; 1870, p. 46.

Tuscaloosa, Alabama.

(Group of *Lampsilis saladoensis*.)

Shell smooth, obovate, inflated, inequilateral, rounded behind, thin, slightly thicker in front; beaks somewhat prominent, nearly smooth; epidermis yellowish olive, shining, rayed; there is a slight carina high up on the posterior slope; pseudocardinals small, lamellar, and oblique; laterals lamellar and curved; anterior cicatrices distinct and well impressed; posterior cicatrices confluent; dorsal cicatrices under the plate, between the pseudocardinals and laterals; nacre bluish white and iridescent.³ Animal unknown.

LAMPSILIS SALADOENSIS Lea.

* *Unio saladoensis* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 305; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 370, pl. LXV, fig. 195; *Obs., VIII, 1860, p. 52, pl. LXV, fig. 195.—B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 586.

* *Margaron (Unio) saladoensis* LEA, Syn., 1870, p. 62.

Rio Salado, New Leon, Mexico.

¹ There is only a single, young, poor specimen in the Lea collection, which may be a valid species, but is quite likely a *L. purpuratus*.

² Only a single, dead, badly worn shell is in Lea's collection. It is something like a young *umbrosus*, but if it really came from Tuscaloosa, Alabama, it probably belongs to the *Alatus* group. I can not decide certainly as to its affinities.

³ Mr. Lea states that a single specimen of *U. saladoensis* was collected by Dr. Berlandier and presented to the Smithsonian Institution by Lieutenant Couch (Obs. VIII, p. 53). I have never found this shell, and have not seen the species, and am somewhat at a loss as to where to place it, but the figure and description would indicate that it is a young shell of a form related to *purpuratus*; its rays, which Dr. Lea says are nearest to *U. amygdalum*, being different from those of any related forms.

(Group of *Lampsilis umbrosus*.)

Shell solid, inflated, short-elliptical, covered with a shining, rather smooth brownish epidermis which often has microscopic, radiating folds; beak sculpture excessively faint, being the merest hint at a double loop; anterior end of the shell suddenly thickened, that of the male scarcely differing from the female; hinge teeth well developed; pseudocardinals strong; laterals large, straight, and club-shaped, naere whitish or purplish. Animal with the inner gills united to the abdominal sac; branchiae large; marsupium occupying the whole outer gills posteriorly, not projecting much below; palpi large, considerably united behind.

† *LAMPSILIS UMBROSUS* Lea.

* *Unio umbrosus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 95; *Jl. Ac. N. Sci. Phila., III, 1857, p. 311, pl. xxx, fig. 26; *Obs., VI, 1857, p. 31, pl. xxx, fig. 26.—

*SOWERBY, Conch. Icon., XVI, 1866, pl. xxxiii, fig. 170.—*B. H. WRIGHT, Check List, 1888.—*P. ETEL, Conch. Sam., III, 1890, p. 171.—*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 587, pl. LXVII, figs. 1, 1a, 1b.

*Margaron (*Unio*) *umbrosus* LEA, Syn., 1870, p. 41.

† *Unio veracruzensis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 150; *Jl. Ac. N. Sci. Phila., VI, 1869, p. 320, pl. LI, fig. 133; *Obs., XII, 1869, p. 80, pl. LI, fig. 133.—*B. H. WRIGHT, Check List, 1888.—*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 605.

*Margaron (*Unio*) *veracruzensis* LEA, Syn., 1870, p. 45.¹

Vera Cruz, Mexico.

† *LAMPSILIS BERLANDIERI* Lea.²

* *Unio berlandieri* LEA, Pr. Ac. Phila., I, 1857, p. 101; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 369, pl. LXV, fig. 194; *Obs. VIII, 1860, p. 51, pl. LXV, fig. 194.—

*REEVE, Conch. Icon., XVI, 1865, pl. xxiii, fig. 108.—*B. H. WRIGHT, Check List, 1888.—*P. ETEL, Conch. Sam. III, 1890, p. 116.—*PILSBRY, Naut. V, 1891, p. 76.—*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 584.

*Margaron (*Unio*) *berlandieri* LEA, Syn., 1870, p. 36.

Northeastern Mexico; southwestern Texas.

† *LAMPSILIS TAMPECOENSIS* Lea.

* *Unio tampecoensis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 24, pl. VII, fig. 18; *Obs., II, 1838, p. 24, pl. VII, fig. 18.—*TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 235.—*HANLEY, Test. Moll., 1842, p. 188; *Biv. Shells, 1843, p. 188.—*CATLOW and REEVE, Conch. Nom., 1845, p. 64.—*CONRAD, Pr. Ac. N. Sci. Phila., VI,

¹The type of *veracruzensis* is in the Smithsonian collection, and is undoubtedly a very young *umbrosus*.

²This species is extremely close to *L. tampecoensis*, and at one time I believed the two to be mere forms of the same thing. The present species is typically heavier and more inflated, as well as lighter colored, and has higher beaks than the latter. According to Hon. J. D. Mitchell, of Victoria, Texas, who has carefully studied these forms when living, *berlandieri* is more active, and while spawning it only buries itself sufficiently to be steady and give the post-ventral portion an elevation, while *tampecoensis* is sedentary in its habits, and buries itself entirely, excepting the extreme posterior point. Letter of September 25, 1895.

1853, p. 258.—* CHENU, Ill. Conch., 1858, pl. XXI, figs. 5, 5a, 5b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* ? KUSTER, Conch. Cab. Unio., 1862, p. 275, pl. XCIII, fig. 1.—* ? SOWERBY, Conch. Icon., XVI, 1867, pl. LVII, fig. 291.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 169.—* FISCHER and CROSSE, Miss. Sci. Pt. 7, II, 1894, p. 586.¹

* *Margarita (Unio) tampecoensis* LEA, Syn., 1836, p. 22; 1838, p. 18.

* *Margaron (Unio) tampecoensis* LEA, Syn., 1852, p. 26; 1870, p. 41.

* *Unio heermanni* LEA, Pr. Ac. N. Sci. Phila., XIII, 1861, p. 392; * JI. Ac. N. Sci. Phila., V, 1862, p. 194, pl. XXVI, fig. 263; * Obs. IX, 1863, p. 16, pl. XXVI, fig. 263.—* ? SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIII, fig. 441.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 154.

* *Margaron (Unio) heermanni* LEA, Syn., 1870, p. 28.²

Northeastern Mexico; southeastern Texas; Honduras.

† LAMPSILIS TECOMATENSIS Lea.

* *Unio tecomatensis* LEA, Pr. Am. Phil. Soc., II, 1841, p. 30; * Tr. Am. Phil. Soc., VIII, 1842, p. 234, pl. XXI, fig. 48; * Obs., III, 1842, p. 72, pl. XXI, fig. 48.—CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XXXI, figs. 6, 6a, 6b.—* B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 589, pl. LXV, fig. 6; LXVI, figs. 4, 4a.

* *Margaron (Unio) tecomatensis* LEA, Syn., 1852, p. 27; 1870, p. 50.

Tecomata River, Mexico.

LAMPSILIS LIVIDUS Simpson.³

* *Unio testudineus* REEVE, Conch. Icon., XVI, 1865, pl. XXII, fig. 101.

* *Unio explicatus* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 594.

Rio Usumasinto, Portugal, according to Reeve, who no doubt intended this for Rio Usumasinta, Mexico, and Guatemala.

† LAMPSILIS EXPLICATUS Morelet.

* *Unio explicatus* MORELET, Test. Noviss, Pt. 1, 1849, p. 28.—FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 594, pl. LXI, fig. 1.

* *Margaron (Unio) semigranosus* LEA, Syn., 1870, p. 34.

Central America.

¹Fischer and Crosse (see above citation) doubt whether Sowerby's figures in Conch. Icon. and Knster's in Conch. Cab. are *tampecoensis*. Though not very accurate figures, I can not refer them equally well to any other species.

²Lea has two specimens of *heermanni* in his collection; the type, which is a young, slightly injured shell, and a very young one. I am certain it is only *tampecoensis*.

³Reeve described and figured this form in the Conchologia, supposing it to be the *testudineus* of Morelet, which is a very different thing. Fischer and Crosse in Mission Scientifique, believing it to be equal to Morelet's *explicatus*, placed it in the synonymy of that species. I have never seen an absolutely authentic specimen of *explicatus*, but if Fischer and Crosse have accurately figured and described it, I am sure it is different from Reeve's species, the latter being evidently more inflated, smoother, differently shaped, and having a great deal higher beaks. This being the case, it becomes necessary to give Reeve's shell a new name. Fischer and Crosse credit the species to Sowerby, but the Plate XXII, in the Conchologia, is credited to Reeve.

⁴A single valve of what is probably this species was received by the United States National Museum from Dr. H. von Ihering, under the name of *Unio explicatus* Morelet.

LAMPASILIS ALIENIGENUS Crosse and Fischer.

* *Unio alienigenus* CROSSE and FISCHER, JI. de Conch., XLI, 1893, p. 294.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 590, pl. LXV, figs. 5, 5a; LXVI, fig. 3.

Goatzalcoalcos, Vera Cruz, Mexico.¹

(Group of *Lampasilis aztecorum*.)

Shell elliptical, oblong, rounded in front and behind, the posterior part being slightly compressed, the whole rather thin; umbonal region inflated; beak sculpture unknown; epidermis brown and rather smooth; pseudocardinals small, laterals curved; naere purple and shining; female shell (the specimen figured) somewhat swollen in post basal region, so that the basal line is slightly incurved.

Animal unknown.

LAMPASILIS AZTECORUM Philippi.

* *Unio aztecorum* PHILIPPI, Zeits. für Mal., IV, 1847, p. 95; *Abbild., III, 1849, p. 109, pl. VI, fig. 2.—* KUSTER, Conch. Cab. Unio, 1862, pp. 2, 84, pl. xcv, fig. 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 145.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 607.

* *Margaron (Unio) aztecorum* LEA, Syn., 1870, p. 44.

(Group of *Lampasilis metallicus*.)

Shell rather small, obovate-rhomboid, inflated, much narrowed and rounded in front, the form approaching that of a *Modiolus*; epidermis a rich coppery bronze, shining; beak sculpture consisting of two faint, rounded loops, the bars of which are inclined to be nodulous; pseudocardinals compressed, not well developed, and sometimes more or less broken into denticles; laterals distinct; pits in the beak cavity numerous and very irregular; naere dark, coppery to purple, with metallic tints, iridescent behind. Animal unknown.

† LAMPASILIS METALLICUS Say.

Unio metallicus SAY, N. Harm., Disseminator (newspaper form), January 15, 1831;

Am. Conch., VI, 1834.— CONRAD, New F. W. Shells, 1834, p. 70.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* L. SAY, New Terr. and Fluv. Shells, 1840, p. 6.—

* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—* PÆTEL, Conch. Sam., III, 1890, p. 159.

†* *Unio cuprinus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 94, pl. XII, fig. 24; ²* Obs., I, 1834, pl. XII, fig. 24.—* HANLEY, Test. Moll., 1842, p. 208; * Biv. Shells, 1843, p. 208, pl. XXII, fig. 7.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* CHENU, Ill. Conch., 1858, pl. XI, figs. 3, 3a, 3b.—* H. and A. ADAMS, Gen. Rec. Moll.,

¹I am considerably in doubt as to the relationship of the last two species. Fischer and Crosse are sure that *alienigenus* is close to *umbrosus*, and their figure and description seem to show that *explicatus* is near to *alienigenus*.

²The Transactions of the Philosophical Society, IV, in which Mr. Lea published his *Unio cuprinus*, which is identical with Mr. Say's species, was issued, according to Scudder, in the latter end of 1831.

II, 1857, p. 492.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXVI, fig. 336.—

* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 149.—

* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 615.

* *Margarita (Unio) cuprinus* LEA, Syn., 1836, p. 39; 1838, p. 25.

* *Margaron (Unio) cuprinus* LEA, Syn., 1852, p. 38; 1870, p. 61.

* *Unio acereus* REEVE, Conch. Icon., XVI, 1856, pl. xxx, fig. 160.¹—* PÆTEL, Conch. Sam., III, 1890, p. 144.

Mexico.

(Group of *Lampsilis gracilis*.)

Shell large, thin, elliptical or slightly obovate, with a high posterior and an anterior wing, not greatly inflated; beaks low; epidermis rather smooth, often feebly rayed, dull colored, but usually glossy; hinge line slightly and rather regularly curved; teeth compressed, pseudocardinals but feebly and often imperfectly developed; naere purplish tinted, rather dull. Shell of the male and female nearly alike, the latter scarcely swollen at post basal region. Animal with the mantle greatly thickened at posterior end, and double and thickened at post base, where it is often crenulate or toothed on its inner border, and has the outer developed into a flap; inner gill united to the abdominal sac throughout; marsupium enormous, composed of a great number of delicate semiradiating ovisacs, projecting far below the inner gills in a semicircle.

† *LAMPASILIS GRACILIS* Barnes.

* *Unio gracilis* BARNES, Am. Jl. Sci., VI, 1823, p. 274.—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 288.—* DESHAYES, An. sans Vert, 2d ed., VI, 1835, p. 559; 3d ed., II, 1839, p. 676; Tr. Element. Conch., 1839, p. 18, pl. xxx, figs. 2, 3.—

* HANLEY, Test. Moll., 1842., p. 174.—* C. B. ADAMS, Thompson's Hist. Vermont, 1842, p. 166; * F. W. and L. Shells of Vermont, 1842, p. 16.—* HANLEY, Biv. Shells, 1843, p. 174, pl. xx, fig. 37.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* DESHAYES, Tr. Elem., II, 1853, p. 217, pl. xxx, figs. 2, 3.—* H.

and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* SOWERBY, Conch. Icon., XVI, 1866, pl. xxxix, fig. 215.—* CALKINS, Pr. Ottawa Ac. Sci., 1874, p. 42.—

* LATCHFORD, Tr. Ottawa F. Nat. Club., 1882, p. 52.—* B. H. WRIGHT, Check List, 1888.

* *Mya gracilis* EATON, Zool. Text Book, 1826, p. 222.

* *Symphynota gracilis* LEA, Tr. Am. Phil. Soc., III, 1830, p. 452.—SHORT and EATON, Transylvania Jl., 1831, p. 80.—* LEA, Obs., I, 1834, p. 66.

* *Margarita (Unio) gracilis* LEA, Syn., 1836, p. 11; 1838, p. 13.

* *Metaptera gracilis* STIMPSON, Shells of N. Eng., 1851, p. 14.

* *Margaron (Unio) gracilis* LEA, Syn. 1852, p. 19; 1870, p. 28.

* † *Anodon gracilis* SOWERBY, Genera, No. XVII, 1823, fig.

* *Lampsilis gracilis* BAKER, Moll. Chicago, Pt. 1, 1898, p. 99, pl. XIX, fig. 1.

* *Unio fragilis* SWAINSON, Zool. Ill., 1st ser., III, pl. CLXXI, 1823.²—* DESHAYES, Encyc. Meth., II, 1830, p. 587.—* SAY, Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 69; * Monog., VI, 1836, p. 55, pl. xxx.—* CHENU, Bib.

¹The description of the plate is dated 1856, and in the contents it is credited to Reeve, although the previous plates bear date of 1865, and those following 1866.

²Swainson's *Unio fragilis* was published about the same time as Barnes's *gracilis*, and as it is impossible for me to say which has priority, I agree with Lea that it is better to adopt Barnes's well-known name.

- Conch., 1st ser., III, 1845, p. 12.—*KUSTER, Conch. Cab. Unio, 1852, p. 19, pl. III, fig. 1.
- **Symphynota fragilis* FERUSSAC, Guer. Mag., 1835, p. 25.
- **Metaptera fragilis* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 260.
- **Unio fragilis* var. *gracilis* PÆTEL, Conch. Sam., III, 1890, p. 153.
- **Mya plana* EATON, Zool. Text-Book, 1826, p. 221.
- **Unio planus* CONRAD, New F. W. Shells, 1834, p. 71.
- **Unio* (*Niäa*) *atrata* SWAINSON, Zool. Ill., 1841, pl. CLXXI.
- **Unio atratus* HANLEY, Test. Moll., 1842, p. 199; *Biv. Shells, 1843, p. 199, pl. XXI, fig. 29.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—*PÆTEL, Conch. Sam., III, 1890, p. 145.

Entire Mississippi River drainage, St. Lawrence system; Red River of the North; Hudson River; eastern Texas.

†*LAMPSILIS LÆVISSIMUS* Lea.

- **Symphynota lævissima*, LEA, Tr. Am. Phil. Soc., III, 1830, p. 444, pl. XIII, fig. 23; *Obs., I, 1834, p. 58, pl. XIII, fig. 23.—SHORT and EATON, Transylvania Jl., 1831, p. 80.
- **Unio lævissima*, DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 558; 3d ed., II, 1839, p. 675.
- **Unio lævissimus* CONRAD, New F. W. Shells, 1834, p. 70.—*HANLEY, Test. Moll., 1842, p. 174; *Biv. Shells, 1843, p. 174, pl. XXI, fig. 41.—*CATLOW and REEVE, Conch. Nom., 1845, p. 60.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—*SOWERBY, Conch. Icon., XVI, 1866, pl. XLVI, fig. 250.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 156.
- **Margarita* (*Unio*) *lævissimus* LEA, Syn., 1836, p. 11; 1838, p. 13.
- **Margaron* (*Unio*) *lævissimus* LEA, Syn., 1852, p. 19; 1870, p. 28.
- **Unio ohioensis* SAY, Am. Conch., VI, 1834.—KUSTER, Conch. Cab., 1866, p. 20, pl. LXIX, fig. 5.
- **Symphynota ohioensis* FERUSSAC, Guer. Mag., 1835, p. 25.
- **Metaptera ohioensis* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 260.
- **Unio discoidens* SOWERBY, Conch. Icon., XVI, 1866, pl. LIII, fig. 275.

Mississippi drainage generally; eastern Texas; southern Michigan; western New York.

†*LAMPSILIS ALABAMENSIS* Conrad.

- **Symphynota inflata* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 99, pl. XIV, fig. 28; Obs., I, 1834, p. 109, pl. XIV, fig. 28.—FERUSSAC, Guer. Mag., 1835, p. 25.
- **Metaptera inflata* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 260.
- **Margarita* (*Unio*) *inflatus* LEA, Syn., 1836, p. 11, 1838, p. 13.
- **Unio inflatus* HANLEY, Test. Moll., 1842, p. 174; *Biv. Shells, 1843, p. 174, pl. XXI, fig. 45.—*CATLOW and REEVE, Conch. Nom., 1845, p. 60.—*KUSTER, Conch. Cab. Unio, 1852, p. 17, pl. II, fig. 1.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—*SOWERBY, Conch. Icon., XVI, 1866, pl. XLV, fig. 246.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 155.
- **Margaron* (*Unio*) *inflatus* LEA, Syn., 1862, p. 28; 1870, p. 28.
- **Unio alabamensis* CONRAD, New F. W. Shells, 1834, p. 67.¹

Alabama and Tombigbee River areas.

¹ New name given by Conrad for *Unio inflatus* Lea, preoccupied by Barnes.

(Group of *Lampsilis amphichænus*.)

Shell rather thin, elongate, elliptical, rounded before, pointed behind, compressed, with a smooth, shining, black, rayless epidermis; beaks not prominent, their sculpture unknown; there is a conspicuous gap at the anterior base, and another very distinctly outlined at the upper part of the posterior end like that of *Schizothærus*; hinge line slightly curved; pseudocardinals imperfect; laterals compressed; nacre purplish; female shell inflated at posterior base.

Animal with large, elliptical palpi; mantle slightly thickened at the edge; anal opening widely separated from the superanal, the latter large; foot large, hatchet-shaped. Several animals were examined, but all were so much decayed that most of the characters could not be made out.

† LAMPSILIS AMPHICHÆNUS Frierson.

* *Unio* (*Lampsilis*) *amphichænus* FRIERSON, Nantilis, XI, 1898, p. 10, pl. I.

(Group of *Lampsilis leptodon*.)

Shell rather thin, elongate, elliptical, compressed, decidedly pointed behind, the point raised above the center of the shell; beaks low, their sculpture very feeble; there is a conspicuous post-dorsal wing in young shells, and sometimes vestiges of an anterior wing; young shell rather dull and faintly rayed; there is a decided anterior basal and posterior gap; hinge teeth very imperfect; pseudocardinals often almost wanting, even in young shells; laterals faint, sometimes wanting in the adult shell; nacre coppery to purplish; female shell not inflated at post base.

Animal with large, very thin, light brown branchiæ, free nearly the whole length of the abdominal sac; palpi large, thin, nearly semi-circular; mantle thin, with a wide crenulate border; marsupium projecting slightly below the inner gills.

† LAMPSILIS LEPTODON Rafinesque.¹

* *Unio* (*Leptodea*) *leptodon* RAFINESQUE, Ann. Gen. Sci. Phys. Brux., 1820, p. 295, pl. LXXX, figs. 5-7.

* *Unio leptodon* SAY, Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 70; Monog., VII, 1836, p. 58, pl. XXXIII.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 12, pl. I, figs. 5-7.—* KUSTER, Conch. Cab. Unio, 1861, p. 197, pl. LXIV, figs. 1, 2.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLVIII, fig. 257.—* PÆTEL, Conch. Sam., III, 1890, p. 157.

* *Symphynota leptodon* FERUSSAC, Guer. Mag., 1835, p. 25.

* *Leptodea leptodon* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 262.

* *Anodon purpurascens* SWAINSON, Zool. Ill., 1st ser., III, pl. CLX, 1823.—* CATLOW and REEVE, Conch. Nom., 1845, p. 67.

* *Unio retum* SAY, New Harm. Disseminator, II, September 23, 1829, p. 293; January 15, 1831 (newspaper form).—* FERUSSAC, Guer. Mag., 1835, p. 28.—* L. W. SAY, Terr. & Fluv. Shells, 1840, p. 5.

¹This is one of the few species which its author has described so distinctly that I feel sure there can be no doubt about it. Besides, his figure, such as it is, is something like the *tenuissimus* of Lea.

- * *Symphynota tenuissima* LEA, Tr. Am. Phil. Soc., III, 1829, p. 453, pl. XI, fig. 21; * Obs. 1, 1834, p. 67, pl. XI, fig. 21.
Symphynota tenuissima SHORT and EATON, Transylvania JI., 1831, p. 80.
 * *Margarita (Unio) tenuissimas* LEA, Syn., 1836, p. 38; 1838, p. 25.
 * *Unio tenuissimus* HANLEY, Test. Moll., 1842, p. 206; * Biv. Shells, 1843, p. 206, pl. XX, fig. 42.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLIV, fig. 240.—* B. H. WRIGHT, Check List, 1888.
 * *Margaron (Unio) tenuissimus* LEA, Syn., 1852, p. 38; 1870, p. 61.

Upper Mississippi River drainage, south to the Tennessee River; Buffalo, New York; southern Michigan; Souris River, Manitoba.

(Group of *Lampsilis scutulatus*.)

Shell rhombic elliptical, compressed, rather thin, with a high posterior and a low anterior wing, with two posterior ridges; the valves often exhibiting fine radiating plications on the posterior part; beaks compressed, sometimes showing the nepionic shell, almost destitute of sculpture, which, when present, consists of faint indications of two rounded loops; epidermis rather dull, sometimes indistinctly rayed; left valve with two pseudocardinals, the anterior a sharp, straight ridge sloping obliquely downward and forward from a point just in front of the beaks, the hinder curiously compressed and showing a tendency to break into denticles; right valve with two sharp, ridge-like, sloping pseudocardinals; laterals slender, compressed, straight or wavy; nacre bluish white to pale violet; dorsal scars few; female shell apparently slightly swollen at the posterior base; animal unknown.

LAMPSILIS SCUTULATUS Morelet.

- * *Unio scutulatus* MORELET, Test. Noviss., I, 1849, p. 30.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 561, pl. LIX, fig. 5; LXVII, fig. 6.

Yucatan.

LAMPSILIS PALUDOSUS Morelet.

- * *Unio paludosus* MORELET, Test. Noviss., I, 1849, p. 30.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1849, p. 559, pl. LIX, fig. 3.

Yucatan.

LAMPSILIS PLANIVALVIS Morelet.

- * *Unio planivalvis* MORELET, Test. Noviss., II, 1851, p. 24.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 560, pl. LIX, fig. 2.¹

Usumasinta River, Guatemala.

† LAMPSILIS DELPHINULUS Morelet.

- * *Unio delphinulus* MORELET, Test. Noviss., I, 1849, p. 31.—* HANLEY, Biv. Shells, 1856, p. 381, pl. XXIII, fig. 60.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLIII, fig. 237.—* B. H. WRIGHT, Check List, 1888.—* PLETET, Conch. Sam., III, 1890, p. 150.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 557, pl. LXIII, figs. 2, 2a, 2b.

- * *Margaron (Unio) delphinulus* LEA, Syn. 1870, p. 28.

Guatemala.

¹ These three may be mere variations of one thing.

LAMPSILIS LARGILLIERTI Philippi.

* *Unio largillierti* PHILIPPI, Zeits. für Mal., IV, 1847, p. 94.¹

Yucatan.

The following are unfigured and unknown to me:

* *Lampsilis fasciola* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 299.

* *Lampsilis fulgens* RAFINESQUE, Cont. Monog., 1831, p. 7.

* *Lampsilis argyratus* RAFINESQUE, Cont. Monog., 1831, p. 7.

Genus PSEUDOSPETHA Simpson, 1900.

(Type, *Spatha tanganyicensis* Smith.)²

Burtonia BOURGUIGNAT, Moll. Fluv. Nyanza, 1883, p. 20.³

Shell compressed, thin, oblique, inequilateral, with a straight dorsal line which ends in a small wing in front and behind, rounded and cut away at the anterior base, produced in the post-basal region, gaping in front and behind, slightly twisted on its axis and usually bent posteriorly to the right or left; a low, often double posterior ridge ends in a biangulate point behind; beaks compressed, the sculpture faint, irregular, concentric ridges which continue over the shell, and at the beaks sometimes become rather sharply pustulous, the glochidium often remaining; epidermis yellowish brown, sometimes delicately rayed; teeth rudimentary, one or more faint compressed pseudocardinals and a single lateral in each valve; muscle scars irregular; nacre coppery or purple, often rayed. Animal unknown.

*PSEUDOSPETHA TANGANYICENSIS Smith.

* *Spatha tanganyicensis* SMITH, Proc. Zool. Soc. Lond., 1880, p. 350, pl. XXXI, fig. 8.

* *Burtonia tanganyicensis* PÆTEL, Conch. Sam., III, 1890, p. 187.—VON MARTENS, Beschalte, 1897, p. 257.

* *Burtonia moineï* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 33; Icon. Mal. Tan., 1888, pl. xxv, fig. 1.

* *Burtonia lavigerina* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 36; Icon., Mal. Tan., 1888, pl. xxiv, figs. 1-4.

* *Burtonia magnifica* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 41; Icon. Mal. Tan., 1888, pl. xxvi, figs. 1-2.

Lakes Tanganyika and Nyanza, Africa.

PSEUDOSPETHA LEOPOLDVILLENSIS Putzeys.

Burtonia leopoldvillensis PUTZEYS, Proc. Verb. Soc. Mal. Belg., 1898, pl. xxviii, fig. 16.

Leopoldville, Congo.

¹ Unfigured and unknown to me. From the description I should think likely it belonged here, but it may not.

² The shells of this group bear a striking likeness to that of *Unio tenuissimus* Lea, in form, texture, color, teeth, and in gaping in front and behind, and I am inclined to believe that they are related to it, and the *Unio myersianus* of Southeastern Asia, to *Unio alatus*, and *U. delphinulus* of Central America.

³ Name preoccupied in birds, 1850.

† PSEUDOSPATHA LIVINGSTONENSIS Bourguignat.

- * *Spatha tanganyicensis* SMITH (part), Proc. Zool. Soc. Lond., 1880, pl. xxxi, fig. 8a; 1881, p. 296, pl. xxxiv, fig. 32.
- * *Burtonia livingstonensis* BOURGUIGNAT, Moll. Fluv. Nyanz., 1883, p. 20.
- * *Burtonia elongata* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 31; Icon. Mal. Tan., 1888, pl. xxv, fig. 3.
- * *Burtonia contorta* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 39; Icon. Mal. Tan., 1888, pl. xxvi, figs. 3-5.

Lake Tanganyika.

PSEUDOSPATHA SUBTRIANGULARIS Bourguignat.

- * *Burtonia subtriangularis* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 35; Icon. Mal. Tan., 1888, pl. xxv, fig. 2.
- * *Burtonia grandidieriana* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 42; Icon. Mal. Tan., 1888, pl. xxvii, figs. 1-3.

Lake Tanganyika.

PSEUDOSPATHA BOURGUIGNATI Bourguignat.¹

- * *Burtonia bourguignati* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 38; Icon. Mal. Tan., 1888, pl. xxvii, figs. 4-5.

Lake Tanganyika.

The following species are unfigured and unknown to me.

- * *Burtonia jouberti* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 40.
- * *Burtonia bridouxii*, BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 40.

All from Lake Tanganyika.

Genus HYRIOPSIS Conrad, 1853.

(Type, *Unio delphinus* Gruner.²)

Hyriopsis CONRAD. Pr. Ac. N. Sci. Phila., 1853, p. 269.

Shell large, compressed, rhomboid-elliptical, dorsally winged and generally biangulate behind, sometimes produced in the post-basal region; beaks low, mostly compressed, the sculpture consisting of numerous concentric ridges nearly parallel with the growth lines, generally extending well on to the disk of the shell, the earlier ones fine, often slightly nodulous or doubly looped; epidermis olive to brown, sometimes faintly rayed; pseudocardinals two or three in the left valve, one to three in the right, when young compressed, but often breaking into irregular denticles when old; laterals long, compressed, two in the left valve and one in the right, sometimes vertically striate, the left valve often having a sort of raised lamellar tooth at or just behind the beak.

¹ Said to be of Joubert in the literature.

² The general make-up of the shells of this genus, *Cyclonys*, and *Cristaria* seems to indicate close relationship to each other, and a not distant one to the subgenus *Proptera* of *Lampsilis*, and I should not be surprised to learn that the marsupium occupies the posterior part of the outer gills in distinctly marked ovisacs.

Beak cavities shallow; dorsal scars numerous, extending from the cavity downward and forward. Anterior muscle scars three, distinct, posterior large, indistinct; front of the shell usually thickened; a row of plications often occurring just below the laterals, as in *Cristaria*.

Animal unknown.

(Group of *Hyriopsis bialatus*.)

Shell somewhat trapezoidal, with a posterior ridge, biangulate behind, with a very high, pointed, triangular, posterior wing, and a smaller, pointed one in front, which projects forward; beaks low, sculpture not seen; epidermis brown; pseudocardinals broken up into denticles; hinge line slightly curved; the three anterior muscle scars separate and deep; dorsal scars not extending greatly downward in front; nacre bluish white.

†HYRIOPSIS BIALATUS Simpson.¹

**Unio delphinus* GRUNER Arch. für Naturg., I, 1841, p. 276, pl. IX, figs. 1, 1a-c.—

* DELESSERT, Rec. Coq. Lam., 1841, pl. XIX, fig. 3, 3a, 3b.—* LEA, Tr. Am. Phil. Soc., VIII, 1843, p. 218, pl. XVII, fig. 35; * Obs., III, 1842, p. 56, pl. XVII, fig. 35.²—* KUSTER, Conch. Cab. Unio, 1852, p. 18, pl. II, figs. 2-4.—* HANLEY, Biv. Shells, 1856, p. 381, pl. XX, fig. 44.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* CHENU, Man., 1859, II, p. 143, fig. 707.—SOWERBY, Conch. Icon., XVI, 1866, pl. XLIII, fig. 238.—* MORELET, Series Conch., IV, 1875, p. 345.—* PÆTEL, Conch. Sam., III, 1890, p. 150.

**Margaron (Unio) delphinus* LEA, Syn. 1852, p. 19; 1870, p. 28.

**Unio megapterus* MORELET, J. de Conch., XII, 1863, p. 159.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

Southeastern Asia, including Cambodia, Siam, and the Malay Peninsula.

(Group of *Hyriopsis cumingii*.)

Shell rather thin, with a high wing posteriorly, and a smaller one in front; beaks greatly compressed, their sculpture at first consisting of fine and later of coarse concentric ridges, extending well on to the disk; corrugations on the posterior slope distinct.

HYRIOPSIS CUMINGII Lea.

**Unio cumingii* LEA, Pr. Ac., N. Sci. Phila., VII, 1852, p. 54.—* LEA, Jl. Ac. N. Sci. Phila., IV, 1860, p. 240, pl. XXXV, fig. 120; * Obs., VII, 1860, p. 58, pl. XXXV, fig. 120.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—

* SOWERBY, Conch. Icon., XVI, 1866, pl. XLIX, fig. 264.—* HEUDE, Conch. Flav. Nank., VIII, pl. LXIV.

**Margaron (Unio) cumingii* LEA, Syn., 1852, p. 19; 1870, p. 28.

China.

¹I regret that Gruner's well-known name *delphinus* was previously used for a *Unio* by Spengler, and the term *megapterus* of Morelet was applied by Rafinesque to a *Metaptera (Unio)* and used by Chenu under the generic name *Unio* before 1863. It will therefore have to have a new name.

²Lea described this species under the above name, not knowing that it had been already applied to the same species by Gruner.

†HYRIOPSIS DELAPORTEI Crosse and Fischer.

* *Unio (Arconaia) delaportei* CROSSE and FISCHER, J. de Conch., XXVI, 1876, p. 327, pl. X, fig. 1; XI, fig. 5.¹

Cambodia; Siam.

(Group of *Hyriopsis myersianus*.)

Shell rather solid, thickened in front; posterior ridge rather low; beaks compressed, the sculpture consisting, apparently, of a few rather faint, irregular ridges, sometimes slightly nodulous.

†HYRIOPSIS MYERSIANUS Lea.

* *Unio myersianus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 92; *Jl. Ac. N. Sci. Phila., III, 1857, p. 290, pl. XXII, fig. 2; *Obs., VI, 1857, p. 10, pl. XXII, fig. 2.—*SOWERBY, Conch. Icon., XVI, 1866, pl. L, fig. 265.—*MORELET, Ser. Conch., IV, 1875, p. 344.—*PÆTEL, Conch. Sam., III, 1890, p. 160.

*Margaron (*Unio*) *myersianus* LEA, Syn., 1870, p. 28.

*† *Unio housei* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 92; *Jl. Ac. N. Sci. Phila., III, 1858, p. 291, pl. XXIII, fig. 3; *Obs., VI, 1857, p. 11, pl. XXIII, fig. 3.—*VON MARTENS, Pr. Zool. Soc. Lond., 1860, p. 14.—*SOWERBY, Conch. Icon., XVI, 1866, pl. XLVIII, fig. 260.—*PÆTEL, Conch. Sam., III, 1890, p. 155.

*Margaron (*Unio*) *housei* LEA, Syn., 1870, p. 28.

Siam; Cambodia.

HYRIOPSIS SUTRANGENSIS L. Morlet.

* *Unio sutrangensis* L. MORLET, Jl. de Conch., XXXVII, 1889, p. 195, pl. IX, fig. 3.

Sutrang River, Siam.

HYRIOPSIS PINCHONIANUS Heude.

* *Unio pinchonianus* HEUDE, Conch. Fluv. Nank, VIII, 1883, pl. LXIII, fig. 12.—PÆTEL, Conch. Sam., III, 1890, p. 163.

China.

(Group of *Hyriopsis vagulus*.)

Shell elliptical, with a rather high, posterior wing, moderately solid, with two slight posterior ridges and a sulcus above them at the juncture of the wing.

HYRIOPSIS VAGULUS Fischer.²

* *Unio subtrigonus* SOWERBY, Conch. Icon., XVI, 1867, pl. LVIII, fig. 292.—*PÆTEL, Conch. Sam., III, 1890, p. 169

*Margaron (*Unio*) *subtrigonus* LEA, Syn., 1870, p. 28.

¹The ventral line of this shell is crooked, having a couple of curves in it, but it is not twisted. Its authors believed *Arconaia* to be only a section of *Unio*, and that the animal was essentially like that of that genus.

²Sowerby published this species as *Unio subtrigonus* in the Conchologia, the name being preoccupied by Deshayes. In the errata he changed it to *Unio (Monocondylæa) cambojensis*, believing it to be the same as Petit's *Pseudodon cambojensis*. It certainly is not that shell, and I agree with Fischer that it is near to the *Unio myersianus*, and accept his name for it.

* *Unio (Monocondylaea) cambojensis* SOWERBY, Conch. Icon., XVI, errata.

* *Pseudodon cambojensis* MORELET, Ser. Conch., IV, 1875, p. 336.

* *Unio lamellatus* HANLEY and THEOBALD, Conch. Ind., 1876, p. 5, pl. IX, fig. 6.

* *Unio ragulus* FISCHER, Bull. Soc. Autun., IV, 1891, p. 223.

Siam.

(Group of *Hyriopsis velthuzeni*.)

Shell somewhat inflated, rather thin, rounded behind, winged; epidermis dark brown with numerous green rays; nacre iridescent, bluish white; anterior scars irregular, rather small; posterior scars rounded; pseudocardinals elongated, one in each valve, in the right valve there is a secondary rudimental one; laterals elongate, arcuate, one in the right valve and two in the left.

† **HYRIOPSIS VELTHUZENI** Schepman.

* *Unio velthuzeni* SCHEPMAN, Notes from Leyden Mus., XVII, 1895, p. 160, pl. IV, figs. 1, 1a—¹ DROUET, Jl. de Conch., XLV, 1897, p. 124.

Mandai River, Borneo.

(Group of *Hyriopsis schlegeli*.)

Shell rather thin, slightly winged behind, narrowly rounded before and angled at umbonial slope; posterior ridge high and rounded, beaks rather low; epidermis rough, blackish; hinge line curved in front, straight behind; pseudocardinals generally entire; plications below the laterals faint; nacre dull, lurid purplish or lead color.

† **HYRIOPSIS SCHLEGELI** von Martens.

* *Unio schlegeli* VON MARTENS, Mal. Blatt., VII, 1861, p. 55.—* KOBELT, Abh. Senck, Nat. Ges., XI, 1879, p. 421, pl. XIV.—* VON MARTENS, Nov. Conch., V, 1879, p. 192, pl. CLVIII, figs. 4, 5.—* VON IHERING, Abh. Senck, Nat. Ges., XVIII, 1893, p. 156.

* *Barbala schlegeli* PÆTEL, Conch. Sam., III, 1890, p. 175.

Japan.

Subgenus **CAUDICULATUS** Simpson, 1900.

(Type, *Unio caudiculatus* von Martens.)

Shell oval, inflated, slightly posteriorly winged, with a well-developed posterior ridge, and above it a deep, wide furrow ending in a broad, well-defined notch on the posterior slope; one pseudocardinal in the right valve and two in the left; two laterals in the left valve and one in the right. Animal unknown.²

HYRIOPSIS CAUDICULATUS Von Martens.

* *Unio caudiculatus* VON MARTENS, Mal. Bl., XIV, 1866, p. 16.—* SCHEPMAN, Notes Leyd. Mus., XVIII, 1896, p. 140.—* DROUET, J. de Conch., XLV, 1897, p. 126.

* *Unio infrarostratus* SCHEPMAN, Notes Leyd. Mus., XVII, 1895, p. 161, pl. IV, fig. 2.

Borneo.

¹According to the photographic figure this shell very closely resembles *Unio alatus* Say.

²I am in doubt as to the relationship of this species.

Genus CHAMBERLAINIA Simpson, 1900.

(Type, *Unio hainesianus* Lea.)

Shell large, very solid in front, thinner behind, round or round obovate, without posterior ridge, alate on the post dorsal part when young; beaks full, sculpture not seen; surface without sculpture excepting a few slightly pustulous corrugations on the posterior slope; epidermis dull, not rayed; hinge curved; pseudocardinals rather small, stumpy, two in the left valve and one in the right; laterals short, remote, the intervening hinge plate rounded, one lateral in the right valve and two in the left, the upper the smaller, and all curved upward along their inner edges; beak cavities not deep; dorsal muscle scars numerous, deep, extending from the cavity downward and forward; area outside the pallial line very wide; nacre lurid bronzy. Animal unknown.

†CHAMBERLAINIA HAINESIANA Lea¹.

* *Unio hainesianus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 92; * JI. Ac. N. Sci. Phil., III, 1857, p. 289, pl. XXI, fig. 1; * Obs. VI, 1857, p. 9, pl. XXI, fig. 1.—

* VON MARTENS, Pr. Zool. Soc. Lond., 1860, Pt. 1, p. 15.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLVII, fig. 251.²—* MORELET, Ser. Conch., IV, 1875, p. 312.—* PÆTEL, Conch. Sam., III, 1890, p. 154.

* Margaron (*Unio*) *hainesianus* LEA, Syn., 1870, p. 28.

* *Unio imperialis* MORELET, Rev. et Mag. Zool., XIV, 1862, p. 480.

Siam.

CHAMBERLAINIA PAVIEI Morelet.

* *Unio paviei* MORELET, J. de Conch., XXXIX, 1891, p. 241, pl. VII, fig. 3.³

Siam.

†CHAMBERLAINIA DUCLERCI Rochebrune.⁴

* *Unio duclerci* ROCHEBRUNE, Bnll. Soc. Phil., VII, 1882, p. 27, pl. I, fig. 2 (in extract).

Mekong River, Siam.

¹ The type consists of a single valve of a young individual. *U. imperialis* according to its author attains a size of 142 mm. in height, 192 mm. in length, and 77 mm. in diameter. I have seen a very fine specimen in the Frederick Stearns collection 6 inches high and 8 in length. It is one of the most magnificent Naiades in the world.

² Credited by Sowerby to the Little Arkansas River.

³ Very close to *C. housei*, but more rounded, and having a higher wing. I do not think this can be the same as Morelet's *Unio pairanus*, described in Latin in the Journal in 1865, p. 227, and unidentifiable.

⁴ It is quite probable that this should form the type of a separate group, but the only specimens I have seen, two badly eroded valves, hardly justify me in separating it.

Genus CRISTARIA Schumacher, 1817.

(Type, *Dipsas plicatus* Leach.)*Barbala* HUMPHREYS, Mus. Col., 1797.*Dipsas* LEACH, Zool. Miscellany I, 1814, p. 119.*Cristaria* SCHUMACHER, Essai Nouv. Syst., 1817, p. 107.*Appius* (LEACH), Menke. Syn., 2d ed., 1828, ? p. 106.*Symphynota* LEA (part), Tr. Am. Phil. Soc., 1830, p. 445. —SWAINSON, Mal., 1840, p. 288.*Dianisotis* RAFINESQUE, Cont. Monog., 1831, p. 7.*Dipsas* VOIGT, Cuvier, Thierreich, III, 1834, p. 493.*Barbata*, SOWERBY, Conch. Man., 1842, p. 81.*Clione* GISTEL, Nat. Sur. Hoh. Schule Bearb., 1848.

Shell generally thin, elliptical, more or less winged posteriorly; beaks rather low, sculptured with fine, somewhat doubly-looped ridges at first, and later with coarse, low, concentric bars, which are nearly parallel with the growth lines; epidermis smooth, often somewhat rayed; hinge teeth imperfect; pseudocardinals feeble or wanting—when present consisting of a single compressed tooth in each valve; laterals single, remote, compressed, sometimes wanting in adult shells.¹

Subgenus CRISTARIA, Schumacher, 1817.

(Type, *Dipsas plicatus* Leach.)

Shell large, thicker in front, strongly alate posteriorly, with two or more slightly developed radial ridges on the posterior slope and a row of plications above them; hinge line regularly curved; dorsal scars forming a row running obliquely downward and forward; anterior scars often blurred, posterior faint.

†CRISTARIA PLICATA Leach.

* *Dipsas plicatus* LEACH, Zool. Miscellany, I, 1815, p. 120, pl. LIII.²—* SOWERBY, Conch. Man., 1839, fig. 142.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 15, pl. VI, fig. 1.—* MORELET, Ser. Conch., IV, 1875, p. 332.—* HEUDE, Conch. Fluv. Nank., 1879, pls. XXXIII, XXXIV, LI, fig. 94.

Mytilus plicatus SOLANDER, in Gray, Ann. of Phil., IX, 1825, p. 27.*Appius plicatus* GRAY, in Menke, Syn. Meth. Moll., 1830, p. 106.* *Margarita* (*Dipsas*) *plicatus* LEA, Syn., 1836, p. 47; 1838, p. 28.* *Margaron* (*Dipsas*) *plicatus* LEA, Syn., 1852, p. 46; 1870, p. 74.* *Unio plicatus* SOWERBY, Conch. Icon., XVII, 1868, pl. LIV., fig. 280.* *Dipsas plicata* TROSCHEL, Arch. für Naturg., XIII, 1874, p. 272.—* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 429, pls. XV–XVII, XVIII, fig. 1.* *Barbala plicata* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 501, pl. CXVII, figs. 4, 4a.* *Anodonta plicata* SCHRENCK, Reis. und F. Am. Lande, II, 1867, p. 704, pl. XXVII, fig. 4.

¹ The animal has been described in Japanese, but the paper containing the description is inaccessible to me.

² This seems to be the first authentic description of this species. It has been claimed that the *Mytilus dubius* of Gmelin is this, but he refers to figures 733, pl. 82, in the eighth volume of Chemnitz, which, I am quite certain, is not the species in question.

- * *Anodonta (Dipsas) plicata* CLESSIN, Conch. Cab. Anodonta, 1876, p. 240, pl. XXI, figs. 1, 2.
- * *Cristaria plicata* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 147.
- * *Cristaria tuberculata* SCHUMACHER, Ess. Nouv. Syst., 1817, p. 140, pl. xx, fig. 2.
- * *Anodonta dipsas* BLAINVILLE, Man. Mal., 1825, p. 538, pl. LXVI, fig. 2.
- * *Symphynota bialata* LEA, Tr. Am. Phil. Soc., 1830, p. 445, pl. xiv, fig. 24; * Obs., I, 1831, p. 59, pl. xiv, fig. 24.
- * *Unio bialata* HANLEY, Test. Moll. 1842, p. 219; * Biv. Shells, 1843, p. 214, pl. xxii, fig. 4.
- * *Barbala bialata* CHENU, Man. 1859, p. 145, fig. 717.—* PÆTEL, Conch. Sam., III, 1890, p. 175.
- * *Unio bialatus* DESHAYES, Tr. Elem. Conch., 1839, p. 19, pl. xxxi, fig. 3.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* DESHAYES, Tr. Elem., II, 1853, p. 218, pl. xxxi, fig. 3.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XLVI, fig. 247.
- * *Anodonta magnifica* CLESSIN, Conch. Cab. Anodonta, 1873, p. 123, pl. xxxv, fig. 1.
- * *Dipsas occidentalis* HEUDE, Conch. Fluv. Nank., IX, 1885, pl. XLVI.
- * *Barbala occidentalis* PÆTEL, Conch. Sam., III, 1890, p. 175.
- * *Barbala plicatula* PÆTEL, Conch. Sam., III, 1890, p. 175.

China; Japan; Amurland, south probably to Cambodia. The form from Cambodia, which is referred to *C. plicata*, is believed by Fischer to be different, and it may be *C. bellua*.

† CRISTARIA BELLUA Morelet.

- * *Anodonta bellua* MORELET, Rev. et Mag. Zool., XVIII, 1866, p. 167; Ser. Conch., IV, 1875, p. 331.

Cambodia.

† CRISTARIA HERCULEA Middendorff.

- Anodonta herculea* MIDDENDORFF, Bull. Phys. Math. Ac. St. Pet., VI, 1848, p. 303; Sib. Reise, II, 1851, p. 278, pl. xxi, fig. 5; xxii, figs. 1, 2; xxvi, figs. 1, 2.¹—* DESHAYES, Bull. Nouv. Arch. Mus., IX, 1873, p. 1, pl. i, fig. 1.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 294.
- * *Anodonta (Dipsas) herculea* CLESSIN, Conch. Cab. Ano., 1875, p. 175, pl. LIX, figs. 1, 2.
- * *Cristaria herculea* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 146.
- * *Barbala herculea* PÆTEL, Conch. Sam., III, 1890, p. 175.
- * *Dipsas plicata* var. *classini* KOBELT, part. Abh. Senck. Nat. Ges., XI, 1879, p. 429, pl. xv.²

East Siberia; Amurland; North China.?

† CRISTARIA SPATIOSA Clessin.

- * *Anodon herculeus* SOWERBY, Conch. Icon., XVII, 1867, pl. III, fig. 7.
- * *Anodonta (Dipsas) spatiosa* CLESSIN, Conch. Cab. Anodonta, 1875, p. 173, pl. LVII, fig. 2.

¹ Plate xxvi, fig. 2, an interior view, is probably *Anodonta beringiana* Middendorff.

² *Craspedodonta smaragdina* ANTON, Verz. der Conch., 1839, p. 16, No. 592; Clessin, Conch. Cab. Ano., 1875, p. 93, pl. xxvii, fig. 2, a badly figured little shell, said to come from North America, may be a young *C. herculea*.

**Cristaria spatiosa* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 145.

**Barbala spatiosa* PÆTEL, Conch. Sam., III, 1890, p. 175.

**Dipsas plicata* var. *japonica* KOBETT, Abh. Senck. Nat. Ges., XI, 1879, p. 429, pl. XLVII.

Japan.

Subgenus **PLETHOLOPHUS** Simpson, 1900.

(Type, *Symphynota discoidea* Lea.)

Shell short-elliptical, lenticular, scarcely thickened in front, with compressed beaks which are sculptured with low, wide, concentric ridges, scarcely winged in front or behind, pointed posteriorly; epidermis smooth, somewhat rayed; hinge teeth very feeble, often nearly wanting; dorsal scars few; muscle scars faint. Animal unknown.

† CRISTARIA DISCOIDEA Lea.

**Symphynota discoidea* LEA, Tr. Am. Phil. Soc., V, 1834, p. 75, pl. XI, fig. 33; * Obs., I, 1834, p. 187, pl. XI, fig. 33.

**Barbala discoidea* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 501.—* CHENU, Man., 1859, II, p. 145, fig. 716.—* PÆTEL, Conch. Sam., 1890, p. 175.

**Dipsas discoidea* KUSTER, Conch. Cab. Unio, 1862, p. 306, pl. c, fig. 1.

**Margarita (Unio) discoideus* LEA, Syn., 1836, p. 47; 1838, p. 28.

**Unio discoideus* HANLEY, Test. Moll., 1842, p. 214; * Biv. Shells, 1843, p. 214, pl. XXI, fig. 57.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.

**Margarita (Unio) discoideus* LEA, Syn., 1852, p. 46; 1870, p. 74.

**Unio tenuis* GRAY, Griffith's Cuv., XII, 1834, p. 601 (index), pl. XXIV, fig. 2.¹

**Anodonta chinensis* KUSTER, Conch. Cab. Ano., 1853, p. 51, pl. XII, fig. 3.

**Unio magnificus* SOWERBY, Conch. Icon., XVI, 1867, pl. LVII, fig. 289.

**Anodonta gruneriana* CLESSIN, Conch. Cab. Ano., 1875, p. 172, pl. LVI, figs. 3, 4.

China.

CRISTARIA PARVULA Heude.²

**Dipeas parvulus* HEUDE (part), Conch. Fluv. Nank., IV, 1878, pl. XXXII, fig. 65.

**Barbala parvula* PÆTEL, Conch. Sam., III, 1890, p. 175.

China.

CRISTARIA REINIANA von Martens.

**Cristaria reiniana* VON MARTENS, Jahrb. Mal. Ges., II, 1875, p. 136, pl. III, fig. 4.

* ? *Dipsas reiniana* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 432, pl. XII, fig. 4; XXI, fig. 2.

**Anodonta reiniana* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 147.³

**Barbata reiniana* PÆTEL, Conch. Sam., III, 1890, p. 175.

Japan.

¹ A brief description is given in the index only. Gray's and Lea's names seem to have been published the same year, and I do not know which appeared first. I use Lea's because it is the one generally known.

² I am doubtful whether this is more than a variety of *C. discoidea*. Heude's figure on plate LIII is a very different thing, which I have called *C. radiata*.

³ Martens says it has distinct cardinals and laterals.

CRISTARIA GENTILIANA Heude.

* *Cristaria gentilianus* HEUDE, Conch. Fluv. Nank., VII, 1881, pl. LIII, fig. 98.

* *Barbala gentiliana* PÆTEL, Conch. Sam., III, 1890, p. 175.

China.

† CRISTARIA SWINHOEI H. Adams.

* *Unio swinhoei* H. ADAMS, Proc. Zool. Soc. Lond., 1866, p. 319.¹—SOWERBY, Conch. Icon., XVI, 1868, pl. XLII, fig. 232.—PÆTEL, Conch. Sam., III, 1890, p. 169.

* *Anodonta swinhoei* H. ADAMS, Proc. Zool. Soc. Lond., 1866, p. 446.—* CLESSIN, Conch. Cab. Anodonta, 1876, p. 125, pl. LXXV, fig. 6.—* PÆTEL, Conch. Sam., III, 1890, p. 185.

* *Anodon swinhoei* SOWERBY, Conch. Icon., XVI, 1868, pl. XXVII, fig. 108.

* *Margaron (Unio) swinhoei* LEA, Syn., 1870, p. 45.

Formosa; Tonkin; Cambodia.

Subgenus CRASSITESTA Simpson, 1900.

(Type, *Cristaria radiata* Simpson.)

Shell rather solid, elliptic-rhomboid, with a rounded posterior ridge, scarcely winged; beaks full, sculpture not seen; epidermis bright, strongly rayed, with two conspicuous light and three dark rays on the posterior slope; hinge line curved; pseudocardinals and laterals reduced to the merest vestiges; dorsal scars only one or two in the cavity of the beaks; adductor scars faint; naere lurid.

† CRISTARIA RADIATA Simpson.²

* *Dipsas parvulus* HEUDE (part), Conch. Fluv. Nank., VII, 1881, pl. LII, fig. 96.

China.

Genus LEPIDODESMA Simpson, 1896.

(Type, *Unio languilati* Heude.)

Lepidodesma SIMPSON, Proc. U. S. Nat. Mus., 1896, p. 311.

Shell large, thin, inflated, with a high, sharp, posterior ridge and a second fainter one above, making the hinder part widely biangulate; beaks very high and full, their sculpture, which extends over the whole shell, consisting of ridges that follow the growth lines; there are two rows of radiating nodules, one on the middle of the disk, and a stronger one on the post ridge; ligament very large, wide, covered with concentric scales; hinge line arched; two pseudocardinals in the left valve, the anterior elongated, running inwardly across the hinge plate, and ending abruptly, with a shorter, fainter one behind it; with two strong laterals, the inner the higher, its edge reflexed upward, and suddenly truncate behind; one low elongated pseudocardinal in the right valve,

¹ This seems to be a *Cristaria* of the *discoidea* section. The cardinals are somewhat better developed than is common among *Cristarias*, but are still quite faint.

² Heude has figured two different species under the name of *Dipsas parvulus*, on plate XXXII. I have therefore given the last-described form the name *radiata*.

and one lateral, which is reflexed upward; epidermis folded into the hinge, and scaly; a single dorsal scar on the inner side of each hinge plate; muscle scars united; pallial line distinct; beak cavities large and wide; nacre very dull. Animal unknown.

† **LEPIDODISMA LANGUILATI** Heude.

* *Unio languilati* HEUDE, J. de Conch., XXII, 1874, p. 116; * Conch. Fluv. Nank., I, 1875, pl. VII.—PÆTEL, Conch. Sam., III, 1890, p. 156.

* *Lepidodesma languilati* SIMPSON, Pr. U. S. Nat. Mus., XVIII, 1896, p. 311.

* *Cristaria megadesma* VON MARTENS, S. B. Nat. Fr., 1875, p. 3; * Mal. Bl., XXII, 1875, p. 187; * Nov. Conch., IV, 1876, p. 152, pl. CXXXV, fig. 1.

* *Barbula megadesma* PÆTEL, Conch. Sam., III, 1890, p. 175.

China.

LEPIDODESMA ALIGERA Heude.

* *Unio languilatus* var. *aligerus* HEUDE, Conch. Fluv. Nank., III, 1877, pl. XVII, fig. 37.—* PÆTEL, Conch. Sam., III, 1890, p. 156.

* *Lepidodesma aligera* SIMPSON, Pr. U. S. Nat. Mus., XVIII, 1896, p. 311.

China.

Genus **PILSBRYOCONCHA** Simpson, 1900.

(Type, *Anodonta exilis* Lea.)

Shell elongated, elliptical, compressed, thin, with a low, posterior wing, narrow and rounded in front, nearly straight below, expanded at the post-basal region, pointed behind. Beaks compressed, sculptured with several coarse, low, irregularly concentric undulations, often slightly doubly looped; surface nearly smooth, shining, yellowish to ashy brownish, sometimes showing very faint rays; hinge straight or slightly curved, reduced to a mere line, often showing a faint, compressed, smooth tooth in front of the beaks, otherwise edentulous; nacre brilliant, silvery and iridescent; muscle scars faint. Animal pure milky white; lobes of the mantle united behind into two short siphons.¹

† **PILSBRYOCONCHA LINGUÆFORMIS** Morelet.

* *Anodonta linguaformis* MORELET, Ser. Conch., IV, 1875, p. 329, pl. XIV, fig. 5.—

* PÆTEL, Conch. Sam., III, 1890, p. 181.

* *Margaron (Unio) linguaformis* LEA, Syn., 1870, p. 48.

Siam; Cambodia.

† **PILSBRYOCONCHA EXILIS** Lea.

* *Anodonta exilis* LEA, Tr. Am. Phil. Soc., VI, 1839, p. 81, pl. XXII, fig. 68; * Obs., II, 1838, p. 81, pl. XXII, fig. 68.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 239.—* HANLEY, Test. Moll., 1842, p. 224; * Biv. Shells, 1843, p. 224.—

* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* MORELET, Ser. Conch., IV, 1875, p. 327.—* CLESSIN, Conch. Cab. Ano., 1875, p. 171, pl. LVI, figs. 6-8.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

¹According to Deshayes, JI. de Conch., 1875, pp. 81-85. He does not give any other characters of the soft parts.

- * *Margarita (Anodonta) exilis* LEA, Syn., 1838, p. 32.
 * *Anodon exilis* CATLOW and REEVE, Conch. Nom., 1845, p. 66.
 * *Margaron (Anodonta) exilis* LEA, Syn., 1852, p. 51; 1870, p. 82.
 * *Monocondylaea exilis* PÆTEL, Conch. Sam., III, 1890, p. 174.
 * *Anodonta siliqua* KÜSTER? Conch. Cab. Ano., 1852, p. 57, pl. XIV, fig. 5.
 * *Anodonta polita* MOUSSON, L. and Suss. Moll. Java, 1849, p. 98, pl. XIX, figs. 2, 3.—
 * MOUSSON, Zeits. für Mal., VII, 1851, p. 46.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 504.—* PÆTEL, Conch. Sam., III, 1890, p. 183.
 * *Margaron (Anodonta) polita* LEA, Syn., 1852, p. 53.
 * *Anodon politus* SOWERBY, Conch. Icon., XVII, 1867, pl. XII, fig. 36.
 * † *Monocondylaea compressa* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 190; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 30, pl. XI, fig. 29; * Obs., XI, 1867, p. 34, pl. XI, fig. 29.—* CLESSIN, Conch. Cab. Ano., 1876, p. 258, pl. LXXX, figs. 3, 4.
 * *Pseudodon compressa* CONRAD, Am. Jl. Conch., I, 1865, p. 233.
 * *Margaron (Monocondylaea) compressa* LEA, Syn., 1870, p. 73.
 * *Spatha compressa* PÆTEL, Conch. Sam., III, 1890, p. 188.
 * *Anodon jarana* SOWERBY, Conch. Icon., XVII, 1867, pl. XI, fig. 33.
 * *Anodon gracilis* SOWERBY? Conch. Icon., XVII, 1867, pl. XIV, fig. 45.
 * *Anodon kellelti* SOWERBY? Conch. Icon., XVII, 1867, pl. XIX, fig. 71.
 * † *Anodonta sempervirens* DESHAYES, Nouv. Arch. de Mus., X, 1874, p. 120, pl. v, figs. 4, 5.—* PÆTEL, Conch. Sam., III, 1890, p. 184.

Southeastern Asia, Sumatra, Java.

† PILSBRYOCONCHA CARINIFERA Conrad.

- * *Anodonta carinifera* CONRAD, Cover of Monography, No. 9, 1837.¹—TROSCHEL, Arch. für Naturg., IV, 1838, Pt. 2, p. 288.—B. H. WRIGHT, Check List, 1888.

Southeastern Asia, probably.

PILSBRYOCONCHA LEMSLEYI Morelet.

- * *Anodonta lemsleyi* MORELET,² Ser. Conch., IV, 1875, p. 328, pl. XIV, fig. 1.—PÆTEL, Conch. Sam., III, 1890, p. 181.

Cambodia, Siam.

PILSBRYOCONCHA SCHOMBURGKI von Martens.

- * *Anodonta schomburgki* VON MARTENS, Proc. Zool. Soc. Lond., 1860, p. 15.—
 * PÆTEL, Conch. Sam., III, 1890, p. 184.

Siam.

Genus MEDIONIDUS Simpson, 1900.

(Type, *Unio conradicus* Lea.)

Medionidus SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 77.

Shell elongated, rather inflated, often arcuate when adult, sometimes having a posterior ridge; dorsal slope and occasionally the posterior

¹There is a specimen of *Pilsbryoconcha* in the Philadelphia Academy which agrees perfectly with Conrad's description, and is probably the type. My attention was called to this by Mr. Pilsbry, who believes this to be the shell Conrad described. It is rather a small species.

²An old, elongated, rather peculiar shell. Conrad's species may be the young of it, and Morelet believes that Martens' *A. schomburgki* is very close to his shell. I have not seen a sufficient amount of material of this genus to determine with certainty whether all these are valid species.

portion of the shell plicately or nodulously wrinkled; epidermis smooth and bright, variegated with broken green rays and blotches; beak sculpture consisting of rather fine, subparallel, often broken ridges in two loops, the anterior rounded, the posterior somewhat angled, occasionally broken up into zigzags;¹ pseudocardinals small, stumpy and somewhat roughened; laterals rather short, slightly curved and club-shaped, remote; dorsal cicatrices few, placed under the hinge just behind the beaks; anterior cicatrices rather deep; posterior cicatrices rounded, large, and well impressed; anterior part of shell somewhat thickened; female shell slightly swollen just behind the middle of the base.

Animal with the inner gills wholly or in part free from the abdominal sac; marsupium occupying the central posterior part of the outer gills, sometimes extending nearly their whole length, consisting of few to many rather large, irregular ovisacs, which are not so distinctly marked out as in *Lampsilis*, but which have rounded bases; mantle much thickened on its lower edge, which is dark colored and sometimes papillous.

(Group of *Medionidus conradicus*.)

Shell small, usually with a well-defined posterior ridge; posterior end and sometimes the greater part of the shell slightly wrinkled or nodulous; epidermis rather smooth, painted with rays broken into irregular arrow-head markings or blotches; pseudocardinals rather small and imperfectly developed; laterals of left valve separated by a narrow, shallow furrow; nacre greenish, purplish, or bluish. The male shell is often decidedly arcuate; that of the female is swollen at or behind the center of the base. Animal having the characters of the genus.

† MEDIONIDUS CONRADICUS Lea.

* *Unio conradicus* LEA, Tr. Am. Phil. Soc., VI, 1834, p. 63, pl. IX, fig. 23; * Obs., I, 1834, p. 175, pl. IX, fig. 23.—* FERUSSAC, Guerin. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 176; * Biv. Shells, 1843, p. 176, pl. XXIII, fig. 22.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* KUSTER, Conch. Cab. Unio, 1861, p. 179, pl. LVI, fig. 5.—* SOWERBY, Conch. Icon., XVI, 1866, pl. LIV, fig. 278.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Margarita (Unio) conradicus* LEA, Syn., 1836, p. 13; 1838, p. 14.

* *Margaron (Unio) conradicus* LEA, Syn., 1852, p. 21.

* *Unio conradicus* CONRAD, Monog., X, 1838, p. 87, pl. XLVII, fig. 3.

* *Margaron (Unio) conradianus* LEA, Syn., 1870, p. 32.

* *Unio conradianus* B. H. WRIGHT, Check List, 1888.

Tennessee River drainage; Cahawba River, Alabama, and probably the entire Alabama River system.

¹The beaks in all the specimens of *Unio subtentus* that I have examined were too badly eroded to make out the character of the sculpture with certainty. In a general way they seemed to be much like those of the *Conradicus* group, only coarser.

† *MEDIONIDUS PARVULUS* Lea.

* *Unio parvulus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 307; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 45, pl. xvi, fig. 43; * Obs., XL, 1867, p. 49, pl. xvi, fig. 43.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) parvulus* LEA, Syn., 1870, p. 32.

Coosa River, Alabama; Chattanooga and Swamp creeks, northwest Georgia.

† *MEDIONIDUS PENICILLATUS* Lea.

* *Unio penicillatus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 171; * Jl. Ac. N. Sci. Phila., IV, 1859, p. 203, pl. xxiii, fig. 85; * Obs., VII, 1859, p. 21, pl. xxiii, fig. 85.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) penicillatus* LEA, Syn., 1870, p. 32.

Chattahoochee and Flint rivers, Georgia.

† *MEDIONIDUS KINGI* B. H. Wright.

Unio kingi B. H. WRIGHT, Nautilus, XII, 1900, p. 138.

Branch of Flint River, Baker County, Georgia.

† *MEDIONIDUS ACUTISSIMUS* Lea.

* *Unio acutissimus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 89, pl. x, fig. 18; * Obs., I, 1834, p. 99, pl. x, fig. 18.—* CONRAD, New F. W. Shells, 1834, p. 67.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* CONRAD, Monog., X, 1838, p. 86, pl. XLVII, fig. 2.—* HANLEY, Test. Moll., 1842, p. 177; * Biv. Shells, 1843, p. 177, pl. xxii, fig. 35.—* CATLOW and REEVE, Conch. Nom., 1845, p. 55.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 244.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. VIII, figs. 3, 3a, 3b.—* SOWERBY, Conch. Icon., XVI, 1866, pl. xxxvi, fig. 189.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 143.

* *Margarita (Unio) acutissimus* LEA, Syn., 1836, p. 14; 1838, p. 14.

* *Margaron (Unio) acutissimus* LEA, Syn., 1852, p. 21; 1870, p. 32.

* *Unio semiplicatus* KUSTER, Conch. Cab. Unio, 1862, p. 279, pl. xciv, fig. 4.¹—* PÆTEL, Conch. Sam., III, 1890, p. 167.

* *Unio rubellinus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 32; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 70, pl. xiii, fig. 51; * Obs., VI, 1858, p. 70, pl. xiii, fig. 51.—* SOWERBY, Conch. Icon., XVI, 1868, pl. xc, fig. 490.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 166.

* *Margaron (Unio) rubellinus* LEA, Syn., 1870, p. 32.

Alabama River system.

(Group of *Medionidus walkeri*.)

Shell rather short, inflated, with a high, posterior ridge; posterior slope strongly corrugated; epidermis dark, slightly clouded; nacre reddish or greenish.

¹There is a *Unio semiplicatus* of Troschel in Wiegman's Archiv. for 1841, p. 180, but I do not know what is. Kuster credits his species to Australia, but it is evidently Lea's shell, no such species being found in Australia.

† MEDIONIDUS WALKERI B. H. Wright.

* *Unio walkeri* B. H. WRIGHT, Nant., XI, 1897, p. 91.

* *Lampsilis walkeri* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 77, pl. 1, fig. 5.

Suwanee River, Florida; Ocklockonee River, Georgia.

(Group of *Medionidus subtentus*.)

Shell elongate-elliptical, scarcely inflated, with an ill-defined posterior ridge, the posterior slope being strongly, corrugately wrinkled; beaks apparently sculptured with coarse, irregular, broken ridges, somewhat doubly looped; epidermis bright, with broken rays which sometimes form irregular patches, but are usually developed into square spots; pseudocardinals rather small, stumpy; laterals rather strong, club-shaped, generally showing traces of vertical striation; nacre dull, bluish-white. The female shell is very slightly inflated just behind the center of the base, the male often becomes arcuate when old.

Animal unknown.

† MEDIONIDUS SUBTENTUS Say.

* *Unio subtentus* SAY, Jl. Ac. N. Sci. Phila., V, 1825, p. 130; Am. Conch., I, No. 2, 1831, pl. XV.—* CONRAD, Am. Jl. Sci., XXVI, 1834, p. 343, pl. I, fig. 1; * New F.W. Shells, 1834, p. 71.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* MÖLLER, Syn. Nov. Gen., 1836, p. 209.—* CONRAD, Monog., X, 1838, p. 85, pl. XLVIII, fig. 1.—* HANLEY, Test. Moll., 1842, p. 176; * Biv. Shells, 1843, p. 176, pl. XX, fig. 34.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 25, pl. IV, figs. 2, 2a, 2b.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* KUSTER, Conch. Cab. Unio, 1856, p. 164, pl. XLVII, fig. 4.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXII, fig. 370.—* B. H. WRIGHT, Check List, 1888.—* PATEL, Conch. Sam., III, 1890, p. 168.

* *Margarita (Unio) subtentus* LEA, Syn., 1836, p. 13; 1838, p. 14.

* *Margaron (Unio) subtentus* LEA, Syn., 1852, p. 21; 1870, p. 32.

* *Unio subtenta* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 555; 3d ed., II, 1839, p. 675.

Tennessee and Cumberland River systems.

Genus NEPHRONAIAS Crosse and Fischer, 1893.

(Type, *Unio plicatulus* Charpentier.)

Nepronaias CROSSE and FISCHER, Miss. Sci., Pt. 7, II, 1893, p. 556.

Shell elliptical, biangulate behind, that of the male showing a tendency to become arcuate with age, the female usually having a post-basal inflation and never arcuate; surface concentrically sculptured; beaks low, with faint broken ridges which show a tendency to fall into two rounded loops; pseudocardinals generally rather compressed, ragged, laterals obliquely ridged; cavity of the beaks rather deep, dorsal muscle scars distinct, running in a line from the beak cavity downward and forward. Animal with the branchiae rounded below, outer the larger behind, inner the larger anteriorly, free from the abdominal sac

for all or part of their length; mantle thickened on the edge, which is often dark colored; anal and branchial openings large and well furnished with papillæ. Nothing is known of the marsupium of any members of this group, the animals examined probably being males; but it is most likely that when gravid the ovules will be found in distinctly marked ovisacs in the posterior part of the outer gills, something as in *Lampsilis*.

(Group of *Nephronaias medellinus*.)

Shell scarcely sulcate, shining, rather brightly rayed, distinctly biangulate behind, solid, swollen; beaks rather full, with delicate sculpture; pseudocardinals solid, stumpy, trigonal; laterals heavy, somewhat club-shaped; nacre rich and soft, silvery or purple; female shell considerably inflated at the post-basal region, and differing somewhat from that of the male.

† NEPHRONAIAS MEDELLINUS Lea.

? *Unio purpuratus* SAY, New Harmony, Diss. (newspaper form), January 15, 1831.¹

* *Unio medellinus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 39, pl. XII, fig. 34; * Obs., II, 1838, p. 39, pl. XII, fig. 34.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 236.—* HANLEY, Test. Moll., 1842, p. 193; * Biv. Shells, 1843, p. 193, pl. XXI, fig. 19; pl. XXIII, fig. 8.—* CATLOW and REEVE, Coneh. Nom., 1845, p. 61.—* CHENU, Ill. Coneh., 1858, pl. XVII, figs. 6, 6a, 6b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Coneh. Cab., 1861, p. 162, pl. XLVI, fig. 5.—* SOWERBY, Coneh. Icon., XVI, 1866, pl. XXXIII, fig. 171.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Coneh. Sam., III, 1890, p. 158.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 603.

* *Margarita (Unio) medellinus*, LEA, Syn., 1836, p. 26; 1838, p. 19.

* *Margaron (Unio) medellinus*, LEA, Syn., 1852, p. 29; 1870, p. 45.

† * *Unio strebeli*, LEA,² Pr. Ac. N. Sci. Phila., X, 1866, p. 133; * JI. Ac. N. Sci. Phila., VI, 1868, p. 318, pl. LI, fig. 131; * Obs., XII, 1869, p. 78, pl. LI, fig. 131.—* B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 595.

* *Margaron (Unio) Strebeli*, LEA, Syn., 1870, p. 53.

State of Vera Cruz, Mexico.

† NEPHRONAIAS SAPOTALENSIS Lea.

* *Unio sapotalensis* LEA, Pr. Am. Phil. Soc., II, 1841, p. 30; * Tr. Am. Phil. Soc., VIII, 1842, p. 233, pl. XXI, fig. 47; * Obs., III, 1842, p. 71, pl. XXI, fig. 47.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Coneh., 1858, pl. XXXIII, figs. 4, 4a, 4b.—* SOWERBY, Coneh. Icon., XVI, 1868, pl. XLI, fig. 495.—* B. H. WRIGHT, Check List.

¹ Believed by Conrad to be the same as Lea's *medellinus*. Fisher and Crosse in Mission Scientifique (see following reference) are doubtful about this, as Say's species was never figured, and his description does not just agree with Lea's species. As the type is not known to be in existence I think under the circumstances, since we can not be sure of what Say described, it is best to use Lea's name.

² The type which is in the collection of the U. S. National Museum is, without doubt, a young *medellinus*.

1888.—* PÆTEL, Conch. Sam., III, 1890, p. 166.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 601, pl. LXVII, figs. 2, 2a.

* MARGARON (*Unio*) *sapotalsensis* LEA, Syn., 1852, p. 29; 1870, p. 45.

Sapotal River, Vera Cruz, Mexico.

(Group of *Nephronaias averyi*.)¹

Shell triangular-ovate, bluntly pointed in front, and biangulate behind, considerably inflated, irregularly sulcate; umbonal region moderately developed; beaks sculptured with fine, irregular ridges which have a tendency to fall into two rounded loops; pseudocardinals slightly compressed, ragged; laterals having feeble, oblique striation; front part of the shell somewhat thickened; posterior part thinner; nacre dull, whitish.

Animal unknown.

† NEPHRONAIAS AVERYI Lea.

* *Unio averyi* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 281; * JI. Ac. N. Sci. Phila., IV, 1860, p. 269, pl. XLIV, fig. 149; * Obs., VII, 1860, p. 87, pl. XLIV, fig. 149.—

* B. H. WRIGHT, Check List, 1888.

* MARGARON (*Unio*) *averyi* LEA, Syn., 1870, p. 36.

Isthmus of Darien.

† NEPHRONAIAS CALDWELLII Lea.

* *Unio caldwellii* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 118; * JI. Ac. N. Sci. Phila., IV, 1860, p. 265, pl. XLIII, fig. 145; * Obs., VII, 1860, p. 83, pl. XLIII, fig. 145.—

* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIX, fig. 477.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 147.

* MARGARON (*Unio*) *caldwellii* LEA, Syn., 1870, p. 35.

Isthmus of Darien.

(Group of *Nephronaias macnielii*.)

Shell solid, slightly sulcate, biangulate behind, covered with a dull, olivaceous epidermis and having faint, bluish green rays; beaks rather full, sculpture feeble, consisting of slightly outlined ridges, somewhat doubly looped, the looped lines becoming swollen or nodulous at their bases; pseudocardinals compressed and ragged; laterals obliquely striated; nacre bluish white, very soft and brilliant, somewhat iridescent posteriorly; male and female shells differing but slightly. Animal unknown.

† NEPHRONAIAS MACNIELII Lea.

* *Unio macnielii* LEA, Pr. Ac. N. Sci. Phila., XIII, 1869, p. 124; * JI. Ac. Nat. Sci. Phila., 1874, p. 25, pl. VIII, fig. 22; * Obs., XIII, 1874, p. 29, pl. VIII, fig. 22.—

* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

Nicaragua.

¹ I have only seen two specimens of Lea's *Unio averyi* and one of his *U. caldwellii*, and I can not be certain as to their systematic position. In the latter the beak sculpture, which is slightly worn, is much of the character of the species of *Nephronaias* generally, and not like that of *Unio*, and though somewhat worn the shell shows traces on the posterior end of faint, bluish green rays. One of the former is slightly more inflated in the post-basal region than the other.

† *NAPHRONAIAS OREGONENSIS* Lea.¹

- * *Unio oregonensis* LEA, Pr. Am. Phil. Soc., V, 1852, p. 252; * Tr. Am. Phil. Soc., X, 1852, p. 275, pl. XXII, fig. 33; * Obs., V, 1852, p. 31, pl. XXII, fig. 33.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* MUSGRAVE, Phot. Conch., 1863, pl. II, fig. 5.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIV, fig. 383.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 162.
 * *Margaron (Unio) oregonensis*, LEA, Syn., 1852, p. 29; 1870, p. 45.

Nicaragua.

† *NEPHRONAIAS ROWELLII* Lea.

- * *Unio rowellii* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 153; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 256, pl. XL, fig. 136; * Obs., VII, 1860, p. 74, pl. XL, fig. 136.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVIII, fig. 471.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 166.
 * *Margaron (Unio) rowellii* LEA, Syn., 1870, p. 35.

Nicaragua; Guatemala; Chagres River, Colombia.

† *NEPHRONAIAS GOASCORANENSIS* Lea.

- * *Unio goascoranensis* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 118; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 258, pl. XLI, fig. 139; * Obs., VII, 1860, p. 76, pl. XLI, fig. 139.—* B. H. WRIGHT, Check List, 1888.
 * *Margaron (Unio) goascoranensis* LEA, Syn., 1870, p. 45.

Goascoran River, Honduras.

† *NEPHRONAIAS DYSONII* Lea.

- * *Unio dysonii* LEA, Pr. Ac. N. Sci. Phila., XI, 1859, p. 152; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 252, pl. XXXIX, fig. 132; * Obs., VII, 1860, p. 70, pl. XXXIX, fig. 132.—* B. H. WRIGHT, Check List, 1888.
 * *Margaron (Unio) dysonii* LEA, Syn., 1870, p. 35.

Honduras; Costa Rica.

† *NEPHRONAIAS TEHUANTEPECENSIS* Crosse and Fischer.

- * *Unio tehuantepecensis* CROSSE and FISCHER, J. de Conch., XLI, 1893, p. 296.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 618, pl. LXV, figs. 3, 3a, 4.

Isthmus of Tehuantepec, Oaxaca, Mexico; Honduras.

(Group of *Nephronaias reticulatus*.)

Shell evenly elliptical, slightly biangulate behind, solid and inflated; surface covered with fine, concentric ridges and having delicate, radiating furrows which cut the sulcations and form them into loops.

¹I formerly believed that this species, with the one preceding and following it, ran together. More recent study, with a large amount of additional material, inclines me to think that perhaps they may be valid species. Under the name of *Unio rowellii* Mr. Lea has in his collection two large *N. macneilii*, one large *rowellii*, three smaller ones and several valves, and two *goascoranensis*, a species which is very close to the rest, but probably as good as most of the Mexican forms. I find specimens which I can scarcely refer with certainty to *macneilii* or *oregonensis*, but which seem to have to some extent the characters of both. This, with a number of the Mexican and Central American forms, is exceedingly puzzling. This species occurs with *N. rowellii* and *macneilii*.

† NEPHRONAIAS RETICULATUS Simpson.

* *Nephronaias reticulatus* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 77, pl. II, fig. 3.

Patook River, Honduras.

(Group of *Nephronaias scamnatus*.)

Shell elliptical to elongate, compressed, strongly sulcate, rather solid, subbiangular behind; that of the female inflated at post-base; epidermis olive to brown, sometimes feebly rayed; beak sculpture delicate and but slightly marked, consisting of faint, parallel ridges, which show a tendency to fall into two rounded loops; hinge rather solid; pseudocardinals stumpy, ragged, laterals straight; front part of the shell solid and suddenly becoming thinner behind; nacre bluish, white, salmon, or purple.

Animal unknown.

† NEPHRONAIAS SCAMNATUS Morelet.

* *Unio scamnatus* MORELET, Test. Nov., No. 1, 1849, p. 30.—* POEY, Mem. Hist. Cuba, II, 1858, p. 417.—* ARANGO, Faun. Mal, Cuba, 1878, p. 144.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 167.

* *Margaron (Unio) scamnatus* LEA, Syn., 1852, p. 40; 1870, p. 35.

* *Unio gundlachi* SOWERBY, Conch. Icon., XVI, 1866, pl. XLVI, fig. 248.

Cuba; Honduras?

† NEPHRONAIAS GUNDLACHI Dunker.

* *Unio gundlachi* DUNKER,¹ Mal. Blatt., V, 1858, p. 228.—* ARANGO, Faun. Mal. Cuba, 1878, p. 144.—* PÆTEL, Conch. Sam., III, 1890, p. 154.

Cuba.

NEPHRONAIAS ÆRUGINOSUS Morelet.²

* *Unio æruginosus* MORELET, Test. Noviss., I, 1849, p. 29.—* B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 596, pl. LXII, figs. 2a, 2a, 2b.

River Michol, at Palenque, State of Chiapas, Mexico.

¹ Lea and Poey regard this as the equivalent of *N. scamnatus* of Morelet, while Arango believes it to be distinct. Unfortunately both species were described in Latin and not figured, but Dunker states that his species is inflated posteriorly and has a bluish nacre, and it is probable that he had before him a female shell when he wrote the description, for one of the Cuban species, which I think is *gundlachi*, has a bluish nacre, and the female shell is decidedly swollen at the post-basal region. In another form, which I presume is *N. scamnatus* of Morelet, the nacre is rather dark, the shell somewhat elongated, and that of the female does not greatly differ from the male.

² I am not positive that this species belongs here, but its form is like that of most of the members of the genus, and Crosse and Fischer state that the beaks in their specimen are entire and smooth. In several of the species of this genus the beak sculpture is very faint, while in *Unio* it is almost always strong. These considerations lead me to place the species here.

NEPHRONAIAS YZABALENSIS Crosse and Fischer.

* *Unio yzabalensis* CROSSE and FISCHER, J. de Conch., XL, 1892, p. 294.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 597, pl. LXIV, figs. 4, 4a, 4b.¹

Lake Ysabal, Guatemala.

† NEPHRONAIAS RUGULOSUS Charpentier in Kuster.

* *Unio rugulosus* CHARPENTIER in KUSTER, Conch. Cab. Unio, 1856, p. 154, pl. XLIV, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 166.²

Habitat.—A shell in the Lea collection, which I take to be this, is marked "Mexico."

NEPHRONAIAS PERSULCATUS Lea.³

* *Unio persulcatus* LEA, Pr. Ac. N. Sci. Phila., XI, 1859, p. 153; * JI. Ac. N. Sci. Phila., IV, 1860, p. 255, pl. XL, fig. 135; * Obs., VII, 1860, p. 73, pl. XI, fig. 135.—* MUSGRAVE, Phot. Conch., 1863, pl. II, fig. 3.—* B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 614.

* *Margaron (Unio) persulcatus* LEA, Syn., 1870, p. 36.

Mexico.

† NEPHRONAIAS CALIMATARUM Morelet.

* *Unio calimatarum* MORELET, Test. Noviss., I, 1849, p. 30.—* ? SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIV, fig. 385.⁴—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 147.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 612.

Margaron (Unio) calimatarum LEA, Syn., 1870, p. 63.

Mexico; Guatemala.

† NEPHRONAIAS TABASCOENSIS Charpentier in Kuster.

* *Unio tabascoensis* CHARPENTIER in KUSTER, Conch. Cab. Unio, 1856, p. 153, pl. XLIV, fig. 3.⁵—* B. H. WRIGHT, Check List, 1888.—* CROSSE and FISCHER, Miss. Sci., Pt. 7, II, 1894, p. 611.

* *Margaron (Unio) tabascoensis* LEA, Syn., 1870, p. 61.

Mexico; Honduras; Cuba?

¹ Probably a mere variety of *aruginosus*.

² Kuster credits this species to Australia and says that it has a purple naere. The naere of all the Australian species is whitish, and the shell is evidently a member of this group.

³ This and the related species are extremely puzzling. Although I have been able to examine a good deal of material, yet I find great variation in the specimens, and few of them accurately agree with the figures and descriptions.

⁴ Fischer and Crosse believe this to be Morelet's species. It looks like a short *cuprinus*, but Sowerby says it is solid, and has a dirty-white naere, while *cuprinus* is rather thin and is copper colored throughout.

⁵ The specimens which I have seen do not wholly agree with Kuster's figure and description, but are nearer to that than anything else. The material in the U. S. National Museum is mostly from Honduras though there are specimens credited to Cuba. Lea referred these to *N. scammatus* Morelet.

† NEPHRONAIAS PLICATULUS Charpentier in Kuster.

* *Unio plicatulus* CHARPENTIER in KUSTER, Conch. Cab. Unio, 1856, p. 154, pl. XLIV, fig. 4.—*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 598.

Mexico.

† NEPHRONAIAS RAVISTELLUS Morelet.

* *Unio ravistellus* MORELET, Test. Noviss., No. 1, 1849, p. 29.—*KUSTER, Conch. Cab. Unio, 1856, pl. XLII, fig. 6.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 165.—*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 609, pl. LXI, figs. 4, 5, and 5a.

*Margaron (*Unio*) *ravistellus* LEA, Syn., 1870, p. 40.

Guatemala.

† NEPHRONAIAS VELLICATUS Reeve.

* *Unio vellicatus* REEVE, Conch. Icon., XVI, 1865, pl. XXII, fig. 103.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 171.—*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 610.

Guatemala.

NEPHRONAIAS MELLEUS Lea.

* *Unio melleus* LEA, Pr. Ac. N. Sci. Phila., XI, 1859, p. 152; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 250, pl. XXXVIII, fig. 129; *Obs. VII, 1860, p. 68, pl. XXXVIII, fig. 129.—*REEVE, Conch. Icon., XVI, 1865, pl. XXI, fig. 92.—*PÆTEL, Conch. Sam., III, 1890, p. 158.

*Margaron (*Unio*) *melleus* LEA, Syn., 1870, p. 35.¹

Mexico or Central America, no doubt.

GENUS GLEBULA Conrad 1853.

(Type, *Unio rotundata* Lamarek.)²

Glebula CONRAD, Pr. Ac. N. Sci. Phila., 1853, p. 268.

Shell solid, much inflated, short elliptical, bluntly pointed and slightly biangulate behind, with a low, posterior ridge; female shell swollen at post base; beaks compressed, the sculpture not seen; epidermis brownish, clothlike; pseudocardinals divided into irregularly radiating, granular lamina, sometimes to the number of a dozen or more in each valve; hinge plate reduced to a mere rounded line behind the pseudocardinals; laterals short, remote, anterior adductor scars deep, smooth; there is in each valve only a single, large, semicircular posterior scar

¹ I am not certain just where this and the preceding species should be placed. This remark can be applied to a number of other forms placed in this genus.

² I have examined a number of animals of the only species of this curious genus, but regret that they were badly decayed, and that none were gravid. In one specimen several ovisacs were distended with gas, and these were separated by a deep sulcus. This fact, the union of the inner gills with the abdominal sac, and the post-basal swelling seen on some shells incline me to believe that it is a highly organized species.

with a straight edge in front; dorsal scars numerous, distinct, placed back of the pseudocardinals; cavity of the beaks not deep or compressed; nacre dull lurid to purplish brown.

Animal with gills nearly equal in size, inner slightly the larger in front, united the whole length to the abdominal sac; ovisacs apparently separated from each other by a sulcus as in *Lampsilis*; palpi very large, wide, wrinkled; mantle having a wide, slightly thickened border, double edged, the inner being beautifully toothed; branchial opening papillose; anal crenulate; foot small; whole animal the color of the nacre.

† GLEBULA ROTUNDATA Lamarck.

* *Unio rotundata* LAMARCK, An. sans Vert., VI, 1819, p. 75.—*LEA, Obs., I, 1834, p. 201.

* *Glebula rotundata* CONRAD, Pr. Ac. N. Sci. Phila., 1853, p. 268.

* *Unio rotundatus* HANLEY, Test. Moll., 1842, p. 201; *Biv. Shells, 1843, p. 201, pl. XXIII, fig. 30.—*CATLOW and REEVE, Conch. Nom., 1845, p. 63.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—*KUSTER, Conch. Cab. Unio, 1861, p. 256, pl. LXXXVI, fig. 4.—*REEVE, Conch. Icon., XVI, 1865, pl. XXIII, fig. 106.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 166.—*CALL, Tr. Ac. Sci. St. Louis, VII, 1895, p. 47, pl. IX.

* *Margarita (Unio) rotundatus* LEA, Syn., 1836, p. 33; 1838, p. 23.

* *Margaron (Unio) rotundatus* LEA, Syn. 347, 1852, p. 34; 1870, p. 55.

* *Unio suborbiculata* LAMARCK, An. sans Vert., VI, 1819, p. 81.—*LEA, Obs., I, 1834, p. 201.—*DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 546; 3d ed., II, 1839, p. 671.

* *Unio suborbiculatus* FERUSSAC, Guer. Mag., 1835, p. 26.

Unio glebulus SAY, Transylvania Jl., IV, 1831, p. 526.—SAY, Am. Conch., No. IV, 1832, pl. XXXIV; No. V, 1832, p. 4 (of cover).—CONRAD, New F. W. Shells, 1834, p. 69.—CHENU, Bib. Conch., 1st ser., III, 1845, p. 46, pl. XI, figs. 10-12.—SOWERBY, Conch. Icon., XVI, pl. LXXIV, fig. 384.

* *Unio glebulus* PÆTEL, Conch. Sam., III, 1890, p. 154.

* *Unio subglobosus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 30, pl. II, fig. 3; *Obs., I, 1834, p. 142, pl. II, fig. 3.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXIV, fig. 321.

* *Unio granadensis* CONRAD,¹ Pr. Ac. N. Sci. Phila., VII, 1855, p. 256.—*B. H. WRIGHT, Check List, 1888.

Louisiana; eastern Texas; Rio Grande River; Escambia County, Florida.

¹ Conrad says it is suboval, with the disks somewhat flattened, with minute, radiating lines extending to the tips of the beaks. The *rotundata* sometimes shows radiating ridges at the beaks in eroded specimens. He also states that the nacre of his species is purple. The description therefore fits Lamarck's species perfectly, though I have never known the species to come from so far west as the Rio Grande—Conrad's locality.

Genus *OBOVARIA* Rafinesque, 1819.(Type, *Unio retusa* Lamarck.)*Obovaria* RAFINESQUE, J. de Phys. Chim. Hist. Nat., 1819, p. 426.¹

Shell short, oval, rounded or retuse, solid, inflated, thick in front, thinner behind, with high beaks which are sculptured with very faint, irregular, often broken and slightly nodulous ridges which show a tendency to fall into two loops, the posterior often open behind; epidermis dull, brownish, silky or clothlike, rarely rayed, rays indistinct; female shell but slightly inflated in the post-basal region, commonly having a shallow furrow or a flattened area at the posterior end; pseudocardinals solid, stumpy; laterals short, club-shaped; anterior and posterior cicatrices deep and distinct; naere bluish white or purple.

Animal with very short gills, the inner united to the abdominal sac throughout; marsupium projecting far below the rest of the branchiæ and occupying the posterior portion of the outer gills, dolabriform or kidney-shaped; mantle with a wide, thickened double-edged border, the inner edge of which is often slightly toothed at its postbasal part.

Subgenus *OBOVARIA* Rafinesque, 1819.(Type, *Unio retusa* Lamarck.)

Shell retrose to short oval; beaks high, central; pseudocardinals rarely parallel with the laterals; cavity of the beaks deep, subcompressed; naere bluish white or purple. Animal with the characters of the genus.

†*OBOVARIA RETUSA* Lamarck.

**Unio retusa* Lamarck, An. sans Vert., VI, 1819, p. 72.—FERUSSAC, Guer. Mag., 1835, p. 28.—DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 534; 3d ed., II, 1839, p. 668.

**Unio retusus* CONRAD, Monog., II, 1836, p. 19, pl. VIII.—*REEVE, Conch. Syst., I, 1841, p. 118, pl. LXXXVIII, fig. 16.—*HANLEY, Test. Moll., 1842, p. 202; *Biv. Shells, 1843, p. 202, pl. XXI, fig. 11.—*CATLOW and REEVE, Conch. Nom., 1845, p. 63.—*KUSTER Conch. Icon. Cab. Unio, 1852, p. 58, pl. XIII, fig. 2.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 256.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXI, fig. 363.—*CALKINS, Pr. Ottawa Ac. Sci., 1874, p. 44.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 165.—*CALL, Tr. Acad. Sci. St. Louis, VII, 1895, p. 45, pl. VIII.

**Margarita (Unio) retusus* LEA, Syn., 1836, p. 34; 1838, p. 23.

**Margaron (Unio) retusus* LEA, Syn., 1852, p. 35; 1870, p. 56.

**Unio torsus* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 149, pl. LVII, figs. 1, 2.

Ohio, Cumberland, and Tennessee river systems.

¹ I am utterly unable to guess as to what Rafinesque's first species (*Obovaria obovalis*) is, and the second (*O. torsa*) is either the *Unio retusa* of Lamarck or a variety of Lea's *U. circulus* with purple naere, I can not be certain which. Which of these two nearly related species it may be, I consider it sufficient to establish the genus, his description being applicable.

† OBOVARIA CIRCULUS Lea.

- * *Unio circulus* LEA, Tr. Am. Phil. Soc., III, 1829, p. 433, pl. ix, fig. 14;¹ * Obs., I, 1834, p. 47, pl. ix, fig. 14.—SHORT and EATON, Transylvania Jl., 1831, p. 77.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 549; 3d ed., II, 1839, p. 672.—* ANTON, Verz. der Conch., 1839, p. 14.—* HANLEY, Test. Moll., 1842, p. 201; * Biv. Shells, 1843, p. 201, pl. xx, fig. 23.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* KUSTER, Conch. Cab. Unio, 1852, p. 41, pl. viii, fig. 2.—* CHENU, Ill. Conch., 1858, pl. xvi, figs. 1, 1a, 1b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* REEVE, Conch. Icon., XVI, 1865, pl. xxvii, fig. 135.—* B. H. WRIGHT, Check List, 1888.
- * *Margarita (Unio) circulus* LEA, Syn., 1836, p. 33; 1838, p. 22.
- * *Margaron (Unio) circulus* LEA, Syn., 1852, p. 34; 1870, p. 55.
- * *Unio subrotundus* FERUSSAC, part, Guer. Mag., 1835, p. 28.²—CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.
- * *Unio subrotundus* Raf. v. *circulus* PÆTEL, Conch. Sam., III, 1890, p. 168.
- * *Mya rotunda* WOOD, Index Test. (Rev.), 1856, p. 199, pl. i, Supp., fig. 1.

Ohio, Tennessee and Cumberland river systems; southeast Louisiana; Alabama and Tombigbee drainage? Reported from Michigan and the St. Lawrence drainage, but the forms in this area are probably *O. leibii*, which is perhaps only a variety of *O. circulus*.

† OBOVARIA TINKERI B. H. Wright.

- * *Unio tinkeri* B. H. WRIGHT, Nautilus, XIII, 1899, p. 7.
- * *Obovaria tinkeri* SIMPSON, Pr. Acad. Nat. Sci. Phila., 1900, p. 78, pl. iv, fig. 3.

Tombigbee River, Alabama; Columbus, Mississippi.

† OBOVARIA LENS Lea.³

- * *Unio lens* LEA, Tr. Am. Phil. Soc. Phila., IV, 1831, p. 80, pl. viii, fig. 10; * Obs. I, 1834, p. 90, pl. viii, fig. 10.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* HANLEY, Test. Moll., 1842, p. 201; * Biv. Shells, 1843, p. 201, pl. xxi, fig. 4.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* CHENU, Ill. Conch., 1856, pl. viii, figs. 10, 10a, 10b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* REEVE, Conch. Icon., XVI, 1865, pl. xxvii, fig. 134.—* SOWERBY, Conch., Icon., XVI, 1866, pl. xxxvii, fig. 200.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 157.
- * *Margarita (Unio) lens* LEA, Syn., 1836, p. 33; 1838, p. 22.
- * *Margaron (Unio) lens* LEA, Syn., 1852, p. 34; 1870, p. 55.
- * ? *Unio levigatus* SAY, Am. Conch., VI, 1834.—CONRAD, New F. W. Shells, 1834, p. 70.—CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.

¹ The figured type has a purple nacre.

² Ferussac makes this = *U. orbiculatus* Hildreth also.

³ This is sometimes considered a synonym of *O. circulus*. The two species are excessively variable in form, and both often have a pink or purplish nacre. While individuals may occasionally be found which can only be referred to either species with doubt, the majority of specimens can readily be determined, and Dr. V. Sterki informs me that he has found differences in the animals of the two. As I have only been able to examine the soft parts of young specimens I have not detected any differential characters in them.

†OBOVARIA LENS var. DEPYGIS Conrad.

**Unio depygis* CONRAD, Am. Jl. Conch., II, 1866, p. 107, pl. x, fig. 1.—B. H. WRIGHT, Check List, 1888.

Ohio River drainage; southern Michigan.

†OBOVARIA UNICOLOR Lea.¹

**Unio unicolor* LEA, Tr. Am. Phil. Soc., IV, 1845, p. 163; *Tr. Am. Phil. Soc., IX, 1848, p. 74, pl. iv, fig. 12; *Obs., IV, 1848, p. 48, pl. iv, fig. 12.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 171.

*Margaron (*Unio*) *unicolor* LEA, Syn., 1852, p. 34; 1870, p. 55.

Mississippi and Alabama, in streams flowing into the Gulf.

†OBOVARIA LEIBII Lea.²

**Unio leibii* LEA, Proc. Acad. Nat. Sci. Phila., VI, 1862, p. 168; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 44, pl. xv, fig. 42; Obs., XI, 1867, p. 48, pl. xv, fig. 42.—*B. H. WRIGHT, Check List, 1888.

*Margaron (*Unio*) *leibii* LEA, Syn., 1870, p. 36.

Lake Erie and streams falling into it; southern Michigan; ?Sequatchie River, Tennessee.

†OBOVARIA ROTULATA B. H. Wright.

**Unio rotulatus* B. H. WRIGHT, Naut., XIII, 1899, p. 22.

**Obovaria rotulata* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 78, pl. iv, fig. 2.

Escambia River, Escambia County, Florida.

Subgenus PSEUDOÖN Simpson, 1900.

(Type, *Unio ellipsis* Lea.)

Shell elliptical, inflated, solid, that of the males slightly pointed at the upper posterior part; epidermis brownish or blackish, rayless or very feebly rayed, beaks anterior; pseudocardinals solid, stumpy, or slightly elongate in age, and showing a tendency toward being parallel with the laterals; cicatrices deep; nacre silvery, iridescent posteriorly.

Animal with mantle having a wide, thickened, double border, the inner edge being toothed throughout below; gills small; marsupium not reaching to the posterior end of the outer branchiæ, though extending quite well forward; ovisacs rather numerous, large and distinct, tinted with purple below; anal and branchial openings finely toothed.

¹The type is a young female, somewhat broken, and besides this Lea has a shell which he has so named that I believe is *O. castaneus*. Under the name of *Unio lens* (Cat. No. 85743, U.S.N.M.) he has a young shell from Jackson, Mississippi, which is a young male *unicolor*. The species grows to the size of *O. lens*, but shows dull rays and is more inflated than that species.

²Dr. Lea states that the type is from Erie County, Michigan. There is no county of that name in the State, and his specimens may be from Erie County, Ohio. I am very doubtful whether it is more than a dwarf variety of *U. circulus*, as specimens have been seen from Tennessee which seem to stand between the two forms.

† OBOVARIA ELLIPSIS Lea.

- * ? *Unio brevalis* SOWERBY, Rec. and Foss. Shells, XVI, fig. (no date).
 * *Unio ellipsis* LEA, Tr. Am. Phil. Soc., III, 1828, p. 268, pl. IV, fig. 4; * Obs., I, 1834, p. 10, pl. IV, fig. 4.—SHORT and EATON, Transylvania JI., 1831, p. 75.—* SAY, Am. Conch. I, No. II, 1831.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 547; 3d ed., II, 1839, p. 672.—* HANLEY, Test. Moll., 1842, p. 188; * Biv. Shells, 1843, p. 188, pl. XXI, fig. 7.—* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 150, pl. LVIII, figs. 1, 2.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 22, pl. IV, figs. 1, 1a, 1b.—* KUSTER, Conch. Cab. Unio, 1852, p. 43, pl. VIII, fig. 4.—* CHENU, Ill. Conch., 1858, pl. XVIII, figs. 1, 1a, 1b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXX, figs. 4, 24.—* CALKINS, Pr. Ottawa Acad. Sci., 1874, p. 42.—* LATCHFORD, Tr. Ottawa Field Nat. Club, 1882, p. 50.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 151.
 * *Margarita (Unio) ellipsis* LEA, Syn., 1836, p. 22; 1838, p. 18.
 * *Margaron (Unio) ellipsis* LEA, Syn., 1852, p. 26; 1870, p. 41.
 * *Unio triangularis* SAY, Am. Conch., VI, 1834.
 * *Unio olivarius* CONRAD, New F. W. Shells, 1834, p. 70.—* FERUSSAC, Guer. Mag., 1835, p. 28.
 * *Unio taitianus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXVII, fig. 338.
 * *Unio pealei* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 191; * JI. Ac. N. Sci. Phila., VIII, 1874, p. 26, pl. VIII, fig. 23; * Obs., XIII, 1874, p. 30, pl. VIII, fig. 23.—* B. H. WRIGHT, Check List, 1888.

Upper Mississippi system as far south as the Tennessee and Arkansas rivers; St. Lawrence drainage area.

† OBOVARIA CASTANEA Lea.

- * *Unio castaneus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 91, pl. XI, fig. 21; ¹ * Obs., I, 1834, p. 101, pl. XI, fig. 21.—* CONRAD, New F. W. Shells, 1834, p. 68.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* HANLEY, Test. Moll., 1842, p. 188; Biv. Shells, 1843, p. 188, pl. XXII, fig. 30.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 246.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XI, figs. 5, 5a, 5b.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 147.
 * *Margarita (Unio) castaneus* LEA, Syn., 1836, p. 22; 1838, p. 18.
 * *Margaron (Unio) castaneus* LEA, Syn., 1852, p. 26; 1870, p. 41.

Streams flowing into the Gulf of Mexico, from the Alabama River west to the Nechez.

The following species are unfigured and indeterminable.

- * *Obovaria cordata* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 312, pl. LXXXII, figs. 6, 7.
 * *Obovaria stegaria* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 312, pl. LXXXII, figs. 4, 5.

¹ The name *castaneus* was applied to some *Unio* which I am unable to determine, by Rafinesque in a continuation of a monograph on the Bivalve Shells of the River Ohio, etc., in October, 1831. According to Scudder, Lea's name was published the latter part of the same year, but I have no means of knowing which appeared first. Under the circumstances I use Lea's name.

* *Obovaria striata* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 311.

* *Obovaria torsa* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 312, pl. LXXXII, figs. 1, 3.

* *Obliquaria retusa* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 306, pl. LXXXI, figs. 19, 20.

Genus *PLAGIOLA* (Rafinesque, 1819) Agassiz.

(Type, *Unio securis* Lea.)

Plagiola RAFINESQUE, J. de Phys. Chim. Hist. Nat., 1819, p. 426.—AGASSIZ, Arch. für Nat., 1852, p. 48, redefined.

Shell triangular ovate, somewhat inflated, solid, with a distinct and often sharp posterior ridge; surface concentrically sculptured; umbonal area somewhat flattened; beaks high, sculptured with delicate, parallel, doubly looped ridges, the anterior loop rounded, the posterior angular; pseudocardinals ragged; laterals club-shaped, straight or slightly curved; cavity of the beaks moderate, often somewhat compressed; nacre silvery; female shell more or less inflated at postbasal region.

Animal with the outer gills narrow in front, wide behind; inner gills wide in front, posteriorly free or united to abdominal sac; marsupium occupying the posterior part of the outer gills, but not extending quite to the hinder end, consisting of well-marked ovisacs which are rounded below; a distinct sulcus extends the whole length of the kidney-shaped marsupium at the inside and outside at some distance above its base; mantle edge thickened and somewhat doubled, in some cases toothed or fringed below.

Subgenus *PLAGIOLA* (Rafinesque, 1819) Agassiz.

Shell solid, surface irregularly concentrically ridged; epidermis smoothish, but here and there wrinkled; painted with larger and smaller scattered rays, which are generally broken into irregular lunate or squarish blotches; hinge heavy and strong; hinge plate wide and flat; female shell smaller than the male, more inflated, and swollen at postbasal region.

Animal with the branchiæ rounded below, inner the larger, wholly or partly free from the abdominal sac; marsupium large, projecting far below the inner gills; mantle very thin, dark on its thickened, fringed double edge; anal opening with very small papillæ.

† *PLAGIOLA SECURIS* Lea.

* ? *Obliquaria (Plagiola) depressa* RAFINESQUE, Ann. Gen. Sci. Phys. Brux., 1820, p. 302, pl. XXXI, figs. 5-7.¹

* *Unio securis* LEA, Tr. Am. Phil. Soc., III, 1829, p. 437, pl. XI, fig. 17; * Obs., I, 1834, p. 437, pl. XI, fig. 17.—* DESHAYES, Encyc. Meth., II, 1830, p. 578.—SHORT and EATON, Transylvania JI., 1831, p. 77.—* DESHAYES, An. sans

¹ Rafinesque's name *depressa*, for what is probably a male of this species, has precedence over Lea's name, but it was used previously by Lamarck for an Australian *Unio*. Rafinesque placed this species in his genus *Obliquaria* and the subgenus *Plagiola*, but after giving the name *Obliquaria depressa* he calls it *U. depressa* in parenthesis.

Vert., 2d ed., VI, 1835, p. 550; 3d ed., II, 1839, p. 673.—*HANLEY, Test. Moll., 1842, p. 184; *Biv. Shells, 1843, p. 184, pl. xx, fig. 51.—*CATLOW and REEVE, Conch. Nom., 1845, p. 64.—*CHENU, Ill. Conch., 1858, pl. xvi, figs. 4, 4a, 4b.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—*CHENU, Man., II, 1859, p. 138, fig. 671.—*SOWERBY, Conch. Icon., XVI, 1868, pl. lxi, fig. 304.—*B. H. WRIGHT, Check List, 1888.

**Margarita (Unio) securis* LEA, Syn., 1836, p. 19; 1838, p. 16.

**Margaron (Unio) securis* LEA, Syn., 1852, p. 24; 1870, p. 37.

**Plagiola securis* SMITH, Bull. U. S. Fish Com., 1899, p. 291, pl. lxxx.

**Unio lineolata* SAY, Am. Conch., VI, 1834.

**Plagiola lineolata* AGASSIZ, Arch. für Nat., I, 1852, p. 48.

**Unio lineolatus* CONRAD, New F. W. Shells, 1834, p. 70.—*FERUSSAC, Guer. Mag. 1835, p. 28.—*CONRAD, Am. Conch. (continuation), VII, pl. lxxviii (no date).—

*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—*KUSTER, Conch. Cab. Unio, 1861, p. 171, pl. liii, figs. 1, 2.

Mississippi drainage south into Arkansas; west into eastern Iowa and Kansas; Tombigbee and Alabama river systems.

Subgenus **AMYGDALONAIAS** Crosse and Fischer. 1893.

(Type, *Unio cognatus* Lea.)

Shell inflated, decidedly truncated at the posterior slope, surface slightly concentrically sculptured; posterior ridge sharp and well defined; epidermis shining, sometimes wrinkled looped, and painted with a beautiful pattern of broken or arrow-marked rays; area of the beaks flattened off in the direction of the axis of the shell, but not compressed; beak sculpture delicate, somewhat broken and doubly looped, the anterior loop rounded, the posterior sharp below, the ribs fading out where they cross the posterior ridge; hinge delicate, pseudo-cardinals rather compressed, high, and ragged; hinge plate narrow; female shell very slightly swollen at post base.

Animal with branchiæ more or less free from the abdominal sac; marsupium consisting of numerous distinct ovisacs, and having a well-marked sulcus extending around it at some distance above its base; mantle thickened and doubled on its edge, which has dark papillæ.

(Group of *Plagiola elegans*.)

Shell short, triangular oval, inflated, with a very sharp posterior ridge, extending from the beaks to the hinder point of the shell; male and female shells scarcely distinguishable, both being much expanded in the basal region.

† **PLAGIOLA ELEGANS** Lea.

**Unio elegans* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 83, pl. ix, fig. 13; *Obs., I, 1834, p. 93, pl. ix, fig. 13.—*HANLEY, Test. Moll., 1842, p. 183; *Biv. Shells, 1843, p. 183, pl. xxi, fig. 33.—*CATLOW and REEVE, Conch. Nom., 1845, p. 58.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU, Ill. Conch., 1858, pl. xv, figs. 3, 3a, 3b; *Manual, II, 1859, p. 138, fig. 672.—*SOWERBY, Conch. Icon., XVI, 1868, pl. lxxiv, p. 380.—*CALKINS, Pr. Ottawa Ac. N. Sci., 1874, p. 42.—*B. H. WRIGHT, Check List, 1888.

**Margarita (Unio) elegans* LEA, Syn., 1836, p. 18; 1838, p. 16.

**Margaron (Unio) elegans* LEA, Syn., 1852, p. 23; 1870, p. 36.

* *Plagiola elegans* BAKER, Moll. Chicago, Pt. 1, 1898, p. 91, pl. XXI, fig. 1.

Unio truncatus SAY, Am. Conch., VI, 1834.—* FERUSSAC, Guer. Mag., 1835, p. 27.—

* CONRAD, Am. Conch. (continuation), Pt. 7, pl. LXVII (no date); * Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.—* KUSTER, Conch. Cab., 1861, p. 174, pl. LV, figs. 2, 3.

* *Unio truncatus* RAFINESQUE var. *elegans*. * PÆTEL, Conch. Sam., III, 1890, p. 170.

Mississippi drainage generally; Michigan; Lake Michigan; southwest to Trinity River, Texas.

(Group of *Plagiola donaciformis*.)

Shell small, elongated, triangular oval; posterior ridge generally well defined but rarely sharp, having broken green rays which form chevrons or blotches on a lighter ground; female shell somewhat inflated in the post-basal region.

† *PLAGIOLA DONACIFORMIS* Lea.

* *Unio donaciformis* LEA, Tr. Am. Phil. Soc., III, 1828, p. 267, pl. IV, fig. 3; * Obs., I, 1834, p. 9, pl. IV, fig. 3.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* DESHAYES, An. sans Vert. 2d ed., VI, 1835, p. 547; 3d ed., II, 1839, p. 672.—* HANLEY, Test. Moll., 1842, p. 183; * Biv. Shells, 1843, p. 183, pl. XXII, fig. 41.—* CATLOW and REEVE, 1845, p. 58.—* CHENU, Ill. Conch., 1858, pl. XI, figs. 4, 4a, 4b.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) donaciformis* LEA, Syn., 1836, p. 18; 1838, p. 16.

* *Margaron (Unio) donaciformis* LEA, Syn., 1852, p. 24; 1870, p. 36.

* *Plagiola donaciformis* BAKER, Moll. Chicago, Pt. 1, 1898, p. 92, pl. XIII, fig. 4.

* *Unio zigzag* LEA, Tr. Am. Phil. Soc., III, 1829, p. 440, pl. XII, fig. 19; * Obs., I, 1834, p. 54, pl. XII, fig. 19.—SHORT and EATON, Transylvania Jl., 1831, p. 79.—* DESHAYES, An. sans Vert. 2d ed., VI, 1835, p. 551; 3d ed., II, 1839, p. 673.—* HANLEY, Test. Moll., 1842, p. 183; * Biv. Shells, 1843, p. 183, pl. XXII, fig. 42.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* KUSTER, Conch. Cab. Unio, 1852, p. 42, pl. VIII, fig. 3.—H. and A. ADAMS, Gen. Rec. Moll., 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. VIII, figs. 8, 8a, 8b; Man., 1859, II, p. 138, fig. 670.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIX, fig. 213.—* CALKINS, Pr. Ottawa Ac. Sci., 1874, p. 46.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 172.

* *Margarita (Unio) zigzag* LEA, Syn., 1836, p. 18; 1838, p. 16.

* *Margaron (Unio) zigzag* LEA, Syn., 1852, p. 24; 1870, p. 36.

* *Unio nervosa* CONRAD, New F. W. Shells, 1834, p. 70.

* *Unio nervosus* SAY, Am. Conch., VI, 1834.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.—* KUSTER, Conch. Cab. Unio, 1861, p. 175, pl. LV, fig. 4.

Mississippi drainage generally; Alabama River area; southwest to the Trinity River, Texas; Michigan. Seems to be gradually replaced in the southwest by *P. macrodon*, and there are specimens which can scarcely be assigned with certainty to either species.

† *PLAGIOLA MACRODON* Lea.

* *Unio macrodon* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 154; * Jl. Ac. N. Sci. Phila., VI, 1862, p. 193, pl. XXVI, fig. 262; * Obs., IX, 1863, p. 15, pl. XXVI, fig. 262.—

* ? SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVIII, fig. 437.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

* *Margaron (Unio) macrodon* LEA, Syn. 1870, p. 37.

Eastern Texas, northward probably into Kansas.

†*PLAGIOLA COGNATA* Lea.

- * *Unio cognatus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 306; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 368, pl. LXV, fig. 193; * Obs., VIII, 1860, p. 50, pl. LXV, fig. 193.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIX, fig. 214.—* B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 602.

* *Margaron (Unio) cognatus* LEA, Syn., 1870, p. 43.

Rio Salado, New Leon, Mexico.

Subgenus *PTYCHODERMA* Simpson, 1900.

(Type, *Unio cyrenoides* Philippi.)

Shell triangularly ovate or rounded, solid, bluntly angled before, more sharply angular behind, with a tolerably well-marked posterior ridge; surface strongly and irregularly concentrically sulcate; beaks small but rather prominent, sculptured with fine, irregular, broken ridges, which are somewhat doubly looped, the front loop being larger and more rounded; epidermis olive to tawny, wrinkled, sometimes having faint rays; pseudocardinals compressed, ragged; hinge plate narrow; laterals short, slightly curved, obliquely striated; muscle scars rather shallow, the posterior round; female shell produced in post-basal region.

Animal with small branchiæ, inner united the whole length to the abdominal sac, wider than the outer in front, narrower behind; marsupium occupying the posterior part of the outer gills in from eleven to twenty distinct ovisacs which are rounded and dark below, the whole having a decided, parallel sulcus inside and out near its base; mantle thin, papery, with a wide, thickened, double edge, which is sometimes crenate; branchial opening large fringed; anal opening smooth or with only the faintest crenulations.¹

(Group of *Plagiola cyrenoides*.)

Shell generally more or less covered with radiating grooves which cut up the concentric sculpture into looped wrinkles or nodules, sometimes having the appearance of dried, wrinkled paint, the sculpture extending into the substance of the shell; epidermis usually tawny or yellowish, rarely showing any vestiges of rays; female shell greatly produced in the post-basal region.

†*PLAGIOLA CYRENOIDES* Philippi.

- * *Unio cyrenoides* PHILIPPI, Zeits. für Mal., IV, 1847, p. 93; * Abbild., III, 1848, p. 49, pl. v, fig. 1.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* KUSTER, Conch. Cab. Unio., 1862, p. 285, pl. xcvi, fig. 1.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 150.

* *Margaron (Unio) cyrenoides* LEA, Syn., 1852, p. 25; 1870, p. 38.

¹ In a female *P. cyrenoides*, which was not gravid, the sulcus near the base of the inside of the marsupium was quite deep, in fact the part above it hung over it in a sort of flap, which would no doubt disappear if it was filled with ova.

*† *Unio newcombianus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 103; *Obs., VI, 1857, p. 32, pl. XXX, fig. 27; *Jl. Ac. N. Sci. Phila., III, 1858, p. 312, pl. XXX, fig. 27.—*B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) newcombianus* LEA, Syn., 1860, p. 36.

* *Unio sagrinatus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXVII, fig. 345.1—PÄTEL, Conch. Sam., III, 1890, p. 166.

Nicaragua; Venezuela? A specimen in the U. S. National Museum from the Morelet collection is marked with the latter locality.

† *PLAGIOLA ENCARPA* Lea.

* *Unio encarpus* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 95; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 294, pl. XLII, fig. 105; *Obs., XII, 1869, p. 54, pl. XLII, fig. 105.—*B. H. WRIGHT, Check List, 1888.

Margaron (Unio) encarpus LEA, Syn., 1870, p. 35.

Nicaragua.

(Group of *Plagiola nicaraguensis*.)

Shell much like that of *cyrenoides*, but simply covered with coarse, irregular, concentric ridges, and having no radiating grooves or wrinkles, faintly rayed in umbonal region. Animal as in *Cyrenoides* group.

† *PLAGIOLA NICARAGUENSIS* Lea.

* *Unio nicaraguensis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 95; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 296, pl. XLIII, fig. 107; *Obs., XII, 1869, p. 56, pl. XLIII, fig. 107.—*B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) nicaraguensis* LEA, Syn., 1870, p. 35.

* *Unio gabbianus* LEA, Pr. Ac., N. Sci. Phila., XII, 1868, p. 95; *Jl. Ac., N. Sci. Phila., VI, 1868, p. 295, pl. XLIII, fig. 106; *Obs., XII, 1869, p. 55, pl. XLIII, fig. 106.—*B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) gabbianus* LEA, Syn., 1870, p. 35.

Nicaragua.

The following species are unfigured and indeterminable.

* *Unio nuculinus* PHILIPPI, Zeits. für Mal., V, 1848, p. 176.

Nicaragua. Probably near *cyrenoides*.

* *Unio (Plagiola?) imbricatus* MÖRCH, Mal. Bl., VII, 1861, p. 205.

Nicaragua. This may equal *encarpus* Lea.

DIGENÆ.

Male shell inflated, truncated posteriorly; female shell compressed, expanded into a broad wing behind the post ridge, the wing being filled with a flap of the mantle.

Genus *TRITOGONIA* Agassiz, 1852.

Tritogonia AGASSIZ, Arch. für Nat., 1852, p. 48.

Shell solid, elongate rhomboid, having a strong, irregular posterior ridge, obliquely truncated behind in the male, in the female this region is somewhat compressed and rounded; base incurved; whole surface

¹ Changed in errata by the author to *U. newcombianus*.

except the rounded wing of the females covered with pustules; beaks rather low, incurved and turned forward over the well-developed lunule, which is elongated and filled with epidermal matter; beak sculpture strong, consisting of irregular, subparallel ridges which are curved upward behind, and fine radiating ridges in front of and behind this; epidermis dark olive; hinge plate rather narrow; pseudocardinals strong, ragged; laterals long and straight, near to the pseudocardinals; adductor scars shallow; cavity of the beaks rather deep and compressed, female shell more compressed than that of the male.

Animal with the inner gills much larger than the outer, generally free for the most part from the abdominal sac; palpi enormous, elongated, united to each other behind, and to the mantle a part of their length; mantle thin, with a thickened, dark, double border, the inner edge often toothed throughout, the base much thickened at the posterior end and folded at the branchial opening; branchial opening large, with numerous crowded papillæ; anal opening smooth or with only fine denticulations; superanal opening long, closed below; in the female there is a thickened flap of the mantle which fills the circular posterior expansion of the shell, and which has a smaller flap inside; foot and abdominal sac large, the latter winged in front.¹

† TRITOGONIA TUBERCULATA Barnes.

* *Unio tuberculatus* BARNES, Am. Jl. Sci., VI, 1823, p. 125, pl. VII, figs. 8a 8b.²—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 282.—SHORT and EATON, Transylvania Jl., 1831, p. 76.—* REEVE, Conch. Syst., I, 1841, p. 118, pl. LXXXVIII, fig. 5.—* HANLEY, Test. Moll., 1842, p. 182; *Biv. Shells, 1843, p. 182, pl. xx, fig. 27.—* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 158, pl. LX, fig. 1.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* REEVE, Elements of Conch., 1860, II, pl. XXXIII, fig. 183.—* CALKINS, Pr. Ottawa Acad. Sci., 1874, p. 45.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 170.

* *Margarita (Unio) tuberculatus* LEA, Syn., 1836, p. 17; 1838, p. 16.

* *Margaron (Unio) tuberculatus* LEA, 1852, p. 23; 1870, p. 34.

* *Mya tuberculata* EATON, Zool. Text-Book, 1826, p. 217.

* *Unio pustulata* SWAINSON, Treat. on Mal., 1840, p. 271, fig. 54d.

* *Unio gigas* SOWERBY,³ Conch. Icon., XVI, 1867, pl. LVI, fig. 287.

¹ I have never seen a specimen among the hundreds examined that had young or ova in the gills. But I have seen spermatozoa in specimens with inflated shells and ova in those with the wing. Mr. H. M. Kelly, of Mount Vernon, Iowa, who has made a very careful study of the anatomy of many of our Unionideæ, assures me that the form with the compressed shell, having the expanded flap behind, is the female.

² It has been claimed that this species was previously described under the name of *Obliquaria verrucosa* by Rafinesque, but I am unable to make anything out of his figure or description; hence I use the name given by Barnes. Barnes's name was used by Rafinesque for the species which the former called *Unio verrucosus*, but as Rafinesque placed his species in *Obliquaria*, and as the generic names of both forms are changed in this work, I do not consider Barnes's name preoccupied.

³ I think that this is a large, rather high male of the species in question. I can not find any description of *Unio gigas* by Swainson, to whom Sowerby credits this species, and the *Megadomus gigas* of Swainson (which, so far as I know, is undescribed) can not be it, as the latter says that his genus, *Megadomus*, has the lateral teeth imperfect, one or none. Treatise on Mal., p. 266.

* *Unio verrucosus* SAY, Am. Conch., VI, 1834. —* CONRAD, New F. W. Shells, 1834, p. 72. —* FERUSSAC, Guer. Mag., 1835, p. 27. —* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 259. —* KUSTER, Conch. Cab., 1862, p. 273, pl. XCII, fig. 1. —* SOWERBY, Conch. Icon., XVI, 1868, pl. I, fig. 4.

* *Tritogonia verrucosa* AGASSIZ, Arch. für. Nat., I, 1852, p. 48.

* *Tritogonia verrucosa* SMITH, Bull. U. S. Fish Com., 1899, p. 291, pl. LXXXI.

† TRITOGONIA TUBERCULATA var. OBESA Simpson.¹

Mississippi drainage generally; streams falling into the Gulf of Mexico from the Alabama system west to central Texas.

† TRITOGONIA CONJUGANS B. H. Wright.

* *Unio conjugans* B. H. WRIGHT, Naut., XIII, 1899, p. 89.

* *Tritogonia conjugans* Simpson, Pr. Ac. N. Sci. Phila., 1900, p. 79, pl. IV, fig. 1.

Hiawassee River, Tennessee. A single male shell of this remarkable species, the type, is in the U. S. National Museum collection. I am not positive that it belongs here, though it probably does.

MESOGENÆ.

Male and female shells alike, short, solid, inflated; embryos occupying a few distinct ovisacs in the center of the outer gills.

Genus CYPROGENIA Agassiz, 1852.

(Type, *Unio irroratus* Lea.)

Cyprogenia AGASSIZ, Arch. für Nat., 1852, p. 47.

Shell solid, inflated, rounded triangular, sometimes slightly retuse, generally a little biangular behind; posterior ridge unusually well developed, especially in the young shell; umbonal region flattened parallel with the axis of the shell, sometimes compressed; beaks curved inward and forward, their sculpture very faint, consisting of slightly doubly looped ridges; sculpture of the shell nodular, radiately wrinkled, or lachrymose; ligament black and conspicuous; lunule distinct and well developed; epidermis shining, painted with a delicate, dark mottling on a light ground; hinge plate wide and flat; pseudocardinals heavy, triangular, blunt and ragged; laterals short, obliquely striated; cavity of the beaks not deep; adductor scars small, well impressed, those at the posterior round; nacre bright and silvery. Animal with inner gills partly free from the abdominal sac, rounded below; outer gills smaller; marsupium consisting of from seven to twenty-three very long, purple ovisacs pendent from near the central base of the outer gills, and formed into a close coil with the ends turned inwardly; branchial opening large with many small papillæ; anal opening smooth.

¹A short, inflated, peculiarly formed shell from Big Eddy, Nechez River, Texas, having very few faint tubercles is in the U. S. National Museum collection, and may be a distinct species. It is certainly worthy of the varietal name here applied.

† CYPROGENIA IRRORATA Lea.

- * *Unio irroratus* LEA, Tr. Am. Phil. Soc., III, 1830, p. 269, pl. v, fig. 5; * Obs., I, 1834, p. 11, pl. v, fig. 5.—SHORT and EATON, Transylvania Journ., 1831, p. 77.—
 * HANLEY, Test. Moll., 1842, p. 181; * Biv. Shells, 1843, p. 181, pl. xx, fig. 25.—
 * CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* CHENU, Ill. Conch., 1858, pl. x, figs. 3, 3a, 3b.—* REEVE, Conch. Icon., XVI, 1864, pl. xii, fig. 44.—* B. H. WRIGHT, Check List, 1888.
 * *Margarita (Unio) irroratus* LEA, Syn., 1836, p. 16; 1838, p. 15.
 * *Margaron (Unio) irroratus* LEA, Syn., 1852, p. 22; 1870, p. 34.
 * *Unio irrorata* DESHAYES, Encyc. Meth., II, 1830, p. 579.—* DESHAYES, An. sans. Vert., 2d ed., VI, 1835, p. 547; 3d ed., II, 1839, p. 672.
 * *Theliderma irrorata*, SWAINSON, Tr. on Mal., 1840, p. 271, fig. 54a.
 * *Unio verrucosus albus* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 281.
 * *Unio stegarius* SAY, Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 71.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* CONRAD, Monog., X, 1838, p. 83, pl. XLVI, fig. 1; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* REEVE, Conch. Icon., XVI, 1864, pl. xi, fig. 45.
 * *Unio stegarius* var. *irroratus* PÆTEL, Conch. Sam., III, 1890, p. 168.

† CYPROGENIA IRRORATA var. PUSILLA Simpson.¹

Ohio, Cumberland, and Tennessee river systems; St. Francis and Saline rivers, Arkansas? Eastern Iowa? The specimens reported from the western localities may be *C. aberti*.

† CYPROGENIA ABERTI Conrad.

- * *Unio aberti* CONRAD, Pr. Ac. N. Sci. Phila., V, 1850, p. 10; * VI, 1853, p. 244; Jl. Ac. N. Sci. Phila., 1854, p. 295, pl. xxvi, fig. 1.—* B. H. WRIGHT, Check List, 1888.
 * *Margaron (Unio) aberti* LEA, Syn. 1870, p. 34.
 * *Unio lamarckianus* LEA, Tr. Am. Phil. Sci., X, 1852, p. 266, pl. xvii, fig. 20; * Obs., V, 1852, p. 22, pl. xvii, fig. 20.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* SOWERBY, Conch. Icon., XVI, 1868, pl. xcii, fig. 498.—* PÆTEL, Conch. Sam., III, 1890, p. 156.
 * *Margaron (Unio) lamarckianus* LEA, Syn., 1852, p. 23.
 * *Unio popenoi* CALL, Bull. Washb. Coll., I, 1885, p. 49, pl. ii.

Kansas; Missouri; Indian Territory; Arkansas.

Genus OBLIQUARIA (Rafinesque 1820) Simpson

(Type, *Obliquaria reflexa* Rafinesque.)

Obliquaria RAFINESQUE, Ann. Gen. Sci. Phys. Brux., 1820, p. 301.

Shell inflated, solid, oval, ending in a tolerably sharp point behind, having a row of large, compressed, longitudinal knobs running from the beaks to the center of the base, those of one valve alternating in position with the knobs of the other, and a well-developed posterior ridge, the space between the ridge and knobs somewhat excavated; posterior slope and sometimes the entire shell more or less corrugately sculptured; beaks prominent, incurved and pointed slightly forward toward a tolerably well-developed lunule; beak sculpture strong, con-

¹A peculiar, dwarfed variety, about one-half the size of the ordinary form and much inflated, from the Green River, Kentucky.

sisting of four or five heavy, parallel ridges, which fall low in front but are curved upward behind; epidermis smooth, generally shining, painted with numerous delicate, wavy, darker, broken rays; pseudo-cardinals strong, direct, and ragged; laterals short, nearly straight; anterior muscle scars small, the sides of the pit smooth, the bottom ragged; front part of the shell very solid, suddenly becoming rather thin just behind the knobs; male and female shells essentially alike.

Animal with small branchiæ, rounded below, inner the larger, free from the abdominal sac in part; marsupium consisting of a few distinctly marked ovisacs (4 to 7), occupying a position just behind the center of the outer gills, projecting far below the rest of the branchiæ, their bases rounded; mantle cut away at the thinner portion of the shell; anal opening smooth or having only minute crenulations.

† OBLIQUARIA REFLEXA Rafinesque.

* *Obliquaria (Quadrula) reflexa* RAFINESQUE, Ann. Gen. Sci. Phys., 1820, p. 306.—

* CHENU, Bib. Conch., 1st Ser., III, 1845, p. 19.

* *Obliquaria reflexa* BAKER, Moll. Chicago, Pt. 1, 1898, p. 89, pl. XIV, fig. 5; XX, fig. 2.

* *Unio reflexus* SAY, Am. Conch., No. VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 71; * Monog., I, 1835, p. 7, pl. IV, fig. 1.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* KUSTER, Conch. Cab. Unio, 1852, p. 52, pl. XI, fig. 2.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 256.—* REEVE, Conch. Icon., XVI, 1864, pl. VI, fig. 23.

* *Unio cornutus* BARNES,¹ Am. J. Sci., VI, 1823, p. 122, pl. IV, figs. 5, 5a, 5b, 5c.—* ? ANTON, Verz. der Conch., 1839, p. 14.—* HANLEY, Test. Moll., 1842, p. 179; Biv. Shells, 1843, p. 179, pl. XX, fig. 30.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CALKINS, Pr. Ottawa Ac. N. Sci., 1874, p. 41.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) cornutus* LEA, Syn., 1836, p. 15; 1838, p. 15.

* *Margaron (Unio) cornutus* LEA, Syn., 1852, p. 22; 1870, p. 33.

* *Mya cornuta* EATON, Zool. Text-book, 1826, p. 216.

* *Theliderma cornuta* SWAINSON, Treat. on Mal., 1840, p. 269.

Unio torulosus SHORT and EATON, Transylvania J., 1831, p. 75.

*†? *Unio philippsi* CONRAD,² Monog., I, 1835, p. 9, pl. V, fig. 1; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—* P. TEL, Conch. Sam., III, 1890, p. 163.

* *Unio phillipsii* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* KUSTER, Conch. Cab. Unio, 1852, p. 50, pl. X, fig. 3.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) phillipsii* LEA, Syn., 1836, p. 15; 1838, p. 15.

* *Margaron (Unio) phillipsii* LEA, Syn., 1852, p. 22; 1870, p. 33.

* *Unio philipsii* HANLEY, Test. Moll., 1842, p. 178; Biv. Shells, 1843, p. 178.

Mississippi drainage; Michigan; Alabama River; southwestwardly to central Texas.

¹ I am sorry to have to place the very appropriate and well-known name of Barnes in the synonymy, but I believe that the description of Rafinesque for his *Obliquaria reflexa* defines this species and nothing else, and although it is not figured, I feel justified in using it.

² A singular shell having the general form of *reflexa*, but the knobs are much broken up. I have seen specimens which agree quite well with Conrad's fine figure, which I have no doubt represents an unusual form of Rafinesque's species, and they hardly seem to me to be a variety. I do not think the so-called type in the Philadelphia Academy of Natural Sciences is valid.

PTYCHOGENÆ.

Male and female shells essentially alike, embryos contained in distinct ovisacs with rounded bases, occupying the entire outer gills, which, when gravid, consist of a series of folds.

Genus *PTYCHOBANCHUS* Simpson, 1900.

(Type, *Unio phaseolus* Hildreth.)

Ptychobanchus SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 79.

Shell triangular, solid, sometimes becoming arcuate in old specimens, umbonal region rather elevated; beak sculpture consisting of faint, somewhat broken ridges which have a tendency to be doubly looped; posterior ridge rounded but well developed; epidermis usually painted with wavy hair line rays or broken, radiating bars, which show a tendency to form square spots; hinge plate rather wide and flat; pseudo-cardinals small, low, triangular and roughened; laterals club-shaped, remote; cavity of the beaks shallow; muscle scars rather deep. Animal with inner gills free all or part of their length from the abdominal sac; marsupium occupying the basal half of the whole length of the outer gills and hanging in from six to twenty beautiful folds; ovisacs distinct, each ending below in an enlarged, rounded bulb which has a colored spot in its center; mantle thin, with a dark, thickened border; branchial opening large, with very minute papillæ or crenulations, sometimes smooth; anal opening crenulate or smooth.

† *PTYCHOBANCHUS PHASEOLUS* Hildreth.

* *Unio phaseolus* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 283.¹—SAY, Am. Conch., No. III, 1830, pl. XXII.—SHORT and EATON, Transylvania Jl., 1831, p. 77.—* HANLEY, Test. Moll., 1842, p. 207; Biv. Shells, 1843, p. 207, pl. xx, fig. 50.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 31, pl. ix, figs. 1-6.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIII, fig. 378.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 163.

* *Margarita (Unio) phaseolus* LEA, Syn., 1836, p. 38; 1838, p. 25.

* *Margaron (Unio) phaseolus* LEA, Syn., 1852, p. 38; 1870, p. 61.

† * *Unio planulatus* LEA, Tr. Am. Phil. Soc., III, 1830, p. 431, pl. ix, fig. 13; * Obs., I, 1834, p. 45, pl. ix, fig. 13.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 549.—* CHENU, Ill. Conch., 1858, pl. XIII, figs. 1, 1a, 1b; * Manual, 1859, II, p. 137, fig. 659.

* *Unio planulata* DESHAYES, An. sans Vert., 3d ed., II, 1839, p. 672.

* *Unio fasciolaris* CONRAD, New F. W. Shells, 1834, p. 69.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* AGASSIZ, Arch. für Naturg., I, 1852, p. 50.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—* KUSTER, Conch. Cab., 1861, p. 172, pl. LIV, figs. 1-4.

* *Unio camelus* LEA,² Tr. Am. Phil. Soc., V, 1834, p. 102, pl. xv, fig. 45; * Obs., I,

¹ Figure left out by the editor.

² Typically this form is quite distinct from the ordinary manifestation of the species, being very solid and much humped, but there seems to be every possible variation to the lighter, more compressed forms, and I do not think it is worthy of a varietal name.

1834, p. 214, pl. xv, fig. 45.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 184; * Biv. Shells, 1843, p. 184, pl. XXI, fig. 54.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* SOWERBY, Conch. Icon., XVI, 1867, pl. LV, fig. 283.—* B. H. WRIGHT, Check List, 1888.—PÆTEL, Conch. Sam., III, 1890, p. 147.

* *Margarita (Unio) camelus* LEA, Syn., 1836, p. 19; 1838, p. 17.

* *Margaron (Unio) camelus* LEA, Syn., 1852, p. 24; 1870, p. 37.

* *Unio compressissimus* LEA,¹ Pr. Am. Phil. Soc., IV, 1845, p. 163; * Tr. Am. Phil. Soc., X, 1848, p. 81, pl. VIII, fig. 23; * Obs., IV, 1848, p. 55, pl. VIII, fig. 23.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Margaron (Unio) compressissimus* LEA, Syn., 1852, p. 24; 1870, p. 37.

Ohio, Tennessee, and Cumberland river systems; peninsula of Michigan; Kansas; Arkansas; Indian Territory; Louisiana. It is quite probable that some of the specimens reported from the southwestern part of the range belong to the next species.

† PTYCHOBANCHUS CLINTONENSIS Simpson.

* *Ptychobanchus clintonensis* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 79, pl. v, fig. 3.

Little Red River, Arkansas; Indian Territory(?); southwest Tennessee(?)

† PTYCHOBANCHUS FOREMANIANUS Lea.

* *Unio foremanianus* LEA, Pr. Am. Phil. Soc., II, 1842, p. 224; * Tr. Am. Phil. Soc., VIII, 1842, p. 247, pl. XXVII, fig. 64; * Obs., III, 1842, p. 85, pl. XXVII, fig. 64.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XXVI, figs. 1, 1a, 1b.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVII, fig. 202.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 153.

* *Margaron (Unio) foremanianus* LEA, Syn., 1852, p. 23; 1870, p. 36.

* *Unio woodwardius* LEA,² Pr. Ac. N. Sci. Phila., IX, 1857, p. 170.

* *Unio woodwardianus* LEA, Jl. Ac. N. Sci. Phila., IV, 1859, p. 199, pl. XXIII, fig. 82; * Obs., 1859, p. 17, pl. XXIII, fig. 82.—* REEVE, Conch. Icon., XVI, 1864, pl. XVI, fig. 73.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 172.

Margaron (Unio) woodwardianus LEA, Syn., 1870, p. 36.³

* *Unio velatus* CONRAD, Jl. Ac. N. Sci. Phila., 1853, p. 298, pl. XXVII, fig. 6; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.

Alabama River drainage.

¹ The type is simply an old, stunted *phaseolus*, with nearly every vestige of epidermis eroded away.

² This is the name originally given by Dr. Lea. He afterward changed it to *woodwardianus*.

³ I formerly thought *foremanianus* and *woodwardianus* distinct, but a large amount of material received from Mr. R. E. Call, from the Cahawba River, Alabama, shows that the two absolutely blend together, there being a great diversity of form, and some specimens show the capillary rays of *foremanianus* and the square spots of *woodwardianus*, while others are of a uniform tawny color.

† PTYCHOBANCHUS TRINACRUS Lea.

* *Unio trinacrus* LEA, Pr. Ac. N. Sci., Phila., V, 1861, p. 59; * Jl. Ac. N. Sci., Phila., V, 1862, p. 86, pl. XII, fig. 235; * Obs., VIII, 1862, p. 90, pl. XII, fig. 235.—

* SOWERBY, Conch. Icon., XVI, pl. LXXXI, fig. 428.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) trinacrus* LEA, Syn., 1870, p. 36.

Coosa River, Alabama.

† PTYCHOBANCHUS GREENI Conrad.

* *Unio greeni* CONRAD, New F. W. Shells, 1834, p. 32, pl. IV, fig. 1, p. 69.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen., 1836, p. 203.—* CONRAD, Monog., VIII, 1837, p. 69, pl. XXXVIII, fig. 2.—* HANLEY, Test. Moll., 1842, p. 191; * Biv. Shells, 1843, p. 191, pl. XXIV, fig. 6.—* CHENU, Bib. Conch., 1st ser., 1845, p. 17, pl. III, fig. 5.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLVII, fig. 253.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 154.

* *Margarita (Unio) greeni* LEA, Syn., 1836, p. 24; 1838, p. 19.

* *Margaron (Unio) greeni* LEA, Syn., 1852, p. 27; 1870, p. 42.

† * *Unio simplex* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 163; * Tr. Am. Phil. Soc., X, 1848, p. 76, pl. v, fig. 15; * Obs., IV, 1848, p. 50, pl. v, fig. 15.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 167.

* *Margaron (Unio) simplex* LEA, Syn., 1852, p. 27; 1870, p. 42.

Black Warrior River, Alabama.

† PTYCHOBANCHUS FLAVESCENS Lea.

* *Unio flavescens* LEA, ¹ Pr. Am. Phil. Soc., IV, 1845, p. 163; Tr. Am. Phil. Soc., X, 1848, p. 72, pl. III, fig. 9; * Obs., IV, 1848, p. 46, pl. III, fig. 9.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) flavescens* LEA, Syn., 1852, p. 27; 1870, p. 43.

* *Unio flavescens* PÆTEL, Conch. Sam., III.

Black Warrior River, Alabama.

ESCHATIGENÆ.

Male and female shells alike; embryos in distinct, short ovisacs, forming a border to the entire outer gills.

Genus DROMUS Simpson, 1900.

(Type, *Unio dromus* Lea.)

Shell solid, rounded-triangular; beaks well forward, rather high; beak sculpture consisting of fine ridges running parallel with the growth lines, the furrows between the ridges interrupted at the posterior ridge; posterior ridge distinct; a series of humps runs from the beaks down to the central part of the base of the shell, which is otherwise sculptured by

¹ I have only seen a single shell of this, and it is not in the best condition. It may be only a form of *greeni*, but it is not so inflated or solid, and it has no rays.

irregular concentric ridges; epidermis beautifully painted by undulated, radiating, broken hair lines or fine maculations; hinge plate wide and flat; pseudocardinals triangular, small and low, ragged; laterals low, short, club-shaped; cavity of the beaks deep and compressed, dorsal muscle scars on the under side of the hinge shelf; adductor scars small; front part of the shell very thick, suddenly becoming thinner at the row of humps; male and female shells much alike.

Animal with the inner branchiæ much the larger, curved below, free from the abdominal sac; marsupium occupying the base of nearly the whole outer branchiæ in numerous rather indistinctly marked ovisacs, bases of the ovisacs slightly rounded; ova very minute; palpi small, curved; branchial opening very large, extending one-third of the way along the base, fringed with many brown papillæ; anal opening very large, finely crenulated.

† DROMUS DROMUS Lea.

- * *Unio dromus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 70, pl. x, fig. 29; * Obs., I, 1834, p. 182, pl. x, fig. 29.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* CONRAD, Monog., X, 1838, p. 84, pl. XLVI, fig. 2.—* HANLEY, Test. Moll., 1842, p. 181; * Biv. Shells, 1843, p. 181, pl. XXIII, fig. 25.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* KUSTER, Conch. Cab. Unio, 1861, p. 172, pl. LIII, figs. 3, 4.—* REEVE, Conch. Icon., XVI, 1864, pl. VIII, fig. 29.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

* *Margarita (Unio) dromus* LEA, Syn., 1836, p. 16; 1838, p. 16.

* *Cyprogenia dromus* AGASSIZ, Arch. für Naturg., I, 1850, p. 48.

* *Margaron (Unio) dromus* LEA, Syn., 1852, p. 23; 1870, p. 34.

Tennessee and Cumberland River systems.

† DROMUS CAPERATUS. Lea.¹

- * *Unio caperatus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; * Tr. Am. Phil. Soc., X, 1848, p. 75, pl. v, fig. 14; * Obs., IV, 1848, p. 75, pl. v, fig. 14.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* REEVE, Conch. Icon., XVI, 1864, pl. v, fig. 19.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 147.

* *Margaron (Unio) caperatus* LEA, Syn., 1852, p. 22; 1870, p. 34.

* *Unio abacoides* HALDEMAN,² Pr. Ac. N. Sci. Phila., III, 1846, p. 75; * Am. Jl. of Sci., II, 1846 (September), p. 274.

Tennessee River system.

DIAGENÆ.

Male and female shells alike; embryos contained in the outer gills in short ovisacs which run crosswise of the branchiæ, and are discharged entire into the water.

¹ Generally this form is more compressed and less humped than *D. dromus*, but I think it is probably only a variety of the latter.

² Haldeman only gave a brief description of his species, and it has never been figured. According to Lea it equals his *U. caperatus*, and I believe he is correct.

Genus STROPHITUS Rafinesque, 1820.

(Type, *Anodonta undulata* Say.)*Strophitus* RAFINESQUE, Ann. Gen. Sci. Phys. Brux., 1820, p. 316.*Uniopsis* AGASSIZ, Arch. für Naturg., 1852, p. 49.

Shell elliptical to rhomboid, inflated, subsolid, pointed or biangulate behind, with a low posterior ridge, which is sometimes double; beaks full, sculpture consisting of a few strong, concentric ridges, which curve sharply upward behind; epidermis rayed or rayless, shining; hinge line incurved in front of the beaks; teeth rudimentary, a vestigial, compressed tooth in each valve, and sometimes a secondary tooth; laterals rarely present; muscle scars shallow.

Animal with the marsupium occupying the whole of the outer gills, consisting of short, horizontal ovisacs which run directly across the gills, and are discharged through the outer wall with the ovules in them; ovules ten to twenty-five in each ovisac, in one or two rows; inner gills the larger, free in part from the abdominal sac, or wholly united; mantle generally bordered behind with square, black spots; branchial opening with numerous papillæ; anal opening papillose or crenulate.¹

† STROPHITUS EDENTULUS Say.

Alasmodonta edentula SAY, N. Harm. Diss., II, No. 22, 1829, p. 340.—* CONRAD, New F. W. Shells, 1834, p. 72.—* L. W. SAY, Terr. and Fluv. Shells, 1840, p. 10.

* *Anodonta edentula* FERUSSAC, Guer. Mag., 1835, p. 25.—* HANLEY, Test. Moll., 1842, p. 218.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CALKINS, Pr. Ottawa Ac., 1874, p. 50.—* CLESSIN, Conch. Cab. Ano., 1873, p. 107, pl. XXX, figs. 5, 6.—* LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 55.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

* *Margarita (Anodonta) edentula* LEA, Syn., 1836, p. 450; 1838, p. 30.

* *Anodon edentula* CATLOW and REEVE, Conch. Nom., 1845, p. 66.—* DE KAY, Zool. N. Y., Pt. 5, 1848, p. 201, pl. XVI, fig. 231.—* HARTMAN and MICHENER, Conch. Cest., 1874, p. 95, fig. 174.

* *Uniopsis edentula* AGASSIZ, Arch. für Naturg., I, 1852, p. 49.

* *Margaron (Anodonta) edentula* LEA, Syn., 1852, p. 49; 1870, p. 79.

* *Strophitus edentulus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.—* BAKER, Moll. Chicago, Pt. 1, 1898, p. 67, pl. XII, figs. 5, 6; XVII, fig. 3.

* *Anodon rugosus* SWAINSON, Zool. Ill., 1st ser., II, pl. xcvi, 1822.

* *Anodonta rugosus* HANLEY, Biv. Shells, 1843, p. 217.

* *Anodonta undulata* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 290.—* CLESSIN, Conch. Cab. Ano., 1873, p. 79, pl. XXVII, figs. 3, 5.

* *Anodon undulata* GOULD, Inv. of Mass., 1841, p. 120, fig. 79; 1870, p. 182, fig. 482.

* *Unio undulata* DESHAYES, Tr. Elem., II, 1853, p. 217, pl. XXX, figs. 8, 9.

* *Anodonta undulatus* SOWERBY, Rich. Faun. Bor. Am., III, 1836, p. 316.—* SOWERBY, Conch. Icon., XVII, 1867, pl. XI, fig. 30.

* *Anodon areolatus* SWAINSON, Zool. Ill., 2d ser., I, 1829, pl. XVIII.

* *Anodonta areolatus* FERUSSAC, Guer. Mag., 1835, p. 25.

¹In this remarkable group of mollusks the changes that take place in the gills when they are used as a marsupium only last as long as they are gravid. When the short ovisacs are discharged the gill becomes an ordinary gill again.

- * *Anodon marginata* FERUSSAC, Guer. Mag., 1835, p. 25.
 * *Anodonta ferussaciana* FERUSSAC, Guer. Mag., 1835, p. 25.
 * † *Margarita (Anodonta) wardiana* LEA, Syn., 1836, p. 50; 1838, p. 30.
 * *Anodonta wardiana* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 46, pl. xiv, fig. 42; * Obs., II, 1838, p. 46, pl. xiv, fig. 42.—* TROSCHER, Arch. für Nat., V, 1839, Pt. 2, p. 238.—* HANLEY, Test. Moll., 1842, p. 220; * Biv. Shells, 1843, p. 220.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* PÆTEL, Conch. Sam., III, 1890, p. 186.—* CATLOW and REEVE, Conch. Nom., 1845, p. 68.—* B. H. WRIGHT, Check List, 1888.
 * *Anodon wardiana* SOWERBY, Conch. Icon., XVII, 1867, pl. xxviii, fig. 114.
 * *Margaron (Anodonta) wardiana* LEA, Syn., 1852, p. 49; 1870, p. 79.
 * *Anodonta virgata* CONRAD, Cover of Mon. No. 5, 1836.
 * *Strophitus virgatus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.
 * *Anodon unadilla* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 199, pl. xv, fig. 228.
 * *Strophitus unadilla* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.
 * *Anodon undulata* GOULD, Inv. Mass., 1841, p. 120, fig. 79; * 1870, p. 182, fig. 79.
 * † *Anodonta tetragona* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 165; * Tr. Am. Phil. Soc., X, 1848, p. 82, pl. viii, fig. 25; * Obs., IV, 1848, p. 56, pl. viii, fig. 25.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* B. H. WRIGHT, Check List, 1888.
 * *Margaron (Anodonta) tetragona* LEA, Syn., 1852, p. 51; 1870, p. 81.
 * *Strophitus tetragona* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.
 * † *Anodonta arkansasensis* LEA, Tr. Am. Phil. Soc., XI, 1852, p. 293, pl. xxix, fig. 56; * Obs., V, 1852, p. 49, pl. xxix, fig. 56.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 176.
 * *Margaron (Anodonta) arkansasensis* LEA, Syn., 1852, p. 50; 1870, p. 80.
 * *Strophitus arkansasensis* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 262.
 * † *Anodonta shæfferiana* LEA, Tr. Am. Phil. Soc., X, 1852, p. 288, pl. xxvi, fig. 50; * Obs., V, 1852, p. 44, pl. xxvi, fig. 50.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* PÆTEL, Conch. Sam., III, 1890, p. 184.
 * *Margaron (Anodonta) shæfferiana* LEA, Syn., 1852, p. 51; 1870, p. 81.
 * *Strophitus shæfferiana* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.
 * *Anodonta shæfferiana* CLESSIN, Conch. Cab. Ano., 1853, p. 243, pl. xvii, figs. 5-7.
 * *Anodon shæfferianus* SOWERBY, Conch. Icon., XVII, 1870, pl. xxxv, fig. 143.
 * *Anodonta shæfferiana* B. H. WRIGHT, Check List, 1888.
 * † *Anodonta showalterii* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 307; * Jl. Ac. N. Sci. Phila., 1862, p. 215, pl. xxxiii, fig. 284; * Obs., IX, 1863, p. 37, pl. xxxiii, fig. 284.—* CLESSIN, Conch. Cab. Ano., 1874, p. 156, pl. li, figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 184.
 * *Anodon showalterii* SOWERBY, Conch. Icon., XVII, 1870, pl. xxvii, fig. 104.
 * *Margaron (Anodonta) showalterii* LEA, Syn., 1870, p. 79.
 * *Alasmodon rhombica* ANTHONY, Am. Jl. Conch., I, 1865, p. 158, pl. xii, fig. 5.
 * *Margaritana rhombica* PÆTEL, Conch. Sam., III, 1890, p. 173.
 * *Anodonta salmonea* CLESSIN, Conch. Cab. Ano., 1873, p. 91, pl. xxiv, figs. 1, 2.

† STROPHITUS EDENTULUS var. PAVONIUS Lea.

- * *Anodonta pavonia* LEA, Tr. Am. Phil. Soc., VI, 1836, p. 78, pl. xxi, fig. 65; * Obs., II, 1838, p. 78, pl. xxi, fig. 65.—* TROSCHER, Arch. für Nat., V, 1839, Pt. 2, p. 239.—* HANLEY, Test. Moll., 1842, p. 218.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* MUSGROVE, Phot. Conch., 1863, pl. i, fig. 9.—* CLESSIN, Conch. Cab. Ano., 1873, p. 156, pl. xxx, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 183.
 * *Margarita (Anodonta) pavonia* LEA, Syn., 1838, p. 30.
 * *Anodon pavonia* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 203, pl. xl, fig. 358.—

* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXII, fig. 128.

* *Margaron (Anodonta) pavonia* LEA, Syn., 1852, p. 50; 1870, p. 79.

* *Strophitus pavonius* BAKER, Moll. Chicago, Pt. 1, 1898, p. 68, pl. III, fig. 5; v, fig. 3.

* *Anodon annulatus* SOWERBY, Conch. Icon., XVII, 1867, pl. XXVIII, fig. 67.

Entire Mississippi drainage; St. Lawrence system and south in streams draining into the Atlantic to North Carolina; north in the British possessions to Lake Winnipeg; southwest to central Texas; Tyner, Alabama.

† STROPHITUS UNDULATUS Say.¹

Anodonta undulata SAY, Nich. Enc., II, 1816, pl. III, fig. 5.—SHORT and EATON, Transylvania Jl., 1831, p. 81.—* FERUSSAC, Guer. Mag., 1835, p. 25.—* C. B. ADAMS, F. W. and L. S. of Vt., 1842, p. 15.—* HANLEY, Test. Moll., 1842, p. 217; Biv. Shells, 1843, p. 217.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* LATCHFORD, Tr. Ot. F. N. Cl., 1882, p. 55.—* B. H. WRIGHT, Check List, 1888.—* H. CARPENTER, Naut., IV., 1890, p. 57.—* PÆTEL, Conch. Sam., III, 1890, p. 186.

* *Margarita (Anodonta) undulata* LEA, Syn., 1836, p. 50; 1838, p. 30.

* *Anodon undulata* CATLOW and REEVE, Conch. Nom., 1845, p. 68.

* *Alasmodonta undulata* C. B. ADAMS, Thompson's Hist. Vt., 1842, p. 165.

* *Margaron (Anodonta) undulata* LEA, Syn., 1852, p. 49; 1870, p. 79.

* *Strophitus undulatus* STIMPSON, Shells of N. Eng., 1851, p. 15.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.

* *Anodonta pennsylvanica* LAMARCK, An. sans Vert., VI, 1819, p. 86.—* STARK, Elem. Nat. Hist., II, 1828, p. 89.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 567; 3d ed., II, 1839, p. 678.—* DELESSERT, Rec. Coq. Lam., 1841, pl. XIII, figs. 4a, 4b.—* CHENU, Ill. Conch., 1858, pl. III, figs. 5, 5a.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.

* *Anodon papyraceus* ANTHONY, Am. Jl. Conch., I, 1865, p. 161, pl. xv, fig. 2.—* SOWERBY, Conch. Icon., XVII, 1867, pl. XIV, fig. 46.

* *Anodonta papyracea* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 183.

* *Margaron (Anodonta) papyracea* LEA, Syn., 1870, p. 81.

* *Anodon quadriplicatus* SOWERBY, Conch. Icon., XVII, 1867, pl. XXVIII, fig. 110.

Northern New England to Virginia, in streams draining into the Atlantic. This species is reported from Manitoba, but it is probable that the specimens from that region are *S. edentulus*.

† STROPHITUS CONNASAUGAENSIS Lea.

* *Margaritana connasaugaensis* LEA, Pr. Ac. N. Sci. Phila., II, 1857, p. 135; * Jl. Ac. N. Sci. Phila., IV, 1859, p. 229, pl. XXXII, fig. 113; * Obs., VII, 1859, p. 47, pl. XXXII, fig. 113.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 172.

* *Margaron (Margaritana) connasaugaensis* LEA, Syn., 1870, p. 69.

* *Unio connasaughensis* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVIII, fig. 474.

* *Margaritana connasaughensis* CLESSIN, Conch. Cab. An., 1875, p. 269, pl. LXXXI, figs. 7, 8.

¹ Much confusion exists concerning this and the preceding species. The *edentulus* is much the larger and solidier shell; the *undulatus* being a small, thin form (Say's figure is only about half an inch long, from a young or very dwarf specimen), usually quite distinctly biangulate behind. The latter is confined to the Atlantic drainage, while the former is found in this area and in the Mississippi Valley.

*†*Margaritana alabamensis* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 41; *Jl. Ac. N. Sci. Phila., V, 1862, p. 104, pl. xvi, fig. 249; *Obs., VIII, 1862, p. 108, pl. xvi, fig. 249.—*CLESSIN, Conch. Cab. Ano., 1876, p. 264, pl. lxxx, fig. 5.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 172.

**Unio alabamensis* SOWERBY, Conch. Icon., XVI, 1868, pl. lxxxii, fig. 433.

**Margaron (Margaritana) alabamensis* LEA, Syn., 1870, p. 68.

Alabama River system.

†STROPHITUS SPILLMANII Lea.

**Margaritana spillmanii* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 138; *Jl. Ac. N. Sci. Phila., V, 1862, p. 105, pl. xvii, fig. 252; *Obs., VIII, 1862, p. 109, pl. xvii, fig. 252.—*CLESSIN, Conch. Cab. Ano., 1876, p. 264, pl. lxxxii, figs. 5, 6.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 174.

**Unio spillmanii* SOWERBY, Conch. Icon., XVI, 1868, pl. lxxxii, fig. 435.

Tombigbee River, Mississippi.

†STROPHITUS TOMBIGBEENSIS Lea.

**Margaritana tombigbeensis* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 138; *Jl. Ac. N. Sci. Phila., V, 1862, p. 107, pl. xviii, fig. 255; *Obs., VIII, 1862, p. 111, pl. xviii, fig. 255.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 174.

**Margaron (Margaritana) tombigbeensis* LEA, Syn., 1870, p. 68.

**Margaritana (Alasmodonta) tombigbeensis* CLESSIN, Conch. Cab. Ano., 1876, p. 268, pl. lxxxi, figs. 3, 4.

Tombigbee River, Mississippi.

†STROPHITUS GESNERII Lea.

**Margaritana gesnerii* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 138; *Jl. Ac. N. Sci. Phila., V, 1862, p. 211, pl. xxxii, fig. 280; *Obs., IX, 1863, p. 33, pl. xxxii, fig. 280.—*B. H. WRIGHT, Check List, 1888.—*SOWERBY, Conch. Icon., XVI, 1868, pl. lxxxiv, fig. 446.

**Margaron (Margaritana) gesnerii* LEA, Syn., 1870, p. 67.

**Margaritana gesneri* PÆTEL, Conch. Sam., III, 1890, p. 173.

Uphaupsee and Swamp creeks, Alabama.

†STROPHITUS ELLIOTTII Lea.

**Margaritana elliottii* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 138; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 226, pl. xxxi, fig. 108; *Obs., VII, 1859, p. 44, pl. xxxi, fig. 108.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 173.

**Margaron (Margaritana) elliottii* LEA, Syn., 1870, p. 68.

**Margaritana (Alasmodonta) elliottii* CLESSIN, Conch. Cab. Ano., 1876, p. 267, pl. lxxxii, figs. 3, 4.

*†*Margaritana elliptica* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 113; *Jl. Ac. N. Sci. Phila., V, 1862, p. 106, pl. xviii, fig. 254; *Obs., VIII, 1862, p. 110, pl. xviii, fig. 254.—B. H. WRIGHT, Check List, 1888.—PÆTEL, Conch. Sam., III, 1890, p. 173.

**Margaron (Margaritana) elliptica* LEA, Syn., 1870, p. 68.

**Margaritana (Alasmodonta) elliptica* CLESSIN, Conch. Cab. Ano., 1876, p. 270, pl. lxxxi, figs. 5, 6.

Columbus, Georgia; Columbus, Mississippi.

?STROPHITUS RADIATUS Conrad.

**Alasmodonta radiata* CONRAD, Am. Jl. Sci., XXV, 1834, p. 341, pl. 1, fig. 10.¹

Small streams of southern Alabama.

HOMOGENÆ.

Male and female shells alike, oval to elongate; beak sculpture coarse; embryos filling the entire outer gills in the form of thick, smooth pads; the ovisacs not separated by sulci.

Genus ANODONTA (Bruguiere em.) Lamarck, 1799.

(Type, *Mytilus cygneus* Linnæus.)

Mya LINNÆUS, part, Syst. Nat., 1758, p. 1158.

Limnaa POLI, Test. Utriusque Sic., I, 1791, p. 31; II, 1795, p. 253.

Anodontites BRUGUIERE, Jl. Hist. Nat., I, 1792, p. 131.

Anodonta LAMARCK, Prodrome Class. Coq., 1799, p. 87.

Anodon OKEN, Lehrb. Nat. Zool., I, 1815, p. 238.

Anodontes CUVIER, Regne An., II, 1817, p. 472.

Shell elliptical, thin, inflated, often slightly winged posteriorly; beak sculpture consisting of rather numerous more or less parallel ridges, usually somewhat doubly looped, and becoming slightly nodulous on the loops; surface generally smooth, shining; hinge edentulous, reduced to a mere line, regularly curved; muscle scars rather faint; nacre dull.

Animal with the marsupium occupying the whole outer gills, when filled forming a smooth, very thick, liver-colored pad; gills free from the abdominal sac from one-half to their entire length; palpi generally large; branchial opening papillose; anal opening without papillæ, though sometimes very slightly crenulate; superanal opening generally small, widely separated from the anal.

(Group of *Anodonta cygnea*.)

Shell very evenly rounded in front, pointed behind, the point elevated above the base, more or less winged on post dorsal part, the line from the posterior part of the wing to the hinder point usually incurved; beaks flattened, the sculpture consisting of numerous more or less concentric ridges, which are sometimes broken up into rather irregular corrugations.

Animal with the inner gills the larger, free from the abdominal sac nearly or quite their whole length; palpi large.

¹I can make nothing out of this, the figure being poor and the description meager. Conrad places it in *Strophitus* in his Synopsis, 1853.

†ANODONTA CYGNEA Linnæus.¹

- **Mytilus cygneus* LINNÆUS,² Syst. Nat., 10th ed., 1758, I, p. 706, No. 1158.—
 *MÜLLER, Vermium, II, 1774, p. 208.—*DA COSTA, Hist. Nat. Test. Brit., 1778, p. 214, pl. XV, fig. 2.—*GMELIN, Syst. Nat., 13th ed., 1778, p. 3335.—
 *SCHRÖTER, Flussconch., 1779, p. 162, pl. III, fig. 1; *Ein. Conch., III, 1783, p. 440.—*OLIVI, Zool. Atr., 1792, p. 125.—*DONOVAN, Brit. Shells, II, 1800, pl. LV.—MONTAGU, Test. Brit., 1803, p. 170.—TURTON, Brit. Faun., 1807, p. 165.—*MATON and RACKETT, Tr. Linn. Soc. Lond., VIII, 1807, p. 109, pl. III, figs. 2, 3.—*SCHUMACHER, Ess. Nouv. Syst., 1817, p. 106.—*DILLWYN, Cat., 1817, p. 315.—TURTON, Conch. Dict. 1819, p. 115.—*SHEPPARD, Tr. Linn. Soc. Lond., XIII, 1820, p. 84, pl. V, fig. 3.—*WOOD, Ind. Test., 1825, p. 58, pl. XII, fig. 32f.—*HANLEY, Ipsa Linn. Conch., 1855, p. 487.—*WOOD, Ind. Test. Rev., 1856, p. 69, pl. XII, fig. 32f.
- **Anodontes cygneus* CUVIER, Regne An., II, 1817, p. 472.
- **Anodon cygneus* TURTON, Conch. Ins. Brit., 1822, p. 239.—*FLEMING, Hist. Brit. Moll., 1828, p. 415.—*TURTON, Man. L. and F. W. Shells, 1831, p. 17, fig. 8.—
 *BROWN, Land and F. W. Conch., 1836, p. 99, pl. XI, figs. 1-3c.—*THOMPSON, Ann. and Mag. N. Hist., VI, 1840, p. 196.—*BROWN, Ill. Rec. Conch., 1844, p. 79, pl. XXVIII, etc.—*TURTON, Man. L. and F. W. Shells, 1857, p. 271, fig. 65, pl. I, fig. 8.—*TATE, L. and F. W. Moll. Brit., 1866, pl. I.—*SOWERBY, Conch. Icon., XVII, 1870, pl. I, fig. 2.
- **Anodonta cygnea* DRAPARNAUD, Hist. Moll. Fr., 1806, p. 134, pl. XII, fig. 1; pl. XI, fig. 6.—*MILLET, Moll. Maine et Loire, 1813, p. 76.—*GAERTNER, Vers. Syst., 1813, p. 38.—*BRARD, Hist. Coq. Paris, 1815, p. 234, pls. IX, X.—*KLEES, Desc. Test., 1818, p. 43.—*LAMARCK, An. sans Vert., VI, 1819, p. 84.—*C. PFEIFFER, L. and S. Moll., 1821, Pt. 1, p. 111, pl. V, fig. 4.—BOSC, Hist. Nat. Coq., 1824, III, p. 142.—*BLAINVILLE, Man., 1825, p. 358, pl. LXVI, fig. 1.—*WAARDENBERG, Com. Hist. Nat. An., 1827, p. 37.—*STARK, Nat. Hist., 1828, II, p. 89.—*GUERIN, Icon. Regne An., II, 1829, pl. XXVIII, fig. 5.—*ROSSMASSLER, Icon., I, 1835, p. 111, pl. III, fig. 67, etc.—*GOUPIÉ, Hist. Moll. Sarthe, 1835, p. 81.—*WYATT, Man. Conch., 1838, p. 68, pl. XI, fig. 2.—*PORRO, Mal. Como., 1838, p. 110.—
 *GRAS, Moll. Isere, 1840, p. 70, pl. VI, fig. 7.—*HANLEY, Test. Moll., 1842, p. 216; *Biv. Shells, 1843, p. 216.—*SCHOLZ, Schleis. L. and W. Moll., 1843, p. 118.—*MORELET, Moll. Port., 1845, p. 100.—*PUTON, Moll. Vosges, 1847, p. 68.—*GASSEIZ, Moll. Ag., 1849, p. 189.—*STEIN, Die Leb. Schneck., 1850,

¹The number of specific names bestowed on the Anodontas of Europe must run up into the thousands. I have gone over the literature as carefully as possible, and large series of specimens from many localities, and I confess that I am absolutely unable to separate these forms specifically. The variations of form, size, color, solidity, and even texture are sufficient for a large number of species if they were not everywhere connected by intermediate examples. The new school of conchologists has named every conceivable variation, and in very many cases distorted individuals. Europe is a densely settled region, and it is possible that the waters of the streams and ponds in many places may be so affected by sewage and other offal as to produce many changes in the mollusks living in them. I shall attempt to recognize a few of the leading forms as varieties, referring them to the author originally describing them, but on account of the enormous amount of literature pertaining to these forms and its confused condition, I can not attempt to trace out these varieties among the descriptions of subsequent writers.

²According to Hanley (Shells of Linnæus, p. 144) the *A. cygnea* var. *cellensis* of Rossmassler (Icon., IV, 1830, pl. XIX, fig. 280) is marked for this in the Linnæan cabinet. It is a large, rather elongated shell, slightly biangulate behind, somewhat cut away on the upper posterior part, and having fairly full beaks.

- p. 100.—*DROUET, Nay. Fr., 1852, p. 5, pl. I; *Rev. et Mag., IV, 1852, p. 55.—
 *DUPUY, Hist. Moll. Fr., 1852, p. 601, pl. xv, fig. 14.—*FORBES and HANLEY,
 Hist. Brit. Moll., II, 1853, p. 155, pl. xxxix, fig. 3; XL, figs. 2, 3; XLI.—*KUS-
 TER, Conch. Cab. Ano., 1853, p. 58, pl. xv.—VON WAHL, Arch. für Natur K.
 Liv., 2d ser., I, 1855, p. 133.—*MOQUIN-TANDON, Moll. Fr., II, 1855, p. 557,
 pl. XLIV.—*VON WAHL, Suss. Biv. Liv., 1855, p. 133.—*NORDENSKIÖLD and
 NYLANDER, Fin. Moll., 1856, p. 90, pl. vii, fig. 78.—H. and A. ADAMS, Gen. Rec.
 Moll., II, 1857, p. 502; III, pl. cxviii, figs. 1, 1a, 1b.—*CHENU, Man., 1859, II,
 p. 145, fig. 718.—*SOWERBY, Ill. Index Brit. Shells, 1859, pl. vii, No. 4.—
 REEVE, Elements of Conch., II, 1860, pl. p, No. 4.—*JEFFREYS, Brit. Conch.,
 I, 1862, p. 41.—*REEVE, L. and F. W. Moll. Br., 1863, p. 215, fig. 1.—*BIELZ,
 Faun. Sieben, 1863, p. 195.—*HOUGHTON, Int. Obs., 1864, p. 68, pl. [I], figs.
 1, 2.—*BROT, Etudes Náyades, Lemán, 1867, p. 28, pl. I, fig. 1; pl. II, figs. 1, 2.—
 *KOBELT, Faun. Nass. Moll., 1871, p. 247.—*WESTERLUND, Faun. S. N. and D.,
 1873, p. 583.—DROUET, Mem. Acad. Dijon, VIII, 1882, p. 25 (reprint).—*L.
 ADAMS, Coll. Manual, 1884, p. 21, pl. II, fig. 1.—*PÆTEL, Conch. Sam., III,
 1890, p. 178.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 199.—*LOCARD,
 Coq. de Fr., 1893, p. 234.—*WESTERLUND, Act. Soc. F. and F. F., XIII, No. 7,
 1897, p. 168.
- Symphynota cygnea* LEA, Tr. Am. Phil. Soc. 1830, p. 456; Obs., I, 1834, p. 70.
 **Margarita (Anodonta) cygnea* LEA, Syn., 1836, p. 48; 1838, p. 29.
 **Anodon cygnea* CROUCH, Ill. Int. Lam., 1827, pl. ix, fig. 6. CATLOW and REEVE,
 Conch. Nom., 1845, p. 66.
 **Margaron (Anodonta) cygnea* LEA, Syn., 1852, p. 47; 1870, p. 76.
 **Anodonta cygnea* GASSEIS, Moll. Ag., 1849, p. 189.
 **Anodonta radiatus* MÜLLER, Vermium, 1774, p. 209.
 **Mytilus zellensis* GMELIN, Syst. Nat., 13th ed., I, 1788, p. 3262.
 **Anodonta zellensis* WAARDENBERG, Hist. Nat. Animalium, 1827, p. 37.
 **Anodonta cellensis* C. PFEIFFER, Deuts. L. and Suss. Moll., I, 1821, p. 110, pl. vi,
 fig. 1.—*BOSC, Hist. Nat. Coq., III, 1824, p. 144.—*ROSSMASSLER, Icon., IV,
 1836, p. 22, pl. XIX, fig. 280.—*SCHOLZ, Schleis. L. and W. Moll., 1843, p. 119.—
 *STABILE, Faun. Lug., 1845, p. 58, pl. II, fig. 69.—*TROSCHEL, Arch. für Nat.,
 XIII, Plt. 1, 1847, p. 272.—*DES MOULINS, Actes. Soc. L. Bord, XVIII, 1852,
 p. 495.—*DROUET, Rev. et Mag., IV, 1852, p. 60.—*DUPUY, Nay. Fr. No. 2, 1852,
 p. 11, pl. II.—*KUSTER, Conch. Cab. Ano., 1853, p. 16, pl. IV, fig. 3; v, figs. 1-4; VI,
 fig. 1.—*BIELZ, Faun. Sieben., 1863, p. 197.—*BROT, Coq. Lemán, 1867, p. 33,
 pl. III, figs. 1-3; IV, figs. 1, 2; VI, fig. 4.—*SCHRENCK, Reis. und F. Am. L.,
 II, 1867, p. 722.—*KOBELT, Faun. Nass. Moll., 1871, p. 251.—*LEHMAN, Die
 Schneck., 1873, p. 297.—*PÆTEL, Conch. Sam., III, 1890, p. 177.—*WESTER-
 LUND, Faun. Pal., II, Pt. 7, 1890, p. 211.
 **Anodonta variabilis* DRAPARNAUD, Tab. Moll. Fr., 1801, p. 108.—FOURNEL, Faun.
 Moselle, I, 1836, p. 487.
 **Mytilus avonensis* MONTAGU, Test. Brit., 1803, p. 172.—*TURTON, Conch. Dict.,
 1819, p. 116.—*WOOD, Ind. Test. Rev., 1856, p. 69, pl. XII, fig. 34.
 **Anodon avonensis*, TURTON, Conch. Ins. Brit., 1822, p. 211.
 **Anodonta avonensis* MOQUIN-TANDON, Moll. France, II, 1855, p. 562, pl. XLVI,
 figs. 7, 8.
 **Mytilus avoensis* TURTON, Brit. Faun., 1807, p. 165.
 **Mytilus stagnalis* TURTON, Brit. Faun., 1807, p. 165.—*DILLWYN, Cat. I, 1817, p.
 316.—TURTON, Conch. Dict., 1819, p. 115.
 **Anodonta stagnalis* BOSC, Hist. Nat. Coq., III, 1824, p. 143.—*WESTERLUND, Faun.
 Pal., II, Pt. 7, 1890, p. 199.—*LOCARD, Coq. Fr., 1893, p. 228.
 **Mytilus fucatus* DILLWYN, Cat. I, 1817, p. 317.
 **Anodonta sulcata* LAMARCK, An. sans Vert., VI, 1819, p. 85.—*NILSSON, Hist. Moll.
 Svec., 1822, p. 113.—*DESHAYES, Enc. Meth., II, 1827, p. 147, pl. cch.—*KLEE-

- BERG, Moll. Bor., 1828, p. 36. — * DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 566; 3d ed., II, 1839, p. 678. — * HANLEY, Biv. Shells, 1843, p. 216.
- * *Anodonta intermedia* LAMARCK, An. sans Vert., VI, 1819, p. 86. — * C. PFEIFFER, Deuts. L. and S. Moll., 1821, Pt. 1, p. 113, pl. VI, fig. 3. — * NILSSON, Hist. Moll. Svec., 1822, p. 117. — * WAARDENBERG, Hist. Nat. Animalium, 1827, p. 37. — * DESHAYES, Enc. Meth., II, 1827, p. 147, pl. CCI, fig. 2. — * PÆTEL, Conch. Sam., III, 1890, p. 180. — * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 74. — * LOCARD, Coq. de Fr., 1893, p. 263.
- * *Anodon intermedia* CATLOW and REEVE, Conch. Nom., 1845, p. 67.
- * *Anodon intermedius* KENYON, Mag. Nat. Hist., I, 1829, p. 426, fig. 185.
- * *Mytilus incrassatus* SHEPPARD, Tr. Linn. Soc. Lond., XIII, 1820, p. 85, pl. v, fig. 4.
- * *Anodonta incrassata* LOCARD, Coq. de Fr., 1893, p. 260.
- * *Mytilus macula* SHEPPARD, Tr. Linn. Soc. Lond., XIII, 1820, p. 86, pl. v, fig. 6.
- * *Anodon paludosus* TURTON, Conch. Ins. Brit., 1822, p. 240, pl. xv, fig. 6.
- * *Anodonta piscinalis* NILSSON, Hist. Moll. Svec., 1822, p. 116. — * ROSSMASSLER, Icon., IV, 1836, p. 23, pl. XIX, fig. 281, etc. — * HANLEY, Biv. Shells, 1843, p. 216. — * GAS-SIES, Moll. Agen., 1849, p. 191, pl. IV, fig. 1. — * DROUET, Rev. et Mag., IV, 1852, p. 285. — * DUPUY, Hist. Moll. Fr., 1852, p. 612, pl. XXI, figs. 17, 18. — * DROUET, Nay. France, No. 5, 1852, p. 11, pl. v, fig. 1. — * DES MOULINS, Actes Soc. L. Bord., XVIII, 1852, p. 496. — * KUSTER, Conch. Cab. Ano., 1853, p. 45, pl. III, figs. 4, 5. — * MOQUIN-TANDON, Moll. Fr., II, 1855, p. 561, pl. XLV, figs. 5, 6; XLVI, figs. 1, 1a. — * KOBELT, Faun. Nass. Moll., 1871, p. 249. — * LEHMAN, Die Schnecken, 1873, p. 300. — * PÆTEL, Conch. Sam., III, 1890, p. 183. — * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 284; * Kong. Svensk. Vet. 1876, XIV, No. 12, p. 66.
- * *Anodonta ventricosa* C. PFEIFFER, Deuts. L. and Suss. Moll., Pt. 2, 1825, p. 30, pl. III, figs. 1-6. — * DROUET, Rev. et Mag., IV, 1852, p. 58; Nay. France, No. 2, 1852, p. 6. — * KUSTER, Conch. Cab. Ano., 1853, p. 56, pl. XI, figs. 3-5. — * BROU, Coq. Fam. Nay. Lem., 1867, p. 32. — * DROUET, Mem. Ac. Dij., 3d ser., VII, 1882, p. 29 (reprint). — * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 203. — * LOCARD, Coq. de Fr., 1893, p. 230.
- Anodonta minima* MILLET, Mem. Soc. Agr., Aug., 1833, p. 241, pl. XII, fig. 2. — * CATLOW and REEVE, Conch. Nom., 1845, p. 67. — * DUPUY, Hist. Moll. Fr., 1852, p. 611, pl. XX, fig. 20. — * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 277. — * LOCARD, Coq. de Fr., 1893, p. 307.
- Anodonta oblonga* MILLET, Mem. Soc. Agr., Aug., 1833, p. 242, pl. XII. — * DROUET, Rev. et Mag., IV, 1852, p. 63; Nay. Fr., No. 2, 1852, p. 14; Hist. Moll. Fr., 1852, p. 65, pl. XVIII, fig. 13. — * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 215. — * LOCARD, Coq. de Fr., 1893, p. 273.
- * *Anodon oblongus* SOWERBY, Conch. Icon., XVII, 1867, pl. xv, fig. 54.
- * *Anodonta attenuata* HELD, Isis, 1836, p. 280.
- * *Anodonta exulcerata* PORRO, Mal. Como., 1838, p. 111, pl. II, fig. 12. — * CATLOW and REEVE, Conch. Nom., 1845, p. 66. — * SOWERBY, Conch. Icon., XVII, 1870, pl. XXXIII, fig. 131. — * CLESSIN, Conch. Cab. Ano., 1874, p. 127, pl. XL, figs. 3, 4. — * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 262. — * LOCARD, Coq. de Fr., 1893, p. 289.
- * *Anodonta coarctata* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 142, pl. LV, fig. 2. — * DUPUY, Hist. Moll. Fr., 1852, p. 616, pl. XX, fig. 51.
- * *Anodonta elongata* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 141, pl. LV, fig. 1. — * PUTON, Moll. Vosges, 1847, p. 69. — * DUPUY, Hist. Moll. Fr., 1852, p. 620, pl. XVI, fig. 16. — * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 307.
- * *Anodonta glabra* STABILE, Faun. Lug., 1845, p. 58, pl. II, fig. 68. — * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 257.
- * *Anodonta lusitana* MORELET, Moll. Port., 1845, p. 103, pl. XII, fig. 1. — * CLESSIN,

- Conch. Cab. Ano., 1873, p. 85, pl. XXIII, fig. 1.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 256.
- * *Anodonta regularis* MORELET, Moll. Port., 1845, p. 100, pl. x.—* CLESSIN, Conch. Cab. Ano., 1873, p. 85, pl. XXIII, fig. 2.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 208.
- * *Anodonta macilenta* MORELET, Moll. Port., 1845, p. 102.—* CLESSIN, Conch. Cab. Ano., 1873, p. 87, pl. XXIV, fig. 2.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 227.—* PÆTEL, Conch. Sam., III, 1890, p. 181.
- * *Anodonta ranarum* MORELET, Moll. Port., 1845, p. 104, pl. XII, fig. 2.—* CLESSIN, Conch. Cab. Ano., 1873, p. 86, pl. XXIII, fig. 3.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 217.
- * *Anodonta atrovirens* PHILIPPI, Menke's Zeits., 1848, p. 130.—* SOWERBY, Conch. Icon., XVII, 1867, pl. XIII, fig. 41.
- * *Anodonta milletii* RAY and DROUET, Rev. Zool., 1848, p. 3, pl. I, figs. 1-2.—* DUPUY, Hist. Moll. Fr., 1852, p. 617, pl. XXI, fig. 16.—* DROUET, Nay. France, 1852, p. 15.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 291.—* LOCARD, Coq. de Fr., 1893, p. 296.
- * *Anodonta grateloupiana* GASSIES, Moll. Agenais, 1849, p. 193, pl. II, figs. 13-15; pl. III, fig. 1, 1B; pl. IV, fig. 2.—* DES MOULINS, Actes Soc. L. Bord., XVIII, 1852, p. 496.—* PÆTEL, Conch. Sam., III, 1890, p. 180.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 303.
- * *Pseudonodonta grateloupiana* BOURGNIGNET, Mat. Moll. Aceph., I, 1881, p. 29.
- * *Anodonta dupuyi* RAY and DROUET, Rev. Zool., 1849, p. 32, pls. I, II.—* DROUET, Rev. et Mag., 1849, p. 11, pl. II, figs. 1-2; * Nay. Fr., 1852, pl. VII.—* DUPUY, Hist. Moll. Fr., 1852, p. 606, pl. XVII, fig. 13.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 234.—* LOCARD, Coq. de Fr., 1893, p. 254.
- Anodonta rayi* DROUET, Cat. Extram., 1849, No. 35.—* DUPUY, Hist. Moll. Fr., 1852, p. 614, pl. XX, fig. 22.—* SOWERBY, Conch. Icon., XVII, 1867, pl. XIX, fig. 72.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 254.—* LOCARD, Coq. de Fr., 1893, p. 276.
- * *Anodonta idrina* SPINELLI, Cat. Moll. Bresc., 1851, p. 19, pl. I, fig. 1.—* CLESSIN, Conch. Cab. Ano., 1875, p. 167, pl. LV, figs. 1, 2.—* KOBETT, Icon., IV, 1876, p. 67, pl. CXX, figs. 1156-1159.—* CLESSIN, Moll. Oest., 1887, p. 712, fig. 484.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 258.
- * *Anodonta arelatensis* DUPUY, Hist. Moll. Fr., 1852, p. 611, pl. XIX, fig. 14.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 264.
- * *Anodontites europæ* LEACH, Syn. Moll. Gt. Br., 1852, p. 326.
- * *Anodonta moulinsiana* DUPUY, Hist. Moll. Fr., 1852, p. 614, pl. XX, fig. 19.—* DES MOULINS, Actes Soc. L. Bord., XVIII, 1852, p. 494.—* DROUET, Nay. Fr., 1854, p. 5.—* KOBETT, Icon., VI, 1879, p. 45, pl. CLXV, fig. 1055.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 217.
- * *Anodonta normandi* DUPUY, Hist. Moll. Fr., 1852, p. 620, pl. XVI, fig. 15.—* SOWERBY, Conch. Icon., XVII, 1870, pl. XXVIII, fig. 109.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 303.
- * *Anodonta parvula* DROUET, Nay. Fr., No. 5, 1852, p. 9, pl. IV, fig. 2.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 265.—* LOCARD, Coq. de Fr., 1893, p. 283.
- * *Anodonta rossmassleriana* DUPUY, Hist. Moll. Fr., 1852, p. 608, pl. XVIII, fig. 14.—* DROUET, Nay. Fr., No. 5 and 6, 1854, p. 6.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 237.—LOCARD, Coq. de Fr., 1893, p. 266.
- * *Anodonta subponderosa* DUPUY, Hist. Moll. Fr., 1852, p. 607, pl. XVII, fig. 14.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 234.—* LOCARD, Coq. de Fr., 1893, p. 252.
- * *Anodonta tenella* KUSTER, Conch. Cab. Ano., 1852, p. 63, pl. IX, fig. 5.
- * *Anodonta scaldiana* DUPUY, Hist. Moll. Fr., 1852, p. 613, pl. XIX, fig. 12.—* KOBETT, Icon., VII, 1880, p. 35, pl. CXCIV, fig. 1960.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 254.—* LOCARD, Coq. de Fr., 1893, p. 272.

- * *Anodonta luxata* KUSTER, Conch. Cab. Ano., 1853, p. 9, pl. III, fig. 1.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 238; LOCARD, Coq. de Fr., 1893, p. 267.
- * *Anodonta callosa* KUSTER, Conch. Cab. Ano., 1853, p. 36, pl. IX, fig. 1.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 270.
- * *Anodonta cariosa* KUSTER, Conch. Cab. Ano., 1853, p. 43, pl. IV, fig. 3; v, fig. 1; x, figs. 1, 2.—* DROUET, Mem. Ac. Dij., VII, 1882, p. 27 (reprint).—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 216.—* LOCARD, Coq. de Fr., 1893, p. 236.
- * *Anodonta arealis* KUSTER, Conch. Cab. Ano., 1853, p. 47, pl. IX, figs. 2-4.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 263.—* LOCARD, Coq. de Fr., 1893, p. 280.
- * *Anodonta charpentieri* KUSTER, Conch. Cab. Ano., 1853, p. 49, pl. XI, figs. 3, 4.—* SOWERBY, Conch. Icon., XVII, 1867, pl. XII, fig. 37.
- * *Anodonta anserirostris* KUSTER, Conch. Cab. Ano., 1853, p. 55, pl. x, fig. 3; XI, figs. 1, 2.—* DROUET, Mem. Ac. Dij., VII, 1882, p. 28.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 214.
- * *Anodonta inornata* KUSTER, Conch. Cab. Ano., 1853, p. 42, pl. III, fig. 6.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 238.—* LOCARD, Coq. de Fr., 1893, p. 267.
- * *Anodonta tumida* KUSTER, Conch. Cab. Ano., 1853, p. 48, pl. XII, figs. 1, 2.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 240.
- * *Anodonta subultrata* KUSTER, Conch. Cab. Ano., 1853, p. 52, pl. XIII, figs. 1, 2.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 258.—* LOCARD, Coq. de Fr., 1893, p. 287.
- * *Anodonta sondermannii* KUSTER, Conch. Cab. Ano., 1853, p. 54, pl. XIII, fig. 4.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 240.
- * *Anodonta opalina* KUSTER, Conch. Cab. Ano., 1853, p. 60, pl. XVI, figs. 1, 2.—* DROUET, Mem. Ac. Dij., VII, 1882, p. 30 (reprint).—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 284.
- * *Anodonta nilssonii* KUSTER, part, Conch. Cab. Ano., 1853, p. 61, pl. XVIII, fig. 2.¹
- * *Anodonta helvetica* BOURGUIGNAT, Rev. et Mag., XV, 1863, p. 22 (pl. XX in vol. XIV, figs. 1, 2).—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 249.
- * *Anodonta melinia* BOURGUIGNAT, Moll. peu. Con., 1863, p. 50, pl. XXVIII.—* BOURGUIGNAT, Rev. et Mag., 1865, pl. XXIV, figs. 1-5.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 229.
- * *Anodonta elachista* BOURGUIGNAT, Moll. peu. Con., 1863, p. 60, pl. XXXI, figs. 12-14.
- * *Anodonta psammia* BOURGUIGNAT, Rev. et Mag., XV, 1863, p. 21 (pl. XXI in vol. XIV, figs. 1-4).—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 269.
- * *Anodonta pictetiana* BROU, Etude Coq. Lem., 1867, p. 45, pl. VIII, figs. 1-3.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 288.
- * *Anodon simplicidus* SOWERBY, Conch. Icon., XVII, 1867, pl. XII, fig. 40.

† ANODONTA CYGNEA var. ANATINA Linnaeus.²

- * *Mytilus anatinus* LINNÆUS, Syst. Nat., 10th ed., I, 1758, No. 1168.—* MÜLLER, Vermium, 1774, p. 207.—* DA COSTA, Hist. Nat. Test. Brit., 1778, p. 2151, pl. XV, fig. 2.—* SCHRÖTER, Flussc., 1779, p. 160, pl. I, figs. 2, 3; * Ein. Conch., III, 1783, p. 442.—* CHEMNITZ, Conch. Cab., VIII, 1785, p. 189, pl. LXXXVI, fig. 763.—* GMELIN, Syst. Nat., 13th ed., 1788, p. 3355.—* OLIVI, Zool. Adr., 1792, p. 125.—* DONOVAN, Brit. Shells, IV, 1802, pl. CXIII.—* MONTAGU, Test. Brit., 1803, p. 171.—* TURTON, Brit. Faun., 1807, p. 165.—* MATON and RACKETT, Tr. Linn. Soc. Lond., VIII, 1807, p. 110, pl. IIIA, figs. 1, 4.—* SCHUMACHER, Ess.

¹ Kuster's two figures on Plate XVII, which he gives for this species, are the *A. footiana* Lea. This figure is *A. cygnea*.

² According to Hanley (Shells of Linnaeus, p. 144), the *Anodonta complanata* in Rossmassler, Icon., Pt. 4, Plate XX, fig. 283, is contained in the box in the Linnaean collection marked *anatina*. This is a rather small, compressed form, slightly angular in front and behind.

- Nouv. Syst., 1817, p. 107, pl. I, fig. 5.—* DILLWYN, Cat., I, 1817, p. 317.—TURTON, Conch. Diet., 1819, p. 116.—* SHEPPARD, Tr. Linn. Soc. Lond., XIII, 1820, p. 85, pl. v, fig. 5.—* WOOD, Ind. Test., 1825, p. 58, pl. XII, fig. 33e.—* HANLEY, Ipsa. Linn. Conch., 1855, pl. 487, pl. II, fig. 1.—* WOOD, Ind. Test. Rev., 1856, p. 69, pl. XII, fig. 33.
- * *Anodon anatinus* TURTON, Conch. Ins. Brit., 1822, p. 240.—* SOWERBY, Rec. and Fos. Sh., XVII, 1823, fig.—* FLEMING, Hist. Brit. Moll., 1828, p. 415.—* REEVE, Conch. Syst., I, 1841, p. 121, pl. xci, figs. 1, 2.
- * *Anodonta anatina* DRAPARNAUD, Hist. Moll. Fr., 1806, p. 135, pl. XII, fig. 2.—* GAERTNER, Vers. Syst., 1813, p. 37.—* MILLET, Moll. Maine et Loire, 1813, p. 75.—KLEES, Desc. Test., 1818, p. 42.—* LAMARCK, An. sans Vert., VI, 1819, p. 85.—* C. PFEIFFER, Deuts. L. und S. Moll., 1821, Pt. 1, p. 112, pl. vi, fig. 2.—* NILSSON, Hist. Moll. Svec., 1822, p. 114.—* BOSCH, Hist. Nat. Coq., III, 1824, p. 143, pl. XXIII, fig. 1.—* WAARDENBERG, Com. Hist. Nat. An., 1827, p. 37.—STARK, Nat. Hist., II, 1828, p. 89.—* KLEEGER, Moll. Bor., 1828, p. 37.—* ROSSMASSLER, Icon., V and VI, 1837, p. 57, pl. XXIX, figs. 417-420.—* PORRO, Mal. Como., 1838, p. 109.—* HANLEY, Test. Moll., 1842, p. 216; Biv. Shells, 1843, p. 216.—* STABILE, Faun. Lug., 1845, p. 57, pl. II, fig. 67.—* PUTON, Moll. Vosg., 1847, p. 68.—* GASSIES, Moll. Ag., 1849, p. 190.—* MIDDENDORFF, Sib. Reise, II, 1851, p. 283, pl. XXI, fig. 4; XXIX, figs. 5, 6.—* DROUET, Rev. et Mag., IV, 1852, p. 244; Nay. Fr., 1852, p. 4, pl. IV, fig. 1.—* DUPUY, Hist. Moll. Fr., 1852, p. 610, pl. XIX, fig. 13.—* FRIELE, Norsk. L. Fersk. Moll., 1853, p. 54.—* VON WAHL, Suss. Biv. Liv., 1855, p. 122.—* MOQUIN-TANDON, Moll. Terr. et Fl. Fr., II, 1855, p. 558, pl. XLV, figs. 1, 2.—* JEFFREYS, Brit. Conch., I, 1862, p. 43.—* SCHRENCK, Reis. und Forsch. Am. L., II, 1867, p. 721.—BROT, Etude Nay. Lem., 1867, p. 38, pl. III, fig. 4; v, fig. 2; VI, figs. 1-3; VII, figs. 1-3.—* LEHMAN, Die Schneck., 1873, p. 303.—* WESTERLUND, Faun. S. N. and D., 1873, p. 587.—* CLESSIN, Conch. Cab. Ano., 1876, p. 83, pl. XIV, figs. 3, 4; XXVI, figs. 1, 2.—* L. ADAMS, Coll. Man., 1884, p. 21, pl. II, fig. 2.—* PÄTEL, Conch. Sam., III, 1890, p. 176.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 225.—* LOCARD, Coq. de Fr., 1893, p. 275.
- * *Anodon anatina* CATLOW and REEVE, Conch. Nom., 1845, p. 66.
- * *Anodonta complanata* ROSSMASSLER, Icon., I, 1835, p. 112, pl. III, fig. 68, IV, p. 24, pl. XX, fig. 283.—SCHOLZ, Schleis. L. und W. Moll., 1843, p. 123.—* KUSTER, Conch. Cab. Ano., 1853, p. 12, pl. III, figs. 2, 3.—* DROUET, Nay. Fr., 1854, p. 8, pl. VII, fig. 1.—* MOQUIN-TANDON, Moll. Terr. et Fluv. Fr., II, 1855, p. 560, pl. XLV, figs. 3, 4.—* VON WAHL, Arch. für Nat. Kund. Liv., 2d ser., I, p. 115.—* WESTERLUND, Faun. S. N. & D., 1873, p. 586.—* CLESSIN, Nach. Mal. Ges., 1874, p. 85; Dents. Ex. Moll., 1876, p. 446, fig. 292.—* KOBELT, Icon., VI, 1879, p. 44, pl. CLXV, fig. 1650-1654.—* DROUET, Mem. Ac. Dij., VII, 1882, p. 24 (reprint).—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 301.—* PÄTEL, Conch. Sam., III, 1890, p. 178.
- * *Anodon complanatus* SOWERBY, Conch. Icon., XVII, 1867, pl. IX, fig. 21

† *ANODONTA CYGNEA* var. *PONDEROSA* C. Pfeiffer.

- * *Anodonta ponderosa* C. PFEIFFER, Dents. L. und S. Moll., 1825, Pt. 2, p. 31, pl. IV, figs. 1-6.—* WAARDENBERG, Com. Hist. Nat. An., 1827, p. 37.—* KLEEGER, Moll. Bor., 1828, p. 37.—* ROSSMASSLER, Icon., IV, 1836, p. 24, pl. XX, fig. 282.—PORRO, Mal. Como., 1838, p. 112.—* HANLEY, Biv. Shells, 1843, p. 216.—SCHOLZ,

¹ Credited to Ziegler Museum by Rossmassler. It was published under the name *Anodonta compressa* in Menke's Synopsis, 1831, but not described.

² According to Clessin in this paper this is the only European species distinct from *A. cygnea*.

Schl. L. & W. Moll., 1843, p. 122.—* DUPUY, Hist. Moll. Fr., 1852, p. 604, pl. XVIII, fig. 12.—* KUSTER, Conch. Cab. Ano., 1853, p. 13, pl. IV, fig. 1.—* DROUET, Nay. Fr., 1854, p. 2, pl. VI.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 233.

† *ANODONTA CYGNEA* var. *ROSTRATA* Rossmassler.¹

* *Anodonta rostrata* ROSSMASSLER, Icon., IV, 1836, p. 25, pl. XX, fig. 284.²—* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—* STABILLE, Faun. Lug., 1845, p. 53, pl. II, fig. 70.—* DROUET, Rev. et Mag., IV, 1852, p. 288; Nay. Fr., 1852, p. 14, pl. V, fig. 2.—* KUSTER, Conch. Cab. Ano., 1853, p. 14, pl. IV, fig. 2.—* BOURGUIGNAT, Rev. et Mag., XV, 1863, p. 23, (also XIV, pl. XXI, fig. 5).—* DROUET, Mem. Ac. Dij., VII, 1882, p. 37 (reprint).—* PÆTEL, Conch. Sam., III, 1890, p. 184.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 246.

* *Anodon rostratus* SOWERBY, Conch. Icon., XVII, 1867, pl. II, fig. 4.

Europe generally; Siberia; southward, perhaps, into Asia Minor. It has been reported from the Amoor River drainage, but I presume that the specimens from that region belong to the group of *A. woodiana*.

ANODONTA RETTERI Drouet.

* *Anodonta retteri* DROUET, Jl. de Conch., 1899, p. 402, fig.

Pond in the palace of the governor of Samarcand, Turkestan; very much like varieties of *A. cygnea*.

† *ANODONTA LUCASI* Deshayes.³

* *Anodonta lucasi* DESHAYES, Hist. Nat. Moll. Alg., 1848, pl. CVIII, figs. 1, 2.—

* MORELET, J. de Conch., II, 1851, p. 359.—* BOURGUIGNAT, Mal. Alg., 1864, II, p. 296, pls. XXIII, XXIV, fig. 1.—* SOWERBY, Conch. Icon., XVII, 1867, pl. XVII, fig. 64.—* PÆTEL, Conch. Sam., III, 1890, p. 181.

* *Margaron (Anodonta) lucasii* LEA, Syn., 1870, p. 78.

* *Anodonta embia* BOURGUIGNAT, Mal. Alg., 1864, II, p. 297, pl. XXV.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

Algiers.

ANODONTA CILICIA Kobelt.

† *Anodonta cilicia* KOBELT, Icon. Ist Supp., 1895, p. 24, pl. VIIA, fig. 1.

Cilicia, north of Tarsus.

ANODONTA NUMIDICA Bourguignat.

* *Anodonta numidica* BOURGUIGNAT, Mal. Alg., 1864, II, p. 298, pl. XXIV, figs. 2-6.—

* PÆTEL, Conch. Sam., III, 1890, p. 182.

* *Anodonta letourneauxi* BOURGUIGNAT, Mal. Alg., 1864, II, p. 299, pl. XXVI, figs. 2-6.—* PÆTEL, Conch. Sam., III, 1890, p. 181.

Algiers.

¹ Credited by Rossmassler to Kokeil Museum.

² There is an *Anodonta rostrata* Held in Isis, 1836, page 280. I do not know whether it or that of Rossmassler was published first, or whether the two are the same or not.

³ Probably a valid species, yet some specimens apparently belonging to it are certainly very much like forms of *cygnea*.

ANODONTA SUBCIRCULARIS Clessin.¹

* *Anodonta subcircularis* CLESSIN, Conch. Cab. Ano., 1873, p. 87, pl. XXII, figs. 3, 4.
Scutari Lake, European Turkey.

ANODONTA VESCOIANA Bourguignat.

- * *Anodonta vescoiana* BOURGUIGNAT, Ann. Mal., II, 1857, p. 34, pl. I, figs. 1, 2.—
* CLESSIN, Conch. Cab. Ano., 1873, p. 100, pl. XXIX, figs. 1, 2.—* KOBELT, Icon.,
VII, 1880, p. 83, pl. CCVII, fig. 2102.—* PÆTEL, Conch. Sam., III, 1890, p. 186.—
* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 201.
* Margaron (*Anodonta*) *vescoiana* LEA, Syn., 1870, p. 81.

Asia Minor.

† ANODONTA BERINGIANA Middendorff.

- * *Anodonta cellensis* SCHRÖTER var. *beringiana* MIDDENDORFF, Sib. Reise, II, 1851,
p. 284, pl. XXVIII, figs. 4-7; XXIX, figs. 1-4.—* CLESSIN, Conch. Cab. Ano., 1875,
p. 211, pl. LXV, figs. 1, 2.
* *Anodonta youconensis* LEA, Pr. Ac. Nat. Sci. Phila., XI, 1867, p. 81.
* *Anodonta youkanensis* LEA, Jl. Ac. Nat. Sci. Phila., VI, 1868, p. 287, pl. XL, fig.
99; Obs., XII, 1869, p. 47, pl. XL, fig. 99.—* CLESSIN, Conch. Cab. Ano., 1874,
p. 133, pl. XLIII, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.
* Margaron (*Anodonta*) *youkanensis* LEA, Syn., 1870, p. 80.
* *Anodonta youkanensis* PÆTEL, Conch. Sam., III, 1890, p. 186.

Kamtschatka; Alaska.²

† ANODONTA KENNERLYI Lea.

- * *Anodonta kennerlyi* LEA, Pr. Ac. Nat. Sci. Phila., IV, 1860, p. 306; * Jl. Ac. Nat.
Sci. Phila., V, 1862, p. 108, pl. XVIII, fig. 256; * Obs., VIII, 1862, p. 112, pl. XVIII,
fig. 256.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890,
p. 181.
* *Anodon kennerlyi* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXIV, fig. 139.
* Margaron (*Anodonta*) *kennerlyi* LEA, Syn., 1870, p. 79.

Puget Sound; British Columbia.

† ANODONTA OREGONENSIS Lea.

- * *Anodonta oregonensis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 80, pl. XXI, fig. 67;
* Obs., II, 1838, p. 80, pl. XXI, fig. 67.—* TROSCHEL, Arch. für Nat., VI, 1839;
II, p. 239.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 264.—* H. and A.
ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1874,
p. 170, pl. LVI, figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch.
Sam., III, 1890, p. 182.
* *Margarita* (*Anodonta*) *oregonensis* LEA, Syn., 1837, p. 30.
* *Anodon oregonensis* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—* GOULD and
CARPENTER, Pr. Zool. Soc. Lond., 1856, p. 218.—* SOWERBY, Conch. Icon.,
XVII, 1867, pl. XVII, fig. 63.
* Margaron (*Anodonta*) *oregonensis* LEA, Syn., 1852, p. 49; 1870, p. 78.
* *Anodon cognata* GOULD, Proc. Bost. Soc. Nat. Hist., III, 1850, p. 294.

¹ It is very hard to say what this is, as it is evidently a diseased shell. Probably a member of the *Cyanea* group.

² There can be no doubt but what this species occurs in Alaska and the northeastern part of Asia, from an examination of specimens in the U. S. National Museum collection from both regions.

- * *Anodonta cognata* GOULD, U. S. Expl. Exp., XII, 1852, p. 435, figs. 546, 546a, 546b; Otia Conch., 1862, p. 187.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 263.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* PÆTEL, Conch. Sam., III, 1890, p. 178.

Washington; Oregon; North California; East to Great Salt Lake.

ANODONTA CALIFORNIENSIS Lea.

- * *Anodonta californiensis* LEA, Tr. Am. Phil. Soc., X, 1852, p. 286, pl. xxv, fig. 47; * Obs., V, 1852, p. 42, pl. xxv, fig. 47.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 263.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* MUSGRAVE, Phot. Conch., 1863, pl. I, fig. 4.—* CLESSIN, Conch. Cab. Ano., 1874, p. 120, pl. xxxix, figs. 4, 5.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 177.
- * *Margaron (Anodonta) californiensis* LEA, Syn., 1852, p. 47; 1870, p. 75.
- * *Anodon californiensis* SOWERBY, Conch. Icon., XVII, 1870, pl. xxviii, fig. 113.

Washington, south to California; east to Idaho and Arizona. Probably a variety of the next.

† ANODONTA NUTTALLIANA Lea.

- * *Anodonta nuttalliana* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 77, pl. xx, fig. 62; * Obs., II, 1838, p. 77, pl. xx, fig. 62.—* TROSCHEL, Arch. für Nat., V, 1839, Pt. 2, p. 238.—* HANLEY, Test. Moll., 1842, p. 216; Biv. Shells, 1843, p. 216.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1874, p. 147, pl. I, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 190, p. 182.
- * *Margarita (Anodonta) nuttalliana* LEA, Syn., 1838, p. 29.
- * *Margaron (Anodonta) nuttalliana* LEA, Syn., 1852, p. 47; 1870, p. 75.
- * *Anodon nuttalliana* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—* GOULD and CARPENTER, Pr. Zool. Soc. Lond., 1856, p. 218.
- * *Anodonta triangularis* TRASK, Pr. Ac. Nat. Sci. Phila., I, 1855, p. 29.
- * *Anodon triangularis*, SOWERBY, Conch. Icon., XVII, 1870, pl. xxix, fig. 56b.

Washington to California.

† ANODONTA WAHLAMETENSIS Lea.

- * *Anodonta wahlametensis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 78, pl. xx, fig. 64; * Obs., II, 1838, p. 78, pl. xx, fig. 64.—* TROSCHEL, Arch. für Nat., V, 1839, Pt. 2, p. 238.—* HANLEY, Test. Moll., 1842, p. 215; Biv. Shells, 1843, p. 215, pl. xxiv, fig. 13.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1874, p. 161, pl. liii, figs. 7, 8.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 186.
- * *Margarita (Anodonta) wahlametensis* LEA, Syn., 1838, p. 29.
- * *Anodon wahlametensis* CATLOW and REEVE, Conch. Nom., 1845, p. 68.—* GOULD and CARPENTER, Pr. Zool. Soc. Lond., 1856, p. 218.—* SOWERBY, Conch. Icon., XVII, 1867, pl. ix, fig. 22.
- * *Margaron (Anodonta) wahlametensis* LEA, Syn., 1852, p. 47; 1870, p. 75.
- * *Anodonta rotundata* TRASK, Pr. Cal. Ac. Sci., I, 1855, p. 29.
- * *Anodon rostratus* SOWERBY, Conch. Icon., XVII, 1867, pl. ii, fig. 4.
- * *Anodonta laosensis* FISCHER,¹ Bull. Soc. H. N. Autun, 1891, p. 219.

Washington; south to southern California; east to Utah.

¹ Sowerby states that his species is from Lao, (Laos?) an error without doubt, as it is certainly *A. wahlametensis*. As the name *rostrata* had been used for an *Anodonta* by Kokeil (or Rossmassler) Fischer changed Sowerby's name to *laosensis*.

ANODONTA COARCTATA Anton.

**Anodonta coarctata* ANTON, Verz. der Conch., 1839, p. 16.—*TROSCHEL, Arch. für Nat., XI, 1845, Pt. 2, p. 323.—*KUSTER, Conch. Cab. Ano., 1853, p. 34, pl. VIII, fig. 2.—*B. H. WRIGHT, Check List, 1888.—*PÄTEL, Conch. Sam., III, 1890, p. 177.—*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 519.

**Anodonta chapalensis* CROSSE and FISCHER, J. de Conch., XL, 1892, p. 295.—

*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 518, pl. LXIV, figs. 8-11.

Lake Chapala, State of Jalisco, Mexico.

†ANODONTA EXILIOR Lea.

**Anodonta exilior* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 188; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 24, pl. VII, fig. 21; *Obs., XIII, p. 28, pl. VII, fig. 21.—

*B. H. WRIGHT, Check List, 1888.—*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 520.

**Anodonta chalcensis* CROSSE and FISCHER, Jl. de Conch., XLI, 1893, p. 110.—

*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 522, pl. LXIV, figs. 7, 7a.

Mexico, mostly in the Pacific drainage.

(Group of *Anodonta dejecta*.)

Shell rather solid, subtrapezoidal, narrower and rounded in front, where it is often considerably thickened, with a slight post-dorsal wing; beaks rather compressed, the sculpture consisting of numerous irregular ridges which have a tendency to become doubly looped and nodulous; epidermis rather smooth and shining, greenish or ashy brown; naere lurid, darker in the cavity of the beaks; muscle scars irregular, not well defined; dorsal scars few and rather deep.

Animal with dark colored branchiae, the inner wider in front, nearly or quite free from the abdominal sac, the outer wrinkled on the outside into longitudinal and vertical folds, all four united to the mantle to the extreme posterior end; palpi very large; mantle with a wide, thickened double edge; branchial opening with numerous papillae; anal opening smooth; superanal opening long, deep, united a long way below.

†ANODONTA DEJECTA Lewis.

**Anodonta dejecta* LEWIS, Field and Forest, I (August and September), 1875, p. 26;¹ *Yarrow's Report, 1875, p. 952.—*B. H. WRIGHT, Check List, 1888.—SIMPSON, Pr. U. S. Nat. Mus., XIX, 1896, p. 372, pl. XXXII, figs. 4, 5.

*†*Anodonta mearnsiana* SIMPSON, Nautilus, VI, 1893, p. 134.

Arizona; southeastern California; northwest Mexico.

(Group of *Anodonta cataracta*.)

Shell thin or solid, inflated, rather evenly rounded in front, generally somewhat biangulate behind, and very slightly winged in the post-

¹Said to be from Arkansas River or its tributaries west of the one hundredth meridian and collected by Dr. H. C. Yarrow, surgeon and naturalist of Wheeler's expedition. This locality is erroneous, as it belongs, no doubt, exclusively to the Pacific drainage.

dorsal region; beaks rather full, the sculpture consisting of a moderate number of concentric ridges or corrugations, which are generally somewhat doubly looped, and often slightly nodulous; epidermis bright and shining, sometimes faintly rayed.

Animal with large branchiæ, the inner larger in front, free in part from the abdominal sac; marsupium occupying the entire outer gills, forming an enormously thick, liver-colored pad when filled; mantle thin, thickened on its edge; branchial papillæ small; anal opening smooth.

† **ANODONTA CATARACTA** Say.¹

Anodonta cataracta SAY, Nich. Enc., 1st ed., II, 1816, pl. III, fig. 4.—* CONRAD, New L. and F. W. Shells, 1834, p. 73.—* FERUSSAC, Guer. Mag., 1835, p. 25.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 566; 3d ed., II, 1839, p. 678; Tr. Elem., I, 1853, Pt. 2, p. 218, pl. XXX, fig. 4.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.—* KUSTER, Conch. Cab. Ano., 1853, p. 33, pl. VII, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 177.—* H. CARPENTER, Nautilus, IV, 1890, p. 56.

* *Unio cataracta* DESHAYES, Tr. Elem. Conch., 1839, p. 18, pl. XXX, fig. 1.

* *Anodon cataracta* CATLOW and REEVE, Conch. Nom., 1845, p. 66. "

* *Mytilus cataractus* EATON, Zool. Text-Book, 1826, p. 221.

* *Anodon cataractus* SOWERBY, Conch. Man., 1839, fig. 152; Conch. Icon., XVII, 1867, pl. IX, fig. 25.

* ? *Mytilus fluviatilis* DILLWYN, Cat., I, 1817, p. 316.—* HANLEY, Ind. Test. Rev., 1856, p. 69.

* *Anodonta fluviatilis* BOSC, Hist. Nat. Coq., 1824, III, p. 143.—* HANLEY, Test. Moll., 1842, p. 217.—* C. B. ADAMS, Thompson's Hist. of Vt., 1842, p. 164; * F. W. and L. S. of Vt., 1842, p. 14.—* HANLEY, Biv. Shells, 1843, p. 217.—* STIMPSON, Shells of N. Eng., 1851, p. 15.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* LATCHFORD, Tr. Ot. F. N. Club, 1882, p. 56.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Anodonta) fluviatilis* LEA, Syn., 1836, p. 51; 1838, p. 30.

* *Anodon fluviatilis* GOULD, Inv. Mass., 1841, p. 117, fig. 80.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 203, pl. XVIII, fig. 234.—* CATLOW and REEVE, Conch. Nom., 1845, p. 66.—* HARTMAN and MICHENER, Conch. Cest., 1874, p. 94, fig. 192.—SOWERBY, Conch. Icon., XVII, 1869, pl. XXII, fig. 86.—* BINNEY, Inv. of Mass., 2d ed., 1870, p. 178, fig. 480.

* *Margaron (Anodonta) fluviatilis* LEA, Syn., 1852, p. 50; 1870, p. 80.

* *Mytilus marginatus* EATON, Zool. Text-Book, 1826, p. 224.

* *Anodonta dignota* FERUSSAC, Guer. Mag., 1835, p. 25.

* † *Anodonta virgulata* LEA,² Pr. Ac. N. Sci. Phila., I, 1857, p. 86; Jl. Ac. N. Sci. Phila., V, 1862, p. 213, pl. XXXII, fig. 282; * Obs., IX, 1863, p. 35, pl. XXXIII, fig. 282.—* CLESSIN, Conch. Cab. Ano., 1874, p. 152, pl. XLIX, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 186.

* *Margaron (Anodonta) virgulata* LEA, Syn., 1870, p. 80.

¹ This species is probably better known by Dillwyn's name, *fluviatilis*. According to Lea and Binney, the first edition of Nicholson's Encyclopedia appeared in 1816, which would give Say's name priority by one year. This edition is extremely rare, and I have never seen it. It is claimed by some authors to have been issued in 1817. Dillwyn refers his species to Gmelin, who states that his shell came from the fresh waters of Europe, and is allied to *anatina*, and it is quite likely it is *A. cygnea*. As there can be no doubt about Say's species, I think it best to use his name.

² I think this is merely a southern form of *cataracta* and scarcely worthy of a varietal name.

- *† *Anodonta williamsii* LEA, Pr. Ac. N. Sci. Phila., VI, 1862, p. 169; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 27, pl. x, fig. 26; *Obs., XI, 1867, p. 31, pl. x, fig. 26.—
 *CLESSIN, Conch. Cab. Ano., 1874, p. 154, pl. LI, figs. 1, 2.—*B. H. WRIGHT, Check List, 1888.—*SOWERBY, Conch. Icon., XVII, 1870, pl. xxviii, fig. 112.
 *Margaron (*Anodonta*) *williamsii* LEA, Syn., 1870, p. 81.
 *† *Anodonta tryonii* LEA, Pr. Ac. N. Sci. Phila., VI, 1862, p. 169; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 28, pl. x, fig. 27; *Obs., XI, 1867, p. 32, pl. x, fig. 27.—
 *CLESSIN, Conch. Cab. Ano., 1874, p. 155, pl. LI, figs. 3, 4.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 186.
 *Margaron (*Anodonta*) *tryonii* LEA, Syn., 1870, p. 80.
 **Anodon tryonii* HARTMAN and MICHENER, Conch. Cest., 1874, p. 95, fig. 195.

Lower St. Lawrence drainage; streams draining into the Atlantic south to North Carolina. This species has been cited from Grand River, Michigan, and from north of Lake Superior. Mr. Bryant Walker states that the former locality is erroneous, and I presume that the latter is.

† ANODONTA MARGINATA Say.¹

- Anodonta marginata* SAY, Nich. Encey., II, 1816, pl. III, fig. 5.—*C. B. ADAMS, Thompson's Hist. Vt., 1842, p. 14.—*STIMPSON, Shells of N. Eng., 1851, p. 15.
 **Anodonta marginatus* HANLEY, Biv. Shells, 1843, p. 217.
 **Anodonta fragilis* LAMARCK, An. sans. Vert., VI, 1819, p. 85.—*FERUSSAC, Guer. Mag., 1835, p. 25.—*DESSAYES, An. sans. Vert., 2d ed., VI, 1835, p. 566; 3d ed., II, 1839, p. 678.—*DELESSERT, Rec. Coq. Lam., 1841, pl. XIII, fig. 2, 1e, 2a, 2b.—
 *HANLEY, Test. Moll., 1842, p. 217; *Biv. Shells, 1843, p. 217.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—*CLESSIN, Conch. Cab. Ano., 1854, p. 73, pl. XVII, figs. 3, 4.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—*CHENU, Ill. Conch., 1858, pl. III, figs. 4, 4a.—*LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 56.—B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 179.
 **Margarita* (*Anodonta*) *fragilis* LEA, Syn., 1836, p. 50; 1838, p. 30.
 **Anodon fragilis* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—*SOWERBY, Conch. Icon., XVII, 1867, pl. XVII, fig. 61.
 **Margaron* (*Anodonta*) *fragilis* LEA, Syn., 1852, p. 49; 1870, p. 78.
 *† *Anodonta lacustris* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 84; *Jl. Ac. N. Sci. Phila., 1860, p. 363, pl. LXII, fig. 188; *Obs., VIII, 1860, p. 45, pl. LXII, fig. 188.—*CLESSIN, Conch. Cab. Ano., 1873, p. 110, pl. XXXIII, figs. 5, 6.—*LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 56.—*B. H. WRIGHT, Check List, 1888.—
 *PÆTEL, Conch. Sam., III, 1890, p. 181.
 **Anodon lacustris* SOWERBY, Conch. Icon., XVII, 1867, pl. XIX, fig. 73.
 **Margaron* (*Anodonta*) *lacustris*, LEA, Syn., 1870, p. 80.
 **Anodon flava* ANTHONY, Am. Jl. Conch., I, 1865, p. 160, pl. XIV, fig. 3.
 **Anodon flava* B. H. WRIGHT, Check List, 1888.
 **Anodon pallida* ANTHONY, Am. Jl. Conch., I, 1865, p. 162, pl. xv, fig. 3.
 **Anodonta pallida*, B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 153.
 **Anodon pallidus* SOWERBY, Conch. Icon., XVII, 1867, pl. XI, fig. 29.
 **Anodonta glandulosa* ANTHONY, Am. Jl. Conch., I, 1865, p. 163, pl. xvi, fig. 3.—
 *B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 180.

¹ I am very sure that this = the *A. fragilis* Lamarck, and not the *fluvialis* Dillwyn, as Dr. Lea believes. Say describes it as very thin and fragile; epidermis green olive, paler on the disk, and greener before (behind), within bluish white, edged with whitish, and from his dimensions and figure it is a smaller shell than Dillwyn's, the whole agreeing better with the *fragilis* of Lamarck.

* *Anodon irisans* ANTHONY, Am. Jl. Conch., I, 1865, p. 163, pl. xvi, fig. 2.

* *Anodonta irisans* B. H. WRIGHT, Check List, 1888.

* *Anodonta irisans* PÆTEL, Conch. Sam., III, 1890, p. 180.

* ? *Anodon hordeum* SOWERBY, Conch. Icon., XVII, 1867, pl. xviii, fig. 66.

* † *Anodonta subcarinata* CURRIER, Am. Jl. Conch., III, 1867, p. 113, pl. vi, fig. 5.—
* B. H. WRIGHT, Check List, 1888.¹

* ? *Anodon subcylindracea* SOWERBY, Conch. Icon., XVII, 1867, pl. xv, fig. 47a.

* *Anodon exilis* SOWERBY, Conch. Icon., XVII, 1869, pl. xxii, fig. 84.

* ? *Anodon pholadiformis* SOWERBY, Conch. Icon., XVII, 1870, pl. xxvii, fig. 106.

St. Lawrence River drainage.

ANODONTA TERES Conrad.

* *Anodonta teres* CONRAD, New F. W. Shells, 1834, p. 47, pl. vii, fig. 2.²—* FERUSSAC Guerin. Mag., 1835, p. 25.—* MÖLLER, Syn. Nov. Gen., 1836, p. 194.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 25, pl. ii, fig. 9.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.

Santee Canal, South Carolina.

† ANODONTA IMPLICATA Say.

Anodonta implicata SAY, New Harm. Diss., II, No. 22, 1829, p. 340.—* CONRAD, New F. W. Shells, 1834, p. 73.—FERUSSAC, Guerin. Mag., 1835, p. 250.—* SAY, New L. and F. W. Shells, 1840, p. 10.—* L. W. SAY, Terr. and Fluv. Shells, 1840, p. 11.—STIMPSON, Shells of N. Eng., 1851, p. 15.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLES-SIN, Conch. Cab. Ano., 1873, p. 78, pl. xix, fig. 3.—* LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 56.—* B. H. WRIGHT, Check List, 1888.—* H. CARPENTER, Nautilus, IV, 1890, p. 57.

* *Anodon implicata* GOULD, Inv. Mass., 1841, p. 118, fig. 78; 2d ed., 1870, p. 180, fig. 481.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 202.—* HARTMAN and MICHENER, Conch. Cest., 1874, p. 94, fig. 193.

* *Margaron (Anodonta) implicata* LEA, Syn., 1852, p. 50; 1870, p. 80.

* ? *Anodon implicatus* SOWERBY, Conch. Icon., XVII, 1867, pl. xiii, fig. 44.³

* † *Anodonta newtonensis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 79, pl. xxi, fig. 66;
* Obs., II, 1838, p. 79, pl. xxi, fig. 66.—* HANLEY, Test. Moll., 1842, p. 219.

* *Margarita (Anodonta) newtonensis* LEA, Syn., 1838, p. 30.

* *Anodon newtonensis* SOWERBY, Conch. Icon., XVII, 1867, pl. xvii, fig. 62.

* *Anodonta newtoniensis* TROSCHEL, Arch. für Naturg., V, 1839, p. 239.—* PÆTEL, Conch. Sam., III, 1890, p. 182.

* *Anodon newtoniana* CATLOW and REEVE, Conch. Nom., 1845, p. 67.

* ? *Anodon excurvata* DE KAY,⁴ Zool. of N. Y., Moll., Pt. 5, 1843, p. 202, pl. xvii, fig. 233.—CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

¹ The type of this is in Mr. Bryant Walker's collection, and seems to be a variety of *marginata*.

² It is hard to say where this belongs, or what it is. The figure shows it a dirty yellowish, without rays, and shaped something like *A. fragilis*, while the description is too brief to properly characterize it.

³ This may be *A. grandis* Say.

⁴ I am not certain whether this is the *implicata* of Say or his *cataracta*. De Kay states that the shell is thin and fragile, and the figure shows it to be brown, green, and yellowish, in irregular bands, and apparently shining and smooth like *cataracta*. He says that it is nearest to *implicata*.

- **Anodonta housatonica* LINSLEY, Am. Jl. Sci., 1845, p. 277.—*STIMPSON, Shells of N. Eng., 1851, p. 15.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—*PÆTEL, Conch. Sam., III, 1890, p. 180.

St. Lawrence drainage; north to Lake Winnipeg; south in streams flowing into the Atlantic to Virginia; not reported from Michigan.

† *ANODONTA HALLENBECKII* Lea.

- **Anodonta hallenbeckii* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 139; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 232, pl. XXXII, fig. 112; *Obs., VII, 1859, p. 50, pl. XXXII, fig. 112.—*CLESSIN, Conch. Cab. Ano., 1873, p. 95, pl. XXVIII, figs. 3, 4.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 180.—*SOWERBY, Conch. Icon., XVII, 1869, pl. XXIII, fig. 89, 483.
 *Margaron (*Anodonta*) *hallenbeckii* LEA, Syn., 1870, p. 78.
 *† *Anodonta gesnerii* LEA, Pr. Ac. N. Sci. Phila., X, 1858, p. 139; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 231, pl. XXXI, fig. 109.—*Obs., VII, 1859, p. 49, pl. XXXI, fig. 109.—*CLESSIN, Conch. Cab. Ano., 1873, p. 95, pl. XXVIII, figs. 1, 2.—*B. H. WRIGHT, Check List, 1888.
 **Anodon gesnerii* SOWERBY, Conch. Icon., XVII, 1867, pl. VII, fig. 15.
 *Margaron (*Anodonta*) *gesnerii* LEA, Syn., 1870, p. 78.
 **Anodonta gessneri* PÆTEL, Conch. Sam., III, 1890, p. 179.¹

Georgia.

† *ANODONTA DOLIARIS* Lea.²

- **Anodonta dolearis* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 193.
 **Anodonta doliaris* LEA, Jl. Ac. N. Sci. Phila., VI, 1866, p. 24, pl. VIII, fig. 23; *Obs., XI, 1867, p. 28, pl. VIII, fig. 23.—*CLESSIN, Conch. Cab. Ano., 1874, p. 130, pl. XLII, figs. 1, 2.—*B. H. WRIGHT, Check List, 1888.
 *Margaron (*Anodonta*) *doliaris* LEA, Syn., 1870, p. 81.

North Carolina.

† *ANODONTA DARIENSIS* Lea.

- **Anodonta dariensis* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 139; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 230, pl. XXVIII, fig. 99; *Obs., VII, 1859, p. 48, pl. XXVIII, fig. 99.—*CLESSIN, Conch. Cab. Ano., 1874, p. 118, pl. XXXVII, figs. 3, 4.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 178.
 **Anodon dariensis* SOWERBY, Conch. Icon., XVII, 1867, pl. XVIII, fig. 70.
 *Margaron (*Anodonta*) *dariensis* LEA, Syn., 1870, p. 80.

Georgia; north Florida.

Group of *Anodonta imbecillis*.

Shell thin, elliptical, sometimes slightly produced near the center of the base, pointed behind, with a slight posterior dorsal wing; area of the beaks generally flattened though often full; beak sculpture consisting of several rather delicate, irregular ridges which show a tendency to break into nodules; epidermis bright and smooth, usually marked

¹ The type of *A. hallenbeckii* is a sulcate shell, somewhat constricted at the central base, while that of *A. gesneri* is not constricted and is smoother. The former may be slightly discolored, and the museum series of shells appears to absolutely connect the two.

² First spelled *dolearis* by Lea, and afterwards changed by him to *doliaris*.

with beautiful, delicate, capillary rays; nacre bluish. Animal sometimes highly colored; gills nearly alike in size, often having a distinct border of different texture; eye-spots on branchial opening generally perceptible.

† **ANODONTA IMBECILLIS** Say.

Anodonta imbecillis SAY, N. Harm. Diss., II, No. 23, 1829, p. 355.—* CONRAD, New F. W. Shells, 1834, p. 73.—* FERUSSAC, Gner. Mag., 1835, p. 25.—* L. SAY, Terr. and Fluv. Shells, 1840, p. 13.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CALKINS, Pr. Dav. Ac. Sci., 1874, p. 47.—* CLESSIN, Conch. Cab. Ano., 1874, p. 70, pl. XLIX, figs. 4, 5; XXV, fig. 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 180.—* STEARNS, Pr. U. S. Nat. Mus., XIV, 1891, p. 106.—* BAKER, Moll. Chicago, Pt. I, 1898, p. 55, pl. v, fig. 1.

* *Margaron (Anodonta) imbecillis* LEA, Syn., 1852, p. 50; 1870, p. 80.

* *Anodon imbecillis* SOWERBY, Conch. Icon., XVII, 1870, pl. XXVII, fig. 102.

* † *Anodonta incerta* LEA, Tr. Am. Phil. Soc., V, 1834, p. 46, pl. VI, fig. 16; * Obs., I, 1834, p. 158, pl. VI, fig. 16.—* FERUSSAC, Gner. Mag., 1835, p. 25.—* HANLEY, Test. Moll., 1842, p. 218.—* PÆTEL, Conch. Sam., III, 1890, p. 180.

* *Margarita (Anodonta) incerta* LEA, Syn., 1836, p. 51; 1838, p. 30.

* *Anodon incerta* CATLOW and REEVE, Conch. Nom., 1845, p. 67.

* *Anodon incertus* SOWERBY, Conch. Icon., XVII, 1867, pl. XVII, fig. 59.

* *Anodon horda* GOULD, Pr. Bost. Soc. N. H., V, 1855, p. 229; * *Otia* Conch., 1862, p. 218.

* *Anodonta hordeum* PÆTEL, Conch. Sam., III, 1890, p. 180.

Entire Mississippi drainage area; south Michigan; North Carolina to Georgia; southwest to Matamoras, Mexico.

† **ANODONTA HENRYANA** Lea.¹

* *Anodonta henryana* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 102; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 373, pl. LXVI, fig. 198; * Obs., VIII, 1860, p. 55, pl. LXVI, fig. 198.—* CLESSIN, Conch. Cab. Ano., 1874, p. 141, pl. XLV, figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 180.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 525.

* *Margaron (Anodonta) henryana* LEA, Syn., 1870, p. 81.

* *Anodon henryana* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXIV, fig. 140.²

Northeast Mexico; southern Texas.

† **ANODONTA GIBBOSA** Say.³

Anodonta gibbosa SAY, Exp. to St. Peters R., 1824, p. 265, pl. XIV, figs. 3, 4.—* FERUSSAC, Gner. Mag., 1835, p. 25; * CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. Ano., 1873, p. 74, pl. XVIII, fig. 3.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

¹Typically very distinct from *imbecillis*, having the dorsal and basal lines quite straight. But there are many intermediate forms which are very hard to place, and it is probably only a variety of the latter.

²*Anodonta impura* Say, (Disseminator, 1829, p. 355) from Mexico has never been figured and was only briefly described. It is quite likely that it is the same as *A. henryana*.

³An exceedingly variable species in form, size, and solidity. *A. dunlapiana* seems to me to be only a slightly elongated, rather solid form.

- **Margarita (Anodonta) gibbosa* LEA, Syn., 1838, p. 31.
 **Anodon gibbosa* CATLOW and REEVE, Conch. Nom., 1845, p. 67.
 **Margaron (Anodonta) gibbosa* LEA, Syn., 1852, p. 51; 1870, p. 81.
 **Anodon gibbosus* SOWERBY, Conch. Icon., XVII, 1867, pl. ix, fig. 23.
 *†*Anodonta couperiana* LEA, Pr. Am. Phil. Soc., I, 1840, p. 289.
 **Anodonta couperiana* LEA, Tr. Am. Phil. Soc., VIII, 1842, p. 227, pl. xx, fig. 146;
 * Obs., III, 1842, p. 65, pl. xx, fig. 46.—* CONRAD, Pr. Ac. N. Sci. Phila., VI,
 1853, p. 263.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN,
 Conch. Cab. Ano., 1874, p. 148, pl. L, figs. 3, 4.—* B. H. WRIGHT, Check List,
 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 433, pl. LXXIV, fig. 1.
 **Margaron (Anodonta) couperiana* LEA, Syn., 1852, p. 51; 1870, p. 81.
 **Anodonta cooperiana* PÆTEL, Conch. Sam., III, 1890, p. 178.

† *ANODONTA GIBBOSA* var. *DUNLAPIANA* Lea.

- *†*Anodonta dunlapiana* LEA, Pr. Am. Phil. Soc., II, 1842, p. 225; * Tr. Am. Phil. Soc.,
 VIII, 1842, p. 248, pl. xxvii, fig. 65; * Obs., III, 1842, p. 86, pl. xxvii, fig. 65.—
 * SOWERBY, Conch. Icon., XVII, 1869, pl. xxiii, fig. 87.—* CLESSIN, Conch.
 Cab. Ano., 1876, p. 220, pl. LXXIV, figs. 1, 2.—* B. H. WRIGHT, Check List,
 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

South Carolina to Florida.

(Group of *Anodonta suborbiculata*.)

Shell large, compressed, suborbicular, somewhat produced near the center of the base, rounded in front and bluntly pointed behind, with a slight posterior dorsal wing; beaks flattened, the sculpture consisting of a few irregular ridges, which are generally broken into nodules, sometimes only corrugated; a well-defined but low posterior ridge separates the smooth, shining area in front from a darker and rougher one behind; epidermis with numerous delicate, capillary rays near the beaks; muscle scars very large; nacre rather coppery and iridescent; pallial line broken, often marked with radiating ridges. Animal unknown.¹

† *ANODONTA SUBORBICULATA* Say.

- Anodonta suborbiculata* SAY, New Harm. Disseminator (newspaper form), January
 29, 1831; Am. Conch. I, No. II, 1831, pl. xi.—* CONRAD, New F. W. Shells,
 1834, p. 73.—* FERUSSAC, Guer. Mag., 1835, p. 25.—* HANLEY, Test. Moll.,
 1842, p. 222; Biv. Shells, 1843, p. 222.—* CHENU, Bib. Conch., 1st ser., III,
 1845, p. 18, pl. iii, figs. 1, 1a.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853,
 p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch.
 Cab. Ano., 1874, p. 160, pl. lii, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—
 * PÆTEL, Conch. Sam., III, 1890, p. 185.
 **Margarita (Anodonta) suborbiculata* SAY, Syn., 1836, p. 52; 1838, p. 31.
 **Anodon suborbiculata* CATLOW and REEVE, Conch. Nom., 1845, p. 68.
 **Margaron (Anodonta) suborbiculata* SAY, Syn., 1852, p. 51; 1870, p. 81.
 **Anodon suborbiculatus* SOWERBY, Conch. Icon., XVII, 1867, pl. v, fig. 11.

Nebraska, Iowa; Illinois; south to Louisiana. Abundant locally.

¹ Closely related to the *imbecillis* group, but differing in form, beak sculpture, the large muscle scars, and the pallial line.

(Group of *Anodonta woodiana*.)¹

Shell oval, inflated, sometimes narrowed in front, pointed behind, and often a little produced at the posterior base, slightly winged in the post-dorsal region; beaks full, sculptured with numerous strong concentric ridges, which are either straight in the center of the disk or slightly doubly looped. There is often a faint posterior ridge, which may be double, making the shell feebly biangulate behind; dorsal scars one or two under the beaks; muscle scars irregular.

Animal with the gills large, equal in size, rounded below, free the greater part of their length from the abdominal sac, somewhat wrinkled; marsupium occupying the entire outer gills, forming thick brown pads when filled; palpi large, subelliptical, free from the mantle a long way; mantle having a wide, thickened edge; branchial opening crowded with dark papillæ; anal opening with no papillæ, but having purple ridges inside; superanal opening closed nearly its whole length.²

† ANODONTA WOODIANA Lea.

**Symphynota woodiana* LEA, Tr. Am. Phil. Soc., V, 1834, p. 42, pl. v, fig. 13; *Obs., I, 1834, p. 154, pl. v, fig. 13.

**Margarita* (*Anodonta*) *woodiana* LEA, Syn., 1836, p. 48; 1838, p. 29.

**Anodon woodiana* CATLOW and REEVE, Conch. Nom., 1845, p. 68.—*SOWERBY, Conch. Icon., XVII, 1870, pl. XXXVI, fig. 149.

**Anodonta woodiana* HANLEY, Test. Moll., 1842, p. 215; *Biv. Shells, 1843, p. 215.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—*CLESSIN, Conch. Cab. Ano., 1875, p. 146, pl. XLVIII, figs. 1, 2.

**Margaron* (*Anodonta*) *woodiana* LEA, Syn., 1852, p. 47; 1870, p. 75.

†*Symphynota magnifica* LEA,³ Tr. Am. Phil. Soc., V, 1834, p. 42, pl. v, fig. 14; Obs., I, 1834, p. 155, pl. v, fig. 14.

**Margarita* (*Anodonta*) *magnifica* LEA, Syn., 1836, p. 48; 1838, p. 29.

**Anodonta magnifica* HANLEY, Test. Moll., 1842, p. 215; Biv. Shells, 1843, p. 215.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—*CHENU, Man., 1859, II, p. 146, fig. 719.—*SCHRENCK, Reis and F. Am. Lande., II, 1867, p. 718, pl. XXVIII, figs. 1, 2.—*PETEL, Conch. Sam., III, 1890, p. 181.

**Anodonta magnifica* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—SOWERBY, Conch. Icon., XVII, 1870, pl. XXV, fig. 96.

**Margaron* (*Anodonta*) *magnifica* LEA, Syn., 1852, p. 47; 1870, p. 75.

*[?]*Anodonta aurata* KUSTER, Conch. Cab. Ano., 1853, p. 30, pl. VI, fig. 2.

¹ Fischer has made a subgenus *Pteranodon* (Man. Conch., p. 1003) for *A. magnifica*, but I do not see any need of giving it and the related species any higher rank than that of a group. The shell and animal differ but little from those of other *Anodontas*.

²A group closely agreeing in general characters, but having enormous individual variation. Père Heude has bestowed specific names on a great number of forms, most of which are, I think; merely mutations of Lea's *woodiana*. Fortunately the U. S. National Museum possesses in the Morelet collection a large series of author's specimens, and the excellent figures in the Conchyliologie Fluviale Nanking enable me to straighten out this difficult group to some extent.

³Larger and rather more finely developed than *A. woodiana*, but the large series of material I have examined absolutely connects the two.

- *[?] *Anodon rotundatus* SWAINSON, Ex. Conch., 2d ed., 1841, pl. XXXVII.
- * *Anodon gibbum* BENSON, Jl. Asiat. Soc., XXIV, 1855, p. 135.
- * *Anodon gibba* SOWERBY, Conch. Icon., XVII, 1867, pl. VI, fig. 13.—HEUDE, Conch. Fl. Nank., VII, 1881, pl. LI, fig. 95.
- * *Margaron (Anodonta) gibba* LEA, Syn., 1870, p. 81.—* VON MARTENS, Nov. Conch., IV, 1876, p. 159, pl. CXXXVI, figs. 6, 7.
- * *Anodonta gibba* CLESSIN, Conch. Cab. Ano., 1875, p. 181, pl. LX, fig. 4.—* PÆTEL, Conch. Sam., III, 1890, p. 179.
- * *Anodon tricostratus* SOWERBY, Conch. Icon., XVII, 1870, pl. XXV, fig. 98.—* PÆTEL, Conch. Sam., III, 1890, p. 186.
- * *Anodonta edulis* HEUDE, J. de Conch., XXII, 1874, p. 117.
- * *Anodon edulis* HEUDE, Conch. F. Nank., I, 1875, pl. VIII, fig. 18.
- * *Anodonta edulis* PÆTEL, Conch. Sam., III, 1890, p. 179.
- * † *Anodon securiformis* HEUDE, Conch. F. Nank., III, 1877, pl. XVIII, fig. 39.
- * *Anodonta securiformis* PÆTEL, Conch. Sam., III, 1890, p. 167.
- * † *Anodon nigricans* HEUDE, Conch. F. Nank., III, 1877, pl. XIX, fig. 41.
- * *Anodonta nigricans* PÆTEL, Conch. Sam., III, 1890, p. 182.
- * † *Anodon piscatorum* HEUDE, Conch. F. Nank., IV, 1878, pl. XXVI, fig. 56.
- * *Anodon elliptica* HEUDE, C. F. Nank., IV, 1878, pl. XXVII, fig. 58.
- * † *Anodon fusca* HEUDE, C. F. Nank., IV, 1878, pl. XXVIII, fig. 59.
- * *Anodon friniana* HEUDE, C. F. Nank., IV, 1878, pl. XXVIII, fig. 60.
- * *Anodon joreti* HEUDE, C. F. Nank., IV, 1878, pl. XXIX, fig. 62.
- * † *Anodon striata* HEUDE, C. F. Nank., IV, 1878, pl. XXX, fig. 63.
- * *Anodonta striata* PÆTEL, Conch. Sam., III, 1890, p. 185.
- * † *Anodon pacifica* HEUDE, C. F. Nank., IV, 1878, pl. XXXII, fig. 66.
- * † *Anodon tumida* HEUDE, C. F. Nank., V, 1879, pl. XXXV, fig. 69.
- * *Anodonta tumida* PÆTEL, Conch. Sam., III, 1890, p. 186.
- * *Anodon subtetragona* HEUDE, C. F. Nank., V, 1879, pl. XXXVI, fig. 70.
- * *Anodon lineata* HEUDE, C. F. Nank., V, 1879, pl. XXXVI, fig. 71.
- * *Anodonta lineata* PÆTEL, Conch. Sam., III, 1890, p. 181.
- * *Anodon irregularis* HEUDE, C. F. Nank., V, 1879, pl. XXXVII, fig. 73.
- * *Anodonta irregularis* PÆTEL, Conch. Sam., III, 1890, p. 180.
- * *Anodon melanochlorea* HEUDE, C. F. Nank., V, 1879, pl. XXXVIII, fig. 74.
- * *Anodonta melanochlorea* PÆTEL, Conch. Sam., III, 1890, p. 182.
- * † *Anodon agricolarum* HEUDE, C. F. Nank., V, 1879, pl. XXXIX, fig. 75.
- * *Anodonta agricolarum* PÆTEL, Conch. Sam., III, 1890, p. 176.
- * † *Anodon mingorum* HEUDE, C. F. Nank., V, 1879, pl. XL, fig. 77.
- * *Anodonta mingorum* PÆTEL, C. Sam., III, 1890, p. 182.
- * *Anodon bigibba* HEUDE, C. F. Nank., VI, 1880, pl. XLI, fig. 78.
- * *Anodonta bigibba* PÆTEL, C. Sam., III, 1890, p. 177.
- * *Anodon despecta* HEUDE, C. F. Nank., V, 1880, pl. XLI, fig. 79.
- * *Anodon obtusa* HEUDE, C. F. Nank., VI, 1880, pl. XLII, fig. 80.
- * *Anodon navicella* HEUDE, C. F. Nank., VI, 1880, pl. XLII, fig. 81.
- * *Anodonta navicula* PÆTEL, C. Sam., III, 1890, p. 182.
- * *Anodon orbicularis* HEUDE, C. F. Nank., VI, 1880, pl. XLII, fig. 82.
- * *Anodonta orbicularis* PÆTEL, C. Sam., III, 1890, p. 182.
- * *Anodon limosa* HEUDE, C. F. Nank., VI, 1880, pl. XLIII, fig. 83.
- * *Anodonta limosa* PÆTEL, C. Sam., III, 1890, p. 181.
- * *Anodon livida* HEUDE, C. F. Nank., VI, 1880, pl. XLIII, fig. 84.
- * *Anodonta livida* PÆTEL, C. Sam., III, 1890, p. 181.
- * *Anodon castanea* HEUDE, C. F. Nank., VI, 1880, pl. XLIII, fig. 85.
- * *Anodonta castanea* PÆTEL, C. Sam., III, 1890, p. 177.
- * *Anodon minuta* HEUDE, C. F. Nank., VI, 1880, pl. XLIV, fig. 86.
- * *Anodonta minuta* PÆTEL, C. Sam., III, 1890, p. 182.
- * † *Anodon succinea* HEUDE, C. F. Nank., VI, 1880, pl. XLIV, fig. 87.
- * *Anodonta succinea* PÆTEL, C. Sam., III, 1890, p. 185.

- * † *Anodon chiniana* HEUDE, C. F. Nank., VI, 1880, pl. XLV, fig. 88.
 * *Anodonta chiniana* PÆTEL, C. Sam., III, 1890, p. 177.
 * *Anodonta scaphydium* HEUDE, C. F. Nank., VI, 1880, pl. XLVI, fig. 89.
 * *Anodonta scaphydium* PÆTEL, C. Sam., III, 1890, p. 184.
 * † *Anodon puerorum* HEUDE, C. F. Nank., VII, 1880, pl. XLVII, fig. 90.
 * *Anodonta puerorum* PÆTEL, C. Sam., III, 1890, p. 183.
 * *Anodon indecora* HEUDE, C. F. Nank., VI, 1880, pl. XLVIII, fig. 91.
 * *Anodonta indecora* PÆTEL, C. Sam., III, 1890, p. 180.
 * *Anodon sorini* HEUDE, C. F. Nank., VII, 1881, pl. XLIX, fig. 92.
 * *Anodonta sorini* PÆTEL, C. Sam., III, 1890, p. 185.
 * *Anodon rosea* HEUDE, C. F. Nank., VII, 1881, pl. I, fig. 93.
 * *Anodonta rosea* PÆTEL, C. Sam., III, 1890, p. 184.
 * *Anodon aubreyi* HEUDE, C. F. Nank., VII, 1881, pl. LII, fig. 97.
 * *Anodonta aubreyi* PÆTEL, C. Sam., III, 1890, p. 176.
 * *Anodon confusa* HEUDE, C. F. Nank., VII, 1881, pl. LIII, fig. 99.
 * *Anodonta confusa* PÆTEL, C. Sam., III, 1890, p. 178.
 * *Anodon obtusata* HEUDE, C. F. Nank., VII, 1881, pl. LIV, fig. 100.
 * *Anodon rubella* HEUDE, C. F. Nank., VII, 1881, pl. LIV, fig. 100 bis.
 * *Anodonta rubella* PÆTEL, C. Sam., III, 1890, p. 184.
 * *Anodon pulchella* HEUDE, C. F. Nank., VII, 1881, pl. LV, fig. 101.
 * *Anodonta pulchella* PÆTEL, C. Sam., III, 1890, p. 183.
 * † *Anodon florida* HEUDE, C. F. Nank., VII, 1881, pl. LV, fig. 102.
 * *Anodonta florida* PÆTEL, C. Sam., III, 1890, p. 179.
 * *Anodonta retusa* HEUDE,¹ J. de Conch., XXXII, 1884, p. 20.—PÆTEL, Conch. Sam., III, 1890, p. 184.
 * *Anodon intermerata* HEUDE, C. F. Nank., IX, 1885, pl. LXVII, fig. 130.
 * *Anodonta intermerata* PÆTEL, C. Sam., III, 1890, p. 180.
 * *Anodon filippiana* HEUDE, C. F. Nank., IX, 1885, pl. LXVIII, fig. 131.
 * *Anodonta filippiana* PÆTEL, C. Sam., III, 1890, p. 183.
 * *Anodon fantozatiana* HEUDE, C. F. Nank., IX, pl. LXIX, fig. 132.
 * *Anodonta fantozatiana* PÆTEL, C. Sam., III, 1890, p. 179.

China; Cambodia; Siam; Amoor River?

† ANODONTA FENOUILII Heude.

- * *Anodon fenouilii* HEUDE, C. F. Nank., IV, 1878, pl. xxxi, fig. 64.

China.

ANODONTA JOURDYI Morlet.

- * *Anodonta jourdyi* MORLET, JI. de Conch., XXXIV, 1886, pp. 76, 288, pl. xv, figs. 1, 1a.—* PÆTEL, C. Sam., III, 1890, p. 180.

Tonkin.

† ANODONTA LAUTA von Martens.

- * *Anodonta lauta* VON MARTENS, S. B. Nat. Fr., 1877, p. 117; Abh. Senck. Nat. Ges., X, 1877, p. 152, fig. 1.—* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 434, pl. XXI, fig. 1; XXII, fig. 1 (var. *rostrata*).—* PÆTEL, C. Sam., III, 1890, p. 191.—* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 150.
 * *Anodonta woodiana* KOBELT,² Abh. Senck. Nat. Ges., XI, 1879, p. 433, pl. xx, fig. 1.—VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 151.

Japan.

¹ New name for *A. obtusa* preoccupied.

² This differs from *A. woodiana* of Lea, which I think has not been found in Japan. I think this is a form of *lauts*.

† ANODONTA JAPONICA von Martens.

**Anodonta japonica* VON MARTENS in Clessin, Conch. Cab. Ano., 1874, p. 144, pl. XLVII, figs. 3, 4.—*KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 436, pl. XXII, fig. 3.—*PÆTEL, Conch. Sam., III, 1890, p. 180.—*VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 150, fig. 2.

**Anodonta cellensis* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 437, pl. XXII, fig. 4.

**Anodonta kobelti* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 152, fig. 1.

**Anodonta haconensis* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 161, fig. 3.

Japan.

ANODONTA LUCIDA Heude

**Anodon lucida* HEUDE, C. F. Nank., III, 1877, pl. xx, fig. 43.

China.

† ANODONTA RIVULARIS Heude.

**Anodon rivularis* HEUDE, C. F. Nank., III, 1877, pl. xx, fig. 44.

**Anodonta rivularis* PÆTEL, C. Sam., III, 1890, p. 184.

China.

† ANODONTA HARLANDI Baird and Adams.

**Anodonta harlandi* BAIRD and ADAMS, Pr. Zool. Soc. Lond., 1867, p. 492, pl. XXVI, figs. 3, 3a.—*PÆTEL, C. Sam., III, 1890, p. 180.

**Anodon harlandi* HEUDE, C. F. Nank., IV, 1878, pl. xxv, fig. 55.

China.

ANODONTA GLOBOSULA Heude.

**Anodon globosula* HEUDE, C. F. Nank., IV, 1878, pl. xxv, fig. 54.

China.

† ANODONTA EUSCAPHYS Heude.

**Anodon euscaphys* HEUDE, C. F. Nank., V, 1879, pl. xxxv, fig. 68.

**Anodonta euscaphys* HEUDE, C. Sam., III, 1890, p. 179.

China.

ANODONTA FLUMINEA Heude.

**Anodon fluminea* HEUDE, C. F. Nank., III, 1877, pl. xx, fig. 42.

**Anodon torrentis* HEUDE, C. F. Nank., IV, 1878, pl. xxix, fig. 61.

**Anodonta fluminea* PÆTEL, C. Sam., III, 1890, p. 179.

China.

ANODONTA PUMILA Heude.

**Anodon pumila* HEUDE, C. F. Nank., V, 1879, pl. xxxvii, fig. 72.

**Anodonta pumila* PÆTEL, C. Sam., III, 1890, p. 183.

China.

ANODONTA DOLIOLUM Heude.

**Anodon doliolum* HEUDE, C. F. Nank. IV, 1878., pl. xxvii, fig. 57.

China.

ANODONTA CALIPYGOS Kobelt.

- **Anodonta calipygos* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 435, pl. XIX, fig. 1.—* PÆTEL, C. Sam., III, 1890, p. 177.—* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 150.

Japan.

ANODONTA UNDULIFERA Clessin.

- **Anodonta undulifera* CLESSIN, Conch. Cab. Ano., 1876, p. 233, pl. LXXVI, figs. 1, 2.

China.

(Group of *Anodonta arcæformis*.)

Shell elliptical, with the dorsal and ventral lines parallel, greatly inflated, rounded in front, pointed and having a slight dorsal wing behind; umbonal region full, but the beaks flattened; the sculpture, consisting of numerous ridges, curved in front and nearly following the growth lines, then running backward in an almost straight line and ending at the low posterior ridge.

Animal unknown.

†ANODONTA ARCÆFORMIS Heude.

- **Anodon arcæformis* HEUDE, C. F. Nank., III, 1877, pl. XIX, fig. 40.
**Anodonta arcæformis* PÆTEL, C. Sam., III, 1890, p. 176.

China.

(Group of *Anodonta grandis*.)

Shell oval, inflated, with a slight post-dorsal wing; umbonal region inflated; beak sculpture consisting of several coarse, irregular, often broken ridges, which are more or less doubly looped and generally nodulous, especially at the bases of the loops; epidermis usually rather smooth.

Animal with large gills, inner the larger, especially in front, free nearly or quite the entire length of the abdominal sac; palpi large and long; mantle much thickened at the border.

†ANODONTA GRANDIS Say.¹

- Anodonta grandis* SAY, N. Harm. Diss., II, 1829, p. 341.—* CONRAD, New F. W. Shells, 1834, p. 73.—* FERUSSAC, Guer. Mag., 1835, p. 25.—* L. W. SAY, Terr. and Fluv. Shells, 1840, p. 12.—* HANLEY, Test. Moll., 1842, p. 220; * Biv. Shells, 1843, p. 220.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. Ano., 1873, p. 96, pl. XXX, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 180.—* BAKER, Moll. Chicago, Pt. 1, 1898, p. 51, pls. II, III, fig. 1; IV, fig. 1.

¹A species having a very wide distribution and abundant in individuals. It is an exceedingly variable form, and a large number of names have been bestowed on its mutations, all of which seem to be closely connected. I have retained some of these in a varietal sense.

- **Margarita (Anodonta) grandis* LEA, Syn., 1836, p. 52; 1838, p. 31.
- **Anodon grandis* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—* SOWERBY, Conch. Icon., XVII, 1870, pl. i, fig. 1.
- **Margaron (Anodonta) grandis* LEA, Syn., 1852, p. 51; 1870, p. 81.
- *†*Anodonta ovata* LEA,¹ Tr. Am. Phil. Soc., VI, 1838, p. 2, pl. II, fig. 2; Obs., II, 1838, p. 2, pl. II, fig. 2.—* TROSCHER, Arch. für Nat., V, 1839; Pt. 2, p. 238.—* HANLEY, Test. Moll., 1842, p. 219.—* CONRAD, Pr. Ac. N. Sci. Phila., 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CALKINS, Pr. Ottawa Ac., 1874, p. 47.—* CLESSIN, Conch. Cab. Ano., 1874, p. 121, pl. XXXIX, figs. 2, 3.—* PÆTEL, Conch. Sam., III, 1890, p. 182.
- **Margarita (Anodonta) ovata* LEA, Syn., 1836, p. 52; 1838, p. 30.
- **Anodon ovata* CATLOW and REEVE, Conch. Nom., 1845, p. 67.
- **Margaron (Anodonta) ovata* LEA, Syn., 1852, p. 50; 1870, p. 80.
- *†*Margarita (anodonta) salmonia* LEA, Syn., 1836, p. 51; 1838, p. 30.²
- *†*Anodonta salmonia* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 45, pl. XIV, fig. 41; * Obs., II, 1838, p. 45, pl. XIV, fig. 41.—* TROSCHER, Arch. für Nat., V, 1839, Pt. 2, p. 238.—* HANLEY, Test. Moll., 1842, p. 218.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 184.
- **Anodon salmonia* CATLOW and REEVE, Conch. Nom., 1845, p. 68.—* SOWERBY, Conch. Icon., XVI, 1867, pl. XIX, fig. 44.
- **Margaron (Anodonta) salmonia* LEA, Syn., 1852, p. 50; 1870, p. 80.
- *†*Anodonta lewisii* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 84; * JI. Ac. N. Sci. Phila., IV, 1860, p. 362, pl. LXII, fig. 187.—* Obs., VIII, p. 44, pl. LXII, fig. 187.—* CLESSIN, Conch. Cab. Ano., 1874, p. 143, pl. XLIX, figs. 5, 6.—* LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 56.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 181.
- **Anodon lewisii* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXV, fig. 142.
- **Margaron (Anodonta) lewisii* LEA, Syn., 1870, p. 80.

† ANODONTA GRANDIS var. FOOTIANA Lea.

- **Anodonta footiana* LEA, Pr. Am. Phil. Soc., I, 1840, p. 289; * Tr. Am. Phil. Soc., VIII, 1842, p. 225, pl. XX, fig. 44; * Obs., III, 1842, p. 63, pl. XX, fig. 44.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1876, p. 227, pl. LXV fig. 5; LXXIV, fig. 3.—* LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 56.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 179.—* BAKER, Moll. Chicago, Pt. 1, 1898, p. 54, pl. III, figs. 2-4; v, fig. 2; VIII, fig. 5.
- **Margaron (Anodonta) footiana* LEA, Syn., 1852, p. 49; 1870, p. 78.
- **Anodon footiana* SOWERBY, Conch. Icon., XVII, 1867, pl. XIV, fig. 48.
- *†*Anodonta marryattana* LEA, Pr. Am. Phil. Soc., I, 1840, p. 289; * Tr. Am. Phil. Soc., VIII, 1842, p. 226, pl. XX, fig. 45; * Obs., III, 1842, p. 64, pl. XX, fig. 45.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1873, p. 72, pl. XXVI, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 181.
- **Margaron (Anodonta) marryattana* LEA, Syn., 1852, p. 51; 1870, p. 82.
- **Anodon marryattanus* SOWERBY, Conch. Icon., XVII, 1870, pl. XXVIII, fig. 111.
- **Anodonta sulcata* KUSTER, Conch. Cab. Ano., 1873, p. 62, pl. XVIII, fig. 1.

¹I have seen Say's type of *A. grandis*, and it appears to be identical with Lea's *ovata*.

²This is an *ovata* which had become diseased internally, probably from the work of a *Trematod*, so that the nacre became roughened or blistered, and generally salmon colored.

- * *Anodonta nilssonii* KUSTER, part, Conch. Cab. Ano., 1853, p. 61, pl. XVII, figs. 3, 4.
- * *Anodon imbricata* ANTHONY, Am. Jl. Conch., I, 1865, p. 159, pl. XIV, fig. 1.
- * *Anodonta imbricata* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 180.
- * *Anodon subinflata* ANTHONY, Am. Jl. Conch., I, 1865, p. 160, pl. XV, fig. 1.
- * *Anodonta subinflata* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 185.
- * ? *Anodon edentulus* SOWERBY, Conch. Icon., XVII, 1867, pl. XVII, fig. 60.
- * *Anodonta houghtonensis* CURRIER, Shell-bearing Moll. Mich., 1881, p. 14, pl. I, fig. 2.—* B. H. WRIGHT, Check List, 1888.

† ANODONTA GRANDIS var. LEONENSIS Lea.

- * *Anodonta leonensis* LEA, Pr. Ac. N. Sci. Phila., VI, 1862, p. 169; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 25, pl. IX, fig. 24; * Obs., XI, 1867, p. 29, pl. IX, fig. 24.—* CLESSIN, Conch. Cab. Ano., 1874, p. 131, pl. XLII, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 181.
- * *Margaron (Anodonta) leonensis* LEA, Syn., 1870, p. 78.

† ANODONTA GRANDIS var. GIGANTEA Lea.¹

- * *Anodonta gigantea* LEA, Tr. Am. Phil. Soc., 1834, p. 1, pl. I, fig. 1; * Obs., II, 1838, p. 1, pl. I, fig. 1.—* TROSCHER, Arch. für Nat., V, 1839, Pt. 2, p. 238.—* HANLEY, Test. Moll., 1842, p. 219.—* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—* KUSTER, Conch. Cab. Ano., 1853, p. 6, pl. I, figs. 1, 2.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 179.
- * *Margarita (Anodonta) gigantea* LEA, Syn., 1836, p. 52; 1838, p. 31.
- * *Margaron (Anodonta) gigantea* LEA, Syn., 1852, p. 50; 1870, p. 81.
- * *Anodon gigantea* SOWERBY, Conch. Icon., XVII, 1867, pl. VIII, fig. 18.
- * *Anodon giganteus* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXVII, fig. 152.
- * † *Anodonta plana*² LEA, Tr. Am. Phil. Soc., V, 1834, p. 48, pl. VII, fig. 18; Obs., I, p. 160, pl. VII, fig. 18.—* FERUSSAC, Guer. Mag., 1835, p. 25.—* HANLEY, Test. Moll., 1842, p. 219.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1874, p. 142, pl. XLVI, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 183.
- * *Margarita (Anodonta) plana* LEA, Syn., 1836, p. 52; 1838, p. 30.
- * *Anodon plana* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 201, pl. XVII, fig. 232.—* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—* SOWERBY, Conch. Icon., XVII, 1869, pl. XXIV, fig. 94.
- * *Margaron (Anodonta) plana* LEA, Syn., 1852, p. 50; 1870, p. 80.
- * *Anodonta declivis* CONRAD, Am. Jl. Sci., XXV, 1834, p. 341, pl. I, fig. 11; New F. W. Shells, 1834, p. 73.—* FERUSSAC, Guer. Mag., 1835, p. 25.—MÖLLER, Syn., Nov. Gen., 1836, p. 194.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.
- * † *Margarita (Anodonta) decora* LEA, SYN., 1836, p. 52; 1838, p. 30.
- * *Anodonta decora* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 64, pl. XX, fig. 63; * Obs., II, 1838, p. 64, pl. XX, fig. 63.—* TROSCHER, Arch. für Nat., V, 1839, Pt. 2, p. 238.—* HANLEY, Test. Moll., 1842, p. 219.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. Ano., 1873, p. 71, pl. XVII, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 178.

¹A variety of *grandis*. The type is a large, heavy form, with a considerably developed posterior ridge, the shell somewhat truncated above and behind the ridge. I have frequently seen such shells which are evidently pathologic.

²This name is written *palna* in above citation; a typographical error, no doubt.

- * *Anodon decora* CATLOW and REEVE, Conch. Nom., 1845, p. 66.—* SOWERBY, Conch. Icon., XVII, 1869, pl. XXI, fig. 83.
- * *Margaron (Anodonta) decora* LEA, Syn., 1852, p. 50; 1870, p. 81.
- *† *Anodonta harpethensis* LEA, Pr. Am. Phil. Soc., I, 1840, p. 289; Tr. Am. Phil. Soc., VIII, 1842, p. 224, pl. XIX, fig. 42; * Obs., III, 1842, p. 62, pl. XIX, fig. 42.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. Ano., 1876, p. 217, pl. LXXII, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 180.
- * *Margaron (Anadonta) harpethensis* LEA, Syn., 1852, p. 50; 1870, p. 81.
- * *Anodon harpethensis* SOWERBY, Conch. Icon., XVII, 1869, pl. XXI, fig. 82.
- * *Anodon subangulata* ANTHONY, Am. Jl. Conch., I, 1865, p. 158, pl. XIII, fig. 1.
- * *Anodonta subangulata*, B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 185.
- * *Anodonta opalina* KUSTER, Conch. Cab. Ano., 1853, p. 60, pl. XVI, figs. 1, 2.
- * *Anodon opalina* ANTHONY, Am. Jl. Conch., I, 1865, p. 159, pl. XIV, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 182.¹
- * *Anodon menielii* ANTHONY, Am. Jl. Conch., II, 1866, p. 144, pl. VI, fig. 1.
- * *Anodon subgibbosa* ANTHONY, Am. Jl. Conch., II, 1866, p. 144, pl. VI, fig. 2.
- * *Anodonta subgibbosa* B. H. WRIGHT, Check List, 1888.—PÆTEL, Conch. Sam., III, 1890, p. 185.
- * *Anodon subgibbosus* SOWERBY, Conch. Icon., XVIII, 1870, pl. XXVII, fig. 107.
- * *Margaron (Anodonta) subglobosa* LEA, Syn., 1870, p. 81.²
- * *Anodon inornata* ANTHONY, Am. Jl. Conch., II, 1866, p. 145, pl. VII, fig. 1.
- * *Anodonta orata* SOWERBY, Conch. Icon., XVII, 1868, pl. XXII, fig. 85.

† ANODONTA GRANDIS var. BENEDICTENSIS Lea.

- * *Symphynota benedictensis* LEA, Tr. Am. Phil. Soc., V., 1834, p. 104, pl. XVI, fig. 48; * Obs., I, 1834, p. 216, pl. XVI, fig. 48.
- * *Anodonta benedictensis* FERUSSAC, Guer. Mag., 1835, p. 25.—* HANLEY, Test. Moll., I, 1842, p. 216.—* C. B. ADAMS, Thompson's Hist. Vt., 1842, p. 164; F. W. and L. S. of Vt., p. 14.—* DE KAY, Zool. N. Y., Pt. 5, 1843, p. 204, pl. XVIII, fig. 235.—* HANLEY, Biv. Shells, 1843, p. 216.—* STIMPSON, Shells of N. Eng., 1851, p. 15.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1875, p. 163, pl. LIV, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 177.
- * *Margarita (Anodonta) benedictensis* LEA, Syn., 1836, p. 28; 1838, p. 29.
- * *Anodon benedictensis* CATLOW and REEVE, Conch. Nom., 1845, p. 66.—* SOWERBY, Conch. Icon., XVII, 1870, pl. XXVI, fig. 99.
- * *Margaron (Anodonta) benedictensis* LEA, Syn., 1852, p. 47.
- * *Margaron (Anodonta) benedictii* LEA, Syn., 1870, p. 75.
- * *Anodonta benedictii* LATCHFORD, Tr. Ottawa F. N. Cl., 1882, p. 55.—* B. H. WRIGHT, Check List, 1888.
- Anodonta cultrata* GOULD, in letter.

Entire Mississippi system; Upper St. Lawrence drainage; Red River of the North; Lake Winnipeg; Manitoba; southwest to Texas; south-east Pennsylvania??

† ANODONTA BEALEI Lea.

- * *Anodonta bealei* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 194; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 26, pl. IX, fig. 25; * Obs., XI, 1867, p. 30, pl. IX, fig. 25.—

¹ Kuster and Anthony appear to have applied the same name independently to this species.

² As of Anthony.

* CLESSIN, Conch. Cab. Ano., 1874, p. 132, pl. XLIV, figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 176.

* *Margaron (Anodonta) bealei* LEA, Syn., 1870, p. 81.

Texas to Kansas.

† ANODONTA TEXASENSIS Lea.¹

* *Anodonta texasensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 113; * JI. Ac. N. Sci. Phila., IV, 1860, p. 366, pl. LXIII, fig. 191; * Obs., VIII, 1860, p. 48, pl. LXIII, fig. 191.—* CLESSIN, Conch. Cab. Ano., 1873, p. 109, pl. XXXIII, figs. 7, 8.—

* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 185.

* *Anodon texasensis* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXVI, fig. 146.

* *Margaron (Anodonta) texasensis* LEA, Syn., 1870, p. 81.

Texas.

† ANODONTA DANIELSII Lea.

* *Anodonta danielsii* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 139; * JI. Ac. N. Sci. Phila., IV, 1860, p. 365, pl. LXIII, fig. 190; * Obs., VIII, 1860, p. 47, pl. LXIII, fig. 190.—* CLESSIN, Conch. Cab. Ano., 1873, p. 108, pl. XXXII, figs. 1, 2.—

* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 178.

* *Margaron (Anodonta) danielsii* LEA, Syn., 1870, p. 78.

* *Anodonta danielsii* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXVI, fig. 148.

* *Margaritana danielsii* PÆTEL, Conch. Sam., III, 1890, p. 173.

Kansas; Indian Territory.

† ANODONTA OPACA Lea.

* *Anodonta opaca* LEA, Tr. Am. Phil. Soc., X, 1852, p. 285, pl. XXV, fig. 46; * Obs., V, 1852, p. 41, pl. XXV, fig. 46.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1875, p. 209, pl. LXIV, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Anodonta) opaca* LEA, Syn., 1852, p. 50; 1870, p. 80.

Louisiana; Mississippi; north to Kansas.

† ANODONTA STEWARTIANA Lea.

* *Anodonta stewartiana* LEA, Tr. Am. Phil. Soc., V, 1834, p. 47, pl. VI, fig. 17; Obs., I, 1834, p. 159, pl. VI, fig. 17.—* FERUSSAC, Gner. Mag., 1835, p. 25.—* HANLEY, Test. Moll., 1842, p. 220; * Biv. Shells, 1843, p. 220.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1874, pl. XLVIII, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 185.

* *Margarita (Anodonta) stewartiana* LEA, Syn., 1836, p. 52; 1838, p. 31.

* *Anodon stewartiana* CATLOW and REEVE, Conch. Nom., 1845, p. 68.

* *Anodon stewartianus* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXII, fig. 133.

* *Margaron (Anodonta) stewartiana* LEA, Syn., 1852, p. 51; 1870, p. 81.

*† *Anodonta virens* LEA, Tr. Am. Phil. Soc., X, 1852, p. 290, pl. XVIII, fig. 53; * Obs., V, 1852, p. 46, pl. XXVIII, fig. 53.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 267.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Anodonta) virens* LEA, Syn., 1852, p. 51; 1870, p. 81.

* *Anodon virens* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXIV, fig. 138.

Arkansas; Louisiana; Texas.

¹ A very young, broken shell, greatly inflated in the post basal region, is all that Lea has in his collection. It may be a distinct species, or a mere variety of *grandis*.

† ANODONTA CORPULENTA Cooper.

* *Anodonta corpulenta* COOPER, App. to Narrative, Exp. Miss. to It. L., 1834, p. 154.—* MÖLLER, Syn. Nov. Gen., 1836, p. 193.—* COOPER, Second App. School. Expl. Exp., 1855, p. 516.—* CALKINS, Pr. Ottawa Ac. Sci., 1874, p. 47.—* B. H. WRIGHT, Check List, 1888.

* *Anodon corpulenta* SOWERBY, Conch. Icon., XVII, 1870, pl. xxxii, fig. 129.

* *Margaron (Anodonta) corpulenta* LEA, Syn., 1870, p. 81.

Missouri River; Upper Mississippi River drainage; east to Indiana; Tyler, Texas. ? Appears to be replaced in the South by *A. stewartiana*.

† ANODONTA LINNEANA Lea.

* *Anodonta linneana* LEA, Tr. Am. Phil. Soc., 1852, p. 289, pl. xxvii, fig. 51;

* Obs., V, 1852, p. 45, pl. lxxvii, fig. 51.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. Ano., 1874, p. 117, pl. xxxvi, figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 181.

* *Margaron (Anodonta) linneana* LEA, Syn., 1852, p. 51; 1870, p. 81.

* *Anodon linneanus* SOWERBY, Conch. Icon., XVII, 1870, pl. xxxv, fig. 144.

Louisiana; Texas.

† ANODONTA GLOBOSA Lea.

* *Margarita (Anodonta) globosa* LEA, Syn., 1836, p. 52.

* *Anodonta globosa* LEA, Pr. Am. Phil. Soc., II, 1841, p. 31; * Tr. Am. Phil. Soc., VIII, 1842, p. 241, pl. xxiv, fig. 56; * Obs., III, 1842, p. 79, pl. xxiv, fig. 56.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. Ano., 1875, p. 213, pl. lxxv, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 180.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 529, pl. lxxviii, figs. 1, 1a.

* *Anodonta globosa* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—* SOWERBY, Conch. Icon., XVII, 1870, pl. xxxv, fig. 141.

* *Margaron (Anodonta) globosa* LEA, Syn., 1852, p. 51; 1870, p. 81.

* † *Anodon nopalatensis* SOWERBY,¹ Conch. Icon., XVII, 1867, pl. xvi, fig. 58.—

* CLESSIN, Conch. Cab. Ano., 1876, p. 223, pl. lxxiii, fig. 3.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 182.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 531.

* *Margaron (Anodonta) nopalatensis* LEA, Syn., 1870, p. 81.

Mexico.

ANODONTA TABASCOENSIS Morelet.

* *Anodonta tabascoensis* MORELET, Jl. de Conch., XXXII, 1884, p. 124.—* PÆTEL, Conch. Sam., III, 1890, p. 185.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 530, pl. lxii, fig. 1.

Tabasco, Mexico.

¹ Lea's *globosa* is a young shell. Sowerby's *nopalatensis* is the adult. There is a magnificent specimen in the collection of the Philadelphia Academy of Natural Sciences, 7½ inches long and 5½ inches high.

ANODONTA MICANS Anthony.

* *Anodonta micans* ANTHONY, Am. Jl. Conch., I, 1865, p. 162, pl. xvi, fig. 1.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Anodonta) micans* LEA, Syn., 1870, p. 78.

Said to come from Texas. I do not know it, nor where it groups, though it seems something like *A. kennicottii*.

ANODONTA LURULENTA Morelet.

* *Anodonta lurulenta* MORELET, Test. Nov. Pt. 1, 1829, p. 28.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 523, pl. LXIV, figs. 6, 6a.

Guatemala; Yucatan.

†ANODONTA KENNICOTTII Lea.

* *Anodonta kennicottii* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 56; * Jl. Ac. N. Sci. Phila., V, 1862, p. 214, pl. xxxiii, fig. 283; * Obs., IX, 1863, p. 16, pl. xxxiii, fig. 283.—* CLESSIN, Conch. Cab. Ano., 1874, p. 153, pl. XLIX, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Anodonta) kennicottii* LEA, Syn., 1870, p. 78.

* † *Anodonta simpsoniana* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 56; * Jl. Ac. N. Sci. Phila., V, 1862, p. 212, pl. xxxii, fig. 281; * Obs., IX, 1863, p. 34, pl. xxxii, fig. 281.—* CLESSIN, Conch. Cab. Ano., 1874, p. 119, pl. xxxvii, figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* P. ETEL, Conch. Sam., III, 1890, p. 184.

* *Margaron (Anodonta) simpsoniana* LEA, Syn., 1870, p. 78.

* *Anodon simpsonianus* SOWERBY, Conch. Icon., XVII, 1870, pl. xxxvii, fig. 153.

* † *Anodonta dallasiana* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 190; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 29, pl. xi, fig. 28; * Obs., XI, 1863, p. 33, pl. xi, fig. 28.—* CLESSIN, Conch. Cab. Ano., 1873, p. 107, pl. xxxiii, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* P. ETEL, Conch. Sam., III, 1890, p. 178.

* *Margaron (Anodonta) dallasiana* LEA, Syn., 1870, p. 78.

Upper and middle St. Lawrence River system; northwest into the Mackenzie drainage.

†ANODONTA PEPINIANA Lea.

* *Anodonta pepinianus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 96, pl. xvi, fig. 51.—

* TROSCHEL, Arch. für Nat., V, 1839, II, p. 239.

* *Anodon pepinianus* SOWERBY, Conch. Icon., XVII, 1870, pl. xxxvi, fig. 150.

* *Anodonta pepiniana* LEA, Obs., II, 1838, p. 96, pl. xvi, fig. 51.—HANLEY, Test. Moll., 1842, p. 220; Biv. Shells, 1843, p. 220.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1874, p. 158, pl. LIII, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—* P. ETEL, Conch. Sam., III, 1890, p. 183.

* *Margarita (Anodonta) pepiniana* LEA, Syn., 1838, p. 30.

* *Margaron (Anodonta) pepiniana* LEA, Syn., 1872, p. 49; 1850, p. 78.¹

Upper and middle St. Lawrence drainage; Lake Winnipeg.

¹ Although this resembles to some extent the so-called *Anodonta modesta*, and is sometimes slightly incurved in front of the beaks, the beak sculpture is that of the *A. grandis* group, and quite different from that of *A. modesta*.

ANODONTA LUGUBRIS Say.

Anodonta lugubris SAY, N. Harm. Diss. II, 1829, p. 340.¹

The following are indeterminate and spurious *Anodontas*:

* *Anodon anatinus* SOWERBY, Rich. Faun. Boreale Am., III, 1836, p. 316.

Saskatchewan River; probably *A. implicata*.

* *Anodonta apollonica* BOURGUIGNAT, Moll. Aceph. Eur., 1881, p. 91.

Asia Minor.

* *Anodonta aureus* PÆTEL, C. Sam., III, 1890, p. 176.

China.

* *Anodonta brandti* DROUET, Jl. de Conch., 1888, p. 108.

* *Anodonta callifera* VON MARTENS, Proc. Zool. Soc., 1860, p. 15.

* *Anodonta chantrei* BOURGUIGNAT, J. de Conch., 1883, p. 187.

* *Anodonta chinensis* PÆTEL, C. Sam., III, 1890, p. 177.

* *Anodon cochlearis* SOWERBY, C. Icon., XVII, 1870, pl. XXXIII, fig. 135. Probably a *glabaris*.

* *Anodonta cornea* PHILIPPI, Menke's Zeits., 1848, p. 130.

* *Anodonta curvatus* JAY, Cat., 1850, p. 27.

Anodon dalei LESSON, Oeuvres Buffon, I, p. 155, pl. III, fig. 1. ?

* *Anodonta dignota* RAFINESQUE, Cont. Mon., 1831, p. 6.

* *Anodonta fragilis* FITZINGER, Syst. Verz., 1833, p. 120.

* *Anodonta gallandi* BOURGUIGNAT, Mat. Aceph. Eur., 1881, p. 95.

* *Anodonta hockingensis* CALL, Am. Nat., 1880, p. 529, Moore, manuscript. According to Call, it is *A. grandis* Say.

* *Anodonta inflata* RAFINESQUE, Cont. Monog., 1831, p. 6.

* *Anodonta laminata* ROCHEBRUNE, Bull. Soc. Philom., VI, 1882, p. 40.

* *Anodonta martensi* CLESSIN, C. Cab. Ano., 1876, p. 181, pl. LXIII, fig. 2.

* *Anodonta ohioensis* RAFINESQUE, An. Gen. Sci. Brux., V, 1820, p. 316.

Anodonta orata STARK, Elements Nat. Hist., II, 1822, p. 90.

* *Anodon pictus* SWAINSON, Ex. Conch., 2d ed., 1841, p. 39.

* *Anodonta polymorpha* LEA, Syn., 1870, p. 84. Credited to Kuster. Where?

* *Anodonta pseudodontopsis* BOURGUIGNAT, Jl. de Conch., 1883, p. 187.

* *Anodontites radiata* VALENCIENNES, Enc. Meth., 1824, p. 147, pl. CCHII, fig. 4 = *Modiolaria nigra*.

* *Anodonta recurvirostra* LEA, Syn., 1870, p. 84; Kuster (?).

* *Anodonta rugifera* DUNKER, Mal. Bl., 1858, p. 225. Probably a *Spatha*.

* *Anodonta schlaflfii* MOUSSON, J. de Conch., 1874, p. 51.

* *Anodonta sedakowi* SIECMASCHKO, Bull. de Ac. Petersb., 1849, p. 225.

* *Anodonta somersi* CALL, Am. Nat., 1880, p. 529, credited to Moore, manuscript, *A. grandis* fide Call.

* *Anodonta subrostrata* PHILIPPI, Mal. Bl., 1869, p. 39.

* *Anodonta subsinuata* PHILIPPI, Mal. Bl., 1869, p. 41. *A. glabaris* probably.

* *Anodonta sulcosa* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503. Said to be of Conrad. Where?

* *Anodon triangularis* SOWERBY, Conch. Icon., XVII, 1867, pl. xv, fig. 56.

* *Anodonta trigona* HELD, Isis., 1836, p. 280, *A. cygnea* probably. The name was used by Spix in 1827.

¹ I do not know what this is. It is said to come from the Cumberland River, and is very likely one of the forms of *A. grandis*. I have omitted the synonymy.

* *Anodonta truncata* KUSTER, Faun. Dalm., 1866, p. 131.

* *Anodonta tunizana* MORELET, J. de Conch., 1864, p. 156.

* *Anodonta viridis* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503, as of Lea. Lea did not describe it.

Genus COLLETOPTERUM Bourguignat, 1881.

(Type, *Anodonta letourneuxi* Bourguignat.)

Colletopterum BOURGUIGNAT, Lettres Malacologiques, 1881, p. 45.

Shell inflated, thin, winged in front and behind, hinge arched, very short, without teeth, but with a feeble lateral lamella shown in pronounced relief; ligament internal.

Animal unknown.

COLLETOPTERUM LETOURNEUXI Bourguignat.

Anodonta letourneuxi BOURGUIGNAT, Annales Mal., I, 1870, p. 76.

* *Anodonta* (*Colletopterum*) *letourneuxi* WESTERLUND, Faun. Pal., II, 1890, Pt. 7, p. 310.

Colletopterum letourneuxi BOURGUIGNAT, Lett. Mal., 1882, p. 46; * Bull. Soc. Mal. Fr., VII, 1890, pl. VI, figs. 1, 2.

Servia, at Belgrade.

COLLETOPTERUM PRÆCLARUM Bourguignat.

Anodonta præclara BOURGUIGNAT, Ann. Mal., I, 1870, p. 78.

* *Anodonta* (*Colletopterum*) *præclara*, WESTERLUND, Faun. Pal., II, 1890, Pt. 7, p. 310.

Colletopterum præclaram BOURGUIGNAT, Lett. Mal., 1882, p. 46.

Bulgaria, at Rutschuk.

COLLETOPTERUM EXIMIUM Bourguignat.

Anodonta eximia BOURGUIGNAT, Ann. Mal., I, 1870, p. 80.

* *Anodonta* (*Colletopterum*) *eximia* WESTERLUND, Faun. Pal., II, 1890, Pt. 7 p. 310.

Wallachia at Giurgevo.

COLLETOPTERUM TANOUSI Bourguignat.

Anodonta tanousi BOURGUIGNAT, Ann. Mal., I, 1870.

* *Anodonta* (*Colletopterum*) *tanousi* WESTERLUND, Faun. Pal., II, 1890, Pt. 7, p. 311.

Servia at Belgrade.¹

Genus GABILLOTIA Servain, 1890.

(Type, *Anodonta pseudodopsis* Locard.)

Gabillotia SERVAIN, Bull. Soc. Mal. Fr., VII, 1890, p. 296.

Shell large, subsolid, subcompressed, more or less rhomboid gaping behind; beaks low, with fine, broken, concentric sculpture; epidermis

¹ I am wholly unacquainted with this genus. All the species are from the Danube, and if the characters are accurately given it may possibly be a valid genus, though Westerlund gives it subgeneric rank under *Anodonta*. It is more than likely, however, that all the so-called species are merely specimens of the well-known *Anodonta cygnea*, having a dorsal wing which conceals the ligament, and something which may be vestiges of lateral teeth. Such examples occur among *Anodontas* in other parts of the world.

shining; teeth rudimentary, one in each valve, sometimes rather sharp pointed and flattened; hinge with occasional vestiges of laterals; epidermal matter mingled with the nacre on the hinge; nacre brilliant, often finely radially ridged, especially at the pallear line.

Animal unknown.

GABILLOTIA PSEUDODOPSIS Locard.

* *Anodonta pseudodopsis* LOCARD, Mal. d'Iacs Tib., 1883, p. 61, pl. XIXbis., figs. 1-3.

* *Gabillotia pseudodopsis*, SERVAIN, Bull. Soc. Mal. Fr., VII, 1890, p. 296, fig.

Syria.

GABILLOTIA LOCARDI Servain.

* *Gabillotia locardi* SERVAIN, Bull. Soc. Mal. Fr., VII, 1890, p. 296, pl. V, figs. 1, 2.

Asia Minor.

GABILLOTIA OPPERTI Bourguignat.

* *Unio opperti* Bourguignat, Rev. et Mag., VIII, 1856, p. 71, pls. VIII, fig. 6; IX, fig. 1.

* *Pseudodon opperti* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 182.

* *Margaritana opperti* PÆTEL, Conch. Sam., III, 1890, p. 173.

* *Margaritana euphratica* KOBELT, Icon., new ed., II, 1886, p. 26, pl. XLV, fig. 266.

Euphrates River.

† GABILLIOTIA EUPHRATICA Bourguignat.

* *Unio euphraticus* BOURGUIGNAT, Test. Nov., 1852, p. 28; * Cat. Rais., 1853, p. 75, pl. IV, figs. 1-3; * Rev. et Mag., VIII, 1856, p. 72.

* *Pseudodon euphratica* CONRAD, Am. Jl. Conch., I, 1865, p. 233.

Margaritana euphratica VON MARTENS, Vorderas Conch., 1874, p.*—KOBELT, Icon., new ed., II, 1886, p. 26, pl. XLV, fig. 266.—* PÆTEL, Conch. Sam., III, 1890, p. 173.

* *Leguminaia euphratica* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 188.

* *Unio churchillianus* SOWERBY, Conch. Icon., XVI, 1868, pl. xcvi, fig. 526.

† GABILLOTIA EUPHRATICA var. CHURCHILLIANUS Bourguignat.

* *Unio churchillianus* BOURGUIGNAT, Rev. et Mag., 1857, p. 18, pl. II, figs. 1-4.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Pseudodon churchillianus* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 183.

† *Monocondylæa rhomboidea* LEA, Pr. Ac. N. Sci., Phila., III, 1859, p. 187; * Jl. Ac. N. Sci., Phila., IV, 1860, p. 263, pl. XLII, fig. 143; * Obs., VII, 1860, p. 81, pl. XLII, fig. 143.

* *Margaron (Monocondylæa) rhomboidea* LEA, Syn., 1870, p. 72.

* *Microcondylæa rhomboidea* PÆTEL, Conch. Sam., III, 1890, p. 175.

* *Leguminaia rhomboidea* KOBELT, Icon., new ed., VI, 1893, p. 94, pl. CLXXVIII, figs. 1124, 1125.

* *Unio rhomboideus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIX, fig. 480.

* *Pseudodon rhomboideus* WESTERLUND, Faun. Pal., II, pl. VII, 1890, p. 183.

Asiatic Turkey.

Genus LEGUMINAIA Conrad, 1863.

(Type, *Monocondylaea mardinensis* Lea.)*Leguminaia* CONRAD, Am. Jl. Conch., I, 1865, p. 233.*Microcondylaea* VEST, Verh. Mitth. Sieben. Ver. Nat., 1866, p. 201.*Microcondylus* DROUET, Jl. de Conch., 1879, p. 138.

Shell rhomboid-elliptical, slightly biangulate behind; posterior ridge faint and double; beaks anterior, sculptured with fine, concentric ridges which are sometimes slightly corrugated, but which show a tendency to form two rounded loops; surface smooth or with very faint corrugations on the posterior slope; epidermis rayless or nearly so; a single tooth in each valve, that in the left under the beak, that in the right in front of it, teeth smooth; epidermal and ligamentary matter mingled with the nacre along the hinge; the two front anterior muscle scars united, the hinder distinct, posterior scars faint; beak cavities shallow; dorsal scars few, irregularly disposed.

Animal with the outer branchiæ adherent to the mantle throughout their entire length; inner free from the abdominal sac. (Drouet.)

Section LEGUMINAIA Conrad, 1865.

(Type, *Monocondylaea mardinensis* Lea.)

Shell somewhat solid, elliptical to rhomboid; beaks rather full; teeth strong, with sometimes a slight secondary tooth in the left valve, well in front.

Group of *Leguminaia mardinensis*.

Characters as in the subgenus.

† LEGUMINAIA MARDINENSIS Lea.

* *Monocondylaea mardinensis* LEA, Pr. Ac. N. Sci., Phila., VIII, 1864, p. 286; * Jl. Ac. N. Sci. Phila., VI, 1869, p. 252, pl. XXX, fig. 67; * Obs., XII, 1869, p. 12, pl. XXX, fig. 67.

* *Leguminaia mardinensis* CONRAD, Am. Jl. Conch., I, 1865, p. 233.—* LOCARD, Arch. Mus. Lyon., III, 1883, p. 250.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 188.—KOBELT, Icon., new ed., VI, 1893, p. 92, pl. CLXXVII, figs. 1122, 1123.

* *Margaron (Monocondylaea) mardinensis* LEA, Syn., 1870, p. 73.

* *Margaritana mardinensis* CLESSIN, Conch. Cab. Ano., 1876, p. 266, pl. LXXXIII, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 173.

LEGUMINAIA MARDINENSIS var. CHANTREI Locard.

* *Leguminaia chantrei* LOCARD, Arch. Mus. Lyon, III, 1883, p. 252, pl. XIXbis, figs. 8-10.—* KOBELT, Icon, 1st sup., 1895, p. 23, pl. IV, fig. 3.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 198.

* *Monocondylaea chantreyi* PÆTEL, Conch. Sam., III, 1890, p. 174.

* *Leguminaia bourguignati* LOCARD, Arch. Mus. Lyon, III, 1883, p. 252, pl. XIXbis, figs. 11-18.—WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 189.

Southern Turkey in Asia.

LEGUMINAIA TRIPOLITANA Bourguignat.¹

- * *Unio tripolitanus* BOURGUIGNAT, Test. Nov., 1852, p. 28; * Cat. Rais., 1853, p. 75, pl. IV, figs. 12, 12a; * Rev. et Mag., 1856, p. 72.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
 * *Pseudodon tripolitanus* CONRAD, Am. Jl. Conch., I, 1865, p. 233.
 * *Margaritana tripolitana* CLESSIN, Conch. Cab. Ano., 1876, p. 265, pl. LXXXV, figs. 6, 7.—* PÆTEL, Conch. Sam., III, 1890, p. 174.
 * *Leguminaia tripolitana* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 190.

Tripoli, Syria.

†LEGUMINAIA WHEATLEYI Lea.

- * *Monocondylwa wheatleyi* LEA, Pr. Ac. N. Sci, Phila., VI, 1862, p. 176; * Jl. Ac. N. Sci., Phila., V, 1863, p. 400, pl. I, fig. 307; * Obs., X, 1863, p. 34, pl. I, fig. 307.
 * *Pseudodon wheatleyi* CONRAD, Am. Jl. Conch., I, 1865, p. 233.
 * *Margaron (Monocondylwa) wheatleyi* LEA, Syn., 1870, p. 72.
 * *Microcondylwa (Margaritana) wheatleyi* CLESSIN, Conch. Cab. Ano., 1875, p. 259, pl. LXXXI, figs. 1, 2.
 * *Myrocondylwa wheatleyi* PÆTEL, Conch. Sam., III, 1890, p. 175.
 * *Leguminaia wheatleyi* LOCARD, Arch. Mus. Lyon, III, 1883, p. 276.—WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 189.

Southern Turkey in Asia.

LEGUMINAIA SAULCYI Bourguignat.

- * *Unio saulcyi* BOURGUIGNAT, Test. Noviss., 1852, p. 27; * Cat. Rais., 1853, p. 74, pl. III, figs. 1-3.—* TRISTRAM, Pr. Zool. Soc. Lond., 1865, p. 544.
 * *Pseudodon saulcyi* CONRAD, Am. Jl. Conch., I, 1865, p. 233.
 * *Margaron (Monocondylwa) saulcyi* LEA, Syn., 1870, p. 72.
 * *Microcondylwa saulcyi* CLESSIN, Conch. Cab. Ano., 1876, p. 260, pl. LXXXVI, figs. 4, 5.—* PÆTEL, Conch. Sam., III, 1890, p. 175.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 190.

Syria.

LEGUMINAIA MICHONII Bourguignat.

- * *Unio michonii* BOURGUIGNAT, Test. Nov., 1852, p. 27; * Cat. Rais., 1853, p. 74, pl. III, figs. 10-12; Rev. et Mag., 1856, p. 72.
 * *Pseudodon michonii* CONRAD, Am. Jl., I, 1865, p. 233.
 * *Margaritana michonii* PÆTEL, Conch. Sam., III, 1890, p. 173.
 * *Leguminaia michonii* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 190.

Jaffa, Syria.

(Group of *Leguminaia locardi*.)

Shell nearly evenly elliptical, somewhat inflated, rather thin, sub-transparent; beaks near the center, rather full, curved forward, with irregular undulations; epidermis bright yellowish brown; nacre brilliant; pseudocardinals very obtuse, rather prominent and thick.

¹ Probably a mere variety of *L. mardineusis*.

LEGUMINAIA LOCARDI Simpson.

**Pseudodon chantrei* LOCARD, Arch. Mus. Lyon, III, 1883, p. 254, pl. XIXbis, figs. 4-7.¹—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 183.

Lake Antioch, Syria.

Section MICROCONDYLÆA Vest, 1866.

(Type, *Anodonta uniopsis* LAMARCK.²)

Shell inequilateral, elongate, subrhomboid or elliptical, compressed, rather thin, with low beaks; teeth rudimentary, much compressed, a single one in each valve; nacre bluish or purplish white.

Animal, as in typical *Leguminaia*.

† LEGUMINAIA UNIOPSIS Lamarck.³

**Anodonta uniopsis* LAMARCK, An. sans Vert., VI, 1819, p. 86.—* HANLEY, Test. Moll., 1842, p. 217; * Biv. Shells, 1843, p. 217.—* DESHAYES, Tr. Element., II, 1853, p. 218, pl. XXX, fig. 4.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* PATEL, Conch. Sam., III, 1890, p. 186.

**Margarita (Anodonta) uniopsis* LEA, Syn., 1836, p. 50; 1838, p. 30.

**Anodon uniopsis* CATLOW and REEVE, Conch. Nom., 1845, p. 68.

**Margaron (Anodonta) uniopsis* LEA, Syn., 1852, p. 49; 1870, p. 78.

**Unio depressa* C. PFEIFFER, Nat. Deuts. Moll., 1825, Pt. 2, p. 32, pl. VIII, figs. 3, 4.⁴

**Leguminaia depressa* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 191.

**Alasmodonta compressa* MENKE, Syn., 1828, p. 106.—* PORRO, Mal. Como., 1838, p. 113.—* STABILE, Faun. Lug., 1845, p. 60, pl. III, fig. 71.

**Unio bonelli* ROSSMÄSSLER, Icon., II, 1835, p. 24, pl. IX, fig. 134.⁵—* HANLEY, Test. Moll., 1842, p. 212; * Biv. Shells, 1843, p. 212, pl. XXIII, fig. 59.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIX, fig. 414.

**Margarita (Margaritana) bonellii* LEA, Syn., 1836, p. 45; 1838, p. 27.

**Margaritana bonellii* KUSTER, Conch. Cab. Unio, 1862, p. 296, pl. XCIX, fig. 1.—

* KOBELT, Icon., 1884, p. 62, pl. XXV, figs. 208, 209; XXVI, fig. 210.

**Margaron (Margaritana) bonellii* LEA, Syn., 1852, p. 43; 1870, p. 72.

**Baphia bonellii* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.

¹Locard described this as a *Pseudodon*, and previously in the same year described a *Leguminaia chantrei*, which seems to me to be a mere variety of *L. mardinensis*. As both are undoubtedly Leguminaias, and as the present species was the last described, I am compelled to change its name.

²Lamarck's brief description of *Anodonta uniopsis* entirely agrees with the shell commonly known as *Microcondylæa* or *Monocondylæa bonelli*. The statement that the ligament passes into the hinge exactly fits this, and this is one of the best generic characters of *Leguminaia*.

³The amount of variation of this species is very great. In many cases the posterior end has evidently been injured by unfavorable environment, and the shell is dwarfed and variously distorted, and sometimes a good deal thickened. Of course every individual variation and distortion of this kind has been seized on by the New School and elevated to specific rank. I have seen only what seemed to me a single species of this subgenus, and nothing which really was worthy of a varietal name.

⁴Credited to Muhlfeld in literature.

⁵Credited by Rossmässler to Ferussac, but without other reference. Lea says Ferussac's manuscript labels, neither of which constitutes a published description.

**Anodon bonelli* SOWERBY, Conch. Icon., XVII, 1867, pl. xi, fig. 34.

**Microcondylaea bonelli* CLESSIN, Conch. Cab. Ano., 1861, p. 257; Moll. Osterr., 1887, p. 719, figs. 486, 487.—* PÆTEL, Conch. Sam. III, 1890, p. 175.

**Microcondylus bonelli* DROUET, Jl. de Conch., xxvi, 1879, p. 138; Union. Italie, 1885, p. 110.

**Margaritana (Microcondylaea) bonellii* KOBELT, Icon., 1884, p. 62, pl. xxv, figs. 208, 209.

**Leguminaria bonellii* WESTERLUND Faun. Pal., II, Pt. 7, 1890, p. 191.

South central Europe.

Genus *LASTENA* Rafinesque, 1820.

(Type, *Lastena lata* Rafinesque.)

Lastena RAFINESQUE, Ann. Gen. Sci. Phys. Brux., 1820, p. 316.

Odatelia RAFINESQUE, Atlantic Jl. and Friend., 1832, p. 154.

Leptodea (Rafinesque) CONRAD, Pr. Ac. N. Sci. Phila., 1853, p. 262.

Shell elongated, subsolid, inequilateral, generally wider in front, rounded-truncate at anterior base, pointed at post-basal region, and having a low posterior ridge, with one or more secondary ridges above it; beaks low, sculpture consisting of a few coarse, irregular, longitudinal folds; epidermis shining, often rayed; a single imperfect tooth occurs in each valve, and sometimes vestiges of laterals; there is one roughened dorsal scar in the shallow beak cavity; muscle scars very large, deep and distinct, the posterior greatly elongated; naere purplish, shading to blue at the edge; palleal line radially ridged.

Animal with very long branchiæ, inner and outer about alike in size and form, projecting free slightly behind, the inner free from the abdominal sac nearly the whole length, brownish throughout; palpi long, large; mantle thickened at edge, brown, black above; branchial opening large, with heavy papillæ; anal opening large, without papillæ; super-anal opening united below; foot very large when living, club-shaped, capable of great extension. No gravid specimens have been seen, but the outer branchiæ are probably used as a marsupium.

† *LASTENA LATA* Rafinesque.

**Anodonta (Lastena) lata* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 317, pl. LXXXII, figs. 17, 18.

**Anodonta lata* FERUSSAC, Guer. Mag., 1835, p. 25.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 26, pl. III, figs. 17, 18.

**Leptodea lata* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 262.

**Anodon lata* SOWERBY, Conch. Icon., XVII, 1867, pl. XIX, fig. 76.

**Unio latus* CONRAD, New F. W. Shells, 1834, p. 70.—* SAY, Am. Conch., VI, 1834.—* KUSTER, Conch. Cab. Unio, 1861, p. 388, pl. LXVII, fig. 5.

Unio dehiscens SAY, N. Harm. Diss., II, 1829, p. 308; * Am. Conch., III, 1830, pl. XXIV.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 554; 3d ed., II, 1839, p. 674.—* HANLEY, Test. Moll., 1842, p. 204; * Biv. Shells, 1843, p. 204, pl. XXI, fig. 36.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 32, pl. IX, figs. 7-9.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXV, fig. 393.

**Margarita (Unio) dehiscens* LEA, Syn., 1836, p. 35; 1838, p. 23.

- * *Hemilastena dehiscens* AGASSIZ, Arch. für Nat., I, 1852, p. 50.
 * *Margaron (Margaritana) dehiscens* LEA, Syn., 1852, p. 43; 1870, p. 69.
 * *Baphia dehiscens* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 499.
 * *Margaritana dehiscens* CLESSIN, Conch. Cab. Ano., 1873, p. 274, pl. XXIV, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 173.
 * *Anodonta dehiscens* PÆTEL, Conch. Sam., III, 1890, p. 178.
 * † *Unio oriens* LEA, Tr. Am. Phil. Soc., IV, 1834, p. 73, pl. VI, fig. 5; Obs., I, 1834, p. 83, pl. VI, fig. 5.—* CHENU, Ill. Conch., 1858, pl. XIV, figs. 2, 2a, 2b.— KUSTER, Conch. Cab., 1861, p. 222, pl. LXXV, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIII, fig. 314.
 * *Unio hildrethi* DELESSERT, Rec. Coq. Lam., 1841, pl. XIX, figs. 4a, 4b.

Ohio, Cumberland, and Tennessee river systems.

Genus *SOLENAIA* Conrad, 1868.

(Type, *Mycetopus emarginatus* Lea.)

Solenaiia CONRAD, Am. Jl. Conch., IV, 1869, p. 249.

Shell greatly elongated, falcate, rather thin, narrower and rounded in front, the upper anterior portion being strongly sulcate, gaping at anterior base and behind, having a strong posterior ridge that ends in a point at the post base; beaks low; sculpture of slightly double-looped concentric ridges; epidermis rayless; hinge line narrow; teeth rudimentary, being mere vestiges of one or more lamellar laterals in each valve; dorsal scars numerous, in a line under the hinge; anterior scars faint; posterior elongated; pallial line with a distinct posterior sinus.

According to Fischer the foot is enormously developed for the purpose of burrowing, as in *Mycetopoda*. It is probable that the mantle is closed behind into two siphons.

(Group of *Solenaiia emarginata*.)

Anterior region much narrowed, drawn out in the direction of the anterior base, marked off from the rest of the shell by a faint ridge.

† *SOLENAIA EMARGINATA* Lea.

- * *Mycetopus emarginatus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 308; * Jl. Ac. N. Sci. Phila., V, 1863, p. 398, pl. I, fig. 305; * Obs., X, 1863, p. 34, pl. I, fig. 305.—SOWERBY, Conch. Icon., XVI, 1868, pl. II, fig. 6.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.—* PÆTEL, Conch. Sam., III, 1890, p. 186.
 * *Platiris (Mycetopus) emarginatus* LEA.
 * *Anodonta emarginatus* CLESSIN, Conch. Cab. Ano., 1875, p. 169, pl. LV, figs. 5, 6.
 * *Mycetopus (Solenaiia) emarginatus* FISCHER, Bull. Soc. d'Autun, IV, 1891, p. 135.

Siam.

†SOLENAIA FALCATA Higgins.

- * *Mycetopus falcatus* HIGGINS, Proc. Zool. Soc. Lond., 1868, p. 179, pl. xiv, fig. 6.¹—SOWERBY, Conch. Icon., XVI, 1868, pl. iv, fig. 9.—* CLESSIN, Conch. Cab. Anz., 1875, p. 204, pl. LXVII, figs. 1, 2.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 8.—* PÆTEL, Conch. Sam., III, 1890, p. 186.

* *Platiris (Mycetopus) falcatus* LEA, Syn., 1870, p. 90.

Southeastern Asia.?

(Group of *Solenaia soleniformis*.)

Shell only slightly narrowed in front, the narrow area not distinctly marked off from the rest of the shell.

†SOLENAIA SOLENIFORMIS Benson.

- * *Anodonta soleniformis* BENSON, Jl. As. Soc. Bengal, V, 1836, p. 749.
 * *Spatha soleniformis* HANLEY and THEOBALD, Conch. Ind., 1876, p. 5, pl. ix, fig. 1.
 * *Mycetopus soleniformis* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 94.
 * *Margarona (Unio) bensoni* LEA, Syn., 1870, p. 57.
 * *Mycetopus bensonianus* PÆTEL, Conch. Sam., III, 1890, p. 186.

Assam, India.

†SOLENAIA OLEIVORA Heude.

- * *Mycetopus oleivorus* HEUDE, Conch. Fluv. Nank., III, 1877, pl. xxii, fig. 46; xxiii, fig. 48.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.—* PÆTEL, Conch. Sam., III, 1890, p. 187.
 * † *Mycetopus recognitus* HEUDE, Conch. Fluv. Nank., III, 1877, pl. xxii, fig. 47.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.—* PÆTEL, Conch. Sam., III, 1890, p. 187.
 * *Mycetopus ceruleus* HEUDE, Conch. Fluv. Nank., IX, 1885, pl. LXXI, fig. 135.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.—* PÆTEL, Conch. Sam., III, 1890, p. 186.
 * *Mycetopus armatus* HEUDE, Conch. Fluv. Nank., IX, 1885, pl. LXX, fig. 133.²
 * *Mycetopus arcuatus* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.—PÆTEL, Conch. Sam., III, 1890, p. 186.
 * *Mycetopus viridis* HEUDE, Conch. Fluv. Nank., IX, 1885, pl. LXXI, fig. 136.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.—* PÆTEL, Conch. Sam., III, 1890, p. 187.
 * *Mycetopus succineus* HEUDE, Conch. Fluv. Nank., IX, 1885, pl. LXX, fig. 184.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

China.

¹Said to come from forest streams near Chyavetas, Upper Amazon, and to have been discovered by Mr. Bartlett. I can not believe that it is a South American species at all, but rather that it is from southeastern Asia, as it is an almost exact miniature copy of *S. emarginatus* Lea from that region, having its texture, and not the soft, shining appearance of the *Mycetopodas*. It was described with a miscellaneous lot of shells from various localities, and that given for this is, I think, erroneous.

²*M. arcuatus* Heude in explanation of plate.

† SOLENAIA IRIDINEA Heude.

* *Mycetopus iridincus* HEUDE, Jl. de Conch., XXII, 1874, p. 117; Conch. Fluv. Nank., I, 1875, pl. VIII, fig. 19.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.—

* PÆTEL, Conch. Sam., III, 1890, p. 187.

* *Mycetopus similis* HEUDE, Conch. Fluv. Nank., III, 1877, pl. XXIII, fig. 50.—

* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

China.

SOLENAIA RIVULARIS Heude.

* *Mycetopus rivularis* HEUDE, Conch. Fluv. Nank., III, 1877, pl. XXIII, fig. 49.—

* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.

China.

SOLENAIA CARINATA Heude.¹

* *Mycetopus carinatus* HEUDE, Conch. Fluv. Nank., III, 1877, pl. XXI, fig. 45.—

* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 11.—* PÆTEL, Conch. Sam., III, 1890, p. 186.

China.

SOLENAIA RUGATA Sowerby.

* *Mycetopus rugatus* SOWERBY, Conch. Icon., XVI, 1868, pl. III, fig. 7.—* SMITH, Jl. de Conch., XXII, 1874, pl. IV, fig. 1.—* CLESSIN, Conch. Cab. Ano., 1876, p. 295, pl. LXVII, fig. 4.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

* *Platiris (Mycetopus) rugatus* LEA, Syn., 1870, p. 90.

Said to come from Victoria River, Australia. I am a little in doubt about the locality.

(Group of *Solenaiia triangularis*.)

Shell short, rounded on post base; very narrow in front.

SOLENAIA TRIANGULARIS Heude.

* *Mycetopus triangularis* HEUDE, Conch. Fluv. Nank., IX, 1885, pl. LXXII, fig. 138.—

* FISHER, Jl. de Conch., XXXVIII, 1890, p. 11.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

China.

Genus GONIDEA Conrad, 1837.

(Type, *Anodonta randalli* Trask, and *Anodonta feminalis* Gould.)

Gonidea CONRAD, Pr. Ac. Nat. Sci. Phila., I, 1857, p. 165.

Shell elongated, subtriangular, much narrowed in front, wide behind, inflated, subsolid, usually with a high, sharp, posterior ridge; beaks rather sharp but not high, the sculpture consisting of a few, strong, concentric bars; epidermis rayless; hinge with a rudimentary pseudo-cardinal and lateral in each valve, though these are sometimes wanting; dorsal scars forming a row within the shallow beak cavity; muscle scars irregular; pallial line with a trace of a sinus behind; nacre lurid to purplish.

¹The type and only known shell is a fossil fragment. It may be a recent species. Proc. N. M. vol. xxii—42

Animal with the outer gills larger than the inner behind, narrower in front, inner free from the abdominal sac, all united to the mantle nearly to their posterior points; mantle double edged; branchial opening large, a few irregular, large and small branched papillæ on it; anal opening large, with rudimentary papillæ; superanal opening closed a long way below; foot very large when living. No ova were found in the branchiæ in numerous specimens taken at different dates and localities.

† GONIDEA ANGULATA Lea.

- * *Anodonta angulata* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 97, pl. XVI, fig. 52; * Obs., II, 1838, p. 97, pl. XVI, fig. 52.—* TROSCHIEL, Arch. für Nat., V, 1839, Pt. 2, p. 239.—* HANLEY, Test. Moll., 1842, p. 222; * Biv. Shells, 1843, p. 222, pl. XXIV, fig. 15.—* GOULD, U. S. Expl. Exp., XII, 1852, p. 436.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 263.—* HAUD A. ADAMS, Gen. Rec. Moll., II, 1857, p. 504.—* CHENU, Man., 1859, II, p. 146, fig. 722.—* REEVE, Elem. Conch., II, 1860, pl. XXXI, fig. 180.—* CLESSIN, Conch. Cab. Ano., 1874, p. 159, pl. LIII, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 176.

* *Margarita (Anodonta) angulata* LEA, Syn., 1838, p. 32.

* *Margaron (Anodonta) angulata* LEA, Syn., 1852, p. 52; 1870, p. 83.

* *Anodon angulata* CALTOW and REEVE, Conch. Nom., 1845, p. 66.

* *Anodon angulatus* SOWERBY, Conch. Icon., XVII, 1867, pl. III, fig. 6.

*† *Anodon feminalis* GOULD, Pr. Bost. Soc. N. Hist., III, 1850, p. 293; * *Otia* Conch., 1862, p. 87.

* *Anodonta feminalis* GOULD, U. S. Expl. Exp., XII, 1852, p. 436, figs. 547, 547a, 547b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

* *Anodonta randalli* TRASK, Pr. Cal. Ac. N. Sci., I, 1855, p. 29.—* PÆTEL, Conch. Sam., III, 1890, p. 177.

* *Anodonta biangulata* SOWERBY, Conch. Icon., XVII, 1869, pl. XXIII, figs. 8, 8a, 8b.

* *Anodonta biangulata* CLESSIN, Conch. Cab. Ano., 1876, p. 234, pl. LXXVII, fig. 3.

Central California; north to British Columbia; east to Idaho.

Genus ANODONTOIDES Simpson (in Baker) 1898.

(Type, *Anodonta ferussaciana* Lea.)¹

Anodontopsis SIMPSON (in Baker), Tr. St. Louis Ac. Sci., VIII, 1898, p. 76; not *Anodontopsis* MCCOY.

Anodontoides SIMPSON (in Baker), Moll. Chicago, 1898, p. 72.

Shell elliptical, inflated, thin, with a faint posterior ridge, sometimes constricted at the center of the base; beaks rather full, with a few coarse, subparallel, concentric ridges, which are curved up rather

¹ The shells of this group are much like those of *Strophitus*, but have different beak sculpture. The character of the marsupia is also different in the two genera. In this group a limited number of ovules was found in the inner gills of one of the few gravid specimens seen, while in *Anodonta* only the outer gills contain ova so far as I have observed. It differs from *Anodonta* in the curving in of the hinge line in front of the beaks, in the beak sculpture, and in having papillæ on the anal opening. Notwithstanding the fact that I have found embryos in all four gills of a specimen of this genus, I have placed it with the Homogenæ rather than with the Tetragenæ because the characters of shell and animal seem to agree better with the former than with the latter.

suddenly behind, and back of these there are fine radiating ridges; epidermis smooth, shining, often rayed; hinge line slightly incurved in front of the beaks, edentulous or bearing the merest rudiments of teeth; muscle scars shallow, irregular; nacre bluish white.

Animal with marsupium occupying the outer and sometimes the four leaves of the branchiæ; ovules more numerous in the outer, the whole pad like; gills large, inner semicircular, free from the abdominal sac, or united to it; branchial opening large, with many small papillæ; anal opening with well-developed papillæ.

† ANODONTOIDES FERUSSACIANUS Lea.

* *Anodonta ferussaciana* LEA, Tr. Am. Phil. Soc., V, 1834, p. 45, pl. VI, fig. 15; * Obs., I, 1834, p. 157, pl. VI, p. 15.—* HANLEY, Test. Moll., 1842, p. 218.—* CATLOW and REEVE, Conch. Nom., 1845, p. 66.—CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* MUSGROVE, Phot. Conch., 1863, pl. I, fig. 15.—* CLESSIN, Conch. Cab. Ano., 1873, p. 75, pl. XXIII, fig. 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

* *Anodon ferussaciana* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 200, pl. XVI, fig. 230.—SOWERBY, Conch. Icon., XVII, 1867, pl. XIV, fig. 51.

* *Margarita (Anodonta) ferussaciana* LEA, Syn., 1836, p. 51; 1833, p. 30.

* *Margaron (Anodonta) ferussaciana* LEA, Syn., 1852, p. 50; 1870, p. 79.

* *Anodontoides ferussacianus* BAKER, Moll. Chicago, Pt. 1, 1898, p. 72, pl. III, fig. 6; v, fig. 2.

* † *Anodonta buchanensis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 47, pl. XIV, fig. 43; * Obs., II, 1838, p. 47, pl. XIV, fig. 43.—* TROSCHEL, Arch. für Nat., V, 1839, Pt. 2, p. 238.—* HANLEY, Test. Moll., 1842, p. 223; Biv. Shells, 1843, p. 223.—* CATLOW and REEVE, Conch. Nom., 1845, p. 66.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 263.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* SOWERBY, Conch. Icon., XVII, 1869, pl. XXIV, fig. 92.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 177.

* *Margarita (Anodonta) buchanensis* LEA, Syn., 1836, p. 54; 1838, p. 32.

* *Margaron (Anodonta) buchanensis* LEA, Syn., 1852, p. 51; 1870, p. 82.

* † *Anodonta argentea* LEA, Pr. Ac. Nat. Sci. Phila., I, 1840, p. 289; * Tr. Am. Phil. Soc., VIII, 1842, p. 223, pl. XIX, fig. 41; * Obs., III, 1842, p. 61, pl. XIX, fig. 41.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. Ano., 1876, p. 218, pl. LXXII, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 176.

* *Margaron (Anodonta) argentea* LEA, Syn., 1852, p. 50; 1870, p. 79.

* *Strophilus argenteus* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 262.

* † *Anodonta ferruginea* LEA, Pr. Am. Phil. Soc., I, 1840, p. 289; * Tr. Am. Phil. Soc., VIII, 1842, p. 225, pl. XIX, fig. 43; * Obs., III, 1842, p. 63, pl. XIX, fig. 43.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* MUSGROVE, Phot. Conch., 1863, pl. I, fig. 2.—* CLESSIN, Conch. Cab. Ano., 1876, p. 219, pl. LXXII, figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

* *Margaron (Anodonta) ferruginea* LEA, Syn., 1852, p. 50; 1870, p. 79.

* † *Anodonta plicata* HALDEMAN, Jl. Ac. Nat. Sci. Phila., VIII, 1842, p. 201.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 264.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Anodonta) plicata* LEA, Syn., 1870, p. 79.

* † *Anodonta denigrata* LEA, Tr. Am. Phil. Soc., X, 1852, p. 285, pl. XXV, fig. 45; * Obs., V, 1852, p. 41, pl. XXV, fig. 45.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN,

Conch. Cab. Ano., 1875, p. 210, pl. LXIV, figs. 4, 5.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 178.

* *Margaron (Anodonta) denigrata* LEA, Syn., 1852, p. 50; 1870, p. 79.

* † *Anodonta oblita* LEA, Tr. Am. Phil. Soc., X, 1-52, p. 290, pl. XXVIII, fig. 52; * Obs., V, 1852, p. 46, pl. XXVIII, fig. 52.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. Ano., 1875, p. 165, pl. LIV, figs. 7, 8.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Anodonta) oblita* LEA, Syn., 1852, p. 50; 1870, p. 79.

* *Anodon subcylindracea* SOWERBY, Conch. Icon., XVII, 1867, pl. XIV, fig. 47.

* *Anodonta subcylindracea* CLESSIN, Conch. Cab. Ano., 1873, p. 92, pl. XXVI, figs. 6, 7.

† ANODONTOIDES FERUSSACIANUS var. SUBCYLINDRACEUS Lea.

* † *Anodonta subcylindracea* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 106, pl. XXIV, fig. 117;

* Obs., II, 1838, p. 106, pl. XXIV, fig. 117.—* TROSCHEL, Arch. für Nat., V, 1839, Pt. 2, p. 239.—* HANLEY, Test. Moll., 1842, p. 223; * Biv. Shells, 1843, p. 223.—* STIMPSON, Shells of N. Eng., 1851, p. 15.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 55.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 185.—* WALTON, Moll. Monroe Co., 1892, p. 17, pl. VIII, fig. 3.

* *Margarita (Anodonta) subcylindracea* LEA, Syn., 1838, p. 32.

* *Anodon subcylindracea* DE KAY, Zool. of N.Y., Pt. 5, 1843, p. 200, pl. XVI, fig. 229.—* CATLOW and REEVE, Conch. Nom., 1845, p. 68.

* *Margaron (Anodonta) subcylindracea* LEA, Syn., 1852, p. 51; 1870, p. 82.

* *Anodontoides subcylindraceus* BAKER, Moll. Chicago, Pt. 1, 1898, p. 74, pl. IV, fig. 4; VI, fig. 1.

ANODONTOIDES FERUSSACIANUS var. MODESTUS Lea.

* † *Anodonta modesta* LEA, Pr. Ac. Nat. Sci. Phila., 1857, p. 84; * JI. Ac. Nat. Sci. Phila., IV, 1860, p. 364, pl. LXIII, fig. 189; * Obs., VIII, 1860, p. 46, pl. LXIII, fig. 189.—* CLESSIN, Conch. Cab. Ano., 1874, p. 153, pl. XLIX, figs. 7, 8.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 182.

* *Margaron (Anodonta) modesta* LEA, Syn., 1879, p. 79.

* *Anodon modestus* SOWERBY, Conch. Icon., XVII, 1867, pl. x, fig. 26.

Mississippi drainage area generally; St. Lawrence River system; Red River of the North; Saskatchewan River; Connecticut. ? The varieties are found in the St. Lawrence system.

Genus PEGIAS Simpson, 1900.

(Type, *Margaritana fabula* Lea.)

Shell small, thickened in front, with a sharp posterior ridge, in front of which is a wide radial depression, ending in a basal sinus; above this ridge is another, making the shell decidedly biangulate and truncate behind; beak sculpture consisting of subconcentric corrugations, generally swollen on the posterior ridge; epidermis decorticated, but showing a few dark, radial rays on the base of the shell; pseudocardinals rather solid; laterals wanting. Animal having the marsupium empty in the specimen examined; inner gills larger in front, free from the abdominal sac, all four truncated behind; palpi large; mantle with square spots on its edge; branchial opening with a few large papillæ; anal opening smooth, not separated from the superanal.

† PEGIAS FABULA Lea.

- * *Margarita (Margaritana) fabula*, LEA, Syn., 1836, p. 46; 1838, p. 28.
- * *Margaritana fabula* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 44, pl. XIII, fig. 39; * Obs., II, 1838, p. 44, pl. XIII, fig. 39.—* TROSCHEL, Arch. für Nat., V, 1839, 1^{te} 2, p. 238.—* KUSTER, Conch. Cab. Unio, 1862, p. 301, pl. c, fig. 3.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 173.
- * *Unio fabula* HANLEY, Test. Moll., 1842, p. 213; Biv. Shells, 1843, p. 213, pl. XXII, fig. 45.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVI, fig. 394.
- * *Margaron (Margaritana) fabula* LEA, Syn., 1852, p. 44; 1870, p. 70.
- * *Micromya fabula* AGASSIZ, Arch. für Nat., I, 1852, p. 47.
- * *Strophitus fabula* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 263.
- * *Baphia fabula* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 499.
- * † *Margaritana curreyana* LEA, Pr. Ac. Nat. Sci. Phila., I, 1840, p. 288; * Tr. Am. Phil. Soc., VIII, 1842, p. 223, pl. XVIII, fig. 40; * Obs., III, 1842, p. 61, pl. XVIII, fig. 40.—* KUSTER, Conch. Cab. Unio, 1862, p. 300, pl. c, fig. 2.—* B. H. WRIGHT, Check List, 1888.
- * *Micromya curreyana* AGASSIZ, Arch. für Nat. I, 1852, p. 47.
- * *Margaron (Margaritana) curreyana* LEA, Syn., 1852, p. 42; 1870, p. 68.
- * *Strophitus curreyana* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 263.
- * *Baphia curreyana* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 499.
- * *Unio curreyana* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIII, fig. 319.
- * *Unio curreyanus* HANLEY, Biv. Shells, 1856, p. 386, pl. XXIV, fig. 10.
- * *Margaritana curreyana* PÆTEL, Conch. Sam., III, 1890, p. 173.

Cumberland and Tennessee river systems.

Genus ARCIDENS Simpson, 1900.

(Type, *Alasmodonta confragosa* Say.)

Shell subsolid, inflated, subrhomboid, with full, high beaks; beak sculpture very strong, consisting of irregular corrugations which fall into two loops, at the bases of which the ridges are swollen into knobs that continue out in two radiating rows on to the disk of the shell; in front of and behind the beaks are many fine, radial wrinkles, the posterior ones being zigzagged; surface of the shell covered with oblique folds and wrinkles; epidermis dark olive, shining; there are two elongated, compressed pseudocardinals in the left valve, the posterior under the beak, and curved upward, cutting off the hinge plate in the right valve, which has a single, compressed pseudocardinal in front; laterals numerous, short, blurred; muscle scars irregular; naere white. Animal with the gills very large, rounded below; inner the larger, free from the abdominal sac; marsupium filling the outer gills, of a peculiar, granular texture; palpi very large, elongated, attached half way to the mantle; mantle with a thick, unspotted border; branchial opening large, papillose; anal opening with rudimentary papillæ.¹

¹ In the only gravid specimen I have seen the marsupium presented a peculiar, striated, granular structure, different from that of any Naiad I know. It was narrowed in the middle, and produced into a sort of lobe behind, but, though the specimen seemed to be a perfectly normal one, I am in doubt whether this peculiar form is constant. I do not think that this granulation is the result of a diseased condition, though it may be.

† ARCIDENS CONFRAGOSUS Say.

- * *Alasmodonta confragosa* SAY, N. Harin. Diss., II, 1829, p. 339; * Am. Conch., I, 1830, pl. XXI; IV, 1832 (cover p. 4).—* CONRAD, New F. W. Shells, 1834, p. 72.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 30, pl. VIII, figs. 1, 1a.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 262.
- * *Margarita (Margaritana) confragosa* LEA, Syn., 1836, p. 43; 1838, p. 27.
- * *Unio confragosa* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 552; 3d ed., II, 1837, p. 674.—* HANLEY, Test. Moll., 1842, p. 210; * Biv. Shells, 1843, p. 210, pl. XXIII, fig. 52.—* DESHAYES, Tr. Elem., 1853, p. 217, pl. XXX, figs. 5-7.
- * *Baphia confragosa* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 500.
- * *Margarita (Margaritana) confragosa* LEA, Syn., 1852, p. 42; 1870, p. 67.
- * *Margaritana confragosa* CALKINS, Pr. Ottawa Acad., 1874, p. 46.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 172.
- * *Unio confragosus* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* SOWERBY, Conch. Icon., XVI, 1867, pl. LX, fig. 299.

Mississippi River and States adjoining it; Ohio River drainage; southwest to the Colorado River, Texas. A specimen is in the Lea collection, said to be from Enterprise, southeastern Alabama, but I am doubtful about the locality.

Genus SYMPHYNOTA Lea, 1829.

(Type, *Symphynota compressa* LEA.)

Symphynota LEA, Tr. Am. Phil. Soc., III, 1829, p. 424.

? *Megadomus* SWAINSON, Tr. on Mal., 1840, p. 266.

? *Complanaria* SWAINSON, Tr. on Mal., 1840, p. 290.

Shell elliptic rhomboid, compressed; beaks low, their sculpture consisting of strong bars; one pseudocardinal in the right valve and two in the left, the hinder somewhat A-shaped, cutting off the hinge plate in the right valve; laterals generally imperfect.

Animal with the gills semicircular below, inner the larger, free nearly the whole length from the abdominal sac; marsupium thick, padlike, filling the outer gills; mantle strongly attached at pallear line; branchial opening papillose; anal opening without papillæ.

Subgenus SYMPHYNOTA Lea, 1829.

Shell smooth, subsolid, shining, rayed; teeth delicate; laterals compressed, moderately developed.

SYMPHYNOTA COMPRESSA Lea.

- * *Symphynota compressa* LEA, Tr. Am. Phil. Soc., 1829, p. 450, pl. XII, fig. 22; * Obs., I, 1834, p. 64, pl. XII, fig. 22.—* FERUSSAC, Guer. Mag., 1835, p. 25.
- * *Complanaria compressa* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 261.
- * *Microcondylaea compressa* PÆTEL, Conch. Sam., III, 1890, p. 175.
- * *Unio compressus* CONRAD, New F. W. Shells, 1834, p. 68; * Monog., VIII, 1837, p. 71, pl. XI, fig. 1.—* C. B. ADAMS, Thompson's Hist. Vt., 1842, p. 166.—* HANLEY, Test. Moll., 1842, p. 174; * Biv. Shells, 1843, p. 174, pl. XX, fig. 39.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 191, pl. XXI, fig. 245.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* KUSTER, Conch. Cab. Unio, 1861, p. 198, pl. LXIV, figs. 3, 4.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXI, fig. 303.

- * *Margarita (Unio) compressus* LEA, Syn., 1836, p. 11; 1838, p. 13.
- * *Complanaria compressus* AGASSIZ, Arch. für. Nat., I, 1852, p. 48.
- * *Complanaria alasmodontina* STIMPSON, Shells of N. Eng., 1851, p. 14.
- * *Unio alasmodontinus* PÆTEL, Conch. Sam., III, 1890, p. 144.
- * *Margaron (Unio) pressus* LEA, Syn., 1852, p. 19; 1870, p. 29.¹
- * *Unio pressus* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* LATCHFORD, Tr. Ottawa F. N. Club., 1882, p. 52.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.
- * *Alasmodonta pressa* BAKER, Tr. Ac. N. Sci. St. L., VIII, 1898, p. 76, fig. ; * Moll. Chicago, Pt. 1, 1898, p. 58, pl. vi, fig. 3; x, fig. 4.

SYMPHYNOTA COMPRESSA var. PLEBIUS C. B. Adams

- * *Unio compressus* var. *plebius* C. B. ADAMS, F. W. and L. Shells of Vt., p. 16.

Ohio and St. Lawrence drainage areas; west to Arkansas, north through Nebraska to Wisconsin; Hudson River.

† SYMPHYNOTA CHARLOTTENSIS Lea.

- * *Unio charlottensis* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 191; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 8, pl. II, fig. 5; * Obs., XI, 1867, p. 12, pl. II, fig. 5.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) charlottensis* LEA, Syn., 1870, p. 51.

Near Charlotte, North Carolina.

† SYMPHYNOTA VIRIDIS Conrad.

- * *Unio viridis* CONRAD, Monog., IV, 1836, p. 35, pl. XVII, fig. 1;² * Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLV, fig. 244.—* B. H. WRIGHT, Check List, 1888.
- * *Margarita (Unio) tappanianus* LEA, Syn., 1836, p. 39; 1838, p. 25.
- * *Unio tappanianus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 62, pl. XVII, fig. 55; * Obs. II, 1838, p. 62, pl. XVII, fig. 55.—* TROSCHER, Arch. für Nat., V, 1839, Pt. 2, p. 236.—* HANLEY, Test. Moll., 1842, p. 209; Biv. Shells, 1843, p. 209, pl. XXI, fig. 38.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 194, pl. XX, fig. 242.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* CHENU, Ill. Conch., 1858, pl. XX, figs. 2, 2a, 2b.—* KUSTER, Conch. Cab. Unio, 1862, p. 270, pl. XCI, fig. 3.—* HARTMAN and MICHENER, Conch. Test., 1874, p. 89, fig. 185.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 169.
- * *Margaron (Unio) tappanianus* LEA, Syn., 1852, p. 39; 1870, p. 62.
- * *Unio tappianus* CATLOW and REEVE, Conch. Nom., 1845, p. 64.
- * † *Unio hyalinus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; * Tr. Am. Phil. Soc., X, 1848, p. 69, pl. II, fig. 4; Obs., IV, 1848, p. 43, pl. II, fig. 4.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* KUSTER, Conch. Cab. Unio, 1861, p. 204, pl. LXVIII, fig. 3.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
- * *Margaron (Unio) hyalinus* LEA, Syn., 1852, p. 39; 1870, p. 62.

¹ Lea changed his name *compressa* to *pressus* because the former had been used for a *Unio* previously by Sowerby. The restoration of the genus *Symphynota* makes the earlier name of Lea valid.

² Credited by Conrad and others to Rafinesque. The description of the latter of *Unio viridis* is not sufficiently clear to distinguish it from some of the other members of this genus, and the locality given is in error for the shell figured by Conrad, which belongs exclusively to the Atlantic drainage, and not to the Ohio River and Kentucky, localities given by Rafinesque.

- *† *Unio pertenuis* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 193; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 8, pl. II, fig. 4; *Obs., XI, 1867, p. 12, pl. II, fig. 4.—* B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) pertenuis* LEA, Syn., 1870, p. 62.

Streams draining into the Atlantic from New York south to North Carolina; Monroe County, Michigan?

† SYMPHYNOTA DECORATA Lea.

- * *Unio decoratus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 257, pl. XIII, fig. 6; *Obs., V, 1852, p. 13, pl. XIII, fig. 6.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—*SOWERBY, Conch. Icon., XVI, 1868, pl. XCI, fig. 496.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 150.

**Margaron (Unio) decoratus* LEA, Syn., 1852, p. 19; 1870, p. 29.

- *† *Unio insolidus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 159; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 40, pl. XIII, fig. 37; *Obs., XIII, 1874, p. 44, pl. XIII, fig. 37.—*B. H. WRIGHT, Check List, 1888.

Virginia to South Carolina.

† SYMPHYNOTA NEGLECTA Lea.

- Unio neglectus* LEA, Desc. 12 Uniones, 1843; *Tr. Am. Phil. Soc., IX, 1846, p. 280, pl. XLII, fig. 10; *Obs., IV, 1848, p. 88, pl. XLII, fig. 10.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*KUSTER, Conch. Cab. Unio, 1861, p. 225, pl. LXXVI, fig. 4.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 160.

**Margaron (Unio) neglectus* LEA, Syn., 1852, p. 33; 1870, p. 54.

Northern Alabama.

† SYMPHYNOTA QUADRATA Lea.

- **Margaritana quadrata* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 41; *Jl. Ac. N. Sci. Phila., II, 1862, p. 210, pl. XXXII, fig. 279; *Obs., IX, 1863, p. 32, pl. XXXII, fig. 279.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 173.

**Margaron (Margaritana) quadrata* LEA, Syn., 1870, p. 68.

- **Margaritana (Alasmodonta) quadrata* CLESSIN, Conch. Cab. Ano., 1876, p. 273, pl. LXXXIII, figs. 5, 6.

Eastern Tennessee.

Subgenus **LASMIGONA** Rafinesque, 1831.

(Type, *Alasmodonta costata* Rafinesque.)¹

Shell subrhomboid, compressed, corrugated behind; beaks low, their sculpture consisting of several coarse ridges which generally fall into two slight loops, and often with radiating ridges in front and behind; epidermis shining; laterals partly developed, consisting of blurred ridges which slope diagonally downward and backward on the hinge plate; cavities of the beaks shallow; dorsal scars faint, irregular.

¹The description of *Alasmodonta costata* of Rafinesque so clearly covers the species described three years later as *Alasmodonta rugosa* by Barnes that the former name must be used.

Animal with the marsupium occupying the whole of the outer gills, very thick, like that of *Anodonta*; inner gills free all or the greater part of their length from the abdominal sac; branchial opening papillose; anal opening generally smooth or only slightly crenulate.

†SYMPHYNOTA COSTATA Rafinesque.

**Alasmodonta costata* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 313, pl. LXXXII, figs. 15, 16.

**Alasmodonta costata* SAY, Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 72.—* FERUSSAC, Guer. Mag., 1835, p. 25

**Complanaria costata* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 261.

**Alasmodonta rugosa* BARNES, Am. Jl. Sci., VI, 1823, p. 278, pl. XIII, fig. 21.—SHORT and EATON, Transylvania Jl., 1831, p. 81.—* C. B. ADAMS, Thompson's Hist. of Vt., 1842, p. 165.—* BAKER, Moll. Chicago, Pt. 1, 1898, p. 57, pl. VII, figs. 1, 2; VIII, figs. 3, 4.

**Mya rugosa* EATON, Zool. Text-Book, 1826, p. 222.

**Margarita (Margaritana) rugosa* LEA, Syn., 1836, p. 44; 1838, p. 27.

**Unio rugosa* HANLEY, Test. Moll., 1842, p. 211; * Biv. Shells, 1843, p. 211, pl. XXI, fig. 8.

**Alasmodon rugosa* DE KAY, Zool. N. Y., Pt. 5, 1843, p. 196, pl. XIV, fig. 226.

**Complanaria rugosa* STIMPSON, Shells of N. Eng., 1851, p. 14.

Margaron (Margaritana) rugosa LEA, Syn., 1852, p. 42; 1870, p. 67.

**Baphia rugosa* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 500.

**Margaritana rugosa* CALKINS, Pr. Ottawa Ac. Sci., 1874, p. 46.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.

**Lasmigona rugosum* RAFINESQUE, Cont. Mon., Og., 1831, p. 5.

**Unio rugosus* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—KUSTER, Conch. Cab. Unio, 1861, p. 200, pl. LXVI, figs. 1-3.—SOWERBY, Conch. Icon., XVI, 1867, pl. LX, fig. 302.

**Alasmodonta hians* FERUSSAC, Guer. Mag., 1835, p. 25.

Mississippi drainage generally; St. Lawrence basin; Manitoba; Hudson River?; Columbus, Mississippi. Texas?

Subgenus **PTEROSYGNA** Rafinesque, 1831.

(Type, *Alasmodonta complanata* Barnes.)

Shell large, ovate-rhomboid, inflated in post basal region; beaks much compressed, their sculpture sharply and strongly doubly looped; epidermis dark, scarcely rayed; teeth very heavy.

Animal with an extremely thick marsupium; palpi large; mantle with small crenulations on lower border; branchial opening large; anal opening small, both slightly papillose.

†SYMPHYNOTA COMPLANATA Barnes.

**Alasmodonta complanata* BARNES, Am. Jl. Sci., VI, 1823, p. 278, pl. XIII, fig. 21.—

* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 289.—* CONRAD, New F. W. Shells, 1834, p. 72.—* SAY, Am. Conch., VI, 1834.—* BAKER, Moll. Chicago, Pt. 1, 1898, p. 60, pl. VIII, figs. 1, 2; IX, figs. 1-4.

**Mya complanata* EATON, Zool. Text-Book, 1826, p. 222.

**Symphynota complanata* LEA, Tr. Am. Phil. Soc., III, 1830, p. 448.—* SHORT and EATON, Transylvania Jl., 1831, p. 80.—* LEA, Obs., I, 1834, p. 62.—FERUSSAC, Guer. Mag., 1835, p. 25.

* *Margarita (Margaritana) complanata* LEA, Syn., 1836, p. 43; 1838, p. 26.

* *Unio complanata* HANLEY, Test. Moll., 1842, p. 210; * Biv. Shells, 1843, p. 210, pl. XXI, fig. 9.

* *Complanaria complanata* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 261.

* *Baphia complanata* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 500.

* *Margaritina complanata* CALKINS, Pr. Ottawa Acad., 1874, p. 46.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 172.

* *Alasmodon complanatus* SOWERBY, Conch. Man., 1842, p. 61, fig. 140.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* KUSTER, Conch. Cab. Unio, 1861, p. 199, pl. LXV, fig. 1.—* SOWERBY, Conch. Icon., XVI, 1866, pl. L, fig. 266.

Complanaria gigas SOWERBY, Conch. Man., 1839, fig. 141, *; 1842, p. 115, fig. 141.

* ? *Megadomus gigas* SWAINSON, Tr. on Mal., 1840, pp. 265, 378.

† SYMPHYNOTA COMPLANATA var. KATHERINÆ Lea.

Unio katherinæ LEA, Syn., 1838, p. 35; ¹ Tr. Am. Phil. Soc., VI, 1839, p. 143.—* TROSCHEL, Arch. für Nat., V, 1839, II, p. 237.

Upper Mississippi drainage, as far south as Arkansas on the west; Ohio River system; upper St. Lawrence and its tributaries; north into the Mackenzie River.

Genus ALASMIDONTA Say, 1818.

(Type, *Monodonta undulata* Say.)²

Monodonta SAY, Nich. Enc., II, 1816.

Alasmidonta SAY, Jl. Ac. N. Sci. Phila., I, 1818, p. 459.

Alasmodonta SAY, Nich. Enc., 3d ed., IV, 1819.

Alasmodon SWAINSON, Tr. on Mal., 1840, p. 382.

Unioopsis SWAINSON, Tr. on Mal., 1840, p. 382.

Shell generally rhomboid, inflated, with a well-developed posterior ridge, which ends in a point behind when it is single or a biangulation when double; beaks full and high, with coarse, concentric or slightly doubly-looped bars; epidermis rayed, shining; hinge with two pseudo-cardinals in the left valve and one in the right; laterals wanting or imperfect; cavity of the beaks deep; dorsal scars under the hinge plate; nacre bluish.

Animal with the marsupium occupying the entire outer gills; gills rounded below, inner much the larger, united to the abdominal sac, or free; mantle generally having square spots on the posterior border; branchial opening papillose; anal opening smooth or crenulate.

Subgenus ALASMIDONTA Say, 1818.

(Type, *Monodonta undulata* Say.)

Shell ovate-rhomboid, solid, inflated, shining, with very strong, generally concentric beak sculpture; pseudocardinals solid, stumpy, some-

¹ This is probably the small, peculiar boreal form of this species, which appears very different from the type.

² Changed by Say afterwards to *Alasmidonta undulata* as the name *Monodonta* was preoccupied. For some reason Say later on called his genus *Alasmodonta*.

what radiately ridged; laterals short, very imperfect, or wanting; beak cavities deep, compressed.

Animal with inner gills wider than the outer throughout; palpi long; marsupium loose and flabby, and having ova scattered throughout it in the only gravid specimen seen.

† *ALASMIDONTA UNDULATA* Say.

Monodonta undulata SAY, Nich. Enc., 1816, pl. III, fig. 3.

* *Alasmidonta undulata* SAY, Jl. Acad. N. Sci. Phila., I, 1818, p. 460.

* *Alasmodonta undulata* SAY, Nich. Enc., IV, 1819, pl. III, fig. 3.—* CONRAD, New F. W. Shells, 1834, p. 73.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* C. B. ADAMS, F. W. and L. S. of Vt., 1842, p. 15.

* *Alasmodon undulata* SWAINSON, Tr. on Mal., 1840, p. 288, fig. 61.—* GOULD, Inv. Mass., 1841, p. 115, fig. 76.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 198, pl. XV, fig. 227.

* *Margarita (Margaritana) undulata* LEA, Syn., 1836, p. 44; 1838, p. 27.

* *Unio undulata* HANLEY, Test. Moll., 1842, p. 211.—* Biv. Shells, 1843, p. 211.

* *Margaritana undulata* KUSTER, Conch. Cab. Unio, 1862, p. 298, pl. XCIX, fig. 4.—* GOULD, Inv. Mass., 1870, p. 176, fig. 478.—* HARTMAN and MICHENER, Conch. Cest., 1874, p. 92, fig. 190.—* LATCHFORD, Tr. Ottawa F. N. Cl., 1882, p. 54.—* H. CARPENTER, Naut., IV, 1890, p. 47.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 174.

* *Margaron (Margaritana) undulata* LEA, Syn., 1852, p. 42; 1870, p. 68.

* *Mya undulata* WOOD, Ind. Test. Rev., 1856, p. 199, pl. I, supp., fig. 5.

* *Baphia undulata* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 499.

* *Unio undulatus* CATLOW and REEVE, Conch. Nom., 1845, p. 65.

* *Strophitus undulatus* STIMPSON, Shells of New Eng., 1851, p. 15.

* † *Unio glabratus* SOWERBY, Rec. and Foss. Shells, XVI, 1823, fig.

Alasmodonta sculptilis SAY, N. Harmony Diss., II, 1829, p. 339.—FERUSSAC, Guer. Mag., 1835, p. 26.—L. W. SAY, Terr. and Fluv. Shells, 1840, p. 9.

* *Strophitus sculptilis* STIMPSON, Shells of New Eng., 1851, p. 15.

* *Unio hians* VALENCIENNES Rec. Obs. Zool., II, 1833, p. 235, pl. LIV, figs. 2a, 2b.

* *Unioopsis radiata* SWAINSON, Tr. on Mal., 1840, p. 289, fig. 62.

* *Unioopsis mytiloides* SWAINSON, Tr. on Mal., 1840, p. 382, fig. 62.

* † *Unio swainsoni* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVI, fig. 396.

* *Margaron (Margaritana) swainsoni* LEA, Syn., 1870, p. 30.

Lower St. Lawrence drainage; southward to North Carolina.

Subgenus *PRESSODONTA* Simpson, 1900.

(Type, *Unio calceolus* Lea.¹)

Shell decidedly rhomboid, surface generally painted with unbroken rays; beak sculpture slightly corrugated; teeth compressed.

Animal with a thick, smooth, pad-like marsupium.

(Group of *Alasmidonta calceola*.)

Shell small, subsolid, rhomboid, inflated, rayed but not shining; beak sculpture varying from strong concentric bars to irregular corrugations, sometimes faintly doubly looped, and with fine radiating

¹The names *Hemiodon*, *Hemiodonta*, and *Hemidon* were applied by Swainson to this group, but he used the first name previously for a division of the Helicidae. It is better to apply a new name.

ridges in front and behind; pseudocardinals rather solid; laterals very faint; marsupium occupying the whole of the outer gills, thick, pad-like; inner gills wholly or partly free from the abdominal sac; mantle bordered with square, black spots.

† *ALASMIDONTA CALCEOLA* Lea.

* *Unio calceolus* LEA, Tr. Am. Phil. Soc., III, 1830, p. 265, pl. III, fig. 1; * Obs., I, 1834, p. 7, pl. III, fig. 1.—* FERUSSAC, Gner. Mag., 1835, p. 26.—* CHENU, III. Conch., 1858, pl. VIII, figs. 2, 2a, 2c.—* KUSTER, Conch. Cab. Unio, 1861, p. 187, pl. LIX, fig. 3.

* *Strophitus calceolus* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 262.

* *Unio calceola* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 546; 3d ed., II, 1839, p. 672.—* HANLEY, Test. Moll., 1842, p. 212; * Biv. Shells, 1843, p. 212, pl. XXII, fig. 29.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVIII, fig. 410.

* *Margarita (Margaritana) calceola* LEA, Syn., 1836, p. 45; 1838, p. 27.

* *Margaron (Margaritana) calceola* LEA, Syn., 1852, p. 43; 1870, p. 68.

* *Margaritana calceola* KUSTER, Conch. Cab. Unio, 1862, p. 299, pl. XCIX, fig. 6.—* B. H. WRIGHT, Check List, 1883.—* PÆTEL, Conch. Sam., III, 1890, p. 172.

* *Baphia calceola* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 500.

* *Alasmodonta truncata* CONRAD, New F. W. Shells, 1834, p. 73.

* *Margarita (Margaritana) deltoidea* LEA, Syn., 1836, p. 44; 1838, p. 27.

* *Margaritana deltoidea* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 43, pl. XIII, fig. 38; Obs., II, 1838, p. 43, pl. XIII, fig. 38.—* TROSCHEL, Arch. für Nat., V, 1839, II, p. 238.—* CHENU, Man., 1859, II, p. 144, fig. 711.—* KUSTER, Conch. Cab., 1862, p. 299, pl. XCIX, fig. 5.—* CALKINS, Pr. Ottawa Ac. Sci., 1874, p. 46.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 173.

* *Unio deltoidea* HANLEY, Test. Moll., 1842, p. 211; * Biv. Shells, 1843, p. 211, pl. XXII, fig. 50.

* *Margaron (Margaritana) deltoidea* LEA, Syn., 1852, p. 42, 1870, p. 67.

* *Strophitus deltoidea* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 263.

* *Baphia deltoidea* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 499.

* *Alasmodonta deltoidea* BAKER, Moll. Chicago, Pt. 1, 1898, p. 63, pl. VI, fig. 2; VII, fig. 4.

* *Unio deltoideus* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVI, fig. 395.

Upper Mississippi drainage; Ohio, Cumberland, and Tennessee rivers; Lower and Middle St. Lawrence systems.

† *ALASMIDONTA MINOR* Lea.

* *Margaritana minor* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 166; Tr. Am. Phil. Soc., X, 1848, p. 82, pl. VIII, fig. 26; * Obs., IV, 1848, p. 56, pl. VIII, fig. 26.—* CHENU, Man., 1859, II, p. 144, fig. 713.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 173.

* *Strophitus minor* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 263.

* *Margaron (Margaritana) minor* LEA, Syn., 1852, p. 42; 1870, p. 67.

* *Baphia minor* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 499.

Tennessee and Cumberland river systems.

(Group of *Alasmodonta heterodon*.)

Shell small, rather thin, rhomboid; beak sculpture very strong, consisting of concentric ridges or irregular corrugations, often slightly

doubly looped and swollen on the well-developed posterior ridge; surface rayed; pseudocardinals compressed, reflexed; laterals single, double, or triple in each valve.

Animal with branchial opening having a few coarse papillæ; anal opening smooth, or with very fine crenulations.

†ALASMIDONTA HETERODON Lea.

- **Unio heterodon* LEA, Tr. Am. Phil. Soc., III, 1830, p. 428, pl. VIII, fig. 11; *Obs., I, 1834, p. 42, pl. VIII, fig. 11.—*CONRAD, New F. W. Shells, 1834, p. 69.—*FERUSSAC, Guer. Mag., 1835, p. 26.—*DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 548; 3d ed., II, 1839, p. 672.—*CONRAD, Monog., X, 1838, p. 90, pl. XLIX, fig. 3.—*HANLEY, Test. Moll., 1842, p. 183; Biv. Shells, 1843, p. 183, pl. XXI, fig. 42.—*CATLOW and REEVE, Conch. Nom., 1845, p. 59.—*CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 250.—*CHENU, Ill. Conch., 1858, pl. XVI, figs. 2, 2a, 2b.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*KUSTER, Conch. Cab., 1861, p. 192, pl. LXI, fig. 2.—*SOWERBY, Conch. Icon., XVI, 1866, pl. XXXV, fig. 184.—*HARTMAN and MICHENER, Conch. Cest., 1874, p. 91, fig. 188.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 155.

**Margarita (Unio) heterodon* LEA, Syn., 1836, p. 18; 1838, p. 16.

**Margaron (Unio) heterodon* LEA, Syn., 1852, p. 24; 1870, p. 36.

Northern New England to Virginia.

ALASMIDONTA DIVERSA Conrad.

- **Unio diversus* CONRAD, Am. Jl. Sci., XXI, 1856, p. 172 (fig. in outline).—B. H. WRIGHT, Check List, 1888.¹

Alabama.

(Group of *Alasmidonta collina*.)

Shell rhomboid, subsolid, with beak sculpture consisting of strong, concentric bars, sometimes slightly doubly looped and swollen on the low, posterior ridge; often with one or more small spines on the disk in front of the posterior ridge; epidermis brown, rayless or feebly rayed; rest marks well defined; teeth rather compressed; laterals lamellar.

Animal unknown.

†ALASMIDONTA COLLINA Conrad.

- **Unio collinus* CONRAD, Monog., VIII, 1837, p. 65, pl. XXXVI, fig. 2; XII, 1840, p. 109, pl. LX, fig. 3.—*TROSCHER, Arch. für Nat., IV, 1838, p. 287.—*CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 247.—*HANLEY, Biv. Shells, 1856, p. 382, pl. XX, fig. 57.—*KUSTER, Conch. Cab. Unio, 1861, p. 188, pl. LIX, fig. 5.—*SOWERBY, Conch. Icon., XVI, 1866, pl. XLIX, fig. 263.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 148.
- **Margaron (Unio) collinus* LEA, Syn., 1852, p. 23; 1870, p. 35.

North Carolina and Virginia.

¹ Conrad says this has a remarkable resemblance to *Unio heterodon*, but is shorter.

(Group of *Alasmodonta holstonia*.)

Shell subsolid, beak sculpture rather strong, decidedly doubly looped, surface brownish, slightly rayed; pseudocardinals delicate; laterals nearly or quite wanting. Animal like that of *A. calceola*.

† *ALASMODONTA HOLSTONIA* Lea.

- * *Margaritana holstonia* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 42, pl. XIII, fig. 37;
 * Obs., II, 1838, p. 42, pl. XIII, fig. 37.—* TROSCHEL, Arch. für Nat., V, 1839, Pt. 2,
 p. 238.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890,
 p. 173.
 * *Margarita (Margaritana) holstonia* LEA, Syn., 1836, p. 46; 1838, p. 28.
 * *Margaron (Margaritana) holstonia* LEA, Syn., 1852, p. 44; 1870, p. 70.
 * *Strophitus holstonia* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.
 * *Baphia holstonia* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 499.
 * *Unio holstonianus* HANLEY, Biv. Shells, 1843, p. 213, pl. XXII, fig. 44.—* CATLOW
 and REEVE, Conch. Nom., 1845, p. 60.—* SOWERBY, Conch. Icon., XVI, 1868, pl.
 LXXVI, fig. 398.
 * *Margaritana holstoniana* KUSTER, Conch. Cab. Unio, 1862, p. 302, pl. c, fig. 4.
 * ? *Unio striatus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVIII, fig. 407.

Tennessee River system; headwaters of the Coosa River.

† *ALASMODONTA GEORGIANA* Lea.¹

- * *Margaritana etowahensis* LEA, Pr. Ac. Nat. Sci. Phila., II, 1858, p. 138; * JI. Ac.
 Nat. Sci. Phila., IV, 1859, p. 227, pl. XXXI, fig. 110; * Obs., VII, 1859, p. 45, pl.
 XXXI, fig. 110.—* B. H. WRIGHT, Check List, 1888.
 * *Margaritana (Alasmodonta) etowahensis* CLESSIN, Conch. Cab. Aro., 1875, p. 270,
 pl. LXXXI, figs. 1, 2.
 * *Margaritana etowahensis* PÆTEL, Conch. Sam., III, 1890, p. 173.
 * *Margaritana georgiana* LEA, Pr. Ac. Nat. Sci. Phila., III, 1859, p. 280.—B. H.
 WRIGHT, Check List, 1888.
 * *Margaron (Margaritana) georgiana* LEA, Syn., 1870, p. 68.
 * *Alasmodonta impressa* ANTHONY, Am. JI. Conch., I, 1865, p. 157, pl. XII, fig. 4.

Tennessee; Etowah River, Georgia.

Subgenus *RUGIFERA* Simpson, 1900.

(Type, *Alasmodonta marginata* Say.)

Shell elongated, rhomboid, inflated, surface brilliantly painted with radiations, which often break into a dappled or splashed pattern of color; posterior slope slightly corrugated; teeth very imperfect; laterals wanting.

Animal with a thick, smooth, pad like marsupium. Mantle with square spots behind.

† *ALASMODONTA MARGINATA* Say

- * *Alasmodonta marginata* SAY, Nich. Enc., 1819, No. 1; Am. Conch., VI, 1834.—
 * CONRAD, New F. W. Shells, 1834, p. 72.—* FERUSSAC, Guer. Mag., 1835, p.
 25.—* STIMPSON, Shells of N. Eng., 1851, p. 15.—* AGASSIZ, Arch. für Nat., I,
 1852, p. 46.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 262.

¹ First called *Margaritana etowahensis* by Dr. Lea, but as that name had previously been used by Conrad its author changed it to *georgiana*.

- * *Alasmodonta marginata* RAFINESQUE, An. Gen. Sci. Brux., V, 1820, p. 317.
 * *Margarita (Margaritana) marginata* LEA, Syn., 1836, p. 43; 1838, p. 27.
 * *Alasmodon marginata* GOULD, Inv. Mass., 1841, p. 116, fig. 77.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 196, pl. xiv, fig. 225.
 * *Unio marginata* HANLEY, Test. Moll., 1842, p. 211; * Biv. Shells, 1843, p. 211, pl. XXI, fig. 23.
 * *Anodonta marginata* C. B. ADAMS, Thompson's Hist. Vt., 1842, p. 164; * F. W. and L. S. of Vt., 1842, p. 14.—* STIMPSON, Shells of N. Eng., 1851, p. 15.
 * *Margaron (Margaritana) marginata* LEA, Syn., 1852, p. 42; 1870, p. 67.
 * *Baphia marginata* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 500.
 * *Margaritana marginata* GOULD, Inv. of Mass., 1870, p. 177, fig. 479.—HARTMAN and MICHENER, Conch. Cest., 1874, p. 93, fig. 191.—* LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 54.—* B. H. WRIGHT, Check List, 1888.—* H. CARPENTER, Naut., IV, 1890, p. 46.—* PÆTEL, Conch. Sam., III, 1890, p. 173.
 * *Unio marginatus* CATLOW and REEVE, Conch. Nom., 1845, p. 61.
 * *Unio varicosa* LAMARCK, An. sans Vert., VI, 1819, p. 78.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 543; 3d ed., II, 1839, p. 671.
 * *Unio cariosa* LAMARCK, An. sans Vert., VI, 1819, p. 80.
 * *Alasmodon corrugata* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 198, pl. XXIV, fig. 259.
 * *Mya rugulosa* WOOD, Ind. Test., 3d ed., 1856, p. 199, pl. I, supp. fig. 7.

Lower St. Lawrence; southward in streams draining into the Atlantic to South Carolina.

† **ALASMIDONTA TRUNCATA** B. H. Wright.¹

- * *Alasmodonta truncata* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 262.
 * *Margaritana marginata* KUSTER, Conch. Cab. Unio, 1862, p. 297, pl. XCIX, fig. 3.—* CALKINS, Pr. Ottawa Acad., 1874, p. 46.
 * *Alasmodonta marginata* BAKER, Moll. Chicago, Pt. 1, 1898, p. 62, pl. IV, fig. 4; VII, fig. 7; XXII, fig. 3.
 * *Margaritana marginata* var. *truncata* B. H. WRIGHT, Naut., XI, 1898, p. 124.
 * *Unio marginatus* SOWERBY, Conch. Icon., XVI, 1866, pl. LI, fig. 267.

Upper Mississippi drainage; Ohio, Cumberland, and Tennessee river systems; Michigan; Upper St. Lawrence drainage.

† **ALASMIDONTA RAVENELIANA** Lea.

- * *Margaritana raveneliana* LEA, Tr. Am. Phil. Soc., V, 1834, p. 106, pl. XVII, fig. 50; * Obs., I, 1834, p. 218, pl. XVII, fig. 50.—* KUSTER, Conch. Cab. Unio, 1862, p. 297, pl. XCIX, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 173.
 * *Alasmodonta raveneliana* FERUSSAC, Guer. Mag., 1835, p. 26.
 * *Margarita (Margaritana) raveneliana* LEA, Syn., 1836, p. 44; 1838, p. 27.
 * *Margaron (Margaritana) raveneliana* LEA, Syn., 1852, p. 42; 1870, p. 68.
 * *Baphia raveneliana* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 500.
 * *Strophitus ravenelians* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 263.
 * *Unio swananoensis* HANLEY, Test. Moll., 1842, p. 211; Biv. Shells, 1843, p. 211, pl. XXIII, fig. 39.—CATLOW and REEVE, Conch. Nom., 1845, p. 64.

¹ Say's manuscript. I believe this to be specifically different from the smaller, thinner, less inflated, biangulate *A. marginata*. The latter is confined to the Atlantic drainage; the former belongs to the Mississippi Valley, but has migrated into the St. Lawrence basin. No description was published of this form that I know of until that of Mr. Wright appeared.

Tennessee and Cumberland river systems.

Unfigured species.

Margaritana etowahensis CONRAD, Pr. Ac. Nat. Sci. Phila., IV, 1849, p. 154.¹

Etowah River, Georgia.

Subgenus *BULLELLA* Simpson, 1900.

(Type, *Margaritana arcula* Lea.)

Shell thin, greatly inflated, somewhat triangular, with a high, sharp posterior ridge; beaks very full, having exceedingly strong, concentric sculpture, extending well on to the disk; pseudocardinals reflexed, compressed.

Animal with the gills large, nearly semicircular below, inner the larger, united the whole length to the abdominal sac; mantle border with square spots.²

†ALASMODONTA ARCULA Lea.

* *Margarita* (*Margaritana*) *arcula* LEA, Syn., 1836, p. 43; 1838, p. 27.

* *Margaritana arcula* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 71, pl. xxii, fig. 69; * Obs., II, 1838, p. 71, pl. xxii, fig. 69.—* TROSCHEL, Arch. für. Nat., V, 1893, Pt. 2, p. 238.—* CHENU, Man., 1859, II, p. 144, fig. 712.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 172

* *Unio arcula* HANLEY, Test. Moll., 1842, p. 210; Biv. Shells, 1843, p. 210, pl. xxii, fig. 49.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLVIII, fig. 259.

* *Margaron* (*Margaritana*) *arcula* LEA, Syn., 1852, p. 42; 1870, p. 67.

* *Alasmodonta arcula* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 262.

* *Baphia arcula* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 500.

* *Margaritana* (*Alasmodonta*) *arcula* CLESSIN, Conch. Cab. Ano., 1876, p. 272, pl. LXXXIII, figs. 7, 8.

Altamaha River, Georgia.

†ALASMODONTA TRIANGULATA Lea.

* *Margaritana triangulata* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 138; * Jl. Ac. N. Sci. Phila., IV, 1859, p. 228, pl. xxxii, fig. 111; * Obs., VII, 1859, p. 46, pl. xxxii, fig. 111.—* KUSTER, Conch. Cab. Unio, 1862, p. 303, pl. c, fig. 6.—* B. H. WRIGHT, Check List, 1888.

* *Unio triangulata* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXX, fig. 414.

* *Margaron*, (*Margaritana*) *triangulata* LEA, Syn., 1870, p. 68.

* *Margaritana triangulatus* PÆTEL, Conch., Sam., III, 1890, p. 174.

South Carolina and Georgia.

The following species are unknown to me:

* *Alasmodon* (*Decurambis*) *atropurpureum* RAFINESQUE, Cont. Monog., 1831, p. 5.

* *Alasmodon* (*Decurambis*) *scriptum* RAFINESQUE, Cont. Monog., 1831, p. 5.

¹ Allied to *ravenelianus* LEA, according to Conrad.

² In the only gravid specimen seen the outer gills were curiously wrinkled and folded longitudinally, though the inner gills were plain. If this were a normal character it would entitle the species to generic rank, but I can not feel certain that it is. *A. triangulata* seems to stand between this and the ordinary *Alasmodontas*.

- * *Alasmodon (Lasmigona) viridis* RAFINESQUE, Cont. Monog., 1831, p. 5.
 * *Alasmodon (Sulcataria) papyraceum* RAFINESQUE, Cont. Monog., 1831, p. 5.
 * *Alasmodon (Sulcataria) badium* RAFINESQUE, Cont. Monog., 1831, p. 5.
 * *Alasmodon (Amblamodon) hians* RAFINESQUE, Cont. Monog., 1831, p. 5.
 * *Margaritana columbeusis* LEA,¹ Pr. Ac. N. Sci. Phila., XI, 1867, p. 81.

Genus HEMILASTENA (Agassiz, 1852) Simpson.

(Type, *Alasmodonta ambigua* Say.)²

Shell small, elongate elliptical, rounded in front and behind, often slightly incurved at the central base; beaks rather sharp, but not full; sculpture consisting of fine, parallel ridges which are looped up in the middle, and open behind; epidermis brownish, rayless; teeth imperfect, a single irregular, compressed tooth in each valve; that of the left under the beak, that of the right in front of it; laterals nearly or quite wanting; anterior end of the shell much thickened; anterior muscle scars united; posterior faint; naere dull whitish.

Animal with the marsupium occupying the whole of the outer branchiæ; ovules very large; ovisacs not separated by a sulcus; inner gills the larger in front, free from the abdominal sac part of the way, all four united to the mantle to the posterior end; mantle with a double border; branchial opening large, with many crowded papillæ; anal opening smooth; animal whitish.

† HEMILASTENA AMBIGUA Say.

- * *Alasmodonta ambigua* SAY, Jl. Ac. N. Sci. Phila., V, 1825, p. 131.—* FERUSSAC, Guer. Mag., 1835, p. 26.
 * *Margaritana ambigua* KUSTER, Conch. Cab. Unio, 1862, p. 300, pl. XCIX, fig. 7.—* PÆTEL, Conch. Sam., III, 1890, p. 173.
 * † *Unio hildrethianus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 36, pl. III, fig. 8; * Obs., I, 1834, p. 148, pl. III, fig. 8.—* HANLEY, Test. Moll., 1842, p. 196; * Biv. Shells, 1843, p. 196, pl. XXIII, fig. 38.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* KUSTER, Conch. Cab. Unio, 1861, p. 206, pl. LXVIII, fig. 8.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVI, fig. 192.—* BAKER, Moll. Chicago, Pt. 1, 1898, p. 71, pl. v, fig. 4.
 * *Margarita (Unio) hildrethianus* LEA, Syn., 1836, p. 28; 1838, p. 20.
 * *Margaron (Margaritana) hildrethianus* LEA, Syn., 1852, p. 43; 1870, p. 69.
 * *Strophitus hildrethiana* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.
 * *Baphia hildrethiana* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 499.
 * *Margaritana hildrethiana* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 173.
 * *Alasmodonta dubia* FERUSSAC, Guer. Mag., 1835, p. 26.

Ohio River system; north to Michigan; west to Iowa; south to Arkansas; east to Tennessee.

¹Dr. Lea does not mention this in his Synopsis.

²This species, the only member of the group, differs from *Margaritana* in the beak sculpture, the thickening of the anterior end of the shell, in the want of muscle scars in the shell cavities, in the color of the animal, and in having the gills united to the mantle posteriorly to their ends, and the shell is not that of a *Unio*, as it differs in the beak sculpture and the teeth. I regret that Lea's well-known name must be displaced for the obscure one of Say, who undoubtedly had this shell before him when he described his *Alasmodonta ambigua*.

Genus MARGARITANA Schumacher, 1817.

(Type, *Mya margaritifera* Linnæus.)*Baphia* MEUSCHEN, Mus. Gevers, 1787, p. 472.*Unio* RETZIUS, part, Diss. Hist. Nov. Test. Gen., 1788, p. 16.*Unio* OKEN, Lehrbuch der Nat., 1815, p. 236.*Margaritana* SCHUMACHER, Essai Nouv. Syst., 1817, p. 137.*Damalis* (LEACH manuscript) GRAY, Pr. Zool. Soc. Lond., 1847, p. 196.*Baphia* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 499.¹

Shell elongated, usually arcuate, rounded in front, almost lacking a posterior ridge; beaks rather low, the sculpture consisting of a few coarse, parallel ridges, which follow the growth lines; epidermis concentrically striate, brownish or blackish; hinge teeth generally imperfect or not fully developed; two more or less perfect pseudocardinals in the left valve, and one in the right, often reduced to mere tubercles; behind these the hinge plate is narrow and rounded for some distance; laterals short, usually imperfect or wholly wanting; cavity of the beaks rather shallow; muscle scars large, those of the anterior roughened, posterior elliptical; nacre generally more or less covered inside the pal-leal line with small muscle scars.

Animal with very long gills, inner wider in front, free for the greater part of their length from the abdominal sac, the two pairs united to their posterior ends, which project backward for some distance unconnected with the mantle; palpi very large, falcate, united half way posteriorly; branchial opening having crowded, often arborescent papillæ arranged in folds; anal opening smooth or crenulate; superanal opening not closed below; marsupium occupying the entire outer gills.² Foot and abdomen small; whole animal dark colored.

† MARGARITANA MARGARITIFERA Linnæus.

Mya margaritifera LISTER, Hist. Anim. Aug. App., 1685, pl. I, fig. 1.—Hist. Conch., 1685, pl. CXLIX, fig. 4.—* LINNÆUS, Systema. Nat., 10th ed., I, 1758, p. 671.—* DA COSTA, Hist. Nat. Brit., 1778, p. 225, pl. xv, fig. 33.—* GMELIN, Syst. Nat., 13th ed., 1792, p. 3219.³—* LINNÆUS, Fauna Suecica, 1761, p. 2130.—KNORR, Vergn., IV, 1769, pl. XXV, fig. 2.—* MÜLLER, Vermes, 1774, p. 210.—PENNANT, Brit. Zool., IV, 1777, pl. XLIII, fig. 18.—* DA COSTA, Hist. Nat. Brit., 1778, p. 225, pl. xv, fig. 3.—* SCHRÖTER, Fluss Conch., 1779, p. 168, pl. IV, fig. 1.—* BORN, Test. Mus. Vind., 1780, p. 21.—* SCHRÖTER, Ein. Conch., 1783, II, p. 606.—* DONOVAN, Brit. Shells, III, 1801, pl. LXXIII.—* MONTAGU, Test. Brit., 1803, p. 33.—* TURTON, Brit. Fauna, 1807, p. 146.—* MATON and

¹ The name *Baphia* Meuschen has precedence, but was never described; it was applied to a miscellaneous lot of bivalves, and is in the plural number. Retzius' first species in *Unio*, the type of a section without laterals, is the *U. margaritifera*, but in 1792 Bruguière in Choix de Mémoires, I, p. 106, fully and carefully redefined the genus *Unio*, restricting it to species with cardinal and lateral teeth.

² According to Van Wahl. I have examined a great many animals taken at different seasons, but have never seen one gravid.

³ Refers to Conchylien Cabinet VI, pl. 1, fig. 5, which is the form we know as *Margaritana margaritifera*.

- RACKETT, Tr. Linn. Soc. Lond., VIII, 1807, p. 40.—* WOOD, Gen. Conch., I, 1815, pl. XXIII, figs. 1-3.—* DILLWYN, Cat., I, 1817, p. 52.—* TURTON, Conch. Dict., 1819, p. 106.—* BINGLEY, Useful Knowledge, III, 1825, p. 245.—* WOOD, Index Test., 1825, p. 12, pl. II, fig. 30.—* EATON, Zool. Text-Book, 1826, p. 216.—* WAARDENBERG, Com. Hist. Nat. Anim., 1827, p. 360.—* CHENU, Bib. Conch., 1st ser., I, 1845, p. 57, pl. XX, figs. 1, 2.—* HANLEY, Ipsa. Linn. Conch., 1855, p. 460.—* HANLEY (WOOD), Ind. Test., 3d ed., 1856, p. 16, pl. II, fig. 30.
- Mya margaritifera* testa ovali oblonga, etc.—* MÖLLER, Zool. Danicæ, 1776, p. 245.—* CHEMNITZ, Conch. Cab., VI, 1782, p. 15, pl. I, fig. 5.
- * *Baphia margaritifera* MEUSCHEN, Mns. Gevers, 1787, p. 472.
- * *Unio margaritifera* DRAPARNAUD, Tab. Moll. Fr., 1801, p. 107; Hist. Moll. Fr., 1806, p. 132, pl. X, figs. 17-19; pl. XI, fig. 5.¹—* GAERTNER, Vers. Syst., 1813, p. 37.—* OKEN, Lehrbuch, 1815, p. 238.—* CUVIER, Règne Animal, II, 1817, p. 473.—* C. PFEIFFER, Nat. Deutsch. Land und Suss. Moll., 1821, Pt. 1, p. 115, pl. V, fig. 11.—* BOSCH, Hist. Nat. Coq., III, 1824, p. 139.—* GRAS, Moll. Isère. Ap., 1840, p. 22.—* FRIELE, Norske L. Fersk Moll., 1853, p. 53.
- * *Margaritana margaritifera* SCHUMACHER, Essai. Nouv. Syst., I, 1817, p. 124, pl. X, fig. 4.—* MICHAUD, Comp. Hist. Moll. Fr., 1831, p. 114.—* SCHOLZ, Schleis L. and W. Moll., 1843, p. 135.—* TROSCHEL, Arch. für Naturg., XIII, Pt. 1, 1847, p. 270; VI, fig. 1.—* MOQUIN-TANDON, Hist. Moll. Fr., 1855, p. 623, pl. XXII, figs. 14-16.—* VON WAHL, Arch. für die Nat. Kunde Liv., 2d ser., I, 1855, p. 118.—* KUSTER, Conch. Cab., 1856, p. 293, pl. XXXVIII; XXXIX, figs. 2-4.—* CHENU, Manual, 1859, II, p. 144, fig. 710.—* WESTERLUND, Faun. Sw. Nor. and Den., 1873, p. 577.—* HARTMAN and MICHENER, Conch. Cest., 1874, p. 91, fig. 189.—* CLESSIN, Deutsch. Ex. Moll., 1876, p. 449, fig. 293.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 173.—* STEARNS, Pr. U. S. Nat. Mus., XIV, 1891, p. 105.—* LOCARD, Coq. de France, 1893, p. 149.—* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 155.—* WESTERLUND, Act. Soc. F. and F. F., XIII, No. 7, 1897, p. 166.
- * *Alasmodonta margaritifera* CONRAD, New F. W. Shells, 1834, p. 72.—FERUSSAC, Guer. Mag., 1835, p. 26.
- * *Margarita (Margaritana) margaritifera* LEA, Syn., 1836, p. 45; 1838, p. 28.
- * *Margaron (Margaritana) margaritifera* LEA, Syn., 1852, p. 43; 1870, p. 69.
- * *Baphia margaritifera* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 499, III, pl. CXVII, figs. 2, 2a.
- * *Unio margaritiferus* RETZIUS, Dis. S. Hist. Nat., 1788, p. 16.—* SPENGLER, Skriv. Nat. Selsk., III, 1793, p. 52.—* TURTON, Conch. Ins. Brit., 1822, p. 241, pl. XVI, fig. 1.—* NILSSON, Hist. Moll. Svec., 1822, p. 103.—* TURTON, Man. L. and F. W. Shells, 1831, p. 19, fig. 9.—* ? ANTON, Verz. der Conch., 1839, p. 15.—* HANLEY, Test. Moll., 1842, p. 213; * Biv. Shells, 1843, p. 213.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* FORBES and HANLEY, Hist. Brit. Moll., II, 1853, p. 146, pl. XXXVIII.—* SOWERBY, Ill. Ind. Brit. Shells, 1859, pl. VII, No. 1.—* TATE, L. and F. W. Moll. Brit., 1866, pl. III, fig. 15.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIV, fig. 325.
- * *Alasmodon margaritiferus* BROWN, L. and F. W. Conch., 1836, p. 112, pl. XXI, fig. 13, pl. XXII, figs. 1-3.—* THOMPSON, Ann. and Mag. Nat. Hist., VI, 1840, p. 197.—* BROWN, Ill. Recent Conch., 1844, p. 83, pls. XXX, figs. 1-4; XXXI, figs. 1, 2; XXXII, figs. 13-15.—* TURTON, Man. L. and F. W. Shells, 1857, p. 277, pl. II, fig. 9.
- * *Margaritana margaritiferus* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 184.
- * *Alasmodon margaritiferum* FLEMING, Hist. Brit. Moll., 1828, p. 417.
- * *Unio margaritifer* var. *minor* ROSSMASSLER, Icon., Pt. 2, 1835, p. 19, pl. IX, fig. 129.

¹The figures on pl. X are of Linnaeus's species; that on pl. XI may be *Unio batavus* Lamarec.

- * *Unio margaritifera* KUSTER, Conch. Cab., 1856, p. 130, pls. XXXVIII, XXXIX.—
 * MOQUIN-TANDON, Moll. Terr. et Fluv. Fr., II, 1855, p. 566, pl. XLVII.—* NORDENSKIÖLD and NYLANDER, Fin. Moll., 1856, p. 86, pl. VI, fig. 74.—* DROUET, Nay. Fr., II, 1857, p. 57, pl. I.—* HESSLING, Perl. und Ihre Perlen, 1859, p. 86, pl. I.—* JEFFREYS, Brit. Conch., I, 1862, p. 37.—* REEVE, L. and F. W. Moll. Brit., 1863, p. 223, fig. 3.—* L. ADAMS, Coll. Man., 1884, p. 19, pl. I, fig. 12.
- * *Unio (Margaritana) margaritifera* SCHRENCK, Reis. und F. im Amur-Lande, II, 1867? p. 700.
- * *Unio auricularis* SPENGLER, Skriv. Nat. Selsk., III, 1793, p. 44.
- * *Unio elongata* LAMARCK, Ann. sans Vert., VI, 1819, p. 70.—* STARK, Nat. Hist., II, 1828, p. 90.—* GRAS, Moll. Isère Ap., 1840, p. 22.—* PUTON, Moll. Vosges., 1847, p. 72.
- * *Damaris elongata* LEACH, Syn. Moll. Gt. Brit., 1852, p. 322.
- * *Margaritana elongata* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 185.—* LOCARD, Coq. de Fr., 1893, p. 149, fig. 162.
- * *Unio elongatus* NILSSON, Hist. Moll. Svec., 1822, p. 106.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXVI, fig. 397.—* PÆTEL, Conch. Sam., III, 1890, p. 151.
- * *Alasmodonta arcuata* BARNES, Am. Jl. Sci., VI, 1823, p. 277, pl. XII, fig. 20.—* C. B. ADAMS, Thompson's Hist. of Verniont, 1842, p. 165, figured; * F. W. and L. S. of Vt., 1842, p. 165, figured.—* HESSLING, Die. Perl. und Ihre Perlen, 1859, p. 205.
- * *Mya arcuata* EATON, Zool. Text Book, 1826, p. 222.
- * *Alasmodon arcuata* GOULD, Inv. of Mass., 1841, p. 113, fig. 75.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 197, pl. XIV, fig. 224.—* MIGHELS, Bost. Jl. N. Hist., 1844, p. 325.
- * *Margaritana arcuata* STIMPSON, Shells of N. Eng., 1851, p. 15.—* KUSTER, Conch. Cab., 1856, p. 293, pl. XXXIX, fig. 1.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 262.—* GOULD, Inv. of Mass., 1870, p. 174, fig. 477.—* PÆTEL, Conch. Sam., III, 1890, p. 172.—* H. CARPENTER, Naut., IV, 1890, p. 35.
- * *Unio sinuata* C. PFEIFFER, Nat. Deutsch. L. und S. W. Moll., Pt. 2, 1825, p. 33, pl. VII, fig. 4.
- * *Unio roissyi* MICHAUD, Comp. Hist. Moll., Fr., 1831, p. 112, pl. XVI, fig. 28.—* GRAS, Moll. Isère Ap., 1840, p. 22.
- * *Margaritana roissyi* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 186.—* LOCARD, Coq. de France, 1893, p. 150.
- * *Unio tristis* MORELET, Moll. Portugal, 1845, p. 107, pl. XIII, fig. 2.²
- * *Margarita (Unio) crassissimus* LEA, Syn., 1836, p. 40; 1838, p. 26.³
- * *Unio (Alasmodonta) dahuricus* MIDDENDORFF, Bull. Phys. Math. Ac. St. Petersburg, IX, 1850, p. ?; * Sib. Reise, II, 1851, p. 275, pl. XXVI, figs. 3-5.
- * *Unio dahuricus* MIDDENDORFF, L. and S. Moll. Sib., 1859, p. 26.—* HESSLING, Perl. und Ihre Perlen, 1859, p. 202.—* WESTERLUND, Kong. Sv. Vet. Ak. Handl., XIV, No. 12, p. 74.
- * *Unio (Margaritana) dahuricus* SCHRENCK, Reis. und Forsh. Amur-Lande, II, 1867, p. 699.
- * *Margaritana dahurica* KOBELT, Abh. Senck. Nat. Ges., XI, 1875, p. 427.—* KOBELT, Faun. Jap. Ext., 1879, p. 143, pl. XIII, figs. 1, 2.—* MIDDENDORFF, Kong. Svensk. Vetensk. Ak. Handl., XIV, No. 12, p. 110.—* PÆTEL, Conch. Sam., III, 1890, p. 173.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 187.

¹ Lamarck thinks this may be the *Mya margaritifera* of Linnæus; Lea is certain that it is.

² The figure looks like an elongated *batarnis*, but Morelet afterwards acknowledged that his shell was probably a young *margaritifera*.

³ The naked name *crassissima* was first applied to this by Klein, Methodi Ostracologiae, 1753, p. 128, pl. x.

*† *Alasmodon falcata* GOULD, Pr. Boston Soc. Nat. Hist., III, 1850, p. 294; * *Otia* Conch., 1862, p. 87;—* U. S. Expl. Exp., XII, 1852, p. 433, figs. 545, 545a, 545b.—CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 262.—PÆTEL, Conch. Sam., III, 1890, p. 173.

* *Unio falcatus* SOWERBY, Conch. Ieon., XVI, 1868, pl. LXXV, fig. 390.

* *Unio (Alasmodonta) complanatus* MIDDENDORFF, Sib. Reise, II, 1851, Pt. 1, p. 273, pl. XXVII, figs. 1-6.³

* *Margaritana complanata* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 187.

* *Unio mongolicus* MIDDENDORFF, Sib. Reise, II, 1851, p. 277, pl. XXVII, figs. 7, 8.—* HESSLING, Perl. und Ihre Perlen, 1859, p. 203.—* SCHRENCK, Reise und F. im Amur-Lande, II, 1867, p. 699.—* WESTERLUND, Kong. Sv. Vet. Ak. Hand, XIV, no. 12, 1876, p. 74.—* PÆTEL, Conch. Sam., III, 1890, p. 159.—* WESTERLUND, Faun. der Pal., II, Pt. 7, 1890, p. 113.

* *Alasmodon yubaensis* TRASK, Pr. Cal. Acad. Sci., I, 1855, p. 30.

*† *Margaritana rareneliana* CHENU., Man., 1859, II, p. 144, fig. 714.³

All Europe except the southernmost portion; northern Asia; Japan; northern North America; Iceland. Its southern limit seems to be about north latitude 40°. It appears to be entirely circumboreal, except that, so far as is known, it is missing in the central part of the North American continent. Wetherby has suggested that it may have been destroyed in this region by the ice of the glacial epoch. It is found in the upper Missouri and in Canada East, New York, and the New England States.

† MARGARITANA CRASSA Retzius.

* *Mya testa crassa*, SCHRÖTER, Flusseeconch., 1779, p. 182, pl. II, fig. 2.

* *Unio crassus* RETZIUS, Diss. Hist. Nat., 1778, p. 17.⁴—SPENGLER, Skriv. Nat. Selsk., III, 1793, p. 56.—* SCHOLZ, Schleis., I. und W. Moll., 1843, p. 130.—* VON WAHL, Arch. Naturkunde Liv., 2d. ser., I, 1855, p. 94.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* DROUET, Moll. Cote d'Or., 1867, p. 101.—* JORDAN, Jahrb. Deutsch. Mal. Ges., VI, 1879, p. 306.—* DROUET, Mem. Acad. Dijon, 3d ser., VII, 1882, p. 13.⁵—* PÆTEL, Conch. Sam., III, 1890, p. 149.

¹ This is undoubtedly *margaritifera*, but in the types the young shells show laterals.

² I have no doubt that this is *Margaritana margaritifera* L. and not *Unio complanatus* Solander as Middendorff supposes. The specimens figured are terribly eroded, and the hinges are in very bad order, but they do not show laterals.

³ Numerous varieties as well as species have been made of this abundant, widespread form. Notwithstanding its enormous distribution, the widest of any Naiad known, I have seen no variation that seems to me to be worthy of a varietal name.

⁴ Retzius' description is in Latin, and consists of ten words, and as he does not figure the species, it would be impossible to determine what he meant only for the fact that he refers to the Flusseeconchylien of Schröter (*Mya testa crassa*, p. 182, pl. II, fig. 2). Schröter's figure is not a very good one, but I can have no doubt that he had before him the large, heavy, black Naiad, with lateral teeth, found in southern Europe. In general it closely resembles the *M. margaritifera* externally, but the epidermis is a more dull black, the beaks are usually higher and more strongly developed, the naere is white, and it has fairly well-developed laterals. European conchologists seem to have quite commonly mistaken heavy, large forms of *Unio batavicus* for Retzius' species, and in a number of cases the *M. margaritifera* has been taken for it.

⁵ Page 13 of reprint; I have not seen the original.

* *Margaron (Unio) crassus* LEA, Syn., 1852, p. 39; 1870, p. 62.¹

* *Unio auricularius* SPENGLER, Skriv. Nat. Selsk., III, 1792, p. 54.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 50.

Unio rugosa POIRET, Coq. Fluv., et Terr. de l'Aisne, 1801, p. 105.²

* *Unio sinuata* LAMARCK, An. sans Vert., VI, 1819, p. 70.³—BLAINVILLE, Man., 1825, p. 539, pl. LXVII, fig. 3.—* DESHAYES, Enc. Méth., II, 1827, p. 151, pl. CCXLVIII, fig. 1; II, 1830, p. 579, pl. CCXLVIII, fig. 1, 1a, 1b.⁴—* SGANZIN, Mem. Hist. Nat. Strasb., III, 1842, II, p. 8.⁵—* PUTON, Moll. Vosges, 1847, p. 72.

* *Unio (Potamida) sinuata* SWAINSON, Tr. on Mal., 1840, p. 268.

* *Unio sinuatus* ROSSMASSLER, Icon., III, 1836, p. 22, pl. XIII, fig. 195.—* GASSIES, Moll. de l'Ag., 1849, p. 198.—* DUPUY, Hist. Moll. Fr., 1852, p. 630, pl. XXIII, fig. 7.—* ROSSMASSLER, Icon., III, 1854, p. 38, pl. LXX, fig. 853, 853a.⁶—* MOQUIN-TANDON, Moll. Terr. and Fluv. Fr., II, 1855, p. 567; III, pl. XLVIII, figs. 1-3.—* DROUET, Nay. Fr., II, 1857, p. 61, pl. II.—* HESSLING, Die Perl. and Ihre Perlen, 1859, p. 184.—* SOWERBY, Conch. Icon., XVI, pl. LXII, fig. 311.—* PÆTEL, Conch. Sam., III, 1890, p. 167.—* LOCARD, Coq. de Fr., 1893, p. 151, fig. 164.

*? *Unio litoralis* C. PFEIFFER, Nat. Deutsch. L. und S. W. Moll., Pt. 1, 1821, p. 117, pl. v, fig. 12.⁷

* *Unio crassissimus* HANLEY, Test. Moll., 1842, p. 209; * Biv. Shells, 1843, p. 209, pl. XXIII, fig. 54.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.

* *Unio gargotta* PHILIPPI, Moll. Sic., 1836, p. 66.—* ROSSMASSLER, Icon., VII and VIII, 1838, p. 26, pl. XXXV, fig. 493.⁸—* DROUET, Jl. de Conch., XXIX., 1881, p. 25.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 156.

Southern Europe; possibly into Asia Minor and southwest Siberia.

† MARGARITANA LAOSENSIS Lea.

* *Unio laosensis* LEA, Pr. Ac. Nat. Sci. Phila., VII, 1863, p. 190; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 63, pl. XXI, fig. 61; * Obs., XI, 1867, p. 67, pl. XXI, fig. 61.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLVII, fig. 256.—* PÆTEL, Conch. Sam., III, 1890, p. 156.

* *Margaron (Unio) laosensis* LEA, Syn., 1870, p. 62.

Unio sula THEOBALD (where?).

Laos Mountains, Cambodia; Siam.

¹ It has often been impossible to determine what species has been referred to by authors under the name *crassus*. In cases where they have cited this species from the north of Europe I have no doubt that they either had *Margaritana margaritifera* or *Unio batavus* before them. In those cases where I could not be certain as to what was meant I have omitted references.

² According to Moquin Tandon this is *U. sinuatus* Lamarck.

³ Lamarck refers to Encyc. Méth., 1797, pl. CCXLVIII, figs. 1a, 1b, which seem to be the *crassus* of Retzius. Lea, who examined Lamarck's type, says he had thought it was the *Margaritana margaritifera*, but that it had lateral teeth. (Obs. I, pp. 35, 198.)

⁴ These figures look something like a heavy inflated *Lampsilis alatus* Say, but are probably the *crassus* of Retzius.

⁵ Given to Sganzin by M. Fabert, chief of battalion of infantry, as having been found at St. Paul in the Isle of Bourbon, but Sganzin very much doubts the locality. Of course it was never found in the Indian Ocean region.

⁶ A magnificent figure of this fine species, and in a, which shows the hinge, the laterals are plainly delineated.

⁷ Dr. Lea is doubtful about this being *litoralis*, and I am sure it is not. It appears more like *M. crassus* than anything else.

⁸ Dr. Lea makes this a synonym of Pfeiffer's *elongatulus*, which is, I believe, quite near to *U. pictorum*. Rossmassler's figure shows a shell very much like *Margaritana margaritifera*, but it has lateral teeth.

† MARGARITANA HEMBELI Conrad.

- * *Unio hembeli* CONRAD, Monog., X, 1838, p. 93, pl. LI, fig. 1.—* HANLEY, Biv. Shells, 1856, p. 383, pl. XXIV, fig. 3.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIII, fig. 172.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 154.
- * *Margarou (Unio) hembeli* LEA, Syn., 1852, p. 21; 1870, p. 32.

Louisiana; Burnt Corn, Alabama.

† MARGARITANA MONODONTA Say.

- Unio monodonta* SAY, N. Harm. Diss., II, 1829, p. 293; * Am. Conch., I, 1830, pl. VI.—SHORT and EATON, Trans. Jl., 1831, p. 79.—* CONRAD, New F. W. Shells, 1834, p. 70.—DESHAYES, An. sans. Vert., 2d ed., VI, 1835, p. 553; 3d ed., II, 1839, p. 674.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 12, pl. II, fig. 1.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.
- * *Alasmodonta monodonta* FERUSSAC, Guer. Mag., 1835, p. 26.
- * *Margaritana monodonta* CONRAD, Pr. Ac. N. Sci., Phila. VI, 1853, p. 262.
- * *Unio monodontus* SAY, Am. Conch., VI, 1834.—* HANLEY, Test. Moll., 1842, p. 210; * Biv. Shells, 1843, p. 210, pl. XXIII, fig. 48.—* H. and A. ADAMS, Gen., Rec. Moll., II, 1857, p. 491.—* KUSTER, Conch. Cab., 1861, p. 221, pl. LXXV, fig. 1.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 159.
- * *Margarita (Unio) monodontus* LEA, Syn., 1836, p. 40; 1838, p. 26.
- * *Margarou (Unio) monodontus* LEA, Syn., 1852, p. 39; 1870, p. 62.
- * *Unio soleuiformis* LEA, Tr. Am. Phil. Soc., LV, 1831, p. 87, pl. x, fig. 17; * Obs., I, 1834, p. 97, pl. x, fig. 17.—* CHENU, Ill. Conch., 1858, pl. XIV, figs. 4, 4a, 4b; * Manual, II, 1859, p. 137, fig. 661.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLV, fig. 243.
- * *Margaritana soleniformis* PÆTEL, Conch. Sam., III, 1890, p. 173.

Ohio; Cumberland, and Tennessee river systems; Illinois; eastern Iowa; Nebraska?

† MARGARITANA DECUMBENS Lea.

- * *Unio decumbens* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 40.—* LEA, Jl. Ac. N. Sci. Phila., 1862, p. 87, pl. XII, fig. 236.—* LEA, Obs., VIII, 1862, p. 91, pl. XII, fig. 236.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXII, fig. 432.—* B. H. WRIGHT, Check List, 1888.
- * *Margarou (Unio) decumbens* LEA, Syn., 1870, p. 62.

Alabama.

Genus UNIO Retzius, 1788.¹

(Type, *Unio tumidus* Retzius.)

- Unio* RETZIUS, Diss. Hist. Nov. Test. Gen., 1788, p. 16.—BRUGUIERE, Choix de Memoirs, I, 1792, p. 106.
- Limnæa* POLI (part), Test. Utr. Sic., I, 1791, p. 31.
- Lymnium* OKEN, Lehrbuch, 1815, p. 237.
- Elliptio* RAFINESQUE, Jl. de Phys. Nat. Hist., 1819, p. 426.
- Mysca* TURTON, Conch. Ins. Brit., 1822, p. 243.

¹This genus was described in a thesis by Laurentius Münter Philipsson under his master, Retzius, in the University of Lund, Sweden, and it is often credited to the former. I am informed by Professor Joh. Chr. Moberg, of Lund, that by a former law or custom of the university the professor was considered the author of all papers which a student under him defended. According to this, Retzius must be credited with the genus. This law was repealed in Lund in 1852.

Canthytia SWAINSON, Tr. on Mal., 1840, p. 278.

Uniomercus CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 268.

Shell inequilateral, oval to elongated, rounded in front and pointed or biangulate behind, with a more or less developed posterior ridge, often becoming slightly arcuate when old; beaks only moderately full, generally sculptured with coarse ridges, which run parallel with the growth lines, or are somewhat doubly looped, sometimes broken and showing fine radiating lines behind; surface smooth, slightly concentrically ridged or pustulous; epidermis generally rather dull colored, rayless or feebly rayed; hinge plate narrow; two pseudocardinals and two laterals in the left valve and one pseudocardinal and one lateral in the right, with rarely a vestige of a second lateral; cavity of the beaks not deep or compressed. Animal having the inner branchiæ free from the abdominal sac for from one-half to their entire length; marsupium occupying the whole length of the outer gills only, forming a thick, smooth pad when filled with young; gills united to the mantle behind to their extreme points, or very nearly so; papillæ on branchial and anal openings unbranched; superanal opening always closed below.

Section LYMNIIUM Oken, 1815.

(Type, *Unio pictorum* Retzius.)

Shell generally smooth; beak sculpture broken, often somewhat corrugated or pustulous; pseudocardinals compressed; beak cavities well excavated, not compressed. Animal highly colored, anal opening crenulate or smooth.

(Group of *Unio pictorum*.)

Shell inflated, elongate, oval, anterior end angled above, swollen a little at posterior base; beaks full, their sculpture consisting of numerous slightly doubly looped bars which often become pustulous; posterior ridge rather low; epidermis smooth, rather bright, sometimes slightly rayed behind; rest periods well marked; pseudocardinals compressed, often a little reflexed, smooth below, those of the left valve partly united; muscle scars smooth; naere whitish to salmon. Animal the same as described for the section.

† UNIO PICTORUM Linnæus.¹

* *Mya pictorum* LINNÆUS, Syst. Naturæ, 10th ed., 1758, I, p. 671; Faun. Suec., 1761, No. 2129.—* MÜLLER, Vermes, 1774, p. 211.—PENNANT, Brit. Zool., IV, 1777? pl. XLIII, fig. 17.—* DA COSTA, Hist. Nat. Brit., 1778, p. 228, pl. xv, fig. 4.—* SCHRÖTER, Flusconch., 1779, p. 178, pl. III, figs. 2, 4, 5.—* BORN, Test. Mus. Vind., 1780, p. 20.—* SCHRÖTER, Ein. Conch., 1783, II, p. 604.—* OLIVI, Zool. Adriatic, 1792, p. 95.—STURM, Deuts. Faun., VI, 1803, 2d ed., p. 19, pls. a, b, c.—* MONTAGU, Test. Brit., 1803, p. 36.—* TURTON, Brit. Fauna,

¹ Hanley says [Ipse Linnæi Conchyliæ, p. 27]: "More Uniones than one are present in the [Linnean] collection, but upon the whole the *U. pictorum* of authors [Rossm. Icon., fig. 196] agrees best with synonymy and description. The figure referred to of Lister is *U. pictorum*; Bonanni's drawing is more doubtful and was possibly meant for *U. tumidus*. The descriptions in Fauna Suecica and Systema are brief and unsatisfactory and might suit either species alike."

1807, p. 116.—*MATON and RACKETT, Tr. Linn. Soc. Lond., VIII, 1807, p. 38.—*WOOD, Gen. Conch., I, 1815, p. 104, pl. XIX, figs. 3, 4.—*DILLWYN, Cat., I, 1817, p. 49.—*TURTON, Conch. Dict., 1819, p. 106.—*WOOD, Index Test., 1825, p. 12, pl. II, fig. 26c; *rev. ed., 1856, p. 15, pl. II, fig. 26.—*CHENU, Bib. Conch., 1st ser., I, 1845, p. 114, pl. XLVII, figs. 8, 9.¹—*HANLEY, Ipsa. Linn. Conch., 1855, p. 460.

**Unio pictorum* RETZIUS, Diss. Hist. Nat., 1778, p. 17.—*SPENGLER, Skriv. Selsk. Nat., III, 1793, p. 59.—*DRAPARNAUD, Tab. Moll. Fr., 1801, p. 106; in part Hist. Moll. Fr., 1806, p. 131, pl. XI, fig. 4.²—*GAERTNER, Vers. Ein. Syst., 1813, p. 36.—*MILLET, Moll. Maine et Loire, 1813, p. 74.—*BRARD, Hist. Coq. Paris, 1815, p. 226, pl. VIII, fig. 1.—BROOKES, Int. to Conch., 1815, p. 51, pl. II, fig. 12.—*CUVIER, Règne Animal, II, 1817, p. 473.—*KLEES, Diss. Test., 1818, p. 45.—*LAMARCK, An. sans Vert., VI, 1819, p. 77.—*C. PFEIFFER, L. and Suss. Moll., Pt. 1, 1821, p. 115, pl. V, figs. 9, 20.—*NILSSON, Hist. Moll. Svee., 1822, p. 111.—*BOSC, Hist. Nat. Coq., III, 1824, p. 139, pl. XXIII, fig. 3.—*BLAINVILLE, Manual, Mal., 1825, p. 539, pl. LXVII, fig. 2.—*CROUCH, Ill. Int. Lamarck, 1827, p. 16, pl. IX, figs. 4, 4a, 4b.—*WAARDENBERG, Com. Hist. Nat., 1827, p. 36.—*KLEEBERG, Moll. Boruss, 1828, p. 38.—STARK, Nat. Hist., II, 1828, p. 90.—*FLEMING, Hist. Brit. Animals, 1828, p. 416.—*GOUPIL, Hist. Moll. Sarthe, 1835, p. 84.—*ROSSMASSLER, Icon., I, 1835, p. 118, pl. III, figs. 71, 71a, 71b; III, 1836, p. 23, pl. XIII, fig. 196; VI, 1837, p. 55, pl. XXIX, fig. 409; IX, 1839, p. 10, pl. XLV, fig. 587-590; XI, 1842, p. 14, pl. LV, fig. 741; XII, 1844, p. 30, pl. LVIII, figs. 762-766; p. 31, pl. LIX, figs. 767-769.—*FOURNEL, Faun. Moselle, I, 1836, p. 486.—*FLEMING, Moll. Animals, 1837, pl. XIV, fig. 51.—*WYATT, Man. Conch., 1838, p. 67, pl. VIII, fig. 6.—*PORRO, Mal. Como., 1838, p. 117.—*TERVER, Moll. Terr. et Fluv., 1839, p. 39.—*ANTON, Verz. der Conch., 1839, p. 15.—*? GRAS, Moll. Isere., 1840, p. 71, pl. I, fig. 8.³—*HANLEY, Test. Moll., 1842, p. 205; *Biv. Shells, 1843, p. 205.—*SCHOLZ, Schleis, L. and W. Moll., 1843, p. 127.—*CATLOW and REEVE, Conch. Nom., 1845, p. 62.—*STABILE, Faun. Lug., 1845, p. 60, pl. III, fig. 73.—*MORELET, Moll. Port., 1845, p. 108.—*BROWN, L. and F. W. Conch., 1845, p. 107, pl. XIX, figs. 1-4.—*PÜTON, Moll. Vosges, 1847, p. 70.—*STEIN, Die Lebend. Schneek., 1850, p. 104, pls. XXIV, XXV, figs. 1, 2.—*MIDDENDORFF, Sib. Reise, II, 1851, p. 276, pl. XXVIII, figs. 1-3.—*LEACH, Syn. Moll. Gt. Brit., 1852, p. 234.—*DUPUY, Hist. M. Fr., 1852, p. 617, pl. XXVI, fig. 20.—*FORBES and HANLEY, Hist. Brit. Moll., II, 1853, p. 142, pl. XXXIX, fig. 1.—*KUSTER, Conch. Cab. Unio, 1854, p. 88, pl. XXIII, figs. 1, 2; pl. XXIV; XXV, figs. 1, 2.—*MOQUIN-TANDON, Moll. Terr. Fluv. Fr., II, 1855, p. 576, pl. I, figs. 8, 10; II, figs. 1, 10.—*VON WAHL, Arch. Naturk. Liv. Est., 2d ser., 1855, p. 105.—*NORDENSKIÖLD and NYLANDER, Fin. Moll., 1856, p. 83, pl. V, fig. 72.—*DROUET, Nay. Fr., II, 1857, p. 103, pl. VIII.—*TURTON, Mau. L. and F. W. Shells, 1857, p. 279, pl. II, fig. 11.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491; III, pl. CXVI, figs. 5, 5a, 5b.—*SOWERBY, Ill. Index Brit. Shells, 1859, No. 2, pl. VII.—*GOODRICH, Ill. Nat. Hist., II, 1859, p. 523, fig.—JEFFREYS, Brit. Conch., I, 1862, p. 34.—*BIELZ, Faun. Sieben, 1863, p. 192.—*REEVE, L. and F. W. Moll. Brit., 1863, p. 221, fig. 2.—*TATE, L. and F. W. Moll. G. Brit., 1866, pl. II, fig. 3.—*? SCHRENCK, Reise und F. im Am. Lande, II, 1867, p. 696.⁴—*KOBELT, Faun. Nass. Moll., 1871, p. 241.—*WESTER-

¹ *U. batavus* perhaps.

² Draparnaud gives figures 1-4 for *pictorum*. One and 2 may possibly be that species, but they look more like *batavus*, and 3 is certainly *batavus*, while 4 is probably *pictorum*.

³ Doubtful, may be *batavus*.

⁴ It is very probable that the specimens from Amur Land and vicinity which have been referred to *U. pictorum* are a smooth variety of *Nodularia douglasii* Gray, a species resembling it in form, but not at all closely related to it.

- LUND, Faun. S. N. and D., 1873, p. 574.—* LEHMAN, Die Schnecken, 1873, p. 290.—* CLESSIN, Deutsche Ex. Moll., 1876, p. 453, fig. 294.—* JORDAN, Jahrb. Deuts. Mal. Ges., VI, 1879, p. 301.—* DROUET, Mem. Acad. Dijon, VII, 1882, p. 10 (reprint).—* L. ADAMS, Coll. Man., 1884, p. 18, pl. 1, fig. 11.—* ? CLESSIN, Moll. Oest., 1887, p. 723.—PÆTEL, Conch. Sam., III, 1890, p. 163.
- * *Baphia pictorum* MEUSCHEN, Mus. Gevers., 1787, p. 472.
- * *Lymnium pictorum* OKEN, Lehrb., 1815, p. 237.
- * *Mysca pictorum* TURTON, Conch. Ins. Brit., 1822, p. 245; * Man. L. and F. W. Shells, Brit. Is., 1831, p. 20, fig. 11.
- * *Margarita (Unio) pictorum* LEA, Syn., 1836, p. 36; 1838, p. 24.
- * *Margaron (Unio) pictorum* LEA, Syn., 1852, p. 36; 1870, p. 58.
- * *Mya angusta subflava*, etc., SCHRÖTER, Fluss. Conch., 1779, p. 184, pl. III, fig. 3; pl. IV, fig. 6.
- * *Mya nodosa* GMELIN, Syst. Nat., 13th ed., 1788, p. 3222.¹—* WOOD, Gen. Conch., I, 1815, p. 110.—* DILLWYN, Cat., I, 1817, p. 54.—* WOOD, Ind. Test., 1825, p. 12, pl. II, fig. 34a; * rev. ed., 1856, p. 16, pl. II, fig. 34.—* PÆTEL, Conch. Sam., III, 1890, p. 161.
- * ? *Unio conus* SPENGLER, Skriv. Selsk. Nat., III, 1793, p. 60.
- * *Mya ovalis* DONOVAN, Brit. Shells, III, 1801, pl. LXXXIX.—* TURTON, Brit. Fauna, 1807, p. 146.
- * ? *Unio oralis* SOWERBY, Rec. and Foss. Shells, No. XVI, 1823, fig. —. * ? REEVE, Conch. Syst., I, 1841, p. 117, pl. LXXXVII, fig. 1.—* CHIENU, Bib. Conch., 1st ser., I, 1845, p. 67, pl. XXIV, figs. 1-3.—* PÆTEL, Conch. Sam., III, 1890, p. 162.
- * *Mya nodulosa* WOOD (part), Gen. Conch., I, 1815, p. 106, pl. XXII, figs. 3, 4.²
- * *Unio nodulosa* LAMARCK, An. sans Vert., VI, 1819, p. 78.
- * *Unio rostrata* LAMARCK, An. sans Vert., VI, 1819, p. 77.—* C. PFEIFFER, Nat. Deuts. L. and Suss. Moll., Pt. 1, 1821, p. 114, pl. v, fig. 8.—* WAARDENBERG, Com. Hist. Nat. Animalium, 1827, p. 36.—* LAMARCK, Enc. Méth., II, 1830, p. 586.—* MICHAUD, Comp. Hist. Moll. Fr., 1831, p. 108, pl. XVI, fig. 25.—* BROWN, L. and F. W. Conch., 1836, p. 109, pl. XX, figs. 1, 2; * Ill. Rec. Conch., 1844, p. 82, pl. XXXII, figs. 9-12.—* PUTON, Moll. Vosges, 1847, p. 70.—* GRAS, Moll. Isere., 1840, p. 71, pl. v, fig. 21.
- * *Unio rostratus* CHIENU, Man., II, 1859, p. 137, fig. 658.—* WESTERLUND, Faun. der Pal., II, Pt. 7, 1890, p. 104.—* PÆTEL, Conch. Sam., III, 1890, p. 166.—* LOCARD, Les Coq. de Fr., 1893, p. 207.—WESTERLUND, Act. Soc. F. and F. F., XIII, No. 7, 1897, p. 165.
- * *Unio manca* LAMARCK, An. sans Vert., VI, 1819, p. 80.
- * *Unio mancus* DROUET, Moll. Côte d'Or, 1867, p. 103.—* WESTERLUND, Faun. der Pal., II, Pt. 7, 1890, p. 85.
- * *Unio limosus* NILSSON, Hist. Moll. Svec., 1822, p. 110.—* ROSSMASSLER, Icon., III, 1836, p. 24, pl. XIII, fig. 199.—* KUSTER, Conch. Cab. Unio, 1854, p. 80, pls. XXI, XXII, XXIII.—* PÆTEL, Conch. Sam., III, 1890, p. 157.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 115.—* LOCARD, Coq. de Fr., 1893, p. 211.—* WESTERLUND, Act. Soc. F. and F. F., XIII, No. 7, 1897, p. 166.
- * *Unio deshayesii* MICHAUD, Comp. Hist. Moll. Fr., 1831, p. 107, pl. XVI, fig. 30.—* ROSSMASSLER, Icon., III, 1836, p. 23, pl. XIII, fig. 197.—* GRAS, Moll. Isere. App., 1840, p. 21.—* BROWN, Ill. Rec. Conch., 1844, p. 81, pl. XXXII, figs. 1-4; L. and F. W. Conch., 1845, p. 108, pl. XX, figs. 3, 4.—* LOCARD, Coq. de Fr., 1893, p. 210.
- * *Unio dubius* FITZINGER, Syst. Verz., 1833, p. 119.
- * *Unio michaudiana* DES MOULINS, Actes Soc. Linn. Bord., VI, 1833, p. 27, plate.
- * *Unio longirostris*, ROSSMASSLER, Icon., III, 1836, p. 26, pl. XIV, fig. 200; XI, 1842,

¹ Gmelin refers to Conch. Cabinet X, p. 347, pl. CLXX, fig. 1650, which is, I am quite certain, a young *U. pictorum*.

² Two species are figured; figs. 1 and 2 are a heavy, inflated Chinese species; figs. 3 and 4 are no doubt *pictorum*.

p. 13, pl. LIV, fig. 738.—*STABILE, Faun. Lug., 1845, p. 60.—*DROUET, Mem. Ac. Dijon, VII, 1882, p. 11 (reprint).—*WESTERLUND, Faun. der Pal., II, Pt. 7, 1890, p. 117.—*PÆTEL, Conch. Sam., III, 1890, p. 158.—*LOCARD, Coq. de Fr., 1893, p. 209.

**Potamida sicula* SWAINSON, Treatise on Mal., 1840, p. 282, fig. 58.

**Unio siculus* HANLEY, Biv. Shells, 1856, p. 383, pl. xx, fig. 19.—*PÆTEL, Conch. Sam., III, 1890, p. 167.

**Unio aradæ* PHILIPPI, Enum. Moll. Sic., III, 1844, p. 49.—*KUSTER, Conch. Cab. Unio, 1854, p. 105, pl. XXVIII, fig. 6.—*KOBELT, Icon., IV, 1876, p. 62, pl. CXVII, figs. 1146, 1147.—*DROUET, Jl. de Conch., XXIX, 1881, p. 25.

**Unio dactylus* MORELET, Moll. Port., 1845, p. 110, pl. XIV, fig. 2.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 111.—*KOBELT, Icon., new ser., VI, 1893, p. 98, pl. CXXX, fig. 1132.

**Uniomucidus* MORELET, Moll. Port., 1845, p. 111.—*KOBELT, Icon., new ser., VI, 1893, p. 98, pl. CLXXX, figs. 1130, 1131.—*PÆTEL, Conch. Sam., III, 1890, p. 160.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 111.

**Unio quinqueannulatus* KUSTER, Conch. Cab. Unio, 1854, p. 93, pl. XXV, figs. 3, 4.—*PÆTEL, Conch. Sam., III, 1890, p. 165.

**Unio pallens* KUSTER, Conch. Cab. Unio, 1854, p. 95, pl. XXV, fig. 5; XXVI, fig. 1.

**Unio viridiflavus* KUSTER, Conch. Cab. Unio, 1854, p. 96, pl. XXVI, figs. 2, 3.—*PÆTEL, Conch. Sam., III, 1890, p. 171.

**Unio petrovichii* KUSTER, Conch. Cab. Unio, 1854, p. 98, pl. XXVI, fig. 5; XXVII, fig. 1.—*PÆTEL, Conch. Sam., III, 1890, p. 163.

**Unio maltzani* KUSTER, Conch. Cab. Unio, 1854, p. 106, pl. XXIX, figs. 1, 2.—*PÆTEL, Conch. Sam., III, 1890, p. 158.

**Unio baltonicus* KUSTER, Conch. Cab. Unio, 1861, p. 231, pl. LXXVIII, fig. 1.—*SERVAIR, Hist. Mal. Bal., 1881, p. 98.

**Unio proechus* BOURGUIGNAT, Rev. et Mag., XIV, 1862, pl. XIX, figs. 1-3; XV, 1863, p. 19, XIX.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 118.

**Unio actephilus* BOURGUIGNAT, Rev. et Mag., XIV, 1862, pl. XIX, figs. 7, 8, XX, fig. 3; XV, 1863, p. 20.

**Unio lawleyanus* GENTILUOMO, Bull. Mal. Ital., I, 1868, p. 54, pl. IV, figs. 1-3.

Europe generally; Siberia east to the Lena River and perhaps farther; south into Asia Minor?; Algiers?

† UNIO PLATYRHYNCHUS Rossmassler.

**Unio platyrhynchus* ROSSMASSLER, Icon., II, 1835, p. 22, pl. IX, fig. 130; V, 1837, p. 20, pl. XXIV, fig. 338.—*PORRO, Mal. Como, 1838, p. 117.—*HANLEY, Test. Moll., 1842, p. 205; *Biv. Shells., 1843, p. 205, pl. XXIII, fig. 53.—*CATLOW and REEVE, Conch. Nom., 1845, p. 62.—*KUSTER, Conch. Cab. Unio, 1854, p. 77, pls. XIX, XX.—*SOWERBY, Conch. Icon., XVI, 1856, pl. XXX, fig. 154.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—*PÆTEL, Conch. Sam., III, 1890, p. 163.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 118.

**Margarita (Unio) platyrhynchus* LEA, Syn., 1836, p. 35; 1838, p. 24.

**Margaron (Unio) platyrhynchus* LEA, Syn., 1852, p. 36; 1870, p. 58.

**Unio fiscalianus* KLECIACH, Atti. Soc. Ital., XV, 1872, p. 92.—*PFEIFFER and KOBELT, Mal. Blatt., XX, 1873, p. 92, fig.

Unio limosus *KOBELT, Icon., new ser., VI, 1893, p. 44, pl. CLVII, figs. 1024-1027.¹

Central and southwestern Europe.

¹ According to Kobelt in above reference, his *platyrhynchus* = the *limosus* of Nilsson. Nilsson does not figure his species in Hist. Moll. Svcc., but refers to pl. v, fig. 10, in Pfeiffer's Land and Susswasser Mollusken, which is plainly a *Unio pictorum*. Kobelt's figures, the latter three certainly, are genuine *platyrhynchus*, which I consider a perfectly good species.

† UNIO PLATYRINCHOIDEUS Dupuy.

Unio platyrhynchoides DUPUY, Cat. Ext. Gall. Test, 1849;¹ * Hist. Moll. Fr., VI, 1852, p. 619, pl. XXVII, fig. 16.—? DROUET, Nay. Fr., II, 1857, p. 100, pl. IX, fig. 1.—PÆTEL, Conch. Sam., III, 1890, p. 163.

* *Unio platyrhynchoides* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 121.

* *Unio platyrhynchoides* SOWERBY, Conch. Icon., XVI, 1868, pl. LXX, fig. 358.

* *Unio danielis* GASSIES, Actes. Soc. Linn. de Bord, XXVI, 1866, p. 132, pl. 1, fig. 8.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 110.—* PÆTEL, Conch. Sam., III, 1890, p. 150.—* LOCARD, Coq. de Fr., 1893, p. 199.

Unio lardelianus PECCHIOLO, Bull. Mal. It., II, 1869, p. 163, pl. v.

South France.

† UNIO ELONGATULUS C. Pfeiffer.

* *Unio elongatula* C. PFEIFFER,² Nat. Dents., L. and S. Moll., II, 1825, p. 35, pl. VIII, figs. 5, 6.—ROSSMASSLER, Icon., II, 1835, p. 23, pl. IX, fig. 132; XII, 1844, p. 27, pl. LVI, fig. 751; VI, 1879, p. 42, pl. CLXII, figs. 1644, 1645.—PORRO, Mal. Como, 1838, p. 114.—CATLOW and REEVE, Conch. Nom., 1845, p. 59.—KUSTER, Conch. Cab. Unio., 1854, p. 104, pl. XXVIII, figs. 4, 5.—DROUET, Nay. Fr., II, 1857, p. 91, pl. VI, fig. 2.—H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXV, fig. 451.—DROUET, J. de Conch., XXVII, 1879, p. 331.—* CLESSIN, Moll. Oest., 1887, p. 735.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 163.—* PÆTEL, Conch. Sam., III, 1890, p. 151.—* LOCARD, Coq. de Fr., 1893, p. 166, fig. 180.

* *Margarita (Unio) elongatulus* LEA, Syn., 1836, 1838, p. 24.

* *Margaron (Unio) elongatulus* LEA, Syn., 1852, p. 37, 1870, p. 59.

Central Europe.

† UNIO TUMIDUS Retzius.

* *Unio tumidus* RETZIUS, Diss. Hist. Nat., 1778, p. 17.—SPENGLER, Skriv. Selsk. Nat., III, 1793, p. 57.—NILSSON, Hist. Moll. Svec., 1822, p. 109.—ROSSMASSLER, Icon., I, 1835, p. 117, pl. III, figs. 70, 70a, 70b; III, 1836, p. 27, pl. XIV, figs. 202–204; VIII, 1838, p. 41, pl. XL, fig. 543; XII, 1844, p. 32, pl. LX, figs. 772–778.³—* BROWN, L. and F. W. Conch., 1836, p. 109, pl. XXI, figs. 8, 9.—* HANLEY, Test. Moll., 1842, p. 205; * Biv. Shells, 1843, p. 205.—* BROWN, Ill. Rec. Conch., 1844, p. 82, pl. XXXII, figs. 5–8.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* STABILE, Faun. Lug., 1845, p. 61, pl. III, fig. 74.—* STEIN, Die Leb. Schnecken, 1850, p. 103.—* DUPUY, Hist. Moll. Fr., 1852, p. 655, pl. XXVIII, fig. 20.—* FORBES and HANLEY, Hist. Brit. Moll., II, 1853, p. 140, pl. XI, fig. 1.—* KUSTER, Conch. Cab. Unio., 1854, p. 71, pls. XVII, XVIII.—* MOQUIN-TANDON, Moll. Terr. Fluv., II, 1855, p. 577, pl. LI, figs. 11, 14.—* VON WAHL, Arch. für Naturkunde Liv., 2d ser., I, p. 115.—* NORDENSKIÖLD and NYLANDER, Fin. Moll., 1856, p. 85, pl. VI, figs. 7, 8.—* GRAY, Thurton's Man., 1857, p. 297.—* DROUET, Nay. Fr., II, 1857, p. 110, pl. IX, fig. 2.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* SOWERBY, Ill. Int. British Shells, 1859, pl. VII, No. 3.—* JEFFREYS, Brit. Conch., I, 1862, p. 32.—* REEVE, L. and F. W. Moll. Brit., 1863, p. 219, fig. 1.—* REEVE, Conch. Icon., XVI, 1865, pl.

¹ Credited by Lea to the above without page reference. I have not seen the paper.

² Pfeiffer credits this to Mühlfeld in *litt.* It is a doubtful species, perhaps a variety of *pictorum*, but is generally smaller, thinner, and more compressed, and is rather more biangulate behind. Pfeiffer's two figures represent somewhat different forms, but they may be the same thing.

³ Fig. 776 is probably a form of *batarus*.

- XXV, fig. 124.—*TATE, L. and F. W. Moll. Brit., 1866, pl. II, fig. 2.—*KOBELT, Faun. Nass. Moll., 1871, p. 239.—*LEHMAN, Die Schnecken, 1873, p. 292.—*WESTERLUND, Faun. S. N. and D., 1873, p. 572.—*CLESSIN, Deuts. Ex. Moll., 1876, p. 458, fig. 299.—*JORDAN, Jahrb. Dents. Mal. Ges., VI, 1879, p. 305.—*DROUET, Union. Russ., 1881, p. 9; *Mem. Ac. Dijon, 3d ser., VII, 1882, p. 12 (reprint).—*L. ADAMS, Coll. Man., 1884, p. 18, pl. L, fig. 10.—*CLESSIN, Moll. Oest., 1887, p. 738.—*PÆTEL, Conch. Sam., III, 1890, p. 170.—*KOBELT, Icon., new ed., VI, 1893, p. 87, pl. CLXXIII, fig. 1115.—*LOCARD, Coq. Fr., 1893, p. 212.—*WESTERLUND, Act. Soc. F. and F. F., XIII, 1897, p. 162.
- **Margarona (Unio) tumidus* LEA, Syn., 1852, p. 36; 1870, p. 58.
- **Unio tumida* C. PFEIFFER, Nat. Deuts. L. and S. Moll., II, 1825, p. 34, pl. VII, figs. 2, 3; pl. VIII, figs. 1, 2.—*KLEEGER, Moll. Bor., 1828, p. 38.—*PUTON, Moll. Vosges, 1847, p. 71.
- **Mya depressa* DONOVAN, Brit. Shells, III, 1801, pl. CI.—*CHENU, Bib. Conch., 1st Ser., I, 1845, p. 71, pl. XXVI, figs. 1-3.
- **Unio depressus* PÆTEL, Conch. Sam., III, 1890, p. 150.
- **Mya orata* DONOVAN, Brit. Shells, IV, 1802, pl. CXXII.¹—*MATON and RACKETT, Tr. Linn. Soc. Lond., VIII, 1807, p. 39.—*WOOD, Gen. Conch., I, 1815, p. 105, pl. XIX, fig. 5.—*DILLWYN, Cat., I, 1817, p. 50.—*TURTON, Conch. Dict., 1819, p. 106.—*WOOD, Ind. Test., 1825, p. 12, pl. II, fig. 27c; *rev. ed., 1856, p. 16, pl. II, fig. 27.—*CHENU, Bib. Conch., 1st ser., I, 1845, p. 82, pl. XXXII, figs. 1-3.
- **Mysca orata* TURTON, Conch. Ins. Brit., 1822, p. 246; Man. Shells Brit. Is., 1831, p. 21, fig. 12.—*SWAINSON, Treat. on Mal., 1840, p. 277, fig. 56.
- **Mysca solida* TURTON, Conch. Ins. Brit., 1822, p. 246, pl. XVI, fig. 2; Man. Shells Brit. Is., 1831, p. 22, fig. 13.
- **Mya ovalis* MONTAGU, Test. Brit., 1803, p. 34.—*FLEMING, Hist. Br. Anim., 1828, p. 416.
- **Unio ovalis* BROWN, L. and F. W. Conch., 1836, p. 101, pl. XVIII, figs. 4, 5; Ill. Rec. Conch., 1844, p. 82, pl. XXXI, figs. 12-14.—*LEACH, Syn. Moll. Gt. Br., 1852, p. 322.
- **Margarita (Unio) ovalis* LEA, Syn., 1836, p. 35; 1838, p. 24.
- **Unio muelleri* ROSSMASSLER, Icon., VIII, 1838, p. 41, pl. XI, fig. 541; XI, 1842, p. 13, pl. LIV, fig. 739.
- **Unio pictorum* BROWN, Ill. Rec. Conch., 1844, p. 81, pl. XXXI, figs. 8-11.

Northern and middle Europe; eastern Siberia.

UNIO TURTONI Payraudeau.

- **Unio turtoni* PAYRAUDEAU, Cat. Moll. Corse, 1826, p. 65, pl. II, fig. 65.²—*PHILIPPI, Moll. Sic., 1836, p. 67.—*ROSSMASSLER, Icon., VII, 1838, p. 25, pl. XXXV, fig. 492.—*CATLOW and REEVE, Conch. Nom., 1845, p. 65.—*REQUIEN, Cat. Coq. Corse, 1848, p. 29.—*DUPUY, Hist. Moll. Franc., 1852, p. 651, pl. XXVII, fig. 17.—*PAULLUCCI, Bull. Soc. Mal. It., V, 1879, p. 107.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 160.—*PÆTEL, Conch. Sam., III, 1890, p. 170.
- **Unio capigliolo* PAYRAUDEAU, Cat. Moll. Corse, 1826, p. 66, pl. II, fig. 4.—*ROSSMASSLER, Icon., V and VI, 1837, p. 22, pl. XXIV, fig. 341; XII, 1844, p. 28, pl. LVII, figs. 755, 756.—*MOQUIN-TANDON, Moll. Terr. Fluv. Fr., II, 1855, p. 574, pl. I, figs. 3, 4.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 146.—*PÆTEL, Conch. Sam., III, 1890, p. 147.
- **Unio capigliolo* var. *bondini* PÆTEL, Conch. Sam., III, 1890, p. 147.

¹ First described by Lister in *Historia Animalium Angliæ*, 1678.

² This has often been confounded with *U. pictorum*, but is generally more rhomboid, higher at the posterior part of the ligament, and more plainly marked at the rest periods than that species.

- * *Unio pictorum* GUERIN, Icon. Règne Anim., II, 1829-1844, pl. XXVIII, fig. 16.—
 * BOURGUIGNAT, Mal. Alg., II, 1864, p. 292, pl. XXII, figs. 6-11.—* REEVE,
 Conch. Icon., XVI, 1865, pl. XXV, fig. 123.
- * *Unio requienii* MICHAUD, Comp. Hist. Nat. Moll. Fr., 1831, p. 106, pl. XVI, fig. 24.—
 * ROSSMASSLER, Icon., III, 1836, p. 24, pl. XIII, fig. 198.—* GRAS, Moll. Isere,
 1840, p. 21.—* STABILE, Faun. Lug., 1845, p. 62, pl. III, fig. 1786.—* GASSIES,
 Moll. Agen., 1849, p. 195, pl. I, figs. 4, 5.—* DUPUY, Hist. M. Fr., 1852, p. 652,
 pl. XXVII, fig. 18.—* KUSTER, Conch. Cab. Unio, 1854, p. 126, pl. XXI, fig. 7;
 XXVI, figs. 1-3; XXXVII, figs. 2-4.—* DROUET, Nay. Fr., II, 1857, p. 93, pl. VII,
 figs. 1-3.—* MOUSSON, Coq. Terr. Fluv. Pal., 1861, p. 66.—* DROUET, Moll. Côte
 d'Or, 1867, p. 102.—* BENOIT, Mus. Cat. Sic., 1881, p. 170.—* CLESSIN, Moll.
 Oest., 1887, p. 731, fig. 494.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 121.—
 * PÆTEL, Conch. Sam., III, 1890, p. 165.
- * *Unio lobata* PHILIPPI, Moll. Sic., 1836, p. 67.
- * *Unio bandanii* ROSSMASSLER, Icon., V, 1837, p. 22, pl. XXIV, fig. 341.
- * *Unio pallens* ROSSMASSLER, Icon., XI, 1842, p. 13, pl. LIV, fig. 740.—* WESTER-
 LUND, Faun. Pal., II, Pt. 7, 1890, p. 114.
- * *Unio hispanus* ROSSMASSLER, Icon., XII, 1844, p. 26, pl. LVI, fig. 747.—* BOUR-
 GUIGNAT, Moll. Pen. Con., 1863, p. 46, pl. XXIV, figs. 1-3.—* BOURGUIGNAT,
 Rev. et Mag., XVII, 1865, p. 344, pl. XXIII, figs. 1-3.—* PÆTEL, Conch. Sam.,
 III, 1890, p. 155.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 139.
- * *Unio alceroni* COMPANYO and MASSOT, Bull. Soc. Pyr. d'Or, VI, Pt. 2, 1845, p.
 234, fig. —* BOURGUIGNAT, Moll. Pen. Con., 1863, p. 49, pl. XXIII, figs. 1-3;
 Rev. et Mag., XVII, 1865, pl. XIX, figs. 1-3.
- * *Unio rousii* DUPUY, Hist. Moll., VI, 1852, p. 653, pl. XVIII, fig. 18.—* MUSGRAVE,
 Phot. Conch., 1863, pl. II, fig. 7.
- * *Unio valentinus* ROSSMASSLER, Icon., III, 1854, p. 37, pl. LXIX, p. 852.—* BOUR-
 GUIGNAT, Moll. Pen. Con., 1863, p. 45, pl. XXVII; Rev. et Mag. Zool., XVII,
 1865, p. 343, pl. XX.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 140.—
 * PÆTEL, Conch. Sam., III, 1890, p. 171.
- * *Margaron (Unio) valentinus* LEA, Syn., 1870, p. 52.
- * *Unio arduisianus* MOQUIN-TANDON, Hist. Moll. Fr., II, 1855, p. 575.—* WESTER-
 LUND, Faun. Pal., II, Pt. 7, 1890, p. 109.—* LOCARD, Coq. de Franc., 1893,
 p. 205.
- * *Unio graellsianus* BOURGUIGNAT, Moll. Pen. Con., 1863, p. 47, pl. XXIII, figs.
 4-7; * Rev. et Mag., 1865, p. 345, pl. XIX, figs. 4-7.—* WESTERLUND, Faun.
 Pal., II, Pt. 7, 1890, p. 140.
- *? *Unio letourneuzi* BOURGUIGNAT, Mal. Alg., 1864, p. 289, pl. XVII, fig. 47.—
 * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 58.—* PÆTEL, Conch. Sam.,
 III, 1890, p. 157.
- * *Unio courquinianus* BOURGUIGNAT, Moll. Pen. Con., 1863, p. 48; * part, Rev.
 et Mag., 1865, p. 346, pl. XXII, figs. 1, 2.
- * *Unio siculus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXI, fig. 364.³
- * *Unio ravoisieri* var. *issericus* KOBELT, Icon., new ser., I, 1884, p. 65, pl. XXVIII,
 fig. 215.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 142.
- * *Unio alexandri* KOBELT, Icon., 1st sup., 1895, p. 14, pl. II, fig. 2.

Entire circummediterranean region.

¹ Credited by Rossmassler to Parreyss. I do not know where the latter described it or whether he described it at all.

² Moquin-Tandon credits this to a letter from Reynies, 1843, and gives a reference to p. 5, pl. I, figs. 6, 7, of the letter. I do not know that this was ever published. He makes it a variety of *requienii*.

³ This is *Potamida sicula* Swainson according to Sowerby. I refer Swainson's shell to *Unio pictorum*.

† UNIO RAVOISIERI Deshayes.¹

* *Unio ravoisieri* DESHAYES, Hist. Nat. Moll. Aceph. Alg., 1848, pl. CVIII, figs. 4-7.—* BOURGUIGNAT, Mal. Alg., 1864, p. 291, pl. XX, figs. 5-10.—* PÆTEL, Conch. Sam., III, 1890, p. 165.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 142.

* *Margaron (Unio) ravoisieri* LEA, Syn., 1870, p. 48.

* *Unio moreleti* DESHAYES, Hist. Moll. Alg., 1848, pl. CIX, figs. 1-4; CXII, fig. 5.—* KOBELT, Icon., new ser., II, 1886, p. 4, pl. XXXIII, fig. 228.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 141.—* PÆTEL, Conch. Sam., III, 1890, p. 160.

* *Margaron (Unio) moreleti* LEA, Syn., 1870, p. 60.

*? *Unio maccarthyanus* BOURGUIGNAT, Moll. Pen. Con., 1886, p. 221, pl. XXXIV, figs. 8-11.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 57.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

Algiers.

† UNIO MUSSOLIANUS Kuster.

* *Unio mussolianus* KUSTER, Conch. Cab. Unio, 1861, p. 244, pl. LXXXII, fig. 1.³—* PÆTEL, Conch. Sam., III, 1890, p. 160.

* *Unio bourguignatianus* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 189; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 54, pl. XVIII, fig. 51; *Obs., XI, 1867, p. 55, pl. XVIII, fig. 51.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 178.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

* *Margaron (Unio) bourguignatianus* LEA, Syn., 1870, p. 39.

* *Unio rasns* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 189; *Jl. Acad. N. Sci. Phila., VI, 1866, p. 50, pl. XVII, fig. 47; *Obs., XI, 1867, p. 54, pl. XVII, fig. 47.—* PÆTEL, Conch. Sam., III, 1890, p. 165.

* *Margaron (Unio) rasns* LEA, Syn., 1870, p. 58.

* *Unio mosulensis* LEA, Pr. Acad. N. Sci. Phila., VII, 1863, p. 190; *Jl. Acad. N. Sci. Phila., VI, 1866, p. 52, pl. XVII, fig. 49; *Obs., XI, 1867, p. 56, pl. XVII, fig. 49.

* *Margaron (Unio) mosulensis* LEA, Syn., 1870, p. 48.

* *Margaritana mosulensis* PÆTEL, Conch. Sam., III, 1890, p. 173.

Assyria.

UNIO HUETI Bourguignat.

* *Unio hueti* BOURGUIGNAT, Rev. et Mag., VII, 1855, p. 332, pl. VIII, figs. 1-4.—VON MARTENS, Vorderas. Conch., 1874, p. 35, pl. VII, fig. 54.—* PÆTEL, Conch. Sam., III, 1890, p. 155.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 171.⁴

Asia Minor.

¹ Doubtfully distinct from *U. turtoni*. It is not quite so rhomboid as that species, which is also found in Algiers, but I have seen much intermediate material which hints at a connection of the two.

² The magnificent work in which this species with other Uniones was figured was never finished, and no descriptions of the *Naiades* were written. The beautiful colored figures are wonderfully characteristic and are accompanied by names.

³ Credited to Parreyss in catalogue.

⁴ This is quite likely only a variety of the preceding, but seems to be thinner and is more evenly elliptical. Von Martens believes it to be the same as Lea's *mosulensis*.

UNIO EUCIRRUS Bourguignat.

- * *Unio eucirrus* BOURGUIGNAT, Mag. Zool., IX, 1857, p. 20, pl. VIII, figs. 4-6.—
 * KOBELT, Icon., VII, 1880, p. 82, pl. CCVI, fig. 2101.—* WESTERLUND, Faun.
 Pal., II, Pt. 7, 1890, p. 171.—* PÆTEL, Conch. Sam., III, 1890, p. 152.
 * Margaron (*Unio eucirrus* LEA, Syn., 1870, p. 46.

Asia Minor. Probably a form of *hueti*.

† UNIO TIGRIDIS Bourguignat.¹

- * *Unio truncatus* SWAINSON, Zool. Ill., 2d ser., I, 1829, pl. X.—* HANLEY, Test. Moll.,
 1842, p. 187; * Biv. Shells., 1843, p. 187.—* CATLOW and REEVE, Conch. Nom.,
 1845, p. 64.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* SOWERBY,
 Conch. Icon., XVI, 1868, pl. LXXXV, fig. 453.
 * *Margarita (Unio) truncatus* LEA, Syn., 1836, p. 21; 1838, p. 18.
 * Margaron (*Unio) truncatus* LEA, Syn., 1852, p. 26; 1870, p. 39.
 * *Unio tigridis* BOURGUIGNAT, Test. Nov. Sanl., 1852, p. 30; * Cat. Rais. Moll., 1853,
 p. 77, pl. IV, figs. 7-9.—* KUSTER, Conch. Cab. Unio, 1861, p. 227, pl. LXXVII,
 fig. 1.—* KOBELT, Icon., new ser., II, 1886, p. 2, pl. XXXII, fig. 226.—* WESTER-
 LUND, Faun. Pal., II, Pt. 7, 1890, p. 175.—* PÆTEL, Conch. Sam., III, 1890, p.
 169.
 * *Unio dignatus* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 189; * Jl. Acad. N. Sci.
 Phila., VI, 1866, p. 51, pl. XVII, fig. 48; * Obs., XI, 1867, p. 55, pl. XVII, fig. 48.—
 * PÆTEL, Conch. Sam., III, 1890, p. 150.
 * Margaron (*Unio) dignatus* LEA, Syn., 1870, p. 39.
 * Margaron (*Unio) tigris* LEA, Syn., 1870, p. 39.
Unio tigris FERUSSAC, manuscript.²
 * ? *Unio kisonis* KOBELT, Icon., 1st sup., 1895, p. 17, pl. VII, figs. 2, 3.³

Asia Minor; Assyria.

UNIO PIETRI Locard.

- * *Unio pietri* LOCARD, Comptes Rendus, XCI, 1880, p. 500; * Arch. Mus. Lyon, III,
 1883, p. 210, pl. XX, figs. 17-19.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890,
 p. 170.
 * *Unio petroi* PÆTEL, Conch. Sam., III, 1890, p. 163.
 * *Unio lorteti* LOCARD, Comptes Rendus, XCI, 1880, p. 502; * Arch. Mus. Lyon, III,
 1883, p. 215, pl. XXI, figs. 7-12.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890,
 p. 173.
 * *Unio tristrami* LOCARD, Arch. Mus. Lyon, III, 1883, p. 209, pl. XX, figs. 15, 16.—
 * PÆTEL, Conch. Sam., III, 1890, p. 170.—* WESTERLUND, Faun. Pal., II, Pt. 7,
 1890, p. 170.
 * *Unio tiberiadensis* LOCARD, Arch. Mus. Lyon, III, 1883, p. 216, pl. XXI, figs. 13-15.—
 * PÆTEL, Conch. Sam., III, 1890, p. 169.—* WESTERLUND, Faun. Pal., II, Pt.
 7, 1890, p. 174.

¹ Swainson gave the name *Unio truncatus* to this species in 1829, but that name had been applied to a *Unio* by Spengler in 1793. Bourguignat credits this to Ferussac manuscript under the name of *Unio tigris*.

² Lea credits this to Ferussac manuscript in museum at Paris. No description has ever been published, so far as I know, though several authors have cited it and credited it to Ferussac. A specimen in the Lea collection which is no doubt authentic is apparently a delicate young *tigridis*.

³ Young shells, and it is very hard to say whether they are *tigridis* or *terminalis*.

- * *Unio prosacrus* LOCARD, Arch. Mus. Lyon, III, 1883, p. 219, pl. XXI, figs. 16, 17.—
 * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 174.
 * *Unio ariacus* LOCARD, Arch. Mus. Lyon, III, 1883, p. 242, pl. XX, figs. 20-23.—
 * PÆTEL, Conch. Sam., III, 1890, p. 145.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 170.
 * *Unio subtigridis* LOCARD, Arch. Mus. Lyon, III, 1883, p. 245, pl. XXI, figs. 18-20.—
 * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 175.—* PÆTEL, Conch. Sam., III, 1890, p. 168.
 * *Unio aremprosthus* LOCARD, Arch. Mus. Lyon, III, 1883, p. 246, pl. XXI, figs. 21-23.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 175.
 * *Unio chantri* LOCARD, Arch. Mus. Lyon, III, 1883, p. 247, pl. XXII, figs. 1-7.—
 * PÆTEL, Conch. Sam., III, 1890, p. 147.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 176.¹

Asia Minor.

UNIO ZABULONICUS Locard.

- * *Unio zabulonicus* LOCARD, Arch. Mus. Lyon, III, 1883, p. 220, pl. XXII, figs. 11-18.—
 * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 177.—* PÆTEL, Conch. Sam., III, 1890, p. 172.
 * *Unio antiochianus* LOCARD, Arch. Mus. Lyon, III, 1883, p. 249, pl. XXII, figs. 14-16.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 176.

Asia Minor.

† UNIO TERMINALIS Bourguignat.

- * *Unio terminalis* BOURGUIGNAT, Test. Noviss., 1852, p. 31; * Cat. Rais. Moll., 1853, p. 76, pl. III, figs. 4-6; * Jl. de Conch., IV, 1853, p. 74, pl. III, figs. 10' 10".—* MOUSSON, Coq. Terr. et Fluv. Pal., 1861, p. 65.—* KOBELT, Icon., IV, 1876, p. 65, pl. CXIX, fig. 115.—* PÆTEL, Conch. Sam., III, 1890, p. 169.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 174.
 * Margaron (*Unio*) *terminalis* LEA, Syn., 1870, p. 39.

Lake Tiberias.

UNIO GRELLOISIANUS Bourguignat.

- * *Unio grelloisianus* BOURGUIGNAT, Rev. et Mag., VIII, 1856, p. 227, pl. XI, figs. 1-4; Moll. Pen. Con., 1863, p. 74, pl. XXIII, figs. 4-7.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, pp. 140, 173.—* PÆTEL, Conch. Sam., III, 1890, p. 154.
 * *Unio lunulifer* BOURGUIGNAT, Rev. et Mag., VIII, 1856, p. 227, pl. XI, figs. 5-8.—
 * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 177.—* PÆTEL, Conch. Sam., III, 1890, p. 158.
 * Margaron (*Unio*) *lunulifer* LEA, Syn., 1870, p. 37.
 * *Unio jordanicus* BOURGUIGNAT, Rev. et Mag., VIII, 1856, p. 228, pl. X, figs. 1-4.—
 * MOUSSON, Coq. Terr. Fluv. Pal., 1861, p. 66.—* PÆTEL, Conch. Sam., III, 1890, p. 156.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 172.
 * Margaron (*Unio*) *jordanicus* LEA, Syn., 1870, p. 44.

Jordan River.

¹All these so-called species of Locard, some of which he credits to the manuscripts of Bourguignat and Lortet, have high beaks placed at some distance from the anterior end, are wedge-shaped and more or less pointed posteriorly. I think it quite probable that all of them, together with the next species, are mere variations of Bourguignat's old *U. terminalis*. Of course Locard has figured every possible variation and distortion and called it a species.

UNIO ELLIPSOIDEUS Locard.

- * *Unio ellipsoideus* LOCARD, Arch. Mus. Lyon., III, 1883, p. 211, pl. XXI, figs. 1-3.¹—
 * WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 171.—* PÆTEL, Conch. Sam., III, 1890, p. 151.
 * *Unio genezarethensis* LOCARD, Arch. Mus. Lyon., III, 1883, p. 213, pl. XXI, figs. 4-6.—* PÆTEL, Conch. Sam., III, 1890, p. 153.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 172.
 * *Unio janberti* LOCARD, Arch. Mus. Lyon., III, 1883, p. 248, pl. XXII, figs. 8-10.—
 * PÆTEL, Conch. Sam., III, 1890, p. 155.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 176.
 * *Unio zabulonius* KOBELT, Icon., new ser., VI, 1893, p. 96, pl. CLXXIX, fig. 1129.
 * *Unio lortiti* KOBELT, Icon., 1st sup., 1895, p. 14, pl. v, fig. 3.
 * *Unio kobelti* ROLLE, Icon., 1st sup., 1895, p. 15, pl. vi, fig. 3.
 * *Unio pietri* KOBELT (part), Icon., 1st sup., 1895, p. 16, pl. vi, figs. 1, 2.
 * *Unio herodes* KOBELT, Icon., 1st sup., 1895, p. 17, pl. vi, fig. 4.

Lake Tiberias.

†UNIO DELICATUS Lea.²

- * *Unio delicatus* LEA, Pr. Acad. N. Sci. Phila., VII, 1863, p. 189.—* JI. Acad. N. Sci. Phila., VI, 1866, p. 58, pl. XIX, fig. 56.—* Obs., XI, 1867, p. 62, pl. XIX, fig. 56.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 178.—* PÆTEL, Conch. Sam., III, 1890, p. 150.
 * Margaron (*Unio*) *delicatus* LEA, Syn., 1870, p. 42.

Orontes River, Syria.

(Group of *Unio littoralis*.)

Shell rather solid, subinflated, rounded rhomboid, with a faint posterior ridge, usually slightly biangulate behind and often becoming arcuate when old; beaks prominent and full; beak sculpture consisting of numerous rather fine, subparallel ridges or corrugations which are sometimes a good deal broken up, and which extend well out on the disk, but begin at the beaks as normal, somewhat coarse *Unio* sculpture, sometimes with fine radial lines posteriorly; pseudocardinals rather solid, subcompressed, smooth below; laterals straight or slightly curved; cavity of the beaks rather deep; muscle scars distinct.

I have never seen the soft parts of any member of this group. Quite a number of descriptions have been published of various nominal species, most of which go into details as to the color of the different parts, but do not give an atom of information as to real characters. The animal is dark or highly colored, and seems to be gravid in summer, and no doubt carries the young in the outer gills alone. Gills large, wider behind, inner the larger, especially in front; mantle thickened at the edges; palpi very large, elliptical, rounded behind, hanging at an angle of 45°; branchial opening large, strongly fringed.³

¹ I think this is a form of *grelloisianus*. Some forms of the latter are very much like *U. tigrinus* Bourguignat, but he says that it has roughened beaks while those of *tigrinus* are always smooth.

² I have only seen the type, a young shell, and its relations are doubtful. The sharp, rather pustulous, beak sculpture, and the shining, yellowish epidermis are like the *pictorum* group, but its form is peculiar. It may not come from Syria at all.

³ From a figure in Mal. d'Algérie, II, pl. XIX.

† UNIO LITTORALIS Lamarck.¹

- * *Unio littoralis* LAMARCK, Syst. An. sans Vert., 1801, p. 114.—* DRAPARNAUD, Tab. Moll. Fr., 1801, p. 101; *Hist. Moll. Fr., 1806, p. 135, pl. x, fig. 20.—*MILLET, Moll. Maine et Loire, 1813, p. 74.—*BRARD, Hist. Coq., 1815, p. 229.—*CUVIER, Règne Animal, II, 1817, p. 473.—*LAMARCK, An. sans Vert., VI, 1819, p. 76.—*DESHAYES, Enc. Méthod., II, 1827, p. 151, pl. CCXLVIII, fig. 2.—*MICHAUD, Comp. Hist. Moll. Fr., 1831, p. 110.—*LEA, Obs., I, 1834, p. 201.—*GOURMIL, Hist. Moll. Sarthe, 1835, p. 83.—*PHILIPPI, Moll. Sic., 1836, p. 66.—*ROSSMASSLER, Icon., V, 1837, p. 21, pl. XXIV, fig. 340.—*TERVER, Cat. Moll. Terr. et Flav., 1839, p. 39.—*SOWERBY, Conch. Man., 1839, fig. 145.—*GRAS, Moll. Isère., 1840, p. 72, pl. v, fig. 20.—*HANLEY, Test. Moll., 1842, p. 201; *Biv. Shells, 1843, p. 201, pl. XXI, fig. 13.—*CATLOW and REEVE, Conch. Nom., 1845, p. 60.—*PUTON, Moll. Vosges., 1847, p. 74.—*GASSIES, Moll. Agenais, 1849, p. 197.—*DUPUY, Hist. Moll. Fr., 1852, p. 632, pl. XXIII, fig. 8; XXIV, figs. 5, 6, 8.—*ROSSMASSLER, Icon., III, 1854, p. 37, pl. LXIX, fig. 850.—*DROUET, Nay. Fr., II, 1857, p. 66, pl. III, figs. 1, 2.—*H. and A. ADAMS, Gén. Rec. Moll., II, 1857, p. 493.—*MOUSSON, Coq. Terr. Fluv. Pal., 1861, p. 64.—*REEVE, Conch. Icon., XVI, 1865, pl. XXII, fig. 98.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 52.—*P.ETEL, Conch. Sam., III, 1890, p. 157.
- * *Unio littoralis* var. *minor* ROSSMASSLER (part), Icon., XI, 1842, p. 14, pl. LV, figs. 743, ? 747.²
- * *Unio littoralis* var. *acarranicus* KOBELT, Icon., IV, 1879, p. 40, pl. CLXI, fig. 1638.
- * *Unio littoralis* var. *piantensis* KOBELT, Icon., VI, 1888, p. 43, pl. CLXIII, fig. 1648.
- * *Margarita (Unio) littoralis* LEA, Syn., 1836, p. 32; 1838, p. 22.
- * *Margaron (Unio) littoralis* LEA, Syn., 1852, p. 34; 1870, p. 54.
- * *Unio littoralis* var. *umbonatus* ROSSMASSLER, Icon., XII, 1844, p. 27, pl. LVI, fig. 754.
- * ? *Unio granosus* SCHUMACHER, Ess. Nouv. Syst., 1817, pl. II, fig. 1.³
- * *Unio brevis* LAMARCK, An. sans Vert., VI, 1819, p. 73.—* LEA, Obs., I, 1834, p. 200.⁴
- * *Unio nana* LAMARCK, An. sans Vert. VI, 1819, p. 76.—* LEA, Obs., I, 1834, p. 202.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 83.
- * *Unio rubens* MENKE, Syn., 1830, p. 149.—* ROSSMASSLER, Icon., V and VI, 1837, p. 56, pl. XXIX, fig. 412.
- * *Unio subtrigona* MICHAUD, Comp. Hist. Moll. Fr., 1831, p. 111.—* GRAS, Moll. Isère., 1840, p. 21.
- * *Unio subtrigonus* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* DUPUY, Hist. Moll. Fr., 1852, p. 634, pl. XXIV, fig. 7.

¹ Lamarck refers this in the Animaux sans Vertébrés first to his *Système An. sans Vertébrés*, published in 1801, and thirdly to Draparnaud (*Hist. Moll. Fr.*, 1806). Draparnaud published this species under the name *Unio littoralis* without a figure in the *Tableau Mollusques de France*, 1801, which appeared, according to Moquin-Tandon, about July 1 of that year. I do not know which has priority, but Lamarck refers to a characteristic figure in the *Encyclopédie Méthodique* (1797), thus fixing the species without a doubt, and as he is most generally considered its author I shall credit it to him.

² Fig. 743 is a small *U. littoralis*. I think 747 is *U. batavus*.

³ Only a partial inside view is given of two valves. Schumacher says it very much resembles *U. corrugatus* Retzius, but is larger, and the teeth are very different. I believe it is a *U. littoralis*.

⁴ Lamarck refers this species to the Isle of France. Sganzin states (*Mém. Soc. Hist. Nat. Strasb.*, 1840-46, p. 8) that the amateurs he consulted in that island assured him that no such mollusk was found there. According to Lea it is *Unio littoralis*.

- * *Unio incurvus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 97, pl. XIII, fig. 27; * Obs., I, 1834, p. 107, pl. XIII, fig. 27.—* CHENU, Ill. Conch., 1858, pl. XI, figs. 1, 1a, 1b.
- Unio draparnaldi* DESHAYES, Desc. Coq. Terr., 1831, p. 38, pl. XIV, fig. 6.—¹ PETEL, Conch. Sam., III, 1890, p. 151.
- * *Unio pianensis* FARINES, Ann. des Sci. Nat., II, 1834, p. 118.—* MÖLLER, Syn. Nov. Gen., 1836, p. 197.—* DUPUY, Hist. Moll. Fr., 1852, p. 635, pl. XXIV, fig. 4.
- * *Unio bigorrensis* MILLET, Guer. Mag., 1843, p. 3, pl. LXIV, fig. 1.
- * *Unio bigorrensis* LOCARD, Coq. de France., 1893, p. 153.
- * *Unio fellmani* DESHAYES, Hist. Nat. Moll. Alg., 1848, pl. CVIII, figs. 8, 9.—* KUSTER, Conch. Cab., 1856, p. 151, pl. XLIV, fig. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* PETEL, Conch. Sam., III, 1890, p. 152.
- * *Margaron (Unio) fellmani* LEA, Syn., 1870, p. 46.
- * *Unio barvandi* DUPUY, Hist. Moll. Fr., 1852, p. 635, pl. XXV, fig. 1.
- * *Unio astierianus* DUPUY, Hist. Moll. Fr., 1852, p. 636, pl. XXIII, fig. 9.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVI, fig. 461.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 53.—* LOCARD, Coq. de France., 1893, p. 153.
- * *Unio cuneatus* ROSSMASSLER, Icon., XIII and XIV, 1854, p. 37, pl. LXIX, fig. 851.—* PETEL, Conch. Sam., III, 1890, p. 149.
- * *Unio rhomboideus* MOQUIN-TANDON, Moll. Terr. Fluv. Fr., II, 1855, p. 568, pl. XLVIII, figs. 4, 9; XLIX, figs. 1, 2.—* BOURGUIGNAT, Mal. Alg., 1864, II, p. 284, pl. XVIII; Rev. et Mag., XVIII, 1866, p. 11.—* LOCARD, Coq. de France., 1893, p. 152, fig. 165.
- * *Unio rothi* BOURGUIGNAT, Moll. Nouv., 1863, p. 41, pl. XX, figs. 1-6; Rev. et Mag., XVII, 1865, p. 337, pl. XVI.—* KOBELT, Icon., VI, 1879, p. 40, pl. CLXI, fig. 1639.—* PETEL, Conch. Sam., III, 1890, p. 166.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 59.
- * *Unio umbonatus* BOURGUIGNAT, Moll. Nouv., 1863, p. 42, pls. XXI, XXII; * Rev. et Mag., XVII, 1865, p. 339, pls. XVII, XVIII.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 54.
- * *Unio subreniformis* BOURGUIGNAT, Moll. Nouv., 1863, p. 43; Rev. et Mag., XVII, 1865, p. 340.—* KOBELT, Icon., IV, 1876, p. 64, pl. CXVIII, fig. 1151.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 54.
- * *Unio ater* REEVE, Conch. Icon., XVI, 1865, pl. XXI, fig. 19.
- * *Unio crassus* REEVE, Conch. Icon., XVI, 1865, pl. XXII, fig. 98.
- * *Unio valentinus* SOWERBY, Conch. Icon., XVI, 1866, pl. XLI, fig. 225.
- * *Unio mauritanicus* BOURGUIGNAT, Moll. Nouv., 1868, p. 317, pl. XLV, figs. 1-5.—* KOBELT, Icon., II, new ser., 1886, p. 5, pl. XXXIII, fig. 230.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 56.
- * *Unio ksibianus* MOUSSON, Mal. Blatt., XXI, 1873, p. 156; * Jahrb. Deuts. Mal. Ges., I, 1874, p. 104, pl. v, fig. 6.—* KOBELT, Icon., IV, new ser., 1876, p. 65, pl. CXIX, fig. 1153.—* PETEL, Conch. Sam., III, 1890, p. 156.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890.
- * *Unio jolyi* KOBELT, Icon., new ser., II, 1886, p. 22, pl. XLI, fig. 256.—* PETEL, Conch. Sam., III, 1890, p. 156.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 57.
- * *Unio maccarthyanus* KOBELT, Icon., new ser., II, 1886, p. 5, pl. XXXIII, fig. 229.
- * *Unio letourneauxi* KOBELT, Icon., new ser., II, 1886, p. 3, pl. XXXII, fig. 227.
- * *Unio lycicus* ROLLE, Icon., 1st sup., 1895, p. 18, pl. II, fig. 1.

Southern Europe; Asia Minor; Assyria; Morocco; Algiers.

† UNIO DELESSERTI Bourguignat.¹

Unio delesserti BOURGUIGNAT, Voy. Mer. Mort., 1852, p. 77; Test. Nov. Saul., 1852, p. 29; * Cat. Rais. Moll., 1853, p. 77, pl. III, figs. 7-9.—* ROTH, Mal. Blatt., 1856,

¹ Approaches *U. littoralis*, and may be only a variety of that, but is not so elongated.

II, p. 57.—* PÆTEL, Conch. Sam., III, 1890, p. 150.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 172.

* *Margaron (Unio) delesserti* LEA, Syn., 1870, p. 46.

Syria.

† UNIO SEMIRUGATUS Lamarck.¹

* *Unio semirugatus* LAMARCK, An. san. Vert., VI. 1819, p. 76.—* DELESSERT, Rec. Coq. Lam., 1841, pl. XII, figs. 6, 6a, 6b.

*† *Unio emesaensis* LEA, Pr. Acad. N. Sci. Phila., VIII, p. 286; Jl. Acad. N. Sci. Phila., VI, 1868, p. 254, pl. XXX, fig. 68; * Obs., XII, 1869, p. 14, pl. XXX, fig. 63.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 61.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

* *Margaron (Unio) emesaensis* LEA, Syn., 1870, p. 57.

†* *Uniosimonis* TRISTRAM, Pr. Zool. Soc. Lond., 1865, Pt. 2, p. 544.²—* LOCARD, Arch. Mus. Lyon, III, 1883, p. 239, pl. XX, figs. 1-3.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 60.—* PÆTEL, Conch. Sam., III, 1890, p. 167.—* KOBELT, Icon., new ser., VI, 1893, p. 91, pl. CLXXVI, fig. 1121; first supp., 1895, p. 18, pl. III, figs. 1-3.

* *Unio luyesi* LOCARD, Arch. Mus. Lyon, III, 1883, p. 205.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

* *Unio galilæi* LOCARD, Arch. Mus. Lyon, III, 1883, p. 206, pl. XX, figs. 10-12.³—WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 60.—* KOBELT, Icon., 1st supp., 1895, p. 20, pl. VII, figs. 4, 5.

* *Unio timius* LOCARD, Arch. Mus. Lyon, III, 1883, p. 207, pl. XX, figs. 13, 14.—* PÆTEL, Conch. Sam., III, 1890, p. 169.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 61.

* *Unio rhomboidopsis* LOCARD, Arch. Mus. Lyon, III, 1883, p. 239, pl. XX, figs. 7-9.—* PÆTEL, Conch. Sam., III, 1890, p. 165.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 61.

* *Unio rollei* KOBELT, Icon., 1st supp., 1895, p. 20, pl. IV, figs. 1, 2.

* *Unio trachæ* KOBELT, Icon., 1st supp., 1895, p. 21, pl. VIIA, fig. 2.⁴

* *Unio wagneri* KOBELT, Icon., 1st supp., 1895, p. 22, pl. VII, figs. 1, 2.

Asia Minor.

† UNIO HOMSENSIS Lea.

* *Unio homsensis* LEA, Proc. Acad. Nat. Sci. Phila., VIII, 1864, p. 285; Jl. Acad. Nat. Sci. Phila., VI, 1868, p. 249, pl. XXIX, figs. 63; * Obs., XII, 1869, p. 9, pl. XXIX, fig. 63.—* LOCARD, Arch. Mus. Lyon., III, 1883, p. 275.—* PÆTEL, Conch. Sam., III, 1890, p. 155.—* WESTERLUND, Faun. Pal. II, Pt. 7, 1890, p. 62.

* *Margaron (Unio) homsensis* LEA, Syn. 1870, p. 31.

Syria.

¹ Lea and Ferussac both referred this to *littoralis* after examining Lamarck's shells. I have seen specimens in the collection of the former from the Jardin des Plantes, Paris, which are, no doubt, authentic, and which, I think, perhaps, are specifically different from *littoralis*, being much shorter and more rounded.

²A specimen of this in the U. S. National Museum collection, from Tristram, labeled as above, is the same as Lea's examples of *semirugatus*.

³Published as *Unio maris-galilæi* in list in Comptes Rendus, XCI, p. 502; but not described.

⁴Some of these approach very close to specimens of *U. littoralis*, especially to the form called *U. rothi* Bourguignat, which I have placed in that species, and I should not be surprised if there was an absolute connection between the short, rounded forms of *U. semirugatus* and the longer rhomboid *U. littoralis*.

UNIO PSEUDONYMUS Simpson.¹

* *Unio hueti* KOBELT, Icon., new ser., II, 1886, p. 22, pl. XI, fig. 225.

Euphrates and Tigris rivers.

† UNIO FERUSSACIANUS Lea.

* *Unio ferussacianus* LEA, Jl. Acad. Nat. Sci. Phila., VI, 1868, p. 255. Footnote to description of *U. emesaensis*.—* LEA, Obs., XII, 1869, p. 15. Footnote.

* *Margaron (Unio) ferussacianus* LEA, Syn., 1870, p. 46.

Bagdad.

UNIO EPISCOPALIS Tristram.

* *Unio episcopalis* TRISTRAM,² Pr. Zool. Soc. Lond., 1865, p. 544.—VON MARTENS Vorderas. Conch., 1874, p. 68.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 62.—* PÆTEL, Conch. Sam., III, 1890, p. 151.—* KOBETT, Icon., VI, new ser., 1893, p. 89, pl. CLXXV, fig. 1119.

Orontes River, Syria.

† UNIO DURIEUI Deshayes.

* *Unio duricui* DESHAYES, Hist. Nat. Moll. Alg. Atlas, 1847, pl. CIX, figs. 5-8.³—BOURGUIGNAT, Mol. Alg., II, 1864, p. 288, pl. XIX, figs. 4-8.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 142.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

* *Margaron (Unio) durieui* LEA, Syn., 1852, p. 39; 1870, p. 48.

* *Unio sitifensis* MORELET, Jl. de Conch., II, 1851, p. 360.

Unio orientalis BOURGUIGNAT, Test. Noviss., 1852, p. 29.

* *Unio bruguierianus* BOURGUIGNAT, Cat. Rais., 1853, p. 78, pl. II, figs. 54-58.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 172.—* PÆTEL, C. Sam., III, 1890, p. 146.

* *Margaron (Unio) bruguierianus* LEA, Syn., 1870, p. 44.

* *Unio vescoi* BOURGUIGNAT, Rev. et Mag., VIII, 1856, p. 74, pl. II, figs. 4-8.⁴—* KOBELT, Icon., IV, 1876, p. 66, pl. CXIX, fig. 1154.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 152.—* PÆTEL, Conch. Sam., III, 1890, p. 171.

* *Margaron (Unio) vescoi* LEA, Syn., 1870, p. 46.

* *Unio schwerzenbachi* BOURGUIGNAT, Rev. et Mag., VIII, 1856, p. 75, pl. VIII, figs. 1-5.—* KUSTER, Conch. Cab. Unio, 1862, p. 266, pl. XC, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XLIV, fig. 241.—* KOBELT, Icon., VII, 1880, p. 81, pl. CCVI, fig. 2099.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 95.

* *Unio prusii* BOURGUIGNAT, Rev. et Mag., VIII, 1856, p. 76, pl. III, figs. 1-4.

* *Unio damascensis* LEA,⁵ Proc. Acad. Nat. Sci. Phila., VII, 1863, p. 190; Jl. Acad.

¹ Kobelt supposed this to be the *U. hueti* of Bourguignat, and figured and described it for that species. It is, however, very distinct, and I therefore am compelled to give it a new name.

² Tristram only describes this in Latin, and does not figure it; and as I have never seen an authentic specimen I am obliged to depend on Kobelt's figure, which agrees well with Tristram's diagnosis. These make it a large, somewhat sulcate form, shaped like *batarus*, but more compressed and having a beautiful purple nacre.

³ There does not seem to be any essential difference between the Algerian and the Syrian shells, which I have referred to this species.

⁴ Bourguignat says he has received numerous specimens of this from Drouet, labeled *U. turcius* Parreyss, but Parreyss's names were only in manuscript, and were never published, and have therefore no scientific standing.

⁵ Only the type is in the Lea collection, and it appears to be a delicate specimen of *duricui*.

Nat. Sci. Phila., VI, 1866, p. 55, pl. XVIII, p. 52; * Obs., XI, 1869, p. 59, pl. XVIII, fig. 52.—* PÆTEL, Conch. Sam., III, 1890, p. 150.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 178.

* *Margaron (Unio) damascensis* LEA, Syn., 1870, p. 52.

* *Unio orontesensis* LEA, Pr. Acad. Nat. Sci. Phila., VII, 1863, p. 190; Jl. Acad. Nat. Sci. Phila., VI, 1866, p. 53, pl. XVIII, fig. 50; * Obs., XI, p. 57, pl. XVIII, fig. 50.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 178.—PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Margaron (Unio) orontesensis* LEA, Syn., 1870, p. 52.

* *Unio orphaensis* LEA, Pr. Acad. Nat. Sci. Phila., VIII, 1864, p. 285; Jl. Acad. Nat. Sci. Phila., VI, 1868, p. 250, pl. XXIX, fig. 64; * Obs., XII, 1869, p. 10, pl. XXIX, fig. 64.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 179.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Margaron (Unio) orphaensis* LEA, Syn., 1870, p. 52.

* *Unio mardinensis* LEA, Pr. Acad. Nat. Sci. Phila., VIII, 1864, p. 286; * Jl. Acad. Nat. Sci. Phila., VI, 1868, p. 252, pl. XXX, fig. 66; * Obs., XII, 1869, p. 12, pl. XXX, fig. 66.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 179.

* *Margaron (Unio) mardinensis* LEA, Syn., 1870, p. 53.

† UNIO DURIEUI var. KULLETHENSIS Lea.

* *Unio kullethensis* LEA, Pr. Acad. Nat. Sci. Phila., VIII, 1864, p. 285; * Jl. Acad. Nat. Sci. Phila., VI, 1868, p. 251, pl. XXIX, fig. 65; * Obs., XII, 1869, p. 11, pl. XXIX, fig. 65.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 178.

* *Margaron (Unio) kullethensis* LEA, Syn., 1870, p. 53.

* *Unio kallethensis* and *kullinthensis* PÆTEL, Conch. Sam., III, 1890, p. 156.

Algiers; Tunis; Asia Minor, Assyria; Southeastern Europe.

† UNIO SYRIACUS Lea.¹

* *Unio syriacus* LEA, Pr. Acad. Nat. Sci. Phila., VII, 1863, p. 189; * Jl. Acad. Nat. Sci. Phila., VI, 1866, p. 56, pl. XIX, fig. 53; * Obs., XI, 1867, p. 60, pl. XIX, fig. 53.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 178.—* PÆTEL, Conch. Sam., III, 1890, p. 169.

* *Margaron (Unio) syriacus* LEA, Syn., 1870, p. 35.

Orontes River, Syria.

UNIO BYTHINICUS Kobelt.

* *Unio bythinicus* KOBELT, Icon. new ser., VI, 1893, p. 96, pl. CLXXIX, fig. 1128.

Asia Minor.

UNIO MICELII Kobelt.

* *Unio micelii* KOBELT, Nach. Mal. Ges., XVI, 1884, p. 182; Icon., new ser., II, 1886, p. 24, pl. XLIII, figs. 260, 261.

Tunis.

† UNIO EMARGINATUS Lea.²

* *Unio emarginatus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 62, pl. IX, fig. 22; * Obs., I, 1834, p. 174, pl. IX, fig. 22.

¹ I have only seen a single, badly eroded specimen, the type, but it seems to differ from anything I know. It probably belongs to this group.

² Lea's only shell, the type, consists of two opposite, odd valves. One of these may possibly be an elongated *batavus*, the other appears distinct.

* *Margarita (Unio) emarginatus* LEA, Syn., 1838, p. 26.

* *Margaron (Unio) emarginatus* LEA, Syn., 1852, p. 39; 1870, p. 62.

Locality unknown. I have no doubt that it belongs somewhere in the circummediterranean region.

† UNIO CARNEUS Kuster.

* *Unio carneus* KUSTER, Conch. Cab. Unio, 1854, p. 103, pl. XXVIII, figs. 1, 2.—

* DROUET, Mem. Acad. Dijon, VII, 1882, p. 23, reprint.—* P.ÉTEL, Conch. Sam., III, 1890, p. 147.

* *Unio gontierii* BOURGUIGNAT, Rev. et Mag., IX, 1857, p. 16, pl. IV, figs. 1-4.—

* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 67.

* *Margaron (Unio) gontieri* LEA, Syn. 1870, p. 44.

* *Unio penchinatus* BOURGUIGNAT, Moll. Pen. Con., 1863, p. 44, pl. XXV; * Rev. et Mag., XVII, 1865, p. 342, pl. XXI.—* KOBELT, Icon., IV, 1876, p. 66, pl. CNIX, fig. 1155.

Southern Europe.

† UNIO BATAVUS Maton and Rackett.

* ? *Mya pictorum* GMELIN, Syst. Nat., 13th ed., 1788, p. 3218¹.—* DONOVAN, Brit. Shells, V, 1803, pl. CLXXIV.—* CHENU, Bib. Conch., 1st ser., I, 1815, p. 114, pl. XLVII, figs. 8, 9.

* *Unio pictorum* DRAPARNAUD, part, Hist. Moll. Fr., 1806, p. 131, pl. XI, figs. 1-4.²

* *Mya batava* MATON and RACKETT,³ Tr. Linn. Soc. Lond., VIII, 1807, p. 37.—

* WOOD, Gen. Conch., I, 1815, p. 303, pl. XIX, figs. 1, 2.—* DILLWYN, Cat., I, 1817, p. 49.—* TURTON, Conch. Dict., 1819, p. 105.—* WOOD, Ind. Test., 1825, p. 12, pl. II, fig. 25b; rev. ed., 1856, p. 15, pl. II, fig. 25.

* *Unio batava* LAMARCK, An. sans Vert., VI, 1819, p. 78.⁴—C. PFEIFFER, Nat. L. and Suss. Moll., Pt. 1, 1821, p. 119, pl. V, fig. 14.—* DESHAYES, Enc. Méth., II, 1827, p. 151, pl. CCXLVIII, fig. 3; II, 1830, p. 584, pl. CCXLVIII, fig. 3.—

* WAARDENBERG, Com. Hist. Nat. An., 1827, p. 36.—* KLEEGER, Moll. Bor., 1828, p. 38.—* FLEMING, Hist. Brit. Moll., 1828, p. 416.—* STARK, Nat. Hist., II, 1828, p. 90.—* MICHAUD, Comp. Hist. Moll. Fr., 1831, p. 109.—* GOUPIL, Hist. Moll. Sarthe, 1835, p. 83.—* FOURNEL, Faun. Moselle, I, 1836, p. 487.—* GRAS, Moll. Isere., 1840, p. 21.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* PUTON, Moll. Vosges, 1847, p. 75.

* *Mya batava* TURTON, Conch. Ins. Brit., 1822, p. 244.—* TURTON, Man. L. and F. W. Shells, Brit. Is., 1831, pl. XX, fig. 10.

* *Unio batavus* NILSSON, Hist. Moll. Svec., 1822, p. 112.—* ROSSMASSLER, Icon., II, 1835, p. 20, pl. VIII, figs. 128, 128a, 128b; III, 1836, pp. 28, 32, pl. XIV, fig. 205; XV, fig. 214; V and VI, 1837, p. 56, pl. XXIX, fig. 414; XI, 1842, p. 14, pl. LV, fig. 745.—* BROWN, L. and F. W. Conch., 1836, p. 111, pl. XVIII, figs. 6-8; XXI, figs. 10, 11.—* ?ANTON, Verz. der Conch., 1839, p. 15.—* HANLEY, Test. Moll., 1842,

¹ Gmelin refers to the figure by Chemnitz in Conchylien Cabinet, VI. This figure is probably a light-colored *batavus*.

² Fig. 3 is certainly *batavus*, and figs. 1 and 2 may be.

³ Maton and Rackett do not figure their species, but refer to several figures. The first is in Ginanni, Opere Postume, 1755, pl. IV, fig. 17. This work I have not seen. The figures referred to in Schröter's Flussconchylien, in Chemnitz, and the Encyclopédie Methodique (pl. 248, fig. 3) are what we understand as *U. batavus*, the latter being the one cited by Lamarck.

⁴ Lamarck does not figure his species, but refers to plate CCXLVIII, fig. 3, in the Encyclopédie Méthodique, which is the species commonly known as *Unio batavus*.

- p. 195; * Biv. Shells, 1843, p. 195.—*SCHOLZ, Schleis. L. and W. Moll., 1843, p. 133.—*BROWN, Ill. Recent Conch., 1844, p. 82, pl. xxxi, figs. 3, 3a, 4-6.—*DUPUY, Hist. Moll. Fr., 1852, p. 638, pl. xxv, figs. 14, 15.—*KUSTER, Conch. Cab. Unio, 1854, p. 121, pl. xxxiii, figs. 4-7; xxxiv, figs. 1, 2.—*VON WAILL, Suss. Biv. Liv., 1855, p. 96.—*MOQUIN-TANDON, Moll. Terr. et Fluv. Fr., II, 1855, p. 571, pl. xlix, figs. 7, 8.—*DROUET, Nay. Fr., II, 1857, p. 79, pl. vi, fig. 1.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*BIELZ, Faun. Sieben, 1863, p. 193.—*BOURGUIGNAT, Mal. Alg., 1864, p. 286, pl. xix, fig. 9; xx, figs. 1-4.—*SOWERBY, Conch. Icon., XVI, 1866, pl. xlii, fig. 234.—*BROT, Coq. Faun. Nay. Leman, 1867, p. 49, pl. ix, figs. 1-5.—*KOBELT, Faun. Nass. Moll., 1871, p. 242.—*LEHMAN, Die Schnecken, 1873, p. 295.—*CLESSIN, Deutsche Ex. Moll., 1876, p. 463, fig. 302.—*DROUET, Mem. Acad. Dijon, VII, 1882, p. 18 (reprint).—*CLESSIN, Moll. Oest., 1887, p. 741.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 74.—*PÆTEL, Conch. Sam., III, 1890, p. 145.—*LOCARD, Coq. de Fr., 1893, p. 167.—*WESTERLUND, Act. Soc. F. and F. F., XIII, No. 7, 1897, p. 162.
- **Margarita (Unio) batavus* LEA, Syn., 1836, p. 26; 1838, p. 20.
- **Margaron (Unio) batavus* LEA, Syn., 1852, p. 30; 1870, p. 47.
- **Unio batavi* var. ? ROSSMASSLER, Icon., V and VI, 1837, p. 55, pl. xxix, fig. 410.
- **Unio nanus* DUPUY, Hist. Moll. Fr., 1852, p. 640, pl. xxv, fig. 16.—*KOBELT, Icon., VI, 1879, p. 42, pl. clxii, figs. 1642, 1643.—*LOCARD, Coq. de Fr., 1893, p. 154, fig. 168.
- **Unio riparia* C. PFEIFFER, Nat. L. and S. Moll., 1821, Pt. 1, p. 118, pl. v, fig. 13.—*KLEBERG, Moll. Bor., 1828, p. 39.
- **Unio riparius* SCHOLZ, Schleis. L. and W. Moll., 1843, p. 129.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 90.—*LOCARD, Coq. de Fr., 1893, p. 164.
- *? *Unio crassus* NILSSON,¹ Hist. Moll. Svec., 1822, p. 108.—*ROSSMASSLER, Icon., II, 1835, p. 19, pl. viii, figs. 126, 127²; V and VI, 1837, p. 55, pl. xxxix, fig. 411.—MOQUIN-TANDON, Moll. Terr. and Fluv. Fr., II, 1855, p. 570; III, pl. xlix, figs. 3, 4.—*NORDENSKIÖLD and NYLANDER, Fin. Moll., 1856, p. 84, pl. vi, fig. 72.—*DROUET, Nay. Fr., II, 1857, p. 76, pl. iv, fig. 2.—*WESTERLUND, Faun. Sw., N. and Den., 1873, p. 569.3.—*LOCARD, Coq. de Fr., 1893, p. 162, fig. 176.—*WESTERLUND, Acta Soc. F. and F. F., XIII, No. 7, 1897, p. 159.
- **Unio crassus* variety *batavus* JORDAN, Jahrb. Deuts. Mal. Ges., VI, 1879, p. 307.
- **Unio ater* NILSSON, Hist. Moll. Svec., 1822, p. 107.—*ROSSMASSLER, Icon., II, 1835, p. 23, pl. ix, fig. 133; VII and VIII, 1838, p. 41, XL, fig. 543.—*SCHOLZ, Schleis. L. and F. W. Moll., 1843, p. 131.—*KUSTER, Conch. Cab. Unio, 1854, p. 114, pl. xxi, figs. 1, 2, 6; xxxii, figs. 1-4.—*MOQUIN-TANDON, Moll. Terr. et Fluv. Fr., II, 1855, p. 570, pl. xlix, figs. 5, 6.—*DROUET, Nay. Fr., II, 1857, p. 72, pl. iv, fig. 1; Mem. Acad. Dijon, VII, 1882, p. 14 (reprint).—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 65.—*PÆTEL, Conch. Sam., III, 1890, p. 145.—*LOCARD, Coq. de Fr., 1893, p. 176.—*WESTERLUND, Act. Soc. F. and F. F., XIII, No. 7, 1897, p. 160.
- **Unio atra* DESHAYES, Encyc. Meth., II, 1830, p. 582.
- Unio rugatus* MENKE, Syn. Moll., 1828, p. 90.—*ROSSMASSLER, Icon., V and VI, 1837, p. 65, pl. xxix, fig. 415.
- **Unio elongata* MICHAUD, Comp. Hist. Moll. Fr., 1831, p. 113, pl. xvi, fig. 29.
- **Unio labacensis* ROSSMASSLER, Icon., II, 1835, p. 21.

¹Nilsson claims that this is Retzius' species, and that it is found in the rivers of Sweden. His Latin description is so brief that nothing can be made out of it. I have no doubt that he had *Unio batavus* before him when he wrote it.

²I believe that this is a rather large *batavus* and the figures are almost exactly like those he gives of that species.

³Claimed to be *crassus* of Retzius, but he gives *U. ater* of Nilsson as one variety and *batavus* Lamarek as another.

- * *Unio decurratus* ROSSMASSLER, Icon., II, 1835, p. 22, pl. IX, fig. 131; V and VI, 1837, p. 21, pl. XXIV, fig. 339.—* KUSTER, Conch. Cab. Unio, 1856, p. 108, pl. XXX, figs. 1, 2.—* MUGRAVE, Phot. Conch., 1863, pl. II, fig. 4.—* SOWERBY, Conch. Icon., XVI, 1867, pl. LVI, fig. 284.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 92.—* PÆTEL, Conch. Sam., III, 1890, p. 150.
- * *Unio reniformis* ROSSMASSLER, Icon., III, 1836, p. 31, pl. XV, fig. 213.—* SCHOLZ, Schleis, L. and W. Moll., 1843, p. 132.—* KUSTER, Conch. Cab. Unio, 1854, p. 110, pl. XXX, figs. 3, 4.—* BOURGUIGNAT, Rev. et Mag., 1865, pl. XXIII, figs. 4-6.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXII, fig. 371.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 80.—* LOCARD, Coq. de Fr., 1893, p. 161.
- * *Unio carinthiacus* ROSSMASSLER, Icon., III, 1836, p. 30, pl. XV, fig. 209.¹—* SOWERBY, Conch. Icon., XVI, 1856, pl. XXX, fig. 157; * PÆTEL, Conch. Sam., III, 1890, p. 147.—* LOCARD, Coq. de Fr., 1893, p. 167.
- * *Unio amnicus* ROSSMASSLER, Icon., III, 1836, p. 31, pl. XV, fig. 212.—* KUSTER, Conch. Cab. Unio, 1856, p. 99, pl. XXVII, fig. 2.—* DROUET, Mem. Acad. Dijon, VIII, 1882, p. 22 (reprint).—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 89.—* LOCARD, Coq. de Fr., 1893, p. 163, fig. 177.
- * *Unio piscinalis* ROSSMASSLER, Icon., III, 1836, p. 30, pl. XV, fig. 210.—* LOCARD, Coq. de Fr., 1893, p. 160.
- * *Unio atrovirens* ROSSMASSLER, Icon., III, 1836, p. 28, pl. XV, figs. 206, 207.—* SCHMIDT, Bull. Imp. Soc. N. H. Moscow, 1840, p.
- * *Unio consentaneus* ROSSMASSLER, Icon., III, 1836, p. 29, pl. XV, fig. 208; VII, 1838, p. 25, pl. XXXV, fig. 491; p. 42, XL, fig. 544; XI, 1842, p. 14, pl. LV, fig. 742.
- * *Unio fusculus* ROSSMASSLER, Icon., III, 1836, p. 30, pl. XV, fig. 211.—* PÆTEL, Conch. Sam., III, 1890, p. 153.—* LOCARD, Coq. de Fr., 1893, p. 159, fig. 173.
- * *Unio?* ROSSMASSLER, Icon., III, 1836, p. 27, pl. XIV, fig. 201.
- * *Unio sterenianus* DROUET, Unionidæ Russ., 1881, p. 14;² Jl. de Conch., XXIX, 1881, p. 24; Supp. Un. Serbie, 1884, pls. I, II.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 66.—* KOBELT (part), Icon., new ser., II, 1886, p. 38, pl. LI, fig. 281.³
- * *Unio bandini* KUSTER, Icon., V and VI, 1837, p. 22, pl. XXIV, fig. 341.
- * *Unio glaucinus* PORRO, Mal. Como., 1838, p. 115.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* STABILE, Faun. Ing., 1845, p. 61, pl. III, fig. 75.—* KOBELT, Icon., new ser., II, 1886, p. 19, pl. XI, fig. 251.—* PÆTEL, Conch. Sam., III, 1890, p. 154.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 90.
- * *Unio gargottæ* ROSSMASSLER, Icon., VII, 1838, p. 26, pl. XXXV, fig. 493.—* MONTEROSATO, Nat. Sic., new ser., 1896, p. 6, fig. 1.
- ? *Unio corrugata* MANDUYT, Moll., Vienna, 1839, p. 8.
- ? *Unio rotundata* MANDUYT, Moll., Vienna, 1839, p. 9.⁴
- ? *Unio gangrenosus* SCHMIDT, Bull. Imp. N. H. Moscow, 1840, p. —.—* KUSTER, Conch. Cab. Unio., 1854, p. 124, pl. XXXIV, figs. 3, 4.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 81.—* PÆTEL, Conch. Sam., III, 1890, p. 153.
- Unio pruinosis* SCHMIDT, Bull. Soc. Nat. Mosc., 1840, p. 445.—* DROUET, Jl. de Conch., XXIX, 1881, p. 248; * Mem. Acad. Dijon, VII, 1882, p. 21 (reprint).—* PÆTEL, Conch. Sam., III, 1890, p. 164.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 82.—* LOCARD, Coq. de Fr., 1893, p. 160.
- * *Unio littoralis* var. *minor* ROSSMASSLER (part), Icon., XI, 1842, p. 14, pl. LV, fig. 744.

¹ Rossmassler credits this and several other species to Zeigler. So far as I am able to learn these are mere manuscript names. Neither Moquin Tandon or Locard give Zeigler as an author in their bibliographies of writers on French mollusks, and Lea refers only to his manuscript. I have not seen his Systematisches Verzeichniss.

² Generally credited to Krinické, who published the naked name in Bull. Nat. Mosc., 1837.

³ Fig. 280, said to be this, is probably *U. tumidus*.

⁴ Both these species of Manduyt are *batavus?* according to Moquin Tandon.

- * *Unio mauca* MILLET, Guer. Mag., 1843, p. 4, pl. LXIV, fig. 2.
- * *Unio maucus* DUPUY, Hist. M. Fr., 1852, p. 642, pl. XXVI, fig. 17.—*LOCARD, Coq. Fr., 1893, p. 156, fig. 170.
- * *Unio moquinianus* DUPUY, Moll. Gers., 1843, p. 80, pl.—*ROSSMASSLER, Icon., XII, 1844, p. 31, pl. LIX, fig. 769.—*KUSTER, Conch. Cab. Unio, 1854, p. 100, pl. XXVII, figs. 3, 4, 5.—*DUPUY, Hist. Moll. Fr., 1852, p. 644, pl. XXVI, fig. 18.—*MOQUIN-TANDON, Moll. Terr. and Fluv. Fr., 1855, p. 573, pl. I, figs. 1, 2.—*DUPUY, Rev. et Mag., VIII, 1856, p. 75.—*DROUET, Nay. Fr., II, 1857, p. 88, pl. VI, fig. 3.—*PÆTEL, Conch. Sam., III, 1890, p. 159.—*WESTERLUND, Fann. Pal., II, Pt. 7, 1890, p. 146.—*LOCARD, Coq. de Fr., 1893, p. 179.
- * *Unio requieni* ROSSMASSLER, Icon., XII, 1844, p. 29, pl. LVII, figs. 757-761.—*KUSTER (part), Conch. Cab. Unio, 1856, p. 126, pl. XXXVI, fig. 2.¹—*MOQUIN-TANDON, Moll. Terr. et Fluv. Fr., II, 1855, p. 574, pl. I, figs. 5-7.—*KOBELT, Icon., VI, 1879, p. 43, pl. CLXIII, fig. 1647.—*LOCARD, Coq. de Fr., 1893, p. 190.
- * *Unio saudri* ROSSMASSLER, Icon., XII, 1844, p. 26, pl. LVI, figs. 748-750.—*KUSTER, Conch. Cab. Unio, 1856, p. 101, pl. XXVII, figs. 6, 7.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIX, fig. 413.—*PÆTEL, Conch. Sam., III, 1890, p. 166.—*WESTERLUND, Fann. Pal., II, Pt. 7, 1890, p. 93.
- ? *Unio alcroni* COMPANYO and MASSOT, Bull. Soc. Agr. Sc. Pyr. d'Or, VI, Pt. 2, 1845, p. 234, fig. 2.—*KOBELT, Icon., IV, 1876, p. 64, pl. CXVIII, fig. 1150.²
- Unio badins* KOKEIL, Mich. Comp., 1831, pl. XVI, fig. 36.—*CATLOW and REEVE, Conch. Nom., 1845, p. 56.—*KOKEIL, Gallenst. Kärnt. Conch., 1852, p. ?.³
- *? *Unio volrichii* MORELET, Moll. Port., 1845, p. 105, pl. XIII, fig. 1.
- Unio jacquemini* DUPUY, Cat. Ext. Gall. Test., 1849, No. 328.—*DUPUY, Hist. M. Fr., 1852, p. 643, pl. XXV, fig. 17.—*KOBELT, Icon., VI, 1879, p. 42, pl. CLXII, fig. 1641.—*WESTERLUND, Fann. Pal., II, Pt. 7, 1890, p. 128.—*LOCARD, Coq. Fr., 1893, p. 188.
- Unio drometi* DUPUY, Cat. Ext. Gal. Test., 1849, No. 327; Hist. Moll. Fr., 1852, p. 639, pl. XXV, fig. 14.
- Unio philippi* DUPUY, Cat. Ext. Gal. Test., 1849, No. 335; Hist. Moll. Fr., 1852, p. 654, pl. XXVIII, fig. 19.
- * *Unio ovalis* DUPUY, Hist. Moll. Fr., 1852, p. 637, pl. XXV, fig. 13.
- * *Unio spinellii* VILLA, in Moll. Bres., 1852, p. 50, fig. D.
- * *Unio moulinsiana* DUPUY, Hist. Moll. Fr., VI, 1852, p. 640, pl. XXIV, fig. 10.—*MOQUIN-TANDON, Hist. Moll. Fr., II, 1855, p. 572.—*LOCARD, Coq. de Fr., 1893, p. 152.
- *? *Unio petterianus* KUSTER, Conch. Cab., 1854, p. 97, pl. XXVII, fig. 4.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*PÆTEL, C. Sam., III, 1890, p. 163.—*WESTERLUND, Fann. Pal., II, Pt. 7, 1890, p. 115.
- * *Margaron (Unio) petterianus* LEA, Syn., 1852, p. 31, 1870, p. 49.
- * *Unio pigliolo* KUSTER (part), Conch. Cab. Unio, 1854, p. 125, pl. XXXIV, figs. 5, 6.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXV, fig. 392.—*DUPUY, Hist. Moll. Fr., 1852, p. 645, pl. XXVI, fig. 19.
- * *Unio heldi* KUSTER, Conch. Cab. Unio, 1854, p. 111, pl. XXX, figs. 5-7.—*DROUET, Jl. de Conch., XXIX, 1881, p. 26.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 80.
- * *Unio luxurians* KUSTER, Conch. Cab. Unio, 1854, p. 119, pl. XXXII, fig. 5.—*WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 94.—*PÆTEL, C. Sam., III, 1890, p. 158.

¹ Kuster gives several figures of what he supposes are *requieni*. Part of them are *turtoui*, and one is possibly *elongatulus* Pfeiffer.

² It is hard to tell just what this is. It is probably a delicate, rather compressed *batavus*.

³ According to Lea and Westerlund this = *batavus*.

* *Unio brevirostris* KUSTER, Conch. Cab. Unio, 1854, p. 120, pl. XXXIII, figs. 1-3.—

* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 96.

* *Unio natolicus* KUSTER, Conch. Cab. Unio, 1856, p. 144, pl. XLII, fig. 4.

* *Unio turcicus* KUSTER, Conch. Cab. Unio, 1862, p. 267, pl. XC, figs. 3, 4.

* *Unio merdiger* REEVE, Conch. Icon., XVI, 1865, pl. XXVIII, fig. 145.¹

Europe; Asia Minor; Northwest Africa.

Section LAPIDOSUS Simpson, 1900.

(Type, *Unio lapidosus* Kobelt.²)

Shell inequilateral, subtriangular ovate, large, solid, slightly inflated, with a double posterior ridge, biangulate behind; beaks high, sculpture not seen; disks strongly, irregularly sulcate; epidermis olive; three thick pseudocardinals in the right valve and two in the left; laterals strong; anterior muscle scars deep, tripartite; posterior scars deep, distinct; nacre rosy white, iridescent behind. Animal unknown.

UNIO LAPIDOSUS Kobelt.

* *Unio lapidosus* KOBELT, Nach Deuts. Mal. Ges., 1893, p. 151; * Icon., new ed., VI, 1893, p. 90, pl. CLXXVI, fig. 1120.—* DROUET, Jl. de Conch., XLIII, 1895, p. 32.

Euphrates River.

Section ELLIPTIO Rafinesque, 1819.

(Type, *Unio crassidens* Lamarek.)

Shell elongated, rhomboid or oval, usually more or less biangulate behind; beak sculpture consisting of a few rather strong ridges, which are nearly parallel to the growth lines or slightly doubly looped; the surface smooth or feebly corrugated.

(Group of *Unio coloratus*.)

Shell somewhat rhomboid, solid, slightly biangulate behind, rather compressed, more or less sculptured with concentric sulcations; beaks moderately prominent, their sculpture unknown; epidermis brownish; teeth strong; laterals rather club-shaped; nacre white or purple. Animal unknown.

?† UNIO COLORATUS Charpentier.

* *Unio coloratus* CHARPENTIER, in Kuster, Conch. Cab. Unio, 1856, p. 155, pl. XLIV, fig. 6.—* PÄTEL, Conch. Sam., III, 1890, p. 148.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 599.

Medellin River, Vera Cruz, Mexico.

¹ Credited to Jay's Catalogue, in which it is not described, but is credited to Waltz.

² This striking shell differs very much from any of the circummediterranean forms and may be entitled to subgeneric rank.

UNIO CALLOSUS Lea.

- * *Unio callosus* LEA, Pr. Am. Phil. Soc. II, 1841, p. 31; *Tr. Am. Phil. Soc., VIII, 1842, p. 239, pl. XXIII, fig. 54; *Obs., III, 1842, p. 77, pl. XXIII, fig. 54.—
 *CONRAD, Pr. Acad. Nat. Sci. Phila., VI, 1853, p. 246.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—*CHENU, Ill. Conch., 1858, pl. XXVI, figs. 4, 4a, 4b.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 147.
 *Margaron (*Unio*) *callosus* LEA, Syn., 1852, p. 33; 1870, p. 33.

Said to come from the Ohio Canal below Columbus. I am sure that the locality is wrong, and it appears to be a Mexican form. The type is not in the Lea collection.

UNIO MEXICANUS Philippi.

- * *Unio mexicanus* PHILIPPI, Zeits. für Mal., IV, 1847, p. 95.—*PHILIPPI, Abbild. und Beschr., III, 1849, p. 110, pl. VI, fig. 3.¹—*KUSTER, Conch. Cab. Unio, 1862, p. 285, pl. XCV, fig. 7.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 159.—*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 593.
 *Margaron (*Unio*) *mexicanus* LEA, Syn., 1870, p. 53.

Mexico.

(Group of *Unio semigranosus*.)

Shell triangular rhomboid, solid, inflated, with a distinct posterior ridge; beaks rather full, their sculpture consisting of numerous somewhat irregular corrugations which pass into the pustulous sculpture of the shell; surface of the valves generally more or less sculptured with chevron-shaped or zigzag ridges or corrugations, which often break into pustules, the posterior slope bearing curved, radiating plications which are likewise sometimes nodulous; epidermis dark, scarcely rayed; pseudocardinals strong, radial, ragged; laterals heavy, obliquely striated; beak cavities not deep, compressed; muscle scars well impressed; nacre purple, with bronzy or coppery shades.

Animal apparently not different from that of other related Unios. I have not seen any with the marsupium filled.

† UNIO PLEXUS Conrad.

- * *Margarita* (*Unio*) *carbonarius* LEA, Syn., 1836, p. 192; 1838, p. 17.²
 * *Unio carbonarius* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 37, pl. XL, fig. 32; *Obs., II, 1838, p. 37, pl. XI, fig. 32.—*TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 236.—
 *HANLEY, Test. Moll., 1842, p. 184; *Biv. Shells, 1843, p. 184, pl. XXII, fig. 10.—*CATLOW and REEVE, Conch. Nom., 1845, p. 57.—*CHENU, Ill. Conch., 1858, pl. XXIII, figs. 1, 1a, 1b.—*PÆTEL, Conch. Sam., III, 1890, p. 147.
 * *Margarita* (*Unio*) *pliciferus* LEA, Syn., 1836, p. 13; 1838, p. 14.
 * *Unio pliciferus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 61, pl. XVII, fig. 53; *Obs., II, 1838, p. 61, pl. XVII, fig. 53.—*TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p.

¹ A miserable figure of some rhomboidal, rather compressed form, probably grouping with *callosus*.

² The names *carbonarius* and *pliciferus* were published two years before *Unio plexus*, but were not accompanied by a description. Part X of the Monography appeared in May, 1838, and the transactions containing Dr. Lea's description of *carbonarius* June 15, according to Scudder. Dr. Lea places his *carbonarius* in the synonymy of *pliciferus*, though the former comes first in his paper.

256.—* HANLEY, Test. Moll., 1842, p. 176; * Biv. Shells, 1843, p. 176, pl. xx, fig. 32.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* KUSTER, Conch. Cab., 1848, p. 142, pl. xlii, fig. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. xxii, figs. 5, 5a, 5b; Manual, 1859, II, p. 142, fig. 702.—* SOWERBY, Conch. Icon., XVI, 1868, pl. lxxv, fig. 387.—* B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., II, Pt. 7, 1894, p. 580, pl. lxxv, fig. 2; lxx, figs. 2, 2a.

Margaron (Unio) pliciferus LEA, Syn., 1852, p. 20; 1870, p. 31.

* *Unio plexus* CONRAD, Monog., X, 1838, p. 89, pl. xlix, figs. 1, 2; * Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 255.

Vera Cruz, Mexico.

† UNIO CROCODILARUM Morelet.

* *Unio crocodilarum* MORELET, Test. Nov., I, 1849, p. 28.—* REEVE, Conch. Icon., XVI, 1864, pl. x, fig. 37.—* PÆTEL, Conch. Sam., III, 1890, p. 149.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 577, pl. lx, fig. 3; lxxvii, fig. 3.

* ? *Unio rusticus* SOWERBY, Conch. Icon., XVI, 1868, pl. lxiv, fig. 324.

Rio Usumacinta, Guatemala.

UNIO MORINI Morelet.

* *Unio morini* MORELET, Test. Nov., II, 1851, p. 24.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 576, pl. lx, fig. 2; lxxvii, fig. 4.

Rio Usumacinta, Guatemala.

UNIO SEMIGRANOSUS von dem Busch.

* *Unio semigranosus* VON DEM BUSCH (in Philippi), Abbild. und Besch., I, 1845, p. 19, pl. i, figs. 1-3.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 257.—* HANLEY, Biv. Shells, 1846, p. 381, pl. xx, fig. 33.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* KUSTER, Conch. Cab. Unio, 1861, p. 252, pl. lxxxv, fig. 1.—* B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 569.

* *Margaron (Unio) semigranosus* LEA, Syn., 1852, p. 20; 1870, p. 34.

* *Unio carbonarius* var. *semigranosus* PÆTEL, Conch. Sam., III, 1890, p. 147.

Mexico, Vera Cruz to Tampico.

† UNIO CORIUM Reeve.¹

* *Unio corium* REEVE, Conch. Icon., XVI, 1864, pl. x, fig. 39.—* PÆTEL, Conch. Sam., III, 1890, p. 149.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 570.

State of Chiapas, Mexico.

UNIO TESTUDINEUS Morelet.

* *Unio testudineus* MORELET, Test. Nov., I, 1849 p. 28.²—FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 571, pl. lxii, fig. 3; lxx, fig. 3.

* *Unio semigranosus* REEVE, Conch. Icon., XVI, 1864, pl. x, fig. 36.

Rio Usumacinta, Guatemala.

¹ Probably a mere variety of *semigranosus*. It is quite likely that when a sufficient amount of material is examined this and some other species of this group will have to be placed in the synonymy.

² Fischer and Crosse think that *testudineus* is perhaps only a local race of *semigranosus*. I think it is quite likely they are right.

† UNIO PSORICUS Morelet.

* *Unio psoricus* MORELET, Test. Nov., Pt. 2, 1851, p. 25.¹—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 572, pl. LXI, fig. 2.

Rio Usamacinta, Guatemala.

(Group of *Unio gibbosus*.)

Shell solid, compressed or inflated, triangular ovate to elongate ovate, arcuate when old, gibbous above, pointed or slightly biangulate behind, with a moderate posterior ridge; beaks low, the sculpture consisting of a few very strong ridges which run parallel with the growth lines; epidermis dull and clothlike; pseudocardinals strong, rough; laterals granular or vertically striated, club-shaped; beak cavities very shallow; one or more slight furrows or ridges occur in the cavity of the shell which run nearly parallel with the laterals; muscle scars very deep and distinct.

Marsupium occupying the entire outer gills; branchiae large, curved below, inner the wider except at the posterior end, free from the abdominal sac only part of their length; mantle very thin, with thickened edges; branchial opening generally large.

† UNIO GIBBOSUS Barnes.

* *Unio nasuta* LAMARCK,² An. sans Vert., VI, 1819, p. 75.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 538; 3d ed., II, 1839, p. 669.

* *Unio nasutus* AGASSIZ, Arch. für Nat., I, 1852, p. 50.

* *Unio gibbosus* BARNES, Am. Jl. Sci., VI, 1823, p. 262, pl. XI, fig. 12.³—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 286.—SHORT and EATON, Transylvania Jl., 1831, p. 78.—* HANLEY, Test. Moll., 1842, p. 207; Biv. Shells, 1843, p. 207, pl. XX, fig. 51.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* KUSTER, Conch. Cab. Unio, 1852, p. 28, pl. IV, figs. 3, 4.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* SOWERBY, Conch. Icon, XVI, 1868, pl. LXXIII, fig. 379.—* CALKINS, Pr. Ottawa Ac. Sci., 1874, p. 42.—* LATCHFORD, Tr. Ottawa F. N. Club, 1882, p. 50.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 153.—* BAKER, Moll. Chicago, 1898, p. 70, pl. XIV, figs. 3, 4; XV, figs. 1-4.

Margarita Unio gibbosus LEA, Syn., 1836, p. 38; 1838, p. 25.

* *Margaron (Unio) gibbosus* LEA, Syn. 1852, p. 38; 1870, p. 61.

* *Mya gibbosa* EATON, Zool. Text-Book, 1826, p. 220.

* *Unio mucronatus* BARNES, Am. Jl. Sci., VI, 1823, p. 266, pl. XIII, fig. 13 (outline).—* AGASSIZ, Arch. für Naturg., I, 1852, p. 50.

* *Mya mucronata* EATON, Zool. Text-Book, 1826, p. 221.

¹ Probably another variety of *semigranosus*.

² This name was given by Lamarck to the species under consideration before that of Barnes, but Say had already used the name *nasutus* for a *Unio*, hence *gibbosus* must be used.

³ In 1820 Rafinesque bestowed the name *Amblesma gibbosa* on some unionoid mollusk which I can not determine. Since then Conrad in 1836, and others after him, have used the specific name as of Rafinesque and applied it to a *Unio*, which they have figured and described. Barnes applied the name in 1823, properly characterizing it, and it must stand, because at that time it had not been used in the genus *Unio*.

- * *Unio dilalatus* SAY,¹ Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 68.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* CONRAD, Monog., V, 1836, p. 42, pl. XXI.—* KUSTER, Conch. Cab. Unio, 1852, p. 38, pl. VI, fig. 4.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.
- * *Unio torulosus* FERUSSAC, Guer. Mag., 1835, p. 28.
- * *Unio arctatus* FERUSSAC, Guer. Mag., 1835, p. 29.
- *† *Unio arctior* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 10, pl. IV, fig. 10;² Obs., II, 1838, p. 10, pl. IV, fig. 10.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 234.—* HANLEY, Test. Moll., 1842, p. 208; * Biv. Shells, 1843, p. 208, pl. XXII, fig. 46.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* KUSTER, Conch. Cab. Unio, 1861, p. 179, pl. LVI, fig. 6.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* CHENU, Ill. Conch., 1858, pl. XXI, figs. 2, 2a, 2b.—* ? SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXV, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.
- * *Margarita (Unio) arctior* LEA Syn., 1836, p. 39; 1838, p. 25.
- * *Margaron (Unio) arctior* LEA, Syn., 1852, p. 38; 1870, p. 71.

† UNIO GIBBOSUS var. ARCUS Conrad.

- * *Unio arcus* CONRAD, Am. Jl. Sci., XXV, 1834, p. 340, pl. I, fig. 8; * New F. W. Shells, 1834, p. 67.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen., 1836, p. 1898.—* HANLEY, Test. Moll., 1842, p. 207; * Biv. Shells, 1843, p. 207, pl. XXIII, fig. 46.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.
- * *Margarita (Unio) arcus* LEA, Syn., 1836, p. 38; 1838, p. 25.
- * *Margaron (Unio) arcus* LEA, Syn., 1852, p. 38; 1870, p. 61.

† UNIO GIBBOSUS var. SUBGIBBOSUS Lea.

- * *Unio subgibbosus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 169; Jl. Ac. N. Sci. Phila., IV, 1858, p. 53, pl. VI, fig. 36; Obs., VI, 1858, p. 53, pl. VI, fig. 36.—* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 432, pl. LXXVIII, fig. 5.
- * *Margaron (Unio) subgibbosus* LEA, Syn., 1870, p. 61.
- * *Unio lazarus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXVIII, fig. 348.
- *† *Unio rufus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 171; Jl. Ac. N. Sci. Phila., IV, 1858, p. 85, pl. XVII, fig. 65; Obs., VI, 1858, p. 85, pl. XVII, fig. 65.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) rufus* LEA, Syn., 1870, p. 61.

† UNIO GIBBOSUS var. ARMATHWAITENSIS B. H. Wright.

- * *Unio gibbosus* var. *armathwaitensis* B. H. WRIGHT, Naut., XI, 1898, p. 123.³

† UNIO GIBBOSUS var. DELICATUS Simpson.⁴

Entire Mississippi drainage; St. Lawrence and its tributaries; Alabama River system; southeast into Florida; southwest to the Guadalupe River, Texas.

¹ As of Rafinesque. I am unable to make out Rafinesque's species.

² Merely a form with white nacre.

³ These three are dwarf varieties of *gibbosus*. All are often more or less humped, and there are many intermediate and connecting specimens.

⁴ A greatly compressed, thin variety, with an evenly elliptical outline and very dark nacre may bear this name.

† UNIO STONENSIS Lea.

* *Unio stonensis* LEA, Pr. Am. Phil. Soc., I, 1840, p. 286; * Tr. Am. Phil. Soc., VIII, 1841, p. 195, pl. VIII, fig. 5; Obs., III, 1842, p. 33, pl. VIII, fig. 5.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XXIX, figs. 7, 7a, 7b.—* SOWERBY, Conch. Icon., XVII, 1868, pl. LXXXVI, fig. 462.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

* *Margaron (Unio) stonensis* LEA, Syn., 1852; 1870, p. 43.

Stone River, Tennessee.

† UNIO LURIDUS Lea.

* *Unio luridus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 273, pl. XX, fig. 29; Obs., V, 1852, p. 29, pl. XX, fig. 29.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251; * H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

* *Margaron (Unio) luridus* LEA, Syn., 1852, p. 30; 1870, p. 48.

Yadkin River, North Carolina; Georgia; Florida.

(Group of *Unio discus*.)

Shell large, solid, compressed, subtriangular; biangulate behind; beak sculpture not seen; beaks high, but not swollen; epidermis dark, rough; teeth heavy; laterals remote, club-shaped; nacre white, yellow or purple.

Animal unknown.

† UNIO DISCUS Lea.

* *Unio discus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 74, pl. XVIII, fig. 57; * Obs., II, 1838, p. 74, pl. XVIII, fig. 51.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 237.—* HANLEY, Test. Moll., 1842, p. 197; * Biv. Shells, 1843, p. 197, pl. XXII, fig. 12.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. XIX, figs. 6, 6a, 6b.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXII, fig. 310.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 151.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 567.

* *Margaron (Unio) discus* LEA, Syn., 1852, p. 31; 1870, p. 50.

* *Unio panacoensis*, VON DEM BUSCH in Philippi, Abbild. und Besch., I, 1843, p. 75, pl. II.—* KUSTER, Conch. Cab. Unio, 1861, pl. LXXXI, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Unio mexicanus* SOWERBY, Conch. Icon., XVI, 1867, pl. LV, fig. 281.

Mexico; Central America.

(Group of *Unio sloatianus*.)

Shell rhomboid, solid, inflated, with a well-defined posterior ridge and a smaller, fainter one above, the posterior end being distinctly biangulate, nearly straight or slightly incurved on the base, rounded in front; beaks only slightly elevated; beak sculpture not observed; surface of the valves covered with somewhat radiately plicate, nodulous corrugations which become less pronounced in front; posterior

slope radially plicate; epidermis dark; pseudocardinals stumpy, radial, granularly striate; laterals double in the left valve, single in the right, with a faint, secondary ridge below; anterior muscle scars deep, nearly smooth; cavity of the beaks moderate, slightly compressed.

Animal unknown.

† UNIO SLOATIANUS Lea.

* *Unio sloatianus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 287; Tr. Am. Phil. Soc.,

VIII, 1842, p. 217, pl. xvi, fig. 33; Obs., III, 1842, p. 55, pl. xvi, fig. 33.—

* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch. 1858, pl. xxxii, figs. 7, 7a, 7b.—* KUSTER, Conch. Cab., 1862, p. 286, pl. xcvi, fig. 3.—

* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 167.

* Margaron (*Unio*) *sloatianus* LEA, Syn., 1852, p. 21; 1870, p. 31.

* *Plectomerus sloatianus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 261.

* *Unio atromarginatus* LEA,¹ Pr. Am. Phil. Soc., I, 1840, p. 288; Tr. Am. Phil. Soc., VIII, 1842, p. 207, pl. xiii, fig. 21; Obs., III, 1842, p. 45, pl. xiii, fig. 21.—

* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 488.—* CHENU, Ill. Conch., 1858, pl. xxviii, figs. 6, 6a, 6b.—* KUSTER, Conch. Cab. Unio, 1861, p. 250, pl. lxxxiv, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam.,

III, 1890, p. 145.

* Margaron (*Unio*) *atromarginatus* LEA, Syn., 1852, p. 19; 1870, p. 29.

* *Plectomerus atromarginatus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 260.

* *Unio aratus* CONRAD, Ann. and Mag., IV, 1849, p. 302; * Pr. Ac. N. Sci. Phila. IV, 1849, p. 154.

* *Unio plectophorus* CONRAD, Jl. Ac. N. Sci. Phila., I, 1850, p. 277, pl. xxxviii, fig. 7.

* *Plectomerus plectophorus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 2.

Chattahoochee and Flint rivers, Georgia.

(Group of *Unio crassidens*.)

Shell generally solid, rather inflated, rhomboid oval, with a sharp, well-developed posterior ridge; beaks prominent, the sculpture consisting of a few coarse ridges running nearly parallel with the growth lines and swollen at the posterior ridge; surface of the valves nearly smooth or only marked by growth lines; posterior slope bearing wrinkled, corrugate, radiate ridges; epidermis often faintly rayed.

Animal having the branchiæ very large, rounded below, inner the larger, free nearly or quite their entire length from the abdominal sac; branchial and anal openings large, with many small papillæ; marsupium occupying the entire outer gills.

† UNIO CRASSIDENS Lamarck.

* *Unio crassidens* LAMARCK, An. sans Vert., VI, 1819, p. 71.—* LEA, Obs., I, 1834,

p. 199.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 532; 3d ed., II, 1839,

p. 667.—* HANLEY, Test. Moll., 1842, p. 184; * Biv. Shells, 1843, p. 184, pl. xx,

fig. 46.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* H. and A. ADAMS,

¹Lea's type of *sloatianus* is an old, arcuate shell, while the type of *atromarginatus* is young and quite different looking. But the young plicate *atromarginatus* gradually changes into the smoother *sloatianus* as it grows older, and specimens of both show the dark bordered nacre, though it is more pronounced in the young shells.

- Gen. Rec. Moll., II, 1857, p. 495.—* CALKINS, Pr. Ot. Ac. Sci., 1874, p. 42.—
 * B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 149.—
 * CALL, Tr. Ac. Sci., St. Louis, VII, 1895, p. 12, pl. II.
 * *Margarita (Unio) crassidens* LEA, Syn., 1836, p. 19; 1838, p. 17.
 * *Margaron (Unio) crassidens* LEA, Syn., 1852, p. 24; 1870, p. 37.
 * ? *Unio nigra* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 291, pl. LXXX, figs. 1-4.
 * *Unio niger* SAY, Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 70.—
 * FERUSSAC, Guer. Mag., 1835, p. 27.—* CONRAD, Monog., VI, 1836, p. 49, pl. XXVI.—* KUSTER, Conch. Cab. Unio, 1852, p. 25, pl. IV, fig. 1.—* AGASSIZ, Arch. für Naturg., I, 1852, p. 50.—* CONRAD, Pr. Acad. N. Sci., Phila., VI, 1853, p. 253.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVIII, fig. 408.
 * ? *Unio cuneatus* BARNES, Am. Jl. Sci., VI, 1823, p. 263.—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 279, fig. 3.—* SHORT and EATON, Transylvania Jl., 1831, p. 78.
 * *Mya cuneata* EATON, Zool. Text Book, 1826, p. 220.
 * ? *Unio discus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXII, fig. 310.
 * *Unio crassus* SOWERBY, Conch. Icon., XVI, 1868, pl. xcv, fig. 520.

Mississippi drainage generally; Alabama and Tombigbee rivers; southeast to the Chattahoochee River. In the southeastern part of its range this species seems to merge into *U. incrassatus*.

† UNIO INCRASSATUS Lea.¹

- * *Unio incrassatus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 286; * Tr. Am. Phil. Soc., VIII, 1840, p. 217, pl. XVI, fig. 34; * Obs., III, 1842, p. 55, pl. XVI, fig. 34.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XXX, figs. 5, 5a, 5b.—* KUSTER, Conch. Cab. Unio, 1861, p. 192, pl. LX, fig. 5.—* REEVE, Conch. Icon., XVI, 1865, pl. xxvi, fig. 127.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
 * *Margaron (Unio) incrassatus* LEA, Syn., 1852, p. 24; 1870, p. 37.

Coosa River system; Chattahoochee and rivers of north Florida; east to Savannah?

† UNIO POLYMORPHUS B. H. Wright.

- * *Unio polymorphus* B. H. WRIGHT, Naut., XIII, 1899, p. 42.

Spanish Creek, Charlton County, Georgia.

† UNIO FORBESIANUS Lea.

- * *Unio forbesianus* LEA, Pr. Am. Phil. Soc., V, 1852, p. 251; * Tr. Am. Phil. Soc., X, 1852, p. 264, pl. XVI, fig. 17; * Obs., V, 1852, p. 20, pl. XVI, fig. 17.—* CONRAD, Pr. Acad. Nat. Sci. Phila., VI, 1853, p. 249.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* KUSTER, Conch. Cab. Unio, 1861, p. 310, pl. LXIX, fig. 6.—* MUSGRAVE, Phot. Conch., 1863, pl. II, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXI, fig. 361.—* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 410, pl. XLIX, fig. 3; L, figs. 2, 3.
 * *Margaron (Unio) forbesianus* LEA, Syn., 1852, p. 24; 1870, p. 37.
 * ? *Unio moussonianus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 268, pl. XVIII, fig. 22; * Obs., V, 1852, p. 24, pl. XVIII, fig. 22.—* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 252.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 160.
 * *Margaron (Unio) moussonianus* LEA, Syn., 1852, p. 29; 1870, p. 46.

¹ I am doubtful whether this is more than a small, rhomboid, southeastern variety of *crassidens*, bearing the same relation to it that *U. subgibbosus* does to *U. gibbosus*.

*† *Unio corvus* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 112; *Jl. Acad. N. Sci., Phila., IV, 1859, p. 217, pl. XXVII, fig. 97; Obs., VII, 1859, p. 35, pl. XXVII, fig. 97.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVIII, fig. 411.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 149.

* *Margaron (Unio) corvus* LEA, Syn., 1870, p. 37.

* *Unio vestitus* LEA, Pr. Ac. N. Sci., Phila., V, 1861, p. 393; *Jl. Ac. N. Sci., Phila., V, 1862, p. 189, pl. XXV, fig. 259; * Obs., IX, 1863, p. 11, pl. XXV, fig. 259.—* B. H. WRIGHT, Check List, 1888.

Margaron (Unio) vestitus LEA, Syn., 1870, p. 50.

Streams from the Congaree River, South Carolina, south to northern Florida.

† UNIO DANIELSII B. H. Wright.

* *Unio danielsii* B. H. WRIGHT, Naut., XIII, 1899, p. 31.

Spring Creek, Decatur County, Georgia.

† UNIO PUSILLUS Lea.¹

* *Unio pusillus* LEA, Pr. Ac. N. Sci. Phila., I, 1840, p. 286; * Tr. Am. Phil. Soc., VIII, 1842, p. 220, pl. XVIII, fig. 36; * Obs., III, 1842, p. 58, pl. XVIII, fig. 36.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 256.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. XXXII, figs. 5, 5a, 5b.—* KUSTER, Conch. Cab., 1861, p. 197, pl. LXIII, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 411, pl. LI, figs. 2, 6.

* *Margaron (Unio) pusillus* LEA, Syn., 1852, p. 31; 1870, p. 48.

*† *Unio buzeus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 261, pl. XV, fig. 13; * Obs., V, 1852, p. 17, pl. XV, fig. 13.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 147.

* *Margaron (Unio) buzeus* LEA, Syn., 1852, p. 29; 1870, p. 46.

*† *Unio anthonyi* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 41; *Jl. Ac. N. Sci. Phila., V, 1862, p. 197, pl. XXVII, fig. 266; * Obs., IX, 1863, p. 19, pl. XXVII, fig. 266.—* B. H. WRIGHT, Check List, 1888.

Margaron (Unio) anthonyi, LEA, Syn., 1870, p. 43.

Abbeville district, South Carolina; south to northern Florida.

† UNIO MERUS Lea.

* *Unio merus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 260, pl. XV, fig. 10; * Obs., V, 1852, p. 16, pl. XV, fig. 10.—* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 252.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 159.

* *Margaron (Unio) merus* LEA, Syn., 1870, p. 54.

*† *Unio castus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 306; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 349, pl. LVII, fig. 174; * Obs., VIII, 1860, p. 31, pl. LVII, fig. 174.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) castus* LEA, Syn., 1870, p. 40.

South Carolina.

¹ The type is a young shell identical with one of Lea's *U. anthonyi*. The type of the latter is a little larger, lighter colored, and less rayed.

UNIO MASONI Conrad.¹

* *Unio masoui* CONRAD, New F. W. Shells, 1834, p. 34, pl. v, fig. 2, p. 70.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* CONRAD, Monog., III, 1836, p. 28, pl. XII, fig. 2.—* MÖLLER, Syn. Nov. Gen., 1836, p. 204.—* HANLEY, Test. Moll., 1842, p. 201; * Bi. Shells, 1843, p. 201, pl. XXIII, fig. 18.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 18, pl. II, fig. 1.—* KUSTER, Conch. Cab. Unio, 1852, p. 34, pl. v, fig. 6.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

* *Margarita (Unio) masoni* LEA, Syn., 1836, p. 33; 1838, p. 23.

* *Margarou (Unio) masoni* LEA, Syn., 1852, p. 34; 1870, p. 55.

Savannah River, Georgia.

† UNIO FRATERNUS Lea.

* *Unio fraternus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 263, pl. XVI, fig. 15; * Obs., V, 1852, p. 19, pl. XVI, fig. 15.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* KUSTER, Conch. Cab. Unio, 1861, p. 201, pl. LXVII, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 153.

* *Margaron (Unio) fraternus* LEA, Syn., 1852, p. 32; 1870, p. 51.

Abbeville district, South Carolina, to Columbus, Georgia.

† UNIO CONGARÆUS Lea.

* *Unio congaræus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 72, pl. VI, fig. 4; Obs., I, 1834, p. 82, pl. VI, fig. 4.—* CONRAD, New F. W. Shells, 1834, p. 68.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* CONRAD, Monog., III, 1836, p. 27, pl. XII, fig. 1.—* HANLEY, Test. Moll., 1842, p. 200; * Biv. Shells, 1843, p. 200, pl. XXII, fig. 31.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* CHENU, Ill. Conch., 1858, pl. III, figs. 5, 5a, 5b.—* SOWERBY, Conch. Icon., XVI, 1867, pl. LIX, fig. 296.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Margarita (Unio) congaræus* LEA, Syn., 1836, p. 32; 1838, p. 22.

* *Margaron (Unio) congaræus* LEA, Syn., 1852, p. 33; 1870, p. 54.

* † *Unio fulvus* LEA,² Tr. Am. Phil. Soc., V, p. 96, pl. XIII, fig. 39; * Obs., I, 1834, p. 208, pl. XIII, fig. 39.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 200; * Biv. Shells, 1843, p. 200, pl. XXIII, fig. 32.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1858, pl. LXXXIX, fig. 483.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 153.

* *Margarita (Unio) fulvus* LEA, Syn., 1836, p. 32; 1838, p. 22.

* *Margaron (Unio) fulvus* LEA, Syn., 1852, p. 33; 1870, p. 54.

Cape Fear River, North Carolina; south to Savannah, Georgia.

¹ I have never seen this species, but should judge from the figure that it groups with *U. crassidens*.

² The type is a young shell, and is not the same as adult shells which are in the Lea collection under that name.

† UNIO DARIENSIS Lea.

* *Unio dariensis* LEA, Tr. Am. Phil. Soc., VIII, 1842, p. 246, pl. XXVI, fig. 61; * Pr. Am. Phil. Soc., II, 1843, p. 225; * Obs., III, 1842, p. 84, pl. XXVI, fig. 61.—* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 248.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* CHENU, Ill. Conch., 1858, pl. XXXIII, figs. 6, 6a, 6b.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCI, fig. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 150.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 413, pl. LIII, fig. 2; LIV, fig. 1.

* *Margaron (Unio) dariensis* LEA, Syn., 1852, p. 33; 1870, p. 54.

Southeast Georgia to north Florida.

† UNIO MONROENSIS Lea.

Unio monroensis LEA,¹ Desc. of 12 sp. of Uniones, 1843, no pagination; * Tr. Am. Phil. Soc., IX, 1845?, p. 279, pl. XLI, fig. 8; * Obs., IV, 1848, p. 37, pl. XLI, fig. 8.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 159.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 410, pl. LI, fig. 1.

* *Margaron (Unio) monroensis* LEA, Syn., 1852, p. 29; 1870, p. 46.

Florida.

† UNIO HARTWRIGHTI² B. H. Wright.

† *Unio hartwrighti* B. H. WRIGHT, Naut., IX, 1896, p. 121, pl. II, figs. 4–6.

Lake Beresford, Florida.

• UNIO HINKLEYI B. H. Wright.

* *Unio hinkleyi* B. H. WRIGHT, Pr. Ac. N. Sci. Phila., 1888, p. 117, pl. IV, fig. 3³;

* Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 423, pl. LXV, fig. 4.

Florida.

¹A puzzling form. The type, the only shell in the Lea collection, is rather solid, and the posterior ridge is curved slightly upward in the middle. Other specimens in the museum are less solid, the posterior ridge is sharper, and straight or curved down in the middle, and there seems to be almost a connection with the very different *U. hartwrighti*.

²This name was applied to another species by Newcomb, and was used by Wright in the Check List, by Pætel and others, but the form to which it was given was believed to be a synonym of something else, and it was never described. Mr. Wright has again used the name for what is probably a valid species.

³The figure of this shell is not at all accurate, being considerably more blunt posteriorly and inflated below than the specimen sent as the type. It approaches *U. monroensis*, but is more elongated, and a series of specimens sent by Mr. Wright almost connects it with *U. dorei*.

† UNIO WEBSTERI B. H. Wright.

* *Unio websteri* B. H. WRIGHT, Pr. Ac. N. Sci. Phila., 1888, p. 113, pl. II, fig. 2¹;

* Check List, 1888.

Florida.

† UNIO DORSATUS Lea.

* *Unio dorsatus* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 160; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 300, pl. XLV, fig. 112; * Obs., XII, 1869, p. 60, pl. XLV, fig. 112.—

* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 411, pl. LI, fig. 7; LI, figs. 1, 2.

* *Margaron (Unio) dorsatus* LEA, Syn., 1870, p. 37.

Catawba River, North Carolina; Florida.

† UNIO WACCAMAWENSIS Lea.²* *Unio waccamawensis* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 193; * Jl. Acad. N. Sci. Phila., VI, 1866, p. 16, pl. v, fig. 14; * Obs., XI, 1867, p. 20, pl. v, fig. 14.—* B. H. WRIGHT, Check List, 1888.* *Margaron (Unio) waccamawensis* LEA, Syn., 1870, p. 36.

Waccamaw Lake, North Carolina.

(Group of *Unio pigerrimus*.)

Shell rather solid, elliptic rhomboid, somewhat inflated, with a low, rounded posterior ridge, sometimes a little arcuate below, biangulate behind; beaks full, their sculpture a few coarse ridges which are curved upward and swollen where they cross the posterior ridge; surface concentrically striate and often sculptured with curved, subradiating or zigzag corrugations which have a tendency to break into nodules; epidermis brownish; hinge rather heavy; pseudocardinals stumpy, granular; laterals club shaped; muscle scars well marked.

Animal unknown.

UNIO PIGERRIMUS Crosse and Fischer.

* *Unio pigerrimus* CROSSE and FISCHER, Jl. de Conch. XLI, 1893, p. 293.—

* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 582, pl. LXV, figs. 1, 1a.

Mexico.

† UNIO MITCHELLI Simpson.

* *Unio mitchelli* SIMPSON (in Dall.), Pr. U. S. Nat. Mus., XVIII, 1896, p. 5; Pr. U. S. Nat. Mus., XIX, 1896, p. 371, pl. XXXII, figs. 1-3.³

Southern Texas to New Leon, Mexico.

¹Mr. Wright has sent to the Museum under the above name two quite different forms, neither of which is like the figure. From an examination of the first one sent, an old, worn shell, and the figure, I believed it to be a form of *U. obesus* (see Proc. U. S. Nat. Mus., XV, 1892, p. 428). Specimens sent since by Mr. Wright and said to be typical approach *U. hartwrighti*, but are more compressed and elongated. These forms are exceedingly puzzling.

²An aberrant species. The shell is inflated, thin, and has a strong posterior ridge. It has characters of both the *Buckleyi* and *Crassidens* groups.

³Close to *pigerrimus* apparently, and may be but a variety of it; it is a lighter, longer shell, is less sculptured, and does not have a rose-colored naere.

† UNIO IHERINGI B. H. Wright.

* *Unio iheringi* B. H. WRIGHT, Naut., XII, 1898, p. 93. — * SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 79, pl. IV, fig. 5.

Southern Texas.

† UNIO SPHENORHYNCHUS Fischer and Crosse.

* *Unio sphenorhynchus* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 617, pl. LXVI, figs. 2, 2a.

Mexico.

(Group of *Unio liebmanni*.)

Shell rather solid, inflated, ovate rhomboid or trapezoidal, with a well-developed posterior ridge, pointed at or near the base behind, but not biangulate, nearly straight or slightly incurved below; beaks full, sculpture not seen; surface smooth or concentrically striate; epidermis dark, scarcely rayed; hinge rather strong; pseudocardinals heavy, compressed; muscle scars distinct; nacre livid to whitish.

Animal unknown.¹

† UNIO LIEBMANNI Philippi.

* *Unio liebmanni* PHILIPPI, Zeits. für Mal., IV, 1847, p. 96; * Abbild. und Besch., III, 1849, p. 109, pl. VI, fig. 1. — * KUSTER, Conch. Cab. Unio, 1862, p. 281, pl. XCIV, fig. 7. — * PÆTEL, Conch. Sam., III, 1890, p. 157. — * FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 616.

* *Margaron (Unio liebmanni)* LEA, Syn., 1870, p. 48.

* *Unio liebmanni* B. H. WRIGHT, Check List, 1888.

Mexico.

† UNIO OPACATUS Crosse and Fischer.

* *Unio opacatus* CROSSE and FISCHER, Jl. de Conch., XLI, 1893, p. 295. — * FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 592, pl. LXVI, figs. 1, 1a.

Mexico.

(Group of *Unio buckleyi*.)

Shell oval, obovate, subtrapezoidal, or elliptical, with a more or less developed posterior ridge, often biangulate behind, compressed or inflated, thin or somewhat solid; beaks generally full, sculptured with several rather strong concentric ridges, which are either nearly parallel with the growth lines or slightly doubly looped; posterior slope often faintly wrinkled; epidermis usually smooth and shining in the younger shells, often rough when old, becoming darker with age, rayed when young. Hinge teeth solid or compressed; nacre mostly brilliant,

¹ This group is exceedingly close to that of *U. buckleyi*, according to the shell characters, and some specimens of *U. liebmanni* are so near others of *U. buckleyi* as to deceive even experts.

of many shades. Animal not differing especially from that of closely related groups.¹

† UNIO BUCKLEYI Lea.

* *Unio buckleyi*, LEA, Desc. of 12 Uniones, 1843 (no pagination); * Tr. Am. Phil. Soc., IX, 1845 ♀, p. 276, pl. XXXIX, fig. 2; * Obs., IV, 1848, p. 34, pl. XXXIX, fig. 2.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* KUSTER, Conch. Cab. Unio, 1861, p. 177, pl. LVI, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIV, fig. 175.—* B. H. WRIGHT, Check List, 1888.—* B. H. and S. H. WRIGHT, Conch. Ex., II, 1888, p. 95.—* PÆTEL, Conch. Sam., III, 1890, p. 146.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 417, pl. LVIII, figs. 6, 7; LIX, figs. 1, 2; LX, fig. 2.

* *Margarou (Unio) buckleyi* LEA, Syn., 1852, p. 30; 1870, p. 48.

* † *Unio buddianus* LEA,² Desc. of 12 Uniones, 1843; Tr. Am. Phil. Soc., IX, 1845, ♀ p. 277, pl. XL, fig. 5; * Obs., IV, 1848, p. 35, pl. XL, fig. 5.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* KUSTER, Conch. Cab. Unio, 1861, p. 249, pl. LXXXIV, fig. 1.—* REEVE, Conch. Icon., XVI, 1865, pl. XX, fig. 88.—* B. H. and S. H. WRIGHT, Conch. Ex., II, 1888, p. 95.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

* *Margarou (Unio) buddianus* LEA, Syn., 1852, p. 32; 1870, p. 52.

* † *Unio dalli* B. H. WRIGHT,³ Pr. Ac. N. Sci. Phila., 1888, p. 119, pl. VI, fig. 1.—* B. H. WRIGHT, Check List, 1888.

† UNIO BUCKLEYI var. ORCUTTII⁴ S. H. Wright.

* *Unio orcuttii* S. H. WRIGHT, West. Am. Sci., IV, 1888, p. 60, 3 figs.⁴

Florida.

† UNIO JAYENSIS Lea.

* *Unio jayensis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 28, pl. IX, fig. 23;⁵ * Obs., II, 1838, p. 28, pl. IX, fig. 23.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 235.—* HANLEY, Test. Moll., 1842, p. 206; * Biv. Shells, 1843, p. 206, pl. XXII, fig. 53.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* CHENU, Ill. Conch., 1858, pl. XXII, figs. 4, 4a, 4b.—* SOWERBY, Conch.

¹The relationship between the *Buckleyi*, *Crassidens*, *Complanatus*, *Downiei* and *Fisherianus* groups is very close. That of *Crassidens* is characterized by inflated shells with a strong posterior ridge, and wrinkles on the posterior slope; of *Buckleyi* by a less developed ridge, the absence of strong wrinkles, and a rich, warm texture of shell; of *Complanatus* by trapezoidal, not greatly inflated shells of dull texture. The species of the *Fisherianus* group have elongated, generally compressed, thin shells, pointed behind, and of the *Downiei* assemblage solid, inflated shells. But there are many species that are quite negative in characters, and others that combine the features of two or more groups, that can not be satisfactorily placed.

²The type is a large, badly eroded shell, apparently a little injured at the upper, anterior part, and is truncated a little there.

³The type is a small, old shell, which has become rough and arcuate.

⁴Credited to Manatee River, Florida, Simpson and Miakka Lake, Florida, Newcomb, by Mr. Wright. I have never found it in the Manatee River, but obtained it in abundance in Miakka Lake.

⁵Although the type is very different from that of *U. buckleyi* the two species approach so closely that there are specimens which I can not name.

Icon., XVI, 1867, pl. LX, fig. 301; LXVII, fig. 343. ?—* PÆTEL, Conch. Sam., III, 1890, p. 155.

* *Margarita (Unio) jayensis* LEA, Syn., 1836, p. 37; 1838, p. 24.

* *Margaron (Unio) jayensis* LEA, Syn., 1852, p. 37.

* † *Unio prasinatus* CONRAD, Am. Jl. Conch., II, 1866, p. 279, pl. xv, fig. 14.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) jayanus* LEA, Syn., 1870, p. 60.

* *Unio jayanus* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 419, pl. LXI, fig. 4.

* † *Unio simpsoni* B. H. WRIGHT¹; Check List, 1888.

* † *Unio marshii* B. H. WRIGHT, Pr. Ac. N. Sci. Phila., 1888, p. 118, pl. v, fig. 2; Check List, 1888.

* † *Unio tryoni* B. H. WRIGHT, Pr. Ac. N. Sci. Phila., 1888, p. 120, pl. vi, fig. 2; Check List, 1888.

Florida.

† UNIO CORUSCUS Gould.

* *Unio coruscus* GOULD, Pr. Bost. Soc. Nat. Hist., I, 1856, p. 15; * *Otia* Conch., 1862, p. 222.—* B. H. and S. H. WRIGHT, Conch. Ex., II, 1888, p. 95.—

* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 419, pl. LXIII, figs. 1, 7.

* *Margaron (Unio) coruscus* LEA, Syn., 1870, p. 48.

† UNIO CORUSCUS var. FRYANUS B. H. Wright.²

Unio fryanus B. H. WRIGHT, Pr. Acad. N. Sci. Phila., 1888, p. 113, pl. II, fig. 1; Check List, 1888.

* † *Unio diazensis* S. H. WRIGHT,³ Nant., X, 1897, p. 5.

Florida.

† UNIO CUNNINGHAMI B. H. Wright.

* *Unio cunninghami* B. H. WRIGHT, Pr. Ac. N. Sci. Phila., XIII, 1883, p. 58, pl. I, figs. 1-4.—* A. F. GRAY, Am. Nat., XVII, 1883, p. 1184.—* S. H. and B. H. WRIGHT, Conch. Ex., II, 1888, p. 105.—* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 422, pl. LXV, fig. 6.

Florida.

† UNIO MICANS Lea.

* *Unio micans* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 85; * Jl. Ac. N. Sci. Phila., V, 1862, p. 59, pl. III, fig. 207; * Obs., VIII, 1862, p. 63, pl. III, fig. 207.—* SOWERBY, Conch. Icon., XVI, 1866, pl. xxxv, fig. 182.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 159.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 423, pl. LXV, fig. 3.

* *Margaron (Unio) micans* LEA, Syn., 1870, p. 45.

* *Unio perlucens* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 193; * Jl. Ac. N. Sci. Phila.,

¹From an examination of specimens sent to the U. S. National Museum by Mr. Wright I referred this to *U. buckleyi*, in Pr. U. S. Nat. Mus., XV, p. 417. The type is now in the U. S. National Museum, and I believe it to be a rather compressed, bright young *U. jayensis*.

²A form of *coruscus* a little less solid and inflated than the type, and quite brilliant.

³This name occurs in B. H. Wright's Check List, but the species was not described until 1897. It scarcely differs from *U. fryanus*.

VI, 1866, p. 18, pl. v, fig. 16; Obs., XI, 1867, p. 22, pl. v, fig. 16.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) perlucens* LEA, Syn., 1870, p. 45.

North Carolina to South Georgia and Florida.

† UNIO PYGMÆUS Lea.¹

* *Unio pygmaeus* LEA, Pr. Am. Phil. Soc., V, 1852, p. 252; * Tr. Am. Phil. Soc., X, 1852, p. 262, pl. xv, fig. 14; * Obs., V, 1852, p. 18, pl. xv, fig. 14; * CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 256.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) pygmaeus* LEA, Syn., 1870, p. 48.

Abbeville District, South Carolina.

† UNIO LEHMANII, S. H. Wright.

* *Unio lehmanii*, S. H. WRIGHT, Naut., X, 1897, p. 138.—* SIMPSON, Pr. Acad. N. Sci. Phila., 1900, p. 80, pl. iv, fig. 9.

St. Marys River, Florida.

† UNIO BURTCHIANUS S. H. Wright.

* *Unio burtchianus* S. H. WRIGHT, Naut., X, 1897, p. 137.—* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 80, pl. iv, fig. 8.

St. Mary's River, Florida.

† UNIO CONFERTUS Lea.²

* *Unio lugubris* LEA, Tr. Am. Phil. Soc., VI, 1834, p. 30, pl. ix, fig. 25; * Obs., II, 1838, p. 30, pl. ix, fig. 25.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 235.—* HANLEY, Test. Moll., 1842, p. 206; * Biv. Shells, 1843, p. 206, pl. xxii, fig. 9.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 53, pl. xv, figs. 1, 1a.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* CHENU, Ill. Conch., 1858, pl. xx, figs. 3, 3a, 3b.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXX, fig. 423.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 158.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 424, pl. LXVI, fig. 4; LXVII, fig. 1.

* *Margarita (Unio) lugubris* LEA, Syn., 1836, p. 37; 1838, p. 24.

* *Margaron (Unio) lugubris* LEA, Syn., 1852, p. 38; 1870, p. 53.

*† *Unio confertus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 103, pl. xvi, fig. 47; * Obs., I, p. 215, pl. xvi, fig. 47.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 200; * Biv. Shells, 1843, p. 200, pl. xxiii, fig. 34.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1866, pl. xxxv, fig. 185.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Margarita (Unio) confertus* LEA, Syn., 1836, p. 31; 1838, p. 22.

* *Margaron (Unio) confertus* LEA, Syn., 1852, p. 33; 1870, p. 53.

*† *Unio geddingsianus* LEA, Pr. Am. Phil. Soc. I, 1840, p. 285; * Tr. Am. Phil. Soc., VIII, 1842, p. 202, pl. xi, fig. 15; * Obs., III, 1842, p. 40, pl. xi, fig. 15.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* CHENU, Ill. Conch., 1858, pl. xxxi, figs. 3, 3a, 3b.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 153.

* *Margaron (Unio) geddingsianus* LEA, Syn., 1852, p. 33; 1870, p. 53.

¹ Only a single broken right valve of this is in the Lea collection.

² The name *lugubris* applied first to this species will have to give way to *confertus* because Say used *lugubris* at an earlier date for a *Unio*.

- * *Unio limatulus* CONRAD, Pr. Ac. N. Sci. Phila., IV, 1849, p. 154; * Ann. and Mag., IV, 1849, p. 301; * Jl. Ac. N. Sci. Phila., I, 1850, p. 276, pl. xxxvii, fig. 9; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 157.
- * *Margaron (Unio) limatulus* LEA, Syn., 1852, p. 32; 1870, p. 52.
- * † *Unio whiteianus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 258, pl. xiv, fig. 8.—* Obs., V, 1852, p. 14, pl. xiv, fig. 8.—* H. and A. ADAMS, Gen., Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 172.
- * *Margaron (Unio) whiteianus* LEA, Syn., 1852, p. 33; 1870, p. 53.
- * *Unio vibex* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 260.¹
- * † *Unio similis* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 169; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 91, pl. xix, fig. 71; * Obs., VI, 1858, p. 91, pl. xix, fig. 71.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 167.
- * *Margaron (Unio) similis* LEA, Syn., 1852, p. 52; 1870, p. 53.
- * † *Unio protensus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 88; * Jl. Ac. N. Sci. Phila., VI, 1858, p. 256, pl. xxxi, fig. 71; * Obs., XII, 1869, p. 16, pl. xxxi, fig. 71.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) protensus* LEA, Syn., 1870, p. 60.
- * † *Unio radiolus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 192; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 21, pl. vi, fig. 18; * Obs., XIII, 1874, p. 25, pl. vi, fig. 18.—* B. H. WRIGHT, Check List, 1888.

North Carolina to Florida.

† UNIO INSULSUS Lea.

- * *Unio insulsus* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 86; * Jl. Ac. N. Sci. Phila., IX, 1862, p. 53, pl. i, fig. 199; * Obs., VIII, 1862, p. 57, pl. i, fig. 199.—* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 422, pl. lxxv, figs. 1, 5.
- * *Margaron (Unio) insulsus* LEA, Syn., 1870, p. 53.
- * †? *Unio lucidus* LEA,² Pr. Ac. N. Sci. Phila., VII, 1863, p. 192; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 9, pl. ii, fig. 6; * Obs., XI, 1867, p. 13, pl. ii, fig. 6.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) lucidus* LEA, Syn., 1870, p. 48.
- * † *Unio cistelliformis* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 192; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 19, pl. vi, fig. 17; * Obs., XI, 1867, p. 23, pl. vi, fig. 17.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) cistelliformis* LEA, Syn., 1870, p. 52.

North Carolina to Florida.

† UNIO OBNUBILUS Lea.

- * *Unio obnubilis* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 169; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 84, pl. xvii, fig. 64; * Obs., VI, 1858, p. 84, pl. xvii, fig. 64.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 424, pl. lxxvi, fig. 3.
- * *Margaron (Unio) obnubilis* LEA, Syn., 1870, p. 53.
- * † *Unio opacus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 169; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 86, pl. xviii, fig. 66; * Obs., VI, 1858, p. 86, pl. xviii, fig. 66.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 161.

¹ Conrad here states that his *vibex* = first *whiteianus* of Lea, and then *zeiglerianus* of Lea. The latter is totally different from *whiteianus*.

² Very near *U. confertus*. It is probably a delicate, light-colored, rayed *insulsus*.

- * *Margaron (Unio) opacus* LEA, Syn., 1870, p. 52.
 *† *Unio aquatus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 170; * JI. Ac. N. Sci. Phila., 1858, p. 89, pl. XIX, fig. 69; * Obs., VI, 1858, p. 89, pl. XIX, fig. 69.—* B. H. WRIGHT, Check List, 1888.
 * *Margaron (Unio) aquatus* LEA, Syn., 1870, p. 52.
 *† *Unio viridicatus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 170; * JI. Ac. N. Sci. Phila., IV, 1858, p. 87, pl. XVIII, fig. 67; * Obs., VI, 1858, pl. XVIII, fig. 67.
 * *Margaron (Unio) viridicatus* LEA, Syn., 1870, p. 52.
 *† *Unio hepaticus* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 154; * JI. Ac. N. Sci. Phila., IV, 1860, p. 348, pl. LVII, fig. 173; * Obs., VIII, p. 30, pl. LVII, fig. 173.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 154.
 * *Margaron (Unio) hepaticus* LEA, Syn., 1870, p. 52.
 *† *Unio santeensis* LEA, Pr. Ac. N. Sci. Phila., I 1871, p. 193.—* JI. Ac. N. Sci. Phila., VIII, 1874, p. 20, pl. VI, fig. 17; * Obs., XIII, 1874, p. 24, pl. VI, fig. 17.—* B. H. WRIGHT, Check List, 1888.
 *† *Unio nolani* B. H. WRIGHT, Pr. Ac. N. Sci. Phila., 1888, p. 116, pl. IV, fig. 11; Check List, 1888.

South Carolina to Florida.

† UNIO FUSCATUS LEA.

- * *Unio fuscatus* LEA, Desc. 12 Uniones, 1843 (no pagination).—* Tr. Am. Phil. Soc., IX, 1845?, p. 277, pl. XL, fig. 4; Obs., IV, 1848, p. 35, pl. XL, fig. 4.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Conch. Cab. Unio, 1848, p. 203, pl. LXVIII, fig. 2.—* REEVE, Conch. Leon., XVI, 1865, pl. XXI, fig. 95.—* PÆTEL, Conch. Sam., III, 1890, p. 153.—* SIMPSON, Pr. U. S. Nat. Mus., XVI, 1892, p. 420, pl. LXIII, figs. 2, 4.—* B. H. WRIGHT, Check List, 1888.
 * *Margaron (Unio) fuscatus* LEA, Syn., 1852, p. 30; 1870, p. 48.

Florida.

† UNIO OCCULTUS LEA.

- * *Unio occultus* LEA, Desc. 12 Uniones, 1843 (no pagination).—Tr. Am. Phil. Soc., IX, 1845?, p. 279, pl. XLI, fig. 7; * Obs., IV, 1848, p. 37, pl. XLI, fig. 7.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* KUSTER, Conch. Cab. Unio, 1861, p. 223, pl. LXXV, fig. 3.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 161.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 420, pl. LXIII, figs. 5, 6.
 * *Margaron (Unio) occultus* LEA, Syn., 1852, p. 48; 1870, p. 48.

Florida.

† UNIO ARCTATUS CONRAD.

- * *Unio arctatus* CONRAD, Am. JI. Sci., XXV, 1834, p. 340, pl. I, fig. 9; * New F. W. Shells, 1834, p. 36, pl. v, fig. 4, p. 67.—* MÖLLER, Syn. Nov. Gen., 1836, p. 200.—* HANLEY, Test. Moll., 1842, p. 207; * Biv. Shells, 1843, p. 207, pl. XXIII, fig. 47.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 19, pl. III, fig. 10.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* KUSTER, Conch. Cab. Unio, 1861, p. 195, pl. LXII, fig. 3.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.
 * *Margarita (Unio) arctatus* LEA, Syn., 1836, p. 38; 1838, p. 25.
 * *Margaron (Unio) arctatus* LEA, Syn., 1852, p. 33; 1870, p. 62.

†UNIO ARCTATUS var. TORTIVUS Lea.

*† *Unio tortivus* LEA,¹ Pr. Am. Phil. Soc., I, 1840, p. 287; Tr. Am. Phil. Soc., VIII, 1842, p. 204, pl. XII, fig. 17; Obs., III, 1842, p. 42, pl. XII, fig. 17.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.—H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* CHENU, Ill. Conch., 1858, pl. XXIX, figs. 6, 6a, 6b.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVI, fig. 193.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 169.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 421, pl. LXIII, fig. 8; LXIV, figs. 1, 3, 4.

*Margaron (*Unio*) *tortivus* LEA, Syn., 1852, p. 33; 1870, p. 53.

*† *Unio tetricus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 170; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 195, pl. XXII, fig. 78; *Obs., VII, 1859, p. 13, pl. XXII, fig. 78.—* KUSTER, Conch. Cab. Unio, 1862, p. 276, pl. XCI, fig. 7; XCIII, fig. 3.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 169.

*Margaron (*Unio*) *tetricus* LEA, Syn., 1870, p. 53.

*† *Unio purpurellus* LEA,² Pr. Ac. N. Sci. Phila., IX, 1857, p. 171; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 19, pl. XXIII, fig. 81; *Obs., VII, 1859, p. 16, pl. XXIII, fig. 81.—* B. H. WRIGHT, Check List, 1888.

*Margaron (*Unio*) *purpurellus* LEA, Syn., 1870, p. 53.

*† *Unio merceri* LEA, Pr. Ac. N. Sci. Phila., VI, 1862, p. 169; *Jl. Ac. N. Sci. Phila., V, 1862, p. 209, pl. XXXI, fig. 278; *Obs., IX, 1863, p. 31, pl. XXXI, fig. 278.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 159.

*Margaron (*Unio*) *merceri* LEA, Syn., 1870, p. 61.

Florida; South Georgia; the type west to the Black Warrior River, Alabama.

†UNIO NIGELLUS Lea.

* *Unio nigellus* LEA, Pr. Am. Phil. Soc., V, 1852, p. 251; *Tr. Am. Phil. Soc., X, 1852, p. 283, pl. XXIV, fig. 42; *Obs., V, 1852, p. 39, pl. XXIV, fig. 42.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 161.

*Margaron (*Unio*) *nigellus* LEA, Syn., 1852, p. 30; 1870, p. 53.

*† *Unio denigratus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 171; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 200, pl. XXIII, fig. 83; *Obs., VII, 1859, p. 18, pl. XXIII, fig. 83.—* B. H. WRIGHT, Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 422, pl. LXV, fig. 1.

*Margaron (*Unio*) *denigratus* LEA, Syn., 1870, p. 52.

Chattahoochee River system; south into Florida.

†UNIO PINEI B. H. Wright.

* *Unio pinei* B. H. WRIGHT, Naut., XI, 1897, p. 40.—* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 80, pl. III, fig. 1.

North Florida.

¹ *Unio arctatus* is a widely distributed, abundant, and variable species, ranging from the Black Warrior River to southern Florida. I have examined a great amount of material from the Black Warrior and Cahawba rivers (collected by Call), from the Flint and Chattahoochee rivers from the Lea and other collections, and I feel reasonably sure that all the species I have united should go together.

² I believe this to be a small, somewhat inflated form of *arctatus*.

† UNIO OSCARI B. H. Wright.

* *Unio oscari* B. H. WRIGHT, Naut., V, 1892, p. 124; IX, 1896, p. 122, pl. II, figs. 1-3.

Florida.

† UNIO HAZELHURSTIANUS Lea.

* *Unio hazelhurstianus* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 166; * Jl. Ac. N. Sci. Phila., IV, 1859, p. 211, pl. XXVI, fig. 92; * Obs., VII, 1859, p. 29, pl. XXVI, fig. 92.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXV, fig. 188.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 154.

* Margaron (*Unio*) *hazelhurstianus* LEA, Syn., 1870, p. 60.

Southern Georgia.

† UNIO BUXTONI B. H. Wright.

* *Unio buxtoni* B. H. WRIGHT, Naut., XI, 1897, p. 55 —* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 80, pl. I, fig. 6.

Marion County, Florida.

† UNIO FERRISSII Marsh.

* *Unio ferrissii* MARSH, Naut., V, 1891, p. 30.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 423, pl. LXVI, figs. 1, 2.

Small creek near Palatka, Florida.

† UNIO DOREI B. H. Wright.

* *Unio dorei* B. H. WRIGHT,² Pr. Ac. N. Sci. Phila., 1888, p. 115, pl. III, fig. 1.

Lake Monroe, Florida.

(Group of *Unio striatulus*.)

Shell subsolid, rhomboid oval, with a well-developed post ridge ending in a point below; beaks full, with numerous subparallel, slightly corrugated, strong ridges; epidermis rayless or faintly rayed, with a few wrinkles on the posterior slope; pseudocardinals stumpy; laterals straight. Animal unknown.

† UNIO STRIATULUS Lea.

* *Unio striatulus* LEA, Pr. Acad. N. Sci. Phila., I, 1857, p. 86; * Jl. Acad. N. Sci. Phila., V, 1862, p. 55, pl. II, fig. 202.—* Obs., VIII, 1862, p. 59, pl. II, fig. 202.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXII, fig. 367.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

* Margaron (*Unio*) *striatulus*, LEA, Syn., 1870, p. 37.

Roanoke River; Welden and Salem, North Carolina.

¹ This was first published in the Joliet Weekly News, a newspaper, May 1, 1891. It is a puzzling form, having something the shape of the variety *orcutti* of *U. buckleyi*, but has plications on the posterior slope such as are found in the *Crassidens* group.

² I do not know where to place this form, of which I have only seen the type. The figure is not very accurate, and I formerly thought it a variety of *U. buckleyi*, but on seeing the type I believe it is not that. It seems to combine characters of the *Buckleyi* and *Crassidens* groups.

† UNIO AMABILIS Lea.

* *Unio amabilis* LEA, Pr. Acad. N. Sci. Phila., IX, 1865, p. 89; * JI. Acad. N. Sci. Phila., VI, 1869, p. 257, pl. XXXI, fig. 72; * Obs., XII, 1869, p. 17, pl. XXXI, fig. 72.—* B. H. Wright, Check List, 1888.

* *Margaron (Unio) amabilis* LEA, Syn., 1870, p. 37.

Butler, Taylor County, Georgia; North Carolina?

UNIO SUBPLANUS Conrad.

Unio subplanus CONRAD, Monog., IX, 1837, p. 73, pl. XLI, fig. 1; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* HANLEY, Biv. Shells, Supp., 1856, p. 383, pl. XXI, fig. 16.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* KUSTER, Conch. Cab. Unio, 1862, p. 272, pl. XCI, fig. 5.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

* *Margaron (Unio) subplanus* LEA, Syn., 1852, p. 33; 1870, p. 54.

* *Unio subplanum* SOWERBY, Conch. Icon., XVI, 1866, pl. XLVII, fig. 252.

North Carolina and Virginia.

UNIO BRIMLEYI S. H. Wright.

* *Unio brimleyi* S. H. WRIGHT, Naut., X, 1897, p. 138; * SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 80, pl. IV, fig. 6.

Neuse River and Irwin's Creek, North Carolina.

(Group of *Unio complanatus*.)

Shell elongate trapezoidal, moderately solid, but little inflated, nearly straight below, with a posterior ridge, usually biangulate behind; beaks not prominent, sculptured with several coarse, parallel ridges which follow the growth lines or are nearly straight. Epidermis sometimes shining and feebly rayed, becoming rough and rayless with age; laterals straight or slightly curved; beak cavities shallow.

Animal with the marsupium occupying all or nearly all of the outer branchiæ; gills long, rounded below, inner much the larger, free nearly or quite their whole length from the abdominal sac.

† UNIO COMPLANATUS (Solander) Dillwyn.¹

Mya complanata SOLANDER, manuscript (no date).—* PORTLAND catalogue, 1786, p. 100, Lot 2190.—* DILLWYN, Cat., I, 1817, p. 51.

* *Unio complanata* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 559; 3d ed., II, 1839, p. 676.

¹ This species was described by Solander in Latin, but the manuscript was never published and is now in the British Museum. Mr. Edgar A. Smith kindly examined it for me and informed me that Dillwyn's description in the catalogue is a translation from Solander. The name *Mya complanata* was used by Lister in Hist. Sive. Conch. (1770, p. 150), but Lister was not a binomial author. The Portland catalogue is anonymous, and a mere sale list. I have done my best to straighten out the synonymy of this variable and exceedingly puzzling group, and have examined nearly all the types and a great amount of material. In some cases, as in the *Buckleyi* group, there seem to be no specific lines whatever.

- * *Unio complanatus* FERUSSAC, Guer. Mag., 1835, p. 26.—* GOULD. Inv. of Mass., 1841, p. 107, figs. 68-70.—* BINNEY, 2d ed., Inv. of Mass., 1870, p. 167, fig. 472.—* C. B. ADAMS, Thompson's Hist. Vt., 1842, p. 167.—* HANLEY, Test. Moll., 1842, p. 199; * Biv. Shells, 1843, p. 199, pl. XXI, fig. 20.—* DE KAY, Zool. of N. Y., Pt. 5, 1843, p. 188, pl. XXII, fig. 246.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* STIMPSON, Shells of N. Eng., 1851, p. 13.—* AGASSIZ, Arch. für Nat., I, 1852, p. 50.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—KUSTER, Conch. Cab. Unio, 1856, p. 137, pl. XLI, fig. 2.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* HARTMAN and MICHENER, Conch. Cest., 1874, p. 86, fig. 181.—* LATCHFORD, Tr. Ottawa Field Nat. Cl., 1882, p. 49.—* B. H. WRIGHT, Check List, 1888.—* H. CARPENTER, Naut., III, 1889, p. 93.—* PÄTEL, Conch. Sam., III, 1890, p. 148.
- * *Margarita (Unio) complanatus* LEA, Syn., 1836, p. 30; 1838, p. 22.
- * *Margaron (Unio) complanatus* LEA, Syn., 1852, p. 32; 1870, p. 51.
- * ? *Unio violaceus* SPENGLER, Skriv. Nat. Selsk. III, 1793, p. 55.
- * ? *Unio purpureus* SAY, Nich. Enc., II, 1816, pl. III, fig. 1; N. Harm. Diss., II, No. 21, 1829, p. 324.—* CONRAD, New F. W. Shells, 1834, p. 71.—* SOWERBY, Rich. Faun. Boreali Am., III, 1836, p. 316.—* L. W. SAY, Terr. and Fluv. Shells, 1840, p. 7.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXVIII, fig. 346.
- * *Mya purpurea* EATON, Zool. Text-Book, 1826, p. 2.
- * ? *Unio purpurea* DESHAYES, Encyc. Méth. II, 1827, p. 151, pl. 249, fig. 5.
- * *Unio rarisulcata* LAMARCK, An. sans Vert., VI, 1819, p. 72.—* LEA, Obs., I, 1834, p. 200.—DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 534.
- * *Unio coarctata* LAMARCK, An. sans Vert., VI, 1819, p. 73.—* LEA, Obs., I, 1834, p. 200.—DESHAYES, Encyc. Méth., II, 1830, p. 581; * An. sans Vert., 2d ed., VI, 1835, p. 535; 3d ed., II, 1839, p. 668.—* ? SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIV, fig. 178.
- * *Unio purpurascens* LAMARCK, An. sans Vert., VI, 1819, p. 73.—DESHAYES, Enc. Méth., II, 1830, p. 581.—* LEA, Obs., I, 1834, p. 200.—DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 535; 3d ed., II, 1839, p. 668.
- * *Unio georgiana* LAMARCK, An. sans Vert., VI, 1819, p. 74.—* LEA, Obs., I, 1834, p. 201.—DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 536; 3d ed., II, 1839, p. 668.
- * *Unio rhombula* LAMARCK, An. sans Vert., VI, 1819, p. 74.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 536; 3d ed., II, 1839, p. 668.
- * *Unio rhombula* LAMARCK, An. sans Vert., VI, 1819, p. 74.—* LEA, Obs., I, 1834, p. 200.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 536; 3d ed., II, 1839, p. 668.—* ? DELESSERT, Rec. Coq. Sam., 1841, pl. XII, fig. 8.
- * *Unio carinifera* LAMARCK, An. sans Vert., VI, 1819, p. 74.—* LEA, Obs., I, 1834, p. 201.—DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 536; 3d ed., II, 1839, p. 668.—* KUSTER, Conch. Cab. Unio, 1861, p. 176, pl. LVI, fig. 1.
- * *Unio glabrata* LAMARCK, An. sans Vert., VI, 1819, p. 75.—* LEA, Obs., I, 1834, p. 201.—DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 537; 3d ed., II, 1839, p. 669.
- * *Unio sulcidens* LAMARCK, An. sans Vert., VI, 1819, p. 77.—* LEA, Obs., I, 1834, p. 202.—DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 539; 3d ed., II, 1839, p. 670.—* ? DELESSERT, Rec. Coq. Sam., 1841, pl. XII, fig. 3.—* CHENU, Ill. Conch., 1858, pl. XII, figs. 5, 5a.
- * *Unio virginiana* LAMARCK, An. sans Vert., VI, 1819, p. 79.—DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 544; 3d ed., II, 1839, p. 671.
- Unio fluviatilis* GREEN, Jl. Mac. Lyceum, 1827, p. 41.
- * ? *Unio raveneli* CONRAD, New F. W. Shells, 1834, p. 39, pl. VI, fig. 4, 1 p. 71.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen., 1836, p. 206.—

¹ It is hard to say what this is. The figure is like *complanatus*, but the shell, which is labeled *raveneli* Conrad and is in the Philadelphia Academy of Natural Sciences, is like *lugubris*.

- * CHENU, Bib. Conch., 1st ser., III, p. 9, pl. III, fig. 8.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 256.—* B. H. WRIGHT, Check List, 1888.
- * *Margarita (Unio) waterensis* LEA, Syn., 1836, p. 31; ¹ 1838, p. 22.
- * *Margaron (Unio) waterensis*, LEA, Syn., 1852, p. 33; 1870, p. 53.
- * *Unio waterensis*, H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—PÆTEL, Conch. Sam., III, 1890, p. 172.
- *† *Unio griffithianus*, LEA, Tr. Am. Phil. Soc., V, 1834, p. 103, pl. XV, fig. 46; Obs., I, 1834, p. 215, pl. XV, fig. 46.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 199; * Biv. Shells, 1843, p. 199, pl. XXIII, fig. 28.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* KUSTER, Conch. Cab. Unio, 1861, p. 208, pl. LXIX, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIX, fig. 449.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 154.
- * *Margarita (Unio) griffithianus*, LEA, Syn., 1836, p. 31; 1838, p. 22.
- * *Margaron (Unio) griffithianus*, LEA, Syn., 1852, p. 33; 1870, p. 52.
- *† *Unio planilaterus*, CONRAD, Monog., XII (no date after 1838), p. 103, pl. LVII, fig. 1; Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 163.
- * *Margaron (Unio) planilaterus*, LEA, Syn., 1870, p. 51.
- *† *Unio fuliginosus*, LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; Tr. Am. Phil. Soc., X, 1848, p. 78, pl. VII, fig. 19.
- *† *Unio rufusculus* LEA, Pr. Am. Phil. Soc., V, 1852, p. 252; * Tr. Am. Phil. Soc., X, 1852, p. 258, pl. XIV, fig. 7; * Obs., V, 1852, p. 14, pl. XIV, fig. 7.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIII, fig. 377.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 166.
- * *Margaron (Unio) rufusculus* LEA, Syn., 1852, p. 33; 1870, p. 54.
- * *Mya rigida* WOOD, Ind. Test. Rev., 1856, p. 200, pl. I, supp. fig. 10.
- *† *Unio abberillensis* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 84; * JI. Ac. N. Sci. Phila., IV, 1858, p. 51, pl. VI, fig. 34; * Obs., VI, 1858, p. 51, pl. VI, fig. 34.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 143.
- * *Margaron (Unio) abberillensis* LEA, Syn., 1870, p. 52.
- *† *Unio contractus* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 86; * JI. Ac. N. Sci. Phila., V, 1862, p. 203, pl. XXIX, fig. 272; * Obs., IX, p. 25, pl. XXIX, fig. 272.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) contractus* LEA, Syn., 1870, p. 61.
- *† *Unio virens* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 169; * JI. Ac. N. Sci. Phila., IV, 1858, p. 80, pl. XVI, fig. 60; * Obs., VI, 1858, p. 80, pl. XVI, fig. 60.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) virens* LEA, Syn., 1870, p. 52.
- *† *Unio savannahensis* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 169; * JI. Ac. N. Sci. Phila., IV, 1858, p. 81, pl. XVI, fig. 61; * Obs., VI, 1858, p. 81, pl. XVI, fig. 61.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) savannahensis* LEA, Syn., 1870, p. 51.
- *† *Unio subflavus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 170; * JI. Ac. N. Sci. Phila., IV, 1858, p. 90, pl. XIX, fig. 70; * Obs., VI, 1858, p. 90, pl. XIX, fig. 70.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) subflavus* LEA, Syn., 1870, p. 52.
- *† *Unio neusensis* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 41; * JI. Ac. N. Sci. Phila., V, 1859, p. 60, pl. IV, fig. 208; * Obs., VIII, 1860, p. 64, pl. IV, fig. 208.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) neusensis* LEA, Syn., 1870, p. 51.

¹ Lea changed the name *raveneli* to *waterensis*, because the name *ravenelianus* had been applied by him to a *Unio*.

- *† *Unio exactus* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 41; *Jl. Ac. N. Sci. Phila., V, 1862, p. 62; *Obs., VIII, 1860, p. 66, pl. iv, fig. 210.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) exactus* LEA, Syn., 1870, p. 43.
- *† *Unio roswellensis* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 165; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 205, pl. xxiv, fig. 87; *Obs., VII, 1859, p. 23, pl. xxiv, fig. 87.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) roswellensis* LEA, Syn., 1870, p. 51.
- *† *Unio postellii* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 165; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 214, pl. xxvi, fig. 94; *Obs., VII, 1859, p. 32, pl. xxvi, fig. 94.—* SOWERBY, Conch. Icon., XVI, 1866, pl. xlvii, p. 255.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) postellii* LEA, Syn., 1870, p. 51.
- *† *Unio baldwinensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 170; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 330, pl. li, fig. 155; *Obs., VIII, 1860, p. 12, pl. li, fig. 155.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) baldwinensis* LEA, Syn., 1870, p. 51.
- *† *Unio raeensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 171; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 331, pl. lii, fig. 156; *Obs., VIII, 1860, p. 13, pl. lii, fig. 156.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) raeensis* LEA, Syn., 1870, p. 52.
- *† *Unio quadratus* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 172; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 338, pl. liv, fig. 163; *Obs., VIII, 1860, p. 20, pl. liv, fig. 163.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) quadratus* LEA, Syn., 1870, p. 54.
- *† *Unio squameus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 391; *Jl. Ac. N. Sci. Phila., V, 1862, p. 200, pl. xxviii, fig. 269; *Obs., IX, 1863, p. 22, pl. xxviii, fig. 269.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) squameus* LEA, Syn., 1870, p. 51.
- * *Unio lugubris* KUSTER, Conch. Cab. Unio, 1861, p. 234, pl. lxxix, fig. 1.
- *† *Unio weldonensis* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 191; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 11, pl. iii, fig. 8; *Obs., XI, 1867, p. 15, pl. iii, fig. 8.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) weldonensis* LEA, Syn., 1870, p. 51.
- *† *Unio gastonensis* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 191; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 20, pl. vi, fig. 18; *Obs., XI, 1867, p. 24, pl. vi, fig. 18.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) gastonensis* LEA, Syn., 1870, p. 51.
- *† *Unio aberrans* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 191; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 7, pl. i, fig. 3; *Obs., XI, 1867, p. 11, pl. i, fig. 3.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) aberrans* LEA, Syn., 1870, p. 52.
- *† *Unio mecklenbergensis* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 191; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 17, pl. v, fig. 15; *Obs., XI, 1867, p. 21, pl. v, fig. 15.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) mecklenbergensis* LEA, Syn., 1870, p. 51.
- *† *Unio raleighensis* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 191; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 6, pl. i, fig. 2; *Obs., XI, 1867, p. 10, pl. i, fig. 2.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) raleighensis* LEA, Syn., 1870, p. 52.
- *† *Unio mediocris* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 192; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 24, pl. vii, fig. 22; *Obs., XI, 1867, p. 28, pl. vii, fig. 22.
- * *Margaron (Unio) mediocris* LEA, Syn., 1870, p. 52.
- *† *Unio indefinitus* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 192; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 15, pl. iv, fig. 12; *Obs., XI, 1867, p. 12, pl. iv, fig. 112.—* B. H. WRIGHT, Check List, 1888.

- * *Margaron (Unio) indefinitus* LEA, Syn., 1870, p. 51.
- *† *Unio humerosus* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 161; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 301, pl. XLV, fig. 113; *Obs., XII, 1869, p. 61, pl. XLV, fig. 113.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) humerosus* LEA, Syn., 1870, p. 51.
- *† *Unio beaverensis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 161; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 297, pl. XLIV, fig. 109.—*Obs., XII, 1869, p. 57, pl. XLIV, fig. 109.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) beaverensis* LEA, Syn., 1870, p. 52.
- * *Unio bleaverenses* PÆTEL, Conch. Sam., III, 1890, p. 146.
- * ? *Unio neglectus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXV, fig. 329.
- * *Unio tortuosus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXV, fig. 330.—* PÆTEL, Conch. Sam., III, 1890, p. 169.
- * *Unio uharensis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 145; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 304, pl. XLVI, fig. 116; *Obs., XII, 1869, p. 63, pl. XLVI, fig. 116.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) uharensis* LEA, Syn., 1870, p. 52.
- *† *Unio nubilus* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 161; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 298, pl. XLIV, fig. 110; *Obs., XII, p. 58, pl. XLIV, fig. 110.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) nubilus* LEA, Syn., 1870, p. 51.
- *† *Unio yadkinensis* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 156; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 32, pl. x, fig. 29; *Obs., XIII, 1874, p. 36, pl. x, fig. 29.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
- *† *Unio amplus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 157; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 39, pl. XIII, fig. 36; *Obs., XIII, 1874, p. 43, pl. XIII, fig. 36.—* B. H. WRIGHT, Check List, 1888.
- *† *Unio ligatus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 157; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 49, pl. XVII, fig. 47; *Obs., XIII, 1874, p. 53, pl. XVII, fig. 47; * B. H. WRIGHT, Check List, 1888.
- *† *Unio subparallelus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 158; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 46, pl. XVI, fig. 44; *Obs., XIII, 1874, p. 50, pl. XVI, fig. 44.—* B. H. WRIGHT, Check List, 1888.
- *† *Unio irwinensis* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 159; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 44, pl. xv, fig. 42; *Obs., XIII, 1874, p. 48, pl. xv, fig. 42.—* B. H. WRIGHT, Check List, 1888.
- *† *Unio curvatus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 159; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 38, pl. XIII, fig. 35; *Obs., XIII, 1874, p. 42, pl. XIII, fig. 35.
- *† *Unio subsquamosus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 160; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 47, pl. XVI, fig. 45; *Obs., XIII, 1874, p. 51, pl. XVI, fig. 45.—* B. H. WRIGHT, Check List, 1888.
- *† *Unio infuscus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 160; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 51, pl. XVII, fig. 49; *Obs., XIII, 1874, p. 55, pl. XVII, fig. 49.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
- *† *Unio ratus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 160; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 52, pl. XVIII, fig. 51; *Obs., XIII, 1874, p. 56, pl. XVIII, fig. 51; * B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 165.
- *† *Unio basalis* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 161; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 48, pl. XVI, fig. 46; *Obs., XIII, 1874, p. 52, pl. XVI, fig. 46.—* B. H. WRIGHT, Check List, 1888.
- *† *Unio subolivaceus* LEA, Pr. Ac. N. Sci. Phila., III, 1873, p. 422; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 57, pl. xx, fig. 56; *Obs., XIII, 1874, p. 61, pl. xx, fig. 56.—* B. H. WRIGHT, Check List, 1888.
- *† *Unio infulgens* LEA, Pr. Ac. N. Sci. Phila., III, 1873, p. 422; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 60, pl. XXI, fig. 59; *Obs., XIII, 1874, p. 64, pl. XXI, fig. 59.—* B. H. WRIGHT, Check List, 1888.

- *† *Unio cirratus* LEA, Pr. Ac. N. Sci. Phila., III, 1873, p. 422; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 53, pl. XIX, fig. 53; *Obs., XIII, 1874, p. 59, pl. XIX, fig. 53.—*B. H. WRIGHT, Check List, 1888.
- *† *Unio corneus* LEA, Pr. Ac. N. Sci. Phila., III, 1873, p. 423; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 59, pl. XX, fig. 58; *Obs., XIII, 1874, p. 63, pl. XX, fig. 58.—*B. H. WRIGHT, Check List, 1888.
- *† *Unio invenustus* LEA, Pr. Ac. N. Sci. Phila., III, 1873, p. 424; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 66, pl. XXII, fig. 62; *Obs., XIII, 1874, p. 70, pl. XXII, fig. 62.—*B. H. WRIGHT, Check List, 1888.
- *† *Arcouaia procancheriaua* PILSBRY, Nat. Canadienne, XX, 1890, p. 171; *Pr. Ac. N. Sci. Phila., 1892, p. 132, pl. VII, figs. 4-6.¹

† UNIO COMPLANATUS var. JEJUNUS Lea.

- * *Unio jejunus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 9, pl. IV, fig. 9; *Obs., II, 1838, p. 9, pl. IV, fig. 9.—*TROSCHER, Arch. für Nat., V, 1839, Pt. 2, p. 234.—*HANLEY, Test. Moll., 1842, p. 199; *Biv. Shells, 1843, p. 199, pl. XXII, fig. 47.—*CATLOW and REEVE, Conch. Nom., 1845, p. 60.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*KUSTER, Conch. Cab. Unio, 1862, p. 265, pl. XC, fig. 1.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXVIII, fig. 347.²—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 155.
- * *Margarita (Unio) jejunus* LEA, Syn., 1836, p. 30; 1838, p. 22.
- * *Margaron (Unio) jejunus* LEA, Syn., 1852, p. 32; 1870, p. 51.
- *† *Unio percoarctatus* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 85; *Jl. Ac. N. Sci. Phila., V, 1862, p. 59, pl. III, fig. 206; *Obs., VIII, 1862, p. 63, pl. III, fig. 206.—*SOWERBY, Conch. Icon., XVI, 1866, pl. LIV, fig. 277.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 163.
- * *Margaron (Unio) percoarctatus* LEA, Syn., 1870, p. 51.

† UNIO COMPLANATUS var. QUADRILATERUS Lea.

- * *Unio squalidus* LEA, in part, Pr. Ac. N. Sci. Phila., VII, 1863, p. 192; Jl. Ac. N. Sci. Phila., VI, 1866, p. 22, pl. VII, fig. 20; Obs., XI, 1867, p. 26, pl. VII, fig. 20.—*B. H. WRIGHT, Check List, 1888.—*SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 429, pl. LXXI, fig. 14.
- * *Margaron (Unio) squalidus* LEA, Syn., 1870, p. 51.
- *† *Unio quadrilaterus* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 192; Jl. Ac. N. Sci. Phila., VI, 1866, p. 5, pl. I, fig. 1; Obs., XI, 1867, p. 9, pl. I, fig. 1.—*B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) quadrilaterus* LEA, Syn., 1870, p. 51.

Atlantic drainage from the St. Lawrence to Georgia; west in the British possessions to Manitoba?

† UNIO CATAWBENSIS Lea.

- * *Unio wheatleyi* LEA, Pr. Ac. N. Sci. Phila., 1857, I, p. 85;³ *Jl. Ac. N. Sci. Phila., V, 1861, p. 54, pl. I, fig. 200; Obs., VIII, p. 58, pl. I, fig. 200.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 147.
- * *Margaron (Unio) catawbensis* LEA, Syn., 1870, p. 52.

¹ Said to come from Yamaska River, China. The type is a twisted *Unio complanatus*, probably from Canada.

² In errata he says this is *depressus* of d'Orbigny, not Lamarek. It is hard to say what it is.

³ Preoccupied by Lea and changed by him to *catawbensis*.

- * *Unio rostrum* LEA, Pr. Ac. N. Sci. Phila., VIII, 1861, p. 391; * Jl. Ac. N. Sci. Phila., V, 1862, p. 201, pl. XXIX, fig. 270; Obs., IX, 1863, p. 23, pl. XXIX, fig. 270.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) rostrum* LEA, Syn., 1870, p. 52.
- * † *Unio oblongus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 158; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 52, pl. XVIII, fig. 50; Obs., XIII, 1874, p. 56, pl. XVIII, fig. 50.

North Carolina.

† UNIO TUOMEYI Lea.

- * *Unio tuomeyi* LEA, Tr. Am. Phil. Soc., X, 1852, p. 256, pl. XIII, fig. 4; * Obs., V, 1852, p. 12, pl. XIII, fig. 4.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 170.
- * *Margaron (Unio) tuomeyi* LEA, Syn., 1852, p. 33; 1870, p. 53.
- * † *Unio barrattii* LEA, Tr. Am. Phil. Soc., X, 1852, p. 256, pl. XIII, fig. 5; * Obs., V, 1852, p. 12, pl. XIII, fig. 5.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) barrattii* LEA, Syn., 1852, p. 37; 1870, p. 245.
- * *Unio barrotti* PÆTEL, Conch. Sam., III, 1890, p. 145.
- * † *Unio pullatus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 262; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 57, pl. VIII, fig. 39; * Obs., VI, 1858, p. 57, pl. VIII, fig. 39.—* KUSTER, Conch. Cab. Unio, 1871, p. 247, pl. LXXXIII, fig. 3.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXVI, fig. 335.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.
- * *Margaron (Unio) pullatus* LEA, Syn., 1870, p. 60.
- * † *Unio sublatus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 169; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 82, pl. XVI, fig. 62; * Obs., VI, 1858, p. 82, pl. XVI, fig. 62.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) sublatus* LEA, Syn., 1870, p. 57.
- * † *Unio fumatus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 171; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 88, pl. XVIII, fig. 68; * Obs., VI, 1858, p. 88, pl. XVIII, fig. 68.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) fumatus* LEA, Syn., 1870, p. 52.
- * † *Unio viridiradiatus* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 154; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 336, pl. LIII, fig. 161; * Obs., VIII, p. 18, pl. LIII, fig. 161.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) viridiradiatus* LEA, Syn., 1870, p. 46.
- * † *Unio viridans* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 170; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 337, pl. LIV, fig. 162; * Obs., VIII, 1860, p. 19, pl. LIV, fig. 162.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) viridans* LEA, Syn., 1870, p. 52.
- * † *Unio hallenbeckii* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 170; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 328, pl. LI, fig. 154; ¹ * Obs., VIII, 1860, p. 10, pl. LI, fig. 154.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) hallenbeckii* LEA, Syn., 1870, p. 52.
- * † *Unio salebrosus* LEA, Pr. Ac. N. Sci. Phila., XI, 1859, p. 170; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 332, pl. LII, fig. 157; * Obs., VIII, 1860, p. 14, pl. LII, fig. 157.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 166.
- * *Margaron (Unio) salebrosus* LEA, Syn., 1870, p. 52.
- * † *Unio verutus* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 471; * Jl. Ac. N. Sci.

¹Extra copies of the paper containing this species were published in December, 1858.

- Phila., IV, 1860, pl. LIII, fig. 160; *Obs., VIII, 1860, p. 17, pl. LIII, fig. 160.—
 * B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 171.
 *Margaron (*Unio*) *verutus* LEA, Syn., 1870, p. 57.
 *† *Unio chathamensis* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 191; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 21, pl. VI, fig. 19; *Obs., XI, 1867, p. 25, pl. VI, fig. 19.—
 * B. H. WRIGHT, Check List, 1888.
 Margaron (*Unio*) *chathamensis* LEA, Syn., 1870, p. 53.
 *† *Unio hastatus* LEA, Pr. Ac. N. Sci. Phila., III, 1873, p. 423; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 56, pl. XIX, fig. 54; *Obs., XIII, 1874, p. 60, pl. XIX, fig. 54.—B. H. WRIGHT, Check List, 1888.
 *† *Unio dooleyensis* LEA, Pr. Ac. N. Sci. Phila., III, 1873, p. 424; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 64, pl. XXII, fig. 60; *Obs., XIII, 1874, p. 68, pl. XXII, fig. 60.—* B. H. WRIGHT, Check List, 1888.
 *† *Unio gesnerii* LEA, Pr. Ac. N. Sci. Phila., III, 1874, p. 424; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 65, pl. XXII, fig. 64; *Obs., XIII, 1874, p. 69, pl. XXII, fig. 61.—
 SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIV, fig. 446.— B. H. WRIGHT, Check List, 1888.

North Carolina to Alabama.

† UNIO OCMULGEENSIS Lea.

- * *Unio ocmulgeensis* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 38; *Jl. Ac. N. Sci. Phila., V, 1862, p. 95, pl. XIV, fig. 243; *Obs., VIII, 1862, p. 99, pl. XIV, fig. 243.—
 SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 424, pl. LXVII, fig. 5.— B. H. WRIGHT, Check List, 1888.
 *Margaron (*Unio*) *ocmulgeensis* LEA, Syn., 1870, p. 57.

Georgia.

† UNIO AQUILUS Lea.

- * *Unio aquilus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 172; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 92, pl. XX, fig. 72; *Obs., VI, 1858, p. 92, pl. XX, fig. 72.—* B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 144.
 *Margaron (*Unio*) *aquilus* LEA, Syn., 1870, p. 60.

South Georgia and northern Florida.

† UNIO ICTERINUS Conrad.

- * *Unio icterinus* CONRAD, New F. W. Shells, 1834, pl. VI, fig. 5, p. 69.—*FERUSSAC, Guer. Mag., 1835, p. 29.—*CONRAD, Monog., IV, 1836, p. 39, pl. XVIII, fig. 2.—
 *MÖLLER, Syn. Nov. Gen., 1836, p. 206.—*CHENU, Bib. Conch., 1st ser., III, 1845, p. 20, pl. I, fig. 5.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—
 * B. H. WRIGHT, Check List, 1888.
 *† *Unio fuliginosus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; *Tr. Am. Phil. Soc., X, 1848, p. 78, pl. VII, fig. 19; *Obs., IV., 1848, p. 52, pl. VII, fig. 19.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 153.
 *Margaron (*Unio*) *fuliginosus* LEA, Syn., 1852, p. 33; 1870, p. 53.
 *† *Unio currierianus* LEA, Tr. Am. Phil. Soc., IX, 1852, p. 263, pl. XVI, fig. 16; Obs., V, 1852, p. 19, pl. XVI, fig. 16.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 150.
 *Margaron (*Unio*) *currierianus* LEA, Syn., 1852, p. 32; 1870, p. 51.
 *† *Unio curatus* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 193; *Jl. Ac. N. Sci. Phila.,

- VI, 1866, p. 23, pl. VII, fig. 21; Obs., XI, 1867, p. 27, pl. VII, fig. 21.—*P. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 149.
- **Margaron (Unio) curatus* LEA, Syn., 1870, p. 46.
- *† *Unio datus* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 161; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 299, pl. XLIV, fig. 3; *Obs., XII, 1869, p. 59, pl. XLIV, fig. 3.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 150.
- **Margaron (Unio) datus* LEA, Syn., 1870, p. 45.
- **Unio ratus* PÆTEL, Conch. Sam., III, 1890, p. 172.
- *† *Unio cuspidatus*, LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 159; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 43, pl. XIV, fig. 50; *Obs., XIII, 1874, p. 47, pl. XIV, fig. 40.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 150.

North Carolina to Georgia.

† UNIO ROANOKENSIS Lea.

- **Margarita (Unio) roanokensis* LEA, Syn., 1836, p. 30; 1838, p. 21.
- * *Unio roanokensis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 27, pl. VIII, fig. 21; Obs., II, 1838, p. 27, pl. VIII, fig. 21.—*TROSCHEL, Arch. für Nat., V, 1839, Pt. 2, p. 235.—*HANLEY, Test. Moll., 1842, p. 199; *Biv. Shells, 1843, p. 199.—*CATLOW and REEVE, Conch. Nom., 1845, p. 63.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 275.—*CHENU, Ill. Conch., 1858, pl. XX, figs. 6, 6a, 6b.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXVI, fig. 341.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam. III, 1890, p. 165.
- **Margaron (Unio)* LEA, Syn., 1852, p. 32; 1870, p. 51.
- *† *Unio macer* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 86; *Jl. Ac. N. Sci. Phila., VI, 1862, p. 202, pl. XXIX, fig. 271; *Obs., IX, 1863, p. 24, pl. XXIX, fig. 271.
- **Margaron (Unio) macer* LEA, Syn., 1870, p. 51.
- *† *Unio latus* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 171; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 334, pl. LIII, fig. 159; *Obs., VIII, 1860, p. 16, pl. LIII, fig. 159.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 157.
- **Margaron (Unio) latus* LEA, Syn., 1870, p. 57.

† UNIO ROANOKENSIS var. NORTHAMPTONENSIS Lea.

- **Unio northamptonensis* LEA,¹ Pr. Ac. N. Sci. Phila., V, 1861, p. 392; *Jl. Ac. N. Sci. Phila., V., 1862, p. 190, pl. XXV, fig. 260; *Obs., IX, 1863, p. 12, pl. XXV, fig. 260.—*B. H. WRIGHT, Check List, 1888.
- **Margaron (Unio) northamptonensis* LEA, Syn., 1870, p. 51.

Connecticut River at Northampton, Massachusetts; south to the Savannah River, Georgia.

† UNIO HOPETONENSIS Lea.

- **Unio hopetonensis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 29, pl. IX, fig. 24; *Obs., II, 1838, p. 29, pl. IX, fig. 24.—*TROSCHEL, Arch. für Nat., VI, 1839, Pt. 2, p. 235.—*HANLEY, Test. Moll., 1842, p. 198; *Biv. Shells, 1843, p. 198, pl. XX, fig. 21.—*CATLOW and REEVE, Conch. Nom., 1845, p. 60.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*CHENU, Ill. Conch., 1858, pl. XX, figs. 5, 5a, 5b.—*KUSTER, Conch. Cab. Unio, 1861, p. 196, pl. LXIII, fig. 1.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXVIII, fig. 349.—*B. H. WRIGHT, Check List,

¹ A large, compressed form, biangulate behind, the posterior point elevated above the base line. Certain specimens are considerably inflated in the post basal region. This is generally taken for *U. complanatus*, but it differs in the characters given above, and seems to be merely a northern variety of *roanokensis*.

1888.—* PÆTEL, Conch. Sam., III, 1890, p. 155.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 412, pl. LII, fig. 3; LIII, fig. 1.

* *Margarita (Unio) hopetonensis* LEA, Syn., 1836, p. 30; 1838, p. 21.

* *Margaron (Unio) hopetonensis* LEA, Syn., 1852, p. 32; 1870, p. 51.

Georgia, in streams flowing into the Atlantic; Santee Canal, South Carolina; northern Florida.

† UNIO LIVINGSTONENSIS LEA.

* *Unio livingstonensis* LEA, Pr. Ac. N. Sci. Phila., VII, p. 192; * JI. Ac. N. Sci. Phila., VI, 1866, p. 14, pl. IV, fig. 11; * Obs., XI, 1867, p. 18, pl. IV, fig. 11.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) livingstonensis* LEA, Syn., 1870, p. 52.

North Carolina to Georgia.

† UNIO INUSITATIS LEA.

* *Unio inusitatis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 171; * JI. Ac. N. Sci. Phila., IV, 1860, p. 333, pl. LII, fig. 158; * Obs., VIII, 1860, p. 15, pl. LII, fig. 158.

* *Margaron (Unio) inusitatus* LEA, Syn., 1870, p. 51.

* *Unio inusitatus* B. H. WRIGHT, Check List, 1888.

Georgia.

† UNIO DIFFERTUS LEA.

* *Unio differtus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 158; * JI. Ac. N. Sci. Phila., VIII, 1874, p. 42, pl. XIV, fig. 39; * Obs., XIII, 1874, p. 46, pl. XIV, fig. 39.—

* B. H. WRIGHT, Check List, 1888.

Georgia.

† UNIO PLANTII LEA.

* *Unio plantii* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 171; * JI. Ac. N. Sci. Phila., IV, 1859, p. 192, pl. XXI, fig. 76; * Obs., VII, 1859, p. 10, pl. XXI, fig. 76.—

* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVII, fig. 473.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., 1888.

* *Margaron (Unio) plantii* LEA, Syn., 1870, p. 43.

Flint River, near Macon, Georgia.

† UNIO STRUMOSUS LEA.

* *Unio strumosus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 158; III, 1873, p. 423;

* JI. Ac. N. Sci. Phila., VIII, 1874, p. 57, pl. XIX, fig. 55; * Obs., XIII, 1874, p. 61, pl. XIX, fig. 55.—* B. H. WRIGHT, Check List, 1888.

Yadkin River, North Carolina.

† UNIO PURUS LEA.

* *Unio purus* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 41; * JI. Ac. N. Sci. Phila., V, 1862, p. 61, pl. IV, fig. 209; * Obs., VIII, 1862, p. 65, pl. IV, fig. 209.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) purus* LEA, Syn., 1870, p. 43.

Neuse River, near Raleigh, North Carolina.

† UNIO SUBNIGER Lea.

* *Unio subniger* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 172; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 196, pl. XXII, fig. 79; *Obs., VII, 1859, p. 14, pl. XXII, fig. 79.—*B. H. WRIGHT, Check List, 1888.

*Margaron (*Unio*) *subniger* LEA, Syn., 1870, p. 52.

Georgia.

† UNIO BURKENSIS Lea.

* *Unio burkensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 112; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 215, pl. XXVII, fig. 95; *Obs., VII, 1859, p. 33, pl. XXVII, fig. 99.—*B. H. WRIGHT, Check List, 1888.

*Margaron (*Unio*) *burkensis* LEA, Syn., 1870, p. 60.

*† *Unio dissimilis* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 161; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 53, pl. XVIII, fig. 52; *Obs., XII, 1874, p. 57, pl. XVIII, fig. 52.—*B. H. WRIGHT, Check List, 1888.

North Carolina; south to Georgia.

† UNIO OBLATUS Lea.

* *Unio oblatus* LEA,¹ Pr. Ac. N. Sci. Phila., VII, 1863, p. 193.

* *Unio oblatus* LEA, Jl. Ac. N. Sci. Phila., VI, 1866, p. 13, pl. IV, fig. 10; *Obs., XI, 1867, p. 17, pl. IV, fig. 10.

*Margaron (*Unio*) *oblatus* LEA, Syn., 1870, p. 60.

Long Creek, Gaston County, North Carolina.

† UNIO ERRANS Lea.

† *Unio paliatus* RAVENEL, letter.²—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.

* *Margarita* (*Unio*) *paliatus* LEA, Syn., 1836, p. 31; 1838, p. 22.

*Margaron (*Unio*) *paliatus* LEA, Syn., 1852, p. 33; 1870, p. 53.

* *Unio palliatus* RAVENEL, Cat., 1875, p. 57.³—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Unio errans* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 262; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 60, pl. IX, fig. 42; *Obs., VI, 1858, p. 60, pl. IX, fig. 42.—* B. H. WRIGHT, Check List, 1888.

*Margaron (*Unio*) *errans* LEA, Syn., 1870, p. 52.

*† *Unio vicinus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 262; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 61, pl. IX, fig. 43; *Obs., VI, 1858, p. 61, pl. IX, fig. 43.—* B. H. WRIGHT, Check List, 1888.

*Margaron (*Unio*) *vicinus* LEA, Syn., 1870, p. 52.

Near Macon, Georgia; South Carolina.

¹ So written in first description. A typographical error, probably. Dr. Lea afterwards changed the name to *oblatus*.

² Dr. Lea credits this to Ravenel's letter, but neither he nor Ravenel ever described it so far as I know.

³ Ravenel here credits this species to Lea. Lea's specimens, received from Ravenel, are marked Santee Canal and South Carolina. They are rather thin and delicate, of a peculiar texture, and have a shining brownish green epidermis.

† UNIO SAGITTIFORMIS Lea.¹

* *Uniosagittiformis* LEA, Tr. Am. Phil. Soc., X, 1852, p. 277, pl. XXII, fig. 35; * Obs., V, 1852, p. 33, pl. XXII, fig. 35.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCII, fig. 499.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 166.

* *Margaron (Unio) sagittiformis* LEA, Syn., 1852, p. 37; 1870, p. 60.

Oconee River, Georgia; Abbeville, South Carolina.

† UNIO ANGUSTATUS Lea.

* *Unio angustatus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 114, pl. XVII, fig. 43; * Obs., I, 1834, p. 124, pl. XVII, fig. 43.—* CONRAD, New F. W. Shells, 1834, p. 67.—* FERUSAC, Guer. Mag., 1835, p. 26.—* CONRAD, Monog., XI, 1838, p. 98, pl. LIV, fig. 2.—* HANLEY, Test. Moll., 1842, p. 204; * Biv. Shells, 1843, p. 204, pl. XXII, fig. 25.—* CATLOW and REEVE, Conch. Nom., 1845, p. 55.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* CHENU, Ill. Conch. 1858, pl. XIV, figs. 1, 1a, 1b.—* KUSTER, Conch. Cab. Unio, 1861, p. 178, pl. LVI, fig. 4.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXII, fig. 372.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.

* *Margarita (Unio) angustatus* LEA, Syn., 1836, p. 35; 1838, p. 23.

* *Margaron (Unio) angustatus* LEA, Syn., 1852, p. 36; 1870, p. 57.

*† *Unio extensus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 31; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 68, pl. XII, fig. 49; * Obs., VI, 1868, p. 67, pl. XII, fig. 49.—* B. H. WRIGHT, Check List, 1888.

* *Margaron extensus* LEA, Syn., 1870, p. 60.

*† *Unio subcylindræus* LEA, Pr. Ac. N. Sci. Phila., III, 1873, p. 422; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 58, pl. XX, fig. 57; * Obs., XIII, 1874, p. 62, pl. XX, fig. 57.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* B. H. WRIGHT, Check List, 1888.

South Carolina to western Georgia.

† UNIO PERSTRIATUS Lea.

* *Unio perstriatus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 255, pl. XII, fig. 3; * Obs., V, 1852, p. 11, pl. XII, fig. 3.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCII, fig. 500.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 163.

* *Margaron (Unio) perstriatus* LEA, Syn., 1852, p. 36; 1870, p. 57.

*† *Unio gracilentus* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 85; * Jl. Ac. N. Sci. Phila., V, 1862, p. 58, pl. III, fig. 205; * Obs., VIII, 1862, p. 62, pl. III, fig. 205.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) gracilentus* LEA, Syn., 1870, p. 60.

*† *Unio perlatus* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 193; * Jl. Ac. N. Sci., VI, 1866, p. 15, pl. IV, fig. 13; * Obs., XI, 1867, p. 19, pl. IV, fig. 12.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) perlatus* LEA, Syn., 1870, p. 60.

North and South Carolina.

¹ A very singular shell, much cut away at the anterior base, and, perhaps, abnormal.

† UNIO NAVICULOIDES Lea.

* *Unio naviculoides* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 170; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 94, pl. xx, fig. 74; * Obs., VI, 1858, p. 94, pl. xx, fig. 74.—B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) naviculoides* LEA, Syn., 1870, p. 60.

Georgia.

† UNIO SORDIDUS Lea.

* *Unio sordidus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 254, pl. xii, fig. 1; * Obs., V, 1852, p. 10, pl. xii, fig. 1.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 1853, p. 257.—

* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

* *Margaron (Unio) sordidus* LEA, Syn., 1852, p. 33; 1870, p. 54.

*† *Unio gibbesianus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 254, pl. xii, fig. 2; * Obs., V, 1852, p. 10, pl. xii, fig. 2.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—

* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) gibbesianus* LEA, Syn., 1852, p. 33; 1870, p. 54.

Abbeville, South Carolina.

† UNIO SPADICEUS Lea.

* *Unio spadiceus* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 86; *Jl. Ac. N. Sci. Phila., V, 1862, p. 55, pl. i, fig. 202; * Obs., VIII, 1862, pl. i, fig. 201.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) spadiceus* LEA, Syn., 1870, p. 48.

North Carolina.

† UNIO STRIGOSUS Lea.

* *Unio strigosus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 287; *Tr. Am. Phil. Soc., VIII, 1843, p. 198, pl. ix, fig. 9; * Obs., III, 1842, p. 36 pl. ix, fig. 9.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857.—* CHENU, Ill. Conch., 1858, pl. xxx, figs. 6, 6a, 6b.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) strigosus* LEA, Syn., 1852, p. 36; 1870, p. 57.

Southwest Georgia; southeast Alabama.

† UNIO LAZARUS Lea.¹

* *Unio lazarus* LEA, Pr. Am. Phil. Soc., V, 1852, p. 251; *Tr. Am. Phil. Soc., XI, 1852, p. 259, pl. xiv, fig. 9; * Obs., V, 1852, p. 15, pl. xiv, fig. 9.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* B. H. WRIGHT, Check List, 1888.—PÆTEL, Conch. Sam., III, 1890, p. 157.

* *Margaron (Unio) lazarus* LEA, Syn., 1852, p. 39; 1870, p. 62.

Abbeville district, South Carolina.

(Group of *Unio downiei*.)

Shell subtrapezoidal, inflated, solid, truncate and more or less triangular behind, sometimes swollen in the postbasal region, with a well-

¹A peculiarly compressed, arcuate form. Some specimens of *U. arctatus* approach it closely.

developed posterior ridge; beaks full, their sculpture not observed; epidermis smooth and shining in the young shell, becoming duller and roughened when old; hinge moderately strong; pseudocardinals radial, roughened; laterals heavy, somewhat remote, curved; there is a more or less developed secondary lateral in the right valve; dorsal scars few, in a row just behind the pseudocardinals and fully exhibited on the inner edge of the shallow beak cavity; naere dull; muscle scars distinct, smooth.

Animal with the marsupium occupying the whole of the outer gills; inner gills free from the abdominal sac a part of their length; palpi rather small; mantle line thick on the border.

† UNIO DOWNIEI Lea.

* *Unio downiei* LEA, Pr. Ac. Nat. Sci. Phila., II, 1858, p. 166; Jl. Ac. Nat. Sci. Phila., IV, 1859, p. 210, pl. xxv, fig. 91; Obs., VII, 1859, p. 28, pl. xxv, fig. 91.—

* SOWERBY, Conch. Icon., XVI, 1868, pl. lxxviii, fig. 350.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 151.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 413, pl. lv, figs. 1-3; lvi, fig. 5.

* *Margaron (Unio) downiei* LEA, Syn., 1870, p. 41.

Southern Georgia; north Florida.

† UNIO GEMINUS Lea.

* *Unio geminus* LEA, Pr. Ac. Nat. Sci. Phila., VIII, 1856, p. 262; * Jl. Ac. Nat. Sci. Phila., 1858, p. 63, pl. x, fig. 45; * Obs., VI, 1858, p. 63, pl. x, fig. 45.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) geminus* LEA, Syn., 1870, p. 41.

*† *Unio satillaensis* LEA, Pr. Ac. Nat. Sci. Phila., II, 1858, p. 166; * Jl. Ac. Nat. Sci. Phila., IV, 1859, p. 216, pl. xxvii, fig. 96; Obs., VII, 1859, p. 34, pl. xxvii, fig. 96.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) satillaensis* LEA, Syn., 1870, p. 41.

Georgia.

† UNIO LECONTIANUS Lea.

* *Margarita (Unio) lecontianus* LEA, Syn., 1836, p. 23; 1838, p. 18.

* *Unio lecontianus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 40, pl. xii, fig. 35; * Obs., II, 1838, fig. 40, pl. xii, fig. 35.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 236.—* HANLEY, Test. Moll., 1842, p. 188; * Biv. Shells, 1843, p. 188, pl. xxii, figs. 11, 51.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 251.—* CHENU, Ill. Conch., 1858, pl. xxiv, figs. 6, 6a, 6b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* SOWERBY, Conch. Icon., XVI, 1866, pl. xxxiii, fig. 173.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 157.

* *Margaron (Unio) lecontianus* LEA, Syn., 1852, p. 26; 1870, p. 41.

* *Unio contrarius* CONRAD, Pr. Ac. Nat. Sci. Phila., IV, 1849, p. 153; * Ann. and Mag., IV, 1849, p. 301; * Jl. Ac. Nat. Sci. Phila., 1850, p. 276, pl. xxxvii, fig. 7; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.

Georgia.

†UNIO SPISSUS Lea,

**Unio spissus* LEA, Pr. Ac. Nat. Sci. Phila., III, 1859, p. 112; *Jl. Ac. Nat. Sci. Phila., IV, 1859, p. 208, pl. xxv, fig. 89; *Obs., VII, 1859, p. 26, pl. xxv, fig. 89.—

*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVIII, fig. 476.—*B. H. WRIGHT, Check List, 1888.—*PATEL, Conch. Sam., III, 1890, p. 168.

*Margaron (*Unio*) *spissus* LEA, Syn., 1870, p. 41.

Georgia.

(Group of *Unio fisherianus*.)

Shell elongated, rather thin, subcompressed, pointed behind and sometimes slightly biangulate; posterior ridge generally well developed; beaks low, the sculpture consisting of a few coarse, slightly irregular ridges which run nearly parallel with the growth lines, generally heavier where they cross the posterior ridge; epidermis sometimes rayed, shining; pseudocardinals usually compressed; laterals long, straight, and lamellar; beak cavities very shallow and containing two or three dorsal scars at some distance behind the beaks; muscle scars well marked, the posterior ones elongated; nacre generally dull. Animal with the gills greatly elongated, inner the larger, more or less free from the abdominal sac; marsupium occupying nearly or quite the whole length of the outer branchiæ; palpi elongated; mantle thin, thicker on the edge.

†UNIO LANCEOLATUS Lea.

**Unio lanceolatus* LEA, Tr. Am. Phil. Soc., 1828, p. 266, pl. III, fig. 2; *Obs., I, 1834, p. 8, pl. III, fig. 2.—*CONRAD, New F. W. Shells, 1834, p. 70.—*FERUSSAC, Guer. Mag., 1835, p. 26.—*CONRAD, Monog. III, 1836, p. 32, pl. xiv., fig. 2.—

*HANLEY, Test. Moll., 1842, p. 204; *Biv. Shells., 1843, p. 204, pl. xx, fig. 60; XXII, fig. 26.—*CATLOW and REEVE, Conch. Nom., 1845, p. 60.—*CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 251.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—*CHENU, Ill., Conch. 1858, pl. VIII, fig. 1, 1a, 1b.—*KUSTER, Conch. Cab. Unio., 1861, p. 204, pl. LXVIII, fig. 4.—*SOWERBY, Conch. Icon., XVI, 1866, pl. XLIII, fig. 236.—*B. H. WRIGHT, Check List, 1888.—*PATEL, Conch. Sam., III, 1890, p. 156.

**Margarita* (*Unio*) *lanceolatus* LEA, Syn., 1836, p. 35; 1838, p. 24.

**Margaron* (*Unio*) *lanceolatus* LEA, Syn., 1852, p. 36; 1870, p. 57.

**Unio lanceolata* DESHAYES, Enc. Méth., II, 1830, p. 585; *An. sans Vert., 2d ed., VI, 1835, p. 547; 3d ed., II, 1839, p. 672.

North Carolina; Virginia.

†UNIO VIRIDULUS Lea.¹

**Unio viridulus* LEA, Pr. Ac. Nat. Sci. Phila., VII, 1863, p. 193; *Jl. Ac. Nat. Sci. Phila., VI, 1866, p. 10, pl. III, fig. 17; *Obs., XI, 1867, p. 14, pl. III, fig. 17.—

*B. H. WRIGHT, Check List, 1888.

**Margaron* (*Unio*) *viridulus* LEA, Syn., 1870, p. 58.

Neuse River, near Raleigh, North Carolina.

¹ A young shell, no doubt, which in texture and appearance is much like *U. lanceolatus*, but is bright green and not so pointed behind.

† UNIO PRODUCTUS Conrad.¹

Unio productus CONRAD, Monog., III, 1836, p. 31, pl. XIV, fig. 1.—* HANLEY, Test. Moll., 1842, p. 205; * Biv. Shells, 1843, p. 205, pl. XXIII, fig. 17.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* KUSTER, Conch. Cab. Unio, 1852, p. 66, pl. XVI, fig. 2.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 255.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.

* *Margarita (Unio) productus* LEA, Syn., 1836, p. 37; 1838, p. 24.

* *Margaron (Unio) productus*, LEA, Syn., 1852, p. 37; 1870, p. 60.

* *Unio barrotti* KUSTER, Conch. Cab. Unio, 1861, p. 189, pl. LIX, fig. 6.

North Carolina; Virginia; Maryland.

† UNIO NASUTULUS Lea. (Emend.)

* *Unio nasutilus* ² LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 192.

* *Unio nasutilus* LEA, Jl. Ac. N. Sci. Phila., VI, 1866, p. 12, pl. III, fig. 9; * Obs., XI, 1867, p. 16, pl. III, fig. 9.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) nasutilus* LEA, Syn., 1870, p. 60.³

Livingston's Creek, Brunswick County, North Carolina.

† UNIO FISHERIANUS Lea.

* *Unio fisherianus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 8, pl. IV, fig. 8; * Obs., II, 1838, p. 8, pl. IV, fig. 8.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 234.—* HANLEY, Test. Moll., 1842, p. 206; * Biv. Shells, 1843, p. 206, pl. XXII, fig. 52.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* CHENU, Ill. Conch., 1858, pl. XX, figs. 4, 4a, 4b.—* KUSTER, Conch. Cab. Unio., 1861, p. 205, pl. LXVIII, fig. 6.—* REEVE, Conch. Icon., XVI, 1865, pl. XXIV, fig. 113.—* HARTMAN and MICHENER, Conch. Cest., 1874, p. 90, fig. 187.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 152.

* *Margarita (Unio) fisherianus* LEA, Syn., 1836, p. 37; 1838, p. 24.

* *Margaron (Unio) fisherianus* LEA, Syn., 1852, p. 37; 1870, p. 60.

* *Unio nasutus* CONRAD (part),⁴ Monog., II, 1838, pl. XVIII, fig. 1.

Virginia; Maryland; Pennsylvania, in the Atlantic drainage.

† UNIO EMMONSII Lea.

* *Unio emmonsii* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 86; * Jl. Ac. N. Sci. Phila., V, 1862, p. 56, pl. II, fig. 203; * Obs., VIII, 1862, p. 60, pl. II, fig. 203.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) emmonsii* LEA, Syn., 1870, p. 60.

North Carolina.

¹ I have seen a large number of specimens of what I suppose is this, none of which exactly agree with Conrad's figure and description, but which I can not refer to anything else.

² So written; no doubt a typographical error.

³ Probably only a variety of *U. productus* Conrad.

⁴ Conrad's first two figures (under fig. 1) are certainly *U. fisherianus*, the third is a female *nasutus*.

† UNIO DISPALANS B. H. Wright.

- * *Unio dispalans* B. H. WRIGHT, Nautilus, 1899, p. 50.

† UNIO SUBINFLATUS Conrad.

- * *Unio subinflatus* CONRAD, Monog., XI, 1838, p. 97, pl. LIV, fig. 1; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

- * *Margaron (Unio) subinflatus* LEA, Syn., 1870, p. 52.

South Georgia; Florida.

† UNIO AHENEUS Lea.

- * *Unio aheneus* LEA, Desc. 12 Uniones, 1843 (no paging); * Tr. Am. Phil. Soc., IX, 1846, p. 280, pl. XLI, fig. 9; * Obs., IV, 1848, p. 38, pl. XLI, fig. 9.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVI, fig. 194.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 431, pl. LXXII, fig. 6.

- * *Margaron (Unio) aheneus* LEA, Syn., 1852, p. 37; 1870, p. 60.

Florida.

† UNIO WALTONI B. H. Wright.

- * *Unio waltoni* B. H. WRIGHT, Pr. Ac. N. Sci. Phila., 1888, p. 114, pl. II, fig. 3; * Check List, 1888.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 431, pl. LXXIII, fig. 7.

Florida.

† UNIO ATTENUATUS Lea.

- * *Unio attenuatus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 157; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 41, pl. XIV, fig. 38; * Obs., XIII, 1874, p. 45, pl. XIV, fig. 38.—* B. H. WRIGHT, Check List, 1888.

Georgia.

† UNIO ROSTRÆFORMIS Lea.

- * *Unio rostræformis* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 262.

- * *Unio rostriformis* LEA, Jl. Ac. N. Sci. Phila., IV, 1858, p. 64, pl. x, fig. 46; * Obs., VI, 1858, p. 64, pl. x, fig. 46.—* B. H. WRIGHT, Check List, 1888.

- * *Margaron (Unio) rostriformis* LEA, Syn., 1870, p. 60.

- * † *Unio maconeusis* LEA, Pr. Ac. N. Sci. Phila., VIII, 1857, p. 172; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 93, pl. XX, fig. 73; * Obs., VI, 1858, p. 93, pl. XX, fig. 73.—* B. H. WRIGHT, Check List, 1888.

- * *Margaron (Unio) maconeusis* LEA, Syn., 1870, p. 60.

Georgia.

UNIO DUTTONIANUS Lea.

- * *Unio duttonianus* LEA, Pr. Ac. N. Sci. Phila., II, 1841, p. 31; * Jl. Ac. N. Sci. Phila., VIII, 1842, p. 236, pl. XXII, fig. 50; * Obs., III, 1842, p. 74, pl. XXII, fig. 50.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* CHENU, Ill. Conch., 1858, pl. XXXII, figs. 4, 4a, 4b.—SOWERBY, Conch. Icon., XVI, 1868, pl. xci, fig. 492.—* B. H. WRIGHT, Check List, 1888.

- * *Margaron (Unio) duttonianus* LEA, Syn., 1852, p. 35; 1870, p. 57.

- * *Unio duttonianus* PÆTEL, C. Sam., III, 1890, p. 151.

Ogeechee Canal, Savannah, Georgia.

† UNIO FOLLICULATUS Lea.

- * *Unio folliculatus* LEA, Tr. Am. Phil. Soc., VI, 1858, p. 38, pl. XI, fig. 33; * Obs., II, 1838, p. 33, pl. XI, fig. 33.—* TROSCHEL, Arch. für Nat., V, 1839, Pt. 2, p. 236.—* HANLEY, Test. Moll., 1842, p. 204; * Biv. Shells, 1843, p. 204, pl. XXII, fig. 55.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* CHENU, Ill. Conch., 1858, pl. XXIII, figs. 4, 4a, 4b.—* KUSTER, Conch. Cab. Unio, 1861, p. 202, pl. LXVII, fig. 4.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCI, fig. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 153.
- * *Margarita (Unio) folliculatus* LEA, Syn., 1836, p. 34; 1838, p. 23.
- * *Margaron (Unio) folliculatus* LEA, Syn., 1852, p. 35; 1870, p. 57.
- * † *Unio exacutus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 159; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 45, pl. xv, fig. 43; * Obs., XIII, 1874, p. 49, pl. xv, fig. 49.—* B. H. WRIGHT, Check List, 1888.
- * † *Unio rostellum* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 160; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 44, pl. xv, fig. 41; * Obs., XIII, 1874, p. 48, pl. xv, fig. 41.—* B. H. WRIGHT, Check List, 1888.

Georgia.

† UNIO SHEPARDIANUS Lea.

- * *Unio shepardianus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 95, pl. XIII, fig. 38; * Obs., I, 1834, p. 207, pl. XIII, fig. 38.—* CONRAD, Monog., VIII, 1837, p. 70, pl. XXXIX.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 203; * Biv. Shells, 1843, p. 203, pl. XXIII, fig. 25.—* KUSTER, Conch. Cab. Unio, 1852, p. 65, pl. XVI, fig. 1.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* CHENU, Man., 1859, II, p. 139, fig. 647.—* REEVE, Conch. Icon., XVI, 1865, pl. XIX, fig. 90.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 167.
- * *Margarita (Unio) shepardianus* LEA, Syn., 1836, p. 34; 1838, p. 23.
- * *Margaron (Unio) shepardianus* LEA, Syn., 1852, p. 35; 1870, p. 57.
- * *Unio sheppardianus* CATLOW and REEVE, Conch. Nom., 1845, p. 64.

Altamaha River, Georgia.

(Group of *Unio popeii*.)

Shell elongated, narrowed in front, and biangulate behind, with the base slightly sinuate, feebly sulcate; beaks not prominent, sculptured with rather fine, somewhat broken, often faint ridges, which have a tendency to fall into two rounded loops; pseudocardinals compressed, high, sharp, ragged; laterals long, slightly curved; cavity of the beaks shallow; cicatrices not deep. The shell is only a little thickened in front, and that of the female is slightly swollen at the posterior base.

Animal with the marsupium occupying the whole length of the outer gills; ovisacs not separated by a sulcus; gills long, inner a little the larger throughout, generally free their whole length from the abdominal sac; palpi enormous, wide, oval, slightly pointed behind, united two-thirds of their length to the mantle; mantle double on its edge; branchial opening large.

† UNIO POPEII Lea.

- * *Unio popeii* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 102; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 372, pl. LXVI, fig. 197; * Obs., VIII, 1860, p. 54, pl. LXVI, fig. 197.
 * *Unio popei* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXI, fig. 430.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.—
 * FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 605.
 * Margaron (*Unio*) *popeii* LEA, Syn., 1870, p. 57.

South Texas; northeast Mexico.

† UNIO POEYANUS Lea.

- * *Unio poeyanus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 85; * Obs., VI, 1857, p. 35, pl. XXXII, fig. 30; * Jl. Ac. N. Sci. Phila., III, 1858, p. 315, pl. XXXII, fig. 30.—
 * SOWERBY, Conch. Icon., XVI, 1868, pl. xc, fig. 486.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 606.
 * Margaron (*Unio*) *poeyanus* LEA, Syn., 1870, p. 57.

Rio de las Balsas, near Coyucan, Mexico.

UNIO VERÆ-PACIS Tristram.

- * *Unio veræ-pacis* TRISTRAM, Proc. Zool. Soc. Lond., 1863, Pt. 3, p. 414.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 171.—
 * FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 620.

Guatemala, Department of Vera Paz.

Section CANTHYRIA Swainson, 1840.

Shell inflated, suboval, spinose, with a high, rather sharp posterior ridge, above which it is somewhat truncated, the posterior slope being sometimes slightly wrinkled; beaks rather compressed, the sculpture not seen; epidermis smooth and shining, variegated with angular blotches; hinge sharply curved at the center; pseudocardinals rather compressed; laterals short, remote, the hinge plate narrowed; beak cavities rather deep. Animal with the marsupium occupying the whole of the outer gills, forming a smooth pad; branchiæ very large, round below, inner the larger, free nearly their whole length from the abdominal sac; palpi large; mantle double on its edge, sometimes with a few papillæ in front of the branchial opening; branchial opening small; supranal opening colored inside.²

- † *Unio spinosus* LEA, Desc. of New Sp. Unio, 1836, colored figs.; * Tr. Am. Phil. Soc., VI, 1838, p. 57, pl. XVI, fig. 50; * Obs., II, 1838, p. 57, pl. XVI, fig. 50.—* JAY, Cat., 1839, p. 113, pl. v, figs. 1, 2.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 236.—* HANTLEY, Test. Moll., 1842, p. 182; * Biv. Shells, 1843, p. 182, pl.

¹ Only a Latin description was given by Tristram, and the species is not identifiable. It is probably close to *U. popeii*, and may possibly be that.

² Notwithstanding the remarkable character of well-developed spines on the only species of this group, *U. spinosus*, the animal even when gravid does not present any striking peculiarities, and is certainly a *Unio*. I have seen one or two specimens of the *spinosus* with only one spine on one valve and none on the other, and one in the Lea collection without spines.

XXIII, fig. 56.—*CATLOW and REEVE, Conch. Nom., 1845, p. 64.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—*KUSTER, Conch. Cab. Unio, 1856, p. 167, pl. XLIX, fig. 1.—*CHENU, Ill. Conch., 1858, pl. XXII, figs. 6, 6a, 6b.—*SOWERBY, Conch. Icon., XVI, 1866, pl. XLIX, fig. 261.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 168.

**Margarita (Unio) spinosus* LEA, Syn., 1836, p. 17, colored plate; 1838, p. 16.

**Margaron (Unio) spinosus* LEA, Syn., 1852, p. 23; 1870, p. 35.

**Canthyria spinosa* SWAINSON, Tr. on Mal., 1840, pp. 276, 378.—*AGASSIZ, Arch. für Nat., I, 1852, p. 44.

**Unio spinosa* GOODRICH, Ill. Nat. Hist., II, 1859, p. 523, fig.

Altamaha River, Georgia.

Section **UNIOMERUS** Conrad, 1853.

(Type, *Unio tetralasmus* Say.)

Shell trapezoid, with a rounded posterior ridge, and pointed or feebly biangulate behind; beaks not prominent, sculptured with 10 to 15 curved, rather strong, concentric ridges, which sweep decidedly upward behind, where they are drawn somewhat closely together; epidermis generally rayless, often clothlike; pseudocardinals usually compressed; laterals delicate, slightly curved; muscle scars large, shallow, nacre generally lurid.

Animal with the marsupium occupying the whole length of the outer branchiæ, pad-like; gills large, inner the larger, free nearly or quite the length of the abdominal sac; mantle generally rather thick, thicker and double on the edge.

† **UNIO TETRALASMUS** Say.

**Unio tetralasmus* SAY, Am. Conch., III, 1830, pl. XXIII.¹—*FERUSSAC, Guer. Mag., 1835, p. 26.—*DESHAYES, An. sans. Vert., 2d ed., VI, 1835, p. 555; 3d ed., II, 1839, p. 674.—*HANLEY, Test. Moll., 1842, p. 198; *Biv. Shells, 1843, p. 198, pl. XXIII, fig. 49.—*CATLOW and REEVE, Conch. Nom., 1845, p. 64.—*CHENU, Bib. Conch., 1st ser., III, 1845, p. 52, pl. x, figs. 1-3.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 169.

¹ An abundant, widespread, and very variable species, and if there were no connecting links it would be easy to make a half dozen species out of it. The first described is the *Unio tetralasmus* Say, an elongated, rather smooth, more or less shining form, of which *excultus*, *jamesianus*, *parallelus*, *suberoceus*, and *symmetricus* seem to be synonymous, varying a little in form, color, and smoothness. The variety *campodon* of Say is quite rhomboid, and is a rather rough, dark shell, and *U. symmetricus* is practically synonymous. Variety *declivis* is peculiarly drawn out at the post basal point. Variety *manubius* is a large, rather smooth form, rounded on the basal line. All these are Southern and Southwestern forms. Variety *sayi* is a yellowish shell, with very distinct rest marks, and has a more northern distribution than the rest, but it absolutely blends into them. Ward sent a description of his *U. sayi* to Tappan, who published it in the American Journal of Science and Arts under date of 1839. In Part XI of the Monography, dated November, 1838, Conrad describes *Unio sayanus* (changed to *sayi* in his notes) and refers it to Ward in the American Journal of Science and Arts of 1839. There is some mistake in these dates. In giving the synonymy it is hard to tell in all cases just what authors have meant.

- **Margarita (Unio) tetralasmus* LEA, Syn., 1836, p. 30; 1838, p. 21.
 **Margaron (Unio) tetralasmus* LEA, Syn., 1852, p. 32; 1870, p. 50.
 **Unio excultus* CONRAD, Monog., XI, 1838, p. 99, pl. LIV, fig. 1; Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—* PÆTEL, Conch. Sam., III, 1890, p. 152.
 *†*Unio parallelus* CONRAD, Pr. Ac. N. Sci. Phila., I, 1841, p. 20; * Jl. Ac. Phila., VIII, 1842, p. 179.¹
 *†*Unio symmetricus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; * Tr. Am. Phil. Soc., X, 1853, p. 73, pl. IV, fig. 11; * Obs., IV, 1848, p. 47, pl. IV, fig. 11.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 169.
 **Margaron (Unio) symmetricus* LEA, Syn., 1852, p. 32; 1870, p. 52.
 *†*Unio porrectus* CONRAD, Jl. Ac. N. Sci. Phila., 1854, p. 296, pl. XXVI, fig. 7.—* B. H. WRIGHT, Check List, 1888.
 *†*Unio subcoccus* CONRAD, Jl. Ac. N. Sci. Phila., VII, 1854, p. 297, pl. XXVII, fig. 1.
 **Unio jamesianus* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 84; * Jl. Ac. N. Sci. Phila., IV, 1853, p. 53, pl. VI, fig. 35; * Obs., VI, 1858, p. 52, pl. VI, fig. 35.—* B. H. WRIGHT, Check List, 1888.
Margaron (Unio) jamesianus LEA, Syn., 1870, p. 50.

† UNIO TETRALASMUS var. CAMPTODON Say.

- **Unio camptodon* SAY, Am. Conch., V, 1832, pl. XIII.—* CONRAD, New F. W. Shells, 1834, p. 68.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* HANLEY, Test. Moll., 1842, p. 198; * Biv. Shells, 1843, p. 198, pl. XXI, fig. 17.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 53, pl. XIV, figs. 2, 2a, 2b.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 246.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXX, fig. 356.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 147.—* STEARNS, Pr. U. S. Nat. Mus., XIV, 1891, p. 105.
 **Margarita (Unio) camptodon* LEA, Syn., 1836, p. 30; 1838, p. 21.
 **Margaron (Unio) camptodon* LEA, Syn., 1852, p. 32; 1870, p. 50.
 *†*Unio geometricus* LEA, Tr. Am. Phil. Soc., V, 1832, p. 38, pl. IV, fig. 10; * Obs., I, 1834, p. 150, pl. IV, fig. 10.

† UNIO TETRALASMUS var. MANUBIUS Gould.

- **Unio manubius* GOULD, Pr. Bost. Soc. Nat. Hist., V, 1856, p. 229; * Otia Conch., 1862, p. 218.—* B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 592.
 **Margaron (Unio) manubius* GOULD, Syn., 1870, p. 54.
 **Unio manubius* PÆTEL, Conch. Sam., III, 1890, p. 158.

† UNIO TETRALASMUS var. DECLIVIS Say.

- Unio declivis* SAY, Transylvania Jl. IV, 1831, p. 527; * Am. Conch., III, 1832, pl. XXXV.—* CONRAD, New F. W. Shells, 1834, p. 68.—* FERUSSAC, Guer. Mag., 1835, p. 26.—* DESHAYES, An. saps. Vert., 2d ed., VI, 1835, p. 556; 3d ed., II, 1839, p. 675.—* HANLEY, Test. Moll., 1842, p. 200; * Biv. Shells, 1843, p. 200, pl. XXIII, fig. 50.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 46, pl. XIII, figs. 1, 1a, 1b.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* KUSTER, Conch. Cab. Unio, 1852, p. 60, pl. XIV, fig. 1.—* CONRAD, Pr. Ac. N. Sci.

¹ This name had been used by Sowerby for a fossil *Unio*, and Conrad subsequently changed it to *porrectus* on that account, but Sowerby had also used that name for a *Unio*.

Phila., VI, 1853, p. 248.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 150.

**Margarita (Unio) declivis* LEA, Syn., 1836, p. 32; 1838, p. 22.

**Margaron (Unio) declivis* LEA, Syn., 1852, p. 33; 1870, p. 54.

UNIO TETRALASMUS var. SAYI Ward.

**Unio sayi* WARD (in Tappan), Am. Jl. Sci., XXXV, 1839, p. 268, pl. III, fig. 1.—

*CONRAD, Monog., XI, 1838?, p. 102, pl. LV, fig. 2.¹—*KUSTER, Conch. Cab. Unio, 1861, p. 246, pl. LXXXIII, fig. 1.—*PÆTEL, Conch. Sam., III, 1890, p. 166.

**Unio sayanus* B. H. WRIGHT, Check List, 1888.

**Unio electrinus* REEVE, Conch. Icon., XVI, 1865, pl. XXV, fig. 121.—*PÆTEL, Conch. Sam., III, 1890, p. 151.

Lower Mississippi drainage north to about latitude 40°; Ohio River system; Alabama River system and southwest through Texas into northern Mexico. Not reported, so far as I know, from the Tennessee and Cumberland rivers.

†UNIO COLUMBENSIS Lea.²

**Unio columbensis* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 31; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 75, pl. XIV, fig. 55; *Obs., VI, 1858, p. 75, pl. XIV, fig. 55.—

*B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) columbensis* LEA, Syn., 1870, p. 50.

**Unio columbiensis* PÆTEL, Conch. Sam., III, 1890, p. 148.

Chattahoochee River, Georgia, west to Little Patsaliga Creek, Alabama.

†UNIO OBESUS Lea.

*?*Unio carolinensis* Bosc,³ Hist. Nat. de Coq., 1824, III, p. 139, pl. XXIII, fig. 2.—

*FERUSSAC, Guer. Mag., 1835, p. 26.—*CHENU, Ill. Conch., 1843, pl. XVIII, fig. 4.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 246.

*†*Unio obesus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 96, pl. XIII, fig. 26; *Obs., I, 1834, p. 106, pl. XIII, fig. 26.—*CONRAD, N. F. W. Shells, 1834, p. 70.—*HAN-

LEY, Test. Moll., 1842, p. 198; *Biv. Shells, 1843, p. 198, pl. XXII, fig. 34.—

*CATLOW and REEVE, Conch. Nom., 1845, p. 61.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—*CHENU, Ill. Conch., 1853, pl. XVIII, figs. 4, 4a, 4b.—

*KUSTER, Conch. Cab. Unio, 1861, p. 196, pl. LXIII, fig. 2.—*REEVE, Conch. Icon., XVI, 1864, pl. XVIII, fig. 84.—*SOWERBY, Conch. Icon., XVI, 1866,

pl. XXXVIII, fig. 212.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 161.—*SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 428, pl.

LXVIII, fig. 6; LXIX, figs. 1, 2, 4; LXXI, fig. 3.

**Margarita (Unio) obesus* LEA, Syn., 1836, p. 30; 1838, p. 21.

**Margaron (Unio) obesus* LEA, Syn., 1852, p. 32; 1870, p. 50.

¹ *Unio sayanus* on plate, *U. sayi* in note.

² This species almost merges into *obesus* on the one hand and the southern forms of *tetralasmus* on the other. The distribution of *obesus* is eastern, that of *tetralasmus* western, while *columbensis* occupies a central position in streams of the Gulf drainage.

³ It is impossible to tell from the very poor figure what this is. Lea thinks it is *Margaritana margaritifera*, but Bosc did not visit any region where that species is found, and the figure seems to show lateral teeth. It is quite probably *U. obesus*. I think Lea's name had better be used.

- * *Unio declivis* CONRAD, Monog., V, 1836, p. 45, pl. XXIII, fig. 1.
- * † *Unio ineptus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 261, pl. XV, fig. 12; * Obs., V, 1852, p. 17, pl. XV, fig. 12.—* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
- * *Margaron (Unio) ineptus* LEA, Syn., 1852, p. 31; 1870, p. 50.
- * † *Unio hebes* LEA, Tr. Am. Phil. Soc., X, 1852, p. 267, pl. XVIII, fig. 21; * Obs., V, 1852, p. 23, pl. XVIII, fig. 21.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 154.
- * *Margaron (Unio) hebes* LEA, Syn., 1852, p. 26; 1870, p. 51.
- * † *Unio rivularis* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* CONRAD, Jl. Ac. N. Sci. Phila., 1854, p. 296.
- * † *Unio cieur* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 39; * Jl. Ac. N. Sci. Phila., V, 1862, p. 93, pl. XIII, fig. 241.—* Obs., VIII, 1862, p. 96, pl. XIII, fig. 241.—B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) cieur* LEA, Syn., 1870, p. 52.
- * *Unio squalidus* LEA, (part) ¹ Pr. Ac. N. Sci. Phila., XV, 1863, p. 192.—* Jl. Ac. N. Sci. Phila., VI, 1866, p. 22, pl. VII, fig. 20; * Obs., XI, 1867, p. 26, pl. VII, fig. 20.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) squalidus* LEA, Syn., 1870, p. 51.
- * † *Unio jewettii* LEA, Pr. Ac. N. Sci. Phila., XI, 1867, p. 81; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 276, pl. XXXVII, fig. 89; * Obs., XII, 1869, p. 36, pl. XXVII, fig. 89.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) jewettii* LEA, Syn., 1870, p. 51.
- * † *Unio paucensis* LEA,² Pr. Ac. N. Sci. Phila., XII, 1868, p. 161; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 302, pl. XLV, fig. 114; * Obs., XII, 1869, p. 62, pl. XLV, fig. 114.—* B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) paucensis* LEA, Syn., 1870, p. 51.
- * † *Unio rivicolus* CONRAD, Am. Jl. Conch., IV, 1868, p. 280, pl. XVIII, fig. 4.—* B. H. WRIGHT, Check List, 1888.

* † **UNIO OBESUS var. BLANDINGIANUS** Lea.

- * *Unio blandingianus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 101, pl. XV, fig. 44;³ * Obs., I, 1834, p. 213, pl. XV, fig. 44.—* FERUSSAC, Gner. Mag., 1835, p. 29.—* CONRAD, Monog., V, 1836, p. 46, pl. XXIII, fig. 2.—* HANLEY, Test. Moll., 1842, p. 200; * Biv. Shells, 1843, p. 200, pl. XXII, fig. 1.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* KUSTER, Conch. Cab. Unio, 1852, p. 36, pl. VI, fig. 2.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXV, fig. 187.—* B. H. WRIGHT, Check List, 1888.—* S. H. WRIGHT, Conch. Ex., II, 1888, p. 95.—* PÆTEL, Conch. Sam., III, 1890, p. 146.
- * *Margarita (Unio) blandingianus* LEA, Syn., 1836, p. 32; 1838, p. 22.
- * *Margaron (Unio) blandingianus* LEA, Syn., 1852, p. 33; 1870, p. 54.

¹ Under this name Dr. Lea has certainly described two species, a heavy quadrate, form biangulate behind, with purple nacre—a variety of *complanatus*—and another which is, no doubt, *U. obesus*, and is the shell figured.

² A rough, apparently diseased *obesus*.

³ A decidedly trapezoidal form with cloth-like, dark epidermis, possibly worthy of a varietal name. The transition to the typical state is, however, very complete.

* † *UNIO OBESUS* var. *PALUDICOLUS* Gould.

* *Unio paludicolus* GOULD, Pr. Bost. Soc. Nat. Hist., II, 1845, p. 53.—* GOULD, Otia. Conch., 1852, p. 197.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) paludicolus* LEA, Syn., 1852, p. 33; 1870, p. 54.

* *Unio paludicolor* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

Southern Virginia southward in streams draining into the Atlantic, and throughout the peninsula of Florida.

† *UNIO BISSELLIANUS* Lea.

* *Unio bisseillianus* LEA, Pr. Ac. N. Sci. Phila., XI, 1867, p. 81; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 277, pl. XXXVII, fig. 90; * Obs., XII, 1869, p. 37, pl. XXXVII, fig. 90.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

* *Margaron (Unio) bisseillianus* LEA, Syn., 1890, p. 50.

Bissell's Pond, Charlotte, North Carolina.

† *UNIO SUBLURIDUS* Simpson.

* *Unio subluridus* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 432, pl. LXXIII, figs. 3, 4.

Orange Springs, Marion County, Florida. I can not be certain as to the relationship of this species, as the beaks are eroded in the only shells I have seen.

Section *MICRONAIAS* Simpson, 1900.

(Type, *Unio aratus* Lea.)

Shell small, oval, a little produced near the posterior base, with a moderate posterior ridge and pointed near the base behind; beaks rather prominent, the sculpture being apparently rather fine, irregularly concentric ridges, having a tendency to fall into two loops; the whole surface strongly and closely concentrically ridged; pseudocardinals compressed, high, slightly curved upward; laterals compressed, curved; beak cavities moderately deep; nacre whitish; anterior muscle scars deep, rough; posterior well marked.

Animal with the marsupium pad-like, occupying all but the extreme posterior part of the outer gills; branchiæ elongated, wider behind, inner the larger throughout, free from the abdominal sac or united to it;¹ palpi large, rounded behind; mantle with thickened border; anal opening crenulate.

¹ In some specimens free throughout; in others entirely united.

† UNIO ARATUS Lea.

* *Unio aratus* LEA, Disc. 12 Uniones, 1843; *Tr. Am. Phil. Soc., IX, 1845, p. 282, pl. XLII, fig. 12; *Obs., IV, 1848, p. 40, pl. XLII, fig. 12.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIV, fig. 320.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 144.

* Margarou (*Unio*) *aratus* LEA, Syn., 1852, p. 28; 1870, p. 35.

Central America.

† UNIO GRANADENSIS Lea.

* *Unio granadensis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 95; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 293, pl. XLII, fig. 103; *Obs., XII, 1869, p. 53, pl. XLII, fig. 103.—* B. H. WRIGHT, Check List, 1888.

* Margarou (*Unio*) *granadensis* LEA, Syn., 1870, p. 35.

Lake Nicaragua.

The following species are of Rafinesque, but I am utterly unable to make anything out of them:

Unio dilatata, elliptica, fasciata, fragilis, levigata, nervosa, viridis, zonata, all in Monograph, 1820; also *Unio bicolor, calendis, castaneus, chloris, diaphanus, fontinalis, fulvus, lasmabrachys, montanus, pallens, platiolus, rivularis, rimosus, teneltus, venus*, all in Continuation of Monograph, 1831.

Other indeterminate *Unionidæ* of Rafinesque are: *Amblema antrosa, costata, gibbosa, olivaria, rubra, torulosa; Lasmonos fragilis; Obovaria obovalis, pachostea; Obliquaria triangularis, attenuata, atrovioleacea, bullata, cliffordiana, cuprea, cyphia, ellipsaria, fasciolaris, flava, flexuosa, lateralis, lincolata, nodulata, obliquata, pusilla, quadrula, sintoxia, sinuata, subrotunda*, all in Monograph, 1820. *Bariosta diploderma, ponderosus, vittatis; Epilobasma biloba; Toxolasma cinerescens, cyclips, flexus*, in Continuation of Monograph, 1831.

The following are also indeterminate and spurious Unios:

Unio tahetianus CATLOW and REEVE, Conch. Nom., 1845, p. 64. Is probably *U. taitianus* Lea.

Unio largillierti PHILIPPI, Menke's Zeits., 1847, p. 94.

Unio musicus SPENGLER, Skriv. Nat. Selsk., III, 1793, p. 67.

Unio radula SAY, N. Harm. Diss., 1829, p. 323.

Unio plombarius VILLA, Disp. Conch. Terr. Fluv., 1841, p. 62.

Unio pequotianus LINSLEY, Am. Jl. Sci., 1845, p. 277.

Unio iridescens CONRAD, Cov. of Monog. No. 11.

Unio angusta LAMARCK, An. sans. Vert., VI, 1819, p. 80.

Unio conus SPENGLER, Skriv. Nat. Selsk., 1792.

Unio delphinus SPENGLER, Skriv. Nat. Selsk., 1793, p. 63.

Unio doumeti LETOURNEUX and BOURGNIGNAT, Prod. Mal. Tunis., p. 163.

Unio distortus BEAN, Ann. and Mag., 1836, p. 376, fig. 53.

Unio uber CONRAD, Am. Jl. Conch., II, p. 279.

Mya oborata SOLANDER, Portland Cat., p. 100.

Genus PLEUROBEMA (Rafinesque, 1820) Agassiz.

(Type, *Unio clava* Lamarck.¹)*Pleurobema* RAFINESQUE, Ann. Gen. Sci. Phys., Brux, 1820, p. 313.

Shell solid, triangular to rhomboid, usually with a prominent umbonal region; beaks at or near the anterior end of the shell, incurved and pointed forward over a small but well developed lunule; beak sculpture coarse, consisting of a few irregular, often broken ridges, which curve upward posteriorly; posterior ridge present, but low and rounded; epidermis showing the rest periods plainly, tawny to olive, often ornamented with rays which show a tendency to break into square spots; hinge rather strong, the plate generally narrow; pseudocardinals triangular, ragged; laterals reaching nearly or quite to the pseudocardinals, double in both valves, in the right valve the inner being smaller; muscle scars deep, the posterior rounded; cavity of the beaks shallow; nacre silvery; male and female shells essentially alike.

Animal having the inner gills much the larger, rounded below, free from the abdominal sac for a part or all of their length; marsupium occupying the entire outer gills, the ovisacs in some cases seeming to be arranged in pairs; animal generally yellowish to salmon red, sometimes more or less brown or blackish.

(Group of *Pleurobema clava*.)

Shell solid, triangular; beaks high, generally anterior; beak sculpture consisting of three or four broken, coarse, irregular ridges; epidermis yellowish or tawny, marked with broken green rays which show a tendency to form square spots; pseudocardinals often somewhat lengthened and more or less parallel with the laterals.

Animal having the inner gills the larger except at the extreme posterior end, free nearly or quite their whole length from the abdominal sac; marsupium occupying the entire outer gills; branchial opening rather large, with small papillæ; anal opening with minute papillæ or crenulations. Animal dirty whitish to salmon.

† PLEUROBEMA CLAVA Lamarck.

* *Unio clava* LAMARCK, An. sans Vert., VI, 1819, p. 74.—* DESHAYES, An. sans Vert., 2ded., VI, 1835, p. 537; 3ded., II, 1839, p. 669.—* CONRAD, Monog., I, 1835, p. 5, pl. III, fig. 1.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* CONRAD, Pr. Ac.

¹ I regret that I have not been able to examine more of the animals of this group, especially those of the gravid females. The shells on the one hand approach very close to those of *Quadrula* in appearance, and to *Unio* on the other hand. But all of them have shallow beak cavities, while those of *Quadrula* are deep, and the embryos of *Pleurobema* are contained in the outer gills only. The shells are generally more solid and more triangular than those of *Unio*, and the pattern of coloring is different from that of either genus. It stands between the two genera. I place *Unio æsopus* and *varicosus* in *Pleurobema* with some hesitation, though in a specimen of the former containing comparatively few embryos there were none in the inner gills, and their beak cavities are shallow.

N. Sci. Phila., VI, 1853, p. 246.—* KUSTER, Conch. Cab. Unio, 1852, p. 39, pl. VII, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIX, fig. 354.

* *Pleurobema clara* AGASSIZ, Arch. für Naturg., I, 1852, p. 49.

* *Unio clarus* REEVE, Conch. Syst., I, 1841, p. 117, pl. LXXXVIII, fig. 3.—* HANLEY, Test. Moll., 1842, p. 187; * Biv. Shells, 1843, p. 187, pl. XX, fig. 52.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 148.—* CALL, Tr. Acad. Sci. St. Louis, VII, 1895, p. 9. pl. I.

* *Margarita (Unio) clavus* LEA, Syn., 1836, p. 22; 1838, p. 18.

* *Margaron (Unio) clavus* LEA, Syn., 1852, p. 26; 1870, p. 40.

*† *Unio patulus* LEA,¹ Tr. Am. Phil. Soc., III, 1829, p. 44, pl. XII, fig. 20; * Obs., I, 1834, p. 55, pl. XII, fig. 20.—* CONRAD, Monog., X, 1838, p. 92, pl. I, fig. 2.—* HANLEY, Test. Moll., 1842, p. 187; * Biv. Shells, 1843, p. 187, pl. XXII, fig. 27.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XVI, figs. 6, 6a, 6b.—* KUSTER, Conch. Cab. Unio, 1861, p. 259, pl. LXXXVII, fig. 5.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Margarita (Unio) patulus* LEA, Syn., 1836, p. 22; 1838, p. 18.

* *Margaron (Unio) patulus* LEA, Syn., 1852, p. 26; 1870, p. 40.

* *Unio cuneatus* SAY, Am. Conch., VI, 1834.—* FERUSSAC, Guer. Mag., 1835, p. 28.

Ohio, Cumberland, and Tennessee river systems. Reported from Iowa City, Iowa; St. Peter's River, Minnesota, and from Nebraska.

† PLEUROBEMA MACULATA Conrad.

* *Unio maculatus* CONRAD, New F. W. Shells, 1834, p. 30, pl. IV, fig. 4, p. 70.—

* FERUSSAC, Guer. Mag., 1835, p. 29.—MÖLLER, Syn. Nov. Gen., 1836, p. 202.—

* HANLEY, Test. Moll., 1842, p. 203; * Biv. Shells, 1843, p. 203.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* CHENU, Bib. Conch., 1st ser., III, 1845,

p. 17, pl. III, fig. 7.—* CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 252.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* KUSTER, Conch. Cab. Unio, 1861, p. 216, pl. LXXII, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

* *Margarita (Unio) maculatus* LEA, Syn., 1836, p. 34; 1838, p. 23.

* *Margaron (Unio) maculatus* LEA, Syn., 1852, p. 24; 1870, p. 38.²

Tennessee River system.

† PLEUROBEMA HOLSTONENSIS Lea.³

* *Unio holstonensis* LEA, Pr. Am. Phil. Soc., I, 1840, p. 288; * Tr. Am. Phil. Soc., VIII, 1842, p. 212, pl. XV, fig. 27; * Obs., III, 1842, p. 50, pl. XV, fig. 27.—* HAN-

¹No specimen in the Lea collection exactly agrees with the figure of this species and Lea gives no exact locality for the type; but from a careful study of his shells and the figure and description I am convinced that it is a somewhat compressed form of the very variable *clava*, which has rather low beaks, that are not placed as far forward as usual. There seems to be a complete set of intermediates.

²Dr. Lea is inclined to believe (see above citation) that Conrad's *Unio maculatus* is the same as his own *ravenelianus*, and if this were the case Lea's name should have precedence. From an examination of specimens in the Philadelphia Academy of Natural Sciences and Conrad's figure and description, I am forced to differ from Dr. Lea, as I consider the species close to Lamarck's *clavus*.

³The type, a young shell, is certainly the same as *Unio lawii*, and there does not seem to be any material difference between it and the types of *U. bellulus* and *mundus*. The figure is hardly accurate.

- LEY, Test. Moll., 1842, p. 213.—*CONRAD, Pr. Ac. N. Sci., Phila., VI, 1853, p. 250.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU, Ill. Conch., 1858, pl. XXXII, figs. 1, 1a, 1b.—*KUSTER, Conch. Cab. Unio, 1862, p. 287, pl. xcvi, fig. 4.—*B. H. WRIGHT, Check List, 1888.
- **Margaron (Unio) holstonensis* LEA, Syn., 1852, p. 25; 1870, p. 38.
- **Unio mundus* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 83; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 40, pl. XIV, fig. 38; *Obs., XI, 1867, p. 44, pl. XIV, fig. 38.—*REEVE, Conch. Icon., XVI, 1864, pl. XVI, fig. 72.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 160.
- **Margaron (Unio) mundus* LEA, Syn., 1870, p. 40.
- † **Unio mundus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 189; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 8, pl. II, fig. 4; *Obs., XIII, 1874, p. 12, pl. II, fig. 4.—*B. H. WRIGHT, Check List, 1888.
- † **Unio pattinoides* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 193; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 16, pl. IV, fig. 12; *Obs., XIII, 1874, p. 20, pl. IV, fig. 12.—*B. H. WRIGHT, Check List, 1888.
- † **Unio bellulus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 161; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 50, pl. XVII, fig. 48; *Obs., XIII, 1874, p. 54, pl. XVII, fig. 48.—B. H. WRIGHT, Check List, 1888.

Tennessee River system.

† PLEUROBEMA BOURNIANUS LEA.

- **Unio bournianus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 288; *Tr. Am. Phil. Soc., VIII, 1842, p. 213, pl. XV, fig. 28; *Obs., III, p. 51, pl. XV, fig. 28.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU, Ill. Conch., 1858, pl. XXVIII, figs. 2, 2a, 2b.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 146.
- **Maragon (Unio) bournianus* LEA, Syn., 1852, p. 25; 1870, p. 39.

Sciota River, near Chillicothe, Ohio.

† PLEUROBEMA EDGARIANUS LEA.

- **Unio edgarianus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 288; *Tr. Am. Phil. Soc., VIII, 1842, p. 214, pl. XV, fig. 30; Obs., III, 1842, p. 52, pl. XV, fig. 30.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU, Ill. Conch., 1858, pl. XXIX, figs. 5, 5a, 5b.—*KUSTER, Conch. Cab., 1861, p. 213, pl. LXX, fig. 5.—*MUSGRAVE, Phot. Conch., 1863, pl. II, fig. 6.—*REEVE, Conch. Icon., XVI, 1864, pl. XV, fig. 65.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 151.
- **Margaron (Unio) edgarianus* LEA, Syn., 1852, p. 25; 1870, p. 39.
- † **Unio obuncus* LEA, Pr. Ac. N. Sci., I, 1871, p. 192; *Jl. Ac. N. Sci., VIII, 1874, p. 9, pl. II, fig. 5; *Obs., XIII, 1874, p. 13, pl. II, fig. 5.—*B. H. WRIGHT, Check List, 1888.
- † **Unio andersonensis* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 155; *Jl. Ac., N. Sci. Phila., VIII, 1874, p. 36, pl. XII, fig. 33; *Obs., XIII, 1874, p. 40, pl. XII, fig. 33.—*B. H. WRIGHT, Check List, 1888.

Tennessee River system.

¹ In the errata Reeve says this is not *mundus*, but that it is nearer *cuneolus* and *mooresianus*. I have compared it with authentic specimens, and it is certainly nearer *mundus* than either of the other mentioned species.

† PLEUROBEMA CUNEOLUS Lea.

* *Unio cuneolus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 286; * Tr. Am. Phil. Soc., VIII, 1842, p. 193, pl. VII, fig. 3; * Obs., III, 1842, p. 31, pl. VII, fig. 3.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XXX, figs. 2, 2a, 2b.—* KUSTER, Conch. Cab. Unio, 1861, p. 182, pl. LXVII, fig. 4.—* REEVE, Conch. Icon., XVI, 1865, pl. XXIII, fig. 107.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 149.

* *Margaron (Unio) cuneolus* LEA, Syn., 1852, p. 24; 1870, p. 38.

†* *Unio clinchensis* LEA, Pr. Ac. N. Sci. Phila., XI, 1867, p. 81; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 278, pl. XXXVII, fig. 91; * Obs., XII, 1869, p. 38, pl. XXXVII, fig. 91.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Margaron (Unio) clinchensis* LEA, Syn., 1870, p. 38.

Tennessee River system.

† PLEUROBEMA LESLEYI Lea.

* *Unio lesleyi* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 306; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 352, pl. LVIII, fig. 177; * Obs., VIII, 1860, p. 34, pl. LVIII, fig. 177.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) lesleyi* LEA, Syn., 1870, p. 40.

Kentucky; Tennessee.

† PLEUROBEMA RAVENELIANA Lea.

* *Unio ravenelianus* LEA,¹ Tr. Am. Phil. Soc., V, 1834, p. 32, pl. III, fig. 5; * Obs., I, 1834, p. 144, pl. III, fig. 5.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 187; * Biv. Shells, 1843, p. 187, pl. XX, fig. 59.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* KUSTER, Conch. Cab. Unio, 1862, p. 269, pl. XCI, fig. 1.—* REEVE, Conch. Icon., XVI, 1864, pl. XVI, fig. 70.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) ravenelianus* LEA, Syn., 1836, p. 22; 1838, p. 18.

* *Margaron (Unio) ravenelianus*, LEA, Syn., 1852, p. 26; 1870, p. 40.

* *Unio rudis* CONRAD, Monog., IX, 1837, p. 76, pl. XLIII, fig. 1.—* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 257.

* *Unio decisus* KUSTER, (part) Conch. Cab. Unio, 1852, p. 41, pl. VIII, fig. 1.

Kentucky; Tennessee; western North Carolina.

† PLEUROBEMA OVIFORMIS Conrad.

* *Unio oviformis* CONRAD, New F. W. Shells, 1834, p. 46, pl. III, fig. 6; p. 70.—

* FERUSSAC, Guer. Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen., 1836, p. 208.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 22, pl. II, fig. 7.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Margaron (Unio) oviformis* LEA, Syn., 1852, p. 26; 1870, p. 40.

Tennessee.

¹ In May, 1834, Conrad published a *Unio raveneli* in New Fresh Water Shells (p. 39) which belongs to the *complanatus* group. Lea's name, *ravenelianus*, appeared in August or September of the same year. Conrad, claiming priority for his name, substituted that of *Unio rudis* for Lea's species. As the names are really different, that of Lea had better stand.

†PLEUROBEMA ACUENS Lea.

- * *Unio acueus* LEA, Pr. Ac. Nat. Sci. Phila., I, 1871, p. 190; * Jl. Ac. Nat. Sci. Phila., VIII, 1874, p. 27, pl. VIII, fig. 24; * Obs., XIII, 1874, p. 31, pl. VIII, fig. 24.—
* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 143.

Tennessee River system.

†PLEUROBEMA ORNATA Lea.

- * *Unio ornatus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 4; * Jl. Ac. N. Sci. Phila., V, 1862, p. 85, pl. XI, fig. 234; * Obs., VIII, 1862, p. 89, pl. XI, fig. 234.—
* PÆTEL, Conch. Sam., III, 1890, p. 162.
* Margaron (*Unio*) *ornatus* LEA, Syn., 1870, p. 57.¹

Alabama.

†PLEUROBEMA APPRESSA Lea.

- * *Unio appressus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 189; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 12, pl. III, fig. 8.—* Obs., III, 1874, pl. III, fig. 8.—* B. H. WRIGHT, Check List, 1888.
* *Unio argenteus* SOWERBY, Conch. Icon., XVI, 1866, pl. xxxvii, fig. 204.
† *Unio tuscumbiensis* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 191; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 11, pl. III, fig. 7; * Obs., XIII, 1874, p. 15, pl. III, fig. 7.—
* B. H. WRIGHT, Check List, 1888.
* *Unio flavidus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 156; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 28, pl. IX, fig. 25; * Obs., XIII, 1874, p. 32, pl. IX, fig. 25.—
* B. H. WRIGHT, Check List, 1888.

Tennessee River system.

†PLEUROBEMA TESSERULÆ Lea.

- * *Unio tesserula* LEA, Pr. Ac. N. Sci. Phila., VI, 1861, p. 392; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 40, pl. XV, fig. 39; * Obs., XI, 1867, p. 44, pl. XIV, fig. 38.—* B. H. WRIGHT, Check List, 1888.
* Margaron (*Unio*) *tesserula* LEA, Syn., 1870, p. 36.

Nolachucky River, Tennessee.

†PLEUROBEMA VALIDA Lea.

- * *Unio validus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 189; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 6, pl. I, fig. 2; * Obs., XIII, 1874, p. 10, pl. I, fig. 2.—* B. H. WRIGHT, Check List, 1888.

Tennessee.

†PLEUROBEMA ABACUS Haldeman.²

- * *Unio abacus* HALDEMAN, Jl. Ac. N. Sci. Phila., VIII, 1842, p. 202.—* CONRAD, Pr. Ac. N. Sci. Phila., VI., 1853, p. 244.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 143.
* Margaron (*Unio*) *abacus* LEA, Syn., 1852, p. 24; 1870, p. 38

Tennessee and southwest Virginia.

¹ The only specimen I have seen (the type) is a young shell, which I can not refer with certainty to any other species.

² As this species was not figured and I have not seen the type, I am unable to be absolutely certain what it is. In the Lea collection are four specimens labeled "*Unio abacus* Hald.," which are close to *appressus*.

† PLEUROBEMA SWORDIANA S. H. Wright.

* *Unio swordianus* S. H. WRIGHT, Naut., XI, 1897, p. 4.¹

* *Pleurobema swordiana*, SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 81, pl. iv, fig. 4.

Head waters of the Tennessee River.

† PLEUROBEMA TUMESCENS Lea.

* *Unio tumescens* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; Tr. Am. Phil. Soc., X, 1848, p. 71, pl. III, fig. 7; Obs., IV, 1848, p. 45, pl. III, fig. 7.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 170.

* *Margaron (Unio) tumescens* LEA, Syn., 1852, p. 25; 1870, p. 38.

† * *Unio radiosus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 192; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 13, pl. III, fig. 9; * Obs., XIII, 1874, p. 17, pl. III, fig. 9.—

* B. H. WRIGHT, Check List, 1888.

Tennessee. The localities, Alexandria, Louisiana, of Lea, for the type, and Ouachita River, Arkansas, given by Call, are open to doubt.

† PLEUROBEMA DOLABELLOIDES Lea.

* *Unio dolabelloides* LEA, Pr. Am. Phil. Soc., I, 1840, p. 288; * Tr. Am. Phil. Soc., VIII, 1842, p. 215, pl. xv, fig. 31; * Obs., III, 1842, p. 53, pl. xv, fig. 51.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* II. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* CHENU, Ill. Conch., 1858, pl. xxix, figs. 4, 4a, 4b.—* KUSTER, Conch. Cab., 1861, p. 214, pl. LXXI, fig. 3.—* SOWERBY, Conch. Icon., XVI, 1866, pl. xxxvii, fig. 205.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

* *Margaron (Unio) dolabelloides* LEA, Syn., 1852, p. 35; 1870, p. 56.

† * *Unio thorntonii*, LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 83; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 38, pl. xiv, fig. 36; * Obs., XI, 1867, p. 42, pl. xiv, fig. 36.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 169.

* *Margaron (Unio) thorntonii* LEA, Syn., 1870, p. 56.

† * *Unio mooresianus* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 83; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 39, pl. xiv, fig. 37; * Obs., XI, 1867, p. 43, pl. xiv, fig. 37.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) mooresianus* LEA, Syn., 1870, p. 39.

* *Unio mooreuianus* PÆTEL, Conch. Sam., III, 1890, p. 159.

† * *Unio recurvatus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 192; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 10, pl. II, fig. 6; * Obs., XIII, 1874, p. 14, pl. II, fig. 6.—* B. H. WRIGHT, Check List, 1888.

† * *Unio circumactus* LEA,² Pr. Ac. N. Sci. Phila., I, 1871, p. 192; * Jl. Ac. N. Sci. Phila., VIII, 1874, pl. IV, fig. 11; Obs., XIII, 1874, p. 19, pl. IV, fig. 11.—* B. H. WRIGHT, Check List, 1888.

Tennessee River drainage.

¹ I am a little in doubt as to the validity of this species.

² The figured type is certainly an adult, and that of *dolabelloides* the young of the same species.

† PLEUROBEMA SUBGLOBATA Lea.

- * *Unio subglobatus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 191; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 7, pl. I, fig. 3; * Obs., XIII, 1874, p. 11, pl. I, fig. 3.—
* B. H. WRIGHT, Check List, 1888.

Cumberland and Tennessee rivers.

† PLEUROBEMA CRUDA Lea.

- * *Unio crudus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 190; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 14, pl. IV, fig. 10; * Obs., XIII, 1874, p. 18, pl. IV, fig. 10.—
* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 149.

Cumberland, French Broad, and Holston rivers, Tennessee; Swamp Creek, Murray County, Georgia.

† PLEUROBEMA BARNESIANA Lea.

- * *Unio barnesianus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 31, pl. x, fig. 26; * Obs., II, 1838, p. 31, pl. x, fig. 26.—* TROSCHEL, Arch. für. Naturg., V, 1839, Pt. 2, p. 235.—* HANLEY, Test. Moll., 1842, p. 185; * Biv. Shells, 1843, p. 185, pl. XXIII, fig. 14.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* CONRAD Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XIX, figs. 2, 2a, 2b.—
* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIV, fig. 180.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 145.
* *Margarita (Unio) barnesiannus* LEA, Syn., 1836, p. 20; 1838, p. 17.
* *Margaron (Unio) barnesiannus* LEA, Syn., 1852, p. 24; 1870, p. 38.
* *Unio ravenelianus* REEVE, Conch. Icon., XVI, 1864, pl. XVI, fig. 70.
† * *Unio tellicoensis* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 155; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 31, pl. x, fig. 28; * Obs., XIII, 1874, p. 35, pl. x, fig. 28.—
* B. H. WRIGHT, Check List, 1888.

Cumberland and Tennessee river systems.

PLEUROBEMA LYONII Lea.

- * *Unio lyonii* LEA, Pr. Ac. N. Sci. Phila., IX, 1865, p. 89; * Jl. Ac. N. Sci. Phila., VI, 1869, p. 259, XXXII, fig. 74; * Obs., XII, 1869, p. 19, pl. XXXII, fig. 74.—
* B. H. WRIGHT, Check List, 1888.
Margaron (Unio) lyonii LEA, Syn., 1870, p. 56.

Tennessee River system.

PLEUROBEMA PUDICA Lea.

- * *Unio pudicus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 92; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 346, pl. LVI, fig. 171; * Obs., VIII, 1860, p. 28, pl. LVI, fig. 171.—
* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXI, fig. 427.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.
* *Margaron (Unio) pudicus* LEA, Syn., 1870, p. 37.
* *Unio subrotundus* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVII, fig. 201.

Tennessee River system.

† PLEUROBEMA BIGBYENSIS Lea.

- * *Unio bigbyensis* LEA, Pr. Am. Phil. Soc., II, 1841, p. 30; * Tr. Am. Phil. Soc., VIII, 1843, p. 237, pl. XXII, fig. 51; * Obs., III, 1842, p. 75, pl. XXII, fig. 51.—

* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XXII, figs. 5, 5a, 5b.—* KUSTER, Conch. Cab. Unio, 1862, p. 279, pl. XCIV, fig. 3.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLI, fig. 227.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

* *Margaron (Unio) bigbyensis* LEA, Syn., 1852, p. 24, 1870, p. 38.

Tennessee River drainage; Indian Territory? Texas? The two last localities very doubtful.

(Group of *Pleurobema decisa*.)

Shell solid, inflated, ovate to elliptical, very inequilateral, somewhat truncated in front and rounded or bluntly pointed behind, wedge-shaped when looked at from above; base slightly angled in front, then nearly straight for two-thirds of its length, from whence it curves to the posterior point; beaks high, curved inward and forward; beak sculpture consisting of a few coarse, irregularly concentric ridges which curve slightly upward behind; epidermis tawny to brownish, rayless, the rest periods very distinctly marked by dark bands; pseudo-cardinals stumpy, ragged, often showing a tendency to elongation in the direction of the axis of the shell.

Animal having the branchiæ rather small, inner the larger, free nearly or quite the entire length of the abdominal sac, marsupium occupying all but the extreme posterior end of the outer gills; branchiæ and anal openings papillose.

† PLEUROBEMA DECISA Lea.

* *Unio decisa* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 92, pl. XII, fig. 23; * Obs., I, 1834, p. 102, pl. XII, fig. 23.—* CONRAD, New F. W. Shells, 1834, p. 68.—* FERUS-SAC, Guer. Mag., 1835, p. 28.—* CONRAD, Monog., I, 1835, p. 6, pl. III, fig. 2.—* HANLEY, Test. Moll., 1842, p. 187; Biv. Shells, 1843, p. 187, pl. XXII, fig. 21.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* KUSTER, Conch. Cab. Unio, 1852, p. 41, pl. VII, fig. 3.¹—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XVIII, figs. 3, 3a, 3b.—* REEVE, Conch. Icon., XVI, 1864, pl. XVI, fig. 71.—B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 150.

* *Margarita (Unio) decisa* LEA, Syn., 1836, p. 21; 1838, p. 18.

* *Margaron (Unio) decisa* LEA, Syn., 1852, p. 26; 1870, p. 40.

* *Unio scalenius* SAY, Am. Conch., VI, 1834.

† * *Unio anaticulus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 40; * Jl. Ac. N. Sci. Phila., 1862, p. 92, pl. XIII, fig. 240; * Obs., VIII, 1862, p. 96, pl. XIII, fig. 240.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVII, fig. 199.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.

* *Margaron (Unio) anaticulus* LEA, Syn., 1870, p. 40.

* *Unio consanguineus* LEA, Pr. Ac. N. Sci. Phila., VI, 1861, p. 60; * Jl. Ac. N. Sci. Phila., V, 1862, p. 67, pl. VII, fig. 217; * Obs., VIII, 1862, p. 71, pl. VII, fig. 217.—* SOWERBY, Conch. Icon., XVI, 1866, pl. LXXVIII, fig. 409.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Margaron (Unio) consanguineus* LEA, Syn., 1870, p. 40.

¹ This figure is probably that of *U. decisa*. That on plate VIII, which he refers to that species, is most likely Lea's *ravenclianus*.

- †* *Unio crebrivittatus* LEA, Pr. Ac. N. Sci. Phila., VI, 1861, p. 60; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 43, pl. xv, fig. 41; *Obs., XI, 1867, p. 47, pl. xv, fig. 41.—*B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) crebrivittatus* LEA, Syn., 1870, p. 40.
- *? *Unio medius* REEVE, Conch. Icon., XVI, 1864, pl. xvii, fig. 77.

Alabama and Tombigbee river systems.

† PLEUROBEMA CHATTANOOGAENSIS Lea.

- * *Unio chattanoogaensis* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 166; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 209, pl. xxv, fig. 90; *Obs., VII, 1859, p. 27, pl. xxv, fig. 90.—*REEVE, Conch. Icon., XVI, 1864, pl. xvi, fig. 69.—B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 148.
- * *Margaron (Unio) chattanoogaensis* LEA, Syn., 1870, p. 40.¹

Alabama River system.

† PLEUROBEMA INTERVENTUS Lea.

- * *Unio interventus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 60; *Jl. Ac. N. Sci. Phila., V, 1862, p. 84, pl. xi, fig. 233; *Obs., VIII, 1862, p. 88, pl. xi, fig. 233.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 156.
- * *Margaron (Unio) interventus* LEA, Syn., 1879, p. 40.
- †* *Unio pallidofulvus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 60; *Jl. Ac. N. Sci. Phila., V, 1862, p. 83, pl. xi, fig. 332; *Obs., VIII, 1862, p. 87, pl. xi, fig. 232.—*B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) pallidofulvus* LEA, Syn., 1870, p. 40.

Cahawba River, Alabama.

† PLEUROBEMA MURRAYENSIS Lea.

- * *Unio murrayensis* LEA, Pr. Ac. N. Sci. Phila., II, 1868, p. 143; *Jl. Ac. N. Sci. Phila., VI, 1869, p. 303, pl. xlvi, fig. 115; *Obs., XII, 1869, p. 62, pl. xlvi, fig. 115.—*B. H. WRIGHT, Check List, 1888.
- * *Margaron (Unio) murrayensis* LEA, Syn., 1870, p. 40.

Coosa River system; Columbus, Georgia?

(Group of *Pleurobema curta*.)

Shell elongate triangular, truncated in front, rounded on the base where it is quite full behind the middle, with a low posterior ridge, the space between the middle of the disk and the ridge flattened or sometimes slightly excavated, rather sharply pointed behind; umbonal region very prominent; beaks well forward; beak sculpture not observed; epidermis rather smooth, brownish olive; pseudocardinals distinct, triangular, radiate; laterals heavy, slightly curved; cicatrices small and deep; nacre iridescent posteriorly. Animal with the gills large, semicircular, the outer slightly larger than the inner, which are free nearly their whole length from the abdominal sac; marsupium occupying the entire length of the outer gills.

¹ Probably did not come from Chattanooga, Tennessee, as Dr. Lea believes, but from some of the streams of northern Georgia or Alabama, which drain southward. According to Hon. T. H. Aldrich, the animal of this and allied species is a brilliant scarlet.

†PLEUROBEMA CURTA Lea.

- * *Unio curtus* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 112; *Jl. Ac. N. Sci. Phila., V, 1862, p. 103, pl. xvii, fig. 253; *Obs., VIII, 1862, p. 107, pl. xvii, fig. 256.—
*B. H. WRIGHT, Check List, 1888.

*Margaron (*Unio*) *curtus* LEA, Syn., 1870, p. 40.

Tombigbee River, Columbus, Mississippi.

(Group of *Pleurobema troscheliana*.)

Shell rather small, solid, rounded-triangular, inflated, somewhat inequilateral, slightly pointed near the posterior base, and rounded angular at the hinder end of the ligament; base of the shell almost evenly rounded; posterior ridge low; beaks high and slightly curved inward and forward over a well-defined lunule; epidermis dull tawny, showing the rest periods and occasionally marked with a radiating row of dark green, squarish spots in front of the posterior ridge; hinge plate wide and flat; pseudocardinals small, triangular, rough; laterals short, curved; muscle scars small, rather deep. Animal with semicircular gills, the inner the larger and free from the abdominal sac half to nearly their whole length; ovisacs of outer gills apparently in pairs.

†PLEUROBEMA TAITIANA Lea.

- **Unio taitianus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 39, pl. iv, fig. 11; *Obs., I, 1834, p. 151, pl. iv, fig. 11.—*FERUSSAC, Guer. Mag., 1835, p. 29.—*HANLEY, Test. Moll., 1842, p. 186; *Biv. Shells, 1843, p. 186, pl. xxiii, fig. 26.—*CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 258.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 169.

**Margarita (Unio) taitianus* LEA, Syn., 1836, p. 21; 1838, p. 18.

*Margaron (*Unio*) *taitianus* LEA, Syn., 1852, p. 25; 1870, p. 38.

Alabama River.

†PLEUROBEMA COR Conrad.

- **Unio mytilloides* CONRAD, Am. Jl. Sci., XXV, 1834, p. 343, pl. i, fig. 7.¹

**Unio cor* CONRAD, New F. W. Shells, 1834, p. 28, pl. iii, fig. 3, p. 68.—*FERUSSAC, Guerin Mag., 1835, p. 29.—*MÖLLER, Syn. Nov. Gen., 1836, p. 202.—*HANLEY, Test. Moll., 1842, p. 187; *Biv. Shells, 1843, p. 187.—*CHENU, Bib. Conch., 1st

¹Conrad says this is between *Unio ellipsis* Lea and *mytilloides* Rafinesque, but that he has no doubt that it is the latter. It is certainly very different from the shell Rafinesque figures and describes as *Pleurobema mytilloides*. (Ann. Gen. Sci. Phys. Brux., XIII, 1820, p. 313, pl. lxxxii, figs. 8-10.) Conrad's *Unio cor* was published in May, 1834, and his *mytilloides* in January of the same year, so that the latter has priority, but as Rafinesque's name was placed under the genus *Pleurobema*, and as I place Conrad's species, which is an entirely different thing, in the same genus, the name *mytilloides* can not be used for it, and it must therefore take the next name proposed, which is Conrad's *cor*.

Dr. Lea admits Rafinesque's *mytilloides* with doubt, and in his collection places under that name specimens which I regard as a rather elongated *Quadrula pyramidata*. Rafinesque's figure represents an elongated shell, almost absolutely straight on the dorsal line, and with the beaks carried far in front of the rest of the shell. I have never seen any thing which at all agrees with it.

ser., III, 1845, p. 16, pl. III, fig. 2.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 247.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 149.

**Margarita (Unio) cor* LEA, Syn., 1836, p. 21; 1838, p. 18.

**Margaron (Unio) cor* LEA, Syn., 1852, p. 26; 1870, p. 39.

† **Unio crapulus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 39; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 42, pl. xv, fig. 40; *Obs., XI, 1867, p. 46, pl. xv, fig. 40.—* B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) crapulus* LEA, Syn., 1870, p. 41.

† **Unio lewisii* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 40; *Jl. Ac. N. Sci. Phila., V, 1862, p. 71, pl. VIII, fig. 220; *Obs., VIII, 1862, p. 75, pl. VIII, fig. 220.—* B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) lewisii* LEA, Syn., 1870, p. 56.

Alabama River system; Flint River, Georgia.

† PLEUROBEMA PEROVATA Conrad.

**Unio perovatus* CONRAD, Am. Jl. of Sci., XXV, 1834, p. 338, pl. I, fig. 3; *† New F. W. Shells, 1834, p. 47, pl. II, fig. 3.¹—FERUSSAC, Guer. Mag., 1835, p. 29.—* MÜLLER, Syn. Nov. Gen., 1836, p. 199; *Test. Moll., 1842, p. 190.—* HANLEY, Biv. Shells, 1843, p. 190.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 22, pl. I, fig. 3.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* B. H. WRIGHT, Check List, 1888.

**Margarita (Unio) perovatus* LEA, Syn., 1836, p. 23; 1838, p. 19.

**Margaron (Unio) perovatus* LEA, Syn., 1852, p. 27; 1870, p. 42.

Prairie Creek (tributary of the Tombigbee), Marengo County, Alabama.

† PLEUROBEMA NUCLEOPSIS Conrad.

Unio nucleopsis CONRAD, Ann. and Mag. Nat. Hist., IV, 1849, p. 301;² *Pr. Ac. N. Sci. Phila., IV, 1849, p. 154; *Jl. Ac. N. Sci. Phila., I, 1850, p. 276, pl. XXXVII, fig. 81.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* KUSTER, Conch. Cab. Unio, 1861, p. 217, pl. LXXIII, fig. 3.—* REEVE, Conch. Icon., XVI, 1864, pl. XVI, fig. 68.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 161.

**Margarita (Unio) nucleopsis* LEA, Syn., 1852, p. 35; 1870, p. 56.

Coosa River system.

† PLEUROBEMA STABILIS Lea.

**Unio stabilis* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 59; *Jl. Ac. N. Sci. Phila., V, 1862, p. 71, pl. VIII, fig. 221; *Obs., VIII, 1862, p. 75, pl. VIII, fig. 221.—* B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) stabilis* LEA, Syn., 1870, p. 38.

† **Unio medius* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 40; *Jl. Ac. N. Sci. Phila., V, 1861, p. 78, pl. x, fig. 227; *Obs., VIII, 1862, p. 82, pl. x, fig. 227.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIII, fig. 375.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

**Margaron (Unio) medius* LEA, Syn., 1870, p. 40.

Coosa River, Alabama.

¹ This figure differs a good deal from that in the American Journal of Science, and it may be taken from a younger, less inflated shell.

² Very close to *P. irrasa*, and it may be that when a sufficient amount of material is examined the two will have to be placed together.

† PLEUROBEMA TROSCHELIANA Lea.

- * *Unio troschelianus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 280, pl. XXIII, fig. 39; * Obs., V, 1852, p. 36, pl. XXIII, fig. 39.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 170.

* *Margaron (Unio) troschelianus* LEA, Syn., 1852, p. 26; 1870, p. 40.

Alabama River system.

PLEUROBEMA IRRASA Lea.

- * *Unio irrasus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 38; * Jl. Ac. N. Sci. Phila., V, 1862, p. 91, pl. XIII, fig. 239; * Obs., VIII, 1862, p. 95, pl. XIII, fig. 239.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) irrasus* LEA, Syn., 1870, p. 38.

Coosa River system.

† PLEUROBEMA ALTA Conrad.

- * *Unio altus* CONRAD, Jl. Ac. N. Sci. Phila., II, 1854, p. 298, pl. XXVII, fig. 5.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) altus* LEA, Syn., 1870, p. 42.

- † * *Unio fibuloides* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 154; Jl. Ac. N. Sci. Phila., IV, 1859, p. 219, pl. XXVIII, fig. 100; * Obs., VII, 1859, p. 37, pl. XXVIII, fig. 100.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XII, fig. 223.—

* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 152.

* *Margaron (Unio) fibuloides* LEA, Syn., 1870, p. 56.

Connasauga River, Georgia.

† PLEUROBEMA HARTMANIANA Lea.

- * *Unio hartmanianus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 307; * Jl. Ac. N. Sci. Phila., V, 1862, p. 73, pl. VIII, fig. 222; * Obs., VIII, 1862, p. 77, pl. VIII, fig. 222.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) hartmanianus* LEA, Syn., 1870, p. 38.

Coosa River, Alabama.

Dr. Lea has shells in his collection from the Clinch River which he places with this species, which I am certain belong to the *Clava* group.

† PLEUROBEMA INSTRUCTA Lea.

- * *Unio instructus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 59; * Jl. Ac. N. Sci. Phila., V, 1862, p. 82, pl. x, fig. 230; * Obs., VIII, 1862, p. 86, pl. x, fig. 230.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) instructus* LEA, Syn., 1870, p. 38.

Cahawba River, Alabama.

(Group of *Pleurobema showalterii*.)

Shell small, rounded triangular, inflated, slightly truncated anteriorly, posterior end arched and ending in a bluntly rounded point at the post-base, posterior ridge well defined; umbonal region full; beaks rather prominent, the sculpture not observed; epidermis brown, not rayed; hinge plate rather wide; pseudocardinals small, triangular,

radial, roughened; laterals short, heavy, nacre white, brilliantly iridescent posteriorly; beak cavities shallow.

Animal apparently having ovisacs in pairs which are slightly wavy and lirate at the base; gills rather large, semicircular, inner much the larger, partly free from the abdominal sac; mantle thin, thickened at edge, and bordered with a dark line.¹

†PLEUROBEMA SHOWALTERII Lea.

* *Unio showalterii* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 307; *Jl. Ac. N. Sci. Phila., V, 1862, p. 73, pl. VIII, fig. 223; Obs., VIII, 1862, p. 77, pl. VIII, fig. 223.—

*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXI, fig. 426.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 167.

*Margaron (*Unio*) *showalterii* LEA, Syn., 1870, p. 55.

Coosa River, Alabama.

(Group of *Pleurobema nux*.)

Shell oval, inflated, moderately solid, with a low posterior ridge, swollen at the post-basal part; posterior end pointed at the termination of the posterior ridge; beaks full, rather high, considerably removed from the anterior end; epidermis pale to dark brown, rest lines very distinct; pseudocardinals small, stumpy, radiate, rough, laterals straight; nacre brilliantly iridescent posteriorly; front part of shell heavy, suddenly becoming thinner behind.

Animal unknown.

†PLEUROBEMA VERA Lea.

**Unio verus* LEA, Pr. Ac. N. Sci. Phila., V, 1860, p. 140; *Jl. Ac. N. Sci. Phila., V, 1862, p. 83, pl. XI, fig. 231; *Obs., VIII, 1862, p. 87, pl. XI, fig. 231.—*B. H. WRIGHT, Check List, 1888.

*Margaron (*Unio*) *verus* LEA, Syn., 1870, p. 38.

Black Warrior and Cahawba rivers, Alabama.

†PLEUROBEMA HAGLERI Frierson.

**Unio (Pleurobema) hagleri* FRIERSON, Nautilus, XIII, 1900, p. 109, pl. II.

North River, Alabama, near Tyner.

†PLEUROBEMA RUBELLA Conrad.

**Unio rubellus* CONRAD, New F. W. Shells, 1834, p. 38, pl. VI, fig. 2, p. 71.—*FERUS-SAC, Guer. Mag., 1835, p. 29.—*MÖLLER, Syn. Nov. Gen., 1836, p. 205.—*HANLEY, Test. Moll., 1842, p. 201; *Biv. Shells, 1843, p. 201.—*CHENU, Bib. Conch., 1st ser., III, 1845, p. 20, pl. II, fig. 2.—*CATLOW and REEVE, Conch. Nom., 1845, p. 63.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 166.

¹ The single species which I place in this group is quite variable. Some specimens approach *Quadrula solida* and *trigona* in general shape, while others are more like *Pleurobema irrasa* and *trosceliana*. I judge from the character of the gills that the marsupium is confined to the outer pair.

**Margarita (Unio) rubellus* LEA, Syn., 1836, p. 33; 1838, p. 28.

**Margaron (Unio) rubellus* LEA, Syn., 1852, p. 34; 1870, p. 55.

† **Uniorudis* CONRAD, ¹ Monog., No. 9, 1837, p. 76, pl. XLIII, fig. 1.—* B. H. WRIGHT, Check List, 1888.

† **Unio pulvinulus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; * Tr. Am. Phil. Soc., X, 1848, p. 81, pl. VIII, fig. 24; * Obs., IV, 1848, p. 55, pl. VIII, fig. 24.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.

**Margaron (Unio) pulvinulus* LEA, Syn., 1852, p. 26; 1870, p. 41.

Black Warrior and Cahawba rivers, Alabama.

† PLEUROBEMA FURVA Conrad.²

**Unio furvus* CONRAD, New F. W. Shells, 1834, p. 39, pl. VI, fig. 3, page 69.—

* FERUSSAC, Guer. Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen., 1836, p. 205.—

* CHENU, Bib. Conch., 1st ser., III, 1845, p. 20, pl. II, fig. 4.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 250.—* B. H. WRIGHT, Check List, 1888.

Black Warrior River, Alabama.

† PLEUROBEMA AVELLANA Simpson.

**Pleurobema avellana* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 81, pl. II, figs. 6, 7.

Cahawba River, Alabama.

† PLEUROBEMA NUX Lea.

**Unio nux* LEA, Tr. Am. Phil. Soc., X, 1852, p. 283, pl. XXIV, fig. 43; * Obs. V, 1852, p. 39, pl. XXIV, fig. 43.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—

* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 161.

**Margaron (Unio) nux* LEA, Syn., 1852, p. 31; 1870, p. 49.

† **Unio cinnamomicus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 39; * Jl. Ac. N. Sci. Phila., V, 1862, p. 100, pl. XVI, fig. 248; * Obs., VIII, 1862, p. 104, pl. XVI, fig. 248.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIII, fig. 438.—* B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) cinnamomicus* LEA, Syn., 1870, p. 49.

Alabama River system.

† PLEUROBEMA PINKSTONI S. H. Wright.

**Unio pinkstoni* S. H. WRIGHT, Nautilus, X, 1897, p. 136.

**Pleurobema pinkstoni* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 81, pl. I, fig. 8.

Alabama River system.

¹ Conrad says that *U. ravenelianus* of Lea is the same as his *U. rudus*, which, according to Ravenel, is found in the French Broad River, though Conrad obtained his species in the Black Warrior. The species of Conrad is, however, quite distinct from that of Lea, the two never occurring in the same drainage system.

² I have seen what I believe are specimens of this species from the collections of the Hon. T. H. Aldrich and Mr. Bryant Walker, taken from the Black Warrior River, Alabama, but I think it not improbable that it may run into *P. rubella*.

†PLEUROBEMA JOHANNIS Lea.

* *Unio johannis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 171; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 340, pl. LV, fig. 168; *Obs., VIII, 1860, p. 25, pl. LV, fig. 168.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) johannis* LEA, Syn., 1870, p. 41.

Alabama River system.

†PLEUROBEMA HANLEYANA Lea.

* *Unio hanleyanus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 279, pl. XXIII, fig. 37; *Obs., V, 1852, p. 35, pl. XXIII, fig. 37.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* REEVE, Conch. Icon., XVI, 1864, pl. XVII, fig. 76.¹—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLVI, fig. 249.—* B. H. WRIGHT, Check List, 1888.—* PÄTEL, Conch. Sam., III, 1890, p. 154.

* *Margaron (Unio) hanleyanus* LEA, Syn., 1852, p. 26; 1870, p. 40.

Coosa River drainage, Georgia and Alabama.

†PLEUROBEMA FLAVIDULUS Lea.

* *Unio flavidulus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 39; *Jl. Ac. N. Sci. Phila., V, 1862, p. 97, pl. xv, fig. 245; *Obs., VIII, 1862, p. 101, pl. xv, fig. 245.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) flavidulus* LEA, Syn., 1870, p. 40.

Columbus, Mississippi.

(Group of *Pleurobema bulbosa*.)

Shell elongate oval, solid, inflated, with a high posterior ridge, ending in a point behind, above which it is feebly wrinkled; beaks full, sculpture not seen; epidermis smooth, dark, rayless; hinge strong; pseudocardinals heavy, torn, sometimes a small third one in the left valve; laterals heavy, club-shaped, granulate, two in the left valve and one and a small secondary lateral in the right. Animal unknown.

†PLEUROBEMA BULBOSA Lea.

* *Unio bulbosus* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 172; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 191, pl. XXI, fig. 75; *Obs., VII, 1859, p. 9, pl. XXI, fig. 75.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) bulbosus* LEA, Syn., 1870, p. 40.

Ocmulgee and Flint rivers, Georgia.

†PLEUROBEMA HARPERI B. H. Wright.

* *Unio harperi* B. H. WRIGHT, Naut., XIII, 1899, p. 6.

* *Pleurobema harperi* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 81, pl. I, fig. 10.

Altamaha and Flint rivers, Georgia; Suwanee River, Florida.²

¹ Reeve states in his errata that this is not *Unio hanleyanus*. The figure agrees fairly well with the type of that species.

² This may prove to be a variety of *P. bulbosa*.

†PLEUROBEMA RECLUSA B. H. Wright.

* *Unio reclusus* B. H. WRIGHT, Naut., XI, 1898, p. 3.

* *Pleurobema reclusa* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 82, pl. I, fig. 2.

Ocklocknee River, Leon County, Florida.

(Group of *Pleurobema brumbyana*.)

Shell oval, inflated, moderately solid, faintly swollen at post basal region; beaks rather prominent, somewhat distant from the anterior end; beak sculpture not seen; disks irregularly, concentrically sculptured; epidermis dark olive to blackish, striate; pseudocardinals subradiate, slightly compressed; beak cavities shallow; nacre iridescent posteriorly, somewhat thickened in front.

†PLEUROBEMA BRUMBYANA Lea.

* *Unio brumleyanus* LEA,¹ Proc. Am. Phil. Soc., II, 1841, p. 82.

* *Unio brumbyanus* LEA, Tr. Am. Phil. Soc., VIII, 1842, p. 245, pl. XXVI, fig. 62;

* Obs., III, 1842, p. 83, pl. XXVI, fig. 62.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

* *Margaron (Unio) brumbyanus* LEA, Syn., 1852, p. 31; 1870, p. 49.

† * *Unio concolor* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 40; * Jl. Ac. N. Sci. Phila., V, 1862, p. 89, pl. XII, fig. 237; * Obs., VIII, 1862, p. 93, pl. XII, fig. 237.—

* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIII, fig. 440.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1870, p. 148.

* *Margaron (Unio) concolor* LEA, Syn., 1870, p. 49.

Alabama River system.

(Group of *Pleurobema argentea*.)

Shell solid, oval to rhomboid, generally compressed, beaks high but not inflated, well removed from the anterior end, with strong, irregular sculpture, which is curved up and swollen behind where the bars are slightly looped; epidermis brownish to straw color, sometimes having a few broken, blotched rays; teeth strong; cicatrices deep and distinct. Animal unknown.²

†PLEUROBEMA SIMULANS Lea.

* *Unio simulans* LEA, Pr. Acad. N. Sci. Phila., I, 1871, p. 190; * Jl. Acad. N. Sci. Phila., VIII, 1874, p. 18, pl. V, fig. 15; * Obs., XIII, 1874, p. 22, pl. V, fig. 15.—

* B. H. WRIGHT, Check List, 1888.

Black Warrior and Cahawba rivers, Alabama; Pine Barren Creek, Escambia County, Florida.

¹ So written in Proceedings; no doubt a typographical error, as the species was named after Brumby.

² The species placed here are puzzling and seem to stand between *Unio* and *Pleurobema*. The color and the beak sculpture incline me to place them with the latter, and some of the species seem to show relationship with such forms as *P. pudica* and *bigbyana*.

† PLEUROBEMA STRODEANA B. H. Wright.

* *Unio strodeanus* B. H. WRIGHT, Nautilus, XII, 1898, p. 5.* *Pleurobema strodeana* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 81, pl. I, fig. 3.

Escambia River, Florida; Flint River, Rhoadsville, Georgia.

† PLEUROBEMA PATSALIGENSIS Simpson

* *Pleurobema patsaligensis* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 82, pl. II, fig. 1.

Little Patsaliga Creek, southeast Alabama.

† PLEUROBEMA FAVOSA Lea.

* *Unio favosus* LEA, Proc. Acad. N. Sci. Phila., VIII, 1856, p. 262; * JI. Acad. N. Sci. Phila., IV, 1858, p. 58, pl. VIII, fig. 40; * Obs., VI, p. 58, pl. VIII, fig. 40.—* SOWERBY, Conch. Icon., XVI, 1868, pl. xc, fig. 488.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 151.* *Margaron (Unio) favosus* LEA, Syn., 1870, p. 38.

Alabama River system.

† UNIO LENTICULARIS Lea.

* *Unio lenticularis* LEA, Pr. Acad. N. Sci. Phila., II, 1872, p. 155; * JI. Acad. N. Sci. Phila., VIII, 1874, p. 30, pl. ix, fig. 27; * Obs., XIII, 1874, p. 34, pl. ix, fig. 27.—

* B. H. WRIGHT, Check List, 1888.

Tellico River, Monroe County, Tennessee; East Tennessee.

† PLEUROBEMA MEREDITHII Lea.

* *Unio meredithii* LEA, Pr. Acad. N. Sci. Phila., II, 1858, p. 40; * JI. Acad. N. Sci. Phila., V, 1862, p. 65, pl. vi, fig. 214; * Obs., VIII, 1862, p. 69, pl. v, fig. 214.—

* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 159.

* *Margaron (Unio) meredithii* LEA, Syn., 1870, p. 35.

Tennessee River system; Black Warrior River, Alabama.

† PLEUROBEMA LITA Lea.

* *Unio litus* LEA, Pr. Acad. N. Sci. Phila., I, 1871, p. 189; * JI. Acad. N. Sci. Phila., VIII, 1874, p. 17, pl. v, fig. 13; * Obs., XIII, 1874, p. 21, pl. v, fig. 13.—* B.

H. WRIGHT, Check List, 1888.

Cahawba and Black Warrior rivers, Alabama.

PLEUROBEMA GEORGIANA Lea.

* *Unio georgianus* LEA, Pr. Am. Phil. Soc., II, 1841, p. 31; * Tr. Am. Phil. Soc., VIII, 1842, p. 235, pl. XXI, fig. 49; * Obs., III, 1842, p. 73, pl. XXI, fig. 49.—

* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. XXXII, figs. 3, 3a, 3b.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 153.

* *Margaron (Unio) georgianus* LEA, Syn., 1852, p. 27; 1870, p. 42.

Stamp Creek, northwest Georgia.

¹ I have only seen a single specimen of this, the type, in bad condition, and I can not be certain where it belongs.

† *PLEUROBEMA PYRIFORMIS* Lea.

- * *Unio pyriformis* LEA, Pr. Ac. N. Sci. Phila., IX, 1857, p. 31; * Jl. Ac. N. Sci. Phila., IV, 1858, p. 69, pl. XII, fig. 50; * Obs., VI, 1858, p. 69, pl. XII, fig. 50.—
* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) pyriformis* LEA, Syn., 1870, p. 40.

Near Columbus, Georgia.

† *PLEUROBEMA MODICA* Lea.

- * *Unio modicus* LEA, Pr. Acad. N. Sci. Phila., IX, 1857, p. 171; * Jl. Acad. N. Sci. Phila., IV, 1859, p. 204, pl. XXIV, fig. 86; * Obs., VII, 1860, p. 22, pl. XXIV, fig. 86.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) modicus* LEA, Syn., 1870, p. 40.

Chattahoochee River, Georgia.

† *PLEUROBEMA STRIATA* Lea.

- * *Unio striatus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 287; * Tr. Am. Phil. Soc., VIII, 1843, p. 203, pl. XII, fig. 16; * Obs., III, 1842, p. 41, pl. XII, fig. 16.—* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* CHENU, Ill. Conch., 1858, pl. XXIX, figs. 1, 1a, 1b.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) striatus* LEA, Syn., 1852, p. 26; 1870, p. 40.

Chattahoochee River, Georgia.

† *PLEUROBEMA GIBBER* Lea.

- * *Unio gibber* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 34, pl. x, fig. 30; * Obs., II, 1838, p. 35, pl. x, fig. 30.—* TROSCHER, Arch. fur Naturg., V, 1839, Pt. 2, p. 236.—
* HANLEY, Test. Moll., 1842, p. 185; * Biv. Shells, 1843, p. 185, pl. XXI, fig. 46.—
* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—
* CHENU, Ill. Conch., 1858, pl. XXIII, figs. 2, 2a, 2b.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XL, fig. 219.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) gibber* LEA, Syn., 1836, p. 20; 1838, p. 17.

* *Margaron (Unio) gibber* LEA, Syn., 1852, p. 24; 1870, p. 37.

Caney Fork, Tennessee.

† *PLEUROBEMA FASSINANS* Lea.

- * *Unio fassinans* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 143; * Jl. Ac. N. Sci. Phila., VI, 1869, p. 305, pl. XLVII, fig. 118; * Obs., XII, p. 65, pl. XLVII, fig. 118.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) fassinans* LEA, Syn., 1870, p. 42.

† *PLEUROBEMA FASSINANS* var. *RHOMBOIDEA* Simpson.¹

Upper Tennessee River drainage.

¹ A form of what I consider *fassinans* is abundant in the Clinch River, in western Virginia, which is considerably more rhomboid than the type and may be called var. *rhomboidea*.

†PLEUROBEMA ARGENTEA Lea.

- **Unio argenteus* LEA, Pr. Am. Phil. Soc., II, 1841, p. 82; *Tr. Am. Phil. Soc., VIII, 1843, p. 242, pl. xxv, fig. 57; *Obs., III, 1842, p. 80, pl. xxv, fig. 57.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—*CHENU, Ill. Conch., 1858, pl. xxxiii, figs. 2, 2a, 2b.—H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*KUSTER, Conch. Cab. Unio, 1861, p. 188, pl. LIX, fig. 4; 1861, p. 225, pl. LXXVI, fig. 3.—*SOWERBY, Conch. Icon., XVI, 1868, pl. xxxvii, fig. 204.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 145.
- *Margaron (*Unio*) *argenteus* LEA, Syn., 1852, p. 26; 1870, p. 40.

Upper Tennessee River drainage.

†PLEUROBEMA ARGENTEA var. PANNOSA Simpson.

- **Pleurobema argentea pannosa* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 82.

A solid, somewhat inflated, roughened form from Hot Springs, Arkansas, may be designated as above. Probably a valid species.

†PLEUROBEMA CONASAUGÆNSIS Lea.

- **Unio conasaugensis* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 155.—*Jl. Ac. N. Sci. Phila., VIII, 1874, p. 33, pl. x, fig. 30; *Obs., XIII, 1874, p. 37, pl. x, fig. 30.—*B. H. WRIGHT, Check List, 1888.

Upper Tennessee River drainage.

†PLEUROBEMA BREVIS Lea.

- **Unio brevis* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 157; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 35, pl. xii, fig. 32; *Obs., XIII, 1874, p. 39, pl. xii, fig. 32.—*B. H. WRIGHT, Check List, 1888.

Tennessee River system; Hardy, Arkansas.

†PLEUROBEMA BREVIS var. SUBELLIPTICA Simpson.

- **Pleurobema brevis subelliptica* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 80.

A solid, inflated form from Hardy, Arkansas, may bear this name. It is probably a distinct species.

†PLEUROBEMA PLANIOR Lea.

- **Unio planior* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 145; *Jl. Ac. N. Sci. Phila., VI, 1869, p. 316, pl. I, fig. 129; *Obs., XII, 1869, p. 76, pl. I, fig. 129.—*B. H. WRIGHT, Check List, 1888.
- *Margaron (*Unio*) *planior* LEA, Syn., 1870, p. 35.

Upper Tennessee River drainage.

†PLEUROBEMA ESTABROOKIANA Lea.

- **Unio estabrookianus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; *Tr. Am. Phil. Soc., X, 1853, p. 77, pl. vi, fig. 17; *Obs., IV, 1848, p. 51, pl. vi, fig. 77.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*B. H. WRIGHT, Check List, 1888.
- *Margaron (*Unio*) *estabrookianus* LEA, Syn., 1852, p. 24; 1870, p. 35.
- **Unio estabrookianus* PÆTEL, Conch. Sam., III, 1890, p. 152.

- * *Unio striatissimus* ANTHONY, Am. Jl. Conch., I, p. 156, 1865, pl. XII, fig. 1.—
 *SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVI, fig. 460.—*PÆTEL, Conch.
 Sam., III, 1890, p. 168.

Upper Tennessee River drainage.

Section *PLETHOBASUS* Simpson, 1900.

(Type, *Unio æsopus* Green.)

Shell large, irregularly oval, inflated, solid, somewhat suddenly swollen at the posterior base; posterior ridge low and rounded; beaks rather high near the anterior end, having a few strong ridges which are curved upward behind; a row of low, irregular tubercles extends from near the beaks to the post basal part of the valves; epidermis tawny brown in the younger shell, becoming darker with age; hinge plate solid, not flattened; pseudocardinals triangular, rough; cavity of the beaks not deep; front part of the shell very heavy; hinder part much thinner.

Animal having the mantle somewhat toothed posteriorly; branchial and anal openings large, the latter smooth or with very faint papillæ; inner gills the larger, rounded below, free nearly or quite their whole length from the abdominal sac; marsupium occupying the entire outer gills.¹

† *PLEUROBEMA* *ÆSOPUS* Green.

- * *Unio æsopus* GREEN, Cont. Mac. Lyceum, I, No. 2, 1827, p. 46, fig. 3.—* HANLEY, Test. Moll., 1842, p. 181; * Biv. Shells, 1843, p. 181, pl. XXIV, fig. 7.—* CATLOW and REEVE, Conch. Nom., 1845, p. 551.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* CALKINS, Proc. Ottawa Acad. N. Sci., 1874, p. 41.—* CALL, Jl. Cinti. Soc. N. H., XVIII, 1896, p. 157, pl. VI.
 * *Margarita (Unio) æsopus* LEA, Syn., 1836, p. 17; 1838, p. 16.
 * *Margaron (Unio) æsopus* LEA, Syn., 1852, p. 23; 1870, p. 34.
 * *Unio cyphia* CONRAD, New F. W. Shells, 1834, p. 68.—* FERUSSAC, Guer. Mag., 1835, p. 27.
 * *Unio cyphius* SAY, Am. Conch., VI, 1834.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* REEVE, Conch. Icon., XVI, 1864, pl. VIII, fig. 28.
 * *Unio cyphas* var. *æsopus* PÆTEL, Conch. Sam., III, 1890, p. 150.
 * *Unio scyphius* KUSTER, Conch. Cab. Unio, 1861, p. 181, pl. LVII, fig. 2.

Ohio, Cumberland, and Tennessee river systems; west to Missouri and Minnesota; Alabama River, according to Call.

¹I am not positive as to the generic position of the two following species, but I am inclined to place them in *Pleurobema*. No other living species of this genus has any development of tubercles, but a great number of fossil forms from the Tertiary of eastern Europe which seem to be most like *Pleurobema* are decidedly pustulous. The heavy, inflated, high beaks and the comparatively shallow beak cavities are characters found in all the species of this genus. In the single gravid specimen I examined (one out of a large number from different localities) the outer gills were not filled very full of ova, though they were found throughout their extent, while the most careful search did not disclose any in the inner gills.

† PLEUROBEMA CICATRICOSA Say.

* *Unio varicosus* LEA, Tr. Am. Phil. Soc., IV, 1829, p. 90, pl. XL, fig. 20;¹ * Obs., I, 1834, p. 100, pl. XI, fig. 20.—* HANLEY, Test. Moll., 1842, p. 181; * Biv. Shells, 1843, p. 181, pl. XXI, fig. 14.—* CATLAW and REEVE, Conch. Nom., 1845, p. 65.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* CHENU, Ill. Conch., 1858, pl. XI, figs. 6, 6a, 6b.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) varicosus* LEA, Syn., 1836, p. 17; 1838, p. 16.

* *Margaron (Unio) varicosus* LEA, Syn., 1852, p. 23; 1870, p. 34.

* *Unio cicatricosus* SAY, N. Harmony Diss., II, No. 19, 1829, p. 292.—* SAY, Am. Conch., VI, 1834.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* L. W. SAY, Terr. and Fluv. Shells, 1840, p. 5.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 246.—* KUSTER, Conch. Cab. Unio, 1861, p. 185, pl. LVIII, fig. 2.—* REEVE, Conch. Icon., XVI, 1864, pl. VIII, fig. 31; XIII, fig. 50.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 148.

* *Unio cicatricosus* var. *varicosus* PETEL, Conch. Sam., III, 1890, p. 148.

Ohio River; Tennessee; Claiborne, Alabama. The last locality I consider doubtful. Lea gives St. Paul, Minnesota, as a habitat, but I do not think it possibly can be.

The following species are described by Rafinesque, but I am unable to make them out.

* *Pleurobema mytiloides* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 313, pl. LXXXII, figs. 8–10.

* *Pleurobema cuneata* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 313.

TETRAGENEÆ.

Male and female shells alike, solid; beak sculpture consisting of coarse, subparallel ridges; beak cavities deep; marsupium filling all four gills, smooth, pad-like.

Genus QUADRULA (Rafinesque, 1820) Agassiz.

(Type, *Quadrula metanевра* Rafinesque.)

Quadrula RAFINESQUE, Ann. Gen. Sci. Phys. Brux., 1820, p. 305.

Rotundaria AGASSIZ, Arch. für Naturg., 1852, p. 48.

Orthonymus AGASSIZ, Arch. für Naturg., 1852, p. 48.

Shell triangular, quadrate or rhomboid, solid, inflated, with rather prominent beaks, which are generally sculptured with a few coarse, irregular, subparallel ridges that are inflated where they cross the posterior ridge; posterior ridge ordinarily well developed; base often incurved in old specimens; disks sculptured or smooth; epidermis usually dull colored, dark and rayless, or feebly rayed; hinge plate

¹ I change Lea's name because Lamarck previously applied the name *Unio varicosa* to what is, no doubt, *Alasmidonta marginata*. Mr. T. G. Lea, of Cincinnati, took many specimens of this species, the shells of which he sent to Dr. Lea, and in several of them he has written in pencil "not charged" or "ovaries charged," with the date, but neither of them seems to have described the animal. I am somewhat at a loss to know where to place this curious form. The young are much like those of *Quadrula solida*, and do not show the swellings until the third or fourth year, and occasionally the adult shell is nearly smooth.

heavy, wide, flattened; pseudocardinals solid, direct, ragged; laterals double in the left and single in the right valve, often with a small secondary lateral below the large one in the right valve; cavity of the beaks deep and compressed; dorsal scars under the hinge plate; male and female shells alike.

Animal having the inner gills the larger, generally free from the abdominal sac the greater part or all of their length; marsupium occupying all four of the gills throughout, the whole smooth and pad-like.

Section CRENODONTA Schlüter.¹

(Type, *Unio plicatus* Say.)

Shell more or less alate; beaks prominent; the surface of the valves usually sculptured with oblique folds; posterior slope generally having smaller radial plications which curve upward behind; epidermis brownish or blackish; anterior muscle scars large, distinct, very shallow, the anterior edge smooth, the rest apparently filled with roughened shelly matter; posterior scars large, shallow, indistinct; escutcheon large and dark.

Animal with the gills generally large, rounded below; inner the larger, usually free nearly or quite the entire length of the abdominal sac,² the two pairs united to the mantle nearly but not quite to the posterior end, having a small portion free; marsupium occupying all the four branchiæ, forming very heavy, thick pads; labial palpi usually large.

(Group of *Quadrula plicata*.)

Shell rounded to subrhomboid; plications usually strong, oblique, though in occasional specimens the surface may be perfectly plane or slightly concentrically sculptured.

¹ Schlüter applied the name *Crenodonta* (Verz. meiner Conch., 1836, p. 33) to a group of Unionidæ, the first species of which was the *Unio plicatus* of Say, but he gave no description of his group and did not designate a type. In 1853 Mörch (Yoldi catalogue, p. 45) used this name without a description or a type, and it has been applied to the plicate Uniones by von Martens (Biologia Centrali-Americana, Mollusca, 1900, p. 479).

² Dr. Lea found in *Quadrula multiplicata* the inner gills generally nearly or entirely free, but in certain specimens they were wholly united. My own experience in examining the animals of this species exactly coincides with his, thus showing that the character of the union of the inner gills with the abdominal sac, or their separation from it, is not a generic character, as Agassiz believed, nor is it even of specific value. Although I have examined thousands of animals belonging to the *Plicata* group of the genus *Quadrula*, taken at various seasons and throughout almost the entire range of the assemblage and of nearly all the species, I have never seen a specimen with ova or embryos in the gills save one in the alcoholic collection of Dr. Lea in the Academy of Natural Sciences at Philadelphia. Other students of the Unionidæ have found these species equally barren. It is probable that they breed only at long intervals but in enormous numbers, a supposition strengthened by the fact that Dr. Lea found about 6,000,000 young in the gills of a single *Unio multiplicatus*.

†QUADRULA PLICATA Say.

* *Unio plicata* SAY,¹ Nich. Enyc., II, 1816, pl. III, fig. 1.

* *Unio (Theliderma) plicata* SWAINSON, Tr. on Mal., 1840, p. 271, fig. 54e.

* *Mya plicata* EATON, Zool. Text-Book, 1826, p. 219.

* *Quadrula plicata* BAKER, Moll. Chicago, Pt. I, 1898, pl. XXV, fig. 1.

* *Unio plicatus* HILDRETH (?), Am. Jl. Sci., XIV, 1828, p. 280.²—SHORT and EATON, Transylvania Jl., 1831, p. 74.—* CONRAD, New F. W. Shells, 1834, p. 71.—* SAY, Am. Conch., VI, 1834.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* HANLEY, Test. Moll., 1842, p. 175; * Biv. Shells, 1843, p. 175, pl. XXI, fig. 21.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* KUSTER, Conch. Cab. Unio, 1856, p. 137, pl. XL, fig. 3.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* REEVE, Conch. Icon., XVI, 1864, pl. II, fig. 5.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 163.

* *Margarita (Unio) plicatus* LEA, Syn., 1836, p. 12; 1838, p. 14.

* *Margaron (Unio) plicatus* LEA, Syn., 1852, p. 20; 1870, p. 30.

* *Plectomerus plicatus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 261.

* *Unio peruriana* LAMARCK, An. sans Vert., VI, 1819, p. 71.³—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 533; 3d ed., II, 1839, p. 667.

* *Unio crassus* BARNES, Am. Jl. Sci., VI, 1823, p. 118.

* *Unio rariplicata* DESHAYES, Enc. Méth., II, 1830, p. 578; * An. sans Vert., 2d ed., VI, 1835, p. 533; 3d ed., II, 1839, p. 667.

* *Unio giganteus* LEA, Obs., II, 1838, p. 35.⁴

* *Unio heros* KUSTER, Conch. Cab. Unio, 1856, p. 136, pl. XI, figs. 1, 2.

†UNIO PLICATA var. HIPPOPÆA Lea.

* *Unio hippopæus* LEA,⁵ Pr. Am. Phil. Soc., IV, 1845, p. 163; * Tr. Am. Phil. Soc., X, 1848, p. 67, pl. I, fig. 1; * Obs., IV, 1848, p. 41, pl. I, fig. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* KUSTER, Conch. Cab., 1861, p. 251, pl. LXXXIV, fig. 3.—* REEVE, Conch. Icon., XVI, 1864, pl. XI, fig. 40.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) hippopæus* LEA, Syn., 1852, p. 21; 1870, p. 31.

* *Unio hippopæus* PÆTEL, Conch. Sam., III, 1890, p. 155.

Upper Mississippi drainage area south to the Tennessee and Arkansas rivers; Red River of the North; Saskatchewan River; Lake Wiinipeg; western Michigan. The variety in eastern Michigan and Lake Erie.

†QUADRULA PERPLICATA Conrad.

* *Unio perplicatus* CONRAD, Pr. Ac. N. Sci. Phila., I, 1841, p. 19; * Jl. Ac. N. Sci. Phila., VIII, 1842, p. 178; * I, 1850, p. 276, pl. XXXVIII, fig. 2.—* H. and A.

¹Say says: "It was found by Mr. Lesueur in Lake Erie, and was communicated by him under the above name." As there is no evidence that Lesueur described the species, and as Say was the one to first properly characterize it, I believe that he must be credited with it.

²Referred to figure 5, but this with a number of other figures in this paper were left out by the editor.

³Lea, who examined Lamarck's types and collection, states that this is the same as Say's *plicata*.

⁴Dr. Lea does not describe this, but only states that it is in Dr. Mitchell's collection. Probably it has never been described.

⁵Apparently a stunted variety of *Quadrula plicata*. Dr. Lea's specimens show a great diversity of characters.

ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—*REEVE, Conch. Icon., XVI, 1864, pl. IX, fig. 35.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 163.

**Margaron (Unio) perplicatus* LEA, Syn., 1852, p. 20; 1870, p. 29.

**Plectomerus perplicatus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 261.

†**Unio atrocostatus* LEA, Tr. Am. Phil. Soc., X, 1848, p. 70, pl. II, fig. 5;¹ *Obs., IV, 1848, p. 44, pl. II, fig. 5.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 145.

**Margaron (Unio) atrocostatus* LEA, Syn., 1852, p. 20; 1870, p. 29.

†**Unio pearlensis* CONRAD, Tr. Am. Phil. Soc., VII, 1855, p. 256.²—*REEVE, Conch. Icon., XVI, 1864, pl. XI, fig. 42.

**Unio perlensis* PÆTEL, Conch. Sam., III, 1890, p. 163.

†**Unio brazosensis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 144;³ *Jl. Ac. N. Sci. Phila., VI, 1869, p. 309, pl. XLVIII, fig. 122; *Obs., XII, 1869, p. 69, pl. XLVIII, fig. 122.—*B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) brazosensis* LEA, Syn., 1870, p. 31.

†**Unio linccumii* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 144; *Jl. Ac. N. Sci. Phila., VI, 1869, p. 312, pl. XLIX, fig. 125; *Obs., XII, 1869, p. 72, pl. XLIX, fig. 125.—*B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) linccumii* LEA, Syn., 1870, p. 31.

**Unio linccurii* PÆTEL, Conch. Sam., III, 1890, p. 157.⁴

†**Unio pauciplicatus* LEA, Pr. Ac. N. Sci. Phila., II, 1872, p. 156; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 29, pl. IX, fig. 26; *Obs., XIII, 1874, p. 33, pl. IX, fig. 26.—*B. H. WRIGHT, Check List, 1888.

†QUADRULA PERPLICATA var. QUINTARDII Cragin.⁵

**Unio quintardii* CRAGIN, Bull. Washb. College, II, 1887, p. 6.—*PILSBRY, Pr. Ac. N. Sci. Phila., 1892, p. 131, pl. VII, figs. 1-3.—*B. H. WRIGHT, Check List, 1888.

Alabama River drainage and streams flowing into the Gulf of Mexico west to central Texas, north to southern Kansas.

†QUADRULA ELLIOTTII Lea.⁶

**Unio elliotii* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 262; *Jl. Ac. N. Sci. Phila., IV, 1858, p. 54, pl. VII, fig. 37; *Obs., VI, 1858, p. 54, pl. VII, fig.

¹This seems to be merely a form of *perplicata*, with a rather rude growth and roughened epidermis.

²A comparatively smooth form of *perplicatus*, according to the type.

³Another smooth form of *perplicatus*.

⁴The above is a fair sample of Pætel's orthography throughout his work.

⁵A right valve of the type was kindly loaned to the writer by Mr. Quintard. It is a rather light brown, glossy shell, and is sculptured with a series of Λ-shaped markings on the disk. Otherwise it is generally like *Q. perplicatus*. I have seen the *Q. undulata* and one or two other species of this group break from the ordinary oblique bars into these peculiar markings, and I am quite sure that in this case the same thing has happened to *Q. perplicata*.

⁶Dr. Lea first called this *Unio atrocostatus*, and then wrote in one of his duplicates *Unio elliotii*. He says that it is very near to *U. perplicatus* Conrad, but that the young of his species have a yellow stain on the posterior part of the nacre not found on any other species. I have, however, found it on young *atrocostatus* in his collection. I am strongly inclined to believe that the *elliotii* is only a peculiar, vigorous variety of *perplicata*, but do not feel quite certain of it, and so let it go as a species with doubt.

37.—*REEVE, Conch. Icon., XVI, 1864, pl. v, fig. 20.—*B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) elliottii* LEA, Syn., 1870, p. 29.

Southern Georgia to Texas.

†QUADRULA UNDULATA Barnes.

**Unio undulatus* BARNES, Am. Jl. Sci., VI, 1823, p. 120, pl. II.—*HILDRETH, Am. Jl. Sci., XIV, 1828, p. 279.—*HANLEY, Test. Moll., 1842, p. 175; *Biv. Shells, 1843, p. 175, pl. XX, fig. 26.—*CATLOW and REEVE, Conch. Nom., 1845, p. 65.—*CHENU, Bib. Conch., 1st ser., III, 1845, p. 24, pl. v, figs. 1, 1a.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU, Manual, 1859, II, p. 143, fig. 704.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVI, fig. 399.—*CALKINS, Pr. Ottawa Acad. Sci., 1874, p. 45.—*B. H. WRIGHT, Check List, 1888.

**Margarita (Unio) undulatus* LEA, Syn., 1836, p. 12; 1838, p. 14.

**Margaron (Unio) undulatus* LEA, Syn., 1852, p. 20; 1870, p. 29.

**Mya undulata* EATON, Zool. Text-Book, 1826, p. 219.

**Unio undulata* VALENCIENNES, Rec. Obs. Zool. Anat., II, 1833, p. 229, pl. LIV, figs. 3, 3a, 3b.—*DESHAYES, Tr. Elem. Conch., 1839, p. 19, pl. XXX, figs. 8, 9.

**Quadrula undulata* BAKER, Moll. Chicago, Pt. I, 1898, p. 82, pl. XXII, figs. 1, 2; XII, fig. 1.

**Unio costatus* SAY, Am. Conch., VI, 1834.—*CONRAD, New F. W. Shells, 1834, p. 68.—*FERUSSAC, Guer. Mag., 1835, p. 27.—*CONRAD, Monog., II, 1836, p. 17, pl. VII.—*KUSTER, Conch. Cab., 1852, p. 54, pl. XI, fig. 4.—*REEVE, Conch. Icon., XVI, 1864, pl. IV, fig. 16.

**Plectomerus costatus* CONRAD, Pr. Acad. N. Sci. Phila., VI, 1853, p. 260.

**Unio plicatus* KUSTER, Conch. Cab., 1856, p. 137, pl. XL, fig. 3.—*CHENU, Manual, 1859, II, p. 143, fig. 706.

**Unio atrocostatus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVII, fig. 404.

†QUADRULA UNDULATA var. LATECOSTATA Lea.

**Unio latecostatus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 163;¹ *Tr. Am. Phil. Soc., X, 1848, p. 68, pl. I, fig. 2; *Obs., IV, 1848, p. 42, pl. I, fig. 2.—*KUSTER, Conch. Cab., 1861, p. 251, pl. LXXXIV, fig. 4.—*B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) laticostatus* LEA, Syn., 1852, p. 21.

**Margaron (Unio) laticostatus* LEA, Syn., 1870, p. 31.

**Unio laticostatus* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—*PÆTEL, Conch. Sam., III, 1890, p. 156.

†QUADRULA UNDULATA var. PILSBRYI Marsh.

**Unio pilsbryi* MARSH, Nautilus, V, 1891, p. 1.²—*PILSBRY, Pr. Ac. N. Sci. Phila., 1892, p. 131, pl. VIII, figs. 7, 8; *Nautilus, VII, 1893, pl. I, figs. 7, 8.³

¹More compressed and having finer, more delicate plications than the type. I have traced it northward into Kansas, where it gradually assumes the ordinary form of *undulata*. Occasional specimens are found in the Northern States which agree very well with the typical *latecostata*.

²I feel quite sure that this is only a peculiar variety of *undulata*.

³A peculiar shell belonging to Mr. J. H. Ferriss, of Joliet, Illinois, from Lake Winipeg, has been examined by the writer. It has unusually strong growth lines and is quite full in the ventral region, and has only the very faintest vestiges of plications below the lower part of the posterior ridge. I regard it as a smooth form of *Q. undulata*, and if any considerable number of such specimens were found it might be worthy of a varietal name.

Mississippi basin generally; St. Lawrence drainage; Red River of the North; Lake Winnipeg; Alabama River system; southwest to western Texas, the varieties in the southwestern area.

QUADRULA DIGITATA Morelet.

- * *Unio digitatus* MORELET, Test. Noviss., Pt. 2, 1851, p. 24.—* B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 563, pl. LX, fig. 1.

Rio Usumacinta, Guatemala.

† QUADRULA TRIUMPHANS B. H. Wright.

- * *Unio triumphans* B. H. WRIGHT, Nautilus, XI, 1898, p. 101.
* *Quadrula triumphans* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 83, pl. III, fig. 3.

Coosa River, St. Clair County, Alabama.

† QUADRULA BOYKINIANA Lea.

- * *Unio boykinianus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 288; * Tr. Am. Phil. Soc., VIII, 1842, p. 208, pl. XIII, fig. 22; * Obs., III, 1842, p. 46, pl. XIII, fig. 22.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* CHENU, Ill. Conch., 1858, pl. XXVII, figs. 2, 2a, 2b.—* KUSTER, Conch. Cab., 1861, p. 181, pl. LVII, fig. 1.—* SOWERBY, Conch. Icon., XVI, 1868, pl. I, fig. 1.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 146.
* *Margaron (Unio) boykinianus* LEA, Syn., 1852, p. 19; 1870, p. 28.
* *Plectomerus boykinianus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 260.
* *Unio boykinianus* PÆTEL, Conch. Sam., III, 1890, p. 145.

Chattahoochee River system; Claiborne, Alabama; Pine Barren Creek, Escambia County, Florida.

† QUADRULA HEROS Say.¹

- Unio heros* SAY, New Harm. Diss., II, No. 19, 1829, p. 291; * Am. Conch., VI, 1831, (part).—* CONRAD, New F. W. Shells, 1834, p. 69.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* CONRAD, Monog., XII, 1840, p. 107, pl. LIX.—* HANLEY, Test. Moll., 1842, p. 175; * Biv. Shells, 1843, p. 175, pl. XXII, fig. 28.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
* *Unio heros* var. *multiplicatus* PÆTEL, Conch. Sam., III, 1890, p. 155.
* *Unio undulatus* SAY, Am. Conch., I, 1831, pl. XVI (April).
† * *Unio multiplicatus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 70, pl. IV, fig. 2 (latter part of the year); * Obs., I, 1834, p. 80, pl. IV, fig. 2.—* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 155, pl. LIX, fig. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* CHENU, Ill. Conch., 1858, pl. IX, figs. 2, 2a, 2b; * Manual, 1859, II, p. 143, fig. 703.—* CALKINS, Pr. Ottawa Acad. Sci., 1874, p. 43.—* B. H. WRIGHT, Check List, 1888.

¹ There has been some dispute as to what name should be applied to the above species. In 1829 Say fully described in the New Harmony Disseminator a species which he called *Unio heros*, but did not figure it. Later in the American Conchology he placed it in the synonymy of *Unio undulatus* Barnes, though the figure which he gives as *undulatus* in that work and to which his description refers is plainly not Barnes's species, but the *heros* described in the Disseminator. In 1831 Dr. Lea described Say's species as *Unio multiplicatus*. Notwithstanding the fact that Say placed his own species in the synonymy, it is perfectly distinct and was properly characterized in his description, and his name will have to take the place of the better known one of Dr. Lea.

† **Margarita (Unio) multiplicatus* LEA, Syn., 1836, p. 12; 1838, p. 14.

**Margaron (Unio) multiplicatus* LEA, Syn., 1852, p. 20; 1870, p. 29.

† **Unio eightsii* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 306; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 367, pl. LXIV, fig. 192; *Obs., VIII, 1860, p. 49, pl. LXIV, fig. 192.—* B. H. WRIGHT, Check List, 1888.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 564.

**Margaron (Unio) eightsii* LEA, Syn., 1870, p. 29.

**Unio eighti* PÆTEL, Conch. Sam., III, 1890, p. 151.

**Unio atrocostatus* REEVE, Conch. Icon., XVI, 1864, pl. iv, fig. 13. (Change in Errata to *Unio heros*.)

Mississippi River system generally; Red River of the North; Tombigbee River, Alabama; southwest to New Leon, Mexico.

† QUADRULA NICKLINIANA Lea.

**Unio nicklinianus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 28, pl. i, fig. 1; *Obs., I, 1834, p. 140, pl. i, fig. 1.—* HANLEY, Test. Moll., 1842, p. 175; *Biv. Shells, 1843, p. 175, pl. XXI, fig. 52.—*CATLOW and REEVE, Conch. Nom., 1845, p. 61.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—*SOWERBY, Conch. Icon., XVI, 1866, pl. LIII, fig. 276.—*B. H. WRIGHT, Check List, 1888.—*FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 565.

**Margarita (Unio) nicklinianus* LEA, Syn., 1836, p. 12; 1838, p. 14.

**Margaron (Unio) nicklinianus* LEA, Syn., 1852, p. 19; 1870, p. 28.

**Plectomerus nicklinianus* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 261.

**Unio nicklinianus* KUSTER, Conch. Cab., 1861, p. 218, pl. LXXIII, fig. 3.—*PÆTEL, Conch. Sam., III, 1890, p. 161.

Mexico; Guatemala.

† QUADRULA STOLLI von Martens.

† **Unio stollii* VON MARTENS, Biol. Centrali-Amer., Mollusca, 1900, p. 492, pl. XXIX, fig. 2.

Rio de las Salinas, Guatemala; Moctezuma River Central America?¹

† QUADRULA NEISLERII Lea.

**Unio neislerii* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 165; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 212, pl. XXVI, fig. 93; *Obs., VII, 1859, p. 30, pl. XXVI, fig. 93.—

*B. H. WRIGHT, Check List, 1888.

**Margaron (Unio) neislerii* LEA, Syn., 1870, p. 29.

Flint River and Macon, Georgia.

¹Very close to *Quadrula nickliniana* Lea. According to von Martens in the above citation the *Unio nickliniana* of Kuster (Conch. Cab. Unio, new ed., p. 218, pl. LXXIII, fig. 3) is not the *nicklinianus* of Lea, but a new species which he calls *Unio stollii*. I can not agree with him, as it seems to me that Kuster's figure represents Lea's species very well. The figure and description of *U. stollii* given by von Martens represent a more elongated and more plicate shell than the typical *nicklinianus* of Lea, but I am doubtful whether it is distinct. Lea's shells of both forms are mostly from the Moctezuma River, which is in Central America according to Lea, but the only river of that name I can find is in New Leon, Mexico. Part of von Martens' work describing the Unionidae of Central America has been received as the Synopsis is going through the press, and for the most part too late to avail myself of the large amount of valuable information it contains.

(Group of *Quadrula trapezoides*.)

Shell rhomboid, inflated, with a high posterior ridge; beak sculpture coarse, irregular corrugations swollen to nodules on the posterior ridge; surface sculptured on posterior half with oblique ridges, which are sometimes corrugated, and with strong corrugations on post slope beak cavities only moderately deep; naere purple.

Animal with the gills slightly rounded below, inner the larger throughout, free from the abdominal sac; palpi enormous, long; branchial opening very large, finely papillose; anal opening with fine papillæ or crenulations.

† *QUADRULA TRAPEZOIDES* Lea.¹

* *Unio crassidens* var. *a*, LAMARCK, An. sans Vert., VI, 1819, p. 71.²

* *Plectomerus crassidens* var. *a*, CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 261.

* *Unio trapezoides* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 69, pl. III, fig. 1; * Obs., I, 1834, p. 79, pl. III, fig. 1.—* CONRAD, New F. W. Shells, 1834, p. 72.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* HANLEY, Test. Moll., 1842, p. 176; * Biv. Shells, 1843, p. 176, pl. XXII, fig. 29.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. x, figs. 2, 2a, 2b; Manual, 1859, II, p. 142, fig. 699.—* KUSTER, Conch. Cab., 1862, p. 274, pl. XCII, fig. 2.—* REEVE, Conch. Icon., XVI, 1864, pl. v, fig. 17.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL Conch. Sam., III, 1890, p. 169.

* *Margarita (Unio) trapezoides* LEA, Syn., 1836; p. 12; 1838, p. 14.

* *Margaron (Unio) trapezoides* LEA, Syn., 1852, p. 21; 1870, p. 31.

Unio interruptus SAY, Transylvania Journal, IV, 1831, p. 525; * Am. Conch., IV, 1832, pl. XXXIII, VI, 1834.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* DESHAYES, An. sans. Vert., 2d ed., VI, 1835, p. 556; 3d ed., II, 1839, p. 675.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 234.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 45, pl. XVII, figs. 1-3.

* *Unio dombeyana* VALENCIENNES, Rec. Obs. Zool., II, 1833, p. 227, pl. LIII, figs. 1, 1a, 1b.

Streams flowing into the Gulf of Mexico, from the Alabama River west to eastern Texas; northward in the Mississippi system to north-west Tennessee.

¹In 1820 Rafinesque applied the name *interrupta* to some, to me, unknown *Unio*, in Ann. Gen. Sci. Phys. Brux., but as he placed his species in the subgenus *Plagiola* of the genus *Obliquaria* and not in *Unio* at all, it does not have priority in the latter genus. In 1831 (December, according to Dr. Lea) Say described in the Transylvania Journal the species which Dr. Lea called *Unio trapezoides* as *Unio interruptus*. In the latter part of the same year (I am unable to ascertain the exact date) Dr. Lea published his name in the Transactions of the American Philosophical Society. As I have found it impossible to determine which name was actually published first I have used the better known—that of Dr. Lea.

²This is *Unio trapezoides* Lea according to Lea, who examined the Lamarckian types. See Synopsis, 4th ed., p. 37, footnote.

Section *QUADRULA* (Rafinesque, 1820) Agassiz.(Type, *Unio cylindricus* Say.)

Shell rhomboid; surface pustulous, with a high, rounded, or sharp posterior ridge; epidermis generally smooth and bright.

(Group of *Quadrula metanевра*.)

Shell quadrate or rhomboid, with a wide, rounded posterior ridge, above which, on the posterior slope is a decided radial furrow; whole surface except the anterior end generally pustulous, the sculpture of the posterior slope being often wrinkled; umbonal region high; epidermis shining, usually painted with a beautiful pattern of triangular spots, and sometimes chevron-shaped lines; hinge strong, the secondary lateral in right valve rather feeble, but there is often a faint third lateral above; cavity of the beaks deep and compressed.

Animal with the marsupium filling all four leaves of the branchiæ, the ova giving it a purplish tint; inner gills free from the abdominal sac for the greater part of their length; mantle bordered with black; branchial opening opposite the lobe of the shell, extending well on to its base; anal opening without papillæ.

†*QUADRULA CYLINDRICA* Say.

- * *Unio cylindricus* SAY, Nich. Encyc., II, 1816, pl. IV, fig. 3.—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 283, figs. 13, 13b.—* SHORT and EATON, Transylvania Jl., 1831, p. 76.—* SAY, Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 68.—* FERUSSAC, Guér. Mag., 1835, p. 27.—* HANLEY, Test. Moll., 1842, p. 182; * Biv. Shells, 1843, p. 182, pl. XX, fig. 31.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* KUSTER, Conch. Cab. Unio, 1861, p. 194, pl. LXII, figs. 1, 2.—* SOWERBY, Conch. Icon., XVI, 1867, pl. LX, fig. 300.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 150.—* CALL, Tr. Acad. Sci. St. Louis, VII, 1895, p. 15, pl. XII.
- * *Margarita (Unio) cylindricus* LEA, Syn., 1836, p. 17; 1838, p. 16.
- * *Margaron (Unio) cylindricus* LEA, Syn., 1852, p. 23; 1870, p. 35.
- * *Orthonymus cylindricus* AGASSIZ, Arch. für. Naturg., I, 1852, p. 48.
- * *Mya cylindrica* EATON, Zool. Text-Book, 1826, p. 219.
- * *Unio (Theliderma) cylindrica* SWAINSON, Treat. on Mal., 1840, p. 271, fig. 54c.
- * *Unio (Euryntia) solenoides* var. *cylindrica* RAFINESQUE, Ann. Gen. Sci. Phys. Brux., V, 1820, p. 298.
- * *Unio naviformis* LAMARCK, An. sans Vert., VI, 1819, p. 75.—* DESHAYES, Encyc. Méth., II, 1830, p. 580.—* VALENCIENNES, Rec. Obs. Zool., II, 1833, p. 233, pl. LIII, fig. 4.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 537; 3d ed., II, 1839, p. 669.—* ANTON, Verz. der. Conch., 1839, p. 14.—* REEVE, Conch. Syst., 1841, p. 118, pl. LXXXIX, fig. 7.¹
- * *Unio rugosus* CHENU, Man., 1859, II, p. 138, fig. 668.

Entire Ohio, Cumberland, and Tennessee river systems; west to Nebraska (Aughey, doubtful); south to Arkansas and Indian Territory.

¹Mr. B. H. Wright, Nautilus, XII, 1898, p. 6, has made a var. *strigillatus* of a compressed, very rough form of *cylindrica*. There seems to me to be an absolute gradation from cylindrical, nearly smooth specimens to this form.

†QUADRULA METANEVRA Rafinesque.

- * *Obliquaria (Quadrula) metanevra* RAFINESQUE, Ann. Gen. Sci. Brux., V, 1820, p. 305, pl. LXXXI, figs. 15, 16.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 18, pl. II, figs. 15, 16.
- * *Unio (Theliderma) metanevra* SWAINSON, Treat. on Mal., 1840, p. 268, figs. 50, 54b.
- * *Unio metanevra* SHORT and EATON, Transylvania Jl., 1831, p. 76.—* CONRAD, New F. W. Shells, 1834, p. 70.—* HANLEY, Test. Moll., 1842, p. 179; * Biv. Shells, 1843, p. 179, pl. XXI, fig. 31.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—* REEVE, Conch. Icon., XVI, 1864, pl. VII, fig. 25.—* PÆTEL, Conch. Sam., III, 1890, p. 159.
- * *Margarita (Unio) metanevra* LEA, Syn., 1836, p. 15; 1838, p. 15.
- * *Unio metanevra* SAY, Am. Conch., VI, 1834.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* CONRAD, Monog., I, 1835, p. 10, pl. v, fig. 2.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CALKINS, Pr. Ottawa Ac. N. Sci., 1874, p. 43.—* B. H. WRIGHT, Check List, 1888.—CALL, Tr. Acad. N. Sci. St. Louis, VII, 1895, p. 28, pl. x.
- * *Margaron (Unio) metanevra* LEA, Syn., 1852, p. 22; 1870, p. 33.
- * *Unio metanevra* CATLOW and REEVE, Conch. Nom., 1845, p. 61.
- * *Unio metanevra* KUSTER, Conch. Cab. Unio, 1852, p. 50, pl. x, fig. 4.—* PÆTEL, Conch. Sam., III, 1890, p. 159.
- *? *Unio nodosus* BARNES, Am. Jl. Sci. VI, 1823, p. 124, pl. vi, figs. 7, 7a, 7b.—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 281.—* CHENU, Manual, 1859, II, p. 138, fig. 663.
- * *Mya nodosa* EATON, Zool. Text-Book, 1826, p. 216.

†QUADRULA METANEVRA var. WARDII Lea.

- * *Unio wardii* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 392; * Jl. Ac. N. Sci. Phila., V, 1862, p. 187, pl. XXIV, fig. 257; * Obs., IX, 1863, p. 9, pl. XXIV, fig. 257.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXII, fig. 434.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 172.
- * *Margaron (Unio) wardii* LEA, Syn., 1870, p. 33.

Mississippi drainage area except its southern portion, extending to the Tennessee and Arkansas rivers.

†QUADRULA TUBEROSA Lea.

- * *Unio tuberosus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 286; * Tr. Am. Phil. Soc., VIII, 1842, p. 210, pl. xiv, fig. 25; * Obs., III, 1842, p. 48, pl. xiv, fig. 25.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. xxviii, figs. 7, 7a, 7b.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 170.
- * *Margaron Unio tuberosus* LEA, Syn., 1852, p. 22; 1870, p. 33.

Cumberland and Tennessee river systems.

¹ By common consent this species—which is a most striking one—has been allowed to stand to the credit of Rafinesque. The outlines of the figures bear some resemblance to the species.

² It is hard to tell from these figures just what Barnes had before him when he described this species. They are as much like *metanevra* as anything.

† *QUADRULA SPARSA* Lea.

* *Unio sparsus* LEA, Pr. Am. Phil. Soc., II, 1841, p. 82;¹ * Tr. Am. Phil. Soc., VIII, 1842, p. 242, pl. xxv, fig. 58; * Obs., III, 1842, p. 80, pl. xxv, fig. 58.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. xxvi, figs. 2, 2a, 2b.—* REEVE, Conch. Icon., XVI, 1864, pl. iv, fig. 14.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

* *Margarou (Unio) sparsus* LEA, Syn., 1852, p. 22; 1870, p. 33.

Holston and Clinch rivers, Tennessee.

† *QUADRULA INTERMEDIA* Conrad.

* *Unio intermedius* CONRAD, Monog., VII, 1836, p. 63, pl. xxxv, fig. 2; * Pr. Ac. N. Sci. Phil., VI, 1853, p. 250.—* HANLEY, Biv. Shells, 1856, p. 381, pl. xx, fig. 28.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* KUSTER, Conch. Cab. Unio, 1861, p. 213, pl. lxx, fig. 6.—* REEVE, Conch. Icon., XVI, 1864, pl. xiii, fig. 48.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 155.

* *Margarou (Unio) intermedius* LEA, Syn., 1852, p. 22; 1870, p. 33.

* *Unio kleinianus* KUSTER, Conch. Cab. Unio, 1861, p. 191, pl. lx, fig. 4.

Tennessee River system.

† *QUADRULA STAPES* Lea.

* *Unio stapes* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 77, pl. vii, fig. 8.—* Obs., I, 1834, p. 87, pl. vii, fig. 8.—* CONRAD, New F. W. Shells, 1834, p. 71.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* CONRAD, Monog., VII, 1836, p. 62, pl. xxxv, fig. 1.—* HANLEY, Test. Moll., 1842, p. 179; * Biv. Shells, 1843, p. 179, pl. xxii, fig. 38.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. xv, figs. 5, 5a, 5b; * Manual, 1859, II, p. 142, fig. 997.—* REEVE, Conch. Icon., XVI, 1864, pl. xiii, fig. 52.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

* *Margarita (Unio) stapes* LEA, Syn., 1836, p. 15; 1838, p. 13.

* *Margarou (Unio) stapes* LEA, Syn., 1852, p. 22; 1870, p. 33.

* *Unio retusus* SAY, Am. Conch., VI, 1834.

* *Unio stegarius* KUSTER, Conch. Cab. Unio, 1861, p. 211, pl. lxx, fig. 3.

Alabama and Tombigbee rivers.

Section *THELIDERMA* (Swainson, 1840) Simpson.²

(Type, *Unio lachrymosus* Lea.)

Shell rounded, quadrate to rhomboid, solid, pustulous; beaks rather prominent; beak sculpture consisting of a few rather coarse, subparallel ridges; anterior end rounded or slightly subtruncate above; base often arcuate; posterior end truncate, high and angled behind the ligament; epidermis scarcely rayed, never pointed; beak cavities rather deep.

¹ It is likely that this is only a small, rather smooth variety of *tuberosus*.

² Swainson's first species of this group in the text of the Treatise is *Unio irroratus*, and after this *metaneurus* and *cylindricus*. In the final classification in the same work he includes in the group *lachrymosus* and *pustulosus*, and by the process of elimination I use the name in order to save coining a new one.

(Group of *Quadrula lachrymosa*.)

Shell quadrate or rhomboid, generally with a sharp, well-developed posterior ridge, in front of which is a wide, shallow, radial excavation; base incurved, posterior end sharply truncate, generally slightly biangular below, high and distinctly angled behind the ligament; posterior tubercles often in somewhat radiating rows; earlier beak sculpture consisting of ridges nearly parallel with the growth lines, but later on becoming doubly looped and blending into the general sculpture of the shell; epidermis smooth; pseudocardinals strong; laterals straight; nacre white.

Animal with all four gills used as a marsupium throughout, inner much the larger, free from the abdominal sac nearly or quite their whole length; palpi very large; branchial opening generally having clustered, often branching papillæ; anal opening smooth; superanal opening closed below.

†*QUADRULA ASPER* Lea.

* *Unio asper* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 85, pl. IX, fig. 15; * Obs., I, 1834, p. 95, pl. IX, fig. 15.—* CONRAD, New F. W. Shells, 1834, p. 67.—* HANLEY, Test. Moll., 1842, p. 179; * Biv. Shells, 1843, p. 179, pl. XXII, fig. 37.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. XV, figs. 4, 4a, 4b; * Manual, 1859, II, p. 142, fig. 695.—* REEVE, Conch. Icon., XVI, 1864, pl. V, fig. 18.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) asper* LEA, Syn., 1836, p. 15; 1838, p. 15.

* *Margaron (Unio) asper* LEA, Syn., 1852, p. 22; 1870, p. 33.

* *Unio nobilis* CONRAD, Jl. Ac. N. Sci. Phila., II, 1854, p. 297, pl. XXVII, figs. 2, 3.

* *Unio turgidus* REEVE, Conch. Icon., XVI, 1864, pl. III, fig. 10.

* *Unio apiculatus* var. *asper* PÆTEL, Conch. Sam., III, 1890, p. 144.

Streams flowing into the Gulf of Mexico from Alabama west to central Texas, and northward to the Verdigris River, Kansas.

†*QUADRULA LACHRYMOSA* Lea.

Unio lachrymosus LEA, Tr. Am. Phil. Soc., III, 1828, p. 272, pl. VI, fig. 8;¹ Obs., I, 1834, p. 14, pl. VI, fig. 8.—* SHORT and EATON, Transylvania Jl., 1831, p. 76.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* HANLEY, Test. Moll., 1842, p. 177; * Biv. Shells, 1843, p. 177, pl. XX, fig. 38.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. XV, figs. 6; 6a, 6b; Manual, 1859, II, p. 142, fig. 698.—* REEVE, Conch. Icon., XVI, 1864, pl. IX, fig. 33.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 156.

* *Margarita (Unio) lachrymosus* LEA, Syn., 1836, p. 14; 1838, p. 15.

* *Margaron (Unio) lachrymosus* LEA, Syn., 1852, p. 21; 1870, p. 32.

* *Unio lachrymosus* KUSTER, Conch. Cab., 1854, p. 70, pl. XVII, fig. 3.

¹Published as a separate in 1827. I have not seen this. Title from Sabin's Bibliotheca America.

- * *Quadrula lachrymosa* BAKER, Moll. Chicago, Pt. I, 1898, p. 83, pl. xxv, fig. 1; XII, fig. 2.
- † * *Unio asperimus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 71, pl. v, fig. 3; ¹ * Obs., I, 1834, p. 81, pl. v, fig. 3.—* HANLEY, Test. Moll., 1842, p. 178; * Biv. Shells, 1843, p. 178, pl. XXI, fig. 12.—* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 156, pl. LIX, figs. 2, 3.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. ix, figs. 1, 1a, 1b.—* B. H. WRIGHT, Check List, 1888.
- * *Margarita (Unio) asperimus* LEA, Syn., 1836, p. 14; 1838, p. 15.
- * *Margaron (Unio) asperimus* LEA, Syn., 1852, p. 21; 1870, p. 33.
- * *Unio quadrulus* SAY, Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 71.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 256.—* KUSTER, Conch. Cab. Unio, 1861, p. 184, pl. LVIII, fig. 1.
- * *Unio quadratus* REEVE, Conch. Icon., XVI, 1864, pl. vi, fig. 24.

† **QUADRULA LACHRYMOSA var. LUNULATA** Pratt.

- * *Unio lunulatus* PRATT, Proc. Day. Acad. Nat. Sci., I, 1876, p. 167, pl. XXXI, fig. 1.²

Entire Mississippi River drainage; various localities in the St. Lawrence basin; Red River of the North; southwest into eastern Texas.

† **QUADRULA COUCHIANA** Lea.

- * *Unio couchianus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 305; Jl. Ac. N. Sci. Phila., IV, 1860, p. 371, pl. LXVI, fig. 196; * Obs., VIII, 1860, p. 53, pl. LXVI, fig. 196.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXI, fig. 429.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 149.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 566.—* SIMPSON, Proc. U. S. Nat. Mus., XIX, 1896, p. 370.
- * *Margaron (Unio) couchianus* LEA, Syn., 1870, p. 54.

Rio Salado, Mexico; southwestern Texas.

† **QUADRULA FRAGOSA** Conrad.

- * *Unio fragosus* CONRAD, Monog., II, 1836, p. 12, pl. vi, fig. 2; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* KUSTER, Conch. Cab. Unio., 1861, p. 173, pl. LV, fig. 1.—* REEVE, Conch. Icon., XVI, 1864, pl. i, fig. 2; VII, fig. 27.—* B. H. WRIGHT, Check List, 1888.—* PETEL, Conch. Sam., III, 1890, p. 153.
- * *Margarita (Unio) fragosus* LEA, Syn., 1836, p. 14; 1838, p. 15.
- * *Margaron (Unio) fragosus* LEA, Syn., 1852, p. 22; 1870, p. 33.
- * *Unio fragosa* CATLOW and REEVE, Conch. Nom., 1845, p. 59.

¹ Dr. Lea separated this from his *lachrymosus* because the latter had tearlike nodules, while those of *asperimus* were simply elevated. The two forms absolutely run together, and specimens may be found in which both varieties of nodules occur. There are forms which are almost smooth, and there is great diversity in the matter of inflation. Dr. Lea finally came to doubt whether the *asperimus* was valid.

² I think that this is a solid, inflated variety of *lachrymosus*. From the figure, Pratt's specimens seem to have few tubercles; but I have seen quite a number of shells from the Ohio River north to Minnesota which resemble it in shape, but are strongly pustulous throughout. I was at first inclined to believe these were varieties of *asper*, but I now regard them as a form of *lachrymosus*, and they may perhaps be referred to Pratt's variety. He describes his species as having a large lunule, but individuals vary greatly in this character.

- * *Unio tragosus* HANLEY, Test. Moll., 1842, p. 178; * Biv. Shells, 1843, p. 178, pl. xx, fig. 40.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.

Ohio, Cumberland, and Tennessee river systems; westward probably to Minnesota, Nebraska, and Kansas.

† *QUADRULA FORSHEYI* Lea.

- * *Unio forsheyi* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 155; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 357, pl. LX, fig. 182; * Obs., VIII, 1860, p. 39, pl. LX, fig. 182.—* REEVE, Conch. Icon., XVI, 1864, pl. VI, fig. 21.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 153.
* *Margaron (Unio) forsheyi* LEA, Syn., 1870, p. 32.

Alabama to Texas, in streams flowing into the Gulf of Mexico.

† *QUADRULA SPECIOSA* Lea.

- * *Unio speciosus* LEA, Pr. Ac. N. Sci. Phila., VI, 1862, p. 168; * Jl. Ac. N. Sci. Phila., V, 1862, p. 207, pl. XXXI, fig. 276; * Obs., IX, 1863, p. 29, pl. XXXI, fig. 276.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIV, fig. 447.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.
* *Margaron (Unio) speciosus* LEA, Syn., 1870, p. 33.

Kansas, south to Texas.

† *QUADRULA APICULATA* Say.

- * *Unio apiculatus* SAY, New Harm. Diss., II, No. 2, 1829, p. 309; * Am. Conch., VI, 1834, pl. LI.²—* CONRAD, New F. W. Shells, 1834, p. 67.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* CONRAD, Monog., IX, 1837, p. 78, pl. XLIV, fig. 1.—* HANLEY, Test. Moll., 1842, p. 178; * Biv. Shells, 1843, p. 178, pl. XXIII, fig. 51.—* CATLOW and REEVE, Conch. Nom., 1845, p. 55.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Manual, 1859, II, p. 142, fig. 696.—* REEVE, Conch. Icon., XVI, 1864, pl. III, fig. 11.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 144.
* *Margarita (Unio) apiculatus* LEA, Syn., 1836, p. 15; 1838, p. 15.
* *Margaron (Unio) apiculatus* LEA, Syn., 1852, p. 22; 1870, p. 33.
* *Unio nobilis* CONRAD (part), Jl. Ac. N. Sci. Phila., 1854, p. 297, pl. XXVII, figs. 2, 3.³

Louisiana to Texas.

† *QUADRULA RUMPHIANA* Lea.

- * *Unio rumphianus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 276, pl. XXII, fig. 34; Obs., V, 1852, p. 32, pl. XXII, fig. 34.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 166.
* *Margaron (Unio) rumphianus* LEA, Syn., 1852, p. 21; 1870, p. 33.
† * *Unio blandianus* LEA,⁴ Pr. Ac. N. Sci. Phila., VIII, 1856, p. 263; * Jl. Ac. N. Sci.

¹ Written *tragosus* in text, no doubt a typographical error, as it is written correctly in the explanation of the plate.

² Say says that *U. asper* Lea is the same as his *apiculatus*. I think he is wrong.

³ One specimen figured (fig. 2) is *Q. apiculata*. The other (fig. 3) is a nondescript, possibly *Q. asper*, or it may be the form of *Q. lachrymosus* which I have placed under var. *lunulatus* Pratt.

⁴ Certain forms of *rumphianus* are very different from *blandianus*, being more inflated, and showing a high, biangulate posterior ridge and a deep radial sinus in front of it, while specimens of the latter are much less inflated and have a low, somewhat rounded, posterior ridge. But the difference between the two seems to be absolutely bridged by intermediate specimens.

Phila., IV, 1858, p. 65, pl. XI, fig. 47; * Obs., VI, 1858, p. 65, pl. XI, fig. 47.—
* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVIII, fig. 405.—* B. H. WRIGHT,
Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 46.

* *Margaron (Unio) blandianus* LEA, Syn., 1870, p. 33.

Western Georgia; Alabama in the Gulf drainage.

(Group of *Quadrula pustulosa*.)

Shell round quadrate, truncated behind, full and angled back of the ligament; beaks high; beak sculpture a few coarse ridges which are swollen at the somewhat rounded posterior ridge; surface generally more or less pustulous but sometimes entirely smooth and in some cases slightly corrugated; epidermis often having a broad, faint, green ray; hinge strong; nacre white or purple.

Animal with the marsupium occupying all four gills throughout; inner gills the larger, free the whole or the greater part of their length from the abdominal sac; branchial opening having clustered, often branching, papillæ.

† *QUADRULA PUSTULOSA* Lea.

* *Unio pustulosus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 76, pl. VII, fig. 7; * Obs., I, 1834, p. 86, pl. VII, fig. 7.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* HANLEY, Test. Moll., 1842, p. 180; * Biv. Shells, 1843, p. 180, pl. XXI, fig. 34.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* CHENU, Ill. Conch., 1858, pl. XXIII, figs. 2, 2a, 2b.—* CALKINS, Pr. Ottawa Acad. N. Sci., 1874, p. 44.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) pustulosus* LEA, Syn., 1836, p. 15; 1838, p. 15.

* *Margaron (Unio) pustulosus* LEA, Syn., 1852, p. 22; 1870, p. 33.

* *Quadrula pustulosa* BAKER, Moll. Chicago, Pt. 1, 1898, p. 86, pl. XXV, fig. 2; XXVIII, fig. 13.

* *Unio verrucosa* VALENCIENNES, Rec. Obs. Zool. Anat., II, 1833, p. 231, pl. LIII, fig. 2.

* *Unio nodulosus* SAY, Am. Conch., VI, 1834.

* *Unio prasinus* CONRAD,¹ New F. W. Shells, 1834, p. 44, pl. III, fig. 1, p. 71.—

* FERUSSAC, Guer. Mag., 1835, p. 29.—* MÖLLER, Syn. Nov. Gen., 1836, p. 208.—* CONRAD, Monog., IX, 1837, p. 79, pl. XLIV, fig. 2.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 22, pl. III, fig. 1.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—* REEVE, Conch. Icon., XVI, 1864, pl. VII, figs. 26, 26a, 26b.—* B. H. WRIGHT, Check List, 1888.

* *Unio bullatus* var. *prasinus* PÆTEL, Conch. Sam., III, 1890, p. 146.

* *Unio bullatus* CONRAD, New F. W. Shells, 1834, p. 68; Monog., X, 1838, p. 82, pl. XLV, fig. 2.—* KUSTER, Conch. Cab. Unio, 1852, p. 47, pl. IX, fig. 3.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* REEVE, Conch. Icon., XVI, 1864, pl. XII, fig. 43.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

* *Unio dorfeuillianus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 73, pl. XVII, fig. 54;

* Obs., II, 1838, p. 73, pl. XVII, fig. 54.—* TROSCHIEL, Arch. für Naturg., V, 1839, II, p. 237.—* HANLEY, Test. Moll., 1842, p. 179; * Biv. Shells, 1843, p. 179, pl. XXIII, fig. 10.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 248.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. XXV, figs. 5, 5a, 5b.—* REEVE, Conch. Icon., XVI, 1864, pl. X, fig. 38.—* B. H. WRIGHT, Check List,

¹ A smooth form of *pustulosus*, according to the type in Philadelphia Academy.

- 1888.—* CALL (part), Tr. Ac. N. Sci. St. Louis, VII, 1895, p. 40, pls. XII, XIII, XIV, XV, figs. 1, 2.
- * *Margarita (Unio) dorfeuillianus* LEA, Syn., 1838, p. 15.
- * *Margaron (Unio) dorfeuillianus* LEA, Syn., 1852, p. 22; 1870, p. 33.
- * *Unio uber* CONRAD, Am. Jl. Conch., II, 1866, p. 279, pl. xv, fig. 16.¹
- †* *Unio schoolcraftensis* LEA, Tr. Am. Phil. Soc., V, 1834, p. 37, pl. III, fig. 9;² *Obs., I, 1834, p. 149, pl. III, fig. 9.—*FERUSSAC, Guer. Mag., 1835, p. 29.—*HANLEY, Test. Moll., 1842, p. 178; *Biv. Shells, 1843, p. 178.—*CATLOW and REEVE, Conch. Nom., 1845, p. 63.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—*SOWERBY, Conch. Icon., XVI, 1868, pl. I, fig. 3; XIII, fig. 47.
- * *Margarita (Unio) schoolcraftensis* LEA, Syn., 1836, p. 15; 1838, p. 15.
- * *Margaron (Unio) schoolcraftensis* LEA, Syn., 1852, p. 22.
- * *Unio bullatus* var. *schoolcraftensis* PÆTEL, Conch. Sam., III, 1890, p. 146.
- * *Margaron (Unio) schoolcraftii* LEA, Syn., 1870, p. 33.
- * *Unio schoolcraftii*, B. H. WRIGHT, Check List, 1888.

†QUADRULA PUSTULOSA var. PERNODOSA Lea.

- * *Unio pernodosus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 163; *Tr. Am. Phil. Soc., X, 1848, p. 71, pl. III, fig. 8; *Obs., IV, 1848, p. 45, pl. III, fig. 8.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 254.—*REEVE, Conch. Icon., XVI, 1864, pl. XII, fig. 46.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 163.
- * *Margaron (Unio) pernodosus* LEA, Syn., 1870, p. 34.
- †* *Unio asperatus* LEA,³ Pr. Ac. N. Sci. Phila., V, 1861, p. 41; *Jl. Ac. N. Sci. Phila., V, 1862, p. 68, pl. VII, fig. 218; *Obs., VIII, 1862, p. 72, pl. VII, fig. 218.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXV, fig. 450.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 145.
- * *Margaron (Unio) asperatus* LEA, Syn., 1870, p. 33.

†QUADRULA PUSTULOSA var. KEINERIANA Lea.

- * *Unio keinerianus* LEA, Pr. Am. Phil. Soc., V, 1852, p. 251; *Tr. Am. Phil. Soc., X, 1852, p. 281, pl. XXIII, fig. 40; *Obs., V, 1852, p. 37, pl. XXIII, fig. 40.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 156.
- * *Margaron (Unio) keinerianus* LEA, Syn., 1852, p. 22; 1870, p. 34.

Entire Mississippi drainage; Michigan; Lake Erie;? Alabama River system; westward to central Texas; the varieties in streams draining into the Gulf of Mexico.

¹ Probably a smooth form of *pustulosus*.

² The type is a nearly smooth, compressed form of *pustulosus*, I am sure. Lea includes with it in his collection a number of strongly *pustulosus*, compressed shells. *Quadrula pustulosa* is a very widespread, abundant, and variable species. A large, solid, inflated, and somewhat quadrate form has been called *Unio dorfeuillianus* by Dr. Lea, and various smooth races have received names, but they so completely blend into the typical form that it does not seem to me they are worthy of varietal names. A somewhat triangular variety, often having a broad, faint ray, is found in the Ohio River and southward which probably equals the variety *pernodosa*.

³ Seems to be about the same as *U. pernodosus*, only a little smoother. These two are more rounded posteriorly, and are doubtfully worthy of a varietal name.

† *QUADRULA COOPERIANA* Lea.

† *Unio cooperianus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 61, pl. VIII, fig. 21; *Obs., I, 1834, p. 173, pl. VIII, fig. 21.—* HANLEY, Test. Moll., 1842, p. 180; *Biv. Shells, 1843, p. 180, pl. XXI, fig. 1.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* KUSTER, Conch. Cab. Unio, 1861, p. 183, pl. LVII, fig. 5.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 149.

* *Margarita (Unio) cooperianus* LEA, Syn., 1836, p. 16; 1838, p. 15.

* *Margaron (Unio) cooperianus* LEA, Syn., 1852, p. 22; 1870, p. 34.

* *Unio striatus* REEVE, Conch. Icon., XVI, 1864, pl. VIII, fig. 30.¹

Ohio, Cumberland, and Tennessee river systems. Reported by Keyes at Muscatine, Iowa, but this is probably an error.

† *QUADRULA MORTONI* Conrad.

* *Unio mortoni* CONRAD, Monog. II, 1836, p. 11, pl. VI, fig. 1.—* KUSTER, Conch. Cab. Unio, 1852, p. 51, pl. XI, fig. 1.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 252.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 160.

† * *Unio turgidus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 11, pl. V, fig. 11; *Obs., II, 1838, p. 11, pl. V, fig. 11.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 234.—* HANLEY, Test. Moll., 1842, p. 180; *Biv. Shells, 1843, p. 180, pl. XXI, fig. 51.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* CHENU, Ill. Conch., 1858, pl. XXV, figs. 1, 1a, 1b.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 170.

* *Margarita (Unio) turgidus* LEA, Syn., 1836, p. 16; 1838, p. 15.

* *Margaron (Unio) turgidus* LEA, Syn., 1852, p. 22; 1870, p. 34.

Lower Mississippi River drainage as far north as western Tennessee and Indian Territory; west into eastern Texas.

† *QUADRULA SPHÆRICA* Lea.

* *Unio sphericus* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 145; *Jl. Ac. N. Sci. Phila., 1869, p. 319, pl. LI, fig. 132; *Obs., XII, 1869, p. 78, pl. LI, fig. 131.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

* *Margaron (Unio) sphericus* LEA, Syn., 1870, p. 33.

Streams of Alabama and Mississippi flowing into the Gulf of Mexico.

† *QUADRULA PUSTULATA* Lea.

* *Unio pustulatus* LEA, Tr. Am. Phil. Soc., 1834, p. 79, pl. VII, fig. 9; *Obs., I, 1834, p. 89, pl. VII, fig. 9.—* HANLEY, Test. Moll., 1842, p. 178; *Biv. Shells, 1843, p. 178, pl. XXII, fig. 36.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. XV, figs. 8, 8a, 8b.—* CALKINS, Pr. Ottawa Ac. N. Sci., 1874, p. 44.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) pustulatus* LEA, Syn., 1836, p. 15; 1838, p. 15.

* *Margaron (Unio) pustulatus* LEA, Syn., 1852, p. 22; 1870, p. 33.

¹ Changed in errata to *cooperianus*.

* *Unio (Theliderma) pustulata* SWAINSON, Treat. on Mal., 1840, p. 271, fig. 54d.

* *Unio nodulatus* SAY, Am. Conch., VI, 1834.—* CONRAD, New F. W. Shells, 1834, p. 70.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* CONRAD, Monog., 1837, p. 80, pl. XLV, fig. 1.—* KUSTER, Conch. Cab., 1861, p. 254, pl. LXXXVI, fig. 1.—

* REEVE, Conch. Icon., XVI, 1864, pl. XIII, fig. 51.

* *Unio nodulatus* RAFINESQUE var. *pustulatus* PÆTEL, Conch. Sam., III, 1890, p. 161.

Ohio, Cumberland, and Tennessee river systems; Mississippi River and tributaries from eastern Iowa south to Louisiana; southern Michigan.

† QUADRULA NODIFERA Conrad.¹

* *Unio nodiferus* CONRAD, Pr. Ac. N. Sci. Phila., I, 1841, p. 19; * Jl. Ac. N. Sci. Phila., VIII, 1842, p. 179; I, 1850, p. 277, pl. XXXVIII, figs. 4, 8; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.

Louisiana and eastern Texas.

† QUADRULA CAHABENSIS Lea.

* *Unio cahabensis* LEA, Pr. Ac. N. Sci. Phila., XXIII, 1871, p. 190; * Jl. Ac. N. Sci. Phila., VIII, 1874, p. 17, pl. v, fig. 14; * Obs., XIII, 1874, p. 21, pl. v, fig. 14.—

* B. H. WRIGHT, Check List, 1888.

Cahawba River, Alabama.

† QUADRULA VALLATA Lea.

* *Unio vallatus* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 145; * Jl. Ac. N. Sci. Phila., 1869, p. 315, pl. I, fig. 128; * Obs., XII, 1869, p. 75, pl. I, fig. 128.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) callatus* LEA, Syn., 1870, p. 34.

Alabama and Black Warrior rivers, Alabama.

† QUADRULA REFULGENS Lea.

* *Unio refulgens* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 145; * Jl. Ac. N. Sci. Phila., VI, 1869, p. 317, pl. LI, fig. 130; * Obs., XII, 1869, p. 77, pl. LI, fig. 130.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) refulgens* LEA, Syn., 1870, p. 34.

* *Unio pustulosus* CALL (part), Tr. Ac. Sci. St. Louis, VII, 1895, p. 42, pl. xv, figs. 3, 4.

Mississippi and probably Alabama.

† QUADRULA HOUSTONENSIS Lea.

* *Unio houstonensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 155; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 358, pl. LX, fig. 183; * Obs., VIII, p. 40, pl. LX, fig. 183.—

* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXI, fig. 425.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) houstonensis* LEA, Syn., 1870, p. 55.

Texas, Louisiana, and southern Arkansas. Verdigris River, Kansas?

¹ There are six shells bearing the name *Unio nodiferus* Conrad in the Philadelphia Academy of Natural Sciences. Part of them are the same as figured in the Journal, and the rest are probably *mortoni*.

†QUADRULA PETRINA Gould.

* *Unio petrinus* GOULD, Pr. Bost. Soc. N. Hist., V, 1855, p. 228; * *Otia* Conch., 1862, p. 218.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) petrinus* LEA, Syn., 1870, p. 55.

Texas. Cragin's localities, Verdigris and Neosho rivers, Kansas, are probably erroneous.

†QUADRULA AUREA Lea.

* *Unio aureus* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 112; * *Jl. Ac. N. Sci. Phila.*, V, 1862, p. 195, pl. XXVI, fig. 264; * *Obs.*, IX, 1863, p. 17, pl. XXVI, fig. 264.—

* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIII, fig. 438.—* B. H. WRIGHT, Check List, 1888.—* PLETZ, Conch. Sam., III, 1890, p. 145.

* *Margaron (Unio) aureus* LEA, Syn., 1870, p. 37.

* *Unio bolli* CALL, Am. Naturalist, XV, 1881, p. 390.¹

Texas.

†QUADRULA PAUPERCULA Lea.

* *Unio pauperculus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 39; * *Jl. Ac. N. Sci. Phila.*, V, 1862, p. 99, pl. xv, fig. 247; * *Obs.*, VIII, 1862, p. 103, pl. xv, fig. 247.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) pauperculus* LEA, Syn., 1870, p. 55.²

Texas.

(Group of *Quadrula infucata*.)

Shell small, rounded rhomboid, with a low, distinct posterior ridge, truncated behind and angled back of the ligament; beaks moderately elevated, subcentral; beak sculpture, four or five coarse, subparallel ridges which curve upward behind, with fine, radial ridges between them and the ligament; surface blackish, sculptured with zigzag or chevron-shaped corrugations which often become pustulous below; posterior slope having radial, broken wrinkles; pseudocardinals small, stumpy; secondary lateral of the right valve present; beak cavities moderately deep, slightly compressed; anterior scars small, smooth; nacre rather dull, purplish.

Animal with the marsupium occupying all four leaves of the gills; inner gills the larger, free from the abdominal sac only part of their length; anal opening without papillæ.³

¹ An inflated form.

² Usually entirely free from sculpture excepting faint corrugations on the posterior slope. A specimen belonging to Mr. William A. Marsh, from Village Creek, Texas, shows slight tuberculation on the disks.

³ I have been puzzled as to the relationship of this group and have been strongly inclined to place it near *chickasawhensis*, but I now believe it is more closely allied to the *Pustulosa* group. Its posterior truncation, the prominent angle behind the ligament, and beak sculpture lead me to place it here. The general sculpture of *Q. petrina* approaches that of these species.

† *QUADRULA INFUCATA* Conrad.

- * *Unio infucatus* CONRAD, New F. W. Shells, 1834, p. 45, pl. III, fig. 2, p. 70.—* MÖLLER, Syn. Nov. Gen., 1836, p. 208.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Test. Moll., 1842, p. 203; * Biv. Shells, 1843, p. 203.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 22, pl. II, fig. 6.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* CONRAD, Pr. Ac. N. Sei. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* KUSTER, Conch. Cab. Unio, 1861, p. 176, pl. LV, fig. 5.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XL, fig. 221.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 155.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 409, pl. XLIX, figs. 1-3, 6.
- * *Margarita (Unio) infucatus* LEA, Syn., 1836, p. 34; 1838, p. 23.
- * *Margaron (Unio) infucatus* LEA, Syn., 1852, p. 20; 1870, p. 30.
- * *Unio securiformis* CONRAD, ¹Ann. and Mag., IV, 1849, p. 300; Pr. Ac. N. Sei. Phila., * I, 1850, p. 275, pl. XXXVII, fig. 1; * IV, 1849, p. 152; * VI, 1852, p. 257.—* PÆTEL, Conch. Sam., III, 1890, p. 167.

Chattahoochee and Flint rivers, Georgia.

† *QUADRULA KLEINIANA* Lea.²

- * *Unio kleinianus* LEA, Pr. Am. Phil. Soc., V, 1852, p. 251; * Tr. Am. Phil. Soc., X, 1852, p. 265, pl. XVII, fig. 18; * Obs., V, 1852, p. 21, pl. XVII, fig. 18.—* CONRAD, Pr. Ac. N. Sei. Phila., VI, 1853, p. 257.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* REEVE, Conch. Icon., XVI, 1864, pl. II, fig. 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 156.
- * *Margaron (Unio) kleinianus* LEA, Syn., 1852, p. 20; 1870, p. 30.
- * *Unio sparsus* KUSTER, Conch. Cab. Unio, 1861, p. 252, pl. LXXXV, fig. 2.

Southern Georgia; northern Florida.

Section *FUSCONAIA* Simpson, 1900.

(Type, *Unio trigonus* Lea.)

Shell round, rhomboid, triangular, or short elliptical, with a moderate posterior ridge; beaks high and full, curved inward and forward, sculptured with a few coarse, parallel ridges which curve upward behind; epidermis dark; surface not sculptured; hinge plate of moderate width; pseudocardinals strong; naere white, salmon, or purple.

Animal having all four gills occupied throughout as a marsupium, filled with pink or purplish ova when gravid; inner gills much the wider in front, free generally from the abdominal sac; outer gills cut away slopingly in front.

(Group of *Quadrula beadleiana*.)

Shell moderately solid, somewhat triangularly rounded, bluntly pointed and sometimes slightly biangular at post base, with a more or less developed posterior ridge, in front of which it is full; beaks not

¹ A rather smooth *infucatus*.

² In the Proceedings of the U. S. National Museum, XV, 1892, p. 404, I united this species with *Unio infucatus* Conrad. Further study of additional material and of the animals which show some differences has induced me to separate the two.

very high; beak sculpture very coarse, concentric ridges;¹ disks irregularly concentrically striate; epidermis rich, dark chestnut; hinge plate rather narrow; pseudocardinals radial, stumpy; cavity of the beaks only moderately deep.

No gravid specimens have been seen, but those examined do not seem to differ from others of the genus.

†QUADRULA CHICKASAWHENSIS Lea.

* *Unio chickasawhensis* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 39; *Jl. Ac. N. Sci. Phila., V, 1862, p. 101, pl. XVI, fig. 250; *Obs., VIII, 1862, p. 105, pl. XVI, fig. 250.—* B. H. WRIGHT, Check List, 1888.

Margaron (Unio) chickasawhensis LEA, Syn., 1870, p. 55.

Mississippi and Louisiana.

†QUADRULA SUCCISSA Lea.

* *Unio succissus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 275, pl. XXI, fig. 32; *Obs., V, 1852, p. 31, pl. XXI, fig. 32.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIII, fig. 174.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 169.—* SIMPSON, Pr. U. S. Nat. Mus., XV, 1892, p. 29, pl. LXXI, fig. 5.

* *Margaron (Unio) succissus* LEA, Syn., 1852, p. 24; 1870, p. 38.

†* *Unio cacao* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 154; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 344, pl. LVI, fig. 169; *Obs., VIII, 1860, p. 26, pl. LVI, fig. 169.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) cacao* LEA, Syn., 1870, p. 54.

West Florida; southern Alabama.

†QUADRULA PUMILA Lea.

* *Unio pumilus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 23, pl. VII, fig. 17; *Obs., II, 1838, p. 23, pl. VII, fig. 17.—* TROSCHER, Arch. für Naturg., V, 1839, Pt. 2, p. 234.—* HANLEY, Test. Moll., 1842, p. 185; *Biv. Shells, 1843, p. 185, pl. XXIII, fig. 13.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 255.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XIX, figs. 1, 1a, 1b.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVI, fig. 198.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 164.

* *Margarita (Unio) pumilus* LEA, Syn., 1836, p. 20; 1838, p. 17.

* *Margaron (Unio) pumilus* LEA, Syn., 1852, p. 24; 1870, p. 37.

- Black River, North Carolina. Only the type shell, a small specimen, not in very good condition, has been seen by me.

†?QUADRULA UTRICULUS Lea.

* *Unio utriculus* LEA, Pr. Am. Phil. Soc., IV, 1845, p. 164; *Tr. Am. Phil. Soc., X, 1848, p. 69, pl. I, fig. 3; Obs., IV, 1848, p. 43, pl. I, fig. 3.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 259.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) utriculus* LEA, Syn., 1870, p. 46.

North Carolina. The specimen in the Lea collection which came from Arkansas is not this species at all.

¹According to Marsh, who has examined the beak sculpture of *Q. askewi*. I have never seen a specimen of the group in which it was not all eroded away.

† *QUADRULA BEADLEIANA* Lea.

- * *Unio beadleianus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 39; * Jl. Ac. N. Sci. Phila., V, 1862, p. 94, pl. XIV, fig. 242; * Obs., VIII, 1862, p. 98, pl. XIV, fig. 242.—* B. H. WRIGHT, Check List, 1888.

* *Margaron* (*Unio*) *beadleianus* LEA, Syn., 1870, p. 55.

Mississippi to eastern Texas.

† *QUADRULA ASKEWI* Marsh.

- * *Unio askewi* MARSH, Nautilus, X, 1896, p. 91; X, 1897, pl. 1, figs. 3, 4.

Western Louisiana; eastern Texas.

(Group of *Quadrula trigona*.)

Shell triangular, generally inflated, with high, full beaks which are incurved and turned forward over a well-developed lunule; anterior end obliquely truncate above, often with a curved, shallow depression in each valve running from the beaks to midway down the anterior end, and forming a sort of secondary lunule; posterior base usually incurved; the posterior ridge ending in a rather sharp point; beak sculpture, a few coarse, concentric ridges turned upward behind and often swollen on the posterior ridge, sometimes becoming finer and broken or irregular on the upper disk; hinge solid but not very wide; pseudocardinals triangular and radial, torn; there is a secondary lateral in the right valve; cavity of the beaks generally deep and compressed; muscle scars small, deep.

Animal with the marsupium occupying the whole of all four brachia; inner gills generally free from the abdominal sac, much wider than the outer in front; outer nearly or quite equaling them in width behind; anal opening distinctly crenulate or papillose.

† *QUADRULA RUBIGINOSA* Lea.

- * *Unio rubiginosus* LEA, Tr. Am. Phil. Soc., III, 1829, p. 427, pl. VIII, fig. 10; * Obs., I, 1834, p. 41, pl. VIII, fig. 10.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 548.—* HANLEY, Test. Moll., 1842, p. 185; * Biv. Shells, 1843, p. 185, pl. XXI, fig. 43.—* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CHENU, Ill. Conch., 1858, pl. XIII, figs. 4, 4a, 4b.—* REEVE, Conch. Icon., XVI, 1865, pl. XXVII, fig. 136.—* CALKINS, Pr. Ottawa Acad. Sci., 1874, p. 44.—* B. H. WRIGHT, Check List, 1888.—* WALTON, Moll. Monroe Co., 1892, p. 16, pl. VIII, fig. 1.

* *Margarita* (*Unio*) *rubiginosus* LEA, Syn., 1836, p. 20; 1838, p. 17.

* *Margaron* (*Unio*) *rubiginosus* LEA, Syn., 1852, p. 24; 1870, p. 37.

* *Unio rubiginosa* DESHAYES, An. sans Vert., 3d ed., II, 1839, p. 672.

* *Quadrula rubiginosa* BAKER, Moll. Chicago, Pt. 1, 1898, p. 77, pl. XIX, fig. 2; XX, fig. 1.

* *Unio flavus* CONRAD, New F. W. Shells, 1834, p. 69.—* SAY, Am. Conch., VI, 1834.—* FERUSSAC, Guer. Mag., 1835, p. 27.—* CONRAD, Monog., IX, 1837, p. 74, pl. XLI, fig. 2.—* KUSTER, Conch. Cab., 1852, p. 61, pl. XIV, fig. 2; p. 265, pl. LXXXIX, fig. 5.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.

* *Unio flavus* var. *rubiginosus* PÄTEL, Conch. Sam., III, 1890, p. 152.

* *Unio trigonus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIV, p. 322.

Entire Mississippi drainage; eastern Texas; St. Lawrence River system; Nelson River and its tributaries.

† *QUADRULA CERINA* Conrad.

- * *Unio cerinus* CONRAD, Monog., XI, 1838, p. 95, pl. LII; * Pr. Ac. N. Sci. Phila., VI, 1853, p. 246.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVII, fig. 468.—
* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 147.

Louisiana; eastern Texas; Mississippi; Alabama; north to Arkansas.

† *QUADRULA HEBETATA* Conrad.

- * *Unio hebetatus* CONRAD, * Jl. Ac. N. Sci. Phila., II, 1854, p. 296, pl. XXVI, fig. 5.—
* B. H. WRIGHT, Check List, 1888.
* *Margaron (Unio) habetatus* LEA, Syn., 1870, p. 38.

Missouri; Tuscaloosa River, Alabama.

† *QUADRULA RUBIDA* Lea.

- * *Unio rubidus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 40; * Jl. Ac. N. Sci. Phila., V, 1862, p. 95, pl. XIV, fig. 244; * Obs., VIII, 1862, p. 99, pl. XIV, fig. 244.—
* B. H. WRIGHT, Check List, 1888.
* *Margaron (Unio) rubidus* LEA, Syn., 1870, p. 35.
† * *Unio negatus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 59; * Jl. Ac. N. Sci. Phila., V, 1862, p. 76, pl. IX, fig. 225; Obs., VIII, 1862, p. 80, pl. IX, fig. 225.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXII, fig. 165.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 160.
* *Margaron (Unio) negatus* LEA, Syn., 1870, p. 35.

Streams flowing into the Gulf of Mexico from Alabama to Louisiana. The shell reported under the name *Unio negatus*, from Kansas, is probably a somewhat sulcate *rubiginosus*.

† *QUADRULA CHUNII* Lea.

- Unio chunii* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 392; * Jl. Ac. N. Sci. Phila., V, 1862, p. 196, pl. XXVII, p. 265; * Obs., IX, 1863, p. 18, pl. XXVII, fig. 265.—
* B. H. WRIGHT, Check List, 1888.—* PÆTEL, III, 1890, p. 148.
* *Margaron (Unio) chunii* LEA, Syn., 1870, p. 38.

Mississippi; west to central Texas north to Arkansas.

† *QUADRULA RIDDELLII* Lea.

- * *Unio riddellii* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 392; * Jl. Ac. N. Sci. Phila., V, 1862, p. 198, pl. XXVII, fig. 267; * Obs., IX, 1863, p. 20, pl. XXVII, fig. 267.—
* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIII, fig. 442.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 165.
* *Margaron (Unio) riddellii* LEA, Syn., 1870, p. 38.

Dallas, Texas.

† *QUADRULA FRIERSONI* B. H. Wright.

- * *Unio friersoni* B. H. WRIGHT, Nautilus, IX, 1896, p. 134, pl. III.

Louisiana; eastern Texas.

† *QUADRULA TRIGONA* Lea.

- * *Unio trigonus*, LEA, Tr. Am. Phil. Soc., IV, 1831, p. 110, pl. XVI, fig. 40; * Obs., I, 1834, p. 120, pl. XVI, fig. 40.—* HANLEY, Test. Moll., 1842, p. 185; * Biv. Shells, 1843, p. 185, pl. XXI, fig. 6.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—
* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 258.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* CHENU, Ill. Conch., 1858, pl. XVI, figs. 7, 7a, 7b.—

- * SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVI, fig. 459.—* CALKINS, Pr. Ottawa Acad. Sci., 1874, p. 45.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 170.
- * *Margarita (Unio) trigonus* LEA, Syn., 1836, p. 18; 1838, p. 17.
- * *Margaron (Unio) trigonus* LEA, Syn., 1852, p. 25; 1870, p. 38.
- * *Quadrula trigona* BAKER, Moll. Chicago, Pt. 1, 1898, p. 76, pl. xv, fig. 5.
- * *Unio undatus* CONRAD, New F. W. Shells, 1834, p. 72.—* SAY, Am. Conch., VI, 1834.
- * *Unio triangularis* KUSTER, Conch. Cab. Unio, 1852, p. 56, pl. XII, fig. 3.
- * *Unio pilaris* REEVE, Conch. Icon., XVI, 1865, pl. XXVII, fig. 138.

Entire Mississippi drainage; Coosa River, Alabama; Michigan and the Upper St. Lawrence drainage.

† *QUADRULA OBLIQUA* Lamarck.

- * *Unio obliqua* LAMARCK, An. sans Vert., VI, 1819, p. 72.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* DESHAYES, An. sans Vert., 2d ed., 1835, p. 534; 3d ed., II, 1839, p. 668.
- * *Margarita (Unio) obliqua* LEA, Syn., 1836, p. 20.
- * *Unio obliquus* HANLEY, Test. Moll., 1842, p. 186; * Biv. 1843, p. 186, pl. xx, fig. 24.—* CATLOW and REEVE, Conch. Nonl., 1845, p. 61.—* KUSTER, Conch. Cab. Unio, 1852, p. 63, pl. xv, fig. 1.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 253.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* CALKINS, Pr. Ottawa Acad. Sci., 1874, p. 44.—* PÆTEL, Conch. Sam., III, 1890, p. 161.—* CALL, Tr. Ac. Sci. St. Louis, VII, 1895, p. 31, pl. iv.
- * *Margarita (Unio) obliquus* LEA, Syn., 1838, p. 17.
- * *Margaron (Unio) obliquus* LEA, Syn., 1852, p. 25; 1870, p. 38.
- * *Unio undatus* BARNES, Am. Jl. Sci., VI, 1823, p. 121, pl. iv, fig. 4.—* HILDRETH, Am. Jl. Sci., XIV, 1828, p. 280.—* CONRAD, Pr. Ac. Nat. Sci. Phila., VI, 1853, p. 259.
- * *Mya undata* EATON, Zool. Text-Book, 1826, p. 219.
- * *Unio mytiloides* SHORT and EATON, Transylvania Jl., 1831, p. 74.
- * *Unio cordatus* CONRAD, Monog., V, 1836, p. 48, pl. xxv.—* KUSTER, Conch. Cab., 1852, p. 57, pl. XIII, fig. 1.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIII, 376.—* B. H. WRIGHT, Check List, 1888.

Ohio, Cumberland, and Tennessee river systems; west in Illinois to the Mississippi; Claiborne, Alabama.

† *QUADRULA COCCINEA* Conrad.

- * *Unio coccineus* CONRAD,² Monog., III, 1836, p. 29, pl. XIII, fig. 1.—LEA, Tr. Am. Phil. Soc., VI, 1838, p. 12, pl. v, fig. 12; * Obs., II, 1838, p. 12, pl. v, fig. 12.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 234.—* HANLEY, Test. Moll., 1842, p. 203; * Biv. Shells, 1843, p. 203, pl. XXII, fig. 54.—* CATLOW and REEVE,

¹ This was figured by Hildreth, but the figures of this species and a number of others were left out by the editor.

² In the Transactions of the American Philosophical Society, VI, 1838, p. 12, pl. v, fig. 12, Lea described this species, and stated that about eighteen months previous Dr. Hildreth had sent him a single specimen under the name *Unio coccineus*; but there is nothing to show that Hildreth had described it. In the Monography in 1836, III, p. 29, pl. XIII, fig. 1, Conrad describes this species under the same name, and also credits it to Hildreth, stating that it was in the collection of the Philadelphia Academy of Natural Sciences under that name. The species must be credited to Conrad, who first described it, though Lea read his description in 1834.

- Conch. Nom., 1845, p. 57.—*KUSTER, Conch. Cab., 1852, p. 49, pl. x, fig. 2.—
 *CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 247.—*H. and A. ADAMS, Gen.
 Rec. Moll., II, 1857, p. 492.—*CHENU, Ill. Conch., 1858, pl. xxv, figs. 4, 4a,
 4b.—*SOWERBY, Conch. Icon., XVI, 1868, pl. xciv, fig. 512.—*CALKINS, Pr.
 Ottawa Acad. Sci., I, 1874, p. 42.—*B. H. WRIGHT, Check List, 1888.—
 *PÆTEL, Conch. Sam., III, 1890, p. 148.
 **Margarita (Unio) coccineus* LEA, Syn., 1836, p. 34; 1838, p. 23.
 **Margaron (Unio) coccineus* LEA, Syn., 1852, p. 35; 1870, p. 56.
 **Quadrula coccinea* BAKER, Moll. Chicago, Pt. 1, 1898, p. 79, pl. xiv, fig. 1; xix,
 fig. 3.
 **Unio rubens* MENKE, Syn., Meth. Moll., 1828, p. 90.
 **Unio catillus* CONRAD, Monog., III, 1836, p. 30, pl. xiii, fig. 2.—*KUSTER, Conch.
 Cab., 1852, p. 64, pl. xv, fig. 2.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853,
 p. 246.—*PÆTEL, Conch. Sam., III, 1890, p. 147.
 **Unio catilus* B. H. WRIGHT, Check List, 1888.
 **Unio gouldianus* WARD,¹ Jay's Catalogue, 3d ed., 1839, p. 24.
 **Unio cuneus* CALL, Tr. Ac. Sci. St. Louis, VII, p. 14.

† *QUADRULA COCCIENEA* var. *PAUPERCULA* Simpson.²

Entire Upper Mississippi drainage; St. Lawrence basin in various localities. The peculiar dwarf variety which is found in this latter area seems almost entitled to specific rank.

† *QUADRULA SOLIDA* Lea.

- Unio solidus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 13, pl. v, fig. 13; *Obs., II, 1838, p. 13, pl. v, fig. 13.—*TROSCHÉL, Arch. für Naturg., V, 1839, Pt. 2, p. 234.—
 *HANLEY, Test. Moll., 1842, p. 186; *Biv. Shells, 1843, p. 186, pl. xxiii, fig. 15.—
 *CATLOW and REEVE, Conch. Nom., 1845, p. 64.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—*CHENU, Ill. Conch., 1858, pl. xxv, figs. 2, 2a, 2b.—*KUSTER, Conch. Cab., 1861, p. 258, pl. lxxxvii, fig. 4.—*REEVE, Conch. Icon., XVI, 1865, pl. xxvii, fig. 133.—*CALKINS, Pr. Ottawa Acad. Sci., 1874, p. 45.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 168.
 **Margarita (Unio) solidus* LEA, Syn., 1836, p. 20; 1838, p. 17.
 **Margaron (Unio) solidus* LEA, Syn., 1852, p. 25; 1870, p. 38.
 *? *Unio cardiacea* DESHAYES, Tr. Elem. de Conch., 1839, p. 19, pl. xxxi, figs. 1, 2.
 **Unio fulgidus* LEA,³ Pr. Am. Phil. Soc., IV, 1845, p. 164; *Tr. Am. Phil. Soc., X, 1848, p. 73, pl. iv, fig. 10; *Obs., IV, 1848, p. 47, pl. iv, fig. 10.—*CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 253.
 **Margaron (Unio) fulgidus* LEA, Syn., 1852, p. 25; 1870, p. 38.
 **Unio obovalis* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 253.

Ohio, Cumberland, and Tennessee river systems; south to Louisiana; west to Arkansas and Kansas; north to Minnesota; east, through Wisconsin, Illinois, and Indiana, to Ohio.

¹ Adopted by Dr. Jay and used in his catalogue, but never described. It is a white nacreous *coccinea*.

² There is a dwarf form, greatly inflated and having the rest periods very plainly marked, found in the St. Lawrence and its tributaries near the Niagara Falls, which may bear the above varietal name.

³ There is only the type, a young shell, in Lea's collection, and I have no hesitation in referring it to *solidus*.

†QUADRULA PLENA Lea.

- * *Unio plenus* LEA, Tr. Am. Phil. Soc., I, 1840, p. 286; *Tr. Am. Phil. Soc., VIII, 1843, p. 211, pl. XIV, fig. 26; *Obs., III, 1842, p. 49, pl. XIV, fig. 26.—*CONRAD, Proc. Ac. N. Sci. Phila., VI, 1853, p. 255.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*KUSTER, Conch. Cab., 1861, p. 264, pl. LXXXIX, fig. 3.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXI, fig. 305.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 163.

**Margaron (Unio) plenus* LEA, Syn., 1852, p. 25; 1870, p. 39.

Ohio, Cumberland, and Tennessee riversystems; southwest to Kansas and Arkansas.

QUADRULA FLEXUOSA Simpson.

Holston River, Tennessee.

Quadrula flexuosa SIMPSON, Proc. Ac. N. Sci. Phila., 1900, p. 83, pl. II, fig. 8.

†QUADRULA PYRAMIDATA Lea.

- **Unio pyramidatus* LEA, Tr. Am. Phil. Soc., IV, 1834, p. 109, pl. XVI, fig. 39; *Obs., I, 1834, p. 119, pl. XVI, fig. 39.—*HANLEY, Test. Moll., 1842, p. 186; *Biv. Shells, 1843, p. 186, pl. XX, fig. 45.—*DESHAYES, Traite Elem. de Conch., II, 1850?, p. 216, pl. XXXI, figs. 1, 2.—*CATLOW and REEVE, Conch. Nom., 1845, p. 63.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU, Ill. Conch., 1858, pl. XVI, figs. 5, 5a, 5b.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXIV, p. 323.—*B. H. WRIGHT, Check List, 1888.

**Margarita (Unio) pyramidatus* LEA, Syn., 1836, p. 21; 1838, p. 17.

**Margaron (Unio) pyramidatus* LEA, Syn., 1852, p. 25; 1870, p. 39.

- **Unio mytiloides* DESHAYES, Enc. Meth., II, 1830, p. 586, pl. CCXLIX, fig. 4.—*SAY, Am. Conch., Pt. 6, 1834.—*FERUSSAC, 1835, p. 28.—*MÖLLER, Syn. Nov. Gen., 1836, p. 209.—*CONRAD, Monog., IV, 1836, p. 41, pl. XX.—*SWAINSON, Treatise on Mal., 1840, p. 267, figs. 52–53.—*CATLOW and REEVE, Conch. Nom., 1845, p. 61.—*KUSTER, Conch. Cab., 1852, p. 59, pl. XIII, fig. 3; 1861, p. 265, pl. LXXXIX, fig. 4.—*R. H. WRIGHT, Check List, 1888.

**Margarita (Unio) mytiloides* LEA, Syn., 1836, p. 21; 1838, p. 17.

**Margaron (Unio) mytiloides* LEA, Syn., 1852, p. 25; 1870, p. 39.

**Unio mytiloides* Rafinesque var. *pyramidatus* PÆTEL, Conch. Sam., III, 1890, p. 160.¹

**Unio cardiacea* GUERIN, Icon. Regne Animal, 1829?, II, pl. XXVIII, fig. 7.

**Unio ruber* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 257.

**Unio obliqua* WOOD, Ind. Test. rev., 1856, p. 200, pl. I, fig. 8.

Ohio, Cumberland, and Tennessee river systems; southwest to Arkansas; west to Nebraska?; north in the Mississippi to Wisconsin.

(Group of *Quadrula subrotunda*.)

Shell solid, rounded, elliptical or ovate, with high beaks curved inward and forward over a distinct lunule; beak sculpture a few coarse

¹Most of the authors who use the name *mytiloides* believe it to be Rafinesque's species, though they figure and describe the *pyramidatus* of Lea. The specimens in the Lea collection which Dr. Lea calls *mytiloides* Rafinesque, are evidently a form of his *pyramidatus*. Rafinesque's figure represents a shell very long and almost straight on the posterior dorsal line, with astonishingly high beaks, and four distinct, equally separated rays on the disk, and he says there are some oblique, black, longitudinal bands on the shell. The Rafinesquian shell seems as near to the *Unio clava* of Lamarek as anything, but it does not agree with that.

ridges curved up behind; posterior ridge feebly developed; epidermis brownish or blackish, and often painted with a few faint rays on the earlier shell which are disposed to break into squarish spots; hinge and teeth strong; secondary lateral in right valve well developed; beak cavities deep, compressed; muscle scars deep. Animal with all four gills used as a marsupium throughout, filled with pink ova; gills large, inner only slightly the larger, free the greater part of their length; anal opening crenulate or papillose.

† *QUADRULA BURSA-PASTORIS* B. H. Wright.

* *Unio bursa-pastoris* B. H. WRIGHT, Nautilus., IX, 1896, p. 133, pl. III.

Clinch and Powell rivers; Virginia and Tennessee.

† *QUADRULA KIRTLANDIANA* Lea.

* *Unio kirtlandianus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 98, pl. XIV, fig. 41; * Obs., I, 1834, p. 210, pl. XIV, fig. 41.—* FERUSSAC, Guer. Mag., 1835, p. 29.—* HANLEY, Biv. Shells, 1843, p. 203, pl. XXIII, fig. 27.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—* KUSTER, Conch. Cab., 1856, p. 168, pl. XLIX, figs. 2, 3; 1861, p. 214, pl. LXXI, fig. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVII, fig. 402.—* B. H. WRIGHT, Cheek List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 156.

* *Margarita (Unio) kirtlandianus* LEA, Syn., 1836, p. 34; 1838, p. 23.

* *Margarita (Unio) kirtlandianus* LEA, Syn., 1852, p. 35; 1870, p. 56.

* *Unio kirklandianus* HANLEY, Test. Moll., 1842, p. 203.¹

† *QUADRULA KIRTLANDIANA* var. *MINOR* Simpson.²

Ohio, Cumberland, and Tennessee river systems; southwest to Arkansas; north to Wisconsin?; east through southern Michigan.

† *QUADRULA SUBROTUNDA* Léa.³

* *Unio subrotundus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 117, pl. XVIII, fig. 45; * Obs., I, 1834, p. 127, pl. XVIII, fig. 45.—* HANLEY, Test. Moll., 1842, p. 203; * Biv. Shells, 1843, p. 203, pl. XX, fig. 58.—* CATLOW and REEVE, Conch. Nom., 1845, p. 64.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* ?CHIENU, III.

¹ A typographical error no doubt. There is a *Unio kirklandianus* recently named by Wright.

² A form is found in the Tennessee drainage which comes nearest, it seems to me, to the *kirtlandianus* of Lea, but is much smaller and more delicate, and is not so strongly colored. Mr. Wetherby believes this to be a form of the nearly allied *subrotundus*, but it seems to me too much compressed for that. Dr. Sterki is of the opinion that the two species should be united. While there are specimens which can hardly be determined, it seems to me that in the main the two are distinct. The small form may be called variety *minor*.

³ The name *subrotunda* was used by Rafinesque in 1820 for some species of *Naiad*, probably a member of the *Circulus* group which I can not determine, and placed in the genus *Obliquaria* by him. Conrad afterwards, in 1834, placed what he supposed was this species in the genus *Unio*. Lea applied the name *subrotundus* to his species in 1831, and being the first to use it for that genus, I think it will have to stand, and Conrad's name become a synonym.

Conch., 1858, pl. xv, figs. 1, 1a, 1b.—* KUSTER, Conch. Cab., 1861, p. 190, pl. LX, fig. 3.—* B. H. WRIGHT, Check List, 1888.

* *Margarita (Unio) subrotundus* LEA, Syn., 1836, p. 34; 1838, p. 23.

* *Margaron (Unio) subrotundus* LEA, Syn., 1852, p. 35; 1870, p. 56.

* ? *Unio brevalis* CROUCH, Ill. Int. to Lamarek, 1827, p. 16, pl. IX, fig. 3.—* ? CATLOW and REEVE, Conch. Nom., 1845, p. 56.

* *Unio personatus* CONRAD, New F. W. Shells, 1834, p. 71.

* *Unio politus* SAY, Am. Conch., VI, 1834.²—* FERUSSAC, Guer. Mag., 1835, p. 28.—

* CONRAD, Monog., VIII, 1837, p. 67, pl. XXXVII, fig. 2.—* KUSTER, Conch. Cab. Unio, 1852, p. 62, pl. XIV, fig. 4.—* CONRAD, Pr. Ac. N. Sci. Phila.,

VI, 1853, p. 255.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXII, p. 168.—

* PÆTEL, Conch. Sam., III, 1890, p. 164.³

Ohio, Cumberland, and Tennessee river systems. Reported from Michigan and the Grand River, Ontario, but it is probable that the material from these localities is not *subrotundus*.

† QUADRULA PILARIS Lea.

* *Unio pilaris* LEA, Pr. Am. Phil. Soc., I, 1840, p. 285; Tr. Am. Phil. Soc., VIII, 1842, p. 209, pl. XIV, fig. 24; Obs., III, 1842, p. 47, pl. XIV, fig. 24.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 255.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* CHENU, Ill. Conch., 1858, pl. XXIX, figs. 3, 3a, 3b.—* KUSTER, Conch. Cab., 1861, p. 255, pl. LXXXVI, fig. 3.—* REEVE, Conch. Icon., XVI, 1865, pl. XXVII, fig. 138.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 163.

* *Margaron (Unio) pilaris* LEA, Syn., 1852, p. 35; 1870, p. 56.

† * *Unio lesueurianus* LEA, Pr. Am. Phil. Soc. I, 1840, p. 286; * Tr. Am. Phil. Soc., VIII, 1842, p. 195, pl. VIII, fig. 6; * Obs., III, 1842, p. 33, pl. VIII, fig. 6.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 251.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* CHENU, Ill. Conch., 1858, pl. XXX, figs. 4, 4a, 4b.—* KUSTER, Conch. Cab. Unio, 1861, p. 215, pl. LXXII, fig. 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 157.

* *Margaron (Unio) lesueurianus* LEA, Syn., 1852, p. 35; 1870, p. 56.

Tennessee and Cumberland river systems; Green River, Kentucky; reported from the Ohio River.

† QUADRULA CUNEUS Conrad.⁴

* *Unio cuneus* CONRAD, Monog., XII, 1840, p. 105; pl. LVIII, fig. 1.—* B. H. WRIGHT, Check List, 1888.

Arkansas; Louisiana; Sabine River, Texas.

¹The figure looks something like *subrotunda*, or it may possibly be *ebenus*. I can not make out Lamarek's *brevalis* (An. sans Vert., VI, 1819, p. 75), which he says came from the Isle de France. Sganzin was told by the people of the island that no *Unio* was found there (Mem. Hist. Nat. Strasb., III, 1846, p. 8).

²Merely listed by Say, but neither Lea nor I have been able to find that it was ever described by him.

³Kirtland states that an oblique sulcation below the superior posterior angle distinguishes the female of this species, as it does the *retusus*, from the male. I have not noticed it.

⁴Mr. Conrad states in the Monography that his shells came from the Little Red River, Arkansas. Our shells in the U. S. National Museum collection are from the same stream, but are not quite so solid, nor are they just the shape of the figure, yet I have no doubt that they are *cuneus*.

†QUADRULA EBENUS Lea.

- * *Unio ebenus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 84, pl. IX, fig. 14; *Obs., I, 1834, p. 94, pl. IX, fig. 14.—*CONRAD, New F. W. Shells, 1834, p. 69.—*HANLEY, Test. Moll., 1842, p. 202; *Biv. Shells, 1843, p. 202, pl. XX, fig. 47.—*CATLOW and REEVE, Conch. Nom., 1845, p. 58.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—*CHENU, Ill. Conch., 1858, pl. XV, figs. 7, 7a, 7b.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXVI, fig. 334.—*CALKINS, Pr. Ottawa Acad. Sci., 1874, p. 42.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 151.
- **Margarita (Unio) ebenus* LEA, Syn., 1836, p. 34; 1838, p. 23.
- **Margaron (Unio) ebenus* LEA, Syn., 1852, p. 35; 1870, p. 56.
- **Unio obovalis* SAY, Am. Conch., VI, 1834.
- **Unio obliquus* CONRAD, Monog., IX, 1837, p. 77, pl. XLIII, fig. 2.—*KUSTER, Conch. Cab. Unio, 1861, p. 215, pl. LXXI, fig. 4.
- **Unio mytiloides* SWAINSON, Treatise on Mal., 1840, p. 270, figs. 52, 53.

Mississippi drainage generally except its western portion; Alabama and Tombigbee rivers; northeast Texas?

†QUADRULA GLOBATA Lea.

- **Unio gl. batus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 191; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 5, pl. I, fig. 1; *Obs., XIII, 1874, p. 9, pl. I, fig. 1.—*B. H. WRIGHT, Check List, 1888.

Holston and Tennessee rivers, Tennessee; Etowah River, Georgia.

(Group of *Quadrula glandacea*.¹)

Shell rhomboid-oval, nearly straight below and slightly biangulate behind, with a well-defined double posterior ridge; beaks probably full; surface slightly and irregularly sulcate; epidermis wrinkled, tawny; hinge strong, the plate somewhat flattened; pseudocardinals triangular; laterals strong, secondary lateral of right valve well developed; beak cavities deep, compressed; muscle scars very deep, smooth; nacre dirty straw color.

†QUADRULA GLANDACEA Lea.

- **Unio glandaceus* LEA, Pr. Ac. N. Sci. Phila., V, 1861, p. 59; *Jl. Ac. N. Sci. Phila., V, 1862, p. 77, pl. IX, fig. 226; *Obs., VIII, p. 81, pl. IX, fig. 226.—*B. H. WRIGHT, Check List, 1888.
- **Margaron (Unio) glandaceus* LEA, Syn., 1870, p. 38.

Coosa and Cahawba rivers, Alabama.

¹I am somewhat at a loss to know where to place this group. The very limited amount of material I have seen is in a badly eroded condition. I can form no idea of the beak sculpture, and I have never seen the soft parts. The texture and coloring of all the shells I have seen are like those of *Pleurobema*, but the very deep beak cavities and the general form are like *Quadrula*.

Section PACHYNAIAS Crosse and Fischer, 1893.

(Type, *Unio spheniopsis* Morelet.)

Shell elongate-triangular, inflated, truncated above and rounded below in front, straight on the base, with a high, sharp posterior ridge; the post base slightly biangulate; beaks full, the sculpture consisting of apparently faint ridges which run parallel with the growth lines; whole surface distinctly concentrically ridged; epidermis without rays; pseudocardinals solid, stumpy, somewhat radiate, roughened; beak cavities rather deep; muscle scars deep, smooth; nacre bluish white, thicker in front.

Animal unknown.

†QUADRULA SPHENIOPSIS Morelet.

* *Unio spheniopsis* MORELET, Test. Nov., I, 1849, p. 29.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 583, pl. LXI, figs. 2, 2a, 2b.

Rio Usumacinta, Guatemala..

†QUADRULA RUGOSOSULCATA Lea.

* *Unio rugososulcatus* LEA, Pr. Ae. N. Sci. Phila., X, 1866, p. 33; * Jl. Ae. N. Sci. Phila., VI, 1868, p. 266, pl. XXXIV, fig. 81; * Obs., XII, 1869, p. 26, pl. XXXIV, fig. 81.—* B. H. WRIGHT, Cheek List, 1888.

* *Margaron (Unio) rugososulcatus* LEA, Syn., 1870, p. 35.

Central America.

Subgenus ROTUNDARIA (Rafinesque, 1821) Simpson.

(Type, *Obliquaria tuberculata* Rafinesque.)

Shell rounded or quadrate, slightly truncate above in front; posterior ridge low; beaks prominent, curved inward and forward over a strongly marked lunule; beak sculpture consisting of numerous (20 to 30) fine, irregular, broken, somewhat concentric corrugations which extend well on to the second year's growth and gradually blend with the regular sculpture; posterior three-fifths of the shell tuberculate; epidermis brown; beak cavities very deep, compressed; nacre violet.

Animal having the gills very large, inner very much wider in front, free the whole length from the abdominal sac; branchial opening immense, with many small papillae; anal opening very large, smooth; superanal opening not at all closed below.¹

(Group of *Quadrula tuberculata*.)

Shell rounded, truncated behind, with a wide depression down the post slope.

¹ I regret that I have never been able to see gravid specimens of either of the species placed here. They are generally placed with the *pustulosus* group, but the strongly developed sulcus on the posterior slope and the very remarkable beak sculpture, which is not at all like that of any of our species, are good distinguishing characters. The enormous anal and branchial openings, and the open superanal opening are characters of some value.

† *QUADRULA TUBERCULATA* Rafinesque.¹

* *Obliquaria (Rotundaria) tuberculata* RAFINESQUE, Ann. Gen. Sei. Brux., V, 1820, p. 103.—* CHENU, Bib. Conch., 1st ser., III, 1845, p. 20.

* *Rotundaria tuberculata* AGASSIZ, Arch. für Naturg., I, 1852, p. 48.

* *Unio tuberculatus* CONRAD, New F. W. Shells, 1834, p. 72.—* SAY, Am. Conch., VI, 1834.—* FERUSSAC, Guer. Mag., 1835, p. 28.—* CONRAD, Monog., V, 1836, p. 43, pl. XXII.—* KUSTER, Conch. Cab. Unio, 1852, p. 45, pl. IX, fig. 1.—* REEVE, Conch. Icon., XVI, 1864, pl. III, figs. 9, 12.

* *Unio verrucosus* BARNES, Am. Jl. Sei.; VI, 1823, p. 123, pl. v, fig. 6.—* SHORT and EATON, Transylvania Jl., 1831, p. 75.—* HANLEY, Test. Moll., 1842, p. 180;

* Biv. Shells, 1843, p. 180, pl. XXI, fig. 24.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* CALKINS, Pr. Ottawa Acad. Sei., 1874, p. 46.—* B. H. WRIGHT, Check List, 1888.—

* PÆTEL, Conch. Sam., III, 1890, p. 171.

* *Margarita (Unio) verrucosus* LEA, Syn., 1836, p. 16; 1838, p. 15.

* *Margaron (Unio) verrucosus* LEA, Syn., 1852, p. 22; 1870, p. 34.

* *Unio verrucosus purpureus* HILDRETH, Am. Jl. Sei., XIV, 1828, p. 281.

* *Mya verrucosa* EATON, Zool. Text Book, 1826, p. 216.

* *Quadrula verrucosa* BAKER, Moll. Chicago, Pt. 1, 1898, p. 85, pl. XXIII.

* *Unio tuberculosa* VALENCIENNES, Rec. Obs. Zool. Anat., II, 1833, p. 232.

Mississippi drainage generally; southern Michigan; San Saba County, central Texas.

† *QUADRULA GRANIFERA* Lea.²

* *Unio graniferus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 69, pl. XIX, fig. 60; * Obs., II, 1838, p. 69, pl. XIX, fig. 60.—* TROSCHEL, Arch. für Naturg., V, 1839, Pt. 2, p. 237.—* HANLEY, Test. Moll., 1842, p. 180; * Biv. Shells, 1843, p. 180, pl. XXIII, fig. 11.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* CONRAD, Pr. Ae. N. Sei. Phila., VI, 1853, p. 250.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* CHENU, Ill. Conch., 1858, pl. XXV, figs. 3, 3a, 3b.—* KUSTER, Conch. Cab. Unio, 1861, p. 212, pl. LXX, fig. 4.—* REEVE, Conch. Icon., XVI, 1864, pl. IX, fig. 34.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 154.

* *Margarita (Unio) graniferus* LEA, Syn., 1838, p. 15.

* *Margaron (Unio) graniferus* LEA, Syn., 1852, p. 22; 1870, p. 34.

† *QUADRULA GRANIFERA* var. *PUSILLA* Simpson.³

Ohio, Cumberland, and Tennessee river systems; northwest to Iowa.

¹ Rafinesque's description is plainly that of one of the pustulous *Uniones* of the Ohio, and the statement that it is truncated posteriorly, wants nodules anteriorly, is three or more inches long, and has a violet naere makes it certain that it is the *verrucosus* of Barnes.

² This form has been considered by many merely at best a variety of the *tuberculata*, and I have so regarded it. It has always a smaller and generally a more inflated shell, higher beaks, and usually a smoother, more olive-colored epidermis than the latter. Its naere is ordinarily more copper colored than that of *tuberculata*. I believe with Mr. Wetherby that it is a valid species.

³ A greatly dwarfed inflated form from the Green River at Mammoth Cave, Kentucky, may bear the above name.

(Group of *Quadrula ostreata*.)

Shell rhomboid, incurved on the base, with a well-developed posterior ridge, somewhat biangulate behind; beak sculpture consisting of a number of fine, concentric ridges showing a tendency to be doubly looped, which gradually change, first to corrugations and then to pustules; pseudocardinals ragged, secondary lateral of right valve but slightly developed.

Animal unknown.

† *QUADRULA OSTREATA* Morelet.

* *Unio ostreatus* MORELET, Test. Noviss., No. 1, 1849, p. 29.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 573, pl. LXIII, fig. 3; LXX, fig. 4.

Rio Usumacinta, Guatemala.

† *QUADRULA USUMASINTÆ* Crosse and Fischer.

* *Unio usumasintæ* CROSSE and FISCHER, J. de Conch., XL, 1892, p. 294.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 574, pl. LXIII, fig. 4; LXVII, fig. 5.

Rio Usumacinta, Guatemala.

† *QUADRULA GUATEMALENSIS* Simpson.

* *Quadrula guatemalensis*, SIMPSON, Pr. Acad. N. Sci. Phila., 1900, p. 83, pl. II, fig. 4.

Rio Usumacinta, Guatemala.

(Group of *Quadrula percompressa*.)

Shell large, triangular to rhomboid, greatly compressed throughout, with a low posterior ridge and a slight secondary ridge above it, making the shell slightly biangulate posteriorly; beaks very prominent, flattened; beak sculpture not seen; surface of the disks covered with rather fine, somewhat lachrymose nodules, corrugate wrinkled on the posterior slope;¹ cavity of the beaks enormously deep and compressed; area outside the palléal line wide in front; shell thickened anteriorly.

† *QUADRULA PERCOMPRESSA* von Martens.

* *Unio percompressus* VON MARTENS, Sitzungs Ber. Nat. Tr. 1887, p. 107.—

* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 576; Biologia Cent-Am. Mollusca, 1900, p. 496, pl. XXXII, figs. 1-3.

Rio de las Salinas, Guatemala.

Subgenus *LAMPROTULA* Simpson, 1900.

(Type, *Chama plumbea* Chemnitz.)

Shell rounded to triangular, inflated, with high beaks; beak sculpture consisting of a few coarse, subparallel ridges which are slightly doubly

¹ Only two left valves of this remarkable species have been seen by the writer, both of which curve to the left in front and to the right behind. It is probable that others may turn to the right, and von Martens' shells are flat.

looped; surface of the shell generally covered with coarse nodules or knobs; posterior slope radially ridged; epidermis dark; two strong laterals in each valve, all vertically striated; cavity of the beaks deep and compressed; nacre mostly whitish, usually radially granularly striated outside the pallial line, having one or more peculiar calluses behind and below the laterals which are iridescent and granularly striate.

Animal unknown.¹

(Group of *Quadrula plumbea*.)

Shell rounded, slightly angulated behind the ligament, solid, moderately inflated, covered with coarse pustules, and having broken, up-curved corrugations on the posterior slope; epidermis light brownish; hinge rather strong, somewhat flattened; laterals vertically striated.²

QUADRULA PLUMBEA Chemnitz.

**Chama plumbea* CHEMNITZ, Conch. Cab., XI, 1795, p. 237, pl. CCIII, figs. 1991, 1992.—*DILLWYN, Cat. I, 1817, p. 215.—*WOOD, Ind. Test. Rev., 1856, p. 52, pl. IX, fig. 6.

Unio plumbeus FERUSSAC, Guer. Mag., 1835, p. 28.—*KUSTER, Conch. Cab. Unio., 1862, p. 289, pl. XCVII, figs. 1, 2.

South East Asia, probably. The locality, "South Sea," given by Chemnitz is, no doubt, erroneous.

†QUADRULA COREANA von Martens.

**Unio coreanus* VON MARTENS, S. B. Nat. Fr., 1886, p. 78.—*PÆTEL, Conch. Sam., II, 1890, p. 149.

Söul, Korea.

(Group of *Q. nodulosa*.)

Shell very solid, oval or rounded, inflated, inequilateral, inequivalve, the valves more or less twisted on their axis or having the appearance as if one of them had been pushed forward on the other, so that a dorsal view shows the outline to be rhomboid; each valve with a posterior ridge, that on the valve pushed forward less developed; beaks high, full, apparently sculptured with a few coarse irregular ridges; posterior slope usually having strong, radiating, upcurved ridges; hinge plate generally wide and flattened; pseudocardinals heavy, somewhat radiate, often with the sockets evenly, radially grooved; laterals strong, heavily vertically ridged; secondary lateral of right valve well developed,

¹ A group of peculiar and striking species which presents a number of quite diverse forms. On account of this diversity, the fact that the beak sculpture is rarely preserved, and that nothing is known of the anatomy, the group is a difficult one to arrange.

² A single, rather young, dead, and somewhat decayed specimen of what I believe is *Q. coreana*, from Söul, Korea, belonging to the collection of F. Stearns has been examined by the writer. The *Chama plumbea* of Chemnitz seems to be near it, but is solidier and shorter.

anterior cicatrices small, deep, posterior shallow; cavity of the beaks enormously deep, compressed; dorsal cicatrices on the under side of the shelf.

Animal unknown.

† *QUADRULA TORTUOSA* Lea.

* *Unio tortuosus* LEA, Pr. Ac. N. Sci. Phila., IX, 1865, p. 76; * JI. Ac. N. Sci. Phila., VI, 1868, p. 286, pl. XXXIX, fig. 98; * Obs., XII, 1869, p. 46, pl. XXXIX, fig. 98.

* *Margaron (Unio) tortuosus* LEA, Syn., 1870, p. 30.

† * *Unio (Lampsilis) subortus* BAIRD and ADAMS, Proc. Zool. Soc. Lond., 1867, p. 491, pl. XXVI, figs. 1, 1a.

* *Unio subortus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVII, fig. 465.—

* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LXI, fig. 119.—* PÆTEL, Conch. Sam., III, 1890, p. 169.

* *Unio retortus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LVII, fig. 109.—* PÆTEL, Conch. Sam., III, 1890, p. 165.

China.

† *QUADRULA TIENTSINENSIS* Crosse and Debeaux.

* *Unio tientsinensis* CROSSE and DEBEAUX, JI. de Conch., III, 1863, p. 257, pl. x, fig. 1.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCI, fig. 514, 514a, 514b.—* PÆTEL, Conch. Sam., III, 1890, p. 169.

* *Margaron (Unio) tientsinensis* LEA, Syn., 1870, p. 30.

China.

QUADRULA ZONATA Heude.

* *Unio zonatus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LXI.—* PÆTEL, Conch. Sam., III, 1890, p. 172.

* *Unio tientsinensis* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LX, fig. 118.

China.

† *QUADRULA FIBROSA* Heude.

* *Unio fibrosus* HEUDE, Conch. Fluv. Nank., II, 1877, pl. XL, fig. 22.¹—* PÆTEL, Conch. Sam., III, 1890, p. 152.

*† *Unio spurius* HEUDE, Conch. Fluv. Nank., II, 1877, pl. XI, fig. 22.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

China.

† *QUADRULA NODULOSA* Wood.

* *Mya nodulosa* WOOD (part), Gen. Conch., I, 1815, p. 106, pl. XXII, figs. 1, 2.²—

* DILLWYN, Catalogue I, 1817, p. 52.—* WOOD, Index Test, 1825, p. 12, pl. II, fig. 29b; * Index Test. Rev., 1856, p. 16, pl. II fig. 29.

* *Unio nodulosus* HANLEY, Test. Moll., 1842, p. 180.—* Biv. Shells, 1843, p. 180.—

* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* H. and A. ADAMS, Gen.

¹ Doubtfully distinct from *zonata*.

² Wood gives four figures; the first and second being a solid, nodulous, Chinese species; the third and fourth seem to be *Unio pictorum*. Lamarek used the name *nodulosa* for a *Unio* (An. sans Vert., VI, 1819, p. 78), and refers to the Encyclopedie Méthodique, 1797, pl. 248, fig. 19, and this figure is no doubt that of *Unio pictorum*. Wood seems to have confounded the two.

Rec. Moll., II, 1857, p. 496.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIII, fig. 439.—* PÆTEL, Conch. Sam., III, 1890, p. 161.

* *Margarita (Unio) nodulosus* LEA, Syn., 1836, p. 16; 1838, p. 15.

* *Margaron (Prisodon) nodulosus* LEA, Syn., 1852, p. 27; 1870, p. 27.

* *Castalia nodulosa* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 509.

† *Unio grandidens* LEA,¹ Pr. Ac. N. Sci. Phila., V, 1862, p. 168; * Jl. Ac. N. Sci. Phila., V, 1862, p. 205, pl. XXX, fig. 274; * Obs., IX, 1863, p. 27, pl. XXX, fig. 274.

* *Margaron (Unio) grandidens* LEA, Syn., 1870, p. 34.

China.

QUADRULA KOUANGENSIS Simpson.²

* *Unio moreletianus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LX, fig. 117.—* PÆTEL, Conch. Sam., III, 1890, p. 160.

Kouang-te-tcheou, China.

†QUADRULA POLYSTICTO-SCRIPTUS Heude.

* *Unio polysticto-scriptus* HEUDE, Conch. Fluv. Nank., II, 1877, pl. XIII, fig. 26.—

* PÆTEL, Conch. Sam., III, 1890, p. 164.

* *Unio scripto-polystictus* HEUDE,³ Conch. Fluv. Nank., II, 1877, pl. XIII, fig. 27.

* *Unio scripto-polystictus* PÆTEL, Conch. Sam., III, 1890, p. 167.

China.

†QUADRULA POLYSTICTUS Heude.

* *Unio polystictus* HEUDE, Conch. Fluv. Nank., II, 1877, pl. XII, fig. 25.—* PÆTEL, Conch. Sam., III, 1890, p. 164.

China.

†QUADRULA SIMILARIS Simpson.

* *Unio affinis* HEUDE, Conch. Fluv. Nank., I, 1875, pl. III, figs. 7, 7a.⁴—* PÆTEL, Conch. Sam., III, 1890, p. 144.

China.

†QUADRULA MICROSTICTIS Heude.

* *Unio microstictis*, HEUDE, Conch. Fluv. Nank., II, 1877, pl. XII, fig. 24; * Jahrb. Deutsch. Mal. Ges., VIII, 1881, p. 122.—* PÆTEL, Conch. Sam., III, 1890, p. 159.

China.

(Group of *Quadrula rochechouarti*.)

Shell rhomboid, with a strong posterior ridge, rounded before, incurved below, and somewhat biangulate on the posterior base, with very strong folds on posterior slope which begin on the posterior ridge as

¹ A single valve (the type) is in the Lea collection, and an opposing matched valve is in that of the U. S. National Museum. Dr. Lea states that they were obtained near Hot Springs, Arkansas, by Major Byrd Powell. I am certain this is a mistake, and that it is a Chinese species. It agrees exactly with Wood's figure of *nodulosa*, is slightly pushed over or twisted on its axis, and has the vertically ridged laterals characteristic of the heavy Chinese forms very strongly developed.

² Heude's name *moreletianus* has been twice preoccupied.

³ Believed by Heude to be a hybrid with *polysticto-scriptus*. I can not see that the two differ sufficiently to be separated.

⁴ The name *affinis* was used long before by Dr. Lea for a *Unio*.

knobs; surface strongly pustulous; beaks rather well forward, sculpture not seen; epidermis concentrically wrinkled, black and rough; pseudo-cardinals strong, somewhat radial, rough, striate; laterals more or less vertically striate, secondary lateral in right valve well developed; beak cavities very deep, compressed; posterior calluses faint.

Animal unknown.

†QUADRULA ROCHECHOUARTI Heude.

* *Unio rochechouarti* HEUDE, Conch. Fluv. Nank., I, 1875, pl. v, fig. 13; *Jahrb. Deutsch. Mal. Ges., VIII, 1881, p. 122.—*PÆTEL, Conch. Sam., III, 1890, p. 165.

China.

(Group of *Quadrula triclava*.)

Shell solid, elongate triangular, obtusely pointed at the posterior base, with a decided posterior ridge, along which runs a row of ponderous knobs; beaks high and far forward; surface strongly nodulous; posterior slope slightly corrugated; epidermis dark chestnut; pseudo-cardinals heavy, subradial, radially striate; laterals vertically striate, secondary lateral of right valve well developed; cavity of the beaks enormously deep and compressed; a granular callus behind the laterals; nacre silvery, radially granularly striate outside the pallial line.

†QUADRULA TRICLAVA Heude.

* *Unio triclavus* HEUDE, Conch. Fluv. Nank., II, 1877, pl. x, figs. 21, 21a.—*PÆTEL, Conch. Sam., III, 1890, p. 170.

China.

(Group of *Quadrula bazini*.)

Shell elongate-trigonal, with a fairly well developed posterior ridge, pointed behind; beaks not high, almost at extreme front of the shell; surface more or less covered with knobs and tubercles, which show a tendency to an arrangement in curved lines; epidermis chestnut to blackish; hinge strong; pseudo-cardinals radial, radiately striate; laterals granular, showing traces of vertical striation; beak cavities very deep, compressed; nacre silvery.

Animal unknown.

†QUADRULA BAZINI Heude.

* *Unio bazini* HEUDE, Conch. Fluv. Nank., II, 1877, pl. ix, fig. 20.—*PÆTEL, Conch. Sam., III, 1890, p. 145.

China.

(Group of *Quadrula lei*.)

Shell obovate, obtusely pointed behind; posterior ridge low; whole surface generally tuberculate or knobbed, with strong, radial, curved

ridges on the posterior slope; beaks rather low, the sculpture coarse, irregular, broken bars, somewhat doubly looped and swollen on the posterior ridge, with strong, radiating ridges behind them; pseudocardinals solid, often compressed in the direction of the axis of the shell; laterals granular; beak cavities moderately deep; one or more granular calluses behind the laterals; nacre white or lurid, slightly radially granular outside the pallial line.

Animal unknown.

† *QUADRULA LEAI* Gray.

* *Unio leai* GRAY, Griff. Cuvier, XII, 1834, p. 600 (index), pl. XXI, fig. 1.¹—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* BENSON, Jl. As. Soc. Calcutta, XXIV, 1855, p. 136 (p. 18, reprint).—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* KUSTER, Conch. Cab. Unio, 1861, p. 232, pl. LXXVIII, fig. 3.—* ? HEUDE, Conch. Fluv. Nank., I, 1875, pl. IV, fig. 10, pl. VI.—* GREDLER, Jahrb. Deutsch. Mal. Ges., VIII, 1881, p. 122.—* VON MARTENS, Sitzber. Nat. Fr., 1885, p. 77.—* PÆTEL, Conch. Sam., III, 1890, p. 157.

* *Margarita (Unio) leaii* LEA, Syn., 1836, p. 17; 1838, p. 16.

* *Margarou (Unio) leaii* LEA, Syn., 1852, p. 23; 1870, p. 34.

* *Unio leaii* HANLEY, Biv. Shells, 1843, p. 182, pl. XXIII, fig. 55.

* *Unio nodulosus* REEVE, Conch. Icon., XVI, 1864, pl. IX, fig. 32.

† *QUADRULA LEAI* var. *LELECI* Heude.

* *Unio leleci* HEUDE, Conch. Fluv. Nank., I, 1875, pl. IV, fig. 12; v, fig. 14.—GREDLER, Jahrb. Deutsch. Mal. Ges., VIII, 1881, p. 122.

* *Unio richthofeni* VON MARTENS, S. B. Nat. Fr. 1875, p. 3; Mal. Blatt, XII, 1875, p. 187; Nov. Conch., IV, 1876, p. 156, pl. CXXXVI, figs. 1-3.—* PÆTEL, Conch. Sam., III, 1890, p. 165.

* *Unio leai* var. *leleci* PÆTEL, Conch. Sam., III, 1890, p. 157.

China; Tonkin; Mekong River.

QUADRULA OVATA Simpson.

Unio vestitus var. *a* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LVIII, fig. 112; LIX, fig. 116.³

QUADRULA CAVEATA Heude.

* *Unio caveatus* HEUDE, Conch. Fluv. Nank., III, 1877, pl. XXIV, fig. 53.—* PÆTEL, Conch. Sam., III, 1890, p. 147.

* *Unio contritus* HEUDE, Conch. Fluv. Nank., VII, 1881, pl. LVI, fig. 103.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Unio quadrangulosus* HEUDE, Conch. Fluv. Nank., VII, 1881, pl. LVI, fig. 104.—* PÆTEL, Conch. Fluv. Nank., III, 1890, p. 164.

China.

¹ A very brief description is given in the index.

² Von Martens, in a note at the close of Vol. IV, places his species in the synonymy of *U. leleci* Heude. I am satisfied that *leleci* is but a variety of *U. leai* Gray.

³ Heude's name is preoccupied by Lea for a *Unio*, I therefore change it as above.

† *QUADRULA CORNUUM LUNÆ* Heude.¹

* *Unio montanus* HEUDE, Conch. Fluv. Nank., I, 1875, pl. iv, fig. 11.—* PÆTEL, Conch. Sam., III, 1890, p. 159.

* *Unio cornuum-lunæ* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LVII, fig. 105.—* PÆTEL, Conch. Sam., III, 1890, p. 149.

* *Unio trisulcatus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LVII, fig. 108.

* *Unio paschalis* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LVIII, fig. 110.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Unio verruculosus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LVIII, fig. 111.—* PÆTEL, Conch. Sam., III, 1890, p. 171.

* *Unio abortivus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LXIII, fig. 124.

* *Unio monticola* HEUDE, Jl. de Conch., XXXII, 1884, p. 19.

QUADRULA CORNUUM LUNÆ var. *CINNAMOMEUS* Gredler.

* *Unio lei* var. *cinnamomens* GREGLER, Jahrb. Deuts. Mal. Ges., VIII, 1881, p. 122, pl. VI, fig. 16.²—* PÆTEL, Conch. Sam., III, 1890, p. 157.

China.

QUADRULA SCRIPTA Heude.

* *Unio scriptus* HEUDE, Conch. Fluv. Nank., I, 1875, pl. III, figs. 8, 8a.—* PÆTEL, Conch. Sam., III, 1890, p. 167.

China.

QUADRULA DIVERGENS Benson.

Unio divergens BENSON, Jl. As. Soc. Beng., XXIX, 1885, p. 137.³

Chusan Island, China.

Subgenus *DISCOMYA* Simpson, 1900.

Shell subsolid, round obovate or subrhomboid, lenticular, rather compressed, widely, faintly biangulate behind, with scarcely any vestige of a posterior ridge; beaks very low, sculpture not seen; front half of the shell densely covered with fine pustules arranged in curved rows in two directions as if engine chased, over which the epidermis is wrinkled. The hinder half is covered with fine, radiating and undulating corrugations, curved upward posteriorly, which are slightly nodulous and show through on the inside of the shell; one slightly compressed pseudo-cardinal in the right valve and two in the left; one delicate, curved

¹ First called *montanus* by Heude, which was several times preoccupied in *Unio*.

² Gredler makes this a variety of *lei*, but I believe it is a rather finely sculptured form of *monticola*. The species varies much in the size of pustules and the degree of its smoothness, and Heude has exalted all these variations to the rank of species. I think it probable that when a large amount of material can be examined it will be found necessary to reduce some of the Chinese forms I have considered as species to the rank of varieties, or even to the synonymy.

³ Benson says this is not very distantly related to *Unio undulatus* Barnes. It has never been figured, so far as I know, but from all I can learn of it it belongs in this group and may be *Q. lei*.

lateral in the right valve and two in the left; beak cavities deep, compressed; nacre lurid; pallial line showing a slight posterior sinus.¹

(Type, *Unio radulosus* Drouet and Chaper.)

†QUADRULA RADULOSA Drouet and Chaper.

**Unio radulosus* DROUET and CHAPER, Mem. Soc. Zool. Fr., V, 1892, p. 150, pl. v, figs. 10-12.—*Jl. de Conch., XLI, 1893, p. 41.

†QUADRULA RADULOSA var. RHOMBOIDEA Simpson.²

Borneo.

The relationship of the following is uncertain.

UNIO NEWENHUISI Schepman.

**Unio newenhuisi* SCHEPMAN, Notes Leyd. Mus., XX, 1898, p. 92, pl. I, figs. 1, 2.—

*DROUET, Jl. de Conch., 1899, p. 406.

Borneo.

May be related to *Unio radulosus* Drouet and Chaper.

Genus SHISTODESMUS Simpson, 1900.

(Type, *Unio lampreyanus* Baird and Adams.)

Shell rather solid, triangular, inflated, truncate above in front, somewhat swollen just behind the center of the base, pointed behind; beaks high; beak sculpture not seen; surface thrown up into strong, very wide, concentric ridges, one or two with each season's growth, and pinched up in the central part to form a radiate row of compressed knobs or spines; epidermis shining, greenish-yellow or brownish, having marvelously delicate, concentric, microscopic liræ, often painted with a few elegant, broken, green rays; there is one large, high, triangular pseudocardinal in the left valve under the beak, a deep, triangular pit in front of it, with an erect, radial lamellar tooth in its center, and a high, lamellar, curved tooth in front of the pit, parallel with the edge of the shell connected with the central tooth. A heavy triangular bifid or trifid pseudocardinal in the right valve stands close to the anterior edge of the shell and is separated from it by a deep, compressed pit; behind this tooth, deep down at its base, is a narrow shelf, and back of this the entire hinge plate is cut out in a deep sinus to the beak; two laterals in each valve; cavity of the beak in left valve deep, somewhat compressed; in the right entirely opened by the sinus; muscle scars small, deep, smooth; pallial line with a sinus at its posterior end; a

¹I know nothing of the anatomy of this curious *Naiad*, which seems to show some relation to such species as *Quadrula microstictus* and related forms. I place it in *Quadrula* with doubt.

²There is a specimen in the U. S. National Museum marked *Unio radulosus* Drouet and Chaper, from Borneo, which is quite rhomboid and but slightly sculptured on the body, though it is plicate behind, which may bear the above name. It is probably a valid species.

beautiful, granularly striate callus is developed behind the laterals; nacre finely, often bifurcately radiate striate outside the parallel line.

Animal unknown.

† SHISTODESMUS LAMPREYANUS Baird and Adams.

- * *Unio (Dysnomia) lampreyanus*, BAIRD and ADAMS, Proc. Zool. Soc. Lond., 1867, p. 491, pl. XXVI, figs. 2, 2a.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVII, fig. 469.—* HEUDE, Conch. Fluv. Nank., VII, 1883, pl. LIX, figs. 114, 114a, 114b.—* P. ET EL, Conch. Sam., III, 1890, p. 156.

Margarou (Unio) lampreyanus LEA, Syn., 1870, p. 30.

SHISTODESMUS SPINOSUS Simpson.

Unio vestitus var. β , HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LIX, fig. 115.¹

Genus GIBBOSULA Simpson, 1900.

(Type, *Mya crassa* Wood.)

Shell very solid, inequilateral, triangular-elliptical, arcuate, pointed behind, with a low, well-marked posterior ridge; beaks rather high; beak sculpture not seen; posterior two-thirds of the shell covered with somewhat radiating corrugations and nodules; posterior slope having strong subradial folds; epidermis dark; hinge plate very wide; pseudo-cardinals not large, stumpy, rough; laterals very heavy, club shaped; beak cavities enormously deep and compressed; anterior cicatrices rather shallow, rough in the bottom; posterior cicatrices deep and distinct; a strong rib runs from the front part of the beak cavity toward the posterior base; nacre whitish.

Animal unknown.

† GIBBOSULA CRASSA Wood.

- * *Mya crassa* WOOD, Gen. Conch., 1815, p. 106, pls. xx, xxi;² * Ind. Test., 1825, p. 12, pl. II, fig. 28b; * Ind. Test. Rev., 1856, p. 16, pl. II, fig. 28.

* *Mya ponderosa* DILLWYN, Dill. Cat., I, 1817, p. 51.

* *Margarita Unio ponderosus* LEA, Syn., 1836, p. 14; 1838, p. 14.

* *Unio ponderosus* HANLEY, Test. Moll., 1842, p. 177; * Biv. Shells, 1843, p. 177.—

* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* H. and A. Adams, Gen. Rec. Moll., II, 1857, p. 497.

* *Margarou (Unio) ponderosus* LEA, Syn., 1852, p. 21; 1870, p. 32.

China.

Genus CUNEOPSIS Simpson, 1900.

(Type, *Unio celliformis* Heude.)

Shell solid, elongated, wide and truncate or rounded in front, tapering to a point behind; often twisted on its axis and curved to the right

¹ Heude made this a variety of his *Unio vestitus* (which I have placed in *Quadrula*. I am satisfied that this is a *Shistodesmus* and distinct from *lampreyanus*, as it seems to have spines instead of knobs. His name *vestitus* is preoccupied in *Unio* by Lea.)

² The name *crassus* had been previously applied to a *Unio* by Retzius, but as Wood's species was never placed in *Unio* under the specific name *crassus*, his name may be used.

or left; beaks rather high, anterior; beak sculpture apparently a few coarse, subparallel nodular ridges curved up behind; posterior ridge low, close to the hinge line, and in front of it the shell is full; surface slightly, irregularly, concentrically striate; epidermis dark, having a peculiar, dull, silky luster; hinge rather narrow; pseudocardinals two in the left valve immediately under the beak, the anterior compressed, high and nearly parallel with the outer edge of the shell, the posterior heavy, joined to the anterior above, with a deep, triangular cavity between; right valve with one large, triangular, anterior pseudocardinal with a deep, triangular pit back of it, and often a small, low, compressed tooth behind the pit; laterals granularly striate, beak cavities rather deep; anterior muscle scars deep, rough at the bottom; posterior scars large, long and oblique; nacre silvery, sometimes radially striate at the edge; pallial line indented into a sinus at its posterior end, above which is a granular, striate callus.

Animal unknown.

†CUNEOPSIS CAPITATUS Heude.

* *Unio capitatus* HEUDE, Jl. de Conch., XXII, 1874, p. 114; * Conch. Fluv. Nank., I, 1875, pl. II, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 147.

China.

†CUNEOPSIS HEUDEI Heude.

* *Unio heudei* HEUDE (Bazin manuscript), Jl. de Conch., XXII, 1874, p. 114.—* PÆTEL, Conch. Sam., III, 1890, p. 155.

* *Unio corderii* HEUDE, Conch. Fluv. Nank., I, 1875, pl. I, fig. 3.¹—* PÆTEL, Conch., Sam., III, 1890, p. 149.

China.

†CUNEOPSIS CELTIFORMIS Heude.

* *Unio celtiformis* HEUDE, Jl. de Conch., XXII, 1874, p. 113; Conch. Fluv. Nank., I, 1875, pl. I, fig. 4.—* PÆTEL, Conch. Sam., III, 1890, p. 147.

China.

†CUNEOPSIS PISCICULUS Heude.

* *Unio pisciculus* HEUDE, Jl. de Conch., XXII, 1874, p. 115; * Conch. Fluv. Nank., I, 1875, pl. II, figs. 6, 6a.—* PÆTEL, Conch. Sam., III, 1890, p. 163.

* *Unio retortus* VON MARTENS, Sitzber. Ges. Nat. Fr., 1875, p. 4; * Mal. Blatt., XXII, 1875, p. 188; Nov. Conch., IV, 1876, p. 158, pl. CXXXVI, figs. 3, 4.²—* PÆTEL, Conch. Sam., III, 1890, p. 165.

China.

†CUNEOPSIS RUFESCENS Heude.

* *Unio rufescens* HEUDE, Jl. de Conch., XXII, 1874, p. 113; Conch. Fluv. Nank., I, 1875, pl. I, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 166.

China.

¹ Changed to *corderi* by Heude in above reference.

² Von Martens places this in the synonymy of *U. pisciculus* in Vol. IV of the Novitates Conchologia.

Subfamily HYRIANÆ Swainson.

(ENDOBRANCHIÆ.)

Male and female shells alike, with beak sculpture radial or zigzag-radial; marsupium occupying the inner gills only.

(ROSANORHAMPHUS.)

Beak sculpture zigzag-radial.

Genus NODULARIA Conrad, 1853.

(Type, *Unio douglasii* Gray.¹)

Nodularia CONRAD, Pr. Ac. N. Sci. Phila., 1853, p. 268.

Shell elliptical to elongated, pointed behind about midway up from the base, the post-basal part produced; beak sculpture variable, irregularly zigzag-radial, often breaking into nodules, and extending in many cases over a part or all of the disk; right valve with two usually compressed pseudocardinals, one above the other, the lower the more elevated, separated by a parallel-sided socket, and having one lateral; left valve with two compressed pseudocardinals, both in front of the beaks, and two laterals; cavity of the beaks moderate, not compressed, anterior muscle scars deep, posterior shallow, nacre white.

Animal (of *N. japonensis* and *N. equitoria*) having the inner gills alone filled throughout their entire length with ova, forming a pad-like marsupium, united to the abdominal sac or free from it.

Section LANCEOLARIA Conrad, 1853.

(Type, *Unio grayanus* Lea.)

Shell ensiform, solid, with a distinct, pinched-up posterior ridge, rounded in front, the posterior end sharp and generally turned a little to the right or left; beaks low, their sculpture nodulous, zigzag, often extending more or less over the surface; pseudocardinals rather stumpy, ragged, striate above, smooth below; anterior muscle scars distinct, the upper round, very deep, appearing as if bored out.

Animal unknown.

† NODULARIA GRAYANA Lea.

* *Unio grayanus* Lea, Tr. Am. Phil. Soc., V, 1834, p. 66, pl. IX, fig. 26; Obs., I, 1834, p. 178, pl. IX, fig. 26.—* REEVE, Conch. Syst., I, 1841, p. 118, pl. LXXXVIII, fig. 4.—* HANLEY, Test. Moll., 1842, p. 177; Biv. Shells, 1843, p. 177, pl. XXIV, fig. 5.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—

* KUSTER, Conch. Cab. Unio, 1856, p. 167, pl. XLVIII, fig. 5.—* H. and A.

¹ A large genus of generally rather small species, very abundant in individuals, and distributed from the Amoor River throughout the mainland of all southeastern Asia and all of Africa except the region bordering on the Mediterranean. Some of the African species are so close to Asiatic forms of the *caruleus* group that they can hardly be separated specifically.

ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Man., 1859, II, p. 142, fig. 700.—* REEVE, Conch. Icon., XVI, 1865, pl. XXV, fig. 191.—* HEUDE, Conch. Fluv. Nank., II, 1877, pl. XVI, fig. 36.—* PÆTEL, Couch. Sam., III, 1890, p. 154.

* *Margarita (Unio) grayanus* LEA, Syn., 1838, p. 14.

* *Margaron (Unio) grayanus* LEA, Syn., 1852, p. 21; 1870, p. 32.

* *Unio grayii* GRIFFITH, Grif. Cuv., XII, 1834, pl. XXI, fig. 3.¹

China.

† NODULARIA GLADIOLUS Heude.

* *Unio gladiolus* HEUDE, C. Fluv. Nank., II, 1877, pl. XVI, fig. 35.—* PÆTEL, Couch. Sam., III, 1890, p. 154.

China.

† NODULARIA TRIFORMIS Heude.

* *Unio triformis* HEUDE, Conch. F. Nank., II, 1877, pl. XVI, fig. 34.

* *Unio distortus* HEUDE, Conch. F. Nank., VIII, 1883, pl. LXII, figs. 122, 122b.—

* PÆTEL, C. Sam., III, 1890, p. 151.

China.

† NODULARIA OXYRHYNCHUS von Martens.

* *Unio oxyrhynchus* VON MARTENS, Mal. Blatt, VII, 1861, p. 57.—* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 420, pl. XIII, figs. 3, 4.—* PFEIFFER, Nov. Conch., V, 1879, p. 192, pl. CLVII, figs. 4, 6.—* PÆTEL, C. Sam., III, 1890, p. 162.—* VON IHERING, Abh. Senck. N. Ges., XVIII, 1893, p. 156.

Japan.

Section CYLINDRICA Simpson, 1900.

(Type, *Nodularia cylindrica* Simpson.)

Shell inflated, solid, cylindrical, smooth, with a rounded posterior ridge; teeth heavy, pseudocardinals radiate, curved.

NODULARIA CYLINDRICA Simpson.

* *Unio grayanus* SCHRENCK,² Reis. und Forsch. im Am. Lande, 1867, p. 694, pl. XXVII, figs. 1-3.

Nodularia cylindrica SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 84.

Amur land.

Section NODULARIA Conrad, 1853.

(Type, *Unio douglasie* Gray.)

Characters the same as the genus.

(Group of *Nodularia douglasie*.)

Shell olive to blackish, subsolid; surface generally more or less covered with nodules arranged in subradiating or chevron-shaped patterns.

¹ I have seen a curious shell in the collection of Mr. Bryant Walker shaped something like the *N. grayana*, but quite heavy and decidedly inflated in the post-basal region. It may be an old *grayana*, but is so eroded that I can not be certain.

² Schrenck supposes this to be the *Unio grayanus* of Lea, but it is very different. It is a large, heavy, cylindrical shell, apparently without sculpture, and the figure does not show plainly whether it has any posterior ridge. A large form between *grayanus* and *oxyrhynchus* is found in Korea, according to von Martens, Sitzber. Nat. Fr., 1885, p. 77.

†*NODULARIA DOUGLASIÆ* Gray.¹

* *Unio douglasæ* GRAY, Griff. An. King., XII, 1833, (p. 601 index, 1834), pl. XXI, fig. 2.—* *PÆTEL*, Conch. Sam., III, 1890, p. 151.

*† *Unio murchisonianus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 33, pl. III, fig. 6; * *Obs.*, I, 1834, p. 145, pl. III, fig. 6.—* HANLEY, Test. Moll., 1842, p. 177; * *Biv. Shells*, 1843, p. 177, pl. XXI, fig. 53.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* KUSTER, Conch. Cab. Unio, 1856, p. 166, pl. XLVIII, figs. 3, 4.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Man., 1859, II, p. 142, fig. 701.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVIII, fig. 207.—* *PÆTEL*, Conch. Sam., III, 1890, p. 160.—* VON INERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 159.

* *Margarita (Unio) murchisonianus* LEA, Syn., 1836, p. 14; 1838, p. 14.

* *Margaron (Unio) murchisonianus* LEA, Syn., 1852, p. 21; 1870, p. 32.

* *Unio osbecki* PHILIPPI, Zeits. für Mal., 1845, p. 164; * *Conch.*, III, 1847, p. 45, pl. III, fig. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* KUSTER, Conch. Cab. Unio, 1861, p. 236, pl. LXXIX, fig. 3.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIX, fig. 353.—* *PÆTEL*, Conch. Sam., III, 1890, p. 162.

* *Margaron (Unio) osbecki* LEA, Syn., 1852, p. 21; 1870, p. 32.

*† *Unio nux-persicus* DUNKER, Zeits. für Mal., 1848, p. 83.²—* MUSGRAVE, Phot. Conch., 1863, pl. I, fig. 10.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIII, fig. 373; * *PÆTEL*, Conch. Sam., III, 1890, p. 161.

* *Margaron (Unio) nux-persicus* LEA, Syn., 1870, p. 32.

*† *Unio wrightii* LEA, Pr. Ac. N. Sci., Phila., IX, 1865, p. 75; * *Jl. Ac. N. Sci. Phila.*, VI, 1868, p. 283, pl. XXXIX, fig. 97; * *Obs.*, XII, 1869, p. 43, pl. XXXIX, fig. 97.

* *Margaron (Unio) wrightii* LEA, Syn., 1870, p. 32.

* *Unio sculptus* DESHAYES, Bull. Nonv. Arch. Mus., IX, 1873, p. 9, pl. I, figs. 3, 3a.—* HEUDE, Conch. F. Nank., III, 1877, pl. XXIV, fig. 51.—* *PÆTEL*, Conch. Sam., III, 1890, p. 167.

* *Unio dactylinus* HEUDE, Conch. F. Nank., 1885, pl. LXV.³

*? *Unio pictorum* var. *longirostris* WESTERLUND, Kong so. vet. Ak. Hand, XIV, No. 12, p. 74.⁴

Unio schrencki WESTERLUND. Where described?

Unio abbreviatus WESTERLUND. Where described?⁵

†*NODULARIA DOUGLASIÆ* var. *SHANGHAIENSIS* Lea.

*† *Unio shanghaiensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 153;⁶ * *Jl. Ac. N. Sci. Phila.*, IV, 1860, p. 242, pl. XXXVI, fig. 121; * *Obs.*, VII, 1860, p. 60, pl. XXXVI, fig. 121.—* REEVE, Conch. Icon., XVI, 1865, pl. XXI, fig. 96.—* *PÆTEL*, Conch. Sam., III, 1890, p. 167.

* *Unio pictorum* var. *amurensis* MOUSSON, *Jl. de Conch.*, XXXV, 1887, p. 26, pl. I, fig. 8.

¹The plate in Griffith's Cuvier on which this species is figured bears date of 1833, and the names of the species are designated on it. The brief description (in the index) was published in 1834, the date of the work. Lea's *Unio murchisonianus* was not published until 1834. This species has an immense distribution and varies greatly in form and degree of sculpture.

²An author's shell in the Lea collection is certainly a young *U. douglasæ*.

³This and *U. firmus* and *chloreus* with others are given as varietal names and lettered.

⁴Credited by Westerlund to Amurland. *Unio pictorum*, or varieties of it, have several times been credited to this region, but no doubt erroneously. This species is always darker when adult than the well-known European species, and has totally different beak sculpture. The two shells often resemble each other closely in form.

⁵Specimens of shells bearing the above names, labeled by Westerlund, are in the U. S. National Museum.

⁶A smooth form of *douglasæ*, possibly worthy of a varietal name.

† *NODULARIA DOUGLASIÆ* var. *NIPPONENSIS* von Martens.¹

- * *Unio nipponensis* VON MARTENS, S. B. Nat. Fr., 1877, p. 119.—* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 422, pl. XII, fig. 3.—* PÆTEL, Conch. Sam., III, 1890, p. 161.—* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 160.

China; Amurland; Japan.

† *NODULARIA DORRI* Wattebled.

- * *Unio dorri* WATTEBLED JI. de Conch., XXXIV, 1886, p. 71, pl. V, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

Cochin China; Anam; Mekong river.

NODULARIA DIGITIFORMIS Sowerby.

- * *Unio digitiformis* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXV, fig. 333.—* PÆTEL, Conch. Sam., III, 1890, p. 150.²

India.

(Group of *Nodularia japonensis*.)

Shell solid, elliptical, subrhomboid or oval, biangulate, rounded or pointed behind and often produced at the posterior base; posterior ridge low or scarcely developed; beaks moderately full; sculpture zig-zag radial; surface of the valves concentrically striate or more or less covered with corrugated chevron-shaped sculpture which often becomes somewhat nodulous; epidermis greenish and sometimes rayed in young shells, becoming brown or blackish when adult.

Animal having the inner gills wider than the outer in front, equal behind, free from the abdominal sac throughout, united to the mantle to the extreme posterior point; palpi rather large, long; mantle double on the border and toothed below; branchial opening large, with well-developed, brown papillæ; anal opening large, with small tubercular papillæ on its inner edge, covered with wart-like papillæ inside; super-anal opening not closed below.³

† *NODULARIA JAPANENSIS* Lea.

- * *Unio japonensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 153; * JI. Ac. N. Sci., Phila., IV, 1860, p. 244, pl. XXXVI, fig. 123; * Obs., VII, 1860, p. 62, pl. XXXVI, fig. 123.—* VON MARTENS, Mal. Bl., VII, 1861, p. 55.—* KUSTER, Conch. Cab. Unio, 1862, p. 227, pl. XCIII, fig. 4.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXX, fig. 420.—* PÆTEL, Conch. Sam., III, 1890, p. 155.—* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 157.

- * Margaron (*Unio*) *japonensis* LEA, Syn., 1870, p. 31.

¹ Most of the Japanese forms of this species are smaller than those found on the mainland, though I have seen nothing that exactly agrees with Kobelt's figure. The ordinary form of *douglasie* is found also in Japan.

² This seems to stand between this group and that of *N. grayanus*.

³ The shells of some of the species of this group are very different from the normal *Nodularia*, being covered with roughish epidermis and apparently lacking sculpture. Others, however, show occasional sculpture, and the young shells have the green epidermis characteristic of the group.

NODULARIA JAPANENSIS var. **JOKOHAMENSIS** von Ihering.

* *Unio japanensis* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 423, pl. XII, figs. 1, 2.¹

* *Unio yokohamensis* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 158.

Japan.

† **NODULARIA HACONENSIS** von Ihering.

* *Unio haconensis* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 161, fig. 3.

Japan.

NODULARIA BRANDTII Kobelt.

* *Unio brandtii* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 426, pl. XXIII, fig. 15.—

* PÆTEL, Conch. Sam., III, 1890, p. 146.—* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 157.²

Japan.

NODULARIA BIWÆ Kobelt.

* *Unio biwæ* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 425, pl. XXIII, figs. 2-4.—

* PÆTEL, Conch. Sam., III, 1890, p. 146.—* VON IHERING, Abh. Senck. Nat. Ges., XVII, 1893, p. 157.

Japan.

NODULARIA OMIENSIS von Heimbürg.

* *Unio omiensis* VON HEIMBURG, Nach. Deuts. Mal. Ges., XVI, 1884, p. 93; * Jahrbuch Mal. Ges., XIV, 1887, p. 2, pl. 1, figs. 2, 3.—* PÆTEL, Conch. Sam., III, 1890, p. 161.

Province of Omi, Japan.

(Group of *Nodularia reiniana*.)

Shell elliptical oval, very solid, truncated in front and somewhat biangulate behind, slightly produced at the posterior base; beak sculpture not known; pseudocardinals heavy, elongated, parallel with the dorsal line. Animal unknown.³

NODULARIA REINIANA Kobelt.

* *Unio reinianus* KOBELT, Abh. Senck. Nat. Ges., XI, 1879, p. 424, pl. XXIII, fig.

1.—* VON IHERING, Abh. Senck. Nat. Ges., XVIII, 1893, p. 157.

Japan.

(Group of *Nodularia sagittaria*.)

Shell elongated, the hinder point above the middle of the shell, with a sharp posterior ridge and a faint ridge above it; beaks rather low,

¹ Kobelt figured this for Lea's species, and it seems to me to be an old, solid, slightly arcuate form, possibly worthy of a varietal name.

² It is quite probable that this and the preceding species are only varieties of *N. japonensis*.

³ The beaks of the specimen figured are greatly eroded. I can not be certain as to the relationship of this curious form, but it probably is a *Nodularia*.

their sculpture irregularly radial; disks finely concentrically striate, not otherwise sculptured; epidermis yellowish olive, cloth-like; pseudo-cardinals greatly elongated, compressed, sometimes dentellate under the beaks; laterals long, compressed. Animal unknown.

† *NODULARIA SAGITTARIA* Lea.

* *Unio sagittarius* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 93; *Jl. Ac. N. Sci. Phila., III, 1857, p. 298, pl. XXVI, fig. 12; *Obs., VI, 1857, p. 18, pl. XXVI, fig. 12.

* *Margaron (Unio) sagittarius* LEA, Syn., 1870, p. 58.

*† *Unio dugasti* MORLET, Jl. de Conch., XL, 1892, p. 86; *XLI, 1893, p. 156, pl. VI, fig. 4.¹

Siam; Cambodia.

(Group of *Nodularia cærulea*.)

Shell elliptical, inflated, pointed about midway up behind, the post-basal region produced, with a well developed posterior ridge; beaks sculptured with numerous fine, radiating riblets, the central ones of which join below, the whole often more or less zigzagged and extending well over the disk; epidermis generally bluish green. Animal unknown.

† *NODULARIA CÆRULEA* Lea.

* *Unio cæruleus* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 95, pl. XIII, fig. 25; *Obs., I, 1834, p. 105, pl. XIII, fig. 25.—*HANLEY, Test. Moll., 1842, p. 194; *Biv. Shells, 1843, p. 194, pl. XX, fig. 49.—*CATLOW and REEVE, Conch. Nom., 1845, p. 56.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*CHENU, III. Conch., 1858, pl. XVI, figs. 3, 3a, 3b.—*KUSTER, Conch. Cab. Unio, 1861, p. 228, pl. LXXVII, fig. 4.—*SOWERBY, Conch. Icon., XVI, 1866, pl. XLII, fig. 230.—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 6, pl. XII, figs. 3, 3a.—*PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Margarita (Unio) cæruleus* LEA, Syn., 1836, p. 26; 1838, p. 20.

* *Margaron (Unio) cæruleus* LEA, Syn., 1852, p. 30; 1870, p. 47.

* *Unio gerbidoni* EYDOUX, Guer. Mag., 1838, p. 9, pl. CXVIII, figs. 2, 2a, 2b.—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 6, pl. XII, fig. 2.—*PÆTEL, Conch. Sam., III, 1890, p. 153.

*† *Unio substriatus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 93; *Obs., VI, 1857, p. 20, pl. XXVI, fig. 14; *Jl. Ac. N. Sci. Phila., III, 1858, pl. XXVI, fig. 14.

* *Margaron (Unio) substriatus* LEA, Syn., 1870, p. 47.

*† *Unio humilis* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 93; *Obs., VI, 1857, p. 16, pl. XXVI, fig. 10; *Jl. Ac. N. Sci. Phila., III, 1858, p. 296, pl. XXVI, fig. 10.—*PÆTEL, Conch. Sam., III, 1890, p. 155.

* *Margaron (Unio) humilis* LEA, Syn., 1870, p. 32.

* *Unio corrianus* KUSTER, Conch. Cab. Unio, 1861, p. 229, pl. LXVII, fig. 5.

*† *Unio leioma* BENSON, Ann. and Mag., 1862, p. 192.—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 6, pl. XII, fig. 6.—*PÆTEL, Conch. Sam., III, 1890, p. 157.

*† *Unio pilatus* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 133; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 281, pl. XXXVIII, fig. 95; *Obs., XII, 1869, p. 41, pl. XXXVIII, fig. 95.

*† *Margaron (Unio) pilatus* LEA, Syn., 1870, p. 47.

*† *Unio evitatus* LEA, Pr. Ac. N. Sci. Phila., X, 1868, p. 133; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 279, pl. XXXVIII, fig. 92; *Obs., XII, 1869, p. 39, pl. XXXVIII, fig. 92.

¹ This group seems to stand in part between section *Lanceolaria* and *Nodularia*.

* *Margaron (Unio) evitatus* LEA, Syn., 1870, p. 47.

* *Unio tirostris* SOWERBY, Conch. Icon., XVI, 1868, pl. LXV, fig. 331.

? *Unio andersonianus* NEVILL, Jl. As. Soc. Beng., XLVI, 1877, p. 40; Researches Yun. Exp. 1877, p. 900, pl. LXXX, figs. 8, 12.¹

India; southeast Asia.

† NODULARIA TERETIUSCULA Philippi.

* *Margarita (Unio) cailliaudi* LEA, Syn., 1838, p. 24.

* *Margaron (Unio) cailliaudi* LEA, Syn., 1852, p. 36; 1870, p. 58.²

* *Unio teretiusculus* PHILIPPI, Conch., III, 1847, p. 45, pl. III, fig. 3.³—* KUSTER, Conch. Cab., 1856, p. 133, pl. XXXV, fig. 5.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXV, fig. 389.—* JICKELL, Faun. N. O. Af., 1874, p. 276, pl. XI, figs. 1-3.—

* PÆTEL, Conch. Sam., III, 1890, p. 169.—* VON MARTENS, Besch. Dents. Ost-Af., 1897, p. 223.

Unio lithophagus ZIEGLER, manuscript, probably.

Upper Nile drainage.

NODULARIA LEDOULXIANA Charmes.

* *Unio ledoulxianus* CHARMES, Bull. Soc. Mal. Fr., II, 1885, p. 173.—* BOURGUIGNAT, Moll. Af. Eq., 1889, p. 194.—* VON MARTENS, Besch., 1897, p. 229.⁴

Central Africa.

NODULARIA GAUDICHAUDI Eydoux.

* *Unio gaudichaudi* EYDOUX, Guer. Mag., 1838, p. 10, pl. CXVIII, fig. 3.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* PÆTEL, Conch. Sam., III, 1890, p. 153.

* *Margaron (Unio) gaudichaudi* LEA, Syn., 1850, p. 32; 1870, p. 50.

Small streams of Bengal.

NODULARIA KERAUDRENI Eydoux.

* *Unio keraudreni* EYDOUX, Guer. Mag., 1838, p. 8, pl. CXVIII, figs. 1, 1a.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* PÆTEL, Conch. Sam., III, 1890, p. 156.

* *Margaron (Unio) keraudreni* LEA, Syn., 1852, p. 30; 1870, p. 46.

Rivers and swamps of Bengal.

¹ I have not seen the figure of *andersonianus*, but Nevill states that it is near *cæruleus*. The *Nodularia cærulea* is an abundant and very widely distributed form, and therefore quite variable in size, form, solidity, color, and sculpture, and it is possible that other so-called species should be merged with it. Some of the above might rank as varieties, but I can hardly separate them.

² Lea credits this name to Ferussac manuscript. So far as I know neither Ferussac or anyone else has ever described it. Lea makes *U. teretiusculus* a synonym of it in his Synopsis.

³ Philippi's figure is a little more rounded posteriorly than are any of the shells I have seen from Africa bearing this name. After a critical comparison of numerous specimens, which are undoubtedly of African origin, I can only place this in the *cæruleus* group (with which it agrees in every essential character) and close to *cæruleus* itself. There are other undoubtedly close relations between species of *Naiades* of the Ethiopian and Oriental regions.

⁴ Unfigured. Von Martens places it in the group of *U. teretiusculus*.

† NODULARIA SHURTLEFFIANA Lea.

- * *Unio shurtleffianus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 94; * Obs., VI, 1857, p. 22, pl. XXVII, fig. 17; * Jl. Ac. N. Sci. Phila., III, 1858, p. 302, pl. XXVII, fig. 17.—
 * PÆTEL, Conch. Sam., III, 1890, p. 167.
 * Margaron (*Unio*) *shurtleffianus* LEA, Syn., 1870, p. 32.

India; Siam.

† NODULARIA OCCATA Lea.

- * *Unio occatus*, LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 307; * Jl. Ac. N. Sci. Phila., VI, 1863, p. 398, pl. L, fig. 304; * Obs., X, 1863, p. 34, pl. L, fig. 304.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIX, fig. 412.—* PÆTEL, Conch. Sam., III, 1890, p. 161.
 * Margaron (*Unio*) *occatus*, LEA, Syn., 1870, p. 31.
 * † *Unio macilentus*, BENSON, Ann. and Mag., X, 1862, p. 187.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 5, pl. X, figs. 2, 4; p. 62, pl. CLIV, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 158.
 * *Unio rugosus*, HANLEY and THEOBALD, Conch. Ind., 1876, p. 62, pl. CLIV, fig. 3.

India.

NODULARIA MOSSAMBICENSIS von Martens.

- * *Unio mossambicensis* VON MARTENS (Peter's manuscript), Mal. Bl., VI, 1860, p. 218, pl. III, figs. 3-5.—* PÆTEL, Conch. Sam., III, 1890, p. 160.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 225, pl. VII, fig. 2.
 * *Unio sennariensis* var. VON MARTENS, Mal. Bl., XXI, 1873, p. 43.
 * *Unio parreysi* (v. d. Busch.) var. *schwiefurthi* VON MARTENS, Nov. Conch., IV, 1876, p. 140, pl. CXXXII, figs. 3-5.¹

Sennaar, northeast Africa; Mosambique, Zambezi River.

† NODULARIA INORNATA Lea.

- * *Unio inornatus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 93; * Obs., VI, 1857, p. 13, pl. XXIV, fig. 6; * Jl. Ac. N. Sci. Phila., III, 1858, p. 293, pl. XXIV, fig. 6.—* VON MARTENS, Pr. Zool. Soc. Lond., 1860, p. 15.—* MORELET, Ser. Conch., IV, 1875, p. 352.—PÆTEL, Conch. Sam., III, 1890, p. 155.
 * Margaron (*Unio*) *inornatus* LEA, Syn., 1870, p. 47.

Siam; Cambodia.

† NODULARIA PACHYSOMA Benson.

- * *Unio pachysoma* BENSON, Ann. and Mag., X, 1862, p. 186.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 6, pl. XII, fig. 1.
 * Margaron (*Unio*) *pachysoma* LEA, Syn., 1870, p. 63.
 * *Unio pachystoma* PÆTEL, Conch. Sam., III, 1890, p. 132.

Assam.

† NODULARIA BONNEAUDI Eydoux.

- * *Unio bonneaudi* EYDOUX, Guer. Mag., 1838, p. 10, pl. CXIX, figs. 1, 1a.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1868,

¹ This species seems to be close to *N. occata*, but is thinner and of delicate texture. I am not certain of its exact affinities. I am sure that it is not a variety of *N. mossambicensis*.

pl. xcv, fig. 515.—* HANLEY and THEOBALD, *Conch. Ind.*, 1876, p. 5, pl. x, fig. 6.—ANDERSON, *Yunnan Exp.*, 1878, p. 900, pl. LXXX, figs. 8-12.—* PÆTEL, *Conch. Sam.*, III, 1890, p. 146.

* *Margaron (Unio) bonneaudi* LEA, *Syn.*, 1852, p. 32; 1870, p. 50.

India.

† NODULARIA TUMIDULA Lea.

* *Unio tumidulus* LEA, *Pr. Ac. N. Sci. Phila.*, VIII, 1856, p. 93; *Obs., VI, 1857, p. 15, pl. xxv, fig. 9; **Jl. Ac. N. Sci. Phila.*, III, 1858, p. 295, pl. xxv, fig. 9.—* VON MARTENS, *Pr. Zool. Soc. Lond.*, 1860, p. 15.—* SOWERBY, *Conch. Icon.*, XVI, 1868, pl. LXXXIX, fig. 482.—* MORELET, *Ser. Conch.*, IV, 1875, p. 351.—* PÆTEL, *Conch. Sam.*, III, 1890, p. 170.

* *Margaron (Unio) tumidulus* LEA, *Syn.*, 1870, p. 39.

Siam, Cambodia.

† NODULARIA ASPERULA Lea.

* *Unio inornatus* REEVE, *Conch. Icon.*, XVI, 1865, pl. xxix, fig. 147.¹

* *Unio asperulus* LEA, *Pr. Ac. N. Sci. Phila.*, X, 1866, p. 133; **Jl. Ac. N. Sci. Phila.*, VI, 1868, p. 280, pl. xxxviii, fig. 94; *Obs., XII, 1869, p. 40, pl. xxxviii, fig. 94.

* *Margaron (Unio) asperulus* LEA, *Syn.*, 1870, p. 31.

* *Margaron (Unio) versus* LEA, *Syn.*, 1870, p. 46.

Siam; Cambodia.

NODULARIA ANCEPS Deshayes.

* *Unio anceps* DESHAYES, *Nouv. Arch. de Mus.*, X, 1874, p. 127, pl. vi, figs. 8-12.—

* PÆTEL, *Conch. Sam.*, III, 1890, p. 144.

Cambodia.

† NODULARIA PUGIO Benson.

* *Unio pugio* BENSON, *Ann. and Mag.*, X, 1862, p. 193.—* SOWERBY, *Conch. Icon.*, XVI, 1868, pl. xcv, fig. 516.—* PÆTEL, *Conch. Sam.*, III, 1890, p. 164.²

Burma.

NODULARIA MICHELOTI L. Morlet.

* *Unio micheloti* L. MORLET, *Jl. de Conch.*, 1886, pp. 77, 291, pl. xiii, figs. 6, 6a.—

* PÆTEL, *Conch. Sam.*, III, 1890, p. 159.

Tonkin.

NODULARIA FLUCTIGER Lea.

* *Unio fluctiger* LEA, *Pr. Ac. N. Sci. Phila.*, III, 1859, p. 152; **Jl. Ac. N. Sci. Phila.*, IV, 1860, p. 250, pl. xxxix, fig. 130; *Obs., VII, 1860, p. 68, pl. xxxix, fig. 130.—

* KUSTER, *Conch. Cab. Unio*, 1861, p. 237, pl. LXXX, fig. 1.—* SOWERBY, *Conch. Icon.*, XVI, 1866, pl. xlii, fig. 299.

* *Margaron (Unio) fluctiger* LEA, *Syn.*, 1870, p. 32.

Southeast Asia, no doubt.

¹ I believe that Reeve's *U. inornatus* = Lea's *U. asperulus*. As Lea had used the name *inornatus* previous to Reeve for a *Unio*, he changed the name of the latter to *U. versus*.

² A shell labeled *Unio bourieri* Nevill in the collection of Frederick Stearns in Detroit is, I believe, the same as *U. pugio* Benson. I do not know if Nevill ever described *U. bourieri* or whether it is only a manuscript name.

†NODULARIA SCOBINATA Lea.

* *Unio scobinata* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 93; * Obs., VI, 1857, p. 19, pl. XXVI, fig. 13; * Jl. Ac. N. Sci. Phila., III, 1858, p. 299, pl. XXVI, fig. 13.—* VON MARTENS, Pr. Zool. Soc. Lond., 1860, p. 15.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIII, fig. 313.—* MORELET, Ser. Conch., IV, 1875, p. 354, pl. XVII, figs. 2-6.—* PÆTEL, Conch. Sam., III, 1890, p. 167.

* *Margaron (Unio) scobinatus* LEA, Syn., 1870, p. 32.

* *Unio mandarinus* MORELET, Jl. de Conch., XII, 1863, p. 159.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

* *Unio pellis-lacerti* MORELET, Jl. de Conch., XIII, 1865, p. 22.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVI, fig. 457.—* MORELET, Ser. Conch., IV, 1875, p. 355, pl. XVII, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Unio venustus* MORELET, Jl. de Conch., XIV, 1866, p. 63.—* PÆTEL, Conch. Sam., III, 1890, p. 171.

* *Unio oblatas* LEA, Syn., 1870, p. 64.¹

Siam; Cochinchina; Cambodia.

NODULARIA GRATIOSA Philippi.²

* *Unio graciosus* PHILIPPI, Conch., I, 1845, p. 20, pl. I, fig. 5.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* ? KUSTER, Conch. Cab. Unio, 1861, p. 239, pl. LXXX, fig. 3.

* *Margaron (Unio) graciosus* LEA, Syn., 1870, p. 32.

India probably; Philippi's locality, New Holland, is certainly an error.

†NODULARIA CRISPATA Gould.

* *Unio crispata* GOULD, Pr. Bost. Soc. Nat. Hist., I, 1843, p. 141; * Otia Conch., 1862, p. 191.

* *Unio crispatus* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 21, pl. XLV, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 149.

* *Margaron (Unio) crispatus* LEA, Syn., 1870, p. 32.

British Burma.

†NODULARIA NUCLEUS Lea.

* *Unio nucleus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 94; * Obs., VI, 1857, p. 26, pl. XXVIII, fig. 21; * Jl. Ac. N. Sci. Phila., III, 1858, p. 303, pl. XXVIII, fig. 21.

* *Margaron (Unio) nucleus* LEA, Syn., 1870, p. 30.

Siam.

†NODULARIA PHASELUS Lea.

* *Unio phaselus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 94; * Obs., VI, 1857, p. 17, pl. XXVI, fig. 11; * Jl. Ac. N. Sci. Phila., III, 1858, p. 297, pl. XXVI, fig. 11.

* *Margaron (Unio) phaselus* LEA, Syn., 1870, p. 32.

Siam.

¹ New name for *Unio venustus* Morelet, preoccupied by Lea.

² Philippi's figure is not very distinct, and I can not be sure of what it is, only that it seems to be a member of this group that I can hardly connect with anything else. He credits it to Parreyss.

NODULARIA SCOBINA Hanley.

- * *Unio scobina* HANLEY, Biv. Shells, 1856, p. 382, pl. XXIII, fig. 40.¹—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 22, pl. XLVI, figs. 2, 3.—* P.ETEL, Conch. Sam., III, 1890, p. 167.

Assam.

(Group of *Nodularia pazii*.)

Shell thin, with the hinder point long drawn out; posterior ridge double, low, and sharp, the dorsal line above it straight or incurved; beak sculpture not seen; surface of the shell smooth, dull olive colored; teeth exceedingly compressed; pseudocardinals long, lamellar.

† NODULARIA PAZII.

- * *Unio pazii* LEA, Pr. Ac. N. Sci. Phila., VI, 1862, p. 176; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 61, pl. XXI, fig. 60; *Obs., XI, 1867, p. 65, pl. XXI, fig. 60.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCII, fig. 502.
* *Margaron (Unio) pazii* LEA, Syn., 1870, p. 39.

China; Siam; Cambodia.

NODULARIA JOURDYI L. Morlet.

- * *Unio jourdyi* MORLET, Jl. de Conch., 1886, p. 289, pl. XIII, figs. 5, 5a.

Tonkin.

† NODULARIA INGALLSIANA Lea.

- * *Unio ingallsianus* LEA, Tr. Am. Phil. Soc., X, 1852, p. 282, pl. XXIV, fig. 41; *Obs., V, 1852, p. 38, pl. XXIV, fig. 41.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* REEVE, Conch. Icon., XVI, 1865, pl. XXVI, fig. 126.—* MORELET, Ser. Conch., IV, 1875, p. 351.—* P.ETEL, Conch. Sam., III, 1890, p. 155.
* *Margaron (Unio) ingallsianus* LEA, Syn., 1852, p. 36; 1870, p. 58.

Cochin-China; Siam; Cambodia.

(Group of *Nodularia trompi*.)

Shell small, subtrapezoidal, wider behind, slightly inflated, with a rounded posterior ridge; beaks full, sculpture not seen; surface covered with very fine, nodulous, somewhat zigzag corrugations, with upcurved posterior ridges on the posterior slope; epidermis reddish brown or black; hinge teeth compressed; pseudocardinals vertically striate.

† NODULARIA TROMPI Drouet and Chaper.

- * *Unio trompi* DROUET and CHAPER, Mem. Soc. Zool. Fr., V, 1892, p. 153, pl. VI, figs. 8-10; Jl. de Conch., XLI, 1893, p. 45.

Borneo.

(Group of *Nodularia olivaria*.)

Shell thin, inflated, blue green or olive green, smooth, shining, with two or more faint ridges on the posterior slope; beaks and disk almost or quite destitute of sculpture; teeth greatly compressed, lamellar.

¹ Credited to Benson manuscript.

† NODULARIA OLIVARIA Lea.

- * *Unio olivarius* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 108, pl. XVI, fig. 38; * Obs., I, 1834, p. 118, pl. XVI, fig. 38.—* HANLEY, Test. Moll., 1842, p. 195; * Biv. Shells, 1843, p. 195, pl. XXII, fig. 32.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. VIII, figs. 11, 11a, 11b.—* KUSTER, Conch. Cab. Unio, 1861, p. 244, pl. LXXXII, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXVI, fig. 195.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 5, pl. X, fig. 1.—PÆTEL, Conch. Sam., III, 1890, p. 161.
- * *Margarita (Unio) olivarius* LEA, Syn., 1836, p. 26; 1838, p. 20; 1852, p. 30; 1870, p. 47.
- * *Unio pumilio* KUSTER, Conch. Cab. Unio, 1862, p. 268, pl. XC, fig. 7.

India.

† NODULARIA NUTTALLIANA Lea.

- * *Unio nuttallianus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 103; * Obs., VI, 1857 p. 30, pl. XXX, fig. 25; Jl. Ac. N. Sci. Phila., III, 1858, p. 310, pl. XXX, fig. 25.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 19, pl. XLI, figs. 5, 6.—* PÆTEL, Conch. Sam., III, 1890, p. 161.
- * *Margaron (Unio) nuttallianus* LEA, Syn., 1870, p. 74.

India.

NODULARIA INVOLUTA Benson.

- * *Unio involutus* BENSON, Hanley Biv. Shells, 1856, p. 385, pl. XXIII, fig. 19.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 19, pl. XLI, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 156.
- * *Margaron (Unio) involutus* LEA, Syn., 1870, p. 37.
- * *Unio involuta* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXIV, fig. 177.

Assam.

Group of *Nodularia contradens*.)

Shell inflated, obovate, rhomboid, with a slight swelling at the post base, and a well-developed posterior ridge; beaks full and high, the sculpture consisting of wavy, zigzag ridges which often extend over the shell as somewhat nodulous corrugations; pseudocardinals obliquely or vertically striate; laterals curved, cavity of the beaks rather shallow.

† NODULARIA CONTRADENS Lea.

- * *Unio contradens* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 75, pl. XVIII, fig. 58; * Obs., II, 1838, p. 75, pl. XVIII, fig. 58.—* TROSCHEL, Arch. für Nat., V, 1839, Pt. 2, p. 237.—* HANLEY, Test. Moll., 1842, p. 209; Biv. Shells, 1843, p. 209, pl. XXII, fig. 8.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* CHENU, Ill. Conch., 1858, pl. XXIX, figs. 4, 4a, 4b.—* REEVE, Conch. Icon., XVI, 1865, pl. XXIX, fig. 149.—* PÆTEL, Conch. Sam., III, 1890, p. 148.
- * *Margarita (Unio) contradens* LEA, Syn., 1838, p. 25.
- * *Margaron (Unio) contradens* LEA, Syn., 1852, p. 39; 1870, p. 46.
- * + *Unio javanus* LEA, Pr. Am. Phil. Soc., I, 1840, p. 285; * Tr. Am. Phil. Soc., VIII, 1842, p. 220, pl. XVIII, fig. 37; * Obs., III, 1842, p. 58, pl. XVIII, fig. 36.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. XXVIII, figs. 4, 4a, 4b.—* KUSTER, Conch. Cab. Unio, 1856, p. 138, pl. XLI, fig. 3; 1861, p. 236, pl. LXXIX, figs. 4, 6.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XC, fig. 489.—* PÆTEL, Conch. Sam., III, 1890, p. 155.

* *Margaron (Unio) javanus* LEA, Syn., 1852, p. 30.

* *Unio exilis* DUNKER, Zeits. für Mal., III., 1846, p. 109.—MOUSSON, L. und S. W. Moll. Java, 1849, p. 92, pl. XVI, fig. 3.

* *Margaron (Unio) exilis* LEA, Syn., 1852, p. 29.

* *Unio mutatus* MOUSSON, L. und S. W. Moll. Java, 1849, p. 92, pl. XVI, figs. 1, 2; Zeits. für Mal., VII, 1851, p. 46; *H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*PÆTEL, Conch. Sam., III, 1890, p. 160.

* *Margaron (Unio) mutatus* LEA, Syn., 1852, p. 29.

* *Unio mederianus* KUSTER,¹ Conch. Cab. Unio, 1861, p. 242, pl. LXXX, fig. 7.—*PÆTEL, Conch. Sam., III, 1890, p. 158.

Java.

† NODULARIA DIMOTA Lea.

* *Unio sumatrensis*, LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 153; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 239, pl. XXXIV, fig. 118; *Obs., VII, 1860, p. 57, pl. XXXIV, fig. 118.—*PÆTEL, Conch. Sam., III, 1890, p. 169.

* *Margaron (Unio) dimotus*, LEA, Syn., 1870, p. 30.—² *PÆTEL, Conch. Sam., III, 1890, p. 150.³

Sumatra.

NODULARIA FISCHERIANIA L. Morlet.

* *Unio fischerianus*, L. MORLET, Jl. de Conch., XXXI, 1883, p. 109, pl. IV, fig. 6.—*PÆTEL, Conch. Sam., III, 1890, p. 152.

Cambodia.

† NODULARIA ANODONTÆFORMIS Tapperone-Caneфри.

* *Unio (Microdontia) anodontaformis* TAPPERONE-CANEFRI, Ann. Mus. Genov., XIX, 1883, p. 295, pl. XI, figs. 3-5.

* *Unio anodontaformis* PÆTEL, Conch. Sam., III, 1890, p. 144.

New Guinea.

† NODULARIA SACELLUS Drouet and Chaper.

* *Unio sacellus* DROUET and CHAPER, Mem. Soc. Zool. Fr., V, 1892, p. 148, pl. V, figs. 4-6; Jl. de Conch., XLI, 1893, p. 43.

Borneo.

† NODULARIA LUGENS Drouet and Chaper.⁴

* *Unio lugens* DROUET and CHAPER, Mem. Soc. Zool. Fr., V, 1892, p. 147, pl. V, figs. 1-3.—*DROUET, Jl. de Conch., XLI, 1893, p. 44.

Borneo.

¹ Kuster credits this to Mousson, Conch. of Java, p. 92. I know of no such work, and the name is not found in Mousson's L. und Suss. W. Moll. von Java.

² Changed to this name by Lea because *sumatrensis* was preoccupied for a *Unio* by Dunker.

³ This species varies much in sculpture, and shows the connection between the nearly smooth *contradens* and the strongly corrugated forms like *rusticus* and *dautzenbergi*. It is really very close to *contradens*.

⁴ I am doubtful whether these two are more than varieties of the same thing. The *N. lugens* is darker colored and seems to lack a little tubercle on the hinge, which is sometimes present in *sacellus*.

† **NODULARIA FULVASTER** Drouet and Chaper.

* *Unio fulvaster* DROUET and CHAPER, Mem. Soc. Zool. de Fr., V, 1892, p. 154, pl. VI, figs. 11-13.—*DROUET, Jl. de Conch., XLI, 1893, p. 45.

Borneo.

NODULARIA SEMIDECORATA L. Morlet.

* *Unio semidecoratus* L. MORLET, Jl. de Conch., XXXVII, 1889, p. 192, pl. VIII, fig. 4.

River Srakeo, Siam.

† **NODULARIA RUSTICA** Lea.

* *Unio rusticus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 93; * Obs., VI, 1857, p. 14, pl. XXV, fig. 7.—*Jl. Ac. N. Sci. Phila., III, 1858, p. 294, pl. XXV, fig. 7.—*VON MARTENS, Pr. Zool. Soc. Lond., 1860, p. 14.—*MORELET, Ser. Conch., IV, 1875, p. 353, pl. XVII, fig. 7.—*PÆTEL, Conch. Sam., III, 1890, p. 166.

* *Margaron (Unio) rusticus* LEA, Syn., 1870, p. 31.

* *Unio paivanus* MORELET, Jl. de Conch., XIII, 1865, p. 227.

* *Unio cambojensis* SOWERBY, Conch. Icon., XVI, 1866, pl. XLII, fig. 231.—*PÆTEL, Conch. Sam., III, 1890, p. 147.

Siam; Cambodia.

† **NODULARIA DAUTZENBERGI** L. Morlet.

* *Unio dautzenbergi* MORLET, Jl. de Conch., XXXVII, 1889, p. 190, pl. VIII, fig. 5.

River Srakeo, Siam.

NODULARIA SOBOLUS Fischer.

* *Unio siamensis* MORLET, Jl. de Conch., XXXVII, 1889, p. 194, pl. VII, fig. 2.

* *Unio sobolus* FISCHER, Bull: Soc. d'Autun., 1891, p. 227.¹

Siam.

NODULARIA VERBECKI Böttger.

* *Unio verbecki* BÖTTGER, Zool. Erg. Nied. Ost. Ind., IV, 1897, p. 89, pl. v, figs. 1, 2, 4, 5.—*PÆTEL, Conch. Sam., III, 1890, p. 171.

Sumatra.

† **NODULARIA ORIENTALIS** Lea.²

* *Unio orientalis* LEA, Pr. Am. Phil. Soc., I, 1840, p. 285; *Tr. Am. Phil. Soc., VIII, 1842, p. 221, pl. XVIII, fig. 38; * Obs., III, 1842, p. 59, pl. XVIII, fig. 38.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—*CHENU, Ill. Conch, 1858, pl. XXXII, figs. 6, 6a, 6b.—*KÜSTER, Conch. Cab. Unio, 1861, p. 241, pl. LXXX, fig. 16.—*VON MARTENS, Mal. Bl., XIV, 1866, p. 15.—*SOWERBY, Conch. Icon., XVI, pl. XCI, fig. 491.—*PÆTEL, Conch. Sam., III, 1890, p. 162.

¹ Changed from *siamensis* to *sobolus* because the former was preoccupied by Lea. Probably an old, smooth *rusticus*.

² I hardly know where to place this, as the only shell I have seen (the type) is evidently young, and has the beaks eroded until I can not make out the sculpture. It seems most likely to be an elongated, somewhat compressed member of the *contradens* group.

* *Margaron (Unio) orientalis* LEA, Syn., 1852, p. 38; 1870, p. 61.

* *Unio productus* MOUSSON, L. und S. W. Moll. Java, 1849, p. 93, pl. xvii, figs. 3-5; Zeits. für Mal., VII, 1851, p. 46.

* *Unio productior* LEA, Syn., 1852, p. 29.—* VON MARTENS, Mal. Bl., XIV, 1866, p. 16.—* II. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.

Java.

Section *RADIATULA*, Simpson.

(Type, *Unio crispisulcatus* Benson.)

Shell rather solid, triangular oval, with high beaks which are but little inflated, not very full at post base, bluntly pointed behind, the beaks and entire surface covered with radiating, occasionally slightly zigzag or divaricate ridges which are cut more or less into nodules or cancellations by concentric sulcations, the sculpture of the posterior slope stronger, and curving upward; pseudocardinals of the left valve 2 to 3, ragged, the anterior larger, two in the right valve, with a parallel sided socket, the larger teeth compressed but rather solid.¹

† *NODULARIA CRISPISULCATA* Benson.

* *Unio crispisulcatus* LEA, Ann. and Mag., X, 1862, p. 193.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLIX, fig. 262.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 6., pl. XI, fig. 5.—* PETEL, Conch. Sam., III, 1890, p. 149.

* *Margaron (Unio) crispisulcatus* LEA, Syn., 1870, p. 31.

Burma.

NODULARIA LIMA Simpson.

* *Unio radula* BENSON,² in Hanley, Biv. Shells, Supp., 1856, p. 382.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 5, pl. x, fig. 3.—* PETEL, Conch. Sam., III, 1890, p. 165.

* *Dysnomia radula* ROCHEBRUNE, Bull. Soc. Phil., VI, 1882, p. 42.

Assam; Cambodia.

Section *CÆLATURA* Conrad, 1853.

(Type, *Unio ægyptiacus* Cailliaud.)

Shell elliptical, pointed or rounded behind, usually slightly produced at the post base; beak sculpture consisting of zigzag ridges which are generally quite pustulous, the sculpture often extending on to the usually rayed disk; teeth compressed; pseudocardinals lamellar.

Animal described under *Nodularia*. (See page 806.)

(Group of *Nodularia ægyptiaca*.)

Characters the same as of the section.

¹Notwithstanding the considerable difference between this group and all other *Nodularias* there are species such as *occatus*, which seem to stand between it and the *cæruleus* group, and show the relationship of the two.

²The name *radula* was used by Say for a *Unio* in 1829. I change Benson's name as above. I am not positive as to the relationship of this species.

† NODULARIA ÆGYPTIACA Cailliaud.

- * *Unio* species nuovo SAVIGNY, Icon. Moli. Egypt, 1813, pl. VII, figs. 3-6.
Unio ægyptiaca CAILLIAUD, Voy. à Méroé, II, 1826, pl. LXI, figs. 6, 7.—* DESHAYES, An. sans Vert., 2d ed., 1835, p. 553; * Enc. Méth., II, 1830, p. 587.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.
 * *Margarita* (*Unio*) *egyptiacus* LEA, Syn., 1838, p. 21.
 * *Margaron* (*Unio*) *egyptiacus* LEA, Syn., 1852, p. 32; 1870, p. 50.
 * *Unio ægyptiacus* HANLEY, Test. Moll., 1842, p. 198; * Biv. Shells, 1843, p. 198, pl. XX, fig. 56.—* KUSTER, Conch. Cab., 1856, p. 157, pl. XLV, fig. 2.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* REEVE, Conch. Icon., XVI, 1865, pl. XXVI, fig. 132.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 179.—* VON MARTENS, Mal. Bl., XIII, 1866, p. 11.—* MORELET, Moll. Terr. et Fluv. 1868, p. 97.—* JICKELI (part) Faun. Moll. N. O. Af., 1874, p. 271, pl. X, figs. 1-8.—* KOBELT, Icon., new ed., 1886, p. 25, pl. XLIV, figs. 262-265.
 * *Unio eueyphus* BOURGUIGNAT, Rev. et. Mag., IX, 1857, p. 19, pl. III, figs. 1-3.—* PÆTEL, Conch. Sam., III, 1890, p. 152.
 * *Unio bourguignati* LANDRAN, Seance Soc. Sci. Nat. Seine, 1864, p. 5, pl. —, figs. 1-3.
 * *Pharaonia bourguignati* ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 113.

Nile system; High Cazamance, West Africa (Vignon); both sides of the equator in Central Africa (Dohrn).

† NODULARIA NILOTICA Cailliaud.

- Unio nilotica* CAILLIAUD, Voy. à Méroé, II, 1826, pl. LXI, figs. 8, 9.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 522; * Enc. Méth., II, 1830, p. 585.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* HANLEY, Test. Moll., 1842, p. 197; * Biv. Shells, 1843, p. 197, pl. XXI, fig. 39.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* VON MARTENS, Mal. Bl., XIII, 1866, p. 13.—* PÆTEL, Conch. Sam., III, 1890, p. 161.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 180.
 * *Margarita* (*Unio*) *niloticus* LEA, Syn., 1836, p. 29; 1838, p. 21.
 * *Margaron* (*Unio*) *niloticus* LEA, Syn., 1852, p. 31; 1870, p. 50.
Unio pumilus ZEIGLER, manuscript, and of authors.
 * *Unio parreyssi* PHILIPPI, Conch., III, 1848, p. 81, pl. v, fig. 6.—* JICKELI, Faun. Moll. N. O. Af., 1874, p. 373.
 * *Unio rugifer* KUSTER, Conch. Cab. Unio, 1856, p. 157, pl. XLV, figs. 3, 4.—* VON MARTENS, Mal. Bl., XIII, 1866, p. 12.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 180.
 * *Unio sennaariensis* KUSTER, Conch. Cab. Unio, 1862, p. 280, pl. XCIV, figs. 5, 6.—* PÆTEL, Conch. Sam., III, 1890, p. 167.
 * *Unio aneus* JICKELI, Faun. Moll. N. O. Af., 1874, p. 274, pl. IX, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 144.
 * *Unio charbonnieri* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 9.—* Icon. Mal. Tan., 1888, pl. XX, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 147.
 * *Unio coulboisi* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 12; * Icon. Mal. Tan., 1888, pl. XX, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 149.
 * *Unio dromauzi* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 17; * Icon. Mal. Tan., 1888, pl. XXII, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

¹This species is often credited to Ferrussac, but I do not know that he ever described it. It is considered by many to be the same as *Nodularia nilotica*, but the comparison of large series of material and authentic specimens in the Lea collection from Cailliaud incline me to separate them. There is, however, great variation in their characters, but *ægyptiaca* is a thinner, more rounded shell than *niloticus*, and has lamellar pseudo-cardinals.

NODULARIA NILOTICA var. GERRARDI von Martens.

* *Unio gerrardi* VON MARTENS, Besch. Ost-Af., 1897, p. 223, pl. VII, fig. 5.

Nile system; Lake Tanganyika; Senegal and High Cazamance (Vignon).

NODULARIA BAGDADENSIS Bourguignat.¹

* *Unio bagdadensis* BOURGUIGNAT, Test. Nov. Saulc., 1852, p. 30; *Cat. Rais., 1853, p. 78, pl. IV, figs. 4-6.

*Margaron (*Unio*) *bagdadensis* LEA, Syn., 1870, p. 46.

Bagdad, on the Tigris River.

NODULARIA RANDABELI Bourguignat.²

* *Unio randabeli* BOURGUIGNAT, Un. and Ir. Tan., 1886, p. 22; *Icon. Mal. Tan., 1888, pl. XXI, figs. 3, 4.—*P.ETEL, Conch. Sam., III, 1890, p. 165.

Lake Tanganyika.

† NODULARIA ESSOENSIS Chaper.

* *Unio essoensis* CHAPER, Bull. Soc. Zool. Fr., X, 1885, p. 481, pl. XI, figs. 8, 9.—

*P.ETEL, Conch. Sam., III, 1890, p. 152.

Assini, west coast of Africa.

NODULARIA DECAMPSIANA Wattebled.³

* *Unio decampsianus* WATTEBLED, Jl. de Conch., XXXII, 1884, p. 132, pl. VII, fig. 1.—

*P.ETEL, Conch. Sam., III, 1890, p. 150.

* *Unio campsianus* P.ETEL, Conch. Sam., III, 1890, p. 147.

Western Soudan.

NODULARIA BELLAMYI Jousseau.

* *Pharaonia bellamyi* JOUSSEAU, Bull. Soc. Zool. Fr., XI, 1886, p. 486, pl. XII, figs. 7a, 7b.

Faraba, on the Niger; High Senegal.

NODULARIA RENEA Jousseau.

* *Renews renew* JOUSSEAU, Bull. Soc. Zool. Fr., XI, 1886, p. 482, pl. XII, figs. 4a, 4b.

* *Renews faidherbi* JOUSSEAU, Bull. Soc. Zool. Fr., XI, 1886, p. 483, pl. XII, figs. 5, 5a.

Niger at Kayon; Senegal River; High Senegal.

¹This appears from the figures and description to be a member of the *Aegyptiac* group of *Nodularia*. It is possible that it came from Asia Minor, but more likely that it is an African species. The *Unio eucyphus* of Bourguignat, which he credits to this region, is believed by Lea to be the *U. aegyptiacus* of Cailliaud, and I agree with him. *Unio bagdadensis* may be merely a peculiar form of the same thing.

²This may possibly be distinct from *N. nilotica*, or it may only be an oddly-formed specimen.

³I can not be positive as to the systematic position of this species, but it probably belongs here.

***NODULARIA FOULADOUGOUENSIS** Jousseau.

- * *Reneus fouladougouensis* JOUSSEAU, Bull. Soc. Zool. Fr., XI, 1886, p. 485, pl. XII, figs. 6, 6a.

Fouladougou, Niger; High Senegal.

† **NODULARIA GABONENSIS** Kuster.

- * *Unio gabonensis* KUSTER, Conch. Cab. Unio, 1862, p. 291, pl. xcvi, fig. 7.—
* PÆTEL, Conch. Sam., III, 1890, p. 153.
Margaron (*Unio*) *gabonensis* LEA, Syn., 1870, p. 47.

Gaboon River, West Africa.

† **NODULARIA ÆQUITORIA** Morelet.

- * *Unio æquitorius* MORELET, Jl. de Conch., XV, 1885, p. 31, pl. II, fig. 9.¹—* PÆTEL, Conch. Sam., III, 1890, p. 144.
* *Unio landanensis* SCHEPMAN, Notes Leyden Mus., VIII, 1891, p. 113, pl. VIII, figs. 3a, 3b.

Congo drainage.

NODULARIA HOREI E. A. Smith.

- * *Unio horei* E. A. SMITH, Ann. and Mag., VI, 1880, p. 429; * Pr. Zool. Soc. Lond., 1881, p. 299, pl. XXXIV, fig. 37.—* CROSSE, Jl. de Conch., XXXIX, 1881, p. 133.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 235.
* *Grandidieria horei* BOURGNIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 6.

Lake Tanganyika.

NODULARIA BORELLII Ancey.

- * *Unio borellii* ANCEY, Bull. Soc. Zool. Fr., VII, 1894, p. 226, fig. 2.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 227.

Lake Nyassa and vicinity.

NODULARIA EMINI von Martens.

- * *Unio emini* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 224, pl. VII, fig. 14.

Victoria Nyanza.

NODULARIA BOHMI von Martens.

- * *Unio bohmi* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 223, pl. VII, fig. 9.

Lake Tanganyika.

* **NODULARIA AMBIFARIA** von Martens.

- * *Unio ambifarius* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 225, pl. VII, fig. 20.

Dar-es-Ssalam, East Africa.

NODULARIA LECHAPTOISI Ancey.

- * *Unio lechaptosi* ANCEY, Bull. Soc. Zool. Fr., VII, 1894, p. 228, fig. 3.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 226.

Shire River, Lake Nyassa.

¹ Doubtfully distinct from *gabonensis*.

NODULARIA LIEDERI von Martens.

* *Unio liederii* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 226, pl. VII, fig. 19.

Lake Nyassa.

NODULARIA CALATHUS Bourgnignat.¹

* *Unio calathus* BOURGNIGNAT, Esp. Ouk. et Tan., 1885, p. 23.

Tanganyika.

NODULARIA BAKOYI Rochebrune.

* *Unio bakoyi* ROCHEBRUNE, Bull. Soc. Phil., 7th ser., VI, 1882, p. 33.

Bakoy, High Senegal.

NODULARIA RADIOTA Charmes.

* *Unio radiotus* CHARMES, Bull. Soc. Mal. Fr., II, 1885, p. 166.

Central Africa.

NODULARIA JULIANI Rang.

* *Unio juliani* RANG, Nouv. Ann. Mus., 1835, p. 309.

Senegal.

(Group of *Nodularia kunenensis*.)

Shell elliptical, subinflated, solid, narrowly biangulate behind, the point of the shell being about midway up the height, the post-base inflated, beaks full, eroded in the specimen figured but no doubt zigzag sculptured; whole surface of the shell covered with wavy corrugations; epidermis brownish; one rather solid, deeply incised pseudocardinal in the right valve, two in the left; nacre yellowish white; muscle scars deep. Animal unknown.

NODULARIA KUNENENSIS Mousson.

* *Unio kunenensis* MOUSSON, Jl. de Conch., XXXV, 1887, p. 300, pl. x.i, fig. 10.—

*SMITH, Ann. and Mag., VIII, 1891, p. 319.

Head of Kunene River, North Ovampo, Southwest Africa.

Section CAFFERIA Simpson, 1900.

(Type, *Unio caffer* Krauss.)

Shell elongated or elliptical, rhomboid and solid when old; beaks full, the sculpture corrugated zigzag, the ridges often extending over the disk; epidermis yellowish-brown to nearly black, dull colored, somewhat sulcate; teeth rather strong, muscle scars deep, well defined.

(Group of *Nodularia caffer*. Characters as in the section.)

¹ This and the following species are unfigured, and are referred by von Martens and others to this group. I know nothing of them. I am a little in doubt as to the affinities of some of the above-figured species which I have not seen, but they are placed here on the excellent authority of von Martens.

† NODULARIA CAFFER Krauss.

- * *Unio caffer* KRAUSS, Sud. Af. Moll., 1848, p. 18, pl. I, fig. 14.—* KUSTER (part), Conch. Cab. Unio, 1856, p. 143, pl. XLII, fig. 2.—* HANLEY, Biv. Shells, 1856, p. 385, pl. XXI, fig. 40.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLI, fig. 226.—* SMITH, Ann. and Mag., VIII, 1891, p. 317.—* PÆTEL, Conch. Sam., III, 1890, p. 147.
- * *Margaron (Unio) caffer* LEA, Syn., 1852, p. 32; 1870, p. 48.
- * ? *Unio schwerzenbachii* BOURGUIGNAT, Rev. et Mag., 1856, p. 71.—* SOWERBY, Conch. Icon., XVI, 1866, pl. XLIV, fig. 241.
- * † *Unio verreauxianus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 94; Obs., VI, 1857, p. 21, pl. XXVI, fig. 16; Jl. Ac. N. Sci. Phila., III, 1858, p. 301, pl. XXVII, fig. 16.—* SOWERBY, Conch. Icon., XVI, 1866, pl. LXIX, fig. 352.—* PÆTEL, Conch. Sam., III, 1890, p. 171.
- * *Margaron (Unio) verreauxianus* LEA, Syn., 1870, p. 36.
- * *Unio verreauxi* PÆTEL, Conch. Sam., III, 1890, p. 171.
- * *Unio navigoliformis* LEA, Pr. Ac. N. Sci. Phila., XI, 1859, p. 152; Jl. Ac. N. Sci. Phila., IV, 1860, p. 248, pl. XXXVII, fig. 124; * Obs., VII, 1860, p. 63, pl. XXXVII, fig. 124.—* REEVE, Conch. Icon., XVI, 1865, pl. XXIV, fig. 114.—* PÆTEL, C. Sam., III, 1890, p. 160.
- * *Margaron (Unio) navigoliformis* LEA, Syn., 1870, p. 31.
- * † *Unio natalensis* LEA, Pr. Ac. N. Sci. Phila., VIII, 1864, p. 113; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 59, pl. XX, fig. 57; * Obs., XI, 1867, p. 63, pl. XX, fig. 57.—* SOWERBY, Conch. Icon., XVI, 1866, pl. LXXI, fig. 362.—* PÆTEL, Conch. Sam., III, 1890, p. 160.
- * *Margaron (Unio) natalensis* LEA, Syn., 1870, p. 32.
- * *Unio rectilinearis* SOWERBY, Conch. Icon., XVI, 1866, pl. LXV, fig. 332.²

† NODULARIA CAFFER var. AFRICANA Lea.

- * *Unio africanus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 94; * Jl. Ac. N. Sci. Phila., III, 1857, p. 300, pl. XXVII, fig. 115; * Obs., VI, 1857, p. 20, pl. XXVII, fig. 15.—* REEVE, Conch. Icon., XVI, 1865, pl. XXII, fig. 100.
- * *Margaron (Unio) africanus* LEA, Syn., 1870, p. 48.
- * *Unio caffer* KUSTER (part), Conch. Cab. Unio, 1856, p. 143, pl. XLII, fig. 3.
- * *Unio niloticus* KUSTER, Conch. Cab. Unio, 1856, p. 158, pl. XLV, fig. 5.

† NODULARIA CAFFER var. VAALENSIS Chaper.

- * *Unio raalensis* CHAPER, Bull. Soc. Zool. Fr., X, 1885, p. 480, pl. XI, figs. 1-3.—* PÆTEL, Conch. Sam., III, 1890, p. 171.

South Africa.

NODULARIA TRAVERSII Pollonera.³

- * *Unio traversii* POLLONERA, Bull. Soc. Mal. It., XIII, 1888, p. 85, pl. III, figs. 14, 15.

Hawash River, Shoa, Africa.

¹According to Smith the above, *U. verreauxianus* Lea, *africanus* Lea, *natalensis* Lea, and *raalensis* Chaper, are all one species. I think he is right, but it seems to me that one or two of these forms are worthy of varietal names. I have seen the types of most and authentic specimens of all of them.

²Said to come from the Columbia River, but it did not.

³Possibly a variety of *N. caffer*.

NODULARIA JICKELI Simpson.

* ? *Unio teretiusculus* JICKELI,¹ Faun. Moll. N. O. Af., 1874, p. 276, pl. XI, figs. 3, 3a, 3b.

Northeast Africa.

NODULARIA ACUMINATA H. Adams.

* *Unio acuminatus* H. ADAMS, Pr. Zool. Soc. Lond., 1866, p. 376.—* PÆTEL, Conch. Sam., III, 1890, p. 143.—* SMITH, Ann. and Mag., X, 1892, p. 127, pl. XII, fig. 12.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 227, pl. VII, figs. 11, 12.

Lake Albert Nyanza.

NODULARIA DEMBEÆ Reeve.

* *Unio dembeæ* REEVE, Conch. Icon., XVI, 1865, pl. XXIX, fig. 153.—* JICKELI, Faun. Moll. Af., 1874, p. 275, pl. IX, figs. 3-4.—* ROSSMASSLER, Bull. Soc. Mal. It., XIII, 1888, p. 84.—* PÆTEL, Conch. Sam., III, 1890, p. 150.

Abyssinia.

NODULARIA LOURDELLI Bourguignat.

* *Unio lourdelli* BOURGUIGNAT, Bull. Soc. Mal. Fr., IV, 1887, p. 271.—* SMITH, Ann. and Mag., X, 1892, p. 128, pl. XII, figs. 13-15.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 228.

Victoria Nyanza Lake.

NODULARIA ABYSSINICA von Martens.

* *Unio abyssinicus* VON MARTENS, Mal. Bl., XIII, 1866, p. 102; * XIV, 1867, p. 17.—* JICKELI, Faun. Moll. N. O. Af., 1874, p. 278, pl. IX, fig. 5; X, fig. 10.—* PÆTEL, Conch. Sam., III, 1890, p. 143.

Abyssinia.

NODULARIA DIMINUTA Lea.

* *Unio diminutis* LEA,² Pr. Ac. N. Sci. Phila., III, 1859 p. 151; * JI. Ac. N. Sci. Phila., 1860, p. 254, pl. XXXIX, fig. 134; * Obs., VII, 1860, p. 72, pl. XXXIX, fig. 134.—* REEVE, Conch. Icon., XVI, 1865, pl. XXVIII, fig. 141.—* JICKELI, Faun. Moll. Af., 1874, p. 280.—* PÆTEL, Conch. Sam., III, 1890, p. 150.
* *Margaron (Unio) diminutus* LEA, Syn., 1870, p. 31.
* *Unio gratosus* PÆTEL, Conch. Sam., III, 1890, p. 154.

East Africa.

The following unfigured species are said to belong here:

* *Unio mandinguorum* ROCHEBRUNE, Bull. Soc. Phil., 7th ser., VI, 1882, p. 34.

Bakoy, High Senegal.

* *Unio zeyheri* MENKE, Zeits. für Mal., VI, 1848, p. 28.

South Africa.

¹ Not *teretiusculus* Philippi, but a larger elongate solid rhomboid form, which is certainly distinct. I therefore name it in honor of Jickeli.

² So in the first published description, probably a typographical error. I am not certain as to the relationships of the species, but presume that it belongs here.

(Group of *Nodularia fissidens*.)

Shell solid, compressed, subelliptical to subrhomboid, with rather full beaks, which have radiate, corrugated sculpture extending out on the disk as wavy sulcations, and changing near the edge to plain sulcations; posterior ridge low, rounded; hinge strong; right valve with two pseudocardinals, the lower the larger, showing a tendency to split up, and having a gape in the plate behind, and having one lateral; two pseudocardinals in the left, somewhat split, and two laterals; muscle scars deep, distinct.

NODULARIA FISSIDENS Böttger.

* *Unio* (*Hyridella*) *fissidens* BÖTTGER, Ber. Senck. Ges., 1886, p. 27, pl. II, figs. 6ab, 7ab.

Kalahari Desert. Southwest Africa.

NODULARIA HYGAPANUS Böttger.

* *Unio* (*Hyridella*) *hygapanus* BÖTTGER, Ber. Senck. Ges., 1886, p. 26, pl. II, fig. 6.

Kalahari Desert. I am not certain as to the relationships of these two species.

Genus HARMANDIA Rochebrune, 1882.

(Type, *Harmandia somboriensis* Rochebrune.)

Harmandia ROCHEBRUNE, Bull. Soc. Phil., VI, 1882, p. 45.

Shell subtrapezoidal, subalate, rather thin, posteriorly compressed, with low beaks and irregularly radial sculpture, a few of the ribs converging at the center of the disk, the rest becoming divaricate and covering the shell, those of the posterior slope springing from the rib on the low, posterior ridge; epidermis gray green; left valve with two arcuate, compressed, elongate pseudocardinals in front and a triangular cardinal under the beak, the right with two pseudocardinals in front and a pit under the beak; there are two nearly straight lamellar laterals in each valve, and a small, up-curved lamella above them near their posterior ends; anterior muscle scars deep; nacre white, brilliant, showing the outside sculpture.

Animal unknown.

HARMANDIA SOMBORIENSIS Rochebrune.

* *Harmandia somboriensis* ROCHEBRUNE, Bull. Soc. Phil., VI, 1882, p. 46, pl. I, figs. 1-3..

Rapids of Sombor sombor, Mekong.

Genus GRANDIDIERIA Bourguignat, 1885.

(Type, *Unio burtoni* Woodward.)

Grandidieria BOURGUIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 6.

Shell small, oval, rounded or rhomboid, solid, much inflated, generally narrowly biangulate behind, often apparently of two forms, one more

inflated in the basal and post-basal parts than the other; beaks high, and pointed, curved forward and inward, very delicately and beautifully sculptured with zigzag liræ, which become finely nodulous and sulcate on the disk, especially in front and behind; posterior ridge well developed, often slightly double; hinge line curved; two pseudocardinals in the right valve separated by a parallel-sided socket, one or two in the left, with often an irregular, triangular, ragged tooth under the beak which is frequently reflexed; one obliquely striate lateral in the right valve and two in the left; nacre of peculiarly soft, rich texture, white, coppery, or purplish, delicately radiate; beak cavities moderate; muscle scars distinct.¹

† **GRANDIDIERIA BURTONI** Woodward.

* *Unio burtoni* WOODWARD, Proc. Zool. Soc. Lond., 1859, p. 348, pl. XLVII, fig. 2;

* Ann. and Mag., V, 1860, p. 338.—* SOWERBY, Conch. Icon., XVI, 1866, pl.

XLVII, fig. 251.—* SMITH, Proc. Zool. Soc. Lond., 1881, p. 297, pl. XXXIV, figs.

33, 33b.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 237.

* *Margaron (Unio) burtoni* LEA, Syn., 1870, p. 31.

* *Grandidieria burtoni* BOURGUIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 6.

* *Grandidieria cyrenopsis* BOURGUIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 9, pl. I, figs. 7-9; * Icon. Mal. Tan., 1888, pl. XIX, figs. 1-3.

GRANDIDIERIA BURTONI var. **SERVAINIANA** Bourguignat.

* *Grandidieria servainiana* BOURGUIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 6.

* *Unio burtoni* var. *servainiana* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 238.

GRANDIDIERIA BURTONI var. **INSIGNIS** Bourguignat.

* *Grandidieria insignis* BOURGUIGNAT, Esp. Ouk. Tan., 1885, p. 16.

* *Unio burtoni* var. *insignis* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 238.

GRANDIDIERIA BURTONI var. **STURANYI** von Martens.

Grandidieria STURANY, Baumann, Durch Massailand, 1894, p. 6, pl. XXIV, fig. 31; XXV, fig. 35.

Unio burtoni var. *sturanyi* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 238.

Lake Tanganyika.

† **GRANDIDIERIA TANGANYICENSIS** Smith.

* *Unio tanganyicensis* SMITH, Proc. Zool. Soc. Lond., 1880, p. 351, pl. XXXI, p. 9, 9a;

* Proc. Zool. Soc. Lond., 1881, p. 298, pl. XXXIV, fig. 35.—* PETEL, Conch. Sam., III, 1890, p. 169.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 240.

* *Grandidieria tanganikana* BOURGUIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 7.

Lake Tanganyika.

¹ Bourguignat, by a most brilliant stroke, placed this group in the *Corbiculidæ*, and stated that he knew twenty-five species, and had no doubt that there would be at least one hundred when a thorough search is made. The group is an exceedingly difficult and variable one, and, judging from the large number of species he has founded on the most trivial variations, we may consider it exceedingly fortunate that he never saw the other seventy-five. I have seen but a limited amount of material, but I am led to believe that two forms of the same species often exist, as in *Lampisilis* and allied genera, and which may be male and female shells. I know nothing of the animal and I am unable to do justice to the genus.

† *GRANDIDIERIA SMITHI* Bourguignat.

* *Unio burtoni* SMITH (part), Proc. Zool. Soc., 1881, pl. XXXIV, fig. 33a.

* *Grandidieria smithi* BOURGUIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 7.

* *Unio burtoni* var. *smithi* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 238.

Lake Tanganyika.

GRANDIDIERIA GRAVIDA Bourguignat.

* *Grandidieria gravida* BOURGUIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 7, pl. I, figs. 1-6; * Icon. Moll. Tan., 1888, p. 41, pl. XVIII, figs. 11-16.

* *Grandidieria rostrata* BOURGUIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 10, pl. I, figs. 10-12; * Icon. Moll. Tan., 1888, p. 41, pl. XVIII, figs. 17-19.

* *Unio rostralis* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 238.

Lake Tanganyika.

† *GRANDIDIERIA THOMSONI* Smith.

* *Unio thomsoni* SMITH, Ann. and Mag., VI, 1880, p. 430; * Proc. Zool. Soc. Lond., 1881, p. 299, pl. XXXIV, fig. 36.—* CROSSE, Jl. de Conch., XXIV, 1881, p. 133.—

* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 240.

* *Grandidieria thomsoni* BOURGUIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 7; * Moll. Gir. Tan., 1885, p. 100.

* *Grandidieria corbicula* BOURGUIGNAT, Not. Prod., 1885, p. 100; * Moll. Gir. Tan., 1885, p. 100; * Icon. Moll. Tan., 1888, p. 43, pl. XIX, figs. 10-12.

* *Grandidieria granulosa* BOURGUIGNAT, Not. Prod., 1885, p. 102; * Icon. Moll. Tan., 1888, p. 43, pl. XIX, figs. 1, 2.

* *Grandidieria singularis* BOURGUIGNAT, Icon. Moll. Tan., 1888, p. 43, pl. XIX, figs. 18-20

Lake Tanganyika.

GRANDIDIERIA RHYNCHONELLA Bourguignat.

* *Grandidieria rhynchonella* BOURGUIGNAT, Icon. Moll. Tan., 1888, p. 43, pl. XIX, figs. 16, 17.

Lake Tanganyika.

GRANDIDIERIA CALLISTA Bourguignat.

* *Grandidieria callista* BOURGUIGNAT, Icon. Moll. Tan., 1888, p. 43, pl. XIX, figs. 13-15.¹

Lake Tanganyika.

GRANDIDIERIA BOURGUIGNATI Joubert.

* *Grandidieria bourguignati* JOUBERT in Bourguignat, Icon. Moll. Tan., 1888, p. 43, pl. XIX, figs. 7-9.

Lake Tanganyika.

GRANDIDIERIA ANCEYI Bourguignat.

* *Unio burtoni* SMITH, (part) Proc. Zool. Soc. Lond., 1881, p. 297, pl. XXXIV, fig. 33a.

* *Grandidieria anceyi* BOURGUIGNAT, Esp. Ouk. et Tan., 1885, p. 15; Moll. Tan., 1888, p. 43, pl. XIX, figs. 4-6.²

Lake Tanganyika.

¹ Probably a strangely distorted form of some other species.

² I am satisfied that Smith's figure 33a is not *burtoni*, but a valid species—*G. anceyi* Bourguignat.

The following are unfigured species of *Grandidieria*:

- G. rotundata* Bourguignat, Not. Prod., 1885, p. 98.
G. mira Bourguignat, Not. Prod., 1885, p. 96.
G. locardiana Bourguignat, Esp. Ouk., 1885, p. 18.
G. incarnata Bourguignat, Moll. Gir., 1885, p. 101.
G. giraudi Bourguignat, Not. Prod., 1885, p. 95.
G. elongata Bourguignat, Moll. Gir., 1885, p. 14.

Genus *PHYSUNIO*, Simpson, 1900.

(Type, *Unio gravidus* Lea.)

Shell thin, irregularly obovate, narrowed in front, decidedly produced at post base, pointed behind and posteriorly winged, with a moderate posterior ridge and often a second or third faint ridge above it; beak sculpture zigzag radial, somewhat disposed in two sets, the one down the posterior ridge slightly nodulous; posterior slope having irregular radial corrugations, the rest of the shell smooth; epidermis often cloth-like, with one or more green rays on the posterior slope; hinge line curved; a single obliquely granularly striate pseudocardinal and generally three laterals in the left valve, and two pseudocardinals and two laterals in the right, all greatly compressed; beak cavities deep; muscle scars irregular; nacre bright, bluish, and iridescent.

Animal unknown.

Section *PHYSUNIO* Simpson, 1900.

(Type, *Unio gravidus* Lea.)

Shell inflated; beak cavities deep and rounded.

† *PHYSUNIO GRAVIDUS* Lea.

- * *Unio gravidus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 93; Obs., VI, 1857, p. 12, pl. XXIV, fig. 5; Jl. Ac. N. Sci. Phila., III, 1858, p. 292, pl. XXIV, fig. 5.—
 * VON MARTENS, Pr. Zool. Soc. Lond., 1860, p. 14.—* SOWERBY, Conch. Icon., XVI, 1866, pl. LII, fig. 271.—* MORELET, Ser. Conch., IV, 1875, p. 347.—
 * PÆTEL, Conch. Sam., III, 1890, p. 154.
 —* Margaron (*Unio gravidus* LEA, Syn., 1870, p. 28.
 * *Lampsilis gravidus* ROCHEBRUNE, Bull. Soc. Phil., VI, 1882, p. 43.
 * *Unio abnormis* MORELET, Rev. et Mag., XIV, 1862, p. 480.
 * *Unio superbus* SOWERBY, Conch. Icon., XVI, 1867, p. LIX, fig. 295.

Siam; Cambodia; Cochin China.

† *PHYSUNIO SUPERBUS* Lea.

- * *Unio superbus* LEA, Desc. 12 New Uniones, 1843 (no pagination); Tr. Am. Phil. Soc., IX, 1845, p. 281, pl. XLII, fig. 11; Obs., IV, 1848, p. 39, pl. XLII, fig. 11.—
 H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* VON MARTENS, Mal. Bl., XIV, 1866, p. 14.—* PÆTEL, Conch. Sam., III, 1890, p. 169.
 * Margaron (*Unio superbus* LEA, Syn., 1852, p. 19; 1870, p. 28.
 * *Unio velaris* HANLEY, Biv. Shells, 1856, p. 385, pl. XXIII, fig. 42.¹
 * *Unio massini* MORELET, Jl. de Conch., XII, 1864, p. 288; Ser. Conch., IV, 1875, p. 348, pl. XV, figs. 1, 3.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

Sumatra; Cochin China.

¹ Credited to Benson manuscript.

PHYSUNIO CROSSEI Deshayes and Julien.

- * *Unio crossei* DESHAYES and JULIEN, N. Arch. Mus., X, 1874, p. 124, pl. VI, figs. 5-7.—* PÆTEL, Conch. Sam., III, 1890, p. 149.

Cambodia; Cochin China.

† PHYSUNIO MICROPTERUS Morelet.

- * *Unio micropterus* MORELET, Jl. de Conch., XIV, 1866, p. 63; Ser. Conch., IV, 1873, p. 349, pl. XV, fig. 7.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

Siam; Cambodia.

PHYSUNIO SEMIALATUS Deshayes and Julien.

- * *Unio semialatus* DESHAYES and JULIEN, N. Arch. Mus., X, 1874, p. 123, pl. VI, figs. 1, 2.¹

Siam; Cambodia.

† PHYSUNIO CAMBODIENSIS Lea.²

- * *Unio cambodiensis* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 103; Jl. Acad. N. Sci. Phil., III, 1857, p. 313, pl. XXX, fig. 28; Obs., VI, 1857, p. 33, pl. XXX, fig. 28.

- * *Margaron (Unio) cambodiensis* LEA, Syn., 1870, p. 42.

Takrong River at Korat, Cambodia:

Section LENS Simpson, 1900.

(Type, *Unio eximius* Lea.)

Shell sublenticular; three laterals of left valve distinct; cavity of the beaks compressed.

† PHYSUNIO EXIMIUS Lea.

- * *Unio eximius* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 93; Obs., VI, 1857, p. 14, pl. XXV, fig. 8; Jl. Ac. N. Sci. Phila., III, 1858, p. 294, pl. XXV, fig. 8.—

- * VON MARTENS, Pr. Zool. Soc. Lond., 1860, p. 15.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIX, fig. 481.—* PÆTEL, Conch. Sam., III, 1890, p. 152.

- * *Margaron (Unio) eximius* LEA, Syn., 1870, p. 28.

- Unio semiquadrata* SOWERBY, Conch. Icon., XVI, 1866, pl. XLVIII, fig. 258.

- * *Unio semiquadratus* PÆTEL, Conch. Sam., III, 1890, p. 167.

Siam; Cambodia.

PHYSUNIO VELARIS Sowerby.

- * *Unio velaris* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXII, fig. 368.³—* PÆTEL, Conch. Sam., III, 1890, p. 171.

Assam.

¹Fischer believes that the above equals *U. micropterus* Morelet (Cat., p. 142). I feel sure he is wrong, and that this is a valid species.

²I have only seen the type of this, a very young shell, but it probably belongs to the *gravidus* group.

³A compressed shell of a uniform yellow color, probably only a variety of *eximius*.

Genus DALLIELLA Simpson 1900.

(Type *Anodonta purpurea* Valenciennes.¹)

Shell subtrapezoidal, generally thin, inflated, with a low, rounded posterior ridge and rather full beaks which have somewhat zigzagged radial sculpture, with a row of chevron-shaped folds running some distance down the posterior ridge, and with fine corrugations in front of the beaks; epidermis somewhat cloth-like; substance of the shell of a peculiar purplish-brown tint; hinge line narrow; teeth imperfectly developed, sometimes reduced to mere rudiments, when present consisting of greatly compressed, feeble, pseudocardinals and laterals; beak cavities rather shallow; dorsal scars one to a few, scattered; muscle scars faint; prismatic border wide.

Animal unknown.

(Group of *Dallielia purpurea*.)

Shell not biangulate behind, rather smooth; nacre coppery purple; teeth very faint, compressed.

† DALLIELLA PURPUREA Valenciennes.

* *Anodonta purpurea* VALENCIENNES, Rec. Obs. Zool., II, 1833, p. 236, pl. XLVIII bis, figs. 3, 3a, 3b.—* HANLEY, Test. Moll., 1842, p. 218.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* VON MARTENS, Mal. Bl., XIV, 1866, p. 12.—* CLESSIN, Conch. Cab. Ano., 1876, p. 77, pl. XIX, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 183.

* *Margarita (Anodonta) purpurea* LEA, Syn., 1836, p. 51; 1838, p. 30.

* *Anodon purpurea* CATLOW and REEVE, Conch. Nom., 1845, p. 67.

* *Margaron (Anodonta) purpurea* LEA, Syn., 1852, p. 50; 1870, p. 79.

* † *Anodonta burroughiana*, LEA, Tr. Am. Phil. Soc., V, 1834, p. 105, pl. XVI, fig. 49; * Obs., I, 1834, p. 217, pl. XVI, fig. 49.—* CLESSIN, Conch. Cab. Ano., 1875, p. 164, pl. LIV, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 177.

* *Margaron (Anodonta) burroughana*, LEA, Syn., 1870, p. 81.

* *Anodon burroughianus*, SOWERBY, Conch. Icon., XVII, 1870, pl. xxvii, fig. 103.

* † *Margarita (Unio) bengalensis* LEA, Syn., 1836, p. 26; 1838, p. 20.

* *Unio bengalensis* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 3, pl. II, fig. 3; * Obs., II, 1838, p. 3, pl. II, fig. 3.—* TROSCHER, Arch. für Nat., V, 1839, pl. II, p. 233.—* HANLEY, Test. Moll., 1842, p. 194; * Biv. Shells, 1843, p. 194, pl. XXI, fig. 50.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* CHENU, Ill. Conch., 1858, pl. XXII, figs. 2, 2a, 2b.—* KUSTER, Conch. Cab. Unio, 1861, p. 228, pl. LXXVII, figs. 2, 3.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

* *Margaron (Unio) bengalensis* LEA, Syn., 1852, p. 30; 1870, p. 47.

* *Anodon bengalensis* SOWERBY, Conch. Icon., XVII, 1867, pl. XIV, fig. 49.

* *Anodonta bengalensis* CLESSIN, Conch. Cab. Ano., 1876, p. 173, pl. LVII, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 177.

* † *Unio vercundus* GOULD, Pr. Bost. S. N. Hist., III, 1850, p. 295; * U. S. Expl. Exp., XII, 1852, p. 431, figs. 541a, 541b, 541c; * Otia Conch., 1862, p. 89.—

¹ Some of the species placed here have been classed with *Anodonta*, but almost any large series of specimens of any species will show variation from edentulous to toothed forms.

*REEVE, Conch. Icon., XVI, 1865, pl. xxv, fig. 125.—*PÆTEL, Conch. Sam., III, 1890, p. 171.

**Unio mauritianus* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 152; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 257, pl. XL, fig. 138; *Obs., VII, 1860, p. 75, pl. XL, fig. 138.

**Margaron (Unio) mauritianus* LEA, Syn., 1870, p. 42.

*? *Anodonta heldii* KUSTER, Conch. Cab. Ano., 1873, p. 64, pl. XIX, fig. 1.

Anodonta chinensis KUSTER? Where?

Philippines. The localities India and Mauritius given for this species are no doubt erroneous.

† DALLIELLA CREPERA Lea.

**Anodonta crepera* LEA, Pr. Zool. Soc. Lond., 1850, p. 198; *Ann. and Mag., VIII, 1851, p. 494.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—*LEA, Jl. Ac. N. Sci. Phila., IV, 1860, p. 238, pl. LXXXIV, fig. 117; *Obs., VII, 1860, p. 56, pl. XXXIV, fig. 117.—*CLESSIN, Conch. Cab. Ano., 1873, p. 97, pl. XXIX, figs. 5, 6.—*PÆTEL, Conch. Sam., III, 1890, p. 178.

*? *Anodon creperus* SOWERBY, Conch. Icon., XVII, No. 16.

Margaron (Anodonta) crepera LEA, Syn., 1852, p. 50; 1870, p. 80.

Luzon Island, Philippines.

† DALLIELLA TENUIS Lea.

**Anodonta tenuis* LEA, Pr. Zool. Soc. Lond., 1850, p. 198; *Ann. and Mag., VIII, 1851, p. 494; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 237, pl. XXXIII, fig. 116; *Obs., VII, 1860, p. 55, pl. XXXIII, fig. 116.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—*MUSGRAVE, Phot. Conch., 1863, pl. I, fig. 8.—*CLESSIN, Conch. Cab. Ano., 1873, p. 99, pl. XXX, figs. 3, 4.—*PÆTEL, Conch. Sam., III, 1890, p. 185.

**Margaron (Anodonta) tenuis* LEA, Syn., 1852, p. 53; 1870, p. 83.

**Anodon tenuis* SOWERBY, Conch. Icon., XVI, 1867, pl. xv, fig. 55.

Luzon Island, Philippines.

† DALLIELLA SUBCRASSA Lea.

**Anodonta subcrassa* LEA, Pr. Zool. Soc. Lond., 1850, p. 198; *Ann. and Mag., VIII, 1851, p. 495; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 236, pl. XXXIII, fig. 115; *Obs., VII, 1860, p. 54, pl. XXXIII, fig. 115.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—*MUSGRAVE, Phot. Conch., 1863, pl. I, fig. 3.—*CLESSIN, Conch. Cab. Ano., 1873, p. 98, pl. XXXI, figs. 1, 2.—*PÆTEL, Conch. Sam., III, 1890, p. 185.

**Margaron (Anodonta) subcrassa* LEA, Syn., 1852, p. 51; 1870, p. 81.

**Anodon subcrassa* SOWERBY, Conch. Icon., XVII, 1867, pl. XIII, fig. 42.

Luzon.

† DALLIELLA GRACILIS Lea.

**Anodonta gracilis* LEA, Pr. Zool. Soc. Lond., 1850, p. 197; *Ann. and Mag., VIII, 1851, p. 193; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 239, pl. XXXIV, fig. 119; *Obs., VII, 1860, p. 57, pl. XXXIV, fig. 119.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—*CLESSIN, Conch. Cab. Ano., 1873, p. 99, pl. XXIX, figs. 3, 4.—*PÆTEL, Conch. Sam., III, 1890, p. 180.

**Margaron (Anodonta) gracilis* LEA, Syn., 1852, p. 51; 1870, p. 82.

Dingle, Isle of Panay, Philippines.

DALLIELLA SERRADELLI Drouet.

* *Unio serradelli* DROUET, Jl. de Conch., XL, 1892, p. 86.

Philippines.

(Group of *Dallielia insularis*.)

Shell subsolid, subrhomboidal biangulate behind, pseudocardinals and laterals blurred, not greatly compressed; nacre lurid, tinged purplish.

† DALLIELLA INSULARIS Drouet.

* *Pseudodon insularis* DROUET, Rev. Biol. Fr., VI, 1894, p. 217, fig. 2.

Borneo; Palawan Island, Philippines.

† DALLIELLA CRASSA Drouet.

* *Pseudodon crassus* DROUET, Jl. de Conch., XL, 1892, p. 93.—*DROUET and CHAPER, Mém. Soc. Zool. de Fr., V, 1892, p. 151, pl. vi, figs. 1-3.

Borneo.

The following is an unfigured species.

* *Unio semmelinki* VON MARTENS, Sitzber. Ges. Nat. Fr. Ber., 1891, p. 111.

Southeast Borneo. Possibly a *Dallielia*.

Genus PSEUDODON Gould, 1844.

(Type, *Anodon salwiniana* Gould.)

Pseudodon GOULD, Pr. Bost. Soc. Nat. His., 1844, p. 161.

Monodontina CONRAD, Pr. Ac. N. Sci. Phila., 1853, p. 269.

Trigonodon CONRAD, Am. Jl. Couch., I, 1865, p. 233.

Monocondylus MORELET, Rev. et Mag., 1866, p. 167.

Pseudodus DE MORGAN, Bull. Soc. Zool. Fr., X, 1885, p. 422.

Shell oval to elongate, having two posterior ridges and often one or two slight, radiating elevations above on the posterior slope, above which there is usually a small wing; beaks rather low, sculpture unknown;¹ epidermis dark, often having a few faint wrinkles on the posterior slope, without rays when adult. There is a single, smooth tooth in each valve, with occasional vestiges of a second; laterals nearly or quite wanting; beak cavities shallow; dorsal scars conspicuous, deep, distinct, 4 to 7 in a row under the hinge; anterior scars irregular; nacre generally dull and lurid.

Animal² having the branchiæ wide and rounded behind, becoming

¹One or two authors in describing species say that the beaks are entirely smooth. It is probable that in most cases they have zigzag radial sculpture. Although I have examined a large number of shells belonging to the group, many of which were young, they were all so badly eroded that the beak sculpture was entirely destroyed.

²Deshayes and Julien figure but do not describe the animal of *P. moreleti* in Nouv. Arch. de Mus., X, pl. v. I am only able to make out part of the characters from this really excellent figure.

narrow in front; palpi enormously long, apparently slender, pointed behind, where they project free for some distance; mantle thin, with a wide, slightly thickened border, faintly papillose behind, there seeming to be but little distinction between anal and branchial openings; anal opening apparently smooth.

Section SUBORBICULUS Simpson, 1900.

(Type, *Monocondylus orbicularis* Morelet.)

Shell compressed, nearly round in outline, feebly biangulate, and having a posterior dorsal wing; beaks low; hinge with a single, compressed, smooth, slightly curved, triangular tooth in each valve.

PSEUDODON ORBICULARIS Morelet.

* *Monocondylus orbicularis* MORELET, Rev. et Mag., XVIII, 1866, p. 167.

* *Pseudodon orbicularis* MORELET, Ser. Conch., IV, 1875, p. 338, pl. XVI, fig. 5.

* *Monocondylaea orbicularis* PÆTEL, Conch. Sam., III, 1890, p. 174.

Siam; Cambodia.

Section TRIGONODON Conrad, 1865.

Shell solid, compressed, rhomboid, round in front, widely and faintly biangulate and rough behind; surface irregularly concentrically wavy sulcate; hinge with a strong triangular pseudocardinal in the right valve, having a decided excavation behind it, and at the posterior side of the pit a faint tooth; there is one strong triangular tooth in the left valve and a vestige of one in front of it at the upper edge of the shell; anterior scars irregular; dorsal scars crowded close to the hinge line.

(Type, *Monocondylaea crebristriata* Anthony.)

† PSEUDODON CREBRISTRIATUS Anthony.

* *Monocondylaea crebristriata* ANTHONY, Am. Jl. Conch., I, 1865, p. 205, pl. XVIII, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 174.

* *Trigonodon crebristriata* CONRAD, Am. Jl. Conch., I, 1865, p. 233.

* *Margaron (Monocondylaea) crebristriata* LEA, Syn., 1870, p. 72.

* *Unio crebristriatus* SOWERBY, Conch. Icon., XVI, 1868, pl. xcv, fig. 517.

* *Pseudodon crebristriatum* HANLEY and THEOBALD, Conch. Ind., 1876, p. 5, pl. IX, fig. 3.

* *Unio vondembuschii* SOWERBY, Conch. Icon., XVI, 1868, pl. xcv, fig. 518.

† PSEUDODON CREBRISTRIATUS var. PEGUENSIS Anthony.¹

* *Monocondylaea peguensis* ANTHONY, Am. Jl. Conch., I, 1865, p. 205, pl. XVIII, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 174.

* *Margaron (Monocondylaea) peguensis* LEA, Syn., 1870, p. 73.

* *Pseudodon crebristriatum* var. *peguensis* HANLEY and THEOBALD, Conch. Ind., 1876, p. 5, pl. IX, fig. 5.

Burma; Pegu; Cambodia?.

¹Typically quite different from *P. crebristriatus*, but our large series seems to show an almost complete connection.

Section PSEUDODON Gould, 1844.

(Type, *Anodon Salrimana* Gould.)

Characters given under generic name.

(Group of *Pseudodon harmandi*.)

Shell rather small, solid, rhomboid, short, inflated; posterior ridge high, only slightly biangulate, faintly corrugated, on the posterior slope.

† PSEUDODON HARMANDI Crosse and Fischer.

* *Pseudodon harmandi* CROSSE and FISCHER, JI. de Conch., XXIV, 1876, p. 331, pl. x, fig. 2.

* *Monocondylæa harmandi* PÆTEL, Conch. Sam., III, 1890, p. 174.

Cambodia.

PSEUDODON AUREUS Heude.

* *Pseudodon aureus* HEUDE, Conch. Flav. Nank., IX, 1885, pl. LXXII, fig. 140.

* *Margaritana aurea* PÆTEL, Conch. Sam., III, 1890, p. 172.

China.

(Group of *Pseudodon rondembuschianus*.)

Shell rather large, somewhat compressed, subsolid, elliptic-rhomboid, wider behind, where it is widely biangulate; teeth compressed.

† PSEUDODON VONDEMBUSCHIANUS Lea.

* *Margaritana rondembuschiana* LEA, Pr. Am. Phil. Soc., I, 1840, p. 288; * Tr. Am. Phil. Soc., VIII, 1842, p. 222, pl. xviii, fig. 39; * Obs., III, 1842, p. 60, pl. xviii, fig. 39.—* KUSTER, Conch. Cab. Unio, 1862, p. 295, pl. xcvi, fig. 3.

* *Margaron (Monocondylæa) rondembuschiana* LEA, Syn., 1852, p. 45; 1870, p. 73.

* *Monocondylæa rondembuschiana* H. and A. ADAMS, Gen. Rec. Moll., II, 1858, p. 501.—* PÆTEL, Conch. Sam., III, 1890, p. 175.

* *Pseudodon rondembuschiana* CONRAD, Am. JI. Conch., I, 1865, p. 233.

* *Monodontina buschiana* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, pp. 269, 449.

* *Unio von buschea* SOWERBY, Conch. Icon., XVI, 1866, pl. LI, fig. 269.

* *Alasmodonta crispata* MOUSSON, L. and Suss. W. Moll. Java, 1849, p. 97, pl. xviii, figs. 1, 2.

* *Margaritana crispata* PÆTEL, Conch. Sam., III, 1890, p. 173.

† * *Monocondylæa planulata* LEA, Pr. Ac. N. Sci. Phila., XI, 1859, p. 187; * JI. Ac. N. Sci. Phila., IV, 1859, p. 262, pl. xlii, fig. 142; * Obs., VII, 1860, p. 80, pl. xlii, fig. 142.—* ? KUSTER, Conch. Cab. Unio, 1862, p. 305, pl. c, fig. 8.

* *Pseudodon planulata* CONRAD, Am. JI. Conch., I, 1865, p. 233.

* *Microcondylæa planulata* PÆTEL, Conch. Sam., III, 1890, p. 175.

* *Margaron (Monocondylæa) planulata* LEA, Syn., 1870, p. 73.

* *Margaritana fragilis* KUSTER, Conch. Cab. Unio, 1862, p. 295, pl. xcvi, fig. 2.

* *Microcondylæa fragilis* PÆTEL, Conch. Sam., III, 1890, p. 175.

* *Monocondylæa rhombica* KUSTER, Conch. Cab. Unio, 1862, p. 304, pl. c, fig. 7.

Java; Sumatra; Borneo; Malacca.

† PSEUDODON ELLIPTICUM Conrad.

* *Pseudodon ellipticum* CONRAD, Am. JI. Conch., I, 1865, p. 352, pl. xxv, fig. 1.

Cambodia.

† PSEUDODON ZOLLINGERI Mousson.

* *Alasmodonta zollingeri* MOUSSON, L. and Suss. Moll. Java, 1849, p. 96, pl. XVIII, fig. 1.

* *Margaritana zollingeri* KUSTER, Conch. Cab. Unio, 1862, p. 294, pl. XCVIII, fig. 1.

* *Pseudodon zollingeri* MORELET, Ser. Conch., IV, 1875, p. 337.

* *Monocondylaea zollingeri* PÆTEL, Conch. Sam., III, 1890, p. 174.

Java.

† PSEUDODON INOSULARIS Gould.

* *Anodon inosularis* GOULD, Pr. Bost. Soc. N. H., I, 1844, p. 160.¹—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 5, pl. IX, fig. 2.

* *Margaron (Monocondylaea) inosularis* LEA, Syn., 1870, p. 73.

* *Margaritana inosularis* PÆTEL, Conch. Sam., III, 1890, p. 173.

River Salwen, British Burma.

† PSEUDODON CUMINGII Lea.

* *Anodonta cumingii* LEA, Pr. Zool. Soc. Lond., 1850, p. 199; *Ann. and Mag., VIII, 1851, p. 495.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—*MUSGRAVE, Phot. Conch., 1863, pl. I, fig. 6.—*PÆTEL, Conch. Sam., III, 1890, p. 178.

* *Margaron (Monocondylaea) cumingii* LEA, Syn., 1852, p. 50; 1870, p. 73.

* *Monocondylaea cumingii* LEA, Jl. Ac. N. Sci. Phila., IV, 1860, p. 235, pl. XXXIII, fig. 114; *Obs., VII, 1860, p. 53, pl. XXXIII, fig. 114.

* *Pseudodon cumingii* CONRAD, Am. Jl. Conch., I, 1865, p. 233.

* *Anodon cumingii* REEVE, Conch. Icon., XVII, 1870, pl. XXXI, fig. 122.

* *Microcondylaea cumingii* CLESSIN, Conch. Cab. Ano., 1876, p. 259, pl. LXXXIII, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 175.

* *Pseudodus cumingii* DE MORGAN, Bull. Soc. Zool. de Fr., X, 1885, p. 422.

Malacca; Perak; Siam; Cambodia.

† PSEUDODON ÆNEOLUS Drouet.

* *Pseudodon æneolus* DROUET, Jl. de Conch., XL, 1892, p. 94.—*DROUET and CHAPER, Mem. Soc. Zool. de Fr., V, 1892, p. 152, pl. VI, figs. 4-7.

Borneo.

† PSEUDODON TUMIDUS Morelet.

* *Monocondylus tumidus* MORELET, Jl. de Conch., XIV, 1866, p. 62.—*MABILLE, Rev. Zool., XXIII, 1872, p. 51, pl. V, figs. 6, 7.

* *Pseudodon tumidus* MORELET, Ser. Conch., IV, 1875, p. 337, pl. XVI, fig. 1.

Siam; Cambodia.

PSEUDODON OVALIS Morlet.

* *Pseudodon oralis* MORLET, Jl. de Conch., XXXVII, 1889, p. 197, pl. VII, fig. 3.

Srakeo River, Siam.

¹Gould did not figure this, but the figure in the Conchologia Indica accurately represents his species, judging from a cotype in Lea's collection, sent him by Dr. Gould. A number of these Pseudodons are very close, and it is quite probable that some of them must be united when sufficient material is studied.

PSEUDODON THOMSONI Morlet.

- * *Pseudodon thomsoni* MORLET, Jl. de Conch., XXXII, 1884, p. 401, pl. XIII, figs. 2, 2a.
 * *Margaritana thomsoni* P.ETEL, Conch. Sam., III, 1890, p. 174.

Cambodia.

† PSEUDODON SALWENIANUS Gould.

- * *Anodon salweniana* GOULD, Pr. Bost. Soc. N. Hist., I, 1844, p. 160.
 * *Anodonta salweniana* GOULD, Otia Conch., 1862, p. 193.
 * *Pseudodon salweniana* CONRAD, Am. Jl. Conch., I, 1865, p. 233.
 * *Monocondylwa salweniana* P.ETEL, Conch. Sam., III, 1890, p. 174.
 * *Unio salwenianus* SOWERBY, Conch. Icon., XVI., 1868, pl. XCIV, fig. 513.
 * *Margaron (Monocondylwa) salweniana* LEA, Syn., 1870, p. 72.
 * *Pseudodon salwenianum* HANLEY and THEOBALD, Conch. Ind., 1876, p. 5, pl. IX, fig. 4.
 * *Unio salwenianus* P.ETEL, Conch. Sam., III, 1890, p. 166.

River Salwen, British Burma.

† PSEUDODON MORELETI Crosse and Fischer.

- * *Monocondylwa tumida* DESHAYES and JULLIEN, Nouv. Arch. de Mus., X, 1874, p. 117, pl. V, figs. 1-3.¹
 * *Pseudodon moreleti* CROSSE and FISCHER, Jl. de Conch., XXV, 1876, p. 330.

Cambodia.

† PSEUDODON CAMBODJENSIS Petit.

- * *Monocondylwa cambodjensis* PETIT, Jl. de Conch., VI, 1865, p. 122.—* P.ETEL, Conch. Sam., III, 1890, p. 174.
 * *Margaron (Monocondylwa) cambodjensis* LEA, Syn., 1870, p. 72.
 * *Pseudodon cambodjensis* CONRAD, Am. Jl. Conch., I, 1865, p. 233.—* MORLET, J. de Conch., XXXII, 1866, p. 266.

Cambodia; Tonkin.

PSEUDODON CHAPERI de Morgan.

- * *Pseudodus chaperi* DE MORGAN, Bull. Soc. Zool. de Fr., X, 1885, p. 423, pl. IX, figs. 1, 2.

Cambodia; Siam.

The following species is unfigured:

PSEUDODON NICOBARICUS Mörch.

- Alasmodonta* (subg. ?) *nicobarica* MÖRCH, Jl. de Conch., XX, 1872, p. 327.²

Nicobar Islands.

¹ This fine, large shell which Deshayes and Jullien supposed to be *P. tumidus* of Morelet is quite a different thing, and Crosse and Fischer have given it a new name.

² Mörch describes the above under the name *Alasmodonta* (subgenus ?) *nicobarica*, and credits it to Fabricius. The *Mytilus nicobariensis* of the latter was only a *nomen nudum*. Mörch says his species is close to *Anodonta zollingeri*.

Section BINEURUS Simpson, 1900.

Shell elongate rhomboid, thin, rounded in front, widely and feebly biangulate behind, having two or more raised radiating lines on the posterior slope, which is somewhat obliquely wrinkled; beaks low; surface finely, irregularly, concentrically grooved; epidermis olive; teeth smooth, compressed; nacre bluish. Animal unknown.

(Type, *Monocondylaea mouhoti* Lea.)

† PSEUDODON MOUHOTI Lea.

* *Monocondylaea mouhoti* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 190.

* *Pseudodon manhoti* CONRAD, Am. Jl. Conch., I, 1865, p. 233.

* *Monocondylaea mouhotiana* LEA, Jl. Ac. N. Sci. Phila., VI, 1866, p. 65, pl. XXI, fig. 62; * Obs., XI, 1867, p. 69, pl. XXI, fig. 62.—* Clessin, Conch. Cab. Ano., 1876, p. 261, pl. LXXXII, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 175.

* *Margaron (monocondylaea) mouhotiana* LEA, Syn., 1870, p. 73.

* *Unio mouhotianus* SOWERBY, Conch. Icon., XVI, 1868, pl. XCII, fig. 503.

* *Pseudodon mouhotianus* MORELET, Ser. Conch., IV, 1875, p. 339.

Cambodia; Siam.

PSEUDODON EXILIS Morelet.

* *Monocondylus exilis* MORELET, Jl. de Conch., XIV, 1866, p. 63.

* *Pseudodon exilis* MORELET, Ser. Conch., IV, 1875, p. 340, pl. XVII, fig. 1.

Perak.

PSEUDODON AVA Theobald.

* *Monocondylaea ava* THEOBALD, Jl. As. Soc. Beng., XLII, Pt. 2, 1873, p. 209, pl. XVII, fig. 15.¹

Mandelay, Burma.

Section NASUS Simpson, 1900.

(Type, *Pseudodon nankingensis* Heude.)

Shell somewhat solid, rather inflated, much elongated, with a well-developed posterior ridge, with a low, faint one above it, bluntly pointed at post-basal part; beaks not high, eroded in the specimens seen; epidermis dark; teeth compressed, high.²

Animal unknown.

† PSEUDODON NANKINGENSIS Heude.

* *Monocondylaea nankingensis* HEUDE, Jl. de Conch., XXII, 1874, p. 116; Conch. Fluv. Nank., I, 1875, pl. IV, fig. 9.

* *Monocondylaea nankingensis* PÆTEL, Conch. Sam., III, 1890, p. 174.

Rivers of Nankin.

¹I am not certain that this belongs in this group.

²I have seen only a single shell of each of the two species of this group, both very badly eroded and decayed, so that I can not give the characters accurately. The animals may show that they are not closely related to the ordinary Pseudodons.

† PSEUDODON SECUNDUS Heude.

* *Pseudodon secundus* HEUDE, Conch. Fluv. Nank., III, 1877, pl. XVIII, fig. 38.

Unio secundus PÆTEL, Conch. Sam., III, 1890, p. 167.

River Hoæ, China.

Section OBOVALIS Simpson, 1900.

(Type, *Pseudodon loomisi* Simpson.)

Shell obovate subinflated, with a low, rounded posterior ridge, and with radiating wrinkles on the posterior slope; disk nearly free from wrinkles; epidermis blackish; there is a high, triangular tooth in each valve; laterals very faint, rounded; nacre coppery.

Animal unknown.

† PSEUDODON LOOMISI Simpson.

* *Pseudodon loomisi* SIMPSON, Pr. Ac. N. Sci. Phila., 1900, p. 84, pl. IV, fig. 7.

Japan.

The following are unfigured or indeterminate species.

Monocondylaea walpolei HANLEY, Proc. Zool. Soc. Lond., 1871, p. 587. Borneo.

Spatha compressa VON MARTEYS, Proc. Zool. Soc. Lond., 1860, p. 66. Siam.

Pseudodon sulcatum ROCHEBRUNE, Bull. Soc. Phil., 7th ser., VI, 1882, p. 41. Mekong River, Cochin China.

Pseudodon pierri ROCHEBRUNE, Bull. Soc. Phil., VI, 1882, p. 41. Cochin China.

Pseudodon mabiliei ROCHEBRUNE, Bull. Soc. Phil., VI, 1882, p. 41. Cochin China, Cambodia.

Pseudodon auodontinus ROCHEBRUNE, Bull. Soc. Phil., VI, 1882, p. 41. Cambodia.

Genus PARREYSIA Conrad, 1853.

(Type, *Unio multidentatus* Philippi.)

Parreysia CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 267.

Shell solid, inflated, oval to subrhomboid, with full, high, zigzag, radially sculptured beaks, the sculpture often extending over the disk; epidermis smooth and bright, sometimes a little rayed, with two irregular pseudocardinals in the left valve which are more or less broken into ragged denticles or are strongly, vertically striate, and two laterals, the lower the larger; right valve with one, sometimes two pseudocardinals, the upper small, compressed, and a few tubercles behind them, with two laterals, the upper the larger; cavity of the beaks rather deep, not compressed; dorsal scars under the hinge, not visible; the two upper anterior muscle scars very deep, confluent, the lower linear; nacre white to salmon, iridescent behind.

Animal unknown.

Subgenus PARREYSIA Conrad, 1853.

Characters the same as the genus.

¹ In the above reference Conrad gives the name *Parreysia* and refers to *Unio multidentatus* Parreys, which may be taken as the type. I do not know that he ever described the group.

(Group of *Parreysia corrugata*.)

Shell having the middle of the basal region swollen, beak sculpture strong, the central ridges generally united at their lower points to form chevron-shaped markings.

† PARREYSIA CORRUGATA Müller.

* *Mya corrugata* MÜLLER,¹ Verni. Terr. et Fluv., 1774, Pt. 2, p. 214; * Besch. Ges. Nat. Ber., IV, 1779, p. 56, pl. IIIb, figs. 7, 8.—* GMELIN,² Sys. Nat., 13th ed., 1788, p. 3221.—* SCHROTER, Fluss. Conch., 1779, p. 181, pl. IX, fig. 3.—* WOOD, Gen. Conch., I, 1815, p. 108, pl. XXIV, figs. 1-3.—* DILLWYN, Cat., I, 1817, p. 52.—MAWE, Linn. Conch., 1823, pl. IV, fig. 3.—* WOOD, Ind. Test., 1825, p. 12, pl. II, fig. 31 a; 1856, rev. ed., p. 16, pl. II, fig. 31.

* *Unio corrugata* LAMARCK,³ An. sans Vert., VI, 1819, p. 78.—* DESHAYES, Enc. Meth., II, 1827, p. 584, pl. CCXLVIII, fig. 8.

* *Unio (Potamida) corrugata* SWAINSON, Tr. on Mal., 1840, p. 268, fig. 51; p. 281, fig. 57.

* *Unio corrugata* RETZIUS, Diss. Hist. Nat., 1778, p. 18.—* SPENGLER, Skriv. Selsk. Nat., III, 1793, p. 68.—* HANLEY, Test. Moll., 1842, p. 197; * Biv. Shells, 1843, p. 197.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, pp. 493, 497.—* KUSTER, Conch. Cab. Unio, 1862, p. 289, pl. XCVII, figs. 3, 4.—* BLANFORD, Jl. As. Soc. Beng., XXXV, 1867, p. 136.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXI, fig. 360.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 21, pl. XLV, figs. 2-5.—* PETEL, Conch. Sam., III, 1890, p. 149.

* *Margarita (Unio) corrugatus* LEA, Syn., 1836, p. 29; 1838, p. 21.

* *Margaron (Unio) corrugatus* LEA, Syn., 1852, p. 20; 1870, p. 30.

* *Mya spuria* GMELIN, Syst. Nat., 13th ed., 1788, p. 322; * WOOD, Gen. Conch., I, 1815, p. 110.—* LAMARCK, An. sans Vert., VI, 1819, p. 80.

Mya gaditana SCHREIBERS, Versuch., 1793, p.

* *Unio multidentatus* PHILIPPI,⁴ Conch., III, 1847, p. 46, pl. III, fig. 4.—* KUSTER, Conch. Cab. Unio., 1862, p. 136, pl. XXXVI, fig. 5.—* PETEL, Conch. Sam., III, 1890, p. 160.

* *Margaron (Unio) multidentatus* LEA, Syn., 1870, p. 50.

* *Unio fulmineus* PHILIPPI,⁵ Conch., III, 1847, p. 46, pl. III, figs. 5, 6.—* KUSTER, Conch. Cab. Unio., 1862, p. 286, pl. XCVI, figs. 2, 3.—* PETEL, Conch. Sam., III, 1890, p. 153.

* *Unio lutens* LEA,⁶ Pr. Ac. Nat. Sci. Phila., VIII, 1856, p. 93.

* *Unio luteus* LEA, Jl. Ac. N. Sci. Phila., III, 1857, p. 291., pl. XXIV, fig. 4; * Obs., VI, 1857, p. 11, pl. XXIV, fig. 4.

¹ Müller's description of *Mya corrugata* is inadequate, but in the Beschäftigungen it is further discussed and characteristic figures are given of the species we know as *Unio corrugatus* of India.

² Refers to Conch. Cab. VI, pl. III, fig. 22, which is probably Müller's shell. Chemnitz refers to it as *Mya corrugata teste orali*, etc. The shell referred to by him in Conch. Cab., X, p. 346, as *Mya corrugata magna*, etc., is a very different thing.

³ Lamarck refers to Enc. Meth. Pt. 19, 1797, pl. CCXLVIII, figs. 6, 8 a, b. I am not at all sure that fig. 6 is the *corrugatus* of Müller.

⁴ Credited by Philippi to Parreyss. I think there can be no doubt that this is the *Mya corrugata* of Müller, as the figure and description entirely agree with the species as we know it.

⁵ Parreyss is the author, according to Philippi. This and *multidentatus* are credited by Philippi to Australia. No shells like these have ever been found there.

⁶ *Lutens* in text, by error no doubt.

- * *Margaron (Unio) luteus* LEA, Syn., 1870, p. 46.
 * ? *Unio semirugatus* CHENU, Ill. Conch., 1858, pl. XII, figs. 2, 2a.
 * *Unio merodabensis* KUSTER, Conch. Cab. Unio, 1861, p. 233, pl. LXXVIII, fig. 4.—
 * BLANFORD, J. As. Soc. Beng., XXXV, 1866, p. 142.—* PÆTEL, Conch. Sam.,
 III, 1890, p. 159.
 * *Unio wynegungansis* HANLEY and THEOBALD,¹ Conch. Ind., 1876, p. 21, pl. XLV,
 fig. 6.
 * *Unio tennenti* HANLEY and THEOBALD, Conch. Ind., 1876, p. 22, pl. XLV, figs. 7-9.—
 * PÆTEL, Conch. Sam., III, 1890, p. 169.
 * *Unio phayresi* THEOBALD, manuscript.

PARREYSIA CORRUGATA var. LÆVIROSTRIS Benson.

- * *Unio lævirostris* BENSON, Ann. and Mag., 1862, p. 192; * Jl. As. Soc. Beng., 1862,
 XXXV, p. 144.—* PÆTEL, Conch. Sam., III, 1890, p. 157.
 * *Unio corrugatus* var. *lævirostris* HANLEY and THEOBALD, Conch. Ind., 1876, p. 21,
 pl. XLV, figs. 5, 6.—* PÆTEL, Conch. Sam., III, 1890, p. 149.

† PARREYSIA CORRUGATA var. NAGPOORENSIS Lea.

- * *Unio nagpoorensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 331; * Jl. Ac. N. Sci.
 Phila., IV, 1860, p. 270, pl. XLV, fig. 150; * Obs., VII, 1860, p. 88, pl. XLV, fig.
 150.—* BLANFORD, Jl. As. Soc. Beng., XXXV, 1866, p. 143.
 * *Margaron (Unio) nagpoorensis* LEA, Syn., 1870, p. 38.
 * *Unio corrugatus* Müll. var. *nagpoorensis* HANLEY and THEOBALD, Conch. Ind.,
 1876, p. 21.—PÆTEL, Conch. Sam., 1890, p. 149.

India; Burma.?

† PARREYSIA WYNEGUNGÆNSIS Lea.

- * *Unio wynegungansis* LEA,² Pr. Ac. N. Sci. Phila., III, 1859, p. 331; * Jl. Ac. N.
 Sci. Phila., IV, 1860, p. 271, pl. XLV, fig. 151; * Obs., VII, 1860, p. 89, pl. XLV,
 fig. 151.—* BLANFORD, Jl. As. Soc. Beng., XXXV, 1866, p. 143.—* SOWERBY,
 Conch. Icon., XVI, 1868, pl. LXVII, p. 339.—* PÆTEL, Conch. Sam., III, 1890,
 p. 172.
 * *Margaron (Unio) wynegungansis* LEA, Syn., 1870, p. 50.

Wynegunga River, Bengal.

† PARREYSIA FAVIDENS Benson.

- * ? *Mya spuria* Wood, Ind. Test., 1825, p. 12, pl. II, fig. 35a.
 * *Unio favidens* BENSON,³ Ann. and Mag., X, 1862, p. 188; * Jl. As. Soc. Beng.,
 XXXV, 1867, p. 138.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 6,
 pl. XI, fig. 1.—* REEVE, Conch. Icon., XVI, 1865, pl. XXVI, fig. 131.—* PÆTEL,
 Conch. Sam., III, 1890, p. 152.
 * *Margaron (Unio) favidens* LEA, Syn., 1870, p. 38.
 * *Unio flavidens* PÆTEL, Conch. Sam., III, 1890, p. 152.
 * *Unio trirostris* MUSGRAVE, Hanley Phot. Conch., 1863, pl. II, fig. 9.—* HANLEY
 and THEOBALD,⁴ Conch. Ind., 1876, p. 6, pl. XI, fig. 6.—* PÆTEL, Conch.
 Sam., III, 1890, p. 170.

¹ I think this is *corrugatus* and not the *wynegungansis* of Lea.

² Heavier, more elongated and pointed behind than *corrugata*, yet probably only a variety of it.

³ Several varieties, *marcens*, *trigona*, *delta*, *chrysis*, *viridula*, and *densa* are given by Benson, which hardly seem to me to be worth mentioning.

⁴ This = *U. tripartitus* of Lea, with a darker epidermis and a more pronounced basal swelling. There is every variation in the coloring and swelling.

† *Unio tripartitus* LEA, Pr. Ac. N. Sci. Phila., VII, 1863, p. 190; *Jl. Ac. N. Sci. Phila., VI, 1866, p. 57, pl. XIX, fig. 55; *Obs., XI, 1867, p. 61, pl. XIX, fig. 55.

**Margaron (Unio) tripartitus* LEA, Syn., 1870, p. 35.¹

**Unio favidens* var. *marcens* BENSON, Ann. and Mag., X, 1862, p. 188.

**Unio marcens* HANLEY and THEOBALD, Conch. Ind., 1876, p. 19, pl. XLII, figs. 4-6.—*PÆTEL, Conch. Sam., III, 1890, p. 158.

**Unio smaragdites* BENSON, Ann. and Mag., X, 1862, p. 190.—*BLANFORD, Jl. As. Soc. Beng., XXXV, 1866, p. 147.—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 5, pl. X, fig. 5.—*PÆTEL, Conch. Sam., III, 1890, p. 167.

PARREYSIA FAVIDENS var. PINAX Benson.

**Unio pinax* BENSON, Ann. and Mag., 1862, p. 192; *Jl. As. Soc. Beng., XXXV, 1866, p. 144.

**Unio favidens* var. *pinax* HANLEY and THEOBALD, Conch. Ind., 1876, p. 6, pl. X, fig. 2.—*PÆTEL, Conch. Sam., III, 1890, p. 152.

PARREYSIA FAVIDENS var. PLAGIOSOMA Benson.

**Unio plagiosoma* BENSON, Ann. and Mag., X, 1862, p. 191; *Jl. As. Soc. Beng., XXXV, 1866, p. 144.

**Unio favidens* var. *plagiosoma* HANLEY and THEOBALD, Conch. Ind., 1876, p. 6, pl. XI, fig. 3.—*PÆTEL, Conch. Sam., III, 1890, p. 152.

India.

† PARREYSIA BHAMOENSIS Theobald.

**Unio bhamoensis* THEOBALD, Jl. As. Soc. Beng., XLII, 1874, Pt. 2, p. 207, pl. XVII, fig. 1.—*? HANLEY and THEOBALD, Conch. Ind., 1876, p. 62, pl. CLV, fig. 2.—*PÆTEL, Conch. Sam., III, 1890, p. 146.

*† *Unio mandalayensis* THEOBALD, Jl. As. Soc. Beng., XLII, 1874, p. 208, pl. XVII, fig. 2.—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 62, pl. CLIV, fig. 4.—*PÆTEL, Conch. Sam., III, 1890, p. 158.

Burma.

† PARREYSIA FEDDENI Theobald.

**Unio feddeni* THEOBALD, Jl. As. Soc. Beng., XLII, 1874, p. 208, pl. XVII, fig. 3.²

Penugunga River, Central India.

† PARREYSIA TAVOYENSIS Gould.

**Unio tavoyensis* GOULD, Pr. Bost. Soc. N. Hist., I, 1843, p. 140.—*CATLOW and REEVE, Conch. Nom., 1845, p. 64.—*KUSTER, Conch. Cab. Unio, 1856, p. 166, pl. XLVIII, fig. 2.—*GOULD, Otia Conch., 1862, p. 190.—*REEVE, Conch. Icon., XVI., 1864, pl. XIII, fig. 49.³—*BLANFORD, Jl. As. Soc. Beng., XXXV, 1866, p. 148.—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 62, pl. CLIV, figs. 6, 7.—*PÆTEL, Conch. Sam., III, 1890, p. 169.

**Margaron (Unio) tavoyensis* LEA, Syn., 1870, p. 31.

**Unio savoyensis* PÆTEL, Conch. Sam., III, 1890, p. 166.

**Unio parma* BENSON, Sowerby Conch. Icon., XVI, 1868, pl. XCV, fig. 514.—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 61, pl. CLIV, fig. 1.—*PÆTEL, Conch. Sam., III, 1890, p. 162.

¹ Lea claims that his species has triple laterals in the left valve. Specimens of *favidens* have the same character, and there is much variation in this particular.

² A young specimen in the U. S. National Museum collection from Morelet's collection is a good deal like *P. favidens*.

³ Gould did not figure this, but Reeve's figure agrees well with the types.

†*PARREYSIA TAVOYENSIS* var. *TRIEMBOLUS* Benson.

- * *Unio triembolus* BENSON, Jl. As. Soc. Beng., XXXV, 1855, p. 144; *Ann. and Mag., X, 1862, p. 190.—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 43, pl. CVII, fig. 2.—*PÆTEL, Conch. Sam., III, 1890, p. 170.

*† *Unio houngdaranicus* TAPPERONE-CANEFRI, Ann. Mus. Civ., VII, 1889, p. 341.

Burma; India.

†*PARREYSIA VULCANA* Hanley.

- * *Unio vulcanus* HANLEY, Pr. Zool. Soc. Lond., 1875, p. 606.—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 62, pl. CLV, fig. 3.—*PÆTEL, Conch. Sam., III, 1890, p. 172.

Burma; Pegu.

†*PARREYSIA FEÆ* Tapperone-Canefri.

- * *Unio feæ* TAPPERONE-CANEFRI, Ann. Mus. Civ. Gen., 2d ser., VII, 1889, p. 340.¹

Meetan River, Burma.

†*PARREYSIA RAJAHENSIS* Lea.

- (* *Unio rajahensis* LEA, Pr. Am. Phil. Soc., II, 1841, p. 30; *Tr. Am. Phil. Soc., VIII, 1842, p. 239, pl. XXIII, fig. 53; *Obs., III, 1842, p. 77, pl. XXIII, fig. 53.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—*CHENU, Ill. Conch., 1858, pl. XXVI, figs. 3, 3a, 3b.—*PÆTEL, Conch. Sam., III, 1890, p. 165.

(* *Margaron (Unio) rajahensis* LEA, Syn., 1852, p. 25; 1870, p. 38.

(* *Unio indicus* SOWERBY, Conch. Icon., XVI, 1866, pl. XL, fig. 222.—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 43, pl. CVII, fig. 1.

(* *Margaron (Unio) indicus* LEA, Syn., 1870, p. 31.

India.

(Group of *Parreysia rugosa*.)

Shell rather solid, elliptical to subtrapezoidal; beaks full, with zig-zag-radial sculpture and fine corrugations and granules over the surface of the shell.

†*PARREYSIA RUGOSA* Gmelin.

- * *Mya rugosa* GMELIN, Syst. Nat., 13th ed., 1788, p. 3222.²—*WOOD, Gen. Conch. I, 1815, p. 109.—*DILLWYN, Cat. I, 1817, p. 53.—*WOOD, Ind. Test., 1825, p. 12, pl. II, fig. 33a; rev. ed., 1856, p. 16, pl. II, fig. 33.

(* *Unio rugosus* KUSTER, Conch. Cab. Unio, 1862, p. 290, pl. XCVII, fig. 5.—*BLANFORD, Jl. As. Soc. Beng., XXXV, 1867, p. 137.—*PÆTEL, Conch. Sam., III, 1890, p. 166.

Coromandel.

¹Not figured. Shells in the U. S. National Museum from Fea show this to be a valid species.

²Gmelin refers to the *Mya corrugata magna*, etc., of Chemnitz, in the Conch. Cab., X, p. 376, pl. CLXX, fig. 1649. There is a shell in Lea's collection from the Vellaur River, southeast India, which seems to be this species.

† PARREYSIA CORBIS Hanley.

* *Unio corbis* HANLEY, Biv. Shells, 1856, p. 386, pl. XXIII, fig. 43.¹—*HANLEY and THEOBALD, Conch. Ind., 1876, p. 22, pl. XLV, fig. 10.

*Margaron (*Unio*) *corbis* LEA, Syn., 1870, p. 50.

Assam.

† PARREYSIA BURMANUS Blanford.

* *Unio burmanus* BLANFORD, Pr. Zool. Soc. Lond., 1869, p. 449.

* *Unio birmanus* HANLEY and THEOBALD, Conch. Ind., 1876, p. 19, pl. XLII, fig. 1.—

*PETEL, Conch. Sam., III, 1890, p. 146

Burma.

† PARREYSIA SIKKIMENSIS Lea.

* *Unio sikkimensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 151; *Jl. Ac. N. Sci. Phila., IV, 1859, p. 251, pl. XXXIX, fig. 131; * Obs., VII, 1860, p. 69, pl. XXXIX, fig. 131.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVI, fig. 400.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 6, pl. XI, fig. 4, p. 44, pl. CVII, figs. 6, 7.—

*PETEL, Conch. Sam., III, 1890, p. 167.

*Margaron (*Unio*) *sikkimensis* LEA, Syn., 1870, p. 64.

Sikkim, India.

† PARREYSIA GOWHATTENENSIS Theobald.

* *Unio gowhattenensis* THEOBALD, Jl. As. Soc. Beng., XLII, 1873, p. 208, pl. XVII, fig. 4.—*PETEL, Conch. Sam., III, 1890, p. 154.

Gowhatti, in Assam.

PARREYSIA SIMPULARIS Heude.

* *Unio modestus* HEUDE,² Conch. Fluv. Nank., II, 1877, pl. XIV, fig. 29.

* *Unio simpularis* HEUDE, Jl. de Conch., XXXII, 1884, p. 20.—*PETEL, Conch. Sam., III, 1890, p. 167.

China.

PARREYSIA TRISULCATA Heude.

* *Unio trisulcatus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LVII, fig. 108.

China.

(Group of *Parreysia leopoldvillensis*.)

Shell rounded rhomboid, somewhat inflated, subsolid, with a rounded post ridge; beaks full, the sculpture strongly and regularly zigzag-radial and extending well over the body of the shell; pseudocardinals heavy, subcompressed, crenate; anterior scars deep.

¹ Credited to Benson manuscript.

² The name *modestus* has been used by authors for a Brazilian *Unio*, but I can not find that it was ever described. Later Kuster used the name for a Chilean species; hence Heude changed his first name to *simpularis*. It is quite probable that this species should be placed in a separate group.

PARREYSIA LEOPOLDVILLENSIS Putzeys.

Unio leopoldvillensis PUTZEYS, Proc. Verb. Soc. Mal. Belg., XXVII, 1898, p. XXVII, figs 12, 13.¹

(Group of *Parreysia bakeri*.)

Shell small, inflated, oval, rounded in front, bluntly pointed behind, inflated at or behind the central base; beaks full, high, sculptured with very strong, zigzag bars which extend part way over the disk and end rather suddenly, below which the shell is smooth; posterior ridge well developed, rounded; two compressed pseudocardinals and one lateral in the right valve, two pseudocardinals and two laterals in the left valve.²

PARREYSIA BAKERI H. Adams.

* *Unio bakeri* H. ADAMS, Pr. Zool. Soc. Lond., 1866, p. 376.—* PÆTEL, Conch. Sam., III, 1890, p. 145.—* SMITH, Ann. and Mag., X, 1892, p. 126, pl. XII, fig. 11.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 231, pl. VII, fig. 6.

Lake Albert Nyanza, Central Africa.

PARREYSIA STUHLMANNI von Martens.

* *Unio stuhlmanni* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 231, pl. VII, fig. 13.

Albert Edward Lake, Central Africa.

NODULARIA HAUTTECOEURI Bourguignat.

* *Unio hauttecoeri* BOURGUIGNAT, Moll. Ny. Ouk., 1883, p. 5, figs. 1-3.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 232, pl. I, fig. 23; VII, fig. 3.

NODULARIA HAUTTECOEURI var. EDWARDSIANA Bourguignat.

* *Unio edwardsianus* BOURGUIGNAT, Moll. Ny. Ouk., 1883, p. 12, figs. 7, 9.

Lake Victoria Nyanza, Central Africa.

PARREYSIA DUPONTI Rochebrune.

* *Reneus duponti* ROCHEBRUNE, Bull. Soc. Zool. Fr., XL, 1881, p. 481.

* *Unio duponti* ROCHEBRUNE, Bull. Soc. Philom., VI, 1882, p. 34.—* BOURGUIGNAT, Moll. Ny. Ouk., 1883, p. 8, figs. 10-12.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

* *Unio bakeri* VON MARTENS, Sitzber. Ges. Nat. Fr., 1879, p. 104.

* *Unio grandidieri* BOURGUIGNAT, Moll. Ny. Ouk., 1883, p. 7, figs. 4-6.—* PÆTEL, Conch. Sam., III, 1890, p. 154.—* VON MARTENS, Besch. Deuts. Ost-Af., 1874, p. 233.

Lake Victoria Nyanza.

PARREYSIA RUELLANI Bourguignat.

* *Unio ruellani* BOURGUIGNAT, Moll. Ny. Ouk. 1883, p. 10, figs. 16-18.—* PÆTEL, Conch. Sam., III, 1890, p. 166.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 235.

Lake Victoria Nyanza.

¹ I have not seen this species, and I place it here with some doubt.

² I adopt this group with practically the same limits as given by von Martens in the Beschalte. Unfortunately I have never seen any species belonging to it.

PARREYSIA MULTICOLOR von Martens.

* *Unio multicolor* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 236, pl. VII. fig. 4.

Lake Victoria Nyanza.

PARREYSIA NGESIANUS von Martens.

* *Unio ngesianus* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 234, pl. VII. fig. 7.

Lake Albert Edward Nyanza.

The following unfigured species are placed in this group by von Martens:

Unio billotianus CHARMES, Bull. Soc. Mal. Fr., II, 1885, p. 170, Bagamoyo, Central Africa.

Unio euphymus CHARMES, Bull. Soc. Mal. Fr., II, 1885, p. 171, Bagamoyo.

Unio dumesleanus CHARMES, Bull. Soc. Mal. Fr., II, 1885, p. 168, Bagamoyo.

(Group of *Parreysia fabagina*.)

Shell very small and solid, inflated, compressed on the sides, rhomboid, with a high, sharp posterior ridge and high beaks, having zigzag-radial sculpture sometimes extended over the disk, the posterior slope furnished with upcurved plications; pseudocardinals heavy, torn, one in the right valve, two in the left; muscular impressions circular, the anterior very deep.

PARREYSIA FABAGINA Deshayes.

* *Unio fabagina* DESHAYES, Nouv. Arch. de Mus., X, 1874, p. 128, pl. VII, figs. 4-6.—

* PÆTEL, Conch. Sam., III, 1890, p. 152.—* FISCHER, Bull. Soc. d'Hist. Nat. Autun., 1891, p. 140.

Mekong River, Cambodia.

PARREYSIA BROTI Deshayes.

* *Unio broti* DESHAYES, Nouv. Arch. de Mus., X, 1874, p. 129, pl. VII, figs. 1-3.—

* CROSSE and FISCHER, Jl. de Conch., XXVI, p. 323.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

Mekong River, Cambodia.

(Group of *Parreysia nyassaensis*.)

Shell small, solid, inflated, triangular to rhomboid, the base straight or even a little arcuate, but sometimes having a slight fullness behind the central part; a high, well-defined posterior ridge ends at the post base and is sometimes slightly double; posterior slope abrupt; beaks high, with zigzag-radial sculpture, the bars approaching and often coalescing behind the center of the disk, the whole shell frequently corrugate sculptured; epidermis greenish, often slightly rayed; teeth solid, much like those of typical *Parreysia*; nacre white or bluish, sometimes microscopically, granularly radiate outside the pallial line; beak cavities not deep; anterior scars distinct.

Animal unknown.

† *PARREYSIA NYASSAENSIS* Lea.

- * *Unio nyassaensis* LEA, Pr. Ac. N. Sci. Phila., VIII, 1864, p. 108; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 33, pl. XII, fig. 32; * Obs., XI, 1867, p. 37, pl. XII, fig. 32.—
 * SMITH, Pr. Zool. Soc. Lond., 1881, p. 298, pl. XXXIV, fig. 34.—* PÆTEL, Conch. Sam., III, 1890, p. 161.—* ANCEY, Bull. Soc. Zool. Fr., VII, 1894, p. 225.—
 * VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 230.
 * Margaron (*Unio nyassaensis* LEA, Syn., 1870, p. 30.
 * *Unio nyassa* SOWERBY, Conch. Icon., XVI, 1868, pl. XLI, figs. 224, 224a, 224b.
 * *Unio nyassanus* BOURGUIGNAT, Bull. Soc. Mal. Fr., VI, 1889, p. 38.
 * *Unio hermosus* BOURGUIGNAT, Bull. Soc. Mal. Fr., VI, 1889, p. 38.—* ANCEY, Bull. Soc. Zool. Fr., VII, 1894, p. 226.¹

Lake Nyassa, Central Africa.

† *PARREYSIA KIRKII* Lea.

- * *Unio kirkii* LEA, Pr. Ac. N. Sci. Phila., VIII, 1864, p. 108; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 32, pl. XII, fig. 30; * Obs., XI, 1867, p. 36, pl. XII, fig. 30.—* BOURGUIGNAT, Bull. Soc. Mal. Fr., VI, 1889, p. 38.—* ANCEY, Bull. Soc. Zool. Fr., VII, 1894, p. 225.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 229.
 * Margaron (*Unio kirkii* LEA, Syn., 1870, p. 30.

Lake Nyassa.

PARREYSIA UJIJENSIS Bourguignat.

- * *Unio nyassaensis* var., SMITH, Pr. Zool. Soc. Lond., 1881, p. 298, pl. XXXIV, fig. 34.²
 * *Grandidieria ujijensis* BOURGUIGNAT, Bull. Soc. Mal. Fr., II, 1885, p. 7.

Lake Tanganyika.

† *PARREYSIA AFERULA* Lea.

- * *Unio aferula* LEA, Pr. Ac. N. Sci. Phila., VIII, 1864, p. 109.
 * *Unio aferulus* LEA, Jl. Ac. N. Sci. Phila., VI, 1866, p. 34, pl. XIII, fig. 34; * Obs., XI, 1867, p. 38, pl. XIII, fig. 34.—* BOURGUIGNAT, Bull. Soc. Mal. Fr., VI, 1889, p. 38.—* ANCEY, Bull. Soc. Zool. Fr., VII, 1894, p. 226.
 * Margaron (*Unio aferulus* LEA, Syn., 1870, p. 30.

Lake Nyassa.

PARREYSIA MONCETI Bourguignat.

- * *Unio monceti* BOURGUIGNAT, Moll. Ny. Ouk., 1883, p. 15, figs. 13-15.—* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 228.

Lake Victoria, Nyanza.

¹ E. A. Smith, in Pr. Zool. Soc. Lond., 1877, p. 719; 1893, p. 640, believes that *Unio kirkii* Lea and *aferulus* Lea are but varieties of *Unio nyassaensis*, and that *Unio hermosus* Bourguignat is identical with it. It is possible that the three forms of Lea may blend together, but the types are so distinct that I do not feel justified in uniting them even as varieties one of another.

² There is some confusion about this shell. The form on pl. XXXIV, which Smith probably refers to in the text as fig. 34b is simply fig. 34 on the plate, and is, I think, a distinct species, which has received the name of *ujijensis* from Bourguignat. There is no fig. 34b on the plate. His 34a, which Smith calls var. *tanganyicensis*, is *U. nyassaensis* Lea. Bourguignat probably alluded to the fig. 34 when he bestowed his name, although he speaks of the var. *tanganyicensis* Smith. These must not be confounded with the original *Unio tanganyicensis* of Smith, which is a very different thing.

PARREYSIA HYPISPRIMNUS von Martens.

* *Unio hypsiprimnus* VON MARTENS, Besch. Deuts. Ost-Af., 1897, p. 230, pl. VII, fig. 1.¹

Lake Nyassa.

(Group of *Parreysia molleuri*.)

Shell subquadrate, solid, inflated, rounded in front, truncated behind, with high beaks, the sculpture of which was not observed, and a strong, sharp, curved posterior ridge; surface concentrically grooved; pseudo-cardinals thick, short, cut up with vertical grooves; laterals curved; nacre white.

PARREYSIA MOLLEURI Morlet.

* *Unio molleuri* MORLET, Jl. de Conch., XXXIX, 1891, p. 242, pl. VII, fig. 4.

Valley of the Mekong River.

Subgenus **AURORA** Simpson, 1900.

(Type, *Unio aureus* Heude.)

Shell small, ovate, subinflated, pointed behind, rather solid, with high, small beaks and strong, corrugated sculpture, which extends over half the disk in somewhat lachrymous, zigzag ridges and nodules, the outer part of the disk apparently having one or more strong, concentric ridges, the posterior slope being radiately ridged and nodulous; epidermis fulvous, beautifully and delicately green radiated; pseudo-cardinals conical, truncate, striate; laterals lamellar; nacre orange.²

PARREYSIA AUROREA Heude.

* *Unio aureus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LVII, fig. 106.--

* PÆTEL, Conch. Sam., III, 1890, p. 145.

China.

Subgenus **PSEUDOBAPHIA** Simpson, 1900.

(Type, *Unio biesianus* Heude.)

Shell large, oval, inflated, gaping in front and behind, rather solid, with full beaks, the beak sculpture not seen, posterior part distinctly biangular; ligament large; epidermis smooth, brownish, somewhat rayed, right valve with a large, irregular pseudocardinal in front of the beak, with a pit in front of it; behind it on the hinge are numerous denticles, and at some distance behind two very faint laterals; left valve with two large pseudocardinals, one behind the other, and two remote

¹I am not positive as to the systematic position of this and the last two species. The figure of the latter is from a very badly eroded shell, and it is difficult to tell what its affinities are.

²Heude's brief Latin description so poorly describes the peculiar species placed here that I can not be sure where it belongs, and that it should not have generic rank. I place it here provisionally.

blurred laterals; two upper anterior muscle scars united, very deep; posterior scars united, the lower long and oblique; beak cavities very large, deep; nacre flesh colored, dull.¹

Animal unknown.

† *PARREYSIA BIESIANA* Heude.

* *Unio biesianus* HEUDE, Conch. Fluv. Nank., II, 1877, pl. XIV, fig. 30.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

China.

Subgenus *ACUTICOSTA* Simpson, 1900.

(Type, *Unio chinensis* Lea.)

Shell pointed behind, produced at the center of the base, inflated, solid, with a sharp, pinched-up, but vanishing posterior ridge, beaks full, the sculpture strong, zigzag radial; epidermis smooth, rayed; pseudocardinals somewhat compressed, vertically ridged, ragged; laterals well developed, strongly obliquely ridged and granular.

† *PARREYSIA CHINENSIS* Lea.

* *Unio chinensis* LEA, Pr. Ac. Nat. Sci. Phila., XII, 1868, p. 150; * Jl. Ac. Nat. Sci. Phila., VI, 1868, p. 325, pl. LIII, fig. 138; * Obs., XII, 1869, p. 85, pl. LIII, fig. 138.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* Margaron (*Unio chinensis* LEA, Syn., 1870, p. 30.

PARREYSIA CHINENSIS var. *SQUAMMOSUS* Heude.

* *Unio sinensis* var. *squamosus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LVIII, fig. 113.

PARREYSIA CHINENSIS var. *LÆVIS* Heude.

* *Unio squamosus* var. *lævis* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LIX, fig. 116.

China.

PARREYSIA RETIARIA Heude.

* *Unio retarius* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LVII, fig. 107.—

* PÆTEL, Conch. Sam., III, 1890, p. 165.

China.

Genus *PTYCHORHYNCHUS* Simpson, 1900.

(Type, *Unio pfisteri* Heude.)

Shell elongate, elliptical, round in front, pointed behind, the point being midway of the height of the shell, solid, slightly inflated, with a faint posterior ridge; beaks rather low, sculpture not seen; the posterior slope sculptured with strong, upcurved ridges, which sometimes extend slightly on to the disk, which is otherwise smooth and shining; left valve with two rather stumpy, roughened pseudocardinals and 1 or 2

¹ Heude says the beaks are corrugated.

somewhat blurred granulous laterals; right valve with a single blunt pseudocardinal, the hinge line in front of it being excavated, and a curious blurred lateral which is sometimes slightly split up; beak cavities shallow; dorsal scars few; nacre whitish. Animal unknown.

(Group of *Ptychobranchus pfisteri*.)

Characters as in the genus.

† PTCHORHYNCHUS PFISTERI Heude.

* *Unio pfisteri* HEUDE, Jl. de Conch., XXII, 1874, p. 112; * Conch. Fluv. Nank., I, 1875, pl. I, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 163.

PTYCHORHYNCHUS PFISTERI var. INSPIRATUS Heude.

* *Unio pfisteri* var. *inspiratus* HEUDE, Conch. Fluv. Nank., II, 1877, pl. XIV, fig. 28.—* PÆTEL, Conch. Sam., III, 1890, p. 155.

Nankin River, China.

† PTYCHORHYNCHUS MEDIASTINUS Heude.¹

* *Unio mediastinus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LXIII, fig. 123.—* PÆTEL, Conch. Sam., III, 1890, p. 158.

China.

(Group of *Ptychorhynchus apicellatus*.)

Shell subrhomboid, more or less biangulate behind.

PTYCHORHYNCHUS APICELLATUS Heude.

* *Unio apicellatus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LXIII, fig. 126.

China.

PTYCHORHYNCHUS SCHOMBURGIANUS Heude.

* *Unio schomburgianus* HEUDE, Jl. de Conch., XL, p. 314; * Conch. Fluv. Nank., IX, 1885, pl. LXXII, fig. 139.—* PÆTEL, Conch. Sam., III, 1890, p. 167.

Isle of Hainan, China.

PTYCHORHYNCHUS INCERTUS Simpson.

* *Unio compressus* HEUDE,² Conch. Fluv. Nank., III, 1877, pl. XXIV, fig. 52.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Unio murinus* HEUDE, Conch. Fluv. Nank., VIII, 1883, pl. LXII, fig. 121.

* *Unio morinus* PÆTEL, Conch. Sam., III, 1890, p. 160.

China.

Genus VIRGUS, Simpson, 1900.

(Type, *Unio beccarianus* Tapperone Canefri.)

Shell rather solid, elongated, inequilateral, compressed, rounded in front, nearly straight below, the dorsal line as far back as the hinder

¹ Probably only a variety of *P. pfisteri*.

² The name *compressus* has been twice used previously for a *Unio*. I change it here to *incertus*.

end of the ligament parallel with the base, behind the ligament the shell is obliquely truncate, the truncation somewhat upcurved, the shell ending in a rounded point at the posterior base; posterior ridge elevated, rounded, with sculpture radiating from it above and below; beaks rather low, radially ridged; pseudocardinals small, generally two in each valve; laterals rather short, club-shaped, one in the right and two in the left valve. Animal unknown.

(Group of *Virgus beccarianus*.)

Shell having the posterior ridge high and rounded, outlined on each side by a sulcus, the radiations from this ridge rather fine; surface concentrically sculptured.

VIRGUS BECCARIANUS Tapperone Canefri.

* *Unio beccarianus* TAPPERONE CANEFRI, ANN. MUS. GENOV., XIX, 1883, p. 291, pl. XI, fig. 2.—* P. ET EL, CONCH. SAM., III, 1890, p. 145.

Fly River, New Guinea.

(Group of *Virgus mattioli*.)

Posterior ridge full, rounded, not outlined by sulci, covered with very strong corrugated ridges which curve outward each way from the center, the remainder of the shell smooth.

VIRGUS MATTIOLI Tapperone Canefri.

* *Unio mattioli* TAPPERONE CANEFRI, ANN. MUS. GENOV., XIX, 1883, p. 292, pl. XI, fig. 1.—* P. ET EL, CONCH. SAM., III, 1890, p. 158.

Fly River, New Guinea.

(Group of *Virgus misoolensis*.)

Posterior ridge merely rounded, with radiating folds scattered over the shell, pseudocardinals two in the left valve and one in the right.

VIRGUS MISOLENSIS Schepman.

* *Unio misoolensis* SCHEPMAN, Notes from Leyd. Mus., XVIII, 1896, p. 259, fig.—* DROUET, JI. de CONCH., XLV, 1897, p. 125.

Misool, Borneo.

Genus CTENODESMA Simpson, 1900.

(Type, *Unio borneensis* Issel.)

Shell subtrapezoid to long elliptical, rather thin, compressed, with a scarcely developed posterior ridge and low beaks whose sculpture is densely zigzag radial, becoming finely corrugated or nodulous on the disk and extending over most of the shell; the disk concentrically striate or sulcate at its outer edge; pseudocardinals one to two in each valve, very diverse in form, but showing a tendency to break up into

denticles; one lateral and sometimes a faint secondary one in the left valve and two in the right; dorsal scars few, diversified in form, and variously disposed in the shallow beak cavities; anterior muscle scars united; nacre whitish.

Animal unknown.

† CTENODESMA BORNEENSIS Issel.

* *Unio plicatulus* LEA,¹ Pr. Ac. Nat. Sci. Phila., III, 1859, p. 152; Jl. Ac. Nat. Sci. Phila., IV, 1860, p. 247, pl. XXXVII, fig. 126; *Obs., VII, 1860, p. 65, pl. XXXVII, fig. 126.—* REEVE, Conch. Icon., XVI, 1865, pl. XXII, fig. 102.—* PÆTEL, Conch. Sam., III, 1890, p. 163.

* *Margaron (Unio) plicatulus* LEA, Syn., 1870, p. 31.

* *Unio borneensis* ISSEL, Moll. Borneo, 1874, p. 113.

* *Unio penisatus* FISCHER and CROSSE,² Miss. Sci., Pt. 7, II, 1894, p. 599.

Borneo.

† CTENODESMA GUPPYI Smith.

* *Unio guppyi* SMITH, Pr. Zool. Soc. Lond., 1885, p. 608, pl. XXXVII, figs. 88a, 88b.—* PÆTEL, Conch. Sam., III, 1890, p. 154.

Shortland Island; Solomon Islands.

Genus RECTIDENS Simpson, 1900.

(Type, *Unio prolongatus* Drouet.)

Shell elongated, with an angle at the anterior dorsal part, pointed and slightly biangulate behind, moderately solid, with two or three posterior ridges, the lower the higher; beaks full, sculpture not observed but probably zigzag radial; the posterior slope, and often part of the disk, covered with faint, granular radiating sculpture; epidermis smooth, olive; one compressed, short, recurved pseudocardinal in the left valve under the beak, with a very long one in front of it, and two laterals; two compressed pseudocardinals in the right valve, and a single lateral; all the teeth straight or nearly so; anterior muscle scars separate; beak cavities rather shallow; only one or two dorsal scars in each valve; nacre white or reddish.

Animal unknown.

† RECTIDENS SUMATRENSIS Dunker.

* *Unio sumatrensis* DUNKER, Zeits. für Mal., IX, 1852, p. 52.—* PFEIFFER, Nov. Conch., II, 1866, p. 152, pl. XXXIX, figs. 10-12.—* VON MARTENS, Mal. Bl., XIV, 1866, p. 15.—* PÆTEL, Conch. Sam., III, 1890, p. 169.

Lake Dana Ivar, Sumatra.

¹Lea's name was preoccupied by Kuster, or Charpentier, for a Mexican species in 1856. The name proposed by Issel in 1874 will have to be used.

²Proposed by Fischer and Crosse, because Lea's name *plicatulus* was preoccupied by Kuster.

† RECTIDENS LINGULATUS Drouet and Chaper.

* *Unio lingulatus* DROUET and CHAPER, Mém. Soc. Zool. Fr., V, 1892, p. 148, pl. v, figs. 7-9.—* DROUET, Jl. de Conch, XLI, 1893, p. 42.

Borneo.

† RECTIDENS PROLONGATUS Drouet.

* *Unio prolongatus* DROUET, Rev. Biol. Nord Fr., VI, 1894, p. 216; * Jl. de Conch., XLIII, 1895, p. 36.—* KOBELT, Abh. Senck. Nat. Ges., XXIV, 1897, p. 87, pl. XI, figs. 1, 2.

Borneo.

RECTIDENS PERAKENSIS de Morgan.

* *Unio perakensis* DE MORGAN, Bull. Soc. Zool. Fr., X, 1885, p. 424, pl. ix, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 163.

Perak.

RECTIDENS PAHANGENSIS Smith.

* *Unio pahangensis* E. A. SMITH, Pr. Mal. Soc. Lond., III, 1899, p. 315, fig.

Pahang River, Malay Peninsula. Exceedingly near to *O. lingulatus* Drouet and Chaper.

Genus LAMELLIDENS Simpson, 1900.

(Type, *Unio marginalis* Lamarck.)

Shell elongate, elliptical, pointed behind, with a slight post-dorsal wing, a low, often biangulate posterior ridge, and generally two sharp, radiating liræ above it; beaks with curved, radiating ridges which sometimes are slightly zigzag and often become almost concentric, but which fade out as they approach each other at the center of the disk; epidermis smooth, generally shining, brownish, often with concentric bands of lighter color, rayless or nearly so; left valve with two compressed pseudocardinals, the front one roughened, and two laterals; right valve with two parallel, lamellar pseudocardinals and one lateral; cavity of the beaks rather shallow; dorsal scars few, deep, distinct, scattered; anterior scars separate; nacre bluish white to straw.

Animal unknown.

Subgenus LAMELLEDENS Simpson, 1900.

(Type, *Unio marginalis* Lamarck.)

Shell dark colored; teeth well developed.

† LAMELLIDENS MARGINALIS Lamarck.

* *Die breite Mahler-Muschel aus Grönland*, SCHRÖTER, Flussc., 1779, p. 181, pl. IX, fig. 1.

* ? *Unio groenlandicus* MÖRCH, Am. Jl. Conch., IV, 1868, p. 38.¹

* ? *Unio testudinarius* SPENGLER, Skriv. Selsk. Nat., III, 1793, p. 65.—* PÆTEL, Conch. Sam., III, 1890, p. 169.

¹ The name *groenlandicus* can not stand for this species, since Schröter did not use his appellation in anything but a descriptive sense. The species is found only in India, no *Naiad* being known from Greenland.

- * ? *Unio truncatus* SPENGLER, Skriv. Selsk. Nat., III, 1793, p. 65.—? SCHRÖTER, Arch. Zool. Zoot. 1802, II, 2 Stück, p. 107.
- * *Unio marginalis* LAMARCK, An. sans Vert., VI, 1819, p. 79.¹—* DESHAYES, Enc. Méth. II, 1827, p. 151, pl. CCXLVII, fig. 1.—* HANLEY, Test. Moll., 1842, p. 206; * Biv. Shells, 1843, p. 206, pl. xx, fig. 53.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* KUSTER, Conch. Cab. Unio, 1861, p. 239, pl. LXXX, fig. 4.—* SOWERBY, Conch. Icon., XVI, 1867, pl. LIX, fig. 297.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 20, pl. XLIII, fig. 2.—* PETEL, Conch. Sam., III, 1890, p. 158.
- * *Margarita (Unio) marginalis* LEA, Syn., 1836, p. 37; 1838, p. 24.
- * *Margaron (Unio) marginalis* LEA, Syn., 1852, p. 38; 1870, p. 60.
- * *Unio anodontina* LAMARCK, An. sans Vert., VI, 1819, p. 80.—* DESHAYES, An. sans Vert., VI, 1835, p. 546; II, 1839, p. 671.
- * *Unio anodontinus* KUSTER, Conch. Cab. Unio, 1861, p. 240, pl. LXXX, fig. 15.—* PETEL, Conch. Sam., III, 1890, p. 144.
- *† *Symphynota bilineata* LEA, Tr. Am. Phil. Soc., IV, 1831, p. 98, pl. XI, fig. 19; * Obs., I, 1834, p. 108, pl. XI, fig. 19.
- * *Margarita (Unio) bilineatus* LEA, Syn., 1836, p. 38; 1838, p. 25.
- * *Unio bilineatus* HANLEY, Test. Moll., 1842, p. 207; * Biv. Shells, 1843, p. 207, pl. XXI, fig. 30.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXI, fig. 365.
- * *Margaron (Unio) bilineatus* LEA, Syn., 1852, p. 38; 1870, p. 61.
- * ? *Unio evanescens* MOUSSON, Moll. Java, 1849, p. 91, pl. XVII, fig. 2.

† LAMELLIDENS MARGINALIS var. OBESUS Hanley and Theobald.

- * *Unio marginalis* var. *obesa* HANLEY and THEOBALD, Conch. Ind., 1876, p. 20, pl. XLIII, fig. 3.
- * ? *Unio corrianus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXVII, fig. 401.

† LAMELLIDENS MARGINALIS var. TRICOLOR Kuster.

- * *Unio tricolor* KUSTER, Conch. Cab. Unio, 1856, p. 156, pl. XLV, fig. 1.—* JICKELI, Faun. Moll. N. O. Af., 1874, p. 275.—* PETEL, Conch. Sam., III, 1890, p. 170.
- * *Unio marginalis* var. *tricolor* HANLEY and THEOBALD, Conch. Ind., 1876, p. 20, pl. XLII, fig. 5.

LAMELLIDENS MARGINALIS var. CANDAHARICUS Hanley and Theobald.

- * *Unio marginalis* var. *candaharica* HANLEY and THEOBALD, Conch. Ind., 1870, p. 20, pl. XLIII, fig. 4.

† LAMELLIDENS MARGINALIS var. CYLINDRICUS Hanley and Theobald.

- * *Unio marginalis* var. *cylindrica* HANLEY and THEOBALD, Conch. Ind., 1876, p. 20, pl. XLIV, fig. 1.

LAMELLIDENS MARGINALIS var. HANLEYI Simpson.

- * *Unio marginalis* var. *corriana* HANLEY and THEOBALD, Conch. Ind., 1876, p. 20, pl. XLIV, fig. 4.²

India; Burma; Pegu; Ceylon; Canton River, China?. I believe that Mousson's *U. evanescens* is *L. marginalis*, but can it be from Java?

¹ I use Lamarck's name for this species because the *U. testudinarius* and *truncatus* were only briefly and imperfectly described, and never figured, their habitats being given as Greenland. Lamarck refers to the figures in Enc. Méth., pl. 247, figs. 1, 1a, 1b, 1c, which very accurately represent the shell we know as *Unio marginalis*.

² Not Lea's *corrianus* at all, but very different. I change the name to *hanleyi*.

† LAMELLIDENS PULCHER Tapperone-Canefri.

* *Unio pulcher* TAPPERONE-CANEFRI, Ann. Mus. Civ., VII, 1889, p. 350.¹

Houngdaran, Burma.

LAMELLIDENS EXANTHEMATICUS Kuster.

* *Unio exanthematicus* KUSTER, Conch. Cab. Unio, 1861, p. 243, pl. LXXXI, fig. 2.—

* PÆTEL, Conch. Sam., III, 1890, p. 152.

East Indies.

† LAMELLIDENS THWAITESII Lea.

Unio thwaitesii LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 152; * JI. Ac. N. Sci. Phila., IV, 1860, p. 246, pl. XXXVII, fig. 125; * Obs., VII, 1860, p. 64, pl. XXXVII, fig. 125.—* REEVE, Conch. Icon., XVI, 1865, pl. XXIII, fig. 105.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 20, pl. XLII, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 169.

* Margaron (*Unio*) *thwaitesii* LEA, Syn., 1870, p. 41.

* *Unio consobrinus* HANLEY and THEOBALD, Conch. Ind., 1876, p. 19, pl. XLI, fig. 7.

Ceylon.

† LAMELLIDENS CONSOBRINUS Lea.

* *Unio consobrinus* LEA,² Pr. Ac. N. Sci. Phila., III, 1859, p. 331; * JI. Ac. N. Sci. Phila., 1860, p. 272, pl. XLV, fig. 152; * Obs., VII, 1860, p. 90, pl. XLV, fig. 152.—

* PÆTEL, Conch. Sam., III, 1890, p. 148.

* Margaron (*Unio*) *consobrinus* LEA, Syn., 1870, p. 46.

* *Unio corbeti* DESCHAMPS, Bull. Soc. Zool. Fr., XVII, 1892, p. 68, fig.³

China; India; Ceylon.

† LAMELLIDENS LAMELLATUS Lea.

* *Unio lamellatus* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 19, pl. VI, fig. 16; * Obs., II, 1838, p. 19, pl. VI, fig. 16.—* TROSCHEL, Arch. für. Nat., V, 1839, Pt. 2, p. 234.—* HANLEY, Test. Moll., 1842, p. 194; Biv. Shells, 1843, p. 194, pl. XXI, fig. 49.—* CATLOW and REEVE, Conch. Nom., 1845, p. 60.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* CHENU, Ill. Conch., 1858, pl. XXII, figs. 3, 3a, 3b.—* SOWERBY, Conch. Icon., XVI, 1866, pl. LI, fig. 272.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 21, pl. XLIV, fig. 7.—* PÆTEL, Conch. Sam., III, 1890, p. 126.

* *Margarita (Unio) lamellatus* LEA, Syn., 1836, p. 26; 1838, p. 20.

* Margaron (*Unio*) *lamellatus* LEA, Syn., 1852, p. 29; 1870, p. 46.

* † *Unio layardii* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 153; * JI. Ac. N. Sci. Phila., IV, 1860, p. 243, pl. XXXVI, fig. 122; * Obs., VII, 1860, p. 61, pl. XXXVI, fig. 122.—* REEVE, Conch. Icon., XVI, 1865, pl. XXIII, fig. 111.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 19, pl. XLI, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 157.

* Margaron (*Unio*) *layardii* LEA, Syn., 1870, p. 46.

India; Ceylon.

¹ Only a Latin description is given. We have two shells in the U. S. National Museum from the type lot, and it is, perhaps, a valid species, close to *marginalis*.

² Possibly only a variety of *thwaitesii*.

³ *Unio mainwaringi* Nevill. Shells have been several times received bearing this name, which seem to me to be only *consobrinus*. I can not find that Nevill ever described the species.

LAMELLIDENS SCUTUM Sowerby.

- * *Unio scutum* SOWERBY, Conch. Icon., XVI, 1868, pl. xciv, fig. 510.¹—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 22, pl. xlvi, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 167.

Tenasserim.

† LAMELLIDENS CORRIANUS Lea.

- * *Unio corrianus* LEA, Tr. Am. Phil. Soc., VI, 1834, p. 65, pl. ix, fig. 25; * Obs., I, 1834, p. 177, pl. ix, fig. 25.—* HANLEY, Test. Moll., 1842, p. 207; * Biv. Shells, 1843, p. 207, pl. xa, fig. 60.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* PÆTEL, Conch. Sam., III, 1890, p. 149.
- * *Margarita (Unio) corrianus* LEA, Syn., 1836, p. 38; 1838, p. 25.
- * *Margarita (Unio) corrianus* LEA, Syn., 1852, p. 38; 1870, p. 61.

India; Burma; Pegu.

† LAMELLIDENS GENEROSUS Gould.

- * *Unio generosus* GOULD, Pr. Bost. S. N. Hist., II, 1847, p. 220; * Otia Conch., 1862, p. 210.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 22, pl. xlvi, fig. 4.—* PÆTEL, Conch. Sam., III, 1890, p. 153.
- * *Margarita (Unio) generosus* LEA, Syn., 1870, p. 29.
- * ? *Unio lamellatus* SOWERBY, Conch. Icon., XVI, 1868, pl. xciv, fig. 511.

Burma; Pegu.

† LAMELLIDENS CANEFRIANUS Simpson.

- * *Unio protensus* TAPPERONE-CANEFRI,² Am. Mus. Civ. Gen., VII, 1889, p. 349.

Prome, Lower Burma.

† LAMELLIDENS JENKINSIANUS Benson.³

- * *Unio jenkinsianus* BENSON, Ann. and Mag., X, 1862, p. 185.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 19, pl. xli, fig. 4.—* PÆTEL, Conch. Sam., III, 1890, p. 155.

Assam.

Subgenus SPATHOPSIS, Simpson. 1900.

(Type, *Anodonta guillaini* Recluz.)

Shell long, elliptical, compressed, beaks rather low, said to bear concentrically roughened ridges, with slight plications in front and behind them; posterior ridge not developed; surface shining, smooth; hinge line narrow, with faint, greatly elongated pseudocardinals and laterals;

¹ Credited by Sowerby to Benson.

² Preoccupied by Lea in 1865 for a North Carolina *Unio*, hence the name is changed as above.

³ This is figured by Hanley and Theobald as a heavy shell, shaped much like an old *Unio buckleyi*, and quite different from the other species of the genus. But there is in the U. S. National Museum collection a specimen which is much thinner than the figured shell, and more nearly approaching *marginalis* in form.

esentcheon deep, triangular; beak cavities shallow; anterior muscle scars elongated; posterior faint; iridescent behind.¹

Animal unknown.

LAMELLIDENS GUILLAINI Recluz.

- * *Anodonta guillaini* RECLUZ, JI. de Conch., I, 1850, p. 55; * Zeits. für Mal., VII, 1851, p. 140.—* CROSSE, JI. de Conch., XXXI, 1883, p. 222, pl. IX, fig. 4.—* PÆTEL, Conch. Sam., III, 1890, p. 180.

Brava, northeast Africa, in the Somali country.

Genus TRAPEZOIDEUS Simpson.

(Type, *Unio foliacea* Gould.)

Shell trapezoid, much compressed, with a low posterior ridge, and but slightly raised though pointed beaks, which seem to be sculptured with irregularly radial ridges arranged in two imperfect chevron-shaped loops; surface concentrically sculptured, often having irregular radial ridges on the posterior slope; epidermis yellowish-green or brownish, with two or more green rays above the posterior ridge; teeth compressed; two pseudocardinals in the left valve, one under the beak, the other in front, often not well separated, and two laterals; right valve with two pseudocardinals and one lateral; pseudocardinals all irregular, often pitted and peculiarly dentellate; laterals granular, showing traces of vertical striation; cavity of the beaks rather deep, compressed; muscle scars shallow, anterior scars separate; nacre soft, creamy, yellowish in the beak cavities.

Animal unknown.

† TRAPEZOIDEUS FOLIACEUS Gould.

- * *Unio foliacea* GOULD, Pr. Bost. Soc. N. Hist., I, 1843, p. 141; * Otia Conch., 1862, p. 191.
 * *Unio foliaceus* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 19, pl. XLII, fig. 3.—* ANDERSON, Yunnan Exp., 1877, p. 900, pl. LXXX, figs. 8-12.—* PÆTEL, Conch. Sam., III, 1890, p. 153.
 * *Margaron (Unio) foliaceus* LEA, Syn., 1852, p. 39; 1870, p. 62.
 * † *Unio peguensis* ANTHONY, Am. JI. Conch., I, 1865, p. 351, pl. XXV, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1868, pl. xcv, fig. 519.—* PÆTEL, Conch. Sam., III, 1890, p. 162.
 * *Margaron (Unio) peguensis* LEA, Syn., 1870, p. 51.

TRAPEZOIDEUS FOLIACEUS var. COMPTUS Deshayes and Jullien.

- * *Unio comptus* DESHAYES and JULLIEN, Nouv. Arch. de Mus., X, 1874, p. 126, pl. VI, figs. 3, 4.
 * *Unio fragiles* NEVILL, JI. As. Soc. Beng., XLVI, 1877, p. 39.—* ANDERSON, Yunnan Exp., 1877, p. 400, pl. LXXX, figs. 8-12.

Burma; Cambodia.

¹ I hardly know where to place this, as the description is lacking in several essential characters. It is probable that the beak sculpture is much like that of *Lamellidens* instead of being properly concentric, and it may be a subgenus of that group.

† **TRAPEZOIDENS MISELLUS** Morelet.

- * *Unio misellus* MORELET, JI. de Conch., XIII, 1865, p. 21; * Ser. Conch., IV, 1875, p. 341, pl. XIV, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 159.
 *† *Unio siamensis* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 133; * JI. Ac. N. Sci. Phila., VI, 1868, p. 279, pl. XXXVIII, fig. 93; * Obs., XII, 1869, p. 39, pl. XXXVIII, fig. 93.
 * *Margaron (Unio) siamensis* LEA, Syn., 1870, p. 57.

TRAPEZOIDEUS PALLEGOIXI Sowerby.

- * *Anodon pallegoixi* SOWERBY,¹ Conch. Icon., XVII, 1867, pl. XVII, fig. 17.
 * *Anodonta pallegoixi* CLESSIN, Conch. Cab. Ano., 1876, p. 210, pl. LXIV, fig. 6.—
 * PÆTEL, Conch. Sam., III, 1890, p. 183.

Siam; Cambodia.

TRAPEZOIDEUS PENINSULARIS Simpson.

- * *Unio sumatrensis* SOWERBY,² Conch. Icon., XVI, 1865, pl. XXVIII, fig. 142.

Sumatra.

TRAPEZOIDEUS EXOLESCENS Gould.³

- * *Unio erolescens* GOULD, Pr. Bost. S. N. Hist., I, 1843, p. 141.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 43, pl. CVII, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 152.
 * *Margaron (Unio) erolescens* LEA, Syn., 1852, p. 32; 1870, p. 51.

Tavoy, Burma.

TRAPEZOIDEUS THECA Benson.

- * *Unio theca* BENSON, Ann. and Mag., X, 1862, p. 186.—* HANLEY and THEOBALD, Conch. Ind., 1876, p. 6, pl. XII, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 169.

Cane River, Bundelkhund, India.

Genus **ARCONAIA** Conrad, 1863.

(Type, *Triquetra lanceolata* Lea.)

- * *Arconaia* CONRAD, Am. JI. Conch., I, 1865, p. 234.

Shell greatly elongated, inflated, solid, twisted on its axis, sometimes straight, but generally having the posterior end curved strongly to the right or left, with a decided posterior ridge on both sides of the shell, ending in a blunt point, with usually a small ridge above on the post slope; beaks rather low, sculpture not seen, but probably zigzag radial;

¹ Sowerby says nothing about teeth in his shell, which is from Siam. The teeth of Lea's specimen of *U. siamensis* are scarcely developed, and a comparison of his shell with Sowerby's figure convinces me that they are very nearly allied. Lea's figure shows the shell biangulate behind, which is an error.

² This is not Lea's *sumatrensis*, nor that of Dunker. I have therefore changed the name to *peninsularis*. I am not certain as to its relations. It is not *U. comptus* Deshayes and Jullien, as Fischer believes.

³ Gould did not figure this, and I have not seen the type. The shell figured as this by Hanley and Theobald in the Conchologia probably belongs here.

often a few nodulous or wavy ridges cross the body of the shell, which is irregularly sulcate; epidermis rayless; two pseudocardinals in the right valve, the upper compressed, separated from the split-up lower one by a parallel-sided pit, two in the left, and two heavy laterals; one lateral in the right and a vestige of one below it; laterals granulated and vertically striate; muscle scars deep, anterior small, the front and upper united, the hinder separate, posterior muscle scars long, oblique; beak cavities very shallow; dorsal scars numerous, placed just on the inner edge of the hinge; nacre white. The lobes of the mantle are not united into siphons, but separated throughout, with only a feeble commissure separating the anal and branchial openings.

(Group of *Arconaia lanceolata*.)

Shell with an anterior wing.

† ARCONAIA LANCEOLATA Lea.

* *Triquetra lanceolata* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 79.

* *Hyria lanceolata* LEA, Pr. Ac. N. Sci. Phila., 1856, p. 300.

* *Arconaia lanceolata* CONRAD, Am. Jl. Conch., I, 1865, p. 234.

* *Triquetra contorta* LEA,¹ Pr. Ac. N. Sci. Phila., VIII, 1856, p. 300; * Obs., VI, 1857, p. 39, pl. XXXIII, fig. 33; * Jl. Ac. N. Sci. Phila., III, 1858, p. 319, pl. XXXIII, fig. 33.

* *Hyria contorta* SOWERBY, Conch. Icon., XVII, 1869, pl. I, figs. 2 a, 2 b.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

* *Margaron (Triquetra) contorta* LEA, Syn., 1870, p. 26.

* *Unio contortus* HEUDE, Conch. Fluv. Nank., II, 1877, pl. xv, fig. 31.

China.

† ARCONAIA MUTICA Heude

* *Unio contortus* var. *muticus* HEUDE, Conch. Fluv. Nank., II, 1877, pl. xv, fig. 32.

China.

(Group of *Arconaia conjungens*.)

Shell moderately solid, slightly twisted and bent, rounded in front; pseudocardinals not so much torn as in the *lanceolata* group; not winged in front.

ARCONAIA CONJUNGENS Heude.

* *Unio contortus* var. *conjungens* HEUDE, Conch. Fluv. Nank., II, 1877, pl. xv, fig. 33.

China.

Genus PSEUDAVICULA Simpson, 1900.

(Type, *Unio johnstoni* Smith.)

Shell rather solid, inflated, with a well-developed anterior and posterior dorsal wing; dorsal line strongly incurved; beaks full, not high; a

¹ The specific name was changed to *contorta* by Lea, because he thought his earlier name, *lanceolata*, was less appropriate.

high, down-curved posterior ridge runs to the posterior basal point, and between this point and the posterior point of the wing the outline is deeply incurved; base and lower part of anterior end rounded; surface slightly sculptured with concentric ridges; beak sculpture probably zigzag rayed; anterior tooth of left valve elongate, slightly corrugated, that of the right double; laterals in left valve double, single in the right, straight, elongated, thin, and prominent; nacre, rose tinted.

Animal unknown.

PSEUDAVICULA JOHNSTONI Smith.

Unio (metaptera) johnstoni SMITH, Pr. Zool. Soc. Lond., 1893, p. 640, pl. LIX, figs. 18-20.¹

Lake Mweru, British Central Africa.

Genus **ARCIDOPSIS** Simpson, 1900.

(Type, *Unio footei* Theobald.)

Shell inequilateral, elongated, with dorsal and ventral lines nearly straight and parallel; anterior end strongly and obliquely truncated from the beaks to the anterior base, and ending in a rather sharp but rounded point below; posterior end obliquely truncated above, rounded below; posterior ridge full, rounded; the shell in front and below rather compressed; surface concentrically and radially ridged; pseudocardinals strong, two in the right valve, one in the left, and supported by a strong, cardinal rib; muscle scars well impressed; parallel line distinct. Animal unknown.²

ARCIDOPSIS FOOTEI Theobald.

* *Unio footei* THEOBALD, Jl. As. Soc. Beng., XLV, 1876, p. 187, pl. XIV, figs. 9, 9a.

Kistna River, India.

The following are indeterminate Oriental Unionidæ:

Unio acrorrhynchus VON MARTENS, S. B. Nat Fr., 1894, p. 214. Korea.

Unio alfierianus BOURGUIGNAT, Voy. Choa, 1885, p. 43. Choa, Central Africa.

Unio antiniloticus BOURGUIGNAT, Bull. Soc. Zool. Fr., XI, 1880, p. 482, pl. XII, fig. 5. High Senegal.

Unio ascia HANLEY, Biv. Shells, 1856, p. 385.

Unio bridouxi BOURGUIGNAT, Un. & Ir. Tan., 1886, p. 13. Lake Tanganyika.

Unio cyamus PHILIPPI, Zeits für Mal., VIII, 1852, p. 125. S. Africa.

Unio despitieri MABILLE, Bull. Soc. Mal. Fr., IV, 1887, p. 162. Tonkin.

Unio dolichorhynchus TAPPERONE CANEFRI, Ann. Mus. Civ. Gen., 1889, p. 348. Mercato de Mandelay.

¹ A remarkable *Naiad*, closely imitating some of the South American *Prisodons* in general appearance, to which it may be nearly allied.

² Unfortunately Theobald's Latin description of *Unio footei* is not at all complete. No laterals are mentioned, and he says nothing of the color of the epidermis or of the nacre. The beaks were too much worn in his specimens to give any characters. The shell resembles some of the *Areas* of the *Barbatia* group, and may not belong to the *Unionidæ* at all.

- Unio forscali* PARREYSS. Where described? Egypt.
Unio gianelli TAPPERONE CANEFRI, Ann. Mus. Civ. Gen., 1889, p. 353. Mercato de Mandelay.
Unio gibbus SPENGLER, Skriv. Selsk. Nat., III, 1793, p. 64. Tranquebar.
Unio gladiator ANCEY, Le Nat., III, 1881, p. 468. Tonkin.
Unio gottschei VON MARTENS, S. B. Nat. Fr., 1894, p. 215. Korea.
Unio grantianus BOURGUIGNAT, Moll. Ny. Ouk., 1883, p. 14. Lake Onkerewe, Central Africa.
Unio guillemeti BOURGUIGNAT, Un. & Ir. Tan., 1886, p. 15. Lake Tanganyika.
Unio hagnei STRUBELL, Nach. Mal. Ges., 1897, p. 10. Sumatra.
Unio hamyanus BOURGUIGNAT, Voy. Choa., 1885, p. 42. Central Ethiopia.
Unio idgi BOURGUIGNAT Moll. Terr. et Fluv. Choa, 1885, p. 39. Choa, Africa.
Unio ilqui SOLEILET, Voy. Choa, 1885, p. 40. Central Ethiopia.
Unio inaequalis ROCHEBRUNE, Bull. Soc. Philom., 1882, p. 44. Mekong.
Unio jaculus ROCHEBRUNE, Bull. Soc. Philom., 1882, p. 44. Mekong.
Unio josseti BOURGUIGNAT, Un. & Ir. Tan., 1886, p. 19. Lake Tanganyika.
Unio jouberti BOURGUIGNAT, Un. & Ir. Tan., 1886, p. 8. Lake Tanganyika.
Unio larigerinus BOURGUIGNAT, Un. & Ir. Tan., 1886, p. 14. Lake Tanganyika.
Unio ligula MOUSSON, L. & S. W. Moll. Java, 1849, p. 94. Java.
Unio longitudinalis ANTON, Verz der Conch. 1839, p. ?
Diplodon ludovicianum ROCHEBRUNE, Bull. Soc. Phil., 1882, p. 43. Mekong.
Unio madagascariensis SGANZIN, Mem. Soc. Hist. Strab., 1846, p. 8. Mahoupa River, Madagascar. No figure.
Unio menardi BOURGUIGNAT, Un. & Ir. Tan., 1886, p. 20. Lake Tanganyika.
Unio meneliki SOLISET, Voy. Choa, 1885, p. 41. Central Ethiopia.
Unio moinei BOURGUIGNAT, Un. & Ir. Tan., 1886, p. 11. Lake Tanganyika.
Unio palembangensis STRUBELL, Nach. Mal. Ges., 1887, p. 10. Sumatra.
Dynomia patonina ROCHEBRUNE, Bull. Soc. Phil., 1882, p. 43. Cochin China.
Unio pliculosus VON MARTENS, S. B. Nat. Fr., 1894, p. 216. Korea.
Unio schweinfurthi VON MARTENS, S. B. Nat. Fr., 1886, p. 127. Egypt.
Unio sitifensis MORELET, Jl. de Conch., 1853, p. 298.
Unio soleilleti BOURGUIGNAT, Voy. Choa, 1885, p. 39. Central Ethiopia.
Loncosilla solenoides RAFINESQUE, Cont. Mon., 1831, p. 7. Jellinghy River, Bengal.
Unio subamgydalinus DROUET, Jl. de Conch., 1895, p. 33. West Africa.
Unio truncatus SPENGLER, Skiv. Selsk. Nat., III, 1793, p. 66. Tranquebar.
Unio verrucifer VON MARTENS, S. B. Nat. Fr., 1894, p. 216. Korea.
Unio vinckeii BOURGUIGNAT, Un. & Ir. Tan., 1886, p. 18. Lake Tanganyika.
Unio visseri BOURGUIGNAT, Un. & Ir. Tan., 1886, p. 21. Lake Tanganyika.
Zaira elegans ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 12.
Zaira poirieri ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 11.
Zaira araneosa ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 11.
Zaira sordida ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 13.
Zaira disciformis ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 10.

The last five from the Kongo. I know nothing whatever of the genus *Zaira*. It may be merely a new name for the section *Calatura* of the genus *Nodularia*. It was proposed by Rochebrune.¹

LAMPHORHAMPHUS.

Male and female shells alike, dull-colored; beak sculpture nearly or quite radial, marsupium occupying the entire inner branchiæ, forming a pad-like mass.

¹ Bull. Soc. Mal. Fr., 1886, p. 1.

Genus TETRAPLODON Spix, 1827.

Castalia LAMARCK, An. sans Vert., VI, 1819, p. 66.*Tetraplodon* SPIX, Test. Fluv. Bras., 1827, p. 32.*Prisodon* LEA, Synopsis, 1852, p. 18.

Shell triangular, solid, inflated, with a high, sharp, posterior ridge, behind which it is decidedly truncated; beaks very full and high, sculpture strictly radial, or with one or two pairs of the central bars coalescing below, the whole extending more or less over the disk as strong ridges; epidermis thick, dark, dull colored; hinge line arched; there is a very strong, compressed pseudocardinal in the left valve, and there are two in the right, all in front of the beaks; behind them are several denticles; one lateral in the right valve and two in the left; all the teeth strongly vertically ridged; beak cavities deep, not compressed; anterior muscle scars small, deep; nacre whitish.

Animal having the labial palpi wider than long; margins of mantle inferiorly open, but united posteriorly to form two siphons, branchial and anal; branchial opening surrounded with papillæ; anal opening smooth; outer branchiæ united to the mantle to their posterior ends; inner united to the abdominal sac; foot tongue-shaped, thick, produced anteriorly. (Troschel.)

(Type, *Tetraplodon pectinatum* Spix.)¹(Group of *Tetraplodon ambiguus* characters, as in the genus.)†TETRAPLODON AMBIGUUS (Lamarck) Sowerby.²*? *Prisodon truncatus* SCHUMACKER, Ess. Nouv. Syst., 1817, p. 139.*? *Castalia ambigua* LAMARCK, An. sans Vert., VI, 1819, p. 67; GUERIN, Icon. Regne. An., 1829, pl. XXVIII.—* WYATT, Man. Conch., 1838, p. 65, pl. XI, fig. 5.—* CHENU, Man., 1859, II, p. 149, fig. 735.—* SOWERBY, Conch. Icon., XVII, 1869, pl. I, fig. 1.* *Unio ambigua* DESHAYES, Tr. Elem. Conch., 1839, p. 18, pl. XXIX, figs. 7-9.* *Unio ambiguus* SOWERBY, Rec. and Fos. Shells, XVI, 1823, fig. d.—* REEVE, Conch. Syst., 1841, p. 117, pl. LXXVII, fig. 2.—* CATLOW and REEVE, Conch. Nonl., 1845, p. 55.—* DESHAYES, Tr. Elem., II, 1853, p. 219, pl. XXIX, figs. 7-9.—* KUSTER, Conch. Cab. Unio, 1856, p. 165, pl. XLVIII, fig. 1.* *Tetraplodon pectinatum* SPIX, Test. Fluv. Bras., 1827, p. 32, pl. XXV, figs. 3-4.*† *Castalia inflata* D'ORBIGNY, Guer. Mag., 1835, p. 43.—* HANLEY, Test. Moll., 1842, p. 173; * Biv. Shells, 1843, p. 173, pl. XXIV, fig. 12.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 598, pl. LXXII, figs. 4-10.*† *Castalia turgida* HUPE, Moll. Nouv., III, 1857, p. 76, pl. XIV, fig. 1.—* SOWERBY, Conch. Icon., XVII, 1869, pl. III, fig. 11.

¹ The name *Castalia* applied by Lamarck to this genus in 1819 can not stand, as it was used in Verme in 1817. I have every reason to suppose that the young in this and all the South American genera of *Unionidae* are carried in the inner gills.

² Schumacher gave only a brief description of his *Prisodon truncatus*, and referred to no figure, and the same can be said of Lamarck regarding his *Castalia ambigua*, both of which are supposed by authors to be the above species. But either of these descriptions apply equally well to several species, therefore they can not be considered authentic.

- *† *Castalia hanleyana* SOWERBY, Conch. Icon., XVII, 1869, pl. I, fig. 5.—* P.ETEL, Conch. Sam., III, 1890, p. 189.
 * *Castalia carolinensis* SOWERBY, Conch. Icon., XVII, 1869, pl. II, fig. 6.—
 * P.ETEL, Conch. Sam., III, 1890, p. 189.
 * *Castalia cordata* STROBEL, Mat. Mal., Pt. I, 1874, p. 75.

Amazon and Rio de la Plata River systems.

† TETRAPLODON QUADRILATERUM d'Orbigny.

- * *Castalia quadrilatera* D'ORBIGNY, Guer. Mag., No. 62, 1835, p. 42.—* HANLEY, Test. Moll., 1842, p. 173; * Biv. Shells, 1843, p. 173.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 599, pl. LXXIII.—* TROSCHEL, Arch. für Nat., XIII, 1847, Pt. 1, p. 222.—* VON IHERING, Arch. für Nat., 1893, p. 89.
 * *Mya ambigua* WOOD, Ind. Test. Rev., 1856, p. 200, pl. I (supp.), fig. 9.
 * *Castalia acuticosta* HUPE, Moll. Nouv., III, 1857, p. 77, pl. XIV, fig. 3.—* SOWERBY, Conch. Icon., XVII, 1869, pl. III, figs. 12, 12a, 12b.—* P.ETEL, Conch. Sam., III, 1890, p. 189.
 * *Castalia cordata* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 509, pl. CXX; III, figs. 2, 2a.—* SOWERBY, Conch. Icon., XVII, 1869, pl. II, fig. 8.—* P.ETEL, Conch. Sam., III, 1890, p. 189.
 * *Castalia latiquadrata* SOWERBY, Conch. Icon., XVII, 1869, pl. II, fig. 10.—* P.ETEL, Conch. Sam., III, 1890, p. 189.

Tropical South America east of the Andes. Perhaps a form of *ambiguus*.

† TETRAPLODON OVATUS Sowerby.

- * *Castalia ovata* SOWERBY, Conch. Icon., XVII, 1869, pl. I, fig. 4.—* P.ETEL, Conch. Sam., III, 1890, p. 189.

Brazil.

† TETRAPLODON RETUSUS Hupe.

- * *Castalia ambigua* SOWERBY, Conch. Man., 1839, fig. 140.
 * *Castalia retusa* HUPE, Moll. Nouv., III, 1857, p. 75, pl. XIV, fig. 2.—* SOWERBY, Conch. Icon., XVII, 1869, pl. I, fig. 2.—* P.ETEL, Conch. Sam., III, 1890, p. 189.

Guiana.

TETRAPLODON QUADRATUS Sowerby.

- * *Castalia quadrata* SOWERBY, Conch. Icon., XVII, 1867, pl. II, figs. 7, 7a, 7b.—
 * P.ETEL, Conch. Sam., III, 1890, p. 189.

TETRAPLODON SCHOMBERGIANUS Sowerby.

- * *Castalia schombergiana* SOWERBY, Conch. Icon., XVII, 1869, pl. XIV, fig. 3.—
 * P.ETEL, Conch. Sam., III, 1890, p. 189.

Guiana.

(Group of *Tetraplodon multisulcatus*.)

Shell small, somewhat compressed, with a moderate posterior ridge, and covered throughout with delicate ridges; teeth more Unionoid than in the typical *Tetraplodon*s.

Animal unknown.

TETRAPLODON MULTISULCATUS Hupe.

* *Castalia ambigua* BLAINVILLE, Man., 1825, p. 539, pl. LXVII, fig. 4.

* *Castalia multisulcata* HUPE, Moll. Nouv., III, 1857, p. 75, pl. XIV, fig. 4.—* SOWERBY, Conch. Icon., XVII, 1869, pl. II, figs. 9, 9a, 9b.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

Brazil.

TETRAPLODON ECARINATUS Mousson.

* *Castalia ecarinata* MOUSSON, Mal. Bl., XVI, 1869, p. 185.—* PFEIFFER, Nov. Conch., IV, 1876, p. 140, pl. CXXXI, figs. 9, 10.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

Puerto Nuevo, Magdalena River, Colombia.

TETRAPLODON CROSSEANUS Hidalgo.

* *Castalia crosseana* HIDALGO, Jl. de Conch., XIII, 1865, pp. 316, 429, pl. XIV, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

Imbabura, Ecuador.

Genus CASTALINA Von Ihering, 1891.

Castalina VON IHERING, Zool. Anzeiger, 1891, p. 478.

Shell somewhat triangular, inflated, but having the sides a little flattened, solid, with a strong posterior ridge, the subtruncate posterior slope rising almost to a wing above; beaks full, high, with nearly strictly radial sculpture; surface slightly, irregularly, concentrically sulcate, sometimes a little corrugated; posterior slope generally plicate or corrugated; epidermis thick, rayless, blackish; hinge plate arched, wide; two to several radial pseudocardinals in each valve; two vertically or obliquely striate laterals in the left valve and one in the right; beak cavities deep; anterior muscle scars deep, united; nacre whitish.

Animal, probably very much like that of *Tetraplodon*, but with the mantle closed or open at the branchial and anal openings.

(Type, *Castalina martensi* von Ihering.)

(Group of *Castalina martensi*.)

Characters as in the genus.

†CASTALINA MARTENSI von Ihering.

* *Castalina martensi* VON IHERING, Zool. Anz., XIV, 1891, p. 477; Arch. für Nat., 1893, p. 81, pl. III, fig. 5.

Southern Brazil.

†CASTALINA NEHRINGI von Ihering.

Castalina nehringi VON IHERING, Zool. Anz., XIV, 1891, p. 477; * Arch. für Nat., 1893, p. 75, pl. III, fig. 4.

Southern Brazil.

† *CASTALINA PSAMMOICA* d'Orbigny.

* *Unio psammoica* d'ORBIGNY, Guer. Mag., 1835, p. 35; * Voy. Am. Mer., 1843, p. 608, pl. LXXI, figs. 4-7.—* KUSTER, Conch. Cab. Unio, 1861, p. 263, pl. LXXXVIII, fig. 4.

† *Castalina psammoica* VON IHERING, Arch. für Nat., 1893, p. 79.

* *Margaron (Unio) psammoicus* LEA, Syn., 1852, p. 19; 1870, p. 30.

* *Unio psammoicus* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCIII, fig. 507.—* PÆTEL, Conch. Sam., III, 1890, p. 164.

Tributaries of the Rio de la Plata.

† *CASTALINA UNDOSA* von Martens.

* *Castalia undosa* VON MARTENS, S. B. Nat. Fr., 1885, p. 148; * Conch. Mitth., III, 1885, p. 19, pl. XLII, figs. 2, 3.—PÆTEL, Conch. Sam., III, 1890, p. 190.—VON IHERING, Arch. für Nat., 1893, p. 84.—* NEHRING, Jl. de Conch., 1894, p. 82.

Southern Brazil.

(Group of *Castalina orbignyi*.)

Shell irregularly rhomboid, somewhat compressed, equilateral, anterior end rounded, posterior truncate, having a wide, shallow, radiating furrow above the rounded posterior ridge which causes a sinuosity in the outline; epidermis brownish yellow, hinge crenate; one pseudocardinal in the left valve and two in the right, which are jagged or somewhat broken up; laterals elongate and striate.

CASTALINA ORBIGNYI Hupe and Deville.¹

* *Unio orbignyi* HUPE and DEVILLE, Rev. et Mag., 1850, p. 645, pl. XVI, fig. 1.—

* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.

* *Margaron (Unio) orbignyi* LEA, Syn., 1852, p. 24; 1870, p. 37.

* *Unio orbignyana* HUPE, Moll. Noniv., III, 1857, p. 83, pl. XVII, fig. 1.

* *Unio d'orbignyanus* SOWERBY, Conch. Icon., XVI, 1868, pl. XCVI, fig. 523.

* *Unio orbignyanus* PÆTEL, Conch. Sam., III, 1890, p. 162.

Upper Amazon.

Genus *CASTALIELLA* Simpson, 1900.

(Type, *Castalia sulcata* Krauss.)

Shell subtriangular, greatly inflated, subsolid and strongly sulcate; beaks high, the sculpture regularly radiate; posterior ridge well defined and sharp; epidermis reddish chestnut; hinge line narrow, arched; there are two vertically striate pseudocardinals in the right valve, separated by a parallel-sided socket, the lower the larger and split, and three in the left standing side by side, the middle one the largest, the lower two torn, with a few small tubercles in each valve behind them; one granular

¹I do not know just where to place this species, never having seen it, and the descriptions not being full in essential details. It is probably a *Castalina*. No mention is made of sculpture, but from the figure it seems to be slightly sulcate.

lateral in the right valve and two in the left, which show traces of vertical striation; beak cavities deep, not compressed; nacre purple.¹ Animal unknown.

† *CASTALIELLA SULCATA* Krauss.

* *Castalia sulcata* KRAUSS, Zeits. für Mal., 1849, p. 99.—* PÆTEL, Conch. Sam., III, 1890, p. 190.

* *Unio kraussii* LEA, Pr. Ac. N. Sci. Phila., VI, 1853, p. 376.²

* *Margaron (Unio) kraussii* LEA, Syn., 1870, p. 35.

Surinam.

Genus *CALLONAIA* Simpson, 1900.

(Type, *Castalia duprei* Recluz.)

Shell triangular, thin, inflated, with very high, full beaks which appear to be without sculpture;³ a very high, sharp posterior ridge extends to the base of the shell, above which it is decidedly truncated; anterior end somewhat pointed above, rounded below; ligament very short; surface nearly smooth but somewhat sulcate anteriorly; epidermis bright greenish yellow, shining; hinge line strongly arched; teeth compressed, high; two pseudocardinals in each valve, all interlocking and side by side; one lateral in the right valve and two in the left, a little ragged and granularly vertically striate; beak cavities very deep, not compressed; muscle scars superficial; nacre brilliant, bluish white, radiately striate posteriorly. Animal unknown.

† *CALLONAIA DUPREI* Recluz.

* *Castalia duprei* RECLUZ, Rev. Zool., 1843, p. 305, pl. xxxv; * Mag. Zool., 1843, p. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 509.—* CHENU, Man., 1859, II, p. 149, fig. 738.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

* *Unio duprei* CATLOW and REEVE, Conch. Nom., 1845, p. 58.

* *Margaron (Prisodon) duprei* LEA, Syn., 1852, p. 18; 1870, p. 27.

* *Castalia dolabella* SOWERBY, Conch. Icon., XVII, 1869, pl. III, figs. 13 a, 13 b, 13 c.—

* PÆTEL, Conch. Sam., III, 1890, p. 189.

Great lakes of Para, Brazil.

¹ This species, of which the type is in the Lea collection, has some of the characters of *Prisodon* and *Castalina*, but I can not satisfactorily refer it to either, and it is certainly not a *Diplodon*. Its strong sulcation, reddish chestnut epidermis, and purple nacre are characters that are not possessed by any South American *Naiades* that I am acquainted with.

² Lea received this shell from Dr. Dunker under the name of *Castalia sulcata* Krauss. Believing it to be a *Unio*, he placed it in that genus, and the name *sulcatus* being preoccupied in *Unio*, he changed it as above.

³ In one of Lea's shells the beaks are almost perfect, and not a sign of sculpture can be seen.

Genus HYRIA Lamarck, 1819.

(Type, *Hyria corrugata* Lamarck.)*Hyria* LAMARCK, An. sans Vert., VI, 1819, p. 81.¹*Triplodon* SPIX, Test. Fluv. Bras., 1827, p. 35.*Naia* SWAINSON, Tr. on Mal., 1840, p. 379.

Shell subrhomboidal, solid, slightly inflated, narrower and dorsally winged in front, and having a post-dorsal wing; posterior ridge often double, causing the shell to be biangulate behind; beaks low, with strong, nearly radial sculpture, the central bars coalescing below, the whole continuing as strong, radial, and zigzag ridges over more or less of the disk; epidermis thick, greenish when young, brownish or blackish when old; there are two or more rather short, compressed pseudo-cardinals in each valve, which are much split up into denticles; one lateral in the right and two in the left valve; teeth often somewhat vertically striated; dorsal scars numerous; nacre whitish.

Animal with mantle lobes united together behind and furnished with two short, contractile siphons. (J. E. Gray.)

Subgenus HYRIA Lamarck, 1819.

(Type, *Hyria corrugata* Lamarck.)

Characters as in the genus.

†HYRIA CORRUGATA Lamarck.

* *Hyria corrugata* LAMARCK, An. sans Vert., VI, 1819, p. 82.—* SOWERBY, Rec. and Foss. Shells, 1823, No. XVI, fig. d; * Conch. Man., 1839, fig. 144.—* REEVE, Conch. Syst., I, 1841, p. 120, pl. xc, fig. 2.—* HANLEY, Test. Moll., 1842, p. 215; * Biv. Shells, 1843, p. 215.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* KUSTER, Conch. Cab. Unio, 1856, p. 140, pl. xli, fig. 1.—* CHENU, Man., 1859, II, p. 149, fig. 733.—* REEVE, Elem. Conch., II, 1860, pl. xxxi, fig. 179.—* SOWERBY, Conch. Icon., XVII, 1869, pl. i, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

* † *Unio corrugata* BLAINVILLE, Man., 1825, p. 539, pl. lxxvii, fig. 1.

¹In 1753 Klein (Testamen Methodi, p. 135, pl. 9, fig. 36) used the name *Triquetra* for the shell which has since been known as *Hyria syrmatophora* Gronovius. In 1817 Schumacher founded the genus *Prisodon* (Essai Nouv. Système, p. 138), in which under section *a* he placed his *P. obliquus*, a smooth species considerably drawn out, and in section *b* he put *P. truncatus*, which is possibly the same as the *Castalia ambigua* of Lamarck. These certainly belong to two genera. Dr. Lea in the Synopsis used Klein's name for the winged shell (*Hyria corrugata* Lamarck), and used the name *Prisodon* for Schumacher's section *b* (*P. truncatus* Schumacher). Lea erred in this, because Klein was not a binomial author, and his name can not be used. In the proceedings of the U. S. National Museum, XVIII, 1896, p. 315, I restored the name of *Prisodon* Schumacher to the first section, pointing out that the *P. obliquus* Schumacher must stand as its type. Two years after Schumacher's name appeared Lamarck published the generic name *Hyria* in the Animaux sans Vertèbres, VI, 1819, p. 81, and placed in it first his *H. avicularis*, which equals Schumacher's *P. obliquus*, and secondly *H. corrugata*. I now believe that these corrugated forms are generically distinct from the smooth ones, and as Schumacher's *Paxyodon* is almost certainly founded on one of the smooth species, Lamarck's name *Hyria* can be used by elimination for the *corrugatus* and allied forms.

- * *Margaron (Triquetra) corrugata* LEA, Syn., 1852, p. 17; 1870, p. 25.
 * *Triquetra corrugata* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 508.
 * ? *Triplodon rugosum* SPIX, Test. Fluv. Bras., 1827, p. 35, pl. XXIX, figs. 1, 2.
 * *Hyria rugosa* DESHAYES, Enc. Méth., II, 1827, p. 151, pl. CCXLVII, fig. 2.
 * *Mya angulata* WOOD, Ind. Test. Sup., 1828, p. 3, pl. I, fig. 12.
 * *Margarita (Unio) angulatus* LEA, Syn., 1836, p. 10; 1838, p. 13.
 * *Hyria exasperata* SOWERBY, Conch. Icon., XVII, 1869, pl. II, fig. 3.—*PÆTEL, Conch. Sam., III, 1890, p. 189.

Eastern Peru to Guiana; south throughout Brazil.

†HYRIA RUGOSISSIMA Sowerby.

- * *Hyria rugosissima* SOWERBY, Conch. Icon., XVII, 1869, pl. III, fig. 5.—*PÆTEL, Conch. Sam., III, 1890, p. 189.

Amazon River.

HYRIA LATIALATA Sowerby.

- * *Hyria latialata* SOWERBY, Conch. Icon., XVII, 1869, pl. II, fig. 4.—*PÆTEL, Conch. Sam., III, 1890, p. 189.

Guiana.

†HYRIA TRANSVERSA Hupe.

- * *Hyria transversa* HUPE, Moll. Nouv., III, 1857, p. 79, pl. XV, fig. 1.—*SOWERBY, Conch. Icon., XVII, 1869, pl. IV, fig. 7.—*PÆTEL, Conch. Sam., III, 1890, p. 189.

Brazil.

Subgenus TRIQUETRANA Simpson, 1900.

(Type, *Unio stevensi* Lea.¹)

Shell compressed, with a feebly developed posterior wing, thickened in front; each valve with about three somewhat divergent pseudo-cardinals which are sometimes nearly smooth and at others break into denticles; one granular lateral in the right valve and two in the left which are sometimes obliquely and faintly vertically striate; external sculpture much as in *Hyria*. Animal unknown.

†HYRIA STEVENSI Lea.

- * *Unio stevensi* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 188; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 22, pl. VII, fig. 19; *Obs., XIII, 1874, p. 26, pl. VII, fig. 19.

Yuruari River, Guiana.

Genus PRISODON Schumacher, 1817.

(Type, *Mya syrmatophora* Meuschen in Gronovius.)

- Triquetra* KLEIN, Tent. Methodi, Ost, 1753, p. 135.
Prisodon SCHUMACHER (part), Essai Nouv. Syst., 1817, p. 138.
Paryodon SCHUMACHER, Essai Nouv. Syst., 1817, p. 139.

Shell aviculiform, solid, somewhat inflated, with a well-developed posterior ridge, the area above it being excavated, its outline from the

¹This shell bears much the same relation to *Hyria* that *Castalina* does to *Tetraplodon*, its teeth being more unionoid than those of *Hyria* and the shell more compressed.

hinder basal point to the end of the post dorsal wing generally incurved; posterior slope having a radial row of plications, as in *Cristaria*; surface slightly concentrically sculptured and having delicate radiating *livæ* throughout, so that the surface is microscopically reticulated; beaks full, but not elevated, without sculpture as far as noticed; epidermis yellowish green or brownish, shining, often bronzy; hinge narrow under the beaks, widening in front and behind; left valve with two or more elongated compressed pseudocardinals which show a tendency to break into denticles, and two laterals; right valve with two or more similar pseudocardinals and a single lateral; laterals vertically, granularly striate.

Animal with the labial palpi triangular, the hinder parts free as in *Unio*, not united posteriorly; branchiæ large, equal in size. (Troschel.)

Subgenus **PRISODON** Schumacher, 1817.

(Type, *Mya syrmatophora* Meuschen.)

Characters as in the genus.

† **PRISODON SYRMATOPHORUS** Meuschen in Gronovius.

* *Triquetra subviridis* KLEIN, Tent. Méth. Ost., 1753, p. 135, pl. ix, fig. 36.¹—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 505; III, pl. cxx, figs. 1, 1a.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

* *Margaron (Triquetra) subviridis* LEA, Syn., 1852, p. 17; 1870, p. 25.

* *Mya syrmatophora* MEUSCHEN in Gronovius, Zooph., 1781, pl. xviii, figs. 1, 2.²—

* GMELIN, Syst. Nat., 1788, No. 3222.—* WOOD, Gen. Conch., I, 1815, p. 110.—

* DILLWYN, Cat., I, 1817, p. 54.—* WOOD, Ind. Test., 1825, p. 12, pl. ii, fig. 36b.

* *Hyria syrmatophora* SOWERBY, Rec. and Foss. Shells, XVI, 1823, fig. d; * Conch. Man., 1839, fig. 143.—* REEVE, Conch. Syst., I, 1841, p. 120, pl. xc, fig. 1.—

* HANLEY, Test. Moll., 1842, p. 214; * Biv. Shells, 1843, p. 214.—* CATLOW and REEVE, Conch. Man., 1845, p. 65.—* TROSCHER, Arch. für Nat., XIII,

Pt. I, 1847, p. 271.—* WOOD, Ind. Test. rev., 1856, p. 17, pl. ii, fig. 36.—* HUPE, Moll. Nouv., III, 1857, p. 79, pl. xv, fig. 2.—* SOWERBY, Conch. Icon., XVII, 1869, pl. v, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

* *Unio syrmatophora* DESHAYES, Tr. Elem., II, 1853, p. 219, pl. xxix, figs. 10, 11.

* *Margarita (Unio) syrmatophorus* LEA, Syn., 1836, p. 11; 1838, p. 13.

* † *Parygodou ponderosus* SCHUMACHER, Ess. Nouv. Syst., 1817, p. 140, pl. xi, fig. 3.

* *Hyria avicularis* var. *b.* LAMARCK, An. sans Vert., VI, 1819, p. 82.

Guiana; Brazil.

† **PRISODON COMPLANATUS** Hupe.

* *Hyria complanata* HUPE, Moll. Nouv., III, 1857, p. 80, pl. xv, fig. 3.—* SOWERBY, Conch. Icon., XVII, 1869, pl. iv, fig. 9.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

* *Hyria syrmatophora* KUSTER, Conch. Cab. Unio, 1856, p. 141, pl. xli, fig. 4.

Guiana.

¹ Klein's species evidently equals the *syrmatophora* of Gronovius, but his name can not be used, as he was not a binomial author.

²Gronovius used only generic names in the Zoophylaceum, and died while the work was in progress. Meuschen finished it and figured the above species and described it in the explanation of the plate.

† PRISODON ALATUS Sowerby.¹

- * *Hyria alata* SOWERBY, Conch. Icon., XVII, 1869, pl. v, fig. 13.—PÆTEL, Conch. Sam., III, 1890, p. 189.

Guiana.

PRISODON CASTELNAUDI Hupe.

- * *Hyria castelnaudi* HUPE, Moll. Nouv., III, 1857, p. 81, pl. XVI, fig. 1.—* SOWERBY, Conch. Icon., XVII, 1869, pl. iv, fig. 8.—* PÆTEL, Conch. Sam., III, 1890, p. 189.²

Brazil.

† PRISODON OBLIQUUS Schumacher.

- * *Prisodon obliquus* SCHUMACHER, Ess. Nouv. Syst., 1817, p. 139, pl. XL, fig. 2.³
 * *Hyria obliqua* PÆTEL, Conch. Sam., III, 1890, p. 189.
 * *Hyria avicularis* LAMARCK, An. sans Vert., VI, 1819, p. 82.—* CROUCH, Ill. Int. Lam., 1827, p. 16, pl. IX, figs. 5 a, b.—* STARK, Nat. Hist., II, 1828, p. 89.—* WYATT, Man. Conch., 1838, p. 67, pl. v, fig. 4.—* DELESSERT, Rec. Coq. Lam., 1841, pl. XII, fig. 9.—* HANLEY, Test. Moll., 1842, p. 214; * Biv. Shells, 1843, p. 214, pl. XXIV, fig. 11.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* TROSCHER, Arch. für Nat., XIII, Pt. 1, 1847, p. 271, pl. VI, fig. 3.—* CHENU, Man., 1859, II, p. 149, fig. 734.—* SOWERBY, Conch. Icon., XVII, 1869, pl. III, figs. 6 a, b.
 * *Unio avicularis* DESHAYES, Enc. Méth., II, 1830, p. 583; * Traite Elem. Conch., 1839, p. 18, pl. XXIX, figs. 10, 11.
 * *Hyria avicularia* GUERIN, Icon. Regne An., 1844, II, pl. XXVIII, fig. 8.
 * *Unio caudatus* WAGNER, Test. Fluv. Bras., 1827, p. 35, pl. XXVII, figs. 1, 2.
 * *Diplodon furcatum* SPIX, Test. Fluv. Bras., 1827, p. 35, pl. XXVII, figs. 1, 2.
 * *Hyria elongata* SWAINSON, Ex. Conch., 1841, p. 29, pl. XXIV.
 * *Triquetra elongata* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 508.

Amazon drainage; southward through Brazil.

PRISODON RECTUS Sowerby.⁴

- * *Hyria recta* SOWERBY, Conch. Icon., XVII, 1868, pl. v, fig. 10.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

South America; locality unknown.

† PRISODON BROWNIANUS Lea.

- * *Unio brownianus* LEA, Tr. Am. Phil. Soc., 1838, p. 108, pl. XXIV, fig. 116; * Obs., II, 1838, p. 108, pl. XXIV, fig. 116.—* TROSCHER, Arch. für Nat., V, 1839, Pt. 2, p. 237.
 * *Margarita (Unio) brownianus* LEA, Syn., 1838, p. 13.
 * *Hyria browniana* HANLEY, Test. Moll., 1842, p. 215; * Biv. Shells, 1843, p. 215.—* CATLOW and REEVE, Conch. Nom., 1845, p. 65.—* SOWERBY, Conch. Icon., XVII, 1869, pl. v, fig. 12.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

¹ This form has the posterior wing greatly produced and curved upward. I doubt whether it is more than a form of *syrrmatophora*.

² This is probably a variety of *obliquus* Schumacher.

³ Only a part of the interior is figured, but from its form I think there need be no doubt as to what the author meant.

⁴ This may be only a peculiar *obliquus*. The dorsal line curves up in the center.

* *Margaron (Triquetra) browniana* LEA, Syn., 1852, p. 17; 1870, p. 26.

* *Triquetra browniana* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 508.

Amazon River.

Subgenus **HYRIANA** Simpson, 1900.

(Type, *Unio ortonii* Lea.)

Shell solid, compressed, elliptical, with a slight posterior dorsal wing and a strong one anteriorly, and a well-defined but low posterior ridge ending in a point well above the base, inflated at post-basal region, distinctly sulcate; beaks low, apparently not rayed; epidermis shining, light yellowish brown; hinge line evenly curved; pseudocardinals numerous, much split into denticles and crooked; two laterals in the left valve and one (no doubt) in the right; beak cavities very shallow; dorsal scars numerous, anterior scars deep; nacre whitish, radially grooved along the palleal line.¹

† **PRISODON ORTONII** Lea.

* *Unio ortonii* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 161; *Jl. Ac. N. Sci. Phila.,

VI, 1868, p. 321, pl. LI, fig. 134; *Obs., XII, 1869, p. 81, pl. LI, fig. 134.—

*PÄTEL, Conch. Sam., III, 1890, p. 162.

* *Margaron (Unio) ortonii* LEA, Syn., 1870, p. 28.

River Napo, Ecuador.

Genus **DIPLODON** Spix, 1827.

(Type, *Diplodon ellipticum* Spix.)

Diplodon SPIX, Test. Fluv. Bras., 1827, p. 33, pl. xxvi.

Shell elliptical, rounded, elongated or trapezoidal, with rather low beaks which are more or less distinctly radially sculptured, the ridges usually curved and approaching below, with a low or scarcely developed posterior ridge; surface slightly concentrically sculptured, sometimes broken into fine nodules or corrugations; epidermis dull, rayless; hinge with two compressed pseudocardinals in the right valve, and one slender lateral, and two compressed pseudocardinals in the left valve, one in front of the other, and two laterals; nacre bluish to white, dull, often blotched; beak cavities shallow; dorsal scars numerous, forming a row in the beak cavity parallel with the hinge line.

Animal with the marsupium occupying nearly the whole length of the inner branchiæ, a few ovules sometimes being found in the outer gills;²

¹ I have seen only a single left valve, the type. The anterior wing of this is gone, and the epidermis is somewhat eroded. The shell has been injured when young, which has probably caused a long, low, radial furrow that runs in front of the posterior ridge. The space occupied by the pseudocardinals is not so long proportionally as in typical *Prisodon*.

² I found a few ovules in the outer gills of *D. acutirostris* Lea, of South America, while the inner gills were full, and Mr. Henry Suter states that the ovules of *D. menziesi* Gray, of New Zealand, occupy the inner gills for the most part.

branchiæ rather large, angular at base, inner much the larger, united their whole length to the abdominal sac; palpi scarcely projecting posteriorly; mantle very thin, thickened on the edges; branchial opening papillose, separated from the smooth anal opening by a strong bridge; superanal opening not closed below.

Subgenus **DIPLODON** Spix, 1827.

Beak sculpture consisting of unbroken ridges, covering the whole beaks.

Animal having the labial palpi rounded below and at posterior base, scarcely projecting behind.

(Type, *Diplodon ellipticum* Spix.)

(Group of *Diplodon delodontus*.)

Shell elliptical to subtrapezoidal, inflated, subsolid, with a low posterior ridge, sometimes having a slight posterior dorsal wing; one or two central pairs of bars of the beak sculpture usually joining below; pseudocardinals sometimes slightly dentellate.

Animal having the characters of the genus.

† **DIPLODON DELODONTUS** Lamarck.

* *Unio delodonta* LAMARCK, An. sans. Vert., VI, 1819, p. 77.—* DELESSERT, Rec. Coq. Lam., 1841, pl. XII, fig. 7.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 605.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* CHENU, Ill. Conch., 1858, pl. XII, figs. 1. 1a.

* *Margarita (Unio) delodontus* LEA, Syn., 1836, p. 26; 1838, p. 20.

* *Unio delodontus* HANLEY, Test. Moll., 1842, p. 194; * Biv. Shells, 1843, p. 194, pl. XXI, fig. 56.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* KUSTER, Conch. Cab. Unio, 1861, p. 234, pl. LXXVIII, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 150.

* *Margaron (Unio) delodontus* LEA, Syn., 1852, p. 29; 1870, p. 46.

* *Unio delodon* STROBEL, Mat. Mal., Pt. 1, 1874, p. 71.

*† *Unio lacteolus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 40, pl. VIII, fig. 19; * Obs., I, 1834, p. 152, pl. VIII, fig. 19.—* D'ORBIGNY, Mag. Zool., 1835, p. 34.

Argentina; Paraguay; Uruguay; southern Brazil.

† **DIPLODON RHOMBEUS** Wagner.

* *Unio rhombeus* WAGNER, Test. Fluv. Bras., 1827, p. 34, pl. XXVIII, figs. 1, 2.—

* HANLEY, Test. Moll., 1842, p. 208; * Biv. Shells, 1843, p. 208, pl. XXII, fig. 20.—

* CATLOW and REEVE, Conch. Nom., 1845, p. 63.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* PÆTEL, Conch. Sam., III, 1890, p. 165.

* *Margarita (Unio) rhombeus* LEA, Syn., 1836, p. 39; 1838, p. 25.

* *Margaron (Unio) rhombeus* LEA, Syn., 1852, p. 38; 1870, p. 61.

Brazil.

† **DIPLODON PAULISTA** von Ihering.

* *Unio paulista* VON IHERING, Arch. für Nat., 1893, p. 93, pl. IV, fig. 7.—* NEHRING, Jl. de Conch., 1894, p. 82.

São Paulo, Brazil.

† *DIPLODON AMPULLACEUS* Lea.

- * *Unio ampullaceus* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 34; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 269, pl. XXXV, fig. 83; * Obs., XII, 1869, p. 29, pl. XXXV, fig. 83.
 * *Margaron (Unio) ampullaceus* LEA, Syn., 1870, p. 53.

South America.

† *DIPLODON APPRIMUS* Lea.

- * *Unio apprimus* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 34; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 263, pl. XXXIII, fig. 78; Obs., XII, 1869, p. 23, pl. XXXIII, fig. 78.
 * *Margaron (Unio) apprimus* LEA, Syn., 1870, p. 46.

South America.

† *DIPLODON WYMANII* Lea.¹

- * *Unio wymanii* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 90; * Jl. Ac. N. Sci. Phila., VI, 1863, p. 381, pl. XLII, fig. 289; * Obs., X, 1863, p. 17, pl. XLII, fig. 289.—
 * SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIV, fig. 449.—* VON MARTENS, Mal. Bl., XV, 1868, p. 193.—* PÆTEL, Conch. Sam., III, 1890, p. 172.
 * *Margaron (Unio) wymanii*, LEA, Syn., 1870, p. 35.

Uruguay River, South America.

DIPLODON EXPANSUS Kuster.²

- * *Unio expansus* KUSTER, Conch. Cab. Unio, 1856, p. 149, pl. XLIII, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 152.

Brazil?

† *DIPLODON PECULIARIS* Lea.

- * *Unio peculiaris* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 33; * Jl. Acad. N. Sci. Phila., VI, 1868, p. 265, pl. XXXIV, fig. 80; Obs., XII, 1869, p. 25, pl. XXXIV, fig. 80.
 * *Margaron (Unio) peculiaris* LEA, Syn., 1870, p. 47.
 * † *Unio paraguayensis* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 34; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 271, pl. XXXV, fig. 85; Obs., XII, 1869, p. 31, pl. XXXV, fig. 85.—* VON MARTENS, S. B. Nat. Fr., 1895, p. 34.
 * *Margaron (Unio) paraguayensis* LEA, Syn., 1870, p. 45.
 * † *Unio nitidulus* KUSTER, Conch. Cab. Unio., 1848, p. 226, pl. LXXVI, fig. 6.—
 * PÆTEL, Conch. Sam., III, 1890, p. 161.

Paraguay.

† *DIPLODON FIRMUS* Lea.³

- * *Unio firmus* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 33; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 267, pl. XXXIV, fig. 82; * Obs., XII, 1869, p. 27, pl. XXXIV, fig. 82.
 * *Margaron (Unio) firmus* LEA, Syn., 1870, p. 45.

DIPLODON FIRMUS var. *BÆTTGERI* von Ihering.

- * *Unio firmus* LEA var. *bættgeri* VON IHERING, Arch. für Nat., 1893, p. 105, pl. IV, fig. 2.—* NEHRING, Jl. de Conch., 1894, p. 83.

Brazil.

¹ Extremely close to *apprimus*, and probably only a young shell of that species.

² I know nothing of this. It looks more like an Australian than a South American form, and is quite likely a *D. australis*. Kuster credits it to Charpentier manuscript.

³ A little longer and more solid than *peculiaris*, but I doubt whether it is distinct. The types of these two and *paraguayensis* are badly worn.

† DIPLODON URUGUAYENSIS Lea.

- * *Unio uruguayensis* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 90; *Jl. Ac. N. Sci. Phila., V, 1863, p. 388, pl. XLV, fig. 298; *Obs., X, 1863, p. 241, pl. XLV, fig. 298.—
 *SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIV, fig. 448.—*PÆTEL, Conch. Sam., III, 1890, p. 171.

*Margaron (*Unio*) *uruguayensis* LEA, Syn., 1870, p. 46.

Uruguay River, South America.

† DIPLODON WHEATLEYANUS Lea.

- * *Unio wheatleyanus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 94; *Obs., VI, 1857, p. 28, pl. XXIX, fig. 23; *Jl. Ac. N. Sci. Phila., 1858, p. 308, pl. XXIX, fig. 23.—
 *SOWERBY, Conch. Icon., XVI, 1868, pl. XL, fig. 487.—*STROBEL, Mat. Mal., Pt. 1, 1874, p. 71.

*Margaron (*Unio*) *wheatleyanus* LEA, Syn., 1870, p. 50.

Argentina.

† DIPLODON RUDUS Lea.

- * *Unio rudus* LEA, Pr. Ac. Nat. Sci. Phila., III, 1859, p. 187; *Jl. Ac. Nat. Sci. Phila., IV, 1860, p. 266, pl. XLIII, fig. 146; Obs., VII, 1860, p. 84, pl. XLIII, fig. 146.—*KUSTER, Conch. Cab. Unio, 1861, p. 261, pl. LXXXVIII, fig. 1.

*Margaron (*Unio*) *rudus* LEA, Syn., 1870, p. 50.

* *Unio rudis* PÆTEL, Conch. Sam., III, 1890, p. 166.

Rio de la Plata.

† DIPLODON GREEFIANUS von Ihering.

- * *Unio greeffianus* VON IHERING, Arch. für Nat., 1893, p. 96, pl. IV, fig. 18.¹—*NEHRING, Jl. de Conch., 1894, p. 82.

São Paulo, Brazil.

† DIPLODON PIGER Lea.

- * *Unio piger* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 90; *Jl. Ac. N. Sci. Phila., V, 1863, p. 388; *Obs., X, p. 23, pl. XLV, fig. 296.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIV, fig. 449.—*PÆTEL, Conch. Sam., III, 1890, p. 163.

*Margaron (*Unio*) *piger* LEA, Syn., 1870, p. 46.

Uruguay River, South America.

† DIPLODON PRUNOIDES Lea.

- * *Unio prunoides* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 150; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 323, pl. LIII, fig. 136; *Obs., XII, 1869, p. 83, pl. LIII, fig. 136.

*Margaron (*Unio*) *prunoides* LEA, Syn., 1870, p. 47.

South America.?

† DIPLODON BESKEANUS Dunker.

- * *Unio beskeanus* DUNKER, Zeits. für Mal., V, 1848, p. 182.²

*Margaron (*Unio*) *beskeanus* LEA, Syn., 1870, p. 61.

Brazil.

¹Credited to Dunker Manuscript.

²There is a specimen in the Lea collection contributed under this name by Dunker himself. It is near *locellus* and *suavidicus*. I do not know that it has ever been figured.

†DIPLODON LOCELLUS Lea.

- * *Unio locellus* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 34; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 264, pl. XXXIII, fig. 79; * Obs., XII, 1869, p. 24, pl. XXXIII, fig. 79.
* *Margaron (Unio) locellus* LEA, Syn., 1870, p. 47.

Argentina.

†DIPLODON SUAVIDICUS Lea.

- * *Unio suavidicus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 95; * Obs., VI, 1857, p. 29, pl. XXIX, fig. 24; * Jl. Ac. N. Sci. Phila., III, 1858, p. 309, pl. XXIX, fig. 24.
* *Margaron (Unio) suavidicus* LEA, Syn., 1870, p. 54.

Amazon.

†DIPLODON BROWNII Lea.

- * *Unio brownii* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 95; * Obs., VI, 1857, p. 27, pl. XXIX, fig. 22; * Jl. Ac. N. Sci. Phila., III, 1858, p. 307, pl. XXIX, fig. 22.
* *Margaron (Unio) brownii* LEA, Syn., 1870, p. 61.

South America?. Dr. Lea reports it from Mocha, Asia (with a question), which is certainly an error.

†DIPLODON LEAI Simpson.¹

- * *Margarita (Unio) modestus* LEA, Syn., 1836, p. 32; 1838, p. 22.
* *Margaron (Unio) modestus* LEA, Syn., 1852, p. 33; 1870, p. 54.
* *Unio modestus* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.

Brazil.

†DIPLODON RHUACOICUS d'Orbigny.

- * *Unio rhuacoica* D'ORBIGNY, Guer. Mag., 1835, p. 35; * Voy. Am. Mer., 1843, p. 606, pl. LXIX, figs. 4, 5.
* *Margaron (Unio) rhuacoicus* LEA, Syn., 1870, p. 54.
* *Unio rhyacoicus* VON MARTENS, Mal. Bl., XV, 1868, p. 195.
* *Monocondylaea rhyacaeca* PATEL, Conch. Sam., III, 1890, p. 147.

Uruguay.

†DIPLODON CHARRUANUS d'Orbigny.

- * *Unio charruana* D'ORBIGNY, Guer. Mag., 1835, p. 35; Voy. Am. Mer., 1843, p. 606, pl. LXXI, figs. 8-14.
* *Margarita (Unio) charruanus* LEA, Syn., 1838, p. 20.

¹ Lea credits this to Ferussac in his Synopsis, but I do not think it has ever been described by that author or anyone else. In 1856 Kuster properly described and figured a species in the Conchylien Cabinet (Vol. Unio, p. 147, pl. XLIII, fig. 2) under the name of *Unio modestus*, which he credits to Charpentier's manuscript, a different shell from that referred to by Lea and the Adams brothers. If the latter was never described the name *modestus* can not be used for it, as it was first properly applied to another species. The shells in Lea's collection may be described as follows: Shell small, subrhomboid, rather solid; beaks full with radial sculpture, the two middle bars coalescing; surface faintly concentrically sculptured, dark olive; two compressed pseudocardinals in each valve which are strongly obliquely striated; two laterals in the left valve and one in the right; anterior muscle scars distinct; naere bluish; beak cavities shallow. Length 28, height 17, diameter 13 mm.

- * *Margaron (Unio) charruanus* LEA, Syn., 1852, p. 29; 1870, p. 54.
 * *Unio charruanus* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCIII, figs. 505, 505a, 505b.—* PÆTEL, Conch. Sam., III, 1890, p. 147.
 * *Unio faba* D'ORBIGNY, Mag. Zool., 1835, p. 35.—* HANLEY, Test. Moll., 1842, p. 197.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 606, pl. LXXI, figs. 8–11.—* HANLEY, Biv. Shell, 1843, p. 197.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* PÆTEL, Conch. Sam., III, 1890, p. 152.
 * *Margarita (Unio) faba* LEA, Syn., 1838, p. 21.
 * *Margaron (Unio) faba* LEA, Syn., 1852, p. 31; 1870, p. 50.

Uruguay.

DIPLODON CAIPIRA von Ihering.

- * *Unio caipira* VON IHERING, Arch. für Nat., 1893, p. 98, pl. IV, fig. 9.—* NEHRING, Jl. de Conch., 1894, p. 83.

Southern Brazil.

† DIPLODON PICEUS Lea.

- * *Unio piceus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 91; * Jl. Ac. N. Sci. Phila., V, 1863, p. 397, pl. XLI, fig. 287; * Obs., X, 1863, p. 15, pl. XLI, fig. 287.
 * *Margaron (Unio) piceus* LEA, Syn., 1870, p. 53.
 * *Unio suppositus* VON IHERING (according to von Ihering), where is it described?

Uruguay River, South America.

† ? DIPLODON WAGNERIANUM Simpson.¹

- * *Unio ellipticus* WAGNER, Test. Fluv. Bras., 1827, p. 33, pl. XXVI, figs. 1, 2.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* KUSTER, Conch. Icon. Unio, 1861, p. 238, pl. LXXX, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIV, fig. 382.—* PÆTEL, Conch. Sam., III, 1890, p. 151.
 * *Margaron (Unio) ellipticus* LEA, Syn., 1852, p. 21; 1870, p. 31.

Brazil.

† DIPLODON ÆTHIOPS Lea.

- * *Unio æthiops* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 91; * Jl. Ac. N. Sci. Phila., V, 1863, p. 377, pl. XLI, fig. 285; Obs., X, 1863, p. 13, pl. XLI, fig. 285.—* VON MARTENS, Mal. Bl., XVI, 1868, p. 196.
 * *Margaron (Unio) æthiops* LEA, Syn., 1870, p. 53.

DIPLODON ÆTHIOPS var. PIRICICABANA von Ihering.

- * *Unio æthiops*, LEA, var. *piricicabana* VON IHERING, Arch. für Nat., 1893, p. 102.—* NEHRING, Jl. de Conch., 1894, p. 83.

Uruguay and Paraguay rivers; southern Brazil.

¹Barnes applied the name *Unio ellipticus* in 1823 to a species which is believed to be *Unio ligamentinus*. Wagner placed this shell in the genus *Unio* in the text, but Spix called it *Diplodon ellipticum* in the plate. I therefore change the name to *wagnerianum*. I have never seen a shell that just agrees with his figure. It is near *æthiopiformis* von Ihering.

† *DIPLODON DIVARICATUS* Lea.¹

* *Unio divaricatus* LEA, Tr. Am. Phil. Soc., V, 1834, p. 64, pl. IX, fig. 24; * Obs., I, 1834, p. 176, pl. IX, fig. 24.—* HANLEY, Test. Moll., 1842, p. 197; * Biv. Shells, 1843, p. 197, pl. XXIII, fig. 36.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—* JICKELI, Faun. Moll. N. O. Af., 1874, p. 280.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

* *Margarita (Unio) divaricatus* LEA, Syn., 1836, p. 29; 1838, p. 21.

* *Margaron (Unio) divaricatus* LEA, Syn., 1852, p. 31; 1870, p. 49.

South America.

† *DIPLODON BINNEYI* Lea.²

* *Unio binneyi* LEA, Tr. Am. Phil. Soc., IV, 1845, p. 165; * Tr. Am. Phil. Soc., X, 1848, p. 77, pl. VI, fig. 18; * Obs., IV, 1848, p. 51, pl. VI, fig. 18.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 245.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

* *Margaron (Unio) binneyi* LEA, Syn., 1852, p. 29; 1870, p. 46.

South America.

(Group of *Diplodon granosus*.)

Shell elliptical to subtrapezoidal, somewhat compressed, rather thin, beaks low, the sculpture irregularly radial, the ridges narrow and sharp, separated by wide, trough-like spaces, the outer in particular often becoming somewhat nodulous or corrugated, the inner approaching and united below; surface sometimes slightly concentrically ridged and granose; teeth delicate, compressed; nacre bluish.³

Animal unknown.

† *DIPLODON GRANOSUS* Bruguiere.

* *Unio granosa* BRUGUIERE, Jl. de Hist. Nat., I, 1792, p. 107, pl. VI, figs. 3, 4.—* LAMARCK, An. sans Vert., VI, 1819, p. 79.—* DESHAYES, Enc. Meth., II, 1827, p. 151, pl. CCXLIX, fig. 2.

* *Unio granosus* HANLEY, Test. Moll., 1842, p. 182; * Biv. Shells, 1843, p. 182, pl. XXIII, fig. 21.—* CATLOW and REEVE, Conch. Nom., 1845, p. 59.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* DROUET, Moll. Guyane, 1859, p. 86.

¹ Credited by Lea to Egypt, but no doubt a South American species, as it has regular radial beak sculpture and appears to belong with this group.

² Dr. Lea believes that this shell came from Alabama, but it certainly did not. It is undoubtedly a South American form. The type, the only shell I have seen is dead and in bad condition; but the beaks have radial sculpture, and I think it belongs here.

³ No wide distinction exists between this and the preceding group, and there are species which might be assigned about as easily to one as the other. In general the forms of this division are more delicate, thinner, and more compressed than those of the *Delodontus* group. There are few clear characters in any of the species. Philippi and others have, it seems to me, unduly multiplied names for these forms, and the lack of suitable material and their extreme simplicity and similarity make it impossible for me to always determine which are valid. The synonymy is in bad shape, but I have done the best I could at straightening it out.

- * *Margarita (Unio) granosus* LEA, Syn., 1836, p. 17; 1838, p. 16.
 * *Margarou (Unio) granosus* LEA, Syn., 1852, p. 23; 1870, p. 34.
 * *Unio multistriatus* LEA,¹ Tr. Am. Phil. Soc., IV, 1831, p. 91, pl. XII, fig. 22; * Obs. I, 1834, p. 101, pl. XII, fig. 22.—* HANLEY, Test. Moll., 1842, p. 176; * Biv. Shells, 1843, p. 176, pl. XX, fig. 35.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* CHENU, Ill. Conch., 1858, pl. XI, figs. 2, 2a, 2b.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXV, fig. 455.—* PÆTEL, Conch. Sam., III, 1890, p. 160.
 * *Margarita (Unio) multistriatus* LEA, Syn., 1836, p. 13; 1838, p. 14.
 * *Unio multistriata* D'ORBIGNY, Voy. Am. Mer., 1843, p. 607.
 * *Unio pfeifferi* DUNKER, Zeits. für Mal., V, 1848, p. 181.—* PFEIFFER, Nov. Conch., II, 1866, p. 151, pl. XXXIX, figs. 4-9.—* PÆTEL, Conch. Sam., III, 1890, p. 163.
 * *Unio psammactinus* PHILIPPI, Conch., III, 1848, p. 79, pl. V, fig. 2.—* KUSTER, Conch. Cab. Unio, 1856, p. 159, pl. XLV, fig. 6.²—* VON MARTENS, Mal. Bl., XV, 1868, p. 194.
 * † *Unio famelicus* GOULD, Pr. Bost. Soc. N. Hist., III, 1850, p. 294;³ U. S. Expl. Exp., XII, 1852, p. 432, figs. 544, 544a, 544b.—* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 249.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* GOULD, Otia. Conch., 1862, p. 88.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXV, fig. 391.—* B. H. WRIGHT, Check List, 1888.
 * *Margaron (Unio) famelicus* LEA, Syn., 1852, p. 34; 1870, p. 54.
 * *Unio niloticus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIII, fig. 374.

Brazil; Guiana.

DIPODON GRANULIFERUS Dunker.

- * *Unio granuliferus* DUNKER, Zeits. für Mal., V, 1848, p. 182.—* PFEIFFER, Nov. Conch., II, 1866, p. 150, pl. XXXIX, figs. 1-3.
 * *Unio granulifer* PÆTEL, Conch. Sam., III, 1890, p. 154.

Province of Rio de Janeiro, Brazil.

† DIPODON EFFULGENS Lea.

- * *Unio effulgens* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 94; * Jl. Ac. N. Sci. Phil., III, 1857, p. 303, pl. XXVIII, fig. 18; * Obs., VI, 1857, p. 23, pl. XXVIII, fig. 18.—PÆTEL, Conch. Sam., III, 1890, p. 151.—* VON IHERING, Arch. für Nat., 1893, p. 106.
 * *Margaron (Unio) effulgens* LEA, Syn., 1870, p. 35.
 * *Unio eurhynchus* KUSTER, Conch. Cab. Unio, 1861, p. 237, pl. LXXIX, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 152.

Brazil.

¹The type presented by Mrs. Mawe is not in the Lea collection, but there are fairly typical examples in it. The species varies much in form and sculpture, some specimens being smooth, and all variations occur to those quite granose throughout. Lea places his species in the synonymy *U. ellipticus* Spix, which I think is an error. The type of *multistriatus* is more elongated than the figure of *U. granosus* in the Encyclopedie Méthodique, but agrees very well with undoubted specimens of this species which I have seen.

²Credited by Kuster to Bronn in litt.

³Credited by Gould to Walla Walla, Oregon. It is a young, rather smooth *granosus* I am sure, and the habitat given is wrong. The type is in the U. S. National Museum.

† DIPLODON RUFOFUSCUS Lea.

* *Unio rufofuscus* LEA, Pr. Ac. N. Sci. Phila., IX, 1865, p. 76; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 282, pl. XXXIX, fig. 96; *Obs., XII, 1869, p. 42, pl. XXXIX, fig. 96.

* *Maragon (Unio) rufofuscus* LEA, Syn., 1870, p. 31.

South America.

DIPLODON DEMARARAENSIS Lea.

* *Unio demararaensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 152; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 253, pl. XXXIX, fig. 133; *Obs., VII, 1860, p. 71, pl. XXXIX, fig. 133.—*REEVE, Conch. Icon., XVI, 1865, pl. XXII, fig. 99.

* *Maragon (Unio) demararaensis* LEA, Syn., 1870, p. 36.

* *Unio demararaensis* PÆTEL, Conch. Sam., III, 1890, p. 150.

Demerara.

† DIPLODON LEPIDIOR Lea.

* *Unio lepidus* LEA,¹ Pr. Ac. N. Sci. Phila., IV, 1860, p. 91; *Jl. Ac. N. Sci. Phila., V, 1863, p. 390, pl. L, fig. 306; *Obs., X, 1863, p. 25, pl. L, fig. 306.

* *Margaron (Unio) lepidior* LEA, Syn., 1870, p. 53.

Uruguay River.

† DIPLODON PARCUS Lea.

* *Unio parvus* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 34; *Jl. Ac. N. Sci. Phila., VI, 1868, p. 262, pl. XXXIII, fig. 77; *Obs., XII, 1869, p. 22, pl. XXXIII, fig. 77.

* *Margaron (Unio) parvus* LEA, Syn., 1870, p. 47.

South America.

† DIPLODON CHILENSIS Gray.

* *Unio chilensis* GRAY, Spic. Zool., 1828, pl. VI, fig. 12.—* PHILIPPI, Conch., III, 1847, p. 9, pl. IV, fig. 2.—* HUPE, Gay's Hist. Chile, VIII, 1854, p. 317.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* KUSTER, Conch. Cab. Unio, 1862, p. 282, pl. XCV, fig. 12.—* SOWERBY, Conch. Icon., XVI, 1867, pl. LVI, fig. 286.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

* *Unio smithii* GRAY and PIDGEON, Griff. Cuv., XII, 1834, p. 600, pl. XX, fig. 3.²—* HANLEY, Biv. Shells, 1843, p. 195, pl. XXIII, fig. 58.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 495.—* PÆTEL, Conch. Sam., III, 1890, p. 167.

* *Margarita (Unio) smithii* LEA, Syn., 1836, p. 27; 1838, p. 20.

* *Margaron (Unio) smithii* LEA, Syn., 1852, p. 30; 1870, p. 48.

* *Unio auratus* PHILIPPI, Conch., III, 1847, p. 9, pl. IV, fig. 1.³—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* PÆTEL, Conch. Sam., III, 1890, p. 145.

* *Margarita (Unio) auratus* LEA, Syn., 1836, p. 31; 1838, p. 22.

* *Margaron (Unio) auratus* LEA, Syn., 1852, p. 33; 1870, p. 53.

* *Unio araucanus* PHILIPPI, Conch., III, 1847, p. 50, pl. IV, fig. 3.—* KUSTER, Conch. Cab. Unio, 1862, p. 283, pl. XCV, fig. 3.

* *Margaron (Unio) araucanus* LEA, Syn., 1870, p. 53.

* *Unio araucana* HUPE, Gay's Hist. Chile, 1854, p. 317.

Chile.

¹ Changed to *lepidior* because the name *lepidus* had been used for a *Unio* by Gould.

² Only the briefest description is given of this in the index. From the figure it seems to be identical with *chilensis*.

³ Generally credited to Swainson, and Philippi refers it to him with doubt. But I can not find that Swainson ever published it.

DIPLODON SOLIDULUS Philippi.

- * *Unio solidulus* PHILIPPI, Mal. Bl., XVI, 1869, p. 46.—* PFEIFFER, Nov. Conch., III, 1869, p. 480, pl. CIII, figs. 9, 10.—* PÆTEL, Conch. Sam., III, 1890, p. 168.

Near Santiago; Chile.

DIPLODON GASSIESI Kuster.

- * *Unio gassiesi* KUSTER(part), Conch. Cab. Unio, 1856, p. 148, pl. XLIII, fig. 4.¹
 * *Unio auratus* REEVE, Conch. Icon., XVI, 1865, pl. XXIX, fig. 150.

Chile.

DIPLODON APLATUS Reeve.²

- * *Unio aplatus* REEVE, Conch. Icon., XVI, 1865, pl. XXVIII, fig. 143.—* PÆTEL, Conch. Sam., III, 1890, p. 144.

Island of Chiloe, Chile.

† DIPLODON MOLINÆ Philippi.

- * *Unio molinæ* PHILIPPI, Conch., III, 1847, p. 50, pl. IV, fig. 4.—* HUPE, Gay's Hist. Chile, VIII, 1854, p. 316.—* KUSTER, Conch. Cab. Unio, 1862, p. 281, pl. XCV, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 159.
 * *Unio auratus* SOWERBY, Conch. Icon., XVI, 1866, pl. XLV, fig. 245.³

Southern Chile.

† DIPLODON MODESTUS Kuster.⁴

- * *Unio modestus* KUSTER, Conch. Cab. Unio, 1856, p. 147, pl. XLIII, fig. 2.
 * *Unio ianthinus* PHILIPPI, Mal. Bl., XVII, 1869, p. 47.—* PFEIFFER, Nov. Conch., III, 1869, p. 485, pl. CIV, figs. 11, 12.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
 * *Unio valdivianus* PHILIPPI, Mal. Bl., XVI, 1869, p. 48.—* PFEIFFER, Nov. Conch., III, 1869, p. 479, pl. CIII, figs. 7, 8.
 * *Unio montanus* PHILIPPI, Mal. Bl., XVI, 1869, p. 48.—* PFEIFFER, Nov. Conch., III, 1869, p. 482, pl. CIV, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 159.

Chile.

† DIPLODON ATRATUS Sowerby.

- * *Unio atratus* SOWERBY, Conch. Man., 1839, fig. 148.⁵
 * *Unio limensis* KUSTER, Conch. Cab. Unio, 1856, p. 146, pl. XLII, fig. 7; XLIII, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 157.
 * *Unio jacobæus* PHILIPPI, Mal. Bl., XVI, 1869, p. 44.—* PFEIFFER, Nov. Conch., III, 1869, p. 478, pl. CII, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
 * *Unio landbecki* PHILIPPI, Mal. Bl., XVI, 1869, p. 45.—* PFEIFFER, Nov. Conch., III, 1869, p. 479, pl. CIII, figs. 5, 6.—* PÆTEL, Conch. Sam., III, 1890, p. 156.

¹ Kuster figures two species, I think, for *gassiesi*. His fig. 3 is, I think, a young *molinæ* Kuster; fig. 4 seems to equal *auratus* of Reeve, not of Philippi or Sowerby.

² Swainson manuscript according to Reeve.

³ Not *U. auratus* of Reeve, figured previously in the same work.

⁴ Credited to Charpentier manuscript by Kuster. See foot note on *Diplodon leai* Simpson, p. 876.

⁵ Not *Unio (Niaa) atrata* Swainson, which equals *Unio gracilis* Barnes. Sowerby's name has precedence.

- * *Unio diplodon* PHILIPPI, Mal. Bl., XVI, 1869, p. 46.—* PFEIFFER, Nov. Conch., III, 1869, p. 483, pl. CIV, figs. 7, 8.—* PÆTEL, Conch. Sam., III, 1890, p. 150.
 *† *Unio dunkerianus* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 94; * Obs., VI, 1857, p. 25, pl. XXIII, fig. 20; * Jl. Ac. N. Sci. Phila., III, 1858, p. 303, pl. XXVIII, fig. 20.—* KUSTER, Conch. Cab. Unio, 1862, p. 290, pl. XCH, fig. 6.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XC, fig. 485.—* PÆTEL, Conch. Sam., III, 1890, p. 151.—* VON IHERING, Arch. für Nat., 1893, p. 109.
 * *Margaron (Unio) dunkerianus* LEA, Syn., 1870, p. 32.

Peru; Chile.

DIPLODON OBTUSUS d'Orbigny.

- * *Unio obtusa* D'ORBIGNY,¹ Gner. Mag., 1835, p. 35; * Voy. Am. Mer., 1843, p. 610.—
 * HUPE, Gay's Hist. Chile, VIII, 1854, p. 316.

Chile.

DIPLODON RHUACONICUS Kuster.²

- * *Unio rhuaconicus* KUSTER, Conch. Cab. Unio, 1856, p. 145, pl. XLII, fig. 5.

Brazil.

† DIPLODON CHILOËNSIS Kuster.

- * *Unio chiloënsis* KUSTER, Conch. Cab., 1856, p. 161, pl. XXXV, fig. 6.—* PÆTEL, Conch. Sam., III, 1890, p. 148.

Island of Chiloe, Chile.

† DIPLODON CASÆBLANCÆ Philippi.

- * *Unio casablancae* PHILIPPI, Zeits. für Mal., V, 1848, p. 176.
 * *Unio casablancae* PFEIFFER, Nov. Conch., III, 1869, p. 481, pl. CIV, figs. 1, 2.—
 * PÆTEL, Conch. Sam., III, 1890, p. 147.
 * *Margaron (Unio) casablancae* LEA, Syn., 1870, p. 53.
 * *Unio longus* PHILIPPI, Mal. Bl., XVI, 1869, p. 44.—* PFEIFFER, Nov. Conch., III, 1869, p. 477, pl. CHII, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 158.
 * *Unio colchaguensis* PHILIPPI, Mal. Bl., XVI, 1869, p. 47.—* PFEIFFER, Nov. Conch., III, 1869, p. 484, pl. CIV, figs. 9, 10.—* PÆTEL, Conch. Sam., III, 1890, p. 148.
 * *Unio foncki* PHILIPPI, Mal. Bl., XV, 1869, p. 49.—* PFEIFFER, Nov. Conch., III, 1869, p. 483, pl. CIV, figs. 5, 6.
 * *Unio foncki* PÆTEL, Conch. Sam., III, 1890, p. 153.

Chile.

DIPLODON FRENZELLII von Ihering.

- * *Unio frenzelli* VON IHERING, Arch. für Nat., 1893, p. 3, pl. IV., fig. 12.

Patagonia and Chile.

DIPLODON MARTENSI von Ihering.

- * *Unio martensi* VON IHERING, Arch. für Nat., 1893, p. 100, pl. IV, fig. 10.

Southern Brazil.

¹Said by d'Orbigny to be of Ferussac, but he does not say where that author described it, and I do not know where. This name has been used for at least four different species of *Unionidae*: As above; in Cuvier's Animal Kingdom (edition without date); for *Unio retusus*, by Potiez and Michaud; and by Dr. Lea. I presume that d'Orbigny is entitled to the name.

²Kuster credits this to "Pfeiffer teste Charpentier." I do not know what it is. It is very different from d'Orbigny's *rhuacoica*.

† DIPLODON CUPRINUS Simpson.

* *Unio fragilis* SOWERBY,¹ Conch. Icon., XVI, 1856, pl. xxx, fig. 155.

Island of Chiloe, Chile.

DIPLODON AUREUS Simpson.

* *Unio auratus* KUSTER, Conch. Cab. Unio, 1856, p. 161, pl. XLVI, fig. 3.²

Chile.

DIPLODON CORIACEUS Dunker.

* *Unio coriaceus* DUNKER, Zeits. für. Mal., V, 1848, p. 181.—* VON IHERING, Arch. für Nat., 1893, p. 115.

Rio Negro, Province of Rio Janeiro, Brazil.

DIPLODON CHILDRENI Gray.

* *Unio childreni* GRAY, Griff. Cuv., XII, 1834, p. 600 (index), pl. xx, fig. 1.—* CATLOW and REEVE, Conch. Icon., 1845, p. 57.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* PETEL, Conch. Sam., III, 1890, p. 148.

Margarita (Unio) childreni LEA, Syn., 1836, p. 25; 1838, p. 19.

* *Margaron (Unio) childreni* LEA, Syn., 1852, p. 28; 1870, p. 44.

South America.

(Group of *Diplodon burroughianus*.)

Shell elliptical, moderately solid, inflated, sharply pointed behind, with a high, well-developed posterior ridge; beaks rather full; the sculpture consisting of regularly radiating, widely separated, sharp ridges, the intervening grooves rounded out; peridermis smooth, dark olive with lighter bands; pseudocardinals compressed, rough, vertically ridged, usually trifid or quadrifid in each valve; laterals curved, compressed, muscle scars smooth; nacre white, silvery.

Animal unknown.

† DIPLODON BURROUGHIANUS Lea.

* *Unio burroughianus*, LEA, Tr. Am. Phil. Soc., V, 1834, p. 67, pl. x, fig. 27; * Obs., I, 1834, p. 179, pl. x, fig. 27.—* D'ORBIGNY, Guer. Mag., 1835, p. 34.—* HANLEY, Test. Moll., 1842, p. 197; * Biv. Shells, 1843, p. 197, pl. xxii, fig. 2.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 609.—* CATLOW and REEVE, Conch. Nom., 1845, p. 56.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* SOWERBY, Conch. Icon., XVI, 1866, pl. xxxii, fig. 169.—* PETEL, Conch. Sam., III, 1890, p. 146.

* *Margarita (Unio) burroughianus*, LEA, Syn., 1836, p. 29; 1838, p. 21.

* *Margaron (Unio) burroughianus*, LEA, Syn., 1852, p. 31; 1870, p. 50.

Parana River, South America.

¹Sowerby's name being preoccupied in *Unio*, I change it to *cuprinus*, which has not been used in *Diplodon*. The date, 1856 on the plate, is perhaps a typographical error for 1866.

²The name *auratus* is preoccupied in *Unio*, by Philippi in the Conchylien. This being a different species, I change it to *aureus*.

† DIPLODON TRIFIDUS Lea.

- * *Unio trifidus*, LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 89; * Jl. Ac. N. Sci. Phila., V, 1863, p. 386, pl. XLIV, fig. 295; * Obs., X, 1863, p. 22, pl. XLIV, fig. 295.
 * *Margaron (Unio) trifidus*, LEA, Syn., 1870, p. 50.

Argentina.

(Group of *Diplodon pazi*.)

Shell subtrapezoidal, rounded in front and strongly biangulate behind, the wavy beak sculpture extending well over the disk; epidermis olive; pseudocardinals ragged, with a tendency to break into denticles.

DIPLODON PAZI Hidalgo.

- * *Castalia pazi* HIDALGO, Jl. de Conch., XVI, 1868, p. 353, pl. XIII, fig. 6.—* PÆTEL, Conch. Sam., III, 1890, p. 189.

Imbabura, Ecuador.

DIPLODON HYLÆAUS d'Orbigny.

- * *Unio hylæa* D'ORBIGNY, Guer. Mag., 1835, p. 36; * Voy. Am. Mer., 1843, p. 607, pl. LXIX, figs. 8, 9.
 * *Margaron (Unio) hylæus* LEA, Syn., 1852, p. 21; 1870, p. 31.
 * *Unio hylæus* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCIII, fig. 506.—* PÆTEL, Conch. Sam., III, 1890, p. 155.
 * *Unio guaraniana* D'ORBIGNY, Guer. Mag., 1835, p. 37; * Voy. Am. Mer., 1843, p. 608, pl. LXIX, figs. 10–12.

Bolivia; Paraguay River.

(Group of *Diplodon parallelipipedon*.)

Shell elongated, subtrapezoidal, inflated, obliquely truncated behind; posterior ridge strong; beaks rather low, sculpture of nearly or quite strictly radial bars with concave spaces between; epidermis dark; pseudocardinals compressed in the young, granular, vertically striate and breaking into denticles in the old shell; laterals long, curved, granular, often vertically striate; anterior scars smooth, distinct.

† DIPLODON PARALLELIPIPEDON Lea.

- * *Unio parallelipipedon* LEA, Tr. Am. Phil. Soc., V, 1834, p. 60, pl. VIII, fig. 20; * Obs., I, 1834, p. 172, pl. VIII, fig. 20.—* D'ORBIGNY, Guer. Mag., 1835, p. 34.—* HANLEY, Test. Moll., 1842, p. 205; Biv. Shells, 1843, p. 205.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 609.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 491.—* KUSTER, Conch. Cab. Unio, 1861, p. 220, pl. LXXIV, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIX, fig. 478.—* PÆTEL, Conch. Sam., III, 1890, p. 162.
 * *Margarita (Unio) parallelipipedon* LEA, Syn., 1836, p. 35; 1838, p. 24.
 * *Margaron (Unio) parallelipipedon* LEA, Syn., 1852, p. 56; 1870, p. 58.
 ? *Unio bonplandi* VALENCIENNES, in continuation of Humboldt's Zool. Obs., 1827.¹

Rio de la Plata system.

¹ According to Lea. I have not seen this paper, and Lea does not cite the page. A specimen with the above label from the Morelet collection = *parallelipipedon* Lea.

†*DIPODON PATAGONICUS* d'Orbigny.

* *Unio patagonica* D'ORBIGNY, Güer. Mag., 1835, p. 37; * Voy. Am. Mer., 1843, p. 610, pl. LXX, figs. 1-4.

* *Margarita (Unio) patagonicus* LEA, Syn., 1838, p. 25.

* *Unio patagonicus* HANLEY, Test. Moll., 1842, p. 208; * Biv. Shells, 1843, p. 208, pl. XXII, fig. 16.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 461.—* REEVE, Conch. Icon., XVI, 1865, pl. XXI, fig. 93.—* STROBEL, Mat. Mal., Pt. 1, 1874, p. 72.—* PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Margaron (Unio) patagonicus* LEA, Syn., 1852, p. 38; 1870, p. 61.

Patagonia.

†*DIPODON ACUTIROSTRIS* Lea.

* *Unio acutirostris* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 34; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 270, pl. XXXV, fig. 84; * Obs., XIII, 1869, p. 30, pl. XXXV, fig. 84.

* *Margaron (Unio) acutirostris* LEA, Syn., 1870, p. 58.

South America.

(Group of *Diplodon quadrans*.)

Shell subtrapezoid, slightly truncate in front, nearly equilateral, greatly inflated, its greatest diameter below the beaks, from which it is wedge-shaped in front and behind, solid, with a decidedly rounded posterior ridge; epidermis brownish, concentrically striate, beaks full but not high, ligament long and slender; pseudocardinals large, compressed, striate and crenulate, double in the right valve, single in the left; laterals curved, crenulate; beak cavities rounded; anterior cicatrices deep, confluent; nacre white.

Animal unknown.

DIPODON QUADRANS Lea.¹

* *Unio quadrans* LEA, Pr. Ac. N. Sci. Phila., IV, 1859, p. 306; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 360, pl. LXI, fig. 185; * Obs., VIII, 1860, p. 42, pl. LXI, fig. 185.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Unio) quadrans* LEA, Syn., 1870, p. 36.

South America (?).

Subgenus *CYCLOMYA* Simpson, 1900.

(Type, *Unio funebris* Lea.)

Shell obovate to suborbicular, narrowed in front, produced just behind the center of the base, generally slightly pointed about the middle of the posterior end, with a scarcely perceptible posterior ridge and a slight dorsal wing behind; beaks high, irregularly radial; hinge line

¹ Only one specimen of this shell is reported, this being in the Wheatley collection, and credited to Texas. The beak sculpture is all eroded away, but I am almost sure it is a South American shell, allied, perhaps, to *D. apprimus*, *uruguayensis*, and possibly to Hupe's *Unio orbigny*.

strongly arched, curved behind and incurved in front of the beaks; the lower pseudocardinal in the right valve largest, often much split up into denticles, pseudocardinals of the left valve variable, dentellate; two lower anterior scars deep, united, the upper very deep, separate.

Animal unknown.

Section CYCLOMYA Simpson, 1900.

(Type, *Unio funebris* Lea.)

† DIPLODON GRATUS Lea.

**Unio gratus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 91; *Jl. Ac. N. Sci. Phila., V, 1863, p. 382, pl. XLIII, fig. 290; *Obs., X, 1863, pl. XLIII, fig. 290.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIV, fig. 444.—*PÆTEL, Conch. Sam., III, 1890, p. 154.

*Margaron (*Unio*) *gratus* LEA, Syn., 1870, p. 56.

Uruguay River, South America.

† DIPLODON PATELLOIDES Lea.

**Unio patelloides* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 89; *Jl. Ac. N. Sci. Phila., V, 1863, p. 383, pl. XLIII, fig. 291; *Obs., X, 1863, p. 19, pl. XLIII, fig. 291.

*Margaron (*Unio*) *patelloides* LEA, Syn., 1870, p. 56.

Amazon River.

† DIPLODON PERÆFORMIS Lea.

**Unio peraformis* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 90; *Jl. Ac. N. Sci. Phila., V, 1863, p. 384, pl. XLIII, fig. 292; *Obs., X, 1863, p. 20, pl. XLIII, fig. 292.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXIV, fig. 443.—*PÆTEL, Conch. Sam., III, 1890, p. 163.

*Margaron (*Unio*) *peraformis* LEA, Syn., 1870, p. 55.

Uruguay River, South America.

† DIPLODON FONTAINIANUS d'Orbigny.

**Unio fontainiana* D'ORBIGNY, Guer. Mag., 1835, p. 36.

**Unio fontainianus* HANLEY, Biv. Shells, 1856, p. 384, pl. XXI, fig. 27.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVII, fig. 466.—*PÆTEL, Conch. Sam., III, 1890, p. 153.—*VON IHERING, Arch. für Nat., 1893, p. 90.

*Margaron (*Unio*) *fontainianus* LEA, Syn., 1870, p. 56.

**Unio fontaineana* D'ORBIGNY, Voy. Am. Mer., 1843, p. 605, pl. LXIX, figs. 6, 7.—*HUPE, Anim. Nouv., 1857, p. 82.

Parana River, southern Brazil.

† DIPLODON ROTUNDUS Wagner.

**Unio rotundus* WAGNER,¹ Test. Fluv. Bras., 1827, p. 34, pl. XXV, figs. 3, 4.—

*MORICAND, Mem. His. Soc. Gen., 1838, p. 49, pl. IV, figs. 12-14.—*KUSTER, Conch. Cab. Unio, 1856, p. 160, pl. XLVI, figs. 1, 2.—*SOWERBY, Conch. Icon., XVI, 1868, pl. LXXII, fig. 369.—PÆTEL, Conch. Sam., III, 1890, p. 166.

*Margaron (*Unio*) *rotundus* LEA, Syn., 1870, p. 56.

Brazil.

¹ *Diplodon rotundum* Spix in explanation of plate.

†DIPLODON DISCULUS Lea.

* *Unio disculus*, LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 91; * Jl. Ac. N. Sci. Phila., V, 1863, p. 385, pl. XLIV, fig. 293; * Obs., X, 1863, p. 21, pl. XLIV, fig. 293.

* *Margaron Unio disculus* LEA, Syn., 1870, p. 55.

Uruguay River, South America.

†DIPLODON FUNEBRALIS Lea.

* *Unio funebris* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 91; * Jl. Ac. N. Sci. Phila., V, 1863, p. 378, pl. XLI, fig. 286; * Obs., X, 1863, p. 14, pl. XLI, fig. 286.—

* SOWERBY, Conch. Icon., XVI, 1867, pl. LVII, fig. 290.—* PÆTEL, Conch. Sam., III, 1890, p. 153.

* *Margaron Unio funebris* LEA, Syn., 1870, p. 55.

Uruguay River.

†DIPLODON PARANENSIS Lea.

* *Unio paranensis* LEA, Tr. Am. Phil. Soc., V, 1834, p. 49, pl. XIV, fig. 42.—* Obs., I, 1834, p. 211, pl. XIV, fig. 42.—* HANLEY, Test. Moll., 1842, p. 202; * Biv. Shells, 1843, p. 202, pl. XXI, fig. 3.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 603.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* KUSTER, Conch. Cab., 1861, p. 253, pl. LXXXV, fig. 3.—* SOWERBY, Conch. Icon., XVI, 1866, pl. LI, fig. 268; * PÆTEL, Conch. Sam., III, 1890, p. 162.

* *Margarita (Unio) paranensis* LEA, Syn., 1836, p. 33; 1838, p. 23.

* *Margaron (Unio) paranensis* LEA, Syn., 1852, p. 34; 1870, p. 55.

*† *Unio nocturnus* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 91; * Jl. Ac. N. Sci. Phila., V, 1863, p. 380, pl. XLII, fig. 288; * Obs., X, 1863, p. 16, pl. XLII, fig. 288.

* *Margaron (Unio) nocturnus* LEA, Syn., 1870, p. 56.

Uruguay and Parana rivers.

Section BULLOIDEUS Simpson, 1900.

(Type, *Unio bulloides* Lea.)

Shell rounded, inflated, thin, nearly equilateral, truncate behind and slightly so before, with a rather sharp posterior ridge and a dorsal wing; beaks full, rather high, regularly radial; epidermis smooth, bronzy olive; pseudocardinals compressed, much elongated, disposed to be split into denticles, two in the right valve and one in the left; laterals two in the left valve and one in the right; dorsal scars few and scattered in the rather shallow cavities; naere bluish.

†DIPLODON BULLOIDES Lea.

* *Unio bulloides* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 187; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 264, pl. XLII, fig. 144; * Obs., VII, 1860, p. 82, pl. LXII, fig. 144.—

* KUSTER, Conch. Cab. Unio, 1861, p. 186, pl. LIX, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVIII, fig. 275.—* PÆTEL, Conch. Sam., III, 1890, p. 146.

Margaron (Unio) bulloides LEA, Syn., 1870, p. 55.

Rio de la Plata.

†DIPLODON SOLISIANUS d'Orbigny.

* *Unio solisiana* D'ORBIGNY, Guer. Mag., 1835, p. 35; * Voy. Am. Mer., 1843, p. 604, pl. LXIX, figs. 1-3.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCIII, fig. 508.

Rio de la Plata and its affluents.

DIPLODON VARIABILIS Maton.

* *Mya variabilis* MATON, Tr. Linn. Soc. Lond., X, 1811, p. 327, pl. XXIV, figs. 4-7.¹—
* WOOD, Gen. Conch., I, 1815, p. 111.—* DILLWYN, Cat., I, 1817, p. 53.—* WOOD, Ind.
Test., 1825, p. 13, pl. II, fig. 38a; rev. ed., 1856, p. 17, pl. III, fig. 38.

* *Unio variabilis* D'ORBIGNY, Voy. Am. Mer., 1843, p. 604, pl. LXXI, figs. 1-3.—

* HANLEY, Test. Moll., 1842, p. 202; * Biv. Shells, 1843, p. 202.—* CATLOW and
REEVE, Conch. Nom., 1845, p. 65.—* H. and A. ADAMS, Gen. Rec. Moll.,
II, 1857, p. 492.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXIV, fig. 381.—

* STROBEL, Mat. Mal., Pt. 1, 1874, p. 69.—* PATEL, Conch. Sam., III, 1890,
p. 171.

* *Margarita (Unio) variabilis* LEA, Syn., 1836, p. 33; 1838, p. 23.

* *Margaron (Unio) variabilis* LEA, Syn., 1852, p. 35; 1870, p. 56.

* *Mytilus matoniana* D'ORBIGNY, Guer. Mag., 1835, p. 35.

* *Unio matonianus* PATEL, Conch. Sam., III, 1890, p. 158.

* ? *Unio membranacea* HANLEY, Biv. Shells, 1843, p. 202, pl. XXII, fig. 6.

* *Unio membranaceus* PHILIPPI, Conch., III, 1848, p. 80, pl. v, fig. 4.—* KUSTER,
Conch. Cab. Unio., 1862, p. 284, pl. xcv, fig. 5.

Rio de la Plata.

Subgenus **HYRIDELLA** Swainson, 1840.

(Type, *Unio australis* Lamarck.)

Beaks rather low, sculpture consisting of curved, generally nodulous ridges, which approach below but usually have a smooth area of shell between them; surface sulcate or sometimes corrugated and nodulous; epidermis rayless; teeth rather delicate, compressed, often somewhat rudimentary.

Animal having the embryos occupying the inner gills for the most part,² which are united for their entire length to the abdominal sac; outer gills pointed below in the middle; palpi triangular; branchial opening papillose; anal opening smooth, not separated from the super anal opening.

Section **HYRIDELLA** Swainson, 1840.

Shell covered with concentric, sulcate sculpture; pseudocardinals well developed, compressed; laterals delicate, sometimes imperfect.

(Group of *Diplodon dorsuosus*.)

Shell somewhat rhomboid, inflated solid, with rather full beaks, the sculpture consisting of very strong, subradial, corrugated ridges which curve toward each other below, those in the center sometimes joining, this sculpture extending well out on the disk; upper part of posterior slope having slight radial folds.

¹I feel certain that Maton has illustrated two species, figs. 4, 6, and 7 being nearly circular, with a rather straight hinge, while fig. 5 is a small shell, is much inflated at the posterior base, and is incurved in front of the breaks. He describes the large shell first. The small one may be the young of *D. patelloides*.

²According to Suter in a letter.

† DIPLODON DORSUOSUS GOULD.

* *Unio dorsuosus* GOULD, Pr. Bost. Soc. N. H., III, 1850, p. 296; * U. S. Expl. Exp., XII, 1852, p. 430, figs. 540, 540a, 540b; * Otia Conch., 1862, p. 89.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* PÆTEL, Conch. Sam., III, 1890, p. 151.

* *Margaron (Unio) dorsuosus* LEA, Syn., 1852, p. 21.

* *Unio napeanensis* CONRAD, Pr. Ac. N. Sci. Phila., V, 1852, p. 10; * Jl. Ac. N. Sci. Phila., 1854, p. 296, pl. XXVI, fig. 4.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 498.—* REEVE, Conch. Icon., XVI, 1865, pl. XXIII, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 160.

* *Margaron (Unio) napeanensis* LEA, Syn., 1852, p. 20; 1870, p. 30.

Nepean River, Australia.

DIPLODON GLENELGENSIS DENNANT.

* *Unio glenelgensis* DENNANT, Pr. Roy. Soc. Vict., X, 1898, p. 112, pl. iv.

Glenelg River, Victoria.

(Group of *Diplodon menziezi*.)

Shell subrhomboid; beak sculpture not strong, consisting of broken, nodulous ridges curving toward each other below, with generally a smooth space between, not extending over the shell.

Animal as in the subgenus.

† DIPLODON MENZIEZI Gray.

* *Unio menziezi* GRAY, In Dieffenbach's N. Z., II, 1843, p. 257.—* ? MUSGRAVE, Phot. Conch., 1863, pl. II, fig. 1.—* HUTTON, N. Z. Moll., 1880, p. 160.—* PÆTEL, Conch. Sam., 1890, p. 159.

* *Unio menziezianus* REEVE, Conch. Icon., XVI, 1865, pl. XXIX, fig. 152.

* *Margaron (Unio) menzieni* LEA, Syn., 1870, p. 46.

* *Unio aucklandica* GRAY, In Dieffenbach's N. Z., II, 1843, p. 257.—* HUTTON, N. Z. Moll., 1880, p. 161.

* ? *Unio aucklandicus* SOWERBY, Conch. Icon., XVI, 1866, pl. XXX, fig. 156.—* PÆTEL, Conch. Sam., III, 1890, p. 145.

* *Margaron (Unio) aucklandicus* LEA, Syn., 1870, p. 45.

† DIPLODON MENZIEZI var. RUGATUS Hutton.¹

* *Unio rugatus* HUTTON, Tr. N. Z. Inst., XVI, 1884, p. 216.—* PÆTEL, Conch. Sam., III, 1890, p. 166.

† DIPLODON MENZIEZI var. HOCHSTELLERI Krauss.

* *Unio hochstelleri* DUNKER, Mal. Bl., VIII, 1862, p. 153.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVI, fig. 463.—* HUTTON, N. Z. Moll., 1880, p. 161.—* PÆTEL, Conch. Sam., III, 1890, p. 155.²

¹ The name *rugatus* was applied previously to a European *Unio* by Rossmassler; but it may be used as a varietal name. This seems to be a small form of *menziezi* which sometimes merges into *hochstelleri*.

² Suter believes this to be only a pathologic form of *menziezi*. He says in a letter to the author that he has seen the same kind of disease among small *rugatus*.

† *DIPODON MENZIEZI* var. *DEPAUPERATUS* Hutton.

* *Unio depauperatus* HUTTON, Tr. N. Z. Inst., XVI, 1884, p. 216.—*PÆTEL, Conch. Sam., III, 1890, p. 150.

New Zealand.

DIPODON WAIKARENSIS Colenso.

* *Unio waikarensis* COLENZO, Tasm. Jl. N. Sci., II, 1841, p. 250, footnote; Tr. N. Z. Inst., XIV, 1882, p. 169.¹

Waikare Lake, New Zealand.

† *DIPODON ZELEBORI* Dunker.

* *Unio zeleborei* DUNKER, Reise der Nov., 1867, p. 15, pl. II, fig. 28.—*HUTTON, N. Z. Moll., 1880, p. 161.—*PÆTEL, Conch. Sam., III, 1890, p. 172.

* *Margaron (Unio) zeleborei* LEA, Syn., 1870, p. 52.

New Zealand.

DIPODON FLYENSIS Tapperone Canefri.

* *Unio flyensis* TAPPERONE CANEFRI, Ann. Mus. Genov., XIX, 1883, p. 293, fig. 1.—*PÆTEL, Conch. Sam., III, 1890, p. 153.

Fly River, New Guinea.

† *DIPODON VITTATUS* Lea.

* *Unio vittatus* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 153; *Jl. Ac. N. Sci. Phila., IV, 1860, p. 249, pl. XXXVIII, fig. 128; *Obs., VII, 1860, p. 67, pl. XXXVIII, fig. 128.—*REEVE, Conch. Icon., XVI, 1864, pl. XXIII, fig. 83.—*PÆTEL, Conch. Sam., III, 1890, p. 171.

* *Margaron (Unio) vittatus* LEA, Syn., 1870, p. 35.

Australia.

† *DIPODON LESSONI* Kuster.

* *Unio lessoni* KUSTER, Conch. Cab., 1856, p. 135, pl. XXXVI, fig. 4.—*PÆTEL, Conch. Sam., III, 1890, p. 157.

* *Unio australis* KUSTER, Conch. Cab., 1861, p. 230, pl. LXXVII, fig. 6.

New South Wales.

† *DIPODON AUSTRALIS* (Lamarck) Hanley.²

*? *Unio australis* LAMARCK, An. sans. Vert., VI, 1819, p. 80.—*?DESHAYES, Enc. Méth., II, 1830, p. 582.—*HANLEY, Test. Moll., 1842, p. 192; *Biv. Shells, 1843, p. 192, pl. XXI, fig. 25.—*CATLOW and REEVE, Conch. Nom., 1845, p. 56.—*PHILIPPI, Conch., III, 1848, p. 81, pl. v, fig. 5.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—*PÆTEL, Conch. Sam., III, 1890, p. 145.

¹ Suter thinks this is a variety of *menziezi*, but Colenso states that the posterior slope is keeled. If this is so, it must be quite different from that species.

² Lamarck's description of this species is wholly inadequate, and would apply about equally well to several species. He refers to no figure, and Lea states that he did not see the type. I credit the species to Hanley, whose figure seems to represent a form of an abundant and variable species commonly referred to Lamarck.

- **Margarita (Unio) australis* LEA, Syn., 1836, p. 25; 1838, p. 19.
 **Hyridella australis* SWAINSON, Tr. on Mal., 1840, p. 285.
 **Margaron (Unio) australis* LEA, Syn., 1852, p. 28; 1870, p. 44.
 *? *Unio depressus* LESSON, Voy. Coquille, 1830, II, p. 427, pl. XV, fig. 5.
 **Margarita (Unio) depressus* LEA, Syn., 1836, p. 32; 1838, p. 22.
 **Margaron (Unio) depressus* LEA, Syn., 1852, p. 33; 1870, p. 54.
 **Unio ambiguus* PHILIPPI, Conch., III, 1847, p. 47, pl. III, fig. 2.¹—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXIX, fig. 355.—* PÆTEL, Conch. Sam., III, 1890, p. 144.
 **Unio shuttleworthi* KUSTER, Conch. Cab. Unio, 1856, p. 152, pl. XLIV, fig. 2.
 **Unio philippianus* KUSTER, Conch. Cab. Unio, 1861, p. 235, pl. LXXIX, fig. 2.
 **Unio balounensis* CONRAD, Pr. Ac. N. Sci. Phila., V, 1850, p. 10; *Jl. Ac. N. Sci. Phila., II, 1854, p. 295, pl. XXVI, fig. 3.
 *? *Unio daniellii* VILLA, Jl. de Conch., XIX, 1871, p. 328.—* SMITH, Jl. Linn. Soc., XVI, 1882, p. 307.—* PÆTEL, Conch. Sam., III, 1890, p. 150.
Unio bednalli TATE. Where? ²

† DIPLODON AUSTRALIS var. LEGRANDI Petterd.

- **Unio depressus* REEVE, Conch. Icon., XVI, 1864, pl. XVIII, fig. 81.
 **Unio legrandi* PETTERD, Pr. R. Soc. Tasm., 1887, p. 22; 1888, p. 81.

Australia; Tasmania.

† DIPLODON JEFFREYSIANUS Lea.

- **Unio jeffreysianus* LEA, Pr. Ac. N. Sci. Phila., I, 1871, p. 188; *Jl. Ac. N. Sci. Phila., VIII, 1874, p. 23, pl. VII, fig. 20; *Obs., XIII, 1874, p. 27, pl. VII, fig. 20.

Australia.

† DIPLODON PROFUGUS Gould.

- **Unio profugus* GOULD, Pr. Bost. Soc. N. II., 1850, p. 295;³ *U. S. Expl. Exp., XII, 1852, p. 429, figs. 543, 543a, 543b.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 494.—*GOULD, Otia Conch., 1862, p. 88.
 **Margaron (Unio) profugus* LEA, Syn., 1852, p. 29.

Hunter's River, New South Wales.

DIPLODON MORETONICUS Reeve.

- **Unio moretonicus* REEVE, Conch. Icon., XVI, 1865, pl. XXIV, fig. 118.—* PÆTEL, Conch. Sam., III, 1890, p. 160.
 **Margaron (Unio) mortonicus* LEA, Syn., 1870, p. 43.

Tasmania.

¹ Credited to Parreyss in litt.

² Specimens received from Mr. Bednall bearing the above name are undoubtedly a form of *D. australis*. I do not know whether it has been described or not.

³ The types are in the U. S. National Museum collection. Lea believes it to be the same as the *australis*. I am inclined to think it distinct.

† *DIPODON CULTELLIFORMIS* Conrad.¹

- *† *Unio depressa* LAMARCK, An. sans. Vert., VII, 1819, p. 79.—*† DELESSERT, Rec. Coq. Lam., 1841, pl. XII, fig. 5.—* CHENU, Ill. Conch., 1858, pl. XII, figs. 4, 4a.
- * *Unio depressus* HANLEY, Test. Moll., 1842, p. 200; * Biv. Shells, 1843, p. 200.—* CATLOW and REEVE, Conch. Nom., 1845, p. 58.—* CONRAD, Jl. Ac. N. Sci. Phila., 1854, p. 295, pl. XXXVI, fig. 2.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* PÆTEL, Conch. Sam., III, 1890, p. 150.
- * *Unio cultelliformis* CONRAD, Pr. Ac. N. Sci. Phila., V, 1850, p. 10.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* PÆTEL, Conch. Sam., III, 1890, p. 149.
- * *Margaron (Unio) cultelliformis* LEA, Syn., 1852, p. 32; 1870, p. 52.
- *† *Unio paramatensis* LEA, Pr. Ac. N. Sci. Phila., VI, 1862, p. 176; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 60, pl. XX, fig. 59; * Obs., XI, 1867, p. 64, pl. XX, fig. 59.
- * *Margaron (Unio) paramattensis* LEA, Syn., 1870, p. 35.

Australia.

† *DIPODON LUTULENTUS* Gould.

- * *Unio lutulentus* GOULD, Pr. Bost. Soc. N. Hist., III, 1850, p. 295; * U. S. Expl. Exp., XII, 1852, p. 428, figs. 542, 542a, 542b, 542c.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 493.—* GOULD, Otia Conch., 1862, p. 88.—* REEVE, Conch. Icon., XVI, 1865, pl. XXV, fig. 122.—* HUTTON, N. Z. Moll., 1880, p. 161.—* PÆTEL, Conch. Sam., III, 1890, p. 158.
- * *Margaron (Unio) lutulentus* LEA, Syn., 1852, p. 32; 1870, p. 52.

New Zealand.

† *DIPODON WILSONII* Lea.

- * *Unio wilsonii* LEA, Pr. Ac. N. Sci. Phila., 1859, p. 153; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 257, pl. XL, fig. 137; * Obs., VII, p. 74, pl. XL, fig. 137.—* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVIII, fig. 474.—* PÆTEL, Conch. Sam., III, 1890, p. 172.
- * *Margaron (Unio) wilsonii* LEA, Syn., 1870, p. 47.
- *† *Unio (Alasmodon) stuarti* ADAMS and ANGAS, Pr. Zool. Soc. Lond., 1863, p. 417.²
- * *Unio stuarti* SOWERBY, Conch. Icon., XVI, 1866, pl. LIV, fig. 279.
- * *Anodon stuarti* SOWERBY, Conch. Icon., XVII, 1867, pl. XXXIV, figs. 136, 136a, 136b.
- * *Margaron (Unio) stuarti* LEA, Syn., 1870, p. 52.
- * *Margaritana stuarti* PÆTEL, Conch. Sam., III, 1890, p. 174.
- * *Anodonta stuarti* PÆTEL, Conch. Sam., III, 1890, p. 185.

Australia.

† *DIPODON EVANSI* Adams and Angas.

- * *Unio (Alasmodon) evansi* ADAMS and ANGAS, Pr. Zool. Soc. Lond., 1864, p. 39.
- * *Unio evansi* SOWERBY, Conch. Icon., XVI, 1867, pl. LVI, fig. 285.—* PÆTEL, Conch. Sam., III, 1890, p. 152.

¹ I can not identify Lamarck's *Unio depressus* with any certainty from his very meager description. Lea believed it to equal the form I have considered as *australis*, but Lamarck's description seems to apply to a more elongated species. Delessert's figure in 1841 probably equals Lea's *paramattensis*. As the name *depressus* was used for a *Unio* in 1801 by Donovan I am obliged to use another for this.

² A specimen of this in the Lea collection from the Stuart expedition is identical with Lea's species.

* *Margaron (Unio) evansi* LEA, Syn., 1870, p. 56.

* *Margaritana evansi* PÆTEL, Conch. Sam., III, 1890, p. 173.

Murray River, Australia

Section CUCUMERIA Conrad, 1853.

(Type, *Unio novæ-hollandiæ* Gray.)

Shell elongated, trapezoidal, widest behind; pseudocardinals irregular, small, not well developed, showing a tendency to break into denticles; laterals feeble; pallial line strongly pitted; nacre much thicker in front.

(Group of *Diplodon novæ-hollandiæ*.)

Shell having the posterior two-thirds covered with irregular nodules, which radiate somewhat from the posterior ridge.

† DIPLODON NOVÆ-HOLLANDIÆ Gray.

* *Unio novæ-hollandiæ* GRAY, Pr. Zool. Soc. Lond., 1834, p. 57.—* MOLLER, Syn. Nov. Gen., 1836, p. 197.—* HANLEY, Test. Moll., 1842, p. 182; * Biv. Shells, 1843, p. 182.—* CATLOW and REEVE, Conch. Nom., 1845, p. 61.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 496.—* PÆTEL, Conch. Sam., III, 1890, p. 161.

* *Margarita (Unio) novæ-hollandiæ* LEA, Syn., 1836, p. 17; 1838, p. 16.

* *Margaron (Unio) novæ-hollandiæ* LEA, Syn., 1852, p. 23; 1870, p. 35.

* † *Unio cucumoides* LEA, Proc. Am. Phil. Soc., 1840, p. 285; * Tr. Am. Phil. Soc., VIII, 1842, p. 192, pl. VII, fig. 2; * Obs., III, 1842 p. 30, pl. VII, fig. 2.—* HANLEY, Biv. Shells, Sup., 1856, p. 382, pl. XXIV, fig. 4.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 497.—* CHENU, Ill. Conch., 1858, pl. XXVII, figs. 3, 3a, 3b.—* KUSTER, Conch. Cab. Unio, 1861, p. 219, pl. LXXIV, fig. 1.—* REEVE, Conch. Icon., XVI, 1865, pl. XIX, fig. 89.—* PÆTEL, Conch. Sam., III, 1890, p. 149.

* *Margaron (Unio) cucumoides* LEA, Syn., 1852, p. 21; 1870, p. 31.

* ? *Unio cumingianus* DUNKER, Zeits. für Mal., 1853, p. 53.

Australia.

(Group of *Diplodon shuttleworthii*.)

Beak sculpture consisting of strong, irregularly radiate, curved, nodulous bars; surface of shell somewhat sulcate, but not nodulous.

† DIPLODON SHUTTLEWORTHII Lea.

* *Unio shuttleworthii* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 94; * Jl. Ac. N. Sci. Phila., III, 1857, p. 304, pl. XXVIII, fig. 19; VI, 1857, p. 24, pl. XXVIII, fig. 19.—

* SOWERBY, Conch. Icon., XVI, 1866, pl. XXXII, fig. 167.

* *Margaron (Unio) shuttleworthii* LEA, Syn., 1870, p. 36.

* ? *Unio mutabilis* REEVE, Conch. Icon., XVI, 1865, pl. XXIV, fig. 112.

* *Unio angasi* SOWERBY, Conch. Icon., XVI, 1867, pl. LV, fig. 282.

* *Margaritana angasi* PÆTEL, Conch. Sam., III, 1890, p. 172.

* *Anodonta angasi* PÆTEL, Conch. Sam., III, 1890, p. 176.

Australia.

† DIPODON MUTABILIS Lea.

* *Unio mutabilis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 152; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 248, pl. XXXVII, fig. 127; * Obs., VII, 1860, p. 66, pl. XXXVIII, fig. 127.—* PÆTEL, Conch. Sam., III, 1890, p. 160.

* Margaron (*Unio*) *mutabilis* LEA, Syn., 1870, p. 53.

Murray River, Australia.

Subgenus LÆVIROSTRIS Simpson, 1900.

(Type, *Unio stagnorum* Dautzenberg.)

Shell irregularly quadrate, thin, inequilateral, compressed, rounded and narrowed in front, somewhat biangulate and truncate behind; beaks rather low, without sculpture;¹ surface finely concentrically grooved; epidermis delicately lamellated and serrate; pseudocardinals small; laterals elongated; anterior muscle impressions small, well marked; posterior scars superficial; nacre bluish-white.

Animal unknown.

DIPODON STAGNORUM Dautzenberg.

* *Unio stagnorum* DAUTZENBERG, Bull. Acad. Belg., XX, 1890, p. 372, pl. 1, figs. 7, 10.

Kongo River, Africa.

The following are unfigured and uncertain species.

* *Unio depressa* D'ORBIGNY, Guer. Mag., 1835, p. 34.²

* *Unio fokkesi* DUNKER, Zeits. für Mal., IX, 1853, p. 54.

Rio de la Plata, Brazil. According to von Ihering, this = *U. wheatleyanus* Lea.

* *Unio koseretzi* CLESSIN, Mal. Bl., X, p. 172.

River Guahyba, Brazil.

* *Unio macropterus* DUNKER, Zeits. für Mal., III, 1846, p. 109.

Brazil.

Unio paraguayanus VON MARTENS, S. B. Ges. Nat. Fr., 1895, p.

* *Unio zealandiens* PÆTEL, C. Sam., III, 1890, p. 172. Said to be of Gray. Where?

Unio damnoica D'ORBIGNY. Where?

Unio guahybae VON IHERING.

Unio athiopiformis VON IHERING.

Unio bishoffi VON IHERING.

Unio sebastanae VON IHERING.

Unio iheringi VON CLESSIN.

¹ According to Dautzenberg the beaks of the remarkable species on which I have founded the above subgenus are completely smooth. It has the appearance of a *Diplodon* in all other respects, according to the figures and description. Dautzenberg does not very carefully describe the teeth.

² According to d'Orbigny (Voy. Am. Mer., p. 611), this species, which he says is the *depressa* of Lamarck, is found in Chile, near Valparaiso. It can not be Lamarck's species.

I do not know where the last five species are described, and they probably have never been published. Wright has named a Texan species *Unio iheringi*.

Family MUTELIDÆ.

Shell usually without sculpture throughout; beaks smooth or but faintly corrugated, never exhibiting the remains of an embryonic shell; with or without teeth, which, when present, are irregularly taxodont, the shell showing vestiges of them in all the genera; nacre soft, richly tinted, generally surrounded with a wide, prismatic border; escutcheon large, distinct; pallial line usually simple, but sometimes having a slight sinus posteriorly.

Animal having the labial palpi large, rounded below, generally without free points, scarcely or not at all united posteriorly; anal and superanal openings not separated; mantle generally closed behind into branchial and anal siphons, which are always separated by a strong bridge; marsupium occupying the inner gills; embryo a lasidium, composed of three segments, the anterior head-like, the median bearing a single shell, the posterior tail-like.

Genus SPATHA Lea, 1838.

(Type, *Anodonta rubens* Lamarck.)

Spatha LEA, Tr. Am. Phil. Soc., VI, 1858, p. 141, footnote.

Shell elliptical; beaks very slightly or not at all sculptured; left valve having a faint, compressed tooth under and in front of the beak, which fits into a corresponding depression in the right valve; the entire hinge area often covered with longitudinally folded epidermal matter; escutcheon dark, sharply triangular; beak cavities shallow, with a single, elongate, deep scar in each; two large anterior muscle scars and two posterior ones, the upper small.

Animal having the palpi longer than wide, semicircular, attached along the upper edge, not united; mantle margin united behind so that the branchial and anal apertures are closed; on the under margin the mantle is entirely open; outer branchiæ united to the mantle to their extremity; inner the larger, free from the abdominal sack; foot tongue shaped, thick; anal and branchial openings separated by a strong bridge. *S. rubens* Lamarck (Troschel).

Subgenus SPATHA Lea, 1838.

(Type, *Spatha rubens* Lamarck.)

Shell solid, having faint concentric beak sculpture.

(Group of *Spatha rubens*.)

Sculpture of the beaks following the growth lines; shell solid.

† *SPATHA RUBENS* Lamarck.

- * *Anodonta rubens* LAMARCK, An. sans. Vert., VI, 1819, p. 85.—* BLAINVILLE, Man., 1825, p. 538.—* CAILLIAUD, Voy. à Méroé, IV, p. 262; Atlas, II, 1826, pl. LX, fig. 12.—* AUDOUIN, Savigny's Exp. de l'Egypte, Coquilles, 1827, pl. VII.¹—* DESHAYES, Enc. Meth., II, 1827, p. 147, pl. CCI, figs. 1, 1b.
- * *Iridina rubens* RANG, Nouv. Ann. Mus., 1885, p. 314.—* LEA, Syn., 1836, p. 56.—* HANLEY, Biv. Shells, 1843, p. 225.—* CATLOW and REEVE, Conch. Nom., 1845, p. 68.—* MORELET, Moll. Terr. et Fluv., 1868, p. 98.
- * *Platiris (Spatha) rubens* LEA, Syn., 1838, p. 33; 1852, p. 55; 1870, p. 89.
- * *Spatha rubens* CLESSIN, Conch. Cab. Ano., 1853, p. 185, pl. VII, fig. 1; 1876, pl. LXI, fig. 1.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, pl. CXIX; III, figs. 3, 3a.—* CHENU, Man., 1859, II, p. 148, fig. 729.—* PÆTEL, Conch. Sam., III, 1890, p. 188.
- * *Anodon rubens* SOWERBY, Conch. Icon., XVII, 1867, pl. II, fig. 5.
- Anodonta clappertoni* KENIG, Duchane and Clapperton's Travels, 1826, p.
- Anodonta splendens* DE CRISTOFORI, Crist. and Jan. Cat., 1832 (?), p.
- * *Iridina solida* ANTON, Verz. der Conch., 1839, p. 16.
- * *Anodonta solida* KUSTER, Conch. Cab. Ano., 1853, p. 50, pl. XII, fig. 1.
- * *Spatha wissmani* VON MARTENS, S. B. Nat. Fr., 1883, p. 73; Conch. Mitth., III, 1885 (?), p. 139, pl. XXVII.
- * *Spatha rotundata* VON MARTENS, Besch., 1897, p. 242, fig. d.²

† *SPATHA RUBENS* var. *CAILLIAUDI* von Martens.

- * *Anodonta rubens* AUDOUIN in SAVIGNY, Icon. Moll. Eg., 1827, pl. VII, fig. 1.
- * *Spatha cailliaudi* VON MARTENS, Mal. Bl., XIII, 1866, p. 9.—* JICKELI, L. and S. W. Moll., 1874, p. 259, pl. VIII, fig. 1.—* KOBELT, Icon., new ed., II, 1886, p. 27, pl. XLVI, fig. 267.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 314.—* PÆTEL, Conch. Sam., III, 1890, p. 188.
- * *Spatha bellamyi* JOUSSEAUME, Bull. Soc. Zool. Fr., XI, 1886, p. 491, pl. XIII, figs. 2, 2a.
- * *Spatha renei* JOUSSEAUME, Bull. Soc. Zool. Fr., XI, 1886, p. 492, pl. XIII, figs. 3, 3a.

Nile; Niger; Senegal; other points in West Africa.

† *SPATHA LEPSII* Jickeli.

- * *Spatha lepsii* JICKELI, Faun. L. and S. W. Moll., 1874, p. 265, pl. IX, fig. 4.—* PÆTEL, Conch. Sam., III, 1890, p. 188.

Upper Egypt: Senegal.

† *SPATHA CHAIZIANA* Rang.

- * *Anodonta chaiziana* RANG, Mem. Aceph. Senegal, (Nouv. Ann. Mns.), 1835, p. 13, pl. XXVIII.³—* MORELET, Moll. Terr. and Fluv., 1868, p. 97.

¹ Savigny's plates were issued without explanation in 1813. In 1827 Audouin published an explanation of them. Von Martens believes this is not the *Spatha rubens* of the Encyc. Méth. (Mal. Bl., 1866). This species is widespread, abundant, and variable, and the material I have examined leads me to believe that it may include either the short, rounded form of the Encyc. Méth. or specimens that are more elongated and even somewhat rhomboid.

² I can not see any essential difference between this and forms of what I believe are *S. rubens*. I believe that the *S. cailliaudi* is only one of these forms.

³ I have only seen the reprint of this paper, which begins with page 1.

* *Margarita (Anodonta) chaiziana* LEA, Syn., 1838, p. 30.

* *Anodon chaiziana* CATLOW and REEVE, Conch. Nom., 1845, p. 66.

* *Margaron (Anodonta) chaiziana* LEA, Syn., 1852, p. 49; 1870, p. 79.

* *Spatha chaiziana* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 507.—* CLESS'N, Conch. Cab. Anc., 1876, p. 187, pl. LXIII, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 188.

Senegal; Niger; Upper Nile; West Africa.

† *SPATHA TRISTIS* Jousseaume.

* *Spatha tristis* JOUSSEAUME, Bull. Soc. Zool. Fr., XI, 1886, p. 497.¹

* *Spatha rochebrunei* JOUSSEAUME, Bull. Soc. Zool. Fr., XI, 1886, p. 494, pl. XIV, figs. 1, 1a.

Upper Senegal.

SPATHA TRAPEZIA von Martens.

* *Spatha trapezia* VON MARTENS, Besch., 1897, p. 243, fig.

SPATHA TRAPEZIA var. *SENILIS* von Martens.

* *Spatha trapezia* var. *senilis* VON MARTENS, Besch., 1897, p. 244.

Lake Victoria Nyanza.

SPATHA DROUETI Chaper.

* *Spatha droueti* CHAPER, Bull. Soc. Zool. de Fr., X, 1885, p. 2, pl. I, figs. 1-3.²—

* PÆTEL, Conch. Sam., III, 1890, p. 188.

Assinie, Africa.

SPATHA MARTENSI STURANY.

Spatha martensi STURANY in Baumann, Durch Massai, 1894, p. 12, pl. XXV, fig. 39.—VON MARTENS, Besch., 1897, p. 244.

Stream falling into Lake Victoria Nyanza. I have not seen Sturany's paper.

SPATHA KIRKI Ancey.

* *Spathella Kirki* ANCEY, Bull. Soc. Zool. Fr., VII, 1894, p. 229, figs. 4-6.—* VON MARTENS, Besch., 1897, p. 244, fig.³

SPATHA KIRKI var. *LIEDERI* von Martens.

* *Spatha kirki* var. *liederi* VON MARTENS, Besch., 1897, p. 245.

Shire River and Lake Nyanza.

SPATHA MABILLET Jousseaume.

* *Spatha mabillet* JOUSSEAUME, Bull. Soc. Zool. de Fr., XI, 1886, p. 495, pl. XIV, figs. 2, 2a.

Senegal River.

¹ Morelet believes this to = *chaiziana*. An author's shell is in the U. S. National Museum, and I am inclined to believe it to be a valid species.

² Only inside and dorsal views, and I can not be sure whether it is a good species or not.

³ This species seems to me to be very close to *S. trapezia* von Martens.

SPATHA ADANSONI Jousseaume.

- * *Spathella adansonii* JOUSSEAUME, Bull. Soc. Zool. Fr., XI, 1886, p. 498, pl. xiv, figs. 4, 4a.

High Senegal.

†SPATHA WAHLBERGI Krauss.

- * *Iridina wahlbergi* KRAUSS, Sud Af. Moll., 1848, p. 19, pl. II, fig. 1.
 * *Platiris (Spatha) wahlbergi* LEA, Syn., 1852, p. 55; 1870, p. 89.
 * *Spatha wahlbergi* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 507.—* VON MARTENS, Mal. Bl., VI, 1860, p. 217.—* CLESSIN, Conch. Cab. Ano., 1876, p. 187, pl. LXIII, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 188.—* VON MARTENS, Besch., 1897, p. 247.
 * *Mutela wahlbergi* SMITH, Ann. and Mag., VIII, 1891, p. 319.
 * † *Spatha natalensis* LEA, Pr. Ac. N. Sci. Phila., VIII, 1864, p. 113; * JI. Ac. N. Sci. Phila., VI, 1866, p. 64, pl. xx, fig. 58; * Obs., XI, 1867, p. 68, pl. xx, fig. 58.—* CLESSIN, Conch. Cab. Ano., 1876, p. 189, pl. LXII, figs. 7, 8.
 * *Platiris (Spatha) natalensis* LEA, Syn., 1870, p. 89.
 * *Spatha hartmanni* VON MARTENS, Mal. Bl., XIII, 1866, p. 10.—* JICKELI, Faun., 1874, p. 263, pl. VIII, fig. 2.—* CLESSIN, Conch. Cab. Ano., 1875, p. 190, pl. LXI, figs. 2, 3.—* PÆTEL, Conch. Sam., III, 1890, p. 188.
 * *Anodon tabula* SOWERBY, Conch. Icon., XVII, 1867, pl. XVIII, fig. 68.
 * *Anodonta tabula* PÆTEL, Conch. Sam., III, 1890, p. 185.

SPATHA WAHLBERGI var BOURGNIGNATI Bourgnignat.

- * *Spatha bourgnignati* BOURGNIGNAT, Esp. Ouk. et Tan., 1885, p. 12.
 * *Spathella bourgnignati* BOURGNIGNAT, Moll. Af. Eq., 1889, p. 197, pl. VIII, fig. 1.
 * *Spathella spatuliformis* BOURGNIGNAT, Moll. Af. Eq., 1889, p. 199, fig.
 * *Spatha wahlbergi spatuliformis* VON MARTENS, Beschalte, 1897, p. 248, pl. VII, fig. 18.
 * *Spathella bloyeti* BOURGNIGNAT, Moll. Af. Eq., 1889, p. 198, pl. VIII, fig. 3.
 * *Spatha bloyeti* VON MARTENS, Besch., 1897, p. 249.

Tropical and southern Africa. The variety is reported from Lake Nyassa.

SPATHA LACUSTRIS Simpson.¹

- * *Spatha anceyi* BOURGNIGNAT, Bull. Soc. Zool. Fr., VII, 1894, p. 231, fig. 7.—* VON MARTENS, Besch., 1897, p. 247.

Lake Nyassa.

†SPATHA NYASSAENSIS Lea.

- * *Spatha nyassaensis* LEA, Pr. Ac. N. Sci. Phila., VIII, 1864, p. 109; * JI. Ac. N. Sci. Phila., VI, 1866, p. 36, pl. XIII, fig. 33; * Obs., XI, 1867, p. 40, pl. XIII, fig. 33.—* CLESSIN, Conch. Cab. Ano., 1876, p. 190, pl. LXII, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 188.—* VON MARTENS, Besch., 1897, p. 246.
 * *Platiris (Spatha) nyassaensis* LEA, Syn., 1870, p. 89.
 * *Spathella nyassaensis* ANCEY, Bull. Soc. Zool. Fr., VI, 1894, p. 228.
 * *Spatha nyassana* BOURGNIGNAT, Bull. Soc. Mal. Fr., VI, 1889, p. 38.

¹ Changed from *S. anceyi*, as there is another species with that name.

SPATHA MARNOI Jickeli.

* *Spatha marnoi* JICKELI, L. and Suss. Moll. N. Ost-Af., 1874, p. 259, pl. VIII, fig. 3.—

* PÆTEL, Conch. Sam., III, 1890, p. 188.

Abyssinia.

SPATHA SUBÆQUILATERA von Martens.

* *Spatha subæquilatera* VON MARTENS, Conch. Mitt., III, 1887, p. 18, pl. XLI, figs. 8, 9; * Besch., 1897, p. 246, pl. VII, fig. 16.

Spatha baumanni STURANY in Baumann, Durch. Massai., 1894, p. 12, pl. XXV, fig. 38.

Lake Victoria Nyanza.

SPATHA SINUATA von Martens.

* *Spatha sinuata* VON MARTENS, S. B. Nat. Fr., 1883, p. 173; Conch. Mitt., II, 1885, p. 190, pl. XXXIV, figs. 5, 6.

Congo.

(Group of *Spatha petersi*.)

Shell rather thin and small, greenish or olive; beak sculpture consisting of faint, more or less oblique, parallel ridges; hinge line narrow; teeth scarcely developed; nacre bluish or greenish. Animal unknown.

† SPATHA PETERSI von Martens.

* *Spatha petersi* VON MARTENS, Mal. Bl., VI, 1860, p. 218, pl. III, figs. 1, 2.—

* DOHRN, Pr. Zool. Soc. Lond., 1864, p. 117.—* VON MARTENS, Besch., 1897, p. 251.

* *Mutela petersi* CLESSIN, Conch. Cab. Ano., 1875, p. 197, pl. LXII, figs. 1, 2.—

* PÆTEL, Conch. Sam., III, 1890, p. 187.—* SMITH, Pr. Mal. Soc. Lond., 1894, p. 167.

Spathella petersi BOURGUIGNAT, Mol. Af. Eq., 1889, p. 197.

* † *Spatha modesta* LEA, Pr. Ac. N. Sci. Phila., VIII, 1864, p. 109; * Jl. Ac. N. Sci. Phila., VI, 1866, p. 37, pl. XIII, fig. 35; * Obs., XI, 1867, p. 41, pl. XIII, fig. 35.

* *Platiris (Spatha) modesta* LEA, Syn., 1870, p. 89.

Mozambique; Zanzibar and Uganda.

† SPATHA ARCUATA Cailliaud.

* *Anodonta arcuata* CAILLIAUD,¹ Voy. à Méroé, IV, 1826, p. 263; II, pl. LX, figs. 4, 5.—

* CATLOW and REEVE, Conch. Nom., 1845, p. 66.—* H. and A. ADAMS, Gen.,

Rec. Moll., II, 1857, p. 503.—* VON MARTENS, Mal. Bl., XIII, 1866, p. 77.—

* ?CLESSIN, Conch. Cab. Ano., 1874, p. 145, pl. XLVII, figs. 1, 2.—* JICKELI, L.

and Suss. Moll. N. Ost-Af. 1874, p. 265.—* PÆTEL, Conch. Sam., III, 1890, p. 176.

* *Margarita (Anodonta) arcuata* LEA, Syn., 1836, p. 54; 1838, p. 32.

* *Iridina arcuata* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 146, pl. LV, fig. 4.

* *Margaron (Anodonta) arcuata* LEA, Syn., 1852, p. 53; 1870, p. 83.

* *Spatha arcuata* JICKELI, Faun., 1874, p. 265.—* PÆTEL, Conch. Sam., III, 1890, p. 188.

Egypt; Abyssinia.

¹ *Arcuta* in The Voyage; a typographical error, I presume.

SPATHA CRYPTORADIATA Putzeys.

- * *Spatha cryptoradiata* PUTZEYS, Proc. Verb. Soc. Mal. Belg., 1898, pl. XXVII, figs. 14, 15.

Leopoldville; Congo.

SPATHA STUHLMANNI von Martens.

- * *Spatha stuhlmanni* VON MARTENS, Besch., 1897, p. 250, fig. ¹

Lake Albert; Nyanza.

SPATHA SUBRENIFORMIS Sowerby.

- * *Anodon soleniformis* SOWERBY, Conch. Icon., XVII, 1867, pl. XIV, fig. 50.
* *Anodonta subreniformis* PÆTEL, Conch. Sam., III, 1890, p. 185.

Lake Nyanza.

SPATHA PFEIFFERIANA Bernardi.

- * *Margaritana pfeifferiana* BERNARDI, Jl. de Conch., IV, 1860, p. 331, pl. XII, figs. 1, 2.—PÆTEL, Conch. Sam., III, 1890, p. 173.
* *Margaron (Margaritana) pfeifferiana* LEA, Syn., 1870, p. 68.

Gaboon; West Africa.

†SPATHA DAHOMEYENSIS Lea.

- * *Anodonta dahomeyensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 154; ²Jl. Ac. N. Sci. Phila., IV, 1859, p. 261, pl. XLI, fig. 141; * Obs., VII, 1860, p. 79, pl. XLI, fig. 141.—* PÆTEL, Conch. Sam., III, 1890, p. 178.—* CLESSIN, Conch. Cab. Ano., 1873, p. 103, pl. XXXI, figs. 5, 6.
* *Anodon dahomeyensis* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXVI, fig. 151.
* *Margaron (Anodonta) dahomeyensis* LEA, Syn., 1870, p. 82.
* † *Anodonta senegalensis* LEA, Pr. Ac. N. Sci. Phila., III, 1859, p. 154; * Jl. Ac. N. Sci. Phila., IV, 1860, p. 260, pl. XLI, fig. 140; * Obs., VII, 1860, p. 78, pl. XLI, fig. 140.—* CLESSIN, Conch. Cab. Ano., 1873, pl. XXXI, figs. 7, 8.—* PÆTEL, Conch. Sam., III, 1890, p. 184.
* *Anodon senegalensis* SOWERBY, Conch. Icon., XVII, 1867, pl. XI, fig. 35; 1870, pl. XXXIII, fig. 130.
* *Margaron (Anodonta) senegalensis* LEA, Syn., 1870, p. 82.
* *Spatha senegalensis* PÆTEL, Conch. Sam., III, 1890, p. 188.
* *Mutelina senegalica* JOUSSEAUME, Bull. Soc. Zool. de Fr., XI, 1886, p. 488.²

West Africa.

SPATHA COMPLANATA Jousseaume.

- * *Mutelina complanata* JOUSSEAUME, Bull. Soc. Zool. de Fr., XI, 1886, p. 489, pl. XIII, figs. 1, 1a.

Upper Senegal; Niger River.

¹ I am doubtful whether this is distinct from *S. arcuata*.

² Jousseaume has the unspeakable assurance to take the *Anodonta senegalensis* of Lea, place it in the so-called genus *Mutelina*, change the specific name to *senegalica*, and credit the whole to Bourguignat.

SPATHA DIVARICATA von Martens.

* *Spatha divaricata* VON MARTENS, Besch., 1897, p. 250, pl. VII, fig. 15.

Lake Victoria; Nyanza.

Subgenus **MONCETIA** Bourguignat, 1885.

(Type, *Moncetia anceyi* Bourguignat.)

Shell long-elliptical, inequilateral, compressed, thick, somber-colored; beaks compressed, smooth, sharp; hinge with a tubercular eminence near the beak of the right valve, with two internal ligaments and three groups of muscular impressions (Bourguignat).¹

Animal unknown.

SPATHA ANCEYI Bourguignat.

* *Moncetia anceyi* BOURGUIGNAT, Esp. Ouk., 1885, p. 34; * Icon. Mal., 1888, pl. xxx, fig. 1.—* VON MARTENS, Besch., 1897, p. 258.

* *Moncetia jouberti* BOURGUIGNAT Un. et Ir., 1886, p. 63; * Icon. Mal., 1888, pl. xxx, fig. 4.

Lake Tanganyika.

SPATHA MONIETI Bourguignat.

* *Moncetia monieti* BOURGUIGNAT, Un. et Ir., 1886, p. 61; * Icon. Mal., 1888, pl. xxx, fig. 7.

* *Moncetia rochebruncana* BOURGUIGNAT, Nouv. Mal., 1886, p. 62; * Icon. Mal., 1888, pl. xxx, fig. 8.

Lake Tanganyika.

SPATHA LAVIGERINA Bourguignat.

* *Moncetia lavigerina* BOURGUIGNAT, Nouv. Mal., 1886, p. 60; * Icon. Mal., 1888, pl. xxx, fig. 6.

* *Moncetia bridouri* BOURGUIGNAT, Un. et Ir., 1886, p. 65; * Icon. Mal., 1888, pl. xxx, fig. 5.

Lake Tanganyika.

Subgenus **ASPATHARIA** Bourguignat, 1885.

(Type, *Margaritana rignoniana* Bernardi.)

Shell elongated rhomboid, somewhat compressed, with a full, rounded posterior ridge, from which curved rows of fine, broken corrugations radiate; epidermis dark olive, wrinkled, rayless; beak sculpture not seen; hinge with a low, slightly elevated ridge in the left valve in front of the beak; dorsal scars small; naere lurid, bluish-green. Animal unknown.

¹This may be a distinct genus, though its characters seem to agree fairly well in most respects with those of *Spatha*. Bourguignat does not describe the muscle scars or tell anything of the naere.

†*SPATHA VIGNONIANA* Bernardi.

* *Margaritana vignoniana* BERNARDI, Jl. de Conch., VII, 1858, p. 302, pl. x, fig. 1.—

* PÆTEL, Conch. Sam., III, 1890, p. 174.

* *Unio vignoniana* REEVE, Conch. Icon., XVI, 1865, pl. xxv, fig. 120.

* *Anodonta vignonana* MUSGRAVE, Phot. Conch., 1863, pl. 1, fig. 1.

* *Anodon vignonanus* REEVE, Conch. Icon., XVII, 1870, pl. xxix, fig. 116.

* *Margaron (Unio) vignonana* LEA, Syn., 1870, p. 32.

* *Margaron (Margaritana) vignouana* LEA, Syn., 1870, p. 67.

* ? *Anodonta rugifera* DUNKER, Mal. Bl., V, 1858, p. 225.¹

Gaboon, West Africa.

SPATHA CORRUGATA Dautzenberg.

* *Spatha corrugata* DAUTZENBERG, Jl. de Conch., XLI, 1893, p. 50, pl. VIII, fig. 5.

River Niari, West Africa.

The following are unfigured and unidentified species.

* *Spatha pangallicensis* ROCHEBRUNE, Bull. Soc. Phil., VI, 1882, p. 33.

High Senegal.

* *Anodonta tawi* RANG., Nouv. Ann. Mus., 1834, p. 297.

Senegal.

* *Spatha baikii* H. ADAMS, Proc. Zool. Soc. Lond., 1866, p. 447.

Niger River.

* *Spatha anataria* JICKELI, Faun. Suss. Moll. N. O. Af., 1874, p. 266.

Egypt.

* *Spatha gancinensis* ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 9.

* *Spatha cornicola* ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 9.

* *Spathella protchi* ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 9.

Kongo River.

* *Chambardia* BOURGUIGNAT in Servain, 1891. New name for the Egyptian *Iridinide*.

Chambardia bourguignati, *C. locardiana*, *C. letourneuxiana*, *C. pharaonum*, *C. rhynchonella*, *C. rhynchoidéa*. Some of these are credited to Bourguignat and some to Servain. I don't know what they are. They may be Moncetias.

Genus *MUTELA* Scopoli, 1777.

(Type, *Iridina exotica* Lamarck.)

Mutela SCOPOLI, Intr. Nat. Hist., 1777, p. 397.

Iridina LAMARCK, An. sans Vert., VI, 1819, p. 88.

Calliscapha SWAINSON, Tr. on Mal., 1840, p. 380.

Eufira GISTEL, Naturg. Hohe Schul., 1848, p.

Shell elongated, slightly inflated, with low, smooth beaks, and a rounded posterior ridge; surface faintly concentrically grooved; epidermis rayless; hinge straight, having generally vestiges of taxodont teeth;

¹ Probably = *S. vignoniana* Bernardi, but there is only a Latin description. Both were published the same year.

dorsal scars an irregular row, extending downward and forward; muscle scars irregular.

Animal having the palpi longer than wide, not united; mantle margin united as far as the foot, so that the branchial and anal openings are closed; outer branchiæ united to the mantle to the extremity; inner entirely united to the foot; foot tongue shaped, somewhat produced anteriorly. *Iridina caelestis* Lea (Troschel.)

(Group of *Mutela exotica*.)

Shell elongate, rhomboid, wider and subtruncated behind, subsolid.

† *MUTELA EXOTICA* Lamarck.

* *Iridina exotica* LAMARCK, An. sans Vert., VI, 1819, p. 89; * Enc. Méth., II, 1827, p. 147, pl. CCIV, fig. 1, 1b; * STARK, Nat. Hist., II, 1828, p. 88.—* LEA, Syn., 1836, p. 56.—* HANLEY, Biv. Shells, 1843, p. 225.—* CATLOW and REEVE, Conch. Nom., 1845, p. 68.—* SOWERBY, Conch. Icon., XVI, 1868, pl. I, fig. 2.—* CLESSIN, Conch. Cab. Ano., 1875, p. 231, pl. LXXI, fig. 1.

* *Anodonta exotica* BLAINVILLE, Man., 1825, p. 538, pl. LXVI, fig. 3.

* *Platiris (Iridina) exotica* LEA, Syn., 1838, p. 33; 1852, p. 54; 1870, p. 88.

* *Pleiodon exoticus* PÆTEL, Conch. Sam., III, 1890, p. 188.

* *Iridina elongata* SOWERBY, Rec. and Fos. Shells, VII, 1821, fig. 1.¹—* OKEN, Isis, 1834, p. 458.—* SOWERBY, Conch. Man., 1839, fig. 150.—* SWAINSON, Tr. on Mal., 1840, p. 286, fig. 60.—* REEVE, Conch. Syst., I, 1841, p. 122, pl. XCII.—* HANLEY, Biv. Shells, 1843, p. 225.—* CATLOW and REEVE, Conch. Nom., 1845, p. 68.—* SOWERBY, Conch. Icon., XVI, 1868, pl. I, fig. 1.

* *Platiris (Spatha) elongata* LEA, Syn., 1838, p. 34.

* *Mutela elongata* PÆTEL, Conch. Sam., III, 1890, p. 187.

* *Iridina striata* OKEN, Isis, 1834, p. 458.

* ? *Mutela soleniformis* BOURGUIGNAT, Esp. Ouk., 1885, p. 25.—* PELSENEER, Bull. Mus. Belg., IV, 1886, p. 109.—* BOURGUIGNAT, Icon. Mal., 1888, pl. XXII, fig. 2.—* VON MARTENS, Besch., 1897, p. 251.

* ? *Mutela bridouri* BOURGUIGNAT, Un. et Ir., 1886, p. 25; * Icon. Mal., 1888, pl. XXII, fig. 1.

Tropical Africa.

† *MUTELA DUBIA* Gmelin.

Le Mutel ADANSON, Hist. Nat. du Sen., 1757, p. 234, pl. XVII, fig. 21.

* *Mytilus dubia* GMELIN, Syst. Nat., 1788, p. 3368.²

* *Anodonta dubia* BOSCH, Hist. Nat. Coq., III, 1824, p. 141.

* *Platiris (Spatha) dubia* LEA, Syn., 1852, p. 55; 1870, p. 89.

* *Mutela dubia* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 506, pl. CXIX; III, figs. 1, 1a.—* CLESSIN, Conch. Cab. Ano., 1873, p. 194, pl. XXV, fig. 3.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

* *Iridina dubia* CHENU, Man., 1859, II, p. 147, fig. 726.

* *Mytilus dubius* DILLWYN, part, Cat., I, 1817, p. 318.—* WOOD, Ind. Test. rev., 1856, p. 69, pl. XII, fig. 36.

West Africa.

¹ I agree with Lea that this = *exotica*. The teeth are not strongly developed.

² Gmelin refers to Adanson's figure. Dillwyn evidently has Gmelin's species and *Cristaria plicata* from China confounded. Two or three of these so-called species of *Mutela* are very close, and large series from different localities would probably show that they run together. There is much variation in the form and the development of hinge tubercles, even in a single species from a given locality.

* *MUTELA NILOTICA* Sowerby.

- * *Iridina nilotica* SOWERBY, Zool. Journal, I, 1835, p. 53, pl. II.—* CAILLIAUD, Voy. à Méroé, 1826, pl. LX, fig. 12.—* AUDOUIN, SAVIGNY, Icon. Moll. Egypt, 1827, pl. VII, fig. 2.—* CROUCH, Ill. Int. Lam., 1827, p. 17, pl. x, fig. 1.—* LEA, Syn., 1836, p. 56.—* HANLEY, Biv. Shells, 1843, p. 225.—* CATLOW and REEVE, Conch. Nom., 1845, p. 68.—* DESHAYES, Tr. Ele., II, 1853, p. 219, pl. XVII, figs. 6, 7.—* CHENU, Man., 1859, II, p. 148, fig. 727.—* SOWERBY, Conch. Icon., XVI, 1868, pl. II, fig. 4.
- * *Platiris (Spatha) nilotica* LEA, Syn., 1838, p. 33.
- * *Mytilus niloticus* WOOD, Ind. Test. Rev., 1856, p. 207, pl. II, sup. fig. 1.
- * *Mutela nilotica* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 506.—* JICKELI, Faun., 1874, p. 259.—* PÆTEL, Conch. Sam., III, 1890, p. 187.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 313.
- * ? *Mutela jouberti* BOURGUIGNAT, Un. et Ir., 1886, p. 28; * Icon. Mal., 1888, pl. XXIII, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 187.
- * ? *Mutela visseri* BOURGUIGNAT, Un. et Ir., 1886, p. 31.—* PÆTEL, Conch. [Sam., III, 1890, p. 187.
- * *Mutela vysseri* BOURGUIGNAT, Icon. Mal., 1888, pl. XXIII, fig. 3.

MUTELA NILOTICA var. *EMINI* von Martens.

- * *Mutela nilotica* var. *emini* VON MARTENS, Besch., 1897, p. 253.

Tropical Africa.

† *MUTELA ANGUSTATA* Sowerby.

- * *Iridina angustata* SOWERBY, Conch. Icon., XVI, 1868, pl. II, fig. 5.
- * *Mutela angustata* JICKELI, Faun. N. O. Af., 1874, p. 268.—* PÆTEL, Conch. Sam., III, 1890, p. 187.—* WESTERLUND, Faun. Pal., II, pl. 7, 1890, p. 312.

Tropical Africa.

† *MUTELA ALATA* Lea.

- * *Spatha alata* LEA, Pr. Ac. N. Sci. Phila., 1864, p. 109; * Jl. Ac. N. Sci. Phila., 1866, p. 35, pl. XII, fig. 31; * Obs., XI, 1867, p. 39, pl. XII, fig. 31.—* PÆTEL, Conch. Sam., III, 1890, p. 188.
- * *Platiris (Spatha) alata* LEA, Syn., 1870, p. 89.
- * *Mutela alata* CLESSIN, Conch. Cab. Ano., 1876, p. 196, pl. LXII, figs. 7, 8.—* ANCEY, Bull. Soc. Zool. de Fr., 1894, p. 232.—* VON MARTENS, Besch., 1897, p. 253.
- * *Burtonia alata* ANCEY, Bull. Soc. Mal. Fr., VI, 1889, p. 38.

Lake Nyassa.

MUTELA SIMPSONI Ancey.

- * *Mutela simpsoni* ANCEY, Bull. Soc. Zool. de Fr., VII, 1834, p. 233, fig. 8.
- * *Mutela alata* var. *simpsoni* VON MARTENS, Besch., 1897, p. 253.

Shire and Karonga rivers, near Lake Nyassa.

MUTELA BOURGUIGNATI Bourguignat.¹

- * *Mutela bourguignati* BOURGUIGNAT, Esp. Ouk., 1885, p. 8.—* SMITH, Ann. and Mag., X, 1892, p. 128, pl. XII, fig. 16.—* VON MARTENS, Besch., 1897, p. 255.

Lake Nyanza.

¹ Credited to Ancey, *in litt.*

(Group of *Mutela rostrata*.)

Shell somewhat inflated, thin, elongated, with a low, rounded posterior ridge ending in a point about midway up from the base; beaks low, smooth; epidermis bluish green, rayless; hinge line edentulous or having only the faintest vestiges of denticles; muscle scars large, shallow, indistinct; nacre bluish, shaded violet.

Animal the same as in typical *Mutela*.

† *MUTELA ROSTRATA* Rang.

* *Iridina rostrata* RANG, Nouv. Ann. Mus., 1835, p. 316.—* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 147, pl. LVI, fig. 1.

* *Spatha rostrata* VON MARTENS, Mal. Bl., XIII, 1866, p. 11.

* *Mutela rostrata* JICKELI, Faun., 1874, p. 269.—* WESTERLUND, Faun. Pal., II, Pt. 7, 1890, p. 312.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

* *Mutelina rostrata* BOURGUIGNAT, Bull. Soc. Zool. de Fr., II, 1886, p. 488.

* *Iridina celestis* LEA, Syn., 1836, p. 57; * Tr. Am. Phil. Soc., VI, 1838, p. 82, pl. XXII, fig. 70; * Obs., II, 1838, p. 82, pl. XXII, fig. 70.—* TROSCHEL, Arch. für Nat., V, 1839, Pt. 2, p. 239.—* HANLEY, Biv. Shells, 1843, p. 225.—* CATLOW and REEVE, Conch. Nom., 1845, p. 68.—* TROSCHEL, Arch. für Nat., XIII, 1847, I, p. 273.—* SOWERBY, Conch. Icon., XVI, 1868, pl. II, fig. 3.

* *Platiris (Spatha) celestis* LEA, Syn., 1838, p. 33; 1852, p. 55; 1870, p. 89.

* *Mutela celestis* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 506.—* CLESSIN, Conch. Cab. Ano., 1876, p. 193, pl. XXV, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

Tropical Africa.

Subgenus *PSEUDOMUTELA* Simpson, 1900.

(Type, *Mycetopus plicatus* Sowerby.¹)

Shell rather thin, elongate, wide, round and gaping behind, cut away and gaping at the anterior base; beaks low; surface rudely sulcate; a series of irregular nodulous projections extends down the low posterior ridge; hinge edentulous, with a kind of faint, broken internal ligament, with a wide prismatic streak behind; muscle scars distinct, united, the anterior protractor, however, separated and irregular; nacre lurid violet, iridescent behind.

† *MUTELA PLICATA* Sowerby.

* *Mycetopus plicatus* SOWERBY, Conch. Icon., XVI, 1868, pl. II, fig. 3; Icon. XVI, 1868, pl. II, fig. 3.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 8.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

* *Platiris (Mycetopus) plicatus* LEA, Syn., 1870, p. 90.

* *Mutela plicata* JICKELI, Faun. Moll. N. Ost-Af., 1874, p. 270.

Locality unknown, but Africa, no doubt. Sowerby credits it to Gray's manuscript in the British Museum.

¹ This is certainly not a *Mycetopoda*, but is evidently nearer *Mutela* than anything. I have placed it in that group as a subgenus, though it may be worthy of generic rank.

The following are unfigured *Mutelas*:

- * *Mutela lavigerina* BOURGUIGNAT, Un. and Ir., 1886, p. 26. Lake Tanganyika.
- * *Mutela moinei* BOURGUIGNAT, Un. et Ir., 1886, p. 27. Lake Tanganyika.
- * *Mutela subdiaphana* BOURGUIGNAT, Moll. Fluv. Ny., 1883, p. 5. Victoria Nyanza.
- * *Mutelina thotoni* ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 7. Kongo.
- * *Mutelina legumen* ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 6. Kongo.
- * *Mutelina mabilli* ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 7. Kongo.
- * *Mutelina paludicola* ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 8.
- * *Mutelina prasina* ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 7. Kongo.
- * *Iridina welwitschii* MORELET, Voy. de Angola, 1868, p. 98.

Genus CHELIDONOPSIS Ancey, 1887.

(Type, *Chelidoneura arietina* Rochebrune.)

Chelidoneura ROCHEBRUNE, S. B. Nat. Fr., 1886, p. 3.

Chelidonopsis ANCEY, Conch. Exchange, II, 1887, p. 22.

Shell elongated, thin, narrowed in front, gaping on the anterior basal part and winged on the front dorsal portion, rounded behind, with a pinched-up posterior ridge which develops into an open or closed tube near its hinder part; the shell ending in a diamond-shaped gap; epidermis shining slightly, concentrically grooved; hinge-line narrow, straight, with faint vestiges of denticles; a deep furrow inside marks the position of the posterior ridge; muscle scars faint; naere brilliant, iridescent.

Animal unknown.

CHELIDONOPSIS ARIETINA Rochebrune.

- * *Chelidoneura arietina* ROCHEBRUNE, Bull. Soc. Mal. Fr., III, 1886, p. 4, pl. I, figs. 1-4.

Gancini, Kongo.

†CHELIDONOPSIS HIRUNDO von Martens.

- * *Spatha hirundo* VON MARTENS, S. B. Nat. Fr., 1881, p. 122; *Conch. Mitth., III, 1885?, p. 139, pl. XXVII; ¹ Conch. Mitth., II, 1883, p. 139, pl. XXVII.

- * *Spatha (Mutela) hirundo* ROCHEBRUNE, Bull. Soc. Mal., Fr., III, 1886, p. 2, pl. I, figs. 5, 6.

Chelidoneura hirundo VON MARTENS, S. B. Nat. Fr., 1886, p. 161, pl. I, figs. 5, 6.

Kongo region.

Genus BRAZZÆA Bourguignat, 1885.

(Type, *Brazzaa anceyi* Bourguignat.²)

Brazzaa BOURGUIGNAT, Esp. Nouv. et Gen. Nouv., 1885, p. 321.

Shell thin, transparent, brilliant wine-colored, elliptical, greatly inflated, with a slight post-dorsal wing, and a high double posterior ridge; beaks smooth, compressed, but the region below them full;

¹ Probably the young of *arietina*.

² This appears to be a valid genus, belonging to the *Mutelidae*. I copy for the most part Bourguignat's description, which leaves out some essential characters. I have never seen any of the species.

hinge edentulous, filiform, with two ligaments, one internal, the other external; there are three groups of muscular impressions and several dorsal scars running in a row from the beak forward and downward; upper border of the left valve projecting over that of the right, forming a sinuous wing.

Animal unknown.

BRAZZÆA ANCEYI Bourguignat.

* *Brazzæa anceyi* BOURGUIGNAT, Esp. Onk. et Tan., 1885, p. 33; * Icon., 1888, pl. XXVIII, figs. 1-4.—* VON MARTENS, Besch., 1897, p. 258.

* *Brazzæa eximia* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 57; * Icon., 1888, pl. XXIX, fig. 5.

Lake Tanganyika.

BRAZZÆA BOURGUIGNATI Bourguignat.

* *Brazzæa bourguignati* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 58; * Icon., 1888, pl. XXVIII, figs. 5, 6.

Lake Tanganyika.

BRAZZÆA ELONGATA Bourguignat.

* *Brazzæa elongata* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 51; * Icon., 1888, pl. XXIX, figs. 2, 3.

Lake Tanganyika.

BRAZZÆA COULBOISI Bourguignat.

* *Brazzæa coulboisi* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 50; * Icon., 1888, pl. XXIX, fig. 1.

Lake Tanganyika.

BRAZZÆA VENTROSA Bourguignat.

* *Brazzæa ventrosa* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 45; * Icon., 1888, pl. XXIX, fig. 4.

Lake Tanganyika.

The following are unfigured species:

Brazzæa randabeli BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 46.

Brazzæa moineti BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 47.

Brazzæa jourdyi BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 48.

Brazzæa charbonnieri BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 52.

Brazzæa lavigerina BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 53.

Brazzæa bridouri BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 54.

Brazzæa newcombiana BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 55.

All from Lake Tanganyika.

Genus PLEIODON Conrad, 1834.

(Type, *Pleiodon oratus* Swainson.)

Pleiodon CONRAD, Jl. Ac. N. Sci. Phila., VII, 1834, p. 178.

Shell elliptical, inflated, solid, with a rounded posterior ridge, and full, smooth beaks; epidermis smooth, shining, generally rayless; hinge

¹ Said by Bourguignat to be of Jonbert *in litt.*

plate wide, nearly straight, set with strong, irregular, taxodont teeth throughout its length; beak cavities moderate, with a row of dorsal scars running obliquely downward and forward; anterior and posterior muscle scars well defined.

Animal with the palpi semilunar, united to the mantle by a straight border, longer than wide; outer gills generally larger than the inner, united their whole length to the mantle; inner united to the abdominal sac; genitalia occupying each side the lateral part of the visceral mass as far forward as the foot; mantle closed below the branchial siphon, and united into branchial and anal siphons; anal opening rather small; branchial opening large, the two separated by a solid bridge which solders together the extremities of the four branchiæ; both orifices have a thickened inner edge, the branchial being furnished with short tubercles or granules; mantle united below posteriorly one-fourth its length; foot large and strong; adductor muscles strong (Pelseneer).

Subgenus PLEIODON Conrad.

Characters as in the genus.

†PLEIODON OVATUS Swainson.

- * *Iridinā ovata* SWAINSON, Phil. Mag., LXI, 1823, p. 112.—* OKEN, Isis, 1834, p. 458.—* REEVE, Conch. Syst., I, 1841, p. 122, pl. XCIII.—* HANLEY, Biv. Shells, 1843, p. 225.—* CATLOW and REEVE, Conch. Nom., 1845, p. 68; * Elements of Conch., II, 1860, pl. XXXIII, fig. 184.—* CLESSIN, Conch. Cab. Ano., 1875, p. 230, pl. LXX, fig. 2; LXXI, fig. 2.
- * *Platiris (Iridina) ovata* LEA, Syn., 1838, p. 33; 1852, p. 54; 1870, p. 88.
- * *Pleiodon ovata* CONRAD, Jl. Ac. Nat. Sci. Phila., VII, 1854, p. 298; * H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 506.
- * *Pleiodon ovatus* CHENU, Man., 1859, II, p. 148, fig. 728.—* SOWERBY, Conch. Icon., XVI, 1866, pl. I, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 188.
- Iridina exotica* CHILDREN, Brande's Jl., XV, 1823, p. ?
- * *Pleiodon macmurtriei* CONRAD, Jl. Ac. Nat. Sci. Phila., VII, 1854, p. 13, pl. XIII; VII, 1854, p. 298.
- * *Margarita (Pleiodon) macmurtriei* LEA, Syn., 1836, p. 55.
- * *Iridina valeus* JAY, Cat., 1850, p. 70.¹
- * *Mutela valeus* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 506.—* PÆTEL, Conch. Sam., III, 1890, p. 187.
- * *Platiris (Iridina) leaii* LEA, Syn., 1852, p. 54; 1870, p. 88.²
- * *Pleiodon leaii* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 506.—* PÆTEL, Conch. Sam., III, 1890, p. 188.
- * *Pleiodon splendens* CONRAD, Jl. Ac. Nat. Sci. Phila., VII, 1854, p. 299.
- * *Iridina splendida* CHENU, Ill. Conch., 1858, pl. I, figs. 2, 2a, 2b, 2c, 2d.

West Africa.

The following are unfigured species of *Pleiodon*:

- * *Pleiodon diolibanus* BOURGUIGNAT, Moll. Egypt and Ab., 1879, p. 47.
- * *Pleiodon elongatus* BOURGUIGNAT, Moll. Egypt and Ab., 1879, p. 47. Senegal River.

¹ Said to be of Parreyss.

² Presented to Dr. Lea by Sowerby as *Iridina leaii*. Lea believes it to be a young *ovatus*.

* *Pleiodon letourneauianus* BOURGUIGNAT, Moll. Egypt and Ab., 1879, p. 48. Senegal River.

* *Pleiodon pachyodon* BOURGUIGNAT, Moll. Egypt and Ab., 1879, p. 43.

Subgenus CAMERONIA Bourguignat, 1879.

(Type, *Iridina spekii* Woodward.)

Anterior end of the hinge plate usually (not always) split up lengthwise into irregular teeth, which bear tubercles on their surfaces.¹

The mantle of *P. spekii* has a decided palleal sinus behind. (Pelseeneer.)

† PLEIODON SPEKII Woodward.

* *Iridina (Pleiodon) spekii* WOODWARD, Pr. Zool. Soc. Lond., 1859, p. 348, pl. XLVII, fig. 2; * Ann. and Mag., V, 1860, p. 337.

* *Pleiodon spekii* SOWERBY, Conch. Icon., XVI, 1866, pl. I, fig. 2.

* *Platiris (Iridina) spekii* LEA, Syn., 1870, p. 88.

* *Cameronia spekii* BOURGUIGNAT, Moll. Egypt and Ab., 1879, p. 43.

* *Iridina spekii* CLESSIN, Conch. Cab. An., 1875, p. 232, pl. LXX, fig. 2.

* *Pleiodon (Cameronia) spekii* CROSSE, Jl. de Conch, XXIX, 1881, p. 130.

* *Pleiodon spekii* SMITH, Proc. Zool. Soc., 1881, p. 296, pl. XXXIV, figs. 31, 31a.—

* PÆTEL, Conch. Sam., III, 1890, p. 188.

* *Mutela (Iridina) spekii* VON MARTENS, Besch., 1897, p. 256.

* *Cameronia gigantea* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 68; * Icon. Mal., 1888, pl. XXXV, fig. 1.

* *Cameronia admirabilis* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 69; * Icon. Mal., 1888, pl. XXXIV, fig. 1.

* *Cameronia coulboisi* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 77; * Icon. Mal., 1888, pl. XXXI, figs. 1, 2.

Cameronia josseti BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 82; * Icon. Mal., 1888, pl. XXXII, fig. 3.

* *Cameronia paradoxa* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 91; * Icon. Mal., 1888, pl. XXXII, fig. 1.

Lake Tanganyika.

PLEIODON LANDEAUI Bourguignat.

* *Cameronia landeui* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 74; * Icon. Mal., 1888, pl. XXXI, fig. 3.²

Lake Tanganyika.

PLEIODON BOURGUIGNATI Bourguignat.³

* *Cameronia bourguignati* BOURGUIGNAT, Esp. Ouk., 1885, p. 26; * Icon. Mal., 1888, pl. XXXIII.

Lake Tanganyika.

The following are unfigured species of *Cameronia*.

* *Cameronia anceyi* BOURGUIGNAT, Esp. Ouk., 1885, p. 30. Lake Tanganyika.

* *Cameronia bridouzi* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 71.

* *Cameronia charbonnieri* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 83. Lake Tanganyika.

¹The splitting up of the anterior end of the hinge is altogether irregular, and may be due to disease, as the hinge is sometimes entire.

²Probably a short, diseased *spekii*.

³Credited to Ancey in litt. by Bourguignat.

- * *Cameronia complanata* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 76. Lake Tanganyika.
- * *Cameronia dromauxi* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 84. Lake Tanganyika.
- * *Cameronia giraudi* BOURGUIGNAT, Nat. Prod., 1885, p. 107. Lake Tanganyika.
- * *Cameronia guillemeti* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 72. Lake Tanganyika.
- * *Cameronia jouberti* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 88. Lake Tanganyika.
- * *Cameronia lavigirina* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 85. Lake Tanganyika.
- * *Cameronia locardiana* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 78. Lake Tanganyika.
- * *Cameronia marioniana* BOURGUIGNAT, Esp. Ouk., 1885, p. 28. Lake Tanganyika.
- * *Cameronia mabilliana* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 86. Lake Tanganyika.
- * *Cameronia moineti* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 89. Lake Tanganyika.
- * *Cameronia pulchella* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 73. Lake Tanganyika.
- * *Cameronia obtusa* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 75. Lake Tanganyika.
- * *Cameronia randebeli* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 90. Lake Tanganyika.
- * *Cameronia revoiliana* BOURGUIGNAT, Moll. Tan., 1885, p. 107. Lake Tanganyika.
- * *Cameronia vyneckii* BOURGUIGNAT, Un. et Ir. Tan., 1886, p. 81. Lake Tanganyika.

Genus MONOCONDYLÆA d'Orbigny, 1835.

(Type, *Monocondylæa paraguayana* d'Orbigny.)

Aplodon SPIX (not of Rafinesque, 1818), Test. Fluv. Bras., 1827, pl. xxv, figs. 1, 2.
Monocondylæa D'ORBIGNY, Guerin Mag., 1835, p. 37.
Spiroconcha PILSBRY, Nautilus, VII, 1893, p. 30.

Shell rounded to obovate, rather solid, with a low, posterior ridge which is generally bordered by two or more dark, radiating bands; epidermis dull olive green to olive brown, cloth like; hinge with two irregular teeth under the beak in the left valve and two in the right, the posterior under the beak, the anterior in front of it and interlocking with those of the left valve, the whole generally more or less tuberculate; nacre soft, silvery, with iridescent shades; anterior scars united; posterior indistinct; prismatic layer wide.

Animal with the marsupium occupying the whole length of the inner branchiæ; gills very large, curved below, inner very much the larger, united the whole length of the abdominal sac; palpi small, round below, attached along their upper edge; mantle thick, greatly thickened at the edge, with a few papillæ in front of the branchial opening below; branchial opening large, with numerous small knob-like papillæ; anal opening large, without papillæ, separated completely from the branchial opening by a bridge; superanal opening not closed below.

(Group of *Monocondylæa guarayana*.)

Shell obovate, inflated, solid, often slightly produced just behind the center of the base; beaks full, turned inward and forward.

† MONOCONDYLÆA GUARAYANA d'Orbigny.

- * *Monocondylæa guarayana* D'ORBIGNY, Guer. Mag., 1835, p. 38; * Voy. Am. Mer., 1843, p. 614, pl. LXVIII, figs. 4-5.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 501; III pl. CXVII, fig. 3.—* CHENU, Man., 1859, II, p. 145, fig. 715.—* PÆTEL, Conch. Sam., III, 1890, p. 174.
- * *Margarita (Monocondylæa) guarayana* LEA, Syn., 1838, p. 28.
- * *Unio guarayana* HANLEY, Test. Moll., 1842, p. 213; * Biv. Shells, 1843, p. 213, pl. XXII, fig. 14.—* SOWERBY, Conch. Icon., XVI, 1868, pl. xcvi, fig. 524.
- * *Margaron (Monocondylæa) guarayana* LEA, Syn., 1852, p. 45; 1870, p. 73.

Bolivia.

MONOCONDYLÆA INERME Spix.

- * *Aplodon inerme* SPIX, Test. Fluv. Bras., 1827, p. 32, pl. xxv, figs. 1-2.—* VON IHERING, Arch. für Nat., 1890, p. 126, pl. IX, figs. 1-3.

South Brazil.

† MONOCONDYLÆA PARCHAPPII d'Orbigny.

- * *Moncondylæa parchappii* D'ORBIGNY, Guer. Mag., 1835, p. 38; * Voy. Am. Mer., 1843, p. 615, pl. LXVIII, figs. 1-3.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 501.—* PÆTEL, Conch. Sam., III, 1890, p. 174.
- * *Margarita (Monocondylæa) parchappii* LEA, Syn., 1838, p. 28.
- * *Unio parchappii* HANLEY, Test. Moll., 1842, p. 212; * Biv. Shells, 1843, p. 212, pl. XXII, fig. 13.—* CATLOW and REEVE, Conch. Nom., 1845, p. 62.
- * *Margaron (Monocondylæa) parchappii* LEA, Syn., 1852, p. 45; 1870, p. 73.
- *† *Monocondylæa pazii* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 34; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 273, pl. XXXVI, fig. 88; * Obs., XII, 1869, p. 36, pl. XXXVI, fig. 88.—* CLESSIN, Conch. Cab. Ano., 1876, p. 251, pl. LXXIX, figs. 6-7.—* PÆTEL, Conch. Sam., III, 1890, p. 174.
- * *Margaron (Monocondylæa) pazii* LEA, Syn., 1870, p. 73.

Argentina.

† MONOCONDYLÆA PARAGUAYANA d'Orbigny.

- * *Monocondylæa paraguayana* D'ORBIGNY, Guer. Mag., 1835, p. 37.—* SOWERBY, Conch. Man., 1839, fig. 149.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 612, pl. LXX, figs. 5-7.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 501.—* CLESSIN, Conch. Cab. Ano., 1876, p. 245, pl. LXXVIII, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 174.
- * *Margarita (Margaritana) paraguayana* LEA, Syn., 1838, p. 27.
- * *Unio paraguayana* HANLEY, Test. Moll., 1842, p. 212; * Biv. Shells, 1843, p. 212, pl. XXII, fig. 17.—* SOWERBY, Conch. Icon., XVI, 1866, pl. LI, fig. 273.
- * *Margaron (Monocondylæa) paraguayana* LEA, Syn., 1852, p. 45; 1870, p. 73.
- * *Unio paraguayanus* CATLOW and REEVE, Conch. Nom., 1845, p. 62.
- *† *Monocondylæa minuana* D'ORBIGNY, Guer. Mag., 1835, p. 37; * Voy. Am. Mer., 1843, p. 612, pl. LXX, figs. 8-10.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 501.—* PÆTEL, Conch. Sam., III, 1890, p. 174.
- * *Margarita (Margaritana) minuana* LEA, Syn., 1838, p. 28.
- * *Margaron (Monocondylæa) minuana* LEA, Syn., 1852, p. 45; 1870, p. 73.
- * *Unio minuanus* HANLEY, Test. Moll., 1842, p. 213; * Biv. Shells, 1843, p. 213, pl. XXII, fig. 18.—* SOWERBY, Conch. Icon., XVI, 1868, pl. xci, fig. 497.

Rio de la Plata drainage.

† MONOCONDYLÆA RETICULATA Moricand.

- * *Monocondylæa reticulata* MORICAND, Rev. et Mag., X, 1858, p. 453, pl. xv, fig. 2.
 * *Margaron (Monocondylæa) reticulata* LEA, Syn., 1870, p. 72.
 * *Margaritana reticulata* PÆTEL, Conch. Sam., III, 1890, p. 173.
 * *Unio reticulatus* SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVII, fig. 458.
 * *Aplodon reticulatus* VON IHERING, Arch. für Nat., 1893, p. 115.

Brazil.

MONOCONDYLÆA COSTULATA Moricand.

- * *Monocondylæa costulata* MORICAND, Rev. et Mag. Zool., X, 1858, p. 453, pl. xv, fig. 1;
 * SOWERBY, Conch. Icon., XVI, 1868, pl. LXXXVIII, fig. 470.—* PÆTEL, Conch. Sam., III, 1890, p. 174.
 * *Margaron (Monocondylæa) costulata* LEA, Syn., 1870, p. 72.

Brazil.

† MONOCONDYLÆA CORRIENTESSENSIS d'Orbigny.

- * *Monocondylæa corrientesensis* D'ORBIGNY, Guer. Mag., 1835, p. 38; * Voy. Am. Mer., 1843, p. 615, pl. LXVII, figs. 8–10.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 501.—* CLESSIN, Conch. Cab. Ano., 1876, p. 246, pl. LXXVIII, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 174.
 * *Margarita (Monocondylæa) corrientesensis* LEA, Syn., 1838, p. 28.
 * *Unio corrientesensis* HANLEY, Test. Moll., 1842, p. 212; * Biv. Shells, 1843, p. 212, pl. XXII, fig. 15.—* CATLOW and REEVE, Conch. Nom., 1845, p. 57.—* SOWERBY, Conch. Icon., XVI, 1868, pl. XCIII, fig. 509.
 * *Margaron (Monocondylæa) corrientesensis* LEA, Syn., 1852, p. 45; 1870, p. 73.

Argentina.

MONOCONDYLÆA JASPIDEA Hupe.

- * *Unio jaspidea* HUPE, An. Nouv., III, 1857, p. 83, pl. xvii, fig. 2.¹
 * *Margaron (Unio?) jaspideus* LEA, Syn., 1870, p. 37.

Amazon.

(Group of *Monocondylæa franciscana*.)

Shell moderately inflated, lenticular, rounded, not very solid; sometimes slightly produced at the post basal region; beaks rather low.

† MONOCONDYLÆA FRANCISCANA Moricand.

- * *Unio (Monocondylæa) franciscana* MORICAND, Mem. Soc. Hist. Nat. Gen., VIII, 1837, p. 39, pl. III, figs. 14–17.—* TROSCHEL, Arch. für Nat., VIII, 1842, Pt. 2, p. 404.
 * *Margarita (Margaritana) franciscana* LEA, Syn., 1838, p. 35.
 * *Margaron (Monocondylæa) franciscana* LEA, Syn., 1852, p. 45; 1870, p. 72.
 * *Monocondylæa franciscana* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 501.—* CLESSIN, Conch. Cab. Ano., 1876, p. 251, pl. LXXIV, figs. 4, 5.—* PÆTEL, Conch. Sam., III, 1890, p. 174.
 * *Aplodon franciscana* VON IHERING, Arch. für Nat., 1893, p. 115.

Brazil.

† MONOCONDYLÆA LENTIFORMIS Lea.

- * *Monocondylæa lentiformis* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 34; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 272, pl. XXXVI, fig. 86; * Obs., XII, 1869, p. 32, pl. XXXVI, fig. 86.—* CLESSIN, Conch. Cab. Ano., 1876, p. 250, pl. LXXIX, figs. 4, 5.—* PÆTEL, Conch. Sam., 1890, p. 174.

¹ The figure shows the teeth, which appear to be those of a *Monocondylæa*.

* *Margaron (Monocondylaea) lentiformis* LEA, Syn., 1870, p. 72.

* *Aplodon lentiformis* VON IHERING, Arch. für. Nat., 1893, p. 67.—* NEHRING, J. de Conch., 1894, p. 82.

Southern Brazil.

The following species is unknown to me:

* *Monocondylaea tamsana* DUNKER, Mal. Bl., V, 1858, p. 226.

Genus IHERINGELLA Pilsbry, 1893.

(Type, *Plagiodon isocardioides* Lea.)

Plagiodon LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 79.

Iheringella PILSBRY, Nautilus, VII, 1893, p. 30.

Shell solid, inflated, rounded to rhomboid in outline, with a more or less developed posterior ridge; beaks high, curved inward and forward, without sculpture; epidermis dull olive, cloth-like; hinge teeth imperfectly developed, nodulous or more or less broken into denticles, there being an irregular tooth in the left valve under the beak, sometimes partially bifid, and two in the right valve, with the pit between them under the beak; anterior muscle scars deep, partially united; posterior scars shallow; nacre soft, bluish silvery, iridescent behind.

Animal unknown.

(Group of *Iheringella isocardioides*.)

Shell rhomboid or *isocardia* shaped, with a high, distinct, strongly curved posterior ridge; beaks decidedly full and turned forward.

† IHERINGELLA ISOCARDIOIDES Lea.

* *Plagiodon isocardioides* LEA, Pr. Ac. N. Sci. Phila., VIII, 1856, p. 79.¹

* *Plagiodon isocardioides* LEA, Obs., VI, 1857, p. 38, pl. XXXII, fig. 32; * Jl. Ac. N. Sci. Phila., III, 1858, p. 318, pl. XXXII, fig. 32.—* KUSTER, Conch. Cab. Unio, 1862, p. 292, pl. XCVII, figs. 8-10.

* *Unio isocardioides* SOWERBY, Conch. Icon., XVI, 1868, pl. xc, fig. 484.—* PÆTEL, Conch. Sam., III, 1890, p. 156.

* *Margaron (Plagiodon) isocardioides* LEA, Syn., 1870, p. 71.

Rio de la Plata; Eastern Peru?

IHERINGELLA SEMISULCATA H. Adams.

* *Monocondylaea (Plagiodon) semisulcata* H. ADAMS, Pr. Zool. Soc. Lond., 1870, p. 376, pl. XXVII, fig. 3.

Eastern Peru.

(Group of *Iheringella rotundata*.)

Shell somewhat rounded, sublenticular, posterior ridge rather low.

† IHERINGELLA ROTUNDATA Mousson.

* *Plagiodon rotundatus* MOUSSON, Mal. Bl., XVI, 1869, p. 187.—* PFEIFFER, Nov. Conch., IV, 1876, p. 139, pl. CXXXI, figs. 8, 9.

South America.

IHERINGELLA BALZANI von Ihering.

* *Plagiodon balzani* VON IHERING, Arch. für Nat., 1893, p. 69, pl. III, fig. 3.

Rio Paraguay; San Paulo, Brazil.

Genus FOSSULA Lea, 1870.

(Type, *Monocondylva fossiculifera* d'Orbigny.)

Fossula LEA, Syn., 1870, p. 72 (footnote).

Shell obovate, inflated, solid, with rather high beaks and a low posterior ridge; epidermis olive-brown, somewhat smooth, sometimes slightly rayed; there is an irregular tooth in the left valve under the beak, and behind it a cavity; in the right valve there is a cavity under the beak, and an irregular tooth in front of and another behind it; the teeth and cavities are generally partly covered with brownish or amber-colored matter, and more or less pitted; anterior cicatrices well marked, united; posterior faint; prismatic border wide.

Animal unknown.

† FOSSULA FOSSICULIFERA d'Orbigny.

* *Monocondylva fossiculifera* D'ORBIGNY, Guer. Mag., 1835, p. 38; * Voy. Am. Mer., 1843, p. 614, pl. LXXX, figs. 5-7.—* H. and. A. ADAMS, Gen. Rec. Moll., II, 1857, p. 501.—* CLESSIN, Conch. Cab. An., 1876, p. 249, pl. XLIX, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 174.

* *Margarita (Margaritana) fossiculifera* LEA, Syn., 1838, p. 28.

* *Unio fossiculifera* HANLEY, Test. Moll., 1842, p. 213; * Biv. Shells, 1843, p. 213, pl. XXII, fig. 19.—* SOWERBY, Conch. Icon., XVI, 1868, pl. xcvi, fig. 521.

* *Margaron (Monocondylva) fossiculifera* LEA, Syn., 1852, p. 45; 1870, p. 73.

* *Fossula fossiculifera* VON IHERING, Arch. für Nat., 1893, p. 64, pl. III, fig. 2.

* *Unio fossiculiferus* CATLOW and REEVE, Conch. Nom., 1845, p. 59.

* *Anodonta fusciculifera* PÆTEL, Conch. Sam., III, 1890, p. 179.

Parana River, South America.

FOSSULA BALZANI von Ihering.

* *Fossula balzani* VON IHERING, Arch. für Nat., 1893, p. 65, pl. III, fig. 1.

Rio Paraguay; Rio Apae, South America.

Genus LEILA Gray, 1840.

(Type, *Anodonta blainvilliana* Lea.)

Columba LEA (not *Columba* Linnaeus, aves, 1758), Tr. Am. Phil. Soc., V, 1833, p. 78.

Leila GRAY, Syn. Brit. Mus., 1840, p. 142.

Shell large, obovate, inflated, with a straight hinge line which is produced into a slight wing before and behind, the posterior end bluntly pointed and somewhat truncate above, with a faint posterior ridge; beaks full, smooth, epidermis olive, generally smooth; hinge edentulous; beak cavities rather shallow; dorsal scars small and numerous, running in a straight line from behind the beaks downward and forward; posterior muscle scars united, large; pallial line generally showing a sinus.¹

¹ Conchologically this genus seems to be very closely related to *Glabaris*, especially to the group of *G. trapesialis*. Some of the shells of that genus have a slight dorsal wing in front and behind, and traces of a pallial sinus.

Animal having the mantle united behind into two short, separate, contractile siphons, according to Gray.¹

† LEILA BLAINVILLEANA Lea.²

* *Anodonta blainvilleana* LEA, Tr. Am. Phil. Soc., V, 1834, p. 77, pl. XII, fig. 35;

* Obs., I, 1834, p. 189, pl. XII, fig. 35.—* HANLEY, Test. Moll., 1842, p. 222;

* Biv. Shells, 1843, p. 222, pl. XXIV, fig. 18.

* *Margarita (Anodonta) blainvilleana* LEA, Syn., 1836, p. 53; 1838, p. 31.

* *Anodon blainvilleana* CATLOW and REEVE, Conch. Nom., 1845, p. 66.

* *Margaron (Anodonta) blainvilleana* LEA, Syn., 1852, p. 52.

* *Leila blainvilleana* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 507; III, pl. CXIX, figs. 4, 4a.—* CHENU, Man., 1859, II, p. 148, fig. 730.

* *Columba blainvilleana* CLESSIN, Conch. Cab. Ano., 1876, p. 253, pl. LXXXVI, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 188.

* *Anodon parishii* GRAY, Pr. Zool. Soc. Lond., 1834, p. 57.³—* MÖLLER, Syn. Nov. Gen., 1836, p. 195.—* CATLOW and REEVE, Conch. Nom., 1845, p. 67.

* *Margarita (Anodonta) parishii* LEA, Syn., 1836, p. 53; 1838, p. 31.

* *Anodonta parishii* HANLEY, Test. Moll., 1842, p. 222; * Biv. Shells, 1843, p. 222.—

* PÆTEL, Conch. Sam., III, 1890, p. 183.

* *Margaron (Anodonta) parishii* LEA, Syn., 1852, p. 52.

* *Leila parishii* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 507.

* *Anodon hians* SOWERBY, Conch. Icon., XVII, 1867, pl. IV, fig. 8.

Peru; Brazil; south to Argentina.

† LEILA ESULA d'Orbigny.

* *Iridina esula* D'ORBIGNY, Guerin Mag., 1835, p. 43; * Voy. Am. Mer., 1843, p. 597.

Anodonta esula JAN, Charpentier and Jan, Cat., 1837, p. 24.⁴

* *Margarita (Anodonta) esula* LEA, Syn., 1838, p. 32.

* *Margaron (Anodonta) esula* LEA, Syn., 1852, p. 52.

* *Margaron (Columba) esula* LEA, Syn., 1870, p. 86.

* *Leila esula* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 507.

* *Anodonta arcuata* HANLEY, Biv. Shells, 1843, p. 223.

* *Leila pulvinata* HUPE, Moll. Nouv., III, 1857, p. 90, pl. XX, fig. 1.

* *Columba pulvinata* CLESSIN, Conch. Cab. Ano., 1876, p. 255, pl. LXXV, figs. 1, 2.—

* PÆTEL, Conch. Sam., III, 1890, p. 188.

* *Anodon pulvinatus* SOWERBY, Conch. Icon., XVII, 1867, pl. V, fig. 10.

* *Leila castelnaudi* HUPE, Moll. Nouv., III, 1857, p. 91, pl. XIX, fig. 1.—* VON MARTENS, Mal. Bl., XV, 1868, p. 201.

* *Anodon castelnaudi* SOWERBY, Conch. Icon., XVII, 1868, pl. XX, fig. 79.

* *Columba castelnaudii* CLESSIN, Conch. Cab. Ano., 1876, p. 254, pl. LXXXIV, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 188.

Bolivia; Brazil; Paraguay.

¹ Ann. and Mag. Nat. Hist., VI, p. 316.

² The volume of the transactions in which Lea's name was published was issued, according to Scudder, in August or September, 1834. Gray read the paper containing the description of his *Anodon parishii* before the Zoological Society on July 8 of the same year, and the date of publication of that volume of the proceedings is 1834. As I do not know just when it appeared, and as Lea's is the better known name, I retain the latter.

³ Gray states in a letter to Lea, November, 1834, that his *A. parishii* and Lea's *blainvilliana* are the same.

⁴ According to Lea. I have not seen this paper. The species is quite commonly credited to Jan, but it was published by d'Orbigny under the name of *Iridina esula* in 1835, and I am not aware that Jan described it earlier than 1837.

LEILA SPIXII von Ihering.

**Anodon giganteus* SPIX (part), Test. Fluv. Bras., 1827, p. 27, pl. XIX, fig. 1 (young).

**Anodonta gigantea* KUSTER, Conch. Cab. Ano. 1853, p. 6, pl. I, fig. 2 (young).

**Columba spixii* VON IHERING, Arch. für Nat., 1890, p. 135, pl. IX, fig. 4.

Amazon River.

Genus GLABARIS Gray, 1847.

(Type, *Anodonta erotica* Lamarck.)

Glabaris GRAY, Pr. Zool. Soc. Lond., 1847, p. 197.

Shell rounded to elongated, inflated, subsolid; beaks full, smooth; epidermis smooth or clothlike, rarely having faint rays; hinge line straight or slightly curved, edentulous, sometimes a little sinuous, the escutcheon distinct and large; nacre soft tinted, the prismatic border wide and well defined.

Animal with the marsupium occupying the whole of the inner branchiæ, which are united their entire length to the abdominal sac; palpi generally semicircular or kidney shaped; attached along their entire upper length, not projecting posteriorly; branchial and anal openings with or without papillæ, not united into siphons in the specimens examined, separated by a strong bridge; superanal opening not closed below.

Section GLABARIS Gray, 1847

(Type, *Anodonta erotica* Lamarck.)

Shell rounded to elliptical; posterior ridge low or wanting.

(Group of *Glabaris patagonicus*.)

Shell solid, inflated, obovate, usually somewhat produced behind the center of the base; beaks quite full, projecting above the hinge line; epidermis olive brown, rather smooth, generally rayless; anterior muscle scars well defined.

Animal with the gills large, rounded below; palpi rather large; mantle thickened at edge and furnished with palpi in front of the branchial opening; branchial opening large, with numerous minute papillæ; anal opening large, with no papillæ.

† GLABARIS PATAGONICUS Lamarck.

**Anodonta patagonica* LAMARCK, An. sans. Vert., VI, 1819, p. 88.—* LAMARCK, Enc.

Méth., II, 1827, p. 147, pl. 203, fig. 1.—* STARK, Nat. Hist., II, 1828, p. 89.—

* HANLEY, Test. Moll., 1842, p. 221; * Biv. Shells, 1843, p. 221.—* KUSTER, Conch.

Cab. Ano., 1853, p. 50, pl. XII, fig. 2.—* H. and A. ADAMS, Gen. Rec. Moll., II,

1857, p. 504.—* VON MARTENS, Mal. Blatt., XV, 1868, p. 198.—* PÆTEL, Conch.

Sam., III, 1890, p. 183.

* *Margarita (Anodonta) patagonica* LEA, Syn., 1836, p. 52; 1838, p. 31.

* *Anodon patagonica* CATLOW and REEVE, Conch. Nom., 1845, p. 67.

* *Margaron (Anodonta) patagonica* LEA, Syn., 1852, p. 51; 1870, p. 82.

- * *Anodon trapezeus* SPIX, Test. Fluv. Bras., 1827, p. 28, pl. xx, fig. 1.
 * *Anodonta trapezeum* D'ORBIGNY, Voy. Am. Mer., 1843, pp. 6, 19.
 * *Anodonta trapezea* KUSTER, Conch. Cab. Ano., 1853, p. 7, fig. 3.
 * *Glabaris trapezea* VON JHERING, Arch. für Naturg., 1893, p. 57.
 * † *Anodonta lato-marginata* LEA, Tr. Am. Phil. Soc., V, 1834, p. 76, pl. xii, fig. 34;
 * Obs., I, 1834, p. 188, pl. xii, fig. 34.—* HANLEY, Test. Moll., 1842, p. 221; * Biv. Shells, 1843, p. 221, pl. xxiv, fig. 14.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 504.—* CHENU, Man., II, 1859, p. 146, fig. 724.—* CLESSIN, Conch. Cab., 1873, p. 76, pl. xxi, figs. 3, 4.
 * *Margarita (Anodonta) lato-marginata* LEA, Syn., 1836, p. 53; 1838, p. 31.
 * *Margaron (Anodonta) lato-marginata* LEA, Syn., 1852, p. 51; 1870, p. 82.
 * *Anodon lato-marginatus* SOWERBY, Conch. Icon., XVII, 1867, pl. ii, fig. 3.
 * *Anodon lati-marginata* CATLOW and REEVE, Conch. Nom., 1845, p. 67.
 * *Anodonta lati-marginata* STROBEL, Mat. Mal., Pt. 1, 1874, p. 67.—* PÆTEL, Conch. Sam., III, 1890, p. 181.
 * *Anodonta membranacea* D'ORBIGNY, Voy. Am. Mer., 1843, p. 616, pl. lxxix, fig. 11.
 * *Anodonta solida* KUSTER, Conch. Cab. Ano., 1853, p. 50, pl. xii, fig. 1.
 * ? † *Anodonta uruguayensis* LEA,¹ Pr. Ac. N. Sci. Phila., IV, 1860, p. 92; * Jl. Ac. N. Sci. Phila., V, 1863, p. 393, pl. xlviii, fig. 302; * Obs., X, 1863, p. 29, pl. xlviii, fig. 302.—* CLESSIN, Conch. Cab., 1874, p. 114, pl. xxxviii, figs. 1, 2.—* SOWERBY, Conch. Icon., XVII, 1870, pl. xxx, fig. 121,
 * *Margaron (Anodonta) uruguayensis* LEA, Syn., 1870, p. 83.
 * *Columbia uruguayensis* PÆTEL, Conch. Sam., III, 1890, p. 188.
 * *Anodonta sinuosa* CLESSIN, Conch. Cab. Ano., 1873, p. 90, pl. xxii, figs. 1, 2.
 * *Anodonta serpentina* CLESSIN, Conch. Cab. Ano., 1876, p. 223, pl. lxxv, figs. 2, 3.—* PÆTEL, Conch. Sam., III, 1890, p. 184.

GLABARIS PATAGONICUS var. FELIX Pilsbry.

- * *Glabaris lato-marginatus* LEA, var. *felix* PILSBRY, Pr. Ac. N. Sci. Phila., 1869, p. 563, pl. xxvi, fig. 8.

† Southern South America, east of the Andes; the variety in Colonia, Uruguay.

GLABARIS CRASSUS Swainson.

- * *Anodon crassus* SWAINSON, Zool. Ill., 1st ser., III, 1823, pl. clxvii.
 * *Margarita (Anodonta) crassa* LEA, Syn., 1836, p. 23; 1838, p. 32.
 * *Anodonta crassa* HANLEY, Test. Moll., 1842, p. 222; * Biv. Shells, 1843, p. 222.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 504.—* PÆTEL, Conch. Sam., III, 1890, p. 178.
 * *Anodon crassa* CATLOW and REEVE, Conch. Nom., 1845, p. 66.
 * *Margaron (Anodonta) crassa* LEA, Syn., 1852, p. 52; 1870, p. 83.

Rio de la Plata.

† GLABARIS WYMANII Lea.

- * *Anodonta wymanii* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 91; * Jl. Ac. N. Sci. Phila., V, 1863, p. 390, pl. xliv, fig. 294; * Obs., X, 1863, p. 26, pl. xliv, fig. 294.—* CLESSIN, Conch. Cab. Ano., 1873, p. 104, pl. xxxii, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 186.

¹ An old worn shell—the type—is the only one in the Lea collection, save a few very young that are doubtfully the same. It is a little more inflated than most specimens that are called *lato-marginata*, but not more than some are, and I hardly think it distinct.

* *Margaron (Anodonta) wymanii* LEA, Syn., 1870, p. 80.

* *Anodon wymani* SOWERBY, Conch. Icon., XVII, 1870, pl. xxx, fig. 117.

Uruguay River, South America.

† GLABARIS SIRIONIS d'Orbigny.

* *Anodonta sirionis* D'ORBIGNY, Mag. Zool., 1835, p. 40.—* HANLEY, Test. Moll., 1842, p. 221; * Biv. Shells, 1843, p. 221.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 615, pl. LXXIV, figs. 4-6; LXXX, figs. 1-4.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 504.—* CLESSIN, Conch. Cab. Ano., 1874, p. 128, pl. XLI, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 184.

* *Margarita (Anodonta) sirionis* LEA, Syn., 1838, p. 31.

* *Anodon sirionis* CATLOW and REEVE, Conch. Nom., 1845, p. 68.

* *Margaron (Anodonta) sirionis* LEA, Syn., 1852, p. 51; 1870, p. 82.

* *Anodonta ferrarisii* D'ORBIGNY, Guér. Mag., 1835, p. 40.

Rio de la Plata

† GLABARIS PAZII Lea.

* *Anodonta pazii* LEA, Pr. Ac. N. Sci. Phila., X, 1866, p. 35; * JI. Ac. N. Sci. Phila., VI, 1868, p. 274, pl. xxxvi, fig. 87; * Obs., XII, 1869, pl. xxxvi, fig. 87.—

* CLESSIN, Conch. Cab. Ano., 1874, p. 139, pl. XLIII, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 183.

* *Margaron (Anodonta) pazii* LEA, Syn., 1870, p. 81.

South America.

† GLABARIS RUBICUNDUS Lea.

* *Anodonta rubicunda* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 92; * JI. Ac. N. Sci. Phila., V, 1863, p. 392, pl. XLVI, fig. 299; * Obs., X, 1863, p. 28, pl. XLVI, fig. 299.—* CLESSIN, Conch. Cab. Ano., 1876, p. 106, pl. xxxii, figs. 5, 6.—* PÆTEL, Conch. Sam., III, 1890, p. 184.

* *Anodon rubicundus* SOWERBY, Conch. Icon., XVII, 1870, pl. xxx, fig. 118.

* *Margaron (Anodonta) rubicunda* LEA, Syn., 1870, p. 75.

Uruguay River, South America.

† GLABARIS ROTUNDUS Spix.

* *Anodon rotundus* SPIX, Test. Fluv. Bras., 1827, p. 28, pl. xx, figs. 2-4.

* *Anodon rotunda* KUSTER, Conch. Cab. Ano., 1853, p. 33, pl. viii, fig. 1.

* *Margaron (Anodonta) rotunda* LEA, Syn., 1870, p. 81.

* *Glabaris rotunda* VON IHERING, Arch. für Nat., 1893, p. 59.

* † *Anodonta cailliaudii* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 308; * JI. Ac. N. Sci. Phila., V, 1863, p. 395, pl. XLV, fig. 297; * Obs., X, 1863, p. 31, pl. XLV, fig. 297.—* CLESSIN, Conch. Cab. Ano., 1873, p. 105, pl. xxxii, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 177.

* *Anodon cailliaudi* SOWERBY, Conch. Icon., XVII, 1867, pl. xii, fig. 38.

* *Margaron (Anodonta) cailliaudii* LEA, Syn., 1870, p. 81.

Brazil.

GLABARIS MEMBRANACEUS Maton.¹

* *Mytilus membranaceus* MATON, Tr. Linn. Soc. Lond., 1811, p. 329, pl. xxiv, figs. 11, 12.

¹ It is difficult to tell what this is, but I believe it to be some species of *Glabaris*, from the fact that Maton says that the hinge is edentulous. The figure (11) plainly shows it to be destitute of radial beak sculpture, though the umbo is perfect. The specimens figured are evidently young.

- * *Margarita (Anodonta) membranacea* LEA, Syn., 1836, p. 22; 1838, p. 23.
 * *Margaron (Anodonta) membranacea* LEA, Syn., 1852, p. 34; 1870, p. 55.
 * *Unio membranaceus* HANLEY, Test. Moll., 1842, p. 202.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 492.—* PÆTEL, Conch. Sam., III, 1890, p. 159.
 * *Unio subtrapezius* PHILIPPI, Zeits. für Mal., IV, 1847, p. 96; * Abbild., III, 1848, p. 80, pl. v, fig. 3.—* KUSTER, Conch. Cab. Unio, 1862, p. 283, pl. xcv, fig. 4.

Rio de la Plata.

† *GLABARIS IHERINGI* Clessin.

- * *Anodonta iheringi* CLESSIN, Mal. Bl., V, 1882, p. 191, pl. IV, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 180.¹

(Group of *Glabaris crispatus*.)

Shell elliptical obovate, slightly produced at the posterior base, and straight or very feebly incurved in front of it; epidermis fuscous or tawny, cloth-like behind, somewhat rayed by more or less incised lines in front, where it is wrinkled like dried paint, the wrinkles often being looped; nacre lurid, bluish, somewhat iridescent.

Animal unknown.

† *GLABARIS CRISPATUS* Bruguiere.

- * *Anodontites crispata* BRUGUIERE, Jl. de Hist. Nat., I, 1792, p. 131.
 * *Anodonta crispata* LAMARCK, An. sans Vert., VI, 1819, p. 86.²—* LEA, Obs., I, 1834, p. 205.—* HANLEY, Test. Moll., 1842, p. 217; * Biv. Shells, 1843, p. 217.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* PÆTEL, Conch. Sam., III, 1890, p. 178.
 * *Margarita (Anodonta) crispata* LEA, Syn., 1836, p. 48; 1838, p. 29.
 * *Anodon crispata* CATLOW and REEVE, Conch. Nom., 1845, p. 66.
 * *Margaron (Anodonta) crispata* LEA, Syn., 1852, p. 47; 1870, p. 75.
 * ? *Anodonta crispa* LAMARCK, Ene. Meth., II, 1827, p. 147, pl. CCIII, fig. 3.
 * † *Anodonta puberula* GOULD, U. S. Expl. Ex., XII, 1852, p. 434, figs. 548, 548a, 548b.³
 * *Anodon reticulatus* SOWERBY, Conch. Icon., XVII, 1867, pl. x, fig. 27.
 * *Margaron (Anodonta) reticulata* LEA, Syn., 1870, p. 80.
 * *Anodonta reticulata* PÆTEL, Conch. Sam., III, 1890, p. 184.

Widely distributed in tropical South America.

GLABARIS PHILIPPIANUS Simpson.⁴

- * *Anodonta subsinuata* PHILIPPI, Mal. Bl., XVI, 1869, p. 41.—* PFEIFFER, Nov. Conch., III, 1869, p. 487, pl. CV, figs. 7, 8.—* PÆTEL, Conch. Sam., III, 1890, p. 185.

Ucayali River, Peru.

¹ This species combines characters of the *Patagonicus* and *Tenebricosus* groups.

² Lamarck refers to Encyclopaëdie Méthodique, pl. CCIII, figs. 3, 3a, 3b.

³ According to Lea's note in pencil on the margin of this description the species = *crispata*. I think he is right.

⁴ The name *subsinuata*, applied to this by Philippi, will have to be placed in the synonymy, as Sowerby used it previously for a *Glabaris* which he placed in *Anodonta*.

† GLABARIS NAPOENSIS Lea.

- * *Anodonta napoensis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 162; * JI. Ac. N. Sci. Phila., VI, 1868, p. 324, pl. LIII, fig. 137; * Obs., XII, 1869, p. 84, pl. LIII, fig. 137.—* CLESSIN, Conch. Cab. Ano., 1873, p. 116, pl. XXXIV, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 182.

* *Margaron (Anodonta) napoensis* LEA, Syn., 1870, p. 80.

River Napo, Ecuador.

† GLABARIS TORTILIS Lea.

- * *Anodonta tortilis* LEA, Tr. Am. Phil. Soc., X, 1852, p. 291, pl. XXVIII, fig. 54; * Obs., V, 1852, p. 47, pl. XXVIII, fig. 54.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1875, p. 166, pl. LV, figs. 7, 8.—* PÆTEL, Conch. Sam., III, 1890, p. 185.

* *Margaron (Anodonta) tortilis* LEA, Syn., 1852, p. 47; 1870, p. 75.

* *Anodon tortilis* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXVII, fig. 154.

Cartagena, Colombia.

† GLABARIS LUTEOLUS Lea.

- * *Anodonta luteola* LEA, Pr. Ac. N. Sci. Phila., II, 1858, p. 118; JI. Ac. N. Sci. Phila., IV, 1860, p. 267, pl. XLIII, fig. 147; * Obs., VII, 1860, p. 85, pl. XLIII, fig. 147.—* CLESSIN, Conch. Cab. Ano., 1874, p. 122, pl. XXXVII, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 181.

* *Margaron (Anodonta) luteola* LEA, Syn., 1870, p. 83.

* *Anodon luteolus* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXIII, fig. 132.

Isthmus of Darien; Nicaragua; Costa Rica.

† GLABARIS SCHOMBURGIANUS Sowerby.

- * *Anodon schomburgianus* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXIV, fig. 137.

* *Anodonta schomburgiana* CLESSIN, Conch. Cab. Ano., 1876, p. 235, pl. LXXVII, fig. 4.

British Guiana.

† GLABARIS STREBELII Lea.

- * *Anodonta strebelii* LEA, Pr. Ac. N. Sci. Phila., XX, 1868, p. 150; * JI. Ac. N. Sci. Phila., VII, 1868, p. 322, pl. LII, fig. 135; * Obs., XII, 1869, p. 82, pl. LII, fig. 135.—* CLESSIN, Conch. Cab. Ano., 1874, p. 138, pl. XLII, figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 185.

* *Margaron (Anodonta) strebelii* LEA, Syn., 1870, p. 79.

Mexico.

† GLABARIS CYLINDRACEUS Lea.

- * *Margarita (Anodonta) cylindracea*, LEA, Syn., 1836, p. 54; 1838, p. 32.

* *Anodonta cylindracea* LEA, Tr. Am. Phil. Soc., VI, 1838, p. 45, pl. XIII, fig. 40;

* Obs., II, 1838, p. 45, pl. XIII, fig. 40.—* TROSCHEL, Arch. für Nat., V, 1839, Pt. 2, p. 238.—* HANLEY, Test. Moll., 1842, p. 223; Biv. Shells, 1843, p. 223.—

* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 263.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1874, p. 161, pl. LIII, figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 178.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 528.

* *Anodon cylindracea* CATLOW and REEVE, Conch. Nom., 1845, p. 66.—* SOWERBY, Conch. Icon., XVII, 1869, pl. XXIV, fig. 93.

* *Margaron (Anodonta) cylindracea* LEA, Syn., 1852, p. 51; 1870, p. 82.

Medellin River, Mexico.

† GLABARIS PUELCHANUS d'Orbigny.

- * *Anodonta puelchana* D'ORBIGNY, Guer. Mag., 1835, p. 40; Voy. Am. Mer., 1843, p. 620, pl. LXXIX, figs. 7-9.—* CHENU, III. Conch., 1858, pl. III, figs. 7, 7a.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* STROBEL, Mal. Argent., 1874.—* CLESSIN, Conch. Cab. Ano., 1874, p. 130, pl. XLI, figs. 5, 6.—* PÆTEL, Conch. Sam., III, 1890, p. 183.
- * *Margaron (Anodonta) puelchana* LEA, Syn., 1852, p. 51; 1870, p. 81.
- * *Anodonta obtusula* HUPE, Moll. Nouv., III, 1857, p. 87, pl. XVII, fig. 3.—* CLESSIN, Conch. Cab. Ano., 1875, p. 214, pl. LXVI, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 182.
- * *Anodon obtusula* SOWERBY, Conch. Icon., XVII, 1868, pl. XX, fig. 81.
- * *Margaron (Anodonta) obtusula* LEA, Syn., 1870, p. 79.
- * *Glabaris obtusula* VON IHERING, Arch. für Nat., 1893, p. 119.

Patagonia, north to the Amazon; west to Bolivia.

† GLABARIS LIMNOICUS d'Orbigny.

- * *Anodonta limnoica*, D'ORBIGNY, Guer. Mag., 1835, p. 40.—* HANLEY, Test. Moll., 1842, p. 218.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 619, pl. LXXIX, figs. 1-3.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. Ano., 1874, p. 129, pl. XLI, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 181.
- * *Anodon limnoica* CATLOW and REEVE, Conch. Nom., 1845, p. 67.
- * *Margarita (Anodonta) limnoica* LEA, Syn., 1838, p. 30.
- * *Margaron (Anodonta) limnoica* LEA, Syn., 1852, p. 50; 1870, p. 79.
- * *Anodonta limnoica* CHENU, III. Conch., 1858, pl. III, figs. 1, 1a, 1b.

Argentina.

GLABARIS LUCIDUS d'Orbigny.¹

- * *Anodonta lucida* D'ORBIGNY, Guer. Mag., 1835, p. 40; * Voy. Am. Mer., 1843, p. 620, pl. LXXIX, fig. 4-6.—* CHENU, III. Conch., 1858, pl. III, figs. 6, 6a, 6b.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1875, p. 208, pl. LXIX, figs. 4, 5.—* PÆTEL, Conch. Sam., III, 1890, p. 181.
- * *Margaron (Anodonta) lucida* LEA, Syn., 1852, p. 51; 1870, p. 82.

Uruguay.

GLABARIS INCARUM Philippi.

- * *Anodonta incarum* PHILIPPI, Mal. Bl., XVI, 1869, p. 40.—* PFEIFFER, Nov. Conch., III, 1869, p. 488, pl. cv., figs. 9-11.—* PÆTEL, Conch. Sam., III, 1890, p. 180.²

Peru.

† GLABARIS HOLTONIS Lea.

- * *Anodonta holtonis* LEA, Pr. Ac. N. Sci. Phila., I, 1857, p. 85; * Jl. Ac. N. Sci. Phila., III, 1857, p. 316, pl. XXXII, fig. 31; * Obs., VI, 1847, p. 36, pl. XXXII, fig. 31.—* CLESSIN, Conch. Cab. Ano., 1874, p. 149, pl. L., figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 180.
- * *Anodon holtonis* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXVI, fig. 147.
- * *Margaron (Anodonta) holtonis* LEA, Syn., 1870, p. 79.

Colombia.

¹ It is not unlikely that this and the two preceding species are variations of one thing.

² Probably a form of *G. puelchanus* d'Orbigny.

(Group of *Glabaris inaequalvis*.)

Shell elliptical, moderately solid and inflated, with a low posterior ridge, slightly produced behind near the base and truncated above the posterior slope; beaks full, smooth; ligament imbedded in a sort of groove which extends to the anterior point of the shell; epidermis slightly roughened, often clothlike and showing the rest periods, having faint radiating lines; nacre bluish, soft, but not brilliant.

Animal unknown.

†GLABARIS INÆQUALVIS Lea.

* *Anodonta inaequalvis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 95; * JI. Ac. N. Sci. Phila., VI, 1868, p. 292, pl. XLIII, fig. 108; * Obs., XII, 1869, p. 52, pl. XLIII, fig. 108.—* CLESSIN, Conch. Cab. Ano., 1874, p. 137, pl. XLIV, figs. 7, 8.—* B. H. WRIGHT, Check List, 1888.

* *Margaron (Anodonta) inaequalvis* LEA, Syn., 1870, p. 83.

*† *Anodonta lenticularis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 95; * JI. Ac. N. Sci. Phila., VI, 1868, p. 290, pl. XLI, fig. 102; * Obs., XII, 1869, p. 50, pl. XLI, fig. 102.—* CLESSIN, Conch. Cab. Ano., 1874, p. 134, pl. XLIII, figs. 5, 6.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 181.

* *Margaron (Anodonta) lenticularis* LEA, Syn., 1870, p. 81.

* *Anodon glabrus* SOWERBY, Conch. Icon., XVII, 1870, pl. xxv, fig. 97.

* *Anodon montezianus* SOWERBY, Conch. Icon., XVII, 1870, pl. xxxvi, fig. 145.

* *Anodonta viridana* CLESSIN, Conch. Cab. Ano., 1876, p. 226, pl. LXXV, fig. 5.—* PÆTEL, Conch. Sam., III, 1890, p. 186.—* FISCHER and CROSSE, Miss. Sci. Pt. 7, II, 1894, p. 521.

Lake Nicaragua; Mexico.

†GLABARIS GRANADENSIS Lea.

* *Anodonta granadensis* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 96; * JI. Ac. N. Sci. Phila., VI, 1868, p. 288, pl. XLI, fig. 100; * Obs., XII, 1869, p. 48, pl. XLI, fig. 100.—* CLESSIN, Conch. Cab. Ano., 1874, p. 134, pl. XLIV, figs. 3, 4.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 180.

* *Margaron (Anodonta) granadensis* LEA, Syn., 1870, p. 81.

Lake Nicaragua, Central America.

†GLABARIS TRAUTWINIANUS Lea.

* *Anodonta trautwiniana* LEA, Tr. Am. Phil. Soc., X, 1852, p. 287, pl. xxvi, fig. 48; * Obs., V, 1852, p. 43, pl. xxvi, fig. 48.

* *Margaron (Anodonta) trautwiniana* LEA, Syn., 1870, p. 81.

* *Anodon trautwinianus* SOWERBY, Conch. Icon., XVII, 1870, pl. xxxiii, fig. 134.

* *Margaron (Anodonta) trautwiniana* LEA, Syn., 1852, p. 51.

* *Anodonta trautwiniana* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1874, p. 112, pl. xxxvi, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 186.

* ? *Anodon oratus* SWAINSON, Ex. Conch., 1841, pl. xxxvi.

? *Anodonta carthagena* ———, Who? Where?

Cartagena, Colombia.

GLABARIS MONTEZUMA Lea.¹

**Anodonta montezuma* LEA, Pr. Am. Phil. Soc., II, 1841, p. 31; *Tr. Am. Phil. Soc., 1842, p. 240, pl. XXIII, fig. 55; *Obs., III, 1842, p. 78, pl. XXIII, fig. 55.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 504.—*CLESSIN, Conch. Cab. Ano., 1876, p. 220, pl. LXXII, figs. 7, 8.—*B. H. WRIGHT, Check List, 1888.—*PÆTEL, Conch. Sam., III, 1890, p. 182.

**Margaron (Anodonta) montezuma* LEA, Syn., 1852, p. 51; 1870, p. 82.

Central America.

(Group of *Glabaris trapesialis*.)

Shell large, inflated, elliptical, narrower in front, with full, rather prominent beaks, a slight wing on the post dorsal part, and sometimes a small one in front of the beaks; epidermis smooth, generally uniform olive green; hinge line straight or slightly sinuous; escutcheon large and conspicuous; nacre bluish silvery, sometimes marked with parallel, wavy, dark lines.

Animal with the marsupium filling the inner branchiæ; gills large, inner the larger; palpi very large, reniform; mantle thin, with a wide, thickened edge; branchial opening small, plicate, but not crenulate or papillose; anal opening large, smooth, separated from the branchial by a strong bridge.

† GLABARIS TRAPESIALIS Lamarck.

**Anodonta trapesialis* LAMARCK, An. sans. Vert., VI, 1819, p. 87.

**Iridina trapesialis* D'ORBIGNY, Guer. Mag., 1835, p. 43².

**Glabaris trapesialis* PILSBY, Pr. Ac. N. Sci. Phila., 1896, p. 563.

**Anodonta trapezialis* BLAINVILLE, Man. de Mal. et Conch., 1825, p. 538, fig. 1.—

*DESHAYES, Enc. Meth., II, 1827, p. 147, pl. CCV, fig. 1.—*WYATT, Man. Conch., 1838, p. 68, pl. XI, fig. 3.—*HANLEY, Test. Moll., 1842, p. 220; *Biv. Shells, 1843, p. 220.—*KUSTER, Conch. Cab. Ano., 1853, p. 31, pl. VII, fig. 4.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 504.—*CLESSIN, Conch. Cab. Ano., 1876, p. 235, pl. LXXVII, fig. 5.

**Anodon trapezialis* CROUCH, Ill. Int. Lam., 1827, p. 16, pl. IX, fig. 7.—*CATLOW and REEVE, Conch. Nom., 1845, p. 68.

**Margarita (Anodonta) trapezialis* LEA, Syn., 1836, p. 53; 1838, p. 31.

**Margaron (Anodonta) trapezialis* LEA, Syn., 1852, p. 52; 1870, p. 82.

**Columba trapesialis* PÆTEL, Conch. Sam., III, 1890, p. 188.

**Anodon giganteus* SPIX (part), Test. Flav. Bras., 1827, p. 27, pl. XIX, fig. 2.

**Anodon gigantea* SOWERBY, Conch. Icon., XVII, 1867, pl. VIII, fig. 18.

**Anodonta gigantea* VON MARTENS, Mal. Bl., XV, 1868, p. 196.

**Columba gigantea* PÆTEL, Conch. Sam., III, 1890, p. 188.

*? *Anodon penicillatus* GRAY,³ Pr. Zool. Soc. Lond., 1834, p. 57.—*MÜLLER, Syn. Nov. Gen., 1836, p. 195.

¹ I have never seen this shell, and can not be positive from the figure and description whether it is a *Glabaris* or an *Anodonta*. I incline to think it the former, and that it belongs in this group.

² = *Anodontites trapesialis* Lamarck according to d'Orbigny in above, but in Voy. Am. Mer. he says it is not that.

³ I have never seen this shell. Dr. Lea believes it to = *trapesialis* Lamarck.

**Anodonta penicillata* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—

*PÆTEL, Conch. Sam., III, 1890, p. 183.

**Anodon susannæ* GRAY in Griffith, Cuv., XII, 1834, pl. XXIV, fig. 1.¹—*SOWERBY, Conch. Icon., XVII, 1867, pl. ix, fig. 21.

**Anodonta susannæ* PÆTEL, Conch. Sam., III, 1890, p. 185.

*†*Anodon ciconia* GOULD, Pr. Bost. Soc. Nat. Hist., IV, 1851, p. 92.²—*?GOULD and CARPENTER, Pr. Zool. Soc. Lond., 1856, p. 202.—*OTIA, Conch., 1862, p. 213.—*PÆTEL, Conch. Sam., III, 1890, p. 177.—*?FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 534, pl. LXVIII, figs. 2, 2a.

**Anodon blainvilliana* SOWERBY, Conch. Icon., XVII, 1867, pl. vi, fig. 2.

† GLABARIS TRAPESIALIS var. ANSERINUS Spix.³

**Anodon anserinus* SPIX, Test. Fluv. Bras., 1827, p. 29, pl. XVII, figs. 1, 2.—*SOWERBY, Conch. Icon., XVII, 1870, pl. XXXI, fig. 125.

**Anodon anserina* CATLOW and REEVE, Conch. Nom., 1845, p. 66.

**Margarita (Anodonta) anserina* LEA, Syn., 1838, p. 31.

**Anodonta anserina* HANLEY, Test. Moll., 1842, p. 222; *Biv. Shells, 1843, p. 222.—*H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 504.—*CHENU, Mau., 1859, II, p. 146.—*CLESSIN, Conch. Cab. Ano., 1873, p. 80, pl. XX, fig. 1.—*PÆTEL, Conch. Sam., III, 1890, p. 176.

**Margaron (Anodonta) anserina* LEA, Syn., 1852, p. 52; 1870, p. 83.

† GLABARIS TRAPESIALIS var. EXOTICUS Lamarck.

**Anodonta exotica* LAMARCK, An. sans Vert., VI, 1819, p. 87.—*STARK, Nat. Hist., II, 1828, p. 89.—*DELESSERT, Rec. Coq. Lam., 1841, pl. XIII, fig. 1.—*HANLEY, Test. Moll., 1842, p. 217; *Biv. Shells, 1843, p. 217.—*D'ORBIGNY, Voy. Am. Mer., 1843, p. 621.—*CHENU, Ill. Conch., 1858, pl. III, fig. 2.—*VON MARTENS, Mal. Bl., XV, 1868, p. 197.—*STROBEL, Mat. Mal., Pt. 1, 1874, p. 66.—*CLESSIN, Conch. Cab. Ano., 1875, p. 178, pl. LVIII, figs. 1, 2.

**Anodon exotica* CATLOW and REEVE, Conch. Nom., 1845, p. 66.

**Columba exotica* PÆTEL, Conch. Sam., III, 1890, p. 188.

† GLABARIS TRAPESIALIS var. SCRIPTUS Sowerby.

**Anodon scriptus* SOWERBY, Conch. Icon., XVII, 1867, pl. iv, fig. 9.⁴

**Anodon subsinuatus* SOWERBY, Conch. Icon., XVII, 1867, pl. vii, fig. 14.

¹This is briefly described in the index, p. 595, as *Anodon susannæ*. In the text the genus is called *Anodontites*.

²Two specimens of this, a large and a smaller one, are in the Gould collection. Lea places the species in the synonymy of *G. trapesialis* Lamarck., to which Lewis objects in a note on the back of Gould's label on account of its locality—Mexico. After carefully comparing these shells with all our *trapesialis* in the Museum collections I am unable to separate the two. Carpenter states (Mazatlan Shells, p. 117) that it is found rather plentifully at Mazatlan. I do not believe that the locality given for Gould's shells is correct (it is referred to Mexico with doubt by its author), and have no doubt that they are from South America. The species alluded to by Carpenter is probably *G. glauca* Valenciennes, which is closely related to *G. trapesialis*.

³*Glabaritis trapesialis* is a very abundant, widely distributed species, and shows a great number of variations. Some of these are probably worthy of varietal names, others seem to be so mixed with various forms that they are scarcely worthy of any designation.

⁴A brownish, somewhat elongated form, much narrowed in front, receiving its name from having dark purplish letter-like marks on the nacre. There is every possible development of this character from shells in which it is a marked feature to those without it at all.

- * *Anodon subsinuatus* SOWERBY, Conch. Icon., XVII, 1867, pl. VIII, fig. 15.
 * *Anodon areolatus* SOWERBY, Conch. Icon., XVII, 1867, pl. x, fig. 28.
 * *Anodonta bahiensis* KUSTER, Conch. Cab. Ano., 1873, p. 94, pl. XX, fig. 2.—*PÆTEL, Conch. Sam., III, 1890, p. 176.
 * *Glabaris bahiensis* VON IHERING, Arch. für Naturg., 1893, p. 115.

GLABARIS TRAPESIALIS var. **MORETONIANUS** Sowerby.

- * *Anodon moretonianus* SOWERBY, Conch. Icon., XVII, 1867, pl. IX, fig. 20.

† **GLABARIS TRAPESIALIS** var. **RIOPLATENSIS** Sowerby.

- * *Anodon rioplatensis* SOWERBY, Conch. Icon., XVII, 1870, pl. XXVI, fig. 101.
 * *Anodonta rioplatensis* CLESSIN, Conch. Cab. Ano., 1875, p. 217, pl. LXIV, fig. 3.—
 * PÆTEL, Conch. Sam., III, 1890, p. 184.
 * ? *Anodon ciconia* SOWERBY, Conch. Icon., XVII, 1870, pl. XXIX, fig. 115 a.

† **GLABARIS TRAPESIALIS** var. **CYGNÆFORMIS** Pilsbry.

- * *Glabaris trapesialis* var. *cygnaformis* PILSBRY, Pr. Ac. N. Sci. Phila., 1896, p. 563, pl. XXVI, fig. 4, 5.

Tropical South America from Brazil and Peru southward.

† **GLABARIS RADIATUS** Spix.¹

- * *Anodon radiatus* SPIX, Test. Fluv. Bras., 1827, p. 31, pl. XXIII, fig. 1.
 * *Margaron (Anodonta) radiatus* LEA, Syn., 1870, p. 83.
 * *Margarita (Anodonta) radiata* LEA, Syn., 1836, p. 51.
 * *Anodon radiata* CATLOW and REEVE, Conch. Nom., 1845, p. 67.
 * *Glabaris radiata* VON IHERING, Arch. für Nat., 1893, p. 115.

Brazil.

† **GLABARIS SIMPSONIANUS** Pilsbry.

- * *Glabaris simpsonianus* PILSBRY, Pr. Ac. N. Sci. Phila., 1896, p. 564, pl. XXVII, fig. 13.

Rio de la Plata.

GLABARIS HERTWIGII von Ihering.

- * *Anodonta hertwigii* VON IHERING, Arch. für Nat., 1890, p. 150, pl. IX, fig. 7.

† **GLABARIS SINUOSUS** Lamarck.

- * *Anodonta sinuosa* LAMARCK, An. sans Vert., VI, 1819, p. 87.—* DESHAYES, Enc. Meth., II, 1827, p. 147, pl. CCHII, fig. 2.—* HANLEY, Test. Moll., 1842, p. 224;
 * Biv. Shells, 1843, p. 224, Pt. 24, fig. 16.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* PÆTEL, Conch. Sam., III, 1890, p. 184.
 * *Margarita (Anodonta) sinuosa* LEA, Syn., 1836, p. 54; 1838, p. 32.
 * *Anodon sinuosa* CATLOW and REEVE, Conch. Nom., 1845, p. 68.
 * *Margaron (Anodonta) sinuosa* LEA, Syn., 1852, p. 53; 1870, p. 83.
 * *Anodon sinuosus* SWAINSON, Ex. Conch., 2d ed., 1841, p. 29, pl. XVI.

Brazil.

¹The specimen figured is a young shell, somewhat narrowed at the posterior end, and quite full at the central base, with broad, faint rays. It may be only a variety of *trapesialis*.

GLABARIS GRIJALVÆ Morelet.

* *Anodonta grijalvæ* MORELET, Jl. de Conch., XXXI, 1884, p. 12.—* PÆTEL, Conch. Sam., III, 1890, p. 180.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 532, pl. LXIX, figs. 1, 1a.

* *Anodonta grijalvæ* PÆTEL, Conch. Sam., III, 1890, p. 180.¹

Tabasco, Mexico.

† GLABARIS GLAUCUS Valenciennes.

* *Anodonta glauca* VALENCIENNES, Rec. Obs. Zool., II, 1833, p. 236, pl. I, fig. 2.—

* FERUSSAC, Gner. Mag., 1835, p. 25.—* DESHAYES, An. sans Vert., 2d ed., VI, 1835, p. 569; 3d ed., II, 1839, p. 679.—* DELESSERT, Rec. Coq. Lam., 1841, pl. XIII, fig. 3.—* HANLEY, Test. Moll., 1842, p. 221; * Biv. Shells, 1843, p. 221.—

* CONRAD, Pr. Ac. N. Sci. Phila., VI, 1853, p. 264.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. An., 1876, p. 222, pl. LXXIII, fig. 1.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 180.—* FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 533, pl. LXIX, figs. 1, 1 a.

* *Margarita (Anodonta) glauca* LEA, Syn., 1836, p. 51; 1838, p. 30.

* *Anodon glauca* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—* GOULD, Pr. Bost. Soc. N. Hist., III, 1850, p. 293.

* *Margaron (Anodonta) glauca* LEA, Syn., 1852, p. 50; 1870, p. 80.

* *Monocondylva glauca* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 501.—* PÆTEL, Conch. Sam., III, 1890, p. 174.

* *Anodon glaucus* SOWERBY, Conch. Icon., XVII, 1870, pl. XXVII, fig. 105.

* ? *Anodonta ciconia* GOULD, Pr. Bost. Soc. N. Hist. IV, 1851, p. 92.—CARPENTER, Maz. Shells, 1857, p. 117.—* ? FISCHER and CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 534, pl. LXVIII, figs. 2, 2a.

* *Anodonta burroughiana* CHENU (part), Ill. Conch., 1858, pl. III.

GLABARIS GLAUCUS var. SINALOENSIS Crosse and Fischer.

* *Anodonta glauca* VALENCIENNES, var. *sinaloensis* CROSSE and FISCHER, Jl. de Conch., XXXI, 1883, p. 219.

Mexico, south to Peru and Brazil.

GLABARIS UMBONATUS Simpson.

Anodon ciconia SOWERBY (part), Conch. Icon., XVII, 1870, pl. XXIX, fig. 115b.²

Locality unknown.

† GLABARIS BRIDGESII Lea.

* *Anodonta bridgesii* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 95; * Jl. Ac. N. Sci. Phila., VI, 1868, p. 291, pl. XLII, fig. 104; * Obs., XII, 1869, p. 51, pl. XLII,

¹ The above atrocity, which is a sample of many names in Pætel, is no doubt intended for *G. grijalvæ*.

² Sowerby gives two figures of what he calls *ciconia*, 115a being most likely a young shell of some form of *trapesialis*, the other a totally different form, apparently a new species of *Glabar* of the *trapesialis* group. As far as the figure shows it may be described as follows: Shell subquadrate with nearly straight dorsal and basal lines, biangulate behind, greatly inflated, with very full, high beaks which rise far above the dorsal line, winged and angulated at the front and hinder dorsal line; color olive-green. Length, 147 mm.; height, 92 mm.; elevation of beaks above dorsal line, 10 mm.

fig. 104.—* CLESSIN, Conch. Cab. Ano., 1874, p. 136, pl. XLV, figs. 1, 2.—* B. H.

WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 177.

* *Margaron (Anodonta) bridgesii* LEA, Syn., 1870, p. 81.

Nicaragua; Honduras.

† GLABARIS JEWITTIANUS Lea.

* *Anodonta jewittiana* LEA, Pr. Ac. N. Sci. Phila., XII, 1868, p. 95.

* *Anodonta jewittii* LEA, Jl. Ac. N. Sci. Phila., VI, 1869, p. 289, pl. XLI, fig. 101;

* Obs., XII, 1869, p. 49, pl. XLI, fig. 101.—* CLESSIN, Conch. Cab. Ano., 1874, p. 135, pl. XLIV, figs. 1, 2.—* B. H. WRIGHT, Check List, 1888.—* PÆTEL, Conch. Sam., III, 1890, p. 180.

* *Margaron (Anodonta) jewittii* LEA, Syn., 1870, p. 81.

Lake Nicaragua.

† GLABARIS FORBESIANUS Lea.

* *Anodonta forbesiana* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 92; * Jl. Ac. N. Sci. Phila., V, 1863, p. 393, pl. XLVII, fig. 301; * Obs., X, 1863, p. 29, pl. XLVII, fig. 301.—* CLESSIN, Conch. Cab. Ano., 1873, p. 115, pl. XXXIV, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

* *Anodon forbesianus* SOWERBY, Conch. Icon., XVII, 1870, pl. XXX, fig. 119.

* *Margaron (Anodon'a) forbesiana* LEA, Syn., 1870, p. 81.

Uruguay; Peru.

† GLABARIS MORICANDII Lea.

* *Anodonta moricandii* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 90; * Jl. Ac. N. Sci. Phila., V, 1863, p. 396, pl. XLIX, fig. 303; Obs., X, 1863, p. 32, pl. XLIX, fig. 303.—* CLESSIN, Conch. Cab. Ano., 1874, p. 114, pl. XXXVIII, figs. 3, 4.—

* PÆTEL, Conch. Sam., III, 1890, p. 182.

* *Anodon moricandii* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXII, fig. 126.

* *Margaron (Anodonta) moricandii* LEA, Syn., 1870, p. 81.

* *Glabaris moricandii* VON IHERING, Arch. für Nat., 1893, p. 115.

* ? *Anodonta exotica* SOWERBY, Conch. Icon., XVII, 1867, pl. XVI, fig. 57.

* *Anodonta angustata* CLESSIN,¹ Conch. Cab. Ano., 1876, p. 226, pl. LXXIV, figs. 6, 7.—

* PÆTEL, Conch. Sam., III, 1890, p. 176.

Brazil.

(Group of *Glabaris georginae*.)

Shell subsolid, obovate, with a decided, curved posterior ridge, and a smaller one above it, the space between the ridges a shallow groove.

GLABARIS GEORGINÆ Griffith.

* *Anodonta georginae* GRIFFITH, Griff. Cuvier, XII, 1834, pl. XIX. Brief description in index.

Rivers of Paraguay.

(Group of *Glabaris trigonus*.)

Shell long, elliptical, solid, inflated, produced at posterior base with smooth, shining epidermis; nacre very bright.

¹ Appears to be a very young *G. moricandii*.

† GLABARIS TRIGONUS Spix.

- * *Anodon trigonus* SPIX, Test. Fluv. Bras., 1827, p. 29, pl. xxii, fig. 2.
 * *Margarita (Anodonta) trigona* LEA, Syn., 1836, p. 51; 1838, p. 30.
 * *Anodon trigona* CATLOW and REEVE, Conch. Nom., 1845, p. 68.
 * *Anodonta trigona* HANLEY, Test. Moll., 1842, p. 218.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 618.—* KUSTER, Conch. Cab. Ano., 1853, p. 9, pl. ii, fig. 5.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* PÆTEL, Conch. Sam., III, 1890, p. 186.
 * *Margaron (Anodonta) trigona* LEA, Syn., 1852, p. 50; 1870, p. 75.
 * *Anodon chiquetana* D'ORBIGNY, Guer. Mag., 1835, p. 41.
 * *Anodonta chiquetana* PÆTEL, Conch. Sam., III, 1890, p. 177.
 * *Anodonta castelnandi* HUPE, Moll. Nouv., III, 1857, p. 88, pl. xviii, fig. 4.—* PÆTEL, Conch. Sam., III, 1890, p. 177.
 * *Anodonta subrostrata* PHILIPPI, Mal. Bl., XVI, 1869, p. 39.—* PFEIFFER, Nov. Conch., III, 1869, p. 486, pl. cv, figs. 1-3.—* PÆTEL, Conch. Sam., III, 1890, p. 185.
 * *Anodonta ucayalensis* PHILIPPI, Mal. Bl., XVI, 1869, p. 40.—* PFEIFFER, Nov. Conch., III, 1869, p. 486, pl. cv, figs. 4-6.—* PÆTEL, Conch. Sam., III, 1890, p. 186.

Brazil; Ecuador; Peru; Bolivia.

† GLABARIS AMAZONENSIS Lea.

- * *Anodonta amazensis* LEA, Pr. Ac. N. Sci. Phila., IV, 1860, p. 89; * Jl. Ac. N. Sci. Phila., V, 1863, p. 395, pl. xlvi, fig. 300; * Obs., X, 1863, p. 31, pl. xlvi, fig. 300.—* CLESSIN, Conch. Cab. Ano., 1874, p. 119, pl. xxxvii, fig. 7.—* PÆTEL, Conch. Sam., III, 1890, p. 176.
 * *Anodon amazonensis* SOWERBY, Conch. Icon., XVII, 1870, pl. xxx, fig. 120.
 * *Margaron (Anodonta) amazonensis* LEA, Syn., 1870, p. 82.

Amazon.

GLABARIS WEDDELLII Hupe.

- * *Anodonta weddellii* HUPE, Moll. Nouv., III, 1857, p. 87, pl. xvii, fig. 5.—* CLESSIN, Conch. Cab. Ano., 1876, p. 214, pl. lxvi, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 186.
 * *Anodon weddellii* SOWERBY, Conch. Icon., XVII, 1868, pl. xx, fig. 80.
 * *Margaron (Anodonta) weddellii* LEA, Syn., 1870, p. 79.

Brazil.

† GLABARIS ELONGATUS Swainson.

- * *Anodon elongatus* SWAINSON, Zool. Ill., 1st Ser., III, pl. clxxvi, 1823.
 * *Margarita (Anodonta) elongata* LEA, Syn., 1836, p. 53; 1838, p. 32.
 * *Anodonta elongata* HANLEY, Test. Moll., 1842, p. 223; * Biv. Shells, 1843, p. 223.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.
 * *Anodon elongata* CATLOW and REEVE, Conch. Nom., 1845, p. 66.
 * *Anodonta solidula* DEVILLE and HUPE, Rev. et Mag. Zool., 1850, p. 644, pl. xvi, fig. 2.—* HUPE, Moll. Nouv., III, 1857, p. 88, pl. xviii, fig. 2.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1876, p. 221, pl. lxxiii, fig. 2.—* PÆTEL, Conch. Sam., III, 1890, p. 185.
 * *Margaron (Anodonta) solidula* LEA, Syn., 1852, p. 53; 1870, p. 83.
 * *Anodon solidula* SOWERBY, Conch. Icon., XVII, 1869, pl. xxiii, fig. 91.
 † * *Anodonta wheatleyi* LEA, Tr. Am. Phil. Soc., X, 1852, p. 287, pl. xxvi, fig. 49; * Obs., V, 1852, p. 43, pl. xxvi, fig. 49.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1874, p. 113, pl. xxxvi, figs. 3, 4.—* PÆTEL, Conch. Sam., III, 1890, p. 186.
 * *Margaron (Anodonta) wheatleyi* LEA, Syn., 1852, p. 51; 1870, p. 82.

- * *Anodon amethystus* SOWERBY, Conch. Icon., XVII, 1869, pl. XXIV, figs. 95, 95a, 95b.¹
 * *Anodonta amethysta* CLESSIN, Conch. Cab. Ano., 1875, p. 180, pl. LX, fig. 3.
 * *Anodon dactylus* SOWERBY, Conch. Icon., XVII, 1867, pl. XIX, fig. 75.
 * *Anodonta dactylus* CLESSIN, Conch. Cab., 1875, p. 175, pl. LVII, fig. 3.—PÆTEL, Conch. Sam., III, 1890, p. 178.
Margaron (Anodonta) dactylus LEA, Syn., 1870, p. 80.

GLABARIS LINGULATUS Hupe.

- * *Anodonta lingulata* HUPE, Moll. Nouv., III, 1857, p. 89.—* CLESSIN, Conch. Cab. Ano., 1875, p. 215, pl. LXVI, fig. 3.—* PÆTEL, Conch. Sam., III, 1890, p. 181.
 * *Anodon lingulata* SOWERBY, Conch. Icon., XVII, 1869, pl. XXIII, fig. 90.
 * *Margaron (Anodonta) lingulata* LEA, Syn., 1870, p. 79.
 * *Glabaris lingulata* VON IHERING, Arch. für Nat., 1893, p. 119.

Paraguay.

† GLABARIS MORTONIANUS Lea.

- * *Anodonta mortoniana* LEA, Tr. Am. Phil. Soc., V, 1834, p. 80, pl. XIII, fig. 37; *Obs., I, 1834, p. 192, pl. XIII, fig. 37.—* HANLEY, Test. Moll., 1842, p. 219.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502.—* CLESSIN, Conch. Cab. Ano., 1874, p. 151, pl. XLVIII, figs. 5, 6.—* PÆTEL, Conch. Sam., III, 1890, p. 182.
 * *Margarita (Anodonta) mortoniana* LEA, Syn., 1838, p. 30.
 * *Anodon mortoniana* CATLOW and REEVE, Conch. Nom., 1845, p. 67.
 * *Margaron (Anodonta) mortoniana* LEA, Syn., 1852, p. 150; 1800, p. 80.

Parana River, South America.

GLABARIS LONGINUS Spix.

- * *Anodon longinus* SPIX, Test. Fluv. Bras, 1827, p. 29, pl. XXII, fig. 1.
 * *Mycetopus longinus* CLESSIN, Conch. Cab. Ano., 1875, p. 202.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 8.
 * *Iridina longina* LEA, Syn., 1836, p. 57.
 * *Anodonta longina* KUSTER, Conch. Cab. Ano., 1853, p. 7, pl. II, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 181.

Brazil; Amazon.

GLABARIS LEOTANDI Guppy.²

- * *Anodonta leotandi* GUPPY, Ann. and Mag., XIV, 1864, p. 243; * XVII, 1866, p. 54; Pr. Sci. Assn. Trinidad, 1872, p. —; * Jl. Conch., VII, 1894, p. 229.
 * *Anodonta leotandi* B. H. WRIGHT, Check List, 1888.
 * *Margaron (Anodonta) leotandi* LEA, Syn., 1870, p. 83.

Trinidad Island, West Indies.

Section STYGANODON von Martens, 1900.

(Type, *Anodonta tenebricosus* Lea.)

Shell subrhomboid, with a thick, dark, rather rough, sombre-colored epidermis, which is sometimes faintly rayed, naere lurid, shaded green; animal unknown.

(Group of *Glabaris tenebricosus*.)

Shell elongate, slightly inflated; base incurved.

¹ Changed to *wheatleyi* Lea, by Sowerby in index.² I am not certain where this groups, as I have never seen the shell, and it has not been figured.

† GLABARIS TENEBRICOSUS Lea.

- * *Anodonta tenebricosa* LEA, Tr. Am. Phil. Soc., V, 1834, p. 78, pl. XII, fig. 36;
 * Obs., I, 1834, p. 190, pl. XII, fig. 36.—* D'ORBIGNY, Guer. Mag., 1835, p. 39.—
 * HANLEY, Test. Moll., 1842, p. 224.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 616.—
 * Hanley, Biv. Shells, 1856, p. 224.—* H. and A. ADAMS, Gen. Rec. Moll., II,
 1857, p. 503.—* CHENU, Man., 1859, p. 146, fig. 720.—* VON MARTENS, Mal. Bl.,
 XV, 1868, p. 200.—* PÆTEL, Conch. Sam., III, 1890, p. 185.
 * *Margarita (Anodonta) tenebricosa* LEA, Syn., 1836, p. 54; 1838, p. 32.
 * *Anodon tenebricosa* CATLOW and REEVE, Conch. Nom., 1845, p. 68.—* SOWERBY,
 Conch. Icon., XVII, 1867, pl. XIII, fig. 43.
 * *Margaron (Anodonta) tenebricosa* LEA, Syn., 1852, p. 53; 1870, p. 83.
 * *Glabaris tenebricosa* VON IHERING, Arch. für Nat., 1893, p. 61.
 * *Anodon tenebricosus* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXI, fig. 123.

Brazil, Ecuador, Peru; south to Argentina.

GLABARIS PASTASANUS Clessin.

- * *Anodonta pastasana* CLESSIN, Mal. Bl. (2), I, 1879, p. 173, pl. XI, fig. 1.

Rio Pastasa, Ecuador.¹

† GLABARIS SOLENIFORMIS d'Orbigny.

- * *Anodonta soleniformis* D'ORBIGNY, Guer. Mag., 1835, p. 41; * Voy. Am. Mer., 1843,
 p. 617, pl. LXXIV, figs. 1, 3.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p.
 503.—* CLESSIN, Conch. Cab. Ano., 1873, p. 111, pl. XXVII, fig. 1.—* PÆTEL,
 Conch. Sam., III, 1890, p. 185.
 * *Margarita (Anodonta) soleniformis* LEA, Syn., 1838, p. 32.
 * *Margaron (Anodonta) soleniformis* LEA, Syn., 1852, p. 53; 1870, p. 83.
 * *Glabaris soleniformis* VON IHERING, Arch. für Nat., 1893, p. 59.
 * *Anodon solenidea* SOWERBY, Conch. Icon., XVII, 1867, pl. XVIII, fig. 65.
 * *Margaron (Anodonta) solenidea* LEA, Syn., 1870, p. 83.

Brazil, south into Argentina.

† GLABARIS CLESSINI Fischer.

- * *Mycetopus plicatus* CLESSIN, Mal. Bl., V, 1882, p. 190, pl. IV, fig. 7.
 * PÆTEL, Conch. Sam., III, 1890, p. 187.
 * *Mycetopus clessini*, FISCHER,² Jl. de Conch., XXXVIII, 1890, p. 8, footnote.
 * *Glabaris nehringi* VON IHERING, Arch. für Nat., 1893, p. 60.

Southern Brazil; southward into Argentina.

GLABARIS BAMBOUSEARUM Morelet.

- * *Anodon bambousearum* MORELET, Test. Nov., II, 1851, p. 24.—* FISCHER and
 CROSSE, Miss. Sci., Pt. 7, II, 1894, p. 527, pl. LXIII, figs. 6, 6a.

Palenque in Chiapas, Mexico.

¹ Probably only a variety of *tenebricosus*.

² Fischer changed the name *plicatus* to *clessini* because the former name had been used by Sowerby for a species which he (Sowerby) placed in *Mycetopus*. Sowerby's shell is a *Mutela*, and that of Clessin a *Glabaris* of the *Tenebricosus* group. Von Ihering applied the name *nehringi* to the above after Fischer had changed it, and was evidently not aware that the French savant had given it a new name.

† GLABARIS SCHRÖTERIANUS Lea.

* *Anodonta schröteriana* LEA, Tr. Am. Phil. Soc., XI, 1852, p. 292, pl. XXIX, fig. 55;

* Obs., V, 1852, p. 48, pl. XXIX, fig. 55.—* HUPE, Moll. Nouv., III, 1857, p. 89, pl. XVIII, fig. 3.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* CLESSIN, Conch. Cab. Ano., 1874, p. 151, pl. XLVII, figs. 5, 6.—* PÆTEL, Conch. Sam., III, 1890, p. 184.

* *Margaron (Anodonta) schröteriana* LEA, Syn., 1852, p. 51; 1870, p. 82.

* *Anodon schröteriana* SOWERBY, Conch. Icon., XVII, 1868, pl. xx, fig. 74.

* *Margaritana schröteriana* PÆTEL, Conch. Sam., III, 1890, p. 173.

Amazon drainage.

(Group of *Glabaris obtusus*.)

Shell short, inflated, thin, feebly rayed, rays often broken; base rather full.

† GLABARIS OBTUSUS Spix.

* *Anodon obtusus* SPIX, Test. Fluv. Bras., 1827, p. 30, pl. XXII, fig. 3.

* *Margarita (Anodonta) obtusa* LEA, Syn., 1836, p. 52; 1838, p. 31.

* *Anodonta obtusa* HANLEY, Test. Moll., 1842, p. 221; Biv. Shells, 1843, p. 221.—* POTIEZ and MICHAUD, Gall. Moll., 1844, p. 144, pl. LV, fig. 3.—* KUSTER, Conch. Cab. Ano., 1853, p. 8, pl. II, figs. 3, 4.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 504.—* MUSGRAVE, Phot. Conch., 1863, pl. I, fig. 7.—* CLESSIN, Conch. Cab. Ano., 1875, p. 170, pl. LVI, figs. 1, 2.—* PÆTEL, Conch. Sam., III, 1890, p. 182.

* *Margaron (Anodonta) obtusa* LEA, Syn., 1852, p. 51; 1870, p. 82.

* *Anodon obtusa* CATLOW and REEVE, Conch. Nom., 1845, p. 67.—* SOWERBY, Conch. Icon., XVII, 1867, pl. XII, fig. 39.

* *Anodonta VALENCIENNES*, Coq. Mar. Biv., 1827?, pl. XLVIII, figs. 3a, 3b.¹

Brazil; Paraguay.

† GLABARIS LITURATUS Spix.

* *Anodon obtusus* var. *b. lituratum* SPIX, Test. Fluv. Bras., 1827, p. 30, pl. XXII, fig. 4.

* *Anodonta litturata* HUPE, Moll. Nouv., 1857, p. 87, pl. XVII, fig. 4.²

* *Anodon litturata* SOWERBY, Conch. Icon., XVII, 1868, pl. xx, fig. 78.

Brazil.

Section VIRGULA Simpson, 1900.

(Type, *Glabaris ensiformis* Spix.)

Shell subsolid to solid, moderately inflated, greatly elongated, straight or falcate, rounded in front, sharply pointed at the posterior base, where the high, sharply defined posterior ridge ends and above which it is somewhat obliquely truncated; beaks not high; epidermis green to olive; nacre brilliant, bluish or purplish, iridescent, rayed with very fine, indistinct ridges; posterior end with a slight sinus.

Animal unknown.

¹ Valenciennes gives no name or description with the above, but the figure is, no doubt, *G. obtusa*.

² Hupe is certain that this is different from *obtusum*, and says that the small angular marks on its surface are very distinct; that it is thinner and less inflated. I doubt whether it is more than a variety of *obtusum*.

† GLABARIS ENSIFORMIS Spix.

* *Anodon ensiformis* SPIX, Test. Fluv. Bras., 1827, p. 31, pl. XXIV, figs. 1, 2.—

* CATLOW and REEVE, Conch. Nom., 1845, p. 66.—* SOWERBY, Conch. Icon., XVII, 1867, pl. XI, fig. 31.

* *Iridina ensiformis* LEA, Syn., 1836, p. 57.

* *Anodonta ensiformis* HANLEY, Test. Moll., 1842, p. 223; * Biv. Shells, 1843, p. 223.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 618, pl. LXXIX, fig. 10.—* KUSTER, Conch. Cab. Ano., 1853, p. 8, pl. II, fig. 12.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 503.—* ? CHENU, Man., 1859, II, p. 146, fig. 721.—* PÆTEL, Conch. Sam., III, 1890, p. 179.

* *Margarita (Anodonta) ensiformis* LEA, Syn., 1838, p. 32.

* *Margaron (Anodonta) ensiformis* LEA, Syn., 1852, p. 51; 1870, p. 82.

Tropical South America.¹† GLABARIS FALSUS Simpson.²

Yuruari River, a branch of the Orinoco.

† GLABARIS LEGUMEN von Martens.

* *Anodonta legumen* VON MARTENS, S. B. Nat. Fr., 1888, p. 65.

Southern Brazil.

The following are unfigured or indeterminate species of *Glabaris*.

* *Anodon brevis* SOWERBY, Conch. Icon., XVII, 1870, pl. XXXI, fig. 124.

La Plata, South America.³

* *Anodon porcifer* GRAY, Pr. Zool. Soc. Lond., 1834, p. 58.

Paraguay.

* *Unio (Anodontes) spixii* D'ORBIGNY, Guer. Mag., 1835, p. 39.⁴

Brazil.

* *Anodon cheeziana* SOWERBY, Conch. Icon., XVII, 1867, pl. xv, fig. 52.

Probably a *Glabaris*.

* *Anodonta nicaragua* PHILIPPI, Zeits. für Mal., V, 1848, p. 130.

Nicaragua.

¹ The three very peculiar species of this section seem to show relationship with *G. wheatleyi* and *G. clessini*.

² Shell elongated, rather thin, subrhomboid and pointed at the posterior base, slightly rayed on the low, rounded posterior ridge; beaks low; epidermis olive green, somewhat cloth-like; hinge line nearly straight; anterior scars deep; escutcheon small, dark; nacre shaded bluish and greenish, lurid purplish in the center, iridescent behind. Length, 76 mm.; height, 21 mm.; diameter, 11 mm. This was supposed by Dr. Lea to be *ensiformis* and I thought it was that until a more careful examination convinced me of its distinctness. It is a smaller, thinner, more rhomboid shell, and has a somewhat cloth-like epidermis, while that of *ensiformis* is smooth. I regret that I can not figure this species, and that I must describe it in a footnote, but I only discovered that it was a distinct species as the Synopsis was about ready for the press.

³ I can not tell what this is. If it is a true *Glabaris*, it would be the type of a group perhaps near *G. tenebricosus*.

⁴ D'Orbigny refers to figures in Spix, but his references are badly mixed up.

**Anodonta aperta* RAFINESQUE, Atl. Jl., No. 4, 1832, p. 134.

Parana River, South America.

**Anodonta atrovireus* PHILIPPI, Zeits. für Mal., V, 1848, p. 130.

**Anodonta carinata* DUNKER, Mal. Bl., V, 1858, p. 225.

Colombia.

**Anodonta cornea* PHILIPPI, Zeits. für Mal., V, 1848, p. 130.

Nicaragua.

**Anodonta giullaini* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 502. Credited to Recluz.

**Anodonta paphos* RAFINESQUE, Atl. Jl. and Friend., 1832, p. 134.

Parana River.

**Anodon pictus* SWAINSON, Ex. Conch., 2d ed., 1841, p. 39.

**Anodonta wallisi* MOUSSON, Mal. Bl., XVI, 1869, p. 188.

**Anodonta tehuan-tepecensis* FISCHER and CROSSE, Miss. Sci., 1894, p. 526.

Tehuantepec. Not yet figured.

**Anodonta hertwigi* VON IHERING.

Where?

**Anodonta bergi* VON IHERING.

Where?

Genus MYCETOPODA d'Orbigny, 1835.¹

(Type, *Mycetopoda soleniformis* d'Orbigny.)

Mycetopoda D'ORBIGNY, Guer. Mag., No. 62, 1835, p. 41.

Mycetopus D'ORBIGNY, Voy. Am. Mer., 1847, p. 600.

Shell thin, elongated, truncate above behind, with a low, posterior ridge and rather flat, smooth or slightly concentrically wrinkled beaks; epidermis smooth, shining, pale greenish-yellow or brownish, rayless; hinge line long, straight, edentulous or showing faint traces of denticles, under a glass, beneath the nacre; nacre soft, bluish-white and iridescent; muscular impressions faint, irregular, the smaller anterior scar above the larger one; beak cavities shallow.

Animal having very long gills, the inner much the larger, united to the abdominal sac throughout their whole length; palpi large, round below, projecting very slightly behind and attached along the whole length of the straight upper border; mantle very thin, slightly thickened at the edges; branchial opening closed below into a short papillose siphon, and separated from the nearly smooth anal opening by a strong bridge; superanal opening not closed below; foot very long, developed at the lower end into a sort of head or button.

¹ So named by its author in the Guerin Magazine, but afterwards changed by him to *Mycetopus* in the Voyage Amerique Meridionale. The genus has been made the type of a separate family by Gill, and was so acknowledged by Pelseneer and others, but it does not seem to me to be separable from the *Mutelidae*. Certain shells under favorable light show slight dentilations along the hinge line, which are, no doubt, vestigial taxodont teeth common to the family; the labial palpi and anal bridge are decidedly mutelid in character, and the great development of the foot is paralleled in *Solenata*, *Lastena*, and to some extent by *Gonidea* among the *Unionidae*.

(Group of *Mycetopoda siliquosa*.)

Beaks in front of the center of the shell; anterior end evenly rounded; basal lines nearly straight; posterior ridge quite low. Animal with the characters of the genus.

† MYCETOPODA SILIQUOSA Spix.

- * *Anodon siliquosus* SPIX, Test. Fluv. Bras., 1827, p. 30, pl. XXIII, fig. 2.
 * *Mycetopoda siliquosus* D'ORBIGNY, Gner. Mag., 1835, p. 41.
 * *Mycetopus siliquosus* D'ORBIGNY, Voy. Am. Mer., 1843, p. 601, pl. LXVII.—* HANLEY, Test. Moll., 1842, p. 224; * Biv. Shells, 1843, p. 224.—* CATLOW and REEVE, Conch. Nom., 1845, p. 69.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 505; III, pl. CXVIII, fig. 1.—* SOWERBY, Conch. Icon., XVI, 1868, pl. I, fig. 2; III, fig. 2a.—* CLESSIN, Conch. Cab. Ano., 1875, p. 200, pl. LXVIII, figs. 2-4; LXIX, fig. 1.—* PÆTEL, Conch. Sam., III, 1890, p. 187.—* VON IHERING, Arch. für Nat., 1893, p. 56.
 * *Platiris (Mycetopus) siliquosus* LEA, Syn., 1852, p. 56; 1870, p. 90.
 * *Platiris (Iridina) siliquosa* LEA, Syn., 1836, p. 56; 1838, p. 34.
 * *Anodonta siliquosa* KUSTER, Conch. Cab. Ano., 1853, p. 35, pl. VIII, fig. 3.
 * *Mycetopoda siliquosa* CHENU, Ill. Conch., 1858, pl. I, figs. 2, 2a, 2b, 2c.

Bolivia, Brazil; south into Argentina.

* MYCETOPODA OCCIDENTALIS Clessin.¹

- Mycetopoda Occidentalis* CLESSIN, Mal. Bl. I, 1879, p. 174, pl. XI, figs. 2, 3.—
 * PÆTEL, Conch. Sam., III, 1890, p. 187.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 8.

Ecuador.

MYCETOPODA STAUDINGERI von Ihering.²

- * *Mycetopus staudingeri* VON IHERING, Arch. für Naturg., 1890, p. 131, figs. A, B.

Upper Amazon in Ecuador and Peru.

MYCETOPODA SUBSINUATA Sowerby.

- * *Mycetopus subsinuatus* SOWERBY, Conch. Icon., XVI, 1868, pl. IV, fig. 10.—* CLESSIN, Conch. Cab. Ano., 1875, p. 205, pl. LXX, fig. 3.—* PÆTEL, Conch. Sam., III, 1890, p. 187.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 8.

Bogota; Colombia; Equador.

† MYCETOPODA PYGMÆA Spix.

- * *Anodon siliquosus* var. *b pygmaeum* SPIX, Test. Fluv. Bras., 1827, p. 30, pl. XXIII, figs. 3, 4.
 * *Mycetopus pygmaeus* SOWERBY, Conch. Icon., XVI, 1868, pl. II, fig. 4.—* CLESSIN, Conch. Cab. Ano., 1875, p. 205, pl. LXX, fig. 3.—* PÆTEL, Conch. Sam., 1890, p. 197.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 8.
 * *Platiris (Mycetopus) pygmaeus* LEA, Syn., 1870, p. 90.
 † *Mycetopus weddellii* HUPE, Moll. Nouv., III, 1857, pl. XX, fig. 2.—* SOWERBY, Conch. Icon., XVI, 1868, pl. II, fig. 5.—* CLESSIN, Conch. Cab. Ano., 1875, p.

¹ Probably a wide variety of *M. siliquosus*.

² I have never seen this species, and the author only gives two rough interior outline figures, but its form is so much like that of *siliquosus* that I somewhat doubt its specific value.

203, pl. LXVI, fig. 6.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 8.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

Brazil; northward to Nicaragua.

MYCETOPODA HUPEANA Clessin.

* *Mycetopus pygmaeus* HUPE, Moll. Nouv., III, 1857, p. 93, pl. XIX, fig. 2.

* *Mycetopus hupeanus* CLESSIN, Conch. Cab. Ano., 1875, p. 206, pl. LXVI, fig. 15.—

* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 8.

Brazil.

(Group of *Mycetopoda ventricosa*.)

Shell rather short and high, inequilateral subrhomboid with a strong angle at the anterior upper point, cut away below in front, slightly sinuous on the base, narrower behind, strongly truncate on posterior slope; posterior ridge wide and lightly curved, hinge line curved.

Animal unknown.

† MYCETOPODA VENTRICOSA d'Orbigny.

* *Mycetopoda ventricosa* D'ORBIGNY, Voy. Am. Mer., 1843, p. 602, pl. LXXII, figs. 1-3.—

* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 505.—* SOWERBY, Conch. Icon., XVI, 1868, pl. III, fig. 8.—* CLESSIN, Conch. Cab. Ano., 1875, p. 202, pl. LXIX, figs. 2, 3.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 8.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

* *Platiris (Mycetopus) ventricosus* LEA, Syn., 1852, p. 56; 1870, p. 90.

Bolivia.

(Group of *Mycetopoda soleniformis*.)

Shell large, rounded in front, but somewhat truncated on the lower anterior part; base line evenly incurved; posterior ridge well developed, curved, truncate on posterior slope; beaks central.

Animal that of the genus.

† MYCETOPODA SOLENIFORMIS d'Orbigny.

* *Mycetopoda soleniformis* D'ORBIGNY, Guer. Mag., 1835, p. 41.—* CHENU, Ill. Conch., 1858, pl. I, figs. 1, 1a, 1b.

* *Platiris (Mycetopus) soleniformis* LEA, Syn., 1838, p. 34; 1852, p. 56; 1870, p. 90.

* *Mycetopus soleniformis* REEVE, Conch. Syst., I, 1841, p. 125, pl. XCIV.—* HANLEY, Test. Moll., 1842, p. 224; * Biv. Shells, 1843, p. 224.—* D'ORBIGNY, Voy. Am. Mer., 1843, p. 601, pl. LXVI.—* CATLOW and REEVE, Conch. Nom., 1845, p. 69.—* H. and A. ADAMS, Gen. Rec. Moll., II, 1857, p. 505; III pl. CXVIII, figs. 2a, 2b.—* CHENU, Man., 1859, II, p. 147, fig. 725.—* REEVE, Elements of Conch., II, 1860, pl. XXXII, fig. 181a, b.—* SOWERBY, Conch. Icon., XVI, 1868, pl. I, fig. 1.—* CLESSIN, Conch. Cab. Ano., 1875, p. 201, pl. LXVIII, fig. 1.—* FISCHER, Jl. de Conch., XXXVIII, 1890, p. 7.—* PÆTEL, Conch. Sam., III, 1890, p. 187.

* *Mycetopus solenoides* SOWERBY, Conch. Man., 1839, fig. 151.

* *Margaron (Anodonta) soleniformis* LEA, Syn., 1852, p. 53.

* *Platiris (Mycetopus) soleniformis* LEA, Syn., 1870, p. 90.

Bolivia; Peru.

¹ 1843 was the date of issue of the last part of the voyage. The description of *Mycetopus* in this work appeared earlier.

BIBLIOGRAPHY.

It is not claimed that the following bibliography is complete or in all respects accurate. Such titles as have been seen by the writer are prefixed with an *. On account of its great length it has been thought best to make considerable abbreviations.

- ADAMI, G. B. Molluschi terrestre e fluviatili viventi nella valle dell' Oglio ossia nelle valli Camonica di Scalve di Borlezza, province di Brescia e Bergamo.
Atti Soc. Pad., V, 1876, Pt. 1.
- *— Nuove Forme Italiane del Genere Unio.
Bull. Soc. Mal. It., VIII, 1882, pp. 129-138. 1 plate.
- *— Novità Malacologiche Recenti.
Bull. Soc. Mal. It., XI, 1886, pp. 202-261. 4 plates.
- *ADAMS, C. B. Shells of Fresh Pond.
Am. Jour. Sci. and Arts, XXXVI, 1839, p. 392.
- *— Catalogue of the Mollusca of Middlebury, Vermont, and vicinity; with observations.
Am. Jour. Sci. and Arts, XL, 1841, pp. 266, 267.
- *— Fresh-water and Land Shells of Vermont.
In *Thompson's History of Vermont*, Pt. 1, 1842.
Published separately under the above title without date.
- Catalogue of Genera and Species of Recent Shells. Collection of C. B. Adams, 1847.
In *Marcy's Exploration of the Red River of Louisiana*, 1852, pp. 242-244.
- *ADAMS, ARTHUR, and ANGAS, GEORGE FRENCH. Descriptions of New Species of Shells chiefly from Australia, in the collection of Mr. Angas.
Proc. Zool. Soc., 1864, pp. 35-40.
- *ADAMS, HENRY. List of the Shells collected by Samnel White Baker during his recent explorations in Central Africa.
Proc. Zool. Soc., 1866, pp. 375, 376.
- *ADAMS, HENRY. List of Land and Fresh-water Shells collected by Mr. E. Bartlett on the Upper Amazons, and on the River Ucayali, eastern Peru, with descriptions of New Species.
Proc. Zool. Soc., 1866, pp. 440-445.
- *— Descriptions of six new species of shells, etc.
Proc. Zool. Soc., 1866, pp. 445-447.
- *— Descriptions of fifteen new species of Land and Fresh-water Shells from Formosa, collected by Robert Swinhoe, etc.
Proc. Zool. Soc., 1866, pp. 316-319, pl. xxxiii.
- *— List of additional species of Land and Fresh-water Shells collected by Mr. E. Bartlett in eastern Peru, with descriptions of new species.
Proc. Zool. Soc., 1870, pp. 374-376.
- *ADAMS, HENRY and ARTHUR. The Genera of Recent Mollusca, arranged according to their organization. 3 vols., 1853-1858.
- *ADAMS, LIONEL ERNEST. The Collector's Manual of British Land and Fresh-water Shells, etc., 1884, pp. 125, 8 plates.
- *— The Mollusca of Northamptonshire.
Jour. Northamp. Soc., VIII, 1895, pp. 265-267.
- *ADANSON, M. Histoire Naturelle du Senegal. Coquillages. Avec la Relation abrégée d' un Voyage fait en ce pays, pendant les années, 1749, 1750, 1751, 1752, et 1753. Paris, 1757, pp. 275, 19 plates.
- AGASSIZ, LOUIS. Mémoire sur les Moules de Mollusques vivants et fossiles.
Mém. Soc. des Sci. Nat. Neuchatel, 1839.
- *— Nomenclator Zoologicus, etc. Soloduri, 1842-1846.

- AGASSIZ, LOUIS. Introduction to the Study of Natural History. New York, 1847.
- Twelve Lectures on Comparative Embryology: Boston, 1849, pp. 22.
- * — Remarks on Fresh-water Shells. *Proc. Bost. Soc. Nat. Hist.*, III, 1850, pp. 301, 356, 357.
- * — Diagnoses of genera of Unionidæ. In *Stimpson's Shells of New England*, 1851.
- * — Letter to Isaac Lea on the Naiades. *Proc. Am. Phil. Soc.*, V, 1852, pp. 219-220.
- * — Ueber die Gattungen unter den Nord Amerikanischen Najaden. *Arch. für Naturg.*, XVIII, 1852, pp. 41-50.
- * — Notes on Unio ligamentinus reversed. *Proc. Bost. Soc. Nat. Hist.*, VII, 1859, pp. 166-167.
- * — Some observations on the rate of increase and other characters of fresh-water shells—Unios. *Proc. Bost. Soc. Nat. Hist.*, VIII, 1861, pp. 100-102.
- * — Methods of Study in Natural History, 1869, pp. 319.
- ALDER, JOSHUA. A Catalogue of the Land and Fresh-water Testaceous Mollusca found in the vicinity of Newcastle-upon-Tyne, etc. *Trans. Nat. Hist. Soc. Northumberland and Durham*, I, 1830, pp. 26-41; II, 1833, pp. 337-342.
- Notes on the Land and Fresh-water Mollusca of Great Britain, with a revised list of species. *Jardine's Mag. of Zool. and Bot.*, II, 1838, pp. 101-119.
- Catalogue of Mollusca of Northumberland and Durham. *Trans. Tyne. Field Nat. Club*, I, 1846-1850, pp. 97-209, 358-365.
- * ALDRICH, TRUMAN H. Partial List of Shells found near Troy, New York. *Rep. of Regents of Univ. of N. Y.*, XXII, 1869, pp. 17-24.
- ALDROVANDUS, ULYSSES. De Reliquis animalibus Exsanguibus utpote de Mollibus Crustaceis Testaceis et Zöophytis, Part IV, Bononiæ, 1606.
- ALTEN, JOHANNES WILHELM. Systematische abhandlungen über die Erd und Fluss-Conchylien, etc., 14 colored plates, Augsburg, 1812.
- AMALITZKI, V. P. К во просу о brevности seruĕst va Unionidæ. [On the probable antiquity of the Unionidæ.] *C. R. Sect. Biol. Soc. Varsovie*, II, 1891, pp. 1-5.
- * — Ueber die Anthracosien der Permformation Russlands. *Palæontogr.*, XXXIX, 1892, pp. 125-215, 5 plates. Separate in Russian.
- A comparison of the Permian Fresh-water Lamellibranchiata from Russia, with those of the Karoo System of South Africa. *Quart. Jour. Geol. Soc.*, LI, 1895, pp. 337-349, 2 plates.
- ANCEY, C. F. Unio gladiator, n. sp. *Le Naturaliste*, III, 1881, p. 468.
- * — On the Generic Name of a Remarkable Bivalve Shell, found in the Congo. *Conch. Exchange*, II, 1888, p. 22.
- * — Résultat des recherches malacologiques de Mgr. Lechaptois sur les bords du lac Nyassa et de la rivière Shiré. *Mém. Soc. Zool. de France*, VII, pp. 217-234, fig. 8.
- ANONYMOUS. A Perfect Description of Virginia, 1649.
- Der Naturforch.*, XVIII, 1782, p. 70.
- * — A Catalogue of the Portland Museum, 17-6.
- * — Sur l'opinion de Treviranus. *Isis*, XX, 1827, p. 752.
- * — Remarks on Dr. Isaac Lea's Memoir in the Transactions of the American Philosophical Society for 1827. *Am. Jour. Sci. and Arts*, XVI, 1829, p. 378.
- Remarks on Mr. Isaac Lea's Monograph of the genus Unio. *Jour. Nat. and Geog. Sci.*, II, 1830, p. 185.
- Criticism. *Am. Jour. Geol. and Nat. Sci.*, I, 1831, p. 370.
- * — Mr. Lea on the Naiades, in the Transactions of the American Philosophical Society. *Am. Jour. Sci. and Arts*, XXII, 1832, pp. 169-178.
- * — Observations on the genus Unio, etc., by I. Lea, in the Transactions of the American Philosophical Society. *Am. Jour. Sci. and Arts*, XXVII, 1835, pp. 371-377.

- *ANONYMOUS. On the Economic Uses of some species of Testacea.
Am. Jour. Sci. and Arts, XXXII, 1837, pp. 235-259.
- *— Observations, etc., from the Transactions of the American Philosophical Society.
Am. Jour. Sci. and Arts, XXXV, 1839, pp. 184-186.
- *— Preparation of Shells for the Cabinet.
Am. Jour. Sci. and Arts, XLII, 1842, p. 391.
- *— Notice of Lea's Observations on the genus *Unio*, etc., VIII.
Am. Jour. Sci. and Arts, XXXIV, 1862, p. 451.
- *— Notice of Lea's Observations on *Unio*, etc., XI.
Am. Jour. Sci. and Arts, XLIII, 1867, p. 411.
- *ANTHONY, JOHN G. On the Byssus of *Unio*, with notes by J. E. Gray, esq.
Ann. Nat. Hist., VI, 1841, p. 77.
- *— List of Land and Fresh-water Shells near Cincinnati, Ohio, 1842, 2d ed., 1843.
- *— Descriptions of New Species of North American Unionidæ.
Am. Jour. Conch., I, 1865, pp. 155-164, pls. XII-XVI.
- *— Descriptions of Two New Species of Monocondylæa.
Am. Jour. Conch., I, 1865, pp. 205-206, pl. xviii.
- *— Descriptions of New Species of Shells.
Am. Jour. Conch., I, 1865, p. 351, pl. xxv, fig. 2.
- *— Descriptions of New American Fresh-water Shells.
Am. Jour. Conch., II, 1866, pp. 144-147, pls. VI, VII.
- *ANTON, HERMANN EDUARD. Verzeichniss der Conchylien, welche sich in der Sammlung von H. E. Anton befinden.
Halle, 1839, pp. 110.
- APATHY, I. Tanúlmaúy a Najadeak Szó vettanaról.
Ert. Term. Kör., XIV, 1885, 121 pp., 4 plates. Abstract in *Biol. Centralbl.*, VII, 1885, pp. 621-630. Abstract in *Jour. Roy. Micr. Soc.*, 1888, pp. 205-206.
- APATHY, I. Studien zur Histologie der Najadeen auf Grund selbstständiger Untersuchungen.
Math. Nat. Ber. Ung., III, 1886, pp. 109-110.
- APGAR, A. C. The Muskrat and the Unio.
Jour. Trenton Soc., I, 1887, pp. 58-59. *Zoologist* (3.), XI, 1887, p. 425.
- ARADAS, ANDR., and MAGGIORE, GIAC. Catalogo ragionato delle conchiglie viventi Sicilia.
Atti Accad. Gioènia, XV, 1839, pp. 53-106, 163-205; XX, 1843, pp. 101-142, 343-360.
- *ARANGO, R. Catalogo de los moluscos terrestres y fluviatiles de la isla de Cuba.
Poey, Report Físico-Nat. de la Isla de Cuba, 1865, pp. 81-112, 123-144, 145-149.
- *ARGENVILLE, ANTONIO JOSEPH DESALLIER D'. La Conchyliologie, ou Histoire Naturelle des Coquilles de mer d'eau douce, terrestres et fossiles; avec un traité de la zoomorphose, 1742. 2 vols., 33 plates. Three editions, the third by Favanne.
- *— Histoire naturelle éclaircie dans une des parties principales, l'Orctologie qui traite des terres, des pierres, des métaux, des minéraux et autres fossiles, etc., Paris, 1753.
- AUBRIOT. Catalogue des Mollusques vivants du département de la Marne, dressé sur les notes de M. Arnould.
Séanc. et Trar. Acad. Rennes, 1846.
- *AUCAPITAINE, HENRI. Mollusques terrestres et de l'eau douce observés dans la Haute Kabylie (versant nord du Djurjura).
Rev. et Mag., XIV, 1862, pp. 144-162.
- AUDOUIN, JEAN VICTOR. Observations pour servir à l'histoire de la formation des Perles.
Mém. Mus., XVII, 1820, p. 174, separate with plate.
- *— Explication des Planches du grand Ouvrage d'Égypte, 1827.
- *AUGHEY, SAMUEL. Catalogue of the Land and Freshwater Shells of Nebraska.
Bull. U. S. Geol. and Geog. Survey of the Territories, III, 1877, pp. 697-704.

- BAER, KARL ERNST VON. Observations sur la génération des monles et sur un système des vaisseaux hydrofères dans ces animaux.
Notiz. aus dem Geb. Nat. und Heilk., 1826, No. 265, p. 1.
- Extrait de Sept Mémoires sur les Entozoaires ou Vers intestinaux des Mollusques.
Bull. Sci. Nat., IX, 1826, No. 103.
- In *Froriep's Notizen*, January, 1828.
- Ueber den Weg, den die Eier unserer Süßwassermuscheln nehmen, um in die Kiemen zu gelangen nebst allgemeinen Bemerkungen über den Bau der Muscheln.
Meckel's Archiv, 1830, pp. 313-352, pl. VII.
- *BAICHÈRE, ED. *Nayades de l'Aube*.
Bull. Soc. Mal. France, VII, 1890, pp. 117-132.
- BAILLIE, ÉMILE. Liste des Unionides récoltées dans la mare de Bouillon, etc.
Bull. Soc. Amis Nat. Rouen, 1898, p. 11.
- *BAILLIE, WILLIAM. Sutherland and Caithness Field Notes.
Jour. of Conch., III, 1882, pp. 297-299.
- *BAIRD, W., and ADAMS, H. Notes upon some interesting Chinese Shells, with a description of two or three New Species of Unionidae, collected at Shanghai, etc.
Proc. Zool. Soc., 1867, pp. 489-492, pl. XXVI.
- *BAKER, FRANK COLLINS. The Molluscan Fauna of Western New York.
Trans. Acad. Sci. St. Louis, VIII, 1898, pp. 71-94, pl. x. 1 figure.
- * — The Mollusca of the Chicago Area. The Pelecypoda.
Nat. Hist. Survey of Chicago Acad. of Sci., Bull. No. 3, Pt. 1. September 1, 1898, pp. 130, 27 plates.
- BALEFOUR. Traité d'embryologie et d'organographie comparatif, 1883.
- BALL, ROBERT. On the occurrence of Fresh-water Shells in some of our Post-Tertiary Deposits in Canada.
Can. Nat. and Geol., VI, 1861, pp. 42, 51.
- *BARBIÉ, AUGUSTE. Catalogue méthodique des Mollusques terrestres et fluviatiles du département de la Côte d'Or, etc.
Mém. Acad. Dijon, 1852, pp. 163-194. 1 plate. Separate, 1854.
- *BARBUT, JAMES. The Genera Vermium, exemplified by figures of the animals contained in the orders of the Intestina and Mollusca Linnæi, drawn from nature, 1783 and 1788.
 Two parts. 25 colored plates.
- BARKER, JOHN. Notes on some Dissections of the Fresh-water Pearl Mussel (Unio) Margaritana margaritifera.
Dublin Quart. Jour. Sci., V, 1865, pp. 62-64. *Proc. Nat. Hist. Soc. Dub.*, IV (1862-65), 1865, pp. 111-114.
- *BARNES, D. W. On the Genera Unio and Alasmiodonta, with Introductory Remarks.
Am. Jour. Sci. and Arts, VI, 1823, pp. 107-127; pp. 258-280, with figures. Extract in *Fer. Bull. Sci. Nat.*, VI, 1825, pp. 122-124.
- * — Reclamation of Unios.
Am. Jour. Sci. and Arts, XIII, 1828, pp. 358-364.
- *BARTRAM, JOHN. Observations concerning the Salt-water Muscle, the Oyster Banks, and the Fresh-water Muscle of Pennsylvania.
Phil. Trans., XLIII, 1744, pp. 157-159.
- *BARTRAM, WILLIAM. Travels through North and South Carolina, Georgia, east and west Florida, etc., 1791.
- BAUDON, AUGUSTE. Catalogue des Mollusques du département de l'Oise.
Mém. Soc. Oise, 1852. Separate, 1853, pp. 20.
- * — Notice sur la ponte de quelques Unio; comparaison entre l'oviducte de ce genre et celui des Anodontes.
Jour. de Conch., IV, 1853, pp. 353-357.
- * — Observation sur la valeur du Pseudidm.
Jour. de Conch., IV, 1853, pp. 277-279.
- BAUMGARTEN. Notizen über die Perlenfischerei u. Perlenzucht im Perlenbache bei Falkenstein.
Korrespbl. d. Zool. Mineral. Ver. Regensburg., 1856, pp. 165-175.

- *BEAN, WILLIAM. Fossil Shells.
Mag. Nat. Hist., IX, 1836, pp. 376-377, 3 figs.
- *BEAUCHAMP, WILLIAM. Distribution of *Unio pressus* (Lea).
Am. Naturalist, XVII, 1883, pp. 433, 434.
- *— Erosion of Fresh-water Shells.
Conch. Ex., No. 1, 1888, p. 49.
- *BECK, H. Index Molluscorum præsentis avi Musei Principis Angustissimi Christiani Frederici. Hafniæ, 1837.
- BIELETZKI, N. Ph. Выецкій. Н. Ф. О Авиже-сѣяхъ ноги и створокъ раковины у нашихъ *Unio*, etc.
Труды общ. испыт. природы имп. харьковск. университ. т. XII, XIII (1879) 1880, стр. 115-132.
- BELL, ROBERT. Catalogue of Animals and Plants, collected and observed on the southeast side of the St. Lawrence, from Quebec to Gaspé, and in the counties of Rimouski, Gaspé, and Bonaventure, in 1858.
Geol. Survey Canada, Rept. of Progress for 1858. Montreal, 1859, pp. 251-255.
- BELL, ROBERT, JR. Notes on the Natural History of the Gulf of St. Lawrence, and the distribution of the Mollusca of Eastern Canada.
Can. Nat. and Geol., IV, 1859, pp. 197-220.
- BELON, P. De Aquatilibus, 1553.
- BENEDEN, P. J., VAN. Sur le sexe des Anodontes, et la signification des Spermatozaires.
Bull. de l'Acad. de Brux., XI, Pt. 2, 1844, pp. 377-385; *l'Institut*, XIII, 1845, No. 598, p. 217; *Isis*, 1848, pp. 1019, 1020.
- BENOIST, ÉMILE. Les Unios de la Gironde, etc.
Actes Soc. Linn. Bord., L, p. lxii-lxiv, 1897.
- *BENOIT, LUIGI. Nuovo catalogo delle conchiglie Terrestri e Fluviatili della Sicilia, etc., 1881, pp. 176.
- BENSON, W. H. Descriptive catalogue of terrestrial and fluviatile Testacea, chiefly from the northeast frontier of Bengal.
Jour. Asiat. Soc. Beng., V, 1836, p. 350; II, p. 741.
- *BENSON, W. H. Chusan Shells, collected by Dr. S. Cantor.
Jour. Asiat. Soc. Beng., XXIV, 1855, pp. 119-141.
Reprinted without date, pp. 22.
Summary in *Ann. Nat. Hist.*, 1842, pp. 119-140.
- *— Descriptions of Indian and Burmese Species of the Genus *Unio*, Retz.
Ann. Nat. Hist., X, 1862, pp. 184-195.
- *BERNARDI, A. C. Description d'espèces nouvelles.
Jour. de Conch., VII, 1858, pp. 301-303, pl. x, fig. 1.
- *— Description d'espèces nouvelles.
Jour. de Conch., IX, 1860, p. 331-332, pl. XII, figs. 1, 2, 4.
- BETTA, EDUARDO DE. Intorno agli Studi per una malacologia terrestre e fluviatile dell'Italia.
Atti Ist. Ven. (5), VI, 1880, pp. 419-431.
- BETTA, EDUARDO DE and MARTINELLI, PIETRO PAOLO. Catalogo dei molluschi terrestri e fluviatili viventi nelle provincie Venete, Verona, 1855, pp. 102.
- *BIELZ, E. A. Fauna der Land- und Süßwasser- Mollusken Siebenbürgens, 1863, p. 206; 2d ed., 1867, p. 216.
- *— Verzeichniss der Mollusken und Conchyliensammlung, 1865.
— Systematisches Verzeichniss der Land- und Süßwasser-Mollusken des Österreichischen Kaiserstaates.
Verhandl. und Mittheil. des Siebenbürg. Vereins für Naturwiss. in Hermannstadt, XVI, 1865.
- *BINGLEY, WILLIAM. Useful knowledge, etc., 3 vols., 1825, London.
- *BINNEY, W. G. Bibliography of North American Conchology previous to the year 1860, Pt. 1, American authors, 1863, Pt. 2, Foreign authors, 1864.
Smithsonian Miscellaneous Collections, V, 1864; IX, 1869.
- BLAINVILLE, HENRI-MARIE DUCROTAY DE. Mollusques.
Dict. Sci. Naturelle, 1814, 1824, 1828.
— Sur la classification méthodique des animaux et établissement d'une nouvelle considération pour y parvenir.
Bull. Soc. Philom., 1814, pp. 175-180.
— Observations sur les branches des Anodontes.
Jour. de Phys., LXXXIX, 1825, p. 127.

- * BLAINVILLE, HENRI-MARIE DUCROTAY DE. *Manuel de Malacologie et de Couchyologie*, 1825, Paris, 2 vols., I, 647 pp.; II, 87 colored plates.
- Note sur l'appareil de la génération dans les Moulettes et les Anodontes. *Nour. Bull. Soc. Philom.*, 1825, pp. 156-160, *Fror. Notizen*, XIII, 1826, pp. 6-10.
- * — Rapport fait à l'Académie des Sciences, sur un mémoire de M. Jacobson, ayant pour titre: Observations sur le développement prétendu des œufs des Moulettes et des Anodontes dans leurs branchies. *Ann. des Sci. Nat.*, XIV, 1828, pp. 22-62. Separate.
- * — Rapport de M. de Blainville sur un mémoire de M. de Quatrefages, intitulé: Sur la vie interbranchiale des petites Anodontes. *Ann. des Sci. Nat.*, IV, 1835, pp. 283-290.
- * BLANCHARD, ÉMILE. Observations sur le Système Nerveux des Mollusques Acéphales testacées ou lamelli-branches. *Ann. des Sci. Nat.*, III, 1845, pp. 321-340, plate XII.
- * BLAND, THOMAS. Note sur l'érosion des coquilles fluviales. *Jour. de Conch.*, IV, 1853, p. 308.
- * BLANFORD, WILLIAM T. Contributions to Indian Malacology, No. VII. List of species of *Unio* and *Anodonta* described as occurring in India, Ceylon, and Burma. *Jour. Asiat. Soc. Beng.*, XXXV, 1867, pp. 134-155.
- * — Descriptions of New Land and Fresh-water Molluscan Species, collected by Dr. John Anderson in Upper Burma and Yunnan. *Proc. Zool. Soc.*, 1869, pp. 444-450.
- Land, Fresh-water, and Estuarine Mollusca. *Brit. Burma Gazetteer*, I, 1880, pp. 115-128; *Rep. Brit. Assn. Adv. Sci.*, 1881, p. 678; *Jour. Asiat. Soc. Beng.*, I, 1881, pp. 267, 271.
- [Estimate of the number of bivalve mollusks in India.] *Rep. Brit. Assn. Adv. Sci.*, 1881, pp. 638, 639.
- * BLOXAM, A. [Land and Fresh-water Shells collected in the neighborhood of Rugby, Warwickshire.] *Mag. Nat. Hist.*, VII, 1834, p. 160.
- * — Pearly Matter in Fresh-water Mussels. *Mag. Nat. Hist.*, VII, 1834, pp. 160, 161.
- BOETIUS. Pearls in *Mya Margaritifera*. *Gem. et Lap. Hist.*, 1609. Edited by Bellenden, Edinburgh.
- * BOETTGER, OSKAR. Armenische und trans-kaukasische Mollusken, aus einer Sendung des Herr Dr. G. Sievers in Tiflis. *Jahrb. Deutsch. Mal. Ges.*, VII, 1880, pp. 151-161.
- * — Sechstes Verzeichniss trans-kaukasischer, armenischer und nordpersischer Mollusken. *Jahrb. Deutsch. Mal. Ges.*, VIII, 1881, pp. 167-261, plates VII-IX.
- Beiträge zur Herpetologie und Malakozoologie Südwest Afrikas. *Ber. d. Senck. naturf. Ges.*, 1886, pp. 3-29, 2 plates.
- * — Malakologische Ergebnisse auf Streifzügen in Thessalien. *Jahrb. Deutsch. Mal. Ges.*, XIII, 1886, pp. 42-73. 2 plates.
- [Cf. Stussiner] Auf Zählung der in Thessalien gesammelten Schnecken und Muscheln. *Jahrb. Deutsch. Mal. Ges.*, XIII, 1886, pp. 50-73, pl. II.
- * — Ad. Strubell's Konchylien aus Java. *Ber. d. Senck. naturf. Ges.*, 1890, pp. 137-173, pls. V, VI, II; pp. 241-312, pls. III, IV.
- * — Zur Mollusken-Fauna russischen Gouvernements Perm und des Gebietes Süd-östlich von Orenburg. *Nach. Deutsch. Mal. Ges.*, 1890, pp. 161-173.
- * — Die Verhältnisszahlen der Hauptschalendimensionen bei den Paläarktischen Art in der Gattung *Unio* Retz. *Nach. Deutsch. Mal. Ges.*, 1893, pp. 65-79; 1898, pp. 33-48.
- * BOJANUS, LUDWIG HEINRICH. Sur les Organes respiratoires et circulatoires des coquillages bivalves en général, et spécialement sur ceux de l'*Anodonta* des Cygnes. *Jour. de Phys.*, LXXXIX, 1819, pp. 108-134, 1 pl.

- * BOJANUS, LUDWIG HEINRICH. Send-schreiben an M. le B^{on} G. Cuvier: Ueber die Athmungs und Kreislaufs-Werkzeuge der zweischaligen Muscheln besonders des *Anodon cygneum*, 1818, 12 pp., 2 plates.
Jour. de Phys., LXXXIX, 1819, p. 108, pl.; *Isis*, 1819, p. 42, pls. 1, II; 1820, p. 404; 1827, p. 752, pl. IX. Jena, 1821, 4 plates.
- BOLTEN, J. F. Museum Boltenianum, 1798; 2d ed., 1819.
- BONANNI, P. Mentis et oculi in observatione animalium Testaceorum curiositate naturae inspectoribus italicis, etc., 1684.
— Testacea Theologia. 1744.
- BONHOMME, JULES. Notice sur les Mollusques bivalves fluviatiles observés jusqu'à ce jour aux environs de Ro-dez.
Mém. Soc. Aveyr., II, 1840, p. 429.
- BONI, CARLO. Le valve del Unio nella Terra mara del Montale.
Ann. Soc. Nat. Modena, VI, 1872, pp. 171-174.
- BORCHERDING, F. Zweiter Nachtrag zur Mollusken-Fauna der nordwest-deutschen Tiefebene.
Abh. ver. Brem., IX, 1884, pp. 141-166, pl. III.
- BORN, IGNATIUS V. Elements de conchyliologie. 1776.
- * — Testacea Musei Cæsarei Vindobonensis quae jussu Mariae Theresiae Augustae disposuit et descripsit. 1780, 18 plates.
- * BOSCH, L. A. C. Histoire naturelle des Coquilles, contenant leur description, les mœurs des animaux qui les habitent et leurs usages. Avec figures dessinées après nature. 5 vols. 43 colored plates, 1804; 2d ed., 1830; 3d ed., without date.
- BOUCHARD, CHAUTEREAUX. Catalogue des Mollusques terrestres et fluviatiles observés jusqu'à ce jour à l'état vivant dans le département du Pas-de-Calais. 1838, 89 pp., 1 plate.
- BOUILLET, J. B. Catalogue des Mollusques et fluviatiles vivants du département du Puy-de-Dôme.
Ann. Aveyr., V, 1832, p. 176. Published separately.
- BOUILLET, J. B. Coquilles fossiles du calcaire d'eau douce du Cantal.
Bull. Soc. Géol. France, VI, 1834, p. — Catalogue des espèces et variétés de Mollusques terrestres et fluviatiles dans la Haute et la Basse-Auvergne suivi (1836) d'un autre catalogue des espèces fossiles recueillies récemment dans les diverses formations tertiaires des mêmes départements. 1836.
- * BOURGUIGNAT, J. R. Testacea Novissima quae Cl. de Sauley in itinere per Orientem annis 1850 et 1851 collegit. Paris, 1852, pp. 31. A preliminary description of shell obtained by F. de Sauley.
* — Description de quelques coquilles provenant de Syrie.
Jour de Conch., IV, 1853, pp. 69-74, pl. III.
- * — Catalogue raisonné des Mollusques terrestres et fluviatiles recueillis par F. de Sauley, pendant son voyage en Orient. Paris, 1853, 96 pp., 4 plates. From Voyage autour de la Mer Morte, par Fr. de Sauley.
- * — Aménités malacologiques.
Rev. et Mag. de Zool., V, 1853, p. 340; VII, 1854, pp. 44, 75, 658; VII, 1855, pp. 23, 317, 556; VIII, 1856, pp. 7, 66, 226, 268, 327, 378, 424, 499. Separate.
- * — Malacologie du lac des Quatre-Cantons et de ses environs.
Rev. et Mag. de Zool., XIV, 1862, pp. 430, 465, pls. XIX-XXI; XV, 1863, p. 5.
- * — Mollusques nouveaux litigieux ou peu connus; mars, 1863, décembre 1868, pp. 324, 46 plates. First appeared in *Rev. et Mag. de Zool.*
- * — Malacologie de l'Algérie et Histoire Naturelle des Animaux Mollusques Terrestres et Fluviatiles, recueillis jusqu'à ce jour dans nos possessions du nord d'Afrique. 2 vols., 1864, pp. 674, 58 plates.
- Malacologie d'Aix-les-Bains (Savoie.) Paris, 1864, pp. 78, 3 plates.
- Malacologie de la Grande-Chartraine (Dauphiné). Paris, 1864. 9 lithographic views, 8 colored plates.
- Recherches sur la distribution géographique des Mollusques terrestres et fluviatiles en Algérie, et dans les régions circonvoisines.
Ann. Sci. Nat., Zool., VI, 1866, p. 313.

- * BOURGUIGNAT, J. R. Aperçu sur la Faune malacologique du Bas-Danube. *Ann. de Mal.* 1870, pp. 37-76.
- * — Description de diverses espèces terrestres et fluviatiles et de différents genres de Mollusques de l'Égypte, de l'Abyssinie, de Zanzibar, du Sénégal et du Centre de l'Afrique. 1879.
- * — Matériaux pour servir à l'Histoire des Mollusques Acéphales du Système Européen, 1880-81, pp. 387.
- Lettres Malacologiques, 1882, with plate.
- * — Histoire Malacologique de l'Abyssinie. *Ann. des Sci. Nat.*, 6th ser., XV, 1883, pp. 47-162. Separate.
- * — Aperçu sur les Unionidæ de la Péninsule Italique, 1883, pp. 117.
- * — Mollusques fluviatiles du Nyanza Onkéréwe, suivis d'une note sur les genres *Cameronia* et *Burtouia* du Tanganyika, 1883, pp. 23, 1 plate.
- * — Espèces nouvelles et genres nouveaux découverts par les rev. pères missionnaires dans les grands lacs africains Onkéréwe et Tanganyika, 1885, pp. 39.
- * — Mollusques Terrestres et Fluviatiles recueillis par M. Paul Soleillet dans son Voyage au Choa, 1885, pp. 45. 1 plate.
- * — Notice prodromique sur les Mollusques terrestres et fluviatiles recueillis par M. Victor Giraud dans la région méridionale du lac Tanganyika, 1885, pp. 110.
- * — Monographie d'un nouveau genre d'Acéphale du Lac Tanganyika. *Bull. Soc. Mal. France*, II, 1885, pp. 1-12, 1 plate.
- * — Unionidæ et Iridinidæ du Lac Tanganyika, 1886, pp. 93.
- * — Mollusques nouveaux de la région du Nyanza Onkéréwe. *Bull. Soc. Mal. France*, IV, 1887, pp. 267-272.
- * — Iconographie malacologique des animaux mollusques fluviatiles du lac Tanganyika, 1888, pp. 82, 35 plates.
- * — Mollusques de l'Afrique équatoriale de Mognedouchou à Bagamoyo et de Bagamoyo au Tanganyika, 1889, pp. 229, 8 plates.
- * BOURGUIGNAT, J. R. Melanidées du lac Nyassa snivis d'un aperçu comparatif sur la faune malacologique du lac, avec cette du grand lac Tanganyika. *Bull. Soc. Mal. France*, VI, 1889, pp. 1-66, plates 1, II.
- * BOUSSUET, FRANCISCUS. Francisci Boussuet Surregiani doctoris Medici de Natura Aquatiliu Caruen in universam Gulielmi Roudeletii doctoris quam de Piscibus marinis scripsit historiam; cum vivis eorum imaginibus emissum. Lugduni, 1558.
- * BOWDICH, T. EDWARD. Elements of Conchology. 2 vols., Paris, 1822, 500 figs.
- * BRARD, C. J. P. Histoire des Coquilles Terrestres et Fluviatiles qui vivent aux environs de Paris, 1815. 10 colored plates.
- * BRAUN, MAX. Über die postembryonale Entwicklung unserer Süßwassermuscheln. *Verh. d. Phys.-Med. Ges. Würz. N. T.*, XIII, 1879; *Sitzber.* 1878, pp. xxiv-xxvii; *Jahrb. Deutsch. Mal. Ges.*, 1878, pp. 307-319; *Arch. Zool. Exp. et Gén.*, VII, 1878, N. and R., p. xlvii. Extract in *Rev. internat. des Sci.*, II, 1878, pp. 634-636.
- [Observations on larvæ of *Anodonta*.] *S. B. Ges. Dorp.*, 1882, pp. 429-431.
- * — Zur Molluskenfauna der Ostsee provinzen. *Nach. Deutsch. Mal. Ges.*, 1883, pp. 174-181.
- Beiträge zur Kenntniss der Fauna baltica, II. Die Land- und Süßwasser-Mollusken der Ostsee-Provinzen. *Arch. Nat. Litv.* 2d ser., IX, 1884, Pt. 5, p. 102.
- * — Die postembryonale Entwicklung der Najaden. *Nach. Deutsch. Mal. Ges.*, 1889, pp. 14-19.
- Ueber parasitische Lamellibranchier zusammenfassender Bericht. *C. B. Bakt. Parasit.*, V, 1889, pp. 241-248; 276-282.
- * BRAZIER, J. Remarks on some fluviatile shells of New South Wales. *Proc. Linn. Soc. N. S. W.*, VII, 1882, pp. 83-86.

- BREVIERE, L. Catalogue des mollusques testacés, terrestres et fluviatiles observés dans le département de la Nièvre, 1880, pp. 30.
- BRIDGMAN, JOHN B. A variety caused by locality.
Quart. Jour. Conch., I, 1874-1878, p. 70.
- BRODIE, JAMES. The Pearls of the Ythan, Aberdeenshire.
Proc. Roy. Phys. Soc., Edinb., III (1862-1866), 1867, pp. 394-396.
- BROECK, ERNST VAN DEN. Excursions découvertes et observations malacologiques faites en Belgique pendant l'année 1870.
Ann. Mal. Belg., V., 1871, pp. 1-64.
- *BRONN, HEINRICH G. System der Urweltlichen Pflanzen-Thiere durch Diagnose, Analyse und Abbildung der Geschlechter erläutert. Heidelberg, 1825.
- *— Lethæa Geognostica, oder Abbildungen und Beschreibungen der für die Gebirgs Formationen bezeichnendsten versteinerungen, with 47 lithographs. 2 vols., text and atlas, 1835-1838.
- Index Palæontologicus, 3 vols., 1849.
- Die Klassen und Ordnungen der Weichthiere (Malacozoa), etc., III, Pt. 1, 1862.
- *BROOKES, SAMUEL. An Introduction to the Study of Conchology, including observations on the Linnæan Genera and on the arrangement of M. Lamarck, 1815, pp. 160, 11 plates.
- *BROOKS, WILLIAM K. Embryology of the Fresh-water Mussels.
Proc. Amer. Assn. Adv. Sci., 1876, pp. 238-240.
- *BROT, A. Étude sur les coquilles de la Famille des Náyades qui habitent le Bassin du Léman.
Ass. Zool. Lem., 1866, pp. 56, 9 plates. Separate.
- *BROWN, ALEXANDER. In Deshayes *Traité Élémentaire*, II, 1850, p. 216.
- *BROWN, A. D. Catalogue of shells in the collection of A. D. Brown, 1861.
- *BROWN, THOMAS. Elements of Conchology, 1816.
- Account of Irish Testacea.
Mem. Werner. Soc., II, 1818, p. 501, pl. XXIV.
- *BROWN, THOMAS. Illustrations of the Recent Conchology of Great Britain and Ireland, 1827, 57 colored plates.
- Description of five new British species of Shells.
Edinb. Jour. Nat. and Geog. Sci., I, 1830, pp. 11-12.
- *— The Conchologist's Text-Book, embracing the arrangements of Lamarck and Linnæus, with a glossary of technical terms, 1833, pp. 180, 19 plates.
- *— Conchology of Great Britain and Ireland, 1839.
- Elements of Fossil Conchology, 1843, p. 91.
- Illustrated Conchology of Great Britain, 1844.
- Article Conchology, *Encyc. Brit.*, 6th ed., VI, no date.
- *BRUGUIÈRE, JEAN GUILLAUME. Encyclopédie Méthodique. VI. Histoire Naturelle des Vers, Pt. 1, 1789; II, 1792.
- *— Sur une nouvelle espèce de Mulette.
Jour. d'Hist. Nat., I, 1792, pp. 103, 109.
- *— Choix de mémoires sur divers objets d'histoire naturelle formant les collections du Journal d'Histoire Naturelle, I, 1792, p. 131.
- *— Sur une nouvelle coquille du genre de l'Anodontite [*Anodonta crispata*].
Jour. d'Hist. Nat., I, 1792, pp. 131-136.
- *BRUMATI, LEONARDO. Catalogo Systematico Conchyliæ terrestre e fluviatile osservate nel territorio de Monfalcon e dall'abate, 1838, pp. 56, 1 lithograph.
- *BRUSINI, SPIRIDIONE. Fauna Molluschi Dalmatica, 1866, pp. 134. 1 plate.
- Ueber die Mollusken-Fauna Oesterreich Ungarns, 1886, p. 30.
- BUCAILLE, E. Catalogue raisonné des Mollusques terrestres et d'eau douce, observés dans le département de la Seine-Inférieure.
Bull. Soc. Rouen, XXVII, 1892, pp. 171-189.
- BUCHOZ, PIERRE-JOSEPH. Collection de Plantes (fossil), 1782.
- *BUTTON, FRED. L. Unionidæ in a Tunnel.
Nautilus, XIII, 1900, p. 130.

- BUVIGNIER, AMAND. Catalogue des Mollusques du département de la Meuse.
Act. Philom. Ferd., 1840, p. 217.
- BUXBAUM, GEBRÜDER. Das Wachsen der Anodonten.
Zool. Gart., XXXI, 1890, pp. 16-18.
- *CAILLAUD, FRÉDÉRIC. Voyage à Méroë et au fleuve Blanc de 1819 à 1822, 4 vols. Mollusques terrestres et fluviatiles, II, IV, 1827.
— Catalogue des Radières, des Annélides, des Cirrhipèdes et des Mollusques marins, terrestres et fluviatiles, recueillis dans le département de la Loire-Inférieure, Nantes, 1865, pp. 323, 5 plates.
- *CALKINS, W. W. The Land and Fresh-water Shells of Lasalle County, Illinois.
Pr. Ottawa Acad. Nat. Sci., 1874, pp. 48, 1 plate.
— Notes on Fresh-water Mollusks found in the vicinity of Chicago, Illinois.
Cinn. Quart. Jour. Sci., I, 1874, pp. 242-244.
— [Unio buckleyi and buddianus united.]
Valley Naturalist, II, 1880, p. —.
- *CALL, R. ELLSWORTH. Mode of Distribution of Fresh-water Mussels.
Am. Naturalist, XII, 1878, pp. 472, 473.
*— Synonymous Unios.
Am. Naturalist, XIII, 1879, pp. 392, 393.
*— Polymorphus Anodontæ.
Am. Naturalist, XIV, 1880, pp. 529, 530;
Jour. of Conch., 1881, p. 186.
*— Unio bolli, new species.
Am. Naturalist, XV, 1881, pp. 390, 391.
*— On the Quaternary and Recent Mollusca of the Great Basin, with descriptions of new forms.
Bull. U. S. Geol. Surv., No. 11, 1884.
*— A Geographic Catalogue of the Unionidæ of the Mississippi Valley.
Bull. Des Moines Acad. Sci., II, 1885, pp. 5-57.
*— Contributions to a Knowledge of the Fresh-water Mollusks of Kansas, 1st to 5th Contributions.
Bull. Washb. Coll. Lab., I, 1885, pp. 49-54, 93-97, 115-124, 177-184.
Proc. N. M. vol. xxii—60
- *CALL, R. ELLSWORTH. Sixth contribution to a knowledge of the Fresh-water Mollusks of Kansas
Sixth Rep. Biol. Surv. Kans., 1887, pp. 11-25, 1 figure.
*— Description of a new species of Unio from Kansas.
Bull. Washb. Coll., 1885, pp. 48, 49.
*— Descriptions of two new species of the genus Unio, from the Ozark region of Missouri.
Proc. U. S. Nat. Mus., 1887, pp. 498-500, 2 plates.
*— Note on the Ctenidium of Unio aberti Conrad.
Am. Naturalist, XXI, 1887, pp. 857-860, figures.
*— The Unionidæ of the Ohio river. (Abstract.)
Proc. Ind. Acad. Sci., IV, (1894) 1895, pp. 139-140.
*— On a small collection of Mollusks from northern Indiana.
Proc. Ind. Acad. Sci., 1895, pp. 246-250.
*— A Revision and Synonymy of the Parvus Group of Unionidæ, with 6 plates.
Proc. Ind. Acad. Sci., 1895, pp. 109-119.
*— Second contribution to a knowledge of Indiana Mollusks.
Proc. Ind. Acad. Sci., 1895, pp. 135-146.
*— A study of the Unionidæ of Arkansas, with incidental reference to their distribution in the Mississippi Valley.
Trans. Acad. Sci. St. Louis, VII, 1895, pp. 1-65, 21 plates. Abstract in *Jour. Roy. Mic. Soc.*, 1895, p. 621.
— Illustrations of little known Unionidæ.
Jour. Cinn. Soc. Nat. Hist., XVIII, 1896, pp. 157-186, 1 plate.
- *CANTOR, T. [List of Shells from Chusan.]
Proc. Zool. Soc., 1842, p. 124.
- *CARLSON, H. P. Shells of Antioch, California, and vicinity.
Proc. Cal. Acad. Sci., IV, 1870, pp. 50-52.
*— On the shells of Truckee River, California, and vicinity.
Proc. Cal. Acad. Sci., IV, 1870, p. 57.

- *CARPENTER, HORACE. The Shell Bearing Mollusca of Rhode Island.
In *Random Notes, Conch. Exchange and Nautilus*. Published separately.
- *CARPENTER, P. P. Catalogue of the Collection of Mazatlan Shells in the British Museum, collected by Frederick Reigen, 1857, pp. 552.
- *— Report on the Present State of our Knowledge with regard to the Mollusca of the West Coast of North America.
Rep. Brit. Assn. Adv. Sci., 1856, pp. 159-368, 4 plates.
- *— Lectures on Mollusks.
Smithsonian Report, 1860, p. 117.
- *— The Mollusks of Western North America, 1872.
Smithsonian Miscellaneous Collection, XII, pp. 446.
- CARPENTER, WILLIAM B. Principles of General and Comparative Physiology. London, 1839.
- *CARUS, C. G. Neue Untersuchungen über die Entwicklungsgeschichte unserer Flussmuscheln, 4 plates.
Nova Acta. Acad. Leop. Carol., XVI, 1832, pp. 1-87. Separate.
- *CASTELNAU, FRANCIS DE. Expedition dans l'Amérique du Sud, 6 vols. Paris, 1857. (See HUPÉ, Animaux Nouveaux ou Rares.)
- *CASTRO, JOSÉ DA SILVA E. Unionidæ nouvelles du Portugal.
Bull. Soc. Mal. France, II, 1885, pp. 277-294.
- CATALOGUE OF BOSTON SOCIETY OF NATURAL HISTORY.
In *Hitchcock's Rep. on Geol. Bot. and Zool. of Mass.*, 1835, pp. 25-28.
- CATLOW, AGNES. Popular Conchology, 1852.
- *CATLOW, AGNES and REEVE, LOVELL. The Conchologists Nomenclator. A catalogue of all the Recent Species of Shells, etc., 1845, pp. 326.
- *CHAPER, M. Note sur quelques Unionidæ de Grèce.
Mém. Soc. Zool. France, VII, 1894, pp. 372-374, 2 plates.
- *— Description de quelques espèces et genres nouveaux de coquilles vivantes de diverses provenances.
Bull. Soc. Zool. France, X, 1885, pp. 42-54. Reprint. pp. 14. 1 plate.
- *CHAPER, M. Description de quelques espèces nouvelles de coquilles vivantes provenant de l'Afrique Australe et d'Assinie.
Bull. Soc. Zool. France, X, 1885, pp. 479-486, pl. xi.
- *CHARMÉS, XAVIER. Unionidæ des environs de Bagamoyo (Zanguebar).
Bull. Soc. Mal. France, II, 1885, pp. 165-174.
- CHARPENTIER, JEAN G. F. DE. Catalogue des Mollusques terrestres et fluviatiles de la Suisse.
Neue Deutschr. Allg. Schweiz. Ges. I, 1837.
Separate.
- CHATIN, M. J. Sur la différenciation du protoplasme dans les fibres nerveuses des Unionides.
Comp. Rend., xciv, 1882, p. 1723-1726.
- *CHEMNITZ, JOHANN HIERONYMUS. Neues systematisches Conchylien-Cabinet, 1769-1795, 11 vols.; plates.
- *CHENU, J. C. Illustrations Conchyliologiques, ou description et figures de toutes les coquilles connues vivantes et fossiles, etc., 4 vols., colored plates, and some text, 1843-1850? Not finished. A large number of plates appeared without accompanying text. Only a few species were described.
- *— Bibliothèque Conchyliologique, in two series, 1845-1855.
- *— Manuel de Conchyliologie et de Paléontologie Conchyliologique, 2 vols., 1859-1862, pp. 835; figures.
- CHERRÉS, COLLARD DES. Catalogue des Testacés terrestres et fluviatiles des environs de Brest et de Quimper (Finistère).
Act. Soc. Linn. Bord., IV, 1830, p. 91.
Separate.
- CHICKERING, J. W., JR. List of Marine, Fresh-water, and Land Shells found in the vicinity of Portland, Maine. [1855 or 1856?]
- CHILDREN, J. G. Translation of Lamarck's Genera.
Braud's Journal, XV, 1823.
- *CHRISTY, R. M. Notes on the Land and Fresh-water Mollusca of Manitoba.
Jour. of Conch., IV, 1855, pp. 339-349.
- *CLAPP, GEORGE H. Mollusca of Allegheny County, Pennsylvania.
Nautilus, VIII, 1895, p. 116.

- * CLESSIN, S. Die Molluskenfauna des Starnberger Sees.
Mal. Blatt., XIX, 1872, pp. 109-125.
 — Studien über die deutschen Species des Genus *Anodonta*, Cuvier.
Correspbl. d. Zool.-Min. Ver. Reg., XXVI, 1872, pp. 82-90, 97-104.
 — Beitrag zur Molluskenfauna der oberbayrischen Seen.
Correspbl. d. Zool.-Min. Ver. Regensb., XXVII, 1873, pp. 56-179.
- * — Die Genera der recenter Süßwasser-Bivalven.
M. T. Ver. Reich, 1874. Separate.
- * — *Anodonta complanata* Z.
Nach. Deutsch. Mal. Ges., 1874, pp. 85, 86.
- * — Die Jahresringe der Süßwasser-Bivalven.
Nach. Deutsch. Mal. Ges., 1874, pp. 25-29.
- * — Die Familie der Najaden.
Mal. Blatt, XXII, 1874, pp. 1-29, 1 plate.
- * — Deutsche Excursions-Mollusken-Faunen, Nürnberg, 1876, pp. 581, figs.
- * — Unionen aus dem mittleren Russland.
Jahrb. Deutsch. Mal. Ges., VII, 1880, pp. 79-82.
- * — Neue Arten.
Mal. Blatt, 1882, pp. 187-193, pl. iv.
- * — *Anodonta Iheringi*, new species.
Mal. Blatt, 1882, p. 193, pl. iv.
- * — Die Molluskfaunen Oesterreich-Ungarns der Schweiz, 1887, 320 pp., figs.
- * — Binnenmollusken aus Süd Brasilien.
Mal. Blatt, X, 1888, pp. 165-174.
- Studien über die deutschen Species des Genus *Anodonta*.
Corres. Blatt. Ver. Regensb., XXXVI, 1872, pp. 82-97.
- CLOQUET, J., and MOQUIN-TANDON, M. Observations sur les perles des Bivalves d'eau douce.
Bull. Soc. Imp. Zool. d'Acclimat., V, 1858, pp. 452-461.
- COCKERELL, T. D. A. The British Naturalist's Catalogue of the Land and Fresh-water Mollusca of the British Isles, etc., 1890, pp. 19.
- COLENZO, W. [*Unio waikarensis* n. sp.]
Tasm. Jl. Nat. Sci., II, 1841, p. —.
- * COLENZO, W. Description of two little-known species of New Zealand Shells.
Trans. New Zealand Inst., XIV, 1882, pp. 168, 169.
- COLLINGE, W. E. The Land and Fresh-water Mollusca of Ingleton Clapham and District.
Naturalist, 1890, pp. 109-114.
 — Note on a *Tumour* in *Anodonta Cygnea*.
Jour. Anat. Phys., XXV, 1890, p. 154.
- * — The Land and Fresh-water Mollusca of Oxfordshire.
Conchologist, 1891, pp. 11-14, 19-23, 39-44, 51-53.
- COMPANYO and MASSOT, PAUL. Description d'une nouvelle espèce de *Mulette*.
Bull. Soc. Agr. Sc. Pyr. Or., VI, 1845, p. 234, fig. 2.
- * CONRAD, TIMOTHY A. Description of Some New Species of Fresh-water Shells from Alabama, Tennessee, etc.
Am. Jour. Sci. and Arts, XXV, 1834, pp. 338-343, 1 plate.
- * — New Fresh-water Shells of the United States. * * * Also a Synopsis of the American Naiades, 1834, pp. 73. 8 colored plates.
- * — Description of a new genus of Fresh-water Shells.
Jour. Acad. Nat. Sci. Phila., VII, 1834, pp. 178-180.
- Appendix to new Fresh-water Shells, 1835.
- * — Monography of the Family Unionidæ or Naiades of Lamarck. Colored plates, I, 1835; II-VII, 1836; VIII, IX, 1837; X, XI, 1838; XII, XIII, no date.
- * — Descriptions of three new species of *Unio*.
Proc. Acad. Nat. Sci. Phila., I, 1841, pp. 19, 20.
- * — Descriptions of three new species of *Unios* from the rivers of the United States.
Jour. Acad. Nat. Sci. Phila., VIII, 1842, pp. 178-180.
- * — Observations on the Geology of a part of East Florida, with a Catalogue of Recent Shells of the Coast.
Am. Jour. Sci. and Arts, II, 1846, pp. 36-45.

- * CONRAD, TIMOTHY A. Notices of Fresh-water Shells, etc., of Rockbridge County, Virginia.
Am. Jour. Sci. and Arts, I, 1846, p. 407.
- *— Descriptions of New Fresh-water Shells.
Ann. and Mag. Nat. Hist., IV, 1849, pp. 300-303.
- *— Descriptions of New Fresh-water and Marine Shells.
Proc. Acad. Nat. Sci. Phila., IV, 1849, pp. 152-155; *Ann. Nat. Hist.*, IV, 1849, p. 300.
- *— Descriptions of New Species of Fresh-water Shells.
Proc. Acad. Nat. Sci. Phila., V, 1850, pp. 10, 11.
- *— Descriptions of New Fresh-water and Marine Shells.
Jour. Acad. Nat. Sci. Phila., 1850, pp. 275-278, 2 plates.
- *— Notes on Shells, with Descriptions of New Species.
Proc. Acad. Nat. Sci. Phila., VI, 1852, pp. 199, 200.
- *— A Synopsis of the Family of Naiades of North America, with Notes, and a Table of some of the genera and subgenera of the Family.
Proc. Acad. Nat. Sci. Phila., VI, 1853, pp. 243-269.
- *— Omissions and Corrections to the Synopsis of North American Naiades.
Proc. Acad. Nat. Sci. Phila., VI, 1853, p. 449.
- *— Notes on Shells, with Descriptions of three Recent and one Fossil Species.
Proc. Acad. Nat. Sci. Phila., VII, 1854, p. 32.
- *— Descriptions of New Species of Unio.
Jour. Acad. Nat. Sci. Phila., 1854, pp. 295-298, 2 plates.
- *— Descriptions of three New Species of Unio.
Proc. Acad. Nat. Sci. Phila., VII, 1855, p. 256.
- *— On a New Species of Unio, and Remarks on Unio viridis.
Am. Jour. Sci. and Arts, XXI, 1856, p. 172.
- *— Descriptions of two New Genera of Shells.
Proc. Acad. Nat. Sci. Phila., 1857, p. 165, 166.
- * CONRAD, TIMOTHY A. Remarks on the genera *Monocordylea* d'Orb, and *Pseudodon* Gould, with a synopsis of the latter.
Amer. Jour. Conch., I, 1865, pp. 232, 233.
- *— Description of a New Genus of Unionidae.
Amer. Jour. Conch., I, 1865, p. 234.
- *— Description of a New Species of *Pseudodon*.
Amer. Jour. Conch., I, 1865, p. 352, fig. 2.
- *— Description of a New Species of Unio.
Am. Jour. Conch., II, 1866, p. 107.
- *— Observations on *Pleiodon Macmurtrii*.
Am. Jour. Conch., III, 1867, p. 4.
- *— Notes on Recent and Fossil Shells, with Descriptions of New Genera.
Am. Jour. Conch., IV, 1868, p. 246-249.
- *— Description of a new Unio and fossil *Goniobasis*.
Am. Jour. Conch., IV, 1868, p. 280.
- *— Notes on Recent Mollusca.
Am. Jour. Conch., V, 1869, pp. 104-108.
- * COOKE, A. H. On the Geographical Distribution of the Land and Fresh-water Mollusca of the Malagasy Region.
Conchologist, II, 1893, pp. 131-139.
- *— Molluscs.
In *The Cambridge Natural History*, III, 1895, pp. 535, figures and maps.
- COOPER, D. A List of Land and Fresh-water Shells of the Environs of London. 1836. Extract from *Flora Metropolitana*.
- COOPER, J. G. Geographical Catalogue of the Mollusca found west of the Rocky Mountains between latitude 33° and 49° north. 1867.
- Catalogue of Mollusks.
Geol. Survey of California, 1867.
- *— Notes on the Fauna of the Upper Missouri.
Am. Naturalist, III, 1870, pp. 294-299.
- *— [Fresh-water Mollusks of the Missouri River.]
Am. Naturalist, III, 1870, pp. 405-422.

- * COOPER, J. G. *The Sylva of Montana*, 1870.
- * — Fresh-water Mollusca of San Francisco County.
Zoe, I, 1890, pp. 196, 197.
- * — On some Pliocene Fresh-water Fossils of California.
Proc. Cal. Acad. Sci., IV, 1894, pp. 166-172, pl. XIX.
- * COOPER, WILLIAM. List of Shells collected by Mr. Schoolcraft in the Western and Northwestern Territory.
- * — Appendix No. 2 to Narrative of an Expedition to the Sources of the Mississippi River in 1820. Resumed and completed by the discovery of its origin in Lake Itasca. Narrative by Henry Schoolcraft, 1855, pp. 515-518.
- * — Report upon Mollusca collected by the Survey. Survey of Minnesota, Nebraska, etc., 1859, p. 381.
Pacific Railroad Survey, XII, Pt. II, *Rep. on Nat. Hist.*, etc., 1859, pp. 369-386.
- COSTA, EMMANUEL MENDES DA. Elements of Conchology, or Introduction to the Knowledge of Shells, 1776, 7 plates.
- * — *Historia Naturalis Testaceorum Britanniae*, or The British Conchology, etc., 1778, 17 colored plates; 2d. ed., 1780.
- * COSTA, O. G. *Catalogo Systematico e ragionato de Testacei delle due Sicilie*, 1829, 2 plates.
- COOPER, J. HAMILTON. Catalogue of the Recent Shells of Georgia. In appendix to *White's Statistics of Georgia*. No date. [1849].
- * CRAGIN, F. W. A New Species of *Unio* from Indian Territory.
Bull. Washb. Coll. Lab., 1887, p. 6.
- * CRAVEN, A. E. On a collection of land and fresh-water shells from the Transvaal and Orange Free State in South Africa, with descriptions of nine new species.
Proc. Zool. Soc., 1880, pp. 614-618, pl. LVII.
- CRISTOFORI, J. DE, and JAN, GEORGES. *Catalogus in IV sectiones divisus verum Naturalium in museo extantium Josephi de Cristofori et Georgii Jan., etc.*, 1832.
- * CROSSE, H. [Remarks on *Anodonta gougetana*, etc.]
Jour. de Conch., 1862, pp. 203, 204.
- * — [Remarks on the number of *Lea's Unionidae*.]
Jour. de Conch., 1863, pp. 310-312.
- * — [On the Rectification of Mr. Conrad's Synopsis.]
Jour. de Conch., 1868, p. 372.
- * — Variétés. Ne Sutor ultra crepidam.
Jour. de Conch., 1870, pp. 251-260.
- * — Variétés. [Reviews of *Conch. Icon.*, etc.]
Jour. de Conch., 1870, pp. 251-260.
- * — Faune Malacologique du Lac Tanganyika.
Jour. de Conch., XXIX, 1881, pp. 105-139, pl. IV.
- * — Supplément à la Faune Malacologique du Lac Tanganyika.
Jour. de Conch., 1881, pp. 277-306.
- * — Note Complémentaire sur l'*Anodonta*.
Jour. de Conch., 1883, pp. 221, 222, pl. IX, fig. 4.
- * — Faune Malacologique, terrestre et fluviatile, de l'île de la Trinité.
Jour. de Conch., 1890, pp. 35-65.
- * — Faune Malacologique, terrestre et fluviatile, de l'île de Cuba.
Jour. de Conch., 1890, pp. 173-335.
- * CROSSE, H., and DEBEAUX, O. Notes sur quelques espèces nouvelles ou peu connues du littoral de l'empire Chinois.
Jour. de Conch., 1863, pp. 253-265.
- * CROSSE, H., and FISCHER, PAUL. Mollusques Fluviales, recueillis au Cambodge, par la mission scientifique française de 1873.
Jour. de Conch., 1876, pp. 313-342, pl. X.
- * — Nouvelles.
Jour. de Conch., 1875, pp. 194-196.
- * — Description d'une variété nouvelle d'*Anodonta*, recueillie par M. A. Forrer, dans l'État de Sinaloa.
Jour. de Conch., 1883, pp. 219-221.
- * — Diagnosis Molluscorum Republicæ Mexicanæ et Guatemalae incolorum.
Jour. de Conch., XL, 1892, p. 294.

- *CROSSE, H., and FISCHER, PAUL. Description d'un Bulimulus et d'un Anodonta nouveaux, provenant du Mexique.
Jour. de Conch., 1893, pp. 31, 32.
- *— Diagnosis Molluscorum novi, Republicæ Mexicanæ incolæ.
Jour. de Conch., 1893, p. 179, 293-297.
- *CROUCH, EDMUND A. An Illustrated Introduction to Lamarck's Conchology, contained in his Histoire Naturelle des Animaux sans Vertèbres, etc., 1827. 22 colored lithographs.
- *CROWTHER, HENRY. Unio margaritifera at Whitby, etc.
Naturalist, IV, 1878, p. 40.
- *CUMING, HUGH. List of Mollusks from Formosa.
Proc. Zool. Soc., 1865, pp. 196, 197.
- CURIER, A. O. List of Shells collected in the Grand River Valley, 1859?
- *— Catalogue of the Mollusca of Grand Rapids, Michigan.
Am. Jour. Conch., I, 1865, pp. 292-296.
- *— Descriptions of New Species of Fresh-Water Shells of Michigan.
Am. Jour. Conch., III, 1867, pp. 112, 113, pl. vi.
- *— List of the Shell-bearing Mollusks of Michigan, especially of Kent and adjoining counties.
Kent Sci. Inst., Grand Rapids, Mich., 1868, pp. 12.
- CUVIER, GEORGES. Tableau élémentaire de l'histoire naturelle des animaux. Paris, 1798.
- *— Le Règne Animal, etc. Paris, 1817, 4 vols.; 2d ed., 1829-30, 5 vols., 20 plates; ed. with colored engravings, no date; Mollusks by G. P. Deshayes, 1 vol. with atlas, 152 plates, ed. by Griffith, 1834.
- *DALL, WILLIAM H. Revision of the Classification of the Mollusca of Massachusetts.
Proc. Bost. Soc. Nat. His., XIII, 1870, pp. 240-257.
- *— On the hinge of Pelecypods and its development, with an attempt toward a better subdivision of the group.
Am. Jour. Sci. and Arts, XXXVIII, 1889, pp. 445-462.
- *— Diagnosis of New Mollusks from the Survey of the Mexican Boundary.
Proc. U. S. Nat. Mus., XVIII, 1895, p. 1.
- DAMBECK, CARL. Die Fluss-Perlenmuschel.
Natur. (Muller) new ser., V, 1879, pp. 171-175.
- *DANA, JAMES D. Manual of Geology. 1 vol., chart, and figs., 1862. 2d ed., 1874; 3d ed., 1880; 4th ed., 1895.
- *DANIEL, FELIX. Faune Malacologique terrestre, fluviatile et marine, des environs de Brest.
Jour. de Conch., 1883, pp. 223-263, 330-391; Supplement *Jour. de Conch.*, 1885, p. 96.
- *DARBYSHIRE, R. D. Anodon herculeus Gerstford?
Jour. of Conch., 1885, pp. 337, 338.
- *DAUTZENBERG, PHIL. Récoltes malacologiques de M. le Capitaine Em. Dorr, dans le Haut-Sénégal et le Soudan Français de 1886 à 1889.
Mém. Soc. Zool. de France, III, 1889, pp. 123-136, 1 plate.
- *— Mollusques Recueillis au Congo par M. E. Dupont, entre l'embouchure du fleuve et le confluent du Kassai.
Bull. Acad. Roy. Belg., XX, 1890, pp. 366-379, plates. Separate.
- *— Description d'une Mollusque nouveau provenant du Congo-Français.
Jour. de Conch., 1893, pp. 50, 51, 166, 167, pl., viii.
- DAVREUX, C. J., Constitution Géognostique de Liège, Bruxelles, 1833.
- DAWSON, G. M. Land and Fresh water Mollusca collected during the summers of 1873-1874 in the vicinity of the Forty-ninth parallel—Lake of the Woods to the Rocky Mountains.
Brit. North Am. Bdry. Com., Rep. on Geol., Montreal, 1875, pp. 347-350.
- DAWSON, J. W. Note on the genus Naïadites as occurring in the coal formation of Nova Scotia; with an appendix by W. Hind.
Quart. Jour. Geol. Soc., I, 1894, pp. 435-442, 1 plate.
- *DEAN, GEORGE W. Distribution of Unionidæ in the three rivers Mahoning, Cuyahoga, and Tuscarawas.
Nautilus, IV, 1890, pp. 20-22.
- *— On the distinguishing characters of Unio radiatus and U. luteolus.
Nautilus, V, 1891, pp. 77, 78.
- *— More about Unio luteolus and U. radiatus.
Nautilus, V, 1892, pp. 112, 113.

- *DEAN, GEORGE W. Catalogue of the Shell-bearing Mollusca of Portage County, Ohio.
Am. Naturalist, XXVI, 1892, pp. 11-23.
- DEBEAUX, O. Catalogue des Mollusques terrestres et fluviatiles du département de Lot-et-Garonne, 1844.
- *— Notice sur les Mollusques vivants observés dans le nord de la Chine par M. Debeaux, etc.
Rev. et Mag. de Zool., 1862, pp. 214-222.
- *DE BERGEN, CARLO AUGUSTO. Classes Conchyliorum auctore Caroli Augusto De Bergen, 1760.
- *DE CAMP, WILLIAM H. List of Shell-bearing Mollusca of Michigan.
Kent Sci. Inst., Grand Rapids, 1881, 13 pp., 1 plate.
- DE HEIDE. Descriptio Anatomica Mytili.
Act. Erud., Leipsic, 1684.
- DE KAY, J. E. Catalogue of the Animals belonging to the State of New York, etc.
Assembly Document, No. 50, 1839, pp. 12, 31-34.
- *— Natural History of New York, Pt. 5. Mollusca, 1843, pp. 271, 40 colored plates.
- *DELESSERT, BENJAMIN. Recueil des Coquilles decrites par Lamarck, dans son Histoire naturelle des Animaux sans Vertèbres et non encore figurées, 1841, 40 folios, colored plates.
- *DENNANT, J. Description of a new Unio from the River Glenelg.
Proc. Roy. Soc. Victoria, X, Pt. 2, 1898, pp. 112, 113, pl. iv.
- *DESCHAMPS, EMILE. Description d'une nouvelle espèce d'Unio.
Bull. Soc. Zool. de France, XVII, 1892, pp. 68, 69, fig.
- *DESHAYES, G. P. Sur l'Iridine, genre de Mollusques acéphales.
Nouv. Bull. Scienc. Soc. Phil., 1826, pp. 60, 61; *Fer. Bull. Sci. Nat.*, IX, 1826, pp. 248-250; XI, 1827, p. 447; *Isis*, 1832, p. 468.
- *— Mémoire anatomique sur l'Iridine du Nil.
Mem. Soc. Hist.-Nat. Paris, III, 1827, pp. 1-16; *Fer. Bull. Sci. Nat.*, XI, 1827, pp. 447-448; *Fror. Not.*, XIV, 1826, pp. 345, 343.
- *DESHAYES, G. P. Encyclopédie Méthodique, Histoire des Vers par Bruguière et Lamarck, complétée par Deshayes, 1830-1832, II, III.
— Tableaux comparatifs de Coquilles Vivantes avec les espèces fossiles, etc., 1830.
- *— Histoire naturelle des Animaux sans vertèbres de Lamarck, 2d ed., 1835.
- *— Mollusques.
Exp. Sci. de Morée, III, 1836, pp. 123, 9 plates.
— Article Conchifera.
Cyc. Anat. et Phys., I, 1836, pp. 694-716.
- *— Traite élémentaire de Conchylogie avec les Applications de cette Science à la Géologie, 1839-1857, 3 vols. text; 1 vol. plates.
- *— Expedition scientifique de l'Algérie, 1847.
- *— Description de quelques espèces de Mollusques nouveaux ou peu connus envoyés de la Chine, par L'Abbé David.
Bull. Nour. Arch. Mus., IX, 1873, pp. 1-14, 3 plates; Pt. 2, 1874, pp. 83-162, 4 plates.
- *— Observations sur les Animaux de deux Nuyades asiatiques.
Jour. de Conch., 1875, pp. 81-85.
- *DESHAYES, G. P., and JULLIEN, M. Mémoire sur les Mollusques nouveaux du Cambodge, envoyés au Museum par le Docteur Jullien.
Nouv. Arch. de Mus., X, 1874, pp. 115-162, 4 plates.
- DESHAYES, G. P., and MILNE-EDWARDS, H. Animaux sans Vertèbres, 3d ed., 1839.
- DESMARS, J. Essai d'un Catalogue des Mollusques terrestres, fluviatiles et marines observés dans l'Île et Vainne, etc., 1873, 94 pp.
- *DES MOULINS, CHARLES. Catalogue des espèces et variétés de Mollusques terrestres et fluviatiles * * * de la Gironde, etc.
Bull. Soc. Linn. Bord., II, 1827, pp. 39-69; Supp. to same, III, 1829, p. 211.
- *— Description d'une espèce d'Unio vivante du Périgord.
Actes Soc. Linn. Bord., VI, 1833, pp. 20-31, 1 plate.

- *DES MOULINS, CHARLES. Considerations générales sur la réforme qu'il conviendrait d'introduire dans la détermination des espèces des genres *Unio* et *Anodonta*.
Actes Soc. Linn. Bord., XI, 1839, 14 pp.; trans. in *Am. Jour. Sci. and Arts*, 1841, pp. 104-116.
 — Catalogue de Mollusques de la Dordogne, 1844.
- *— Mollusques terrestres et fluviatiles
 * * * de la Gironde.
Actes Soc. Linn. Bord., XVII, 1851, pp. 421-437; XVIII, pp. 492-499.
- *— DEVILLE, E., and HUPÉ, H. Description de quelques coquilles nouvelles provenant de l'expédition de M. de Castelnau.
Rev. et Mag. de Zool., 1850, pp. 631-644, 2 plates.
- DEWEY, C. List of Naiades found in western New York, and sent to the State collection at Albany, etc.
Ninth Ann. Rep. Regents Univ. N. Y., 1856, pp. 32-38.
 — Catalogue of the Fresh-water Shells furnished to the University of New York.
Ninth Ann. Rep. Regents Univ. N. Y., Appendix, 1856, p. 31.
- *DIEFFENBACH, ERNEST. Travels in New Zealand; with contributions to the Geography, Geology, Botany, and Natural History, etc., 1843, 2 vols.
- *DILLWYN, LEWIS WESTON. A Descriptive Catalogue of Recent Shells, etc., 1817, 2 vols.
- DIXON, JOHN. *Unio margaritiferns*.
Naturalist (London), I, 1865, pp. 81, 82.
- DOERING, A. Moluscos in: Informe oficial de la Expedición al Rio Negro (Patagonia), etc., Buenos Aires, 1881.
- *DOHRN, H. List of the Shells collected by Captain Speke during his second journey through Central Africa.
Proc. Zool. Soc., 1864, pp. 116-118.
- DOKIC, —. Mollusques lamellibranches de la Serbie, 1883.
- *DONOVAN, E. The Natural History of British Shells, etc., 1799-1804, 5 vols. 180 colored plates.
 Translated and reprinted by Chenu, in *Bib. Conch.*, 1845.
- *D'ORBIGNY, ALCIDE. Synopsis Terrestrialium et Fluviatilium Molluscorum in suo per Americannum Meridionalum itinero.
Mag. de Zool., 1835, p. 37.
- *— Mollusques de Cuba.
 In *Sagra, Hist. Nat. l'île de Cuba*, 2 vols., text, 1841-1853; atlas, 1842.
- *— Voyage dans l'Amérique Méridionale, 1835-1843. One vol. text; 1 of colored plates.
 — Paléontologie de l'Amérique Méridionale, 1842.
 — Considérations sur la station normale comparative des mollusques bivalves, 1843.
- *— Paléontologie Française. Paris, 1840-1861. 7 vols. text; 7 vols. plates.
- *— Mollusques vivants et fossiles, 10 vols. 1845.
- *— Prodrôme de Paléontologie, 3 vols. 1850-1852.
 — List of shells of South America in the British Museum, 1854.
- *DOUVILLE, H. Lamellibranches.
Annuaire Géol. Univ., VI, 1890, pp. 897-920.
- *DRAPARNAUD, J. P. R. Tableau des Mollusques Terrestres et Fluviatiles de la France, 1801, pp. 116.
- *— Histoire Naturelle des Mollusques Terrestres et Fluviatiles de la France, 1805.
- DROGEMÜLLER II. Die Fluss Perlenmuschel und die Wiederhebung der deutscher Perlen-fischerei.
Circ. deutsch. Fisch.-Ver., 1887, pp. 136-142.
- *DROUET, HENRI. Études sur les Anodontes de l'Aube.
Rev. et Mag. de Zool., pp. 51, 244, 285, 362, 527, etc., 1852-1854, 7 plates.
- *— Observations sur deux Anodontes.
Jour de Conch., 1856, pp. 123-129.
 — Études sur les Naiades de la France, pt. 2.
Mém. Acad. Aube, XXI, 1857. Separate.
- *— Essai sur les Mollusques Terrestres et Fluviatiles de la Guyane Française, 1859, pp. 110, 4 plates.
- *— Mollusques Terrestres et Fluviatiles de la Côte d'Or.
Mém. Acad. Dijon, 1866, pp. 33-154. Separate.

- *DROUET, HENRI. Unionidæ nouveaux ou peu connus.
Jour. de Conch., 1879, pp. 137-142, 327-333; 1881, pp. 22-31, 244-254; 1888, pp. 103-111; 1892, pp. 86-94; 1893, pp. 36-50, 167-178; 1895, pp. 26-40; 1897, pp. 124-136; 1899, pp. 402-411.
- *— Unionidæ de la Russie, 1881, pp. 35.
- Unionidæ de la Serbie, 1881, pp. 40. Reprinted in *Mém. Acad. Dijon*, 1882.
- Unionidæ de l'Italie, 1883, 125 pp.
- Unionidæ du bassin du Rhône. *Mém. Acad. Dijon*, 1889, pp. 27-114, 3 plates. Separate.
- Description de deux Unios nouveaux du bassin del Oronte. *Rev. Biol. Nord Fr.*, V, 1893, pp. 285-288, figs.
- *— Unionidæ de l'Espagne. *Mém. Acad. Dijon*, IV, 1894, pp. 143-226, 2 plates.
- *— DROUET, H., and CHAPER, M. Voyage de M. Chaper à Bornéo, Unionidæ.
Mém. Soc. Zool. de France, V, 1892, pp. 145-155, 2 plates; *Jour. de Conch.*, 1892, pp. 93, 94.
- *DUBOIS, CHARLES. Au Epitome of Lamarck's arrangement of Testacea, etc., 1825.
- DUMERIL, C., and DE BLAINVILLE, M. D. Rapport sur un mémoire de M. Jacobson: Observations sur le développement prétendu des œufs des mulettes, etc.
Ann. Sci. Nat. XIV, 1828, p. 22; *Isis*, 1830, pp. 217-222.
- DUMONT, FRANÇOIS, and MORTILLET, GABR. Histoire des Mollusques terrestres et d'eau douce vivants et fossiles de la Savoie, etc., 1852-1854, pp. 270.
- Catalogue critique et Malacostatique des Mollusques de Savoie, etc., 1857.
- *DUNKER, WILHELM. Diagnosis Molluscorum novorum.
Zeits. für Mal., III, 1846, pp. 108-112; V, 1848, pp. 177-186.
- *— Einige neue Species der Naiaden. *Mal. Blatt*, 1858, pp. 225-229.
- *DUPUY, L'ABBÉ. Essai sur les Mollusques terrestres et fluviatiles * * * du département du Gers, 1843, pp. 140, 1 plate.
- *— Histoire naturelle des Mollusques terrestres et d'Eau Douce qui vivent en France, 1847-1852, pp. 737, 31 plates.
- Catalogus Extramarinorum Gallie Testaceorum, etc., 1849, 2d ed., 1852.
- *— Note sur quelques Mollusques trouvés à Barbotan.
Jour. de Conch., 1877, pp. 15-23.
- Catalogue des Mollusques testacés terrestres et d'eau douce qui vivent à la Preste.
Bull. Soc. Hist. Nat. Toulouse, 1879. Separate.
- De la recherche des Mollusques terrestres et d'eau douce et des moyens de se les procurer.
Rev. Soc. Hist. Nat. Toulouse, 1878, 2d ed., 1381.
- D'URBAN, W. S. M. Catalogue of animals and plants of the valley of the River Rouge, etc.
Col. Surv. Can., Report of Progress for 1858 (1859), pp. 241-243.
- *DUVERNOY, M. Sur le système nerveux des Mollusques acéphales.
Comptes Rendus, 1845, pp. 482-484.
- EARLE, JOHN MILTON. List of Land and Fresh-water Shells in Massachusetts, so far as they have been ascertained.
Rep. on Geol. Bot. and Zool. of Mass., by E. Hitchcock, 1835, pp. 22-23.
- *EATON, AMOS. Zoological Text-Book, etc., 1826.
- *EDITOR. Notice of Index to Lea's Observations.
Am. Jour. Sci. and Arts, XIV, 1868, p. 129.
- *— One Year of Science.
N. Y. Tribune Extra, No. 2, 1874.
- Note on the circumboreal distribution of certain fresh-water mussels.
Am. Naturalist, XVII, 1883, pp. 203-204.
- *EICHWALD, EDUARD. Naturhistorische Skizze von Lithauen, Volhynien und Podolien, etc., 1830.
- Die Thier- und Pflanzen-Reste.
Bull. Acad. St. Petersb., 1840.
- *— Fauna Caspio-Caucasia, etc., 1841, pp. 264.

- EICHWALD, EDUARD. *Urwelt Russland.*
Bull. Acad. St. Petersb., 1840-1842.
- ENCYCLOPEDIA AMERICANA, Article Conchology, 1835.
- *ESMARK, BIRGITHE. On the Land and Fresh-water Mollusks of Norway.
Jour. of Conch., VI, 1886, pp. 90-100.
- *ESMARK, BIRGITHE, and HOYER, Z. A. Die Land-und Süßwassermollusken des arktischen Norwegens.
Mal. Blatt., 1886, pp. 84-123, 2 plates.
- *EYDOUX, FORT. Mollusques du voyage de la Favorite.
Mag. de Zool., 1838, p. 118, pl. CXIV-CXIX.
- FAGOT, PAUL. Catalogue des Mollusques des petites Pyrénées de la Haute-Garonne, etc.
Bull. Soc. Hist. Nat. Toulouse, XI, 1877. Separate.
- Matériaux pour la Faune malacologique terrestre, des eaux douces et des eaux saumâtres de l'Aude.
Bull. Soc. Hist. Nat. Toulouse, XIII, 1879. Separate.
- Histoire malacologique des Pyrénées françaises.
Bull. Soc. Hist. Nat. Toulouse, XIV, 1880. Separate.
- Mollusques terrestres et d'eau douce de la vallée d'Aulus.
Bull. Soc. Agr. Pyr. Or., XIV, 1880. Separate.
- FAGOT, PAUL, and DE NANSOUTY. Mollusques des Hautes-Pyrénées, etc., [without date].
- FAGOT, PAUL, and MALFOSSE. Catalogue des Mollusques terrestres et fluviatiles vivants dans le département de la Lozère.
Bull. Soc. Hist. Nat. Toulouse, XI, 1854. Separate.
- *FARINES, J. Description de trois espèces nouvelles de coquilles vivantes du département des Pyrénées-Orientales.
Ann. des Sci. Nat., 1834, pp. 118-123;
Jour. Inst., Paris, 1834, pp. 268, 269;
Bull. Soc. Philom., 1835, I, p. 29.
- FARWICK, B. Verzeichniss der Weichtiere Westfalens und Lippe-Deilmolds.
J. B. Westf. Prov., 1876, pp. 49-57.
- FAUSSEK, V. Ueber den Parasitismus den Anodonta-Larven in der Fischhaut.
Biol. Cent., XV, 1895, pp. 115-125, figs.; *Jour. Roy. Mic. Soc.*, 1895, pp. 521, 522.
- FAVANNE DE MONTCERVILLE. *Diet. Hist. Natur.*, 1775.
- *— La Conchyliologie, ou Histoire naturelle des coquilles.
Hist. Nat. D'Argenville, 1780. 2 vols. text; 1 vol. plates.
- *— Catalogue systématique et raisonné; description du magnifique cabinet, etc., 1784.
- FAVART-D'HERBIGNY, L'ABBÉ. Dictionnaire d'histoire naturelle qui concerne les coquillages de mer, etc., 1775.
- *FERUSSAC, ANDRÉ ÉTIENNE, BARON DE. Tableau systématique des animaux, mollusques, etc., 1819-1834.
- *— Concordance systématique pour les mollusques terrestres et fluviatiles de la Grande-Bretagne, etc.
Jour. de Phys., XC, 1820, pp. 212-217, 281-302. Separates.
- Histoire naturelle des mollusques terrestres et fluviatiles, etc.
Ann. Gén. Sci. Phys., 1820, pp. 319-348.
- Tableaux systématiques de l'embranchement des mollusques.
Isis, 1824; *Lett. Aus.*, 1824, pp. 1-52.
- *— Observations sur la génération des moules, etc.
Bull. Sci. Nat. et Géol., IX, 1826.
- *— Observations adressées en forme de lettre à MM. Th. Say, C. S. Rafinesque, Is. Lea, S. P. Hildreth, T. A. Conrad, et E. A. Poulson, etc.
Guer. Mag., 1835, pp. 1-36.
- *FERUSSAC, ANDRÉ ÉTIENNE, BARON DE. Catalogue de la collection de Coquilles formée par feu le Baron D'Audebard de Ferussac, 1837, pp. 23.
- Article Anodonta.
Dict. class. d'hist. nat., I, 1822, p. 397.
- *FERUSSAC (PÈRE), JEAN BAPTISTE LOUIS, BARON DE. Essai d'une méthode conchyliologique appliqué aux mollusques fluviatiles et terrestres, etc., 1807.
- Manuscript in Museum at Paris, unpublished.

- * FERUSSAC, JEAN BAPTISTE LOUIS DE and DESHAYES, G. P. Histoire naturelle générale et particulière des mollusques terrestres et fluviatiles, etc. Atlas of 217 plates, 2 vols. text; 1820-1851.
- FEUCHTWANGER, LEWIS. Popular Treatise on Gems, Pearls, etc., 1859.
- * FISCHER DE WALDHEIM, GOTTHELF. Bibliographia Palæontologica Animalium Systematica, 1834.
- FISCHER, K. Die Flussperlenmuschel im Regierungsbezirk.
Verh. Ver. Rheinl., XIV, 1888, pp. 292-294; Abstract in *Jour. Roy. Mic. Soc.*, 1889, p. 373.
- FISCHER, L. Naturgeschichte Livlands, 1791.
- * FISCHER, PAUL. Sur les Nayaides de la France, par H. Drouet, pt. 2.
Jour. de Conch., 1857, pp. 195-199.
- * — Observations sur les organes visuels des Nayaides.
Jour. de Conch., 1857, pp. 203-205.
- * — Mémoire sur les coquilles fossiles des terrains d'eau douce du sud-ouest de la France.
Jour. de Conch., 1868, pp. 286-289.
- * — Manuel de Conchyliologie et Paléontologie conchyliologique, etc., 1880-1887.
- * — Observations sur les genres *Mycetopus* et *Soleuaia*.
Jour. de Conch., 1890, pp. 1-14.
- * — Note sur la faune conchyliologique terrestre et fluviatile de l'île d'Hainan.
Jour. de Conch., 1890, pp. 96-100; 1892, pp. 313-315.
- * — Catalogue et distribution géographique des mollusques terrestres, fluviatiles et marines d'une partie de l'Indo-Chine.
Bull. Soc. Hist. N. Aut., IV, 1891, pp. 87-276. Separate.
- * FISCHER, PAUL, and CROSSE, H. Mission scientifique au Mexique et dans l'Amérique Centrale, Pt. 7, II, 1894.
- FITTON, —. Strata below the Chalk, 1836.
- * FITZGERALD, F. R. The Land and Fresh-water Mollusca of Harrowgate and District.
Jour. of Conch., VI, 1889, pp. 18-30.
- * FITZINGER, L. I. Systematisches Verzeichniss der im Erzherzogthum Oesterreich, etc., 1833.
- FLEISCHMANN, A. Die Bewegung des Fusses der Lamellibranchiaten.
Zeit. Wiss. Zool., XLII, 1885, pp. 367-431.
- * FLEMING, John. History of British Animals, 2 vols., 1828.
- * — Molluscons Animals, including Shell Fish.
Article Mollusks.
Encyclopædia Britannica, 7th ed., 1837, pp. 246. Separate.
- FLEMMING, W. Ueber die ersten Entwicklungserscheinungen am Ei der Teichmuschel.
Arch. f. Mukrosk. Anat., X, 1874, pp. 257-293; *J. B. Deutsch. Mal. Ges.*, 1874, pp. 270, 271.
- Studien in der Entwicklungsgeschichte der Najaden.
Stizb. Math. Nat. Akad. Wien, LXXI, 1875, pp. 132, 4 plates. Separate.
- Notiz zur Entwicklungsgeschichte der Najaden.
Zeits. Wiss. Zool., XXVI, 1876, pp. 355-360.
- * FOLIN, L. DE. Observations sur une espèce nouvelle d'Unio de France.
Le Naturaliste, X, 1888, pp. 273, 274, 4 figures.
- Observations sur *Unio moriscottei* n. sp.
Le Naturaliste, XI, 1889, p. 245, figure.
- Note sur l'*Anodonta piscinalis*, var. *vitula*.
Le Naturaliste, XI, 1889, p. 243.
- FOLIN, L. and BERILLON. Unios of Basses Pyrénées.
Bull. Soc. Bay., 1875, pl. —.
- * FONTANNES, F. Note sur la Découverte d'un *Unio* Plisse dans le Miocène du Portugal, 1883.
- FORBES, EDWARD. Malacologia Monensis, 1838.
- Estuary Beds and Oxford Clay.
Jour. Geol. Soc., VII, 1851, p. 104.
- * FORBES, EDWARD, and HANLEY, SYLVESTER. A History of British Mollusca and their Shells. 4 vols., colored plates, 1853.
- FOREL, F. A. Einige Beobachtung über Entwicklung des zelligen Muskelgewebes, 1886.
- Über die Entwicklung der Najaden.
Wurzburg Nat. Zeits., 1866, 1867.

- FOREL, F. A. Sur l'épithélium vibratile des branches des Nayaides.
Act. d. l. Soc. Helv. d. Sci. Nat. Neuch., 1866, pp. 94, 95.
- Recherches sur le développement des Naiades.
Bull. Soc. Vaud., 1866-1868, pp. 210, 211.
- Quelques observations sur l'accroissement de la Coquille des Anodontes.
Verh. Schw. Nat. Ges., 1868, pp. 90, 91.
- FORSKÅL, P. Voyage dans le Levant, 1775.
- *FOURNEL, D. H. L. Faune de la Moselle, etc. 2 vols., 1836.
- FOWLER, W. W. Large Specimen of Unio.
Science Gossip, XIII, 1877, p. 67.
- FRAUENFELD, GEORG RITTER VON. Fauna Dalmatica, 1856.
- Zoologische Miscellen, V.
Verh. Zool. bot. Ges. Wien, XV, 1865, pp. 525-536, 3 plates.
- FRIEDEL, E. Über die verwendung der Süßwasser muschel thiere als Schweinfutter in Nord-Deutschland.
Verh. Berl. Ges., 1873, pp. 22-23; *Zool. Gart.*, XIX, 1878, p. 276.
- *FRIELE, JOACHIM. Norske Land- og Ferskvands Mollusker som findes i Omegnen af Christiania og Bergen, 1853, pp. 65.
- *FRIERSON, LORRAINE S. Conchological Notes from Louisiana.
Nautilus, XI, 1897, pp. 3, 4.
- *— Unio (*Lampsilis*) *amphichænus*, n. sp.
Nautilus, XI, 1898, pp. 109, 110, pl.—
- *— How Uniones emigrate.
Nautilus, XII, 1899, pp. 139, 140.
- *— The Unionidae of De Soto Parish, Louisiana.
The Gulf Fauna and Flora, Bulletin, June, 1900, pp. 6-12.
- *FURTADO, A. Coquilles terrestres et fluviatiles de l'Exploration Africaine de MM. Capello et Ivens.
Jour. de Couch., 1886, pp. 138-152, 2 plates.
- *GABB, W. M. List of mollusks inhabiting the neighborhood of Philadelphia.
Proc. Acad. Nat. Sci. Phila., 1861, pp. 306-310.
- *GABB, W. M. Palæontology.
Geol. Survey of Cal., I, 1864, p. 182.
- GAERTNER, GOTTFRIED. Versuch einer Systematischen Beschreibung, etc.
Ann. Wetter., III, 1813, p. 281.
- GALLENSTEIN, T. VON. Studien ans der Najaden fauna des Isonvogegebietes, 1894, 49 pp.
- GALTON, JOHN C. The Anatomy of the River Mussel.
Pop. Sci. Rev., IX, 1870, pp. 240-256, 1 plate.
- GARMAN, H. A preliminary report on the animals of the waters of the Mississippi Bottoms, etc., 1889.
- GARNER, ROBERT. On the Nervous System of Mollusks.
Trans. Linn. Soc. Lond., XVII, 1834, p. 485.
- *— On the Anatomy of Lamellibranchiate Conchiferous Animals.
Proc. Zool. Soc., 1836, pp. 12-16; *Isis*, 1838, p. 820; *Trans. Zool. Soc. Lond.*, II, 1841, p. 87, 2 plates.
- Notes on anatomy and pearls of *Alasmodon margaritifera*.
Proc. Zool. Soc., XI, 1872, pp. 426-428.
- *GASKOIN—[Account of a Unio from Australia, etc.]
Proc. Zool. Soc., 1850, pp. 243, 244; *Ann. Nat. Hist.*, 1852, p. 498.
- *GASSIES, J. B. Tableau méthodique et descriptif des Mollusques Terrestres et d'Eau douce de l'Agenais, 1849, pp. 209, 4 colored plates.
- Du progrès de la malacologie en France, etc., 1858, 17 pp.
- *— Catalogue raisonné des Mollusques terrestres et d'eau douce de la Gironde, 1859, pp. 74.
- Catalogue des Mollusques terrestres et d'eau douce du département du Lot-et-Garonne.
Rev. Trar. Soc. Ag. Sc. et Arts. d'Agen, 2d ser., I, 1861, separate; Supp. in *Bull. Soc. Eorda*, III, 1878, p. 33.
- *— Malacologie terrestre et d'eau douce de la région intra-littorale de l'Aquitaine.
Actes Soc. Linn. Bord., XXVI, 1866, pp. 109-136, plate, separate.
- *GAY, CLAUDIO. Historia física y política de Chile segun documentos, etc., VIII, Mollusks by Hupé, 1854.
- *GEGENBAUR, CARL. Elements of comparative anatomy, 1878.

- *GEINITZ, HANS BRUNO. Grundriss der Verst., etc., 1846.
- GENTILUOMO, C. *Uniolaidleyanus* n. sp.
Bull. Soc. Mal. It., II, 1868, pp. 54-58.
- * — Catalogo dei Mollusche terrestri e fluviatili della Toscana.
Rivista Bib. It., 1868, pp. 67-99;
Bull. Soc. Mal. It., I, 1868, pp. 1-100, plate.
- GEOFFREY, E. L. *Traité sommaire des coquilles tant fluviatiles que terrestres aux environs de Paris*, 1767.
- *Recueil de Coquilles fluviatiles et terrestres qui se trouvent aux environs de Paris*, etc., 1776, 3 plates.
- GERMAIN, LOUIS. *Sur deux unions sous le nom rotundatus*.
Bull. Soc. Amis N. Rouen, 1898, pp. 70-73.
- *GERSTFELDT, G. Ueber Land- und Siisswasser-Mollusken Sibiriens und des Amur-Gebietes.
Mem. Sav. Etr., IX, 1850, pp. 508-535, separate, 1859, pp. 44, 1 plate.
- GERVAIS, PAUL. *Liste des Mollusques terrestres et fluviatiles de la France*, 1847.
- GESNER, CONRAD. *Nomenclator Aquatibum*, 1560.
- GEVE, NICHOLAS. *Monatliche Belustigungen im Reiche*, etc., 1755, 24 colored plates.
- GIBBES, LEWIS R. *Molluscons animals of South Carolina*.
Geol. South Carolina, 1848, pp. XIX-XXII. Appendix.
- GINANNI, GIUS. *Opere postume del Conte Giuseppe Ginanni, Testacei Marittima paludosa e terrestre dell Adriatico*, etc., 1755.
- GIRARD, CHARLES. *Catalogue of Recent Shells and other Mollusca found in the District of Columbia*, etc.
Proc. Nat. Inst., I, 1856, pp. 79-82.
- *GISTEL, JOHANNES. *Natürgeschichte des Thierreichs*. 1 vol., 236 pp; atlas, 32 plates, 1848.
- *GMELIN, JEAN FREDERIC. *Caroli a Linnæus, Systema naturæ per regna*, etc., 13th ed., 6 vols., 1788, 1792.
- GODET, PAUL. *Note sur les Anodontes du lac de Neuchâtel*.
Bull. Soc. Nat. Neuch., VI, 1864, pp. 71-75.
- GODET, PAUL. *Les Anodontes du canton de Neuchâtel*.
Bull. Soc. Nat. Neuch., IX, 1873, pp. 145-151.
- GOETTE, A. *Bemerkungen über die Embryonalentwicklung der Anodonta piscinalis*.
Zeit. Wiss. Zool., II, 1891, pp. 158-168.
- GOLDFUSS, O. *Petrificationen Deutschlands*, II, 1837.
- *Beiträge zur Mollusken-Fauna der Mansfelder Seen*, etc.
Nach. Deuts. Mal. Ges., 1894, pp. 43-64.
- *GOODRICH, S. C. *Illustrated Natural History*, etc., II, 1859.
- GOULD, AUG. A. *Lamarck's Genera of Shells, with a Catalogue of Species*. Translated. 1833.
- *Catalogue of Maine Shells*.
First Rep. Geol. Maine, 1837.
- *Results of an Examination of the Shells of Massachusetts*.
Jour. Bost. Soc. Nat. Hist., 1841, pp. 488, 489.
- * — *Shells of Massachusetts and their Geographical Distribution*.
Jour. Bost. Soc. Nat. Hist., III, 1841, pp. 483-494.
- * — *Report on the Invertebrata of Massachusetts*, 1841, pp. 373, woodcuts and plates. 2d ed., by W. G. Binney, 1870.
- * — *Descriptions of Shells from Tavoy, Burmah*.
Proc. Bost. Soc. Nat. Hist., I, 1843, pp. 139-141.
- * — *Descriptions of two species of Anodon from the River Salween, etc*.
Proc. Bost. Soc. Nat. Hist., I, 1841, p. 160.
- * — *Descriptions of New Shells*.
Proc. Bost. Soc. Nat. Hist., I, 1844, p. 139.
- * — *Descriptions of Recent Shells collected by Mr. John Bartlett in the Everglades of Florida*.
Proc. Bost. Soc. Nat. Hist., II, 1845, p. 53.
- *Expedition Shells, described for the work of the United States Exploring Expedition*.
Proc. Bost. Soc. Nat. Hist., II, 1846.
- * — [Description of Shells from Tavoy, British Burmah.]
Proc. Bost. Soc. Nat. Hist., II, 1847, pp. 218-221.

- * GOULD, AUG. A. Descriptions of new shells.
Proc. Bost. Soc. Nat. Hist., II, 1848, p. 218.
- *— Descriptions of shells found in Connecticut, named by Rev. J. H. Linsley.
Am. Jour. Sci. and Arts, VI, 1848, pp. 233-236.
- *— Catalogue of shells, with descriptions of new species.
Lake Superior, etc., by Louis Agassiz, 1850, pp. 243-245.
- *— Descriptions of new species of shells.
Proc. Bost. Soc. Nat. Hist., III, 1851, pp. 292-296.
- Descriptions of new species of shells.
Proc. Bost. Soc. Nat. Hist., IV, 1851, pp. 87-92.
- *— Mollusca and Shells of the United States Exploring Expedition, one vol. and atlas, 1852, pp. 510.
- *— Descriptions of shells from the Gulf of California and the Pacific coasts of Mexico and California.
Jour. Bost. Soc. Nat. Hist., VI, 1853, pp. 374-407.
- A System of Natural History, 1853.
- *— [Descriptions of shells.]
Proc. Bost. Soc. Nat. Hist., V, 1856, pp. 228-229; VI, 1856, pp. 11-16.
- *— [On the geographical distribution of Unios.]
Proc. Bost. Soc. Nat. Hist., VII, 1859, p. 167.
- *— *Otia Conchologica*, 1862.
- * GOULD, A. A., and CARPENTER, PHILIP. Descriptions of shells from the Gulf of California and the Pacific coasts of Mexico and California, Pt. II.
Proc. Zool. Soc., 1856, pp. 198-229.
- * GOUPII, C. J. Histoire des Mollusques terrestres et fluviatiles observés dans le Département de la Sarthe, etc., 1835.
- GRÆLLS, M. P. Catálogo de los Moluscos terrestres y de agua dulce observados en España, etc., 1846, 23 pp., 1 plate.
- * GRANDIDIER, ALFRED. Mollusques de l'Ousaghara, de l'Oukami, etc.
Bull. Soc. Mal. Fr., IV, 1887, pp. 185-194.
- GRANT, U. S. Notes on the molluscan fauna of Minnesota.
Geol. and Nat. Hist. Surv. Minn., XVI, 1888, pp. 481-484.
- * GRAS, ALBIN. Description des Mollusques fluviatiles et terrestres du Département de l'Isère, 1840; appendix, 1840, pp. 109.
- *— Description des Mollusques fluviatiles et terrestres de la France, 1846.
- GRATELOUP, J. P. S. DE. Tableau des Coquilles fossiles.
Act. Soc. Linn. Bordeaux, V, 1829.
- Tableau méthodique des Mollusques terrestres et fluviatiles de Dax.
Act. Soc. Linn. Bordeaux, 1829, pp. 43, 87, 143.
- Essai sur la Nourriture des Mollusques, 1857.
- Essai sur la Distribution géographique de la Gironde, 1858.
- * GRATELOUP, J. P. S. DE, and RAULIN, VICTOR. Catalogue des Mollusques terrestres et fluviatiles vivants et fossiles dans les différentes régions et contrées de la France, etc., 1855.
- * GRAY, A. F. Description of *Unio borealis*.
Trans. Ottawa Field Nat. Club, III, 1882, p. 53, plate.
- *— On the eastern range of *Unio pressus*.
Am. Naturalist, XVII, 1883, pp. 204, 205.
- *— Remarks on the distribution of *Margaritana margaritifera*.
Am. Naturalist, XVII, 1883, pp. 324-326.
- GRAY, JOHN EDWARD. New British species of Mollusca.
Lond. Med. Rep., XV, 1821, p. 239.
- A list and description of some species of shells not taken notice of by Lamarck, 1825.
Griffith's Cuvier, XII, 1834.
- *— *Spicilegia Zoologica*, 1830.
- *— Some observations on the economy of molluscous animals, and on the structure of their shells.
Phil. Trans., 1833, pp. 771-819.
- *— [Characters of new species of shells.]
Proc. Zool. Soc., 1834, pp. 57-72.

- GRAY, JOHN EDWARD. Review of Lea's Observations on the genus *Unio*.
Mag. de Zool., I, 1837, pp. 284, 285.
- Synoptical Catalogue of the Shells in the British Museum, 1837.
- *— The animal of Hyria.
Ann. Nat. Hist., VI, 1840, p. 316.
- A manual of the land and fresh-water shells of the British Islands by W. Turton, new ed., 1840.
- *— On the Byssus of *Unio* by John G. Anthony, with notes.
Ann. Nat. Hist., VI, 1841, p. 77.
- *— In Dieffenbach's Travels in New Zealand, 1843.
- *— A list of the genera of recent Mollusca, etc.
Proc. Zool. Soc., XV, 1847, pp. 129-214.
- *— A revision of the arrangement of the families of bivalve shells.
Ann. Nat. Hist., XIII, 1854, pp. 408-418.
- Guide to Mollusca, 1857.
- On a new genus of Mytilidæ, and on some distorted forms which occur among bivalve shells.
Proc. Zool. Soc., 1858, pp. 90-92.
- *GRAY, MARIA EMMA. Figures of molluscos animals, 5 vols., 1842-1859.
- *GREDLER, VINCENT. Zur Conchylien Fauna von China.
Jahrb. Deutsch. Mal. Ges., VIII, 1881, pp. 10-35, 110-232, plate; *Nach. Deutsch. Mal. Ges.*, XV, 1889, pp. 168-178; XVI, 1890, pp. 33-41, 145-153.
- GREEN, JACOB. Some remarks on the Unios of the United States, with a description of a new species.
Contributions to Maclurian Lyceum, I, 1827, pp. 41-47.
- Notes on the American shells figured in the Supplement to the Index Testaceologicus.
Trans. Albany Inst., I, 1830, pp. 134-136.
- Notes of a naturalist.
Cab. Nat. Hist., II, 1832, pp. 290, 291.
- GREENE, C. Catalogue of the land and fresh-water shells in the county of Suffolk.
Suff. Inst. Arch. and N. H., VII, 1891, Pt. 3.
- *GREISBACH, HERM. ADOLPH. Über den Bau des Bojanussehen Organes der Teich Muschel.
Arch. für Naturg., 1877, pp. 63-107.
- Über das Gefäß-system und die Wasseraufnahme bei den Najaden und Mytiliden.
Biol. Cent., II, 1882, pp. 305-309; 573-577.
- *GRIFFITH, EDWARD, and PIDGEON, EDWARD. The Mollusca and Radiata, arranged by the Baron Cuvier, with supplementary additions, etc.
Cuvier's Animal Kingdom, 1834, pp. 434, 40 plates. Text by J. E. Gray.
- GRIFFITHS, A. B. Researches on the problematical organs of the Invertebrata, etc.
Proc. Roy. Soc. Edinb., XIV, 1888, pp. 230-238, figures.
- GRIFFITHS, A. B., and FELLOWS, H. Chémico-biological examination of the organs of Bojanus in Anodonta.
Chem. News, LI, 1885, p. 241.
- GROGNOT, —. Mollusques testacés fluviatiles et terrestres du Département de Saône-et-Loire, 1869, pp. 23.
- *GRONOVIVS, LAURENT THEODORE. Zoophylacium Gronovianum, etc., 1781.
- *GRUNER, E. CH. L. Einige neue Land und Süsswasser Conchylien.
Arch. für Naturg., 1841, pp. 276-278.
- *GUALTIERI, NICOLAI. Index Testarum Conchyliorum, etc., 1742, 2 vols.
- *GUÉRIN-MÉNEVILLE, F. E. Iconographie du Règne Animal de G. Cuvier. 3 vols., 1844, colored plates.
- *GUÉRIN-MÉNEVILLE, F. E., and DEBEAUX, O. Notice sur les Mollusques vivants observés dans le nord de la Chine, etc.
Rev. et Mag. de Zool., 1862, pp. 214-222.
- *GUNTHER, ALBERT. Zoological literature.
Zool. Record, I., 1864, pp. 201, 244; II, 1865, p. 293; III, 1866, 1867.
- *— Remarks on Morelet's work on African land and fresh-water mollusks.
Zool. Record, 1868, p. 483.
- *GUPPY, R. LECHIMERE. On the terrestrial and fluviatile Mollusca of Trinidad.
Ann. Nat. Hist., XVII, 1866, pp. 42-56.

- *GUPPY, R. LECHMERE. The land and freshwater mollusks of Trinidad.
Jour. of Conch., VII, 1894, pp. 210-231.
- *GYSSER, AUGUST. Vergleichende Zusammenstellung der Molluskenfauna der beiden äussersten nordöstlichen und südwestlichen Grenzländer, etc.
Mal. Blatt., XII, 1865, pp. 78-91.
- *H., ———. Notice of Lea's Observations, etc.
Am. Jour. Sci. and Arts, XXXVI, 1863, p. 139.
- HAGENMULLER, ———. Catalogue des Mollusques terrestres et fluviatiles d'Alsace.
Bull. Soc. Colmar, 1871-1872, p. 135. Separate.
- HALAVÁT, G. A. Szegedi Kút artezi Kút Magyar Föld int. Evkőn, IX, 1891, pp. 79-97.
- *HALDEMAN, S. S. Remarks on Unios.
Proc. Acad. Nat. Sci. Phila., I, 1841, p. 104.
- *—— [Note on colors of *U. tappanianus*.]
Proc. Acad. Nat. Sci. Phila., I, 1841, p. 104.
- *—— Description of five new species of North American fresh-water shells.
Jour. Acad. Nat. Sci. Phila., VIII, 1842, pp. 200-202.
- *—— [Communication correcting nomenclature.]
Proc. Acad. Nat. Sci. Phila., 1842, pp. 176, 177.
- Mollusca.
Rupp's History of Lancaster Co[unty], 1844, p. 479.
- *—— Enumeration of the recent fresh-water Mollusca common to North America and Europe, etc.
Jour. Bost. Soc. Nat. Hist., IV, 1844, pp. 468-484.
- *—— Description of *Unio abacoides*, a new species.
Proc. Acad. Nat. Sci. Phila., III, 1846, p. 75; *Ann. Nat. Hist.*, XVIII, 1846, p. 430; *Am. Jour. Sci. and Arts*, II, 1846, p. 274.
- *—— [Western Uniones in the Susquehanna.]
Proc. Acad. Nat. Sci. Phila., III, 1846, p. 15.
- HALDEMAN, S. S. The Iconographic Encyclopedia of Science, Literature, and Art, etc., additions to translation, II, 1857, pp. 49-90.
- *—— On Nomenclature.
Am. Naturalist, III, 1870, p. 159.
- *HAMONVILLE, Baron d'. Les Monts perlières de Billiers.
Bull. Soc. Zool. de France, XIX, 1894, pp. 140-142.
- HANHAM, A. W. List of Land and Fresh-water Shells of the Hamilton District, etc.
Jour. Ham. Assn., V, 1889, pp. 80-91.
- *—— A List of the Land and Fresh-water Shells of Manitoba.
Nautilus, XIII, 1899, pp. 1-6.
- HANITSCH, R. Die Wasseranfnahme bei Cyclos und Anodonta.
Inaugural Dissertation, 1884, 38 pp.
- *HANLEY, SYLVANUS. The Young Conchologist's Book of Species, 1840.
- *—— Testaceous Mollusca, Lamarck's Species of Shells, 1842, pp. 224, 3 colored plates.
- *—— An Illustrated and Descriptive Catalogue of Recent Bivalve Shells, Appendix to Index Testaceologicus, 1842-1856.
- History of British Mollusca, 1853.
- Conchological Miscellany, 1854; 2d ser., 1864.
- *—— *Ipsa Linnaei Conchyli*, 1855.
- *—— Description of a new *Monocondylaea*.
Proc. Zool. Soc., 1871, pp. 587-588.
- Description of new Land and Fresh-water Shells from India.
Proc. Zool. Soc., 1875, pp. 605-607.
- *HANLEY, SYLVANUS, and THEOBALD, WILLIAM. Conchologia Indica, etc., 1870.
- HARDY, JOHN. On the Occurrence of *Unio tumidus* in the Manchester District.
Proc. Lit. Soc. Manch., XII, 1873, p. 117.
- HARTING, J. E. Rambles in Search of Shells, 1875.
- Land and Fresh-water Mollusks of Sussex.
Zoologist, II, 1878, pp. 84-94, 122-126, 161-166, 180-181.
- HARTMAN, WILLIAM D. A Classified Catalogue of the Terrestrial and Fluviatile Mollusks of Chester County, Pennsylvania, 1853.

- *HARTMAN, WILLIAM D. Illustrated Catalogue of the Land and Fresh-water Shells of Chester County, Pennsylvania.
Village Record, 1895, p. 7, figs.
- *HARTMAN, WILLIAM D., and MICHENER, EZRA. *Conchologia Cestrica*, etc., 1874, 114 pp., figs.
- HARTMANN, J. D. W. *System der Erd und Süßwasser Gasteropoden Europas*, etc., 1821.
- HARTMANN, J. D. W. *System der Erd und Flüß Mollusken der Schweiz*, etc.
Steinmüller's Neue Alpina, 1821; *Fer. Bull.*, 1825, p. 271.
- HARTOG, MARCUS M. On the Organ of Bojanus in Anodon.
Jour. Anat. Phys., XIII, 1879, p. 400, 578.
- *HAYDEN, F. V. Notes on the Geology of Wyoming and Colorado Territories.
Proc. Am. Phil. Soc., XI, 1869, pp. 25-56.
- Report on the Geology of Colorado and New Mexico, 1869.
- *— Report of the United States Geological Survey of the Territories, IX, 1876, pp. 112-115.
- Geology of Tertiary Formations of Dakota and Nebraska.
Jour. Acad. Nat. Sci. Phila., VII, 1869.
- *HAZAY, JULIUS. Die Molluskenfauna von Budapest.
Mal. Blatt., III, 1881, pp. 1-69, 160-182; IV, pp. 43-224, plates.
- *— Die "Nouvelle Ecole," beleuchtet durch Dr. Georg Servain's *Histoire malacologique du lac Balatan*.
Mal. Blatt., VI, 1883, pp. 179-197.
- HECART. *Catalogue des Coquilles terrestres et fluviatiles des environs de Valenciennes*, 1833.
- *HEDLEY, C., and MUSSON, C. T. On a Collection of Land and Fresh-water Shells from Queensland.
Proc. Linn. Soc. N. S. W., VI, 1891, pp. 551-564, figs.
- *HEDLEY, C., and SUTER, H. Reference List of the Land and Fresh-water Mollusca of New Zealand.
Proc. Linn. Soc. N. S. W., VII, 1892, pp. 613-665.
- *HEIMBURG, H. VON. *Diagnosen neuer Arten*.
Nach. Deutsch. Mal. Ges., XVI, 1884, pp. 92-95.
- *— *Abbildung und Beschreibung neuer Arten*.
Jahrb. Deutsch. Mal. Ges., XIV, 1887, pp. 1-4, plate.
- *HELD, FREDERICH. *Aufzählung der in Bayern Mollusken*.
Isis, VI, 1836, pp. 271-282.
- HERBIGNY, FAVART D'. *Dictionnaire d'Histoire naturelle qui concerne les Testacées*, etc., II, 1775.
- HERBST. *Einleitung*, 1787.
- *HERMANNSEN, A. N. *Indices generum Malacozoorum*, etc. 2 vols., 1846-1847; Supplement, 1852.
- *HESSLING, THEODOR VON. *Die Perlenmuscheln und ihre Perlen*, 1859, 8 plates.
- *Ueber die Befruchtung der Flussperlenmuschel*, etc.
Froriep's Notizen, 1860, pp. 161-168.
- *HEUDE, R. P. *Diagnoses Molluscorum fluminibus provincia Nankingensis*, etc.
Jour. de Conch., 1874, pp. 112-118.
- *— *Conchyliologie fluviatile de Nanking*, 1875-1886.
- *— *Rectifications de nomenclature*.
Jour. de Conch., 1884, p. 20.
- *HEY, W. C. *Fresh-water mussels in the Ouse and Foss*.
Jour. of Conch., III, 1882, pp. 268-273.
- HIBBERT, SAMUEL. *On the Fresh-water Limestone of Burdie House*, etc.
Trans. Roy. Soc. Edinburgh, XIII, 1836, pp. 169-282.
- *HIDALGO, J. GONZALEZ. *Diagnoses Castaliae novæ*.
Jour. de Conch., 1865, p. 316, 429-430, figs.
- *— *Obras malacologicas*, pt. 3.
Mem. Ac. Madrid, XVI, 1893, pp. 33-432.
- HIGGINS, FRANK. *A Catalogue of the Shell-bearing species of Mollusca, of Columbus, Ohio*, etc., 1858.
- HIGGINS, H. H. [Local Fresh-water and Land Mollusks around Liverpool.]
Proc. Liv. Field Club, 1890, pp. 11-33.

- * HIGGINS, THOMAS. Description of Six New Species of Shells.
Proc. Zool. Soc., 1868, pp. 178-180, plate.
- * HILDRETH, S. P. Observations on and Descriptions of the Shells found in the Muskingum River, etc.
Am. Jour. Sci. and Arts, XIV, 1828, pp. 276-291, 2 plates.
- * — Habits of Uniones.
Am. Jour. Sci. and Arts, XXXI, 1836, p. 62.
- * — Disease among Shellfish.
Am. Jour. Sci. and Arts, XXXII, 1837, p. 97.
- Catalogue of Fresh-water Shells found near Marietta, Ohio. No date.
- HINCKS, WILLIAM. Molluscous Animals.
Canadian Jour. Ind. Sci. and Art, 1867, p. 392; extract in *Am. Jour. Conch.*, 1868, pp. 161, 162.
- HIND, WHEELTON. On the Affinities of Anthracoptera and Anthracomya.
Quart. Jour. Geol. Soc., XLIX, 1893, pp. 249-275, plates.
- A Monograph on Carbonicola, Anthracomya Naiadites, etc.
Pal. Soc. Mon., Pt. 1, 1894, 80 pp. 11 plates; Pt. 2, 1895, pp. 81-170, 8 plates.
- * HINKLEY, A. A. New locality for *Unio ellipsis*.
Nautilus, VIII, 1894, p. 82.
- HISLOP. Tertiary Deposits in the East Indies.
Jour. Geol. Soc., XVI, 1860, p. 174.
- HODGE, JAMES THATCHER. Article Conchology.
Appleton's New American Cyclopaedia, V., 1859.
- HOLANDRE. Faune du département de la Moselle, etc., 1836.
- HOLZINGER, J. M. Notes on the Mollusca of Wenona County.
Geol. and Nat. His. Surv. Minn., XVI, 1888, pp. 485-491.
- HOME, EVERARD. The digestive organs of Fresh-water Mussels, 1814.
- On the production and formation of pearls.
Phil. Trans., III, 1826, p. 338; *Isis*, 1836, p. 580.
- Ueber die Fortpflanzung der Anster Flussmuschel.
Hensingen's Zeits. Phys., 1827, pp. 391-396.
- * HOME, EVERARD. On the Propagation of Oysters and the Large Fresh-water Mussel.
Phil. Trans., CXVII, 1827, pp. 39-48, 4 plates. *Ferussac's Bull. Sci. Nat.*, XIII, 1828, pp. 252-254.
- HÖRNES. Die Fossilien Mollusken, etc.
Geol. Jahrb., 1851-1852.
- * HOUGHTON, W. On the Parasitic Nature of the Fry of *Anodonta cygnea*.
Quart. Jour. Roy. Mic. Soc., 1862, pp. 162-168.
- * — The Swan Mussel and its Anatomy.
Intellectual Observer, VI, 1864, pp. 67-75, 1 plate.
- HUBBARD, EBER W. Catalogue of Shells of Ohio. No date.
- HUEBER, LEOPOLD V. Zur Naturgeschichte der Unionen.
Jahrb. Deutsch. Natur. Landes-Kart., 1871, pp. 151-157.
- * HUMBOLDT, ALEX. DE, and BONPLAND, A. Recueil d'observations de zoologie et d'anatomie comparée, etc., II, 1833.
- HUMPHREY, GEORGE. Museum Caloniannum, 1797. From a manuscript by Hovass.
- HUNT, WILLIAM. The Cabinet of Nature and Philosophy. Book 1, 1833, 4 plates.
- * HUPÉ, H. In Gay's *Historia física y política de Chile*, VIII, 1854.
- * — Mollusques, in *Animaux nouveaux ou Rares*, 1857, pp. 103, 20 plates.
- HUTTON, FREDERICK WALLASTON. Catalogue of the Tertiary Mollusca of New Zealand in the collection of the Colonial Museum, 1873, pp. 48.
- * — Manual of the New Zealand Mollusca, 1880.
- * — On the Fresh-water Lamellibranchs of New Zealand.
Trans. New Zeal. Inst., XIV, 1881, pp. 148-150, plate.
- * — List of the Fresh-water Shells of New Zealand.
Proc. Linn. Soc. N. S. W., VII, 1882.
- * — Notes on some Marine Mollusca, with descriptions of new species.
Trans. New Zeal. Inst., XVI, 1884, pp. 212-216.
- * — The Mollusca of the Parcora and Oamaru systems of New Zealand.
Proc. Linn. Soc. N. S. W., 1886, pp. 205-237.

- * IHERING, HERM. von. Über die Entwicklungsgeschichte der Najaden.
Sitzb. Nat. Ges. Leip., 1874, pp. 3-8;
Jahrb. Deutsch. Mal. Ges., 1874, p. 272.
- * — Revision der von Spix in Brasilien gesammelten Najaden.
Arch. für Naturg., 1890, pp. 117-170, plate.
- * — Anodonta und Glabaris.
Zool. Anzeiger., 1891-92, pp. 1-5. Separate.
- * — Zur Kenntniss der Gattung Cris-taria.
Nach. Deutsch. Mal. Ges., 1892, pp. 1-14.
- * — Najaden von S. Paulo und die geo-graphische Verbreitung der Süß-wasser-Faunen von Süd-Amerika.
Arch. für Naturg., 1893, pp. 45-140, 2 plates, separate; abstract in *Jour. Roy. Mic. Soc.*, 1893, p. 165.
- * — Die Süßwasserbivalven Japans.
Abh. Senck. Ges., XVIII, 1894, pp. 145-166, plate. Separate.
- * — Os Unionidos da Florida.
Revista Mus. Paul., I, 1895, pp. 207-222.
- INGERSOLL, E. [Colorado Mollusks.]
Bull. U. S. Geol. Surv., 1875, pp. 128-136.
- * — On a collection of Mollusks from Utah and Colorado.
Proc. Dar. Acad. Sci., 1876-77, pp. 130-134.
- Investigator. Remarks, etc.
Monthly Am. Jour. Geol. and Nat. Sci., 1832, pp. 537-549.
- ISHIKAWA, C. [Introduction to the anatomy of animals.] Tokyo, 1892, pp. 35, 6 plates.
- ISSEL, A. Dei Molluschi raccolti nella provincia di Pisa.
Mem. Soc. It. S. Nat., II, 1866.
- Appendice al Catalogo Molluschi di Pisa.
Atti Soc. It., XV, 1872, pp. 21.
- * — Molluschi Borneensi, etc.
Ann. Mus. Genov., VI, 1874, pp. 366-478, 3 plates. Separate.
- * — Delle Linei ornate di fascie e delle Anodonta, etc.
Bull. Soc. Mal. It., II, 1876, pp. 50-52.
- ISSEL, A. Molluschi terrestri e d'acqua dolce Tunisia, etc.
Ann. Mus. Genov., XV, 1880, pp. 259-282, figs.
- * — Materiali per lo studio della Fauna Tunisina, etc.
Ann. Mus. Genov., II, 1885, pp. 5-15, figures.
- IWAKAWA, T. [Freshwater Mollusca of Japan.]
Zool. Mag. Tokyo, VII, 1895.
- * JACKSON, C. T. [Shells found at Lake Superior.]
Proc. Bost. Soc. Nat. Hist., I, 1844, p. 198.
- Catalogue of Shells of the Alle-ghanies.
The Mountain, 1860, p. 154.
- * JACKSON, R. T. Phylogeny of the Pelecypoda, etc.
Mem. Bost. Soc. Nat. Hist., IV, 1890, pp. 277-400, 8 plates, figures.
- JACOBSON, L. L. [On Sex in Unio.]
Bull. des Sci. Nat., 1827.
- Unders ögelsar til noermere Oplysning, etc.
K. Dansk. Selsk. Nat. Afh., 1828, pp. 251-297; *Bid. Au. Phys. Cop.*, III, 1828, p. 281.
- Anatomy og Physiologie.
K. Dansk. Selsk. Nat., 1828, pp. 249-362; *Mem. Ac. Cop.*, 1828, p. 249; *Jour. Phys.*, 1820, p. 318.
- JACQUEMINE, ÉMILE. Guide du Voya-geur à Arles, 1835.
- JAN. In Charpentier's Catalogue des Mollusques de la Suisse, 1837.
- * JAPP, WILLIAM. Stray notes on the Pearl Mussel.
Scottish Nat., VIII, 1886, pp. 62-64; II, new ser., 1886, pp. 113-115.
- * JAY, JOHN C. A Catalogue of Recent Shells, etc., 1835, 2d ed., 1836; 3d ed., 1839.
- Report on the Shells collected by the Japan Expedition, etc.
Rep. of Perry Exp., II, 1856, p. 291.
- JEFFREYS, C. A Catalogue of British Land and Freshwater Mollusca, 1887.
- * JEFFREYS, JOHN GWYN. British Conchology, 5 vols., 1862-69.
- * JELSKI, C. Sur la Faune Malacologique de Kieff.
Jour. de Conch., 1863, pp. 129-137.

- JICKELI, C. F. Diagnosen neuer Mollusken, etc.
Mal. Blatt, XX, 1873, pp. 99-108.
- Abnorme Schlossbildung bei *Spaetha hartmanni*.
Nach. Deutsch. Mal. Ges., 1873, pp. 69, 70.
- * — Reisebericht.
Mal. Blatt., XX, 1873, pp. 1-20, 109-151; XXI, pp. 109-151.
- * — Fauna der Land- und Süßwasser-Mollusken, Nord-Ost-Afrikas.
Verh. Kais.-Leop. Ak. Dr., XXXVII, 1874, pp. 1-350, 11 plates. Separate.
- * — Land- und Süßwasserconchylien Nord-Ost-Afrikas.
Jahrb. Deutsch. Mal. Ges., VIII, 1881, pp. 336-340.
- JOANNIS, L. DE. Étude sur les Náyades du département de Maine-et-Loire.
Ann. Soc. Linn., 1859, pp. 261-295, 12 plates.
- JOB, AUGUSTE. Catalogue des Mollusques terrestres et fluviatiles dans le département de la Moselle.
Bull. Soc. Hist. Nat. Metz, 1844, 1 plate. Separate.
- JOB, NUMA. Note sur des Anodonta cygnea, etc.
Ann. des Sci. Nat., 1845, pp. 373, 374.
- * JOHNSON, C. W. Ellwood Pleas.
Nautilus., XI, 1898, pp. 114, 115.
- * JOHNSON, GEORGE. An Introduction to Conchology, etc., 1850.
- * JOHNSTON, R. M. [Observations on the Variability of the Tasmanian Unio.]
Proc. Roy. Soc. Tasm., 1888, pp. 84, 90, 3 plates.
- * — Provisional Aid to the Study of the Tasmanian Mollusca.
Proc. Roy. Soc. Tasm., 1890, pp. 57-151, 1 plate.
- JONSTON, JOH. Historia Naturalis, etc., 1657.
- JORDAN, A. Catalogue of the British Mollusca, 1839.
- * JORDAN, HERMANN. Die Mollusken der preussischen Oberlausitz.
Jahrb. Deutsch. Mal. Ges., VI, 1879, pp. 291-384.
- Einfluss der bewegten Wassers auf die Gestaltung der Muscheln aus der Familie Najades.
Biol. Centr., I, 1881, pp. 392-399.
- JORDAN, HERMANN. Die Binnenmollusken der nördlich gemäßigten Länder von Europa, etc.
Nova Acta Acad. Leo. Car., XLV, 1884, pp. 181-402, 8 plates.
- * JOUSSEAU, F. Faune malacologique des environs de Paris.
Bull. Soc. Zool. de France, 1876-1883.
- * — Coquilles du Haut-Sénégal.
Bull. Soc. Zool. de France, 1876, pp. 471-502, 3 plates.
- * — Mollusques recueillis à Ceylon par M. Simon, etc.
Mém. Soc. Zool. de France, 1891, pp. 264-330, 1 plate.
- JURINE, L. Catalogue des mollusques trouvés aux environs de Genève, 1817.
- * KAMMERER. Die Conchylien im Cabinet des Herrn Erbprinzen, etc., 1786.
- KARSTEN. Museum Leskeanum, 1789.
- * KAWAIL, H. La pêche des perles en Livonie.
Ann. Soc. Mal. Belg., VII, 1872, pp. 38-46.
- * KEBER, GOTHARD AUG. FERD. De Nervis Concharum, 1837.
- * — Beiträge zur Anatomie und Physiologie, etc., 1851.
- Bisherige Ansichten der Teichmuscheln.
Fror. Tagsber., 477, 1852, pp. 275-278.
- * KEEP, JOSIAH. West Coast Shells, etc., 1888.
- * KELLY, H. M. A Statistical Study of the Parasites of the Unionidae.
Bull. Ill. Lab. N. H., V, 1899, pp. 399-418.
- KENNICOTT, ROBERT. Catalogue of Animals observed in Cook County, Illinois.
Trans. Ill. Ag. Soc., I, 1855, p. 577.
- * KENYON, JOSEPH. Remarks on British Land and Freshwater Shells.
Mag. Nat. Hist., I, 1829, p. 424.
- * KEW, H. W. The Dispersal of Shells, etc., with a preface by Alfred Russell Wallace, 1893, 291 pp.
- * KEYES, CHARLES R. On the Fauna of the Lower Coal Measures of Central Iowa.
Proc. Acad. Nat. Sci. Phila., 1888, pp. 222-246.
- * — An Annotated Catalogue of the Mollusca of Iowa.
Bull. Essex Inst., XX, 1888, pp. 61-83.

- * KICKX, JEAN (fils). Synopsis Molluscorum Brabantiae, etc., 1830, pp. 97.
- * KIRSCH, PHILIP H. Biennial Report of the Fish Commissioner of Ohio, 1896, pp. 147.
- * KIRTLAND, JARED P. Observations on the Sexual Characters of the Naiades. *Am. Jour. Sci. and Arts*, XXVI, 1834, pp. 117-120, figs.; *Wieg. Arch.*, 1836, I, p. 236.
- Report on the Zoology of Ohio, 1838.
- * — Sexual Differences and Habits of Naiades. *Am. Jour. Sci. and Arts*, XXXIX, 1840, p. 164.
- * — Fragments of Natural History. *Am. Jour. Sci. and Arts*, XXXIX, 1840, pp. 164-168.
- * — Remarks on the Sexes and Habits of some Acephalous Bivalve Mollusca. *Proc. Am. Ass. Adv. Sci.*, V, 1851, pp. 85-91.
- KLEBER, F. Zur controverse über die Befruchtung des Flus muschel, etc. *Arch. f. Anat.*, 1869, pp. 284-306.
- * KLEEBERG, G. Molluscorum Bornsiorum Synopsis. Dissert. inang. Regiomonti, 1828, pp. 43.
- * KLEES, JOANNES G. Dissertatio Inauguralis Zoologica, etc., 1818, pp. 45.
- * KLEIN, JACOBUS THEODORUS. Tentamen Methodi Ostracologicae, etc., 1753, 12 plates.
- KLIPSTEIN. Zur Geologischen Kenntniss, etc., *Geol. Jahrb.*, 1843.
- * KLODEN, K. F. Die Versteinerungen Branderberg, etc., 1834.
- KNAUTH, KARL. Perle in Unio rostratus. *Zool. Gart.*, 1896, p. 155.
- KNORR, GEORGE WOLFGANG. Vergnügen der Augen und des Gemüths, in Vorstellung einer Sammlung von Muscheln und andern Geschöpfen, etc., 1757-1772.
- * KOBELT, W. Zur Entwicklung der Najaden. *Nach. Deutsch. Mal. Ges.*, 1870, p. 149.
- * — Fauna der Nassauischen Mollusken, 1871, pp. 286.
- Catalog der im Europäischen Faunagebiet, etc., 1871, pp. 150; new ed. 1881, pp. 294.
- * KOBELT, W. Die nordschleswigsche Perlenmuschel. *Mal. Blatt.*, 1872, pp. 142-147.
- * — Die Geographische Verbreitung der Mollusken. *Jahrb. Deutsch. Mal. Ges.*, 1878, pp. 170-185; 1880, pp. 1-30, 241-286.
- Rossmässler's Iconographie der europäischen Land- und Süßwasser-Mollusken, from 1876-1899, not completed.
- * — Fauna Japonica Extramarina. *Abh. Senck. Nat. Ges.*, 1878, 1879, pp. 171, 23 plates.
- * — [Margaritana margaritifera found in Anticosti.] *Nach. Deutsch. Mal. Ges.*, 1883, p. 93.
- * — Neue Tanisische Unioneu. *Nach. Deutsch. Mal. Ges.*, 1884, pp. 182, 183.
- * — Eine Nassauische Perlenmuschel. *Nach. Deutsch. Mal. Ges.*, 1886, pp. 88-90.
- Erster Nachtrag zur Fauna der Nassauischen Mollusken. *Jahrb. Nassau. Ver.*, 1886, pp. 70-103. Separate. *Abst. Nach. Mal. Ges.*, 1886, p. 153.
- Unios y Anodontas nuevos de la fauna española. *An. Soc. Esp.*, XVI, 1887, pp. 435-439.
- * — Die deutschen Bivalven ein Vorschaltz zu gemeinsamer Arbeit. *Nach. Deutsch. Mal. Ges.*, XX, 1888, pp. 47-50.
- * — Die Bivalven Nieder-Andalusiens. *Nach. Deutsch. Mal. Ges.*, 1888, pp. 16-30.
- * — Diagnosen neuer Palaearctischer Arten. *Nach. Deutsch. Mal. Ges.*, 1893, pp. 150-153.
- Zweiter Nachtrag zur Fauna der Nassauischen Mollusken. *Jahrb. Nassau. Ver.*, 1894, pp. 83-89, 1 plate.
- * — Land- und Süßwasser-Konchylien. *Abh. Senck. Nat. Ges.*, XXIV, 1897, pp. 17-92, 8 plates.
- KOELREUTER, J. T. Observationes anatomico-physiologicae Mytili cygnei, etc. *Nora Acta Ac. Pet.*, 1788, pp. 206-239.

- KOENIG, EMANUEL. Denham and Clapperton's Travels in Africa, 1826.
- *KOROID, C. A. Michigan Unionidae.
Am. Naturalist, XXXII, 1898, pp. 956, 957.
- *KOKEIL, FRED. In Rossmässler's Iconographic.
— In Gallenstein's Kärnten Conchylien. 1852.
- *KONINCK, L. DE. Description des Animaux fossiles, etc., 2 vols. 1842, 1844.
- *KRAUSS, FERDINAND. Die Süd Afrikanischen Mollusken, etc., 1848, pp. 146, 6 plates.
- *— Eine Neue Art Castalia.
Zeits. für Mal., 1849, p. 99.
- *— Reise der Österreichischen Fregatte Novara, 1867.
- *KRELINGER, CARL. Systematisches verzeichniss, etc., 1870.
- *KRUGER, J. F. Urvweltliche Naturgeschichte, etc , 1825.
- KRYNICKI, J. Conchyliia tam Terrestria quam Fluvialitia, etc.
Bull. Soc. Nat. Mosc., X, 1837, p. 50.
- *KUSTER, H. C. In Systematisches Conchylien Cabinet (1848), 1852-1862.
— Die Binnen Mollusken Fauna von Triest, etc.
Ber. Ges. Bamberg, IX, 1871, pp. 86-101.
- KUTSCHIG, —. Fauna Dalmatica, 1866.
- LABECHE, DE. Catalogue des Mollusques de Genève.
Zool. Jour., I, 1824, pp. 97-168, 4 plates.
- *LACAZE-DUTHIERS, H. Recherches sur les Organes Genitaux des Acephales, etc.
Ann. des Sci. Nat., 1854, pp. 155-248.
- *— Mémoire sur l'Organe de Bojanus des Acephales, etc.
Ann. des Sci. Nat., 1855, pp. 267-319.
- *— Observations sur l'hermaphroditisme des Anodontes.
Ann. des Sci. Nat., 1855, pp. 381, 382.
- *— Le Développement des Branchies Acephales, etc.
Ann. des Sci. Nat., 1856, pp. 5-47.
- *LALLEMANT, CH., and SERVAIN, G. Catalogue des Mollusques de Janlgonne, 1869.
- *LAMARCK, J. B., in Encyclopédie Méthodique, 1791-1832.
- *LAMARCK, J. B. Prodrôme d'une nouvelle classification des Coquilles.
Mém. Soc. Hist. Paris, 1799, pp. 63-91.
- *— Système des Animaux sans Vertèbres, etc., 1801.
— In *Philosophie Zoologique*, 1805, p. 328; 1808, p. 138.
— Extrait d'un Cours de Zoologie, etc., 1812.
— Histoire Naturelle des Animaux sans Vertèbres, 1815-1822.
— Exposition des considerations relatives à l'histoire naturelle des Animaux, etc.
Philosophie Zoologique, I, 1830, p. 318.
- LAMOROUX, JEAN VIET. Unio antistrophodonta bei Nancy.
Neue Jahrbuch, p. 444, 1837.
- LAMPREY, JONES. Notes on Geology of the Great Plains.
Jour. China Br. Roy. As. Soc. Shang., 1865, p. 16.
— Notes on the Geology of China.
Jour. China Br. Soc. Shang., 1866.
- LANDRIN, ARMAND. Lu à la Seance du Decembre, 1864, de la Soc. Sci. Nat. Seine-et-Oise, pp. 1-6, 2 plates.
- *LANGER, KARL. Das Gefäss-System der Teichmuschel.
Sitz. Math. Nat., 1854.
- *LAPHAM, I. A. Systematic Catalogue of the Animals of Wisconsin, 1851.
— Catalogue of the Mollusca of Wisconsin.
Trans. Wis. Ag. Soc., 1852, pp. 367-370.
- *— List of the Shells of Wisconsin.
Proc. Acad. Nat. Sci. Phila., 1860, pp. 154-156.
- *LARKEN, E. P. Does Unio spin a byssus?
Science, III, 1884, p. 302.
- LASKEY, JAMES. Account of North British Testacea.
Mem. Wern. Soc., 1811, pp. 370-417, 1 plate.
- *LATCHFORD, F. R. Notes on the Ottawa Unionidae.
Trans. Ottawa Field Nat. Club, 1882, pp. 48-57.
- *LATREILLE, PIERRE ANDRÉ. Esquisse d'une distribution générale des Mollusques, etc.
Ann. Soc. Nat., 1824, p. 317, 1 plate.

- *LATTER, OSWALD H. Notes on Anodon and Unio.
Proc. Zool. Soc., 1891, pp. 52-59, 1 plate. Separate. Abstract in *Jour. Roy. Mic. Soc.*, 1891, p. 455.
- *LEA, ISAAC. Description of six new species of Unios, etc.
Trans. Am. Phil. Soc., 1827, pp. 259-273, 4 plates. Separate.
- *— Observations on the Genus Unio, etc., 1829, pp. 71, 12 plates.
- *— Description of a New Genus of Naiades, etc.
Trans. Am. Phil. Soc., 1829, pp. 403-457, 8 plates.
- *— Observations on the Naiades, etc.
Trans. Am. Phil. Soc., 1831, pp. 63-121, 16 plates.
- *— Observations on the Genus Unio, etc., 1832, pp. 133.
- *— Observations on the Naiades, etc.
Trans. Am. Phil. Soc., 1834, pp. 23-119, 19 plates.
- *— Observations on the Genus Unio, I, 1834, 233 pp, 46 plates.
- *— A Synopsis of the Family of Naiades, 1836; 2d ed., 1838; 3d ed., 1852; 4th ed., 1870.
- Description of a New Unio (spinosus), 1836.
- Neue Gattungen von Najaden.
Zeits. für Zool. Ver. Am., 1837, pp. 43-53, 1 plate.
- *— Description of New Fresh-water and Land Shells.
Trans. Am. Phil. Soc., 1838, pp. 1-154, 34 plates.
- *— Observations, etc., II, 1838, pp. 152, 29 plates.
- *— Descriptions of New Fresh-water and Land Shells.
Proc. Am. Phil. Soc., 1840, pp. 284-289.
- *— Continuation of above.
Proc. Am. Phil. Soc., 1841, pp. 30-34, 81-83.
- *— Description of new Fresh-water and Land Shells.
Trans. Am. Phil. Soc., 1842, pp. 163-252, 23 plates.
- *— Observations, etc., III, 1842, pp. 88, 23 plates.
- *— Continuation of paper on New Shells.
Proc. Am. Phil. Soc., 1842, pp. 224, 225.
- *LEA, ISAAC. [Communication.]
Proc. Acad. Nat. Sci. Phila., 1843, pp. 215, 216.
- Description of twelve new species of Uniones.
Proc. Am. Phil. Soc., IV, 1843, p. 11. Abstract in advance, no pagination.
- *— Names of twelve new Uniones.
Proc. Am. Phil. Soc., 1844, p. 8.
- *— Descriptions of new Fresh-water and Land Shells.
Proc. Am. Phil. Soc., 1845, pp. 162-168.
- *— Description of new Fresh-water and Land Shells.
Trans. Am. Phil. Soc., 1845, pp. 275-282, 4 plates.
- *— Descriptions of new Fresh-water and Land Shells.
Trans. Am. Phil. Soc., 1848, pp. 67-101, 9 plates.
- *— Observations, etc., IV, 1848, pp. 101, 13 plates.
- *— Descriptions of five new Anodontæ, etc.
Proc. Zool. Soc., 1850, pp. 197-199; *Ann. Nat. Hist.*, VIII, 1851, pp. 493-496.
- *— Exhibition of large Unios, etc.
Proc. Am. Phil. Soc., 1850, p. 153.
- *— On a communication from Professor Agassiz.
Proc. Am. Phil. Soc., 1851, pp. 187-189.
- *— On the great size of certain Naiades, etc.
Proc. Am. Phil. Soc., 1851, pp. 191-193.
- *— Correspondence between Mr. Lea and Prof. Agassiz, etc.
Proc. Am. Phil. Soc., 1851, pp. 218-221.
- *— Descriptions of new species of the family Unionidæ.
Proc. Am. Phil. Soc., 1852, pp. 251, 252; *Trans. Am. Phil. Soc.*, 1852, pp. 253-294, 18 plates.
- *— Observations, etc., V, 1852, p. 62, 19 plates.
- *— On a remarkably large *Dipsas plicatus*.
Proc. Acad. Nat. Sci. Phila., 1852, p. 52.
- *— On the period of maturing and ejecting young in the family Unionidæ.
Proc. Acad. Nat. Sci. Phila., 1852, p. 54.

- *LEA, ISAAC. Description of a new Syn-
phynote Unio.
Proc. Acad. Nat. Sci. Phila., 1852,
pp. 54, 55.
- *— Definition of the term species and
remarks on classification, etc.
Proc. Acad. Nat. Sci. Phila., 1852,
p. 57, 58.
- *— Description of new species of Unio-
nidae.
Proc. Acad. Nat. Sci. Phila., 1852,
pp. 251, 252.
- *— Notice of the *Mya nodulosa*, Wood.
Proc. Acad. Nat. Sci. Phila., 1853,
p. 368.
- *— On the *Castalia sulcata*, Krauss, etc.
Proc. Acad. Nat. Sci. Phila., 1853,
pp. 376, 377.
- *— Rectification of Conrad's Synopsis
of the Naiades, etc.
Proc. Acad. Nat. Sci. Phila., 1854,
pp. 236-249.
- *— Description of a new subgenus of
Naiades.
Proc. Acad. Nat. Sci. Phila., 1856,
p. 79.
- *— Description of a new *Triquetra*.
Proc. Acad. Nat. Sci. Phila., 1856,
p. 79.
- *— Description of new Fresh-water
Shells from California.
Proc. Acad. Nat. Sci. Phila., 1856,
pp. 80, 81.
- *— Description of twenty-five new
Exotic Uniones.
Proc. Acad. Nat. Sci. Phila., 1856,
pp. 92-95.
- *— Description of four new Exotic
Uniones.
Proc. Acad. Nat. Sci. Phila., 1856,
p. 103.
- *— Description of the *Byssus* in Unio.
Proc. Acad. Nat. Sci. Phila., 1856,
pp. 213, 214.
- *— Description of eleven new Uniones
from Georgia.
Proc. Acad. Nat. Sci. Phila., 1856,
pp. 262, 263.
- *— [Communication on *Triquetra lan-
ceolata*, etc.]
Proc. Acad. Nat. Sci. Phila., 1856,
pp. 300, 301.
- *— Remarks on the visual organs of
Unios.
Proc. Acad. Nat. Sci. Phila., 1857,
p. 17; *Jour. de Conch.*, 1857, p. 203.
- *LEA, ISAAC. Description of thirteen
new Uniones from Georgia.
Proc. Acad. Nat. Sci. Phila., 1857,
pp. 31, 32.
- *— Description of six new Uniones from
Alabama.
Proc. Acad. Nat. Sci. Phila., 1857,
p. 83.
- *— Description of eight new Naiades.
Proc. Acad. Nat. Sci. Phila., 1857,
p. 84.
- *— Description of three new Naiades.
Proc. Acad. Nat. Sci. Phila., 1857,
p. 85.
- *— Description of twelve new Naiades
from North Carolina.
Proc. Acad. Nat. Sci. Phila., 1857,
pp. 85, 86.
- *— [Remarks on *Unio spinosus*.]
Proc. Acad. Nat. Sci. Phila., 1857,
p. 101.
- *— Description of six new Fresh-water
and Land Shells of Texas and Ta-
maulipas.
Proc. Acad. Nat. Sci. Phila., 1857,
pp. 101, 102.
- *— Description of four new Exotic
Uniones, etc., 1857.
- *— [Proposed change of name of two
Unios.]
Proc. Acad. Nat. Sci. Phila., 1857,
p. 104.
- *— Descriptions of twenty-seven new
Uniones from Georgia.
Proc. Acad. Nat. Sci. Phila., 1857,
pp. 169-172.
- *— [Remarks on plates of Naiades of
Siam.]
Proc. Acad. Nat. Sci. Phila., 1857,
p. 205.
- *— Description du sous-genre *Plagio-
don*, etc.
Jour. de Conch., 1857, pp. 207, 208.
- *— Descriptions of Exotic Genera and
Species of Unionidae.
Jour. Acad. Nat. Sci. Phila., 1857,
pp. 289-321, 13 plates.
- *— Observations, etc., VI, Pt. 1, 1857,
pp. 34, 13 plates.
- *— Description du *Byssus* dans le Genre
Unio.
Jour. de Conch., 1857, p. 200.
- *— Remarks on Unionidae of Nebraska
Territory, etc.
Proc. Acad. Nat. Sci. Phila., 1858,
pp. 13-14.

*LEA, ISAAC. Descriptions of new Unios from Tennessee, etc.

Proc. Acad. Nat. Sci. Phila., 1858, pp. 40, 41.

*—[Remarks on some remarkable Unionidæ.]

Proc. Acad. Nat. Sci. Phila., 1858, p. 114.

*—Description of four new Freshwater Mollusks from Darien.

Proc. Acad. Nat. Sci. Phila., 1858, p. 118.

*—[Remarks on reproduction of *Unio rubiginosus*.]

Proc. Acad. Nat. Sci. Phila., 1858, p. 135.

*—Descriptions of seven new Margaritanas and four new Anodontas.

Proc. Acad. Nat. Sci. Phila., 1858, pp. 138, 139.

*—Descriptions of twelve new Uniones of the United States.

Proc. Acad. Nat. Sci. Phila., 1858, pp. 165, 166.

*—[Remarks on *Triqueta contorta*.]

Proc. Acad. Nat. Sci. Phila., 1858,

p. 179.

*—Descriptions of the embryonic forms of thirty-eight species of Unionidæ.

Jour. Acad. Nat. Sci. Phila., 1858, pp. 43-50, 1 plate, separate.

*—New Unionidæ of the United States.

Jour. Acad. Nat. Sci. Phila., 1858, pp. 51-95, 15 plates.

*—Descriptions of Exotic Genera and Species of Unionidæ.

Jour. Acad. Nat. Sci. Phila., 1858, pp. 289-321, 13 plates.

*—Observations, etc., VI, Pt. 2, 1858, 52 pp., 16 plates.

*—Observations, etc., VI, 1858, 97 pp., 29 plates.

*—Descriptions of eight new Unionidæ from Georgia, etc.

Proc. Acad. Nat. Sci. Phila., 1859, pp. 112, 113.

*—Descriptions of twenty-one new Exotic Unionidæ.

Proc. Acad. Nat. Sci. Phila., 1859, pp. 151-154.

*—Descriptions of two new Uniones from Georgia.

Proc. Acad. Nat. Sci. Phila., 1859, p. 154.

*LEA, ISAAC. Descriptions of seven new Uniones from South Carolina, etc.

Proc. Acad. Nat. Sci. Phila., 1859, pp. 154, 155.

*—Descriptions of twelve new Uniones from Georgia.

Proc. Acad. Nat. Sci. Phila., 1859, pp. 170-172.

*—[On dead shells of *Anodonta lewisii*.]

Proc. Acad. Nat. Sci. Phila., 1859, p. 177.

*—[On *Unio kleinianus*.]

Proc. Acad. Nat. Sci. Phila., 1859, p. 177.

*—Descriptions of four new Exotic Unionidæ.

Proc. Acad. Nat. Sci. Phila., 1859, p. 187.

*—New Unionidæ of the United States.

Jour. Acad. Nat. Sci. Phila., 1859, pp. 191-233, 12 plates.

*—Observations, etc., VII, 1859, Pt. 1, 43 pp., 12 plates.

*—[Change of name of *Margaritana etowahensis*.]

Proc. Acad. Nat. Sci. Phila., 1859, p. 280.

*—Description of a new *Unio* from Darien.

Proc. Acad. Nat. Sci. Phila., 1859, p. 281.

*—[Remarks on four Unios, etc.].

Proc. Acad. Nat. Sci. Phila., 1859, pp. 329-330.

*—Descriptions of three new Exotic Uniones.

Proc. Acad. Nat. Sci. Phila., 1859, p. 331.

*—Remarks on the number of species of Unionidæ, etc.

Proc. Acad. Nat. Sci. Phila., XII, 1860, p. 3.

*—Observations, etc., VII, 1860, Pt. 2, pp. 38, 13 plates.

*—Observations, etc., VII, 1860, pp. 93, 25 plates.

*—[Remarks on Abnormal Shells.]

Proc. Acad. Nat. Sci. Phila., 1860, pp. 51-53.

*—Descriptions of eight new Unionidæ from Georgia, etc., 1860, pp. 24.

- *LEA, ISAAC. [On the number of Unionidæ in North America.]
Proc. Acad. Nat. Sci. Phila., 1860, p. 85.
- *— [On letter of Dr. Lewis.]
Proc. Acad. Nat. Sci. Phila., 1860, pp. 88, 89.
- *— Descriptions of four new Unionidæ from Brazil, etc.
Proc. Acad. Nat. Sci. Phila., 1860, pp. 89, 90.
- *— Descriptions of fifteen new Uruguayan Unionidæ.
Proc. Acad. Nat. Sci. Phila., 1860, pp. 90-92.
- *— Descriptions of five new Uniones from North Alabama.
Proc. Acad. Nat. Sci. Phila., 1860, pp. 92, 93.
- *— [On a new form of Palpi of Anodonta from the Uruguay River.]
Proc. Acad. Nat. Sci. Phila., 1860, p. 97.
- *— Descriptions of two new Uniones from Georgia.
Proc. Acad. Nat. Sci. Phila., 1860, p. 305.
- *— Descriptions of three new Uniones from Mexico.
Proc. Acad. Nat. Sci. Phila., 1860, pp. 305, 306.
- *— Descriptions of seven new Unionidæ from the United States.
Proc. Acad. Nat. Sci. Phila., 1860, pp. 306, 307.
- *— Descriptions of six new Unionidæ from Alabama.
Proc. Acad. Nat. Sci. Phila., 1860, p. 307.
- *— Descriptions of three new Exotic Unionidæ.
Proc. Acad. Nat. Sci. Phila., 1860, pp. 307, 308, 331; 1865, pp. 75, 76; extract in *Am. Jour. Conch.*, I, 1865, pp. 354, 355.
- *— Descriptions of Exotic Unionidæ.
Jour. Acad. Nat. Sci. Phila., 1860, pp. 235-273, 13 plates.
- *— New Unionidæ of the United States, etc.
Jour. Acad. Nat. Sci. Phila., 1860, pp. 327-374, 16 plates.
- *— Observations, etc., VIII, Pt. 1, 1860, pp. 56, 16 plates.
- *LEA, ISAAC. Descriptions of twenty-five new Unionidæ from Georgia, Alabama, Mississippi, Tennessee, and Florida.
Proc. Acad. Nat. Sci. Phila., 1861, pp. 38-41.
- *— Descriptions of two new Anodontæ from Arctic America.
Proc. Acad. Nat. Sci. Phila., 1861, p. 56.
- *— Descriptions of twelve new Uniones from Alabama.
Proc. Acad. Nat. Sci. Phila., 1861, pp. 59, 60.
- *— Descriptions of eleven new Unios from the United States.
Proc. Acad. Nat. Sci. Phila., 1861, pp. 391-393.
- Remarks on the number of Unionidæ, etc., 1862.
- *— New Unionidæ of the United States.
Jour. Acad. Nat. Sci. Phila., 1861, pp. 53-109, 18 plates.
- *— Observations, etc., VIII, 1862, Pt. 2, pp. 56, 18 plates.
- *— Observations, etc., VIII, 1862, pp. 115, 34 plates.
- *— Descriptions of ten new Unionidæ of the United States.
Proc. Acad. Nat. Sci. Phila., 1862, pp. 168, 169.
- *— Descriptions of two new Exotic Uniones, etc.
Proc. Acad. Nat. Sci. Phila., 1863, p. 176.
- Description of a new Genus, etc., 1862.
- *— Checklist of the Shells of North America. Unionidæ.
Smithsonian Miscellaneous Collection, II, pp. 52, 1862. Separate.
- *— New Unionidæ of the United States.
Jour. Acad. Nat. Sci. Phila., 1862, pp. 53-109, 18 plates.
- *— New Unionidæ of the United States, etc.
Jour. Acad. Nat. Sci. Phila., 1862, pp. 187-216, 10 plates.
- *— Observations, etc., IX, 1863, pp. 180, 16 plates.
- *— Descriptions of eleven new Exotic Unionidæ.
Proc. Acad. Nat. Sci. Phila., 1863, pp. 189, 190.

- *LEA, ISAAC. Description of a new Unio and a Monocondylea.
Proc. Acad. Nat. Sci. Phila., 1863, p. 190.
- *— Descriptions of twenty-four new Unionidæ of the United States.
Proc. Acad. Nat. Sci. Phila., 1863, pp. 191-194.
- *— New Exotic Unionidæ.
Jour. Acad. Nat. Sci. Phila., 1863, pp. 377-400, 10 plates.
- *— Descriptions of the soft parts of one hundred and forty-three species, and some Embryonic Forms of Unionidæ, etc.
Jour. Acad. Nat. Sci. Phila., 1863, pp. 401-456.
- *— Observations, etc., X, 1863, pp. 94, 10 plates.
- *— Descriptions of six new Unionidæ from Lake Nyassa.
Proc. Acad. Nat. Sci. Phila., 1864, pp. 108, 109.
- *— Descriptions of two new Unionidæ from South Africa.
Proc. Acad. Nat. Sci. Phila., 1864, p. 113.
- *— Descriptions of six new Western Asiatic Unionidæ.
Proc. Acad. Nat. Sci. Phila., 1864, pp. 285, 286.
- *— Descriptions of three new Exotic Unionidæ.
Proc. Acad. Nat. Sci. Phila., 1865, pp. 75, 76.
- *— Descriptions of eight new Unios from the United States.
- Descriptions of fourteen new Melanidæ, etc., 1866, pp. 32.
Appendix IV [to Bibliography of the North American Conchology, by W. G. Binney.]
Smithsonian Miscellaneous Collection, IX, 1869, pp. 289-294.
- *— Description of twelve new Unionidæ from South America.
Proc. Acad. Nat. Sci. Phila., 1866, pp. 33-35.
- *— Description of five new Unios.
Proc. Acad. Nat. Sci. Phila., 1866, p. 133.
- *— New Unionidæ Melanidæ, etc.
Jour. Acad. Nat. Sci. Phila., 1867, pp. 5-65, 21 plates; *Am. Jour. Conch.*, 1867, pp. 84, 85. Separate.
- *LEA, ISAAC. Observations, etc., XI, 1867, pp. 146, 24 plates.
- *— Index to Volumes I-XI of Observations, 1867, pp. 63.
- *— Descriptions of five new Unionidæ, etc.
Proc. Acad. Nat. Sci. Phila., 1867, p. 81.
- *— Descriptions of nine new Unios from Lake Nicaragua.
Proc. Acad. Nat. Sci. Phila., 1868, pp. 94-96.
- *— Description of sixteen new Unionidæ, etc.
Proc. Acad. Nat. Sci. Phila., 1868, pp. 143-145.
- *— Notes on some singular Chinese Unios.
Proc. Acad. Nat. Sci. Phila., 1868, pp. 145, 146; *Am. Jour. Conch.*, 1868, p. 253.
- *— Descriptions of four new Exotic Unionidæ.
Proc. Acad. Nat. Sci. Phila., 1868, p. 150.
- *— Descriptions of seven new Unios from North Carolina.
Proc. Acad. Nat. Sci. Phila., 1868, pp. 160, 161.
- *— Description of two new Unionidæ from Ecuador.
Proc. Acad. Nat. Sci. Phila., 1868, pp. 161, 162.
- *— Descriptions of Unionidæ from the Lower Cretaceous of New Jersey.
Proc. Acad. Nat. Sci. Phila., 1868, pp. 162-164.
- Descriptions of twelve new South American Unionidæ, 1868.
- *— New Unionidæ, Melanidæ, etc., of the United States.
Jour. Acad. Nat. Sci. Phila., 1868, pp. 249-302, 17 plates; pp. 303-343, 9 plates.
- *— Observations, etc., XII, 1869, pp. 105, 25 plates.
- *— Index to Volume XII, etc., 1869, pp. 23.
- *— Descriptions of six new Freshwater Shells.
Proc. Acad. Nat. Sci. Phila., 1869, pp. 124, 125.
- *— Are Unios sensitive to light?
Am. Jour. Sci. and Arts, XLVII, 1869, pp. 430, 431; *Ann. Nat. Hist.*, III, 1869, p. 466.

- *LEA, ISAAC. Descriptions of three new Exotic Unionidae.
Proc. Acad. Nat. Sci. Phila., 1871, p. 188.
- *— Descriptions of twenty new Uniones of the United States.
Proc. Acad. Nat. Sci. Phila., 1871, pp. 189-193.
- *— Descriptions of twenty-nine Unionidae from the United States.
Proc. Acad. Nat. Sci. Phila., 1872, pp. 155-161.
- *— Rectification of Conrad's Synopsis, etc., 1872.
- *— Description of seven new Unionidae, etc.
Proc. Acad. Nat. Sci. Phila., 1874, pp. 422, 423.
- *— Descriptions of three new Uniones from the United States.
Proc. Acad. Nat. Sci. Phila., 1873, p. 424.
- *— Description of fifty-two species of Unionidae.
Jour. Acad. Nat. Sci. Phila., 1874, pp. 5-54, 18 plates; pp. 55-59, 4 plates.
- *— Observations, etc., XIII, 1874, pp. 75, 22 plates.
- *— Index to Volumes I-XIII, Observations, etc., 1874, 29 pp.
- LEA, ISAAC, and BINNEY, W. G. Explorations in Nebraska, etc.
Rep. on Geol. and Nat. Hist., of Nebraska, by F. V. Hayden, 1858-9.
- *LEACH, WILLIAM ELDFORD. The Zoological Miscellany, 1815.
- *— A synopsis of the Mollusca of Great Britain, 1852. Completed by J. E. Gray.
- LEE, W. S. How the Muskrat opens the Unio.
Jour. Trenton Soc., I, 1886, p. 8.
- *LEENWENHOEK, ANT. VON. Arcana Naturæ delecta, etc., 1695.
- LE GOUVIERIE, T. La pêche aux perles dans la Loire, 1880.
- *LEHMAN, R. Die Schnecken and Muscheln, 1873.
- *LEIDIG, —. In Mollusken der Main, 1866.
- *LEIDY, JOS. [On the Parasites of Anodonta fluviatilis.]
Proc. Acad. Nat. Sci. Phila., 1837, p. 18.
- *LEIDY, JOS. On the reproduction and Parasites of Anodonta fluviatilis.
Proc. Acad. Nat. Sci. Phila., 1883, pp. 44, 45; *Science*, 1883, p. 150; *Nach. Deutsch. Mal. Ges.*, 1883, p. 116.
- LELIEVRE, A. Clef dichotomique pour la détermination des Mollusques du Nord de la France.
Bull. Sci. Dép. Nord, 1878, pp. 84, 85, 142-153, 178-183.
- *LEMON, JAMES H. Interglacial Shells at Toronto.
Nautilus, XII, 1898, pp. 6, 7.
- LESKE. Anfangs der Naturgeschichte, 1784.
- LESSER, CREUSE DE. Statistique du département de l'Hérault, 1824.
- *LESSER, FREDERIC CHRISTIAN. Testaceo Theologia, etc., 2 vols., 1744, 1776.
- *LESSON, R. P. Voyage de la Coquille. Zool., II, 1730.
— Synopsis Méthodique des Mollusques, 1830.
- LETOURNEUX, ARISTIDE. Mollusques terrestres et fluviatiles de Lamaloules-Bains, 1877.
- LETOURNEUX, TACITE. Catalogue des Mollusques du département de la Vendée, et particulièrement dans Fontenay le Comte.
Rev. et Mag. Zool., XX., 1869, pp. 49-64, 105-109, 145-148, 193-203. Separate.
- *LETOURNEUX, A., and BOURGUIGNAT, J. R. Prodrome de la Malacologie de Tunisie, 1887.
- LEVILLE, J. B. T. Manuel d'histoire naturelle des Oiseaux, des Poissons, Paris, 1800.
- *LEWIS, JAMES. [List of Mollusca in Little Lakes, Otsego County, New York.] In letter to Dr. Lea.
Proc. Acad. Nat. Sci. Phila., 1856, pp. 259, 260.
- *— [On Alasmodonta marginata Say.]
Proc. Bost. Soc. Nat. Hist., V, 1854, pp. 120-124.
- *— Notes on Mollusca.
Proc. Bost. Soc. Nat. Hist., 1855, pp. 121-124.
- *— [On the Shells of New York].
Proc. Bost. Soc. Nat. Hist., 1856, pp. 295-298.

- *LEWIS, JAMES. Shells in Herkimer and Otsego counties.
Proc. Bost. Soc. Nat. Hist., 1856, pp. 1-4.
- *— [Erosion of Unios.]
Proc. Bost. Soc. Nat. Hist., 1857, pp. 149-151.
- *— Catalogue of the Mollusks in the vicinity of Mohawk, etc.
Proc. Acad. Nat. Sci. Phila., 1860, pp. 17-19.
- *— Colors of Unionidæ.
Proc. Bost. Soc. Nat. Hist., 1860, pp. 245, 246; *Proc. Acad. Nat. Sci. Phila.*, 1860, p. 88.
- *— Letter in relation to habits and growth of Unionidæ.
Proc. Acad. Nat. Sci. Phila., 1861, pp. 57, 58.
- *— On the Shells of Michigan described by A. O. Currier.
Am. Jour. Conch., IV, 1868, pp. 81-85.
- *— Remarks on the Mollusks of the Valley of the Mohawk.
Am. Jour. Conch., 1868, pp. 241-245.
- *— Catalogue of the Shells of the Coosa River, Alabama.
Am. Jour. Conch., 1869, pp. 166-169.
- *— On the Shells of the Holston River.
Am. Jour. Conch., 1870, pp. 216-226.
- *— Land and Fresh-water Shells of New York.
Bull. Buffalo Soc. Nat. Hist., 1874, pp. 127-142.
- *— Fresh-water and Land Shells of Alabama.
Fauna of Ala., 1876, pp. 1-42, 4 supp.
- *— Unionidæ of Ohio and Alabama.
Proc. Acad. Nat. Sci. Phila., 1877, pp. 26-36.
- *— On Unio subrostratus, Say.
Proc. Acad. Nat. Sci. Phila., 1879, pp. 273-275.
- *— Suggestions to collectors. [No date.]
- LIESVILLE, DE. Catalogue des Mollusques aux environs d'Alençon, 1856.
- LILLIE, F. R. Account of the Embryology of Unio complanata.
Jour. Morph., VIII, 1893, pp. 569-578, 1 plate.
- The Embryology of the Unionidæ.
Jour. Morph., X, 1895, pp. 1-100, 6 plates.
- *LILLIE, F. R. On the Origin of the Centers of the First Cleavage Spindle in Unio complanata.
Science, V, 1897, pp. 389, 390.
- Centrosome and Sphere in the Egg of Unio.
Zool. Bull., I, 1898, pp. 265-274, figs.
- LINK, H. F. Beschreibung der Naturalien-Sammlung, III, 1808.
- LINNEUS, CAROL. Wästgötha Resa. Stockholm, 1746.
- *— Systema Naturæ, etc., 10th ed., 1758, 2 vols.; 12th ed. 1766-67, 3 vols.
- Fauna Suecica sistens Animalii, etc., 1761.
- *LINSLEY, JAMES H. Catalogue of the Shells of Connecticut.
Am. Jour. Sci. and Arts, XLVIII, 1845, pp. 271-286. Separate.
- *LISTER, MARTIN. Historiæ Animalium Anglicæ, etc., 1678. Appendix to same, 1685.
- *— Historiæ Conchyliorum, etc., 1685.
- *— De Cochleis tam Terrestribus quam Fluvialibus, etc., 1685.
- Dissert. Anat. Tertia, 1685.
- *— Historiæ sive Synopsis Methodicæ Conchyliorum, 1685-1770.
- Conchyliorum bivalvum utriusque, etc., 1690.
- *LLOYD, R. M. On the specific distinctions of Anodonta anatina.
Ann. and Mag. Zool., 1870, p. 65.
- *LOCARD, ARNOULD. Malacologie lyonnaise, etc.
Ann. Soc. Ag. Lyons, 1877. Separate.
- Études sur les variations Malacologiques, etc.
Ann. Soc. Ag. Lyons, 1880, p. 567.
- Catalogue des Mollusques du Département de l'Ain.
Mém. Acad. Lyon, 1881. Separate.
- Catalogue des Mollusques de Lagny.
Ann. Soc. Linn. Lyon, 1881. Separate.
- Distribution et répartition des Mollusques, etc., 1882.
- *— Catalogue général des Mollusques de France, 1882.
- *— Malacologie des Lacs de Tiberiade, etc.
Arch. Mus. Lyon, III, 1883, pp. 192-292, plates.

- *LOCARD, ARNOULD. Description de quelques Anodontes nouveaux, etc., 1884.
- Contributions à la Faune malacologique française.
Ann. Soc. Ag. Lyon, 1884. Separate.
- Description de deux Náyades nouvelles, etc.
Bull. Soc. Rouen, 1885, p. 1.
- Catalogne des espèces françaises, etc.
Ann. Soc. Zool. Lyon, 1889, pp. 111-269.
- Matériaux pour servir à l'histoire malacologique française.
Bull. Soc. Mal. de France, 1889, pp. 283-307.
- *— Révision des espèces françaises, Margaritana et Unio, etc., 1889.
- *— Révision des espèces françaises, Pseudanodonta et Anodonta, 1890, pp. 240.
- *— Les Coquilles des eaux douces et saumâtres de France.
Ann. Soc. Ag. Soc. et Ind., 7th ser., 1893, pp. 5-196. Separate.
- Description de quelques Unionidæ nouveaux, etc.
Bull. Soc. Elbenf, 1894, pp. 49-62.
- Description de deux nouvelles espèces de Pseudanodonta.
Rev. Sci. Bourb., 1894, pp. 64-68, figs.
- *— Coquilles terrestres des eaux douces et saumâtres de Portugal.
Arch. Mus. Hist. Nat. Lyon, VII, 1899, pp. — 303.
- *LORTET, M. Dragages profonds dans le lac de Tiberiade, etc.
Comptes Rendus, 1880, pp. 500-502.
- LOUDON. Remarks on Lea's Observations.
Mag. Nat. Hist., 1835, p. 126.
- *LOVEN, SVEN. Bidrag till Kännedom, etc.
Kongl. Vit. Akad. Hand., 1848, pp. 329-435, 6 plates.
- [On Branchiæ of Mollusca, etc.]
Mém. Acad. de Stockholm, 1848, p. ?.
- *— Berattelse om fram Stegne i Molluskernes, etc., 1852.
- LOW, FRANZ. Die Flussperlenmuscheln-fischerei in der Moldau in Böhmen.
Verh. d. Zool.-bot. Ges. Wien., 1859, pp. 333-336.
- *LUID, EDWARD. Lithophylæi Britannici Iconographia, etc., 1699.
- LYELL, CHARLES. Second Visit to the United States, 1849.
- *— Manual of Geology. Various editions.
- *MABILLE, JULES. Histoire malacologique du bassin parisien, 1870.
- *— Sur quelques Mollusques du Tonkin.
Bull. Soc. Mal. de France, IV, 1887, pp. 73-164, 4 plates.
- *MABILLE, J., and LE MESLE, GEORGES. Observations sur la Faune malacologique de Cochinchine, etc.
Jour. de Conch., 1866, pp. 117-138, 1 plate.
- MCCOY. Descriptions of Palæozoic Fossils, 1855.
- MACE, J. A. Essai d'un Catalogue des Mollusques de Cherbourg et Valognes.
Séanc. Congrès Sci. Fr., 1861, pp. 241-288. Separate.
- MACGILLIVRAY, W. Mollusks of Aberdeen, 1843.
- A History of the Molluscos Animals of Scotland, etc., 1844.
- MCLELLAN, A. Third Report on the Shells of Stirlingshire.
Trans. Stir. Soc., 1894-95, pp. 87-89.
- MACMUNN, C. A. On the Chromatology of the Blood in some Invertebrates.
Quart. Jour. Mic. Sci., 1885, pp. 469-490, 2 plates.
- MALTZAN, HERM. VON. Zur Kenntniss unserer Anodonten.
Arch. d. Vereins f. Nat. Meck., 1869, pp. 170-178.
- Systematisches Verzeichniss der Mollusken, etc.
Arch. d. Vereins f. Nat. Meck., XXI, 1872, pp. 64-95.
- [Anodonta eygnea and piscinalis, etc.]
Arch. d. Vereins f. Nat. Meck., XXI, 1872, pp. 91, 92.
- MAIZINE, F. DE. Essai sur la Faune malacologique du Belgique, etc., 1867.
- MANDUYT. Tableau indicatif des Mollusques du Département de la Vienne, 1839.
- *MANGILI, C. G. Zootomische de Conchiglia Bivalvi, 1804.

- MANSELL-PLEYDELL, J. C. Ornithology and conchology of the county of Dorset, 1874.
- The Land and Fresh-water Mollusca of Dorsetshire.
Proc. Dors. N. Hist. Club, 1885, pp. 76-127, 128-183, 6 plates.
- *MANTELL, GIDEON A. The Geology of the southeast of England, 1833.
- *— On the Unionidæ of the River of the Country of the Ignanodon.
Am. Jour. Sci. and Arts, XLVII, 1844, pp. 402-406.
- *— Geology of the Isle of Wight, etc., 1847.
- *MARAVIGNA, M. C. Mémoires pour servir à l'histoire naturelle de la Sicile, 1838.
- *MARCEL DE SERRES. Géognosie de Terrains Tertiaires, etc., 1829.
- MARQUAND, E. D. The Land and Fresh-water Shells of Guernsey.
Rep. Guernsey Soc., 1894, pp. 356-359.
- MARSH, WILLIAM A. New American Unionidæ.
Aledo Republican, November 18, 1896, Newspaper.
- *— Descriptions of two new species of Unio from Arkansas.
Nautilus, 1891, pp. 1, 2.
- *— Descriptions of two new Unios from Florida.
Nautilus, 1891, pp. 29, 30.
- *— Brief Notes on the Land and Fresh-water Shells of Mercer County, Illinois.
Conch. Exchange, LII, 1887, 1888, pp. 42, etc.
- *MARSHALL, WILLIAM B. Beaks of Unionidæ, etc.
Bull. N. Y. State Mus., II, 1890, pp. 169-181, plate.
- *— Preliminary List of New York Unionidæ.
Bull. N. Y. State Mus., I, 1892, pp. 17.
- MARSHALL, A. M., and HURST, C. H. A Junior Course of Practical Zoology, 1887.
- *MARSIGLI, L. F. Description du Danube, etc., 1749.
- *MARTENS, E. VON. Ueber die Binnemollusken des Mittleren und Südlichen Norwegens.
Mal. Blatt, 1856, pp. 69-117.
- *MARTENS, E. VON. Reisebemerkungen über einige Binnenschnecken Italiens.
Mal. Blatt, 1857, pp. 120-155.
- *— Verzeichniss der von Prof. Peters in Mossambique, etc.
Mal. Blatt, 1859, pp. 211-221.
- *— Die Japanische Binnenschnecken im Leidner Museum.
Mal. Blatt, 1861, pp. 32-61.
- *— On the Mollusca of Siam.
Proc. Zool. Soc., 1860, pp. 61-78.
- *— Mörch, Synopsis Molluscorum Danie.
Mal. Blatt, 1865, pp. 16-26.
- *— Bourguignat, Malacologie du lac des Quatre Cantons.
Mal. Blatt, 1865, pp. 41-45.
- *— Übersicht der Land und Süßwasser Mollusken des Nil Gebietes.
Mal. Blatt, 1865, pp. 177-207; 1866, pp. 1-21, 91-114.
- Ueber die Mollusken Fauna Württemburgs.
Jahrb. Ver. Nat. Wurt., 1865, pp. 178-217.
- *— Ueber die Mexicanischen Binnenschnecken, etc.
Mal. Blatt, 1865, pp. 1-78, 151-153, plate.
- *— Ueber einige Afrikanische Binnenschnecken.
Mal. Blatt, 1866, pp. 91-110, 2 plates.
- *— Ueberblick der Najaden des indischen Archipels.
Mal. Blatt, 1867, p. 10.
- *— Ueber einige Muscheln des Oberen Nil gebietes.
Mal. Blatt, 1867, pp. 17-20.
- *— Die Süßwasser Mollusken Japans.
Arch. für Naturg., 1868, p. 125.
- *— Ueber Südbrasilische Land- und Süßwasser Mollusken, etc.
Mal. Blatt, 1868, pp. 169-217.
- *— Conchologische Notizen.
Mal. Blatt, 1869, pp. 81-84.
- *— Conchylien aus dem Oberen Nil gebiet.
Mal. Blatt, 1870, pp. 32-36.
- *— Unterscheidung der Anodonten.
Nach. Deutsch. Mal. Ges., 1871, p. 138.
- Über das vorkommen von Unio sinnatus, etc.
Sitz. Ber. Ges. Nat. Fr., 1872, pp. 101-103.

- *MARTENS, E. VON. Mollusken an der
Mundung des Jensei.
Nach. Deutsch. Mal. Ges., 1872, pp.
71-73.
- *— Zusammenstellung der von Dr.
Georg Schweinfurth in Afrika, etc.
Mal. Blatt, 1873, pp. 37-46.
- Critical list of the Mollusca of
New Zealand, etc., 1873.
- *— Die Binnenmollusken Venezuelas,
etc.
Festschrift Gesell. Nat. Freunde,
1873, pp. 157-225, 2 plates. Separate.
- Ueber vorderasiatische conchylien,
etc., 1874.
- *— Binnen-Mollusken aus Mittlern
China.
Mal. Blatt, 1875, pp. 185-188; *Sitz.
Ber. Nat. Fr.*, 1875, p. 3; *Proc. Zool.
Soc.*, 1875, p. 606.
- *— *Cristaria reiniana*.
Jahrb. Deutsch. Mal. Ges., 1875, p. 136.
- Über die von Ehrenberg auf seiner
Reise durch Russland, etc.
Sitz. Ber. Nat. Fr., 1875, pp. 88-96.
- *— Über einige neue ostasiatische Land
und Süßwasser Conchylien.
Sitz. Ber. Nat. Fr., 1875, pp. 2-4.
- Die von Prof. Dr. R. Bucholz, in
Westafrika, etc.
Mon. Akad. Wiss. Ber., 1876, pp.
253-274, 5 plates.
- Uebersicht über die von Hilgendorf
und Dönitz in Japans, etc.
Sitz. Ber. Nat. Fr., 1877, pp. 97-123.
- *— Mehrerlei ausländische Conchylien
arten, etc.
Sitz. Ber. Nat. Fr., 1877, pp. 99-105.
- *— Uebersicht der von ihm von 1843
bis 1847 in Mossambique gesammel-
ten Mollusca.
Mon. Akad. Wis. Ber., 1879, pp. 727-
749.
- Über die Vertheilung der Land-
schnecken im Europäischen Russ-
land.
Sitz. Ber. Nat. Fr., 1878, pp. 20-26.
- *— Aufzählung der von Dr. Alex.
Brandt in Russisch Armenien gesam-
melten Mollusken.
Mélanges Biol. St. Petersb., X, 1880,
pp. 379-400; *Bull. Pet. Ak.*, 1880, pp.
142-158.
- [Tubercles on beaks of *Unio cras-
sus* and *tumidus*.]
Sitz. Ber. Nat. Fr., 1881, p. 94.
- *MARTENS, E. VON. Zwei Binnenconchy-
lien aus Angola.
Sitz. Ber. Nat. Fr., 1881, pp. 121-125.
- *— Binnenconchylien aus Angola und
Loango.
Jahrb. Deutsch. Mal. Ges., 1882, pp.
243-250.
- *— Conchologische Mittheilungen, I,
1881; II, 1881-1885.
- *— Einige centralafrikanische Con-
chylien, etc.
Sitz. Ber. Nat. Fr., 1883, pp. 71-74.
- *— Zeigte ferner einige der von Dr.
Gottsche in Japan und Korea, etc.
Sitz. Ber. Nat. Fr., 1885, pp. 76-80.
- Ueber brasilische Land- und Süß-
wasser-Mollusken.
Sitz. Ber. Nat. Fr., 1885, pp. 147, 148.
- Afrikanische Binnenmollusken.
Conch. Mitt., 1881-1885, pp. 181-
190, 1 plate.
- *— Subfossile Süßwasser-conchylien
aus Ägypten.
Sitz. Ber. Nat. Fr., 1886, pp. 126-129.
- *— Vorlegung eines lebenden exam-
plares von *Unio tumidus*.
Sitz. Ber. Nat. Fr., 1887, pp. 105, 106.
- *— Vorlegung einiger Süßwassermu-
scheln aus Guatemala.
Sitz. Ber. Nat. Fr., 1887, pp. 106-108.
- *— Zwei neue Schneckenarten, etc.
Sitz. Ber. Nat. Fr., 1888, pp. 64, 65.
- *— Sudafrikanische Landschnecken.
Sitz. Ber. Nat. Fr., 1889, pp. 160-165.
- *— *Spatha subæquilatera* new species.
Conch. Mitt., 1889, p. 18, fig.
- Mittheilungen über die Fauna von
Ost-Afrika.
Sitz. Ber. Nat. Fr., 1890, pp. 181-184.
- *— Von Dr. F. Stuhlmann auf der Ex-
pedition Emin Pasha's, etc.
Sitz. Ber. Nat. Fr., 1891, pp. 13-18.
- *— Ueber die Süßwasser-mollusken
des malayischen Archipels.
Sitz. Ber. Nat. Fr., 1891, pp. 109-112.
- *— *Unio* an welchen ein Stück von der
Wirbelgegend abgesprengt.
Sitz. Ber. Nat. Fr., 1893, p. 169.
- *— Neue Süßwasser-Conchylien aus
Korea.
Sitz. Ber. Nat. Fr., 1894, pp. 207-217.
- *— Mollusken von Paraguay.
Sitz. Ber. Nat. Fr., 1895, pp. 33-35.
- *— Beschalte Weichthiere Deutsch Ost-
Afrikas, 1897, pp. 308, 7 plates, figures.

- *MARTENS, E. VON. Zoologische Ergebnisse einer Reise in niederländische Ost-Indien, herausgegeben, etc., 1897.
- * — Conchologische Miscellen, III. *Arch. für Naturg.*, I, 1899, pp. 26-48, pls. III-VII.
- * — *Biologia Centrali-Americana. Zoologia, Mollusca*, 1900.
- MARTINI, FREDERICH HEINRICH WILHELM. Abhandlung von den Conchylien der Süßwasser. *Berliner Mag.*, 1767, pp. 113-158, etc.
- Verzeichniss Konchylien, 1773.
- *MARTINI, F. H. W. and CHEMNITZ, J. H. Neues systematisches Conchylien-Cabinet, etc., 10 vols., 1769-1829.
- MARTYN, THOMAS. Universal Conchologist, 1784, 4 vols., with plates.
Another edition entitled, *Figures of Nondescript Shells*, 1789, 2 vols.
- MASSON, C. T. Large-sized Anodonta and Unio.
Naturalist, 1884, p. 112.
- MASSOT. Énumération des mollusques du département des Pyrénées-Orientales.
Bull. Soc. Agr. Pyr. Or., 1877. Separate.
- *MATHERON, PHILIPPE. Catalogue des fossiles du département des Bouches-du-Rhône, 1842.
- *MÂTON, WILLIAM GEORGE. Descriptions of seven new species of Testacea. *Trans. Linn. Soc. Lond.*, 1811, p. 325.
- *MATON, WILLIAM GEORGE, and RACKETT, THOMAS. A Descriptive Catalogue of the British Testacea. *Trans. Linn. Soc. Lond.*, 1807, pp. 17-250, 6 plates.
- *MAWE, JOHN. The Linnaean System of Conchology, etc., 1823.
- MAYO. Lessons on Shells in a Pestalozzian School, etc., 1832, 1846, 1856.
- *MEEK, SETH EUGENE. A list of Fishes and Mollusks collected in Arkansas and Indian Territory in 1894.
Bull. U. S. Fish Com., 1895, pp. 341-349.
- *MENKE, CARL THEODOR. Synopsis Methodica Molluscorum, etc., 1828.
— [Catalogue of the Great Shell collection of Baron Walsburg], Pymont, 1829.
- *MENKE, CARLO THEODORO. Molluscorum Novæ Hollandiæ specimen, etc., 1843.
- * — Ein Süd-africanischer Unio. *Zeits. für Mal.*, 1848, pp. 28, 29.
- * — [Notice of Histoire Naturelle des Mollusques, by Moquin-Tandon.] *Mal. Blatt.*, 1857, pp. 1-16.
- MERMET, C. Histoire des Mollusques dans les Pyrénées occidentales. *Act. Soc. Sc. Pau*, 1843, separate.
- *MERY, L. Remarques sur la Moule des estangs.
Mém. Ac. Sci. Paris, 1710, pp. 408-426.
- *MEUSCHEN, FRED. CHRISTIAN. Schediasma Systematis Testaceorum Museum Geversianum, etc., 1787.
- *MICHAUD, ANDRÉ LOUIS GASPARD. Complément de l'Histoire Naturelle des Mollusques de France, 1831.
- * — Description de quelques espèces nouvelles de Coquilles Fossiles. *Guer. Mag. Zool.*, 1837, pls. 81-85.
- MIDDENDORFF, A. TH. VON. Anodonta herculea n. sp.
Bull. Phys. Math. Acad. St. Pétr., 1848, pp. 303, 304.
- * — Malacozologia Rossica, 2 pts., 1847-49.
- Sibirische Reise, II, 1851, 347 pp., 22 plates.
- *MIGHELS, J. W. Catalogue of the Marine fluviatile and terrestrial Shells of Maine.
Jour. Bost. Soc. Nat. Hist., 1844, pp. 308-344.
- MILACHEVITCH, C. Études sur la faune des Mollusques de Moscow.
Bull. Mosc., 1881, p. 215-241.
- *MILES, MANLY. Catalogue of the Mollusks of Michigan.
Rep. Geol. Surr. Mich., 1860, p. 235.
- MILLER, C. A. Remarks on Unio sayi and Unio camptodon, etc.
Cinn. Quart. Jour. Sci., I, 1874, pp. 244-247.
- MILLER, K. Die Schalthiere des Bodensees.
Schrift. Gesch. Bod., IV, 1873, pp. 123-134, 2 plates; Extract by von Martens in *Jahrb. Deutsch. Mal. Ges.*, 1875.
- * — Die Binnenmollusken von Ecuador, II *Mal. Blatt.*, 1879, pp. 117-199, 12 plates.

- *MILLET, P. A. Mollusques terrestres et fluviatiles dans le département de Maine-et-Loire, 1813, 2d ed., in *Act. Soc. Linn. Bord.*, VI, 1833, pp. 114-139; 3d ed. in same, 1854, separate.
 — Description de deux nouvelles Anodontes.
Mém. Soc. Ag. Ang., 1833, p. 241.
 — Description de quelques nouvelles Mollusques de France.
Mag. Zool., 1843, 2 plates.
- MILNE-EDWARDS, ALPHONSE. Éléments de Zoologie, 1834.
- *MILNES, HERBERT. List of the Land and Freshwater Shells of Derbyshire.
Jour. Conch., 1893, pp. 274-288.
- MINDEN. Über die Flussperlenmuschel, etc.
Schrift. Phys. Ges. Kön., 1867.
- MOBIUS, K. Eine echte Perle von ungewöhnlicher Form und Färbung.
Sitz. Ber. Nat. Fr., 1892, pp. 1-3, fig.
- MOLLENDORFF, O. F. VON. Beiträge zur Fauna Bosniens, Görlitz, 1873, 73 pp., plate.
- * — Zur Binnenmolluskenfauna von Nord China.
Jahrb. Deutsch. Mal. Ges., 1881, pp. 33-43, figs.
- MONCKTON, H. W. Zoological report on Land and Freshwater Shells round Wellington College.
Rep. Welling. Soc., 1888, pp. 52-53.
- *MONTAGU, GEORGE. Testacea Britanica, Natural history of British Shells, marine, land and fresh water, 2 vols., 1803, 16 colored plates, supplement, 1808.
- *MONTEROSATO, ALLERY DI. Note intorno alle Najadi Siciliano.
Naturalista Siciliano, 1896, p. 1, figs.
- MONTFORT, DENYS DE. Conchyliologie systématique, etc., 2 vols., 1808-1810, figs.
- MONTFORT, DENYS DE; ROISSY, FELIX, and LEMAN. Histoire naturelle des Mollusques animaux sans vertèbres et à sang blanc, 6 vols., 1802-1805.
- *MOORE, J. E. S. The Mollusks of the Great African Lakes.
Quart. Jour. Mic. Sci., 1898, pp. 159-204, 4 plates.
- MOORE, R. D., and BUTLER, A. W. Land and Freshwater Mollusca in Franklin County, Ind.
Bull. Brookville Soc., 1885, pp. 41-44.
- *MOQUIN-TANDON, ALFRED. Mémoire sur quelques Mollusques nouveaux pour la faune de Toulouse.
Mém. Acad. Sci. Toulouse, 1843, p. 167. Separate.
- * — Note relative à l'oviducte des Unios et Anodontes.
Jour. de Conch., 1853, pp. 410-411.
- * — Histoire naturelle des Mollusques terrestres et fluviatiles de France, etc., 1855, 2 vols. and atlas.
- MÖRCH, O. A. L. Catalogus Conchyliorum qui reliquit C. P. Kjerulf, 1850, 37 pp., 2 plates.
- Catalogus Conchyliorum quæ reliquit D. Alphonso d'Aguirra et Gadecomes de Yoldi, 1852-3, 252 pp.
- * — Beiträge zur Molluskenfauna Central Amerikas.
Mal. Blatt, 1861, p. 170.
- * — [On Link's Genera of Shells.]
Proc. Zool. Soc., 1862, pp. 226-228.
- * — Synopsis Molluscorum Danicæ, etc.
Land. Forsk. blod. Naturh. For., 1864; *Mal. Blatt*, 1864, p. 16.
- * — Ueber Pinna fluviatilis.
Mal. Blatt, 1865, pp. 110-116.
- * — Land and Fresh-water Mollusca of Greenland.
Am. Jour. Conch., 1868, pp. 25-40, 2 plates.
- * — Land and Fresh-water Mollusks of Iceland.
Am. Jour. Conch., 1869, pp. 41-45.
- * — Catalogue des Mollusques des anciennes colonies danoises du Bengal.
Jour. de Conch., 1872, pp. 303-345.
- *MORELET, ARTHUR. Description des Mollusques du Portugal, 1845, 1 vol., 14 plates.
- * — Testacea novissima insulæ Cubanæ, etc., Pt. I, 1849; II, 1851.
- * — Appendice à la conchyliologie de l'Algérie, etc.
Jour. de Conch., 1851, pp. 351-361.
- * — Catalogue des Mollusques de l'Algérie.
Jour. de Conch., 1853, pp. 280-305.
- * — Séries Conchyliologiques, etc., 1858-1875.
- * — Diagnoses Testarum Indo-sinarum.
Rev. et Mag. Zool., 1862, p. 477.

- *MORELET, ARTHUR. Descriptions de coquilles inédites.
Jour. de Conch., 1864, pp. 155-159, 286-290.
- *— Récifications et additions à la Faune malacologique de l'Indo-Chine.
Jour. de Conch., 1865, pp. 19-23, 225-228.
- *— Description d'espèces appartenant à la Faune malacologique de l'Indo-Chine.
Jour. de Conch., 1866, pp. 62-68.
- *— Diagnoses de coquilles nouvelles de l'Indo-Chine.
Rev. et Mag. Zool., 1866, p. 165.
- *— Mollusques terrestres et fluviatiles. Voyage du Dr. Fred Welwitsch, 1868.
- *— Observations critiques sur quelques Paludines de l'Indo-Chine.
Jour. de Conch., 1869, pp. 403-408.
- Notice sur les coquilles recueillies sur les côtes de l'Abyssinie.
Ann. Mus. Civ. Gén., 1872.
- *— Révision des Mollusques terrestres et fluviatiles du Portugal.
Jour. de Conch., 1877, pp. 242-261.
- *— La Faune malacologique du Maroc.
Jour. de Conch., 1880, pp. 1-83.
- *— Deux Naiades inédites du Tabasco.
Jour. de Conch., 1884, pp. 122-125.
- *— Coquilles terrestres et fluviatiles de l'Afrique Équinoxiale.
Jour. de Conch., 1885, pp. 20-33, plate.
- *MORGAN, J. Mollusques terrestres et fluviatiles du Pérak, etc.
Bull. Soc. Zool. de France, 1885, pp. 353-428, 5 plates.
- MORGAN, J. B. The Land and Freshwater Shells of Montgomeryshire.
Coll. Powys-land Club, XXI, 1887, pp. 103-106.
- *MORICAND, STEFANO. Mémoire sur les coquilles envoyées de Bahia par M. Blanchet.
Mém. Soc. Phys. Gén., 1836, 1 plate; 1st Sup., 1836, 1 plate; 2d Sup., 1839, 1 plate; 3d Sup., 1845, 1 plate.
- Nouveaux Naiades du Brésil.
Rev. et Mag. Zool., 1858, pp. 453-454, 1 plate.
- *— Note sur quelques espèces nouvelles de coquilles terrestres, pp. 50, 3 colored plates. No date. [1833.]
- *MORLET, L. Diagnoses Molluscorum novarum Tonkini.
Jour. de Conch., 1886, pp. 75-78.
- *— Catalogue des Mollusques des environs de Neuf-Brisach, Colmar, et Belfort.
Jour. de Conch., 1871, pp. 34-59. Separate.
- *— Description d'espèces de coquilles recueillies par M. Pavie, au Cambodge.
Jour. de Conch., 1883, pp. 104-110, plate; 1884, pp. 386-403.
- *— Liste des coquilles recueillies au Tonkin, par M. Jourdy.
Jour. de Conch., 1886, pp. 257-295, 4 plates.
- *— Catalogue des coquilles recueillies, par M. Pavie, dans le Cambodge, etc.
Jour. de Conch., 1889, pp. 121-200.
- *— Contributions à la Faune malacologique de l'Indo-Chine.
Jour. de Conch., 1891, pp. 230-254, 1 plate.
- *— Diagnoses Molluscorum in Indo-China collectarum.
Jour. de Conch., 1892, pp. 82-86.
- *— Descriptions d'espèces nouvelles provenant de l'Indo-Chine.
Jour. de Conch., 1893, pp. 153-157.
- *MORRIS, JOHN A. Catalogue of British Fossils, etc., 1843.
- *MORSE, E. S. Observations on the Terrestrial Pulmonifera of Maine, etc.
Journ. Portland Soc. Nat. Hist., I, 1864, pp. 1-63, 10 plates.
- *— Salt and Freshwater Clams.
Am. Naturalist, 1869, pp. 21-26.
- MORTILLET, G. DE. Catalogue des coquilles des environs de Nice.
Bull. Soc. Nat. Savoie, 1851.
- Catalogue des coquilles des Iles Britanniques, etc., 1853.
- *MORTON, SAMUEL G. Organic remains.
Am. Jour. Sci. and Arts, XXIX, 1836, pp. 149-154.
- *MOULINS, CHARLES DES. Catalogue des espèces de Mollusques du département de la Gironde.
Bull. Soc. Linn. Bord., 1827, p. 39; supp., 1829, p. 211.
- *— Description d'une nouvelle Unio du Périgord.
Actes Soc. Linn. Bord., 1830, p. 158; 1833, p. 20.

- *MOULINS, CHARLES DES. Considérations générales, etc.
Actes Soc. Linn. Bord., 1839, p. 3.
- Mollusques à ajouter au Catalogue de la Gironde.
Actes Soc. Linn. Bord., 1851, p. 421.
- *MOUSSON, ALBERT. Die Land- und Süßwasser-Mollusken von Java, 1849, 22 plates.
- *— Coquilles terrestres et fluviatiles recueillies dans l'Orient, 1859, 71 pp.
- *— Coquilles recueillies par M. le Prof. J. R. Roth, dans son dernier voyage en Palestine, 1861.
- *— Notiz über einige von Herrn Gustav Wallis aus dem Nördlichen Südamerika, zurückgebrachte Mollusken.
Mal. Blatt., 1869, pp. 170-189.
- *— Diagnosen neuer Mollusken aus West Marocco.
Mal. Blatt., 1873, pp. 149-157.
- *— Coquilles recueillies par M. le Dr. A. Schläfli en Orient.
Jour. de Conch., 1874, pp. 160.
- Bemerkungen über die von Hrn. Dr. von Fritsch und Dr. Reiu aus West Morocco, etc.
Jahrb. Deutsch. Mal. Ges., 1874, pp. 81-107, 2 plates.
- *— Coquilles recueillies dans le Sud-Ouest de l'Afrique.
Jour. de Conch., 1887, pp. 291-301, plate.
- *— Coquilles recueillies par M. Louis Grasser, dans le bassin de l'Amour.
Jour. de Conch., 1887, pp. 10-32.
- *MÜLLER, OTHO FREDERIC. Vermium Terrestrialium et Fluviatilium, Succincta historia, etc., 2 vols., 1773-4.
- Von zweien wenig bekannter Muscheln.
Beschäftigungen der Berlinischen Gesellschaft naturforschender Freunde, IV, 1779, pp. 55-59, pls. IIIa, IIIb.
- *— Zoologiæ Danicæ Prodomus, seu animalium Danicæ et Norwegiæ, etc., 1776, XXXII, 282 pp.
- *MÜLLER, TH. Synopsis novarum generum, etc., 1836.
- MÜNSTER, GEORGE. Beiträge zur Petri-factenkunde, 1839.
- *MURCHISON, RODERICK IMPEY, and LYELL, CHARLES. On the Tertiary Freshwater formations of Aix, 1821.
Proc. Geol. Soc., 1829, pp. 150-152.
- *MURRAY, ADOLPHUS. Fundamenta Testaceologie, 1771.
- *MUSGRAVE, SAMUEL. Photographic Conchology of the Conchological Miscellany of Hanley. 1863.
- MUSSEY, R. D. Catalogue of the Shells of the Ohio Valley.
Proc. Nat. Inst., 1843, p. 297.
- NANSOUTY, DE. Catalogue des Mollusques du département des Basses Pyrénées, etc.
Bull. Soc. Nat. Toulouse, 1872.
- *NEHRING, A. Einige Notizen über Säugethiere und Flussmusscheln von Piracicaba.
Sitz. Ber. Nat. Fr., 1885, pp. 122-127.
- *— Ueber Najadeu von Piracicaba in Brasilien.
Sitz. Ber. Nat. Fr., 1893, pp. 159-167.
- NELSON, W., and TAYLOR, J. W. Annotated List of the Mollusca of York-shire.
Tr. Yorks. Union, 1891, pp. 65-80.
- NEUMAN, ED. Die Mollusken fauna des Königreichs Sachsen.
Nach. Deutsch. Mal. Ges., 1893, pp. 47-63.
- NEUMANN, A. C. De Anodontarum et Unionum oviducta, 1827.
- NEUMAYR, M. Ueber die Herkunft der Unioniden.
Sitz. Ber. Akad. Wien. Abth., 1889, pp. 5-26, 3 plates; Abstract in *Jour. Roy. Mic. Soc.*, 1889, p. 498, and in *Ann. and Mag. Nat. Hist.*, 1889, p. 372.
- *NEUWYLER, M. Die Generations Organa von Unio und Anodonta, etc.
Nat. Ges. Zur., 1841, pp. 1-324, plates; *Isis*, 1841, pp. 218-220; *Fror. Notizen*, 1841, pp. 134-136.
- *NEVILL, G. List of the Mollusca brought back by Dr. J. Anderson from Yunnan, etc.
Jour. As. Soc. Beng., 1874, pp. 14-41.
- *— Anatomical and Zoological Researches during the Yunnan Expedition, 1877.
Yunnan Expedition, pp. 878-903, 1 plate.
- NEWMANN, JOH. GOTTFR. Land und Wasser Mollusken, 1832, 1833.

- *NEWTON, R. BULLEN. Egyptian newer Tertiary Shells.
Geol. Mag. Lond., VI, 1899, pp. 402-407, pls. XIX, XX.
- *NICKLIN, P. H. Conchological Observations on Lamarck's Naiades.
Trans. Am. Phil. Soc., 1830, p. 395.
- *—Translation of Des Moulin's "Considerations Générales."
Am. Jour. Sci. and Arts, XLI, 1841, p. 104.
- NICOLAS, H. Faune Malacologique Française, etc.
Mém. Acad. Vaucluse, 1891, pp. 139-150, 2 plates.
- NILSSON, SVEN. Historia Molluscorum Sveciæ, etc., 1822, 124 pp.
- *NITSCHKE, H. Die Süßwasser perlen auf der internationalen Fischerie ausstellung in Berlin, 1880.
Nach. Deutsch. Mal. Ges., 1882, pp. 49-64.
- NOLL, C. T. Unsere Flussmuscheln, etc.
Ber. Senck. Nat. Ges., 1870, pp. 33-64.
- *NORDENSKIÖLD, A. E., and NYLANDER, A. E. Finlands Mollusken, 1856.
- NORQUET, DE. Catalogue des Mollusques du Département du Nord, 1872.
- NOULET, JEAN BAPTISTE. Précis analytique de l'Histoire Naturelle des Mollusques dans le bassin sous-Pyrénéen, 1834.
- Coquilles Fossiles d'eau douce du Sud-ouest de la France, 1854.
- Mémoire sur les Coquilles des terrains d'eau douce Sud-ouest de la France, 1868.
- *NYLANDER, OLAFO. Fresh-water Shells of Northeast Maine.
Nautilus, 1897, pp. 9-12.
- OGERION, F. Description d'une Nouvelle espèce d'Anodonta.
Rev. et Mag. Zool., 1861, pp. 115-117.
- *OKEN, LAURENT. Lehrbuch der Naturgeschichte. Pt. 3, Zoologie. 2 vols., 40 plates, 1815, 1816.
Isis, 1817-47, 40 vols.
- *—Auszug aus dem Essai d'un nouveau Système von Schumacher.
Isis, 1826, pp. 706-733.
- [Note on Anodonta Unio and Iridina.]
Isis, 1832, p. 468.
- *OKEN, LAURENT. W. Swainson über Iridina.
Isis, 1834, p. 458; *Oldham Naturalist*, X, 1885, p. 181.
- *OLIVI, G. Zoologia Adriatica, etc., 1792.
- OLIVIER, GUILLAUME A. Voyage dans l'empire Ottoman, l'Égypte et la Perse, 1801-1807, 6 vols. and atlas.
- *OSBORN, HENRY LESLIE. Observations on the parasitism of Anodonta plana, etc.
Zool. Bull., 1898, pp. 301-310, figures.
- *— Observations on the Anatomy of Platyspis, etc.
Zool. Bull., 1898, pp. 55-67.
- OWSJANNIKOW, P. Ueber Blutkörperchen. 1. Die Blutkörperchen... der Teichmuschel.
Bull. Acad. St. Pétersburg, 1895, pp. 372-375, figs.
- *PÆTEL, FR. Catalog der Conchylien-Sammlung, etc., 1890, 3 vols.
- PAGENSTECHE, H. A. Ueber junge Fische in den Kiemen von Unio pictorum.
Ver. Nat. Med. Heid., 1865-1868, pp. 4-6.
- *PALADILHE, M. Nouvelles Miscellanees Malacologiques.
Rev. et Mag. Zool., 1866-1869, pp. 38-41. Separate.
- PANESCORSE, FERDINAND. Prodrome d'histoire naturelle du département du Var, 1853.
- *PARREYSS, LUDWIG. Systematisches Verzeichniss der Oesterreich. Land- und Fluss-Conchylien.
Freunde der Natur., 1850, pp. 97-102.
- Manuscripts, often referred to but never published, according to Bourguignat.
Rev. et Mag. Zool., 1856, p. 74.
- PARTSCH, PAUL. In Jahrbuch der k. k. geologische reichsanstalt, 1837, p. 424.
- PASCAL, LOUIS. Catalogue des Mollusques du département de la Haute-Loire, etc.
Arch. Miss. Sci. and Litt., 1877. Separate.
- PAULUCCI, MME. Matériaux pour servir à l'étude de la Faune Malacologique de l'Italie et de Milan, 1878.
- Faune Malacologica in Calabria, 1879.

- *PAULUCCI, MME. Studio sopra alcune specie del genere Unio.
Bull. Soc. Mal. It., 1879, pp. 107-111.
- PAYAU-DUMOULIN, DE. Notice sur les coquilles dans le département de la Haute-Loire, etc., 1862.
- *PAYRAUDEAU, B. C. Catalogue Descriptif et Méthodique des Annelides et des Mollusques de l'île de Corse, etc., 1826, pp. 174, colored plates.
- *PEARCE, S. S., and MAYFIELD, A. The Land and Fresh-water Mollusca of East Norfolk.
Jour. de Conch., 1894, pp. 391-404.
- PECAGLIA, L. Aggiunte al Catalogo dei Molluschi di Modena, etc.
Bull. Soc. Mal. It., 1894, pp. 113-117.
- PECCHIOLE, V. Unio lardereliani n. sp.
Bull. Soc. Mal. It., 1869, pp. 163-168.
- *PECHAUD, M. J. Anodontes nouvelles de France.
Bull. Soc. Mal. de France, 1884, pp. 189-196.
- PECK, R. H. The minute structure of the gills of Lamellibranchiate Mollusca.
Quart. Jour. Mic. Sci., 1897, pp. 43-66, 4 plates.
- *PELSENER, PAUL. Notice sur les Mollusques recueillis par Capitaine Storms dans la région du Tanganyika.
Bull. Mus. Hist. Nat. Belg., 1886, pp. 101-128, figures; separate.
- PENNANT, Thomas. British Zoology, etc., 1777-1812.
- PERKINS, George H. Molluscan Fauna of New Haven, etc.
Proc. Bost. Soc. Nat. Hist., 1869, pp. 154.
- PERRET, J. B. Catalogue des Mollusques d'Aix, etc., 1834; 2d ed., 1841.
- PETAU. Observations sur des Moules d'étang, dans les coquilles on a trouvé des perles.
Hist. et Mém. Ac. Sci., 1769, pp. 23, 24.
- PETCH, T. The Land and Fresh-water Mollusca of West Norfolk.
Trans. Norf. Soc., 1895, pp. 61-67.
- *PETERS, W. Uebersicht der von ihm von 1843 bis 1847 in Mossambique gesammelten Mollusca, bearbeitet von Hrn. Prof. Dr. E. von Martens.
Monatsber. Akad. Berl., 1879, pp. 727-749.
- *PETIT DE LA SAUSSAYE, S. Sur le nombre des Unionidæ du Nord de l'Amérique.
Jour. de Conch., 1861, pp. 149-151.
- * ——— Note sur le genre Monocondylæa de d'Orbigny, etc.
Jour. de Conch., 1865, pp. 15-19.
- PETIVER, JACOBUS. Gazophylacii Naturæ, etc., 1702-1711.
- *PETTERD, W. F. Contributions for a Systematic Catalogue of the Aquatic Shells of Tasmania, Pt. 1.
Proc. Roy. Soc. Tasm., 1888, pp. 60-83, 4 plates; separate.
- *PFEIFFER, CARL. Naturgeschichte deutschen Land- und Süßwasser-Mollusken, 3 vols., colored plates, 1821-1828.
- *PFEIFFER, L. Schluss der Bemerkungen von Hrn. Alb. Mousson zu seinen "Land- und Süßwasser-Mollusken von Java."
Zeits. für Mal., 1851, p. 46.
- * ——— [Notice of Lea's Synopsis and Observations on the Genus Unio, etc.]
Zeits. für Mal., 1853, p. 61.
- * ——— [Notice of Dupuy's Naiades of France.]
Mal. Blatt, 1854, pp. 253, 264.
- * ——— [Notice of Rossmassler's Iconographie.]
Mal. Blatt, 1854, pp. 230-252.
- * ——— Novitates Conchologicae, 5 vols., 1854-1879, colored plates.
- * ——— [Notice of Drouet's Naiades of France.]
Mal. Blatt, 1855, pp. 188-190.
- * ——— [Notice of Proceedings of the Academy of Natural Sciences of Philadelphia.]
Mal. Blatt, 1865, pp. 58-60.
- PFLUG. Über Perlenmuscheln aus dem Fichtelgebirge.
Ber. Ober. Ges. Nat. Heilk., 1876, pp. 89, 90.
- *PHILIPPI, RUDOLPH ARMAND. Enumeratio Molluscorum Siciliae, etc., 2 vols, 1836-1844.
- * ——— Abbildungen und Beschreibungen neuer oder wenig gekannter Conchylien, 3 vols, 1842-1851, plates.
- * ——— Descriptiones testaceorum chinensium, etc.
Zeits. für Mal., 1845, pp. 161-166.

- *PHILIPPI, RUDOLPH ARMAND. *Centuria altera Testaceorum novarum, etc.*
Zeits. für Mal., 1847, pp. 71-77, 84-96, 113-127; 1848, pp. 99-112, 123-150, 151-176, 186-192; 1849, pp. 123-176.
- *— [Remarks on Mousson's Shells of Java.]
Zeits. für Mal., 1850, pp. 17-21.
- *— Diagnoses Molluscorum Pernanum.
Mal. Blatt, 1869, pp. 32-42.
- *— In Novitates Conchologicæ, 1869.
- *— Bemerkungen über die Chilenischen Unionen.
Mal. Blatt., 1869, pp. 43-49.
- *PHILLIPS, JOHN. Illustrations of the Geology of Yorkshire, 1829; 2d ed., 1835.
- *— Manual of Geology, etc., 1855.
- *PHILIPPSSON, LAURENTIUS MUNTER. *Dissertatio historico naturalis nova Testaceorum, etc.*, 1788.
- PICARD, CASIMIR. *Histoire de Mollusques du département de la Somme.*
Bull. Soc. Linn. Nord, 1840, p. 150.
- *Mémoire sur les déviations dans le genre Unio, etc.*
Bull. Soc. Linn. Nord, 1840, p. 339, 8 plates. Separate.
- PICOT, F. J. *Traité de Paléontologie*, 1853-6, 4 vols.
- PILKINGTON, Mrs. *Goldsmith's Natural History abridged, etc.*, 1810.
- *PILSBRY, H. A. Critical notes on eastern Texas Unionidæ.
Nautilus, 1891, pp. 74-77.
- *— A new species of Arconaia.
Nautilus, 1891, pp. 127, 128; *Naturaliste Canadien*, 1891, pp. 171, 172.
- *— Notes on a collection of Shells from Tabasco, Mexico.
Proc. Acad. Nat. Sci. Phila., 1892, pp. 338-341, plate.
- *— New and unfigured Unionidæ.
Proc. Acad. Nat. Sci. Phila., 1892, pp. 128, 129, 131, 132, 3 plates.
- *— Notes on the genera of Unionidæ and Muteilidæ.
Nautilus, 1893, p. 30.
- *— New species of Fresh-water Mollusks from South America.
Proc. Acad. Nat. Sci. Phila., 1896, pp. 561-565, 2 plates.
- *— Notes on some Southern Mexican Shells.
Nautilus, XIII, 1900, pp. 139-141.
- *PINI, N. Note Malacologiche sulle Fauna Italiani.
Atti Soc. It., 1884, pp. 79-87, figures.
- *PLEAS, E. Shells of Henry Co[unty], Indiana.
Nautilus, 1893, pp. 68-70.
- PLINIUS. *Naturalis Historiæ*, Book IX, A. D. 80.
- *PLUMMER, JOHN T. Scraps in Natural History.
Am. Jour. Sci. and Arts, XLVI, 1844, pp. 236-249.
- Catalogue of Shells of Wayne Co[unty], Indiana.
Am. Jour. Sci. and Arts, XLVI, 1845, p. 93.
- POE, EDGAR ALLEN. *Conchologist's First Book*, 1840, pp. 166.
- *POEY FELIPE. *Memorias sobre la Historia Natural de la Isla de Cuba*, 2 vols., 1851 and 1856-1858.
- *POHLIG, HANS. *Maritime Unionen.*
Palæontographica, 1880. Separate.
- *Unio aus dem Rhein bei Bonn.*
Sitz. Ber. Nied. Ges., 1886, pp. 91, 92.
- *— Die Land-, Süßwasser- und Binnenconchylien des Nördlichen Persien.
Jahrb. Deutsch. Mal. Ges., 1887, pp. 118-121; *Zool. Record*, XXII, p. 18.
- *POIRET, J. L. M. *Reise in die Barbarey, etc.*, 1789.
- *Coquilles dans le département de l'Aisne, etc.*, 1801.
- *POLI, J. X. *Testacea utriusque Siciliae, etc.*, 1791-1827.
- *POLLONERA, CARLO. *Molluschi della Scioa, etc.*
Bull. Soc. Mal. It., 1888, pp. 49-86, 2 plates.
- *Elenco dei Molluschi in Piemonte.*
Mus. Zool. Anat. Comp. Torino, 1889, p. ?.
- *PONTOPPIDON, ERIC. *Histoire Naturelle de la Norwege*, 1753. Translated and published in London, 1755.
- *PORRO, CARLO. *Conchyliarum terrestrium et fluviatiliumque Italiae catalogus*, 1837.
- *Malacologia della provincia Comasca*, 1838.
- *Catalogus Conchyliarum terrestrium fluviatiliumque Europæ*, 1839.
- POSNER, C. *Über den Bau der Najadenkieme, etc.*
Arch. Mik. Anat., 1875, pp. 517-560, 2 plates.

- *POTIEZ, VALÉRY LOUIS VICTOR, and MICHAUD, ANDRÉ LOUIS GASPARD. *Galerie des Mollusques*, etc., 2 vols., 1838, 1844, plates.
- POULSEN, C. M. *Bornholms Land- og Ferskvands Bløddyr*.
Id. Med., 1874, nos. 13, 14, pp. 189-281.
- POULSON, C. A. A Monograph of the Bivalve Shells of the Ohio, 1832. Translated from C. S. Rafinesque.
- POUPART, FRANÇOIS. *Remarques sur les Coquillages*, etc.
Mém. Ac. Sci. Paris, 1706, pp. 51-61.
- PRATT, W. H. List of Shells at Davenport, Iowa.
Proc. Dar. Acad., 1876 pp. 165-167.
- *— Description of a Unio Shell, etc.
Proc. Dar. Acad., 1876, pp. 167, 168, figs.
- *— The Shell Beds of the Vicinity of Davenport.
Proc. Dar. Acad., 1878 pp. 156-162.
- PRATTEN, L. M. *Unio margaritifera*, etc.
Naturalist, London, 1865, p. 220.
- PRESCOTT, W. Catalogue of the Shells of Massachusetts, 1842.
- *PREVOST, J. L. Note sur la génération des Mulettes.
Ann. des Sci. Nat., 1825, pp. 323, 324; 1826, pp. 447-455; *Nouv. Bull. Soc. Philom.*, 1825, p. 78; *Fer. Bull.*, 1825, p. 299.
- *— De la génération chez les Mollusques des peintres.
Mém. Soc. Phys. et d'Hist. Gén., 1825, pp. 57, 58; *Isis*, 1834, p. 857, 858.
Ann. des Sci. Nat., 1826, p. 447.
- PULTENEY, B. Catalogue of Birds, Shells, and Plants of Dorsetshire, etc., 1799; 2d ed., 1813.
- PUSCH, GEORGE GOTTLIEB. *Polens Palæontologie*, 1837.
- PUTON, ERNEST. *Essai sur les Mollusques des Vosges*.
Stat. Dépt. des Vosges, 1847. Separate.
- Note sur l'Unio ater de Nilsson.
Ann. Soc. Anat. Vosg., 1853, p. 4.
- *PUTZEYS, DR. Diagnoses de quelques Coquilles Nouvelles provenant de l'Etat Indépendant du Congo.
Proc. Verb. Séanc. Soc. Mal. Belg., XXVII, 1898, pp. xxv-xxviii, figs.
- *QUATREFAGES, ARMAND DE. *Mémoire sur la vie intra-branchiale des petites Anodontes*.
Ann. des Sci. Nat., 1836, pp. 321-336.
- *— *Embryogénie des Unio*.
Comptes Rendus, 1847, pp. 82-86; 1849, p. 229.
- QUILTER, H. New variety of *Unio tumidus*.
Science Gossip, 1886, p. 68.
- *RABL, CARL. *Über die Entwicklungsgeschichte der Mählermuschel*, etc.
Jen. Zeits., 1876, pp. 310-393. 3 plates. Separate.
- *— *Bemerkungen über den Bau der Najadenkieme*.
Jen. Zeits., 1877, pp. 349-354, 1 plate.
- RAFINESQUE, CONSTANTINE SAMUEL. *Discoveries in Natural History*.
Am. Monthly Mag., III, 1818, p. 354; IV, 1818, pp. 42, 107.
- *— *Prodrome de 70 nouveaux genres d'Animaux découverts dans l'intérieur des Etats-Unis*, etc.
Jour. de Phys. Chim. et His. Nat., 1819, pp. 423-428.
- *— *Monographie des coquilles bivalves fluviales de la Rivière Ohio*. Contenant douze Genres et soixante huit Espèces.
Ann. Gén. Sci. Phys. Brux., 1820, pp. 287-322, 2 plates; separate; also translated by C. A. Poulson.
- *— Continuation of a Monograph of the Bivalve Shells of the River Ohio, etc., 1820.
- *— On eight Asiatic bivalve fluvial shells.
App. to Continuation of Monog., 1831.
- *— Two new bivalve fluvial shells of South America.
Atl. Jour. and Friend, 1832, p. 154.
- *— *Odatelia*, new genus of North American shells.
Atl. Jour. and Friend, 1832, p. 154.
- *— The complete writings of Constantine Samuel Rafinesque on recent and fossil conchology, by William G. Binney and George W. Tryon, 1864.
- *RANG, SANDER. *Manuel de l'histoire naturelle des Mollusques*, etc., 1829, pp. 390, 6 plates.
- *— *Mémoire sur quelques Acéphales d'eau douce du Sénégal*.
Nouv. Ann. Mus., 1835, pp. 297-321, 2 plates; separate.

- RANKIN, W. M. Ueber das Bojanische Organ der Teichmuschel.
Jen. Zeits., 1890, pp. 227-267, 2 plates.
- RASPAIL, F. V. Note sur la parturition vivipare les moules de rivières, 1828.
- Observations microscopiques sur les moules, etc., 1828.
- Mikroskopische Untersuchungen über die Flussmuscheln.
Hensinger Zeits., 1828, p. 118.
- *— Réponse à M. Baer, etc.
Isis, 1829, p. 556.
- Revue zoologique sur la génération des bivalves.
Ann. des Sci. Obs., 1829, p. 107, figs.
- RATHKE, J. Om Dammuslingen.
Skrift. Nat. Selsk. Kiob., 1797, pp. 139-179, plates; *Tillaeg*, 1798, pp. 173-185.
- [On the parasites of Naiades.]
Soc. d'Hist. Nat. Cop., 1797, p. 139.
- *RAVENEL, EDMUND. Catalogue of Recent and Fossil Shells, 1834, pp. 20.
- RAWITZ, B. Die Mantelrand der Acephalen, etc.
Jen. Zeits., 1890, pp. 549-631, 4 plates.
- RAY, JULES, and DROUET, HENRI. Nouvelle Anodonta.
Rev. Zool., 1848, pp. 235-239.
- Notice sur les Anodontes
Rev. Soc. Cur., 1848. Separate.
- Description d'une nouvelle Anodonte.
Rev. et Mag. de Zool., 1849, pp. 29-34. Separate.
- Catalogue des Mollusques de la Champagne, etc.
Rev. et Mag. de Zool., 1851. Separate.
- READ, WILLIAM. Catalogue of the Shells of Massachusetts, 1845.
- *RÉAUMUR, R. A. F. DE. De la formation et de l'accroissement des Coquilles, etc.
Mém. Acad. Sci. Paris, 1709, pp. 364-400, 2 plates.
- *RECLUZ, C. A. Description de deux Coquilles nouvelles.
Rev. Zool., 1842, pp. 305-307; *Mag. de Zool.*, 1843, pt. 77, 78.
- *— Description d'une nouvelle Anodonte.
Jour. de Conch., 1850, pp. 55, 56.
- REES, A. Encyclopedia. Supp., Pt. 1, p. 79. Article Conchology, 1810.
- *REEVE, LOVELL. Conchologia Systematica, 2 vols., 1841, 1842, pp. 522, 300 colored plates.
- *— [Tabula Methodica.]
Proc. Zool. Soc., 1841, pp. 72-77.
- *— Conchologia Iconica, 20 vols., 2,727 colored plates, 1843-1878; completed by Sowerby. Unios in XVI, Anodontas in XVII.
- *— Elements of Conchology, etc., 1860, colored plates.
- *— The Land and Fresh-water Mollusks of the British Isles, 1863, pp. 275, figs.
- REGELSPERGER, G. Mollusques terrestres et de l'eau douce recueillis aux environs de Bern, etc.
Mittheil. Ges. Bern, 1883, pp. 31-42. Separate.
- REINHARDT, O. Verzeichniss der weichtiere der Provinz Brandenburg.
Nach. Deutsch. Mal. Ges., 1886, p. 155.
- REMAK. Ueber ein selbständiges Darmnerven-System, Berlin, 1847.
- RENGARTEN, LUDWIG VON. Dissertation de Anodontæ vasorum systematica, etc., 1854.
- RENIERI, S. A. Tavole alla Classif., 1807.
- *REQUIEN, E. Catalogue des Coquilles de l'île de Corse, 1848.
- *RETZIUS, D. M. ANDR. J. Dissertatio historico naturalis, etc., 1788. See Philippson.
- REYNES, PAUL DE. Lettre à M. Moquin-Tandon sur quelques Mollusques, etc., 1843.
- *RHOADS, SAMUEL. On a collection of Pennsylvania Mollusks from the Ohio River system, etc.
Nautilus, 1899, pp. 133-138.
- RIABININE, J. [The Naiades of the Governments of Kharkoff and Pultova.]
Trudni Khark. Univ., 1887, pp. 159-204, 4 plates.
- *RIGACCI, GIOVANNI. Cataloga della Conchiglie, etc., 1866.
- RIMMER, R. The Land and Fresh-water Shells of the British Isles, etc., 1880, pp. 240.
- ROBERTS, G. Anodonta cygnea and anatina.
Science Gossip, 1888, p. 280.
- *— Observations on the Unionidæ.
Science Gossip, 1888, pp. 127-129.

- *ROCHEBRUNE, A. T. DE. Sur Quelques espèces du Haut-Sénégal.
Bull. Soc. Philom., 1881-82, pp. 33-35.
- *— Documents sur la faune malacologique de la Cochinchine et du Cambodge.
Bull. Soc. Philom., 1882, pp. 35-74; Supp., pp. 99-118.
- *— Sur une nouvelle Unio des Mékong.
Bull. Soc. Philom., 1883, pp. 26-31.
- *— Sur Quelques Lamellibranches nouveaux du Congo, etc.
Bull. Soc. Mal. de France, 1886, pp. 1-14, plate.
- ROEBUCK, W. D. Annotated List of the Mollusca of Lincolnshire.
Naturalist, 1887, pp. 245-272.
- Census of Scottish Land and Fresh-water Mollusca.
Proc. Roy. Phys. Soc. Edinburgh 1889-90 (1891) pp. 437-503.
- *ROEMER, FERDINAND. Texas, 1849.
— Die Kreide von Texas, 1852.
— Die Versteinerungen Oolithengebirges, 1835-6.
— Die Versteinerungen des nord-deutschen Kreidegebirges, etc., 1840-41.
- ROEMER, H. Über Spatha Caillaudi, etc.
Sitz. Ber. Ges. Nat. Freunde, 1875, pp. 21, 22.
- ROGENS, T. On the Land and Fresh-water Shells of Tasmania.
Proc. Manch. Soc., 1880, pp. 101-103.
- *ROGERS, WILLIAM B. Analyses of Shells.
Am. Jour. Sci. and Arts, XXVI, 1834, pp. 361-365.
- ROSSY. See Montfort Denys.
- *ROLLESTON, GEORGE. On the Aquiferous and Oviducal System in the Lamellibranchiate Mollusks.
Ann. Nat. Hist., 1862, pp. 446-458.
- RONDELETIUS, Gulielmus. Universa Aquatiliu Historia, 1554, 1555.
- *— Histoire entière des Poissons, 1558.
- ROSEN, O. Essai d'une description de la faune Malacozoologique Transcaspienne, etc.
Cong. Intern. Zool., 1892. II. Pt. 2, pp. 171-178.
- *ROSSMÄSSLER, E. A. Iconographie der Land- und Süßwasser-Mollusken, etc., 1835; not yet completed.
- ROSSMÄSSLER, E. A. Ueber Unio Mulleri.
Arch. der Freunde Meck., 1851, p. 201.
- *— Bemerkungen über die Europäischen Najaden.
Zeits. für Mal., 1853, p. 10.
- *— Ueber eine Fauna Molluscorum Extramarinorum Europæ, etc.
Zeits. für Mal., 1853, p. 33.
- *— Unio littoralis.
Zeits. für Mal., 1853, pp. 92, 93.
- *— [Notice of Pt. 2 of Études sur les Náyades de France, by Drouet.]
Mal. Blatt., 1857, pp. 225-228.
- *ROTH, J. R. Spicilegium Molluscorum Orientalium, etc.
Mal. Blatt., 1855, pp. 17-58, 2 plates.
- RUSCHENBERGER, W. S. W. Natural History. Book 4, 1850. Conchology, September, 1850.
- RUSSELL, JOHN LEWIS. Familiar notice of some of the Shells of Essex County, Massachusetts.
Jour. Essex Co. Nat. His. Soc., 1839, pp. 49-76.
- Conspectus of Shells in Essex County, etc.
Jour. Essex Co. Nat. His. Soc., 1852, pp. 126-133.
- *SABATIER, A. Anatomie de la Moule commune.
Ann. des Sci. Nat., 1877, pp. 1-132, 9 plates.
- *SAENGER, EDWARD B. The Fresh-water Shells of Cooper's Creek, Australia.
Am. Naturalist, 1883, p. 1184, fig.
- SAGER, ABRAHAM. Report of Dr. Sager, zoologist of the Geological Survey of Michigan, 1899.
- SAMMLER, C. F. Dei Flussperlemuschel.
Mitth. Voigt. Naturk., 1870, pp. 19-39.
- *SAMPSON, F. A. Shells of Pettis County, Missouri.
Bull. Sedalia Soc., 1885, pp. 16-28.
- *SANDBERGER, C. L. FRIDOLIN. Die Conchylien des Mainzer Tertiärbeckens, 1863, pp. 339, 35 plates.
- *— Ueber Unio sinnatus Lamarck, etc.
Mal. Blatt., 1873, pp. 95-99.
- *SARGENT, H. E. Annotated list of the Mollusca of Clearwater, Wright County, Minnesota.
Nautilus, 1895, pp. 87-90, 125-128.

- SARS, G. O. Bidrag til kundskaben om Norges Arktiske Fauna, 1878, 466 pp., 52 plates.
- SARS, MICHAEL. Beskriveleise og jagt-lagelser Bergen, 1825.
- SATTER, HANNS. Verbreitungs- u. Verwandtschaftsverhältniss der Unioniden Kärntens.
Jahres. d. Acad. Naturwiss. Fer., 1879, pp. 56-62.
- *SAVIGNY, MARIE J. C. DE. Iconographie des Description de l'Égypte Mollusques 1813?, 14 plates.
- SAY, LUCY W. Terrestrial and Fluvatile Shells, 1840.
- *SAY, THOMAS. Article Conchology in the American edition of Nicholson's Encyclopedia, 1st ed., 1816; 2d ed., 1818; 3d ed., 1819. Separate as Descriptions of Land and Fresh-water Shells of the United States.
- * — Description of a New Genus of Fresh-water Bivalve Shells.
Jour. Acad. Nat. Sci. Phila., 1818, p. 454.
- Narrative of an Expedition to the source of St. Peter's River, etc., under Stephen Long.
Long's Exp., II, 1824, pp. 256-265.
- * — Description of some New Fresh-water and Land Shells of the United States.
Jour. Acad. Nat. Sci. Phila., 1825, pp. 118-131.
- Descriptions of new Terrestrial and Fluvatile Shells of North America.
New Harmony Disseminator, 1829, pp. 291-293, 308, 309, 323, 324, 339-341, 355, 356, January 15, 1831; January 29, 1831.
- * — Descriptions of several New Shells, etc.
Transylvania Journal, 1831, pp. 525-528. Separate.
- * — American Conchology, etc. Parts 1-6, 1830-1834, no pagination; Part 7, no date [1834?], issued by Conrad.
- Catalogue of Exotic Shells in my Cabinet, 1833.
- * — An attempt to exhibit the Synonymy of the Western North American Species of Unio and Alasmodonta, 1834.
- The complete writings of Thomas Say on the Conchology of the United States; edited by W. G. Binney, 1853, pp. vi, 252, 75 plates.
- *SAYN, GUSTAVE. Catalogue de Mollusques du département de la Drome.
Bull. Soc. Mal. de France, 1889, pp. 67-81.
- *SCACCHI, A. Catalogus Conchyliorum Regni Neapolitani, etc., 1836; 2d ed., 1857.
- *SCHARDT, M. LE DR. Coquilles de trois Unio batavus, trouvés vivant dans le lac Léman.
Arch. Sci. Nat., 1887, pp. 220, 221.
- SCHARFF, R. F. The Irish Land and Fresh-water Mollusca.
Irish Nat., I, pp. 45-47, etc.; separate.
- ? Vorlage eines Wasserläufers welcher an einem Bein von einer grossen Anodonta, etc.
Sitz. Ber. Nat. Fr., 1888, pp. 121, 122.
- * — Some remarks on the Distribution of British Land and Fresh-water Mollusca.
Conchologist, 1892, pp. 1-6.
- SCHEPMAN, M. M. Zoological Researches in Liberia.
Notes Leyden Mus., 1888, pp. 245-252, plate.
- * — A new Species of Unio.
Notes Leyden Mus., 1891, pp. 113-114, figure.
- * — The Mollusca of the Dutch Scientific Borneo Expedition, etc.
Notes Leyden Mus., 1895, pp. 145-162, plates.
- * — Note on Unio infrarostratus.
Notes Leyden Mus., 1896, p. 140.
- * — Description of a New Unio.
Notes Leyden Mus., 1896, pp. 259, 260, figure.
- * — Description of a New Unio from Borneo.
Notes Leyden Mus., XX, 1898, p. 92, pl. I, figs. 1, 2.
- *SCHIERIOLZ, CARL. Zur Entwicklungsgeschichte der Teich- und Flussmuscheln, 1878, pp. 71, 3 plates.
Abstract in *Zeit. Wiss. Zool.*, 1878, p. 492.
- * — Entwicklung der Unioniden
Denk. Kais. Akad. Wiss., 1888, pp. 183-214; abstract in *Anz. Akad. Wien*, 1888, pp. 126.
- SCHLAEFLER. Coquilles dans l'Orient, 1859.

- SCHLICHTER, H. Einiges über *Anodonta mutabilis*, etc.
Jah. Ver. Nat. Württ., 1886, pp. 348-350.
- SCHLOTTERBECK, PHIL. JAC. *Observatio Physicæ Cochleis quibusdam*, etc.
Act. Helveticus, 1762, pp. 275-286, plate.
- *SCHMIDT, ADOLF. Beiträge zur Malakologie, 1857.
- SCHMIDT, ED. O. Beiträge zur Anatomie und Physiologie der Naiden.
Muller, Archiv., 1846, pp. 406-420.
- SCHMIDT, FERD. J. Conchylien-Sammlungen, 1818.
- Einige Beobachtungen über die verschiedenen in Krain, etc.
Bull. Soc. Imp. Mosc., 1840, pp. 430-453.
- SCHMIDT, F. Vorläufiger Bericht über untersuchungen der postembryonalen Entwicklung von *Anodonta*.
Sitz. Ber. Ges. Dorp., 1885, pp. 303-307.
- Beitrag zur Kenntniss der postembryonalen Entwicklung der Naiden.
Arch. für Naturg., 1885, pp. 201-234, plates; Abstract in *Jour. Roy. Mic. Soc.*, 1885, p. 222.
- *SCHOLTZ, H. Schlesiens Land- und Wasser-Mollusken, etc., 1843; Supp., 1853.
- *— Neueste Beobachtungen im Gebiete der Molluskenfauna Schlesiens.
Zeits. für Mal., 1845, pp. 97-110.
- SCHREIBER, KARL VON. Versuch einer vollständigen Conchylien, etc., 1793, 2 vols.
- *SCHRENCK, LEOPOLD V. Übersicht der Land- und Süßwasser-Mollusken Livlands, 1847.
Bull. Soc. Imp. Mos., 1848, p. 135.
- *— Reisen und Forschungen im Amurlande, etc., 1867, pp. 976, 17 plates.
- SCHROCKINGER, J. VON. Oesterreich'schäuse-tragende Bauch für See- und Muschelthiere.
Verh. Zool. Ges. Wien, 1865, pp. 303-324.
- SCHRÖDER, E. G. C. *Dissertatio de natura*, etc., 1761.
- SCHRÖD, H. Ueber der Bourgnigat'sche methode der messung der Acephalen.
Schr. Ver. Harz, 1888, pp. 11-18.
- *SCHRÖDER, RICHARD. Notice sur quelques Unionides, etc.
Bull. Soc. Mal. de France, 1885, pp. 209-230.
- *SCHRÖTER, JOHANN SAMUEL. Versuch einer systematischen Abhandlung über die Erdkonchylien, etc., 1771-1779.
- *— Die Geschichte der Flussconchylien, etc., 1779.
- *— Einleitung in die Conchylien-Kenntniss nach Linne, 1783-1786, 3 vols.
- *SCHUMACHER, FREDERIC CHRISTIAN. *Essai d'un nouveau Système des Habitations des vers Testacés*, 1817.
- *SCHWEIGER, A. F. Handbuch der Naturgeschichte, etc., 1820, pp. 776.
- *SCOPOLI, JOANNES ANT. *Introductio ad Historiam Naturalem*, etc., 1777.
- SCOTT, T. Notes on the Mollusca of Greenock, etc.
Proc. Nat. Hist. Soc. Glasgow., 1887, pp. 279-285.
- *SEBA, ALBERTUS. *Locupletissimum naturalium Thesauri*, etc., 1734, 4 vols.
- SEMMLER, C. F. Die Flussperlenmuschel *Unio margaritifera*.
Mitth. Voigt. Ver. Reich., 1870, p. 19.
- *SERVAIR, GEORGES. *Histoire Malacologique du Lac Balatan*, etc., 1881, pp. 125, 8 plates.
- *— *Histoire des Mollusques Acéphales des environs de Frankfort*, 1882, pp. 68.
- *— *Unios et Anodontas du Lac de Zurich*.
Bull. Soc. Mal. de France, 1885, pp. 323-352.
- *— *Histoire Malacologique du Lac de Grandlien*, etc.
Bull. Soc. Mal. de France, 1887, pp. 237-266.
- *— *Aperçu sur la Faune des Mollusques des environs de Hambourg*.
Bull. Soc. Mal. de France, 1888, pp. 287-340, 2 plates.
- *— *Des Acéphales du Système Européen*.
Bull. Soc. Mal. de France, 1890, pp. 281-323, 3 plates.
- *SGANZIN, VICTOR. *Catalogue des Coquilles trouvés aux îles de France, de Bourbon, et de Madagascar*.
Mém. Soc. Mus. Strasb., 1840-1846, pp. 1-30.

- SHAEFFER, DAVID H. A complete list of Land and Fresh-water Shells in the Vicinity of Cincinnati. No date.
- *SHARP, BENJAMIN. Remarks on the phylogeny of the Lamellibranchiata. *Proc. Acad. Nat. Sci. Phila.*, 1888, pp. 121-124.
- *SHARSWOOD, WILLIAM. On the discovery of the sensitiveness to light possessed by Unios. *Am. Jour. Sci. and Arts*, XLIX, 1870, pp. 422, 423.
- *SHELDON, D. S. [Shells new to the Davenport List.] *Proc. Dav. Acad. Nat. Sci.*, 1877, p. 143.
- SHEPPARD, MRS. On the recent Shells which characterize Quebec and its environs. *Trans. Lit. Hist. Soc. Quebec*, 1829, pp. 188-198.
- *SHEPPARD, REVETT. On two new species of *Mytilus*. *Trans. Linn. Soc. Lond.*, 1820, pp. 83-87.
- *— Description of seven new British Shells. *Trans. Linn. Soc. Lond.*, 1823, p. 148-170.
- *SHIMEK, B. *Margaritana hildrethiana*. *Conch. Exchange*, 1888, p. 114.
- *SHORT, C. W., and EATON, H. H. Notes on Western Botany and Conchology. *Transylvania Journal of Medicine*, 1831, pp. 74-82.
- *SHRUBSOLE, G. W. On the erosion of Freshwater Shells. *Jour. of Conch.*, 1886, pp. 66-71.
- SHUMARD, B. F. A catalogue of Fluvial Shells belonging to the family of Naiades, of Jefferson County, Ky. *Nat. and Jour. Ag., etc.*, 1846, pp. 105, 106.
- SHUMARD, G. C. Notes of additional species of shells. *Exploration of Red River of Louisiana*, by R. B. Marcy, 1854, pp. 244, 245.
- SHUTTLEWORTH, ROBERT JAMES. Ueber die Land- und Süsswasser-Mollusken von Corsica. *Mitth. Nat. Ges. Bern*, No. 2, 1843, pp. 9-21; No. 3, p. 17.
- Diagnosen neuer Mollusken. *Mitth. Ges. Bern*, 1852, pp. 137-146, 193-208, 289-304.
- *SIBBALD, ROBERTS. *Scotia Illustrata*, etc., 1634.
- SIEBOLD, C. TH. E. VON. [On sexuality in mussels.] *Arch. für Naturg. (Müller's)*, 1837, p. 51.
- Ueber den unterschied der Schalenbildung, etc. *Arch. für Nat. (Müller's)*, 1837, pp. 415, 416.
- Observations sur un organe énigmatique. *Arch. für Nat. (Müller's)*, *Ann. des Sci. Nat.*, 1838, p. 319.
- SIEMASCHKO, MARC VON. Bemerkungen über einige Land- und Süsswasser-Mollusken Russlands. *Bull. Phys. Math. Ac. Pet.*, 1849, pp. 225-240.
- SILVA Y CASTRO, J. Contributions à la Faune Malacologique du Portugal. *Jour. Sci. Lisbon*, 1883, pp. 121-152.
- *SIMPSON, C. T. Contributions to the Mollusca of Florida. *Proc. Dav. Acad.*, 1887-89 pp. 45-72, 63*-72*.
- *— Notes on some Indian Territory Shells. *Proc. U. S. Nat. Mus.*, 1888, pp. 449-454.
- *— On the means of distribution of Unionidæ in the Southeastern United States. *Nautilus*, 1891, pp. 15-17.
- *— Notes on Unionidæ. *Nautilus*, 1891, pp. 86-88.
- *— Notes on the Unionidæ of Florida and the Southeastern States. *Proc. U. S. Nat. Mus.*, 1892, pp. 405-436, 26 plates.
- *— On a Revision of the American Unionidæ. *Nautilus*, 1892, pp. 78-80.
- *— A Review of von Ihering's Classification of the Unionidæ and Mutelidæ. *Nautilus*, 1893, pp. 17-21.
- *— On some Fossil Unios and other Shells from the Drift at Toronto, etc. *Proc. U. S. Nat. Mus.*, 1893, pp. 591-595.
- *— On the Relationships and Distribution of the North American Unionidæ, etc. *Am. Naturalist*, 1893, pp. 353-358.
- *— A new Anodonta, etc. *Nautilus*, 1893, pp. 134, 135.

- *SIMPSON, C. T. Types of Anodonta dejecta rediscovered.
Nautilus, 1894, pp. 52, 53.
- *— On the validity of the genus *Margaritana*.
Am. Naturalist, 1895, pp. 336-344.
- *— Note on *Unio oregonensis*.
Nautilus, 1895, pp. 116-118, 130.
- *— *Unio ochraceus* and *cariosus*.
Nautilus, 1895, p. 122, figs.
- *— Description of Four New Triassic Unios, etc.
Proc. U. S. Nat. Mus., 1896, pp. 381-385, figs.
- *— On the Mississippi Valley *Unio*-nidae found in the St. Lawrence and Atlantic Drainage Areas.
Am. Naturalist, 1896, pp. 379-384.
- *— The Classification and Geographical Distribution of the Pearly Fresh-water Mussels.
Proc. U. S. Nat. Mus., 1896, pp. 295-343, map.
- *— *Unio mitchelli*, Simpson, new species. In Diagnoses of New Mollusks from the Survey of the Mexican Boundary, by W. H. Dall.
Proc. U. S. Nat. Mus., 1896, p. 5, 6.
- *— [Notes upon *Unio* and *Anodonta*.] In Report on the Mollusks collected by the International Boundary Commission, [etc.] by W. H. Dall.
Proc. U. S. Nat. Mus., 1896, pp. 370-374, pl. 32, figs. 1-5.
- *— Notes on the Classification of the Unios.
Nautilus, 1897, pp. 18-23.
- *SIMPSON, GEORGE B. Anatomy and Physiology of *Anodonta fluviatilis*.
Ann. Rep. N. Y. State Mus., XXXV, 1884, pp. 169-191, 11 plates.
- SLAVIK, ALFR. Monographie der Land- und Süsswasser Mollusken Böhmens.
Arch. für Naturg. Landes Böh., 1869, pp. 79-132, 5 plates.
- SŁOSARSKI, A. Matériaux pour la Faune Malacologique du royaume de Pologne.
Bull. Soc. Zool. de France, 1876, pp. 291-299.
- SMITH, EDGAR A. Mollusca in: The Zoology of the Voyage of H. M. S. *Erebus* and *Terror*, etc., 1874.
- *SMITH, EDGAR A. On the Shells of Lake Nyassa, etc.
Proc. Zool. Soc., 1877, pp. 712-722, 2 plates. See *Nach. Deutsch. Mal. Ges.*, 1878, pp. 85-86.
- *— Diagnosis of New Shells from Lake Tanganyika, etc.
Ann. Nat. Hist., 1880, pp. 425-430.
- *— On the Shells of Lake Tanganyika and the neighborhood of Ujiji, etc.
Proc. Zool. Soc., 1880, pp. 344-352.
- *— On the Fresh-water Shells of Australia.
Jour. Linn. Soc. London, 1882, pp. 255-317, 3 plates.
- *— On a collection of Shells from the Solomon Islands.
Proc. Zool. Soc., 1885, pp. 588-609.
- *— Note on *Anodon herculeus* Gerstford.
Jour. of Conch., 1885, p. 366.
- *— On the Shells of the Albert Nyanza, obtained by Dr. Emin Pasha.
Proc. Zool. Soc., 1888, pp. 52-56, figs.
- *— Diagnoses of New Shells from Lake Tanganyika.
Ann. and Mag. Nat. Hist., 1889, pp. 173-175.
- *— List of Land and Fresh-water Shells collected by Dr. Emin Pasha in Central Africa, etc.
Ann. and Mag. Nat. Hist., 1890, pp. 146-168, 2 plates.
- *— On the Mollusca of British Central Africa.
Proc. Zool. Soc., 1891, pp. 309, 310.
- *— Notes on African Mollusca.
Ann. and Mag. Nat. Hist., 1891, pp. 317-324.
- *— On the Shells of the Victoria Nyanza or Lake Oukeréwé.
Ann. and Mag. Nat. Hist., 1892, pp. 121-128, plate.
- *— On a collection of land and fresh-water shells from H. H. Johnston, from British Central Africa.
Proc. Zool. Soc., 1893, pp. 632-641, 1 plate.
- A list of the land and fresh-water Mollusca collected by J. W. Gregory in East Africa, etc.
Proc. Mal. Soc. London, 1894, pp. 163-168, figs.

- *SMITH, EDGAR A. Description of *Unio pahangensis* n. sp. from the River Pahang.
Proc. Mal. Soc. Lond., III, 1899, pp. 315-316, fig.
- *SMITH, HUGH M. The Pearl-Button Industry of the Mississippi River.
Scientific American, August 5, 1899, pp. 86, 87, figs. Separate.
- *— The Mussel Fishery and Pearl-Button Industry of the Mississippi River.
Bull. U. S. Fish Com., 1898 (1899), pp. 289-314, 31 plates.
- SMITH, SIDNEY. Large pearl in *Unio margaritifera*.
Naturalist, London, 1866, p. 133.
- SOLANDER, DANIEL. Manuscripts in the British Museum and unpublished.
- SOLEILLET, P. Mollusques Terrestres et Fluviales recueillis par P. Soleillet dans son voyage au Choa.
In *Mollusques Terrestres et Fluviales au Choa*, by J. R. Bourguignat, pp. 39-44, 1885.
- *SOLLAS, W. J. On the Origin of Fresh-water Faunas.
Sci. Trans. Roy. Soc. Dublin, 1884, pp. 87-118.
- SORDELLI. [On a pearl from *Anodonta piscinalis* containing a small insect.]
Bull. Mal., V, 1872, p. 12, figs.
- *SOURBIEU, EDMOND. Espèces nouvelles pour la Faune française.
Bull. Soc. Mal. de France, 1887, pp. 233-236.
- SOWERBY, JAMES. Mineral Conchology, 1812.
- *SOWERBY, GEORGE BRETTINGHAM. The Genera of Recent and Fossil Shells, 1821-1834? Undated, plates and pages unnumbered.
- *— Description of a new species of *Iridina*.
Zool. Jour., I, 1824, p. 53.
- *— Some observations on the Lamarckian Naiades, etc.
Zool. Jour., I, 1824, pp. 53-55.
- Catalogue of Shells in the Tankerville Collection, 1825.
- *— A Conchological Manual, 1839; 2d ed., 1842.
- Illustrated index of British Shells; containing figures of all the recent species, with names and other information, London, 1859; 2d ed., 1887.
- *SOWERBY, GEORGE BRETTINGHAM.
Continuation of Reeve's *Conchologia Iconica*, XVI, XVII, 1865-1870.
- SOWERBY, JAMES DE CARLE. Mollusca in Richardson's *Fauna Borcali Americana*, III, 1836.
- SPENGLER, LORENZ. In *Schrift. der Berlin. Ges. Nat. Fr.*, 1793, III.
- *— *Skrivter af Naturhistorie Selskabet*, 5 vols., 1790-1799.
- *SPINELLI, GIO BATTISTA. Catalogo dei Molluschi della provincia Brescina, 1856.
- *— Catalogo dei Molluschi in Venezia, etc., 1869.
- *SPIX, J. B., and WAGNER, J. A. Testacea Fluvialia Braziliana, etc., 1827.
- *SQUYER, HOMER. List of Shells from Mingsville, Montana.
Nautilus, 1894, pp. 63-65.
- *STABILE, GIUSEPPE. Fauna Elvetica delle Conchiglie del Luganese, 1846, pp. 68, 3 plates.
- *— *Prospetto Molluschi di Lugano*, 1859.
- STANDEN. Lancashire Land and Fresh-water Mollusca.
Naturalist, 1887, pp. 155-176. Separate.
- *STARK, JOHN. Elements of Natural History, 2 vols., 1828.
- STATUTI, A. Catalogo dei Molluschi nella provincia Romana.
Bull. Soc. Mal. It., 1882, pp. 1-128.
- *STEARNS, R. E. C. Rambles in Florida.
Am. Naturalist, 1869, pp. 397-405.
- *— On the History and Distribution of the Fresh-water Mussels, etc.
Proc. Cal. Acad. Sci., 1882, pp. 1-21, 2 plates.
- *— On the Shells of the Colorado Desert, etc.
Am. Naturalist, 1883, pp. 1014-1020, fig.
- *— List of North American Land and Fresh-water Shells received from the Department of Agriculture.
Proc. U.S. Nat. Mus., 1891, pp. 95-106.
- *— Notes on recent collections of North American Shells from the Department of Agriculture.
Proc. U. S. Nat. Mus., 1894, pp. 743-755.
- *STEFANESSEN, SABBA. Études sur les Terrains tertiaires de Roumanie.
Mém. Soc. Géol. Fr., 1896, pp. 147, 10 plates.

- * STEIN, J. P. E. Frederick. Die lebenden Schnecken und Muscheln, etc., 1850, 120 pp., 3 plates.
- СТЕПАНОВЪ, П. Описаніе МЯГКОТ, etc., 1870.
- [Über den Parasitismus der Najaden] 1871.
- * STERKI, V. Die Mollusken der Umgebung von New Philadelphia, Ohio. *Nach. Deutsch. Mal. Ges.*, 1886, pp. 10-25, 54-58.
- * — Zur Fauna von Ohio. *Nach. Deutsch. Mal. Ges.*, 1887, pp. 178-184.
- * — A byssus in Unio. *Nautilus*, 1891, pp. 73, 74, 90, 91.
- The Land and Fresh-water Mollusca in the vicinity of New Philadelphia, Ohio, 1894, 14 pp.
- * — Some notes on the Genital Organs of Unionidae, etc. *Nautilus*, 1895, pp. 91-94.
- * — Some observations on the Genital Organs of Unionidae, etc. *Nautilus*, 1898, pp. 18-21, 28-32.
- * — Anodonta imbecillis hermaphroditie. *Nautilus*, 1898, pp. 87, 88.
- * STIMPSON, WILLIAM. Shells of New England, etc., 1851, pp. 56, 2 plates.
- STOLICZKA, F. Palaeontologia Indica, Cretaceous Fauna of Southern India, III. The Pelecypoda. 1871, 538 pp., 50 plates.
- STOLTERFOHT, JO. JAC. Unilogia physico-medica, 1700. 6 vols.
- STREBEL, H. Ein Beitrag zur Fauna Mexikanischer Land- und Süßwasser-Conchylien. *Abh. Ver. Hamb.*, VI, 1874, pp. 69.
- STRICKLAND, HUGH E. A List of Some Fresh-Water and Land Shells of Hanley-on-Thames. *Mag. Nat. Hist.*, 1834, p. 494.
- * STROBEL, PELLEGRINO. Materiali per una Malacostatistica de Terra e di Acqua dolce dell'Argentina Meridionale, 1874.
- * STRODE, W. S. Mollusks of Thompson's Lake, Illinois. *Nautilus*, IV, 1891, pp. 133, 134.
- * — Mollusks of Spoon River, Illinois. *Nautilus*, V, 1891, pp. 61-63.
- * — Destruction of Anodonta corpulenta, etc. *Nautilus*, 1891, pp. 89-80.
- * STRODE, W. S. The Unionidae of Spoon River, etc. *Am. Naturalist*, 1892, pp. 495-501.
- * — Unios, very large. *Nautilus*, 1896, p. 115.
- * — Unio collecting. *Nautilus*, 1898, p. 8.
- * STRUBELL, BRUNO. Neue Süßwasser-Conchylien aus Sumatra und Java. *Nach. Deutsch. Mal. Ges.*, 1897, pp. 8-12.
- STUDER, S. Systematisches Verzeichniss Schweizer Conchylien, etc. *Nat. Anz. Allg. Sch.*, 1820, pp. 83-94.
- STUHLMANN, DR. FRANZ. Berichte über eine mit Unterstützung der Königlich-n Reichlichen Reise nach Ost Afrika, etc. *Kgl. Preuss. Akad. Wiss.*, 1888, pp. 801-815; 1889, pp. 451-466.
- Mit Emin Pasha ins Herz von Afrika. (Berlin, Dietrich Reimer), p. 731. [Notiz von Prof. Dr. E. von Marten's Über die Vertheilung der Süßwasser-Mollusken in Central-Africa, 1894.]
- STUMP, F. C. Dissection of the Fresh-water Mussel. *Tr. Manchester Micr. Soc.*, 1888, pp. 85-86.
- STURANY, DR. RUDOLPH. Ueber die Mollusken Fauna Centralafrikas, etc. In *Durch Massailand* by E. Bauman. 1894, pp. 297-322, 2 plates.
- STURM, JAKOB. Deutschland Fauna in Abbildungen nach der Natur mit Beschreibungen. 8 vols., 1803-1809, colored plates.
- * STUSSINER, J. Malakologische Ergebnisse auf Streifzügen in Thessalien, etc. *Jahrb. Deutsch. Mal. Ges.*, 1886, pp. 42-73, 2 plates.
- * STUTCHBURY, SAMUEL. On a new genus of Fossil Shells. *Ann. Nat. Hist.*, VIII, 1842, pp. 481-485, 2 plates.
- SULLIVANT, JOSEPH. Catalogue of Shells and Minerals, etc., 1838.
- * SURBECK, GEORG. Die Molluskenfauna des Vierwaldstättersees. 1899. *Revue Suisse de Zoologie*, VI, pp. 429-552, 2 plates and map.
- SUTER, H. Beiträge zur Schweizerischen Mollusken-Fauna. *Mal. Blatt*, 1889, pp. 1-26.

- *SUTER, H. Liste synonymique et bibliographique des Mollusques terrestres et fluviatiles de la Nouvelle Zélande. *Jour. de Conch.*, 1893, pp. 220-293, plate.
- * — Check-list of the New Zealand Land and Fresh-water Mollusca. *Trans. New Zealand Inst.*, 1894, pp. 139-154.
- *SWAINSON, WILLIAM. A Catalogue of the Rare and Valuable Shells of the collection of Mrs. Bligh, etc. Appendix, pp. 58. Appendix, pp. 20, 2 plates, [1822?]
- * — Zoological Illustrations, 6 vols.: 1st ser., 1820-1823; 2d ser., 1829-1833.
- Remarks on Iridina, etc. *Phil. Mag.*, 1823, pp. 112, 113; *Isis*, 1834, p. 458.
- * — The Elements of Modern Conchology, etc., 1835.
- * — A Treatise on Malacology, etc., 1840.
- * — Exotic Conchology, 1821-1822; 2d ed., 1841, edited by Hanley.
- SWAMMERDAM, JOANNES. *Biblia Naturæ sive historie insectorum Lugduni Batavorum*, 2 vols., 1737 and 1738.
- * — *Bibel der Natur*, 1752.
- *TAPPAN, BENJAMIN. Description of some New Shells. *Am. Jour. Sci. and Arts*, XXXV, 1839, pp. 268-270.
- *TAPPARONE-CANEVRI, C. Fauna Malacologica delle Nuova Guinea. *Ann. Mus. Civ. Stor. Nat. Genoa*, 1883, pp. 1-313, 11 plates.
- * — Molluschi terrestri e d'acqua dolce. Viaggio di Leonardo Fea in Birmania, etc. *Ann. Mus. Civ. Stor. Nat. Genoa*, 1889, pp. 295-359, 3 plates.
- TASLE. Catalogue des Mollusques du Morbihan. *Bull. Poly. Morbihan*, 1864. Separate.
- *TATE, RALPH. A plain and easy account of the Land and Fresh-water Mollusks of Great Britain, etc., 1866, pp. 244, figs.
- * — On the Land and Fresh-water Mollusca of Nicaragua. *Am. Jour. Conch.*, 1869, pp. 151-162. *Proc. N. M. vol. xxii*—63
- *TATE, RALPH. [Unio stuarti in rain-water holes at Umbum, Australia, is eaten by natives.] *Trans. Roy. Soc. Adelaide*, 1880, p. 171.
- * — A Revision of the Recent Lamellibranch and Palliobranch Mollusca of South Australia. *Trans. and Pr. and Rep. of Roy. Soc. S. Austr.*, 1887, pp. 76-111; *Supplement in Trans. of same*, 1887, pp. 67-69.
- *TATE, R., and BRAZIER, J. Check list of the Fresh-water Shells of Australia. *Proc. Linn. Soc. N. S. W.*, 1881, pp. 552-569.
- TAUCH, LEOPOLD. Ueber einige Conchylien aus dem Tanganyika-See, etc. *Sitz. Akad. Wiss. Wien.*, 1894, pp. 56-70.
- *TAYLOR, G. W. Preliminary Check list of the Land and Fresh-water Mollusks of Canada. *Ottawa Nat.*, 1892, pp. 33-37.
- * — Land and Fresh-water Shells in the Rocky Mountains. *Nautilus*, 1893, pp. 85, 86.
- * — The Land and Fresh-water Shells of Alberta. *Ottawa Nat.*, 1895, pp. 173-178.
- *TAYLOR, JOHN. Unusually large Unio pictorum. *Jour. of Conch.*, 1884, p. 224.
- *TAYLOR, J. W. Unio pictorum var. latior in Lancaster. *Jour. of Conch.*, 1888, p. 331.
- * — A Monograph of the Land and Fresh-water Mollusca of the British Isles. Introduction, 1894.
- TAYLOR, J. W., and ROEBUCK, W. D. Authenticated Materials toward a Land and Fresh-water Molluscan Fauna of Ireland. *Proc. Roy. Irish Academy*, 1888, pp. 672-692.
- *TAYLOR, J. W., ROEBUCK, W. DENISON, and NELSON, W. Conchological Society's list of British Land and Fresh-water Mollusca. *Jour. of Conch.*, IV, 1883, pp. 45-52.
- TAYLOR, W. H., and SHIVERICK, C. F. Catalogue of Shells of the county of Bristol, Massachusetts, [1840?].
- *TENISON-WOODS, J. E. On the Anatomy and Life History of Mollusks peculiar to Australia. *Jour. and Proc. Roy. Soc. N. S. W.*, 1888, pp. 106-187, 14 plates.

- *TENISON-WOODS, J. E. Malaysian Land and Freshwater Mollusca.
Proc. Linn. Soc. N. S. W., (2) III, 1888, pp. 1003-1095, plates XXVII-XXX.
- *TERVER, M. V. Catalogne des Mollusques Terrestres et Fluviatiles dans les possessions Françaises au nord de l'Afrique, 1839.
- *THEOBALD, WILLIAM. Descriptions of new species of Unionidæ.
Jour. Asiat. Soc. Beng., 1874, pp. 207-209.
- Descriptions of New Land and Fresh-water Shells from India and Burmah.
Jour. Asiat. Soc. Beng., 1876, pp. 183-189, plate.
- THOMPSON, F. C. Report to the British Association on the Fauna of Ireland, 1843.
- The anatomy of the Fresh-water Mussel.
Jour. Northumb. Soc., IV, 1887, pp. 82-86, 1 pl.
- *THOMPSON, WILLIAM. Catalogue of Land and Fresh-water Mollusca of Ireland.
Ann. Nat. Hist., 1840, pp. 16, 109, 194-206.
- *TODD, J. E. Chelydra versus Unio.
Ann. Nat. Hist., 1883, p. 428.
- TOTTEN, JOSEPH G. List of Shells on and near the Coast of New England.
Rept. on Geol., Min., Bot., and Zool. of Mass., by E. Hitchcock, 1833, pp. 560-562.
- *TRASK, J. B. [California Fresh-water Shells.]
Proc. Cal. Acad. Sci., 1855, pp. 28-30.
- TREVIRANUS, GOTTLIEB REINHOLD. Ueber die Fortpflanzung der Anodonta.
Zeits. Phys., 1824, pp. 36-41; *Nachtrag*, 1828, pp. 153, 154.
- Mémoire sur les Organes Génératifs.
Zeits. Phys., 1824.
- Memorandum sur les Organes de la Génération des Mollusques.
Ferussac's Bull., 1825, pp. 285-290.
- Appendice aux Observations Anodontes.
Zeits. Phys., 1828.
- *TRISTRAM, H. B. Supplemental Catalogue of Terrestrial and Fluvialile Mollusks collected in Guatemala, etc.
- *TRISTRAM, H. B. Report on the Terrestrial and Fluvialile Mollusca of Palestine.
Proc. Zool. Soc., 1865, pp. 530-545.
- * — The Survey of Western Palestine, the Flora and Fauna of Palestine, 1884.
Proc. Zool. Soc., 1863, pp. 411-414.
- TROOST, G. Seventh Annual Report on the Geology of Tennessee, 1844, pp. 40-42.
- TROSCHER, F. H. Die Familie der Nadjaden. A great number of notices of papers on Unionidæ.
In *Archiv für Naturgeschichte*, 1837-1869.
- TRUE, JOSEPH. Shells gathered about Salem, Massachusetts, 1858.
- *TRYON, GEORGE W. Catalogue of Mollusca collected by Professor Sheldon at Davenport, Iowa.
Am. Jour. Conch., 1865, pp. 68-70.
- * — [Various Reviews.]
Am. Jour. Conch., 1866-1869.
- * — Notes on Mollusca collected by F. V. Hayden in Nebraska.
Am. Jour. Conch., 1868, pp. 150, 151.
- * — Structural and Systematic Conchology, 3 vols., 1882-1884, 140 plates.
- TUFTS, SAMUEL, JR. A List of Shells collected at Swampscott, etc.
Proc. Essex Inst., 1856, pp. 25-32.
- Catalogue of Shells deposited in the State Cabinet.
Cat. Mass. State Cab. Appendix, 1859, pp. 85-88.
- TURTON, WILLIAM. Translation of Linneus Systema Naturæ, 1802.
- * — British Fauna, etc., 1807.
- * — A Conchological Dictionary of the British Islands, 1819, pp. 272, 28 plates.
- * — Conchylia Insularum Britannienm, 1822, colored plates.
- * — A Manual of the Land and Fresh-water Shells of the British Islands, 1831, pp. 143, colored plates. New editions by John E. Gray, 1840, 1857.
- ULICHUY. [On Parasites on Anodonta.]
Arch. für Naturg., 1878, pp. 215-217, pl. VI.
- UNGER, F. W. De Anodonta anatina.
Diss. Vindobonæ, 1827.

- UPSON, J. B.
In *Our Home and Science Gossip*, 1881.
- *VALENCIENNES, ACHILLE. Coquilles Marines Bivalves de l'Amérique Equinoxiale recueillis pendant le Voyage de MM. De Humboldt et Bonpland.
In *Recueil d'Observations*, etc., 1833.
- *— Coquilles Univalves Terrestres et Fluviales.
In *Recueil d'Observations*, 1833.
- *VALENTIN, B. BERNHARD. Amphitheatrum Zootomicum, etc., 1720.
- VALENTYN, FRANZ. Voyage de Amboyne, 1754, 76 pp., 18 plates.
- *VANDERHOOM, J. Handbook of Zoology, 1856, 1858.
- *VAN HALEN, M. Liste des Mollusques dans la Mer du Nord et de l'île Nordernay, etc.
Ann. Gén. Sci. Phys., 1820, p. 374.
- *VAUGHAN, T. W. Mollusks of Dorcheat Bayou and Lake Bisteneau, etc.
Nautilus, V, 1892, pp. 109-111.
- *— Notes on Mollusks from northwestern Louisiana, etc.
Am. Naturalist, 1893, pp. 944-961.
- VERNEUIL, E. Mémoires de Géologie sur la Crimée, 1837.
- VEST, WILH. V. Über die Abreibung der Werbel bei den Süßwassermuscheln, etc.
Verh. Mitth. Sieb., 1862, pp. 105-110.
- Über Margaritana bonelli Fer.
Verh. Mitth. Sieb., 1866, pp. 193-201; *Nachtrag*, 1867, pp. 202-204.
- VINCENTIUS. Speculum Naturale, 1473, 1495.
- *VIGNON, M. Catalogues des Mollusques Terrestres et Fluviales Occidentales d'Afrique.
Bull. Soc. Mal. de France, 1888, pp. 65-76.
- VILLA, ANT. Benacensis Conchyliorum, 1841.
- Catalogo dei Molluschi della Lombardia.
Notiz. Nat. Civ. Lomb., 1844.
- Description d'un Unio de l'Australie.
Jour. de Conch., 1871, p. 328.
- *VILLA, A. and G. B. Dispositio Conchyliorum in Collectione A. et G. B. Villa, 1841.
- *VILLA, A. and G. B. Specie e varietà di Molluschi della Lombardia, etc., 1871.
- VIZE, J. E. Land and Fresh-water Shells, etc.
Tr. Wool. Nat. Field Club, 1892, pp. 180-188.
- *VOGT, CARL. On Some Inhabitants of the Fresh-water Mussels.
Ann. Nat. Hist., 1850, pp. 450-454.
- VOGT, C., and YUNG. Traité d'anatomie comparée, etc., 1888.
- VOIT, CARL. Allgemeine Betrachtungen aus der chemischen Untersuchung der Perlmuschelauszug.
Fror. Notizen, 1860, pp. 289-295.
- VON HÜBER. Zur Naturgeschichte der Unionen.
J. B. Mus. Karuth., 1871, pp. 151-157, plate.
- *VON WAHL, EDUARD. Die Süßwasser-Bivalven Livlands.
Arch. für Naturkunde Liv., 1855, pp. 75-148. Separate.
- *WAARDENBERG, HENRICO GULIELMI. Commentatio de Historia Naturali Animalium Molluscorum, etc., 1827.
- WAILES, B. L. C. Mollusca of the Mississippi.
Geol. Rep. of Miss., 1854, p. 338, 339.
- WALDERDORFF, COUNT DE. Systematisches Verzeichniss der im Kreise Cattaro im Südlichen Dalmatien, etc.
Verh. Zool. Bot. Ges. Wien, 1864, pp. 503-514.
- *WALKER, BRYANT. Catalogue of the Shell-bearing Mollusca of Michigan.
Jour. of Conch., 1879, pp. 325-337.
- *— On the occurrence of Unio complanatus in Michigan.
Nautilus, 1889, pp. 16, 17.
- *— The Shell-bearing Mollusca of Michigan.
Nautilus, 1892, pp. 13-19, 31-35, 42-47, 63-67, plate; 1893, pp. 135-141.
- *— Shells of Saginaw Valley, Michigan.
Nautilus, 1894, pp. 125-129.
- *— A month with the Michigan Fish Commission.
Nautilus, 1895, pp. 2-6.
- *— The distribution of the Unionidæ in Michigan, 1898.
- *WALTON, JOHN. The Mollusca of Monroe County, New York.
Proc. Roch. Acad. Nat. Sci., 1892, pp. 3-19, 8 plates. Separate.
- WARREN, J. The Conchologist, 1834.

- *WATTEBLED, G. Catalogue des Mollusques de Monlins.
Jour. de Conch., 1881, pp. 316-333. Separate.
- *— Description d'une nouvelle Unio provenant du Soudan occidental.
Jour. de Conch., 1884, p. 132, fig.
- *— Description de Mollusques inédits de l'Annam, etc.
Jour. de Conch., 1886, pp. 54-71, 3 plates.
- WEBB, W. M. *Anodonta cygnea* Linn.
Science Gossip, 1888, pp. 235, 236.
- *WEINLAND, DAVID F. [Remarks on the Corrosion of Shells of Fresh-water Clams.]
Proc. Bost. Soc. Nat. Hist., 1857, pp. 57, 58.
- WESTERLUND, C. A. *Malakozoologie* Kalmarlan, etc.
Ofvers. af Kongl. Vet. Ak. For., 1865, p. 537.
- *Sveriges Land- och Sötvatten-Mollusker*, 1865.
- *— *Beschreibungen und Kritik neuer Mollusken*.
Mal. Blatt, 1867, pp. 199-206.
- *Exposé Critique des Mollusques de la Suède et de la Norvège*.
Nor. Act. Ups., 1871, pp. 80-96.
- *— *Fauna Molluscorum Terrestrium et Fluvatilium Sveciæ, Norvege et Daniæ, etc.*, 1873, p. 651.
- *Fauna Europæa Molluscorum Extramarinorum*. *Prodromus*, 1876, 1878.
- *— *Sibiriens Land- och Sötvatten-Mollusker*.
Kong. Sv. Vet. Ak. Hand., 1876, XIV, No. 12. Separate.
- *Sveriges, Norges, Danmarks och Finlands Land- och Sötvatten-Mollusker*, 1884.
- *— *Fauna der in der Paläaretischen Region, Lebenden Binnen-Conchylien, etc.*, 2 vols., 1884-1890.
- *— *Spicilegium Malacologicum; neue Binnen-Conchylien aus der Paläarktischen Region*.
Nach. Deutsch. Mal. Ges., 1893, pp. 116-132.
- *— *Synopsis Molluscorum Extramarinorum Scandinaviæ, etc.*
Act. Soc. Faun. and Fl. Fin., 1897, pp. 1-238.
- WESTERLUND, C. A., and BLANC, H. *Aperçu sur la Faune Malacologique de la Grèce*, 1879.
- WESTERN ACADEMY. *Catalogue of the Unios, etc., of the Ohio River and its tributaries*, 1849.
- *WETHERBY, A. G. *Margaritana dehiscens* Say.
Am. Naturalist, 1878, p. 254.
- *— *On the Geographical Distribution of Certain Fresh-water Mollusks, etc.*
Jour. Cinn. Soc. Nat. Hist., 1880, pp. 317-324.
- *— *The Distribution of Margaritana margaritifera*.
Am. Naturalist, 1882, pp. 675, 676.
- WHEATLEY, CHARLES M. *Catalogue of Shells of the United States, with localities*, 1842; 2d ed., 1845.
- *— *Revision of M. Petit's Catalogue of the genus Monocondylæa*.
Am. Jour. Conch., 1865, pp. 65-67.
- *WHEELER, CHARLES LE ROY. *Remarks on the Unio Muddle*.
Nautilus, 1893, pp. 9, 10.
- *WHITE, C. A. *Note on the Shell Structure of Certain Naiades*.
Am. Jour. Sci. and Arts, XLV., 1868, pp. 400, 401; *Ann. Nat. Hist.*, 1868, pp. 485, 486.
- *— [Uniones found in] *Kjoekkenmøddings in Iowa*.
Am. Naturalist, 1869, pp. 54, 55.
- *— *Are Unios sensitive to light?*
Am. Jour. Sci. and Arts, 1869, pp. 280-282; *Ann. Nat. Hist.*, 1869, p. 399.
- *— *A Review of the Non-Marine Fossil Mollusca of North America*.
Ann. Rep. Dir. U. S. Geol. Surv., 1883, pp. 403-488, 32 plates.
- *— *Notes on the Invertebrate Fauna of the Dakota Formation, etc.*
Proc. U. S. Nat. Mus., 1894, pp. 131-138.
- *WHITEAVES, J. F. *On the Land and Fresh-water Mollusca of Lower Canada*.
Can. Naturalist, 1863. Separate. pp. 31, wood-cuts.
- *— *Notes on the recent discovery of large Unio-like Shells in the Coal Measures of South Joggins, Nova Scotia*.
Trans. Roy. Soc. Can., 1893, pp. 111-121, fig.

- *WHITEAVES, J. F. Notes on Recent Canadian Unionidæ.
Can. Record, 1895, pp. 250-263. Separate.
- * — Additional notes on Recent Canadian Unionidæ.
Can. Record, 1895, pp. 365, 366. Separate.
- *WHITEMORE, T. J. [Shells of Schuyler's Lake, New York.]
Proc. Bost. Soc. Nat. Hist., 1859, pp. 150, 151.
- *WIEGMANN, A. FR. AUGUST. In *Archiv für Naturgeschichte*, 1835-1840.
- WILLIAMS, J. W. A Tumour in the Fresh-water Mussel.
Jour. Anat. Phys., 1889, pp. 306, 307.
— Land and Fresh-water Shells, etc., 1889.
- *WILLIAMSON, W. C. On the Limestones of Manchester, London, and Edinburgh.
Philos. Mag., 1836, pp. 241-249, etc.
- WITTER, F. M. List of the Shells of Iowa.
Quart. Jour. of Conch., 1878, pp. 385-394.
- * — *Unio luteolus* and its allied forms.
Jour. of Conch., 1880, pp. 173-175.
- WOEHRMANN, S. VON. Ueber die Systematische Stellung der Trigoniden, etc.
Jahrb. Geol. Reich., 1893, pp. 1-28, 2 plates.
- WOHLBEREDT, O. Nachtrag zu dem Verzeichniss der im der Preussischen Oberlausitz, etc.
Abh. Ges. Görlitz, 1893, pp. 167-180.
- WOLF, JOHANN. Einige Beobachtungen über die Entenmiesmuschel *Mytilus anatinus*.
Voigt's Mag., 1803, pp. 23-30.
- *WOOD, S. V. [Lacustrine Formations among the Tertiary Beds, etc.]
Mag. Nat. Hist., 1834, pp. 274, 275.
- *WOOD, WILLIAM. Observations on the hinges of British Bivalve Shells.
Trans. Linn. Soc. Lond., 1802, pp. 154-176.
- * — General Conchology, etc., 1815, colored plates.
- * — Index Testaceologicus, etc., 1818; subsequent editions, 1825, supplement, 1828; 2,300 figs. Edited by Sylvanus Hanley, 1856.
- *WOOD, WILLIARD M. [Japanese Anodonta in San Francisco markets.]
Nautilus, 1892, pp. 51, 52.
- *WOOD, W. M., and RAYMOND, W. T. Mollusks of San Francisco.
Nautilus, 1891, pp. 54-58.
- WOODS, J. E. TENISON. On the Fresh-water Shells of Tasmania.
Proc. Roy. Soc. Tasm., 1875, pp. 66-82; Abstract in *Jour. de Conch.*, 1875, pp. 80-87, 203.
— On some new Marine Mollusca.
Trans. Roy. Soc. Vict., 1881, pp. 80-83, plate.
- WOODWARD, SAMUEL. British Organic Remains, 1830.
- *WOODWARD, SAMUEL P. An outline of the Geology of Norfolk, 1833.
- * — A Manual of the Mollusca, etc., 1851-56. Several subsequent editions; also translated into French.
— Speke's Journey in Africa.
Blackwood's Mag., November, 1859.
- * — On some Fresh-water Shells from Central Africa.
Ann. Nat. Hist., V, 1860, p. 337.
- *WRIGHT, BERLIN H. A new *Unio* from Florida.
Proc. Acad. Nat. Sci. Phila., 1883, p. 58, figs.
- * — Description of new species of Unionides from Florida.
Proc. Acad. Nat. Sci. Phila., 1888, pp. 113-120, 5 plates.
- * — Check List of North American Unionidæ, etc., 1888.
- * — A new Florida *Unio*.
Nautilus, V, 1892, pp. 124, 125.
- * — The *Unio* muddle.
Nautilus, VI, 1893, pp. 113-116.
- * — Notes on *Unio coruscus* Gould.
Nautilus, VI, 1893, pp. 126, 127.
- * — New Florida Unionides.
Nautilus, IX, 1896, pp. 121, 122, plate.
- * — New American Unionidæ.
Nautilus, IX, 1896, pp. 133-135.
- * — New Unionides.
Nautilus, XI, 1897, pp. 55, 56.
- * — A New Undulate *Unio* from Alabama.
Nautilus, XI, 1898, pp. 101, 102.
- * — Description of a New *Unio*.
Nautilus, XI, 1898, pp. 111, 112.
- * — New Varieties of Unionidæ.
Nautilus, XI, 1898, pp. 123, 124.
- * — New Unionidæ.
Nautilus, XII, 1898, p. 5.

- *WRIGHT, BERLIN H. A New Unio.
Nautilus, XII, 1898, pp. 32, 33.
- * — A New Unio from Texas.
Nautilus, XII, 1898, p. 93.
- * — New Southern Unios.
Nautilus, XIII, 1899, pp. 6-8, 22, 23,
31, 42-43, 50, 69, 89, 90., 1900. pp.
138, 139.
- *WRIGHT, S. HART. A New Unio.
West. Am. Scientist, 1888, p. 60,
plate.
- * — Unionidæ of G[eorgi]a, Ala[bama],
S[outh] C[arolina], and L[ouisian]a,
in South Florida.
Nautilus, IV, 1891, p. 125.
- * WRIGHT, S. HART, and B. H. Notes on
the Unionidæ of Southern Florida.
Conch. Exchange, II, 1888, pp. 67,
95, 111, 112.
- * — Contributions to a knowlede of
United States Unionidæ.
Nautilus, X, 1897, pp. 136-139; XI,
1897, pp. 4, 5.
- * WYATT, THOMAS. A Manual of Con-
chology, etc., 1838, pp. 191, 36 plates.
— Synopsis of Natural History, etc.,
1849, 49 plates.
- * WYMAN, JEFFRIES. Freshwater Shell-
heaps of Florida.
Am. Naturalist, 1869, pp. 393-403.
— Freshwater Shell Mounds on the
St. John's River, Florida.
Mem. Peabody Acad., Salem, 1875,
94 pp., 9 plates.
- YARROW, H. C. Report upon the collec-
tions of Mollusca made in portions of
Colorado, Utah, New Mexico, and
Arizona.
In *Report on Geog. and Geol. Sur-
veys west of 100th Mer.*, V, 1875, by
George M. Wheeler, pp. 923-954.
- YUNG, E. Relations de l'organe de Bo-
janus chez les Mollusques lamelli-
branches, etc.
Arch. Sci. Nat., 1887, pp. 428, 429.
- ZIEGLER. Manuscripts, no date.
- ZIETEN. C. H. v., Die Versteinerungen
Württembergs, 1830-33, 12 parts, 72
plates.

INDEX TO SYNOPSIS OF THE NAIADES.

[Italicised figures, as 749, etc., are used to indicate the principal references.]

	Page.		Page.
Abaeoides (Unio) Haldeman.....	615	Æsopus (Pleurobema) Green.....	509, 764
Abaeus (Margaron) Lea.....	749	Æsopus (Unio) Green.....	745, 764
Abaeus (Pleurobema) Lea.....	749	Æthiopiformis (Unio) von Ihering.....	894
Abaeus (Unio) Haldeman.....	749	Æthiops (Diplodon) Lea.....	877
Abbevillensis (Margaron) Lea.....	722	Æthiops (Margaron) Lea.....	877
Abbevillensis (Unio) Lea.....	722	Æthiops piriecabana (Diplodon) von	
Abbreviatus (Unio) Westerlund.....	808	Ihering.....	877
Aberrans (Margaron) Lea.....	723	Æthiops piriecabana (Unio) von Ihering.....	877
Aberrans (Unio) Lea.....	723	Æthiops (Unio) Lea.....	877
Aberti (Cyprogenia) Conrad.....	610	Aferula (Parreysia) Lea.....	848
Aberti (Margaron) Lea.....	610	Aferulus (Margaron) Lea.....	848
Aberti (Unio) Conrad.....	610	Aferulus (Unio) Lea.....	848
Ablatus (Unio) Lea.....	730	Affinis (Lampsilis) Lea.....	537
Abnormis (Unio) Morelet.....	830	Affinis (Margaron) Lea.....	537
Abortivus (Unio) Heude.....	802	Affinis (Unio) Heude.....	799
Abruptus (Unio) Say.....	540	Affinis (Unio) Lea.....	537
Abyssiniea (Nodularia) von Martens.....	826	Affinis (Unio) Sowerby.....	531, 534
Abyssinieus (Unio) von Martens.....	826	Africanus (Margaron) Lea.....	825
Aerorhynchus (Unio) von Martens.....	861	Africanus (Unio) Lea.....	825
Actephilus (Unio) Bourguignat.....	683	Agricolarum (Anodon) Heude.....	638
Acuens (Pleurobema) Lea.....	749	Aheneus (Margaron) Lea.....	736
Aeuens (Unio) Lea.....	749	Aheneus (Unio) Lea.....	736
Aeuminata (Nodularia) H. Adams.....	826	Alabamensis (Lampsilis) Conrad.....	574
Aeuminatus (Unio) H. Adams.....	826	Alabamensis (Margaritana) Lea.....	619
Aeuticosta (Castalia) Hupé.....	864	Alabamensis (Margaron) Lea.....	619
Aeuticosta Simpson.....	850	Alabamensis (Unio) Conrad.....	574
Aeutrostris (Diplodon) Lea.....	872, 885	Alabamensis (Unio) Sowerby.....	619
Aeutrostris (Unio) Lea.....	885	Alata (Burtonia) Aneey.....	904
Aeutissimus (Margarita) Lea.....	590	Alata (Hyria) Sowerby.....	871
Aeutissimus (Margaron) Lea.....	590	Alata (Lymnadia) Swainson.....	567
Aeutissimus (Medionidus) Lea.....	590	Alata (Metaptera) Stimpson.....	567
Aeutissimus (Unio) Lea.....	590	Alata (Mutela) Lea.....	904
Adansonii (Spatha) Jousseume.....	898	Alata (Mya) Eaton.....	567
Adansonii (Spathella) Jousseume.....	898	Alata (Mysea) Swainson.....	567
Admirabilis (Cameronia) Bourguignat.....	909	Alata (Platiris) Lea.....	904
Ægyptiaca (Nodularia) Cailliaud.....	820, 821	Alata simpsoni (Mutela) von Martens.....	904
Ægyptiaca (Margarita) Lea.....	821	Alata (Spatha) Lea.....	904
Ægyptiaca (Margaron) Lea.....	821	Alata (Symphynota) Lea.....	567
Ægyptiacus (Unio) Cailliaud.....	820, 821, 822	Alata (Unio) Lamarek.....	567
Æneolus (Pseudodon) Drouet.....	837	Alatus (Lampsilis) Say.....	567, 678
Æneus (Unio) Jekeli.....	821	Alatus (Margarita) Lea.....	567
Æquatus (Margaron) Lea.....	717	Alatus (Margaron) Lea.....	567
Æquatus (Unio) Lea.....	717	Alatus poulsoni (Lampsilis) Conrad.....	568
Æquatoria (Nodularia) Morelet.....	806, 823	Alatus (Prisodon) Sowerby.....	871
Æquatorius (Unio) Morelet.....	823	Alatus (Unio) Say.....	566, 567, 577, 581
Aereus (Unio) Reeve.....	573	Alasmodonta Say.....	506, 510, 514, 666
Aeruginosus (Nephronaias) Morelet.....	595	Alasmodon Swainson.....	666
Aeruginosus (Unio) Morelet.....	595	Alasmodonta Say.....	666
Æsopus (Margarita) Lea.....	764	Alasmodontina (Complanaria) Stimpson.....	663
Æsopus (Margaron) Lea.....	764	Alasmodontine.....	501

	Page.		Page.
Alasmodontinus (Unio) Pætel.....	663	Angasi (Anodonta) Pætel.....	893
Aleroni (Unio) Companyo and Massot ..	686, 699	Angasi (Margaritana) Pætel.....	893
Alexandri (Unio) Kobelt.....	686	Angasi (Unio) Sowerby.....	893
Alfieriianus (Unio) Bourguignat.....	861	Angulata (Anodonta) Lea.....	658
Alienigenus (Lampsis) Crosse and Fischer.....	572	Angulata (Gonidea) Lea.....	658
Alienigenus (Unio) Crosse and Fis- cher.....	572	Angulata (Margarita) Lea.....	658
Algira (Lepidodesma) Heude.....	587	Angulata (Margaron) Lea.....	658
Alta (Pleurobema) Conrad.....	756	Angulata (Mya) Wood.....	869
Altilis (Lampsis) Conrad.....	529	Angulatus (Margarita) Lea.....	869
Altilis (Margarita) Lea.....	529	Angusta subflava (Mya) Sehroter.....	682
Altilis (Margaron) Lea.....	529	Angusta (Unio) Lamarek.....	744
Altilis (Unio) Conrad.....	529	Angustata (Anodonta) Clessin.....	927
Altilis (Unio) Reeve.....	532	Angustata (Iridina) Sowerby.....	904
Altus (Margaron) Lea.....	756	Angustata (Mutela) Sowerby.....	904
Altus (Unio) Conrad.....	756	Angustatus (Margarita) Lea.....	731
Amabilis (Margaron) Lea.....	720	Angustatus (Margaron) Lea.....	731
Amabilis (Unio) Lea.....	720	Angustatus (Unio) Lea.....	731
Amazonensis (Anodonta) Lea.....	928	Annulatus (Anodon) Sowerby.....	618
Amazonensis (Glabaris) Lea.....	928	Anodon Oken.....	620
Amazonensis (Margaron) Lea.....	928	Anodonta (Bruguere) Lamarek.....	501, 506, 513, 514, 620, 658, 832, 919, 923
Ambifaria (Nodularia) von Martens.....	823	Anodonta Valenciennes.....	931
Ambifarius (Unio) von Martens.....	823	Anodontæformis (Nodularia) Tapperone- Canefri.....	818
Ambigua (Alasmodonta) Say.....	673	Anodontæformis (Unio) Tapperone-Can- efri.....	818
Ambigua (Castalia) Blainville.....	865	Anodontes Cuvier.....	620
Ambigua (Castalia) Lamarek.....	863, 868	Anodontina (Unio) Lamarek.....	855
Ambigua (Castalia) Sowerby.....	864	Anodontinæ Swainson.....	501
Ambigua (Hemilastena) Say.....	673	Anodontinus (Pseudodon) Roehbrunck.....	840
Ambigua (Margaritana) Kuster.....	673	Anodontites Bruguere.....	620, 624
Ambigua (Mya) Wood.....	864	Anodontoides floridensis (Lampsis) Lea.....	544
Ambigua (Unio) Deshayes.....	863	Anodontoides (Lampsis) Lea.....	543, 544
Ambiguus (Diplodon).....	863	Anodontoides (Margarita) Lea.....	543
Ambiguus (Tetraplodon) Sowerby.....	863	Anodontoides (Margaron) Lea.....	543
Ambiguus (Unio) Philippi.....	891	Anodontoides Simpson.....	506, 514, 658
Amblamodon Rafinesque.....	673	Anodontoides (Unio) Lea.....	503, 509, 543
Amethystus (Anodon) Sowerby.....	929	Anodontoides (Unio) Reeve.....	544
Amnienus (Unio) Rossmüssler.....	698	Anodontopsis Simpson.....	658
Amœnus (Lampsis) Lea.....	555	Anserina (Margarita) Lea.....	924
Amœnus (Margaron) Lea.....	555	Anserina (Margaron) Lea.....	924
Amœnus (Unio) Lea.....	555	Anserinus (Anodon) Spix.....	924
Amphiehanus (Lampsis) Frierson.....	575	Anserirostris (Anodonta) Kuster.....	625
Amphiehanus (Unio) Frierson.....	575	Anthonyi (Margaron) Lea.....	708
Amplus (Unio) Lea.....	724	Anthonyi (Unio) Lea.....	708
Ampullæus (Diplodon) Lea.....	874	Antinilotius (Unio) Bourguignat.....	861
Ampullæus (Margaron) Lea.....	874	Antiochianus (Unio) Locard.....	689
Ampullæus (Unio) Lea.....	874	Antrosa (Amblema) Rafinesque.....	744
Ampudalonia Crosse and Fischer.....	604	Aperta (Anodonta) Rafinesque.....	933
Amygdalum (Lampsis) Lea.....	561, 562	Apieinus (Lampsis) Lea.....	551
Amygdalum (Margaron) Lea.....	561	Apieinus (Margaron) Lea.....	551
Amygdalum (Unio) Lea.....	561, 569	Apieinus (Unio) Lea.....	551
Anataria (Spatha) Jekeli.....	902	Apieellatus (Ptychorhynchus) Heude.....	851
Anatieulus (Margaron) Lea.....	752	Apieellatus (Unio) Heude.....	851
Anatieulus (Unio) Lea.....	752	Apieellatus (Margarita) Lea.....	778
Anatina (Anodonta) Draparnaud.....	626	Apieellatus (Margaron) Lea.....	778
Anatinus (Anodonta) Sowerby.....	648	Apieellatus (Unio) Lea.....	778
Anatinus (Mytilus) Linnæus.....	625	Apieellatus (Unio) Lea.....	778
Aneeps (Nodularia) Deshayes.....	814	Aplatus (Diplodon) Reeve.....	881
Aneeps (Unio) Deshayes.....	814	Aplatus (Unio) Reeve.....	881
Aneeyi (Brazzæa) Bourguignat.....	906, 907	Aplodon Spix.....	910
Aneeyi (Cameronia) Bourguignat.....	909	Apollonia (Anodonta) Bourguignat.....	648
Aneeyi (Grandidieria) Bourguignat.....	829	Applus Leach.....	583
Aneeyi (Moneetia) Bourguignat.....	901	Appressa (Pleurobema) Lea.....	749
Aneeyi (Spatha) Bourguignat.....	898, 901		
Andersonensis (Unio) Lea.....	747		
Andersonianus (Unio) Nevill.....	812		

	Page.		Page.
Appressus (Unio) Lea.....	749	Arietina (Chelidonopsis) Rochebrune....	906
Apprimus (Diplodon) Lea.....	874, 885	Arkansasensis (Anodonta) Lea.....	617
Apprimus (Margaron) Lea.....	874	Arkansasensis (Lampsilis) Lea.....	557
Apprimus (Unio) Lea.....	874	Arkansasensis (Strophitus) Conrad.....	617
Approximus (Lampsilis) Lea.....	537	Arkansasensis (Unio) Lea.....	557
Approximus (Margaron) Lea.....	537	Armatus (Mycetopus) Heude.....	656
Approximus (Unio) Lea.....	537	Arquatus (Unio) Conrad.....	545
Aquillus (Margaron) Lea.....	727	Ascia (Unio) Hanley.....	861
Aquillus (Unio) Lea.....	727	Askewi (Quadrula) Marsh.....	785, 786
Arada (Unio) Philippi.....	683	Askewi (Unio) Marsh.....	786
Araneosa (Zaira) Rochebrune.....	862	Aspatharia Bourguignat.....	901
Aratus (Margaron) Lea.....	744	Asper (Margarita) Lea.....	776
Aratus (Unio) Conrad.....	706	Asper (Margaron) Lea.....	776
Aratus (Unio) Lea.....	743, 744	Asper (Quadrula) Lea.....	776
Araucanus (Margaron) Lea.....	880	Asper (Unio) Lea.....	776
Araucanus (Unio) Philippi.....	880	Asperatus (Margaron) Lea.....	780
Arceformis (Anodonta) Heude.....	641	Asperatus (Unio) Lea.....	780
Arceformis (Margarita) Lea.....	519	Asperimus (Margarita) Lea.....	777
Arceformis (Margaron) Lea.....	519	Asperimus (Margaron) Lea.....	777
Arceformis (Truncilla) Lea.....	519	Asperimus (Unio) Lea.....	777
Arceformis (Unio) Lea.....	519	Asperula (Nodularia) Lea.....	814
Arcidens Simpson.....	506, 514, 661	Asperulus (Margaron) Lea.....	814
Arcidopsis Simpson.....	506, 515, 861	Asperulus (Unio) Lea.....	814
Arconia Conrad.....	506, 515, 580, 859	Astierianus (Unio) Dupuy.....	692
Arctatus (Margarita) Lea.....	717	Ater (Unio) Lea.....	568
Arctatus (Margaron) Lea.....	717	Ater (Unio) Nilsson.....	697
Arctatus tortivus (Unio) Lea.....	718	Ater (Unio) Reeve.....	692
Arctatus (Unio) Conrad.....	717	Atrata (Unio) Swainson.....	881
Arctatus (Unio) Ferussac.....	704	Atrata (Unio) Swainson.....	574
Arctior (Margarita) Lea.....	704	Atratus (Unio) Sowerby.....	881
Arctior (Margaron) Lea.....	704	Atrocostatus (Margaron) Lea.....	768
Arctior (Unio) Lea.....	704	Atrocostatus (Unio) Lea.....	768
Arcuata (Alasmodonta) Barnes.....	676	Atrocostatus (Unio) Reeve.....	771
Arcuata (Anodonta) Hupé.....	915	Atrocostatus (Unio) Sowerby.....	769
Arcuata (Iridina) Potiez and Michaud.....	899	Atromarginatus (Margaron) Lea.....	706
Arcuata (Margarita) Lea.....	899	Atromarginatus (Plectomerus) Conrad.....	706
Arcuata (Margaritana) Stimpson.....	676	Atromarginatus (Unio) Lea.....	706
Arcuata (Margaron) Lea.....	899	Atropurpureum (Alasmodon) Rafinesque.....	672
Arcuata (Mya) Eaton.....	676	Atroviolacea (Obliquaria) Rafinesque.....	744
Arcuata (Spatha) Cailliaud.....	899	Atrovirens (Anodonta) Philippi.....	624, 933
Arcuatus (Mycetopus) Fischer.....	656	Atrovirens (Unio) Rossmasser.....	698
Arcula (Alasmodonta) Lea.....	672	Attenuata (Anodonta) Held.....	623
Arcula (Alasmodonta) Clessin.....	672	Attenuata (Obliquaria) Rafinesque.....	744
Arcula (Baphia) H. and A. Adams.....	672	Attenuatus (Unio) Lea.....	736
Arcula (Margaritana) Lea.....	672	Aubreyi (Anodon) Heude.....	639
Arcula (Margaron) Lea.....	672	Aucklandica (Unio) Gray.....	889
Arcula (Unio) Hanley.....	672	Aucklandicus (Margaron) Lea.....	889
Arcus (Margarita) Lea.....	704	Aurata (Anodonta) Kuster.....	637
Arcus (Margaron) Lea.....	704	Auratus (Margarita) Lea.....	880
Arcus (Unio) Conrad.....	704	Auratus (Margaron) Lea.....	880
Ardusianus (Unio) Moquin-Tandon.....	686	Auratus (Unio) Kuster.....	883
Arealis (Anodonta) Kuster.....	625	Auratus (Unio) Philippi.....	880
Arelatensis (Anodonta) Dupuy.....	624	Auratus (Unio) Reeve.....	881
Aremprosthus (Unio) Locard.....	689	Auratus (Unio) Sowerby.....	881
Areolatus (Anodon) Sowerby.....	925	Aurea (Margaritana) Pætel.....	836
Areolatus (Anodon) Swainson.....	616	Aurea (Quadrula) Lea.....	783
Argentea (Anodonta) Lea.....	659	Aureus (Anodonta) Pætel.....	648
Argentea (Margaron) Lea.....	659	Aureus (Diplodon) Simpson.....	883
Argentea pannosa (Pleurobema) Simpson.....	763	Aureus (Margaron) Lea.....	783
Argentea (Pleurobema) Lea.....	760, 763	Aureus (Pseudodon) Heude.....	836
Argenteus (Margaron) Lea.....	763	Aureus (Unio) Lea.....	783
Argenteus (Strophitus) Conrad.....	659	Auricularius (Unio) Spengler.....	676, 678
Argenteus (Unio) Lea.....	763	Aurora Simpson.....	849
Argenteus (Unio) Sowerby.....	749	Aurorea (Parreysia) Heude.....	849
Argyratus (Lampsilis) Rafinesque.....	577	Auroreus (Unio) Heude.....	849
Arietina (Chelidonura) Rochebrune.....	906	Australis (Diplodon) Lamarek.....	890, 891

	Page.		Page.
Australis (Ilyridella) Swainson	891	Batava (Mysca) Turton	606
Australis (Lampsilis) Simpson	544	Batavi (Unio) Rossmässler	697
Australis legrandi (Diplodon) Petterd	891	Batavus (Margarita) Lea	697
Australis (Margarita) Lea	891	Batavus (Margaron) Lea	697
Australis (Margaron) Lea	891	Batavus (Unio) Maton and Rackett	513,
Australis (Unio) Hanley	890	675, 677, 678, 681, 691, 696, 697	
Australis (Unio) Kuster	890	Baumannii (Spatha) Sturany	899
Australis (Unio) Lamarck	888	Bazini (Quadrula) Heude	800
Ava (Monocondylea) Theobald	839	Bazini (Unio) Heude	800
Ava (Pseudodon) Theobald	839	Beadleiana (Quadrula) Lea	784, 786
Avellana (Pleurobema) Simpson	753	Beadleianus (Margaron) Lea	786
Averillii (Unio) B. H. Wright	560	Beadleianus (Unio) Lea	785
Averyi (Nephronaias) Lea	593	Bealei (Anodonta) Lea	644
Averyi (Unio) Lea	593	Bealei (Margaron) Lea	564, 645
Avicularis (Hyria) Lamarck	871	Bealei (Unio) Lea	564
Avicularis (Unio) Deshayes	871	Beaverensis (Margaron) Lea	724
Avicularis var. b. (Hyria) Lamarck	870	Beaverensis (Unio) Lea	724
Avonensis (Anodonta) Moquin-Tandon	622	Beccarianus (Unio) Tapperone-Canefri	851, 852
Avonensis (Mytilus) Montagu	622	Beccarianus (Virgus) Tapperone-Canefri	852
Avonensis (Mytilus) Turton	622	Bednalli (Unio) Tate	891
Axiacus (Unio) Locard	689	Bellamyi (Nodularia) Jousseau	822
Aztecum (Lampsilis) Philippi	572	Bellamyi (Pharaonia) Jousseau	822
Aztecum (Margaron) Lea	572	Bellamyi (Spatha) Jousseau	896
Aztecum (Unio) Philippi	572	Bellua (Anodonta) Morelet	584
Badium (Alasmodon) Rafinesque	673	Bellua (Cristaria) Morelet	584
Badius (Unio) Kokeil	699	Bellulus (Unio) Lea	746, 747
Bagdadensis (Margaron) Lea	822	Benedictensis (Anodonta) Ferussac	644
Bagdadensis (Nodularia) Bourguignat	822	Benedictensis (Margarita) Lea	644
Bagdadensis (Unio) Bourguignat	822	Benedictensis (Margaron) Lea	644
Bahiensis (Anodonta) Kuster	925	Benedictensis (Symphynota) Lea	644
Bahiensis (Glabaris) von Ihering	925	Benedictii (Anodonta) Latchford	644
Baikii (Spatha) H. Adams	902	Benedictii (Margaron) Lea	644
Bairdianus (Margaron) Lea	563	Bengalensis (Anodon) Sowerby	832
Bairdianus (Unio) Lea	563	Bengalensis (Margarita) Lea	832
Bakeri (Parreysia) H. Adams	846	Bengalensis (Margaron) Lea	832
Bakeri (Unio) H. Adams	846	Bengalensis (Unio) Lea	832
Bakeri (Unio) von Martens	846	Bensoni (Margaron) Lea	656
Bakoyi (Nodularia) Rochebrune	824	Bensonianus (Mycetopus) Pætel	656
Bakoyi (Unio) Rochebrune	824	Bergi (Anodonta) von Ihering	933
Baldwinensis (Margaron) Lea	723	Beringiana (Anodonta) Middendorff	584, 628
Baldwinensis (Unio) Lea	723	Berlandieri (Lampsilis) Lea	570
Baletonicus (Unio) Kuster	683	Berlandieri (Margaron) Lea	570
Balonnensis (Unio) Conrad	891	Berlandieri (Unio) Lea	570
Balzani (Fossula) von Ihering	914	Beskeanus (Diplodon) Dunker	875
Balzani (Iheringella) von Ihering	914	Beskeanus (Margaron) Lea	875
Balzani (Plagiodon) von Ihering	914	Beskeanus (Unio) Dunker	875
Bambousearum (Anodon) Morelet	930	Bhamoensis (Parreysia) Theobald	843
Bambousearum (Glabaris) Morelet	930	Bhamoensis (Unio) Theobald	843
Bandini (Unio) Kuster	698	Bialata (Barbala) Chenu	584
Bandinii (Unio) Rossmässler	686	Bialata (Symphynota) Lea	584
Baphia H. and A. Adams	674	Bialata (Unio) Hanley	584
Baphiæ Meuschen	674	Bialatus (Hyriopsis) Simpson	579
Barbala Humphreys	501, 583	Bianguilata (Anodon) Sowerby	658
Barbata Sowerby	583	Bianguilatus (Lampsilis) Lea	533
Barbatia	861	Bianguilatus (Margaron) Lea	533
Barnesiana (Pleurobema) Lea	751	Bianguilatus (Unio) Lea	533
Barnesianus (Margarita) Lea	751	Bi-caelatus (Unio) Reeve	548
Barnesianus (Margaron) Lea	751	Bicolor (Unio) Rafinesque	744
Barnesianus (Unio) Lea	751	Biemarguata (Truncilla) Lea	524
Barrandii (Unio) Dupuy	692	Biemarginatus (Margaron) Lea	524
Barrattii (Margaron) Lea	726	Biemarginatus (Unio) Lea	524
Barrattii (Unio) Lea	726	Bicsiana (Parreysia) Heude	850
Barrotii (Unio) Kuster	735	Biesianus (Unio) Heude	850
Barrotii (Unio) Pætel	726	Bigbyana (Pleurobema)	760
Basalis (Unio) Lea	724	Bigbyensis (Pleurobema) Lea	751
Batava (Mya) Maton and Rackett	696	Bigbyensis (Margaron) Lea	752

	Page.		Page.
Bigbyensis (Unio) Lea	751	Bourguignati (Pseudospatha) Bourgui-	
Bigerrensis (Unio) Millet	692	gnat	578
Bigibba (Anodon) Heude	638	Bourguignati (Spatha) Bourguignat	898
Bigorrensis (Unio) Locard	692	Bourguignati (Spathella) Bourguignat ..	898
Bilineata (Margarita) Lea	855	Bourguignati (Unio) Landran	821
Bilineata (Symphynota) Lea	855	Bourguignatianus (Margaron) Lea	687
Bilineatus (Margaron) Lea	855	Bourguignatianus (Unio) Lea	687
Bilineatus (Unio) Hanley	855	Bournianus (Margaron) Lea	747
Billotianus (Unio) Charnes	847	Bournianus (Pleurobema) Lea	747
Biloba (Epilobasma) Rafinesque	744	Bournianus (Unio) Lea	747
Bineurus Simpson	839	Bouvieri (Unio) Nevill	814
Binneyi (Diplodon) Lea	878	Boydianus (Margaron) Lea	536
Binneyi (Margaron) Lea	878	Boydianus (Unio) Lea	536
Binneyi (Unio) Lea	878	Boykiniana (Quadrula) Lea	770
Binominatus (Lampsilis) Simpson	528	Boykinianus (Margaron) Lea	770
Birmanus (Unio) Hanley and Theobald ..	845	Boykinianus (Plectomerus) Conrad	770
Bischoffi (Unio) von Ihering	894	Boykinianus (Unio) Lea	770
Bissellianus (Margaron) Lea	743	Bracteatus (Lampsilis) Gould	543
Bissellianus (Unio) Lea	743	Bractcatus (Margaron) Lea	543
Biwæ (Nodularia) Kobelt	810	Bractcatus (Unio) Gould	543
Biwæ (Unio) Kobelt	810	Brandti (Anodonta) Drouet	648
Blainvilleana (Anodon) Sowerby	924	Brandtii (Nodularia) Kobelt	810
Blainvilleana (Anodonta) Lea	914, 915	Brandtii (Unio) Kobelt	810
Blainvilleana (Columba) Clessin	915	Brazosensis (Margaron) Lea	768
Blainvilleana (Leila) Lea	915	Brazosensis (Unio) Lea	768
Blainvilleana (Margarita) Lea	915	Brazzea Bourguignat	506, 515, 906
Blainvilleana (Margaron) Lea	915	Brevialis (Unio) Crouch	792
Blandianus (Margaron) Lea	779	Brevialis (Unio) Lamarck	691
Blandianus (Unio) Lea	778	Brevialis (Unio) Sowerby	602
Blandingianus (Margarita) Lea	742	Breviculus brittsi (Lampsilis) Simpson ..	533
Blandingianus (Margaron) Lea	742	Breviculus (Lampsilis) Call	533
Blandingianus (Unio) Lea	742	Breviculus (Unio) Call	533
Bleaverensis (Unio) Pætel	724	Brevidens (Margarita) Lea	517
Bloyeti (Spatha) von Martens	898	Brevidens (Margaron) Lea	517
Bloyeti (Spathella) Bourguignat	898	Brevidens (Truncilla) Lea	517
Bohmi (Nodularia) von Martens	823	Brevidens (Unio) Lea	517
Bohmi (Unio) von Martens	823	Brevirostris (Unio) Kuster	700
Bolli (Unio) Call	783	Brevis (Anodon) Sowerby	932
Bonelli (Anodon) Sowerby	654	Brevis (Pleurobema) Lea	763
Bonelli (Leguminaia) Westerlund	654	Brevisubelliptica (Pleurobema) Simpson ..	763
Bonelli (Margaritana) Kobelt	654	Brevis (Unio) Lea	763
Bonelli (Microcondylaea) Clessin	654	Bridgesii (Anodonta) Lea	926
Bonelli (Microcondylus) Drouet	654	Bridgesii (Glabaris) Lea	926
Bonelli (Unio) Rossmässler	653	Bridgesii (Margaron) Lea	927
Bonellii (Baphia) H. and A. Adams	653	Bridoux (Brazzea) Bourguignat	907
Bonellii (Margarita) Lea	653	Bridoux (Burtonia) Bourguignat	578
Bonellii (Margaron) Lea	653	Bridoux (Cameronia) Bourguignat	909
Bonneaudi (Margaron) Lea	814	Bridoux (Monetia) Bourguignat	901
Bonneaudi (Nodularia) Eydoux	813	Bridoux (Mutela) Bourguignat	903
Bonneaudi (Unio) Eydoux	813	Bridoux (Unio) Bourguignat	861
Bonplandi (Unio) Valenciennes	884	Brimleyi (Unio) S. H. Wright	720
Borealis (Lampsilis) Gray	535	Brittsi (Lampsilis) Simpson	533
Borealis (Unio) Gray	535	Broti (Parreysia) Deshayes	847
Borellii (Nodularia) Ancy	823	Broti (Unio) Deshayes	847
Borellii (Unio) Ancy	823	Browniana (Triquetra) H. and A. Adams ..	872
Borneensis (Ctenodesma) Issel	853	Brownianus (Margarita) Lea	871
Bornensis (Unio) Issel	852, 853	Brownianus (Margaron) Lea	872
Bourguignati (Brazzea) Bourguignat	907	Brownianus (Prisodon) Lea	871
Bourguignati (Burtonia) Bourguignat	578	Brownianus (Unio) Lea	871
Bourguignati (Cameronia) Bourguignat ..	909	Brownii (Diplodon) Lea	876
Bourguignati (Chambardia)	902	Brownii (Margaron) Lea	876
Bourguignati (Grandidieria) Joubert	829	Brownii (Unio) Lea	876
Bourguignati (Leguminaia) Locard	651	Bruguierianus (Margaron) Lea	694
Bourguignati (Mutela) Bourguignat	904	Bruguierianus (Unio) Bourguignat	694
Bourguignati (Pharaonia) Rochebrune ..	821	Brumbyana (Pleurobema) Lea	760
Bourguignati (Pleiodon) Bourguignat	909	Brumbyanus (Margaron) Lea	760

	Page.		Page.
Brumbyanus (Unio) Lea	760	Ceruleus (Margaron) Lea	811
Brumleyanus (Unio) Lea	760	Ceruleus (Mycetopus) Heude	656
Buchanensis (Anodonta) Lea	659	Ceruleus (Unio) Lea	811
Buchanensis (Margarita) Lea	659	Caffer africana (Nodularia) Lea	825
Buchanensis (Margaron) Lea	659	Caffer (Margaron) Lea	825
Buckleyi (Margaron) Lea	713	Caffer (Nodularia) Krauss.	824, 825
Buckleyi oreutti (Unio) S. H. Wright	713	Caffer (Unio) Krauss.	824, 825
Buckleyi (Unio) Lea	504, 712, 713, 719, 857	Caffer vaalensis (Nodularia) Chaper	825
Buddianus (Margaron) Lea	713	Cafferia Simpson	824
Buddianus (Unio) Lea	713	Cahabensis (Quadrula) Lea	732
Bulbosa (Pleurobema) Lea	759	Cahabensis (Unio) Lea	782
Bulbosus (Margaron) Lea	759	Caillaudi (Spatha) von Martens.	896
Bulbosus (Unio) Lea	759	Cailliaudii (Anodonta) Lea	918
Bullata (Obliquaria) Rafinesque	744	Cailliaudii (Margarita) Lea	812
Bullatus prasinus (Unio) Pætel	779	Cailliaudii (Margaron) Lea	812, 918
Bullatus schoolcraftensis (Unio) Pætel	780	Caipira (Diplodon) von Ihering	877
Bullatus (Unio) Conrad	779	Caipira (Unio) von Ihering	877
Bullella Simpson	672	Calathus (Nodularia) Bourguignat	824
Bulloides (Diplodon) Lea	887	Calathus (Unio) Bourguignat	824
Bulloides (Margaron) Lea	887	Calceola (Alasmidonta) Lea	667, 668
Bulloides (Unio) Lea	887	Calceola (Baphia) H. and A. Adams.	668
Bulloideus Simpson	887	Calceola (Margarita) Lea	668
Burkensis (Margaron) Lea	730	Calceola (Margaritana) Kuster	668
Burkensis (Unio) Lea	730	Calceola (Margaron) Lea	668
Burmanus (Parreysia) Blanford	845	Calceolus (Strophitus) Conrad	668
Burmanus (Unio) Blanford	845	Calceolus (Unio) Lea	667, 668
Burroughiana (Anodonta) Chenu	926	Caldwellii (Margaron) Lea	593
Burroughiana (Anodonta) Lea	832	Caldwellii (Nephronaias) Lea	593
Burroughianus (Diplodon) Lea	833	Caldwellii (Unio) Lea	593
Burroughianus (Margarita) Lea	883	Calendis (Unio) Rafinesque	744
Burroughianus (Margaron) Lea	883	Californiensis (Anodonta) Lea	629
Burroughianus (Unio) Lea	883	Californiensis (Margaron) Lea	629
Bursa-pastoris (Quadrula) B. H. Wright	791	Caliginosus (Margaron) Lea	547
Bursa-pastoris (Unio) B. H. Wright	791	Caliginosus (Unio) Lea	547
Burtchianus (Unio) S. H. Wright	715	Calimatarum (Nephronaias) Morelet	596
Burtoni (Grandidieria) Woodward	828	Calimatarum (Unio) Morelet	596
Burtoni insignis (Grandidieria) Bourguignat	828	Calipygos (Anodonta) Kobelt	641
Burtoni insignis (Unio) von Martens	828	Callifera (Anodonta) von Martens	648
Burtoni (Margaron) Lea	828	Calliscapha Swainson	902
Burtoni servainiana (Grandidieria) Bourguignat	828	Callista (Grandidieria) Bourguignat	829
Burtoni servainiana (Unio) von Martens	828	Callonia Simpson	506, 507, 515, 867
Burtoni smithi (Unio) von Martens	829	Callosa (Anodonta) Kuster	625
Burtoni sturanyi (Grandidieria) von Martens	828	Callosus (Margaron) Lea	701
Burtoni sturanyi (Unio) von Martens	825	Callosus (Unio) Lea	701
Burtoni (Unio) Smith	829	Cambodgensis (Margaron) Lea	838
Burtoni (Unio) Woodward	827, 828	Cambodgensis (Monocondylaea) Petit	838
Burtonia Bourguignat	577	Cambodgensis (Pseudodon) Petit	838
Buschiana (Monodontina) Conrad	836	Cambodiensis (Margaron) Lea	831
Buxeus (Margaron) Lea	708	Cambodiensis (Physunio) Lea	831
Buxeus (Unio) Lea	708	Cambodiensis (Unio) Lea	831
Buxtoni (Unio) B. H. Wright	719	Cambojensis (Pseudodon) Morelet	581
Bythinicus (Unio) Kobelt	695	Cambojensis (Unio) Sowerby	581, 819
Byssanodonta	501	Camelopardalis (Unio) Sowerby	542
Cacao (Margaron) Lea	785	Camelopardalis (Lampsilis) Lea	542
Cacao (Unio) Lea	785	Camelopardalis (Margaron) Lea	542
Caelata (Micromya) Conrad	525	Camelopardalis (Unio) Lea	542
Caelatura	862	Camelus (Margarita) Lea	613
Caelatura Conrad	820	Camelus (Margaron) Lea	613
Caelatus (Margarita) Lea	526	Camelus (Unio) Lea	612
Caelatus (Margaron) Lea	526	Cameronia Bourguignat	909
Caelatus (Unio) Conrad	525	Campsianus (Unio) Pætel	822
Cerulea (Nodularia) Lea	811, 812	Camptodon (Margarita) Lea	740
Ceruleus (Margarita) Lea	811	Camptodon (Margaron) Lea	740
		Camptodon (Unio) Say	740
		Canadensis (Margaron) Lea	527
		Canadensis (Unio) Lea	527

	Page.		Page.
Canefricanus (Lamellidens) Simpson	857	Castelnaudi (Prisodon) Hupe	871
Canthyrta Swainson	680, 738	Castelnaudii (Columba) Clessin	915
Capax (Lampsilis) Green	529, 530	Castus (Margaron) Lea	708
Capax (Margaron) Lea	530	Castus (Unio) Lea	708
Capax (Unio) Green	529	Catacraeta (Anodonta) Say	630, 631
Caperatus (Dromus) Lea	615	Catacraeta (Unio) Deshayes	631
Caperatus (Margaron) Lea	615	Catacraetus (Mytilus) Eaton	631
Caperatus (Unio) Lea	615	Catawbensis (Margaron) Lea	725
Capigliolo bondini (Unio) Pætel	685	Catawbensis (Unio) Lea	725
Capigliolo (Unio) Kuster	699	Catillus (Unio) Conrad	789
Capigliolo (Unio) Payraudeau	685	Catilus (Unio) B. H. Wright	789
Capillaris (Unio) Lea	522	Caudatus (Unio) Wagner	871
Capillus (Unio) Say	525	Caudiculatus (Hyriopsis) von Martens	581
Capitatus (Cuneopsis) Heude	805	Caudiculatus Simpson	581
Capitatus (Unio) Heude	805	Caudiculatus (Unio) von Martens	581
Capseformis (Margarita) Lea	524	Caveata (Quadrula) Heude	801
Capseformis (Margaron) Lea	524	Caveatus (Unio) Heude	801
Capseformis (Trunella) Lea	524	Cellensis (Anodonta) C. Pfeiffer	622
Capseformis (Unio) Lea	524	Cellensis (Anodonta) Kobelt	640
Carbonarius (Margarita) Lea	701	Cellensis beringiana (Anodonta) Schröter	628
Carbonarius semigranosus (Unio) Pætel	702	Celtiformis (Cuneopsis) Heude	805
Carbonarius (Unio) Lea	701	Celtiformis (Unio) Heude	804, 805
Cardiacea (Unio) Deshayes	789	Cerina (Quadrula) Conrad	787
Cardiacea (Unio) Guérin	790	Cerinus (Unio) Conrad	787
Cardium oecidens (Unio) Pætel	527	Chaiziana (Anodonta) Rang	896
Cardium (Unio) Conrad	527	Chaiziana (Margarita) Lea	897
Cardium ventriosus (Unio) Pætel	527	Chaiziana (Margaron) Lea	897
Carinata (Anodonta) Dunker	933	Chaiziana (Spatha) H. and A. Adams	897
Carinata (Mya) Eaton	539	Chaiziana (Spatha) Rang	896
Carinata (Solenalia) Heude	657	Chaleoensis (Anodonta) Crosse and Fischer	630
Carinatus (Mycetopus) Heude	657	Chambardia Bourguignat	902
Carinatus (Unio) Barnes	539	Chamberlainia Simpson	505, 506, 511, 514, 532
Carinifera (Anodonta) Conrad	588	Chantrei (Anodonta) Bourguignat	648
Carinifera (Pilsbryonecha) Conrad	588	Chantrei (Leguminalia) Locard	651, 653
Carinifera (Unio) Lamarek	721	Chantrei (Monocondylaea) Pætel	651
Carinthiaeus (Unio) Rossmässler	698	Chantrei (Pseudodon) Locard	653
Cariosa (Anodonta) Kuster	625	Chantri (Unio) Locard	689
Cariosa (Lampsilis) Stimpson	529	Chapalensis (Anodonta) Crosse and Fischer	630
Cariosa (Mya) Eaton	529	Chaperi (Pseudodon) de Morgan	833
Cariosa (Unio) Lamarek	529, 671	Chaperi (Pseudodus) de Morgan	838
Cariosus (Lampsilis) Say	528	Charbonnieri (Brazzea) Bourguignat	907
Cariosus (Margarita) Lea	529	Charbonnieri (Cameronia) Bourguignat	909
Cariosus (Margaron) Lea	529	Charbonnieri (Unio) Bourguignat	821
Cariosus (Unio) Say	528	Charlottensis (Margaron) Lea	663
Carneus (Unio) Kuster	696	Charlottensis (Symphynota) Lea	663
Carolinensis (Castalia) Sowerby	864	Charlottensis (Unio) Lea	663
Carolinensis (Unio) Bose	741	Charpentieri (Anodonta) Kuster	625
Carthagenia (Anodonta)	922	Charruana (Unio) d'Orbigny	876
Carunculina Simpson	563	Charruanus (Diplodon) d'Orbigny	876
Casablancae (Margaron) Lea	882	Charruanus (Margarita) Lea	876
Casablancae (Diplodon) Philippi	882	Charruanus (Margaron) Lea	877
Casablancae (Unio) Philippi	882	Chathamensis (Margaron) Lea	727
Castalia Lamarek	863	Chathamensis (Unio) Lea	727
Castaliella Simpson	515, 866	Chattanoogaensis (Margaron) Lea	753
Castalina von Ihering	506, 515, 865, 866, 867, 869	Chattanoogaensis (Pleurobema) Lea	753
Castanea (Anodon) Heude	638	Chattanoogaensis (Unio) Lea	753
Castanea (Obovaria) Lea	602	Cheeziana (Anodon) Sowerby	932
Castaneus (Margarita) Lea	602	Chelidonura Rochebrune	906
Castaneus (Margaron) Lea	602	Chelidonopsis Ancey	506, 515, 906
Castaneus (Obovaria) Lea	601	Chickasawhensis (Margaron) Lea	785
Castaneus (Unio) Lea	602	Chickasawhensis (Quadrula) Lea	785
Castaneus (Unio) Rafinesque	744	Chickasawhensis (Unio) Lea	785
Castelnaudi (Anodon) Sowerby	915	Childreni (Diplodon) Gray	883
Castelnaudi (Anodonta) Hupé	928	Childreni (Margarita) Lea	883
Castelnaudi (Hyria) Hupé	871	Childreni (Margaron) Lea	883
Castelnaudi (Leila) Hupé	915		

	Page.		Page.
Childreni (Unio) Gray	883	Coarctata (Anodonta) Potiez and Mi-	
Childreni (Unio) Hanley	534	chaud	623
Chilensis (Diplodon) Gray	880	Coarctata (Unio) Lamarek	721
Chilensis (Unio) Gray	880	Coccinea paupereula (Quadrula) Simpson ..	789
Chiloënsis (Diplodon) Kuster	882	Coccinea (Quadrula) Conrad	788
Chiloënsis (Unio) Kuster	882	Coccineus (Margarita) Lea	789
Chinensis (Anodonta) Kuster	585, 833	Coccineus (Margaron) Lea	789
Chinensis (Anodonta) Pætel	648	Coccineus (Unio) Conrad	788
Chinensis levis (Parreysia) Heude	850	Cochlearis (Anodon) Sowerby	648
Chinensis (Margaron) Lea	850	Cocoduensis (Unio) Reeve	547
Chinensis (Parreysia) Lea	850	Cœlestis (Iridina) Lea	905
Chinensis squamosus (Parreysia)		Cœlestis (Platiris) Lea	905
Heude	850	Cognata (Anodon) Gould	628, 629
Chinensis (Unio) Lea	850	Cognata (Plagiola) Lea	606
Chinifana (Anodon) Heude	639	Cognatus (Margaron) Lea	606
Chiquetana (Anodon) d'Orbigny	928	Cognatus (Unio) Lea	604, 606
Chlorens (Unio) Heude	808	Colchaguensis (Unio) Philippi	882
Chloris (Unio) Rafinesque	744	Colletopterum Bourguignat	506, 513, 514, 649
Chunii (Margaron) Lea	787	Collina (Alasmodonta) Conrad	669
Chunii (Quadrula) Lea	787	Collinus (Margaron) Lea	669
Chunii (Unio) Lea	787	Collinus (Unio) Conrad	669
Churchillanus (Unio) Bourguignat	650	Coloradoensis (Lampsilis) Lea	568
Churchillianus (Unio) Sowerby	650	Coloradoensis (Margaron) Lea	568
Cicatricosa (Pleurobema) Say	765	Coloradoensis (Unio) Lea	568
Cicatricosus (Unio) Say	765	Coloratus (Unio) Charpentier	700
Cicatricosus varicosus (Unio) Pætel	765	Columba Lea	914
Ciconia (Anodon) Sowerby	925, 926	Columbensis (Margaritana) Lea	673
Ciconia (Anodonta) Gould	924, 926	Columbensis (Margaron) Lea	741
Cieur (Margaron) Lea	742	Columbensis (Unio) Lea	741
Cieur (Unio) Lea	742	Columbensis (Unio) Pætel	741
Cilicia (Anodonta) Kobelt	627	Compactus (Margaron) Lea	518
Cincinnatiensis (Margaron) Lea	523	Compactus (Unio) Lea	518
Cinecinnatiensis (Unio) Lea	523	Complanaria Swainson	662
Cinerescens (Toxolasma) Rafinesque	744	Complanata (Alasmodonta) Barnes	665
Cinnamomicus (Margaron) Lea	758	Complanata (Anodonta) Rossmässler	625, 626
Cinnamomicus (Unio) Lea	758	Complanata (Baphia) H. and A. Adams ..	666
Circulus (Margarita) Lea	600	Complanata (Cameronia) Bourguignat ..	910
Circulus (Margaron) Lea	600	Complanata (Complanaria) Conrad	666
Circulus (Obovaria) Lea	600	Complanata (Hyria) Hupé	870
Circulus (Unio) Lea	599, 600, 61	Complanata Katherinae (Symphynota) Lea ..	666
Circumactus (Unio) Lea	750	Complanata (Margarita) Lea	666
Cirratus (Unio) Lea	725	Complanata (Margaritana) Calkins	666
Cistelliformis (Margaron) Lea	716	Complanata (Margaritana) Westerlund ..	677
Cistelliformis (Unio) Lea	716	Complanata (Mutelina) Jousseau	900
Claibornensis (Lampsilis) Lea	537	Complanata (Mya) Dillwyn	720
Claibornensis (Margarita) Lea	537	Complanata (Mya) Eaton	665
Claibornensis (Margaron) Lea	537	Complanata (Spatha) Jousseau	900
Claibornensis (Unio) Lea	537	Complanata (Symphynota) Barnes	665
Clappertoni (Anodonta) Kenig	896	Complanata (Unio) Hanley	666
Clarkianus (Lampsilis) Lea	532	Complanatus jejunos (Unio) Lea	725
Clarkianus (Margaron) Lea	532	Complanatus (Margarita) Lea	721
Clarkianus (Unio) Lea	532	Complanatus (Margaron) Lea	721
Clava (Pleurobema) Lamarek	509, 510, 745, 746	Complanatus (Prisodon) Hupé	870
Clava (Unio) Lamarek	745	Complanatus quadrilaterus (Unio) Lea ..	725
Clavus (Margarita) Lea	746	Complanatus (Unio) Dillwyn	677, 720, 725, 728
Clavus (Margaron) Lea	746	Complanatus (Unio) Middendorff	677
Clavus (Unio) Reeve	746	Compressa (Alasmodonta) Menke	653
Clessini (Glabaris) Fischer	930, 932	Compressa (Anodonta) Menke	626
Clessini (Mycetopus) Fischer	930	Compressa (Complanaria) Conrad	662
Cliffordiana (Obliquaria) Rafinesque	744	Compressa (Margaron) Lea	588
Clinchensis (Margaron) Lea	748	Compressa (Microcondylæa) Pætel	662
Clinchensis (Unio) Lea	748	Compressa (Monocondylæa) Lea	588
Clintonensis (Ptychobranhus) Simpson ..	613	Compressa plebia (Symphynota) C. B.	
Clyone Gistel	583	Adams	663
Coarctata (Anodonta) Anton	630	Compressa (Pseudodon) Conrad	588

	Page.		Page.
Compressa (Spatha) Patet	588	Contortus muticus (Unio) Heude	860
Compressa (Spatha) von Martens	840	Contortus (Unio) Heude	860
Compressa (Symphynota) Lea	662	Contractus (Margaron) Lea	722
Compressissimus (Margaron) Lea	613	Contractus (Unio) Lea	722
Compressissimus (Unio) Lea	613	Contradens (Margarita) Lea	817
Compressus (Margarita) Lea	663	Contradens (Margaron) Lea	817
Compressus (Unio) Conrad	662	Contradens (Nodularia) Lea	817
Compressus (Unio) Heude	851	Contradens (Unio) Lea	817
Comptus (Unio) Deshayes and Jullien	858	Contrarius (Unio) Conrad	537, 733
Conasaugænsis (Pleurobema) Lea	763	Contrarius (Lampsilis) Conrad	537
Conasaugænsis (Unio) Lea	763	Contritus (Unio) Heude	801
Concavus (Margaron) Lea	561	Conus (Unio) Spengler	682, 744
Concavus (Unio) Lea	561	Cooperiana (Quadrula) Lea	781
Concestatator (Lampsilis) Lea	543	Cooperianus (Margarita) Lea	781
Concestatator (Margaron) Lea	548	Cooperianus (Margaron) Lea	781
Concestatator (Unio) Lea	548	Cooperianus (Unio) Lea	781
Concolor (Margaron) Lea	760	Copei (Margaron) Lea	549
Concolor (Unio) Lea	760	Copei (Unio) Lea	549
Confertus (Margarita) Lea	715	Cor (Margarita) Lea	755
Confertus (Margaron) Lea	715	Cor (Margaron) Lea	755
Confertus (Unio)	716	Cor (Pleurobema) Conrad	754
Confertus (Unio) Lea	715	Cor (Unio) Conrad	754
Confragosa (Alasmodonta) Say	661, 662	Corbeti (Unio) Deschamps	856
Confragosa (Baphia) H. and A. Adams	662	Corbicula (Grandidieria) Bourguignat	829
Confragosa (Margarita) Lea	662	Corbiculidæ	828
Confragosa (Margaritana) Calkins	662	Corbis (Margaron) Lea	845
Confragosa (Margaron) Lea	662	Corbis (Parreysia) Hanley	845
Confragosa (Unio) Deshayes	662	Corbis (Unio) Hanley	845
Confragosus (Arcidens) Say	662	Cordata (Castalia) H. and A. Adams	864
Confusa (Anodon) Heude	639	Cordata (Obovaria) Rafinesque	602
Congaræus (Margarita) Lea	709	Cordatus (Unio) Conrad	788
Congaræus (Margaron) Lea	709	Corderii (Unio) Heude	805
Congaræus (Unio) Lea	709	Coreana (Quadrula) von Martens	797
Conjugans (Tritogonia) B. H. Wright	609	Coreanus (Unio) von Martens	797
Conjugans (Unio) B. H. Wright	609	Coriaceus (Diplodon) Dunker	833
Conjungens (Arconia) Heude	860	Coriaceus (Unio) Dunker	883
Connasaugænsis (Margaritana) Lea	618	Corium (Unio) Reeve	702
Connasaugænsis (Margaron) Lea	618	Cornea (Anodonta) Philippi	648
Connasaugænsis (Strophitus) Lea	618	Corneola (Spatha) Rochebrune	902
Connasaughensis (Unio) Sowerby	618	Corneus (Unio) Lea	725
Connesaugensis (Margaritana) Clessin	618	Cornuta (Mya) Eaton	611
Conradianus (Margaron) Lea	589	Cornuta (Thecliderma) Swainson	611
Conradianus (Unio) Lea	589	Cornutus (Margarita) Lea	611
Conradicus (Margarita) Lea	589	Cornutus (Margaron) Lea	611
Conradicus (Margaron) Lea	589	Cornutus (Unio) Barnes	504, 611
Conradicus (Medionidus) Lea	589	Cornuum-lunæ cinnamomeus (Quadrula)	
Conradicus (Unio) Conrad	589	Heude	802
Conradicus (Unio) Lea	509, 588, 589	Cornuum-lunæ (Quadrula) Heude	802
Con sanguineus (Margaron) Lea	752	Cornuum-lunæ (Unio) Heude	802
Consanguineus (Unio) Lea	752	Corpulenta (Anodonta) Cooper	646
Consentaneus (Unio) Rossmässler	698	Corpulenta (Margaron) Lea	646
Consobrinus (Lamellidens) Lea	856	Correyana (Margaritana) Patet	661
Consobrinus (Unio) Hanley and Theobald	856	Corrianus (Lamellidens) Lea	857
Consobrinus (Unio) Lea	856	Corrianus (Margarita) Lea	857
Conspicuus (Unio) Lea	536	Corrianus (Margaron) Lea	857
Constrictus (Lampsilis) Conrad	551	Corrianus (Unio) Kuster	811
Constrictus (Margaron) Lea	551	Corrianus (Unio) Lea	857
Constrictus (Unio) Conrad	551	Corrianus (Unio) Sowerby	855
Contiguus (Margaron) Lea	548	Corrientesensis (Margarita) Lea	912
Contiguus (Unio) Lea	548	Corrientesensis (Margaron) Lea	912
Contorta (Burtonia) Bourguignat	578	Corrientesensis (Monocondylæa) d'Or-	
Contorta (Hyria) Sowerby	860	bigny	912
Contorta (Margaron) Lea	860	Corrugata (Alasmodon) De Kay	671
Contorta (Triquetra) Lea	860	Corrugata (Hyria) Lamarck	868
Contortus conjungens (Unio) Heude	860	Corrugata laevirostris (Parreysia) Benson	842

	Page.		Page.
Corrugata (Margaron) Lea	841, 869	Crassus (Margarita) Lea	539
Corrugata (Mya) Müller	841	Crassus (Margaron) Lea	678
Corrugata nagpoorensis (Parreysia) Lea	842	Crassus (Pseudodon) Drouet	834
Corrugata (Parreysia) Müller	841	Crassus (Unio) Barnes	767
Corrugata (Spatha) Dautzenberg	902	Crassus (Unio) Conrad	540
Corrugata (Triquetra) H. and A. Adams	869	Crassus (Unio) Kuster	536
Corrugata (Unio) Blainville	868	Crassus (Unio) Nilsson	697
Corrugata (Unio) Lamarek	841	Crassus (Unio) Reeve	692
Corrugata (Unio) Manduyt	698	Crassus (Unio) Retzius	677
Corrugatus lævirostris (Unio) Hanley and Theobald	842	Crassus (Unio) Say	539
Corrugatus (Margarita) Lea	841	Crassus (Unio) Sowerby	540, 707
Corrugatus nagpoorensis (Unio) Hanley and Theobald	842	Craviovensis (Psilunio)	510
Corrugatus (Unio) Retzius	691	Crebristriata (Margaron) Lea	835
Coruneulina	563	Crebristriata (Monoeondylæa) Anthony	835
Coruneus fryanus (Unio) B. H. Wright	714	Crebristriata (Trigonodon) Conrad	835
Coruseus (Margaron) Lea	714	Crebristriatus peguensis (Pseudodon) Anthony	835
Coruseus (Unio) Gould	714	Crebristriatus (Pseudodon) Anthony	835
Corvinus (Margaron) Lea	566	Crebristriatus (Unio) Sowerby	835
Corvinus (Unio) Lea	566	Crebrivittatus (Margaron) Lea	753
Corvuneulus (Lampsilis) Lea	566	Crebrivittatus (Unio) Lea	753
Corvuneulus (Margaron) Lea	566	Crenodonta Schlüter	766
Corvuneulus (Unio) Lea	566	Crepera (Anodonta) Lea	833
Corvus (Margaron) Lea	708	Crepera (Dallielia) Lea	833
Corvus (Unio) Lea	708	Crepera (Margaron) Lea	833
Costata (Alasmodonta) Rafinesque	664	Creperus (Margarita) Lea	552
Costata (Alasmodonta) Say	665	Creperus (Margaron) Lea	552
Costata (Amblema) Rafinesque	744	Creperus (Unio) Lea	552
Costata (Complanaria) Conrad	665	Cresserus (Unio) Hanley	552
Costata (Symphynta) Rafinesque	665	Crispa (Anodonta) Lamarek	919
Costatus (Plectomerus) Conrad	769	Crispata (Alasmodonta) Mousson	836
Costatus (Unio) Say	769	Crispata (Anodonta) Lamarek	919
Costulata (Margaron) Lea	912	Crispata (Anodontites) Bruguiere	919
Costulata (Monoeondylæa) Morieand	912	Crispata (Margarita) Lea	919
Couehiana (Quadrula) Lea	777	Crispata (Margaritana) Pætel	836
Couehianus (Margaron) Lea	777	Crispata (Margaron) Lea	919
Couehianus (Unio) Lea	777	Crispata (Nodularia) Gould	815
Coulboisi (Brazzeia) Bourguignat	907	Crispata (Unio) Gould	815
Coulboisi (Cameronia) Bourguignat	909	Crispatus (Glabaris) Bruguiere	919
Coulboisi (Unio) Bourguignat	821	Crispatus (Margaron) Lea	815
Couperiana (Anodonta) Lea	636	Crispisuleata (Nodularia) Benson	820
Couperiana (Margarita) Lea	636	Crispisuleatus (Margaron) Lea	820
Couperiana (Margaron) Lea	636	Crispisuleatus (Unio) Benson	508, 820
Courquinianus (Unio) Bourguignat	686	Cristaria Schumacher	505, 506, 514, 578, 583
Crapulus (Margaron) Lea	755	Croeatus (Margaron) Lea	531
Crapulus (Unio) Lea	755	Croeatus (Unio) Lea	531
Crassa (Dallielia) Drouet	834	Crocodilarum (Unio) Morelet	702
Crassa (Gibbosula) Wood	804	Cromwellii (Margaron) Lea	566
Crassa (Margarita) Lea	917	Cromwellii (Lampsilis) Lea	565
Crassa (Margaritana) Retzius	513, 677	Cromwellii (Unio) Lea	565
Crassa (Margaron) Lea	917	Crosseana (Castalia) Hidalgo	865
Crassa (Mya) Eaton	539	Crosseanus (Tetraplodon) Hidalgo	865
Crassa (Mya) Wood	804	Crossei (Physunio) Deshayes and Jullien	831
Crassidens (Margarita) Lea	707	Crossei (Unio) Deshayes and Jullien	831
Crassidens (Margaron) Lea	707	Cruda (Pleurobema) Lea	751
Crassidens (Unio) Lamarek	503, 504, 700, 706, 709	Crudus (Unio) Lea	751
Crassidens (Unio) Sowerby	539	Cryptoradiata (Spatha) Putzeys	900
Crassidens var. <i>a</i> (Plectomerus) Conrad	772	Cylindricus (Margaron) Say	773
Crassidens var. <i>a</i> (Unio) Lamarek	772	Ctenodesma Simpson	506, 515, 852
Crassissimus (Margarita) Lea	676	Cueumeria Conrad	893
Crassissimus (Unio) Hanley	678	Cueumoides (Margaron) Lea	893
Crassitesta Simpson	586	Cueumoides (Unio) Lea	893
Crassus (Anodon) Swainson	917	Cultelliformis (Diplodon) Conrad	892
Crassus batavus (Unio) Jordan	697	Cultelliformis (Unio) Conrad	892
Crassus (Glabaris) Swainson	917	Cultrata (Anodonta) Gould	644
		Cumberlandianus (Margarita) Lea	553

	Page.		Page.
Cumberlandianus (Margaron) Lea	553	Cylindracea (Margaron) Lea	920
Cumberlandicus (Margaron) Lea	554	Cylindrellus (Lampisilis) Lea	565
Cumberlandicus (Unio) Lea	509, 553	Cylindrellus (Margaron) Lea	565
Cumingianus (Unio) Dunker	893	Cylindrellus (Unio) Lea	565
Cumingii (Anodonta) Lea	837	Cylindrica (Mya) Eaton	773
Cumingii (Hyriopsis) Lea	579	Cylindrica (Quadrula) Say	773
Cumingii (Margaron) Lea	579, 837	Cylindrica Simpson	807
Cumingii (Microcondylæa) Clessin	837	Cylindricus (Margarita) Say	773
Cumingii (Monocondylæa) Lea	837	Cylindricus (Orthonymus) Agassiz	773
Cumingii (Pseudodon) Lea	837	Cylindricus (Unio) Say	773
Cumingii (Pseudodus) de Morgan	837	Cyphia (Obliquaria) Rafinesque	744
Cumingii (Unio) Lea	579	Cyphla (Unio) Conrad	764
Cuneata (Mya) Eaton	707	Cyphas asopus (Unio) Pætel	764
Cuneata (Pleurobema) Rafinesque	765	Cyphius (Unio) Say	764
Cuneatus (Unio) Barnes	707	Cyprogenia Agassiz	506, 510, 514, 609
Cuneatus (Unio) Rossmässler	692	Cyrenoides (Margaron) Lea	606
Cuneatus (Unio) Say	746	Cyrenoides (Plagiola) Philippi	606
Cuneatus (Unio) Swainson	517	Cyrenoides (Unio) Philippi	606
Cuneolus (Margaron) Lea	748	Cyrenopsis (Grandidieria) Bourgingnat	828
Cuneolus (Pleurobema) Lea	748	Dactylinus (Unio) Heude	808
Cuneolus (Unio) Lea	748	Dactylus (Anodon) Sowerby	929
Cuneopsis Simpson	506, 509, 514, 804	Dactylus (Margaron) Lea	553, 929
Cuneus (Quadrula) Conrad	792	Dactylus (Unio) Lea	553
Cuneus (Unio) Call	789	Dactylus (Unio) Morelet	683
Cunninghami (Unio) B. H. Wright	714	Dahomeyensis (Anodonta) Lea	900
Cuprea (Obliquaria) Rafinesque	744	Dahomeyensis (Margaron) Lea	900
Cuprinus (Diplodon) Simpson	883	Dahomeyensis (Spatha) Lea	900
Cuprinus (Margarita) Lea	573	Dahuricus (Margaritana) Kobelt	676
Cuprinus (Margaron) Lea	573	Dahuricus (Unio) Middendorff	676
Cuprinus (Unio) Lea	572	Dalei (Anodonta) Buffon	648
Curatus (Margaron) Lea	728	Dallasiana (Anodonta) Lea	647
Curatus (Unio) Lea	727	Dallasiana (Margaron) Lea	647
Curreyana (Baphia) H. and A. Adams	661	Dalli (Unio) B. H. Wright	713
Curreyana (Margaritana) Lea	661	Dalliella Simpson	506, 515, 832, 834
Curreyana (Margaron) Lea	661	Damalis Leach	674
Curreyana (Micromya) Agassiz	661	Damascensis (Margaron) Lea	695
Curreyana (Strophitus) Conrad	661	Damascensis (Unio) Lea	694
Curreyana (Unio) Sowerby	661	Damnoica (Unio) d'Orbigny	894
Curti (Pleurobema) Lea	753, 754	Daniells (Unio) Gassies	684
Curtus (Margaron) Lea	754	Daniellii (Unio) Villa	891
Curtus (Unio) Lea	754	Danielsii (Anodonta) Lea	645
Curvatus (Anodonta) Jay	648	Danielsii (Margaritana) Pætel	645
Curvatus (Unio) Lea	724	Danielsii (Margaron) Lea	645
Cuspidatus (Unio) Lea	728	Danielsii (Unio) B. H. Wright	708
Cuvierianus (Margaron) Lea	727	Dariensis (Anodonta) Lea	634
Cuvierianus (Unio) Lea	727	Dariensis (Margaron) Lea	634, 710
Cyanus (Unio) Philippi	861	Dariensis (Unio) Lea	710
Cyclips (Toxolasma) Rafinesque	744	Datus (Margaron) Lea	728
Cyclips (Unio) Ferussac	540	Datus (Unio) Lea	728
Cyclomya Simpson	578, 885, 886	Dautzenbergi (Nodularia) L. Morlet	819
Cyanea anatina (Anodonta) Linnaeus	625	Dautzenbergi (Unio) L. Morlet	819
Cyanea (Anodonta) Linnaeus	513, 620	Davilai (Unio)	510
	621, 626, 627, 631, 649	Decampsiana (Nodularia) Wattebled	822
Cyanea (Anodonta) Rossmässler	625	Decampsianus (Unio) Wattebled	822
Cyanea cellensis (Anodonta)	621	Decisa (Pleurobema) Lea	752
Cyanea (Margarita) Lea	622	Decisus (Margarita) Lea	752
Cyanea (Margaron) Lea	622	Decisus (Margaron) Lea	752
Cyanea ponderosa (Anodonta) C. Pfeiffer	626	Decisus (Unio) Kuster	748
Cyanea rostrata (Anodonta) Rossmässler	627	Decisus (Unio) Lea	752
Cyanea (Symphynota) Lea	622	Declivis (Anodonta) Conrad	643
Cygneus (Anodontes) Cuvier	621	Declivis (Margarita) Lea	643, 741
Cygneus (Mytilus) Linnaeus	620, 621	Declivis (Margaron) Lea	741
Cygnæa (Anodonta) Gassies	622	Declivis (Unio) Conrad	742
Cylindracea (Anodonta) Lea	920	Declivis (Unio) Say	740
Cylindracea (Glabaris) Lea	920	Decora (Anodonta) Lea	643
Cylindracea (Margarita) Lea	920	Decora (Margarita) Lea	643

	Page.		Page.
Decora (Margaron) Lea	644	Deviata (Truncilla) Reeve	524
Decorata (Symphynota) Lea	664	Deviatus (Margaron) Lea	524
Decoratus (Margaron) Lea	664	Deviatus (Unio) Reeve	524
Decoratus (Unio) Lea	664	Diagenæ	514, 615
Decumbens (Margaritana) Lea	679	Dianisotis Rafinesque	583
Decumbens (Margaron) Lea	679	Diaphanus (Unio) Rafinesque	744
Decumbens (Unio) Lea	679	Diazensis (Unio) S. H. Wright	714
Decurvatus (Unio) Rossmässler	698	Diespiter (Unio) Mabilie	861
Dehiscens (Anodonta) Patet	655	Differtus (Unio) Lea	729
Dehiscens (Baphia) H. and A. Adams	655	Difficilis (Margaron) Lea	554
Dehiscens (Hemilastena) Agassiz	655	Difficilis (Unio) Lea	554
Dehiscens (Margarita) Lea	654	Digeneæ	514, 607
Dehiscens (Margaritana) Clessin	655	Digitata (Quadrula) Morelet	770
Dehiscens (Margaron) Lea	655	Digitatus (Unio) Morelet	770
Dehiscens (Unio) Say	654	Digitiformis (Nodularia) Sowerby	809
Dejecta (Anodonta) Lewis	630	Digitiformis (Unio) Sowerby	809
Delaportei (Hyriopsis) Crosse and Fischer	580	Dignatus (Margaron) Lea	688
Delaportei (Unio) Crosse and Fischer	580	Dignatus (Unio) Lea	688
Dellesserti (Margaron) Lea	693	Dignota (Anodonta) Ferussac	631
Dellesserti (Unio) Bourguignat	692	Dignota (Anodonta) Rafinesque	648
Delicatus (Margaron) Lea	690	Dilatata (Unio) Rafinesque	744
Delicatus (Unio) Lea	690	Dilatatus (Unio) Say	704
Delodonta (Unio) Lamarck	873	Diminuta (Nodularia) Lea	826
Delodontus (Diplodon) Lamarck	873	Diminutus (Margaron) Lea	826
Delodontus (Margarita) Lea	873	Diminutus (Unio) Lea	826
Delodontus (Unio) Sowerby	539	Dimota (Nodularia) Lea	818
Delphinulus (Lampsilis) Morelet	576	Dimotus (Margaron) Lea	818
Delphinulus (Unio) Morelet	576, 577	Diolibanus (Pleiodon) Bourguignat	908
Delphinus (Margaron) Lea	579	Diploderma (Bariosta) Rafinesque	744
Delphinus (Unio) Grüner	579	Diplodon Spix	506, 515, 867, 872, 873, 883
Delphinus (Unio) Spengler	744	Diplodon (Unio) Philippi	882
Deltoidea (Alasmodonta) Baker	668	Dipsas (Anodonta) Blainville	584
Deltoidea (Baphia) H. and A. Adams	668	Dipsas Leach	583
Deltoidea (Margarita) Lea	668	Dipsax Voigt	583
Deltoidea (Margaritana) Lea	668	Discoidea (Barbala) H. and A. Adams	585
Deltoidea (Margaron) Lea	668	Discoformis (Zaira) Rochebrune	862
Deltoidea (Strophitus) Conrad	668	Discoidea (Cristaria) Lea	585
Deltoidea (Unio) Hanley	668	Discoidea (Dipsas) Kuster	585
Delumbis (Lampsilis) Conrad	531	Discoidea (Margarita) Lea	585
Delumbis (Unio) Conrad	531	Discoidea (Symphynota) Lea	585
Delumbis (Unio) Kuster	559	Discoideus (Margaron) Lea	585
Demararaensis (Diplodon) Lea	880	Discoideus (Unio) Hanley	585
Demararaensis (Margaron) Lea	880	Discoideus (Unio) Sowerby	574
Demararaensis (Unio) Lea	880	Discomya Simpson	802
Dembeæ (Nodularia) Reeve	826	Discrepans (Margaron) Lea	554
Dembeæ (Unio) Reeve	826	Discrepans (Unio) Lea	554
Denigrata (Anodonta) Lea	659	Disculus (Diplodon) Lea	887
Denigrata (Margaron) Lea	660	Disculus (Margaron) Lea	887
Denigratus (Margaron) Lea	718	Disculus (Unio) Lea	887
Denigratus (Unio) Lea	718	Disens (Margaron) Lea	705
Depauperatus (Unio) Hutton	890	Discus (Unio) Lea	705
Depressa (Mya) Donovan	685	Discus (Unio) Sowerby	707
Depressa (Leguminifolia) Westerlund	653	Dispalans (Unio) B. H. Wright	736
Depressa (Obliquaria) Rafinesque	603	Dispanus (Unio) Lea	549
Depressa (Unio) C. Pfeiffer	653	Dispar (Lampsilis) Lea	561
Depressa (Unio) d'Orbigny	894	Dispar (Margaron) Lea	561
Depressa (Unio) Lamarck	603, 892	Dispar (Unio) Lea	561
Depressus (Margarita) Lea	891	Dissimilis (Unio) Lea	730
Depressus (Margaron) Lea	891	Distans (Unio) Anthony	534
Depressus (Unio) Lamarck	892	Distortus (Unio) Bean	714
Depressus (Unio) Lesson	891	Distortus (Unio) Heude	807
Depressus (Unio) Patet	685	Divaricata (Spatha) von Martens	901
Depressus (Unio) Reeve	891	Divaricatus (Diplodon) Lea	878
Depygis (Unio) Conrad	601	Divaricatus (Margaron) Lea	878
Deshayesi (Unio) Michaud	682	Divaricatus (Unio) Lea	878
Despecta (Anodon) Heude	638	Divergens (Quadrula) Benson	802

	Page.		Page.
Diversens (Unio) Benson	802	Duclerei (Chamberlainia) Rochebrune ..	582
Diversa (Alasmidonta) Conrad	669	Duclerei (Unio) Rochebrune	582
Diversus (Unio) Conrad	669	Dugasti (Unio) Morlet	811
Dolabella (Castalia) Sowerby	867	Dumesleanus (Unio) Charnes	847
Dolabelloides (Margaron) Lea	750	Dunkerianus (Unio) Lea	882
Dolabelloides (Pleurobema) Lea	750	Dunlapiana (Anodonta) Lea	635, 636
Dolabelloides (Unio) Lea	750	Duponti (Parreysia) Rochebrune	846
Dolabraformis (Lampsilis) Lea	529	Duponti (Reneus) Rochebrune	846
Dolabraformis (Margarita) Lea	529	Duponti (Unio) Rochebrune	846
Dolabraformis (Margaron) Lea	529	Duprei (Callonaiia) Recluz	867
Dolabraformis (Unio) Lea	529	Duprei (Castalia) Recluz	867
Dolabraformis (Unio) Sowerby	527	Duprei (Margaron) Lea	867
Dolearis (Anodonta) Lea	634	Dupuyi (Anodonta) Ray and Drouet	624
Doliaris (Anodonta) Lea	634	Durieui kullethensis (Unio) Lea	695
Doliaris (Lampsilis) Lea	533	Durieui (Margaron) Lea	694
Doliaris (Margaron) Lea	533, 634	Durieui (Unio) Deshayes	694
Doliaris (Unio) Lea	533	Duttonianus (Margaron) Lea	736
Dolichorhynchus (Unio) Tapperone- Canefri	861	Duttonianus (Unio) Lea	736
Doliolum (Anodonta) Heude	640	Dysnomia (Agassiz)	521
Dolosus (Margaron) Lea	569	Dysonii (Nephronaias) Lea	594
Dolosus (Unio) Lea	568	Dysonii (Margaron) Lea	594
Dombeyana (Unio) Valenciennes	772	Dysonii (Unio) Lea	594
Donaciformis (Margarita) Lea	605	Ebenus (Margarita) Lea	793
Donaciformis (Margaron) Lea	605	Ebenus (Margaron) Lea	793
Donaciformis (Plagiola) Lea	605	Ebenus (Quadrula) Lea	793
Donaciformis (Unio) Lea	605	Ebenus (Unio) Lea	509, 793
Dooleyensis (Unio) Lea	727	Ecarinata (Castalia) Mousson	865
Dorei (Unio) B. H. Wright	710, 719	Ecarinatus (Tetraplodon) Mousson	865
Dorfeuillanus (Margarita) Lea	780	Edentula (Alasmodonta) Say	616
Dorfeuillanus (Margaron) Lea	780	Edentula (Anodonta) Ferussac	616
Dorfeuillanus (Unio)	779, 780	Edentula (Anodonta) Say	503, 504
Dorri (Nodularia) Wattebled	809	Edentula (Margarita) Lea	616
Dorri (Unio) Wattebled	809	Edentula (Margaron) Lea	616
Dorsatus (Margaron) Lea	711	Edentula (Uniopsis) Agassiz	616
Dorsatus (Unio) Lea	711	Edentulus (Anodon) Sowerby	643
Dorsuosus (Diplodon) Gould	888, 889	Edentulus Pavonius (Strophitus) Lea	617
Dorsuosus (Margaron) Lea	889	Edentulus (Strophitus) Say	616
Dorsuosus (Unio) Gould	889	Edgarianus (Margaron) Lea	747
Douglasia nipponensis (Nodularia) von Martens	809	Edgarianus (Pleurobema) Lea	747
Douglasia (Nodularia) Gray	807, 808, 861	Edgarianus (Unio) Lea	747
Douglasia Shanghaiensis (Unio) Lea	808	Edulis (Anodonta) Heude	638
Douglasia (Unio) Gray	806, 807, 808	Efulgens (Diplodon) Lea	879
Doumeti (Unio) Letourneux and Bour- guignat	744	Efulgens (Margaron) Lea	879
Downiei (Margaron) Lea	733	Efulgens (Unio) Lea	879
Downiei (Unio) Lea	732, 733	Egyptiacus (Unio) Hanley	821
Draparnaldi (Unio) Deshayes	692	Eighti (Unio) Pætel	771
Dromauxi (Cameronia) Bourguignat	910	Eightisi (Margaron) Lea	771
Dromauxi (Unio) Bourguignat	821	Eightisi (Unio) Lea	771
Dromus (Cyprogenia) Agassiz	615	Elachista (Anodonta) Bourguignat	625
Dromus (Dromus) Lea	615	Electrinus (Unio) Reeve	741
Dromus (Margarita) Lea	615	Elegans (Margarita) Lea	604
Dromus (Margaron) Lea	615	Elegans (Margaron) Lea	604
Dromus Simpson	510, 514, 614	Elegans (Plagiola) Lea	604, 605
Dromus (Unio) Lea	614, 615	Elegans (Unio) Lea	604
Droueti (Spatha) Chaper	897	Elegans (Zaira) Rochebrune	862
Droueti (Unio) Dupuy	699	Elliotii (Margaritana) Lea	619
Dubia (Alasmodonta) Ferussac	673	Elliotii (Margaron) Lea	619, 769
Dubia (Anodonta) Bose	903	Elliotii (Quadrula) Lea	763
Dubia (Iridina) Chenu	903	Elliotii (Strophitus) Lea	619
Dubia (Mulela) Gmelin	903	Elliotii (Unio) Lea	768
Dubia (Mytilus) Gmelin	583, 903	Ellipsaria (Obliquaria) Rafinesque	744
Dubia (Platiris) Lea	903	Ellipsarius (Unio) Say	539
Dubius (Unio) Fitzinger	682	Ellipsiformis (Lampsilis) Conrad	557
		Ellipsiformis (Unio) Conrad	557
		Ellipsis (Margarita) Lea	602
		Ellipsis (Margaron) Lea	602

	Page.		Page.
Ellipsis (Obovaria) Lea	540, 602	Estabrookiana (Pleurobema) Lea	763
Ellipsis (Unio) Lea	601, 602, 754	Estabrookianus (Margaron) Lea	763
Ellipsoideus (Unio) Locard	690	Estabrookianus (Unio) Lea	763
Elliptica (Anodon) Heude	638	Esula (Iridina) d'Orbigny	915
Elliptica (Margaritana) Lea	619	Esula (Leila) d'Orbigny	915
Elliptica (Margaron) Lea	619	Esula (Margarita) Lea	915
Elliptica (Mya) Eaton	539	Esula (Margaron) Lea	915
Elliptica (Unio) Rafinesque	744	Etawahensis (Margaritana) Conrad	672
Ellipticum (Diplodon) Spix	872, 873, 877	Etawahensis (Margaritana) Lea	670
Ellipticum (Pseudodon) Conrad	836	Eucirrus (Margaron) Lea	688
Ellipticus (Margaron) Lea	877	Eucirrus (Unio) Bourguignat	683
Ellipticus (Unio) Barnes	589	Eucyphus (Unio) Bourguignat	821, 822
Ellipticus (Unio) Wagner	877, 879	Eufira Gistel	902
Elliptio Rafinesque	679, 700	Euphratica Churchillianus (Gabillotia) Bourguignat	650
Elongata (Anodonta) Potiez & Michaud	623	Euphratica (Gabillotia) Bourguignat	650
Elongata (Brazzea) Bourguignat	907	Euphratica (Leguminaria) Westerlund	650
Elongata (Burtonia) Bourguignat	578	Euphratica (Margaritana) Kobelt	650
Elongata (Damaris) Leach	676	Euphratica (Pseudodon) Conrad	650
Elongata (Grandidieria) Bourguignat	830	Euphymus (Unio) Charmes	847
Elongata (Hyria) Swainson	871	Eurhynchus (Unio) Kuster	879
Elongata (Iridina) Sowerby	903	Europe (Anodonta) Leach	624
Elongata (Margarita) Lea	928	Eurynia Rafinesque	534
Elongata (Mutela) Pætel	903	Euscaphys (Anodonta) Heude	640
Elongata (Platiris) Lea	903	Evanescens (Unio) Mousson	855
Elongata (Triquetra) H. and A. Adams	871	Evansi (Diplodon) Adams and Angas	892
Elongata (Unio) Goodrich	536	Evansi (Margaritana) Pætel	893
Elongata (Unio) Lamarck	676	Evansi (Margaron) Lea	893
Elongata (Unio) Michaud	697	Evansi (Unio) Adams and Angas	892
Elongatula (Unio) C. Pfeiffer	684	Evitatus (Margaron) Lea	812
Elongatulus (Margarita) Lea	684	Evitatus (Unio) Lea	811
Elongatulus (Margaron) Lea	684	Exactus (Margaron) Lea	723
Elongatulus (Unio) C. Pfeiffer	513-684	Exactus (Unio) Lea	723
Elongatus (Anodon) Swainson	928	Exactus (Unio) Lea	737
Elongatus (Glabaris) Swainson	928	Exanthematicus (Lamellidens) Kuster	856
Elongatus (Pleiodon) Bourguignat	908	Exanthematicus (Unio) Kuster	856
Emarginata (Solenaria) Lea	655	Exasperata (Hyria) Sowerby	869
Emarginatus (Anodonta) Clessin	655	Excavatus (Lampisilis) Lea	528
Emarginatus (Margarita) Lea	696	Excavatus (Unio) Lea	528
Emarginatus (Margaron) Lea	696	Excultus (Unio) Conrad	740
Emarginatus (Mycetopus) Lea	655	Excurvata (Anodonta) De Kay	633
Emarginatus (Platiris) Lea	655	Exiguus (Margaron) Lea	560
Emarginatus (Solenaria) Lea	656	Exiguus (Unio) Lea	560
Emarginatus (Unio) Lea	695	Exilior (Anodonta) Lea	630
Embia (Anodonta) Bourguignat	627	Exilis (Anodon) Catlow and Reeve	588
Emesensis (Margaron) Lea	693	Exilis (Anodon) Sowerby	633
Emesensis (Unio) Lea	693	Exilis (Anodonta) Lea	587
Emini (Nodularia) von Martens	823	Exilis (Margarita) Lea	588
Emini (Unio) von Martens	823	Exilis (Margaron) Lea	588, 818
Emmonsii (Margaron) Lea	735	Exilis (Monocondylæa) Pætel	588
Emmonsii (Unio) Lea	735	Exilis (Monocondylus) Morelet	839
Encarpa (Plagiola) Lea	607	Exilis (Pilsbryconcha) Lea	557
Encarpus (Margaron) Lea	607	Exilis (Pseudodon) Morelet	839
Encarpus (Unio) Lea	607	Exilis (Unio) Dunker	818
Endobranchiæ	515, 806	Eximia (Anodonta) Bourguignat	649
Ensiformis (Anodon) Spix	932	Eximia (Brazzea) Bourguignat	907
Ensiformis (Glabaris) Spix	931, 932	Eximium (Colletopterum) Bourguignat	649
Ensiformis (Iridina) Lea	932	Eximius (Margaron) Lea	831
Ensiformis (Margarita) Lea	932	Eximius (Physunio) Lea	831
Ensiformis (Margaron) Lea	932	Eximius (Unio) Lea	831
Episcopalis (Unio) Tristram	694	Exobranchiæ	514
Errans (Margaron) Lea	730	Exolescens (Margaron) Lea	859
Errans (Unio) Lea	730	Exolescens (Trapezoideus) Gould	859
Eschatigenæ	514, 614	Exolescens (Unio) Gould	859
Essoensis (Nodularia) Chaper	822	Exotica (Anodon) Sowerby	927
Essoensis (Unio) Chaper	822	Exotica (Anodonta) Blainville	903
Estabrookianus (Unio) Pætel	763		

	Page.		Page.
Exotica (Anodonta) Lamarck.....	916	Favidens pinax (Parreysia) Benson.....	843
Exotica (Columba) Pætel.....	924	Favidens pinax (Unio) Hanley and Theo- bald.....	843
Exotica (Iridina) Children.....	908	Favidens plagiosoma (Parreysia) Benson.....	843
Exotica (Iridina) Lamarck.....	902, 903	Favidens plagiosoma (Unio) Hanley and Theobald.....	843
Exotica (Mutela) Lamarck.....	903	Favidens (Unio) Lea.....	842
Exotica (Platiris) Lea.....	903	Favosa (Pleurobema) Lea.....	761
Exoticus (Anodonta) Lamarck.....	924	Favosus (Margaron) Lea.....	761
Exoticus (Pleiodon) Pætel.....	903	Favosus (Unio) Lea.....	761
Expansus (Diplodon) Kuster.....	874	Fea (Parreysia) Tapperone-Canefri.....	844
Expansus (Unio) Kuster.....	874	Fea (Unio) Tapperone-Canefri.....	844
Explicatus (Lampsilis) Morelet.....	571	Feddeni (Parreysia) Theobald.....	843
Explicatus (Unio) Morelet.....	571	Feddeni (Unio) Theobald.....	843
Extensus (Margaron) Lea.....	731	Fellmani (Margaron) Lea.....	692
Extensus (Unio) Lea.....	731	Fellmani (Unio) Deshayes.....	692
Exulcerata (Anodonta) Porro.....	623	Feminalis (Anodonta) Gould.....	657, 658
Faba (Margarita) Lea.....	877	Fenouilli (Anodonta) Heude.....	639
Faba (Margaron) Lea.....	877	Ferrarisii (Anodonta) d'Orbigny.....	918
Faba (Unio) d'Orbigny.....	877	Ferrissii (Unio) Marsh.....	719
Fabaceus (Margaron) Lea.....	549	Ferruginea (Anodonta) Lea.....	659
Fabaceus (Unio) Lea.....	549	Ferruginea (Margaron) Lea.....	659
Fabagina (Parreysia) Deshayes.....	847	Ferussaciana (Anodonta) Ferussac.....	617
Fabagina (Unio) Deshayes.....	847	Ferussaciana (Anodonta) Lea.....	504, 658, 659
Fabalis (Margaron) Lea.....	525	Ferussaciana (Margarita) Lea.....	659
Fabalis (Micromya) Lea.....	525	Ferussaciana (Margaron) Lea.....	659
Fabalis (Margarita) Lea.....	525	Ferussacianus (Anodontoides) Lea.....	659
Fabalis (Unio) Lea.....	524, 525	Ferussacianus (Margaron) Lea.....	694
Fabula (Baphia) H. and A. Adams.....	661	Ferussacianus modestus (Anodontoides) Lea.....	660
Fabula (Margarita) Lea.....	661	Ferussacianus subeylindraceus (Anodon- toides) Lea.....	660
Fabula (Margaritana) Lea.....	660, 661	Ferussacianus (Unio) Lea.....	694
Fabula (Margaron) Lea.....	661	Fibrosa (Quadrula) Heude.....	798
Fabula (Micromya) Agassiz.....	661	Fibrosus (Unio) Heude.....	798
Fabula (Pegias) Lea.....	661	Fibuloides (Margaron) Lea.....	756
Fabula (Strophitus) Conrad.....	661	Fibuloides (Unio) Lea.....	756
Fabula (Unio) Hanley.....	661	Filippiana (Anodon) Heude.....	639
Faidherbi (Reneus) Jousseume.....	822	Firmus boettgeri (Diplodon) von Ihering.....	874
Falcata (Alasmodon) Gould.....	677	Firmus boettgeri (Unio) von Ihering.....	874
Falcata (Solenaia) Higgins.....	656	Firmus (Diplodon) Lea.....	874
Falcatus (Mycetopus) Higgins.....	656	Firmus (Margaron) Lea.....	874
Falcatus (Platiris) Lea.....	656	Firmus (Unio) Heude.....	808
Falcatus (Unio) Sowerby.....	677	Firmus (Unio) Lea.....	874
Fallaciosus (Lampsilis) Simpson.....	544	Fiscallianus (Unio) Kleciach.....	683
Fallax (Margaron) Lea.....	548	Fischeriana (Nodularia) L. Morlet.....	818
Fallax (Unio) Lea.....	548	Fischerianus (Unio) L. Morlet.....	818
Falsus (Glabaris) Simpson.....	932	Fisherianus (Margarita) Lea.....	735
Famelicus (Margaron) Lea.....	879	Fisherianus (Margaron) Lea.....	735
Famelicus (Unio) Gould.....	879	Fisherianus (Unio) Lea.....	534, 546, 735
Fantoziana (Anodon) Heude.....	639	Fissidens (Nodularia) Böttger.....	827
Fasciata (Unio) Rafinesque.....	744	Fissidens (Unio) Böttger.....	827
Fasciatus (Unio) Say.....	539	Flarens (Unio) Pætel.....	614
Fasciola (Lampsilis) Rafinesque.....	577	Flava (Anodon) Anthony.....	632
Fasciolaris (Obliquaria) Rafinesque.....	744	Flava (Obliquaria) Rafinesque.....	744
Fasciolaris (Unio) Conrad.....	612	Flavescens (Margaron) Lea.....	614
Fasciolus (Unio) Conrad.....	556	Flavescens (Ptychobranchus) Lea.....	614
Fasciolus (Unio) Ferussac.....	527	Flavescens (Unio) Lea.....	614
Fasciolus (Unio) Say.....	532	Flavidens (Unio) Pætel.....	842
Fassinans (Margaron) Lea.....	762	Flavidulus (Margaron) Lea.....	759
Fassinans (Pleurobema) Lea.....	762	Flavidulus (Pleurobema) Lea.....	759
Fassinans rhomboidea (Pleurobema) Simpson.....	762	Flavidulus (Unio) Lea.....	749
Fassinans (Unio) Lea.....	762	Flavus rubiginosus (Unio) Pætel.....	786
Fatuus (Lampsilis) Lea.....	553	Flavus (Unio) Conrad.....	786
Fatuus (Margaron) Lea.....	553	Flexuosa (Dysnomia) Agassiz.....	521
Fatuus (Unio) Lea.....	553		
Favidens marces (Unio) Benson.....	843		
Favidens (Margaron) Lea.....	842		
Favidens (Parreysia) Benson.....	842, 843		

	Page.		Page.
Flexuosa (Obliquaria) Rafinesque	744	Fouladougouensis (Nodularia) Jousseau-	
Flexuosa (Quadrula) Simpson	790	ume	823
Flexuosus (Unio) Conrad	521	Fouladougouensis (Reneus) Jousseau.	823
Flexus (Toxolasma) Rafinesque	744	Fragilis (Anodonta) Fitzinger	648
Florentina (Truncilla) Lea	524	Fragilis (Anodonta) Lamarck	632, 633
Florentinus (Margaron) Lea	524	Fragilis gracilis (Unio) Pætel	574
Florentinus (Unio) Lea	524	Fragilis (Lasmonus) Rafinesque	744
Florida (Anodon) Heude	639	Fragilis (Margarita) Lea	632
Floridensis (Margaron) Lea	544	Fragilis (Margaritana) Kuster	836
Floridensis (Unio) Lea	544	Fragilis (Margaron) Lea	632
Floridensis (Unio) S. H. Wright	560	Fragilis (Metaptera) Conrad	574
Fluetiger (Margaron) Lea	814	Fragilis (Microcondylæa) Pætel	836
Fluctiger (Nodularia) Lea	814	Fragilis (Symphynota) Ferussac	574
Fluctiger (Unio) Lea	814	Fragilis (Unio) Nevill	858
Fluminea (Anodonta) Heude	640	Fragilis (Unio) Rafinesque	744
Fluviatilis (Anodonta) Bose	631	Fragilis (Unio) Sowerby	883
Fluviatilis (Margarita) Lea	631	Fragilis (Unio) Swainson	573
Fluviatilis (Margaron) Lea	631	Fragosa (Quadrula) Conrad	777
Fluviatilis (Mytilus) Dillwyn	631	Fragosus (Margarita) Lea	777
Fluviatilis (Unio) Green	721	Fragosus (Margaron) Lea	777
Flyensis (Diplodon) Tapperone-Canefri ..	890	Fragosus (Unio) Conrad	777
Flyensis (Unio) Tapperone-Canefri	890	Franciscana (Aplodon) Lea	912
Fokkesi (Unio) Dunker	894	Franciscana (Margarita) Lea	912
Foliacea (Unio) Gould	858	Franciscana (Margaron) Lea	912
Foliaceus comptus (Trapezoides) Des-		Franciscana (Monocondylæa) Moricand ..	912
hayes and Jullien	858	Franciscana (Unio) Moricand	912
Foliaceus (Margaron) Lea	858	Fraternus (Margaron) Lea	709
Foliaceus (Trapezoides) Gould	858	Fraternus (Unio) Lea	709
Foliata (Truncilla) Hildreth	521	Frenzelli (Diplodon) von Ihering	882
Foliatus (Margarita) Lea	521	Frenzelli (Unio) von Ihering	882
Foliatus (Margaron) Lea	521	Frierioni (Quadrula) B. H. Wright	787
Foliatus (Unio) Hildreth	521	Frierioni (Unio) B. H. Wright	787
Folliculatus (Margarita) Lea	737	Friniana (Anodon) Heude	638
Folliculatus (Margaron) Lea	737	Fryanus (Unio) B. H. Wright	714
Folliculatus (Unio) Lea	737	Fucatus (Margaron) Lea	542
Foncki (Unio) Philippi	882	Fucatus (Mytilus) Dillwyn	622
Fontaineana (Unio) d'Orbigny	886	Fucatus (Unio) Lea	542
Fontainiana (Unio) d'Orbigny	886	Fulgens (Lampsilis) Rafinesque	577
Fontainianus (Diplodon) d'Orbigny	886	Fulgidus (Margaron) Lea	789
Fontainianus (Margaron) Lea	886	Fulgidus (Unio) Lea	789
Fontanus (Unio) Conrad	547	Fuliginosus (Margaron) Lea	727
Fóntinalis (Unio) Rafinesque	744	Fuliginosus (Unio) Lea	722, 727
Footei (Arcidopsis) Theobald	861	Fuligo (Unio) Lea	551
Footei (Unio) Theobald	861	Fulmineus (Unio) Philippi	841
Footiana (Anodonta) Lea	625, 642	Fulvaster (Nodularia) Drouet and Cha-	
Footiana (Margaron) Lea	642	per	819
Forbesiana (Anodonta) Lea	927	Fulvaster (Unio) Drouet and Chaper	819
Forbesiana (Margaron) Lea	707, 927	Fulvus (Margarita) Lea	709
Forbesianus (Glabaris) Lea	927	Fulvus (Margaron) Lea	709
Forbesianus (Unio) Lea	707	Fulvus (Unio) Lea	709
Foremanianus (Margaron) Lea	613	Fulvus (Unio) Rafinesque	744
Foremanianus (Ptychobanchus) Lea	613	Fumatus (Margaron) Lea	726
Foremanianus (Unio) Lea	613	Fumatus (Unio) Lea	726
Formosus (Unio) Lea	517	Funki (Unio) Pætel	882
Forscalli (Unio) Parreys	862	Funeralis (Diplodon) Lea	887
Forsheyi (Margaron) Lea	778	Funeralis (Margaron) Lea	887
Forsheyi (Quadrula) Lea	778	Funeralis (Unio) Lea	885, 886, 887
Forsheyi (Unio) Lea	778	Fureatum (Diplodon) Spix	871
Fossiculifera (Anodonta) Pætel	914	Furva (Pleurobema) Conrad	758
Fossiculifera (Fossula) d'Orbigny	914	Furvus (Unio) Conrad	758
Fossiculifera (Margarita) Lea	914	Fusca (Anodon) Heude	638
Fossiculifera (Margaron) Lea	914	Fuscatus (Margaron) Lea	717
Fossiculifera (Monocondylæa) Lea	914	Fuscatus (Unio) Lea	717
Fossiculiferus (Unio) Catlow and Reeve ..	914	Fusconaia Simpson	784
Fossula Lea	507, 515, 914	Fusculus (Unio) Rossmässler	698

	Page.		Page.
Gabbianus (Margaron) Lea	607	Gibbosa (Amblema) Rafinesque.....	703, 744
Gabbianus (Unio) Lea.....	607	Gibbosa (Anodonta) Say	635
Gabillotia Servain.....	506, 514, 649	Gibbosa dunlapiana (Anodonta) Lea	636
Gabonensis (Margaron) Lea.....	823	Gibbosa (Dysnomia) Agassiz	523
Gabonensis (Nodularia) Kuster.....	823	Gibbosa (Margarita) Lea.....	636
Gabonensis (Unio) Kuster.....	823	Gibbosa (Margaron) Lea.....	636
Gaditana (Mya) Schreibers	841	Gibbosa (Mya) Eaton	703
Galilaei (Unio) Locard.....	693	Gibbosula Simpson.....	506, 509, 514, 804
Gallandi (Anodonta) Bourguignat.....	648	Gibbosus areus (Unio) Conrad	707
Gancinensis (Spatha) Rochebrune.....	902	Gibbosus armathwaitensis (Unio) B. H. Wright	704
Gangrenosus (Unio) Schmidt	698	Gibbosus delicatus (Unio) Simpson	704
Gargottæ (Unio) Philippi.....	678	Gibbosus (Margarita) Lea	703
Gargottæ (Unio) Rossmässler	698	Gibbosus (Margaron) Lea	703
Gassiesi (Diplodon) Kuster.....	881	Gibbosus perobliquus (Unio) Conrad.....	520
Gassiesi (Unio) Kuster	881	Gibbosus perplexus (Unio) Patel	523
Gastonensis (Margaron) Lea	723	Gibbosus subgibbosus (Unio) Lea	704
Gastonensis (Unio) Lea	723	Gibbosus (Unio)	510
Gaudichaudi (Margaron) Lea	812	Gibbosus (Unio) Barnes.....	504, 545, 703, 707
Gaudichaudi (Nodularia) Eydoux	812	Gibbosus (Unio) Ferussac.....	523
Gaudichaudi (Unio) Eydoux.....	812	Gibbum (Anodon) Benson	638
Geddingsianus (Margaron) Lea	715	Gibbus (Unio) Spengler	862
Geddingsianus (Unio) Lea	715	Gigantea (Anodonta) Lea.....	643
Geminus (Margaron) Lea	733	Gigantea (Cameronia) Bourguignat.....	909
Geminus (Unio) Lea	733	Gigantea (Columba) Patel.....	923
Generosus (Lamellidens) Gould.....	857	Gigantea (Margarita) Lea	643
Generosus (Margaron) Lea.....	857	Gigantea (Margaron) Lea	643
Generosus (Unio) Gould	857	Giganteus (Anodon) Spix	916, 923
Genearethensis (Unio) Locard	690	Giganteus (Unio) Lea	767
Genthii (Margaron) Lea.....	551	Gigas (Complanaria) Sowerby	666
Genthii (Unio) Lea.....	551	Gigas (Megadomus) Swainson.....	608, 666
Gentiliana (Barbala) Heude	586	Gigas (Unio) Sowerby	608
Gentiliana (Cristaria) Heude	586	Giraudi (Cameronia) Bourguignat	910
Genuinus (Margaron) Lea	548	Giraudi (Grandidieria) Bourguignat	890
Genuinus (Unio) Lea.....	548	Giullaini (Anodonta) H. and A. Adams..	933
Geometricus (Unio) Lea.....	740	Glabaris Gray	501,
Georgiana (Alasmidonta) Lea.....	670	507, 515, 648, 914, 916, 918, 919, 923, 926, 930, 932	
Georgiana (Margaritana) Lea	670	Glaber (Margarita) Lea	554
Georgiana (Margaron) Lea	670	Glaber (Margaron) Lea.....	554
Georgiana (Pleurobema) Lea	761	Glaber (Unio) Lea.....	554
Georgianæ (Anodonta) Griffith	927	Glabra (Anodonta) Stable	623
Georgianæ (Glabaris) Griffith	927	Glabrata (Unio) Lamarek	721
Georgianus (Margaron) Lea.....	761	Glabratus (Unio) Sowerby	667
Georgianus (Unio) Lea.....	761	Glabus (Anodon) Sowerby	922
Georgina (Unio) Lamarek.....	721	Gladiator (Unio) Aney	862
Gerbidoni (Unio) Eydoux.....	811	Gladiolus (Nodularia) Heude	807
Gerhardtii (Lampsilis) Lea.....	532	Gladiolus (Unio) Heude	807
Gerhardtii (Margaron) Lea	532	Glandacea (Quadrula) Lea.....	793
Gerhardtii (Unio) Lea.....	532	Glandaceus (Margaron) Lea	793
Germanus (Lampsilis) Lea	565	Glandaceus (Unio) Lea	793
Germanus (Margaron) Lea	565	Glandulosa (Anodonta) Anthony	632
Gerrardi (Unio) von Martens	822	Glans (Lampsilis) Lea	565
Gesneri (Anodonta) Patel.....	634	Glans (Margarita) Lea	565
Gesnerii (Anodonta) Lea	634	Glans (Margaron) Lea.....	565
Gesnerii (Margaritana) Lea	619	Glans (Unio) Lea	565
Gesnerii (Margaron) Lea.....	619, 634	Glaucia (Anodonta) Valenciennes.....	926
Gesnerii (Strophitus) Lea	619	Glaucia cinaloensis (Anodonta) Crosse and Fischer	926
Gesnerii (Unio) Lea	727	Glaucia (Margarita) Lea	926
Gianelli (Unio) Tapperone-Canefri.....	862	Glaucia (Margaron) Lea	926
Gibba (Margaron) Lea	638	Glaucia (Monocondylæa) H. and A. Adams	926
Gibber (Margarita) Lea	762	Glaucinus (Unio) Porro	698
Gibber (Margaron) Lea	762	Glaucus cinaloensis (Glabaris) Crosse and Fischer	926
Gibber (Pleurobema) Lea	762	Glaucus (Glabaris) Valenciennes.....	924, 926
Gibber (Unio) Lea.....	762		
Gibbesianus (Margaron) Lea.....	732		
Gibbesianus (Unio) Lea	732		

	Page.		Page.
Glebula Conrad	506, 514, 597	Granosus (Margaron) Lea.....	879
Glebulus (Unio) Say.....	598	Granosus (Unio) Schumacher.....	691
Glencgensis (Diplodon) Dennent.....	889	Grantianus (Unio) Bourguignat.....	862
Glencgensis (Unio) Dennent.....	889	Granulatus (Margaron) Lea.....	565
Globata (Quadrula) Lea.....	793	Granulatus (Truncella) Rafinesque.....	518
Globatus (Unio) Lea.....	793	Granulatus (Unio) Lea.....	565
Globosa (Anodonta) Lea.....	646	Granuliferus (Diplodon) Dunker.....	879
Globosa (Margarita) Lea.....	530, 646	Granuliferus (Unio) Dunker.....	879
Globosa (Margaron) Lea.....	646	Granulosa (Grandidieria) Bourguignat.....	829
Globosa (Symphynota) Lea.....	530	Gratiosa (Nodularia) Philippi.....	815
Globosula (Anodonta) Heude.....	640	Gratiosus (Margaron) Lea.....	815
Globosus (Unio) Hanley.....	530	Gratiosus (Unio) Pætel.....	826
Globulus (Unio) Pætel.....	598	Gratiosus (Unio) Philippi.....	815
Goascoranensis (Margaron) Lea.....	594	Grateloupiana (Anodonta) Gassies.....	624
Goascoranensis (Nephronaia) Lea.....	594	Grateloupiana (Pseudanodonta) Bourguignat.....	624
Goascoranensis (Unio) Lea.....	594	Gratus (Diplodon) Lea.....	886
Gonidea (Conrad).....	506, 514, 657	Gratus (Margaron) Lea.....	886
Gontieri (Unio) Bourguignat.....	696	Gratus (Unio) Lea.....	886
Gotteschei (Unio) von Martens.....	862	Gravida (Grandidieria) Bourguignat.....	829
Gouldianus (Unio) Ward.....	789	Gravidus (Lampsilis) Rochebrune.....	830
Gouldii (Lampsilis) Lea.....	569	Gravidus (Margaron) Lea.....	830
Gouldii (Margaron) Lea.....	569	Gravidus (Physunio) Lea.....	830
Gouldii (Unio) Lea.....	569	Gravidus (Unio) Lea.....	830
Gowhattenensis (Parreysia) Theobald.....	845	Gravis (Mya) Wood.....	539
Gowhattenensis (Unio) Theobald.....	845	Gravis (Unio) Pætel.....	539
Gracilentus (Margaron) Lea.....	731	Grayana (Nodularia) Lea.....	806
Gracilii (Unio) Lea.....	731	Grayanus (Margarita) Lea.....	807
Gracillor (Lampsilis) Lea.....	559	Grayanus (Margaron) Lea.....	807
Gracillor (Margaron) Lea.....	559	Grayanus (Nodularia).....	809
Gracillor (Unio) Lea.....	559	Grayanus (Unio) Lea.....	806
Gracilis (Anodon) Sowerby.....	573, 588	Grayanus (Unio) Schrenck.....	807
Gracilis (Anodonta) Lea.....	833	Grayii (Unio) Griffith.....	807
Gracilis (Dallisia) Lea.....	833	Greecianus (Diplodon) von Ihering.....	875
Gracilis (Lampsilis) Barnes.....	573	Greecianus (Unio) von Ihering.....	875
Gracilis (Margarita) Lea.....	573	Grelloisianus (Unio) Bourguignat.....	689
Gracilis (Margaron) Lea.....	573, 833	Greeni (Margarita) Lea.....	614
Gracilis (Metaptera) Stimpson.....	573	Greeni (Margaron) Lea.....	614
Gracilis (Mya) Eaton.....	573	Greeni (Ptychobranchus) Conrad.....	614
Gracilis (Symphynota) Lea.....	573	Greeni (Unio) Conrad.....	614
Gracilis (Unio) Barnes.....	573, 881	Griffithianus (Margarita) Lea.....	722
Gracilsianus (Unio) Bourguignat.....	686	Griffithianus (Margaron) Lea.....	722
Granadensis (Anodonta) Lea.....	922	Griffithianus (Unio) Lea.....	722
Granadensis (Glabaris) Lea.....	922	Grijalve (Anodonta) Morelet.....	926
Granadensis (Margaron) Lea.....	744, 922	Grijalve (Glabaris) Morelet.....	926
Granadensis (Unio) Conrad.....	598	Grijolœ (Anodonta) Pætel.....	926
Granadensis (Unio) Lea.....	744	Grœnlandicus (Unio) Mörch.....	854
Grandidens (Margaron) Lea.....	799	Grunciana (Anodonta) Clessin.....	585
Grandidens (Unio) Lea.....	799	Gnahybœ (Unio) von Ihering.....	894
Grandidieri (Unio) Bourguignat.....	846	Guaraniana (Unio) d'Orbigny.....	884
Grandidieria Bourguignat.....	506, 515, 827	Guarayana (Margarita) Lea.....	911
Grandidieriana (Burtonia) Bourguignat.....	578	Guarayana (Margaron) Lea.....	911
Grandis (Anodonta) Say.....	633, 641, 647, 648	Guarayana (Monocondylea) d'Orbigny.....	910, 911
Grandis benedictensis (Anodonta) Lea.....	644	Guarayana (Unio) Hanley.....	911
Grandis footiana (Anodonta) Lea.....	642	Guatemalensis (Quadrula) Simpson.....	796
Grandis gigantea (Anodonta) Lea.....	643	Gubernaculum (Unio) Reeve.....	523
Grandis leonensis (Anodonta) Lea.....	643	Guillaini (Anodonta) Recluz.....	857, 858
Grandis (Margarita) Lea.....	642	Guillaini (Lamellidens) Recluz.....	833
Grandis (Margaron) Lea.....	642	Guillemeti (Cameronia) Bourguignat.....	910
Granifera pusilla (Quadrula) Simpson.....	795	Guillemeti (Unio) Bourguignat.....	862
Granifera (Quadrula) Lea.....	795	Gundlachi (Nephronaia) Dunker.....	595
Graniferus (Margarita) Lea.....	795	Gundlachi (Unio) Dunker.....	595
Graniferus (Margaron) Lea.....	795	Gundlachi (Unio) Sowerby.....	595
Graniferus (Unio) Lea.....	795	Guppyi (Ctenodesma) Smith.....	853
Granosa (Unio) Brugulere.....	878, 879	Habetatus (Margaron) Lea.....	787
Granosus (Diplodon) Bruguiere.....	878	Haconensis (Anodonta) von Ihering.....	640
Granosus (Margarita) Lea.....	879		

	Page.		Page.
Haeonensis (Nodularia) von Ihering.....	810	Hereulea (Anodonta) Middendorff.....	584
Haeonensis (Unio) von Ihering.....	810	Hereulea (Barbala) Patet.....	584
Hagleri (Pleurobema) Frierson.....	757	Hereulea (Cristaria) Middendorff.....	584
Hagleri (Unio) Frierson.....	757	Hereuleus (Anodon) Sowerby.....	584
Hagnei (Unio) Strubell.....	862	Hermosus (Unio) Bourguignat.....	848
Hainesiana (Chamberlainia) Lea.....	582	Hierodes (Unio) Kobelt.....	690
Hainesianus (Margaron) Lea.....	582	Heros multiplicatus (Unio) Patet.....	770
Hainesianus (Unio) Lea.....	582	Heros (Quadrula) Say.....	770
Haleianus (Lampsilis) Lea.....	563, 564	Heros (Unio) Kuster.....	767
Haleianus (Margaron) Lea.....	564	Heros (Unio) Say.....	770, 771
Haleianus (Unio) Lea.....	564	Hertwigii (Anodonta) von Ihering.....	925
Hallenbeckii (Anodonta) Lea.....	634	Hertwigii (Glabaris) von Ihering.....	925
Hallenbeckii (Margaron) Lea.....	634, 726	Heterodon (Alasmidonta) Lea.....	668, 669
Hallenbeckii (Unio) Lea.....	726	Heterodon (Margarita) Lea.....	669
Hamyanus (Unio) Bourguignat.....	862	Heterodon (Margaron) Lea.....	669
Hanleyana (Castalia) Sowerby.....	864	Heterodon (Unio) Lea.....	669
Hanleyana (Pleurobema) Lea.....	759	Heterogene.....	514, 516
Hanleyanus (Unio) Lea.....	759	Heudei (Cuneopsis) Heude.....	805
Harlandi (Anodonta) Baird and Adams.....	640	Heudei (Unio) Heude.....	805
Harmandi (Monocendylaea) Patet.....	836	Hians (Alasmodon) Rafinesque.....	673
Harmandi (Pseudodon) Crosse and Fischer.....	836	Hians (Alasmodonta) Ferussac.....	665
Harmandia Rochebrune.....	506, 515, 827	Hians (Anodon) Sowerby.....	915
Harperi (Pleurobema) B. H. Wright.....	759	Hians (Unio) Valenciennes.....	667
Harperi (Unio) B. H. Wright.....	759	Higginsii (Lampsilis) Lea.....	540
Harpethensis (Anodonta) Lea.....	644	Higginsii (Margaron) Lea.....	540
Harpethensis (Margaron) Lea.....	644	Higginsii (Unio) Lea.....	540
Hartmaniana (Margaron) Lea.....	756	Hildrethi (Unio) Delessert.....	655
Hartmaniana (Pleurobema) Lea.....	756	Hildrethiana (Baphia) H. and A. Adams.....	673
Hartmanianus (Unio) Lea.....	756	Hildrethiana (Margaritana) B. H. Wright.....	673
Hartmanni (Spatha) von Martens.....	898	Hildrethiana (Strophitus) Conrad.....	673
Hartwrighti (Unio) B. H. Wright.....	710, 711	Hildrethianus (Margarita) Lea.....	673
Hastatus (Unio) Lea.....	727	Hildrethianus (Margaron) Lea.....	673
Hautteeœuri edwardsiana (Parreysia) Bourguignat.....	846	Hildrethianus (Unio) Lea.....	673
Hautteeœuri (Parreysia) Bourguignat.....	846	Hinkleyi (Unio) B. H. Wright.....	710
Hautteeœuri (Unio) Bourguignat.....	846	Hippopæus (Margaron) Lea.....	767
Haysiana (Scalenaria) Agassiz.....	520	Hippopæus (Unio) Lea.....	767
Haysiana (Truncella) Lea.....	520	Hippopæus (Unio) Patet.....	767
Haysianus (Margarita) Lea.....	520	Hirundo (Chelidoneura) von Martens.....	906
Haysianus (Margaron) Lea.....	520	Hirundo (Chelidonopsis) von Martens.....	906
Haysianus (Unio) Lea.....	520	Hirundo (Spatha) von Martens.....	906
Hazelhurstianus (Margaron) Lea.....	719	Hispanus (Unio) Rossmasser.....	686
Hazelhurstianus (Unio) Lea.....	719	Hochstetieri (Unio) Dunker.....	889
Hebes (Margaron) Lea.....	742	Hoekingensis (Anodonta) Call.....	648
Hebes (Unio) Lea.....	742	Holstonensis (Margaron) Lea.....	747
Hebetata (Quadrula) Conrad.....	787	Holstonensis (Pleurobema) Lea.....	746
Hebetatus (Unio) Conrad.....	787	Holstonia (Alasmidonta) Lea.....	670
Heermannii (Margaron) Lea.....	571	Holstonia (Baphia) H. and A. Adams.....	670
Heermannii (Unio) Lea.....	571	Holstonia (Margarita) Lea.....	670
Heldi (Anodonta) Kuster.....	833	Holstonia (Margaritana) Lea.....	670
Heldi (Unio) Kuster.....	699	Holstonia (Margaron) Lea.....	670
Helix.....	503	Holstoniana (Margaritana) Kuster.....	670
Helvetica (Anodonta) Bourguignat.....	625	Holstonianus (Unio) Hanley.....	670
Hembeli (Margaritana) Conrad.....	679	Holtonis (Anodonta) Lea.....	921
Hembeli (Margaron) Lea.....	679	Holtonis (Glabaris) Lea.....	921
Hembeli (Unio) Conrad.....	679	Holtonis (Margaron) Lea.....	921
Hemidon Swainson.....	667	Homogene.....	514, 620, 658
Hemilastena Agassiz.....	506, 514, 673	Homsensis (Margaron) Lea.....	693
Hemiodon Swainson.....	667	Homsensis (Unio) Lea.....	693
Hemiodonta Swainson.....	667	Hopetonensis (Margarita) Lea.....	729
Henryana (Anodonta) Lea.....	635	Hopetonensis (Margaron) Lea.....	729
Henryana (Margaron) Lea.....	635	Hopetonensis (Unio) Lea.....	728
Hepaticus (Margaron) Lea.....	717	Horda (Anodon) Gould.....	635
Hepaticus (Unio) Lea.....	717	Hordeum (Anodon) Sowerby.....	633
		Horei (Grandidieria) Bourguignat.....	823
		Horei (Nodularia) E. A. Smith.....	823

	Page.		Page.
Houghtonensis (Anodonta) Currier.....	643	Inerassatus (Mytilus) Sheppard.....	623
Hongddaranienus (Unio) Tapperone-Can-		Inerassatus (Unio) Lea.....	707
efri.....	844	Ineurvus (Unio) Lea.....	692
Houstonia (Anodonta) Linsley.....	634	Indecora (Anodon) Heude.....	639
Housel (Margaron) Lea.....	580	Indefinitus (Margaron) Lea.....	724
Housei (Unio) Lea.....	580	Indefinitus (Unio) Lea.....	723
Houstonensis (Margaron) Lea.....	782	Indicus (Margaron) Lea.....	844
Houstonensis (Quadrula) Lea.....	782	Indicus (Unio) Sowerby.....	844
Houstonensis (Unio) Lea.....	782	Ineptus (Margaron) Lea.....	742
Hueti (Unio) Bourguignat.....	687	Ineptus (Unio) Lea.....	742
Hueti (Unio) Kobelt.....	694	Inerme (Aplodon) Spix.....	911
Humerosus (Margaron) Lea.....	724	Inerme (Monocondylaea) Spix.....	911
Humerosus (Unio) Lea.....	724	Inflata (Anodonta) Rafinesque.....	648
Humilis (Margaron) Lea.....	811	Inflata (Castalia) d'Orbigny.....	863
Humilis (Unio) Lea.....	811	Inflata (Metaptera) Conrad.....	574
Hupeana (Mycetopoda) Clessin.....	935	Inflata (Mya) Eaton.....	534
Hupeanus (Mycetopus) Clessin.....	935	Inflata (Symphynta) Lea.....	574
Hyalinus (Margaron) Lea.....	663	Inflatus (Margarita) Lea.....	574
Hyalinus (Unio) Lea.....	663	Inflatus (Margaron) Lea.....	574
Hydianus (Lampsilis) Lea.....	536	Inflatus (Unio) Barnes.....	534
Hydianus (Margarita) Lea.....	536	Inflatus (Unio) Conrad.....	568
Hydianus (Margaron) Lea.....	536	Inflatus (Unio) Hanley.....	574
Hydianus (Unio) Kuster.....	534	Infrarostatus (Unio) Schepman.....	581
Hydianus (Unio) Lea.....	536	Infucata (Quadrula) Lea.....	783, 784
Hygapanus (Nodularia) Böttger.....	827	Infucatus (Margarita) Lea.....	784
Hygapanus (Unio) Böttger.....	827	Infucatus (Margaron) Lea.....	784
Hylea (Unio) d'Orbigny.....	884	Infucatus (Unio) Conrad.....	784
Hyleus (Diplodon) d'Orbigny.....	884	Infulgens (Unio) Lea.....	724
Hyleus (Margaron) Lea.....	884	Infusus (Unio) Lea.....	724
Hyria Lamarck.....	506, 515, 868	Ingallsiana (Nodularia) Lea.....	816
Hyriana Simpson.....	572	Ingallsianus (Margaron) Lea.....	816
Hyrianae.....	501, 515, 806	Ingallsianus (Unio) Lea.....	816
Hyridella Swainson.....	888	Inornata (Anodonta) Anthony.....	644
Hypsiprinnus (Parreysia) von Martens.....	849	Inornata (Anodonta) Kuster.....	625
Hypsiprinnus (Unio) von Martens.....	849	Inornata (Nodularia) Lea.....	813
Hyriopsis Conrad.....	505, 506, 511, 514, 578	Inornatus (Margaron) Lea.....	813
Ianthinus (Unio) Philippi.....	881	Inornatus (Unio) Lea.....	813
Icterinus (Unio) Conrad.....	727	Inornatus (Unio) Reeve.....	814
Idgi (Unio) Bourguignat.....	862	Inoscularis (Anodon) Gould.....	837
Idrina (Anodonta) Spinelli.....	624	Inoscularis (Margarita) Pætel.....	837
Iheringella Pilsbry.....	507, 515, 913	Inoscularis (Margaron) Lea.....	837
Iheringi (Anodonta) Clessin.....	919	Inoscularis (Pseudodon) Gould.....	837
Iheringi (Glabaris) Clessin.....	919	Insignis (Grandidieria) Bourguignat.....	828
Iheringi (Unio) Clessin.....	894	Insolidus (Unio) Lea.....	664
Iheringi (Unio) B. H. Wright.....	712, 895	Instructa (Pleurobema) Lea.....	756
Iliqi (Unio) Soleillet.....	862	Instructus (Margaron) Lea.....	756
Imbecillis (Anodonta) Say.....	634, 635	Instructus (Unio) Lea.....	756
Imbecillis (Margaron) Lea.....	635	Insularis (Dallia) Drouet.....	834
Imbricata (Anodonta) Anthony.....	643	Insularis (Pseudodon) Drouet.....	834
Imbricatus (Unio) Moreh.....	607	Insulsus (Margaron) Lea.....	716
Imperialis (Unio) Morelet.....	582	Insulsus (Unio) Lea.....	716
Implicata (Anodonta) Say.....	633	Intercedens (Margaron) Lea.....	548
Implicata (Margaron) Lea.....	633	Intercedens (Unio) Lea.....	548
Impressa (Alasmodonta) Anthony.....	670	Intermedia (Anodonta) Lamarck.....	623
Impura (Anodonta) Say.....	635	Intermedia (Quadrula) Conrad.....	775
Inæqualis (Unio) Rochebrune.....	862	Intermedius (Margaron) Conrad.....	775
Inæqualis (Anodonta) Lea.....	922	Intermedius (Unio) Conrad.....	775
Inæqualis (Glabaris) Lea.....	922	Intermerata (Anodon) Heude.....	639
Inæqualis (Margaron) Lea.....	922	Interrupta (Truncella) Agassiz.....	518
Inearnata (Grandidieria) Bourguignat.....	830	Interruptus (Margarita) Lea.....	541
Inearum (Anodonta) Philippi.....	921	Interruptus (Margaron) Lea.....	541
Inearum (Glabaris) Philippi.....	921	Interruptus (Unio) Conrad.....	518
Incerta (Anodonta) Lea.....	635	Interruptus (Unio) Lea.....	541
Incerta (Margarita) Lea.....	635	Interruptus (Unio) Say.....	772
Incertus (Ptychorhynchus) Simpson.....	851	Intervetus (Pleurobema) Lea.....	753
Incrassatus (Margaron) Lea.....	707	Intervetus (Unio) Lea.....	753

	Pagc.		Page.
Inusitatis (Unio) Lea.....	729	Jewittii (Anodonta) Lea	927
Inusitatus (Margaron) Lea.....	729	Jewittii (Margaron) Lea.....	742
Inusitatus (Unio) B. H. Wright.....	729	Jewittii (Unio) Lea.....	742
Invenustus (Unio) Lea.....	725	Jickeli (Nodularia) Simpson.....	826
Involuta (Nodularia) Benson.....	817	Juliani (Nodularia) Rang.....	824
Involutus (Margaron) Lea.....	817	Juliani (Unio) Rang.....	824
Involutus (Unio) Benson.....	817	Johannis (Margaron) Lea.....	759
Iridesceus (Unio) Conrad.....	744	Johannis (Pleurobema) Lea.....	759
Iridina Lamarck.....	902	Johannis (Unio) Lea.....	759
Iridinae.....	501	Johnstoni (Pseudavicula) Smith.....	861
Iridinea (Solenia) Heude.....	657	Johnstoni (Unio) Smith.....	860, 861
Iridineus (Mycetopus) Heude.....	657	Jolyi (Unio) Kobelt.....	692
Irisans (Anodon) Anthony.....	633	Jonesii (Margaron) Lea.....	554
Iris (Lampsilis) Lea.....	551, 552, 558	Jonesii (Unio) Lea.....	554
Iris (Margarita) Lea.....	552	Jordanicus (Margaron) Lea.....	689
Iris (Margaron) Lea.....	552	Jordanicus (Unio) Bourguignat.....	689
Iris (Unio) Lea.....	552	Joretii (Anodon) Heude.....	638
Irrasa (Pleurobema) Lea.....	756, 757	Josseti (Cameronia) Bourguignat.....	909
Irrasus (Margaron) Lea.....	756	Josseti (Unio) Bourguignat.....	862
Irrasus (Unio) Lea.....	756	Jouberti (Burtonia) Bourguignat.....	578
Irregularis (Anodon) Heude.....	638	Jouberti (Cameronia) Bourguignat.....	910
Irrorata (Cyprogenia) Lea.....	610	Jouberti (Moncettia) Bourguignat.....	901
Irrorata Pusilla (Cyprogenia) Simpson.....	610	Jouberti (Mutela) Bourguignat.....	904
Irrorata (Theliderma) Swainson.....	610	Jouberti (Unio) Bourguignat.....	862
Irrorata (Unio) Deshayes.....	610	Jourdyi (Anodonta) Heude.....	639
Irroratus (Margarita) Lea.....	610	Jourdyi (Brazzaea) Bourguignat.....	907
Irroratus (Margaron) Lea.....	610	Jourdyi (Nodularia) L. Morlet.....	816
Irroratus (Unio) Lea.....	504, 609, 775	Jourdyi (Unio) L. Morlet.....	816
Irwiniensis (Unio) Lea.....	724	Kallethensis (Unio) Pæstel.....	695
Isocardioides (Iheringella) Lea.....	913	Katherinae (Unio) Lea.....	666
Isocardioides (Margaron) Lea.....	913	Keinerianus (Margaron) Lea.....	780
Isocardioides (Plagiodon) Lea.....	913	Keinerianus (Unio) Lea.....	780
Isocardioides (Unio) Sowerby.....	913	Kelletti (Anodon) Sowerby.....	588
Jacobeus (Unio) Philipp.....	881	Kleiniana (Quadrula) Lea.....	784
Jacquemini (Unio) Dupuy.....	699	Kennerlyi (Anodonta) Lea.....	628
Jaculus (Unio) Rochebrune.....	862	Kennerlyi (Margaron) Lea.....	628
Jamesianus (Margaron) Lea.....	740	Kennicottii (Anodonta) Lea.....	647
Jamesianus (Unio) Lea.....	740	Kennicottii (Margaron) Lea.....	647
Japanensis jokohamensis (Nodularia)		Keraudreni (Margaron) Lea.....	812
von Ihering.....	810	Keraudreni (Nodularia) Eydoux.....	812
Japanensis (Margaron) Lea.....	809	Keraudreni (Unio) Eydoux.....	812
Japanensis (Nodularia) Lea.....	806, 809	Kingi (Medionidus) B. H. Wright.....	590
Japanensis (Unio) Lea.....	809	Kingi (Unio) B. H. Wright.....	590
Japanensis (Unio) Kobelt.....	810	Kirki liederi (Spatha) von Mariens.....	897
Japonica (Anodonta) von Martens.....	640	Kirki (Spatha) Ancey.....	897
Jaspideus (Margaron) Lea.....	912	Kirki (Spathella) Ancey.....	897
Jaspidea (Monocondylea) Hupé.....	912	Kirkii (Margaron) Lea.....	848
Jaspidea (Unio) Hupé.....	912	Kirkii (Parreysia) Lea.....	848
Jauberti (Unio) Locard.....	690	Kirkii (Unio) Lea.....	848
Javana (Anodon) Sowerby.....	588	Kirtlandiana minor (Quadrula) Simpson.....	791
Javanus (Margaron) Lea.....	818	Kirtlandiana (Quadrula) Lea.....	791
Javanus (Unio) Lea.....	817	Kirklandianus (Lampsilis) B. H. Wright.....	557
Jayanus (Margaron) Lea.....	714	Kirklandianus (Margarita) Lea.....	791
Jayanus (Unio) B. H. Wright.....	714	Kirklandianus (Margaron) Lea.....	791
Jayensis (Margarita) Lea.....	714	Kirklandianus (Unio) B. H. Wright.....	557
Jayensis (Margaron) Lea.....	714	Kirklandianus (Unio) Hanley.....	791
Jayensis (Unio) Lea.....	713	Kirtlandianus (Unio) Lea.....	791
Jeffreyianus (Diplodon) Lea.....	891	Kisonis (Unio) Kobelt.....	688
Jeffreyianus (Unio) Lea.....	891	Kleiniana (Quadrula) Lea.....	784
Jejunus (Margarita) Lea.....	725	Kleinianus (Margaron) Lea.....	784
Jejunus (Margaron) Lea.....	725	Kleinianus (Unio) Knuster.....	775
Jejunus (Unio) Lea.....	725	Kleinianus (Unio) Lea.....	784
Jenkinsianus (Lamellidens) Benson.....	857	Kobelti (Anodonta) von Ihering.....	640
Jenkinsianus (Unio) Benson.....	857	Kobelti (Unio) Rolle.....	690
Jewittiana (Anodonta) Lea.....	927	Koseretzii (Unio) Clessin.....	894
Jewittianus (Glabaris) Lea.....	927	Kouangensis (Quadrula) Simpson.....	799

	Page.		Page.
Kraussii (Margaron) Lea.....	867	Lardellianus (Unio) Pechchloli.....	684
Kraussii (Unio) Lea.....	867	Largillierti (Lampsilis) Philippi.....	577
Ksibianus (Unio) Mousson.....	692	Largillierti (Unio) Philippi.....	577, 744
Kullethensis (Margaron) Lea.....	695	Lasmabrachys (Unio) Rafinesque.....	744
Kullethensis (Unio) Lea.....	695	Lasmignona Rafinesque.....	664, 673
Kullinthenensis (Unio) Pætel.....	695	Lastena Rafinesque.....	506, 514, 654, 933
Kunenensis (Nodularia) Mousson.....	824	Lata (Anodonta) Rafinesque.....	654
Kunenensis (Unio) Mousson.....	824	Lata (Lastena) Rafinesque.....	654
Labacensis (Unio) Rossmassler.....	697	Lata (Leptodea) Conrad.....	654
Lachrymosa lunulata (Quadrula) Pratt.....	777	Lateralis (Obliquaria) Rafinesque.....	744
Lachrymosa (Quadrula) Lea.....	776, 778	Latialata (Hyria) Sowerby.....	589
Lachrymosus (Margarita) Lea.....	776	Laticostatus (Margaron) Lea.....	769
Lachrymosus (Margaron) Lea.....	776	Laticostatus (Margaron) Lea.....	769
Lachrymosus (Unio) Lea.....	775	Laticostatus (Unio) H. and A. Adams.....	769
Lacteolus (Unio) Lea.....	873	Laticostatus (Unio) Lea.....	769
Lacustris (Anodonta) Lea.....	632	Lati-marginata (Anodon) Catlow and Reeve.....	917
Lacustris (Margaron) Lea.....	632	Latiquadrata (Castalia) Sowerby.....	864
Lacustris (Spatha) Simpson.....	898	Latiradiatus (Unio) Conrad.....	541
Lævigata (Unio) Say.....	600	Latissima (Unio) Rafinesque.....	545
Lævirostris Simpson.....	894	Latissimus (Unio) Sowerby.....	527
Lævirostris (Unio) Benson.....	842	Lato-marginata (Anodonta) Lea.....	917
Lævisimus (Lampsilis) Lea.....	574	Lato-marginata (Margarita) Lea.....	917
Lævisimus (Margarita) Lea.....	574	Lato-marginata (Margaron) Lea.....	917
Lævisimus (Margaron) Lea.....	574	Lato-marginatus felix (Glabaris) Pilsbry.....	917
Lævisimus (Symphynta) Lea.....	574	Latus (Margaron) Lea.....	728
Lævisimus (Unio) Conrad.....	574	Latus (Unio) Conrad.....	654
Lamareckianus (Margaron) Lea.....	610	Latus (Unio) Lea.....	728
Lamareckianus (Unio) Lea.....	610	Lauta (Anodonta) von Martens.....	639
Lamellatus (Lamellidens) Lea.....	856	Lavigerina (Brazzaea) Bourguignat.....	907
Lamellatus (Margarita) Lea.....	856	Lavigerina (Burtonia) Bourguignat.....	577
Lamellatus (Margaron) Lea.....	856	Lavigerina (Cameronia) Bourguignat.....	910
Lamellatus (Unio) Hanley and Theobald.....	581	Lavigerina (Mutela) Bourguignat.....	906
Lamellatus (Unio) Lea.....	856	Lavigerina (Spatha) Bourguignat.....	901
Lamellatus (Unio) Sowerby.....	857	Lavigerinus (Unio) Bourguignat.....	862
Lamellidens Simpson.....	506, 508, 515, 854, 858	Lawii (Unio) Lea.....	746, 747
Laminata (Anodonta) Rochebrune.....	648	Lawleyanus (Unio) Gentiluomo.....	683
Lamphorhamphus.....	815, 862	Layardii (Unio) Lea.....	856
Lampreyanus (Margaron) Lea.....	804	Lazarus (Unio) Lea.....	732
Lampreyanus (Shitodesmus) Baird and Adams.....	804	Lazarus (Unio) Sowerby.....	704
Lampreyanus (Unio) Baird and Adams.....	803, 804	Leal cinnamomicus (Unio) Heude.....	802
Lamproscapha.....	501	Leal (Dipodoni) Simpson.....	876, 881
Lamprotula Simpson.....	796	Leal leleci (Quadrula) Heude.....	801
Lampsilis Rafinesque.....	505,	Leal leleci (Unio) Pætel.....	801
	506, 511, 514, 526, 578, 589, 592, 828	Leal (Margarita) Lea.....	801
Lanceolaria Conrad.....	806, 811	Leal (Margaron) Lea.....	801
Lanceolata (Arconala) Lea.....	860	Leal (Quadrula) Gray.....	801, 802
Lanceolata (Hyria) Lea.....	860	Leal (Unio) Gray.....	801
Lanceolata (Triquetra) Lea.....	859, 860	Leali (Iridina) Sowerby.....	908
Lanceolatus (Margarita) Lea.....	734	Leali (Platiris) Lea.....	908
Lanceolatus (Margaron) Lea.....	734	Leali (Pleiodoni) H. and A. Adams.....	908
Lanceolatus (Unio) Lea.....	734	Lechaptosis (Nodularia) Ancey.....	823
Landanensis (Unio) Schepman.....	823	Lechaptosis (Unio) Ancey.....	823
Landbecki (Unio) Philippi.....	881	Lecontianus (Margarita) Lea.....	733
Landeaul (Cameronia) Bourguignat.....	909	Lecontianus (Margaron) Lea.....	733
Landeaul (Pleiodoni) Bourguignat.....	909	Lecontianus (Unio) Lea.....	537, 733
Langulati (Lepidodesma) Heude.....	587	Ledouxliana (Nodularia) Charnes.....	812
Langulati (Unio) Heude.....	586, 587	Ledouxlianus (Unio) Charnes.....	812
Langulatus aligerus (Unio) Heude.....	587	Leeai (Unio) Hanley.....	801
Laosensis (Anodonta) Fischer.....	629	Legrandi (Unio) Petterd.....	891
Laosensis (Margaritana) Lea.....	678	Legumen (Anodonta) von Martens.....	932
Laosensis (Margaron) Lea.....	678	Legumen (Glabaris) von Martens.....	932
Laosensis (Unio) Lea.....	678	Legumen (Mutellina) Rochebrune.....	906
Lapidus Simpson.....	700	Leguminata Conrad.....	506, 513, 514, 651
Lapidus (Unio) Kobelt.....	700	Lehmani (Unio) S. H. Wright.....	715
Lapillus (Unio) Say.....	525	Leibil (Margaron) Lea.....	601

	Page.		Page.
Leibii (Obovaria) Lea	600, 601	Liebmanni (Unio) Philippi	712
Leibii (Unio) Lea	601	Liederi (Nodularia) von Martens	824
Leila Gray	501, 507, 515, 914	Liederi (Unio) von Martens	824
Lefoma (Unio) Benson	811	Lienosus constrictus (Unio) Conrad	551
Leleci (Unio) Heude	801	Lienosus (Lampsilis) Conrad	547
Lemsleyi (Anodonta) Morelet	588	Lienosus (Margarita) Lea	547
Lemsleyi (Pilsbryconcha) Morelet	588	Lienosus (Margaron) Lea	547
Le Mutel Adanson	903	Lienosus unicastatus (Lampsilis) B. H. Wright	547
Lenior (Margaron) Lea	518	Lienosus (Unio) Conrad	547
Lenior (Truncilla) Lea	518	Ligamentina (Unio) Lamarck	539, 877
Lenior (Unio) Lea	518	Ligamentinus gibbus (Lampsilis) Simpson	540
Lenis (Unio) Conrad	527	Ligamentinus (Lampsilis) Lamarck	539
Lenis (Unio) Lea	518	Ligamentinus (Margaron) Lea	539
Lens depygis (Obovaria) Conrad	601	Ligatus (Unio) Lea	724
Lens (Margarita) Lea	600	Ligula (Unio) Mousson	862
Lens Margaron Lea	600	Lima (Nodularia) Simpson	820
Lens (Obovaria) Lea	600	Limatulus (Margaron) Lea	716
Lens Simpson	831	Limatulus (Unio) Conrad	716
Leis (Unio) Lea	600, 601	Limensis (Unio) Kuster	881
Lenticularis (Anodonta) Lea	922	Limnæa Poli	620, 679
Lenticularis (Margaron) Lea	922	Limnoica (Anodonta) d'Orbigny	921
Lenticularis (Pleurobema) Lea	761	Limnoica (Glabaris) d'Orbigny	921
Lenticularis (Unio) Lea	761	Limnoica (Margarita) Lea	921
Lentiformis (Aplodon) von Ihering	913	Limnoica (Margaron) Lea	921
Lentiformis (Margaron) Lea	913	Limosa (Anodon) Heude	638
Lentiformis (Monocondylæa) Lea	912	Limosus (Unio) Kobelt	683
Leonensis (Anodonta) Lea	643	Limosus (Unio) Nilsson	682
Leonensis (Margaron) Lea	643	Lincecumii (Margaron) Lea	768
Leopoldvillensis (Burtonia) Putzeys	577	Lincecumii (Unio) Lea	768
Leopoldvillensis (Parreysia) Putzeys	845, 846	Lindsleyi (Margaron) Lea	542
Leopoldvillensis (Pseudospatha) Putzeys	577	Lindsleyi (Unio) Lea	542
Leopoldvillensis (Unio) Putzeys	846	Lineata (Anodon) Heude	638
Leotandi (Anodonta) Guppy	929	Lineata (Unio) Valenciennes	536
Leotandi (Glabaris) Guppy	929	Lineatus (Margaron) Lea	528
Leotandi (Margaron) Lea	929	Lincecurii (Unio) Patel	768
Lepidior (Margaron) Lea	880	Lineatus (Unio) Lea	528
Lepidodesma Simpson	506, 514, 586	Lineolata (Obliquaria) Rafinesque	744
Lepidus (Margaron) Lea	561	Lineolata (Plagiola) Agassiz	604
Lepidus (Unio) Gould	561	Lineolata (Unio) Say	604
Lepidus (Unio) Lea	880	Lineolatus (Unio) Conrad	604
Lepsi (Spatha) Jickeli	896	Linguaeformis (Anodonta) Morelet	587
Lesleyi (Margaron) Lea	748	Linguaeformis (Margaron) Lea	554, 587
Lesleyi (Pleurobema) Lea	743	Linguaeformis (Pilsbryconcha) Morelet	587
Lesleyi (Unio) Lea	748	Linguaeformis (Unio) Lea	554
Lessoni (Diplodon) Kuster	890	Lingulata (Anodonta) Hupe	929
Lessoni (Unio) Kuster	890	Lingulata (Glabaris) Hupe	929
Lesueurianus (Margaron) Lea	792	Lingulata (Margaron) Lea	929
Lesueurianus (Unio) Lea	792	Lingulatus (Rectidens) Drouet and Chaper	854
Leptodea (Rafinesque) Conrad	654	Lingulatus (Unio) Drouet and Chaper	854
Leptodon (Lampsilis) Rafinesque	575	Linneana (Anodonta) Lea	646
Leptodon (Leptodea) Conrad	575	Linneana (Margaron) Lea	646
Leptodon (Symphynota) Ferussac	575	Lita (Pleurobema) Lea	761
Leptodon (Unio) Rafinesque	575	Lithophagus (Unio) Ziegler	812
Letourneauxiana (Pleiodon) Bourguignat	909	Litoralis (Margarita) Lea	691
Letourneuxi (Anodonta) Bourguignat	627, 649	Litoralis (Margaron) Lea	691
Letourneuxi (Colletopterum) Bourguignat	649	Litoralis (Unio) C. Pfeiffer	678
Letourneuxi (Unio) Bourguignat	686	Litoralis umbonatus (Unio) Rossmässler	691
Letourneuxi (Unio) Kobelt	692	Littoralis acarranicus (Unio) Kobelt	691
Letourneuxiana (Chambardia)	902	Littoralis minor (Unio) Rossmässler	691, 698
Levigata (Unio) Rafinesque	744	Littoralis pianensis (Unio) Kobelt	691
Lewisii (Anodonta) Lea	642	Littoralis (Unio) Lamarck	690, 691, 692, 693
Lewisii (Margaron) Lea	755	Litturata (Anodonta) Hupe	931
Lewisii (Unio) Lea	642, 755	Lituratus (Glabaris) Spix	931
Liebmanni (Margaron) Lea	712	Litus (Unio) Lea	761

	Page.		Page.
Livida (Anodon) Heude.....	638	Lutulentus (Unio) Gould.....	892
Lividus (Lampsilis) Simpson.....	571	Luxata (Anodonta) Kuster.....	625
Livingstonensis (Burtonia) Bourguignat ..	578	Luxurians (Unio) Kuster	699
Livingstonensis (Margaron) Lea	729	Luynes (Unio) Locard.....	693
Livingstonensis (Pseudospatha) Bour-		Lyceus (Unio) Rolle.....	692
guignat.....	573	Lyonii (Margaron) Lea.....	751
Livingstonensis (Unio) Lea	729	Lyonii (Pleurobema) Lea	751
Lobata (Unio) Philippi.....	686	Lyonii (Unio) Lea.....	751
Locardi (Gabbillota) Scrvain.....	650	Lymnium Oken	679, 650
Locardi (Leguminaria) Simpson	652, 653	Lymnoica (Anodonta) Chenu.....	921
Locardiana (Cameronia) Bourguignat ...	910	Mabiliei (Pseudodon) Rochebrune.....	840
Locardiana (Chambardia)	902	Mabiliei (Spatha) Jousseau.....	897
Locardiana (Grandidieria) Bourguignat ..	830	Mabilii (Mutellina) Rochebrune.....	906
Locellus (Margaron) Lea.....	876	Mabiliana (Cameronia) Bourguignat ...	910
Locellus (Diplodon) Lea.....	876	Maccarthianus (Unio) Kobelt.....	692
Locellus (Unio) Lea.....	876	Maccarthianus (Unio) Bourguignat	687
Longina (Iridina) Lea	929	Macer (Margaron) Lea	728
Longinus (Anodon) Spix.....	929	Macer (Unio) Lea	728
Longinus (Glabaris) Spix.....	929	Macilenta (Anodonta) Morelet.....	624
Longinus (Mycetopus) Clessin	929	Macilentus (Unio) Benson	813
Longirostris (Unio) Rossmässler	682	Macmurtrei (Margarita) Lea	908
Longitudinatus (Unio) Anton.....	862	Macmurtrei (Pleodon) Conrad.....	908
Longus (Unio) Philippi	882	Macnielli (Nephronaias) Lea.....	593, 594
Loomisi (Pseudodon) Simpson	840	Macnielli (Unio) Lea.....	593
Lorteti (Unio) Kobelt	690	Maconensis (Margaron) Lea.....	736
Lorteti (Unio) Locard	688	Maconensis (Unio) Lea.....	736
Lourdelli (Nodularia) Bourguignat	826	Macrodon (Margaron) Lea	605
Lourdelli (Unio) Bourguignat.....	826	Macrodon (Plagiola) Lea.....	605
Lucasi (Anodonta) Deshayes.....	627	Macrodon (Unio) Lea	605
Lucasii (Margaron) Lea.....	627	Macropterus (Unio) Dunker.....	894
Lucida (Anodonta) d'Orbigny.....	921	Macula (Mytilus) Sheppard.....	623
Lucida (Anodonta) Heude.....	640	Maculata (Pleurobema) Conrad.....	746
Lucida (Glabaris) d'Orbigny.....	921	Maculatus (Margarita) Lea	746
Lucida (Margaron) Lea.....	921	Maculatus (Margaron) Lea.....	746
Lucidus (Margaron) Lea	716	Maculatus (Unio) Conrad	746
Lucidus (Unio) Lea	716	Madagascariensis (Unio) Sganzin.....	862
Ludovicianum (Diplodon) Rochebrune..	862	Magnifica (Anodonta) Clessin	584
Lugens (Nodularia) Drouet and Chaper..	818	Magnifica (Anodonta) Lea.....	637
Lugens (Unio) Drouet and Chaper.....	818	Magnifica (Burtonia) Bourguignat.....	577
Lugubris (Anodonta) Say	643	Magnifica (Margarita) Lea.....	637
Lugubris (Margarita) Lea.....	715	Magnifica (Margaron) Lea.....	637
Lugubris (Margaron) Lea	715	Magnifica (Symphynota) Lea	637
Lugubris (Unio) Kuster	723	Magnificus (Unio) Sowerby	585
Lugubris (Unio) Lea	715	Mainwaringi (Unio) Nevill	856
Lugubris (Unio) Say	568	Maltzani (Unio) Kuster	683
Lunulatus (Unio) Pratt	777	Manca (Unio) Lamarck	682
Lunulifer (Margaron) Lea	689	Manca (Unio) Millet	699
Lunulifer (Unio) Bourguignat	689	Mandarinus (Unio) Lea	815
Luridus (Margaron) Lea	705	Mandelayensis (Unio) Theobald	843
Luridus (Unio) Lea	705	Mandinguorum (Unio) Rochebrune	826
Lurulenta (Anodonta) Morelet.....	647	Manhoti (Pseudodon) Conrad.....	839
Lusitana (Anodonta) Morelet	623	Manubius (Margaron) Lea	740
Lutens (Unio) Lea	841	Manubius (Unio) Gould.....	740
Luteola (Anodonta) Lea	920	Manubius (Unio) Patel	740
Luteola (Margaron) Lea	920	Marcens (Unio) Hanley and Theobald... 843	
Luteola (Unio) Lamarck	534	Mardinensis chantrei (Leguminaria) Lo-	
Luteolus (Glabaris) Lea	920	card	651
Luteolus (Lampsilis) Lamarck	534	Mardinensis (Leguminaria) Lea	651, 652, 653
Luteolus (Margarita) Lea	534	Mardinensis (Margaritana) Clessin	651
Luteolus (Margaron) Lea	534	Mardinensis (Margaron) Lea.....	651, 695
Luteolus rosaceus (Lampsilis) De Kay...	535	Mardinensis (Monocondylaea) Lea.....	651
Luteolus (Unio) Sowerby	539	Mardinensis (Unio) Lea.....	695
Luteus (Margaron) Lea	842	Margaritana Schumacher	501, 506, 513, 514, 674
Luteus (Unio) Lea.....	841	Margaritifera minor (Unio) Rossmässler ..	675
Lutulentus (Diplodon) Gould.....	892	Margaritifera (Unio) Kuster.....	676
Lutulentus (Margaron) Lea.....	892	Margaritifera (Alasmodonta) Conrad....	675

	Page.		Page.
Margaritifera (Baphia) H. and A. Adams.	675	Mauritanus (Unio) Lea	833
Margaritifera (Baphia) Meuschen.	675	Menfelli (Anodonta) Anthony	644
Margaritifera (Margarita) Lea	675	Mearnsi (Lampsilis) Simpson	564
Margaritifera (Margaritana) Linnæus	513, 674, 677, 678, 741	Mearnsiana (Anodonta) Simpson	630
Margaritifera (Margaron) Lea	675	Mecklenbergensis (Margaron) Lea	723
Margaritifera (Mya) Linnæus	674, 675, 676	Mecklenbergensis (Unio) Lea	723
Margaritifera (Unio) Draparnaud.	675	Medellinus (Margarita) Lea	592
Margaron Lea	501	Medellinus (Margaron) Lea	592
Marginaliscandaharica (Unio) Hanley and Theobald.	855	Medellinus (Nephronaias) Lea	592
Marginalis candaharicus (Lamellidens) Hanley and Theobald.	855	Medellinus (Unio) Lea	592
Marginalis corriana (Unio) Hanley and Theobald.	855	Mederianus (Unio) Kuster.	818
Marginalis cylindricus (Lamellidens) Hanley and Theobald.	855	Mediastinus (Ptychorhynchus) Heude.	851
Marginalis cylindricus (Unio) Hanley and Theobald.	855	Mediastinus (Unio) Heude.	851
Marginalis hanleyi (Lamellidens) Simp- son	855	Mediocris (Margaron) Lea	723
Marginalis (Lamellidens) Lamarek.	854	Mediocris (Unio) Lea	723
Marginalis (Margarita) Lea	855	Medionidus Simpson.	506, 510, 514, 533
Marginalis (Margaron) Lea	855	Medius (Margaron) Lea	755
Marginalis obesa (Unio) Hanley and The- obald.	855	Medius (Unio) Lea	755
Marginalis obesus (Lamellidens) Hanley and Theobald.	855	Medius (Unio) Reeve	753
Marginalis tricolor (Lamellidens) Kuster.	855	Megadesma (Barbala) Pætel.	587
Marginalis (Unio) Lamarck.	854, 855	Megadesma (Cristaria) von Martens	587
Marginata (Alasmidonta) Say.	670, 671, 765	Megadomus Swainson	608, 662
Marginata (Alasmodonta) Say.	670	Megapterus (Unio) Morelet	579
Marginata (Anodon) Ferussac	617	Melanochoorea (Anodon) Heude.	635
Marginata (Anodonta) C. B. Adams	671	Melinia (Anodonta) Bourguignat.	625
Marginata (Anodonta) Say.	632	Melinus (Unio) Conrad.	536
Marginata (Baphia) H. and A. Adams.	671	Melleus (Margaron) Lea	597
Marginata (Margarita) Lea	671	Melleus (Nephronaias) Lea	597
Marginata (Margaritana) Gould	671	Melleus (Unio) Lea	597
Marginata (Margaron) Lea	671	Mellinus (Unio) Pætel	536
Marginata truncata (Margaritana) B. H. Wright	671	Membranacea (Anodonta) d'Orbigny	917
Marginata (Unio) Hanley	671	Membranacea (Unio) Hanley	888
Marginatus (Mytilus) Eaton	631	Membranaceus (Glabaris) Maton	913
Marginis (Margaron) Lea	566	Membranaceus (Margarita) Lea	919
Marginis (Unio) Lea	566	Membranaceus (Margaron) Lea	919
Marioniana (Cameronia) Bourguignat.	910	Membranaceus (Mytilus) Maton	918
Maris-galilei (Unio) Locard.	693	Membranaceus (Unio) Hanley	919
Marnoi (Spatha) Jickel	899	Menardi (Unio) Bourguignat	862
Marryattana (Anodonta) Lea	642	Meneliki (Unio) Solillet.	862
Marryattana (Margaron) Lea	642	Menkianus (Margarita) Lea	541
Marsili (Unio) B. H. Wright	714	Menkianus (Margaron) Lea	541
Martensi (Anodonta) Clessin	648	Menkianus (Unio) Lea	541
Martensi (Castalina) von Ihering	865	Menzieni (Margaron) Lea	889
Martensi (Diplodon) von Ihering	882	Menziezi depauperatus (Diplodon) Hut- ton	890
Martensi (Spatha) Sturany.	897	Menziezi (Diplodon) Gray	872, 839
Martensi (Unio) von Ihering.	882	Menziezi hochstellersi (Diplodon) Dunker	839
Masoni (Margarita) Lea	709	Menziezi rugatus (Diplodon) Hutton	839
Masoni (Margaron) Lea	709	Menziezi (Unio) Gray	889
Masoni (Unio) Conrad	709	Menzieziana (Unio) Reeve	889
Massini (Unio) Morelet	830	Merceri (Margaron) Lea	718
Matoniana (Mytilus) d'Orbigny.	888	Merceri (Unio) Lea	718
Matonianus (Unio) Pætel	888	Merdiger (Unio) Reeve	700
Mattioli (Unio) Tapperone Canefri	852	Meredithii (Margaron) Lea	761
Mattioli (Virgus) Tapperone Canefri.	852	Meredithii (Pleurobema) Lea	761
Mauritanicus (Unio) Bourguignat	692	Meredithii (Unio) Lea	761
Mauritanus (Margaron) Lea	833	Merodabensis (Unio) Kuster	842
		Merus (Margaron) Lea	708
		Merus (Unio) Lea	708
		Mesogenæ	514, 609
		Metallicus (Lampsilis) Say.	572
		Metallicus (Unio) Say.	572
		Metaneurus (Unio) Kuster	774
		Metanever (Margarita) Lea	774
		Metanever (Margaron) Lea	774
		Metanevra (Obliquaria) Rafinesque	774

	Page.		Page.
Metanevra (Quadrula) Rafinesque...	773, 774, 776	Modicellus (Margaron) Lea	518
Metanevra (Unio) Swainson	774	Modicellus (Unio) Lea	518
Metanevra wardii (Quadrula) Lea	774	Modicus (Margaron) Lea	762
Metanevrus (Unio) Rafinesque	503, 504	Modicus (Unio) Lea	762
Metaniver (Unio) Catlow and Reeve	774	Modioliformis (Lampsis) Lea	558, 559
Metaplatea (Truncilla) Rafinesque	518	Modioliformis (Margarita) Lea	559
Metaptera (Metaptera) Rafinesque	567	Modioliformis (Margaron) Lea	559
Metaptera Rafinesque	579	Modioliformis (Unio) Lea	559
Metastriata (Truncilla) Conrad	519	Mœstus (Lampsis) Lea	565
Metastriatus (Unio) Conrad	519	Mœstus (Margaron) Lea	565
Mexicanus (Margaron) Lea	701	Mœstus (Unio) Lea	565
Mexicanus (Unio) Philippi	701	Moineti (Brazzaea) Bourguignat	907
Mexicanus (Unio) Sowerby	705	Moineti (Burtonia) Bourguignat	577
Micans (Anodonta) Anthony	647	Moineti (Cameronia) Bourguignat	910
Micans (Margaron) Lea	647, 714	Moineti (Moncetia) Bourguignat	901
Micans (Unio) Lea	714	Moineti (Mutela) Bourguignat	906
Micellii (Unio) Kobelt	695	Moineti (Spatha) Bourguignat	901
Michaudiana (Unio) Des Moulins	682	Moineti (Unio) Bourguignat	862
Micheloti (Nodularia) L. Morlet	814	Molinæ (Diplodon) Philippi	881
Micheloti (Unio) L. Morlet	814	Molinæ (Unio) Philippi	881
Michonii (Leguminaia) Bourguignat	652	Molleuri (Parreysia) Morlet	849
Michonii (Margaritana) Pætel	652	Molleuri (Unio) Morlet	849
Michonii (Pseudodon) Conrad	652	Monceti (Parreysia) Bourguignat	848
Michonii (Unio) Bourguignat	652	Monceti (Unio) Bourguignat	848
Microcondylæa Vest	651, 653	Moncetia Bourguignat	901
Microcondylus Drouet	651	Mongolicus (Unio) Middendorff	677
Micromya Agassiz	506, 514, 524	Monocondylæa d'Orbigny	501, 507, 515, 910, 912
Micronaias Simpson	743	Monocondylus Morelet	834
Micropterus (Physunio) Morelet	831	Monodonta (Alasmiodonta) Ferussac	679
Micropterus (Unio) Morelet	831	Monodonta (Margaritana) Say	679
Microstictis (Quadrula) Heude	799, 803	Monodonta Say	666
Microstictis (Unio) Heude	799	Monodonta (Unio) Say	679
Milletii (Anodonta) Ray and Drouet	624	Monodontina Conrad	834
Mingorum (Anodon) Heude	638	Monodontus (Margarita) Lea	679
Minima (Anodonta) Millet	623	Monodontus (Margaron) Lea	679
Minor (Alasmiodonta) Lea	668	Monroensis (Margaron) Lea	710
Minor (Baphia) H. and A. Adams	668	Monroensis (Unio) Lea	710
Minor (Lampsis) Lea	562	Montanus (Unio) Heude	802
Minor (Margaritana) Lea	668	Montanus (Unio) Philippi	881
Minor (Margaron) Lea	562, 668	Montanus (Unio) Rafinesque	744
Minor (Strophitus) Conrad	668	Montezianus (Anodon) Sowerby	922
Minor (Unio) Lea	562	Montezuma (Anodonta) Lea	923
Minuana (Margarita) Lea	911	Montezuma (Glabaris) Lea	923
Minuana (Margaron) Lea	911	Montezuma (Margaron) Lea	923
Minuana (Monocondylæa) d'Orbigny	911	Monticola (Unio) Heude	802
Minuanus (Unio) Hanley	911	Moorenianus (Unio) Pætel	750
Minuta (Anodon) Heude	638	Mooresianus (Margaron) Lea	750
Mira (Grandidieria) Bourguignat	830	Mooresianus (Unio) Lea	750
Misellus (Trapezoides) Morelet	859	Moquinianus (Unio) Dupuy	699
Misoolensis (Unio) Schepman	852	Moreleti (Margaron) Lea	687
Misoolensis (Virgus) Schepman	852	Moreleti (Pseudodon) Crosse and Fischeh.	834, 838
Mississippiensis (Margaron) Lea	547	Moreleti (Unio) Deshayes	687
Mississippiensis (Unio) Conrad	547	Moreletianus (Unio) Heude	799
Mitchelli (Unio) Simpson	711	Moretonicus (Diplodon) Reeve	891
Modesta (Anodonta) Lea	647, 660	Moricandii (Anodonta) Lea	927
Modesta (Margaron) Lea	660	Moricandii (Glabaris) Lea	927
Modesta (Platiris) Lea	899	Moricandii (Margaron) Lea	927
Modesta (Spatha) Lea	899	Morini (Unio) Morelet	702
Modestus (Diplodon) Kuster	881	Morinus (Unio) Heude	851
Modestus (Margarita) Lea	876	Mortoni (Quadrula) Conrad	781
Modestus (Margaron) Lea	876	Mortoni (Unio) Conrad	781
Modestus (Unio) H. and A. Adams	876	Mortoniana (Anodonta) Lea	929
Modestus (Unio) Heude	845	Mortoniana (Margarita) Lea	929
Modestus (Unio) Kuster	881	Mortoniana (Margaron) Lea	929
Modica (Pleurobema) Lea	762	Mortonianus (Anodon) Sowerby	925
Modicella (Truncilla) Lea	518	Mortonianus (Glabaris) Lea	929

	Page.		Page.
Mossambicensis (Nodularia) von Martens.	813	Myersianus (Margaron) Lea	580
Mossambicensis (Unio) von Martens	813	Myersianus (Unio) Lea	577, 580
Mossulensis (Margaritana) Patet	687	Mysea Turton	679
Mosulensis (Margaron) Lea	687	Mytilloides (Pleurobema) Rafinesque	754
Mosulensis (Unio) Lea	687	Mytilloides (Unio) Conrad	754
Mouhoti (Monocondylæa) Lea	839	Mytiloides (Margarita) Lea	790
Mouhoti (Pseudodon) Lea	839	Mytiloides (Margaron) Lea	790
Mouhotiana (Monocondylæa) Lea	839	Mytiloides (Pleurobema) Rafinesque	765
Mouhotianus (Margaron) Lea	839	Mytiloides pyramidatus (Unio) Patet	790
Mouhotianus (Pseudodon) Morelet	839	Mytiloides (Unio) Deshayes	790
Mouhotianus (Unio) Sowerby	839	Mytiloides (Unio) Swainson	667
Moulinsiana (Unio) Dupuy	699	Mytiloides (Unio) Short and Eaton	788
Moulinsiana (Anodonta) Dupuy	624	Mytiloides (Unio) Swainson	793
Moussonianus (Margaron) Lea	707	Nagpoorensis (Margaron) Lea	842
Moussonianus (Unio) Lea	707	Nagpoorensis (Unio) Lea	842
Mucidis (Unio) Morelet	683	Nala Swainson	868
Mucronata (Mya) Eaton	703	Naiades . . . 501, 502, 503, 505, 507, 512, 513, 558, 687, 867	
Mucronatus (Unio) Barnes	703	Nana (Unio) Lamarck	691
Muelleri (Unio) Rossmässler	685	Nankingensis (Monocondylæa) Heude	839
Mühlfeldianus (Lampsilis) Lea	555	Nankingensis (Pseudodon) Heude	839
Mühlfeldianus (Margarita) Lea	555	Nanus (Unio) Dupuy	697
Mühlfeldianus (Margaron) Lea	555	Napoensis (Anodonta) Lea	920
Mühlfeldianus (Unio) Lea	555	Napoensis (Glabaris) Lea	920
Multicolor (Parreysia) von Martens	847	Napoensis (Margaron) Lea	920
Multicolor (Unio) von Martens	847	Nashvillianus (Margarita) Lea	547
Multidentatus (Margaron) Lea	841	Nashvillianus (Margaron) Lea	547
Multidentatus (Unio) Philippi	840, 841	Nashvillianus (Unio) Lea	546
Multiplicata (Quadrula)	766	Nashvilliensis (Unio) Lea	547
Multiplicatus (Margarita) Lea	771	Nasuta (Eurynea) Agassiz	546
Multiplicatus (Margaron) Lea	771	Nasuta (Mya) Eaton	546
Multiplicatus (Unio) Lea	503, 770	Nasuta (Unio) Lamarck	703
Multiradiatus (Lampsilis) Lea	532	Nasutilus (Unio) Lea	735
Multiradiatus (Margarita) Lea	532	Nasutulus (Margaron) Lea	735
Multiradiatus (Margaron) Lea	532	Nasutulus (Unio) Lea	735
Multiradiatus (Unio) Lea	532	Nasutus (Lampsilis) Say	545
Multiradiatus (Unio) Sowerby	534	Nasutus (Margarita) Lea	546
Multistriatus (Margarita) Lea	879	Nasutus (Margaron) Lea	546
Multistriatus (Unio) Lea	879	Nasutus (Unio) Conrad	735
Multisulcata (Castalia) Hupé	865	Nasutus (Unio) Say	545
Multisulcatus (Tetraplodon) Hupé	864, 865	Nasus Simpson	839
Mundus (Margaron) Lea	747	Natalensis (Margaron) Lea	825
Mundus (Unio) Lea	746, 747	Natalensis (Platiris) Lea	898
Murchisonianus (Margarita) Lea	808	Natalensis (Spatha) Lea	898
Murchisonianus (Margaron) Lea	808	Natalensis (Unio) Lea	825
Murchisonianus (Unio) Lea	808	Natolicus (Unio) Kuster	700
Murinus (Unio) Heude	851	Navicella (Anodon) Heude	638
Murrayensis (Margaron) Lea	753	Navicula (Anodonta) Patet	638
Murrayensis (Pleurobema) Lea	753	Naviculoides (Margaron) Lea	732
Murrayensis (Unio) Lea	753	Naviculoides (Unio) Lea	732
Muscus (Unio) Spengler	744	Naviformis (Unio) Lamarck	773
Mussolianus (Unio) Kuster	687	Navigoliformis (Margaron) Lea	825
Mutabilis (Diplodon) Lea	894	Navigoliformis (Unio) Lea	825
Mutabilis (Margaron) Lea	894	Nebulosus (Lampsilis) Conrad	553
Mutabilis (Unio) Lea	894	Nebulosus (Unio) Conrad	553
Mutabilis (Unio) Reeve	893	Negatus (Margaron) Lea	787
Mutatus (Unio) Mousson	818	Negatus (Unio) Lea	787
Mutela Scopoli 501, 505, 506, 902, 903, 930		Neglecta (Symphynota) Lea	664
Mutelidæ 501, 502, 505, 507, 512, 515, 895, 906, 933		Neglectus (Margaron) Lea	664
Mutelina	900	Neglectus (Unio) Lea	664
Mutica (Arconia) Heude	860	Neglectus (Unio) Sowerby	724
Mya Linnæus	530, 620	Nehringi (Castalina) von Ihering	865
Mycetopina	502	Nehringi (Glabaris) von Ihering	930
Mycetopoda d'Orbigny	507, 515, 905, 933	Neislerii (Margaron) Lea	771
Mycetopodidæ	501	Neislerii (Quadrula) Lea	771
Mycetopus d'Orbigny	501, 930, 933, 935	Neislerii (Unio) Lea	771
Myersianus (Hyriopsis) Lea	580	Nephronaias Crosse and Fischer	506, 514, 591

	Page.		Page.
Nervosa (Unio) Rafinesque.....	744	Nodulatus (Unio) Say.....	782
Nervosus (Unio) Say.....	605	Nodulosa (Castalia) H. and A. Adams....	799
Neusensis (Margaron) Lea.....	722	Nodulosa (Mya) Wood.....	682, 798
Neusensis (Unio) Lea.....	722	Nodulosa (Quadrula) Wood.....	797, 798
Newcombiana (Brazzæa) Bourguignat....	907	Nodulosa (Unio) Lamarek.....	682
Newcombianus (Margaron) Lea.....	607	Nodulosus (Margarita) Lea.....	799
Newcombianus (Unio) Lea.....	607	Nodulosus (Margaron) Lea.....	799
Newenhuisi (Unio) Schepman.....	803	Nodulosus (Unio) Hanley.....	793
Newtonensis (Anodonta) Lea.....	633	Nodulosus (Unio) Reeve.....	831
Newtonensis (Margarita) Lea.....	633	Nodulosus (Unio) Say.....	779
Newtoniensis (Anodonta) Troschel.....	633	Nolani (Unio) Lea.....	717
Nexus (Unio) Say.....	519	Nopalatensis (Anodonta) Sowerby.....	646
Ngesianus (Parreysia) von Martens.....	847	Normandi (Anodonta) Dupuy.....	624
Ngesianus (Unio) von Martens.....	847	Northamptonensis (Unio) Lea.....	728
Nicaragua (Anodonta) Philippi.....	932	Notatus (Margarita) Lea.....	554
Nicaraguensis (Margaron) Lea.....	607	Notatus (Margaron) Lea.....	554
Nicaraguensis (Plagiola) Lea.....	607	Notatus (Unio) Conrad.....	556
Nicaraguensis (Unio) Lea.....	607	Notatus (Unio) Lea.....	554
Nicklianus (Unio) Kuster.....	771	Novæ-hollandiæ (Diplodon) Gray.....	893
Nickliniana (Quadrula) Lea.....	771	Novæ-hollandiæ (Margarita) Lea.....	893
Nicklinianus (Margarita) Lea.....	771	Novæ-hollandiæ (Margaron) Gray.....	893
Nicklinianus (Margaron) Lea.....	771	Novæ-hollandiæ (Unio) Gray.....	893
Nicklinianus (Plectomerus) Conrad.....	771	Novi-eboraci (Margarita) Lea.....	552
Nicklinianus (Unio) Lea.....	771	Novi-eboraci (Margaron) Lea.....	552
Nicobarica (Alasmodonta) Mörch.....	838	Novi-eboraci (Unio) Lea.....	552
Nicobaricus (Mytilus) Fabricius.....	838	Nubilis (Margaron) Lea.....	724
Nicobaricus (Pseudodon) Mörch.....	838	Nubilis (Unio) Lea.....	724
Nigellus (Margaron) Lea.....	718	Nucleopsis (Margarita) Lea.....	755
Nigellus (Unio) Lea.....	718	Nucleopsis (Pleurobema) Conrad.....	755
Nigerrimus (Lampsilis) Lea.....	551	Nucleopsis (Unio) Conrad.....	755
Nigerrimus (Margaron) Lea.....	551	Nucleus (Margaron) Lea.....	815
Nigerrimus (Unio) Lea.....	551	Nucleus (Nodularia) Lea.....	815
Nigra (Modiolaria) Lea.....	648	Nucleus (Unio) Lea.....	815
Nigra (Unio) Rafinesque.....	707	Nuculinus (Unio) Philippi.....	607
Nigricans (Anodon) Heude.....	638	Numidica (Anodonta) Bourguignat.....	627
Nigrinus (Margaron) Lea.....	560	Nuttalliana (Anodonta) Lea.....	629
Nigrinus (Unio) Lea.....	560	Nuttalliana (Margarita) Lea.....	629
Nigritus (Diplodon) Simpson.....	881	Nuttalliana (Margaron) Lea.....	629
Nilotica emini (Mutela) von Martens.....	904	Nuttalliana (Nodularia) Lea.....	817
Nilotica gerrardi (Nodularia) von Martens	822	Nuttallianus (Margaron) Lea.....	817
Nilotica (Iridina) Sowerby.....	904	Nuttallianus (Unio) Lea.....	817
Nilotica (Mutela) Sowerby.....	904	Nux (Margaron) Lea.....	758
Nilotica (Nodularia) Cailliaud.....	821, 822	Nux (Pleurobema) Lea.....	757, 758
Nilotica (Platiris) Lea.....	904	Nux (Unio) Kuster.....	524
Nilotica (Unio) Cailliaud.....	821	Nux (Unio) Lea.....	758
Niloticus (Margarita) Lea.....	821	Nux-persicus (Margaron) Dunker.....	808
Niloticus (Margaron) Lea.....	821	Nux-persicus (Unio) Dunker.....	808
Niloticus (Unio) Kuster.....	825	Nyassæ (Unio) Sowerby.....	848
Niloticus (Unio) Sowerby.....	879	Nyassaensis (Margaron) Lea.....	848
Nilssonii (Anodonta) Kuster.....	625, 643	Nyassaensis (Parreysia) Lea.....	847, 848
Nipponensis (Unio) von Martens.....	809	Nyassaensis (Platiris) Lea.....	898
Nitens (Margaron) Lea.....	549	Nyassaensis (Spatha) Lea.....	898
Nitens (Unio) Lea.....	549	Nyassaensis (Spathella) Ancey.....	898
Nitidulus (Unio) Kuster.....	874	Nyassaensis (Unio) Lea.....	848
Nobilis (Unio) Conrad.....	776, 778	Nyassaensis (Unio) Smith.....	848
Nocturnus (Margaron) Lea.....	887	Nyassana (Spatha) Bourguignat.....	898
Nocturnus (Unio) Lea.....	887	Nyassanus (Unio) Bourguignat.....	848
Nodifera (Quadrula) Conrad.....	782	Obesus blandinianus (Unio) Lea.....	742
Nodiferus (Unio) Conrad.....	782	Obesus (Margarita) Lea.....	741
Nodosa (Mya) Eaton.....	774	Obesus (Margaron) Lea.....	741
Nodosa (Mya) Gmelin.....	682	Obesus paludiculus (Unio) Gould.....	745
Nodosus (Unio) Barnes.....	774	Obesus (Unio) Lea.....	711, 741
Nodularia Conrad.....	506,	Obfuscus (Margaron) Lea.....	559
	515, 806, 807, 809, 810, 811, 822, 862	Obfuscus (Unio) Lea.....	559
Nodulata (Obliquaria) Rafinesque.....	744	Oblatus (Margaron) Lea.....	730
Nodulatus pustulatus (Unio) Pætel.....	782	Oblatus (Unio) Lea.....	730, 815

	Page.		Page.
Obliqua (Hyria) Pætel	871	Ocmulgeensis (Margaron) Lea	727
Obliqua (Margarita) Lea	788	Ocmulgeensis (Unio) Lea	727
Obliqua (Quadrula) Lamarck	788	Odatella Rafinesque	654
Obliqua (Unio) Wood	790	Ogecheensis (Lampsilis) Conrad	550
Obliquaria Rafinesque ... 506, 510, 514, 610, 772, 791		Ogecheensis (Unio) Conrad	550
Obliquata (Obliquaria) Rafinesque	744	Ohioensis (Anodonta) Rafinesque	648
Obliquatus (Unio) Say	520	Ohioensis (Metaptera) Conrad	574
Obliquiradiatus (Unio) Reeve	536	Ohioensis (Symphynota) Ferussac	574
Obliquus (Margaron) Lea	788	Ohioensis (Unio) Say	574
Obliquus (Prisodon) Schumacher	868, 871	Oleivora (Solenais) Heude	656
Obliquus (Unio) Conrad	793	Oleivorus (Mycetopus) Heude	656
Obliquus (Unio) Hanley	788	Olivaria (Amblema) Rafinesque	744
Obliquus (Unio) Potiez and Michaux	523	Olivaria (Nodularia) Lea	816, 817
Oblita (Anodonta) Lea	660	Olivarius (Margarita) Lea	817
Oblita (Margaron) Lea	660	Olivarius (Unio) Conrad	602
Oblonga (Anodonta) Millet	623	Olivarius (Unio) Lea	817
Oblongata (Mya) Wood	536	Omiensis (Nodularia) von Heimbürg	810
Oblongus (Unio) Lea	726	Omiensis (Unio) von Heimbürg	810
Obnubilus (Margaron) Lea	716	Opaca (Anodonta) Lea	645
Obnubilus (Unio) Lea	716	Opaca (Margaron) Lea	645
Obovalis (Obovaria) Rafinesque	599, 744	Opacatus (Unio) Crosse and Fischer	712
Obovalis Simpson	840	Opacus (Margaron) Lea	717
Obovalis (Unio) Conrad	789	Opacus (Unio) Lea	716
Obovalis (Unio) Say	793	Opalina (Anodon) Anthony	644
Obovaria Rafinesque	506, 510, 514, 599	Opalina (Anodonta) Kuster	625
Obovata (Mya) Solander	744	Opalinus (Unio) Anthony	552
Obovatus (Lampsilis) Lea	549, 550	Opperti (Gabillotta) Bourguignat	650
Obscurus (Margarita) Lea	550	Opperti (Margaritana) Pætel	650
Obscurus (Margaron) Lea	550	Opperti (Pseudodon) Westerlund	650
Obscurus (Unio) Lea	549	Opperti (Unio) Bourguignat	650
Obtusa (Anodon) Heude	638	Oratus (Unio) Conrad	529
Obtusa (Anodonta) Heude	639	Orbicularis (Anodon) Heude	638
Obtusa (Cameronia) Bourguignat	910	Orbicularis (Monocondylæa) Pætel	835
Obtusa (Margarita) Lea	931	Orbicularis (Monocondylus) Morelet	835
Obtusa (Margaron) Lea	931	Orbicularis (Pseudodon) Morelet	835
Obtusa (Unio) d'Orbigny	882	Orbicularis (Lampsilis) Hildreth	510
Obtusata (Anodon) Heude	639	Orbiculatus (Margarita) Lea	540
Obtusula (Anodonta) Hupe	921	Orbiculatus (Margaron) Lea	540
Obtusula (Glabaris) von Ihering	921	Orbiculatus (Unio) Hildreth	540, 600
Obtusula (Margaron) Lea	921	Orbignyana (Unio) Hupé	866
Obtusulus (Anodon) Spix	931	Orbigny (Castalina) Hupé and Deville	866
Obtusulus (Diplodon) d'Orbigny	882	Orbigny (Margaron) Lea	866
Obtusulus (Glabaris) Spix	931	Orbigny (Unio) Hupé	885
Obtusulus (Margaron) Lea	537	Orbigny (Unio) Hupé and Deville	866
Obtusulus (Unio) Lea	537	Oregonensis (Anodonta) Lea	628
Obtusulus var. b. lituratum (Anodon) Spix	931	Oregonensis (Margarita) Lea	628
Obuncus (Unio) Lea	747	Oregonensis (Margaron) Lea	594, 628
Occata (Nodularia) Lea	813	Oregonensis (Nephronaias) Lea	594
Occatus (Unio) Lea	813	Oregonensis (Unio) Lea	594
Occidens (Margarita) Lea	527	Oriens (Unio) Lea	655
Occidens (Margaron) Lea	527	Oriens (Unio) Sowerby	544
Occidens (Unio) Lea	527	Orientalis (Margaron) Lea	820
Occidentalis (Barbala) Pætel	584	Orientalis (Nodularia) Lea	819
Occidentalis (Dipsas) Heude	584	Orientalis (Unio) Bourguignat	694
Occidentalis (Lampsilis) Conrad	551	Orientalis (Unio) Lea	819
Occidentalis (Margaron) Lea	551	Ornata (Pleurobema) Lea	749
Occidentalis (Mycetopoda) Clessin	934	Ornatus (Margaron) Lea	749
Occidentalis (Unio) Conrad	551	Ornatus (Unio) Lea	749
Occultus (Margaron) Lea	717	Ornatus (Unio) Sowerby	528
Occultus (Unio) Lea	717	Orontesensis (Margaron) Lea	695
Ochracea (Margarita) Lea	531	Orontesensis (Unio) Lea	695
Ochracea (Mya) Eaton	530	Orphaensis (Margaron) Lea	695
Ochracea (Symphynota) Lea	531	Orphaensis (Unio) Lea	695
Ochraceus (Lampsilis) Say	530	Orthonymus Agassiz	765
Ochraceus (Margaron) Lea	530	Ortonii (Prisodon) Lea	872
Ochraceus (Unio) Say	529, 530	Ortonii (Unio) Lea	872

	Page.		Page.
Osbecki (Margaron) Lea	808	Tallescens (Unio) Lea	537
Osbecki (Unio) Philippi	808	Palliatius (Unio) Ravenel	730
Oseari (Unio) B. H. Wright	719	Pallida (Anodon) Anthony	632
Ostreata (Quadrula) Morelet	796	Pallidofulvus (Margaron) Lea	753
Ostreatus (Unio) Morelet	796	Pallidofulvus (Unio) Lea	753
Othealoogensis (Margaron) Lea	521	Paludicola (Mutelina) Rochebrune	906
Othealoogensis (Truncella) Lea	521	Paludicolor (Unio) Conrad	743
Othealoogensis (Unio) Lea	521	Paludicolus (Margaron) Lea	743
Ovalis (Margarita) Lea	685	Paludicolus (Unio) Gould	743
Ovalis (Mya) Donovan	682	Paludosus (Anodon) Turton	623
Ovalis (Mya) Montagu	685	Paludosus (Lampsilis) Morelet	576
Ovalis (Pseudodon) Morelet	837	Paludosus (Unio) Morelet	576
Ovalis (Unio) Brown	685	Panacoensis (Unio) von dem Busch	705
Ovalis (Unio) Dupuy	699	Pangalliensis (Spatha) Rochebrune	902
Ovalis (Unio) Sowerby	682	Paphos (Anodonta) Rafinesque	933
Ovata (Eglia) Swainson	530	Papyracea (Margaron) Lea	618
Ovata (Anodonta) Lea	642	Papyraceum (Alasmodon) Rafinesque	673
Ovata (Anodonta) Sowerby	644	Papyraceus (Anodon) Anthony	618
Ovata (Anodonta) Stark	648	Papyraceus (Lampsilis) Gould	562
Ovata (Castalia) Sowerby	864	Papyraceus (Margaron) Lea	563
Ovata (Iridina) Swainson	908	Papyraceus (Unio) Gould	562
Ovata (Margarita) Lea	642	Paradoxa (Cameronia) Bourguignat	909
Ovata (Margaron) Lea	642	Paraguayana (Margarita) Lea	911
Ovata (Mya) Donovan	685	Paraguayana (Margaron) Lea	874, 911
Ovata (Mya) Eaton	530	Paraguayana (Monocondylaea) d'Or-	
Ovata (Mysea) Turton	685	bigny	910, 911
Ovata (Platiris) Lea	908	Paraguayana (Unio) Hanley	911
Ovata (Quadrula) Simpson	801	Paraguayanus (Unio) von Martens	894
Ovata (Unio) Deshayes	527	Paraguayensis (Margaron) Lea	874
Ovatus (Anodon) Sowerby	922	Paraguayensis (Unio) Lea	874
Ovatus (Lampsilis) Say	530	Parallelipipidon (Diplodon) Lea	834
Ovatus (Margarita) Lea	530	Parallelipipidon (Margarita) Lea	884
Ovatus (Margaron) Lea	530	Parallelipipidon (Margaron) Lea	884
Ovatus ornatus (Unio) Conrad	528	Parallelipipidon (Unio) Lea	884
Ovatus (Pleiodon) Swainson	907, 908	Parallelus (Unio) Conrad	740
Ovatus (Tetraplodon) Sowerby	864	Paramattensis (Margaron) Lea	892
Ovatus (Unio) Say	526, 530	Paramattensis (Unio) Lea	892
Ovatus var. b (Unio) Lamarek	529	Paranensis (Diplodon) Lea	887
Oviformis (Margaron) Lea	748	Paranensis (Margarita) Lea	887
Oviformis (Pleurobema) Conrad	748	Paranensis (Margaron) Lea	887
Oviformis (Unio) Lea	748	Paranensis (Unio) Lea	887
Oxyrhynehus (Nodularia) von Martens	807	Parchappii (Margarita) Lea	911
Oxyrhynehus (Unio) von Martens	807	Parchappii (Monocondylaea) d'Orbigny	911
Ozarkensis (Lampsilis) Call	557	Parchappii (Margaron) Lea	911
Ozarkensis (Unio) Call	557	Parchappii (Unio) Hanley	911
Pachostea (Obovaria) Rafinesque	744	Pareus (Diplodon) Lea	880
Pachynaia Crosse and Fischer	794	Pareus (Margaron) Lea	880
Pachyodon (Pleiodon) Bourguignat	909	Pareus (Unio) Lea	880
Pachysoma (Margaron) Lea	813	Parishii (Anodon) Gray	915
Pachysoma (Nodularia) Benson	813	Parishii (Leila) H. and A. Adams	915
Pachysoma (Unio) Benson	813	Parishii (Margarita) Lea	915
Pachystoma (Unio) Petel	813	Parishii (Margaron) Lea	915
Pacifica (Anodon) Heude	638	Parma (Unio) Benson	843
Pahrangensis (Rectidens) Smith	854	Parreysi sehweinfurthi (Unio) von Mar-	
Pahrangensis (Unio) Smith	854	tens	813
Paivanus (Unio) Morelet	582, 819	Parreysia Conrad	506, 508, 515, 840
Palembangensis (Unio) Strubell	862	Parreysi (Unio) Philippi	821
Pallatus (Margarita) Lea	730	Parva (Mya) Eaton	564
Paliatus (Margaron) Lea	730	Parvula (Anodonta) Drouet	624
Paliatus (Unio) Ravenel	730	Parvula (Barbala) Petel	585
• Pallegoixi (Anodon) Sowerby	859	Parvula (Cristaria) Heude	585
Pallegoixi (Trapezoideus) Sowerby	859	Parvulus (Dipsas) Heude	585, 586
Pallens (Unio) Kuster	683	Parvulus (Margaron) Lea	590
Pallens (Unio) Rafinesque	744	Parvulus (Medionidus) Lea	590
Pallens (Unio) Rossmässler	686	Parvulus (Unio) Lea	590
Pallescens (Margaron) Lea	538	Parvus (Lampsilis) Barnes	564

	Page.		Page.
Parvus (Margarita) Lea.....	564	Pellucidus (Unio) Lea.....	562
Parvus (Margaron) Lea.....	564	Penchinatianus (Unio) Bourguignat.....	696
Parvus (Unio) Barnes.....	564	Penicillatus (Anodon) Gray.....	923
Parvus (Unio) Conrad.....	563	Penicillatus (Margaron) Lea.....	590
Pasehalis (Unio) Heude.....	802	Penicillatus (Medionidus) Lea.....	590
Pastasana (Anodonta) Clessin.....	930	Penicillatus (Unio) Lea.....	590
Pastasanus (Glabaris) Clessin.....	930	Peninsularis (Trapezoides) Simpson.....	859
Patagonia (Anodonta) Lamarek.....	916	Penisatus (Unio) Fischer and Crosse.....	853
Patagonia (Margarita) Lea.....	916	Penita (Truncella) Conrad.....	518
Patagonia (Margaron) Lea.....	916	Penitus (Margarita) Lea.....	518
Patagonia (Unio) d'Orbigny.....	885	Penitus (Margaron) Lea.....	518
Patagonicus (Diplodon) d'Orbigny.....	885	Penitus (Unio) Conrad.....	518
Patagonicus felix (Glabaris) Pilsbry.....	917	Pennsylvania (Anodonta) Lamarek.....	618
Patagonicus (Glabaris) Lamarek.....	916	Pepiniana (Anodonta) Lea.....	647
Patagonicus (Margarita) Lea.....	885	Pepiniana (Margarita) Lea.....	647
Patagonicus (Margaron) Lea.....	885	Pepiniana (Margaron) Lea.....	647
Patelloides (Diplodon) Lea.....	886, 888	Pequotianus (Unio) Linsley.....	744
Patelloides (Margaron) Lea.....	886	Peræformis (Diplodon) Lea.....	836
Patelloides (Unio) Lea.....	886	Peræformis (Margaron) Lea.....	836
Pattinoides (Unio) Lea.....	747	Peræformis (Unio) Lea.....	886
Patsaligensis (Pleurobema) Simpson.....	761	Perakensis (Rectidens) de Morgan.....	854
Patulus (Margarita) Lea.....	746	Perakensis (Unio) de Morgan.....	854
Patulus (Margaron) Lea.....	746	Percoarctatus (Margaron) Lea.....	725
Patulus (Unio) Lea.....	746	Percoarctatus (Unio) Lea.....	725
Pauciplicatus (Unio) Lea.....	768	Perecompressa (Quadrula) von Martens.....	736
Paulista (Diplodon) von Ihering.....	873	Perecompressus (Unio) von Martens.....	736
Paulista (Unio) von Ihering.....	873	Perdix (Lampsilis) Lea.....	512
Paulus (Lampsilis) Lea.....	566	Perdix (Margarita) Lea.....	512
Paulus (Margaron) Lea.....	566	Perdix (Margaron) Lea.....	512
Paulus (Unio) Lea.....	566	Perdix (Unio) Lea.....	512
Paupercula (Quadrula) Lea.....	783	Perdix (Unio) Reeve.....	541
Pauperculus (Margaron) Lea.....	783	Perlatus (Margaron) Lea.....	731
Pauperculus (Unio) Lea.....	783	Perlatus (Unio) Lea.....	731
Paviei (Chamberlainia) Morelet.....	582	Perlensis (Unio) Patel.....	768
Paviei (Unio) Morelet.....	582	Perlucens (Margaron) Lea.....	715
Pavonia (Anodonta) Lea.....	617	Perlucens (Unio) Lea.....	714
Pavonia (Margarita) Lea.....	617	Permiseens (Lampsilis) Lea.....	569
Pavonia (Margaron) Lea.....	618	Permiseens (Margaron) Lea.....	569
Pavonina (Dysnomia) Rochebrune.....	862	Permiseens (Unio) Lea.....	569
Pavonius (Strophitus) Baker.....	618	Pernodosus (Margaron) Lea.....	780
Pawensis (Unio) Lea.....	742	Pernodosus (Unio) Lea.....	780
Paxyodon Schumacher.....	501, 868, 869	Perobliquus (Unio) Conrad.....	520
Pazi (Castalia) Hidalgo.....	884	Perovalis (Lampsilis) Conrad.....	531
Pazi (Diplodon) Hidalgo.....	884	Perovalis (Margarita) Lea.....	531
Pazii (Anodonta) Lea.....	913	Perovalis (Margaron) Lea.....	531
Pazii (Glabaris) Lea.....	913	Perovalis (Unio) Conrad.....	531
Pazii (Margaron) Lea.....	816, 911, 913	Perovalis (Unio) Sowerby.....	532
Pazii (Monocondylæa) Lea.....	911	Perovata (Pleurobema) Conrad.....	755
Pazii (Nodularia) Lea.....	816	Perovatus (Margarita) Lea.....	755
Pazii (Unio) Lea.....	816	Perovatus (Margaron) Lea.....	755
Pealei (Unio) Lea.....	602	Perovatus (Unio) Conrad.....	755
Pearlensis (Unio) Conrad.....	763	Perpastus (Lampsilis) Lea.....	532
Pectitus (Unio) Conrad.....	520	Perpastus (Margaron) Lea.....	532
Pectinatum (Tetraplodon) Spix.....	863	Perpastus (Unio) Lea.....	532
Pectorosus (Margarita) Lea.....	543	Perpictus (Margaron) Lea.....	554
Pectorosus (Unio) Conrad.....	542	Perpictus (Unio) Lea.....	554
Peculiaris (Diplodon) Lea.....	874	Perplexa cincinnatiensis (Truncella) Lea.....	523
Peculiaris (Margaron) Lea.....	874	Perplexa rangiana (Truncella) Lea.....	523
Peculiaris (Unio) Lea.....	874	Perplexa (Truncella) Lea.....	522
Pegias Simpson.....	506, 514, 660	Perplexus (Margarita) Lea.....	522
Peguensis (Margaron) Lea.....	835, 853	Perplexus (Margaron) Lea.....	523
Peguensis (Monocondylæa) Anthony.....	835	Perplexus (Truncella) Rafinesque.....	518
Peguensis (Unio) Anthony.....	853	Perplexus (Unio) Lea.....	522
Pellis-lacerti (Unio) Morelet.....	815	Perplexus (Unio) Rafinesque.....	522
Pellucidus (Lampsilis) Lea.....	562	Perplicata (Quadrula) Conrad.....	767, 768
Pellucidus (Margaron) Lea.....	532	Perplicata quintardii (Quadrula) Cragin.....	768

	Page.		Page.
Perplicatus (Margaron) Lea.....	768	Pictorum (Baphie) Meuschen.....	682
Perplicatus (Plectomerus) Conrad.....	768	Pictorum, etc. (Unio) Chemnitz.....	535
Perplicatus (Unio) Conrad.....	767	Pictorum longirostris (Unio) Westerlund.....	808
Perpurpureus (Margaron) Lea.....	558	Pictorum (Lymnium) Oken.....	682
Perpurpureus (Lampsilis) Lea.....	558	Pictorum (Margarita) Lea.....	682
Perpurpureus (Unio) Lea.....	558	Pictorum (Margaron) Lea.....	682
Perradiatus (Margaron) Lea.....	532	Pictorum (Mya) Gmelin.....	696
Perradiatus (Unio) Lea.....	532	Pictorum (Mya) Linnaeus.....	680
Personata (Scalcarnia) Agassiz.....	522	Pictorum (Mysea) Turton.....	682
Personata (Truncella) Say.....	522	Pictorum (Unio) Brown.....	685
Personatus (Margarita) Lea.....	522	Pictorum (Unio) Draparnaud.....	696
Personatus (Margaron) Lea.....	522	Pictorum (Unio) Guerin.....	686
Personatus (Unio) Conrad.....	792	Pictorum (Unio) Linnaeus.....	503,
Personatus (Unio) Say.....	522	513, 678, 680, 682, 685, 686, 798, 808	
Perstriatus (Margaron) Lea.....	731	Pictorum (Unio) Retzius.....	681
Perstriatus (Unio) Lea.....	731	Pictus (Anodon) Swainson.....	648, 933
Persulcatus (Margaron) Lea.....	596	Pictus (Lampsilis) Lea.....	542
Persulcatus (Nephronaias) Lea.....	596	Pictus (Margarita) Lea.....	542
Persulcatus (Unio) Lea.....	596	Pictus (Margaron) Lea.....	542
Pertenuis (Margaron) Lea.....	664	Pictus (Unio) Sowerby.....	539
Pertenuis (Unio) Lea.....	664	Pierri (Pseudodon) Rochebrune.....	840
Peruviana (Unio) Lamarck.....	767	Pietri (Unio) Kobelt.....	690
Petersi (Mutchia) Clessin.....	899	Pietri (Unio) Locard.....	683
Petersi (Spathella) Bourguignat.....	899	Piger (Diplodon) Lea.....	875
Petrina (Quadrula) Gould.....	783	Piger (Margaron) Lea.....	875
Petrinus (Margaron) Lea.....	783	Piger (Unio) Lea.....	875
Petrinus (Unio) Gould.....	783	Pigerrimus (Unio) Crosse and Fischer.....	711
Petroi (Unio) Patel.....	688	Pilaris (Margaron) Lea.....	792
Petrovichii (Unio) Kuster.....	683	Pilaris (Quadrula) Lea.....	792
Petterianus (Margaron) Lea.....	699	Pilaris (Unio) Reeve.....	788
Petterianus (Unio) Kuster.....	699	Pilatus (Unio) Lea.....	811
Pfeifferi (Unio) Dunker.....	879	Pilea (Simpson).....	522
Pfeifferiana (Margaritana) Bernardi.....	900	Pileus (Margarita) Lea.....	522
Pfeifferiana (Margaron) Lea.....	900	Pileus (Margaron) Lea.....	522
Pfeifferiana (Spatha) Bernardi.....	900	Pileus (Unio) Lea.....	522
Pfisteri inspiratus (Ptychorhynchus)		Pilsbryi (Unio) Marsh.....	769
Heude.....	851	Pilsbryconcha Simpson.....	505, 506, 511, 514, 587
Pfisteri inspiratus (Unio) Heude.....	851	Pinax (Unio) Benson.....	843
Pfisteri (Ptychorhynchus) Heude.....	851	Pinchonianus (Hyriopsis) Heude.....	580
Pfisteri (Unio) Heude.....	850, 851	Pinchonianus (Unio) Heude.....	580
Pharaoonum (Chambardia).....	902	Pinei (Unio) B. H. Wright.....	718
Phaselus (Margaron) Lea.....	815	Pinguis (Lampsilis) Lea.....	540
Phaselus (Nodularia) Lea.....	815	Pinguis (Margaron) Lea.....	540
Phaselus (Unio) Lea.....	815	Pinguis (Unio) Lea.....	540
Phaseolus (Margarita) Lea.....	612	Pinkstoni (Pleurobema) S. H. Wright.....	758
Phaseolus (Margaron) Lea.....	612	Pinkstoni (Unio) S. H. Wright.....	758
Phaseolus (Ptychobranchus) Hildreth.....	612	Piscatorum (Anodon) Heude.....	638
Phaseolus (Unio) Hildreth.....	503, 612	Pisciculus (Cuneopsis) Heude.....	805
Phayresi (Unio) Theobald.....	842	Pisciculus (Unio) Heude.....	805
Philippi (Unio) Dupuy.....	699	Piscinalis (Anodonta) Nilsson.....	623
Philippianus (Glabaris) Simpson.....	919	Piscinalis (Unio) Rossmässler.....	698
Philippianus (Unio) Kuster.....	891	Placitus (Margaron) Lea.....	536
Philipsii (Unio) Hanley.....	611	Placitus (Unio) Lea.....	536
Philipsii (Margarita) Lea.....	611	Plagiodon Lea.....	913
Phillipsi (Margaron) Lea.....	611	Plagiola (Rafinesque).....	506, 510, 514, 603, 772
Phillipsi (Unio) Catlow and Reeve.....	611	Plagiosoma (Unio) Benson.....	843
Phillipsii (Unio) Conrad.....	611	Plana (Anodonta) Lea.....	643
Phillipsii (Unio) Reeve.....	523	Plana (Margarita) Lea.....	643
Pholadiformis (Anodon) Sowerby.....	633	Plana (Margaron) Lea.....	643
Physunio Simpson.....	506, 515, 830	Planus (Lampsilis) Lea.....	556
Pianensis (Unio) Farines.....	692	Planus (Margaron) Lea.....	556
Piceus (Diplodon) Lea.....	877	Planus (Unio) Lea.....	556
Piceus (Margaron) Lea.....	877	Planicostatus (Lampsilis) Lea.....	553
Piceus (Unio) Lea.....	877	Planicostatus (Margaron) Lea.....	553
Pietetiana (Anodonta) Brot.....	625	Planicostatus (Unio) Lea.....	553
Pictorum amurensis (Unio) Mousson.....	808	Planior (Margaron) Lea.....	763

	Page.		Page.
Planior (Pleurobema) Lea	763	Pliculosus (Unio) von Martens.....	862
Planior (Unio) Lea	763	Plombarius (Unio) Villa	744
Planilaterus (Margaron) Lea	722	Plumbea (Chama) Chemnitz.....	797
Planilaterus (Unio) Conrad	722	Plumbea (Quadrula) Chemnitz.....	797
Planivalvis (Lampsilis) Morelet.....	576	Plumbeus (Unio) Ferussac.....	797
Planivalvis (Unio) Morelet.....	576	Poeyanus (Margaron) Lea.....	738
Plantii (Margaron) Lea	729	Poeyanus (Unio) Lea.....	738
Plantii (Unio) Lea	729	Poirieri (Zaira) Rochebrune	862
Planulata (Margaron) Lea	836	Polita (Anodonta) Mousson.....	588
Planulata (Microcondylaea) Patet.....	836	Polita (Margaron) Lea	588
Planulata (Monocondylaea) Lea.....	836	Politus (Anodon) Sowerby	588
Planulata (Pseudodon) Conrad	836	Politus (Unio) Say	792
Planulatus (Unio) Lea	612	Polymorpha (Anodonta) Lea	648
Planus (Unio) Conrad.....	574	Polymorphus (Unio) B. H. Wright.....	707
Platolus (Unio) Rafinesque.....	744	Polystictio-scriptus (Quadrula) Heude.....	799
Platiris Lea	501	Polystictio-scriptus (Unio) Heude	799
Platyrhynchus (Margarita) Lea.....	683	Polystictus (Quadrula) Heude	799
Platyrhynchus (Margaron) Lea	683	Polystictus (Unio) Heude	799
Platyrhynchus (Unio) Rossmässler	683	Ponderosa (Anodonta) C. Pfeiffer.....	626
Platyrhynchoideus (Unio) Westerlund ..	684	Ponderosa (Mya) Dillwyn.....	804
Platyrhynchoideus (Unio) Dupuy	513, 684	Ponderosus (Bariosta) Rafinesque.....	744
Platyrhynchoideus (Unio) Sowerby	684	Ponderosus (Margarita) Lea	804
Pleasii (Lampsilis) Marsh.....	583, 557	Ponderosus (Margaron) Lea.....	804
Pleasii (Unio) Marsh	557	Ponderosus (Paxyodon) Schumacher	870
Plectophorus (Plectomerus) Conrad	706	Ponderosus (Unio) Hanley.....	804
Plectophorus (Unio) Conrad	706	Popeii (Margaron) Lea	738
Pleiodon Conrad	501, 506, 515, 907, 908	Popeii (Unio) Lea	737, 738
Plena (Quadrula) Lea.....	790	Popenoi (Unio) Call.....	610
Plenus (Margaron) Lea	790	Poreifer (Anodon) Gray.....	932
Plenus (Unio) Lea	790	Porphyreus (Lampsilis) Lea.....	538
Plethobasus Simpson.....	764	Porphyreus (Margaron) Lea	538
Pletholophus Simpson.....	585	Porphyreus (Unio) Lea.....	538
Pleurobema Rafinesque	506,	Porrectus (Unio) Conrad.....	740
509, 510, 514, 745, 754, 760, 774, 793		Postellii (Margaron) Lea	723
Plexus (Unio) Conrad.....	701, 702	Postellii (Unio) Lea.....	723
Plicata (Anodonta) Haldeman.....	659	Poulsoni (Symphynota) Ferussac	568
Plicata (Anodonta) Schrenck.....	583	Poulsoni (Unio) Conrad.....	568
Plicata (Barbala) H. and A. Adams.....	583	Poulsoni (Unio) Sowerby	568
Plicata clessini (Dipsas) Kobelt.....	584	Powellii (Lampsilis) Lea.....	538
Plicata (Cristaria) Leach.....	583, 584, 903	Powellii (Margaron) Lea	538
Plicata hippopæa (Quadrula) Lea.....	767	Powellii (Unio) Lea.....	538
Plicata japonica (Dipsas) Kobelt.....	585	Præclara (Anodonta) Bourguignat.....	649
Plicata (Margaron) Lea	659	Præclarum (Colletopterum) Bourguignat.....	649
Plicata (Mutela) Sowerby.....	905	Prælonga (Eurynea) Stimpson.....	545
Plicata (Mya) Eaton	767	Prælongus (Unio) Barnes	545
Plicata (Quadrula) Say	766, 767	Prasina (Mutelina) Rochebrune	906
Plicata (Unio) Say	504, 766, 767	Prasinatus (Unio) Conrad.....	714
Plicatula (Barbala) Patet.....	584	Prasinus (Unio) Conrad.....	779
Plicatulus (Margaron) Lea.....	853	Prattii (Lampsilis) Lea.....	550
Plicatulus (Nephronaias) Charpentier	597	Prattii (Margaron) Lea.....	550
Plicatulus (Unio) Charpentier	591, 597	Prattii (Unio) Lea.....	550
Plicatulus (Unio) Lea	853	Pressa (Alasmodonta) Baker.....	663
Plicatus (Appius) Gray.....	583	Pressodonta Simpson.....	667
Plicatus (Dipsas) Leach.....	583	Pressus (Margaron) Lea	663
Plicatus (Margarita) Lea.....	583, 767	Pressus (Unio) H. and A. Adams.....	663
Plicatus (Margaron) Lea	583, 767	Prevostianus (Lampsilis) Lea.....	559
Plicatus (Mycetopus) Clessin.....	930	Prevostianus (Margaron) Lea.....	550
Plicatus (Mycetopus) Sowerby	905	Prevostianus (Unio) Lea.....	550
Plicatus (Mytilus) Solander.....	583	Prevostianus (Unio) Sowerby	560
Plicatus (Platiris) Lea	905	Prisodon Lea	863
Plicatus (Plectomerus) Conrad.....	767	Prisodon Schumacher	501,
Plicatus (Unio) Kuster	769	506, 507, 515, 867, 868, 869, 870, 872	
Plicatus (Unio) Sowerby	583	Productus (Unio) Lea.....	820
Pliciferus (Margarita) Lea	701	Productus (Margarita) Lea.....	735
Pliciferus (Margaron) Lea.....	702	Productus (Margaron) Lea	735
Pliciferus (Unio) Lea.....	701	Productus (Unio) Conrad	735

	Page.		Page.
Prodictus (Unio) Mousson.....	820	Pulcher (Margarita) Lea.....	541
Prodictus (Unio) Bourguignat.....	683	Pulcher (Margaron) Lea.....	541
Profugus (Diplodon) Gould.....	891	Pulcher (Unio) Lea.....	541
Profugus (Margaron) Lea.....	891	Pulcher (Unio) Tapperone-Canefri.....	856
Profugus (Unio) Gould.....	891	Pullatus (Margaron) Lea.....	726
Prolongatus (Reticiens) Drouet.....	854	Pullatus (Unio) Lea.....	726
Prolongatus (Unio) Drouet.....	853, 854	Pullus (Lampsilis) Conrad.....	566
Propinqua (Truncella) Lea.....	523	Pullus (Margaron) Lea.....	566
Propinquus (Margaron) Lea.....	523	Pullus (Unio) Conrad.....	566
Propinquus (Unio) Lea.....	523	Pulvinata (Columba) Clessin.....	915
Proprius (Lampsilis) Lea.....	548	Pulvinata (Leila) Hupé.....	915
Proprius (Margaron) Lea.....	548	Pulvinatus (Anodon) Sowerby.....	915
Proprius (Unio) Lea.....	548	Pulvinulus (Margaron) Lea.....	758
Prostera Rafinesque.....	511, 566, 578	Pulvinulus (Unio) Lea.....	758
Prosacrus (Unio) Locard.....	689	Pumila (Anodonta) Heude.....	640
Protechi (Spathella) Rochebrune.....	902	Pumila (Quadrula) Lea.....	785
Protensus (Margaron) Lea.....	716	Pumilio (Unio) Kuster.....	817
Protensus (Unio) Lea.....	716	Pumilus (Margaron) Lea.....	785
Protensus (Unio) Tapperone-Canefri.....	857	Pumilus (Unio) Lea.....	785
Provancheriana (Arconata) Pilsbry.....	725	Pumilus (Unio) Ziegler.....	821
Proximus (Margaron) Lea.....	551	Punctatus (Lampsilis) Lea.....	542
Proximus (Unio) Kuster.....	556	Punctatus (Margaron) Lea.....	542
Proximus (Unio) Lea.....	550	Punctatus (Unio) Lea.....	542
Pruinosus (Unio) Schmidt.....	698	Puniceus (Lampsilis) Haldeman.....	548
Prunoides (Diplodon) Lea.....	875	Puniceus (Margaron) Lea.....	548
Prunoides (Margaron) Lea.....	875	Puniceus (Unio) Haldeman.....	548
Prunoides (Unio) Lea.....	875	Purpurascens (Anodon) Swainson.....	575
Prusli (Unio) Bourguignat.....	694	Purpurascens (Unio) Lamarek.....	721
Psammaetinus (Unio) Philippi.....	879	Purpuratus (Lampsilis) Lamarek.....	568, 569
Psammitta (Anodonta) Bourguignat.....	625	Purpuratus (Margarita) Lea.....	568
Psammoica (Castalina) d'Orbigny.....	866	Purpuratus (Margaron) Lea.....	568
Psammoica (Unio) d'Orbigny.....	866	Purpuratus (Unio) Lea.....	568
Psammoicus (Margaron) Lea.....	866	Purpurea (Anodonta) Valenciennes.....	832
Pseudavicula Simpson.....	506, 515, 860	Purpurea (Dallia) Valenciennes.....	832
Pseudobaphia Simpson.....	849	Purpurea (Margarita) Lea.....	832
Pseudodon Gould.....	506, 515, 834, 836	Purpurea (Margaron) Lea.....	832
Pseudodontopsis (Anodonta) Bourguignat.....	648	Purpurea (Mya) Eaton.....	721
Pseudodontopsis (Anodonta) Locard.....	649, 650	Purpurellus (Margaron) Lea.....	718
Pseudodontopsis (Gabiollia) Locard.....	650	Purpurellus (Unio) Lea.....	718
Pseudodus de Morgan.....	834	Purpureus (Unio) Say.....	721
Pseudomutela Simpson.....	905	Purpuratus (Unio) Say.....	592
Pseudonymus (Unio) Simpson.....	694	Purus (Margaron) Lea.....	729
Pseudon Simpson.....	601	Purus (Unio) Lea.....	729
Pseudospatha Simpson.....	505, 506, 514, 577	Pusilla (Obliquaria) Rafinesque.....	744
Psilunio.....	510	Pusillus (Margaron) Lea.....	708
Psoricus (Unio) Morelet.....	703	Pusillus (Unio) Lea.....	708
Pteranodon Fischer.....	637	Pustulata (Margaron) Lea.....	781
Pterocygnia Rafinesque.....	665	Pustulata (Quadrula) Lea.....	781
Ptychobrancheus Simpson.....	506, 510, 514, 612	Pustulata (Unio) Swainson.....	608
Ptychorhynchus Simpson.....	506, 515, 350	Pustulatus (Margarita) Lea.....	781
Ptychoderma Simpson.....	606	Pustulatus (Unio) Lea.....	781, 782
Ptychogenæ.....	514, 612	Pustulosa keineriana (Quadrula) Lea.....	780
Puberula (Anodonta) Gould.....	919	Pustulosa pernodosa (Quadrula) Lea.....	780
Pudica (Pleurobema) Lea.....	751, 760	Pustulosa (Quadrula) Lea.....	510, 779, 780
Pudicus (Margaron) Lea.....	751	Pustulosus (Margarita) Lea.....	779
Pudicus (Unio) Lea.....	751	Pustulosus (Margaron) Lea.....	779
Puelchana (Anodonta) d'Orbigny.....	921	Pustulosus (Unio) Call.....	782
Puelchana (Margaron) Lea.....	921	Pustulosus (Unio) Lea.....	779
Puelchanus (Glabaris) d'Orbigny.....	921	Pybasii (Margaron) Lea.....	549
Puerorum (Anodon) Heude.....	639	Pybasii (Unio) Lea.....	549
Pugio (Nodularia) Benson.....	814	Pygmæa (Mycetopoda) Spix.....	934
Pugio (Unio) Benson.....	814	Pygmæus (Margaron) Lea.....	715
Pulchella (Anodon) Heude.....	639	Pygmæus (Mycetopus) Hupé.....	935
Pulchella (Cameronia) Bourguignat.....	910	Pygmæus (Mycetopus) Sowerby.....	934
Pulcher (Lamellidens) Tapperone-Canefri.....	856	Pygmæus (Platiris) Lea.....	934
		Pygmæus (Unio) Lea.....	715

	Page.		Page.
Pyramidata (Quadrula) Lea.....	510, 754, 790	Rajahensis (Unio) Lea.....	844
Pyramidatus (Margarita) Lea.....	790	Raleighensis (Margaron) Lea.....	723
Pyramidatus (Margaron) Lea.....	790	Raleighensis (Unio) Lea.....	723
Pyramidatus (Unio) Lea.....	790	Ranarum (Anodonta) Morelet.....	624
Pyriformis (Margaron) Lea.....	762	Randabeli (Brazzea) Bourguignat.....	907
Pyriformis (Pleurobema) Lea.....	762	Randabeli (Cameronia) Bourguignat.....	910
Pyriformis (Unio) Lea.....	762	Randabeli (Nodularia) Bourguignat.....	822
Quadrangulosus (Unio) Heude.....	801	Randabeli (Unio) Bourguignat.....	822
Quadrans (Diplodon) Lea.....	885	Randalli (Anodonta) Trask.....	657, 658
Quadrans (Margaron) Lea.....	885	Rangianus (Margarita) Lea.....	523
Quadrans (Unio) Lea.....	885	Rangianus (Margaron) Lea.....	523
Quadrata (Castalia) Sowerby.....	864	Rangianus (Unio) Lea.....	523
Quadrata (Margaritana) Lea.....	664	Rariplicata (Unio) Deshayes.....	767
Quadrata (Margaron) Lea.....	664	Rarisulcata (Unio) Lamarck.....	721
Quadrata (Symphynta) Lea.....	664	Rasus (Margaron) Lea.....	687
Quadratus (Margaron) Lea.....	723	Rasus (Unio) Lea.....	687
Quadratus (Tetraplodon) Sowerby.....	864	Ratus (Unio) Lea.....	724
Quadratus (Unio) Lea.....	723	Raveneli (Unio) Conrad.....	721, 748
Quadratus (Unio) Reeve.....	777	Raveneliana (Alasmodonta) Lea.....	671
Quadrilatera (Castalia) d'Orbigny.....	864	Raveneliana (Alasmodonta) Ferussac.....	671
Quadrilaterum (Tetraplodon) d'Orbigny.....	864	Raveneliana (Baphia) H. and A. Adams.....	671
Quadrilaterus (Margaron) Lea.....	725	Raveneliana (Margaritana) Chenu.....	677
Quadrilaterus (Unio) Lea.....	725	Raveneliana (Margaritana) Lea.....	671
Quadruplicatus (Anodon) Sowerby.....	618	Raveneliana (Pleurobema) Lea.....	748
Quadrula (Obliquaria) Rafinesque.....	744	Ravenelianus (Margarita) Lea.....	671, 748
Quadrula (Rafinesque).....	505, 506, 508, 510, 514, 745, 765, 773, 793, 805	Ravenelianus (Margaron) Lea.....	671, 748
Quadrulus (Unio) Say.....	777	Ravenelianus (Strophitus) Conrad.....	671
Quinqueannulatus (Unio) Kuster.....	683	Ravenelianus (Unio) Lea.....	748, 758
Quintardii (Unio) Cragin.....	768	Ravenelianus (Unio) Reeve.....	751
Radians (Margaron) Lea.....	554	Ravistellus (Margaron) Lea.....	597
Radians (Unio) Lea.....	554	Ravistellus (Nephronalis) Morelet.....	597
Radiata (Alasmodonta) Conrad.....	620	Ravistellus (Unio) Morelet.....	597
Radiata (Anodonta) Valenciennes.....	648	Ravoisieri issericus (Unio) Kobelt.....	686
Radiata (Cristaria) Simpson.....	585, 586	Ravoisieri (Margaron) Lea.....	687
Radiata (Margarita) Lea.....	925	Ravoisieri (Unio) Deshayes.....	687
Radiata (Mya) Gmelin.....	535	Rayi (Anodonta) Dronet.....	624
Radiata (Unio) Lamarck.....	528, 535	Reclusa (Pleurobema) B. H. Wright.....	760
Radiata (Unio) Swainson.....	667	Reclusus (Unio) B. H. Wright.....	760
Radiatula Simpson.....	820	Recognitus (Mycetopus) Heude.....	656
Radiatus (Anodon) Spix.....	925	Recta (Hyria) Sowerby.....	871
Radiatus (Anodonta) Müller.....	622	Recta (Unio) Lamarck.....	534, 544
Radiatus conspiuus (Lampsilis) Lea.....	536	Rectidens Simpson.....	506, 515, 853
Radiatus (Glabaris) Spix.....	925	Rectilinearis (Unio) Sowerby.....	825
Radiatus (Lampsilis) Gmelin.....	534, 535, 536	Rectus (Lampsilis) Barnes.....	545
Radiatus (Margarita) Lea.....	535	Rectus (Lampsilis) Lamarck.....	544
Radiatus (Margaron) Lea.....	535, 925	Rectus (Margarita) Lea.....	545
Radiatus (Strophitus) Conrad.....	620	Rectus (Margaron) Lea.....	545
Radiatus (Unio) De Kay.....	552	Rectus (Prisodon) Sowerby.....	871
Radiatus (Unio) Hildreth.....	527	Rectus (Unio) Lamarck.....	509
Radiolus (Unio) Lea.....	716	Recurvatus (Unio) Lea.....	750
Radiosus (Unio) Lea.....	750	Recurvirostris (Anodonta) Lea.....	648
Radiota (Nodularia) Charmes.....	824	Reevianus (Lampsilis) Lea.....	538
Radiotus (Unio) Charmes.....	824	Reevianus (Margaron) Lea.....	538
Radula (Dysnomia) Rochebrune.....	820	Reevianus (Unio) Lea.....	538
Radula (Unio) Benson.....	820	Reflexa (Obliquaria) Rafinesque.....	610, 611
Radula (Unio) Say.....	744	Reflexus (Unio) Say.....	611
Radulosa (Quadrula) Dronet and Chaper.....	803	Refulgens (Margaron) Lea.....	782
Radulosa rhomboidea (Quadrula) Simpson.....	803	Refulgens (Quadrula) Lea.....	782
Radulosus (Unio) Dronet and Chaper.....	803	Refulgens (Unio) Lea.....	782
Racensis (Margaron) Lea.....	723	Regularis (Anodonta) Morelet.....	624
Racensis (Unio) Lea.....	723	Regularis (Margaron) Lea.....	556
Rajahensis (Margaron) Lea.....	844	Regularis (Unio) Lea.....	555
Rajahensis (Parreysia) Lea.....	844	Regularis (Unio) Sowerby.....	531
		Reiniana (Anodonta) von Thering.....	585
		Reiniana (Barbata) Pætel.....	585
		Reiniana (Cristaria) von Martens.....	585

	Page.		Page.
Reiniana (Dipsas) Kobelt	585	Rivularis (Anodonta) Heude	640
Reiniana (Nodularia) Kobelt	570	Rivularis (Mycetopus) Heude	657
Reinianus (Unio) Kobelt	510	Rivularis (Solcnaia) Heude	657
Renea (Nodularia) Jousseaume	822	Rivularis (Unio) Conrad	742
Renei (Spatha) Jousseaume	896	Rivularis (Unio) Rafinesque	744
Reueus (Reneus) Jousseaume	822	Roanokensis (Margarita) Lea	728
Reniformis (Unio) Rossmässler	698	Roanokensis (Margaron) Lea	728
Requeni (Unio) Michaud	686	Roanokensis northamptonensis (Unio)	
Requeni (Unio) Rossmässler	699	Lea	728
Retiaria (Parreysia) Heude	850	Roanokensis (Unio) Lea	728
Retiarius (Unio) Heude	850	Rochebruneana (Monectia) Bourguignat	901
Reticulata (Margaritana) Patet	912	Rochebrunei (Spatha) Jousseaume	897
Reticulata (Margaron) Lea	912, 919	Rochechouarti (Quadrula) Heude	799, 800
Reticulata (Monocondylaea) Moricand	912	Rochechouarti (Unio) Heude	800
Reticulatus (Anodon) Sowerby	919	Roissyi (Margaritana) Westerlund	676
Reticulatus (Aplodon) von Ihering	912	Roissyi (Unio) Michaud	676
Reticulatus (Nephronaias) Simpson	595	Rollei (Unio) Kobelt	693
Reticulatus (Unio) Sowerby	912	Rosaceus (Unio) Conrad	531
Retortus (Unio) Heude	798	Rosaceus (Unio) De Kay	535
Retortus (Unio) von Martens	805	Rosanorhamphus	515, 806
Retteri (Anodonta) Drouet	627	Rosea (Anodon) Heude	639
Retusa (Anodon) Heude	639	Rossmässleriana (Anodonta) Dupuy	624
Retusa (Castalia) Hupe	864	Rostellum (Unio) Lea	737
Retusa (Obliquaria) Rafinesque	603	Rostreiformis (Unio) Lea	736
Retusa (Obovaria) Lamarck	599	Rostralis (Unio) von Martens	829
Retusa (Unio) Lamarck	599, 775	Rostrata (Anodonta) Held	627
Retusus (Margarita) Lea	599	Rostrata (Anodonta) Rossmässler	627
Retusus (Margaron) Lea	599	Rostrata (Grandieria) Bourguignat	829
Retusus (Tetraplodon) Hupé	864	Rostrata (Iridina) Rang	905
Retusus (Unio) Say	775	Rostrata (Mutela) Rang	905
Revoiliana (Cameronia) Bourguignat	910	Rostrata (Mutelina) Bourguignat	905
Rhombeus (Diplodon) Wagner	873	Rostrata (Spatha) von Martens	905
Rhombeus (Margarita) Lea	873	Rostrata (Unio) Lamarck	682
Rhombeus (Margaron) Lea	873	Rostrata (Unio) Valenciennes	546
Rhombica (Alasmodon) Anthony	617	Rostratus (Anodon) Sowerby	629
Rhombica (Margaritana) Patet	617	Rostriformis (Margaron) Lea	736
Rhombica (Monocondylaea) Kuster	836	Rostriformis (Unio) Lea	736
Rhomboida (Leguminaia) Kobelt	650	Rostrum (Margaron) Lea	726
Rhomboida (Margaron) Lea	650	Rostrum (Unio) Lea	726
Rhomboida (Microcondylaea) Patet	650	Roswellensis (Margaron) Lea	723
Rhomboida (Monocondylaea) Lea	650	Roswellensis (Unio) Lea	723
Rhomboides (Pseudodon) Westerlund	650	Rothi (Unio) Bourguignat	692, 693
Rhomboides (Unio) Moquin-Tandon	692	Rotulata (Obovaria) B. H. Wright	601
Rhomboides (Unio) Sowerby	650	Rotulatus (Unio) B. H. Wright	601
Rhomboidopsis (Unio) Locard	693	Rotunda (Margaron) Lea	918
Rhombula (Unio) Lamarck	721	Rotunda (Mya) Wood	600
Rhuacoica (Unio) d'Orbigny	876	Rotundaria Agassiz	765
Rhuacoicus (Diplodon) d'Orbigny	876	Rotundaria Rafinesque	794
Rhuacoicus (Margaron) Lea	876	Rotundata (Anodonta) Trask	629
Rhuaconicus (Diplodon) Kuster	882	Rotundata (Glebulula) Lamarck	598
Rhuaconicus (Unio) Kuster	882	Rotundata (Grandieria) Bourguignat	830
Rhyacæa (Monocondylaea) Patet	876	Rotundata (Ihringella) Mousson	913
Rhynchoidea (Chambardia)	902	Rotundata (Spatha) von Martens	896
Rhynchonella (Chambardia)	902	Rotundata (Unio) Lamarck	597, 598
Rhynchonella (Grandieria) Bourguignat		Rotundata (Unio) Manduyt	698
nat	829	Rotundatus (Anodon) Swainson	638
Richthofeni (Unio) von Martens	801	Rotundatus (Margarita) Lea	598
Riddellii (Margaron) Lea	787	Rotundatus (Margaron) Lea	598
Riddellii (Quadrula) Lea	787	Rotundatus (Plagiodon) Mousson	913
Riddellii (Unio) Lea	787	Rotundatus (Unio) Lamarck	541
Ridibundus (Unio) Say	520	Rotundus (Anodon) Spix	918
Rigida (Mya) Wood	722	Rotundus (Diplodon) Wagner	886
Rimosus (Unio) Rafinesque	744	Rotundus (Glabaris) Spix	918
Rioplatensis (Anodon) Sowerby	925	Rotundus (Margaron) Lea	886
Riparia (Unio) C. Pfeiffer	697	Rotundus (Unio) Wagner	886
Rivicolus (Unio) Conrad	742	Rousii (Unio) Dupuy	686

	Page.		Page.
Rovirosal (Lampsilis) Pilsbry	568	Rugosissima (Hyria) Sowerby	869
Rovirosal (Unio) Pilsbry	568	Rugosulcata (Quadrula) Lea	794
Rowellii (Margaron) Lea	594	Rugosulcatus (Margaron) Lea	794
Rowellii (Nephronaias) Lea	594	Rugosulcatus (Unio) Lea	794
Rowellii (Unio) Lea	594	Rugosum (Lasmigona) Rafinesque	665
Rubella (Anodon) Heude	639	Rugosum (Triplodon) Spix	869
Rubella (Pleurobema) Conrad	757, 758	Rugosus (Anodon) Swainson	616
Rubellinus (Margaron) Lea	590	Rugosus (Unio) Chenu	773
Rubellinus (Unio) Lea	590	Rugosus (Unio) Hanley and Theobald	813
Rubellus (Margarita) Lea	758	Rugosus (Unio) Kuster	844
Rubellus (Margaron) Lea	758	Rugulosa (Mya) Wood	671
Rubellus (Unio) Conrad	757	Rugulosus (Nephronaias) Charpentier	596
Rubens (Anodonta) Lamarek	895, 896	Rugulosus (Unio) Charpentier	596
Rubens cailliaudi (Spatha) von Martens	896	Rumphianus (Margaron) Lea	778
Rubens (Iridina) Rang	896	Rumphianus (Quadrula) Lea	778
Rubens (Platiris) Lea	896	Rumphianus (Unio) Lea	778
Rubens (Spatha) Lamarek	895, 896	Rustica (Nodularia) Lea	819
Rubens (Unio) Menke	691, 789	Rusticus (Margaron) Lea	819
Ruber (Unio) Conrad	790	Rusticus (Unio) Lea	819
Rubicunda (Anodonta) Lea	918	Rusticus (Unio) Sowerby	702
Rubicunda (Margaron) Lea	918	Rutersvillensis (Margaron) Lea	547
Rubicundus (Glabaris) Lea	918	Rutersvillensis (Unio) Lea	547
Rubida (Quadrula) Lea	787	Rutilans (Margaron) Lea	560
Rubidus (Margaron) Lea	787	Rutilans (Unio) Lea	560
Rubidus (Unio) Lea	787	Saccatus (Unio) Kuster	524
Rubiginosa (Quadrula) Lea	510, 786	Sacculus (Unio) Reeve	524
Rubiginosus (Margarita) Lea	786	Sacellus (Nodularia) Drouet and Chaper	818
Rubiginosus (Margaron) Lea	786	Sacellus (Unio) Drouet and Chaper	818
Rubiginosus (Unio) Lea	786	Sageri (Unio) Conrad	545
Rubra (Amblema) Rafinesque	744	Sagittaria (Nodularia) Lea	810, 811
Rudis (Unio) Conrad	748, 758	Sagittarius (Margaron) Lea	811
Rudis (Unio) Pætel	875	Sagittarius (Unio) Lea	811
Rudus (Diplodon) Lea	875	Sagittiformis (Margaron) Lea	731
Rudus (Margaron) Lea	875	Sagittiformis (Unio) Lea	731
Rudus (Unio) Conrad	758	Sagrinateus (Unio) Sowerby	607
Rudus (Unio) Lea	875	Saladoensis (Lampsilis) Lea	569
Ruellani (Parreysia) Bourguignat	846	Saladoensis (Margaron) Lea	569
Ruellani (Unio) Bourguignat	846	Saladoensis (Unio) Lea	569
Rufescens (Cuneopsis) Heude	805	Salebrosus (Margaron) Lea	726
Rufescens (Unio) Heude	805	Salebrosus (Unio) Lea	726
Rufofuscus (Diplodon) Lea	880	Salmonia (Anodonta) Clessin	617
Rufofuscus (Margaron) Lea	722, 880	Salmonia (Anodonta) Lea	642
Rufofuscus (Unio) Lea	880	Salmonia (Margarita) Lea	642
Rufus (Margaron) Lea	704	Salmonia (Margaron) Lea	642
Rufus (Unio) Lea	704	Salweniana (Anodon) Gould	836, 838
Rufusculus (Unio) Lea	722	Salweniana (Margaron) Lea	838
Rugata (Solenais) Heude	657	Salwenianus (Pseudodon) Gould	838
Rugatus (Mycetopus) Heude	657	Salwenianus (Unio) Sowerby	838
Rugatus (Platiris) Lea	657	Sampsonii (Truncilla) Lea	523
Rugatus (Unio) Hutton	889	Sampsonii (Unio) Lea	523
Rugatus (Unio) Menke	697	Sandri (Unio) Rossmassler	699
Rugifer (Unio) Kuster	821	Santeensis (Unio) Lea	717
Rugifera (Anodonta) Dunker	648, 902	Sapotalensis (Margaron) Lea	593
Rugifera Simpson	670	Sapotalensis (Nephronaias) Lea	592
Rugosa (Alasmodonta) Barnes	664, 665	Sapotalensis (Unio) Lea	592
Rugosa (Baphia) H. and A. Adams	665	Satillaensis (Margaron) Lea	733
Rugosa (Complanaria) Stimpson	665	Satillaensis (Unio) Lea	733
Rugosa (Hyria) Deshayes	869	Satur (Margaron) Lea	528
Rugosa (Margarita) Lea	665	Satur (Unio) Lea	527
Rugosa (Margaritana) Calkins	665	Sauleyi (Leguminaia) Bourguignat	652
Rugosa (Margaron) Lea	665	Sauleyi (Margaron) Lea	652
Rugosa (Mya) Eaton	665	Sauleyi (Microcondylea) Clessin	652
Rugosa (Mya) Gmelin	844	Sauleyi (Pseudodon) Conrad	652
Rugosa (Parreysia) Gmelin	844	Sauleyi (Unio) Bourguignat	652
Rugosa (Unio) Hanley	665	Savannahensis (Margaron) Lea	722
Rugosa (Unio) Poiret	678	Savannahensis (Unio) Lea	722

	Page.		Page.
Savoyensis (Unio) Pætel	843	Securis (Margarita) Lea	604
Saxeus (Margaron) Lea	547	Securis (Margaron) Lea	604
Saxeus (Unio) Conrad	547	Securis (Plagiola) Lea	603
Sayatus (Unio) Conrad	739, 741	Securis (Unio) Lea	603
Sayi (Unio) Ward	739, 741	Sedakowi (Anodonta) Siemaschko	648
Scaldiana (Anodonta) Dupuy	624	Semialatus (Physunio) Deshayes and Jullien	831
Scalcnaria (Rafinesque)	519	Semialatus (Unio) Deshayes and Jullien	831
Scalenius (Unio) Say	752	Semidecorata (Nodularia) L. Morlet	819
Seamnatus (Margaron) Lea	595	Semidecoratus (Unio) L. Morlet	819
Seamnatus (Nephronaias) Morelet	595	Semigranosis (Margaron) Lea	702
Seamnatus (Unio) Morelet	595	Semigranosis (Unio) Lea	571
Seaphydium (Anodon) Heude	639	Semigranosis (Unio) Reeve	702
Schæfferiana (Strophitus) Conrad	617	Semigranosis (Unio) von dem Buseh	701, 702
Schlaflii (Anodonta) Mousson	648	Semplicatus (Unio) Kuster	590
Schlegeli (Barbala) Pætel	531	Semiquadrata (Unio) Sowerby	831
Schlegeli (Hyriopsis) von Martens	581	Semirugatus (Unio) Chenu	842
Schlegeli (Unio) von Martens	581	Semirugatus (Unio) Lamarek	693
Schæfferiana (Anodon) Sowerby	617	Semisulcata (Iheringella) H. Adams	913
Schombergiana (Castalia) Sowerby	864	Semisulcata (Monocondylæa) H. Adams	913
Schomburgianus (Anodon) Sowerby	920	Semmelfinki (Unio) von Martens	834
Schomburgianus (Glabaris) Sowerby	920	Sempervivens (Anodonta) Deshayes	588
Schomburgianus (Ptychorhynchus) Heude	851	Senegalensis (Anodonta) Lea	900
Schomburgianus (Tetraplodon) Sowerby	864	Senegalensis (Margaron) Lea	900
Schomburgianus (Unio) Heude	851	Senegalensis (Spatha) Pætel	900
Schomburgki (Anodonta) von Martens	588	Senegalia (Mucilina) Jousseume	900
Schomburgki (Pilsbryconcha) von Mar- tens	588	Sennaaricensis (Unio) Kuster	821
Schoolcraftensis (Margarita) Lea	780	Sennariensis (Unio) von Martens	813
Schoolcraftensis (Margaron) Lea	780	Serpentina (Anodonta) Clessin	917
Schoolcraftensis (Unio) Lea	780	Serradelli (Dallielia) Drouet	834
Schoolcraftii (Margaron) Lea	780	Serradelli (Unio) Drouet	134
Schoolcraftii (Unio) B. H. Wright	780	Servainiana (Grandidieria) Bourguignat	628
Schrenkei (Unio) Westerlund	808	Shæfferiana (Anodonta) Lea	617
Schröteriana (Anodonta) Lea	931	Shæfferiana (Margaron) Lea	617
Schröteriana (Margaritana) Pætel	931	Shafferiana (Anodonta) B. H. Wright	617
Schröteriana (Margaron) Lea	931	Shanghaiensis (Unio) Lea	808
Schröterianus (Glabaris) Lea	931	Shefferiana (Anodonta) Clessin	617
Schweinfurthi (Unio) von Martens	862	Shepardianus (Margarita) Lea	737
Schwerzenbachi (Unio) Bourguignat	694	Shepardianus (Margaron) Lea	737
Scitulus (Margaron) Lea	554	Shepardianus (Unio) Lea	737
Scitulus (Unio) Lea	554	Sheppardianus (Unio) Pætel	737
Scobina (Nodularia) Hanley	816	Shistodesma Simpson	506, 514, 803, 804
Scobina (Unio) Hanley	816	Showalterii (Anodonta) Lea	617
Scobinata (Nodularia) Lea	815	Showalterii (Margaron) Lea	617, 757
Scobinatus (Margaron) Lea	815	Showalterii (Pleurobema) Lea	756, 757
Scobinatus (Unio) Lea	815	Showalterii (Unio) Lea	757
Scripta (Quadrula) Heude	802	Shurtleffiana (Nodularia) Lea	813
Scripto-polystictus (Unio) Heude	799	Shurtleffianus (Margaron) Lea	813
Scripto-polystictus (Unio) Pætel	799	Shurtleffianus (Unio) Lea	813
Scriptum (Alasmodon) Rafinesque	672	Shuttleworthii (Diplodon) Lea	893
Scriptus (Anodon) Sowerby	921	Shuttleworthii (Margaron) Lea	893
Scriptus (Unio) Heude	802	Shuttleworthii (Unio) Kuster	891
Seutum (Lamellidens) Sowerby	857	Shuttleworthii (Unio) Lea	893
Seutum (Unio) Sowerby	857	Siamensis (Margaron) Lea	859
Sculptilis (Alasmodonta) Say	667	Siamensis (Unio) Lea	859
Sculptilis (Strophitus) Stimpson	667	Siamensis (Unio) Morelet	819
Seulptus (Unio) Deshayes	808	Sicula (Potamida) Swainson	683, 686
Seutulatus (Lampsilis) Morelet	576	Siculus (Unio) Hanley	683
Seutulatus (Unio) Morelet	576	Siculus (Unio) Sowerby	686
Seyphus (Unio) Kuster	761	Sikkimensis (Margaron) Lea	845
Sebastianæ (Unio) von Ihering	894	Sikkimensis (Parreysia) Lea	845
Secundus (Pseudodon) Heude	840	Sikkimensis (Unio) Lea	845
Secundus (Unio) Pætel	841	Siliqua (Anodonta) Kuster	588
Securiformis (Anodon) Heude	638	Siliquoidea (Lampsilis) Stimpson	534
Securiformis (Unio) Conrad	781	Siliquoidea (Mya) Eaton	534
		Siliquoides (Unio) Kuster	534

Page.		Page.
Silicoideus (Unio) Barnes	534	Soleniformis (Anodonta) d'Orbigny..... 930
Siliquosa (Mycetopoda) Spix.....	934	Soleniformis (Glabaris) d'Orbigny..... 930
Siliquosus (Anodon) Spix.....	934	Soleniformis (Margarita) Lea..... 930
Siliquosus (Mycetopus) d'Orbigny.....	934	Soleniformis (Margaritana) Patet..... 679
Siliquosus (Platiris) Lea.....	934	Soleniformis (Margaron) Lea..... 930, 935
Siliquosus var. b. pygmaeus (Anodon) Spix.....	934	Soleniformis (Mutela) Bourguignat..... 903
Simplex (Margaron) Lea.....	614	Soleniformis (Mycetopoda) d'Orbigny... 935
Simplex (Unio) Lea.....	614	Soleniformis (Mycetopus) Fischer..... 656
Similaris (Quadrula) Simpson.....	799	Soleniformis (Mycetopus) Reeve..... 935
Similis (Margaron) Lea.....	716	Soleniformis (Platiris) Lea..... 935
Similis (Mycetopus) Heude.....	657	Soleniformis (Solenaia) Benson..... 656
Similis (Unio) Lea.....	716	Soleniformis (Spatha) Hanley and Theo- bald..... 656
Simonis (Unio) Tristram.....	693	Soleniformis (Unio) Lea..... 679
Simplicidus (Anodon) Sowerby.....	625	Solida (Anodonta) Kuster..... 917
Simpsoni (Mutela) Ancey.....	904	Solida (Iridina) Kuster..... 896
Simpsoni (Unio) B. H. Wright.....	714	Solida (Mysea) Turton..... 685
Simpsoniania (Anodonta) Lea.....	647	Solida (Quadrula) Lea..... 757, 765, 789
Simpsoniania (Margaron) Lea.....	647	Solidula (Anodonta) Deville and Hupe... 928
Simpsonianus (Glabaris) Pilsbry.....	925	Solidula (Margaron) Lea..... 928
Simularis (Parresia) Heude.....	845	Solidulus (Diplodon) Philippi..... 881
Simularis (Unio) Heude.....	845	Solidulus (Unio) Philippi..... 881
Simulans (Pleurobema) Lea.....	760	Solidus (Margarita) Lea..... 789
Simulans (Unio) Lea.....	760	Solidus (Margaron) Lea..... 789
Simus (Lampsilis) Lea.....	556	Solidus (Unio) Lea..... 789
Simus (Margarita) Lea.....	556	Solisiana (Unio) d'Orbigny..... 887
Simus (Margaron) Lea.....	556	Solisianus (Diplodon) d'Orbigny..... 887
Simus (Unio) Lea.....	556	Somboriensis (Harmandia) Rochebrune... 827
Sinensis squamosus (Unio) Heude.....	850	Somersi (Anodonta) Call..... 648
Singleyanus (Lampsilis) Marsh.....	563	Sondermannii (Anodonta) Kuster..... 625
Singleyanus (Unio) Marsh.....	563	Sordida (Zaira) Rochebrune..... 862
Singularis (Grandideria) Bourguignat.....	829	Sordidus (Margaron) Lea..... 732
Sintoxia (Obliquaria) Rafinesque.....	744	Sordidus (Unio) Lea..... 732
Sinuata (Obliquaria) Rafinesque.....	744	Sorini (Anodon) Heude..... 639
Sinuata (Spatha) von Martens.....	899	Sowerbyanus (Margarita) Lea..... 521
Sinuata (Unio) C. Pfeiffer.....	676	Sowerbyanus (Margaron) Lea..... 521
Sinuata (Unio) Lamarck.....	678	Sowerbyanus (Unio) Lea..... 520
Sinuosa (Anodonta) Clessin.....	917	Spadiceus (Margaron) Lea..... 732
Sinuosa (Anodonta) Lamarck.....	925	Spadiceus (Unio) Lea..... 732
Sinuosa (Margarita) Lea.....	925	Sparsa (Quadrula) Lea..... 775
Sinuosa (Margaron) Lea.....	925	Sparsus (Margaron) Lea..... 775
Sinuosus (Glabaris) Lamarck.....	925	Sparsus (Unio) Kuster..... 784
Sirionis (Anodonta) d'Orbigny.....	918	Sparsus (Unio) Lea..... 775
Sirionis (Glabaris) d'Orbigny.....	918	Sparus (Margaron) Lea..... 555
Sirionis (Margarita) Lea.....	918	Sparus (Unio) Lea..... 555
Sirionis (Margaron) Lea.....	918	Spatha Lea..... 506, 515, 895, 901
Sitifensis (Unio) Morelet.....	694, 862	Spathopsis Simpson..... 857
Sloatianus (Margaron) Lea.....	706	Spathuliformis (Spathella) Bourguignat... 898
Sloatianus (Plectomerus) Conrad.....	706	Spatiosa (Anodonta) Clessin..... 584
Sloatianus (Unio) Lea.....	705, 706	Spatiosa (Barbala) Patet..... 585
Smaragdina (Craspedodonta) Anton.....	584	Spatiosa (Cristaria) Clessin..... 584, 585
Smaragdites (Unio) Benson.....	843	Spatulatus (Lampsilis) Baker..... 557
Smithi (Grandideria) Bourguignat.....	829	Spatulatus (Margaron) Lea..... 557
Smithii (Margarita) Lea.....	880	Spatulatus (Unio) Lea..... 557
Smithii (Margaron) Lea.....	880	Spatulatus (Unio) Sowerby..... 556
Smithii (Unio) Gray and Pidgeon.....	880	Speciosa (Quadrula) Lea..... 778
Sobolus (Nodularia) Fischer.....	819	Speciosus (Margaron) Lea..... 778
Sobolus (Unio) Fischer.....	819	Speciosus (Unio) Lea..... 778
Soleileti (Unio) Bourguignat.....	862	Spekii (Cameronia) Bourguignat..... 909
Solenaia Conrad.....	506, 514, 655, 933	Spekii (Iridina) Woodward..... 909
Solenidea (Anodon) Sowerby.....	930	Spekii (Mutela) von Martens..... 909
Solenidea (Margaron) Lea.....	930	Spekii (Platiris) Lea..... 909
Solenoides cylindrica (Unio) Rafinesque.....	773	Spekii (Pleiodon) Woodward..... 909
Solenoides (Loncosilla) Rafinesque.....	862	Sphaerica (Quadrula) Lea..... 781
Solenoides (Mycetopus) Sowerby.....	935	Sphaericus (Margaron) Lea..... 781
Soleniformis (Anodonta) Benson.....	656	Sphaericus (Unio) Lea..... 781

	Page.		Page.
Spheniopsis (Quadrula) Morelet.....	794	Stolli (Unio) von Martens.....	771
Spheniopsis (Unio) Morelet.....	794	Stonensis (Margaron) Lea.....	705
Sphenorhynchus (Unio) Fischer and Crosse.....	712	Stonensis (Unio) Lea.....	705
Spillmanii (Margaritana) Lea.....	533, 619	Stramineus (Lampsilis) Conrad.....	538
Spillmanii (Margaron) Lea.....	532	Stramineus (Margarita) Lea.....	538
Spillmanii (Strophitus) Lea.....	619	Stramineus (Unio) Conrad.....	538
Spillmanii (Unio) Lea.....	532	Strebelii (Anodonta) Lea.....	920
Spillmanii (Unio) Sowerby.....	619	Strebelii (Glabaris) Lea.....	920
Spinelli (Unio) Villa.....	699	Strebelii (Margaron) Lea.....	592, 920
Spinosa (Canthyrta) Swainson.....	739	Strebelii (Unio) Lea.....	592
Spinus (Margarita) Lea.....	739	Striata (Anodon) Heude.....	638
Spinus (Margaron) Lea.....	739	Striata (Iridina) Oken.....	903
Spinus (Shistodesmus) Simpson.....	804	Striata (Obovaria) Rafinesque.....	603
Spinus (Unio) Lea.....	738	Striata (Pleurobema) Lea.....	762
Spissus (Margaron) Lea.....	734	Striatissimus (Unio) Anthony.....	764
Spissus (Unio) Lea.....	734	Striatulus (Margaron) Lea.....	719
Spixii (Columba) von Ihering.....	916	Striatulus (Unio) Lea.....	719
Spixii (Leila) von Ihering.....	916	Striatus (Margaron) Lea.....	762
Spixii (Unio) d'Orbigny.....	932	Striatus (Unio) Lea.....	762
Spixoeoncha Pilsbry.....	910	Striatus (Unio) Reeve.....	781
Splendens (Anodonta) de Cristofori.....	896	Striatus (Unio) Sowerby.....	548, 670
Splendens (Pleiodon) Conrad.....	908	Strigillatus cylindrica (Unio) B. H. Wright.....	773
Splendida (Iridina) Chenu.....	908	Strigosus (Margaron) Lea.....	732
Splendidus (Lampsilis) Lea.....	531	Strigosus (Unio) Lea.....	732
Splendidus (Margarita) Lea.....	531	Strodeana (Pleurobema) B. H. Wright.....	761
Splendidus (Margaron) Lea.....	531	Strodeanus (Unio) B. H. Wright.....	761
Splendidus (Unio) Lea.....	531	Strophitus Rafinesque.....	506, 514, 616 658
Spuria (Mya) Gmelin.....	841	Strumosus (Unio) Lea.....	729
Spuria (Mya) Wood.....	842	Stuarti (Anodon) Sowerby.....	892
Spurius (Unio) Heude.....	798	Stuarti (Margaritana) Pætel.....	892
Squalidus (Margaron) Lea.....	725, 742	Stuarti (Unio) Sowerby.....	893
Squalidus (Unio) Lea.....	725, 742	Stuhlmanni (Parreysia) von Martens.....	816
Squameus (Margaron) Lea.....	723	Stuhlmanni (Spatha) von Martens.....	900
Squameus (Unio) Lea.....	723	Stuhlmanni (Unio) von Martens.....	846
Squammosus lævis (Unio) Heude.....	350	Stygianodon von Martens.....	929
Stabilis (Margaron) Lea.....	755	Suavidicus (Diplodon) Lea.....	876
Stabilis (Pleurobema) Lea.....	755	Suavidicus (Margaron) Lea.....	876
Stabilis (Unio) Lea.....	755	Suavidicus (Unio) Lea.....	876
Stagnalis (Anodonta) Bosc.....	622	Subæquilatera (Spatha) von Martens.....	899
Stagnalis (Margaron) Lea.....	560	Subamygdalinus (Unio) Drouet.....	862
Stagnalis (Mytilus) Turton.....	622	Subangulata (Anodon) Anthony.....	644
Stagnalis (Unio) Conrad.....	560	Subangulatus (Lampsilis) Lea.....	556
Stagnorum (Diplodon) Dautzenberg.....	894	Subangulatus (Margaron) Lea.....	556
Stagnorum (Unio) Dautzenberg.....	894	Subangulatus (Unio) Lea.....	556
Stapes (Margarita) Lea.....	775	Subangulatus (Unio) Sowerby.....	560
Stapes (Margaron) Lea.....	775	Subearinata (Anodonta) Currier.....	633
Stapes (Quadrula) Lea.....	775	Subcircularis (Anodonta) Clessin.....	628
Stapes (Unio) Lea.....	775	Subcrassa (Dallia) Lea.....	833
Staudingeri (Mycetopoda) von Ihering.....	934	Subcrassa (Margaron) Lea.....	833
Staudingeri (Mycetopus) von Ihering.....	934	Suberoceus (Unio) Conrad.....	740
Stearnsii (Unio) B. H. Wright.....	562	Subcylindracea (Anodon) Sowerby.....	633 660
Stegaria (Obovaria) Rafinesque.....	602	Subcylindracea (Anodonta) Lea.....	660
Stegarius irroratus (Unio) Pætel.....	610	Subcylindracea (Margarita) Lea.....	660
Stegarius (Unio) Kuster.....	775	Subcylindracea (Margaron) Lea.....	660
Stegarius (Unio) Say.....	610	Subcylindraceus (Anodontoides) Baker.....	660
Stevenianus (Unio) Drouet.....	698	Subcylindraceus (Unio) Lea.....	731
Stevensii (Hyria) Lea.....	869	Subdiaphana (Mutela) Bourguignat.....	906
Stevensii (Unio) Lea.....	869	Subellipsis (Margaron) Lea.....	560
Stewardsoni (Margaron) Lea.....	521	Subellipsis (Unio) Lea.....	560
Stewardsoni (Truncilla) Lea.....	521	Subflavus (Margaron) Lea.....	722
Stewardsoni (Unio) Lea.....	521	Subflavus (Unio) Lea.....	722
Stewartiana (Anodonta) Lea.....	645	Subgibbosa (Anodon) Anthony.....	644
Stewartiana (Margarita) Lea.....	645	Subgibbosus (Margaron) Lea.....	704
Stewartiana (Margaron) Lea.....	645	Subgibbosus (Unio) Lea.....	704, 707
Stolli (Quadrula) von Martens.....	771	Subglobata (Pleurobema) Lea.....	751

	Page.		Page.
Subglobatus (Unio) Lea.....	751	Subviridis (Triquetra) Klein.....	870
Subglobosa (Margaron) Lea.....	644	Succinea (Anodon) Heude.....	638
Subglobosus Unio) Lea.....	598	Succineus (Mycetopus) Heude.....	656
Subinflata (Anodonta) B. H. Wright.....	643	Succissa (Quadrula) Lea.....	785
Subinflatus (Margaron) Lea.....	736	Succissus (Margaron) Lea.....	785
Subinflatus (Unio) Conrad.....	736	Succissus (Unio) Lea.....	785
Sublatus (Margaron) Lea.....	726	Sudus (Lampsilis) Lea.....	561
Sublatus (Unio) Lea.....	726	Sudus (Margaron) Lea.....	561
Subluridus (Unio) Simpson.....	743	Sudus (Unio) Lea.....	561
Subluxata (Anodonta) Kuster.....	625	Sula (Unio) Theobald.....	678
Subniger (Margaron) Lea.....	730	Suleata (Anodonta) Kuster.....	642
Subniger (Unio) Lea.....	730	Suleata (Anodonta) Lamarek.....	622
Subolivaceus (Unio) Lea.....	724	Suleata (Castalia) Krauss.....	867
Suborbiculata (Anodonta) Say.....	636	Suleata (Castaliella) Krauss.....	867
Suborbiculata (Margarita) Lea.....	636	Suleata (Truncella) Lea.....	520
Suborbiculata (Margaron) Lea.....	636	Suleata var. deliata (Truncella) Simpson.....	520
Suborbiculata (Unio) Lamarek.....	598	Suleataria Rafinesque.....	673
Suborbiculus Simpson.....	835	Suleatum (Pseudodon) Roehebrune.....	840
Subovatus (Margarita) Lea.....	527	Suleatus (Margarita) Lea.....	520
Subovatus (Margaron) Lea.....	527	Suleatus (Unio) Lea.....	519, 520
Subovatus (Unio) Lea.....	527	Suleidens (Unio) Lamarek.....	721
Subparallelus (Unio) Lea.....	724	Suleosa (Anodonta) H. and A. Adams.....	648
Subplanus (Margaron) Lea.....	720	Sumatrensis (Unio) Lea.....	818
Subplanus (Unio) Conrad.....	720	Sumatrensis (Rectidens) Dunker.....	853
Subponderosa (Anodonta) Dupuy.....	624	Sumatrensis (Unio) Dunker.....	853
Subreniformis (Anodon) Sowerby.....	900	Sumatrensis (Unio) Sowerby.....	859
Subreniformis (Spatha) Sowerby.....	900	Superbus (Margaron) Lea.....	830
Subreniformis (Unio) Bourguignat.....	692	Superbus (Physunio) Lea.....	830
Subrostrata (Anodonta) Philippi.....	648, 928	Superbus (Unio) Sowerby.....	830
Subrostratus (Lampsilis) Say.....	546	Superiorensis (Lampsilis) Marsh.....	535
Subrostratus (Unio) Kuster.....	552	Superiorensis (Unio) Marsh.....	535
Subrostratus (Unio) Say.....	546	Suppositus (Unio) von Ihering.....	877
Subrotunda (Obliquaria) Rafinesque.....	744	Susanna (Anodon) Gray.....	924
Subrotunda (Quadrula) Lea.....	790, 791	Sutrangensis (Hyriopsis) Morelet.....	530
Subrotundus circulus (Unio) Pætel.....	600	Sutrangensis (Unio) Morelet.....	580
Subrotundus (Margarita) Lea.....	792	Swainsoni (Unio) Sowerby.....	667
Subrotundus (Margaron) Lea.....	792	Swainsoni (Margaron) Lea.....	667
Subrotundus (Unio) Ferrussae.....	600	Swananoensis (Unio) Hanley.....	671
Subrotundus (Unio) Lea.....	791	Swinhoei (Anodonta) H. Adams.....	586
Subrotundus (Unio) Sowerby.....	751	Swinhoei (Cristaria) H. Adams.....	586
Subsinuata (Anodonta) Philippi.....	648, 919	Swinhoei (Margaron) Lea.....	586
Subsinuata (Mycetopoda) Sowerby.....	934	Swinhoei (Unio) H. Adams.....	586
Subsinuatus (Anodon) Sowerby.....	924, 925	Swordiana (Pleurobema) S. H. Wright.....	750
Subsinuatus (Mycetopus) Sowerby.....	934	Swordianus (Unio) S. H. Wright.....	750
Subsquamosus (Unio) Lea.....	724	Symmetrieus (Margaron) Lea.....	740
Substriatus (Margaron) Lea.....	811	Symmetrieus (Unio) Lea.....	739, 740
Substriatus (Unio) Lea.....	811	Symphynota Lea.....	506, 514, 583, 662, 663
Subtentus (Margarita) Lea.....	591	Syriaeus (Margaron) Lea.....	695
Subtentus (Margaron) Lea.....	591	Syriaeus (Unio) Lea.....	695
Subtentus (Medionidus) Say.....	591	Syrmatophora (Hyria) Gronovius.....	868
Subtentus (Unio) Say.....	589, 591	Syrmatophora (Hyria) Sowerby.....	870
Subtetrгона (Anodon) Heude.....	638	Syrmatophora (Mya) Meuschen.....	870
Subtetrгона (Unio) Michaud.....	691	Syrmatophora (Unio) Deshayes.....	870
Subtigridis (Unio) Locard.....	689	Syrmatophorus (Margarita) Lea.....	870
Subtrapezius (Unio) Hanley.....	919	Syrmatophorus (Prisodon) Meuschen.....	870
Subtriangularis (Burtonia) Bourguignat.....	578	Tabaseoensis (Anodonta) Morelet.....	646
Subtriangularis (Pseudospatha) Bourguignat.....	578	Tabaseoensis (Margaron) Lea.....	596
Subtortus (Unio) Baird and Adams.....	798	Tabaseoensis (Nephronaia) Charpentier.....	596
Subtrigonus (Margaron) Lea.....	580	Tabaseoensis (Unio) Charpentier.....	596
Subtrigonus (Unio) Sowerby.....	580	Tabula (Anodon) Sowerby.....	898
Subvexa (Anodonta) Conrad.....	533	Tæniatus (Lampsilis) Conrad.....	541
Subvexa (Lampsilis) Conrad.....	533	Tæniatus (Margarita) Lea.....	541
Subvexa (Margarita) Lea.....	534	Tæniatus (Unio) Conrad.....	541
Subvexa (Margaron) Lea.....	534	Tætiatus (Unio) Catlow and Reeve.....	744
Subviridis (Margaron) Lea.....	870	Taitiana (Pleurobema) Lea.....	754
		Taitianus (Margarita) Lea.....	754

	Page.		Page.
Taitianus (Margaron) Lea.....	754	Teretiusculus (Unio) Jickeli.....	826
Taitianus (Unio) Lea.....	744, 754	Teretiusculus (Unio) Philippi.....	812
Taitianus (Unio) Sowerby.....	602	Terminalis (Margaron) Lea.....	689
Tampecoensis (Lampsilis) Lea.....	570	Terminalis (Unio) Bourguignat.....	689
Tampecoensis (Margarita) Lea.....	571	Tesserula (Margaron) Lea.....	749
Tampecoensis (Margaron) Lea.....	571	Tesserula (Pleurobema) Lea.....	749
Tampecoensis (Unio) Lea.....	570	Tesserula (Unio) Lea.....	749
Tamsana (Monocondylea) Dunker.....	913	Testa crassa (Mya) Schröter.....	677
Tanganikana (Grandidieria) Bourguignat.....	828	Testudinarius (Unio) Spengler.....	854
Tanganyicensis (Burtonia) Patet.....	577	Testudineus (Unio) Morelet.....	702
Tanganyicensis (Grandidieria) Smith.....	828	Testudineus (Unio) Reeve.....	571
Tanganyicensis (Pseudospatha) Smith.....	577	Tetragene.....	514, 658, 765
Tanganyicensis (Spatha) Smith.....	577, 578	Tetragona (Anodonta) Lea.....	617
Tanganyicensis (Unio) Smith.....	828, 848	Tetragona (Margaron) Lea.....	617
Taniatus (Unio) Patet.....	541	Tetragona (Strophitus) Conrad.....	617
Tanousi (Anodonta) Bourguignat.....	649	Tetralasmus campodon (Unio) Say.....	740
Tanousi (Collopterygia) Bourguignat.....	649	Tetralasmus declivis (Unio) Say.....	740
Tappanianus (Margarita) Lea.....	663	Tetralasmus manubius (Unio) Gould.....	740
Tappanianus (Margaron) Lea.....	663	Tetralasmus (Margarita) Lea.....	740
Tappanianus (Unio) Lea.....	663	Tetralasmus (Margaron) Lea.....	740
Tappianus (Unio) Catlow and Reeve.....	663	Tetralasmus sayi (Unio) Ward.....	741
Tavoyensis (Margaron) Lea.....	843	Tetralasmus (Unio) Say.....	739
Tavoyensis (Parreysia) Gould.....	843	Tetraplodon Spix.....	536, 515, 863, 869
Tavoyensis triembolus (Parreysia) Benson.....	844	Tetricus (Margaron) Lea.....	718
Tavoyensis (Unio) Gould.....	843	Tetricus (Unio) Lea.....	718
Tawi (Anodonta) Rang.....	902	Texasensis (Anodonta) Lea.....	645
Tecomatensis (Lampsilis) Lea.....	571	Texasensis compressus (Lampsilis) Simpson.....	564
Tecomatensis (Margaron) Lea.....	571	Texasensis (Lampsilis) Lea.....	563, 564
Tecomatensis (Unio) Lea.....	571	Texasensis (Margaron) Lea.....	563, 645
Tehuantepecensis (Anodonta) Fischer and Crosse.....	933	Texasensis (Unio) Lea.....	563
Tehuantepecensis (Nephronalax) Crosse and Fischer.....	594	Theca (Trapezoides) Benson.....	859
Tehuantepecensis (Unio) Crosse and Fischer.....	594	Theca (Unio) Benson.....	859
Tellicoensis (Unio) Lea.....	751	Theliderma Swainson.....	775
Tenebricosa (Anodonta) Lea.....	929, 930	Thomsoni (Grandidieria) Smith.....	829
Tenebricosa (Margarita) Lea.....	930	Thomsoni (Margaritana) Patet.....	838
Tenebricosa (Margaron) Lea.....	930	Thomsoni (Pseudodon) Morelet.....	838
Tenebricosus (Glabaris) Lea.....	930, 932	Thomsoni (Unio) Smith.....	829
Tenebricus (Margaron) Lea.....	549	Thorntonii (Margaron) Lea.....	750
Tenebricus (Unio) Lea.....	549	Thorntonii (Unio) Lea.....	750
Tenebrosus (Unio) Conrad.....	536	Thottoni (Mutelina) Roehbrune.....	906
Tenella (Anodonta) Kuster.....	624	Thwaitesii (Lamellidens) Lea.....	856
Teneltus (Unio) Rafinesque.....	744	Thwaitesii (Margaron) Lea.....	856
Tener (Lampsilis) Lea.....	555	Thwaitesii (Unio) Lea.....	856
Tener (Margaron) Lea.....	555	Tiberiadensis (Unio) Locard.....	688
Tener (Unio) Lea.....	555	Tientsinensis (Quadrula) Crosse and Debeaux.....	798
Tenerus (Margarita) Lea.....	559	Tientsinensis (Unio) Crosse and Debeaux.....	798
Tenerus (Margaron) Lea.....	559	Tientsinensis (Unio) Heude.....	798
Tenerus (Unio) Ravenel.....	559	Tigridis (Unio) Bourguignat.....	688, 690
Tenerus (Unio) Simpson.....	551	Tigris (Margaron) Lea.....	688
Tennenti (Unio) Hanley and Theobald.....	842	Tinius (Unio) Locard.....	693
Tennesseensis (Margaron) Lea.....	541	Tinker (Obovaria) B. H. Wright.....	600
Tennesseensis (Unio) Lea.....	541	Tinkeri (Unio) B. H. Wright.....	600
Tenuis (Anodonta) Lea.....	833	Tombigbeensis (Margaritana) Lea.....	619
Tenuis (Dallia) Lea.....	833	Tombigbeensis (Margaron) Lea.....	619
Tenuis (Margaron) Lea.....	833	Tombigbeensis (Strophitus) Lea.....	619
Tenuis (Unio) Gray.....	585	Topekaensis (Margaron) Lea.....	547
Tenuissima (Symphynota) Lea.....	576	Topekaensis (Unio) Lea.....	547
Tenuissimus (Margarita) Lea.....	516	Torrentis (Anodonta) Heude.....	640
Tenuissimus (Margaron) Lea.....	516	Torsa (Obovaria) Rafinesque.....	599, 603
Tenuissimus (Unio) Lea.....	576, 577	Torsus (Unio) Potiez and Michaud.....	599
Teres (Anodonta) Conrad.....	633	Tortilis (Anodonta) Lea.....	920
Teres (Unio) Say.....	543	Tortilis (Glabaris) Lea.....	920
Teretiuscula (Nodularia) Philippi.....	812	Tortilis (Margaron) Lea.....	920
		Tortivus (Margaron) Lea.....	718

	Page.		Page.
Tortivus (Unio) Lea.....	718	Trifidus (Margaron) Lea.....	884
Tortuosa (Quadrula) Lea.....	798	Trifidus (Unio) Lea.....	884
Tortuosus (Margaron) Lea.....	798	Triformis (Nodularia) Heude.....	807
Tortuosus (Unio) Lea.....	798	Triformis (Unio) Heude.....	807
Tortuosus (Unio) Sowerby.....	724	Trigona (Anodonta) Held.....	648
Torulosa (Amblema) Rafinesque.....	744	Trigona (Margarita) Lea.....	788, 928
Torulosus (Unio) Conrad.....	523	Trigona (Margaron) Lea.....	928
Torulosus (Unio) Ferrussac.....	704	Trigona (Quadrula) Lea.....	510, 757, 787, 928
Torulosus (Unio) Short and Eaton.....	611	Trigonodon Conrad.....	834, 835
Trabalis (Lampsilis) Conrad.....	553	Trigonus (Glabaris) Spix.....	927, 928
Trabalis (Unio) Conrad.....	558	Trigonus (Margaron) Lea.....	788
Trachæa (Unio) Kobelt.....	693	Trigonus (Unio) Lea.....	503, 504, 509, 784, 787
Tragosus (Unio) Hanley.....	778	Trigonus (Unio) Sowerby.....	786
Transversa (Hyria) Hupe.....	869	Trinaerus (Margaron) Lea.....	614
Trapesialis (Anodonta) Lamarck.....	923	Trinaerus (Ptychobranchus) Lea.....	614
Trapesialis (Anodontites) Lamarck.....	923	Trinaerus (Unio) Lea.....	614
Trapesialis anserinus (Glabaris) Spix.....	924	Tripartitus (Margaron) Lea.....	843
Trapesialis cygnaeformis (Glabaris) Pilsbry.....	925	Tripartitus (Unio) Lea.....	842, 843
Trapesialis exotica (Glabaris) Lamarck.....	924	Triplodon Spix.....	868
Trapesialis (Glabaris) Lamarck.....	914, 923, 924	Tripolitana (Leguminaia) Bourguignat.....	652
Trapesialis (Iridina) d'Orbigny.....	923	Tripolititanus (Margaritana) Clessin.....	652
Trapesialis moretonianus (Glabaris) Sowerby.....	925	Tripolititanus (Pseudodon) Conrad.....	652
Trapesialis rioplatensis (Glabaris) Sowerby.....	925	Tripolititanus (Unio) Bourguignat.....	652
Trapesialis scriptus (Glabaris) Sowerby.....	924	Triqueter (Unio) Short and Eaton.....	517
Trapezea (Glabaris) von Ihering.....	917	Triquetra Klein.....	869
Trapezeus (Anodon) Spix.....	917	Triquetra (Truncilla) Rafinesque.....	516, 517, 519
Trapezia senilis (Spatha) von Martens.....	897	Triquetra Simpson.....	869
Trapezia (Spatha) von Martens.....	897	Trirostris (Unio) Musgrave.....	842
Trapezialis (Anodonta) Blainville.....	923	Trirostris (Unio) Sowerby.....	812
Trapezialis (Columba) Pætel.....	923	Tristis (Spatha) Jousseume.....	897
Trapezialis (Margarita) Lea.....	923	Tristis (Unio) Morelet.....	676
Trapezialis (Margaron) Lea.....	923	Tristrami (Unio) Locard.....	688
Trapezoides (Margarita) Lea.....	772	Trisulcata (Parreysia) Heude.....	845
Trapezoides (Margaron) Lea.....	772	Trisulcatus (Unio) Heude.....	802, 845
Trapezoides (Quadrula) Lea.....	772	Tritogonia Agassiz.....	506, 510, 514, 607
Trapezoides (Unio) Lea.....	772	Triumphans (Quadrula) B. H. Wright.....	770
Trapezoideus Simpson.....	506, 515, 858	Triumphans (Unio) B. H. Wright.....	770
Trautwiniana (Anodonta) Lea.....	922	Trompi (Nodularia) Drouet and Chaper.....	816
Trautwiniana (Margaron) Lea.....	922	Trompi (Unio) Drouet and Chaper.....	816
Trautwinianus (Glabaris) Lea.....	922	Troostensis (Margarita) Lea.....	558
Traversii (Nodularia) Pollonera.....	825	Troostensis (Margaron) Lea.....	558
Traversii (Unio) Pollonera.....	825	Troostensis (Unio) Lea.....	558
Triangularis (Anodonta) Sowerby.....	648	Troostensis (Unio) Sowerby.....	531, 558
Triangularis (Anodonta) Trask.....	629	Troostii (Margaron) Lea.....	558
Triangularis (Margarita) Lea.....	517	Troostii (Unio) B. H. Wright.....	558
Triangularis (Margaron) Lea.....	517	Troschelliana (Pleurobema) Lea.....	754, 756, 757
Triangularis (Mya) Eaton.....	517	Troschellianus (Margaron) Lea.....	756
Triangularis (Mycetopus) Heude.....	657	Troschellianus (Unio) Lea.....	756
Triangularis (Obliquaria) Rafinesque.....	744	Trossulus (Lampsilis) Lea.....	562
Triangularis (Solenia) Heude.....	657	Trossulus (Margaron) Lea.....	562
Triangularis (Unio) Barnes.....	517	Trossulus (Unio) Lea.....	562
Triangularis (Unio) Kuster.....	788	Troutwiniana (Anodonta) H. and A. Adams.....	922
Triangularis (Unio) Say.....	602	Troutwiniana (Margaron) Lea.....	922
Triangulata (Alasmidonta) Lea.....	672	Truncata (Alasmodonta) B. H. Wright.....	671
Triangulata (Margaritana) Lea.....	672	Truncata (Alasmodonta) Conrad.....	668
Triangulata (Margaron) Lea.....	672	Truncata (Anodonta) Kuster.....	649
Triangulata (Unio) Sowerby.....	672	Truncatus elegans (Unio) Rafinesque.....	605
Triclava (Quadrula) Heude.....	800	Truncatus (Margarita) Lea.....	688
Triclavus (Unio) Heude.....	800	Truncatus (Margaron) Lea.....	688
Tricolor (Unio) Kuster.....	855	Truncatus (Prisodon) Schumacher.....	863, 868
Tricostatus (Anodon) Sowerby.....	638	Truncatus (Unio) Say.....	605
Triembolus (Unio) Benson.....	844	Truncatus (Unio) Spengler.....	855, 862
Trifidus (Diplodon) Lea.....	884	Truncatus (Unio) Swainson.....	688
		Truncilla Rafinesque.....	506, 511, 514, 516, 517
		Tryoni (Unio) B. H. Wright.....	714

	Page.		Page.
Tryonif (Anodonta) Lea	632	Undulata (Alasmodonta) Say	667
Tryonif (Margaron) Lea	632	Undulata (Anodon) Gould	617
Tuberculata (Cristaria) Schumacher	584	Undulata (Anodonta) Hildreth	616
Tuberculata (Mya) Eaton	608	Undulata (Anodonta) Say	616, 618
Tuberculata obesa (Tritogonia) Simpson	609	Undulata (Baphia) H. and A. Adams	667
Tuberculata (Obliquaria) Rafinesque	794, 795	Undulata latecostata (Quadrula) Lea	769
Tuberculata (Quadrula) Rafinesque	794, 795	Undulata (Margarita) Lea	618, 667
Tuberculata (Rotundaria) Agassiz	795	Undulata (Margaritana) Kuster	667
Tuberculata (Tritogonia) Barnes	608	Undulata (Margaron) Lea	618, 667
Tuberculatus (Margarita) Lea	608	Undulata (Monodonta) Say	666, 667
Tuberculatus (Margaron) Lea	608	Undulata (Mya) Eaton	769
Tuberculatus (Unio) Barnes	509, 608	Undulata (Mya) Wood	667
Tuberculatus (Unio) Conrad	795	Undulata pilsbryi (Quadrula) Marsh	769
Tuberculosa (Unio) Valenciennes	795	Undulata (Quadrula) Barnes	768, 769
Tuberosa (Quadrula) Lea	774	Undulata (Unio) Deshayes	616
Tuberosus (Margaron) Lea	774	Undulata (Unio) Hanley	667
Tuberosus (Unio) Lea	774	Undulatus (Margarita) Lea	769
Tumescens (Margaron) Lea	750	Undulatus (Margaron) Lea	769
Tumescens (Pleurobema) Lea	750	Undulatus (Strophitus) Say	618
Tumescens (Unio) Lea	750	Undulatus (Strophitus) Stimpson	667
Tumida (Anodon) Heude	638	Undulatus (Unio) Barnes	769, 802
Tumida (Anodonta) Kuster	625	Undulatus (Unio) Say	770
Tumida (Monocondylæa) Deshayes and Jullien	838	Undulifera (Anodonta) Clessin	641
Tumidula (Nodularia) Lea	814	Unicolor (Margaron) Lea	601
Tumidulus (Margaron) Lea	814	Unicolor (Obovaria) Lea	601
Tumidulus (Unio) Lea	814	Unicolor (Unio) Lea	601
Tumidus (Margaron) Lea	685	Unicostatus (Unio) B. H. Wright	547
Tumidus (Monocondylus) Morelet	837	Unio Retzius	501, 502, 503, 506, 510, 513, 514, 518, 530, 539, 579, 593, 602, 644, 679, 703, 738, 740, 745, 760, 772, 791, 792, 798, 799, 801, 804, 845, 851, 857, 867, 877, 880, 883, 892
Tumidus (Pseudodon) Morelet	837	Uniomercus Conrad	680, 739
Tumidus (Unio) Retzius	513, 530, 679, 680, 684, 685, 698	Unionidæ	501, 502, 503, 504, 505, 507, 511, 513, 514, 515, 516, 535, 744, 861, 863, 882, 933
Tunizana (Anodonta) Morelet	649	Unioninæ	501, 514, 516
Tuomeyi (Margaron) Lea	726	Uniopsis Agassiz	616
Tuomeyi (Unio) Lea	726	Uniopsis (Anodonta) Lamarek	653
Turcicus (Unio) Kuster	700	Uniopsis (Leguminaia) Lamarek	653
Tureicus (Unio) Parreys	694	Uniopsis (Margarita) Lea	653
Turgida (Castalia) Hupe	863	Uniopsis (Margaron) Lea	653
Turgidulus (Margaron) Lea	524	Uniopsis Swainson	666
Turgidulus (Unio) Lea	524	Upsoni (Unio) Marsh	539
Turgidus (Margarita) Lea	781	Uruguayensis (Anodonta) Lea	917
Turgidus (Margaron) Lea	781	Uruguayensis (Columbia) Patel	917
Turgidus (Unio) Lea	781	Uruguayensis (Diplodon) Lea	875, 885
Turgidus (Unio) Reeve	776	Uruguayensis (Margaron) Lea	875, 917
Turtoni (Unio) Payraudeau	685, 687	Uruguayensis (Unio) Lea	875
Tusembiensis (Unio) Lea	749	Usumasintæ (Quadrula) Crosse and Fischer	796
Uber (Unio) Conrad	744, 780	Usumasintæ (Unio) Crosse and Fischer ..	796
Ucayalensis (Anodonta) Philippi	928	Utriculus (Margaron) Lea	785
Uharensis (Margaron) Lea	724	Utriculus (Quadrula) Lea	785
Uharensis (Unio) Lea	724	Utriculus (Unio) Lea	785
Umbonatus (Glabaris) Simpson	926	Ujijiensis (Grandidieria) Bourguignat ..	848
Umbonatus (Unio) Bourguignat	692	Ujijiensis (Parreysia) Bourguignat	848
Umbrans (Margaron) Lea	549	Vaalensis (Unio) Chaper	825
Umbrans (Unio) Lea	549	Vagulus (Hyriopsis) Fischer	580
Umbrosus (Lampisilis) Lea	570	Vagulus (Unio) Fischer	581
Umbrosus (Margaron) Lea	570	Valdivanus (Unio) Philippi	881
Umbrosus (Unio) Lea	570	Valentinus (Margaron) Lea	686
Unadilla (Anodon) De Kay	617	Valentinus (Unio) Rossmässler	686
Unadilla (Strophitus) Conrad	617	Valentinus (Unio) Sowerby	692
Undata (Mya) Eaton	788	Valcus (Iridina) Jay	908
Undatus (Unio) Barnes	788	Valida (Pleurobema) Lea	749
Undatus (Unio) Conrad	788	Validus (Unio) Lea	749
Undosa (Castalia) von Martens	866	Vallata (Quadrula) Lea	782
Undosa (Castalina) von Martens	866		
Undulata (Alasmodonta) Say	666, 667		
Undulata (Alasmodonta) C. B. Adams ..	618		

	Page.		Page.
Vallatus (Margaron) Lea.....	782	Verrucosa (Tritogonia) Smith.....	609
Vallatus (Unio) Lea.....	782	Verrucosa (Tritogonia) Agassiz.....	609
Vanuxemensis (Lampisilis) Lea.....	548, 549	Verrucosa (Unio) Valenciennes.....	779
Vanuxemensis (Margarita) Lea.....	549	Verrucosus albus (Unio) Hildreth.....	610
Vanuxemensis (Margaron) Lea.....	549	Verrucosus (Margarita) Lea.....	795
Vanuxemensis (Unio) Conrad.....	549	Verrucosus (Margaron) Lea.....	795
Vanuxemensis (Unio) Lea.....	549	Verrucosus purpureus (Unio) Hildreth.....	795
Vanuxemensis (Unio) Sowerby.....	558	Verrucosus (Unio) Barnes.....	795
Vanuxemi (Margaron) Lea.....	549	Verrucosus (Unio) Say.....	608
Vanuxemii (Unio) B. H. Wright.....	549	Verruculosus (Unio) Heude.....	802
Variabilis (Anodonta) Draparnaud.....	622	Versus (Margaron) Lea.....	814
Variabilis (Diplodon) Maton.....	888	Verus (Margaron) Lea.....	757
Variabilis (Margarita) Lea.....	888	Verus (Unio) Lea.....	757
Variabilis (Margaron) Lea.....	888	Verutus (Margaron) Lea.....	727
Variabilis (Mya) Maton.....	888	Verutus (Unio) Lea.....	726
Variabilis (Unio) d'Orbigny.....	888	Vescoi (Margaron) Lea.....	694
Varicosa (Pleurobema).....	509	Vescoi (Unio) Bourguignat.....	694
Varicosa (Unio) Lamarck.....	671	Vescoiana (Anodonta) Bourguignat.....	628
Varicosus (Margarita) Lea.....	765	Vescoiana (Margaron) Lea.....	628
Varicosus (Margaron) Lea.....	765	Vesicularis (Lampisilis) Lea.....	562
Varicosus (Unio) Lea.....	765	Vesicularis (Unio) Lea.....	562
Vatus (Unio) Patel.....	728	Vestitus (Margaron) Lea.....	708
Vaughanians (Lampisilis) Lea.....	550	Vestitus (Unio) Lea.....	708, 804
Vaughanians (Margarita) Lea.....	550	Vestitus var. a (Unio) Heude.....	801
Vaughanians (Margaron) Lea.....	550	Vestitus var. b (Unio) Heude.....	804
Vaughanians (Unio) Lea.....	550	Vibex (Lampisilis) Conrad.....	559
Vaughanians (Unio) Sowerby.....	546	Vibex (Margarita) Lea.....	560
Velaris (Physunio) Sowerby.....	831	Vibex (Margaron) Lea.....	560
Velaris (Unio) Hanley.....	830	Vibex nigrinus (Lampisilis) Lea.....	560
Velaris (Unio) Sowerby.....	831	Vibex (Unio) Conrad.....	559, 716
Velatus (Unio) Conrad.....	613	Vicinus (Margaron) Lea.....	730
Vellicatus (Nephronalae) Reeve.....	597	Vicinus (Unio) Lea.....	730
Velthuzeni (Hyriopsis) Schepman.....	581	Vignoniiana (Anodonta) Musgrave.....	902
Velthuzeni (Unio) Schepman.....	581	Vignoniiana (Margaritana) Bernardi.....	901, 902
Velum (Unio) Say.....	575	Vignoniiana (Margaron) Lea.....	902
Ventricosa (Anodonta) C. Pfeiffer.....	623	Vignoniiana (Spatha) Bernardi.....	902
Ventricosa (Mycetopoda) d'Orbigny.....	935	Vignoniiana (Unio) Reeve.....	902
Ventricosa (Mycetopus) d'Orbigny.....	935	Villosus (Lampisilis) B. H. Wright.....	562
Ventricosus (Lampisilis) Barnes.....	526, 527, 529	Villosus (Unio) B. H. Wright.....	562
Ventricosus (Margarita) Lea.....	526	Vinekei (Unio) Bourguignat.....	862
Ventricosus (Margaron) Lea.....	526	Violaceus (Unio) Spengler.....	721
Ventricosus (Mya) Eaton.....	526	Virens (Anodonta) Lea.....	645
Ventricosus (Platiris) Lea.....	935	Virens (Margaron) Lea.....	645, 722
Ventricosus satur (Lampisilis) Lea.....	527	Virens (Unio) Lea.....	722
Ventricosus (Unio) Barnes.....	526, 527, 528, 533	Vireseens (Lampisilis) Lea.....	544
Ventrosa (Brazzeia) Bourguignat.....	907	Vireseens (Margaron) Lea.....	544
Venus (Unio) Rafinesque.....	744	Vireseens (Unio) Lea.....	544
Venustus (Lampisilis) Lea.....	543	Virgata (Anodonta) Conrad.....	617
Venustus (Margarita) Lea.....	543	Virgatus (Strophitus) Conrad.....	617
Venustus (Margaron) Lea.....	543	Virginiana (Unio) Lamarck.....	536, 721
Venustus (Unio) Morelet.....	815	Virgulata (Anodonta) Lea.....	631
Venustus (Unio) Sowerby.....	539	Virgulata (Margaron) Lea.....	631
Vera (Pleurobema) Lea.....	757	Virgulus Simpson.....	931
Veraacruzensis (Margaron) Lea.....	570	Virgus Simpson.....	506, 515, 851
Veraacruzensis (Unio) Lea.....	570	Viridana (Anodonta) Clessin.....	922
Verae-pacis (Unio) Tristram.....	738	Viridans (Margaron) Lea.....	726
Verbecki (Nodularia) Böttger.....	819	Viridans (Unio) Lea.....	726
Verbecki (Unio) Böttger.....	819	Viridicatus (Margaron) Lea.....	717
Verecundus (Unio) Gould.....	832	Viridicatus (Unio) Lea.....	717
Verreauxi (Unio) Patel.....	825	Viridiflavus (Unio) Kuster.....	683
Verreauxianus (Margaron) Lea.....	825	Viridiradiatus (Margaron) Lea.....	726
Verreauxianus (Unio) Lea.....	825	Viridiradiatus (Unio) Lea.....	726
Verrucifer (Unio) von Martens.....	862	Viridis (Alasmodon) Rafinesque.....	673
Verrucosa (Mya) Eaton.....	795	Viridis (Anodonta) H. and A. Adams.....	619
Verrucosa (Obliquaria) Rafinesque.....	608	Viridis (Mycetopus) Heude.....	656
Verrucosa (Quadrula) Baker.....	795	Viridis (Symphynota) Conrad.....	663

	Page.		Page.
Viridis (Unio) Conrad.....	663	Wheatleyi (Microcondylæa) Clessin.....	652
Viridis (Unio) Ferussac.....	529	Wheatleyi (Monocondylæa) Lea.....	652
Viridis (Unio) Rafinesque.....	744	Wheatleyi (Pseudodon) Conrad.....	652
Viridulus (Margaron) Lea.....	734	Wheatleyi (Unio) Lea.....	725
Viridulus (Unio) Lea.....	734	Whiteianus (Margaron) Lea.....	716
Visseri (Mutela) Bourguignat.....	904	Whiteianus (Unio) Lea.....	716
Visseri (Unio) Bourguignat.....	862	Williamsii (Anodonta) Lea.....	632
Vittatis (Bariosta) Rafinesque.....	744	Williamsii (Margaron) Lea.....	632
Vittatus (Diplodon) Lea.....	890	Wilsonii (Diplodon) Lea.....	892
Vittatus (Margaron).....	890	Wilsonii (Margaron) Lea.....	892
Vittatus (Unio) Lea.....	890	Wilsonii (Unio) Lea.....	892
Von buschea (Unio) Sowerby.....	836	Wissmani (Spatha) von Martens.....	896
Vondembuschiana (Margaritana) Lea....	836	Wolwiehii (Unio) Morelet.....	699
Vondembuschiana (Monocondylæa) H. and A. Adams.....	836	Woodiana (Anodonta) Kobelt.....	639
Vondembuschianus (Pseudodon) Lea....	836	Woodiana (Anodonta) Lea.....	637
Vondembuschii (Unio) Sowerby.....	835	Woodiana (Margarita) Lea.....	637
Vuleana (Parreysia) Hanley.....	844	Woodiana (Margaron) Lea.....	637
Vulcanus (Unio) Hanley.....	844	Woodiana (Symphynota) Lea.....	637
Vyneklî (Cameronia) Bourguignat.....	910	Woodwardianus (Margaron) Lea.....	613
Vysseri (Mutela) Bourguignat.....	904	Woodwardianus (Unio) Lea.....	613
Waccamawensis (Margaron) Lea.....	711	Woodwardius (Unio) Lea.....	613
Waccamawensis (Unio) Lea.....	711	Wrightii (Margaron) Lea.....	808
Wagneri (Unio) Kobelt.....	693	Wrightii (Unio) Lea.....	808
Wagnerianum (Diplodon) Simpson.....	877	Wymanii (Anodonta) Lea.....	917
Wahlametensis (Anodonta) Lea.....	629	Wymanii (Diplodon) Lea.....	874
Wahlametensis (Margarita) Lea.....	629	Wymanii (Glabaris) Lea.....	917
Wahlametensis (Margaron) Lea.....	629	Wymanii (Margaron) Lea.....	874, 918
Wahlbergi bourguignati (Spatha) Bour- guignat.....	898	Wymanii (Unio) Lea.....	874
Wahlbergi (Iridina) Krauss.....	898	Wynegungaensis (Parreysia) Lea.....	842
Wahlbergi (Mutela) Smith.....	898	Wynegungaensis (Unio) Hanley and Theobald.....	842
Wahlbergi (Platiris) Lea.....	898	Wynegungaensis (Unio) Lea.....	842
Wahlbergi (Spatha) Krauss.....	898	Yadkinensis (Unio) Lea.....	724
Wahlbergi spathuliformis (Spatha) von Martens.....	898	Yokohamensis (Unio) von Ihering.....	810
Waikarensis (Diplodon) Colenso.....	890	Youconensis (Anodonta) Lea.....	628
Waikarensis (Unio) Colenso.....	890	Youkanensis (Anodonta) Lea.....	628
Walkerî (Medionidus) B. H. Wright.....	590, 591	Youkanensis (Margaron) Lea.....	628
Wallisi (Anodonta) Mousson.....	933	Yubaensis (Alasmodon) Trask.....	677
Walpolei (Monocondylæa) Hanley.....	840	Yzabalensis (Nephronalæ) Crosse and Fischer.....	596
Waltoni (Unio) B. H. Wright.....	736	Yzabalensis (Unio) Crosse and Fischer ..	596
Wardiana (Anodonta) Lea.....	617	Zabulonius (Unio) Kobelt.....	690
Wardiana (Margarita) Lea.....	617	Zabulonius (Unio) Locard.....	689
Wardiana (Margaron) Lea.....	617	Zaira Roehbrune.....	862
Wardii (Margaron) Lea.....	774	Zealandicus (Unio) Pætel.....	894
Wardii (Unio) Lea.....	774	Zeiglerianus (Unio) Lea.....	550
Waterensis (Margarita) Lea.....	722	Zezebori (Diplodon) Dunker.....	890
Waterensis (Margaron) Lea.....	722	Zezebori (Margaron) Dunker.....	890
Waterensis (Unio) H. and A. Adams.....	722	Zezebori (Unio) Dunker.....	890
Websteri (Unio) B. H. Wright.....	711	Zellensis (Anodonta) Waardenberg.....	622
Weddellii (Anodonta) Hupé.....	928	Zellensis (Mytilus, Gmelin.....	622
Weddellii (Glabaris) Hupé.....	928	Zeyheri (Unio) Menke.....	826
Weddellii (Margaron) Lea.....	928	Zieglerianus (Margaron) Lea.....	550
Weddellii (Mycetopus) Hupé.....	934	Zieglerianus (Margarita) Lea.....	550
Weldonensis (Margaron) Lea.....	723	Zigzag (Margarita) Lea.....	605
Weldonensis (Unio) Lea.....	723	Zigzag (Margaron) Lea.....	605
Welwitsehii (Iridina) Morelet.....	906	Zigzag (Unio) Lea.....	605
Wheatleyanus (Diplodon) Lea.....	875	Zollingeri (Alasmodonta) Mousson.....	837
Wheatleyanus (Margaron) Lea.....	875	Zollingeri (Anodonta) Lea.....	838
Wheatleyanus (Unio) Lea.....	875	Zollingeri (Margaritana) Kuster.....	837
Wheatleyi (Anodonta) Lea.....	928	Zollingeri (Monocondylæa) Pætel.....	837
Wheatleyi (Glabaris).....	932	Zollingeri (Pseudodon) Mousson.....	837
Wheatleyi (Leguminaia) Lea.....	652	Zonata (Quadrula) Heude.....	793
Wheatleyi (Margaron) Lea.....	652, 928	Zonata (Unio) Rafinesque.....	744
		Zonatus (Unio) Heude.....	798