 

Digitized by the Internet Archive
in 2014

3 9007 0289 8917 6

H. Patrick Henry '38.

MS

ACTE

1820

8

WILD PLANTS OF CANADA

WILD PLANTS OF CANADA

A FLORA, WITH DESCRIPTIVE KEY OF
FAMILIES REPRESENTED

BY

H. B. SPOTTON, M.A., LL.D., F.L.S.

FORMERLY INSPECTOR OF HIGH SCHOOLS FOR ONTARIO

AND

A. COSENS, M.A., PH.D.

FORMERLY SCIENCE MASTER, PARKDALE COLLEGIATE INSTITUTE, TORONTO

AND

T. J. IVEY, M.A.

LATE SCIENCE MASTER, HARBORD COLLEGIATE INSTITUTE, TORONTO

TORONTO

W. J. GAGE & COMPANY, LIMITED

1935

QK

201

S65

cop. 2

Strawie

ALL RIGHTS RESERVED

PREFACE

Since the last edition of Dr. Spotton's *Flora of the Wild Plants of Canada* was issued, so many changes in nomenclature and classification have been made that a new edition of the book has been considered advisable.

While the general plan of the *Flora* has not been altered, several differences will be noticed in the new edition. These are concerned chiefly with the nomenclature and the general arrangement of the subject matter; in this connection it may be stated that the *Key* has been entirely re-written, and fuller detail added to make its use easier for the beginner in Botany. In the body of the book many species of plants from Ontario have been added. The majority of these are from the southwestern part of the province. Also a large number of additional species from Western Canada have been included; this will make the *Flora* even more useful than it has been to the botanists of the Prairie Provinces.

As the necessary information is not available no attempt has been made to indicate the range of the various species. Plants reported from the Prairie Provinces, formerly marked "N.W." are now indicated by the abbreviation "Pr. Prov." while those from the eastern part of Canada are marked "Atl. Prov." as before.

While, in general, the *Flora* is limited to the commonly occurring wild plants, many introduced species of rather local range have been included, especially when these are known to be noxious.

The classification and nomenclature followed in the present revision are, with few exceptions, those of the 7th edition of Dr. Gray's *New Manual of Botany*. Other descriptive works have been consulted, and among those found especially useful should be mentioned *The Forest Trees of Ontario* by Dr. J. H. White. To these authors and the following:—Britton and Brown, Dodge, Sargent, Macoun, and others, we gratefully acknowledge our indebtedness.

Sincere thanks are extended to all who assisted in the work of revision, but more particularly to the Botanical Department of the University of Toronto and to Mr. F. J. A. Morris, Collegiate Institute, Peterborough, Ont.

In spite of care, omissions and mistakes, more or less serious, may have occurred. The revisers will greatly appreciate having their attention drawn to these.

A.C.

Toronto, October, 1926.

T.J.I.

Some Familiar Wild Flowers and Mushrooms, Ferns, and Grasses and Some More Wild Flowers, both by Magistrate James Edmund Jones, B.A., may be used with advantage in connection with this text. Both are very fully illustrated with photographic reproductions of the greater number of the wild flowers dealt with in *Wild Plants of Canada*. Copies may be obtained from the National Boys' Work Board, 21 Dundas Square, Toronto, or from W. J. Gage & Co., Limited, 84 Spadina Avenue, Toronto.

TABLE OF CONTENTS

KEY TO THE FAMILIES.....	1
THALLOPHYTES.....	1
BRYOPHYTES.....	1
FLORA.....	21
PTERIDOPHYTES.....	21
SPERMATOPHYTES.....	36
GYMNOSPERMS.....	36
ANGIOSPERMS.....	39
MONOCOTYLEDONS.....	39
DICOTYLEDONS.....	69
GLOSSARY.....	277
ABBREVIATIONS OF PROPER NAMES.....	287
INDEX.....	291

WILD PLANTS OF CANADA

DESCRIPTIVE KEY

DIVISION I. THALLOPHYTES

Including (1) Algae—chlorophyll-bearing plants, most of which live in water. Pond scums, sea weeds, etc.

(2) Fungi—plants without chlorophyll, living on organic matter. Molds, bacteria, mushrooms, etc.

DIVISION II. BRYOPHYTES

Including (1) Hepaticae—plants with expanded body, bright green in colour or with leaves arranged in two rows on opposite sides of a stem; the whole plant flattened in appearance. Liverworts.

(2) Musci—plants with mostly erect stems, on which the greenish leaves are spirally arranged; spores produced in spore-cases borne usually on slender stalks. Mosses.

(The above Divisions are not included in the *Flora*.)

DIVISION III. PTERIDOPHYTES

Plants without true flowers, reproducing by spores and embracing clubmosses, horsetails, ferns and related plants.

A. Small floating plants or plants with floating leaves and delicate creeping rhizomes, which root in the mud.

Small floating plants which do not root in mud.

Leaves apparently 2-ranked.

SALVINIACEAE, 31

Plants with delicate creeping rhizomes, rooting in mud.

Leaves 4-parted, long-petioled and commonly floating.

MARSILEACEAE, 31

- A. Plants not growing in water, or, if growing in water, without floating leaves and delicate creeping root-stocks.
- B. Stems with sheathed joints, rush-like and mostly hollow. Spores produced in a terminal cone. EQUISETACEAE, 32
- B. Stems without sheathed joints, neither rush-like nor hollow.
- C. Leaves usually very narrow, with axillary sporangia.
 - D. Acaulescent; leaves long, narrow, in a rosette; mostly aquatic. ISOËTACEAE, 36
 - D. Stem elongate; leaves small, densely crowded along the stem.
 - E. Spores of two kinds, small and large, in different sporangia; plants 1-2 inches high. SELAGINELLACEAE, 35
 - E. Spores of one kind, small; sporangia all alike; plants usually much larger, 2-8 inches high. LYCOPODIACEAE, 33
- C. Ferns or fern-like plants; leaves (fronds) usually broad, without axillary sporangia.
 - F. Fertile frond or a fruiting part differing from a sterile; sporangia without ring or annulus, or with mere traces of one.
 - G. Rootstock erect, not elongate; frond consisting of two parts, the uppermost fertile or fruiting and the lower sterile and leaf-like; sporangia borne in two ranks. OPHIOGLOSSACEAE, 30
 - G. Rootstock not erect, either elongate or stout; fertile and sterile fronds quite unlike or clearly distinct; sporangia globose, densely crowded, but not 2-ranked. OSMUNDACEAE, 29
 - F. Fertile fronds and sterile fronds essentially alike (except in *Onoclea*); sporangia usually not globose, but, if globose, 2-ranked; the sporangia opening in all cases by an incomplete elastic ring or annulus. POLYPODIACEAE, 21

DIVISION IV. SPERMATOPHYTES

Plants with true flowers, reproducing by seeds.

SUBDIVISION I. GYMNOSPERMS

Ovules borne on an open scale, not inclosed. Trees or shrubs with needle-shaped, awl-shaped, or scale-like, mostly evergreen leaves.

- A. Fruit a cone or bluish and berry-like; leaves not 2-ranked, but appearing so in *Tsuga* and *Abies*. PINACEAE, 37
- A. Fruit red and berry-like; leaves spreading into two ranks. TAXACEAE, 36

SUBDIVISION II. ANGIOSPERMS

Ovules borne in a closed ovary.

CLASS I. MONOCOTYLEDONS

Parts of the flowers usually in threes or sixes, never in fives, leaves parallel-veined, rarely net-veined. Herbaceous except in species of *Smilax*.

- A. Small, free-floating plants, without true leaves, the single frond of each plant bearing flowers and simple roots or none. LEMNACEAE, 48
- A. Plants rooted in the ground, except in a very few cases and possessing stems and true leaves.
 - B. Perianth absent or consisting of scales, bristles, or chaffy bracts.
 - C. Flowers inclosed by chaff-like bracts or glumes; stems jointed; leaves narrow and sheathing. Stems terete or compressed, usually hollow; leaf sheaths split. GRAMINEAE, 46
 - Stems mostly somewhat triangular, usually solid; sheaths closed. CYPERACEAE, 46
 - C. Flowers not inclosed in scale-like or chaff-like bracts. Immersed aquatics; flowers perfect, monoecious or dioecious and not in globose heads. NAJADACEAE, 41
- Plants in damp or marshy places; flowers on a spadix, usually surrounded by a prominent spathe. ARACEAE, 46
- Aquatic or marsh plants; flowers not perfect, either monoecious or dioecious, in cylindrical spikes or globose heads. TYPHACEAE, 40

Aquatic or marsh plants; flowers in white or lead-coloured, woolly heads at the summits of striated scapes (2-12 inches high). *ERIOCAULACEAE*, 49

B. Perianth present, herbaceous and coloured, greenish in *Juncaginaceae* and in *Smilax*.

D. Perianth free from the ovary (ovary superior).

Marsh plants with sheathing, strongly veined leaves; the 3 petals white; the numerous carpels separate in a head or ring. *ALISMACEAE*, 44

Marsh plants with rush-like or grass-like leaves; the 3 petals greenish or greenish-yellow; 3-6 carpels more or less united. *JUNCAGINACEAE*, 44

Mostly terrestrial plants; perianth of 6 (in one case 4) similar divisions, coloured alike except in *Trillium*; one stamen in front of each part of the perianth. *LILIACEAE*, 53

Aquatic plants with perfect flowers subtended by a spathe; perianth of 6 blue or yellow divisions; corolla 2-lipped or tubular; stamens only 3 or 6 (3 long and 3 short). *PONTEDERIACEAE*, 50

Grass-like or sedge-like plants, with greenish or purplish-brown flowers; the 6 divisions of the perianth small and glume-like. *JUNCACEAE*, 50

Rush-like aquatics with perfect flowers in dense heads; perianth with the 3 outer divisions green and glume-like, the 3 inner yellow; stamens 3. *XYRIDACEAE*, 49

D. Perianth united to the ovary (ovary inferior).

Aquatic plants; only the pistillate flowers above water, the staminate ones submersed. *HYDROCHARITACEAE*, 45

Twining plants from knotty rootstocks; leaves heart-shaped; pod with 3 large wings. *DIOSCOREACEAE* 60

Plants with irregular flowers; stamens 1-2, gynandrous; seeds innumerable. *ORCHIDACEAE*, 61

Plants with showy flowers, mostly blue; leaves 2-ranked, equitant; stamens 3, opposite the outer segments of the perianth. *IRIDACEAE*, 60

Herbs with hexandrous flowers on a scape from a solid bulb. *AMARYLLIDACEAE*, 60

CLASS II. DICOTYLEDONS

Parts of the flowers mostly in fours or fives; leaves net-veined. Herbs, shrubs, or trees.

A. Corolla absent (apetalous), calyx present or absent.

B. Flowers in catkins.

C. Staminate (sterile) or pistillate (fertile) flowers in catkins or heads but not both.

Herbs with mostly opposite, palmately lobed or divided leaves; fruit a glandular achene.

CANNABINEAE (URTICACEAE), 81

Trees with alternate pinnate leaves; fruit a dry drupe, with a hard nutshell and a 4-lobed seed. (Juglans has polypetalous fertile flowers.)

JUGLANDACEAE, 74

Trees with alternate simple leaves; fruit a 1-seeded nut contained in a cup or involucre.

FAGACEAE, 78

C. Both staminate and pistillate flowers in catkins or heads.

Trees or shrubs with alternate simple leaves; fruit 1-celled, many-seeded, the seeds tufted with silky down.

SALICACEAE, 70

Trees with large, alternate, palmately lobed leaves; flowers in spherical heads; fruit club-shaped, 1-seeded nutlets.

PLATANACEAE, 129

Shrubs with resinous-dotted, fragrant leaves; fertile flowers, 1 under each bract; fruit in the form of drupe-like nuts, usually waxy-coated.

MYRICACEAE, 73

Trees or shrubs with alternate simple leaves; fertile flowers, 2 or 3 under each bract; fruit a 1-celled, 1-seeded nut.

BETULACEAE, 75

Trees with ovate simple leaves; the pistillate catkin forming a fleshy aggregate fruit from the enlarged calyces of the several flowers.

MOREAE (URTICACEAE), 80

B. Flowers not in catkins.

D. Calyx absent as well as the corolla.

- Marsh herbs with fragrant white flowers in a nodding spike; pistil of 3 or more carpels, separate or nearly so. PIPERACEAE, 70
- Immersed aquatics with sessile, whorled, finely dissected leaves; style 1. CERATOPHYLLACEAE, 101
- Aquatic herbs with opposite entire leaves; ovary 4-celled; styles 2. CALLITRICHACEAE, 160
- Shrubby evergreens with flowers in terminal heads, each in the axil of a scaly bract; fruit a small drupe. COREMA (EMPETRACEAE), 161
- D. Calyx present.
- E. Calyx free from the ovary (calyx inferior).
- F. Ovaries more than 1.
- G. Ovaries separate.
- Herbs with compound or lobed leaves; calyx 4-parted; stamens on the calyx; carpels 1-4. ROSACEAE, 129
- Mostly herbs or woody climbers; sepals usually coloured or petal-like; stamens on the receptacle. RANUNCULACEAE, 102
- Shrubs with prickly stems and pinnate leaves with transparent dots; fruit and leaves contain a pungent aromatic oil. ZANTHOXYLUM (RUTACEAE), 156
- G. Ovaries united into a compound pistil with more than 1 cell.
- Small herbs; sepals 5; carpels 5, united to form a 5-celled, 5-pointed capsule. (In *Sedum* the 5 carpels are separate.) CRASSULACEAE, 125
- Tall smooth herbs with unpleasant odour and large petioled leaves; sepals 5, white; ovary 10-celled, 10-seeded; fruit a purple berry. PHYTOLACCACEAE, 94
- Herbs with usually milky juice; no true calyx but a calyx-like involucre surrounding the flowers (none in *Acalypha*); ovary 3-celled, 3-seeded, protruding on a pedicel. EUPHORBIACEAE, 158

- Prostrate herbs forming mats; leaves spatulate and whorled; sepals 5, white on the inside; ovary 3-celled, many-seeded.
AIZOACEAE, 94
- Low fleshy herbs, growing on the seashore; ovary 1-celled, 5-valved, few-seeded.
GLAUX (PRIMULACEAE), 197
- Herbs having flowers with 2 stamens and a silicle, 2-celled, flattened contrary to the partition.
LEPIDIUM (CRUCIFERAE), 116
- Herbs with opposite entire leaves and stems that are usually swollen at the joints; styles 2-5, ovary usually 1-celled.
CARYOPHYLLACEAE, 95
- Trees with opposite leaves; flowers in clusters; ovary 2-celled; fruit two 1-seeded samaras joined.
ACERACEAE, 164
- Shrubs with simple leaves and greenish flowers; ovary 3-celled, 3-seeded, forming a black, berry-like drupe.
RHAMNUS (RHAMNACEAE), 166
- Trees with simple alternate leaves; styles 2; ovary 1-2-celled; fruit a 1-seeded samara winged all around.
ULMEAE (URTICACEAE), 80
- Shrubby evergreens with heath-like aspect and flowers solitary in the axils of the leaves; fruit a small drupe.
EMPETRUM (EMPETRACEAE), 161
- F. Ovary only 1, 1-celled, 1-seeded.
- Herbs having stems with swollen joints and stipules sheathing the stems (except in Eriogonum).
POLYGONACEAE, 84
- Herbs with watery juice and simple leaves; flowers in spikes or racemes; style 1; fruit an achene.
URTICAEAE (URTICACEAE), 80
- Small tufted herbs with opposite entire leaves and scarious stipules; flowers solitary or in small cymes; fruit a 1-seeded utricle.
ILLECEBRACEAE, 94

Weedy herbs with exstipulate leaves and greenish or reddish flowers, spiked and interspersed with scarious bracts.

AMARANTHACEAE, 93

Chiefly herbs with exstipulate leaves, greenish flowers without bracts.

CHENOPODIACEAE, 89

Trees with opposite pinnate leaves; calyx small, 4-cleft, regular or reduced to a mere disk; fruit a 1-2-seeded samara.

OLEACEAE, 199

Trees with alternate simple leaves; flowers upon branches of the same year; fruit a dark-purple drupe.

CELTIDAE (URTICACEAE), 80

Aromatic trees or shrubs with alternate simple leaves; flowers dioecious; fruit a blue or red drupe.

LAURACEAE, 112

Shrubs with alternate simple leaves and tough bark; flowers perfect, appearing before the leaves; fruit a red drupe.

THYMELAEACEAE, 175

E. Calyx united to the ovary (calyx superior).

Small smooth herbs with decumbent forking stems and inconspicuous, greenish-yellow flowers; stamens 8-10, very short and inserted on a fleshy disk; capsule 1-celled.

CHRYSOPLERIUM (SAXIFRAGACEAE), 126

Aquatic herbs; leaves entire or when immersed usually much-dissected; stamens 1-8; capsule 1-4-celled, each cell 1-seeded.

HALORAGIDACEAE, 181

Smooth herbs in wet places; petals none or small and reddish when growing out of water; stamens 4; capsule short, 4-lobed, many-seeded.

LUDVIGIA (ONAGRACEAE), 178

Dwarf plants, parasitic on the twigs of conifers; the slender, scattered or clustered stems from $\frac{1}{2}$ - $\frac{3}{4}$ inch in height.

LORANTHACEAE, 84

Low herbs or twining shrubs with large leaves, which are entire, petioled, and cordate; capsule 6-celled.

ARISTOLOCHIACEAE, 84

Low herbs with greenish-white flowers in umbel-like clusters; ovary 1-celled, forming a 1-seeded nutlet.

SANTALACEAE, 83

Shrubs with silvery-scurfy leaves; calyx 4-parted;
fruit fleshy or berry-like. ELAEAGNACEAE, 176

Trees with horizontal branches, and entire
obovate leaves, shiny when old; flowers
greenish; fruit ovoid, bluish-black.
NYSSA (CORNACEAE), 188

A. Both calyx and corolla present.

H. Corolla of separate petals.

I. Stamens more than 10 (rarely 9-10 in Polanisia),
and more than twice as many as the petals.

J. Calyx entirely free from the ovary (calyx
inferior).

K. Carpels several or many, wholly distinct or
slightly united at the base.

Aquatic herbs with peltate leaves.

BRASENIA (NYMPHAEACEAE), 101

Woody twiners with leaves peltate near the
edge; flowers in small whitish clusters;
fruit a black drupe with a crescent-shaped
seed. MENISPERMACEAE, 110

Herbs with usually lobed or much-dissected
leaves; stamens hypogynous. RANUNCULACEAE, 102

Herbs with alternate stipulate leaves; flowers
with numerous monadelphous stamens
forming a column united at the base to the
petals. MALVACEAE, 168

Plants with alternate, usually stipulate leaves;
stamens perigynous. ROSACEAE, 129

Trees or shrubs with smooth leaves notched
at the apex; flowers greenish-yellow, showy;
fruit an elongated dry cone. MAGNOLIACEAE, 109

Small trees with entire leaves; calyx of 3
sepals; corolla of 6 petals in 2 rows; fruit
long, dark brown, and edible. ANONACEAE, 110

K. Carpels one, or if more, united into a com-
pound pistil, as shown in the number of
its cells, placentae, styles, or stigmas.

L. Pistil of a single carpel (simple).

Herbs with 2-3-ternately compound leaves;
fruit a red or white berry.

ACTAEA (RANUNCULACEAE), 103

Herbs bearing 1 or 2 large, 5-9-parted
leaves, and a nodding white flower from
the fork; fruit a large, yellow, fleshy
berry.

PODOPHYLLUM (BERBERIDACEAE), 111

Trees or shrubs with simple, mostly serrate
leaves; fruit a fleshy edible drupe with
bony stone.

PRUNUS (ROSACEAE), 130

L. Pistil of several carpels united (compound).

Plants with entire dotted leaves; stamens
in 3 or more clusters.

HYPERICACEAE, 169

Herbs with milky or coloured juice; sepals 2
(rarely 3), caducous.

PAPAVERACEAE, 113

Herbs with palmate leaves of 3 leaflets;
flowers cruciform; stamens not tetradynamous.

POLANISIA (CAPPARIDACEAE), 123

Herbs or low shrubs with simple leaves, the
lower usually opposite, the upper alter-
nate; sepals 5, unequal.

CISTACEAE, 170

Bog plants with pitcher-shaped leaves.

SARRACENIACEAE, 124

Herbs with alternate stipulate leaves;
flowers with numerous monadelphous
stamens, forming a column united at the
base to the petals; pod 5-celled.

HIBISCUS (MALVACEAE), 168

Aquatic herbs with cordate leaves and
yellow or white flowers; petals numerous
and stamen-like.

NYMPHAEACEAE, 101

Trees with large, cordate, serrate leaves;
flowers yellowish, in small cymes on a
peduncle to which is attached a mem-
branous bract.

TILIACEAE, 167

J. Calyx united to the ovary (calyx superior).

Fleshy plants, which are globular, or flattened
and jointed, leafless and prickly.

CACTACEAE, 175

Herbs with fleshy entire leaves; calyx only partly adnate to the ovary; sepals 2, petals 5, stamens perigynous. PORTULACACEAE, 100

Shrubs or small trees with alternate stipulate leaves; stamens perigynous; fruit a pome. POMEAE (ROSACEAE), 131

I. Stamens not more than twice as many as the petals.

M. Stamens of the same number as the petals and opposite them.

Herbs with alternate leaves, stipulate or with dilated bases; sepals and petals usually in 2 rows of 3 each; fruit a berry or pod. BERBERIDACEAE, 111

Herbs with fleshy entire leaves; calyx inferior; sepals 2, petals 5. PORTULACACEAE, 100

Maritime herbs with simple radical leaves; flowers regular, pentamerous; calyx plaited; ovary 1-celled, 1-seeded. PLUMBAGINACEAE, 196

Leafy-stemmed plants with yellow regular flowers, which occasionally have separate petals. LYSIMACHIA (PRIMULACEAE), 197

Plants with alternate, stipulate, many-cleft leaves; stamens perigynous. CHAMAERHODOS (ROSACEAE), 130

Shrubs climbing by tendrils; calyx minute; ovary 2-4-celled; fruit a 2-celled berry. VITACEAE, 166

Shrubs with simple leaves; stamens perigynous; ovary 3-celled, 3-seeded; fruit a pod or drupe. RHAMNACEAE, 166

M. Stamens not of the same number as the petals, or if of the same number, alternate with them.

N. Corolla irregular.

Delicate smooth herbs with compound dissected leaves; corolla spurred or saccate at the base; stamens in two sets; ovary compound, 1-celled. FUMARIACEAE, 114

Herbs with 5 sepals, a 1-spurred or gibbous corolla of 5 petals; 5 hypogynous stamens and fruit a 1-celled, 3-valved pod. VIOLACEAE, 171

Succulent herbs with yellowish or orange flowers; sepals and petals coloured alike, one of former spurred; ovary 5-celled; fruit a pod bursting elastically. BALSAMINACEAE, 165

Plants with simple exstipulate leaves; the lower of 3 petals keel-shaped and often crested on the back; ovary forming a 2-celled, 2-seeded pod. POLYGALACEAE, 157

Herbs, shrubs, or trees with usually papilionaceous corollas; stamens usually 10, monadelphous, diadelphous, or rarely distinct; carpel 1-celled, forming a legume. LEGUMINOSAE, 141

N. Corolla regular or nearly so.

O. Calyx entirely free from the ovary (calyx inferior).

P. Stamens inserted on the receptacle (hypogynous).

Q. Leaves simple.

Herbs with pungent juice; petals 4, cruciform; stamens tetradynamous; fruit a silique or silicle (except in *Raphanus*). CRUCIFERAE, 115

Herbs or low shrubs with simple leaves, the lower usually opposite, the upper alternate; sepals 5, unequal. CISTACEAE, 170

Glandular-haired bog herbs with radical leaves; flowers in a one-sided raceme on a naked scape. DROSERACEAE, 124

Plants with entire dotted leaves; pod 1-celled, seeds numerous. HYPERICACEAE, 169

Herbs with stems usually swollen at the joints; leaves opposite and entire; styles 2-5. CARYOPHYLLACEAE, 95

Herbs with sessile, entire, exstipulate leaves and pentamerous flowers; stamens monadelphous at the base; pod 10-celled, 10-seeded. LINACEAE, 154

Plants with lobed or divided, stipulate leaves and pentamerous flowers; deeply lobed ovary with an elongated axis from which, at maturity, the 1-seeded carpels separate.

GERANIUM (GERANIACEAE), 155

Herbs or shrubs with leaves which are sometimes evergreen, but in one instance the entire plant is whitish or reddish; anthers opening by terminal pores.

ERICACEAE, 190

Small herbs with narrow radical leaves and naked, 1-flowered scapes, bearing small greenish flowers.

MYOSURUS (RANUNCULACEAE), 103

Q. Leaves compound.

Herbs with pungent juice; petals 4, cruciform; stamens tetradynamous; fruit a silicle or silique.

CRUCIFERAE, 115

Herbs with pinnate leaves and pentamerous flowers; ovary as in Geranium above.

ERODIUM (GERANIACEAE), 155

Low herbs with sour juice and palmate leaves with 3 obcordate leaflets; stamens 10; styles 5, distinct.

OXALIDACEAE, 155

Low herbs with small flowers solitary in the axils; leaves of 3-5 lanceolate leaflets; sepals and petals 3, stamens 6.

LIMNANTHACEAE, 161

Plants with ovate leaflets, downy when young; flowers greenish; fruit a 2-seeded samara or a pod.

RUTACEAE, 156

P. Stamens not hypogynous.

R. Stamens inserted on the calyx (perigynous) or inserted on a disk in the bottom of the calyx tube.

S. Leaves simple.

Herbs or shrubs with opposite and alternate, exstipulate leaves; styles

or stigmas 2 (4 in *Parnassia*);
 carpels fewer than the sepals;
 stamens perigynous. SAXIFRAGACEAE, 125

Small herbs with symmetrical flowers;
 sepals 4-5; carpels 5, united to
 form a 5-celled, 5-pointed capsule,
 or distinct. CRASSULACEAE, 125

Herbs or slightly woody plants with
 entire, mostly whorled leaves;
 flowers purple; calyx inclosing but
 free from the ovary; stamens
 perigynous; stigma 1. LYTHRACEAE, 177

Shrubs with leaves having minute,
 fugacious stipules; stamens on a
 calyx disk; fruit 3-5-lobed, seeds
 in a red aril. CELASTRACEAE, 163

Trees or shrubs with opposite, palm-
 ately lobed leaves (or compound
 in *A. Negundo*); fruit two 1-seeded
 samaras, which separate. ACERACEAE, 164

S. Leaves compound.

Shrubs with opposite, pinnate,
 stipulate leaves of 3-5 leaflets; a
 large disk in the bottom of the
 calyx tube; fruit a bladdery,
 inflated, 2-3-horned capsule.
 STAPHYLEACEAE, 164

Trees or shrubs with regular pen-
 tamerous flowers; stamens on a
 disk, styles or stigmas 3.
 ANACARDIACEAE, 162

R. Stamens attached at the base of the petals (epipetalous) or free from them in *Nemopanthus*.

Shrubs with small flowers, the parts
 mostly in 4's or 6's; fruit a red, 4-8-
 seeded, berry-like drupe. AQUIFOLIACEAE, 162

O. Calyx united to the ovary (calyx superior).

T. Stamens inserted on the calyx (perigy-
 nous).

Herbs with fleshy entire leaves; calyx partly adnate to the ovary, sepals 2, petals 5. PORTULACA (PORTULACACEAE), 100

Aquatic herbs; leaves crowded, often whorled and when immersed dissected; petals 4 (or none); stamens 4-8; stigmas 4; fruit nut-like, 4-celled, deeply 4-lobed.

MYRIOPHYLLUM (HALORAGIDACEAE), 181

Herbs with symmetrical perfect flowers; petals 2-4; stamens as many or twice as many as the petals; style 1, stigma 2-4-lobed or capitate. ONAGRACEAE, 178

Herbs with square, wing-angled stems and opposite, 3-7-ribbed leaves; the petals purplish; stamens 8; style 1, stigma 1. MELASTOMACEAE, 177

Tendrill-bearing herbs with small, whitish, monoecious flowers and prickly fruit. CUCURBITACEAE, 240

Herbs or shrubs with opposite and alternate, exstipulate leaves; styles or stigmas 2; carpels fewer than the sepals. SAXIFRAGACEAE, 125

Thorny shrubs or trees with simple stipulate leaves; petals on a disk in the calyx tube; stamens 5-25; fruit a reddish or yellowish pome containing 1-5 bony nutlets. CRATAEGUS (ROSACEAE), 131

Shrubs with alternate, wavy-toothed, simple leaves; flowers yellow, appearing in late autumn; petals 4, strap-shaped; stamens 8; styles 2, stigmas 2. HAMAMELIDACEAE, 129

T. Stamens not inserted on the calyx, but on the ovary or on a disk which covers the ovary (epigynous).

Herbs with small flowers chiefly in compound umbels; calyx minute, petals 5;

stamens 5; styles 2, surrounded by a fleshy disk; fruit dry, 2-seeded.

UMBELLIFERAE, 183

Plants with alternate compound leaves; flowers in simple umbels or panicles; styles usually more than 2; fruit a drupe.

ARALIACEAE, 182

Erect shrubs with 4-sided branchlets and opposite serrate leaves; fruit 3-5-lobed, seeds in a red aril.

EVONYMUS (CELASTRACEAE), 163

Shrubs or trees (rarely herbs) with simple leaves; petals 4 or 5; style 1; fruit a 1-2-seeded drupe.

CORNACEAE, 188

H. Corolla with petals more or less united.

U. Calyx entirely free from the ovary (calyx inferior).

V. Stamens more numerous than the lobes of the corolla.

Herbs, shrubs, or trees with usually papilionaceous corollas; stamens usually 10, monadelphous, diadelphous, or rarely distinct; carpel 1-celled forming a legume. LEGUMINOSAE, 141

Delicate smooth herbs with compound dissected leaves; corolla spurred or saccate at the base; ovary compound, 1-celled. FUMARIACEAE, 114

Herbs with alternate stipulate leaves; flowers with numerous monadelphous stamens united at the base to the petals. MALVACEAE, 168

Plants in some cases with evergreen leaves; in Rhododendron the stamens sometimes as few as the lobes of the corolla, anthers opening by terminal pores. ERICOIDEAE (ERICACEAE), 190

Plants with simple exstipulate leaves; the lower of 3 petals, keel-shaped and often crested on the back; ovary forming a 2-celled, 2-seeded pod. POLYGALACEAE, 157

Low herbs with sour juice; leaves palmate with 3 obcordate leaflets; stamens 10; styles 5, distinct. OXALIDACEAE, 155

V. Stamens not more numerous than the corolla lobes.

W. Stamens of the same number as the corolla lobes.

X. Stamens opposite the corolla lobes.

Herbs with simple leaves and regular flowers; stamens inserted on the corolla; style 1, stigma 1, ovary 1-celled.

PRIMULACEAE, 197

Maritime herbs with simple radical leaves and regular pentamerous flowers; calyx plaited; ovary 1-celled, 1-seeded.

PLUMBAGINACEAE, 196

X. Stamens alternate with the corolla lobes.

Y. Ovaries more than 1, or, if 1, deeply 4-lobed.

Plants with acrid milky juice and pentamerous and pentandrous flowers; anthers of the stamens slightly adherent to the stigma, filaments distinct; ovaries 2.

APOCYNACEAE, 202

Plants with milky juice and flowers in umbels; anthers of the stamens connected with the stigma, filaments united.

ASCLEPIADACEAE, 203

Herbs with square stems; opposite aromatic leaves and didynamous stamens or only 2; fruit consisting of 4 seed-like nutlets around the base of the style.

LABIATAE, 213

Usually rough-hairy herbs with alternate leaves and regularly 5-lobed corolla (except in *Echium*); stamens 5, epipetalous; fruit as in Labiatae.

BORAGINACEAE, 208

Y. Ovary only 1, not deeply lobed.

Z. Ovary 1-celled.

Hairy herbs with lobed or divided leaves and pentamerous and pentandrous flowers.

HYDROPHYLLACEAE, 207

Smooth herbs with bitter juice and simple leaves (in *Menyanthes* of 3 leaflets).

GENTIANACEAE, 200

Z. Ovary 2-10-celled.

Tall, usually woolly herbs with sessile leaves; corolla rotate, somewhat irregular, with 5 stamens inserted on its tube; ovary 2-celled.

VERBASCUM (SCROPHULARIACEAE), 223

Herbs with ribbed radical leaves and small spiked flowers on a naked scape; stamens 4 (rarely 2) attached to corolla; capsule 2-celled, the top coming off like a lid. PLANTAGINACEAE, 232

Herbs with regular, pentamerous and pentandrous flowers; corolla salverform; ovary 3-celled, style 3-lobed. POLEMONIACEAE, 206

Chiefly twining or trailing herbs with pentamerous and pentandrous flowers, capsule usually 2-celled, 2-4-seeded; style 2-cleft.

CONVOLVULACEAE, 205

Rank-scented herbs with regular pentamerous and pentandrous flowers (*Hyocyamus* somewhat irregular); style and stigma 1; fruit a 2-5-celled, many-seeded capsule or berry.

SOLANACEAE, 220

Plants with leaves which are sometimes evergreen (*Pterospora* is purplish-brown); (petals nearly separate in *Ledum*); anthers opening by terminal pores or longitudinal slits; style 1, capsule 5-celled.

ERICACEAE, 190

Shrubs having small flowers with parts mostly 4-6; stamens attached to the base of the petals or free, style 1; fruit a red, 4-6-seeded, berry-like drupe.

AQUIFOLIACEAE, 162

W. Stamens fewer than the corolla lobes.

AA. Corolla regular.

Plants with opposite leaves and somewhat
unequally 5-lobed corolla; stamens
didynamous; ovary 4-celled but not 4-
lobed. VERBENACEAE, 212

Herbs with ribbed radical leaves and
small spiked flowers on a naked scape;
stamens 4 (rarely 2); capsule 2-celled,
top coming off like a lid. PLANTAGINACEAE, 232

Herbs with square stems and opposite
aromatic leaves; stamens 2; fruit con-
sisting of 4 seed-like nutlets around the
base of the style. LYCOPUS (LABIATAE), 213

AA. Corolla irregular.

Herbs with square stems and opposite
aromatic leaves; stamens didynamous
or only 2; fruit consisting of 4 seed-like
nutlets around the base of the style.
LABIATAE, 213

Herbs growing in water or wet places;
calyx and corolla 2-lipped, the latter
personate and spurred at the base;
stamens 2. LENTIBULARIACEAE, 230

Herbs parasitic on roots and destitute of
green foliage; stamens didynamous;
corolla ringent and more or less 2-lipped.
OROBANCHACEAE, 231

Herbs with 2-lipped corolla (nearly regular
in Veronica); stamens 4, didynamous or
rarely 2. SCROPHULARIACEAE, 223

Herbs growing in water or wet places;
leaves opposite, entire; flowers purplish;
calyx 5-cleft; corolla 2-lipped; stamens 2.
ACANTHACEAE, 232

Herbs with coarsely toothed, ovate leaves;
calyx and corolla 2-lipped, the latter
purplish or rose-coloured. PHRYMACEAE, 232

U. Calyx united to the ovary (calyx superior).

BB. Stamens more or less united by their anthers.

Tendrill-bearing herbs with small, whitish, monoecious flowers and prickly fruit. CUCURBITACEAE, 240

Flowers in a dense head on a common receptacle, surrounded by an involucre; fruit a seed-like achene. COMPOSITAE, 243

Herbs with acrid milky juice and scattered flowers; corolla irregular, 5-lobed and split down one side. LOBELIACEAE, 242

BB. Stamens not united together in any way.

CC. Stamens inserted on the corolla (epipetalous).

Coarse prickly herbs with opposite or whorled, exstipulate leaves; flowers in dense involucre heads; stamens 4. DIPSACACEAE, 240

Herbs with opposite exstipulate leaves; flowers in cymose clusters; calyx limb of plumose bristles; stamens 1-3, fewer than the corolla lobes. VALERIANACEAE, 239

Herbs or shrubs with opposite stipulate leaves or whorled exstipulate leaves; stamens alternate with the mostly 4 lobes of the corolla. RUBIACEAE, 233

Shrubs, rarely herbs, with opposite exstipulate leaves (*Viburnum* sometimes with appendages which are evidently stipules); corolla tubular or rotate. CAPRIFOLIACEAE, 236

CC. Stamens free from the corolla or nearly so.

Herbs with milky juice and scattered flowers; corolla regular, 5-lobed; stamens 5. CAMPANULACEAE, 241

Somewhat shrubby plants; anthers opening by pores; an edible, berry-like fruit. ERICACEAE, 190

FLORA

DIVISION III. PTERIDOPHYTES*

(Ferns and Fern Allies)

Plants not producing true flowers, but reproducing themselves by means of spores instead of seeds, the spores consisting merely of simple cells, and not containing an embryo.

POLYPODIA'CEAE (FERN FAMILY)

Leafy plants with rootstocks. Spores borne in sporangia (spore cases), these collected into various kinds of clusters called sori. Each sporangium encircled by an elastic jointed ring. Sori usually covered by a delicate membrane called the indusium.

Synopsis of the Genera

1. **Polypo'dium.** Fruit-dots round, attached to the back of the fronds on the ends of the free veins. No indusium.
2. **Adian'tum.** Fruit-dots marginal, the edge of the frond being reflexed so as to form an indusium. Midrib of the pinnules close to the lower edge or altogether wanting. Stipe black and shining. All the pinnules distinct and generally minutely stalked. Veins free.
3. **Pte'ris.** Fruit-dots marginal. Indusium formed by the reflexed edge of the frond. Midrib of the pinnules in the centre and prominent. Veins free. Stipe light-coloured.
4. **Pellae'a.** Fruit-dots marginal, covered by a broad indusium, formed by the reflexed margin of the frond. Small ferns with once- or twice-pinnate fronds, the fertile ones very much like the sterile, but with narrower divisions. Stipe shining, purple.
5. **Cryptogram'ma.** Fruit-dots roundish or elongated, attached to the free-forking veins. Margins of the fertile segments at first reflexed and meeting at the midrib, but opening out at maturity. Low ferns, with smooth, 2-3-pinnate, tufted fronds; the fertile ones taller and with narrower divisions.

*Where the accent immediately follows the vowel, the vowel is pronounced long; where it follows the consonant, the vowel in the accented syllable is short.

6. **Asple'nium.** Fruit-dots elongated (rarely curved), on veins on the back of the pinnules, oblique to the midrib, but only on the upper side of the vein (rarely attached to both sides). Indusium attached to the vein by one edge, the other edge free. Veins free.
7. **Woodwar'dia.** Fruit-dots elongated, on cross-veins parallel to the midrib, forming a chain-like row on each side of the latter. Indusium as in the last. Veins reticulated.
8. **Scolopen'drium.** Fruit-dots elongated, occurring in pairs on contiguous veinlets, the free edges of the two indusia facing each other, so that the sori appear to be single, with an indusium split down the centre. Veins free. Frond simple, ribbon-shaped, about an inch broad, generally wavy-margined.
9. **Camptoso'rus.** Fruit-dots elongated, those near the base of the midrib double, as in Scolopendrium; others single, as in Asplenium. Fronds simple, $\frac{1}{2}$ to $\frac{3}{4}$ inch wide at the heart-shaped base, and tapering into a long and narrow point; growing in tufts on limestone rocks, and commonly rooting at the tip of the frond, like a runner. Veins reticulated.
10. **Polys'tichum.** Fruit-dots orbicular, covered by a circular indusium on a centrally placed stalk. Fronds tufted, chiefly firm or leathery in texture.
11. **Phegop'teris.** Fruit-dots roundish, borne on the back of the veinlet below the apex, rather small. Indusium none. Veins free. Fronds triangular in outline, in one species twice-pinnatifid, with a winged rhachis, and in the other in three petioled, spreading divisions, the divisions once- or twice-pinnate.
12. **Aspid'ium.** Fruit-dots round. Indusium evident, flat, orbicular or kidney-shaped, fixed by the centre, opening all around the margin. Veins free. Generally rather large ferns with tufted fronds from once- to thrice-pinnate.
13. **Cystop'teris.** Fruit-dots round. Indusium not depressed in the centre, but rather raised, attached to the frond not by the centre, but by the edge partly under the fruit-dot, and generally breaking away on the side towards the apex of the pinnule, and becoming reflexed as the sporangia ripen. Fronds slender and delicate, twice- or thrice-pinnate.
14. **Onocle'a.** Round fruit-dots partially covered by delicate hood-shaped indusia. Fertile fronds erect, rigid, with contracted, pod-like divisions. Sterile fronds leaf-like.
15. **Wood'sia.** Small tufted ferns. Fruit-dots round. Indusium very thin and delicate, attached by its base all around under

the sorus, the top at length bursting into more or less narrow segments. Veins free.

16. **Dickso'nia**. Fruit-dots round, very small, each on a recurved toothlet on the upper margin of the lobes of the pinnules usually one to each lobe. Sporangia on an elevated globular receptacle, and inclosed in a cup-shaped indusium open at the top and partly adherent to the reflexed toothlet of the frond. Fronds minutely glandular or hairy, 2-3 feet high, ovate-lanceolate in outline, pale green, very thin, without chaff.

1. POLYPO'DIUM L. POLYPODY

1. **P. vulga're** L. Fronds evergreen, 4-10 inches long, deeply pinnatifid, the lobes obtuse and obscurely toothed. Sori large, midway between the margin and the midrib.—Common on shady rocks.

2. ADIAN'TUM L. MAIDENHAIR

1. **A. peda'tum** L. Stipe upright, black and shining. The frond forked at the top of the stipe, the two branches of the fork recurved, and each bearing on its inner side several slender, spreading divisions, the latter with numerous thin, pinnatifid pinnules which look like the halves of pinnules, owing to the midrib being close to the lower edge. Upper margin of the pinnules cleft.—Common in rich woods.

3. PTE'RIS L. BRAKE. BRACKEN

1. **P. aquili'na** L. Stipe stout and erect. Frond large and divided into 3 large, spreading divisions at the summit of the stipe, the branches twice-pinnate, the lower pinnules more or less pinnatifid, the lobes margined all around with the indusium.—Common in thickets and on dry hillsides.

4. PELLAE'A Link. CLIFF BRAKE

1. **P. atropurpu'rea** Link. Small ferns, 6-15 inches high, the stipe dark purple and shining. Frond pale, once- or (below) twice-pinnate, the divisions broadly linear or oblong (or the sterile sometimes oval), stalked at the base. Stipes clustered.—Dry rocks.

5. CRYPTOGRAM'MA R. Br. ROCK BRAKE

1. **C. acrostichoi'des** R.Br. Fertile fronds with linear segments; sterile fronds with ovate-oblong, serrulate segments. Stipes straw-coloured and scaly.—Rock crevices.

2. *C. Stelle'ri* Prantl. Fronds 3-6 inches high, slender, of few pinnae, the lower ones once- or twice-pinnatifid into 3-5 divisions, those of the fertile fronds narrower than those of the sterile ones. Stipe polished, brownish, darker at the base.—Shady limestone rocks; not common.

3. *C. den'sa* Diels. Fronds 3-8 inches high, nearly alike; segments of the sterile fronds lance-linear; stipes purplish brown.—Vertical limestone cliffs.

6. ASPLE'NIUM L. SPLEENWORT

A. Indusium straight, attached to the upper side of the vein.

1. *A. Trichom'anes* L. (MAIDENHAIR SPLEENWORT.) A very delicate little fern growing in tufts. Fronds 3-6 inches long, linear in outline, pinnate, the little pinnae oval and unequal-sided, about $\frac{1}{5}$ inch long. The stipes thread-like, purplish-brown and shining. This species is evergreen.—Shaded rocks.

2. *A. vir'ide* Hudson. Resembling *A. Trichomanes*, but less rigid, and the stipe brownish at the base, becoming green upwards. Pinnae roundish-ovate or ovate-rhomboid, short-stalked, crenately toothed.—Shaded limestone.

3. *A. platyneu'ron* Oakes. Evergreen. Fronds 9-18 inches high, upright, linear, oblanceolate in outline, fertile ones much the taller, simply pinnate. Pinnae mostly alternate, sessile, finely serrate, the base auricled on the upper or both sides. Stipe and rhachis blackish-purple and shining.—Rocky open woods.

4. *A. acrostichoi'des* Sw. Not evergreen. Fronds 2-3 feet high, pinnate, the pinnae linear-lanceolate in outline, 3-5 inches long, deeply pinnatifid, each of the crowded lobes bearing 3-6 pairs of oblong fruit-dots.—Rich woods.

5. *A. angustifo'lium* Michx. Not evergreen. Fronds simply pinnate, somewhat resembling *Polystichum acrostichoides*, but very smooth and thin, and larger. Pinnae entire or crenulate, short-stalked. Fruit-dots linear, crowded.—Rich woods; not common.

B. Indusium curved, delicate, often crossing the vein, thus appearing reniform.

6. *A. Filix - fe'mina* Bernh. (LADY FERN.) Fronds 1-3 feet high, broadly lanceolate in outline, twice-pinnate, the pinnae lanceolate

in outline, and the pinnules confluent by a narrow margin on the rhachis of the pinna, doubly serrate. Indusium curved, often shaped something like a horse-shoe, owing to its crossing the vein and becoming attached to both sides of it.—Rich woods.

7. WOODWARDIA Smith. CHAIN FERN

1. *W. virginica* Smith. Fronds 2–3 feet high, pinnate; pinnae lanceolate, pinnatifid. Veins forming a single row of meshes next the midrib.—Wet swamps.

8. SCOLOPENDRIUM Adans. HART'S TONGUE

1. *S. vulgare* Smith. Frond simple, with an auricled, heart-shaped base, bright green, a foot or more in length, and an inch or more in width.—Shaded ravines and limestone cliffs; not very common.

9. CAMPTOSORUS Link. WALKING-LEAF

1. *C. rhizophyllus* Link. A curious little fern, growing in tufts on shaded limestone rocks. Frond simple, tapering, often rooting at the tip.—Not very common.

10. POLYSTICHUM Roth.

1. *P. acrostichoides* Schott. (CHRISTMAS FERN.) Fronds narrowly oblong or lanceolate, simply pinnate; pinnae linear-lanceolate with a pointed lobe on the upper side of the slightly stalked base. At maturity the fruit-dots are spread over the entire surface of the pinnae.—Rich woods.

2. *P. Lonchites* Roth. Not unlike No. 1, but the fronds are narrower, and longer, more rigid and with hardly any stipe. Pinnae densely spinulose-toothed.—Rocky woods; not common.

3. *P. Braunii* Fée. Fronds twice-pinnate, oblong-lanceolate, narrowing gradually towards the base. Pinnules ovate or oblong, truncate and nearly rectangular at the base (the lower short-stalked), beset with long and soft as well as chaffy hairs. Indusium fixed by the centre. Fronds evergreen, very chaffy on the stalk and rhachis.—Ravines and deep woods, Atl. Prov.

11. PHEGopteris Fée. BEECH FERN

1. *P. polypodioides* Fée. Fronds triangular, longer than broad, 4–6 inches long, hairy on the veins, twice-pinnatifid, the rhachis winged. The pinnae sessile, linear-lanceolate in outline,

the lowest pair deflexed and standing forwards. Fruit-dots small and all near the margin. Stipes rather longer than the fronds, from a slender creeping rootstock.—Rich woods.

2. *P. hexagonop'tera* Fée. Fronds triangular, generally broader than long, 7–12 inches broad. Pinnæ lanceolate; the lowest very large, their divisions elongated and pinnatifid, the basal divisions decurrent on the main rhachis and forming a many-angled wing. Fruit-dots not exclusively near the margin.—Rich woods.

3. *P. Dryop'teris* Fée. (OAK FERN.) Fronds broadly triangular in outline, primarily divided into 3 triangular, spreading, petioled divisions, smooth, the three divisions once- or twice-pinnate. Fronds from 4 to 6 inches wide. Fruit-dots near the margin. The whole plant delicate, and light green in colour.—Rich woods, common.

4. *P. Robertia'na* A. Br. Closely resembling *P. Dryopteris*, but differs in that the fronds are minutely glandular and somewhat rigid, and the lowest pinnæ on the lower side of the lateral divisions are proportionally smaller.—Rocky hillsides, Ont. and Pr. Prov.

12. ASPID'IUM Swartz. SHIELD FERN. WOOD FERN

A. *Stipes not chaffy.*

1. *A. Thelyp'teris* Swartz. Fronds tall and narrow, lanceolate in outline, pinnate, the pinnæ deeply pinnatifid, nearly at right angles to the rhachis, linear-lanceolate in outline, the margins of the lobes strongly revolute in fruit. Stipe over a foot long, and usually longer than the frond.—Common in low wet places.

2. *A. noveboracen'se* Swartz. Fronds much lighter in colour than the preceding, tapering towards both ends, pinnate, the pinnæ deeply pinnatifid, much closer together than in No. 1, and not at right angles with the rhachis. Veins simple. Lower pinnæ short and deflexed.—Swamps and thickets.

B. *Stipes chaffy.*

3. *A. spinulo'sum* Swartz. Stipes slightly chaffy or scaly. Fronds large, ovate-lanceolate in outline, twice-pinnate, the pinnules deeply pinnatifid (nearly pinnate) and spiny-toothed. Pinnæ triangular-lanceolate in outline.

Var. *intermedium* D. C. Eaton. Margin of the indusium denticulate and beset with minute stalked glands, the few scales of the stipes pale brown with a dark centre at the base, and the lower pinnae unequal-sided.—Very common in Canadian woods.

Var. *dilatatum* Hook. Scales of the stipe brown with a dark centre at the base, the frond broader, ovate or triangular-ovate in outline, often twice-pinnate, and the indusium smooth and naked.

4. *A. Boottii* Tuckerman. Fronds elongated-oblong or elongated-lanceolate, somewhat narrowed at the base; indusium minutely glandular; scales of the stipe pale brown.

5. *A. fragrans* Swartz. Fronds 4–12 inches high, fragrant, narrowly lanceolate, with narrowly oblong, pinnately parted pinnae, their divisions nearly covered beneath by very large, thin indusia.—Rocks, Atl. Prov. and northward.

6. *A. cristatum* Swartz. Stipes chaffy with broad scales. Fronds large, linear-lanceolate in outline, pinnate, the pinnae deeply pinnatifid, the upper ones triangular-lanceolate in outline, the lower considerably broader, the lobes cut-toothed. Fruit-dots large and conspicuous, half-way between the midrib of the lobe and the margin.—Swamps.

Var. *Clintonia* D. C. Eaton. Distinguished chiefly by its greater size and more numerous pinnae and segments.

7. *A. Goldianum* Hook. A fine fern, 2–4 feet in height, the large fronds growing in a circular cluster from a chaffy rootstock. Frond ovate or ovate-oblong in outline, pinnate, the pinnae deeply pinnatifid, 6–9 inches long, broadest in the middle, the lobes slightly scythe-shaped, finely serrate. Fruit-dots large, near the midrib of the lobe.—Rich moist woods.

8. *A. Filix-mas* Swartz. (MALE FERN.) Fronds lanceolate, very chaffy at the base, twice-pinnate except that the upper pinnules run together. Pinnae linear-lanceolate, tapering from the base to the summit. Pinnules very obtuse, the basal ones incisely lobed. Fruit-dots rather closer to the midvein than to the margin. Indusium convex, persistent.—Rocky woods.

9. *A. margin'le* Swartz. Stipes very chaffy at the base. Fronds ovate-oblong in outline, twice-pinnate, the pinnae lanceolate in outline, broadest above the base. Pinnules crenate-margined. Fruit-dots large, close to the margin.—Rich woods, mostly on hill-sides.

13. CYSTOP'TERIS Bernh. BLADDER FERN

1. *C. bulbif'era* Bernh. Frond lanceolate-attenuate (1-2 feet), narrow and very delicate, twice-pinnate, the pinnae nearly at right angles to the rhachis. Rhachis and pinnae usually with bulblets beneath. Pinnules toothed.—Shady moist ravines in calcareous soil.

2. *C. frag'ilis* Bernh. Frond only 4-8 inches long, with a stipe of the same length, 2-3-pinnate. Rhachis winged.—Shady cliffs.

14. ONOCLE'A L.

1. *O. sensib'ilis* L. (SENSITIVE FERN.) Fronds scattered; sterile long-stalked, pinnatifid; fertile contracted, bipinnate, the pinnules rolled into berry-like bodies.—Common in wet grounds.

2. *O. Struthiop'teris* Hoffm. (OSTRICH FERN.) Sterile fronds with lower pinnae gradually much shorter than the upper ones; pinnae deeply pinnatifid. Fertile fronds with pod-like or necklace-shaped pinnae.—Common in low wet grounds.

15. WOOD'SIA R. Brown

1. *W. ilven'sis* R. Brown. (RUSTY WOODSIA.) Stalks indistinctly jointed at some distance above the base. Fronds oblong-lanceolate, 2-6 inches long, rather smooth and green above, thickly clothed below with bristly rusty chaff, pinnate; the pinnae oblong, blunt, sessile, pinnately parted with the segments indistinctly crenate. Fruit-dots near the margin.—Exposed rocks, Atl. Prov. and northward.

2. *W. alpi'na* S. F. Gray. Stalks jointed. Fronds narrowly oblong-lanceolate. Sparingly hairy beneath with chaffy hairs, pinnate; the pinnae triangular-ovate, pinnately lobed, the lobes few and almost entire.—Ravines, Atl. Prov. and northward.

3. *W. glabel'la* R. Brown. Stalks jointed as in the two preceding species. Frond linear, very delicate, smooth, pinnate. Pinnae roundish-ovate, the lower somewhat distant, crenately lobed.—Moist rocks, Atl. Prov. and northward.

4. *W. obtu'sa* Torr. Stalks not jointed. Frond broadly lanceolate, beset with small glandular hairs, once- or nearly twice-pinnate. Pinnae pinnately parted. Segments of pinnae crenately toothed.—Cliffs and rocky places; rare.

5. *W. scopuli'na* D. C. Eaton. Like No. 4 but has shorter and smoother fronds with shortened lower pinnae. The indusium, which consists of hair-like divisions, is hidden below the sorus.

16. DICKSONIA L'Her. DICKSONIA

1. *D. punctilob'ula* Gray. Pleasantly odorous, hay-scented.—Moist shady places.

OSMUNDA'CEAE (FLOWERING FERN FAMILY)

Sporangia naked, globular, pedicelled, reticulated, with no ring or with mere traces of one, and opening by a longitudinal slit. Rootstocks creeping.

1. OSMUNDA L. FLOWERING FERN

Fertile fronds or fertile portions of the frond much contracted, bearing naked sporangia, which are globular, short-pedicelled, and opening by a longitudinal slit to discharge the spores. Frond tall and upright, 1-2-pinnate, from thick rootstocks.

1. *O. rega'lis* L. (FLOWERING FERN.) Fronds twice-pinnate, fertile at the top, very smooth, pale green. Sterile pinnules oblong-oval, finely serrate towards the apex, 1-2 inches long, either sessile or short-stalked, usually oblique and truncate at the base.—Swamps along streams and lake-margins.

2. *O. Claytonia'na* L. (INTERRUPTED FERN.) Sterile fronds large, pinnate, pale green, densely white-woolly when unfolding from the bud; a few of the middle pinnae fertile. Sterile pinnae deeply pinnatifid, the lobes entire; fertile pinnae entirely pinnate. Fertile fronds taller than the sterile; sporangia greenish.—Low grounds.

3. *O. cinnamo'mea* L. (CINNAMON FERN.) Fertile fronds distinct from the sterile ones, and appearing earlier, contracted, twice-pinnate, covered with cinnamon-coloured sporangia. Sterile fronds taller, rusty-woolly when young, smooth afterwards, once-pinnate, the pinnae deeply pinnatifid. The long sterile fronds in a cluster, with the fertile ones in the centre.—Low grounds.

OPHIOGLOSSA'CEAE (ADDER'S TONGUE FAMILY)

Sporangia naked, not reticulated, opening by a transverse slit. Roostocks erect.

Synopsis of the Genera

1. *Botrych'ium*. Sporangia in compound spikes, distinct in double rows, and opening by a transverse slit. Sterile part of the frond compound. Veins free.
2. *Ophioglos'sum*. Sporangia coherent in 2 ranks on the edges of a simple spike. Sterile part of the frond simple. Veins reticulated.

1. BOTRYCH'TUM Swartz. MOONWORT. GRAPE FERN

A. *Sterile portion of the frond not fleshy.*

1. *B. virginia'num* Swartz. Fronds tall, in some cases from a common rootstock; sterile segment sessile above the middle of the plant, broadly triangular, ternate, these divisions 1-2-pinnate, then once- or twice-pinnatifid; fertile segment 2-3-pinnate.—Rich woods, everywhere.

B. *Sterile portion of the frond more or less fleshy.*(1) *Sterile segment sessile or nearly so.*

2. *B. Luna'ria* Swartz. Sterile segment very fleshy, nearly sessile, oblong, simply pinnate, with 5-15 lunate or fan-shaped, fleshy divisions. Fertile segment 2-3-pinnate.—Rare.

3. *B. sim'plex* Hitchcock. Fronds small, seldom 6 inches high, the sterile segment borne near the middle of the plant, sometimes at the summit or near the base, short-petioled, fleshy, simple, and roundish, or pinnately lobed, with roundish lobes decurrent on the broad and flat, indeterminate rhachis. Veins all forking from the base.

4. *B. lanceola'tum* Angstroem, var. *angustisegmen'tum* Pease and Moore. Frond 3-10 inches high. Sterile part closely sessile at the top of the slender common stalk, very slightly fleshy, triangular, ternately twice-pinnatifid, with acute, toothed lobes. Veinlets branching from the continuous midvein. The fertile part twice- or thrice-pinnate.—Atl. Prov. and Ont.

5. *B. ramo'sum* Aschers. Resembling the preceding, but the sterile segment not quite sessile, somewhat fleshy, pinnate to

twice-pinnatifid, with obtuse lobes. Midvein broken up into forking veinlets.—Damp woods.

(2) *Sterile segment long-stalked.*

6. *B. obliquum* Muhl. Fleshy, 4–16 inches high, sparingly hairy. Sterile segment attached near the base of the plant, broadly triangular, variously decompound, with stalked divisions. Fertile segment erect, 2–4-pinnate.—Not uncommon. Pastures.

7. *B. ternatum* Swartz, var. *intermedium* D.C. Eaton. Stout, very fleshy, nearly glabrous. Habit much like the preceding. Sterile frond large, somewhat triangular with numerous usually obtuse divisions.—Sandy pastures and open woods.

2. OPHIOGLOS'SUM L. ADDER'S TONGUE

1. *O. vulgatum* L. Sterile part of the frond ovate or elliptical-oblong, 2–3 inches long, rather fleshy, sessile, near the middle of the stalk; the latter 6–12 inches high. Midvein indistinct or none, the main veins forming a loose network.—Bogs and grassy meadows.

MARSILEA'CEAE

Plants rooted in mud, with slender creeping rootstocks. Sporocarps borne on peduncles which rise from the rhizome near the insertion of the leaves.

1. MARSIL'EA L.

Leaves compound of 4 leaflets. Sporocarps with 2 teeth near the base, 2-celled.

1. *M. vestita* Hook and Grev. Leaves compound of 4 horizontal leaflets, which are broadly wedge-shaped. Peduncles very short, free from the petiole. Sporocarp solitary, about $\frac{1}{5}$ inch long.

SALVINIA'CEAE

Small, floating, moss-like plants with branching axis, covered with minute, 2-lobed, imbricated leaves. Sporocarps in pairs beneath the stem, very soft and thin-walled, the smaller ones acorn-shaped, containing a single macrospore, the larger ones globose, containing several microspores.

1. AZOL'LA Lam.

1. *A. carolinia'na* Willd. Somewhat deltoid in outline, much branched, of reddish aspect. Floating on quiet waters of Lake Ontario, not common.

EQUISETA'CEAE (HORSETAIL FAMILY)

Somewhat rush-like plants from branching rootstocks; the striated stems usually hollow, jointed and sheathed at the joints; the fertile stem terminated by a conical or spike-like, spore-bearing body.

1. EQUISE'TUM L. HORSETAIL

Rootstocks usually blackish, often tuber-bearing, with toothed sheaths at the nodes. Stem erect, simple or branched, jointed ridged and covered with a coat of silex; nodes solid, sheathed; internodes mostly with a large central cavity, a smaller one under each groove (vallecular) and a still smaller one under each ridge (carinal). Branches, when present, in whorls from the nodes. Sporangia borne on stalked sporophylls in a terminal cone. Spores numerous, round, green, and each provided with 4 hygroscopic elaters.

A. Annual-stemmed, not surviving the winter.

(1) *Fruiting in spring, the fertile stems pale or brownish, and earlier than the herbaceous sterile stems or branches.*

1. *E. arven'se* L. (COMMON HORSETAIL.) Fertile stems not branching, 4-10 inches high, pale brown, sheaths 8-12-toothed; sterile stems slender, taller, sometimes prostrate with many 4-angular, solid, green branches.—Moist sandy or gravelly soil, common.

2. *E. pratense* Ehrh. Stems more slender, with 3-sided, simple branches shorter than in *E. arvense*. Stem-sheaths with short, ovate-lanceolate teeth (those of the branches 3-toothed.) The fertile stems produce branches, when older, except at the top, which perishes after fructification.—Low meadows, Atl. Prov. and northward.

3. *E. sylvat'icum* L. Fertile stems branching. Branches compound, with loose sheaths, those of the stem having 8-14 somewhat blunt teeth, while those of the branches have 4-5 (of the branchlets 3) lance-pointed, diverging teeth. Top withering away after fructification.—Damp shady places, common.

(2) *Fruiting in summer. Stems all similar, or the fertile contemporaneous with the sterile ones.*

4. *E. palus'tre* L. Stems 10–18 inches high, slender, very deeply 5–9-grooved, ridges narrow, the lance-awl-shaped teeth whitish-margined, branches hollow, 4–7-angled.—Atl. Prov. and northward.

5. *E. fluviat'ile* L. (PIPES.) Stem 1–5 feet high, slightly many-furrowed, simple or branched near the middle, branches hollow, 4–6-angled. Sheaths appressed, with 10–22 dark-brown, acute, short, rigid teeth.—In shallow water or in mud.

B. Stems evergreen (except No. 7), all similar, not branching or only slightly so.

6. *E. hyema'le* L. (SCOURING RUSH.) Stems stout and tall, 1–3 feet high. Sheaths elongated, tight, with two black rings separated by an ashy one, and about 20 narrow linear teeth, 1-keeled at the base, and with awl-shaped, deciduous points.—Wet banks.

7. *E. laeviga'tum* A. Br. Stems 1–4 feet high, rather slender, pale green, 14–30-grooved, the ridges almost smooth. Sheath slightly enlarged upward, with a black girdle at the base of the white-margined teeth.—Pr. Prov.

8. *E. variega'tum* Schleicher, var. *jesu'pi* A. A. Eaton. Stem slender, in tufts, with 5–10 grooves, ascending, 6–18 inches high. Sheaths loose, green, variegated with black above, 5–10-toothed; teeth black, with white borders and long tips that wither.—Shores and wet banks.

9. *E. scirpoi'des* Michx. Stems slender, very numerous in a tuft, filiform, 3–6 inches high, curving, mostly 6-grooved, solid at the centre. Sheaths 3-toothed.—Wooded hillsides and moist banks.

LYCOPODIA'CEAE (CLUB-MOSS FAMILY)

Chiefly moss-like and perennial, evergreen plants; often with long, running, branching stems and crowded, lanceolate or awl-shaped, persistent, entire leaves; the sporangia 1-celled, solitary in the axils of the leaves. Spores all of one kind.

1. LYCOPO'DIUM L. CLUB-MOSS

Spore cases in the axils of the ordinary leaves or in those of the leaves forming a fruiting spike. Spores in the form of a yellow, inflammable powder.

A. Spore-cases in the axils of the ordinary dark-green, rigid, lanceolate leaves.

1. *L. Sela'go* L. Rootstock short and slender. Stems erect and rigid, forking, forming a level-topped cluster. Leaves uniform, glossy, pale green, sharp.—Atl. Prov. and northward.

2. *L. lucid'ulum* Michx. Stems less rigid than the last, forking and ascending. Leaves at first spreading, then deflexed, arranged in alternate zones of longer and shorter leaves, the latter usually bearing sporangia in their axils.

B. Spore-cases only in the axils of the upper bract-like leaves, forming a spike.

(1) *Leaves of the spike not very different from those of the stem.*

3. *L. inunda'tum* L. A low plant with weak, creeping, sterile stems, and solitary, erect, fertile stems 1-5 inches high, bearing a short thick spike. Stem-leaves lanceolate, acute, soft, spreading, mostly entire, those of the spike closely resembling them.—Sandy bogs.

Var. *Bigelo'vii* Tuckerm. Stems taller and leaves more upright and denticulate.—Atl. Prov.

(2) *Leaves of the catkin-like spike quite different from those of the stem.*

4. *L. annot'inum* L. Stems creeping, 1-4 feet long. Branches 4-9 inches high, once- or twice-forked. Spike solitary, sessile, the leaves of it yellowish and scale-like, ovate or heart-shaped, the others spreading or reflexed, rigid, pointed, nearly entire, pale green.—Cold woods.

5. *L. obscu'rum* L. (GROUND PINE.) Rootstock creeping underground, nearly leafless. Stems much resembling little hemlocks, 6-9 inches high; numerous fan-like, spreading branches with shining, lanceolate, entire leaves. Spikes sessile, nearly as in No. 4, 1-3 on each plant.—Moist woods.

Var. *dendroi'deum* D.C. Eaton. Similar to the species but fruiting spikes 4-10 on each plant.

6. *L. clava'tum* L. (CLUB-MOSS.) Stem creeping or running extensively. Spikes mostly in pairs, sometimes in fours or single, raised on a slender peduncle (4-6 inches long). Leaves linear, awl-shaped, bristle-tipped.—Dry woods.

7. *L. complana'tum* L. Stem creeping extensively, near the surface. Branches flattened, fan-like and spreading, forking above, the branchlets crowded. Leaves awl-shaped, small, in 4 ranks. Spikes 2-4, cylindrical, on a slender peduncle.—Dry woods; mostly with evergreens.

8. *L. trista'chyum* Pursh. Stem pale glaucous. Rootstock deeper than No. 7. Branches more numerous, erect, narrower. Peduncles bear 1-many spikes.—Dry rocky or sandy soil northward.

SELAGINELLA'CEAE

Small leafy plants, terrestrial. Stem branching and short. Leaves 4-6-ranked. Spore-cases 1-celled, solitary, axillary or borne on the upper surface of the leaf at its base and enwrapped in its margins. Spores of two kinds, the large ones (macrospores) developing prothallia which bear archegonia only, and the small ones (microspores) giving rise to antheridia only.

1. SELAGINEL'LA Beauv.

Terrestrial. Stem slender. Leaves small. Spore-cases very small, axillary, some containing the orange-coloured, powdery microspores, and others with 3 or 4 larger macrospores.

1. *S. rupe'tris*. Spring. A little, moss-like, branched, rigid evergreen growing in dense tufts 1-3 inches high. Leaves appressed, similar, awl-shaped, with a grooved keel and tipped with a bristle. Spikes 4-cornered.—Exposed dry rocks.

2. *S. selaginoi'des* Link. Sterile stems prostrate, small and slender, 1-2 inches high; the fertile larger, taller, and unbranched. Leaves lanceolate, spreading and somewhat larger on the fertile stems.

3. *S. a'pus* Spring. Stems tufted, creeping, branched, not rigid. Leaves of two kinds, the larger spreading and ovate, the smaller appressed and lanceolate.—Low shady places.

ISOËTA'CEAE (QUILLWORT FAMILY)

Small aquatic plants with corm-like stem, from which grow a cluster of elongated, taper-pointed leaves. Spores of two kinds in axillary sporangia, in the dilated bases of the leaves.

1. ISO'ËTES L. QUILLWORT

Growing in water. Stem corm-like. Leaves elongated and rush-like. Spore-cases large, enwrapped by the spreading bases of the leaves.

1. *I. echinos'pora* Durieu, var. *Braun'ii* Engelm. (QUILLWORT.) A small, aquatic, grass-like plant, with a corm-like stem, bearing 15-30 slender leaves dilated at the base. The sporangia axillary, partly enwrapped by the thin edges of the excavated bases of the leaves, beset with small spinules.—Lakes and ponds.

DIVISION IV. SPERMATOPHY'TES

Plants with perfect flowers or with either stamens or pistil developed. Usually reproduced by seeds.

SUBDIVISION I. GYM'NOSPERMS

Trees or shrubs with needle-like, linear or scale-like leaves, that are mostly evergreen. Ovules not inclosed in an ovary. Cotyledons 2-many.

TAXA'CEAE (YEW FAMILY)

Shrubs with solitary flowers; ovule naked, surrounded by a cup-shaped disk, which becomes pulpy and berry-like in fruit, inclosing the nut-like seed. Berry red.

1. TAX'US L. YEW

Flowers chiefly dioecious. Leaves evergreen, mucronate, rigid, scattered. A low straggling shrub, usually in the shade of other evergreens.

1. *T. canaden'sis* Marsh. (AMERICAN YEW. GROUND HEMLOCK.) A low straggling shrub. Leaves green on both sides. Berry globular, red.

PINA'CEAE (PINE FAMILY)

Trees or shrubs with resinous juice and mostly monoecious flowers in catkins. Fruit a cone or berry-like.

TRIBE I. ABIE'TEAE (PINE FAMILY PROPER)

Fruit a true cone, the imbricated scales in the axils of bracts. Ovules 2 on the inside of each scale at the base, seeds winged.

A. Cones not ripening till the second year.

1. **Pi'nus.** Leaves needle-shaped, 2-5 in a cluster, evergreen, in the axil of a thin scale. Sterile catkins in spikes at the bases of the new shoots, consisting of many almost sessile anthers spirally inserted on the axis. Cones more or less woody. Cotyledons of the embryo 3-12.

*B. Cones ripening the first year.**(1) Cones pendulous, bracts smaller than the scales.*

2. **Pic'ea.** Leaves sessile, keeled on both sides, needle-shaped, rigid, not appearing 2-ranked.
3. **Tsu'ga.** Leaves petioled, flat, whitened beneath, spirally arranged, but appearing 2-ranked.

(2) Cones erect, the bracts longer than the scales.

4. **A'bies.** Leaves spirally arranged but appearing 2-ranked, sessile, flat, with the midrib prominent on the whitened lower surface.
5. **Larix.** Leaves needle-shaped, soft, clustered or fascicled on lateral spurs of last year's wood, many in each bundle, those on the new shoots solitary and spirally arranged. Leaves fall off in the autumn.

TRIBE II. CUPRESS'EAE (CYPRESS FAMILY)

Fertile flowers of only a few scales, these not in the axils of bracts, forming in fruit either an oblong cone or a sort of false berry, owing to the thickening of the scales.

A. Flowers monoecious. Fruit a small cone.

6. **Thu'ja.** Leaves, some awl-shaped, others scale-like, closely imbricated on the flat branches. Catkins oblong, terminal.

B. *Flowers mostly dioecious. Fruit berry-like, black with a bloom.*

7. *Junip'erus*. Leaves awl-shaped or scale-like, sometimes of both shapes, on the same plant, evergreen, in whorls of 3 or opposite.

1. PINUS L. PINE

1. *P. resino'sa* Ait. (RED PINE.) Leaves in twos, slender, dark green, 4-6 inches long. Cone sessile, about 2 inches long. Bark rather smooth, reddish.

2. *P. Banksia'na* Lambert. (JACK PINE.) Leaves in twos, (very rarely in threes) $\frac{1}{2}$ -1 $\frac{1}{4}$ inches long. Cones conical, usually curved, smooth and hard, about 1 $\frac{1}{2}$ inches long.—Barren soil. A northern species.

3. *P. rig'ida* Mill. (PITCH PINE.) Resembles the former species, but the leaves are in threes and 3-4 inches long.

4. *P. Stro'bus* L. (WHITE PINE.) Leaves in fives, very slender, about 4 inches long. Bark smooth except on old trees, not reddish. Cones about 5 inches long.—Common.

2. PIC'EA Link. SPRUCE

1. *P. maria'na* BSP. (BLACK SPRUCE.) Branchlets rusty-hairy. Leaves needle-shaped, 4-sided, about $\frac{1}{2}$ inch long. Cones about 1 inch long, pointed, on curved stalks, persistent; scales stiff with minutely toothed edges.

2. *P. canaden'sis* BSP. (WHITE SPRUCE.) Branchlets usually smooth. Leaves as in No. 1. Cones about 2 inches long, cylindrical, deciduous, scales flexible with entire edges.

3. TSU'GA Carrière. HEMLOCK

1. *T. canaden'sis* Carr. (HEMLOCK SPRUCE.) Leaves flat, about $\frac{1}{2}$ inch long, lighter beneath, pointing out in two directions, i.e., right and left on each side of the branch, obtuse. Cones hanging, persistent, about $\frac{3}{4}$ inch long.

4. A'BIES Hill. FIR

1. *A. balsa'mea* Miller. (BALSAM FIR.) Leaves flat, the lower surface whitish and the midrib prominent, crowded, pointing mostly right and left on the branches. Cones erect on the upper sides of the branches, violet-coloured, and at maturity the scales become detached from the central axis.—Damp woods and swamps.

5. LARIX Adans. LARCH

1. *L. larici'na* Koch. (TAMARACK.) A slender, very graceful tree, well marked by the arrangement of its leaves in clusters on the lateral branches. The leaves are needle-like, soft, and about 1 inch in length. Cones $\frac{1}{2}$ - $\frac{3}{4}$ inch long, reddish, ovoid, with rounded scales.

6. THU'JA L. ARBOR VITAE

1. *T. occidenta'lis* L. (ARBOR VITAE. WHITE CEDAR.) Small evergreen trees with foliage in flat sprays. Leaves scale-like, opposite and decussate in 4 rows on the 2-edged branches. Cones composed of a few pointless scales arranged like the leaves.

7. JUNIP'ERUS L. JUNIPER

1. *J. commu'nis* L., var. *depres'sa* Pursh. (COMMON JUNIPER.) A spreading shrub with ascending or decumbent stems, growing in circular patches on poor sandy or rocky soil. Leaves in whorls of 3, glaucous above, white-striped beneath, prickly-pointed. Fruit berry-like, bluish-black, with a bloom.

2. *J. horizonta'lis* Moench. A prostrate, often creeping shrub with 2 sorts of leaves, awl-shaped and scale-like, these are usually opposite in arrangement. Fruit berry-like on short recurved peduncles.

3. *J. virginia'na* L. (RED CEDAR.) A shrub or small tree. The small, scale-like leaves have the opposite decussate arrangement of the White Cedar, but the longer, sharp-pointed leaves, when present, are usually in whorls of 3, like the juniper. Fruit berry-like, small, erect. Heart-wood red and odorous.

SUBDIVISION II. AN'GIOSPERMS

Trees, shrubs, or herbs with leaves that are not usually evergreen. Ovules inclosed in an ovary. Cotyledons 1-2.

CLASS I. MONOCOTYLE'DONS

Parts of the flowers usually in threes, never in fives. Leaves parallel-veined, rarely net-veined.

TYPHA'CEAE (CAT-TAIL FAMILY)

Aquatic or marsh herbs with sessile, linear, sword-shaped leaves, erect or floating; monoecious flowers, either in separate heads or on different parts of the same spike or spadix, but without a spathe; floral envelopes absent or represented by a few scales. Fruit an achene, 1-seeded.

Synopsis of the Genera

1. *Ty'pha*. Flowers in a very dense and long, cylindrical, terminal spike, the upper ones staminate, the lower pistillate, the ovaries long-stalked and surrounded by copious bristles forming the down of the fruit. Leaves sword-shaped, erect, sheathing the stem below.
2. *Sparga'nium*. Flowers in separate globular heads along the upper part of the stem, the higher ones staminate, the lower ones pistillate, each ovary sessile and surrounded by a few scales not unlike a calyx. Both kinds of heads leafy-bracted. Leaves flat or triangular, sheathing the stem with their bases.

1. TYPHA L. CAT-TAIL FLAG

1. *T. latifo'lia* L. (COMMON CAT-TAIL.) Stem 5-8 feet high. Leaves flat. No space between the staminate and pistillate parts of the spike.—Marshy places.

2. *S. angustifo'lia* L. (NARROW-LEAVED or SMALL CAT-TAIL.) Leaves channelled towards the base, narrowly linear. The two parts of the spike usually with an interval between them.

2. SPARGANIUM L. BUR-REED

1. *S. eurycar'pum* Engelm. Stem erect, stout, 2-4 feet high. Leaves mostly flat on the upper side, keeled and hollow-sided on the lower. Heads several, the pistillate an inch across in fruit. Nutlets or achenes with a broad, abruptly pointed top.—Borders of slow waters and ponds.

2. *S. sim'plex* Huds. Stem slender, erect, 6-24 inches high. Leaves more or less channelled and 3-angled, about $\frac{1}{3}$ inch wide. Fertile heads about $\frac{1}{2}$ inch broad in fruit. Stigma linear.—Borders of ponds, etc.

3. *S. america'num* Nutt., var. *androc'ladum* Fernald and Eames. Only 1 or 2 weak branches from the lower part of the flower-cluster.

4. *S. angustifolium* Michx. Very slender, with leaves floating, long, narrow, and flat. Inflorescence simple, and fruiting heads smaller than in No. 1.

5. *S. minimum* Fries. Usually floating, with very slender stems, and thin, flat, narrow leaves. Fertile heads only 1 or 2. Stigma oval. Fruit oblong-obovate, pointed, somewhat triangular.

NAJADACEAE (PONDWEED FAMILY)

Immersed aquatic herbs, with jointed stems and sheathing stipules. Leaves flat, immersed or floating. Flowers perfect or imperfect, inconspicuous, naked, or with a free, scale-like calyx. Ovaries 1-celled, 1-seeded, solitary or 2-4, and distinct.

Synopsis of the Genera

A. *Flowers perfect, in spikes or clusters. Leaves alternate.*

1. *Potamogeton*. Flowers with sepals, stamens, and ovaries each 4.

B. *Flowers monoecious or dioecious, axillary, naked, monandrous. Leaves opposite except in No. 4.*

2. *Najas*. Flowers dioecious. Ovaries solitary and naked. Stigmas 2 or 4, awl-shaped. Staminate flower (of 1 stamen) inclosed in a membranous spathe. Stems floating. Leaves linear, opposite, somewhat crowded into whorls, sharp-toothed, sessile and spreading at the base.

3. *Zannichellia*. Flowers monoecious, usually both kinds in the same axil. Ovaries 2-5, from a cup-shaped involucre. Leaves linear-thread-form, entire.

4. *Zostera*. Flowers monoecious, the two kinds naked and sessile, arranged alternately in 2 vertical rows on the inner side of a leaf-like, inclosed spadix. Sterile flowers of single, sessile, 1-celled anthers; the fertile of single, ovate-oblong ovaries. Stigmas 2, bristle-form, deciduous. Stem creeping.

1. POTAMOGETON L. PONDWEED

A. *Leaves of 2 sorts, the floating ones with a dilated petioled blade, different in form from the submersed ones.*

1. *P. natans* L. Submersed leaves grass-like or capillary; upper stipules very long, acute. Spikes cylindrical, all out of the water. Stem hardly branched. Floating leaves long-petioled, elliptical, with a somewhat heart-shaped base, with a blunt apex, 21-29-nerved.

2. *P. epihy'drus* Raf. Stem compressed. Submersed leaves linear, 2-5 inches long, 2-ranked, 5-nerved; stipules obtuse. Floating leaves short-petioled, chiefly opposite, oblong, 11-17-nerved. Spikes all above water.

3. *P. dimor'phus* Raf. Stems very slender. Floating leaves when present oval to lanceolate, about as long as the petiole; submersed leaves narrowly linear, or the upper ones broad-linear, or lance-oblong. Emerged flowers in many-flowered spikes; submersed flowers usually solitary. Fruit either winged and 4-5-toothed, or wingless and entire.

4. *P. alpi'nus* Balbis. Stem simple. Submersed leaves almost sessile, lanceolate and lance-oblong; floating leaves (often wanting) wedge-ob lanceolate, narrowed into a short petiole, 11-17-nerved. Spike dense, many-flowered. Fruit obovate, lenticular, with acute margin, and pointed with the long style.

5. *P. america'nus* C. and S. Stem often branching below. Submersed leaves very long, lanceolate and lance-linear, 7-15-nerved; floating leaves lance-oblong, or oblong-elliptical, long-petioled, 17-23-nerved. Spike dense. Fruit obliquely obovate, 3-keeled when dry, the middle one winged above.—In streams mostly.

6. *P. amplifo'lius* Tuckerman. Submersed leaves large, lanceolate or oval, acute at each end, recurved, wavy; stipules long and tapering. Floating leaves large, oblong or lance-ovate, or slightly cordate, long-petioled, 30-50-nerved.

7. *P. heterophyl'us* Schreb. Stem slender, very much branched below. Submersed leaves lanceolate or linear-lanceolate, acuminate or cuspidate, 3-7-nerved, upper ones petioled, lower sessile; floating leaves variable, with short blunt points, 9-15-nerved. Stipules obtuse.

8. *P. angustifo'lius* Berchtold and Presl. Much-branched at the base. Upper leaves long-petioled and sometimes emerged; the others nearly sessile, all usually numerous, wavy, and shining. Resembling the next species, but smaller.

B. Leaves all submersed and similar, mostly sessile, membranaceous and dilated. Stipules obtuse, becoming loose.

9. *P. lu'cens* L. Stem thick, branching. Leaves petioled, oval or lanceolate, mucronate, frequently shining. Fruit roundish, compressed, with blunt edges, slightly keeled.

10. *P. praelon'gus* Wulfen. Stem very long and branching. Leaves all submersed and similar, lanceolate, half-clasping, with a boat-shaped cavity at the end. Spikes loose-flowered with very long peduncles. Fruit sharply keeled when dry. Stem white.—Ponds and large rivers.

11. *P. perfolia'tus* L. Leaves all submersed, varying in width from orbicular to lanceolate, clasping by a heart-shaped base. Stem branching.

12. *P. Richardso'nii* Rydb. Has long-lanceolate, acuminate leaves. Peduncles thickened upward.

C. Leaves all submersed and similar, mostly sessile, linear or bristle-like.

13. *P. zosterifo'lius* Schum. Leaves linear, grass-like, sessile, abruptly pointed, with three large nerves and many fine ones. Stem branching, wing-flattened. Stipules free from the sheathing base of the leaf. Spikes cylindrical.

14. *P. folio'sus* Raf. Stem filiform, flattish and much-branched. Leaves narrowly-linear, acute, obscurely 3-nerved. Spikes capitate.

15. *P. pusil'lus* L. Stem slender, somewhat flattish, often much-branched. Leaves narrow- or setaceous-linear, 1-3 nerved, with translucent glands on each side at the base.

16. *P. Frie'sii* Rupr. Like the last, but the stem less branching, and the leaves broader, often 5-nerved.

17. *P. pectina'tus* L. Leaves bristle-shaped. Stem repeatedly forking, filiform. Spikes interrupted, on long slender peduncles. Stipules united with the sheathing base of the leaf.

18. *P. Robbin'sii* Oakes. Leaves narrowly lanceolate or linear, crowded in 2 ranks, recurved, serrulate, many-nerved. Stems rigid with numerous branches. Fruit keeled with a broadish wing. Stipules united with the sheathing base of the leaf.

2. NAJ'AS L. NAIAD

1. *N. flex'ilis* Rostk and Schmidt. Stems very slender. Leaves very narrowly linear, minutely serrulate.

3. ZANNICHEL'LIA L. HORNED PONDWEED

1. *Z. palus'tris* L. Fruit obliquely oblong, flattish and somewhat incurved. Style half as long as the nutlet.

4. ZOSTERA L. EEL-GRASS. GRASS-WRACK

1. *Z. mari'na* L. Leaves obscurely 3-5-nerved.—Atl. sea-coast, in shoal water.

JUNCAGINA'CEAE (ARROW GRASS FAMILY)

Marsh herbs with greenish perfect flowers in spikes or racemes. Sepals and petals 3 each, stamens 3-6, carpels 3-6, somewhat united but separating at maturity. Leaves either grass-like or terete and rush-like.

Synopsis of the Genera

1. *Triglo'chin*. Flowers small, in a spike or close raceme, without bracts. Carpels united to the top; when ripe, splitting away from a central persistent axis. Leaves rush-like.
2. *Scheuchze'ria*. A low bog herb, with a creeping jointed rootstock and grass-like leaves. Stamens 6. Carpels 3, globular, nearly distinct.

1. TRIGLO'CHIN L. ARROW GRASS

1. *T. palus'tris* L. A slender, rush-like plant, 6-18 inches high. Carpels 3, awl-pointed at the base, splitting away from below upwards. Spike or raceme slender, 3-4 inches long.

2. *T. marit'ima* L. The whole plant is stouter than No. 1, and the carpels are usually 6 in number.—Atl. sea-coast and saline marshes.

2. SCHEUCHZE'RIA L. SCHEUCHZERIA

1. *S. palus'tris* L. Stem zigzag. Flowers in a loose terminal raceme, with sheathing bracts.—Bogs.

ALISMA'CEAE (WATER PLANTAIN FAMILY)

Marsh herbs with flowers having 3 distinct sepals and 3 distinct petals, pistils either apocarpous or separating at maturity into distinct carpels, and hypogynous stamens 6-many. Flowers on scapes or scape-like stems in loose racemes or panicles; calyx green, persistent; corolla white. Leaves mostly heart-shaped or arrow-shaped, petiolate and strongly nerved.

Synopsis of the Genera

1. *Alis'ma*. Flowers perfect. Stamens usually 6. Carpels numerous in a ring. Leaves all radical. Scapes with whorled paniced branches.

2. *Sagitta'ria*. Flowers monoecious, sometimes dioecious. Stamens numerous. Carpels numerous, in more or less globular heads. Leaves arrow-shaped, but varying greatly. Flowers mostly in whorls of 3 on the scapes, the sterile ones uppermost.

1. ALIS'MA L. WATER PLANTAIN

1. *A. Planta'go-aquat'ica* L. Leaves long-petioled, mostly oblong-heart-shaped, but often narrower, 3-9-nerved or ribbed, and with cross veinlets between the ribs. Flowers small, white, in a large and loose, compound panicle.—Low and marshy places, often growing in the water.

2. SAGITTA'RIA L. ARROW-HEAD

A. *Filaments narrow, longer than the anthers.*

1. *S. latifo'lia* Willd. Plant glabrous. Leaves ovate, acute, usually sagittate, with acute lobes. Petals large, white. Achenes irregular, obovate, with horizontal beak. Very variable, includes several ill-defined forms.

B. *Filaments very short, with enlarged, mostly glandular base.*

2. *S. heterophyl'la* Pursh. Scape weak and at length procumbent. Achenes with a long erect beak. Leaves lanceolate or lance-ovate, entire, or with one or two narrow, basal, sagittate appendages.

3. *S. gramin'ea* Michx. Scape very slender, erect. Leaves varying from ovate-lanceolate to linear, scarcely ever sagittate. Achenes small and almost beakless.

HYDROCHARITA'CEAE (FROG'S-BIT FAMILY)

Aquatic herbs, with dioecious or polygamo-dioecious flowers on scape-like peduncles from a kind of spathe of one or two leaves, the perianth in the fertile flowers of 6 pieces united below into a tube, which is adherent to the ovary. Stigmas 3. Fruit ripening under water.

Synopsis of the Genera

1. *Elode'a*. Growing under water, the pistillate flowers alone coming to the surface. Stem leafy and branching. Perianth of the fertile flowers with a 6-lobed, spreading limb, the tube attached to the thread-like style. Leaves crowded, pellucid, 1-nerved, sessile, whorled in threes or fours, or opposite. Stamens 3-9.

2. *Vallisneria*. Stemless plants. Nothing but the pistillate flowers above the surface, these on scapes of great length, and after fertilization drawn below the surface by the spiral coiling of the scapes. Tube of the perianth not prolonged. Leaves linear, thin, long and ribbon-like.

(In both genera the staminate flowers break off spontaneously and float on the surface around the pistillate ones, shedding their pollen upon them.)

1. *ELODEA* Michx. WATER-WEED

1. *E. canadensis* Michx. Leaves linear to oblong, about $\frac{1}{4}$ inch in length. Common in slow waters.

2. *VALLISNERIA* L. TAPE-GRASS. EEL-GRASS

1. *V. spiralis* L. Leaves 1-4 feet long.—Common in slow waters.

GRAMIN'EAE (GRASS FAMILY)

CYPERA'CEAE (SEDGE FAMILY)

(For the details of the numerous genera and species of these two large families, the student is referred to *Gray's New Manual of Botany* or to any other advanced work.)

ARA'CEAE (ARUM FAMILY)

Herbs with pungent juice and simple or compound leaves, these sometimes net-veined and hence suggesting that the plants may be Dicotyledons. Spadix usually accompanied by a spathe. Flowers either without a perianth of any kind, or with 4-6 sepals. Fruit usually a berry.

Synopsis of the Genera

A. *Leaves not linear. Flowers without perianth of any sort.*

1. *Arisaema*. Flowers mostly dioecious, collected on the lower part of the spadix only. Spathe (in our common species) arched over the spadix. Scape from a solid bulb. Leaves compound, net-veined, sheathing the scape below with their petioles. Berries bright red.
2. *Peltandra*. Flowers monoecious, covering the whole spadix; anthers above, ovaries below. Spathe convolute throughout, wavy on the margin, mostly green. Leaves arrow-shaped. Scapes from a root of thick fibres. Fruit a fleshy green berry, 1-3-seeded.

3. *Calla*. Flowers (at least the lower ones) perfect, covering the whole spadix. Spathe open and spreading, with a white upper surface, tipped with an abrupt point. Scape from a creeping rootstock. Leaves not net-veined, simple, heart-shaped. Fruit a red berry.

B. *Leaves not linear. Flowers with a perianth of 4 sepals.*

4. *Symplocar'pus*. Leaves all radical, very large and veiny, appearing after the spathes, which are close to the ground and are produced very early in spring. Flowers perfect, their ovaries immersed in the spadix, the latter globular and surrounded by the shell-shaped spathe. Sepals hooded. Stamens 4. Fruit consisting of the soft enlarged spadix in which the seeds are sunk.

C. *Leaves linear, sword-shaped. Spadix on the side of the scape. Flowers with a perianth of 6 sepals.*

5. *Ac'orus*. Scape 2-edged, resembling the leaves, the cylindrical spadix borne on one edge. Sepals hollowed. Stamens 6. The leaf-like prolongation of the scape forms a sort of narrow spathe.

1. ARISAE'MA Mart. INDIAN TURNIP

1. *A. tryphyl'lum* Schott. (INDIAN TURNIP. JACK-IN-THE-PULPIT.) Leaves usually 2, each of 3 ovate, coarsely net-veined leaflets. Spadix club-shaped, blunt, and shorter than the spathe, which is pale green, often variegated with purplish and whitish stripes, and which arches over the spadix.

2. *A. Dracon'tium* Schott. (GREEN DRAGON.) Leaf usually solitary, pedately divided into 7-11 oblong-lanceolate, pointed leaflets. Spathe convolute, pointed; the slender point of the spadix extending beyond it.

2. PELTAN'DRA Raf. ARROW ARUM

1. *P. virgin'ica* Kunth. Root of thick tufted fibres. Scape 12-18 inches high. Staminate part of the spadix much longer than the pistillate.—Shallow water; apparently rare.

3. CALLA L. WATER ARUM

1. *C. palus'tris* L. (MARSH CALLA.) A low bog herb with petioles and scape from 4-8 inches high and spathe 1-2½ inches long and 1-1½ inches wide.

4. SYMPLOCAR'PUS Salisb. SKUNK CABBAGE

1. *S. foet'idus* Nutt. Leaves many, 1-2 feet long, often 1 foot wide, ovate-heart-shaped, short-petioled. Spathe purplish and yellowish, incurved. Plant with skunk-like odour.—Bogs and wet places; not common northward.

5. AC'ORUS L. SWEET FLAG. CALAMUS

1. *A. Cal'amus* L. Scape much prolonged beyond the spadix.—Swamps and wet places.

LEMNA'CEAE (DUCKWEED FAMILY)

Very small plants floating about freely on the surface of ponds and ditches, consisting merely of a little frond, commonly with a single root or a tuft of roots from the lower surface, and producing minute monoecious flowers from a cleft in the edge of the frond, or from the upper surface. Fruit a 1-7-seeded utricle. The flowers are rarely to be seen.

Synopsis of the Genera

1. *Lem'na*. Flowers from a cleft in the edge of the frond, usually 2 of a single stamen each, and 1 of a simple pistil, the three surrounded by a spathe. Root only one, or rarely wanting.
2. *Spirode'la*. Flowers as in *Lemna*. Fronds 7-12-nerved, with several rootlets.
3. *Wolff'ia*. Flowers from the centre of the upper surface of the frond, only 2; one of a single stamen, the other of a simple pistil. Roots none. The smallest of flowering plants.

1. LEM'NA L. DUCKWEED

1. *L. trisul'ca* L. Fronds oblong to oblong-lanceolate, $\frac{1}{4}$ inch or more long, narrowed at the base into a slender stalk, denticulate at the tip, obscurely 3-nerved. Rootlet single, often wanting.—Ponds and springy places.

2. *L. minor* L. Fronds round to elliptic-obovate, very small. Rootlet single.—Stagnant waters.

2. SPIRODE'LA SCHLEID

1. *S. polyrhi'za* Schleid. Fronds round-obovate, green above, purplish beneath, mostly 7-nerved, somewhat smaller than in *L. trisulca*. Rootlets several.—Common in ponds.

3. WOLFFIA Horkel

1. *W. columbia'na* Karsten. Globular or nearly so, light green all over, not dotted.—Stagnant waters, usually rather below the surface.

2. *W. puncta'ta* Griseb. Oblong, deep green above, pale below, dotted all over with brown.—With the last, but floating.

ERIOCAULA'CEAE (PIPEWORT FAMILY)

Aquatic or marsh herbs with clustered, radical, grass-like leaves, in which the large cells are clearly visible. The solitary naked scape bears a dense head of small flowers.

1. ERIOCAUL'ON L. PIPEWORT

1. *E. articula'tum* Morong. A slender plant with a naked scape from 2 inches to 2 feet high, growing in water or along the margins of our northern ponds. Leaves short, awl-shaped, in a tuft at the base. Flowers monoecious, both staminate and pistillate in a small woolly head at the summit of the scape. Perianth double; the outer set or calyx of 2-3-keeled sepals; the inner set or corolla tubular in the sterile flowers and of 2-3 separate petals in the fertile ones. Scape usually 7-striate. The head (except the beard) lead-coloured.

XYRIDA'CEAE (YELLOW-EYED GRASS FAMILY)

Rush-like herbs with equitant leaves sheathing the base of a naked scape, terminated by a head of perfect 3-androus flowers, with glumaceous calyx and coloured corolla. Capsule 3-valved and 1-celled, containing several or many seeds.

1. XY'RIS L. YELLOW-EYED GRASS

1. *X. monta'na* Ries. Scape small and very slender, 2-9 inches high, not bulbous at the base. Flowers in the axils of scale-like bracts densely imbricated in a head. Sepals 3. Petals 3, mostly yellow. Fertile stamens 3, with 3 alternating sterile filaments bearded at the apex.—Peat-bogs and wet sandy shores, Atl. Prov. and Ontario northward.

PONTEDERIA'CEAE (PICKEREL-WEED FAMILY)

Aquatic herbs with perfect flowers from a spathe. Leaves erect, heart-shaped.

Synopsis of the Genera

1. *Pontede'ria*. Inflorescence a many-flowered spike. Perianth 2-lipped. Stamens 6.
2. *Heteranthe'ra*. Flowers solitary. Stamens 3.

1. PONTEDERIA L. PICKEREL-WEED

1. *P. corda'ta* L. A stout plant growing in shallow water, sending up a scape bearing a single large, arrow-heart-shaped, blunt leaf, and a spike of violet-blue flowers with a spathe-like bract. Perianth 2-lipped, the 3 upper divisions united, the 3 lower spreading, the whole revolute-coiled after flowering, the fleshy base inclosing the fruit. Stamens 6, 3 of them exerted on long filaments, the rest short.

Var. *angustifo'lia* Torr. Has narrow, scarcely cordate leaves.

2. HETERANTHE'RA R. and P. WATER STAR-GRASS

1. *H. du'bia* MacM. A grass-like herb, wholly under water, only the small yellowish flowers reaching the surface, the latter single, from spathes. Perianth salver-shaped, regular. Stamens 3, anthers sagittate.

JUNCA'CEAE (RUSH FAMILY)

Grass-like or sedge-like plants, with small regular flowers. Perianth greenish and glumaceous, of 6 divisions in 2 sets of 3 each. Stamens 6 (occasionally 3). Style 1. Stigmas 3. Pod 3-celled, or 1-celled with 3 placentae on the walls. The plants of the Family are not of any very great interest to the young student, and the determination of the species is rather difficult. A brief description of a few of the most common plants of this Family is given here, as an easy introduction to its study with the aid of more advanced text-books.

Synopsis of the Genera

1. *Lu'zula*. Plants less than 1 foot high. Leaves linear or lance-linear, flat, usually hairy. Pod 1-celled, 3-seeded. Flowers in umbels or in spikes. Plants usually growing in dry ground.

2. *Jun'cus*. Plants always smooth, growing in water or wet soil. Flowers small, greenish or brownish, paniced or clustered. Pod 3-celled, many-seeded.

1. LU'ZULA DC. WOOD-RUSH

1. *L. saltuen'sis* Fernald. Flowers umbelled, long-peduncled, brown-coloured. Sepals pointed.—Shady banks.

2. *L. campes'tris* DC. Flowers (light brown) in 4-12 spikes, the spikes umbelled. Sepals bristle-pointed.—Fields and woods.

2. JUNCUS L. RUSH

A. *Scapes single and leafless, but with sheaths at the base. Flowers in sessile panicles, apparently from the side of the scape, owing to the involucre leaf being similar to and continuing the scape.*

1. *J. effu'sus* L. (COMMON or SOFT RUSH.) Scape 2-4 feet high, soft and pliant, furnished at the base with merely leafless sheaths, the inner sheaths awned. Panicle many-flowered. Flowers small, greenish, only 1 on each pedicel. Stamens 3. Pod greenish-brown, triangular-obovate, not pointed.—Marshes.

2. *J. filifor'mis* L. Very slender scape (1-2 feet high), fewer flowers than No. 1, and 6 stamens in each. Pod greenish, broadly ovate, and short-pointed. No leaves.

3. *J. bal'ticus* Willd., var. *littora'lis* Engelm. Scape rigid, 2-3 feet high. No leaves. Panicle loose. Flowers brownish. Pod elliptical, somewhat triangular, obtuse but pointed, deep brown.

B. *Stem leafy at the base or throughout; the leaves flat or channelled but never knotted. Panicle terminal.*

4. *J. styg'ius* L., var. *america'nus* Buch. Scape slender, 1-3-leaved below, naked above. Leaves thread-like, hollow, not knotted. Heads 1 or 2, of 3-4 flowers, about as long as the dry, awl-pointed, sheathing bract. Stamens 6.—Peat-bogs, Atl. Prov.

5. *J. Gree'nii* Oakes and Tuckerm. Stem slender, simple, tufted. Leaves nearly terete, deeply channelled on the inner side. Flowers solitary, paniced. The principal leaf of the involucre usually much longer than the panicle. Pod ovoid-oblong, obtuse, longer than the acute sepals. Seeds ribbed and delicately cross-lined.—Wet sandy places.

6. *J. Vase'yi* Engelm. Stem slender, rigid, tufted, leafy below, 12–30 inches high, not branching. Leaves nearly terete, very slightly channelled. Panicle longer than the involucre leaf. Flowers greenish or light brown, few, often 1-sided. Pod oblong, retuse, greenish-brown.—Wet meadows, Pr. Prov.

7. *J. Gerard'i* Loisel. (BLACK GRASS.) Stems rigid, 1–2 feet high. Panicle contracted, usually longer than the involucre leaf. Flowers brown and green. Sepals incurved, as long as the mucronate capsule. Seeds obovate, delicately ribbed and cross-lined.—Salt marshes, Atl. Prov.

8. *J. longis'tylis* Torr. and Gr. Flowers in heads. Heads few, or rarely single. Flowers greenish with brown lines.—Pr. Prov.

9. *J. bufo'nus* L. Stem leafy, slender, 3–9 inches high, branching from the base. Panicle terminal, spreading. Flowers greenish, single on the pedicels. Sepals awl-pointed, the outer set much longer than the inner and than the blunt pod. Stamens 6.—Ditches along roadsides.

10. *J. ten'uis* Willd. Stems leafy below, wiry, 9–18 inches high, simple tufted. Panicle loose, shorter than the slender involucre leaves. Flowers greenish, single on the pedicels; the sepals longer than the blunt pod. Seeds white-pointed at both ends.—Open low grounds.

C. *Stem leafy; the leaves terete or laterally compressed, knotted by internal cross-partitions. Panicle terminal, the flowers mostly in heads.*

(1) *Stamens 6.*

11. *J. pelocar'pus* E. Meyer. Stems slender and erect, 6–18 inches high. Leaves few, thread-like, slightly knotted. Flowers greenish with red, single or in pairs in the forks and along one side of the branches of the panicle, and often with accompanying tufts of leaves. The 3 inner sepals longer than the outer ones, but shorter than the oblong, taper-beaked, 1-celled pod. Seeds obovate, short-pointed.

12. *J. articula'tus* L. Stems 9–15 inches high, tufted from a short creeping rootstock, with 1 or 2 slender leaves, bearing crowded, 3–8-flowered heads in short spreading panicles. Capsule deep brown.—Wet places, Atl. Prov.

13. *J. alpi'nus* Villars., var. *insignis* Fries. Stems erect, 9–18 inches high, with 1 or 2 slender leaves. Branches of the meagre panicle erect, bearing numerous distant, greenish or brownish heads of 3–6 flowers each. Outer sepals mucronate or cuspidate, and longer than the rounded inner ones. Pod short-pointed, light-brown. Seeds spindle-shaped.—Lake- and river-margins, common.

14. *J. nodo'sus* L. Stem erect, 6–15 inches high, slender, from a creeping, slender and tuber-bearing rootstock, usually with 2 or 3 slender leaves. Heads few, 8–20-flowered, and overtopped by the involucre leaf. Flowers brown. Pod slender, taper-pointed, 1-celled. Seeds obovate, mucronate.—Common.

15. *J. megaceph'alus* M. A. Curtis. Large 30–80-flowered heads.

(2) *Stamens 3. Seeds tailed.*

16. *J. brevicauda'tus* Fernald. Stems slender, 9–18 inches high, tufted, bearing 2 or 3 leaves. Panicle somewhat erect, contracted; the heads 3–5-flowered, deep brown. Pod prismatic, abruptly pointed, deep brown. Seeds slender, with short tails.—A very late-flowering species.

17. *J. canaden'sis* J. Gay. Taller than No. 16, with many-flowered heads.

LILIA'CEAE (LILY FAMILY)

Herbs, distinguished as a whole by their regular and symmetrical flowers, having a 6-leaved perianth (but 4-leaved in the genus *Maianthemum*), free from the usually 3-celled ovary, and as many stamens as divisions of the perianth (one before each) with 2-celled anthers. Fruit a pod or berry, generally 3-celled. The outer and inner divisions of the perianth coloured alike, except in the genus *Trillium*.

Synopsis of the Genera

A. *Leaves net-veined.*

1. *Tril'lium*. Leaves 3, in a whorl at the top of the stem. Divisions of the perianth in 2 sets, the outer green, the inner coloured.
2. *Mede'ola*. Leaves in 2 whorls, the lower near the middle of the stem, and consisting of 5–9 leaves, the upper of (generally) 3 small leaves, near the summit. Stem tall, covered with loose wool. Flowers small, in an umbel. Divisions of the perianth alike, greenish-yellow, recurved. Anthers turned

outwards. Styles thread-shaped. Berry globular or nearly so, dark purple.

3. *Smi'lax*. Shrubby or herbaceous plants with alternate, ribbed, petioled leaves and small, greenish-yellow, dioecious flowers in umbels.

B. *Leaves straight-veined, linear, grass-like, alternate. Stem simple or tufted. Styles 3.*

4. *Zygad'enus*. Flowers rather large, perfect or polygamous, greenish-white, in a few-flowered panicle; the divisions of the perianth each with a conspicuous obcordate spot or gland on the inside, near the narrowing base. Stem smooth and glaucous, from a coated bulb.

5. *Tofie'ldia*. Flowers small, perfect, greenish-white, in a terminal raceme or spike, which, however, develops from above downward; the pedicels in clusters of 3, from little involucre of 3 bracts. Pod triangular. Roots fibrous. Stem leafy at the base only, in our species sticky. Leaves 2-ranked, equitant.

C. *Leaves straight-veined, but broad (not grass-like), alternate. Stem not from a bulb. Style one, or more or less divided into 3 above.*

(1) *Leaves of the perianth completely separate (polyphyllous).*

6. *Uvula'ria*. Stem leafy, forking above. Flowers yellow, at least an inch long, drooping, lily-like, usually solitary (but occasionally in pairs) at the end of the forks of the stem. Style deeply 3-cleft. Pod triangular. Leaves clasping-perfoliate.
7. *Oake'sia*. Flowers resembling those of *Uvularia*, but the stems are acutely angled, and the leaves are sessile or partly clasping.
8. *Clinto'nia*. Stemless, the naked scape sheathed at the base by 2, 3, or 4 large, oblong or oval, ciliate leaves. Flowers few, greenish-yellow, in an umbel at the top of the scape. Filaments long and slender. Style long, the stigmas hardly separate. Berry blue.
9. *Dis'porum*. Downy low herbs, branching above. Flowers greenish, bell-shaped, rather large, solitary or in pairs, drooping on terminal slender peduncles. Stigmas 3. Leaves ovate-oblong, taper-pointed, closely sessile, downy underneath. Berry oblong or obovate, red.
10. *Strep'topus*. Stem leafy and forking. Flowers small, not quite in the axils of the ovate clasping leaves, on slender peduncles which are bent near the middle. Anthers arrow-shaped, 2-horned at the apex.

11. **Vera'trum.** Stem tall and stout, very leafy to the top, from a thickened base, producing coarse fibrous roots. Flowers yellowish-green, in a pyramidal panicle of dense, spike-like racemes. Perianth somewhat spreading, slightly adherent to the ovary. Filaments of the stamens recurved. Anthers kidney-shaped. Styles 3.

(2) *Leaves of the perianth more or less united (gamophyllous).*

12. **Smilaci'na.** Flowers small, white, in a terminal raceme. Perianth 6-parted, spreading. Style short and thick. Stigma obscurely lobed. Filaments slender.
13. **Maian'themum.** Resembling Smilacina, but the perianth is 4-parted; stamens 4 and stigma 2-lobed.
14. **Polygon'atum.** Flowers small, greenish, nodding, mostly in pairs in the axils of the nearly sessile leaves. Perianth cylindrical, 6-lobed at the summit, the 6 stamens inserted on or above the middle of the tube. Stem simple, from a long and knotted rootstock. Leaves glaucous beneath.
15. **Al'etris.** Bitter herbs with a cluster of lanceolate leaves at the base. Flowers small, in a spike-like raceme.
16. **Aspar'agus.** Flowers small, greenish-yellow, axillary, on jointed pedicels. Perianth 6-parted, spreading above, the 6 stamens on its base. Stigma 3-lobed. Berry globular, 3-celled. Stem much-branched. Leaves thread-like. Rootstock thick and matted.

D. *Leaves straight-veined, not grass-like. Stem usually from a coated or scaly bulb. Style 1, not divided above, but the stigma sometimes 3-lobed.*

17. **Lil'ium.** Stem leafy, from a scaly bulb, the leaves often whorled or crowded. Anthers at first erect, at length versatile. Style long, rather club-shaped. Stigma 3-lobed. Pod oblong. Flowers large and showy, one or more.
18. **Erythro'nium.** Herbs with two smooth fleshy leaves, the petioles of which sheathe the base of the usually 1-flowered scape. Bulb deep.
19. **Allium.** Scape naked, from a coated bulb. Flowers in an umbel, from a spathe. Style thread-like. Strong-scented plants.
20. **Hemerocal'lis.** Scape not from a bulb, but from a fleshy-fibrous root. Perianth large, orange-yellow, lily-like, the short tube inclosing the ovary, the spreading limb 6-parted. Stamens 6, on the perianth. Leaves long, linear, keeled, in 2 ranks, at the base of the scape. Flowers several, bracted.

1. TRILLIUM L. WAKE-ROBIN

1. *T. grandiflorum* Salisb. (LARGE WHITE TRILLIUM.) Leaves sessile, longer than broad. Peduncle erect. Petals white (rose-coloured when old), obovate.—Rich woods.

2. *T. erectum* L. (PURPLE TRILLIUM.) Leaves sessile, about as broad as long. Peduncles erect. Petals dull purple or occasionally greenish-white, ovate.—Rich woods.

3. *T. cernuum* L. Leaves sessile or nearly so, broadly rhomboid, abruptly pointed. Peduncle recurved under the leaves. Petals white, oblong-ovate, acute.—Chiefly eastward.

4. *T. undulatum* Willd. (PAINTED TRILLIUM.) Leaves distinctly petioled, rounded at the base. Petals pointed, white, with purple stripes inside at the base.—Not uncommon northward in damp woods and low grounds.

2. MEDEOLA L. INDIAN CUCUMBER-ROOT

1. *M. virginiana* L. Stem 1-3 feet high.—Rich woods.

3. SMILAX L. GREEN BRIER. CAT BRIER

1. *S. hispida* Muhl. Stem woody, densely covered below with long weak prickles. Leaves large, ovate or heart-shaped, pointed, thin, 5-9-nerved. Peduncles of the axillary umbels much longer than the petioles. Berry black.—Moist thickets.

2. *S. rotundifolia* L., var. *quadrangularis* Wood. Stem woody, it and the branches armed with scattered prickles. Branches 4-angular. Peduncles not longer than the petioles. Leaves ovate, broader than long, slightly cordate. Berry blue-black.

3. *S. herbacea* L. (CARRION FLOWER.) Stem herbaceous, not prickly. Leaves ovate-oblong and heart-shaped, 7-9-ribbed, long-petioled, mucronate. Flowers carrion-scented. Berry bluish-black.—Meadows and river-banks.

4. *S. ecirrata* Wats. Stem herbaceous, erect, without tendrils or only at the top. Lower leaves bract-like; upper ovate-orbicular.

4. ZYGADENUS Michx. ZYGADENE

1. *Z. chloranthus* Richards. Not uncommon in bogs and beaver-meadows northward. Leaves flat and pale.

5. TOFIE'LDIA Hudson. FALSE ASPHODEL

1. *T. glutino'sa* Pers. Stem and pedicels very sticky with dark glands. Leaves short.—Lake Huron coast.

6. UVULA'RIA L. BELLWORT

1. *U. perfolia'ta* L. Leaves clasping-perfoliate. Rootstock short.—Rich woods.

7. OAKE'SIA Wats.

1. *O. sessilifo'lia* Wats. Leaves sessile or partly clasping, lance-oblong. Rootstock creeping.—Chiefly eastward.

8. CLINTO'NIA Raf. CLINTONIA

1. *C. borea'lis* Raf. Umbel 2-7-flowered. Leaves 5-8 inches long. Perianth pubescent outside.—Damp woods, often under evergreens.

9. DIS'PORUM Salisb.

1. *D. lanugino'sum* Nichols. Leaves taper-pointed. Fruit oblong, pointed.—Rich woods, Western Ontario.

2. *D. trachycar'pum* B. and H. Whitish perianth, ovate to oblong-lanceolate leaves, and broadly obovate fruit.—Pr. Prov.

10. STREP'TOPUS Michx. TWISTED-STALK

1. *S. ro'seus* Michx. Flowers rose-purple. Leaves green both sides, finely ciliate. Stigma 3-cleft.—Damp woods.

2. *S. amplexifo'lius* DC. Flowers greenish-white. Leaves very smooth, glaucous beneath. Stigma entire.—Chiefly Atl. Prov. and Pr. Prov.

11. VERA'TRUM L. FALSE HELLEBORE

1. *V. vir'ide* Ait. (AMERICAN WHITE HELLEBORE. INDIAN POKE.) Leaves broadly oval, pointed, sheath-clasping. Segments of the perianth ciliate-serrulate.—Swamps and low grounds, Atl. Prov. and Eastern Townships, Que.

12. SMILACI'NA Desf. FALSE SOLOMON'S SEAL

1. *S. racemo'sa* Desf. (FALSE SPIKENARD.) Raceme compound. Stem pubescent, 2 feet high. Leaves many, oblong, taper-pointed, ciliate. Berries speckled with purple.—Rich woods and thickets.

2. *S. stella'ta* Desf. Raceme simple. Stem nearly smooth, 1-2 feet high. Leaves 7-12, oblong-lanceolate, slightly clasping. Berries black.—Moist woods and copses.

3. *S. trifo'lia* Desf. Raceme simple. Stem low (3-6 inches), glabrous. Leaves usually 3, oblong, the bases sheathing. Berries red.—Bogs.

13. MAIAN'THEMUM Wiggers

1. *M. canaden'se* Desf. Distinguished at once by the 4-parted perianth and the 4 stamens. Raceme simple. Stem 3-5 inches high. Leaves usually 2, but sometimes 3.—Moist woods.

14. POLYGON'ATUM Hill. SOLOMON'S SEAL

1. *P. biflo'rum* Ell. (SMALLER SOLOMON'S SEAL.) Stem slender, 1-3 feet high. Leaves ovate-oblong or lance-oblong. Peduncles mostly 2-flowered. Filaments hairy.—Rich woods.

2. *P. commuta'tum* Dietrich. (GREAT SOLOMON'S SEAL.) Occasionally met with westward and south-westward. The stem is taller and stouter than in the last, the peduncles are several-flowered, and the filaments are not hairy.

15. AL'ETRIS L. COLIC-ROOT. STAR-GRASS

1. *A. farino'sa* L. Upper leaves bract-like. Flowers white, perianth tubular and roughened.

16. ASPAR'AGUS L. ASPARAGUS

1. *A. officina'lis* L. (GARDEN ASPARAGUS.) Escaped from gardens in a few places.

17. LIL'TUM L. LILY

1. *L. philadel'phicum* L. (WILD ORANGE-RED LILY.) Divisions of the perianth narrowed into claws below, not recurved at the top. Flowers erect, 1-3, orange, spotted with purple inside. Leaves linear-lanceolate, the upper mostly in whorls of 5-8.—Sandy soil.

2. *L. canaden'se* L. (WILD YELLOW LILY.) Divisions of the perianth recurved above the middle. Flowers nodding, few, orange, spotted with brown inside. Leaves remotely whorled, 3-ribbed.—Swamps and wet meadows.

3. *L. super'bum* L. (TURK'S-CAP LILY.) Divisions of the perianth very strongly recurved. Flowers nodding, often numerous,

in a pyramidal raceme, bright orange, dark-purple-spotted within. Lower leaves whorled, 3-ribbed or nerved. Stem taller than either of the first two, 3-7 feet.—Rich low grounds, commoner southward and south-westward.

18. ERYTHRO'NIUM L. DOG'S-TOOTH VIOLET

1. *E. america'num* Ker. (YELLOW ADDER'S TONGUE.) Perianth light yellow, sometimes spotted at the base.—Copses and rich meadows.

2. *E. al'bidum* Nutt. (WHITE DOG'S-TOOTH VIOLET.) Leaves less blotched than the last. Perianth pinkish-white.—Not common.

19. ALLIUM L. ONION. LEEK

A. Capsule not crested.

1. *A. tricoc'cum* Ait. (WILD LEEK.) Leaves 2 or 3, flat, lance-oblong, 5-9 inches long, 1-2 inches wide, appearing in early spring and withering before the flowers are developed. Sepals white. Pod strongly 3-lobed. Scape 9 inches high.—Rich woods.

2. *A. Schoenop'rasum* L., var. *sibir'icum* Hartm. Leaves linear, hollow. Scape naked, or leafy at the base. Flowers rose-purple, in a globular umbel. Sepals lanceolate, pointed. Ovary not crested.—Margin of rivers, Atl. Prov. and Pr. Prov.

3. *A. canaden'se* L. (WILD GARLIC.) Leaves narrowly linear. Ovary crested with 6 teeth. Umbel few-flowered, often bearing a head of bulbs instead of flowers. Sepals pale rose-colour.—Along river-banks; rather rare.

B. Capsule crested with short projections.

4. *A. cer'nuum* Roth. (WILD ONION.) Scape naked, angular, nodding at the apex, bearing a loose or drooping umbel of rose-coloured flowers. Leaves linear, flattened, sharply keeled. Capsule 6-crested.—Pr. Prov.

5. *A. stella'tum* Ker. Scape terete, 6-18 inches high, slender, bearing an erect umbel. Stamens and style exserted. Bulb-coats membranous. Capsule 6-crested.—Pr. Prov. Not very common.

6. *A. reticula'tum* Don. Scape 3-8 inches high. Bulbs densely and coarsely fibrous-coated. Stamens not exserted. Capsule crested.—Pr. Prov.

20. HEMEROCAL'IS L. DAY-LILY

1. *H. ful'va* L. (COMMON DAY-LILY.) Inner divisions of the tawny orange perianth wavy and obtuse.—Escaped from gardens occasionally.

DIOSCOREA'CEAE (YAM FAMILY)

Twining plants with ovate petioled leaves and small, regular, dioecious flowers.

1. DIOSCORE'A L. YAM

1. *D. villo'sa* L. (WILD YAM-ROOT.) A slender twiner, with knotted rootstocks, and net-veined, heart-shaped, 9-11-ribbed, petioled leaves. Flowers dioecious, small, in axillary racemes. Stamens 6. Pod with three large wings.—Reported only from the warm and sheltered valley lying between Hamilton and Dundas, Ont., the banks of the Thames at London, Ont., and Niagara Falls.

AMARYLLIDA'CEAE (AMARYLLIS FAMILY)

Bulbous and scape-bearing herbs, with linear flat root-leaves, and regular and perfect, 6-androus flowers, the tube of the petal-like, 6-parted perianth adherent to the 3-celled ovary. Lobes of the perianth imbricated in the bud. Style single. Anthers introrse.—Represented with us by one species of the genus

1. HYPOX'IS L. STAR-GRASS

1. *H. hirsu'ta* Coville. A small herb sending up a slender scape from a solid bulb. Leaves linear, grass-like, longer than the umbellately 1-4-flowered scape. Perianth hairy and greenish outside, yellowish within, 6-parted nearly down to the ovary. Stamens 6, sagittate. Pod indehiscent, crowned with the withered perianth.—Meadows and open woods.

IRIDA'CEAE (IRIS FAMILY)

Herbs with equitant leaves and perfect flowers. The 6 petal-like divisions of the perianth in 2 (similar or dissimilar) sets of 3 each; the tube adherent to the 3-celled ovary. Stamens 3, distinct or monadelphous, opposite the 3 stigmas, and with anthers extrorse, that is, on the outside of the filaments, facing the divisions of the perianth and opening on that side. Flowers usually showy, from leafy bracts.

Synopsis of the Genera

1. *Iris*. The 3 outer divisions of the perianth reflexed, the 3 inner erect and smaller. Stamens distinct, the anther of each concealed under a flat and petal-like, arching stigma. The styles below adherent to the tube of the perianth. Pod 3-angled. Flowers blue, large and showy. Leaves sword-shaped or grass-like.
2. *Sisyrinchium*. The 6 divisions of the perianth alike, spreading. Stamens monadelphous. Stigmas thread-like. Pod globular, 3-angled. Stems 2-edged. Leaves grass-like. Flowers blue, clustered, from 2 leafy bracts. Plants low and slender.

1. IRIS L. FLEUR-DE-LIS

1. *I. versicolor* L. (LARGER BLUE FLAG.) Stem stout and leafy, from a thickened rootstock. Leaves sword-shaped. Flowers violet-blue, 2 or 3 inches long. Petals much shorter than the sepals.—Wet places.

2. *I. prismatica* Pursh. (SLENDER BLUE FLAG.) Distinguished from the last by its very slender stem and narrowly linear leaves. Tube of the flower very short.—Atl. sea-coast.

3. *I. lacustris* Nutt. (LAKE DWARF IRIS.) Stem low, 3–6 inches high. Inner petals nearly equal to the outer. Tube of the perianth slender, less than an inch long, dilated upwards, rather shorter than the divisions of the perianth. Leaves lanceolate, 3–5 inches long.—Shore of Lake Huron.

2. SISYRINCHIUM L. BLUE-EYED GRASS

1. *S. angustifolium* Mill. A pretty little plant, rather common in moist meadows among grass. The divisions of the delicate blue perianth obovate, notched at the end, and bristle-pointed from the notch. The spathe solitary. Roots fibrous.

2. *S. gramineum* Curtis. Scape 6–18 inches high, and usually bearing 2 or more peduncled spathes.

ORCHIDACEAE (ORCHIS FAMILY)

Herbs well marked by the peculiar arrangement of the stamens, these being gynandrous, that is, borne on or adherent to the stigma or style. There is also usually but a single stamen, of two rather widely separated anthers, but in the last genus of the following list there are 2 distinct stamens, with the rudiment of a third at the back of the stigma.

The Orchids, as a rule, require the aid of insects to convey the pollinia, or pollen-masses, to the stigma, but occasionally it happens that when the anther-cells burst open the pollinia fall forward and dangle in front of the viscid stigma beneath, being sooner or later driven against it either by the wind or by the head of some insect in pursuit of honey. In all cases where the student meets with an Orchid in flower, he should, by experiment, endeavour to make himself acquainted with the method of its fertilization.

The Orchis family is a very large one, there being probably as many as 7,000 different species, but the greater number are natives of tropical regions. Many of them are epiphytes, or air-plants, deriving their support chiefly from the moisture of the air, through their long aerial roots, which never reach the ground. In many species, the perianth, and particularly the labellum, or lip, assumes the most fantastic shapes, making the plants great favourites for hot-house cultivation. In Canada the representatives of this great Order, though not very numerous, are among the most interesting and beautiful of our wild flowers. They are, as a rule, bog plants, and will be found in flower in early summer.

Synopsis of the Genera

- (a) *Fertile anther only one.*
- (b) *Lip with a spur underneath.*
- 1. **Orchis.** The 3 sepals and 2 of the petals erect and arching over the centre of the flower; the lip turned down. The 2 glands or viscid disks of the base of the pollen-masses inclosed in a little pouch just over the concave stigma. Leaves 1 or 2, large. Flowers few, in a spike.
- 2. **Habena'ria.** The lateral sepals usually spreading. The glands or viscid disks of the pollen-masses not inclosed in a covering. Flowers in spikes.
- (b) *Lip without a conspicuous spur.*
- (c) *Plant green. Leaves one or more.*
- (d) *Stem produced from a solid bulb or corm.*
- (e) *Lip saccate.*
- 3. **Calyp'so.** Flower solitary, large, showy, variegated with purple, pink, and yellow. Lip large, inflated, sac-shaped, 2-pointed under the apex. Scape short, from a solid bulb, with a single ovate or slightly heart-shaped leaf below.
- (e) *Lip not saccate.*

4. **Calopo'gon.** Ovary not twisted, the lip consequently turned towards the stem. Flowers large, pink-purple, 2-6 at the summit of the scape; the lip spreading at the outer end, and beautifully bearded above with coloured hairs. Leaf grass-like, only one. Pollen-masses 4.
5. **Arethu'sa.** Flower solitary, large, rose-purple. Lip dilated, recurved, spreading at the end. Sepals and petals lanceolate, nearly alike, arching over the column. Pollen-masses 4. Scape low, sheathed, from a globular solid bulb, with a single linear-nerved leaf hidden in the sheaths of the scape.
6. **Microsty'lis.** Small herbs from solid bulbs; the scape bearing a single leaf and a raceme of minute greenish flowers. Column very small, terete, with 2 teeth at the top, and the anther between them. Petals thread-like or linear, spreading.
7. **Lip'aris.** Small herbs from solid bulbs; the low scape bearing 2 radical leaves and a raceme of a few greenish flowers. Column elongated, incurved, margined at the apex. Petals thread-like or linear, spreading. Anther lid-like.
8. **Aplec'trum.** Somewhat like No. 14, but the perianth is not gibbous below, and the rootstock, instead of being coral-like, is slender, and produces each year a solid bulb or corm. Lip with 3 ridges on the palate. Scape with 3 greenish sheaths below.
 (d) *Stem not produced from a bulb or corm.*
 (f) *Lip saccate.*
9. **Sera'pias.** Stem 1-2 feet high, stout, leafy. Leaves broadly ovate, the upper narrower. Sepals and petals nearly equal, spreading. Lip deeply concave at the base, constricted, and somewhat joined in the middle, dilated and petaloid above. Anther sessile behind the stigma, on a slender jointed base. Ovaries reflexed at maturity.
10. **Epipact'is.** Flowers very much as in *Spiranthes*, but the lip sac-shaped, and without protuberance at the base. Leaves white-veiny, in a tuft at the base of the scape.
 (f) *Lip not saccate.*
11. **Pogo'nia.** Flower solitary, irregular, large, sweet-scented, pale rose-colour or white. Column club-shaped. Lip crested and fringed. Pollen-masses 2. Stem 6-9 inches high, with a single oval or lance-oblong leaf near the middle, and a smaller one, or bract, near the flower.
12. **Spiran'thes.** Spike (of white or whitish flowers) more or less spirally twisted. Sepals and petals narrow and generally connivent. Lip oblong, the lower part embracing the column, and with a protuberance on each side of the base.

13. *Liste'ra*. Flowers small, greenish or brownish-purple, in a spike or raceme. Stem bearing a pair of opposite, sessile, roundish leaves near the middle. Lip flat, mostly drooping, 2-lobed at the apex.
(c) *Plant not green (yellowish or purplish.) Leaves wanting.*
14. *Corallorhi'za*. Brownish or yellowish plants, with the small dull flowers in spikes or scapes, which are leafless or have mere sheaths instead of leaves. Rootstocks branching and coral-like. Perianth gibbous or slightly spurred below. Lip with 2 ridges on the inner part of the face.
(a) *Fertile anthers two. Lip large, sac-like.*
15. *Cypripe'dium*. Lip a large inflated sac, into the mouth of which the style is declined. Sepals and the other petals much alike, the former apparently only 2, two of them being generally united into one under the lip. Leaves large, many-nerved. Flowers solitary or few.

1. ORCHIS L. ORCHIS

1. *O. spectab'ilis* L. (SHOWY ORCHIS.) Scape 4-angled, 4-7 inches high, bearing a few flowers in a spike. The arching upper lip pink-purple, the labellum white, undivided; each flower in the axil of a leaf-like bract, longer than the flower. Leaves 2, radical, shining, oblong-obovate.

2. *O. rotundifo'lia* Banks. Spike few-flowered, loose. Flowers rose-purple, the lip usually white, spotted with purple, 3-lobed, the middle larger and notched, longer than the slender spur. Stem 5-9 inches high, naked and scape-like above, bearing a single roundish leaf at the base.—Bogs and wet woods.

2. HABENA'RIA Willd. REIN ORCHIS. FRINGED ORCHIS

1. *H. clavella'ta* Spreng. Spike few-flowered, the flowers very small, greenish-white. Lip wedge-shaped, truncate and 3-toothed at the apex. Spur slender, longer than the ovary, curved upwards. Stem less than a foot high, slender, with one oblanceolate leaf below and 2 or 3 much smaller ones above.—Wet woods.

2. *H. fla'va* Gray. Stem 10-20 inches high. Spike of small greenish flowers at first dense, with the bracts longer than the flowers, at length long and loose. Lip oblong, almost truncate at the tip; a tooth on each side at the base, and a nasal protuberance on the face. Spur slender, club-shaped. Leaves ovate-oblong or oblong-lanceolate, the upper ones gradually narrowing and passing into bracts.—Wet places.

3. *H. bractea'ta* R. Br. Spike many-flowered, close. Flowers small, greenish. Lip oblong-linear, 2-3-lobed at the tip, much longer than the very short and sac-like spur. Stem 6-12 inches high, leafy, the lower leaves obovate, the upper oblong or lanceolate, gradually reduced to bracts much longer than the flowers.

4. *H. hyperbo'rea* R. Br. Spike many-flowered, long and dense. Flowers small, greenish. Lip lanceolate, entire, about the same length as the slender incurved spur. Stem 6-24 inches high, very leafy, the leaves lanceolate and erect.—Bogs and wet woods.

5. *H. dilata'ta* Gray. Not unlike No. 4, but more slender and with linear leaves and white flowers, and lip with a dilated base.

6. *H. obtusa'ta* Richardson. Stem 4-10 inches high. The solitary leaf is obovate or spatulate-oblong. Spike few-flowered, the flowers greenish-white. Upper sepal broad and rounded, the others and the petals lance-oblong. Lip entire, deflexed, as long as the tapering and curving spur.—Bogs.

7. *H. Hooke'ri* Torr. Spike many-flowered, strict. Flowers yellowish-green, the lip lanceolate, pointed, incurved; petals lance-awl-shaped. Spur slender, acute, nearly an inch long. Stem scape-like above, 2-leaved at the base, the leaves orbicular or elliptical.—Woods.

8. *H. orbicula'ta* Torr. Spike many-flowered, loose and spreading. Flowers greenish-white. Lip narrowly linear, obtuse. Spur curved, about an inch long, thickened towards the apex. Scape 2-leaved at the base, the leaves very large, orbicular or elliptical and lying flat on the ground, shining above, silvery beneath.—Rich woods.

9. *H. blephariglot'tis* Torr. (WHITE FRINGED ORCHIS.) Spike many-flowered, open. Flowers white, very handsome; the lip ovate-lanceolate, fringed, but not lobed, at the apex. Spur long and narrow. Stem a foot high; the leaves oblong or lanceolate, the bracts shorter than the ovaries.—Peat-bogs, etc.

10. *H. leucophae'a* Gray. (GREENISH FRINGED ORCHIS.) Spike as in the last, but the flowers greenish or yellowish-white. Petals obovate, minutely cut-toothed. Lip 3-parted above the stalk-like base, the divisions fan-shaped, fringed. Spur gradually thickened downwards, longer than the ovary. Stem leafy, 2-4 feet high. Leaves oblong-lanceolate; bracts a little shorter than the flowers.—Wet meadows.

11. *H. lac'era* R. Br. (RAGGED FRINGED ORCHIS.) Like the last, but the petals are oblong-linear and entire. The divisions of the lip also are narrow, and the fringe is less copious.—Bogs and rich woods.

12. *H. psycodes* Sw. (PURPLE FRINGED ORCHIS.) Spike cylindrical, many-flowered, the flowers pink-purple, fragrant. Lip fan-shaped, 3-parted above the stalk-like base, the divisions fringed. Spur curved, somewhat thickened downward, very long.—Low grounds.

13. *H. fimbria'ta* R. Br. Resembling the last, but the flowers 3 or 4 times larger, and the petals toothed down the sides. Divisions of the large lip more fringed. Spike loosely flowered.—Wet meadows, chiefly eastward.

14. *H. cilia'ris* R. Br. (YELLOW FRINGED ORCHIS.) Similar to No. 10. Leaves mostly oblong-lanceolate. Flowers orange-yellow. Lip oblong, densely fringed.

15. *H. unalascen'sis* Wats. A slender plant with oblanceolate radical leaves, which wither early. Lip long hastate. Spur club-shaped, shorter than the ovary.

3. CALYP'SO Salisb. CALYPSO

1. *C. bulbo'sa* Oakes. A beautiful little plant growing in mossy bogs. The lip woolly inside; the petals and sepals resembling each other, lanceolate, sharp-pointed. Column winged.

4. CALOPO'GON R. Br. CALOPOGON

1. *C. pulchel'us* R. Br. Leaf linear. Scape a foot high. Flowers an inch across.—Bogs.

5. ARETHU'SA L. ARETHUSA

1. *A. bulbo'sa* L. A beautiful little bog plant, bearing a single large flower (rarely 2), with the oblong lip bearded-crested on the face.

6. MICROSTY'LIS Eaton. ADDER'S MOUTH

1. *M. monophyl'los* Lindl. Leaf sheathing the base of the stem, ovate-elliptical. Raceme spiked, long and slender. Lip pointed, entire.—Cold swamps.

2. *M. unifo'lia* BSP. Leaf near the middle of the stem, ovate, clasping. Raceme short. Lip 3-toothed.—Damp woods, not so common as the last.

7. LIP'ARIS Richard. TWAYBLADE

1. *L. Loese'lii* Richard. Lip yellowish-green, mucronate, shorter than the unequal petals and sepals. Leaves elliptical-lanceolate or oblong, keeled.—Bogs.

8. APLEC'TRUM Torr. PUTTY-ROOT. ADAM-AND-EVE

1. *A. hyema'le* Torr. Scape a foot high. Leaf solitary, ovate, pointed. Perianth greenish-brown; lip whitish, 3-lobed.—Rich mould in woods.

9. SERA'PIAS L.

1. *S. Hellebori'ne* L. Plants 1-2 feet high. Leaves ovate-lanceolate, strongly nerved. Calyx and corolla greenish-purple, lip darker.—Reported first from Montreal but is now not uncommon at Toronto and has also been found in other Ontario localities.

10. EPIPAC'TIS Boehm. RATTLESNAKE PLANTAIN

1. *E. re'pens* Crantz, var. *ophioi'des* A. A. Eaton. Flowers in a loose, 1-sided spike. Lip saccate with a recurved tip. Scape 5-8 inches high. Leaves thickish, petioled, intersected with whitish veins.—Woods, usually under evergreens.

2. *E. tessela'ta* A. A. Eaton. Stem 6-8 inches high, with mostly ovate-lanceolate leaves faintly or distinctly white-marked. Spike loose, not 1-sided; lip saccate with a distinct tip.

3. *E. pubes'cens* A. A. Eaton. Spike not 1-sided, densely many-flowered. Plant rather larger than the last, and the leaves more strongly white-veined. Lip strongly saccate.

4. *E. decip'iens* Ames. Leaves are less white-veiny than the preceding, and the column is continued above the stigma into a conspicuous, awl-shaped beak. The lip, also, is hardly, if at all, saccate below.—North-western part of Ontario.

11. POGO'NIA Juss. POGONIA

1. *P. ophioglossoi'des* Ker. A bog plant. Sepals and petals nearly equal and alike. Root of thick fibres.

12. SPIRAN'THES Richard. LADIES' TRESSES

1. *S. lu'cida* Ames. Flowers white, in 3 ranks, forming a narrow spike. Lip oblong, yellowish on the face, not contracted in the middle, wavy-crisped at the blunt apex. Stem nearly naked. Leaves oblong or lance-oblong.—Moist banks.

2. *S. Romanzoffia'na* Chamisso. Spike dense, oblong or cylindrical. Flowers white, in 3 ranks in the spike. Lip ovate-oblong, contracted below the wavy recurved apex. Sepals and petals all connivent above. Beak of the stigma 2-horned, short. Stem 5–18 inches high, leafy below, leafy-bracted above; the leaves oblong-lanceolate or linear.—Cool bogs.

3. *S. cer'nua* Richard. Resembling the last, but the lower sepals are not upturned or connivent with the upper. Margins of the lower lip much incurved. Gland of the stigma linear, in a long and slender beak.—Wet places.

4. *S. grac'ilis* Beck. Flowers in a single, spirally-twisted rank at the summit of the very slender scape. Leaves with blades all in a cluster at the base, mostly withering before the flowers appear, ovate or oblong. Scape 8–18 inches high.—Sandy plains and pine barrens.

13. LISTE'RA R. Br. TWAYBLADE

1. *L. corda'ta* R. Br. Racemes crowded; pedicels not longer than the ovary. Lip linear, 2-cleft. Column very short. Leaves orbicular-ovate, somewhat heart-shaped.—Damp cold woods.

2. *L. convallarioi'des* Torr. Racemes loose and slender; pedicels longer than the ovary. Lip wedge-oblong, 2-lobed. Column longer than the last.—Damp thickets.

14. CORALLORRHIZA R. Br. CORAL-ROOT

1. *C. trif'ida* Chatelain. Flowers small; the lip whitish, 3-lobed above the base. Spur very small. Stem slender, brownish-yellow, with a few-flowered spike.—Swamps.

2. *C. macula'ta* Raf. Spike many-flowered. Stem purplish, stout. Lip deeply 3-lobed. Spur more prominent than in No. 1.—Dry woods.

3. *C. stria'ta* Lindl. Spike crowded, of numerous large flowers, all the parts of the perianth strikingly marked with 3 dark lines. Lip not lobed. Spur none, but the base of the perianth gibbous.—Rich woods; not common.

15. CYPRIPE'DIUM L. LADY'S SLIPPER. MOCCASIN-FLOWER

A. *The three sepals separate.*

1. *C. arieti'num* R. Br. (RAM'S-HEAD LADY'S SLIPPER.) The smallest species. Stem slender, 6-10 inches high, leafy. Leaves 3 or 4, elliptical-lanceolate, nearly smooth. Lip only $\frac{1}{2}$ inch, red-and-whitish-veiny, prolonged at the apex into a deflexed point.—Swamps; rare.

B. *Two sepals united into one piece under the lip.*

2. *C. parviflo'rum* Salisb. (SMALLER YELLOW LADY'S SLIPPER.) Stem leafy to the top, 1-3-flowered. Lip yellow, flattish above, rather less than an inch long. Sepals and petals wavy-twisted, brownish, pointed, longer than the lip.—Bogs and wet woods.

Var. *pubes'cens* Knight. (LARGER YELLOW LADY'S SLIPPER.) Lip rounded above, larger than the species, but the two are not sufficiently distinct.

3. *C. hirsu'tum* Mill. (SHOWY LADY'S SLIPPER.) Stem 1-2 feet high, 1-2-flowered. Lip very large, white, pinkish in front. Sepals and petals rounded, white, not longer than the lip.—Bogs.

4. *C. can'didum* Muhl. (SMALL WHITE LADY'S SLIPPER.) Stem 6-8 inches high, 1-flowered. Leaves oval, sharp-pointed. Lip white, streaked with purple at the base. Sepals and petals greenish and spotted.

5. *C. acaul'e* Ait. (STEMLESS LADY'S SLIPPER.) Scape naked, 2-leaved at the base, 1-flowered. Lip rose-purple, split down the whole length in front, veiny. Sepals and petals greenish.—Dry or moist woods, under evergreens.

6. *C. passeri'num* Richards. Stem about 8 inches high. Leaves lanceolate. Upper sepals yellowish, lip spherical, pink, spotted with deep crimson within.

CLASS II. DICOTYLE'DONS

Embryo with 2 (rarely more) cotyledons. Parts of the flower usually in fours or fives. Leaves net-veined. Wood in circular layers between the pith and the bark.

PIPERA'CEAE (PEPPER FAMILY)

Herbs with jointed stems and alternate entire leaves. Flowers in spikes; calyx and corolla absent; ovary 1-celled.

1. SAURU'RUS L. LIZARD'S TAIL

1. *S. cer'nuus* L. Herbs with jointed branching stems, 2 feet high. Leaves petioled, heart-shaped, with converging ribs. Flowers white, fragrant, in a dense terminal spike, nodding at the end, each flower with a lanceolate bract. Flowers perfect, but entirely destitute of calyx and corolla. Stamens usually 6 or 7, with long, slender, white filaments. Carpels 3 or 4, slightly united at the base.—Shallow water or swamps.

SALICA'CEAE (WILLOW FAMILY)

Trees or shrubs with dioecious flowers, both sorts in catkins, one under each scale of the catkin. Neither calyx nor corolla present. Fruit 1-celled, many-seeded, the seeds furnished with tufts of down. This Family comprises the Willows and Poplars.

Synopsis of the Genera

1. *Sa'lix*. Trees or shrubs with mostly long and pointed leaves and slender branches. Bracts or scales of the catkins entire or toothed. Stamens mostly 2 under each bract, but in one or two species as many as 5 or 6. Stigmas short. Catkins appearing before or with the leaves.
2. *Pop'ulus*. Trees with broad and more or less heart-shaped leaves. Bracts of the catkins irregularly toothed or cut. Stamens 8–30, or even more, under each scale. Stigmas long. Catkins long and drooping, preceding the leaves.

1. SA'LIX L. WILLOW

A. *Catkins borne on the ends of the short, lateral, leafy branchlets. Scales yellowish, deciduous. Filaments hairy below. Trees or large shrubs with taper-pointed leaves.*

1. *S. ni'gra* Marshall. (BLACK WILLOW.) A tree with a roughish black bark, growing along streams. Leaves narrowly lanceolate, tapering at each end, serrate, smooth, green on both sides. Stamens 3–6. Capsule slender-pedicelled. Sterile catkins long and narrow.

2. *S. amygdaloi'des* Anders. (PEACH-LEAVED WILLOW.) A tree with lanceolate or ovate-lanceolate leaves, pale or glaucous

beneath, and with long slender petioles. Fertile catkins becoming very loose from the lengthening of the pedicels. Stamens 3-6.—Pr. Prov.

3. *S. lu'cida* Muhl. (SHINING WILLOW.) A shrub or small bushy tree, growing along streams. Leaves ovate-oblong or narrower, with a long tapering point, shining on both sides, serrate. Stamens mostly 5. Scales of catkins dentate, hairy at the base. Sterile catkins densely flowered, showy.

4. *S. frag'ilis* L. (CRACK WILLOW.) A tall and handsome tree with brittle twigs. Leaves lanceolate, long-pointed, pale or glaucous beneath, 3-6 inches long. Stamens mostly 2, rarely 3-4. Capsule short-pedicelled. (Introd.)

5. *S. alba* L. Leaves ashy-gray or white on both sides owing to silky hairs, lanceolate, long and slender-pointed, finely serrate. Stamens 2. Capsule sessile or nearly so. (Introd.)

Var. *vitelli'na* Koch. The old leaves are glabrous and bluish-white beneath. The twigs are yellow or reddish in colour.

6. *S. longifo'lia* Muhl. (SAND-BAR WILLOW.) A shrub or small tree, varying greatly in size, growing along streams in sandy or gravelly places, and often forming dense clumps. Leaves linear-lanceolate, very long, tapering towards both ends, nearly sessile, serrate with a few spreading teeth, grayish-hairy when young. Stamens 2.

B. Catkins lateral or terminal. Scales coloured at the tip, persistent. Stamens 2, the filaments not hairy. Shrubs or small trees.

(1) *Capsules woolly.*

7. *S. dis'color* Muhl. (GLAUCOUS WILLOW.) A shrub or small tree, 8-15 feet high, growing in low grounds and along streams. Leaves lanceolate or ovate-lanceolate, irregularly toothed in the middle of the margin, entire at each end, smooth and bright green above, white-glaucous beneath, when young. Stipules moon-shaped, toothed. Catkins sessile, very early in spring before the leaves. Scales very dark and hairy.

Var. *erioceph'ala* Anders. Densely flowered and very silky catkins. Leaves somewhat pubescent even when old.

8. *S. rostra'ta* Richards. A good-sized shrub, chiefly in moist situations. Leaves wrinkled, oblong or obovate-lanceolate, barely

toothed, downy above, very veiny, hairy and glaucous beneath. Stipules semi-lunar, toothed. Capsule at length raised on a very slender stalk, catkins appearing with the leaves.

9. *S. hu'milis* Marshall. (PRAIRIE WILLOW.) A grayish shrub, 3-8 feet high, growing usually in dry or barren places. Leaves oblanceolate, pointed, the lowest obovate, slightly downy above, thickly so beneath. Stipules semi-ovate or moon-shaped, with a few teeth, shorter than the petioles. Catkins ovoid, sessile, before the leaves, naked at the base. Scales dark red or brownish.

10. *S. seric'ea* Marsh. Large shrubs growing in wet places. Leaves narrowly lanceolate, silky-pubescent when young, finely serrate. Fertile catkins densely flowered; capsule ovoid, round-tipped, pedicel about as long as the scale.

11. *S. petiola'ris* Smith. (PETIOLED WILLOW.) A low shrub on sandy river-banks. Leaves narrowly lanceolate, finely and evenly serrate, smooth, silky when young. Catkins with a few small, leaf-like bracts at the base. Scales of the fertile catkins acute, very hairy. Capsule tapering, silky, stalked.

12. *S. can'dida* Flügge. (HOARY WILLOW.) A shrub not more than 3 or 4 feet high, growing in bogs and wet places; the twigs and leaves clothed with a web-like wool, giving the whole plant a whitish aspect. Leaves lanceolate, narrow, with somewhat revolute margins. Stipules small, lanceolate, toothed. Catkins cylindrical. Anthers red.

(2) *Capsules glabrous.*

13. *S. corda'ta* Muhl. (HEART-LEAVED WILLOW.) A shrub or small tree, growing in wet grounds. Leaves lanceolate, not always heart-shaped, sharply serrate, smooth, green both sides. Catkins cylindrical, rather slender, leafy-bracted at the base, the sterile ones silky.

14. *S. balsamif'era* Barratt. A small, much-branched shrub. Young twigs shining - chestnut on the sunny side. Leaves ovate-lanceolate, usually slightly cordate at base, at first very thin and of a reddish colour, at length rigid, dark green above, and paler and conspicuously reticulate-veined beneath, slightly serrate, with slender petioles. Sterile catkins very silky, with a few bracts at the base; fertile catkins leafy-peduncled, becoming very

loose in fruit. Capsules long-pedicelled.—Atl. Prov. and Northern Canada.

15. *S. pedicella'ris* Pursh. Low shrub, 1–3 feet high, growing in swamps and peat-bogs. Leaves elliptic-obovate, an inch long, entire, smooth, somewhat coriaceous when mature, revolute, reticulated, pale or glaucous beneath. Fertile catkins loosely few-flowered, on long leafy peduncles. Capsules glabrous, on slender pedicels.

2. *POP'ULUS* L. POPLAR

1. *P. tremuloi'des* Michx. (AMERICAN ASPEN.) A tree with a greenish-white bark, and roundish-heart-shaped leaves, continually in a state of agitation, due to the lateral compression of the petiole, and the consequent susceptibility of the leaf to the least motion of the air. Teeth of the leaves small.

2. *P. grandidenta'ta* Michx. (LARGE-TOOTHED ASPEN.) Roundish-ovate leaves with large, irregular, sinuate teeth.

3. *P. balsamif'era* L. (BALSAM POPLAR.) A tall tree, growing in swamps and along streams; the large buds varnished with resinous matter. Leaves ovate, tapering, finely serrate with rounded teeth, whitish beneath. Stamens very numerous.

4. *P. can'dicans* Ait. (BALM OF GILEAD.) Leaves broader and more or less heart-shaped, with lower surfaces and petioles hairy. (Intro.)

5. *P. deltoi'des* Marsh. (COTTONWOOD.) A tree with broadly deltoid leaves, slightly heart-shaped, serrate with incurved teeth. Young branches slightly angled, at length round. Fertile catkins very long, the scales cut-fringed, not hairy.

MYRICA'CEAE (SWEET-GALE FAMILY)

Shrubs with monoecious or dioecious flowers, both sterile and fertile ones collected in short catkins or heads. Sepals none or rudimentary; petals none. Leaves with resinous dots, usually fragrant. Fruit a 1-seeded, dry, drupe-like nut, usually coated with little waxy grains.

1. *Myri'ca*. Flowers dioecious or monoecious, catkins lateral, each bract with a pair of bractlets underneath. Stamens in the sterile flowers 2–8. Ovary solitary in the fertile flowers, 1–2-celled, tipped with 2 thread-like stigmas, and surrounded by 2–8 small scales at the base.

1. MYRTICA L.

1. *M. Gale* L. (SWEET GALE.) A shrub 3-4 feet high. Leaves wedge-lanceolate, serrate towards the apex, pale. The small nuts in crowded heads, and winged by the 2 scales.—Bogs.

2. *M. cerifera* L. (BAYBERRY. WAX-MYRTLE.) Leaves oblong-lanceolate, entire, or wavy-toothed towards the apex, shining and sprinkled with resinous dots on both sides, fragrant. Sterile catkins scattered. Nuts naked, bony, and covered with white wax.—Sandy soil near the coast, Atl. Prov.

3. *M. asplenifolia* L. (SWEET FERN.) A shrub 1-2 feet high, sweet-scented. Leaves linear-lanceolate in outline, deeply pinnatifid, the lobes numerous and rounded.—Dry soil; especially in pine barrens.

JUGLANDACEAE (WALNUT FAMILY)

Trees with alternate pinnate leaves and no stipules. Flowers monoecious. Sterile flowers in catkins. Fertile flowers solitary or in small clusters, with a regular, 3-4-lobed calyx adherent to the ovary. Fruit a sort of drupe, the fleshy outer layers at length becoming dry and forming a husk, the inner layer hard and bony and forming a nut-shell. Seed solitary in the fruit, very large and 4-lobed. This Family comprises the Walnuts, Butternuts, and Hickories.

Synopsis of the Genera

1. *Juglans*. Sterile flowers in long, drooping, solitary catkins from the previous year's wood. Filaments of the numerous stamens very short. Fertile flowers single or in small clusters on peduncles at the ends of the branches. Calyx 4-toothed, with small petals at the sinuses. Styles and stigmas 2, the latter fringed. Exocarp (epicarp + mesocarp) drying without splitting. Shell of the nut very rough and irregularly furrowed.
2. *Ca'rya*. Sterile flowers in slender clustered catkins. Stamens 3-10, with very short filaments. Fertile flowers in small clusters at the ends of the branches. Calyx 4-toothed; no petals. Stigmas 2 or 4, large. Husk (involucre) 4-valved, drying and splitting away from the very smooth and bony nut-shell.

1. JU'GLANS L. WALNUT

1. *J. cine'rea* L. (BUTTERNUT.) Leaflets oblong-lanceolate, pointed, serrate. Petioles and branchlets clammy. Fruit ellipsoidal, clammy.—Rich woods.

2. *J. ni'gra* L. (BLACK WALNUT.) Leaflets ovate-lanceolate, taper-pointed, serrate. Petioles downy but not clammy. Fruit spherical. Wood a darker brown than in the Butternut.—Rich woods.

2. CA'RYA Nutt. HICKORY

1. *C. ova'ta* K. Koch. (SHELL-BARK or SHAG-BARK HICKORY.) Leaflets 5-7, the lower pair much smaller than the others. Husk of the fruit thick and woody, splitting completely into 4 valves. Nut flattish-globular, mucronate, shell thinnish. Bark of the trunk rough, scaling off in rough strips.—Rich woods.

2. *C. lacinio'sa* Loud. (BIG SHELL-BARK.) Resembles No. 1, but leaflets are 7-9. Fruit ovoid, 4-ribbed above the middle. Husk very thick. Nut large ($1\frac{1}{4}$ -2 inches long), thick-walled, usually pointed at both ends.

3. *C. alba* K. Koch. (WHITE-HEART HICKORY or MOCKER NUT.) Leaflets 7-9. Bark close, rough but not shaggy, and not scaling off on the old trunks. Husk as in the last. Catkins, shoots, and lower surface of the leaves tomentose when young. Nut ovoid or globular, very thick-walled.—Sparingly found in the Niagara district and south-westward.

4. *C. cordifor'mis* K. Koch. (SWAMP HICKORY or BITTERNUT.) Leaflets 7-11. Husk of the fruit thin and friable, splitting half-way down, or slowly to the base. Nut spherical, short-pointed. Bark smooth, not scaling off.—Moist ground.

5. *C. gla'bra* Spach. (PIG-NUT. BROOM HICKORY.) Leaflets 5-7. Shoots, etc., glabrous. Husk as in the last. Nut thick-shelled, oblong or oval.

BETULA'CEAE (BIRCH FAMILY)

Trees or shrubs with monoecious, rarely dioecious, flowers without corolla, both sorts in catkins, 2 or 3 flowers under each scale or bract of the catkin. Ovary 2-celled and 2-ovuled, but in fruit only 1-celled and 1-seeded. Fruit a small nut. Stigmas 2, long and slender. Twigs and leaves often aromatic.

Synopsis of the Genera

1. *Bet'ula*. Sterile catkins long and pendulous, formed during summer and expanding the following spring; each flower consisting of a calyx of one small scale to which is attached 4 short filaments; 3 flowers under each scale of the catkin.

Fertile catkins stout, oblong, the scales or bracts 3-lobed and with 2 or 3 flowers under each; each flower without a calyx, consisting only of a naked ovary, becoming a winged nutlet in fruit. Bark easily coming off in sheets.

2. **Al'nus.** Catkins much as in *Betula*, but each fertile and sterile flower has a distinct, 3-5-parted calyx. Catkins solitary or clustered at the ends of leafless branchlets or peduncles. Nutlets wingless or with a thin wing.
3. **Cor'yus.** Sterile flowers in drooping catkins. No calyx. Stamens 8 (with 1-celled anthers), and 2 small bractlets under each bract. Fertile flowers in a small scaly head; 1 ovary, surmounted by 2 long red stigmas, under each scale, and accompanied by a pair of bractlets which, in fruit, enlarge and form a leaf-like or tubular, fringed or toothed involucre closely enveloping each nut. Sterile catkins from the axils of the previous year. Fertile flowers terminating the new shoots.
4. **Os'trya.** Sterile flowers in drooping catkins. Calyx wanting. Stamens several under each bract, but not accompanied by bractlets. Fertile flowers in short catkins, 2 under each bract, each ovary tipped with 2 long stigmas, and surrounded by a tubular bractlet, which, in fruit, becomes a greenish-white, inflated bag, having the small nut in the bottom.
5. **Carpi'nus.** Sterile flowers in drooping catkins. Calyx wanting. Stamens several under each bract; no bractlets. Fertile flowers much as in *Ostrya*, but the bractlet subtending the ovary is not tubular but open, and in fruit becomes a 3-lobed leaflet with a small, ribbed, ovate nut in its axil.

1. BET'ULA L. BIRCH

1. **B. lenta** L. (CHERRY BIRCH. SWEET or BLACK BIRCH.) Bark of the trunk dark brown, close, aromatic; that of the twigs bronze-coloured. Wood rose-coloured. Leaves ovate, with somewhat heart-shaped base, doubly serrate, pointed, short-petioled. Fruiting catkins sessile, thick, oblong-cylindrical.—Atl. Prov. and reported from Ont.

2. **B. lu'tea** Michx. f. (YELLOW or GRAY BIRCH.) Bark of the trunk yellowish-gray, somewhat silvery, scaling off in thin layers. Leaves oblong-ovate, hardly at all heart-shaped. Fruiting catkins thicker and shorter than in No. 1.—Moist woods.

3. **B. populifo'lia** Marsh. (AMERICAN WHITE BIRCH. GRAY BIRCH.) Leaves very tremulous on slender petioles, triangular, very taper-pointed, nearly truncate at the base, smooth and shining except when young. Bark of trunk white, less separable than in *Canoe Birch*.—Poor soil, Atl. Prov.

4. *B. papyrif'era* Marsh. (PAPER OR CANOE BIRCH.) Bark of the trunk white, easily separating in sheets. Leaves ovate, taper-pointed, heart-shaped, long-petioled, coarsely serrate. Fruiting catkins cylindrical, usually hanging on slender peduncles.

5. *B. pu'mila* L. (LOW BIRCH.) A shrub with brownish bark, not glandular. Leaves ovate or roundish, pale beneath; veinlets on both surfaces finely reticulated. Catkins mostly erect, on short peduncles.—Bogs and low grounds, northward.

2. AL'NUS Hill. ALDER

1. *A. inca'na* Moench. (SPECKLED OR HOARY ALDER.) A shrub or small tree, growing in thickets in low grounds along streams. Leaves oval or ovate, rounded at the base, serrate, whitish beneath. Flowers preceding the leaves in early spring, from clustered catkins formed the previous summer and remaining naked over winter. Fruit orbicular, wingless.

2. *A. cris'pa* Pursh. (GREEN OR MOUNTAIN ALDER.) A shrub 3-8 feet high, along mountain streams. Leaves oval to ovate, slightly heart-shaped. Flowers appearing with the leaves, the staminate catkins having remained naked during the winter, the pistillate inclosed in a scaly bud. Fruit with a thin wing.—Northward.

3. COR'YLUS L. HAZEL-NUT. FILBERT

1. *C. america'na* Walt. (WILD HAZEL-NUT.) Leaves roundish-heart-shaped. Involucre spreading above, leaf-like and cut-toothed.

2. *C. rostra'ta* Ait. (BEAKED HAZEL-NUT.) A rather common shrub, easily distinguished from No. 1 by the involucre, which is prolonged into a narrow tube much beyond the nut, and is densely bristly-hairy.

4. OS'TRYA Scop. HOP HORNBEAM. IRONWOOD

1. *O. virginia'na* K. Koch. (IRON-WOOD.) A slender tree with brownish furrowed bark. Leaves oblong-ovate, taper-pointed, sharply doubly serrate. Fertile catkin like a hop in appearance. Wood very hard and close.—Rich woods.

5. CARPI'NUS L. HORNBEAM

1. *C. carolinia'na* Walt. (BLUE OR WATER BEECH.) Small trees with furrowed trunks and close, smooth, gray bark. Leaves ovate-oblong, pointed, doubly serrate.—Resembling a beech in general aspect.

FAGA'CEAE (BEECH FAMILY)

Shrubs or trees with alternate simple leaves, deciduous stipules, and monoecious flowers, without petals. Sterile flowers in catkins (but in Beech in small heads); the fertile ones solitary or clustered, and furnished with an involucre which forms a scaly cup or a bur surrounding the nut.

Synopsis of the Genera

1. *Quer'cus*. Sterile flowers with a 2-8-parted calyx including few or several stamens with slender filaments. Fertile flowers scattered or somewhat clustered, each in a scaly involucre or cupule. Nut (acorn) rounded, the base inclosed by the cupule.
2. *Casta'nea*. Sterile flowers in long slender catkins. Calyx 6-parted. Fertile flowers usually 3 in each involucre, the latter prickly, forming a bur. Calyx 6-lobed. Stigmas bristle-shaped. Nuts inclosed (mostly 2 or 3 together) in the prickly, 4-valved involucre, flattened when there are more than one.
3. *Fa'gus*. Sterile flowers in a small head on drooping peduncles. Calyx bell-shaped. Fertile flowers usually in pairs in the involucre, which consists of awl-shaped bractlets grown together at the bases. Calyx lobes awl-shaped. Nuts 3-angled, generally in pairs in the bur-like, 4-valved cupule. Bark close, smooth, and light gray.

1. QUER'CUS L. OAK

A. *Acorns ripening the first year, and therefore borne on the new shoots. Lobes or teeth of the leaves not bristle-pointed.*

1. *Q. alba* L. (WHITE OAK.) A large tree. Leaves (when mature) smooth, bright green above, whitish beneath, obliquely cut into few or several oblong, mostly entire lobes. The oblong nut much larger than the saucer-shaped, rough cupule.—Rich woods.

2. *Q. macrocar'pa* Michx. (BUR OAK. MOSSY-CUP WHITE OAK.) A medium-sized tree. Leaves obovate, deeply lobed on the lower half, smooth above, pale or downy beneath. Acorn broadly ovoid, half or altogether covered by the deep cup, the upper scales of which taper into bristly points, making a fringed border. Cup varying greatly in size, often very large.—Rich soil.

3. *Q. bi'color* Willd. (SWAMP WHITE OAK.) A tall tree. Leaves sinuate-toothed, but hardly lobed, wedge-shaped at the base, downy or hoary beneath, the main veins 6-8 pairs. Cup

nearly hemispherical, about half as long as the oblong-ovoid acorn, sometimes with a fringed border. Peduncles in fruit longer than the petiole.—Low grounds.

4. *Q. Muhlenbergii* Engelm. (CHESTNUT OAK.) A tall tree with thin bark that flakes off. Leaves oblong-lanceolate, slender-petioled, acute, sharply toothed. Nut globular to ovoid. Cup shallow and thin, covering about one-half the nut.

5. *Q. prinoides* Willd. (DWARF CHESTNUT OAK.) A low shrub, 3–9 feet high, with small leaves on rather short petioles; otherwise like No. 4, but with deeper cups.

B. *Acorns ripening the second year, and therefore borne on the previous year's wood, below the leaves of the season. Lobes or teeth of the leaves bristle-pointed.*

6. *Q. velutina* Lam. (QUERCITRON. YELLOW-BARKED OR BLACK OAK.) Leaves usually deeply pinnatifid, slender-petioled, rather rounded at the base, rusty-downy when young, smooth and shining above when mature, often slightly pubescent beneath, turning brownish, orange or dull red in autumn. Cup somewhat hemispherical with a conical base. Acorn rather small. The bark dark brown and rough, and yellow or orange inside.

7. *Q. rubra* L. (RED OAK.) A large tree. Leaves moderately pinnatifid, turning dark red in the autumn. Cup saucer-shaped, sessile or nearly so, very much shorter than the oblong-ovoid acorn.

8. *Q. palustris* Muench. (PIN OAK.) A medium-sized tree. Cup flat-saucer-shaped, very much shorter than the ovoid-globose acorn, which is about $\frac{1}{2}$ inch long. Leaves deeply pinnatifid, with divergent lobes and rounded sinuses.

2. CASTANEA Hill. CHESTNUT

1. *C. dentata* Borkh. (CHESTNUT.) A large tree. Leaves oblong-lanceolate, pointed, coarsely and sharply serrate, acute at the base. Nuts 1–3 in each bur.—South-western Ontario.

3. FAGUS L. BEECH

1. *F. grandifolia* Ehrh. (AMERICAN BEECH.) A very common tree in rich woods, the branches horizontal. Leaves oblong-ovate, taper-pointed, toothed, the very straight veins terminating in the teeth.

URTICA'CEAE (NETTLE FAMILY)

Herbs, shrubs, or trees, with monoecious or dioecious (or, in the Elms, sometimes perfect) flowers without corolla but with a regular calyx free from the 1-2-celled ovary, which becomes a 1-seeded fruit. Stamens opposite the lobes of the calyx.

TRIBE I. UL'MEAE

Trees with alternate simple leaves, and deciduous small stipules. Flowers often perfect. Styles 2. Fruit a samara winged all around.

1. **Ul'mus.** Flowers in lateral clusters, earlier than the leaves, purplish or greenish-yellow. Calyx bell-shaped, 4-9-cleft. Stamens 4-9; the filaments long and slender. Anthers extrorse. Ovary 1-2-celled, but the samara only 1-seeded. Stigmas 2.

TRIBE II. CELTID'EAE

2. **Cel'tis.** Flowers greenish, polygamous, the pistillate solitary or in pairs, appearing with the leaves. Calyx 5-6-parted, persistent. Stamens 5-6. Anthers introrse. Stigmas 2, long and pointed and recurved. Ovary 1-ovuled. Fruit a drupe.

TRIBE III. MO'REAE (BREAD-FRUIT AND FIG FAMILY)

Flowers monoecious or dioecious, crowded in catkin-like spikes or heads, the whole pistillate catkin becoming an aggregate fruit from the enlargement of the calyx in the several flowers. Calyx 4-parted. Stamens 4. Ovary 2-celled, 1-cell eventually disappearing. Styles 2.

3. **Mo'rus.** Pistillate and staminate flowers in separate catkins. Trees with milky juice and rounded leaves. Staminate spikes slender.

TRIBE IV. URTIC'EAE (NETTLE FAMILY)

Herbs with watery juice and opposite or alternate leaves, often beset with stinging hairs. Flowers monoecious or dioecious, in spikes or racemes. Stamens as many as the sepals. Style only 1. Ovary 1-celled. Fruit an achene.

4. *Urti'ca*. Leaves opposite. Plant beset with stinging hairs. Sepals 4 in both sterile and fertile flowers. Stamens 4. Stigma a small sessile tuft. Achene flat inclosed between the 2 larger sepals. Flowers greenish.
5. *Laport'ea*. Leaves alternate. Plant beset with stinging hairs. Sepals 5 in the sterile flowers, 4 in the fertile, 2 of them much smaller than the other 2. Stigma awl-shaped. Achene flat, very oblique, reflexed on its winged pedicel.
6. *Pil'ea*. Leaves opposite. Whole plant very smooth and semi-transparent. Sepals and stamens 3-4. Stigma a sessile tuft.
7. *Boehme'ria*. Leaves mostly opposite. No stinging hairs. Sepals and stamens 4 in the sterile flowers. Calyx tubular or urn-shaped in the fertile ones, and inclosing the achene. Stigma long and thread-like.
8. *Parieta'ria*. Leaves alternate, entire, 3-ribbed. No stinging hairs. Flowers polygamous, in involucrate-bracted, cymose, axillary clusters. Calyx of the pistillate flowers tubular or bell-shaped, 4-lobed. Stigma tufted. Staminate flowers nearly as in the last.

TRIBE V. CANNABIN'EAE (HEMP FAMILY)

Rough herbs with watery juice and tough bark. Leaves opposite and palmately divided or lobed. Flowers dioecious. Sterile ones in compound racemes; stamens 5; sepals 5. Fertile ones in crowded clusters; sepal only 1, embracing the achene. Stigmas 2.

9. *Can'nabis*. A rather tall, rough plant with palmately compound leaves of 5-7 linear-lanceolate, serrate leaflets. Fertile flowers spiked-clustered.
10. *Hu'mulus*. Leaves 3-5-lobed. Plant climbing. Fertile flowers in a short spike, forming a membranaceous catkin in fruit.

1. UL'MUS L. ELM

1. *U. ful'va* Michx. (RED or SLIPPERY ELM.) Flowers nearly sessile. Leaves very rough above, taper-pointed. Buds downy with rusty hairs. A medium-sized tree with mucilaginous inner bark.

2. *U. america'na* L. (AMERICAN or WHITE ELM.) Leaves not rough above, abruptly pointed. Flowers on drooping pedicels. Buds glabrous. A large ornamental tree with drooping branchlets. —Moist woods.

3. *U. racemo'sa* Thomas. (ROCK OR CORK ELM.) Resembling the last, but the bud-scales are downy-ciliate, the branches corky, and the flowers racemed.

2. CEL'TIS L. NETTLE-TREE. HACKBERRY

1. *C. occidenta'lis* L. (SUGARBERRY.) A medium-sized tree of Elm-like aspect. Leaves reticulated, ovate, taper-pointed, serrate, more or less oblique at the base. Fruit as large as a pea, dark purple when ripe, the flesh thin.—Low grounds; a few trees here and there through Ontario.

3. MO'RUS L. MULBERRY

1. *M. ru'bra* L. (RED MULBERRY.) Leaves heart-ovate, rough above, downy beneath, pointed. Fruit red, turning dark purple, long.—Niagara district, and south-westward.

2. *M. alba* L. (WHITE MULBERRY.) Leaves smooth and shining. Fruit white-pinkish.—South-western Ontario.

4. URTI'CA L. NETTLE

1. *U. grac'ilis* Ait. Stem slender, 2-6 feet high. Leaves ovate-lanceolate, pointed, serrate, 3-5-nerved from the base, nearly smooth, the long petioles with a few bristles. Flower-clusters in slender spikes.—Moist ground and along fences.

2. *U. dioic'a* L. (STINGING NETTLE.) Plant bristly with very stinging hairs. Leaves ovate, cordate, very deeply serrate. Spikes branching.—Waste places.

3. *U. u'rens* L. Leaves elliptical or ovate, coarsely and deeply serrate with spreading teeth, petioled. Flower-clusters 2 in each axil, composed of both staminate and pistillate flowers.—Waste grounds, Atl. Prov.

5. LAPORTEA Gaudichaud. WOOD-NETTLE

1. *L. canaden'sis* Gaudichaud. Stem 2-3 feet high. Leaves large, ovate, long-petioled, a single 2-cleft stipule in the axil.—Moist woods.

6. PIL'EA Lindl. RICHWEED. CLEARWEED

1. *P. pu'mila* Gray. Stem 3-18 inches high. Leaves ovate, coarsely-toothed, 3-ribbed.—Cool moist places.

7. BOEHMERIA Jacq. FALSE NETTLE

1. *B. cylindrica* Sw. Stem 1-3 feet high, smoothish. Leaves ovate-oblong or ovate-lanceolate, serrate, 3-nerved, long-petioled. Stipules separate.—Moist shady places.

8. PARIETARIA L. PELLITORY

1. *P. pennsylvanica* Muhl. A low annual, simple or sparingly branched, minutely downy. Leaves oblong-lanceolate, thin, veiny, roughish with opaque dots.—Usually in crevices of limestone rocks; not very common.

9. CANNABIS L. HEMP

1. *C. sativa* L. (HEMP.) Common everywhere along roadsides and in waste places.

10. HUMULUS L. HOP

1. *H. lupulus* L. (COMMON HOP.) A twining perennial. Leaves heart-shaped, mostly 3-5-lobed, petioled. Calyx of fertile flower a single sepal. In fruit the calyx, achene, etc., sprinkled with yellow resinous grains, which give the hop its taste and smell.

SANTALACEAE (SANDALWOOD FAMILY)

Low, herbaceous or partly woody plants (with us) with perfect flowers, these greenish-white, in terminal or axillary, corymbose clusters. Calyx bell-shaped or urn-shaped, 4-5-cleft, adherent to the 1-celled ovary, lined with a 5-lobed disk, the stamens on the edge of the latter between its lobes and opposite the lobes of the calyx, to which the anthers are attached by a tuft of fine hairs. Fruit nut-like, crowned with the persistent calyx lobes.

1. COMANDBRA Nutt. BASTARD TOAD-FLAX

1. *C. umbellata* Nutt. Stem 8-10 inches high, leafy. Leaves oblong, pale green, an inch long. Flower-clusters at the summit of the stem. Calyx tube prolonged and forming a neck to the fruit. Style slender.—Dry soil.

2. *C. livida* Richardson. Peduncles axillary, slender, several-flowered. Leaves oval, alternate, almost sessile. Fruit pulpy when ripe, red.—Boggy barrens near the Atl. coast, and Pr. Prov.

3. *C. pal'lida* A. DC. Leaves glaucous, linear to narrowly lanceolate, acute. Fruit ovoid, sessile or on short stout pedicels.—Pr. Prov.

4. *C. Richardsia'na* Fernald. Flowering stems 4–10 inches high, very leafy, leaves green, lanceolate-ovate; inflorescence a few-flowered corymb.—Pr. Prov.

LORANTHA'CEAE (MISTLETOE FAMILY)

Dwarf plants, parasitic on the twigs of Conifers; the stems slender, scattered or clustered.

1. ARCEUTHO'BIUM BIEB.

1. *A. pusil'lum* Peck. Dwarf, $\frac{1}{2}$ – $\frac{3}{4}$ inch in height, greenish-brown in colour. On *Picea*.

ARISTOLOCHIA'CEAE (BIRTHWORT FAMILY)

Herbs with perfect flowers, the tube of the 3-lobed calyx adherent to the 6-celled, many-seeded ovary. Leaves heart-shaped or kidney-shaped, on long petioles from a thick rootstock. Stamens 12 or 6. Flowers solitary. Calyx dull-coloured, the lobes valvate in the bud.

1. AS'ARUM L. WILD GINGER

1. *A. canadense* L. Radiating stigmas 6. Leaves only a single pair, kidney-shaped, and rather velvety, the peduncle in the fork between the petioles, close to the ground. Rootstock aromatic. Calyx brown-purple inside, the spreading lobes pointed. Corolla wanting or very rudimentary.—Rich woods.

2. ARISTOLO'CHIA L.

1. *A. macrophyl'la* Lam. (DUTCHMAN'S PIPE.) A glabrous twining shrub, with large, kidney-shaped leaves. Flowers reddish-purple with the calyx tube curved like a Dutchman's pipe. (Introd.)

POLYGONA'CEAE (BUCKWHEAT FAMILY)

Herbs, well marked by the stipules of the alternate leaves being in the form of membranous sheaths above the usually swollen joints of the stem (these obsolete in one Genus). Flowers usually

perfect. Calyx 4-6-parted. Corolla wanting. Stamens 3-9, inserted on the base of the calyx. Stigmas 2 or 3. Ovary 1-celled, with a single ovule rising from the base, forming a little nutlet.

Synopsis of the Genera

A. *Flowers involucrate. Stamens 9. Stipules obsolete.*

1. *Eriog'onum*. Involucre 4-8-toothed, the flowers exserted. Calyx 6-parted, coloured (yellow in ours), inclosing the achene.

B. *Flowers not involucrate. Stamens 4-8.*

2. *Polyg'onum*. Sepals 5 (occasionally 4), often coloured and petal-like, persistent, embracing the 3-angled (or sometimes flattish) nutlet or achene. Flowers in racemes or spikes, or sometimes in the axils.
3. *Ru'mex*. Sepals 6, the 3 outer ones herbaceous and spreading in fruit, the 3 inner (called valves) somewhat petal-like and, after flowering, convergent over the 3-angled achene, often with a grain-like projection on the back. Stamens 6. Styles 3. Flowers usually in crowded whorls, the latter in paniced racemes.
4. *Fagopy'rum*. Calyx 5-parted, petal-like. Stamens 8, with 8 yellow glands between them. Styles 3. Achenes 3-angled. Flowers white, in panicles. Leaves triangular-heart-shaped or halberd-shaped.

1. ERIOG'ONUM Michx.

1. *E. fla'vum* Nutt. Woolly throughout, a few inches high. Leaves oblanceolate. Umbel of 3-9 rays, on a naked peduncle. Flowers yellow, silky.—Pr. Prov.

2. POLYG'ONUM L. KNOTWEED

A. *Flowers along the stem, inconspicuous, greenish-white, nearly sessile in the axils of the small leaves. Sheaths cut-fringed or torn.*

1. *P. marit'imum* L. (COAST KNOTGRASS.) Prostrate with stout stems, glaucous. Leaves thick, oval to narrowly oblong. Flowers in the axils of leaves, clustered. Stipules large, silvery. Stamens 8. Achenes smooth and shining, projecting above the calyx.—Sea-coast.

2. *P. avicula're* L. (KNOTGRASS. GOOSEGRASS.) A weed everywhere in yards and waste places. Stem prostrate and spreading. Stamens 5-8. Achene 3-sided, dull. Stigmas 3. Leaves sessile, lanceolate or oblong.

3. *P. erectum* L. A stouter weed than No. 2, erect. Leaves broader, elliptical. Achene dull.

4. *P. ramosissimum* Michx. Erect or ascending, yellowish-green. Leaves lanceolate to linear, acute, very small above. Sepals mostly 6; stamens 3-6. Achene smooth and shining.—Chiefly westward; sandy places.

5. *P. tenue* Michx. Stem slender, upright, sparingly branched, angled. Leaves sessile, narrowly linear, very acute, 3-nerved. Stamens 8. Achene dull black.—Dry soil and rocky places.

B. *Flowers in terminal spikes or racemes, mostly rose-coloured or pinkish, occasionally greenish.*

(1) *Leaves not heart-shaped or arrow-shaped.*

6. *P. lapathifolium* L. Sheaths not fringed, stem nearly smooth, 3-6 feet high. Leaves long, tapering from near the base to a narrow point, rough on the midrib and margins. Spikes oblong to linear and erect or nearly so. Stamens 6. Styles 2. Achene flat or hollow-sided.—In muddy places along streams and ponds.

7. *P. pennsylvanicum* L. Sheaths not fringed. Stem 1-3 feet high, the upper branches and the peduncles bristly with stalked glands. Spikes thick, erect, short. Flowers rose-coloured. Stamens 8. Achene flat, orbicular.—Low open grounds.

8. *P. persicaria* L. (LADY'S THUMB.) Sheaths with a somewhat ciliate border. Stem nearly smooth, a foot or more in height. Leaves with a dark blotch on the middle of the upper surface. Spikes dense, erect, on naked peduncles. Stamens 6. Achene flat or 3-angled, according as the stigmas are 2 or 3, shining.—Very common near dwellings in moist ground.

9. *P. amphibium* L. (WATER PERSICARIA.) Spike of flowers dense, oblong, showy, rose-red. Stem floating in shallow water or rooting in soft mud, nearly glabrous. Leaves long-petioled, often floating. Sheaths not bristly-fringed. Stamens 5. Stigmas 2.—In shallow water, mostly northward.

Var. *Hartwrightii* Bissell. Distinguished from *P. amphibium* by its foliaceous and ciliate sheaths.—Muddy margins of ponds and lakes.

10. *P. Muhlenbergii* Watson. Differs from the last in being rough with appressed hairs all over.—Ditches.

11. *P. hydropiperoides* Michx. (MILD WATER-PEPPER.) Stem slender, 1–3 feet high, in shallow water. Leaves narrow. Sheaths hairy and fringed with long bristles. Spikes slender, erect, pale rose-coloured or whitish. Stamens 8. Stigmas 3. Achene 3-angled, shining.—In shallow water.

12. *P. acre* HBK. (WATER SMARTWEED.) Sheaths fringed with bristles. Leaves transparent-dotted. Stem rooting at the decumbent base, 2–4 feet high, in shallow water or muddy soil. Leaves narrow, taper-pointed. Spikes slender, erect, pale rose-coloured. Stamens 8. Achene 3-angled, shining.—Muddy soil or shallow water.

13. *P. Hydropiper* L. (COMMON SMARTWEED or WATER-PEPPER.) Sheaths and leaves as in the last, the leaves, however, larger, acrid. Spikes slender, nodding, greenish. Sepals glandular-dotted. Stamens 6. Achene dull.—Wet places.

14. *P. virginianum* L. Calyx greenish, unequally 4-parted. Stamens 5. Styles 2, persistent on the flat achene. Flowers in long and slender naked spikes. Stem upright, nearly smooth. Leaves ovate or ovate-lanceolate, taper-pointed, rough-ciliate. Sheaths hairy and fringed.—Thickets, in rich soil.

(2) *Leaves heart-shaped or sagittate. Sheaths much longer on one side than on the other.*

15. *P. arifolium* L. (HALBERD-LEAVED TEAR-THUMB.) Has grooved stem, halberd-shaped, long-petioled leaves, flowers in short loose racemes, 6 stamens, and a flattish achene.

16. *P. sagittatum* L. (ARROW-LEAVED TEAR-THUMB.) Stem 4-angled, the angles beset with reflexed minute prickles, by which the plant is enabled to climb. Leaves arrow-shaped. Stamens usually 8. Achenes 3-angled.—Common in low grounds.

17. *P. Convolvulus* L. (BLACK BINDWEED.) Stem twining, not prickly but roughish; the joints naked. Flowers in loose paniced racemes, 3 of the calyx lobes rigid in fruit. Leaves heart-shaped and partly halberd-shaped. Not climbing so high as the next.—Cultivated grounds and waste places.

18. *P. scandens* L. (CLIMBING FALSE BUCKWHEAT.) Stem twining high, smooth; sheaths naked, 3 of the calyx lobes winged in fruit.—Moist thickets.

19. *P. cilino'de* Michx. Stem twining, minutely downy. Sheaths fringed at the base with reflexed bristles.—Sandy pine woods and rocky hills.

20. *P. vivip'arum* L. A small smooth perennial, with bulb-like stem, lanceolate leaves, and pinkish flowers in an erect, spike-like cluster. Stamens 8, styles 3.

3. RU'MEX L. DOCK. SORREL

A. Herbage not sour; the leaves not halberd-shaped.

1. *R. Britan'nica* L. (GREAT WATER DOCK.) Growing in marshes. Stem erect, stout, 5-6 feet high. Leaves lanceolate; not wavy-margined or heart-shaped, often over a foot long. Flowers nodding on thread-like pedicels. Valves nearly orbicular, finely net-veined, each with a grain on the back.—Wet places.

2. *R. pal'lidus* Bigel. (WHITE DOCK.) May be looked for in marshes on the sea-coast and far northward. Leaves narrowly lanceolate or oblong. The whorls of flowers are dense and form a very conspicuous spike, owing to the great size of the whitish grains on the back of the valves.

3. *R. verticilla'tus* L. (SWAMP DOCK.) Leaves lanceolate or oblong-lanceolate, not wavy, the lowest often heart-shaped. Stem tall. Fruit-bearing pedicels slender, club-shaped, abruptly reflexed, several times longer than the fruiting calyx. Valves dilated-rhomboid, strongly wrinkled, each bearing a very large grain.—Swamps, common.

4. *R. cris'pus* L. (YELLOW DOCK.) Leaves with strongly wavy or curly margins, lanceolate. Whorls of flowers in long, wand-like racemes. Valves grain-bearing.—Cultivated soil and waste places.

5. *R. obtusifo'lius* L. (BITTER DOCK.) Lowest leaves oblong-heart-shaped, obtuse, only slightly wavy-margined; the upper oblong-lanceolate, acute. Whorls loose, distinct. Valves somewhat halberd-shaped, deeply toothed at the base, usually one only grain-bearing.—Waste grounds.

6. *R. sanguin'eus* L. Leaves oblong-lanceolate, often fiddle-shaped, wavy-margined. Whorls distant, in long, slender, leafless spikes. Valves narrowly oblong, obtuse, entire. Veins of the leaf red or green.—Atl. Prov. chiefly.

7. *R. persicarioides* L. (GOLDEN DOCK.) 1–2 feet high, slightly pubescent, much-branched. Leaves linear-lanceolate, wavy-margined, the lower auricled or heart-shaped at the base. Flowers in whorls forming leafy spikes. Valves oblong, lance-pointed, each bearing 2–3 long bristles on each side, and a large grain on the back.—Seashore, Atl. Prov. and Ont.

8. *R. venosus* Pursh. Stems erect, 1–2 feet high. Leaves ovate-lanceolate; stipules dilated. Flower panicle dense and short, almost sessile. Valves rose-coloured, cordate, without grains.—Pr. Prov.

9. *R. occidentalis* Wats. Stems smooth, erect, often tinged with purple. Leaves large, somewhat flat. Valves ovate-orbicular, without grains or with only one.—Pr. Prov.

10. *R. mexicanus* Meisn. Stems erect. Leaves lanceolate to narrowly oblong. Panicle dense, its branches usually upright. Valves with brown grains.—Pr. Prov.

B. Herbage sour; leaves halberd-shaped.

11. *R. Acetosella* L. (FIELD OR SHEEP SORREL.) Stem 6–12 inches high. Leaves narrow-lanceolate, the lower halberd-shaped. Flowers dioecious, in a terminal naked panicle. Achene naked.—A very common weed in poor soil.

3. *FAGOPYRUM* L. BUCKWHEAT

1. *F. esculentum* Moench. (BUCKWHEAT.) Old fields and copses, remaining after cultivation. Achene smooth and shining.

2. *F. tartaricum* Gaertn. Very small flowers, and a dull roughish achene.—Escaped from cultivation in a few places.

CHENOPODIA'CEAE (GOOSEFOOT FAMILY)

Homely herbs, with more or less succulent leaves (chiefly alternate), and small greenish flowers mostly in interrupted spikes. Corolla wanting. Stamens usually as many as the lobes of the calyx and opposite them. Ovary 1-celled and 1-ovuled, forming an achene or utricle in fruit. Stigmas mostly 2.

Synopsis of the Genera

1. *Chenopodium*. Weeds with (usually) mealy leaves, and very small, perfect, greenish, sessile flowers in small, panicked, spiked

- clusters. Calyx 5-cleft, more or less enveloping the fruit, and sometimes becoming fleshy and berry-like. Stamens mostly 5, filaments slender.
2. *Monol'epis*. A low annual, glabrous or somewhat mealy, with small fleshy leaves. Sepal only 1, bract-like. Stamen 1. Styles 2. Seed vertical, flattened.
 3. *Cyclolo'ma*. A much-branched, coarse herb, with very small, scattered, sessile flowers in open panicles. Flowers perfect or pistillate. Calyx 5-cleft, the concave lobes strongly keeled, at length with a broad and continuous, horizontal wing. Stamens 5; styles 3. Seed horizontal, flat.
 4. *At'riplex*. Flowers monoecious or dioecious, the staminate with a regular calyx, in spiked clusters; the pistillate without a calyx, but with a pair of appressed bracts.
 5. *Corisper'mum*. Flowers all perfect, single, and sessile in the axils of the upper leaves, usually forming a spike. Calyx of a single delicate sepal. Low herbs, with linear, 1-nerved leaves.
 6. *Salicor'nia*. Low saline plants with fleshy, leafless, jointed stems and opposite branches. Flowers perfect, densely spiked, 3 together sunk in each hollow of the fleshy axis of the spike. Calyx small and bladder-like, with a toothed or torn margin, at length spongy and narrowly wing-bordered. Stamens 1 or 2. Styles 2, united at the base.
 7. *Suae'da*. Fleshy saline plants with alternate, terete, linear leaves. Flowers perfect, sessile in the axils of leafy bracts. Calyx 5-parted, very fleshy. Stamens 5, the anthers exerted. Stigmas 2 or 3. Seed horizontal.
 8. *Sal'sola*. A diffusely branching annual, with alternate, awl-shaped, prickly-pointed leaves. Flowers perfect, sessile, with 2 bractlets. Calyx 5-parted, persistent, its divisions at length horizontally winged on the back. Seed horizontal.

1. CHENOPO'DIUM L. GOOSEFOOT. PIGWEED

1. *C. album* L. (LAMB'S QUARTERS.) Stem upright, 1-3 feet high. Leaves varying from rhombic-ovate to lanceolate, more or less toothed, mealy, as are also the dense flower-clusters.—Extremely common in cultivated soil.

2. *C. ur'bicum* L. Rather pale and only slightly mealy, 1-3 feet high, branches erect. Leaves triangular, acute, coarsely and sharply many-toothed. Spikes erect, crowded in a long and narrow racemose panicle.—Waste places in towns.

3. *C. hy'bridum* L. (MAPLE-LEAVED GOOSEFOOT.) Bright green. Stem widely branching, 2-4 feet high. Leaves thin,

large, triangular-heart-shaped, sinuate-angled, the angles extended into pointed teeth. Panicles loose, leafless. Plant with a rank unpleasant odour.—Waste places.

4. *C. Bo'trys* L. (JERUSALEM OAK.) Not mealy, but sticky; low, spreading, sweet-scented. Leaves deeply sinuate, slender-petioled. Racemes in loose divergent corymbs.—Roadsides; escaped from gardens.

5. *C. ambrosioides* L. (MEXICAN TEA.) Not mealy, but sticky. Leaves slightly petioled, wavy-toothed or nearly entire. Spikes densely flowered.—Streets of towns.

6. *C. glaucum* L. (OAK-LEAVED GOOSEFOOT.) Somewhat glaucous-mealy, 5-12 inches high, spreading. Leaves sinuately pinnatifid-toothed, oblong, obtuse. Clusters small, in axillary spikes. Seed vertical, exserted, with sharp edges.

7. *C. Bo'nus-Henricus* L. (GOOD-KING-HENRY.) Stout, erect, 1-2 feet high. Leaves broadly triangular-hastate, slightly sinuate or entire. Flowers somewhat densely paniculate-spiked. Seed vertical, exserted, with blunt edges. Not common.

8. *C. capita'tum* Aschers. (STRAWBERRY BLITE.) Stem ascending, branching. Leaves smooth, triangular. Flower-clusters interruptedly spiked and in fruit resemble strawberries, owing to the bright-red, fleshy calyces.

9. *C. hu'mile* Hook. A dwarf form with lanceolate to spatulate leaves. Flowers crowded in axillary clusters.—Pr. Prov.

10. *C. ru'brum* L. (COAST BLITE.) Stem angled, much-branched. Leaves thickish, acuminate, the upper linear-lanceolate. Flowers-clusters scattered in axillary leafy spikes. Stamens 1-2.—Pr. Prov., in saline soil.

11. *C. Boscia'num* Moq. Stems slender, erect and branched. Leaves oblong to lanceolate, tapering to a slender petiole. Flowers solitary or in small clusters, pericarp easily detached from the seed.

12. *C. leptophyl'um* Nutt. Stems simple or branched. Leaves linear, entire, with short petioles. Flower-clusters in dense or interrupted spikelets; calyx lobes keeled; pericarp easily detached.

2. MONOL'EPIS Schrad.

1. *M. Nuttallia'na* Wats. Branched from the base. Leaves slender and hastate, passing into leaf-like bracts; little if at all mealy.—Pr. Prov.

3. CYCLOLO'MA Moquin. WINGED PIGWEED

1. *C. atriplicifolium* Coult. Diffuse, 6–15 inches high, light green or sometimes purple.

4. AT'RIPLEX L. ORACHE

1. *A. patula* L. Erect or prostrate, scurfy, green or rather hoary. Leaves varying from triangular or halberd-shaped to lance-linear, petioled.

Var. *hastata* Gray. The lower leaves, at least, broadly triangular-hastate, often toothed.—Atl. Prov. and Pr. Prov.

Var. *littoralis* Gray. Slender, with leaves linear-lanceolate to linear.—Waste places.

2. *A. Nuttallii*, Watson. A shrubby, densely appressed-scurfy perennial with oblong-spatulate to narrowly oblanceolate, entire leaves.—Pr. Prov.

3. *A. argentea* Nutt. A low, branching, gray-scurfy plant with deltoid, hastate leaves. Sterile flowers in spikes, the fertile in axillary clusters; bracts of the fertile flowers united, their margins deeply toothed.—Pr. Prov.

5. CORISPER'MUM L. BUG-SEED

1. *C. hyssopifolium* L. Somewhat hairy when young, pale. Stamens 1 or 2. Styles 2. Fruit oval, flat.—Sandy beaches, Western and South-western Ontario, and Pr. Prov.

6. SALICOR'NIA L. GLASSWORT. SAMPHIRE

1. *S. europoea* L. (SAMPHIRE.) Flowers perfect, in threes, embedded in hollows on the thickened upper joints, forming an elongated narrow spike. Calyx small and bladder-like, its margin toothed. Stamens 1 or 2.—Salt marshes, Atl. Prov. and Pr. Prov.

2. *S. mucronata* Bigel. Thick spikes and mucronate-pointed scales. Stem turning red when old.—Atl. sea-coast.

7. SUAE'DA Forskal. SEA BLITE

1. *S. linearis* Moq. (SEA BLITE.) A branching fleshy herb, with alternate, roundish, linear leaves. Flowers perfect, sessile in the axils of leafy bracts on slender branchlets. Sepals very thick. Stamens 5, with anthers exerted.—Atl. Prov.

8. SAL'SOLA L. SALTWORT

1. *S. Ka'li* L., var. *tenuifo'lia* G.F.W. Mey. (RUSSIAN THISTLE.) Flowers perfect, sessile, with two bractlets, single in axils of leaves. Calyx 5-parted, inclosing the depressed fruit. Stamens 5. A branching plant with alternate, filiform, prickly-pointed leaves.

AMARANTHA'CEAE (AMARANTH FAMILY)

Homely weeds, a good deal like the plants of the last Family, but the flower-clusters are interspersed with dry and chaff-like (sometimes coloured), persistent bracts, usually 3 to each flower. Corolla wanting.

Synopsis of the Genera

1. *Amaran'thus*. Flowers monoecious or polygamous, all with a calyx of 3 or 5 distinct, erect sepals.
2. *Acni'da*. Flowers dioecious; calyx none in the pistillate flowers.

1. AMARAN'THUS L. AMARANTH

1. *A. panicula'tus* L. Reddish flowers in terminal and axillary, slender spikes, the bracts awn-pointed.—In the neighbourhood of gardens.

2. *A. retroflex'us* L. (PIGWEEED.) Flowers greenish, in spikes, forming a stiff panicle. Leaves a dull green, long-petioled, ovate, wavy-margined. Stem erect.—Common in cultivated soil.

3. *A. hyb'ridus* L. Similar to No. 2, but smoother, with slender spikes, and more or less zigzag; bracts with fairly long awns.

4. *A. graeci'zans* L. Flowers greenish, in small, close, axillary clusters. Stem low and spreading.—Roadsides.

5. *A. blitoid'es* Wats. Like No. 4 but prostrate. Spikelets contracted, bracts ovate-oblong, acute.

2. ACNI'DA L.

1. *A. cannabi'na* L. A tall smooth herb, with lanceolate or oblong-ovate, alternate leaves on long petioles, and small clusters of greenish flowers in interrupted spikes. Fruit indehiscent.—Wet places.

2. *A. tubercula'ta* Moq. Has the flowers in the lower part of the stem in close clusters in the axils of the leaves. Fruit dehiscing irregularly.

PHYTOLACCA'CEAE (POKEWEED FAMILY)

Herbs with alternate leaves and perfect flowers. The ovary is composed of several carpels in a ring, forming a berry in fruit.

1. PHYTOLAC'CA L. POKEWEED

1. *P. decan'dra* L. (COMMON POKE.) Calyx of 5 rounded white sepals. Corolla wanting. Ovary green, of ten 1-seeded carpels united in a ring. Styles 10, short and separate. Stamens 10. Fruit a crimson or purple, 10-seeded berry. Stem very tall and stout, smooth. Flowers in long racemes opposite the leaves. —Sandy soil.

ILLECEBRA'CEAE (KNOTWORT FAMILY)

Small, diffuse or tufted herbs with mostly opposite and entire leaves, scarious stipules, and a 4-5-toothed or parted, coriaceous, persistent calyx. Petals wanting. Stamens perigynous, as many as the lobes of the calyx and opposite them, or fewer. Style 2-cleft at the apex. Fruit a 1-seeded utricle.

1. PARONYCH'IA Adans. WHITLOW-WORT

1. *P. sessiliflo'ra* Nutt. Flowers terminal, solitary, sessile. Stems densely matted or tufted, from a woody root. The dry silvery stipules 2-cleft. Sepals oblong-linear, concave, awned at the apex.—Pr. Prov.

AIZOA'CEAE (ICE-PLANT FAMILY)

A miscellaneous group, embracing plants formerly included in Caryophyllaceae and Portulacaceae; differing, however, from true representatives of these in having partitions in the ovary. Petals wanting in our genus.

1. MOLLU'GO L. CARPET-WEED

1. *M. verticilla'ta* L. A prostrate, much-branched herb, growing in patches. Leaves spatulate, apparently verticillate. Flowers on long axillary pedicels, clustered into a sort of umbel. Sepals 5, white inside. Petals none. Stamens mostly 3. Styles 3. Pod 3-celled, 3-valved.

CARYOPHYLLACEAE (PINK FAMILY)

Herbs with opposite (occasionally whorled) and entire leaves, the stems usually swollen at the joints. Flowers regular, with the parts mostly in fives, occasionally in fours. Petals sometimes absent. Stamens not more than twice as many as the petals. Styles 2-5, stigmatic along the inner side. Pod usually 1-celled, with the seeds attached to the base, or to a column which rises from the centre of the cell.

Synopsis of the Genera

A. *Sepals united into a tube or cup. Petals and stamens borne on the stalk of the ovary; petals with long narrow claws.*

1. *Sapona'ria*. Calyx cylindrical or 5-angled. Styles 2.
2. *Sile'ne*. Calyx 5-toothed. Styles 3.
3. *Lych'nis*. Calyx 5-toothed. Styles 5, alternate with the petals.
4. *Agrostem'ma*. Calyx 5-toothed. Styles 5, opposite the petals.

B. *Sepals separate to the base or nearly so. Petals without claws, they and the stamens inserted at the base of the sessile ovary.*
Low herbs.

(1) *Stipules none.*

5. *Arena'ria*. Petals not cleft at the apex. Styles usually 3. Pod splitting into 3 or 6 valves.
6. *Stella'ria*. Petals 2-cleft at the apex, or none. Pod splitting to the base into twice as many valves as there are styles. Styles generally 3.
7. *Ceras'tium*. Petals 2-cleft, or notched. Styles mostly 5. Pod opening at the apex by 10 teeth.
8. *Sagi'na*. Petals 4 or 5, or none, entire. Stamens as many or twice as many. Styles 4 or 5. Pod 4-5-valved.

(2) *Stipules present.*

9. *Spergula'ria*. Styles usually 3. Pod 3-valved, short. Leaves filiform or linear, opposite.
10. *Sper'gula*. Styles 5. Valves of the pod opposite the sepals. Leaves thread-like, whorled.

1. SAPONA'RIA L. SOAPWORT

1. *S. officina'lis* L. (BOUNCING BET.) A stout perennial with rose-coloured or pinkish flowers clustered in corymbs. Petals

with an appendage at the top of the claw. Leaves 3-5-ribbed, the lower ovate, upper lanceolate. Pod raised on a short stalk. Styles 2.—Old gardens and roadsides.

2. *S. Vacca'ria* L. (COMMON COW-HERB.) Annual, glabrous. Flowers pale red, in corymbed cymes. Petals without an appendage. Calyx 5-angled, wing-angled in fruit.—Waste places.

2. SILE'NE L. CATCHFLY. CAMPION

1. *S. latifo'lia* Britton and Rendle. (BLADDER CAMPION.) Pale or glaucous, very smooth. Stem erect, a foot high. Leaves ovate-lanceolate. Calyx much-inflated, purple-veined. Stamens and styles exserted.—Not common westward.

2. *S. antirrhi'na* L. (SLEEPY CAMPION.) Stem slender, simple or slightly branching above, a portion of the upper internodes sticky. Leaves linear or lanceolate. Flowers small, pink or purplish, opening only for a short time in sunshine. Calyx ovoid, shining, not inflated.—Dry soil.

3. *S. noctiflo'ra* L. (NIGHT-FLOWERING CATCHFLY.) Stems very sticky, pubescent. Lower leaves spatulate, upper lanceolate. Flowers few, peduncled. Calyx tube with awl-shaped teeth. Petals white or whitish, 2-parted. Opening only at night or in cloudy weather.—A very common weed in cultivated grounds.

4. *S. Arme'ria* L. (SWEET-WILLIAM CATCHFLY.) Glaucous. Leaves ovate-lanceolate. Flowers pink or white, in flat cymes. Calyx club-shaped.—Escaped from gardens in some places.

5. *S. virgin'ica* L. (FIRE PINK.) Occurs in South-western Ontario, and may be recognized by its crimson petals, and bell-shaped calyx, nodding in fruit.

6. *S. Menzie'sii* Hook. Stem weak and forked. Small white flowers in cymose clusters. The calyx pubescent with viscid hairs. Petals 2-cleft.—Pr. Prov.

7. *S. acaul'is* L. (MOSS CAMPION.) A very small, tufted, moss-like perennial, 1-2 inches high, with linear crowded leaves. Petals purple or rarely white.—Atl. Prov.

3. LYCH'NIS L. CAMPION

1. *L. al'ba* Mill. (WHITE CAMPION.) Viscid - pubescent. Flowers commonly dioecious, white or pinkish, fragrant, opening at evening.—Waste grounds in a few localities.

2. *L. dioi'ca* L. Resembles the last, but has red (rarely white) flowers without odour, opening in the morning.—Rare.

4. AGROSTEM'MA L.

1. *A. Githa'go* L. (CORN COCKLE.) Plant clothed with long, soft, appressed hairs. Leaves linear. Calyx lobes extremely long, very much like the upper leaves, surpassing the purple petals.—Wheat-fields.

5. ARENA'RIA L. SANDWORT

1. *A. serpyllifo'lia* L. (THYME-LEAVED SANDWORT.) Much-branched, 2–6 inches high, roughish-pubescent. Leaves small, ovate, acute. Petals white, hardly as long as the sepals. Sepals pointed, 3–5-nerved. Pod pointed, 6-toothed.—Sandy fields.

2. *A. stricta* Michx. Stem erect, or diffusely spreading from a small root. Leaves awl-shaped or bristle-form, the upper ones reduced to 1-nerved bracts, crowded in the axils. Cyme diffuse, many-flowered. Sepals pointed, 3-ribbed, half as long as the white petals.—Rocky fields.

3. *A. lateriflo'ra* L. Stem erect, slender, minutely pubescent. Leaves oval or oblong, $\frac{1}{2}$ –1 inch long. Peduncles usually three-flowered. Sepals obtuse. Petals white, large, twice as long as the sepals. Flower $\frac{1}{3}$ inch across when fully expanded.—Gravelly shores.

4. *A. groenlan'dica* Spreng. Stems densely tufted, filiform, with soft, filiform-linear leaves. Flowering stems erect, smooth, 2–4 inches high, few-flowered. Sepals oblong, obtuse, nerveless.—Atl. sea-coast.

5. *A. peploi'des* L. Very fleshy stems and leaves, the latter somewhat clasping. —Occurs eastward towards the sea-coast.

6. STELLA'RIA L. CHICKWEED. STARWORT

1. *S. me'dia* Cyrill. (COMMON CHICKWEED.) Stems branching, decumbent, soft and brittle, marked lengthwise with one or two pubescent lines. Lower leaves on hairy petioles, ovate. Flowers small, white. Petals cleft, shorter than the sepals.—Extremely common in damp grounds and old gardens.

2. *S. longifo'lia* Muhl. (LONG-LEAVED STITCHWORT.) Stems branching, very weak and brittle, supporting themselves on other

plants. Leaves linear. Pedicels of the flowers long, slender, and spreading, deflexed. Petals white, longer than the 3-nerved sepals.—Low grassy banks of streams.

3. *S. ion'gipes* Goldie. (LONG-STALKED STITCHWORT.) Leaves somewhat rigid, ascending, linear-lanceolate, acute, broadest at the base. Cyme terminal, few-flowered, the long pedicels erect, scaly-bracted. Petals longer than the sepals. Seeds smooth.

4. *S. gramin'ea* L. Like the last, but the leaves broadest above the base, the pedicels widely spreading, and the seeds strongly but finely rugose. (Intro. from Europe.)

5. *S. uligino'sa* Murr. (SWAMP STITCHWORT.) Stems weak, decumbent or diffuse. Leaves lanceolate or oblong, veiny. Petals and ripe pods as long as the sepals. Seeds roughened. Cymes naked, becoming lateral.—Swamps and rills, eastward.

6. *S. crassifo'lia* Ehrh. Stems diffuse or erect, weak. Leaves rather fleshy, lanceolate to oblong, those of flowering branches smaller and thinner. Petals longer than sepals or wanting. Seeds rugose-roughened. Flowers terminal or in the forks of stem or branches.—Wet places.

7. *S. borea'lis* Bigelow. (NORTHERN STITCHWORT.) Stem erect or spreading, weak, forking. Leaves broadly-lanceolate to ovate-oblong. Petals 2-5, shorter than the sepals or wanting. Cyme leafy. Seeds smooth.—Wet places.

8. *S. humifu'sa* Rottb. Low, spreading or creeping. Leaves fleshy, ovate or oblong. Pedicels axillary or terminal, on leafy stems or branches. Petals a little longer than the sepals. Seeds smooth.—Atl. Prov.

7. CERAS'TIUM L. MOUSE-EAR CHICKWEED

1. *C. visco'sum* L. (LARGER MOUSE-EAR CHICKWEED.) Stem ascending, hairy and somewhat clammy. Leaves ovate or obovate, obtuse. Flowers at first in close clusters. Pedicels not longer than the sepals. Petals shorter than the calyx.—Not common, sometimes confounded with No. 2.

2. *C. vulga'tum* L. (COMMON MOUSE-EAR CHICKWEED.) Stems hairy, viscid, spreading. Leaves lanceolate-oblong, rather acute. Flowers in loose cymes. Pedicels longer than the sepals. Petals usually equalling the calyx.—Common; fields and copses.

3. *C. arven'se* L. (FIELD CHICKWEED.) Stem decumbent at the base, pubescent, slender, 4-8 inches high. Leaves linear, or linear-lanceolate, often fascicled in the axils, longer than the lower internodes. Petals obcordate, more than twice as long as the calyx. Pod $\frac{1}{2}$ to $\frac{2}{3}$ longer than the calyx. Cyme few-flowered.

Var. *oblongifo'lium* Holl. and Britt. Taller, pubescent. Leaves oblong or oblong-lanceolate. Pod twice as long as the calyx.

4. *C. nu'tans* Raf. Stems very clammy-pubescent and branching diffusely. The loose and open cymes many-flowered. Leaves lance-oblong. Pods nodding on the stalks, curved upwards, thrice the length of the calyx.—In places where water lies in spring.

8. SAGI'NA L. PEARLWORT

1. *S. procum'bens* L. (PEARLWORT.) A low matted herb with narrowly linear leaves. Flowers small, terminal, with their parts in fours, rarely in fives. Petals shorter than the ovate obtuse sepals, or none. Pod many-seeded, 4-5-valved. Top of peduncle often bent into a hook.—Damp places, Atl. Prov.

2. *S. nodo'sa* Fenzl. A low tufted herb. The lower leaves thread-form; the upper short, awl-shaped, with clusters of minute ones in their axils. Parts of flower in fives, the stamens sometimes ten. Petals much longer than sepals. Flowers terminal. Pods as in *S. procumbens*.—Wet sandy shores, Atl. Prov.

9. SPERGULA'RIA J. and C. Presl. SAND SPURREY

1. *S. ru'bra* J. and C. Presl. Leaves linear, flat, hardly fleshy. Stipules lanceolate. Stems usually glandular-pubescent near the summit. Calyx rather longer than the pink-red corolla, and small pod. Seeds rough with projecting points.—Dry sandy ground, Atl. Prov.

2. *S. mari'na* Griseb. More fleshy than *S. rubra*, usually pubescent. Stipules ovate. Leaves terete. Sepals a little shorter than the pod. Petals pale. Seeds usually roughened with points.—Sea-coast, Atl. Prov.

3. *S. me'dia* C. Presl. Much-branched, glabrous. Leaves linear, fleshy, somewhat terete. Petals white. Pod about twice as long as the sepals; seeds usually smooth.—Sea-coast and Pr. Prov.

10. SPER'GULA L. SPURREY

1. *S. arven'sis* L. (CORN SPURREY.) An annual herb resembling a Buda, with numerous thread-like leaves in whorls. Flowers white, in paniced cymes. Pod 5-valved; seeds minutely roughened.—Grain fields, Atl. Prov.

PORTULACA'CEAE (PURSLANE FAMILY)

Herbs with fleshy, entire, exstipulate leaves, and regular, hypogynous or perigynous flowers. Sepals 2. Petals 5. Stamens mostly 5–20. Styles 3–8, united below. Pod 1-celled, few or many-seeded.

Synopsis of the Genera

1. *Portulac'a*. Stamens 7–12. Pod opening by a lid, many-seeded.
2. *Clayto'nia*. Stamens 5. Pod 3-valved, 3–6-seeded.
3. *Mon'tia*. Stamens usually 3. Petals 5, three of them somewhat smaller. Pod 3-valved and 3-seeded.

1. PORTULAC'A L. PURSLANE

1. *P. olera'cea* L. (COMMON PURSLANE.) A low fleshy herb, very smooth, with obovate or wedge-shaped leaves. Calyx 2-cleft, partly adnate to the ovary, the sepals keeled. Petals 5, yellow, fugacious.—A common pest in gardens.

2. CLAYTO'NIA L. SPRING-BEAUTY

1. *C. virgin'ica* L. Leaves linear-lanceolate, 3–6 inches long.
2. *C. carolinia'na* Michx. Leaves ovate-lanceolate or oblong, tapering at the base. In both species the corolla is rose-coloured, with dark veins. The stem springs from a small tuber and bears two opposite leaves and a loose raceme of flowers.—Rich woods in early spring.

3. MON'TIA L.

1. *M. fonta'na* L. A small, spreading, somewhat fleshy herb, with opposite spatulate leaves. Sepals 2. Petals 5, three of them somewhat smaller. Stamens usually 3, borne on the claws of the smaller petals. Pod 3-valved and 3-seeded.—Wet places, Atl. Prov.

CERATOPHYLLA'CEAE (HORNWORT FAMILY)

Aquatic herbs growing under water in ponds and slow streams. Leaves whorled, finely dissected.

1. CERATOPHYLLUM L. HORNWORT

1. *C. demersum* L. Flowers minute, axillary, sessile, monocious, without calyx or corolla, but with an 8–12-cleft involucre. Staminate flowers of 12–24 stamens with large sessile anthers. Pistillate flowers of a single 1-celled ovary, forming an achene, beaked with the slender style. Embryo with 4 cotyledons.

NYMPHAEACEAE (WATER-LILY FAMILY)

Aquatic herbs with cordate or peltate, usually floating, leaves. Floating flowers on long emerged peduncles. Petals and stamens generally numerous.

Synopsis of the Genera

1. *Brasenia*. Sepals and petals each 3 (occasionally 4). Stamens 12–18. Leaves oval, peltate.
2. *Castalia*. Sepals 4. Petals numerous, white, imbricated in many rows, gradually passing into stamens, hypogynous, or epigynous. Stamens epigynous. Stigmas radiating as in a poppy-head.
3. *Nymphaea*. Sepals 5–6, yellow. Petals many, small and stamen-like. Stamens under the ovary.

1. BRASENIA Schreber. WATER-SHIELD

1. *B. Schreberi* Gmel. Stems and under-surface of the leaves coated with jelly. Leaves oval, 2 inches across, peltate. Flowers small, purplish.—Ponds and slow-flowing streams.

2. CASTALIA Salisb. WATER-LILY

1. *C. odorata* Woodville and Wood. (SWEET-SCENTED WATER-LILY.) Leaves orbicular, cleft at the base to the petiole, 5–9 inches wide, often crimson underneath. Flower very sweet-scented, white, sometimes pink-tinted.—Ponds and slow streams.

2. *C. tuberosa* Greene. (TUBER-BEARING WATER-LILY.) Leaves larger and more prominently ribbed than in No. 1, reniform-orbicular, green on both sides, rarely purplish beneath. Flower not at all, or only slightly, sweet-scented. Rootstocks producing tubers,

which come off spontaneously.—Mostly in slow waters opening into Lake Ontario.

3. NYMPHAE'A L. YELLOW POND-LILY

1. *N. ad'vena* Ait. (COMMON YELLOW POND-LILY.) Leaves floating, or emersed and erect, thickish, roundish or oblong, cordate. Sepals 6.—Stagnant water.

Var. *minus*. More slender, with smaller leaves and flowers. Probably a hybrid between this and the next.

2. *N. microphyl'la* Pers. (SMALL YELLOW POND-LILY.) Floating leaves usually not more than 2 inches across, the sinus very narrow or closed; the radical ones very thin and with a wide obtuse sinus. Flowers hardly an inch across. Sepals 5.—Northward, in slow waters.

RANUNCULA'CEAE (CROWFOOT FAMILY)

Herbs or woody climbers with an acrid colourless juice. Parts of the flower separate from each other. Corolla sometimes wanting. Stamens numerous. Pistil (with one or two exceptions) apocarpous. Fruit an achene, follicle, or berry. Leaves exstipulate, with the blades usually dissected, and petioles spreading at the base.

Synopsis of the Genera

1. *Clem'atis*. Real petals none or stamen-like. Coloured sepals 4 or more, valvate in the bud. Fruit an achene, with the long and feathery style attached. Leaves all opposite. Plant climbing by the bending of the petioles.
2. *Anemo'ne*. Petals none or stamen-like. Coloured sepals 5-numerous, imbricated in the bud. Achenes many, in a head, pointed or tailed, not ribbed. Stem-leaves opposite or whorled, forming an involucre remote from the flower.
3. *Hepat'ica*. Petals none. Coloured sepals 6–12, whitish or bluish. Achenes many, not ribbed. Leaves all radical. An involucre of 3 leaves close to the flower, and liable to be mistaken for a calyx.
4. *Thalic'trum*. Petals none. Coloured sepals 4 or more, greenish. Achenes several, angled or grooved. No involucre. Stem-leaves alternate, decom-pound. Flowers in panicles or corymbs, mostly dioecious.
5. *Anemonel'la*. Petals none. Sepals 5–10, large, white. Achenes 4–12, strongly ribbed. Stem-leaves in a whorl just below the flowers. Radical leaves 1 (or 2), compound. Flowers one or more (in an umbel).

6. **Myosu'rus.** Petals 5, white, with slender claws. Sepals 5, spurred at the base. Stamens 5-20. Achenes in a long slender spike. Small plants with tufted narrow root-leaves, and 1-flowered scapes. Flowers small.
7. **Ranun'culus.** Sepals 5, deciduous. Petals generally 5, each with a pit or little scale on the inside of the claw. Achenes many, in heads, short-pointed. Stem-leaves alternate. Flowers solitary or corymbed, mostly yellow, rarely white.
8. **Cal'tha.** Petals none. Sepals 5-9, yellow. Fruit a many-seeded follicle. Leaves large, glabrous, heart-shaped or kidney-shaped, mostly crenate. Stem hollow and furrowed.
9. **Cop'tis.** Sepals 5-7, white, deciduous. Petals 5-7, yellow, with slender claws, and somewhat tubular at the apex. Carpels 3-7, on slender stalks. Fruit a follicle. Flowers on naked scapes. Leaves radical, shining, divided into three wedge-shaped leaflets, sharply toothed. Rootstocks golden yellow, bitter.
10. **Aquile'gia.** Sepals 5, coloured. Petals 5, each a long hollow spur. Carpels 5. Follicles erect, many-seeded. Flowers very showy, terminating the branches. Leaves decom-pound.
11. **Delphin'ium.** Sepals 5, unequal, petal-like, the upper one produced backward into a spur. Petals 4, irregular, the upper pair spurred, and projecting backward into the spur of the upper sepal (rarely only 2). Flowers blue or whitish, in terminal racemes. Leaves palmately dissected.
12. **Actae'a.** Sepals 4-5, caducous. Petals 4-10, with slender claws. Stamens many, with long filaments. Fruit a many-seeded berry. Flowers white, in a short, thick raceme. Leaves decom-pound, leaflets sharply toothed.
13. **Cimicif'uga.** Sepals 4-5, caducous. Petals several, small, two-horned at the apex. Stamens numerous. Carpels 1-8, becoming pods. Flowers in long, plume-like racemes.
14. **Hydras'tis.** Petals none. Flowers solitary. Sepals 3, petal-like, greenish-white, caducous. Stamens numerous. Carpels 12 or more, forming a head of crimson, 1-2-seeded berries in fruit. Stem low, from a knotted yellow rootstock. Leaves simple, lobed.

1. CLEM'ATIS L. VIRGIN'S BOWER

1. **C. virginia'na** L. (COMMON VIRGIN'S BOWER.) A woody-stemmed climber. Flowers in paniced clusters, often dioecious, white. Leaves of 3 ovate leaflets, which are cut or lobed. Feathery tails of the achenes very conspicuous in the autumn.—Along streams and in swamps.

2. *C. ligusticifolia* Nutt. Very much like the last, but the leaves have usually 5 leaflets.—Pr. Prov.

3. *C. verticillaris* DC. Shrubby climber. Peduncles bearing single large flowers, with thin, wide-spreading, bluish-purple sepals. Tails of the achenes plumose. Leaves of three leaflets, which are entire, or sparingly toothed or lobed.—Rocky places.

2. ANEMONE L. ANEMONE

A. Styles long and hairy, forming feathery tails in fruit.

1. *A. patens* L., var. *Wolfgangiana* Koch. (PRAIRIE ANEMONE or PASQUE FLOWER.) Villous with long silky hairs. Flower solitary, erect, appearing before the leaves. Sepals 5–7, purplish or white, an inch or more in length, spreading at maturity. Involucre sessile, its lobes numerous, all united at the base to form a shallow cup. Leaves radical, the blades deeply cut into 3 well-marked divisions, the middle one stalked, and all again dissected into narrow lobes. Feathery tails of the carpels 2 inches long. A low plant, flowering in early spring.—Pr. Proy.

B. Styles short, not plumose.

2. *A. parviflora* Michx. (SMALL-FLOWERED ANEMONE.) Stem 6–12 inches high, 1-flowered. Sepals 5 or 6, white. Involucre 2–3-leaved far below the flower. Head of carpels woolly, globular. Root-leaves small, 3-parted, their divisions crenately lobed.—Rocky river-margins.

3. *A. multifida* Poir. (MANY-CLEFT ANEMONE.) Silky-hairy, 6–15 inches high. Principal involucre 2–3-leaved, far below the flower and bearing one naked and one or two 2-leaved peduncles. Leaves of the involucre short-petioled, twice or thrice 3-parted and cleft, their divisions linear. Sepals red, greenish-yellow, or whitish. Head of carpels spherical or oval, woolly.—Rocky river-margins, etc.

4. *A. cylindrica* Gray. (LONG-FRUITED ANEMONE.) Carpels very numerous, in an oblong woolly head about 1 inch long. Peduncles 2–6, long, upright, leafless. Stem-leaves (involucre) in a whorl, twice or thrice as many as the peduncles, long-petioled. Sepals 5, greenish-white. Plant about two feet high, clothed with silky hairs.—Dry woods.

5. *A. virginia'na* L. (VIRGINIAN ANEMONE.) Much like the last, but larger. Stem-leaves (involucre) 3, each 3-parted. Also, the central peduncle only is naked, the others having each a pair of leaves about the middle, from whose axils other peduncles occasionally spring. Sepals 5, greenish. Head of carpels oval or oblong.—Dry rocky woods and river-banks.

6. *A. canadensis* L. Carpels fewer and the head not woolly, but pubescent and spherical. Stem-leaves sessile, primary ones 3 in a whorl, but only a pair of smaller ones on each side of the flowering branches. Radical leaves 5-7-parted. Sepals 5, obovate, large and white. Plant hairy, scarcely a foot high.—Low meadows.

7. *A. quinquefo'lia* L. (WOOD ANEMONE. WIND-FLOWER.) Plant not more than 6 inches high, nearly smooth, 1-flowered. Stem-leaves in a whorl of 3, long-petioled, 3-5-parted. Sepals 4-7, oval, white, or often purplish on the back.—Moist places.

8. *A. riparia* Fernald. Resembling No. 4, but the leaflets are thinner, and the plant is less pubescent. Sepals 5, large, white or rarely reddish. Fruit-head sub-cylindrical.

3. HEPATICA Hill. LIVER-LEAF. HEPATICA

1. *H. acutifolia* DC. (SHARP-LOBED HEPATICA.) Leaves with 3 (sometimes 5) acute lobes, appearing after the flowers. Petioles silky-hairy.—Woods in spring.

2. *H. trifolia* Chaix. (ROUND-LOBED HEPATICA.) Leaves with 3 rounded lobes appearing after the flowers, those of the involucre also obtuse.—Dry rich woods in spring.

4. THALICTRUM L. MEADOW-RUE

1. *T. dioicum* L. (EARLY MEADOW-RUE.) Stem smooth, pale and glaucous, 1-2 feet high. Flowers dioecious, in ample panicles, purplish or greenish; the yellow anthers drooping and very conspicuous. Leaves alternate, decomposed; leaflets with 5-7 rounded lobes.—Woods.

2. *T. polygamum* Muhl. (TALL MEADOW-RUE.) Stem smooth or nearly so, 2-6 feet high. Leaves sessile; leaflets much like No. 1. Flowers white, in compound panicles; anthers not drooping; filaments club-shaped.—Low wet meadows, and along streams.

3. *T. revolutum* DC. (PURPLISH MEADOW-RUE.) Stem mostly purplish, 2-4 feet high. Stem-leaves sessile or nearly so; leaflets veiny beneath, often glandular-pubescent, and so distinguished from No. 2. Flowers resembling those of No. 1.

5. ANEMONELLA Spach.

1. *A. thalictroides* Spach. (RUE ANEMONE.) Stem low, 4-10 inches high. Stem-leaves all in a whorl at the top. Flowers usually several in an umbel (by which character this plant is easily distinguished from Wood Anemone, which it otherwise resembles). Flower, stem, and 1 or 2 radical leaves rising from a cluster of fleshy tuberous roots.

6. MYOSURUS L. MOUSETAIL

1. *M. minimus* L. Scapes 2-6 inches high. Leaves linear-spatulate. Achenes blunt, in a spike 1-2 inches long when ripe.

7. RANUNCULUS L. CROWFOOT. BUTTERCUP

1. *R. aquatilis* L., var. *capilla-ceus* DC. (WHITE WATER CROWFOOT.) Foliage under water, filiform, usually collapsing when withdrawn from the water. Flowers white, floating, each petal with a little pit on the inside of the claw.—Stagnant pools and slow-flowing streams.

2. *R. circinatus* Sibth. Much like No. 1, but the emerged leaves are sessile, and do not collapse when withdrawn from the water.

3. *R. Cymbalaria* Pursh. (SEA-SIDE CROWFOOT.) Low, smooth, spreading by runners which take root at the joints. Leaves long-petioled, roundish, crenate, rather fleshy. Petals 5-8, yellow. Carpels striate, in an oblong head.

4. *R. delphinifolius* Torr. (YELLOW WATER CROWFOOT.) Like No. 1, but larger, and with yellow flowers. When creeping in the mud, the leaves are round-kidney-shaped, and more or less deeply lobed and toothed.—Ponds and ditches.

5. *R. Flammula* L., var. *repens* Meyer. (CREEPING SPEARWORT.) Stem reclining, rooting at the joints, only 3-6 inches long. Leaves linear, entire, remote. Flowers yellow, $\frac{1}{4}$ inch broad.—Sandy and gravelly shores of ponds and rivers.

6. *R. rhomboides* Goldie. Stem erect, low (3-8 inches), hairy; root-leaves roundish or rhombic-ovate, mostly crenate; lowest stem-leaves similar or 3-5 lobed, the upper nearly sessile and deeply cut into linear lobes. Petals large, exceeding the calyx; achenes orbicular, with a minute beak, in a globular head.—Dry plains, in early summer.

7. *R. affinis* R. Br. (ROUGH-FRUITED CROWFOOT.) Taller than No. 6, more or less pubescent. Root-leaves petioled, usually pedately multifid; stem-leaves nearly sessile, with broadly linear lobes. Petals light yellow, about $\frac{1}{2}$ inch long. Achenes, often hairy, with recurved beaks, forming an oblong head.—Pr. Prov.

8. *R. abortivus* L. (SMALL-FLOWERED CROWFOOT.) Petals shorter than the reflexed calyx. Stem erect, smooth or pubescent, slender. Radical leaves roundish, kidney-shaped, crenate, petiolate; stem-leaves 3-5-parted, sessile. Carpels in a globular head, each with a minute curved beak.—Shady hillsides and wet pastures.

9. *R. sceleratus* L. (CURSED CROWFOOT.) Petals about the same length as calyx. Stem thick, hollow, smooth. Radical leaves 3-lobed; stem-leaves 3-parted, uppermost almost sessile. Head of carpels oblong.—Wet ditches.

10. *R. recurvatus* Poir. (HOOKED CROWFOOT.) Petals shorter than reflexed calyx. Stem hirsute, with stiff spreading hairs. Radical and cauline leaves about alike, long-petioled. Head of carpels globular, each with a long recurved beak.—Woods.

11. *R. pennsylvanicus* L. f. (BRISTLY CROWFOOT.) Petals not longer than the reflexed calyx. Stem erect, hirsute. Leaves ternately divided, divisions of leaves stalked, unequally 3-cleft. Head of carpels oblong, achenes with straight beaks, and so easily distinguished from No. 10.—Wet places.

12. *R. hispidus* Michx. Resembling the last species, but with few-leaved, ascending or reclining stems, not always hirsute. Root a cluster of stout fibres. Calyx hardly reflexed, soon deciduous, much shorter than the petals. Achenes strongly margined, with straight beaks; in a globular or oval head.

13. *R. Macounii* Britton. Stem reclining or ascending, not always hirsute. Calyx only slightly reflexed, sepals shorter than the petals. Beaks of achenes stout and straight.

14. *R. septentrionalis* Poir. Petals much longer than the calyx. Early-flowering stems ascending, putting forth long runners during the summer. Leaves ternate, divisions generally stalked, deeply and sharply lobed, petioles appressed-pubescent. Achenes large, compressed, strongly margined, in globular heads, and with long flat beaks. Peduncles furrowed.—Wet places.

15. *R. repens* L. Much resembling the last in habit, but smaller, and the leaves not so deeply and sharply cut. Flowering later. Leaves often blotched with white. The style shorter than in No. 14, and stigmatic along the whole inner side, persistent.

16. *R. bulbosus* L. (BULBOUS CROWFOOT or BUTTERCUP.) Petals much longer than calyx. Stem erect, about 1 foot high, from bulb-like base. Flowers an inch broad, on furrowed peduncles.—Pastures. Rather rare.

17. *R. acris* L. (TALL CROWFOOT or BUTTERCUP.) Much taller than No. 16. Petals much longer than the calyx. Stem upright, no bulb at the base. Peduncles not furrowed.

18. *R. fascicularis* Muhl. (EARLY CROWFOOT.) Petals much longer than calyx. Plant 5–9 inches high, erect, pubescent with silky hairs. Radical leaves appearing pinnate, the terminal division long-stalked, the lateral ones sessile. Root a bundle of thickened fleshy fibres.—Rocky woods and fields in spring.

8. CAL'THA L. MARSH MARIGOLD

1. *C. palustris* L. (MARSH MARIGOLD.) Stem about a foot high, hollow, round, forking, very glabrous. Flowers golden yellow, 1–1½ inches broad.—Swamps and wet meadows. A very conspicuous plant in early spring.

9. COP'TIS Salisb. GOLDTHREAD

1. *C. trifolia* Salisb. (THREE-LEAVED GOLDTHREAD.) Low and stemless. Scapes 1-flowered, with a single small bract above the middle. Petals much smaller than the sepals.—On logs and about stumps in cedar swamps.

10. AQUILE'GIA L. COLUMBINE

1. *A. canadensis* L. (WILD COLUMBINE.) Stem branching, a foot or more in height, smooth. Leaves decompose; leaflets in threes. Flowers nodding, scarlet outside, yellow within.—Rocky woods and thickets.

2. *A. vulga'ris* L. (GARDEN COLUMBINE.) This species has escaped from cultivation in some places. Spurs hooked. Flowers blue, purple, or whitish.

11. DELPHIN'IUM L. LARKSPUR

1. *D. azu'reum* Michx. (PRAIRIE LARKSPUR.) Carpels 3, the pods erect. Lobes of the leaves numerous, narrowly linear. Raceme strict, but not dense. Spur usually curved upwards.—Pr. Prov.

2. *D. Consol'ida* L. (FIELD LARKSPUR.) Escaped from gardens in a few places. The pistil is single, and the flowers are scattered on the spreading branches. Petals 2, united.

12. ACTAE'A L. BANEERRY

1. *A. ru'bra* Willd. Raceme short, breadth and length being about the same. Pedicels slender. Berries red.—Rich woods.

2. *A. alba* Mill. (WHITE BANEERRY.) Raceme longer than broad. Pedicels thickened in fruit, cherry-coloured. Berries white.—Same localities as No. 1.

13. CIMICIF'UGA L. BUGBANE

1. *C. racemo'sa* Nutt. (BLACK SNAKEROOT.) Stem 3-6 feet high. Resembling a tall Actaea, but easily distinguished by its plume-like raceme of white flowers.—Rich woods.

14. HYDRAS'TIS Ellis. ORANGEROOT. YELLOW PUCCOON

1. *H. canaden'sis* L. (GOLDEN SEAL.) A low plant, bearing a single radical leaf, and a pair of cauline ones near the summit of the simple stem. Flower solitary, terminal, greenish-white. Leaves rounded, cordate, 5-7-lobed, very large when fully grown.—Wet meadows, in early summer.

MAGNOLIA'CEAE (MAGNOLIA FAMILY)

Trees with alternate entire or lobed (not toothed) leaves, the buds of which are covered by the stipules. Sepals 3, coloured, deciduous. Petals 6-9, deciduous. Stamens hypogynous, indefinite, separate; anthers adnate. Carpels numerous, in many rows on an elongated receptacle. Fruit resembling a cone.

1. MAGNO'LIA L.

1. *M. acumina'ta* L. (CUCUMBER TREE.) Leaves thin, oblong-ovate, acutely pointed. Flowers greenish-yellow, bell-shaped. Fruit long, cylindrical, when young resembling a small cucumber.

2. LIRIODEN'DRON L. TULIP TREE

1. *L. Tulipif'era* L. A large and stately tree, growing to a great height in many parts of the western peninsula of Ontario. Leaves smooth, large, truncate, or with a shallow notch at the end. Flowers large, showy, solitary; petals greenish-yellow, marked with orange. Fruit a dry cone, 2-3 inches long, which, at maturity, separates into dry, winged, indehiscent carpels.

ANONA'CEAE (CUSTARD APPLE FAMILY)

Trees or shrubs with alternate, exstipulate, entire leaves, and solitary, axillary, perfect, hypogynous flowers. Sepals 3. Petals 6, in two sets, deciduous. Stamens numerous. Carpels few or many, fleshy in fruit.

1. ASIM'INA Adans. NORTH AMERICAN PAPAW

1. *A. tril'oba* Dunal. (COMMON PAPAW.) Found only in the Niagara Peninsula. A small tree, not unlike a young beech in appearance, and forming thickets near Queenston Heights. Flowers reddish-purple, appearing before the leaves; the three outer petals much larger than the three inner ones. Fruit 2-3 inches long, edible.

MENISPERMA'CEAE (MOONSEED FAMILY)

Woody twiners, with peltate alternate leaves and small dioecious flowers. Sepals and petals yellowish-white, usually 6 of each, the petals in front of the sepals. Stamens numerous. Fruit a drupe, in appearance something like a small grape, with moon-shaped seeds.

1. MENISPER'MUM L. MOONSEED

1. *M. canaden'se* L. (CANADIAN MOONSEED.) A twining plant, found, though not abundantly, in low grounds in rich woods. It may be easily recognized by its usually 7-angled leaves, which are peltate near the edge. Fruit bluish-black.

BERBERIDA'CEAE (BARBERRY FAMILY)

Herbs (or shrubs) with alternate petiolate leaves. Sepals and petals in fours, sixes, or eights (except in the genus *Podophyllum*), with the petals in front of the sepals. Stamens (except in *Podophyllum*) as many as the petals, one before each. Anthers usually opening by 2 valves at the top. Fruit berry-like, or a pod.

Synopsis of the Genera

A. *Petals and stamens 6.*

1. **Ber'beris.** Shrub with yellow wood and yellow flowers in drooping racemes. Stamens irritable. Petals 6, obovate, concave, each with two glandular spots inside at the base. Fruit a sour berry, oblong, scarlet.
2. **Caulophyl'lum.** A purplish herb, flowering in early spring. Petals thick, much shorter than the sepals. Leaves decom-pound.

B. *Petals 6-9. Stamens 8-18.*

3. **Podophyl'lum.** Petals 6-9. Stamens 12-18. Anthers not opening by uplifting valves. Fruit a large berry. Leaves peltate.
4. **Jefferso'nia.** Petals and stamens mostly 8. Anthers opening by uplifting valves. Pod opening by a lid. Leaves radical, divided into 2 leaflets.

1. BER'BERIS L. BARBERRY

1. **B. vulga'ris** L. (COMMON BARBERRY.) Shrub. Leaves on the fresh shoots of the season mostly reduced to branched spines, from whose axils proceed the next year close clusters of obovate-oblong, bristly-toothed leaves, with short, jointed petioles, and many-flowered racemes. Sepals, petals, and stamens 6 each. Outside of sepals are 2-6 bractlets. Petals yellow. Fruit an oblong, sour, scarlet berry.—Cultivated grounds.

2. CAULOPHYLLUM Michx. BLUE COHOSH

1. **C. thalictroi'des** Michx. (BLUE COHOSH.) Plant 1-2 feet high, very glaucous and dull purple when young. Flowers yellowish-green, in a terminal small raceme, appearing in spring before the decom-pound leaves are developed. Sepals 6, with 3 little bractlets at their base. Petals 6, thick and somewhat kidney-shaped, much smaller than the sepals. Stamens 6, one before each petal. Ovary bursting soon after the flowering, and leaving the

two drupe-like seeds naked on their rather thick stalks. Fruit bluish, $\frac{1}{4}$ inch across.—Rich woods.

3. *PODOPHYLLUM* L. MAY APPLE. MANDRAKE

1. *P. peltatum* L. Stem about 1 foot high. Flowerless stems with one large, 7–9-lobed, umbrella-like leaf, peltate in the centre; the flowering ones with two leaves, peltate near the edge, the flower nodding from the fork. Sepals 6, caducous. Petals 6–9, large and white. Stamens 12–18. Fruit large, oval, yellowish, edible.—Found in patches in rich woods.

4. *JEFFERSONIA* Barton. TWIN-LEAF

1. *J. diphylla* Pers. A low plant, flowering in early spring; the solitary white flowers on naked scapes. Sepals 4, fugacious. Petals 8. Stamens 8. Ovary pointed. Stigma 2-lobed. Pod pear-shaped, the top forming a lid. Leaves radical, long-petioled; the blades divided into two leaflets with the outer margins lobed.—Woods.

LAURACEAE (LAUREL FAMILY)

Trees or shrubs with spicy-aromatic bark and leaves, the latter simple (often lobed), alternate and marked with small transparent dots (visible under a lens). Sepals 6, petal-like. Petals wanting. Flowers dioecious or polygamo-dioecious. Stamens in sterile flowers 9, inserted at the base of the calyx. Anthers opening by uplifting valves. Ovary in fertile flowers free from the calyx, 1-celled, with a single ovule hanging from the top of the cell. Style and stigma 1. Fruit a 1-seeded drupe.

1. *SASSAPARILLA* Nees. SASSAPARILLA

1. *S. variifolium* Ktze. A small or moderate-sized tree with yellowish or greenish-yellow twigs and either oval-to-oblong, entire, or oval-to-obovate, 2–3-lobed leaves. Flowers greenish-yellow, in naked corymbs, appearing with the leaves in the axils of the latter. Drupe blue, on a reddish pedicel. The 9 stamens in 3 rows, the 3 inner each with a pair of yellow glands at the base of the filament. Anthers 4-celled, 4-valved.—Rich woods, in Southern and Western Ontario.

2. *BENZOIN* Fabric. WILD ALLSPICE. FEVER-BUSH

1. *B. aestivale* Nees. (SPICE-BUSH.) A nearly smooth shrub with oblong-obovate leaves, pale beneath. Flowers honey-yellow

in lateral, umbel-like clusters, before the leaves. Stamens very much as in *Sassafras*, but the anthers are 2-celled and 2-valved. Pistillate flowers with 15–18 rudiments of stamens. Drupe red.—Damp woods, in early spring.

PAPAVERACEAE (POPPY FAMILY)

Herbs with milky or coloured juice and alternate leaves without stipules. Flowers polyandrous, hypogynous. Sepals 2, caducous. Petals 4–12. Stamens numerous, anthers introrse. Fruit a 1-celled pod (in Poppy imperfectly many-celled), with numerous seeds.

Synopsis of the Genera

1. *Chelidonium*. Juice yellow. Petals 4, crumpled in the bud, yellow.
2. *Sanguinaria*. Juice red. Petals 8–12, not crumpled in the bud, white.
3. *Papaver*. Juice milky. Petals mostly 4. Ovary imperfectly many-celled. Stigmas united into a radiate sessile crown.

1. CHELIDONIUM L. CELANDINE

1. *C. majus* L. Petals 4, deciduous, crumpled in the bud. Juice of the plant yellow. Flower-buds nodding. Flowers small, yellow, in a kind of umbel. Fruit a smooth, 1-celled, slender pod, from which the two valves fall away, leaving the parietal placentas as a slender framework, with the seeds attached.—Waste places.

2. SANGUINARIA L. BLOOD-ROOT

1. *S. canadensis* L. Petals 8–12, not crumpled in the bud. Flower-buds not nodding. A stemless plant, with a thick rhizome which emits a red juice when cut, and sends up in early spring a single rounded, 5–7-lobed, thickish leaf, and a 1-flowered scape. Flowers white.—Rich woods.

3. PAPAVER L. POPPY

1. *P. somniferum* L. (COMMON POPPY.) Smooth and glaucous. Leaves clasping, wavy, cut-toothed. Pod globose. Petals white or purple, the buds nodding.—Waste places and old gardens.

2. *P. Rhoeas* L. (CORN POPPY.) Sparingly hispid. Leaves deeply pinnatifid, the lobes cut-toothed. Pod globular, smooth.

Corolla 2-4 inches in diameter, scarlet, often with a dark centre.—Waste-heaps, Atl. Prov.

3. *P. du'bium* L. Smaller than the last, and the lobes of the leaves narrower. Pod oblong, narrowed at the base. Corolla a paler red.—Ballast-heaps, Atl. Prov.

FUMARIA'CEAE (FUMITORY FAMILY)

Smooth herbs with brittle stems, watery juice, dissected leaves, and irregular flowers. Sepals 2, very small. Corolla flattened and closed, of 4 petals, the two inner united by their tips over the anthers of the 6 stamens. Stamens in two sets of 3 each; filaments often united; the middle anther of each set 2-celled, the others 1-celled. Fruit a 1-celled pod.

Synopsis of the Genera

1. *Adlu'mia*. Corolla 2-spurred. Petals all permanently united. Plant climbing.
2. *Dicen'tra*. Corolla 2-spurred. Petals slightly united, easily separated. Not climbing.
3. *Coryd'alis*. Corolla 1-spurred. Fruit a slender pod, many-seeded.
4. *Fuma'ria*. Corolla 1-spurred. Fruit a globular, 1-seeded nutlet, indehiscent.

1. ADLU'MIA Raf. CLIMBING FUMITORY

1. *A. fungo'sa* Greene. A smooth vine, climbing by the petioles of its decompound leaves. Flowers in axillary pendulous clusters, pale pink.—Low and shady grounds, and rocky woods.

2. DICEN'TRA Bernh. DUTCHMAN'S BREECHES

1. *D. Cuculla'ria* Bernh. (DUTCHMAN'S BREECHES.) Leaves all radical, multifid; these and the slender scapes rising from a bulb-like rhizome of coarse grains. Flowers several in a raceme, whitish, spurs divergent, elongated, acute, straight.—Rich woods.

2. *D. canaden'sis* Walp. (SQUIRREL CORN.) Underground shoots bearing small yellow tubers, something like grains of corn. Leaves very much as in No. 1. Corolla merely heart-shaped; spurs very short and rounded. Flowers greenish-white, rose-tinged, fragrant.—Rich woods.

3. CORYD'ALIS Medic. CORYDALIS

1. *C. aur'ea* Willd. (GOLDEN CORYDALIS.) Stems low and spreading. Leaves dissected. Flowers in simple racemes, golden yellow, the outer petals keeled, but not crested on the back. Spur about $\frac{1}{2}$ as long as the corolla. Pods pendulous.—Rocky river-margins and burnt woods.

Var. *occidenta'lis* Engelm. Larger flowers with spurs almost as long as their bodies.—Pr. Prov.

2. *C. flav'ula* DC. (YELLOW CORYDALIS.) Stems low and spreading. Flowers pale yellow, the outer petals wing-crested on the back; crest 3-4-toothed.

3. *C. semper'virens* Pers. (PALE CORYDALIS.) Stems upright, 1-4 feet high. Flowers in compound racemes, purplish, tipped with yellow. Pods erect.—Rocky woods.

4. FUMA'RIA L. FUMITORY

1. *F. officina'lis* L. (COMMON FUMITORY.) Corolla flesh-coloured, tipped with crimson. Flowers small, in dense racemes or spikes.—Waste places near dwellings.

CRUCIF'ERAE (CRESS FAMILY)

Herbs with a pungent watery juice, alternate leaves without stipules, and regular hypogynous flowers in racemes or corymbs. Pedicels without bractlets. Sepals 4, deciduous. Petals 4, forming a cross-shaped corolla, rarely apetalous. Stamens 6, two of them shorter, or only 2. Fruit usually a silique or silicle.

Synopsis of the Genera

A. *Pod dehiscent by 2 valves, usually much longer than broad.*

1. *Radic'ula*. Flowers white or yellow. Pod terete, oblong-linear or ellipsoid. Seeds in two rows in each cell, globular, without a wing.
2. *Barbare'a*. Flowers yellow. Pod somewhat 4-sided, the valves strongly 1-nerved. Seeds in one row in each cell.
3. *Denta'ria*. Flowers white or pale purple. Pod lanceolate, flat. Seeds wingless, on broad seed-stalks. Stem-leaves 2 or 3 in a whorl; stem naked below. Rootstock toothed or tuberous.

4. **Cardam'ine.** Flowers white or rose-coloured. Pod linear or lanceolate, flat, the valves nerveless. Seeds wingless, on slender seed-stalks. Stem leafy below.
5. **Ar'abis.** Flowers white, rose-coloured, or rarely yellow. Pod linear or elongated, flattened, the valves usually with a distinct mid-rib. Stem leafy.
6. **Erys'imum.** Flowers yellow (or creamy). Pod linear, distinctly 4-sided. Stigma broadly 2-lobed. Pedicels of the pods usually diverging from the stem. Leaves simple, not clasping. Pubescent with appressed hairs.
7. **Sisym'brium.** Flowers yellow or white, small. Pods awl-shaped or 4-6-sided, the valves 1-3-nerved. Stigma small.
8. **Hes'peris.** Flowers large, purple. Pod terete, elongated. Stigma lobes narrow, erect.
9. **Bras'sica.** Flowers yellow. Pod linear or oblong, nearly terete, or 4-sided, with a distinct beak extending beyond the end of the valves. Seeds usually in one row.
10. **Diplo'tax'is.** Sepals, petals, and stamens almost as in *Brassica*. Pods slender, on slender spreading pedicels. Beak conical, usually short. Seeds in two rows.

B. Pod dehiscent by 2 valves, comparatively short.

(1) *Silicle compressed parallel with the broad partition or globular.*

11. **Lesquerel'la.** Flowers yellow. Plant densely hoary with stellate hairs. Pods ovate or globose. Seeds flattened, in two rows in each cell.
12. **Dra'ba.** Flowers white. Pod flat; seeds several or many, 2 rows in each cell. Low herbs.
13. **Alys'sum.** Flowers pale yellow or white. Pod orbicular, flat, 2-4-seeded.
14. **Camel'ina.** Flowers yellow. Pod pear-shaped, pointed; valves 1-nerved.

(2) Silicle compressed contrary to the narrow partition.

15. **Capsel'la.** Flowers white. Pod obcordate-triangular; valves boat-shaped, wingless. Seeds numerous.
16. **Thlas'pi.** Flowers white. Pod obovate or obcordate, winged. Seeds several.
17. **Lepid'ium.** Flowers white or greenish. Pod roundish, very flat; the valves boat-shaped and winged. Seeds solitary.
18. **Coron'opus.** Flowers whitish, very small. Pod 2-seeded, the valves wrinkled and separating at maturity as 2 closed, one-seeded nutlets. Prostrate diffuse herbs, with a disagreeable characteristic odour.

C. Pod indehiscent.

19. *Caki'le*. Flowers purplish. Pod 2-jointed, fleshy. Leaves fleshy.
20. *Raph'anus*. Flowers yellow, turning whitish or purplish. Pod elongated, several-seeded, mostly constricted between the seeds.
21. *Nes'lia*. Flowers very much as in the last. Pod nearly globular, hardly $\frac{1}{12}$ inch across, usually only 1-celled and 1-seeded, wrinkled, tipped with the slender style.

1. RADIC'ULA Hill. WATER CRESS

1. *R. Nastur'tium-aquat'icum* B. and R. (WATER CRESS.) Flowers white. Stem spreading and rooting. Leaves pinnate; leaflets 3-11, roundish or oblong, nearly entire. Pods oblong-linear.—Ditches and streamlets.

2. *R. palus'tris* Moench. (MARSH CRESS.) Flowers yellow. Stem erect. Leaves pinnately parted, the lobes cut-toothed. Pods ovoid.—Wet places.

Var. *his'pida* Robinson. Hairy, pods nearly globular.

3. *R. aquat'ica* Robinson. (LAKE CRESS.) Flowers white. An aquatic plant, with the submerged leaves finely dissected; the leaves out of the water oblong, and either entire, serrate, or pinnatifid. Pods ovoid, 1-celled.

4. *R. Armora'cia* Robinson. (HORSERADISH.) Has escaped from gardens in many places. Flowers white. Root-leaves very large, oblong, and generally crenate; stem-leaves lanceolate. Pods globular. Roots very large.

5. *R. trachycar'pum* Gray. Flowers white. Stem nearly glabrous, erect, branching. Leaves lyrate-pinnatifid. Pods oblong-linear, somewhat roughened, soon recurved, shorter than the awl-shaped style.—Pr. Prov., in beds of pools and streams.

6. *R. amphib'ium* R. Br. Flowers white. Stem tall, erect, branching above, sparingly pubescent. Leaves lyrate-pinnatifid, rough-pubescent, the lobes coarsely toothed. Pods several times longer than the style, but much shorter than the pedicels.—Wet places.

2. BARBARE'A R. Br. WINTER CRESS

1. *B. vulga'ris* R. Br. (YELLOW ROCKET.) Stem smooth, 1-3 feet high. Lower leaves lyrate, the terminal division round

and generally large, and 1–5 pairs of lateral ones (the latter sometimes absent); upper leaves obovate, more or less pinnatifid at the base. Pods linear, erect, or slightly spreading.

2. *B. stric'ta* Andrz. Foliage similar to No. 1. Flowers pale yellow and the pods appressed.—Pr. Prov.

3. DENTA'RIA L. TOOTHWORT. PEPPER-ROOT

1. *D. diphy'l'a* Michx. (TWO-LEAVED TOOTHWORT.) Flowers white. Stem-leaves 2, nearly opposite, ternately divided. Rootstock toothed, pleasantly pungent to the taste.—Rich woods.

2. *D. lacinia'ta* Muhl. (LACINIATE TOOTHWORT.) Flowers purplish, or white. Stem-leaves 3, in a whorl. Rootstock jointed, scarcely toothed.—Rich woods.

4. CARDAM'INE L. BITTER CRESS

1. *C. bulbo'sa* BSP. (SPRING CRESS.) Flowers white, sepals greenish. Stem tuberous at the base. Lower leaves round-cordate; upper nearly lanceolate; all somewhat angled or toothed.—Wet meadows.

2. *C. Douglas'sii* Britton. Similar to No. 1, but the sepals are purple-tinged, and the petals are rose-purple.

3. *C. praten'sis* L. (CUCKOO-FLOWER. LADY'S SMOCK.) Flowers white or rose-coloured, showy. Stem from a short rootstock. Leaves pinnate, leaflets 7–15, those of the lower leaves rounded and stalked, entire or nearly so.—Bogs.

4. *C. hirsu'ta* L. (SMALL BITTER CRESS.) Flowers white, small. Root fibrous. Leaves pinnate, leaflets 5–11, the terminal leaflets largest. Pods erect, slender.—Wet places.

5. AR'ABIS L. ROCK CRESS

A. Seeds in one row in each cell.

1. *A. lyra'ta* L. (LOW ROCK CRESS.) Flowers white, petals twice as long as calyx. Stem branching from base. Radical leaves clustered, pinnatifid, the terminal lobe largest; stem-leaves scattered, linear, with tapering base. Pods slender, erect, and spreading.—Rocky or sandy shores.

2. *A. hirsu'ta* Scop. (HAIRY ROCK CRESS.) Flowers greenish-white, small, petals slightly longer than calyx. Stem-leaves many,

rough, sagittate; often a dense rosette at base of stem. Pods erect, straight. Stems 1-2 feet high, 2 or 3 from the same root.—Rocky shores and dry plains.

3. *A. laeviga'ta* Poir. (SMOOTH ROCK CRESS.) Flowers white, rather small. Leaves linear or lanceolate, entire or slightly toothed, sagittate, clasping. Pods long and narrow, recurved-spreading. Stem glaucous, 1-2 feet high. Easily recognized by the pods.—Dry hillsides.

4. *A. canaden'sis* L. (SICKLE-POD.) Flowers whitish, with linear petals, about twice the length of the calyx. Stem-leaves pointed at both ends, downy, the lower ones toothed. Pods 2-3 inches long, scythe-shaped, hanging. Stem 2-3 feet high. A striking plant when the pods are fully formed.—Dry woods and ravines.

5. *A. denta'ta* T. and G. Flowers whitish, petals about the same length as the calyx. Leaves oblong, sharply toothed, those of the stem half-clasping. Pods very slender, widely spreading.

6. *A. gla'bra* Bernh. (TOWER MUSTARD.) Flowers yellowish-white. Petals scarcely longer than the calyx. Stem 2-4 feet high, glaucous. Cauline leaves ovate-lanceolate or oblong, clasping with sagittate base. Pods long and very narrow, on erect pedicels. Easily recognized by its strict habit.

B. Seeds in two distinct rows in each cell.

7. *A. Drummon'di* Gray. Flowers white or rose-coloured. Petals twice as long as the calyx. Stem 1-2 feet high, smooth above. Cauline leaves lanceolate or oblong-linear, with sagittate base; lowest leaves spatulate and toothed. Pods long and flat; the pedicels not so strictly erect as in the last species.—Rocky banks of streams.

8. *A. Holboel'lii* Hornem. Petals white or rose-coloured, occasionally purplish, becoming reflexed, twice as long as the sepals. Stem 6-24 inches high, usually simple, more or less hairy. Lowest leaves small, spatulate, somewhat toothed, with margined petioles; upper sessile, sagittate, partly clasping, linear-lanceolate. Pods reflexed. Style none.—Pr. Prov.

6. ERYSIMUM L. TREACLE MUSTARD

1. *E. cheiranthoides* L. (WORM-SEED MUSTARD.) Flowers yellow, inconspicuous. Stem slender, branching. Leaves lanceolate, scarcely toothed, roughish with appressed pubescence. Pods small and short, on slender diverging pedicels.—Waste wet places.

2. *E. asperum* DC. (PRAIRIE ROCKET.) Flowers showy, bright orange-yellow, rarely purple. Stem stout, 1–2 feet high, simple, hoary, with minute appressed hairs. Leaves lanceolate to linear. Pods long, ascending on stout spreading pedicels.—Dry soil, Pr. Prov.

3. *E. parviflorum* Nutt. Stem low and simple, hoary. Leaves all linear-oblong, densely clustered at the base of the stem. Flowers small, sulphur-yellow. Pods long, narrow, ascending, on short pedicels.—Pr. Prov.

4. *E. orientale* R. Br. Stem slender and branching, 1–2 feet high. Leaves grayish-green, oblong to oval, slightly clasping; radical ones spatulate. Flowers pale yellow or cream-coloured, small.

7. SISYMBRIUM L. HEDGE MUSTARD

1. *S. officinale* Scop. (HEDGE MUSTARD.) Flowers yellow, small. Leaves runcinate. Stem 1–2 feet high, with spreading branches. Pods awl-shaped, close pressed to the stem.—A very common roadside weed.

2. *S. canescens* Nutt. (TANSY MUSTARD.) Flowers yellowish, very small. Leaves twice pinnatifid, hoary with short branching hairs, the divisions toothed. Pods acute at each end, shorter than the slender spreading pedicels.—Pr. Prov.

3. *S. Sophia* L. Hoary. Leaves dissected into very numerous small leaflets. Pods very slender, about an inch long, ascending.—Scarce.

4. *S. altissimum* L. Tall, branching. Loosely pubescent below, smooth above. Leaves deeply pinnatifid, the lobes narrowly linear. Pods very long (3–4 inches) and slender, spreading.—An introduced weed which has now become a pest in the Pr. Prov.

5. *S. incisum* Engelm., var. *Hartwegianum* Watson. Pinnate leaves, the leaflets narrowly oblong and toothed. Pods only $\frac{1}{8}$ – $\frac{1}{4}$ inch long, in a crowded raceme.—Pr. Prov.

8. HES'PERIS L. ROCKET

1. *H. matrona'lis* L. (GARDEN ROCKET.) Tall, leaves lanceolate, acuminate, serrate, usually petiolate. Flowers large, purple. Pods 2-4 inches long, spreading.—Waste places.

9. BRAS'SICA L. CABBAGE. MUSTARD. TURNIP

1. *B. arven'sis* Ktze. (CHARLOCK.) Flowers bright yellow. Stem 1-2 feet high, branching, it and the leaves hairy. Upper leaves rhombic. Pods about $1\frac{1}{2}$ inches long, at least $\frac{1}{3}$ occupied by a stout flat beak.—Too common in our grain fields.

2. *B. ni'gra* Koch. (BLACK MUSTARD.) Flowers sulphur-yellow. Stem 3-6 feet high, round, hirsute and branching. Lower leaves lyrate.—Pods much shorter than in the last and the beak smaller.—Fields and waste places.

3. *B. campes'tris* L. Stem tall and erect, it and the leaves smooth when old. Lower leaves ovate or elliptical, coarsely toothed, on long petioles; upper ones narrow and clasping. Pods an inch long or more, on ascending pedicels; beak about $\frac{1}{3}$ the length of the pod.—A common weed in Manitoba.

10. DIPLTAX'IS DC.

1. *D. mura'lis* DC. Branching from near the base. Smooth or sparingly hispid. Leaves oblanceolate, shallowly and bluntly toothed or pinnatifid. Petals pale yellow. Pods over an inch long, on slender spreading pedicels half as long as the pods. Valves 1-nerved.—Atl. Prov.

2. *D. tenuifo'lia* DC. Like the last, but the leaves are deeply sinuate-pinnatifid, with narrow lobes. Pedicels of the pods over an inch long.

11. LESQUEREL'IA Wats. BLADDER-POD

1. *L. argen'tea* MacM. Pubescent. Leaves linear-oblanceolate. Pods hanging on recurved pedicels.—Pr. Prov.

12. DRA'BA L. WHITLOW GRASS

1. *D. arab'isans* Michx. Hoary-pubescent. Flowers white. Leaves lanceolate or oblanceolate to ovate, entire or sparingly toothed. Pods elliptic-lanceolate, much-twisted, on short divergent pedicels. Style short.—Dry rocks.

2. *D. nemoro'sa* L. A small slender plant with leaves about $\frac{3}{4}$ inch long. Flowers whitish. Leaves oblong, or somewhat lanceolate, hairy, sessile, serrate. Racemes elongated. Pods elliptical-oblong, half as long as the wide-spreading pedicels, pubescent or smooth. Style none.—Pr. Prov.

3. *D. carolinia'na* Walt. Flowers white. Stem 1-5 inches high. Leaves obovate, entire. Raceme very short. Pods broadly linear, smooth, longer than the ascending pedicels. Style none.—Rocks; Southern Ontario.

13. ALYS'SUM L. ALYSSUM

1. *A. alyssoides* L. A dwarf hoary annual, with linear-spatulate leaves. Calyx persistent. Pod 4-seeded, sharp-edged.—Rather rare; abundant at Queenston Heights.

14. CAMELINA Crantz. FALSE FLAX

1. *C. sativa* Crantz. (COMMON FALSE FLAX.) Flowers yellowish. Stem 1-2 feet high, straight, erect, branching. Leaves lanceolate, sagittate. Pods pear-shaped, large, margined; style slender.—In flax fields.

15. CAPSULA Medic. SHEPHERD'S PURSE

1. *C. Bursa-pastoris* Medic. Flowers small, white. Root-leaves clustered, pinnatifid; stem-leaves clasping, sagittate.—A very common weed.

6. THLASPI L. PENNY CRESS

1. *T. arven'se* L. (FIELD PENNY CRESS. FRENCH WEED.) A low smooth plant with undivided radical leaves, and stem-leaves sagittate and clasping. Pods half an inch broad, deeply notched at the top.—Waste places.

17. LEPIDUM L. PEPPER GRASS

1. *L. virgin'icum* L. (WILD PEPPER GRASS.) Flowers small; petals present, white. Stem 1-2 feet high. Leaves lanceolate, the upper linear or lanceolate and entire, the lower toothed or pinnatifid, tapering towards the base. Pods marginless or nearly so, oval or orbicular.—Railways and roadsides.

2. *L. apetalum* Willd. Distinguished from No. 1 by the usual absence of petals, and the pods being minutely winged at the top.—Dry sandy fields.

3. *L. rudera'le* L. Petals always absent. More branched than the preceding. Lower leaves bi-pinnatifid.

4. *L. campes'tre*. R. Br. Well distinguished from other species by its sagittate clasping leaves. Pods ovate, winged.—Rather rare.

5. *L. sati'vum* L. (GARDEN CRESS.) Leaves variously divided and cut, with numerous roundish winged pods, and flowers white or rose-coloured.—Escaped from cultivation in some places.

6. *L. Dra'ba* L. An obscurely hoary perennial. Upper leaves auricled, lower sessile, oblong. Pods ovate, wingless, slightly heart-shaped, the style conspicuous.—Abundant about Clifton, Ont.

18. CORON'OPUS Ludwig. WART CRESS. SWINE CRESS

1. *C. did'ymus* Sm. Pods notched at the apex.—Atl. Prov.

2. *C. procum'bens* Gilibert. Pods not notched.—Atl. Prov.

19. CAKI'LE Ludwig. SEA-ROCKET

1. *C. eden'tula* Hook. (AMERICAN SEA-ROCKET.) Flowers purplish. Leaves obovate, fleshy, wavy-toothed. Pod fleshy, 2-jointed.—Seashore, and borders of the Great Lakes.

20. RAPH'ANUS L. RADISH

1. *R. Raphanis'trum* L. (WILD RADISH.) Pods linear or oblong, tapering, indehiscent, long-beaked, necklace-form when ripe. Style long. Leaves rough, lyrate. Petals yellow, veiny, turning whitish or purplish.—An introduced weed; chiefly eastward.

21. NESLIA Desv. BALL MUSTARD

1. *N. panicula'ta* Desv. Flowers yellow, in small panicles. Leaves oblong, sagittate-clasping.

CAPPARIDA'CEAE (CAPER FAMILY)

Herbs (in Canada) with an acrid watery juice, and alternate, palmately compound leaves. Flowers cruciform. Stamens 6 or more. Pod like that of a Crucifer, but only 1-celled.

Synopsis of the Genera

1. *Polanis'ia*. Stamens 8 or more.
2. *Cleo'me*. Stamens 6.

1. *POLANIS'IA* Raf. *POLANISIA*

1. *P. grave'olens* Raf. A strong-scented herb with a viscid hairy stem. Leaflets 3, oblong. Flowers in terminal racemes. Sepals 4. Petals 4, yellowish-white, narrowed below into long claws. Stamens 8-12, exserted. Pod glandular-pubescent, 2 inches long, linear.—Sandy lake-shores.

2. *P. trachysper'ma* Torr. and Gray. Has larger flowers and strongly exserted stamens, and a sessile pod.—Pr. Prov.

2. *CLEO'ME* L.

1. *C. serrula'ta* Pursh. Stem 2-3 feet high, glabrous. Flowers showy, rose-coloured or white, in leafy-bracted racemes. Leaves of 3 lanceolate or oblong leaflets. Stipe of the pod as long as the pedicel of the flower.—Dry clay soil, Pr. Prov.

SARRACENIA'CEAE (PITCHER-PLANT FAMILY)

Bog plants, easily distinguished by their pitcher-shaped leaves, all radical.

1. *SARRACE'NIA* L. SIDE-SADDLE FLOWER

1. *S. purpu'rea* L. (PURPLE SIDE-SADDLE FLOWER. HUNTS-MAN'S CUP.) Leaves hollow, with a wing on one side, purple-veined, curved, with hood erect and open. Sepals 5, coloured, with 3 small bractlets at base. Petals 5, fiddle-shaped, curved over centre of flower, deep purple. Ovary 5-celled, globose, the short style expanding above into a 5-angled umbrella, with a hooked stigma at each angle. Flowers on naked scapes, nodding.—Bogs.

DROSERA'CEAE (SUNDEW FAMILY)

Low, glandular-hairy, marsh herbs with circinate, tufted, radical leaves, the glands of which exude clear, sticky, glittering drops. The regular hypogynous flowers are borne on a naked scape. Sepals, petals, and stamens, 5 each, anthers turned outwards. Styles 3-5, deeply 2-parted. Pod 1-celled, 3-valved.

1. *DROS'ERA* L. SUNDEW

1. *D. rotundifo'lia* L. (ROUND-LEAVED SUNDEW.) Flowers small, white, in a 1-sided, nodding raceme. Leaves orbicular or nearly so, abruptly narrowed into the hairy petiole, clothed with reddish glandular hairs.—Bogs.

2. *D. longifo'lia* L. (LONGER-LEAVED SUNDEW.) Oblong spatulate leaves gradually narrowed into erect naked petioles.—Bogs; not common.

3. *D. linea'ris* Goldie. Resembles the last, but the leaves are linear and the flowers sometimes pinkish.

CRASSULA'CEAE (ORPINE FAMILY)

Succulent herbs (except in one genus), chiefly differing from Saxifragaceae in having symmetrical flowers, the sepals, petals, and carpels being the same in number, and the stamens usually twice as many.

Synopsis of the Genera

1. *Pen'thorum*. Not succulent. The carpels united, forming a 5-celled pod.

2. *Se'dum*. Succulent. Carpels distinct.

1. PEN'THORUM L. DITCH STONE-CROP

1. *P. sedoi'des* L. Not succulent. Sepals 5. Petals 5, if any; sometimes wanting. Stamens 10. Pod 5-angled, 5-horned, and 5-celled. Leaves scattered, lanceolate, acute at both ends. A homely weed with greenish-yellow flowers in a loose cyme.—Wet places.

2. SE'DUM L. STONE-CROP. ORPINE

1. *S. acre* L. (MOSSY STONE-CROP.) Leaves very thick and succulent, crowded, very small. Sepals and petals 4–5. Petals yellow. Stamens 8–10. A spreading, moss-like plant, which has escaped from cultivation in many places.—Roadsides.

2. *S. purpu'reum* Tausch. (LIVE-FOR-EVER.) Stems tall and stout. Leaves oval, toothed. Flowers in compound cymes, petals purple. Sepals, petals, and carpels 5 each. Stamens 10. (Introd. from Europe.)

3. *S. ro'seum* Scop. (ROSE-ROOT.) Stems 5–10 inches high. Flowers dioecious, greenish-yellow or purplish. Stamens mostly 8, other parts in fours.—Rocky shores, Atl. Prov.

SAXIFRAGA'CEAE (SAXIFRAGE FAMILY)

Herbs or shrubs, distinguished from Rosaceae chiefly in having opposite as well as alternate leaves, and usually no stipules; stamens only as many or twice as many as the (usually 5) petals;

and the carpels fewer than the sepals (mostly 2), and usually more or less united with each other. Stamens and petals generally inserted on the calyx.

Synopsis of the Genera

1. **Ri'bes.** Shrubs, sometimes prickly, with alternate and palmately veined and lobed leaves, which are plaited in the bud. Calyx 5-lobed, the tube adherent to the ovary (superior). Petals 5, small, inserted on the calyx. Stamens 5. Styles 2. Fruit a many-seeded berry.
2. **Parnas'sia.** Smooth herbs with entire and chiefly radical leaves, and solitary flowers terminating the long scapes. Petals 5, large, veiny, each with a cluster of sterile filaments at the base. Proper stamens 5. Stigmas 4. Pod 4-valved. Calyx free from the ovary.
3. **Saxif'raga.** Herbs with clustered root-leaves. Flowers at first in close cymes. Calyx lobes hardly adherent to the ovary. Petals 5. Stamens 10. Styles 2. Fruit a pair of follicles, slightly united at the base.
4. **Mitel'la.** Low and slender herbs with round-heart-shaped, radical leaves, those on the scape (if any) opposite. Flowers in terminal racemes. Calyx 5-lobed, adherent to the base of the ovary. Petals 5, slender, pinnatifid. Stamens 10, short. Styles 2. Pod 2-beaked, but 1-celled.
5. **Tiarel'la.** Slender herbs with radical, heart-shaped leaves, and leafless scapes, bearing a simple raceme of flowers. Calyx bell-shaped, 5-parted. Petals 5, entire. Stamens 10, long and slender. Styles 2. Pod 2-valved, the valves unequal.
6. **Heuche'ra.** Perennial herbs with round-heart-shaped, radical leaves; those on the stem (if any) alternate. Greenish or purple flowers, clustered in a long narrow panicle. Petals 5, small, spatulate, entire. Stamens 5. Styles 2. Pod 2-beaked.
7. **Chrysosple'nium.** Small and smooth herbs, with mostly opposite roundish leaves. Calyx tube adherent to the ovary. Petals none. Stamens twice as many as the calyx lobes (8-10), inserted on a conspicuous disk. Styles 2. Pod 2-lobed.

1. RIBES L. CURRANT. GOOSEBERRY

1. **R. Cynos'bati** L. (WILD GOOSEBERRY.) Stem with small thorns at the bases of the leaves, the latter downy, on slender petioles, roundish-heart-shaped, 3-5-lobed. Peduncles slender, 2-3-flowered. Berry covered with long prickles, rarely smooth.—Open woods and clearings.

2. *R. oxyacanthoides* L. (SMALL WILD GOOSEBERRY.) Stems with very short thorns or none. Peduncles very short, 1-2-flowered. Stamens shorter than, or equalling, the broadly oblong calyx lobes. Berry small, smooth.—Low grounds.

3. *R. rotundifolium* Michx. Like the last, but the stamens are longer than the narrowly oblong-spatulate calyx lobes.—Chiefly eastward.

4. *R. lacustris* Poir. (SWAMP GOOSEBERRY.) Shrubby. Young stems prickly, and thorny at the bases of the leaves. Leaves cordate, deeply 3-5-lobed, the lobes deeply cut. Racemes 4-9-flowered, slender, nodding. Fruit bristly.—Swamps and wet woods.

5. *R. floridum* L'Her. (WILD BLACK CURRANT.) Stems and fruit without prickles or thorns. Leaves resinous-dotted, sharply 3-5-lobed, doubly serrate. Racemes many-flowered, drooping. Calyx bell-shaped. Fruit black, smooth.—Woods.

6. *R. triste* Pall. (SWAMP RED CURRANT.) A low shrub with straggling stems. Leaves obtusely 3-5-lobed. Racemes from lateral buds separate from the leaf-buds, drooping. Calyx flat. Fruit red, smooth.—Bogs and wet woods.

7. *R. prostratum* L'Her. (FETID CURRANT.) Stems reclined. Leaves deeply cordate, 5-7-lobed, smooth, the lobes ovate, acute, doubly serrate. Racemes erect, slender, the flowers greenish. Fruit pale red, glandular-bristly.—Cold damp woods and rocks.

8. *R. hudsonianum* Richards. Like the last, but with white flowers crowded in the erect raceme, and darker and smooth fruit.—Pr. Prov. Also in Central Ontario.

2. PARNASSIA L. GRASS OF PARNASSUS

1. *P. caroliniana* Michx. Petals sessile, very veiny. Sterile filaments 3 in each set. Leaves ovate or rounded, usually only one low down on the stalk. Flower an inch across, the petals much longer than the calyx.—Beaver meadows and wet banks.

2. *P. palustris* L. Sterile filaments 9-15 in each set. Scapes 3-10 inches high. Flower nearly an inch across, the petals much longer than the sepals. Leaves cordate.—Sandy banks.

3. *P. parviflora* DC. Petals but little longer than the sepals. Sterile filaments about 7 in each set. Leaves ovate or oblong.—Atl. Prov. and Pr. Prov.

3. SAXIF'RA L. SAXIFRAGE

1. *S. virginien'sis* Michx. (EARLY SAXIFRAGE.) Stem 4-9 inches high. Scape clammy. Leaves obovate, crenately toothed, thickish. Petals white, oblong, twice as long as the sepals.—Rocks and hillsides.

2. *S. tricuspidata* Rottb. Stems tufted, 4-8 inches high. Leaves alternate, oblong or spatulate, with 3 rigid sharp teeth at the end. Petals yellow.—Pr. Prov.

3. *S. Aizo'on* Jacq. Scape 5-10 inches high. Leaves thick, spatulate, with white, finely-toothed margins. Petals cream-coloured, obovate, often spotted at the base.—Moist rocks, Atl. sea-coast.

4. MITEL'LA L. MITREWORT. BISHOP'S CAP

1. *M. diphylla* L. (TWO-LEAVED MITREWORT.) Stem hairy. Leaves cordate, 3-5-lobed, those on the scape 2, opposite, nearly sessile. Flowers white.—Rich woods.

2. *M. nu'da* L. (NAKED-STALKED MITREWORT.) Stem small and delicate. Leaves kidney-shaped, doubly crenate. Scape leafless, few-flowered. Flowers greenish.—Deep woods, on moss-covered logs, etc.

5. TIAREL'LA L. FALSE MITREWORT

1. *T. cordifolia* L. Scapes leafless, 5-12 inches high. Leaves heart-shaped, sharply toothed, sparsely hairy above, downy beneath. Petals white, oblong.—Rich woods.

6. HEUCHE'RA L. ALUM-ROOT

1. *H. americana* L. (COMMON ALUM-ROOT.) Stems 2-3 feet high, glandular and short-hairy. Flowers small, in a loose panicle. Petals equalling or longer than the calyx lobes. Stamens and style exserted.

2. *H. hispida* Pursh. Larger flowers in a very narrow panicle, and taller stems, with long spreading hairs. Stamens short, but soon exserted.—Pr. Prov.

7. CHRYSOSPLE'NIUM L. GOLDEN SAXIFRAGE

1. *C. americanum* Schwein. A low and delicate smooth herb, with spreading and forking stems. Leaves opposite, round and slightly crenate. Flowers greenish-yellow, inconspicuous, nearly sessile in the forks.—Shady wet places.

HAMAMELIDA' CEAE (WITCH-HAZEL FAMILY)

Tall shrubs with alternate simple leaves and deciduous stipules. Flowers in clusters or heads, often monoecious. Calyx 4-parted, adherent to the base of the ovary, the latter of 2 united carpels. Fruit a 2-beaked, 2-celled, woody pod, opening at the top. Petals 4, strap-shaped, inserted on the calyx. Stamens 8, 4 of them anther-bearing, the remainder reduced to scales. The only genus with us is

1. HAMAME'LIS L. WITCH-HAZEL

1. *H. virginiana* L. Leaves obovate or oval, crenate or wavy-toothed, pubescent. Flowers yellow, appearing late in the autumn. —Damp woods.

PLATANA'CEAE (PLANE-TREE FAMILY)

Large trees with palmately lobed leaves; the base of the petioles inclosing the buds of the next season. Monoecious flowers, the sterile and fertile in separate spherical heads.

1. PLAT'ANUS L. PLANE-TREE. BUTTONWOOD

1. *P. occidentalis* L. (AMERICAN PLANE-TREE or SYCAMORE.) A fine large tree found in South-western Ontario. Leaves alternate, rather scurfy when young, palmately lobed or angled, the lobes sharp-pointed; stipules sheathing. Flowers monoecious, both sterile and fertile ones in catkin-like heads, without calyx or corolla, but with small scales intermixed. Ovaries in the fertile flowers club-shaped, tipped with the thread-like, simple style, and downy at the base. Fertile heads solitary, on slender peduncles. The greenish-gray bark separates into thin plates.

ROSA'CEAE (ROSE FAMILY)

Herbs, shrubs, or trees with alternate, usually stipulate leaves and regular flowers. The petals (mostly 5) and stamens (mostly more than 10) inserted on the edge of a disk which lines the calyx tube.

There are no poisonous plants in this family. It includes many species, both wild and cultivated, that bear fleshy, edible fruits; among these are the Hawthorn, Choke-cherry, June-berry, Apple, Pear, Plum, etc.

Synopsis of the Genera

TRIBE I. PRU'NEAE

Carpel solitary, becoming a drupe, entirely free from the calyx, the latter deciduous. Ovules 2, but seed solitary as a rule. Trees or shrubs with simple leaves and deciduous stipules.

1. **Pru'nus**. Flowers perfect. Petals and calyx lobes 5. Fruit a drupe.

TRIBE II. RO'SEAE

Carpels few or many, free from the persistent calyx, becoming achenes, follicles, or drupe-like in fruit.

2. **Physocar'pus**. Carpels 1-5, inflated, ovules 2-4; stamens 30-40.
3. **Spirae'a**. Carpels mostly 5, forming follicles in fruit. Calyx 5-cleft, short. Petals obovate, similar.
4. **Gille'nia**. Carpels and fruit as in Spiraea. Calyx elongated, 5-toothed. Petals slender, dissimilar.
5. **Agrimo'nia**. Carpels 2, forming achenes inclosed in the hardened calyx tube. Calyx armed with hooked bristles. Flowers yellow, in slender spikes.
6. **Alchemil'la**. Carpels 1-4, forming achenes inclosed in the persistent calyx tube. Petals none. Stamens 1-4. Calyx tube inversely conical, the limb 4-parted, with 4 alternating bractlets. Low herbs, with palmately lobed leaves.
7. **Sanguisor'ba**. Carpels 1-3. Achene (mostly solitary) inclosed in the dry, 4-angled, closed calyx tube. Petals none. Lobes of the top-shaped calyx 4, petal-like, spreading. Stigma tufted. Tall herbs, with pinnate leaves and a dense white spike of small flowers, often polygamous or dioecious.
8. **Ge'um**. Carpels numerous, 1-ovuled, becoming dry achenes, the persistent styles becoming tails, plumose or naked, and straight or jointed. Calyx lobes with 5 alternating bractlets.
9. **Waldstei'nia**. Carpels 2-6, forming achenes. Leaves radical, of 3 wedge-form leaflets. Bractlets of the calyx minute and deciduous. Flowers yellow, on bracted scapes.
10. **Potentil'la**. Carpels numerous, forming achenes heaped on a dry receptacle, the style not forming tails. Lobes of the calyx with 5 alternating bractlets.
11. **Chamaer'hodos**. Carpels 5-10, on a dry receptacle. Petals white, obovate. Stamens 5, opposite the petals. Calyx campanulate, 5-cleft. Small, glandular-pubescent herbs with flowers in forked cymes, and many-cleft leaves.

12. *Fraga'ria*. Flowers as in *Potentilla*, but receptacle becoming fleshy or pulpy and scarlet in fruit. Leaves all radical, of 3 leaflets. Low plants, producing runners.
13. *Dalibar'da*. Carpels 5-10, each 2-ovuled, forming nearly dry drupelets. Calyx 5-6-parted, 3 of the divisions larger than the others, and toothed. Calyx without bracts, persistent, inclosing the fruit. Leaves radical, round-heart-shaped. Flowers white, on scapes.
14. *Ru'bus*. Carpels numerous, 2-ovuled, forming drupelets heaped on the receptacle. Fruit edible. Calyx without bracts.
15. *Ro'sa*. Carpels numerous, 1-ovuled, forming achenes inclosed in the fleshy calyx tube.

TRIBE III. PO'MEAE

Carpels inclosed in and adnate to the fleshy calyx tube, forming a pome in fruit. Styles often united below. Trees or shrubs.

16. *Cratae'gus*. Calyx tube urn-shaped, becoming thick and fleshy in fruit, inclosing and combined with the 1-5 carpels. Fruit a pome, but drupe-like, containing 1-5 bony nutlets. Thorny shrubs. Flowers generally white.
17. *Py'rus*. Fruit a pome or berry-like, the 2-5 carpels or cells of a papery or cartilaginous texture, each 2-seeded. Shrubs or trees.
18. *Amelan'chier*. Pome berry-like, 10-celled, i.e., with twice as many cells as styles. Petals narrow. Otherwise as in *Pyrus*. Shrubs or small trees, not thorny.

1. PRU'NUS L. PLUM. CHERRY

1. *P. ni'gra* Ait. (WILD PLUM.) A thorny tree, 8-10 feet high, with orange or red drupes $\frac{1}{2}$ inch or more in diameter; and ovate, conspicuously pointed, coarsely or doubly serrate, veiny leaves. Flowers white, appearing before the leaves, in umbel-like, lateral clusters.—Woods and river-banks.

2. *P. pu'mila* L. (DWARF CHERRY. SAND CHERRY.) A small trailing shrub, 6-18 inches high. Leaves obovate-lanceolate, tapering to the base, toothed near the apex, pale beneath. Flowers in umbels of 2-4, appearing with the leaves. Fruit ovoid, dark red, as large as a good-sized pea.—Crevices of rocks, and sand-beaches and plains.

3. *P. pennsylvan'ica* L.f. (WILD RED CHERRY. PIN CHERRY.) A tree 20-30 feet high, or shrubby. Leaves oblong-lanceolate, sharply serrate, green both sides. Flowers (appearing with the

leaves) in large clusters, the pedicels elongated. Fruit globular, as large as a red currant, very sour.—Rocky thickets, and in old windfalls.

4. *P. virginia'na* L. (CHOKE-CHERRY.) A good-sized shrub or small tree, 4–10 feet high. Leaves oval, oblong, or obovate, finely and sharply serrate, abruptly pointed. Flowers in short erect racemes, appearing after the leaves. Fruit red, becoming darker, very astringent.—Woods and thickets.

5. *P. serot'ina* Ehrhart. (WILD BLACK CHERRY.) A large tree with reddish-brown branches. Leaves smooth, varying from oval to ovate-lanceolate, taper-pointed, serrate, with short and blunt, incurved teeth, shining above. Flowers in long racemes. Fruit purplish-black, edible.—Woods and thickets.

2. *PHYSOCAR'PUS* Maxim. NINE-BARK

1. *P. opulifo'lius* Maxim. (NINE-BARK.) Shrub 3–7 feet high, the old bark separating in thin layers. Leaves broadly ovate or cordate, 3-lobed, doubly crenate, smooth. Flowers white, in umbel-like corymbs terminating the branches. Follicles 2–5, inflated, purplish.—River-banks.

3. *SPIRAE'A* L. MEADOW-SWEET

1. *S. salicifo'lia* L. (COMMON MEADOW-SWEET.) Shrub 2–3 feet high, nearly smooth. Leaves wedge-lanceolate, doubly serrate. Flowers white or rose-coloured, in a dense terminal panicle.—Low grounds along streams.

2. *S. tomento'sa* L. (DOWNY MEADOW-SWEET.) Deep rose-coloured flowers, and stems and under-surface of leaves densely woolly. Occurs eastward towards the sea-coast, and in the northern counties of Ontario.

4. *GILLE'NIA* Moench. INDIAN PHYSIC

1. *G. trifolia'ta* Moench. (BOWMAN'S ROOT.) Herb with 3-foliolate leaves; the leaflets ovate-oblong, pointed, rather coarsely serrate; stipules small, awl-shaped, entire. Flowers white or rose-coloured, in loose, few-flowered corymbs.—Rich woods.

5. *AGRIMO'NIA* L. AGRIMONY

1. *A. gryposep'ala* Wallr. (COMMON AGRIMONY.) Stem herbaceous, hairy, 2–3 feet high. Leaves interruptedly pinnate, larger

leaflets 5-7, oblong-ovate, coarsely serrate. Petals yellow, twice as long as the calyx. Stamens 5-15.—Borders of woods.

2. *A. parviflora* Ait. (SMALL-FLOWERED AGRIMONY.) Crowded lanceolate leaflets, 11-19, with smaller ones intermixed, and small petals.

3. *A. striata* Michx. Stem erect, usually unbranched below the flower-cluster, softly pubescent. Leaflets usually narrowed to a point and somewhat ribbed, resinous-dotted on the lower surface. Calyx strongly deflexed in fruit.—Pr. Prov.

6. *ALCHEMILLA* L. LADY'S MANTLE

1. *A. vulgaris* L. Leaves large, several-lobed, serrate. Flowers small, yellowish-green, in corymbose terminal clusters.—Atl. sea-coast.

7. *SANGUISORBA* L. BURNET

1. *S. canadensis* L. Stamens 4, long-exserted, white. Stem 3-6 feet high. Leaflets numerous, heart-shaped, ovate or oblong-lanceolate, coarsely serrate.—Bogs and wet meadows, Atl. Prov.

8. *GEUM* L. AVENS

1. *G. canadense* Jacq. (WHITE AVENS.) Stem 2 feet high, slender, branching, smoothish or downy. Root-leaves pinnate, the cauline ones 3-divided, lobed, or only toothed. Petals white, as long as the calyx. Achenes bristly, tipped with the hooked lower joint of the style, the upper joint falling away. Receptacle of the fruit bristly.—Low rich woods and thickets.

2. *G. virginianum* L. Stem stout, bristly-hairy. Leaves nearly as in No. 1. Petals white, shorter than the calyx. Receptacle of the fruit nearly smooth.—Meadows and thickets; not common.

3. *G. macrophyllum* Willd. Bristly-hairy, stout. Root-leaves interruptedly pinnate, with a very large, round-heart-shaped, terminal leaflet. Stem-leaves with 2-4 minute lateral leaflets, the terminal 3-cleft, with wedge-form, rounded lobes. Petals yellow, longer than the calyx. Receptacle nearly naked.—Atl. Prov. chiefly.

4. *G. strictum* Ait. (YELLOW AVENS.) Stem 2-3 feet high, rather hairy. Root-leaves interruptedly pinnate; stem-leaves 3-5-foliate, leaflets obovate or ovate. Petals yellow, longer than calyx. Receptacle of the fruit downy. Achenes tipped with the hooked style.—Dry thickets.

5. *G. riva'le* L. (WATER OR PURPLE AVENS.) Petals purplish-yellow; calyx brown-purple. Flowers nodding, but the fruiting heads upright. The upper joint of the style feathery, persistent. Stem simple, 2 feet high. Root-leaves lyrate; stem-leaves few, 3-foliolate, lobed.—Bogs and wet places.

6. *G. triflo'rum* Pursh. Stem about a foot high, soft-hairy. Flowers 3 or more, on long peduncles, purple. Styles not jointed, feathery, at least 2 inches long in the fruit.—Dry hills and thickets. Not common.

9. WALDSTEI'NIA Willd. BARREN STRAWBERRY

1. *W. fragarioi'des* Tratt. A low plant, 4–6 inches high. Leaflets 3, broadly wedge-form, crenately toothed. Scapes several-flowered. Petals yellow, longer than the calyx.—Dry woods and hillsides.

10. POTENTIL'LA L. CINQUE-FOIL. FIVE-FINGER

A. Styles thickened and glandular towards the base. Inflorescence cymose.

(1) *Style arising from near the base of the carpel. Petals whitish or yellowish.*

1. *P. argu'ta* Pursh. Stem stout, 1–2 feet high, brownish-hairy. Leaves pinnate, of 7–11 oval, serrate leaflets, downy underneath. Flowers in dense cymose clusters. Petals yellowish or cream-coloured, deciduous. Stamens about 30. Plant clammy above.—Dry thickets.

(2) *Style arising from the top of the carpel. Petals yellow.*

2. *P. monspeliensis* L., var. *norveg'ica* Rydb. (NORWAY CINQUE-FOIL.) Stem erect, hairy, branching above. Leaves palmate, of 3 leaflets; leaflets obovate-oblong, coarsely serrate. Flowers in cymose clusters. Calyx large. Petals pale yellow, small, not longer than the sepals. Stamens about 15.—Fields and low grounds.

3. *P. riva'lis* Nutt., var. *millegra'na* Watson. More slender and branching than the last, softly villous. Leaves all of 3 serrate leaflets. Cymes loose. Calyx small. Petals yellow, minute. Stamens 10–20, rarely 5.—Pr. Prov.

4. *P. paradox'a* Nutt. A plant of spreading or decumbent habit, with pinnate leaves of 5–11 leaflets, loose leafy cymes, small petals, as long as the sepals, and achenes with an appendage at the base.

5. *P. pennsylvan'ica* L. Perennial; more or less white-woolly. Leaflets 5–9, white-woolly beneath, less so above, cut-pinnatifid, the segments linear. Stamens about 25.—Chiefly eastward and Pr. Prov.

Var. *strigo'sa* Pursh. Silky-woolly throughout, 6–12 inches high. Leaflets deeply pinnatifid, the margins of the narrow lobes revolute. Cyme short and close.—Pr. Prov.

B. *Styles filiform, not glandular at the base. Inflorescence cymose.*

(1) *Style arising from the top of the carpel.*

6. *P. recta* L. Perennial, tall, sparingly villous, glandular-puberulent. Leaves palmate, of 5–7 leaflets, the latter cut-pinnatifid. Flowers yellow, large, in a broad cyme.

7. *P. Nuttal'lii* Lehm. Stems several, about 2 feet high. Leaves palmate, leaflets 5–9, usually oblanceolate, very deeply toothed. Flowers yellow, numerous in compact cymes.—Pr. Prov.

8. *P. Hippia'na* Lehm. Densely white-tomentose throughout. Stem slender. Leaves pinnate, of 5–11 leaflets, diminishing regularly down the petiole. Leaflets sharply toothed at least at the apex. Stamens 20.—Pr. Prov.

Var. *pulcher'rima* Watson. Upper surfaces of the crowded leaflets green and pubescent.—Pr. Prov. (Macoun).

9. *P. effu'sa* Dougl. Tomentose throughout, with scattered villous hairs. Stems diffusely branched above, 4–12 inches high. Leaflets 5–11, interruptedly pinnate. Carpels 10.—Pr. Prov.

10. *P. Platten'sis* Nutt. Stems decumbent. Pubescence appressed silky-villous throughout. Leaflets 7–13, crowded, deeply cut-pinnatifid into linear segments. Flowers few. Carpels many.—Pr. Prov. (Macoun).

11. *P. grac'ilis* Dougl. Stems 2–3 feet high. Leaves palmate, of about 7 leaflets, the latter serrate, tomentose beneath, green above. Carpels many.—Pr. Prov. (Macoun.)

Var. *flabellifor'mis* Torr. and Gray. Very deeply pinnatifid leaflets.—Pr. Prov. (Macoun.)

Var. *rig'ida* Watson. Villous, but not tomentose, and usually tall and stout.—Pr. Prov.

12. *P. pilo'sa* Willd. Distinguished from the preceding species by its tall rigid stems and light yellow flowers.—Niagara Falls, not common. (Macoun.)

13. *P. argen'tea* L. (SILVERY CINQUE-FOIL.) Stem ascending, branched at summit, white-woolly. Leaves palmate, of 5 leaflets, the latter deeply serrate towards the apex, with revolute margins, and woolly beneath. Petals yellow, longer than the sepals.—Dry fields and roadsides.

14. *P. interme'dia* L. Stems stouter than the last, grayish-hairy. Leaves palmately compound; leaflets 3–5, mostly oblanceolate, dentate, grayish-hairy beneath. Flowers yellow, in loose leafy cymes, petals $\frac{1}{4}$ inch long.

15. *P. humifu'sa* Nutt. Stems decumbent, 2–4 inches long, slender. Densely white-tomentose. Leaflets 5, serrate at the rounded or truncate apex with 3 or 5 teeth.—Pr. Prov. (Macoun.)

(2) *Style arising from the side of the carpel.*

16. *P. frutico'sa* L. (SHRUBBY CINQUE-FOIL.) Stem erect, shrubby, 1–3 feet high, much-branched. Leaves pinnate, of 5–7 leaflets, closely crowded, entire, silky, especially beneath. Flowers numerous, large, yellow, terminating the branches.—Bogs.

17. *P. tridenta'ta* Ait. (THREE-TOOTHED CINQUE-FOIL.) Common eastward towards the sea-coast. Stem 4–6 inches high, somewhat woody. Leaves rigid, palmate, of 3 wedge-shaped leaflets, 3-toothed at the apex. Petals white.

18. *P. palus'tris* Scop. (MARSH FIVE-FINGER.) Stem ascending. Leaves pinnate, of 5–7 oblong-lanceolate, crowded, deeply serrate leaflets, whitish beneath. Calyx an inch broad, dark purple inside. Petals purple.—Bogs.

C. Styles filiform, lateral. Peduncles axillary, solitary, 1-flowered.

19. *P. Anseri'na* L. (SILVER-WEED.) A low plant, creeping with slender runners. Leaves all radical, interruptedly pinnate; leaflets 9–19, serrate, green above, silvery-silky beneath. Flowers solitary, on long, scape-like peduncles, bright yellow.—River- and lake-margins.

20. *P. canadensis* L. (CANADA CINQUE-FOIL.) Stem prostrate or ascending, silky-hairy. Leaves palmate, of 5 leaflets, the latter serrate towards the apex. Flowers solitary. Petals yellow, larger than the sepals.—Dry soil.

11. CHAMAE/RHODOS Bunge.

1. *C. erecta* Bunge. Stem slender, erect, 2–12 inches high, branching above. Radical leaves many-cleft, forming a rosette; cauline ones 3–5-cleft.—Pr. Prov.

12. FRAGA/RIA L. STRAWBERRY

1. *F. virginiana* Duchesne. Achenes deeply imbedded in pits on the surface of the fleshy receptacle; calyx erect after flowering. Leaflets firm.

2. *F. vesca* L. Achenes not sunk in pits, but merely on the surface of the receptacle; calyx spreading or reflexed. Leaflets thin.

13. DALIBAR/DA Kalm. DALIBARDA

1. *D. repens* L. Stems tufted, downy. Whole plant with something of the aspect of a violet. Flowers of two kinds, white petaliferous and fertile apetalous.—Low woods.

14. RU/BUS L. BRAMBLE

1. *R. odoratus* L. (PURPLE FLOWERING-RASPBERRY.) Shrubby, 3–5 feet high. Branches, peduncles, and calyx clammy with glandular hairs. Flowers large and handsome, rose-purple. Leaves large, broadly ovate, 3–5-lobed, the lobes acute, minutely toothed. Fruit flat.

2. *R. parviflorus* Nutt. Distinguished from No. 1 by its oval white petals, and its 5-lobed, coarsely toothed leaves.—Pr. Prov.

3. *R. Chamaemorus* L. (CLOUD-BERRY. BAKED-APPLE BERRY.) A low herb with dioecious flowers. Stem with simple branches, without prickles, 2–3-leaved, bearing one large white flower. Leaves simple, kidney-form, 5-lobed, serrate.—Sphagnous swamps, chiefly eastward.

4. *R. triflorus* Richardson. (DWARF RASPBERRY.) Stems ascending or trailing, a foot high, not prickly. Leaflets 3–5, nearly smooth, rhombic-ovate, acute at both ends, doubly serrate.

Peduncle usually 3-flowered. Petals white or pink; sepals reflexed. Fruit red.—Cedar-swamps.

5. *R. idae'us* L. Stems upright, beset with stiff straight bristles. Leaflets 3–5, oblong-ovate, pointed, cut-serrate, whitish beneath. Sepals velvety. Fruit light red.—Hillsides and thickets.

Var. *aculeatis'simus* Regel and Tiling. (WILD RED RASPBERRY.) Calyx with bristly prickles.

6. *R. occidenta'lis* L. (BLACK RASPBERRY. BLACK CAP.) Stem glaucous, recurved, armed with hooked prickles. Leaflets 3, ovate, pointed, coarsely serrate, white-downy beneath. Fruit purplish-black.—Borders of fields, especially where the ground has been burned over.

7. *R. villo'sus* Ait. (HIGH BLACKBERRY.) Stem shrubby, furrowed, erect or reclining, armed with hooked prickles. Leaflets 3–5, unequally serrate, the terminal one conspicuously stalked. Lower surface of the leaflets hairy and glandular. Flowers racemed, numerous, large and white. Fruit oblong, black.

Var. *humifu'sus* T. and G. Trailing and smaller, and the flowers are less numerous.—Borders of thickets.

8. *R. alleghenien'sis* Porter. Old canes purplish, 3–6 feet tall; prickles stout, almost straight, found only on the old canes. Leaves ovate-lanceolate, acute or acuminate. Pedicels and peduncles glandular-pubescent. Fruit usually sub-cylindric.

9. *R. canaden'sis* L. (LOW BLACKBERRY.) Stem shrubby, extensively trailing, slightly prickly. Leaflets chiefly 3, oval or ovate-lanceolate, nearly smooth, sharply serrate. Flowers in racemes.—Thickets and rocky hills.

10. *R. his'pidus* L. (RUNNING SWAMP-BLACKBERRY.) Occurs occasionally in low meadows. Stem prostrate, with small reflexed prickles, sending up at intervals the short flowering shoots. Leaflets mostly 3, smooth and shining. Fruit of few grains, red or purple.

15. RO'SA L. ROSE

A. *Styles cohering in a protruding column, as long as the stamens.*

1. *R. setig'era* Michx. Stem climbing. Prickles nearly straight. Leaflets, 3–5, ovate. Petals deep rose-coloured, changing to white.—Borders of thickets and along fences.

B. Styles separate, included within the calyx tube; sepals spreading after flowering, and deciduous.

2. *R. caroli'na* L. (SWAMP ROSE.) Stem 1-7 feet high, erect, armed with stout, straight or somewhat curved prickles. Leaflets 5-9, very finely serrate, mostly narrowly oblong. Stipules long and very narrow. Flowers in corymbs, numerous. Calyx and globular calyx tube beset with glandular bristles.—Wet places.

3. *R. virginia'na* Mill. Stem 1-6 feet high, armed with stout, more or less hooked prickles. Leaflets 5-9, coarsely serrate, smooth and often shining above, rather thick. Peduncles 1-3-flowered.—Dry soil, or borders of swamps, chiefly eastward.

4. *R. hu'milis* Marsh. Low, more slender and less leafy than the last, with straight slender spines. Stipules narrow. Leaflets thin and pale. Outer sepals always more or less lobed.—Mostly in sandy soil.

5. *R. nit'ida* Willd. Low. Stem and branches usually thickly covered with prickles interspersed with straight slender spines. Stipules mostly dilated. Leaflets bright green and shining, mostly narrowly oblong. Flowers generally solitary. Sepals entire.—Margins of swamps, Atl. Prov.

6. *R. rubigino'sa* L. (SWEET BRIER. EGLANTINE.) Stem tall. Prickles numerous, the larger hooked, the smaller awl-shaped. Leaflets 5-7, doubly serrate, glandular beneath, aromatic. Flowers mostly solitary. Fruit pear-shaped or obovate.—Roadsides and fields.

C. Styles separate; sepals erect and connivent after flowering, persistent.

(1) *Fruit globose.*

7. *R. blanda* Ait. (EARLY WILD ROSE.) Stem 1-3 feet high. Prickles (if any) few and scattered, straight. Leaflets 5-7, mostly oblong-lanceolate, cuneate at the base, not resinous, simply serrate. Sepals hispid, not lobed. Peduncles 1-3-flowered.—Rocks and rocky shores.

8. *R. pratin'cola* Greene. Stems low, very prickly. Stipules narrow. Leaflets 7-11, broadly elliptical to oblong-lanceolate, wedge-shaped at the base, simply serrate, not resinous. Outer sepals lobed, rarely hispid.—Pr. Prov.

(2) *Fruit oblong-ovate to oblong.*

9. *R. acicula'ris* Lindl. Stems 1-4 feet high, often very prickly. Leaflets 5-7, elliptical to oblong-lanceolate, the teeth serrulate. Flowers large, solitary. Sepals not lobed. Fruit $\frac{1}{2}$ -1 inch long.—Shores of Lake Superior and westward.

16. *CRATAEGUS* L. HAWTHORN

1. *C. coccin'ea* L. Shrubs or small trees with occasional spines. Leaves elliptic-ovate, doubly-serrate or lobed, the teeth glandular-tipped; corymbs and calyx villous; stamens about 10. Fruit globose to ellipsoidal, dark reddish-brown, pubescent.

2. *C. macracan'tha* Lodd. Spines numerous, curved, 1-4 inches long. Leaves ovate to obovate, doubly serrate, with fine teeth, cuneate at the base. Flowers about 1 inch wide, in many-flowered corymbs; calyx villous; stamens about 10. Fruit sub-globose, dark cherry-red.

3. *C. mol'lis* Scheele. Round-topped trees with spines 1-2 inches long. Leaves broadly ovate, usually slightly cordate at the base, acutely lobed. Corymbs of flowers densely hairy; stamens about 20. Fruit sub-globose, scarlet.

4. *C. puncta'ta* Jacq. Flat-topped trees with straight spines 1-3 inches long. Leaves obovate to oblong with strongly hollowed veins above. Flowers about 1 inch broad; calyx lobes pubescent; stamens about 20. Fruit ellipsoidal, yellow or red.

5. *C. Crus-gal'li* L. A shrub or low tree with strong, straight, chestnut-brown spines 1-6 inches long. Leaves obovate to elliptical, sharply serrate, except near the base. Corymbs of flowers glabrous; stamens about 10. Fruit ovoid to globose, about $\frac{1}{2}$ inch wide, greenish to dull-red.

6. *C. Dougla'sii* Lindl. Trees or shrubs with reddish twigs and spines less than 1 inch long. Leaves ovate to obovate, doubly serrate and lobed, except near the base. Corymbs glabrous; stamens 10-20. Fruit ellipsoidal, black and sweet.—Pr. Prov.

17. *PYRUS* L. PEAR. APPLE

1. *P. corona'ria* L. (AMERICAN CRAB-APPLE.) A small tree, with ovate, serrate, simple leaves, tomentose beneath. Flowers in umbel-like cymes. Styles woolly and cohering at the base. Fruit a greenish apple.—Toronto and westward.

2. *P. arbutifo'lia* L. (CHOKE-BERRY.) A shrub, with oblong or oblanceolate, finely serrate, simple leaves, tomentose beneath. Flowers in compound cymes. Fruit berry-like, nearly globular, red.—Swamps.

3. *P. melanocar'pa* Willd. Nearly smooth throughout, with large black fruit.—Swamps.

4. *P. america'na* DC. (AMERICAN MOUNTAIN ASH.) A small tree with odd-pinnate leaves of 13–15 leaflets, the latter lanceolate, taper-pointed, sharply serrate, bright green. Fruit scarlet, berry-like, not larger than peas. Flowers in flat cymes.—Swamps and cool woods.

5. *P. sitchen'sis* Piper. Differs from the last in having oblong, oval, or lance-ovate, obtuse leaflets, smaller cymes, and larger flowers and berries.—Pr. Prov.

18. AMELAN'CHIER Medic. JUNE-BERRY. SASKATOON-BERRY

1. *A. canaden'sis* Medic. (SHADBUSH. SERVICE-BERRY.) A tree 10–30 feet high, with a purplish, berry-like, edible fruit. Leaves ovate to ovate-oblong, slightly cordate, pointed, very sharply serrate, nearly or soon glabrous. Bracts and stipules silky-ciliate. Flowers large, in drooping, nearly glabrous racemes. Petals oblong. Pedicels of the berries long.—Open woodlands.

Var. *rotundifolia* Torr. and Gray. Broader leaves.

2. *A. oblongifo'lia* Roem. A small tree (6–10 feet) with the young leaves and racemes densely white-tomentose. Leaves broader and generally rounded at the base. Flowers in shorter and denser racemes, and pedicels of the fruit shorter. Petals oblong-spatulate.—Low grounds.

3. *A. oligocarp'a* Roem. A shrub 2–4 feet high. Leaves thin, oblong, acute at both ends. Petals oblong-obovate. Fruit dark purple with a dense bloom.—Cold swamps.

4. *A. alnifo'lia* Nutt. A shrub 3–8 feet high. Leaves mostly broadly elliptical and obtuse, coarsely toothed towards the apex. Raceme short and rather dense. Petals wedge-shaped-oblong.—Pr. Prov.

LEGUMINO'SAE (PULSE FAMILY)

Herbs, shrubs, or trees, mostly with compound, alternate, stipulate leaves, and papilionaceous or sometimes regular corollas. Stamens usually 10 (rarely 5), monadelphous, diadelphous, or distinct. Fruit a legume.

Synopsis of the Genera

A. *Flowers papilionaceous. Stamens 10 (rarely 5), monadelphous or diadelphous.*

(1) *Not climbing or twining. Pod not jointed.*

1. **Lupi'nus.** Leaves palmately compound, leaflets 7-9. Flowers in terminal racemes. Stamens monadelphous.
2. **Trifo'lium.** Leaves of 3 leaflets. Flowers in heads. Stamens diadelphous.
3. **Medica'go.** Leaves pinnate, of 3 leaflets. Flowers in axillary spikes or racemes. Pod curved or coiled. Stamens diadelphous.
4. **Melilo'tus.** Leaves pinnate, of 3 leaflets, the leaflets toothed. Flowers in slender axillary racemes. Pod ovoid, wrinkled, 1-2-seeded. Stamens diadelphous.
5. **Hosack'ia.** Leaves pinnate, of 1-3 leaflets, stipules gland-like, minute. Flowers yellow, often turning reddish, solitary on axillary peduncles bearing each a leafy bract. Pod linear, several-seeded, partitioned between the seeds.
6. **Robin'ia.** Trees. Leaves odd-pinnate, often with spines for stipules, and the leaflets with small stipules. Flowers in hanging axillary racemes. Pod margined on one edge. Stamens diadelphous.
7. **Astrag'alus.** Leaves odd-pinnate, leaflets numerous. Flowers in dense axillary spikes. Corolla long and narrow, the keel not tipped with a sharp point. Pod turgid, one or both sutures projecting into the cell, thus partially or wholly dividing the cavity. Stamens diadelphous.
8. **Oxyt'ropis.** Low perennials with very short tufted stems from a hard rootstock, covered with scaly stipules. Leaves pinnate, of many leaflets. Flowers nearly as in Astragalus, but the keel is tipped with a sharp appendage. Peduncles scape-like.
9. **Glycyrrhi'za.** Leaves odd-pinnate, glandular-viscid. Flowers whitish, in axillary spikes, like those of Astragalus. Root long, perennial, sweet. Pods few-seeded, clothed with rough glands or short hooked prickles.
10. **Tephro'sia.** Hoary perennials. Stem simple. Flowers yellowish-white marked with purple, in a terminal, dense, oblong raceme. Wings coherent with the keel. Pod linear, flat, several-seeded.
11. **Psora'lea.** Leaves palmate, of 3-5 leaflets. Stamens diadelphous (sometimes monadelphous). Flowers spiked or racemed, mostly bluish, sometimes white. Pod as long as the calyx, 1-seeded. Calyx, pods, etc., usually glandular-dotted.

12. **Amor'pha.** Leaves odd-pinnate, of many leaflets, glandular-dotted. Corolla of one petal only, the standard infolding the stamens and style. Flowers violet or purple, in dense terminal spikes. Stamens 10, monadelphous at the base only, distinct above.
13. **Petaloste'mum.** Leaves odd-pinnate, crowded, glandular-dotted. Flowers small, in dense terminal heads or spikes. Corolla hardly papilionaceous. Stamens only 5. Petals on thread-like claws, 4 of them on the tube of filaments, the fifth on the bottom of the calyx. Pod 1-2-seeded.

(2) *Climbing by tendrils at the ends of the leaves.*

14. **Vic'ia.** Leaves abruptly pinnate, the leafstalk prolonged into a tendril. Flowers axillary. Style filiform, hairy at the apex. Pod 2-several-seeded. Stamens diadelphous.
15. **Lath'yrsus.** Leaves as in *Vicia*. Style flattish, flattened above, and hairy down the side opposite the free stamen. Stamens diadelphous.

(3) *Twining herbs.*

16. **A'pios.** A twining herb. Leaves pinnate, of 5-7 leaflets. Keel of the flower slender and coiled inward. Flowers brown-purple, in dense racemes. Stamens diadelphous.
17. **Strophosty'les.** A twining herb. Leaves pinnate, of 3 leaflets. Keel long, strongly incurved. Standard recurved-spreading. Style bearded along the upper side. Pod terete. Seeds oblong, pubescent. Stamens diadelphous.
18. **Amphicar'pa.** A low and slender twiner, the stem clothed with brownish hairs. Leaves pinnate, of 3 leaflets. Flowers polygamous, those of the upper racemes perfect, those near the base fertile, with the corolla inconspicuous or none. Stamens diadelphous.

(4) *Pods transversely jointed, the joints reticulated.*

19. **Desmo'dium.** Leaves pinnate, of 3 leaflets. Calyx 2-lipped. Flowers purple or purplish, in axillary or terminal racemes. Pod flat, the lower margin deeply lobed, thus making the pod jointed, roughened with hooked hairs, causing the pods to adhere to the clothing, etc. Stamens diadelphous.
20. **Hedys'arum.** Leaves pinnate, of numerous leaflets. Calyx 5-cleft. Pod flat, several-jointed, the joints roundish, reticulated, not rough. Flowers purple to white, deflexed, in racemes. Stamens diadelphous.
21. **Lespede'za.** Leaves pinnate, of 3 leaflets. Calyx 5-cleft. Pod flat, oval or roundish, occasionally 2-jointed, but only 1-seeded. Flowers sometimes polygamous. Stamens diadelphous.

B. Flowers papilionaceous. Stamens 10, all separate.

22. *Baptis'ia*. Leaves palmate, of 3 leaflets. Flowers yellow. Stamens all separate. The keel-petals nearly separate. Racemes terminating the bushy branches. Pods inflated.
23. *Thermop'sis*. Leaves palmate, of 3 leaflets, and with broad, leaf-like stipules. Stamens all separate. Flowers yellow. Pod flat, linear, curved.

C. Flowers not papilionaceous or only slightly so; polygamous dioecious or perfect. Stamens 3-10, all separate. Trees.

24. *Gledit'sia*. Thorny trees, with abruptly once- or twice-pinnate leaves. Flowers greenish, inconspicuous, in small spikes. Two lower petals somewhat united. Stamens separate, 3-10.
25. *Gymnoc'ladus*. Not thorny. Large trees with doubly pinnate leaves, the leaflets vertical. Pod very long and broad. Stamens 10. Petals on the summit of the long calyx tube, whitish.
26. *Cer'cis*. Small trees with heart-shaped, simple, entire leaves. Flowers perfect. Calyx 5-toothed; corolla imperfectly papilionaceous. Pod oblong, containing many almost circular seeds.

1. LUPINUS L. LUPINE

1. *L. peren'nis* L. (WILD LUPINE.) Stem erect, somewhat hairy. Leaflets 7-9, oblanceolate. Flowers racemed, purplish-blue; calyx deeply 2-lipped. Pods very hairy.—Sandy soil.

2. *L. argen'teus* Pursh., var. *argophyl'lus* Watson. Petals blue or cream-coloured, and calyx decidedly spurred at the base. Plant hoary-pubescent.—Pr. Prov.

3. *L. King'ii* Watson. Seeds only 2 or 3. Plant low, villous with white soft hairs. Racemes short, few-flowered, on long slender peduncles.—Pr. Prov.

2. TRIFO'LIUM L. CLOVER. TREFOIL

1. *T. arven'se* L. (RABBIT-FOOT OR STONE CLOVER.) Stem erect, 4-12 inches high, branching. Heads ovoid, of whitish flowers, oblong, very silky and soft. Calyx teeth fringed with long silky hairs.—Dry fields.

2. *T. prat'en'se* L. (RED CLOVER.) Stems and leaves somewhat hairy, the latter marked with a pale spot on the upper side. Flowers purplish, in dense sessile heads.—Pastures.

3. *T. me'dium* L. (ZIGZAG CLOVER.) Like the last, but with smoother zigzag stems, and leaves without the pale spot. Heads mostly stalked.—Atl. Prov.

4. *T. re'pens* L. (WHITE CLOVER.) Smooth, creeping. Heads of white flowers rather loose.—Fields everywhere.

5. *T. hy'bridum* L. (ALSIKE CLOVER.) Resembles the last, but the flowers are rose-tinted, and the stems erect or ascending.

6. *T. reflex'um* L. (BUFFALO CLOVER.) Only in South-western Ontario, in the neighbourhood of the Detroit River. Heads large, on naked peduncles; standard rose-red, wings and keel whitish. Flowers reflexed when old.

7. *T. agra'rium* L. (YELLOW or HOP CLOVER.) Flowers yellow, reflexed when old. Leaflets obovate-oblong, all 3 from the same point. Stem 6–12 inches high.—Sandy fields.

8. *T. procum'bens* L. (LOW HOP CLOVER.) Flowers yellow, reflexed when old. Leaflets wedge-obovate, the lateral ones at a short distance from the terminal one. Stem smaller than in No. 7, spreading.—Sandy fields.

3. *MEDICA'GO* L. MEDICK

1. *M. lupuli'na* L. (BLACK MEDICK.) Stem procumbent, downy. Leaflets obovate, toothed at the apex. Flowers yellow. Pods kidney-shaped.—Waste places.

2. *M. sati'va* L. (LUCERNE or ALFALFA.) Upright, has leaflets much as in No. 1, purple flowers in a long raceme, and spirally twisted pods.—Cultivated fields.

3. *M. arab'ica* Huds. (SPOTTED MEDICK.) Spreading or procumbent. Leaflets obcordate, with a purple spot. Flowers yellow. Peduncles 3–5-flowered. Pods compactly spiral, furrowed on the thick edge, bristly.—Atl. sea-coast.

4. *M. his'pida* Gaertn. Pods loosely spiral, deeply reticulated, with a thin keeled edge; otherwise like the last.—Atl. sea-coast.

4. *MELILO'TUS* Hill. SWEET CLOVER

1. *M. officina'lis* Lam. (YELLOW MELILOT.) Stem erect, 2–4 feet high. Leaflets obovate-oblong. Flowers yellow. Pod drooping, 1–2-seeded.—Waste places.

2. *M. alba* Desr., (WHITE MELILOT.) Much like No. 1, but has white flowers.

5. HOSACKIA Douglas

1. *H. america'na* Piper. An annual about 1 foot high, more or less silky-villous, sometimes glabrous. Leaves nearly sessile; leaflets varying from ovate to lanceolate. Flowers yellow or reddish, in umbels. Calyx teeth linear, much longer than the tube.—Pr. Prov.

6. ROBINIA L. LOCUST TREE

1. *R. Pseu'do-Aca'cia* L. (COMMON LOCUST.) Racemes slender, loose. Flowers white, fragrant. Branches glabrous. A large tree.

2. *R. visco'sa* Vent. (CLAMMY LOCUST.) Racemes crowded. Flowers white, with a reddish tinge. Branchlets and leaf-stalks clammy. Smaller than No. 1.

7. ASTRAG'ALUS L. MILK-VETCH

NOTE.—All the following species except Nos. 2, 7, 8, and 11 belong to the Prairie Provinces.

A. Pod completely 2-celled, being divided lengthwise by the intrusion of one or both sutures.

(1) *Pod thick and fleshy, sessile.*

1. *A. caryocar'pus* Ker. (GROUND PLUM.) A pale perennial, with low leafy stems, and minutely pubescent. Flowers violet, large, in short, spike-like racemes. Leaflets about 21, elliptical-oblong.

(2) *Pod coriaceous or cartilaginous, sessile.*

2. *A. canaden'sis* L. (CANADIAN MILK-VETCH.) Stem erect, 1-4 feet high, somewhat pubescent (or glabrous). Leaflets 21-27, linear-oblong, comparatively large. Flowers greenish-yellow, in long dense spikes. Pod and ovary glabrous.—Common from Quebec to the Rocky Mountains.

3. *A. adsur'gens* Pall. Stem 4-18 inches high, ascending or decumbent, grayish with appressed pubescence. Leaflets about 21, narrowly oblong. Flowers purplish, medium-sized, in dense short spikes. Pod finely pubescent, triangular-compressed, with a dorsal furrow.

4. *A. hypoglot'tis* L. Stem slender, 6-24 inches long, diffusely procumbent or ascending, sparingly pubescent. Leaflets 15-21. Flowers violet, in capitate clusters. Calyx copiously pubescent. Pod silky-villous, with a dorsal furrow.

B. *Pod* incompletely 2-celled.**(1) *Pod sessile*.**

5. *A. missourien'sis* Nutt. Stem very short, prostrate or matted, hoary-silky-pubescent. Leaflets 5–15 or more. Flowers large, violet or purplish, in few-flowered heads on scape-like peduncles. Pod obcompressed (flattened contrary to the usual way), nearly straight, pubescent.

(2) *Pod* with a very short stalk.

6. *A. lotiflorus* Hook. Stems very short, hoary with close pubescence. Leaflets 7–13. Flowers yellowish, in few-flowered heads. Pod oblong-ovate, acute at the base, hoary.—Rare.

7. *A. eucosmus* Robinson. Stems 12–18 inches high, grayish-puberulent. Leaflets 11–15, oval to linear-oblong. Flowers pale blue or purple, in a long, 1-sided raceme. Pod lens-shaped, gray-pubescent, grooved on the back.—Atl. Prov.

(3) *Pod* long-stalked.

8. *A. alpinus* L. Stems diffuse, 6–12 inches high, smooth or hairy-pubescent. Leaflets 13–25, oval or oblong. Racemes short, many-flowered, the flowers violet-purple. Pod narrowly oblong, black-pubescent, triangular-turgid, deeply grooved on the back, pendent.—Atl. Prov.

9. *A. racemosus* Pursh. Stem stout, 1–2 feet high. Leaflets 13–25. Flowers white and hanging. Pod straight, narrow, about 1 inch long, deeply grooved on the back.

C. *Pod* strictly 1-celled, coriaceous or cartilaginous.**(1) *Leaves* pinnate, of many leaflets.****(a) *Pod sessile*.**

10. *A. pectinatus* Dougl. Stem a foot high, grayish-pubescent, the branches striate and angled. Leaflets numerous, nearly filiform. Flowers large, white. Pod short, thick-cartilaginous, pendulous, cuspidate, smooth, with a very thick dorsal suture.

(b) *Pod* with a very short stalk.

11. *A. neglectus* Sheldon. Stem erect, nearly smooth, 1–2 feet high. Leaflets 11–21, comparatively large, minutely hoary

beneath. Flowers white, in short spikes. Pod inflated, smooth, slightly furrowed on both sides.—Ontario and Quebec.

12. *A. flexuosus* Dougl. Stem 1–2 feet high, ascending, grayish-pubescent. Leaflets numerous, narrow. Flowers white or purplish, small, in loose racemes. Pod very finely pubescent, not furrowed.

(c) *Pod exerted on its stalk.*

13. *A. bisulcatus* Gray. Stem stout, a foot or more high, with short rigid pubescence. Leaflets numerous. Flowers violet, in dense, spike-like racemes. Pod straight, narrowly oblong, the furrow of the ventral surface divided lengthwise into two by the projecting suture.

14. *A. tenellus* Pursh. Stem slender, somewhat glabrous. Leaflets 13–21, rather narrow. Flowers yellowish-white, tinged with purple, in loosely few-flowered racemes, the peduncles not exceeding the leaves. Stipules mostly connate. Pod 7–20-seeded, reflexed.

15. *A. aborigino-rum* Rich. Stems slender, rigid, clustered, hoary, pubescent or somewhat villous. Leaflets 7–13. Flowers white or bluish, the keel violet, in short, spike-like racemes. Pod with a straight dorsal edge (not furrowed), and a rounded ventral edge.

(2) *Leaves apparently palmate, mostly of 3 leaflets.*

16. *A. triphyllus* Pursh. A low, stemless, glossy-silky perennial with crowded, long-petioled leaves, mostly having 3 lanceolate leaflets, and whitish, crowded, sessile flowers. Pod conical-ovate, villous.

8. OXYTROPIS DC.

A. Leaves simply pinnate.

1. *O. campes'tris* DC., var. *johannen'sis* Fernald. Flowers white or yellowish, often tinged with purple or violet, in short spikes on naked scapes. Pods ovate or oblong, of a thin or papery texture. Leaves pinnate, of many leaflets.—Gravelly river-margins, Pr. Prov. and Atl. Prov.

B. Leaflets numerous, mostly in fascicles of 3 or 4 along the axis.

2. *O. Lambert'i* Pursh. Silvery-silky-pubescent. Flowers large, purple, violet, or white. Scapes often a foot high. Pods silky-

pubescent, cartilaginous, about $\frac{1}{2}$ inch long, imperfectly 2-celled.
—Pr. Prov.

3. *O. splen'dens* Dougl. Silvery-silky-villous, 6–12 inches high. Flowers erect-spreading. Pod erect, hardly surpassing the villous calyx.—Pr. Prov.

9. GLYCYRRHI'ZA L. LIQUORICE

1. *G. lepido'ta* Pursh. (WILD LIQUORICE.) Two to three feet high. Leaflets 15–19, sprinkled with little scales when young.
—Pr. Prov.

10. TEPHRO'SIA Pers. HOARY PEA

1. *T. virginia'na* Pers. (GOAT'S RUE. CATGUT.) Stem 1–2 feet high, simple, leafy to the top. Leaflets 17–29, linear-oblong. Roots long and slender and tough.—Sandy soil.

11. PSORA'LEA L.

1. *P. argophyl'la* Pursh. Silvery-silky all over. Leaves of 3–5 leaflets, the latter elliptical-lanceolate. Spikes of flowers interrupted. Roots not tuberous.—Pr. Prov.

2. *P. lanceola'ta* Pursh. Smooth or nearly so. Leaflets 3. Flowers small, in short spikes. Roots not tuberous.—Pr. Prov.

3. *P. esculen'ta* Pursh. Rough-hairy all over. Leaflets 5, obovate or lance-oblong. Spikes dense. Roots tuberous.—Pr. Prov.

12. AMOR'PHA L. FALSE INDIGO

1. *A. canes'cens* Pursh. (LEAD-PLANT.) White-downy all over, 1–3 feet high. Leaflets less than $\frac{1}{2}$ inch long, crowded. Spikes generally clustered. Pods 1-seeded.—Pr. Prov.

2. *A. microphyl'la* Pursh. Differs from the last in being lower and nearly smooth. Spikes usually solitary.—Pr. Prov.

3. *A. frutico'sa* L. A tall shrub. Leaflets larger and not so crowded as in the other species. Pods 2-seeded.—River-banks, Pr. Prov.

13. PETALOSTE'MUM Michx. PRAIRIE CLOVER

1. *P. purpu'reum* Rydb. Leaflets 5, very narrow. Calyx silky-hoary. Corolla rose-purple.—Pr. Prov.

2. *P. can'didum* Michx. Leaflets 7–9, lanceolate or linear-oblong. Calyx smooth or nearly so. Corolla white.—Dry prairies, Pr. Prov.

14. *VICIA* L. VETCH. TARE

1. *V. sati'va* L. (COMMON VETCH OR TARE.) Stem simple, somewhat pubescent. Leaflets 10-14, varying from obovate-oblong to linear. Flowers purple, large, one or two together, sessile in the axils, or nearly so.—Cultivated fields and waste grounds.

2. *V. tetrasper'ma* Moench. Peduncles long and slender, 1-2-flowered. Flowers small. Calyx teeth unequal. Corolla bluish. Pod 4-seeded, smooth. Leaflets about 4 pairs.—Atl. Prov.

3. *V. Crac'ca* L. (TUFTED VETCH.) Downy-pubescent. Leaflets 20-24, oblong-lanceolate, strongly mucronate. Peduncles long, bearing a dense, 1-sided raceme of blue flowers, bent downward in the spike, and turning purple before withering.—Borders of thickets, and pastures.

4. *V. carolinia'na* Walt. Smooth. Leaflets 8-12, oblong. Peduncles bearing a rather loose raceme of whitish flowers, the keel tipped with blue.—Low grounds and river-banks.

5. *V. america'na* Muhl. Smooth. Leaflets 10-14, oval or ovate-oblong, very veiny. Peduncles 4-8-flowered, flowers purple.—Moist places.

Var. *angustifo'lia* Nees. A low form, with linear leaflets.—Common in Pr. Prov.

6. *V. hirsu'ta* S. F. Gray. Stem weak. Leaflets 12-16, linear. Peduncles 3-6-flowered. Pods hairy, 2-seeded.

15. *LATH'YRUS* L. EVERLASTING PEA

1. *L. marit'imus* Bigel. (BEACH PEA.) Stem stout, about a foot high. Leaflets 8-16, oval or obovate. Stipules broadly halberd-shaped, about as large as the leaflets. Flowers large, purple.—Sea-coast, and shores of the Great Lakes.

2. *L. veno'sus* Muhl. (VEINY EVERLASTING PEA.) Stem 2-3 feet high. Leaflets 10-14. Stipules very small, slender, half-arrow-shaped. Flowers numerous.—Shady banks, Pr. Prov.

3. *L. ochroleu'cus* Hook. (PALE EVERLASTING PEA.) Stem slender. Leaflets 6-8, smooth and glaucous. Stipules half-heart-shaped, large. Corolla yellowish-white.

4. *L. palus'tris* L. (MARSH EVERLASTING PEA.) Stem slender, wing-margined. Leaflets 4-8, lanceolate, linear, or narrowly oblong,

sharply mucronate. Stipules small, half-arrow-shaped. Corolla blue-purple.—Moist places.

16. A'PIOS Ludwig. GROUND-NUT. WILD BEAN

1. *A. tubero'sa* Moench. Flowers brown-purple. Rootstocks with tuber-like enlargements.—A common twining plant in low grounds.

17. STROPHOSTYLES Ell.

1. *S. hel'vola* Britton. Annual. Stems 1–6 feet long. Leaflets ovate to oblong-ovate, often with a rounded lobe at the base. Flowers greenish-white and purplish, few, sessile, clustered on the long peduncles.

18. AMPHICAR'PA Ell. HOG PEA-NUT

1. *A. monoic'a* Ell. Leaflets 1–2 inches long. Flowers white or purplish, in racemes; calyx about $\frac{1}{6}$ inch long.—Moist thickets and river-banks.

2. *A. Pitche'ri* T. and G. Differs from No. 1 in having leaflets $\frac{1}{2}$ –1 inch in length, and calyx one-half as long again ($\frac{1}{4}$ inch).

19. DESMO'DIUM Desv. TICK-TREFOIL

A. Pod raised on a stalk much surpassing the calyx, the latter slightly toothed. Stipules bristle-form.

1. *D. nudiflo'rum* DC. Stem smooth, 4–8 inches high. Leaves crowded at the summit of sterile stems, leaflets bluntish. Flowers in a terminal raceme or panicle, on a scape which rises from the root. Leaflets broadly ovate.

2. *D. grandiflo'rum* DC. Stem pubescent. Leaves all crowded at the summit of the stem, from which the raceme or panicle arises. Leaflets conspicuously pointed.—Rich woods.

3. *D. pauciflo'rum* DC. Leaves scattered along the low ascending stems; leaflets rhombic-ovate, rather blunt. Racemes few-flowered, terminal.—Rich woods.

B. Pod raised on a stalk hardly surpassing the calyx, the latter deeply cleft. Stipules ovate, taper-pointed.

4. *D. rotundifolium* DC. Stem prostrate, soft-hairy. Leaflets orbicular. Flowers purple. Pods indented on both edges.—Dry sandy woods.

C. Pod hardly, if at all, stalked.

5. *D. canes'cens* DC. Stem covered with hairs some of which are hooked, others spreading and glutinous. Leaflets ovate, white and veiny beneath. Pod composed of about 5 somewhat triangular joints.

6. *D. bracteo'sum* DC. Stem tall, erect, very smooth. Leaflets ovate-lanceolate, taper-pointed, very large, green on both sides. Flowers and bracts large. Pod 4-6-jointed.—Thickets.

7. *D. illinoen'se* Gray. Has been found in the Thames valley. Leaflets obtuse and roughish, grayish beneath, with prominent veins. Pod about an inch long, 3-5-jointed, indented on both margins. Otherwise nearly like the last.

8. *D. panicula'tum* DC. Stem slender, nearly smooth. Leaflets oblong-lanceolate, tapering to a blunt point. Flowers medium-sized. Pod 3-5-jointed, the joints triangular. Racemes paniced.—Rich woods.

9. *D. Dille'nii* Darlengt. Distinguished from the last by the pubescent stem and finely pubescent leaflets, the latter oblong or oblong-ovate.—Dry and open thickets.

10. *D. canaden'se* DC. Stem erect, hairy, tall, furrowed. Leaflets oblong-lanceolate ($1\frac{1}{2}$ -3 inches long), with many straightish veins. Flowers large, about $\frac{1}{2}$ inch long, in dense racemes. Joints of the pod roundish.—Dry woods.

11. *D. sessilifo'lium* T. and G. Stem pubescent, 1-4 feet high. Leaves with very short petioles, the leaflets linear to linear-oblong. Flowers small, in dense racemes. Pods few-jointed.

12. *D. obtu'sum* DC. Stem ascending, slender, hairy. Leaflets round-ovate ($\frac{1}{2}$ -1 inch long). Flowers small, in loose racemes.—Dry thickets.

20. HEDYS'ARUM L.

1. *H. borea'le* Nutt. Leaflets 13-21, oblong or lanceolate. Stipules scaly, united opposite the petiole. Calyx 5-cleft, the awl-shaped lobes nearly equal. Pod of 3-4 flattened, roundish joints, easily separated.—Rocky banks, Atl. Prov. and Pr. Prov.

21. *LESPEDA'ZA* Michx. BUSH-CLOVER

A. *Flowers of two sorts; the larger perfect, the smaller pistillate and usually apetalous, mingled with the others.*

1. *L. viola'cea* Pers. Stems upright, branched. Leaflets varying from oblong to linear, downy underneath. Flowers violet-purple.—Dry borders of woods.

2. *L. virgin'ica* Britton. Stems erect, 1–4 feet high. Leaves crowded, usually long-petioled, leaflets linear or linear-oblong. Flowers violet-purple, short-peduncled, and crowded.

3. *L. frutes'cens* Britton. Stems erect, only about $\frac{1}{2}$ the height of the preceding. Leaves long-petioled, leaflets oval-oblong. Flowers as in No. 2, but the peduncles are of various lengths.

B. *All the flowers perfect, in close spikes or heads.*

4. *L. hir'ta* Hornem. Stem erect, wand-like, tall, pubescent. Leaflets roundish or oval, pubescent. Spikes dense, on peduncles longer than the leaves. Corolla yellowish-white, with a purple spot on the standard.

5. *L. capita'ta* Michx. Peduncles and petioles short. Leaflets varying from oblong to linear, silky underneath. Flowers in dense heads; corolla as in No. 1. Calyx much longer than the pod.—Both species are found in dry soil.

22. *BAPTIS'IA* Vent. FALSE INDIGO

1. *B. tincto'ria* R. Br. (WILD INDIGO.) Smooth and slender, 2–3 feet high, branching. Leaves nearly sessile. Leaflets wedge-obovate, turning black on drying. Flowers yellow.—Dry soil.

23. *THERMOP'SIS* R. Br.

1. *T. rhombifo'lia* Rich. A low perennial, with short, few-flowered racemes terminating the branches. Stipules as long as the petioles. Pods recurved, 10–14-seeded.—Pr. Prov.

24. *GLEDIT'SIA* L. HONEY-LOCUST

1. *G. triacan'thos* L. Thorns stout, often triple or compound. Pods linear, often more than a foot long, with pulp between the flat seeds.—Common in cultivation.

25. GYMNOCLADUS Lam. KENTUCKY COFFEE-TREE

1. *G. dioica* Koch. Leaves 2-3 feet long, without stipules; leaflets ovate. Pod 6-10 inches long and 2 inches broad.—Established in a few localities.

26. CER'CIS L. REDBUD. JUDAS TREE

1. *C. canadensis* L. (REDBUD.) Leaves pointed. Flowers red-purple, somewhat pea-like, appearing in umbellate clusters before the leaves.

LINA'CEAE (FLAX FAMILY)

Herbs with entire exstipulate leaves (but sometimes with glands in place of stipules), and regular hypogynous flowers. Sepals, petals, stamens, and styles, 5 each. Filaments united at the base. Pod 10-celled, 10-seeded. Our only genus is

1. LINUM L. FLAX

1. *L. virginianum* L. (VIRGINIA FLAX.) Flowers yellow, small ($\frac{1}{4}$ inch long), scattered. Stem erect, it and the spreading branches terete. Leaves lanceolate and acute, the lower obtuse and opposite. No glands, styles distinct.—Dry soil.

2. *L. striatum* Walt. Branches wing-angled, broader leaves and more crowded flowers than No. 1. The whole plant is stouter.—Shores of the Great Lakes.

3. *L. sulcatum* Riddell. Yellow flowers twice as large as the preceding, wing-angled branches, and a pair of dark glands in place of stipules. Sepals strongly 3-nerved, with rough bristly margins. Styles united to the middle.—Dry soil, rare.

4. *L. rigidum* Pursh. Glaucous. Flowers yellow, large. Sepals lanceolate, glandular-serrulate. Branches rigid, angled. Stipular glands usually present. Styles united.—Pr. Prov.

5. *L. Lewisii* Pursh. Flowers blue, on long peduncles. Perennial, glabrous and glaucous, leaves linear, acute. Styles distinct. Pod ovate.—Pr. Prov.

6. *L. usitatissimum* L. (COMMON FLAX.) Annual. Flowers blue, sepals ciliate. Leaves alternate, linear-lanceolate, acute, 3-veined.—Cultivated grounds.

OXALIDA'CEAE (WOOD-SORREL FAMILY)

Low herbs with an acid juice and alternate compound leaves, the 3 leaflets obcordate and drooping in the evening. Flowers very much the same in structure as in the following Family, but the fruit is a 5-celled pod. Styles 5, separate.

1. OX'ALIS L. WOOD-SORREL

1. *O. Acetose'l'a* L. (WHITE WOOD-SORREL.) Scape 1-flowered. Petals white, with reddish veins.—Cold woods.

2. *O. cornicu'l'a* L. (YELLOW WOOD-SORREL.) Annual, or perennial by running underground shoots, sparingly pubescent with spreading hairs. Peduncles 2-6-flowered, longer than the leaves. Petals yellow. Pod elongated, erect in fruit.

3. *O. stric'ta* L. Pubescent, with appressed hairs. Stem decumbent. Peduncles usually 2-flowered, pedicels at length deflexed.—Pr. Prov.

GERANIA'CEAE (GERANIUM FAMILY)

Herbs (often strong-scented) with symmetrical pentamerous flowers, the filaments usually united at the base, and glands on the receptacle alternate with the petals. Stigmas 5. Carpels 5, each 2-ovuled, 1-seeded, they and the lower part of the styles attached to an axis which rises from the receptacle. In the fruit the styles split away from the axis, carrying the carpels with them.

Synopsis of the Genera

1. *Gera'nium*. Stamens 10, rarely 5, all with anthers.
2. *Ero'dium*. Stamens with anthers, only 5.

1. GERA'NIUM L. CRANESBILL

1. *G. macula'tum* L. (WILD CRANESBILL.) Perennial. Stem erect, hairy, about a foot high. Leaves 5-7 parted, the wedge-shaped divisions lobed and cut. Flowers purple, 1 inch across. Petals entire, bearded on the claw, much longer than the long-pointed sepals.—Open woods and fields.

2. *G. inci'sum* Nutt. Perennial. Villous and glandular-pubescent. Leaves 5-7-cleft nearly to the base, the lobes cut into narrow divisions. Petals deep purple.—Pr. Prov.

3. *G. carolinia'num* L. (CAROLINA CRANESBILL.) Not perennial. Stems usually decumbent, hairy. Sepals awn-pointed, as long as the notched, rose-coloured or whitish petals.—Waste places.

4. *G. Robertia'num* L. (HERB ROBERT.) Not perennial. Stems reddish, spreading, pubescent; branches weak. Leaves 3-divided, or pedately 5-divided, the divisions twice pinnatifid. Sepals awned, much shorter than the reddish-purple petals. Plant with a very strong odour.—Shaded ravines and moist woods.

5. *G. pusil'lum* Burm, f. (SMALL-FLOWERED CRANESBILL.) Not perennial. Stem procumbent, slender, minutely pubescent. Leaves rounded, kidney-shaped, deeply 5-7-cleft, the divisions wedge-shaped. Sepals awnless, about the same length as the purplish petals. Stamens 5.—Waste places.

2. ERO'DIUM L'Her. STORK'S BILL

1. *E. cicuta'rium* L'Her. Stem low and spreading, hairy. Leaves pinnate, the leaflets sessile, 1-2-pinnatifid. Peduncles several-flowered. Styles, when they separate from the beak, bearded on the inside.—Not common.

RUTA'CEAE (RUE FAMILY)

Shrubs with compound, transparently dotted leaves, having an acrid taste. Flowers (with us) dioecious or polygamous, appearing before the leaves. Stamens hypogynous, as many as the petals.

Synopsis of the Genera

1. *Zanthox'ylum*. Flowers dioecious. Ovaries separate, forming pods. Leaflets more than 3.
2. *Pte'lea*. Flowers polygamous. Fruit a samara, winged all around. Leaflets 3.

1. ZANTHOX'YLUM L. PRICKLY ASH

1. *Z. america'num* Mill. (NORTHERN PRICKLY ASH. TOOTHACHE TREE.) A prickly shrub, with yellowish-green flowers in dense umbels in the axils. Sepals none. Petals 4-5. Stamens in the sterile flowers 4-5. Carpels 3-5, forming fleshy, 1-2-seeded pods. Fruit very pungent and aromatic. Leaves pinnate, 4-5 pairs, with an odd one at the end.—Forming thickets in low grounds along streams.

2. PTE'LEA L. SHRUBBY TREFOIL. HOP-TREE

1. *P. trifolia*'ta L. A tall shrub with greenish-white, small, polygamous flowers in compound terminal cymes. Sepals and petals 3-5. Stamens as many. Ovary 2-celled. Stigmas 2. Fruit a 2-celled and 2-seeded samara, winged all around. Leaves of 3 ovate, pointed leaflets.

POLYGALA'CEAE (MILKWORT FAMILY)

Herbs with entire exstipulate leaves and irregular hypogynous flowers, with petals in some cases almost if not quite separate. Stamens 6 or 8, monadelphous or diadelphous, the anthers 1-celled and opening at the top by a pore. Pod 2-celled and 2-seeded, flattened contrary to the partition.

1. POLYG'ALA L. MILKWORT

Sepals 5, the upper one and the two lower ones small and often greenish, the 2 lateral ones (called wings) larger and coloured like petals. Petals 3, connected with each other and with the tube of filaments, the lower one keel-shaped, and usually fringed or crested at the top. Style prolonged and curved.

1. *P. verticilla*'ta L. Flowers small, greenish-white, in slender spikes. Stems 4-8 inches high, much-branched. Stem-leaves linear, 4-5 in a whorl, the upper ones scattered.—Dry soil.

2. *P. Sen*'ega L. (SENECA SNAKEROOT.) Flowers greenish-white, in a solitary, cylindrical, close spike. Stems several, from a hard knotty rootstock, 6-12 inches high. Leaves lanceolate, with rough margins, alternate.—Dry hillsides and thickets.

3. *P. polyg*'ama Walt. Flowers rose-purple, showy, fringed, in a many-flowered raceme. Stems 5-8 inches high, tufted and very leafy, the leaves linear-oblong or oblanceolate. Whitish fertile flowers on underground runners.—Dry soil.

4. *P. paucifo*'lia Willd. (FRINGED POLYGALA.) Flowers rose-purple, rarely white, very showy, fringed, only 1-3 in number. Stems 1-4 inches high, from long underground runners, which also bear concealed fertile flowers. Upper leaves ovate, crowded at the top of the stem, lower leaves scale-like.—Dry woods.

5. *P. sanguin*'ea L. Flowers usually bright red-purple, but sometimes pale. Corolla inconspicuously crested. Flowers in

dense globular heads, at length oblong. True petals mostly shorter than the wings, the latter broadly ovate, closely sessile. Stem leafy to the top; leaves oblong-linear.—Sandy and moist places.

EUPHORBIA'CEAE (SPURGE FAMILY)

Plants with usually milky juice and monoecious flowers, represented in Canada chiefly by the two following genera:

Synopsis of the Genera

1. **Euphor'bia**. Calyx and corolla both absent, but with an involucre resembling a calyx or corolla surrounding numerous staminate and 1-pistillate flower. Stamen 1.
2. **Acaly'pha**. Calyx 4- or 3-5-parted; corolla absent, but without an involucre; stamens 8 or more.

1. EUPHOR'BIA L. SPURGE

Flowers monoecious, the sterile and fertile ones both destitute of calyx and corolla, but both contained in the same 4-5-lobed, cup-shaped involucre which resembles a calyx, and therefore the whole will probably at first sight be taken for a single flower. Sterile flowers numerous, each of a single naked stamen from the axil of a minute bract. Fertile flower only 1 in the middle of each involucre; ovary 3-lobed, soon protruded on a long pedicel; styles 3, each 2-cleft. Peduncles terminal, often umbellate. Plants with a milky juice.

A. Leaves all similar and opposite, short-petioled, green or blotched with brown above, furnished with awl-shaped or scale-like, often fringed stipules. Stems spreading or prostrate, much-forked. Involucres in terminal or lateral clusters, or one involucre in each fork, the involucre invariably with 4 (mostly petal-like) glands in the sinuses.

1. **E. polygonifo'lia** L. Leaves entire, oblong-linear, mucronate, very smooth. Stipules bristly-fringed. Peduncles in the forks. Glands of the involucre very small, not petal-like. Pods obtusely angled.—Shores of the Great Lakes, in sandy or gravelly places.

2. **E. serpens** HBK. Leaves entire, round-ovate, very small and smooth. Stipules membranaceous, triangular. Peduncles longer than the petioles, in loose clusters. Glands of the small

involucre with minute crenulate appendages. Stems thread-like, prostrate. Pods acutely angled. Seeds smooth.

3. *E. serpyllifo'lia* Pers. Leaves obovate-oblong, narrowed at the oblique base. Stipules lanceolate, fringed. Peduncles in loose lateral clusters. Stems prostrate-spreading. Seeds 4-angled, slightly cross-wrinkled, often pitted.—Sandy and alluvial soil, Pr. Prov.

4. *E. glyptosper'ma* Engel. Leaves serrulate towards the apex, linear-oblong, very unequal at the base. Stipules lanceolate, cut into bristles. Peduncles as long as the petioles, in dense lateral clusters. Glands in the small involucre with crenulate appendages. Stems erect-spreading. Pods sharply angled. Seeds sharply 4-angled, with 5 or 6 transverse wrinkles.—Gravelly soil.

5. *E. macula'ta* L. Leaves serrulate, oblong-linear, somewhat pubescent, with a brownish blotch in the centre, very oblique at the base. Peduncles in dense lateral clusters. Glands of the involucre with reddish, petal-like attachments. Pods sharply angled.—Roadsides.

6. *E. Pres'lii* Guss. Stem ascending. Leaves serrate, often with a red spot or with red margins, oblique at the base, ovate-oblong or oblong-linear. Peduncles in cymes at the ends of the branches. Glands of the involucre with white or occasionally reddish, petal-like attachments. Pod smooth, obtusely angled.—Cultivated soil.

7. *E. hirsu'ta* Wiegand. Inclined to be procumbent. Stems hirsute and branched. Leaves oblong to ovate. Seeds black with semi-transparent envelope, 4-angled; the concave or flat side obscurely wrinkled.—Dry sandy soil.

B. *Only the uppermost or floral leaves whorled or opposite. Stems erect. Stipules none. Involucres 5-lobed; inflorescence umbelliform, in the forks of the branches, and terminal.*

8. *E. corolla'ta* L. Stem erect, glabrous, 1-3 feet high. Conspicuous for the 5 bright-white, false lobes of the involucre, resembling petals; the true lobes very small.—Gravelly or sandy soil.

C. *Involucres chiefly in terminal umbels, and their glands always without petal-like attachments. Leaves without stipules or blotches, those of the stem alternate or scattered, the floral ones usually of a different shape, and whorled or opposite.*

9. *E. platyphyl'la* L. Umbel 5-rayed. Stem erect, 8–18 inches high. Upper stem-leaves lance-oblong, acute, serrulate, the uppermost heart-shaped, the floral ones triangular-ovate and cordate. Pod warty.—Shores of the Great Lakes.

10. *E. Heliosco'pia* L. Umbel first 5-rayed, then with 3, and finally merely forked. Stem ascending, 6–12 inches high. Leaves all obovate, rounded or notched at the apex, finely serrate. Pods smooth.—Along the Great Lakes.

11. *E. Cyparis'sias* L. Densely clustered stem, 6–12 inches high, crowded linear stem-leaves (the floral ones round-heart-shaped), and a many-rayed umbel.—Escaped from gardens in some localities.

12. *E. Peplus* L. 6–12 inches high. Umbel 3-rayed, then forking. Glands long-horned. Seeds ash-coloured, 2-grooved on the inner face, and pitted on the back.—Rather rare, waste places.

2. ACALY'PHA L. THREE-SEEDED MERCURY

1. *A. virgin'ica* L. Flowers monoecious, both kinds having a calyx, the staminate 4-parted, the pistillate 3–5-parted; no involucre. Staminate flowers very small, in spikes, with 1–3 pistillate flowers at the base in the axil of a large, leaf-like, 5–9-lobed bract. Stamens 8–16, monadelphous at the base, the anther-cells hanging from the apex of the filament. Styles 3, the stigmas cut-fringed, usually red. Pod separating into 3 globular carpels. A nettle-like weed, with ovate, sparsely serrate, alternate, long-petioled leaves.—Fields and open places.

CALLITRICHA'CEAE (WATER STARWORT FAMILY)

Small aquatic plants with entire leaves and without calyx and corolla. Fruit nut-like, flattened, 4-lobed and 4-celled.

1. CALLIT'RICHE L. WATER STARWORT

1. *C. palus'tris* L. Amphibious. Floating leaves obovate, tufted; submersed leaves linear. Flowers monoecious, axillary, usually between a pair of bracts. Sterile flower a single stamen; fertile flower a single pistil with a 4-celled ovary. Leaves beset with stellate scales.

2. *C. autumnalis* L. Growing under water. May be distinguished from No. 1 by its leaves being retuse and all linear from a broader base, and its flowers without bracts.

EMPETRA'CEAE (CROWBERRY FAMILY)

Low shrubby evergreens, resembling Heaths as to leaves and general aspect. Flowers dioecious or polygamous. Calyx somewhat petal-like or none. Corolla wanting. Ovary 3-9-celled, berry-like in fruit.

Synopsis of the Genera

1. *Em'petrum*. Flowers scattered and solitary in the axils. Sepals 3, somewhat petal-like.
2. *Core'ma*. Flowers in terminal heads. Calyx none.

1. EM'PETRUM L. CROWBERRY

1. *E. ni'grum* L. (BLACK CROWBERRY.) A slender procumbent shrub with the foliage and aspect of a Heath. Flowers polygamous, inconspicuous in the axils of the leaves. Calyx 3 petal-like sepals. Corolla wanting. Stamens 3. Fruit a black, berry-like drupe.—Atl. Prov. and Que.

2. CORE'MA Don. BROOM CROWBERRY

1. *C. Conrad'ii* Torr. (BROOM CROWBERRY.) Closely resembling the preceding. Flowers dioecious or polygamous, collected in terminal heads, each in the axil of a scaly bract, having no true calyx, but with 5 or 6 thin, dry bractlets under each. Stamens 3 or 4 with slender filaments. Drupes small, juiceless when ripe.—Atl. Prov.

LIMNANTHA'CEAE (FALSE MERMAID FAMILY)

Herbs with regular symmetrical flowers, the parts in threes or sixes. Leaves pinnate, exstipulate.

1. FLOERK'EA Willd. FALSE MERMAID

1. *F. proserpinacoi'des* Willd. A low tender annual with very small, solitary flowers on axillary peduncles. Leaves alternate, pinnate, of 3-5 lanceolate leaflets.—Marshes and river-banks.

ANACARDIA'CEAE (CASHEW FAMILY)

Trees or shrubs with a milky or resinous juice, and alternate leaves without dots or stipules. Sepals, petals, and stamens, each 5. Fruit a 1-seeded drupelet. The petals and stamens inserted under the edge of a disk which surrounds the base of the ovary.

1. RHUS L. SUMACH

1. *R. typhi'na* L. (STAGHORN SUMACH.) A small tree, 10-30 feet high, with densely soft-hairy branches and stalks. Flowers greenish-white, polygamous, forming a terminal thyrse. Fruit globular, covered with crimson hairs. Leaves pinnate, leaflets 11-31, oblong-lanceolate, serrate, pointed.—Dry hillsides.

2. *R. glabra* L. (SMOOTH SUMACH.) Smooth, glaucous, and seldom exceeding 5 feet in height.

3. *R. Toxicoden'dron* L. (POISON IVY. POISON OAK.) Shrub about a foot high, smooth, often climbing by rootlets. Leaves 3-foliolate, leaflets rhombic-ovate, notched irregularly, crenulate or entire. Flowers polygamous, in slender axillary panicles. Plant poisonous to the touch.

Var. *rad'icans* Torr. Climbs high.

4. *R. Ver'nix* L. (POISON ELDER.) A tall shrub, smooth or nearly so. Leaves odd-pinnate; leaflets 7-13, obovate-oblong, entire. Greenish-white flowers as in No. 3.—Swamps.

5. *R. canaden'sis* Marsh. (FRAGRANT SUMACH.) A shrub 2-3 feet high, or more, with 3-foliolate leaves, sweet-scented when crushed, and catkin-like spikes of pale-yellow flowers appearing before the leaves.—Dry rocky banks.

Var. *triloba'ta* Gray. Small leaflets and an unpleasant odour.—Rather common in Pr. Prov.

AQUIFOLIA'CEAE (HOLLY FAMILY)

Shrubs or small trees with small, axillary, polygamous or dioecious flowers, the parts mostly in fours or sixes. Calyx very minute, free from the ovary. Stamens alternate with the petals, attached to their base or free from them (in *Nemopanthus*), the corolla being almost polypetalous. Anthers opening lengthwise. Stigma nearly sessile. Fruit a berry-like, 4-8-seeded drupe.

1. ILEX L. HOLLY

1. *I. verticilla'ta* Gr. (BLACK ALDER. WINTERBERRY). A shrub with the greenish flowers in sessile clusters, or the fertile ones solitary. Parts of the flowers mostly in sixes. Fruit bright red. Leaves alternate, obovate, oval, or wedge-lanceolate, pointed, veiny, serrate.—Swamps and low grounds.

2. NEMOPAN'THUS Raf. MOUNTAIN HOLLY

1. *N. mucrona'ta* Trel. A branching shrub with gray bark and alternate, oblong, nearly entire, smooth leaves on slender petioles. Flowers on long, slender, axillary peduncles, mostly solitary. Petals 4-5, oblong-linear, distinct. Fruit light red.—Moist woods.

CELASTRA'CEAE (STAFF-TREE FAMILY)

Shrubs with simple stipulate leaves, alternate or opposite, and small regular flowers, the sepals and petals both imbricated in the bud. Stamens 4-5, alternate with the petals, and inserted on a disk which fills the bottom of the calyx. Pods orange or crimson when ripe.

Synopsis of the Genera

1. *Evon'y mus*. Flowers perfect. Sepals 4 or 5, united at the base, and forming a flat calyx. Branchlets 4-sided; leaves opposite. Flowers in axillary cymes. Stamens inserted on a disk which covers the ovary.
2. *Celas'trus*. Flowers polygamous. Petals and stamens 5. Calyx lined by a cup-shaped disk. Leaves alternate. Flowers in terminal racemes.

1. EVON'YMUS L. SPINDLE-TREE

1. *E. obova'tus* Nutt. (STRAWBERRY BUSH.) A low, rather straggling shrub, trailing, with rooting branches and short-petioled or sessile leaves, obovate, obtuse. Flowers greenish, with the parts generally in fives. Pods rough-warty, depressed, crimson when ripe.—Wooded river-banks and low grounds.

2. *E. atropurpu'reus* Jacq. (BURNING BUSH.) May be distinguished from No. 1 by its greater size (4-8 feet high), its long-petioled, oval, pointed leaves, purplish flowers with parts mostly in fours, and smooth pods.

2. CELASTRUS L. STAFF-TREE

1. *C. scandens* L. (WAX-WORK. CLIMBING BITTER-SWEET.)

A twining smooth shrub, with oblong-ovate, serrate, pointed leaves. Flowers small, greenish, in terminal racemes. Pods orange-coloured. These burst in autumn and display a scarlet pulpy aril, presenting a highly ornamental appearance.—Twining over bushes on river-banks and in thickets.

STAPHYLEACEAE (BLADDER NUT FAMILY)

Shrubs or small trees with perfect flowers and opposite pinnate leaves with deciduous stipules. Lobes of the whitish calyx, petals, and stamens each 5. Fruit a 2-3-horned, 2-3-celled, inflated capsule.

1. STAPHYLEA L. BLADDER NUT

1. *S. trifolia* L. (AMERICAN BLADDER NUT.) Shrub, 4-9 feet high. Leaflets 3, ovate, pointed. Flowers white, in drooping racemes, at the ends of the branchlets.—Thickets and hillsides.

ACERACEAE (MAPLE FAMILY)

Trees or shrubs with opposite, compound or lobed leaves, and usually unsymmetrical and often irregular flowers which are polygamous or dioecious. Sepals and petals mostly 5, both imbricated in the bud. Petals sometimes wanting. Stamens 3-12, inserted on a fleshy disk (sometimes wanting), which fills the bottom of the calyx tube. Ovary 2-celled, with 2 ovules in each cell. Fruit 2 separable, 1-seeded samaras or keys.

1. ACER L. MAPLE

1. *A. pennsylvanicum* L. (STRIPED MAPLE.) A small tree, 10-20 feet high, with light-green bark striped with dark lines. Leaves 3-lobed at the apex, finely and sharply doubly serrate, the lobes taper-pointed. Flowers greenish, in terminal racemes, appearing after the leaves. Samaras large, with divergent wings.—Rich woods.

2. *A. spicatum* Lam. (MOUNTAIN MAPLE.) A shrub or small tree, 4-8 feet high, growing in clumps in low grounds. Leaves 3-lobed, coarsely serrate, the lobes taper-pointed. Flowers greenish, appearing after the leaves, in dense upright racemes. Fruit with small, widely diverging or erect wings.

3. *A. sac'charum* Marsh. (SUGAR MAPLE.) A fine tree with 3-5-lobed leaves, breadth and length the same, dull above, a paler green underneath, the rather narrow sinuses rounded, and the lobes sparingly sinuate-toothed, the petioles without stipules. Flowers greenish-yellow, drooping on slender hairy pedicels, appearing at the same time as the leaves. Calyx fringed on the margin. Petals none.

Var. *nigrum* Torr. and Gray. Distinguished from the ordinary form by its paler and more pubescent leaves, the petioles with adnate foliaceous stipules.—Rich woods.

4. *A. sacchari'num* L. (WHITE OR SILVER MAPLE.) Leaves deeply 5-lobed, the sinuses rather acute, silvery-white underneath, the divisions narrow, sharply toothed. Flowers in erect clusters, greenish-yellow, appearing much before the leaves; petals none. Samara very large, woolly when young.—River-banks and low grounds.

5. *A. ru'brum* L. (RED MAPLE.) Leaves 3-5-lobed, sinuses acute. Flowers red, appearing much before the leaves. Petals linear-oblong. Samara small and smooth, on drooping pedicels. A smaller tree than No. 4, with reddish twigs, and leaves turning bright crimson in the autumn.—Swamps.

6. *A. Negun'do* L. (BOX ELDER.) Leaves pinnate, leaflets usually 3-7. Calyx minute, 4-5-cleft, disk absent. Petals none. Staminate flowers in rather dense clusters on slender pedicels, the pistillate ones in drooping racemes. Wings of the samaras incurved.—Pr. Prov. and naturalized in Ontario.

BALSAMINA'CEAE (BALSAM FAMILY)

Smooth herbs with succulent stems and simple exstipulate leaves. Flowers irregular, the sepals and petals coloured alike, one of the coloured sepals spurred. Stamens 5, coherent above. Pod bursting elastically, and discharging its seeds with considerable force.

1. IMPA'TIENS L. TOUCH-ME-NOT. JEWEL-WEED

1. *I. biflo'ra* Walt. (SPOTTED TOUCH-ME-NOT.) Flowers orange-coloured, spotted with reddish brown. Sac longer than broad, conical tapering into a long recurved spur.—Cedar swamps and along streams.

2. *I. pal'lida* Nutt. (PALE TOUCH-ME-NOT.) Flowers pale yellow, sparingly dotted with brown. Sac dilated, broader than long, ending in a short spur.—Wet places.

RHAMNA'CEAE (BUCKTHORN FAMILY)

Shrubs with simple stipulate leaves, and small, regular, perigynous, greenish or whitish flowers. Stamens opposite the petals, and with them inserted on the margin of a fleshy disk which lines the calyx tube. Fruit a berry-like drupe, or a pod.

Synopsis of the Genera

1. *Rham'nus*. Petals minute, or none. Drupe berry-like. Calyx and disk free from the ovary.
2. *Ceano'thus*. Petals white, long-clawed, hooded. Fruit dry, dehiscent. Calyx and disk adherent to the base of the ovary.

1. RHAM'NUS L. BUCKTHORN

1. *R. alnifo'lia*, L'Her. A low erect shrub, not thorny, with oval, acute, serrate leaves, and apetalous flowers. Fruit a 3-seeded berry.—Swamps.

2. CEANO'THUS L. NEW JERSEY TEA

1. *C. america'nus* L. A shrubby plant with downy branches and ovate, 3-ribbed, serrate leaves. Flowers in white clusters at the summit of the naked flower-branches. Sepals and petals white, the latter hooded, and with slender claws. Pedicels also white.—Dry hillsides.

2. *C. ova'tus* Desf. Leaves narrowly oval or elliptical-lanceolate, finely serrate, and glabrous or nearly so. The flowers, also, are larger than in No. 1.

VITA'CEAE (VINE FAMILY)

Shrubs climbing by tendrils, with small greenish flowers in paniced clusters, opposite the leaves. Stamens as many as the petals and opposite them. Calyx minute. Petals 4 or 5, hypogynous or perigynous, very deciduous. Fruit a berry, 1-4-seeded. Leaves palmately veined, or compound.

Synopsis of the Genera

1. *Vitis*. Leaves simple, heart-shaped, and variously lobed.
2. *Psed'era*. Leaves compound-digitate, of 5 serrate leaflets.

1. VITIS L. GRAPE

1. *V. aestiva*'lis Michx. (SUMMER OR PIGEON GRAPE.) Leaves and branches woolly. Berries rather large, black, with a bloom.—Moist thickets.

2. *V. bic'olor* Le Conte. (SUMMER GRAPE.) Resembles No. 1, but the branchlets are nearly or quite glabrous and the leaves glaucous, soon becoming glabrous beneath.

3. *V. cordifo'lia* Michx. (FROST GRAPE.) Leaves smooth or nearly so, bright green on both sides, not lobed or slightly 3-lobed, heart-shaped, sharply serrate. Stipules small. Berries small, black or shining.—Banks of streams.

4. *V. vulpi'na* L. (RIVER-BANK OR FROST GRAPE.) Differs from No. 3 in having larger stipules (about $\frac{1}{4}$ inch long). Leaves usually 3-lobed. Berries blue, with a bloom.

2. PSED'ERA Neck. VIRGINIA CREEPER

1. *P. quinquefo'lia* Greene. A common woody vine in low grounds. Leaves digitate, of 5 oblong-lanceolate leaflets. Tendrils with sucker-like disks at the end, by which they cling to walls, trunks of trees, etc. Fruit a small black berry.

TILIA'CEAE (LINDEN FAMILY)

Trees with fibrous bark, soft and white wood, and heart-shaped and serrate leaves, with deciduous stipules. Flowers in small cymes hanging on an axillary peduncle, to which is attached a leaf-like bract. Sepals deciduous. The only Canadian genus is

1. TILIA L. BASSWOOD. WHITEWOOD

1. *T. america'na* L. (BASSWOOD.) A fine tree, in rich woods. Flowers yellow or cream-coloured, very fragrant. Leaves smooth and green on both sides, obliquely cordate or truncate at the base, sharply serrate. Sepals 5. Petals 5. Fruit a globular nut, 1-celled, 1-2-seeded.

MALVA'CEAE (MALLOW FAMILY)

Herbs with palmately veined, alternate, stipulate leaves. Flowers regular. Calyx valvate. Corolla convolute in the bud. Sepals 5, united at the base. Petals 5. Stamens numerous, monadelphous, in a column, united to the base of the petals; anthers 1-celled. Carpels united in a ring, separating after ripening. Seeds kidney-shaped.

Synopsis of the Genera

1. **Malva.** Carpels without beaks, 1 seeded. A circle of 3 bractlets at the base of the calyx. Stigmas occupying the inner face of the style.
2. **Malvas'trum.** Carpels without beaks, 1-seeded. Involucel of 3 bractlets, or none. Stigmas terminal, capitate. Low hoary herbs.
3. **Abu'tilon.** Carpels 2-beaked, 2-9-seeded. No circle of bractlets.
4. **Hibis'cus.** Column of stamens naked and 5-toothed at the apex. Pod 5-celled, many-seeded. Involucel of many bractlets.

1. MALVA L. MALLOW

1. **M. rotundifo'lia** L. (ROUND-LEAVED MALLOW.) Stems several, procumbent, from a stout tap-root. Leaves long-petioled, round-heart-shaped, crenate, crenately lobed. Petals obcordate, whitish, streaked with purple, twice as long as the sepals.—Waysides and cultivated fields.

2. **M. sylves'tris** L. (HIGH MALLOW.) Stem erect, 2-3 feet high. Leaves sharply 5-7-lobed. Petals purple, 3 times as long as the sepals.—Near dwellings.

3. **M. cris'pa** L. (CURLED MALLOW.) A tall erect annual with round and angled, toothed and crisped leaves, and small sessile flowers crowded in the axils.—Escaped from old gardens.

4. **M. moscha'ta** L. (MUSK MALLOW.) Stem erect, 1 foot high. Stem-leaves 5-parted, the divisions cleft. Flowers large and handsome, rose-coloured or white, on short peduncles, crowded on the stem and branches in the upper axils.—Roadsides near gardens.

2. MALVAS'TRUM Gray. FALSE MALLOW

1. **M. coccin'eum** Gray (PRAIRIE MALLOW.) A low hoary perennial with rose-pink flowers in spikes or racemes. Leaves 5-parted or pedate. Petals much larger than the sepals.—Pr. Prov.

3. ABU'TILON Mill. INDIAN MALLOW

1. *A. Theophras'ti* Medic. (VELVET-LEAF.) Stem 2-4 feet high, branching. Leaves velvety, round-cordate, long-pointed. Corolla yellow.—Not common; near gardens.

4. HIBIS'CUS L. ROSE MALLOW

1. *H. Moscheu'tos* L. (SWAMP ROSE MALLOW.) A tall perennial with very large and showy, pink or white flowers, in late summer. Calyx not inflated. Leaves ovate, pointed, toothed; the lower 3-lobed, all white-downy beneath.—Marshes.

2. *H. Trio'num* L. (BLADDER KETMIA.) A low hairy annual. Calyx inflated in fruit, and 5-winged. Corolla sulphur-yellow, with dark centre.—Escaped from gardens.

HYPERICA'CEAE (ST. JOHN'S-WORT FAMILY)

Herbs or shrubs with opposite, entire, dotted leaves, and no stipules. Flowers regular, hypogynous, mostly yellow. Sepals 5, persistent. Petals 5, deciduous. Stamens mostly numerous, and usually in 3 or more clusters. Styles 3-5, sometimes united. Pod 1-5-celled. Seeds numerous.

1. HYPER'ICUM L. ST. JOHN'S-WORT

A. Pods 3-celled. Styles 3, separate. Petals usually with black dots.

1. *H. perfora'tum* L. (COMMON ST. JOHN'S-WORT.) Stem much-branched, producing runners at the base, slightly 2-edged. Leaves linear-oblong, with transparent dots, easily observed by holding the leaf up to the light. Petals deep yellow. Flowers in open leafy cymes.—Fields.

2. *H. puncta'tum* Lam. (CORYMBED ST. JOHN'S-WORT.) Stem rounded, not so branching as No. 1. Leaves with both black and transparent dots, oblong, somewhat clasping. Flowers small, pale yellow, crowded, corymbed.—Damp woods and wet places.

3. *H. prolif'icum* L. Leafy bushy plants with 2-edged branches and oblong, obtuse leaves. Flowers yellow; stamens more than 12.

4. *H. virgin'icum* L. (MARSH ST. JOHN'S-WORT.) Stem smooth. Leaves oblong or oval, clasping, often purple-veined, obtuse, conspicuously dotted beneath. Flowers flesh-coloured, in the axils, and at the summit of the stem. Stamens mostly 9 in 3

clusters, styles 3, pod 3-celled and purplish. The whole plant is of a purplish hue.—Marshes.

B. Pod 5-celled. Styles more or less united. Stamens very many, in 5 clusters, if clustered at all.

5. *H. As'cyron* L. (GREAT ST. JOHN'S-WORT.) Stem 3–5 feet high. Leaves 2–3 inches long, somewhat clasping. Flowers very large, the yellow petals about an inch long, and narrowly obovate. Stamens showy. Pod conical, large.—Along streams; not common.

6. *H. Kalmia'num* L. (KALM'S ST. JOHN'S-WORT.) Shrubby, a foot or more in height; leaves linear-lanceolate, crowded, revolute on the margins, thickly punctate, and sessile. Flowers 1 inch or more across, in clusters, petals yellow.—Niagara Falls and westward.

C. Pod 1-celled.

7. *H. ellip'ticum* Hook. (ELLIPTICAL-LEAVED ST. JOHN'S-WORT.) Stem about 1 foot high, not branched. Leaves spreading, elliptical-oblong, obtuse, thin. Flowers rather few, showy, in a nearly naked cyme. Pod purple, ovoid, obtuse. Petals yellow.—Banks of streams, eastward.

8. *H. mu'tilum* L. (SMALL ST. JOHN'S-WORT.) Stem slender, branching above, hardly a foot high. Leaves 5-nerved. Cymes leafy at the base. Flowers small, not $\frac{1}{4}$ inch across.—Low grounds.

9. *H. canaden'se* L. (CANADA ST. JOHN'S-WORT.) Stem upright, 6–15 inches high, with branches erect. Leaves linear or linear-lanceolate, 3-nerved at the base, the upper ones acute, sessile. Cymes naked. Pod much longer than the calyx. Flowers small, deep yellow.—Wet sandy places.

10. *H. gentianoi'des* BSP. With somewhat wiry, erect, close branches. Leaves awl-shaped, scale-like, appressed. Flowers minute, sessile, and continuous throughout the season.

CISTA'CEAE (ROCK ROSE FAMILY)

Herbs or low shrubs with simple entire leaves and regular polyandrous flowers. Calyx persistent, usually of 3 large and 2 smaller sepals. Petals 5 or 3, convolute in the bud. Stamens 3–20. Pod 1-celled, 3-valved. Seeds on 3 parietal projections.

Synopsis of the Genera

1. *Helian'themum*. Petals 5, fugacious, or none. Style none.
2. *Hudso'nia*. Petals 5, fugacious. Style long and slender.
3. *Le'chea*. Petals 3, persistent. Style none.

1. *HELIAN'THEMUM* Mill. ROCK ROSE

1. *H. canaden'se* Michx. (FROST-WEED.) Flowers of two sorts, some solitary, with large yellow corolla and many stamens, the petals lasting but one day after the flower opens; others small, clustered in the axils of the leaves, and apetalous or with small petals. Leaves lanceolate, downy beneath.—Sandy places.

2. *HUDSO'NIA* L. HUDSONIA

1. *H. tomento'sa* Nutt. (DOWNY HUDSONIA.) Hoary. Leaves oval or narrowly oblong, short, close-pressed, or imbricated. Flowers small, sessile, yellow, very numerous.—A little, heath-like shrub, on the shores of the Great Lakes and the River St. Lawrence.

2. *H. ericoi'des* L. Downy, but greenish. Leaves slender, awl-shaped. Flowers on slender naked stalks.—Dry soil, Atl. coast.

3. *LE'CHEA* L. PINWEED

1. *L. minor* L. (SMALLER PINWEED.) Flowers inconspicuous, purplish, loosely racemose, on distinct pedicels. Stem slender, rough with appressed scattered hair, producing radical shoots. Leaves scattered, linear. Pods the size of a pin's head.—Dry soil.

2. *L. villo'sa* Ell. Stem 1–2 feet high, stout, very leafy, villous with spreading hairs, producing prostrate branches at the base. Leaves elliptical, mucronate. Flowers densely crowded, on very short pedicels.—Dry soil.

VIOLA'CEAE (VIOLET FAMILY)

Herbs with alternate stipulate leaves. Flowers irregular, the lower of the 5 petals being spurred or gibbous. Sepals 5, persistent. Stamens 5, hypogynous, the anthers more or less coherent, and surrounding the pistil. Fruit a 1-celled pod, splitting into 3 valves. Seeds in 3 rows on the walls of the ovary.

Synopsis of the Genera

1. *Vi'ola*. Sepals auricled at the base. Lower petal usually distinctly spurred. Stamens only slightly, if at all, coherent; the 2 lower ones spurred. Besides ordinary flowers, apetalous, cleistogamous flowers on short stems are produced later.
2. *Hyban'thus*. Sepals not auricled. Lower petal saccate at the base. Stamens completely united and sheathing the ovary. Plant 1-2 feet high. Flowers of one kind only.

1. VI'OLA L. VIOLET

A. *Stemless Violets; leaves and scapes all from rootstocks.*

(1) *Flowers white.*

1. *V. blan'da* Willd. (SWEET WHITE VIOLET.) Lower petal streaked with purple. Leaves round-heart-shaped or reniform, usually acute, glabrous except for a few hairs on the upper surface. Petals beardless. Flower sweet-scented.—Swamps and wet meadows, in spring.

2. *V. renifo'lia* Gray. Leaves much larger and more pubescent than those of the preceding, and abruptly pointed. Petals beardless.—Dry cedar swamps, and ravines in rich woods.

3. *V. pal'lens* Brainerd. Leaves glabrous on both sides, round-heart-shaped, mostly obtuse. Lateral petals with a few hairs. Flowers sweet-scented.

4. *V. lanceola'ta* L. (LANCE-LEAVED VIOLET.) Flowers white. Petals beardless. Leaves lanceolate, erect, tapering into a long, margined petiole, nearly entire.—Often in bogs.

5. *V. primulifo'lia* L. (PRIMROSE-LEAVED VIOLET.) Flowers white, lateral petals usually somewhat bearded. Leaves ovate or oblong, heart-shaped, or abrupt at the base.—Damp or dry ground; Atl. Prov.

(2) *Flowers blue or purple.*

6. *V. peda'ta* L. (BIRD-FOOT VIOLET.) Nearly smooth. Rootstock short, thick and erect. Leaves all deeply cut into 3-5 mostly linear segments, the lateral divisions 2-3-parted. Flower about an inch across; stigma large and not beaked.—Pr. Prov.

7. *V. pedatif'ida* G. Don. Very much like the last, but the flowers are smaller and of a deeper blue, and the stigma is beaked.—Pr. Prov.

8. *V. Selkirk'ii* Pursh. (GREAT-SPURRED VIOLET.) A small and delicate plant with round-cordate leaves, slightly pubescent on the upper surface, distinguished by the slender rootstock, and the very large spur, thickened at the end. The pale-violet petals also are beardless.—Damp shady places.

9. *V. palus'tris* L. (MARSH VIOLET.) Very similar in foliage, etc., to No. 1, but the leaves are glabrous, the flowers are pale lilac, the spur is short, and the rootstock is jointed.—Wet swamps amongst moss.

10. *V. palma'ta* L. (COMMON BLUE VIOLET.) Leaves on very long petioles, cordate or reniform, sides folded inwards when young, the later ones variously 5-9-lobed or parted. Lateral petals bearded. Flowers blue or rarely white. Spur short and thick.—Low grounds everywhere.

11. *V. cuculla'ta* Ait. Leaves merely crenate, heart-shaped, glabrous or nearly so. Flowers violet-blue, rarely white, on long petioles; lateral petals bearded.—Wet places, common.

12. *V. odora'ta* L. (ENGLISH SWEET VIOLET.) Flowers very fragrant. Leaves cordate, pubescent.—Escaped from gardens in some places.

13. *V. sagitta'ta* Ait. Plant glabrous. Leaves lanceolate, long-petioled, the blades 1-3 inches long, with toothed lobes at the base, somewhat sagittate.—Moist fields.

14. *V. fimbriat'ula* Sm. Rootstock stout, mostly erect. Leaves ovate to ovate-oblong, acute, pubescent; petioles nearly equalling the blades or shorter. Flowers violet-purple.—Dry sandy hill-sides.

15. *V. aff'nis* Le Conte. Smooth. Leaves heart-shaped, tapering, with crenate-to-serrate margins. Petals violet with white base.

16. *V. soro'ria* Willd. Resembling No. 11, but the leaves and petioles are more or less pubescent. Flowers are violet, rarely white. Spurred petal nearly glabrous.—Moist meadows.

17. *V. septentriona'lis* Greene. Leaves and petioles hairy. Flowers violet to white and spurred petal villous.

(3) *Flowers yellow.*

18. *V. rotundifo'lia* Michx. (ROUND-LEAVED VIOLET.) Leaves round-ovate, cordate, repand-crenulate, about an inch wide at flowering, increasing later to 3 or 4 inches, and then flat on the ground, shining above. Lateral petals bearded and marked with brown lines. Spur very short.—Cold woods, chiefly eastward.

B. *Leafy-stemmed Violets.*(1) *Flowers yellow.*

19. *V. pubes'cens* Ait. (DOWNY YELLOW VIOLET.) Plant downy, 6–12 inches high. Leaves broadly cordate, coarsely serrate; stipules large, entire. Lower petals veined with purple. Spur very short.—Rich woods.

20. *V. scabrius'cula* Schwein. Smaller and less pubescent, often nearly smooth.

21. *V. Nuttal'lii* Pursh. Low, densely pubescent, or sometimes nearly glabrous. Leaves oblong-ovate or oblong, obtuse, entire or obscurely sinuate, decurrent on the petiole; stipules mostly narrow, entire.—Dry soil, Pr. Prov.

(2) *Flowers not yellow.*

22. *V. canaden'sis* L. (CANADA VIOLET.) Tall, often a foot high. Leaves large, cordate, serrate, pointed. Petals white inside, purplish outside, lower petal tinged with yellow and marked with dark lines. Spur very short.—Flowering all summer.

23. *V. arena'ria* D.C. Stem low, in tufts. Leaves thick, densely pubescent, ovate, slightly cordate, and with obtuse apex. Flowers blue, with short spur.—Sandy soil, Pr. Prov.

24. *V. adun'ca* Gray. Nearly or quite glabrous. Leaves round-ovate, rarely cordate. Flowers blue, spur long.—Dry soil, Pr. Prov.

25. *V. consper'sa* Reichenb. Rootstock branched. Leaves glabrous, round-cordate or reniform, crenate. Flowers pale violet, rarely white. Leafy stems and flowers numerous.—Common.

26. *V. stria'ta* Ait. (PALE VIOLET.) Stem angular, 6–10 inches high. Leaves cordate, finely serrate; stipules fringed-toothed. Spur thickish, much shorter than the cream-coloured or white petals.—Low grounds.

27. *V. rostra'ta* Pursh. (LONG-SPURRED VIOLET.) Distinguished at once by its extremely long, nearly straight spur. Stems numerous. Leaves round to ovate, heart-shaped, serrate. Petals violet-coloured.

28. *V. Rafines'quii* Greene. (PANSY.) Stipules large, leaf-like and lyrate-pinnatifid. Stem angled and branched. Leaves roundish. Petals variable in colour, about as long as the sepals.—Dry soil.

2. *HYBAN'THUS* Jacq. GREEN VIOLET

1. *H. con'color* Spreng. A homely herb with oblong entire leaves pointed at each end, and 1-3 small, greenish-white flowers in the axils, on short recurved pedicels. Pod an inch long.—Rare.

CACTA'CEAE (CACTUS FAMILY)

Very fleshy and commonly leafless plants; the stems globose or columnar and angled, or of flattened joints, mostly prickly. Flowers solitary, sessile. The numerous sepals and petals adherent to the 1-celled ovary. Style 1, stigmas many. Fruit a 1-celled berry with many seeds on the walls.

Synopsis of the Genera

1. *Mamilla'ria*. Stems globose or oval, covered with spine-bearing tubercles, the flowers between these.
2. *Opun'tia*. Stems of flattened joints, bearing very small, awl-shaped leaves with clusters of bristles in their axils.

1. *MAMILLA'RIA* Haw.

1. *M. vivip'ara* Haw. Stems 1-5 inches high, the tubercles bearing bundles of 5-8 reddish-brown spines, surrounded by numerous grayish ones. Flowers purple.—Pr. Prov.

2. *OPUN'TIA* Mill. PRICKLY PEAR

1. *O. Rafines'quii* Engel. Stem prostrate, deep green. Bristles reddish-brown. Spines few. Flowers yellow, sometimes with reddish centre.—Point Pelee.

THYMELAEAE'CEAE (MEZEREUM FAMILY)

Shrubs with tough, leather-like bark and entire leaves. Flowers perfect, without corolla. Calyx tubular or spreading, resembling a corolla, pale yellow. Stamens in our species twice as many as the

lobes of the calyx. Style thread-like, or short. Stigma capitate. Ovary 1-celled, 1-ovuled, free from the calyx. Fruit a berry-like drupe.

Synopsis of the Genera

1. *Dir'ca*. Calyx tubular. Style thread-like.
2. *Daph'ne*. Calyx lobes spreading. Styles short.

1. DIR'CA L. LEATHERWOOD. MOOSEWOOD

1. *D. palus'tris* L. A branching shrub, 2-5 feet high, with curious jointed branchlets and nearly oval leaves on short petioles. Flowers in clusters of 3 or 4, preceding the leaves. Filaments exserted, half of them longer than the others.—Damp woods.

2. DAPH'NE L. MEZEREUM

1. *D. Meze'reum* L. A low shrub with purple, rose-coloured, or whitish flowers, preceding the lanceolate leaves in early spring.—Escaped from cultivation in a few places.

ELAEAGNA'CEAE (OLEASTER FAMILY)

Shrubs with perfect or dioecious flowers, and leaves which are scurfy on the under-surface. The calyx tube in the fertile flowers becomes fleshy and incloses the ovary, forming a berry-like fruit. Corolla wanting.

Synopsis of the Genera

1. *Elaeag'nus*. Flowers perfect. Stamens 4. Leaves alternate.
2. *Shepherd'ia*. Flowers dioecious. Stamens 8. Leaves opposite.

1. ELAEAG'NUS L.

1. *E. argen'tea* Pursh. (SILVERBERRY.) Shrub 6-12 feet high, the young branches covered with rusty scales. Leaves elliptical to lanceolate, silvery-scurfy. Flowers many, deflexed, silvery outside, pale-yellow within, fragrant.—Pr. Prov.

2. SHEPHERD'IA Nutt. SHEPHERDIA

1. *S. canaden'sis* Nutt. A shrub 3-6 feet high. Calyx in sterile flowers 4-parted. Stamens 8. Calyx in fertile flowers urn-shaped, 4-parted. Berries yellow to reddish, nauseous. Branchlets brown-scurfy. Leaves opposite, entire, ovate, green above, silvery-scurfy beneath, the small flowers in their axils.—Gravelly banks of streams and lakes.

2. *S. argen'tea* Nutt. (BUFFALO BERRY.) A shrub 6–18 feet high with twigs often ending in thorns. Leaves oblong, silvery-scurfy on both sides. Fruit ovoid, scarlet to yellowish, sour and edible.

LYTHRA'CEAE (LOOSESTRIFE FAMILY)

Herbs or slightly woody plants, with opposite or whorled entire leaves, without stipules. Calyx inclosing but free from the ovary. Petals and stamens inserted on the calyx. Flowers axillary or whorled. Style 1. Stigma capitate.

Synopsis of the Genera

1. *Lyth'rum*. Petals mostly 6. Stamens mostly 6 or 12. Flowers purple, solitary in the axils, or forming an interrupted spike. Calyx teeth with projections in the sinuses. Pod oblong, 2-celled.
2. *Dec'odon*. Petals 5 (rarely 4). Stamens twice as many, in two sets. Pod 3–5-celled.

1. LYTH'RUM L. LOOSESTRIFE

1. *L. ala'tum* Pursh. Tall and wand-like. Flowers solitary in the upper axils. Calyx teeth often shorter than the projections between them. Petals deep purple. Stamens usually 6.

2. *L. Salica'ria* L. (SPIKED LOOSESTRIFE.) More or less downy and tall. Flowers purple, crowded, and forming an interrupted spike. Stamens twice as many as the petals, in two sets.

2. DEC'ODON J. F. Gmel. SWAMP LOOSESTRIFE

1. *D. verticilla'tus* Ell. Stems curving, 2–6 feet long, 4–6-sided. Leaves lanceolate, mostly whorled. Flowers purple, in the axils of the upper leaves. Calyx bell-shaped, with 5–7 erect teeth, with supplementary projections between them. Stamens 10, exserted, 5 longer than the rest.—Swamps.

MELASTOMA'CEAE (MELASTOMA FAMILY)

Low herbs with opposite, 3–5-ribbed leaves. Calyx tube adherent to the ovary, the limb 4-cleft. Petals 4, showy, convolute in the bud. Stamens 8, with 1-celled anthers opening by a pore at the apex; these and the petals inserted on the calyx. Style and stigma 1. Pod 4-celled, many-seeded; seeds coiled.

1. RHEXIA L. DEER-GRASS. MEADOW-BEAUTY

1. *R. virginica* L. Stem square, wing-angled. Leaves oval-lanceolate. Petals purple.—Shores of the Muskoka Lakes.

ONAGRA'CEAE (EVENING PRIMROSE FAMILY)

Herbs with perfect and symmetrical flowers, the parts of the latter in twos or fours. Calyx tube adherent to the ovary, and often prolonged above it. Petals and stamens inserted on the calyx. Style 1. Stigmas 2 or 4 or capitate.

Synopsis of the Genera

1. *Circae'a*. Petals 2, obcordate. Stamens 2. Stigma capitate. Fruit bur-like, 1-2-seeded, beset with hooked bristles. Delicate low plants with opposite leaves and very small, white flowers in racemes.
2. *Epilo'bium*. Petals 4. Stamens 8. Calyx tube hardly prolonged beyond the ovary. Fruit a linear pod, many-seeded, the seeds provided with tufts of downy hairs.
3. *Oenoth'e'ra*. Petals 4. Stamens 8. Stigma 4-lobed or discoid. Flowers yellow (white in one species). Calyx tube much prolonged. Pods cylindrical or club-shaped. Seeds without tufts.
4. *Gaur'a*. Petals 4, clawed and unequal. Stamens 8. Stigma 4-lobed, with a cup-like border. Calyx tube much prolonged. Fruit hard and nut-like, ribbed, indehiscent or nearly so, few-seeded. Flowers rose-coloured or white, turning scarlet.
5. *Ludvig'ia*. Petals 4, or none. Stamens 4. Calyx tube not prolonged. Stigma capitate.

1. CIRCAE'A L. ENCHANTER'S NIGHTSHADE

1. *C. lutetia'na* L. Stem 1-2 feet high. Leaves opposite, ovate, slightly toothed. No bracts under the pedicels. Fruit roundish, bristly hairy, 2-celled.—Rich woods.

2. *C. alpi'na* L. Stem low and delicate (3-8 inches). Leaves cordate, thin, coarsely toothed. Minute bracts under the pedicels. Fruit club-shaped, soft-hairy, 1-celled.—Deep low woods.

2. EPILO'BIUM L. WILLOW-HERB

1. *E. angustifo'lium* L. (GREAT WILLOW-HERB.) Stem 3-6 feet high, simple. Leaves lanceolate, scattered. Flowers purple,

rarely white, very showy, in a terminal raceme or spike. Stamens and style deflexed. Stigma of 4 long lobes.—Newly cleared land.

2. *E. hirsutum* L. Stem tall, erect, and densely soft-hairy, with opposite, lance-oblong leaves. Flowers smaller than in No. 1, rose-purple, forming a leafy short raceme. Stamens and style erect.

3. *E. paniculatum* Nutt. Glabrous, or pubescent above. Stem varying from 1 to 2 feet high, branching above. Leaves narrowly linear, mostly alternate and fascicled. Flowers few, small, terminating the spreading, slender and almost leafless branches. Stamens and style erect. Stigma club-shaped.—Colpoy's Bay, Lake Huron, and Pr. Prov.

4. *E. densum* Raf. Stem 1–2 feet high, erect, slender, branching above, hoary-pubescent. Leaves linear-lanceolate, nearly entire. Flowers small, corymbed at the ends of the branches, purplish or white. Petals erect. Stigma club-shaped.—Bogs.

5. *E. molle* Torr. Occasionally met with. It differs from No. 4 chiefly in having the leaves crowded, broader, and their points more obtuse. The petals are rose-coloured.—Bogs.

6. *E. palustre* L. Stem low, slender, and simple (about a foot high), finely pubescent. Leaves erect or ascending, sessile, linear to linear-lanceolate, obtuse, with revolute margins. Seeds roughened with points.—Atl. sea-coast and Pr. Prov.

7. *E. coloratum* Muhl. Stem 1–2 feet high, nearly smooth, but with 2–4 hairy lines decurrent from the leaves, the latter lanceolate or ovate-lanceolate, serrate, with conspicuous petioles. Flowers small, more or less nodding, corymbed. Petals pink, deeply notched. Coma of the seed brownish.—Not so common as the next.

8. *E. adenocaulon* Haussk. More glandular-pubescent than the last. Leaves abruptly contracted into short petioles. Flowers erect. Coma of the seed much lighter in colour.—Common in wet places.

3. OENOTHE'RA L. EVENING PRIMROSE

A. Stigma-lobes linear.

1. *O. bien'nis* L. (COMMON EVENING PRIMROSE.) Stem 2–4 feet high, hairy. Leaves oblong-lanceolate. Flowers yellow, odorous, in a leafy spike, opening in the evening or in cloudy weather. Pods oblong, narrowing towards the top.—Waste places.

Var. *grandiflo'ra* Ait. Much larger flowers, and petals as long as the calyx tube.

2. *O. pal'lida* Lindl. Flowers white, changing to rose-colour, nodding in the bud. Stem white, and commonly glabrous.—Pr. Prov.

3. *O. pu'mila* L. (SMALL EVENING PRIMROSE.) Stem low, 5–12 inches high, smooth or nearly so. Leaves spatulate, lanceolate or oblanceolate. Pods nearly sessile, club-shaped, 4-angled. Flowers pale yellow, opening in sunshine.—River- and lake-margins.

4. *O. tril'oba* Nutt. Nearly glabrous. Leaves petioled, pinnatifid or wavy-toothed. Flowers white or rose-colour. Pods ovoid, 4-angled, strongly-winged.

B. *Stigma discoid.*

5. *O. serrula'ta* Nutt. A low and slender plant, with linear-to-lanceolate leaves. Calyx tube broadly funnel-form. Petals obovate.—Pr. Prov.

4. GAUR'A L.

1. *G. coccin'ea* Pursh. Hoary and very leafy, 6–12 inches high. Leaves linear-lanceolate. Flowers small, in simple spikes, pinkish, turning red.—Pr. Prov.

2. *G. bien'nis* L. Pubescent, 4–6 feet high. Leaves oblong-lanceolate. Flowers in long spikes. Fruit 4-angled, ribbed and downy.

5. LUDVIG'IA L. FALSE LOOSESTRIFE

1. *L. palus'tris* Ell. (WATER PURSLANE.) Stems floating or creeping in the mud of ditches or river-margins, smooth. Leaves ovate, opposite, tapering into a slender petiole. Flowers sessile, solitary, usually without petals. Pod 4-sided.

2. *L. alternifo'lia* L. (SEED-BOX.) Stem branching, about 3 feet high, nearly smooth. Flowers yellow, conspicuous, peduncled in the upper axils. Leaves alternate, pointed at both ends. Capsules wing-angled.

3. *L. polycar'pa* Short and Peter. Stem 1–3 feet high, erect and branching, but producing runners at the base. Flowers small, sessile in the axils, with greenish petals or none. Leaves acute at both ends, those of the runners oblong-spatulate. Bractlets at the base of the capsules linear-awl-shaped.

HALORAGIDA'CEAE (WATER MILFOIL FAMILY)

Aquatic or marsh plants with small inconspicuous flowers, sessile in the axils of the leaves or bracts. Calyx tube adherent to the ovary, the latter 2-4-celled (in *Hippuris* of a single carpel). Limb of the calyx minute or none. Petals small or none. Stamens 1-8. Fruit indehiscent, a single seed in each cell.

Synopsis of the Genera

1. *Myriophyl'lum*. Flowers monoecious or polygamous, with the parts in fours. Petals sometimes none. Stamens 4 or 8. Immersed leaves pinnately dissected into capillary divisions. Fruit 4-celled, 4-lobed.
2. *Hippu'ris*. Flowers perfect. Stamen, style, and carpel only one. Petals none. Leaves entire, linear, acute; in whorls of 8 or 10. Fruit 1-celled, 1-seeded.
3. *Proserpina'ca*. Flowers perfect, the parts in threes. Petals none. Stems creeping at the base. Leaves alternate, the immersed ones pinnately dissected. Fruit 3-angled, 3-celled.

1. MYRIOPHYL'LUM L. WATER MILFOIL

1. *M. spica'tum* L. Stamens 8. Bracts ovate, entire, shorter than the flowers. Leaves in whorls of 3 or 4. Flowers greenish, in terminal spikes. Stems very long.—Deep water.

2. *M. verticilla'tum* L. Stamens 8. Leaves finely dissected and whorled as in No. 1. Bracts pectinate-pinnatifid, much longer than the flowers, and the spikes therefore leafy. Stem 2-4 feet long.—Stagnant water.

3. *M. heterophyl'lum* Michx. Stamens 4. Lower leaves dissected, in whorls of 4 or 5. Bracts ovate or oblanceolate, entire or finely toothed, crowded, the lower ones pinnatifid. Stem stout.—Stagnant or slow water.

4. *M. tenel'lum* Bigel. Flowering stems nearly leafless. Bracts small, entire. Flowers alternate, monoecious. Stamens 4.—Borders of ponds.

2. HIPPU'RIS L. MARE'S TAIL

1. *H. vulga'ris* L. A perennial aquatic with jointed erect stem.—Muddy margins of ponds and streams.

3. PROSERPINA'CA L. MERMAID-WEED

1. *P. palus'tris* L. (MERMAID-WEED.) Low herb. Stem creeping at base. Leaves alternate, lanceolate, sharply serrate. Petals none. Stamens 3. Fruit nut-like, 3-seeded.—Wet swamps.

ARALIA'CEAE (GINSENG FAMILY)

Herbs (with us) differing from the next Family chiefly in having, as a rule, more than 2 styles, and the fruit a drupe. The umbels, also, are either single, or corymbed, or paniced. Flowers often polygamous.

Synopsis of the Genera

1. *Ara'lia*. Leaves alternate or only 1. Carpels 5; fruit black. Umbels compound.
2. *Pa'nax*. Leaves whorled. Carpels 2-3, fruit red or yellow. Umbels single.

ARA'LIA L. WILD SARSAPARILLA

1. *A. racemo'sa* L. (SPIKENARD.) Umbels in a large compound panicle. Stem 2-3 feet high, widely branching. Leaves very large and decomposed; leaflets ovate-cordate, doubly serrate. Roots aromatic.—Rich woods.

2. *A. his'pida* Vent. (BRISTLY SARSAPARILLA. WILD ELDER.) Stem 1-2 feet high, bristly, leafy, somewhat shrubby at the base. Umbels 2-7, corymbed. Leaves twice-pinnate. Leaflets sharply serrate. Fruit black.—Rocky or sandy woods.

3. *A. nudicaul'is* L. (WILD SARSAPARILLA.) True stem very short, sending up a naked scape bearing 3 or 4 long-peduncled umbels at the summit, and one long-petioled leaf, ternately divided, and with 5 leaflets on each division. Root horizontal, aromatic.—Rich woods.

PA'NAX L. GINSENG

1. *P. quinquefo'lium* L. (GINSENG.) Leaves in a whorl of 3 at the summit of the stem, the latter a foot high. Leaflets mostly 5, long-stalked. Root stout and usually forked. Often cultivated.—Rich woods.

2. *P. trifo'lium* L. Stem 4-6 inches high. Leaves in a whorl of 3 at the summit, but the leaflets usually only 3, and sessile. Root globular or a tuber and deep in the ground.—Rich woods.

UMBELLIFERAE (PARSLEY FAMILY)

Herbs with small flowers mostly in compound umbels. Calyx tube grown fast to the surface of the ovary; calyx teeth usually minute or none. The 5 petals and 5 stamens inserted on a disk which crowns the ovary. Styles 2. Fruit dry, 2-seeded. Stems hollow. Leaves usually much-cut. Many species are poisonous.

Synopsis of the Genera

A. Seeds flat (not hollow) on the inner face.

1. **Hydrocot'yle.** Umbels simple, or one springing from the summit of another, axillary. Flowers white. Stem slender and creeping. Leaves round-kidney-shaped. Fruit ovoid, flattened laterally.
2. **Sanic'ula.** Umbels irregular (or compound), the greenish flowers capitate in the umbellets. Leaves palmately lobed or parted. Calyx evident. Fruit globular, covered with hooked prickles.

(In the Genera which follow, the umbels are regularly compound.)

3. **Dauc'us.** Stem bristly. Leaves 2-3-pinnate, or pinnatifid. Bracts of the involucre pinnatifid, very long. Flowers whitish. Fruit oblong, flattened dorsally, ribbed, the ribs bristly.
4. **Heracle'um.** Stem 3-4 feet high, woolly and grooved. Leaves 1-2-ternately compound. Flowers white, the outer corollas larger than the others. Fruit wing-margined at the junction of the carpels, flattened dorsally. Carpels 5-ribbed on the back.
5. **Pastina'ca.** Stem smooth, grooved. Leaves pinnate. Flowers yellow, all alike. Fruit as in No. 4.
6. **Loma'tium.** Low acaulescent herbs, with spindle-shaped roots and twice-pinnate leaves. No involucre; involucels of scarious-margined (often purplish), lanceolate bractlets. Flowers white or yellow. Fruit round, winged on the sides.
7. **Oxyp'olis.** Stem smooth. Leaves pinnate, of 3-9 rather narrow leaflets. Flowers white. Calyx teeth evident. Fruit broadly winged, flat, 5-ribbed on the back.
8. **Angel'ica.** Stem smooth, stout, purple. Leaves 2-3-ternately compound. Flowers greenish-white. Fruit smooth, flattish on the back, double-wing-margined, each carpel with 3 ribs on the back.
9. **Conioseli'num.** Stem smooth. Leaves 2-3-pinnately compound, the petioles inflated. Flowers white. Fruit doubly

wing-margined, and with 3 narrow wings on the back of each carpel.

10. **Thas'pium.** Stem smooth. Leaves 1-3-ternately divided. Flowers yellow. Calyx teeth conspicuous. Fruit not flattened, 10-winged or ribbed.
11. **Aethu'sa.** Leaves 2-3-ternately compound, the divisions finely dissected. No involucre: involucels of long narrow bractlets. Flowers white. Fruit ovate-globose, the carpels with 5 thick sharp ribs.
12. **Ligus'ticum.** Stem smooth, from large aromatic roots. Leaves 2-ternate, coarsely toothed. Flowers white. Fruit oblong to ovate, with prominent acute ribs, having broad spaces between them.
13. **Taenid'ia.** Stem slender, smooth and glaucous. Leaves 2-3-ternately compound. Flowers yellow. Involucre and involucels mostly wanting. Fruit oblong, flattened laterally.
14. **Ziz'ia.** Smooth plants with mostly ternately divided leaves and toothed leaflets. Calyx teeth prominent. Flowers golden yellow. Fruit ovate to oblong.
15. **Cicu'ta.** Stem streaked with purple, stout. Leaves 2-3-pinnately compound. Flowers white. Calyx teeth evident. Fruit a little contracted at the sides, the carpels strongly 5-ribbed.
16. **Ca'rum.** Stem erect, slender, smooth. Leaves pinnately decompound, the divisions filiform. Roots tuberous. Flowers white. Ribs of the ovate or oblong fruit inconspicuous.
17. **Sium.** Stem grooved. Leaves simply pinnate. Flowers white. Fruit as in No. 10.
18. **Cryptotae'nia.** Stem smooth. Leaves 3-foliolate. The umbels with very unequal rays. Flowers white. Fruit linear-oblong.

B. Inner face of each seed hollowed lengthwise.

19. **Bupleu'rum.** Stem smooth. Leaves perfoliate, ovate, entire. No involucre: involucels of 5 very conspicuous, ovate, mucronate bractlets. Flowers yellow. Fruit oblong.
20. **Muse'nium.** Stem short, branching from the base. Flowers yellow. Leaves 2-pinnatifid. Roots spindle-shaped. No involucre; involucels 1-sided, of a few narrow bractlets. Ribs of the ovate fruit slightly prominent.
21. **Osmorhi'za.** Leaves large, 2-3-ternately compound. Flowers white. Fruit linear-oblong, angled, tapering downwards into a stalk-like base. Ribs of the carpels bristly upwards.
22. **Coni'um.** Leaves large, decompound. Flowers white. Fruit ovate, flattened at the sides, 5-ribbed, the ribs wavy.

C. Inner face of each seed curved inwards at top and bottom.

23. *Erigeni'a*. Stem low and smooth. Leaves 2-3-ternately divided. Umbels 3-rayed, small. Flowers white. Fruit twin. Carpels nearly kidney-form.

1. HYDROCOTYLE L. WATER PENNYWORT

1. *H. america'na* L. Stem spreading and creeping, very slender. Leaves kidney-shaped, crenate, slightly lobed, shining. Umbels 3-5-flowered, inconspicuous, in the axils of the leaves.—Shady wet places.

2. SANIC'ULA L. SANICLE. BLACK SNAKEROOT

1. *S. marilan'dica* L. Leaves 5-7-parted. Staminate flowers very numerous, and on slender pedicels. Styles long, recurved.—Rich woods.

2. *S. canaden'sis* L. Leaves 3-5-parted. A few staminate flowers among the perfect ones, and on very short pedicels. Styles shorter than the prickles of the fruit.—Low rich woods, not so common.

3. DAUC'US L. CARROT

1. *D. Caro'ta* L. (CARROT.) A common weed in old fields and waste places. In fruit the umbel becomes hollow like a bird's nest.

4. HERACLE'UM L. COW PARSNIP

1. *H. lana'tum* Michx. Umbels large and flat. Petioles of the leaves spreading and sheathing. Leaves very large; leaflets broadly heart-shaped, deeply lobed.—Low wet meadows.

5. PASTINA'CA L. PARSNIP

1. *P. sati'va* L. (COMMON PARSNIP.) Found wild in old fields and along roadsides. Leaves pinnate; leaflets ovate, shining above.

6. LOMA'TIUM Raf.

1. *L. orienta'le* Coult. and Rose. Pubescent. Peduncles 3-8 inches high. Flowers white, in early spring.—Pr. Prov.

2. *L. daucifo'lium* Coult. and Rose. Yellow flowers.—Pr. Prov.

7. OXYP'OLIS Raf. COWBANE

1. *O. rigid'ior* Coult. and Rose. Calyx 5-toothed. Involucre almost none; involucels of many small bractlets.—Sandy swamps.

8. ANGELICA L. ANGELICA

1. *A. atropurpurea* L. (GREAT ANGELICA.) Stem very tall (4-6 feet) and stout, dark purple. Whole plant strong-scented. Petioles much-inflated at the base.—Marshes and low river-banks.

2. *A. Gmel'ini* DC. Stem slightly downy at the top. Involucels about as long as the umbellets. Plant but little aromatic.—Rocky coasts, Atl. Prov.

9. CONIOSELI'NUM Fischer. HEMLOCK PARSLEY

1. *C. chinen'se* BSP. Stem 2-4 feet high. Petioles much inflated. Leaflets of the involucels awl-shaped.—Swamps.

10. THAS'PIUM Nutt. MEADOW-PARSNIP

1. *T. aur'eum* Nutt. Stem 1-2 feet high, angular-furrowed. Leaflets oblong-lanceolate, sharply serrate. Flowers deep yellow. Fruit with 10 winged ridges.—Dry or rich woods.

2. *T. barbino'de* Nutt. Loosely branched, pubescent on the joints. Leaflets ovate to lanceolate, coarsely cut-serrate. Flowers light yellow.

11. AETHU'SA L. FOOL'S PARSLEY

1. *A. Cyna'pium*, L. A poisonous annual, with an unpleasant odour, found occasionally in cultivated grounds.

12. LIGUS'TICUM L. LOVAGE

1. *L. scoth'icum* L. (SCOTCH LOVAGE.) Stem 1-2 feet high, simple. Leaflets ovate. Fruit narrowly oblong.—Salt marshes, Atl. sea-coast.

13. TAENID'IA Drude.

1. *T. integer'rima* Drude. Stem slender, 1-2 feet high, from a stout rootstock. Involucels none. Plant strong-scented.—Rocky hillsides.

14. ZIZ'IA Koch.

1. *Z. aur'ea* Koch. Leaves 2-3-ternate, except a few at the top; radical with long petioles; leaflets ovate-lanceolate. Fruit oblong, about $\frac{1}{4}$ inch long.—Pr. Prov

2. *Z. corda'ta* DC. Stem-leaves 1 or 5-ternate; leaflets ovate-lanceolate, serrate; radical leaves usually long-petioled and cordate, only rarely lobed or divided. Fruit ovate, less than $\frac{1}{4}$ inch long.—Pr. Prov.

15. CICH'ITA L. WATER HEMLOCK

1. *C. macula'ta* L. (SPOTTED COWBANE. BEAVER-POISON.) Stem stout, 3-6 feet high, purplish, smooth. Leaflets ovate-lanceolate, coarsely serrate, pointed.—Swamps and low grounds.

2. *C. bulbif'era* L. Easily distinguished from No. 1 by the slender stem bearing clusters of bulblets in the axils of the upper leaves. The leaflets, also, are linear.—Swamps and low grounds.

16. CA'RUM L. CARAWAY

1. *C. Car'vi* L. (CARAWAY.) Escaped from cultivation in many places.

17. SIUM L. WATER PARSNIP

1. *S. cicutaefo'lium* Schrank. Stem 2-3 feet high, smooth, furrowed. Leaflets varying from linear to oblong, sharply pointed and serrate. Fruit oblong or ovate, with prominent ribs.—Borders of marshes usually in the water.

18. CRYPTOTAE'NIA DC. HONEWORT.

1. *C. canaden'sis* DC. Stem 1-2 feet high, slender. Leaflets large, ovate, doubly serrate. No involucre.—Rich woods and thickets.

19. BUPLEU'RUM L. THOROUGH-WAX

1. *B. rotundifo'lium* L. Fields and ballast-heaps.—Atl. Prov.

20. MUSE'NIUM Nutt.

1. *M. divarica'tum* Nutt. Decumbent. Leaves glabrous and shining, the divisions confluent with the winged rhachis. Rather ill-smelling herbs, with yellow flowers.—Pr. Prov.

21. OSMORHI'ZA Raf. SWEET CICELY

1. *O. longisty'lis* DC. (SMOOTHER SWEET CICELY.) Stem reddish, nearly smooth. Leaflets sparingly pubescent, short-pointed. Styles slender, nearly as long as the ovary, recurved.—Rich woods.

2. *O. Clayto'ni* Clarke. (HAIRY SWEET CICELY.) Whole plant hairy. Leaflets taper-pointed. Styles very short, conical.—Rich woods.

22. CONI'UM L. POISON HEMLOCK

1. *C. macula'tum* L. Stem smooth, spotted. Leaflets lanceolate, pinnatifid, pale green, with an offensive odour when bruised.

Involucels one-sided. Inner face of the seed marked with a deep groove.—Waste places.

23. ERIGENIA Nutt. **HARBINGER-OF-SPRING**

1. **E. bulbo'sa** Nutt. Stem 4-6 inches high, from a tuber deep in the ground, producing 2 leaves, the lower radical. Leaflets much incised. Flowers few.—Alluvial soil.

CORNA'CEAE (DOGWOOD FAMILY)

Shrubs or trees (rarely herbs) with simple leaves. Calyx tube adherent to the 1-2-celled ovary, the limb of the calyx inconspicuous. Petals and stamens all epigynous. Style 1. Fruit a 1-2-seeded drupe.

Synopsis of the Genera

1. **Cor'nus**. Flowers perfect (the parts in fours), either forming cymes, or in close heads surrounded by a showy involucre resembling a corolla. Leaves mostly opposite. Stigma flat or capitate.
2. **Nys'sa**. Flowers dioeciously polygamous, the parts in fives. Leaves alternate, mostly crowded at the ends of the branchlets. Stigma elongated.

1. COR'NUS L. **CORNEL**. **DOGWOOD**

A. Flowers in a close head, surrounded by a showy involucre of 4 white bracts. Fruit red.

1. **C. canadensis** L. (**BUNCH-BERRY**.) Stem simple, 5 or 6 inches high. Upper leaves crowded and apparently whorled, in fours or sixes (rarely opposite), ovate, the lower scale-like. Leaves of the whitish involucre ovate. Flowers greenish white.—Rich woods.

2. **C. flor'ida** L. (**FLOWERING DOGWOOD**.) A small tree, with opposite, ovate, pointed leaves. Leaves of the white involucre notched at the apex, very showy.—Rocky woods.

B. Flowers (white) in flat cymes. No involucre. Fruit blue or white.

3. **C. circina'ta** L'Her. (**ROUND-LEAVED DOGWOOD**.) A shrub 4-6 feet high, with greenish, warty, dotted branches. Leaves opposite, broadly oval, white-woolly beneath. Fruit light blue.—Rich woods.

4. *C. Amo'mum* Mill. (SILKY CORNEL.) A large shrub with purplish branches. Leaves opposite, narrowly ovate or oblong, silky beneath. Branchlets often rusty. Fruit light blue. Distinguished from No. 3 by the colour of the branches and the much smaller leaves.—Low wet grounds.

5. *C. stolonif'era* Michx. (RED-OSIER DOGWOOD.) A shrub forming clumps by the production of suckers or stolons, 3–6 feet high. Branches bright red-purple, smooth. Leaves opposite, ovate, roughish, whitish beneath. Fruit white or whitish.—Low wet grounds.

6. *C. panicula'ta* L'Her. (PANICLED CORNEL.) A shrub, 4–8 feet high, with erect, gray, and smooth branches. Flowers white, very numerous. Leaves opposite, ovate-lanceolate, taper-pointed. Cymes convex. Fruit white, on red stalks.—Thickets and river-banks.

7. *C. asperifo'lia* Michx. (ROUGH-LEAVED DOGWOOD.) Branches brownish, the branchlets rough-pubescent. Leaves opposite, rather small, oblong or ovate; rough above, downy beneath. Petals rather long. Fruit white.—Reported by Macoun as common on Point Pelee.

8. *C. Baile'yi* Coult. and Evans. Branchlets brownish and pubescent, not rough. Leaves ovate-lanceolate, pubescent both sides. Petals short. Fruit pure white.

9. *C. alternifo'lia* L. (ALTERNATE-LEAVED CORNEL.) A large shrub or small tree, with alternate greenish branches streaked with white. Leaves mostly alternate, oval, acute at the base, pointed, crowded at the ends of the branches. Flowers yellowish, in loose cymes. Fruit deep blue, on reddish stalks.—Thickets.

2. NYS'SA L. PEPPERIDGE. SOUR GUM

1. *N. sylvat'ica* Marsh. A middle-sized tree with horizontal branches, reported from Niagara and South-western Ontario. Leaves varying from elliptic to obovate. Sterile flowers in dense axillary clusters, with small calyx, and small fleshy petals or none. Stamens mostly 10, but sometimes fewer. Fertile flowers in clusters of 3–8, at the summit of a slender axillary peduncle, larger than the staminate ones. Style revolute. Ovary 1-celled. Drupe ovoid, bluish-black, about $\frac{1}{2}$ inch long.

ERICA'CEAE (HEATH FAMILY)

Chiefly shrubs, distinguished by the anthers opening, as a rule, by a pore at the top of each cell. Stamens free from the corolla, as many or twice as many as its lobes. Leaves simple and usually alternate. Corolla in some cases polypetalous.

Synopsis of the Genera

SUBFAMILY I. VACCINOI'DEAE (WHORTLEBERRY SUBFAMILY)

Calyx tube adherent to the ovary. Fruit a berry crowned with the calyx teeth.

1. **Gaylussa'cia.** Stamens 10, the anthers opening by a pore at the apex. Corolla tubular, ovoid, the border 5-cleft. Berry 10-celled, 10-seeded. Flowers white with a red tinge. Leaves covered with resinous dots. Branching shrubs.
2. **Vaccin'ium.** Stamens 8 or 10, the anthers prolonged upwards into tubes with a pore at each apex. Corolla deeply 4-parted and revolute, or cylindrical with the limb 5-toothed. Berry 4-celled, or more or less completely 10-celled. Flowers white or reddish, solitary or in short racemes. Shrubs.
3. **Chiog'enes.** Stamens 8, each anther 2-pointed at the apex. Corolla bell-shaped, deeply 4-cleft. Limb of the calyx 4-parted. Flowers very small, nodding from the axils, with 2 bractlets under the calyx. Berry white, 4-celled. A trailing slender evergreen.

SUBFAMILY II. ERICOI'DEAE (HEATH SUBFAMILY)

Calyx free from the ovary. Shrubs or small trees. Corolla gamopetalous, except in No. 11.

4. **Arctostaph'ylos.** Corolla urn-shaped, the limb 5-toothed, revolute. Stamens 10, the anthers each with 2 reflexed awns on the back. Fruit a berry-like drupe, 5-10-seeded. A trailing, thick-leaved evergreen, with nearly white flowers.
5. **Epigae'a.** Corolla salver-shaped, hairy inside, rose-coloured. Stamens 10; filaments slender; anthers awnless, opening lengthwise. Calyx of 5 pointed and scale-like, nearly distinct sepals. A trailing evergreen, bristly with rusty hairs. Flowers rose-coloured.
6. **Gaulthe'ria.** Corolla ovoid or slightly urn-shaped, 5-toothed, nearly white. Stamens 10, the anthers 2-awned. Calyx 3-cleft, closing the pod and becoming fleshy and berry-like in fruit. Stems low and slender, leafy at the summit.

7. **Chamaedaph'ne.** Corolla cylindrical, 5-toothed. Stamens 10, the anther-cells tapering into beaks with a pore at the apex, awnless. Calyx of 5 overlapping sepals, and 2 similar bractlets. Pod with a double pericarp, the outer of 5 valves, the inner cartilaginous and of 10 valves. A low shrub with rather scurfy leaves and white flowers.
8. **Androm'eda.** Corolla globular-urn-shaped, 5-toothed. Calyx of 5 nearly distinct, valvate sepals, without bractlets. Stamens 10; the filaments bearded; the anther-cells each with a slender awn. A low shrub with white flowers in a terminal umbel.
9. **Callu'na.** Corolla bell-shaped, 4-parted, persistent, becoming scarious. A low evergreen shrub, with numerous minute opposite leaves. Flowers rose-coloured or white, in mostly 1-sided racemes.
10. **Kal'mia.** Corolla broadly bell-shaped, with 10 pouches receiving as many anthers. Shrubs with showy, rose-purple flowers.
11. **Le'dum.** Calyx 5-toothed, very small. Corolla of 5 obovate and spreading, nearly distinct petals. Stamens 5-10. Leaves evergreen, with revolute margins, covered beneath with rusty wool.
12. **Rhododen'dron.** Corolla irregular (in our species), nearly an inch long, 2-lipped, the upper lip 3-lobed, the lower of 2 oblong-linear, curved, nearly or quite distinct petals. Stamens 10 (or sometimes 5), as long as the rose-coloured corolla. A shrub with alternate, oblong, somewhat pubescent leaves.—Atl. Prov.

SUBFAMILY III. PYROLOI'DEAE (PYROLA SUBFAMILY)

Calyx free from the ovary. Corolla polypetalous. More or less herbaceous evergreens.

13. **Pyr'ola.** Calyx 5-parted. Petals 5, concave. Stamens 10. Stigma 5-lobed. Leaves evergreen, clustered at the base of an upright, scaly-bracted scape, which bears a simple raceme of nodding flowers.
14. **Mone'ses.** Petals 5, orbicular, spreading. Stamens 10. Stigma large, peltate, with 5 narrow radiating lobes. Plant having the aspect of a *Pyrola*, but the scape bearing a single terminal flower.
15. **Chimaph'ila.** Petals 5, concave, orbicular, spreading. Stamens 10. Stigma broad and round, the border 5-crenate. Low plants with running, underground shoots, and thick, shining, sharply serrate, somewhat whorled leaves. Flowers corymbed or umbelled on a terminal peduncle.

SUBFAMILY IV. *MONOTROPOIDEAE* (INDIAN PIPE SUBFAMILY)

16. *Monotropa*. Smooth reddish or whitish plants parasitic on roots, bearing scales instead of leaves. Flowers solitary, terminal or in scaly racemes.
17. *Pterospora*. A purplish-brown, clammy-pubescent plant, parasitic on the roots of pines. Stem simple. Flowers numerous, nodding, white, forming a raceme.

1. *GAYLUSSACIA* HBK. HUCKLEBERRY

1. *G. bacca'ta* C. Koch. (BLACK HUCKLEBERRY.) Fruit black, without a bloom. Racemes short, 1-sided, in clusters. Leaves oval or oblong. Branching shrub, 1-3 feet high.—Low grounds.

2. *G. dumo'sa* Torr. and Gr. (DWARF HUCKLEBERRY.) Fruit black, insipid. Racemes long, with leaf-like, persistent bracts. Leaves obovate, oblong, mucronate.—Sandy low ground, Atl. Prov.

2. *VACCINIUM* L. CRANBERRY. BLUEBERRY

1. *V. Oxycoc'cos* L. (SMALL CRANBERRY.) A creeping or trailing, very slender, shrubby plant with ovate, acute, evergreen leaves only $\frac{1}{4}$ inch long, the margins revolute. Corolla rose-coloured, 4-parted, the lobes reflexed. Anthers 8. Stem 4-9 inches long. Berry only about $\frac{1}{4}$ inch across, often speckled with white.—Bogs.

2. *V. macrocar'pon* Ait. (LARGE OR AMERICAN CRANBERRY.) Different from No. 1 in having prolonged stems (1-3 feet long) and the flowering branches lateral. The leaves also are oblong-elliptic and nearly twice as large, and the berry is fully $\frac{1}{2}$ inch broad.—Bogs.

3. *V. Vi'tis-Idaea* L. A low plant with erect branches from tufted creeping stems. Leaves evergreen, obovate, with revolute margins, shining above, dotted with blackish bristly points beneath. Corolla bell-shaped, 4-lobed. Anthers 8-10. Flowers in a short bracted raceme. Berry dark red and acid.—Northward and eastward.

4. *V. pennsylvan'icum* Lam. (DWARF BLUEBERRY.) Stem 6-15 inches high, the branches green, angled and warty. Corolla cylindrical, bell-shaped, 5-toothed. Anthers 10. Flowers in short racemes. Leaves lanceolate or oblong, serrulate with

bristly-pointed teeth, smooth and shining on both sides. Berry blue or black, with or without a bloom.—Dry plains and woods.

5. *V. canadense* Kalm. (CANADIAN BLUEBERRY.) Stem 1–2 feet high. Leaves oblong-lanceolate or elliptical, entire, downy both sides, as are also the branchlets.—A very common Canadian species.

6. *V. vacillans* Kalm. (LOW BLUEBERRY.) Stem 1–2 feet high, glabrous, with yellowish-green branchlets. Leaves obovate or oval, very pale or dull and glaucous, at least beneath. Corolla between bell-shaped and cylindraceous, the mouth somewhat contracted. Berries blue with a bloom.—Dry soil.

7. *V. corymbose* L. (SWAMP BLUEBERRY.) A tall shrub (3–10 feet) growing in swamps and low grounds, with leaves varying from ovate to elliptical-lanceolate; berries black with a bloom, many-seeded and ripening late.

Var. *amoenum* Gray. Leaves bristly-ciliate, green both sides, shining above.

Var. *palidum* Gray. Leaves mostly glabrous, pale or whitish, glaucous at least underneath, and serrulate with bristly teeth.

8. *V. atrococcum* Heller. Leaves entire, downy, or woolly underneath, as well as the branchlets. Berries shiny-black, without a bloom.

9. *V. stamineum* L. (DEER-BERRY. SQUAW HUCKLEBERRY.) Stem diffusely branching, 2–3 feet high. Leaves ovate or oval, pale, whitish beneath. Corolla open-bell-shaped, 5-lobed, greenish-white or purplish. Anthers 2-awned on the back, much exserted. Flowers slender-pedicelled. Berries greenish or yellowish, large.

3. CHIOG'ENES Salisb. CREEPING SNOWBERRY

1. *C. hispidula* Torr. and Gr. Leaves very small, ovate and pointed, on short petioles, the margins revolute. The lower surface of the leaves and the branches clothed with rusty bristles. Berries bright white.—Bogs and cool woods.

4. ARCTOSTAPH'YLOS Adans. BEARBERRY

1. *A. Uva-ursi* Spreng. Flowers in terminal racemes. Leaves alternate, obovate or spatulate, entire, smooth. Berry red.—Bare hillsides.

5. *EPIGAE'A* L. GROUND LAUREL. TRAILING ARBUTUS

1. *E. re'pens* L. (MAYFLOWER.) Flowers in small axillary clusters from scaly bracts. Leaves evergreen, rounded and heart-shaped, alternate, on slender petioles. Flowers very fragrant.—Dry woods in early spring.

6. *GAULTHE'RIA* L. AROMATIC WINTERGREEN

1. *G. procum'bens* L. (TEABERRY. WINTERGREEN.) Flowers mostly single in the axils, nodding. Leaves obovate or oval, obscurely serrate, evergreen. Berry bright red, edible.—Cool woods, chiefly in the shade of evergreens.

7. *CHAMAEDAPH'NE* Moench. LEATHER-LEAF

1. *C. calycula'ta* Moench. Flowers in 1-sided, leafy racemes. Leaves oblong, obtuse, flat.—Bogs.

8. *ANDROM'EDA* L. ANDROMEDA

1. *A. polifo'lia* L. Stem smooth and glaucous, 6–18 inches high. Leaves oblong-linear, with strongly revolute margins, generally white beneath.—Bogs.

9. *CALLU'NA* Salisb. HEATHER

1. *C. vulga'ris* Hull. (HEATHER.) A low evergreen shrub with numerous opposite, minute leaves, mostly auricled at the base. Flowers axillary or terminating very short shoots, forming close racemes mostly one-sided, rose-coloured or white. Calyx of 4 sepals. Corolla 4-parted, bell-shaped. Calyx and corolla both persistent and becoming dry. Stamens 8. Capsule 4-celled.—Found sparingly in a few places on the coast of Nova Scotia and Newfoundland.

10. *KAL'MIA* L. AMERICAN LAUREL

1. *K. polifo'lia* Wang. (PALE LAUREL.) A straggling shrub about a foot high, with few-flowered, terminal corymbs. Branchlets 2-edged. Leaves opposite, oblong, the margins revolute. Flowers $\frac{1}{2}$ inch across.—Bogs.

2. *K. angustifo'lia* L. (SHEEP LAUREL. LAMBKILL.) Leaves opposite or in threes, oblong, obtuse, petioled. Corymbs lateral, many-flowered. Pod depressed. Pedicels recurved in fruit.—Bogs and damp barren grounds, abundant eastward.

11. LE'DUM L. LABRADOR TEA

1. *L. groenlan'dicum* Oeder. Flowers white, in terminal umbel-like clusters. Leaves elliptical or oblong. Stamens 5, or occasionally 6 or 7.—Bogs.

12. RHODODEN'DRON L. ROSE BAY. AZALEA

1. *R. canaden'se* BSP. Corolla irregular, nearly an inch long, two-lipped; the upper lip 3-lobed; the lower lip of 2 oblong-linear, curved, nearly or quite distinct petals. Stamens 10, as long as the rose-coloured corolla. Leaves alternate, oblong, somewhat pubescent. Shrub.—Bogs and damp barrens, from Montreal eastward.

13. PYR'OLA L. WINTERGREEN. SHIN-LEAF

A. *Styles straight, narrower than the 5-rayed stigma.*

1. *P. mi'nor* L. Leaves roundish, slightly crenulate, thickish, usually longer than the margined petiole. Raceme not one-sided. Flowers white or rose-colour. Style short and included in corolla.—Cold woods, Atl. Prov.

2. *P. secun'da* L. Easily recognized by the flowers of the dense raceme being all turned to one side. Leaves ovate. Style long, protruding.—Rich woods.

Var. *obtusa'ta* Turcz. Orbicular leaves. 3-8-flowered.—Peat-bogs and swamps.

B. *Style declined, the apex curved upward. Stigma narrower than the ring-like apex of the style.*

3. *P. america'na* Sweet. Leaves orbicular, thick, shining, usually shorter than the petiole. Calyx lobes lanceolate. Flowers white.—Moist woods.

4. *P. asarifo'lia* Michx. Leaves round-reniform, and petals mostly rose-coloured.

Var. *incarna'ta* Fernald. Leaves obovate to suborbicular, rounded at the base. Flowers rose-purple.

5. *P. ellip'tica* Nutt. (SHIN-LEAF.) Leaves elliptical, thin, dull, usually longer than the margined petiole. Flowers greenish-white.—Rich woods.

6. *P. chloran'tha* Swartz. Leaves small, roundish, and dull, petals converging, greenish-white, and anther-cells contracted below the pore into a distinct neck or horn.—Open woods.

14. MONE'SES Salisb. ONE-FLOWERED PYROLA

1. *M. uniflo'ra* Gr. Leaves thin, rounded, veiny, and serrate. Scape 2-4 inches high, bearing a single white or rose-coloured flower.—Deep woods.

15. CHIMAPH'ILA Pursh. PIPSISSEWA

1. *C. umbella'ta* Nutt. (PRINCE'S PINE.) Leaves wedge-lanceolate, acute at the base. Peduncles 4-7-flowered. Corolla rose- or flesh-coloured.—Dry woods.

2. *C. macula'ta* Pursh. (SPOTTED WINTERGREEN.) Leaves ovate-lanceolate, obtuse at the base, the upper surface variegated with white.—Dry woods.

16. MONOT'ROPA L. INDIAN PIPE. PINE-SAP

1. *M. uniflo'ra* L. (INDIAN PIPE. CORPSE PLANT.) Smooth, waxy-white, turning black in drying.—Dark rich woods.

2. *M. Hypop'itys* L. A tawny or reddish parasitic plant with several flowers in a scaly raceme. Somewhat pubescent. Sepals bract-like. Stigma ciliate. Style longer than the ovary, hollow. Pod globular or ovoid.—Oak and pine woods.

17. PTEROS'PORA Nutt. PINE-DROPS

1. *P. andromede'a* Nutt. Calyx 5-parted. Corolla ovate, urn-shaped, 5-toothed, persistent. Stamens 10. Stigma 5-lobed. Pod 5-lobed, 5-celled.—Usually under pines in dry woods.

PLUMBAGINA'CEAE (LEADWORT FAMILY)

Maritime herbs with regular pentamerous flowers, a plaited calyx, 5 stamens opposite the lobes (or separate petals) of the corolla, and a 1-celled and 1-seeded ovary.

1. LIMO'NIUM Hill.

1. *L. carolinia'num* Britton. (MARSH ROSEMARY.) A maritime herb with a thick, woody, astringent root, and oblong, spatulate, or obovate-lanceolate, radical leaves, tipped with a deciduous bristle. Flowers lavender-coloured, paniced on branching scapes. Calyx funnel-form, membranaceous. Corolla of 5 nearly or quite distinct petals, with the 5 stamens severally borne on their bases. Ovary 1-celled and 1-ovuled.—Salt marshes, Atl. Prov.

PRIMULA'CEAE (PRIMROSE FAMILY)

Herbs with regular perfect flowers, well marked by having a stamen before each petal or lobe of the corolla and inserted on the tube. Ovary 1-celled, the placenta rising from the base. Style 1, stigma 1.

Synopsis of the Genera

A. *Stemless. Leaves all in a cluster from the root.*

1. **Prim'ula.** Flowers in an umbel at the summit of a simple scape. Corolla salver-shaped or funnel-form, open at the throat. Stamens 5, included.
2. **Andros'ace.** Flowers very small, white, in an umbel at the summit of a scape. Corolla salver-shaped or funnel-form, constricted at the throat. Stamens included.
3. **Dodeca'theon.** Flowers showy, rose-coloured or white, in an umbel at the summit of a scape. Corolla reflexed, 5-parted. Stamens exserted, connivent in a slender cone.

B. *Stems leafy. Corolla rotate, wanting in Glaux.*

4. **Trienta'lis.** Leaves in a whorl at the summit of a slender erect stem. Calyx usually 7-parted, the lobes pointed. Corolla usually 7-parted, spreading, without a tube. Filaments united in a ring below. Flowers usually only one, white and star-shaped.
5. **Lysima'chia.** Leafy-stemmed. Flowers yellow, axillary or in a terminal raceme. Calyx usually 5-parted. Corolla wheel-shaped, mostly 5-parted, and sometimes polypetalous.
6. **Steirone'ma.** Differs from No. 5 in having 5 slender staminodia between the fertile stamens and in the absence of dots on the foliage and petals.
7. **Glaux.** A fleshy herb with purplish and white, axillary flowers. Corolla wanting, the calyx petal-like.
8. **Anagal'lis.** Low and spreading. Leaves opposite or whorled, entire. Flowers variously coloured, solitary in the axils. Calyx 5-parted. Corolla wheel-shaped, 5-parted. Filaments bearded.
9. **Sam'olus.** Smooth and spreading, 6-10 inches high. Corolla bell-shaped, 5-parted, with 5 sterile filaments in the sinuses. Calyx partially adherent to the ovary. Flowers very small, white, racemed. Leaves alternate.

1. PRIM'ULA L. PRIMROSE. COWSLIP

1. **P. farino'sa** L. (BIRD'S-EYE PRIMROSE.) Lower surface of leaves covered with a white mealiness. Corolla lilac with a yellow centre.

2. *P. mistassin'ica* Michx. Leaves not mealy. Corolla flesh-coloured, the lobes obcordate.

2. ANDROS'ACE L.

1. *A. septentriona'lis* L. Almost glabrous, 2-10 inches high. Leaves lanceolate to oblong-lanceolate, narrowed at the base, toothed. Bracts of the involucre awl-shaped.—Pr. Prov.

2. *A. occidenta'lis* Pursh. A low annual, only slightly pubescent. Leaves oblong-ovate, entire, sessile. Corolla white, shorter than the leafy calyx tubes.—Pr. Prov.

3. DODECA'THEON L. AMERICAN COWSLIP

1. *D. Mea'dia* L. (SHOOTING-STAR.) Smooth. Leaves oblong or spatulate. Flowers nodding on slender pedicels.—Pr. Prov.

4. TRIENTA'LIS L. CHICKWEED WINTERGREEN

1. *T. america'na* Pursh. (STAR FLOWER.) Leaves thin and veiny, lanceolate, tapering towards both ends. Petals pointed.—Moist woods.

5. LYSIMA'CHIA L. LOOSESTRIFE

1. *L. thyrsiflo'ra* L. (TUFTED LOOSESTRIFE.) Flowers in spike-like clusters from the axils of a few of the upper leaves. Petals lance-linear, purplish-dotted, usually as many minute teeth between them. Leaves scale-like below, the upper lanceolate, opposite, sessile, dark-dotted.—Wet swamps.

2. *L. terres'tris* BSP. Flowers on slender pedicels in a long terminal raceme. Petals lance-oblong, streaked with dark lines. Leaves opposite, lanceolate, acute at each end, sessile, dark-dotted.—Low grounds.

3. *L. quadrifo'lia* L. Flowers on long slender peduncles from the axils of the upper leaves. Petals streaked. Leaves mostly in whorls of 4 or 5, ovate-lanceolate, dark-dotted.—Sandy soil.

4. *L. nummula'ria* L. (MONEYWORT.) Escaped from gardens in a few places. Stem trailing and creeping, smooth. Leaves roundish, small, opposite. Peduncles axillary, 1-flowered. Petals not streaked or dotted.—Damp places.

6. STEIRONE'MA Raf.

1. *S. cilia'tum* Raf. Flowers nodding on slender peduncles from the upper axils. Petals not streaked or dotted. Leaves opposite,

not dotted, ovate-lanceolate, pointed, cordate at the base, on long fringed petioles.—Low grounds.

2. *S. quadriflorum* Hitchc. Petals not streaked or dotted. Stem-leaves sessile, narrowly linear, 2–4 inches long, the margins sometimes revolute. Stem 4-angled.—Moist soil.

3. *S. lanceolatum* Gray. Stem erect, 1–2 feet high. Leaves lanceolate to oblong and linear, narrowed into a short, margined petiole, or the lowest short and broad on long petioles, not dotted. Petals not streaked or dotted.

7. GLAUX L.

1. *G. maritima* L. (SEA MILKWORT.) A fleshy herb, with usually opposite, oblong, entire, sessile leaves. Flowers single in the axils, nearly sessile. Calyx bell-shaped, 5-cleft, purplish and white. Corolla wanting. Stamens 5, on the base of the calyx. Capsule 5-valved, few-seeded.—Seashore, Atl. Prov.

8. ANAGALLIS L. PIMPERNEL

1. *A. arvensis* L. (COMMON PIMPERNEL.) Petals obovate, fringed with minute teeth, mostly bluish or purplish. Flowers closing at the approach of rain. Leaves ovate, sessile.—Sandy fields and garden soil.

9. SAMOLUS L. WATER PIMPERNEL. BROOK-WEED

1. *S. floribundus* HBK. Stem slender, diffusely branched. The slender pedicels each with a bractlet at the middle.—Wet places, not common.

OLEACEAE (OLIVE FAMILY)

Trees with pinnate leaves, and polygamous, dioecious or monocious flowers without petals, and sometimes also without a calyx; stamens usually 2, with large oblong anthers. Fruit a 1–2-seeded samara. Flowers insignificant, from the axils of the previous year's leaves.

1. FRAXINUS L. ASH

A. Leaflets with petioles.

1. *F. americana* L. (WHITE ASH.) Fruit winged from the apex only, the base cylindrical. Branchlets and petioles smooth and glabrous. Calyx very minute, persistent. Leaflets 7–9, stalked.—Rich woods.

2. *F. pennsylvan'ica* Marsh. (RED ASH.) Branchlets and petioles softly pubescent. Leaflets 5-9, almost entire, usually pale beneath. Fruit acute at the base, 2-edged, and gradually expanding into the long wing above. Calyx as in No. 1.

Var. *lanceola'ta* Sarg. (GREEN ASH.) Glabrous throughout. Fruit as in No. 2. Leaflets 5-9, often serrate towards the tip, bright green both sides.—Along streams.

3. *F. quadrangula'ta* Michx. (BLUE ASH.) Branchlets often square, smooth. Leaflets 7-11, sharply serrate, green both sides. Calyx wanting. Fruit narrowly oblong, of the same width at both ends, often notched at the apex, wing-margined.

B. Lateral leaflets sessile.

4. *F. ni'gra* Marsh. (BLACK or SWAMP ASH.) Branchlets and petioles smooth. Leaflets 7-11, sessile, serrate, green. Fruit winged all around. Calyx wanting.—Swamps.

GENTIANA'CEAE (GENTIAN FAMILY)

Smooth herbs, distinguished by having a 1-celled ovary with seeds on the walls, either in lines or on the whole inner surface. Leaves mostly opposite, simple, and sessile, but in one Genus (*Menyanthes*) alternate and compound. Stamens as many as the lobes of the regular corolla, inserted on its tube and alternate with them. Stigmas 2. Calyx persistent. Juice colourless and bitter.

Synopsis of the Genera

1. *Frase'ra*. Corolla wheel-shaped, 4-parted, fringed glandular spot on each lobe. Flowers light greenish-yellow, with small, purple-brown spots.
2. *Hale'nia*. Corolla 4-lobed, the lobes all spurred at the base. Flowers yellowish or purplish, somewhat cymose.
3. *Gentia'na*. Corolla not spurred, 4-5-lobed, mostly funnel-form or bell-shaped, generally with teeth or folds in the sinuses of the lobes. Stigmas 2, persistent. Pod oblong. Seeds innumerable. Flowers showy, in late summer and autumn.
4. *Menyan'thes*. A bog plant. Corolla short, funnel-form, 5-lobed, densely white-bearded on the upper face. Leaves alternate, compound, of 3 oval leaflets. The flowers in a raceme at the summit of a naked scape, white or tinged with pink.

5. *Nymphoides*. An aquatic with simple, round-heart-shaped, floating leaves on long petioles. Corolla white, wheel-shaped, 5-parted, bearded on the margins only. Flowers in an umbel borne on the petiole.

1. *FRASE'RA* Walt. AMERICAN COLUMBO

1. *F. carolinien'sis* Walt. Tall and showy. Leaves whorled, mostly in fours. Root thick. Flowers numerous in a pyramidal panicle.—Dry soil.

2. *HALE'NIA* Bork. SPURRED GENTIAN

1. *H. deflex'a* Griseb. Stem erect, 9–18 inches high. Leaves 3–5-nerved, those at the base of the stem oblong-spatulate, petioled; the upper acute and sessile or nearly so. Spurs of the corolla curved.

3. *GENTIA'NA* L. GENTIAN

1. *G. crini'ta* Froel. (FRINGED GENTIAN.) Corolla funnel-form, 4-lobed, the lobes fringed on the margins; no plaited folds in the sinuses. Flowers sky-blue, solitary, on long naked stalks, terminating the stem or simple branches. Ovary lanceolate. Leaves lance-shaped or ovate-lanceolate.—Low grounds.

2. *G. proc'era* Holm. (SMALLER FRINGED GENTIAN.) Distinguished from No. 1 by the shorter or almost inconspicuous fringe of the corolla, the linear or lance-linear leaves, and the elliptical ovary.—Moist grounds.

3. *G. quinquefo'lia* L. (FIVE-FLOWERED GENTIAN.) Corolla tubular-funnel-form, pale blue, no folds in sinuses. Calyx 5-cleft, lobes awl-shaped. Lobes of the corolla triangular-ovate, bristle-pointed. Anthers separate. Stem slender and branching, a foot high, the branches racemed or paniced, about 5-flowered at the summit.—Dry hillsides.

4. *G. affi'nis* Griseb. Stems 1–1½ feet high, clustered and leafy. Leaves linear to oblong, rough on the margins. Flowers blue, numerous to few, in raceme-like clusters; corolla funnel-form with spreading lobes, between which are plaits with appendages almost or as long as the lobes.

5. *G. puber'ula* Michx. Stems erect or ascending, 8–16 inches high, minutely rough above. Leaves rigid, lanceolate to linear-lanceolate, 1–2 inches long. Flowers mostly clustered. Calyx lobes lanceolate, much shorter than the bell-funnel-form, open, bright-blue corolla.

6. *G. flav'ida* Gray. (WHITISH GENTIAN.) Corolla inflated-club-shaped, at length open, 5-lobed, lobes about twice as long as the toothed appendages in the sinuses. Flowers greenish-white or yellowish, sessile, crowded in a terminal cluster. Anthers usually cohering. Leaves lance-ovate, with a clasping, heart-shaped base.—Low grounds.

7. *G. Sapona'ria* L. Stems smooth. Leaves ovate-lanceolate or obovate. Calyx lobes linear, not shorter than the tube, half the length of the corolla. Lobes of the light-blue corolla obtuse and broad; appendages 2-cleft and finely toothed.

8. *G. Andrews'ii* Griseb. (CLOSED GENTIAN.) Corolla inflated-club-shaped, closed at the mouth, the apparent lobes being really the large, fringed-toothed appendages. Flowers blue, rarely white, in a close, sessile, terminal cluster. Anthers cohering. Leaves ovate-lanceolate from a narrower base.—Low grounds.

9. *G. Amarel'la* L., var. *acu'ta* Herder. Corolla somewhat funnel-form, mostly blue, its lobes entire, acute, with a fringed crown at their base. Calyx lobes (4-5) lanceolate or linear, foliaceous.—Atl. Prov. chiefly.

10. *G. linea'ris* Froel., var. *latifolia* Gray. Flowers in a terminal cluster with a leafy involucre. Corolla blue, narrow-funnel-form, with roundish-ovate lobes, and broad appendages. Leaves sessile, oblong-linear to ovate-lanceolate, smooth. Seeds winged.—Boggy places, Atl. Prov.

4. MENYAN'THES L. BUCKBEAN

1. *M. trifolia'ta* L. A common plant in bogs and wet places. The bases of the long petioles sheathe the lower part of the scape, or thick rootstock, from which they spring. Plant about a foot high.

5. NYMPHOIDES Hill. FLOATING HEART

1. *N. lacuno'sum* Fernald. In shallow waters.

APOCYNACEAE (DOGBANE FAMILY)

Herbs or slightly shrubby plants with milky juice, opposite, simple, entire leaves, and regular, pentamerous and pentandrous flowers with the lobes of the corolla convolute in the bud. Dis-

tinguished by having 2 separate ovaries, but the 2 stigmas united. Calyx free from the ovaries. Stamens inserted on the corolla. Anthers converging around the stigmas. Seeds with a tuft of down on the apex.

1. APOCYNUM L. DOGBANE

1. *A. androsaemifolium* L. (SPREADING DOGBANE.) The corolla bell-shaped, 5-cleft, pale rose-coloured, the lobes turned back. Branches of the stem widely forking. Flowers fragrant, nodding, in loose, rather spreading cymes. Leaves ovate, petioled, dark green above, smooth. Fruit 2 long and slender, diverging pods.—Banks and thickets.

2. *A. cannabinum* L. (INDIAN HEMP.) Lobes of the greenish-white corolla not turned back. Branches erect. Leaves ovate-oblong to lanceolate, usually pale green. Cymes closer than in No. 1, and the flowers much smaller, and erect.—Along streams, very variable.

Var. *pubescens* DC. Pedicels pubescent and leaves white-pubescent beneath.

ASCLEPIADA'CEAE (MILKWEED FAMILY)

Herbs with milky juice and opposite or whorled (rarely scattered), simple, entire leaves, and regular, pentamerous and pentandrous flowers. Pods, seeds, and anthers as in the last Family, but the anthers are more closely connected with the stigma, the (reflexed) lobes of the corolla are valvate in the bud, the pollen is in waxy masses, and the (monadelphous) short filaments bear 5 curious, hooded bodies behind the anthers. Flowers in umbels.

Synopsis of the Genera

1. *Asclepias*. Corolla reflexed, deeply 5-parted. A crown of 5 hooded, fleshy bodies with an incurved horn rising from the cavity of each hood and resting on the stamen tube, which is inserted on the corolla. Leaves mostly opposite or whorled.
2. *Acera'tes*. Flowers in dense umbels. Corolla reflexed or merely spreading. Crown as in No. 1, but no incurved horn. Leaves mostly alternate, often opposite, petioles short or entirely absent.—Plants growing in dry soil.

1. ASCLE'PIAS L. MILKWEED

A. Flowers greenish, yellowish, or white, or merely purplish-tinged.

1. *A. syri'aca* L. (COMMON MILKWEED.) Stem tall and stout. Leaves oval or oblong, short-petioled, pale green, 4-8 inches long. Flowers dull greenish-purple. Pods ovate, soft-spiny, woolly.—Mostly in dry soil; very common.

2. *A. specio'sa* Torr. Finely white-woolly or becoming glabrous, the many-flowered umbel and calyx densely woolly. Leaves oval to oblong, slightly cordate. Corolla lobes purplish. Hoods with a long lanceolate appendage at the summit.—Pr. Prov.

3. *A. phytolacoi'des* Pursh. (POKE MILKWEED.) Stem tall and smooth. Leaves broadly ovate, acute at both ends, short-petioled. Pedicels loose and nodding, very long and slender. Corolla greenish, with the hooded appendage white. Pods minutely downy, but not warty.—Moist thickets.

4. *A. ovalifo'lia* Decaisne. Low, soft-downy. Leaves ovate to lance-oblong, acute, short-petioled, soft-pubescent beneath. Umbels loosely 10-18-flowered. Pedicel slender. Corolla lobes greenish-white, slightly tinged with purple outside. Hoods yellowish, with a small horn, obtuse, entire.—Pr. Prov.

5. *A. quadrifo'lia* Jacq. Stem slender, 1-2 feet high, mostly leafless below, with one or two whorls of four in the middle, and one or two pairs of ovate or ovate-lanceolate, taper-pointed leaves. Corolla lobes pale pink; hoods white.—Not common, but abundant near Toronto.

6. *A. verticilla'ta* L. Slender stems and filiform-linear leaves with revolute margins, 3-6 in a whorl. Corolla lobes greenish-white.

B. Flowers red.

7. *A. incarna'ta* L. (SWAMP MILKWEED.) Stem tall, leafy, branching, and smooth. Leaves oblong-lanceolate, acute, obscurely heart-shaped at the base. Flowers rose-purple. Pods very smooth and glabrous.—Swamps and low grounds.

C. Flowers orange.

8. *A. tubero'sa* L. (BUTTERFLY-WEED. PLEURISY-ROOT.) Stem very leafy, branching above, rough-hairy. Leaves linear or oblong-

lanceolate, chiefly scattered. Corolla greenish-orange, with the hoods bright orange-red. Pods hoary. Almost destitute of milky juice.—Dry hillsides and fields.

2. ACERA'TES Ell. GREEN MILKWEED

1. *A. viridiflo'ra* Ell. Stems ascending, 1–2 feet high, minutely soft-downy, becoming smoothish. Leaves oval to oblong, often opposite. The compact umbels of greenish flowers nearly sessile, lateral, many-flowered.—Dry soil.

Var. *lanceola'ta* Gray. Lanceolate leaves.

Var. *linea'ris* Gray. Elongated-linear leaves, and low stems. Umbels often solitary.—Pr. Prov.

2. *A. florida'na* Hitchc. Usually minutely rough-hairy. Leaves usually alternate, linear to lanceolate. Umbels few, lateral or terminal on peduncles as long as the pedicels.

CONVOLVULA'CEAE (CONVOLVULUS FAMILY)

Chiefly twining or trailing herbs, with alternate leaves and regular flowers. Sepals 5, imbricated. Corolla 5-plaited or 5-lobed and convolute in the bud. Stamens 5. Ovary 2-celled.

Synopsis of the Genera

1. *Convol'vulus*. Leafy, green plants. Calyx inclosed in 2 large leafy bracts, or in one species without bracts. Corolla funnel-form, the border obscurely lobed. Pod 4-seeded. Stigmas 2, linear to oblong.
2. *Ipomoe'a*. Leafy green plants. Stigmas 2-lobed or entire.
3. *Cus'cuta*. Leafless, parasitic, slender twiners with yellowish or reddish stems, attaching themselves to the bark of other plants. Flowers small, mostly white, clustered. Corolla bell-shaped. Stamens with a fringed appendage at their base.

1. CONVOLVULUS L. BINDWEED

1. *C. se'pium* L. (HEDGE BINDWEED.) Stem mostly twining. Leaves halberd-shaped. Peduncles 4-angled. Corolla commonly rose-coloured.—Moist banks.

2. *C. spithamae'us* L. Stem soft-pubescent, low and simple, upright or ascending, not twining, 6–12 inches high. Leaves oblong, more or less heart-shaped at the base. Corolla white.—Dry soil.

3. *C. arven'sis* L. (BINDWEED.) Stem twining or procumbent and low, propagating by running rootstocks. Leaves ovate-oblong, sagittate, the lobes acute. Corolla white, or tinged with red.

2. IPOMOE'A L. MORNING GLORY

1. *I. pandura'ta* G.F.W. Mey. (WILD POTATO VINE.) Stems long and stout, from a large root. Leaves sometimes fiddle-shaped. Penduncles 1-5-flowered. Corolla funnel-form, white, tinged on the inside with purple.

3. CUS'CUTA L. DODDER

1. *C. Grono'vii* Willd. Stems resembling coarse threads, spreading themselves over herbs and low bushes. Corolla lobes obtuse, spreading. Capsule globose, abruptly pointed. Flowers in loose paniced cymes.—Wet shady places.

2. *C. arven'sis* Beyrich. Stems pale and slender, low. Flowers in dense clusters. Capsule depressed-globose. Corolla lobes acute, with inflexed points. Stamen scales deeply fringed.—Dry ridges, Pr. Prov.

3. *C. Epil'inum*. Weihe. (FLAX DODDER.) Stems very slender, low. Flowers sessile, in dense scattered heads. Corolla short-cylindrical, hardly exceeding the calyx, persistent around the capsule. Stamen scales short and broad. Capsule globose, circumscissile.—Atl. Prov. (Introd.)

POLEMONIA'CEAE (POLEMONIUM FAMILY)

Herbs with regular, pentamerous and pentandrous flowers, but a 3-celled ovary and 3-lobed style. Lobes of the corolla convolute in the bud. Calyx persistent.

Synopsis of the Genera

1. *Phlox*. Corolla salver-form. Leaves opposite, entire.
2. *Gil'ia*. Corolla tubular-funnel-form or salver-shaped, very slender. Leaves alternate, entire.

1. PHLOX L. PHLOX

1. *P. divarica'ta* L. Corolla salver-shaped, with a long tube. Stamens short, unequally inserted. Stem ascending from a prostrate base, somewhat clammy. Leaves oblong-ovate. Flowers

lilac or bluish, in a spreading, loosely flowered cyme. Lobes of the corolla mostly obcordate.—Moist rocky woods.

2. *P. pilo'sa* L. Stem nearly erect, not prostrate. Leaves lanceolate or linear, tapering to a sharp point. Lobes of the pink-purple corolla obovate, entire.

3. *P. subula'ta* L. The Moss Pink of the gardens. Escaped from cultivation in some places. Stem creeping and tufted in broad mats. Leaves mostly linear-lanceolate. Flowers mostly rose-coloured, in cymes.—Dry grounds.

4. *P. Hood'ii* Richards. Forms broad dense mats or tufts, 2–4 inches high. Leaves awl-shaped. Flowers white, sessile and solitary at the ends of the branches.

2. GIL'IA Ruiz and Pavon

1. *G. linea'ris* Gray. A branching herb with alternate, linear-lanceolate or oblong, sessile and entire leaves. Corolla salver-form, with stamens unequally inserted in its narrow tube, lilac-purple to nearly white. Ovules solitary.

HYDROPHYLLA'CEAE (WATERLEAF FAMILY)

Herbs with alternate, cut-toothed or lobed leaves, and regular, pentamerous and pentandrous flowers, having a 1-celled ovary with the seeds on the walls (parietal). Style 2-cleft. Flowers mostly in 1-sided cymes which uncoil from the apex.

Synopsis of the Genera

1. *Hydrophyl'lum*. Stamens exserted. Calyx unchanged in fruit.
2. *Ellis'ia*. Stamens included. Calyx enlarged in fruit.

1. HYDROPHYL'LUM L. WATERLEAF

1. *H. virginia'num* L. Corolla bell-shaped, the 5 lobes convolute in the bud; the tube with 5 folds down the inside, one opposite each lobe. Stamens and style exserted, the filaments bearded below. Stem smoothish. Leaves pinnately cleft into 5–7 divisions, the latter ovate-lanceolate, pointed, cut-toothed. Calyx lobes very narrow, bristly-ciliate. Flowers white or pale blue. Peduncles longer than the petioles of the upper leaves. Rootstocks scaly-toothed.—Moist woods.

2. *H. canadense* L. Differs from the last in having the leaves palmately 5-7-lobed, and rounded; the peduncles shorter than the petioles; the calyx lobes nearly smooth. Flowers whitish.—Rich woods.

3. *H. appendiculatum* Michx. Stem, pedicels, and calyx hairy. Stem-leaves palmately 5-lobed and rounded, the lowest leaves pinnately divided. Calyx with a small reflexed appendage in each sinus; corolla violet or purple. Stamens sometimes not exerted.—Rich woods.

2. ELLISTA L.

1. *E. Nyctelea* L. A delicate branching annual, minutely or sparingly roughish-hairy. Leaves pinnately parted into 7-13 narrow, sparingly cut-toothed divisions. Peduncles 1-flowered, solitary in the forks or opposite the leaves. Flowers small, whitish. Capsule pendulous.—Pr. Prov.

BORAGINACEAE (BORAGE FAMILY)

Herbs with alternate entire leaves and a deeply 4-lobed ovary, forming 4 seed-like nutlets, but the corolla is regularly 5-lobed, with 5 stamens inserted upon its tube.

Synopsis of the Genera

A. *Corolla without any scales in the throat.*

1. *Echium*. Corolla with a funnel-form tube and a spreading border of 5 somewhat unequal lobes. Stamens exerted, unequal. Flowers bright blue, with a purplish tinge, rarely whitish, in racemed clusters. Plant bristly.

B. *Corolla with 5 scales completely closing the throat.*

2. *Symphytum*. Corolla tubular-funnel-form with short spreading lobes; scales awl-shaped. Flowers yellowish-white, in nodding, raceme-like clusters, the latter often in pairs. Nutlets nearly smooth. Coarse hairy herbs.
3. *Lappula*. Nutlets prickly on the margin. Corolla salver-shaped, lobes rounded; scales short and blunt. Flowers blue, small, in leafy-bracted racemes. Plant rough-hairy.
4. *Cynoglossum*. Nutlets prickly all over. Corolla funnel-form; scales blunt. Flowers red-purple or pale blue, in racemes which are naked above, but usually leafy-bracted below. Strong-scented, coarse herbs.

5. *Lycop'sis*. Corolla slightly irregular funnel-shaped, the scales in the throat bristly. Flowers blue, in leafy, raceme-like clusters. Nutlets rough-bristly, as is the whole plant.

C. Corolla open, the scales or folds not sufficient to completely close the throat.

6. *Merten'sia*. Corolla bell-funnel-shaped, mostly with 5 small glandular folds in the open throat. Calyx deeply 5-cleft. Style long and thread-form. Nutlets smooth or wrinkled. Flowers purplish-blue or white, in loose and short, raceme-like clusters, only the lower one leafy-bracted. Pedicels slender.

7. *Onosmo'dium*. Corolla tubular, the 5 lobes acute and erect or converging. Anthers mucronate; filaments very short. Style thread-form, much exserted. Flowers greenish- or yellowish-white. Rather tall, stout plants, shaggy with spreading bristly hairs, or rough with short appressed bristles. Nutlets smooth.

8. *Lithosper'mum*. Corolla funnel-form or salver-shaped, the 5 lobes of the spreading limb rounded. Anthers almost sessile. Root mostly red. Flowers small and almost white, or large and deep yellow, scattered, spiked and leafy-bracted. Nutlets smooth or wrinkled.

9. *Myoso'tis*. Corolla salver-shaped, with a very short tube, the lobes convolute in the bud; scales or appendages of the throat blunt and arching. Flowers blue, in (so-called) racemes without bracts. Low plants, mostly soft-hairy. Nutlets smooth.

1. E'CHIUM L. VIPER'S BUGLOSS

1. *E. vulga're* L. (BLUE-WEED.) Stem erect, 2 feet high. Leaves sessile, linear-lanceolate. Flowers showy, in lateral clusters, the whole forming a long narrow raceme.—Roadsides.

2. SYM'PHYTUM L. COMFREY

1. *S. officina'le* L. (COMMON COMFREY.) Stem winged above by the decurrent bases of the leaves, branched. Leaves ovate-lanceolate or lanceolate.—Moist soil; escaped from gardens.

3. LAP'PULA Moench. STICKSEED

1. *L. echina'ta* Gilibert. A very common roadside weed, 1-2 feet high, hispid, branching above. Leaves lanceolate, rough. Nutlets warty on the back, with a double row of prickles on the margin. Pedicels stout, not deflexed. Calyx becoming foliaceous.

2. *L. Redow'skii* Greene, var. *occidentalis* Rydb. Erect, hispid, 1-2 feet high. Leaves linear-lanceolate. Calyx becoming leaf-like, pedicels not deflexed; nutlets bearing a single row of stout flattened prickles.

3. *L. floribunda* Greene. Stem strict, 2 feet high or more, not hispid. Leaves oblong- to linear-lanceolate, the lowest tapering into margined petioles. Racemes numerous, rather strict. Pedicels slender, deflexed in fruit. Nutlets rough and margined with a close row of flat, awl-shaped prickles.—Chiefly Pr. Prov.

4. *L. virginiana* Greene. (BEGGAR'S LICE.) Stem 2-4 feet high, not hispid. Lower leaves round-ovate or cordate, slender-petioled; upper ones tapering at both ends. Racemes widely spreading. Pedicels slender, deflexed in fruit. Nutlets prickly over the whole back.—Open woods and thickets.

5. *L. deflexa* Garcke, var. *americana* Greene. Widely branched, 1-3 feet high. Leaves oblong to lanceolate. Flowers small, in loose racemes; pedicels deflexed in fruit. Nutlets with a few prickles along the middle.—Pr. Prov.

4. CYNOGLOSSUM L. HOUND'S TONGUE

1. *C. officinale* L. (COMMON HOUND'S TONGUE.) Flowers red-purple. Upper leaves lanceolate, sessile. Stem soft-pubescent. Nutlets rather flat.—A common weed in fields and along roadsides.

2. *C. virginicum* L. (WILD COMFREY.) Flowers pale blue. Stem roughish with spreading hairs. Leaves few, lanceolate-oblong, clasping. Racemes corymbed, raised on a long naked peduncle.—Rich woods.

5. LYCOPHIS L. BUGLOSS

1. *L. arvensis* L. (SMALL BUGLOSS.) Very rough-bristly, 1 foot high. Leaves lanceolate.—Dry or sandy fields.

6. MERTENSIA Roth. LUNGWORT

1. *M. paniculata* Don. Filaments broad and short. Nutlets wrinkled when dry. Plant more or less hairy, erect, loosely branched. Leaves ovate to ovate-lanceolate, ribbed.—Pr. Prov.

2. *M. maritima* S. F. Gray. (SEA LUNGWORT.) Corolla white, trumpet-shaped, conspicuously 5-lobed, throat crested. Leaves fleshy, glaucous, ovate to spatulate. Stems spreading, often decumbent. Filaments longer and narrower than the anthers. Nutlets shining.—Atl. sea-coast.

7. ONOSMO'DIUM Michx. FALSE GROMWELL

1. *O. hispidis'simum* Mackenzie. Stem stout, 3-4 feet high. Leaves ovate-lanceolate, acute. Lobes of the corolla ovate-triangular, very hairy outside.—Banks of streams.

2. *O. virgina'num* A. DC. Stem slender, 1-2 feet high. Leaves narrowly oblong. Lobes of the corolla lance-awl-shaped, sparingly bearded outside with long bristles.—Banks and hillsides; not common.

8. LITHOSPER'MUM L. GROMWELL. PUCCOON

A. Corolla almost white. Nutlets wrinkled, gray.

1. *L. arven'se* L. (CORN GROMWELL.) Stem 6-12 inches high, erect. Leaves lanceolate or linear.—Waste grounds and in wheat-fields.

B. Corolla deep yellow. Nutlets smooth and shining.

2. *L. Gmeli'ni* Hitchc. (HAIRY PUCCOON.) Stem 1-2 feet high, hispid. Stem-leaves lanceolate or linear; those of the flowering branches ovate-oblong, ciliate. Flowers peduncled. Corolla woolly at the base inside.—Dry woods.

3. *L. canes'cens* Lehm. (HOARY PUCCOON. ALKANET.) Stem 6-15 inches high, soft-hairy. Leaves linear-oblong, obtuse. Corolla naked at the base inside. Flowers sessile. Limb of the corolla smaller, and the calyx shorter, than in No. 2.—Open woods and plains.

4. *L. angustifo'lium* Michx. Erect or diffusely branched from the base, somewhat hoary. Leaves all linear. Flowers pedicelled, leafy-bracted, of two sorts: the earlier large and conspicuous, the later small and pale.—Pr. Prov.

C. Corolla greenish-white or cream-coloured. Nutlets smooth and shining, mostly white.

5. *L. officina'le* L. (COMMON GROMWELL.) Much-branched above. Leaves broadly lanceolate, acute. Corolla exceeding the calyx.—Roadsides and fields.

6. *L. latifo'lium* Michx. Loosely branched above. Leaves ovate and ovate-lanceolate, mostly taper-pointed. Corolla shorter than the calyx.—Borders of woods.

9. MYOSO'TIS L. FORGET-ME-NOT

1. *M. scorpioi'des* L. (FORGET-ME-NOT.) Stem ascending from a creeping base, about a foot high, smoothish, loosely branched. Leaves rough-pubescent. Calyx open in fruit, its lobes much shorter than its tube. Corolla sky-blue, with a yellow eye. Pedicels spreading.—Wet places.

2. *M. laxa* Lehm. Stem very slender, decumbent, from thread-like, subterranean shoots. Pubescence all appressed. Calyx open in fruit, its lobes as long as its tube. Corolla paler blue.—Wet places.

3. *M. arven'sis* Hill. Stem erect or ascending, hirsute. Calyx closing in fruit. Corolla blue, rarely white. Pedicels spreading in fruit and longer than the 5-cleft, equal calyx. Racemes naked at the base.—Fields.

4. *M. virgin'ica* BSP. Differs from the last in having a very small, white corolla, pedicels erect in fruit, and the racemes leafy at the base. The calyx, also, is unequally 5-toothed, somewhat 2-lipped and hispid.—Dry hills.

VERBENA'CEAE (VERVAIN FAMILY)

Herbs with opposite leaves, didynamous stamens, and corolla irregularly 5-lobed, and salver-form. Ovary 4-celled (when ripe splitting into 4 nutlets), but not 4-lobed.

1. VERBE'NA L. VERVAIN

1. *V. hasta'ta* L. (BLUE VERVAIN.) Stem 3–5 feet high. Leaves oblong-lanceolate, taper-pointed, serrate. Spikes of purple flowers dense, erect, corymbed or paniced.—Low meadows and fields.

2. *V. urticaefo'lia* L. (NETTLE-LEAVED VERVAIN.) Stem tall. Leaves oblong-ovate, acute, coarsely serrate. Spikes of small white flowers very slender, loosely paniced.—Fields and roadsides.

3. *V. angustifo'lia* Michx. Stem low. Leaves narrowly lanceolate, tapering at the base, sessile, roughish, slightly toothed. Flowers purple, in a crowded spike.—Dry soil.

4. *V. bracteo'sa* Michx. Stem spreading or procumbent, hairy. Leaves wedge-lanceolate, cut-pinnatifid or 3-cleft. Spikes single, thick, sessile, leafy-bracted, the bracts longer than the small purple flowers.

LABIATAE (MINT. FAMILY)

Herbs with square stems, opposite leaves (mostly aromatic), didynamous (or in one or two genera diandrous) stamens, a 2-lipped or irregularly 4- or 5-lobed corolla, and a deeply 4-lobed ovary, forming in fruit 4 nutlets or achenes.

Synopsis of the Genera

A. *Stamens 4, curved upwards, parallel, exserted from a deep notch on the upper side of the 5-lobed corolla.*

1. **Teu'crium.** Calyx 5-toothed. The four upper lobes of the corolla nearly equal, with a deep notch between the upper 2; the lower lobe much larger. Flowers pale purple.
2. **Isan'thus.** Calyx bell-shaped, 5-cleft, almost equalling the small, pale-blue corolla. Lobes of the corolla almost equally spreading. Stamens only slightly exserted.

B. *Stamens 4, the outer or lower pair longer, or only 2 with anthers, straight and not converging in pairs. Anthers 2-celled.*

(1) *Corolla almost equally 4-lobed, quite small.*

3. **Men'tha.** Calyx equally 5-toothed. Upper lobe of the corolla rather the broadest, and sometimes notched. Stamens 4, of equal length, not convergent. Flowers either in terminal spikes or in head-like, whorled clusters, often forming interrupted spikes. Corolla purplish or whitish.
4. **Lyc'opus.** Calyx teeth 4 or 5. Stamens 2, the upper pair, if any, without anthers. Flowers white, in dense axillary clusters.

(2) *Corolla evidently 2-lipped, but the lobes nearly equal in size; the tube not bearded inside. Stamens with anthers 2.*

5. **Hedeo'ma.** Calyx 2-lipped, bulging on the lower side of the base, hairy in the throat; 2 stamens with good anthers, and 2 sterile filaments with false anthers. Low odorous plants, with bluish flowers in loose axillary clusters.

(3) *Corolla 2-lipped, the lower of the 5 lobes much larger than the other 4; the tube with a bearded ring inside. Stamens 2 (occasionally 4), much exserted.*

6. **Collinso'nia.** Calyx ovate, enlarged and turned down in fruit, 2-lipped. Corolla elongated, the lower lip toothed or fringed. Strong-scented plants with yellowish flowers on slender pedicels in terminal paniced racemes.

(4) *Corolla evidently 2-lipped. Stamens with anthers 4.*

7. **Hysso'pus.** Calyx tubular, 15-nerved, equally 5-toothed. Corolla blue-purple, short; upper lip erect, flat, obscurely notched, the lower 3-cleft, with the middle lobe larger and 2-cleft. Stamens exserted, diverging. Flowers in small clusters, crowded in a spike. Branches simple, wand-like.
8. **Pycnan'themum.** Calyx short-tubular, 10-13-nerved, equally 5-toothed. The whitish or purplish flowers in small dense heads, forming terminal corymbs. Aromatic plants, with narrow rigid leaves crowded and clustered in the axils.
9. **Sature'ja.** Calyx bell-shaped, or tubular. Aromatic plants, with narrow leaves and purplish, spiked flowers. Some species have anthers approximate in pairs.

C. *Stamens only 2, parallel; the anthers only 1-celled. Corolla 2-lipped.*

10. **Monar'da.** Calyx tubular, nearly equally 5-toothed, hairy in the throat. Corolla elongated, strongly 2-lipped, the upper lip narrow. Stamens with long protruding filaments, each bearing a linear anther on its apex. Flowers large, in whorled heads surrounded by bracts.
11. **Blephil'ia.** Resembles Monarda, but the calyx is 2-lipped, the upper lip 3-toothed, the lower 2-toothed. Flowers bluish-purple in dense globular whorls.

D. *Stamens 4, the upper or inner pair longer. Anthers approximate in pairs (except in Agastache). Corolla 2-lipped.*

12. **Nep'eta.** Calyx obliquely 5-toothed. Anthers approaching each other in pairs under the inner lip of the corolla, the cells of each anther divergent.
13. **Dracoceph'alum.** Calyx straight, 5-toothed, the upper tooth much the largest. Whorls of bluish flowers with awn-toothed or fringed, leafy bracts, in a crowded head or spike.
14. **Agas'tache.** Calyx obliquely 5-toothed. Stamens exserted, the upper pair declined, the lower ascending, so that the pairs cross. Anther-cells parallel. Tall herbs with small flowers in interrupted terminal spikes.

E. *Stamens 4, the lower or outer pair longer (except in Leonurus). Anthers approximate in pairs. Corolla 2-lipped.*

15. **Melis'sa.** Calyx with the upper lip flattened and 3-toothed, the lower 2-cleft. Corolla nearly white, the tube recurved-ascending. Stamens curved and conniving under the upper lip. Flower-cluster loose, few-flowered, one-sided, with few bracts resembling the leaves.

16. **Physoste'gia.** Calyx not 2-lipped, 5-toothed or lobed, thin and membranaceous, inflated-bell-shaped in fruit. Anther-cells parallel. Flowers large and showy, rose-coloured variegated with purple, opposite, in terminal leafless spikes.
17. **Prunel'la.** Calyx 2-lipped, flat on the upper side, closed in fruit; the upper lip 3-toothed, the lower 2-cleft. Filaments 2-toothed at the apex, the lower tooth bearing the anther. Flowers violet, in a close terminal spike or head, which is very leafy-bracted.
18. **Scutella'ria.** Calyx 2-lipped, short, closed in fruit, the lips rounded and entire, the upper with a projection on the back. Corolla blue or violet, the tube elongated and somewhat curved. Anthers of the lower stamens 1-celled, of the upper 2-celled. Flowers solitary in the axils of the upper leaves, or in axillary or terminal, 1-sided racemes.
19. **Marru'bium.** Calyx 10-toothed, the teeth spiny and recurved after flowering. Stamens 4, included in the corolla tube. Whitish woolly plants with small white flowers in head-like whorls.
20. **Galeop'sis.** Calyx 5-toothed, the teeth spiny. The middle lobe of the lower lip of the corolla inversely heart-shaped, the palate with 2 teeth at the sinuses. Stamens 4, the anthers opening cross-wise. Flowers purplish, in axillary whorls.
21. **Sta'chys.** Calyx 5-toothed, beset with stiff hairs, the teeth spiny, diverging in fruit. Stamens 4, the outer pair turned down after discharging their pollen. Flowers purple, crowded in whorls, these at length forming an interrupted spike.
22. **Leonu'rus.** Calyx 5-toothed, the teeth spiny, and spreading when old. The middle lobe of the lower lip of the corolla narrowly oblong-obovate, entire. Flowers pale purple, in close whorls in the axils of the cut-lobed leaves. Nutlets sharply 3-angled.
23. **La'mium.** Calyx tubular-bell-shaped, 5-nerved, with 5 nearly equal, awl-pointed teeth. Corolla dilated at the throat; upper lip arched and narrowed at the base, the middle lobe of the lower lip notched at the apex and narrowed at the base. Herbs with purple flowers in few or several whorls or heads.
24. **Ballo'ta.** Calyx nearly funnel-form, 10-ribbed, with a spreading, 5-toothed border. Whorls of purplish flowers dense. Anthers exserted. Plant erect, hairy (but green).

1. TEU'CRUM L. GERMANDER

1. **T. canadense** L. (AMERICAN GERMANDER. WOOD SAGE.)
 Stem 1-3 feet high, downy. Leaves ovate-lanceolate, serrate,

short-petioled, hoary beneath. Flowers in a long spike.—Low grounds.

2. *T. occidenta'le* Gr. Loosely pubescent. Calyx villous with viscid hairs.

2. ISAN'THUS Michx. FALSE PENNYROYAL

1. *I. brachia'tus* BSP. A low, branching, clammy-pubescent annual. Leaves lance-oblong, 3-nerved, nearly entire. Peduncles axillary, 1-3-flowered.—Gravelly soil.

3. MEN'THA L. MINT

1. *M. spica'ta* L. (SPEARMINT.) Flowers in a narrow terminal spike. Leaves ovate-lanceolate, wrinkled, veiny, unequally serrate, usually sessile.—Wet places.

2. *M. piperi'ta* L. (PEPPERMINT.) Flowers in loose interrupted spikes. Leaves ovate or ovate-oblong, acute, petioled. Plants smooth.—Wet places.

3. *M. arven'sis* L. Flowers in globular clusters in the axils of the leaves, the uppermost without flowers. Leaves oblong to ovate with rounded bases, pubescent, closely serrate.—Atl. Prov.

Var. *canaden'sis* Briquet. The lanceolate to oblong-lanceolate leaves are more pubescent.

Var. *glabra'ta* Fernald. Smoothish, with a rather pleasanter odour.

4. *M. genti'lis* L. (WHORLED MINT.) Flowers in globular clusters in the axils of leaves; the uppermost axils not flower-bearing. Leaves petioled, ovate, sharply serrate. Calyx with very slender teeth.—Atl. Prov.

4. LYC'OPUS L. WATER HOREHOUND

1. *L. virgin'icus* L. (BUGLE-WEED.) Calyx teeth 4, bluntish. Stems obtusely 4-angled, 6-18 inches high, producing slender runners from the base. Leaves ovate-lanceolate, toothed.—Moist places.

2. *L. america'nus* Muhl. Calyx teeth 5, sharp-pointed. Stem sharply 4-angled, 1-3 feet high. Leaves varying from cut-toothed to pinnatifid.—Wet places.

3. *L. uniflorus* Michx. Resembles No. 1, but is generally more slender and smooth. Rootstocks and stolons tuberous. Leaves lanceolate to oblong, narrowed at both ends. Corolla lobes wide-spreading.

4. *L. rubellus* Moench. Also resembles No. 1 but has long slender calyx teeth.

5. *L. lucidus* Turcz., var. *americanus* Gray. Calyx teeth 5, very acute. Corolla hardly exceeding the calyx. Stem strict, short, 2-3 feet high. Leaves lanceolate or oblong-lanceolate, sessile or nearly so, sharply and coarsely serrate.—Pr. Prov.

5. HEDEOMA Pers. MOCK PENNYROYAL

1. *H. pulegioides* Pers. (AMERICAN PENNYROYAL). Stem 5-8 inches high, branching, hairy. Leaves oblong-ovate, petioled, obscurely serrate. Whorls few-flowered. Plant with a pungent aromatic odour.—Open woods and fields.

2. *H. hispida* Pursh. Leaves sessile, linear, and entire. Calyx ciliate and hispid.—Not common.

6. COLLINSONIA L. HORSE BALM

1. *C. canadensis* L. (RICHWEED. STONE-ROOT.) Stem smooth or nearly so, 1-3 feet high. Leaves serrate, pointed, petioled, 3-6 inches long.—Rich woods.

7. HYSSOPUS L. HYSSOP

1. *H. officinalis* L. Escaped from gardens in a few localities. Leaves lanceolate or linear, entire.

8. PYCNANTHEMUM Michx. MOUNTAIN MINT. BASIL

1. *P. virginicum* Durand and Jackson. Stem 2 feet high, smoothish or minutely pubescent. Leaves lanceolate or lance-linear, entire. Heads downy. Calyx teeth short. Lips of corolla very short.—Dry soil.

2. *P. pilosum* Nutt. Hoary with loose pubescence. Leaves oblong-lanceolate, denticulate. Bracts and calyx teeth villous-pubescent. Heads larger than in No. 1.

9. SATURJA L. SAVORY

1. *S. hortensis* L. (SUMMER SAVORY.) Stem pubescent. Clusters few-flowered.—Escaped from gardens in a few localities.

2. *S. vulga'ris* Fritsch. (BASIL.) Stem hairy, erect, 1-2 feet high. Flower-clusters dense. Leaves ovate, nearly entire, petioled.—Thickets.

3. *S. gla'bra* Fernald. Smooth, 5-9 inches high. Leaves narrowly oblong. Clusters few-flowered, the flowers on slender naked pedicels. Bracts linear or oblong.—Wet limestone rocks.

10. MONAR'DA L. HORSE MINT

1. *M. did'yma* L. (OSWEGO TEA.) Corolla bright red, very showy. The large outer bracts tinged with red.—Along shaded streams.

2. *M. fistulo'sa* L. (WILD BERGAMOT.) Corolla purplish, the outer bracts somewhat purplish.—Dry and rocky banks and woods.

3. *M. mol'lis* L. Flesh-coloured or lilac corolla.—Common in the Pr. Prov.

11. BLEPHIL'IA Raf.

1. *B. cilia'ta* Raf. Finely pubescent. Leaves oblong-ovate, short-petioled, whitish beneath. Corolla hairy.

12. NEP'ETA L. CAT MINT

1. *N. Cata'ria* L. (CATNIP.) Flowers in cymose clusters. Stem erect, downy, branching. Leaves oblong, crenate, whitish beneath. Corolla whitish dotted with purple.—Roadsides.

2. *N. hедера'cea* Trevisan. (GROUND IVY.) Creeping and trailing. Leaves round-kidney-shaped, crenate, green both sides. Corolla light blue.—Damp waste grounds.

13. DRACOCEPH'ALUM L. DRAGON-HEAD

1. *D. parviflo'rum* Nutt. Stem erect, 8-20 inches high, leafy. Leaves ovate-lanceolate, cut-toothed, petioled. Corolla bluish, small and slender.

14. AGAS'TACHE Clayt. GIANT HYSSOP

1. *A. nepetoi'des* Ktze. Smooth or nearly so. Leaves coarsely crenate-toothed. Calyx teeth ovate, rather obtuse. Corolla greenish-yellow.—Borders of woods.

2. *A. scrophulariaefo'lia* Ktze. Calyx teeth lanceolate. Corolla purplish.

3. *A. Foenic'ulum* Ktze. Leaves ovate, acute, and glaucous-white underneath.—Pr. Prov.

15. MELIS'SA L. BALM

1. *M. officina'lis* L. (COMMON BALM.) Upright, branching, pubescent. Leaves broadly ovate, crenate-toothed, lemon-scented.—Escaped from gardens in a few places.

16. PHYSOSTE'GIA Benth. FALSE DRAGON-HEAD

1. *P. virginia'na* Benth. Stem smooth, wand-like. Lower leaves oblong-ovate, upper lanceolate. Corolla an inch long, funnel-form, the throat inflated; upper lip slightly arching, the lower 3-parted, spreading, small.—Wet banks.

2. *P. parviflo'ra* Nutt. Stem 1-2 feet high, simple below. Spikes dense; calyx teeth blunt.

17. PRUNEL'LA L. SELF-HEAL

1. *P. vulga'ris* L. (COMMON HEAL-ALL.) A low plant with oblong-ovate, petioled leaves. Clusters 3-flowered, the whole forming a close, terminal, elongated head.—Woods and fields everywhere.

18. SCUTELLA'RIA L. SKULL-CAP

1. *S. galericula'ta* L. Flowers blue, $\frac{3}{4}$ inch long, solitary in the axils of the upper leaves. Stem nearly smooth, 1-2 feet high.—Wet places.

2. *S. par'vula* Michx. Flowers blue, $\frac{1}{4}$ inch long, solitary in the upper axils. Stem minutely downy, 2-6 inches high. Lowest leaves round-ovate, the upper narrower, mostly entire. Stolon necklace-form.—Dry banks.

3. *S. lateriflo'ra* L. Flowers blue, $\frac{1}{3}$ inch long, in 1-sided racemes. Stem upright, much-branched, 1-2 feet high.—Wet places.

19. MARRU'BIUM L. HOREHOUND

1. *M. vulga're* L. Leaves round-ovate, crenate-toothed. Calyx with 5 long and 5 short teeth, recurved.—Escaped from gardens in some places.

20. GALEOP'SIS L. HEMP NETTLE

1. *G. Tetra'hit* L. (COMMON HEMP NETTLE.) Stem bristly-hairy, swollen below the joints. Leaves ovate, coarsely serrate. Corolla often with a purple spot on the lower lip.—Waste places and fields.

21. STA'CHYS L. HEDGE NETTLE

1. *S. palus'tris* L. Stem 2-3 feet high, 4-angled, the angles beset with stiff reflexed hairs or bristles. Leaves sessile, or the lower short-petioled, oblong or ovate-lanceolate, crenately serrate, downy. Calyx hispid. Upper lip of the corolla pubescent.—Wet grounds.

2. *S. tenuifo'lia* Willd., var. *aspera* Fernald. Taller than the last, the leaves nearly all distinctly petioled. Calyx mostly glabrous. Corolla glabrous throughout.—Wet grounds.

22. LEONU'RUS L. MOTHERWORT

1. *L. Cardi'aca* L. (COMMON MOTHERWORT.) Stem tall. Leaves long-petioled, the lower palmately lobed, the upper 3-cleft. Upper lip of the corolla bearded.—Near dwellings.

23. LA'MIUM L. DEAD NETTLE

1. *L. amplexicaul'e* L. Leaves rounded, deeply crenate-toothed, the upper ones clasping.—In gardens in some places.

2. *L. purpu'reum* L. Leaves rounded or oblong, heart-shaped, crenate-toothed, all petioled.—A garden escape.

3. *L. macula'tum* L. Taller. Leaves ovate-heart-shaped, often with a white spot above. Flowers rather large, purplish, with hairs at the base inside.—A garden escape.

24. BALLO'TA L. FETID HOREHOUND

1. *B. ni'gra* L. (BLACK HOREHOUND.) Leaves ovate, toothed. Calyx teeth longer than the tube of the purplish corolla.

SOLANA'CEAE (NIGHTSHADE FAMILY)

Rank-scented herbs (or one species shrubby) with colourless bitter juice, alternate leaves, and regular, pentamerous and pentandrous flowers, but a 2-celled (in *Nicandra* 3-5-celled) ovary, with the placentae in the axis. Fruit a many-seeded berry or pod.

Synopsis of the Genera

1. *Sola'num*. Corolla wheel-shaped, 5-lobed, the margins turned inwards in the bud. Anthers conniving around the style, the cells opening by pores at the apex; filaments very short. The larger leaves often with an accompanying smaller one. Fruit a berry.

2. *Phys'alis*. Calyx 5-cleft, enlarging after flowering, becoming at length much-inflated, and inclosing the berry. Corolla between wheel-shaped and funnel-form. Anthers separate, opening lengthwise. Plant clammy-pubescent.
3. *Nican'dra*. Calyx 5-parted, 5-angled, the divisions rather arrow-shaped, enlarged and bladder-like in fruit, inclosing the 3-5-celled, globular, dry berry. A smooth herb, 2-3 feet high, with pale-blue flowers.
4. *Lyc'ium*. Corolla funnel-form or tubular. Fruit a small berry, the calyx persistent but not inflated. A shrubby plant with long drooping branches and greenish-purple flowers on slender peduncles fasciated in the axils.
5. *Hyoscy'amus*. Fruit a pod, the top coming off like a lid. Calyx urn-shaped, 5-lobed, persistent. Corolla funnel-form, oblique, the limb 5-lobed, dull-coloured and veiny. Plant clammy-pubescent.
6. *Datu'ra*. Fruit a large, prickly, naked pod. Calyx long, 5-angled, not persistent. Corolla very large, funnel-form, strongly plaited in the bud, with 5 pointed lobes. Stigma 2-lipped. Rank-scented weeds, with the showy flowers in the forks of the branching stems.
7. *Nicotia'na*. Fruit a pod, inclosed in the calyx. Calyx tubular-bell-shaped, 5-cleft. Corolla dull greenish-yellow, funnel-form, plaited in the bud. Leaves large. Flowers racemed or paniced.

1. SOLA'NUM L. NIGHTSHADE

1. *S. Dulcama'ra* L. (BITTER-SWEET.) Stem somewhat shrubby and climbing. Leaves ovate and heart-shaped, the upper halberd-shaped, or with 2 ear-like lobes at the base. Flowers violet-purple, in small cymes. Berries red.—Near dwellings and in moist grounds.

2. *S. triflo'rum* Nutt. A low, spreading herb, only slightly, if at all, pubescent. Leaves pinnatifid with rounded angles between the lobes. Corolla white; berries green and globular.

3. *S. ni'grum* L. (COMMON NIGHTSHADE.) Stem low and spreading, branched. Leaves ovate, wavy-toothed. Flowers small, white, drooping in umbel-like, lateral clusters. Berries black.—Fields and damp grounds.

4. *S. rostra'tum* Dunal. A prickly herb with large yellow flowers and sharp anthers.

2. *PHYS'ALIS* L. GROUND CHERRY

1. *P. visco'sa* L. Corolla greenish-yellow, brownish in the centre. Anthers yellow. Leaves ovate or heart-shaped, mostly toothed. Berry orange, sticky.—Sandy soil.

2. *P. heterophyl'la* Nees. A much-branched perennial, usually viscid-hairy. Leaves oblong to ovate, rounded or cordate at the base. Anthers generally yellow.

3. *P. subglabra'ta* Mackenzie and Bush. Leaves ovate or ovate-oblong. Corolla brownish, spotted in the centre with violet, calyx persistent and globose at maturity. Berry large, reddish or purple.

4. *P. grandiflo'ra* Hook. Corolla white, large, with a woolly ring in the throat. Anthers tinged with blue or violet.

3. *NICAN'DRA* Adans. APPLE OF PERU

1. *N. Physalo'des* Pers. Leaves ovate, sinuate-toothed or angled. Flowers solitary on axillary and terminal peduncles.—Escaped from gardens in some places.

4. *LYC'IUM* L. MATRIMONY VINE

1. *L. halimifo'lium* Mill. Common about dwellings. Berry ovoid, orange-red.

5. *HYOSCY'AMUS* L. HENBANE

1. *H. ni'ger*, L. (BLACK HENBANE.) Escaped from gardens in some localities. Corolla dull yellowish, netted with purple veins. Leaves clasping, sinuate-toothed. A strong-scented and poisonous herb.

6. *DATU'RA* L. STRAMONIUM. THORN APPLE

1. *D. Stramo'nium*, L. (COMMON THORN APPLE.) Stem green. Corolla white, 3 inches long. Leaves ovate, sinuate-toothed.—Roadsides.

2. *D. Tat'ula* L. (PURPLE THORN APPLE.) Stem purple. Corolla pale violet-purple.

6. *NICOTIA'NA* L. TOBACCO

1. *N. rus'tica*, L. (WILD TOBACCO.) Old fields and in gardens.

SCROPHULARIA'CEAE (FIGWORT FAMILY)

Herbs distinguished by a 2-lipped or more or less irregular corolla, stamens usually 4 and didynamous, or only 2 (or in *Verbascum*, 5) and a 2-celled and usually many-seeded ovary. Style 1; stigma entire or 2-lobed.

Synopsis of the Genera

A. Corolla wheel-shaped, and only slightly irregular.

1. *Verbas'cum*. Stamens (with anthers) 5. Flowers in a long terminal spike. Corolla 5-parted, nearly regular. Filaments (or some of them) woolly.
2. *Veron'ica*. Stamens only 2; filaments long and slender. Corolla mostly 4-parted, nearly or quite regular. Pod flattish. Flowers solitary in the axils, or forming a terminal raceme or spike.

B. Corolla 2-lipped, or tubular and irregular.

- (1) *Upper lip of the corolla embracing the lower in the bud, except occasionally in Mimulus.*
3. *Lina'ria*. Corolla personate, with a spur beneath. Stamens 4. Flowers yellow, in a crowded raceme.
4. *Scrophula'ria*. Corolla tubular, somewhat inflated, 5-lobed; the 4 upper lobes erect, the lower one spreading. Stamens with anthers 4, the rudiment of a fifth in the form of a scale on the upper lip of the corolla. Flowers small and dingy, forming a narrow terminal panicle. Stem 4-sided.
5. *Collins'ia*. Corolla 2-cleft, the short tube saccate on the upper side; the middle lobe of the lower lip sac-like and inclosing the 4 declined stamens; the upper lip 2-cleft, the lobes partly turned backwards. Fifth stamen rudimentary. Leaves opposite. Flowers blue and white, in umbel-like clusters.
6. *Chelo'ne*. Corolla inflated-tubular. Stamens 4, with woolly filaments and anthers, and a fifth filament without an anther. Flowers white, in a close, terminal spike.
7. *Pentste'mon*. Corolla 2-lipped, gradually widening upwards. Stamens 4, with a fifth sterile filament, the latter yellow-bearded. Flowers white or purplish, in a loose panicle.
8. *Mim'ulus*. Calyx 5-angled and 5-toothed. Upper lip of the corolla erect or reflexed-spreading, the lower spreading, 3-lobed. Stamens 4, alike, no rudiment of a fifth. Stigma 2-lipped. Flowers blue or yellow, solitary on axillary peduncles.

9. *Grati'ola*. Corolla tubular and 2-lipped. Stamens with anthers only 2, included. Flowers with a yellowish tube, on axillary peduncles, solitary. Style dilated at the apex.
10. *Ilysan'thes*. Corolla tubular and 2-lipped. Stamens with anthers only 2, included; also a pair of filaments which are 2-lobed but without anthers. Flowers purplish, axillary. Style 2-lipped at the apex.

(2) *Lower lip of the corolla embracing the upper in the bud.*

11. *Gerar'dia*. Corolla funnel-form, swelling above, the 5 spreading lobes more or less unequal. Stamens 4, strongly didynamous, hairy. Style long, enlarged at the apex. Flowers purple or yellow, solitary on axillary peduncles, or sometimes forming a raceme.
12. *Castille'ja*. Corolla tubular and 2-lipped, its tube included in the tubular and flattened calyx; the upper lip long and narrow and flattened laterally, the lower short and 3-lobed. Stamens 4, didynamous. Floral leaves scarlet (rarely yellow) in our species. Corolla pale yellow.
13. *Orthocarp'us*. Corolla tubular and 2-lipped. Calyx tubular-campanulate, 4-cleft. Upper lip of the corolla much narrower than the inflated lower one. Flowers golden yellow, in a dense spike.
14. *Euphra'sia*. Calyx 4-cleft. Upper lip of the corolla erect, 2-lobed, the sides turned back; the lower spreading. Stamens 4, under the upper lip. Very small herbs, with whitish or bluish spiked flowers.
15. *Odonti'tes*. Calyx 4-cleft. Upper lip of the corolla entire, the sides not turned back. Flowers small, rose-red, in loose spikes. Small herbs with opposite sessile leaves.—Atl. sea-coast chiefly.
16. *Rhinan'thus*. Calyx flat, greatly inflated in fruit, 4-toothed. Upper lip of the corolla arched, flat, with a minute tooth on each side below the apex. Stamens 4. Flowers yellow, solitary in the axils, nearly sessile, the whole forming a crowded, 1-sided spike.
17. *Pedicula'ris*. Calyx split in front, not inflated in fruit. Corolla 2-lipped, the upper lip arched or hooded, incurved, flat, 2-toothed under the apex. Stamens 4. Pod flat, somewhat sword-shaped.
18. *Melampy'rum*. Calyx 4-cleft, the lobes sharp-pointed. Corolla greenish-yellow; upper lip arched, compressed, the lower 3-lobed at the apex. Stamens 4; anthers hairy. Pod 1-4-seeded, flat, oblique. Upper leaves larger than the lower ones and fringed with bristly teeth at the base.

1. VERBAS'CUM L. MULLEIN

1. *V. Thap'sus* L. (COMMON MULLEIN.) A tall and very woolly herb, with the simple stem winged by the decurrent bases of the leaves. Flowers yellow, forming a dense spike.—Fields and roadsides everywhere.

2. *V. Blatta'ria* L. (MOTH MULLEIN.) Stem slender, nearly smooth. Lower leaves petioled, doubly serrate; the upper partly clasping. Flowers whitish with a purple tinge, in a loose raceme. Filaments all violet-bearded.—Roadsides; not common northward.

3. *V. Lychni'tis* L. (WHITE MULLEIN.) Plant covered with a powdery woolliness. Leaves ovate, greenish above. Flowers yellow, rarely white, in a paniced cluster.

2. VERON'ICA L. SPEEDWELL

1. *V. america'na* Schweinitz. (AMERICAN BROOKLIME.) Flowers pale blue, in opposite axillary racemes. Leaves lanceolate to ovate, mostly petioled, thickish, serrate. Pod swollen.—A common plant in brooks and ditches.

2. *V. Anagal'lis-aquat'ica* L. (WATER SPEEDWELL.) Much like No. 1, but the leaves are sessile, with a heart-shaped base.

3. *V. scutella'ta* L. (MARSH SPEEDWELL.) Flowers pale blue, in racemes, chiefly from alternate axils. Leaves sessile, linear, opposite, hardly toothed. Racemes 1 or 2, slender and zig-zag. Flowers few. Pods very flat, notched at both ends.—Bogs.

Var. *villo'sa* Schumacher. Hairy stem and leaves.

4. *V. officina'lis* L. (COMMON SPEEDWELL.) Flowers light blue. Stem prostrate, rooting at base, pubescent. Leaves short-petioled, obovate-elliptical, serrate. Racemes dense, chiefly from alternate axils. Pod obovate-triangular, strongly flattened, notched.—Hillsides and open woods.

5. *V. Chamae'drys* L. Flowers pale blue. Stem pubescent, at least in 2 lines, from a creeping base. Leaves nearly sessile, ovate or cordate, incisely crenate. Racemes axillary, loosely flowered. Pod triangular-obcordate.

6. *V. serpyllifo'lia* L. (THYME-LEAVED SPEEDWELL.) Plant only 2 or 3 inches high. Flowers whitish or pale blue, in a loose terminal raceme. Stem nearly smooth, branched at the creeping base. Leaves obscurely crenate, the lowest petioled. Pod flat, notched.—Roadsides and fields.

7. *V. peregrina* L. (NECKWEED.) Flowers whitish, solitary in the axils of the upper leaves, short-pedicelled. Corolla shorter than the calyx. Stem 4-9 inches high, nearly smooth. Pod orbicular, slightly notched.—Waste places and cultivated grounds.

8. *V. arvensis* L. (CORN SPEEDWHEEL.) Flowers (blue) as in No. 7, but the stem is hairy, and the pod inversely heart-shaped.—Cultivated soil.

9. *V. agrestis* L. (FIELD SPEEDWELL.) Leaves round or ovate, crenate, petioled. Flowers small, in the axils of the ordinary leaves, long-pedicelled. Seeds few, cup-shaped.—Sandy fields, Atl. Prov.

10. *V. Tournefortii* C.C. Gmel. Leaves as in the last. Flowers (blue) large, long-pedicelled, in the axils of the ordinary leaves. Calyx lobes widely spreading in fruit. Pod obcordate-triangular, broadly notched, many-seeded.—Atl. Prov.

3. LINA'RIA Hill. TOAD FLAX

1. *L. vulgaris* Hill. (TOAD FLAX. BUTTER AND EGGS.) Leaves crowded, linear, pale green. Corolla pale yellow, with a deeper yellow or orange-coloured palate.—Roadsides.

2. *L. canadensis* Dumont. (WILD TOAD FLAX.) A slender herb, with linear, entire, alternate leaves. Flowers blue, small, in a naked terminal raceme. Spur of corolla curved, filiform.—Sandy soil.

3. *L. Cymbalaria* Mill. A delicate, little, trailing plant, with thickish, 3-5-lobed leaves on long petioles, and small, yellow and purple flowers.—A garden escape.

4. SCROPHULA'RIA L. FIGWORT

1. *S. marilandica* L. Stem smooth, 3-4 feet high. Leaves ovate or oblong, the upper lanceolate, serrate.—Damp thickets. (There appear to be two forms, one with broad leaves and greenish-brown flowers, and the other with narrow leaves and greenish-yellow flowers.)

2. *S. leporella* Bicknell. Closely resembles No. 1, but the flower-cluster is more slender and elongated, and the rudimentary stamen is yellowish-green.—Pr. Prov.

5. COLLIN'SIA Nutt.

1. *C. verna* Nutt. Slender, 6-20 inches high. Lower leaves ovate, upper ovate-lanceolate, clasping. Corolla (blue and white) twice as long as the calyx. Peduncles long.

2. *C. parviflora* Lindl. Smaller. Corolla (blue) very small, scarcely longer than the calyx. Peduncles short.—Chiefly Pr. Prov.

6. CHELO'NE L. TURTLE-HEAD

1. *C. gla'bra* L. Stem smooth, erect and branching. Leaves short-petioled, lance-oblong, serrate, opposite. Bracts of the flowers concave.—Wet places.

7. PENTSTE'MON Ait. BEARD-TONGUE

1. *P. hirsu'tus* Willd. Stem 1-3 feet high, pubescent; the panicle more or less clammy. Throat of the corolla almost closed. Stem-leaves lanceolate, clasping.—Dry soil.

2. *P. gra'cilis* Nutt. Stem 1 foot high or less, glabrous or nearly so, viscid-pubescent above. Corolla tubular-funnel-form, the throat open, lilac-purple or whitish.—Pr. Prov.

3. *P. acumina'tus* Dougl. Stem 6-20 inches high, stout, glabrous and glaucous. Leaves thick, the lower obovate or oblong, the upper lanceolate to broadly ovate or clasping. Panicle narrow, leafy below.—Pr. Prov.

4. *P. crista'tus* Nutt. Pubescent, and viscid-pubescent above. Leaves from linear-lanceolate to oblong. Corolla funnel-form, the lower lip long-villous within. Sterile filament copiously yellow-bearded.—Pr. Prov.

5. *P. al'bidus* Nutt. Stems several, usually erect. Leaves broadly linear. Flowers in a raceme-like cluster; corolla about 1 inch long, white, often tinged with purple, throat dilated; lobes of its limb roundish, large and widely spreading.—Pr. Prov.

6. *P. proc'erus* Dougl. Is at once recognized by the interrupted, spike-like panicle, which consists of 2 to 5 dense verticillate clusters. Corolla narrow, the lower lip conspicuously bearded within.—Pr. Prov.

8. MIM'ULUS L. MONKEY FLOWER

1. *M. rin'gens* L. Stem square, 1-2 feet high. Corolla blue, an inch long. (A white-flowered variety is sometimes met with.) Leaves oblong or lanceolate, clasping.—Wet places.

2. *M. glabra'tus* HBK., var. *Jame'sii* Gray. Stem creeping at the base. Corolla yellow, small. Leaves roundish or kidney-shaped, nearly sessile. Calyx inflated in fruit.—In cool springs.

9. GRATI'OLA L. HEDGE HYSSOP

1. *G. virginia'na* L. Stem 4–6 inches high, clammy with minute pubescence above. Sterile filaments minute or none. Leaves lanceolate. Peduncles slender.—Moist places.

2. *G. aur'ea* Muhl. Nearly glabrous. Sterile filaments slender, tipped with a little head. Corolla usually golden yellow.

10. ILYSAN'THES Raf. FALSE PIMPERNEL

1. *I. du'bia* Barnhart. Stem 4–8 inches high, much-branched, diffusely spreading. Leaves ovate, rounded or oblong, the upper partly clasping.—Wet places.

11. GERAR'DIA L. GERARDIA

1. *G. purpu'rea* L. (PURPLE GERARDIA.) Corolla rose-purple. Leaves linear, acute, rough-margined. Flowers an inch long, on short peduncles.—Low grounds.

2. *G. pauper'cula* Britton. Smoother, with smaller and lighter-coloured corolla.

3. *G. tenuifo'lia* Vahl. (SLENDER GERARDIA.) Corolla rose-purple. Leaves linear, acute. Flowers about $\frac{1}{2}$ inch long, on long, thread-like peduncles.—Dry woods.

4. *G. fla'va* L. (DOWNY GERARDIA.) Corolla yellow, woolly inside. Stem 3–4 feet high, finely pubescent. Leaves oblong or lance-shaped, the upper entire, the lower usually more or less pinnatifid, downy-pubescent.—Woods.

5. *G. virgin'ica* BSP. (SMOOTH GERARDIA.) Corolla yellow, woolly inside. Stem 3–6 feet high, smooth and glaucous. Lower leaves twice-pinnatifid, the upper pinnatifid or entire, smooth.—Woods.

6. *G. pedicula'ria* L. (CUT-LEAVED GERARDIA.) Nearly smooth. Flowers nearly as in Nos. 4 and 5. Stem 2–3 feet high, very leafy, much-branched. Leaves pinnatifid, the lobes cut and toothed.—Thickets.

12. CASTILLEJA Mutis. PAINTED CUP

1. *C. coccinea* Spreng. (SCARLET PAINTED CUP.) Calyx 2-cleft, yellowish. Stem pubescent or hairy, 1-2 feet high. The stem-leaves nearest the flowers 3-cleft, the lobes toothed, bright scarlet. (A yellow-bracted form occurs on the shore of Lake Huron.)—Sandy soil.

2. *C. palida* Spreng., var. *septentrionalis* Gray. Calyx equally cleft, divisions 2-cleft. Upper lip of corolla decidedly shorter than the tube. Lower leaves linear; upper broader, mostly entire; the floral oblong or obovate, greenish-white, varying to yellowish, purple, or red.—Atl. Prov.

3. *C. sessiliflora* Pursh. Calyx more deeply cleft in front, the narrow lobes deeply 2-cleft. Leaves mostly 3-5 cleft, the floral ones similar, and not coloured.—Pr. Prov.

4. *C. minima* Dougl. Calyx about equally cleft before and behind, the lobes 2-cleft. Corolla over an inch long, the upper lip linear, longer than the tube. Leaves lanceolate or linear, entire, the floral ones usually bright red.—Pr. Prov.

13. ORTHOCARPUS Nutt.

1. *O. luteus* Nutt. Stem about 1 foot high, pubescent, sometimes viscid. Leaves linear to lanceolate, sometimes 3-cleft. Bracts not coloured.—Pr. Prov.

14. EUPHRASIA L. EYEBRIGHT

1. *E. officinalis* L. Rather common on the lower St. Lawrence and the sea-coast. Flowers white, with purple veins. Lowest leaves crenate, those next the flowers bristly-toothed. (Reported by Macoun.)

Var. *Tartarica* Benth. Pale-purple flowers. (Reported by Macoun.)

15. ODONTITES LUDWIG

1. *O. rubra* Gilib. Stem branching, scabrous-pubescent. Leaves oblong-lanceolate, coarsely and remotely serrate.—Atl. sea-coast.

16. RHINANTHUS L. YELLOW RATTLE

1. *R. Cristagalli* L. (COMMON YELLOW RATTLE.) Localities much the same as those of Euphrasia. Seeds broadly winged, rattling in the inflated calyx when ripe.

17. *PEDICULARIS* L. LOUSEWORT

1. *P. canadensis* L. (COMMON LOUSEWORT. WOOD BETONY.) Stems clustered, simple, hairy. Lowest leaves pinnately parted. Flowers in a short spike.—Copses and banks.

2. *P. lanceolata* Michx. Nearly simple, smooth, upright stem, and oblong-lanceolate, cut-toothed leaves. Calyx 2-lobed, leafy-crested. Pod ovate.—Grassy swamps.

3. *P. Furbishiae* Watson. Leaves pinnately parted, and the short oblong divisions pinnately cut, or (in the upper) serrate. Calyx lobes 5. Upper lip of corolla straight and beakless.—River-banks, Atl. Prov.

18. *MELAMPYRUM* L. COW WHEAT

1. *M. lineare* Lam. Leaves lanceolate, short-petioled; the lower ones entire.—Open woods.

LENTIBULARIA'CEAE (BLADDERWORT FAMILY)

Small aquatic or marsh herbs, with a 2-lipped calyx and a personate corolla with a spur or sac underneath. Stamens 2. Ovary as in Primulaceae.

1. *UTRICULARIA* L. BLADDERWORT

1. *U. vulgaris* L. (GREATER BLADDERWORT.) Immersed leaves crowded, finely dissected into capillary divisions, furnished with small air-bladders. Flowers yellow, several in a raceme on a naked scape. Corolla closed; the spur conical and shorter than the lower lip.—Ponds and slow waters.

2. *U. intermedia* Hayne. Immersed leaves 4 or 5 times forked, the divisions linear-awl-shaped, minutely bristle-toothed on the margin, not bladder-bearing, the bladders being on leafless branches. Stem 3–6 inches long. Scape very slender, 3–6 inches long, bearing few yellow flowers. Upper lip of the corolla much longer than the palate; the spur closely pressed to the broad lower lip.—Shallow waters.

3. *U. cornuta* Michx. Spur awl-shaped, turned downwards and outwards, and the lower lip of the corolla helmet-shaped. Flowers yellow. Leaves awl-shaped.—Not uncommon in northern parts of Ontario.

4. *U. clandestina* Nutt. Stems and scapes slender. Leaves hair-like, bearing small bladders. Corolla yellow; lower lip 3-lobed, longer than the thick blunt spur. Submerged stems bearing cleistogamous flowers.—Ponds, Atl. Prov.

5. *U. gibba* L. Scape only 1–3 inches high, 1–2-flowered, with very slender, short branches at the base, bearing capillary, root-like leaves and scattered bladders. Corolla yellow, the lower lip with sides reflexed. Spur very thick and blunt, conical, gibbous.—Shallow water.

6. *U. purpurea* Walt. Capillary leaves petioled, whorled, bearing many bladders. Flowers 2–4 on almost naked scapes. Pedicels erect. Corolla violet-purple, with a 3-lobed lip.

7. *U. resupinata* B. D. Greene. Leaves with small lobes. Flowers showy, solitary on a 2-bracted scape and transverse at its summit; corolla purplish and 2-parted.

2. PINGUICULA L. BUTTERWORT

1. *P. vulgaris* L. A small and stemless perennial growing on damp rocks. Scapes 1-flowered. Leaves entire, ovate or elliptical, soft-fleshy, clustered at the root. Upper lip of the calyx 3-cleft, the lower 2-cleft. Corolla violet, the lips very unequal, the palate open, and hairy or spotted.

OROBANCHACEAE (BROOM-RAPE FAMILY)

Parasitic herbs, destitute of green foliage. Corolla more or less 2-lipped. Stamens didynamous. Ovary 1-celled with 2 or 4 parietal placentae, many-seeded.

1. EPIFAGUS Nutt. BEECH-DROPS

1. *E. virginiana* Bart. A yellowish-brown, branching plant, parasitic on the roots of beech trees. Flowers racemose or spiked; the upper sterile, with long corolla; the lower fertile, with short corolla.

2. CONOPHOLIS Wallroth. SQUAW-ROOT

1. *C. americana* Wallroth. A chestnut-coloured or yellow plant found in clusters in oak woods in early summer, 3–6 inches high and rather less than an inch in thickness. The stem covered with fleshy scales so as to resemble a cone. Flowers under the upper scales; stamens projecting.

3. OROBAN'CHE L. NAKED BROOM-RAPE. CANCER-ROOT

1. *O. uniflo'ra* L. Plant yellowish-brown. Flower solitary at the top of a naked scape. Stem subterranean or nearly so, short and scaly. Scapes 3-5 inches high. Calyx 5-cleft, the divisions lance-awl-shaped. Corolla with a long curved tube and 5-lobed border, and 2 yellow-bearded folds in the throat. Stigma 2-lipped. —Woods, in early summer.

2. *O. fascicula'ta* Nutt. Scaly stem erect, and rising 3 or 4 inches above the ground, mostly longer than the crowded peduncles; parasitic on *Artemisia*, etc.—Pr. Prov.

ACANTHA'CEAE (ACANTHUS FAMILY)

Herbs (with us), with opposite leaves, diandrous (or didynamous) stamens inserted on the tube of the 2-lipped corolla, and a 2-celled and several-seeded capsule. Seeds flat, supported by hooked projections of the placentae. Flowers commonly much-bracted. Calyx 5-cleft.

1. DIANTHE'RA L. WATER WILLOW

1. *D. america'na* L. A perennial herb growing in water or wet places, with entire leaves and purplish flowers in oblong, dense, long-peduncled, axillary spikes. Corolla deeply 2-lipped, the upper erect, notched; the lower spreading, 3-parted. Stamens 2, the anther-cells separated. Pod obovate, flat, 4-seeded.

PHRYMA'CEAE (LOPSEED FAMILY)

Flowers opposite in loose slender spikes, reflexed in fruit; calyx cylindrical, 2-lipped, the upper lip of 3 slender teeth; corolla 2-lipped. Ovary 1-celled, 1-seeded.

1. PHRY'MA L. LOPSEED

1. *P. Leptostach'ya* L. Corolla purplish or pale rose-coloured. Stem slender and branching, 1-2 feet high. Leaves ovate, coarsely toothed.—Woods and thickets.

PLANTAGINA'CEAE (PLANTAIN FAMILY)

Herbs with the leaves all radical, and the flowers in a close spike at the summit of a naked scape. Calyx of 4 sepals, persistent. Corolla 4-lobed, thin and membranaceous, spreading. Stamens 4,

usually with long filaments, inserted on the corolla. Pod 2-celled, the top coming off like a lid. Leaves ribbed. The principal genus is

1. PLANTA'GO L. PLANTAIN. RIBGRASS

1. *P. ma'jor* L. (COMMON PLANTAIN.) Spike long and slender. Leaves 5-7-ribbed, ovate or slightly heart-shaped, with channelled petioles. Pod 7-16-seeded, circumscissile near the middle.—Moist ground about dwellings.

2. *P. Ruge'lii* Decaisne. Leaves as in No. 1, but the petioles are crimson at the base. Spikes elongate. Pods cylindrical 4-9-seeded, circumscissile much below the middle.

3. *P. lanceola'ta* L. (RIBGRASS. ENGLISH PLANTAIN.) Spike thick and dense, short. Leaves 3-5-ribbed, lanceolate or lanceolate-oblong. Scape grooved, long and slender.—Dry fields and banks.

4. *P. me'dia* L. Resembling No. 3, but the leaves are oblong-ovate, downy, and the flowers are fragrant.

5. *P. corda'ta* Lam. Tall and glabrous. Leaves ovate, cordate, 4-10 inches long. Bracts round, ovate, fleshy. Pod 2-4-seeded.

6. *P. elonga'ta* Pursh. Pubescent. Leaves very narrow, linear to filiform, entire. Pod ovoid, short, 4-seeded, about as long as the calyx.—Pr. Prov.

7. *P. decip'iens* Barneoud. Has very narrow and slender spike, and linear fleshy leaves. Found on the sea-coast and lower St. Lawrence.

8. *P. eriop'oda* Torr. Usually a mass of yellowish wool at the base. Leaves thickish, oblanceolate to obovate, with stout short petioles. Pod never more than 4-seeded.—Atl. sea-coast and Pr. Prov.

9. *P. Pursh'ii* R. and S. White with silky wool. Leaves oblong-linear to filiform. Spike very dense, woolly.—Dry soil, Pr. Prov.

RUBIA'CEAE (MADDER FAMILY)

Herbs or shrubs, chiefly distinguished by the presence of stipules between the opposite entire leaves, or by the leaves being in whorls without stipules. Calyx superior. Stamens alternate with the (mostly 4) lobes of the corolla, and inserted on its tube. Ovary 2-4-celled.

Synopsis of the Genera

1. *Ga'lium*. Leaves in whorls. Slender weak herbs with square stems. Calyx teeth inconspicuous. Corolla 4-parted, wheel-shaped. Styles 2. Fruit twin, separating into two 1-seeded carpels.
2. *Sherard'ia*. Leaves in whorls. Stems square, slender, procumbent. Corolla funnel-form, 4-5-lobed. Calyx lobes lanceolate. Flowers blue or pinkish, with a gamophyllous involucre.
3. *Cephalan'thus*. Leaves opposite. Shrubs with the flowers in a globular, peduncled head. Lobes of calyx and corolla each 4. Style very slender, much-protruded. Stigma capitate.
4. *Mitchel'la*. Leaves opposite. Shining, trailing, evergreen herbs, with flowers in pairs, the ovaries united. Lobes of calyx and corolla each 4, the corolla bearded inside. Style 1. Stigmas 4. Fruit a red, 2-eyed berry.
5. *Housto'nia*. Leaves opposite. Lower and slender erect herbs, with the flowers in small terminal clusters or solitary. Lobes of calyx and corolla each 4. Style 1. Stigmas 2.

1. GA'LIUM BEDSTRAW. CLEANERS

1. *G. Apari'ne* L. (CLEAVERS. GOOSEGRASS.) Leaves about 8 in a whorl, lanceolate, rough-margined. Peduncles 1-3-flowered, axillary. Fruit covered with hooked prickles. Low grounds.

2. *G. triflo'rum* Michx. (SWEET-SCENTED BEDSTRAW.) Leaves chiefly 6 in a whorl, elliptical-lanceolate, bristle-pointed. Peduncles 3-flowered, terminating the branches. Fruit covered with hooked prickles.—Woods.

3. *G. pilo'sum* Ait. Leaves in whorls of 4, hairy, oval. Peduncles 2- or 3-forked.

4. *G. lanceola'tum* Torr. (WILD LIQUORICE.) Leaves all in whorls of 4 each, lanceolate, tapering at the apex, more or less 3-nerved. Peduncles mostly 1-forked. Flowers few or several, remote; corolla yellowish to purple. Fruit covered with hooked prickles.

5. *G. circae'zans* Michx. Similar to No. 4, but the leaves are obtuse instead of tapering, and the corolla is greenish.—Woods.

6. *G. kamtschat'icum* Steller. Leaves orbicular to oblong-ovate, and corolla yellowish-white and glabrous.—Mountain woods, Atl. Prov.

7. *G. asprel'lum* Michx. (ROUGH BEDSTRAW.) Leaves in whorls of 6, or 4 or 5 on the branchlets, elliptical-lanceolate, very rough on the edges and midrib. Stem weak, 3-5 feet high, leaning upon and clinging to bushes by recurved prickles on its edges. Flowers numerous in paniced clusters. Fruit not rough.—Thickets.

8. *G. trif'idum* L. (SMALL BEDSTRAW.) Leaves in whorls of 4-6. Stem 6-18 inches high, roughened on the edges, as are the leaves usually. Flowers few, in threes, not paniced. Corolla lobes generally in threes. Fruit smooth.

9. *G. tincto'rium* L. Stem taller and stouter, with nearly smooth angles. Leaves mostly in fours, linear-lanceolate, 1-nerved. Peduncles 3-7-flowered. Corolla lobes and stamens 4. Fruit smooth.

10. *G. borea'le* L. (NORTHERN BEDSTRAW.) Leaves in whorls of 4, linear-lanceolate, 3-nerved. Flowers very numerous, crowded in a narrow and compact terminal panicle. Stem erect and rigid, 1-3 feet high. Fruit generally hispid.—Rocky thickets and river-banks.

11. *G. Mollu'go* L. Stems erect, usually smooth. Leaves in eights or on the branchlets in sixes, oblanceolate to linear. Flowers white, in loose panicles.

12. *G. ve'rum* L. (YELLOW BEDSTRAW.) Stem erect. Leaves usually in eights, linear, soon deflexed. Flowers yellow, in dense panicles.

2. SHERARD'IA L.

1. *S. arven'sis* L. Sparingly naturalized.

3. CEPHALAN'THUS L. BUTTON BUSH

1. *C. occidenta'lis* L. A smooth shrub growing in swamps, with ovate, petioled, pointed leaves, which are opposite or in whorls of 3. Easily recognized by the globular head of white flowers.

4. MITCHEL'LA L. PARTRIDGE BERRY

1. *M. re'pens* L. Leaves round-ovate, shining, sometimes with whitish lines.—Common in dry woods.

5. HOUSTO'NIA L. HOUSTONIA

1. *H. purpu'rea* L. Stems tufted, 3-6 inches high. Leaves varying from roundish-ovate to lanceolate, 3-5-ribbed, sessile.—Woodlands.

2. *H. longifo'lia* Gaertn. Lower, with thinner, oblong-lanceolate to linear leaves.—Chiefly Pr. Prov.

3. *H. caeru'lea* L. (BLUETS. INNOCENCE.) A slender herb with erect stems. A single flower on each slender peduncle. Leaves oblong-spatulate. Corolla light blue to nearly white, with a yellowish eye and a long tube.—Moist grassy places.

CAPRIFOLIA'CEAE (HONEYSUCKLE FAMILY)

Shrubs, rarely herbs, with the calyx tube adherent to the ovary, the corolla borne on the ovary, and the stamens as many as the petals (1 less in *Linnaea*) borne on the tube of the corolla. Leaves opposite and without stipules, but some species of *Viburnum* have appendages resembling stipules. Fruit a berry, drupe, or pod.

Synopsis of the Genera

A. Corolla tubular, sometimes 2-lipped. Style slender.

1. *Linnae'a*. A trailing or creeping herb with evergreen, oval, crenate leaves and slender, scape-like peduncles which fork at the top into 2 pedicels, each of which bears a nodding, narrowly bell-shaped, purplish flower. Stamens 4, 2 shorter than the others.
2. *Symphoricar'pos*. Upright, branching shrubs with oval, entire, short-petioled leaves. Flowers in interrupted spikes at the ends of the branches, rose-coloured. Corolla bell-shaped, 4-5-lobed, with as many stamens. Berries large and white, 4-celled, but only 2-seeded.
3. *Lonic'ra*. Upright or twining shrubs with entire leaves. Corolla funnel-form, more or less irregular, 5-lobed, often with a projection on one side at the base. Berry several-seeded.
4. *Diervil'la*. Low upright shrubs with ovate, pointed, serrate leaves. Calyx tube tapering towards the top, the teeth slender. Flowers light yellow, peduncles mostly 4-flowered. Corolla funnel-form, nearly regular. Pods slender-pointed.
5. *Trios'teum*. Coarse herbs. Lobes of the calyx leaf-like. Flowers brownish-purple, sessile in the axils of the leaves. Corolla bulging at the base, 5-lobed. Fruit a 3-seeded, orange-coloured drupe.

B. Corolla rotate or urn-shaped, regular, 5-lobed. Flowers white, in broad cymes. Stigma sessile or nearly so.

6. *Sambu'cus*. Upright shrubs with pinnate leaves, the leaflets serrate. Corolla spreading urn-shaped, 5-lobed. Stigmas 3.

Fruit purple or red, a juicy, berry-like drupe, with 3 seed-like stones.

7. *Viburnum*. Upright shrubs with simple leaves, and white flowers in flat, compound cymes. Corolla 5-lobed. Stigmas 1-3. Fruit a 1-seeded drupe.

1. LINNAEA L. TWIN-FLOWER

1. *L. borealis* L., var. *americana* Rehder.—Cool mossy woods and swamps.

2. SYMPHORICARPOS Ludwig. SNOWBERRY

1. *S. racemosus* Michx. (SNOWBERRY.) Shrub with leaves oblong-orbicular, green both sides. Corolla bearded inside. Flowers in a rather loose spike.

Var. *pauciflorus* Robbins. Low, leaves whitened beneath, diffusely branched, and spreading, with two or three flowers only, in the axils of the uppermost leaves.—Dry rocky hillsides.

2. *S. occidentalis* Hook. (WOLFBERRY.) Flowers in denser spikes than the last, and with larger and more funnel-form corolla and longer stamens. Corolla much-bearded within.—Pr. Prov.

3. LONICEA L. HONEYSUCKLE. WOODBINE

1. *L. dioica* L. (SMALL HONEYSUCKLE.) Twining shrub, 2-4 feet high, with smooth leaves which are glaucous beneath, the upper ones connate-perfoliate; corolla yellowish-purple.—Rocky banks.

2. *L. hirsuta* Eaton. (HAIRY HONEYSUCKLE.) Stem twining high. Leaves not glaucous, very large, downy-hairy, the upper ones connate-perfoliate. Flowers in close whorls; corolla greenish-yellow, clammy-pubescent.—Deep thickets.

3. *L. Sullivanii* Gray. Twining. At length very glaucous-whitened. Leaves oval and ovate-oblong, mostly connate on the flowering stems; the united leaves form an orbicular disk. Corolla pale yellow.—Pr. Prov.

4. *L. glaucescens* Rydb. Less glaucous than No. 3. Leaves pubescent only beneath; the connate ones form a somewhat oblong or rhombic disk.—Ont. and Pr. Prov.

5. *L. canadensis* Marsh. (FLY HONEYSUCKLE.) A branching, straggling shrub, with thin, oblong-ovate, ciliate leaves. Peduncles axillary, filiform, shorter than the leaves, each 2-flowered at the top. Corolla greenish-yellow, almost spurred at the base. The two red berries separate.—Damp woods.

6. *L. caerulea* L., var. *villosa* T. and G. (MOUNTAIN FLY HONEYSUCKLE.) Smaller (1-2 feet high), and with upright branches. Leaves oval. Ovaries united into one blue, edible berry.

7. *L. oblongifolia* Hook. (SWAMP FLY HONEYSUCKLE.) A shrub with upright branches and oblong leaves. Peduncles long and slender, 2-flowered. Corolla deeply 2-lipped, yellowish-white. Berries reddish-purple, united at the base.—Swamps and low grounds.

8. *L. involucrata* Banks. Is at once recognized by the involucre of 4 leaf-like bracts under the 2 flowers. Corolla yellowish, viscid-pubescent. Berries dark purple, distinct.—Woods and banks of streams, Atl. Prov. and Pr. Prov.

4. DIERVILLA Mill. BUSH HONEYSUCKLE

1. *D. lonicera* Mill.—Rocky woods and clearings.

5. TRIOSTEUM L. FEVERWORT

1. *T. perfoliatum* L. A coarse herb, 2-4 feet high, soft-hairy. Leaves oval, narrowed at the base. Fruit orange-coloured.—Old clearings and thickets.

6. SAMBUCUS L. ELDER

1. *S. canadensis* L. (COMMON ELDER.) Shrub 5-10 feet high, in clumps; pith white. Leaflets 5-11, oblong. Cymes flat. Fruit black-purple.—Open grounds, and along streams.

2. *S. racemosa* L. (RED-BERRIED ELDER.) May be distinguished from No. 1 by its warty bark, brown pith, 5-7 leaflets, convex or pyramidal cymes, and red berries.—Rocky woods.

7. VIBURNUM L. ARROW-WOOD. LAURESTINUS

1. *V. Lenta* L. (SWEET VIBURNUM. SHEEP-BERRY. NANNY-BERRY.) A small tree with ovate, finely-serrate, pointed leaves, with long and margined petioles. Cyme sessile. Fruit bluish-black.—Along streams.

2. *V. cassinoides* L. (WITHE-ROD.) A smooth shrub with somewhat scurfy shoots and tall straight stems. Leaves thickish, entire or wavy-toothed, dotted beneath. Cymes with short peduncles, about 5-rayed. Fruit blue-black with a bloom.—Cold swamps.

3. *V. pubes'cens* Pursh. (DOWNY ARROW-WOOD.) A straggling shrub, not more than 4 feet high, with small, ovate, coarsely serrate leaves, the lower surface soft-downy. Cymes small. Fruit oblong, dark purple.—Rocky places.

4. *V. acerifo'lium* L. (MAPLE-LEAVED ARROW-WOOD. DOCK-MACKIE.) A shrub 3-6 feet high, with greenish bark. Leaves 3-lobed, 3-ribbed, soft-downy beneath. Stipular appendages bristle-shaped. Cymes small, on long peduncles. Fruit red, becoming black.—Thickets and river-banks.

5. *V. Op'ulus* L., var. *america'num* Ait. (CRANBERRY TREE. HIGH-BUSH CRANBERRY.) An upright shrub, 5-10 feet high, with strongly 3-lobed leaves, broader than long, the lobes spreading and pointed. Cymes peduncled. Marginal flowers of the cyme very large and neutral. Stipular appendages conspicuous. Fruit red, pleasantly acid.—Low grounds.

6. *V. pauciflo'rum* Raf. A low shrub. Leaves 5-ribbed at the base, serrate, with 3 short lobes at the summit. Cyme few-flowered. Stamens shorter than the corolla. Fruit red, sour, with a very flat stone.—Cold woods.

7. *V. alnifo'lium* Marsh. (HOBBLE-BUSH.) A straggling shrub with reclining branches. Leaves large, round-ovate, heart-shaped at the base, serrate, many-veined, the veins underneath and the stalks and branchlets very rusty-scurfy. Stipular appendages conspicuous. Cymes sessile, very broad and flat, with very conspicuous, neutral flowers on the margin.—Moist woods.

VALERIANA'CEAE (VALERIAN FAMILY)

Herbs with opposite exstipulate leaves and small cymose flowers. Calyx tube adherent to the ovary, the latter 3-celled, but only one of these fertile. Stamens 1-3, fewer than the lobes of the corolla. Style slender. Stigmas 1-3.

1. VALERIA'NA L. VALERIAN

1. *V. uligino'sa* Rydb. Not uncommon in cedar-swamps. Root fibrous. Calyx limb consisting of several bristles rolled inwards in the flower, but expanding in fruit. Corolla gibbous at the base, rose-coloured or white. Stamens 3. Root-leaves ovate or oblong, usually entire; stem-leaves pinnate, leaflets 5-11. Stem erect, striate, 1-5 feet high.

2. *V. edulis* Nutt. Root spindle-shaped, large. Flowers whitish, in a long and narrow, interrupted panicle, nearly dioecious. Stem-leaves deeply pinnatifid; root-leaves spatulate and lanceolate.—Low grounds.

DIPSACA'CEAE (TEASEL FAMILY)

Herbs with the flowers in heads, surrounded by a many-leaved involucre, as in the **Compositae**, but the stamens are distinct. Leaves opposite.

1. DIP'SACUS L. TEASEL

1. *D. sylvestris* Huds. (WILD TEASEL.) A stout, coarse, prickly plant, not unlike a thistle in appearance. Flowers in oblong, very dense heads, bluish. Corolla 4-cleft. Stamens 4, on the corolla. Bracts among the flowers terminating in a long awn. Leaves generally comate.—Roadsides and ditches.

CUCURBITA'CEAE (GOURD FAMILY)

Herbs, climbing by tendrils. Flowers monoecious. Calyx tube adherent to the 1-3-celled ovary. Corolla commonly more or less gamopetalous. Stamens usually 3, united by their tortuous anthers, and often also by the filaments. Leaves alternate, palmately lobed or veined.

Synopsis of the Genera

1. **Sic'yos.** Flowers greenish-white, small; the staminate corymboid, the pistillate clustered in a head on a long peduncle. Corolla 5-cleft, with a spreading border. Style slender; stigmas 3. Ovary 1-celled, 1-seeded. Fruit dry and indehiscent, prickly, bur-like in appearance.
- 2 **Echinocys'tis.** Flowers whitish, small; the staminate in long compound racemes, the pistillate in small clusters from the same axils. Corolla 6-parted. Stigma broad, almost sessile. Ovary 2-celled, 4-seeded. Fruit fleshy, becoming dry, clothed with weak prickles.

1. SIC'YOS L. STAR CUCUMBER

1. *S. angulatus* L. A clammy-hairy weed in damp yards. Leaves roundish-heart-shaped, 5-angled or lobed.

2. ECHINOCYS'TIS Torr. and Gray. WILD BALSAM APPLE

1. *E. loba'ta* Torr. and Gray. Climbing high about dwellings. Leaves deeply and sharply 5-lobed. The oval fruit 2 inches long.

CAMPANULA'CEAE (CAMPANULA FAMILY)

Herbs with milky juice, differing from the following Family chiefly in having a regular, 5-lobed corolla (bell-shaped or wheel-shaped), separate stamens (5), and 2 or more (with us, 3) stigmas.

Synopsis of the Genera

1. *Campan'ula*. Calyx 5-cleft. Corolla generally bell-shaped, 5-lobed. Pod short.
2. *Specula'ria*. Calyx 5-lobed. Corolla nearly wheel-shaped, 5-lobed. Pod prismatic or oblong.

1. CAMPAN'ULA L. BELL-FLOWER

1. *C. rotundifo'lia* L. (HAREBELL.) Flowers blue, loosely paniced, on long slender peduncles, nodding. Stem slender, branching, several-flowered. Root-leaves round-heart-shaped; stem-leaves linear. Calyx lobes awl-shaped.—Shaded banks.

2. *C. aparinoi'des* Pursh. (MARSH BELL-FLOWER.) Flowers white or nearly so, about $\frac{1}{3}$ inch long. Stem very slender and weak, few-flowered, angled, roughened backwards. Leaves linear-lanceolate. Calyx lobes triangular.—Wet places in high grass. This plant has the habit of a *Galium*.

3. *C. america'na* L. (TALL BELL-FLOWER.) Flowers light blue, about an inch across, crowded in a leafy spike. Corolla deeply 5-lobed. Style long and curved. Stem 3–6 feet high, simple. Leaves ovate or ovate-lanceolate, taper-pointed, serrate.—Moist rich soil.

4. *C. rapunculoi'des* L. Flowers nodding, single in the axils of bracts, forming a raceme. Stem-leaves pointed, ovate-lanceolate, serrate; the lower cordate, ovate, long-petioled.—Atl. Prov.

2. SPECULA'RIA Fabricius. VENUS'S LOOKING-GLASS

1. *S. perfolia'ta* A. DC. Flowers purplish-blue, only the latter or upper ones expanding. Stem hairy, 3–20 inches high. Leaves roundish or ovate, clasping. Flowers solitary or 2 or 3 together in the axils.—Sterile open ground.

LOBELIA'CEAE (LOBELIA FAMILY)

Herbs with milky acrid juice, alternate leaves, and loosely racemed flowers. Corolla irregular, 5-lobed, the tube split down one side. Stamens 5, syngenesious, and commonly also monadelphous, free from the corolla. Calyx tube adherent to the many-seeded ovary. Style 1.

1. LOBELIA L. LOBELIA

1. *L. cardinalis* L. (CARDINAL FLOWER.) Corolla large, deep red. Stem simple, 2-3 feet high, smooth. Leaves oblong-lanceolate, slightly toothed. Bracts of the flowers leaf-like.—Low grounds.

2. *L. syphilitica* L. (GREAT LOBELIA.) Corolla rather large, light blue. Stem hairy, simple, 1-2 feet high. Leaves thin, acute at both ends, serrate. Calyx lobes half as long as the corolla, the tube hemispherical. Flowers in a dense spike or raceme.—Low grounds.

3. *L. inflata* L. (INDIAN TOBACCO.) Flowers small, $\frac{1}{8}$ inch long, pale blue. Stem leafy, branching, 8-18 inches high, pubescent. Leaves ovate or oblong, toothed. Pods inflated. Racemes leafy.—Dry fields.

4. *L. spicata* Lam. Flowers small, $\frac{1}{8}$ inch long, pale blue. Stem slender, erect, simple, 1-3 feet high, minutely pubescent below. Leaves barely toothed, the lower spatulate or obovate, the upper reduced to linear bracts. Racemes long and naked.—Sandy soil.

5. *L. Kal'mii* L. Flowers small, $\frac{1}{8}$ inch long, light blue. Stem low, 4-18 inches high, very slender. Pedicels filiform, as long as the flowers, with 2 minute bractlets above the middle. Leaves mostly linear, the radical ones spatulate and the upper ones reduced to bristly bracts.—Wet rocks and banks, chiefly northward.

6. *L. Dortman'na* L. (WATER LOBELIA.) Small leaves, all tufted at the root, and a hollow scape 5 or 6 inches long with a few small, light-blue, pedicelled flowers at the summit.—Borders of ponds in Muskoka.

COMPOS'ITAE (COMPOSITE FAMILY)

Flowers in a dense head on a common receptacle, and surrounded by an involucre. Calyx tube adherent to the ovary, its limb either obsolete or forming a pappus of few or many bristles or chaffy scales, etc. Corolla either tubular or with one side much-prolonged (strap-shaped or ligulate). Stamens usually 5, on the tube of the corolla, their anthers united (syngenesious). Style 2-cleft.

The heads of flowers present some variety of structure. All the flowers of a head may be tubular; or only the central ones or disk-flowers, as they are then called, may be tubular, whilst those around the margin, then known as ray-flowers, are ligulate or strap-shaped. Or again, all the flowers may be strap-shaped. It is not unusual, also, to find a mixture of perfect and imperfect flowers in the same head.

The bracts which are often found growing on the common receptacle among the florets are known as the chaff. When these bracts are entirely absent, the receptacle is said to be naked. The leaves of the involucre are called its scales or bracts.

Artificial Synopsis of the Genera

SERIES I. TUBULIFLO'RAE

Heads either altogether without strap-shaped corollas, or the latter, if present, forming only the outer circle (the ray). Ray-flowers, when present, always without stamens, and often without a pistil also.

I. a. Ray-flowers entirely absent.

A. *Scales of the involucre in many rows, bristly-pointed, or fringed.*

(1) *Florets all perfect.*

1. Cir'sium. Leaves and scales of the involucre prickly. Pappus of long plumose bristles. Receptacle with long soft bristles among the florets. Flowers reddish-purple.
2. Car'duus. Resembling Cirsium, but the bristles of the pappus are not plumose.
3. Onopor'dum. Leaves and scales of the involucre prickly. Heads much as in Cirsium, but the receptacle naked, and deeply honeycombed. Pappus of long bristles, not plumose. Stem winged by the decurrent bases of the leaves. Flowers purple.

4. **Arc'tium.** Leaves not prickly, but the scales of the globular involucre tipped with hooked bristles. Pappus of many short, rough bristles. Receptacle bristly. Flowers purple, rarely white.
- Verno'nia.** One species has scales of the involucre somewhat bristly.
- (2) *Marginal florets often sterile, and much larger than the others, forming a kind of false ray. Both kinds of florets in one head.*
5. **Centaur'ea.** Leaves not prickly. Scales of the involucre fringed. Pappus double and bristly, or very short or none. Receptacle bristly.
6. **Cni'cus.** Leaves spiny. Flowers yellow. Pappus of 10 teeth; 10 long bristles and 10 shorter ones.
- (3) *Sterile and fertile florets in separate heads, i.e., monoecious. Fruit a completely closed involucre (usually bristly) containing only one or two florets, these heads sessile in the axils of the bracts or upper leaves. Sterile heads with more numerous florets in flattish involucre, and forming racemes or spikes. Pappus none.*
7. **Xan'thium.** Fertile florets only 2 together in burs with hooked prickles, clustered in the axils. Sterile heads in short spikes above them, the scales of their involucre in one row only, but not united together.
8. **Ambro'sia.** Fertile florets single, in a closed involucre armed with a few spines at the top. Sterile heads in racemes or spikes above, the scales of their involucre in a single row and united into a cup.
9. **Franse'ria.** Inflorescence much like that of Ambrosia, but the fruiting involucre is armed with a number of long, flat and thin spines.

B. *Scales of the involucre without bristles of any kind.*

(1) *Marginal florets without stamens.*

- (a) *Pappus none or minute. Receptacle naked. Very strong-scented herbs.*
10. **Tanace'tum.** Flowers yellow, in numerous corymbed heads. Scales of the involucre dry, imbricated. Pappus 5-lobed. Leaves dissected.
11. **Artemis'ia.** Flowers yellowish or dull purplish, in numerous small heads which are panicked or racemed. Scales of the involucre with dry and scarious margins, imbricated. Achenes with narrow top.

(b) *Pappus of all the florets bristly. Receptacle naked.*

12. **Erecht'ites.** Flowers whitish. Scales of the involucre in a single row, linear, with a few bractlets at the base. Corolla of the marginal florets very slender. Pappus copious, of fine, soft, white hairs. Heads corymbed. Erect and coarse herbs.
13. **Gnaph'alium.** Flowers whitish or yellowish. Scales of the involucre yellowish-white, in many rows, dry and scarious, woolly at the base. Outer corollas slender. Pappus a single row of rough bristles. Flocculent-woolly herbs.
14. **Anaph'alis.** Resembles *Antennaria*. Pappus not thickened above and not at all united at the base.
15. **Antenna'ria.** Very much like *Gnaphalium* in appearance, being white-woolly, but the heads are usually dioecious, and the bristles of the pappus thicker in the sterile florets.

(2) *All the florets in the head perfect.*

15. **Antenna'ria**, with dioecious heads, may be looked for here. See previous paragraph.

Bidens. One or two species have no rays.

Sene'cio. One species is without rays.

16. **Lia'tris.** Flowers handsome, rose-purple. Receptacle naked. Pappus of long and slender bristles, plumose or rough. Achenes slender, 10-ribbed. Lobes of the corolla slender. Stem wand-like, leafy, from a corm or tuber. Leaves narrow or grass-like.
 17. **Verno'nia.** Flowers purple. Scales of the involucre (with us) with slender bristly tips. Receptacle naked. Pappus double, the outer minute, the inner capillary. Branches of the style long and slender, minutely bristly.
 18. **Eupato'rium.** Flowers white or purple. Receptacle naked. Pappus of slender, hair-like bristles, smooth or nearly so, in a single row. Achenes 5-angled. Heads in corymb. Leaves whorled, connate, or opposite.
- (3) *Marginal florets pistillate and fertile, those in the centre of the head staminate and sterile. Receptacle chaffy. Pappus wanting.*
19. **Caca'lia.** Flowers white or whitish. Heads 5-flowered, rather large, in flat corymb. Involucre 5-leaved. Receptacle naked, bearing a scale-like, pointed appendage in the centre. Corolla deeply 5-cleft. Pappus of numerous capillary bristles.
 20. **I'va.** Marginal pistillate florets 1-5, with very small, tubular corolla or none. Staminate florets with funnel-form, 5-toothed corolla, receptacle with narrow chaff. Anthers nearly separate. Achenes very short. Coarse plants with small, greenish-white, nodding heads.

I. b. Rays or strap-shaped corollas around the margin of the head.

A. *Pappus of hairy-like bristles. Receptacle naked.*
Rays pistillate.

21. **Sene'cio.** Rays yellow, or in one species none. Scales of the involucre in a single row, or with a few bractlets at the base. Pappus very fine and soft. Heads corymbose or solitary. Leaves alternate.
22. **In'ula.** Rays yellow, numerous, very narrow, in a single row. Outer scales of the involucre leaf-like. Anthers with two tails at the base. Stout plants, with large alternate leaves which are woolly beneath.
23. **Chrysop'sis.** Rays yellow, many. Disk yellow. Pappus double, the outer of very small, chaffy bristles, the inner capillary. Low hairy herbs.
24. **Aplopap'pus.** Rays yellow, many. Disk yellow. Pappus simple, of unequal bristles.
25. **Solida'go.** Rays yellow, few, as are also the disk-florets. Involucre oblong, scales of unequal lengths, appressed. Achenes many-ribbed. Heads small, in compound racemes or corymbs. Stems usually wand-like. Leaves alternate.
26. **Ar'nica.** Rays yellow. Disk yellow. Scales of the bell-shaped involucre in two rows, lanceolate, erect. Receptacle flat, minutely fringed. Achenes slender. Leaves opposite. Heads showy, with long peduncles. Stem simple.
27. **Tussila'go.** Rays yellowish, in several rows, very narrow. Involucre nearly simple. Pappus copious, soft. A low perennial with creeping rootstocks, sending up a scaly scape in early spring bearing one head, and producing later heart-shaped, angled or toothed leaves, woolly when young.
28. **Petasi'tes.** Rays whitish or purplish. Heads in a corymb, fragrant. Scales of the involucre in a single row. Heads somewhat dioecious, the staminate with one row of pistillate ray-flowers, the pistillate with ray-flowers in many rows. Woolly herbs with large leaves, all radical, and sheathing scaly bracts on the scape.
29. **Aster.** Rays white, purple, or blue, never yellow, but the disk generally yellow. Pappus of numerous fine, roughish bristles (in one species double). Achenes flattish. Heads corymbed or racemose. Flowering in late summer.
30. **Erig'eron.** Rays and disk as in Aster, but the rays very narrow, and usually in more than one row. Scales of the involucre in one or two rows, nearly of equal length. Pappus of long bristles with shorter ones intermixed, simple in some species. Heads corymbed or solitary. Leaves generally sessile.

B. *Pappus not of hair-like bristles, but altogether wanting or consisting of a few chaffy scales or teeth, or only a minute crown.*

(1) *Receptacle naked.*

31. **Grinde'lia.** Rays yellow, many. Disk yellow. Heads large, Pappus of 2 or 3 rigid awns, falling off early. Scales of the involucre with spreading tips. Coarse herbs with sessile serrate leaves.
32. **Gutierre'zia.** Rays yellow, few. Disk-florets 3 or 4. Heads numerous, small, crowded. Pappus of a few short, chaffy scales. A low, partly shrubby plant, with narrowly linear, entire, alternate, leaves.
33. **Hele'nium.** Rays yellow, wedge-shaped, 3-5-cleft at the summit. Scales of the involucre reflexed, awl-shaped. Pappus of 5-8 chaffy scales, 1-nerved, the nerve usually extending to a point. Leaves alternate, decurrent on the angled stem. Heads corymbed, showy.
34. **Chrysan'themum.** Rays white; disk yellow. Disk-corollas with a flattened tube. Pappus none. Heads single, or corymbed. Involucre broad and flat, the scales with scarious margins. Receptacle flattish.
35. **Bellis.** Rays white or pink, numerous. Scales of the involucre in about 2 rows. Receptacle conical, naked. Low herbs with solitary heads on scapes.—Escaped from cultivation.
36. **Matrica'ria.** Rays white, or none, disk yellow. Receptacle conical. Achenes 3-5-ribbed. Pappus a short crown.

(2) *Receptacle chaffy.*

37. **An'themis.** Rays white. Disk yellow. Ray-florets pistillate or neutral. Pappus none or a minute crown. Receptacle conical, more or less chaffy. Herbs with strong odour, and pinnately dissected leaves. Heads terminal.
38. **Rudbeck'ia.** Rays yellow, usually long; disk dark purple, or, in one species, greenish-yellow. Scales of the involucre leaf-like. Receptacle conical. Pappus none, or only a minute crown. Ray-florets neutral.
39. **Lep'achys.** Rays yellowish, drooping; disk grayish. Scales of the involucre few, small and spreading. Pappus none or of 2 teeth.
40. **Helian'thus.** Rays yellow, neutral. Receptacle flattish or convex. Chaff persistent, and embracing the 4-sided achenes. Pappus deciduous, of 2 thin scales. Coarse herbs.
41. **Actinom'eris.** Rays 2-8 (or none), irregular, yellow, neutral. Receptacle small, convex. Scales of the involucre few, soon deflexed. Achenes flat, obovate, spreading when ripe. Pappus of 2 or 3 smooth awns, persistent. Tall branching perennials with mostly decurrent leaves.

42. **Coreop'sis.** Rays mostly 8, yellow (sometimes brown-tinted), neutral. Receptacle flat. Involucre double, each of about 8 scales, the outer leaf-like and spreading, the inner rather membranaceous and appressed. Achenes flat, compressed parallel with the scales of the involucre, 2-toothed or sometimes naked. Heads corymbed.
43. **Gaillard'ia.** Rays yellow (sometimes purplish), numerous, 3-cleft. Receptacle convex, the chaff bristly or awl-shaped. Achenes top-shaped, 5-ribbed. Pappus of several long, thin scales, awn-tipped. Heads showy, fragrant, terminal, on scape-like stems.
44. **Bi'dens.** Rays yellow, few; but 2 species are without rays. Scales of the involucre in 2 rows, the outer large and leaf-like. Ray-florets neutral. Achenes crowned with 2 or more stiff awns which are barbed backwards.
45. **Heliop'sis.** Rays yellow, 10 or more, pistillate. Scales of the involucre in 2 or 3 rows, the outer leaf-like. Receptacle conical; chaff linear. Achenes smooth, 4-angled. Pappus none.
46. **Achille'a.** Rays white (occasionally pinkish), few. Receptacle flattish. Pappus none. Achenes margined. Heads small, in flat corymb. Leaves very finely dissected.
47. **Polym'nia.** Rays whitish-yellow, wedge-form, shorter than the involucre, few in number. Scales of the involucre in 2 rows, the outer leaf-like, the inner small and partly clasping the achenes. Pappus none. Coarse clammy herbs with an unpleasant odour.
48. **Sil'phium.** Coarse herbs with resinous juice and yellow flowers. Flat involucre, with bracts in 2 rows, the outer with leaf-like tips. Pappus none or of 2 teeth joined to the margins of the achenes.
49. **Galinso'ga.** Heads numerous, small. Rays few, pistillate, white. Disk yellow. Bracts of the involucre few, ovate. Pappus usually of cut-fringed scales. Annuals.

SERIES II. LIGULIFLO'RAE

Corolla strap-shaped in all the florets of the head. All the florets perfect. Herbs with milky juice and alternate leaves.

50. **Krig'ia.** Flowers yellow. Pappus double, the outer short, of many minute, chaffy scales, the inner of many long, capillary bristles. Low perennials branching from the base, with single small heads on scapes.
51. **Lap'sana.** Flowers yellow, 8-12 in a head. Scales of the involucre 8, in a single row, with a short outer series. Pappus none. Stem slender. Heads small, in loose panicles.

52. Cicho'rium. Flowers bright blue, showy. Scales of the involucre in 2 rows, the outer of 5 short scales, the inner of 8-10 scales. Pappus chaffy. Heads sessile, 2 or 3 together.
53. Leon'todon. Flowers yellow. Involucre with bractlets at the base. Pappus of plumose bristles, these broader at the base. Heads borne on branching scapes. Leaves radical.
54. Hiera'cium. Flowers yellow. Scales of the involucre more or less imbricated. Pappus a single row of tawny, hair-like, rough bristles. Heads corymböse.
55. Cre'pis. Like Hieracium, but the pappus is of copious, white and soft, capillary bristles.
56. Prenan'thes. Stems leafy. Flowers yellowish or greenish-white often tinged with purple; heads nodding. Involucre of 5-14 scales in a single row, with a few bractlets below. Pappus copious, of brownish or yellowish, rough bristles. Achenes short, blunt.
57. Lygodes'mia. Stems nearly leafless. Flowers rose-purple. Involucre with bractlets below. Pappus white. Heads erect. Achenes long, tapering.
58. Agos'eris. Flowers yellow. Head large, solitary, borne on a scape. Leaves radical, elongated, linear, tufted. Achenes 10-ribbed. Involucre loosely imbricated in 2 or 3 rows.
59. Tarax'acum. Flowers yellow, on slender, naked, hollow scapes. Achenes prolonged into a slender, thread-like beak, on which is borne the copious white pappus. Leaves all radical, runcinate.
60. Lactu'ca. Flowers pale yellow or purplish. Florets about 20, in the head. Scales of the involucre in 2 or more rows of unequal length. Achenes with long thread-form or short and thick beaks, and a pappus of very soft, white or tawny bristles. Heads numerous, paniced. Tall smooth herbs.
61. Son'chus. Flowers pale yellow. Heads many-flowered, enlarging at the base. Achenes without beaks. Pappus very soft and white. Tall glaucous herbs with spiny-toothed leaves.
62. Tragopo'gon. Flowers yellow or purple. Heads large. Involucre of about 12 lanceolate, rather fleshy scales in one row, somewhat united at the base. Achenes with long tapering beaks. Pappus of plumose bristles, 5 of these longer and naked at the summit. Leaves entire, straight-veined, clasping.

1. CIR'SIUM Hill. COMMON THISTLE

1. C. lanceola'tum Hill. (COMMON BULL THISTLE.) All the scales of the involucre prickly-pointed. Leaves decurrent, pinnati-

fid, the lobes prickly-pointed, rough above, woolly with webby hairs beneath.—Fields and roadsides everywhere.

2. *C. undula'tum* Spreng. White-woolly throughout. Low and stout, leafy. Scales tipped with short prickles. Leaves undulate-pinnatifid.—Pr. Prov.

3. *C. pu'milum* Spreng. Differs from the last in being pubescent but not woolly. The leaves, also, are not pinnatifid. Stem 1-flowered.—Lake Huron.

4. *C. dis'color* Spreng. The inner scales of the involucre not prickly. Stem grooved. Leaves prickly, green above, white-woolly beneath. Flowers pale purple. Whole plant with a whitish aspect.—Dry thickets.

5. *C. mu'ticum* Michx. (SWAMP THISTLE.) Scales of the webby involucre hardly prickly, and not spreading. Stem very tall, and smoothish, and sparingly leafy. Heads single or few.—Swamps and low woods.

6. *C. arven'se* Scop. (CANADA THISTLE.) Scales of the involucre with reflexed points. Leaves prickly, smooth both sides, or slightly woolly beneath. Roots extensively creeping. Heads small and numerous.—Fields and roadsides.

2. CAR'DUUS L. PLUMELESS THISTLE

1. *C. nu'tans* L. (MUSK THISTLE.) Leaves decurrent, sinuate, spiny. Heads solitary, drooping. Flowers purple.—Atl. Prov.

3. ONOPOR'DUM L. SCOTCH THISTLE

1. *O. Acan'thium* L. A coarse, branching herb, 3-8 feet high, with woolly stem and leaves. Bristles of the pappus united at the base into a ring.—Roadsides and old fields; not common.

4. ARC'TIUM L. BURDOCK

1. *A. mi'nus* Bernh. (COMMON BURDOCK.) A coarse plant with very large, cordate leaves with hollow petioles, and numerous small, globular heads of purple flowers. The involucre forms a bur, which clings to one's clothing or to the hair of animals.—Near dwellings, mostly in manured soil.

2. *A. Lap'pa* L. Differs from No. 1 in having larger greenish flower-heads, $1\frac{1}{4}$ inches across, and the hooked scales of the involucre more spreading; also the petioles of the leaves are not hollow.

5. CENTAUR'EA L. STAR THISTLE

1. *C. Cy'anus* L. (BLUEBOTTLE.) An old garden plant, found occasionally along roadsides. False rays very large. Scales of the involucre fringed. Leaves linear, entire or nearly so. Stem erect. Heads single at the ends of the branches, the flowers bluish. Pappus very short.

2. *C. ni'gra* L. (KNAPWEED.) No false rays. Scales of the globular involucre black-fringed. Leaves lanceolate, mostly entire, rough. Flowers purple. Pappus short or none.—Waste places, chiefly eastward.

6. CNI'CUS L. BLESSED THISTLE

1. *C. benedic'tus* L. Flowers yellow. Pappus double, of 10 long outer bristles and 10 short inner ones. A low, branching annual, with clasping, cut leaves, and large, sessile, leafy-bracted heads.—Atl. Prov.

7. XAN'THIUM L. CLOTBUR

1. *X. echina'tum* Murr. (COMMON COCKLEBUR.) Stem rough, not prickly or spiny. Leaves broadly triangular, and somewhat heart-shaped, long-petioled. Fruit a hard, 2-celled bur, nearly an inch long, clothed with stiff hooked prickles, the two beaks of the fruit long and usually incurved.—Low river-banks and waste places.

2. *X. spino'sum* L. (SPINY CLOTBUR.) Stem armed with conspicuous, straw-coloured, triple, slender spines, at the bases of the lanceolate, short-petioled leaves, the latter white-woolly beneath.—Town of Dundas, Ontario; the seeds having been brought in wool from South America.

3. *X. canaden'se* Mill. Similar in habit to No. 1. Leaves broadly ovate, cordate, indistinctly 3-lobed and coarsely toothed. Burs oblong, smooth, with two slightly incurved beaks, and scattered, slender, hooked prickles.

4. *X. commu'ne* Britton. Similar to No. 3, but the beaks are more hooked and the prickles more numerous.—Pr. Prov.

8. AMBRO'SIA L. RAGWEED

1. *A. artemisiifo'lia* L. (HOGWEED.) Stem erect, 1-3 feet high, branching, hairy. Leaves twice-pinnatifid, the lobes linear,

paler beneath.—Waste places everywhere, but not so common northward.

2. *A. trifida* L. (GREAT RAGWEED.) Stem stouter than No. 1, 2–4 feet high. Leaves opposite, deeply 3-lobed, the lobes oval-lanceolate and serrate.—Low grounds.

3. *A. psilostachya* DC. Similar in habit to No. 1, but the leaves are once-pinnatifid and thicker, and the rootstock is creeping.—Pr. Prov.

9. FRANSE'RIA Cav.

1. *F. acanthicar'pa* Coville. Low and diffuse, hairy. Leaves bipinnatifid, at least the lower ones.—Pr. Prov.

10. TANACE'TUM L. TANSY

1. *T. vulga're* L. (COMMON TANSY.) A very strong-scented herb, 2–4 feet high, smooth. Leaves twice-pinnate, the lobes serrate, as are also the wings of the petiole. Heads densely corymbed.

Var. *crispum* DC. Easily distinguished by its crisper and more incised leaves.—Old gardens and roadsides near dwellings.

2. *T. huronen'se* Nutt. Hairy when young. Heads usually few, and much larger than in *T. vulgare*. Pistillate flowers flattened, instead of terete as in *T. vulgare*.—River-margins, Atl. Prov.

11. ARTEMIS'IA L. WORMWOOD

1. *A. canadensis* Michx. Stem smooth or sometimes hoary with silky down, erect, usually brownish. Lower leaves twice-pinnatifid, the lobes linear.—Shores of the Great Lakes.

2. *A. cauda'ta* Michx. Lobes of the dissected leaves filiform, and the heads small, in a wand-like, long panicle. Disk-flowers perfect but sterile; marginal florets fertile.—Atl. Prov. and Pr. Prov.

3. *A. dracunculoi'des* Pursh. Glabrous or slightly hoary. Stems 2–4 feet high, branching. Leaves mostly entire, linear, sometimes 3-cleft. Heads very numerous, in a compound panicle.—Pr. Prov.

4. *A. glauc'a* Pall. A smaller plant than the last, 1–2 feet high, of stricter aspect. Leaves linear to oblong-lanceolate.—Pr. Prov.

5. *A. vulga'ris* L. (COMMON MUGWORT.) Stem tall, and branching above. Leaves green and smooth above, white-woolly

beneath, pinnatifid, the lobes linear-lanceolate. Heads small, erect, in panicles. Flowers purplish.—Old fields near dwellings.

6. *A. ludoviciana* Nutt. White-woolly throughout. Leaves lanceolate, the upper entire, the lower lobed or toothed, the upper surface sometimes green and smoothish. Heads small, mostly sessile in narrow panicles.—Pr. Prov.

7. *A. bien'nis* Willd. Glabrous; stem strict, 1–3 feet high. Leaves once- or twice-pinnately parted, the lobes linear, acute. Heads in very short, axillary spikes or clusters, crowded in a narrow leafy panicle. Florets all fertile.

8. *A. Absinthium* L. (COMMON WORMWOOD.) Somewhat shrubby, 2–3 feet high. Whole plant silky-hoary. Stem angular-branched, the branches with drooping extremities. Leaves 2–3-pinnately divided, the lobes lanceolate. Heads nodding.—Escaped from gardens in some places.

9. *E. frigida* Willd. (PASTURE SAGE-BRUSH.) A low plant, growing in tufts, white-silky. Leaves dissected into narrowly linear divisions. Heads globose, in racemes.—Pr. Prov.

10. *A. ca'na* Pursh. (SAGE-BRUSH.) Slightly shrubby, 1–2 feet high, much-branched, silvery-hoary. Leaves narrow, tapering to both ends, small, mostly entire. Heads clustered in a leafy narrow panicle, few-flowered.—Pr. Prov.

12. ERECTITES Raf. FIREWEED

1. *E. hieracifolia* Raf. Stem tall, grooved. Leaves sessile, lanceolate, cut-toothed, and auricled.—Common in places recently over-run by fire.

13. GNAPHALIUM L. CUDWEED

1. *G. decurrens* Ives. (EVERLASTING.) Stem erect, 2 feet high, clammy-pubescent, white-woolly on the branches. Heads corymbd. Leaves linear-lanceolate, partly clasping, decurrent.—Fields and hillsides.

2. *G. polycephalum* Michx. (COMMON EVERLASTING.) Stem erect, 1–2 feet high, white-woolly, fragrant. Heads corymbd. Leaves lanceolate, tapering at the base, not decurrent.—Old pastures and woods.

3. *G. uligino'sum* L. (LOW CUDWEED.) Stem spreading, 3-6 inches high, white-woolly. Leaves linear. Heads small in crowded terminal clusters subtended by leaves.—Low grounds.

4. *G. sylvat'icum* L. Erect, usually 9-12 inches high. Leaves linear. Heads axillary, nearly sessile, forming an erect leafy spike. Scales obtuse with a brown bar across each near the top.—Atl. Prov.

14. ANAPH'ALIS DC. EVERLASTING

1. *A. margarita'cea* B. and H. (PEARLY EVERLASTING.) Stem in clusters, downy. Leaves linear-lanceolate, taper-pointed, sessile, tomentose on both sides. Scales of the involucre pearly-white. Heads in corymbs.—Along fences and in open woods.

15. ANTENNA'RIA Gaertn.

1. *A. plantaginifo'lia* Richardson. (PLANTAIN-LEAVED EVERLASTING.) Stem scape-like, 4-12 inches high. Radical leaves spatulate or obovate, 3-nerved; stem-leaves few, linear. Heads small, in a crowded corymb. Involucre green or purplish; styles crimson.—Old pastures and woods.

2. *A. neodioic'a* Greene. Stems slender from mats. Basal leaves 1-nerved, obovate; stem-leaves scale-like, linear. Styles pale.

3. *A. petaloid'ea* Fernald. Stout, 8-16 inches high; stolons procumbent, not leafy. Heads in a loose corymb; involucre brownish or greenish with mostly petaloid bracts.

16. LIA'TRIS Schreb. BLAZING STAR

1. *L. cylindra'cea* Michx. Stem wand-like, 6-18 inches high. Leaves linear, rigid, generally 1-nerved. Heads few, cylindrical. Scales with abruptly mucronate tips. Pappus plumose.—Sandy fields and thickets.

2. *L. spica'ta* Willd. Stem stout and rigid, 2-5 feet high, very leafy. Leaves linear, erect, the lowest 3-5-nerved. Heads crowded in a long spike. Scales obtuse. Pappus not obviously plumose.—Low grounds.

3. *L. scario'sa* Willd. Heads many-flowered, large, few or many. Pappus not obviously plumose. Stem 1-5 feet high, stout, pubescent or hoary. Leaves lanceolate. Scales of the involucre obovate or spatulate, often with coloured tips or edges.

4. *L. puncta'ta* Hook. Heads 4-6-flowered, usually many in a dense spike. Pappus plumose. Stem 10-30 inches high, stout. Scales of the involucre acuminate. Leaves narrowly linear.—Pr. Prov.

17. *VERNO'NIA* Schreb. IRON-WEED

1. *V. altis'sima* Nutt. Tall. Leaves lanceolate to lance-oblong. Heads in an open cyme, the involucre purplish.—Fields and roadsides.

2. *V. noveboracen'sis* Willd. Leaves lanceolate to oblong, pubescent beneath. Heads 30-40-flowered; pappus purplish; bracts of the involucre ovate, with filiform tips.

3. *V. illinoen'sis* Gleason. Leaves oblong-lanceolate. Heads about 40-flowered. Bracts of the involucre rounded, without conspicuous tips; pappus purple.

18. *EUPATO'RIMUM* L. THOROUGHWORT

1. *E. purpu'reum* L. (JOE PYE WEED. TRUMPET-WEED.) Stem tall, 1-6 feet high, and simple. Leaves petioled, 3-6 in a whorl. Flowers purplish or flesh-coloured. Heads in dense corymbs.—Low grounds.

2. *E. perfoli'atum* L. (BONESET.) Stem 1-4 feet high, hairy. Leaves rugose, connate-perfoliate, tapering. Flowers whitish. Corymbs very large.—Low grounds.

3. *E. urticaefo'lium* Reichard. (WHITE SNAKEROOT.) Stem very smooth, commonly branching, 2-3 feet high. Leaves opposite, petioled, broadly ovate, pointed, coarsely serrate. Flowers white, in corymbs.—Low rich woods.

19. *CACA'LIA* L. INDIAN PLANTAIN

1. *C. tubero'sa* Nutt. Stem angled and grooved, tall, 2-6 feet high, from a thick root. Leaves thick, 5-7-nerved, the lower lance-ovate or oval tapering into long petioles.

20. *I'VA* L. MARSH ELDER

1. *I. axilla'ris* Pursh. Stem 1-2 feet high, the nodding heads mostly solitary in the axils of the upper leaves. Leaves small, obovate to oblong-linear, entire, sessile.—Pr. Prov.

2. *I. xanthifo'lia* Nutt. Stem 3-5 feet high, the narrow, spike-like clusters of heads forming a naked panicle. Leaves large, broadly ovate, long-petioled, serrate.—Pr. Prov.

21. SENE'CIO L. GROUNDSEL

1. *S. vulga'ris* L. (COMMON GROUNDSEL.) Ray-florets wanting. Stem low, branching, nearly glabrous. Leaves pinnatifid and toothed, clasping. Flowers yellow, terminal.—Cultivated and waste grounds.

2. *S. visco'sus* L. Ray-florets very minute. Stem viscid-pubescent and strong-scented. Leaves twice-pinnatifid. Flowers bright yellow.—Atl. Prov.

3. *S. aur'eus* L. (GOLDEN RAGWORT. SQUAW-WEED.) Rays 8-12. Stem smooth, or woolly when young, 1-2 feet high. Root-leaves simple, rounded, usually cordate, crenately-toothed, long-petioled. Stem leaves sessile, lanceolate, deeply pinnatifid. Heads in a corymb nearly like an umbel.—Swamps, often in gardens.

4. *S. obova'tus* Muhl. Root-leaves round-obovate, gradually tapering into a winged petiole, and stem-leaves pinnatifid and sessile.

5. *S. Balsam'itae* Muhl. Root-leaves oblong, spatulate, or lanceolate, serrate.

6. *S. integer'rimus* Nutt. Rays conspicuous. Stem woolly-pubescent when young, soon smoothish and green. Leaves oblong-lanceolate or oblong, the upper bract-like. Heads in a naked corymb, the scales green-tipped.—Pr. Prov.

7. *S. ca'nus* Hook. Low, persistently woolly. Leaves small, spatulate to oblong. Rays conspicuous.—Pr. Prov.

8. *S. Jacobae'a* L. (COMMON RAGWEED. STINKING WILLIE.) Flowers golden yellow, the heads forming a spreading corymb. Stem erect, 2-3 feet high, branching, glabrous or somewhat cottony. Leaves numerous, lyrate, bipinnatifid, the lower with broad segments, the upper with linear divisions, all glabrous.—A troublesome weed in the Atl. Prov. Rare westward.

9. *S. platten'sis* Nutt. Stems 2 or 3 from the base or one. Lower leaves petiolate, roundish to oblong-lanceolate, crenate to pinnatifid, glabrate. Inflorescence cymose; rays yellow.

10. *S. palus'tris* Hook. Stem stout, erect, covered with loose hairs. Leaves oblong-lanceolate, irregularly toothed; the upper clasping with heart-shaped base. Heads corymbose; rays yellowish; pappus long, white, copious.—Pr. Prov.

22. *IN'ULA* L. ELE CAMPANE

1. *I. Hele'nium* L. (COMMON ELE CAMPANE.) Stem stout, 2-5 feet high. Root-leaves very large, ovate, petioled. Stem-leaves clasping. Heads large, yellow. Rays numerous, narrow.—Roadsides.

23. *CHRYSOP'SIS* Nutt. GOLDEN ASTER

1. *C. villo'sa* Nutt. Hirsute and villous-pubescent. Leaves narrowly oblong, hoary, bristly-ciliate towards the base. Stem branching, the branches terminating in the single heads.—Dry plains, Pr. Prov.

24. *APLOPAP'US* Cass.

1. *A. spinulo'sus* DC. Low, perennial, branches very minutely hoary-pubescent. Leaves narrow, pinnately or tripinnately lobed, the lobes and teeth bristly, as are also the scales of the involucre.—Dry plains, Pr. Prov.

2. *A. lanceola'tus* Torr. and Gr. Stem simple, with a tuft of coriaceous radical leaves, and bearing 3 to 15 heads. Rays 20 to 50. Scales of the involucre in 3 or 4 unequal series, lanceolate, with greenish tips and whitish base.—Marshy plains, Pr. Prov.

3. *A. acau'lis* Gray, var. *glabra'tus* Eaton. Stems depressed-tufted, from a woody rootstock, glabrous or nearly so. Leaves rigid, persistent, crowded below, a few only on the scape-like, flowering stems, which occasionally branch above and bear 2 or 3 heads; bracts not green-tipped.—Dry gravelly ridges, Pr. Prov.

25. *SOLIDA'GO* L. GOLDEN-ROD

A. Heads clustered in the axils of the feather-veined leaves.

1. *S. squarro'sa* Muhl. Stem stout, 2-5 feet high, simple, hairy above. Scales of the involucre with reflexed herbaceous tips. Leaves large, oblong, serrate, veiny; the lower tapering into a long winged petiole, the upper sessile and entire. Heads in racemose clusters, the whole forming a dense, leafy, interrupted, compound spike.—Rocky woods.

2. *S. bic'olor* L. Stem hoary-pubescent, usually simple. Leaves oval-lanceolate, acute at both ends; the lower oval and tapering into a petiole, serrate. Heads in short racemes in the upper axils, the whole forming an interrupted spike or compound raceme. Ray-florets whitish.—Dry banks and thickets.

3. *S. his'pida* Muhl. Similar to No. 2, but the leaves are oblanceolate to obovate, and the rays are orange-yellow in colour.

4. *S. latifo'lia* L. Stem smooth, angled, zigzag, 1-3 feet high. Leaves broadly ovate or oval, strongly and sharply serrate, pointed at both ends. Heads in very short, axillary clusters. Rays 3 or 4.—Cool woods.

5. *S. cae'sia* L., var. *axilla'ris* Gray. Stem smooth, terete, glaucous, slender, usually branching above. Leaves smooth, lanceolate, pointed, serrate, sessile. Heads in very short clusters in the axils of the leaves.—Rich woods and hillsides.

*B. Racemes terminal, erect, loosely thyrsoid, not one-sided
Leaves feather-veined.*

6. *S. racemo'sa* Green, var. *Gillma'ni* Fernald. Stem low, 6-12 inches high, usually smooth; the heads, peduncles, etc., mostly glutinous. Leaves lanceolate or oblanceolate, serrate or entire, the radical ones petiolate, obtuse, and serrate at the apex.—Rocky banks, not common.

7. *S. macrophyl'la* Pursh. Stems stout, simple, pubescent near the summit. Leaves thin, ovate, with sharp projecting teeth, the lower ones abruptly contracted into long, margined petioles. Heads large, in an oblong raceme, loose and thin, long-pointed. Rays 8-10, long.—Wooded hillsides, chiefly Atl. Prov.

C. Heads in a compound corymb terminating the simple stem, not at all racemose.

8. *S. ohioen'sis* Riddell. Very smooth throughout. Stem slender, reddish, leafy. Radical leaves very long (often a foot), slightly serrate towards the apex, tapering into long, margined petioles; stem-leaves oblong-lanceolate, entire, sessile.—Wet grassy shores of Red Bay, Lake Huron.

9. *S. rig'ida* L. Rough and rather hoary. Stem stout, 2-5 feet high, very leafy. Leaves oval or oblong.—Dry soil, Pr. Prov.

D. Heads in one-sided racemes, spreading or recurved. Leaves not 3-ribbed, but sometimes obscurely 3-nerved.

10. *S. semper'virens* L. Stem stout. Leaves long, lanceolate, thickish, smooth, entire, obscurely 3-nerved. Racemes short, in a terminal panicle. Heads large, showy.—Salt marshes and sea-shores.

11. *S. puber'ula* Nutt. Stem and panicle minutely hoary. Stem-leaves lanceolate, acute, tapering to the base, somewhat smooth, the lowermost spatulate, sparingly toothed. Heads not large, crowded in compact short racemes, which form a long, dense, terminal panicle. Rays 10–14.—Barren soil, Atl. Prov.

12. *S. uligino'sa* Nutt. Smooth. Stem simple, strict. Leaves lanceolate, pointed, smooth, the lower tapering into winged petioles, finely but sparingly serrate, or entire. Racemes crowded and appressed in a close, wand-like panicle. Heads middle-sized. Rays 5–6, small.—Peat-bogs and wet places, Atl. Prov.

13. *S. specio'sa* Nutt. Somewhat resembles the last, but the leaves are oval or ovate, and the stem is taller and stouter (3–6 feet). The rays, also, are larger.—Copses, Atl. Prov.

14. *S. pat'ula* Muhl. Stem tall, smooth, 4-angled. Lower leaves long, 4–6 inches, ovate, serrate, acute, and rough on the upper surface; upper leaves smaller, sharp-pointed. Racemes short and spreading.

15. *S. jun'cea* Ait. Whole plant smooth, 1–4 feet high, rigid, branching above. Lower leaves oval or elliptical-lanceolate, serrate with spreading teeth, pointed, tapering into winged and ciliate petioles; upper ones lanceolate. Racemes very dense, naked, at length elongated and recurved.—Woods and banks.

16. *S. argu'ta* Ait. Stem smooth, angled or furrowed. Leaves large and thin, ovate, doubly serrate; the upper elliptical-lanceolate. Racemes much shorter and looser than in No. 15, and the rays much larger.—Moist woods and thickets.

17. *S. rugo'sa* Mill. Stem rough-hairy, about a foot high. Leaves ovate-lanceolate or oblong, coarsely serrate, veiny, often rugose. Racemes panicked, spreading.—Borders of fields and copses.

18. *S. neglec'ta* Torr. and Gr. Stem smooth, 2–3 feet high, stout. Leaves thickish, smooth both sides, the upper oblong-lanceolate, nearly entire, the lower ovate-lanceolate or oblong,

sharply serrate, tapering into a petiole. Heads rather large. Racemes short and dense, at first erect and scarcely one-sided, at length spreading.—Swamps.

E. *Racemes one-sided and recurved, and the leaves plainly 3-ribbed.*

19. *S. nemoralis* Ait. Stem minutely and closely hoary-pubescent, simple or corymbed. Leaves more or less hoary, obscurely serrate or entire; the lower oblanceolate, somewhat crenate, and tapering into a petiole. Racemes numerous, dense, at length recurved, forming a large panicle.—Dry fields.

20. *S. missouriensis* Nutt. Stem smooth. Leaves linear-lanceolate, tapering to both ends, with rough margins. Racemes densely crowded.—Pr. Prov.

21. *S. canadensis* L. Stem minutely pubescent or glabrous, tall and stout. Leaves lanceolate, serrate, pubescent beneath, rough above. Panicle exceedingly large.—Very common along fences and in moist thickets.

22. *S. serotina* Ait. Stem smooth, stout. Leaves usually lanceolate, taper-pointed, sharply serrate, except at the base, smooth both sides, rough-ciliate. Rays 7–14, rather long. Panicle large, pubescent.—Open thickets and meadows.

Var. *gigantea* Gray. Very tall and the leaves more or less pubescent beneath.—Thickets and low grounds.

F. *Inflorescence a flat-topped corymb.*

23. *S. graminifolia* Salisb. Stem pubescent above, much-branched. Leaves linear-lanceolate, the nerves (3–5) and margins rough-pubescent. Heads in dense corymbed clusters, giving a decidedly characteristic aspect to this species.—Low lake- and river-margins.

26. ARNICA L.

1. *A. mollis* Hook. Soft-hairy. Stem leafy to the top, bearing 1–5 heads. Leaves thin and veiny, toothed; the upper ovate-lanceolate, sessile; the lower narrow, tapering to a margined petiole.—Atl. Prov.

2. *A. foliosa* Nutt. Lanceolate leaves, the upper partly clasping and the lower with tapering bases connate. Stem strict.—Pr. Prov.

3. *A. fulgens* Pursh. Less leafy, low, yellowish-pubescent or villous. Stem simple, bearing a single head as a rule. Leaves thickish, narrowly oblong to lanceolate, the cauline only 1 or 2 pairs.

27. TUSSILA'GO L. COLT'S FOOT

1. *T. Farfara* L. Wet places, chiefly eastward, but also in Ontario.

28. PETASI'TES Hill. SWEET COLT'S FOOT

1. *P. palma'tus* Gray. Leaves rounded, somewhat kidney-shaped, palmately 5-7-lobed, the lobes toothed and cut.—Cedar-swamps and bogs; wet banks of streams.

2. *P. sagitta'tus* Gray. Leaves deltoid-oblong to reniform-hastate, repand-dentate, white-tomentose beneath.—Swamps, Pr. Prov.

29. ASTER L. STARWORT. ASTER

A. *Leaves, at least the lower ones, heart-shaped and petioled.*

1. *A. divarica'tus* L. Rays 6-9, white or nearly so. Heads in corymbs. Stems slender, 1-2 feet high, zigzag. Leaves thin, smoothish, sharp-pointed, coarsely serrate, all the lower ones on slender naked petioles.—Woodlands.

2. *A. macrophyl'lus* L. Rays white or bluish. Stem stout, 2-3 feet high. Leaves thickish, rough, finely serrate, the lower long-petioled, large and cordate, especially on the sterile plants. Inflorescence somewhat viscid-glandular. Heads in closer corymbs than in No. 1.—Woodlands.

3. *A. azu'reus* Lindl. Rays 10-20, bright blue. Heads racemed or panicled. Stem roughish, erect, racemose-compound above. Leaves entire or nearly so, rough; the lower ovate-lanceolate, on long petioles; the upper lanceolate or linear, sessile. The latest flowering of our Asters.—Dry soil.

4. *A. Shortii* Lindl. Rays bright blue. Stem slender, nearly smooth. Leaves lanceolate to ovate-lanceolate; all but the uppermost cordate and on naked petioles.

5. *A. undula'tus* L. Rays bright blue. Heads racemed or panicled. Stem hoary with close pubescence, spreading. Leaves with somewhat wavy margins, entire or nearly so, ovate or ovate-lanceolate, roughish above, downy beneath; the lowest cordate, on

margined petioles; the upper with winged short petioles clasping at the base, or sessile.—Dry woods.

6. *A. cordifo'lius* L. Rays pale blue or nearly white. Heads small, profuse, paniced. Stem much-branched, nearly glabrous. Leaves thin, sharply serrate, the lower on slender ciliate petioles. Scales of the inversely conical involucre all appressed, and tipped with very short, green points.—Woods and along fences.

7. *A. Lowriea'nus* Porter, var. *lanceola'tus* Porter. Resembles No. 6. Leaves glaucous beneath, smooth, often greasy; the lower lanceolate to ovate, truncate or narrowed at the base.

8. *A. sagittifo'lius* Wedemeyer. Rays pale blue or purple. Heads small, in dense compound racemes or panicles. Stem smooth or nearly so, erect, with ascending branches. Leaves conspicuously serrate, ovate-lanceolate, pointed, pubescent, the lowest on long, margined petioles, the upper lanceolate or linear, pointed at both ends. Scales of the oblong involucre linear, tapering into awl-shaped, slender and loose tips.—Thickets and along fences.

9. *A. Lindleya'nus* Torr. and Gr. Rays pale violet. Stem rather stout, smooth or sparsely pubescent. Leaves conspicuously serrate. Root-leaves and lowest stem-leaves ovate, more or less cordate, with margined petioles; uppermost sessile, and pointed at both ends. Heads rather small, in a loose thyrse or panicle. Scales linear, green-tipped.—Open barren grounds.

B. Upper leaves all sessile or clasping by a heart-shaped base; lower ones not heart-shaped.

10. *A. lae'vis* L. Rays large, sky-blue. Very smooth throughout. Heads in a close panicle. Leaves lanceolate or ovate-lanceolate, chiefly entire, rough on the margins, the upper ones clasping by an auricled base.—Dry woods.

11. *A. oblongifo'lius* Nutt. Rays violet-purple; minutely glandular-pubescent. Leaves oblong-lanceolate, partly clasping. Scales of the involucre nearly equal, linear, appressed.

12. *A. no'vae-an'gliae* L. Rays many, narrow, violet-purple, showy; heads large. Involucre of many slender, equal scales, apparently in a single row, clammy. Stem stout, 3-8 feet high, hairy, corymbd above. Leaves very numerous, lanceolate, entire,

clasping by an auricled base, pubescent.—River-banks and borders of woods.

13. *A. punic'eus* L. Rays long, lilac-blue. Stem stout, tall and hairy. Scales of the involucre narrowly linear, loose, in about two rows, not glandular-hairy. Stem 3–6 feet high, stout, rough-hairy, usually purple below. Leaves oblong-lanceolate, clasping by an auricled base, sparingly serrate in the middle, rough above, smooth beneath, pointed.—Swamps; usually clustered.

14. *A. tardiflorus* L. Rays pale violet. Stem glabrous or nearly so, 1–2 feet high. Leaves lanceolate or oblong-lanceolate, acuminate, with tapering and somewhat auricled base, slightly if at all clasping. Heads few, as a rule. Involucre loose, the outer scales leaf-like.—Atl. Prov.

C. None of the leaves heart-shaped; those of the stem sessile, tapering at the base.

15. *A. rad'ula* Ait. Rays light violet. Stem simple, with a corymb at the top. Radical leaves all tapering into margined petioles. Stem-leaves sessile, oblong-lanceolate, pointed, sharply serrate in the middle, very rough both sides, rugose. Scales of the bell-shaped involucre oblong, appressed, with slightly spreading, herbaceous tips.—Low grounds, Atl. Prov.

16. *A. ericoi'des* L. Rays white or nearly so. Stem smooth or nearly so, 1–3 feet high, the branchlets or peduncles racemose on the upper side of the spreading branches. Lowest leaves oblong-spatulate, the others linear-lanceolate or linear-awl-shaped.—Barren soil, and margins of lakes and rivers.

Var. *villosus* Torr. and Gray. Hairy stems and leaves.

17. *A. multiflorus* Ait. Rays white, 10–20. Stem pale or hoary with minute pubescence, 1 foot or more in height, bushy. Leaves crowded, linear, with rough margins; the upper partly clasping. Heads small ($\frac{1}{4}$ inch long), crowded on the racemose branches. Scales of the involucre with spreading green tips.—Dry soil.

18. *A. commuta'tus* Gray. Similar to *A. multiflorus* except that the heads are longer ($\frac{1}{3}$ inch long) and the rays more numerous (20–30).—Pr. Prov.

purple
aster

19. *A. dumo'sis* L. Rays purple to blue. Stem smooth or nearly so. Leaves linear or oblong, entire. Scales of the involucre with abrupt green tips.

20. *A. Tradescan'ti* L. Rays white or purplish. Scales of the involucre narrowly linear, in 3 or 4 rows. Heads small, very numerous, in 1-sided, close racemes on the branches. Stem 2-4 feet high, much-branched, smooth. Leaves linear-lanceolate, with long slender points, the larger ones with a few remote teeth in the middle.—Low grounds.

21. *A. lateriflo'rus* Britton. Rays pale blue or whitish. Involucre nearly as in No. 20. Stem more or less pubescent, much-branched. Heads small, in loose, 1-sided racemes on the spreading branches. Leaves lanceolate, acute at each end, sharply serrate in the middle.—Low grounds.

22. *A. panicula'tus* Lam. Rays pale blue or whitish. Scales of the involucre linear-awl-shaped. Stem stout, smooth or nearly so, with numerous leafy branches. Heads medium-sized, scattered, loosely paniculate. Leaves smooth, oblong to lanceolate, tapering at both ends, the lower serrate.—Moist and shady banks.

23. *A. jun'ceus* Ait. Rays light purple or white. Stem slender, 1-3 feet high, simple, with few small heads, or loosely branching. Leaves narrow, entire, or the lower sparingly denticulate. Scales of the involucre small, narrow, in 2 or 3 rows.—Bogs and wet places.

24. *A. vimin'eus* Lam. Rays white or nearly so. Stem 2-5 feet high, smooth, bushy. Leaves linear or narrowly lanceolate, the larger ones sparingly serrate in the middle with fine teeth. Heads very numerous, in 1-sided racemes on short branchlets. Scales of the involucre narrowly linear, in 3 or 4 rows.—Moist banks.

25. *A. nemora'lis* Ait. Rays lilac-purple, elongated. Stem slender and leafy, the upper branches terminating in 1-flowered, nearly naked peduncles. Leaves small, rigid, narrowly lanceolate, nearly entire, with revolute margins.—Swamps, Atl. Prov. and Muskoka.

26. *A. ptarmicoi'des* Torr. and Gr. Rays pure white. Stems clustered, generally a foot high, each bearing a flat corymb of small heads. Leaves linear-lanceolate, acute, rigid, entire, mostly 1-3-nerved, with rough margins.—Dry or gravelly hills. Our earliest Aster.

27. *A. acumina'tus* Michx. Rays white or faintly purple. Stem about 2 feet high, somewhat hairy, zigzag, paniced-corymbose at the top. Leaves large, thin, oblong, lanceolate, pointed, coarsely toothed towards the apex, entire at the base.—Cool sandy woods; mostly eastward.

28. *A. umbella'tus* Mill. Pappus double, the inner of long capillary bristles, the outer of short and rigid bristles. Rays white. Stem smooth, leafy to the top, tall, simple. Leaves lanceolate, long-pointed. Heads small, very numerous, in compound flat corymbs.—Moist thickets.

30. ERIG'ERON L. FLEABANE

1. *E. canaden'sis* L. (HORSE-WEED. BUTTER-WEED.) Rays white, but very inconspicuous, shorter than their tubes. Heads very small, numerous, in paniced racemes. Pappus simple. Stem 1-5 feet high, erect and wand-like, bristly-hairy. Leaves linear, mostly entire, the radical lobed.—Common in burnt woods and new clearings.

2. *E. acris* L., var. *asteroi'des* DC. Rays purplish or bluish, about the same length as the copious simple pappus. Heads several or many, small, at length corymbose, hirsute. Stem 10-20 inches high, pubescent or smoothish. Leaves mostly lanceolate, entire. A set of pistillate flowers within the circle of ray-flowers.—Atl. Prov. and Pr. Prov.

3. *E. pulchel'lus* Michx. (ROBIN'S PLANTAIN.) Rays bluish-purple, numerous. Heads medium-sized, few, on slender corymbose peduncles. Pappus simple. Stem hairy, producing offsets from the base. Radical leaves spatulate or obovate, toothed above the middle; stem-leaves oblong, few, sessile or partly clasping, entire.—Thickets.

4. *E. hyssopifo'lius* Michx. Rays rose-purple or whitish. Stem slightly pubescent, slender, 6-12 inches high, from slender rootstocks. Leaves very many, short, linear. Heads small, terminating the slender naked branches. Pappus simple.—Atl. sea-coast.

5. *E. caespito'sus* Nutt. Stem dwarf, tufted, from a stout rootstock, more or less hoary-pubescent. Rays white, 40-50, narrow.—Pr. Prov.

6. *E. philadelphicus* L. (COMMON FLEABANE.) Rays rose-purple, very numerous and narrow. Heads small, few, in corymbs. Pappus simple. Stem hairy, with numerous stem-leaves. Radical leaves spatulate and toothed; the upper ones clasping by a heart-shaped base, entire.—Moist grounds.

7. *E. ramo'sus* BSP. (DAISY FLEABANE.) Rays white, conspicuous, numerous. Pappus plainly double. Stem and leaves roughish with minute appressed hairs, or nearly smooth. Lower leaves spatulate and slender-petioled, entire or nearly so, the upper lanceolate, scattered.—Dry fields and meadows.

8. *E. glabell'us* Nutt. Rays purple, very many, much longer than the hoary-hispid involucre. Stem 6–15 inches high, stout, smooth below, bearing 1–7 large heads on the naked summit. Pappus double. Leaves smooth but ciliate, the upper oblong-lanceolate and pointed, sessile or clasping; the lower petiolate, spatulate.—Pr. Prov.

9. *E. an'nuus* Pers. (LARGER DAISY FLEABANE.) Rays white, tinged with purple. Pappus double. Stem stout, rough with spreading hairs. Leaves coarsely toothed; the lower ovate, tapering into a margined petiole; the upper ovate-lanceolate. Heads corymbed.—Fields and meadows.

31. GRINDE'LIA Willd.

1. *G. squarro'sa* Dunal. Leaves spatulate to linear-oblong. Heads large, terminating the leafy branches, involucre very resinous.—Dry prairies, Pr. Prov. and Ont.

32. GUTIERRE'ZIA Lag.

1. *G. Saro'thræ* B. and R. Not more than 8 inches high.—Dry plains, Pr. Prov.

33. HELE'NIUM L. SNEEZE-WEED

1. *H. autumnna'le* L. (SNEEZE-WEED.) Stem nearly smooth. Leaves lanceolate, toothed. Disk yellow, globular.—Low river- and lake-margins.

34. CHRYSAN'THEMUM L. OX-EYE DAISY

1. *C. Leucan'themum* L. (OX-EYE DAISY. WHITE-WEED.) Stem erect, naked above. Bearing a single large head. Leaves

pinnatifid or cut-toothed, the lowest spatulate, the others partly clasping.—Pastures and old fields.

2. *C. Parthe'nium* Bernh. (FEVERFEW.) Stem branching, leafy. Leaves twice-pinnately divided, the divisions ovate, cut. Heads corymbed.—Escaped from gardens.

3. *C. Balsam'ita* L., var. *tanacetoi'des* Boiss. (COSTMARY.) A garden-escape, smooth, with pleasant odour. Leaves pale, oblong, somewhat toothed. Heads small, pale yellow; rays wanting.

35. *BELLIS* L. DAISY

1. *B. perennis* L. The true Daisy, a native of the Old World, is a low stemless herb. It is an uncommon garden escape. The heads are many-flowered with numerous pistillate rays. The scales of the involucre equal, in about 2 rows, herbaceous. Receptacle conical. Pappus wanting.

36. *MATRICA'RIA* L. WILD CHAMOMILE

1. *M. inodo'ra* L. Leaves twice-pinnately divided into very narrow lobes. Heads large, naked-peduncled, the rays many and long.—Chiefly Atl. Prov.

2. *M. suave'olens* Buchenau. (PINEAPPLE-WEED.) A low plant with the odour of pineapple when bruised. Leaves 2-3-pinnately parted into narrow lobes. Heads rayless, without pappus, and somewhat conical.

37. *AN'THEMIS* L. CHAMOMILE

1. *A. Cot'ula* L. (MAY-WEED.) Stem branching. Leaves thrice-pinnate, finely dissected. Odour disagreeable. Rays soon reflexed.—Roadsides everywhere.

2. *A. arven'sis* L. (CORN CHAMOMILE.) Resembling the last, but the leaves are not so finely dissected, and the odour not so unpleasant.—Atl. Prov., rare.

38. *RUDBECK'IA* L. CONE-FLOWER

1. *R. lacinia'ta* L. Rays linear, 1-2 inches long, drooping. Disk greenish-yellow. Stem tall, smooth, branching. Lowest leaves pinnate, of 5-7-lobed leaflets; upper ones 3-5-parted, or the uppermost undivided and generally ovate. Heads terminal, long-peduncled.—Swamps.

2. *R. hir'ta* L. (BLACK-EYED SUSAN.) Rays bright yellow. Disk purplish-brown. Stem very rough-hairy, naked above, bearing single large heads. Leaves 3-ribbed, the lowest spatulate, narrowed into a petiole, the upper ones sessile.—Meadows.

39. LEP'ACHYS Raf.

1. *L. columna'ris* T. and G. Rays yellow or purplish; disk grayish. Receptacle oblong. Chaff thickened and bearded at the tip. Pappus none or of 2 teeth. Stem branching from the base, 1-2 feet high. Leaves pinnately divided, the divisions 5-9, narrow. Heads single, on the naked branches.

2. *L. pinna'ta* T. and G. Rays light yellow, drooping and long (2 inches). Hoary with minute pubescence. Leaflets 3-7, lanceolate, acute.

40. HELIAN'THUS L. SUNFLOWER

A. Annuals. Leaves alternate. Receptacle flat. Disk brownish.

1. *H. an'nuus* L. (COMMON SUNFLOWER.) Tall, rough. Leaves 3-ribbed, ovate, serrate. Scales of the involucre long-pointed, ciliate.—Escaped from cultivation.

2. *H. petiola'ris* Nutt. More slender, 1-3 feet high. Leaves narrow, mostly entire. Scales seldom ciliate.—Pr. Prov.

B. Perennials. Receptacle convex. Lower leaves usually opposite.

(1) *Disk dark.*

3. *H. scaber'rimus* Ell. Stem tall and stout, rough. Leaves very thick and rigid, rough both sides, oblong-lanceolate, pointed at both ends, the lowest oval, 3-nerved. Rays 20-25. Pappus of 2 large and often several small scales.—Pr. Prov.

(2) *Disk yellow.*

4. *H. Nuttall'ii* Torr. and Gr. Stem slender, smooth, simple. Leaves lanceolate or linear. Pappus-scales long and narrow.—Pr. Prov.

5. *H. strumo'sus* L. Stem 3-6 feet high, smooth below. Leaves broadly lanceolate, rough above and whitish beneath, pointed, serrate with small appressed teeth, short-petioled. Rays about 10.—Moist copses and low grounds.

6. *H. divarica'tus* L. Stem 1-4 feet high, smooth, simple or forking above. Leaves all opposite, widely spreading, sessile, rounded or truncate at the base, ovate-lanceolate, 3-nerved, long-pointed, serrate, rough on both sides. Heads few, on short peduncles. Rays about 12.—Open thickets and dry plains.

7. *H. decapet'alus* L. Stem 3-6 feet high, branching, smooth below, rough above. Leaves thin, green on both sides, ovate, coarsely serrate, pointed, abruptly contracted into short, margined petioles. Rays usually 10.—Thickets and river-banks.

8. *H. gigante'us* L. Stem tall, hairy or rough, branching above. Leaves lanceolate, pointed, serrate, very rough above, hairy below, narrowed and ciliate at the base. Heads somewhat corymbed, not large. Disk yellow; rays pale yellow, 15-20.—Low grounds.

9. *H. Maximilia'ni* Schrad. Habit similar to No. 8, but usually not branching above. Heads terminal and axillary, with longer and more rigid bracts.

10. *H. tubero'sus* L. (JERUSALEM ARTICHOKE.) Escaped from cultivation in some places. It is at once recognized by its tubers.

41. ACTINOM'ERIS Nutt.

1. *A. alternifo'lia* DC. Stem hairy, tall, commonly winged above. Leaves oblong to ovate-lanceolate, pointed at both ends.

42. COREOP'SIS L. TICKSEED

1. *C. tincto'ria* Nutt. Stem smooth, 2-3 feet high. Leaves once- or twice-pinnately divided, the lobes narrow. Pappus none. Achenes oblong. Rays yellow, brown-tinted.—Pr. Prov.

2. *C. verticilla'ta* L. Glabrous. Leaves sessile, divided into 3 sessile leaflets (appearing whorled), these pinnately dissected into very narrow divisions.—Damp places.

3. *C. trip'teris* L. Smooth. Stem tall, 3-8 feet high, corymbed above. Leaves petioled, pinnately divided, the divisions lanceolate, acute, entire. Disk turning brownish.—Damp places.

43. GAILLARD'IA Foug.

1. *G. arista'ta* Pursh. Rough-hairy, about 2 feet high. Leaves lanceolate to oblanceolate, entire to coarsely pinnatifid. Disk brownish-purple.—Dry soil, Pr. Prov.

44. *BIDENS* L. BUR-MARIGOLD

1. *B. frondo'sa* L. (COMMON BEGGAR-TICKS.) Rays very small, yellow. Achenes flat, wedge-obovate, ciliate on the margins with bristles pointing upwards, 2-awned. Stem tall, branched. Leaves thin, long-petioled, pinnately 3-5-divided, the leaflets ovate-lanceolate, pointed, serrate.

2. *B. vulga'ta* Greene. Stem tall, often 4 feet high. Leaves pinnatifid. Heads large, $\frac{1}{2}$ –1 inch in diameter, with outer bracts sometimes longer than the disk. Rays pale yellow.—Pr. Prov.

3. *B. conna'ta* Muhl. (SWAMP BEGGAR-TICKS.) Rays usually none. Achenes flat, narrowly wedge-shaped, 2-4 awned, ciliate with minute bristles pointing downwards. Stem 1-2 feet high, smooth. Leaves lanceolate, pointed, serrate, tapering and connate at the base, the lowest often 3-parted and decurrent on the petiole. —In shallow water and low grounds.

4. *B. como'sa* Wiegand. Stem glabrous, stout, 2 feet high. Leaves elliptical on winged petioles, serrate. Rays wanting; corollas 4-toothed, pale yellow; achenes olive or brown; awns nearly $\frac{3}{4}$ as long as the achene.

5. *B. cer'nua* L. (SMALLER BUR-MARIGOLD.) Rays short, pale yellow. Achenes flat, wedge-obovate, 4-awned, ciliate with bristles pointing downwards. Stem nearly smooth, 5-10 inches high. Leaves all simple, lanceolate, unequally serrate, hardly connate. Heads nodding.—Wet places.

6. *B. lae'vis* BSP. (LARGE BUR-MARIGOLD.) Rays an inch long, showy, golden yellow. Achenes wedge-shaped, 2-4-awned, bristly downwards. Stem smooth, 6-30 inches high, erect or ascending. Leaves lanceolate, tapering at both ends, rarely connate, regularly serrate.—Swamps and ditches.

7. *B. Beck'ii* Torr. (WATER MARIGOLD.) Aquatic. Stems long and slender. Immersed leaves dissected into fine, hairy-like divisions; those out of water lanceolate, slightly connate, toothed. Rays showy, golden yellow, larger than the involucre. Achenes linear, bearing 4-6 very long awns barbed towards the apex.—Ponds and slow streams.

8. *B. trichosperm'a* Britton, var. *tenuil'oba* Britton. (TICKSEED SUNFLOWER.) Stem smooth, branching. Rays golden-yellow.

Leaves short-petioled, pinnately divided, the segments serrate, and narrowly linear. Achenes narrowly wedge-oblong, 2-toothed.—Swamps.

45. *HELIOP'SIS* Pers. OX-EYE

1. *H. helianthoides* Sweet. Stem smooth, slender, branching. Leaves ovate-lanceolate, acute, sharply serrate, on slender petioles. Heads showy; peduncles elongated.—Dry open thickets.

2. *H. scabra* Dunal. Roughish, especially the leaves. Pappus of 2 or 3 teeth, or a mere chaff-like border.—Pr. Prov.

46. *ACHILLE'A* L. YARROW

1. *A. millefolium* L. (MILFOIL.) Stems simple. Leaves dissected into fine divisions. Corymb flat-topped. Rays only 4 or 5, short.—Fields and along fences; very common.

2. *A. Ptarmica* L. (SNEEZE-WORT.) Leaves simple, lance-linear, serrate. Corymb loose. Rays 8–12, much longer than the involucre.—Atl. Prov.

47. *POLYMNIA* L. LEAF-CUP

1. *P. canadensis* L. A coarse, clammy-hairy herb. Lower leaves opposite, petioled, pinnatifid; the upper alternate, angled or lobed. Heads small; rays pale yellow.—Shaded ravines.

48. *SILPHIUM* L. ROSIN-PLANT

1. *S. perfoliatum* L. (CUP-PLANT.) Stem stout, square, 4–8 feet high. Leaves ovate, coarsely toothed, the upper ones united by their bases.—South-western Ontario.

2. *S. terebinthinaeum* Jacq. (PRAIRIE DOCK.) Stem tall, round, naked above, smooth. Radical leaves sometimes 2 feet long, rough-hairy, coarsely serrate, on slender petioles. Heads small, loosely paniced.—Open woods and grassy banks.

3. *S. trifoliatum* L. Stem slender, smooth, often glaucous. Stem-leaves lanceolate, in whorls of 3 or 4, the uppermost opposite; none of the leaves large.

49. *GALINSO'GA* R. and P.

1. *G. parviflora* Cav., var. *hispidula* DC. Plant 1 foot or more in height. Leaves ovate, thin, 3-nerved, toothed. Pubescence of stiff hairs. Pappus-scales in disk-florets whitish and bristle-tipped.—Roadsides and waste places.

50. KRIG'IA Schreber. DWARF DANDELION

1. *K. amplexicaulis* Nutt. (CYNTHIA.) Roots fibrous. Stem-leaves 1-2, oblong or lanceolate-spatulate, clasping, mostly entire, the radical ones on short winged petioles. Peduncles 2-5.

51. LAP'SANA L. NIPPLEWORT

1. *L. communis* L. Very slender and branching. Leaves angled or toothed. Heads small, loosely paniced.—Borders of springs.

52. CICHORIUM L. SUCCORY. CICHORY

1. *C. In'tybus* L. Stem-leaves oblong or lanceolate, partly clasping; radical ones runcinate.—Roadsides and waste places.

53. LEONTODON L. FALL DANDELION

1. *L. autumnalis* L. (FALL DANDELION.) Leaves lanceolate, laciniate-toothed or pinnatifid. Scape branched.—Roadsides and waste places.

54. HIERACIUM L. HAWKWEED

1. *H. canadense* Michx. (CANADA HAWKWEED.) Heads large. Stem simple, leafy, corymbed, 1-3 feet high. Peduncles downy. Leaves ovate-oblong with a few coarse teeth, somewhat hairy, sessile, or the uppermost slightly clasping. Achenes tapering towards the base.—Dry banks and plains.

2. *H. vulgatum* Fries. Tall slender stems, bearing few to many small stem-leaves and large, ovate-lanceolate, basal leaves, which are usually purple-blotched and remotely toothed. Heads one to several and 1 inch or more wide.

3. *H. scabrum* Michx. (ROUGH HAWKWEED.) Heads small. Stem stout, 1-3 feet high, rough-hairy, corymbose. Peduncles or involucre densely pubescent and usually covered with dark glands. Achenes not tapering.—Sandy woods and thickets.

4. *H. Gronovii* L. (HAIRY HAWKWEED.) Heads small. Stem wand-like, leafy and very hairy below, naked above, forming a long and narrow panicle. Achenes with a very tapered summit.—Dry soil.

5. *H. venosum* L., (RATTLESNAKE-WEED.) Scape smooth and naked (or bearing one leaf), and a loose corymb of very slender peduncles.

6. *H. panicula'tum* L. Stem slender, leafy, diffusely branched, hairy only below. Heads very small, in a loose panicle, on slender diverging pedicels, 10-20-flowered. Achenes short, not tapering above. Leaves lanceolate, acute at both ends.—Open woods.

7. *H. auranti'acum* L. (DEVIL'S PAINT-BRUSH.) A hirsute species, with clustered heads of deep-orange-red flowers on a simple peduncle, leafy at the base. Leaves oblanceolate.

55. CRE'PIS L.

1. *C. runcina'ta* Torr. and Gr. Stem 1-2 feet high, glabrous. Radical leaves obovate-oblong to oblong-lanceolate, with short lobes or teeth. Cauline leaves absent or very small. Involucre pubescent, sometimes it and the upper part of the scape glandular.—Pr. Prov.

56. PRENAN'THES L. RATTLESNAKE-ROOT

1. *P. alba* L. (WHITE LETTUCE.) Heads 8-12-flowered. Pappus deep cinnamon-coloured. Stem 2-4 feet high, smooth and glaucous, corymbose-paniculate. Leaves triangular-halberd-shaped, or 3-5-lobed, the uppermost oblong and undivided.—Rich woods.

2. *P. altiss'ima* L. (TALL WHITE LETTUCE.) Heads 5-6-flowered. Pappus pale straw-coloured. Stem taller but more slender than in No. 1, with a long leafy panicle at the summit. Leaves usually ovate-heart-shaped, frequently 3-5-lobed.—Rich woods.

3. *P. racemo'sa* Michx. Heads about 12-flowered. Involucre and peduncles hairy. Stem wand-like, smooth. Leaves oval or oblong-lanceolate, slightly toothed. Heads crowded in a long and narrow, interruptedly spiked panicle. Pappus straw-coloured; flowers purplish.

4. *P. mainen'sis* Gray. Leaves as in the last, but the radical ones ovate and more abruptly narrowed to the short petiole. Heads 8-12-flowered, persistently drooping on slender pedicels.—Atl. Prov.

57. LYGODES'MIA Don.

1. *L. jun'cea* Don. Much-branched from the base, the branches closely erect and rigid, rush-like, terminating in erect heads of pinkish flowers. Leaves small, the lower lance-linear, the upper scale-like.—Pr. Prov.

58. AGOS'ERIS Raf.

1. *A. cuspidata* Steud. Scape a foot high. Leaves lanceolate, tapering to a sharp point, entire, woolly on the margins. Achenes beakless.—Pr. Prov.

2. *A. glauca* Steud. Scape 1-2 feet high. Leaves varying from entire to dentate or laciniate. Achenes long-beaked.—Pr. Prov.

59. TARAX'ACUM Ludwig. DANDELION

1. *T. officinale* Weber. (COMMON DANDELION.) Outer involucre reflexed. Leaves runcinate.—Fields everywhere.

60. LACTU'CA L. LETTUCE

1. *L. canadensis* L. (WILD LETTUCE.) Heads numerous, in a long and narrow, naked panicle. Stem stout, smooth, hollow, 4-9 feet high. Leaves mostly runcinate, partly clasping, pale beneath; the upper entire. Achenes longer than their beaks.—Borders of fields and thickets.

Var. *montana* Britton. Entire oblanceolate leaves.

2. *L. scariola* L. (PRICKLY LETTUCE.) Stem below sparingly bristly. Leaves vertical, spinulose-denticulate, oblong or lanceolate, prickly on the midrib below. Heads small, 6-12-flowered.—Waste places.

Var. *integra* Gren. and Godr. Leaves oblong and only the lowest, if any, pinnatifid.

3. *L. integrifolia* Bigel. Stem 3-6 feet high; leaves all undivided, entire or slightly toothed. Flowers pale yellow, cream-colour, or purple—Dry soil.

4. *L. hirsuta* Muhl. Leaves runcinate, the midrib beneath often sparingly bristly-hairy. Flowers yellowish-purple, rarely white.—Dry soil.

5. *L. pulchella* DC. (BLUE LETTUCE.) Stems simple, 1-3 feet high. Leaves oblong-lanceolate, sessile, glabrous, entire or pinnatifid. Heads large; the erect peduncles forming a raceme. Flowers blue; achenes with a short beak.—Pr. Prov.

6. *L. spicata* Hitchc. Stem tall and very leafy. Heads in a dense compound panicle. Flowers bluish to cream-colour.—Borders of damp woods, and along fences.

61. SON'CHUS L. SOW-THISTLE

1. *S. olera'ceus* L. (COMMON SOW-THISTLE.) Stem-leaves runcinate, slightly toothed with soft spiny teeth, clasping; the auricles acute.—Manured soil about dwellings.

2. *S. asper* Hill. (SPINY-LEAVED SOW-THISTLE.) Leaves hardly lobed, fringed with soft spines, clasping; the auricles rounded. Achenes margined.—Same localities as No. 1.

3. *S. arven'sis* L., (FIELD SOW-THISTLE.) Large, bright-yellow flowers and bristly involucre and peduncles. Perennial with creeping rootstocks.

62. TRAGOPO'GON L. GOAT'S BEARD. SALSIFY

1. *T. pratensis* L. (YELLOW GOAT'S BEARD.) Spreading westward along the railway lines. Flowers yellow. Peduncle little thickened below the head.

2. *T. porrifo'lius* L. (SALSIFY.) Stem 2-3 feet high. Peduncle thickened below the head. Flowers purple.

GLOSSARY

A

Abortive, defective or barren.
Acaulescent, apparently without a stem.
Achene, a dry, indehiscent, 1-seeded fruit, with the pericarp free from the seed.
Achlamydeous, without calyx and corolla.
Acicular, needle-shaped.
Acuminate, with a long tapering extremity.
Acute, sharp at the end.
Adnate, grown fast to.
Alternate, not opposite.
-androus, at the end of a word, refers to stamens.
Angiospermous, having seeds inclosed in an ovary.
Annual, lasting one year or season only.
Anterior, in front; away from the axis; the same as *lower* and *outer*.
Anther, the pollen-bearing part of the stamen.
Apetalous, without petals.
Appressed, lying close and flat.
Aquatic, growing in water.
Aril, an extra coat or appendage of certain seeds.
Ascending, growing upwards in a slanting direction.
Attenuate, tapering and becoming very slender.
Auricle, an ear-like appendage or lobe.
Auriculate, having auricles.

Awl-shaped, narrowed upwards from the base to a sharp rigid point.
Awn, a bristle-like appendage.
Axil, the angle on the upper side of a leaf where it joins the stem.
Axile, central; in the axis.
Axillary, growing from an axil.
Axis, the central line.

B

Baccate, berry-like.
Barbed, with short sharp points, usually pointing backwards.
Beaked, tipped with a distinct beak or point.
Bearded, having long or stiff hairs.
Berry, a pulpy fruit like a currant or gooseberry.
Bi-, a prefix meaning twice, or two.
Biennial, lasting two years or seasons.
Bifid, two-cleft.
Bilabiate, two-lipped.
Bladdery, thin and inflated.
Blade, the broad part of a leaf.
Bract, a modified leaf near a flower or flower-cluster.
Bractlet, a secondary bract.
Bristle, a stiff hair.
Bud, an undeveloped branch or flower.
Bulb, a fleshy underground mass, consisting chiefly of leaf-scales or coats.
Bulblet, a small bulb, especially one borne on the stem.

Bulbous, having the appearance of a bulb.

C

Caducous, falling off very early.

Calyx, the outer set of flower-leaves, often coloured like a corolla.

Campanulate, bell-shaped.

Canescent, grayish-hoary.

Capitate, like a head.

Capsule, any dry, dehiscent, syncarpous fruit.

Carinate, having a keel-like ridge or projection.

Carpel, one of the parts which, whether separate or united, make up the pistil.

Cartilaginous, firm and tough.

Caryopsis, a grain or fruit like that of a grass.

Catkin, a slender, spike-like cluster of usually imperfect flowers, as in Willow, etc.

Caulescent, having a manifest stem.

Cauline, belonging to the stem.

Cespitose, growing in tufts.

Chaff, a small thin scale or bract, becoming dry and membranous.

Channelled, grooved lengthwise.

Ciliate, with hairs on the edge.

Cinereous, ash-coloured.

Circumscissile, opening by a horizontal circular line.

Clavate, club-shaped, thicker at the top.

Cleistogamous, applied to certain flowers which are fertilized in the bud, without opening.

Cleft, cut to about the middle.

Climbing, ascending by laying hold of neighbouring objects by means of tendrils, etc.

Cohesion, the union of similar parts.

Coma, a tuft of hairs on a seed.

Compressed, flattened.

Conduplicate, folded together lengthwise.

Confluent, blending together into one.

Coniferous, cone-bearing.

Connate, grown together.

Connective, the middle portion of an anther connecting the cells.

Connivent, converging, approaching each other.

Convolute, rolled up lengthwise.

Cordate, indented at the base.

Coriaceous, leathery in texture.

Corm, a solid bulb, like that of Indian-Turnip.

Corolla, the inner of the two sets of the perianth.

Corymb, a flat-topped flower-cluster with pedicels arising at different points on the stem, the outer flowers opening first.

Corymbose, in corymbs; corymb-like.

Cotyledons, the seed-leaves.

Creeping, trailing on the ground and striking root at intervals.

Crenate, with rounded teeth.

Crenulate, finely crenate.

Crested, bearing an appendage like a crest.

Crown, an appendage in the throat of a corolla.

Culm, the peculiar stem of a grass or sedge.

Cuneate, wedge-shaped, the broad end upwards.

Cuspidate, tipped with an abrupt but distinctly tapering point.

Cylindraceous, somewhat cylindrical.

Cyme, a flat-topped flower-cluster, with the central flowers opening first.

Cymose, bearing cymes, or cyme-like.

D

Deciduous, falling off; not persistent.

Decomound, more than once compound or divided.

Decumbent, reclining.

Decurrent, applied to a leaf when the lobes at the base extend down the sides of the stem.

Decussate, with opposite pairs at right angles to each other.

Deflexed, turned abruptly downwards.

Dehiscent, splitting open to allow the contents to escape.

Deltoid, triangular, the base downwards.

Dentate, toothed, the teeth pointing outwards.

Denticulate, minutely dentate.

Depressed, flattened from above.

Di-, twice or two.

Diadelphous, with the filaments cohering in two sets.

Diandrous, having two stamens.

Dichotomous, forking regularly by pairs.

Dicotyledonous, having two cotyledons.

Didynamous, of stamens, when in two pairs of different lengths.

Diffuse, spreading.

Digitate, compound, the members arising from the same point.

Dioecious, of flowers when the staminate and pistillate ones are on separate plants.

Discoid, of composite flowers when ray-florets are absent.

Disk, a development of the receptacle around the base of the pistil; the central part of the head of a composite flower as distinct from the ray.

Dissected, cut up into many segments.

Dissepiment, a partition in an ovary or fruit.

Distinct, not united.

Divergent, spreading apart.

Divided, lobed to the base.

Dorsal, relating to the back of any organ.

Drupe, a fleshy fruit with the seed (or occasionally seeds) inclosed in a hard shell, as a plum or cherry.

E

Elliptical, oval.

Emarginate, slightly notched at the end.

Embryo, the undeveloped plant in the seed.

Endocarp, the innermost layer of the pericarp.

Endogenous, applied to stems whose wood does not grow in rings, but in scattered bundles.

Entire, without indentations of any kind.

Epicarp, the outermost layer of the pericarp.

Epigynous, growing on the ovary.

Epiphyte, an air-plant, whose roots do not reach the ground.

Equitant, applied to such leaves as those of Iris, which are folded lengthwise, each *astride* of the next one within.

Erect, upright.

Excurrent, applied to stems which can be readily traced through to the top, as in Pine.

Exogenous, applied to stems whose wood grows in layers or rings.

Exserted, thrust out beyond the line of the enveloping organ, as stamens out of a corolla.

Extrorse, facing outwards.

F

Fascicle, a close bundle.

Fertile, applied to flowers having pistils.

Fibrous, thread-like.

Filament, the stalk of the stamen.

Filiform, thread-shaped; long and slender.

Floccose, soft-woolly.

Foliaceous, leaf-like.

-foliate, relating to leaves.

-foliolate, relating to leaflets.

Follicle, a dehiscent fruit of one carpel, splitting down one side only.

Free, not growing fast to any other organ.

Friable, crumbling easily.

Frond, the leaf of a Fern, and some other cryptogams.

Fruit, the ripened ovary, along with any adherent part.

Fugacious, falling away very early.

Fusiform, spindle-shaped, thicker in the middle than at either end.

G

Galea, a helmet-shaped piece of a perianth, as the upper lip of some labiate corollas.

Gamopetalous, having the petals united together.

Gamophyllous, having the pieces of the perianth united.

Gibbous, bulging outwards on one side.

Glabrate, somewhat glabrous; becoming glabrous.

Glabrous, smooth.

Gland, a secreting structure of any kind. Any protuberance having the appearance of such an organ.

Glandular, bearing glands; or gland-like.

Glaucous, whitened with a bloom.

Globose, nearly spherical.

Glumaceous, having glumes; or glume-like.

Glume, a chaffy bract near a grass-flower.

Granular, composed of small, grain-like pieces.

Gymnospermous, having seeds which are not inclosed in an ovary.

Gynandrous, of stamens which are borne on the style.

H

Habit, the general aspect of a plant.

Halberd-shaped, with spreading lobes at the base.

Hastate, the same as halberd-shaped.

Head, a dense cluster of sessile flowers on a very short axis.

Heart-shaped, with an indentation at the base.

Herb, a plant with little or no wood in its stem.

Herbaceous, having the character of an herb; leaf-like in colour and texture.

Hilum, the scar showing where a seed was attached.

Hirsute, pubescent with coarse hairs.

Hispid, with stiff bristly hairs.

Hoary, grayish-white with fine close pubescence.

Hooded, shaped like a hood.

Hypogynous, inserted on the receptacle below the ovary, and free from the latter.

I

Imbricate, overlapping, as the bracts of the involucre in most composite flowers, having one piece wholly internal and one wholly external, as in the calyx and corolla of many flowers.

Immersed, wholly under water.

Incised, sharply cut.

Included, not at all protruded beyond the surrounding envelope.

Indefinite, many in number.

Indehiscent, not splitting open so as to discharge the contents.

Indigenous, native.

Inferior, lower; outer; of a calyx when below and free from the ovary; of an ovary when attached to the calyx.

Inflorescence, the arrangement of the flowers or flower-clusters.

Inserted, attached to.

Introrse, turned inwards.

Involucel, a secondary involucre.

Involucre, a circle of bracts below a flower or flower-cluster.

Involute, rolled inwards.

Irregular, with parts differing in size or shape.

K

Keel, a central longitudinal ridge; the two united petals in the front of a papilionaceous flower.

Kidney-shaped, of a leaf with large rounded lobes at the base.

L

Labiate, having two lips.

Laciniate, cut into narrow pointed lobes.

Lamella, a thin flat plate.

Lanceolate, rather narrow and tapering from the base to the point.

Lateral, referring to or borne on the side.

Lax, loose.

Leaflet, one of the pieces of the blade of a compound leaf.

Legume, a dehiscent fruit of a single carpel, which opens as a rule by two seams.

Lenticular, shaped like a double-convex lens.

Ligulate, strap-shaped.

Limb, the spreading part of a petal or corolla, etc.

Linear, long and narrow, with both ends alike or nearly so.

Lobe, any segment of an organ.

Lyrate, pinnatifid with the terminal lobe very large and rounded.

M

Marginal, along or near the edge.

Membranaceous, thin, somewhat transparent.

Mesocarp, the middle layer of the pericarp.

Midrib, the central or main vein of a leaf.

Monadelphous, with all the filaments grown together.

Monocotyledonous, having only one cotyledon in the embryo.

Monoecious, having staminate and pistillate flowers on different parts of the same plant.

Mucronate, tipped with a minute sharp point.

Multifid, cut into many segments.

N

Naked, without a covering, as of chaff, hairs, etc.

Nerve, a longitudinal line, as on the calyx of Catnip.

Node, the place on a stem from which a leaf springs.

Numerous, indefinite in number.

Nut, an indehiscent fruit of one cell and one seed with a hard shell, usually the product of a compound pistil.

Nutlet, a body like a little nut.

O

Obcompressed, compressed from above instead of from side to side.

Obcordate, indented at the apex.

Oblanceolate, narrow and tapering towards the base.

Oblique, unequal-sided, slanting.

Oblong, longer than broad, with rather straight sides, and both ends alike.

Obovate, egg-shaped, with the narrow end down.

Obsolete, not evident.

Obtuse, blunt.

Ocreate, having sheathing stipules.

Operculate, having a lid.

Orbicular, circular in outline.

Oval, somewhat longer than broad, with rounded sides and both ends alike.

Ovary, the lower part of the pistil containing the ovules.

Ovate, egg-shaped, the broad end down.

Ovoid, a solid with an ovate outline.

P

Palate, an upward projection of the lower lip of a personate corolla, closing the throat.

Palmate, with veins or leaflets radiating from the end of the petiole.

Panicle, an irregularly compound inflorescence of pedicelled flowers.

Panicked, resembling a panicle, borne in a panicle.

Papilionaceous, having a standard, wings, and keel, as in leguminous corollas.

Pappus, the tuft of hairs or bristles answering to the calyx-limb in many composite flowers.

Parasitic, living upon other living plants.

Parietal, borne on the walls.

Parted, very deeply cleft.

Pectinate, cut into comb-like, narrow teeth.

Pedate, palmately divided or parted, with the basal segments 2-cleft.

Pedicel, a secondary flower-stalk; the support of a single flower in a cluster.

Peduncle, the support of a solitary flower or of a flower-cluster.

Peltate, shield-shaped; attached by the lower surface.

Pendulous, hanging down.

Perennial, lasting year after year.

Perfect, having both stamens and pistil in the same flower.

Perfoliate, with the stem passing through the leaf.

Perianth, the floral envelopes.

Pericarp, the matured ovary.

Perigynous, inserted on the calyx.

Persistent, remaining after other parts have fallen away.

Personate, applied to a labiate corolla with the throat closed by an upward projection of the lower lip.

Petal, one of the pieces of a corolla.

Petaloid, petal-like.

Petiolate, having a petiole.

Petiole, a leaf-stalk.

Pilose, with soft hairs.

Pinna, a primary division of a frond.

Pinnate, with veins or leaflets proceeding from each side of a midrib.

Pinnatifid, deeply pinnately cleft.

Pinnule, a secondary division of a frond; one of the divisions of a pinna.

Pistil, the seed-bearing organ of the flower.

Pistillate, having pistil but no stamens.

Pitted, marked with small pits or depressions.

Placenta, the ridge or projection in the ovary to which the ovules are attached.

Plicate, or **Plaited**, folded into plaits.

Plumose, feathery; with fine hairs on each side of the axis.

Pod, any dry dehiscent fruit.

Pollen, the grains produced in the anther.

Pollinium (plural *pollinia*), a mass of pollen, as in Milkweeds and Orchids.

Polygamous, producing staminate, pistillate, and perfect flowers on one plant.

Polypetalous, having petals separate from each other.

Pome, an inferior syncarpous fruit consisting largely of the fleshy receptacle.

Posterior, the side next the axis.

Procumbent, lying flat on the ground.

Prostrate, the same as procumbent.

Puberulent, minutely pubescent.

Pubescent, covered with fine short hairs.

Punctate, showing transparent dots when held up to the light.

Pungent, acrid to the taste.

R

Raceme, a flower-cluster of lateral flowers, each on a pedicel of its own.

Racemose, in racemes; resembling a raceme.

Radiate, spreading from a centre; bearing ray-florets.

Radical, proceeding from the base of the stem.

Ray, the branch of an umbel: the circle of marginal florets in a composite flower, as distinguished from the disk.

Receptacle, the enlarged top of the flower-stalk.

Recurved, curved backwards.

Reflexed, bent backwards or downwards.

Regular, with all the parts of the organ of the same size and shape.

Reniform, kidney-shaped.

Repand, with a margin resembling that of an expanded umbrella.

Reticulate, in the form of a network.

Retuse, with a shallow notch at the end.

Revolute, rolled backwards from the margin.

Rhachis, the axis of a frond or compound leaf.

Rhizome, an underground stem; a rootstock.

Rib, one of the main veins of a leaf.

Ringent, wide open, gaping.

Rootstock, an underground stem.

Rotate, wheel-shaped.

Rudimentary, but slightly developed.

Rugose, wrinkled.

Runcinate, with teeth on the margin pointing backwards.

Runner, a thread-like, prostrate branch, proceeding from the base of a stem, and rooting at the extremity.

S

Saccate, sac-shaped.

Sagittate, arrow-shaped, the basal lobes directed downwards.

Salver-shaped, with a border spreading at right angles to the tube.

Samara, a winged fruit, as that of the Ash.

Scabrous, rough.

Scape, a peduncle, usually leafless, rising from the ground or near it.

Scarious, thin, dry, and membranaceous.

Scurf, small, rusty-looking scales on the epidermis.

Segment, one of the parts of a divided leaf.

Sepal, a division of a calyx.

Septum, a partition.

Serrate, toothed, the teeth directed towards the apex.

Serrulate, finely serrate.

Sessile, without a stalk.

Setaceous, bristle-like.

Setose, beset with bristles.

Sheath, a tubular envelope.

Sheathing, inclosing as with a sheath.

Shrub, a woody perennial smaller than a tree.

Silicle, a short and broad silique.

Silique, the peculiar pod of a cruciferous flower.

Simple, of one piece.

Sinuate, wavy.

Sinus, the indentation between two lobes.

Sorus, a cluster of sporangia.

Spadix, a spike on a fleshy axis.

Spathe, a bract, subtending or enveloping a spadix.

Spatulate, gradually narrowed downwards from a rounded apex.

Spicate, in the form of a spike.

Spike, a cluster of sessile flowers on a more or less elongated axis.

Spikelet, a small or secondary spike.

Spindle-shaped, larger in the middle than at either end.

Spine, a sharp woody outgrowth of the stem.

Spinose, spine-like, or beset with spines.

Sporangium, a spore-case.
Sporocarp, the fruit-case of certain Cryptogams.
Spur, a hollow projection.
Squarrose, having spreading tips.
Stamen, a pollen-bearing organ.
Staminodia, sterile stamens or bodies resembling stamens but without anthers.
Standard, the upper petal of a papilionaceous corolla.
Stellate, star-shaped.
Sterile, not productive, applied to pistils and stamens.
Stigma, the upper end of the pistil, adapted for the reception of pollen.
Stigmatic, stigma-like.
Stipe, the leaf-stalk of a Fern: the stalk supporting a pistil in certain flowers.
Stipular, relating to stipules.
Stolon, a branch which roots.
Stoloniferous, bearing stolons.
Striate, marked with fine longitudinal lines.
Strict, rigid and upright.
Strigose, beset with appressed, sharp, straight and stiff hairs.
Style, the narrow part of a pistil between the ovary and the stigma.
Stylododium, a disk-like expansion at the base of a style, as in umbelliferous plants.
Sub-, a prefix meaning "some-what".
Subulate, awl-shaped.
Succulent, juicy.
Suffrutescent, slightly shrubby.
Suffruticose, low and woody, diminiutively shrubby.
Sulcate, grooved or furrowed.

Superior (calyx), attached to the ovary.
Suture, a seam or line of dehiscence.
Symmetrical, with sets having the same number of parts each, or a multiple of that number.

T

Tail, any slender prolongation.
Terete, cylindrical.
Terminal, at the apex.
Ternate, in threes.
Tetradynamous, with four long stamens and two short ones.
Throat, the entrance to the tube of a calyx or corolla.
Thyrse, a compact panicle.
Thyrseoid, like a thyrse.
Tomentose, woolly.
Transverse, across.
Tri-, a prefix meaning three or thrice.
Trifoliolate, having three leaflets.
Truncate, as if cut off square across the end.
Tuber, the enlarged end of an underground stem.
Tubercle, a small, tuber-like body.
Tuberous, having the appearance of a tuber.
Tunicated, having coats one within another.
Twining, winding spirally about a support.

U

Umbel, a flower-cluster with radiating pedicels.
Umbellate, umbel-like.
Umbellet, a secondary umbel.
Undulate, wavy.

Urceolate, urn-shaped.

Utricle, an indehiscent, 1-seeded fruit, with a thin loose pericarp.

V

Valvate, with the edges meeting but not overlapping.

Valve, one of the pieces into which a capsule splits.

Veins, the threads of fibro-vascular tissue running through the substance of a leaf.

Ventral, belonging to the inferior or lower side.

Vernation, the folding of a leaf in the bud.

Versatile, attached by the centre (of the anther) and so able to swing about.

Vertical, upright.

Verticillate, whorled.

Villous, with long soft hairs.

Viscid, sticky.

W

Whorl, a circle of leaves around a stem.

Wing, a thin expansion bordering any organ; one of the side petals of a papilionaceous corolla.

Woolly, with long matted hairs.

ABBREVIATIONS OF PROPER NAMES

A. Br.....	Braun, Alexander.	Hausk.....	Hausknecht, Carl.
Adans.....	Adanson, Michel.	Haw.....	Haworth, Adrian H.
A. DC.	De Candolle, Alphonse.	HBK.....	Humboldt, F. Alex von,
Ait.....	Aiton, William.		Bonpland, A., and
Anders.....	Anderson, Nils Johan.		Kunth, C. S.
Andrz.....	Andrzejowski, Anton L.	Hitchc.....	Hitchcock, Albert Spear.
Aschers.....	Ascherson, Paul.	Hook.....	Hooker, William Jackson.
		Hornem.....	Hornemann, Jens Wilken.
		Huds.....	Hudson, William.
B. and H....	Bentham, George, and		
	Hooker, Joseph Dalton.	Jacq.....	Jacquin, Nicolaus Joseph.
Beauv.....	Beauvois, A.M.F.J.P.		
Bernh.....	Bernhardi, Johann Jacob.	Ktze.....	Kuntze, Otto.
Bieb.....	Bieberstein, F. A. Marschall		
	von.	L.....	Linnaeus, C.
Bigel.....	Bigelow, Jacob.	Lag.....	Lagasca, Mariano.
Boiss.....	Boissier, Edmond.	Lam.....	Lamarck, Jean Baptiste.
Borkh.....	Borkhausen, M. B.	L'Her.....	L'Heritier de Brutelle, C. L.
BSP.....	Britton, N. L., Sterns,	Lehm.....	Lehmann, J. G.
	E. E., Pöggendorf, J. F.	Lindl.....	Lindley, John.
		Lodd.....	Loddiges, Conrad.
Cav.....	Cavanilles, Antonio José.	Loud.....	Loudon, John Claudius.
Cyrill.....	Cirillo, Domenico.		
		MacM.....	MacMillan, Conway.
DC.....	De Candolle, Augustin P.	Marsh.....	Marshall, Humphrey.
Desf.....	Desfontaines, René L.	Maxim.....	Maximowicz, Carl Johann.
Desr.....	Desrousseau.	Medic.....	Medicus, Fried. Casimir.
Doug.....	Douglas, David.	Mey. G.F.W.	Meyer, G. F. Wilhelm.
Dumont.....	Du Mont de Courset,	Michx.....	Michaux, André.
	G.L.M.	Mill.....	Miller, Philip.
		Moq.....	Moquin-Tandon, Alfred.
Ehrh.....	Ehrhart, Friedrich.	Muench.....	Muenchhausen, Otto
Ell.....	Elliott, Stephen.		Freiherr von.
Engelm.....	Engelmann, George.	Muhl.....	Muhlenberg, G. H. E.
		Murr.....	Murray, Johann Andreas.
Foug.....	Fougeroux, Auguste D.		
Froel.....	Froelich, Joseph A.	Neck.....	Necker, Noel Joseph de.
		Nutt.....	Nuttall, Thomas.
Gaertn.....	Gaertner, Joseph.	Pall.....	Pallas, Peter Simon.
Gmel.....	Gmelin, Samuel Gottlieb.	Pers.....	Persoon, Christian Hend-
Grev.....	Greville, Robert Kaye.		rick.
Griseb.....	Grisebach, Heinrich.	Poir.....	Poiret, Jean L. Marie.
Guss.....	Gussoni, Giovanni.		

Raf.	Rafinesque—Schmaltz,
C. S.	T. and G. . . .	Torrey, John and Gray,
Asa.
R. Br.	Brown, Robert.	Torr.	Torrey, John.
Reichenb. . . .	Reichenbach, H. G. L.	Trel.	Trelease, William.
Roem.	Roemer, M. J.	Turcz.	Turczaninow, Nicolaus.
Rottb.	Rottboell, Christen Fries.		
Rydb.	Rydberg, Per Axel.	Vent.	Ventenat, Étienne Pierre.
Salisb.	Salisbury, Richard Anthony.	Wallr.	Wallroth, K. F. W.
Sarg.	Sargent, Charles Sprague.	Walp.	Walpers, W. G.
Scop.	Scopoli, Johann A.	Walt.	Walter, Thomas.
Sibth.	Sibthorp, John.	Wang.	Wangenheim, F. A. J. von.
Sm.	Smith, James Edward.	Wats.	Watson, Sereno.
Spreng.	Sprengel, Kurt.	Willd.	Willdenow, Carl Ludwig.
Steud.	Steudel, Ernst Gottlieb.		

INDEX

- Abies, 38.
- Abieteeae, 37.
- Abutilon, 169.
- Acalypha, 160.
- ACANTHACEAE, 232.
- Acanthus Family, 232.
- Acer, 164.
- ACERACEAE, 164.
- Acerates, 205.
- Achillea, 271.
- Acnida, 93.
- Acorus, 48.
- Actaea, 109.
- Actinomeris, 269.
- Adam-and-Eve, 67.
- Adder's Mouth, 66.
- Adder's Tongue, 31.
- Adder's Tongue Family, 30.
- Adiantum, 23.
- Adlumia, 114.
- Aethusa, 186.
- Agastache, 218.
- Agoseris, 274.
- Agrimonia, 132.
- Agrimony, 132.
- Agrostemma, 97.
- AIZOACEAE, 94.
- Alchemilla, 133.
- Alder, 77.
- Aletris, 58.
- Alfalfa, 145.
- Alisma, 44.
- ALISMACEAE, 44.
- Alkanet, 211.
- Allium, 59.
- Alnus, 77.
- Alum-root, 128.
- Alyssum, 122.
- Amaranth, 93.
- Amaranth Family, 93.
- Amaranthus, 93.
- AMARANTHACEAE, 93.
- AMARYLLIDACEAE, 60.
- Amaryllis Family, 60.
- Ambrosia, 251.
- Amelanchier, 141.
- American Brooklime, 225.
- American Columbo, 201.
- American Cowslip, 198.
- American Laurel, 194.
- American White Hellebore, 57.
- Amorpha, 149.
- Amphicarpa, 151.
- ANACARDIACEAE, 162.
- Anagallis, 199.
- Anaphalis, 254.
- Androsace, 198.
- Andromeda, 194.
- Anemone, 104.
- Anemonella, 106.
- Angelica, 186.
- ANGIOSPERMS, 39.
- ANONACEAE, 110.
- Antennaria, 254.
- Anthemis, 267.
- Apios, 151.
- Aplectrum, 67.
- Aplopappus, 257.

- APOCYNACEAE, 202.
 Apocynum, 203.
 Apple, 140.
 Apple of Peru, 222.
 AQUIFOLIACEAE, 162.
 Aquilegia, 108.
 Arabis, 118.
 ARACEAE, 46.
 Aralia, 182.
 ARALIACEAE, 182.
 Arbor Vitae, 39.
 Arceuthobium, 84.
 Arctium, 250.
 Arctostaphylos, 193.
 Arenaria, 97.
 Arethusa, 66.
 Arisaema, 47.
 Aristolochia, 84.
 ARISTOLOCHIACEAE, 84.
 Arnica, 260.
 Aromatic Wintergreen, 194.
 Arrow Arum, 47.
 Arrow Grass, 44.
 Arrow-head, 45.
 Arrow-wood, 238.
 Artemisia, 252.
 Arum Family, 46.
 Asarum, 84.
 ASCLEPIADACEAE, 203.
 Asclepias, 204.
 Ash, 199.
 Asimina, 110.
 Asparagus, 58.
 Aspen, 73.
 Aspidium, 26.
 Asplenium, 24.
 Aster, 261.
 Astragalus, 146.
 Atriplex, 92.
 Avens, 133.
 Azalea, 195.
 Azolla, 32.
 Baked-apple Berry, 137.
 Ball Mustard, 123.
 Ballota, 220.
 Balm, 219.
 Balm of Gilead, 73.
 Balsam Family, 165.
 BALSAMINACEAE, 165.
 Baneberry, 109.
 Baptisia, 153.
 Barbarea, 117.
 Barberry, 111.
 Barberry Family, 111.
 Barren Strawberry, 134.
 Basil, 217.
 Basswood, 167.
 Bastard Toad-flax, 83.
 Bayberry, 74.
 Beach Pea, 150.
 Bearberry, 193.
 Beard-tongue, 227.
 Beaver-poison, 187.
 Beech, 79.
 Beech Family, 78.
 Beech-drops, 231.
 Beech Fern, 25.
 Bedstraw, 234.
 Beggar's Lice, 210.
 Beggar-ticks, 270.
 Bell-flower, 241.
 Bellis, 267.
 Bellwort, 57.
 Benzoin, 112.
 BERBERIDACEAE, 111.
 Berberis, 111.
 Bergamot, 218.
 Betula, 76.
 BETULACEAE, 75.

Bidens, 270.
Bindweed, 205.
Birch, 76.
Birch Family, 75.
Birthwort Family, 84.
Bishop's Cap, 128.
Bitter Cress, 118.
Bitter-sweet, 221.
Black Alder, 163.
Blackberry, 138.
Black Bindweed, 87.
Black Cap, 138.
Black Grass, 52.
Black Horehound, 220.
Black-eyed Susan, 268.
Black Mustard, 121.
Black Snakeroot, 109, 185.
Bladder Campion, 96.
Bladder Fern, 28.
Bladder Ketmia, 169.
Bladder Nut, 164.
Bladder-pod, 121.
Bladderwort, 230.
Bladderwort Family, 230.
Blazing Star, 254.
Blephilia, 218.
Blessed Thistle, 251.
Blite, 91.
Blood-root, 113.
Blue Ash, 200.
Blue Beech, 77.
Blueberry, 192.
Bluebottle, 251.
Blue Cohosh, 111.
Blue Flag, 61.
Bluets, 236.
Blue-eyed Grass, 61.
Blue Lettuce, 274.
Blue-weed, 209.
Boehmeria, 83.

Boneset, 255.
Borage Family, 208.
BORAGINACEAE, 208.
Botrychium, 30.
Bouncing Bet, 95.
Bowman's Root, 132.
Box Elder, 165.
Bracken, 23.
Brake, 23.
Bramble, 137.
Brasenia, 101.
Brassica, 121.
Bread-fruit and Fig Family, 80.
Bristly Sarsaparilla, 182.
Brooklime, 225.
Brook-weed, 199.
Broom Crowberry, 161.
Broom-rape Family, 231.
Buckbean, 202.
Buckthorn, 166.
Buckthorn Family, 166.
Buckwheat, 89.
Buckwheat Family, 84.
Buffalo Berry, 177.
Bugbane, 109.
Bug-seed, 92.
Bugle-weed, 216.
Bugloss, 210.
Bunch-berry, 188.
Bupleurum, 187.
Burdock, 250.
Bur-marigold, 270.
Burnet, 133.
Burning Bush, 163.
Bur-reed, 40.
Bush-clover, 153.
Bush Honeysuckle, 238.
Butter and Eggs, 226.
Buttercup, 106.
Butterfly-weed, 204.

- Butternut, 74.
 Butter-weed, 265.
 Butterwort, 231.
 Button Bush, 235.
 Buttonwood, 129.

 Cabbage, 121.
 Cacalia, 255.
 CACTACEAE, 175.
 Cactus Family, 175.
 Cakile, 123.
 Calamus, 48.
 Calla, 47.
 Callitriche, 160.
 CALLITRICHACEAE, 160.
 Calluna, 194.
 Calopogon, 66.
 Caltha, 108.
 Calypso, 66.
 Camelina, 122.
 Campanula, 241.
 CAMPANULACEAE, 241.
 Campanula Family, 241.
 Campion, 96.
 Camptosorus, 25.
 Cancer-root, 232.
 Cannabineae, 81.
 Cannabis, 83.
 Caper Family, 123.
 CAPPARIDACEAE, 123.
 CAPRIFOLIACEAE, 236.
 Capsella, 122.
 Caraway, 187.
 Carduus, 250.
 Cardamine, 118.
 Cardinal Flower, 242.
 Carpet-weed, 94.
 Carpinus, 77.
 Carrion Flower, 56.
 Carrot, 185.

 Carum, 187.
 Carya, 75.
 CARYOPHYLLACEAE, 95.
 Cashew Family, 162.
 Castalia, 101.
 Castanea, 79.
 Castilleja, 229.
 Cat Brier, 56.
 Catchfly, 96.
 Catgut, 149.
 Cat Mint, 218.
 Catnip, 218.
 Cat-tail Family, 40.
 Cat-tail Flag, 40.
 Caulophyllum, 111.
 Ceanothus, 166.
 Cedar, 39.
 Celandine, 113.
 CELASTRACEAE, 163.
 Celastrus, 164.
 Celtideae, 80.
 Celtis, 82.
 Centaurea, 251.
 Cephalanthus, 235.
 Cerastium, 98.
 CERATOPHYLLACEAE, 101.
 Ceratophyllum, 101.
 Cercis, 154.
 Chain Fern, 25.
 Chamaedaphne, 194.
 Chamaerhodos, 137.
 Chamomile, 267.
 Charlock, 121.
 Chelidonium, 113.
 Chelone, 227.
 CHENOPODIACEAE, 89.
 Chenopodium, 90.
 Cherry, 131.
 Chestnut, 79.
 Chickweed, 97.

- Chickweed Wintergreen, 198.
Chimaphila, 196.
Chiogenes, 193.
Choke-berry, 141.
Choke-cherry, 132.
Chrysanthemum, 266.
Christmas Fern, 25.
Chrysopsis, 257.
Chrysosplenium, 128.
Cichorium, 272.
Cichory, 272.
Cicuta, 187.
Cimicifuga, 109.
Cinnamon Fern, 29.
Cinque-foil, 134.
Circaea, 178.
Cirsium, 249.
CISTACEAE, 170.
Claytonia, 100.
Clearweed, 82.
Cleavers, 234.
Clematis, 103.
Cleome, 124.
Cliff Brake, 23.
Climbing Bitter-sweet, 164.
Clintonia, 57.
Clotbur, 251.
Cloud-berry, 137.
Clover, 144.
Club-moss, 34.
Club-moss Family, 33.
Cnicus, 251.
Coast Blite, 91.
Cockle, 97.
Cocklebur, 251.
Cohosh, 111.
Colic-root, 58.
Collinsia, 227.
Collinsonia, 217.
Colt's Foot, 261.
Columbine, 108.
Comandra, 83.
Common Horsetail, 32.
Comfrey, 209.
COMPOSITAE, 243.
Composite Family, 243.
Cone-flower, 267.
Conioselinum, 186.
Conium, 187.
Conopholis, 231.
CONVOLVULACEAE, 205.
Convolvulus, 205.
Convolvulus Family, 205.
Coptis, 108.
Corallorrhiza, 68.
Coral-root, 68.
Corema, 161.
Coreopsis, 269.
Corispermum, 92.
CORNACEAE, 188.
Corn Cockle, 97.
Cornel, 188.
Corn Spurrey, 100.
Cornus, 188.
Coronopus, 123.
Corpse Plant, 196.
Corydalis, 115.
Corylus, 77.
Costmary, 267.
Cottonwood, 73.
Cowbane, 185.
Cow-herb, 96.
Cow Parsnip, 185.
Cowslip, 197.
Cow Wheat, 230.
Crab-apple, 140.
Cranberry, 192.
Cranberry Tree, 239.
Cranesbill, 155.
CRASSULACEAE, 125.

- Crataegus, 140.
Creeping Snowberry, 193.
Crepis, 273.
Cress Family, 115.
Crowberry, 161.
Crowberry Family, 161.
Crowfoot, 106.
Crowfoot Family, 102.
CRUCIFERAE, 115.
Cryptogramma, 23.
Cryptotaenia, 187.
Cuckoo-flower, 118.
Cucumber Tree, 110.
CUCURBITACEAE, 240.
Cudweed, 253.
Cup-plant, 271.
Cupresseae, 37.
Currant, 126.
Cuscuta, 206.
Custard Apple Family, 110.
Cycloloma, 92.
Cynoglossum, 210.
Cynthia, 271.
CYPERACEAE, 46.
Cypress Family, 37.
Cypripedium, 69.
Cystopteris, 28.

Daisy, 267.
Daisy Fleabane, 266.
Dalibarda, 137.
Dandelion, 274.
Daphne, 176.
Datura, 222.
Daucus, 185.
Day-lily, 60.
Dead Nettle, 220.
Decodon, 177.
Deer-berry, 193.
Deer-grass, 178.

Delphinium, 109.
Dentaria, 118.
Desmodium, 151.
Devil's Paint-brush, 272.
Dianthera, 232.
Dicentra, 114.
Dicksonia, 29.
DICOTYLEDONS, 69.
Diervilla, 238.
Dioscorea, 60.
DIOSCOREACEAE, 60.
Diploaxis, 121.
DIPSACACEAE, 240.
Dipsacus, 240.
Dirca, 176.
Disporum, 57.
Ditch Stone-crop, 125.
Dock, 88.
Dockmackie, 239.
Dodder, 206.
Dodecatheon, 198.
Dogbane, 203.
Dogbane Family, 202.
Dog's-tooth Violet, 59.
Dogwood, 188.
Dogwood Family, 188.
Downy Arrow-wood, 239.
Draba, 121.
Dracocephalum, 218.
Dragon-head, 218.
Drosera, 124.
DROSERACEAE, 124.
Duckweed, 48.
Duckweed Family, 48.
Dutchman's Breeches, 114.
Dutchman's Pipe, 84.
Dwarf Dandelion, 272.

Echinocystis, 241.
Echium, 209.

- Eel-grass, 44, 46.
 Eglantine, 139.
 ELAEAGNACEAE, 176.
 Elaeagnus, 176.
 Elder, 238.
 Elecampane, 257.
 Ellisia, 208.
 Elm, 81.
 Elm Family, 80.
 Elodea, 46.
 EMPETRACEAE, 161.
 Empetrum, 161.
 Enchanter's Nightshade, 178.
 Epigaea, 194.
 Epilobium, 178.
 Epipactis, 67.
 Epifagus, 231.
 EQUISETACEAE, 32.
 Equisetum, 32.
 Erechites, 253.
 ERICACEAE, 190.
 Ericoideae, 190.
 Erigenia, 188.
 Erigeron, 265.
 ERIOCAULACEAE, 49.
 Eriocaulon, 49.
 Eriogonum, 85.
 Erodium, 156.
 Erythronium, 59.
 Erysimum, 120.
 Eupatorium, 255.
 Euphorbia, 158.
 EUPHORBIACEAE, 158.
 Euphrasia, 229.
 Evening Primrose, 179.
 Evening Primrose Family, 178.
 Everlasting, 253, 254.
 Everlasting Pea, 150.
 Evonymus, 163.
 Eyebright, 229.
 FAGACEAE, 78.
 Fagopyrum, 89.
 Fagus, 79.
 Fall Dandelion, 272.
 False Asphodel, 57.
 False Dragon-head, 219.
 False Flax, 122.
 False Gromwell, 211.
 False Hellebore, 57.
 False Indigo, 149, 153.
 False Lettuce, 274.
 False Loosestrife, 180.
 False Mallow, 168.
 False Mermaid Family, 161.
 False Mitrewort, 128.
 False Nettle, 83.
 False Pennyroyal, 216.
 False Pimpernel, 228.
 False Solomon's Seal, 57.
 False Spikenard, 57.
 Fetid Horehound, 220.
 Fever-bush, 112.
 Feverfew, 267.
 Feverwort, 238.
 Figwort, 226.
 Figwort Family, 223.
 Filbert, 77.
 Fir, 38.
 Fire Pink, 96.
 Fireweed, 253.
 Five-finger (Cinque-foil), 134.
 Flax, 154.
 Flax Dodder, 206.
 Flax Family, 154.
 Fleabane, 265.
 Fleur-de-lis, 61.
 Floating Heart, 202.
 Floerkea, 161.
 Flowering Fern, 29.
 Flowering Fern Family, 29.

- Fool's Parsley, 186.
Forget-me-not, 212.
Fragaria, 137.
Franseria, 252.
Fraseria, 201.
Fraxinus, 199.
French Weed, 122.
Frog's-bit Family, 45.
Frost-weed, 171.
Fumaria, 115.
FUMARIACEAE, 114.
Fumitory, 115.
Fumitory Family, 114.

Gaillardia, 269.
Galeopsis, 219.
Galinsoga, 271.
Galum, 234.
Garden Cress, 123.
Garlic, 59.
Gaultheria, 194.
Gaura, 180.
Gaylussacia, 192.
Gentian, 201.
Gentiana, 201.
GENTIANACEAE, 200.
Gentian Family, 200.
GERANIACEAE, 155.
Geranium, 155.
Geranium Family, 155.
Gerardia, 228.
Germander, 215.
Geum, 133.
Giant Hyssop, 218.
Gillenia, 132.
Gilia, 207.
Ginseng, 182.
Ginseng Family, 182.
Glasswort, 92.
Glaux, 199.

Gleditsia, 153.
Glycyrrhiza, 149.
Gnaphalium, 253.
Goat's Beard, 275.
Goat's Rue, 149.
Golden Aster, 257.
Golden Ragwort, 256.
Golden-rod, 257.
Golden Saxifrage, 128.
Golden Seal, 109.
Goldthread, 108.
Good-King-Henry, 91.
Gooseberry, 126.
Goosefoot, 90.
Goosefoot Family, 89.
Goosegrass, 234.
Gourd Family, 240.
GRAMINEAE, 46.
Grape, 167.
Grape Fern, 30.
Grass of Parnassus, 127.
Grass-wrack, 44.
Gratiola, 228.
Great Angelica, 186.
Green Ash, 200.
Green Brier, 56.
Green Dragon, 47.
Green Milkweed, 205.
Green Violet, 175.
Grindelia, 266.
Gromwell, 211.
Ground Cherry, 222.
Ground Hemlock, 36.
Ground Ivy, 218.
Ground Laurel, 194.
Ground-nut, 151.
Ground Pine, 34.
Groundsel, 256.
Gutierrezia, 266.

Gymnocladus, 154.
GYMNOSPERMS, 36.

Habenaria, 64.
Hackberry, 82.
Halenia, 201.
HALORAGIDACEAE, 181.
HAMAMELIDACEAE, 129.
Hamamelis, 129.
Harbinger-of-spring, 188.
Harebell, 241.
Hart's Tongue, 25.
Hawkweed, 272.
Hawthorn, 140.
Hazel-nut, 77.
Heal-all, 219.
Heather, 194.
Heath Family, 190.
Hedeoma, 217.
Hedge Bindweed, 205.
Hedge Hyssop, 228.
Hedge Mustard, 120.
Hedge Nettle, 220.
Hedysarum, 152.
Helenium, 266.
Helianthemum, 171.
Helianthus, 268.
Heliopsis, 271.
Hemerocallis, 60.
Hemlock, 38.
Hemlock Parsley, 186.
Hemlock Spruce, 38.
Hemp, 83.
Hemp Family, 81.
Hemp Nettle, 219.
Henbane, 222.
Hepatica, 105.
Heracleum, 185.
Herb Robert, 156.
Hesperis, 121.

Heteranthera, 50.
Heuchera, 128.
Hibiscus, 169.
Hickory, 75.
Hieracium, 272.
High-bush Cranberry, 239.
Hippurus, 181.
Hoary Pea, 149.
Hoary Puccoon, 211.
Hobble-bush, 239.
Hog Pea-nut, 151.
Hogweed, 251.
Holly, 163.
Holly Family, 162.
Honey-locust, 153.
Honeysuckle, 237.
Honeysuckle Family, 236.
Honewort, 187.
Hop, 83.
Hop Hornbeam, 77.
Hop-tree, 157.
Horehound, 219.
Hornbeam, 77.
Horned Pondweed, 43.
Hornwort, 101.
Hornwort Family, 101.
Horse Balm, 217.
Horse Mint, 218.
Horseradish, 117.
Horsetail, 32.
Horsetail Family, 32.
Horse-weed, 265.
Hosackia, 146.
Hound's Tongue, 210.
Houstonia, 235.
Huckleberry, 192.
Hudsonia, 171.
Humulus, 83.
Huntsman's Cup, 124.
Hybanthus, 175.

- Hydrastis, 109.
 HYDROCHARITACEAE, 45.
 Hydrocotyle, 185.
 HYDROPHYLLACEAE, 207.
 Hydrophyllum, 207.
 Hyoscyamus, 222.
 HYPERICACEAE, 169.
 Hypericum, 169.
 Hypoxis, 60.
 Hyssop, 217.
 Hyssopus, 217.

 Ice-plant Family, 94.
 Ilex, 163.
 ILLECEBRACEAE, 94.
 Ilysanthes, 228.
 Impatiens, 165.
 Indian Cucumber-root, 56.
 Indian Hemp, 203.
 Indian Mallow, 169.
 Indian Physic, 132.
 Indian Pipe, 196.
 Indian Plantain, 255.
 Indian Poke, 57.
 Indian Tobacco, 242.
 Indian Turnip, 47.
 Innocence, 236.
 Interrupted Fern, 29.
 Inula, 257.
 Ipomoea, 206.
 IRIDACEAE, 60.
 Iris, 61.
 Iris Family, 60.
 Iron-weed, 255.
 Iron-wood, 77.
 Isanthus, 216.
 ISOËTACEAE, 36.
 Isoëtes, 36.
 Iva, 255.

 Jack-in-the-pulpit, 47.
 Jack Pine, 38.
 Jeffersonia, 112.
 Jerusalem Artichoke, 269.
 Jerusalem Oak, 91.
 Jewel-weed, 165.
 Joe Pye Weed, 255.
 Judas Tree, 154.
 Juglans, 74.
 JUGLANDACEAE, 74.
 JUNCACEAE, 50.
 JUNCAGINACEAE, 44.
 Juncus, 51.
 June-berry, 141.
 Juniper, 39.
 Juniperus, 39.

 Kalmia, 194.
 Kentucky Coffee-tree, 154.
 Knapweed, 251.
 Knotgrass, 85.
 Knotweed, 85.
 Knotwort Family, 94.
 Krigia, 272.

 LABIATAE, 213.
 Labrador Tea, 195.
 Lactuca, 274.
 Ladies' Tresses, 68.
 Lady Fern, 24.
 Lady's Mantle, 133.
 Lady's Slipper, 69.
 Lady's Thumb, 86.
 Lady's Smock, 118.
 Lake Cress, 117.
 Lambkill, 194.
 Lamb's Quarters, 90.
 Lamium, 220.
 Laportea, 82.
 Lappa, 250.

- Lappula, 209.
Lapsana, 272.
Larch, 39.
Larix, 39.
Larkspur, 109.
Lathyrus, 150.
LAURACEAE, 112.
Laurel Family, 112.
Laurestinus, 238.
Lead-plant, 149.
Leadwort Family, 196.
Leaf-cup, 271.
Leather-leaf, 194.
Leatherwood, 176.
Lechea, 171.
Ledum, 195.
Leek, 59.
LEGUMINOSAE, 141.
Lemna, 48.
LEMNACEAE, 48.
LENTIBULARIACEAE, 230.
Leontodon, 272.
Leonurus, 220.
Lepachys, 268.
Lepidium, 122.
Lespedeza, 153.
Lesquerella, 121.
Lettuce, 274.
Liatris, 254.
Liguliflorae, 248.
Ligusticum, 186.
LILIACEAE, 53.
Lilium, 58.
Lily, 58.
Lily Family, 53.
LIMNANTHACEAE, 161.
Limonium, 196.
LINACEAE, 154.
Linaria, 226.
Linden Family, 167.
Linnaea, 237.
Linum, 154.
Liparis, 67.
Liquorice, 149.
Liriodendron, 110.
Listera, 68.
Lithospermum, 211.
Live-for-ever, 125.
Liver-leaf, 105.
Lizard's Tail, 70.
Lobelia, 242.
LOBELIACEAE, 242.
Lobelia Family, 242.
Locust Tree, 146.
Lomatium, 185.
Loniceria, 237.
Loosestrife, 177, 198.
Loosestrife Family, 177.
Lopseed, 232.
LORANTHACEAE, 84.
Lousewort, 230.
Lovage, 186.
Lucerne, 145.
Ludvigia, 180.
Lungwort, 210.
Lupine, 144.
Lupinus, 144.
Luzula, 51.
Lychnis, 96.
Lycium, 222.
LYCOPODIACEAE, 33.
Lycopodium, 34.
Lycopsis, 210.
Lycopus, 216.
Lygodesmia, 273.
Lysimachia, 198.
LYTHRACEAE, 177.
Lythrum, 177.
Madder Family, 233.

- Magnolia, 110.
MAGNOLIACEAE, 109.
Magnolia Family, 109.
Maianthemum, 58.
Maidenhair, 23.
Maidenhair Spleenwort, 24.
Male Fern, 27.
Mallow, 168.
Mallow Family, 168.
Malva, 168.
MALVACEAE, 168.
Malvastrum, 168.
Mamillaria, 175.
Mandrake, 112.
Maple, 164.
Mare's Tail, 181.
Marrubium, 219.
Marsh Calla, 47.
Marsh Cress, 117.
Marsh Elder, 255.
Marsh Marigold, 108.
Marsh Rosemary, 196.
Marsh St. John's-wort, 169.
MARSILEACEAE, 31.
Marsilea, 31.
Matricaria, 267.
Matrimony Vine, 222.
May Apple, 112.
Mayflower, 194.
May-weed, 267.
Meadow-beauty, 178.
Meadow-parsnip, 186.
Meadow-rue, 105.
Meadow-sweet, 132.
Medeola, 56.
Medicago, 145.
Medick, 145.
Melampyrum, 230.
Melastoma Family, 177.
MELASTOMACEAE, 177.
Melilot, 145.
Melilotus, 145.
Melissa, 219.
MENISPERMACEAE, 110.
Menispermum, 110.
Mentha, 216.
Menyanthes, 202.
Mermaid-weed, 182.
Mertensia, 210.
Mexican Tea, 91.
Mezereum Family, 175.
Microstylis, 66.
Milfoil, 271.
Milk-vetch, 146.
Milkweed, 204.
Milkweed Family, 203.
Milkwort, 157.
Milkwort Family, 157.
Mimulus, 227.
Mint, 216.
Mint Family, 213.
Mistletoe Family, 84.
Mitchella, 235.
Mitella, 128.
Mitrewort, 128.
Moccasin-flower, 69.
Mock Pennyroyal, 217.
Mocker Nut, 75.
Mollugo, 94.
Monarda, 218.
Moneses, 196.
Moneywort, 198.
Monkey Flower, 227.
MONOCOTYLEDONS, 39.
Monolepis, 91.
Monotropa, 196.
Monotropoideae, 192.
Montia, 100.
Moonseed, 110.
Moonseed Family, 110.

Moonwort, 30.
 Moosewood, 176.
 Moreae, 80.
 Morning Glory, 206.
 Morus, 82.
 Moss Campion, 96.
 Moss Pink, 207.
 Mossy Stone-crop, 125.
 Motherwort, 220.
 Mountain Ash, 141.
 Mountain Holly, 163.
 Mountain Maple, 164.
 Mountain Mint, 217.
 Mouse-ear Chickweed, 98.
 Mousetail, 106.
 Mugwort, 252.
 Mulberry, 82.
 Mullein, 225.
 Musenium, 187.
 Musk Mallow, 168.
 Musk Thistle, 250.
 Mustard, 121.
 Myosotis, 212.
 Myosurus, 106.
 Myrica, 74.
 MYRICACEAE, 73.
 Myriophyllum, 181.
 Naiad, 43.
 NAJADACEAE, 41.
 Najas, 43.
 Naked Broom-rape, 232.
 Nanny-berry, 238.
 Neckweed, 226.
 Nemopanthus, 163.
 Nepeta, 218.
 Neslia, 123.
 Nettle, 82.
 Nettle Family, 80.
 Nettle-tree, 82.

New Jersey Tea, 166.
 Nicandra, 222.
 Nicotiana, 222.
 Nightshade, 221.
 Nightshade Family, 220.
 Nine-bark, 132.
 Nipplewort, 272.
 Nymphaea, 102.
 NYMPHAEACEAE, 101.
 Nymphoides, 202.
 Nyssa, 189.
 Oak, 78.
 Oakesia, 57.
 Oak Family, 78.
 Oak Fern, 26.
 Odontites, 229.
 Oenothera, 179.
 OLEACEAE, 199.
 Oleaster Family, 176.
 Olive Family, 199.
 ONAGRACEAE, 178.
 Onion, 59.
 Onoclea, 28.
 Onopordum, 250.
 Onosmodium, 211.
 OPHIOGLOSSACEAE, 30.
 Ophioglossum, 31.
 Opuntia, 175.
 Orache, 92.
 Orange-root, 109.
 ORCHIDACEAE, 61.
 Orchis, 64.
 Orchis Family, 61.
 OROBANCHACEAE, 231.
 Orobanche, 232.
 Orpine, 125.
 Orpine Family, 125.
 Orthocarpus, 229.
 Osmorhiza, 187.

- Osmunda, 29.
 OSMUNDACEAE, 29.
 Ostrich Fern, 28.
 Ostrya, 77.
 Oswego Tea, 218.
 OXALIDACEAE, 155.
 Oxalis, 155.
 Ox-eye, 271.
 Ox-eye Daisy, 266.
 Oxypolis, 185.
 Oxytropis, 148.

 Painted Cup, 229.
 Panax, 182.
 Papaver, 113.
 PAPAVERACEAE, 113.
 Papaw, 110.
 Parietaria, 83.
 Parnassia, 127.
 Paronychia, 94.
 Parsley Family, 183.
 Parsnip, 185.
 Partridge Berry, 235.
 Pasque Flower, 104.
 Pastinaca, 185.
 Pasture Sage-brush, 253.
 Pear, 140.
 Pearlwort, 99.
 Pedicularis, 230.
 Pellaea, 23.
 Pellitory, 83.
 Peltandra, 47.
 Penny Cress, 122.
 Pennyroyal, 216, 217.
 Penthorum, 125.
 Pentstemon, 227.
 Pepper Family, 70.
 Pepperidge, 189.
 Pepper Grass, 122.
 Peppermint, 216.
 Pepper-root, 118.
 Petalostemum, 149.
 Petasites, 261.
 Phegopteris, 25.
 Phlox, 206.
 Phryma, 232.
 PHRYMACEAE, 232.
 Physalis, 222.
 Physocarpus, 132.
 Physostegia, 219.
 Phytolacca, 94.
 PHYTOLACCACEAE, 94.
 Picea, 38.
 Pickerel-weed, 50.
 Pickerel-weed Family, 50.
 Pig-nut, 75.
 Pigweed, 90.
 Pilea, 82.
 Pimpernel, 199.
 PINACEAE, 37.
 Pine, 38.
 Pineapple-weed, 267.
 Pine-drops, 196.
 Pine Family, 37.
 Pine-sap, 196.
 Pinguicula, 231.
 Pink Family, 95.
 Pinus, 38.
 Pinweed, 171.
 PIPERACEAE, 70.
 Pipes, 33.
 Pipewort, 49.
 Pipewort Family, 49.
 Pipsissewa, 196.
 Pitch Pine, 38.
 Pitcher-plant Family, 124.
 Plane-tree, 129.
 Plane-tree Family, 129.
 PLANTAGINACEAE, 232.
 Plantago, 233.

- Plantain, 233.
 Plantain Family, 232.
 PLATANACEAE, 129.
 Platanus, 129.
 Pleurisy-root, 204.
 Plum, 131.
 PLUMBAGINACEAE, 196.
 Plumeless Thistle, 250.
 Podophyllum, 112.
 Pogonia, 67.
 Poison Elder, 162.
 Poison Hemlock, 187.
 Poison Ivy, 162.
 Poison Oak, 162.
 Poke Milkweed, 204.
 Pokeweed, 94.
 Pokeweed Family, 94.
 Polanisia, 124.
 POLEMONIACEAE, 206.
 Polemonium Family, 206.
 Polygala, 157.
 POLYGALACEAE, 157.
 POLYGONACEAE, 84.
 Polygonatum, 58.
 Polygonum, 85.
 Polymnia, 271.
 POLYPODIACEAE, 21.
 Polypodium, 23.
 Polypody, 23.
 Polystichum, 25.
 Pomeae, 131.
 Pond-lily, 102.
 Pondweed, 41.
 Pondweed Family, 41.
 Pontederia, 50.
 PONTEDERIACEAE, 50.
 Poplar, 73.
 Poppy, 113.
 Poppy Family, 113.
 Populus, 73.
 Portulaca, 100.
 PORTULACACEAE, 100.
 Potamogeton, 41.
 Potentilla, 134.
 Prairie Clover, 149.
 Prairie Dock, 271.
 Prairie Rocket, 120.
 Prenanthes, 273.
 Prickly Ash, 156.
 Prickly Pear, 175.
 Primrose, 197.
 Primrose Family, 197.
 Primula, 197.
 PRIMULACEAE, 197.
 Prince's Pine, 196.
 Proserpinaca, 182.
 Pruneeae, 130.
 Prunella, 219.
 Prunus, 131.
 Psedera, 167.
 Psoralea, 149.
 Ptelea, 157.
 PTERIDOPHYTES, 21.
 Pteris, 23.
 Pterospora, 196.
 Puccoon, 211.
 Pulse Family, 141.
 Purslane, 100.
 Purslane Family, 100.
 Putty-root, 67.
 Pycnanthemum, 217.
 Pyrola, 195.
 Pyroloideae, 191.
 Pyrus, 140.
 Quercitron, 79.
 Quercus, 78.
 Quillwort, 36.
 Quillwort Family, 36.

- Radicula, 117.
 Radish, 123.
 Ragweed, 251.
 Ragwort, 256.
 RANUNCULACEAE, 102.
 Ranunculus, 106.
 Raphanus, 123.
 Raspberry, 137.
 Rattlesnake Plantain, 67.
 Rattlesnake-root, 273.
 Rattlesnake-weed, 272.
 Redbud, 154.
 Rein Orchis, 64.
 RHAMNACEAE, 166.
 Rhamnus, 166.
 Rhexia, 178.
 Rhinanthus, 229.
 Rhododendron, 195.
 Rhus, 162.
 Ribes, 126.
 Ribgrass, 233.
 Richweed, 82, 217.
 Robinia, 146.
 Robin's Plantain, 265.
 Rock Brake, 23.
 Rock Cress, 118.
 Rock Elm, 82.
 Rocket, 121.
 Rock Rose, 171.
 Rock Rose Family, 170.
 Rosa, 138.
 ROSACEAE, 129.
 Rose, 138.
 Roseae, 130.
 Rose Family, 129.
 Rose Bay, 195.
 Rose Mallow, 169.
 Rose-root, 125.
 Rosin-plant, 271.
 RUBIACEAE, 233.
 Rubus, 137.
 Rudbeckia, 267.
 Rue Family, 156.
 Rumex, 88.
 Rush, 51.
 Rush Family, 50.
 Russian Thistle, 93.
 Rusty Woodsia, 28.
 RUTACEAE, 156.
 Sage-brush, 253.
 Sagina, 99.
 Sagittaria, 45.
 St. John's-wort, 169.
 St. John's-wort Family, 169.
 SALICACEAE, 70.
 Salicornia, 92.
 Salix, 70.
 Salsify, 275.
 Saltwort, 93.
 Salsola, 93.
 SALVINIACEAE, 31.
 Sambucus, 238.
 Samolus, 199.
 Samphire, 92.
 Sandalwood Family, 83.
 Sand Spurrey, 99.
 Sandwort, 97.
 Sanguinaria, 113.
 Sanguisorba, 133.
 Sanicle, 185.
 Sanicula, 185.
 SANTALACEAE, 83.
 Saponaria, 95.
 Sarracenia, 124.
 SARRACENIACEAE, 124.
 Saskatoon-berry, 141.
 Sassafras, 112.
 Satureja, 217.
 Saururus, 70.

- Savory, 217.
Saxifraga, 128.
SAXIFRAGACEAE, 125.
Saxifrage, 128.
Saxifrage Family, 125.
Scheuchzeria, 44.
Scolopendrium, 25.
Scotch Thistle, 250.
Scouring Rush, 33.
Scrophularia, 226.
SCROPHULARIACEAE, 223.
Scutellaria, 219.
Sea Blite, 92.
Sea Milkwort, 199.
Sea-rocket, 123.
Sedum, 125.
Seed-box, 180.
Selaginella, 35.
SELAGINELLACEAE, 35.
Self-heal, 219.
Seneca Snakeroot, 157.
Senecio, 256.
Sensitive Fern, 28.
Serapias, 67.
Service-berry, 141.
Shadbush, 141.
Shag-bark Hickory, 75.
Sheep-berry, 238.
Sheep Laurel, 194.
Shepherdia, 176.
Shepherd's Purse, 122.
Sherardia, 235.
Shield Fern, 26.
Shin-leaf, 195.
Shrubby Trefoil, 157.
Sickle-pod, 119.
Sicyos, 240.
Side-saddle Flower, 124.
Silene, 96.
Silphium, 271.
Silverberry, 176.
Silver-weed, 136.
Sisymbrium, 120.
Sisyrinchium, 61.
Sium, 187.
Skull-cap, 219.
Skunk Cabbage, 48.
Smartweed, 87.
Smilacina, 57.
Smilax, 56.
Snakeroot, 109, 157, 185, 255.
Sneeze-weed, 266.
Sneeze-wort, 271.
Snowberry, 193, 237.
Soapwort, 95.
SOLANACEAE, 220.
Solanum, 221.
Solidago, 257.
Solomon's Seal, 58.
Sonchus, 275.
Sorrel, 88.
Sour-gum, 189.
Sow-thistle, 275.
Sparganium, 40.
Spearmint, 216.
Spearwort, 106.
Specularia, 241.
Speedwell, 225.
Spergula, 100.
Spergularia, 99.
SPERMATOPHYTES, 36.
Spice-bush, 112.
Spikenard, 182.
Spindle-tree, 163.
Spiny Clotbur, 251.
Spiraea, 132.
Spiranthes, 68.
Spirodela, 48.
Spleenwort, 24.
Spotted Cowbane, 187.

Spring-beauty, 100.
Spring Cress, 118.
Spruce, 38.
Spurge, 158.
Spurge Family, 158.
Spurred Gentian, 201.
Surrey, 100.
Squaw Huckleberry, 193.
Squaw-root, 231.
Squaw-weed, 256.
Squirrel Corn, 114.
Stachys, 220.
Staff-tree, 164.
Staff-tree Family, 163.
Staphylea, 164.
STAPHYLEACEAE, 164.
Star Cucumber, 240.
Star Flower, 198.
Star-grass, 58, 60.
Star Thistle, 251.
Starwort, 97, 261.
Steironema, 198.
Stellaria, 97.
Stickseed, 209.
Stinking Willie, 256.
Stitchwort, 98.
Stone-crop, 125.
Stone-root, 217.
Stork's Bill, 156.
Stramonium, 222.
Strawberry, 137.
Strawberry Blite, 91.
Strawberry Bush, 163.
Streptopus, 57.
Strophostyles, 151.
Suaeda, 92.
Succory, 272.
Sugarberry, 82.
Sumach, 162.
Summer Savory, 217.

Sundew, 124.
Sundew Family, 124.
Sunflower, 268.
Swamp Dock, 88.
Swamp Loosestrife, 177.
Sweet Brier, 139.
Sweet Cicely, 187.
Sweet Clover, 145.
Sweet Colt's Foot, 261.
Sweet Fern, 74.
Sweet Flag, 48.
Sweet Gale, 74.
Sweet-gale Family, 73.
Swine Cress, 123.
Sycamore, 129.
Symphoricarpos, 237.
Symphytum, 209.
Symplocarpus, 48.

Taenidia, 186.
Tamarack, 39.
Tanacetum, 252.
Tansy, 252.
Tansy Mustard, 120.
Tape-grass, 46.
Taraxacum, 274.
Tare, 150.
TAXACEAE, 36.
Taxus, 36.
Teaberry, 194.
Tear-thumb, 87.
Teasel, 240.
Teasel Family, 240.
Tephrosia, 149.
Teucrium, 215.
Thalictrum, 105.
Thaspium, 186.
Thermopsis, 153.
Thistle, 250.
Thorn, 140.

- Thorn Apple, 222.
 Thorough-wax, 187.
 Thoroughwort, 255.
 Thlaspi, 122.
 Three-seeded Mercury, 160.
 Thuja, 39.
 THYMELAEACEAE, 175.
 Tiarella, 128.
 Tick-trefoil, 151.
 Tickseed, 269.
 Tilia, 167.
 TILIACEAE, 167.
 Toad Flax, 226.
 Tobacco, 222.
 Tofieldia, 57.
 Toothache-tree, 156.
 Toothwort, 118.
 Touch-me-not, 165.
 Tower Mustard, 119.
 Trailing Arbutus, 194.
 Tragopogon, 275.
 Treacle Mustard, 120.
 Trefoil, 144.
 Trientalis, 198.
 Trifolium, 144.
 Triglochin, 44.
 Trillium, 56.
 Triosteum, 238.
 Trumpet-weed, 255.
 Tsuga, 38.
 Tubuliflorae, 243.
 Tulip Tree, 110.
 Turnip, 121.
 Turtle-head, 227.
 Tussilago, 261.
 Twayblade, 67, 68.
 Twin-flower, 237.
 Twin-leaf, 112.
 Twisted-stalk, 57.
 Typha, 40.
 TYPHACEAE, 40.
 Ulmeae, 80.
 Ulmus, 81.
 UMBELLIFERAE, 183.
 Urtica, 82.
 URTICACEAE, 80.
 Urticeae, 80.
 Utricularia, 230.
 Uvularia, 57.
 Vaccinium, 192.
 Vaccinoideae, 190.
 Valerian, 239.
 Valeriana, 239.
 VALERIANACEAE, 239.
 Valerian Family, 239.
 Vallisneria, 46.
 Velvet-leaf, 169.
 Venus's Looking-glass, 241.
 Veratrum, 57.
 Verbascum, 225.
 Verbena, 212.
 VERBENACEAE, 212.
 Vernonia, 255.
 Veronica, 225.
 Vervain, 212.
 Vervain Family, 212.
 Vetch, 150.
 Viburnum, 238.
 Vicia, 150.
 Vine Family, 166.
 Viola, 172.
 VIOLACEAE, 171.
 Violet, 172.
 Violet Family, 171.
 Viper's Bugloss, 209.
 Virginia Creeper, 167.
 Virgin's Bower, 103.

VITACEAE, 166.

Vitis, 167.

Wake-robin, 56.

Waldsteinia, 134.

Walking-leaf, 25.

Walnut, 74.

Walnut Family, 74.

Wart Cress, 123.

Water Arum, 47.

Water Beech, 77.

Water Cress, 117.

Water Hemlock, 187.

Water Horehound, 216.

Waterleaf, 207.

Waterleaf Family, 207.

Water-lily, 101.

Water-lily Family, 101.

Water Marigold, 270.

Water Milfoil, 181.

Water Milfoil Family, 181.

Water Parsnip, 187.

Water Pennywort, 185.

Water-pepper, 87.

Water Persicaria, 86.

Water Pimpernel, 199.

Water Plantain, 45.

Water Plantain Family, 44.

Water Purslane, 180.

Water-shield, 101.

Water Star-grass, 50.

Water Starwort, 160.

Water Starwort Family, 160.

Water-weed, 46.

Water Willow, 232.

Wax-myrtle, 74.

Wax-work, 164.

White Lettuce, 273.

White Snakeroot, 255.

White-weed, 266.

Whitewood, 167.

Whitlow Grass, 121.

Whitlow-wort, 94.

Whortleberry Subfamily, 190.

Wild Allspice, 112.

Wild Balsam Apple, 241.

Wild Bean, 151.

Wild Bergamot, 218.

Wild Chamomile, 267.

Wild Comfrey, 210.

Wild Elder, 180.

Wild Ginger, 84.

Wild Indigo, 153.

Wild Liquorice, 234.

Wild Potato Vine, 206.

Wild Radish, 123.

Wild Sarsaparilla, 182.

Willow, 70.

Willow Family, 70.

Willow-herb, 178.

Wind-flower, 105.

Winterberry, 163.

Winter Cress, 117.

Wintergreen, 194.

Witch-hazel, 129.

Witch-hazel Family, 129.

Withe-rod, 238.

Wolfberry, 237.

Wolffia, 49.

Wood Betony, 230.

Woodbine, 237.

Wood Fern, 26.

Wood-nettle, 82.

Wood-rush, 51.

Wood Sage, 215.

Wood-sorrel, 155.

Wood-sorrel Family, 155.

Woodsia, 28.

Woodwardia, 25.

Worm-seed Mustard, 120.

Wormwood, 252.

Xanthium, 251.

XYRIDACEAE, 49.

Xyris, 49.

Yam, 60.

Yam Family, 60.

Yarrow, 271.

Yellow Adder's Tongue, 59.

Yellow-eyed Grass, 49.

Yellow-eyed Grass Family, 49.

Yellow Pond-lily, 102.

Yellow Puccoon, 109.

Yellow Rattle, 229.

Yellow Rocket, 117.

Yew, 36.

Yew Family, 36.

Zannichellia, 43.

Zanthoxylum, 156.

Zizia, 186.

Zostera, 44.

Zygadene, 56.

Zygadenus, 56.

1. Complete flower - four sets of floral organs present.
2. Incomplete - one or more of the sets wanting.
3. Perfect - both stamens and pistils present.
4. Imperfect - when either is wanting. If only stamens present - staminate; if only pistils present - pistillate.
5. Regular - when the parts of each set of floral organs are alike in shape and size.
6. Irregular - when the parts of each set of floral organs are not alike in shape and size.
7. Monoecious (one household) - stamens and pistils in separate flowers on same plant - plant is monoecious.
8. Dioecious (two households) - stamens and pistils in separate flowers on different plants - plant is dioecious.
9. Cohesion - union of like parts.
10. Adhesion - union of unlike parts.

CALYX.

11. Gamosepalous - sepals joined.
12. Polysepalous - sepals separate.
13. Inferior (calyx) - when attached to the receptacle below the ovary and free from it.
14. Superior (calyx) - when attached to the ovary.

COROLLA.

15. Gamopetalous - petals joined.
16. Polypetalous - petals separate.
17. Hypogynous - parts attached to the receptacle below the ovary.
18. Perigynous - parts arising from a cup surrounding the ovary (i.e. when they are inserted in the calyx).
19. Epigynous - when the parts arise from the top of the ovary.

ANDROECIUM.

20. Monadelphous - stamens united by their filaments into one group, anthers free.

21. Diadelphous - stamens united by their filaments into two groups.

22. Syngenesious - stamens united by their anthers, filaments free.

23. Hypogynous, perigynous, epigynous.

24. Epipetalous - on the petals.

25. Tetradynamous - 6 stamens, 4 long and 2 short.

26. Didynamous - 4 stamens, 2 long and 2 short.

27. Androus - term applied to free stamens - pentandrous, hexandrous and polyandrous.

GYNAECIUM.

28. Apocarpous - carpels free.

29. Syncarpous - carpels united.

30. Monocarpous - one carpel.

Date Due

OCT 13 1971	STEACIE	DEC 10 1973	
OV 5 - 1971	STEACIE	DEC 12 1975	
OCT 22 1971	STEACIE	DEC 19 1975	
.....			
STEACIE	NOV 21 1972		
STEACIE	NOV 1 1972		
STEACIE	NOV 28 1972		
STEACIE	NOV 8 1972		
STEACIE	DEC 27 1972		
STEACIE	DEC 20 1972		
STEACIE	JUN 3 1973		
STEACIE	MAY 2 1973		
STEACIE	DEC 27 1973		

64

CAT. NO. 23 233

PRINTED IN U.S.A.

14522 02