Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

109
P69P
copy

THE PLANT DISEASE REPORTER

Issued By

THE PLANT DISEASE SURVEY

Division of Mycology and Disease Survey

BUREAU OF PLANT INDUSTRY, SOILS, AND AGRICULTURAL ENGINEERING

AGRICULTURAL RESEARCH ADMINISTRATION

UNITED STATES DEPARTMENT OF AGRICULTURE

SUPPLEMENT 163

✓
ROOTROTS AND LEAFSPOTS OF GRAINS AND GRASSES IN THE
NORTHERN GREAT PLAINS AND WESTERN STATES

Supplement 163

June 15, 1946

The Plant Disease Reporter is issued as a service to plant pathologists throughout the United States. It contains reports, summaries, observations, and comments submitted voluntarily by qualified observers. These reports often are in the form of suggestions, queries, and opinions, frequently purely tentative, offered for consideration or discussion rather than as matters of established fact. In accepting and publishing this material the Division of Mycology and Disease Survey serves merely as an informational clearing house. It does not assume responsibility for the subject matter.

PLANT DISEASE REPORTER SUPPLEMENT

Issued by

THE PLANT DISEASE SURVEY
DIVISION OF MYCOLOGY AND DISEASE SURVEY

Plant Industry Station

Beltsville, Maryland

ROOTROTS AND LEAFSPOTS OF GRAINS AND GRASSES
IN THE NORTHERN GREAT PLAINS AND WESTERN STATES¹

Roderick Sprague²

Plant Disease Reporter
Supplement 163

June 15, 1946

INTRODUCTION

As delimited by the title, this annotated list contains at least mention of all the parasites known by the writer to attack Gramineae in the western United States. Since he has experienced difficulty in staying abreast of the literature on these some 200 species of fungi on nearly 500 hosts, the writer has assembled this material primarily for his own use. While the resulting preliminary work is not a text, it is somewhat more than a list and it seems that it may have some value to a number of western experiment station workers and field men in the routine diagnosis and study of the lesser known leafspot and rootrot diseases of grasses and cereals. Since brevity has been necessary it appeared to the writer that some of the better known parasites

¹ Cooperative investigations between the Divisions of Cereal Crops and Diseases, Forage Crops and Diseases, and Soils, Fertilizers, and Irrigation, Bureau of Plant Industry, Soils, and Agricultural Engineering, Agricultural Research Administration; and the Division of Nurseries, Soil Conservation Service, U. S. Department of Agriculture; and the North Dakota Agricultural Experiment Station.

² Pathologist, Division of Cereal Crops and Diseases, and Collaborator, Division of Forage Crops and Diseases, Bureau of Plant Industry, Soils, and Agricultural Engineering, Mandan, N. Dakota.

could be given less space than some of the lesser known ones, as information on the latter frequently is buried in obscure journals. Most of the important diseases of cereals have been well covered in such standard texts as Heald's Manual (316)³, Owens (497), Chester (103), and Dickson (160). This article, then, may be useful as a supplement to available texts.

This outline is the result of work started about 15 years ago at Corvallis, Oregon, and is partly a direct outgrowth of several check lists issued in 1935, 1937, and 1942. The writer felt that the re-grassing program, started during and after the dry years of the last decade, would eventually require more information on the fungi of Gramineae than was then available. There were others soon interested in studying this neglected field. The recent and current work, for instance, of Fischer on fundamental taxonomy of the smuts of grasses has added a great deal of needed data on this group. Research in the Canadian provinces on rootrots, the studies made at Iowa State College on Pythium spp., and the publication of such texts as Grove (296, 297) and Sampson and Western (572) have aided workers during recent years. There is a new publication by Garrett (257), which gives a modern and understandable summary of the epidemiology of rootrots in general. The recent text by Gilman (262) is also helpful in determining common soil molds associated with parasitic root fungi, as are texts by Thom (705), Jensen (362), Boedijn (55), and Clements and Shear (117), and the Check-list by Weiss (755-757).

The region covered herein is by no means thoroughly explored from our viewpoint. However, during the past decade various workers have collected and studied intensively in widely scattered parts of it. Probably the greatest impetus to this work has been the increase in knowledge of the hosts through the publication of Hitchcock's manual of the grasses (326) and the increase in the number of workers who are able to determine the hosts with a fair degree of accuracy. We therefore find that the western part of the United States is, in general, as well covered as the eastern part. Seymour (599) contains about 2,820 citations (hosts x fungi) of parasitic fungi for all of North America on Gramineae, while the unpublished list of the writer and Fischer includes 3257 citations for the western area alone. Since Seymour's list appears to contain very few western citations other than rusts and smuts it is apparent that the recent investigations have greatly expanded the host range on Gramineae in this once neglected region.

This article includes reference to about 221 species and sub-divisions of species of parasitic fungi found growing naturally in the region. There are 13 Phycomycetes, 24 Ascomycetes (if we include 15 species of Phyllachora), 8 Basidiomycetes, and 162 Fungi Imperfecti, as well as a few undetermined or non-parasitic troubles. In addition, brief mention

3

Figures in parentheses refer to "Literature cited," p. 215.

is made of associated non-parasitic saprophytes which are sometimes such early invaders of dead tissue that they have been considered as weakly parasitic in some cases. The following table indicates the number of species of leafspot and rootrot-causing fungi reported on Gramineae in the several western States.

Number of species found in:	Class of fungi					Total
	<u>Phyco- mycetes</u>	<u>Asco- mycetes</u>	<u>Basidio- mycetes</u>	<u>Fungi Imperfecti</u>	<u>Miscellany</u>	
North Dakota	8	11	3	79	4	105
South Dakota	7	9	1	49	0	66
Minnesota	6	7	4	43	2	62
Iowa	7	7	2	35	1	52
Nebraska	5	10	1	42	0	58
New Mexico	1	9	0	7	0	17
Arizona	1	5	0	5	2	18
Colorado	1	5	0	13	1	20
Utah	0	3	2	10	1	16
Wyoming	2	3	2	40	0	47
Montana	8	5	4	45	0	62
Idaho	0	4	3	30	2	39
Washington	5	6	4	67	2	84
Oregon	6	7	4	97	1	115
California	3	9	1	47	3	63
Nevada	0	0	0	1	0	1
Alaska	0	1	0	5	0	6

It is evident from the table that Oregon, North Dakota, and Washington have been the most thoroughly explored. Some States such as Nevada have very few fungi reported on Gramineae other than rusts and smuts. Of course climate influences these records tremendously. The coast region of Oregon, Washington, and northern California is highly favorable to the development of leafspot fungi while some of the Rocky Mountain areas also harbor a large number of them. The arid plains have few leaf spots but many rootrot fungi and late-summer leaf spots occur in the Northern Great Plains. The wooded and swamp areas of Minnesota contain numerous leaf spots.

An intensive survey has been conducted in Washington and North Dakota on exotic hosts in the large grass nurseries at Pullman and Mandan. Therefore the greatest host range (fungi x hosts) occurs in North Dakota with 1,369 citations to all fungi including smuts and rusts. Washington has 971 citations, Oregon has 912 (mostly on native species), South Dakota 497, Montana 460, Minnesota 445, California 437, Nebraska 331, Iowa 327, Wyoming 282, Colorado 264 (mostly rusts and smuts), Idaho 262, Utah 152, New Mexico 133, Arizona 133, Alaska 40, and Nevada 39. This indicates approximately which States need additional survey information.

While the majority of the cereal diseases are well covered in plant pathological literature, many of the fungi causing grass diseases are scarcely mentioned except for the original mycological description. Therefore it is still necessary for workers with these fungi to be acquainted with the location of these descriptions as well as where the type material or comparable exsiccata may be found. Material available from the western area is not plentiful among the earlier collections. David Griffith's Western American Fungi contains numerous smut fungi but few of the fungi that fall within the bounds of this study. Brenckle's Fungi Dakotensis, Ellis and Tverhart's and associates' various collections, and Cooke's recent Mycobiota series contain some collections, while Solheim's Mycoflora Saximontanensis is also useful. Most of the material dealt with by the writer is filed either at Oregon State College, Corvallis, Oregon; or is in the Mycological Collections of the Bureau of Plant Industry, Soils, and Agricultural Engineering, U. S. Department of Agriculture, Beltsville, Maryland. Some of this material is scanty and of use for record only, but an attempt was made to preserve a fairly complete record of recent collections.

In this article, the fungi are listed alphabetically under the classes: (1) Phycomycetes, page 105, (2) Ascomycetes, page 111, (3) Basidiomycetes, page 116, and (4) Fungi Imperfecti, page 120. No attempt has been made to give an exhaustive list of references, but instead only those are included that are considered necessary for a working basis. It has not been possible to check all references against the originals and therefore considerable use has been made of abstract journals.

In discussing the various fungi, usually brief summaries are given of

symptoms, host range, and sometimes morphology; and less often of taxonomy.

PHYCOMYCETES

APHANOMYCES SPP. Root Necrosis.

Pure cultures of Aphanomyces have been isolated from oats growing in acid soil in northern Minnesota, eastern Montana, and in the coast region of Oregon. Scattered isolates have also been obtained from meadow fescue (Festuca elatior L.), red fescue (F. rubra L.), and green needle grass (Stipa viridula Trin.) from Mandan, North Dakota, and from Setaria viridis (L.) Beauv. mingled with oats in eastern Montana. The only place where there was very much evidence of damage was in the very red, very acid soil in Lincoln County, Oregon, and even in this instance the fungus was associated with so many other organisms that its exact role was not readily determinable (Sprague, 639). The culture from northern Minnesota appears to be A. camptostylus Drechs., which is a somewhat delicate-appearing fungus originally isolated from diseased oat roots at Sauk City, Wisconsin, by M. B. Linford. The writer worked with this fungus in 1926 at Madison and has found that some of his recent isolates were similar in appearance. No study has been made with these forms in North Dakota as yet, because they appear to be less important than a number of other fungi. However, as species of Aphanomyces are somewhat difficult to isolate, they may be more prevalent than our occasional isolates would indicate. They are evidently of more importance in northern Minnesota and in the coast region in Oregon.

References: Drechsler (181, figs. 9-11); Sprague (639).

PHYTOPHTHORA SPP., Rot.

Phytophthora cactorum (Leb. and Cohn) Schroet. and some undetermined forms are sometimes found associated with soft rots of the leaves and roots of grains and grasses. While corn is the most frequently reported host (Sideris, 606), the following have been noted: quack grass (Minnesota), Triticum durum Desf., Agropyron sibiricum (Willd.) Beauv., Hordeum jubatum L., and Stipa viridula Trin. in North Dakota, Lolium multiflorum Lam. (Oregon), Poa vaseyochloa Scribn. (Oregon), and Avena fatua L. (Oregon and California). This fungus needs abundant moisture to be able to develop parasitically.

McMurphy (450) noted a species similar to Phytophthora calocasiae Rac. (Tucker, 716) on oats in California. The infected areas were at first yellowish, then whitish, and finally brown or reddish brown. The fragmentary material seen by the writer on wild oats (Avena fatua) near The Dalles, Oregon, and in northern coastal California were only

water-soaked leaves in contact with the soil. The North Dakota isolations were from roots of seedlings and were usually associated with other fungi. The most abundant material that the writer has obtained was in stunted plants of barnyard grass growing near Belle Fourche, South Dakota. This material appeared to be P. cactorum. A similar fungus was isolated from corn roots near Newell, South Dakota, at the same time. In the greenhouse at Mandan artificial inoculations with this fungus (inoculum added to soil at seeding time) caused 88 percent loss in Turghai proso, only 6 percent loss in crested wheatgrass, scarcely a trace in wheat, 70 percent in Ladak alfalfa, 71 percent in blue grama, and only a trace in corn. Its injury was confined to seed rot at the temperature employed (48° to 70° F). In general, Phytophthora is not important on Gramineae.

References: McMurphy (450); Sideris (606); Tucker (716, 717).

(?) SYNCHYTRIUM SP., Purple Leaf Speckle.

In 1942, the writer (656) stated that this fungus was a "primitive unilocular body or cluster of large cells imbedded in a leaf spot." There is no appreciable gall formation and for that reason it was doubted that the fungus was a species of Synchytrium. It appeared to be nearest to Eurychasmidium (Karling, 377) of the Ectrogellaceae. However, recent illustrations by Cook (124) of species of Synchytrium from Louisiana more strongly suggest that this fungus could be placed in Synchytrium. We have not been able to find mention of a similar fungus on grass unless it be the somewhat vague Protomyces rhizobius Trail, which occurs on Poa annua L. in Scotland. There is, however, little or no mycelium present in the material from Oregon and Washington. Protomyces is placed in either the Phycomycetes or the Ascomycetes, depending on the authority.

In Oregon and Washington, the fungus is relatively common but seldom destructive during open, wet, winter weather in the Willamette Valley of Oregon, and the coastal region of both States. It is sometimes found on bent grass lawns at Corvallis. It occurs on Agrostis alba L., A. canina L., A. exarata Trin., A. hallii Vasey, A. palustris Huds., A. tenuis Sibth., Anthoxanthum odoratum L., Bromus carinatus Hook. and Arn., Dactylis glomerata L., Elymus glaucus Buckl., Festuca rubra var. commutata Gaud., Poa annua L., P. juncifolia Scribn., P. pratensis L., and P. secunda Presl. Representative material is filed at Oregon State College and at Harvard University.

References: Cook (124); Karling (377); Sprague (655, 656).

PYTHIUM ARISTOSPORUM Vanterpool
Root Browning of Grains, Seedling Blight of Grasses.

This fungus is closely related to the much more abundant Pythium arrhenomanes Drechs. In the United States it has been isolated from wheat, crested wheatgrass (Agropyron cristatum (L.) Gaertn.), orchard grass (Dactylis glomerata L.), barnyard grass (Echinochloa crusgalli (L.) Beauv.), and tall meadow oatgrass (Arrhenatherum elatius (L.) Beauv.) at Bozeman, Montana, on barley at Dickinson, North Dakota, and on this host and oats near Eugene, Oregon, and on slender wheatgrass (Agropyron trachycaulum (Lk.) Malte) at Mandan, North Dakota. Like P. arrhenomanes this species produces a wet-appearing pebbly growth on potato-dextrose agar, but the oogonia with their numerous antheridia do not disintegrate quite as fast as do those of P. arrhenomanes. The oogonia also appear to be larger in P. aristosporum.

References: Sleeth (614); Vanterpool (739, Fig. 2, Pl. XXII; 743); Vanterpool and Sprague (744).

PYTHIUM ARRHENOMANES Drechs.
Root Browning of Grains, Seedling Blight of Grasses.

Root browning is characterized by dark brown lesions on the small roots of small grains (wheat, oats, barley) in late May and June in the Northern Great Plains and adjacent regions. The plants when severely infected show a pale green color in contrast with healthy ones. This fungus is very widespread in all of the northern plains States and also causes a serious seedling blight of sorghum in the Sacramento Delta region of California. It is the well-known cause of Milo disease in the Great Plains. The most serious aspect of this fungus is its effect on spring-seeded grasses in North Dakota, South Dakota, Nebraska, and Minnesota. It causes a blight of the young plants, which wither and die about 6 weeks after seeding. During some years, almost 100 percent kills have occurred in crested wheatgrass, slender wheatgrass, and smooth brome (Bromus inermis Leyss.) at Brookings, South Dakota, and in places in North Dakota. The host range is very great (Vanterpool and Sprague 744). Curzi (134) reports a similar disease on cereals in Italy but the identity of the Italian fungus has not yet been determined.

Drechsler (185) contends that his Pythium arrhenomanes (179) is distinct from P. graminicolum Subr. (692) because, "the close mycelial connection between oogonium and antheridium, very frequent in P. graminicolum, is rare in P. arrhenomanes; and in parallel cultures the sturdy, more substantial membranous parts of the sexual apparatus of the former species remain clearly discernible long after the evanescent antheridial envelopes and supporting branches of the latter have become nearly or wholly invisible." On the basis of this distinction the material in the Northern Great Plains is P. arrhenomanes, except P. aristosporum which has the long-lived traits of P. graminicolum in

pure culture. Both P. arrhenomanes and P. graminicolum are recognized by Matthews (440).

Root browning in Saskatchewan (Vanterpool, 740, 742) is partly checked by judicious use of phosphate fertilizers wherein a proper balance of nitrogen is maintained. Hanson (306) has obtained striking results in the control of rootrot at Lamberton, Minnesota, by the use of phosphate or phosphate and nitrogen fertilizers on wheat. The writer obtained similar results in the coast region of Oregon, but in the Northern Great Plains where most areas are not yet deficient in phosphorus and where nitrogen is not seriously deficient, fertilizing has not had much effect on root browning, except that recent trials with ammonium phosphate (16-20-0) indicate that this material may have some effect on the parasite as well as on the host. Partial control of the rootrot complex in wheat with 100 lbs. of ammonium phosphate per acre in trials at McCanna, North Dakota, have shown particular promise (Sprague, annual report, 1945).

Since Pythium arrhenomanes is restricted to Gramineae, use of non-grass plants in rotation reduces its injury. Among cereals oats are most tolerant; proso millets (Panicum miliaceum) or sorghums usually are less desirable to grow. While fallow does not appear to increase the amount of Pythium it creates the same effect by starving out other organisms, permitting the Pythium forms temporary ascendancy. Small-seeded grasses should not be seeded on fallow if it can be avoided, at least not in the spring of the year. Crested wheat grass, however, can be sown on fallow under most conditions. Fall seeding is preferred to spring seeding where conditions permit. Even winter seeding of crested wheatgrass is preferred if periods of open weather occur as is sometimes the case in North Dakota.

References: Andrews (14); Bowman et al. (61); Branstetter (63) Buchholtz (73); Carpenter (92); Connors and Savile (123); Curzi (134); Drechsler (179, 185); Edgerton, Tins, and Mills (194); Elliott et al. (204, figs. 1-8, 17, 21); Flor (232); Hanson (306); Ho and Melhus (330); Ho et al. (331); Ho and Koepper (329); Hoffmaster (333); Johann (364); Kendrick and Briggs (380); Leukel (403, fig. 1); Matthews (440); Melchers (455); Melchers and Lowe (456, 457); Melhus, Martin, and Murphy (460); Middleton (465); Rands and Dopp (523, 524, 525); Roldan (539, 540); Sallans (567); Simmonds, Russell, and Sallans (611); Sprague (658, 661, 663); Sprague and Atkinson (670); Subramaniam (692); Vanterpool (737, 738, 739, 740, 741, 742, 743); Vanterpool and Sprague (744); Vanterpool and Truscott (745); Wagner (749).

PYTHIUM DEBARYANUM Hesse, P. IRREGULARE Buis., P. ULTIMUM Trow
Seed rot, Damping-off and Root Necrosis.

Although two of the originally recognized hosts of P. debaryanum were Panicum miliaceum L. (proso millet) and Zea mays L. (Hesse, 325), this fungus is little recognized as a parasite of Gramineae. However, Eriksson (219) and also Gram and Rostrup (281) report it on barley, and there are a number of citations on cane crops, including corn, (Ho, 328). In the Northern Great Plains, P. debaryanum and related species cause considerable seed rot in wet soil in the spring and may cause damping-off or seedling injury to some small-seeded grasses (Sprague, 665). Sometimes when cereals are planted in low wet ground or when old moldy seed is planted in heavy ground Pythium will cause seed rot. The injury in most cases in both grasses and cereals occurs just as the seed sprouts. The rootlets are unable to emerge. In the greenhouse under artificial conditions some grasses have the roots reduced to yellow or brown stubs even after emergence. There are various strains of these fungi that will attack some hosts more readily than others. Some will attack certain cereals, such as wheat, barley, and oats, more severely than others; some strains from grasses will attack many species of grasses but are mildly parasitic on non-grass crops; while in other cases the reverse is true. In general, however, the host range is so great that crop rotation to control this group would be very difficult to work out. Since legumes, notably alfalfa, are very susceptible to P. debaryanum (Buchholtz, 79) and fallow seems to favor it, the control of P. debaryanum becomes even more difficult than that of P. arrhenomanes, which is confined to the grass family. However, P. debaryanum is usually not an important parasite on wheat and usually does not cause the complete destruction on grasses that seedling blight (P. arrhenomanes) does. Buchholtz (78) has shown that seedling blight symptoms are produced by P. arrhenomanes (called P. graminicolum by him). We have found that recovery from P. debaryanum is very rapid and from P. arrhenomanes very slow, in many grasses, in the recovery period after the June seedling blight days have passed. P. debaryanum is, therefore, mainly important as a seed rot but can cause some root necrosis. Oats are somewhat more susceptible to P. debaryanum, especially during cool, wet weather (Welch, 758; Gram and Rostrup, 281; Beaumont, 29).

Pythium debaryanum, P. ultimum, and P. irregulare belong to the sphaerosporangial group of Pythium spp., that is the sporangia are spherical or lemon-shaped. On the other hand, P. aristosporum and P. arrhenomanes produce irregular, lobed or lobulate swollen bodies, from which stalked spherical sporangia develop later. P. debaryanum is by far the most common of the sphaerosporangial forms on Gramineae in North Dakota. P. ultimum is common in the Palouse region near Pullman, Washington, in the heavy Palouse silt loam in early season. P. irregulare is scattered but not uncommon on oats and brome grasses (B. inermis, B. tectorum L.). It is distinguished from the other two by its irregular oogonial wall with prominent projections. P. debaryanum produces zoospores from the sporangia, but P. ultimum germinates

direct by sending out germ tubes from the sporangia. In addition, in P. ultimum the antheridia arise from the oogonial stalk while in P. debaryanum the several antheridia are likely to be produced from distinct, distant hyphae. P. vexans DBY. is another sphaerosporangial species sometimes encountered on grasses and cereals, more or less readily distinguished by its broadly appressed antheridial attachment. Also, P. hypogynum Middleton (464) probably occurs in North Dakota on wheat (Sprague isolation notes). This fungus has the antheridium growing up through the oogonial stalk. Still another related species, P. iwayamai Ito (Ito and Tokunaga 355), was said to cause a snow rot disease of cereals in Japan (Iwayama, 356), but this has not been seen in the United States. Biraghi (48) reported P. polymorphon Sideris (606), which may be the same as P. irregulare, on wheat from Italy.

References: Beaumont (29); Biraghi (48); Buchholtz (78, 79); Butler (87); Drechsler (178); Eddins (192); Eriksson (219); Gram and Rostrup (281); Grandfield, Lefebvre, and Metzger (282); Hesse (325); Ho (328); Ho and Melhus (330); Höhnke (335); Ito and Tokunaga (355); Iwayama (356); Matthews (440, pl. 17, fig. 17-21, 22, 24, 25); McLaughlin et al. (449); Melhus et al. (461); Meredith (462, 463); Middleton (464, 465); Miura (471); Petri (513); Ramakrishnan (522); Sideris (606); Sprague (658, 663, 664); Spychneva (677, 678); Van Luijk (736); Vanterpool (743); Welch (758, 759).

PYTHIUM MONOSPERMUM Pringsh., Root Necrosis.
(P. complens A. Fischer)

Pythium monospermum was isolated from mildly necrotic roots of Turghai proso millet at Mandan, North Dakota, and from Sporobolus cryptandrus (Torr.) A. Gray near Buffalo, South Dakota. It has filamentous or semi-lobulate branched sporangia. An inoculation trial at Mandan with the Mandan isolate caused 62 percent loss in crested wheatgrass, 100 percent loss in alfalfa and blue grama (Bouteloua gracilis (H.B.K.) Lag.) 31 percent in wheat, 28 percent in oats, and only 4 percent in corn. It should be added that Drechsler (177) could not infect cabbage heads with P. monospermum, but with the culture at Mandan, 98 percent pre-emergence seedling kill was obtained in Chinese cabbage and 100 percent in tomatoes. P. monospermum appears to be rare in the Northern Great Plains.

References: Drechsler (177); Matthews (440); Middleton (465).

PYTHIUM PERIILUM Drechs., Root Necrosis.

What appears to be Pythium periilum was isolated from the roots of quack grass (Agropyron repens (L.) (Beauv.) in a gray, glacial soil near Bemidji, Minnesota. The oogonia were similar to those illustrated by

Drechsler in one of his remarkable drawings (186, fig. 5). The culture from the Bemidji material died before its pathogenicity could be tested. Stevenson and Rands (685) and Rands and Dopp (524) stated that it was feebly parasitic on sugar cane roots. Further study is needed with this Minnesota form.

References: Drechsler (183, 186); Rands and Dopp (524); Stevenson and Rands (685).

PYTHIUM ROSTRATUM Butl., Root Necrosis.

Pythium rostratum was isolated in May 1942 from roots of Avena sativa L. at Pullman, Washington. It is generally considered to be a soil saprophyte and no data is available on whether it is parasitic on oats. It was associated with mild root necrosis.

References: Butler (86); Matthews (440, pl. 17); Middleton (465).

PYTHIUM TARDICRESCENS Vanterpool, Root Browning.

Vanterpool found Pythium tardicrescens moderately common in Saskatchewan, Canada, but it has been found only twice in the United States, once on Avena fetua L. at Pullman, Washington, and once on barnyard grass at Graceton, Minnesota. It produces a firm, dark brown rot or root browning, and in our trials has about the same host range and relative pathogenicity as the other lobulate sporangial species such as P. arrhenomanes. The oogonia generally average smaller than P. arrhenomanes and have a characteristic sub-globulate content. The fungus produces a weak scanty growth in culture on corn-meal agar and only a few oogonia mature.

References: Sprague (658; 663); Vanterpool (739, fig. 1, pl. XIII; 740; 743).

ASCOMYCETES

DOTHICHLOË LIMITATA Diehl, Leaf Disease.
(D. atramentosa (Berk. and Curt.) Atk.)

Diehl (170) described Dothichloë limitata as having white conidial masses on the upper leaf surfaces of infected leaves. While the fungus is largely found in warm temperate areas, it ranges as far north as North Dakota where it occurs on Calamagrostis inexpansa A. Gray. Diehl believes that this fungus is systemic and that it causes sterility in

the grasses it attacks. Low temperatures inhibit its fructification and no doubt limit its northward distribution.

References: Chardon (102, pl. 14, fig. 12); Diehl (168, pl. 2, D, E, and fig. 3, B; 170, pl. 1).

DOTHIDELLA ARISTIDAE (Schw.) Ell. and Ev.

(See Septogloeum oxysporum Sacc., Bomm. and Rouss. in the Fungi Imperfecti, p. 170).

EPICHLÖE TYPHINA (Fr.) Tul., Cat-tail Disease or Choke.

Epichloë typhina forms a cylindrical collar of creamy, later dark orange stroma around the culms of various grasses, causing some sterility. In North Dakota it has been found on Canada bluegrass (Poa compressa L.), Kentucky bluegrass (P. pratensis), Sandberg bluegrass (P. secunda), western wheatgrass (Agropyron smithii Rydb.), slender wheatgrass (A. trachycaulum), Virginia wild-rye (Elymus virginicus L.) and Junegrass (Koeleria cristata (L.) Pers.). At Pullman, Washington, it is found on Agropyron spicatum (Pursh) Scribn. and Sm., A. trachycaulum, Elymus canadensis L., E. dahuricus Turcz., and Poa stenantha Trin. It occurs also at Whitehall, Montana, on Agropyron smithii. It occurs on timothy and Kentucky bluegrass in Nebraska, on slender wheatgrass in Montana, on Junegrass in South Dakota, on A. inerme (Scribn. and Sm.) Rydb. at Athol, Idaho, and on Hystrix patula Moench in Minnesota. It is more spectacular than important in most cases.

References: Sampson (568, 569); Stevens (682).

GIBBERELLA ZEAE (Schw.) Petch. See Fusarium graminearum, Schw. p. 181.

LEPTOSPHAERIA HERPOTRICHOIDES De N., Secondary Rootrot.

Leptosphaeria herpotrichoides was, for many years, believed to be the main cause of straw-breaker footrot of cereals, especially wheat, in Europe. Research has shown that it is actually only a questionable, weak parasite (Sprague, 620; 622; Sprague and Fellows, 671) and it is seldom mentioned in recent literature, Cercospora herpotrichoides Fron having replaced it as the recognized cause of the straw-breaker disease in Europe and the Pacific Northwest.

Leptosphaeria herpotrichoides was found by the writer on dead culms of rye near Leonard, North Dakota, in 1940 and was reported on wheat from Peone Prairie, Washington, by McKinney (446, p. 29).

References: Delacroix (147); Henry and Foster (324); Mangin (432); McKinney (446); Sprague (620, 622); Sprague and Fellows (671).

MYCOSPHAERELLA TULASNEI (Jancz.) Rothers, Leaf Mold
(Cladosporium herbarum Lk.)

The conidial stage (Cladosporium herbarum) is an omnipresent saprophyte. There are some indications that in early spring it is parasitic on the leaves of certain grasses, notably western wheatgrass at Mandan and Idaho fescue (Festuca idahoensis Elmer) in the far west. It produces brown flecks on the young leaves and forms a brown to olive-black mold on any of the above-ground parts after the plant dies. There is some doubt about whether it is parasitic on the young leaves of western wheatgrass at Mandan, North Dakota, because it usually follows early spring low-temperature conditions. In addition to the early season condition, it is often associated with leaf rusts and with such leaf parasites as Septoria nodorum Berk. It is seldom isolated from the roots of cereals and grasses.

The conidiophores of this species form dense tufts of velvety olive-black color. The conidiophores are more or less erect, septate, sparsely branched; spores often in chains of 2 or 3, sub-cylindric, pale olive, 1-(2-3) septate, 10-15 x 4-7 μ . Heald and Ruehle (317) presented evidence that the microsporous stage, Hormodendrum cladosporioides Sacc, is not conclusively part of the cycle of M. tulasnei. They verified (Ruehle 547, Heald and Ruehle, 317) the work of Janczewski (360) who determined that M. tulasnei was the perithecial stage of C. herbarum. The writer obtained M. tulasnei in a pure culture of C. herbarum in 1943, isolated from Stipa viridula in western North Dakota.

Ruehle in his work with fruit rots (547) reported the perithecia of Mycosphaerella tulasnei as black with thick walls, typically broadly flask-shaped with a short neck, erumpent, 150-250 x 100-150 μ ; asci cylindrical, fusoid or slightly tapering at the ends, 80-120 x 15-20 μ ; ascospores oblong, bluntly pointed, 1-septate, upper cell somewhat larger than the lower one, 18-28 x 6- 8.5 μ .

Cladosporium herbarum is probably one of several species of this genus on Gramineae, but without sufficient critical modern study, the writer follows precedent in assigning them all to C. herbarum. C. graminum Cda. is frequently used also. Its distinction from C. herbarum is not clear. Schnellhardt and Heald (586) report C. malorum Ruehle as common on wheat seeds in Washington.

Cladosporium herbarum is probably not important in its role as mold on shocked or standing grain but seed treatment with New Improved Ceresan is recommended, as its presence often indicates occurrence of more virulent forms also.

Mycosphaerella tassiana (De N.) Johans. has been found on grasses in the far west but it is evidently not related to the C. herbarum complex. Cash (97) listed M. longissima Fekl. on Bromus secalinus L. from Sitka, Alaska.

References: Bennett (31); Bockmann (54); Cash (97); Heald and Ruehle (317, fig. 9; 13, Q, P, Q); Janczewski (360); Ruehle (546, 547); Schnellhardt and Heald (586).

OPHIOBOLUS GRAMINIS Sacc., Take-all.

In the region covered by this paper, take-all occurs in Oregon, Washington, Idaho, and northern California. During the historically open winter of 1933-34, it was found on the Idaho prairies near Nezperce, where it occurs but is so inhibited during most years as to be unrecognized. In addition to western Oregon, it occurs also in trace amounts in the Grand Ronde Valley, Oregon. In 1930 it was found in irrigated spring wheat at Jerome, Idaho (Sprague, 621).

Take-all appears to be native in the Pacific Northwest, although it was not noticed until 1901 at Albany, Oregon (Cordley, 129)⁴. It occurs in virgin soils on Holcus lanatus L., high in the hills of the Coast Range in Oregon and has been found on Bromus orcuttianus Vasey in the Siskiyou Mts., Oregon. It is particularly common on annual fescues (Festuca dertonensis (All.) Aschers. and Graebn., F. megalura Nutt., F. myuros L.) in the Willamette Valley, in Newberg sandy loam, and in the greenhouse at Corvallis, Oregon, it is a nuisance in unsterilized soil. While O. graminis is important mainly on winter wheat and winter barley, it has been collected also on winter oats in acid soil in coastal Oregon and California. It was relatively common on these hosts during the year 1933-34 and was found in the Willamette Valley as well as along the coast. A trace of take-all was noted on Poa canbyi (Scribn.) Piper at Pullman, Washington. Fraser, Simmonds, and Russell (238,a) believed that O. graminis was indigenous to north-central Alberta.

The work of Fellows, Garrett, and the Canadian and Australian workers, has made O. graminis one of the most thoroughly studied fungi in the cereal rootrot complex. It needs plenty of study as it is a very uncertain parasite, here today and gone tomorrow. The worst infection the writer has ever encountered was in a plot of wheat seeded on land that had been in orchard for two decades or more.

Garrett's text (257) deals thoroughly with this fungus.

4

The still unlabeled specimens of this 1901 material were seen by the writer in 1933. They were definitely Ophiobolus graminis.

References: Åkerman et al. (3); Adam and Colquhoun (5); Broadfoot (69, 70, 71); Brommelhues (76); Buddin and Garrett (80); Bussman (85); Carne and Campbell (91); Clark (115); Cordley (129); Dana (137); Davis, R. J. (145); Fellows (223-228); Fellows and Ficke (230); Foster and Henry (236); Fraser (238); Fraser, Simmonds, and Russel (238,a); Garrett (242-258); Garrett and Dennis (259); Glynne (263); Griffiths (294); Henry (323); Hynes (346); Kirby (383); Laar, van de (395); Lal (396); Ludbrook (414); Ludwig and Henry (415); Mackie (423); Mangin (432); McAlpine (443); McKinney (446); McKinney and Davis (447); Melchers and Sewell (458); Moritz (474, 475); Müller-Kögler (478); Nattrass (480); Noll (490); Osborn (496); Padwick (498, 499); Padwick and Henry (501); Robertson (537); Russell (548, 549); Samuel (573); Sanford (578); Sanford and Broadfoot (579); Simmonds (608); Simmonds, Russell and Sallans (611); Sprague (621); Stevens (683); Stumbo et al. (691); Turner (720, 721); Walker (750); White (764, 765); Winter (769-772, 774).

PHYLLACHORA SPP., Tar Spots.

The black, usually elongate, tar-like spots on grass leaves caused by various species of Phyllachora are classed as leaf spots. However, they represent a specialized group and as such have been carefully monographed recently by Orton (495). Since virtually all of the western material, including that of the writer, is discussed by Orton it is deemed satisfactory to refer this group to his work rather than discuss the group individually here. The publication (*Mycologia*) is readily available.

The tar spots are widespread and sometimes moderately important in reducing foliage, particularly on species of Muhlenbergia and on Elymus canadensis in our area. Studies on the life history of these forms is needed, particularly in reference to associated conidial and pycnidial forms.

Reference: Orton (495).

PYRENOPHORA BROMI (Died.) Drechs.

(The writer prefers to discuss this fungus under its conidial stage, Helminthosporium bromi Died., as it is one of a large number of species in this genus.)

SCLEROTINIA HOMOECARPA F. T. Bennett, Sclerotinia Turf Rot.

Bennett (38) in England concluded that his earlier described Rhizoctonia monteithianum F. T. Bennett (36) was a species of Sclerotinia. R. monteithianum was said to be the same as the small Rhizoctonia turf or greens disease known generally as dollar spot (Monteith and Dahl, 472).

Since dollar spot occurs on the eastern edge of our area, S. homoeocarpa is included until critical work is available to verify or reject this classification. Probably the same fungus or fungi occur west of the Cascades in Oregon and Washington.

References: Bennett (36, 38); Keil (379); Monteith and Dahl (472); Struble (690); Tervet (704); Tyson (725).

BASIDIOMYCETES

ENTYLOMA SPP., Blister Smut on Leaves of Bluegrasses

The blister smuts of Poa spp. should be mentioned here because they can be and have been confused with tar spot fungi (Phyllachora). The stroma-like sub-epidermal chlamydospores form gray-black blister-like areas on the leaves of Poa pratensis, P. compressa, P. juncifolia, and P. annua in Oregon and adjacent Washington. The conidial phase of the fungus forms a fine, dried, amorphous series of dots scattered seriate-ly along the surface of the blisters. The Ramularia-like sporidia of Entyloma irregulare Johans. from Poa annua germinated readily on potato-dextrose agar and produced flesh-color mucose masses of sporidia.

Entyloma spp. (E. bingenensis Zundel, in herb.) occurs abundantly on Poa vaseyochloa along the rocky warm slopes near Bingen, Washington, in late February or usually in March, as soon as growth starts and the first flowers appear (cfr. O.S.C. 8079, collected Feb. 25, 1935).

The two species, Entyloma crastophilum Sacc. and E. irregulare, are widely distributed but are especially prevalent, as mentioned, in western Oregon and Washington. They occur in the northern Great Plains and Minnesota. (?) E. spragueanum Zundel (782) from North Dakota is a fragment, needing further study. The entire group needs study.

References: Sprague (655, 656) Zundel (782).

CORTICIUM FUCIFORME (Berk.) Wakef., Pink Patch.

The fungus known generally as Corticium fuciforme but whose taxonomic status appears to be most uncertain has been reported by the writer from western Oregon (647). It develops on chewings fescue lawns during rainy periods in early spring. It also occurs on wild annual fescues, particularly Festuca myuros L. in Oregon.

References: Erwin (220, 221); Sprague (647).

RHIZOCTONIA SOLANI Kuehn,

Sharp Eyespot, Sheath Spot, Rootrot and Stem Canker (Strawbreaker), Brown Patch.

D. P. Rogers (538) places the basidial stage of R. solani in Pellicularia filamentosa (Pat.) Rogers based on Hypochnus filamentosus Pat. He gives Corticium vagum var. solani Burt ex Rolfs as a synonym, but C. vagum Berk. and Curt. is recognized as a common mold on woodland debris and is assigned to Pellicularia vaga (Berk. and Curt.) Rogers ex Linder. Because most of our material has not been connected with P. filamentosa, it is easier to employ the well-known R. solani for the purpose of this report.

R. solani is prevalent in most soils, particularly acid ones. It attacks a large number of grass hosts during wet weather in the spring. The damage usually is slight, mostly mild root necrosis or slight stunting; or sometimes elliptical light-colored eyespots with brown borders occur on the sheaths or culm bases. Under favorable conditions, the fungus may be rather destructive on cereals, causing deep-seated lesions at the ground line, which under stress of wind cause straw-breaking at this point. This condition was noted in one variety of wheat at Mandan in 1943 (Sprague, 663). An exceptionally virulent race of R. solani causes considerable damage in acid soils in the coast region in Oregon (Sprague, 623, 633, 663). The coastal fungus was erroneously assigned to an utterly different fungus by the writer in 1934 (623) although he had, before receiving mislabeled Italian material, properly listed this fungus as R. solani. The Oregon material from the coast is definitely a distinct race, both in its very strong pathological traits and the light, somewhat fluffy type of growth it makes in culture. The Oregon strain or race may be similar to Italian material, in fact the culture which the writer sent to Italy looked very much like the one which was later sent to him from northern Italy. Glynne and Ritchie (268) report similar symptoms on wheat from England and Wales. They believe that Oort's "sharp eye spot" (494) is the same fungus. They mention that Blair, working with some Canadian strains of R. solani, found some that caused stem girdling. They state that sharp eyespot does not increase with the frequency of wheat or barley in the rotations. It is as common in the first wheat crop as it is on old, arable land. The writer has long recognized that this fungus is difficult to control by crop rotation because of its ability to develop on so many hosts. In some places in Oregon cereals have been employed to clean up the soil in preparation for seeding Rhizoctonia-susceptible crops in following seasons. This may not be entirely successful. It is interesting to note that much of the detailed study in recent years on racial differences in R. solani (Houston, 341) has not considered the forms on Gramineae, the most widespread host family.

Hynes describes "purple patch" disease of cereals in Australia due to Rhizoctonia solani (344, 347) wherein small to extensive areas in a field have stunted, stiffened, slightly purplish plants with rotted

roots. Dillon Weston and Garrett (171) discuss a similar condition in England and state that the culm is not attacked as has been discussed by Glynne, Sprague, and also Oort as noted above. Samuel and Garrett (574) reported Rhizoctonia on cereals in Australia as early as 1932, and Dana (137) mentions it from western Washington in 1919.

Foëx and Rosella (235) described Sclerotium constantini Foëx and Rosella on wheat, which is the same as the stem-canker-causing strain of R. solani (Sprague, 663).

In addition to R. solani and its several races, the writer (663) has mentioned other species of this genus that need taxonomic study.

Brown patch is a case of an attack by Rhizoctonia solani in a crowded host colony wherein the fungus is able to spread from a central point, leaving dead brown areas of grass. Brown patch is important in lawns in Minnesota and in the coast region of Oregon and Washington, as well as elsewhere. In Oregon, R. solani is a common leaf rot of a large number of species of grasses during the humid winter months. It was a serious pest in spray chamber inoculation experiments at Corvallis during the years the writer was studying Septoria leaf spots of grasses. It was particularly destructive to leaves of Holcus lanatus, Agrostis alba, A. tenuis, A. palustris, and barley.

References: Bertus (44); Blair (51, 52); Dana (137); Dickinson (158); Dillon Weston and Garrett (171); Foëx and Rosella (235); Garrett (257); Glynne and Ritchie (268); Hearn (320); Hynes (344, 347); Jaarsveld (358); Matz (442); Cort (494); Peltier (508); Peyronel (515); Rogers (538); Ryker (553); Samuel and Garrett (574); Sanford (577); Shaw and Ajrekar (602); Sprague (623, 633, 645, 663); Voorhees (747).

SCLEROTIUM RHIZODES A uers

Most of the western material originally assigned to Sclerotium rhizodes (Hungerford, 343) or to S. fulvum Fr. (Young, 777) has been shown to belong to Typhula spp. (Remsberg, 532, 533). One specimen on leaves of Agrostis alba collected in the Yellowstone National Park, Wyoming, in 1941 by Fischer may be left in S. rhizodes for the present. The sclerotia were black, differing from the tawny to hazel brown sclerotia of Typhula itoana Imai and the small chestnut brown ones of T. idahoensis. In a footnote to one of her articles, Remsberg (532) states that H. H. Whetzel lists, in unpublished data, S. rhizodes under Rhizoctonia.

References: Hungerford (343); Kreitlow (391); Remsberg (532, 533); Stirrup (687); Stout (688, figs. 1-8); Young (777).

TYPHULA SPP., Snow Molds.

After many years of confusion, the snow mold of winter cereals in certain areas in Idaho and Montana have been shown to be caused by two species of Typhula, T. itoana Imai and T. idahoensis Remsberg (Remsberg, 532, 533). The fungi develop on over-wintered rosettes of wheat and certain grasses. After the snow has melted the plants are found covered with a moldy white fungous growth and the tissue is filled with small reddish brown (T. idahoensis) or dark brown (T. itoana) sclerotia. These fungi occur in Idaho (Hungerford, 343; Remsberg, 532, 533) and Montana (Young, 777 P. idahoensis), while T. itoana is recognized in the eastern United States (Wernham, 761) and Minnesota (Tervet, 703). The writer (656) noted sclerotia similar to T. idahoensis on Bromus tectorum L. and P. mollis L. on High Prairie, Washington. Snow mold material on wheat sent from Badger Mt. near Waterville, Washington, appeared to have a conidial stage that resembled that of Sclerotinia graminearum Elen. (Solkina, 618), which is a winter parasite of grain in Russia. Later reports indicate, however, that this fungus is a species of Typhula.

Remsberg noted that T. graminum Karst. is distinct from the Idaho fungi and that Sclerotium fulvum is the same as T. itoana and not related to T. graminum, as shown by differences in the structure of the sclerotia of each. T. borealis Ekstrand, which was incompletely described, is apparently the same as Remsberg's T. idahoensis.

References: Ekstrand (201, 202); Heald (315); Hungerford (343); Imai (350, 351); Müller (477); Remsberg (532, 533); Remsberg and Hungerford (534); Solkina (618); Sprague (656); Tervet (703); Volk (746); Wernham (761); Young (777) Young and Morris (778).

NON-SPORULATING BASIDIOMYCETES, Snow Mold and Root Necrosis.

Broadfoot and Cormack (73) report a snow mold-like gray fungus on over-wintered grasses, winter wheat, and alfalfa in Alberta. This fungus is very parasitic at low temperatures. They state, "This pathogen grows readily on most nutrient media at temperatures from 0° to 18° C., producing a fluffy, white sterile mycelium. Cardinal temperatures for growth are -4°, 15°, and 26° C. Sclerotia, fruiting bodies, and spores have not been found on the diseased host plants, and many attempts to produce them in culture under varying conditions of temperature, light, moisture, and nutrient media have failed. The characteristic clamp connections of the mycelium and the occasional production of small rudimentary sporophores on culture media indicate that the fungus is a basidiomycete."

Another non-fruiting basidiomycete has been isolated from crested wheatgrass (Agropyron cristatum) in Minnesota by Andrews (15), and from

A. sibiricum, A. elongatum (Host.) Beauv., Elymus canadensis, and crested wheatgrass at Mandan, North Dakota, (Sprague, 663), and on slender wheatgrass (A. trachycaulum) near Medora, North Dakota. It has been isolated also from crested wheatgrass collected near Wibaux, Montana. This fungus is a weak parasite on grasses, causing a slow root necrosis and yellowing of the seedlings. It develops during the summer and fall, as far as it is known at present. It produces a white cottony growth on potato-dextrose agar.

References: Andrews (15); Broadfoot and Cormack (73); Sprague (663).

FUNGI IMPERFECTI

Because there are so many species of Fungi Imperfecti on Gramineae and because some of them have such extensive host ranges, the entire group represents a very complex problem, considered from the viewpoint of the plant pathologist as well as the mycologist. It will be obvious to the reader that considerable fundamental work needs to be done with this heterogeneous group.

In order to bring related fungus genera together, the following grouping is followed:

- A Spores borne in pycnidia-----(1) Sphaeropsidales, p. 120
- AA Spores borne in acervuli----- (2) Melanconiales, p. 168
- AAA Spores borne otherwise----- (3) Moniliales, p. 172

1. SPHAEROPSIDALES

ASCOCHYTA SPP., Leaf Spots.

The writer and A. G. Johnson have prepared but not yet published (674) a taxonomic study of the species of Ascochyta that occur on Gramineae in the western region. It appears desirable to submit the key to these forms, which will appear in a later publication. Ascochyta is considered here as including species with hyaline spores and also those with yellow spores (Ascochyta and Ascochyta), as well as certain species previously assigned to Diplodina, which genus is not distinct from Ascochyta (see also Diedicke, 166).

The characters principally used in segregating the fungi studied in this report are included in the following key:

- A. Spores finally yellow or yellow-brown, pycnidia dark.....
Sect. Ascochyta
- B. Spores definitely fusoid, ends somewhat pointed; strongly yellow-brown, 18-27 x 5-7.7 μ . On Hordeum vulgare and Holcus lanatus Ascochyta sp.
- BB. Spores irregularly cylindrical, sometimes 2-septate, 17-26 x 6.2-7 μ , tardily yellow. On Avena.....
Ascochyta avenae Petr.
- BBB. Spores sub-cylindric, ends rounded, 14-19 x 4.5-6.2 μ in summer material (larger in type). On Agropyron and Elymus
Ascochyta agropyrina (Fairm.) Trott.
- AA. Spores hyaline or only chlorinous, rarely yellow, pycnidia brown
- B. Spores sometimes over 4.2 μ in diameter
- C. Spores very broad, as much as 8 μ diameter
- D. Pycnidia 80-120 μ , spores ovate fusoid, ends acute, 15-19 x 6-8 μ . On Panicum virgatum L.
Ascochyta sp.
- DD. Pycnidia 190-200 μ , spores sub-cylindric, ends obtuse, 16-23 x 6.5-9 μ
- E. Spores 16-19 x 5-7 μ . On Sporobolus airoides (Torr.) Torr. A. stipae Died.
- EE. Spores 17-23 x 6.5-9 μ . On Stipa sp.
A. stipae Died.
- CC. Spores 6-7 μ wide A. boutelouae Fairm.
- CCC. Spores (in our material) typically less than 6.5 μ broad
- D. Spores short cylindric or ovate, 11-18 x 3.5-5 μ . On Distichlis Diplodina graminea Sacc.
- DD. Spores cylindric or oblong, 16-23 x 4-5.3 μ . On Festuca, Phalaris, Pleuropogon, Stipa, etc.
A. graminicola var. brachypodii Trail
- DDD. Spores spindleform, ends acute, 17-21 x 4.2-5 μ , finally subcylindric. On Andropogon and Sorghastrum
A. graminicola var. brachypodii Trail

BB. Spores seldom exceeding 4 μ in diameter

C. Spores 11-20 x 2.5-4 μ , scarcely narrowly cylindrical

D. Spots pale brown or with red borders, usually scattered

E. Spores mostly 11-16 x 2.5-4 μ

F. Spores 11-16 x 2.5-4 μ , ovate or sub-cylindric A. graminicola Sacc.

FF. Spores 11-13 x 3.2-3.9 μ , short cylindric A. sorghi Sacc.

EE. Spores typically longer, cylindrical, 14-20 x 2.5-4 μ A. graminicola var. holci Sacc.

DD. Spots chocolate brown, often numerous, spores cylindrical and similar to A. graminicola var. holci or shorter, but pycnidia larger. On Bromus spp.
A. graminicola var. diedickeana Baudys and Picb.

CC. Spores 11-24 x 1.5-3.0 μ , narrowly cylindrical to sublanceolate

D. Spores sometimes sublanceolate, typically with 1-2 small oil drops adjacent to the cross wall, 12-24 x 2-2.8 μ , in pale spots on Poa spp.
one-septate phase of Septoria oudemansii Sacc.

DD. Spores typically cylindrical

E. In brown lesions, spores 11-18 x 2.4-3.1 μ
A. desmazieri Cav.

EE. In paler lesions, spores 11-18 x 1.5-3 μ in very pale brown thin-wall pycnidia
A. elymi Tehon and Daniels

EEE. In tawny spots in brown pycnidia, spores very narrow, 11-16 x 1.5-2.2 μ . On Phleum pratense L.
Ascochyta sp.

ASCOCHYTA sp. on Hordeum and Holcus, Leaf Spot and Leaf Rot.

Over-wintered leaves of barley and velvet grass (Holcus lanatus) in

Klickitat County, Washington, show pale gray emarginate spots with prominent black pycnidia. The spores are light brown, fusoid-cylindric, often pointed at both ends, $18-24 \times 5-6.8 \mu$, and have coarse granular contents. It is fairly common on High Prairie, Washington, but does not appear to be of very much importance (Sprague and Johnson, 674). It appears in the spring of the year just after the snow has left the rosettes of the over-wintering plants. What appears to be the same species occurs on Agropyron smithii in North Dakota.

ASCOCHYTA sp. on Panicum, Leaf Rot.

Our material of this fungus consists only of a fragment on frost-injured leaves of Panicum virgatum L. and on P. dichotomiflorum Michx. from North Dakota. The spores are short, thick, lemon-shape with sharp ends and slight constriction at the septum. It is probably saprophytic or very weakly parasitic (Sprague and A. G. Johnson 674).

ASCOCHYTA sp. on Phleum pratense L.

This fungus occurs on leaves of timothy in a swamp near Niawa, Minnesota. The tawny leaf spots contain brown pycnidia with very narrow spores $11-16 \times 1.5-2.2 \mu$.

References: Sprague and A. G. Johnson (674); Sprague (668).

ASCOCHYTA AGROPYRINA (Fairm.) Trott., Leaf Spots.
(Ascochyta agropyrina Fairm.)

This very weak parasite or saprophyte has yellowish-brown spores that would place it in Ascochyta, but Trotter (Saccardo, 558, V. 25) places this genus as a sub-genus in Ascochyta which appears to be logical. A. agropyrina is widely scattered and common but never occurs in great abundance in any material. It has been found on Agropyron bakeri E. Nels. in New Mexico (Fairman, 222); on A. caninum (L.) Beauv. (from Russian seed), A. cristatum, A. michnoi Roshev, A. semicostatum (Steud.) Nees, A. subsecundum (Lk.) Hitchc., A. trachycaulum, A. trichophorum (Lk.) Richt., Elymus canadensis, and E. virginicus in North Dakota; A. cristatum and A. smithii in Montana; and on A. semicostatum from Pullman, Washington. The material from Pullman has fuscous spots with red to vinaceous borders but the spores are pale yellow and with the stout cylindrical shape of A. agropyrina.

References. Fairman (222); Saccardo (558) v.25; Sprague and A. G. Johnson (674); Trotter (713).

ASCOCHYTA BOUTELOUAE Fairm., Leaf Mold.

Scarcely parasitic, a relatively common fungus on the leaves of blue grama in central North Dakota and in New Mexico. The spores in this species are thicker than those of A. graminicola, somewhat near A. graminicola brachypodii but the black, often sub-superficial (saprophytic), tardily ostiolate pycnidia, together with a hint of smokiness in the spores, distinguish A. boutelouae from A. graminicola var. brachypodii (Sprague and A. G. Johnson, 674). A. boutelouae is scarcely distinguishable morphologically from Diplodina graminea Sacc. (Sprague and A. G. Johnson, 674). Fairman (222) described this from New Mexico.

ASCOCHYTA DESMAZIERI Cav., Leaf Blotch.
(Septoria lolii West. and Diplodina lolii Zimm.)

The fungus causes a brown or brownish purple spot on the leaves of Lolium perenne L. and L. multiflorum in Oregon, Washington, and California. It is sometimes destructive. The spores are truly cylindrical and readily distinguished from those of the A. graminicola complex. Its symptoms are readily distinguished from the speckled gray blotch caused by Septoria tritici var. lolicola Sprague and A. G. Johnson (665), but can be confused with those of S. loligena Sprague. It should, in fact, be compared critically with this fungus.

References: Cavara (99); Sprague (655; 665); Sprague and A. G. Johnson (674); Zimmerman (781, (p. 101).

ASCOCHYTA ELYMI Tehon and Daniels, Leaf Spot.

Tehon and Daniels (701) distinguish this species from A. graminicola by its fragile pycnidia as contrasted with the brown ones of A. graminicola. The writer has found A. elymi on seedlings of Elymus canadensis near Solen, North Dakota. Traces of it occur on Elymus spp. in the Rocky Mountains and possibly on Agropyron repens near Eugene, Oregon.

ASCOCHYTA GRAMINICOLA Sacc. and Varieties, Leaf Spots.

Davis' summary of the common parasite, A. graminicola, is generally accepted (Davis, 140). While the fungus usually is limited to a few red-bordered brown or fawn-colored spots on the lower leaves or leaf tips, the fungus is sometimes aggressively parasitic. Such material has been seen on Hierochloë odorata (L.) Beauv. in North Dakota and South Dakota. In the far west it is common or more often scattered on a number of grasses during the rainy season in winter and spring, while east of the Rockies it develops during rainy periods in the summer

(Sprague, 655, 656).

Some 14 varieties have been described for A. graminicola, but these include some "varieties" which are readily referable to Darluca filum (Biv.) Cast. In fact, it is conservative to say that a majority of the specimens in herbaria that are assigned to A. graminicola are D. filum or sometimes immature stages of Hendersonia, Stagonospora, or Sentoria. Among the recognizable varieties of A. graminicola, var. holci is considered to have somewhat larger spores than true A. graminicola and much of our western material would fall under this variety if it is recognized. A more obvious form, var. brachypodii Trail (708), occurs on Phalaris arundinacea L. at Mandan, North Dakota; on Lolium perenne L. from Troutdale, Oregon, on Stipa comata Trin. and Rupr. in Montana, on Andropogon furcatus Muhl. at Mandan, North Dakota, and on Pleuropogon refractus (A. Gray) Benth. in Oregon (Sprague and Johnson 674). These all tend to have fusoid spores, 15-17 x 5 μ , while the species proper and var. holci have narrower spores that are nearly cylindrical, 10-20 x 2.5-4 μ .

Var. diedickeana Baudyš and Picb. (26) produces brown-bordered lesions with paler centers, and in severe cases dark brown or nearly black general scalding of affected green parts. It occurs on Bromus carinatus in Oregon, on leaves of B. inermis in Minnesota, and on roots of B. inermis in North Dakota (see also Broadfoot, 72). It occurs on Hordeum nodosum L. at Hunter, Alaska. In addition to its distinct symptoms and parasitic tendencies, this variety is recognizable to some extent by its cylindrical to fusiform spores, 11-17 x 2.5-3.1 μ , which are narrower than those of var. holci.

Ascochyta graminicola is indeed a complex, and further life history studies may show that some of these morphologically similar forms are growth stages of entirely different fungi.

References: Baudyš and Picbauer (26); Broadfoot (72); Davis (140); Sprague (655,656); Sprague and A. G. Johnson (674); Trail (708).

ASCOCHYTA SORGHII Sacc., Leaf Spot.

What can be called this fungus was collected on sorghum at Brookings, South Dakota, in 1912 by C. R. Ball and determined by the writer 30 years later. The spores are small and cylindrical, 11-12.5 x 3.2-3.9 μ .

ASCOCHYTA STIPAE Died., Leaf Spot.

A prominent maroon to vinaceous leaf spot occurs on Stipa sp. on Mt. Shasta, California. The broad spores are 17.5-24 x 6.5-9 μ . Another collection on Sporobolus airoides from Pullman, Washington, has smaller spores but they are similar in shape to those on Stipa (Sprague and

Johnson, 674).

ASCOCHYTELLA AVENAE Petr., Leaf Spot.

This fungus which was placed in still another genus (Ascochyella) by Petrak (510, p. 107) belongs in the yellow-spored section or subgenus Ascochyula of Ascochyta. It was found once in late winter on Avena byzantina C. Koch (red oats) near Troutdale, Oregon, and on A. fatua (wild oats) at Pullman, Washington (Sprague and A. G. Johnson, 674).

CONIOTHYRIUM PSAMMAE Oud., Leaf Spot.

Coniothyrium psammae occurs scatteringly along the sea coast in Oregon on Calamagrostis nutkaensis (Presl) Steud. The spots are large, elliptical, light buff with prominent vinaceous borders. The pycnidia are few and contain brown elliptical spores, 10-13 x 3-4 μ .

Reference: Sprague (652).

DILOPHOSPORA ALOPECURI (Fr.) Fr., Twist Disease.

The leaves and more often the heads are attacked. The fungus causes a black, charred-appearing crust accompanied by deformation and twisting of the head and adjacent leaves. Affected heads are frequently killed in the boot. The fungus develops abundant black masses of pycnidia, which produce spores with characteristic bristle-like appendages at both ends. The spores are hyaline, 8-15 x 1.6-2.5 μ ; and the appendages are 5 to 7 μ , less often 10 μ long and 0.5 μ thick at the base, 3-4 branched with some branches forked or sometimes tri-furcate.

Atanasoff (18) has detailed a study wherein he suggests that the spores are carried by Tylenchus tritici (Steinb.) Bast. to the growing points of the plant. This appears to be discounted by many writers. Certainly in the Pacific Northwest the twist disease does very well without the assistance of nematodes in most cases.

Some workers, including Rainio, claim that Nastigosporium album Riess and D. alopecuri are part of the same life cycle. This is utterly incorrect (Sampson and Western, 57C; Sprague, 64C). No proof of connection with Dilophia graminis Sacc. has been obtained in this country either.

Dilophospora alopecuri is common in Oregon and Washington on Holcus lanatus and Sitanion jubatum J. G. Sm. and it occurs on Agropyron inerme in eastern Washington; on A. smithii in Idaho; on A. spicatum in Montana; on A. trachycaulum in Yellowstone Park, Wyoming, and Roosevelt Park,

North Dakota, on Melica bulbosa Geyer in Yellowstone Park; and on Poa secunda at Logan, Utah. Most of the collections have been made recently by Fischer and the writer but the fungus has been present in the West for an indefinite period. Jackson collected it in Oregon in April, 1915 (O.S.C. 10,530). Bessey (45) reported it from Wisconsin in 1906.

Bevilacqua (46) recommends that the wheat be cut high, the stubble burned, and the seed treated.

Reference: Atanasoff (18); Bessey (45); Bevilacqua (46); Grove (296); Kotte (389); Raino (520); Sampson and Western (570); Schaffnit and Wieben (585); Sprague, 640).

DIPLODIA ZEAE (Schw.) Lév., Dry Rot of Corn.

This dry rot of corn ears has been well discussed and illustrated by several workers in Nebraska (Heald et al., 318), Iowa (Durrell, 189), and the eastern States. The fungus also causes sheath and leaf spots and stalk rots, and the fungus in the seed causes seedling blight. The disease is important in Iowa, eastern Nebraska, and Minnesota but disappears in the West although it probably occurs in favored, more humid areas.

References: Bijl (47); Burrill and Barrett (84); Clayton (116); Durrell (189, 190); Eddins and Voorhees (193); Heald (312); Heald et al. (318); Hoadley (332); Holbert et al. (336); Holbert, Hoppe, and Smith (337); Hoppe (338); Kent (381); Kinsel (382); Koehler and Holbert (388); Larsh (398); McNew (451); Raleigh (521); Semeniuk (590-593); Semeniuk et al. (596, 597); Ullstrup (728); Valteau et al. (735); Wilson (767).

DIPLODINA GRAMINEA Sacc., Leaf Mold.

Material collected by the writer at Davis Junction, California, on Cynodon dactylon (L.) Pers. shows sub-superficial pycnidia scattered among Helminthosporium cynodontis Merig. The spores of the Ascochyta are capsular, one-septate, clear hyaline but with a hint of smokiness in mass, 11-18 x 3.9-5 μ (Sprague and Johnson, 674). Diehl (169) reported the fungus from South Carolina in 1935, the first report for the western hemisphere. The fungus is probably common. Nothing is known as to its pathogenicity. See also A. cynodontis Unam. (730).

HENDERSONIA spp., Molds.

Some of the common invaders of dead or dying tissue are species of Hendersonia Berk. em. Sacc. Therefore, in routine examination of diseased plants, workers are confronted with scattered pycnidia belonging

to species of this genus. The brown spores of Hendersonia are broadened, not filiform nor elongate-vermiform as in some species of Phaeoseptoria (Sprague, 659). Since this group does not contain any species that are definitely parasitic it is not considered necessary to discuss them in detail. In addition the taxonomy of this group is in great confusion, at least from the viewpoint of literature available on the western forms. The writer has an uncompleted manuscript dealing with about 16 species in the Western States (667). They are particularly common on overwintering grasses in the Pacific Northwest and Rocky Mountain areas and on grass in the late fall in the Northern Great Plains.

Among the common species of Hendersonia are H. simplex Schroet., H. culmicola Sacc., H. crastophila Sacc., H. calamovilfae Petr., H. stipae-pennatae Fautr., and H. calospora Fautr. Several less common or undescribed species also occur.

The following is a simplified key to these species:

- A. Spores when mature typically 7-septate.
 - B. Pycnidia very large, 180-300 μ in diameter
 - C. Spores falcate or falcate-fusiform, 28-40 x 3.8-4.4 μ
..... Wojnowicia graminis
 - CC. Spores fusiform, vermiform, coarse, somewhat constricted at the septa, 35-53 x 3.5-5 μ H. calospora Fautr.
 - BB. Pycnidia smaller, 80-220 μ in diameter
 - C. Spores regularly cylindrical, ends blunt, cross walls closely adjacent, 20-29 x 4-6 μ
H. stipae-pennatae Fautr.
 - CC. Spores more or less fusiform, sub-cylindrical; ends or at least one end somewhat to definitely tapered; septa not particularly closely adjacent
 - D. Spores as much as 7 μ wide ... H. calamovilfae Petr.
 - DD. Spores mostly 3-5 μ wide
 - E. Spores brown, fusiform or falcate, flattened on one side, constricted at the septa
 - F. Spores fusiform, pycnidia rarely with hairs on the ostiolar end
H. crastophila Sacc.

- FF. Spores fusiform-falcate or falcate, pycnidia with scattered hairs on the ostiolar end
Wojnowicia graminis (McAlp.) Sacc. and D. Sacc.
 (Confusable with H. crastophila when immature)
- EE. Spores lighter colored, small paired regularly placed prominent guttulae, often constricted at septa, sub-cylindrical, ends conical, not stiffly flattened on one end Hendersonia sp.
- EEE. Spores shorter than in EE, rarely constricted at septa, more often 5- than 7-septate
H. culmicola Sacc.
- AA. Spores when mature typically 5-septate or less (see also H. culmicola above)
- B. Spores sometimes more than 3-septate, cylindrical
- C. Spores 3-, 5-, to 7-septate, 30-38 x 3.1-4.2 μ
H. culmicola Sacc.
- CC. Spores 3- to 4-septate, 20-27 x 3.2-3.4 μ
H. culmicola var. intermedia Sacc.
- BB. Spores not more than 3-septate
- C. Spores with stout episporous, 6.2-7.8 μ wide
Hendersonia sp.
- CC. Spores with thin walls, less than 5 μ wide
- D. Spores 28-48 x 3.8-4.3 μ Hendersonia sp.
- DD. Spores less than 30 μ long or approximately that
- E. Spores short-cylindrical, some at least 4 μ wide
- F. Spores 22-29 x 2.8-4.6 μ
H. agropyri-repentis Oud.
- FF. Spores 15-20 x 4.3-4.8 μ
Hendersonia sp. on Danthonia intermedia Vasey
- EE. Spores sub-cylindric to fusiform, not over 3-5 μ wide
- F. Spores often 1-septate, finally 3-septate, sub-lanceolate at first, 17-22 x 2-2.6 μ
H. culmicola var. minor Sacc.

FF. Spores mostly 3-septate, sub-bacillar, 14-26
x 2.3-3.1 μ H. simplex. Schroet.

The closely related species Wojnowicia graminis is included in the key for comparison.

References: Grove (297); Saccardo (558, v. 3, 10, 11, 12, 13, 14, 16, 18, 22, 25); Sprague (667).

MACROPHOMA PHLEI Tehon and Stout, Leaf Mold.

That was probably saprophytic material of M. phlei was seen at Mandan, North Dakota, on Elymus triticoides Buckl. on October 1, 1943 (B.P.I. 80,893). Spots were diffuse straw-color, or more often absent. The pycnidia were black, depressed, globose, 120-135 x 120-204 μ bearing elliptical to elongate ovate spores, which were slightly larger at one end, with pure white opaque contents, 28-39 x 8.5-11.1 μ , mean size 32 x 9.4 μ . This compares with 18-26 x 6.4-7.7 μ for the type of M. phlei, which indicates that the Mandan material has somewhat larger spores but is probably the same species. M. phlei has also been seen on Agrostis palustris from Waldport, Oregon, with spores 24-31 x 9.5-11 μ .

References: Sprague (666); Tehon and Stout (702, p. 188).

MACROPHOMINA PHASEOLI (Maubl.) Ashby, Charcoal Rot.
(Sclerotium bataticola Taub.)

Charcoal rot develops in the stalks of corn and sorghum in late season, causing general decay and weakness with the resultant breaking over of diseased plants. The numerous small black bodies form in large numbers. These are said to be prosenchymatous-filled pycnidia or sclerotium-like bodies of the Sphaeropsidaceous Macrophomina phaseoli. M. phaseoli occurs on corn in California (Mackie, 426) and recently has been given considerable publicity in Nebraska (Livingston, 409, 410), Iowa (Semeniuk, 594) and adjacent states. It has not been isolated, as yet, from North Dakota material, although a sclerotial form superficially resembling it has been obtained from roots of several grass crops at Mandan.

References: Ashby (17); Dunlap (188); Haigh (304); Hoffmaster et al. (334); Livingston (409, 410); Schwarze (587, p. 148-149); Semeniuk (594, 595); Taubenhaus (699); Tehon and Boewe (700); Uppal et al. (733).

PHAEOSCEPTRIA Spp., Leaf Molds and Secondary Invaders.

Species of this genus sometimes are encountered on dead or dying leaf and culm tissue of grasses. Phaeosceptoria belongs to the Sclerosporeae,

has coarse black pycnidia and light brown multiseptate spores. A key to all known forms of this genus on grasses and a discussion of their taxonomy has recently appeared (Sprague, 659) and it seems unnecessary to repeat this information here as the group is of little economic importance.

PHOMA SP.

An undetermined species of Phoma was reported as important on Agropyron smithii Rydb. in Utah by O'Gara (493).

PHOMA TERRESTRIS Hansen, Pink Root.

The same fungus that attacks the roots of onions, causing pink root, is able to attack the roots of cereals and grasses (Kreutzer, 392). Johann (365) and Sprague (662) verified this and also isolated the fungus from Gramineae. It occurs in many species of grasses and cereals in the Northern Great Plains but is not very parasitic.

P. terrestris causes a pink to carmine-lake coloration of the roots. It sometimes causes mild seed rot and slight rootrot. In culture, it produces a slow-growing mounded gray colony with pink to magenta tints in the substrate. It does not fruit readily on potato-dextrose agar.

References: Carvajal (96); Johann (365); Kreutzer (392); O'Gara (493); Sprague (662).

PHYLLOSTICTA

Five species of Phyllosticta, not known to be phases of more complex fungi, may be distinguished by the following key:

A. Pycnidia in spots with pale centers

B. Spots small, white, spores 11-15 x 2.5-4.5 μ
P. rogleri Sprague

BB. Spots irregular, pale to discolored, with red to purple borders, spores 4-7 x 2-3 μ P. sorghina Sacc.

AA. Pycnidia in gray or vague spots

B. Spots gray, elongate, spores bacillar, 2.3-4.6 x 1-1.4 μ
P. owensii Sprague

BB. Spots vague on dead leaves, spores bacillar, 5-9 x 1-1.6 μ
 P. anthoxella Sprague

BBB. Spots cream color to pale yellow, emarginate, spores oval to elliptic, 6.5-8 x 3-4 μP. avenophila Tehon and Daniels⁵

PHYLLOSTICTA ANTHOXELLA Sprague, Leaf Spot.

The small black pycnidia are scattered over moldy dry leaves of An-thoxanthum odoratum L. collected at Triangle Lake, Oregon (Sprague, 652, fig. 1, e). Little is known about this fungus, which appears to be of slight importance.

PHYLLOSTICTA OWENSII Sprague, Gray Leaf Streak.

Also a minor parasite or mold on living, dead or salt-spray injured leaves of Dactylis glomerata in the vicinity of Waldport, Oregon (Sprague, 652). It is an associate of Scolecotrichum graminis Fckl.

PHYLLOSTICTA ROGLERI Sprague, White Leaf Spot.

Known only on Digitaria sanguinalis (L.) Scop. from Albia, Iowa. (Sprague, 652, fig. 1, f).

PHYLLOSTICTA SORGHINA Sacc., Eyespot.
 (Phoma insidiosa Tassi)

The lesions are irregular to sub-circular, pale straw to isabelline color spots with narrow purple to red borders, 3-8 mm in diameter. They develop during the summer rainy season and are associated with various bacterial and possibly non-parasitic spots on Setaria viridis in North Dakota (Sprague, 652, fig. 1, g).

SELENOPHOMA

The genus Selenophoma is characterized by pycnidia with non-septate, sickle shape (falcate), hyaline spores (Maire, 428).

Key to the species:

A. Spores acute, boomerang or arcuate

⁵

This fungus occurs in Illinois and Mississippi but is likely to be found in the area covered in this report.

- B. Spores arcuate, typically 10-16 x 1.1-1.8 μ
S. everhartii (Sacc. and Syd.) Sprague and A. G. Johnson
- BB. Spores lunate to boomerang, typically 18-30 x 2-4 μ
S. donacis (Pass.) Sprague and A. G. Johnson
- BBB. Spores lunate to sub-arcuate, typically 12-24 x 1.5-2.7 μ ...
S. donacis var. stomaticola (Bauml.) Sprague and A.G. Johnson
- BBBB. Spores lunate, 17-25 x 2-3.2 μ . On Bromus
S. bromigena (Sacc.) Sprague and A. G. Johnson
- AA. Spores obtuse, sub-reniform-falcate
S. obtusa Sprague and A. G. Johnson

SELENOPHOMA BROMIGENA (Sacc.) Sprague and A. G. Johnson

The common host, Bromus inermis, is severely damaged by this fungus which forms gray, irregular to sub-circular spots on the leaves in early spring. It is especially prevalent in the Red River valley and in Minnesota but occurs westward to Pullman, Washington. It is not uncommon on the prairies in Idaho but has not yet been noted in Oregon or California. It appears to be rather largely confined to B. inermis growing in the plains countries. It occurs in Hungary (Krypt. Exs. Palatino, No. 1465) but has not been given much recognition outside the United States.

Selenophoma bromigena differs from S. donacis somewhat vaguely in that the spores of S. bromigena are regularly lunate, sharp-pointed, rarely boomerang shape. An undated packet of a fungus called Phlyctaena bromi by Clements (Crypt. F. Coloradensium) on Bromus ciliatus L. appears to be a saprophytic phase of S. bromigena on an unusual host. Allison (9) found that S. bromigena was confined to B. inermis. Field observations show that there are very few exceptions to this. S. bromigena has been collected in trace quantities on B. carinatus at Glenrock, Wyoming, Pullman, Washington, and Mandan, North Dakota.

References: Allison (3-10); Darley (138); Saccardo (557); Sprague and A. G. Johnson (673, 675, 676).

SELENOPHOMA DONACIS (Pass.) Sprague and A. G. Johnson.
(Septoria oxyspora Penz. and Sacc.)

This fungus is commonly known as Septoria oxyspora, less often as Septoria donacis Pass., an earlier name. While it was described on Arundo donax L., a giant reed, it occurs also on Phalaris arundinacea in Washington and North Dakota; rarely on Phragmites communis Trin. in

North Dakota; (Septoria curva Karst.) on Elymus condensatus Presl in Washington, North Dakota, and Oregon; on E. flavescens Scribn. and Sm. in Washington; on Panicum virgatum in the Great Plains; and on Stipa columbiana Macoun, S. comata, S. richardsoni Lk., S. spartea Trin., and S. viridula in the Great Plains and westward.

Selenophoma donacis on Arundo, Phalaris, and Elymus has spores averaging around $20-21 \times 3 \mu$ and more or less boomerang shape. Spores of S. donacis on Panicum are the same length as those on Arundo but are recognizable because of their narrower shape. They are $21.5 \times 2.2 \mu$, mean size, in one collection. Those on Stipa are stubbier, $18.3 \times 2.8 \mu$, average, and the material on Phragmites has thicker, short spores, $16-18 \times 3-4 \mu$ but typically boomerang shape.

Most of the lesions are straw color with tawny, fuscous, or lilac tinted borders, and contain small obscure pycnidia. Very often the spots fade, leaving the pycnidia in unspotted tissue on the culms, sheaths or leaves.

S. donacis has not been studied so thoroughly as its variety stomaticola, and therefore cannot be segregated into races. On cultural characters, it is indicated that there is a distinct race on Elymus flavescens from the sandhills near Hanford, Washington; one on Arundo from California; and probably the same race on Phalaris in the northern United States; another on Panicum; and probably two on Stipa. The culture from E. flavescens from Hanford, Washington, for instance, produces an ivory color growth quite different from the congo pink to testaceous rose mycelia from California material of Arundo donax. However, a race with narrower spores isolated from Panicum virgatum from material collected at Coburn, North Dakota, was very similar to the culture from Arundo (Sprague and A. G. Johnson, 675). Until there is more evidence to the contrary, all of these groups are placed under one species because the morphological differences between them are not fundamental.

References: Grove (296); Passerini (504); Penzig (509); Sprague and A. G. Johnson (673, 675).

SELENOPHOMA DONACIS var. STOMATICOLA (Bauml.) Sprague and A. G. Johnson. (Phyllosticta stomaticola Bauml.)

The complex that we place under this variety has, in general, spores that are $3-5 \mu$ shorter than those of the species and are distinctly narrower. The problem is not so simple as that, however, because spores in some groups, such as that on Agropyron, approach or equal in size those of S. donacis on Panicum, while in others, as that on Sporobolus, they are close to the small-spored S. everhartii (Sacc. and Syd.) Sprague and A. G. Johnson. One group on Poa (cfr. Septoria nebulosa Rostr.) has short spores with sharper tips, while material on

Danthonia has evenly lunate spores with softly tipped ends. It is, nevertheless, difficult to segregate these groups into species on any satisfactory basis. Because of our present incomplete knowledge, it appears more practical to assign all of them to the earliest available name as a variety under S. donacis.

The symptoms induced by this variety are of two kinds, light color spots with pale, tawny, red or lilac borders, or no spots at all with the minute pycnidia scattered over straw-color areas. On Calamovilfa, in the Dakotas, the spots are particularly pale, while on Danthonia californica Bol. in Oregon, the spots are a "frog eye" type with a prominent red or maroon border. The lesions on rye (Secale cereale L.) are fuscous to straw color without prominent borders while the pycnidia are more prominent against the light background than on most other hosts.

Pure culture studies, plus examination of a large number of collections on western grasses, have shown that var. stomaticola is divisible into 11 major groups or major races as follows:

Race 1, on Danthonia californica, is closest to Septoria lunata Grove (296) with the regularly lunate spores with soft-pointed tips. It produces a light gray crustose colony with scant mycelium and opalescent grayish white spore droplets. The medium was tinted rosy-vinaceous. The fungus is restricted to this species of host.

Race 2, on Dactylis glomerata and Koeleria cristata from Pullman, Washington. The isolates from these two hosts are identical, each produces a lavender-tinted colony.

Race 3, on Festuca idahoensis from Lyle, Washington, has the appearance of race 2 in pure culture but the spores are narrower and resemble those of the next race.

Race 4, on Agropyron and Elymus spp. has spores that average slightly narrower and longer than those on most other hosts of this variety and they form a definite group.

Race 5, on Poa spp. The native species of Poa from the middle Columbia River Valley harbor a race that is similar in pure culture to race 2, but not identical with it, and with spores that are thicker in their middle portions and sharper at their ends.

Race 6, on Poa spp. from Pullman, Washington, and the Big Horn Mountains, Wyoming, tends to form cream color, soon tan, brown, creosote, and finally black colonies. These seem to represent a distinct group.

Race 7, on Poa pratensis from Quad Creek, Montana, is akin to

race 6 but the burnt-orange, later brown colonies plus somewhat boomerang shape spores indicate that this is at least a localized variant. Races 5 to 7 are similar to Rhabdospora groenlandica Lind (407), particularly race 7 from the sub-alpine Rockies.

Race 8, on Arrhenatherum elatius (type host of Phyllosticta stomaticola) from Bozeman, Montana, differs from race 2 in lesser tendency to form lavender color in agar, and in the production of dirty tan, mucose conidial masses in early stages of growth. It is close to race 6.

Race 9, on Phleum pratense, is distinct from race 1 in host range but its relation to the others is undetermined. This fungus was described by Lind (406) as Septoria culmifida Lind. However, it is morphologically indistinguishable from most of the S. donacis stomaticola complex.

Race 10, on Sporobolus asper (Michx.) Kunth, has not been cultured and little is known of its relationship with the other races. Since it has some boomerang-shape spores it somewhat resembles S. donacis but the spores are narrower.

Race 11, on rye (Secale cereale), also needs study as soon as viable material is available. The pycnidia are prominent but not large, 95-111 x 72-85 μ , and the spores are 12-18 x 2-3.5 μ , mean size 14.5 x 2.4 μ , in one collection, 16 x 2.7 μ in another. This fungus has been found in an area north of Jamestown, North Dakota, and also in Minnesota. The symptoms may be confused with those caused by bacterial leaf blight and therefore the disease on rye may be more widespread than is known at the present. It has no relation to Septoria falcispora Demidova (148) (non Bubák).

Selenophoma donacis var. stomaticola has the following host range in the various States: Agropyron albicans Scribn. and Sm. (N. Dak.); A. dasystachyum (Hook.) Scribn. (Oreg.); A. repens (Wash., N. Dak.); A. riparium Scribn. and Sm. (N. Dak.); A. smithii (N. Dak., S. Dak., Mont., Wyo.); A. spicatum (N. Dak., Wyo., Wash.); A. subsecundum (Mont.); A. trachycaulum (Mont., N. Dak.); Arrhenatherum elatius (Mont.); Calamovilfa longifolia (Hook.) Scribn. (N. Dak., S. Dak.); Dactylis glomerata (Oreg., Idaho, Wash.); Danthonia californica (Oreg.); Elymus canadensis (N. Dak.); E. giganteus Vahl (Wash.); E. glaucus (Wash., Oreg.); E. junceus Fisch. (N. Dak.); E. virginicus (N. Dak.); Festuca idahoensis (Wash.); F. ovina L. (Wash.); Hordeum brevisubulatum (Trin.) Lk. (N. Dak.); H. vulgare (Oreg.); Koeleria cristata (N. Dak., Wyo., S. Dak., Mont., Wash., Idaho); Melica harfordii Boland (Calif.); Oryzopsis hymenoides (Roem. and Schult.) Ricker (N. Dak., Mont., Calif., Oreg.); Phleum pratense (Alaska, N. Dak.); Poa alpina L. (Mont.); P. ampla Merr. (Wash., Mont., Oreg.); P. arctica R. Br. (Mont.); P. compressa

(N. Dak.); P. epilis Scribn. (Wyo.); P. interior Rydb. (Mont.); P. juncifolia (Oreg., Wash.); P. nervosa (Hock.) Vasey (Mont., Wyo.); P. pratensis (Mont., N. Dak., Wyo., Wash.); P. secunda (N. Dak., Wyo., Mont.); Puccinellia nuttalliana (Schult.) Hitchc. (N. Dak.); Secale cereale (N. Dak., Minn.); Sitanion hystrix (Nutt.) J. G. Sm. (Wash., Oreg.); Sporobolus asper (N. Dak., Mont., Iowa); S. cryptandrus (N. Dak.).

Some of these have not been given racial classification as little is known about them. The entire problem is a tremendous one, but since members of the group, with a few exceptions, are minor parasites, the need for study is not pressing. The group is almost unknown to plant pathologists although extremely common. Very often the small spherical pycnidia (frequently only 40 μ in diameter) contain but few spores, which are not easily crushed out of the pycnidia. These scattered spores when seen floating free in the mount may be easily mistaken for spores of Colletotrichum.

References: Bäumler (27); Bisby et al. (49); Bubák (77); Demidova (148); Grove (296); Lind (406, 407); Lobik (411); Petrak (511); Rostrup (545); Sprague and A. G. Johnson (673, 674, 676).

SELENOPHOMA EVERHARTII (Sacc. and Syd.) Sprague and A. G. Johnson.
(Septoria everhartii Sacc. and Syd.)

The pycnosporos of this species are smaller than those of S. donacis var. stomaticola and in most of the material are more arcuate, being beautifully crescent-shape with sharp points. There is reason for assigning this species to the S. donacis complex because all sizes of spores are found on the various hosts from the large ones of S. donacis on Arundo to the small ones on Agrostis. In final analysis, it has been decided to separate S. everhartii from var. stomaticola, although most of our herbarium material was originally assigned to S. donacis. Sprague and A. G. Johnson (675) concluded that the S. donacis complex was sufficiently confusing without adding the small-spored group to it (676). The spores are 11-15 x 1.1-1.3 μ , sometimes 10-20 x 1-2.4 μ .

The symptoms caused by S. everhartii are similar to those produced by S. donacis. On Calamagrostis canadensis (Michx.) Beauv. the lesions are elongate, white with vinaceous to vague borders. Much of the material on Agrostis and on alpine material in semi-arid regions in general is obscure, with minute pycnidia on stramineous, dead plants. On Aristida and on Bouteloua curtipendula (Michx.) Torr. gray to tawny or paler, eye-spot lesions with vinaceous borders occur, often intermingled with saprophytic material from the preceding year.

In pure cultures on potato dextrose agar, S. everhartii shows as much racial variation as S. donacis var. stomaticola. Thus, an isolate from Agrostis scabra Willd. (Ft. Totten, North Dakota) has gray and rose

colored leathery mycelia with lavender color in the substratum. From Trisetum spicatum (L.) Richt. (Beartooth Pass, Montana), the mycelium is gray and creosote color over pale lavender; while on agar an isolate from Bouteloua curtipendula (western, North Dakota) produces a crustose off-white mycelium with no substratum color, very different in appearance from the others. A culture from Deschampsia atropurpurea (Wahl.) Scheele (also at Beartooth Pass, in Wyoming) is similar to that from T. spicatum but with gray and isabelline mycelium. On D. danthonioides (Trin.) Munro (Pullman, Washington), the mycelium was burnt-orange mixed with pink and black and with an orange substratum color, quite different from any of the others. A recent isolate from Muhlenbergia racemosa (Michx.) B. S. P. (Mandan, North Dakota) formed a black carbonaceous growth. This isolate when incubated on sterilized brome stems over winter produced small black perithecia (80-90 μ) belonging to the Mycosphaerellaceae. A few contained fascicled asci containing 8 ellipsoidal ascospores 7-9 x 2-2.6 μ . Unfortunately, it could not be determined whether these were slightly immature and only bi-guttulate (Ascospora) or somewhat more immature and ultimately one-septate (Mycosphaerella); nor could the spores be germinated.

Selenophoma everhartii occurs on the following hosts: Agrostis diegoensis Vasey (Calif.); A. scabra (N. Dak.); Aristida longiseta Steud. (N. Dak.); A. oligantha Michx. (Iowa); Bouteloua curtipendula (N. Dak.); Calamagrostis canadensis (this is the type host from Wisconsin of Septoria calamagrostidis Ell. and Ev. (non (Lib.) Sacc.) from which Septoria everhartii Sacc. and Sydow was derived); C. montanensis Scribn. (N. Dak.); C. scribneri Beal (Wyo.); Deschampsia atropurpurea (Wyo., Mont.); D. caespitosa (L.) Beauv. (Mont. and Wyo.); D. danthonioides (Wash.); D. elongata (Hook.) Munro (E. Wash.); Festuca kingii (S. Wats.) Cassidy (Mont.); F. ovina (Mont. and Wash.); F. ovina var. brachyphylla (Schult.) Piper (Wyo. Mont.); F. rubra (Mont.); Muhlenbergia cuspidata (Torr.) Rydb. (N. Dak.); M. racemosa (N. Dak.); Sphenopholis obtusata (Michx.) Scribn. (N. Dak.); Trisetum spicatum (Mont., Wyo.).

Some of the specimens that have been placed in Selenophoma everhartii may be immature stages of S. donacis var. stomaticola, in fact one reason for the long hesitation in assigning these to a separate species was the fact that fungi on Agrostis and Calamagrostis tend to be smaller on that host than the same species on some other hosts.

Selenophoma everhartii is the most common species on alpine and sub-alpine plants and is abundant on the dry culms of grasses in the plains country in late summer.

References: Ellis and Everhart (214); Rostrup (545); Saccardo (558); Sprague and A. G. Johnson (675, 676).

SELENOPHOMA OBTUSA Sprague and A. G. Johnson.

This fungus is a variant of the S. donacis complex but in most collections it is distinguishable from that group by its short, thick, almost reniform spores averaging 13-17 x 2.5-4 μ . In pure culture it is distinguishable from most of the others by prominent black pycnidial aggregates scattered in salt and pepper pattern over a light isabelline to very pale flesh background. While the spore exudate is pale lavender, the strong rosy tints of S. donacis and the vinaceous to lavender ones of var. stomaticola are less prominent. In its early stages of growth, an isolate from Sitanion hystrix, Mt. Shasta, California, (W. B. Cooke) produced a wet-leathery, faintly flesh color to isabelline colony. A culture isolated from Agropyron inerme collected at Washtucna, Washington (G. W. Fischer) showed the same type of growth in early stages, but later, mucose, tan, and still later, brown masses of spores formed over the earlier growth. This case was similar in appearance to race 6 of var. stomaticola and this collection may be misassigned. While typical material on Sitanion hystrix appears to be very distinct, other material on Elymus glaucus and on S. hystrix is variable, some close to var. stomaticola, others to S. obtusa. Certainly our S. obtusa is not completely separable from the whole complex.

The host range of Selenophoma obtusa is as follows: Agropyron inerme (Wash.); Elymus condensatus (Wash.); E. glaucus (Wash., Calif.); Poa arida Vasey (N. Dak.); Sitanion hanseni (Scribn.) J. G. Sm. (Calif.); S. hystrix (type locality is Mt. Shasta, Calif.; Mont.); Stipa richardsoni (Mont., Wyo.); Stipa sp. (Calif.). Selenophoma obtusa occurs throughout the range of S. donacis from Mandan to Mt. Shasta but is typically a fungus of the high mountains and desert. It has been collected for instance at White Bluffs, Washington, in a region having 3 inches of average annual precipitation.

References: Sprague and A. G. Johnson (675, 676).

SEPTORIA

Most of the species included here have just been the subject of a detailed study by the writer (Sprague, 665) and therefore it does not appear necessary to discuss them in detail. It is, however, believed desirable to give the following key to the species:

A. Pycnospores 0- to 2-septate, less often 3-septate

B. Pycnidia prominent, as much as 180-250 μ diameter

C. Spores filiform, 36-50 x 1.4-1.6 μ
Septoria poliomela Syd.

- CC. Spores obclavate filiform, 33-65 x 1.2-2.5 μ
S. bromi Sacc.
- CCC. Spores obclavate, 28-50 x 2.5-4.5 μ
S. infuscans (Ell. and Ev.) Sprague
- BB. Pycnidia intermediate, 100-180 μ in diameter
 - C. Spores filiform, 2-septate, 40-56 x 1.8-2.4 μ
S. elymi Ell. and Ev. on Elymus canadensis var. robustus
 (Scribn. and Sm.) Mackenz. and Bush
 - CC. Spores elongate-fusiform, 26-38 x 2-2.5 μ
S. spartinae (Trel.) Sprague
 - CCC. Spores elongate-fusiform, 24-42 x 2.5-4 μ , pycnidia
 sometimes less than 100 μ diameter
S. calamovilfae Petr.
 - CCCC. Spores bacillar, 1-septate, 3-septate in one Mandan, N.
 Dak., collection S. oudemansii Sacc.
 - CCCCC. Spores filiform, faintly septate, 50-70 x 1-1.5 μ
S. cenchrina J. J. Davis
- BBB. Pycnidia small, obscure, 50-120 μ diameter
 - C. Pycnidia typically 50-80 μ , spores 0- to 1-septate
S. triseti Speg. em. Sprague
 - CC. Pycnidia typically 80-120 μ , spores 0- to 2-septate
S. tenella Cke. and Ell.
- AA. Spores typically more than 2-septate
 - B. Pycnospores usually less than 50 μ long
 - C. Spores averaging more than 2 μ wide
 - D. Spores bacillar
 - E. Spores seldom more than 30 μ long
S. nodorum Berk.
 - EE. Spores frequently more than 30 μ long
 - F. Produces a pale-colored leaf spot, spores
 3-septate, 25-50 x 2.5-3.5 μ , on rye
S. secalis Prill. and Del.

- FF. Produces dark brown leaf spots, spores
3-septate, 25-50 x 2.7-4.1 μ , on Lolium
S. loligena Sprague
- FFF. Produces a pale-colored leaf spot, spores 1-
to 3-septate, 37-62 x 2.6-3.4 μ , on Stipa,
Agrostis
S. secalis var. stipae Sprague
- FFFF. Produces a brown leaf spot, spores 3-septate,
22-38 x 2.8-3.8 μ , on Melica
S. melicae Pass.
- FFFFF. Produces a pale leaf spot, brown-bordered,
spores 3- to 4-septate, 25-45 x 3-4 μ , on
Avena S. avenae Frank
- DD. Spores narrowly obclavate, apex more or less pointed
- E. Pycnidia 100-140 μ , on Sporobolus
S. andropogonis f. sporobolicola Sprague
- EE. Pycnidia 75-100 μ , on Andropogon
S. andropogonis J. J. Davis
- EEE. Pycnidia 130-160 μ , on Sphenopholis
S. andropogonis J. J. Davis
- CC. Spores less than 2 μ wide
- D. Pycnidia golden-brown, ellipsoid
- E. Pycnophores 4-7 μ long, spores 35-40 x 1.3-1.6 μ
S. macropoda Pass.
- EE. Pycnophores 3-4 μ long, spores 30-50 x 1.2-2.1 μ
S. elymi Ell. and Ev.
- DD. Pycnidia creosote-brown, subglobose to ellipsoid,
spores 14-46 x 1.3-2 μ
S. passerinii Sacc.
- BB. Spores frequently more than 50 μ long
- C. Spores narrowly filiform, about 1 μ wide
S. stipina Died.
- CC. Spores more than 1 μ wide
- D. Spores often more than 2.7 μ wide

E. Spores averaging less than or approximately 65 μ long

F. Spores strictly hyaline

G. Spores 2- to 4-septate, stiffly curved, obclavate-filiform, pycnophores 4-9 μ long (in part)
S. infuscans (Ell. and Ev.) Sprague

GG. Spores mostly 3-septate, somewhat whip-like, obclavate-filiform, pycnophores 5-11 μ long
S. bromi var. phalaricola Sprague

GGG. Spores 3-septate, obclavate-cylindric to narrower S. agropyrina Lob.

GGGG. Spores 3- to 5-septate, narrowly obclavate, up to 73 μ long on Stipa comata
S. andropogonis f. sporobolicola

FF. Spores somewhat chlorinous
S. mississippiensis Sprague

EE. Spores averaging more than 65 μ long

F. Pycnidia prominent, as much as 320 μ in diameter

G. Spores seldom more than 5-septate, lance-shape, obclavate
S. jaculella Sprague

GG. Spores 3- to 7-septate, obclavate, apex elongated
S. arctica Berk. and Curt.

FF. Pycnidia smaller, commonly less than 150 μ diameter

G. Pycnophores narrowly ampulliform, spores long-filiform
S. pacifica Sprague

GG. Pycnophores cylindrical, spores obclavate-filiform
S. munroae Ell. and Barth.

DD. Spores less than 2.7 μ wide but more than 1 μ wide

- E. Pycnidia brown, cells compact, frequently elongated
- F. Pycnophores awl-shape, 2-5 μ long
 - G. Spores very narrowly obclavate-filiform
 - H. On Agrostis exarata
S. calamagrostidis (Lib.) Sacc.
 - HH. On Koeleria cristata
S. calamagrostidis f. koeleriae
 (Cocc. and Mor.) Sprague
 - GG. Spores filiform, less often very narrowly obclavate-filiform
 - H. Spores typically sinuous
S. calamagrostidis on Agrostis palustris
 - HH. Spores usually slightly curved, but less often sinuous
S. calamagrostidis on Trisetum spp.
- FF. Pycnophores short, papillate, 2-3 μ long ...
S. macropoda on Poa howellii Vasey and Scribn.
- EE. Pycnidia golden-brown, cells moderately compact, variable in shape, not uniformly elongated
 - F. Spores seldom over 1.7 μ wide
S. macropoda var. septulata (Gonz. Frag.) Sprague
 - FF. Some spores more than 1.7 μ wide
 - G. Spores never 8- to 9-septate
 - H. Pycnophores 2- to 6 μ long
S. macropoda var. grandis Sprague
 - HH. Pycnophores 4 to 13 μ long
 - I. Spores typically sinuous
S. tritici f. avenae (Desm.) Sprague
 - II. Spores variously curved, less often sinuous

J. Microspores present, macrospores 25-90 x 1.4-2.6 μ , on wheat
S. tritici Rob.

JJ. Microspores not known

K. Pycnosporos up to 85 μ long, on Lolium
S. tritici var. lolicola
 Sprague and A. G. Johnson

KK. Pycnosporos up to 105 μ long, on Holcus
S. tritici f. holci
 Sprague

GG. Spores 3- to 9-septate, up to 95 μ long.
S. andropogonis var. sorghastri H. C.
 Greene and Sprague

GGG. Spores usually 5-septate, up to 60 μ long
S. quinquesepitata Sprague

SEPTORIA AGROPYRINA Lob., Leaf Spot and Blotch.

This species is a very common but relatively unimportant component of the leaf spot complex on Elymus and Agropyron spp. in the Great Plains. Its range, however, extends west to Oregon. It apparently occurs widely in Asia and adjacent Europe. It is usually found as a saprophyte on the tips of dead or nearly dead leaves but some collections show it associated with a brown or fuscous blotch. The spores are coarse, obclavate to sub-cylindric, and are apically pointed in most specimens. They average 40-65 x 2.5-4.0 μ and are borne in large brown pycnidia. It is difficult to distinguish this fungus from a slender phase of Septoria infuscans on the one hand and from Stagonospora arenaria Sacc. on the other. Its pointed spores more or less distinguish it from Stagonospora arenaria, while its development on potato dextrose agar (it forms a cottony buff colony) contrasts sharply with the scanty pellicular growth of S. infuscans. While typical S. infuscans does not appear to extend east of Yellowstone National Park, S. agropyrina does occur with Stagonospora arenaria west to Oregon.

Septoria agropyrina is readily segregated from most material of S. elymi, as the latter has much smaller spores and produces a yeasty colony on potato dextrose agar. There is, however, a phase of S. elymi that is the same as S. elymicola Died. which may sometimes be intermediate between typical S. elymi and the more slender spored collections

of S. agropyrina. This phase usually occurs on Elymus canadensis or more often on E. canadensis robustus. This phase has been seen from Iowa and North Dakota while Hardison sent us material from Michigan.

Septoria agropyrina is further briefly discussed in connection with associated forms. It should be pointed out that critical comparison with European types is needed in this group.

References: Lobik (411); Sprague (665).

SEPTORIA ANDROPOGONIS J. J. Davis, Leaf Spot.

This fungus and its various forms and variety is very common in the Great Plains region. While the species proper on Andropogon furcatus Muhl. has not been collected closer than Kansas (Lefebvre, see Sprague, 660) and Wisconsin (Davis, 139, type area), it probably occurs on this host within the area covered by this article.

Gray elongate lesions are formed with moderately prominent pycnidia more or less seriatly arranged. The spores are characteristically obclavate with tapering blunt bases and narrowly sharpened apices (Sprague, 660, fig. 1, b). They are usually 3-septate, 30-51 x 2-3.9 μ , and the contents are hyaline to yellowish.

One collection on Sphenopholis obtusata from Devil's Lake, North Dakota, appears to be S. andropogonis rather than S. quinqueseptata Sprague, which also occurs on this host.

References: Davis (139); Sprague (660).

SEPTORIA ANDROPOGONIS var. SORGHASTRI H. C. Greene and Sprague, Leaf Spot.

This differs from the species in the much longer and multiseptate spores. Except for their extreme length they are fundamentally the same type of spore on a closely related genus. The first collection of this fungus on Sorghastrum nutans (L.) Nash was made by Rev. J. M. Bates, many years ago, at Long Pine, Nebraska, but the specimen was undetermined until Greene and the writer compared notes with some recently collected Wisconsin material and concluded that the fungus was only an unusual phase of S. andropogonis.

References: Greene (290); Sprague (665).

SEPTORIA ANDROPOGONIS forma SPOROBOLICOLA Sprague, Leaf Spot and Blotch.

The type of this occurs as a non-striate circular spot on Sporobolus

heterolepis A. Gray, but similar material has recently been found on Stipa comata at Mandan, North Dakota, which forms a blotch at first, later a light-colored spot. The spores of the type on Sporobolus are slightly narrower than those of Sentoria andropogonis proper, while those on Stipa are somewhat longer ($40-73 \times 2.4-3.4 \mu$), but also proportionately slightly narrower. This character, plus the somewhat larger, light-colored pycnidia in f. sporobolicola, readily segregates the form from the species. Except for these differences they all belong in the one species. At present f. sporobolicola is confined to the Agrostidae; while the species proper and its var. sorghastri occur on the Andropogoneae except for the Devil's Lake collection on Sphenopholis of the Aveneae. The possibly related species S. quinquesepata also occurs on Sphenopholis, while another related but we believe distinct species, S. mississippiensis, occurs on Muhlenbergia of the Agrostideae. It would appear that this group has adapted itself to a considerable range of tribes under the favorable conditions of association in the plains country. Since much of this information has been obtained only recently it is expected that further forms will be noted.

Forma sporobolicola was recently identified also on Stipa viridula from South Dakota. It is likely much more common than collections indicate, as much of the material on Stipa is not fruiting.

References: Sprague (660, 665).

SEPTORIA ARCTICA Berk. and Curt., Leaf Spot.

The type of this species was collected on Dupontia fischeri R. Br., on an island off the Siberian coast in the Bering Straits. It occurs also on Calamagrostis nutkaensis near Seaside, Oregon, and apparently in southern Alaska and northern Europe (Sprague, 665). The spots are ill-defined brown to dull gray lesions; pycnidia are large ($140-250 \mu$); spores are hyaline obclavate-scolecosporous with tapering rounded bases and sharply pointed apices. The spores in the type are 2- to 3-septate or more, $60-80 \times 2.8-3.6 \mu$, straight to stiffly curved, and somewhat resemble spores of S. jaculella Sprague on Bromus spp. On C. nutkaensis, the spores are 3- to 7-septate, $70-89 \times 3.5-4.6 \mu$. It is suspected that this species is more widespread and abundant than collections indicate. It appears to be distinctly parasitic.

References: Berkeley and Curtis (42); Sprague (665).

SEPTORIA AVENAE Frank, Speckled Blotch of Oats.

(Ascigerous stage, Leptosphaeria avenaria G. F. Weber.)

A collection on some late-seeded green-manure oats on the Northern Great Plains Field Station, Mandan, North Dakota, represents the only

collection of this fungus from the western area, although it is common in Wisconsin and northward and is also probably present in Minnesota.

The 3-septate bacillar spores of this fungus are close to Stagonospora. Since the perithecial stage is known (Weber, 752), the fungus can be referred to that stage and the questionable status of the imperfect stage is not so important as if its perithecial stage were unknown. Routine workers will recognize the pycnidial stage as a Septoria, although it is probably closer to the narrow-spore Stagonospora forms that produce cottony mycelium from germinating spores on potato dextrose agar, as contrasted with the slimy yeasty (mucose) conidial masses that most grass-inhabiting species of Septoria form under the same conditions.

References: Frank (237); Weber (752).

SEPTORIA BROMI Sacc., Leaf Spot.

Lesions on brome grasses are obscure, sometimes evident as elongate or elliptical spots, but often the infected leaves turn yellow, then dry up and become brown. The pycnidia are only moderately prominent, 60-240 μ , black-brown, and the spores are characteristically whip-like, usually 2-septate, 33-65 x 1.2-2.5 μ , or sometimes stouter in Oregon material, but not to be confused with the javelin-like spores of S. jaculella. The host range includes: Bromus commutatus Schrad. (Oreg., Idaho); B. inermis (Mandan, N. Dak., Oreg., Wash.); B. japonicus Thurb. (Wash., Wyo.); B. mollis (Oreg., Wash.); B. racemosus L. (Oreg.); and B. secalinus L. (Oreg., Idaho, Iowa). This fungus is seldom serious in the West.

References: Sprague (665); Weber (754).

SEPTORIA BRÖMI var. PHALARICOLA Sprague, Buff Spot.

This fungus occurs in buff lesions on the leaves of reed canary grass at Astoria, Oregon. The spores are obclavate-filiform, 3-septate instead of 2-septate as in S. bromi, 37-66 x 1.7-2.9 μ , borne in large pycnidia 150-180 x 204-276 μ . This species may be distinguished from certain other pyrenidium-bearing leaf spots of reed canary grass as follows:

- A. Spots pale with narrow fuscous or tinted border, spores non-septate, falcate Selenophoma donacis (Pass.) Sprague and A. G. Johnson
- AA. Spots buff, yellow, or obsolete

- B. Spots pale or absent, spores 0- to 1-septate, falcate, from the Argentine
Septoria macrostoma Speg. (Sprague, 665, fig. 19, c)
- BB. Spots buff
- C. Spores 1-septate, hyaline, 20-25 x 1.3 μ , from Italy ..
Septoria phalaridis Cocc. and Mor. (120)
- CC. Spores 3-septate, hyaline, obclavate-filiform, from Oregon and probably Italy
Septoria bromi var. phalaricola Sprague (665)
- BBB. Spots tawny blotches, spores 38-71 x 4.5-6.0 μ (when mature), 3- to 8-septate (3-septate phase 25-40 x 2.7-4.5 μ), from Middle West and the Great Plains and in Europe
Stagonospora foliicola (Bres.) Eub. (See under Stagonospora below)

SEPTORIA BROMI var. PHALARIDIS Trail (Trail 709) is PHAEOSEPTORIA PHALARIDIS (Trail) Sprague (Sprague 659)

References: Cocconi and Morini (120); Saccardo (554); Sprague (659, 665); Trail (709)

SEPTORIA CALAMAGROSTIDIS (Lib.) Sacc., Leaf Spot.

The lesions are gray to straw color, usually not abundant, and occur most commonly during late winter or early spring in Oregon. The pycnidia are strongly flattened, dark brown, 50-140 x 50-180 μ in lateral diameter and 50-80 μ vertically, and are characteristically formed of compactly intertwined cells 1.7 μ in diameter. The spores are filiform, 0- to 5-septate, 25-71 x 1.0-2.0 μ , often sinuous. The host range of this species is as follows: Agrostis diegoensis Vasey (Oreg.); A. exarata (Oreg., Alaska); Trisetum canescens Buckl. (Oreg.); T. cernuum Trin. (Oreg.). The fungus on Trisetum usually has stiffer straighter spores than on Agrostis but one collection on T. canescens from Douglas County, Oregon, has spores identical with those on Agrostis. This form on Trisetum is confined to species of that genus on which it is very parasitic. It is considered to be only a race of S. calamagrostidis. Such species as S. caballeroi Gonz. Frag. (270), S. caballeroi var. panicei Gonz. Frag. (272), and S. graminum f. triseti-loeflingianii Cab. (88) are synonyms.

Our studies indicate that material on Agrostis palustris may be called race 1, on A. exarata, race 2; and on Trisetum spp., race 3.

Other fungi that may be confused are the shorter-spored Septoria triseti (common on A. tenuis but not on A. pelustris, the reverse of the situation with S. calamagrostidis), and the bacillar spored S. secalis var. stipae which sometimes occurs on Agrostis.

References: Anderson (11); Caballero (88); Cocconi and Morini (120); Gonzalez Fragoso (270, 272); Saccardo, Peck, and Trelease (559); Sprague (642, 665)

SEPTORIA CALAMAGROSTIDIS (Lib.) Sacc. f. KOELERIAE (Cocc. and Mor.)
Sprague, Leaf Spot. (S. koeleriae Cocc. and Mor.)

This fungus differs from S. calamagrostidis in its stiff needle-like spores, 35-60 x 1.1-2 μ , that are readily distinguishable from the species proper. It is confined to Koeleria.

Septoria calamagrostidis f. koeleriae occurs from North Dakota to Oregon and Washington. The fuscous spots are relatively abundant in spring on Koeleria cristata in the eastern and mountainous parts of Oregon and Washington.

References: Cocconi and Morini (119); Sprague (665).

SEPTORIA CALAMOVILFAE Petr., Leaf Spot.

The type locality of this species is Kulm, North Dakota, where it occurs on Calamovilfa longifolia (Hook.) Scribn. The writer has found it at Faith, South Dakota, and near Colony, Wyoming. The spores are smoky-hyaline, spindle-shape, 24-42 x 2.5-3.7 μ , 0- to 2-, sometimes 3-septate. It causes a white or pale gray spot on the leaves and sheaths.

Reference: Petrak (510, p. 86).

SEPTORIA CENCHRINA J. J. Davis, Leaf Spot.

Reported on Cenchrus pauciflorus Benth. from Minnesota as well as from the type area at Spring Green, Wisconsin. This species has filiform, continuous or faintly septate spores, 30-100 x 1.5-3.0 μ , borne in proportionately small pycnidia, 60-115 μ . The longer spores (Davis, 142) protrude through the wide ostiole.

References: Davis (142, p. 296).

SEPTORIA ELYMI Ell. and Ev., Speckled Leaf Blotch
(Septoria agropyri Ell. and Ev.)

This fungus causes pale gray to tan or fuscous lesions which, with the numerous pycnidia, produce the effect of speckled blotch. The spores are smaller than those of S. tritici, 25-50 x 1.2-2.1 μ , 0- to 3-septate. A form of this fungus on Elymus canadensis var. robustus in Iowa, Wisconsin, and Michigan has 2-septate larger spores, 40-56 x 1.8-2.4 μ . The form on Agropyron spp. (Ellis and Everhart, 213) has spores too nearly like those of S. elymi Ell. and Ev. (212) to be worthy of recognition as a separate species, therefore S. agropyri Ell. and Ev. (214) is considered a synonym of S. elymi. The host range also is partly interchangeable.

The host range of Septoria elymi is as follows: Agropyron inerme (Oreg.); A. repens (Oreg., Wash., Mont., N. Dak., Minn., Iowa, Nebr.); A. smithii (N. Dak., S. Dak.); A. spicatum (Wash., Wyo.); A. subsecundum (Mont., N. Dak.); A. trachycaulum (N. Dak.); Elymus canadensis (N. Dak., S. Dak., Minn.); E. canadensis var. robustus (Iowa); E. glaucus (Oreg., Wash.); E. triticoides (Calif.)

S. elymi is readily distinguished by its smaller spores from S. infuscans, S. pacifica, Stagonospora arenaria, and Septoria agropyrina, all of which occur on Elymus and Agropyron.

References: Ellis and Everhart (212-214); Sprague (665); Weber (754).

SEPTORIA INFUSCANS (Ell. and Ev.) Sprague, Brown Leaf Blotch.
(Cylindrosporium infuscans Ell. and Ev.)

Elymus condensatus and sometimes E. glaucus (Oregon and Washington), Agropyron spicatum (Oregon), and Elymus triticoides (Oregon, California) are heavily blotched by this fungus in the western States as far east as Yellowstone National Park, Wyoming. The type material and much of that from the Columbia Basin of Oregon and Washington has obclavulate, blunt-base pycnosporos tapering to a stubby point at the apex, 1- to 3-septate, 28-50 x 2.5-4.5 μ (Cooke, 126, no. 113). However, some material, particularly that on E. triticoides, has obclavulate-filiform spores 1- to 4-septate, 40-75 x 1.7-2.7 μ , resembling those of S. elymi-europaei Jaap. In pure culture, both kinds produce small gray pellicular colonies with imbedded pycnidia. One isolate of a narrower form from E. condensatus, Washtucna, Washington, produced an off-white mucose growth that spread somewhat more rapidly than most of the isolates, and this may be different.

East of the range of S. infuscans, a large spored fungus in white stramineous or fuscous lesions occurs on Elymus and Agropyron. This is referable to S. agropyrina Lob. (non Unamuno, 731). It produces a buff

cottony growth, similar to Stagonospora arenaria Sacc., in pure culture.

S. infuscans frequently has microspores, $5-7 \times 1-1.5 \mu$, narrowly ovate, short cylindric, often curved.

References: Cooke, W. B. (126, No. 113); Ellis and Everhart (216); Jaap (357); Sprague (649, 665); Unamuno (731).

SEPTORIA JACULELLA Sprague, Western Brome Speckle.

This fungus has been known for only a few years, but it occurs on a leaf of Bromus carinatus collected by David Douglas in 1832, probably from southern Oregon. The fungus invades living tissue very slowly so that the lesions are vague, mottled or blotched, partly owing to the fact that green, brown, and yellow tissues are intermingled with mature pycnidia. Sometimes streaks of dead tissue are formed.

The pycnidia are large; the diamond blue hyaline spores are stiffly javelin or lance-shaped, $45-90 \times 1.9-4.0 \mu$, with numerous microspores present in some collections. Efforts to obtain the fungus in pure culture have been negative; the spores will germinate on potato dextrose agar, but they soon die.

The host range is as follows: Bromus carinatus (Oreg., Wash; Trude, Idaho); B. ciliatus L. (Bradshaw Mts., Ariz.); B. laevipes Shear (Calif.) B. rigidus Roth (Oreg., Wash.), and B. tectorum (Wash.).

References: Cooke (127, no. 114); Sprague (650, 665).

SEPTORIA LOLIGENA Sprague, Brown Leaf Spot.
(S. lolii (Cast.) Sacc.)

Causes a deep chocolate-brown leaf spot surrounded by lighter areas. The spores are bacillar, clear hyaline, 3-septate, $28-45 \times 2.7-4.2 \mu$, and are therefore similar to those of S. secalis or to S. avenae and not to those of S. tritici var. lolicola, which has much narrower spores.

Besides the original material from France, this fungus occurs in California on Lolium multiflorum. It was collected in the San Francisco area by W. B. Cooke recently (Sprague, 665).

References: Castagne (98); Sprague (665).

SEPTORIA MACROPODA Pass., Leaf Blotch.
(Septoria annua Ell. and Ev., S. poae-annuae Bres.)

The common leaf blotch on Poa annua and on P. kelloggii Vasey (Oregon),

P. howellii (Oregon), and P. nervosa is abundant in the northern United States wherever the hosts occur and humid conditions prevail during the growing season. It is obtainable any month of the year in coastal Oregon but is particularly prevalent in late winter. The spores are very slender, $30-40 \times 1-1.5 \mu$.

References: Bresadola (67); Cocconi (59); Ellis and Everhart (215); Gilman and Archer (262); Sprague (665).

SEPTORIA MACROPODA var. GRANDIS Sprague, Leaf Blotch.

The spores of this variety are larger than those of S. macropoda, some of them approaching Septoria tritici, $40-70 \times 1.5-2.4 \mu$. It occurs in the interior of the United States from the Cascade Mountains in Oregon, Washington, and California to the center of North Dakota. The host range of this variety is as follows: Poa ampla (Wash.); P. arida (Faith, S. Dak.); P. canbyi (Mont.); P. cusickii Vasey (Mont.); P. nevadensis Vasey (Oreg., Calif.); P. pratensis (Wyo.); P. scabrella (Thurb.) Benth. (Wash.); P. secunda (Oreg., Wash., S. Dak., Wyo., Mont., N. Dak.); P. vaseyochloa (Wash.). The collection on P. pratensis was more or less accidental; the common form on that host is var. septulata to be discussed later. Septoria macropoda var. grandis is very widespread in the high range country in the very early spring and has been collected as early as December in Klickitat County, Washington. As soon as the weather warms up and becomes dry in Oregon and Washington the spots lose their identity and are difficult to see. Some racial differences were noted in this variety.

References: Sprague (656, 665).

SEPTORIA MACROPODA var. SEPTULATA (Gonz. Frag.) Sprague, Leaf Blotch.
(S. poae-annuae var. septulata Gonz. Frag.)

Septoria macropoda var. septulata is cosmopolitan throughout the northern United States as the cause of a dark gray or brown spot irregular in shape on the leaves of Poa pratensis. It is readily recognizable from the species by its needle-like spores, which are $40-60 \times 1.3-1.7 \mu$ and are narrower than those of var. grandis. While these forms are similar enough to be placed in one species, they are sufficiently distinct from each other both morphologically and ecologically to be recognized as subdivisions. The var. septulata also occurs on Poa compressa and on P. nervosa.

References: Gonzalez Fragoso (273, 274); Sprague (657, 665); Weber (754).

SEPTORIA MELICAE Pass., Leaf Blotch.

This fungus was found in the grass nursery at the Northern Great Plains Field Station, Mandan, North Dakota, in August 1943. It caused a brown leaf spot on Melica scabrosa Trin. This is another of the complex of closely related species with 3-septate bacillar spores, in this case, 22-38 x 2.8-3.8 μ . It is slightly less robust appearing than S. avenae. There is no report on this fungus for North America except that Schizachne purpurascens (Torr.) Swallen is listed as a host in Seymour's list (599). The writer (666) mentions it in a recent article.

SEPTORIA MISSISSIPPIENSIS Sprague, Leaf Spot.

This fungus causes a buff eyespot with prominent brown borders on Muhlenbergia mexicana (L.) Trin. in Minnesota and on M. asperifolia (Nees and Mey.) Parodi in North Dakota. The spores are like those of S. andropogonis (see also Karsten, 378, on S. bromi var. alopecuri Karst.) except that they are chlorine yellow instead of hyaline to faintly yellow, and the contents appear much different as can be readily seen by comparing the two species (Sprague, 660, fig. 1, a,b,c). S. mississippiensis does not appear to be very abundant.

References: Karsten (378); Sprague (660, 665).

SEPTORIA MUNROAE Ell. and Barth, Leaf Spot.

The lesions on Munroa squarrosa (Nutt.) Torr. in Kansas and Colorado are straw color and obscure but the seriatly arranged pycnidia are abundant. Spores are 80-110 x 2.5-3.0 μ and resemble certain species of Cercospora. This species has not been studied to any extent.

References: Ellis and Bartholomew (208); Sprague (665).

SEPTORIA NODORUM Berk., Glume Blotch.

Typical spores of the fungus causing glume blotch of wheat are short, cylindrical, 0- to 3-septate, 15-30 x 2.3 μ . This species is widely known in Europe and in the northern United States and Canada on wheat and wheat relatives as a leaf spot and later, towards harvest, as a blotch on the glumes of the maturing heads and on culm parts, particularly the nodes. In Germany, the fungus is often known as Macrophoma hennebergii (Kuehn) Berl. and Vogl. (43) or sometimes by the earlier name of Phoma hennebergii Kuehn (393). Petrak and Sydow (512) placed it in Stagonospora hennebergii (Kuehn) Petr. and Syd. A study of material from Germany, from North Dakota, and from adjacent regions shows that true S. nodorum is associated with forms that have somewhat larger spores, 25-45 x 2.5-4.3 μ . In addition material on Elymus and

Agropyron is confused with Septoria agropyrina Lob. but in the latter the apical end of the longer spores tends to be somewhat pointed and less cylindrical. Just where Stagonospora arenaria Sacc. ends and S. nodorum begins in some of the collections is also an open question. In fact there is some question as to whether S. nodorum is not a species of Stagonospora, as pointed out by Petrak and Sydow (512).

After segregating what appears to be Septoria agropyrina and Stagonospora arenaria from Septoria nodorum, the host range of the latter is as follows: Agropyron repens (N. Dak.); A. trachycaulum (N. Dak.); Elymus excelsus Turcz. (N. Dak.); E. sibiricus L. (N. Dak.); Glyceria elata (Nash) Hitchc. (Cornucopia, Oreg.); G. pauciflora Presl (Cascade Mts., Wash.); G. striata (Lam.) Hitchc. (Hoover, Oreg.); Hystrix patula (Mim.); Poa palustris (Rough Lock Falls, S. Dak.); Poa pratensis (Oreg.); Triticum aestivum L. (wheat) (Mont., N. Dak., S. Dak., Nebr., Minn., Iowa); T. spelta L. (Alsea Valley, Oreg.); Stipa viridula (Mandan, N. Dak.); T. dicoccum Schrank (N. Dak.).

The collections on Glyceria represent a fungus that may warrant separation from Septoria nodorum. The spots are large, buff, later paler, and the spores appear narrower than typical S. nodorum. The apices are somewhat pointed but the spore shape is more slender and not at all like that of S. agropyrina, which is a much coarser fungus. The writer (638) referred the first collection, from Hoover, Oregon, to Stagonospora glyceriae Roum. and Fautr., but better material has since shown that this was incorrect. Host range studies are needed with this species or form to determine its relationship to the group.

Septoria nodorum produces a buff or light-colored cottony mycelium in potato-dextrose agar and superficially resembles isolates of Stagonospora arenaria and of most of the other bacillar-spored forms. In producing a cottony growth, it differs fundamentally from the true Septoria group with filiform spores, which mainly produce masses of mucose conidia in pure culture from germinating spores.

References: Alfaro (6); Berkeley (41); Berlese and Voglino (43); Bockmann (53); Diedicke (167); Grove (295,296); Kuehn (393); Luchetti (412); Machacek (419); Passerini (503); Petrak and Sydow (512); Rosen (544); Sprague (665); Sydow (695).

SEPTORIA OUDEMANSII Sacc., Leaf Spot.

Light purple, finally straw color lesions occur on Poa pratensis and P. compressa during wet weather in Oregon and Washington. The lesions on these hosts and on P. canbyi, P. secunda, P. juncifolia, and P. nevadensis from the Rocky Mountains and the Northern Great Plains are obscure, sometimes vague blotches, more often faded to neutral straw color. The disease is not important, although it was severe at Mandan in August 1943. In western Oregon and Washington it occurs during the

rainy period in winter and early spring. Griffith (293) indicates that the disease is important on Poa pratensis leaves in Kentucky during the cool, wet season. Seymour (599) lists S. oudemansii on Poa flexuosa var. elongata Blytt and on Hierochloë alpina (Sw.) Roem. and Schult., probably from northeast America. The first record from this country appears to be a collection made by the writer in Klickitat County, Washington, in February 1935, although he had the fungus in pure culture as early as 1932 but did not determine its identity until later. In 1935, he reported it on Poa pratensis in two places in western Oregon as Ascochyta sp. (Sprague, 629) and in 1942 corrected this to Septoria oudemansii Sacc. (Sprague, 655, 656). He also found it in material sent to him from Michigan for determination (Hardison, 310). Host range additions were noted by the writer in 1942 (Fischer, et al., 231). The fungus therefore appears to be firmly established in literature as a recognized parasite of bluegrasses in the northern United States.

In first classifying the fungus as a species of Ascochyta, the writer (629) placed the fungus where its most common spore phase should be placed. In all but one collection of several dozen seen, the spores are 12-24 x 1.7-2.8 μ , cylindrical but typically slightly thicker at the central septum. Microspores range from 4 x 2 μ up to typical Ascochyta macrospores. That this Ascochyta is potentially a species of Septoria is shown by one collection made at Mandan in rainy weather in August, 1943. The spores are 3-septate, 35-55 x 3-4 μ , mingled with spores 11-18 x 2.5 μ . In pure culture they are identical. Therefore, S. oudemansii is left in Septoria even though its common spore condition is that of a slender Ascochyta phase.

References: Allescher (7); Fischer, et al. (231); Griffith (293); Hardison (310); Seymour (599); Sprague (629, 655, 656, 665).

SEPTORIA PACIFICA Sprague, Leaf Elotch.

The spots are fuscous, soon straw color, the pycnidia prominent and seriatly arranged. The spores are very slender, 1- to 2-septate, 60-110 x 1.5-3.5 μ , and therefore are too long and narrow for any phase of S. infuscans or of Stagonospora arenaria. Septoria pacifica is found on coarse beach grass, Elymus mollis Trin., along the seashore in Oregon and also on E. condensatus var. pubens Piper in central California (H. W. Johnson). A stray isolate of apparently the same species was obtained from E. glaucus in plots at Mandan, North Dakota.

The fungus produces a pale caenstone color colony, which eventually becomes covered with small-mounded, black, stromatic tissue which exudes white spore masses. The colonies, except for the color of the spore exudate, resemble the common form of S. infuscans.

Reference: Sprague (665).

SEPTORIA PASSERINII Sacc., Leaf Blotch.

Fuscos, somewhat indefinite lesions are formed on barley and related grasses. The leaves later turn yellow, then brown after death. The pycnidia are only moderately conspicuous against the strongly fuscous lesions. The pycnidia are creosote-brown, mostly subglobose, scarcely erumpent, 90-140 μ in diameter and up to 100 μ deep. The walls are relatively thin, 5 to 10 μ , composed of strongly pigmented cells and an inner layer of polyhedral cells. The pycnophores are somewhat filiform to subulate, 3.4-5 x 1.2-1.8 μ . The pycnospores are of two kinds: (1) Macrospores which are broadly filiform, obtusely pointed at both ends, rarely curved, 1- sometimes 3-septate, 22-44 x 1.5-2.2 μ ; but saprophytic material on Hordeum nodosum from Sydney, Montana, had spores 45-70 x 1.8-2.4 μ , although typical material on the same host in the same general locality had spores 30-45 x 1.6-2.0 μ . Material on Sitanion hystrix from Mt. Hood, Oregon, had narrowly obclavate to nearly filiform spores 14-46 x 1.3-2.0 μ (mostly 40 x 1.4 μ). (2) Microspores, which are much smaller, bacteria-like, 3-6 x 0.3-0.6 μ , but occur in typical pycnidia of the species. Microspores are common on H. nodosum in Oregon and have been noted on Hordeum distichon L. from the Kiev District, Russia.

Septoria microspora Ell. on Hystrix patula (Indiana) has spores 9-12 x 0.7-1.2 μ (Ellis, 207) and is likely only a microspore stage of S. passerinii. Recently macrospores referable to S. passerinii were found on Hystrix patula at Cotton Lake, Minnesota. (Sprague, 669).

S. passerinii is rare on cultivated barley (Hordeum vulgare) in the far West but is common on Hordeum nodosum in Oregon, Washington, Wyoming, and Montana, and is very prevalent on Hordeum jubatum in the Dakotas. From observations made in eastern South Dakota and from the earlier collections of A. G. Johnson (368) it is apparent that barley is commonly parasitized by this fungus in the area.

References: Ellis (207); A. G. Johnson (368); Passerinii (505); Saccardo and Trotter (560); Sprague (665); Weber (754).

SEPTORIA POLICMELA Syd., Leaf Spot.

Forms gray to straw color spots on the sides and tips of the leaves of Deschampsia caespitosa (western Oregon) and tan to straw color lesions on the lower leaves of D. danthonioides in Oregon and Washington (Sprague, 655, 656). The spores are clear hyaline, strongly curved, 36-50 x 1.4-1.6 μ , C- to 2-septate on D. danthonioides. On D. caespitosa the spores are 1- to 3-septate, needle-like, tapering towards each end or slightly blunt at one end, 33-47 x 1.5-2.0 μ , mostly about 40 x 1.8 μ . In the type collection on Aira caryophyllea L. from Madeira (Sydow, 694) the spores were 15-30 (50) x 1.5-2.5 μ (Sprague, 665, fig. 13,b), mostly 20-30 x 1.5-2.2 μ . The same fungus probably occurs in Morocco

(Maire and Werner, 429).

The fungus produced a white to pale cream color, slow growing mounded yeasty colony on potato dextrose agar which soon became covered with a short-nap, velvety, gray mycelium, similar to S. elymi (Sprague, 665).

References: Maire and Werner (429); Sprague (655, 656, 665); Sydow (694).

SEPTORIA QUINQUESEPTATA Sprague, Leaf Spot.

The type specimen from Mandan, North Dakota, occurs on Sphenopholis obtusata (665). The spores are uniformly filiform, 5-septate with obtuse bases and sub-acute apices, 48-58 x 1.9-2.4 μ , borne in pycnidia with small ostioles, 98-115 μ , on saprophytic, straw color material without any spotting.

Since the species was described, another collection on this host has been made near Devil's Lake, North Dakota, which seems to be Septoria andropogonis. The spores are 30-38 x 2.3-3.8 μ , mostly 2.8 μ wide. The pycnidia are 130-160 μ in diameter, which is atypical for most material of S. andropogonis (but see f. sporobolicola). The spots are small linear, white, finally fading to straw color. S. quinqueseptata occurs also on Koeleria cristata at Mandan, North Dakota (666). Further study may show that S. quinqueseptata is a form of S. andropogonis, which is becoming recognized as a somewhat polymorphic species on a wide range of hosts.

References: Sprague (665, 666).

SEPTORIA SECALIS Prill. and Del., Leaf Blotch.

The lesions on rye (Secale cereale L.) are yellow brown, the pycnidia are brown with walls of thick layers of light brown, rectangular cells, which give rise to creeping hyphae terminating in prominent subulate to almost fusiform pycnophores. These pycnophores are frequently bent or semi-sinuous and 5.5-8 x 1.9-2.2 μ (Sprague, 665). The spores are hyaline, 3-septate, narrowly cylindrical, 25-50 x 2.0-3.5 μ , mostly 35 x 2.7 μ and therefore slightly longer than the bare minimum of length to width ratio (10 to 1) permitted in Septoria. The microspore stage has spores about 10 x 0.5 μ and has been described as Phoma secalina Jancz. (S. secalina (Jancz.) Sacc.). S. secalis occurs in Iowa and has been collected near Washington, D. C. It was abundant in Minnesota, North Dakota, and South Dakota in 1944 as far west as Mandan. It was not serious in any fields that were examined. This species has not been found on rye in the Far West.

References: Janczewski (360); Sprague (665); Weber (754).

SEPTORIA SECALIS var. STIPAE Sprague, White Leaf Spot.

The lesions eventually become white or finally stramineous. The variety has spores 37-50 x 2.6-3.1 μ , appearing not only longer but also slimmer than the species proper on rye. The pycnidia are globose, 130-190 μ , with an erumpent ostiole which is somewhat more prominent than in the species. The variety is common on Stipa viridula in the Northern Great Plains and a similar fungus occurs on Agrostis scabra in the same region. The summer material on Stipa is often 1-septate and somewhat larger and proportionately more slender than Septoria nodorum but confusable at times. S. nodorum occurs on Stipa viridula in the region also. The same one-celled phase occurs in Oregon on Agrostis hallii. Also, one collection on Stipa williamsii Scribn. in plots at Mandan, North Dakota, had 1-septate spores only, 21-31 x 2.8-3.8 μ , and with microspores 5-10 x 0.5-0.7 μ .

Cross inoculation studies are needed to determine the host range of this variety and its relation to the species. In fact, a critical study is needed of the bacillar-spored group, including Septoria avenae, S. secalis, S. nodorum, S. melicae, S. loligena, and others, to determine whether some of them are not taxonomically unnecessary. In the writer's earlier investigations (665), this group was not available for adequate host range studies.

Septoria andropogonis f. sporobolicola, which occurred abundantly on Stipa comata at Mandan in 1944 has pointed, proportionately more slender spores.

References: Sprague (652, 665, 666).

SEPTORIA SPARTINAE (Trel.) Sprague, Leaf Spot.

Spots on the living leaves are elliptical to striate, straw to buff color, with obscure, tardily erumpent pycnidia 100 to 140 μ wide and 70 to 90 μ deep, deeply immersed, black, heavy walled, darker around the ostiole. The pycnophores are prominent, cuspidate to narrowly pyriform, arising from hyaline pycnophore initials along the floor of the pycnidium. The pycnidia are somewhat "Rhabdospora-like", that is, dark and heavy-walled, because they are buried in the silicified, thick leaves of the host. The hyaline, mostly 2-septate spores of material from Utah, are narrowly elongate-fusiform, tapering to a relatively sharp apex and a more gradually blunted or truncate base, 26-38 x 2.0-2.5 μ . Material from Wisconsin has spores that range up to 3.0 μ wide. Septoria spartinae occurs on Spartina gracilis Trin. (Utah) and on S. pectinata Lk. (S. Dak., Wis.). While it was originally described by Trel. (1885) as a species of Ascochyta, it is obviously closer to Septoria.

References: Davis (140); Sprague (665); Trel. (1885, p. 121; p. 17

of reprint).

SEPTORIA STIPINA Died., Leaf Spot.

The lesions on the dead necrotic leaves of Stipa columbiana var. nelsoni (Scribn.) Hitchc. in eastern Oregon are vague and the material scanty. The pycnidia are prominent, dark brown, erumpent, ostiolate, finally collapsed, 100-120 x 150-200 μ , composed of compacted, creosote-brown crushed cells. The spores are straight, very narrow, 1- to 4- (mostly 3-) septate, 39-63 x 0.8-1.1 μ (average 52 x 0.95 μ) (Sprague, 665). The only specimen seen in the United States is this fragmentary one from the Blue Mountains of Oregon. It appears to be closer to S. stipina than any other fungus encountered. S. stipina was described from material gathered at Kashmir, India, on Stipa sp. (Sydow and Butler, 696). It has dilute brown pycnidia, 90-120 μ in diameter, with prominent ostioles and filiform, curved to flexuous, hyaline spores 25-40 x 0.5 μ . The Oregon material therefore has longer, stiffer spores borne in larger pycnidia. Other species to consider are: S. stipae Died. (167), which, by virtue of priority became S. capillatae Trott. (Sacc. 558 v. 25, p. 430.), but Grove (296) indicated that S. stipae Died. had irregularly bent spores allied to S. lunata Grove and is therefore likely a species of Selenophoma in the Sel. donacis var. stomaticola complex. S. stipae Trabut (707) has superficial pycnidia and may be a species of an aseptate-spored Septoriopsis Gonz. Frag. (271).

References: Diedicke (167); Gonzalez Fragoso (271); Grove (296); Sprague (665); Sydow and Butler (696).

SEPTORIA TENELLA Cke. and Ell., Leaf Blotch.

Small, vague, gray to greasy brown spots form on the filiform leaves of the fescues. On some hosts, the lesions are stramineous and indefinite. The pycnidia are creosote-brown, 60-120 x 95-130 μ , more or less elliptical in cross section, composed of tightly crushed, oblong to polyhedral cells averaging 0.5 to 3.0 μ wide (or rarely more). The pycnophores are characteristically cylindrical, or very narrowly subulate, averaging 1 μ wide and up to 4 μ long. Sometimes the hyphae are scarcely different from the small globose to angular pycnophore initials. This species is characterized by the small size of the cells composing the pycnidial wall and by the small size of the pycnophores (Sprague, 665, fig. 12, e, k). The pycnosporos of S. tenella are exceedingly variable, 5-70 x 0.8-2.1 μ , but most commonly they are filiform, somewhat curved, 0- to 2-septate and 25-45 x 1-1.5 μ (Sprague 665, fig. 12).

Spores on Festuca dertonensis from western Oregon vary from 28-41 x 0.8-1.2 μ to 7-17 x 1.0-1.5 μ and even 9-12 x 1.4-1.7 μ in adjacent

pycnidia, while on F. rubra var. commutata all spore types are intermingled and found to some extent in the same pycnidium. The most common material and the most widespread occurs on F. octoflora Walt. from Idaho to Minnesota, with spores 0- to 2-septate, 44-67 x 1.6-2.1 μ . In May 1941, this fungus caused a speckled lesion on the basal leaves and culms of almost 100 percent of this grass over vast areas in the Dakotas. It was at that time the most abundant parasite on any grass in the region.

Material on F. rubra commutata from Oregon appears to be a variant similar to Septoria festucae Died. However, the material on F. dertonensis could be assigned to S. tenella, S. festucae, or Ascochyta sp., depending on the particular pycnidium examined. The material from the Great Plains is more typically and uniformly like S. tenella. It is no doubt racially distinct from the polymorphic West Coast material. S. tenella also occurs on F. ovina in the Yellowstone Park.

In pure culture, the isolates from F. rubra commutata produced a peculiar slimy cocoa brown mass of conidia, which contained many Ascochyta-like spores and also filiform spores plus many variants or freaks. Later the conidial masses, through germination of the spores, produced a somewhat carbonized surface to the colony. The isolations from near Grassy Butte, North Dakota, from F. octoflora were different in that mucose mounded, flesh pink colonies similar to Septoria tritici were formed.

A collection of a species of Phyllosticta on F. subulata had spores 4.5-11 x 1.4-2.0 μ and was assigned as being close to Septoria tenella but possibly different (Sprague, 665).

References: Cooke and Ellis (125); Diedicke (165); Sprague (665); Tehon and Daniels (701).

SEPTORIA TRIVETI Speg. em. Sprague, Leaf Mottle.

This fungus that causes a spot on Agrostis tenuis is commoner than the larger spored fungus S. calamagrostidis, which does not attack A. tenuis but is common on A. palustris. S. triveti also occurs on A. alba, A. exarata, and A. castellana Boiss. and Reut. in Oregon. S. triveti causes gray, later fuscous or brown spots, and the spots are more abundant and larger than those caused by S. calamagrostidis. The spores are 18-43 x 1.1-2.0 μ , 0- to 1-septate. It is confined, as far as known, to western Oregon and Washington in the United States.

References: Spegazzini (619); Sprague (665).

SEPTORIA TRITICI Rob., Speckled Leaf Blotch.

Septoria tritici causes a gray speckled blotch on the leaves of wheat and other Triticum spp. The numerous pycnidia are golden brown, sub-stomatal, ostiolate, usually distinctly flattened, ellipsoid, 60 to 200 μ in diameter, averaging usually 100 to 150 μ . The thin walls are composed of 2 to 5 outer layers of golden brown, oblong, box-shape, thin-walled cells; then inside that, 1 to 4 intermediate layers of polyhedral compacted cells; and inside this a thinner area of hyaline tissue that gives rise to narrowly ampulliform pycnophores. Grove (295, 296) lists Septoria graminum Desm. var. crassipes Grove on leaves of wheat and states that this variety has short ampulliform cuspidate sporophores measuring 10-13 x 2.5-3 μ . Grove's illustrations (295) indicate that he had the same fungus as the one on wheat in the western United States. The western material has flask-shape to almost cylindrical pycnophores measuring 5-13 x 1.3-3 μ (Sprague, (665, fig. 1).

Septoria tritici produces both microspores and macrospores. The macrospores are the more prevalent; they are 1- to 5-septate, narrowly obclavate-filiform, hyaline, 22-98 x 1.4-2.1 μ , mostly 45-80 x 1.6-2.0 μ . Summer material; often erroneously called S. graminum Desm., tends to more nearly filiform spores, shorter and with fewer septa.

Microspores are curved or hooked, aseptate, 5-9 x 0.3-1 μ , and are borne in small pycnidia 45-50 μ . The spores are so minute that they have no doubt been mistaken for bacteria. In pycnidia, 55-70 x 60-80 μ , sometimes both microspores and macrospores occur. Microspores may be produced on the tips of ampulliform pycnophores or on the tips of aborted macrospores (Sprague, 665, fig. 1). The microspore stage has been reported only from Oregon but has likely been overlooked elsewhere.

Septoria tritici occurs on Triticum aestivum, T. spelta, T. dicoccum var. farrum Bayle, and T. turgidum L. It is an important parasite on winter wheat in the moister parts of Oregon, Washington, and California and sometimes is serious in Nebraska and Kansas. It is less common east of the Cascade Mountains in Oregon and Washington and becomes decreasingly important eastward until it is virtually nonexistent in the Spring Wheat Belt of the Northern Great Plains. In parts of Oregon and Washington winter rosettes of wheat are 100 percent covered with the salt and pepper lesions of S. tritici and this injury cannot but have a depressing effect on the host, even though the fungus often disappears entirely when warm, drier weather comes in spring and summer.

It has been shown by Weber (753) that S. graminum does not occur on wheat in Wisconsin, while the writer reported that S. tritici was the only filiform-spored species present on wheat in the United States (665) and that S. graminum (642) was apparently limited to Brachypodium spp. and does not, as far as known, occur in this country. Most of the species that the writer assigned to synonymy are obvious synonyms, but

one, S. briosiana Mor. (473), is less obvious. In this form microspores (9-11 x 0.5-0.75 μ) are borne on more or less abortive macrospores (32-40 x 1.5-2.0 μ). Similar conditions have been noted in this country.

Septoria tritici produces mucose, flesh color colonies, on potato-dextrose agar, which later become carbonaceous, finally subcottony.

Roberge, not Desmazières (153), should be credited with the naming of this species.

References: Baudyš and Picbauer (25); Beach (28); Cormier (130); Desmazières (153-155); Grove (295, 296); Luthra et al. (416, 417); Morini (473); Rivera (536); Rosella (542); Saccardo (555, 556); Sprague (642, 665); Unamuno (731); Weber (753).

SEPTORIA TRITICI f. AVENAE (Desm.) Sprague, Speckled Leaf Blotch.

At one time the writer (626) concluded that S. tritici on oats was nothing more than a physiologic race of the species, but further study disclosed that the form on oats is distinguishable by its more sinuous spores averaging slightly narrower than those on wheat. Forma avenae is limited to oats and does not attack wheat or any known grass. The form occurs on Avena sativa, A. byzantina (red oats), and A. fatua (wild oats) in Oregon and Washington, and in Europe, New Zealand, and Argentina. It is seldom abundant except during moist open weather in winter in the Willamette Valley and adjacent areas. Red oats and wild oats are particularly susceptible.

References: Desmazières (155); Sprague (626, 665).

SEPTORIA TRITICI f. HOLCI Sprague, Speckled Leaf Blotch.

Cross inoculation trials show that this fungus is confined to Holcus lanatus. It is common in wet weather in the winter and early spring in western Oregon and Washington. The spots are irregular, eventually umber color, with numerous pycnidia. The spores are long, filiform, 1- to 4-septate, 35-105 x 1.3-2.0 μ , and are narrower and longer than typical S. tritici on wheat.

Reference: Sprague (665).

SEPTORIA TRITICI var. LCLICOLA Sprague and A. G. Johnson, Speckled Leaf Blotch.

The lesions are at first green or yellowish but later turn fuscous to deep brown. The pycnidia are obscure because of the dark color of the older lesions. They are subglobose to globose, sometimes ellipsoid,

80-150 x 90-180 μ , mostly 120-150 x 120-150 μ in diameter. The spores are hyaline, sometimes with coarse protoplasm, sometimes containing small oil drops, which in slightly immature specimens are as large as 1 μ in diameter. The spores usually are slightly enlarged at the base, curved or sinuous, sometimes whip-like, 0- to 5-septate, commonly 1- to 3-septate, 21-85 x 1.3-2.8 μ (Sprague, 665, fig. 2). On Lolium perenne and less often on L. multiflorum, the spores are often more variable, with short, 1-septate spores occurring with the larger ones. They differ from those of S. tritici on wheat in the slightly more whip-like appearance due to the slightly enlarged bases and somewhat more tapering distal portions.

Septoria tritici var. lolicola occurs on L. perenne and L. multiflorum in western Oregon and in France. The early taxonomy of this fungus is greatly confused. Desmazières, no. 2169 in his *Fungi Cryptogames de France* (1851), described S. graminum var. lolii Mont., which now appears to be a smaller spored fungus, Ascochyta desmazieri Cav. (99). But var. lolicola was also present in this material, as well as Rhabdospora lolii Cast. (98). The latter fungus was transferred to Septoria under the name S. lolii by Saccardo (in *Sylloge Fungorum* v. 10), who failed to note S. lolii West. (763) which latter is the same as A. desmazieri. The fungus called R. lolii Castagne, according to rules of nomenclature, was nameless and hence was transferred to S. loligena by the writer. In all this welter of names, the filiform spored Septoria also escaped without a name and was therefore called S. tritici var. lolicola.

References: Castagne (98); Cavara (99); Desmazières (156); Sprague (665); Westendorp (763).

STAGONOSPORA SPP.

Key to species:

A. Spores over 8 μ wide (30-45 x 8-11 μ)
Stagonospora simplicior Sacc. and Berl.

AA. Spores narrower

B. Spores somewhat boat-shape, 20-35 x 6-8 μ
St. subseriata Desm.

BB. Spores more or less cylindrical

C. Spores somewhat fusiform, 15-30 x 2-3.2 μ

D. Spores somewhat truncated at base
St. agrostidis f. angusta Sprague

DD. Spores more fusiform than truncated St. bromi A. L. Sm. and Ramsb.

CC. Spores definitely cylindrical

D. Spores 25-70 x 2.5-5 μ St. arenaria Sacc.

DD. Spores 30-80 x 4-9 μ St. foliicola (Bres.) Bub.

This genus Stagonospora is separable from Septoria with difficulty, because on grasses there is a group of fungi, some placed in Septoria, some in Stagonospora, that have spores about 30-40-60 μ long x 3-4-6 μ wide and therefore lie on the border between filiform and cylindrical. Some workers consider Stagonospora as having coarse spores similar in all respects to Hendersonia, except for color, and similar also to Hendersonia in being mainly weak parasites or saprophytes (Kirschstein, 384). There are exceptions to this classification that make its value questionable. In general, the writer uses the length-width ratio of 10 to 1 as the division point between cylindrical and filiform. Since the length of the spores of many of these species range from about 9 to 1 to 13 to 1, they become difficult to classify. In such cases they are left in the genus to which they are customarily referred; e.g. Septoria nodorum, Stagonospora arenaria, Septoria avenae, and Stagonospora arrhenatheri A. L. Sm. and Ramsb. The forms with filiform spores produce yeasty masses of conidia on germinating on potato-dextrose agar while those with cylindrical spores are more likely to produce cottony mycelium and produce spores in true pycnidia. However, some produce small crustose colonies in which pycnidia are embedded in a matrix (Septoria infuscans), so that at least three groups are recognizable. If we place these intermediate forms in Septoria, the genus should be segregated into three sections as follows:

A. Spores filiform or narrowly obclavate, producing conidia from germinating spores on potato-dextrose agar.....Section Filisporae
..... (Various typical species of Septoria)

AA. Spores clavulate or coarser, producing pycnidia in crustose colonies on potato-dextrose agar..... Section Clavulosporae
(Species such as Septoria infuscans)

AAA. Spores narrowly cylindrical, producing spores in true pycnidia in pure culture on potato-dextrose agar.... Section Cylindrosporae
(Species such as Septoria nodorum, S. avenae, S. loligena, S. secalis, S. melicae, S. oudemansii and S. agropyrina)

STAGONOSPORA AGROSTIDIS f. ANGUSTA Sprague, Leaf Blotch.

This is an obscure fungus associated with other fungi on leaves of

Stipa viridula in western North Dakota (Sprague, 666). It has fusiform spores but the bases are characteristically sub-obtuse or truncated. The spores are small, 15-27 x 2.2-3.2 μ , borne in globose, golden-brown pycnidia, 120-280 μ in diameter. It is probably of slight importance economically. Morphologically it is closest to St. agrostidis Syd. (697, p. 4). The spores differ in shape from those of Septoria nodorum.

STAGONOSPORA ARENARIA Sacc., Purple-brown Blotch.

In well-developed but not old-weathered material, the spots or blotches are purple-brown, purple or nearly sepia or deep brown. Older material may be light straw color with tawny borders or old material may be completely faded to neutral straw color. The lesions are sometimes numerous, as on Fluminea festucae (Willd) Hitchc., or sometimes scattered and obscure, as on Elymus. The lesions do not retain their dark color very long but fade in the center, paling to the neutral straw color mentioned above. St. arenaria occurs on several species of Elymus and Agropyron and morphologically similar material occurs on Cinna latifolia (Trevir.) Griseb., Oryzopsis hymenoides, and Fluminea festucae in the western area. On Elymus and Agropyron, this fungus merges with Septoria nodorum, S. avenae, and S. agropyrina, as mentioned previously, and there exists some uncertainty as to where to draw species lines in this polymorphic group. Since the publication of an earlier article (Sprague, 653), it has been determined that the fungus on Phalaris is an immature or small-spored phase of St. foliicola (Sprague, 666; and below under St. foliicola). It is now apparent, also, that St. arrhenatheri should not be classed with St. arenaria but with Septoria avenae, from which even St. arenaria is doubtfully distinct on some hosts.

Host range studies are needed with this group in order to determine racial affinities of St. arenaria. It is no doubt a complex of races.

References: Sprague (653, 665, 666).

STAGONOSPORA BROMI A. L. Sm. and Ramsb., Purple-Brown Blotch.

This fungus causes a purple-brown blotch on the leaves of brome grasses. The disease is very destructive in Michigan (Hardison, 310) and the writer has collected it on Bromus carinatus in grass plots at Park River, North Dakota. He also recently found it on B. vulgaris (Hook.) Shear near Lake Itasca, Minnesota.

The fungus has 3-septate fusiform spores, mostly 17-24 x 3 μ . Its life history should be studied in comparison with Septoria nodorum, but it appears to be different. St. bromi is much more parasitic than St. agrostidis f. angusta and the spores are clearly hyaline and with less truncate bases.

References: Grove (296); Hardison (310).

STAGONOSPORA FOLIICOLA (Bres.) Bub., Tawny Blotch.
(St. vexata var. foliicola Bres., St. vexata var. baldingeriae Sacc.)

This is an active parasite on the leaves of Phalaris arundinacea in North Dakota, South Dakota, Iowa, Minnesota, and Wisconsin. It also occurs on P. californica Hook. and Arn. in Marin County, California. In field plots at Mandan, almost all of the leaves are covered with the tawny blotch in late season. Some difference in susceptibility was noted among plants in this nursery.

In the Mandan material, the spores are hyaline, multiseptate, and very large, 30-80 x 5-9 μ , obclavate-filiform to obclavate-cylindrical. (Sprague, 666). All of the material that has been seen previously was immature and resembled somewhat fusiform material of St. arenaria (Sprague, 653). From a study of developing material at Mandan during the summer months, it is now evident that these collections are clearly part of the life cycle of the coarse-spored St. foliicola. The inclusion of this material in St. intermixta (Cke.) Sacc. by J. J. Davis (144, p. 73), who had only the developing phase, had confused the writer in earlier studies. The fungus produces a white to pale pink, somewhat weak cottony growth on potato-dextrose agar.

STAGONOSPORA SIMPLICIOR Sacc. and Berl., Leaf Mold.

This species is, for the most part, a saprophytic leaf mold. The symptoms consist of vague stramineous or gray areas on dead leaves. Pycnidia are more or less gregarious, 100-140 μ in diameter in our material, black, globose, ostiolate; and the pycnospores are cylindrical or slightly bent, obtuse, slightly or scarcely constricted at the septa, 3-septate, with coarse contents, often one huge hyaline inclusion in each cell; spores are hyaline or faintly yellow, borne on blunt pycnophores, 28-35 x 8-9 μ on Calamovilfa longifolia in North Dakota as compared with 32-38 x 8-9 μ on Phragmites communis in the type from France. Our local material, collected in September on dead leaves, appears to be saprophytic. However, St. simplicior var. andropogonis Sacc. was described from North Dakota (Dr. J. F. Brenckle). Greene sent material of a fungus on Sorghastrum nutans from Madison, Wisconsin, that he determined as this variety, and it appears to be parasitic, causing elliptical spots on living leaves. St. simplicior is likely to prove rather widespread on coarse grasses in our region but, except for the Wisconsin material, it has not, as mentioned, been shown to be associated with living host parts.

STAGONOSPORA SUBSERIATA (Desm.) Sacc., Leaf Mold.

This species has boat-shaped spores $20-40 \times 4-7 \mu$ borne in pycnidia on various hosts in the western United States (Sprague, 652). It occurs on Poa pratensis and Phleum pratense in northern Minnesota. The species appears to be saprophytic on most of the hosts seen. St. subseriata var. maculata Grove (296) is an active parasite, however, on orchard grass in England (Grove, 296), Pennsylvania (Sprague, 652), and Maryland.

References: Grove (296); Sprague (652).

STAGONOSPORA SPP., Molds.

A considerable number of species of Stagonospora occur on dead culms, leaves, and sheaths of grasses. These are particularly common on coarse marsh grasses such as Phragmites communis. So many species have been described on this grass, for instance, that it is practically impossible for most workers, and that includes the writer, to determine these with available facilities. Since most or all of these fungi are saprophytes their interest is purely mycological and of no importance to the pathologist.

WOJNOWICIA GRAMINIS (McAlp.) Sacc. and D. Sacc., Secondary Rootrot.

This fungus was removed from Hendersonia because of short hairs on the neck of the pycnidium. As a diagnostic character this has doubtful merit.

W. graminis is an associate of Cercospora herpotrichoides in the prairies of the Columbia Basin but also occurs beyond this region into the dry land wheat regions of the West. It occurs commonly in Kansas, Colorado, and westward. It is not abundant in North Dakota in the spring wheat region, but is sometimes isolated from the roots or occurs on necrotic leaves of quack grass, orchard grass, and certain other species. It has been noted on about 29 species of Gramineae in the West.

Wojnowicia graminis, while scarcely parasitic, is of some economic importance because it weakens the culms of ripening grain, causing them to lodge prematurely. This condition is common in the Palouse in areas not affected by Cercospora herpotrichoides. The effect of weak parasites and saprophytes on lodging of ripened grain has not been given sufficient attention. Carroll (95), in Eire, discusses this problem to some extent.

The spores of W. graminis are yellow brown, slightly flattened on one side, curved spindleform, $24-38 \times 3-5 \mu$, usually 7-septate. An isolate from the roots of Agropyron repens in North Dakota had spores measuring

43-51 x 4.5-6 μ but otherwise typical.

References: Broadfoot (68); Carroll (95); Ludbrook (413); McKinney and Johnson (448); Sprague (630, 667).

2. MELANCONIALES

COLLETOTRICHUM GRAMINICOLUM (Cos.) G. W. Wils., Anthracose.
(C. cereale Manns)

Colletotrichum graminicolum is said to attack the roots, stems, leaves, and spikes of cereals and grasses, causing spots and injury generally known as anthracnose in which the black spots due to pustules (acervuli) produce a somewhat scab-like effect on the diseased surface. The fungus may develop saprophytically in late season with little evidence of its presence except for setae which give the diseased part a slightly moldy appearance.

In our area it occurs on Agropyron desertorum (Fisch.) Schult. A. repens, A. trachycaulum, Agrostis alba, Alopecurus aequalis Sobol., A. pratensis L., Andropogon furcatus, Anthoxanthum odoratum, Arrhenatherum elatius, Avena fatua, A. hookeri Scribn., A. sativa, Bromus purgans L., Calamagrostis inexpansa, C. rubescens Buckl., Dactylis glomerata, Digitaria sanguinalis, Festuca octoflora Walt., Hordeum vulgare, Lolium multiflorum, L. perenne, Oryzopsis micrantha (Trin. and Rupr.) Thurb., Phleum pratense, Poa pratensis, Schizachne purpurascens (Torr.) Swallen, Secale cereale, Sorghum vulgare var. sudanense (Piper) Hitchc., Sphenopholis obtusata, Sporobolus heterolepis, Trisetum cernuum, and Triticum aestivum. The fungus is particularly common on Anthoxanthum and Arrhenatherum in the coast region of Oregon. While common in Minnesota, it is less prevalent in the drier regions to the west.

The acervuli have prominent black setae 60-120 x 6-8 μ . The conidia are spindle-form to falcate, 18-26 x 3-4 μ , borne on short conidiophores, 12-6 x 1-2 μ .

Bolley (1913) reported Colletotrichum on the roots of wheat in North Dakota. The illustration indicates that he was dealing with a fungus that the writer has isolated in large numbers from diseased roots of many cereals and grasses. The cultures produce pink or orange slimy yeasty (mucose) masses of kidney-shape or falcate spores 4-8 x 1.4-2.5 μ . The colonies later develop black leathery wrinkled mycelium in which Phoma-like bodies sometimes occur, or less often there may be strands of pseudo-setae which never develop into true horny setae. Inoculations with a number of these isolates have been negative, although in one case an isolate obtained in August was parasitic on several hosts. Subsequent trials with this were negative and the end result in doubt. The writer has called these cultures Gloeosporium

spp. Whether they are a phase of C. graminicolum is doubtful. Certainly they are different in culture from "C. cereale" of Selby and Manns (589). Selby and Manns obtained true mycelium development on glucose agar without mucose development of spore masses. Setae were sparingly produced and the spores were falcate, not dissimilar to those on the host. The North Dakota Gloeosporium resembles in its development some strains of Colletotrichum gloeosporioides Penz. in pure culture (Burger, 83) although the spores of the latter are larger.

References: Böning (60); Bolley (56); Burger (83); Chowdhury (105); Fischer, et al. (231); Koehler (386); Sanford (576); Selby and Manns (589); Wilson (766); Winter (773).

GLOEOSPORIUM GRAMINUM Rostr., Anthracnose.

What answers to the description of this fungus was found on leaves of Poa pratensis in two localities in Benton County, Oregon, (Sprague, 637). It may be related to the Gloeosporium-like fungus reported under Colletotrichum graminicolum above.

PSEUDODISCOSIA AVENAE Sprague and A. G. Johnson, Red Leather Leaf

In western Oregon and Washington, entire fields of winter oats sometimes show a brilliant red color in February and March. In other cases, the disease is confined to small scattered patches in which the leaves nearly all show a general reddish or reddish-brown color. All but the innermost leaves of a plant may be affected. These lesions cause a leathery, fawn to red or brown appearance. An excess of red anthocyanin pigmentation in the uninjured portions of diseased leaves gives the reddish-brown color of affected plants. The diseased plants are stunted, compact, and rigid; and the leaves are slightly rolled.

This fungus is widespread on common and red oats in the vicinity of Corvallis, Oregon, in the winter and early spring and may also be found as far east as southwestern Klickitat County, Washington. The type locality is an isolated high prairie in the latter region near the Columbia River gorge. In this area the diseased plants of Winter Turf (Gray Winter) oats showed 40 percent reduction in yield over the healthy ones. The disease is of some importance, although the oats recover to a considerable extent as soon as warm weather arrives in the spring. "Red leather leaf" is almost unrecognized as a definite disease, even at Corvallis, the symptoms being confused with red leaf symptoms in general (Sprague, 644). Many specimens of this disease are mailed to the Oregon Agricultural Experiment Station. It occurs only on fall-seeded oats.

Recently Bremer (64) reported this fungus from Turkey where it appears to be indigenous.

Pseudodiscosia avenae produces the fusiform, basally ciliated spores on very short conidiophores compacted together on a poorly developed stroma in the epidermis. There is no apparent rupture of the leaf epidermis as in typical acervuli, and in assigning this originally to the Melanconiales, the describers considered the stroma as closer to the Melanconiales than to the Moniliales (Sprague and A. G. Johnson, 672). There still appears to be no better place to assign this species.

References: Bremer (64); Sprague (625, 628, 632, 641, 644); Sprague and A. G. Johnson (672).

SEPTOGLOEUM SP.

This fungus causes dark brown to black lesions on the pedicels and foliage of Festuca idahoensis Elmer near Trude, Idaho. The spores are large, stoutly club-shape to cylindrical with blunt bases and tapering apices, which are sometimes pointed. They are usually aseptate, hyaline, 44-53 x 4.6-5.6 μ . This fungus is distinct from any species known to the writer. Until final study can be made with prepared material the fungus is assigned to Septogloeum rather than Septoria.

Reference: Sprague (669).

SEPTOGLOEUM OXYSPORUM Sacc., Bomm., and Rouss., Blotch and Char Spot.

The young lesions on the leaves are tawny with yellow margins, circular at first but soon elliptical to elongate. On Arrhenatherum elatius the fungus rarely forms black stromata, but on other hosts the lesions caused by a similar if not identical fungus become covered with dull-black, charcoal-like streaks, 2-5 mm wide and often several times as long. While the charcoal spots range from nearly circular to elongate, they are typically elongate, tapering to a point at each end. The center of the lesion is often paler than the periphery, being pale gray or isabelline from accumulation of conidia in some instances. Sometimes pycnidia occur as small black dots at the margin of the lesion, but more often, if they are present, they are obscured by the charcoal-like stroma and are not visible to the naked eye.

Spores are borne in stromatic tissue in obscure acervuli or in pycnidia in the center of the lesions, conidia arising from hyaline, globular or cuspidate conidiophores in great, grayish masses; conidia yellow to subhyaline, fusoid, often flattened slightly on one side, subtruncate at the base, tapering to an obtusely pointed apex. The conidia are 0- to 3-septate, mostly 2-septate, 17-38 x 2.5-5.9 μ ; on Agrostis, they are 21-31 x 3.2-4.4 μ ; on Arrhenatherum, 24-33 x 3.5-4.7 μ ; on Calamagrostis, 22-27 x 3.5-4 μ ; on Distichlis, 20-25 x 4.3-5 μ with relatively short, thick spores ranging from fusoid to nearly cylindrical; and on Elymus and Agropyron, they are variable, 17-38 x 2.5-5.0 μ .

Pycnidia in the char spots are subglobose, brown, more or less imbedded in the creosote brown stromatic tissue, or separate at the margin of the lesions, 80-160 μ in diameter, with variable ostioles sometimes 60 μ in diameter. The pycnosporos are similar to the conidia but are hyaline. Associated perithecia in the stroma are globose and up to 250 μ in diameter and are sometimes filled with hyaline prosenchyma. The perithecial walls are composed of creosote-brown compacted polygonal cells. The asci consist of short fascicles without paraphyses but in material seen to date on Elymus condensatus none of it is mature enough to determine.

Synonyms:

Mastigosporium album var. athrix Eriks. Fungi par. scan. exsicc. Fasc. VIII, 351-400. 1891. (Bot. Centralbl. 47: 296-298. 1891).

Fusoma biseptatum Sacc. Krieger, W. Fungi sax. exsicc. Fasc. XIV, no. 683. 1891

F. triseptatum Sacc. Sylloge Fungorum v. 10. 1892

Fusarium osiliense Bres. and Vest. Vestergren, Verzeichniss nebst Diagnosen und kritische Bemerkungen zu meinem Exsiccatenwerke "Micromycetes rariores selecti." Fasc. VII-X. Botan. Notis. 1900: 27-44. 1900

Septogloeum athrix (Eriks.) Sprague (640)

The host range of Septogloeum oxysporum includes the following hosts listed by States: Agropyron spicatum (Wyo.); A. trachycaulum (N. Dak., S. Dak.); Agrostis hallii (Oreg.), A. alba (Wyo.); A. scabra (N. Dak.); Arrhenatherum elatius (Oreg.); Bromus ciliatus (Colo.); Calamagrostis inexpansa (N. Dak.); Distichlis stricta (Torr.) Rydb. (N. Dak.); Elymus condensatus (Calif., Wyo., Mont., Wash.); E. glaucus (Utah); E. triticoides (Calif.); Muhlenbergia asperifolia (N. Dak.).

This fungus is not of great importance but at times is epiphytotic on grasses in humid areas.

There is still reason to question whether true S. oxysporum that occurs on Arrhenatherum is the same species as the char spot fungus found on other hosts. Life history studies are certainly needed. The char spot fungus with its Septogloeum-like stage is associated with an ascomycete which is close, taxonomically, to Dothidella aristidae. The entire charspot-tarspot situation is still further confused when the discussion by Greene (289, pp. 86-87) is noted, regarding Davisiella elymina (J. J. Davis) Petr. and other fungi. Members of the group Davisiella are either conidial phases of Phyllachora spp., or are associated saprophytes or even parasites. They appear to form part of the same over all problems as does the char spot and Septogloeum complex. The writer agrees with Greene that likely more than one fungus is involved in the whole Middle Western and Far Western situation.

References: Bommer and Rousseau (57); Greene (289); Sprague (651).

SEPTOGLOEUM SPARTINAE (Ell. and Ev.) Wr.
(Fusarium spartinae Ell. and Ev.)

This fungus has not been seen by the writer. The sporodochia are pale-yellow, floccose; conidia oblong elliptical to oblong fusoid, 1-3 septate, 12-15 x 3-4 μ . Saccardo (558, v. 18, p. 675) lists it on Spartina stricta (= S. leiantha Benth.) from Pacific Grove, California. It is listed by Seymour (599) on S. alternifolia Loisel. var. glabra (Muhl.) Fern. (= S. alternifolia Loisel).

References: Ellis and Everhart (216, p. 14); Saccardo (558, v. 18, p. 674-675); Seymour (599); Weiss (757, p. 547); Wollenweber and Reinking (776, p. 336).

3. MONILIALES

ALTERNARIA SPP. (and STEMPHYLIUM SPP.), Black Molds.

Species of Alternaria occur as saprophytes on aerial parts of many plants. They are one of the components of seed mold of wheat, called "black germ" in Siberia, "puntatura" in Italy, "moucheture" in Morocco, and such names as "kernel smudge", "black point" (in part), seed blight or "durum blight" in the United States and Canada. A melanistic (black skin pigment) browning of wheat lemmas is termed "Alternaria blotch" by Johnson and Hagborg (374).

Almost all careful research on Alternaria molds on wheat shows that they are saprophytic or at most, scarcely parasitic. They are economically important in that their presence may cause dockage losses to the grower, particularly in North Dakota durum wheat used for macaroni (Brentzel, 65).

The taxonomy of this group is in some confusion. Generally A. tenuis Nees is recognized as the common mold on Gramineae while Bockmann (54), among others, includes A. peglionii Curzi and A. circinans (Berk. and Curt.) Bolle.⁴

A. tenuis is isolated frequently from dead roots of old grass and cereal plants and less often from roots of seedlings. Our trials do not show any reason to believe that the fungus is parasitic to any extent on germinating seeds (Sprague, 663). This was shown by E. C. Johnson (372) 32 years ago.

Because A. tenuis is often associated with more parasitic species, such as Helminthosporium sativum Pam., King, and Bakke, moldy wheat seed should be treated with New Improved Ceresan at 1/2 oz. per bushel,

⁴ The writer is indebted to David R. Sumstine, Carnegie Museum, Pittsburgh, 13, Pa., for determination of the cultures of Alternaria isolated from Gramineae.

and grass seed with New Improved Ceresan at 1 oz. or Semesan at 4 oz. per bushel.

References: Bockmann (54); Bolley (56); Brentzel (65); Christensen and Stakman (113); Curzi (134); Elliott (206); Groves and Skolko (298, 299); Hagborg (303); Hanson and Christensen (309); Henry (321); Johnson, E. C. (372); Johnson, T. and Hagborg (374); Machacek and Greaney (420); Peyronel (514); Rosella (543); Sallans (562, 563); Wiltshire (768); Ziling (780).

ANSATOSPORA BROMI (Sprague) Sprague, Blotch.
(Cercospora bromi Sprague)

Gray to sordid, elliptical or striate spots with broader dark brown or umber margins occur on Bromus rigidus Roth in two widely separated areas in Oregon and in southwestern Washington, and also on B. vulgaris in coastal Oregon.

This fungus is notable for the peculiar secondary conidia, which are attached and down-deflected from the first or second basal cell of the conidia. Evidence from study of material on B. vulgaris indicates that this species (A. bromi) may not be the cause of the spot on that host. On the basis of Newhall's study (481) of a disease of celery our fungus was placed in Newhall's newly created genus Ansatospora (Sprague, 666).

References: Newhall (481); Sprague (634, 655, 666).

CEPHALOSPORIUM ACREMONIUM Cda., Black-Bundle Disease.

This fungus occurs in the eastern part of the Great Plains and in California. It is thought to be the cause of a black fibrovascular bundle condition, which is accompanied by development of nubbins in corn, multiple ears at one node, barrenness and purple discoloration of the plant. The fungus is seed and seedling-borne. It was well described by Reddy and Holbert (530).

References: Corda (128); Harris (311); Koehler (385); Koehler and Holbert (388); Reddy and Holbert (530).

CERCOSPORA SPP.

This genus has been the subject of part-time investigations by Dr. Charles Chupp for many years. The reader is referred to his mimeographed article (114). One of our species, C. bromi, would on the basis of Chupp's suggestions to Newhall (481) belong in the genus Ansatospora Newhall to which we have referred it (Sprague, 666) (see above). The few species of Cercospora that occur on western grasses are briefly

mentioned.

References: Chupp (114); Newhall (481); Sprague (666).

CERCOSPORA AGROSTIDIS Atk., Leaf Spot.

This fungus forms broadly elliptical, very light brown lesions 3-5 mm long with broad dull red borders on leaves of Agrostis in Alabama. Some fragmentary material on Sphenopholis obtusata from North Dakota, sent by the writer to Dr. Chupp, was assigned to C. agrostidis. The conidia are hyaline to tinted, 1- to 7-septate, terete, straight or slightly curved, 10-60 x 2.5 μ .

References: Atkinson (20); Underwood and Earle (732, p. 141).

CERCOSPORA FUSIMACULANS Atk., Brown Blotch.

The lesions are amphigenous, light brown, bordered by dark brown, broadly fusoid or elliptical, frequently confluent, 3-4 mm long.

The conidiophores are epiphyllous, fasciculate, olive-reddish-brown, straight, sub-geniculate or nodulose, sparingly denticulate towards the apex, septate, 50-100 x 4-4.5 μ , bearing small, hyaline, 3- to 4-septate conidia that taper a little toward each end, 25-40 x 2 μ .

The rather obscure blotch symptoms have prevented this fungus from being recognized widely. It occurs on Panicum pacificum Hitchc. and Chase on the exposed slope of Mary's Peak, Oregon, (O.S.C. 9898), and occurs in the Great Plains on P. virgatum (N. Dak.), in Iowa on P. dichotomiflorum, and in Alabama on P. dichotomum L.

It is reported also on barnyard grass (Echinochloa crus-galli) from Iowa (Gilman and Archer, 262). In this connection, the writer recently sent to Dr. Chupp material of a spot on barnyard grass found along the Missouri River bottoms, near Mandan, North Dakota, (September, 1944). Dr. Chupp replied as follows: "I examined the Cercospora on Echinochloa crusgalli (L.) Beauv. which did not resemble the Davis type (of C. echinochloae) at all. For instance, C. echinochloae has more or less cylindric conidia, straight to strongly curved, 1-7 but mostly 2-3 septate, 3-5 x 20-55 μ . The conidiophores are hypophyllous, short, 14-5.5 x 10-25 μ . Your specimen has acicular to obclavate conidia, almost straight, multiseptate, 2-4 x 30-120 μ , and the conidiophores are amphigenous, 4-5 x 25-85 μ . In fact, according to my key, it meets the characters of C. sorghi E. and E. (Jour. Mycol. 3:15. 1887)." Dr. Chupp suggested that it would be important to determine if the fungus involved would attack sorghum.

References: Gilman and Archer (262); Underwood and Earle (732).

CERCOSPORA SEMINALIS Ell. and Ev., False Smut.

It is questionable whether this should be listed in a discussion of leaf spots because this particular fungus attacks the flowers of buffalo grass (Buchloë dactyloides (Nutt.) Engelm.), resulting in sterile seed balls containing the brown powder (conidia) of the fungus. The conidia are acrogenous, sub-hyaline, obclavate, granulose, 3- to 5-septate, 80-110 x 6-7 μ .

False smut is credited with being an important factor in reducing the yield of viable seed. Since re-grassing programs in some areas depend heavily on buffalo grass, false smut is important. Buffalo grass seed is usually pre-soaked before planting. Whether viable conidia of the fungus have any detrimental effect on the seedlings is not known, but no isolations have been made that indicate such is the case.

References: Ellis and Everhart (210).

CERCOSPORA SETARIAE Atk., Brown Leaf Spot.

This fungus produces a dark brown elliptical leaf spot with an indefinite pale border. It is reported on Setaria lutescens (Weigel) F. T. Hubb from Iowa. Tehon and Daniels (701) use the following key to distinguish three species of Cercospora on Setaria:

Conidiophores long, 50-120 μ

Spores 20-150 x 4-5 μ C. setariae Atk.

Spores 45 x 1.5 μ C. striaeformis Wint.

Conidiophores short, 17-40 μ C. setariicola Tehon and Daniels

References: Atkinson (20); Tehon and Daniels (701).

CERCOSPORA SETARIICOLA Tehon and Daniels, Brown-Bordered Leaf Spot.

The old spots are elongate-elliptic with a cinereous center and a sharply delimited brown border. The spores are 4- to 12-septate, 30-90 x 3-4 μ . The writer has collected this fungus on Setaria lutescens at McCanna, North Dakota.

References: Tehon and Daniels (701).

CERCOSPORA SORGHI Ell. and Ev., Leaf Spot.

This is reported as a leaf spot on Johnson grass (Sorghum halepense (L.) Pers.) from Nebraska. It is common on sorghums farther south. The spores are said to be hyaline, $70-80 \times 3 \mu$ (Ellis and Everhart, 209). (See also discussion under C. fusimaculans above).

CERCOSPORELLA Spp.

Until recent years, there were no species of Cercospora recognized on grasses. There are at present four known for the region. They may be distinguished by the following key:

- A. Attacks base of culm causing foot rot, spores $30-80 \times 1.5-3.5 \mu$
Cercospora herpotrichoides Fron
- AA. Attacks leaves causing leaf spot or scald
 - B. Causes an eyespot lesion on leaves of Poa pratensis, spores $45-90 \times 3.6-5 \mu$ C. poagens Sprague
 - BB. Causes a scald or non-eyespot type of leaf spot
 - C. Conidia subulate, $20-35 \times 2.5-4.3 \mu$
C. subulata Sprague
 - CC. Conidia filiform, $40-105 \times 1.5-3 \mu$ C. holci Sprague

CERCOSPORELLA HERPOTRICHOIDES Fron, Cercospora Footrot.

In the United States, the Cercospora footrot is restricted to some high prairies in the winter wheat area in the Columbia Basin, where the precipitation is 14 to 25 inches and the climate not severe.

Fawn-color to white elliptical lesions with brown borders form at the ground line on the stems in early spring. Stromatic mycelium eventually covers the base of the culms with black, charred-appearing hyphae. The culms break over at this point. Seriously diseased fields have characteristically pale green leaves and smaller heads than normal. Later, so much of the grain may be broken over and tangled together that harvesting is very difficult.

Winter wheat and wheat relatives (Triticum spp.) and also barley (Hordeum spp.) are susceptible, while rye is more resistant and oats highly so. Aegilops spp. (Sprague, 655, 656), crested wheatgrass, Agropyron inerme, A. riparium Scribn. and Sm., Bromus carinatus, B. inermis, B. tectorum, Koeleria cristata, Poa secunda Presl, and

Sitanion hystrix have all been found carrying the fungus.

Cercospora herpotrichoides is greatly favored by open warm wet winters, which induce excessive growth in winter wheat or barley.

Olympia barley, Rex, Hymar, or Holland wheat (High Prairie, Washington) are preferred over bearded Turkey types of wheat in areas where this footrot occurs. Short straw is preferred to tall. Seeding should be delayed in the fall for a few days past the optimum date to reduce the chances of excess vegetative growth. Calcium cyanamid, 100 lbs. per acre increased yield.

References: Adam (4); Foëx and Rosella (233, 234); Fron (240); Glynne (264-267); Gorter (280); Heald (314); Moritz and Bockmann (476); Oort (494); Saxby (584); Sprague (620, 622, 624, 631, 635, 643); Sprague and Fellows (671).

CERCOSPORELLA HOLCI Sprague, Buff Scald or Tan Leaf Spot.

This fungus is very common in western Oregon and Washington on Holcus lanatus. The spots are tan, yellow-bordered to darker buff or tawny, elliptical or extensive scald-like lesions on any or all leaves of a plant. They develop in mid- to late- winter or early spring. In severe cases the injury resembles frost scalding or snow injury.

Conidia (1- to 9-septate, 40-103 x 1.5-3 μ) are produced in great numbers in winter, but in spring the conidia may be entirely absent. Such late spring lesions have been called nonparasitic by early workers who had not collected the fungus in the cold wet winter weather. This fungus is probably one of the reasons why velvet grass is not a satisfactory crop for pastures in western Oregon.

Reference: Sprague (634, fig.1).

CERCOSPORELLA POAGENA Sprague, Cercospora Eyespot of Kentucky Bluegrass

An eyespot lesion caused by this fungus has appeared in two places in western Oregon on Kentucky bluegrass. The lesions are circular to sometimes elongate, brown with yellowish border, often straw color in the center and hence forming an eyespot type of injury.

The spores are hyaline, elongate, broadly filiform to obclavate with bases blunt, tapering, with the apices either abruptly acuminate or attenuate at the tips, 4- to 7-septate, 45-90 x 3.6-5 μ . It is known only from Forest Grove and East Corvallis, Oregon, and occurs in mid-early spring.

Reference: Sprague (657).

CERCOSPORELLA SUBULATA Sprague, Leaf Blast.

This fungus was originally described from a few gray to olive buff scald-like lesions on a herbarium sheet of a specimen of Melica subulata (Griseb.) Scribn. collected in the Ochoco National Forest in eastern Oregon. Since then it has been found on Deschampsia caespitosa (L.) Beauv. in the Montana mountains, in three places in western Oregon on Festuca rubra and on Melica bulbosa in the Big Horn Mts., Wyoming.

In pure culture on potato-dextrose agar an isolate from F. rubra produced a slow-growing fawn-color cottony growth that formed a few scattered conidia for a short time before staling.

The fungus is somewhat of a mystery as it appears to be both scarce and widespread, that is, it is never found in any abundance but has been found over a wide area.

The conidia have a characteristic, subulate shape with a whip-like distal portion. The conidia are 20-35 x 2.5-4.3 μ , mostly 28-33 x 2.8-3.5 μ . This fungus is now considered distinct from Cercospora (Sprague, 668, in ms.).

References: Sprague (634, fig. 2, A, 668).

CLADOSPORIUM HERBARUM Lk., Mold.

See Mycosphaerella tulasnei (Jancz.) Rothers, its ascigerous stage.

CURVULARIA SPP., Mold.

These fungi sometimes cause molds, seed rots, leaf spots, seedling blights, and black kernels. In general they are saprophytic molds on dead plant parts.

Curvularia Boed. (55) has 3- to 4-celled, brown, fusiform spores, typically curved or bent, with one or two of the central cells somewhat enlarged. The species are universal molds on countless kinds of plants and on debris in the soil. Because their classification is still in a state of flux they are considered here as a unit. Groves and Skolko (300) have presented the latest information on the group and the reader is referred to their paper. On the basis of the ratio of length to width of spore, almost all of our material from the Northern Great Plains falls in the species Curvularia geniculata (Tracy and Earle) Boed., to which we have been referring specimens for several years. The spores of this species are brown, 3- to usually 4-septate, and 24-46 x 8-14 μ in pure culture. Our material resembles an Australian species, which Hynes (348) has reported as C. ramosa (Bainier) Boed. The spores from one of Hynes' collections average 32.6 x 13 μ , and those from

another, $35.3 \times 13.3 \mu$. According to the method used by Groves and Skolko, both of these would belong to the "geniculata" group and close to C. geniculata, but they place C. ramosa as a synonym under C. inaequalis (Shear) Boed. C. inaequalis has proportionately somewhat stouter spores than C. geniculata. Groves and Skolko report C. inaequalis on seeds of grasses and certain other crops in Canada. As they state, the problem is by no means solved and needs further study. We prefer, for the present, to leave all of the "geniculata" material in C. geniculata although admitting that there are distinct differences in isolates.

Besides the great number of isolates of Curvularia geniculata obtained from the roots and seeds of many cereals and grasses in our area, we have isolated another species, confined largely to Echinochloa crus-galli. The spores are olive-green rather than brown, and are very short and thick, sometimes almost triangular in outline. These have been referred to C. trifolii (Kauff.) Boed., but need to be compared with several species of Helminthosporium for a possible earlier name. C. trifolii belongs in the 3-septate "lunata" group. C. lunata (Wakk.) Boed. itself is an exceedingly abundant mold in the tropics and probably occurs on rice and some weeds within the bounds of this report. It is a nuisance on rice grains in the southern United States.

Wernham and Kirby (762) reported Curvularia sp. as associated with a "going-out" or "melting-out" disease of Poa annua and Agrostis palustris (Metropolitan bent) during hot weather. This fungus grew best at 30°C and was inhibited if the temperature dropped below 25°C .

Bunting (81) did not find that C. lunata was conclusively parasitic on corn in the Gold Coast of Africa.

Hynes (345) found that there were 11 races of C. ramosa (Helminthosporium-M) associated with wheat in Australia. Hynes believed that this species or races of it were more virulent than either H. sativum or H. tetramera McK. (C. spicifera ?). Hynes' forms of C. ramosa differed considerably, one from another, in pure culture.

In the Northern Great Plains the writer (663) has found that most isolates of C. geniculata were nonparasitic on seedlings of many grasses and cereals; in a few cases, however, isolates were parasitic on certain grasses, but scarcely so on cereals. One culture, 367 B-6, isolated from seedlings of Agropyron sibiricum in mid-April, caused injury to roots of crested wheatgrass (A. cristatum), and moderate injury to sudan grass. Most isolates caused some injury, mostly pre-emergence killing, to proso millet and blue grama when added to the soil at seeding time. In general, however, C. geniculata appears to be a saprophyte on the roots, culms, leaves, and inflorescences of Gramineae and other plants in the northern United States.

In culture Curvularia geniculata produces a luxuriant dark gray to

nearly black, more or less stringy growth on potato dextrose agar. Other isolates are velvety black, similar to some isolates of H. sativum, and such isolates produce quantities of spores.

Cralley and Tullis (131), Matsuura (441), as well as Boedijn (55), report that C. lunata could cause a seedling blight of rice. Martyn (437) reported C. lunata on brown discolored palea of flowering heads of rice from British Guiana. He was able to cause the condition by inoculating with spores of C. lunata and incubating under humid conditions.

Nigam (482) isolated C. lunata from red spots on sorghum leaves and found that the fungus produced, on 2 percent rice agar, alternate zones of pink and white mycelium, pink in daylight and white in darkness. In continuous darkness the colonies were a dense, bottle-green color.

Sanitary measures to inhibit these molds, particularly in warmer climates, would appear to be difficult. However, such places as rice straw stacks (Martin 435) may be sources of abnormally heavy infection and can be eliminated. Martin and Altstatt (436) also recommend burning all weeds and volunteer rice around fields, keeping late seedings away from early ones so that spores are not spread during threshing of early crops, and keeping the grain and shocks dry to avoid heating.

When C. geniculata is very common on moldy grass seed, it might be desirable to treat with New Improved Ceresan 1/2 to 1 oz. per bushel depending on the size of the seed (a bushel of small seeds involves a great deal more surface than a bushel of wheat). Most of the writer's trials have not shown great benefit from treatment of range grass seeds except those infected with Helminthosporium sativum.

References: Bainer (21); Boedijn (55); Bunting (81,82); Cralley and Tullis (131); Groves and Skolko (300); Henry (321); Hynes (345, 348); Lefebvre and Johnson (400); Martin (435); Martin and Altstatt (436); Martyn (437); Mason (439); Matsuura (441); Nigam (482); Sprague (663); Tullis (718, 719); Wernham and Kirby (762).

FUSARIUM SPP.

The number of species of Fusarium that occur on Gramineae is not great, but the artificial system of classification in general vogue has multiplied the number of their subdivisions so as to represent a considerable array of fungi. Since some of these fungi, notably F. oxysporum Schlecht., are only weak parasites, it is more practical for the cereal and grass pathologist to recognize the species, and, in many instances, to ignore most of the subdivisions. Thus, in the case of such a complex as F. oxysporum, we prefer to use the all-inclusive F. oxysporum Schlecht. em. Snyder and Hansen, and, for all practical purposes, not explore the multiplicity of forms that are involved, inasmuch as this and certain other species are of comparatively little

consequence as parasites of cereals.

The student of Fusarium will naturally refer to Wollenweber's Fusarium autographice delineata (775), Wollenweber and Reinking's text (776), or the work of Raillo (518, 519). In North America, Snyder and Hansen (616) and Gordon (276, 277) have been followed. Gordon and the writer (278) have listed the common species of Fusarium isolated from Gramineae in the Northern Great Plains and this will serve as a guide to most of the species found in the plains country and westward. A key to these forms follows:

Key to Species of Fusarium on Western Grasses
(as grown on potato-dextrose agar in diffuse light at 70° F)

- A. Typically producing red, pink, or wine color mycelium (sometimes color is scanty or absent in group with pip-shape Sporotrichum microspores)
 - B. Producing at least a few Sporotrichum-type microspores
 - C. Macrospores 1- to 3-septate F. poae (Pk.) Wr.
 - CC. Macrospores 1- to 5-septate
F. sporotrichioides Sherb.
 - BB. No Sporotrichum-type microspores produced
 - C. Colony typically rosy pink; spore masses bright carrot color, often absent; macrospores somewhat acuminate at distal end, 5-septate average 45 x 4.1 μ , 7-septate 55 x 4.5 μ . (The common pink Fusarium on the roots of plants in the semi-arid Plains country)
F. scirpi var. acuminatum (Ell. and Ev.) Wr.
 - CC. Colony white and rose or pink; spore masses carrot color; spores very narrow, 3-septate average 34 x 3.1 μ , 5-septate average 50 x 3.7 μ . (Common on the heads of Gramineae in moister regions)
F. avenaceum (Fr.) Sacc.
 - CCC. Colonies bright carmine; spore masses orange red, often profuse; spores stout, not acuminate, 3-septate average 29 x 6.1 μ , 5-septate 38 x 6.6 μ , and 7-septate 46 x 7 μ . (Usually strongly parasitic)
F. culmorum (W. G. Sm.) Sacc.
 - CCCC. Colonies carmine or darker; conidial masses bright orange; spores longer than in F. culmorum, proportionately narrower, 3-septate average 41 x 4.3 μ , 5-septate average 51 x 4.9 μ , 7-septate 73 x 5.4 μ . (Strongly parasitic) F. graminearum Schw.

- CCCC. Colonies white and bright rose; spores small, scarcely
heeled, 1- to 3-septate, 1-septate average $16 \times 2.8 \mu$,
3-septate $25 \times 3 \mu$ F. nivale (Fr.) Ces.
- AA. Typically producing a peach-color, later buff to nearly brown
mycelium; macrospores typically acuminate on distal end
- B. Macroconidia 3- to 5-septate. (Common on grass roots)
F. equiseti (Cda.) Sacc.
- BB. Macroconidia 3- to 7-septate, slightly more acuminate at
apex than F. equiseti. (Appears to be much less common
than F. equiseti, and very similar)
F. scirpi Lamb. and Fautr.
- AAA. Typically producing white or bluish mycelia
- B. Microspores produced in chains; growth generally white on
potato-dextrose agar F. moniliforme Sheldon
(The spores of var. subglutinans are produced in heads or
false heads which are clusters of spores adhering temporar-
ily)
- BB. Microspores not produced in chains
- C. Colonies white, soon showing lavender, blue, purplish,
or less often pink or even isabelline in the depth of the
medium; mostly aseptate microspores, $7-10 \times 2.1-3 \mu$; term-
inal and intercalary chlamydospores numerous. (Very com-
mon in roots of grasses in the Plains country)
F. oxysporum Schlecht. em. Snyder and Hansen
- CC. Colonies becoming intensely blue in the depth of the
medium; spore masses dirty cream to brownish; macrospores
long cylindrical, fusiform, 3-septate $28-42 \times 4.1-6.2 \mu$,
5-septate mostly $42-51 \times 5-6.3 \mu$. (Saprophytic)
F. solani (Mart.) Appel and Wr.

This key is useful only in a general classification of the rootrot,
mold, and scab fungi on Gramineae but has some value in aiding workers
who still class all Fusarium fungi as "spp."

References: Bennett (39); Carrera (93,94); De Haan (146); Gordon (276,
277); Gordon and Sprague (278); Padwick (500); Raillo (518, 519); Sim-
monds (6C9); Snyder and Hansen (616); Wollenweber and Reinking (776).

FUSARIUM AVENACEUM (Fr.) Sacc., Mold and Stem Rot.

This species apparently prefers marine or moist climates. It sometimes occurs as a parasite on rust spores on grasses in the western United States during rainy periods. The writer has seen it on a stem canker of orchard grass in the Cascade Mountains, Oregon. It has not been found on any rootrot material from North Dakota, although it occurs not far north of there in Canada (Gordon, 275). Johnston and Greaney (375) present evidence to indicate that it is a weak parasite. The fungus is readily recognized by its narrow macrospores.

References: Bennett (32); Gordon (275); Gorlenko (279); Johnston and Greaney (375); Wollenweber and Reinking (776, fig. 2, p. 52).

FUSARIUM CULMORUM (W. G. Sm.) Sacc., Rootrot and Head Mold or Scab.

Fusarium culmorum causes preemergence killing, stunting, and brown rootrot of cereals and grasses. The bases of some plants such as oats are sometimes covered with pink spore masses. It sometimes causes pink head mold or scab, and may even cause a small red-bordered sheath and leaf spot on wheat in the early season, as noted in the Palouse area of Washington. While the fungus is scarce in the Northern Great Plains, it is very common from central Montana westward. In the coast region, it and crown rust (Puccinia coronata Cda.) make the growing of oats somewhat hazardous. It is important east of the Great Plains also. In the Pacific Northwest it is an associate of Cercospora herpochrysa and of Helminthosporium sativum (dry land rootrot). Ledingham (399) found that F. culmorum was not inhibited by H. sativum but that the latter was inhibited by F. culmorum. There is increasing evidence that the various components of "common rootrot" are not entirely compatible.

Fusarium culmorum rootrot is one of the causes of red leaf in oats in Oregon and Washington (Sprague, 644). This symptom is particularly apparent in the Willamette Valley and in the vicinity of Moro, Oregon. Inoculation trials show that isolates of this fungus from North Dakota and Oregon are destructive parasites, far more parasitic than F. scirpi var. acuminatum.

The name Fusarium culmorum has been a catch-all for any pink or carmine tinted fungus in the United States. Very often F. scirpi var. acuminatum has been called F. culmorum, and the name has been used even for the entirely distinct F. equiseti and F. oxysporum. This accounts for the tendency to consider F. culmorum as a quasi-parasite of slight importance in some areas. Quite to the contrary, it is essentially as parasitic as F. graminearum; and where it occurs in quantity, it is to be reckoned with as a major problem, as it is in Canada and in the Pacific Northwest.

References: Bennett (32); Blair (50); Broadfoot (68, 69); Foster and Henry (236); Greaney (283); Ledingham (399); Rose (541); Sadasivan (561); Samuel and Greaney (575); Shen (603); Simmonds (607, 608); Sprague (663); Sprague and Fellows (671); Tyner (722, 723); Vosbein (748); Walker (750).

FUSARIUM EQUISETI (Cda.) Sacc. and F. SCIRPI Lamb. and Fautr., Secondary Rootrots.

These two similar species belong in the Gibbosum group which have the distal end of the macrospores pinched or narrowed. F. equiseti represents about 50 percent of all Fusarium spp. isolated from roots of Gramineae in April to June in the Northern Great Plains and is exceeded in numbers during the summer only by F. oxysporum. It first produces a peach color or apricot-tinted, cottony colony that later turns buff or medium brown. Microspores and later macrospores are produced in considerable numbers when grown in north light on potato dextrose agar. Putty color masses of macrospores, often without pseudopionnotes, are formed later. Neither of these species appears to be parasitic on cereal seedlings in trials at Mandan, North Dakota. (Sprague, 663). Bennett (34), however, found English strains of Fusarium scirpi var. pallens F. T. Bennett that were mildly parasitic on wheat and barley, while Johnston and Greaney (375) found that F. equiseti was moderately parasitic on older cereal plants. The isolates from the Northern Great Plains usually do not even cause destructive seed rots although at times they may reduce stands of blue grama to some extent, as do many weak parasites and molds (Sprague, 663).

References: Bennett (34, 37); Gordon and Sprague (276); Johnston and Greaney (375); Sprague (658, 663).

FUSARIUM GRAMINEARUM Schw., Seedling Blight, Scab.
(Ascigerous stage, Gibberella zeae (Schw.) Petch)

This well known cause of scab in cereals has been the subject of intensive study in the Middle West. An outline of recent studies up to 1940 is given by Dickson (160).

Fusarium graminearum is especially prevalent in southern Minnesota (Hanson, 305, 307) and is important westward into eastern South Dakota and the Red River Valley in extreme eastern North Dakota on spring wheat and barley. It is isolated occasionally from the roots of grasses in North Dakota west of the region where it occurs on cereals. It has been obtained from roots of Agropyron subsecundum, A. cristatum, and Agrostis sp. (P. I. 131,533) in western North Dakota, and from leaves of Poa pratensis near Jackson Hole, Wyoming, collected during very rainy weather in August 1941. It was isolated from barnyard grass and Hordeum nodosum growing in a moist ditch at Newell, South Dakota. It has not been reported authentically from the Far West. All cases attributed to

it in western Oregon have proved to be F. culmorum. Connors (122) indicates that it is rare in Manitoba.

References: Bennett (33); Connors (122); Dickson (159-161); Dickson et al., (162); Eide (200); Goddard (269); Hanson (305, 307); Henry (322); Ho and Melhus (330); Immer and Christensen (352); MacInnes (422); McIndoe (444); Melhus (459); Pearson (507); Senn (598); Sprague (663); Tu (714, 715).

FUSARIUM MONILIFORME Sheldon, Corn Ear Mold, Seed Rot.
(Ascigerous stage, Gibberella fujikuroi (Saw.) Wr.)

While this is a common mold on corn ears in the South and Middle West, it is somewhat less abundant in the Northern Great Plains, although Edwards (195) found it to be common on corn seed from Nebraska, and Hoppe (339) reported it as abundant even into Colorado. Boyle (62) reported that Ray isolated it from corn stalks from Pasco, Washington. It has been isolated by the writer from the roots of corn, sorghums, millets, and a number of grasses during the summer time in North Dakota. Our inoculation results indicate that it is a weak seedling parasite on small seeded grasses.

Recently Edwards (197), working with F. moniliforme and F. moniliforme var. subglutinans Wr. and Reinking, found that because the corn seed contained sufficient nutrient for independent seedling growth for 10 to 12 days, soil fertility did not influence the action of the parasite on the plant during this critical period.

References: Boyle (62); Dickson (160); Edwards 195-198); Ho (327); Hoppe (339); Koehler (385); Koehler and Dungan (387); Koehler and Holbert (388); Leonian (402); Sprague (663); Ullstrup (726, 728).

FUSARIUM NIVALE (Fr.) Ces., Snow Mold.
(Ascigerous stage, Calonectria graminicola (Berk. and Br.) Wr.)

In the United States, the status of snow mold due to Fusarium spp. has been in considerable confusion. It is said to be a common cause of turf injury in the Far West and from Minnesota eastward. Just how much actually is due to F. nivale is not known for certain. Fusarium snow mold is very common in Oregon during open rainy winter weather. Small, more or less circular areas, with a fringe of white or pinkish mass of mycelium and spores, are sometimes very conspicuous in bent grass lawns (Agrostis tenuis, A. palustris). Rhizoctonia solani and Fusarium culmorum, among other fungi, are prominent, but a species of Fusarium with a preponderance of 1-septate heel-less spores is most common, and this, at least, is referable to F. nivale but culture work and critical study are certainly needed.

A similar fungus with small falcate spores occurs on overwintering, lax leaves of Bromus tectorum, B. carinatus, B. rigidus Roth, Hordeum murinum L., and H. nodosum in the Columbia Gorge region of Oregon and Washington.

It is not unlikely that Fusarium nivale has been confused with F. oxysporum, although the species seem to develop at somewhat different seasons of the year in Oregon and Washington. In the eastern United States, recent work has shown that part of the snow mold is caused by Typhula itoana.

What is called Fusarium nivale occurs on Agrostis alba, A. palustris, A. tenuis, B. commutatus, Festuca rubra, F. rubra var. commutata, Poa annua and P. pratensis in Oregon; on A. palustris in Washington; and on Hordeum vulgare, Secale cereale, Agrostis alba, A. canina, A. tenuis, A. palustris and Festuca rubra in Minnesota. F. nivale var. majus Wr. (Wollenweber and Reinking, 776, p. 43, fig. 2, a) has been observed on winter leaves of Agrostis palustris in Lincoln County, Oregon.

References: Andren (12, 13); Bennett (35); Dahl (136); Dennis (151); Markvich (434); Monteith and Dahl (472); Nilsson (483); Sprague (655, 656); Wernham (761); Wollenweber and Reinking (776).

FUSARIUM OXYSPORUM (Schlecht.) emended Snyder and Hansen, Secondary Rootrot and Seed Rot.

While this fungus is very common on the roots of Gramineae, particularly during the summer, it does not appear to be very parasitic. The writer obtained moderate seed rot in some small-seeded grasses such as blue grama and in proso millet but on cereals the fungus was scarcely parasitic (Sprague, 663). Johnston and Greaney (375) obtained the same results on cereals. Simmonds and Ledingham (610) report F. oxysporum as nonparasitic on wheat and barley and slightly parasitic on oats.

The pink coloration in dead roots of oats and to some extent barley and wheat is largely due to a saprophytic development of F. oxysporum (Sprague, 663).

References: Gordon (276); Gordon and Sprague (278); Johnston and Greaney (375); Simmonds and Ledingham (610); Snyder and Hansen (616).

FUSARIUM POAE (Pk.) Wr., Silver Top, Secondary Rootrot, False Scab.

Poa pratensis is affected with a disease called silver top. The heads turn white and die prematurely. They are covered with a pale buff mold growth of Fusarium poae, associated with a species of mite (Pediculopsis graminum (Reuter) in some cases. Heald (313) showed a similar association of Fusarium and mite in bud rot of carnation, while Stewart and

Hodgkins (686) reported it on carnation and blue grass.

F. poae occurs on a head blight of oats in the coast region of Oregon. Wollenweber determined an isolate from this material as F. poae (Sprague, 636). The culture on potato dextrose agar was buff-colored and produced many pip-shape Sporotrichum-type spores. The fungus appeared to be a secondary parasite and was usually found on sterile spikelets.

In the Northern Great Plains, F. poae is not abundant but it is sometimes isolated from the roots of Bouteloua gracilis, Bromus inermis, Panicum capillare L., P. miliaceum, and Zea mays. It is more prevalent in Minnesota, particularly northern Minnesota, on Poa pratensis, P. palustris, and Muhlenbergia racemosa. It was found associated with wooly aphids on smooth brome sheaths at Mandan, North Dakota. It occurs on timothy in Iowa (Gilman and Archer, 262).

The material in pure culture at Mandan developed white to powdery buff colonies that soon showed rosy tints. When the Sporotrichum stage is present in abundance the mycelium has a readily recognized loose powdery appearance.

Our inoculation trials show that some isolates of Fusarium poae are comparatively weak parasites or saprophytes. A few strains cause some seed rot in blue grama and millets. F. poae appears to be a relatively minor mold on Gramineae.

References: Gilman and Archer (262); Gordon and Sprague (278); Heald (313); Sprague (636, 663); Stewart and Hodgkins (686).

FUSARIUM SCIRPI VAR. ACUMINATUM (Ell. and Ev.) Wr., Pink Fusarium Mold, Secondary Rootrot.

This common pink mold causes yellow-brown lesions on the roots of some grasses and cereals but is usually associated with other fungi without any specific symptoms. It is a common saprophyte on practically all Gramineae in the western United States but causes some seed rot in spring and fall on small-seeded crops. It is only a weak parasite on wheat and barley and is apparently nonparasitic on oats.

This variety differs from Fusarium scirpi in the rose-pink, firm colony that is produced on potato dextrose agar. The mycelium tends to be coarse and breaks into characteristic stiff fragments in microscope mounts. It often produces no spores, although some isolates form carrot-red masses of macrospores slightly smaller than those of F. scirpi. Five-septate macrospores average $45 \times 4.1 \mu$, and 7-septate, $55 \times 4.3 \mu$. The curved spores have the typical Gibbosum shape, narrow at the apex. This fungus is readily distinguished from F. culmorum and F. graminearum by its firm felty growth with rose-pink coloration instead of the carmine and yellow-buff looser mycelium of the two scab-forming species,

F. culmorum and F. graminearum. F. scirpi var. acuminatum is the common pink Fusarium of the prairie country. Less common are F. scirpi var. compactum Wr. and F. scirpi var. longipes (Wr. and Reinking) Wr., which Gordon determined as being present in North Dakota (Gordon and Sprague, 278). The former is distinguished by closely spaced septa in the macroconidia, and the latter by its exceptionally long macroconidia.

F. scirpi var. acuminatum apparently is favored by cool wet weather in March and April when it develops on the roots of overwintering perennial grasses and on young seedlings and sprouting seeds. It is a very common saprophyte following Cercospora herpotrichoides in the Pacific Northwest, and is one of the components of dry land rootrot of winter wheat in the Great Plains. It also occurs in acid soils in the peat area of northern Minnesota and westward, and is common in soil of relatively high alkalinity. Cultures of this fungus isolated from red soil in Wyoming were particularly pale in color and evidently were a local race. Sometimes other cultures from various areas have shown marked buff colorations but these colorations are frequently induced by bacterial associates in the cultures. The writer's inoculation trials (663) have shown considerable variation in racial traits among the isolates, some being mildly to moderately parasitic, others saprophytic or essentially so.

References: Fellows (229); Gordon (276); Gordon and Sprague (278); Sprague (663).

FUSARIUM SPOROTRICHIOIDES Sherb., Secondary Rootrot, False Scab.

This fungus is readily confused with Fusarium poae. It has the same general host range and produces Sporotrichum-type microspores. The microspores of F. poae are 0- to 1-septate, citron shape or pip shape, mostly $6.6-14 \times 4.7-6.4 \mu$, while those of F. sporotrichioides are usually longer, $5-28 \times 2.2-3.8 \mu$. Macrospores of F. poae are 3-septate, $18-35 \times 3.5-5 \mu$; in F. sporotrichioides they are 3- 5-septate, $25-56 \times 3.4-4.5 \mu$.

Fusarium sporotrichioides is sometimes isolated early in the season from roots of Gramineae in the Northern Great Plains. It occurs on the roots of blue grama, side oats grama (Bouteloua curtipendula), smooth brome, Elymus junceus, Festuca rubra, Hordeum jubatum, Panicum capillare, and P. miliaceum in the area. It disappears rather completely during the growing season, but reappears as a pink mold or false scab on wheats and barley at or near harvest time. It is a minor component of "durum blight" but appears to be saprophytic in these instances. In general, tests by the writer (Sprague, 663) indicate that F. sporotrichioides is even less parasitic than F. poae. Its chief importance lies in the fact that growers are sometimes docked for "scabby" wheat when the grain is carrying only the relatively harmless false scab organism, F. sporotrichioides. F. poae appears to be more abundant in Manitoba than F.

sporotrichioides (Gordon, 275, 276), but in North Dakota the reverse has been true during recent years. F. poae is more common in north-eastern North Dakota and in northern Minnesota, thus indicating relationship to the Manitoba situation.

References: Gordon (276); Gordon and Sprague (278); Sprague (663); Wollenweber and Reinking (776).

HELMINTHOSPORIUM

This complex and important genus was well presented about 20 years ago by Drechsler (176) although his work was limited to the Eastern States. At present the genus is being re-studied in that area and to some extent in the Plains region. The work with rootrots in the Northern Great Plains (Sprague, 663), has disclosed a number of forms that need critical taxonomic study.

HELMINTHOSPORIUM AVENAE Eidam, Helminthosporium Leaf Blotch of Oats (Leaf-stripe disease of oats in Scotland (Dennis, 149)).

In summer, in the Northern Great Plains, the spots on oat leaves may be poorly defined brown areas merging into yellowish red or orange lesions. In winter or in cool weather, in Oregon, they are more frequently definite spots of red or purple anthocyanin shades surrounded by haloes of yellow. As they get older, the spots fade to straw color in the center. Dennis (149) reported pre-emergence killing in seedlings from seed-borne mycelia or spores in Scotland. Survivors showed leaf stippling. Later, spotting developed, often secondarily in late season, particularly on second growth oats.

The fungus requires cool moist growing conditions such as prevail in the winter, early spring, and fall in the coast region of Oregon, Washington, and in parts of the British Isles. In Oregon, H. avenae was isolated from the bases of diseased plants but appeared to have washed down the culm and developed as an associate of H. sativum, Ophiobolus graminis, Fusarium culmorum, Pythium spp., and Aphanomyces sp., which thrived in the highly acid soil of the coastal region where the diseased condition prevailed (Sprague, 639). H. avenae is somewhat lighter gray in pure culture than most isolates of H. sativum and it tends to form bracketed growths in test tube cultures, somewhat distinct from any formed by H. sativum. The spores are very different also; they are olivaceous or paler, cylindrical or slightly swollen in the middle, rounded at the ends, 4- to 6-septate, 80-110 x 15-16 μ , borne on scattered, stout cylindrical, multiseptate, fuliginous conidiophores, 150-200 x 9-12 μ .

While this fungus causes a minor leaf spot in many places, it is very serious in Scotland and Ireland and moderately important in western

Oregon. Seed treatment with New Improved Ceresan at 1/2 oz. per bushel is very necessary where the disease is serious. Varietal resistance was not noted in Scotland although the varieties Sovereign and Ascot were particularly susceptible. In Oregon, winter-hardy oats such as Support or Gray Winter are more tolerant to the disease than less winter-hardy ones such as Red Rustproof or Kanota of the red oat group (Avena byzantina). Wild oats (Avena fatua) carries the disease in Oregon, California, and North Dakota; and it occurs also on A. barbata Brot. at San Diego, California, and on Arrhenatherum elatius on Mt. Diablo, California.

References: Dennis (149, 150); Drechsler (176); Eidam (199); Ito and Kuribayashi (354); Muskett (479); O'Brien and Dennis (491); Rathsclag (526); Sprague (639).

HELMINTHOSPORIUM BROMI Died., Leaf Spot of Brome.
(Ascigerous stage, Pyrenophora bromi (Died.) Drechs.)

Minute dark brown or black spots with a yellowish halo appear on the first leaves in early spring in the northern United States. The spots elongate, become as much as 2 x 6 mm in size and, through coalescence, may cover much of a leaf area. The leaves then become yellow, wither, and die. The fungus is favored by cool weather, with temperatures ranging from 2° C to 10° C, and becomes less conspicuous as the season advances. It is very common on Bromus inermis in North Dakota, particularly in the Red River Valley (Weniger, 760). It occurs also, more or less incidentally, on the roots of this grass in the Northern Great Plains and has been collected also on leaves of B. carinatus at Wawawai, Washington; on B. japonicus Thurb. at Alzada, Montana, and nearby in Wyoming; on B. tectorum near Underwood, Washington; and on B. rigidus near Tumwater, Oregon.

The perithecial stage, Pyrenophora bromi, is well known (Drechsler, 176). Chamberlain and Allison (100, 101) recently reported that P. bromi was an important parasite on brome in Wisconsin during cool wet spring weather. The perithecia are considered to be more important in overwintering the fungus than the conidial stage, which latter does not live over winter. Varietal resistance was noted.

References: Chamberlain and Allison (100, 101); Diedicke (163); Drechsler (176, pl. 8, 9); Weniger (760)

HELMINTHOSPORIUM CALIFORNICUM Mackie and Paxton, Spot Blotch.

The writer (639) believes that this is a form of H. sativum, similar to if not identical with material studied by the writer in Lincoln County, Oregon. Mackie and Paxton described the fungus as a leaf spot parasite of barley in central California (Mackie and Paxton, 427; Mackie, 424). In recent years the name has disappeared from current literature.

HELMINTHOSPORIUM CYCLOPS Drechs., Leaf Spot.

The spots on Danthonia californica in the Willamette Valley, Oregon, are scattered, dark brown or black, and small, usually less than 0.5 x 2 mm in diameter.

Drechsler (176, pp. 729-730, pl. 32, 33) described this fungus on D. spicata (L.) Beauv. from Lisbon Falls, Maine. The Oregon collections (Sprague, 655) represent the only other report of this fungus as far as our records show. Neither the fungus nor its host is of very much economic importance.

HELMINTHOSPORIUM CYNODONTIS Marig., Leaf Mold.

Forms an effuse olive to fuliginous blotch on dry leaves of Bermuda grass (Cynodon dactylon) in the Bay region of California and southward. Drechsler (176, pp. 719-720, pl. 27) reports it also on Eleusine indica (L.) Gaertn. and Muhlenbergia mexicana in Florida and near Washington, D. C., and into New York state. It is practically omnipresent in the South but outside of disfiguring lawns it is not believed to cause much damage. Material seen at Davis Junction, California, in 1939 would indicate that it might be of more importance.

HELMINTHOSPORIUM DEMATIOIDEUM Bub. and Wróbl., Leaf Mold.

This fungus causes a leaf mold of sweet vernalgrass (Anthoxanthum odoratum L.) in northwestern Oregon. Its parasitic nature is not known. Drechsler (176, pp. 683-685, pl. 14) reported it on Agrostis alba and A. perennans (Walt.) Tuckerm. from the Atlantic coast.

HELMINTHOSPORIUM DICTYOIDES Drechs., Fescue Net Blotch.

A leaf net blotch characterized by a minute but well defined reticulate design on brown discolorations. The lesions are numerous and extensive. The leaves eventually wither and die, usually progressively from the leaf tips.

This fungus appears to be of some importance in the Far West. It was collected at Bingen, Washington, on Festuca occidentalis Hook. and recently a trace of it was found in some material of chewings fescue from northwestern Oregon.

Drechsler (174, 176, p. 677-679, pl. 11) reported it as the most important parasite of meadow fescue (F. elatior) in the eastern United States, where it is so common that it serves as a ready means of identifying the host in its vegetative stage.

HELMINTHOSPORIUM ERYTHROSPILUM Drechs., Red Leaf Spot.

This species has not been reported from the area covered by this report, but is likely to occur in the eastern part. The report of H. stenacrum Drechs. from Iowa possibly should be rechecked with this fungus. H. erythrospilum causes a true leaf spot on Agrostis, not a mold or streak as does H. stenacrum. During moist weather, it forms numerous straw-color leaf spots with a reddish-brown border, 0.5-2.5 mm. During prolonged periods of rainy weather the spots more or less overlap, forming pseudozonate patterns. They are surrounded by a water-soaked area, which presents a halo-spot effect. After dry weather returns the leaves wither and brown.

Reference: Drechsler (184).

HELMINTHOSPORIUM GIGANTEUM Heald and Wolf, Zonate Eyespot.

This fungus was described from Texas. While it is generally confined to the southern part of the United States it occurs as far north as Michigan and Illinois. It has not been reported in the range of this article but it is likely to occur. The fungus is noted for its huge fragile conidia and for the destructive zonate leaf spot (eyespot) that it produces on many grasses, notably Bermuda grass.

References: Drechsler (176, 180, 182); Heald and Wolf (319).

HELMINTHOSPORIUM GRAMINEUM Rab., Barley Stripe.

The symptoms of this important disease are detailed by Drechsler (176). Previous to heading time, the leaves become streaked with yellow bands that sometimes traverse the entire length of the leaf. The streaks later become yellowish brown or brown and the plants die prematurely. The leaves eventually become ragged or shredded. Heads may be completely sterile and the undeveloped kernels more or less brown.

Barley stripe is not important west of the Red River Valley except in certain areas in California and scattered spots in western Oregon, and sometimes in the Palouse region of Washington and Idaho.

The use of New Improved Ceresan, 1/2 oz. per bushel, has reduced the seriousness of the disease. Infection of the seed is favored by high humidity at and following flowering time. The disease may be carried over as mycelium penetrating from the glumes into the seed coat, or it may be transmitted by conidia on the seed, or wind-borne. Infection of the seedling occurs at sprouting time and is favored by cold, wet soil.

Christensen and Graham (112) found that H. gramineum was polymorphic and multiracial. They found about 125 races showing some morphological

differences in some cases and many variations in cultural characters. About 20 races could be distinguished by their relative virulence on 16 varieties of barley.

Shands (600) found that H. gramineum was somewhat adaptable to varying temperature conditions. Barley will grow at cool temperatures (12-16° C) as will H. gramineum, but the latter also developed well at cooler temperatures and also at 20° C.

The following varieties have been more or less resistant to stripe: Glabron, Trebi, Spartan, and Wisconsin barbles.

References: Åberg (2); Arny (16); Christensen and Graham (112); Dickson (160); Drechsler (176, p. 650-656); Johnson (373); Leukel, Dickson and Johnson (404); Majdrakoff (430); Misikado (486); Paxton (506); Reddy and Burnett (529); Shands (600); Shands and Arny (601); Smith (110); Suneson and Santoni (693); Weniger (760).

HELMINTHOSPORIUM HADROTRICHOIDES Ell. and Ev., Mold.

Medium gray, linear streaks followed by withered, sometimes shredded leaves or vague spots on dead leaves. Material, probably saprophytic, found by the writer on leaves of Eragrostis cilianensis (All.) Lk. from Miles City, Montana, was determined by C. L. Lefebvre as H. hadrotrichoides. We also isolated the same fungus from the roots of this grass growing in sandy soil at Mandan, North Dakota.

References: Drechsler (176, pp. 710-711, pl. 23); Ellis and Everhart (211); Mitra and Mehta (470).

HELMINTHOSPORIUM HALODES Drechs., Mold.

This was originally described on Distichlis spicata (L.) Greene from the Atlantic coast, but it appears to be a fairly common saprophytic mold, causing secondary seed rot or possibly rootrot in Azopyron repens and Muhlenbergia japonica Steud. in plots at Mandan, North Dakota, and in Panicum miliaceum, Poa spp., Schedonnardus paniculatus (Nutt.) Trel., Setaria italica (L.) Beauv. and Setaria viridis in North Dakota, South Dakota, and Minnesota. The symptoms produced by H. halodes are vague, dark discolored, more or less bluish areas on the leaves, sheaths and culms, also a black mold on the inflorescence. The species is not recognized as being associated with any particular rootrot symptoms.

In culture the dark spores are somewhat longer and less "kneed" than those of Curvularia geniculata although some strains are confusable, when immature, with the 4-septate phase of this species.

Mitra (468, p. 287) described H. halodes var. tritici Mitra on

root-rotted wheat from India. The conidia are cylindrical or elliptical with the distal end abruptly rounded and the proximal end tending to taper to an acute base with a prominent, protruding hilum. The spores are widest at their middle, straight or slightly curved, 2- to 9-septate (average 6), light yellowish brown to smoky brown, the two end cells being slightly lighter, $23-73 \times 13-20 \mu$, mean $50 \times 16.5 \mu$. The description resembles some of the material isolated from roots of various grasses at Mandan, North Dakota.

Most of the isolates of H. halodes have been saprophytes or have produced mild seed rots (Sprague, 663).

References: Drechsler (176, pp. 707-709, pl. 22, 23); Mitra (468); Sprague (663).

HELMINTHOSPORIUM INCONSPICUUM Var. BUCHLOËS Ell. and Ev., Mold.

Material that C. L. Lefebvre refers to this fungus causes a mold on withered leaves of Buchloë dactyloides, Bouteloua gracilis, and B. curtipendula in the Northern Great Plains. The mold is associated with vague, brown blotches and is sometimes isolated from roots of older plants.

Pammel, King, and Bakke (502) illustrated H. inconspicuum var. buchloës as having small, heavy-walled spores. Technically the fungus apparently is undescribed (nomen nudum). Nothing is known about its economic importance.

References: Pammel, King, and Bakke (502).

HELMINTHOSPORIUM MONOCERAS Drechs., Spotblotch.

Forms dark brown or chocolate color spots, $0.3-1 \times 1-3 \text{ mm}$, on the upper leaf blades. The lower leaves become withered and are covered with similar, larger elliptical spots, sometimes $1.5 \times 5 \text{ mm}$. The older spots on the dead leaves are faded to a dull medium brown. The leaf sheaths and the tissue at the base of the plants are a diffuse brown. Conidiophores appear after the death of tissue.

This fungus occurs on barnyard grass (Echinochloa crusgalli), but a similar fungus has sometimes been isolated from the roots of Panicum miliaceum and of Setaria viridis in the Northern Great Plains. (See also H. setariae Saw.).

The spores are yellowish when young but later darken to the same color as those of H. sativum, thus contrasting with the much paler ones of H. turcicum Pass. The spores are $40-150 \times 15-22 \mu$, typically straight, and usually widest at the middle segment, gradually tapering toward the tip to

1/3 or 1/2 of the maximum width, then bluntly rounded, tapering towards the base to approximately 1/6 of the median diameter, the contours then curving gently into the protruding hilum, 3- to 10-septate, scarcely constricted.

Schweinfurth and de Thuemen (588) list H. flexuosum Cda. (Brachysporium flexuosum (Cda.) Sacc. on the culms and inflorescence of Echinochloa crusgalli in Egypt. Possibly this is the same fungus as Curvularia trifolii mentioned earlier in this article.

An undetermined species of Helminthosporium is listed from the vicinity of Sydney, Australia, on E. crusgalli (Noble et al., 488).

References: Drechsler (176, p. 704-707, pl. 20, 21); Noble et al. (488); Schweinfurth and de von Thuemen (588).

HELMINTHOSPORIUM POAE Baudyš, Mold.

Somewhat fragmentary material on Poa trivialis L. from Astoria, Oregon, (O.S.C. 10,330) and on Poa secunda from Klickitat County, Washington (O.S.C. 4) did not appear to belong in H. vagans Drechs. and was assigned with considerable hesitation to H. poae (Baudyš, 24), a little known species described from central Europe. According to a footnote by Drechsler (176, p. 688) this fungus has 2- to 6-septate spores, 36 to 73 μ long. The spots are yellowish with a dark brown margin. Drechsler compares his H. vagans not only with H. poae but also with Napicladium gramineum Pk., which latter is a synonym of Scolecotrichum graminis Fekl. (Sprague, 657). H. poae certainly needs critical study. Possibly it too may have some well known earlier name.

References: Baudyš (24); Drechsler (176, p. 688); Sprague (637, 655-657).

HELMINTHOSPORIUM SATIVUM Pam., King, and Bakke, Spotblotch, Rootrot, Seedling Blight, Black Point, and Dryland Rootrot.

The seedling phase of the disease is characterized by a slow, dark rot of the coleoptile region with injury spreading later to the cortex and small feeder roots. The seedlings of small grasses may be wiped out early or even killed before emergence (pre-emergence blight), but the injury is less spectacular than that caused by Pythium arrhenomanes. Such hosts as barley, which show stunting or killing of lower leaves, very often recover later with slight apparent ill effects if growing conditions are optimum for the host.

After the seedling phases have been passed, leaf spotting may occur. This is a brown blotch rather than a definite spot or stripe. Abundant spotting in late season on barley or wheat speeds up premature ripening of the grain. The fruiting bodies of the fungus may give the maturing

plants a blotchy or olivaceous cast. Barley seed and glumes show dark brown stains that are accompanied by shriveling of the kernels. The basal portions of kernels are more likely to be discolored. Black point of wheat and other grains is partly due to H. sativum, and most of the injury that occurs at seeding time in the Northern Great Plains from seed-borne parasites is due to this fungus.

In winter wheat, injury due to Helminthosporium sativum is vague, particularly on wheat grown in the semi-arid regions. The more or less recognized term, "dryland rootrot" (or footrot), is used to designate a particularly vague type of injury in which the plants show pronounced stunting in more or less circular to irregular areas in the fields. The crowns of these plants, when bisected with a sharp knife, uniformly show a conical area of brown, prematurely dead or necrotic tissue. These areas of greatest injury are frequently associated with shallow soil, hard pan under layers, or other conditions unfavorable for the growth of the plant. In the dry, winter wheat regions, occasional white heads (sterile heads) develop in winter wheat showing dryland rootrot. These are particularly common in eastern Oregon and are a phase of premature killing of plants by the almost omnipresent H. sativum.

Helminthosporium sativum occurs on many hosts, but it is most important as a rootrot and kernel smudge or blight of wheat and barley. It has been reported on dozens of species of grasses as well as on oats, corn, and wheat relatives. In the Northern Great Plains, it has been isolated from the roots of 79 species of Gramineae. It is especially common on species of Elymus, Agropyron, Bromus, and Hordeum, and less abundant on species of Poa, which are resistant to some extent. The host range as established by seedling and seed inoculation trials in the greenhouse is very extensive.

H. sativum is world-wide in distribution in the temperate zone. It occurs in very acid soil (pH 4.8) in a region of very heavy rainfall (70-100 inches yearly) in Oregon, and is omnipresent in the arid and semi-arid parts of the western United States. While it is an important parasite of wheat and barley, it takes second place as a parasite of grass roots when compared with Pythium arrhenomanes. Its importance on grasses may have been somewhat exaggerated in the past. However, because it is so universally distributed, it cannot be dismissed as a minor parasite even on grasses.

As a component of dryland rootrot of fall-sown wheat, it is not considered as important as it is on spring grain. Its effect is, nevertheless, important in spite of its obscure nature.

Helminthosporium sativum has been studied in great detail by a number of workers. Christensen has shown that it possesses a large number of physiologic races; also, it is generally recognized as being readily influenced by environment. Changes in light or moisture, or the presence or absence of antagonists can change a single isolate to almost

unrecognizably different sub-cultures. While H. sativum is able to thrive in cool and moist weather, often it causes greatest injury in drought periods that follow. The diseased plants have weakened root systems and are unable to supply the plant with water and nutrients as readily as do healthy plants with larger root systems.

Seed treatment (New Improved Ceresan, 1/2 oz. per bushel) is effective insofar as it destroys seed-borne H. sativum on cereals but soil-borne spores and mycelium are scarcely affected by seed treatment.

References: Andrews (14); Beckwith (30); Bolley (56); Brentzel (65, 66); Broadfoot (69, 71, 72); Broadfoot and Tyner (74, 75); Christensen (106-109); Christensen and Davies (110, 111); Christensen and Stakman (113); Dosdall (172); Dosdall and Christensen (173); Drechsler (176, pp. 690-704, pl. 17-19); Greaney (284); Greaney and Bailey (285); Greaney and Machacek (286); Greaney, Machacek and Johnston (287); Greaney and Wallace (288); Gries (292); Hanson and Christensen (308, 309); Henry (321, 322); Hynes (345, 348, 349); Johnson (372); Kuribayashi (394); Ledingham (399); Machacek (418); Machacek and Greaney (421); McKinney (445, 446); Mead (453, 454); Mitra (467, 468); O'Gara (492); Pammel, King, and Bakke (502); Pinck and Allison (516); Richardson (535); Russell (550); Russell and Ledingham (551); Russell and Sallans (552); Sallans (563-567); Sanford (578); Sanford and Broadfoot (580); Sanford and Cormack (581); Simmonds, Russell, and Sallans (611); Simmonds and Sallans (612, 613); Smith (615); Sprague (639, 654, 655-661); Stakman (679); Stevens (683, 684); Stover (689); Tropova (712); Tyner and Broadfoot (724); Valleau (734); Weniger (760); Ziling (780).

HELMINTHOSPORIUM SETARIAE Saw., Spotblotch and Secondary Rootrot.

(H. setariae Lind)

(Ascigerous stage, Ophiobolus setariae Ito and Kuribay.).

This species is sometimes isolated from the roots of Setaria viridis and less frequently from S. italica in North Dakota. In Japan (Ito and Kuribayashi, 354) the fungus produces abundant dark brown spots or leaf blight. In North Dakota such spotting is less common but in the late season it frequently occurs, mixed with spots caused by H. sativum.

In the description of Ophiobolus setariae Ito and Kuribayashi (354), the conidial stage, H. setariae, is listed as having dark olivaceous spores that are fusiform, obclavate ellipsoidal, mostly somewhat curved, slightly broader at or somewhat below the middle, tapering towards both ends, 5- to 10-septate, 40-120 x 10-18 μ , with fragile walls.

Pure cultures resemble H. sativum macroscopically but the spores are narrower than those of H. sativum. A. G. Johnson determined New Jersey material (Haenseler, 302) as belonging to H. setariae. It was some leaf spot material on Setaria italica, collected at Beltsville, Maryland, sent by Dr. Johnson that permitted determination of Mandan and North

Dakota material as H. setariae, and not as H. turcicum to which the writer had been inclined to assign it. Inoculation trials at Mandan show that H. setariae, when added as a sand-bran inoculum to electrically sterilized soil, is somewhat pathogenic to Setaria spp., Echinochloa crusgalli, Panicum miliaceum, and Bouteloua gracilis, but is less parasitic on Triticum aestivum and Agropyron cristatum, and only moderately parasitic to Sorghum vulgare L. Indications at present are that the fungus is not important in North Dakota.

References: Elveden (218); Haenseler (302); Huang (342); Ito (353); Ito and Kuribayashi (354); Lind (408); Sawada (583).

HELMINTHOSPORIUM SICCANS Drechs., Brown Blight.

The leaf blight of Lolium perenne and L. multiflorum caused by this fungus is very common in the Willamette Valley, Oregon, in winter and in early spring. It is particularly prominent on uncut lawns, where the leaves in certain areas are nearly all killed back to their bases. The leaf spots are numerous, dark brown, measuring 0.1-0.3 x 0.2-1 mm, and so abundant that much of the leaf surface may be covered with a pseudo-reticulate scorching or browning.

The conidia of H. siccans are never dark olivaceous, as in H. sativum, but are subhyaline or light fuliginous, later becoming yellow, brownish, or brownish olivaceous. The spores are typically straight or slightly curved, usually subcylindrical or tapering slightly or markedly toward the apex, 35-130 x 14-20 μ .

H. siccans is one of the most important diseases of L. multiflorum in the eastern United States (Drechsler, 176, pp. 679-682, pl. 12). Diedicke (164) reported a species of Helminthosporium on L. perenne. H. siccans occurs also in Scotland (Dennis and Foister, 152, p. 274) and England (Sampson and Western, 571).

HELMINTHOSPORIUM STENACRUM Drechs., Leaf Mold.

The symptoms are indefinite and the conidiophores develop on withered leaves. Oregon material on Agrostis tenuis appears to be saprophytic. H. stenacrum is reported on A. alba from Iowa.

The spores in this species are subhyaline to yellowish, 53-135 x 15-23 μ , subcylindrical with hemispherical or hemi-ellipsoidal ends, or widest somewhat below the middle and tapering moderately towards the ends. The apical portion of the spore is sometimes narrowed and prolonged. The spores are 1- to 11-septate, scarcely or not constricted at the septa, the peripheral wall is thin and the dark hilum is included within its contour (Drechsler, 176, pp. 682-683). Usually it will not sporulate on nutrient agar.

HELMINTHOSPORIUM TERES Sacc., Net Blotch.
(Ascigerous stage, Pyrenophora teres (Died.) Drechs.)

The dark brown spots or streaks on barley leaves are about 1 mm long, at first, finally 20 to 25 mm long but narrow. Some streaks of brown pigment occur transversely that break the pattern into a dark brown reticulate design within the area of diffused brown, hence the term "net blotch". The blotch tends to form near the leaf tips. These blotches may sometimes occur on the hulls.

Net blotch usually is a minor disease; although widespread it is seldom serious. It is not common in the Far West, and is most frequent in the northern United States east of the Great Plains.

The spores of H. teres are similar to those of H. avenae, subhyaline at first, later greenish fuliginous or yellowish brown but lighter than in H. gramineum, thin-walled, with a retracted hilum, 1- to 10-septate, 30-175 x 15-22 μ .

Partial control is obtained by treating seed with New Improved Ceresan, 1/2 oz. per bushel, or by reducing the number of overwintering perithecia by sanitation methods.

References: Atanasoff and Johnson (19); Böning and Wallner (59); Drechsler (176, pp. 656-663, pl. 2,3); Johnson (367); Ravn (527).

HELMINTHOSPORIUM TETRAMERA McKinney, Rootrot and Mold.

McKinney (446) described this species as the cause of a mild rootrot of wheat in Texas. Conidiophores are dark olivaceous to brown, simple or compound, with septa 5 to 50 μ apart and with the conidia produced at irregular intervals. Conidia are chiefly 4-celled, borne in clusters of 2 to 3 to 50 or more, dark olivaceous to brown, usually rather symmetrical in shape, tapering towards the rounded ends, 20-41 x 10.2-20.4 μ , frequently 30-34 x 10.2-13.6 μ .

The writer has isolated a similar fungus from cereals and grasses in North Dakota, South Dakota, Montana, Wyoming, and Nebraska. The spores on potato-dextrose agar are dark olive smoky, relatively thin-walled, regularly elongate, elliptical, 3-septate, 23-31 x 8-12 μ , borne on simple or sparsely branched conidiophores. A number of attempts to produce rootrot lesions or even seed rot on various Gramineae with this fungus have been unsuccessful. The local fungus appears to be Brachycladium spiciferum according to a letter received from J. E. Machacek in 1941. B. spiciferum was transferred to Curvularia spicifera (Bainier) Boed., but without much justification as the regular elliptical spores are not in the least Curvularia-like. The fungus has small spores but could be placed in Helminthosporium or Brachysporium,

according to Machacek. He believes that B. spiciferum differs from H. tetramera. The spores of the former may be smaller than in H. tetramera, and the non-parasitic nature of the former may indicate specific differences between the two. In our case we question whether H. tetramera proper has been found in the Northern Great Plains.

Brachycladium spiciferum is usually isolated from overwintering roots and crowns of perennial grasses and from roots and sprouting seeds of grasses and cereals in the early season and late fall. It produces a luxuriant dark gray-black colony on potato-dextrose agar. Spores are produced in moderate quantities but seldom in large numbers. McKinney stated that H. tetramera produced long, simple, or branched sclerotia composed of hard white pseudoparenchyma with an outer black layer or rind. Many hyphal strands developed from these rinds, but no conidia were found on them. However, McKinney did have conidia in pure culture, as he stated earlier, and there seemed to be little if any change in spore characters when the fungus was cultured artificially. Since B. spiciferum produces a readily growing gray-black cottony colony that resembles Alternaria tenuis Nees and shows none of the sclerotial development of McKinney's cultures, the two forms are possibly distinct. Machacek mentioned in a letter to the writer (May 2, 1941) that he had seen material on oats and barley from eastern Canada that is closer to H. tetramera than to B. spiciferum, although, as he points out, Hynes (348) has referred Australian material to C. spicifera while Mason has referred it to H. tetramera. The tendency during recent years appears to be to lump the two species together. While no critical taxonomic report is available, the group had been under study by Machacek, until current economic conditions interfered. Crosier and Weimer (132), who also had the aid of Machacek, concluded that C. spicifera was a saprophyte, although they did obtain mild seedling injury on Poa pratensis, P. compressa, P. trivialis, and Agrostis alba (A. palustris). Machacek points out in his letter to the writer that in Crosier and Weimer's paper the legends on the illustrations of C. spicifera and C. gemiculata are transposed. Thomas (706) also failed to obtain injury with H. tetramera (C. spicifera). In 1937 (Anonymous, 1), B. spiciferum was reported as an associate of R. solani in a brown patch of lawns in New South Wales. Bensaude (40) obtained consistently negative results on wheat in Portuguese East Africa with a fungus also referred to H. tetramera. Some further study with the Texas material is indicated as being desirable.

References: Anonymous (1); Bensaude (40); Crosier and Weimer (132); Hynes (345, 348); McKinney (446); Thomas (706).

HELMINTHOSPORIUM TRISEPTATUM Drechs., LEAF MOLD.

This causes a gray mold on dead or withering leaves of Holcus lanatus in the humid parts of coastal Oregon and Washington. It was also found on moldy leaves of Agrostis exarata at Wells (now Camp Adair), Oregon.

In New York, Horsfall (349) reported leaf tip killing on Agrostis alba and orchard grass. Drechsler points out the contrast between the vague lesions caused by this fungus on red top (A. alba) and the definite leaf spots caused by H. erythrospilum (Drechsler, 184).

The spores of H. triseptatum are dark, olivaceous, ellipsoid or short cylindrical with hemispherical ends, 2- to 3-septate, 35-50 x 15-21 μ . The peculiar thickenings of the conidiophores, as shown by Drechsler, are diagnostic (Drechsler, 176, pp. 685-686, pl. 14).

HELMINTHOSPORIUM TRITICI-REPENTIS Died., Leaf Blight.
(Ascigerous stage, Pyrenophora tritici-repentis (Died.) Drechs.

Drechsler (176, pp. 667-670, pl. 5-6) describes the symptoms on Agropyron repens as being vague, manifested in a gradual fading and necrosis of the leaves, which finally become gray and heavily covered with conidiophores and later with perithecia, particularly on dead culms. In Oregon, definite brown to fuliginous spots and streaks occur on the leaves of A. repens and these are followed later by general necrosis. On Elymus hirsutus Presl the dark brown spots were very abundant on all the leaves of some plants along the Santiam River, Oregon, in May.

In the Northern Great Plains saprophytic development is sometimes great on wheat leaves during rainy periods in later season; however, little evidence has been seen of parasitic activity in this region. On wheat the fungus is usually associated with Septoria nodorum and other fungi. Fungi similar to H. tritici-repentis are isolated occasionally from the roots of cereals and grasses in the Northern Great Plains. Isolates from the roots of crested wheatgrass cause preemergence injury in crested wheatgrass, blue grama, proso, Italian millet, Japanese millet, and slight rootrot in wheat (Sprague, annual report, manuscript, 1944).

The conidia are subhyaline, straight cylindrical, 1- to 9-septate, 45-175 x 12-21 μ . Drechsler (176) states that "the most distinctive peculiarity....is found in the shape of the basal segment which....in profile is remotely suggestive of the horizontal aspects of the head of a snake, while the distal end usually is rounded off in hemispherical form."

References: Barrus (22); Connors (121); Drechsler (176); Johnson (369); McRae (452); Mitra (469); Nisikado (486); Raabe (517).

HELMINTHOSPORIUM TURCICUM Pass., Leaf Blast, Seed Rot.
(H. inconspicuum Cke. and Ell.)

A gray-green leaf blight of corn was seen in some experimental plots

in Lincoln County, Oregon, some years ago, and it occurs also on Sorghum vulgare var. sudanense in California. H. turcicum is otherwise unreported west of the Rockies. Chilton (104) reported H. turcicum on seeds of sudan grass from South Dakota, while the writer has isolated the fungus from the roots of corn and sorghum in North Dakota at various times.

The conidia are typically subhyaline to yellowish brown, very thin-walled, straight or slightly curved, widest near the middle and tapering decidedly towards the ends, somewhat cone-shaped at the hilum end, with the apical end rounded off but not beaked, 1- to 8-septate, 45-140 x 15-25 μ . The hilum scar is minute but it is distinctly protruded. The fungus develops a luxuriant gray-black aerial mycelium on potato-dextrose agar.

This fungus depends on high humidity, long rainy periods, and a maturing host for its most severe action. Because it develops in late season, its effect is often more spectacular than serious. Sherbakoff and Mayer (604) reported a black ear rot of corn as due to H. turcicum in Tennessee, but no doubt it is H. carbonum (Ullstrup, 729). Chilton has shown that H. turcicum is seed-borne on sudan grass and potentially destructive. The disease is considered to be serious in the Tropics and it interferes with seed production in the United States, particularly in sweet corn plantings in the East. Its importance on the roots of corn, sorghum, sudan grass, proso millet, and species of Setaria in North Dakota is slight. The identity of these forms remains in doubt. Some of them are atypical for H. turcicum.

Sanitary measures that reduce inoculum help to control this disease. Ducomet (187) recommended wider spacing to increase aeration. Seed treatment with New Improved Semesan Jr. kills seed-borne spores. Recent work by Ullstrup (727-729) evidences the complexity of the Helminthosporium problem on corn.

References: Campi (90); Chilton (104); Drechsler (176, pp. 712-718, pl. 24, 25); Ducomet (187); Ellett (203); Elliott and Jenkins (205); Gentner (260); Lefebvre and Sherwin (401); Mitra (466); Nisikado (485); Nisikado and Miyake (487); Reinking (531); Sherbakoff and Mayer (604); Ullstrup (727-729); Valteau (734); Zhavoronkov (799).

HELMINTHOSPORIUM VAGANS Drechs., Bluegrass Leaf Spot.

The leaf spots are circular to elongate with prominent reddish-brown borders darkening to nearly black in the center and, later, some becoming lighter in the center. Occasionally the spots are prevalent, particularly near the crown, but sometimes they are scattered over the upper leaves.

The disease occurs on Poa compressa but is most abundant on Poa pratensis. It also occurs on Poa arida Vasey in plots at Mandan, North

Dakota. It is seldom serious in the Pacific Northwest, and even less so in most of the Northern Great Plains, but appears to be definitely more important from Minnesota and Iowa eastward.

Since H. vagans is favored by rainy weather it develops most actively during late winter and early spring in the Pacific Northwest and during late summer and fall in the Northern Great Plains. At Mandan, North Dakota, moderate injury occurs during September or even October.

Horsfall (340) found that copper-lime dust inhibited this fungus.

References: Drechsler (175, 176, p. 686-688, pl. 15); Horsfall (340); Sampson and Western (571); Sprague (657).

HELMINTHOSPORIUM SPP. on Various Hosts.

In addition to the instances cited a number of collections were either too fragmentary or too recently seen to justify cataloguing. Most of these collections have been forwarded to A. G. Johnson and C. L. Lefebvre for study.

The following are included in undetermined material:

Helminthosporium (catenarium Drechs.) on Beckmannia syzigachne (Steud.) Fern., from West Fargo, North Dakota, June 20, 1944. Mingled with various leaf molds and Physoderma sp.

Helminthosporium sp. on Sporobolus neglectus Nash, Mandan, North Dakota, September 4, 1944.

Helminthosporium sp. on Gastridium ventricosum (Gouan) Schinz and Thell., a fragment from the southern part of the Willamette Valley, Oregon, showing a small circular white leaf spot.

Helminthosporium sp., a leaf blotch on Muhlenbergia asperifolia from Ft. Totten, North Dakota.

The isolates from the roots of warm season hosts including Panicum spp., Setaria spp., sorghums, and corn need further study. No doubt most of the isolations from Setaria that are not assignable to H. sativum belong in H. setariae, but some few are close to H. turcicum. Most of these forms are distinctly less parasitic than H. sativum.

HETEROSPORIUM AVENAE Oud., Leaf Mold.

This mold is not uncommon on overwintering plants. While it is usually saprophytic, the writer has seen material on Avena sativa from Oregon;

on Elymus condensatus from Oregon and Washington, and from Jackson, Wyoming, and on Phalaris arundinacea from Nebraska; that appeared to show evidence of parasitic activity on the part of Heterosporium avenae. It is usually found in the early season. Conidiophores of this fungus on western material resemble those of Scolecotrichum graminis Fekl., but the cylindrical spores are echinulate.

Solheim (617) reported Heterosporium avenae on leaves, stems, and spikes of wheat from Laramie, Wyoming, collected in August 1930. The writer noted it on leaves of overwintering wheat in Klickitat County, Washington, some years ago. A similar species was also seen on leaves of Bromus carinatus from Mt. Hood, Oregon.

HETEROSPORIUM PHLEI Greg., Leaf Spot.

The spots on Phleum pratense are small, oval, 0.5 to 1 mm wide and 1-4 mm long, rarely larger. The center of the spot is light mauve, with a narrow madder-violet or dark nigrosin-violet margin fading to brown or sometimes intensifying to black. Sometimes intervening tissue becomes yellow, and severely infected leaves may die. The scattered, sharply defined spots usually distinguish this parasite from other spots on timothy. Spores are rarely produced in the field. The conidia are 1- to 6-celled, chiefly 1-septate, not constricted at the septa, echinulate, pale to dark brown, 13-57 x 6-14 μ , averaging 23 x 9 μ .

Jacques (359) found that the fungus grew rapidly on potato-dextrose agar. The colony was bordered by a white margin, surrounding the olive-green center in the outer portion of which four or five zones of different shades of green could be recognized. A certain amount of white aerial mycelium gave the center a grayish tinge. The medium showed a reddish or purplish coloration for some distance around the margin of the colony.

In the original description, Gregory (291) reported that the fungus grew slowly in pure culture and did not sporulate until a thallus of considerable size was formed. Jacques (359), working with this strain, found that the fungus produced spores abundantly in pure culture after only 4-5 days when the thallus was only 12 mm in diameter.

Jacques stated that Heterosporium phlei is possibly merely a physiologic race of some earlier-described species, such as H. avenae Oud., H. graminis McAlp., H. hordei Bub. or H. phragmitis Sacc.

Heterosporium phlei occurs on timothy in Oregon, Washington, and Idaho to a limited extent, particularly in prairie areas and along the coast where the rainfall is moderate to heavy. It is more prevalent in Minnesota and appears to be important at times in New York and in the Eastern States in general. The spores are able to germinate at from 3° to 33° C, with the optimum at 24° C. Since the disease has

been noted in February, March, July, and August, the fungus is evidently adaptable to a wide range of temperature. H. phlei also occurs on Phleum phleoides (L.) Karst. at Pullman, Washington.

Horsfall (340) indicated that relatively satisfactory control could be obtained with Kolodust and somewhat less control with 300-mesh dusting sulfur. In most instances it would not pay to dust for this leaf spot alone.

References: Gregory (291); Horsfall (340, fig. 18, 19); Jacques (359).

MASTIGOSPORIUM SPP.

Key to species:

A. Spores with apical appendages M. album Riess
(Not reported from U. S.)

AA. Spores muticate, 3-septate

B. Spores 25-32 x 4.5-5.9 μ M. cylindricum Sprague

BB. Spores 29-90 x 9-17 μ
M. rubricosum (Dearn. and Barth.) Sprague

Most keys include only Mastigosporium album, which is readily traceable by its simple or forked, elongate, hyaline apical appendages, and for anyone not familiar with the large clear hyaline, 3-septate, cylindrical spores of M. rubricosum its determination is likely to be difficult with any available keys.

MASTIGOSPORIUM CYLINDRICUM Sprague, Brown Spot.

This very distinct species is known only from a restricted forested area in the Coast range of Oregon, on Bromus vulgaris. The brown spots are elliptical to elongate, finally confluent and mottled. The mycelium is mostly endophytic, somewhat coalesced beneath the upper leaf surface, coarse, hyaline or lightly tinted, producing short, stout conidiophores from which spores are developed by expansion of the distal portion and eventual abscission.

Reference: Sprague (648, fig. 1,b).

MASTIGOSPORIUM RUBRICOSUM (Dearn. and Barth.) Sprague, Eye Spot.
(M. calvum (Ell. and Ev.) Sprague).

On Dactylis glomerata, dark purple brown flecks are formed, which enlarge and become elliptical spots with ashy-gray to fawn color centers, 1-8 mm long, somewhat restricted by the parallel leaf veins. Larger lesions show various shades of gray, ashy, or light fawn color with more or less purple, red, or ochre borders.

On species of Agrostis, the lesions usually are broader, elliptical, light brown with fawn centers, becoming an eyespot or frog-eye spot with a broad red, or red and yellow margin.

On Calamagrostis and Trisetum the lesions are the same as on Agrostis, or are small brown flecks, 1- to 3-mm in diameter, with small gray centers, or are sometimes surrounded by fawn or pale buff-color tissue.

The large navicular, crystal hyaline, 3-septate conidia are formed by expansion of the hyphal tips. They differ from M. album in the complete absence of the simple or forked aseptate filiform apical appendages.

In the humid parts of Oregon, Washington, and Alaska, this is one of the most serious leaf spots, being particularly destructive to orchard grass, red top, and creeping bent (Agrostis palustris), and in the mountains it is very abundant on Calamagrostis canadensis.

This fungus develops during open winter weather in Oregon and Washington, and during rainy or foggy weather throughout the growing season. High humidity and dense stands favor its development. In Oregon it occurs commonly on lawns in late winter.

The Oregon fungus from Agrostis alba was found to be a distinct race from that on Dactylis glomerata (Sprague 640, 648). There also are differences in susceptibility of several species of Agrostis to the fungus (Sprague, 640).

References: Bondarzeva-Monteverde (58); Dennis and Foister (152, p. 276; Sampson and Western (570); Sprague (640, 648).

NAPICLADIUM ARUNDINACEUM (Cda.) Sacc., Black Leaf Mold.

This black leaf mold of Phragmites communis has been reported from Nebraska, and material from Pelican Rapids, Minnesota, and Devils Lake, North Dakota, collected by the writer, appears to be this species. Large streaks or coalesced areas form on the leaf tips of growing plants, sometimes apparently killing considerable tissue. Yellow or necrotic areas on some leaves indicate that the fungus possibly may be somewhat systemic. There does not appear to be very much written on

this fungus. It is evidently not uncommon in marshy areas in Minnesota and North Dakota.

NIGROSPORA ORYZAE (Berk. and Br.) Petch, Dry Rot of Corn.
(Basisporium gallarum Moll.)

This is mainly a cob rot of corn. The cob tends to shred at the base or shank and sometimes the kernels are somewhat shriveled. Infected ears show black spores around the area where the ear was broken from the shank or at the base of kernels. Yellow kernels are often bleached or discolored yellow brown. The fungus may attack injured seed and rot it in the soil if the soil remains cold and wet after planting.

References: Durrell (191); Koehler (385); Mason (438); Reddy (528); Savulescu and Rayss (582, 582,a); Standen (680, 681).

CVULARIA HORDEI (Cav.) Sprague, Leaf Spot.
(Ophiocladium hordei Cav.)

In Europe this fungus causes an obscure spot on the leaves of barley, but in Wisconsin (Davis, 143, 144; Greene, 289), South Dakota, and Wyoming it occurs on the leaves of Phalaris arundinacea. In material collected along creeks in the Black Hills, South Dakota, and adjacent Wyoming village of Moxee, the writer noted prominent straw color to buff, later pale brown spots, which finally became covered with the chalky fruiting material of the fungus. The spots were amphigenous, elongate, striate, often 1-2 x 10-30 mm in diameter, finally more or less confluent and covering or killing most of the leaf surface. The spots were emarginate or were surrounded by a diffuse yellow-buff area. The conidiophores were in compact fascicles in neat rows between the veins of the leaf, emerging from stomata and arising from abundant yellow to mostly hyaline compacted or serpentine hyphae. The conidiophores were 3-4.5 μ in diameter, 20-40 μ tall, tortuous, twisted or remarkably serpentine particularly at the apex. The conidia were produced near the apex on the side of the conidiophores and, on breaking off, a slight hilum was left at the base of the elliptical to elongate ovate spores. The spores were hyaline, one-celled with an evident wall which appeared faintly roughened, 11-15.5 x 6-7.8 μ .

Jørstad (376) mentions that Ophiocladium hordei produces dark spots, bearing a white or pale pink coating produced by the conidiophores on barley leaves.

References: Cavara (99, p. 26); Davis (143, 144); Greene (289, p. 89); Jørstad (376); Sprague (666).

OVULARIA LOLII Volkart, Eyespot of Lolium.

The spores of this fungus are ellipsoid to ovoid-piriform, 15-18 x 10-11 μ , rarely 10-19 x 7-13 μ , borne in ashy spots with reddish borders. The size of the spores would indicate that this species is different from O. pulchella (Ces.) Sacc. but the writer referred one of his meager Oregon collections on L. perenne to O. pulchella and another to O. lolii, indicating that both species occur on this host, or more likely, that both belong in one variable species. The status of O. pulchella var. lolii-italici Ferr. has not been determined.

References: Saccardo (558, v. 18, p. 531); Sprague (627, 655).

OVULARIA PULCHELLA (Ces.) Sacc., Leaf Spot.

This fungus causes scattered spots on leaves and culms of Agrostis alba, A. palustris, Arrhenatherum elatius, Festuca megalaria Nutt., F. myuros, Holcus lanatus, and Lolium marschalli Stev. in western Oregon, on Agrostis alba and A. tenuis in Utah, and on Bromus carinatus and Glyceria elata (Nash) Hitchc. in southwestern Washington.

The spots are usually elliptical. Those on fescues are brown or tan color, often on the stem or sheath; those on broms are gray; those on species of Agrostis, Lolium, and Holcus are red bordered, of an eyespot type. Except on annual fescues, the disease is seldom common. Fescue grasses that are growing in thick stands in sandy or open-wooded areas sometimes have a considerable percentage of the plants showing one to several spots.

Ovularia pulchella (Ces.) Sacc. (558, v. 4, p. 145) was originally described as Ramularia pulchella by Cesati (99a), and its description later amplified by Fresenius (239) as causing a purple leaf spot of Dactylis glomerata. The spores are few, crystal hyaline, ovoid, 8-12 μ long. Davis described O. pulchella var. agropyri J. J. Davis on Agropyron trachycaulum (A. tenerum) from Wisconsin (Davis, 141) as having spherical to oval spores 9-12 x 6-9 μ . Later he appeared to have considered this as being scarcely distinguishable from the species proper (Davis, 144, p. 102). See also O. pusilla (Ung.) Sacc. (558, v. 4, p. 140).

References: Cesati (99a); Davis (141, p. 714; 144, p. 102); Fresenius (239); Saccardo (558, v. 4).

PENICILLIUM SPP., Seed Rots.

Corn, sorghum, and to some extent wheat, Agropyron cristatum, and other grasses are injured in the soil before emergence by reduction in the

starchy content of the seeds before the seedlings are developed. Most of the injury occurs during prolonged rainy periods in April or May (and in 1942 even into June) in the Northern Great Plains, or earlier in the Pacific Northwest. Penicillium expansum Lk. em. Thom is the most common blue mold on these seeds, but a green mold caused by P. oxalicum Currie and Thom (133) appears to be an important species on seed corn in the Middle West. This species invades the embryo, causing yellowing of the leaves of the seedling and a gradual dying and drying of infected seedlings before the sixth leaf has unfolded (Johann, 363; Johann et al. 366). The dead plants, which remain stiffly upright, show little or no decay of the roots. The fungus is essentially a saprophyte. The injury is apparently due to oxalic acid, which kills the plant tissue in advance of the fungus.

Seed treatment (New Improved Semesan Jr.) will help reduce the endosperm-destroying molds. Proper curing of corn seed is necessary. In the drier parts of the Northern Great Plains, seed molds usually are less important and if clean local seed is used, seed treatment of field corn does not often show any benefit. However, sweet corn and pop corn usually show definite benefit from seed treatment in this area.

References: Currie and Thom (133); Diachun (157); Johann (363); Johann et al. (366); Leukel and Martin (405); Marchionatto (433); Noll (489); Thom (705).

PIRICULARIA GRISEA (Cke.) Sacc., Elast.

Gray, ashy or water-soaked spots are formed on the leaves and culms. The conidiophores rise from the stomata in groups of 2 to 5, are simple or rarely branched, fuliginous, septate, and bear conidia in terminal scorpioid cymes. The conidia are ovate, 2-septate, 24-29 x 10-12 μ .

P. grisea is common on Setaria millets in North Dakota and South Dakota and also occurs on certain other hosts in the United States. It is not important in the Northern States, as it is in the warmer regions. The closely related, if not identical, P. oryzae Briosi and Cav. on rice (Oryza sativa L.) has been intensively studied by the Japanese.

References: Schwartze (587, p. 120, figs. 735, 736); Tanaka (698).

RHYNCHOSPORIUM SPP.

Key to species:

Conidia apically obliquely-beaked..... R. secalis (Cud.) J. J. Davis

Conidia cylindrical..... R. orthosporum Caldwell

RHYNCHOSPORIUM ORTHOSPORUM Caldwell, Scald.

The spots on the leaves are lenticular at first, coalescing to form irregular water-soaked lesions, which later become gray, surrounded by a brown margin. In some cases, on Dactylis glomerata, the spots fade to light straw color.

The fungus is very destructive on Alopecurus pratensis L. at Corvallis, Oregon. It is moderately common on D. glomerata in that region, and occurs there on Agrostis alba, Elymus glaucus, Lolium perenne, and L. multiflorum also. While R. orthosporum differs distinctly from R. secalis in the absence of an oblique beak at the apex of the spore, there are sometimes collections in Oregon that are more or less intermediate between the two species. Both species occur on L. multiflorum.

R. orthosporum occurs on Calamagrostis canadensis in the Yellowstone National Park, Wyoming, and along Spearfish Creek in the Black Hills, South Dakota. Therefore, adding Caldwell's type on D. glomerata from Wisconsin, the fungus is scattered well across the Northern United States.

The material on Elymus glaucus, restricted to an area in Linn and Marion Counties, Oregon, may be an undescribed species or a variety with somewhat distinct morphology.

References: Caldwell (89); Sprague (655).

RHYNCHOSPORIUM SECALIS (Oud.) J. J. Davis, Scald.

The lesions are at first dark bluish-gray, water-soaked in appearance, later the collapsed tissue becomes light gray with a dark brown margin, sometimes irregularly zonate, often completely killing "scalded" leaves.

This fungus causes a serious disease of barley in Oregon, Washington, and California. Losses in winter barley in western Oregon and California often average 10 percent and losses up to 30 percent of the crop are reported. Even when all the leaves on some varieties have been 100 percent scalded the plants recover and give fair yields. This has erroneously led observers to believe that the temporary loss of foliage is unimportant. Wiebe, Caldwell (89, p. 176), and others have found this to be untrue. This disease is gradually becoming recognized as worthy of major consideration.

Besides barley, R. secalis occurs more or less spasmodically on a number of grasses and cereals in Oregon and Washington and scatteringly eastward as follows:

Oregon: Agropyron dasystachyum (Hook.) Scribn., A. repens, Agrostis

alba, Elymus glaucus, Hordeum jubatum, H. murinum, Lolium multiflorum,
L. perenne, Secale cereale.

Washington: Agropyron repens, A. semicostatum, A. trachycaulum, Bromus carinatus, B. inermis, Elymus canadensis, E. condensatus, E. glaucus,
E. junceus, E. triticoides, Secale cereale.

Idaho: Agropyron subsecundum.

California: Agropyron repens, Secale cereale.

Minnesota: A. repens.

South Dakota: Phalaris arundinacea.

Wyoming: P. arundinacea.

North Dakota: A. repens, P. arundinacea.

The collections on Phalaris arundinacea are less strongly beaked than most of the other collections (see B.P.I. 80,456). The spores are 13-18 x 2.3-2.8 μ .

In his detailed study, Caldwell (89) recognized six highly specialized physiologic races, which were distinguished by their ability to attack one of six hosts: rye, barley, quack grass, smooth brome, Canada wild-rye, or Hordeum jubatum. None of these, except the race from Canada wild-rye, could attack a host species of a genus other than that from which it was isolated. The race from Canada wild-rye could attack Agropyron trachycaulum, which is more or less to be expected as fungi attacking Agropyron trachycaulum or other members of the genus would likely also attack some members of the related genus Elymus. No doubt, were all of the additional hosts tested for racial affinities, several unrecognized strains would now be found.

References: Bartels (23); Caldwell (89); Davis (141, p. 713); A. G. Johnson and Mackie (371); Mackie (425).

SCOLECOTRICHUM GRAMINIS Fckl., Leaf Streak.

The shape, color, and extent, of the lesions caused by this fungus vary with age and host species. In its most common condition it occurs as elongate gray, brown-ochraceous to stramineous streaks on the leaves with aniline to sulphine yellow borders in which the black clusters of conidiophores appear as minute dots in seried arrangement. Younger stages show water-soaked circular to elliptical lesions which are deep olive gray in the morning when dew-wet, and deep dull gray when dry. These spots become brown, purple-brown to ocher with gray centers, with

a tendency to form streaks gradually as the leaves die.

This fungus is readily distinguished by the prominent, black densely fascicled conidiophores arranged in parallel rows as black dots along the surface. Without a hand lens, they are sometimes mistaken for pycnidia. Less frequently they may be plentiful enough to give a sooty mold aspect to the underside of the leaves. The spores are Indian-club shaped, i.e., fusoid-obclavate, 35-45 x 8-10 μ , 1- or sometimes 2-septate, olive brown.

The correct name of this fungus has never been settled to the satisfaction of all workers, although most of them use the old name, S. graminis, which Fuckel proposed in 1863 (241, No. 134). Horsfall (340) suggested that all available evidence indicated that Cercospora graminis (Fckl.) Horsfall would be a satisfactory name for the fungus. He stated that Von Höhnelt (Centrbl. Bak. 2:60: 1-26. 1924) considered Scolecotrichum as probably invalid. Until the matter is settled, the writer uses the name that is recognized by most workers. Synonyms for the fungus include: S. compressum Allesch. (Hedw. 35(2): 34. 1896); Cercospora graminicola Tracy and Earle (Torr. Bot. Club 22: 179. 1895); Passalora graminis (Fckl.) Hoehn. (Centrbl. Bakt. 2:60: 1-26. 1924); Passalora dactylina Pass. (In Lindau, G. Rab. Krypt. Fl. 1:8: 793-799. 1907); S. graminis var. nanum Sacc. (In Höhnelt, F. von. Centrbl. Bakt. 2 etc. 1924); Napicladium gramineum Pk. (Sprague, 657) and probably Cercospora poae Baudyš and Picb. 25); (Chupp, in letter to Sprague, 657).

Scolecotrichum graminis is known to occur on at least 104 species of Gramineae in the United States, 94 of them being included in this report. Of these, 10 are species of Agropyron; 6, of Agrostis; 4, of Alopecurus; 8 of Bromus; 9, of Elymus; and 13, of Poa; as well as 44 species of 22 other genera. The host range on Gramineae is almost unlimited. Horsfall listed only 28 host species in 1930; Seymour, about 27 for the entire United States. Still further study will no doubt add many more to the host range of this omnipresent species. S. graminis is particularly common on the Pacific Coast, but is abundant in the Plains country as well. It ranks as one of our most important leaf spot diseases of timothy, orchard grass, blue grass, tall meadow oats grass, and red top. Sometimes it is less serious than it appears to be because the fungus sporulates profusely in late season and develops saprophytically on many hosts on which it is only mildly parasitic. It is frequently secondary to other fungi. It is not important on cereals but sometimes occurs on rye in the coast region of Oregon. Once it was found on 40 percent of the leaves of rye in early May near Alsea, Oregon, (Sprague, 641) and on another occasion it was found on rye heads in the same general area. Guarch (301) reported S. graminis var. brachypoda Speg. on rye in Uruguay, and Jankowska (361) lists it from Poland on rye, and also as causing slight injury to wheat. Shitakova-Poussakova (605) reports the disease as widely distributed and of moderate importance on rye in the Soviet Union. She found some varietal resistance. Landaluze (397) reports S. graminis as a secondary parasite of rye in a

hilly region in Spain where the soil was pH 5.4. This condition is similar to that encountered near Alsea, Oregon, where the soil is highly acid (pH 4.9-5.5) and very deficient in nitrogen and phosphorus. Probably the injury on rye is not representative of strong parasitism in many cases.

Tabulation of about 250 collections of S. graminis shows that this fungus may be found any month of the year, but in Oregon its greatest activity is from March to July with a secondary development in the fall after the rains start again. Injury is greatest on older plants in spring. In the Great Plains, the fungus develops in May and June, but profusely fruiting material has been collected from May until late September.

This fungus needs critical study to determine its physiological status. Morphological variants occur (as on Poa ampla Merr. in Klickitat County, Washington), but whether they are anything more than variations due to host and humidity is not known.

Information on the control of this disease is scanty. Certainly grass weeds such as Hordeum jubatum and Beckmannia syzigachne probably aid in spreading it. Burning over fields no doubt reduces the injury from this fungus, which is able to increase its inoculum through saprophytic or semi-saprophytic development while the host is dormant or necrotic in very early or late season.

References: Fischer, et al. (231); Fuckel (241); Guarch (301); Horsfall (340); Jankowska (361); A. G. Johnson and Hungerford (370); Krause (390); Landaluze (397); Nilsson-Ehle (484); Shitakova-Roussakova (605); Sprague (641, 657); Trelease (709).

SCOLECOTRICHUM MACULICOLUM ELL. and Kell., Leaf Spot.

Prominent, elliptical, dirty-white spots form on the broad, living leaves of Phragmites communis. Originally described from Kansas (Ellis and Kellerman, 217), this fungus has been collected several times in one locality on Young's Bay, near Astoria, Oregon. The fungus has not been studied critically nor compared closely with S. graminis. It has spores 20-22 x 8-11 μ borne on conidiophores 40 x 4-5 μ . S. graminis has spores 35-45 x 8-10 μ (Saccardo 558 v. 4: 348) borne on conidiophores, 90-100 x 6-8 μ .

SPOROTRICHUM SP.

This fungus causes an obscure leaf killing on Agropyron spicatum, A. inerme, and A. subsecundum in Washington, Idaho, and Montana. The stromata are barely visible to the naked eye as fine, white stippling on dried leaf parts. Spores, which are intimately associated with the

branched septate hyphae, are 3-8 x 2.5-6 μ and secondary spores are produced by budding.

References: Sprague (668, 669).

STEMPHYLIUM SPP. (See Alternaria.)

Stemphylium botryosum Wallr. (Ascigerous stage, Pleospora herbarum (Pers.) Rab.) is the most abundant species, but it is of slight importance.

References: (Wiltshire, 768); (Groves and Skolko, 298).

LITERATURE CITED

1. ANONYMOUS. Plant diseases. Notes contributed by the Biological Branch. Agr. Gaz. N. S. Wales 48: 573-577. illus. 1937.
(Rev. Appl. Mycol. 17: 186. 1938).
2. ÅBERG, EVERT. Effect of vernalization on the development of stripe in barley. Phytopathology 35: 367-369. 1945
3. ÅKERMAN, Å., L. GRANHALL, and H. HAGANDER. Undersökningar över olika förfrukters inverkan å Veteskörden, särskilt med hänsyn till angrepp av rotdödare (Ophiobolus graminis). (Investigations on the influence of different preceding crops on the wheat harvest, especially with reference to the attack of the root-killer (Ophiobolus graminis). Sveriges Utsädesför. Tidskr. 1934, 44: 386-408. 1935. (German summary). (Rev. Appl. Mycol. 14: 351, 1935).
4. ADAM, D. B. On the occurrence of Cercosporiella herpotrichioides Fron in South Australia. Austral. Inst. Agr. Sci. Jour. 6: 48-51. 1940
5. _____ and T. T. CCLQUHOUN. The spread of take-all through the soil. Austral. Inst. Agr. Sci. Jour. 2: 172-174. 1936.
(Rev. Appl. Mycol. 16: 307. 1937).
6. ALFARO, A. Una septoriosis del Trigo. Bol. de Patol. Vég. y Ent. Agr. 9: 205-211, illus. 1940. (Rev. Appl. Mycol. 20: 249-250, 1941).
7. ALLESCHER, A. Fungi Imperfecti: hyaline-sporige sphaeriodeen. In RABENHORST, L., Kryptogamenflora von Deut., Oester. und Schweiz. 2, Aufl. 1 (Abt. 6): 1016 pp. 1901. (Leipzig).
8. ALLISON, J. L. Studies on Septoria bromigena. Phytopathology 29: 1. (Abs.) 1939
9. _____ Studies of monosporous cultures of Septoria bromigena. Phytopathology 29: 554-556. 1939
10. _____ Selenophoma bromigena leaf spot on Bromus inermis. Phytopathology 35: 233-240. 1945.
11. ANDERSON, J. P. Some Alaska fungi. Iowa Acad. Sci. Proc. 27: 99-108. [1920].
12. ANDRÉN, F. Några resultat från betningsförsök. Växtskyddsnotiser, Växtskyddsanst., Stockh., 1939 (4-5): 75-82. 1939. (Rev. Appl. Mycol. 19: 267, 1940).
13. _____ Resultatet från betningsförsök. Växtskyddsnotiser, Växtskyddsanst., Stockh., 1940: 3-7. 1940
14. ANDREWS, E. A. Seedling blight and root rot of grasses in Minnesota. Phytopathology 33: 234-239. 1943
15. _____ The pathogenicity of a non-sporulating Basidiomycete on grasses in Minnesota. Phytopathology 34: 352-353. 1944
16. ARMY, D. C. Physiologic specialization in Helminthosporium gramineum Rabh. Phytopathology 35: 571-572. 1945
17. ASHBY, S. F. Macrophomina phaseoli (Maubl.) comb. nov. The pycnidial stage of Rhizoctonia bataticola (Taub.) Butl. Brit. Mycol. Soc. Trans. 12: 141-147. 1927.

18. ATANASOFF, D. The Dilophospora disease of cereals. *Phytopathology* 15: 11-40. 1925
19. _____ and JOHNSON, A. G. Treatment of cereal seeds by dry heat. *Jour. Agr. Res.* 18: 379-390, illus. 1920
20. ATKINSON, G. F. Some Cercosporae from Alabama. *Jour. Elisha Mitchell Sci. Soc.* 8: 33-66. 1891
21. BAINIER, M. G. *Mycothèque de l'école de Pharmacie*, XXIV. Soc. Mycol. de France, *Bul. Trimest.* 24: 73-84. 1908
22. BARRUS, M. F. Yellow-spot disease of wheat in New York State. U. S. Dept. Agr. Bu. Plant Indus. *Plant Dis. Rptr.* 26: 246. 1942
23. BARTELS, F. Studien über Marssonina graminicola. *Forsch. auf dem Geb. Pflanzenkrank. u. Immunität Pflanzenr.* 5: [73] - 114. illus. 1928
24. BAUDYŠ, E. Ein beitrage zur kenntnis der mikromyceten in Böhmen. *Lotos* 63: 103-112; 64: 11-29, 42-64, 80-85, illus. 1915-16
25. _____ and PICBAUER, R. *Fungi novi vel minus cogniti Pars I.* *Acta Soc. Sci. Nat. Morav.* I: 299-304. 1924
26. _____ *Fungi novi vel minus cogniti Pars III.* *Acta Soc. Sci. Nat. Morav.* 3: 29-36, illus. 1926
27. BAUMLER, J. A. *Mykologische Notizen.* Wien. II. *Osterr. Bot. Ztschr.* 39: 289-290. 1890
28. BEACH, W. S. Biologic specialization in the genus Septoria. *Amer. Jour. Bot.* 6: 1-33, illus. 1919
29. BEAUMONT, A. Damping off in oats. *Seale-Hayne Agr. Col. Pam.* 21: 9. 1927
30. BECKWITH, T. D. Root and culm infections of wheat by soil fungi in North America. *Phytopathology* 2: 90. 1912
31. BENNETT, F. T. On Cladosporium herbarum: the question of its parasitism, and its relation to "thinning out" and "deaf ears" in wheat. *Ann. Appl. Biol.* 15: 191-212, illus. 1928 (Rev. *Appl. Mycol.* 7: 709. 1928.)
32. _____ On two species of Fusarium, Fusarium culmorum (W. G. Sm.) Sacc. and Fusarium avenaceum (Fries) Sacc. as parasites of cereals. *Ann. Appl. Biol.* 15: 213-244, illus. 1928. (Rev. of *Appl. Mycol.* 7: 710. 1928).
33. _____ Gibberella saubinetii (Mont.) Sacc. on British cereals. *Ann. Appl. Biol.* 17: 43-58. illus. 1930 (Rev. *Appl. Mycol.* 9: 516)
34. _____ Fusarium species on British cereals. The Gibbosum Group. I. F. scirpi Lamb. et Fautr. *Ann. Appl. Biol.* 19: 21-34. illus. 1932.
35. _____ Fusarium species on British cereals. Fusarium nivale (Fr.) Ces. Calonectria graminicola (Berk. and Br.) Wr. *Ann. Appl. Biol.* 20: 272-290. illus. 1933
36. _____ Rhizoctonia disease of turf. *Gard. Chron.* 97: 129. illus. 1935. (Rev. *Appl. Mycol.* 14: 449. 1935).
37. _____ Fusarium species on British cereals. *Ann. Appl. Biol.* 22: 479-507. illus. 1935, b (Rev. *Appl. Mycol.* 15: 6. 1936).
38. _____ Dollar spot disease of turf and its causal organism

- Sclerotinia homeocarpa n. sp. Ann. Appl. Biol. 24: 236-257. 1937. (Rev. Appl. Mycol. 16: 681-682. 1927).
39. BENNETT, F. T. Fusarium disease of cereals. Jour. Min. Agr. 45: 1115-1118. illus. (Rev. Appl. Mycol. 18: 446. 1939).
 40. BENSAUDE, MATHILDE. L'Helminthosporium tetramera McK. sur Blé à Angola. Compt. Rend. Soc. de Biol. 103: 1265-1266. 1930.
 41. BERKELEY, M. J. Disease in the wheat crop. Gard. Chron. 1845, p. 601. 1845
 42. _____ and M. A. CURTIS. Characters of new fungi collected in the north Pacific exploring expedition by Charles Wright. Amer. Acad. Arts and Sci. Proc. 4: 111-130. 1860.
 43. BERLESE, A. M., and PIETRO VOGLINO. Sopra un nuova genere di Funghi Sferopsidei. 32 pp., illus. 1886. (Extract from Soc. Veneto-Trentino di Sci. Nat., Padova. Atti 10, Fasc. 1).
 44. BERTUS, I. S. A sclerotial disease of maize due to Rhizoctonia solani. Ceylon Dept. Agr. Yearbook, pp. 44-46. 1927
 45. BESSEY, E. A. Dilophospora alopecuri. Jour. Mycol. 12: 57-58. illus. 1906
 46. BEVILACQUA, I. La micosi del Grano. Istria Agr. n. s. 15: 317-319. 1935
 47. BILJ, P. A. van der. A study on the "dry-rot" disease of maize caused by Diplodia zeae (Schw.) Lév. Union So. Africa Dept. Agr. Sci. Bul. 7, 60 pp., illus. 1916
 48. BIRAGHI, A. Ricerche sulla micoflora radicicola del Grano. Bol. Staz. Pat. Veg., Roma, n. s. 16: 147-154. illus. 1936
 49. BISBY, G. R., A. H. R. BULLER, J. Dearness, W. P. Fraser, R. C. RUSSELL, and H. T. GUSSOW. The fungi of Manitoba and Saskatchewan. Canad. Natl. Res. Council Mem., 189 pp. (Ottawa), illus. 1938
 50. BLAIR, I. D. The foot-rot disease of wheat. New Zeal. Jour. Agr. 52: 129-137. illus. 1936
 51. _____ Studies on the growth in soil and the parasitic action of certain Rhizoctonia solani isolates from wheat. Canad. Jour. Res. Sect. C, 20: 174-185. illus. 1942
 52. _____ Behavior of the fungus Rhizoctonia solani Kühn in the soil. Ann. Appl. Biol. 30: 118-127. 1943. (Expt. Sta. Rec. 90: 205. 1944)
 53. BOCKAMP, HANS. Ein Beitrag zur Biologie und wirtschaftlichen Bedeutung des Erregers der Braunfleckigkeit des Weizens: Macrophoma hennebergii (Kühn). Angew. Bot. 14: 79-86. 1932.
 54. _____ Die Schwärzspilze des Getreides unter besonderer Berücksichtigung ihrer Pathogenität und des Vorkommens von Rassen innerhalb der Gattungen Cladosporium Link und Alternaria Nees. Angew. Bot. 15: 308-385. 1933
 55. BOEDIJN, K. B. Ueber einige phragmosporen Dematiaceen. (On some phragmosporous Dematiaceae.) Buitenzorg Jard. Bot. Bul. Ser. III, 13: 120-134. illus. 1933
 56. BOLLEY, H. L. Wheat: Soil troubles and seed deterioration. Causes of soil sickness in wheat lands. Possible methods of control. Cropping methods with wheat. N. Dak. Agr. Expt. Sta.

- Bul. 107, 96 pp., illus. 1913
57. BONIER, E., and M. ROUSSEAU. Contributions a la flore mycologique de Belgique. Soc. Roy. de Bot. de Belg. Mem. (Bul.) 29: 205-302. 1890
 58. BONDARZEVA-MONTEVERDE, V. N. K Mikoflore Orlovskoi gubernii. Dva noviek narzritniek gribka. [On the Microflora of Lower Orlovski. Two new parasitic fungi.] Materialy po Mikologicheskomu Obsledovaniiu Rossii [Materials on Mycological Inspection in Russia.] 5, No. 1, 4 pp. 1921. [In Russian. Description of fungi in Latin.]
 59. BÖNING, K., and F. WALLNER. Versuche zur Bekämpfung der Netzfleckenkrankheit der Gerste (Helminthosporium teres Sacc.). Prakt. Bl. f. Pflanzenbau u. Pflanzenschutz 12: 219-229. illus. 1934. (Rev. Appl. Mycol. 14: 159. 1935).
 60. _____ Welke, Fusskrankheit, und andere Schädigungen an Mais durch Colletotrichum graminicolum (Ces.) Wilson. Phytopath. Ztschr. 9: 99-100. illus. 1936. (Rev. Appl. Mycol. 15: 494. 1936).
 61. BOWMAN, D. H., J. H. MARTIN, L. E. MELCHERS, and J. H. PARKER. Inheritance of resistance to Pythium root rot in Sorghum. Jour. Agr. Res. 55: 105-115. illus. 1937
 62. BOYLE, L. W. Diseases of oats in Western Washington and Oregon. U. S. Dept. Agr. Bur. Plant Indus. Plant Dis. Rptr. 28: 1121-1122. 1944. (Processed)
 63. BRANSTETTER, B. B. Corn root rot studies. Missouri Agr. Expt. Sta. Res. Bul. 113, 79 pp., illus. 1927
 64. BREYER, H. An American oat disease found in western Anatolia. Phytopathology 33: 165-167. 1943
 65. BRENTZEL, W. E. Blight and other damage of durum wheat in 1940. 2. Effects of blights and other damage on durum and hard wheat seed. N. Dak. Agr. Expt. Sta. Bul. 296: 9-14. illus. 1941
 66. _____ The black point disease of wheat. N. Dak. Agr. Expt. Sta. Bul. 330, 14 pp., illus. 1944
 67. BRESADOLA, G. Fungi aliquot Saxonici novi lecti a Cl. W. Krieger. Hedwigia 39: 325-328. 1900
 68. BROADFOOT, W. C. On the pathogenicity of Wojnowicia graminis. Phytopathology 23: 1001-1002. 1933
 69. _____ Studies on foot and root rot of wheat I. Effect of age of the wheat plant upon the development of foot and root rot. Canad. Jour. Res. Sect. C, 8: 483-491. illus. 1933
 70. _____ Studies on foot and root rot of wheat II. Cultural relationships on solid media of certain microorganisms in association with Ophiobolus graminis Sacc. Canad. Jour. Res. Sect. C, 8: 545-552. 1933
 71. _____ Studies on foot and root rot of wheat III. Effect of crop rotation and cultural practise on the development of foot rot of wheat. Canad. Jour. Res. Sect. C, 10: 95-113. 1934

72. BROADFOOT, W. C. Studies on foot and root rot of wheat. IV. Effect of crop rotation and cultural practises on the relative prevalence of Helminthosporium sativum and Fusarium spp. as indicated by isolations from wheat plants. Canad. Jour. Res. Sect. C, 10: 115-124. 1934
73. _____, and M. W. CORMACK. A low-temperature Basidiomycete causing early spring killing of grasses and legumes in Alberta. Phytopathology 31: 1058-1059. illus. 1941
74. _____, and L. E. TYNER. Studies on foot and root rot of wheat, V. The relation of phosphorus, potassium, nitrogen, and calcium nutrition to the foot- and root-rot disease of wheat caused by Helminthosporium sativum P. K. and B. Canad. Jour. Res. Sect. C, 16: 125-134. illus. 1938
75. _____, and _____ Studies on foot- and root-rot of wheat VI. Methods of securing infection of wheat seedlings for study in nutrient solutions. Canad. Jour. Res. Sect. C, 16: 253-261. 1938
76. BROMMELHUES, MARIA. Die wechselseitige Beeinflussung von Pilzen und die Bedeutung der Pilzkonkurrenz für das Ausmass der Schädigung an Weizen durch Ophiobolus graminis Sacc. Zentbl. f. Bakt., Abt. 2, 92: 81-116. illus. 1935 (Rev. Appl. Mycol. 14: 688-689. 1935)
77. BUBÁK, F. In HANDEL-MAZETTI, H. F. von. Ergebnisse einer botanischen Reise in das Pontische Randgebirge in Sandschak Trapezunt. Ann. K. K. Naturhist. Hofmus. 23: 105. 1909
78. BUCHHOLTZ, W. F. Gross pathogenic effects of Pythium graminicolum, Pythium debaryanum and Helminthosporium sativum on seedlings of crested wheatgrass. Phytopathology 32: 2. (Abs.). 1942
79. _____ Influence of cultural factors on alfalfa seedling infection by Pythium debaryanum Hesse. Iowa Agr. Expt. Sta. Res. Bul. 296. 24 pp., illus. 1942
80. BUDDIN, W., and S. D. GARRETT. Seasonal occurrence of the take-all disease of wheat. Ann. Appl. Biol. 28: 74. 1941. (Rev. Appl. Mycol. 20: 250. 1941).
81. BUNTING, R. H. Local cereal diseases in the records of the mycological Division. Year-Book Gold Coast Dept. of Agr., 1926, Bul. 7: 25-27. illus. 1927. (Rev. Appl. Mycol. 7: 231. 1928).
82. _____ Fungi affecting Gramineous plants of the Gold Coast. Gold Coast Dept. Agr. Bul. 10, i-iii + 51 pp., illus. 1928.
83. BURGER, O. F. Variations in Colletotrichum gloeosporioides. Jour. Agr. Res. 20: 723-736. illus. 1921.
84. BURRILL, T. J., and J. T. BARRETT. Ear rots of corn. Ill. Agr. Expt. Sta. Bul. 133: 64-109. illus. 1909
85. BUSSMAN, B. Untersuchungen über die Virulenz von Ophiobolus graminis Sacc. Phytopath. Ztscht. 9: 571-581. illus. 1936. (Rev. Appl. Mycol. 16: 305-306. 1937).
86. BUTLER, E. J. An account of the genus Pythium and some

- Chytridiaceae. India Dept. Agr. Mem., Bot. Ser., 1 (5), 161 pp. illus. 1907
87. BUTLER, E. J. Pythium debaryanum Hesse. India Dept. Agr. Mem., Bot. Ser., 5 (5): 262-267, illus. 1913.
 88. CABALLERO, A. Adiciones a la micoflora española. R. Soc. Españ. de Hist. Nat. Bol. 28: 421-430. illus. 1928.
 89. CALDWELL, RALPH M. Rhynchosporium scald of barley, rye, and other grasses. Jour. Agr. Res. 55: 175-188. illus. 1937
 90. CAMPI, M. D. Helminthosporium turcicum Pass. en la Republica Argentina. Lilloa 4: 5-32. 1939. (English summary).
 91. CARNE, W. M., and J. G. C. CAMPBELL. Take-all of wheat and similar diseases of cereals. West Austral. Dept. Agr. Bul. 119, 10 pp. 1924.
 92. CARPENTER, C. W. Predisposing factors in Pythium root rot. Hawaii. Planters' Rec. 38: 279-338. 1934.
 93. CARRERA, C. J. M. El género Fusarium en la Republica Argentina. Estudio e identificación de algunas especies. Physis 15: 21-77. illus. 1939.
 94. _____ El género Fusarium en la República Argentina. Estudio y clasificación sistemática. (Segunda contribucion). Rev. Argentina de Agron. 7: 277-296. illus. 1940
 95. CARROLL, P. T. Some factors influencing lodging in cereals. Jour. Dept. Agr. Eire 40: 280-285. 1943
 96. CARVAJAL, FERNANDO. Phoma terrestris on sugar-cane roots in Louisiana. Phytopathology 35: 744. 1945
 97. CASH, EDITH K. Alaskan fungi. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 18: 74-88. 1934. [Processed.]
 98. CASTAGNE, L. Supplement au catalogue des plantes qui croissent naturellement aux environs de Marseilles. 135 pp. illus. 1851. Aix. Imper. de Nicotol Pardigon
 99. CAVARA, F. Ueber einige parasitische Pilze auf dem Getreide. Ztschr. f. Pflanzenzucht. 3: 21, 26. 1893
 - 99a. CESATI, V. de. Ramularia pulchella sp. nov. In Rabenhorst, Klotzsch. Herb. Mycol. No. 1781. Also Bot. Zeit. XI, p. 238. 1853.
 100. CHAMBERLAIN, DONALD W., and J. LEWIS ALLISON. A brown leaf spot on Bromus inermis Leyss caused by Pyrenophora bromi (Died.) Drechsler. Phytopathology 34: 997-998. (Abs.). 1944.
 101. _____, and _____ The brown leaf spot on Bromus inermis caused by Pyrenophora bromi. Phytopathology 35: 241-248. 1945.
 102. CHARDON, CARLOS E. A contribution to our knowledge of the Pyrenomycetes of Porto Rico. Mycologia 13: 279-300. illus. 1921.
 103. CHESTER, K. STARR. The nature and prevention of plant disease. 584 pp. 1942. The Blakiston Co., Philadelphia
 104. CHILTON, S. J. P. The occurrence of Helminthosporium turcicum in the seed and glumes of Sudan grass. Phytopathology 30: 533-536. 1940

105. CHOWDHURY, S. C. A disease of Zea mays caused by Colletotrichum graminicolum (Ces.) Wils. Indian Jour. Agr. Sci. 6: 833-843. illus. 1936
106. CHRISTENSEN, J. J. Studies of the parasitism of Helminthosporium sativum. Minn. Agr. Expt. Sta. Tech. Bul. 11, 42 pp., illus. 1922.
107. _____ Physiologic specialization and mutation in Helminthosporium sativum. Phytopathology 15: 785-795. 1925.
108. _____ Physiologic specialization and parasitism of Helminthosporium sativum. Minn. Agr. Expt. Sta. Tech. Bul. 37, 101 pp., illus. 1926.
109. _____ Association of micro-organisms in relation to seedling injury arising from infected seed. Phytopathology 26: 1091-1105. 1936.
110. _____, and F. R. DAVIES. Nature of variation in Helminthosporium sativum. Mycologia 29: 85-99. 1937.
111. _____, and _____ Variation in Helminthosporium sativum induced by a toxic substance produced by Bacillus mesentericus. Phytopathology 30: 1017-1033. 1940.
112. _____, and T. W. GRAHAM. Physiologic specialization and variation in Helminthosporium gramineum Rab. Minn. Agr. Expt. Sta. Tech. Bul. 95, 40 pp., illus. 1934.
113. _____, and E. C. STAKMAN. Relation of Fusarium and Helminthosporium in barley seed to seedling blight and yield. Phytopathology 25: 309-327. 1935.
114. CHUPP, CHARLES. Cercospora species and their host genera. Plant Pathology Dept. Cornell Univ., Ithaca, New York. 23 pp. 1937. [Processed]
115. CLARK, F. E. Experiments toward the control of the take-all disease of wheat and the Phymatotrichum root rot of cotton. U. S. Dept. Agr. Tech. Bul. 835, 27 pp., illus. 1942
116. CLAYTON, E. E. Diplodia ear-rot disease of corn. Jour. Agr. Res. 34: 357-371. 1927
117. CLEMENTS, F. E., and C. L. SHEAR. The general of fungi. i-viii + 496 pp., illus. 1931. H. W. Wilson Co., New York
118. COCCONI, G. Un pizzico di funghi nuovi. R. Accad. delle Sci. dell'Ist. Bologna, Mem. della Cl. di Sci. Fis., Ser. 5, 6: 149-158. illus. 1896-97.
119. _____, and F. MORINI. Enumerazione dei funghi della provincia di Bologna, Secunda Cent. R. Accad. delle Sci. dell'Ist. Bologna, Mem. della Cl. di Sci. Fis., Ser. 4, 5: 273-300. illus. 1883
120. _____, and _____ Enumerazione dei funghi della provincia di Bologna, Terza Cent. R. Accad. delle Sci. dell'Ist. Bologna, Mem. della Cl. di Sci. Fis., Ser. 4, 6: 371-400. 1884
121. CONNERS, I. L. Seventeenth Annual Report of the Canadian Plant Disease Survey, 1937. i-xi + 87 pp. 1938. [Processed]
122. _____ Twenty-first annual report of the Canadian Plant Disease Survey, 1941. i-xviii + 102 pp. 1942. [Processed]

123. CONNERS, I. L., and D. B. O. SAVILE. Twenty-second annual report of the Canadian Plant Disease Survey, 1942. i-xix + 110 pp. 1943. [Processed]
124. COOK, M. T. Species of Synchytrium in Louisiana. III. The development and structure of the galls. Mycologia 37: 715-740. 1945.
125. COOKE, M. C. and J. B. ELLIS. New Jersey fungi. Grevillea 8: 11-16. 1879.
126. COOKE, WM. BRIDGE. Septoria infuscans (E. and E.) Sprague comb. nov. Mycobiota of North America 113. 1940. (Exsiccati.)
127. _____ Septoria jacucella [jaculella] Sprague sp. nov. Mycobiota of North America 114. 1940. (Exsiccati.)
128. CORDA, A. C. I. Icones fungorum hucusque cognitorum. t. 3, 55. Prague, illus. 1839.
129. CORDLEY, A. B. A foot rot of wheat. Oreg. Agr. Expt. Sta. Ann. Rpt. 14: 66-67. 1902
130. CORMIER, P. Le Septoria graminum, champignon parasite du blé. Acad. d'Agr. de France Compt. Rend. 18: 671-672. 1932.
131. CRALLEY, E. M., and E. C. TULLIS. Effect of seed treatments on seedling emergence, severity of seedling blight, and yield of rice. Ark. Agr. Expt. Sta. Bul. 345, 24 pp. 1937
132. CROSIER, W., and D. WEIMER. Some fungi associated with grass seed. Assoc. Off. Seed Anal. No. Amer. Proc. 1939, p. 120-124. illus. 1940.
133. CURRIE, J. N., and C. THOM. An oxalic acid producing Penicillium. Jour. Biol. Chem. 22: 287-293. 1915. (Also Science n. s. 42: 952. 1915).
134. CURZI, D. M. La pintatura delle cariossidi di frumento e una nuova specie di Alternaria. Riv. di Patol. Veg. 16: 125-136. 1926.
135. _____ Una nuova grave malattia del granturco. Rend. Accad. Lincei, ser. 6, 10: 306-308. 1929. (Rev. Appl. Mycol. 9: 174-175. 1930).
136. DAHL, A. S. Snowmold of turf grasses caused by Fusarium nivale. Phytopathology 24: 197-214. 1934.
137. DANA, B. F. A preliminary note on foot rot of cereals in the Northwest. Science n. s. 50: 484-485. 1919.
138. DARLEY, E. F. Spore germination of Selenophoma bromigena. Phytopathology 31: 953-954. illus. 1941.
139. DAVIS, J. J. Notes on parasitic fungi in Wisconsin--I. Wis. Acad. Sci., Arts, Letters, Trans. 18: 78-92. 1915.
140. _____ North American Ascochytae. Wis. Acad. Sci., Arts, Letters, Trans. 19: 655-670. 1919
141. _____ Notes on parasitic fungi in Wisconsin, VI. Wis. Acad. Sci., Arts, Letters, Trans. 19: 705-727. 1919
142. _____ Notes on parasitic fungi in Wisconsin, XI. Wis. Acad. Sci., Arts, Letters, Trans. 21: 287-302. 1924.
143. _____ Notes on parasitic fungi in Wisconsin, XIX. Wis. Acad. Sci., Arts, Letters, Trans. 27: 185. 1932.

144. DAVIS, J. J. Parasitic fungi of Wisconsin, 157 pp. Madison, Wis. 1942
145. DAVIS, R. J. Studies on Ophiobolus graminis Sacc. and the take-all disease of wheat. Jour. Agr. Res. 31: 801-825. 1925.
146. DE HAAN, J. T. Untersuchungen über das Auftreten der Keimlingsfusariose bei Gerste, Hafer, Mais und Reis. Phytopath. Ztschr. 10: 235-305. illus. 1937.
147. DELACROIX, G. Sur le piétin des céréales. Bul. Soc. Mycol. France. 17: 136-144. illus. 1901
148. DEMIDOVA, Z. Nabliudeniiia nad vi idami Septoria na zlakakh. [Studies on species of Septoria on grasses.] Mater. po Mikol. i Fitopat. [Mater. Mycol. and Phytopath.] 5: 133-157. illus. 1926.
149. DENNIS, R. W. G. Studies in morphology and biology of Helminthosporium avenae. Brit. Mycol. Soc. Trans. 18: 223-238. 1933.
150. _____ Notes on the occurrence of Pyrenophora avenae Ito in Scotland. Brit. Mycol. Soc. Trans. 19: 288-290. 1935.
151. _____ Seed borne diseases in Scottish seed oats. Scot. Farmer, July 24, p. 7, 1943.
152. _____, and C. E. FOISTER. List of diseases of economic plants recorded in Scotland. Brit. Mycol. Soc. Trans. 25: 266-306. 1942
153. DESMAZIERES, J. B. H. J. Neuvième notice sur quelques plantes cryptogames. Ann. Sci. Nat. Bot. Sér. 2, 17: 91-(107). 1842
154. _____ Dixième notice sur quelques plantes cryptogames. Ann. Sci. Nat. Bot. Sér. 2, 19: 335-373. 1843
155. _____ Quatorzième notice sur les plantes cryptogames. Ann. Sci. Nat. Bot. Sér. 3, 8: 9-37. 1847.
156. _____ Plantes Cryptogames de France (Exsiccati). (Ed. 1). Fasc. 44, no. 2169. 1851
157. DIACHUN, STEPHEN. The effect of some soil factors on Penicillium injury of corn seedlings. Phytopathology 29: 231-241. 1939
158. DICKINSON, LAWRENCE S. The effect of air temperature on the pathogenicity of Rhizoctonia solani parasitizing grasses on putting-green turf. Phytopathology 20: 597-608. 1930.
159. DICKSON, JAMES G. Influence of soil temperature and moisture on the development of the seedling-blight of wheat and corn caused by Gibberella saubinetii. Jour. Agr. Res. 23: 837-870. illus. 1923.
160. _____ Outline of diseases of cereal and forage crop plants of the northern part of the United States. i-vii + 259 pp. 1939. Burgess Pub. Co., Minneapolis, Minn.
161. _____ Scab of wheat and barley and its control. U. S. Dept. Agr. Farmers' Bul. 1599, 22 pp., illus. 1942.
162. _____, S. H. ECKERSON, and K. P. LINK. The nature of resistance to seedling blight of cereals. Natl. Acad. Sci. Proc. 9: 434-439. 1923.
163. DIEDICKE, H. von. Ueber den Zusammenhang zwischen Pleospora- und Helminthosporium-Arten I. Centrbl. Bakt. etc., Abt.

- 2, Bd. 9: 317-329. illus. 1902.
164. DIEDICKE, H. von. Ueber den Zusammenhang zwischen Pleospora-
und Helminthosporium-Arten II. Centrbl. Bakt., etc., Abt.
2, Bd. 11: 52-59. 1903.
 165. _____ Die Gattung Septoria. Ann. Mycol. (Sydow)
10: 477-487. 1912.
 166. _____ Die Abteilung Hyalodidymae der Sphaerioideen.
Ann. Mycol. (Sydow) 10: 135-152. 1912
 167. _____ Kryptogamen Flora der Mark Brandenburg.
Pilze VII. Sphaeropsideae, Melanconieae. 9. 962. pp.,
illus. 1915. (Leipzig).
 168. DIEHL, W. W. Conidial fructifications in Balansia and Dothi-
chloë. Jour. Agr. Res. 41: 761-766. illus. 1930.
 169. _____ Diplodina graminea Sacc. in South Carolina. U. S.
Dept. Agr. Bur. Plant Indus. Plant Dis. Rptr. 19: 246. 1935.
[Processed.]
 170. _____ Identity and parasitism of a species of Dothichloë.
Jour. Agr. Res. 58: 947-954. illus. 1939
 171. DILLON WESTON, W. A. R., and S. D. GARRETT. Rhizoctonia solani
associated with a root rot of cereals in Norfolk. Ann. Appl.
Biol. 30: 79. 1943.
 172. DOSDALL, LOUISE. Factors influencing the pathogenicity of Hel-
minthosporium sativum. Minn. Agr. Expt. Sta. Tech. Bul. 17,
47 pp., illus. 1923.
 173. _____, and J. J. CHRISTENSEN. Variations in the length
of spores of Helminthosporium sativum P. K. & B. under differ-
ent conditions of growth. Phytopathology 13: 50. (Abs.).
1923.
 174. DRECHSLER, CHARLES. Net blotch of meadow fescue caused by an
undescribed species of Helminthosporium. Phytopathology 12:
35. (Abs.). 1922
 175. _____ A new leaf spot of Kentucky blue grass
caused by an undescribed species of Helminthosporium. Phy-
topathology 12: 35. (Abs.). 1922
 176. _____ Some graminicolous species of Helminthospor-
ium: I. Jour. Agr. Res. 24: 641-740. illus. 1923
 177. _____ Pythium infection of cabbage heads. Phy-
topathology 15: 482-485, illus. 1925
 178. _____ Pythium ultimum and Pythium debaryanum.
Phytopathology 17: 54-55. (Abs.) 1927.
 179. _____ Pythium arrhenomanes n. sp., a parasite
causing maize root rot. Phytopathology 18: 873-875, illus.
1928
 180. _____ Zonate eyespot of grasses caused by Hel-
minthosporium giganteum. Jour. Agr. Res. 37: 473-492, illus.
1928
 181. _____ The beet water mold and several related root
parasites. Jour. Agr. Res. 38: 309-361, illus. 1929
 182. _____ Occurrence of the zonate-eyespot fungus

- Helminthosporium giganteum on some additional grasses. Jour. Agr. Res. 39: 129-[136], illus. 1929
183. DRECHSLER, CHARLES. Some new species of Pythium. Wash. Acad. Sci. Jour. 20: 398-418. 1930.
 184. _____ A leaf spot of bent grasses caused by Helminthosporium erythrospilum n. sp. Phytopathology 25: 344-361. 1935.
 185. _____ Pythium graminicolum and P. arrhenomanes. Phytopathology 26: 676-684. 1936.
 186. _____ Three species of Pythium associated with root rots. Phytopathology 30: 189-213. illus. 1940
 187. DUCOMET, V. La brulure du maïs dans le Sud-Ouest. Jour. d'Agr. Prat. n.s. 5: 507-511. illus. 1903.
 188. DUNLAP, A. A. Lodging of sorghum in Texas. U. S. Dept. Agr. Bur. Plant Indus. Plant Dis. Rptr. 22: 402-403. [Processed] 1938
 189. DURRELL, L. W. Dry rot of corn. Iowa Agr. Expt. Sta. Res. Bul. 77, pp. 345-376. 1923
 190. _____ A preliminary study of fungous action as the cause of down corn. Phytopathology 15: 146-154. 1925.
 191. _____ Basisporium dry rot of corn. Iowa Agr. Expt. Sta. Res. Bul. 84, p. 139-160, illus. 1925.
 192. EDDINS, A. H. Corn diseases in Florida. Fla. Agr. Expt. Sta. Bul. 210, 35 pp., illus. 1930.
 193. _____, and R. K. VORHEES. Physalospora zeicola on corn and its taxonomic and host relationships. Phytopathology 23: 63-72. 1933.
 194. EDGERTON, C. W., E. C. TIMS, and P. J. MILLS. Relation of species of Pythium to the root-rot disease of sugar cane. Phytopathology 19: 549-564, illus. 1929.
 195. EDWARDS, E. T. Internal grain infection and Kernel rot in the 1938 American maize crop. Jour. Austral. Inst. Agr. Sci. 6: 25-31. 1940
 196. _____ The biological antagonism of Gibberella fujikuroi and Gibberella fujikuroi var. subglutinans to Trichoderma viride, with notes on the pathological effects of the latter fungus on maize. Jour. Austral. Inst. Agr. Sci. 6: 91-100, illus. 1940
 197. _____ The relation of mineral nutrition to seedling blight infection in maize. Jour. Austral. Inst. Agr. Sci. 7: 147-154, illus. 1941
 198. _____ Internal grain infection in Maize due to Gibberella fujikuroi and Gibberella fujikuroi var. subglutinans. Jour. Austral. Inst. Agr. Sci. 7: 74-82, illus. 1941
 199. EIDAM, E. Das vorkommen der Fleckenkrankheit auf Gersten- und auf Haferblättern. Der Landw. Bd. 27, p. 509. 1891
 200. EIDE, C. J. The pathogenicity and genetics of Gibberella saubinetii (Mont.) Sacc. Minn. Agr. Expt. Sta. Tech. Bul. 106, 67 pp. illus. 1935.
 201. EKSTRAND, H. Trädklubba på vintersäd Sklerotiesjuka på fodergräs (Typhula on winter cereals, Sclerotial disease on forage grass).

- Växtskyddsnotiser, Statens Växtskyddsanst., Stockholm, 1: 3-5, illus. 1937
202. EKSTRAND, H. Några ekonomiskt viktiga sjukdomar på höstsäd och vallväxter. Medd. Växtskyddsanst., Stockholm, 25, 23 pp. 1938. (German summary). (Rev. Appl. Mycol. 18: 298-299. 1939).
203. ELLETT, C. W. Leaf blight of corn. Phytopathology 33: 407-408. 1943.
204. ELLIOTT, CHARLOTTE, L. E. MELCHERS, C. L. LEFEBVRE, and F. A. WAGNER. Pythium rot of milo. Jour. Agr. Res. 54: 797-834. illus. 1937.
205. _____, and M. T. JENKINS. Helminthosporium turcicum leaf blight of corn. Phytopathology 35: 485. (Abs.). 1945
206. ELLIOTT, JOHN A. Taxonomic characters of the genera Alternaria and Macrosporium. Amer. Jour. Bot. 4: 439-476. 1917
207. ELLIS, J. B. Descriptions of some new species of fungi. Jour. Mycol. 7: 274-278. 1893.
208. _____, and E. BARTHOLOMEW. New species of fungi from various localities. Jour. Mycol. 8: 173-178. 1902.
209. _____, and B. M. EVERHART. Additions to Cercospora, Gloeosporium and Cylindrosporium. Jour. Mycol. 3: 15. 1887
210. _____, and _____. Additions to Ramularia and Cercospora. Jour. Mycol. 4: 4. 1888. (or Sacc. Syll. Fung. 10:656. 1892).
211. _____, and _____. New species of fungi from various localities. Jour. Mycol. 4: 44-46. 1888
212. _____, and _____. New species of fungi. Jour. Mycol. 7: 130-135. 1892
213. _____, and _____. New North American fungi from various localities. Acad. Nat. Sci. Phila. Proc. 1893: 128-172. 1894.
214. _____, and _____. New species of fungi from various localities. Torrey Bot. Club Bul. 25: 501-514. 1898
215. _____, and _____. New species of fungi from various localities. Torrey Bot. Club Bul. 27: 571-578. 1900
216. _____, and _____. New species of fungi from various localities. Jour. Mycol. 8: 11-19. 1902.
217. _____, and W. A. KELLERMAN. Synopsis of the North American Species of Xylaria poronia. Jour. Mycol. 3: 108. 1887
218. ELVEDEN, VISCOUNT. A contribution to the investigation into the results of partial sterilisation of the soil by heat. J. Agr. Sci. 11: 197-210. 1921
219. ERIKSSON, J. Fungoid diseases of agricultural plants (English edition). 1912. p. 33.
220. ERWIN, L. E. A grass destroying fungus new to America. R. I. Agr. Expt. Sta. 46th Ann. Rpt. (Contrib. 449). pp. 89-92. 1934.
221. _____. Pathogenicity and control of Corticium fuciforme. R. I. Stat. Col. Bul. 278, 34 pp., illus. 1941.
222. FAIRMAN, C. E. New or noteworthy ascomycetes and lower fungi

- from New Mexico. *Mycologia* 10: 258. (239-264). 1918
223. FELLOWS, H. The influence of oxygen and carbon dioxide on the growth of Ophiobolus graminis in pure culture. *Jour. Agr. Res.* 37: 349-355, illus. 1928.
224. _____ Some chemical and morphological phenomena attending infection of the wheat plant by Ophiobolus graminis. *Jour. Agr. Res.* 37: 647-661. 1928.
225. _____ The infestation of soil with Ophiobolus graminis and its subsequent increase and spread in the soil. *Phytopathology* 27: 956. (Abs.) 1937
226. _____ Effect of climatic conditions on the prevalence of Ophiobolus graminis in the field. *Phytopathology* 27: 956. (Abs.) 1937
227. _____ Interrelation of take-all lesions on the crowns, culms, and roots of wheat plants. *Phytopathology* 28: 191-195. 1938.
228. _____ Effect of certain environmental conditions on the prevalence of Ophiobolus graminis in the soil. *Jour. Agr. Res.* 63: 715-726. Dec. 15, 1941.
229. _____ Unpublished annual reports dealing with rootrots of cereals in the Great Plains. Filed in the Div. of Cereal Crops and Diseases, Bur. of Plant Industry, Soils, and Agr. Eng., U. S. Dept. of Agr. Beltsville, Md. 1925-44.
230. _____, and C. H. FICKE. Soil infestation by Ophiobolus graminis and its spread. *Jour. Agr. Res.* 58: 505-519. 1939.
231. FISCHER, GEORGE W., R. SPRAGUE, HOWARD W. JOHNSON, and J. R. HARDISON. Host and pathogen indices to the diseases observed on grasses in certain western states during 1941. U. S. Dept. Agr., Bur. Plant Indus. Plant Dis. Rptr. Suppl. 137, pp. 87-144. 1942
232. FLOR, H. H. Relation of environmental factors to growth and pathogenicity of Pythium isolated from roots of sugar cane. *Phytopathology* 20: 319-328, illus. 1930.
233. FOEX, E. and E. ROSELLA. "Recherches sur le piétin." *Ann. Epiphyties.* 16: 51-82. 1930
234. _____ and _____ Quelques observations sur le piétin du blé. *Rev. Path. Veg. et Ent. Agr.* 18: 133-42. 1931.
235. _____ and _____ Un Sclerotium parasite du blé. *Ann. Sci. Nat. Bot., Sér. X*, 19: 221-231. 1937
236. FOSTER, W. R., and A. W. HENRY. Overwintering of certain cereal pathogens in Alberta. *Canad. Jour. Res. Sect. C*, 15: 547-559. 1937
237. FRANK, B. Die neuen Deutschen Getreidepilze. *Deut. Bot. Gesell. Ber.* 13: 61-65. 1895
238. FRASER, W. P. "Take-all" of wheat in western Canada. *Phytopathology* 14: 347 (Abs.). 1924
- 238a. _____, P. M. SIMMONDS, and R. C. RUSSELL. The take-all disease in Canada. *Phytopathology* 16: 80-81. (Abs.) 1926
239. FRESENIUS, G. Ramularia pulchella Cesati. *Beiträge zur Mykologie*,

- p. 90, illus. 1863.
240. FROM, G. Contribution a l'étude de la maladie "pied noir des céréales" ou "maladie du piétin". Ann. Sci. Agron. Franç. et Étrang. (4) 1: 3-29. 1912. (Rev. in Bul. of Bur. Agr. Intell. Plant Dis. Int. Inst. Agr. 3: 1054-1056, 1912).
 241. FÜCKEL, L. Fungi Rhenani Exsiccati. Hedwigia 2: 134. 1863. (And in Symbolae Mycol. 1869).
 242. GARRETT, S. D. Factors affecting the severity of take-all. I. The importance of soil micro-organisms. So. Austral. Dept. Agr. Jour. 37: 664-674. illus. 1934.
 243. _____ Factors affecting the severity of take-all. II. Soil temperature. So. Austral. Dept. Agr. Jour. 37: 799-805. illus. 1934.
 244. _____ Factors affecting the severity of take-all. III. The climatic factors. So. Austral. Dept. Agr. Jour. 37: 976-983. 1934.
 245. _____ Soil conditions and the take-all disease of wheat. Ann. Appl. Biol. 23: 667-699.
 246. _____ Take-all or whiteheads disease of wheat and barley and its control. Roy. Agr. Soc. Jour. 98: 24-34. 1937.
 247. _____ Soil conditions and the take-all disease of wheat II. The relation between soil reaction and soil aeration. Ann. Appl. Biol. 29: 747-751. 1937.
 248. _____ Soil conditions and the take-all disease of wheat. III. Decomposition of the resting mycelium of Ophiobolus graminis in infected wheat stubble buried in the soil. Ann. Appl. Biol. 25: 744-766. illus. 1938.
 249. _____ Soil conditions and the take-all disease of wheat. IV. Factors limiting infection by ascospores of Ophiobolus graminis. Ann. Appl. Biol. 26: 47-55. illus. 1939.
 250. _____ Temporary leys and the take-all and whiteheads disease of wheat and barley. Jour. Minist. Agr. 47: 134-135. 1940.
 251. _____ Utilization of nitrogen by Ophiobolus graminis. Nature (London) 145: 108-109. 1940.
 252. _____ Soil conditions and the take-all disease of wheat. V. Further experiments on the survival of Ophiobolus graminis in infected wheat stubble buried in the soil. Ann. Appl. Biol. 27: 199-204. 1940.
 253. _____ Soil conditions and the take-all disease of wheat. VI. The effect of plant nutrition upon disease resistance. Ann. Appl. Biol. 28: 14-18. 1941. (Rev. Appl. Mycol. 20: 250, 1941).
 254. _____ Soil conditions and the take-all disease of wheat. VII. Survival of Ophiobolus graminis on the roots of different grasses. Ann. Appl. Biol. 28: 325-332. 1941.
 255. _____ The take-all disease of cereals. Tech. Commun. Bur. Soil Sci. (Harpden) 41, 40 pp. illus. 1942.
 256. _____ Competition for nitrogen between the take-all

- fungus and the roots of crop plants. *Nature* (London) 152: 417-418. 1943.
257. GARRETT, S. D. Root disease fungi. *Ann. Crypt. et Phytopath.* 1, xii + 177 pp. The Chronica Botanica Co., Waltham, Mass. 1944
258. _____ Soil conditions and the take-all disease of wheat, VIII. Further experiments on the survival of Ophiobolus graminis in infected wheat stubble. *Ann. Appl. Biol.* 31: 186-191. 1944.
259. _____, and R. W. G. DENNIS. Notes on the occurrence of Ophiobolus graminis Sacc. var. avenae Turner in Scotland in 1942. *Brit. Mycol. Soc. Trans.* 26: 146-147. 1943.
260. GENTER, C. F. A study of the inheritance of susceptibility to leaf blight of corn. Ohio State Univ. Abs. Doct. Diss. No. 40, pp. 97-107. illus. 1942.
261. GILMAN, JOSEPH C. A manual of soil fungi, 392 pp., illus. The Collegiate Press, Inc., Ames, Ia. 1945.
262. _____, and ANDREW W. ARCHER. The fungi of Iowa parasitic on plants. *Iowa State College Jour. Sci.* 3: 299-502. 1929
263. GLYNNE, MARY D. Incidence of take-all on wheat and barley on experimental plots at Woburn. *Ann. Appl. Biol.* 22: 225-235. illus. 1935
264. _____ Some new British records of fungi on wheat. Cercospora herpotrichoides Fron, Gibbellina cerealis Pass., and Ophiobolus herpotrichus (Fr.) Sacc. *Brit. Mycol. Soc. Trans.* 20: 120-122. 1936
265. _____ III. Eyespot lodging of wheat caused by Cercospora herpotrichoides Fron. *Agr. Prog.* 16: 1-5. 1939
266. _____ Eyespot lodging of wheat caused by Cercospora herpotrichoides Fron. *Sci. Progr.* 16: 70-74. 1939
267. _____ Cercospora herpotrichoides Fron causing eyespot of wheat in Great Britain. *Ann. Appl. Biol.* 29: 254-264. illus. 1942
268. _____, and WENDY M. RITCHIE. Sharp eyespot of wheat caused by Corticium (Rhizoctonia) solani. *Nature* (London) 152: 161.
269. GODDARD, M. Studies on variation in Gibberella saubinetii (Mont.) Sacc. (Fusarium graminearum Schwabe). *Mo. Bot. Gard. Ann.* 26: 99-164. illus. 1939
270. GONZALEZ FRAGOSO, R. Nueva contribucion a la flora micologica del guadarrama, teleomicetos y deuteromicetos. *Trabajos Mus. Nac. Ciencias Nat. Ser. Bot. n. 7*: 1-80, illus. 1914
271. _____ Hongos parásitos de la florula hispalense, nuevos ó poco conocidos. *Bot. R. Soc. Españ. de Hist. Nat. Bot.* 15: 120-132. illus. 1915
272. _____ Algunos micromicetos más de los alrededores de Melilla (Marruecos) recolectados por el profesor d. a. Cabellero. *Soc. Bot. R. Españ. de Hist. Bot.* 17: 82-83. 1917
273. _____ Datos para la deuteromicetologia Catalana.

- Barcelona R. Acad. de Ciéncia y Artes, Mem. 15: 3-41 (429-467). 1920
274. GONZALEZ FRAGOSO, R. Hongos de España Ser. 2. Brotéria Ser. Bot. 22: 67-68. 1926
275. GORDON, W. L. Species of Fusarium isolated from field crops in Manitoba. Reprinted from Volume II. Proceedings of the World's Grain Exhibition and Conference, Canada, 1933. p. 298-299. 1933
276. _____ Fusarium species associated with diseases of cereals in Manitoba. Phytopathology 19: 7-8 (Abs.) 1939
277. _____ The occurrence of Fusarium species in Canada I. Species of Fusarium isolated from farm samples of cereal seed in Manitoba. Canad. Jour. Res. Sect. C, 22: 282-286.
278. _____, and SPRAGUE, R. Species of Fusarium associated with root rots of the Gramineae in the Northern Great Plains. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 25: 168-180. (Processed) 1941.
279. GORLENKO, M. V. (Evaluation of spring wheat varieties for resistance to Fusarium-induced diseases, powdery mildew, and brown rust) (Trans. title). Summ. sci. Res. Wk. Inst. Prot. Leningr. 1935, p. 141-143. 1936
280. GORTER, G. J. M. A. Eye-spot lodging disease of wheat. Farming in So. Africa 16: 81, 110. illus. 1941
281. GRAM, E. and S. ROSTRUP. Oversigt over sygdomme hos Landbrugets og Havebrugets Kulturplanter i 1921. Tidsskr. for Planteavl 28: 185-246. 1922
282. GRANDFIELD, C. O., C. L. LEFEEVRE, and W. H. METZGER. Relation between fallowing and the damping-off of alfalfa seedlings. Amer. Soc. Agron. Jour. 27: 800-806. 1935
283. GREANEY, F. J. The effect of phosphate deficiencies on infection of wheat by Fusarium culmorum. Canad. Jour. Res. Sect. C, 16: 27-37. 1938
284. _____ Influence of time, rate, and depth of seeding on the incidence of root rot in wheat. Phytopathology 36: 252-263. 1946
285. _____, and D. L. BAILEY. Studies in Cereal Diseases II. Root-rots and foot-rots of wheat in Manitoba. Dom. of Canad. Dept. Agr. Bul. 85 N. S. 32-pp. illus. 1927
286. _____, and J. E. MACHACEK. The production of a white fertile saltant of Helminthosporium sativum by means of ultra violet radiation. Phytopathology 23: 379-383. 1933
287. _____, and _____, and C. L. JOHNSTON. Varietal resistance of wheat and oats to root rot caused by Fusarium culmorum and Helminthosporium sativum. Sci. Agr. 18: 500-523. 1938
288. _____, and H. A. H. WALLACE. Varietal susceptibility to kernel smudge in wheat, Sci. Agr. 24: 126-134. 1943
289. GREENE, H. C. Notes of Wisconsin Fungi. II. Wis. Acad. Sci. Arts, Letters, Trans. 34: 83-98. 1942.
290. _____ Notes on Wisconsin parasitic fungi. IV. Farlowia 1: 569-581. July, 1944

291. GREGORY, C. T. Heterosporium leaf spot of timothy. *Phytopathology* 9: 576-580. 1919
292. GRIES, GEORGE A. The effect of plant-decomposition products on root diseases. *Phytopathology* 33: 1111-1112. (Abs.) 1943
293. GRIFFITH, ROBERT B. A leaf disease of Kentucky bluegrass. *Phytopathology* 33: 745. 1943
294. GRIFFITHS, R. L. Take-all: Incidence and control on the lighter soils of the Mallee. *Jour. Dept. Agr. S. Austral.* 36: 774-778. 1933
295. GROVE, W. B. Some Septorias on wheat. *Gard. Chron.* 60: 194, 210. 1916
296. _____ British stem- and leaf-fungi (Coelomycetes). Vol. I. *Sphaeropsidales*, 488 pp. illus. (Cambridge, England) 1935
297. _____ British stem- and leaf-fungi (Coelomycetes). Vol. II, 407 pp., illus. (Cambridge, England) 1937
298. GROVES, J. W. and A. J. SKOLKO. Notes on seed-borne fungi. I. *Stemphyllium*. *Canad. Jour. Res., Sect. C*, 22: 190-199. 1944
299. _____ and _____ Notes on seed-borne fungi. II. *Alternaria*. *Canad. Jour. Res. Sect. C*, 22: 217-234. 1944
300. _____ and _____ Notes on seed-borne fungi. III. *Canad. Jour. Res. Sect. C*, 23: 94-104. 1945
301. GUARCH, A. M. Comunicaciones fitopatologicas. *Rev. Fac. Agron. Univ. Montevideo* 23: 9-20. 1941. (*Rev. Appl. Mycol.* 21: 129, 1942)
302. HAENSELER, C. M. Helminthosporium leaf spot on millet in New Jersey. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 25: 486. [Processed] 1941
303. HAGBORG, W. A. F. Black chaff, a composite disease. *Canad. Jour. Res.* 14: 347-359. 1936
304. HAIGH, J. C. Macrophomina phaseoli (Maubl.) Ashby. The pycnidial stage of Rhizoctonia bataticola (Taub.) Butler. *Trop. Agr. (Ceylon)* 70: 77-79. 1928
305. HANSON, E. W. Results from disease garden and [wheat] root rot studies. In report of the Sixth Hard Spring Wheat Conference, pp. 58-59. Minneapolis, Minn. [Processed] 1939
306. _____ Root rots [wheat]: Results of fertilizer studies. In report of the Milling and Baking Meeting and Seventh Hard Spring Wheat Conference, pp. 34-37. Minneapolis, Minn. [Processed] 1944
307. _____ Studies on [wheat] scab in Minnesota. In report of the Milling and Baking Meeting and Seventh Hard Spring Wheat Conference, pp. 38-39. Minneapolis, Minn. [Processed] 1944
308. _____ and J. J. CHRISTENSEN. Insects in relation to root rot and basal stem rot of cereals. *Phytopathology* 30: 7-8. 1940
309. _____ and _____ Studies on the "Black Point" disease of wheat in the United States. *Phytopathology* 33: 1112-1113. (Abs.) 1943.

310. HARDISON, J. R. Grass diseases in Michigan in 1941. U. S. Dept. Agr., Bur. of Plant Indus., Plant Dis. Rptr. 26: 67-75. 1942
311. HARRIS, M. R. The relationship of Cephalosporium acremonium to the black bundle disease of corn. Phytopathology 26: 965-980. 1936
312. HEALD, F. D. New and little known plant diseases in Nebraska. Science n.s. 23: 624. 1906
313. _____ The bud-rot of carnation. Nebr. Agr. Expt. Sta. Bul. 103, p. 3-31. 1908
314. _____ Foot rot of wheat. Wash. Agr. Expt. Sta. Buls. 155, p. 38 (1920); 167, pp. 39-40 (1922); 175, p. 36 (1922); 180, pp. 42-44 (1923); 187, p. 71 (1924,a). 1920-24
315. _____ Sclerotium rhizodes Auers. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. Suppl. 35: 270. 1924
316. _____ Manual of Plant Diseases, 2nd Ed., pp. 1-953. McGraw-Hill Book Co. 1933
317. _____, and G. D. RUEHLE. The rots of Washington apples in cold storage. Wash. Agr. Expt. Sta. Bul. 253, pp. 25-27. illus. 1931
318. _____, E. M. WILCOX, and VENUS W. POOL. The life-history and parasitism of Diplodia zeae (Schw.) Lév. Nebr. Agr. Expt. Sta. Rpt. 22: 1-7. 1909
319. _____ and F. A. WOLF. New species of Texas fungi. Mycologia 3: 5-22. 1911
320. HEARN, J. J. Jr. Rhizoctonia solani Kühn and the brown patch disease of grass. Tex. Acad. Sci. Trans. 26: 41-42. (Abs.) 1942
321. HENRY, A. W. Root-rot of wheat. Minn. Agr. Expt. Sta. Tech. Bul. 22, 71 pp. 1924
322. _____ The natural microflora of the soil in relation to the foot-rot problem of wheat. Canad. Jour. Res. 4: 69-77. 1931
323. _____ Influence of soil temperature and soil sterilization on the reaction of wheat seedlings to Ophiobolus graminis. Canad. Jour. Res. 7: 198-203. 1932
324. _____ and W. R. FOSTER. Leptosphaeria foot-rot of wheat in Alberta. Phytopathology 19: 689-690. 1929
325. HESSE, RUDOLPH. Pythium debaryanum, ein endophytischer Schmarotzer in den Geweben der Keimlinge der Leinditter, der Rüben, des Spargels und einiger anderer landwirtschaftlicher Kulturpflanzen. Inaug. Dissert. Göttingen, 76 pp., illus. 1874
326. HITCHCOCK, A. S. Manual of the grasses of the United States. U. S. Dept. Agr. Misc. Pub. 200, 1040 pp. 1935
327. HO, W. C. Soil inhabiting fungi attacking the roots of maize. Iowa State Col. Jour. Sci. 16: 72-74. 1941
328. _____ Soil-inhabiting fungi attacking the roots of maize. Iowa Agr. Expt. Sta. Res. Bul. 332, p. 403-446. 1944
329. _____, and JAMES M. KOEPFER. Host response of maize seedlings to Pythium graminicolum (Phytopathology 32: 9. (Abs.) 1942

330. HO, W. C. and I. E. MELHUS. Succession of soil-inhabiting fungi attacking the roots of maize. *Phytopathology* 30: 10. (Abs.) 1940
331. _____, C. H. MEREDITH, and I. E. MELHUS. Pythium graminicola Subr. on barley. Iowa Agr. Expt. Sta. Res. Bul. 287. 1941
332. HOADLEY, ALFRED D. A study of the nature of resistance of dent corn to Diplodia zeae (Schw.) Lev. stalk rot. Maryland Univ. Abs. Theses 39: 11-13. 1942
333. HOFFMASTER, DONALD E. The sorghum root- and- stalk-rot complex in Oklahoma. *Phytopathology* 32: 9 (Abs.) 1942.
334. _____, J. H. McLAUGHLIN, WINFIELD W. RAY, and K. STARR CHESTER. The problem of dry rot caused by Macrophomina phaseoli (= Sclerotium bataticola). *Phytopathology* 33: 1113-1114 (Abs.) 1943
335. HOHNK, W. A new parasitic Pythium. *Mycol.* 24: 489-507. 1932
336. HOLBERT, J. R., W. L. BURLISON, B. KOEHLER, C. M. WOODWORTH, and G. H. DUNGAN. Corn root, stalk, and ear rot diseases and their control through seed selection and breeding. Ill. Agr. Expt. Sta. Bul. 255, pp. 239-478. illus. 1924
337. _____, P. E. HOPPE, and A. L. SMITH. Some factors affecting infection with and spread of Diplodia zeae in the host tissue. *Phytopathology* 25: 1113-1114. 1935
338. HOPPE, P. E. Intraspecific and interspecific aversion in Diplodia. *Jour. Agr. Res.* 53: 671-680. 1936
339. _____ Relative prevalence and geographic distribution of various ear rot fungi in the 1939 corn crop. U. S. Dept. Agr., Bur. Plant Indus., Pl. Dis. Rptr. 24: 210-213. [Processed] 1940
340. HORSFALL, JAMES G. A study of meadow-crop diseases in New York. Cornell Agr. Expt. Sta. Mem. 130. 139 pp. 1930
341. HOUSTON, B. R. Culture types and pathogenicity of isolates of Corticium solani. *Phytopathology* 35: 371-393. 1945
342. HUANG, T. W. The influence of Helminthosporium setariae upon the growth and yield of Chinese corn. *Entom. and Phytopath.* Hangchow, China 3: 234-238. (In Chinese) 1935
343. HUNGERFORD, C. W. A serious disease of wheat caused by Sclerotium rhizodes in Idaho. *Phytopathology* 13: 463-464. 1923
344. HYNES, H. J. 'Purple patch' of wheat and oats. A disease caused by the fungus Rhizoctonia solani. *Agr. Gaz. N. S. Wales* 44: 879-883, illus. 1933
345. _____ Helminthosporium root-rot of wheat and other cereals. Part I Economic importance, symptoms and causal organisms. Part 2 Physiologic specialisation in Helminthosporium spp. *Sci. Bul. Dept. Agr. N. S. Wales* 47, 39 pp. illus. 1935
346. _____ Technical notes. Studies on 'take-all' of wheat. I. *Jour. Austral. Inst. Agr. Sci.* 3: 43-48. 1937
347. _____ Studies on Rhizoctonia root-rot of wheat and oats. *Sci. Bul. Dept. Agr. N. S. Wales* 58, 42 pp. illus. 1937

348. HYNES, H. J. Species of Helminthosporium and Curvularia associated with root rot of wheat and other graminaceous plants. Jour. and Proc. Roy. Soc. N. S. Wales 70: 378-391, illus. 1937 (Rev. Appl. Mycol. 16: 735. 1937).
349. _____ Studies on Helminthosporium root-rot of wheat and other cereals. Part 3. Factors influencing infection. Part 4 The control problem. Sci. Bul. Dept. Agr. N. S. Wales 61, 67 pp., illus. 1938. (Rev. Appl. Mycol. 17: 805. 1938).
350. IMAI, SANSHI. On the Clavariaceae of Japan (Trans. title). Sapporo Nat. Hist. Soc. Trans. 11: 38-44. 1929
351. _____ On the causal fungus of the Typhula-blight of gramineous plants (Trans. title). Japanese Jour. Bot. 8: 5-18. 1936. (Rev. Appl. Mycol. 15: 347. 1936)
352. IMMER, F. R. and J. J. CHRISTENSEN. Studies on susceptibility of varieties and strains of barley to Fusarium and Helminthosporium kernel blight when tested under muslin tents or in nurseries. Jour. Amer. Soc. Agron. 35: 515-522. 1943
353. ITO, S. On some new ascigerous stages of the species of Helminthosporium parasitic on cereals. Imper. Acad. Japan, Proc. 6: 352-355. 1930
354. _____, and K. KURIBAYASHI. The ascigerous forms of some graminicolous species of Helminthosporium in Japan. Jour. Facul. Agr. Hokkaido Imp. Univ., Sapporo, 29: 85-125 (English). 1931
355. _____, and Y. TOKUNAGA. Notae mycologicae Asiae Orientalis I. Sapporo Nat. Hist. Soc. Trans. 14: 11-33. 1935. (Rev. Appl. Mycol. 15: 57-58. 1936).
356. IWAYAMA, S. On a new snow-rot disease of cereal plants caused by Pythium sp. Pamphlet issued by Agr. Exp. Sta. Toyama-Ken. (Japan) 20 pp., illus. 1933
357. JAAP, OTTO. Beitrage pilzflora der Schweiz. Ann. Mycol. (Sydow) 5: 246-272. 1907
358. JAARVELD, ALIDA. Der Einfluss verschiedener Bodenpilze auf die Virulenz von Rhizoctonia solani Kühn. Phytopath. Ztschr. 14: 1-75. illus. 1942. (Rev. Appl. Mycol. 22: 495. 1943).
359. JACQUES, J. EMILE. Studies in the genus Heterosporium. Contrib. Inst. Bot. Univ. Montreal 39: 36-38, illus. 1941
360. JANCZEWSKI, E. Cladosporium herbarum i jego najpospolisze na zbozu towarzysze (Recherches sur le Cladosporium herbarum et ses compagnons habituels sur les cereales.) Bul. Internat. Acad. Sci. Cracovie 27: 187-208. 1894
361. JANKOWSKA, KRZYSTYNA. Spostrzennia nad wystepowaniem chorab roslin uprawnych w woj. Lubelskiem w latach 1927 i 1928. Mem. Inst. Nat. Polonais d'Economie Rurale a Pulaway 9: 574-595. 1929
362. JENSEN, C. N. Fungus flora of the soil. Cornell Univ. Agr. Exp. Sta. Bul. 315: 413-501. 1912
363. JOHANN, HELEN. Penicillium injury to corn seedlings. Phytopathology 18: 239-242. 1928
364. _____ Grated carrot agar favorable for studies of Pythium. Phytopathology 18: 710. 1928

365. JOHANN, HELEN. Phoma terrestris in the roots of mature maize plants. Phytopathology 33: 526-528. 1943
366. _____, JAMES R. HOLBERT, and J. G. DICKSON. Further studies on Penicillium injury to corn. Jour. Agr. Res. 43: 757-790, illus. 1931
367. JOHNSON, A. G. The ascigerous stage of Helminthosporium teres Sacc. Phytopathology 4: 408. (Abs.) 1914
368. _____ Septoria on barley. Phytopathology 6: 117. (Abs.) 1916
369. _____ Helminthosporium tritici-vulgaris on wheat in Maryland. U. S. Dept. Agr., Bur. Plant. Indus. Plant Dis. Rptr. 26: 246-247. 1942
370. _____, and C. W. HUNGERFORD. Scolecotrichum graminis on timothy, orchard grass and other grasses. Phytopathology 7: 69. (Abs.) 1917
371. _____, and W. W. MACKIE. Evidence of disease resistance in barley to attacks of Rhynchosporium. Phytopathology 10: 54 (Abs.) 1920
372. JOHNSON, E. C. A study of some imperfecti fungi isolated from wheat, oat and barley plants. Jour. Agr. Res. 1: 475-489. illus. 1914
373. JOHNSON, THORVALDUR. Studies on the pathogenicity and physiology of Helminthosporium gramineum Rab. Phytopathology 15: 797-804. 1925
374. _____, and W. A. F. HAGEBORG. Brown necrosis and Alternaria blotch of wheat. Sci. Agr. 22: 746-760, illus. 1942
375. JOHNSTON, C. L. and F. J. GREANEY. Studies on the pathogenicity of Fusarium species associated with root rot of wheat. Phytopathology 32: 670-684. 1942
376. JØRSTAD, I. Beretning om plantesykdommer i land- og hagebruket. VI. Sykdommer på korn- og engvekster. [Norway] Landbr. Dir. Beret. 1929. Tillegg C. Beret. Skadeinsekt. og Plantesykd. 84 pp. 1930
377. KARLING, JOHN S. The simple holocarpic biflagellate Phycomycetes. (New York). 123 pp. 1942
378. KARSTEN, P. A. Symbolae ad mycologiam fennicam XVI. Medd. Soc. Faun. Fl. Fenn. 11: 148-161. 1884
379. KEIL, H. L. New fungicide developments for turf. Greenkeepers' Rptr. 12: 5-6. 1944
380. KENDRICK, J. B., and F. N. BRIGGS. Pythium root rot of milo and the development of resistant varieties. Calif. Agr. Expt. Sta. Bul. 629, 18 pp. 1939
381. KENT, G. C. The production of spores of Diplodia zeae in culture. Phytopathology 32: 12. (Abs.) 1942
382. KINSEL, KATHERINE. Carbohydrate utilization by the corn Diplodias. Phytopathology 27: 1119-1120. 1937
383. KIRBY, R. S. The take-all disease of cereals and grasses caused by Ophiobolus cariceti (Berkeley and Broome) Saccardo. Cornell Univ. Agr. Expt. Sta. Memoir 88, 45 pp. 1925
384. KIRSCHSTEIN, W. Über neue, seltene und kritische Ascomyceten und Fungi Imperfecti, II. Ann. Mycol. 37: 88-140. 1939

385. KOEHLER, B. Natural mode of entrance of fungi into corn ears and some symptoms that indicate infection. Jour. Agr. Res. 64: 421-422. illus. 1942
386. _____ Disease threatening broom corn production in Illinois. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 27: 70-73. 1943. [Processed]
387. _____, and G. H. DUNGAN. Disease infection and field performance of bin- and hanger-dried seed corn. Amer. Soc. Agron. Jour. 32: 768-781. illus. 1940
388. _____, and JAMES R. HOLBERT. Combatting corn diseases in Illinois. Ill. Agr. Expt. Sta. Cir. 484, 36 pp. 1938
389. KOTTE, W. Die Federbuschsporen-Krankheit des Getreides. Nachr. Schädlingsbekampf. Leverkusen, 9 (4): 170-174. illus. 1934
390. KRAUSE, F. Scolecotrichum graminis Fuck. f. Avenae Erikss. Centbl. f. Bakt. Abt. II, 25: 102-106. 1909
391. KREITLOW, K. W. Sclerotium rhizodes on grasses in Pennsylvania. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 26: 360-361. 1942. [Processed]
392. KREUTZER, W. A. Host-parasite relationships in pink root of Allium cepa. II. The action of Phoma terrestris on Allium cepa and other hosts. Phytopathology 31: 907-915. 1941
393. KUEHN, JULIUS. Phoma hennebergii nov. spec. Landw. Versuchsst. 21: 193-194. 1878
394. KURIBAYASHI, K. The ascigerous stage of Helminthosporium sativum [Trans. title]. Sapporo Nat. Hist. Soc. Trans. 10: 138-145. 1929
395. LAAR, J. H. J. van de. (Investigations on Ophiobolus graminis Sacc. and Ophiobolus herpotrichus (Fr.) Sacc. and on the diseases of Triticum vulgare Vill. and other Gramineae caused by fungi.) [Trans. title] (French summary). illus. Wageningen. 160 pp. 1931
396. LAL, A. Interaction of soil micro-organisms with Ophiobolus graminis Sacc., the fungus causing the take-all disease of wheat. Ann. Appl. Biol. 26: 247-261. illus. 1939
397. LANDALUZE, P. URQUIJO. (In) Memoria de los trabajos por la Estación de Fitopatología agrícola de la Coruña. Anos 1939-40. Publ. Estac. Fitopat. Agr. Coruña 14, 35 pp., illus. 1941
398. LARSH, HOWARD W. Relative prevalence of Diplodia zeae and Diplodia macrocarpa on corn. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 22: 159-162. 1938. [Processed]
399. LEDINGHAM, R. J. Observations on antagonism in inoculation tests of wheat with Helminthosporium sativum P. K. and B. and Fusarium culmorum (W. G. Sm.) Sacc. Sci. Agr. 22: 688-697. 1942
400. LEFEBVRE, C. L., and H. W. JOHNSON. Collections of fungi, bacteria, and nematodes of grasses. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 25: 556-579. 1941. [Processed]
401. _____, and HELEN S. SHERWIN. Races of Helminthosporium turcicum. Phytopathology 35: 487. (Abs.). 1945
402. LEONIAN, L. H. The pathogenicity of and variability of Fusarium moniliforme from corn. West Va. Agr. Expt. Sta. Bul. 248, 16 pp., illus. 1932

403. LEUKEL, R. W. Chloropicrin as a disinfectant for plant beds. *Phytopathology* 32: 1034-1036. illus. 1942
404. _____, J. G. DICKSON, and A. G. JOHNSON. Effects of certain environmental factors on stripe disease of barley and the control of the disease by seed treatment. U. S. Dept. Agr. Tech. Bul. 341, 39 pp. 1933
405. _____, and J. H. MARTIN. Seed rot and seedling blight of sorghum. U. S. Dept. Agr. Tech. Bul. 839, 26 pp. 1943
406. LIND, J. Bemerkenswerte Pilzfunde in Dänemark. *Ann. Mycol.* (Sydow) 5: 272. 1907
407. _____ Fungi (Micromycetes) collected in Arctic North America (King William Land, King Point and Herschell Isl.) by the Gjöa Expedition under Captain Roald Amundsen 1904-1906. *Vidensk.-Selsk. Christiania, Skr. I. Math.-Nat.* 1909 (9): 1-25. illus. 1910
408. _____ Danish Fungi as represented in the Herbarium of E. Rostrup (No. 3117, see tab. IX). Copenhagen. 1913
409. LIVINGSTON, J. E. Charcoal rot of corn and sorghum in Nebraska. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 26: 50-52. 1942. [Processed]
410. _____ Charcoal rot -- threat to Nebraska corn and sorghums. *Nebr. Crop Impr. Assoc.* 33d Ann. Rpt. 1942. In *Nebr. State Board Agr. Ann. Rpt.* 1942: p. 206. 1942
411. LOBIK, A. I. (Materials for mycological flora of Terek district) (Trans. title). *Morbi Plantarum (Bolezni Rastenii)* 17: 178. illus. 1928
412. LUCHETTI, G. Su di alcune micosi delle spighe di frumento e sulla biologia di alcuni funghi da queste isolati. *Annali Fac. Agr. R. Univ. Pisa* n. s. 2, 337-461. illus. 1939
413. LUDEROCK, W. V. Wojnowicia graminis (McAlp.) Sacc. and D. Sacc. in relation to foot rot of wheat in Australia. *Commonwealth of Australia (Council for Sci. and Ind. Res.) Bul.* 103, 23 pp. 1936. (Melbourne)
414. _____ Root amputation experiments with wheat under dry conditions, in relation to attack by Ophiobolus graminis, *Jour. Council Sci. Industr. Res. Austral.* 15 (2): 129-134. 1942
415. LUDWIG, R. A., and A. W. HENRY. Studies on the microbiology of recontaminated sterilized soil in relation to its infestation with Ophiobolus graminis Sacc. *Canad. Jour. Res. Sect. C*, 21: 348-350. illus. 1943
416. LUTHRA, J. C., A. SATTAR, and M. A. GHANI. A comparative study of species of Septoria occurring on wheat. *Indian Jour. Agr. Sci.* 7: 271-289. illus. 1937
417. _____, _____, and _____ Perpetuation and control of Septoria disease of wheat in the Punjab. *Agr. Live-Stk. (India)* 8: 17-25. 1938
418. MACHACEK, J. E. An estimate of loss in Manitoba from common root rot of wheat. *Sci. Agr.* 24: 70-77. 1943

419. MACHACEK, J. E. The prevalence of Septoria on cereal seed in Canada. *Phytopathology* 35: 51-53. 1945
420. _____, and F. J. GREANEY. The "black-point" or "kernel smudge" disease of cereals. *Canad. Jour. Res. Sect. C*, 16: 84-113. 1938
421. _____, and _____. Further experiments on the control by seed disinfection of root-rotting fungi in wheat. *Phytopathology* 31: 379-394. 1941
422. MAC INNES, J. The growth of the wheat scab organism in relation to hydrogen-ion concentration. *Phytopathology* 12: 290-294. 1922
423. MACKIE, W. W. Footrot or Ophiobolus in California. *Phytopathology* 13: 561-562. 1923
424. _____ Inheritance of resistance to Helminthosporium californicum in a cross between Chevalier barley, a resistant variety, and Abyssinian, a susceptible variety. *Phytopathology* 16: 764. (Abs.) 1926
425. _____ Inheritance of resistance to barley scald. *Phytopathology* 19: 1141. (Abs.). 1929
426. _____ A hitherto unreported disease of maize and beans. *Phytopathology* 22: 637-644. 1932
427. _____, and G. E. PAXTON. A new disease of cultivated barley in California caused by Helminthosporium californicum n. sp. *Phytopathology* 14: 124. (Abs.). 1924
428. MAIRE, R. Contributione à l'étude de la flore mycologique de l'Afrique du Nord. *Bul. Soc. Bot. France* 53: 187. 1906
429. _____, and R. WERNER. Catalogue raisonne des Champignons connus jusqu' ici au Mem. Soc. Sci. Nat. du Maroc. *Mem.* 45: 148 pp. 1937 (1938).
430. MAJDRAKOFF, V. P. Versuche mit der Streifenkrankheit der Gerste. *Bot. Archiv.* 34: 337-362. 1932
431. MALAN, C. E. La Calonectria graminicola (Berk. and Brme.) Wr. sulle cariosidi di Segala della zona alpina (Alpi Cozie). *Nuovo Gior. Bot. Ital. n. s.* 46: 323-326. 1939.
432. MANGIN, L. Sur le piétin ou maladie du pied du blé. *Bul. Soc. Mycol. de France* 15: 210-239. 1899
433. MARCHIONATTO, J. B. El "verdin" de maiz (Penicillium viridicatum). *Buenos Aires Univ. Facult. de Agron. y Vet. Rev.* 9: 159-169. 1942
434. MARKEVICH, N. P. Perezimovka i Porazhaemost Snezhnoi Plesenyu Ekotipov Ozimoi Pshenitsy. (The overwintering and injuring by the snowy mould of the winter wheat ecotypes). *Bul. Plant Protect. (Leningrad)* 1 (20): 119-121. [In Russian]. 1939. (Rev. Appl. Mycol. 19: 398)
435. MARTIN, A. L. Rice straw stacks as a source of infection with the black kernel disease. *U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr.* 23: 247-249. (Processed) 1939

436. MARTIN, A. L., and G. E. ALTSTAT. Black kernel and white tip of rice. Tex. Agr. Expt. Sta. Bul. 584, 14 pp., illus. 1940
437. MARTYN, E. B. Report on the Botanical and Mycological Division for the year 1935. Div. Rept. Dept. Agr. Brit. Guiana, 1935, p. 89-92. 1936
438. MASON, E. W. On species of the genus Nigrospora Zimmermann, recorded on Monocotyledons. Brit. Mycol. Soc. Trans. 12: 152-165. 1927
439. _____ Annotated account of fungi received at the Imperial Bureau of Mycology. List II (Fascicle 1). The Imperial Bur. Mycol. Kew. 43 pp. 1928
440. MATTHEWS, V. D. Studies on the genus Pythium. 136 pp. Univ. of North Carolina Press, Chapel Hill, illus. 1931
441. MATSUURA, I. Comparative studies on four Hyphomycetes pathogenic to rice seedlings. (Japanese). Jour. Microbiol. Soc. 21: 1551-1572. 1927
442. MATZ, J. The Rhizoctonias of Porto Rico. Jour. Dept. Agr. Porto Rico 5, 30 pp. illus. 1921
443. MC ALPINE, D. Take-all and white-heads in wheat. Jour. Dept. Agr. Victoria 2 (5): 410-426. 1904. Reprinted as Victoria Dept. Agr. Bul. 9, 20 pp.
444. MC INDOE, D. G. The inheritance of the reaction of maize to Gibberella saubinetii. Phytopathology 21: 615-639. 1931
445. MC KINNEY, H. H. Influence of soil temperature and moisture on infection of wheat seedlings by Helminthosporium sativum. Jour. Agr. Res. 26: 195-218. 1923
446. _____ Foot-rot diseases of wheat in America. U. S. Dept. Agr. Bul. 1347, 40 pp., 1925
447. _____, and R. J. DAVIS. Influence of soil temperature and moisture on infection of young wheat plants by Ophiobolus graminis. Jour. Agr. Res. 31: 827-840. 1925
448. _____, and A. G. JOHNSON. Wojnowicia graminis (McAlp.) Sacc. and D. Sacc. on wheat in the United States. Phytopathology 11: 505-506. 1921
449. MC LAUGHLIN, J. H., J. HARVEY, and I. E. MELHUS. The response of some field crops on soil treated with chloropicrin. Iowa State Col. Jour. Sci. 17: 213-220. 1943.
450. MC MURPHY, J. A Phytophthora on oats. Science n. s. 43: 534. 1916
451. MC NEW, G. L. Crown infection of corn by Diplodia zeae. Iowa Agr. Expt. Sta. Res. Bul. 216: 187-222. 1937
452. MC RAE, W. Report of the Imperial Mycologist. Pusa Imper. Inst. Agr. Res. Sci. Rpts., 1930-31, p. 73-86. 1932
453. MEAD, H. W. Host-parasite relationships in a seed-borne disease of barley caused by Helminthosporium sativum Pammel, King and Bakke. Canad. Jour. Res. Sect. C, 20: 501-523. 1942
454. _____ Environmental relationships in a seed-borne disease of barley caused by Helminthosporium sativum Pammel, King and Bakke. Canad. Jour. Res. Sect. C, 20: 525-538. 1942.
455. MELCHERS, L. E. On the cause of the milo disease. Phytopathology 32: 640-641. 1942

456. MELCHERS, L. E., and A. E. LOWE. The reaction of sorghum varieties and hybrids to milo disease. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. Suppl. 126, pp. 165-175. 1940. [Processed]
457. _____, and _____. The development of sorghums resistant to milo disease. Kans. Agr. Expt. Sta. Tech. Bul. 55, 24 pp. 1943
458. _____, and M. C. SEWELL. The effect of tillage, fertilizers, and rotations on the spread of wheat foot-rot. Phytopathology 16: 81. (Abs.). 1926
459. MELHUS, I. E., Pathology and mycology of corn. Rept. Iowa Agr. Expt. Sta. 1942-43, Part II, pp. 52-57. 1944 (?). (Rev. Appl. Mycol. 23: 331-333).
460. _____, J. N. MARTIN, and H. C. MURPHY. Botany and Plant Pathology Section. Rep. Iowa Agr. Expt. Sta. 1942-43, Part I, pp. 125-145, 1943 (Rev. Appl. Mycol. 23 (9): 330-331. 1944) illus.
461. _____, _____, C. S. REDDY, W. F. BUCHHOLTZ, and H. C. MURPHY. Botany and plant pathology section. Iowa Agr. Expt. Sta. 1939-40 Rpt., Pt. I, pp. 132-151. 1940
462. MEREDITH, CLIFFORD H. Phycomycetes in Iowa soil. Phytopathology 28: 15-16. (Abs.) 1938
463. _____ A quick method of isolating certain phycomycetous fungi from soil. Phytopathology 30: 1055-1056. 1940
464. MIDDLETON, JOHN T. Root rot of barley caused by Pythium hypogynum n. sp. Phytopathology 31: 863. (Abs.) 1941
465. _____ The taxonomy, host range and geographic distribution of the genus Pythium. Torrey Bot. Club Mem., 20, No. 1, 171 pp. illus. 1943
466. MITRA, M. Helminthosporium spp. on cereals and sugar cane in India, Part. I. (Diseases of Zea mays and Sorghum vulgare caused by species of Helminthosporium). India Dept. Agr. Mem. Bot. Ser. 5: 219-242, illus. 1923
467. _____ Saltation in the genus Helminthosporium. Brit. Mycol. Soc. Trans. 16: 115-127, illus. 1931
468. _____ A comparative study of species and strains of Helminthosporium on certain Indian cultivated crops. Brit. Mycol. Soc. Trans. 15: 254-293, illus. 1931. (Rev. Appl. Mycol. 10: 758-759, 1931).
469. _____ A leaf spot disease of wheat caused by Helminthosporium tritici-repentis Died. Indian Jour. Agr. Sci. 4: 692-700, illus. 1934
470. _____ and MEHTA, P. R. Diseases of Eleusine coracana Gaertn., and E. aegyptiaca Desf. caused by species of Helminthosporium. Indian Jour. Agr. Sci. 4: 943-975. 1934
471. MIURA, M. Manshu Shuyo Nosakumotsu no Byogai (Diseases of Principal Crops in Manchuria) (Koshurei) So. Manchuria Ry. Co. Agr. Expt. Sta. Rpt. No. 11, i-ii + 56 pp. (In Japanese).
472. MONTEITH, JOHN JR. and ARNOLD S. DAHL. Turf diseases and their control. Bul. U. S. Golf Assoc. Green Sect. 12: 85-186. 1932
473. MORINI, F. Alcune osservazioni sopra una nuova malattia del frumento. Nuovo Gior. Bot. Ital. 18: 32-43. 1886

474. MORITZ, O. Entstehungsbedingungen und Verhütungsmöglichkeiten der Ophiobolose des Weizens. Vorläufige Mitteilung. Deut. Pflanzenschutzdienst Nachrichtenbl. 11: 100-101. 1931
475. _____ Weitere Studien über die Ophiobolose des Weizens. Biol. Reichsanst. f. Land. u. Forstw. Arb. 20: 27-48, illus. 1932
476. _____ and H. BOCKMANN. Einleitende Studien über Cercospora herpotrichoides Fron. Angew. Bot. 15: 409-419, illus. 1933
477. MÜLLER, K. D. Über das Auftreten der Typhula-fäule am Wintergerste und Roggen in der Provinz Sachsen. Deut. Landw. Presse .. 57: 234. 1930
478. MÜLLER-KÖGLER, E. Untersuchungen über die Schwarzbeinigkeit des Getreides und den Wirtspflanzenkreis ihres Erregers (Ophiobolus graminis Sacc.) Biol. Reichsanst. f. Land. u. Forstw. Arb. Berl. 22: 271-319, illus. 1938
479. MUSKETT, A. E. A study of the epidemiology and control of Helminthosporium disease of oats. Ann. Bot. n. s. 1: 763-784, illus. 1937
480. NATTRASS, R. M. Take-all disease of cereals (Ophiobolus graminis). E. Afr. Agr. Jour. 8: 133-135, illus. 1943
481. NEWHALL, A. G. A serious storage rot of celery caused by the fungus Anasopora macrospora n. gen. Phytopathology 34: 92-105. 1944
482. NIGAM, B. S. Physiology of zonation---effect of light and temperature on zonation in Acrothecium lunatum Wakker. Jour. Indian Bot. Soc. 15: 115-123, illus. 1936
483. NILSSON, F. Råg försök och rågförädling vid Sveriges Utsädesförening Västernorrlandsfilial. Sveriges Utsädesför. Tidskr. 1: 4-30. 1940
484. NILSSON-EHLE, H. Iakttagelser öfver hafresorters olika mottaglighet för Scölecotrichum-eller fläcksjukan., Tidskr. för Landtmän 29: 817-820; 827-834; 845-848. 1908
485. NISIKADO, Y. Temperature relations to the growth of graminicolous species of Helminthosporium: I. Effect of temperature on the vegetative and reproductive growth of Helminthosporium turcicum Passer. and Helminthosporium maydis Nisikado et Miyake Ber. O'Hara Inst. Landwirts. Forsch. in Kuraschiki 3: 349-377, (English). 1927
486. _____ Studies on the Helminthosporium diseases of Gramineae in Japan (Trans. title). O'Hara Inst. Agr. Research, Spec. Rpt. No. 4, 394 pp., illus. 1928. (English summary in Ohara Inst. f. Landw. Forsch. Ber. 4: 111-126, illus. 1929. (E. S. R. 64: 448. 1931).
487. _____ and C. MIYAKE. Studies on two helminthosporium diseases of maize caused by Helminthosporium turcicum Passerini and Ophiobolus heterostrophus Drechsler (H. maydis Nisikado et Miyake). O'Hara Inst. f. Landw. Forsch. Ber. 3: 221-226. 1926
488. NOBLE, R. J., H. J. HYNES, F. C. McCLEERY, and W. A. BIRMINGHAM. Plant diseases recorded in New South Wales. N. S. Wales Dept. Agr. Sci. Bul. 46, 18 pp. 1934
489. NOLL, A. Experimentelle Untersuchungen über den Befall des Weizens mit Penicillium. Phytopath. Ztschr. 9: 147-186, illus. 1936. (Rev. Appl. Mycol. 15: 640).

490. NOLL, W. El pietin del Trigo (Ophiobolus graminis Sacc.) en el Uruguay. Informe preliminar. Arch. Fitotec. Uruguay 3: 96-101, illus. (English and German summaries). 1938
491. O'BRIEN, G. D. and R. W. G. DENNIS. The Helminthosporium disease of oats. The west of Scotland Agr. Col. Res. Bul. 3: 74 pp. illus. 1933.
492. O'GARA, P. J. A Podosporiella disease of germinating wheat. Phytopathology 5: 323-326, illus. 1915
493. _____ A Phoma disease of western wheatgrass. Science n. s. 43: 110-111. 1916
494. OORT, A. J. P. De Oogvlékkenziekte van de granen veroorzaakt door Cercospora herpotrichoides Fron. (The eye-spot disease of cereals caused by Cercospora herpotrichoides Fron) Tijdschr. over Plantenziekten 42 (7): 179-234. illus. (English summary) 1936
495. ORTON, C. R. Graminicolous species of Phyllachora in North America. Mycologia 36 (1): 18-53. 1944
496. OSBORN, T. G. B. Take-all in oats. Jour. Dept. Agr. So. Austral. 22: 519. 1919
497. OWENS, C. E. Principles of plant pathology. 929 pp., illus. John Wiley and Sons, Inc. 1928
498. PADWICK, G. W. Influence of wild and cultivated plants on the multiplication, survival and spread of cereal foot-rotting fungi in the soil. Canad., Jour. Res. Sect. C. 12: 575-589. 1935
499. _____ Biologic strains of Ophiobolus graminis Sacc. Ann. Appl. Biol. 23: 45-46. 1936
500. _____ The genus Fusarium. VI. A recent attempt at mass revision. Indian Jour. Agr. Sci. 11: 663-674. 1941
501. _____ and A. W. HENRY. The relation of species of Agropyron and certain other grasses to the foot-rot problem of wheat in Alberta. Canad. Jour. Res. Sect. C. 8: 349-363. 1933
502. PAMEL, L. H., C. M. KING, and A. L. BAKKE. Two barley blights, with comparison of species of Helminthosporium upon cereals. Iowa Agr. Expt. Sta. Bul. 116, pp. 178-190. illus. 1910
503. PASSERINI, G. La nebbia dei cereali (abstract). Nuovo Gior. Bot. Ital. 9: 111. 1877
504. _____ (In Rabh. Fungi. Eur. n. 2452). 1879
505. _____ Funghi parmensi enumerati. Atti della Soc. Crittog. Ital. (Milano) 2: 20-47. 1879
506. PAXTON, G. E. Studies on Helminthosporium species found on cultivated barley in California. Phytopathology 12: 446-447. (Abs.) 1922
507. PEARSON, N. L. Parasitism of Gibberella saubinetii on corn seedlings. Jour. Agr. Res. 43: 569-596. 1931
508. PELTIER, G. L. Parasitic Rhizoctonias in America. Ill. Agr. Expt. Sta. Bul. 189, pp. 279-390. 1916
509. PENZIG, O. II Fungi della mortola. Atti R. Istit. Venet. Sec. 6a, 2, p. 652. 1884
510. PETRAK, F. Mykologische notizen VIII. Ann. Mycol. (Sydow) 23: 86, 92-93, 107. 1925
511. _____ Mykologische notizen X. Ann. Mycol. (Sydow) 27: 324 (392). 1929
512. _____ and H. SYDOW. Kritisch-systematische Originaluntersuchungen über Pyrenomyzeten, Sphaeropsiden und Melanconieen. Ann. Mycol. (Sydow) 23: 209 (211). 1925
513. PETRI, L. Un'estesa infezione di Pythium su piante di Grano. R. Staz. Patol. Veg. Bol. N.S. 10: 285-301, illus. 1930
514. PEYRONEL, B. La "puntatura" dello scudetto nelle cariossidi del frumento. [Roma] R. Staz. di Patol. Veg. Bol. 6: 10-25. 1926

515. PEYRONEL, B. Il "mal del piede" dei cereali. R. Staz. di Patol. Veg. Bol. 6: 285-336. 1926. (Rev. Appl. Mycol. 6: 342. 1927).
516. PINCK, L. A. and F. E. ALLISON. The synthesis of lignin-like complexes by fungi. Soil. Sci. 57: 155-161. 1944
517. RAABE, A. Helminthosporium tritici-vulgaris Nisikado, Erreger einer Blattfleckenkrankheit des Weizens. Phytopath. Ztschr. 10: 111-112. 1937. (Rev. Appl. Mycol. 16: 664. 1937)
518. RAILLO, A. I., Mme. Diagnosticheskaja otsenka morfologicheskikh i kultural nykh priznakov u vidov roda Fusarium. (Diagnostic estimation of morphological and cultural characters of species in the genus Fusarium). Trudy po Zashch. Rast. II. Ser. Fitopat. (Bul. Plant Protect. II. Ser. Phytopath.) No. 7, 99 pp., illus., 1935
519. _____ Fusarium -- Taxonomy of the genus Fusarium and a method for the determination of the species belonging to it. Acta Inst. Bot. Acad. Sci. U. S. S. R. Ser. II. (Pl. Cryptogamae), 1936, 3, pp. 803-857, illus. 1936. (English summary)
520. RAINIO, A. J. Ueber die Dilophospora-Krankheit von Phleum pratense L. und Alopecurus pratensis L. Valtion Maatalouskoe-toiminnam Julkaisu. 87: 5-29. 1936
521. RALEIGH, W. P. Infection studies of Diplodia zeae (Schw.) Lév. and control of seedling blights of corn. Iowa Agr. Expt. Sta. Res. Bul. 124, 91-121 pp., 1930
522. RAMAKRISHNAN, T. S. Root-rot of sugarcane. Current Science 10: 254-255. illus. 1941
523. RANDS, R. D., and E. DOFP. Variability in Pythium arrhenomanes in relation to root rot of sugarcane and corn. Jour. Agr. Res. 49: 189-(222). illus. 1934
524. _____, and _____ Pythium root rot of sugarcane. U. S. Dept. Agr. Tech. Bul. 666, 95 pp., illus. 1938
525. _____, and _____ Influence of certain harmful soil constituents on severity of Pythium root rot of sugar cane. Jour. Agr. Res. 56: 53-68. 1938
526. RATHSCHLAG, H. Studien über Helminthosporium avenae. Phytopath. Ztschr. 2: 469-492. 1930. (Rev. Appl. Mycol. 10: 233-235. 1930).
527. RAVN, F. KØLPIN. Nogle Helminthosporium-arter Og de af dem fremkaldte sygdomme hos byg og havre. Bot. Tidsskr., Bd. 23, pp. 101-322. illus. 1900
528. REDDY, C. S. Resistance of dent corn to Basisporium gallarum Moll. Iowa Agr. Expt. Sta. Res. Bul. 167. pp. 1-40. 1933
529. _____, and L. C. BURNETT. Development of seed treatments for the control of barley stripe. Phytopathology 20: 367-390. 1930
530. _____, and J. R. HOLBERT. The black-bundle disease of corn. Jour. Agr. Res. 27: 177-205. 1924.
531. REINKING, O. A. Philippine economic-plant diseases. Philippine Jour. Sci., A, 13: 165-274. illus. 1918
532. REMSBERG, RUTH E. Studies in the genus Typhula. Mycologia 32: 52-96. 1940
533. _____ The snow molds of grains and grasses caused by Typhula itoana and Typhula idahoensis. Phytopathology 30: 178-180. 1940

534. REMSBERG, RUTH E., and C. W. HUNGERFORD. Certain sclerotium diseases of grains and grasses. *Phytopathology* 23: 863-874. 1933.
535. RICHARDSON, J. K. Studies on root rot of corn in Ontario. *Canad. Jour. Res., Sect. C*, 20: 241-256. 1942. (Processed)
536. RIVERA, V. Osservazioni sopra la recettività di alcune varietà di frumento per la Septoria graminum Desm. *R. Staz. di Patol. Veg. Bol. n. s.* 8: 248-257. 1923
537. ROBERTSON, H. T. Maturation of foot and root tissue in wheat plants in relation to penetration by Ophiobolus graminis Sacc. *Sci. Agr.* 12: 575-592. 1932
538. ROGERS, D. P. The genus Pellicularia (Thelephoraceae). *Farlowia* 1: 95-118. 1943
539. ROLDAN, E. F. The occurrence of Pythium root rot disease of maize and sugar cane in the Philippine Islands. *Philippine Agr.* 19: 327. 1930
540. _____ Pythium root disease of corn in the Philippine Islands. *Philippine Agr.* 21: 165-176. 1932
541. ROSE, JESSIE P. Fusarium culmorum var. leteus, a cause of disease in cereals and grasses. *Phytopathology* 14: 50. (Abs.). 1924
542. ROSELLA, E. Sur deux cas de Septoriose au Maroc. *Rev. Path. Vég. et Ent. Agr.* 16: 211-213. 1929
543. _____ Quelques observations sur la moucheture des céréales. *Rev. Path. Vég. et Ent. Agr.* 17: 338-344. 1930
544. ROSEN, H. R. Septoria glume blotch of wheat. *Ark. Agr. Expt. Sta. Bul.* 175, 17 pp., 1921
545. ROSTRUP, F. G. E. *Fungi Groenlandiae. Oversigt over Grønlands Svampe (Meddleser om Grønlands) Kjøbenhavn III*, pp. 515-590. 1888
546. RUEHLE, G. D. Cladosporium species from apple fruit and the perfect stage of Cladosporium herbarum Lk. *Phytopathology* 20: 854. (Abs.). 1930
547. _____ New apple-rot fungi from Washington. *Phytopathology* 21: 1141-1152. illus. 1931
548. RUSSELL, R. C. Studies of take-all and its causal organism, Ophiobolus graminis Sacc. *Dom. Canada Dept. Agr. Bul.* 170, n. s., 64 pp. 1934
549. _____ Pathogenicity tests with cultures of Ophiobolus graminis Sacc. *Sci. Agr.* 19: 622-699. 1939
550. _____ The relative importance, from the pathological standpoint, of two types of smudge on wheat kernels. *Sci. Agr.* 23: 365-375. 1943
551. _____, and R. J. LEDINGHAM. Wheat seed testing from the pathological standpoint with special reference to embryo exposure. *Sci. Agr.* 21: 761-775. 1941
552. _____, and B. J. SALLANS. The effect of phosphatic fertilizers on common rootrot. *Sci. Agr.* 21: 44-51. 1940
553. RYKER, T. C. The Rhizoctonia disease of Bermuda grass, sugar cane, rice and other grasses in Louisiana. *Internatl. Soc. Sugar Cane Technol. Proc. Sixth Congr.* 1938, pp. 198-201, illus. 1939

554. SACCARDO, P. A. *Fungi Veneti novi vel critici mycologiae*.
Venetae Addendi VII. *Michelia* 1: 133-221. 1878
555. _____ *Notae mycologicae*. Series XV. *Fungi ex Gallia, Germania, Italia, Melita (Malta), Mexico, India, Japonia*.
Ann. Mycol. (Sydow) 11: 14-21. 1913
556. _____ *Notae mycologicae*. Series XVII. VI. *Fungi melitensis*.
Ann. Mycol. (Sydow) 11: 560-565. 1913
557. _____ *Notae mycologicae*. Ser. XIX. *Ann. Mycol. (Sydow)* 13: 128. 1915
558. _____ *Sylloge Fungorum*: v. 3 (1884); v. 4 (1886);
v. 10 (1892); v. 11 (1895); v. 12 (1897); v. 13 (1898); v. 14 (1899); v. 16 (1902); v. 18 (1906); v. 22 (1913); v. 25 (1931).
Pavia, Italy. 1884-1931
559. _____, C. H. PECK, and WM. TRELEASE. The fungi of
Alaska. *Harriman Alaska Expedition, Cryptogamic Botany* 5:
13-64. illus. 1904
560. _____, and ALEX. TROTTER. *Fungi Tripolitani*. *Ann. Mycol. (Sydow)* 11: 417. 1913
561. SADASIVAN, T. S. Succession of fungi decomposing wheat straw in
different soils with special reference to *Fusarium culmorum*.
Ann. Appl. Biol. 26: 497-508. 1939
562. SALLANS, B. J. *Alternaria* studies. Rpt. Dom. Bot. for the year
1929. Canada Dept. Agr. Div. Bot. pp. 100-108. 1931
563. _____ A study of the root-rot problem of wheat and bar-
ley caused by *Helminthosporium sativum* in Saskatchewan. Rpt.
Dom. Bot. for the year 1930. Canada Dept. Agr. Div. Bot. pp.
82-84. 1931
564. _____ Methods of inoculation of wheat with *Helmintho-
sporium sativum* P. K. and B. *Sci. Agr.* 13: 515-527. 1933
565. _____ The use of water by wheat plants when inoculated
with *Helminthosporium sativum*. *Canad. Jour. Res., Sect. C.*,
18: 178-198. 1940
566. _____ The relationship of weeds to losses caused by
common rootrot in wheat. *Sci. Agr.* 20: 632-637. illus.
1940
567. _____ The importance of various roots to the wheat
plant. *Sci. Agr.* 23: 17-26. illus. 1942
568. SAMPSON, KATHLEEN. Diseases of grasses. Some observations on
Epichloë typhina (Fr.) Tul. *Gard. Chron. (London)* 85: 280-
281. 1929
569. _____ The systemic infection of grasses by *Epi-
chloë typhina* (Pers.) Tul. *Brit. Mycol. Soc. Trans.* 17: 30-
47. 1933. (Rev. Appl. Mycol. 13: 169. 1934)
570. _____, and J. H. Western. Notes on the supposed
connexion between *Mastigospirium album* Riess and *Dilophospora
alopecuri* (Fr.) Fr. *Brit. Mycol. Soc. Trans.* 22: 168-173.
illus. 1938.
571. _____, and _____ Two diseases of grasses
caused by species of *Helminthosporium* not previously recorded
in Britain. *Brit. Mycol. Soc. Trans.* 24: 255-263. illus.
1940

572. SAMPSON, KATHLEEN, and J. H. WESTERN. Diseases of British grasses and herbage legumes. i-vii + 85 pp., illus. 1941. (Cambridge, England)
573. SAMUEL, G. Whiteheads or take-all in wheat. Jour. Min. Agr. 44: 231-241. 1937
574. ———, and S. D. GARRETT. Rhizoctonia solani on cereals in South Australia. Phytopathology 22: 827-836. 1932
575. ———, and F. J. GREANEY. Some observations on the occurrence of Fusarium culmorum on wheat. Brit. Mycol. Soc. Trans. 21: 114-117. 1937
576. SANFORD, G. B. Colletotrichum graminicolum (Ces.) Wils. as a parasite of the stem and root tissues of Avena sativa. Sci. Agr. 15: 370-376. 1935
577. ———. Studies on Rhizoctonia solani Kuehn. III. Racial differences in pathogenicity. Canad. Jour. Res., Sect. C., 16: 53-64. 1938
578. ———. Research on certain soil-borne diseases as affected by other micro-organisms. Sci. Agr. 19: 609-615. 1939
579. ———, and W. C. BROADFOOT. Studies of the effects of other soil-inhabiting micro-organisms on the virulence of Ophiobolus graminis Sacc. Sci. Agr. 11: 512-528. 1931
580. ———, and ———. On the prevalence of pathogenic forms of Helminthosporium sativum and Fusarium culmorum in the soil of wheat fields and its relation to the root rot problem. Canad. Jour. Res. Sect. C, 10: 264-274. 1934
581. ———, and M. W. CORMACK. Variability in association effects of other soil fungi on the virulence of Helminthosporium sativum on wheat seedlings. Canad. Jour. Res. Sect. C, 18: 562-565. 1940
582. SAVULESCU, T., and T. RAYSS. Contribution a la connaissance de la biologie de Nigrospora oryzae (B. et Br.) Petch parasite du maïs. Recueil de Travaux cryptogamiques dédiés à Louis Mangin. Paris. Mus. d'Hist. Nat., p. 233-240. 1931. (Reprint 8pp.)
- 582a. ———, and ———. Influence des conditions extérieures sur le développement de Nigrospora oryzae (B. et Br.) Petch, parasite du maïs en Roumanie. (Paris) Acad. des Sci. Compt. Rend. 194: 1262-1265. 1932
583. SAWADA, K. Diseases of crops in Formosa, II. Diseases of Italian millet. Formosa Agr. Bul. 64, pp. 15-19. 1912. (In Japanese)
584. SAXBY, S. H. Eye-spot in wheat. New Zeal. Jour. Agr. 66: 257-261. illus. 1943
585. SCHLEFFNIT, E., and M. WIEBEN. Untersuchungen über den Erreger der Federbuschsporenkrankheit (Dilophospora alopecuri). Forsch. auf dem Geb. Pflanzenkrank. u. Immunität im Pflanzenr. 5: 5-38. 1928
586. SCHNELLEHARDT, OTTO, and F. D. HEALD. The spore load of market wheat. Amer. Micros. Soc. Trans. 55: 281-285. 1936

587. SCHWARZE, CARL A. The parasitic fungi of New Jersey. N. J. Agr. Expt. Sta. Bul. 313, 226 pp., illus. 1917
588. SCHWEINFURTH, G., and F. de von THUEMEN. Fungi Egyptiaci. Grevillea 6: 102-104. 1878
589. SELBY, A. D., and T. F. MANNS. Studies in diseases of cereals and grasses. Ohio Agr. Expt. Sta. Bul. 203, pp. 185-236. illus. 1909
590. SEMENIUK, G. Growth stimulation of Diplodia zeae. Phytopathology 30: 21-22. (Abs.). 1940
591. _____ Seedling infection of maize by Diplodia zeae in steamed and nonsteamed soil. Phytopathology 30: 22. (Abs.). 1940
592. _____ Development of Diplodia zeae and Gibberella saubinetii in maize pith following stalk inoculations. Phytopathology 31: 20. (Abs.). 1941
593. _____ Comparative reactions of single crosses of dent maize to Diplodia zeae. Phytopathology 32: 16. (Abs.) 1942
594. _____ Charcoal-rot (Sclerotium bataticola) of maize, new to Iowa. Iowa Acad. Sci. Proc. (1942) 49: 256. (Abs.). 1942
595. _____ Seedling infection of dent maize by Sclerotium bataticola Taub. Phytopathology 34: 838-843. 1944.
596. _____, C. S. REDDY, I. E. MELHUS, E. W. LINDSTROM, and G. F. Sprague. Pathology and mycology of corn. Iowa Agr. Expt. Sta. 1942-43 Rpt., Part II, p. 52-57. 1944. (Rev. Appl. Mycol. 23: 331-333. 1944)
597. _____, _____, W. E. LOOMIS, E. V. LINDSTROM, and G. F. SPRAGUE. Pathology and mycology of corn. Iowa Agr. Expt. Sta. 1940-41 Rpt., Part II, p. 54-58, illus. 1941
598. SENN, P. H. The effect of the sugary gene in corn on resistance to seedling blight caused by Gibberella saubinetii. Phytopathology 22: 675-697. 1932
599. SEYMOUR, A. B. Host index of the fungi of North America. Harvard Univ. Press, Cambridge, Mass. 732 pp. 1929
600. SHANDS, H. L. Temperature studies on stripe of barley. Phytopathology 24: 364-383. 1934
601. _____, and D. C. ARNY. Stripe reaction of spring barley varieties. Phytopathology 34: 572-585. 1944
602. SHAW, F. J. F., and S. L. AJREKAR. The genus Rhizoctonia in India. India Dept. Agr. Mem. Bot. Ser. 7: 177-194. 1915
603. SHEN, C. I. Soil conditions and the Fusarium culmorum seedling blight of wheat. Ann. Appl. Biol. 27: 323-329. 1940
604. SHERBAKOFF, C. D., and L. S. MAYER. Black ear rot of corn. Phytopathology 27: 207. 1937
605. SHITALKOVA-ROUSAKOVA, Mme. A. A. The susceptibility of rye varieties to the spot disease (Sclerotium graminis Fuckel) under conditions of the Otrada-Kuban selection farm. Lenin-grad Plant Protect, 1939, 18: 71-76, illus. 1939. (Rev. Appl. Mycol. 18: 587. 1939)

606. SIDERIS, C. P. Taxonomic studies in the family Pythiaceae II. Pythium. Mycologia 24: 14-61. 1932
607. SIMMONDS, P. M. Seedling blight and foot-rots of oats caused by Fusarium culmorum (W. G. Sm.) Sacc. Canada Dept. Agr. Bul. 105., n. s., 43 pp. 1928
608. _____ A review of the investigations conducted in western Canada on root rots of cereals. Sci. Agr. 19: 565-582. 1939
609. _____ Rootrots of cereals. Bot. Rev. 7: 308-332. 1941
610. _____, and R. J. LEDINGHAM. A study of fungous flora of wheat roots. Sci. Agr. 18: 49-59. 1937
611. _____, R. C. RUSSELL, and B. J. SALLANS. A comparison of different types of root rot of wheat by means of root excavation studies. Sci. Agr. 15: 680-700. 1935
612. _____, and B. J. SALLANS. Root amputations in the study of root-rots of cereals. Proc. Canad. Phytopath. Soc. 1: 33-43. 1930
613. _____, and _____ Further studies on amputations of wheat roots in relation to diseases of the root system. Sci. Agr. 13: 439-448. 1933
614. SLEETH, B. Agar medium and technique for isolating Pythium free of bacteria. Phytopathology 35: 1030-1031. 1945
615. SMITH, N. J. G. Observations of the Helminthosporium disease of cereals in Britain. Ann. Appl. Biol. 16: 236-260. 1929
616. SNYDER, W. C. and H. N. HANSEN. The species concept in Fusarium. Amer. Jour. Bot. 27: 64-67. 1940
617. SOLHEIM, W. G. Mycoflora Saximontanensis Exsiccata. Cent. I. Wyo. Univ. Pub. Bot. 1 (8): 219-232. 1934
618. SOLKINA, Mme. A. F. A study of the cycle of development of the fungus Sclerotinia graminearum Elen. Leningrad Plant Protect., 1939, 18: 100-108, illus. 1939. (English summary)
619. SPEGAZZINI, C. Fungi Fuegiana. Acad. Nac. de Cien. Córdoba. Bol. 11: 133-311. 1887
620. SPRAGUE, R. Cercospora herpotrichoides Fron, the cause of the Columbia Basin footrot of winter wheat. Science n. s., 74: 51-53. 1931
621. _____ The distribution of cereal footrots in the Pacific Northwest. Northwest Science 5: 10-12. 1931
622. _____ The relative importance of Cercospora herpotrichoides and of Leptosphaeria herpotrichoides as parasites of winter cereals. Phytopathology 24: 167-168. 1934
623. _____ Preliminary note on another foot rot of wheat and oats in Oregon. Phytopathology 24: 946-948. 1934
624. _____ The association of Cercospora herpotrichoides with the Festuca consociation. Phytopathology 24: 669-676. 1934
625. _____ Observations on diseases of Gramineae in Oregon and adjacent parts of Washington during the open winter of 1933-34. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 18: 24-25. 1934. (Processed)

626. SPRAGUE, R. A physiologic form of Septoria tritici on oats. Phytopathology 24: 133-143. 1934
627. _____ Notes on some diseases of rye grasses in western Oregon. U. S. Dept. Agr. Bur. Plant Indus. Plant Dis Rptr. 19: 19. 1935. (Processed)
628. _____ A new leaf spot on oats. Northwest Sci. 9: 15. 1935
629. _____ A preliminary check list of the parasitic fungi on cereals and other grasses in Oregon. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 19: 156-186. 1935. (Processed)
630. _____ Wojnowicia graminis as a very weak, secondary parasite of winter cereal crops. Phytopathology 25: 405-415. 1935
631. _____ Relative susceptibility of certain species of Gramineae to Cercospora herpotrichoides. Jour. Agr. Res. 53: 659-670. 1936
632. _____ Leaf reddening in winter oats in western Oregon. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 20: 114-115. 1936. (Processed)
633. _____ A further note on the fungus causing a white foot rot of wheat and oats. Phytopathology 27: 798-800. 1937
634. _____ Undescribed species of Cercospora and Cercospora on certain grasses in Oregon and Washington. Mycologia 29: 199-206. illus. 1937
635. _____ Influence of climatological factors in the development of Cercospora foot rot of winter wheat. U. S. Dept. Agr. Cir. 451, 39 pp., illus. 1937
636. _____ Fusarium poae on spring oats in Oregon. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 21: 87-88. (Processed)
637. _____ Additions to the check list of the parasitic fungi on cereals and other grasses in Oregon (1935-1937). U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 21: 412-422. 1937. (Processed)
638. _____ New or noteworthy parasitic species of fungi imperfecti in Oregon. Mycologia 29: 426-433. 1937
639. _____ Soil-borne cereal diseases in coastal Oregon. Northwest Science 12: 74-80. 1938. (Rev. Appl. Mycol. 19: 74)
640. _____ Two Mastigosprium leaf spots on Gramineae. Jour. Agr. Res. 57: 287-299. illus. 1938
641. _____ Notes on diseases of cereals and grasses in Oregon and adjacent counties in Washington during the spring of 1938. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 22: 174-175. 1938. (Processed)
642. _____ The status of Septoria graminum. Mycologia 30: 672-678. illus. 1938
643. _____ Cercospora foot rot of winter grains in the Pacific Northwest. U. S. Dept. Agr., Bur. Plant Indus. (Unnumbered Pub.), 13 pp. 1939. (Processed)

644. SPRAGUE, R. Leaf reddening in oats in Oregon. Oreg. Agr. Expt. Sta. Cir. Inform., no. 208, 3 pp., 1939. (Processed)
645. _____ Cereal diseases in Oregon and adjacent Washington in 1939. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 23: 220-221. 1939. (Processed)
646. _____ Foot rots and root rots of small grains in Oregon. Oreg. Agr. Expt. Sta. Cir. Inform., no. 207, 6 pp. 1939 (Supersedes Cir. Inform. no. 99) (Processed)
647. _____ Pink patch of fescue grasses in Oregon. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 24: 164. 1940. (Processed)
648. _____ A third species of Mastigosporium on Gramineae. Mycologia 32: 43-45. illus. 1940
649. _____ Septoria infuscans (E. and E.) Sprague Comb. Nov. In Cooke, Wm. Bridge. Mycobiota of North America No. 113 (Exsiccati). 1940
650. _____ Septoria jacucella [jaculella] Sprague sp. nov. In Cooke, Wm. Bridge. Mycobiota of North America No. 114 (Exsiccati). 1940
651. _____ A blotch and char-spot of western grasses. Northwest Science 15: 81-85. illus. 1941
652. _____ Some leaf spot fungi on western Gramineae. Mycologia 33: 655-665. 1941
653. _____ Stagonospora arenaria on grasses. Mycologia 33: 371-379. illus. 1941
654. _____ Cereal root-rot investigations and control factors. N. Dak. Agr. Expt. Sta. Bimo. Bul. 3: 19-22. 1941
655. _____ A revised check list of the parasitic fungi on cereals and other grasses in Oregon. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. Suppl. 134, 36 pp., 1942. (Processed)
656. _____ An annotated list of the parasitic fungi on cereals and other grasses in Klickitat County, Washington. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 26: 228-240. 1942. (Processed)
657. _____ Cercospora eyespot of Kentucky bluegrass. Phytopathology 32: 737-738. illus. 1942
658. _____ Root rots of Gramineae in the Northern Great Plains, 1940-43. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 27: 248-250. 1943. (Processed)
659. _____ The genus Phaeoseptoria on grasses in the Western Hemisphere. Mycologia 35: 483-491. 1943
660. _____ The status of Septoria alopecuri and some related species. Mycologia 35: 259-263. illus. 1943
661. _____ Can we increase the yield of cereals by reducing root rot injury? N. Dak. Agr. Expt. Sta. Bimo. Bul. 6: 20-25. 1943
662. _____ Phoma terrestris on Gramineae in the Northern Great Plains. Phytopathology 34: 129-131. 1944
663. _____ Root rots of cereals and grasses in North Dakota. N. Dak. Agr. Expt. Sta. Bul. 332, 32 pp., illus. 1944

664. SPRAGUE, R. Notes on diseases of cereals and grasses in Minnesota and North Dakota. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 28: 703-704. 1944. (Processed)
665. _____ Septoria diseases of Gramineae in the western United States. Oreg. State [Agr.] Col. Monog. Bot. 6, 151 pp., illus. 1944 [1946] (Issued January 1946)
666. _____ Additions to the Fungi Imperfecti on grasses in the United States. Mycologia 38: 52-64. illus. 1946
667. _____ Notes on the genus Hendersonia on western grasses. (In ms.) 1946
668. _____ Some leaf spot fungi on western Gramineae, II. (In ms.) 1946
669. _____ A Sporotrichum disease of Agropyron species. (In ms.) 1946
670. _____, and R. E. ATKINSON. Cross inoculations with Pythium arrhenomanes from cereals and grasses in the Northern Great Plains. Phytopathology 32: 17. (Abs.). 1942
671. _____, and H. FELLOWS. Cercospora foot rot of winter cereals. U. S. Dept. Agr. Tech. Bul. 428, 24 pp., illus. 1934
672. _____, and A. G. JOHNSON. A new Pseudodiscosia. Mycologia 28: 181-185. 1936
673. _____, and _____ Selenophoma on grasses. Mycologia 32: 415. 1940
674. _____, and _____ Ascochyta leaf spots of cereals and grasses in western America. (In ms.). 1945
675. _____, and _____ Selenophoma species on western grasses. (In ms.) 1945
676. _____, and _____ Selenophoma on grasses, II. Mycologia 37: 638-639. 1945
677. SPYSHNEVA, N. N. (SPESCHNEV, N. N.) Materialy dlia izucheniia mikologicheskoi flory Kavkaza. I, Gribnye parazity Goviiskogo uiezda. (Materials for the study of the mycological flora of the Caucasus. I, Fungus parasites of the Goriisk region.). Tiflis Bot. Sad Trudy (Tiflis Bot. Gard. Rpt.) 1: 65-78. 1895
678. _____ Materialy dlia izucheniia mikologicheskoi flory Kavkaza. II. Gribnye parazity Kakhetii (Materials for the study of the mycological flora of the Caucasus. II, Fungus parasites of Kakhetii.). Tiflis Bot. Sad. Trudy (Tiflis Bot. Gard. Rpt.) 2: 199-266. 1897
679. STARMAN, LOUISE J. A Helminthosporium disease of wheat and rye. Minn. Agr. Expt. Sta. Bul. 191, 23 pp. 1920
680. STANDEN, J. H. Chemical and physical characteristics of maize cobs in relation to the growth of Nigrospora oryzae. Phytopathology 34: 315-323. 1944
681. _____ Nigrospora oryzae (B. and Br.) Petch on maize. Phytopathology 35: 552-564. 1945
682. STEVENS, F. L. The fungi which cause plant disease. 754 p., illus. New York, 1913

683. STEVENS, F. L. Foot-rot disease of wheat -- Historical and Bibliographic. Bul. Ill. Nat. Hist. Surv. 13: 259-286. 1919
684. _____ The Helminthosporium foot-rot of wheat, with observations on the morphology of Helminthosporium and on the occurrence of saltation in the genus. Ill. Nat. Hist. Surv. Bul. 14: 77-185. illus. 1922
685. STEVENSON, J. A., and R. D. RANDS. An annotated list of the fungi and bacteria associated with sugarcane and its products. Hawaiian Planters' Rec. 42: 247-314. 1938
686. STEWART, F. C., and H. E. HODCKINS. The Sporotrichum bud-rot of carnations and the silver top of June grass. New York Agr. Expt. Sta. Tech. Bul. 7: 83-119. 1908
687. STIRRUP, H. H. Sclerotium rhizodes Auersw. in England. Brit. Mycol. Soc. Trans. 16: 308. 1922
688. STOUT, A. B. A Sclerotium disease of blue joint and other grasses. Wis. Agr. Expt. Sta. Res. Bul. 18, pp. 207-253, illus. 1911
689. STOVER, W. G. The relation of soil temperature to the development of seedling blight of corn caused by Helminthosporium spp. Phytopathology 12: 30. (Abs.). 1922
690. STRUBLE, F. BEN. Plot technique for disease-control studies on fine turf. Phytopathology 33: 528-530. 1943
691. STUMBO, C. R., P. L. GAINES, and F. E. CLARK. Microbiological and nutritional factors in the take-all disease of wheat. Jour. Agr. Res. 64: 653-665. 1942
692. SUBRAMANIAM, L. S. Root rot and sclerotial disease of wheat. Pusa Imp. Inst. Agr. Res. Bul. 177, 7 pp., illus. 1928
693. SUNESON, COIT A., and SYLVIA C. SANTONI. Barley varieties resistant to stripe, Helminthosporium gramineum Rabh. Am. Soc. Agron. Jour. 35: 736-737. 1943
694. SYDOW, H. Novae fungorum species XXV. Ann. Mycol. (Sydow) 35: 244-286. 1937
695. _____ Mycotheca Germanica Fasc. LXV-LXVIII (No. 3201-3400). Ann. Mycol. (Sydow) 38: 453-484. 1940
696. _____, and E. J. BUTLER. Fungi Indiae Orientalis, Pars V. Ann. Mycol. (Sydow) 14: 177-220. 1916
697. _____, and P. SYDOW. Beiträge zur Kenntniss der Pilzflora der Mark Brandenburg. III. Hedwigia, Beibl. 39: (1)-(6). 1900
698. TANAKA, T. New Japanese Fungi. Notes and translations -- VIII. Mycologia 12: 25-32. 1920
699. TAUBENHAUS, J. J. The black rots of the sweet potato. Phytopathology 3: 159-166. 1913
700. TEHON, L. R., and G. H. BOEWE. Charcoal rot in Illinois. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 23: 312-325. illus. 1939. (Processed)
701. _____, and E. Y. DANIELS. Notes on the parasitic fungi of Illinois -- III. Mycologia 19: 110-129. 1927
702. _____, and G. L. STOUT. Notes on the parasitic fungi of Illinois -- IV. Mycologia 21: 180-196. 1929
703. TERVET, I. W. Turf injuries in Minnesota. Minn. Agr. Expt. Sta. Rpt. (Abs.)., p. 46. 1941. (Expt. Sta. Rec. 86: 792-793. 1942)

704. TERVET, I. W. Control of dollar spot in Minnesota. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 26: 495-496. 1942. (Processed)
705. THOM, CHARLES. The Penicillia. 644 pp. (Baltimore, Md.) 1930
706. THOMAS, K. M. Detailed administration report of the government mycologist, Madras, for the year 1930-39. 30 pp., illus. 1939
707. TRABUT, L. Étude sur l'halfa (Stipa tenacissima). Algeria, 90 pp., illus. 1889 (Paris)
708. TRAIL, J. W. H. New Scotch microfungi. Scot. Nat. 9: 88. 1887
709. _____ Revision of Scotch Sphaeropsidaceae and Melanconieae. Scot. Nat. 9: 231. 1887
710. TRELEASE, WM. Preliminary list of Wisconsin parasitic fungi. Wis. Acad. Trans. 6: 106-144. 1885. (Reprint, 40 pp.)
711. _____ A spot disease of orchard grass, Scolecotrichum graminis Fckl. U. S. Commissioner Agr. Rpt. 1886: 129-131. 1887
712. TROPOVA, A. T. Vliianie vozvrata vesennikh kholodov na porazhenie iarovoi pshenitsy grihom Helminthosporium sativum P. K. et B. (Influence of a return of spring cold weather on injuries of spring wheat by Helminthosporium sativum P. K. et B.) Dokl. Vses. Akad. S.-Kh. Nauk V.I. Lenina (Lenin Acad. Agr. Sci. U.S.S.R.) 1940 (10): 24-28. 1940. (In Russian).
713. TROTTER, ALEX. In Saccardo, P. A., Sylloge Fungorum 25: 345. 1931.
714. TU, CHIH. Physiologic specialization in Fusarium spp. causing headblight of small grains. Phytopathology 19: 143-154. illus. 1929
715. _____ Physiologic specialization in Fusarium spp. causing headblight of small grains. Minn. Agr. Expt. Sta. Tech. Bul. 74, 27 pp., illus. 1930
716. TUCKER, C. M. The taxonomy of the genus Phytophthora. Mo. Agr. Expt. Sta. Res. Bul. 153, 208 pp. 1931
717. _____ The distribution of the genus Phytophthora. Mo. Agr. Expt. Sta. Res. Bul. 184, 80 pp. 1933
718. TULLIS, E. C. Fungi isolated from discolored rice kernels. U. S. Dept. Agr. Tech. Bul. 540, 11 pp., illus. 1936
719. _____ Diseases of rice. U. S. Dept. Agr. Farmers' Bul. 1854, 17 pp. 1940
720. TURNER, ELIZABETH M. Ophiobolus graminis Sacc. var. avenae var. n., as the cause of take all or whiteheads of oats in Wales. Brit. Mycol. Soc. Trans. 24: 269-281. 1940
721. _____ The reaction of oats to different strains of Ophiobolus graminis. Brit. Mycol. Soc. Trans. 24: 267. (Abs.) 1940
722. TYNER, L. E. The effect of crop debris on the pathogenicity of cereal root-rotting fungi. Canad. Jour. Res. Sect. C, 18: 289-306. illus. 1940
723. _____ Some factors affecting the virulence of artificial inoculum of Helminthosporium sativum P. K. and B. and of Fusarium culmorum (W. G. Sm.) Sacc. Canad. Jour. Res. Sect. C., 19: 42-48. 1941

724. TYNER, L. E., and W. C. BROADFOOT. Field tests of the differential reaction of wheat varieties to root rot. *Sci. Agr.* 24: 153-163. 1943
725. TYSON, J. Management of bent grass lawns. *Mich. Agr. Expt. Sta. Cir. Bul.* 156, 18 pp., illus. 1936
726. ULLSTRUP, A. J. The occurrence of Gibberella fujikuroi var. subglutinans in the United States. *Phytopathology* 26: 685-693. illus. 1936
727. _____ Two physiologic races of Helminthosporium maydis in the corn belt. *Phytopathology* 31: 508-521. illus. 1941
728. _____ Diseases of dent corn in the United States. U. S. Dept. Agr. *Cir.* 674, 34 pp., illus. 1943
729. _____ Further studies on a species of Helminthosporium parasitizing corn. *Phytopathology* 34: 214-222. illus. 1944
730. UNAMUNO, LUIS M. Hongos microscopicos de los alrededores de la vid (Burgos). *Soc. Españ. de Hist. Nat. Bol.* 29: 387-402. 1929
731. _____ Hongos microscopicos de los alrededores de caudete (Albacete). *Soc. Españ. de Hist. Nat. Bol.* 30: 379-390. 1930
732. UNDERWOOD, L. M., and F. S. EARLE. A preliminary list of Alabama fungi. *Ala. Agr. Expt. Sta. Bul.* 80, p. 111-283. 1897
733. UPPAL, B. N., K. G. KOLHATKAR, and M. K. PATEL. Blight and hollow stem of sorghum. *Indian Jour. Agr. Sci.* 6: 1323-1334. illus. 1936
734. VALLEAU, W. D. Seed transmission of Helminthosporium of corn. *Phytopathology* 25: 1109-1112. 1935
735. _____, P. E. KARRAKER, and E. M. JOHNSON. Corn root rot -- a soil-borne disease. *Jour. Agr. Res.* 33: 453-476. illus. 1926.
736. VAN LUIJK, A. Untersuchungen über Krankheiten der Gräser. *Meded. Phytopath. Lab. Willie Commelin Scholten, Baarn (Holland)*, 13: 1-22. illus. 1934
737. VANTERPOOL, T. C. Toxin formation by species of Pythium parasitic on wheat. Reprinted from *World's Grain Exhibition and Conference Proc.* v. 2, pp. 294-298, illus. 1933
738. _____ Studies on browning root rot of cereals: III, Phosphorus-nitrogen relations of infested fields; IV, Effects of fertilizer amendments; V, Preliminary plant analyses. *Canad. Jour. Res. Sect. C*, 13: 221-250. illus. 1935
739. _____ Some species of Pythium parasitic on wheat in Canada and England. *Ann. Appl. Biol.* 25: 528-543. illus. 1938
740. _____ Root-rot diseases of cereals. *Saskatchewan Univ., Col. Agr. Ext. Bul.* 99, (4 pp.). 1939
741. _____ Present knowledge of browning root rot of wheat with special reference to its control. *Sci. Agr.* 20: 735-749. 1940

742. VANTERPOOL, T. C. Studies on browning root rot of cereals: VI, further contributions on the effects of various soil amendments on the incidence of the disease in wheat. *Canad. Jour. Res. Sect. C*, 18: 240-257. illus. 1940
743. _____ Pythium root rot of grasses. *Sci. Agr.* 22: 674-687. 1942
744. _____, and R. SPRAGUE. Pythium arrhenomanes on cereals and grasses in the Northern Great Plains. *Phytopathology* 32: 327-328. 1942
745. _____, and J. H. L. TRUSCOTT. Studies on browning root rot of cereals. II. Some parasitic species of Pythium and their relation to the disease. *Canad. Jour. Res.* 6: 68-93. illus. 1932
746. VOLK, A. Untersuchungen über Typhula graminum Karst. *Ztschr. f. Pflanzenkr.* 47: 338-365. 1937
747. VOORHEES, R. K. Sclerotial rot of corn caused by Rhizoctonia zeae n. sp. *Phytopathology* 24: 1290-1303. 1934
748. VOSBEIN, O. Können wir uns gegen die Fusskrankheit schützen? (Can we guard against foot rot?). *Mitt. Landw. Berl.* 1, 39, p. 827. 1935
749. WAGNER, F. A. Reaction of sorghums to the root, crown, and shoot rot of milo. *Amer. Soc. Agron. Jour.* 28: 643-654. illus. 1936
750. WALKER, A. G. The colonization of buried wheat straw by soil fungi, with special reference to Fusarium culmorum. *Ann. Appl. Biol.* 28: 333-350. 1941
751. _____ Rhizomatous grass weeds and Ophiobolus graminis (Sacc.). *Ann. Appl. Biol.* 32: 177-178. 1945
752. WEBER, G. F. Septoria diseases of cereals. I. Speckled blotch of oats caused by Leptosphaeria. *Phytopathology* 12: 449-470. 1922
753. _____ Septoria diseases of cereals. II. Septoria diseases of wheat. *Phytopathology* 12: 537-585. 1922
754. _____ III. Septoria diseases of rye, barley, and certain grasses. *Phytopathology* 13: 1-23. 1923
755. WEISS, F. Check list revision. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 27: 118-126; 175-184; 188-193. 1943
756. _____ Check list revision. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 28: 1138-1147; 1162-1164. 1944
757. _____ Check list revision. U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. 29: 34-39; 68-76; 132-139; 186-194; 222-228; 274-282; 328-337; 374-377; 416-422; 468-474; 506-513; 542-548; 576-583; 624-631; 660-668. 1945
758. WELCH, AARON. Pythium injury of oats. *Phytopathology* 30: 27. 1940
759. _____ Further studies on Pythium injury of oats. *Phytopathology* 32: 19. 1942
760. WENIGER, WANDA (Revised by W. E. Brentzel). Diseases of grain and forage crops in North Dakota. *N. Dak. Agr. Expt. Sta. Bul.* 255, 97 pp., illus. 1932 (Revision of Bul. 166, May, 1923).

761. WERNHAM, C. C. New facts about eastern snowmold. *Phytopathology* 31: 940-943. 1941
762. _____, and R. S. KIRBY. A new turf disease. *Phytopathology* 31: 24. 1941
763. WESTENDORP, G. D. Cinquième notice sur quelques hypoxylees inédites, ou nouvelles pour la flore de la Belgique. *Bul. l'Acad. Roy. des Sci., des Lettres et des Beaux-Arts de Belgique*, Ser. 2, 2: 554-579. illus. 1857
764. WHITE, N. H. Physiological studies of the fungus Ophiobolus graminis Sacc. I, Growth factor requirements. *Jour. Council Sci. and Indus. Res. (Austral.)* 14: 137-146. 1941
765. _____ The genetics of Ophiobolus graminis Sacc. I. Heritable variations for culture color and pathogenicity. *Jour. Council Sci. and Indus. Res. (Austral.)* 15: 118-124. illus. 1942
766. WILSON, G. W. The identity of the anthracnose of grasses in the United States. *Phytopathology* 4: 106-112. 1914
767. WILSON, WILLIAM E. Physiological studies on two species of Diplodia parasitic on corn. *Phytopathology* 32: 130-140. 1942
768. WILTSHIRE, S. P. The original and modern conceptions of Stemphylium. *Brit. Mycol. Soc. Trans.* 21: 211-239. illus. 1938
769. WINTER, G. Der Einfluss der physikalischen Bodenstruktur auf den Infektionsverlauf bei der Ophiobolose des Weizens. *Ztschr. Pflanzenkrank.* 49: 513-559. illus. 1939
770. _____ Untersuchungen über den Einfluss biotischer Faktoren auf die Infektion des Weizens durch Ophiobolus graminis. *Ztschr. Pflanzenkrank.* 50: 113-134. 1940
771. _____ Die Infektion des Weizens durch Ophiobolus graminis als Funktion der Temperatur. *Ztschr. Pflanzenkrank.* 50: 444-459. illus. 1940
772. _____ Weitere untersuchungen über den Einfluss der Bodenstruktur auf die Infektion des Weizens durch Ophiobolus graminis. *Zentbl. f. Bakt. Abt. 2*, 101: 364-388. 1940
773. _____ Ein neuer Fusskrankheitserreger an Weizen, Gerste, Roggen und Hafer (Colletotrichum graminicolum (Ges.) Wils.). *Phytopath. Ztschr.* 13: 282-292. 1940
774. _____ Der Einfluss partieller Sterilisation des Bodens auf die Entwicklung der Laufhyphen von Ophiobolus graminis. *Phytopath. Ztschr.* 14: 204-302. illus. 1942. (*Rev. Appl. Mycol.* 24: 94-95. 1945)
775. WOLLENWEBER, H. W. Fusarium autographice delineata. 1917. (See *Ann. Mycol. (Sydow)* 15: 1-56)
776. _____, and O. A. REINKING. Die Fusarien, ihre Beschreibung, Schadwirkung, und Bekämpfung. P. Parey, Berlin, 355 pp., illus. 1935
777. YOUNG, P. A. Sclerotium blight of wheat. *Phytopathology* 27: 1113-1118. 1937
778. _____, and H. E. MORRIS. Sclerotium root rot, (Typhula graminum Karst.). U. S. Dept. Agr., Bur. Plant Indus., Plant Dis. Rptr. Suppl. 69: 127. 1929. (Processed)

779. ZHAVORONKOV, I. P. Niekotoryia nabliudeniiia na Helminthosporium turcicum Pass. Mater. po Mikol. i Fitopat. (Mater. Mycol. and Phytopath.) 1: 42-50. illus. 1915
780. ZILING, M. K. Black germ of wheat. Siberian Sci. Inst. for Cer. Ind. Omsk. p. 15-39. 1932
781. ZIMMERMANN, H. Verzeichnis der Pilze aus der Umgebung von Eisgrub, II. In Verh. Naturforsch. Verein. Brünn (1913) 52, pp. 66-128, illus. 1914
782. ZUNDEL, GEORGE L. Studies on the Ustilaginales of the World. II. Mycologia 34: 123-127. 1942

HOST INDEX

- Aegilops, 176
 Agropyron, 121, 134, 135, 144,
 150, 154, 165, 170, 196, 211,
 212
 --- albicans Scribn. and Sm.,
 136
 --- bakeri E. Nels., 123
 --- caninum (L.) Beauv., 123
 --- cristatum (L.) Gaertn., 106,
 107, 108, 110, 119, 120, 123,
 176, 179, 184, 198, 201, 206
 --- dasystachyum (Hook.) Scribn.,
 136, 210
 --- desertorum (Fisch.) Schult.,
 168
 --- elongatum (Host.) Beauv., 120
 --- inerme (Scribn. and Sm.)
 Rydb., 112, 126, 139, 150, 176,
 213
 --- michnoi Roshev, 123
 --- repens (L.) Beauv., 105, 110,
 124, 136, 150, 154, 167, 168,
 193, 201, 210, 211
 --- riparium Scribn. and Sm.,
 136, 176
 --- semicostatum (Steud.) Nees,
 123, 211
 --- sibiricum (Willd.) Beauv.,
 105, 120, 179
 --- smithii Rydb., 112, 113, 123,
 126, 131, 136, 150
 --- spicatum (Pursh) Scribn. and
 Sm., 112, 126, 136, 150, 171,
 213
 --- subsecundum (Lk.) Hitchc.,
 123, 136, 150, 184, 211, 213
 --- trachycaulum (Lk.) Malte, 107,
 112, 120, 123, 126, 136, 150,
 154, 168, 171, 208, 211
 --- trichophorum (Lk.) Richt.,
 123
 Agrostis, 137, 138, 141, 148, 149,
 170, 174, 184, 192, 206, 208,
 212
 --- alba L., 106, 118, 160, 168,
 171, 186, 191, 198, 200, 201,
 206, 208, 210, 211, 212
 --- canina L., 106, 186
 (Agrostis) castellana Boiss. and
 Reut., 160
 --- diegoensis Vasey, 138, 148
 --- exarata Trin., 106, 143, 148,
 160, 200
 --- hallii Vasey, 106, 158, 171
 --- palustris Huds., 106, 118,
 130, 143, 148, 149, 160, 179,
 185, 186, 200, 206, 208
 --- perennans (Walt.) Tuckerm.,
 191
 --- scabra Willd., 137, 138, 158,
 171
 --- tenuis Sibth., 106, 118, 149,
 160, 185, 186, 198, 208
 Aira caryophyllea L., 156
 Alopecurus, 212
 --- aequalis Sobol., 166
 --- pratensis L., 166, 210
 Andropogon, 121, 141
 --- furcatus Muhl., 125, 145, 168
 Anthoxanthum, 168
 --- odoratum L., 106, 132, 168,
 191
 Aristida, 137
 --- longiseta Steud., 138
 --- oligantha Michx., 138
 Arrhenatherum, 168, 170, 171, 190
 --- elatium (L.) Beauv., 107, 136,
 168, 170, 171, 208, 212
 Arundo, 134, 137
 --- donax L., 133, 134
 Avena, 121, 141
 --- barbata Brot., 190
 --- byzantina C. Koch, 126, 162,
 169, 190
 --- fatua L., 105, 111, 126, 162,
 168, 190
 --- hookeri Scribn., 168
 --- sativa L. (see also oats), 111,
 162, 168, 169, 190, 203
 Barley (see also Hordeum vulgare),
 107, 109, 114, 117, 118, 122,
 156, 176, 177, 184, 186, 187,
 188, 190, 192, 195, 196, 199,
 200, 207, 210, 211

- Beckmannia syzigachne* (Steud.)
 Fern., 203, 213
Bouteloua curtipendula (Michx.)
 Torr., 137, 138, 188, 194
 --- *gracilis* (H.B.K.) Lag., 106,
 110, 124, 179, 184, 186, 187,
 188, 194, 198, 201
Brachypodium, 161
Bromus, 122, 146, 165, 190, 196,
 208, 212
 --- *carinatus* Hook. and Arn.,
 106, 125, 131, 151, 165, 176,
 186, 190, 204, 208, 211
 --- *ciliatus* L., 155, 151, 171
 --- *commutatus* Schrad., 147, 186
 --- *inermis* Leyss., 107, 109,
 125, 133, 147, 176, 187, 188,
 190, 211
 --- *japonicus* Thurb., 147, 190
 --- *laevipes* Shear, 151
 --- *mollis* L., 119, 147
 --- *orcuttianus* Vasey, 114
 --- *purgans* L., 168
 --- *racemosus* L., 147
 --- *rigidus* Roth, 151, 173, 186,
 190
 --- *secalinus* L., 114, 147
 --- *tectorum* L., 109, 119, 151,
 176, 186, 190
 --- *vulgaris* (Hook.) Shear, 165,
 173, 205
Buchloë dactyloides (Nutt.)
 Engelm., 175, 194

Calamagrostis, 138, 170, 206
 --- *canadensis* (Michx.) Beauv.,
 137, 138, 206, 210
 --- *inexpansa* A. Gray, 111, 168,
 171
 --- *montanensis* Scribn., 138
 --- *nutkaensis* (Presl) Steud.,
 126, 146
 --- *rubescens* Buckl., 168
 --- *scribneri* Beal, 138
Calamovilfa, 136
 --- *longifolia* (Hook.) Scribn.,
 136, 149, 166
Cenchrus pauciflorus Benth., 149
Cinna latifolia (Trevir.) Griseb.,
 165

 Corn, see *Zea mays*
Cynodon dactylon (L.) Pers., 127,
 191, 192, 201

Dactylis glomerata L., 106, 107,
 132, 135, 136, 167, 168, 183,
 201, 206, 208, 210, 212
Danthonia, 135
 --- *californica* Bol., 135, 136,
 191
 --- *intermedia* Vasey, 129
 --- *spicata* (L.) Beauv., 191
Deschampsia atropurpurea (Wahl.)
 Scheele, 138
 --- *caespitosa* (L.) Beauv., 138,
 156, 178
 --- *danthonioides* (Trin.) Munro,
 138, 156
 --- *elongata* (Hook.) Munro, 138
Digitaria sanguinalis (L.) Scop.,
 132, 168
Distichlis, 121, 170
 --- *spicata* (L.) Greene, 193
 --- *stricta* (Torr.) Rydb., 171
Dupontia fischeri R. Br., 146

Echinochloa crusgalli (L.) Beauv.,
 106, 107, 111, 174, 179, 184,
 194, 195, 198
Eleusine indica (L.) Gaertn., 191
Elymus, 121, 124, 134, 135, 144,
 150, 153, 165, 170, 196, 211,
 212
 --- *canadensis* L., 112, 115, 120,
 123, 124, 136, 145, 150, 211
 --- --- *var. robustus* (Scribn.
 and Sm.) Mackenz. and Bush.,
 140, 145, 150
 --- *condensatus* Presl, 134, 139,
 150, 171, 204, 211
 --- --- *var. pubens* Piper, 155
 --- *dahuricus* Turcz., 112
 --- *excelsus* Turcz., 154
 --- *flavescens* Scribn. and Sm.,
 134
 --- *giganteus* Vahl., 136
 --- *glaucus* Buckl., 106, 136, 139,
 150, 155, 171, 210, 211
 --- *hirsutus* Presl, 201
 --- *junceus* Fisch., 136, 188, 211

- (*Elymus*) *mollis* Trin., 155
 --- *sibiricus* L., 154
 --- *triticoideus* Buckl., 130, 150, 171, 211
 --- *virginicus* L., 112, 123, 136
Eragrostis cilianensis (All.) Lk., 193
- Festuca*, 116, 121, 159, 191, 208
 --- *dertonensis* (All.) Aschers. and Graebn., 114, 159, 160
 --- *elatior* L., 105, 191
 --- *idahoensis* Elmer, 113, 135, 136, 170
 --- *kingii* (S. Wats.) Cassidy, 138
 --- *megalura* Nutt., 114, 208
 --- *myuros* L., 114, 116, 208
 --- *occidentalis* Hook., 191
 --- *octoflora* Walt., 160, 168
 --- *ovina* L., 136, 138, 160
 --- --- var. *brachyphylla* (Schult.) Piper, 138
 --- *rubra* L., 105, 138, 178, 186, 188
 --- --- var. *commutata* Gaud., 106, 160, 186
 --- *subulata* Trin., 160
Fluminea festucacea (Willd.) Hitchc., 165
- Gastridium ventricosum* (Cowan) Schinz and Thell, 203
Glyceria elata (Nash) Hitchc., 154, 208
 --- *pauciflora* Presl, 154
 --- *striata* (Lam.) Hitchc., 154
- Hierochloë alpina* (Sw.) Roem. and Schult., 155
 --- *odorata* (L.) Beauv. 124
Holcus, 122, 144, 208
 --- *lanatus* L., 114, 118, 121, 122, 126, 162, 177, 200, 208
Hordeum, 122, 156, 176, 196
 --- *brevisubulatum* Lk., 136
 --- *distichon* L., 156
 --- *jubatum* L., 105, 156, 188, 211, 213
 --- *murinum* L., 186, 211
- (*Hordeum*) *nodosum* L., 125, 156, 184, 186
 --- *vulgare* L. (see also barley), 121, 136, 156, 168, 186
Hystrix patula (Scribn. and Sm.) Rydb., 112, 154, 156
- Koeleria*, 149
 --- *cristata* (L.) Pers., 112, 135, 136, 143, 149, 157, 176
- Lolium*, 141, 144, 208
 --- *marschalli* Stev., 208
 --- *multiflorum* Lam., 105, 124, 151, 163, 168, 198, 210, 211
 --- *perenne* L., 124, 125, 163, 168, 198, 208, 210, 211
- Melica*, 141
 --- *bulbosa* Geyer, 127, 178
 --- *harfordii* Boland, 136
 --- *scabrosa* Trin., 153
 --- *subulata* (Griseb.) Scribn., 178
- Muhlenbergia*, 115, 146
 --- *asperifolia* (Nees and Mey.) Parodi, 153, 171, 203
 --- *cuspidata* (Torr.) Rydb., 138
 --- *japonica* Steud., 193
 --- *mexicana* (L.) Trin., 153, 191
 --- *racemosa* (Michx.) B.S.P., 138, 187
- Munroa squarrosa* (Nutt.) Torr., 153
- Oats* (see also *Avena*), 105, 107, 108, 109, 110, 111, 114, 146, 162, 169, 176, 183, 186, 187, 189, 196, 200
- Oryza sativa* L., 179, 180, 209
Oryzopsis hymenoides (Roem. and Schult.) Ricker, 136, 165
 --- *micrantha* (Trin. and Rupr.) Thurb., 168
- Panicum*, 123, 134, 203
 --- *capillare* L., 187, 188
 --- *dichotomiflorum* Michx., 123, 174
 --- *dichotomum* L., 174

- (*Panicum*) *miliaceum* L., 106, 108,
 109, 110, 179, 186, 187, 188,
 193, 194, 198, 201, 202
 --- *pacificum* Hitchc. and Chase,
 174
 --- *virgatum* L., 121, 123, 134,
 174
Phalaris, 121, 134, 165
 --- *arundinacea* L., 125, 133, 134,
 147, 166, 204, 207, 211
 --- *californica* Hook. and Arn.,
 166
Phleum phleoides (L.) Karst., 205
 --- *pratense* L., 112, 122, 123,
 136, 167, 168, 187, 204, 212
Phragmites, 134
 --- *communis* Trin., 133, 134, 166,
 167, 206, 213
Pleuropogon, 121, 125
 --- *refractus* (A. Gray) Benth.,
 125
Poa, 116, 122, 134, 135, 196, 212
 --- *alpina* L., 136
 --- *ampla* Merr., 136, 152, 213
 --- *annua* L., 106, 116, 151, 179,
 186
 --- *arctica* R. Br., 136
 --- *arida* Vasey, 139, 152, 202
 --- *canbyi* (Scribn.) Piper, 114,
 152, 154
 --- *compressa* L., 112, 116, 136,
 152, 154, 200, 202
 --- *cusickii* Vasey, 152
 --- *epilis* Scribn., 137
 --- *flexuosa* var. *elongata* Blytt,
 155
 --- *howellii* Vasey and Scribn.,
 143, 152
 --- *interior* Rydb., 137
 --- *juncifolia* Scribn., 106, 116,
 137, 154
 --- *kelloggii* Vasey, 151
 --- *nervosa* (Hook.) Vasey, 137,
 152
 --- *nevadensis* Vasey, 152, 154
 --- *palustris* L., 154, 187
 --- *pratensis* L., 106, 112, 116,
 135, 137, 152, 154, 155, 167,
 168, 169, 176, 177, 184, 186,
 187, 200, 202
 (*Poa*) *scabrella* (Thurb.) Benth.,
 152
 --- *secunda* Presl., 106, 112, 127,
 137, 152, 154, 176, 195
 --- *stenantha* Trin., 112
 --- *trivialis* L., 195, 200
 --- *vaseyochloa* Scribn., 105, 116,
 152
Puccinellia nuttalliana (Schult.)
 Hitchc., 137
 Rice, see *Oryza sativa*
 Rye, see *Secale cereale*
Schedonnardus paniculatus (Nutt.)
 Trel., 193
Schizachne purpurascens (Torr.)
 Swallen, 153, 168
Secale cereale L., 112, 135, 136,
 137, 140, 157, 168, 176, 186,
 211, 212, 213
Setaria, 175, 185, 198, 201, 202,
 203, 209
 --- *italica* (L.) Beauv., 193, 197,
 201
 --- *lutescens* (Weigel) F.T.Hubb,
 175
 --- *viridis* (L.) Beauv., 105, 132,
 193, 194, 197
Sitanion hanseni (Scribn.) J. G.
 Sm., 139
 --- *hystrix* (Nutt.) J. G. Sm.,
 137, 139, 156, 177
 --- *jubatatum* J. G. Sm., 126
Sorghastrum, 121
 --- *nutans* (L.) Nash, 145, 166
Sorghum, 107, 108, 125, 130, 174,
 176, 180, 185, 202, 203, 208
 --- *halepense* (L.) Pers., 176
 --- *vulgare* L., 198
 --- --- var. *sudanense* (Piper)
 Hitchc., 168, 179, 202
Spartina alternifolia Loisel., 172
 --- --- var. *glabra* (Muhl.) Fern.,
 172
 --- *gracilis* Trin., 158
 --- *leiantha* Benth., 172
 --- *pectinata* Lk., 158
 --- *stricta*, 172
Sphenopholis, 141, 146

(*Sphenopholis*) *obtusata* (Michx.)
 Scribn., 138, 145, 157, 168,
 174
Sporobolus, 134, 141, 146
 --- *airoides* (Torr.) Torr., 121,
 125
 --- *asper* (Michx.) Kunth, 136,
 137
 --- *cryptandrus* (Torr.) A. Gray,
 110, 137
 --- *heterolepis* A. Gray, 146, 168
 --- *neglectus* Nash, 203
Stipa, 121, 125, 134, 139, 141,
 146, 158, 159
 --- *columbiana* Macoun 134
 --- --- *var. nelsoni* (Scribn.)
 Hitch., 159
 --- *comata* Trin. and Rupr., 125,
 134, 142, 146, 158
 --- *richardsoni* Lk., 134, 139
 --- *spartea* Trin., 134
 --- *viridula* Trin., 105, 113, 134,
 146, 154, 158, 165
 --- *williamsii* Scribn., 158
Trisetum, 143, 148, 206

(*Trisetum*) *canescens* Buckl., 148
 --- *cernuum* Trin., 148, 168
 --- *spicatum* (L.) Richt., 138
Triticum, 153, 161, 176, 196
 --- *aestivum* L., 154, 161, 168,
 176, 198
 --- *dicoccum* Schrank, 154
 --- --- *var. farrum* Beyle, 161
 --- *durum* Desf., 105, 172, 188
 --- *spelta* L., 154, 161
 --- *turgidum* L., 161

Wheat, 106, 107, 108, 109, 110,
 112, 114, 117, 118, 119, 127,
 144, 153, 154, 161, 162, 163,
 167, 168, 172, 176, 177, 179,
 183, 184, 186, 187, 188, 194,
 195, 196, 199, 200, 201, 204,
 208, 212

Zea mays L. (corn), 105, 106,
 109, 110, 127, 130, 173, 179,
 185, 187, 201, 202, 203, 207,
 208, 209

FUNGUS INDEX

- Alternaria*, 172, 200
 --- *circinans* (Berk. and Curt.)
 Bolle, 172
 --- *peglionii* Curzi, 172
 --- *tenuis* Nees, 172, 200
Ansatospora, 173
 --- *bromi* (Sprague) Sprague, 173
Aphanomyces, 105, 189
 --- *camptostylus* Drechs., 105
Ascochyta, 120, 121, 122, 123,
 127, 155, 158, 160
 --- *agropyrina* (Fairm.) Trott.,
 121, 123
 --- *boutelouae* Fairm., 121, 124
 --- *cynodontis* Unam., 127
 --- *desmazieri* Cav., 122, 124,
 163
 --- *elymi* Tehon and Daniels, 122,
 124
 --- *graminicola* Sacc. (and var-
 ieties), 121, 122, 124, 125
 --- *sorgi* Sacc., 122, 125
 --- *stipae* Died., 121, 125
Ascochyta, 120, 126
 --- *avenae* Petr., 121, 126
Ascochyta, 120, 121, 123, 126
 --- *agropyrina* Fairm., 121, 123
Ascomycetes, 102, 103, 111
Ascospora, 138
Basidiomycetes, 102, 103, 116
 ---, Non-sporulating, 119
Basisporium *gallarum* Moll., 207
Brachycladium *spiciferum* Bainier,
 199, 200
Brachysporium, 199
 --- *flexuosum* (Cda.) Sacc., 195

Calonectria *graminicola* (Berk.
 and Br.) Wr., 185
Cephalosporium *acremonium* Cda.,
 173
Cercospora 153, 173, 174, 175
 --- *agrostidis* Atk., 174
 --- *bromi* Sprague, 173
 --- *echinoclaoe* J. J. Davis, 174
 --- *fusimaculans* Atk., 174, 176
 --- *graminicola* Tracy and Earle,
 212
 (*Cercospora*) *graminis* (Fckl.)
 Horsfall, 212
 --- *poae* Baudyš and Picb., 212
 --- *seminalis* Ell. and Ev., 175
 --- *setariae* Atk., 175
 --- *setariicola* Tehon and Daniels,
 175
 --- *sorgi* Ell. and Ev., 174, 176
 --- *striaeformis* Wint., 175
Cercospora, 176
 --- *herpotrichoides* Fron, 112,
 167, 176, 183, 188
 --- *holci* Sprague, 176, 177
 --- *poagena* Sprague, 176, 177
 --- *subulata* Sprague, 176, 178
Cladosporium *graminum* Cda., 113
 --- *herbarum* Lk., 113, 114, 178
 --- *malorum* Ruehle, 113
Colletotrichum, 137, 168
 --- *cereale* Manns, 168, 169
 --- *gloeosporioides* Penz., 169
 --- *graminicum* (Ces.) G. W.
 Wils., 168, 169
Coniothyrium *psammae* Oud., 126
Corticium *fuciforme* (Berk.)
 Wakef., 116
 --- *vagum* Berk. and Curt., 117
 --- --- *var. solani* Burt ex Rolfs,
 117
Curvularia, 178, 179
 --- *geniculata* (Tracy and Earle)
 Boed., 178, 179, 180, 193, 200
 --- *inaequalis* (Shear) Boed., 179
 --- *lunata* (Wakk.) Boed., 179,
 180
 --- *ramosa* (Bainier) Boed., 178,
 179
 --- *spicifera* (Bainier) Boed.,
 179, 199, 200
 --- *trifolii* (Kauff.) Boed., 179,
 195
Cylindrosporium *infusans* Ell.
 and Ev., 150

Darluca *filum* (Biv.) Cast., 125
Davisiella *elymina* (J. J. Davis)
 Petr., 171
Dilophia *graminis* Sacc., 126

Dilophospora alopecuri (Fr.) Fr.,
126

Diplodia zese (Schw.) Lév., 127

Diplodina, 120

--- *graminea* Sacc., 121, 124, 127

--- *lolii* Zimm., 124

Dothichloë atramentosa (Berk. and
Curt.) Atk., 111

--- *limitata* Diehl, 111

Dothidella aristidae (Schw.) Ell.
and Ev., 112, 171

Entyloma, 116

--- *bingenensis* Zundel, 116

--- *crastophilum* Sacc., 116

--- *irregulare* Johans., 116

--- (?) *spragueanum* Zundel, 116

Epichloë typhina (Fr.) Tul., 112

Eurychasmidium, 106

Fungi Imperfecti, 102, 103, 120

Fusarium, 180, 184, 185, 186

--- *avenaceum* (Fr.) Sacc., 181,
183

--- *culmorum* (W. G. Sm.) Sacc.,
181, 183, 185, 187, 188, 189

--- *equiseti* (Cda.) Sacc., 182,
183, 184

--- *graminearum* Schw., 112, 181,
183, 184, 187, 188

--- *moniliforme* Sheldon, 182, 185

--- --- *var. subglutinans* Wr. and
Reinking, 182, 185

--- *nivale* (Fr.) Ces., 182, 185

--- --- *var. majus* Wr., 186

--- *osiliense* Bres. and Vest.,
171

--- *oxysporum* Schlecht. em. Snyder
and Hansen, 180, 182, 183, 184,
186

--- *poae* (Pk.) Wr., 181, 186,
188, 189

--- *scirpi* Lamb. and Fautr., 182,
184, 187

--- --- *var. acuminatum* (Ell.
and Ev.) Wr., 181, 183, 187

--- --- *var. compactum* Wr., 188

--- --- *var. longipes* (Wr. and
Reinking) Wr., 188

--- --- *var. pallens* F. T. Ben-
nett, 184

(*Fusarium*) *solani* (Mart.) Appel
and Wr., 182

--- *spartinae* Ell. and Ev., 172

--- *sporotrichioides* Sherb., 181,
188

Fusoma biseptatum Sacc., 171

--- *triseptatum* Sacc., 171

Gibberella fujikuroi (Saw.) Wr.,
185

--- *zese* (Schw.) Petch, 112, 184

Gloeosporium sp., 168, 169

--- *graminum* Rostr., 169

Helminthosporium, 179, 189, 195,
198, 199, 202, 203

--- *avenae* Eidam, 189, 199

--- *bromi* Died., 115, 190

--- *californicum* Mackie and Pax-
ton, 190

--- *carbonum* Ullstrup, 202

--- *catenarium* Drechs., 203

--- *cyclops* Drechs., 191

--- *cynodontis* Marig., 127, 191

--- *denatioides* Bub. and Tréb.,
191

--- *dictyoides* Drechs., 191

--- *erythrospilum* Drechs., 192,
201

--- *flexuosum* Cda., 195

--- *giganteum* Heald and Wolf, 192

--- *gramineum* Rab., 192, 199

--- *hadrotrichoides* Ell. and Ev.,
193

--- *halodes* Drechs., 193

--- --- *var. tritici* Mitra, 193

--- *inconspicuum* Cke. and Ell.,
201

--- --- *var. buchloës* Ell. and
Ev., 194

--- M., 179

--- *monoceras* Drechs., 194

--- *poae* Baudyš, 195

--- *sativum* Pam., King, and Bakke,
172, 179, 180, 183, 189, 190,
194, 195, 197, 198, 203

--- *setariae* Lind., 197

--- *setariae* Saw., 194, 197, 203

--- *siccans* Drechs., 198

--- *stenacrum* Drechs., 192, 198

--- *teres* Sacc., 199

- (*Helminthosporium*) *tetramera*
 McKinney, 179, 199
 --- *triseptatum* Drechs., 200
 --- *tritici-repentis* Died., 201
 --- *turcicum* Pass., 194, 198,
201, 203
 --- *vagans* Drechs., 195, 202
Hendersonia, 125, 127, 164, 167
Heterosporium avenae Oud., 203,
204
 --- *graminis* McAlp., 204
 --- *hordei* Bub., 204
 --- *phlei* Greg., 204
 --- *phragmitis* Sacc., 204
Hormodendrum cladosporioides
 Sacc., 113
Hypochnus filamentosus Pat., 117

Leptosphaeria avenaria G. F.
 Weber, 146
 --- *herpotrichoides* De N., 112

Macrophoma hennebergii (Kuehn)
 Berl. and Vogl., 153
 --- *phlei* Tehon and Stout, 130
Macrophomina phaseoli (Maubl.)
 Ashby, 130
Mastigosporium, 205
 --- *album* Reiss, 126, 205, 206
 --- --- *var. athrix* Ericks., 171
 --- *calvum* (Ell. and Ev.) Sprague,
 206
 --- *cylindricum* Sprague, 205
 --- *rubricosum* (Dearn. and Barth.)
 Sprague, 205, 206
Melanconiales, 168
Moniliales, 172
Mycosphaerella, 138
 --- *longissima* Fckl., 114
 --- *tassiana* (De N.) Johans., 114
 --- *tulasnei* (Jancz.) Rothers,
113, 178

Napicladium arundinaceum (Cda.)
 Sacc., 206
 --- *gramineum* Pk., 195, 212
Nigrospora oryzae (Berk. and Br.)
 Fetch, 207
Non-sporulating basidiomycetes,
 119

Ophiobolus graminis Sacc., 114,
189
 --- *setariae* Ito and Kuribayashi,
 197
Ophiocladium hordei Cav., 207
Ovularia hordei (Cav.) Sprague,
207
 --- *lolii* Völkart, 208
 --- *pulchella* (Ces.) Sacc., 208
 --- --- *var. agropyri* J. J. Davis,
 208
 --- --- *var. lolii-italici* Ferr.,
 208
 --- *pusilla* (Ung.) Sacc., 208

Passalora dactylina Pass., 212
 --- *graminis* (Fckl.) Hoehn., 212
Pellicularia filamentosa (Pat.)
 Rogers, 117
 --- *vaga* (Berk. and Curt.) Rogers
 ex Linder, 117
Penicillium spp., 208
 --- *expansum* Lk., 209
 --- *oxalicum* Currie and Thom, 209
Phaeoseptoria, 128, 130, 148
 --- *phalaridis* (Träfl) Sprague,
 148
Phlyctaena bromi, 133
Phoma, 131, 168
 --- *hennebergii* Kuehn, 153
 --- *insidiosa* Tassi, 132
 --- *secalina* Jancz., 157
 --- *terrestris* Hansen, 131
Phycomycetes, 102, 103, 105
Phyllachora, 115, 116, 171
Phyllosticta, 131, 160
 --- *anthoxella* Sprague, 132
 --- *avenophila* Tehon and Daniels,
 132
 --- *owensii* Sprague, 131, 132
 --- *rogleri* Sprague, 131, 132
 --- *sorghina* Sacc., 131, 132
 --- *stomaticola* Bauml., 134, 136
Physoderma sp., 203
Phytophthora spp., 105
 --- *cactorum* (Leb. and Cohn)
 Schroet., 105, 106
 --- *calocasiae* Rac., 105
Piricularia grisea (Cke.) Sacc.,
209

- (Piricularia) oryzae Briosi and Cav., 209.
 Pleospora herbarum (Pers.) Rab., 214
 Protomyces rhizobius Trail, 106
 Pseudodiscosia avenae Sprague and A. G. Johnson, 169
 Puccinia coronata Cda., 183
 Pyrenophora bromi (Died.) Drechs., 115, 190
 --- teres (Died.) Drechs., 199
 --- tritici-repentis (Died.) Drechs., 201
 Pythium, 107, 108, 109, 189
 --- aristosporum Vanterpool, 107, 109
 --- arrhenomanes Drechs., 107, 109, 111, 195, 196
 --- complens A. Fischer, 110
 --- debaryanum Hesse, 109
 --- graminicolum Subr., 107, 108, 109
 --- hypogynum Middleton, 110
 --- irregulare Buis., 109
 --- iwayama Ito, 110
 --- monospermum Pringsh., 110
 --- peridium Drechs., 110
 --- polymorphon Sideris, 110
 --- rostratum Butl., 111
 --- tardicrescens Vanterpool, 111
 --- ultimum Trow, 109
 --- vexans D By., 110
 Ramularia pulchella Ces., 208
 Rhabdospora groenlandica Lind, 136
 --- lolii Cast., 163
 Rhizoctonia, 118
 --- monteithianum F. T. Bennett, 115
 --- solani Kuehn, 117, 185, 200
 Rhynchosporium, 209
 --- orthosporum Caldwell, 209, 210
 --- secalis (Oud.) J. J. Davis, 209, 210
 Sclerotinia graminearum Elen., 119
 --- homoeocarpa F. T. Bennett, 115
 Sclerotium bataticola Taub., 130
 --- constantini Foëx and Rosella, 118
 --- fulvum Fr., 118, 119
 --- rhizodes Auers., 118
 Scolecotrichum, 212
 --- compressum Allesch., 212
 --- graminis Fekl., 132, 195, 204, 211, 213
 --- --- var. brachypoda Speg., 212
 --- --- var. nanum Sacc., 212
 --- maculicolum Ell. and Kell., 213
 Selenophoma, 132, 159
 --- bromigena (Sacc.) Sprague and A. G. Johnson, 133
 --- donacis (Pass.) Sprague and A. G. Johnson, 133, 134, 136, 137, 139, 147
 --- --- var. stomaticola (Baüml.) Sprague and A. G. Johnson, 133, 134, 137, 138, 139, 159
 --- everhartii (Sacc. and Syd.) Sprague and A. G. Johnson, 133, 134, 137
 --- obtusa Sprague and A. G. Johnson, 133, 139
 Septogloeum, 170, 171
 --- athrix (Eriks.) Sprague, 171
 --- oxysporum Sacc., Bomm. and Rouss., 112, 170
 --- spartinae (Ell. and Ev.) Wr., 172
 Septoria, 118, 125, 139, 147, 154, 157, 160, 164, 170
 --- agropyri Ell. and Ev., 150
 --- agropyrina Lob., 142, 144, 150, 154, 164, 165
 --- endropogonis J. J. Davis, 141, 145, 146, 153, 157
 --- --- f. sporobolicola Sprague, 141, 142, 145, 157, 158
 --- --- var. sorghastri H. C. Greene and Sprague, 144, 145, 146
 --- annua Ell. and Ev., 151
 --- arctica Berk. and Curt., 142, 146
 --- avenae Frank, 141, 146, 151, 153, 158, 164, 165

- (Septoria) briosiana Mor., 162
 --- bromi Sacc., 140, 147
 --- --- var. alopecuri Karst.,
 153
 --- --- var. phalaricola Sprague,
 142, 147, 148
 --- --- var. phalaridis Trail,
 148
 --- caballeroi Gonz. Frag., 148
 --- --- var. panicei Gonz. Frag.,
 148
 --- calamagrostidis Ell. and Ev.,
 138
 --- calamagrostidis (Lib.) Sacc.,
 143, 148, 149, 160
 --- --- f. koeleriae (Cocc. and
 Mor.) Sprague, 143, 149
 --- calamovillae Petr., 140, 149
 --- capillatae Trott., 159
 --- cenchrina J. J. Davis, 140,
 149
 --- culmifida Lind., 136
 --- curva Karst., 134
 --- donacis Pass., 133
 --- elymi Ell. and Ev., 140, 141,
 144, 150, 157
 --- elymi-europaei Jaap, 150
 --- elymicola Died., 144
 --- everhartii Sacc. and Syd.,
 137
 --- falcispora Demidova, 136
 --- festucae Died., 160
 --- graminum Desm., 161
 --- --- f. triseti-loeflingianii
 Cav., 148
 --- --- var. crassipes Grove,
 161
 --- --- var. lolii Mont., 163
 --- infuscans (Ell. and Ev.)
 Sprague, 140, 142, 144, 150,
 155, 164
 --- jaculella Sprague, 142, 146,
 147, 151
 --- koeleriae Cocc. and Mor.,
 149
 --- loligena Sprague, 124, 141,
 151, 158, 163, 164
 --- lolii (Cast.) Sacc., 151,
 163
 --- lolii West., 124, 163
 (Septoria) lunata Grove, 135, 159
 --- macropoda Pass., 141, 143,
 151, 152
 --- --- var. grandis Sprague,
 143, 152
 --- --- var. septulata (Gonz.
 Frag.) Sprague, 143, 152
 --- macrostoma Speg., 148
 --- melicae Pass., 141, 153, 158,
 164
 --- microspora Ell., 156
 --- mississippiensis Sprague, 142,
 146, 153
 --- munroae Ell. and Barth., 142,
 153
 --- nebulosa Rostr., 134
 --- nodorum Berk., 113, 140, 153,
 158, 164, 165, 201
 --- oudemansii Sacc., 122, 140,
 154, 164
 --- oxyspora Penz. and Sacc., 133
 --- pacifica Sprague, 142, 150,
 155
 --- passerinii Sacc., 141, 156
 --- phalaridis Cocc. and Mor.,
 148
 --- poae-annuae Bres., 151
 --- --- var. septulata Gonz. Frag.,
 152
 --- poliomela Syd., 139, 156
 --- quinqueseptata Sprague, 144,
 145, 146, 157
 --- secalina (Jancz.) Sacc., 157
 --- secalis Prill and Del., 140,
 151, 157, 158, 164
 --- --- var. stipae Sprague, 141,
 149, 158
 --- spartinae (Trel.) Sprague,
 140, 158
 --- stipae Died., 159
 --- stipae Trabut, 159
 --- stipina Died., 141, 159
 --- tenella Cke. and Ell., 140,
 159
 --- triseti Speg. em. Sprague,
 140, 149, 160
 --- tritici Rob., 144, 150, 152,
 160, 161, 162, 163
 --- --- f. avenae (Desm.) Sprague,
 143, 162

- (*Septoria*) *tritici* f. *holci*
 Sprague, 144, 162
 --- var. *lolicola* Sprague
 and A. G. Johnson, 124, 144,
 151, 162
Septoriopsis, 159
Sphaeropsidales, 120
Sporotrichum sp., 213
Stagonospora, 125, 147, 163, 167
 --- *egrostidis* Syd., 165
 --- f. *angusta* Sprague, 163,
164, 165
 --- *arenaria* Sacc., 144, 150,
 151, 154, 155, 164, 165, 166
 --- *arrhenatheri* A. L. Sm. and
 Ramsb., 164, 165
 --- *bromi* A. L. Sm. and Ramsb.,
 164, 165
 --- *foliicola* (Bres.) Bub., 148,
 164, 165, 166
 --- *glyceriae* Roum. and Fautr.,
 154
 --- *hennebergii* (Kuehn) Petr.
 and Syd., 153
 --- *intermixta* (Cke.) Sacc., 166
- (*Stagonospora*) *simplicior* Sacc.
 and Berl., 163, 166
 --- var. *andropogonis* Sacc.,
 166
 --- *subseriata* Desm., 163, 167
 --- var. *maculata* Grove, 167
 --- *vexata* var. *baldingeriae* Sacc.,
 166
 --- var. *foliicola* Bres., 166
Stemphylium, 172, 214
 --- *botryosum* Wallr., 214
Synchytrium, 106
- Typhula*, 118, 119
 --- *borealis* Ekstrand, 119
 --- *graminum* Karst., 119
 --- *idahoensis* Remsberg, 118, 119
 --- *itoana* Imai, 118, 119, 186
- Wojnowicia graminis* (McAlp.) Sacc.
 and D. Sacc., 128, 129, 130,
167