Harmer, S. F. Subantarctic whales. (Abstract)

H

(1919)

HARVARD UNIVERSITY.

LIBRARY

OF THE

MUSEUM OF COMPARATIVE ZOÖLOGY
62.335

GIFT OF

Thomas Barbour

October 18. 1923.

ROYAL INSTITUTION OF GREAT BRITAIN.

WEEKLY EVENING MEETING.

Friday, May 16, 1919.

SIR JAMES CRICHTON-BROWNE, J.P. M.D. LL.D. F.R.S.,
Treasurer and Vice-President, in the Chair.

S. F. HARMER, Sc.D. F.R.S., *M.R.I.*, Director of the Natural
History Departments of the British Museum.

Subantarctic Whales and Whaling.*

[ABSTRACT.]

THE history of Whaling in Northern waters, and of the hunting of Sperm Whales in warmer seas, has often been written, and some of the principal facts relating to these subjects are matters of common knowledge. There is reason to believe that the existence of a whaling industry, which was inaugurated just outside the South Polar Circle after the commencement of the present century, is by no means generally known. Although Captain Cook, Sir James Ross and others had many years before reported the presence of whales in those latitudes, no practical advantage was taken of the information under fourteen years ago; and since that date the industry has eclipsed in importance all that had been done previously, even when the Greenland Whale "fishery" was at its height.

In 1892 Capt. C. A. Larsen left Norway for the far South, which he reached in the October of that year. No whaling was done, and an Expedition to the same regions, fitted out by Capt. Svend Foyn in the next year, was also unproductive of whales, mainly for the reason that it had been intended to hunt Right Whales and Sperm Whales. The Norwegian captains brought home, however, a very vivid impression of the enormous number of whales frequenting Subantarctic waters; and the fact that they at first made no further ventures was due to the profitable nature of the whaling in the neighbourhood of the Norwegian coasts. Dr. W. S. Bruce had simultaneously (1892) accompanied four vessels of the Dundee whaling fleet to the Antarctic, and had been similarly impressed with the abundance of whales in these waters. A meeting was shortly afterwards held in the rooms of the Royal Scottish Geographical Society at Edinburgh, to advocate the use of the modern Norwegian methods of whaling in the South; but the proposal was not carried, and no practical steps were taken.

Capt. Larsen subsequently became the Commander of the

* Published by permission of the Trustees of the British Museum.

"Antarctic," the exploring vessel of Dr. O. Nordenskjöld's Swedish South Polar Expedition, 1901-1903. The "Antarctic" was wrecked; and Capt. Larsen, on his return journey, found himself at Buenos Aires, where in 1904 he founded the *Compania Argentina de Pesca*, the first Whaling Company which undertook operations in the far South. This Company commenced work at South Georgia in 1905, while the South Shetland Islands were visited, with the same object, in 1906, and the South Orkney Islands in 1911. The operations have proved so successful that there are now numerous companies whaling at South Georgia and the South Shetlands, a large proportion being Norwegian. While the most successful whalers in the Greenland industry, from the seventeenth to the nineteenth Centuries, were the British and the Dutch, the Norwegians have almost a monopoly of the art at the present time, and nearly all the skilled workers are of that nationality.

The older whalers hunted with hand-harpoons from small boats, provided with sails and oars, which were launched from the parent ship on sighting a whale. The objects of their chase were principally the Greenland Whale, the Atlantic Right Whale and the Sperm Whale; and they were unable to attack and capture the larger and swifter Rorquals.

About 1865 the Norwegian whaling Captain Svend Foyn invented the modern whaling-gun, which was fitted with an explosive tip and with a barbed harpoon carrying a strong rope. The explosion was regulated so as to occur immediately after the harpoon hit the whale, which is sometimes killed at once, and in any case is severely injured by a successful shot. The gun is carried in the bow of a steam-whaler, which chases the animal until a favourable opportunity for shooting occurs. These methods have revolutionised whaling, and there is now no whale which is too large to be captured.

In the prosperous days of the Greenland whale "fishery," 1437 whales were caught by 76 ships in 1814—an average of not quite twenty whales to each vessel—and this is mentioned by Scoresby (1820) as a specially good year. At the present day the number of whales caught by a single vessel, during the whaling season of six months, may rise to over 300; and the total number caught off South Georgia and the South Shetlands together has exceeded 10,000 in one year. Bearing in mind the universal history of whaling in the past—a period of prosperity succeeded by a rapid decline and a final abandonment of the industry—the question arises whether there is not a serious danger that Subantarctic whaling will have a similar experience.

The question of the disappearance of the whales is not merely a sentimental one, though Zoologists would naturally view their extermination with deep concern on scientific grounds. The plea for their preservation may be strengthened, however, by emphasising the fact that they are of the highest economic importance. The

baleen or whalebone of the Right Whales is a material of much practical utility for many purposes; but its importance is almost negligible compared with that of the oil which is derived from the blubber and other parts of whales. Whale-oil can be readily transformed into soap and glycerine, while it is possible to prepare from it a fat which is perfectly inodorous and is utilised in the manufacture of margarine. During the War it has been of vital importance. Enormous quantities of glycerine derived from it have been used for the manufacture of explosives, and it has been hardly less important in its relation to the food-supply. If whale-oil had not been obtainable, the glycerine which was essential for our National security must have been derived from other animal fats or from vegetable oils; and the shortage of fat required as food would have been very serious. After the oil has been extracted on the whaling grounds, the remainder of the carcass may be dried and ground down into "guano," which is valuable as a fertiliser for crops and is also utilised for the preparation of cattle-foods. It is not always possible to carry out this part of the process; and an enormous waste of valuable material may result from this omission. Since it is well known that the flesh of Cetacea is fit for human food, it is by no means impossible that a part of the enormous quantity of meat which might be obtained from them may be so utilised in the future.

Although a few Right Whales and Sperm Whales are captured by the Subantarctic whalers, who occasionally kill some of the smaller Cetacea as well, the industry in these waters is almost entirely confined to three species of the larger whales. Of these the Humpback rarely exceeds 55 feet in length, the Fin Whale is not much more than 85 feet, while the Blue Whale, probably the largest animal that has ever existed, is sometimes over 100 feet long. In the first few years of Subantarctic whaling, the Humpback constituted nearly the whole catch, even more than 96 per cent. in 1910-11, when 5299 individuals of this species were captured off South Georgia in the six months of the principal whaling season. In 1912-13 the number of Humpbacks caught in the same locality, in the corresponding six months, fell to 2251 (about 53 per cent. of the total catch), and in 1913-14 it was reduced to 474 (about 18 per cent.). This diminution, which has persisted to the present time, has been due largely to a reduction of the number of Humpbacks frequenting South Georgia; but it has been partly caused by an increase in the size of the whaling vessels and of the strength of the tackle employed, enabling the whalers to hunt the larger kinds, which naturally yield more oil and other products than the comparatively small Humpback. The whaling industry thus depends at present almost entirely on the Fin Whale and the Blue Whale. It is noteworthy that the Fin Whale, of intermediate size, first rose to prominence on the decline of the Humpback; but the gigantic Blue Whale has now surpassed it and has become the favourite object of the whalers' pursuit.

There is at present no certain evidence of the reduction in numbers of Fin Whales and Blue Whales.

In explaining the reduced number of Humpbacks frequenting the whaling grounds, the whalers rely on the hypothesis that individuals of this species are of a timid nature, and are readily frightened away from a locality by pursuit. There is probably some truth in this view, but it is at least possible that the reduction is due to a diminution in number of the total stock of Humpbacks. Whales are migratory animals, and their movements are almost certainly influenced by two causes : (1) the distribution of their food-supply ; (2) the position of their breeding grounds. The Plankton-organisms on which the Whalebone Whales subsist are present in vast quantities in Polar waters during the summer, and the whales are accordingly found there at this period. Towards the end of the summer or the beginning of autumn most of them forsake high latitudes. The Southern Humpback executes extensive migrations northwards, along the coasts of the great Southern Continents, to the neighbourhood of the Equator and even beyond it. It is in these warmer waters that it is known to breed, and it subsequently proceeds southwards in the ensuing spring. One of the most alarming facts about this species is that it has been extensively hunted along the coasts of Africa, South America and elsewhere. Although it is not possible to assert positively that the South Georgia Humpbacks are thus affected, there are strong reasons for believing that this is the case ; and it would thus follow that this species is persecuted in Subantarctic waters during the summer, and further north during other parts of the year. Remembering that the old whalers reduced the Greenland Whale almost to the point of extermination by the use of what may now be regarded as primitive methods, and that a similar fate has befallen the once flourishing whaling industries of other localities, it thus appears that there are the most urgent reasons for seeking to afford some immediate measure of protection to this and other species of whales.

In devising methods for the protection of animals, the principle of saving them from being hunted during their breeding season has been found specially effective. It is very difficult to get complete information on this subject with regard to whales, but one of the ways in which a conclusion may be reached is the examination of foetal records, a method which has already been adopted with some success by Guldberg and others. By the study of a relatively large mass of statistics which has been supplied to the British Museum (Natural History) by the Whaling Companies operating off South Georgia, I have found it possible to arrive at certain definite results which amplify or correct those of previous observers. These are specially clear in the case of the Fin Whale, which has provided the largest number of records. It is found that in a given month there is a particular length of foetus which has the greatest frequency ;

that in the next month a greater length is most common ; and so on for several of the succeeding months, after which the whaling season comes practically to an end, and the number of records is inadequate to give a normal result. Although there are certain irregularities needing explanation in the graphs constructed from the statistics, the general result has been arrived at that, in each of the three species principally hunted at South Georgia, pairing takes place with greatest frequency at a certain period of the year, and that a normal curve of pairing can be drawn. This result gives in the main a satisfactory explanation of the statistical records. The season when pairing is at its height falls, in each case, outside the period when whaling is actively carried on in the far South ; and the important conclusion is reached that if the whales are to be protected during their breeding season, it must probably be done in regions of the world further North than South Georgia. The validity of the Southern figures, which have no doubt been roughly recorded by the whalers, has been confirmed by obtaining corresponding results from the examination of the statistical foetal records of Northern whales.

It can hardly be doubted that protective measures of some kind are urgently necessary now, or will at least become so in the near future ; although it is by no means certain what form they should take. The British Government is in some respects in a specially favourable position with regard to this matter, since all the important whaling grounds of the Subantarctic region belong to the Dependencies of the Falkland Islands and lie in its jurisdiction. It is satisfactory to be able to conclude with the statement that the Government is fully alive to the necessity of taking steps before it is too late, and that an Interdepartmental Committee is at present engaged, under the auspices of the Colonial Office, in framing a scheme for an Expedition which is to investigate the whaling problem on the spot, with the view of obtaining information on which legislation may be based.

[S. F. H.]

Photomount
Pamphlet
Binder
Gaylord Bros.
Makers
Syracuse, N. Y.
PAT. JAN 21, 1908

MCZ ERNST MAYR LIBRARY

3 2044 128 397 270

M