Botany Library

580.7

F45 Library of

(2)

Wellesley College

PRESENTED BY

The Museum

127483

580.7
745

PUBLICATIONS
OF THE
FIELD MUSEUM OF NATURAL
HISTORY

BOTANICAL SERIES

VOLUME II

CHICAGO, U. S. A.
1900-1916.*

*For actual dates of issue of each paper see Table of Contents.

FIELD COLUMBIAN MUSEUM.

PUBLICATION 136.

BOTANICAL SERIES.

VOL. II, No. 7.

PRÆNUNCIÆ BAHAMENSES—II.

CONTRIBUTIONS TO A FLORA OF THE
BAHAMIAN ARCHIPELAGO.

BY

CHARLES FREDERICK MILLSPAUGH, M.D.

Curator, Department of Botany.

CHICAGO, U. S. A.

August, 1909.

Issued

August 6, 1909.

Digitized by the Internet Archive
in 2014

PRÆNUNCIÆ BAHAMENSES — II.

CONTRIBUTIONS TO A FLORA OF THE BAHAMIAN ARCHIPELAGO.

CHARLES F. MILLSPAUGH, M.D.

Since the issue of the first of these papers* the co-workers upon the Flora of the Bahamian Archipelago have continued their investigations as outlined in that publication and have extended the survey to embrace Andros, Eleuthera, Little San Salvador, Cat Island, Conception Island, Rum Cay, Watling's, Atwood's Cay (Samana), Long Island, Great Ragged Island, Crooked Island, Fortune Island, Turk's Islands and the Caicos and Cay Sal Groups.

The collections have been increased to some 12,000 sheets and embrace the field work of the following collectors in the localities indicated under each:

Brace — Mr. Louis J. K. Brace, in addition to the field work outlined on page 138, has been commissioned by the New York Botanical Garden and this Museum to make three further explorations. From these he has returned over 1,800 numbers as follows: Rum Cay (3919-3993); Fortune Island (3994-4179); Acklin's Island (4260-4471); Crooked Island (4533-4776); and Andros (4876-5377 and 6657-7139). In the Andros work he not only traversed the East side but also made his way around to the difficult swash region of the West coast, and penetrated the dividing channel across the southern third of the Island. Mr. Brace has also continued his search for interesting plants on New Providence (7141-7161 and 7905-7910). Our collections of his plants have been further enriched through securing his personal set of the series he collected in 1886 which formed the basis of Gardiner & Brace's Plants of the Bahama Islands.†

*Field Col. Mus. Bot., 2: 137-184. Feb. 17. 1906.

†Gardiner & Brace per Dolley in Proc. Acad. Nat. Sci. Phila., 1889, 349-426.

Britton — Dr. N. L. Britton, see page 138.

Britton & Brace — Dr. N. L. Britton and Mr. Louis J. K. Brace, see page 139.

Britton & Millspaugh — Dr. N. L. Britton and the author, in addition to the field work outlined on pages 139-140, undertook, in 1907, a very exhaustive exploration of the Out Islands, embracing the following localities: Eleuthera, from the Glass Window to Harbor Island (5376-5404); Glass Window to Gregorytown (5405-5440); Governor's Harbor and vicinity (5441-5542); Rock Sound and vicinity (5543-5590); Miller's and Bannermantown and vicinities (5591-5613) and in the vicinity of the extreme southern point of the island (5614-5656). Little San Salvador (5657-5701). Cat Island, from Orange Creek around the extreme north end of the island, down the northeast shore to Cotton Point and across (5702-5791); the vicinity of The Bight and across the island and back (5792-5945); and Port Howe and vicinity (5946-5986). Conception Island (5987-6043). Watling's Island, Cockburntown vicinity (6044-6144); from Cockburntown across Little and Great Lakes to the Light on the east coast (6145-6169); Graham's Harbor south along the east coast to Columbus' Monument (6170-6188); Graham's Harbor around the north end of the island to Cockburntown (6189-6224). Long Island, vicinity of Clarencetown and across the island to the west shore, and on Thatch and Strachan's Cays (6225-6359); and the extreme north end of the island at Cape St. Maria (6360-6370). New Providence in the vicinity of Nassau (5354-5366).

Coker — Prof. William C. Coker, see page 140.

Cooper — Dr. William Cooper, see page 140.

Curtiss — Mr. A. H. Curtiss, see page 140.

Earle — Prof. F. S. Earle, see page 140.

E. G. Britton — Mrs. N. L. Britton, in addition to her field work noted on page 141, accompanied her husband and the author as far as Harbor Island and remained there, as a base, while working that island and the extreme north end of Eleuthera (6370-6527). Returning to New Providence she did further discriminating field work on that island (6529-6656) while awaiting the return of the Out Island Expedition.

Hitchcock — Prof. Albert S. Hitchcock, see page 141.

Through a later purchase of Prof. Hitchcock's "Florida Herbarium" this Museum secured 133 duplicate sheets of his Bahamian collections, these are in addition to his original collection which is still a loan to this Museum by the Missouri Botanical Garden.

Howe — Dr. M. A. Howe, in addition to his field work on the algae of the

Bahamas as indicated on page 141, accompanied Mr. Percy Wilson on the expedition to the southern islands (see Wilson) whence he returned a large amount of interesting and critical additional material.

Millsbaugh — The author, see page 141.

Nash & Taylor — Dr. George V. Nash and Mr. Norman Taylor, see page 142.

Northrop — Mrs. Alice (Rich) Northrop, see page 142.

Rothrock — Prof. J. T. Rothrock made, in the interests of the University of Pennsylvania, an expedition to the British West Indies in his 51-ton Yawl "White Cap" during the winter of 1890-91. On this expedition he collected 4 days on New Providence; 2 on Cat Island; a part of one day on Watling's; 2 days on Crooked Island; 2 on Fortune Island, and a day on Great Inagua. His collecting was done independently from that of Prof. Hitchcock, who accompanied him. The first set was deposited in the herbarium of the University of Pennsylvania; the second, comprising 162 sheets, he has kindly placed in the herbarium of this Museum.

Wight — Mr. Alex. E. Wight, under the patronage of the Gray Herbarium, Cambridge, Mass., made a Bahamian collection of 281 numbers from December, 1904, to May, 1905. He collected on New Providence and Hog Island (1-226 and 271-274), and on Andros at Mangrove Cay, Fresh Creek, Calabash Bay, and Small Hope (227-270). The resulting plants were determined by the author and the initial set retained at this Museum (with the exception of about six numbers in the Orchidaceae which are in the herbarium of Prof. Oakes Ames). Duplicate sets have been deposited in the Gray Herbarium and the herbarium of The New York Botanical Garden. [This paragraph should substitute that concerning Mr. Wight's collections on page 142.]

Wilson — Mr. Percy Wilson, of the New York Botanical Garden, on a commission from that institution and this Museum, and accompanied by Dr. Howe, made an exploration of the southeastern islands from November 22d to December 29th, 1907. His field was as follows: Cat Island, at The Bight (7163-7197) and the southwest end of the island (7198-7202). Watling's Island, at Cockburntown (7203-7225); Graham's Harbor (7226-7258); and the extreme southeast and southwest ends (7259-7336 and 7337-7359). Atwood's Cay (Samana) (7360-7424 and 7903-7904). Mariguana, at a point ten miles west of Abraham Bay (7425-7451); Abraham Bay and vicinity (7452-7540); five miles west of Southeast Point (7541-7557) and at Southeast Point (7558-7589). The Caicos Islands: on South Caicos (7590-

7693); Pine Cay (7694-7697); North Caicos at Kew and vicinity (7698-7748); Providenciales (7749-7752) and West Caicos (7753-7763); Little Inagua, at the western end (7764-7782). Castle Island (7783-7802). The Ragged Cays, at Great Ragged Island (7803-7869) and Hog Cay (7870-7882). The Exuma Chain, at Harvest Cay (7883-7898); and Rose Island (7899-7902). From May 13 to June 7, 1909, he explored the Cay Sal Bank, spending four days on Anguilla Islands (7932-8078) and one day each on Salt Cay (8079-8124), Water Cay (8132-8159), and Elbow Cay (8125-8131). On the trip out he spent one day at Riding Rocks (7912-7924), and one on Orange Cay (7925-7931), of the Bimini Group; and on his return to Nassau: six days collecting on New Providence (8168-8247, 8323-8343, 8370-8413, 8434-8436) and four on Hog Island (8248-8322, 8344-8369, 8414-8433). The collections are deposited in duplicate in the herbaria of The New York Botanical Garden and this Museum.

Islands Represented in these Collections.

- Abaco — *Brace, Coker.*
Acklin's Island — *Brace.*
Allen's Cay (Abaco) — *Brace.*
Andros — *Brace, Coker, Northrop, Wight.*
Anguilla Island — *Wilson.*
Atwood's Cay (Samana) — *Howe, Wilson.*
Big Golden Cay (Andros) — *Wight.*
Bimini, North — *Brace, Howe, Millspaugh.*
Bimini, South — *Brace, Howe, Millspaugh.*
Broad Rock Cay (Andros) — *Brace.*
Caicos, North — *Howe, Wilson.*
Caicos, South — *Howe, Wilson.*
Caicos, West — *Howe, Wilson.*
Castle Island — *Howe, Wilson.*
Cat Cay, North (Biminis) — *Brace, Howe, Millspaugh.*
Cat Cay, South (Biminis) — *Howe, Millspaugh.*
Cat Island — *Britton & Millspaugh, Hitchcock, Howe, Wilson, Rothrock.*
Cave Cay (Exuma Chain) *Britton & Millspaugh, Howe.*
Conception Island — *Britton & Millspaugh.*
Crab Cay (Watling's Isl.) — *Britton & Millspaugh.*
Crooked Island — *Brace, Hitchcock, Rothrock.*
Delectable Cay (Acklin's) — *Brace.*
Eleuthera — *Britton & Millspaugh, Coker, E. G. Britton, Hitchcock, Rothrock.*
Elbow Cay (Abaco) — *Brace, Coker.*
Elbow Cay (Cay Sal Bank) — *Wilson.*
Exuma — *Britton & Millspaugh, Howe.*
Fortune Island (or Long Cay) — *Brace, Hitchcock, Rothrock.*
Frozen Cay (Berry Is.) — *Britton & Millspaugh, Howe.*
Galiot Cay, Great (Exuma Chain) — *Britton & Millspaugh, Howe.*
Galiot Cay, Little (Exuma Chain) — *Britton & Millspaugh, Howe.*
Garden Cay (Great Bahama) — *Brace.*
George's Island (Eleuthera) — *Coker.*
Goat Cay (Berry Is.) *Britton & Millspaugh.*
Goat Cay (Andros) — *Brace.*
Great Bahama Island — *Brace, Britton & Millspaugh, Howe.*
Green Cay — *Coker.*
Green Turtle Cay (Abaco) — *Brace.*

- Guana, Great (Exuma Chain) — *Britton & Millspaugh, Howe*.
Gun Cay (Biminis) — *Howe, Millspaugh*.
Harbor Cay, Great (Berry Is.) — *Britton & Millspaugh, Howe*.
Harbor Cay, Little (Berry Is.) — *Britton & Millspaugh, Howe*.
Harbor Island — *E. G. Britton*.
Harvest Cay (Exuma Chain) *Howe, Wilson*.
High Point Cays (Andros) — *Brace*.
Hog Cay (Ragged Group) — *Howe, Wilson*.
Hog Island (New Providence) — *Brace, Britton, E. G. Britton, Britton & Brace, Britton & Millspaugh, Northrop, Wight, Wilson*.
Inagua — *Hitchcock, Nash & Taylor, Rothrock*.
Inagua, Little — *Howe, Nash & Taylor, Wilson*.
Isaac, Great — *Brace*.
Joulter's Cays (Andros) — *Howe, Millspaugh*.
Lignum Vitae Cay (Berry Is.) — *Britton & Millspaugh, Howe*.
Little San Salvador — *Britton & Millspaugh*.
Long Bay Cays (Andros) — *Brace*.
Long Cay, see Fortune Island.
Long Island — *Coker, Britton & Millspaugh*.
Mangrove Cay (Andros) — *Brace, Coker, Wight*.
Mangrove Cay, Little (Andros) — *Brace, Coker*.
Man o'War Cay (Abaco) — *Brace*.
Mariguana — *Howe, Wilson*.
New Providence — *Brace, Britton, E. G. Britton, Britton & Brace, Britton & Millspaugh, Coker, Cooper, Curtiss, Earle, Hitchcock, Howe, Millspaugh, Northrop, Rothrock, Wight, Wilson*.
No Harbor Cay (Rose Island) — *Britton & Millspaugh*.
Orange Cay — *Wilson*.
Pigeon Cay (Abaco) — *Brace*.
Pine Cay (Caicos) — *Howe, Wilson*.
Providenciales — *Howe, Wilson*.
Ragged Island, Great — *Howe, Wilson*.
Riding Rocks — *Wilson*.
Rose Cay (Andros) — *Brace*.
Rose Island — *Britton & Millspaugh, Howe, Northrop, Wilson*.
Rum Cay — *Brace, Coker*.
Salt Cay (Cay Sal Bank) — *Wilson*.
Salt Cay (New Providence) — *Northrop*.
Samana, see Atwood's Cay.
San Salvador, Little — *Britton & Millspaugh*.
Sheep Cay (Inagua) — *Nash & Taylor*.

- Ship Channel Cay (Exuma Chain) — *Britton & Millspaugh, Howe.*
Silver Cay (New Providence) — *Howe, Millspaugh.*
Spanish Cay (Abaco) — *Brace.*
Stocking Island (Exuma Chain) — *Britton & Millspaugh, Howe.*
Strahan's Cay (Long Isl.) — *Britton & Millspaugh, Howe.*
Sturup Cay (Berry Is.) — *Britton & Millspaugh, Howe.*
Thatch Cay (Long Isl.) — *Britton & Millspaugh.*
Turk Island, Grand — *Nash & Taylor.*
Water Cay (Cay Sal Bank) — *Coker, Wilson.*
Watling's Island — *Britton & Millspaugh, Coker, Hitchcock, Howe, Rothrock, Wilson.*
Whale Cay (Berry Is.) — *Britton & Millspaugh, Howe.*
Wide Opening, Cay North of (Exuma Chain) — *Britton & Millspaugh, Howe.*
-

The following, chronologically arranged, additional publications, based upon critical examination of the above collections, have appeared since the issue of *Prænuncia I* and may be noted in connection with the list on page 144:

- "The Polyporaceae of North America—IX." W. A. Murrill in Bull. Torrey Club, 31: 604. *Sesia striata* (Sw.) Murr. (Jan. 9, 1905.)
"Additions to the Flora of Subtropical Florida." John K. Small in Bull. N. Y. Bot. Gard., 3: 424, 437. *Linum Curtissii*, *Scutellaria longiflora*. (Jan. 27, 1905.)
"The Polyporaceae of North America—XIII." W. A. Murrill in Bull. Torrey Club, 32: 648, 651, 652. *Cariolus membranaceus* (Sw.) Pat., *C. pinisitus* (Fr.) Pat., *C. arenicolor* (B. & C.) Murr. (Jan. 22, 1906.)
"*Prænunciæ Bahamenses*—I." C. F. Millspaugh in Pub. Field Mus. Bot. Ser., 2: 137–184. (Feb. 17, 1906.)
"Contributions to the Flora of the Bahama Islands—III." N. L. Britton in Bull. N. Y. Bot. Gard., 4: 137–143. (Mar. 19, 1906.)
"The Genus *Vernonia* in the Bahamas." Henry A. Gleason in Bull. Torrey Club, 33: 183–188. (Apr. 7, 1906.)
"*Tectaria minima* sp. nov." Lucien M. Underwood in Bull. Torrey Club, 33: 199–200. (Apr. 7, 1906.)
"Report on the Continuation of the Botanical Exploration of the Bahama Islands." N. L. Britton in Jour. N. Y. Bot. Gard., 8: 71–81. (Apr., 1907.)
"*Epidendrum cochleatum* L." Geo. V. Nash in Bull. Torrey Club, 34: 117. (May 7, 1907.)

- "Agave Seen by Columbus found." Popular account of the Britton & Millspaugh expedition of 1907. "Discovery," 1: 30-32. (June, 1907.)
- "Oxalidaceae." John K. Small in N. Am. Flora, 25: 43. *Ionoxalis intermedia* (A. Rich.) Small. (Aug. 24, 1907.)
- "Erythroxylaceae." N. L. Britton in N. Am. Flora, 25: 61, 63, 65. *Erythroxylon rotundifolium* Lun., *E. areolatum* L., *E. reticulatum* Northrop, *E. obovatum* Macf. (Aug. 24, 1907.)
- "Linaceae." John K. Small in N. Am. Flora, 25: 72, 74, 75. *Cathartolinum Curtissii*, *C. corallicola*, *C. Bracei*, *C. lignosum*. (Aug. 24, 1907.)
- "Contributions to the Flora of the Bahama Islands—IV." N. L. Britton in Bull. N. Y. Bot. Gard., 5: 311-318. (Oct. 26, 1907.)
- "Phycological Studies—III." Marshall A. Howe in Bull. Torrey Club, 34: 503, 512. *Halimeda simulans*, *Avrainvillea sordida*. (Dec. 17, 1907.)
- "Opuntionidae of North America." N. L. Britton. Britton & Rose in Smithson. Misc. Coll., 50: 513-514. (Feb. 20, 1908.)
- "New Species of Uredineae." J. C. Arthur in Bull. Torrey Club, 34: 587-588. *Prospodium bahamense* sp. nov. (Feb. 27, 1908.)
- "Report on the Botanical Exploration of the Bahama and Caicos Islands." M. A. Howe and Percy Wilson in Jour. N. Y. Bot. Gard., 9: 41-50. (Mar. 1908.)
- "Two Bahamian Species of *Evolvulus*." Homer D. House in Bull. Torrey Club, 35: 89-90. (Mar. 9, 1908.)
- "Studies in North American *Convolvulaceae*." Homer D. House in Bull. Torrey Club, 35: 103. (Apr. 20, 1908.)
- "The Genus *Ernodea* Swartz: A Study of Species and Races." N. L. Britton in Bull. Torrey Club, 35: 204-208. (Apr. 29, 1908.)
- "The North American Species of the Genus *Ipomoea*." Homer D. House in N. Y. Acad. Sci., 18: 216. *Ipomoea carolina*, *Ipomoea gracilis*. (May 11, 1908.)
- "New West Indian *Lejeunea*." Alexander W. Evans in Bull. Torrey Club, 35: 383. *Brachiolejeunea bahamensis*. (Aug. 26, 1908.)
- "Studies in West Indian Plants—II." N. L. Britton in Bull. Torrey Club, 35: 564. *Harrisia Brookii*. (Jan. 2, 1909.)
- "Phycological Studies—IV." Marshall A. Howe in Bull. Torrey Club, 36: 84-99. *Neomeris mucosa*, *N. Cokerii*; *Acetabulum pusillum*, *A. polyphysoides*; *A. p. deltoideum*; *Halimeda lachrymosa*; *Udotea spinulosa*. (Mar. 4, 1909.)

OBSERVATIONS AND NEW SPECIES.

The genus *Dondia*, in so far as it is represented in our Bahamian collections, may be considered as follows:

DONDIA Adans. Fam. 2:261. 1763.

Type species: *Chenopodium altissimum* L. Sp. Pl. 221. 1753.

Stamens as long or longer than the calyx; anthers exserted.

Leaves 3-5 cm. elongated-linear; sepals strongly hooded.

1. **D. linearis.**

Leaves 4-7 mm. narrowly-oblong; sepals simply inflexed.

2. **D. Wilsonii.**

Stamens shorter than the calyx; anthers included.

Leaves 1-3 cm. linear; sepals carinate-hooded.

3. **D. carinata.**

Leaves 2-4 mm. oblong to ovate-spatulate;
sepals simply inflexed.

4. **D. insularis.**

1. **Dondia linearis** (Ell.) Heller Cat. N. A. Pl. 69. 1900.

Salsola linearis Ell. Carol. 1:332. 1821.

Saline borders and maritime rocks. Andros, at Red Bays, *Northrop* 455. Eleuthera, Rock Sound vicinity, *Britton & Millspaugh* 5563:—North Carolina to South Florida and Texas; Cuba.

Referred by Mrs. Northrop to *D. fruticosa* (Forsk.) an Asiatic species.

2. **Dondia Wilsonii** sp. nov.

Annua? ramis ad rhizomam brevam divergentibus 1-3 dm. longis; foliis separatis inapproximatis anguste-oblongis 4-7 mm. longis 0.7-1 mm. crassis ad apicem rotundatis, breve petiolatis vel subsessilis; flores 1-2 in axillam foliorum supremis, 1.5-2 mm. latis; sepalis ovatis, acutis, apice inflexis; stamina in longitudinem prope sepalas, filamentae antheram tres partes longior, cellulae antherarum ad apicem contiguuae. Fructus ignotus.

Margins of salt pans. South Caicos Island, *Percy Wilson* 7616 **Type.**

3. **Dondia carinata** sp. nov.

Herba perennis; ramis ad basim lignosis, erectis, 1.5-2.5 dm longis; foliis approximatis, anguste-linearis, 1.3 cm. longis, 0.8-1 mm. crassis, apiculatis, ad basim attenuatis, amplexicaulibus; inflorescentia fere unica in axilas supremis foliorum bracteacearum depositis, 2-2.5 mm. latis; sepalis scaphoideis, obtusis, apice cucullatis carinatis; stamina sepalas medio longo, cellulae antherarum plus minus didymae filamentae aequalia; semina nigerrima nitida, 1.4 x 1 mm.

Borders of savannas. South Bimini, margin of a mangrove swale, *Millspaugh 2361*. New Providence, near Nassau, *Northrop 150 Type, 194*. Inagua, near Blakeville, *Nash & Taylor 1121*:—Cuba, *C. Wright 2030* in Herb New York Botanical Garden; on coast near the mouth of the Bueyraca, *Britton & Wilson 54*.

Referred by Mrs. Northrop to *D. linearis* (Ell.) Heller.

4. **Dondia insularis** Britton Bull. N. Y. Bot. Gard. 4: 138. 1906.

In a salina. Grand Turk Island, *Nash & Taylor 3873 Type*.

This species has somewhat the habitual appearance of the South Florida *D. conferta* Small, which, however, has larger flowers and filaments longer than the sepals.

The genus *Portulaca*, in so far as it is represented in our Bahamian collections, may be considered as follows:

PORTULACA L. Sp. Pl. 445. 1753.

Type species: *Portulaca oleracea* L. infra.

Leaves flat.

1. **P. oleracea.**

Leaves subterete.

Flowers yellow; seeds brown.

2. **P. phaeosperma.**

Flowers purple; seeds black.

3. **P. gagatosperma.**

1. **Portulaca oleracea** L. Sp. Pl. 445. 1753.

Cultivated soil, waste places, and maritime rocks. Gun Cay, *Millspaugh 2320*. Andros, Cormorant Cay, *Northrop 658*; Road to Morgan's Bluff, *Brace 6675*. Eleuthera, Savanna Sound, *Hitchcock*; Governor's Harbor vicinity, *Britton & Millspaugh 5457, Hitchcock*. Fortune Island, *Brace 4148*. Crooked Island, Marine View Hill, *Brace 4773*. Inagua, *Hitchcock*. Far more common than the actual collections indicate:—All temperate and tropic regions.

2. **Portulaca phaeosperma** Urban, Symb. Ant. 4:233. 1905.

Dry, rocky, and cultivated soil. Great Bahama, West end, *Brace 3529*. Great Sturup Cay, *Britton & Millspaugh 2258*. Andros, Conch Sound, *Northrop 580*; Fresh Creek, *Wight 260*; Mangrove Cay, *Brace 4871, 4920*; Pure Gold, *Brace 5047*. New Providence, Farringdon Road, *Britton & Brace 196*. Eleuthera, Savanna Sound, *Hitchcock*; Governor's Harbor, *Hitchcock*. Great Galiot Cay, *Britton & Millspaugh 2858*. Watling's Island, *Hitchcock*. Long Island, Clarendetown vicinity, *Britton & Millspaugh 6256*. Fortune Island, *Hitchcock*. Acklin's Island, Jumbie Hole, *Brace 4452*. Mariguana, Southeast Point, *Wilson 7566*. Doubtless throughout the Archipelago:—South Florida; Cuba to St. Croix; Jamaica.

Referred by Mrs. Northrop to *P. halimoides*, and by Profs. Hitchcock and Coker to *P. pilosa*.

3. *Portulaca gagatosperma* sp. nov.

Annua suffruticosa diffusa; ramis numerosis nodibus lanato-pilosis; foliis alternis brevis 6–10 mm. longis 1.5–2 mm. latis subcylindricis acuminatis breve petiolatis; inflorescentia terminalia flores ad basim lanatis pedicelis longis, lobae calycis late ovatis apiculatis non carinatis, petala purpureis emarginatis, stylis elongatis longitudae tertia 3–4-partitis; capsula 4 mm. lata infra media circumcissis, semina 0.6 mm. diam. nigerissima rugae stellulatae conjunctae in centro non mamillatae.

In waste soil and the mud of pond borders. Great Bahama, at Eight Mile Rocks, *Britton & Millspaugh 2411*. New Providence, *Brace 368* (Anno 1880); Lake Cunningham, *Britton & Brace 637*, Inagua, *Nash & Taylor 1064* **Type**; *Hitchcock*. Grand Turk Island, *Nash & Taylor 3791*:—Endemic.

Referred by Prof. Hitchcock to *P. pilosa*.

Cassia tora Linn. Sp. Pl., 376. 1753.

The only specimen of this species that I have seen from our region is the one collected by Prof. *Hitchcock* at Governor's Harbor, Eleuthera. This was referred by him to *C. obtusifolia*.

Cassia bicapsularis Linn. Sp. Pl., 376. 1753.

The first return of this species from the Bahamas is Mr. *Wilson's 7739* collected in the vicinity of Kew, North Caicos.

Kallstroemia maxima (Linn.) W. & A. Prodr., 145. 1834.

Rarely seen in the Bahamas. Our only specimens are: *Britton & Brace 784* from waste places at Nassau, New Providence; *Rothrock*, and *Hitchcock* from Fortune Island.

§ EUPHORBIEAE.

As concerns this area the *Euphorbieae* may be considered as follows:

Male and female flowers together in an involucre.

Involucre calceiform, glands internal. 1. *Pedilanthus*.

Involucre campanuliform, glands external.

Glands with petaloid appendages.

Leaves inaequilateral, oblique at the base.

2. *Chamaesyce*.

Glands without petaloid appendages.

Herbs or suffrutices, leafy throughout.

Leaves alternate below, opposite above,

inflorescence solitary or cymose. 3. *Poinsettia*.

Leaves scattered or whorled,
 inflorescence umbelliform.
 Trees, leafy only at tips of branches.
 Shrubs or trees, without true leaves.
 Stems woody, articulate.
 Stems fleshy, ribbed, or tuberculate,
 often with spines or thorns.

4. *Tithymalus*.
5. *Euphorbiodendron*.
6. *Arthrothamnus*.
7. *Euphorbia*.

1. **PEDILANTHUS** Poit. Ann. Mus. Par., 19:390. 1812.

Type species: *Euphorbia tithymaloides* Linn. Sp. Pl., 453. 1753.

1. **Pedilanthus tithymaloides** (Linn.) Poit. Ann. Mus. Par., 19:390. 1812.

Euphorbia tithymaloides Linn. Sp. Pl., 453. 1753.

About dwellings and escaped to coppices. Andros and Cat Islands southward to Grand Turk Island:—South Florida, West Indies, Mexico, Central and South America, Old World Tropics and Sub Tropics.

2. **CHAMAESYCE** S. F. Gray, Nat. Arr. Brit. Pl., 2:260. 1821.

Type species: *Euphorbia peplis* Linn. Sp. Pl., 652. 1753.

Suffrutices, leaves thick, mostly entire.

Leaves ovate.

Inflorescence terminal, clustered.

Plants glabrous.

Plants densely hairy.

Inflorescence axillary, solitary.

Leaves glabrous

entire, stipules deltoid acicular-bristled.

stipules aristate, not bristled.

dentate, at least at the apex.

Leaves canescent, entire.

Leaves linear-oblong, glabrous.

Herbaceous, leaves thin, mostly serrate.

Leaves glabrous.

Inflorescence clustered.

Leaves oblong, acute, sharply serrate.

broadly falcate, seeds red.

narrowly falcate, seeds black.

Inflorescence solitary.

Leaves ovate, blunt, dentate, or entire.

Capsules glabrous.

Capsules hairy on the angles.

Leaves hairy.

Inflorescence clustered.

Leaves ovate-lanceolate, sharply serrate, acute.

1. *buxifolia*.

2. *cayensis*.

3. *Wilsonii*.

4. *lecheoides*.

5. *exumensis*.

6. *Bracei*.

7. *vaginulata*.

8. *hypericifolia*.

9. *brasiliensis*.

10. *Blodgettii*.

11. *prostrata*.

12. *hirta*.

Leaves ovate, crenate-dentate, obtuse. 13. **Berteriana**.
Inflorescence solitary.

Leaves orbicular or rotund, entire or apically dentate.

14. **Brittonii**.

1. **Chamaesyce buxifolia** (Lam.) Small, Fl. S. E. U. S., 712. 1903.

Euphorbia buxifolia Lam. Encyc., 2:421. 1786.

Maritime sands throughout the Archipelago:—Coasts of the Gulf of Mexico and the Caribbean Sea.

2. **Chamaesyce cayensis** (Millsp.). comb. nov.

Euphorbia cayensis Millsp. Torrya, 4:172. 1904.

Coastal whitelands of Joulter's Cays and the Berry Islands:—Endemic.

3. **Chamaesyce Wilsonii** sp. nov.

Glabra purpureo-cinerea diffusoprostratis multo-remosisimis, ramis tenuissimis divaricatis, foliis carnosius ovatis basi oblique auriculocordatis apice obtusis margine integris, stipulis deltoideis blepharociliatis, involucris in dichotomia superiori terminalibusque solitariis cylindro-campanulatis brevissime pedicelatis extus glabris intus dense crispulo-pilosis lobis aristatis glandulis 4 orbiculatis convexis appendices rudimentariis glandulis quintus lobo triangularibus amplior representavit, stylis brevibus stigmata bilobis, capsulae ovatis coccis laeve carinatis, seminibus triangularibus albis angulis acutis faciem laevis.

Plants rosulate prostrate 14–20 cm. in diameter; leaves 2–4 mm. long, 1.5–2.5 mm. broad, seeds 1 mm. x. 7 mm.

Whitelands of Castle Island, *Wilson* 7795, type in herb. Field Museum, sheet no. 221,916:—Endemic.

4. **Chamaesyce lecheoides** (Millsp.). comb. nov.

Euphorbia lecheoides Millsp. Field Mus. Bot., 2:163. 1906.

Scrublands of Inagua and Grand Turk Island:—Endemic.

5. **Chamaesyce exumensis** sp. nov.

Glabra purpureo- aut virido-cinerea, caulibus multo-ramosisimis erectis vel declinato-prostratis denudatis ramisque strictis ad articulos brevissimos nodosis, ramis secundariis subtenuis divaricatis, foliis crassis ovatis petiolatis ad basim obliquo-cordatis ad apicem obtusis vel acutis margine integris saepe remote-dentatis praesertim ad apicem discerni potest, stipulis aristatis, involucris in dichotomia superiori terminalibusque solitariis turbinatis brevi pedicelatis extus glabris intus dense barbatis lobis triangularibus ad apicem setaceis glandulis 4 ovatis planis appendice alba integra glandulis quintus lobo triangularibus amplior representavit, staminibus 5 filamentae ad basim cristato-barbatis stylis brevibus profunde bifidis, capsulae ovatae coccis acute-carinatis, seminibus triangularibus roseo-cinereis farinosis angulis prominens fere rotundis faciem transverse et leve anastomososo-rugosis.

Plants 30 cm. to 1.5 m. tall; leaves 3–10 mm. long, 2–6 mm. broad; seeds 1.1 mm. long, .8 mm. broad. The plants have much the general appearance of *C. cayensis* though the likeness entirely disappears on intimate examination.

Coastal sands and whitelands and sand filled pockets in rocks. Great Exuma, on Stocking Island, *Britton & Millspaugh 3071 type* in herb. Field Museum sheet No. 174,208. Eleuthera, at the southernmost end, *Britton & Millspaugh 5620*. Little San Salvador, *Britton & Millspaugh 5666*. Cat Island, the Bight vicinity, *Britton & Millspaugh 5804*. Ship Channel Cay, *Britton & Millspaugh 2747*. Cay north of Wide Opening, *Britton & Millspaugh 2777, 2779*. Conception Island, *Britton & Millspaugh 5987, 5996*. Watling's Island, Cockburntown vicinity and at Graham's Harbor, *Britton & Millspaugh 6143, 7239*. Rum Cay, at Port Nelson, *Coker 452; Brace 3927*. Long Island, Clarendetown vicinity, *Britton & Millspaugh 6323*. Great Ragged Island, *Wilson 7807*. Crooked Island, Landrail Point, *Brace 4588*. Fortune Island, on the south side, *Brace 4179*.—Endemic.*

Referred to by Prof. Coker as *Euphorbia cayensis* Millsp.

In Praenuncia I, I remarked upon this plant under *Euphorbia flexuosa* Kth. Since then I have had an opportunity of examining the type of Kunth's species and found, as I apprehended, that our species is plainly distinct from his, which, in reality, is a true *Chamaesyce buxifolia*.

6. **Chamaesyce Bracei** (Millsp.) comb. nov.

Euphorbia Bracei Millsp. Field Mus. Bot., 2:159. 1906.

Whitelands of Abaco and Man-o-War Cay:—Endemic.

7. **Chamaesyce vaginulata** (Griseb.) comb. nov.

Euphorbia vaginulata Griseb. Fl. Brit. W. I., 52. 1859.

Rocky whitelands. Inagua, Little Inagua and Grand Turk Island:—Endemic.

8. **Chamaesyce hypericifolia** (Linn.) comb. nov.

Euphorbia hypericifolia Linn. Sp. Pl. 454. 1753.

In open and grassy situations in good soils throughout the archipelago:—Bermuda, the southern United States, West Indies, Mexico to South America. Old World tropics.

9. **Chamaesyce brasiliensis** (Lam.) Small, Fl. S. E. U. S., 712. 1903.

Euphorbia brasiliensis Lam. Dict., 2:423. 1790.

Open places and grassy coverts of New Providence and Eleuthera:—South Florida to Arizona and southward to Brazil; the West Indies.

The most apparent characters separating this species from its near

congeners are: its tenuous branchlets, short-ciliate stipules, and its black seeds with but two ridges transversing the dorsal facets.

10. **Chamaesyce Blodgettii** (Engelm.). Small, *ibid.*

Euphorbia Blodgettii Engelm. Hitchc. in Rep. Mo. Bot. Gard., 4:126. 1893.

On rocks, sands, whitelands, and in pot holes throughout the archipelago from the southern coast of Great Bahama southward:—South Florida, Jamaica, and the Caymans.

11. **Chamaesyce prostrata** (Ait.) Small, *ibid.* 713.

Euphorbia prostrata Ait, Hort. Kew, 2:139. 1789.

Grassy places on New Providence and Grand Turk Island:—Bermuda; North Carolina to Missouri and southward to Brazil and Peru; the West Indies.

12. **Chamaesyce hirta** (Linn.) comb. nov.

Euphorbia hirta Linn. Sp. Pl., 454 ante. 1753

Euphorbia pilulifera Linn. Sp. Pl., 454 post. 1753.

Both the erect and the prostrate form (*Euphorbia pilulifera* var. *procumbens* Boiss.) are frequent in cultivated grounds and along paths from Andros and New Providence to Inagua:—Bermuda; South Florida; the West Indies and Mexico; American and Old World tropics.

The species exists in three forms: the ascending or erect form (*Euphorbia pilulifera* and *hirta* Linn., *E. globulifera* Kunth., *E. capitata* Lam.); the same but with purple-tinged or maculate leaves (*E. pilulifera discolor* Engelm.); and the decumbent or prostrate form (*E. pilulifera procumbens* Boiss., *E. procumbens* D. C., *E. ophthalmica* Pers., *E. oblitterata* Jacq., *E. gemella* Lag.). I cannot concede even varietal rank to these forms, especially as plants exist in this herbarium having *E. pilulifera* and *E. pilulifera procumbens* springing from the same rootstock; and others with *E. pilulifera* and *E. pilulifera discolor* on the same stem. All plants in our region having multicellular amber-colored hairs and ovate, acute, sharply serrate leaves, are the species.

13. **Chamaesyce Berteriana** (Balb.) comb. nov.

Euphorbia Berteriana Balbis, in Spr. Syst., 3:794. 1826.

Redlands of Exuma, Long Island, Atwood's Cay, Mariguana and Castle Island:—Hayti and Porto Rico southward to Guadeloupe and Martinique.

14. **Chamaesyce Brittonii** (Millsp.) comb. nov.

Euphorbia Brittonii Millsp. Field Mus. Bot., 2:159. 1906.

Whitelands of New Providence near Nassau:—Endemic.

3. **POINSETTIA** R. Grah., Edinb. N. Phil. Jour., 20:412. 1836.

Type species: *Euphorbia pulcherrima* Willd. ex Klotzsch in Otto & Dietr. Allg. Gartenz. 2:27. 1834.

1. **Poinsettia heterophylla** (Linn.) Kl. & Gke., Tricocc., 104. 1860.
Euphorbia heterophylla Linn. Sp. Pl., 649. 1753.

Grassy places in good soil, throughout the archipelago:—Bermuda; Illinois to Montana southward to Florida, the West Indies, Mexico, Central America, and South America tropics.

From thorough field examination in a large number of localities I am convinced that the supposed *E. havanensis* Willd. of South Florida and the Bahamas is nothing more than a form of the above species. Not only is this form very frequently found intermixed with the species but the two occur often upon the same plant. There are in this herbarium several sheets that show two forms of this species growing upon the same stem. Of these the Bahamian instances are as follows:

{	<i>Poinsettia havanensis</i> (Willd.)	<i>Euphorbia havanensis</i> Willd. Boiss.,
	and	D. C. Prodr., 15:73 1862.
	<i>Poinsettia graminifolia</i> (Mx.)	<i>Euphorbia graminifolia</i> Mx. Fl. Bor. Am., 2:210. 1803.

Collected at the southeast end of Watling's Island by Mr. Percy Wilson 7317 together with the next:

{	<i>Poinsettia havanensis</i> (Willd.)
	and
{	<i>Poinsettia cyathophora</i> Kl. & Gke., Tricocc., 104. 1859.

Collected by Mr. Wilson with the last and under the same field number; and from Abaco and Great Bahama by Mr. Brace 1503, 3499.

{	<i>Poinsettia heterophylla</i> (Linn.) Kl. & Gke.
	and
{	<i>Poinsettia prunifolia</i> Kl. & Gke. ibid.

Collected on New Providence at Nassau by Mr. Curtiss, 73.

{	<i>Poinsettia heterophylla</i> (Linn.) Kl. & Gke.
	and
{	<i>Poinsettia havanensis</i> (Willd.).

Collected on Eleuthera near Governor's Harbor by Britton & Mills-paugh 5536; this colony also showed a large number of pure *heterophylla* and of pure *havanensis*.

4. **TITHYMALUS** Adans. Fam., 2:355. 1763.

Type species: *Euphorbia Peplus* L. Sp. Pl., 456. 1753.

1. **Tithymalus trichotomus** (Kth.) Kl. & Gke., Tricocc., 81. 1860.

Euphorbia trichotoma (Kth.) in H. B. K. Nov. Gen. et Spec., 2:60. 1817.

Coastal sands on Allen's Cay, Great Bahama and Andros:--South Florida; Cuba, and the Cayman Islands.

5. EUPHORBIODENDRON gen. nov.

Arbores, arbusculæ vel frutices latex ex vulneribus profunderunt. Ramis crassiusculis inferne denudatis cicatricosis superne ad ramis terminalibus foliosis. Folia sparsa, integra sæpe ampla. Cymæ corymbosæ 1-pleurocephalæ ex axillis supremis vel subterminalibus. Involucrum majusculum; glandulis quinis; styli inferne plus minus longe coaliti, longiusculi bifidi, apice rarius incrassati. Semina laevia semper ecarunculata. § *Laurifoliae* Boiss. in D. C. Prodr., 15:105.

1. *Euphorbiodendron gymnonotum* (Urban) comb. nov.

Euphorbia gymnonota Urban, Symb. Antill., 4:396. 1908.

Coppices. Watling's Island, and Fortune Island to Inagua:—Endemic.

Referred to in *Praenuncia*—I as *Euphorbia punicea*; also by Grisebach, Hitchcock, and Coker.

The other species in this genus are:

EUPHORBIODENDRON PUNICEUM (Sw.) comb. nov.

Euphorbia punicea Sw., Prod. Veg. Ind. Occ., 76. 1788.

Poinsettia punicea Kl. & Gke., Tricocc. 102. 1860.

Habitat:—Jamaica.

EUPHORBIODENDRON HELENÆ (Urban) comb. nov.

Euphorbia Helenæ Urban, Symb. Antill., 4:393. 1908.

Euphorbia punicea Griseb., Fl. Br. W. I., 53. 1862.

Habitat:—Cuba.

EUPHORBIODENDRON TROYANUM (Urban) comb. nov.

Euphorbia troyana Urban loc. cit., 394.

Habitat:—Jamaica.

EUPHORBIODENDRON FULVUM (Stapf) comb. nov.

Euphorbia fulva Stapf, Bul. Kew., 294. 1907.

Euphorbia elastica Alt. & Rose, Bul. Inst. Med. Nac. Mex., 1:1905
non Jumelle.

Habitat:—Mexico.

EUPHORBIODENDRON CALYCVLATUM (Kth.) comb. nov.

Euphorbia calyculata Kunth in H. B. K. nov. Gen. et Sp., 2:59. 1817.

Tithymalus calyculatus Kl. & Gke., Tricocc., 81. 1860.

Habitat:—Mexico.

EUPHORBIODENDRON LATAZI (Kth.) comb. nov.

Euphorbia Latazi Kunth, loc. cit., 58.

Habitat:—New Grenada.

EUPHORBIODENDRON CESTRIFOLIUM (Kth.) comb. nov.

Euphorbia cestrifolia Kth. loc. cit., 59.

Habitat:—Peru.

EUPHORBIODENDRON CUBENSIS (Boiss.) comb. nov.

Euphorbia cubensis Boiss. in D. C. Prodr., 15:1265. 1864.

Habitat:—Cuba.

EUPHORBIODENDRON DUSSII (Kr. et Urban) comb. nov.

Euphorbia Dussii Kr. & Urban, in Duss. Fl. Ant. Franc., 44. 1897.

Habitat:—Martinique.

EUPHORBIODENDRON GUDOTI (Boiss.) comb. nov.

Euphorbia Gudoti Boiss., loc. cit., 106. 1862.

Habitat:—New Grenada.

EUPHORBIODENDRON LAURIFOLIUM (Juss.) comb. nov.

Euphorbia laurifolia Juss. in Lam. Dict., 2:418. 1790.

Habitat:—Peru.

EUPHORBIODENDRON SINCLAIRIANUM (Bth.) comb. nov.

Euphorbia Sinclairiana Benth. in Bot. Voy. Sulph., 163. 1844.

Habitat:—Panama region.

6. **ARTHROTHAMNUS** Kl. & Gke. Monatsb. Akad. Berl., 251. 1859.

Type species: *Euphorbia tirucalli* L. Sp. Pl., 452. 1753.

1. **Arthrothamnus cassythoides** (Boiss.) comb. nov.

Euphorbia cassythoides Boiss. Cent. Euph., 20. 1860.

Sandy soil. Andros:—Cuba.

7. **EUPHORBIA** Linn. Sp. Pl., 450. 1753.

Type species: *Euphorbia antiquorum* Linn. Sp. Pl., 450. 1753.

1. **Euphorbia lactea** Haw. Pl. Succ., 127. 1812.

Widely escaped from cultivation and in many places forming dense thickets. New Providence, Cat Island, Watling's, Long Island and Inagua:—West Indies and Mexico. Naturalized from India and the Maluccas.

Referred by Mrs. Northrop and Prof. Hitchcock to *Euphorbia antiquorum*.

Croton Cascarilla.

On account of the doubt that has surrounded this species, as well as the interest in Cascarilla bark economically, it has been the object of special investigation by all of our collectors. No plant agreeing with the plate of Catesby has been found. Wherever we have seen the bark gathered for the market, or for native use, it has been called Sweetwood bark and was taken from the stems and branches of *Croton eluteria* (L.) Sw. (*Clusia eluteria* Linn.). Linnaeus himself says, in his *Flora*

Zeylanica, that Sweet bark and Cascarilla are the same; and in his *Materia Medica* that Cascarilla bark is the *Ricinoides elacagni folio* of Catesby. Catesby says, in the place referred to by Linnaeus (Carolina 2:46), "*An Ricinoides Aleagini folio*. The Ilathera bark. These shrubs grow plentifully on most of the Bahama Islands; seldom above ten Feet high, and rarely so big as a Man's Leg" etc. On page xxxviii he refers to the natives collecting Ilathera bark as a means of support, and to the fact that Cascarilla bark is another common name just as frequently used among the natives.

The remarks of Catesby, his description, and the references of Linnaeus all indicate plainly that he depicts in his plate 46 the plant from which the common product is gathered. This must bear the name *Croton eluteria* (Linn.) Sw. As to the drawing, that has proved so misleading to all authors since its appearance on account of the narrow leaves and long petioles depicted, one need only examine his plates 40 (*Rhus Metopium*) and 42 (*Jacaranda caerulea*) of vol. i, to be assured of his inaccuracy in the matter of proportions.

There remain not the least doubt but that Linnaeus' *Clutia Cascarilla* is synonymous with his *Clutia Eluteria*, and that his *Croton Cascarilla* of the second edition of the *Species Plantarum*, and Bennett's *Croton Cascarilla* of the *Journal of the Linnean Society*, 4:30, 1860, (of which I have had the opportunity to examine the type) are synonymous with the *Croton linearis* of Jacquin.

The synonymy of the two species is as follows:

Croton eluteria (L.) Sw. *Prod. Veg. Ind. Occ.*, 100. 1788.

Clutia eluteria Linn. *Sp. Pl.*, 1042 (ante). 1753.

Clutia Cascarilla Linn. *Sp. Pl.*, 1042 (post). 1753.*

Croton Cascarilla Benn. *Jour. Linn. Soc.*, 4:30. 1860, as to references.

Once common in the Bahamas but becoming quite scarce.

Croton linearis Jacq. *Pl. Carib.*, 32. 1760.

Croton cascarilla Linn. *Sp. Pl.*, ed. 2:1424. 1763. Excluding reference to Catesby.

Croton cascarilla Benn. *Jour. Linn. Soc.*, 4:30. 1860 as to his specimen and the description.

Croton Fergusonii Small, *Fl. Southeastern U.S.*, 695. 1908.

Croton cascarilla var. *linearis* Willd. *Sp. Pl.*, 4: 532. 1805.

A very common coastal species showing a multitude of races in the Bahamas, the extremes of leaf form being:

*Linnaeus draws his description of this species from Catesby's plate 46. vol. ii, and errs in his statement "Habitat in Carolina." Catesby says "Bahama."

Leaves linear 4.5 cm. x 1.5 mm. Eleuthera, *Britton & Millspaugh 5551*.

Leaves linear-oblong 5.5 cm. x 0.8 cm. Watling's Island, *idem 6164*.

Leaves oblong-lanceolate 3.5 cm. x 1 cm.	} On same plant but different branches. Watling's, <i>idem 6177</i> .
Leaves elliptic-oblong 4 cm. x 1.5 cm.	
Leaves obovate 2.2 cm. x 1.3 cm.	

Croton bahamensis sp. nov.

Frutex 1–2 m. alt. sub-di-trichotome ramosus piperi-aromaticus ramis ramulisque teretis infra glabris supra stellato (albo) tomentosus, foliis petiolis limbum 4-plo brevioribus penninervio basi 2-patellari glanduligeris lanceolatis acuminatis mucronatis ad basim breviter oblique-rotundatis, supra viridis ex equae distans stellato-pubescentibus subtus dense stellato-tomentosus, stellae ad centro granularibus, margine subintegris vel crenato-dentatis cum tota sinu stipitato-glandulosis, stipulis fimbriatis stipitato-glandulosis; racemis terminalibus subdensifloris, calycis foem. laciniis oblongis tota stellato-pilosis stylis ad basi 4-fidis monile-villosis stigmae circinnatae, calycis masc. non glanduligeris, petala alba cymbiforma apice minute fimbriata staminibus 35–50; capsulis globosis profunde sulcatis infra glabris supra et in sulcam longe pilosis, seminibus nigro-fuscis coccinelliformis, rugae laeve costaeformae, carunculo gilvo-cereo.

Near *C. humilis*. Leaves 3–7 cm. long, 0.6–1.8 cm. broad; capsules 4 mm. diameter; seeds 3 mm. long, 2 mm. broad.

Open pastures and in thickets bordering openings. Eleuthera, near Rock Sound, *Britton & Millspaugh 5574 type* (4 sheets of same in herb. Field Museum Nos. 198402–3–4–5). Cat Island, Port Howe vicinity, *Britton & Millspaugh, 5954, Hitchcock, Rothrock 566*. Conception Island, *Britton & Millspaugh 6021*. Watling's Island, Graham's Harbor to Columbus' Monument, *Britton & Millspaugh 6176*, Southeast End, *Wilson 7330*. Mariguana, Abraham Bay vicinity, *Wilson 7486*. Rum Cay, Road to Sclaters, *Brace 3979*. Long Island, old pasture near Clarendetown, *Britton & Millspaugh 6240*, and in a thicket nearby, *6246*. Fortune Island, on Hanna Hill, *Brace 4064*, and Cove Road to the southside *4165*. Acklin's Island, at Spring Point, *Brace 4293*.

Referred to in Praenuncia I and by Prof. Hitchcock as *C. humilis*.

Centaurium Brittonii Millspaugh & Greenman, sp. nov.

Herba annua pusilla erecta 5–15 cm. alta glaberrima a basi plerumque ramosissima; caulibus quadrangularibus et plus minusve anguste alatis

subdichotomo-ramosis; foliis oppositis sessilibus, primis oblongo-spathulatis, ceteris oblongo-lanceolatis vel linearis 3–12 mm. longis 15mm. latis plerumque acutis integris; floribus longe pedicellatis tetrameris, pedicellis gracilis usque ad 2 cm. longis; calyce circiter 5 mm. longo, laciniis lineari-attenuatis tubo multo longioribus; corolla alba 6–10 mm. longa, lobis elliptico-oblongis 4–4.5 mm. longis retusis vel irregulariter et minute dentatis; stigmatibus bilamellatis; capsula elliptico-lanceolata circiter 8 mm. longa e calyce persistente exsertis; semina subsphaerica foveolato-reticulata.

A low, much-branched annual having much the aspect of *CENTAUURIUM DIVARICATUM* (Schaffner) Millsp. & Greenm. comb. nov. (*Erythraea divaricata* Schaffn. ex. Schlecht. Bot. Zeit., 1855, p. 920), but readily distinguished by its profuse subdichotomous branching and small white flowers; from *C. tetramerum* (Schiede) Eastw., a species of similar habit and tetramerous flowers, *C. Brittonii* differs in having white flowers with retuse or minutely dentate corolla-lobes and smaller capsules.

Sandy roadsides and rocky scrublands. Watling's Island, north-west of Cockburntown, March 15, 1907, *Britton & Millspaugh 6224* (hb. Field Mus. cat. No. 198651) **Type.** Great Exuma, near Georgetown, Feb. 22, 1905, *Britton & Millspaugh 3125*. Eleuthera, vicinity of Eleuthera Point, Feb. 24, 1907, *Britton & Millspaugh 5632*.

Heliotropium Nashii sp. nov.

Caulibus suffruticosis prostratis diffusis ramosissimis glabris; foliis sessilis cum pila griseo strigosis laxè vestitis oppositis ascendentibus ovatis acuminatis 1.5–2 mm. longis 0.8–1 mm. latis internodiis aequalibus margine integris non revolutis; inflorescentia in axillis supremis solitariis, flores sessilis vel subsessilis, sepalae ovatis acuminatis inaequalis vix a foliis recens discerni potest, corollae albae 2 mm. longae lobis ovatis acuminatis inaequalibus patentibus tubum tres partes brevior; staminibus sessilis ad tubo corollae supramedium coalitis; stylus ovarium semel brevior crassis erectis, stigma incrassata peltata 4-lobata apice producto mamillata; drupa ovata 1 mm. diam. plane sulcatim in carpellae quatuor, seminae cuneiformae facies duo planis tertius (dorsalis) convexas.

Scrublands. Inagua, near Mathewtown, *Nash & Taylor, 1011* **Type;** *Hitchcock, Dec. 3, 1890*.

Referred by Prof. Hitchcock to *H. microphyllum* Sw.

The genus *Varronia*, in so far as it is represented in our Bahamian collections, may be considered as follows:

VARRONIA P. Br. Hist. Jam., 172. 1756.

Type species: *Lantana corymbosa* L. Sp. Pl., 628. 1753.

Flowers in globose heads.

Calyx teeth filiform; leaves coarsely dentate. 1. **V. globosa.**

Calyx teeth deltoid; leaves entire or few toothed. 2. **V. bahamensis.**

Flowers in spikes.

Leaves linear-oblong; filaments pilose at the base. 3. *V. Brittonii*.
Leaves spatulate-obovate; filaments not pilose. 4. *V. lucayana*.

1. *Varronia globosa* Jacq. Enum. Pl. Carib., 14. 1760.

Cordia globosa H. B. K. Nov. Gen. et Sp., 3: 76. 1818.

Scrublands and old fields. Andros, at Kemp Bay, *Brace* 5028. Watling's at the north end, *Britton & Millspaugh* 6208. Long Island, near Clarendon, *Britton & Millspaugh* 6223, *Coker* 501:—Florida Keys: Cuba to Porto Rico: Jamaica; Grand Cayman. Also credited to the French Antilles, Venezuela, and Panama.

2. *Varronia bahamensis* (Urban), comb. nov.

Cordia bahamensis Urban Symb. Ant., 1: 392. 1899.

Coppices, scrublands, pinelands, and whitelands. Man-o-War Cay, *Brace* 1586. Abaco, at Marsh Harbor, *Brace* 1601. Great Bahama, at Barnett's Point, *Britton & Millspaugh* 2645; at Eight Mile Rocks, *Brace* 3733, and at West End, *Brace* 3536. Andros, at Nichol's Town, *Northrop* 376, *Brace* 6719, 6842; along road to Conch Sound, *Brace* 6822; Mastic Point vicinity, *Brace* 7095. New Providence, at Waterloo, *E. G. Britton* 6620; at Fox Hills, *Wight* 194; near Carmichael, *Eggers* 4193; near Nassau, *Hitchcock*, *Coker* 77, 106, *Curtiss* 78; Blue Hills. *Millspaugh* 2075; at South Beach, *Britton* 47, *Millspaugh* 2119; along Fox Hills road, *Britton & Brace* 373. Eleuthera, along path from Harbor Island to Spanish Wells and from Harbor Island to the Bluff, *E. G. Britton* 6494, 6516; along the path from the Glass Window to Harbor Island, *Britton & Millspaugh* 5384, 5401; Governor's Harbor vicinity, *Britton & Millspaugh* 5456, *Hitchcock*; Rock Sound vicinity, *Britton & Millspaugh* 5580. Conception Island, *Britton & Millspaugh* 5993. Cat Island, Orange Creek vicinity, *Britton & Millspaugh* 5761; at Port Howe, *Hitchcock*; at the southwest end, *Wilson* 7199. Watling's, Cockburntown vicinity, *Britton & Millspaugh* 6058, *Wilson* 7209; at the southeast end, *Wilson* 7264; *Hitchcock*. Rum Cay, near Port Nelson, *Coker* 443, *Brace* 3972. Great Guana Cay, *Britton & Millspaugh* 2893. Great Exuma, near Georgetown, *Britton & Millspaugh* 2924. Long Island, Clarendon vicinity, *Britton & Millspaugh* 6296. Crooked Island, at Landrail Point, *Brace* 4648. Fortune Island, *Hitchcock*, *Brace* 4038. Mariguana, Abraham Bay vicinity, *Wilson* 7498. Inagua, *Hitchcock*:—Endemic.

The species includes a large number of races the extremes represented in the collections being: *Brace's* 3536, from the west end of Great Bahama, with leaves 1.5–2.5 cm. long and 0.4–0.7 cm. wide and on young shoots 5.5–6.5 cm. long and 1.5–2 cm. broad; and *Britton &*

Millspaugh 6058, from Watling's near Cockburntown, with leaves 4.4–5 cm. long and 2–3 cm. broad and on young shoots 7–8 cm. long by 3.5–4 cm. broad.

The species is referred by Mrs. Northrop to *Cordia lima* R. & S., and by Prof. Hitchcock to *Cordia globosa* H. B. K. and *Cordia* sp.

3. **Varronia Brittonii** sp. nov.

Ramis teretibus tenuis glabris junioribus pulverentis; foliis 1.5–3 cm. longis 2.5–4 mm. latis, linearo-oblongis ad basim cuneatis ad apicem truncatis vel rotundatis raro acutis infrequens crenato-dentatis, pagina minute scabris et resino punctatis; pedunculis terminalibus spicis aequalibus vel brevioribus; calyce pulverentis corollam duplo brevioribus, limbus 4–5 dentatis dentibus deltoideus, corollae 5-lobatis lobis deltoideis vel triangulo-ovatis inaequaliter et diversiter repando-dentatis; filamentae ad tubo corollam infra medium coalitis ad basim pilosis antheram longioribus, antherae in corollam inclusit; drupa 2–2.5 mm. longis in calyce persistens plane inclusis.

The species differs from *V. bahamensis* in its spicate inflorescence, leaf form, size and texture and all its minor characters. It differs from *V. angustifolia* Desv. (*Cordia angustifolia* R. & S.) which has [type seen] narrowly lanceolate, acute, petiolate leaves, papillose-scabrous above, woolly beneath and dentate throughout the margin.

Coppices and scrublands. Great Bahama, at Eight Mile Rocks, *Britton & Millspaugh 2481* Type; *Brace 3710*. Andros at Fresh Creek, *Northrop 619*; at Deep Creek, *Brace 5210*. Eleuthera, at Governor's Harbor, *Britton & Millspaugh 5426*; at Gregorytown, *Coker 378*. Cat Island, Orange Creek vicinity, *Britton & Millspaugh 5737*. Long Island, near Clarendon, *Britton & Millspaugh 6306*; *Coker 496*.—Cuba, *C. Wright 3114* in Herb. Torrey, New York Botanical Garden. This sheet has two plants attached thereto, one being this species, the other *V. leptoclada* (Urban & Britton) (*Cordia leptoclada* Urban & Britton in Urban Symb., 5:478).

The collections show a race with somewhat narrower leaves than those of the type.

Referred by Prof. Coker to *Cordia cylindristachya* R. & S., and by Mrs. Northrop to *Cordia angustifolia* R. & S.

4. **Varronia lucayana** sp. nov.

Frutices humilis diffusis 1–1.8 m. altus, ramis teretibus glabris cicutrici foliorum delapsorum valide scabris, ramules junior nonnihil ferrugineo-pulverentis; foliis 0.7–2 cm. longis 0.4–0.8 cm. latis obovato-spathulatis ad basim coarctatis, apice truncatis inaequaliter repando-dentatis, utrinque pagina puberulo-scabris et resino-punctatis, petiolae 1–1.5 mm. longis ferrugineo-pulverentis; inflorescentia spicata ramulorum junior terminalia, spicae 1–1.5 cm. longa pauciflora pedunc-

ulæ 0.5–0.8 cm. longa; calyce corollam circa tres partes brevioribus, limbus 5-dentatis dentae inaequalis obtuso-triangularis: corollae glabra alba, limbus inaequaliter 5-lobatus, lobis ovatis crenato-dentatis; filamentae ad tubo corollam supra medium coalitis ad basim non pilosis antheram brevioribus, antherae in corollam inclusit; drupa 2.5–3 mm. longa in calyce persistens semisepultus.

Rocky plains. Acklin's Island, at Spring Point, *Brace* 4245. Mariguana, near Abraham Bay and at South East Point, *Wilson* 7533, 7586. South Caicos Island, *Wilson* 7672. Inagua, on James' Hill, *Nash & Taylor* 1175. **Type**:—Endemic.

Psychotria undata Jacq. Hort. Schoenb., 3: 5. 1798.

We had several opportunities, on the north end of Watling's Island, to compare living specimens of this species with *P. bahamensis* Millsp.** The leaves of *P. undata* were, in every case, strongly wavy throughout their length, while those of *P. bahamensis* were always plane. *P. undata* is a much larger shrub and has deeper green leaves.

Catesbaea foliosa sp. nov.

§ *Erectiflorae*. Frutex aculeatus vel eaculeatus ramis validis cortice laxis albogriseis ramulis dense et minute resino-setulosis, foliis crassis fuscoviridis lineari-spathulatis vel obovatis sub-sessilis, apice obtusis raro mucronulatis margine integris revolutis supra vernicosis, flores pedicellatis, calycis dentibus subulatis obtusis explicatis minute setulosis, bacca alba globosa, seminibus rubidis ovatis planis utrinque facies ad centro umbonatis.

A thick branched, spreading shrub 1 to 2 m. high; leaves 0.8 to 1 cm. long, 2 to 6 mm. broad; fruit 3 mm. in diameter, the persistent sepals 1 mm. long; seeds 1.7 mm. long, 1.5 mm. broad.

Scrublands. Mariguana, Abraham Bay vicinity, *Wilson* 7507; five miles west of Southeast Point, *idem* 7555. Crooked Island, road to Stopper Hill, *Brace* 4785; *Hitchcock*.* Fortune Island, Forsyth Road, *Brace* 4210, Acklin's Island, at Abercorn, *idem* 4473. Inagua, Miner's Tent to Balsam Hill, *Nash & Taylor* 1272.* Grand Turk Island, *idem* 3771.* Caicos Islands, on West Caicos, *Wilson* 7761 **type**, in herb. Field Museum sheet No. 221880.

Referred to by Prof. Hitchcock as *C. parviflora*.

Callicarpa Hitchcockii sp. nov.

Frutex scandens ramis divaricatis pallido-gilvis ramulis cum cymis dense furfuraceo-ferrugineo-tomentosis, foliis bicoloribus crassis anguste oblanceolatis ad basim attenuatis ad apicem obtusis valide petiolatis, margine revolutis integris supranitidis ad nerviam profunde sulcatis infra prominente nervatis dense stellato-lanatis; cymis supra-

* Referred, in Praenuncia—I, to *C. campanulata*. ** Bull. N. Y. Bot. Gard., 3: 451. 1905.

axillaribus corymbosis submultifloris pedunculo petiolum bis superante folio triplo brevioribus, calyce glabris resino-punctatis obscuriter vel nequaquam dentatis; bacca glabra minute rugosa et resino-punctata.

A climbing shrub 2–3 m. high with a strong odor of fenugreek. Leaves 2–2.5 cm. long, 5–7 mm. broad, petioles 4 mm. long.

It differs strongly from *C. fulva* Rich. the type of which has ovate-lanceolate acuminate sharply dentate leaves up to 10 cm. long and 3.5 cm. wide (fide Briquet); from *C. ferruginea* Sw. which has lanceolate membranous leaves also dentate or dentate-serrate; and from *C. cubensis* Urban which has ovate or elliptic-ovate leaves woolly beneath (not stellate) 3 to 6 cm. long, 1.5 to 3.5 cm. broad and are subtruncate or rotund at the base.

Shrubbery edge of a rocky plain back of the settlement of Port Howe, Cat Island, *Britton & Millspaugh 5946*; The Bight, on the edge of a coppice back of The Point, *ibid. 5819*; and in the edge of a coppice over the hill East of the last station, *ibid. 5913* **type**. Prof. Hitchcock's specimens were collected at or near the first station mentioned above, in October, 1890. Mr. Brace also returns the species from a savanna on the west side of Andros Island across from Mastic Point (7100), and from a pine barren near Mastic Point itself (6965):—Endemic.

Since I included this species under *C. fulva* Rich. in *Praenuncia*–I, I have had the kind assistance of Professors C. de Candolle, Le Compte and Briquet in establishing its status through comparisons of our material in their herbaria, Prof. Briquet has the type of Richard's *C. fulva* in Linden's Cuban 2066 and has favored me with a sketch of the same which satisfies me of my error in first considering our material to be that species. Prof. Robinson, Gray Herbarium, Cambridge, Mass., has also kindly loaned this Museum the Wright sheets of *Callicarpa*, from Cuba, from each of which our species is distinct.

***Solanum didymacanthum* Millsp. Field Mus. Bot., 2: 183. 1909.**

After persistent search Dr. Britton succeeded in finding this species, both in flower and ripe fruit, on Columbus' Point, near Port Howe, Cat Island, Prof. Hitchcock's and Prof. Rothrock's original station. The doubtful mark may now be removed from line 12 of the original description reading: "corolla alba (?)" and the words "Bacca ignota" are to be replaced by: Fructae sessilae globosae aurantiaceae translucentae basi et apice saepe compressis sepalis persistente non longiore.

***Cestrum pallidum* Lam. Encycl., 1: 688. 1783.**

One of Prof. Hitchcock's Cat Island sheets, recently secured by this Museum, is this species. The two sheets in the herbarium of the Missouri Botanical Garden are, however, *C. bahamense* Britton.

NATIVE PLANT NAMES.*

Above all	<i>Tecoma bahamensis</i>
Abraham bush	<i>Phyllanthus epiphyllanthus</i>
Almond	<i>Catappa Catappa</i>
Bamboo	<i>Agave bahamana</i>
Bastard Torch	<i>Ocotea Catesbyana</i>
Bay Bean	<i>Canavalia</i> sps.
Bay Cedar	<i>Suriana maritima</i>
Bay Geranium	<i>Ambrosia hispida</i>
Bay Hops	<i>Ipomoea pes-caprae</i>
Bay Lavender	<i>Tournefortia gnaphaloides</i>
Bay Marigold	<i>Borrichia arborescens</i>
Bay Rush**	<i>Zamia</i> sps. <i>Echites umbellata</i>
Bay top	<i>Coccothrinax argentea</i>
Bay Wormwood	<i>Croton linearis</i>
Bean, bay	<i>Canavalia</i> sps.
Beefwood	<i>Torrubia longifolia</i>
Benny	<i>Sesamum indicum</i>
Black Buttonwood	<i>Conocarpus erecta</i>
Black Ebony	<i>Pera bumeliaefolia</i>
Black-eyed Susan	<i>Abrus precatorius</i>
Black Mangrove	<i>Avicennia nitida</i>
Black Torch	<i>Erithalis fruticosa</i> , <i>Amyris elemifera</i>
Blolly	<i>Torrubia longifolia</i>
Boar-hog bush	<i>Callicarpa Hitchcockiana</i>
Bow pidgeon	<i>Coccoloba Krugii</i>
Boxwood	<i>Jacaranda caerulea</i>
Brier	<i>Anthacanthus acicularis</i>
Brier, China	<i>Smilax bahamensis</i>
Broom brush	<i>Evolvulus arbuscula</i>
Buffalo top	<i>Thrinax microcarpa</i>
Bullrush	<i>Uniola spicata</i>
Bull's wood	<i>Hypelate trifoliata</i>
Butter bough	<i>Exothea paniculata</i>
Buttercup	<i>Turnera ulmifolia</i>
Buttonwood	<i>Conocarpus erecta</i>

*Only such names as have been given to us personally, by natives while confronting the plant indicated, are included in this list.

**This appellation belongs properly to *Zamia* sps. The only locality in which it is applied to *Echites*, so far as we know, is on Cat Island where a starch, similar to that made from *Zamias*, is manufactured from its tubers.

Buttonwood, black	<i>Conocarpus erecta</i>
Buttonwood, white	<i>Conocarpus sericea</i>
Candlewood	<i>Dodonea Ehrenbergii</i> , <i>Phialanthus myrtilloides</i>
Cane, wild	<i>Panicum divaricatum</i>
Canker berry	<i>Solanum bahamense</i>
Cascarilla bark	<i>Croton linearis</i>
Cassava wood	<i>Dipholis salicifolia</i>
Cathartic bark	<i>Thevetia Thevetia</i>
Cat's paw	<i>Solanum didymacanthum</i>
Cedar, bay	<i>Suriana maritima</i>
Chew stick	<i>Gouania domingensis</i>
China	<i>Smilax Beyrichii</i>
China brier	<i>Smilax bahamensis</i>
Christmas vine *	<i>Ipomoea antillana</i>
Cinnamon	<i>Pimenta Pimenta</i>
Cinnamon bark	<i>Canella Winterana</i>
Cinnecord	<i>Acacia choriophylla</i>
Coco Plum	<i>Chrysobalanus</i> sps.
Coffee, wild	<i>Psychotria</i> sps.
Cow bush	<i>Helicteres spiralis</i>
Crabwood	<i>Coccoloba Krugii</i> , † <i>Gymnanthes lucida</i>
Darling Plum	<i>Reynosia septentrionalis</i>
Day lily	<i>Hymenocallis caribaea</i>
Devil's Pumpkin	<i>Passiflora cuprea</i>
Dildo	<i>Pilocereus</i> and <i>Cephalocereus</i> sps.
Dogwood	<i>Ichthyomethia piscipula</i>
Ebony, black	<i>Pera bumeliaefolia</i>
Elemi, gum	<i>Terebinthus Simaruba</i>
Feather bed	<i>Maba caribaea</i>
Frogwood ‡	<i>Guettarda Krugii</i>
Gardena	<i>Cakile aequalis</i>
Geranium, bay	<i>Ambrosia hispida</i>
Goldenrod	<i>Lantana bahamensis</i>
Granny bush	<i>Croton linearis</i>
Grape, sea	<i>Coccoloba uvifera</i>
Guava, wild	<i>Eugenia bahamensis</i>
Gum Elemi	<i>Terebinthus Simaruba</i>
Gunwood	<i>Tecoma bahamensis</i>

*Used in decorating churches at Christmas.

† "Crab, he like eat um berries, much."

‡ "Frog, he like climb um up."

Hard head	<i>Phyllanthus epiphyllanthus</i>
Haulback	<i>Mimosa bahamensis</i>
Hippo	<i>Asclepias curassavica</i>
Hog bush	<i>Rhachicallis maritima</i>
Hog Palmetto	<i>Pseudophoenix Sargentii</i> .
Honeysuckle	<i>Diapedium assurgens</i>
Hops, bay	<i>Ipomoea pes-caprae</i>
Horse bush	<i>Gundlachia domingensis</i>
Horseflesh	<i>Lysiloma Sabicu</i>
Horseradish tree	<i>Moringa Moringa</i>
Hurricane grass	<i>Fimbristylis glomerata</i>
Jackmadá	<i>Eupatorium villosum</i>
Jerusalem Parsley	<i>Chenopodium spathulatum</i>
Jimbay	<i>Lucaena glauca</i>
Joe wood	<i>Jacquinia keyensis</i>
Lavender, bay	<i>Tournefortia gnaphaloides</i>
Lightwood	<i>Lasiocroton</i>
Lignum vitae	<i>Guaicum officinale</i>
Lime, wild	<i>Fagara Fagara</i>
Link vine	<i>Vanilla</i> sps.
Logwood	<i>Haematoxylon campechianum</i>
Love vine	<i>Cuscuta</i> sps.
Madeira	<i>Swietenia Mahogani</i>
Mahogany	<i>Swietenia Mahogani</i>
Manchineel	<i>Hippomane mancinella</i> , <i>Excoecaria lucida</i>
Mangrove	<i>Rhizophora Mangle</i>
Mangrove, black	<i>Avicennia nitida</i>
Mangrove, white	<i>Laguncularia racemosa</i>
Marigold	<i>Stemmodontia bahamensis</i>
Marigold, bay	<i>Borrichia arborescens</i>
Mastic	<i>Sideroxylon foetidissimum</i>
Milk bush	<i>Euphorbia cayensis</i>
Mistletoe	<i>Dendropemon</i> sps.
Mosquito bush	<i>Cassia angustisiliqua</i>
Moujean tea	<i>Lantana balsamifera</i>
Nickers	<i>Guilandina</i> sps.
Old Man's beard	<i>Dendropogon usneoides</i>
Olive	<i>Picrodendron macrocarpum</i>
Pain-in-back	<i>Trema Lamarckiana</i>
Palmetto, hog	<i>Pseudophoenix Sargentii</i>
Parsley, Jerusalem	<i>Chenopodium spathulatum</i>

Pea, pidgeon	Cajanus Cajan
Pepper bush	Croton bahamensis
Pigeon pea	Cajanus Cajan
Pigeon plum	Coccoloba and Chrysobalanus sps.
Plum, coco	Chrysobalanus sps.
Plum, darling	Reynosia septentrionalis
Plum, pigeon	Coccoloba and Chrysobalanus sps.
Poison bush	Hippomane mancinella
Poison wood	Metopium Metopium
Pond top	Sabal Palmetto
Pork bush	Cakile aequalis
Pork-and-doughboy *	Bumelia loranthifolia, Acacia acuífera
Prickly apple	Catesbaea spinosa
Princewood	Exostemma caribaeum
Quinine	Ammocallis rosea
Ram's horn	Pithecolobium keyense
Sage, wild	Lantana sps.
Sapodilla, wild	Mimusops sps.
Satinwood	Fagara flava
Sea Grape	Coccoloba uvifera
Seven year Apple	Genipa clusiaefolia
Shanks	Salmea pterobioides
Shepherd's needle	Bidens leucantha
Silver top	Coccothrinax argentea
Slag	Typha domingensis
Snakeroot	Picramnia pentandra
Stopper	Eugenia axillaris
Stopper, white	Calyptanthus pallens
Strong-back	Bourreria havanensis
Sweet potato	Ipomoea batatas
Sweet William	Ammocallis rosea
Sweetwood bark	Croton eleuteria
Tea, Moujean	Lantana balsamifera
Thyme, wild	Rhachicallis maritima
Torch, bastard	Ocotea Catesbyana
Torch, black	Erithalis fruticosa, Amyris elemifera
Turk's head	Melocactus sps.
What-o'clock †	Jacaranda caerulea
White Buttonwood	Conocarpus sericea

* The thorns are used in lieu of forks in eating pork and dumplings.

† Our guide could not explain reference.

White Mangrove	<i>Laguncularia racemosa</i>
White Stopper	<i>Calyptranthes pallens</i>
Whitewood	<i>Drypetes keyensis</i>
Wild Cane	<i>Panicum divaricatum</i>
Wild Coffee	<i>Psychotria</i> sps.
Wild Guava	<i>Eugenia bahamensis</i>
Wild Lime	<i>Fagara Fagara</i>
Wild Sage	<i>Lantana</i> sps.
Wild Sapodilla	<i>Mimusops</i> sps.
Wild Thyme	<i>Rhachicallis maritima</i>
Woe vine	<i>Cassytha americana</i>
Woman's tongue*	<i>Lysiloma Sabicu</i>
Wormwood, bay	<i>Croton linearis</i>

* Said to allude to the noise made by the pods in a high wind.

INDEX.

PRÆNUNCIÆ BAHAMENSES I & II.*

- Abena jamaicensis* 179.
Acetabulum polyphysoides 296.
 polyphysoides deltoideum 296.
 pusillum 295.
Achyranthes aspera obtusifolia 147.
 indica 147.
Acalypha alopecuroides 156.
 ostreaefolia 156.
 setosa 156.
Adelia Bernardia 155.
Alernanthera maritima 147.
 muscioides 147.
 paronychioides 147.
 repens 147.
Amaranthus crassipes 146.
 emarginatus 146.
 hybridus 146.
 paniculatus 146.
 polygonoides 146.
 spinosus 146.
 tristis 146.
 viridis 147.
Argythamnia argentea 154.
 candicans 154.
 candicans 155.
 lucayana 154.
 sericea 155.
Arthrothamnus cassythoides 306.
Avicennia nitida 183.
Avrainvillea sordida 296.
Bernardia Bernardia 155.
 carpinifolia 155.
 dichotoma 155.
 mexicana 155.
Bonania cubana 157.
 emarginata 157.
Bouchea Ehrenbergii 178.
 prismatica 178.
Brachioleunea bahamensis 296.
Callicarpa fulva 180.
 Hitchcockii 312.
 lancifolia 181.
Cariolus arenicolor 295.
 membranaceus 295.
 pinisitus 295.
Cassia bicapsularis 299.
 tora 299.
Catesbaea campanulata 167.
 fasciculata 167.
 foliosa 312.
 parviflora 167, 312.
 parviflora septentrionalis 167.
 spinosa 167.
Cathartolinum Bracei 296.
 corallicola 296.
 Curtissii 296.
 lignosum 296.
Centaurium Brittonii 308.
 divaricatum 309.
Cestrum pallidum 313.
Chamaesyce Berteriana 303.
 Blodgettii 303.
 Bracei 302.
 brasiliensis 302.
 Brittonii 303.
 buxifolia 301.
 cayensis 301.
 exumensis 301.
 hirta 303.
 hypericifolia 302.
 lecheoides 301.
 prostrata 303.
 vaginulata 302.
 Wilsonii 301.
Chenopodium altissimum 297.
Chiococca alba 170.
 parvifolia 170.
 pinetorum 171.
 racemosa 170.
Citharexylum bahamense 180.
 Berterii 179.
Citharexylum caudatum 179.
 lucidum 179.
 quadrangulare 179.
 villosum 180.
Clerodendron aculaetum 182.
Cordia angustifolia 311.
 bahamensis 310.
 cylindristachya 311.
 globosa 310.
 leptoclada 311.
 lima 311.
Croton bahamensis 308.
 Cascarilla 152, 306.
 Cascarilla 153.
 Cascarilla linearis 153.
 discolor 152.
 eluteria 152, 307.
 Fergusonii 307.
 flocculosus 152.
 Hjalmarssonii 153.
 homolepidus 152.
 humilis 153.
 humilis 308.
 linearis 153, 306.
 lobatus 153.
 lucidus 154.
 niveus 152.
 rosmarinifolius 154.
Dondia carinata 297.
 fruticosa 297.
 insularis 298.
 linearis 297.
 linearis 298.
 Wilsonii 297.
Drypetes diversifolia 151.
 keyensis 151.
 lateriflora 152.
Duranta Plumieri 180.
 repens 180.
Epidendron cochleatum 295.
Erithalis fruticosa 109.
 odorifera 169.

*New species and combinations in **black face** type.

- Erythraea divaricata* 309.
Erythroxylon areolatum 296.
 obovatum 296.
 reticulatum 296.
 rotundifolium 296.
Euphorbia antiquorum 163.
 306.
 Berteriana 158.
 Berteriana 303.
 Blodgettii 158.
 Blodgettii 303.
 Bracei 159.
 Bracei 302.
 brasilensis 159.
 brasilensis 302.
 Brittonii 159.
 Brittonii 303.
 buxifolia 160.
 buxifolia 301.
 buxifolia flexuosa 161.
 calyculata 305.
 cassythoides 161.
 cassythoides 306.
 cayensis 161.
 cayensis 162, 301.
 cestrifolia 305.
 cubensis 306.
 Dussii 306.
 elastica 305.
 flexuosa 161.
 flexuosa 302.
 fulva 305.
 gemella 303.
 geniculata 162.
 graminifolia 304.
 Gudoti 306.
 gymnonota 305.
 hirta 303.
 havanensis 162.
 havanensis 304.
 Helenae 305.
 heterophylla 162.
 heterophylla 304.
 heterophylla graminifolia
 162.
 hypericifolia 163.
 hypericifolia 302.
 lactea 163, 306.
 Latazi 305.
 laurifolia 306.
 lecheoides 163.
 lecheoides 301.
 obliterata 164.
 obliterata 303.
 ophthalmica 303.
 peplis 300.
 Peplus 304.
 pilulifera 164.
 pilulifera 303.
 pilulifera discolor 303.
 pilulifera oblitterata 164.
Euphorbia pilulifera procum-
 bens 164, 303.
 procumbens 303.
 prostrata 164.
 prostrata 303.
 prunifolia 162.
 pulcherrima 304.
 punicea 164.
 punicea 305.
 Sinclairiana 306.
 tirucalli 306.
 tithymaloides 300.
 trichotoma 164.
 trichotoma 304.
 troyana 305.
 vaginulata 165.
 vaginulata 302.
EUPHORBIOIDENDRON 305.
 calyculatum 305.
 cestrifolium 305.
 cubensis 306.
 Dussii 306.
 fulvum 305.
 Gudoti 306.
 gymnonotum 305.
 Helenae 305.
 Latazi 305.
 laurifolium 306.
 puniceum 305.
 Sinclairianum 306.
 troyanum 305.
Exostemma caribaeum 165.
Excoecaria Sagraei 157.
Galium hispidulum 174.
Genipa clusiaefolia 166.
Ghinia curassavica 174.
 verbenacea 174.
Guettarda elliptica 168.
 Krugii 168.
 scabra 168.
Gymnanthes lucida 157.
Halimeda lachrymosa 296.
 simulans 296.
Hamelia patens 167.
Harrisia Brookii 296.
Heliotropium microphyllum
 309.
 Nashii 309.
Hemidiodia ocimifolia 173.
Hippomane Mancinella 157.
Hura crepitans 158.
Ionoxalis intermedia 296.
Ipomoea carolina 296.
 gracilis 296.
Iresine celosioides 148, 149.
 inaguensis 149.
 keyensis 148.
 paniculata 148.
Ischnia verbenacea 174.
Jatropha curcas 156.
 gossypifolia 157.
Kallstroemia maxima 299.
Lantana bahamensis 174.
 balsamifera 175.
 camara 175.
 camara 174.
 corymbosa 175.
 crocea 175.
 demutata 175.
 involutrata 175.
 odorata 176.
 ovatifolia 176.
Lasiocroton macrophyllus 156.
Laugeria densiflora 169.
Linum Curtissii 295.
Lippia canescens 177.
 nodiflora 177.
 nodiflora 178.
 reptans 177.
 stoechadifolia 178.
Lithophila muscoides 147.
 vermicularis 147.
Manihot Apii 157.
 Manihot 157.
Mercurialis annua 155.
Morinda Roioe 173.
Myristicophyllum ligustrifolium
 172.
 pubescens 172.
 undatum 172.
NASHIA 176.
 inaguensis 177.
Neomeris Cokerii 296.
 mucosa 296.
Ovieda fragrans 183.
Pedilanthus angustifolius 165.
 tithymaloides 165, 300.
Pera bumelifolia 156.
 bumeliaefolia parvifolia
 156.
Pettitia domingensis 181.
 Poeppigii 181.
Phialanthus myrtilloides 170.
Philoxerus vernicularis 148.
Phyllanthus angustifolius 149.
 bahamensis 150.
 distichus 150.
 epiphyllanthus 150.
 niruri 150.
 radicans 151.
 saxicola 151.
Poinsettia cyathophora 304.
 graminifolia 304.
 heterophylla 304.
 prunifolia 304.
 punicea 305.
Portulaca gagatosperma 299.
 halimoides 299.
 oleracea 298.
 phacosperma 298.
 pilosa 299.
Priva echinata 179.

- Priva lappulacea 179.
 Prosopodium bahamense 296.
PSEUDOCARPIDIUM 181.
 avicennioides 182.
 ilicifolium 182.
 Wrightii 182.
 Psychotria bahamensis 172.
 lanceolata 172.
 ligustrifolia 172.
 pubescens 172.
 undata 173, 312.
 Rachicallis *americana* 165.
 maritima 165.
 rupestris 165.
 Randia aculeata 166.
 Relbunium hypocarpum 174.
 Ricinus communis 156.
 Salsola *linearis* 297.
 Savia bahamensis 149.
 Scolosanthus bahamensis 171.
 Scutellaria longiflora 295.
 Securinea acidothermus 149.
 Sesia striata 295.
 Solanum *didymacanthum* 183,
 313.
 Spermacoce aspera 173.
 tenuior 174.
 Strumpfia maritima 171.
 Tectaria minima 205.
 Tithymalus *calyculatus* 305.
 trichotomus 304.
 Udotea spinulosa 296.
 Valerianodes *fruticosa* 178.
 jamaicensis 178.
 Varronia *bahamensis* 310.
 Brittonii 311.
 globosa 310.
 leptoclada 311.
 lucayana 311,
 Verbena *curassavica* 174.
 Vitex *avicennioides* 182.
 ilicifolia 182.

QK1.F42:11

SCIII

3 5002 00143 1159

Millspaugh, Charles Frederick
Contributions to North American Euphorbi

QK1.F42:1-2

SCIII

3 5002 00144 8799

Millspaugh, Charles Frederick
Plantae Utowanae. Plants collected in Be

QK1.F42:4

SCIII

3 5002 00143 0920

Greenman, Jesse More
Studies in the genus Citharexylum.

WELLESLEY COLLEGE LIBRARY

3 5002 03631 6292

QK1.F42:5

SCIII

3 5002 00244 1702

Millspaugh, Charles Frederick
Flora of the sand keys of Florida, by Ch

QK1.F4 2:10

SCIII

3 5002 00405 2200

Millspaugh, Charles Frederick
Contributions to North American Euphorbi

QK1.F42:3

SCIII

3 5002 00405 2481

Millspaugh, Charles Frederick
Praenunciae bahamenses— I. Contribution

QK1.F42:6

SCIII

3 5002 00405 0345

Greenman, Jesse More
New or noteworthy spermatophytes from Me

QK1.F42:7

SCIII

3 5002 00405 2507

Millspaugh, Charles Frederick
Praenunciae Bahamenses— II. Contributio

QK1.F42:8

SCIII

3 5002 00405 0139

Greenman, Jesse More
II. Diagnoses of new species and notes o

QK1.F42:9

SCIII

3 5002 00405 2572

Millspaugh, Charles Frederick
II. Two new stonecrops from Guatemala.