Historic, Archive Document

Do not assume content reflects current
scientific knowledge, policies, or practices.

az SD76
A1484

United States
Department of
Agriculture

National
Agricultural
Library

Beltsville,
Maryland 20705

and

United States
Environmental
Protection Agency

Office of Pesticide
Programs

Bibliographies
and Literature
of Agriculture
Number 96

September 1990

ct

The Protection of Small Grains, Other Than Wheat, Rice, or Sorghums, 1979 - April 1990

Citations from AGRICOLA Concerning Diseases and other Environmental Considerations

United States
Department of
Agriculture

National
Agricultural
Library

Beltsville,
Maryland 20705

and

United States
Environmental
Protection Agency

Office of Pesticide
Programs

Bibliographies
and Literature
of Agriculture
Number 96

September 1990

The Protection of Small Grains, Other Than Wheat, Rice, or Sorghums, 1979 - April 1990

Citations from AGRICOLA Concerning Diseases and other Environmental Considerations

Compiled and Edited by

Charles N. Bebee
National Agricultural Library

National Agricultural Library
Beltsville, Maryland
1990

National Agricultural Library Cataloging Record:

Beebe, Charles N.

The protection of small grains, other than wheat, rice, or sorghums,
1979-April 1990 : citations from AGRICOLA concerning diseases and
other environmental considerations.

(Bibliographies and literature of agriculture ; no. 96)

1. Grain -- Diseases and pests -- Bibliography. 2. Plants, Protection of --
Bibliography. I. Title.

aZ5076.A1U54 no.96

FOREWORD

This is the 34th volume in a series of commodity-oriented environmental bibliographies resulting from a memorandum of understanding between the U.S. Department of Agriculture, National Agricultural Library (USDA-NAL), and the U.S. Environmental Protection Agency, Office of Pesticide Programs (EPA-OPP).

This close working relationship between the two agencies will produce a series of bibliographies which will be useful to EPA in the regulation of pesticides, as well as to any researcher in the field of plant or commodity protection. The broad scope of information contained in this series will benefit USDA, EPA, and the agricultural community as a whole.

The sources referenced in these bibliographies include the majority of the latest available information from U.S. publications involving commodity protection throughout the growing and processing stages for each agricultural commodity.

We welcome the opportunity to join this cooperative effort between USDA and EPA in support of the national agricultural community.

JOSEPH H. HOWARD, Director
National Agricultural Library

DOUGLAS D. CAMPT, Director
Office of Pesticide Programs

INTRODUCTION

The citations in this bibliography, The Protection of Small Grains Other Than Wheat, Rice, or Sorghums, are selected from the AGRICOLA (AGRICultural OnLine Access) database limited to those produced by North American authors. They cover articles or monographic publications added to the database from 1979 - April 1990.

This is the 34th bibliography in a series of commodity-oriented listings of citations from AGRICOLA jointly sponsored by the U.S. Department of Agriculture, National Agricultural Library (USDA-NAL), and the U.S. Environmental Protection Agency, Office of Pesticide Programs (EPA-OPP). Additional volumes issued recently include The Protection of Cotton, 1985 - 1989, The Protection of Soybeans, 1985 - 1989, The Protection of Small Fruits and Berries, The Protection of Grapes and Cherries, The Protection of Ornamental Plants, The Protection of Farm Animals, and The Protection of Wildlife and Vertebrate Pest Control. The 1990 volumes include The Protection of Tropical and Subtropical Fruits, The Protection of Small Grains (other than Wheat, Rice or Sorghums), The Protection of Cucurbits, The Protection of Minor Vegetable Crops, The Protection of Beans, Peas, and Lentils, and The Protection of Forestry.

Entries in the bibliography are subdivided into a series of section headings used in the contents of the Bibliography of Agriculture. Each item appears under every section heading assigned to the cited document. A personal author index is also included in the publication and a site index to plants follows the personal author index.

The U.S. Environmental Protection Agency contact for this project is Richard B. Peacock, Office of Pesticides and Toxic Substances.

Any comments or questions concerning this bibliography may be addressed to the compiler and editor:

Charles N. Bebee
Special Services Branch
USDA-NAL, Room 1402
Beltsville, MD 20705
(301) 344-3875

EPA BIBLIOGRAPHY

The Protection of Small Grains, 1979 - April 1990

(Other Than Wheat, Rice, Sorghums)

Contents

	<u>Item</u> <u>Number</u>
Research	1 - 2
Meteorology and Climatology	3 - 6
History	7 - 8
Legislation	9 - 25
Farm Organization and Management	26 - 30
Distribution and Marketing	31 - 32
Plant Production - General	33 - 36
Plant Production - Horticultural Crops	37 - 41
Plant Production - Field Crops	42 - 173
Plant Production - Range	174 - 176
Plant Breeding	177 - 418
Plant Ecology	419 - 430
Plant Structure	431 - 449
Plant Nutrition	450 - 493
Plant Physiology and Biochemistry	494 - 820
Plant Taxonomy and Geography	821 - 827
Protection of Plants	828 - 837
Pests of Plants - General and Misc.	838 - 847
Pests of Plants - Insects	848 - 941
Pests of Plants - Nematodes	942 - 950
Plant Diseases - General	951 - 974
Plant Diseases - Fungal	975 - 1220
Plant Diseases - Bacterial	1221 - 1234
Plant Diseases - Viral	1235 - 1259
Plant Diseases - Physiological	1260 - 1271
Miscellaneous Plant Disorders	1272 - 1370
Protection of Plant Products - General and Misc.	1371 - 1385
Protection of Plant Products - Insects	1386 - 1430
Weeds	1431 - 1611
Pesticides - General	1612 - 1689
Soil Biology	1690 - 1693
Soil Chemistry and Physics	1694 - 1706
Soil Fertility - Fertilizers	1707 - 1748
Soil Cultivation	1749 - 1775
Soil Erosion and Reclamation	1776 - 1779
Forestry Related	1780
Entomology Related	1781 - 1796
Animal Ecology	1797 - 1798
Animal Nutrition	1799 - 1802
Animal Taxonomy and Geography	1803 - 1804

Pests of Animals - General and Misc.	1805
Animal Diseases - Fungal	1806
Animal Disorders - Physical Trauma	1807
Aquaculture Related	1808 - 1809
Agricultural Engineering	1810
Farm Equipment	1811 - 1812
Drainage and Irrigation	1813 - 1817
Food Science, Field Crop	1818 - 1819
Food Processing	1820
Food Storage, Field Crop	1821 - 1826
Food Contamination and Toxicology	1827 - 1831
Food Contamination, Field Crop	1832 - 1842
Food Contamination, Horticultural Crop	1843
Food Composition, Dairy	1844
Food Composition, Field Crop	1845 - 1851
Feed Processing and Storage	1852
Feed Contamination and Toxicology	1853 - 1857
Feed Composition	1858 - 1860
Physiology of Human Nutrition	1861
Pollution	1862 - 1870
Mathematics and Statistics	1871 - 1878
Documentation	1879
Human Medicine, Health, and Safety	1880

Index

Page

Author Index	227 - 241
Site Index	242 - 245

EPA BIBLIOGRAPHY

RESEARCH

0001

Determination of chlorsulfuron residues in grain, straw, and green plants of cereals by high-performance liquid chromatography.

Slates, R.V. JAFCA. Washington : American Chemical Society. Journal of agricultural and food chemistry. Jan/Feb 1983. v. 31 (1). p. 113-117. ill. 1 p. ref. (NAL Call No.: 381 J8223).

0002

Shortcut for bluetongue research.

AGREA. Senft, D. Washington, D.C. : The Administration. Agricultural research - U.S. Department of Agriculture, Agricultural Research Service. Mar 1989. v. 37 (3). p. 22. (NAL Call No.: DNAL 1.98 AG84).

METEOROLOGY AND CLIMATOLOGY

0003

Effect of planting dates on spring barley varieties, San Juan Branch Station, 1974-1979 (New Mexico, performance, yield).

Gregory, E.J. Arnold, R.N. Las Cruces : The Station. Research report - New Mexico, Agricultural Experiment Station. June 1982. June 1982. (478). 12 p. 8 ref. (NAL Call No.: 100 N465R).

0004

The effects of preharvest rain.

Derera, N.F. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 1-14. ill., maps. Includes references. (NAL Call No.: DNAL SB189.6.P73).

0005

The modification of the foliage structure and radiation absorption in a plant canopy by wind.

Hipps, L.E. Baiden, E. Boston : The Society, 1985. 17th Conference on Agricultural and Forest Meteorology and seventh Conference on Biometeorology and Aerobiology, May 21-24, 1985, Scottsdale, Ariz. : preprint volume / sponsored by the American Meteorological Society. p. 127-128. Includes references. (NAL Call No.: DNAL S600.2.C6 1985).

0006

The potential effect of windbreaks, barriers and residues on the development of small grain plants.

Freckleton, M.E. Bozeman, Mont. : Montana State University, Cooperative Extension Service. Great Plains Agriculture i.e. Agricultural Council publication. Paper presented at the "International Symposium on Windbreak Technology," June 23-27, 1986, Lincoln, Nebraska. 1986. (117). p. 139. (NAL Call No.: DNAL S27.A3).

HISTORY

0007

Field crops.

Arny, D.C. Dubuque, Iowa : Kendall/Hunt Pub. Co., c1986. With one foot in the furrow : a history of the first seventy-five years of the Department of Plant Pathology at the University of Wisconsin-Madison / edited by Paul H. Williams, Melissa Marosy. p. 164-174. ill. (NAL Call No.: DNAL SB732.54.U6W5).

0008

History of benzimidazole use and resistance.

Smith, C.M. St. Paul, Minn. : ARS Press, American Phytopathological Society, 1988. Fungicide resistance in North America / Charles J. Delp, editor. p. 22-24. (NAL Call No.: DNAL SB951.F88).

LEGISLATION

0009

Method for screening bacteria and application thereof for field control of Pythium spp. on small grain crops.

Weller, D.M. Cook, R.J.; Becker, J.O.
Washington, D.C.? : The Department. Abstract:
A method for screening bacteria to select strains which will suppress Pythium spp. in small grain crops under field conditions and a method for applying field-suppressive bacteria to suppress Pythium spp. in a commercial setting are described. Four Pseudomonas strains which passed the screen test are disclosed. United States Department of Agriculture patents. Copies of USDA patents are available for a fee from the Commissioner of Patents and Trademarks, U.S. Patents and Trademarks Office, Washington, D.C. 20231. Mar 3, 1987. (4,647,533). 1 p. ill. Includes references. (NAL Call No.: DNAL aT223.V4A4).

0010

Registration of Arizona 8501 barley germplasm for disturbed land reclamation.

CRPSAY. Day, A.D. Ludeke, K.L.; Ottman, M.J.
Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 387. Includes references. (NAL Call No.: DNAL 64.8 C883).

0011

Registration of barley yellow dwarf virus tolerant barley composite cross XLIV germplasm.

CRPSAY. Crosslin, J.M. Carroll, T.W.; Hockett, E.A.; Zaske, S.K. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 387-388. Includes references. (NAL Call No.: DNAL 64.8 C883).

0012

Registration of 'Florida 401' rye.

CRPSAY. Pfahler, P.L. Barnett, R.D.; Luke, H.H. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 836. Includes references. (NAL Call No.: DNAL 64.8 C883).

0013

Registration of four pairs of greenbug-resistant vs. susceptible near-isolines of winter barley germplasms.

CRPSAY. Carver, B.F. Morgan, G.H.; Edwards, L.H.; Webster, J.A. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1988. v. 28 (6). p. 1034-1035. Includes references. (NAL Call No.: DNAL 64.8 C883).

0014

Registration of 'Heartland' barley.

CRPSAY. Therrien, M.C. Irvine, R.B.; Campbell, K.W.; Wolfe, R.I. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1985. v. 25 (6). p. 1124. (NAL Call No.: DNAL 64.8 C883).

0015

Registration of 'Hercules' oat.

CRPSAY. Marshall, H.G. Kolb, F.L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1988. v. 28 (4). p. 719. Includes references. (NAL Call No.: DNAL 64.8 C883).

0016

Registration of 'Hitchcock' barley.

CRPSAY. Schmidt, J.W. Dreier, A.F.; Dofing, S.M. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1985. v. 25 (6). p. 1123. (NAL Call No.: DNAL 64.8 C883).

0017

Registration of 'Kelly' oat.

CRPSAY. Reeves, D.L. Hall, L. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 385-386. (NAL Call No.: DNAL 64.8 C883).

0018

Registration of 'Lamont' barley.

CRPSAY. Wesenberg, D.M. Robbins, G.S. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 373. Includes references. (NAL Call No.: DNAL 64.8 C883).

0019

Registration of 'Multiline E76' and 'Multiline E77' oats.

CRPSAY. Frey, K.J. Browning, J.A.; Simons, M.D. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1985. v. 25 (6). p. 1125. Includes 1 references. (NAL Call No.: DNAL 64.8 C883).

0020

Registration of 'Samson' barley.

CRPSAY. Helm, J.H. Dyson, D.H.; Stewart, W.M. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 384. (NAL Call No.: DNAL 64.8 C883).

0021

Registration of 'Simpson' oat.

CRPSAY. Graham, W.D. Jr. Morton, B.C. Jr.; Kingsland, G.C.; Gambrell, R.H. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 386. (NAL Call No.: DNAL 64.8 C883).

0022

Registration of 'Venus' barley.

CRPSAY. Brown, A.R. Johnson, J.W.; Rothrock, C.S.; Bruckner, P.L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1988. v. 28 (4). p. 718-719. Includes references. (NAL Call No.: DNAL 64.8 C883).

0023

Registration of 'Virden' barley.

CRPSAY. Therrien, M.C. Irvine, R.B.; Campbell, K.W.; Wolfe, R.I. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 374. Includes references. (NAL Call No.: DNAL 64.8 C883).

0024

Registration of 'Webster' oat.

CRPSAY. Frey, K.J. Simons, M.D.; Michel, L.J.; Murphy, J.P.; Browning, J.A. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 374-375. Includes references. (NAL Call No.: DNAL 64.8 C883).

0025

Registration of Webster oat isolines as parental lines.

CRPSAY. Frey, K.J. Simons, M.D.; Michel, L.J.; Murphy, J.P.; Browning, J.A. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 386-387. Includes references. (NAL Call No.: DNAL 64.8 C883).

FARM ORGANIZATION AND MANAGEMENT

0026

734-739. Includes references. (NAL Call No.: DNAL 79.8 W41).

The effect of ryegrass and rye seeding rate on forage yield planted alone and in mixtures in south Alabama.

Kee, D.D. Duffy, P.D.; Ward, C.Y. Lexington, Ky. : The Conference. Proceedings of the Forage and Grassland Conference. 1988. p. 221-225. Includes references. (NAL Call No.: DNAL SB193.F59).

0027

Effects of pasture management systems on cow-calf productivity on loessial soils in Northeast Louisiana.

LAXBA. Coombs, D.F. Bartleson, J.L.; Rogers, R.L.; Saxton, A.M.; Huffman, D.C.; Alison, M.W. Baton Rouge, La. : The Station. Bulletin - Louisiana Agricultural Experiment Station. Oct 1989. (815). 19 p. Includes references. (NAL Call No.: DNAL 100 L93 (1)).

0028

Evaluation of three wild oat herbicides in combination with broadleaf herbicides for the control of weeds in spring barley and the economic value of the crop.

WSWPA. Stewart, V.R. Keener, T.K. Reno, Nev. : The Society. Proceedings - Western Society of Weed Science. 1987. v. 40. p. 104-105. (NAL Call No.: DNAL 79.9 W52).

0029

A management tool for small grain producers...the wild oat staging card (Weeds, herbicide application timing, Montana).

Dyer, W. Fay, P.; Rardon, P.; Stewart, V. Bozeman : The Station. Capsule information series - Montana Agricultural Experiment Station. June 1983. June 1983. (30). 5 p. ill. (NAL Call No.: S83.M6).

0030

Volunteer barley (*Hordeum vulgare*) interference in canola (*Brassica campestris* and *B. napus*).

WEESA6. O'Donovan, J.T. Sharma, A.K.; Kirkland, K.J.; De St Remy, E.A. Champaign, Ill. : Weed Science Society of America. The yield potential and the effect on yield loss of canola of different densities of volunteer barley were investigated at three locations in western Canada. Field studies were conducted from 1982 to 1986. Rectangular hyperbolic models based on data pooled over years, locations, and canola cultivars, and incorporating different densities of volunteer barley and canola accurately portrayed field responses in most instances. Results indicated that volunteer barley severely reduced canola yield. However, financial losses due to reduced canola yield were partly offset by the volunteer barley crop. Weed science. Nov 1988. v. 36 (6). p.

DISTRIBUTION AND MARKETING

0031

In-transit fumigation of truck-ship containers with hydrogen phosphide--a feasibility study (Insect control, stored product pests, cereals).

Jay, E. Davis, R.; Zehner, J.M. New Orleans : The Region. Advances in agricultural technology. AAT-S - United States, Dept. of Agriculture, Agricultural Research Service, Southern Region. Apr 1983. Apr 1983. (28). 13 p. Includes references. (NAL Call No.: aS21.A75U7).

0032

Why the world doesn't want our grain.

McDonald, D. Philadelphia : The Journal. Farm journal. Oct 1985. v. 109 (12). p. 13-17. ill. (NAL Call No.: DNAL 6 F2212).

PLANT PRODUCTION – GENERAL

0033

Effect of Basic-H on vegetable and agronomic crops and soil fertility at Pt. MacKenzie.
AGBOBD. Laughlin, W.M. Smith, G.R.; Peters, M.A. Fairbanks : The Station. Agroborealis - Alaska Agricultural and Forestry Experiment Station, University of Alaska-Fairbanks. July 1987. v. 19 (1). p. 31-33. Includes references. (NAL Call No.: DNAL S33.E2).

0034

The effects of phosphate on early growth and maturity.

AGJDAT. Noll, C.F. Madison, Wis. : American Society of Agronomy. A number of investigators have found that the use of phosphates produced a more rapid growth of the roots of seedlings and several have claimed that, in the case of small grains, phosphates promote tillering. Russell states that in England superphosphates cause rapid early growth of turnips and swedes. The evidence is not conclusive that these effects from phosphate are more pronounced than from the other fertilizer elements where the latter are the limiting factors for crop yields. Fertilizer tests have quite generally, if not always, shown that the use of phosphatic fertilizers induces earlier ripening of the grain crops on soils low in phosphorus. Similar effects have been noted with cabbage and with cotton. In the case of tomatoes, the use of phosphates at the Pennsylvania Experiment Station, has been accompanied by a much greater growth of stalk and larger total yield, but also by later ripening. The effects of different phosphatic fertilizers seem to vary with the availability of the phosphates as shown by the responses in crop yields. However, at the Ohio Experiment Station, acid phosphate had more influence in promoting earliness than either steamed bonemeal or basic slag, altho all these phosphates gave approximately the same yield. Applied in moderate rates, the soluble phosphates, as a rule, have shown a more pronounced influence in hastening maturity than the same quantity of phosphoric acid in rock phosphate, the influence in earliness varying in degree with the response in yield. Increasing the rates of application of phosphatic fertilizers above the needs of the crop as indicated by yields has not been accompanied by further increases in earliness. Though phosphates show a more marked influence in promoting earliness than the other fertilizer elements, yet moderate amounts of nitrogen and potash, where these are needed for the production of crops, have a similar effect. Lime, also, if applied to soils having a high lime requirement, may induce earli. Agronomy journal. Mar 1923. v. 15 (3). p. 87-99. (NAL Call No.: DNAL 4 AM34P).

0035

Requirements and developments in small grain seed treatments.

PLDRA. Phipps, P.M. St. Paul, Minn. : American Phytopathological Society. Plant disease. Nov 1985. v. 69 (11). p. 1009-1010. (NAL Call No.: DNAL 1.9 P69P).

0036

Weed control by means of chemical sprays.

AGJDAT. Bolley, H.L. Madison, Wis. : American Society of Agronomy. Agronomy journal. 1910. v. 1. p. 159-168. (NAL Call No.: DNAL 4 AM34P).

PLANT PRODUCTION - HORTICULTURAL CROPS

0037

The effects of using copper for mitigating histosol subsidence on: 1. The yield and nutrition of oats and lettuce grown on histosols, mineral sublayers, and their mixtures.

Levesque, M.P. S.D.S.C.A. Mathur, S.P. Baltimore : Williams & Wilkins. Soil science. Feb 1983. v. 135 (2). p. 88-100. Includes references. (NAL Call No.: 56.8 S03).

0038

Postemergence control of an oat cover crop and broadleaf weeds in direct-seeded nursery beds.
HJHSA. Warmund, M.R. Alexandria, Va. : American Society for Horticultural Science. HortScience. Aug 1987. v. 22 (4). p. 603-605. Includes references. (NAL Call No.: DNAL SB1.H6).

0039

Root-lesion nematodes, potassium deficiency, and prior cover crops as factors in apple replant disease.
JOSH.B. Merwin, I.A. Stiles, W.C. Alexandria, Va. : The Society. Growth chamber evaluations of soil from an orchard replant site showed severe stunting of 'Northern Spy' apple (*Malus domestica* Borkh.) seedlings grown in field soil (FS) compared with pasteurized soil (PS) from the same site. The FS : PS seedling dry weight ratio of 0.44 indicated a serious replant problem. Leaf nutrient content was generally higher in PS than FS seedlings. Multiple-regression analysis indicated that leaf K and root-lesion nematode (*Pratylenchus penetrans* Filipjev) primary inoculum accounted for 75% of the variation in FS seedling dry weight. Apple seedling dry weight in FS was 97% of that in PS following a marigold (*Tagetes patula* L. cv. Sparky) cover crop, and 75% following oats (*Avena sativa* L. cv. Saia). Root-lesion nematodes were nearly eliminated from the plots with marigold. Other cover crops and weed-free fallow period were less effective in controlling apple replant disease and/or phytonematodes. Journal of the American Society for Horticultural Science. Sept 1989. v. 114 (5). p. 728-732. Includes references. (NAL Call No.: DNAL 81 S012).

0040

Seasonal establishment of bermudagrass using plastic and straw mulches.
AGJOAT. Sowers, R.S. Welterlen, M.S. Madison, Wis. : American Society of Agronomy. Bermudagrass *Cynodon dactylon* (L.) Pers. is normally established vegetatively during the early summer in the transition zone, to allow sufficient establishment time before the onset of freezing conditions in the fall. Clear polyethylene covers and straw mulches cause changes in the turfgrass microenvironment that may influence the rate of sprig establishment. The objective of this study was to evaluate the

effects of barley (*Hordeum vulgare* L.) straw and clear polyethylene plastic mulch (0.04-mm thickness) on the establishment of 'Midiron', 'Tufcote', and 'Vamont' bermudagrass from sprigs. Separate tests were conducted in the fall of 1983 and 1984 and the summer of 1984 and 1985. Summer spriggings were made in May, June, and July, months generally recommended for planting in the transition zone. Late-season spriggings were made in August, September, and October. Plantings were made on a Sassafrass sandy loam (fine-loamy, siliceous, mesic Typic Hapludult). Summer bermudagrass establishment was reduced under straw and polyethylene covers. Injury under plastic occurred to plantings made in May and June, which were exposed to mean soil temperatures above 41 degrees C during the 8-wk cover period. In contrast, plastic mulch stimulated early fall bermudagrass growth and delayed dormancy. August and September plantings remaining under plastic throughout the winter exhibited high winter survival in comparison to unmulched or straw mulched turf, and were nearly 100% established by 1 July of the following year. Spriggings under plastic planted after 20 September exhibited poor establishment by 1 July the following year. These studies showed that plastic covers can be used to extend the establishment season of bermudagrass into the fall; however, plastic covers are detrimental to summer bermudagrass establishment. Agronomy journal. Jan/Feb 1988. v. 80 (1). p. 144-148. Includes references. (NAL Call No.: DNAL 4 AM34P).

0041

Treatment of currants and cherries to prevent spot diseases / L.H. Pammel, G.W. Carver. Squirrel-tail grass or wild barley, (*Hordeum jubatum* L.) / by L.H. Pammel. The chemical composition of Squirrel-tail grass / J.B. Weems, W.H. Heileman.
Pammel, L. H. 1862-1931. Carver, George Washington, 1864?-1943.; Weems, J. B.; Heileman, W. H. Ames, Iowa : Experiment Station. Iowa Agricultural College, 1895. Caption title. p. 289-321 : ill. ; 23 cm. Includes bibliographical references. (NAL Call No.: DNAL 100 Io9 no.30).

PLANT PRODUCTION - FIELD CROPS

0042

Andre spring barley (Two-rowed, long-auned variety, performance, disease resistance).
Morrison, K.J. Ullrich, S.E.; Nilan, R.A.; Brauen, S.E.; Muir, C.E.; Reisenauer, P. Pullman, Wash. : The Service. Extension Bulletin - Washington State University, Cooperative Extension Service. Mar 1984. Mar 1984. (1249). 5 p. (NAL Call No.: 275.29 W27P).

0043

Annual Grass control in flax (Linum usitatissimum) with quizalofop.
WETEE9. Friesen, G.H. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Apr 1988. v. 2 (2). p. 144-146. Includes references. (NAL Call No.: DNAL SB610.W39).

0044

Availability studies upon high potash nitrate.
AGJOAT. Allison, R.V. Madison, Wis. : American Society of Agronomy. Experimental studies upon the availability of the nitrogen and potash of High Potash Nitrate have been carried out under both green-house and field conditions. The results in pot culture indicated the value of the nitrogen and potash of this compound as being fully equivalent to that derived from sodium nitrate and potassium sulfate. Because of the fact that the field plots responded strongly to applications of nitrogen and not at all to potash, it may be assumed that the nitrogen of the High Potash Nitrate was as effective as that of sodium nitrate. Agronomy journal. Jan 1924. v. 16 (1). p. 26-30. (NAL Call No.: DNAL 4 AM34P).

0045

Barley.
AGRYA. Foster, E. Prentice, N. Madison, Wis. : American Society of Agronomy. Agronomy. 1987. (28). p. 337-396. ill. Includes references. (NAL Call No.: DNAL 4 AM392).

0046

Barley (Hordeum vulgare) response to herbicides applied at three growth stages.
WETEE9. Martin, D.A. Miller, S.D.; Alley, H.P. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Jan 1988. v. 2 (1). p. 41-45. Includes references. (NAL Call No.: DNAL SB610.W39).

0047

Barley production on Alabama farms.
Mask, P.L. Hagan, A.K. Auburn, Ala. : The Service. Circular ANR - Alabama Cooperative Extension Service, Auburn University. In Subseries: Agriculture and Natural Resources. Agronomy. Oct 1988. (517). 2 p. (NAL Call No.: DNAL S544.3.A2C47).

0048

Barley variety performance in Michigan.
MUCBA. Copeland, L.O. Freed, R.D.; Leep, R.H.; Wolfe, D. East Lansing, Mich. : The Service. Extension bulletin E - Cooperative Extension Service, Michigan State University. Apr 1986. (1313,rev.). 2 p. (NAL Call No.: DNAL 275.29 M58B).

0049

Barley vs. oat companion crops. I. Forage yield and quality response during alfalfa establishment.
CRPSAY. Brink, G.E. Marten, G.C. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 1060-1067. Includes references. (NAL Call No.: DNAL 64.8 C883).

0050

Bowers barley.
Leep, R. H. Copeland, L. O.; Grafius, J. E. & A new spring barley for feed. Document available from: Michigan State Univ., Bulletin Office, P.O. Box 231, East Lansing, Michigan 48824 1979. This discusses the history of Bowers barley, it's time and rate of seeding, seed quality and treatment, weed control, fertilization, and harvesting. 1 sheet : ill. (NAL Call No.: Document available from source.). (NAL Call No.: Extension Bulletin E-1314).

0051

Breeding winter oats for the South.
AGJOAT. Stanton, T.R. Madison, Wis. : American Society of Agronomy. A nursery for the isolation and development of cold-resistant, high-yielding, and otherwise desirable strains of winter oats has been maintained at the Arlington Experiment Farm since 1907. A more or less standardized nursery system is used consisting of series of single 5-foot plant or head rows, duplicated 15-foot rows, and duplicated blocks or plats of three 15-foot rows each. Two strains of Winter Turf, C. I. Nos. 435-4 and 541-4, have been developed which during the 10-year period, 1916-25, inclusive, have outyielded the unselected stock by about 8 bushels to the acre. Definite winter types of Fulghum have been isolated and are being studied. In these forms the plants show a

(PLANT PRODUCTION - FIELD CROPS)

decided spreading or winter habit in early growth and have dark green, rather narrow leaves. The selections apparently are more cold-resistant, and are superior in yield to the original Fulghum. Several high-yielding strains of winter oats have been developed from a Winter Turf-Aurora case. In these strains the earliness and excellent kernel characters of the Aurora variety have been combined with the winter resistance of the Winter Turf oat to a highly satisfactory degree. Two of the most promising selections have been named Lee and Custis. The maintenance of varietal identity in such standard oats as fulghum and Red Rustproof is one of the most pressing practical problems in connection with the growing of oats in the South. Selected strains of these varieties are available of these varieties are available and should be substituted for the nondescript, so-called Red Rustproof oats which are grown in many districts of the South. The selection work with Fulghum was considerably expanded in the fall of 1925 at the Arlington Experiment Farm. In addition, 500 selections from Fulghum were sown at Tifton, Ga., for the development of strains especially adapted to the southern portion of the cotton belt. Numerous selections from hybrids between Fulghum and hardy winter varieties, such as Culberson. *Agronomy journal*. Sept 1926. v. 18 (9). p. 804-814. ill. (NAL Call No.: DNAL 4 AM34P).

0052

Broadleaf weed control in barley with sulfonyl urea herbicides.
WAEBA. Miller, S.D. Krall, J.M. Laramie : The Station. Bulletin B - Wyoming, Agricultural Experiment Station. 1986. (885). p. 150-151. (NAL Call No.: DNAL 100 W99 (1)).

0053

The Cereal rusts /edited by William R. Bushnell and Alan P. Roelfs. --.
Bushnell, William R.; Roelfs, A. P. Orlando : Academic Press, c1984-1985. 2 v. : ill. ; 24 cm. Includes bibliographies and indexes. (NAL Call No.: DNAL SE741.R8C44).

0054

Cereal vegetation as a source of forage.
AGRYA. Wedin, W.F. Hoveland, C. Madison, Wis. : American Society of Agronomy. *Agronomy*. 1987. (28). p. 83-99. Includes references. (NAL Call No.: DNAL 4 AM392).

0055

The comparative attractiveness of various small grains to the chinch bug /by Curtis Benton and W.P. Flint.
Benton, Curtis, 1898-. Flint, W. P. 1882-. Washington, D.C. : U.S. Dept. of Agriculture, 1938. Caption title.~ Contribution from Bureau

of Entomology and Plant Quarantine in cooperation with the Illinois Agricultural Experiment Station. 8 p. ; 23 cm. (NAL Call No.: DNAL 1 Ag84C no.508).

0056

Comparative winter hardiness of barley varieties /by G.A. Wiebe and David A. Reid.
Wiebe, G. A.(Gustav A.), 1899-. Reid, David A. 1914-. Washington : U.S. Dept. of Agriculture, 1958. 20 p. : ill., maps ; 23 cm. Includes bibliographical references. (NAL Call No.: DNAL 1 Ag84Te no.1176).

0057

Conservation-tillage and residue-management systems for interior Alaska.
AGBOB. Siddoway, F.H. Lewis, C.E.; Cullum, R.F. Fairbanks : The Station. *Agroborealis - Alaska Agricultural Experiment Station*, Fairbanks. Includes lists of species. July 1984. v. 16 (2). p. 35-40. ill. Includes 5 references. (NAL Call No.: DNAL S33.E2).

0058

Control of nitrate leaching with winter annual cover crops.
McCracken, D. Lexington, Ky. : The Department. Soil science news & views - Cooperative Extension Service and University of Kentucky, College of Agriculture, Department of Agronomy. 1989. v. 10 (2). 2 p. (NAL Call No.: DNAL S591.55.K4S64).

0059

Crop rotations and manure versus agricultural chemicals in dryland grain production.
JSWCA3. Sahs, W.W. Lesoing, G. Ankeny, Iowa : Soil Conservation Society of America. *Journal of soil and water conservation*. Nov/Dec 1984. v. 40 (6). p. 511-516. Includes 27 references. (NAL Call No.: DNAL 56.8 J822).

0060

C4 grasses and cereals growth, development, and stress response /C. Allan Jones. --.
Jones, C. Allan. New York : Wiley, c1985. "A Wiley-Interscience publication."~ Includes index. xi, 419 p. : ill. ; 24 cm. Bibliography: p. 333-412. (NAL Call No.: DNAL QK495.G74J66).

(PLANT PRODUCTION - FIELD CROPS)

0061

Damage to barley spikes resulting from impact momentum similar to small sized hail.
AGJ0AT. Ferguson, H. Jones, A.J.; Tsai, K.J. Madison, Wis. : American Society of Agronomy. Agronomy journal. Nov/Dec 1987. v. 79 (6). p. 1015-1018. Includes references. (NAL Call No.: DNAL 4 AM34P).

0062

Development rate and growth duration of oats in response to delayed sowing.
AGJ0AT. Colville, D.C. Frey, K.J. Madison, Wis. : American Society of Agronomy. Agronomy journal. May/June 1986. v. 78 (3). p. 417-421. Includes references. (NAL Call No.: DNAL 4 AM34P).

0063

Disease-resistant and hardy oats for the South /by T.R. Stanton and F.A. Coffman.
Stanton, T. R. 1885-. Coffman, Franklin A. 1892-. Washington, D.C. : U.S. Dept. of Agriculture, 1943. 10 p. : ill. ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1947).

0064

Disease-resistant and hardy oats for the South by T.R. Stanton and F.A. Coffman . --.
Stanton, T. R. Washington, D.C. : U.S. Dept. of Agriculture, 1943. 10 p. : ill. --. (NAL Call No.: DNAL Fiche S-70 no.1947).

0065

Dormant embryo morphology, seedling morphogenesis and tiller development of Himalaya barley / Richard H. Mullenax .
Mullenax, Richard H. Pullman, Wash. : Washington Agricultural Experiment Stations, College of Agriculture, Washington State University, 1967. Cover title. 6 p. : ill. ; 28 cm. Bibliography: p. 6. (NAL Call No.: DNAL 100 W27T no.55).

0066

Drills and seeders for heavy residues and untilled soils (Small grain planting equipment, minimum tillage farming, Kansas).
Powell, G.M. Manhattan : The Service. L - Cooperative Extension Service, Kansas State University. June 1982. June 1982. (634). 8 p. ill. (NAL Call No.: 275.29 K13LE).

0067

Effect of ammonium and nitrate on growth and yield of barley on acid soils.
CSOSA2. Malhi, S.S. Nyborg, M.; Caldwell, C.D.; Hoyt, P.B.; Leitch, R.H. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. May/Sept 1988. v. 19 (7/12). p. 1049-1063. Includes references. (NAL Call No.: DNAL S590.C63).

0068

Effect of burning cereal straw on soil properties and grain yields in Saskatchewan (Effects on physical, chemical and biological characteristics of the soil and on grain yields).
Biederbeck, V.O. Campbell, C.A.; Bowren, K.E.; Schnitzer, M.; McIver, R.N. Madison, Wis., The Society. Journal - Soil Science Society of America. Soil Science Society of America. Jan/Feb 1980. v. 44 (1). p. 103-111. ill. 42 ref. (NAL Call No.: 56.9 S03).

0069

The effect of defence reactions on the energy balance and yield of resistant plants (Barley, Erysiphe graminis hordei).
Smedegaard-Petersen, V. New York, Plenum Press. NATO advanced study institutes series. Series A. Life sciences. 1982. v. 37. p. 299-315. ill. Includes 31 ref. (NAL Call No.: QH301.N32).

0070

The effect of different times of plowing small-grain stubble in eastern Colorado /by O.J. Grace.
Grace, Oliver J. 1880-. Washington, D.C. : U.S. Dept. of Agriculture, 1915. Caption title. 15 p. : charts ; 24 cm. (NAL Call No.: DNAL 1 Ag84B no.253).

0071

Effect of erect leaf angle on grain yield in barley.
CRPSAY. Tunland, L. Chapko, L.B.; Wiersma, J.V.; Rasmusson, D.C. Madison, Wis. : Crop Science Society of America. Crop science. Jan/Feb 1987. v. 27 (1). p. 37-40. Includes references. (NAL Call No.: DNAL 64.8 C883).

0072

The effect of ethephon (2-chloroethyl) phosphonic acid on lodging and yield of small grain cereal crops.
Schwartz, T.K.PPGD. Coffin, R.H.; Evans, W.F.; Nash, R.L.; Rao, K.P. Lake Alfred : The Society. Proceedings annual meeting - Plant Growth Regulator Society of America. 1982.

(PLANT PRODUCTION - FIELD CROPS)

1982. (9th). p. 89-95. Includes references.
(NAL Call No.: S8128.P5).

0073

Effect of mefluidide on rye forage yield and quality.

Breman, J.W. Fletcher, J.H. S.I. : The Society. Proceedings - Soil and Crop Science Society of Florida. 1987. v. 46. p. 35-38. Includes references. (NAL Call No.: DNAL 56.9 S032).

0074

Effect of nitrogen nutrition on quality of agronomic crops.

Deckard, E.L. Tsai, C.Y.; Tucker, T.C. Madison, Wis. : American Society of Agronomy, 1984. Nitrogen in crop production : proceedings, symposium, 25-27 May, 1982, Sheffield, Alabama / spon. by National Fertilizer Development Center of Tennessee Valley Authority ... et al. ; Roland D. Hauck. Literature review. p. 601-615. Includes references. (NAL Call No.: DNAL S651.N57).

0075

Effect of plant growth regulators and nitrogen on the agronomic performance of small grain crops.

Leary, W.P. PPGD. Oplinger, E.S. Lake Alfred : The Society. Proceedings annual meeting - Plant Growth Regulator Society of America. 1983. 1983. (10th). p. 277-286. Includes references. (NAL Call No.: S8128.P5).

0076

Effect of planting dates on spring barley varieties, San Juan Branch Station, 1974-1979 (New Mexico, performance, yield).

Gregory, E.J. Arnold, R.N. Las Cruces : The Station. Research report - New Mexico, Agricultural Experiment Station. June 1982. June 1982. (478). 12 p. 8 ref. (NAL Call No.: 100 N465R).

0077

The effect of powdery mildew on the response of barley to applied nitrogen.

Young, K.J. Khan, T.N. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 177-182. Includes references. (NAL Call No.: DNAL SB191.B2A9 1987).

0078

The effect of residual copper levels on the nutrition and yield of oats grown in microplots on three organic soils.

CSOSA2. Belanger, A. Levesue, M.; Mathur, S.P. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. Jan 1986. v. 17 (1). p. 85-96. Includes 12 references. (NAL Call No.: DNAL S590.C63).

0079

The effect of ryegrass and rye seeding rate on forage yield planted alone and in mixtures in south Alabama.

Kee, D.D. Duffy, P.D.; Ward, C.Y. Lexington, Ky. : The Conference. Proceedings of the Forage and Grassland Conference. 1988. p. 221-225. Includes references. (NAL Call No.: DNAL S8193.F59).

0080

Effect of seeding rate and time of N application on oat grain yield.

JPRAEN. Ahmadi, M. Wiebold, W.J.; Beuerlein, J.E. Madison, Wis. : American Society of Agronomy. Journal of production agriculture. July/Sept 1988. v. 1 (3). p. 242-244. Includes references. (NAL Call No.: DNAL S539.5.J68).

0081

The effect of selection in pure-line oat work.

AGJDAT. Spragg, F.A. Madison, Wis. : American Society of Agronomy. Agronomy journal. 1912. v. 4. p. 81-83. (NAL Call No.: DNAL 4 AM34P).

0082

Effect of slope position on the microclimate, growth, and yield of barley.

AGJDAT. Whitman, C.E. Hatfield, J.L.; Reginato, R.J. Madison, Wis. : American Society of Agronomy. Agronomy journal. Sept/Oct 1985. v. 77 (5). p. 663-669. Includes references. (NAL Call No.: DNAL 4 AM34P).

0083

The effect of tillage method, crop sequence, and date of seeding upon the yield and quality of cereals and other crops grown under dry-land conditions in north-central Montana /by M.A. Bell.

Bell, M. A. 1897-. Bozeman, Mont. : Montana State College, Agricultural Experiment Station, 1937. 118 p. ; 23 cm. Bibliography: p. 92. (NAL Call No.: DNAL 100 M76 (1) no.336).

(PLANT PRODUCTION - FIELD CROPS)

0084

The effect of trace elements on alfalfa and oats in Minnesota /John M. MacGregor and John F. Mulvehill.

MacGregor, John M. Mulvehill, John F. St. Paul : University of Minnesota, Agricultural Experiment Station, 1955. 16 p. : maps ; 23 cm. Bibliography: p. 16. (NAL Call No.: DNAL 100 M66 (3) no.213).

0085

The effect of weed and invertebrate pest management on alfalfa establishment in oat stubble.

Bahler, C.C. Byers, R.A.; Stout, W.L. Ankeny, Iowa : Soil Conservation Society of America, c1987. The role of legumes in conservation tillage systems / J.F. Power, editor. Paper presented at the "National Conference on the Role of Legumes in Conservation Tillage Systems", April 27-29, 1987, University of Georgia, Athens, Georgia. p. 76-77. Includes references. (NAL Call No.: DNAL SB203.R6).

0086

Effects of by-product sulfuric acid on phyto availability of nutrients in irrigated calcareous, saline-sodic soils (Barley, *Hordeum vulgare*).

Cates, R.L. Jr. Haby, V.A.; Skogley, E.D.; Ferguson, H. Madison, Wis. : American Society of Agronomy. Journal of environmental quality. Apr/June 1984. v. 13 (2). p. 252-256. Includes references. (NAL Call No.: QH540.U6).

0087

Effects of chloride fertilizer and systemic fungicide seed treatments on common root rot of barley.

PLDRA. Shefelbine, P.A. Mathre, D.E.; Carlson, G. St. Paul, Minn. : American Phytopathological Society. Plant disease. July 1986. v. 70 (7). p. 639-642. Includes 22 references. (NAL Call No.: DNAL 1.9 P69P).

0088

Effects of forage legumes on yield and nitrogen uptake by a succeeding barley crop.

AAREEZ. Abernethy, R.H. Bohl, W.H. New York : Springer. Applied agricultural research. 1987. v. 2 (2). p. 97-102. Includes references. (NAL Call No.: DNAL S539.5.A77).

0089

Effects of N, P, and K (nitrogen, phosphorus, potassium) fertilization on barley grown in a newly cleared subarctic soil (Alaska).

Michaelson, G.J. Loynachan, T.E.; Woding, F.J.; Mitchell, G.A. Madison, Wis. : American Society of Agronomy. Agronomy journal. July/Aug 1982. v. 74 (4). p. 694-699. ill., map. 16 ref. (NAL Call No.: 4 AM34P).

0090

Effects of nitrogen fertilization on manganese concentration and yield of barley and oats (*Hordeum vulgare*, *Avena sativa*).

Petrie, S.E. Jackson, T.L. Madison, Wis. : The Society. Journal - Soil Science Society of America. Mar/Apr 1984. v. 48 (2). p. 319-322. Includes references. (NAL Call No.: 56.9 S03).

0091

Effects of nitrogen fertilizer rate, seeding rate, and row spacing on semidwarf and conventional height spring oat.

CRPSAY. Marshall, H.G. Kolb, F.L.; Roth, G.W. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1987. v. 27 (3). p. 572-575. Includes references. (NAL Call No.: DNAL 64.8 C883).

0092

Effects of nitrogen rate and placement on two malting barley varieties (Comparisons, Minnesota).

Varvel, G.E. MXMRA. Severson, R.K. St. Paul : The Station. Miscellaneous publication - University of Minnesota, Agricultural Experiment Station. 1983. 1983. (2 rev.). p. 63-64. (NAL Call No.: S1.M52).

0093

Effects of planting date on development of net blotch epidemics in winter barley in Pennsylvania.

PLDRA. Delserone, L.M. Cole, H. Jr. St. Paul, Minn. : American Phytopathological Society. Plant disease. May 1987. v. 71 (5). p. 438-441. Includes references. (NAL Call No.: DNAL 1.9 P69P).

0094

The effects of using copper for mitigating histosol subsidence on: 1. The yield and nutrition of oats and lettuce grown on histosols, mineral sublayers, and their mixtures.

Levesque, M.P. SOSCA. Mathur, S.P. Baltimore : Williams & Wilkins. Soil science. Feb 1983. v. 135 (2). p. 88-100. Includes references. (NAL

Call No.: 56.8 S03).

0095

Effects of various salts on barley growth /H.E. Dregne.

Dregne, H. E. University Park, N.M. : Agricultural Experiment Station, New Mexico State University, 1962. Caption title. 5, 1 p. : charts ; 26 cm. (NAL Call No.: DNAL 100 N465R no.62).

0096

Establishment of legumes using no-till seedings with and without herbicides.

AAREEZ. Guillard, K. Allinson, D.W. New York : Springer. Applied agricultural research. 1987. v. 1 (5). p. 281-288. Includes references. (NAL Call No.: DNAL S539.5.A77).

0097

Ethephon application and nitrogen fertilization of irrigated winter barley in an arid environment.

AGJOAT. Pearson, C.H. Golus, H.M.; Tindall, T.A. Madison, Wis. : American Society of Agronomy. Lodging in barley (*Hordeum vulgare* L.) can result in grain loss, lower seed quality, increased disease, high grain moisture, reduced harvest efficiency, and lower milling grade. Application of N fertilizer to achieve high barley yields often results in increased lodging. Consequently, N applications are often reduced in order to lessen the potential for lodging. Use of plant growth regulators may permit higher N application rates without increased lodging. Field studies were conducted in an irrigated arid environment during 1986 and 1987 on Fruita sandy clay loam (fine-loamy, mixed, mesic Typic Haplargid) soil. Ethephon (2-chloroethyl) phosphonic acid was applied at 0.42 kg a.i. ha⁻¹ to irrigated winter barley that received 56, 112, 168, and 224 kg ha⁻¹ of spring-applied N. The 1986 data indicated that, in the absence of lodging, ethephon did not affect barley grain yields. Ethephon increased grain yields in 1987 in association with reduced lodging. Plant height response to ethephon was similar across N rates. Ethephon shortened internode lengths on the upper portion of the culm and reduced lodging of winter barley grown at high N rates. Ethephon did not affect spikes m⁻², kernel mass, kernels spike⁻¹, or seed N content in either year. Application of higher N rates than traditionally used in irrigated winter barley increased grain yield and seed N content. In years when lodging occurred, ethephon was effective as an antilodging agent. *Agronomy journal*. Sept/Oct 1989. v. 81 (5). p. 7171-719. Includes references. (NAL Call No.: DNAL 4 AM34P).

0098

Field experiments with oats, 1890 / George E. Morrow, Thomas F. Hunt . Milk and butter tests / G.E. Morrow . Cream raising by dilution / G.E. Morrow, E.H. Farrington . The hessian fly / S.A. Forbes . Canada thistles : their extermination / T.J. Burrill .
Morrow, G. E. 1840-. Hunt, Thomas F. 1862-; Morrow, G. E. 1840-; Morrow, G. E. 1840-; Farrington, E. H. 1860-; Forbes, Stephen Alfred, 1844-1930.; Burrill, Thomas Jonathan, 1839-1916. Champaign, Ill. : University of Illinois Agricultural Experiment Station, 23 cm. Caption title. p. 353 -388 : ill. ; 23 cm. (NAL Call No.: DNAL 100 I165 no.12).

0099

For horses: Kelly.

SDFHA. Leslie, J. Brookings, S.D. : The Station. South Dakota farm and home research - South Dakota, Agricultural Experiment Station. 1984. v. 35 (2). p. 14. ill. (NAL Call No.: DNAL 100 S082S).

0100

Genetic and environmental effects on rye grain production in Florida.

Pfahler, P.L. Barnett, R.D.; Luke, H.H. S.1. : The Society. *Proceedings - Soil and Crop Science Society of Florida*. 1985. v. 44. p. 167-170. Includes 16 references. (NAL Call No.: DNAL 56.9 S032).

0101

Genetic and temperature effects on rye germination and seedling development.

Sullivan, B.P. Peahler, P.L. S.1. : The Society. *Proceedings - Soil and Crop Science Society of Florida*. 1985. v. 44. p. 205-208. Includes 10 references. (NAL Call No.: DNAL 56.9 S032).

0102

Genotypic variability in response of oat to delayed sowing.

AGJOAT. Baltenberger, D.C.C. Frey, K.J. Madison, Wis. : American Society of Agronomy. *Agronomy journal*. Sept/Oct 1987. v. 79 (5). p. 813-816. Includes references. (NAL Call No.: DNAL 4 AM34P).

(PLANT PRODUCTION - FIELD CROPS)

0103

Grain-filling duration and yield in spring barley.

CRPSAY, Metzger, D.D. Czaplewski, S.J.; Rasmusson, D.C. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1984. v. 24 (6). p. 1101-1105. Includes 11 references. (NAL Call No.: DNAL 64.8 C883).

0104

Grain pesticides American Association of Cereal Chemists. --.

Washington, D.C.? : The Association, 1985. Title from container. 4 videocassettes (362 min.) : sd., col. : 1/2 in. (NAL Call No.: DNAL Videocassette no.21).

0105

Grass weed control in California cereals: update and review.

WSWPA, Mitich, L.W. Kyser, G.B. Reno : The Society. Proceedings - Western Society of Weed Science. Paper presented at the annual meeting of the Western Society of Weed Science, March 18-20, 1986, San Diego, California. 1986. v. 39. p. 169-172. (NAL Call No.: DNAL 79.9 W52).

0106

Grow disease-resistant oats /by T.R. Stanton and F.A. Coffman.

Stanton, T. R. 1885-. Coffman, Franklin A. 1892-. Washington, D.C. : U.S. Dept. of Agriculture, 1943. 13 p. : ill. ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1941).

0107

Grow disease-resistant oats by T.R. Stanton and F.A. Coffman. --.

Stanton, T. R. Washington, D.C. : U.S. Dept. of Agriculture, 1949. 13 p. : ill. --. (NAL Call No.: DNAL Fiche S-70 no.1941 1949).

0108

Growth and yield of barley isopopulations differing in solute potential.

CRPSAY, Grumet, R. Albrechtsen, R.S.; Hanson, A.D. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1987. v. 27 (5). p. 991-995. Includes references. (NAL Call No.: DNAL 64.8 C883).

0109

Influence of weed control treatments on soybean cultivars in an oat-soybean rotation.

AGJOAT, Burnside, D.C. Moomaw, R.S. Madison, Wis. : American Society of Agronomy. Agronomy journal. Nov/Dec 1984. v. 76 (6). p. 887-890. Includes 13 references. (NAL Call No.: DNAL 4 AM34P).

0110

Insects on small grains and their control (Includes cultural control).

Coppock, S. Burton, R. Stillwater : The Service. DSU extension facts - Cooperative Extension Service, Oklahoma State University. Mar 1983. Mar 1983. (7176 rev.). 4 p. ill. (NAL Call No.: S544.3.0505).

0111

Interactions between calcium amendments and phosphate on the response of alfalfa and barley growing on acid soils.

CSDSA2. Soon, Y.K. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. May/Sept 1988. v. 19 (7/12). p. 1343-1353. Includes references. (NAL Call No.: DNAL S590.C63).

0112

Isogenic heading date effects on yield component development in 'Titan' barley.

CRPSAY, Smail, V.W. Eslick, R.F.; Hockett, E.A. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 1023-1029. Includes references. (NAL Call No.: DNAL 64.8 C883).

0113

Jaycee spring oats.

Purdue University - Cooperative Extension Service. 1967. This publication examines Jaycee, C17971, as an acceptable spring growing oat for the southern one fourth of Indiana and is recommended for production in the rest of Indiana. Document available from: Mailing Room, Ag. Administration Bldg., Purdue University, West Lafayette, IN 47907. 1 sheet. (NAL Call No.: ID-69).

0114

Kline barley: a new winter feed grain for Georgia and the Southeast.

GARRA, Brown, A.R. Johnson, J.W.; Cunfer, B.M.; Morey, D.D. Athens, Ga. : The Stations. Research report - University of Georgia, College of Agriculture, Experiment Stations. May 1985. (469). 6 p. Includes references. (NAL Call No.: DNAL S51.E22).

(PLANT PRODUCTION - FIELD CROPS)

0115

Leaf hopper injurious to cereal and forage crops /by Herbert Osborn.
Osborn, Herbert, 1856-. Washington, D.C. : U.S. Dept. of Agriculture, 1932. Caption title.~ "Contribution from Bureau of Entomology.". 34 p. : ill., maps ; 23 cm. Bibliography: p. 33-34. (NAL Call No.: DNAL 1 Ag84C no.241).

0116

Legume growth and residual effects on oat (Avena sativa L.) production.
CSOSA2. Guillard, K. Allinson, D.W. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. Apr 1985. v. 16 (4). p. 375-383. Includes 13 references. (NAL Call No.: DNAL S590.C63).

0117

Losses in harvesting and threshing grain.
AGUOAT. Bracken, A.F. Madison, Wis. : American Society of Agronomy. Agronomy journal. Aug 1925. v. 17 (8). p. 508-514. (NAL Call No.: DNAL 4 AM34P).

0118

Measured crop performance: small grain, 1984.
NUSRA. Bowman, D.T. Raleigh, N.C., The Station. Research report - Dept. of Crop Science, North Carolina State University, Agricultural Experiment Station. Includes statistical data. July 1984. (96). 26 p. maps. (NAL Call No.: DNAL 100 N8122).

0119

Measured crop performance: small grain, 1985.
NUSRA. Bowman, D.T. Raleigh, N.C., The Station. Research report - Dept. of Crop Science, North Carolina State University, Agricultural Experiment Station. Includes statistical data. July 1985. (101). 29 p. maps. (NAL Call No.: DNAL 100 N8122).

0120

Methods of eradicating buckthorn (Rhamnus) susceptible to crown rust (Puccinia coronata) of oats /by S.M. Dietz and L.D. Leach.
Dietz, S. M. 1893-. Leach, L. D. 1900-. Washington, D.C. : U.S. Dept. of Agriculture, 1930. Caption title.~ "Contribution from Bureau of Plant Industry.". 16 p. : ill., charts, maps ; 24 cm. Bibliography: p. 15. (NAL Call No.: DNAL 1 Ag84C no.133).

0121

Methods of seeding oats drilling and broadcasting /by Albert N. Hume, O.D. Center and Leonard Hegnauer.
Hume, Albert N. 1878-. Center, O. D. 1872-; Hegnauer, Leonard, 1872-. Urbana, Ill. : University of Illinois Agricultural Experiment Station, 1909. Cover title. p. 297 -312 : ill. ; 23 cm. (NAL Call No.: DNAL 100 I16S no.136).

0122

Miniature thrasher and separator.
AGUOAT. Mackie, W.W. Hoffman, A.H. Madison, Wis. : American Society of Agronomy. Agronomy journal. Jan 1924. v. 16 (1). p. 57-60. (NAL Call No.: DNAL 4 AM34P).

0123

Modification of nutritional quality by environment and production practices.
AGRYA. Sander, D.H. Allaway, W.H.; Olson, R.A. Madison, Wis. : American Society of Agronomy. Agronomy. 1987. (28). p. 45-82. Includes references. (NAL Call No.: DNAL 4 AM392).

0124

Modifications of plant growth and ash content as effected by acids added to soils.
AGUOAT. Carr, R.H. Haverkamp, H.G. Madison, Wis. : American Society of Agronomy. A comparison has been made of the effect upon the growth of soybean and buckwheat plants of adding an organic (acetic) and an inorganic (sulfuric) acid to a silt and a clay-loam soil. It has been found that both acids seriously interfere with plant growth, but that acetic was more deleterious than sulfuric acid. This is believed to be due to the effect of acetic acid in supplying more toxic aluminum and iron, etc., to the plant and in decreasing calcium and magnesium to a greater extent than was done by the sulfuric acid. In addition two other series of acids (phosphoric and silicic) were used, because they make compounds with aluminum and iron which seem to be harmless to the growing plant, regardless of the fact that the root is surrounded by a soil which is highly acid. The good growth of both kinds of plants obtained, even when rather large amounts of these latter acids were added, indicates that relatively insoluble compounds of aluminum and iron, etc., were formed and that the acid by itself did no harm. Agronomy journal. Apr 1924. v. 16 (4). p. 278-283. (NAL Call No.: DNAL 4 AM34P).

(PLANT PRODUCTION - FIELD CROPS)

0125

Montana small grains integrated crop and pest management 1984-85 field summary.
Bozeman, Mont. : Cooperative Extension Service
Montana State University, 1986-. Cover title.-
"February, 1986."~ Caption title: Integrated
crop and pest management, 1984-85. v. : map ;
28 cm. (NAL Call No.: DNAL 275.29 M76C no.1332
etc.).

0126

**Mosaics of winter oats and their control in the
southeastern states /by H.H. McKinney ... et
al. .**
McKinney, Harold Hall, 1889-. Washington, D.C.
: U.S. Dept. of Agriculture, 1949. Caption
title. 17 p. : ill. ; 23 cm. Includes
bibliographical references. (NAL Call No.: DNAL
1 Ag84C no.809).

0127

**Narrow row soybean production in untilled oat
stubble.**
AGJDAT. Burnside, D.C. Moomaw, R.S. Madison,
Wis. : American Society of Agronomy. Agronomy
journal. Jan/Feb 1985. v. 77 (1). p. 36-40.
Includes 11 references. (NAL Call No.: DNAL 4
AM34P).

0128

**Natural selection in a doubled-haploid mixture
and a composite cross of barley.**
CRPSAY. Patel, J.D. Reinbergs, E.; Mather,
D.E.; Choo, T.M.; Sterling, J.D.E. Madison,
Wis. : Crop Science Society of America. Crop
science. May/June 1987. v. 27 (3). p. 474-479.
Includes references. (NAL Call No.: DNAL 64.8
C883).

0129

**Nematode populations in a rye/soybean
succession after four years of no tillage
management.**
Post, T.J. Gallaher, R.N.; Dickson, D.W.
Gainesville, Fla. : The Station. Agronomy
research report AY - Agricultural Experiment
Stations, University of Florida. 1983? .
(84-6). 5 p. Includes references. (NAL Call
No.: DNAL S540.A2F62).

0130

**New varieties of oats from bond crosses
resistant to Victoria Blight /by T.R. Stanton.**
Stanton, T. R. 1885-. Washington, D.C. : U.S.
Dept. of Agriculture, 1948. 7 p., 1 folded leaf
: ill. ; 26 cm. Bibliography: p.6-7. (NAL Call
No.: DNAL 1 Ag84C no.795).

0131

Nitrate accumulation in plants.
Jensen, E.H. Reno, Nev. : College of
Agriculture, University of Nevada-Reno, Nevada
Cooperative Extension. Fact sheet - College of
Agriculture, University of Nevada-Reno, Nevada
Cooperative Extension. 1987? . (87-32). 3 p.
(NAL Call No.: DNAL S544.3.N3C66).

0132

**Nitrate content in oats /by B.J. Kolp ... et
al. .**
Kolp, B. J. 1928-. Laramie, Wyo. : University
of Wyoming, Agricultural Experiment Station,
1963. 12 p. ; 23 cm. Bibliography: p. 12. (NAL
Call No.: DNAL 100 W99 (1) no.409).

0133

**Nitrogen and yield as related to water use of
spring barley (*Hordeum vulgare*, seasonal
evapotranspiration, New Mexico).**
Kallsen, C.E. AGJDAT. Sammis, T.W.; Gregory,
E.U. Madison : American Society of Agronomy.
Agronomy journal. Jan/Feb 1984. v. 76 (1). p.
59-64. Includes references. (NAL Call No.: 4
AM34P).

0134

**Nitrogen harvest index in oats: its
repeatability and association with adaptation.**
CRPSAY. Rattunde, H.F. Frey, K.J. Madison, Wis.
: Crop Science Society of America. Crop
science. May/June 1986. v. 26 (3). p. 606-610.
Includes references. (NAL Call No.: DNAL 64.8
C883).

0135

Norwind barley.
Copeland, L. O. Grafius, J. E.; Hart, L.
P. & MSU Ag Facts. Document available from:
Michigan State Univ., Bulletin Office, P.O. Box
231, East Lansing, Michigan 48824 1980. This
discusses Norwind barley, a new variety of
winter barley released by Michigan State
University in 1978, including disease
resistance, description, and seed availability.
1 sheet : ill. (NAL Call No.: Document
available from source.). (NAL Call No.:
Extension Bulletin E-1379).

0136

Oats.
KAEB. Harper, J.N. Peter, A.M. Lexington : The
Station. Bulletin - Kentucky, Agricultural
Experiment Station. Documents available from:
Agriculture Library, Agricultural Science
Center - North, University of Kentucky,
Lexington, Ky. 40546-0091. Apr 1902. (99). p.

(PLANT PRODUCTION - FIELD CROPS)

63-78. (NAL Call No.: DNAL 100 K41 (2)).

0137

Oats (Production, breeding, composition).
Youngs, V.L. Peterson, D.M.; Brown, C.M. St. Paul, Minn. : American Association of Cereal Chemists. Advances in cereal science and technology. 1982. Literature review. v. 5. p. 49-105. ill. Includes references. (NAL Call No.: TS2120.A3).

0138

Oats research.
Reeves, D.L. Brookings, S.D. : The Station. Plant science pamphlet - Plant Science Dept., Agricultural Experiment Station, South Dakota State University. In the series analytic: 1986 Annual progress report, Northeast Research Station, Watertown, South Dakota. Jan 1987. (100). p. 7. (NAL Call No.: DNAL S541.5.S8P5).

0139

Oats research.
Reeves, D.L. Brookings, S.D. : The Station. Plant science pamphlet - Plant Science Dept., Agricultural Experiment Station, South Dakota State University. In the series analytic: 1987 Annual Progress Report--Northeast Research Station, Watertown, South Dakota. Jan 1988. (5). p. 37-38. (NAL Call No.: DNAL S541.5.S8P5).

0140

On the prevention of oat smut and potato scab / R.A. Moore .
Moore, R. A. 1861-1941. Madison, Wis. : University of Wisconsin, Agricultural Experiment Station, 1903. Cover title. 23 p. : ill. ; 23 cm. (NAL Call No.: DNAL 100 W75 (1) no.98).

0141

Path coefficient analysis of correlation between stress and barley yield components.
Campbell, W.F. Wagenet, R.J.; Bamatraf, A.M.; Turner, D.L. Madison, Wis., American Society of Agronomy. Agronomy journal. Nov/Dec 1980. v. 72 (6). p. 1012-1016. ill. 15 ref. (NAL Call No.: 4 AM34P).

0142

Performance of cereal crops in the Tanana River Valley of Alaska, 1978-1979 (Varieties, yields, resistance).
Wooding, F.J. McBeath, J.H. Fairbanks : The Station. Circular - Alaska Agricultural Experiment Station. Dec 1979. Dec 1979. (34). 23 p. (NAL Call No.: 100 AL122E).

0143

Performance of cereal crops in the Tanana Valley of Alaska, 1980 (Varieties, yields, disease resistance).
Wooding, F.J. McBeath, J.H.; Hanscom, J.T.; Van Veldhuizen, R.M. Fairbanks : The Station. Circular - Alaska Agricultural Experiment Station. Dec 1980. Dec 1980. (36). 29 p. (NAL Call No.: 100 AL122E).

0144

Performance of cereal crops in the Tanana Valley of Alaska, 1981 (Varieties, yields, disease resistance).
Wooding, F.J. McBeath, J.H.; Hanscom, J.T.; Van Veldhuizen, R.M.; Delucchi, G.M. Fairbanks : The Station. Circular - Alaska Agricultural Experiment Station. Mar 1982. Mar 1982. (42). 27 p. (NAL Call No.: 100 AL122E).

0145

Performance of cereal crops in the Tanana Valley of Alaska, 1982 (Variety trials, disease resistance comparisons, yields).
Wooding, F.J. McBeath, J.H.; Frost, S.; Hanscom, J.T.; Van Veldhuizen, R.M. Fairbanks : The Station. Circular - Alaska Agricultural Experiment Station. Feb 1983. Feb 1983. (44). 28 p. (NAL Call No.: 100 AL122E).

0146

Pest control in forages and small grains 1980 / J. D. Doll ... (et al.). -
WI. Doll, J. D. Madison University of Wisconsin, Cooperative Extension Programs 1980. Cover title. 37 p. : ill. 28 cm. (NAL Call No.: S544.3.W6W53 no. 1981).

0147

Physiological changes associated with three cycles of recurrent selection for grain yield improvement in oats.
CRPSAY. Payne, T.S. Stuthman, D.D.; McGraw, R.L.; Bregitzer, P.P. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 734-736. Includes references. (NAL Call No.: DNAL 64.8 C883).

(PLANT PRODUCTION - FIELD CROPS)

0148

Plant response to topsoil thickness on an eroded loess soil.

JSWCA3. Mielke, L.N. Schepers, J.S. Ankeny, Iowa : Soil Conservation Society of America. Journal of soil and water conservation. Jan/Feb 1986. v. 41 (1). p. 59-63. Includes 15 references. (NAL Call No.: DNAL 56.8 J822).

0149

The potential effect of windbreaks, barriers and residues on the development of small grain plants.

Freckleton, M.E. Bozeman, Mont. : Montana State University, Cooperative Extension Service. Great Plains Agriculture i.e. Agricultural Council publication. Paper presented at the "International Symposium on Windbreak Technology," June 23-27, 1986, Lincoln, Nebraska. 1986. (117). p. 139. (NAL Call No.: DNAL S27.A3).

0150

Profitable small grain production in the Texas Gulf Coast.

TAEBA. Miller, T.D. Livingston, S. College Station, Tex. : The Station. B - Texas Agricultural Experiment Station. Nov 1987. (1587). 8 p. (NAL Call No.: DNAL 100 T31S (1)).

0151

Reduced tillage systems for Montana (Small grain production, includes herbicides and pesticides application guidelines).

Rardon, P. Bozeman : The Service. Bulletin - Cooperative Extension Service. Montana State University. Mar 1983. Mar 1983. (1286). 28 p. 111. (NAL Call No.: 275.29 M76C).

0152

Reduced yield of spring oat cultivars by cereal leaf beetles (*Oulema melanopus*).

Webster, J.A. Smith, D.H.; Hoxie, R.P. Peoria, Ill., The Administration. Abstract: In this research, we determined grain yield loss of oat cultivars commonly grown in cereal leaf beetle infested areas of Michigan. We measured yield loss by comparing grain yield and panicle weights from infested plots with those of uninfested plots of three locations. Two out of three tests showed significant losses in kg/ha grain yield, with an average loss of 17 percent and a range of 4 to 20 percent. The average larval density was one larva per two stems with a range of 0 to 1.2 percent. The average larval density was one larva per two stems with a range of 0 to 1.2 larvae per stem. Of the yield components, no significant losses were noted in fertile tillers (except in one test where red leaf virus was present), but significant losses were found in kernels per panicle and kernel

weight. 'Korwood' and 'Orbit' were the highest yielding cultivars, while 'Clintland 64' was the lowest. Agricultural research results; ARR-NC - United States, Dept. of Agriculture, Science and Education Administration. Oct 1980. Oct 1980. (6). 6 p. 111. 10 ref. (NAL Call No.: aS21.A75U75).

0153

Relation of dust fungicides to flow of small grains through drills and to drill injury /by R.W. Leukel.

Leukel, R. W. 1888-. Washington, D.C. : U.S. Dept. of Agriculture, 1930. Caption title.~ "Contribution from Bureau of Plant Industry." 10 p. ; 23 cm. Includes bibliographical references. (NAL Call No.: DNAL 1 Ag84C no.119).

0154

Response to spring barley (*Hordeum vulgare*) to herbicides.

WETEE9. Clay, S.A. Thill, D.C.; Cochran, V.L. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Jan 1988. v. 2 (1). p. 68-71. Includes references. (NAL Call No.: DNAL SB610.W39).

0155

Salinity, irrigation frequency, and fertilization effects on barley growth.

Wagenet, R.J. Campbell, W.F.; Bamatraf, A.M.; Turner, D.L. Madison, Wis., American Society of Agronomy. Agronomy journal. Nov/Dec 1980. v. 72 (6). p. 969-974. 111. 23 ref. (NAL Call No.: 4 AM34P).

0156

Small grain production and utilization in Georgia.

Lee, R.D. Bullock, D.; Segars, W.I.; Bullock, F.; Suber, E.F.; Hudson, R.D.; Thompson, S.S.; Hammond, C.; Johnson, J.T.; Owsley, W.F. Athens, Ga. : The Service. Bulletin - Cooperative Extension Service, University of Georgia, College of Agriculture. Feb 1986. (929). 18 p. (NAL Call No.: DNAL 275.29 G29B).

0157

Soil fertility and defoliation effects with arrowleaf clover and nitrogen fertilizer equivalence of crimson-arrowleaf clover combinations (*Trifolium vesiculosum*, *Trifolium incarnatum*, *Secale cereale*, rhizobium, nitrogen fixation by forage legumes, Oklahoma).

Lynd, J.Q. AGJDAT. Hanlon, E.A. Jr.; Odell, G.V. Jr. Madison : American Society of Agronomy. Agronomy journal. Jan/Feb 1984. v. 76 (1). p. 13-16. 111. Includes references. (NAL Call No.:

(PLANT PRODUCTION - FIELD CROPS)

4 AM34P).

0158

Spring interseeding of winter rye with cover crops.

JSWCA3. Edwards, L.M. Ankeny, Iowa : Soil Conservation Society of America. Journal of soil and water conservation. May/June 1986. v. 41 (3). p. 190-191. Includes 5 references. (NAL Call No.: DNAL 56.8 J822).

0159

Stripe resistance and yield of smooth-awned barley hybrids / R.G. Shands ... et al. .

Shands, R. G. 1903-. Madison, Wis. : Agricultural Experiment Station of the University of Wisconsin, 1933. Cover title. 22 p. : ill. ; 23 cm. Bibliography: p. 22. (NAL Call No.: DNAL 100 W75 (2) no.116).

0160

Student's method for interpreting paired experiments.

AGJ0AT. Love, H.H. Brunson, A.M. Madison, Wis. : American Society of Agronomy. Too close an analogy should not be drawn between the determination of exact constants and the results of comparative agronomic experiments. Bessel's and Peter's formulas are not adapted to calculating the degree of significance of a difference between two varieties or two comparative treatments in experiments in which the variability within each variety or treatment is high because they extend through a series of years or because of place effect. For observations which naturally arrange themselves in pairs, Student's method is a better method with which to determine the probability of the difference, and deserves a larger application in the interpretation of agronomic investigations. Agronomy journal. Jan 1924. v. 16 (1). p. 60-68. (NAL Call No.: DNAL 4 AM34P).

0161

Studies on the effect of nitrogen applied to oats at different periods of growth.

AGJ0AT. Gericke, W.F. Madison, Wis. : American Society of Agronomy. Agronomy journal. Nov 1922. v. 14 (8). p. 312-320. (NAL Call No.: DNAL 4 AM34P).

0162

Unit 8--growing small grain.

Delke, E. A. Fenster, W. E.; Chambers, G. R.; Radford, J. D. & Management of crops and soils in NC and NE Minnesota correspondence course. Document available from: University of Minnesota, Bulletin Room, 1420 Eckles Avenue, St. Paul, Minnesota 55108 1983. Provides

information to recognize the management factors that will help you attain best small grain yields on your farm and understand which small grains will grow best on your farm. 7 p. (NAL Call No.: Document available from source.). (NAL Call No.: EF 662).

0163

Volunteer Jerusalem artichoke (*Helianthus tuberosus*) interference and control in barley (*Hordeum vulgare*).

WETEE9. Wall, D.A. Friesen, G.H. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Jan/Mar 1989. v. 3 (1). p. 170-172. Includes references. (NAL Call No.: DNAL SB610.W39).

0164

Water stress: role in differential aluminum tolerance of barley genotypes.

AGABB. Krizek, D.T. Foy, C.D. Madison, Wis. : American Society of Agronomy. Agronomy abstracts. Includes abstract. 1981. (73rd). p. 181-182. (NAL Call No.: DNAL 241 AM39).

0165

Weed control and its impact on cereal production (Herbicides, yield increases, Tunisia).

Ettounsi, L. Corvallis : The Station. Special report - Agricultural Experiment Station, Oregon State University. 1982. French text p. 124-137. List of common weed species. 1982. (668). p. 321-331. maps. (NAL Call No.: 100 OR3M).

0166

Weed-control evaluations in no-till soybeans (*Glycine max*) double-cropped with rye (*Secale cereale*) (Georgia).

Banks, P.A. GARRA. Kvien, J.S. Athens : The Stations. Research report - University of Georgia, College of Agriculture, Experiment Stations. July 1983. July 1983. (431). 6 p. Includes references. (NAL Call No.: S51.E22).

0167

Weed control in conservation tillage systems--small grains.

Wicks, G.A. Champaign, Ill. : Weed Science Society of America. Monograph series of the Weed Science Society of America. Literature review. 1985. (2). p. 77-92. Includes references. (NAL Call No.: DNAL SB610.M65).

(PLANT PRODUCTION - FIELD CROPS)

0168

Weed management in dryland cereal production with special reference to the Near East.
PPRBA. Kukula, S.T. Rome : World Reporting Service on Plant Diseases and Pests, FAO. Plant protection bulletin. 1986. v. 34 (3). p. 133-138. Includes references. (NAL Call No.: DNAL 421 P692).

0169

White grubs in cereal and forage crops and their control / T.R. Chamberlin and C.L. Fluke .
Chamberlin, Thomas R. 1889-. Fluke, C. L. 1891-. Madison : Agricultural Experiment Station, University of Wisconsin, 1947. Cover title. 15 p. : ill. ; 23 cm. (NAL Call No.: DNAL 100 W75 no.159).

0170

Wild oats control in barley.
WAEBA. Miller, S.D. Lauer, J. Laramie : The Station. Bulletin B - Wyoming, Agricultural Experiment Station. 1986. (885). p. 138-139. (NAL Call No.: DNAL 100 W99 (1)).

0171

Winter oats for the South /by T.R. Stanton and F.A. Coffman.
Stanton, T. R. 1885-. Coffman, Franklin A. 1892-. Washington, D.C. : U.S. Dept. of Agriculture, 1951. "Revision of and supersedes Farmers' bulletin 1947, Disease-resistant and hardy oats in the South.". 19 p. : ill. ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.2037).

0172

Year-round forage production with sodseeded cool-season annuals and summer perennial grasses, 1982-1983.
Allen, M. Friesner, D.; Joqari, F.; Mason, L. Baton Rouge? , La. : The Station. Annual progress report - Southeast Research Station, Louisiana Agricultural Experiment Station. 1983. p. 75-82. (NAL Call No.: DNAL S67.E22).

0173

Yield losses in spring barley caused by cereal aphids (Homoptera: Aphididae) in South Dakota.
JEENAI. Kieckhefer, R.W. Kantack, B.H. College Park, Md. : Entomological Society of America. Journal of economic entomology. June 1986. v. 79 (3). p. 749-752. Includes references. (NAL Call No.: DNAL 421 J822).

PLANT PRODUCTION - RANGE

0174

Effects of pasture management systems on cow-calf productivity on loessial soils in Northeast Louisiana.

LAXBA. Coombs, D.F. Bartleson, J.L.; Rogers, R.L.; Saxton, A.M.; Huffman, D.C.; Alison, M.W. Baton Rouge, La. : The Station. Bulletin - Louisiana Agricultural Experiment Station. Oct 1989. (815). 19 p. Includes references. (NAL Call No.: DNAL 100 L93 (1)).

0175

Forage establishment: weather effects on stubble vs. fallow and fall vs. spring seedling.

JRMGA. Hart, R.H. Dean, J.G. Denver, Colo. : Society for Range Management. Journal of range management. May 1986. v. 39 (3). p. 228-230. Includes references. (NAL Call No.: DNAL 60.18 J82).

0176

Mountain meadow (*Hordeum jubatum*, Weed control, Wyoming).

Alley, H.P. Vore, R.E.; Humburg, N.E. Laramie, Wyo., The Station. Research journal - University of Wyoming, Agricultural Experiment Station. Jan 1982. Jan 1982. (172). p. 19-27. (NAL Call No.: S131.E22).

PLANT BREEDING

0177

Aberrant ratio revisited.

Nelson, O.E. New York : Alan R. Liss. UCLA symposia on molecular and cellular biology. Paper presented at the "Symposium on Molecular and Cellular Biology of Plant Stress," April 15-21, 1984, Keystone, Colorado. 1985. v. 22. p. 1-12. Includes references. (NAL Call No.: DNAL QH506.U34).

0178

Adapted varieties.

Graham, W.D. Gooden, D.T. Clemson, S.C. : The Service. Circular - Clemson University, Cooperative Extension Service. In series analytic: Small grain production guidelines for South Carolina, 1988-89. Aug 1988. (463,rev.). p. 29-38. (NAL Call No.: DNAL 275.29 S08E).

0179

Agronomic evaluation of oat cultivars with substituted Avena fatua and A. sterilis cytoplasms.

CRPSAY. Rines, H.W. Halstead, R.P. Madison, Wis. : Crop Science Society of America. Genetic diversity in cytoplasms may potentially reduce the vulnerability of a crop to an epiphytotic and also increase variability available for trait improvement. The objectives of this study were to increase the cytoplasmic diversity available in cultivated oat Avena sativa L.) and to test effects of substituted wild oat cytoplasms on grain yield and other agronomic traits. Cytoplasm substitution lines were developed by five to six backcrosses of four oat cultivars into cytoplasms of five A. fatua L. and three A. A. sterilis wild oat accessions. Two to six lines of various alloplasmic combinations were compared to their respective euplasmic A. sativa recurrent parents in hill plot tests at two locations in 1985 and one location in 1986. The A. fatua and A. sterilis cytoplasms generally did not affect grain yield, aboveground biomass production, height, heading date, harvest index, or seed weight. One alloplasmic combination, among 24 tested, consistently yielded significantly less than its euplasmic A. sativa parent, but none yielded significantly more. The few instances where particular alloplasmic combinations differed from their euplasmic A. sativa parents may have resulted from residual nuclear contributions from the cytoplasm donor. The general lack of detrimental effects of substituted A. fatua and A. sterilis cytoplasms indicate that they may be useful for broadening the cytoplasmic diversity of oat. Crop science. Sept/Oct 1988. v. 28 (5). p. 805-809. Includes references. (NAL Call No.: DNAL 64.8 C883).

0180

Allelic relationships of genes for multinodosum mutants in barley (Hordeum vulgare L.).

Ahuja, S.L. Sethi, G.S. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Aug 31, 1987. v. 17. p. 63-67. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0181

Allelic relationships of genes for naked caryopsis mutants in barley (Hordeum vulgare L.).

Ahuja, S.L. Sethi, G.S. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Aug 31, 1987. v. 17. p. 58-62. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0182

Allelic relationships of genes for short-awned, awnless and blue aleurone characters in barley (Hordeum vulgare L.).

Ahuja, S.L. Sethi, G.S. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Aug 31, 1987. v. 17. p. 55-57. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0183

Allelism of genes in the M1-a locus (Barley, powdery mildew resistance).

Giese, H. Jensen, H.P.; Jorgensen, J.H. Fort Collins, Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1980. v. 10. p. 22-24. 3 ref. (NAL Call No.: QK495.G74B34).

0184

Aluminum tolerance of spring rye inbred lines.

Aniol, A. Hill, R.D.; Larter, E.N. Madison, Wis., Crop Science Society of America. Crop science. Mar/Apr 1980. v. 20 (2). p. 205-208. 11 ref. (NAL Call No.: 64.8 C883).

0185

Andre spring barley (Two-rowed, long-awned variety, performance, disease resistance).

Morrison, K.U. Ullrich, S.E.; Nilan, R.A.; Brauen, S.E.; Muir, C.E.; Reisenauer, P. Pullman, Wash. : The Service. Extension Bulletin - Washington State University, Cooperative Extension Service. Mar 1984. Mar 1984. (1249). 5 p. (NAL Call No.: 275.29 W27P).

0186

Assessment of barley accessions PI 382313, PI 382474, PI 382915, and PI 382976 for stem rust resistance.
CRPSAY. Jedel, P.E. Metcalfe, D.R.; Martens, J.W. Madison, Wis. : Crop Science Society of America. New sources of resistance to *Puccinia graminis* Pers. f. sp. *tritici* Erikss. and Henn. in barley (*Hordeum vulgare* L.) should be identified and incorporated into breeding programs to ensure the continued resistance of barley to this pathogen. The objective of this study was to identify resistance to stem rust in barley accessions from the USDA Wiebe collection and determine the relationship of this resistance to resistance conferred by the T gene. Four accessions, PI 382313, PI 382474, PI 382915, and PI 382976, were selected as having good field resistance at the Winnipeg Agriculture Canada Research Station, Glenlea rust nursery in 1986. The field resistance of these four accessions was reconfirmed in the 1987 rust nursery and was found to be as effective as, or more effective than, that conferred by T. Seedling and adult plant tests were conducted in controlled environments with six races (C5, C10, C17, C25, C35, and C53). In seedling tests, PI 382313 was resistant or moderately resistant to all six races. Both PI 382474 and PI 382915 were moderately susceptible to C35 and susceptible to the other five races. The PI 382976 was moderately resistant to C35 and C53, and moderately susceptible or susceptible to the other races. In adult plant tests, PI 382474 was resistant to C35 and PI 382313 was resistant to C17. Both PI 382915 and PI 382979 displayed little resistance as adults to the six races. Genetic studies with PI 382313 indicated that the field resistance of this accession was conferred by a single dominant gene that different from T. Crop science. Nov/Dec 1989. v. 29 (6). p. 1473-1477. Includes references. (NAL Call No.: DNAL 64.8 C883).

0187

Association of a mildew resistance gene JM1h in Hanna 906 with chromosome 6.
Hayashi, J. Heta, H. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Oct 10, 1985. v. 15. p. 46-47. (NAL Call No.: DNAL QK495.G74B34).

0188

Association of host cytoplasm with reaction to *Puccinia coronata* in progeny of crosses between wild and cultivated oats.
PLDRA. Simons, M.D. Robertson, L.D.; Frey, K.J. St. Paul, Minn. : American Phytopathological Society. Plant disease. Nov 1985. . v. 69 (11). p. 969-971. Includes 26 references. (NAL Call No.: DNAL 1.9 P69P).

0189

The attempted transfer of disease resistance from *Hordeum bulbosum* L., to *H. vulgare* L.
Pickering, R.A. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 5-8. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0190

Attempts to locate powdery mildew resistance gene M1-(La) to a barley chromosome.
Jensen, H.P. Helms Jorgensen, J.; Jensen, J. Fort Collins, Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1982. v. 12. p. 65-68. 3 ref. (NAL Call No.: QK495.G74B34).

0191

Barley breeding.
AGRYA. Anderson, M.K. Reinbergs, E. Madison, Wis., Soil Science Society of America. Agronomy. A series of monographs - American Society of Agronomy. Literature review. 1985. (26). p. 231-268. Includes references. (NAL Call No.: DNAL 4 AM392).

0192

Barley composite cross CC XXXV-A as a source of both specific and slow rusting resistance against leaf rust.
CRPSAY. Andres, M.W. Wilcoxson, R.D. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 273-275. Includes references. (NAL Call No.: DNAL 64.8 C883).

0193

Barley yellow dwarf symptom severity in oat affected by plant growth stage at infection and plot type.
CRPSAY. Goulart, L.R. Ohm, H.W.; Foster, J.E. Madison, Wis. : Crop Science Society of America. Lack of a standard method for evaluation of barley yellow dwarf (BYD) symptoms may result in inconsistency for genetic evaluation. The BYD symptoms were determined in oat, *Avena sativa* L., subsequent to virus infection at three plant growth stages and with two types of field plots. Five treatments: noninoculated; and inoculated at the two- to three-leaf (transplanted), three-leaf, four- to five-leaf, and stem elongation stages (direct seeded) were whole plots. Five oat cultivars were subplots. Two types of plots, hill and space planted, were whole units across the five cultivars. Seven traits were measured: BYD symptoms, plant height, heading date, grain yield and yield components. The plot types were not significantly different and there was a high

(PLANT BREEDING)

correlation between the types of plot for these characters. Cultivar rank was similar at all growth stages, indicating that sensitive cultivars may be severely damaged by BYDV infection at any stage of development. Small differences among cultivars for tolerance to BYDV were detected, subsequent to infection at the stem elongation stage under conditions of moderate temperatures and well distributed rainfall during the growing season. The plants that were transplanted after inoculation at the two- to three-leaf stage were more severely affected by BYDV than plants that were seeded in the field and inoculated at the three-leaf stage. Differences among cultivars were most distinct for BYD symptom, kernel weight, grain yield, number of kernels per panicle and plant height. Inoculation at the three-leaf stage resulted in maximum differences among cultivars and hill plots involved minimal effort for BYDV evaluation. Crop science. Nov/Dec 1989. v. 29 (6). p. 1412-1416. Includes references. (NAL Call No.: DNAL 64.8 C883).

0194

Betaine accumulation: metabolic pathways and genetics.

Hanson, A.D. Grumet, R. New York : Alan R. Liss. UCLA symposia on molecular and cellular biology. Paper presented at the "Symposium on Molecular and Cellular Biology of Plant Stress," April 15-21, 1984, Keystone, Colorado. 1985. v. 22. p. 71-92. ill. Includes references. (NAL Call No.: DNAL QH506.U34).

0195

Biotype C and E greenbugs: distribution in the Texas rolling plains and damage to four small grain varieties (Schizaphis graminum).

Puterka, G.J. Slosser, J.E.; Gilmore, E.C. College Station, Tex., Southwestern Entomological Society. The Southwestern entomologist. Mar 1982. v. 7 (1). p. 4-8. map. Includes 5 ref. (NAL Call No.: QL461.S65).

0196

Breeding cereal grains for resistance to witchweed.

Ramaiah, K.V. Boca Raton, Fla. : CRC Press, c1987-. Parasitic weeds in agriculture / editor, Lytton J. Musselman. Literature review. p. 227-242. Includes references. (NAL Call No.: DNAL SB611.P34).

0197

Breeding for pest resistance.

AGRYA. Sharp, E.L. Madison, Wis., Soil Science Society of America. Agronomy. A series of monographs - American Society of Agronomy. Literature review. 1985. (26). p. 313-333. Includes references. (NAL Call No.: DNAL 4 AM392).

0198

Breeding for preharvest sprouting tolerance.

Derera, N.F. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 111-128. ill. Includes references. (NAL Call No.: DNAL SB189.6.P73).

0199

Breeding oats for food and feed: conventional and new techniques and materials.

Burrows, V.D. St. Paul, Minn. : American Association of Cereal Chemists, c1986. Oats : chemistry and technology / edited by Francis H. Webster. p. 13-46. ill. Includes references. (NAL Call No.: DNAL TP435.O502).

0200

Breeding winter oats for the South.

AGUOAT. Stanton, T.R. Madison, Wis. : American Society of Agronomy. A nursery for the isolation and development of cold-resistant, high-yielding, and otherwise desirable strains of winter oats has been maintained at the Arlington Experiment Farm since 1907. A more or less standardized nursery system is used consisting of series of single 5-foot plant or head rows, duplicated 15-foot rows, and duplicated blocks or plats of three 15-foot rows each. Two strains of Winter Turf, C. I. Nos. 435-4 and 541-4, have been developed which during the 10-year period, 1916-25, inclusive, have outyielded the unselected stock by about 8 bushels to the acre. Definite winter types of Fulghum have been isolated and are being studied. In these forms the plants show a decided spreading or winter habit in early growth and have dark green, rather narrow leaves. The selections apparently are more cold-resistant, and are superior in yield to the original Fulghum. Several high-yielding strains of winter oats have been developed from a Winter Turf-Aurora case. In these strains the earliness and excellent kernel characters of the Aurora variety have been combined with the winter resistance of the Winter Turf oat to a highly satisfactory degree. Two of the most promising selections have been named Lee and Custis. The maintenance of varietal identity in such standard oats as fulghum and Red Rustproof is one of the most pressing practical problems in connection with the growing of oats in the South. Selected strains of these varieties are available of these varieties are available and should be substituted for the nondescript, so-called Red Rustproof oats which are grown in many districts of the South. The selection work with Fulghum was considerably expanded in the fall of 1925 at the Arlington Experiment Farm. In addition, 500 selections from Fulghum were sown at Tifton, Ga., for the development of strains especially adapted to the southern portion of the cotton belt. Numerous selections from hybrids between Fulghum and hardy winter varieties, such as Culberson. Agronomy journal. Sept 1926. v. 18 (9). p. 804-814. ill. (NAL Call No.: DNAL 4 AM34P).

0201

Can we lead the pathogen astray? (Breeding cereals for resistance to fungus diseases). Wolfe, M.S. Barrett, J.A. St. Paul, Minn., American Phytopathological Society. Plant disease. Feb 1980. v. 64 (2). p. 148-151, 153-155. ill. 13 ref. (NAL Call No.: 1.9 P69P).

0202

Canopy light and tiller mortality in spring barley.

Lauer, J.G. Simmons, S.R. Madison, Wis. : Crop Science Society of America. Mortality of tillers in barley (*Hordeum vulgare* L.) is an important developmental event affecting spike number and yield potential of the crop. It has traditionally been thought that increased shading of young tillers by the developing canopy initiates the premature senescence of tillers. A series of field experiments were conducted in 1982, 1983, and 1984 at St. Paul, MN on a Waukegan silt loam soil (fine-silty over sandy or sandy skeletal, mixed, Typic Hapludoll) to test this hypothesis. Degree of shading within the crop canopy was measured in relation to the position of tiller leaves during the vegetative phase of development for three genotypes. The decline in rate of leaf appearance on nonsurviving primary tillers, monitored as an early indicator of tiller senescence, was noted within 3 to 4 wk after crop emergence and before appreciable shading of tillers occurred. This indicates that tiller mortality was not initiated by lack of light. However, after main stem elongation began, tillers soon became heavily shaded suggesting the possibility that shading plays an auxiliary role in the senescence of barley tillers. We discuss the possibility that changes in light quality early in the growing season may be important for initiating the senescence of tillers. Crop science. Mar/Apr. 1989. v. 29 (2). p. 420-424. Includes references. (NAL Call No.: DNAL 64.8 C883).

0203

Characteristic cellular responses as expression of genes for resistance to Erysiphe graminis f. sp. hordei in barley (Hordeum vulgare).

Koga, H. PHYTA. Toyoda, H.; Mayama, S.; Shishiyama, J.; Hiura, U. St. Paul : American Phytopathological Society. Phytopathology. June 1983. v. 73 (6). p. 907-910. Includes references. (NAL Call No.: 464.8 P56).

0204

Chitinase locus (Chi 1) assigned to barley chromosome 1 (7H) by immunoblotting.

Hejgaard, J. Bjorn, S.E. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Aug 31, 1987. v. 17. p. 92-93. ill. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0205

Chromosome damage induced by artificial seed aging in barley. II. Types of chromosomal aberrations at first mitosis (Hordeum vulgare). Murata, M. Tsuchiya, T.; Roos, E.E. Chicago, Ill., University of Chicago Press. Botanical gazette. Mar 1982. v. 143 (1). p. 111-116. ill. 1 p. ref. (NAL Call No.: 450 B652).

0206

Cloning and nitrate induction of nitrate reductase mRNA.

PNASA. Cheng, C.L. Dewdney, J.; Kleinhofs, A.; Goodman, H.M. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. Sept 1986. v. 83 (18). p. 6825-6828. ill. Includes references. (NAL Call No.: DNAL 500 N21P).

0207

Common root rot of spring barley: relative susceptibility of six-rowed cultivars.

NDFRA. Stack, R.W. Fargo, N.D. : The Station. North Dakota farm research - North Dakota, Agricultural Experiment Station. July/Aug 1986. v. 44 (1). p. 5-7. ill. Includes references. (NAL Call No.: DNAL 100 N813B).

0208

A comparison of the effect of salt on polypeptide and translatable mRNAs in roots of a salt-tolerant and a salt-sensitive cultivar of barley.

PLPHA. Harkman, W.J. Fornari, C.S.; Tanaka, C.K. Rockville, Md. : American Society of Plant Physiologists. The effect of salt stress on polypeptide and mRNA levels in roots of two barley (*Hordeum vulgare* L.) cultivars differing in salt tolerance (cv CM 72, tolerant; cv Prato, sensitive) was analyzed using two-dimensional polyacrylamide gel electrophoresis. Preliminary experiments indicated that germination of Prato was inhibited significantly in the presence of NaCl, but growth of the surviving Prato seedlings was not substantially different from that of CM 72. Fluorographs of two-dimensional gels containing in vivo labeled polypeptides or in vitro translation products were computer analyzed to identify and quantitate changes that resulted when plants were grown in the presence of 200 millimolar NaCl for 6 days. The patterns of in vi vo labeled polypeptides and in vitro products of CM 72 and Prato were qualitatively the same. Salt caused quantitative changes in numerous polypeptides and translatable mRNAs, but, overall, the changes were relatively small. Salt did not induce the synthesis of unique polypeptides or translatable mRNAs and did not cause any to disappear. Because of the similarities of the two cultivars with respect to growth and polypeptide patterns and the slight changes in polypeptide and translation product levels

(PLANT BREEDING)

caused by salt, specific polypeptides or translatable mRNAs that are related to salt tolerance in barley could not be identified. Plant physiology. Aug 1989. v. 90 (4). p. 1444-1456. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0209

Compatibility of Lysiphlebus testaceipes (Hymenoptera: Braconidae) with greenbug (Homoptera: Aphididae) biotypes "C" and "E" reared on susceptible and resistant oat varieties (Schizaphis graminum).
Salto, C.E. EVETB. Eikenbary, R.D.; Starks, K.J. College Park : Entomological Society of America. Environmental entomology. Apr 1983. v. 12 (2). p. 603-604. Includes references. (NAL Call No.: QL461.E532).

0210

Coordinator's report: anthocyanin genes. Stock list of ant mutants kept at the Carlsberg laboratory.
Jende-Strid, B. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 74-79. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0211

Coordinator's report: chromosome 5.
Jensen, J. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 61-63. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0212

Coordinator's report: chromosome 5 (Hordeum, barley, hordein component controlled by locus).
Jensen, J. Fort Collins : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1983. v. 13. p. 94-97. Includes references. (NAL Call No.: QK495.G74B34).

0213

Coordinator's report: disease and pest resistance genes.
Jorgensen, J.H. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 65-69. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0214

Cytoplasmic effects on plant traits in interspecific matings of Avena (Cultivars, oats).
Robertson, L.D. Frey, K.J. Madison : Crop Science Society of America. Crop science. Jan/Feb 1984. v. 24 (1). p. 199-204. Includes references. (NAL Call No.: 64.8 C883).

0215

Cytoplasmic male sterility in barley. VIII. Lipoygenase activity and anther amino nitrogen in the msml-Rfm1a system.
Ahokas, H. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Jan 1982. v. 69 (1). p. 268-272. ill. 33 ref. (NAL Call No.: 450 P692).

0216

Designation of two new greenbug (Homoptera: Aphididae) biotypes G and H.
JEENAI. Puterka, G.J. Peters, D.C.; Kerns, D.L.; Slosser, J.E.; Bush, L.; Worrall, D.W.; McNew, R.W. Lanham, Md. : Entomological Society of America. Journal of economic entomology. Dec 1988. v. 81 (6). p. 1754-1759. Includes references. (NAL Call No.: DNAL 421 J822).

0217

Development and characterization of friable, embryogenic oat callus.
CRPSAY. Bregitzer, P. Somers, D.A.; Rines, H.W. Madison, Wis. : Crop Science Society of America. Friable, embryogenic (FE) callus cultures are desired for genetic manipulations at the cellular level. The goal of this research was to develop and characterize such cultures for oat (Avena spp.). Nonfriable (NF) oat callus derived from immature embryos was plated on a modified MS medium containing 60 g L-1 sucrose and no hormones. This treatment resulted in the development of distinct somatic embryos that germinated to form complete plants when placed on a modified MS medium containing 20 g L-1 sucrose and no hormones. Embryogenic sectors isolated from 26-wk-old NF callus were visually selected during repeated subculture on a modified MS medium containing 2 mg L-1, 2,4-dichlorophenoxyacetic acid (2,4-D) and 20 g L-1 sucrose to produce FE callus. Transfer of FE callus to a modified MS medium containing 60 g L-1 sucrose and no hormones induced the maturation of somatic embryos. Subsequent transfer of this FE callus to modified MS medium containing 20 g L-1 sucrose and no hormones allowed the germination of some of these embryos. Plants were regenerated from FE callus lines for more than 78 wk after FE callus establishment. Friable embryogenic callus also was initiated directly from immature embryos of three genotypes and seedling mesocotyls of two genotypes. Genotypic variation in this response was noted. The FE oat cultures represent an improved callus type

for in vitro genetic manipulation of oat. Crop science. May/June 1989. v. 29 (3). p. 798-803. ill. Includes references. (NAL Call No.: DNAL 64.8 C883).

0218

Development of a barley germ plasm resistant to the seed transmission of barley stripe mosaic virus (Abstract only).

Carroll, T.W. Zaske, S.K.; Hockett, E.A. St. Paul, Minn., American Phytopathological Society. Phytopathology. Aug 1981. v. 71 (8). p. 865. (NAL Call No.: 464.8 P56).

0219

Differential aluminum tolerance of winter barley varieties and selections in associated greenhouse and field experiments.

AGJOAT. Reid, D.A. Jones, G.D.; Armiger, W.H.; Foy, C.D.; Koch, E.J.; Starling, T.M. Madison, Wis. : American Society of Agronomy. Agronomy journal. Mar/Apr 1969. v. 61 (2). p. 218-222. ill. Includes references. (NAL Call No.: DNAL 4 AM34P).

0220

Differential mRNA transcription during salinity stress in barley.

PNASA. Ramagopal, S. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. Jan 1987. v. 84 (1). p. 94-98. ill. Includes references. (NAL Call No.: DNAL 500 N21P).

0221

Diploid-tetraploid comparisons in rye. III. Temperature effects on seedling development.

CRPSAY. Sullivan, B.P. Pfahler, P.L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 795-799. Includes 22 references. (NAL Call No.: DNAL 64.8 C883).

0222

Disease severity and grain yield in barley multilines with resistance to powdery mildew.

CRPSAY. Kolster, P. Munk, L.; Stolen, O. Madison, Wis. : Crop Science Society of America. Superiority of yield in cultivar mixtures may be ascribed to specific and nonspecific resistance against leaf pathogens and to differential reactions of components to environment. This study was conducted to investigate the significance of specific resistance on powdery mildew (*Erysiphe graminis* DC.: Fr. f. sp. *hordei*) severity and grain yield in barley (*Hordeum vulgare* L.) multilines and isolines, each carrying different genes for specific resistance to powdery mildew.

Multilines were composed of equal proportions of four or five isolines. In the multilines, disease levels were 50 to 80% lower than the mean level for the components grown in pure stands. The advantage of mixing increased with increasing mean level of resistance in the component isolines. Grain yield in both multilines and isolines increased with the mean level of resistance, with most resistant multilines producing higher grain yields than predicted by the mean level of the components in pure stands. The results indicate that components with higher levels of resistance gained the most benefit from utilizing them in a multiline. Crop science. Nov/Dec 1989. v. 29 (6). p. 1459-1463. Includes references. (NAL Call No.: DNAL 64.8 C883).

0223

Durable resistance of barley cultivars to the nematode *Heterodera avenae*.

Person-Dedryver, F. NASSD. New York : Plenum Press. NATO advanced study institutes series. Series A. Life sciences. 1983. v. 55. p. 207-210. Includes references. (NAL Call No.: QH301.N32).

0224

Effect of environment and genotype and their interaction on pathogenicity of *Ustilago hordei* (causing the covered smut of barley). I. Parasite-environment effects.

Emara, Y.A. Freake, G.W. Washington, D.C., American Genetic Association. The Journal of heredity. July/Aug 1981. v. 72 (4). p. 261-263. 15 ref. (NAL Call No.: 442.8 AM3).

0225

Effect of erect leaf angle on grain yield in barley.

CRPSAY. Tunglund, L. Chapko, L.B.; Wiersma, J.V.; Rasmusson, D.C. Madison, Wis. : Crop Science Society of America. Crop science. Jan/Feb 1987. v. 27 (1). p. 37-40. Includes references. (NAL Call No.: DNAL 64.8 C883).

0226

Effect of genetically and environmentally induced heading date differences on yield and adaptation of an isogenic barley pair.

CRPSAY. Smail, V.W. Eslick, R.F.; Hockett, E.A. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 889-893. Includes references. (NAL Call No.: DNAL 64.8 C883).

(PLANT BREEDING)

0227

The effect of genotypes on the feed value of barley straws (North Dakota).
Erickson, D.D. JANS. Meyer, D.W.; Foster, A.E.
Champaign : American Society of Animal Science.
Journal of animal science. Nov 1982. v. 55 (5).
p. 1015-1026. 15 ref. (NAL Call No.: 49 J82).

0228

Effect of host genotype unit area on epidemic development of crown rust following focal and general inoculations of mixtures of immune and susceptible oat plants.

PHYTA. Mundt, C.C. Leonard, K.J. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Oct 1985. v. 75 (10). p. 1141-1145. Includes 26 references. (NAL Call No.: DNAL 464.8 P56).

0229

Effect of planting dates on spring barley varieties, San Juan Branch Station, 1974-1979 (New Mexico, performance, yield).

Gregory, E.J. Arnold, R.N. Las Cruces : The Station. Research report - New Mexico, Agricultural Experiment Station. June 1982. June 1982. (478). 12 p. 8 ref. (NAL Call No.: 100 N465R).

0230

The effect of selection in pure-line oat work.
AGJOAT. Spragg, F.A. Madison, Wis. : American Society of Agronomy. Agronomy journal. 1912. v. 4. p. 81-83. (NAL Call No.: DNAL 4 AM34P).

0231

Effect of sexual and asexual reproduction on race abundance in cereal rust fungus populations (*Puccinia graminis* f. sp. *tritici*, *Puccinia coronata*, *Puccinia recondita*, genetic variation).

Groth, J.V. PHYTA. Roelfs, A.P. St. Paul : American Phytopathological Society. Phytopathology. Nov 1982. v. 72 (11). p. 1503-1507. 20 ref. (NAL Call No.: 464.8 P56).

0232

An effective method developing inversions on a specific chromosome.

Makino, T. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Aug 31, 1987. v. 17. p. 103-104. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0233

Effectiveness of utilizing various leaf characteristics as selection criteria in breeding for drought tolerance in barley / by Marvin L. Boerboom. -

Boerboom, Marvin L. Ann Arbor, Mich. University Microfilms International 1978. Thesis--North Dakota State University, 1977. Facsimile produced by microfilm-xerography. viii, 128 leaves. Bibliography: leaves 92-99. (NAL Call No.: DISS 77-29,178).

0234

The effects of benzyladenine, cycloheximide, and cordycepin on wilting-induced abscisic acid and proline accumulations and abscisic acid- and salt-induced proline accumulation in barley leaves.

PLPHA. Stewart, C.R. Voetberg, G.; Rayapati, P.J. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Nov 1986. v. 82 (3). p. 703-707. Includes references. (NAL Call No.: DNAL 450 P692).

0235

Effects of cycloheximide treatments on prophase and metaphase cells of a (2X) *Hordeum vulgare* X (2X) *Hordeum bulbosum* hybrid (Barley).

Wheatley, W.G. Kasha, K.J. Fort Collins, Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1982. v. 12. p. 72-74. 111. 2 ref. (NAL Call No.: QK495.G74B34).

0236

Effects of dried sewage sludge on forage production from barley genotypes in the Sonoran Desert.

Day, A.D. Thompson, R.K.; Swingle, R.S. Superior : University of Arizona. Desert plants. 1986. v. 8 (1). p. 17-19. Includes references. (NAL Call No.: DNAL QK938.D4D4).

0237

The effects of genes controlling barley leaf and sheath waxes on agronomic performance in irrigated and dryland environments (*Eceriferum*, mutation).

Baenziger, P.S. CRPSA. Wesenberg, D.M.; Sicher, R.C. Madison : Crop Science Society of America. Crop science. Jan/Feb 1983. v. 23 (1). p. 116-120. Includes references. (NAL Call No.: 64.8 C883).

0238

Effects of natural selection in advances generations of barley composite cross II (Hordeum vulgare, yield components).
 Hockett, E.A.CRPSA. Eslick, R.F.; Qualset, C.O.; Dubbs, A.L.; Stewart, V.R. Madison : Crop Science Society of America. Crop science. July/Aug 1982. v. 23 (4). p. 752-756. Includes references. (NAL Call No.: 64.8 C883).

0239

Effects of nitrogen fertilizer rate, seeding rate, and row spacing on semidwarf and conventional height spring oat.
 CRPSAY. Marshall, H.G. Kolb, F.L.; Roth, G.W. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1987. v. 27 (3). p. 572-575. Includes references. (NAL Call No.: DNAL 64.8 C883).

0240

Effects of plant density on the performance of 10 barley cultivars (Genotype-environment interaction, yield).
 Baker, R.J.CRPSA. Briggs, K.G. Madison : Crop Science Society of America. Crop science. Nov/Dec 1982. v. 22 (6). p. 1164-1167. 6 ref. (NAL Call No.: 64.8 C883).

0241

Effects of regional releases of resistant wheats on the population dynamics of the cereal leaf beetle (Coleoptera: Chrysomelidae) (Oulema melanopus).
 Lampert, E.P.AESAA. Haynes, D.L.; Sawyer, A.J.; Jokinen, D.P.; Wellso, S.G. College Park : The Society. Annals of the Entomological Society of America. Nov 1983. v. 76 (6). p. 972-980. Includes references. (NAL Call No.: 420 EN82).

0242

Erysiphe graminis resistance in the Hordeum murinum complex (Powdery mildew, wild barley, genetic aspects).
 Giles, B. Barrett, J. Fort Collins : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1983. v. 13. p. 78-82. Includes references. (NAL Call No.: QK495.G74E34).

0243

Evaluation of sources of resistance in oats to Puccinia coronata by use of many fungus cultures.
 Michel, L.J.PIAIA. Simons, M.D. Cedar Falls : The Academy. The Proceedings of the Iowa Academy of Science. Dec 1983. v. 90 (4). p. 144-146. Includes references. (NAL Call No.:

500 I093).

0244

Field crops.
 Arny, D.C. Dubuque, Iowa : Kendall/Hunt Pub. Co., c1986. With one foot in the furrow : a history of the first seventy-five years of the Department of Plant Pathology at the University of Wisconsin-Madison / edited by Paul H. Williams, Melissa Marosy. p. 164-174. ill. (NAL Call No.: DNAL SB732.54.U6W5).

0245

Field evaluation of seedling root length selection in oats (Avena sativa, drought stress).
 Barbour, N.W. Murphy, C.F. Madison : Crop Science Society of America. Crop science. Jan/Feb 1984. v. 24 (1). p. 165-169. Includes references. (NAL Call No.: 64.8 C883).

0246

For horses: Kelly.
 SDFHA. Leslie, J. Brookings, S.D. : The Station. South Dakota farm and home research - South Dakota, Agricultural Experiment Station. 1984. v. 35 (2). p. 14. ill. (NAL Call No.: DNAL 100 S0825).

0247

Generation mean analysis of inheritance of resistance to barley yellow dwarf in crosses involving bates spring oats.
 PLDRA. Gellner, J.L. Sechler, D.T. St. Paul, Minn. : American Phytopathological Society. Plant disease. Aug 1986. v. 70 (8). p. 795-797. Includes references. (NAL Call No.: DNAL 1.9 P69P).

0248

Genes conditioning resistance of Hordeum spontaneum (wild relative of barley) to Erysiphe graminis f. sp. hordei (barley powdery mildew).
 Moseman, J.G. Baenziger, P.S.; Kilpatrick, R.A. Madison, Wis., Crop Science Society of America. Crop science. Mar/Apr 1981. v. 21 (2). p. 229-232. 24 ref. (NAL Call No.: 64.8 C883).

0249

Genetic analysis of a mutant resistant to barley yellow mosaic virus.
 Ukai, Y. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1984. v. 14. p. 31-33. Includes references. (NAL Call No.:

(PLANT BREEDING)

QK495.G74B34).

0250

Genetic analysis of changes in scald resistance in barley composite cross V (*Rhynchosporium secalis*, *Hordeum vulgare*).

Jackson, L.F. Webster, R.K.; Allard, R.W.; Kahler, A.L. St. Paul, Minn., American Phytopathological Society. *Phytopathology*. Aug 1982. v. 72 (8). p. 1069-1072. 16 ref. (NAL Call No.: 464.8 P56).

0251

Genetic analysis of resistance to barley yellow dwarf virus in hybrids between *Avena sativa* 'Lamar' and virus-resistant lines of *Avena sterilis* (Oats).

Landry, B. Comeau, A.; Minvielle, F.; St.Pierre, C.A. Madison, Wis. : Crop Science Society of America. *Crop science*. Mar/Apr 1984. v. 24 (2). p. 337-340. ill. Includes references. (NAL Call No.: 64.8 C883).

0252

Genetic and temperature effects on rye germination and seedling development.

Sullivan, B.P. Peahler, P.L. S.I. : The Society. *Proceedings - Soil and Crop Science Society of Florida*. 1985. v. 44. p. 205-208. Includes 10 references. (NAL Call No.: DNAL 56.9 S032).

0253

Genetic control of ethirimol resistance in a natural population of *Erysiphe graminis* f.sp. *hordei* (Barley powdery mildew).

Hollomon, D.W. Laramie, The Station. *Science monograph - University of Wyoming, Agricultural Experiment Station*. May 1981. v. 71 (5). p. 536-540. 20 ref. (NAL Call No.: S131.E2).

0254

Genetic control of glycinebetaine level in barley.

CRPSAY. Grument, R. Isleib, T.G.; Hanson, A.D. Madison, Wis. : Crop Science Society of America. *Crop science*. July/Aug 1985. v. 25 (4). p. 618-623. ill. Includes references. (NAL Call No.: DNAL 64.8 C883).

0255

Genetic dissection of a powdery mildew resistance gene in barley.

Jorgensen, J.H. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. *Barley genetics newsletter*. Aug 31, 1987. v. 17. p. 99-100. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0256

Genetic factors governing resistance and susceptibility of oats to *Puccinia coronata* Corda var. *avenae*, F. and L. Race 57 /by Verne C. Finkner.

Finkner, Verne C. Ames, Iowa : Agricultural Experiment Station, Iowa State College, 1954. p. 1040-1063. 1 leaf of plates : ill. (1 col.); 22 cm. Bibliography: p. 1063. (NAL Call No.: DNAL 100 Io9 no.411).

0257

Genetic studies of resistance to barley yellow mosaic virus of German winter barley cultivars.

Friedt, W. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. *Barley genetics newsletter*. Oct 10, 1985. v. 15. p. 58-61. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0258

Genetic variability for herbicide reaction in plant populations (Wild oat, *Avena barbata*, *Avena fatua*, *godetia*, *Clarkia williamsonii*, toxicity, Sierra Nevada in California).

Price, S.C. WEESA. Hill, J.E.; Allard, R.W. Champaign : Weed Science Society of America. *Weed science*. Sept 1983. v. 31 (5). p. 652-657. ill. Includes references. (NAL Call No.: 79.8 W41).

0259

Genetic variability for pathogenicity, isozyme, ribosomal DNA and colony color variants in populations of *Rhynchosporium secalis*.

GENTA. McDermott, J.M. McDonald, B.A.; Allard, R.W.; Webster, R.K. Baltimore, Md. : Genetics Society of America. Samples of *Rhynchosporium secalis* were collected from two experimental barley populations known to carry a diverse array of alleles for resistance to this fungal pathogen. Classification of 163 isolates for four putative isozyme systems, a colony color dimorphism and 20 ribosomal DNA restriction fragment length variants revealed 49 different multilocus phenotypes (haplotypes). The six most common haplotypes differed significantly in pathogenicity. Genetic analyses of the data indicated that effective population sizes of the fungus were very large, that the effects of genetic drift were small, and that negligible recombination occurred in the populations

studied. Frequency dependent selection was suggested as an explanation for the maintenance of variation in pathogenicity in the fungus. Genetics. July 1989. v. 122 (3). p. 561-565. ill. Includes references. (NAL Call No.: DNAL 442.8 G28).

0260

Genotype X year interaction for length and rate of grain filling in oats (Environmental stresses).

Wych, R.D. McGraw, R.L.; Stuthman, D.D. Madison, Crop Science Society of America. Crop science. Sept/Oct 1982. v. 22 (5). p. 1025-1028. 20 ref. (NAL Call No.: 64.8 C883).

0261

Genotypic variability in response of oat to delayed sowing.

AGJ0AT. Baltenberger, D.C.C. Frey, K.J. Madison, Wis. : American Society of Agronomy. Agronomy journal. Sept/Oct 1987. v. 79 (5). p. 813-816. Includes references. (NAL Call No.: DNAL 4 AM34P).

0262

Genotypic variation for glycinebetaine accumulation by cultivated and wild barley in relation to water stress.

Ladyman, J.A.R. CRPSA. Ditz, K.M.; Grumet, R.; Hanson, A.D. Madison : Crop Science Society of America. Crop science. May/June 1983. v. 23 (3). p. 465-468. ill. Includes references. (NAL Call No.: 64.8 C883).

0263

Geographical distribution of diazinon sensitive varieties in barley.

Takeda, K. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Aug 31, 1987. v. 17. p. 90-93. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0264

Grain-filling duration and yield in spring barley.

CRPSAY. Metzger, D.D. Czaplewski, S.J.; Rasmusson, D.C. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1984. v. 24 (6). p. 1101-1105. Includes 11 references. (NAL Call No.: DNAL 64.8 C883).

0265

Gramine (indole alkaloid) in barley forage--effects of genotype and environment.

Hanson, A.D. Traynor, P.L.; Ditz, K.M.; Reicosky, D.A. Madison, Wis., Crop Science Society of America. Crop science. Sept/Oct 1981. v. 21 (5). p. 726-730. ill. 21 ref. (NAL Call No.: 64.8 C883).

0266

Greenbug (Homoptera: Aphididae) antibiosis tests in growth chambers: design of experiments and optimum sample size.

EVELEX. Inayatullah, C. Webster, J.A.; Nguyen, H.T. College Park, Md. : Entomological Society of America. Environmental entomology. June 1986. v. 15 (3). p. 581-584. Includes references. (NAL Call No.: DNAL QL461.E532).

0267

Growth and yield of barley isopopulations differing in solute potential.

CRPSAY. Grumet, R. Albrechtsen, R.S.; Hanson, A.D. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1987. v. 27 (5). p. 991-995. Includes references. (NAL Call No.: DNAL 64.8 C883).

0268

Half-seed determination of hordeins associated with known M1-a alleles conferring race-specific resistance to barley powdery mildew (Erysiphe graminis f. sp. hordei).

Hash, C.T. Jr. Blake, T.K. Fort Collins, Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1981. v. 11. p. 74-76. ill. 4 ref. (NAL Call No.: QK495.G74B34).

0269

Heterodera avenae: virulence and resistance.

NASSD. Andersen, K. Andersen, S. New York, N.Y. : Plenum Press. NATO advanced science institutes series : Series A : Life sciences. In the series analytic: Cyst nematodes / edited by F. Lamberti and C.E. Taylor. 1986. v. 121. p. 277-285. Includes references. (NAL Call No.: DNAL QH301.N32).

0270

Histology of the relation between minor and major genes for resistance of barley to leaf rust (Puccinia hordei on Hordeum vulgare).

Niks, R.E. PHYTA. Kuiper, H.J. St. Paul : American Phytopathological Society. Phytopathology. Jan 1983. v. 73 (1). p. 55-59. 25 ref. (NAL Call No.: 464.8 P56).

(PLANT BREEDING)

0271

Hordein mutants for barley Hrd loci.
Netsvetayev, V.P. Sozinov, A.A. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Oct 10, 1985. v. 15. p. 51-55. ill. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0272

Hormonal regulation of alpha-amylase gene transcription in wild oat (*Avena fatua* L.) aleurone protoplasts.
PLPHA. Zwar, J.A. Hookey, R. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Feb 1986. v. 80 (2). p. 459-463. ill. Includes 25 references. (NAL Call No.: DNAL 450 P692).

0273

Hybrid barley research: history and potential.
PMASA. Hockett, E.A. S.L. : The Academy. Proceedings of the Montana Academy of Sciences. Literature review. 1987. v. 47. p. 27-35. Includes references. (NAL Call No.: DNAL 500 M762).

0274

Identification of three new loci which control male sterility of barley.
Frankowiak, J.D. Hockett, E.A. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 11-13. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0275

In vitro culture of barley: a method to study *Rhynchosporium secalis* disease and select plants resistant to the toxin, Rhynchosporoside (*Rhynchosporium secalis*, *Hordeum vulgare*).
Branchard, M. New York : Praeger, 1982. Variability in plants regenerated from tissue culture / edited by E.D. Earle, Y. Demarly. p. 343-350. ill. Includes references. (NAL Call No.: QK840.V37).

0276

In vitro selection for resistance.
Foroughi-Wehr, B. Friedt, W.; Schuchmann, R.; Kohler, F.; Wenzel, G. Hingham, Mass. : Martinus Nijhoff Publishers. Advances in agricultural biotechnology. 1986. (20). p. 35-44. ill. Includes references. (NAL Call No.: DNAL S494.5.B563A39).

0277

Individual crown selection for resistance to freezing stress in winter oats (Germplasm, Cold hardiness).
Marshall, H.G. Kolb, F.L. Madison, Wis., Crop Science Society of America. Crop science. May/June 1982. v. 22 (3). p. 506-510. ill. 8 ref. (NAL Call No.: 64.8 C883).

0278

Induction of alcohol dehydrogenase and lactate dehydrogenase in hypoxically induced barley.
PLPHA. Good, A.G. Crosby, W.L. Rockville, Md. : American Society of Plant Physiologists. In barley (*Hordeum vulgare* L.), alcohol dehydrogenase (ADH) and lactate dehydrogenase (LDH) are induced by anaerobiosis in both aleurone layers and roots. Under aerobic conditions, developing seeds of cv Himalaya accumulate ADH activity, which survives seed drying and rehydration. This activity consists almost entirely of the ADH1 homodimer. Activity of LDH also increases during seed development, but the level of activity in dry or rehydrated seeds is very low, indicating that this enzyme may not be involved in anaerobic glycolysis during the initial stages of germination. In contrast to ADH, the LDH isozymes present in developing seeds are similar to those found in uninduced and induced roots. Developmental expression of ADH and LDH was monitored from 0 to 24 days postgermination. Neither activity was induced to any extent in the germinating seeds; however, both enzymes were highly induced by anoxia in root tissue during development. Based on gel electrophoresis, this increase in activity results from the differential expression of different Adh and Ldh genes in root tissue. The changes in ADH and LDH activity levels were matched by changes in the amount of these particular proteins, indicating that the increase in activity results from de novo synthesis of these two proteins. The level of inducible LDH activity in an ADH1- mutant was not found to differ from cv Himalaya. We suggest that although the ADH- plants are more susceptible to flooding, they are not capable of responding to the lack of ADH1 activity by increasing the amount of LDH activity in root tissue. Plant physiology. July 1989. v. 90 (3). p. 860-866. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0279

Induction of lactate dehydrogenase isozymes by oxygen deficit in barley root tissue.
PLPHA. Hoffman, N.E. Bent, A.F.; Hanson, A.D. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Nov 1986. v. 82 (3). p. 658-663. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0280

Induction of resistance to *Erysiphe graminis* f. sp. *hordei* in near-isogenic barley lines.
PHYTA. Cho, B.H. Smedegaard-Petersen, V. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Mar 1986. v. 76 (3). p. 301-305. ill. Includes 16 references. (NAL Call No.: DNAL 464.8 P56).

0281

The influence of ammonium and chloride on potassium and nitrate absorption by barley roots depends on time of exposure and cultivar.
PLPHA. Bloom, A.J. Finazzo, J. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1986. v. 81 (1). p. 67-69. Includes 22 references. (NAL Call No.: DNAL 450 P692).

0282

Influence of genotype and environment on kernel discoloration of Midwestern malting barley.
PLDRA. Miles, M.R. Wilcoxson, R.D.; Rasmusson, D.C.; Wiersma, J.; Warnes, D. St. Paul, Minn. : American Phytopathological Society. Plant disease. June 1987. v. 71 (6). p. 500-504. Includes references. (NAL Call No.: DNAL 1.9 P69P).

0283

Influence of weed control treatments on soybean cultivars in an oat-soybean rotation.
AGJDAT. Burnside, O.C. Moomaw, R.S. Madison, Wis. : American Society of Agronomy. Agronomy journal. Nov/Dec 1984. v. 76 (6). p. 887-890. Includes 13 references. (NAL Call No.: DNAL 4 AM34P).

0284

Inheritance of a second source of greenbug resistance in barley.
CRPSAY. Merkle, O.G. Webster, J.A.; Morgan, G.H. Madison, Wis. : Crop Science Society of America. Genetic studies were conducted to determine the inheritance of resistance to biotype E greenbug, *Schizaphis graminum* (Rondani), in PI 426756, a barley (*Hordeum vulgare* L.) line previously identified as resistant to this pest. An additional objective was to determine whether the genes governing resistance in PI 426756 and 'Post' are allelic or nonallelic. Results from greenbug-infested F2 and F3 seedlings in the greenhouse indicated that the resistance in each line is governed by a single dominant gene, and that these two genes are nonallelic and independent. The discovery of another gene for greenbug resistance is important because greenbug resistance in all current U.S. commercial barley cultivars appears to trace back to the same single dominant gene present in Post. The

gene symbol Rsg2b was assigned to this gene; whereas, the gene symbol formerly designated Grb was modified to Rsg1a. Crop science. Mar/Apr 1987. v. 27 (2). p. 241-243. Includes references. (NAL Call No.: DNAL 64.8 C883).

0285

Inheritance of blast resistance in two-rowed barley.
PLDIDE. Yaegashi, H. St. Paul, Minn. : American Phytopathological Society. Plant disease. July 1988. v. 72 (7). p. 608-610. ill. Includes references. (NAL Call No.: DNAL 1.9 P69P).

0286

Inheritance of dwarfness in three oat crosses and relationship of height to panicle and culm length (Lodging resistance).
Marshall, H.G. Murphy, C.F. Madison, Wis., Crop Science Society of America. Crop science. Mar/Apr 1981. v. 21 (2). p. 335-338. 15 ref. (NAL Call No.: 64.8 C883).

0287

Inheritance of resistance in oats to two biotypes of the greenbug (*Schizaphis graminum*).
Boozaya-Angoon, D. Starks, K.J.; Edwards, L.H.; Pass, H. College Park, Md., Entomological Society of America. Environmental entomology. Aug 1981. v. 10 (4). p. 557-559. 6 ref. (NAL Call No.: QL461.E532).

0288

Inheritance of resistance to barley leaf stripe.
Skou, J.P. Haahr, V. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 50-52. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0289

Inheritance of resistance to *Puccinia graminis* f. sp. *secalis* in barley (Breeding for disease resistance).
Steffenson, B.J. Wilcoxson, R.D.; Roelfs, A.P. St. Paul, Minn. : American Phytopathological Society. Plant disease. Sept 1984. v. 68 (9). p. 762-763. Includes 9 references. (NAL Call No.: 1.9 P69P).

(PLANT BREEDING)

0290

The inheritance of smut resistance in crosses of certain varieties of oats.

AGJOAT. Barney, A.F. Madison, Wis. : American Society of Agronomy. The data presented indicate that some varieties of oats contain one factor pair for resistance to loose smut, that other varieties contain two independent factor pairs, while in other varieties three factor pairs may be concerned with resistance. Selection within seventy families from two crosses indicated that resistant families can be isolated in the second generation which breed true to that characteristic thru the F3, F4 and F5 generations. Likewise, highly susceptible families can be isolated which breed true. Such results show the possibility of obtaining desirable types of smut resistant oats from crosses between resistant and susceptible varieties. The accompanying illustrations (Figures 1, 2, 3 and 4) are presented to indicate the response to smut exposure exhibited by two resistant and two susceptible families. Agronomy journal. Apr 1924. v. 16 (4). p. 283-291. (NAL Call No.: DNAL 4 AM34P).

0291

Inheritance of subcrown internode length in a winter barley cross (Winter hardiness, optimum depth of crown formation).

Dofing, S.M. Schmidt, J.W. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1984. v. 24 (4). p. 692-694. ill. Includes references. (NAL Call No.: 64.8 C883).

0292

Interrelationships among three major loci controlling heading date of spring barley when grown under short daylengths.

CRPSAY. Gallagher, L.W. Belhadri, M.; Zahour, A. Madison, Wis. : Crop Science Society of America. Early maturing cultivars of barley (*Hordeum vulgare* L.) would enhance yield stability in lowland, low rainfall areas of Morocco. The objective of this work was to study the inheritance of heading date in crosses involving six spring barleys developed in the Sacramento Valley of California, USA and the Yaqui Valley of Sonora, Mexico. Data were gathered under short daylengths at 32 degrees C 15' N Lat with December sowing in Morocco. Frequency distributions of parental, F1, F2, and backcross generations indicated a three-locus model with recessive and duplicate dominant epistasis accounting for most of the phenotypic variability. Recessive gene interaction resulted in plants that were about 19 to 35 days earlier than later counterparts with the dominant allele present. Among the later genotypes, the presence of a dominant allele at either of the duplicate loci in the genotypes not homozygous for extreme earliness resulted in increased earliness of 5 to 14 days over the duplicate double recessive depending upon modifiers. A genotype for each parent was suggested; however, the three-locus model did

not account for the 4- to 6-day differences between two pairs of parents. Apparently, modifying loci or different alleles were present in some parents. Crop science. Mar/Apr 1987. v. 27 (2). p. 155-160. Includes references. (NAL Call No.: DNAL 64.8 C883).

0293

List of barley genetic stocks.--Supplementary list No. 1 (to Masterlist of Barley Genes): genes for reaction to *Erysiphe graminis hordei* (Powdery mildew).

Sogaard, B. Jorgensen, J.H. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Includes addresses for sources of genetic stock seeds. Aug 31, 1987. v. 17. p. 120-134. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0294

Localization of induced genes for powdery mildew resistance.

Robbelen, G. Heun, M. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Oct 10, 1985. v. 15. p. 4-7. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0295

Low ribonuclease I activity prior to cold acclimation in freeze selected winter barley.

CRPSAY. Kenefick, D.G. Blake, T.K. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1986. v. 26 (6). p. 1099-1103. Includes 14 references. (NAL Call No.: DNAL 64.8 C883).

0296

Maintenance of low Cl⁻ concentrations in mesophyll cells of leaf blades of barley seedlings exposed to salt stress.

PLPHA. Huang, C.X. Van Steveninck, R.F.M. Rockville, Md. : American Society of Plant Physiologists. The concentrations of vacuolar Na⁺ and Cl⁻ in the epidermal and mesophyll cells of the leaf blade and sheath of *Hordeum vulgare* seedlings (cv California Mariout and Clipper) were measured by means of quantitative electron probe X-ray microanalysis. A preferential accumulation of Cl⁻ in vacuoles of epidermal cells in both blade and sheath and a low level in mesophyll cells of the blade were evident in plants grown in full strength Johnson solution. The concentration of Cl⁻ in the mesophyll cells of the blade remained at a low level after exposure to 50 or 100 millimolar NaCl for 1 day or to 50 millimolar for 4 days, while at the same time the concentration of Cl⁻ in the epidermis and mesophyll of the sheath showed a dramatic increase. Clipper generally contained more Cl⁻ in the mesophyll cells of the blade than

California Mariout. A greater accumulation of Na^+ in the mesophyll of the sheath relative to that of the blade was only apparent after treatment with 100 millimolar NaCl for 1 day or 50 millimolar for 4 days. These results confirm the suggestion that sheath tissue is capable of accumulating excess Cl^- (and to a lesser extent Na^+) and suggest that the site of regulation of Cl^- concentration in the barley leaf is located in the mesophyll cells of the blade. Plant physiology. Aug 1989. v. 90 (4). p. 1440-1443. Includes references. (NAL Call No.: DNAL 450 P692).

0297

Male sterile facilitated recurrent selection populations for developing broad-based resistance by major and minor effect genes (Barley, genetic manipulation).
Sharp, E.I. Bockelman, H.E.; Eslick, R.F. Minneapolis, Minn. : Published for the Congress by Burgess Pub., c1981. Proceedings of symposia : IX International Congress of Plant Protection, Washington, D.C., U.S.A., August 5-11, 1979 / editor, Thor Kommedahl. p. 193-194. (NAL Call No.: SB951.I5 1979).

0296

Marker genes for cereal cyst nematode resistance.
Ellis, S.E. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 249-250. (NAL Call No.: DNAL SB191.B2A9 1987).

0299

Mass rearing the bird cherry oat aphid, *Rhopalosiphum padi* (L.).
PIACA. Kudagamage, C. Foster, J.E. Indianapolis, Ind. : The Academy. Proceedings of the Indiana Academy of Science. Includes abstract. 1985. v. 94. p. 304. (NAL Call No.: DNAL 500 IN2).

0300

Measured crop performance: small grain, 1984.
NUSRA. Bowman, D.T. Raleigh, N.C., The Station. Research report - Dept. of Crop Science, North Carolina State University, Agricultural Experiment Station. Includes statistical data. July 1984. (96). 26 p. maps. (NAL Call No.: DNAL 100 N8122).

0301

Measured crop performance: small grain, 1985.
NUSRA. Bowman, D.T. Raleigh, N.C., The Station. Research report - Dept. of Crop Science, North Carolina State University, Agricultural Experiment Station. Includes statistical data. July 1985. (101). 29 p. maps. (NAL Call No.: DNAL 100 N8122).

0302

Measuring disease progress in pure and mixed stands of plant cultivars.
PHYTA. Gumpert, F.M. St. Paul, Minn. : American Phytopathological Society. A measure is presented that quantifies disease progress of fungal pathogens during the exponential phase of the epidemic. This measure, referred to as rate of disease increase (p), represents the factor by which the number of infection units is multiplied from one day to another after the initial infection waves have damped. The rate of disease increase depends on four pathotype/cultivar-specific parameters (latent period, infectious period, infection efficiency, spore production rate) and two nonspecific parameters (deposition frequency and autodeposition frequency). Three applications are given. First, it is used to determine the levels of partial resistance of barley cultivars to leaf rust. The calculated p values were highly correlated both with total spore production per unit leaf area ($r = 0.97$) and with disease score in the field ($r = 0.81$). The rate of disease increase is particularly useful if the epidemiological parameters display compensatory effects. The second application deals with the long-term composition of pathotype mixtures. The pathotype with the largest p value will predominate in the long run. This is again exemplified by using barley leaf rust data. The third application concerns disease control strategies. A host stand should be composed so that the corresponding predominant pathotype has a p value smaller than the prevailing pathotype of any other composition. This concept does not presuppose selection against unnecessary genes for virulence. A condition is given under which a cultivar mixture may be more beneficial than each of its components grown in pure stands, and this condition is illustrated by a simple example of two pathotypes and two cultivars. Phytopathology. Sept 1989. v. 79 (9). p. 969-973. Includes references. (NAL Call No.: DNAL 464.8 P56).

0303

Mechanical mass selection methods for improvement of oat groat percentage.
CRPSAY. Souza, E.J. Sorrells, M.E. Madison, Wis. : Crop Science Society of America. Three methods of mechanical mass selection for seed density, aspiration (ASP), gravity table selection (GT), and gravity table selection with a speeded polishing pretreatment (P-GT), were evaluated and compared for effectiveness of increasing caryopsis (groat) percentage and

(PLANT BREEDING)

test weight in oat (*Avena sativa* L.). Selection using these methods in two heterogeneous F5 populations resulted in absolute differences in the groat percentage between high and low density treatments of 0.3 and 0.5% for ASP and P-GT, respectively. The GT selection resulted in a lower groat percentage in the high density treatment. Selection methods interacted with populations and seed size classes. These interactions were due to different levels of variation in the populations for seed size and shape. Two cycles of selection for high groat percentage beginning in F3 segregating bulk populations showed little overall improvement. Selection increased the number of fertile florets per spikelet. This increase in fertility was due primarily to a 22% increase in tertiary florets per panicle per cycle of selection. Tertiary floret number exhibited a negative genotypic correlation with groat percentage, which limited improvement for groat percentage. There also were significant changes in plant height, days to flowering, and harvest index in these experiments with direction and magnitude dependent on the individual population. Effectiveness of mechanical mass selection for groat percentage could be increased by avoiding selection for characters such as tertiary kernels which have undesirable correlated effects on milling quality. *Crop science*. July/Aug 1988. v. 28 (4). p. 618-623. Includes references. (NAL Call No.: DNAL 64.8 C883).

0304

A method for rating freezing inhibitor activity of barley genotypes using seed and crown tissue mucilages (Winter hardiness).

Olien, C.R. AR-NC. Kuhna, A.S. Madison, Wis., *Crop Science Society of America. Crop science*. July/Aug 1980. v. 20 (4). p. 537-539. ill. 9 ref. (NAL Call No.: 64.8 C883).

0305

Methods in breeding cereals for rust resistance.

AGJOAT. Johnson, E.C. Madison, Wis. : American Society of Agronomy. *Agronomy journal*. 1910. v. 2. p. 76-80. (NAL Call No.: DNAL 4 AM34P).

0306

Metribuzin absorption and translocation in two barley (*Hordeum vulgare*) cultivars.

WEESA6. Gawronski, S. Haderlie, L.C.; Stark, J.C. Champaign, Ill. : Weed Science Society of America. *Weed science*. July 1986. v. 34 (4). p. 491-495. ill. Includes 13 references. (NAL Call No.: DNAL 79.8 W41).

0307

Models explaining the specificity and durability of host resistance derived from the observations on the barley-Puccinia hordei system.

Parlevliet, J.E. NASSD. New York : Plenum Press. NATO advanced study institutes series. Series A. Life sciences. 1983. v. 55. p. 57-80. Includes references. (NAL Call No.: QH301.N32).

0308

Molecular and cellular biology associated with endosperm mobilization in germinating cereal grains.

Fincher, G.B. Palo Alto, Calif. : Annual Reviews, Inc. Annual review of plant physiology and plant molecular biology. Literature review. 1989. v. 40. p. 305-346. ill. Includes references. (NAL Call No.: DNAL QK1.A57).

0309

Molecular biology of salinity stress: preliminary studies and perspectives.

Ramagopal, S. New York : Plenum Press, c1987. Tailoring genes for crop improvement : an agricultural perspective / edited by George Bruening ... et al. . p. 111-119. ill. Includes references. (NAL Call No.: DNAL SB123.57.C66 1986).

0310

Molecular cloning and sequence of cDNA encoding the plasma membrane proton pump (H⁺-ATPase) of *Arabidopsis thaliana*.

PNASA. Harper, J.F. Surowy, T.K.; Sussman, M.R. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. Feb 1989. v. 86 (4). p. 1234-1238. ill. Includes references. (NAL Call No.: DNAL 500 N21P).

0311

The morphological and physiological response of slender oat (*Avena barbata*) to the herbicides barban and difenzoquat.

WEESA6. Price, S.C. Hill, J.E.; Allard, R.W. Champaign, Ill. : Weed Science Society of America. *Weed science*. Jan 1988. v. 36 (1). p. 60-69. Includes references. (NAL Call No.: DNAL 79.8 W41).

0312

Morphological and physiological variation in wild oat (*Avena fatua*, dormancy, herbicide resistance, varieties, weeds).

Miller, S.D. Nalewaja, J.D.; Mulder, C.E.G. Madison, Wis., American Society of Agronomy. *Agronomy journal*. Sept/Oct 1982. v. 74 (5). p.

771-775. ill. 11 ref. (NAL Call No.: 4 AM34P).

0313

Morphological changes associated with three cycles of recurrent selection for grain yield improvement in oat.

CRPSAY. Bregitzer, P.P. Stuthman, D.D.; McGraw, R.L.; Payne, T.S. Madison, Wis. : Crop Science Society of America. To assess the potential for improving grain yield is via indirect morphological trait selection, the original and third cycle parents from a recurrent selection program for grain yield in oat (*Avena sativa* L.) were compared. Evaluation was based on hill plots grown in three field environments. This selection increased grain yield 13.5% and total dry matter 15.9%; thus, there was no increase in harvest index. All morphological structures measured increased in size. Grain yield was not closely or consistently related to any individual morphological trait, nor to the photosynthetic area of the flag leaf and panicle. Analysis of allometric relationships (the size of one plant structure relative to another) indicated that the size of morphological structures within a genotype tended to be proportional, but variability in allometric relationships was present. Therefore, selection for altered allometric relationships may be successful, but construction of an ideotype based on certain levels of expression of individual morphological traits, or photosynthetic areas, is not likely to be effective in increasing grain yield. Crop science. Mar/Apr 1987. v. 27 (2). p. 165-168. Includes references. (NAL Call No.: DNAL 64.8 C883).

0314

Multiline breeding (Cereal varieties, disease resistance).

Frey, K.J. New York : Academic Press, 1982. Plant improvement and somatic cell genetics / edited by I.K. Vasil, W.R. Scowcroft, K.J. Frey. p. 43-71. ill. Includes references. (NAL Call No.: SB123.P563).

0315

Multiple inoculation technique for evaluating resistance of single barley seedlings to three fungi (*Rhynchosporium secalis*, *Puccinia hordei*, *Erysiphe graminis hordei*, breeding for disease resistance).

Kilpatrick, R.A. Baenziger, P.S.; Moseman, J.G. St. Paul, Minn., American Phytopathological Society. Plant disease. June 1981. v. 65 (6). p. 504-506. 8 ref. (NAL Call No.: 1.9 P69P).

0316

Mutant selection.

Flick, C.E. Evans, D.A. New York : Macmillan Publishing Company. Plant-microbe interactions : molecular and genetic perspectives. 1984. v. 1. p. 168-192. Includes references. (NAL Call No.: DNAL QR351.P53).

0317

Natural selection in a doubled-haploid mixture and a composite cross of barley.

CRPSAY. Patel, J.D. Reinbergs, E.; Mather, D.E.; Choo, T.M.; Sterling, J.D.E. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1987. v. 27 (3). p. 474-479. Includes references. (NAL Call No.: DNAL 64.8 C883).

0318

Near-isogenic barley lines with genes for resistance to powdery mildew.

CRPSAY. Kolster, P. Munk, L.; Stolen, O.; Lohde, J. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 903-907. Includes references. (NAL Call No.: DNAL 64.8 C883).

0319

New barley developed (Scio variety, resistant to lodging).

Corvallis, The Station. Oregon's agricultural progress - Oregon Agricultural Experiment Station. Spring/Summer 1982. v. 28/29 (4/1). p. 14. (NAL Call No.: 100 OR30R).

0320

New mutants in the genetic male sterile barley collection.

Hockett, E.A. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 70-73. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0321

New sources of resistance to *Puccinia hordei* in barley land race cultivars.

PHYTAJ. Yahyaoui, A.H. Sharp, E.L.; Reinhold, M. St. Paul, Minn. : American Phytopathological Society. Phytopathology. July 1988. v. 78 (7). p. 905-908. Includes references. (NAL Call No.: DNAL 464.8 P56).

(PLANT BREEDING)

0322

New varieties of oats from bond crosses resistant to Victoria Blight /by T.R. Stanton. Stanton, T. R. 1885-. Washington, D.C. : U.S. Dept. of Agriculture, 1948. 7 p., 1 folded leaf : ill. ; 26 cm. Bibliography: p.6-7. (NAL Call No.: DNAL 1 Ag84C no.795).

0323

Nitrate accumulation in plants.

Jensen, E.H. Reno, Nev. : College of Agriculture, University of Nevada-Reno, Nevada Cooperative Extension. Fact sheet - College of Agriculture, University of Nevada-Reno, Nevada Cooperative Extension. 1987? . (87-32). 3 p. (NAL Call No.: DNAL S544.3.N3C66).

0324

Nitrogen harvest index in oats: its repeatability and association with adaptation. CRPSAY, Rattunde, H.F. Frey, K.J. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1986. v. 26 (3). p. 606-610. Includes references. (NAL Call No.: DNAL 64.8 C883).

0325

Oats (Production, breeding, composition).

Youngs, V.L. Peterson, D.M.; Brown, C.M. St. Paul, Minn. : American Association of Cereal Chemists. Advances in cereal science and technology. 1982. Literature review. v. 5. p. 49-105. ill. Includes references. (NAL Call No.: TS2120.A3).

0326

Oats research.

Reeves, D.L. Brookings, S.D. : The Station. Plant science pamphlet - Plant Science Dept., Agricultural Experiment Station, South Dakota State University. In the series analytic: 1987 Annual Progress Report--Northeast Research Station, Watertown, South Dakota. Jan 1988. (5). p. 37-38. (NAL Call No.: DNAL S541.5.S8P5).

0327

Pathogen fitness in cereal mildews (Plant breeding, disease resistance).

Wolfe, M.S. NASSD. Barrett, J.A.; Slater, S.E. New York : Plenum Press. NATO advanced study institutes series. Series A. Life sciences. 1983. v. 55. p. 81-100. Includes references. (NAL Call No.: QH301.N32).

0328

Pathogenic variation in field populations of barley mildew.

Chan, K. Boyd, W.J.R. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 242-245. (NAL Call No.: DNAL SB191.B2A9 1987).

0329

Performance of cereal crops in the Tanana River Valley of Alaska, 1978-1979 (Varieties, yields, resistance).

Wooding, F.J. McBeath, J.H. Fairbanks : The Station. Circular - Alaska Agricultural Experiment Station. Dec 1979. (34). 23 p. (NAL Call No.: 100 AL122E).

0330

Performance of cereal crops in the Tanana Valley of Alaska, 1980 (Varieties, yields, disease resistance).

Wooding, F.J. McBeath, J.H.; Hanscom, J.T.; Van Veldhuizen, R.M. Fairbanks : The Station. Circular - Alaska Agricultural Experiment Station. Dec 1980. (36). 29 p. (NAL Call No.: 100 AL122E).

0331

Performance of cereal crops in the Tanana Valley of Alaska, 1981 (Varieties, yields, disease resistance).

Wooding, F.J. McBeath, J.H.; Hanscom, J.T.; Van Veldhuizen, R.M.; Delucchi, G.M. Fairbanks : The Station. Circular - Alaska Agricultural Experiment Station. Mar 1982. (42). 27 p. (NAL Call No.: 100 AL122E).

0332

Performance of cereal crops in the Tanana Valley of Alaska, 1982 (Variety trials, disease resistance comparisons, yields).

Wooding, F.J. McBeath, J.H.; Frost, S.; Hanscom, J.T.; Van Veldhuizen, R.M. Fairbanks : The Station. Circular - Alaska Agricultural Experiment Station. Feb 1983. (44). 28 p. (NAL Call No.: 100 AL122E).

0333

Performance of small grain varieties in Louisiana, 1987-88.

Harrison, S.A. Boquet, D.J.; Colyer, P.D.; Groth, D.E.; Habetz, R.J.; Hallmark, W.B.; Hutchinson, R.L.; Moore, S.H.; Rabb, J.L. Baton Rouge, La. : The Department. Report of projects - Louisiana Agricultural Experiment Station, Department of Agronomy. 1988. p. 80-83. (NAL Call No.: DNAL 100 L936).

0334

The physiological basis of nitrogen redistribution during grain filling in cereals.
Dalling, M.U. Rockville, Md. : American Society of Plant Physiologists, c1985. Exploitation of physiological and genetic variability to enhance crop productivity / edited by James E. Harper, Lawrence E. Schrader, and Robert W. Howell. Literature review. p. 55-71. ill. Includes 45 references. (NAL Call No.: DNAL SB189.4.E97).

0335

Physiological changes associated with three cycles of recurrent selection for grain yield improvement in oats.
CRPSAY. Payne, T.S. Stuthman, D.D.; McGraw, R.L.; Bregitzer, P.P. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 734-736. Includes references. (NAL Call No.: DNAL 64.8 C883).

0336

Plant cell culture and somatic cell genetics of cereals and grasses.
Vasil, I.K. New York : Academic Press, 1982. Plant improvement and somatic cell genetics / edited by I.K. Vasil, W.R. Scowcroft, K.J. Frey. Literature review. p. 179-203. Includes references. (NAL Call No.: SB123.P563).

0337

Plastid transcription activity and DNA copy number increase early in barley chloroplast development.
PLPHA. Baumgartner, B.J. Rapp, J.C.; Mullet, J.E. Rockville, Md. : American Society of Plant Physiologists. Plastid transcription activity and DNA copy number were quantified during chloroplast development in the first foliage leaf in dark-grown and illuminated barley (*Hordeum vulgare* L.) seedlings. Primary foliage leaves of seedlings given continuous illumination from 2 days post-imbibition reached a final mean length of 15 centimeters at 6.5 days, whereas primary leaves of dark-grown seedlings required 7 days to reach a similar length. Dividing cells were observed in the basal 0.5 to 1 centimeter of primary leaves until 5.5 days post-imbibition. Plastids isolated from cells located in the basal meristem of 4-day-old seedlings were small (approximately 2 micrometers in diameter), exhibited low transcription activity and contained approximately 130 copies of plastid DNA per organelle. Cell size increased from 18 to 60 micrometers in a 1 to 3 centimeter region located adjacent to the leaf basal meristem. In this region, transcriptional activity per plastid increased 10-fold and DNA copy number increased from 130 to 210. Plastid transcriptional activity declined rapidly in illuminated plants with increasing leaf cell age and plastid DNA copy number also declined

but with a slower time course. In dark-grown seedlings plastid transcriptional activity declined more slowly than in illuminated plants while DNA copy number remained constant with increasing cell age. These data show that plastid transcriptional activity and DNA copy number increase early in chloroplast development and that transcriptional activity per DNA template varies up to 5-fold during barley leaf biogenesis. *Plant physiology*. Mar 1989. v. 89 (3). p. 1011-1018. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0338

Powdery mildew resistance gene M1-a8 (Reg1h8) in northwest European spring barley varieties (*Erysiphe graminis hordei*, *Hordeum*).
Jorgensen, J.H. Jensen, H.P. Fort Collins : Dept. of Agronomy, Colorado State University. *Barley genetics newsletter*. Apr 10, 1983. v. 13. p. 51-53. Includes references. (NAL Call No.: QK495.G74B34).

0339

A preliminary note on the inheritance of rust resistance in oats.
AGUDAT. Garber, R.J. Madison, Wis. : American Society of Agronomy. *Agronomy journal*. Jan 1921. v. 13 (1). p. 41-43. ill. Includes references. (NAL Call No.: DNAL 4 AM34P).

0340

A preliminary study of the inheritance of rust resistance in oats.
AGUDAT. Parker, J.H. Madison, Wis. : American Society of Agronomy. Pedigree lines of two oat varieties, Burt and Sixty-Day, together with a large number of F2 generation hybrids between these varieties, were studied in relation to their rust resistance. Most of the inoculations were made on seedlings, but enough were made on plants at time of heading to show that the results were similar. The rusts used were the crown rust of oats, *Puccinia lolii avenae* McAlpine, and the stem rust of oats, *Puccinia graminis avenae* Erikss, and Henn. Burt and Sixty-Day and all the hybrids of these two varieties so far tested were found to be entirely susceptible to stem rust. All plants to Sixty-Day also were uniformly susceptible to crown rust. Of 223 inoculated plants of Burt, 48 were classified as resistant, 152 as intermediate, and 23 as susceptible. Each of the five hybrid families contained, in the F2 generation, some plants showing a high degree of resistance to crown rust and others which were as susceptible as plants of the Sixty-Day parent. In other words, there was definite segregation. There was, however, a rather large number of plants which were classified as intermediate and which showed varying degrees of resistance. The numerical results of inoculations made in the F2 hybrids were as follows: seedlings classified as resistant no. 81 (17.3%); seedlings classified as

(PLANT BREEDING)

intermediate no. 61 (13.0%); seedlings classified as susceptible no. 326 (69.7%); total no. 468 (100.0%). The fact that there were so many more susceptible than resistant plants indicates that susceptibility to crown rust in this cross is partially dominant, while resistance is recessive. These contrasted characters are not thought to be due to environmental conditions or to differences in the metabolism of the host plants, but to definite genetic factors. Nonhereditary factors may of course influence or modify their expression. Rust resistance and susceptibility hardly can be considered as simple characters or as being determined by a single factor difference. The F2 gene. *Agronomy journal*. Jan 1920. v. 12 (1). p. 23-38. plates. Includes references. (NAL Call No.: DNAL 4 AM34P).

0341

Proanthocyanidin-free barley for brewing: progress in breeding for high yield and research tool in polyphenol chemistry.
TOMBA. Wettstein, D. von. Nilan, R.A.; Ahrenst-Larsen, B.; Erdal, K.; Ingversen, J.; Jende-Strid, B.; Kristiansen, K.N.; Larsen, J.; Outtrup, H.; Ullrich, S.E. Madison, Wis. : The Association. Technical quarterly - Master Brewers Association of the Americas. 1985. v. 22 (2). p. 41-52. Includes references. (NAL Call No.: DNAL 390.9 M39T).

0342

Production of ochratoxin A in barley by *Aspergillus ochraceus* and *Penicillium viridicatum*: effect of fungal growth, time, temperature, and inoculum size (Mycotoxins).
Hagblom, P. Washington, D.C., American Society for Microbiology. Applied and environmental microbiology. May 1982. v. 43 (5). p. 1205-1207. Includes 10 ref. (NAL Call No.: 448.3 AP5).

0343

A quest for cereal varieties resistant to the minor cereal diseases in eastern Oregon.
Kolding, M.F. Corvallis, Or., The Station. Special report - Oregon Agricultural Experiment Station. June 1981. June 1981. (623). p. 35-42. (NAL Call No.: 100 OR3M).

0344

Rapid electrophoresis of oat (*Avena sativa* L.) prolamins from single seeds for cultivar identification.
CECHAF. Hansen, A.E. Nassuth, A.; Altosaar, I. St. Paul, Minn. : American Association of Cereal Chemists. Cereal chemistry. Mar/Apr 1988. v. 65 (2). p. 153-154. ill. Includes references. (NAL Call No.: DNAL 59.8 C33).

0345

Reaction of *Hordeum bulbosum* L. to Japanese races of powdery mildew (*Erysiphe graminis* f. sp. *hordei*) (that attack *Hordeum vulgare*, barley, genetic transfer of resistance).
Prasad, G. Yasuda, S.; Konishi, T. Fort Collins : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1983. v. 13. p. 60-62. Includes references. (NAL Call No.: QK495.G74B34).

0346

Receptivity, incubation period, and lesion size as criteria for screening barley genotypes for resistance to *pyrenophora teres*.
PHYTAJ. Nutter, F.W. Jr. Pederson, V.D. St. Paul, Minn. : American Phytopathological Society. Phytopathology. May 1985. v. 75 (5). p. 603-606. Includes 12 references. (NAL Call No.: DNAL 464.8 P56).

0347

Recurrent selection for tolerance to barley yellow dwarf virus in oat.
CRPSAY. Baltenberger, D.E. Ohm, H.W.; Foster, J.E. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1988. v. 28 (3). p. 477-480. Includes references. (NAL Call No.: DNAL 64.8 C883).

0348

Reduced accumulation of ABA during water stress in a molybdenum cofactor mutant of barley.
PLPHA. Walker-Simmons, M. Kudrna, D.A.; Warner, R.L. Rockville, Md. : American Society of Plant Physiologists. A barley (*Hordeum vulgare* L.) mutant (Az34) has been identified with low basal levels of abscisic acid (ABA) and with reduced capacity for producing ABA in response to water stress. The mutation is in a gene controlling the molybdenum cofactor resulting in a pleiotropic deficiency in at least three molybdoenzymes, nitrate reductase, xanthine dehydrogenase, and aldehyde oxidase. The mutant was found to lack aldehyde oxidase activity with several substrates including: (a) ABA aldehyde, a putative precursor of ABA; (b) an acetylenic analog of ABAaldehyde; and (c) heptaldehyde. Elevating the growth temperature from 18 to 26 degrees C caused mutant leaves to wilt and brown. Desiccation of mutant leaves was prevented by applying ABA. These results indicate that ABA biosynthesis at some developmental stages is dependent upon a molybdoenzyme which may be an aldehyde oxidase. Plant physiology. June 1989 v. 90 (2). p. 728-733. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0349

Registration of Arizona 8501 barley germplasm for disturbed land reclamation.
CRPSAY. Day, A.D. Ludeke, K.L.; Ottman, M.U. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 387. Includes references. (NAL Call No.: DNAL 64.8 C883).

0350

Registration of barley yellow dwarf virus tolerant barley composite cross XLIV germplasm.
CRPSAY. Crosslin, J.M. Carroll, T.W.; Hockett, E.A.; Zaske, S.K. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 387-388. Includes references. (NAL Call No.: DNAL 64.8 C883).

0351

Registration of CM 221 semidwarf buckwheat germplasm.
CRPSAY. Campbell, C.G. Madison, Wis. : Crop Science Society of America. Crop science. Jan/Feb 1987. v. 27 (1). p. 151. Includes references. (NAL Call No.: DNAL 64.8 C883).

0352

Registration of 'Diamond' barley.
CRPSAY. Kaufmann, M.L. Kibite, S. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1985. v. 25 (4). p. 706. Includes references. (NAL Call No.: DNAL 64.8 C883).

0353

Registration of 'Florida 401' rye.
CRPSAY. Pfahler, P.L. Barnett, R.D.; Luke, H.H. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 836. Includes references. (NAL Call No.: DNAL 64.8 C883).

0354

Registration of four pairs of greenbug-resistant vs. susceptible near-isolines of winter barley germplasms.
CRPSAY. Carver, B.F. Morgan, G.H.; Edwards, L.H.; Webster, J.A. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1988. v. 28 (6). p. 1034-1035. Includes references. (NAL Call No.: DNAL 64.8 C883).

0355

Registration of four stem rust and crown rust resistant oat germplasm lines.
CRPSAY. Rothman, P.G. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1984. v. 24 (6). p. 1217-1218. Includes 6 references. (NAL Call No.: DNAL 64.8 C883).

0356

Registration of 'Heartland' barley.
CRPSAY. Therrien, M.C. Irvine, R.B.; Campbell, K.W.; Wolfe, R.I. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1985. v. 25 (6). p. 1124. (NAL Call No.: DNAL 64.8 C883).

0357

Registration of 'Hercules' oat.
CRPSAY. Marshall, H.G. Kolb, F.L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1988. v. 28 (4). p. 719. Includes references. (NAL Call No.: DNAL 64.8 C883).

0358

Registration of 'Hitchcock' barley.
CRPSAY. Schmidt, J.W. Dreier, A.F.; Dofing, S.M. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1985. v. 25 (6). p. 1123. (NAL Call No.: DNAL 64.8 C883).

0359

Registration of 'Jasper' oat.
CRPSAY. Kibite, S. Kaufmann, M.L.; Allen, H.T. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1987. v. 27 (5). p. 1089. Includes references. (NAL Call No.: DNAL 64.8 C883).

0360

Registration of 'Kelly' oat.
CRPSAY. Reeves, D.L. Hall, L. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 385-386. (NAL Call No.: DNAL 64.8 C883).

0361

Registration of Kline barley.
CRPSAY. Brown, A.R. Morey, D.D.; Johnson, J.W.; Cunfer, B.M. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1985. v. 25 (4). p. 706-707. Includes 1 references. (NAL Call No.: DNAL 64.8 C883).

(PLANT BREEDING)

0362

Registration of 'Lamont' barley.
CRPSAY. Wesenberg, D.M. Robbins, G.S. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 373. Includes references. (NAL Call No.: DNAL 64.8 C883).

0363

Registration of 'Lancer' oats.
CRPSAY. Reeves, D.L. Hall, L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1985. v. 25 (4). p. 710. Includes 2 references. (NAL Call No.: DNAL 64.8 C883).

0364

Registration of leaf rust resistant barley composite cross XLI germplasm (*Hordeum vulgare*).
Bockelman, H.E. CRPSAY. Reinhold, M.; Sharp, E.L.; Eslick, R.F. Madison : Crop Science Society of America. Crop science. Nov/Dec 1983. v. 23 (6). p. 1224-1225. Includes references. (NAL Call No.: 64.8 C883).

0365

Registration of 'Multiline E76' and 'Multiline E77' oats.
CRPSAY. Frey, K.J. Browning, J.A.; Simons, M.D. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1985. v. 25 (6). p. 1125. Includes 1 reference. (NAL Call No.: DNAL 64.8 C883).

0366

Registration of net blotch resistant barley composite cross XLV germplasm.
CRPSAY. Bockelman, H.E. Sharp, E.L.; Harrabi, M.M.; Cherif, M. Madison, Wis. : Crop Science Society of America. Crop science. Jan/Feb 1988. v. 28 (1). p. 199. (NAL Call No.: DNAL 64.8 C883).

0367

Registration of 'Noble' barley.
CRPSAY. Helm, J.H. Salmon, D.F.; Dyson, D.H.; Stewart, W.M. Madison, Wis. : Crop Science Society of America. Crop science. Jan/Feb 1989. v. 29 (1). p. 235. (NAL Call No.: DNAL 64.8 C883).

0368

Registration of oat germplasms IA H676, IA H677, and IA H681 resistant to the crown fungus.
CRPSAY. Michel, L.J. Simons, M.D. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1985. v. 25 (4). p. 716-717. Includes 4 references. (NAL Call No.: DNAL 64.8 C883).

0369

Registration of powdery mildew resistant barley composite cross XLII germplasm (*Hordeum vulgare*, *Erysiphe graminis*).
Bockelman, H.E. CRPSAY. Sharp, E.L.; Eslick, R.F.; Ramage, R.T. Madison : Crop Science Society of America. Crop science. Nov/Dec 1983. v. 23 (6). p. 1225. Includes references. (NAL Call No.: 64.8 C883).

0370

Registration of 'Proat' oat.
CRPSAY. Stuthman, D.D. Rothman, P.G.; Wilcoxson, R.D.; Rines, H.W. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1987. v. 27 (4). p. 816. (NAL Call No.: DNAL 64.8 C883).

0371

Registration of 'Ray' barley.
CRPSAY. Lafever, H.N. Madison, Wis. : Crop Science Society of America. Crop science. Jan/Feb 1988. v. 28 (1). p. 187. (NAL Call No.: DNAL 64.8 C883).

0372

Registration of 'Samson' barley.
CRPSAY. Helm, J.H. Dyson, D.H.; Stewart, W.M. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 384. (NAL Call No.: DNAL 64.8 C883).

0373

Registration of scald and net blotch resistant barley composite cross XLIII germplasm (*Hordeum vulgare*).
Bockelman, H.E. CRPSAY. Eslick, R.F.; Sharp, E.L. Madison : Crop Science Society of America. Crop science. Nov/Dec 1983. v. 23 (6). p. 1225-1226. Includes references. (NAL Call No.: 64.8 C883).

0374

Registration of 'Simpson' oat.

CRPSAY. Graham, W.D. Jr. Morton, B.C. Jr.; Kingsland, G.C.; Gambrell, R.H. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 386. (NAL Call No.: DNAL 64.8 C883).

0375

Registration of six germplasm sources of cereal leaf beetle resistant hard red spring wheats (Reg. No. GP 132 to GP 137).

Smith, D.H. Jr. AR-NC. Webster, J.A.; Everson, E.H. Madison, Wis., Crop Science Society of America. Crop science. May/June 1980. v. 20 (3). p. 420. (NAL Call No.: 64.8 C883).

0376

Registration of three oat germplasm lines resistant to the crown rust fungus.

CRPSAY. Simons, M.D. Michel, L.J.; Frey, K.J. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1987. v. 27 (2). p. 369. Includes references. (NAL Call No.: DNAL 64.8 C883).

0377

Registration of two cereal leaf beetle resistant barley germplasms (*Oulema melanopus*).

Smith, D.H. Jr. Webster, J.A.; Grafius, J.E. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1984. v. 24 (4). p. 828. Includes references. (NAL Call No.: 64.8 C883).

0378

Registration of 'Venus' barley.

CRPSAY. Brown, A.R. Johnson, J.W.; Rothrock, C.S.; Bruckner, P.L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1988. v. 28 (4). p. 718-719. Includes references. (NAL Call No.: DNAL 64.8 C883).

0379

Registration of 'Virden' barley.

CRPSAY. Therrien, M.C. Irvine, R.B.; Campbell, K.W.; Wolfe, R.I. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 374. Includes references. (NAL Call No.: DNAL 64.8 C883).

0380

Registration of 'Webster' oat.

CRPSAY. Frey, K.J. Simons, M.D.; Michel, L.J.; Murphy, J.P.; Brownning, J.A. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 374-375. Includes references. (NAL Call No.: DNAL 64.8 C883).

0381

Registration of Webster oat isolines as parental lines.

CRPSAY. Frey, K.J. Simons, M.D.; Michel, L.J.; Murphy, J.P.; Browning, J.A. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 386-387. Includes references. (NAL Call No.: DNAL 64.8 C883).

0382

Registration of 'Willis' winter barley.

CRPSAY. Sorrells, M.E. Jensen, N.F. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1989. v. 29 (4). p. 1086. Includes references. (NAL Call No.: DNAL 64.8 C883).

0383

Registration of 'Wysor' barley.

CRPSAY. Starling, T.M. Roane, C.W.; Camper, H.M. Jr.; Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1987. v. 27 (6). p. 1306-1307. (NAL Call No.: DNAL 64.8 C883).

0384

Relation between kernel row number and crude protein content of the grain in barley (Effects of two-rowed alleles on agronomic characters, physiological cause).

Yasuda, S. Moriya, I. Fort Collins, Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1982. v. 12. p. 45-46. (NAL Call No.: QK495.G74B34).

0385

A relationship between ice-nucleation-active bacteria, freeze damage, and genotype in oats.

PHYTAU. Marshall, D. St. Paul, Minn. : American Phytopathological Society. Phytopathology. July 1988. v. 78 (7). p. 952-957. Includes references. (NAL Call No.: DNAL 464.8 P56).

(PLANT BREEDING)

0386

Relationship between subcrown internode length and winter survival in winter barley.

CRPSAY. Dofing, S.M. Schmidt, J.W. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1985. v. 25 (4). p. 690-692. Includes references. (NAL Call No.: DNAL 64.8 C883).

0387

Resistance of cereal crops to aphids: role of allelochemicals.

ACSMC. Corcuera, L.J. Argandona, V.H.; Zuniga, G.E. Washington, D.C. : The Society. ACS Symposium series - American Chemical Society. 1987. (330). p. 129-135. Includes references. (NAL Call No.: DNAL QD1.A45).

0388

Resistance of *Hordeum spontaneum* collected in Israel to infection with *Erysiphe graminis hordei* (Barley, powdery mildew, germplasm, disease).

Moseman, J.G. CRPSAY. Nevo, E.; Zohary, D. Madison : Crop Science Society of America. Crop science. Nov/Dec 1983. v. 23 (6). p. 1115-1119. maps. Includes references. (NAL Call No.: 64.8 C883).

0389

Resistance of wild barley accessions from Israel to leaf rust collected in the USA and Israel.

CRPSAY. Manisterski, J. Treeful, L.; Tomerlin, J.R.; Anikstern, Y.; Moseman, J.G.; Wahl, I.; Wilcoxson, R.D. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 727-730. Includes 22 references. (NAL Call No.: DNAL 64.8 C883).

0390

Resistance to covered smut in varieties and hybrids of oats.

AGJOAT. Gaines, E.F. Madison, Wis. : American Society of Agronomy. Of 210 varieties and selections of oats tested for resistance to covered smut, *Ustilago levis* (K. & S.) Magn., during the past eight years, 21 were smut free, the others ranging from slight infection to extreme susceptibility in which nine-tenths of the panicles were black masses of smut spores. Markton, one of the immune selections, is a thin hulled, white oat of sativa type, and is one of the best yielding varieties tested in recent years. It has been increased and grown commercially since 1924. The inheritance of the immunity of Red Rustproof has been studied in four crosses with susceptible varieties. The crosses with Black Tartarian and Abundance indicate that Red Rustproof carries three dominant factors for immunity, any one of which

prevents the production of covered smut spores. In crosses with Large and Chinese Hulless, one factor apparently does not give complete dominance in hulless segregates, but otherwise the prepotency of the factors for immunity is similar in all four crosses. Of 56 F3 plants selected from smut-free rows, 45 produced only smut-free plants in the F4 generation. From the results already obtained in varietal testing and breeding for disease resistance in field crops, it is to be hoped that the major diseases will in time be successfully controlled by the development and introduction of resistant or immune varieties. Agronomy journal. Dec 1925. v. 17 (12). p. 775-789. (NAL Call No.: DNAL 4 AM34P).

0391

Resistance to disease and maximum potential yield.

Vanderplank, J.E. New York : Gordon and Breach Science Pub., c1988. Experimental and conceptual plant pathology / edited by R.S. Singh ... et al. . p. 593-599. Includes references. (NAL Call No.: DNAL SB731.E97).

0392

Response of current Midwestern soybean cultivars to late planting.

CRPSAY. Raymer, P.L. Bernard, R.L. Madison, Wis. : Crop Science Society of America. Soybean Glycine max (L.) Merr. is grown in the Midwestern USA primarily as a full-season crop and only to a limited extent as a double crop following small grains. Development of cultivars specifically adapted to later planting dates commonly associated with double-crop production has been suggested as a means to expand double-crop hectareage in this area. To determine if currently used soybean cultivars differ in their adaptation to late planting and if any specific plant traits are related to improved performance under late-planted conditions, 16 soybean cultivars were evaluated at both conventional (May) and late (late June to early July) planting dates in 1979, 1980, and 1981. Cultivar by planting date interactions were found for days to maturity, height at maturity, seed quality, and seed mottling, but not for yield, days to flowering, height at flowering, lodging, and weight per 100 seeds. All cultivars suffered substantial and similar yield reductions when planted late. Phenotypic correlation coefficients of cultivar performance between the two planting dates were positive and highly significant for all plant traits measured. The relationship of yield with various plant traits varied greatly from year to year and no differences in these relationships were observed between the two planting dates. These results do not furnish any evidence to justify a separate breeding program for the development of double-crop cultivars adapted to the Midwest. The lack of a strong cultivar by planting date interaction for yield and the lack of any strong associations of specific plant characteristics with yield in a

late-planted environment imply that testing in a conventional early-planted environment will be effective in identifying lines that perform well in either full-season or double-crop environments. Crop science. Sept/Oct 1988. v. 28 (5). p. 761-764. Includes references. (NAL Call No.: DNAL 64.8 C883).

0393

Responses of two-, three-, and four-component barley mixtures to a variable pathogen population.

CRPSAY. McDonald, B.A. Allard, R.W.; Webster, R.K. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1988. v. 28 (3). p. 447-452. Includes references. (NAL Call No.: DNAL 64.8 C883).

0394

Root development and lodging resistance in oats / Dale T. Sechler.

Sechler, Dale Truman, 1926-. Columbia, Mo. : University of Missouri, College of Agriculture, Agricultural Experiment Station, 1961. Cover title. 38 p. : ill. ; 23 cm. Bibliography: p. 37. (NAL Call No.: DNAL 100 M693 (3) no.769).

0395

Seed dormancy in wild and weedy relatives of cereals.

Key, J.M. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 15-25. ill. Includes references. (NAL Call No.: DNAL SB189.6.P73).

0396

Selection of barley for slow rusting resistance to leaf rust in epidemics of different severity.

CRPSAY. Andres, M.W. Wilcoxson, R.D. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1986. v. 26 (3). p. 511-514. Includes 25 references. (NAL Call No.: DNAL 64.8 C883).

0397

Sources of genes resistant to Puccinia hordei in barley (leaf rust, Hordeum vulgare, cultivars and lines).

Sharp, E.L. Reinhold, M. St. Paul, American Phytopathological Society. Plant disease. Nov 1982. v. 66 (11). p. 1012-1013. 14 ref. (NAL Call No.: 1.9 P69P).

0398

Soybean cultivar response to reduced tillage systems in northern dryland areas.

AGJOAT. Deibert, E.J. Madison, Wis. : American Society of Agronomy. Information on response of soybean Glycine max (L.) Merr. cultivars to reduced tillage systems in northern dryland areas is limited. A 4-yr field study (1984 to 1987) was conducted to evaluate the effect of tillage system, weed control method, and cultivar maturity on soybean seed yield variables. An early and a late-maturing soybean cultivar were grown on a Fargo clay (fine, montmorillonitic frigid Vertic Haplaquoll) on established tillage plots. Tillage systems included conventional (moldboard plow) and three reduced tillage systems (sweep, intertill, and no-till) with herbicides or herbicides plus cultivation for weed control. Climatic conditions resulted in differences among years in seed yield, seed weight, seed moisture, seed oil concentration, and seed oil yield. These seed variables were not significantly influenced by tillage system, weed control method, or cultivar maturity when grown in rotation with barley (Hordeum vulgare L.), but showed significant interactions. Cultivation for weed control depressed seed yield and weight of only the early cultivar. Early plant water stress (June and July) lowered yield of the early cultivar more than the late cultivar. Early cultivar no-till yields (1240 kg ha⁻¹) were greater than tilled system yields (average 1070 kg ha⁻¹), while late cultivar yields were similar among systems (average 1420 kg ha⁻¹). An early maturing cultivar performed similarly to a late-maturing cultivar irrespective of tillage system unless early plant water stress was encountered. Fall application of granular herbicide provided good weed control, but cultivation for weed control was not beneficial for the yields parameters measured. Agronomy journal. July/Aug 1989. v. 81 (4). p. 672-676. Includes references. (NAL Call No.: DNAL 4 AM34P).

0399

Sterility in rye.

AGJOAT. Stroman, G.N. Madison, Wis. : American Society of Agronomy. Agronomy journal. June 1923. v. 15 (6). p. 253-254. (NAL Call No.: DNAL 4 AM34P).

0400

Sterility of rye.

AGJOAT. Leith, B.D. Madison, Wis. : American Society of Agronomy. Agronomy journal. Mar 1925. v. 17 (3). p. 129-132. (NAL Call No.: DNAL 4 AM34P).

(PLANT BREEDING)

0401

Strategies to select biochemical mutants with potential use in genetic manipulation of plants.

Jacobs, M. Negrutiu, I.; Cammaerts, D.; Cattoir-Reynaerts, A.; Dirks, R.; Dolferus, R.; Famelaer, Y. New York : Alan R. Liss. UCLA symposia on molecular and cellular biology. Paper presented at the "Symposium on Plant Genetics," April 13-19, 1985, Keystone, Colorado. 1985. v. 35. p. 685-700. Includes references. (NAL Call No.: DNAL QH506.U34).

0402

Student's method for interpreting paired experiments.

AGJOAT. Love, H.H. Brunson, A.M. Madison, Wis. : American Society of Agronomy. Too close an analogy should not be drawn between the determination of exact constants and the results of comparative agronomic experiments. Bessel's and Peter's formulas are not adapted to calculating the degree of significance of a difference between two varieties or two comparative treatments in experiments in which the variability within each variety or treatment is high because they extend through a series of years or because of place effect. For observations which naturally arrange themselves in pairs, Student's method is a better method with which to determine the probability of the difference, and deserves a larger application in the interpretation of agronomic investigations. Agronomy journal. Jan 1924. v. 16 (1). p. 60-68. (NAL Call No.: DNAL 4 AM34P).

0403

Studies of inheritance in crosses between Bond, Avena byzantina, and varieties of A. sativa /H.K. Hayes, M.B. Moore, E.C. Stakman. Hayes, H. K. 1884-. Moore, M. B. 1905-; Stakman, E. C. 1885-1979. St. Paul : University of Minnesota, Agricultural Experiment Station, 1939. Cover title. 38 p. : ill. ; 23 cm. Bibliography: p. 37-38. (NAL Call No.: DNAL 100 M66 (3) no.137).

0404

Sugar composition and freezing tolerance in barley crowns at varying carbohydrate levels. CRPSAY. Livingston, D.P. III. Olien, C.R.; Freed, R.D. Madison, Wis. : Crop Science Society of America. During hardening, carbohydrate content in winter cereals is affected by light intensity, which in turn affects freezing survival. To determine whether total carbohydrate content affects composition of individual sugars and whether a particular composition is related to freezing tolerance, three barley (*Hordeum vulgare* L.) cultivars, differing in carbohydrate composition and in response to a controlled freezing test, were hardened under six light levels. At each level, total crown sugars were measured by

water/ethanol extraction and high-pressure liquid chromatography and at four light levels plants were freeze tested. Carbohydrate composition changed depending on total carbohydrate content, and maximum freezing resistance was not found at the maximum simple sugar level. In addition, the relationship between kill temperature and total carbohydrate was different for the three cultivars. A simple relationship between carbohydrate composition and freezing tolerance was not found, but results suggest that the three cultivars may allocate total carbohydrate differently for cryoprotection. Crop science. Sept/Oct 1989. v. 29 (5). p. 1266-1270. Includes references. (NAL Call No.: DNAL 64.8 C883).

0405

Survey of barley germplasm for betaine-accumulating potential in controlled environments and in the field (*Hordeum vulgare*, *Hordeum spontaneum*, genotypes).

Ladyman, J.A.R. Ditz, K.M.; Hanson, A.D. East Lansing, Mich., The Laboratory. Annual report - Michigan State University, MSU/DOE Plant Research Laboratory. 1981. 1981 (16th). p. 64-66. 3 ref. (NAL Call No.: OK1.M5).

0406

Techniques for selection of cold hardiness in cereals (Oats).

Marshall, G.H. Olien, C.R.; Everson, E.H. Boca Raton, Fla., CRC Press. Analysis and improvement of plant cold hardiness. 1981. 1981. p. 139-159. 30 ref. (NAL Call No.: SB781.A52).

0407

Tillering dynamics of winter barley as influenced by cultivar and nitrogen fertilizer: a field study (*Hordeum vulgare*, *Hordeum distichon* L., tillers survival).

Garcia del Moral, L.F. Ramos, J.M.; Recalde, L. Madison : Crop Science Society of America. Crop science. Jan/Feb 1984. v. 24 (1). p. 179-181. ill. Includes references. (NAL Call No.: 64.8 C883).

0408

Tillering in barley: genotype, row spacing, and seeding rate effects (Tiller mortality, abortion, axillary shoots).

Simmons, S.R. Rasmusson, D.C.; Wiersma, J.V. Madison, Wis., Crop Science Society of America. Crop science. July/Aug 1982. v. 22 (4). p. 801-805. 10 ref. (NAL Call No.: 64.8 C883).

0409

Tolerances of oat cultivars to an acid soil high in exchangeable aluminum.
 JPNUDS. Foy, C.D. Smith, D.H. Jr.; Briggie, L.W. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Tenth International Plant Nutrition Colloquium," August 4-9, 1986, Beltsville, Maryland. 1987. v. 10 (9116). p. 1163-1174. ill. Includes references. (NAL Call No.: DNAL QK867.J67).

0410

Transfer of field resistance to Puccinia coronata from Avena sterilis to cultivated oats by backcrossing.
 PHYTAJ. Simons, M.D. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Mar 1985. v. 75 (3). p. 314-317. Includes 26 references. (NAL Call No.: DNAL 464.8 P56).

0411

Un8 allele for loose smut resistance associated with necrosis in embryos of infected barley.
 PHYTAJ. Gabor, B.K. Thomas, P.L. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Apr 1987. v. 77 (4). p. 533-538. ill. Includes references. (NAL Call No.: DNAL 464.8 P56).

0412

Use of phytotoxins in selection of disease resistant mutants in tissue culture.
 ISJRA6. Gengenbach, B.G. Rines, H.W. Ames, Iowa : Iowa State University. Iowa state journal of research. May 1986. v. 60 (4). p. 449-476. ill. Includes references. (NAL Call No.: DNAL 470 I09).

0413

Using cDNA probes to identify barley yellow dwarf virus-resistant barley lines.
 Heath, R. Van de Velde, R.; Holloway, P. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 247-248. (NAL Call No.: DNAL SB191.B2AS 1987).

0414

Variation of resistance to powdery mildew in wild barley, Hordeum spontaneum.
 Fukuyama, T. Heta, H.; Takeda, K. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Nov 10, 1986. v. 16. p. 15-17. (NAL Call No.: DNAL QK495.G74B34).

0415

Varietal variation of water sensitivity in Asian barley varieties.
 Takeda, K. Fukuyama, T. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Oct 10, 1985. v. 15. p. 15-16. (NAL Call No.: DNAL QK495.G74B34).

0416

Victorin as a tool for elucidating the molecular mechanism of disease specificity in victoria blight of oats.
 Macko, V. Wolpert, T.J.; Acklin, W.; Arigoni, D. Columbia, Mo. : The Interdisciplinary Plant Biochemistry and Physiology Program. Current topics in plant biochemistry and physiology : Proceedings of the ... Plant Biochemistry and Physiology Symposium held at the University of Missouri, Columbia. 1987. v. 6. p. 55-58. Includes references. (NAL Call No.: DNAL QK861.P55).

0417

Volunteer barley (Hordeum vulgare) interference in canola (Brassica campestris and B. napus).
 WEESA6. O'Donovan, J.T. Sharma, A.K.; Kirkland, K.J.; De St Remy, E.A. Champaign, Ill. : Weed Science Society of America. The yield potential and the effect on yield loss of canola of different densities of volunteer barley were investigated at three locations in western Canada. Field studies were conducted from 1982 to 1986. Rectangular hyperbolic models based on data pooled over years, locations, and canola cultivars, and incorporating different densities of volunteer barley and canola accurately portrayed field responses in most instances. Results indicated that volunteer barley severely reduced canola yield. However, financial losses due to reduced canola yield were partly offset by the volunteer barley crop. Weed science. Nov 1988. v. 36 (6). p. 734-739. Includes references. (NAL Call No.: DNAL 79.8 W41).

0418

Water stress enhances expression of an alpha-amylase gene in barley leaves.
 PLPHA. Jacobsen, J.V. Hanson, A.D.; Chandler, P.C. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Feb 1986. v. 80 (2). p. 350-359. ill. Includes 33 references. (NAL Call No.: DNAL 450 P692).

0419

Allelopathic activity of rye (*Secale cereale* L.).

Barnes, J.P. Putnam, A.R.; Burke, B.A. New York, N.Y. : John Wiley & Sons, c1986. The Science of allelopathy / edited by Alan R. Putnam and Chung-Shih Tang. Literature review. p. 271-286. ill. Includes references. (NAL Call No.: DNAL QK898.A43S34).

0420

Conjugation of allelochemicals by plant. Enzymatic glucosylation of salicylic acid by *Avena sativa*.

ACSMC. Balke, N.E. Davis, M.P.; Lee, C.C. Washington, D.C. : The Society. ACS Symposium series - American Chemical Society. 1987. (330). p. 214-227. Includes references. (NAL Call No.: DNAL QD1.A45).

0421

The effect of sigma-aminolevulinic acid on red-light-induced leaf unrolling (Barley). Sundqvist, C. Briggs, W.R. Washington, D.C., The Institute. Year book - Carnegie Institution of Washington. 1979/1980. 1979/1980. (79th). p. 136-138. ill. 4 ref. (NAL Call No.: 500 C21).

0422

Effects of ambient temperature and nitrogen supply on content of four nitrogen fractions in barley plants / by Ikbalur Rashid Chowdhury. -. Chowdhury, Ikbalur Rashid, 1939-. 1970. Thesis (Ph.D.)--North Dakota State University, 1970. Photocopy. Ann Arbor, Mich. : University Microfilms, 1971. xiii, 204 leaves ; 21 cm. Bibliography: leaves 162-175. (NAL Call No.: DISS 71-5,508).

0423

Effects of deletions in the N-terminal basic arm of brome mosaic virus coat protein on RNA packaging and systemic infection.

JOVIAM. Sacher, R. Ahlquist, P. Washington, D.C. : American Society for Microbiology. The first 25 amino acids of brome mosaic virus (BMV) coat protein include 8 basic and no acidic residues and are implicated in binding the encapsidated RNA. Using infectious transcripts from BMV RNA3 cDNA clones, we modified this region of the coat gene. A coat protein mutant with the first 25 amino acids deleted failed to direct either packaging of viral RNA in protoplasts or systemic infection of whole barley plants. Neither symptoms, virions, nor viral RNA was detectable in plants inoculated with this mutant or a mutant with a frameshift mutation in the coat gene. Mutants with the normal start codon changed to AAG or with the first eight codons deleted allowed translation to start at a downstream AUG,

resulting in a deletion of the first 7 amino acids of the mature wild-type coat protein. These mutants not only packaged viral RNA in protoplasts but directed symptomatic, systemic infections that developed with normal speed and degree of spread within the host. The AUG-to-AAG point substitution did not revert to the wild type after long-term culture in planta. Wild-type BMV virions were also found to contain small amounts of a protein that coelectrophoresed with the truncated coat protein produced by the viable AAG and eight-codon-deletion mutants. This minor coat protein species presumably arose by infrequent translation initiation at the second AUG in the wild-type coat protein gene. Absence of encapsidation-competent coat protein appeared to stimulate production of nonstructural proteins in protoplast infections. Journal of virology. Nov 1989. v. 63 (11). p. 4545-4552. ill. Includes references. (NAL Call No.: DNAL QR360.J6).

0424

Factors affecting survival of winter oats / by Franklin A. Coffman .

Coffman, Franklin A. 1892-. Washington : U.S. Dept. of Agriculture, 1965. 28 p. : 1 map ; 23 cm. Literature cited: p. 27-28. (NAL Call No.: DNAL 1 Ag84Te no.1346).

0425

Natural occurrence of the mycotoxin viomellein in barley and the associated quinone-producing penicillia.

Hald, B.APMBA. Christensen, D.H.; Krogh, P. Washington : American Society for Microbiology. Applied and environmental microbiology. Dec 1983. v. 46 (6). p. 1311-1317. ill. Includes references. (NAL Call No.: 448.3 AP5).

0426

Oat stem vascular size in relation to kernel number and weight. 1. Controlled environment (*Avena sativa*).

Housley, T.L. Peterson, D.M. Madison, Wis., Crop Science Society of America. Crop science. Mar/Apr 1982. v. 22 (2). p. 259-263. ill. Includes 15 ref. (NAL Call No.: 64.8 C883).

0427

Pulse domestication and cereal domestication: how different are they?

ECBOA. Zohary, D. Bronx, N.Y. : New York Botanical Garden. Economic botany. Jan/Mar 1989. v. 43 (1). p. 31-34. Includes references. (NAL Call No.: DNAL 450 EC7).

0428

Response of round-headed buckwheat to summer burning.

NOSCA. Rickard, W.H. Pullman, Wash. : Washington State University Press. Northwest science : official publication of the Northwest Scientific Association. Aug 1989. v. 63 (4). p. 144-145. Includes references. (NAL Call No.: DNAL 470 N81).

0429

Rye residues contribute weed suppression in no-tillage cropping systems (Agroecosystems, biomass).

Barnes, J.P.JCECD. Putnam, A.R. New York : Plenum Press. Journal of chemical ecology. Aug 1983. v. 9 (8). p. 1045-1057. ill. Includes references. (NAL Call No.: QD415.A1J6).

0430

Stimulatory and inhibitory effects of light upon auxin biosynthesis (Avena, oats).

Wolf, F.T. Nashville, Tenn. : The Academy. Journal of the Tennessee Academy of Science. Jan/Apr 1984. v. 59. p. 25-27. Includes references. (NAL Call No.: 500 T25A).

PLANT STRUCTURE

0431

Cell membrane permeability and ultrastructural effects of difenzoquat on wild oats (*Avena fatua*).
WEESA6. Thai, K.M. Jana, S.; Fowke, L.C. Champaign, Ill. : Weed Science Society of America. Effects of difenzoquat on wild oats grown under controlled environmental conditions were studied. Seedling height and fresh weight were significantly reduced 5 days after postemergence treatment. Dose-dependent increase in cell membrane permeability was detected after a 12-h exposure to the herbicide. Scanning electron micrography showed normal leaf hairs and cuticular wax but swollen guard cells 10 days after treatment. Ultrastructural changes occurred before the visible symptoms. The primary effect of difenzoquat appears to be the disruption of the tonoplast and plasmalemma. The tonoplast showed greater damage than the plasmalemma. Secondary effects included damage to mitochondria and chloroplasts. Mitochondria were swollen and often ruptured, but the effect did not increase with the duration of exposure. Chloroplasts became spherical in shape, and their contents were also affected. The changes included: accumulation and then disappearance of starch granules, swelling of frets, fusion of granal thylakoids, detachment and then rupture of the outer membrane of the envelope, and clumping of ribosomes. By contrast, natural senescence caused greater injury of the plasmalemma than the tonoplast, a marked increase in size of plastoglobuli, and loss of starch grains without early accumulation. Weed science. Jan 1989. v. 37 (1). p. 98-106. ill. Includes references. (NAL Call No.: DNAL 79.8 W41).

0432

Changes in endogenous gibberellins and the metabolism of (3H)-GA4 after geostimulation in shoots of the oat plant (*Avena sativa*) (Recovery from lodging).
Pharis, R.P. Legge, R.L.; Noma, M.; Kaufman, P.B.; Ghosheh, N.S.; LaCroix, J.D.; Heller, K. Rockville, Md.; American Society of Plant Physiologists. Plant physiology. May 1981. v. 67 (5). p. 892-897. ill. 26 ref. (NAL Call No.: 450 P692).

0433

Characterization of the nuclear DNA of *Hordeum vulgare* root hairs: amplification disappears under salt stress.
AJBDAA. Murry, L.E. Christianson, M.L.; Alfinito, S.H.; Garger, S.J. Columbus, Ohio : Botanical Society of America. American journal of botany. Dec 1987. v. 74 (12). p. 1779-1786. Includes references. (NAL Call No.: DNAL 450 AM36).

0434

Effect of erect leaf angle on grain yield in barley.
CRPSAY. Tungland, L. Chapko, L.B.; Wiersma, J.V.; Rasmusson, D.C. Madison, Wis. : Crop Science Society of America. Crop science. Jan/Feb 1987. v. 27 (1). p. 37-40. Includes references. (NAL Call No.: DNAL 64.8 C883).

0435

The effects of iron and light treatments on chloroplast composition and ultrastructure in iron-deficient barley leaves (*Hordeum vulgare*).
Pushnik, J.C. Miller, G.W. New York ; Basel : Marcel Dekker, 1982. Iron nutrition and interactions in plants : Brigham Young University, August 12-14, 1981 / edited by S.D. Nelson ... (et al.). p. 311-321. ill. 26 ref. (NAL Call No.: QK867.J67 v. 5, nos. 4-7).

0436

Heat shock causes destabilization of specific mRNAs and destruction of endoplasmic reticulum in barley aleurone cells.
PNASA. Belanger, F.C. Brod1, M.R.; Ho, T.H.D. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. Mar 1986. v. 83 (5). p. 1354-1358. ill. Includes references. (NAL Call No.: DNAL 500 N21P).

0437

Inheritance of subcrown internode length in a winter barley cross (Winter hardiness, optimum depth of crown formation).
Dofing, S.M. Schmidt, J.W. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1984. v. 24 (4). p. 692-694. ill. Includes references. (NAL Call No.: 64.8 C883).

0438

Localization and pattern of graviresponse across the pulvinus of barley *Hordeum vulgare*.
PLPHA. Brock, T.G. Lu, C.R.; Ghosheh, N.S.; Kaufman, P.B. Rockville, Md. : American Society of Plant Physiologists. Pulvini of excised stem segments from barley (*Hordeum vulgare* cv Larker') were pretreated with 1 millimolar coumarin before gravistimulation to reduce longitudinal cell expansion and exaggerate radial cell enlargement. The cellular localization and pattern of graviresponse across individual pulvini were then evaluated by cutting the organ in cross-section, photographing the cross-section, and then measuring pulvinus thickness and the radial width of cortical and epidermal cells in enlargements of photomicrographs. With respect to orientation during gravistimulation, we designated the uppermost point of the cross-section 0 degrees and the lowermost point

180 degrees. A gravity-induced increase in pulvinus thickness was observable within 40 degrees of the vertical in coumarin-treated pulvini. In the upper halves of coumarin-treated gravistimulated pulvini, cells in the inner cortex and inner epidermis had increased radial widths, relative to untreated gravistimulated pulvini. In lower halves of coumarin-treated pulvini, cells in the central and outer cortex and in the outer epidermis showed the greatest increase in radial width. Cells comprising the vascular bundles also increased in radial width, with this pattern following that of the central cortex. These results indicate (a) that all cell types are capable of showing a graviresponse, (b) that the graviresponse occurs in both the top and the bottom the responding organ, and (c) that the magnitude of the response increases approximately linearly from the uppermost point to the lowermost. These results are also consistent with models of gravitropism that link the pattern and magnitude of the graviresponse to graviperception via statolith sedimentation. Plant physiology. Oct 1989. v. 91 (2). p. 744-748. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0439

The modification of the foliage structure and radiation absorption in a plant canopy by wind. Hipps, L.E. Baiden, E. Boston : The Society, 1985. 17th Conference on Agricultural and Forest Meteorology and seventh Conference on Biometeorology and Aerobiology, May 21-24, 1985, Scottsdale, Ariz. : preprint volume / sponsored by the American Meteorological Society. p. 127-128. Includes references. (NAL Call No.: DNAL S600.2.C6 1985).

0440

Molecular and cellular biology associated with endosperm mobilization in germinating cereal grains.

Fincher, G.B. Palo Alto, Calif. : Annual Reviews, Inc. Annual review of plant physiology and plant molecular biology. Literature review. 1989. v. 40. p. 305-346. ill. Includes references. (NAL Call No.: DNAL QK1.A57).

0441

Morphological and histological effects of three grass selective herbicides on developing wild oat (*Avena fatua*) stems.

WEESA6. Jain, R. Born, W.H. vanden. Champaign, Ill. : Weed Science Society of America. Three grass selective herbicides, sethoxydim, fluazifop, and haloxyfop, applied to wild oat plants at the five-leaf stage inhibited growth and induced chlorosis in leaves. Young and actively growing tissues were affected first. Stem elongation in wild oat was inhibited within 2 days of treatment with sethoxydim and within 5 days of treatment with fluazifop or haloxyfop. At these same observation times,

internodes that were elongating rapidly at the time of treatment were constricted at the base. These symptoms were followed by necrosis of the internode tissue. Histological examination of the affected internodes indicated that the herbicides inhibited cell division in very young internodes and inhibited both cell division and cell elongation in slightly older internodes. Initial injury occurred in the epidermal, cortical, and procambium cells of the peripheral regions of the stems located at the base of the affected internodes. Necrosis then progressed to the center of the stem tissue and all cells in the internodes were killed within 14 days of treatment. All three herbicides caused similar morphological and histological effects on developing wild oat stems. Weed science. July 1989. v. 37 (4). p. 575-584. ill. Includes references. (NAL Call No.: DNAL 79.8 W41).

0442

New mutants in the genetic male sterile barley collection.

Hockett, E.A. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 70-73. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0443

Physiological and ultrastructural studies of oat membranes treated with *Helminthosporium victoriae* toxin / by Vernon Edward Gracen, Jr.

-. Gracen, Vernon Edward, 1945-. 1970. Thesis (Ph.D.)--University of Florida, 1970. Photocopy. Ann Arbor, Mich. : University Microfilms, 1971. ix, 89 leaves ; 21 cm. Bibliography: leaves 84-88. (NAL Call No.: DISS 71-248).

0444

Quantitative observations on the pattern of synergid degeneration in barley.

AJBOA. Mogensen, H.L. Baltimore, Md. : Botanical Society of America. American journal of botany. Nov/Dec 1984. v. 71 (10). p. 1448-1451. ill. Includes references. (NAL Call No.: DNAL 450 AM36).

0445

The response of wild oat (*Avena fatua*) and *Avena sterilis* accessions to photoperiod and temperature (Morphology, seed dormancy, growth rate).

Somody, C.N. Nalewaja, J.D.; Miller, S.D. Champaign : Weed Science Society of America. Weed science. Mar 1984. v. 32 (2). p. 206-213. ill. Includes references. (NAL Call No.: 79.8 W41).

(PLANT STRUCTURE)

0446

Structural changes occurring in nuclei of barley root cells in response to a combined effect of salinity and ageing (*Hordeum vulgare*).

Werker, E. AJBOA. Lerner, H.R.; Weimberg, R.; Poljakoff-Mayber, A. Baltimore : Botanical Society of America. American journal of botany. Feb 1983. v. 70 (2). p. 222-225. ill. Includes references. (NAL Call No.: 450 AM36).

0447

A study of stem inflation in wild buckwheat, *Eriogonum inflatum* (Insect damage, anatomy).
Stone, A.M. Mason, C.T. Jr. Superior, Ariz., University of Arizona. Desert plants. Nov 1979. v. 1 (2). p. 77-81. ill. 9 ref. (NAL Call No.: QK938.D4D4).

0448

Weed control in small grains /by F.W. Slife ... et al. .

Slife, F. W. 1923-. Urbana, Ill. : University of Illinois, College of Agriculture, Extension Service in Agriculture and Home Economics. 1950. Cover title. 11 p. : ill., plans ; 23 cm. (NAL Call No.: DNAL 275.29 I162c no.658).

0449

Wild oats: one of the worst weeds.

CRSOA. Puy, D. van der. Madison, Wis. : American Society of Agronomy. Crops and soils magazine. Aug/Sept 1986. v. 38 (9). p. 13-14. ill. (NAL Call No.: DNAL 6 W55).

PLANT NUTRITION

0450

Barley, potato and brome grass chemical composition unchanged by use of a multipurpose wetting agent (The active ingredients of which are alcohol ethoxylates, propylene glycol, and dimethylpolysiloxane, fertilizer, uptake).
Laughlin, W.M. AK-AR-NC. Smith, G.R.; Peters, M.A. Fairbanks, The Station.
Agroborealis. Alaska. Agricultural Experiment Station, Fairbanks. Jan 1980. v. 12 (1). p. 29-30. ill. 7 ref. (NAL Call No.: S33.E2).

0451

Characteristics of boron absorption by barley plants / by Ahmed Abd Elhamid Mohamed Elseewi.
Elseewi, Ahmed Abd Elhamid Mohamed, 1937-.
1968. Thesis (Ph.D.)--University of California, Riverside, 1968. Photocopy. Ann Arbor, Mich. : Univeristy Microfilms, 1970. ix, 102 leaves : ill. ; 21 cm. Bibliography: leaves 96-102. (NAL Call No.: DISS 69-10,096).

0452

The control of nutrient uptake rates in relation to the inorganic composition of plants Memna .
Glass, A.D.M. Siddiqi, M.Y. New York : Praeger.
Advances in plant nutrition. 1984. v. 1. p. 103-147. ill. Includes 168 references. (NAL Call No.: DNAL QK867.A48).

0453

Effect of Cu on the distribution of P, Ca, and Fe in barley plants.
SOSCAK. Brown, J.C. Foy, C.D. Baltimore, Md. : Williams & Wilkins. Soil science. Dec 1964. v. 98 (6). p. 362-370. ill. Includes references. (NAL Call No.: DNAL 56.8 S03).

0454

Effect of heavy metals on the calcium absorption by intact barley roots.
Veltrup, W. New York, Marcel Dekker. Journal of plant nutrition. 1981. v. 3 (1/4). p. 225-231. ill. 17 ref. (NAL Call No.: QK867.J67).

0455

Effect of legume cover crops and tillage on soil water, temperature, and organic matter.
Utomo, M. Blevins, R.L.; Frye, W.W. Ankeny, Iowa : Soil Conservation Society of America, c1987. The role of legumes in conservation tillage systems / J.F. Power, editor. Paper presented at the "National Conference on the Role of Legumes in Conservation Tillage Systems", April 27-29, 1987, University of Georgia, Athens, Georgia. p. 5-6. Includes

references. (NAL Call No.: DNAL SB203.R6).

0456

Effect of manganese and soil pH on the iron content of crops grown on Podzol soils (Hordeum vulgare, barley, Pisum sativum, peas).
Gupta, U.C.JPNUD. New York : Marcel Dekker.
Journal of plant nutrition. 1982. v. 5 (10). p. 1229-1239. 14 ref. (NAL Call No.: QK867.J67).

0457

Effect of NaCl (sodium chloride) on some trace metals in barley (Salinity, interactions, cobalt, lithium, strontium).
Wallace, A. Cha, J.W.; Mueller, R.T. New York, Marcel Dekker. Journal of plant nutrition. 1980. v. 2 (1/2). p. 115-117. ill. 5 ref. (NAL Call No.: QK867.J67).

0458

Effect of nitrogen nutrition on quality of agronomic crops.
Deckard, E.L. Tsai, C.Y.; Tucker, T.C. Madison, Wis. : American Society of Agronomy, 1984. Nitrogen in crop production : proceedings, symposium, 25-27 May, 1982, Sheffield, Alabama / spon. by National Fertilizer Development Center of Tennessee Valley Authority ... et al. ; Roland D. Hauck. Literature review. p. 601-615. Includes references. (NAL Call No.: DNAL S651.N57).

0459

Effect of potassium fertilizers on malting barley infected with common root rot.
AGJ0AT. Timm, C.A. Goos, R.J.; Johnson, B.E.; Sobolik, F.J.; Stack, R.W. Madison, Wis. : American Society of Agronomy. Agronomy journal. Jan/Feb 1986. v. 78 (1). p. 197-200. Includes references. (NAL Call No.: DNAL 4 AM34P).

0460

Effect of potassium nutrition on sink intensity and duration (Cereal grains).
Haeder, H.E. Bern : International Potash Institute, 1980. Physiological aspects of crop productivity : proceedings of the 15th Colloquium of the International Potash Institute held in Wageningen, The Netherlands. p. 185-194. ill. 26 ref. (NAL Call No.: SB185.5.C6 1980).

(PLANT NUTRITION)

0461

The effect of residual copper levels on the nutrition and yield of oats grown in microplots on three organic soils.

CSOSA2. Belanger, A. Levesue, M.; Mathur, S.P. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. Jan 1986. v. 17 (1). p. 85-96. Includes 12 references. (NAL Call No.: DNAL S590.C63).

0462

Effect of sample pretreatment on extractable soil potassium.

CSOSA2. Haby, V.A. Sims, J.R.; Skogley, E.O.; Lund, R.E. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. 1988. v. 19 (1). p. 91-106. Includes references. (NAL Call No.: DNAL S590.C63).

0463

The effect of surface applied soil amendments on barley root growth in an acid subsoil.

CSOSA2. Wright, R.J. Hern, J.L.; Baligar, V.C.; Bennett, D.L. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. Feb 1985. v. 16 (2). p. 179-192. ill. Includes 26 references. (NAL Call No.: DNAL S590.C63).

0464

The effect of thiosulfate on phosphorus availability and uptake by plants.

JPNUDS. Morden, G. Soper, R.; Huzel, V.; Swan, M. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Oct 1986. v. 9 (10). p. 1315-1321. Includes references. (NAL Call No.: DNAL QK867.J67).

0465

Effect of zinc deficiency on the accumulation of boron and other mineral nutrients in barley.

SSSJD4. Graham, R.D. Welch, R.M.; Grunes, D.L.; Cary, E.E.; Norvell, W.A. Madison, Wis. : The Society. Soil Science Society of America journal. May/June 1987. v. 51 (3). p. 652-657. Includes references. (NAL Call No.: DNAL 56.9 S03).

0466

Effect of zinc, phosphorus, and root-zone temperature on nutrient uptake by barley.

SSSJD4. Schwartz, S.M. Welch, R.M.; Grunes, D.L.; Cary, E.E.; Norvell, W.A.; Gilbert, M.D.; Meredith, M.P.; Sanchirico, C.A. Madison, Wis. : The Society. Soil Science Society of America journal. Mar/Apr 1987. v. 51 (2). p. 371-375. ill. Includes references. (NAL Call No.: DNAL 56.9 S03).

0467

Effects of NaCl on metabolic heat evolution rates by barley roots.

PLPHA. Criddle, R.S. Hansen, L.D.; Breidenbach, R.W.; Ward, M.R.; Huffaker, R.C. Rockville, Md. : American Society of Plant Physiologists. The effect of salinity stress on metabolic heat output of barley (*Hordeum vulgare* L.) root tips was measured by isothermal microcalorimetry. Several varieties differing in tolerance to salinity were compared and differences quantified. Two levels of inhibition by increasing salt were found. Following the transition from the initial rate to the first level, inhibition remained at about 50% with further increases in salt concentration up to 150 millimolar. The concentration of salt required to inhibit to this level was cultivar dependent. At higher concentrations (greater than 150 millimolar) of salt, metabolism was further decreased. This decrease was not cultivar dependent. The decreased rate of metabolic heat output at the first transition could be correlated with decreases in uptake of NO₃⁻, NH₄⁺, and Pi that occurred as the salt concentration was increased. The high degree of dependence of the inhibition of metabolic heat output on NaCl concentration points to a highly cooperative reaction responsible for the general inhibition of metabolism and nutrient uptake. The time required to attain the first level of salt inhibition is less than 20 minutes. Inhibition of root tips was not reversible by washing with salt free solutions. In addition to revealing these features of salt inhibition, isothermal microcalorimetry is a promising method for convenient and rapid determination of varietal differences in response to increasing salinity. Plant physiology. May 1989. v. 90 (1). p. 53-58. Includes references. (NAL Call No.: DNAL 450 P692).

0468

Effects of nitrogen rate and placement on two malting barley varieties (Comparisons, Minnesota).

Varvel, G.E. MXMRA. Severson, R.K. St. Paul : The Station. Miscellaneous publication - University of Minnesota, Agricultural Experiment Station. 1983. 1983. (2 rev.). p. 63-64. (NAL Call No.: S1.M52).

0469

Effects of nitrogen rate and placement on two malting barley varieties (in Northwest Minnesota).

Varvel, G.E. Severson, R.K. St. Paul : The Station. Miscellaneous publication - University of Minnesota, Agricultural Experiment Station. 1982. 1982. (2). p. 70-71. (NAL Call No.: S1.M52).

0470

Effects of P and K (phosphorus and potassium) on cereal crop diseases (Mineral deficiencies). Piening, L.J. Atlanta, Ga., Potash & Phosphate Institute. Better crops with plant food. Winter 1981/1982. v. 66. p. 36-37, 39. (NAL Call No.: 6 B46).

0471

The effects of using copper for mitigating histosol subsidence on: 1. The yield and nutrition of oats and lettuce grown on histosols, mineral sublayers, and their mixtures. Levesque, M.P. S.O.S.C.A. Mathur, S.P. Baltimore : Williams & Wilkins. Soil science. Feb 1983. v. 135 (2). p. 88-100. Includes references. (NAL Call No.: 56.8 S03).

0472

Evidence for a specific uptake system for iron phytosiderophores in roots of grasses. PLPHA. Romheld, V. Marschner, H. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Jan 1985. v. 80 (1). p. 175-180. ill. Includes 29 references. (NAL Call No.: DNAL 450 P692).

0473

Exploitation of soil potassium in layered profiles by root systems of cotton and barley. SSSJD4. Gulick, S.H. Cassman, K.G.; Grattan, S.R. Madison, Wis. : The Society. On vermiculitic soils of the San Joaquin Valley, cotton (*Gossypium hirsutum* L.) often responds to K fertilization where other annual crops do not. These soils have markedly higher available K in surface soil than subsoil. A pot study was conducted to compare rooting patterns, K uptake, and dry matter production of cotton and barley (*Hordeum vulgare* L.) in soil profiles with layered K availability. Topsoil (0.15 cmolc K kg⁻¹) was layered to each of six depths above subsoil (0.09 cmolc K kg⁻¹) in 17-L pots with a total soil depth of 45 cm in all treatments. Two irrigation regimes were imposed and nutrients other than K were adequately supplied. For both crops, plant dry matter and K uptake increased linearly with increased topsoil depth but K uptake per unit increase in topsoil depth was 6.5- and 3.6-fold greater by barley than for cotton with frequent and infrequent irrigation, respectively. Increased K uptake per unit increase in topsoil depth reflected coincident root and K distribution: barley root length density (RLD) was 2.7 times greater than RLD of cotton in topsoil layers but little different in subsoil layers. Poor exploitation of topsoil layers by the cotton root system was attributed to greater sensitivity to low soil water potential. A root system with little compensatory root development in the surface soil when the subsoil is low in nutrients may limit K uptake

and crop productivity on layered soils in the San Joaquin Valley and may require management systems designed to promote more congruent root and nutrient distribution. Soil Science Society of America journal. Jan/Feb 1989. v. 53 (1). p. 146-153. ill. Includes references. (NAL Call No.: DNAL 56.9 S03).

0474

The formation of delta-aminolevulinate, a precursor of chlorophyll, in barley and the role of iron (*Hordeum vulgare*, iron requirements). Miller, G.W. Denney, A.; Pushnik, J.; Yu, M.H. New York ; Basel : Marcel Dekker, 1982. Iron nutrition and interactions in plants : Brigham Young University, August 12-14, 1981 / edited by S.D. Nelson ... (et al.). p. 289-300. 25 ref. (NAL Call No.: QK867.J67 v. 5, nos. 4-7).

0475

Fractionation of selenium in barley and rye-grass. JPNUDS. Gissel-Nielsen, G. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Tenth International Plant Nutrition Colloquium," August 4-9, 1986, Beltsville, Maryland. 1987. v. 10 (9/16). p. 2147-2152. ill. Includes references. (NAL Call No.: DNAL QK867.J67).

0476

The functions of calcium in plant nutrition. Hanson, J.B. New York : Praeger. Advances in plant nutrition. 1984. v. 1. p. 149-208. ill. Includes references. (NAL Call No.: DNAL QK867.A48).

0477

Growth and composition of alfalfa fertilized in greenhouse trials with deproteinized juice from low and high saponin alfalfa and from oat herbage (Phytotoxicity). Welch, D.A. Smith, D.; Soberalske, R.M.; Ream, H.W. New York, Marcel Dekker. Journal of plant nutrition. 1979. v. 1 (2). p. 151-170. ill. 15 ref. (NAL Call No.: QK867.J67).

0478

The influence of ammonium and chloride on potassium and nitrate absorption by barley roots depends on time of exposure and cultivar. PLPHA. Bloom, A.J. Finazzo, J. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1986. v. 81 (1). p. 67-69. Includes 22 references. (NAL Call No.: DNAL 450 P692).

(PLANT NUTRITION)

0479

Inoculation effects on growth, nodulation, cytosol components and nitrogen fixation of narrowleaf and hairy vetch (*Rhizobium leguminosarum*, *Vici angustifolia*, *Vici villosa*, *Secale cereale*).

Lynd, J.Q. CSOSA. McNew, R.W. New York : Marcel Dekker. Communications in soil science and plant analysis. 1983. v. 14 (5). p. 411-426. ill. Includes references. (NAL Call No.: S590.C63).

0480

An iron chelating compound released by barley roots in response to Fe-deficiency stress.

JPNUDS. Jolley, V.D. Brown, J.C.; Blaylock, M.U. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Jan 1988. v. 11 (1). p. 77-91. Includes references. (NAL Call No.: DNAL QK867.J67).

0481

Legume growth and residual effects on oat (*Avena sativa* L.) production.

CSOSA2. Guillard, K. Allinson, D.W. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. Apr 1985. v. 16 (4). p. 375-383. Includes 13 references. (NAL Call No.: DNAL S590.C63).

0482

Long term residual effects of applied selenium on the selenium uptake by plants (Cereals, forages, Canada).

Gupta, U.C. Winter, K.A. New York, Marcel Dekker. Journal of plant nutrition. 1981. v. 3 (1/4). p. 493-502. 19 ref. (NAL Call No.: QK867.J67).

0483

Mineral nutrition of plants exposed to salinity stress.

Lauchli, A. Davis : University of California, Davis?, 1981? . A Conference on biosalinity : the problem of salinity in agriculture : a joint conference of Egyptian, Israeli and American scientists, Univ. of California, Davis, September 1-4, 1981 / organized and. p. 63-69. ill. Includes 8 references. (NAL Call No.: DNAL S619.S24C6).

0484

Mutual influence of certain crops in relation to nitrogen.

AGJOAT. Kellerman, K.F. Wright, R.C. Madison, Wis. : American Society of Agronomy. Agronomy journal. July/Oct 1914. v. 6 (4/5). p. 204-210. Includes references. (NAL Call No.: DNAL 4

AM34P).

0485

Nitrification and total nitrogen as affected by crops, fertilizers, and copper sulfate.

AGJOAT. Jensen, C.A. Madison, Wis. : American Society of Agronomy. The accumulation of nitrates in the soils in the Arkansas River Valley in Colorado, which promised in 1910 and 1911 to become a serious factor in sugar beet growth, became less in 1912 and 1913. In the work in 1911, mustard appeared to have some effect in checking the accumulation of nitrate in the field. Copper sulfate at the rate of 100 pounds per acre on fallow was also effective in checking nitrification, reducing the average seasonal accumulation to about 60 percent of the amount found in the check plot. Molasses on fallow decreased nitrification about 25 percent as compared with the check, but the molasses-treated plot showed a little more nitrates than the plots cropped to cane and oats. Manure on fallow gave a slightly higher accumulation of nitrates than the fallow check. The waste lime on fallow caused strong nitrate accumulation, being more effective in this regard than any other treatment. In general, active nitrification did not set in until the first part of June. From then until the middle of July it was strongest and then suddenly decreased and became very feeble until the end of the experiment, August 17, regardless of the field treatments. The differences in the average seasonal accumulations of nitrates could not have been due entirely to the differences in beet yields from the plots. The mustard plot contained less total nitrogen than any of the others. The fallow plots receiving copper sulfate and molasses contained less total nitrogen than the other fallow plots. The fallow plots with waste lime and manure each contained less total nitrogen than the plot fallowed with nothing added. The reverse was true as regards nitrates. In general, there seemed to be an inverse relation between the amounts of total nitrogen. In the work in 1912, in the presence of a vigorously growing beet crop, the only treatment which showed decided increases in nitrification were cyanamid and manure plus ammonium sulfate. Aside from considerable weekly variations, there. Agronomy journal. Jan/Feb 1916. v. 8 (1). p. 10-22. Includes references. (NAL Call No.: DNAL 4 AM34P).

0486

Nitrogen and yield as related to water use of spring barley (*Hordeum vulgare*, seasonal evapotranspiration, New Mexico).

Kallsen, C.E. AGJOAT. Sammis, T.W.; Gregory, E.J. Madison : American Society of Agronomy. Agronomy journal. Jan/Feb 1984. v. 76 (1). p. 59-64. Includes references. (NAL Call No.: 4 AM34P).

0487

Oxygen and carbon dioxide fluxes from barley shoots depend on nitrate assimilation.
 PLPHA. Bloom, A.J. Caldwell, R.M.; Finazzo, J.; Warner, R.L.; Weissbart, J. Rockville, Md. : American Society of Plant Physiologists. A custom oxygen analyzer in conjunction with an infrared carbon dioxide analyzer and humidity sensors permitted simultaneous measurements of oxygen, carbon dioxide, and water vapor fluxes from the shoots of intact barley plants (*Hordeum vulgare* L. cv Steptoe). The oxygen analyzer is based on a calciazirconium sensor and can resolve concentration differences to within 2 microliters per liter against the normal background of 210,000 microliters per liter. In wild-type plants receiving ammonium as their sole nitrogen source or in nitrate reductase-deficient mutants, photosynthetic and respiratory fluxes of oxygen equaled those of carbon dioxide. By contrast, wild-type plants exposed to nitrate had unequal oxygen and carbon dioxide fluxes: oxygen evolution at high light exceeded carbon dioxide consumption by 26% and carbon dioxide evolution in the dark exceeded oxygen consumption by 25%. These results indicate that a substantial portion of photosynthetic electron transport or respiration generates reductant for nitrate assimilation rather than for carbon fixation or mitochondrial electron transport. Plant physiology. Sept 1989. v. 91 (1). p. 352-356. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0488

Peptides related to phytosiderophore secretion by Fe-deficient barley roots.
 JPNUDS. Mori, S. Hachisuka, M.; Kawai, S.; Takagi, S.; Kishi-Nishizawa, N. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Fourth International Symposium on Iron Nutrition and Interactions in Plants," July 6-9, 1987, University of New Mexico, Albuquerque. June/Nov 1988. v. 11 (6/11). p. 653-662. Includes references. (NAL Call No.: DNAL QK867.J67).

0489

Physiological responses of barley leaves of foliar applied urea-ammonium nitrate.
 CRPSAY. Turley, R.H. Ching, T.M. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 987-993. Includes references. (NAL Call No.: DNAL 64.8 C883).

0490

The rate of absorption of nitrate of soda by oats and cotton when applied at different stages of plant growth.
 AGJOAT. Appleton, W.H. Helms, H.B. Madison, Wis. : American Society of Agronomy. Experiments were conducted in the greenhouse to

study the rate of absorption of nitrate nitrogen by oats and cotton when applied as nitrate of soda at different stages of plant growth. The general plan of the experiment was to add nitrate to a series of cultures at a certain stage of growth. Then, at intervals of a few days, part of the cultures were harvested and the amount of nitrate absorbed was determined by plant analysis and also by determining the nitrate content of the soil. Cultures that did not receive nitrate were used as checks to indicate how much nitrogen the plant secured from the soil. With oats these studies were made at four stages of plant growth. With cotton the study was made at three stages of plant growth. The results may be summarized as follows: When sodium nitrate at the rate of 400 pounds per acre was applied to oats 14 days after planting, absorption of the nitrate was very slow for three weeks. After the third week absorption increased and all nitrate was absorbed by the close of the seventh week. When the nitrate was applied to oats at later stages of growth, the rate of absorption was more rapid. Nitrate applied 42, 70, and 92 days after planting was completely absorbed in 20, 14, and 10 days, respectively. With both oats and cotton there was a close correlation between the rate of growth and the rate of nitrate absorption. Sodium nitrate at the rate of 600 pounds per acre was applied to cotton 14, 40, and 61 days after planting. Absorption of the nitrate was complete in 36, 14, and 11 days, respectively. The results of both experiments indicate that the loss of soluble nitrogenous fertilizer by leaching may be reduced by delaying the application until the crop will absorb it rapidly. Agronomy journal. Oct 1925. v. 17 (10). p. 596-605. (NAL Call No.: DNAL 4 AM34P).

0491

Soil fertility and defoliation effects with arrowleaf clover and nitrogen fertilizer equivalence of crimson-arrowleaf clover combinations (*Trifolium vesiculosum*, *Trifolium incarnatum*, *Secale cereale*, rhizobium, nitrogen fixation by forage legumes, Oklahoma).
 Lynd, J.Q. AGJOAT. Hanlon, E.A. Jr.; Odell, G.V. Jr. Madison : American Society of Agronomy. Agronomy journal. Jan/Feb 1984. v. 76 (1). p. 13-16. ill. Includes references. (NAL Call No.: 4 AM34P).

0492

Strategy I and strategy II mechanisms affecting iron availability to plants may be established too narrow or limited.
 JPNUDS. Brown, J.C. Jolley, V.D. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Fourth International Symposium on Iron Nutrition and Interactions in Plants," July 6-9, 1987, University of New Mexico, Albuquerque. June/Nov 1988. v. 11 (6/11). p. 1077-1098. Includes references. (NAL Call No.: DNAL QK867.J67).

(PLANT NUTRITION)

0493

Utilization of phosphorus from barley residues
in calcareous soils /by W.H. Fuller and R.N.
Rogers.

Fuller, Wallace Hamilton. Rogers, R. N. 1927-
Tucson, Ariz. : University of Arizona, 1951.

Cover title. p. 53-77 : ill. ; 23 cm.

Bibliography: p. 77. (NAL Call No.: DNAL 100
Ar4 (2) no.123).

PLANT PHYSIOLOGY AND BIOCHEMISTRY

0494

Accumulation of plastoquinone A during low temperature growth of winter rye (*Secale cereale*, cold acclimation).
Griffith, M. Elfman, B.; Camm, E.L. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Mar 1984. v. 74 (3). p. 727-729. ill. Includes references. (NAL Call No.: 450 P692).

0495

An adaptive response of rye to freezing (Winter hardiness, cryoprotection, *Secale cereale*, *Hordeum vulgare*).
Olien, C.R. Madison : Crop Science Society of America. Crop science. Jan/Feb 1984. v. 24 (1). p. 51-54. ill. Includes references. (NAL Call No.: 64.8 C883).

0496

Alteration of membrane sterols of barley roots by DNP.
JCLBA3. Jackson, P.C. Grunwald, C. New York, N.Y. : Rockefeller University Press. The Journal of cell biology. Includes abstract. Nov 1986. v. 103 (5,pt.2). p. 223a. (NAL Call No.: DNAL 442.8 J828).

0497

Altered growth response to exogenous auxin and gibberellic acid by gravistimulation in pulvini of *Avena sativa*.
PLPHA. Brock, T.G. Kaufman, P.B. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1988. v. 87 (1). p. 130-133. Includes references. (NAL Call No.: DNAL 450 P692).

0498

Aluminum-induced changes in calmodulin.
NASSD. Haug, A. Weis, C. New York, N.Y. : Plenum Press. NATO advanced science institutes series : Series A : Life sciences. In the series analytic: Molecular and cellular aspects of calcium in plant development / edited by A. J. Trewavas. 1986. v. 104. p. 19-25. Includes references. (NAL Call No.: DNAL QH301.N32).

0499

Aluminum tolerance of spring rye inbred lines.
Aniol, A. Hill, R.D.; Larter, E.N. Madison, Wis., Crop Science Society of America. Crop science. Mar/Apr 1980. v. 20 (2). p. 205-208. ill. 11 ref. (NAL Call No.: 64.8 C883).

0500

Analysis of midwinter freezing stress (Barley, hardiness, injuries).
Olien, C.R. Boca Raton, Fla., CRC Press. Analysis and improvement of plant cold hardiness. 1981. 1981. p. 35-59. 65 ref. (NAL Call No.: SB781.A52).

0501

Application of ¹⁴C-labeled herbicides in lysimeter studies.
WEESA6. Fuhr, F. Champaign, Ill. : Weed Science Society of America. Weed science. Presented at the "Symposium on Assessment of Methodology for Field Evaluation of Herbicide Behavior in Soils," Weed Science Society of America, February 9, 1984, Miami, Florida. 1985. v. 33 (suppl.2). p. 11-17. ill. Includes 39 references. (NAL Call No.: DNAL 79.8 W41).

0502

Autoradiographic investigation of the site of betaine synthesis in water-stressed barley leaves (*Hordeum vulgare*).
Giddings, T.H. Hanson, A.D. East Lansing, Mich., The Laboratory. Annual report - Michigan State University, MSU/DOE Plant Research Laboratory. 1981. 1981 (16th). p. 66-67. 2 ref. (NAL Call No.: QK1.M5).

0503

Barley.
AGRYA. Foster, E. Prentice, N. Madison, Wis. : American Society of Agronomy. Agronomy. 1987. (28). p. 337-396. ill. Includes references. (NAL Call No.: DNAL 4 AM392).

0504

Barley, potato and brome grass chemical composition unchanged by use of a multipurpose wetting agent (The active ingredients of which are alcohol ethoxylates, propylene glycol, and dimethylpolysiloxane, fertilizer, uptake).
Laughlin, W.M. AK-AR-NC. Smith, G.R.; Peters, M.A. Fairbanks, The Station. Agroborealis. Alaska. Agricultural Experiment Station, Fairbanks. Jan 1980. v. 12 (1). p. 29-30. ill. 7 ref. (NAL Call No.: S33.E2).

0505

Barley vs. oat companion crops. I. Forage yield and quality response during alfalfa establishment.
CRPSAY. Brink, G.E. Marten, G.C. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 1060-1067. Includes references. (NAL Call No.: DNAL 64.8 C883).

0506

Betaine accumulation: metabolic pathways and genetics.

Hanson, A.D. Grumet, R. New York : Alan R. Liss. UCLA symposia on molecular and cellular biology. Paper presented at the "Symposium on Molecular and Cellular Biology of Plant Stress," April 15-21, 1984, Keystone, Colorado. 1985. v. 22. p. 71-92. ill. Includes references. (NAL Call No.: DNAL QH506.U34).

0507

Betaine synthesis and accumulation in barley during field water stress.

Hitz, W.D. Ladyman, J.A.R.; Hanson, A.D. Madison, Wis., Crop Science Society of America. Crop science. Jan/Feb 1982. v. 22 (1). p. 47-54. ill. Includes 24 ref. (NAL Call No.: 64.8 C883).

0508

Betaine synthesis from radioactive precursors in attached, water-stressed barley leaves.

Hanson, A.D. Scott, N.A. Bethesda, Md., American Society of Plant Physiologists. Plant physiology. Aug 1980. v. 66 (2). p. 342-348. ill. 23 ref. (NAL Call No.: 450 P692).

0509

Buffer capacities of leaves, leaf cells, and leaf cell organelles in relation to fluxes of potentially acidic gases.

PLPHA. Pfan, H. Heber, U. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. June 1986. v. 81 (2). p. 597-602. Includes 38 references. (NAL Call No.: DNAL 450 P692).

0510

Calcium transport in protoplasts isolated from ml-o barley isolines resistant and susceptible to powdery mildew.

PLPHA. Wrona, A.F. Spanswick, R.M.; Aist, J.R. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Dec 1988. v. 88 (4). p. 1157-1162. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0511

Canopy light and tiller mortality in spring barley.

CRPSAY. Lauer, J.G. Simmons, S.R. Madison, Wis. : Crop Science Society of America. Mortality of tillers in barley (*Hordeum vulgare* L.) is an important developmental event affecting spike number and yield potential of the crop. It has traditionally been thought that increased shading of young tillers by the developing

canopy initiates the premature senescence of tillers. A series of field experiments were conducted in 1982, 1983, and 1984 at St. Paul, MN on a Waukegan silt loam soil (fine-silty over sandy or sandy skeletal, mixed, Typic Hapludoll) to test this hypothesis. Degree of shading within the crop canopy was measured in relation to the position of tiller leaves during the vegetative phase of development for three genotypes. The decline in rate of leaf appearance on nonsurviving primary tillers, monitored as an early indicator of tiller senescence, was noted within 3 to 4 wk after crop emergence and before appreciable shading of tillers occurred. This indicates that tiller mortality was not initiated by lack of light. However, after main stem elongation began, tillers soon became heavily shaded suggesting the possibility that shading plays an auxiliary role in the senescence of barley tillers. We discuss the possibility that changes in light quality early in the growing season may be important for initiating the senescence of tillers. Crop science. Mar/Apr. 1989. v. 29 (2). p. 420-424. Includes references. (NAL Call No.: DNAL 64.8 C883).

0512

Cell membrane permeability and ultrastructural effects of difenzoquat on wild oats (*Avena fatua*).

WEESA6. Thai, K.M. Jana, S.; Fowke, L.C. Champaign, Ill. : Weed Science Society of America. Effects of difenzoquat on wild oats grown under controlled environmental conditions were studied. Seedling height and fresh weight were significantly reduced 5 days after postemergence treatment. Dose-dependent increase in cell membrane permeability was detected after a 12-h exposure to the herbicide. Scanning electron microscopy showed normal leaf hairs and cuticular wax but swollen guard cells 10 days after treatment. Ultrastructural changes occurred before the visible symptoms. The primary effect of difenzoquat appears to be the disruption of the tonoplast and plasmalemma. The tonoplast showed greater damage than the plasmalemma. Secondary effects included damage to mitochondria and chloroplasts. Mitochondria were swollen and often ruptured, but the effect did not increase with the duration of exposure. Chloroplasts became spherical in shape, and their contents were also affected. The changes included: accumulation and then disappearance of starch granules, swelling of frets, fusion of granal thylakoids, detachment and then rupture of the outer membrane of the envelope, and clumping of ribosomes. By contrast, natural senescence caused greater injury of the plasmalemma than the tonoplast, a marked increase in size of plastoglobuli, and loss of starch grains without early accumulation. Weed science. Jan 1989. v. 37 (1). p. 98-106. ill. Includes references. (NAL Call No.: DNAL 79.8 W41).

0513

Cereal cyst nematode (*Heterodera avenae*) on oats. II. Early root development and nematode tolerance.

JONEB. Volkmar, K.M. Lake Alfred, Fla. : Society of Nematologists. Journal of nematology. July 1989. v. 21 (3). p. 384-391. Includes references. (NAL Call No.: DNAL QL391.N4J62).

0514

Changes in composition of chlorophylls, carotenoids, and prenylquinones in green seedlings of *Hordeum* (barley) and *Raphanus* (radishes) induced by the herbicide San 6706--an effect possibly antagonistic to phytochrome action.

Kleudgen, H.K. New York, Academic Press. Pesticide biochemistry and physiology. Dec 1979. v. 12 (3). p. 231-238. ill. 36 ref. (NAL Call No.: SB951.P49).

0515

Changes in endogenous gibberellins and the metabolism of (3H)-GA4 after geostimulation in shoots of the oat plant (*Avena sativa*) (Recovery from lodging).

Pharis, R.P. Legge, R.L.; Noma, M.; Kaufman, P.B.; Ghosheh, N.S.; LaCroix, J.D.; Heller, K. Rockville, Md., American Society of Plant Physiologists. Plant physiology. May 1981. v. 67 (5). p. 892-897. ill. 26 ref. (NAL Call No.: 450 P692).

0516

Changes in K, Rb, and Na transport to shoots after anoxia.

PLPHA. Brauer, D. Leggett, J.E.; Egli, D.B. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Jan 1987. v. 83 (1). p. 219-224. Includes references. (NAL Call No.: DNAL 450 P692).

0517

Characteristics of boron absorption by barley plants / by Ahmed Abd Elhamid Mohamed Elseewi.

Elseewi, Ahmed Abd Elhamid Mohamed, 1937-. 1968. Thesis (Ph.D.)--University of California, Riverside, 1968. Photocopy. Ann Arbor, Mich. : University Microfilms, 1970. ix, 102 leaves : ill. ; 21 cm. Bibliography: leaves 96-102. (NAL Call No.: DISS 69-10,096).

0518

Characteristics of cadmium and zinc in four soils treated with sewage sludge.

JEVQAA. Mullins, G.L. Sommers, L.E. Madison, Wis. : American Society of Agronomy. Journal of environmental quality. Oct/Dec 1986. v. 15 (4). p. 382-387. Includes references. (NAL Call No.: DNAL QH540.J6).

0519

Characteristics of injury and recovery of net N03- transport of barley seedlings from treatments of NaCl.

PLPHA. Klobus, G. Ward, M.R.; Huffaker, R.C. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Aug 1988. v. 87 (4). p. 878-882. Includes references. (NAL Call No.: DNAL 450 P692).

0520

The characteristics of secondary dormant wild oat (*Avena fatua* L.) seed.

WSWPA. Fellows, G.M. Fay, P.K.; Foley, M.E. Reno : The Society. Proceedings - Western Society of Weed Science. 1985. v. 38. p. 105-107. (NAL Call No.: DNAL 79.9 W52).

0521

Characterization of the nuclear DNA of *Hordeum vulgare* root hairs: amplification disappears under salt stress.

AJBOAA. Murry, L.E. Christianson, M.L.; Alfinito, S.H.; Garger, S.J. Columbus, Ohio : Botanical Society of America. American journal of botany. Dec 1987. v. 74 (12). p. 1779-1786. Includes references. (NAL Call No.: DNAL 450 AM36).

0522

Cloning and nitrate induction of nitrate reductase mRNA.

PNASAA. Cheng, C.L. Dewdney, J.; Kleinhofs, A.; Goodman, H.M. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. Sept 1986. v. 83 (18). p. 6825-6828. ill. Includes references. (NAL Call No.: DNAL 500 N21P).

0523

A comparative spin-label study of isolated plasma membranes and plasma membranes of whole cells and protoplasts from cold-hardened and nonhardened winter rye.

PLPHA. Windle, J.J. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Dec 1988. v. 88 (4). p. 1388-1396. Includes references. (NAL Call No.: DNAL 450 P692).

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

0524

Comparative systemic translocation of several xenobiotics and sucrose (in soybean and barley).

Martin, R.A. Edgington, L.V. New York, Academic Press. Pesticide biochemistry and physiology. Oct 1981. v. 16 (2). p. 87-96. ill. 19 ref. (NAL Call No.: SB951.P49).

0525

Comparative winter hardiness of barley varieties /by G.A. Wiebe and David A. Reid. Wiebe, G. A.(Gustav A.), 1899-. Reid, David A., 1914-. Washington : U.S. Dept. of Agriculture, 1958. 20 p. : ill., maps ; 23 cm. Includes bibliographical references. (NAL Call No.: DNAL 1 Ag84Te no.1176).

0526

Comparison of selenium treatments of crops in the field.

Gissel-Nielsen, G. Passaic, N.J. : Humana Press. Biological trace element research. Sept 1986. v. 10 (3). p. 209-213. Includes references. (NAL Call No.: DNAL QP534.B56).

0527

A comparison of the effect of salt on polypeptides and translatable mRNAs in roots of a salt-tolerant and a salt-sensitive cultivar of barley.

PLPHA. Hurkman, W.U. Fornari, C.S.; Tanaka, C.K. Rockville, Md. : American Society of Plant Physiologists. The effect of salt stress on polypeptide and mRNA levels in roots of two barley (*Hordeum vulgare* L.) cultivars differing in salt tolerance (cv CM 72, tolerant; cv Prato, sensitive) was analyzed using to-dimensional polyacrylamide gel electrophoresis. Preliminary experiments indicated that germination of Prato was inhibited significantly in the presence of NaCl, but growth of the surviving Prato seedlings was not substantially different from that of CM 72. Fluorographs of two-dimensional gels containing in vivo labeled polypeptides or in vitro translation products were computer analyzed to identify and quantitate changes that resulted when plants were grown in the presence of 200 millimolar NaCl for 6 days. The patterns of in vivo labeled polypeptides and in vitro products of CM 72 and Prato were qualitatively the same. Salt caused quantitative changes in numerous polypeptides and translatable mRNAs, but, overall, the changes were relatively small. Salt did not induce the synthesis of unique polypeptides or translatable mRNAs and did not cause any to disappear. Because of the similarities of the two cultivars with respect to growth and polypeptide patterns and the slight changes in polypeptide and translation product levels caused by salt, specific polypeptides or translatable mRNAs that are related to salt

tolerance in barley could not be identified. Plant physiology. Aug 1989. v. 90 (4). p. 1444-1456. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0528

Comparison of the lipid composition of oat root and coleoptile plasma membranes. Lack of short-term change in response to auxin. PLPHA. Sandstrom, R.P. Cleland, R.E. Rockville, Md. : American Society of Plant Physiologists. The total lipid composition of plasma membranes (PM), isolated by the phase partitioning method from two different oat (*Avena sativa* L.) tissues, the root and coleoptile, was compared. In general, the PM lipid composition was not conserved between these two organs of the oat seedling. Oat roots contained 50 mole percent phospholipid, 25 mole percent glycolipid, and 25 mole percent free sterol, whereas comparable amounts in the coleoptile were 42, 39, and 19 mole percent, respectively. Individual lipid components within each lipid class also showed large variations in the root PM was more than double the activity in the coleoptile. Treatment of coleoptile with auxin for 1 hour resulted in no detectable changes in PM lipids or extractable ATPase activity. Differences in the PM lipid composition between the two tissues that may define the limits of ATPase activity are discussed. Plant physiology. July 1989. v. 90 (3). p. 1207-1213. Includes references. (NAL Call No.: DNAL 450 P692).

0529

Competency for graviresponse in the leaf-sheath pulvinus of *Avena sativa*: onset to loss.

AJBDA. Brock, T.G. Kaufman, P.B. Columbus, Ohio : Botanical Society of America. American journal of botany. Nov 1988. v. 75 (11). p. 1672-1677. Includes references. (NAL Call No.: DNAL 450 AM36).

0530

The control of nutrient uptake rates in relation to the inorganic composition of plants Memna .

Glass, A.D.M. Siddiqi, M.Y. New York : Praeger. Advances in plant nutrition. 1984. v. 1. p. 103-147. ill. Includes 168 references. (NAL Call No.: DNAL QK867.A48).

0531

Coordinator's report: anthocyanin genes. Stock list of ant mutants kept at the Carlsberg laboratory.

Jende-Strid, B. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 74-79. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0532

Correlation between the lipid composition and the responsiveness of Avena sativa stem segments to gibberellic acid (Oats).

Jusaitis, M. Paleg, L.G.; Aspinall, D. Rockville : American Society of Plant Physiologists. Plant physiology. Nov 1982. v. 70 (5). p. 1486-1494. 17 ref. (NAL Call No.: 450 P692).

0533

Correlations between potassium uptake and hydrogen efflux in barley varieties. A potential screening method for the isolation of nutrient efficient lines.

Glass, A.D.M. Siddiqi, M.Y.; Giles, K.I. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Aug 1981. v. 68 (2). p. 457-459. ill. 15 ref. (NAL Call No.: 450 P692).

0534

Crop response to sludge loading rates.

BCYCDK. Day, A.D. Solomon, M.A.; Ottman, M.U.; Taylor, B.B. Emmaus, Pa. : J.G. Press. BioCycle. Aug 1989. v. 30 (8). p. 72-75. Includes references. (NAL Call No.: DNAL 57.8 C734).

0535

Cytokinin biochemistry in relation to leaf senescence. I. The metabolism of 6-benzylaminopurine and zeatin in oat leaf segments (Avena sativa).

Tao, G.Q. Letham, D.S.; Palni, L.M.S.; Summons, R.E. New York : Springer. Journal of plant growth regulation. 1983. v. 2 (2). p. 89-102. Includes references. (NAL Call No.: QK745.J6).

0536

Cytokinin inhibition of respiration in mitochondria from six plant species (Cereals and vegetables).

Miller, C.D. Washington, D.C., The Academy. Proceedings of the National Academy of Sciences of the United States of America. Aug 1980. v. 77 (8). p. 4731-4735. ill. 26 ref. (NAL Call No.: 500 N21P).

0537

Cytokinin-like activities of nucleocyclitols.

JPGRDI. Carceller, M. Cadenas, R.A. New York, N.Y. : Springer. Journal of plant growth regulation. 1988. v. 7 (3). p. 153-159. Includes references. (NAL Call No.: DNAL QK745.J6).

0538

Cytoplasmic male sterility in barley. VIII. Lipoygenase activity and anther amino nitrogen in the msml-Rfm1a system.

Ahokas, H. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Jan 1982. v. 69 (1). p. 268-272. ill. 33 ref. (NAL Call No.: 450 P692).

0539

C4 grasses and cereals growth development, and stress response /C. Allan Jones. --.

Jones, C. Allan. New York : Wiley, c1985. "A Wiley-Interscience publication."~ Includes index. xi, 419 p. : ill. ; 24 cm. Bibliography: p. 333-412. (NAL Call No.: DNAL QK495.G74J66).

0540

Daily growth of the oat kernel and effect on germination of immaturity and controlled low temperatures /Ernest Gordon Booth.

Booth, Ernest Gordon, 1899-. St. Paul : University Farm, 1929? . Cover title.~ Originally presented as: Thesis (Ph. D)--University of Minnesota, 1928. 42 p. : ill., charts ; 23 cm. Bibliography: p. 41-42. (NAL Call No.: DNAL 100 M66 (3) no.62).

0541

Determining vigor of natural and planted stands of sea oats (Uniola paniculata) on the Texas Gulf Coast.

Baker, R.L. Dahl, B.E. Austin, Tex., Southwestern Association of Naturalists. The Southwestern naturalist. May 21, 1981. v. 26 (2). p. 117-123. ill. 17 ref. (NAL Call No.: 409.6 S08).

0542

Development and characterization of friable, embryogenic oat callus.

CRPSAY. Bregitzer, P. Somers, D.A.; Rines, H.W. Madison, Wis. : Crop Science Society of America. Friable, embryogenic (FE) callus cultures are desired for genetic manipulations at the cellular level. The goal of this research was to develop and characterize such cultures for oat (Avena spp.). Nonfriable (NF) oat callus derived from immature embryos was plated on a modified MS medium containing 60 g L-1 sucrose and no hormones. This treatment resulted in the development of distinct somatic embryos that germinated to form complete plants when placed on a modified MS medium containing 20 g L-1 sucrose and no hormones. Embryogenic sectors isolated from 26-wk-old NF callus were visually selected during repeated subculture on a modified MS medium containing 2 mg L-1, 2,4-dichlorophenoxyacetic acid (2,4-D) and 20 g L-1 sucrose to produce FE callus. Transfer of FE callus to a modified MS medium containing 60

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

g L⁻¹ sucrose and no hormones induced the maturation of somatic embryos. Subsequent transfer of this FE callus to modified MS medium containing 20 g L⁻¹ sucrose and no hormones allowed the germination of some of these embryos. Plants were regenerated from FE callus lines for more than 78 wk after FE callus establishment. Friable embryogenic callus also was initiated directly from immature embryos of three genotypes and seedling mesocotyls of two genotypes. Genotypic variation in this response was noted. The FE oat cultures represent an improved callus type for in vitro genetic manipulation of oat. Crop science. May/June 1989. v. 29 (3). p. 798-803. ill. Includes references. (NAL Call No.: DNAL 64.8 C883).

0543

Development at cold-hardening temperatures: membrane assembly and organization.

Huner, N.P.A. Krol, M.; Williams, J.P.; Maissan, E.; Krupa, Z. Boca Raton, FL : CRC Press, c1989. Low temperature stress physiology in crops / editor, Paul H. Li. Literature review. p. 53-65. ill. Includes references. (NAL Call No.: DNAL SB781.L68).

0544

Development at cold-hardening temperatures. The structure and composition of purified rye light harvesting complex II.

PLPHA. Krupa, Z. Huner, N.P.A.; Williams, J.P.; Maissan, E.; James, D.R. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1987. v. 84 (1). p. 19-24. Includes references. (NAL Call No.: DNAL 450 P692).

0545

Development rate and growth duration of oats in response to delayed sowing.

AGJOAT. Colville, D.C. Frey, K.J. Madison, Wis. : American Society of Agronomy. Agronomy journal. May/June 1986. v. 78 (3). p. 417-421. Includes references. (NAL Call No.: DNAL 4 AM34P).

0546

Developmental effects of (the herbicide) Sandoz 6706 on activities of enzymes of phenolic and general metabolism in barley shoots grown in the dark or under low or high intensity light. Blume, D.E. McClure, J.W. Bethesda, Md., American Society of Plant Physiologists. Plant physiology. Feb 1980. v. 65 (2). p. 238-244. ill. 24 ref. (NAL Call No.: 450 P692).

0547

Diallel analyses of some developmental morpho-physiological and agronomic traits on barley, *Hordeum vulgare* L. Emend. Lam. / by Yap, Thoo Chai. -.

Yap, Thoo Chai, 1940-. 1970. Thesis (Ph.D.)--University of Saskatchewan, 1970. Photocopy. Ann Arbor, Mich. : University Microfilms, 1971. xiii, 133 leaves : ill. ; 21 cm. Bibliography: leaves 115-120. (NAL Call No.: DISS 71-20,635).

0548

Diclofop-methyl increases the proton permeability of isolated oat-root tonoplast.

PLPHA. Ratterman, D.M. Balke, N.E. Rockville, Md. : American Society of Plant Physiologists. Diclofop-methyl (methyl ester of 2- 4-(2',4'-dichlorophenoxy)phenoxy propionate; 100 micromolar) and diclofop (100 micromolar) inhibited both ATP- and PPi-dependent formation of H⁺ gradients by tonoplast vesicles isolated from oat (*Avena sativa* L., cv Dal) roots. Diclofop-methyl (1 micromolar) significantly reduced the steady-state H⁺ gradient generated in the presence of ATP. The ester (diclofop-methyl) was more inhibitory than the free acid (diclofop) at pH 7.4, but this relative activity was reversed at pH 5.7. Neither compound affected the rate of ATP or PPi hydrolysis by the proton-pumping enzymes. Diclofop-methyl (50, 100 micromolar), but not diclofop (100 micromolar), accelerated the decay of nonmetabolic H⁺ gradients established across vesicle membranes. Diclofop-methyl (100 micromolar) did not collapse K⁺ gradients across vesicle membranes. Both the (+)- and (-)-enantiomers of diclofop-methyl dissipated nonmetabolic H⁺ gradients established across vesicle membranes. Diclofop-methyl, but not diclofop (each 100 micromolar), accelerated the decay of H⁺ gradients imposed across liposomal membranes. These results show that diclofop-methyl causes a specific increase in the H⁺ permeability of tonoplast. Plant physiology. Oct 1989. v. 91 (2). p. 756-765. Includes references. (NAL Call No.: DNAL 450 P692).

0549

Differences between effects of undissociated and anionic 2, 4-dinitrophenol on permeability of barley roots (*Hordeum vulgare*).

Jackson, P.C. Rockville : American Society of Plant Physiologists. Plant physiology. Nov 1982. v. 70 (5). p. 1373-1379. 21 ref. (NAL Call No.: 450 P692).

0550

Differences in steady-state net ammonium and nitrate influx by cold- and warm-adapted barley varieties.

Bloom, A.J. Chapin, F.S. III. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Nov 1981. v. 68 (5). p. 1064-1067. ill. 24 ref. (NAL Call No.: 450 P692).

0551

Differences in the physical properties of native and partially degraded phytochrome as probed by their differential sensitivity to permanganate oxidation (*Avena sativa*, oats).
Baron, D. PLPHA. Epel, B.L. Rockville : American Society of Plant Physiologists. Plant physiology. Oct 1983. v. 73 (2). p. 471-474. Includes references. (NAL Call No.: 450 P692).

0552

Differential aluminum tolerance of winter barley varieties and selections in associated greenhouse and field experiments.
AGUDAT. Reid, D.A. Jones, G.D.; Armiger, W.H.; Foy, C.D.; Koch, E.J.; Starling, T.M. Madison, Wis. : American Society of Agronomy. Agronomy journal. Mar/Apr 1969. v. 61 (2). p. 218-222. ill. Includes references. (NAL Call No.: DNAL 4 AM34P).

0553

Differential thermal analysis of the freezing of water in leaves of cold-hardened and nonhardened Puma rye (*Secale cereale*).
Lindstrom, D.M. Huner, N.P.A.; Carter, J.V. Chicago : University of Chicago Press. Botanical gazette. June 1983. v. 144 (2). p. 234-239. Includes references. (NAL Call No.: 450 B652).

0554

Diploid-tetraploid comparisons in rye. III. Temperature effects on seedling development.
CRPSAY. Sullivan, B.P. Pfahler, P.L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 795-799. Includes 22 references. (NAL Call No.: DNAL 64.8 C883).

0555

Distribution of potassium between vacuole and cytoplasm in response to potassium deficiency.
NASSD. Storey, R. Leigh, R.A. New York, N.Y. : Plenum Press. NATO advanced science institutes series : Series A : Life sciences. In the series analytic: Plant vacuoles: their importance in solute compartmentation in cells and their applications in plant biotechnology /

edited by B. Marin. Proceedings of a Workshop, July 6-11, 1986, Sophia-Antipolis, France. 1987. 134. p. 95-98. ill. Includes references. (NAL Call No.: DNAL QH301.N32).

0556

Dormant embryo morphology, seedling morphogenesis and tiller development of Himalaya barley / Richard H. Mullenax .
Mullenax, Richard H. Pullman, Wash. : Washington Agricultural Experiment Stations, College of Agriculture, Washington State University, 1967. Cover title. 6 p. : ill. ; 28 cm. Bibliography: p. 6. (NAL Call No.: DNAL 100 W27T no.55).

0557

Drought and freezing tolerance and adaptation in plants: some evidence of near equivalences (*Rye, Secale cereale*).
Siminovitch, D. CRYBA. Cloutier, Y. New York : Academic Press. Cryobiology. Aug 1983. v. 20 (4). p. 487-503. ill. Includes references. (NAL Call No.: QH324.C7).

0558

Early effects of salinity on nitrate assimilation in barley seedlings.
PLPHA. Aslam, M. Huffaker, R.C.; Rains, D.W. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Oct 1984. v. 76 (2). p. 321-325. ill. Includes 31 references. (NAL Call No.: DNAL 450 P692).

0559

Effect of abscisic acid on ion transport and translocation in *Hordeum vulgare* seedlings.
TISAA. Riedell, W.E. Schmid, W.E. Springfield : The Academy. Transactions of the Illinois State Academy of Science. Includes list of species. 1984. v. 77 (3/4). p. 241-247. Includes references. (NAL Call No.: DNAL 500 IL6).

0560

Effect of calcium and phosphorus content of various soil series in western Washington upon the calcium and phosphorus composition of oats, red clover and white clover / by Henry F. Holtz.
Holtz, Henry F., 1880-1931. Pullman, Wash. : State College of Washington, Agricultural Experiment Station, 1930. Cover title. 45 p. ; 23 cm. Bibliography: p. 31-33. (NAL Call No.: DNAL 100 W27E no.243).

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

0561

Effect of cold acclimation on intracellular ice formation in isolated protoplasts (*Secale cereale*, rye).

Dowgert, M.F. PLPHA. Steponkus, P.L. Rockville : American Society of Plant Physiologists. Plant physiology. Aug 1983. v. 72 (4). p. 978-988. Includes references. (NAL Call No.: 450 P692).

0562

Effect of Cu on the distribution of P, Ca, and Fe in barley plants.

SOSCAK. Brown, J.C. Foy, C.D. Baltimore, Md. : Williams & Wilkins. Soil science. Dec 1964. v. 98 (6). p. 362-370. ill. Includes references. (NAL Call No.: DNAL 56.8 S03).

0563

The effect of ethephon (2-chloroethyl) phosphonic acid on lodging and yield of small grain cereal crops.

Schwartz, T.K. PPGGD. Coffin, R.H.; Evans, W.F.; Nash, R.L.; Rao, K.P. Lake Alfred : The Society. Proceedings annual meeting - Plant Growth Regulator Society of America. 1982. (9th). p. 89-95. Includes references. (NAL Call No.: SB128.P5).

0564

Effect of genetically and environmentally induced heading date differences on yield and adaptation of an isogenic barley pair.

CRPSAY. Smail, V.W. Eslick, R.F.; Hockett, E.A. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 889-893. Includes references. (NAL Call No.: DNAL 64.8 C883).

0565

Effect of heavy metals on the calcium absorption by intact barley roots.

Veltrup, W. New York, Marcel Dekker. Journal of plant nutrition. 1981. v. 3 (1/4). p. 225-231. ill. 17 ref. (NAL Call No.: QK867.J67).

0566

Effect of inorganic orthophosphate on in vitro activity of NADH-nitrate reductase isolated from 2-row barley leaves.

PLPHA. Oji, Y. Ryoma, Y.; Wakiuchi, N.; Okamoto, S. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Mar 1987. v. 83 (3). p. 472-474. Includes references. (NAL Call No.: DNAL 450 P692).

0567

Effect of leaf extracts of mustard and barley on growth behavior of some phylloplane microfungi (*Brassica campestris*, *Hordeum vulgare*).

Singh, D.B. Rai, B. Bronx, N.Y., The Club. Bulletin of the Torrey Botanical Club. Oct/Dec 1981. v. 108 (4). p. 419-421. Includes 10 ref. (NAL Call No.: 451 T63B).

0568

Effect of nitrogen nutrition on endosperm protein synthesis in wild and cultivated barley grown in spike culture.

PLPHA. Corke, H. Atsmon, D. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. June 1988. v. 87 (2). p. 523-528. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0569

The effect of sigma-aminolevulinic acid on red-light-induced leaf unrolling (Barley).

Sundqvist, C. Briggs, W.R. Washington, D.C., The Institute. Year book - Carnegie Institution of Washington. 1979/1980. 1979/1980. (79th). p. 136-138. ill. 4 ref. (NAL Call No.: 500 C21).

0570

Effect of triadimefon and triadimenol on growth of various plant species as well as on gibberellin content and sterol metabolism in shoots of barley seedlings (Fungicides, retardation).

Buchenauer, H. Rohner, E. New York, Academic Press. Pesticide biochemistry and physiology. Feb 1981. v. 15 (1). p. 58-70. ill. 36 ref. (NAL Call No.: SB951.P49).

0571

Effect of vanadate on bean leaf movement, stomatal conductance, barley leaf unrolling, respiration, and phosphatase activity.

Saxe, H. Rajagopal, R. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Oct 1981. v. 68 (4). p. 880-884. 24 ref. (NAL Call No.: 450 P692).

0572

Effect of volatile compounds, nutrients, and source of sclerotia on eruptive sclerotial germination of *Sclerotium rolfsii*.

PHYTAJ. Punja, Z.K. Jenkins, S.F.; Grogan, R.G. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Nov 1984. v. 74 (11). p. 1290-1295. ill. Includes 29 references. (NAL Call No.: DNAL 464.8 P56).

0573

Effect on buckwheat protein quality of seed germination and changes in trypsin inhibitor content (*Fagopyrum esculentum*). Ikeda, K. Arioka, K.; Fujii, S.; Kusano, T.; Oku, M. St. Paul, Minn. : American Association of Cereal Chemists. Cereal chemistry. May/June 1984. v. 61 (3). p. 236-238. Includes references. (NAL Call No.: 59.8 C33).

0574

Effectiveness of utilizing various leaf characteristics as selection criteria in breeding for drought tolerance in barley / by Marvin L. Boerboom. -. Boerboom, Marvin L. Ann Arbor, Mich. University Microfilms International 1978. Thesis--North Dakota State University, 1977. Facsimile produced by microfilm-xerography. viii, 128 leaves. Bibliography: leaves 92-99. (NAL Call No.: DISS 77-29,178).

0575

Effects of ambient temperature and nitrogen supply on content of four nitrogen fractions in barley plants / by Ikbalur Rashid Chowdhury. -. Chowdhury, Ikbalur Rashid, 1939-. 1970. Thesis (Ph.D.)--North Dakota State University, 1970. Photocopy. Ann Arbor, Mich. : University Microfilms, 1971. xiii, 204 leaves ; 21 cm. Bibliography: leaves 162-175. (NAL Call No.: DISS 71-5,508).

0576

The effects of anions on PPI-dependent proton transport in tonoplast vesicles from oat roots. NASSD. Pope, A.J. Leigh, R.A. New York, N.Y. : Plenum Press. NATO advanced science institutes series : Series A : Life sciences. In the series analytic: Plant vacuoles: their importance in solute compartmentation in cells and their applications in plant biotechnology / edited by B. Marin. Proceedings of a Workshop, July 6-11, 1986, Sophia-Antipolis, France. 1987. 134. p. 199-204. Includes references. (NAL Call No.: DNAL QH301.N32).

0577

The effects of benzyladenine, cycloheximide, and cordycepin on wilting-induced abscisic acid and proline accumulations and abscisic acid- and salt-induced proline accumulation in barley leaves. PLPHA. Stewart, C.R. Voetberg, G.; Rayapati, P.J. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Nov 1986. v. 82 (3). p. 703-707. Includes references. (NAL Call No.: DNAL 450 P692).

0578

The effects of cytokinins of auxin and growth in tobacco tissue cultures and in oat seedlings and a search for an auxin-transfer RNA complex / by William R. Jordon. -. Jordon, William R. (William Roy), 1944-. 1971. Thesis (Ph.D.)--University of Wisconsin, 1971. Photocopy. Ann Arbor, Mich. : University Microfilms, 1972. vi, 153 leaves ; 21 cm. Includes bibliographies. (NAL Call No.: DISS 71-25,728).

0579

Effects of different endomycorrhizal fungi on five host plants grown on calcined montmorillonite clay (Apple, asparagus, leek, strawberry, oats). Plenchette, C. Furian, V.; Fortin J.A. Alexandria, Va., The Society. Journal of the American Society for Horticultural Science. July 1982. v. 107 (4). p. 535-538. 16 ref. (NAL Call No.: 81 S012).

0580

Effects of DNP on root membrane lipids (Barley). Jackson, P.C. St John, J.B. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Apr 1981. Abstract only. v. 67 (4). p. 65. (NAL Call No.: 450 P692).

0581

Effects of NaCl on metabolic heat evolution rates by barley roots. PLPHA. Criddle, R.S. Hansen, L.D.; Breidenbach, R.W.; Ward, M.R.; Huffaker, R.C. Rockville, Md. : American Society of Plant Physiologists. The effect of salinity stress on metabolic heat output of barley (*Hordeum vulgare* L.) root tips was measured by isothermal microcalorimetry. Several varieties differing in tolerance to salinity were compared and differences quantified. Two levels of inhibition by increasing salt were found. Following the transition from the initial rate to the first level, inhibition remained at about 50% with further increases in salt concentration up to 150 millimolar. The concentration of salt required to inhibit to this level was cultivar dependent. At higher concentrations (greater than 150 millimolar) of salt, metabolism was further decreased. This decrease was not cultivar dependent. The decreased rate of metabolic heat output at the first transition could be correlated with decreases in uptake of NO_3^- , NH_4^+ , and P_i that occurred as the salt concentration was increased. The high degree of dependence of the inhibition of metabolic heat output on NaCl concentration points to a highly cooperative reaction responsible for the general inhibition of metabolism and nutrient uptake. The time required to attain the first level of salt inhibition is less than 20 minutes. Inhibition of root tips was not

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

reversible by washing with salt free solutions. In addition to revealing these features of salt inhibition, isothermal microcalorimetry is a promising method for convenient and rapid determination of varietal differences in response to increasing salinity. *Plant physiology*. May 1989. v. 90 (1). p. 53-58. Includes references. (NAL Call No.: DNAL 450 P692).

0582

Effects of NaCl (sodium chloride) on proline synthesis and utilization in excised barley leaves (*Hordeum vulgare*).

Buhl, M.B. PLPHA. Stewart, C.R. Rockville : American Society of Plant Physiologists. *Plant physiology*. July 1983. v. 72 (3). p. 664-667. Includes references. (NAL Call No.: 450 P692).

0583

Effects of open-air fumigation with sulphur dioxide on the growth and yield of winter barley.

NEPHA. McLeod, A.R. Roberts, T.M.; Alexander, K.; Cribb, D.M. New York, N.Y. : Cambridge University Press. *The New phytologist*. Literature review. May 1988. v. 109 (1). p. 67-78. ill. Includes references. (NAL Call No.: DNAL 450 N42).

0584

The effects of phosphate on early growth and maturity.

AGJOAT. Noll, C.F. Madison, Wis. : American Society of Agronomy. A number of investigators have found that the use of phosphates produced a more rapid growth of the roots of seedlings and several have claimed that, in the case of small grains, phosphates promote tillering. Russell states that in England superphosphates cause rapid early growth of turnips and swedes. The evidence is not conclusive that these effects from phosphate are more pronounced than from the other fertilizer elements where the latter are the limiting factors for crop yields. Fertilizer tests have quite generally, if not always, shown that the use of phosphatic fertilizers induces earlier ripening of the grain crops on soils low in phosphorus. Similar effects have been noted with cabbage and with cotton. In the case of tomatoes, the use of phosphates at the Pennsylvania Experiment Station, has been accompanied by a much greater growth of stalk and larger total yield, but also by later ripening. The effects of different phosphatic fertilizers seem to vary with the availability of the phosphates as shown by the responses in crop yields. However, at the Ohio Experiment Station, acid phosphate had more influence in promoting earliness than either steamed bonemeal or basic slag, altho all these phosphates gave approximately the same yield. Applied in moderate rates, the soluble phosphates, as a rule, have shown a more pronounced influence in hastening maturity

than the same quantity of phosphoric acid in rock phosphate, the influence in earliness varying in degree with the response in yield. Increasing the rates of application of phosphatic fertilizers above the needs of the crop as indicated by yields has not been accompanied by further increases in earliness. Though phosphates show a more marked influence in promoting earliness than the other fertilizer elements, yet moderate amounts of nitrogen and potash, where these are needed for the production of crops, have a similar effect. Lime, also, if applied to soils having a high lime requirement, may induce earli. *Agronomy journal*. Mar 1923. v. 15 (3). p. 87-99. (NAL Call No.: DNAL 4 AM34P).

0585

Effects of powdery mildew (*Erysiphe graminis* f.sp. *hordei*) and water stress on CO₂ (carbon dioxide) exchange in uninfected leaves of barley.

Williams, G.M. Ayres, P.G. Rockville, Md., American Society of Plant Physiologists. *Plant physiology*. Sept 1981. v. 68 (3). p. 527-530. 25 ref. (NAL Call No.: 450 P692).

0586

The effects of salt on the pattern of protein synthesis in barley roots.

PLPHA. Hurkman, W.J. Tanaka, C.K. Rockville, Md. : American Society of Plant Physiologists. *Plant physiology*. Mar 1987. v. 83 (3). p. 517-524. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0587

The effects of salt stress on polypeptides in membrane fractions from barley roots.

PLPHA. Hurkman, W.J. Tanaka, C.K.; DuPont, F.M. Rockville, Md. : American Society of Plant Physiologists. *Plant physiology*. Dec 1988. v. 88 (4). p. 1263-1273. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0588

Effects of seed moisture content, gamma rays, and unmoderated fission neutrons on various biological end-points in barley, *Hordeum vulgare* L. emend Lam. / by Gurubasavappa Shivashankar. -.

Shivashankar, Gurubasavappa, 1934-. 1971. Thesis (Ph.D.)--University of Tennessee, 1971. Photocopy of typescript. Ann Arbor: University Microfilms, 1972. xvi, 119 leaves ; 21 cm. Bibliography: leaves 90-95. (NAL Call No.: DISS 72-15,549).

0589

Effects of seeding time on lipid content and fatty acid composition of buckwheat grains.
JAFCAU. Taira, H. Akimoto, I.; Miyahara, T. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. Jan/Feb 1986. v. 34 (1). p. 14-17. Includes references. (NAL Call No.: DNAL 381 J8223).

0590

Effects of soil moisture on the vegetative growth of wild oat (*Avena fatua*) (Water stress, stomatal diffusion, Manitoba).
Akey, W.C. Morrison, I.N. Champaign, Ill. : Weed Science Society of America. Weed science. Sept 1984. v. 32 (5). p. 625-630. ill. Includes 14 references. (NAL Call No.: 79.8 W41).

0591

Effects of sulfur supply on the yield, composition, and quality of grain from cereals, oilseeds, and legumes.
ACSTD. Randall, P.J. Wrigley, C.W. St. Paul, Minn. : American Association of Cereal Chemists. Advances in cereal science and technology. Literature review. 1986. v. 8. p. 171-206. ill. Includes references. (NAL Call No.: DNAL TS2120.A3).

0592

Effects of tabtoxinine-beta-lactam on nitrogen metabolism in *Avena sativa* L. roots.
PLPHA. Knight, T.U. Durbin, R.D.; Langston-Unketer, P.J. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Dec 1986. v. 82 (4). p. 1045-1050. Includes references. (NAL Call No.: DNAL 450 P692).

0593

Effects of tiller removal on spring barley.
CRPSAY. Chafai El Alaoui, A. Simmons, S.R.; Crookston, R.K. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 305-307. Includes references. (NAL Call No.: DNAL 64.8 C883).

0594

Effects of 2,4-dinitrophenol on membrane lipids of roots (*Hordeum vulgare* var. *trebi*).
Jackson, P.C. St John, J.B. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1982. v. 70 (3). p. 858-862. 20 ref. (NAL Call No.: 450 P692).

0595

Efficacy of postharvest herbicides on Russian thistle (*Salsola iberica*) control and seed germination.
WEESA6. Young, F.L. Whitesides, R.E. Champaign, Ill. : Weed Science Society of America. Weed science. July 1987. v. 35 (4). p. 544-559. Includes references. (NAL Call No.: DNAL 79.8 W41).

0596

End product storage in cereals.
Jenner, C.F. New York : Alan R. Liss. Plant biology. In the series analytic: Phloem Transport / edited by J. Cronshaw, W.J. Lucas and R.T. Giaquinta. Proceedings of an International Conference, August 18-23, 1985, Asilomar, California.~ Literature review. 1986. v. 1. p. 561-572. Includes references. (NAL Call No.: DNAL QH301.P535).

0597

Enhancement of nitrate uptake and growth of barley seedlings by calcium under saline conditions.
PLPHA. Ward, M.R. Aslam, M.; Huffaker, R.C. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Feb 1986. v. 80 (2). p. 520-524. Includes 28 references. (NAL Call No.: DNAL 450 P692).

0598

Estimates of free and bound indole-3-acetic acid and zeatin levels in relation to regulation of apical dominance and tiller release in oat shoots (*Avena sativa*).
Harrison, M.A.JPGRDI. Kaufman, P.B. New York : Springer. Journal of plant growth regulation. 1983. v. 2 (3). p. 215-223. ill. Includes references. (NAL Call No.: QK745.U6).

0599

Ethylene as an effector of wound-induced resistance to cellulase in oat leaves (*Avena sativa*).
Geballe, G.T. Galston, A.W. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1982. v. 70 (3). p. 788-790. 31 ref. (NAL Call No.: 450 P692).

0600

Evaluation of chlorophyll fluorescence parameters for an intact-plant herbicide bioassay.
CRPSAY. Shaw, D.R. Peeper, T.F.; Nofziger, D.L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 756-760. Includes references. (NAL Call

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

No.: DNAL 64.8 C883).

0601

Evaluation of varieties and selections of barley for disease resistance and winter hardiness in southeastern United States /by J.G. Moseman.

Moseman, J. G. 1921-. Washington : U.S. Dept. of Agriculture, 1956. 33 p. ; 23 cm. Literature cited: p. 32-33. (NAL Call No.: DNAL 1 Ag84Te no.1152).

0602

Evidence for a specific uptake system for iron phytosiderophores in roots of grasses.

PLPHA. Romheld, V. Marschner, H. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Jan 1985. v. 80 (1). p. 175-180. ill. Includes 29 references. (NAL Call No.: DNAL 450 P692).

0603

Examination of North American bison saliva for potential plant growth regulators (Oats, cucumber).

Detling, J.K. Ross, C.W.; Walmsley, M.H.; Hilbert, D.W.; Bonilla, C.A.; Dyer, M.I. New York, Plenum Press. Journal of chemical ecology. Mar 1981. v. 7 (2). p. 239-246. ill. 27 ref. (NAL Call No.: QD415.A1J6).

0604

Expression of hydrogenase activity in barley (Hordeum vulgare L.) after anaerobic stress.

ABBIA. Torres, V. Ballesteros, A.; Fernandez, V.M. New York, N.Y. : Academic Press. Archives of biochemistry and biophysics. Feb 15, 1986. v. 245 (1). p. 174-178. Includes 21 references. (NAL Call No.: DNAL 381 AR2).

0605

The extent to which weeds modify the transpiration of cereals /by A.L. Bakke and H.H. Plagge.

Bakke, Arthur Laurence, b. 1886. Plagge, H. H. 1894-. Ames, Iowa : Agricultural Experiment Station, Iowa State College of Agriculture and Mechanic Arts, 1926. p. 212-239 : charts ; 22 cm. Bibliography: p. 238-239. (NAL Call No.: DNAL 100 Io9 no.96).

0605

Factors affecting survival of winter oats / by Franklin A. Coffman .

Coffman, Franklin A. 1892-. Washington : U.S. Dept. of Agriculture, 1965. 28 p. : 1 map ; 23 cm. Literature cited: p. 27-28. (NAL Call No.: DNAL 1 Ag84Te no.1346).

0607

Factors of wild oat (Avena fatua) interference on spring barley (Hordeum vulgare) growth and yield.

WEESA6. Morishita, D.W. Thill, D.C. Champaign, Ill. : Weed Science Society of America. Weed science. Jan 1988. v. 36 (1). p. 37-42. Includes references. (NAL Call No.: DNAL 79.8 W41).

0608

Fate of tagged urea N (nitrogen) in the field with different methods of N and organic matter placement (Broadcast and banded, barley).

Tomar, J.S. Soper, R.J. Madison, Wis., American Society of Agronomy. Agronomy journal. Nov/Dec 1981. v. 73 (6). p. 991-995. 18 ref. (NAL Call No.: 4 AM34P).

0609

Field evaluation of seedling root length selection in oats (Avena sativa, drought stress).

Barbour, N.W. Murphy, C.F. Madison : Crop Science Society of America. Crop science. Jan/Feb 1984. v. 24 (1). p. 165-169. Includes references. (NAL Call No.: 64.8 C883).

0610

Fluorescence properties indicate that photosystem II reaction centers and light-harvesting complex are modified by low temperature growth in winter rye.

PLPHA. Griffith, M. Huner, N.P.A.; Kyle, D.J. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Oct 1984. v. 76 (2). p. 381-385. ill. Includes 26 references. (NAL Call No.: DNAL 450 P692).

0611

Fluorescence induction transients in the process of development and senescence of barley leaves during prolonged influence of blue or red light.

SOPPA. Rozhkovskii, A.D. Bukhov, N.G.; Voskresenskaya, N.P. New York, N.Y. : Consultants Bureau. Soviet plant physiology. Translated from: Fiziologiya rastenii, p. 1046-1054. (450 F58). Nov/Dec 1985 (pub. 1986). v. 32 (6, pt. 1). p. 795-801. Includes 17

references. (NAL Call No.: DNAL 450 F58AE).

0612

The formation of delta-aminolevulinate, a precursor of chlorophyll, in barley and the role of iron (*Hordeum vulgare*, iron requirements).

Miller, G.W. Denney, A.; Pushnik, J.; Yu, M.H. New York ; Basel : Marcel Dekker, 1982. Iron nutrition and interactions in plants : Brigham Young University, August 12-14, 1981 / edited by S.D. Nelson ... (et al.). p. 289-300. 25 ref. (NAL Call No.: QK867.J67 v. 5, nos. 4-7).

0613

Fractionation of selenium in barley and rye-grass.

JPNUDS. Gissel-Nielsen, G. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Tenth International Plant Nutrition Colloquium," August 4-9, 1986, Beltsville, Maryland. 1987. v. 10 (9/16). p. 2147-2152. ill. Includes references. (NAL Call No.: DNAL QK867.J67).

0614

Freeze-induced changes in soluble carbohydrates of rye.

CRPSAY. Olien, C.R. Lester, G.E. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1985. v. 25 (2). p. 288-290. Includes 9 references. (NAL Call No.: DNAL 64.8 C883).

0615

Freezing resistance and cold acclimation in turfgrasses (*Perennial rye*, *Lolium perenne*, *Kentucky bluegrass*, *Poa pratensis*, *chewings fescue*, *Festuca rubra*).

Rajashekar, C.HJHSA. Tao, D.; Li, P.H. Alexandria : American Society for Horticultural Science. HortScience. Feb 1983. v. 18 (1). p. 91-93. ill. Includes references. (NAL Call No.: SB1.H6).

0616

The functions of calcium in plant nutrition. Hanson, J.B. New York : Praeger. Advances in plant nutrition. 1984. v. 1. p. 149-208. ill. Includes references. (NAL Call No.: DNAL QK867.A48).

0617

Genetics, phenotypes, agronomic and malting performance of glossy sheath mutants in barley, *Hordeum vulgare* L. / by Wayne Lucas McProud. -. McProud, Wayne Lucas, 1945-. 1971. Thesis (Ph.D.)--Montana State University, 1971. Photocopy. Ann Arbor, Mich. : University Microfilms, 1972. ix, 72 leaves ; 21 cm. Bibliography: leaves 70-72. (NAL Call No.: DISS 72-8,888).

0618

Genotype X year interaction for length and rate of grain filling in oats (Environmental stresses).

Wych, R.D. McGraw, R.L.; Stuthman, D.D. Madison, Crop Science Society of America. Crop science. Sept/Oct 1982. v. 22 (5). p. 1025-1028. 20 ref. (NAL Call No.: 64.8 C883).

0619

Genotypic variability in response of oat to delayed sowing.

AGJQAT. Baltenberger, D.C.C. Frey, K.J. Madison, Wis. : American Society of Agronomy. Agronomy journal. Sept/Oct 1987. v. 79 (5). p. 813-816. Includes references. (NAL Call No.: DNAL 4 AM34P).

0620

Genotypic variation for glycinebetaine accumulation by cultivated and wild barley in relation to water stress.

Ladyman, J.A.R. CRPSA. Ditz, K.M.; Grumet, R.; Hanson, A.D. Madison : Crop Science Society of America. Crop science. May/June 1983. v. 23 (3). p. 465-468. ill. Includes references. (NAL Call No.: 64.8 C883).

0621

Germination and emergence of little barley (*Hordeum pusillum*).

Fischer, M.L. Stritzke, J.F.; Ahring, R.M. Champaign : Weed Science Society of America. Weed science. Nov 1982. v. 30 (6). p. 624-628. 16 ref. (NAL Call No.: 79.8 W41).

0622

Grain-filling duration and yield in spring barley.

CRPSAY. Metzger, D.D. Czaplowski, S.J.; Rasmusson, D.C. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1984. v. 24 (6). p. 1101-1105. Includes 11 references. (NAL Call No.: DNAL 64.8 C883).

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

0623

Gramine accumulation in leaves of barley during high-temperature stress (*Hordeum vulgare*).
Hanson, A.D. Ditz, K.M.; Singletary, G.W.; Leland, T.; Reicosky, D.A. East Lansing, Mich., The Laboratory. Annual report - Michigan State University, MSU/DOE Plant Research Laboratory. 1981. 1981 (16th). p. 69-70. 5 ref. (NAL Call No.: QK1.M5).

0624

Gramine accumulation in leaves of barley grown under high-temperature stress (*Hordeum vulgare*).

Hanson, A.D. PLPHA. Ditz, K.M.; Singletary, G.W.; Leland, T.J. Rockville : American Society of Plant Physiologists. Plant physiology. Apr 1983. v. 71 (4). p. 896-904. ill. Includes references. (NAL Call No.: 450 P692).

0625

Gramine (indole alkaloid) in barley forage--effects of genotype and environment.

Hanson, A.D. Traynor, P.L.; Ditz, K.M.; Reicosky, D.A. Madison, Wis., Crop Science Society of America. Crop science. Sept/Oct 1981. v. 21 (5). p. 726-730. ill. 21 ref. (NAL Call No.: 64.8 C883).

0626

Growth and yield of barley isopopulations differing in solute potential.

CRPSAY. Grumet, R. Albrechtsen, R.S.; Hanson, A.D. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1987. v. 27 (5). p. 991-995. Includes references. (NAL Call No.: DNAL 64.8 C883).

0627

The growth of subterranean clover established with barley / by Anthony Alexander Mc Gowan. -

Mc Gowan, Anthony Alexander, 1932-. 1970. Thesis (Ph.D.)--University of California, Davis, 1970. Photocopy. Ann Arbor, Mich. : University Microfilms, 1971. ix, 194 leaves ; 21 cm. Bibliography: leaves 158-169. (NAL Call No.: DISS 71-15,543).

0628

Growth stages in cereals--Zadoks and Feekes comparison (Chart showing comparison of scales).

Atlanta : Potash & Phosphate Institute. Better crops with plant food. Winter 1983/1984. v. 68. p. 11. (NAL Call No.: 6 B46).

0629

Heat shock causes destabilization of specific mRNAs and destruction of endoplasmic reticulum in barley aleurone cells.

PNASA. Belanger, F.C. Brodl, M.R.; Ho, T.H.D. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. Mar 1986. v. 83 (5). p. 1354-1358. ill. Includes references. (NAL Call No.: DNAL 500 N21P).

0630

Heavy metal inhibition of barley root plasma membrane-bound Ca (calcium) -ATPase and its reversal by monovalent cations (*Hordeum vulgare*).

Caldwell, C.R. Suhayda, C.G.; Haug, A. East Lansing, Mich., The Laboratory. Annual report - Michigan State University, MSU/DOE Plant Research Laboratory. 1981. 1981 (16th). p. 72-73. (NAL Call No.: QK1.M5).

0631

Hordein gene expression in a low protein barley cultivar.

PLPHA. Dailey, J.E. Peterson, D.M.; Osborn, T.C. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Oct 1988. v. 88 (2). p. 450-453. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0632

Hordein mutants for barley Hrd loci.

Netsvetaev, V.P. Sozinov, A.A. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Oct 10, 1985. v. 15. p. 51-55. ill. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0633

Hormonal regulation of alpha-amylase gene transcription in wild oat (*Avena fatua* L.) aleurone protoplasts.

PLPHA. Zwar, J.A. Hooley, R. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Feb 1986. v. 80 (2). p. 459-463. ill. Includes 25 references. (NAL Call No.: DNAL 450 P692).

0634

Hormonal regulation of lateral bud (tiller) release in oats (*Avena sativa* L.).

Harrison, M.A. Kaufman, P.B. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Dec 1980. v. 66 (6). p. 1123-1127. ill. 19 ref. (NAL Call No.: 450 P692).

0635

Hormonal regulation of protein synthesis in barley aleurone layers / by Warren H. Evins. -. Evins, Warren H., 1944-. 1971. Thesis (Ph.D.)--Michigan State University, 1971. Photocopy. Ann Arbor, Mich. : University Microfilms, 1972. ix, 92 leaves ; 21 cm. Bibliography: leaves 86-92. (NAL Call No.: DISS 71-23,182).

0636

Hormonal regulation of the development of protease and carboxypeptidase activities in barley aleurone layers. PLPHA. Hammerton, R.W. Ho, T.H.D. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Mar 1986. v. 80 (3). p. 692-697. ill. Includes 26 references. (NAL Call No.: DNAL 450 P692).

0637

How cereal grass shoots perceive and respond to gravity. AUBOA. Kaufman, P.B. Brock, T.G.; Song, I.; Rho, Y.B.; Ghosheh, N.S. Baltimore, Md. : Botanical Society of America. American journal of botany. Sept 1987. v. 74 (9). p. 1446-1457. ill. Includes references. (NAL Call No.: DNAL 450 AM36).

0638

Hypochlorite disinfection influences the GA3-induced synthesis of alpha-amylase isozymes by isolated barley aleurone layers. PLPHA. Tittle, F.T. Goudey, J.S.; Spencer, M.S. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Feb 1988. v. 86 (2). p. 510-511. Includes references. (NAL Call No.: DNAL 450 P692).

0639

Illumination of eight-day-old dark-grown barley seedlings activates chloroplast protein synthesis; evidence for regulation of translation initiation. Gamble, P.E. Klein, R.R.; Mullet, J.E. New York, N.Y. : Alan R. Liss. Plant biology. In the series analytic: Photosynthesis / edited by W.R. Briggs. Proceedings of the C.S. French Symposium, July 17-23, 1988, Stanford, California. 1989. v. 8. p. 285-298. ill. Includes references. (NAL Call No.: DNAL QH301.P535).

0640

Individual crown selection for resistance to freezing stress in winter oats (Germplasm, Cold hardiness). Marshall, H.G. Kolb, F.L. Madison, Wis., Crop Science Society of America. Crop science. May/June 1982. v. 22 (3). p. 506-510. ill. 8 ref. (NAL Call No.: 64.8 C883).

0641

Indole-3-acetic acid sensitization of phytochrome-controlled growth of coleoptile sections (Avena sativa). Shinkle, J.R. Briggs, W.R. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. June 1984. v. 81 (12). p. 3742-3746. Includes references. (NAL Call No.: 500 N21P).

0642

Induction of a specific N-methyltransferase enzyme by long-term heat stress during barley leaf growth. PLPHA. Leland, T.J. Hanson, A.D. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Oct 1985. v. 79 (2). p. 451-457. ill. Includes 23 references. (NAL Call No.: DNAL 450 P692).

0643

Induction of barley leaf urease. PLPHA. Chen, Y.G. Ching, T.M. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Mar 1988. v. 86 (3). p. 941-945. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0644

Induction of lactate dehydrogenase isozymes by oxygen deficit in barley root tissue. PLPHA. Hoffman, N.E. Bent, A.F.; Hanson, A.D. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Nov 1986. v. 82 (3). p. 658-663. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0645

The influence of aliphatic alcohols on leaf senescence (Oat seedlings). Satler, S.O. Thimann, K.V. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1980. v. 66 (3). p. 395-399. ill. 28 ref. (NAL Call No.: 450 P692).

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

0646

The influence of ammonium and chloride on potassium and nitrate absorption by barley roots depends on time of exposure and cultivar. PLPHA. Bloom, A.J. Finazzo, J. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1986. v. 81 (1). p. 67-69. Includes 22 references. (NAL Call No.: DNAL 450 P692).

0647

The influence of copper on the metabolism of malic acid in intact barley roots (*Hordeum distichon*). Veltrup, W. JPNUD. New York : Marcel Dekker. Journal of plant nutrition. 1981. v. 4 (2). p. 217-230. 20 ref. (NAL Call No.: QK867.J67).

0648

The influence of gibberellic acid and temperature on the growth rate of *Avena sativa* stem segments (Oats). Jusaitis, M. Paleg, L.G.; Aspinall, D. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Aug 1982. v. 70 (2). p. 532-539. ill. 28 ref. (NAL Call No.: 450 P692).

0649

Influence of light and CO₂ (carbon dioxide) on nitrate assimilation by barley seedlings. Aslam, M. Huffaker, R.C.; Rains, D.W.; Rao, K.P. New York, Plenum Press. Genetic engineering of symbiotic nitrogen fixation and conservation of fixed nitrogen. 1980 (pub. 1981). 1980 (pub. 1981). p. 541-546. Includes 2 p. ref. (NAL Call No.: QR89.7.S95).

0650

Inheritance of subcrown internode length in a winter barley cross (Winter hardiness, optimum depth of crown formation). Dofing, S.M. Schmidt, J.W. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1984. v. 24 (4). p. 692-694. ill. Includes references. (NAL Call No.: 64.8 C883).

0651

Inhibited mitotic entry is the cause of growth inhibition by cinmethylin. WEESA6. El-deek, M.H. Hess, F.D. Champaign, Ill. : Weed Science Society of America. Weed science. Sept 1986. v. 34 (5). p. 684-688. Includes references. (NAL Call No.: DNAL 79.8 W41).

0652

Interaction between senescence and wounding in oat leaves. PLPHA. Giridhar, G. Thimann, K.V. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1985. v. 78 (1). p. 29-33. Includes 23 references. (NAL Call No.: DNAL 450 P692).

0653

Interaction of indoleacetic acid and gibberellic acid in the short-term growth kinetics of oat stem segments (*Avena sativa*). Adams, P.A. PLPHA. Ross, M.A. Rockville : American Society of Plant Physiologists. Plant physiology. Nov 1983. v. 73 (3). p. 566-568. ill. Includes references. (NAL Call No.: 450 P692).

0654

Interactions between ethylene, CO₂, and ABA and GA₃-induced amylase synthesis in barley aleurone tissue. PLPHA. Tittle, F.L. Spencer, M.S. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Apr 1986. v. 80 (4). p. 1034-1037. Includes 22 references. (NAL Call No.: DNAL 450 P692).

0655

Interrelationships among three major loci controlling heading date of spring barley when grown under short daylengths. CRPSAY. Gallagher, L.W. Belhadri, M.; Zahour, A. Madison, Wis. : Crop Science Society of America. Early maturing cultivars of barley (*Hordeum vulgare* L.) would enhance yield stability in lowland, low rainfall areas of Morocco. The objective of this work was to study the inheritance of heading date in crosses involving six spring barleys developed in the Sacramento Valley of California, USA and the Yaqui Valley of Sonora, Mexico. Data were gathered under short daylengths at 32 degrees C 15' N Lat with December sowing in Morocco. Frequency distributions of parental, F₁, F₂, and backcross generations indicated a three-locus model with recessive and duplicate dominant epistasis accounting for most of the phenotypic variability. Recessive gene interaction resulted in plants that were about 19 to 35 days earlier than later counterparts with the dominant allele present. Among the later genotypes, the presence of a dominant allele at either of the duplicate loci in the genotypes not homozygous for extreme earliness resulted in increased earliness of 5 to 14 days over the duplicate double recessive depending upon modifiers. A genotype for each parent was suggested; however, the three-locus model did not account for the 4- to 6-day differences between two pairs of parents. Apparently, modifying loci or different alleles were present in some parents. Crop science. Mar/Apr

1987. v. 27 (2). p. 155-160. Includes references. (NAL Call No.: DNAL 64.8 C883).

0656

Involvement of plasma membrane alterations in cold acclimation of winter rye seedlings (*Secale cereale* L. cv Puma) (Sterol and protein composition changes).

Uemura, M. Yoshida, S. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. July 1984. v. 75 (3). p. 818-826. ill. Includes 34 references. (NAL Call No.: 450 P692).

0657

Iron reduction and uptake by grapevine roots (*Vitis* spp., also *Avena sativa*, oats). Varanini, Z. Maggioni, A. New York ; Basel : Marcel Dekker, 1982. Iron nutrition and interactions in plants : Brigham Young University, August 12-14, 1981 / edited by S.D. Nelson ... (et al.). p. 521-529. 9 ref. (NAL Call No.: QK867.J67 v. 5, nos. 4-7).

0658

Isogenic heading date effects on yield component development in 'Titan' barley. CRPSAY. Smail, V.W. Eslick, R.F.; Hockett, E.A. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 1023-1029. Includes references. (NAL Call No.: DNAL 64.8 C883).

0659

Isolation and identification of some phytotoxic compounds from aqueous extracts of rye (*Secale Cereale* L.).

JAFCAU. Shilling, D.G. Jones, L.A.; Worsham, A.D.; Parker, C.E.; Wilson, R.F. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. July/Aug 1986. v. 34 (4). p. 633-638. Includes references. (NAL Call No.: DNAL 381 J8223).

0660

Kinetics of the uptake of ¹⁴C-labelled chlorinated benzenes from soil by plants. EESAD. Topp, E. Scheunert, I.; Korte, F. Duluth, Minn. : Academic Press. Ecotoxicology and environmental safety. Apr 1989. v. 17 (2). p. 157-166. Includes references. (NAL Call No.: DNAL QH545.A1E29).

0661

Lamellar-to-hexagonalII phase transitions in the plasma membrane of isolated protoplasts after freeze-induced dehydration. PNASA. Gordon-Kamm, W.J. Steponkus, P.L. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. Oct 1984. v. 81 (20). p. 6363-6377. ill. Includes 34 references. (NAL Call No.: DNAL 500 N21P).

0662

Latency of plasma membrane H⁺-ATPase in vesicles isolated by aqueous phase partitioning: increased substrate accessibility or enzyme activation. PLPHA. Sandstrom, R.P. deBoer, A.H.; Lomax, T.L.; Cleland, R.E. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Nov 1987. v. 85 (3). p. 693-698. Includes references. (NAL Call No.: DNAL 450 P692).

0663

Leaf water potential, relative water content, and diffusive resistance as screening techniques for drought resistance in barley. AGJOAT. Matin, M.A. Brown, J.H.; Ferguson, H. Madison, Wis. : American Society of Agronomy. Rapid drought resistance screening techniques could accelerate selection of improved cultivars for semiarid areas. This study was conducted to determine if total leaf water potential, leaf relative water content, and leaf diffusive resistance could be used to differentiate between barley (*Hordeum vulgare* L.) cultivars differing in apparent drought resistance. Resistant and susceptible cultivars were selected by regressing individual cultivar yields against site mean yield for numerous yield trials. Individual plants were grown in cone-tainers (plastic tubes 25 mm in diameter and 123 mm long). The lower 30 mm of the cone-tainers were imbedded in sand so that roots emerging from the bottom of the cone-tainers were in the capillary fringe above a water table in the sand. Fifty plants were grown at a time in a sand bed, five replications each of five resistant and five susceptible barley cultivars of either two- or six-row head types. At the three-leaf growth stage total leaf water potential was measured on the third leaf of each plant under low stress conditions (predawn). Water was then drained from the sand bed to stress the plants. Leaf diffusive resistance (LDR) was measured at 1000 and 1300 h each day as stress developed. When the PM diffusive resistance reached about 1000 s m⁻¹, relative water content (RWC) of the second leaf was determined. All measurements were repeated in three different trials for each head type. Relative water content and total leaf water potential differentiated between drought-resistant and drought-susceptible groups of two- and six-rowed barley at the 0.01 probability level in every trial. Leaf diffusive resistance

differentiated between resistant and susceptible groups at the 0.01 or 0.05 level under low and high stress conditions. Neither RWC nor LDR allowed separation of cultivars within resistant or susceptible groups, but total leaf water potential did show differences between cultivars within groups. These techniques provide an ea. Agronomy journal. Jan/Feb 1989. v. 81 (1). p. 100-105. Includes references. (NAL Call No.: DNAL 4 AM34P).

0664

Lipid and fatty acid composition of three winter barley cultivars differing in winterhardiness / by Mohammed Abdul Ali. - Ali, Mohammed Abdul, 1941-. 1971. Thesis (Ph.D.)--University of Wyoming, 1971. Photocopy of typescript. Ann Arbor: University Microfilms, 1972. v. 32 leaves ; 21 cm. Bibliography: leaves 28-31. (NAL Call No.: DISS 72-18,322).

0665

Lipid composition of plasma membranes and endomembranes prepared from roots of barley (*Hordeum vulgare* L.). Effects of salt. PLPHA. Brown, D.J. DuPont, F.M. Rockville, Md. : American Society of Plant Physiologists. Membrane fractions enriched in endoplasmic reticulum (ER), tonoplast and Golgi membranes (TG) and plasma membranes (PM) were prepared from barley (*Hordeum vulgare* L. cv CM 72) roots and the lipid compositions of the three fractions were analyzed and compared. Plants were grown in an aerated nutrient solution with or without 100 millimolar NaCl. Each membrane fraction had a characteristic lipid composition. The mole per cent of the individual phospholipids, glycolipids, and sterols in each fraction was not altered when roots were grown in 100 millimolar NaCl. The ER had the highest percentages of phosphatidylinositol and phosphatidylcholine of the three fractions (7 and 45 mole per cent, respectively, of the total lipid). The TG contained the highest percentage of glycosylceramide (13 mole per cent). The PM had the highest percentage of phosphatidylserine (3 mole per cent) and nearly equal percentages of phosphatidylethanolamine (15 mole per cent and phosphatidylcholine (18 mole per cent). The most abundant sterols in membranes prepared from barley roots were stigmasterol (10 mole per cent), sitosterol (50 mole per cent), and 24 zeta-methylcholesterol (40 mole per cent of the total sterol). Salt-treated plants contained a slightly higher percentage of stigmasterol than controls. The percentage of stigmasterol increased with age and a simple cause and effect relationship between salt treatment and sterol composition was not observed. Plant physiology. July 1989. v. 90 (3). p. 955-961. Includes references. (NAL Call No.: DNAL 450 P692).

0666

Localization and pattern of graviresponse across the pulvinus of barley *Hordeum vulgare*. PLPHA. Brock, T.G. Lu, C.R.; Ghosheh, N.S.; Kaufman, P.B. Rockville, Md. : American Society of Plant Physiologists. Pulvini of excised stem segments from barley (*Hordeum vulgare* cv Larker') were pretreated with 1 millimolar coumarin before gravistimulation to reduce longitudinal cell expansion and exaggerate radial cell enlargement. The cellular localization and pattern of graviresponse across individual pulvini were then evaluated by cutting the organ in cross-section, photographing the cross-section, and then measuring pulvinus thickness and the radial width of cortical and epidermal cells in enlargements of photomicrographs. With respect to orientation during gravistimulation, we designated the uppermost point of the cross-section 0 degrees and the lowermost point 180 degrees. A gravity-induced increase in pulvinus thickness was observable within 40 degrees of the vertical in coumarin-treated pulvini. In the upper halves of coumarin-treated gravistimulated pulvini, cells in the inner cortex and inner epidermis had increased radial widths, relative to untreated gravistimulated pulvini. In lower halves of coumarin-treated pulvini, cells in the central and outer cortex and in the outer epidermis showed the greatest increase in radial width. Cells comprising the vascular bundles also increased in radial width, with this pattern following that of the central cortex. These results indicate (a) that all cell types are capable of showing a graviresponse, (b) that the graviresponse occurs in both the top and the bottom the responding organ, and (c) that the magnitude of the response increases approximately linearly from the uppermost point to the lowermost. These results are also consistent with models of gravitropism that link the pattern and magnitude of the graviresponse to gravi-perception via statolith sedimentation. Plant physiology. Oct 1989. v. 91 (2). p. 744-748. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0667

Longitudinal and transverse profiles of K⁺ and Cl⁻ concentration in 'low-' and 'high-salt' barley roots. NEPHA. Huang, C.X. Van Steveninck, R.F.M. New York, N.Y. : Cambridge University Press. The New phytologist. Aug 1989. v. 112 (4). p. 415-480. ill. Includes references. (NAL Call No.: DNAL 450 N42).

0668

Low ribonuclease I activity prior to cold acclimation in freeze selected winter barley. CRPSAY. Kenefick, D.G. Blake, T.K. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1986. v. 26 (6). p. 1099-1103. Includes 14 references. (NAL Call No.: DNAL 64.8 C883).

0669

Maintenance of low Cl⁻ concentrations in mesophyll cells of leaf blades of barley seedlings exposed to salt stress.
 PLPHA. Huang, C.X. Van Steveninck, R.F.M. Rockville, Md. : American Society of Plant Physiologists. The concentrations of vacuolar Na⁺ and Cl⁻ in the epidermal and mesophyll cells of the leaf blade and sheath of *Hordeum vulgare* seedlings (cv California Mariout and Clipper) were measured by means of quantitative electron probe X-ray microanalysis. A preferential accumulation of Cl⁻ in vacuoles of epidermal cells in both blade and sheath and a low level in mesophyll cells of the blade were evident in plants grown in full strength Johnson solution. The concentration of Cl⁻ in the mesophyll cells of the blade remained at a low level after exposure to 50 or 100 millimolar NaCl for 1 day or to 50 millimolar for 4 days, while at the same time the concentration of Cl⁻ in the epidermis and mesophyll of the sheath showed a dramatic increase. Clipper generally contained more Cl⁻ in the mesophyll cells of the blade than California Mariout. A greater accumulation of Na⁺ in the mesophyll of the sheath relative to that of the blade was only apparent after treatment with 100 millimolar NaCl for 1 day or 50 millimolar for 4 days. These results confirm the suggestion that sheath tissue is capable of accumulating excess Cl⁻ (and to a lesser extent Na⁺) and suggest that the site of regulation of Cl⁻ concentration in the barley leaf is located in the mesophyll cells of the blade. Plant physiology. Aug 1989. v. 90 (4). p. 1440-1443. Includes references. (NAL Call No.: DNAL 450 P692).

0670

Mechanical properties of the plasma membrane of isolated plant protoplasts. Mechanism of hyperosmotic and extracellular freezing injury (*Secale cereale*, rye seedlings, changes in area and membrane contraction in lysis).
 Wolfe, J. PLPHA. Steponkus, P.L. Rockville : American Society of Plant Physiologists. Plant physiology. Feb 1983. v. 71 (2). p. 276-285. ill. 28 ref. (NAL Call No.: 450 P692).

0671

Mechanical study of the deformation and rupture of the plasma membranes of protoplasts during osmotic expansions.
 JMBB80. Wolfe, J. Dowgert, M.F.; Steponkus, P.L. New York, N.Y. : Springer. The Journal of membrane biology. 1986. v. 93 (1). p. 63-74. ill. Includes references. (NAL Call No.: DNAL QH573.J6).

0672

The mechanics of injury to isolated protoplasts following osmotic contraction and expansion.
 PLPHA. Dowgert, M.F. Wolfe, J.; Steponkus, P.L. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Apr 1987. v. 83 (4). p. 1001-1007. Includes references. (NAL Call No.: DNAL 450 P692).

0673

Mesocotyl growth of etiolated seedlings of *Avena sativa* and *Zea mays* in relation to light and diffusible auxin.
 JPGRDI. Kessler, R.W. Gerhold, L.S.; Muir, R.M. New York, N.Y. : Springer. Journal of plant growth regulation. 1985. v. 4 (1). p. 11-18. Includes references. (NAL Call No.: DNAL QK745.J6).

0674

Mesophyll resistances to SO₂ fluxes into leaves.
 PLPHA. Pfanz, H. Martinoia, E.; Lange, O.L.; Heber, U. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Dec 1987. v. 85 (4). p. 922-927. Includes references. (NAL Call No.: DNAL 450 P692).

0675

A method for rating freezing inhibitor activity of barley genotypes using seed and crown tissue mucilages (Winter hardiness).
 Olien, C.R. AR-NC. Kuhna, A.S. Madison, Wis., Crop Science Society of America. Crop science. July/Aug 1980. v. 20 (4). p. 537-539. ill. 9 ref. (NAL Call No.: 64.8 C883).

0676

A model for the regulation of K⁺ influx, and tissue potassium concentrations by negative feedback effects upon plasmalemma influx.
 PLPHA. Siddiqi, M.Y. Glass, A.D.M. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1986. v. 81 (1). p. 1-7. Includes 32 references. (NAL Call No.: DNAL 450 P692).

0677

Modification of flag leaf senescence and yield characters in barley (*Hordeum vulgare* L.) by gibberellic acid and kinetin.
 JPGRDI. Mishra, S.D. Gaur, B.K. New York, N.Y. : Springer. Journal of plant growth regulation. 1985. v. 4 (2). p. 63-70. Includes references. (NAL Call No.: DNAL QK745.J6).

0678

Modifications of plant growth and ash content as effected by acids added to soils.
AGJ0AT. Carr, R.H. Haverkamp, H.G. Madison, Wis. : American Society of Agronomy. A comparison has been made of the effect upon the growth of soybean and buckwheat plants of adding an organic (acetic) and an inorganic (sulfuric) acid to a silt and a clay-loam soil. It has been found that both acids seriously interfere with plant growth, but that acetic was more deleterious than sulfuric acid. This is believed to be due to the effect of acetic acid in supplying more toxic aluminum and iron, etc., to the plant and in decreasing calcium and magnesium to a greater extent than was done by the sulfuric acid. In addition two other series of acids (phosphoric and silicic) were used, because they make compounds with aluminum and iron which seem to be harmless to the growing plant, regardless of the fact that the root is surrounded by a soil which is highly acid. The good growth of both kinds of plants obtained, even when rather large amounts of these latter acids were added, indicates that relatively insoluble compounds of aluminum and iron, etc., were formed and that the acid by itself did no harm. *Agronomy journal*. Apr 1924. v. 16 (4). p. 278-283. (NAL Call No.: DNAL 4 AM34P).

0679

Modifications of sulfhydryl groups on phytochrome and their influence on physicochemical differences between the red- and far-red-absorbing forms.
PLPHA. Smith, W.O. Jr. Cyr, K.L. Rockville, Md. : American Society of Plant Physiologists. *Plant physiology*. May 1988. v. 87 (1). p. 195-200. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0680

Molecular and cellular biology associated with endosperm mobilization in germinating cereal grains.
Fincher, G.B. Paio Alto, Calif. : Annual Reviews, Inc. Annual review of plant physiology and plant molecular biology. Literature review. 1989. v. 40. p. 305-346. ill. Includes references. (NAL Call No.: DNAL QK1.A57).

0681

The molecular biology of barley storage protein synthesis.
Cameron-Mills, V. Brandt, A.; Ingversen, J. New York, Academic Press, 1980. *Cereals for food and beverages : recent progress in cereal chemistry and technology*, edited by George E. Inglett, Lars Munck. International Conference on Cereals for Food and Beverages (1979 : Copenhagen). p. 339-364. ill. 33 ref. (NAL Call No.: TX557.I57 1979).

0682

Molecular cloning and sequence of cDNA encoding the plasma membrane proton pump (H⁺-ATPase) of *Arabidopsis thaliana*.
PNASA. Harper, J.F. Surowy, T.K.; Sussman, M.R. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. Feb 1989. v. 86 (4). p. 1234-1238. ill. Includes references. (NAL Call No.: DNAL 500 N21P).

0683

Morphological and histological effects of three grass selective herbicides on developing wild oat (*Avena fatua*) stems.
WEESA6. Jain, R. Born, W.H. vanden. Champaign, Ill. : Weed Science Society of America. Three grass selective herbicides, sethoxydim, fluazifop, and haloxyfop, applied to wild oat plants at the five-leaf stage inhibited growth and induced chlorosis in leaves. Young and actively growing tissues were affected first. Stem elongation in wild oat was inhibited within 2 days of treatment with sethoxydim and within 5 days of treatment with fluazifop or haloxyfop. At these same observation times, internodes that were elongating rapidly at the time of treatment were constricted at the base. These symptoms were followed by necrosis of the internode tissue. Histological examination of the affected internodes indicated that the herbicides inhibited cell division in very young internodes and inhibited both cell division and cell elongation in slightly older internodes. Initial injury occurred in the epidermal, cortical, and procambium cells of the peripheral regions of the stems located at the base of the affected internodes. Necrosis then progressed to the center of the stem tissue and all cells in the internodes were killed within 14 days of treatment. All three herbicides caused similar morphological and histological effects on developing wild oat stems. *Weed science*. July 1989. v. 37 (4). p. 575-584. ill. Includes references. (NAL Call No.: DNAL 79.8 W41).

0684

The morphological and physiological response of slender oat (*Avena barbata*) to the herbicides barban and difenzoquat.
WEESA6. Price, S.C. Hill, J.E.; Allard, R.W. Champaign, Ill. : Weed Science Society of America. *Weed science*. Jan 1988. v. 36 (1). p. 60-69. Includes references. (NAL Call No.: DNAL 79.8 W41).

0685

Morphological and physiological variation in wild oat (*Avena fatua*, dormancy, herbicide resistance, varieties, weeds).
Miller, S.D. Nalewaja, J.D.; Mulder, C.E.G. Madison, Wis., American Society of Agronomy. *Agronomy journal*. Sept/Oct 1982. v. 74 (5). p.

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

771-775. ill. 11 ref. (NAL Call No.: 4 AM34P).

0686

Multiple actions of abscisic acid in senescence of oat leaves.

JPGRDI. Zhi-Yi, T. Veierskov, B.; Park, J.; Thimann, K.V. New York, N.Y. : Springer. Journal of plant growth regulation. 1988. v. 7 (4). p. 213-226. Includes references. (NAL Call No.: DNAL QK745.J6).

0687

NaCl induces a Na⁺/H⁺ antiport in tonoplast vesicles from barley roots.

PLPHA. Garbarino, J. DuPont, F.M. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Jan 1988. v. 86 (1). p. 231-236. Includes references. (NAL Call No.: DNAL 450 P692).

0688

Natural selection in a doubled-haploid mixture and a composite cross of barley.

CRPSAY. Patel, J.D. Reinbergs, E.; Mather, D.E.; Choo, T.M.; Sterling, J.D.E. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1987. v. 27 (3). p. 474-479. Includes references. (NAL Call No.: DNAL 64.8 C883).

0689

Nature of photooxidative events in leaves treated with chlorosis-inducing herbicides (Secale cereale, winter rye seedlings).

Feierabend, J. Winkelhusener, T. Rockville : American Society of Plant Physiologists. Plant physiology. Nov 1982. v. 70 (5). p. 1277-1282. 30 ref. (NAL Call No.: 450 P692).

0690

New mutants in the genetic male sterile barley collection.

Hockett, E.A. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 70-73. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0691

Nitrate accumulation in plants.

Jensen, E.H. Reno, Nev. : College of Agriculture, University of Nevada-Reno, Nevada Cooperative Extension. Fact sheet - College of Agriculture, University of Nevada-Reno, Nevada Cooperative Extension. 1987? . (87-32). 3 p. (NAL Call No.: DNAL S544.3.N3C66).

0692

Nitrate content in oats /by B.J. Kolp ... et al..

Kolp, B. J. 1928-. Laramie, Wyo. : University of Wyoming, Agricultural Experiment Station, 1963. 12 p. ; 23 cm. Bibliography: p. 12. (NAL Call No.: DNAL 100 W99 (1) no.409).

0693

Nitrate uptake into barley (Hordeum vulgare) plants. A new approach using ³⁵ClO₃⁻ (chlorate) as an analog for NO₃⁻ (nitrate).

Deane-Drummond, C.E. Glass, A.D.M. Rockville, American Society of Plant Physiologists. Plant physiology. July 1982. v. 70 (1). p. 50-54. 27 ref. (NAL Call No.: 450 P692).

0694

Nitrate utilization by nitrate reductase-deficient barley mutants (Nutritional deficiencies).

Warner, R.L. Kleinhofs, A. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Apr 1981. v. 67 (4). p. 740-743. 111. 22 ref. (NAL Call No.: 450 P692).

0695

Nitrogen and yield as related to water use of spring barley (Hordeum vulgare, seasonal evapotranspiration, New Mexico).

Kallsen, C.E. AGJOAT. Sammis, T.W.; Gregory, E.J. Madison : American Society of Agronomy. Agronomy journal. Jan/Feb 1984. v. 76 (1). p. 59-64. Includes references. (NAL Call No.: 4 AM34P).

0696

Nitrogen harvest index in oats: its repeatability and association with adaptation. CRPSAY. Rattunde, H.F. Frey, K.J. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1986. v. 26 (3). p. 606-610. Includes references. (NAL Call No.: DNAL 64.8 C883).

0697

Oat leaf volatiles: possible insect attractants.

Buttery, R.G. Ling, L.C.; Wellso, S.G. Washington, D.C., American Chemical Society. Journal of agricultural and food chemistry. July/Aug 1982. v. 30 (4). p. 791-792. 10 ref. (NAL Call No.: 381 J8223).

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

0698

Oat lipids and lipid-related enzymes.

Youngs, V.L. St. Paul, Minn. : American Association of Cereal Chemists, c1986. Oats : chemistry and technology / edited by Francis H. Webster. Literature review. p. 205-226. ill. Includes references. (NAL Call No.: DNAL TP435.0502).

0699

Oat stem vascular size in relation to kernel number and weight. I. Controlled environment (*Avena sativa*).

Housley, T.L. Peterson, D.M. Madison, Wis., Crop Science Society of America. Crop science. Mar/Apr 1982. v. 22 (2). p. 259-263. ill. Includes 15 ref. (NAL Call No.: 64.8 C883).

0700

Oat stem vascular size in relation to kernel number and weight. II. Field environment (*Avena sativa*).

Peterson, D.M. Housley, T.L.; Luk, T.M. Madison, Wis., Crop Science Society of America. Crop science. Mar/Apr 1982. v. 22 (2). p. 274-278. Includes 22 ref. (NAL Call No.: 64.8 C883).

0701

Oats (Production, breeding, composition).

Youngs, V.L. Peterson, D.M.; Brown, C.M. St. Paul, Minn. : American Association of Cereal Chemists. Advances in cereal science and technology. 1982. Literature review. v. 5. p. 49-105. ill. Includes references. (NAL Call No.: TS2120.A3).

0702

Oats tolerant of *Pseudomonas syringae* pv. *tabaci* contain tabtoxinine-beta-lactam-insensitive leaf glutamine synthetases.

PLPHA. Knight, T.J. Bush, D.R.; Langston-Unkefer, P.J. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Oct 1988. v. 88 (2). p. 333-339. Includes references. (NAL Call No.: DNAL 450 P692).

0703

On the uptake, metabolism and retention of (3H) (hydrogen isotope) gibberellin A1 by barley aleurone layers at low temperatures.

Keith, B. Boal, R.; Srivastava, L.M. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Nov 1980. v. 66 (5). p. 956-961. ill. 20 ref. (NAL Call No.: 450 P692).

0704

Origin, taxonomy, and related species.

AGRYA. Bothmer, R. von. Jacobsen, N. Madison, Wis., Soil Science Society of America. Agronomy. A series of monographs - American Society of Agronomy. Literature review. 1985. (26). p. 19-56. maps. Includes references. (NAL Call No.: DNAL 4 AM392).

0705

Osmotic adjustment and growth of barley genotypes under drought stress.

CRPSAY. Blum, A. Madison, Wis. : Crop Science Society of America. Osmotic adjustment is receiving increasing attention as a probable component of drought resistance in crop plants, including barley (*Hordeum vulgare* L.). This work was done to study a possible association between osmotic adjustment and growth under drought stress. Twenty-six diverse barley genotypes were grown in nutrient solutions in the growth chamber. Drought stress was imposed by adding polyethylene glycol 6000 to the nutrient solution producing a -0.54 MPa solution water potential as compared with controls of pure nutrient solution with a -0.02 MPa of solution potential. Percent growth reduction by drought stress was calculated from growth rates measured under stress and in the controls. Osmotic adjustment was calculated as the difference in osmotic potential at full turgor between stress and controls. Leaf water potentials of genotypes grown under drought stress ranged from -1.37 to -1.51 MPa, but did not differ significantly. For the same leaf water potential, genotypes differed significantly in osmotic adjustment which ranged from -0.17 to 0.46 MPa. Growth reduction by drought stress differed significantly among genotypes with a range of 78.1%. Growth reduction by drought stress was negatively associated with osmotic adjustment across genotypes ($r = -0.80$). The association was nonlinear, indicating that growth reduction decreased only as osmotic adjustment increased to about 0.2 MPa or more. These results suggest that induced osmotic adjustment under drought stress may be an important component of drought resistance in barley growth, and it should be evaluated when all genotypes are subjected to the same leaf water status. Crop science. Jan/Feb 1989. v. 29 (1). p. 230-233. Includes references. (NAL Call No.: DNAL 64.8 C883).

0706

Osmotic stress-induced polyamine accumulation in cereal leaves. I. Physiological parameters of the response (Includes wild oats, *Avena fatua*, and peas, water stress).

Flores, H.E. Galston, A.W. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1984. v. 75 (1). p. 102-109. ill. Includes 33 references. (NAL Call No.: 450 P692).

0707

Osmotic stress-induced polyamine accumulation in cereal leaves. II. Relation to amino acid pools (Oats).

Flores, H.E. Galston, A.W. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1984. v. 75 (1). p. 110-113. Includes 29 references. (NAL Call No.: 450 P692).

0708

Oxygen and carbon dioxide fluxes from barley shoots depend on nitrate assimilation.

PLPHA. Bloom, A.J. Caldwell, R.M.; Finazzo, J.; Warner, R.L.; Weissbart, J. Rockville, Md. : American Society of Plant Physiologists. A custom oxygen analyzer in conjunction with an infrared carbon dioxide analyzer and humidity sensors permitted simultaneous measurements of oxygen, carbon dioxide, and water vapor fluxes from the shoots of intact barley plants (*Hordeum vulgare* L. cv Steptoe). The oxygen analyzer is based on a calceazirconium sensor and can resolve concentration differences to within 2 microliters per liter against the normal background of 210,000 microliters per liter. In wild-type plants receiving ammonium as their sole nitrogen source or in nitrate reductase-deficient mutants, photosynthetic and respiratory fluxes of oxygen equaled those of carbon dioxide. By contrast, wild-type plants exposed to nitrate had unequal oxygen and carbon dioxide fluxes: oxygen evolution at high light exceeded carbon dioxide consumption by 26% and carbon dioxide evolution in the dark exceeded oxygen consumption by 25%. These results indicate that a substantial portion of photosynthetic electron transport or respiration generates reductant for nitrate assimilation rather than for carbon fixation or mitochondrial electron transport. Plant physiology. Sept 1989. v. 91 (1). p. 352-356. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0709

Partial characterization of fusaric acid (a phytotoxin isolated from the fungus *Fusicoccum amygdali*) binding to receptor sites on oat root membranes.

Stout, R.G. Cleland, R.E. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1980. v. 66 (3). p. 353-359. ill. 27 ref. (NAL Call No.: 450 P692).

0710

Phase transitions in liposomes formed from the polar lipids of mitochondria from chilling-sensitive plants.

PLPHA. Raison, J.K. Orr, G.R. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. July 1986. v. 81 (3). p. 807-811. Includes 26 references. (NAL Call No.: DNAL 450 P692).

0711

Phosphorylation of *Avena* phytochrome in vitro as a probe of light-induced conformational changes.

JBCHA3. Wong, Y.S. Cheng, H.C.; Walsh, D.A.; Lagarias, J.C. Baltimore, Md. : American Society of Biological Chemists. The Journal of biological chemistry. Sept 15, 1986. v. 261 (26). p. 12089-12097. ill. Includes references. (NAL Call No.: DNAL 381 J824).

0712

Photoassimilate partitioning by tillers and individual tiller leaves in field-grown spring barley.

CRPSAY. Lauer, J.G. Simmons, S.R. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 279-282. Includes references. (NAL Call No.: DNAL 64.8 C883).

0713

Photosynthesis, leaf resistances, and ribulose-1,5-bisphosphate carboxylase degradation in senescing barley leaves.

Friedrich, J.W. Huffaker, R.C. Bethesda, Md., American Society of Plant Physiologists. Plant physiology. June 1980. v. 65 (6). p. 1103-1107. ill. 26 ref. (NAL Call No.: 450 P692).

0714

Physiological and ultrastructural studies of oat membranes treated with *Helminthosporium victoriae* toxin / by Vernon Edward Gracen, Jr.

Gracen, Vernon Edward, 1945-. 1970. Thesis (Ph.D.)--University of Florida, 1970. Photocopy. Ann Arbor, Mich. : University Microfilms, 1971. ix, 89 leaves ; 21 cm. Bibliography: leaves 84-88. (NAL Call No.: DISS 71-248).

0715

Physiological aspect of mugineic acid, a possible phytosiderophore of graminaceous plants (Barley, iron uptake).

Takagi, S. Nomoto, K.; Takemoto, T. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. 1984. Presented at the "Second International Symposium on Iron Nutrition and Interactions in Plants," August 2-5, 1983, Utah State University, Logan. v. 7 (1/5). p. 469-477. ill. Includes references. (NAL Call No.: QK867.J67).

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

0716

Physiological aspects of enzymes during grain development and germination.

Duffus, C.M. St. Paul, Minn. : American Association of Cereal Chemists, c1987. Enzymes and their role in cereal technology / edited by James E. Kruger, David Lineback, Clyde E. Stauffer. Literature review. p. 83-116. ill. Includes references. (NAL Call No.: DNAL TP434.E58).

0717

The physiological basis of nitrogen redistribution during grain filling in cereals.
Dalling, M.J. Rockville, Md. : American Society of Plant Physiologists, c1985. Exploitation of physiological and genetic variability to enhance crop productivity / edited by James E. Harper, Lawrence E. Schrader, and Robert W. Howell. Literature review. p. 55-71. ill. Includes 45 references. (NAL Call No.: DNAL SB189.4.E97).

0718

Physiological changes associated with three cycles of recurrent selection for grain yield improvement in oats.

CRPSAY. Payne, T.S. Stuthman, D.D.; McGraw, R.L.; Bregitzer, P.P. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 734-736. Includes references. (NAL Call No.: DNAL 64.8 C883).

0719

Physiological control of arginine decarboxylase activity in K-deficient oat shoots.

PLPHA. Young, N.D. Galston, A.W. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Oct 1984. v. 76 (2). p. 331-335. ill. Includes 13 references. (NAL Call No.: DNAL 450 P692).

0720

Physiological interaction, its mathematical expression (*Avena sativa* L., diuron, phorate).

Drury, R.E. Champaign, Ill., Weed Science Society of America. Weed science. Sept 1980. v. 28 (5). p. 573-579. ill. 22 ref. (NAL Call No.: 79.8 W41).

0721

Physiological mechanism of the auxin-induced increase in light sensitivity of phytochrome-mediated growth responses in *Avena coleoptile* sections.

PLPHA. Shinkle, J.R. Briggs, W.R. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Oct 1985. v. 79 (2). p.

349-356. Includes 27 references. (NAL Call No.: DNAL 450 P692).

0722

Physiological responses of barley leaves of foliar applied urea-ammonium nitrate.

CRPSAY. Turley, R.H. Ching, T.M. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 987-993. Includes references. (NAL Call No.: DNAL 64.8 C883).

0723

Physiology and development.

AGRYA. Wych, R.D. Simmons, S.R.; Warner, R.L.; Kirby, E.U.M. Madison, Wis., Soil Science Society of America. Agronomy. A series of monographs - American Society of Agronomy. Literature review. 1985. (26). p. 103-125. Includes references. (NAL Call No.: DNAL 4 AM392).

0724

Physiology of sprouting resistance.

King, R.W. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 27-60. ill. Includes references. (NAL Call No.: DNAL SB189.6.P73).

0725

Plant bioregulators in cereal crops: action and use (Growth regulators, chlormequat chloride, CCC, mepiquat chloride, ethephon).

Jung, J. Washington, D.C. : The Society. ACS Symposium series - American Chemical Society. 1984. 1984. (257). p. 29-43. ill. Includes references. (NAL Call No.: OD1.A45).

0726

Plant cell culture and somatic cell genetics of cereals and grasses.

Vasil, I.K. New York : Academic Press, 1982. Plant improvement and somatic cell genetics / edited by I.K. Vasil, W.R. Scowcroft, K.J. Frey. Literature review. p. 179-203. Includes references. (NAL Call No.: SB123.P563).

0727

Plant response to topsoil thickness on an eroded loess soil.

JSWCA3. Mielke, L.N. Schepers, J.S. Ankeny, Iowa : Soil Conservation Society of America. Journal of soil and water conservation. Jan/Feb 1986. v. 41 (1). p. 59-63. Includes 15 references. (NAL Call No.: DNAL 56.8 J822).

0728

Plant stress and polyamine metabolism induction of arginine decarboxylase in stressed oat leaves (putrescine, mineral- deficiency, ionic-stress) /Nevin Dale Young.

Young, Nevin Dale. 1984. Thesis (Ph.D)--Yale University, 1984. vi, 179 p. : ill. Bibliography: p. 166-179. (NAL Call No.: DNAL DISS 85-09,796).

0729

Plasma membrane lipid alterations associated with cold acclimation of winter rye seedlings (*Secale cereale* L. cv Puma).

PLPHA. Lynch, D.V. Steponkus, P.L. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Apr 1987. v. 83 (4). p. 761-767. Includes references. (NAL Call No.: DNAL 450 P692).

0730

Plasma membrane lipid alterations following cold acclimation: possible relevance to freeze tolerance.

Lynch, D.V. Steponkus, P.L. New York : Plenum Press, c1987. The metabolism structure, and function of plant lipids / edited by Paul K. Stumpf, J. Brian Mudd, and W. David Nes. Paper presented at the "Seventh International Symposium on Plant Lipids," held July 27-August 1, 1986, University of California, Davis, California. p. 213-215. Includes references. (NAL Call No.: DNAL QK898.L56155 1986).

0731

Plastid transcription activity and DNA copy number increase early in barley chloroplast development.

PLPHA. Baumgartner, B.J. Rapp, J.C.; Mullet, J.E. Rockville, Md. : American Society of Plant Physiologists. Plastid transcription activity and DNA copy number were quantified during chloroplast development in the first foliage leaf in dark-grown and illuminated barley (*Hordeum vulgare* L.) seedlings. Primary foliage leaves of seedlings given continuous illumination from 2 days post-imbibition reached a final mean length of 15 centimeters at 6.5 days, whereas primary leaves of dark-grown seedlings required 7 days to reach a similar length. Dividing cells were observed in the basal 0.5 to 1 centimeter of primary leaves until 5.5 days post-imbibition. Plastids isolated from cells located in the basal meristem of 4-day-old seedlings were small (approximately 2 micrometers in diameter), exhibited low transcription activity and contained approximately 130 copies of plastid DNA per organelle. Cell size increased from 18 to 60 micrometers in a 1 to 3 centimeter region located adjacent to the leaf basal meristem. In this region, transcriptional activity per plastid increased 10-fold and DNA copy number increased from 130 to 210. Plastid

transcriptional activity declined rapidly in illuminated plants with increasing leaf cell age and plastid DNA copy number also declined but with a slower time course. In dark-grown seedlings plastid transcriptional activity declined more slowly than in illuminated plants while DNA copy number remained constant with increasing cell age. These data show that plastid transcriptional activity and DNA copy number increase early in chloroplast development and that transcriptional activity per DNA template varies up to 5-fold during barley leaf biogenesis. Plant physiology. Mar 1989. v. 89 (3). p. 1011-1018. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0732

Polysomes from winter rye seedlings grown at low temperature. I. Size class distribution, composition, and stability.

PLPHA. Laroche, A. Hopkins, W.G. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Nov 1987. v. 85 (3). p. 648-654. Includes references. (NAL Call No.: DNAL 450 P692).

0733

The potato scab /by Roland Thaxter. The proteids or albuminoids of the oat-kernel /by Thomas B. Osborne.

Thaxter, Roland, 1858-1932. Osborne, Thomas B. 1859-1929. New Haven : Connecticut Agricultural Experiment Station, 1890. Cover title. 8 p. ; 24 cm. (NAL Call No.: DNAL 100 C76St (1) no.105).

0734

Preharvest sprouting damage and sprouting tolerance: assay methods and instrumentation. Mares, D.J. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Denera. Literature review. p. 129-170. ill. Includes references. (NAL Call No.: DNAL SB189.6.P73).

0735

Proanthocyanidin-free barley for brewing: progress in breeding for high yield and research tool in polyphenol chemistry. TQMB. Wettstein, D. von. Nilan, R.A.; Ahrenst-Larsen, B.; Erdal, K.; Ingversen, J.; Jende-Strid, B.; Kristiansen, K.N.; Larsen, J.; Outtrup, H.; Ullrich, S.E. Madison, Wis. : The Association. Technical quarterly - Master Brewers Association of the Americas. 1985. v. 22 (2). p. 41-52. Includes references. (NAL Call No.: DNAL 390.9 M39T).

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

0736

Propagation of sea oats (*Uniola latifolia*).
Wilsey, M.H. Boulder, Colo. : International Plant Propagators' Society. The Plant propagator. June 1985. v. 31 (2). p. 15. (NAL Call No.: DNAL 81 P692).

0737

Protoporphyrinogen oxidation in chloroplasts and plant mitochondria, a step in heme and chlorophyll synthesis (Barley seedlings, *Hordeum vulgare*).

Jacobs, J.M.ABBIA. Jacobs, N.U.; De Maggio, A.E. New York : Academic Press. Archives of biochemistry and biophysics. Oct 1, 1982. v. 218 (1). p. 233-239. 26 ref. (NAL Call No.: 381 AR2).

0738

Pulse domestication and cereal domestication: how different are they?.

ECBDA. Zohary, D. Bronx, N.Y. : New York Botanical Garden. Economic botany. Jan/Mar 1989. v. 43 (1). p. 31-34. Includes references. (NAL Call No.: DNAL 450 EC7).

0739

Purification and characterization of gibberellic acid-induced cysteine endoproteases in barley aleurone layers.

PLPHA. Koehler, S. Ho, T.H.D. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1988. v. 87 (1). p. 95-103. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0740

Putrescine and acid stress. Induction of arginine decarboxylase activity and putrescine accumulation by low pH (*Avena sativa*, oats, *Pisum sativum*, peas).

Young, N.D.PLPHA. Galston, A.W. Rockville : American Society of Plant Physiologists. Plant physiology. Apr 1983. v. 71 (4). p. 767-771. Includes references. (NAL Call No.: 450 P692).

0741

P700 chlorophyll a-protein--purification, characterization, and antibody preparation (*Hordeum vulgare*, barley, *Glycine max*, soybeans, *Nicotiana tabacum*, tobacco, *Petunia X hybrida*, *Lycopersicon esculentum*, tomatoes, *Chlamydomonas reinhardtii*, Algae).

Vierling, E.PLPHA. Alberte, R.S. Rockville : American Society of Plant Physiologists. Plant physiology. July 1983. v. 72 (3). p. 625-633. ill. Includes references. (NAL Call No.: 450 P692).

0742

Quantitative measurement and evaluation of seasonal plant water deficits in barley / by Ray Eugene Jensen. -.

Jensen, Ray Eugene. 1971. Thesis (Ph.D.)--North Dakota State University, 1971. Photocopy of typescript. Ann Arbor: University Microfilms, 1972. xviii, 178 leaves ; 21 cm. Bibliography: leaves 151-155. (NAL Call No.: DISS 72-6,540).

0743

Quantitative observations on the pattern of synergid degeneration in barley.

AJBDA. Mogensen, H.L. Baltimore, Md. : Botanical Society of America. American journal of botany. Nov/Dec 1984. v. 71 (10). p. 1448-1451. ill. Includes references. (NAL Call No.: DNAL 450 AM36).

0744

Radiotracer evidence implicating phosphoryl and phosphatidyl bases as intermediates in betaine synthesis by water-stressed barley leaves.

Hitz, W.D. Rhodes, D.; Hanson, A.D. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Oct 1981. v. 68 (4). p. 814-822. 20 ref. (NAL Call No.: 450 P692).

0745

A rapid and simple procedure for purification of indole-3-acetic acid prior to GC-SIM-MS analysis.

PLPHA. Chen, K.H. Miller, A.N.; Patterson, G.W.; Cohen, J.D. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Mar 1988. v. 86 (3). p. 822-825. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0746

Rapid induction of Na⁺/H⁺ exchange activity in barley root tonoplast.

PLPHA. Garbarino, J. DuPont, F.M. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Jan 1989. v. 89 (1). p. 1-4. Includes references. (NAL Call No.: DNAL 450 P692).

0747

A rapid method for the determination of barley seed viability.

PIACA. Marks, G.C. Bloom, W.W.; Boyle, J.G. Indianapolis, Ind. : The Academy. Proceedings of the Indiana Academy of Science. 1985. v. 94. p. 117-119. Includes references. (NAL Call No.: DNAL 500 IN2).

0748

Rapid reduction by IAA of malondialdehyde levels in avena coleoptiles, a possible effect on lipid peroxidation.

BBRCA. Dhindsa, R.S. Amaral, A.C.; Cleland, R.E. New York, N.Y. : Academic Press. Biochemical and biophysical research communications. Nov 30, 1984. v. 125 (1). p. 76-81. Includes 20 references. (NAL Call No.: DNAL 442.8 B5236).

0749

The rate of absorption of nitrate of soda by oats and cotton when applied at different stages of plant growth.

AGJOAT. Appleton, W.H. Helms, H.B. Madison, Wis. : American Society of Agronomy. Experiments were conducted in the greenhouse to study the rate of absorption of nitrate nitrogen by oats and cotton when applied as nitrate of soda at different stages of plant growth. The general plan of the experiment was to add nitrate to a series of cultures at a certain stage of growth. Then, at intervals of a few days, part of the cultures were harvested and the amount of nitrate absorbed was determined by plant analysis and also by determining the nitrate content of the soil. Cultures that did not receive nitrate were used as checks to indicate how much nitrogen the plant secured from the soil. With oats these studies were made at four stages of plant growth. With cotton the study was made at three stages of plant growth. The results may be summarized as follows: When sodium nitrate at the rate of 400 pounds per acre was applied to oats 14 days after planting, absorption of the nitrate was very slow for three weeks. After the third week absorption increased and all nitrate was absorbed by the close of the seventh week. When the nitrate was applied to oats at later stages of growth, the rate of absorption was more rapid. Nitrate applied 42, 70, and 92 days after planting was completely absorbed in 20, 14, and 10 days, respectively. With both oats and cotton there was a close correlation between the rate of growth and the rate of nitrate absorption. Sodium nitrate at the rate of 600 pounds per acre was applied to cotton 14, 40, and 61 days after planting. Absorption of the nitrate was complete in 36, 14, and 11 days, respectively. The results of both experiments indicate that the loss of soluble nitrogenous fertilizer by leaching may be reduced by delaying the application until the crop will absorb it rapidly. Agronomy journal. Oct 1925. v. 17 (10). p. 596-605. (NAL Call No.: DNAL 4 AM34P).

0750

Reduced accumulation of ABA during water stress in a molybdenum cofactor mutant of barley.

PLPHA. Walker-Simmons, M. Kudrna, D.A.; Warner, R.L. Rockville, Md. : American Society of Plant Physiologists. A barley (*Hordeum vulgare* L.) mutant (Az34) has been identified with low basal levels of abscisic acid (ABA) and with

reduced capacity for producing ABA in response to water stress. The mutation is in a gene controlling the molybdenum cofactor resulting in a pleiotropic deficiency in at least three molybdoenzymes, nitrate reductase, xanthine dehydrogenase, and aldehyde oxidase. The mutant was found to lack aldehyde oxidase activity with several substrates including: (a) ABA aldehyde, a putative precursor of ABA; (b) an acetylenic analog of ABA aldehyde; and (c) heptaldehyde. Elevating the growth temperature from 18 to 26 degrees C caused mutant leaves to wilt and brown. Desiccation of mutant leaves was prevented by applying ABA. These results indicate that ABA biosynthesis at some developmental stages is dependent upon a molybdoenzyme which may be an aldehyde oxidase. Plant physiology. June 1989 v. 90 (2). p. 728-733. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0751

Relation between kernel row number and crude protein content of the grain in barley (Effects of two-rowed alleles on agronomic characters, physiological cause).

Yasuda, S. Moriya, I. Fort Collins, Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1982. v. 12. p. 45-46. (NAL Call No.: QK495.G74B34).

0752

Relationship between fructose 2,6-bisphosphate and carbohydrate metabolism in darkened barley primary leaves.

PLPHA. Baysdorfer, C. Sicher, R.C.; Kremer, D.F. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. July 1987. v. 84 (3). p. 766-769. Includes references. (NAL Call No.: DNAL 450 P692).

0753

Relationship between stress-induced ABA and proline accumulations and ABA-induced proline accumulation in excised barley leaves.

PLPHA. Stewart, C.R. Voetberg, G. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Sept 1985. v. 79 (1). p. 24-27. Includes 17 references. (NAL Call No.: DNAL 450 P692).

0754

Relationship between subcrown internode length and winter survival in winter barley.

CRPSAY. Dofing, S.M. Schmidt, J.W. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1985. v. 25 (4). p. 690-692. Includes references. (NAL Call No.: DNAL 64.8 C883).

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

0755

Relationship to lipophilicity to influx and efflux of triazine herbicides in oat roots.
PCBPB. Balke, N.E. Price, T.P. Duluth, Minn. Academic Press. Pesticide biochemistry and physiology. Mar 1988. v. 30 (3). p. 228-237. Includes references. (NAL Call No.: DNAL SB951.P49).

0756

The response of barley aleurone layers to gibberellic acid includes the transcription of new sequences.
Bernal-Lugo, I. Beachy, R.N.; Varner, J.E. New York, Academic Press. Biochemical and biophysical research communications. Sept 30, 1981. v. 102 (2). p. 617-623. ill. 15 ref. (NAL Call No.: 442.8 B5236).

0757

Response of soft chess (*Bromus mollis*) and slender oat (*Avena barbata*) to simulated drought cycles.
Ewing, A.L.URMGA. Menke, J.W. Denver : Society for Range Management. Journal of range management. July 1983. v. 36 (4). p. 415-418. ill. Includes references. (NAL Call No.: 60.18 J82).

0758

The response of wild oat (*Avena fatua*) and *Avena sterilis* accessions to photoperiod and temperature (Morphology, seed dormancy, growth rate).
Somody, C.N. Nalewaja, J.D.; Miller, S.D. Champaign : Weed Science Society of America. Weed science. Mar 1984. v. 32 (2). p. 206-213. ill. Includes references. (NAL Call No.: 79.8 W41).

0759

Responses of *Avena* (oat) coleoptiles to suboptimal fusaric acid: kinetics and comparisons with indoleacetic acid.
Rubinstein, B. Cleland, R.E. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1981. v. 68 (3). p. 543-547. 17 ref. (NAL Call No.: 450 P692).

0760

Responses of winter cereals to various low temperature stresses.
Pomeroy, M.K. Andrews, C.J. Columbia, Mo. : The University, 1983. Current topics in plant biochemistry and physiology, 1983 : proceedings, inaugural Plant Biochemistry and Physiology Symposium held, Univ. Missouri-Columbia, 7-9 April 1982 / editorial

committee: D.D. Randall ... (et al.). v. 2 p. 96-106. ill. Includes 22 references. (NAL Call No.: QK861.P55 1982).

0761

Role of benzoxazinones in allelopathy by rye (*Secale cereale* L.).
JCECD. Barnes, J.P. Putnam, A.R. New York, N.Y. : Plenum Press. Journal of chemical ecology. Apr 1987. v. 13 (4). p. 889-906. Includes references. (NAL Call No.: DNAL QD415.A1J6).

0762

Role of endogenous growth regulators in seed dormancy of *Avena fatua*. I. Short chain fatty acids (Wild oats).
Metzger, J.D. Sebesta, D.K. Rockville : American Society of Plant Physiologists. Plant physiology. Nov 1982. v. 70 (5). p. 1480-1485. 25 ref. (NAL Call No.: 450 P692).

0763

Role of endogenous plant growth regulators in seed dormancy of *Avena fatua* (Wild oats).
Metzger, J.D.PLPHA. Rockville : American Society of Plant Physiologists. Plant physiology. Nov 1983. v. 73 (3). p. 791-795. Includes references. (NAL Call No.: 450 P692).

0764

The role of hormone transport and metabolism in apical dominance in oats.
BOGAA. Harrison, M.A. Kaufman, P.B. Chicago, Ill. : University of Chicago Press. Botanical gazette. Sept 1984. v. 145 (3). p. 293-297. ill. Includes references. (NAL Call No.: DNAL 450 B652).

0765

Root development and lodging resistance in oats /Dale T. Sechler.
Sechler, Dale Truman, 1926-. Columbia, Mo. : University of Missouri, College of Agriculture, Agricultural Experiment Station, 1961. Cover title. 38 p. : ill. ; 23 cm. Bibliography: p. 37. (NAL Call No.: DNAL 100 M693 (3) no.769).

0766

Root excision decreases nutrient absorption and gas fluxes.
PLPHA. Bloom, A.J. Caldwell, R.M. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Aug 1988. v. 87 (4). p. 794-796. Includes references. (NAL Call No.: DNAL 450 P692).

0767

Salinity effects on rye grain yield, quality, vegetative growth, and emergence.

AGJOAT. Francois, L.E. Donovan, T.J.; Lorenz, K.; Maas, E.V. Madison, Wis. : American Society of Agronomy. Although current rye (*Secale cereale* L.) grain production is concentrated mainly in the northern half of the USA and Canada, some rye grain is grown in the arid southwest. Soils in this area are, or have the potential to become, highly saline from the application of saline irrigation water. Since there is neraly a complete lack of information about the response of rye grown under saline conditions, a 2-yr field plot study was conducted. Six salinity treatments were imposed on a Holtville silty clay (clayey over loamy, montmorillonitic calcareous, hyperthermic Typic Torrifluvent) by irrigating with Colorado River water artificially salinized with NaCl and CaCl₂ (1:1 by weight). Electrical conductivities of the irrigation waters were 1.1, 4.0, 8.0, 12.1, 16.0, and 20.1 dS m⁻¹ the first year, and 1.1, 3.9, 7.5, 11.6, 15.6, and 19.8 dS m⁻¹ the second year. Grain yield and vegetative growth were measured. Relative grain yield of two cultivars, Maton and Bonel, was unaffected up to a soil salinity of 11.4 dS m⁻¹ (electrical conductivity of the saturation extract;k(e)). Each unit increase in salinity above 11.4 dS m⁻¹ reduced yield by 10.8%. These results place rye in the salt-tolerant category. Yield reduction was attributed primarily to reduced spike weight and individual seed weight rather than spike number. Bread quality decreased slightly with increasing levels of salinity. Straw yield was more sensitive to salinity than was grain yield. Plant emergence was determined in greenhouse sand cultures. Both cultivars were slightly less salt tolerant during plant emergence than during subsequent stages of growth. *Agronomy journal*. Sept/Oct 1989. v. 81 (5). p. 707-712. Includes references. (NAL Call No.: DNAL 4 AM34P).

0768

Salinity stress induced tissue-specific proteins in barley seedlings.

PLPHA. Ramagopal, S. Rockville, Md. : American Society of Plant Physiologists. *Plant physiology*. June 1987. v. 84 (2). p. 324-331. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0769

Seed dormancy in *Avena fatua*. I. Induction of germination by mechanical injury (Wild oats).

Hsiao, A.I. McIntyre, G.I.; Hanes, J.A. Chicago : University of Chicago Press. *Botanical gazette*. June 1983. v. 144 (2). p. 217-222. Includes references. (NAL Call No.: 450 B652).

0770

Seed dormancy in wild and weedy relatives of cereals.

Key, J.M. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 15-25. ill. Includes references. (NAL Call No.: DNAL SB189.6.P73).

0771

Seedling competition between mountain rye, 'Hycrest' crested wheatgrass, and downy brome.

JRMGA. Buman, R.A. Monsen, S.B.; Abernethy, R.H. Denver, Colo. : Society for Range Management. *Journal of range management*. Jan 1988. v. 41 (1). p. 30-34. Includes references. (NAL Call No.: DNAL 60.18 J82).

0772

Simultaneous consideration of tissue and substrate potassium concentration in K⁺ (potassium ion) uptake kinetics: a model (With the example of barley).

Siddiqi, M.Y. Glass, A.D.M. Rockville, Md., American Society of Plant Physiologists. *Plant physiology*. Jan 1982. v. 69 (1). p. 283-285. 25 ref. (NAL Call No.: 450 P692).

0773

Sodium and potassium fluxes and compartmentation in roots of *Atriplex* and oat.

PLPHA. Mills, D. Robinson, K.; Hodges, T.K. Rockville, Md. : American Society of Plant Physiologists. *Plant physiology*. July 1985. v. 78 (3). p. 500-509. Includes 37 references. (NAL Call No.: DNAL 450 P692).

0774

Soil tests for copper, iron, manganese, and zinc in histosols: 4. Selection on the basis of soil chemical data and uptakes by oats, carrots, onions, and lettuce.

SOSCAK. Mathur, S.P. Levesque, M.P. Baltimore, Md. : Williams & Wilkins. *Soil science*. Dec 1989. v. 148 (6). p. 424-432. Includes references. (NAL Call No.: DNAL 56.8 S03).

0775

Solubilization of the fusicoccin receptor and a protein kinase from highly purified plasma membrane from oat roots.

NASSD. De Boer, A.H. Lomax, T.L.; Sandstrom, R.P.; Cleland, R.E. New York, N.Y. : Plenum Press. NATO advanced science institutes series : Series A : Life sciences. In the series analytic: Membrane receptors, dynamics, and energetics / edited by K.W.A. Wirtz. Proceedings of a NATO Advanced Study Institute,

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

August 17-30, 1986, Spetsai, Greece. 1987. v. 133. p. 181-190. Includes references. (NAL Call No.: DNAL QH301.N32).

0776

Sources of ammonium in oat leaves treated with tabtoxin or methionine sulfoximine (*Avena sativa*).

Frantz, T.A. Peterson, D.M.; Durbin, R.D. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Feb 1982. v. 69 (2). p. 345-348. Includes 35 ref. (NAL Call No.: 450 P692).

0777

Sporulation of oat-cultured *Physarum polycephalum*. I. Sporulation competence of dark starved plasmodia (Fungi).

Hosoda, E. Bronx, N.Y., New York Botanical Garden. Mycologia. July/Aug 1981. v. 73 (4). p. 689-696. 111. 18 ref. (NAL Call No.: 450 M99).

0778

Sporulation of oat-cultured *Physarum polycephalum*. II. Sporulation competence of slow desiccation-induced sclerotia (Fungi).

Hosoda, E. MYCOA. Bronx : The New York Botanical Garden. Mycologia. Jan/Feb 1983. v. 75 (1). p. 14-21. 16 ref. (NAL Call No.: 450 M99).

0779

Steady state proline levels in salt-shocked barley leaves.

PLPHA. Voetberg, G. Stewart, C.R. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Nov 1984. v. 76 (3). p. 567-570. 111. Includes 20 references. (NAL Call No.: DNAL 450 P692).

0780

Stem reserves and the response of barley to elevated post-anthesis temperature.

Cornish, P.S. Lockley, P. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 258. (NAL Call No.: DNAL SB191.B2A9 1987).

0781

Sterility in rye.

AGJDAT. Stroman, G.N. Madison, Wis. : American Society of Agronomy. Agronomy journal. June 1923. v. 15 (6). p. 253-254. (NAL Call No.: DNAL 4 AM34P).

0782

Sterility of oats /by Charlotte Elliott.

Elliott, Charlotte, 1883-. Washington, D.C. : U.S. Dept. of Agriculture, 1922. Caption title.~ "March 6, 1922."~ "Professional paper.". 8 p., 4 p. of plates ; 24 cm. (NAL Call No.: DNAL 1 Ag84B no.1058).

0783

Sterility of rye.

AGJDAT. Leith, B.D. Madison, Wis. : American Society of Agronomy. Agronomy journal. Mar 1925. v. 17 (3). p. 129-132. (NAL Call No.: DNAL 4 AM34P).

0784

Stimulation of glutathione synthesis in photorespiring plants by catalase inhibitors.

PLPHA. Smith, I.K. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Dec 1985. v. 79 (4). p. 1044-1047. Includes 17 references. (NAL Call No.: DNAL 450 P692).

0785

Stimulatory and inhibitory effects of light upon auxin biosynthesis (*Avena*, oats).

Wolf, F.T. Nashville, Tenn. : The Academy. Journal of the Tennessee Academy of Science. Jan/Apr 1984. v. 59. p. 25-27. Includes references. (NAL Call No.: 500 T25A).

0786

Storage protein accumulation in 'Scio' barley seed as affected by late foliar applications of nitrogen.

CRPSAY. Turley, R.H. Ching, T.M. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 778-782. Includes references. (NAL Call No.: DNAL 64.8 C883).

0787

Structural changes in thylakoid proteins during cold acclimation and freezing of winter rye (*Secale cereale* L. cv. Puma).

Griffith, M. Brown, G.N.; Huner, N.P.A. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Aug 1982. v. 70 (2). p. 418-423. 111. 35 ref. (NAL Call No.: 450 P692).

0788

Studies of self-fertilization in rye /by H.E. Brewbaker.
 Brewbaker, H. E. 1895-. St. Paul : University Farm, 1926. Cover title.~ Originally presented as Thesis (Ph. D.)--University of Minnesota, 1926. 40 p. : ill. ; 23 cm. Bibliography: p. 30-32. (NAL Call No.: DNAL 100 M66 (3) no.40).

0789

Studies of the regulation of nitrate influx by barley seedlings using 13N03-1.
 PLPHA. Siddiqi, M.Y. Glass, A.D.M.; Ruth, T.U.; Fernando, M. Rockville, Md. : American Society of Plant Physiologists. Using 13N03-, effects of various N03- pretreatments upon N03- influx were studied in intact roots of barley (*Hordeum vulgare* L. cv Klondike). Prior exposure to roots to N03- increased N03- influx and net N03- uptake. The 'induction' of N03- uptake was dependent both on time and external N03- concentration (N030). During induction influx was positively correlated with root N03-. In the postinduction period, however, N03- influx declined as root N03- increased. It is suggested that induction and negative feedback regulation are independent processes: induction appears to depend upon some critical cytoplasmic N03-; removal of external N03- caused a reduction of 13N03- influx even though mean root N03- remained high. It is proposed that cytoplasmic N03- is depleted rapidly under these conditions resulting in 'deinduction' of the N03- transport system. Beyond 50 micromoles per gram N03-, 13N03- influx was negatively correlated with root N03-. However, it is unclear whether root N03- per se or some product(s) of N03- assimilation are responsible for the negative feedback effects. Plant physiology. July 1989. v. 90 (3). p. 806-813. Includes references. (NAL Call No.: DNAL 450 P692).

0790

Studies on plant absorption of Fe (iron) humate (Extraction from Brunizem soil, *Hordeum vulgare*, barley).
 Orioli, G.A. New York ; Basel : Marcel Dekker, 1982. Iron nutrition and interactions in plants : Brigham Young University, August 12-14, 1981 / edited by S.D. Nelson ... (et al.). Abstract only. p. 413. (NAL Call No.: QK867.J67 v. 5, nos. 4-7).

0791

Studies on the effect of nitrogen applied to oats at different periods of growth.
 AGJOAT. Gericke, W.F. Madison, Wis. : American Society of Agronomy. Agronomy journal. Nov 1922. v. 14 (8). p. 312-320. (NAL Call No.: DNAL 4 AM34P).

0792

Studies with 4,6-dioxoheptanoic acid on etiolated and greening barley leaves (Varieties).
 Meller, E. Gassman, M.L. Rockville, Md., American Society of Plant Physiologists. Plant physiology. June 1981. v. 67 (6). p. 1065-1068. ill. 18 ref. (NAL Call No.: 450 P692).

0793

Sugar composition and freezing tolerance in barley crowns at varying carbohydrate levels.
 CRPSAY. Livingston, D.P. III. Olien, C.R.; Freed, R.D. Madison, Wis. : Crop Science Society of America. During hardening, carbohydrate content in winter cereals is affected by light intensity, which in turn affects freezing survival. To determine whether total carbohydrate content affects composition of individual sugars and whether a particular composition is related to freezing tolerance, three barley (*Hordeum vulgare* L.) cultivars, differing in carbohydrate composition and in response to a controlled freezing test, were hardened under six light levels. At each level, total crown sugars were measured by water/ethanol extraction and high-pressure liquid chromatography and at four light levels plants were freeze tested. Carbohydrate composition changed depending on total carbohydrate content, and maximum freezing resistance was not found at the maximum simple sugar level. In addition, the relationship between kill temperature and total carbohydrate was different for the three cultivars. A simple relationship between carbohydrate composition and freezing tolerance was not found, but results suggest that the three cultivars may allocate total carbohydrate differently for cryoprotection. Crop science. Sept/Oct 1989. v. 29 (5). p. 1266-1270. Includes references. (NAL Call No.: DNAL 64.8 C883).

0794

Survey of barley germplasm for betaine-accumulating potential in controlled environments and in the field (*Hordeum vulgare*, *Hordeum spontaneum*, genotypes).
 Ladyman, J.A.R. Ditz, K.M.; Hanson, A.D. East Lansing, Mich., The Laboratory. Annual report - Michigan State University, MSU/DOE Plant Research Laboratory. 1981. 1981 (16th). p. 64-66. 3 ref. (NAL Call No.: QK1.M5).

0795

Synthesis and degradation of barley nitrate reductase (*Hordeum vulgare*).
 Somers, D.A. PLPHA. Kuo, T.M.; Kleinhofs, A.; Warner, R.L.; Oaks, A. Rockville : American Society of Plant Physiologists. Plant physiology. Aug 1983. v. 72 (4). p. 949-952. ill. Includes references. (NAL Call No.: 450 P692).

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

0796

Techniques for selection of cold hardiness in cereals (Oats).

Marshall, G.H. Olien, C.R.; Everson, E.H. Boca Raton, Fla., CRC Press. Analysis and improvement of plant cold hardiness. 1981. 1981. p. 139-159. 30 ref. (NAL Call No.: SB781.A52).

0797

Temperature requirements for mountain rye, Hycrest crested wheatgrass, and downy brome germination.

JRMGA. Buman, R.A. Abernethy, R.H. Denver, Colo. : Society for Range Management. Journal of range management. Jan 1988. v. 41 (1). p. 35-39. Includes references. (NAL Call No.: DNAL 60.18 J82).

0798

Three alcohol dehydrogenase genes in wild and cultivated barley: characterization of the products of variant alleles (Hordeum vulgare, Hordeum spontaneum).

Hanson, A.D. Brown, A.H.D. New York, N.Y. : Plenum Press. Biochemical genetics. June 1984. v. 22 (5/6). p. 495-515. ill. Includes references. (NAL Call No.: QR73.B5).

0799

Tillering dynamics of winter barley as influenced by cultivar and nitrogen fertilizer: a field study (Hordeum vulgare, Hordeum distichon L., tillers survival).

Garcia del Moral, L.F. Ramos, J.M.; Recalde, L. Madison : Crop Science Society of America. Crop science. Jan/Feb 1984. v. 24 (1). p. 179-181. ill. Includes references. (NAL Call No.: 64.8 C883).

0800

Tolerances of oat cultivars to an acid soil high in exchangeable aluminum.

JPNUDS. Foy, C.D. Smith, D.H. Jr.; Briggie, L.W. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Tenth International Plant Nutrition Colloquium," August 4-9, 1986, Beltsville, Maryland. 1987. v. 10 (9116). p. 1163-1174. ill. Includes references. (NAL Call No.: DNAL QK867.J67).

0801

Toxic factors in acid soils.

Foy, C.D. Burns, G.R. Washington, D.C. : National Plant Food Institute. Plant food review. Fall 1964. v. 10 (3). p. 2-3, 16. ill. (NAL Call No.: DNAL 57.8 P694).

0802

Transformation of the cryobehavior of rye protoplasts by modification of the plasma membrane lipid composition.

PNASA. Steponkus, P.L. Uemura, M.; Balsamo, R.A.; Arvinte, T.; Lynch, D.V. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. Dec 1988. v. 85 (23). p. 9026-9030. ill. Includes references. (NAL Call No.: DNAL 500 N21P).

0803

Transpiration efficiency of oat.

AGJOAT. Ehlers, W. Madison, Wis. : American Society of Agronomy. Dry matter production, water use, and their ratio (i.e., the water use efficiency WUE of crops) vary among species and with climatic conditions. This study was undertaken to determine the WUE of oat (*Avena sativa* L.) because recent field evaluations for this crop are not available. Oat was grown on loess-derived silt loam fine-silty, mixed, mesic Typic Hapludalf near Goettingen, West Germany, during 4 yr. Periodically, shoot and root dry matter were determined. Water use within five consecutive growth stages was evaluated by the soil water budget approach. Balanced evapotranspiration (ET), including evaporation from soil (E), water intercepted by leaves (I), and transpiration from plants (T), was divided into its components, E, I, and T by calculation procedures. The term E was approximated from meteorological factors, leaf area index (LAI), and soil water tension, and I from daily precipitation and LAI. Shoot and total oat dry matter including roots were linearly related to cumulative T, IT, and ET. The X-intercept of the regression line was nearest to zero, when X was IT. Therefore, I was regarded to be part of "productive" water use. Including I and considering only shoot dry matter, WUE(IT) was 4.1 kg m⁻³. When considering shoots and roots, WUE(IT) increased to 4.5 kg m⁻³. The relation between dry matter production and IT was improved by relating IT to the potential evapotranspiration rate or the saturation deficit (delta epsilon) of the air. As compared with other crops, WUE(IT) of oat appeared to be lower. This study provided evidence that normalization of IT by delta epsilon requires the appropriate evaluation of delta epsilon for the daylight period when transpiration occurs. *Agronomy journal*. Sept/Oct 1989. v. 81 (5). p. 810-817. Includes references. (NAL Call No.: DNAL 4 AM34P).

0804

Twenty-four-hour induction of freezing and drought tolerance in plumules of winter rye seedlings by desiccation stress at room temperature in the dark.

Siminovitch, D. Cloutier, Y. Rockville, Md., American Society of Plant Physiologists. *Plant physiology*. Jan 1982. v. 69 (1). p. 250-255. ill. 25 ref. (NAL Call No.: 450 P692).

0805

(NAL Call No.: DNAL 450 P692).

Uptake and subcellular compartmentation of gibberellin A1 applied to leaves of barley and cowpea.

Ohlrogge, J.B. AR-NRRC. Garcia-Martinez, J.L.; Adams, D.; Rappaport, L. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1980. v. 66 (3). p. 422-427. ill. 33 ref. (NAL Call No.: 450 P692).

0806

The uptake, distribution and metabolism of four organic chemicals by soybean plants and barley roots.

ETOCK, McFarlane, C. Nolt, C.; Wickliff, C.; Pfleeger, T.; Shimabuku, R.; McDowell, M. Elmsford : Pergamon Press. Environmental toxicology and chemistry. 1987. v. 6 (11). p. 847-856. ill. Includes references. (NAL Call No.: DNAL QH545.A1E58).

0807

Uptake of phenylalanine into isolated barley vacuoles is driven by both tonoplast adenosine triphosphatase and pyrophosphatase: evidence for a hydrophobic L-amino acid carrier system. PLPHA. Homeyer, U. Litek, K.; Huchzermeyer, B.; Schultz, G. Rockville, Md. : American Society of Plant Physiologists. The uptake of phenylalanine was studied with vacuole isolated from barley mesophyll protoplasts. The phenylalanine transport exhibited saturation kinetics with apparent Km-values of 1.2 to 1.4 millimolar for ATP- or PPi-driven uptake; Vmax apparent was 120 to 140 nanomoles Phe per milligram of chlorophyll per hour (1 milligram of chlorophyll corresponds to 5 X 10⁶ vacuoles). Half-maximal transport rates driven with ATP or PPi were reached at 0.5 millimolar ATP or 0.25 millimolar PPi. ATP-driven transport showed a distinct pH optimum at 7.3 while PPi-driven transport reached maximum rates at pH 7.8. Direct measurement of the H⁺-translocating enzyme activities revealed Km apparent values of 0.45 millimolar for ATPase (EC 3.6.1.3) and 23 micromolar for pyrophosphatase (PPase) (EC 3.6.1.1). In contrast to the coupled amino acid transport, ATPase and PPase activities had relative broad pH optima between 7 to 8 for ATPase and 8 to 9 for PPase. ATPase as well as ATP-driven transport was markedly inhibited by nitrate while PPase and PPi-coupled transport was not affected. The addition of ionophores inhibited phenylalanine transport suggesting the destruction of the electrochemical proton potential difference while the rate of ATP and PPi hydrolysis was stimulated. The uptake of other lipophilic amino acids like L-Trp, L-Leu, and L-Tyr was also stimulated by ATP. They seem to compete for the same carrier system. L-Ala, L-Val, D-Phe, and D-Leu did not influence phenylalanine transport suggesting a stereospecificity of the carrier system for L-amino acids having a relatively high hydrophobicity. Plant physiology. Apr 1989. v. 89 (4). p. 1388-1393. ill. Includes references.

0808

Use of a mathematical model to determine the fate of atrazine in barley (*Hordeum vulgare*). WEESA6. Raynaud, S. Bastide, J.; Coste, C. Champaign, Ill. : Weed Science Society of America. Weed science. Nov 1985. v. 33 (6). p. 906-912. Includes 28 references. (NAL Call No.: DNAL 79.8 W41).

0809

Use of 31P NMR to assess effects of DNP on ATP levels in vivo in barley roots. PLPHA. Jackson, P.C. Pfeffer, P.E.; Gerasimowicz, W.V. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Aug 1986. v. 81 (4). p. 1130-1133. Includes 11 references. (NAL Call No.: DNAL 450 P692).

0810

Utilization of microbial siderophores in iron acquisition by oat. PLPHA. Crowley, D.E. Reid, C.P.P.; Szaniszlo, P.J. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. July 1988. v. 87 (3). p. 680-685. Includes references. (NAL Call No.: DNAL 450 P692).

0811

Vacuolar deposition of ascorbate-derived oxalic acid in barley. Wagner, G.J. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Mar 1981. v. 67 (3). p. 591-593. 15 ref. (NAL Call No.: 450 P692).

0812

Varietal variation of water sensitivity in Asian barley varieties. Takeda, K. Fukuyama, T. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Oct 10, 1985. v. 15. p. 15-16. (NAL Call No.: DNAL QK495.G74B34).

0813

Victorin-induced changes in carbohydrate metabolism in oat leaves / by Carroll David Rawn. - Rawn, Carroll David. Ann Arbor, Mich. University Microfilms International 1978. Thesis--University of Kentucky, 1974. Facsimile produced by microfilm-xerography. ix, 79, 3 leaves. Bibliography: leaves 73-78. (NAL Call

(PLANT PHYSIOLOGY AND BIOCHEMISTRY)

No.: DISS 75-18,508).

0820

5,6-dichloroindole-3-acetic acid as a potent auxin and 5,7-dichloroindole-3-isobutyric acid as potent anti-auxin: their synthesis and biological activities.

PPGGD. Hatano, T. Katayama, M.; Marumo, S. Lake Alfred, Fla. : The Society. Proceedings annual meeting - Plant Growth Regulator Society of America. 1987. (14th). p. 353-358. Includes references. (NAL Call No.: DNAL SB128.P5).

0814

Water stress enhances expression of an alpha-amylase gene in barley leaves.

PLPHA. Jacobsen, J.V. Hanson, A.D.; Chandler, P.C. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Feb 1986. v. 80 (2). p. 350-359. ill. Includes 33 references. (NAL Call No.: DNAL 450 P692).

0815

Wild oat (*Avena fatua*) and spring barley (*Hordeum vulgare*) growth and development in monoculture and mixed culture.

WEESA6. Morishita, D.W. Thill, D.C. Champaign, Ill. : Weed Science Society of America. Weed science. Jan 1988. v. 36 (1). p. 43-48. Includes references. (NAL Call No.: DNAL 79.8 W41).

0816

Wild oat (*Avena fatua*) seed environment and germination (North Dakota).

Somody, C.N. Nalewaja, J.D.; Miller, S.D. Champaign, Ill. : Weed Science Society of America. Weed science. July 1984. v. 32 (4). p. 502-507. ill. Includes 11 references. (NAL Call No.: 79.8 W41).

0817

Wound-induced resistance to cellulase in oat leaves (*Avena sativa*).

Geballe, G.T. Galston, A.W. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1982. v. 70 (3). p. 781-787. ill. 20 ref. (NAL Call No.: 450 P692).

0818

Zinc-inhibited electron transport of photosynthesis in isolated barley chloroplasts.

Tripathy, B.C. Mohanty, P. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Dec 1980. v. 66 (6). p. 1174-1178. ill. 25 ref. (NAL Call No.: 450 P692).

0819

4-Amino-5-hexynoic acid--a potent inhibitor of tetrapyrrole biosynthesis in plants.

PLPHA. Elich, T.D. Lagarias, J.C. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Nov 1988. v. 88 (3). p. 747-751. Includes references. (NAL Call No.: DNAL 450 P692).

PLANT TAXONOMY AND GEOGRAPHY

0821

Geographical distribution of diazinon sensitive varieties in barley.

Takeda, K. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Aug 31, 1987. v. 17. p. 90-93. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0827

Wild oat identification and control.

Strand, Oliver E. Document available from: University of Minnesota, Bulletin Room, 1420 Eckles Avenue, St. Paul, Minnesota 55108 1981. Explores identification and control for wild oat. 1 sheet : ill. (NAL Call No.: Document available from source.)(NAL Call No.: Fs No.9).

0822

Origin, taxonomy, and related species.

AGRYA. Bothmer, R. von. Jacobsen, N. Madison, Wis., Soil Science Society of America. Agronomy. A series of monographs - American Society of Agronomy. Literature review. 1985. (26). p. 19-56. maps. Includes references. (NAL Call No.: DNAL 4 AM392).

0823

Pulse domestication and cereal domestication: how different are they?.

ECBOA. Zohary, D. Bronx, N.Y. : New York Botanical Garden. Economic botany. Jan/Mar 1989. v. 43 (1). p. 31-34. Includes references. (NAL Call No.: DNAL 450 EC7).

0824

Rapid electrophoresis of oat (*Avena sativa* L.) prolamins from single seeds for cultivar identification.

CECHAF. Hansen, A.E. Nassuth, A.; Altosaar, I. St. Paul, Minn. : American Association of Cereal Chemists. Cereal chemistry. Mar/Apr 1988. v. 65 (2). p. 153-154. ill. Includes references. (NAL Call No.: DNAL 59.8 C33).

0825

Seed dormancy in wild and weedy relatives of cereals.

Key, J.M. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 15-25. ill. Includes references. (NAL Call No.: DNAL SB189.6.P73).

0826

Weed control in conservation tillage systems--small grains.

Wicks, G.A. Champaign, Il. : Weed Science Society of America. Monograph series of the Weed Science Society of America. Literature review. 1985. (2). p. 77-92. Includes references. (NAL Call No.: DNAL SB610.M65).

PROTECTION OF PLANTS

0828

Ag facts: Barley diseases.

MUCBA. Clayton, J.L. Hart, L.P. East Lansing, Mich. : The Service. Extension bulletin E - Cooperative Extension Service, Michigan State University. June 1986. (1982). 8 p. Includes references. (NAL Call No.: DNAL 275.29 M58B).

0829

Ag facts: Oat diseases.

MUCBA. Clayton, J.L. Hart, L.P. East Lansing, Mich. : The Service. Extension bulletin E - Cooperative Extension Service, Michigan State University. June 1986. (1981). 3 p. Includes references. (NAL Call No.: DNAL 275.29 M58B).

0830

Ag facts: Rye diseases.

MUCBA. Clayton, J.L. Hart, L.P. East Lansing, Mich. : The Service. Extension bulletin E - Cooperative Extension Service, Michigan State University. June 1986. (1980). 3 p. Includes references. (NAL Call No.: DNAL 275.29 M58B).

0831

Compendium of barley diseases /D.E. Mathre, editor.

Mathre, Don E., 1938-. St. Paul, Minn. : American Phytopathological Society ; Bozeman : Dept. of Plant Pathology, Montana State University, 1982. v, 78 p., 16 p. of plates : ill. (some col.) ; 28 cm. Includes bibliographies and index. (NAL Call No.: DNAL SB608.B2M37).

0832

Montana small grain problem diagnostic guide and disease index.

Inglis, D. (comp.). Bozeman, Mont. : The Service. Circular - Montana State University, Cooperative Extension Service. June 1986. (1301). 15 p. (NAL Call No.: DNAL 275.29 M76CI).

0833

Pest control in forages and small grains--1981 (Includes control of weeds, insects, diseases).

Doll, J.D. Wedberg, J.L.; Grau, C.R.; Doersch, R.E. Madison, Wis., The Programs. Publication - Cooperative Extension Programs, University of Wisconsin Extension. June 1981. June 1981. (A1981). 36 p. (NAL Call No.: S544.3.W6W53).

0834

Preservation and storage of grains, seeds, and their by-products cereals, oilseeds, pulses, and animal feed /edited by J.L. Multon ; preface by A.M. Reimbert ; translated from the French by D. Marsh ; reread by A.J. Eydt.

Multon, J. L. 1938-. New York, N.Y. : Lavoisier Pub., c1988. Translated from French.~ Translation of: Conservation et stockage des grains et graines et produits d'origine.~ Originally published: Lavoisier : Technique et documentation, 1982. xxiv, 1095 p. : ill. ; 25 cm. Includes bibliographies and index. (NAL Call No.: DNAL SB190.C6513).

0835

Profitable small grain production in the Texas Gulf Coast.

TAEB. Miller, T.D. Livingston, S. College Station, Tex. : The Station. B - Texas Agricultural Experiment Station. Nov 1987. (1587). 8 p. (NAL Call No.: DNAL 100 T31S (1)).

0836

Subsurface drip irrigation of cotton and small grains.

Tollefson, S. St. Joseph, Mich. : American Society of Agricultural Engineers, c1985. Drip/trickle irrigation in action : proceedings of the Third International Drip/Trickle Irrigation Congress, November 18-21, 1985, Centre Plaza Holiday Inn, Fresno, California, USA. p. 887-895. Includes 7 references. (NAL Call No.: DNAL S612.I5 1985).

0837

Using crop diversity to manage pest problems: some California examples.

Flint, M.L. Roberts, P.A. Greenbelt, Md. : Institute for Alternative Agriculture. Moderate to large scale California growers (as well as small scale ones) manipulate cropping patterns in a number of ways to reduce pest problems. Crop rotation, which can be defined as diversifying crops over time, is used to manage selected pests, primarily weeds, pathogens, and nematodes. As a substitute for pesticides, crop rotation has been most rewarding in the control of nematodes; sugarbeet cyst and root knot nematode examples are detailed. Some pests that invade fields from nearby areas can be managed by modifying adjacent cropping patterns or practices; Pierce's disease of grapes, sugarbeet yellows and border harvesting of alfalfa are given as examples. Finally, multiple crops can be grown within a single field or orchard. Although this approach is not widely practiced by many California growers, two examples of systems where intercropping has been shown to limit pest numbers without the use of pesticides are described: intercropping of cotton with alfalfa and companion planting oats when seeding alfalfa. These examples show that using crop diversification to manage pests

(PROTECTION OF PLANTS)

is feasible, but growers must be strongly motivated to make the necessary changes in cropping patterns. Most of the systems that have been widely adopted are those for which few other economically feasible methods were available. American journal of alternative agriculture. Fall 1988. v. 3 (4). p. 163-167. Includes references. (NAL Call No.: DNAL S605.5.A43).

PESTS OF PLANTS - GENERAL AND MISC.

0838

Breeding for pest resistance.

AGRYA. Sharp, E.L. Madison, Wis., Soil Science Society of America. Agronomy. A series of monographs - American Society of Agronomy. Literature review. 1985. (26). p. 313-333. Includes references. (NAL Call No.: DNAL 4 AM392).

0839

Can goose damage to grain fields be prevented through methiocarb-induced aversive conditioning?

WLSBA. Conover, M.R. Bethesda, Md. : The Society. Wildlife Society bulletin. Summer 1989. v. 17 (2). p. 172-175. Includes references. (NAL Call No.: DNAL SK357.A1W5).

0840

Compensation for vertebrate pest damage (Cereals, forage, livestock, bees, Alberta, Canada).

Gurba, J.B.PVPCB. Davis, Calif. : University of California. Proceedings ... Vertebrate Pest Conference. June 1982. June 1982. (10th). p. 90-94. ill. Includes references. (NAL Call No.: SB950.A1V4).

0841

Coordinator's report: disease and pest resistance genes.

Jorgensen, J.H. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 65-69. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0842

Effect of grazing by Canada geese on the winter growth of rye.

JWMAA9. Conover, M.R. Bethesda, Md. : Wildlife Society. Journal of wildlife management. Jan 1988. v. 52 (1). p. 76-80. Includes references. (NAL Call No.: DNAL 410 J827).

0843

Methods for assessing the efficiency of aerial spraying control operations on quelea colonies and roosts (Bird control, damage to the smaller cereal grain crops in Africa).

Elliott, C.C.H. Philadelphia : American Society for Testing and Materials, 1981. Vertebrate pest control and management materials : proceedings of the Third Symposium on Test Methods for Vertebrate Pest Control and Management Materials, Fresno, Calif., 7 March 1980 : a symposium. p. 62-73. ill. 10 ref. (NAL Call No.: SB950.A2S95 1980).

0844

Pest control in forages and small grains.

Doll, J. D. Grau, C. R.; Doersch, R. E.; Wedberg, J. L. 1981. This publication discusses the various methods of pest control in forages and small grains. Document available from: Agricultural Bulletin Building, 1535 Observatory Drive, University of Wisconsin, Madison, WI 53706. 36 p. (NAL Call No.: A 1981).

0845

The rodent problem in Latin America.

Elias, D.J. Fall, M.W. Boca Raton, Fla. : CRC Press, c1988. Rodent pest management / editor, Ishwar Prakash. Literature review. p. 13-28. maps. Includes references. (NAL Call No.: DNAL SB994.R6R65).

0846

Suggested methods for determining the efficacy of vertebrate control agents in developing countries (Cereal crop losses, bird, rodent pests, Africa, Philippines).

Bruggers, R.L. Jackson, W.B. Philadelphia : American Society for Testing and Materials, 1981. Vertebrate pest control and management materials : proceedings of the Third Symposium on Test Methods for Vertebrate Pest Control and Management Materials, Fresno, Calif., 7 March 1980 : a symposium. p. 15-28. ill. 31 ref. (NAL Call No.: SB950.A2S95 1980).

0847

Third international symposium on pre-harvest sprouting in cereals / edited by James E. Kruger and Donald E. LaBerge. -.

Kruger, James E.; LaBerge, Donald E. Boulder, Colo. Westview Press 1983. xii, 312 p. : ill. ; 24 cm. Includes bibliographies and index. (NAL Call No.: SB608.G6T45).

PESTS OF PLANTS - INSECTS

0848

Alabama small grains damaged by Hessian fly.
HARAA. Estes, P.M. Johnson, W.; Little, J.; Moore, D. Auburn, Ala. : The Station. Highlights of agricultural research - Alabama, Agricultural Experiment Station. Fall 1985. v. 32 (3). p. 16. ill. (NAL Call No.: DNAL 100 AL1H).

0849

Aphid control on small grains (Schizaphis graminum, Rhopalosiphum padi, Macrosiphum (Sitobion) avenae, Acyrthosiphon (Metopolophium) dirhodum, Rhopalosiphum maidis, Washington).
Pike, P. Retan, A. Pullman, Wash., The Service. Extension Bulletin - Washington State University, Cooperative Extension Service. Mar 1982. Mar 1982. (1001). 4 p. ill. (NAL Call No.: 275.29 W27P).

0850

Aphid feeding behavior: relationship to barley yellow dwarf virus resistance in Agropyron species.
PHYTAJ. Shukle, R.H. Lampe, D.J.; Lister, R.M.; Foster, J.E. St. Paul, Minn. : American Phytopathological Society. Phytopathology. May 1987. v. 77 (5). p. 725-729. ill. Includes references. (NAL Call No.: DNAL 464.8 P56).

0851

Assessment of experimental designs for greenbug (Homoptera: Aphididae) antixenosis tests.
JEENAI. Webster, J.A. Inayatullah, C. College Park, Md. : Entomological Society of America. Journal of economic entomology. Aug 1988. v. 81 (4). p. 1246-1250. Includes references. (NAL Call No.: DNAL 421 J822).

0852

Biological and small grain resistance experiments with biotype E greenbug (Schizaphis graminum).
Daniels, N.E. Chedester, L.D. College Station, Tex., The Station. PR - Texas Agricultural Experiment Station. May 1981. May 1981. (3870). 7 p. ill. 11 ref. (NAL Call No.: 100 T31P).

0853

Biological observations of Menochilus sexmaculatus (potential biological control agent), reared on Schizaphis graminum (A major pest of small grain crops).
Campbell, R.K. Farris, T.N.; Perring, T.M.; Leonard, M.E.; Cartwright, B.O.; Eikenbary, R.D. College Park, Md., The Society. Annals of the Entomological Society of

America. Entomological Society of America. Mar 15, 1980. v. 73 (2). p. 153-157. ill. 20 ref. (NAL Call No.: 420 EN82).

0854

Biology and control of cereal leaf beetle using natural enemies (authors: S.P. Briggs ... (et al.). -.
Briggs, S. P. (Blacksburg, Va.) Chemical, Drug and Pesticide Unit, Virginia Tech (1980?). Pesticide Applicator Training collection. 56 slides : col. + 1 sound cassette (17:50 min.) + 1 script. (NAL Call No.: Slide no.60).

0855

Biotype C and E greenbugs: distribution in the Texas rolling plains and damage to four small grain varieties (Schizaphis graminum).
Puterka, G.J. Slosser, J.E.; Gilmore, E.C. College Station, Tex., Southwestern Entomological Society. The Southwestern entomologist. Mar 1982. v. 7 (1). p. 4-8. map. Includes 5 ref. (NAL Call No.: QL461.S65).

0856

Cereal leaf beetle.
Matthew, D.L. Jr. Edwards, C.R. Lafayette : The Service. Publication E - Purdue University, Cooperative Extension Service. Jan 1979. (20). 2 p. ill. (NAL Call No.: DNAL SB844.I6P8).

0857

Cereal leaf beetle.
Matthew, David L. Jr. Edwards, Richard C. & Field crops insects. Document available from: Purdue University, Publication Mailing Room, 301 South Second Street, Lafayette, Indiana 47905 1979. Includes cereal leaf beetle host plants, type of damage, life cycle and control measures. 1 sheet : ill. (NAL Call No.: Document available from source.)(NAL Call No.: E-20).

0858

The cereal leaf beetle in North Carolina.
Robertson, R.L. Van Duyn, J.W.; Brandenburg, R.L. Raleigh, N.C. : The Service. AG - North Carolina Agricultural Extension Service, North Carolina State University. Mar 1988. (324,rev.). 5 p. maps. (NAL Call No.: DNAL S544.3.N6N62).

(PESTS OF PLANTS - INSECTS)

0859

The cereal leaf beetle in North Carolina (Life cycle, biological control, chemical control). Robertson, R.L. Van Duyn, J.W. Raleigh, N.C. : The Service. AG - North Carolina Agricultural Extension Service, North Carolina State University. July 1983. July 1983. (324). 4 p. ill., maps. (NAL Call No.: S544.3.N6N62).

0860

The cereal leaf beetle in Utah. Karren, J.B. Logan, Utah : The Service. Extension leaflet - Cooperative Extension Service, Utah State University. Feb 1986. (212). 8 p. ill. Includes references. (NAL Call No.: DNAL 275.29 UT1L).

0861

Cereal leaf beetle (*Oulema melanopus*) pupation under controlled temperatures and relative humidities (Insect migration, barley). Wellso, S.G. Hoxie, R.P. College Park, Md., Entomological Society of America. Environmental entomology. Feb 1981. v. 10 (1). p. 58-61. 6 ref. (NAL Call No.: QL461.E532).

0862

Cereal leaf beetle under control (Biological control by wasps). Wellso, S.G. Madison, Wis., American Society of Agronomy. Crops and soils magazine. Aug/Sept 1982. v. 34 (9). p. 7-9. ill. (NAL Call No.: 6 W55).

0863

The comparative attractiveness of various small grains to the chinch bug /by Curtis Benton and W.P. Flint. Benton, Curtis, 1898-. Flint, W. P. 1882-. Washington, D.C. : U.S. Dept. of Agriculture, 1938. Caption title.~ Contribution from Bureau of Entomology and Plant Quarantine in cooperation with the Illinois Agricultural Experiment Station. 8 p. ; 23 cm. (NAL Call No.: DNAL 1 Ag84C no.508).

0864

Compatibility of *Lysiphlebus testaceipes* (Hymenoptera: Braconidae) with greenbug (Homoptera: Aphididae) biotypes "C" and "E" reared on susceptible and resistant oat varieties (*Schizaphis graminum*). Salto, C.E. EVETB. Eikenbary, R.D.; Starks, K.J. College Park : Entomological Society of America. Environmental entomology. Apr 1983. v. 12 (2). p. 603-604. Includes references. (NAL Call No.: QL461.E532).

0865

Constant and fluctuating temperature effects on developmental rates and life table statistics of the greenbug (Homoptera: Aphididae). JEENAI. Walgenbach, D.D. Elliott, N.C.; Kieckhefer, R.W. College Park, Md. : Entomological Society of America. Journal of economic entomology. Apr 1988. v. 81 (2). p. 501-507. Includes references. (NAL Call No.: DNAL 421 J822).

0866

Designation of two new greenbug (Homoptera: Aphididae) biotypes G and H. JEENAI. Puterka, G.J. Peters, D.C.; Kerns, D.L.; Slosser, J.E.; Bush, L.; Worrall, D.W.; McNew, R.W. Lanham, Md. : Entomological Society of America. Journal of economic entomology. Dec 1988. v. 81 (6). p. 1754-1759. Includes references. (NAL Call No.: DNAL 421 J822).

0867

Developing small grains resistant to the cereal leaf beetle / by James A. Webster and David H. Smith, Jr. . Webster, James A. Smith, David H. Washington : U.S. Dept. of Agriculture, Agricultural Research Service, 1983. Cover title.~ Distributed to depository libraries in microfiche.~ "Issued August 1983"--T.p. verso. 9 p. : ill. ; 28 cm. Bibliography: p. 9- 10 . (NAL Call No.: DNAL 1 Ag84Te no.1673).

0868

Development of a model for on-line control of the cereal leaf beetle (*Oulema melanopus* (L.)) /by Winston Cordell Fulton. --. Fulton, Winston Cordell. 1978. Thesis (Ph. D.)--Michigan State University, 1978. Photocopy. Ann Arbor, Mich. : University Microfilms International, 1986. ix, 130 leaves : ill. ; 21 cm. Bibliography: leaves 127-130. (NAL Call No.: DNAL DISS 79-00.692).

0869

Effect of barley residue on *Rhizoctonia* crown rot of sugarbeet (*Rhizoctonia solani*, Wyoming). Fernandez, J.A. Laramie, Wyo., The Station. Research journal - University of Wyoming. Agricultural Experiment Station. Jan 1982. Jan 1982. (171). p. 15-17. (NAL Call No.: S131.E22).

(PESTS OF PLANTS - INSECTS)

0870

Effect of culture-host preconditioning on greenbug response to different plant species (*Schizaphis graminum*, cereal pest).
Wilson, R.L. Starks, K.J. College Station, Tex., Southwestern Entomological Society. The Southwestern entomologist. Sept 1981. v. 6 (3). p. 229-232. 6 ref. (NAL Call No.: QL461.S65).

0871

The effect of weed and invertebrate pest management on alfalfa establishment in oat stubble.
Bahler, C.C. Byers, R.A.; Stout, W.L. Ankeny, Iowa : Soil Conservation Society of America, c1987. The role of legumes in conservation tillage systems / J.F. Power, editor. Paper presented at the "National Conference on the Role of Legumes in Conservation Tillage Systems", April 27-29, 1987, University of Georgia, Athens, Georgia. p. 76-77. Includes references. (NAL Call No.: DNAL SB203.R6).

0872

Effects of insecticides on invertebrate predators and their cereal aphid (Hemiptera: Aphididae) prey: laboratory experiments (*Metopolophium dirhodum*).
Brown, K.C. EVETB. Lawton, J.H.; Shires, S.W. College Park : Entomological Society of America. Environmental entomology. Dec 1983. v. 12 (6). p. 1747-1750. Includes references. (NAL Call No.: QL461.E532).

0873

Effects of regional releases of resistant wheats on the population dynamics of the cereal leaf beetle (Coleoptera: Chrysomelidae) (*Oulema melanopus*).
Lampert, E.P. AESAA. Haynes, D.L.; Sawyer, A.J.; Jokinen, D.P.; Wellso, S.G. College Park : The Society. Annals of the Entomological Society of America. Nov 1983. v. 76 (6). p. 972-980. Includes references. (NAL Call No.: 420 EN82).

0874

Effects of rye cover crop management on seedcorn maggot (Diptera: Anthomyiidae) populations in soybeans.
EVETEX. Hammond, R.B. College Park, Md. : Entomological Society of America. Environmental entomology. Oct 1984. v. 13 (5). p. 1302-1305. Includes references. (NAL Call No.: DNAL QL461.E532).

0875

An evaluation of greenbug resistance in oats (*Schizaphis graminum*, Texas).
Chedester, L.D. SENTD. Michels, G.J. Jr. College Station : Southwestern Entomological Society. The Southwestern entomologist. Sept 1982. v. 7 (3). p. 166-169. 9 ref. (NAL Call No.: QL461.S65).

0876

Feeding behavior, development, and damage by biotypes B, C, and E of *Schizaphis graminum* (Homoptera: Aphididae) on 'Wintermalt' and 'Post' barley.
EVETEX. Peters, D.C. Kerns, D.; Puterka, G.J.; McNew, R. College Park, Md. : Entomological Society of America. Abstract: Greenbug, *Schizaphis graminum* (Rondani), biotypes B, C, and E, were monitored on susceptible 'Wintermalt' and resistant 'Post' barley to determine if there were behavioral differences among the biotypes when they fed on resistant and susceptible barley. All biotype-cultivar combinations were tested by observing feeding behavior, fecundity, and damage. Greenbug biotype B had shorter nonfeeding times on the host plant before probing than did biotypes C and E. Biotype B appeared to be less successful in feeding from the phloem of susceptible 'Wintermalt' but was relatively more successful than biotypes C and E on the resistant 'Post'. Development, reproduction, and particularly dry weight of biotype B were inferior to biotypes C and E on both resistant and susceptible plants. However, biotype B caused the greatest overall reduction in chlorophyll content of infested leaves and should be considered the most virulent but least biologically fit of the three biotypes in these comparisons. Environmental entomology. June 1988. v. 17 (3). p. 503-507. Includes references. (NAL Call No.: DNAL QL461.E532).

0877

Feeding responses of *Rhopalosiphum padi* (Homoptera: Aphidae) to barley yellow dwarf virus resistant and susceptible barley varieties.
EVETEX. Ullman, D.E. Qualset, C.O.; McLean, D.L. Lanham, Md. : Entomological Society of America. The probing and feeding behavior of an important vector of barley yellow dwarf virus (BYDV), the bird cherry-oat aphid, *Rhopalosiphum padi* was electronically monitored on barley varieties with and without the Yd2 gene, a gene that imparts BYDV resistance to certain cereal cultivars. The presence of the Yd2 gene was shown to have no significant effect on the time elapsed to the first sieve element contact, total number of sieve element contacts, or duration of phloem sieve element ingestion by *R. padi*. Furthermore, aphid probing and feeding was not greatly affected by BYDV infection in susceptible plants. These data demonstrate that the mechanisms underlying BYDV resistance in plants with the Yd2 gene do not involve factors significantly influencing

(PESTS OF PLANTS - INSECTS)

plant-aphid interactions that invoke alterations in aphid probing or feeding behavior. In addition, the number of sieve element contacts and duration of ingestion from sieve element cells by *R. padi* indicate that this aphid should be highly efficient in acquiring and transmitting the phloem-delimited BYDV to 'California Mariout' barley, with or without the Yd2 gene. *Environmental entomology*. Dec 1988. v. 17 (6). p. 988-991. Includes references. (NAL Call No.: DNAL QL461.E532).

0878

Field experiments with oats, 1890 / George E. Morrow, Thomas F. Hunt . Milk and butter tests / G.E. Morrow . Cream raising by dilution / G.E. Morrow, E.H. Farrington . The hessian fly / S.A. Forbes . Canada thistles : their extermination / T.J. Burrill .
Morrow, G. E. 1840-. Hunt, Thomas F. 1862-;
Morrow, G. E. 1840-; Morrow, G. E. 1840-;
Farrington, E. H. 1860-; Forbes, Stephen Alfred, 1844-1930.; Burrill, Thomas Jonathan, 1839-1916. Champaign, Ill. : University of Illinois Agricultural Experiment Station, 23 cm. Caption title. p. 353-388 : ill. ; 23 cm. (NAL Call No.: DNAL 100 I16S no.12).

0879

Grasshopper control in relation to cereal and forage cropby W.R. Walton. --.
Walton, W. R. Washington, D.C. : U.S. Dept. of Agriculture, 1922. 18 p. : ill. --. (NAL Call No.: DNAL Fiche S-70 no.747 1922).

0880

Grasshopper control in relation to cereal and forage crops /by W.R. Walton.
Walton, W. R. 1873-. Washington, D.C. : U.S. Dept. of Agriculture, 1916. Cover title.~
"Contribution from the Bureau of Entomology.". 20 p. : ill. ; 23 cm. Bibliography: p. 19-20. (NAL Call No.: DNAL 1 Ag84F no.747).

0881

Grasshoppers and their control /by J.R. Parker.
Parker, J. R. 1884-. Washington, D.C. : U.S. Dept. of Agriculture, 1939. "Supersedes Farmers' bulletin 1691: How to control grasshoppers in cereal and forage crops.". 38 p. : ill., maps ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1828).

0882

The greenbug (*Destructive pests of small grains, Schizaphis graminum*, Washington).
Pike, K.S. Klostermeyer, E.C.; Retan, A.H.; Peterson, V.; Young, J.O. Pullman, Wash., The Service. Extension Bulletin - Washington State University, Cooperative Extension Service. Mar 1982. Mar 1982. (1003). 2 p. ill. (NAL Call No.: 275.29 W27P).

0883

Greenbug (Homoptera: Aphididae) antibiosis tests in growth chambers: design of experiments and optimum sample size.
EVETEX. Inayatullah, C. Webster, J.A.; Nguyen, H.T. College Park, Md. : Entomological Society of America. *Environmental entomology*. June 1986. v. 15 (3). p. 581-584. Includes references. (NAL Call No.: DNAL QL461.E532).

0884

Greenbug-resistance studies with small grains /by H.L. Chada ... et al..
Chada, Harvey L. College Station, Tex. : Texas Agricultural Experiment Station, 1961. 18 p. : ill. ; 28 cm. Bibliography: p. 18. (NAL Call No.: DNAL 100 T31S (1) no.982).

0885

Helping bugs control bugs (*Oulema melanopus*, the cereal leaf beetle, biological control).
Effron, D. APHIS. Bronx, N.Y., The Garden Society, New York Botanical Garden. Garden (New York). July/Aug 1980. v. 4 (4). p. 12-13, 30-32. ill. (NAL Call No.: SB403.G3).

0886

The Hessian fly in Oregon (*Mayetiola destructor*, cereal pests, control).
Fisher, G. Pike, K.; Rickman, R.; Maxwell, D. Corvallis, Or., The Service. Extension circular - Oregon State University, Extension Service. Sept 1981. Sept 1981. (1093). 4 p. ill., map. (NAL Call No.: 275.29 OR32C).

0887

How to control billbugs destructive to cereal and forage crops /A.F. Satterthwait.
Satterthwait, A. F. 1879-. Washington, D.C. : U.S. Dept. of Agriculture, 1932. Originally issued Jan. 1919. 22 p. : ill. ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1003 1932).

(PESTS OF PLANTS - INSECTS)

0888

How to control billbugs destructive to cereal and forage crops /A.F. Satterthwait.
Satterthwait, A. F. 1879-. Washington, D.C. : U.S. Dept. of Agriculture, 1919. Cover title.~ "Contribution from the Bureau of Entomology.". 23 p. : ill. ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1003).

0889

How to control billbugs destructive to cereal and forage crops by A.F. Satterthwait . --.
Satterthwait, A. F. Washington, D.C. : U.S. Dept. of Agriculture, 1932. 22 p. : ill. --. (NAL Call No.: DNAL Fiche S-70 no.1003 1932).

0890

How to control grasshoppers in cereal and forage crops /by J.R. Parker and W.R. Walton.
Parker, J. R. 1884-. Walton, W. R. 1873-. Washington, D.C. : U.S. Dept. of Agriculture, 1938. Originally issued Apr. 1932. 17 p. : ill., plans ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1691 1938).

0891

How to control grasshoppers in cereal and forage crops /by J.R. Parker and W.R. Walton.
Parker, J. R. 1884-. Walton, W. R. 1873-. Washington, D.C. : U.S. Dept. of Agriculture, 1932. "This bulletin is a revision of and supersedes Farmers' bulletin 747: Grasshopper control in Relation to Cereal and Forage Crops.". 14 p. : ill., plans ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1691).

0892

How to control grasshoppers in cereal and forage crops by J.R. Parker and W. R. Walton, and R.L. Shotwell . --.
Parker, J. R. Washington, D.C. : U.S. Dept. of Agriculture, 1938. 17 p. : ill. --. (NAL Call No.: DNAL Fiche S-70 no.1691 1938).

0893

Influence of rye-cover crop management on soybean foliage arthropods.
EVETEX. Smith, A.W. Hammond, R.B.; Stinner, B.R. College Park, Md. : Entomological Society of America. Environmental entomology. Feb 1988. v. 17 (1). p. 109-114. Includes references. (NAL Call No.: DNAL QL461.E532).

0894

The inheritance and mechanism of resistance of barley to cereal leaf beetle, *Oulema melanopus* L. / by Chung Lee. --.
Lee, Chung, 1943-. 1970. Thesis (Ph.D.)--Michigan State University, 1970. Photocopy. Ann Arbor, Mich. : University Microfilms, 1971. ix, 82 leaves ; 21 cm. Bibliography: leaves 73-76. (NAL Call No.: DISS 70-20,486).

0895

Inheritance of a second source of greenbug resistance in barley.
CRPSAY. Merkle, D.G. Webster, J.A.; Morgan, G.H. Madison, Wis. : Crop Science Society of America. Genetic studies were conducted to determine the inheritance of resistance to biotype E greenbug, *Schizaphis graminum* (Rondani), in PI 426756, a barley (*Hordeum vulgare* L.) line previously identified as resistant to this pest. An additional objective was to determine whether the genes governing resistance in PI 426756 and 'Post' are allelic or nonallelic. Results from greenbug-infested F2 and F3 seedlings in the greenhouse indicated that the resistance in each line is governed by a single dominant gene, and that these two genes are nonallelic and independent. The discovery of another gene for greenbug resistance is important because greenbug resistance in all current U.S. commercial barley cultivars appears to trace back to the same single dominant gene present in Post. The gene symbol Rsg2b was assigned to this gene; whereas, the gene symbol formerly designated Grb was modified to Rsg1a. Crop science. Mar/Apr 1987. v. 27 (2). p. 241-243. Includes references. (NAL Call No.: DNAL 64.8 C883).

0896

Inheritance of resistance in oats to two biotypes of the greenbug (*Schizaphis graminum*).
Boozaya-Angoon, D. Starks, K.J.; Edwards, L.H.; Pass, H. College Park, Md., Entomological Society of America. Environmental entomology. Aug 1981. v. 10 (4). p. 557-559. 6 ref. (NAL Call No.: QL461.E532).

0897

Insect control: small grain.
Kantack, B.H. Kreckhefer, R. Brookings, S.D. : The Service. FS - South Dakota State University, Cooperative Extension Service. Jan 1984. (818). 2 p. (NAL Call No.: DNAL 275.29 S085FS).

(PESTS OF PLANTS - INSECTS)

0898

Insect management.

Chapin, J.W. Sullivan, M.J. Clemson, S.C. : The Service. Circular - Clemson University, Cooperative Extension Service. In series analytic: Small grain production guidelines for South Carolina, 1988-89.~ Includes statistical data. Aug 1988. (463,rev.). p. 21-28. (NAL Call No.: DNAL 275.29 S08E).

0899

Insects on small grains and their control.

Coppock, S. Burton, R.; Pitts, J.T. Stillwater, Okla. : The Service. OSU extension facts - Cooperative Extension Service, Oklahoma State University. June 1984. (7176,rev.). 6 p. ill. (NAL Call No.: DNAL S544.3.0505).

0900

Insects on small grains and their control.

Coppock, S. OK-AR-SD. Burton, R. Stillwater, Okla., The Service. OSU extension facts - Cooperative Extension Service, Oklahoma State University. Oklahoma State University. Cooperative Extension Service. June 1979. June 1979. (7176). 4 p. ill. (NAL Call No.: S544.3.0505).

0901

Insects on small grains and their control (Includes cultural control).

Coppock, S. Burton, R. Stillwater : The Service. OSU extension facts - Cooperative Extension Service, Oklahoma State University. Mar 1983. Mar 1983. (7176 rev.). 4 p. ill. (NAL Call No.: S544.3.0505).

0902

Laboratory evaluation of new insecticides for control of redbanded cutworm larvae (*Euxoa ochrogaster*, pest of cereal and other crops in Canada and the United States).

McDonald, S. College Park, Md., Entomological Society of America. Journal of economic entomology. Oct 1981. v. 74 (5). p. 593-596. 9 ref. (NAL Call No.: 421 J822).

0903

Laboratory insecticide bioassays, *Melanoplus sanguinipes*, Bozeman, Montana, Winters 1976-77 and 1977-78 (Rye).

Mazuranich, P.C. AR-W-AR-W. Onsager, J.A. Reprints. United States. Dept. of Agriculture. Science and Education Administration. Agricultural Research. (NAL Call No.: aS21.A8U5/AR).

0904

Leaf hopper injurious to cereal and forage crops /by Herbert Osborn.

Osborn, Herbert, 1856-. Washington, D.C. : U.S. Dept. of Agriculture, 1932. Caption title.~ "Contribution from Bureau of Entomology.". 34 p. : ill., maps ; 23 cm. Bibliography: p. 33-34. (NAL Call No.: DNAL 1 Ag84C no.241).

0905

Mass rearing the bird cherry oat aphid, *Rhopalosiphum padi* (L.).

PIACA. Kudagamage, C. Foster, J.E. Indianapolis, Ind. : The Academy. Proceedings of the Indiana Academy of Science. Includes abstract. 1985. v. 94. p. 304. (NAL Call No.: DNAL 500 IN2).

0906

A mathematical model for evaluating insect resistance in plants.

JKESA. Webster, J.A. Inayatullah, C.; Fargo, W.S. Lawrence, Kan. : The Society. Journal of the Kansas Entomological Society. Includes abstract. July 1985. v. 58 (3). p. 564. (NAL Call No.: DNAL 420 K13).

0907

Montana small grains integrated crop and pest management 1984-85 field summary.

Bozeman, Mont. : Cooperative Extension Service Montana State University, 1986-. Cover title.~ "February, 1986."~ Caption title: Integrated crop and pest management, 1984-85. v. : map ; 28 cm. (NAL Call No.: DNAL 275.29 M76C no.1332 etc.).

0908

New insect poses threat to western cereal crops.

Washington, D.C. : National Agricultural Aviation Association. Agricultural aviation. May 1988. v. 15 (3). p. 12-14. ill. (NAL Call No.: DNAL S494.5.A3W3).

0909

Oat leaf volatiles: possible insect attractants.

Buttery, R.G. Ling, L.C.; Wellso, S.G. Washington, D.C., American Chemical Society. Journal of agricultural and food chemistry. July/Aug 1982. v. 30 (4). p. 791-792. 10 ref. (NAL Call No.: 381 J8223).

(PESTS OF PLANTS - INSECTS)

0910

Pest control in forages and small grains -- 1982.

Doll, J. D. Wedberg, J. L.; au, C. R.; Kenney, J. F. 1981. This publication gives information on pest control in forages and small grains. Document available from: University of Wisconsin, Agricultural Bulletin Building, 1535 Observatory Drive, Madison, Wisconsin 53706. 41 p. (NAL Call No.: Not available at NAL.). (NAL Call No.: A1981).

0911

Pesticide screening test for the southern chinch bug (*Stenotaphrum secundatum*, grass and cereal pest).

Crocker, R.L. Simpson, C.L. College Park, Md., Entomological Society of America. Journal of economic entomology. Dec 15, 1981. v. 74 (6). p. 730-731. Includes 4 ref. (NAL Call No.: 421 J822).

0912

Pests not known to occur in the United States or of limited distribution. 80. A click beetle.

Whittle, K. Hyattsville, Md. : The Service. APHIS 81 - U.S. Department of Agriculture, Animal and Plant Health Inspection Service. Sept 1987. (50). 5 p. ill., maps. Includes references. (NAL Call No.: DNAL aSB599.A3U5).

0913

Potato insecticides and fungicides in 1902 ; Oat smut and its prevention / Chas. D. Woods . Woods, Chas. D. 1856-1925. Orono : Maine Agricultural Experiment Station, 1902. Cover title. p. 198-212 ; 23 cm. (NAL Call No.: DNAL 100 M28S (1) no.87).

0914

Protecting field and forage crops from green cloverworm.

MUCBA. Ruppel, R.F. Parker, K.A. East Lansing, Mich. : The Service. Extension bulletin E - Cooperative Extension Service, Michigan State University. Apr 1985. (1793). 2 p. ill. (NAL Call No.: DNAL 275.29 M58B).

0915

Protecting field and forage crops from potato leafhopper.

MUCBA. Ruppel, R.F. Parker, K.A. East Lansing, Mich. : The Service. Extension bulletin E - Cooperative Extension Service, Michigan State University. Apr 1985. (1794). 3 p. ill. (NAL Call No.: DNAL 275.29 M58B).

0916

Reduced yield of spring oat cultivars by cereal leaf beetles (*Oulema melanopus*).

Webster, J.A. Smith, D.H.; Hoxie, R.P. Peoria, Ill., The Administration. Abstract: In this research, we determined grain yield loss of oat cultivars commonly grown in cereal leaf beetle infested areas of Michigan. We measured yield loss by comparing grain yield and panicle weights from infested plots with those of uninfested plots of three locations. Two out of three tests showed significant losses in kg/ha grain yield, with an average loss of 17 percent and a range of 4 to 20 percent. The average larval density was one larva per two stems with a range of 0 to 1.2 percent. The average larval density was one larva per two stems with a range of 0 to 1.2 larvae per stem. Of the yield components, no significant losses were noted in fertile tillers (except in one test where red leaf virus was present), but significant losses were found in kernels per panicle and kernel weight. 'Korwood' and 'Orbit' were the highest yielding cultivars, while 'Clintland 64' was the lowest. Agricultural research results; ARR-NC - United States, Dept. of Agriculture, Science and Education Administration. Oct 1980. Oct 1980. (6). 6 p. ill. 10 ref. (NAL Call No.: aS21.A75U75).

0917

Registration of four pairs of greenbug-resistant vs. susceptible near-isolines of winter barley germplasms.

CRPSAY. Carver, B.F. Morgan, G.H.; Edwards, L.H.; Webster, J.A. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1988. v. 28 (6). p. 1034-1035. Includes references. (NAL Call No.: DNAL 64.8 C883).

0918

Registration of two cereal leaf beetle resistant barley germplasms (*Oulema melanopus*).

Smith, D.H. Jr. Webster, J.A.; Grafius, J.E. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1984. v. 24 (4). p. 828. Includes references. (NAL Call No.: 64.8 C883).

0919

Resistance of cereal crops to aphids: role of allelochemicals.

ACSMC. Corcuera, L.J. Argandona, V.H.; Zuniga, G.E. Washington, D.C. : The Society. ACS Symposium series - American Chemical Society. 1987. (330). p. 129-135. Includes references. (NAL Call No.: DNAL QD1.A45).

(PESTS OF PLANTS - INSECTS)

0920

Small grain insect management guidelines number 1 - aphids / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-031).

0921

Small grain insect management guidelines number 2 - armyworms / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"July 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-032).

0922

Small grain insect management guidelines, number 3 -- Cereal leaf beetles / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-033).

0923

Small grain insect management guidelines number 4 - fall armyworms / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-034).

0924

Small grain insect management guidelines, number 5 -- natural enemies / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-035).

0925

Small grain insect management guidelines. 1. Aphids.
McPherson, R.M. Blacksburg, Va. : The Service. Publication - Virginia Cooperative Extension Service. May 1983. (444-031). 2 p. ill. (NAL Call No.: DNAL S544.3.V8V52).

0926

Small grain insect management guidelines. 2. Armyworms.
McPherson, R.M. Blacksburg, Va. : The Service. Publication - Virginia Cooperative Extension Service. July 1983. (444-032). 2 p. ill. (NAL Call No.: DNAL S544.3.V8V52).

0927

Small grain insect management guidelines. 3. Cereal leaf beetles.
McPherson, R.M. Blacksburg, Va. : The Service. Publication - Virginia Cooperative Extension Service. May 1983. (444-033). 2 p. ill. (NAL Call No.: DNAL S544.3.V8V52).

0928

Small grain insect management guidelines. 4. Fall armyworms.
McPherson, R.M. Blacksburg, Va. : The Service. Publication - Virginia Cooperative Extension Service. May 1983. (444-034). 2 p. ill. (NAL Call No.: DNAL S544.3.V8V52).

0929

Small grain insect management guidelines. 5. Natural enemies.
McPherson, R.M. Blacksburg, Va. : The Service. Publication - Virginia Cooperative Extension Service. May 1983. (444-035). 2 p. ill. (NAL Call No.: DNAL S544.3.V8V52).

0930

Small grain insects / Ohio State University, Cooperative Extension Service, Columbus, Ohio. 1980. This publication discusses the cereal leaf beetle, hessian fly, and aphids. A table of several insects and insecticide recommendations is included. Document available from: Ext. Office of Information, Ohio State University, 2120 Fyffe Road, Columbus, OH 43210. 1 sheet : ill. (NAL Call No.: Not available at NAL.). (NAL Call No.: Field Ent Series 15).

(PESTS OF PLANTS - INSECTS)

0931

Sowing date and cereal aphid (*Macrosiphum avenae*, *Rhopalosiphum*, *Schizaphis graminum*, *Metopolophium dirhodum*) infestations and damage to barley in southwestern Quebec.
Ba-Angood, S.A. Stewart, R.K. College Park, Md., Entomological Society of America. Journal of economic entomology. June 1980. v. 73 (3). p. 462-464. ill. 4 ref. (NAL Call No.: 421 J822).

0932

A study of stem inflation in wild buckwheat, *Eriogonum inflatum* (Insect damage, anatomy).
Stone, A.M. Mason, C.T. Jr. Superior, Ariz., University of Arizona. Desert plants. Nov 1979. v. 1 (2). p. 77-81. ill. 9 ref. (NAL Call No.: QK938.D4D4).

0933

Understanding some insect pests of cereal grains...pale western and army cutworm (*Agrotis orthogonia*, *Euxoa auxiliaris*, chemical control, Montana).
Morrill, W.L. Jensen, G. Bozeman : The Station. Capsule information series - Montana Agricultural Experiment Station. June 1982. June 1982. (27). 4 p. ill. Includes references. (NAL Call No.: S83.M6).

0934

Use of aluminum-foil and oat-straw mulches for controlling aster leafhopper, *Macrosteles fascifrons* (Homoptera: Cicadellidae), and aster yellows in carrots.
GRLEA. Setiawan, D.P. Ragsdale, D.W. East Lansing, Mich. : Michigan Entomological Society. The Great Lakes entomologist. Summer 1987. v. 20 (2). p. 103-109. Includes references. (NAL Call No.: DNAL QL461.M5).

0935

Webworms injurious to cereal and forage crops and their control /by Geo. G. Ainslie.
Ainslie, George G. 1886-1930. Washington, D.C. : U.S. Dept. of Agriculture, 1922. Cover title.~ "Contribution from the Bureau of Entomology.". 16 p. : ill., 1 map ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1258).

0936

Webworms injurious to cereal and forage crops and their control by Geo. G. Ainslie . --.
Ainslie, George G. Washington, D.C. : U.S. Dept. of Agriculture, 1922. 16 p. : ill., map --. (NAL Call No.: DNAL Fiche S-70 no.1258).

0937

White grubs in cereal and forage crops and their control / T.R. Chamberlin and C.L. Fluke .
Chamberlin, Thomas R. 1889-. Fluke, C. L. 1891-. Madison : Agricultural Experiment Station, University of Wisconsin, 1947. Cover title. 15 p. : ill. ; 23 cm. (NAL Call No.: DNAL 100 W75 no.159).

0938

Wireworms destructive to cereal and forage crops /J.A. Hyslop.
Hyslop, J. A. 1884-. Washington, D.C. : U.S. Dept. of Agriculture, 1916. Caption title.~ "Contribution from the Bureau of Entomology.". 12 p. : ill. ; 23 cm. Bibliography: p.11-12. (NAL Call No.: DNAL 1 Ag84F no.725).

0939

Wireworms destructive to cereal and forage cropsby J.A. Hyslop. --.
Hyslop, J. A. Washington, D.C. : U.S. Dept. of Agriculture, 1916. 12 p. : ill. --. Bibliography: p. 11-12. (NAL Call No.: DNAL Fiche S-70 no.725).

0940

Yield losses and host selection of cereal leaf beetles (*Oulema melanopus*) in resistant and susceptible spring barley.
Webster, J.A. AR-NC. Smith, D.H. Jr. Madison, Wis., Crop Science Society of America. Crop science. Nov/Dec 1979. v. 19 (6). p. 901-904. ill. 12 ref. (NAL Call No.: 64.8 C883).

0941

Yield losses in spring barley caused by cereal aphids (Homoptera: Aphididae) in South Dakota.
JEENAI. Kieckhefer, R.W. Kantack, B.H. College Park, Md. : Entomological Society of America. Journal of economic entomology. June 1986. v. 79 (3). p. 749-752. Includes references. (NAL Call No.: DNAL 421 J822).

PESTS OF PLANTS - NEMATODES

0942

Anhydrobiosis in *Pratylenchus penetrans* and *Tylenchorhynchus* n. sp. in cultivated soils cropped to winter rye.

JONEB. Townshend, J.L. Raleigh, N.C. : Society of Nematologists. Journal of nematology. Apr 1987. v. 19 (2). p. 164-171. Includes references. (NAL Call No.: DNAL QL391.N4J62).

0943

Biological control of nematodes: status and prospects in agricultural IPM systems.

Van Gundy, S.D. Orlando, Fla. : Academic Press, 1985. Biological control in agricultural IPM systems / edited by Marjorie A. Hoy, Donald C. Herzog. Paper presented at the "Symposium on Biological Control in Agricultural Integrated Pest Management Systems" June 4-6, 1984, held at the Citrus Research and Education Center, University of Florida, at. p. 467-478. Includes references. (NAL Call No.: DNAL SB933.3.B548).

0944

Cereal cyst nematode (*Heterodera avenae*) on oats. II. Early root development and nematode tolerance.

JONEB. Volkmar, K.M. Lake Alfred, Fla. : Society of Nematologists. Journal of nematology. July 1989. v. 21 (3). p. 384-391. Includes references. (NAL Call No.: DNAL QL391.N4J62).

0945

Durable resistance of barley cultivars to the nematode *Heterodera avenae*.

Person-Dedryver, F. NASSD. New York : Plenum Press. NATO advanced study institutes series. Series A. Life sciences. 1983. v. 55. p. 207-210. Includes references. (NAL Call No.: QH301.N32).

0946

***Heterodera avenae*: virulence and resistance.**

NASSD. Andersen, K. Andersen, S. New York, N.Y. : Plenum Press. NATO advanced science institutes series : Series A : Life sciences. In the series analytic: Cyst nematodes / edited by F. Lamberti and C.E. Taylor. 1986. v. 121. p. 277-285. Includes references. (NAL Call No.: DNAL QH301.N32).

0947

Marker genes for cereal cyst nematode resistance.

Ellis, S.E. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 249-250. (NAL Call

No.: DNAL SB191.B2A9 1987).

0948

Nematode populations in a rye/soybean succession after four years of no tillage management.

Post, T.J. Gallaher, R.N.; Dickson, D.W. Gainesville, Fla. : The Station. Agronomy research report AY - Agricultural Experiment Stations, University of Florida. 1983? . (84-6). 5 p. Includes references. (NAL Call No.: DNAL S540.A2F62).

0949

Pests not known to occur in the United States or of limited distribution. 70. A cereal cyst nematode.

Friedman, W. Hyattsville, Md. : The Service. APHIS 81 - U.S. Department of Agriculture, Animal and Plant Health Inspection Service. Sept 1985. (46). p. 115-123. ill., maps. Includes references. (NAL Call No.: DNAL aSB599.A3U5).

0950

Pests not known to occur in the United States or of limited distribution. 71.

Friedman, W. Hyattsville, Md. : The Service. APHIS 81 - U.S. Department of Agriculture, Animal and Plant Health Inspection Service. Sept 1985. (46). p. 125-132. ill., maps. Includes references. (NAL Call No.: DNAL aSB599.A3U5).

PLANT DISEASES - GENERAL

0951

Andre spring barley (Two-rowed, long-auned variety, performance, disease resistance). Morrison, K.J. Ullrich, S.E.; Nilan, R.A.; Brauen, S.E.; Muir, C.E.; Reisenauer, P. Pullman, Wash. : The Service. Extension Bulletin - Washington State University, Cooperative Extension Service. Mar 1984. Mar 1984. (1249). 5 p. (NAL Call No.: 275.29 W27P).

0952

Barley diseases and their control by R.W. Leukel . --. Leukel, R. W. Washington, D.C. : U.S. Dept. of Agriculture, 1955. 28 p. : ill. --. (NAL Call No.: DNAL Fiche S-70 no.2089).

0953

Breeding for pest resistance. AGRYA. Sharp, E.L. Madison, Wis., Soil Science Society of America. Agronomy. A series of monographs - American Society of Agronomy. Literature review. 1985. (26). p. 313-333. Includes references. (NAL Call No.: DNAL 4 AM392).

0954

Coordinator's report: disease and pest resistance genes. Jorgensen, J.H. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 65-69. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0955

Disease-resistant and hardy oats for the South /by T.R. Stanton and F.A. Coffman. Stanton, T. R. 1885-. Coffman, Franklin A._1892-. Washington, D.C. : U.S. Dept. of Agriculture, 1943. 10 p. : ill. ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1947).

0956

Disease-resistant and hardy oats for the South by T.R. Stanton and F.A. Coffman . --. Stanton, T. R. Washington, D.C. : U.S. Dept. of Agriculture, 1943. 10 p. : ill. --. (NAL Call No.: DNAL Fiche S-70 no.1947).

0957

Evaluation of varieties and selections of barley for disease resistance and winter hardiness in southeastern United States /by J.G. Moseman. Moseman, J. G. 1921-. Washington : U.S. Dept. of Agriculture, 1956. 33 p. ; 23 cm. Literature cited: p. 32-33. (NAL Call No.: DNAL 1 Ag84Te no.1152).

0958

Field crops. Arny, D.C. Dubuque, Iowa : Kendall/Hunt Pub. Co., c1986. With one foot in the furrow : a history of the first seventy-five years of the Department of Plant Pathology at the University of Wisconsin-Madison / edited by Paul H. Williams, Melissa Marosy. p. 164-174. ill. (NAL Call No.: DNAL SB732.54.U6W5).

0959

For horses: Kelly. SDFHA. Leslie, J. Brookings, S.D. : The Station. South Dakota farm and home research - South Dakota, Agricultural Experiment Station. 1984. v. 35 (2). p. 14. ill. (NAL Call No.: DNAL 100 S082S).

0960

Grow disease-resistant oats /by T.R. Stanton and F.A. Coffman. Stanton, T. R. 1885-. Coffman, Franklin A._1892-. Washington, D.C. : U.S. Dept. of Agriculture, 1943. 13 p. : ill. ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1941).

0961

Grow disease-resistant oats by T.R. Stanton and F.A. Coffman . --. Stanton, T. R. Washington, D.C. : U.S. Dept. of Agriculture, 1949. 13 p. : ill. --. (NAL Call No.: DNAL Fiche S-70 no.1941 1949).

0962

Inquiry concerning smut of oats /A.D. Shamel. Shamel, A. D. 1878-. Urbana, Ill. : University of Illinois Agricultural Experiment Station, 1900. Caption title. 3 p. : ill. ; 23 cm. (NAL Call No.: DNAL 275.29 IL62C no.25).

(PLANT DISEASES - GENERAL)

0963

Measured crop performance: small grain, 1984.
NUSRA. Bowman, D.T. Raleigh, N.C., The Station.
Research report - Dept. of Crop Science, North
Carolina State University, Agricultural
Experiment Station. Includes statistical data.
July 1984. (96). 26 p. maps. (NAL Call No.:
DNAL 100 N8122).

0964

Measured crop performance: small grain, 1985.
NUSRA. Bowman, D.T. Raleigh, N.C., The Station.
Research report - Dept. of Crop Science, North
Carolina State University, Agricultural
Experiment Station. Includes statistical data.
July 1985. (101). 29 p. maps. (NAL Call No.:
DNAL 100 N8122).

0965

Modeling disease-loss systems in cereals.
Teng, P.S. Minneapolis, Minn. : Published for
the Congress by Burgess Pub., c1981.
Proceedings of symposia : IX International
Congress of Plant Protection, Washington, D.C.,
U.S.A., August 5-11, 1979 / editor, Thor
Kommehahl. p. 122-127. ill. Includes 3 ref.
(NAL Call No.: SB951.15 1979).

0966

**Oat diseases and their control / by M.D. Simons
and H.C. Murphy .**
Simons, M. D. 1925-. Murphy, H. C. 1902-.
Washington, D.C. : Agricultural Research
Service, U.S. Dept. of Agriculture in
cooperation with Iowa Agriculture and Home
Economics Experiment Station, 1968. Cover
title. iv, 15 p. : ill. (some col.) ; 26 cm.
(NAL Call No.: DNAL 1 Ag84Ah no.343).

0967

**Oat diseases and their control by M.D. Simons
and H.C. Murphy . --.**
Simons, M. D. Washington, D.C. : Agricultural
Research Service, U.S. Dept. of Agriculture in
cooperation with Iowa Agriculture and Home
Economics Experiment Station, 1968. iv, 15 p. :
ill. --. (NAL Call No.: DNAL Fiche S-85
no.343).

0968

**On the prevention of oat smut and potato scab
/ R.A. Moore .**
Moore, R. A. 1861-1941. Madison, Wis. :
University of Wisconsin, Agricultural
Experiment Station, 1903. Cover title. 23 p. :
ill. ; 23 cm. (NAL Call No.: DNAL 100 W75 (1)
no.98).

0969

**A quest for cereal varieties resistant to the
minor cereal diseases in eastern Oregon.**
Kolding, M.F. Corvallis, Or., The Station.
Special report - Oregon Agricultural Experiment
Station. June 1981. June 1981. (623). p. 35-42.
(NAL Call No.: 100 OR3M).

0970

**Stripe resistance and yield of smooth-awned
barley hybrids / R.G. Shands ... et al. .**
Shands, R. G. 1903-. Madison, Wis. :
Agricultural Experiment Station of the
University of Wisconsin, 1933. Cover title. 22
p. : ill. ; 23 cm. Bibliography: p. 22. (NAL
Call No.: DNAL 100 W75 (2) no.116).

0971

**Studies on seed treatments for cereal crops
/W.E. Brentzel.**
Brentzel, W. E. 1889-. Fargo : Agricultural
Experiment Station, North Dakota Agricultural
College, 1944. 19 p. : ill., 1 map ; 22 cm.
Includes bibliographical references. (NAL Call
No.: DNAL 100 N813 no.331).

0972

Treatment of oats for smut.
Urbana, Ill. : University of Illinois
Agricultural Experiment Station, 1900. Caption
title. 3 p. : ill. ; 23 cm. (NAL Call No.:
DNAL 275.29 IL62C no.20).

0973

Treatment of oats for smut /by A.D. Smith.
Shamel, A. D. 1878-. Urbana, Ill. : University
of Illinois Agricultural Experiment Station,
1900. Caption title. 2 p. ; 23 cm. (NAL Call
No.: DNAL 275.29 IL62C no.29).

0974

Treatment of oats for smut /by Albert N. Hume.
Hume, Albert N. 1878-. Urbana, Ill. :
University of Illinois Agricultural Experiment
Station, 1908. Cover title.~ "January 1908.". 3
p. ; 23 cm. (NAL Call No.: DNAL 275.29 IL62C
no.89 1908).

PLANT DISEASES - FUNGAL

0975

Aflatoxin decomposition in various soils.
JPFCD2. Angle, J.S. New York, N.Y. : Marcel Dekker. Journal of environmental science and health. Part B. Pesticides, food contaminants, and agricultural wastes. 1986. v. 21 (4). p. 277-288. Includes references. (NAL Call No.: DNAL TD172.J61).

0976

Agronomic characteristics and disease resistance of winter barleys tested in Missouri, 1943 to 1948 /J.M. Poehlman.
Poehlman, John Milton, 1910-. Columbia, Mo. : University of Missouri, College of Agriculture, Agricultural Experiment Station, 1949. 28 p. ; 22 cm. Bibliography: p. 27-28. (NAL Call No.: DNAL 100 M693 (3) no.442).

0977

Allelism of genes in the M1-a locus (Barley, powdery mildew resistance).
Giese, H. Jensen, H.P.; Jorgensen, J.H. Fort Collins, Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1980. v. 10. p. 22-24. 3 ref. (NAL Call No.: QK495.G74B34).

0978

Assessment of barley accessions PI 382313, PI 382474, PI 382915, and PI 382976 for stem rust resistance.
CRPSAY. Jedel, P.E. Metcalfe, D.R.; Martens, J.W. Madison, Wis. : Crop Science Society of America. New sources of resistance to Puccinia graminis Pers. f. sp. tritici Erikss. and Henn. in barley (Hordeum vulgare L.) should be identified and incorporated into breeding programs to ensure the continued resistance of barley to this pathogen. The objective of this study was to identify resistance to stem rust in barley accessions from the USDA Wiebe collection and determine the relationship of this resistance to resistance conferred by the T gene. Four accessions, PI 382313, PI 382474, PI 382915, and PI 382976, were selected as having good field resistance at the Winnipeg Agriculture Canada Research Station, Glenlea rust nursery in 1986. The field resistance of these four accessions was reconfirmed in the 1987 rust nursery and was found to be as effective as, or more effective than, that conferred by T. Seedling and adult plant tests were conducted in controlled environments with six races (C5, C10, C17, C25, C35, and C53). In seedling tests, PI 382313 was resistant or moderately resistant to all six races. Both PI 382474 and PI 382915 were moderately susceptible to C35 and susceptible to the other five races. The PI 382976 was moderately resistant to C35 and C53, and moderately susceptible or susceptible to the other races. In adult plant tests, PI 382474 was resistant to C35 and PI 382313 was resistant to C17. Both

PI 382915 and PI 382979 displayed little resistance as adults to the six races. Genetic studies with PI 382313 indicated that the field resistance of this accession was conferred by a single dominant gene that different from T. Crop science. Nov/Dec 1989. v. 29 (6). p. 1473-1477. Includes references. (NAL Call No.: DNAL 64.8 C883).

0979

Association of a mildew resistance gene JM1h in Hanna 906 with chromosome 6.
Hayashi, J. Heta, H. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Oct 10, 1985. v. 15. p. 46-47. (NAL Call No.: DNAL QK495.G74B34).

0980

Association of host cytoplasm with reaction to Puccinia coronata in progeny of crosses between wild and cultivated oats.
PLDRA. Simons, M.D. Robertson, L.D.; Frey, K.J. St. Paul, Minn. : American Phytopathological Society. Plant disease. Nov 1985. v. 69 (11). p. 969-971. Includes 26 references. (NAL Call No.: DNAL 1.9 P69P).

0981

The attempted transfer of disease resistance from Hordeum bulbosum L., to H. vulgare L.
Pickering, R.A. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 5-8. Includes references. (NAL Call No.: DNAL QK495.G74B34).

0982

Attempts to locate powdery mildew resistance gene M1-(La) to a barley chromosome.
Jensen, H.P. Helms Jorgensen, J.; Jensen, J. Fort Collins, Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1982. v. 12. p. 65-68. 3 ref. (NAL Call No.: QK495.G74B34).

0983

Barley composite cross CC XXXV-A as a source of both specific and slow rusting resistance against leaf rust.
CRPSAY. Andres, M.W. Wilcoxson, R.D. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1986. v. 26 (2). p. 273-275. Includes references. (NAL Call No.: DNAL 64.8 C883).

(PLANT DISEASES - FUNGAL)

0984

Barley smuts.

Johnson, H.G. Bissonnette, H.L. 1981. This publication gives information on the control of Barley Smut. Document available from: Bulletin Room, 3 Coffey Hall, 1420 Eckles Avenue, University of Minnesota, St. Paul, Minnesota 55108. 1 sheet : ill. (NAL Call No.: Not available at NAL.). (NAL Call No.: Plant Pathology Fact Sheet No.6).

0985

Calcium transport in protoplasts isolated from ml-o barley isolines resistant and susceptible to powdery mildew.

PLPHA. Wrona, A.F. Spanswick, R.M.; Aist, J.R. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Dec 1988. v. 88 (4). p. 1157-1162. ill. Includes references. (NAL Call No.: DNAL 450 P692).

0986

Can we lead the pathogen astray? (Breeding cereals for resistance to fungus diseases).

Wolfe, M.S. Barrett, J.A. St. Paul, Minn., American Phytopathological Society. Plant disease. Feb 1980. v. 64 (2). p. 148-151, 153-155. ill. 13 ref. (NAL Call No.: 1.9 P69P).

0987

Carbon-13 and proton nuclear magnetic resonance spectral assignments of deoxynivalenol and other mycotoxins from Fusarium graminearum (Contaminants of cereal crops).

Blackwell, B.A. Greenhalgh, R.; Bain, A.D. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. Sept/Oct 1984. v. 32 (6). p. 1078-1083. ill. Includes 35 references. (NAL Call No.: 381 J8223).

0988

Cell wall appositions and plant disease resistance: acoustic microscopy of papillae that block fungal ingress (Erysiphe graminis hordei, on barley).

Israel, H.W. Wilson, R.G.; Aist, J.R.; Kunoh, H. Washington, D.C., The Academy. Proceedings of the National Academy of Sciences of the United States of America. National Academy of Sciences. Apr 1980. v. 77 (4). p. 2046-2049. ill. 26 ref. (NAL Call No.: 500 N21P).

0989

Cephalosporium stripe disease of cereals.

WUEXA. Murray, T.D. Pullman, Wash. : The Service. Extension bulletin - Washington State University, Cooperative Extension Service. In subseries: Plant Diseases. Apr 1987. (1434). 2 p. ill. (NAL Call No.: DNAL 275.29 W27P).

0990

Cereal grain seed treatments for Montana and their potential economic impact /by D.E.

Mathre, W.G. Heid, Jr., and R.H. Johnston. Mathre, Don E., 1938-. Heid, Walter George, 1932-; Johnston, R. H. Bozeman, Montana : Montana Agricultural Experiment Station, Montana State University, 1973. 17 p. ; 28 cm. (NAL Call No.: DNAL 100 M76 (1) no.669).

0991

The Cereal rusts /edited by William R. Bushnell and Alan P. Roelfs. --.

Bushnell, William R.; Roelfs, A. P. Orlando : Academic Press, c1984-1985. 2 v. : ill. ; 24 cm. Includes bibliographies and indexes. (NAL Call No.: DNAL SB741.R8C44).

0992

Cereal smuts and the disinfection of seed grain /Harry B. Humphrey and Alden A. Potter.

Humphrey, Harry Baker, 1873-. Potter, Alden A. Washington, D.C. : U.S. Dept. of Agriculture, 1918. Cover title.~ "Contribution from the Bureau of Plant Industry.". 28 p. : ill. ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.939).

0993

Cereal smuts and their control by R.W. Leukel and V.F. Tapke. --.

Leukel, R. W. Washington, D.C. : U.S. Dept. of Agriculture, 1954. 28 p. : ill. --. (NAL Call No.: DNAL Fiche S-70 no.2069).

0994

Characteristic cellular responses as expression of genes for resistance to Erysiphe graminis f. sp. hordei in barley (Hordeum vulgare).

Koga, H. PHYTA. Toyoda, H.; Mayama, S.; Shishiyama, J.; Hiura, U. St. Paul : American Phytopathological Society. Phytopathology. June 1983. v. 73 (6). p. 907-910. Includes references. (NAL Call No.: 464.8 P56).

(PLANT DISEASES - FUNGAL)

0995

Characterization in isogenic lines of oat crown rust resistance genes from four sources / by Bhisham Pal Singh. -
Singh, Bhisham Pal, 1941-. 1971. Thesis (Ph.D.)--Iowa State University, 1971. Photocopy of typescript. Ann Arbor: University Microfilms, 1972. iii, 187 leaves ; 21 cm. Bibliography: leaves 160-174. (NAL Call No.: DISS 72-5,257).

0996

Chemical control of barley spot blotch with foliar sprays, 1981 (Cochliobolus sativus, Hordeum vulgare).
Luz, W.C. da.FNETD. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 46. (NAL Call No.: 464.9 AM31R).

0997

Chemical seed treatments for control of barley leaf stripe in California.
PLDRA, Gordon, T.R. Webster, R.K.; Jackson, L.F.; Hall, D.H. St. Paul, Minn. : American Phytopathological Society. Plant disease. June 1985. v. 69 (6). p. 474-477. Includes 10 references. (NAL Call No.: DNAL 1.9 P69P).

0998

Chemical treatment to control barley net blotch, 1982 (Pyrenophora teres, Hordeum vulgare).
Luz, W.C. da.FNETD. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 46. (NAL Call No.: 464.9 AM31R).

0999

Chitinase locus (Chi 1) assigned to barley chromosome 1 (7H) by immunoblotting.
Hejgaard, J. Bjorn, S.E. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Aug 31, 1987. v. 17. p. 92-93. ill. Includes references. (NAL Call No.: DNAL QK495.G74B34).

1000

Coevolution of host and pathogen populations in the Hordeum vulgare-Rhynchosporium secalis pathosystem.
PNASA, McDonald, B.A. McDermott, J.M.; Allard, R.W.; Webster, R.K. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. May 1989. v. 86 (10). p. 3924-3927. Includes references. (NAL Call No.: DNAL 500 N21P).

1001

Common root rot of spring barley: relative susceptibility of six-rowed cultivars.
NDFRA. Stack, R.W. Fargo, N.D. : The Station. North Dakota farm research - North Dakota, Agricultural Experiment Station. July/Aug 1986. v. 44 (1). p. 5-7. ill. Includes references. (NAL Call No.: DNAL 100 N813B).

1002

Comparative resistance of varieties of oats to crown and stem rusts /by L.W. Durrell and John H. Parker.
Durrell, L. W. 1888-. Parker, John H. 1891-. Ames, Iowa : Agricultural Experiment Station, Iowa State College of Agriculture and Mechanic Arts, 1920. p. 28-56, 56a-56d : ill., charts ; 22 cm. Bibliography: 56c-56d. (NAL Call No.: DNAL 100 Io9 no.62).

1003

A comparison of certain combinations of oat varieties as crown rust differentials /by M.D. Simons and H.C. Murphy.
Simons, M. D. 1925-. Murphy, H. C. 1902-. Washington : U.S. Dept. of Agriculture, 1955. Cover title. 22 p. ; 23 cm. Literature cited: p. 22. (NAL Call No.: DNAL 1 Ag84Te no.1112).

1004

Control of barley covered smut, seed decay and seedling blight, 1984.
FNETD. Pederson, V.D. s.l. : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1985. v. 40. p. 168. (NAL Call No.: DNAL 464.9 AM31R).

1005

Control of barley covered smut, seed decay and seedling blight, 1985.
FNETD. Pederson, V.D. s.l. : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1986. v. 41. p. 124. (NAL Call No.: DNAL 464.9 AM31R).

1006

Control of barley covered smut with experimental seed treatment, 1979 (Barley (spring) (Hordeum vulgare 'Purcell'), covered smut; Ustilago hordei).
Johnston, R.H. Mathre, D.E. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 173. (NAL Call No.: 464.9 AM31R).

(PLANT DISEASES - FUNGAL)

1007

Control of barley covered smut with seed treatments, 1980 (Barley (*Hordeum vulgare* 'Purcell'), covered smut; *Ustilago hordei*). Johnston, R.H. Metz, S.G.; Mathre, D.E. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1981. v. 36. p. 152-153. (NAL Call No.: 464.9 AM31R).

1008

Control of barley covered smut, 1978 (Barley *Hordeum vulgare* 'Resibee'), covered smut; *Ustilago horedi*). Alcock, K.T. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 172. (NAL Call No.: 464.9 AM31R).

1009

Control of barley covered smut, 1979 (Barley (*Hordeum vulgare* 'Dickson'), covered smut; *Ustilago hordei*, seed decay and seedling blight; *Fusarium*, *Pythium* and *Helminthosporium* spp.). Pederson, V.D. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 173-174. (NAL Call No.: 464.9 AM31R).

1010

Control of barley covered smut, 1980 (Barley (*Hordeum vulgare* (Dickson') covered smut; *Ustilago hordei*, seed decay and seedling blight; *Fusarium*, *Pythium* and *Helminthosporium* spp.). Pederson, D.V. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1981. v. 36. p. 153-154. (NAL Call No.: 464.9 AM31R).

1011

Control of barley loose smut, seed decay and seedling blight, 1979 (Barley (*Hordeum vulgare* 'Larker'), loose smut; *Ustilago nuda*, seed decay and seedling blight; *Fusarium*, *Pythium* and *Helminthosporium* spp.). Pederson, V.D. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 174-175. (NAL Call No.: 464.9 AM31R).

1012

Control of barley loose smut, seed decay and seedling blight, 1980 (Barley (*Hordeum vulgare* 'Larker'), loose smut; *Ustilago nuda*, seed decay and seedling blight; *Fusarium*, *Pythium* and *Helminthosporium* spp.). Pederson, V.D. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1981. v. 36. p. 154-155. (NAL Call No.: 464.9 AM31R).

1013

Control of barley loose smut, seed decay and seedling blight, 1981 (Barley (*Hordeum vulgare* 'Larker'), loose smut; *Ustilago nuda*, seed decay and seedling blight; *Fusarium*, *Pythium* and *Helminthosporium* spp.). Pederson, V.D. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1982. v. 37. p. 162-163. (NAL Call No.: 464.9 AM31R).

1014

Control of barley loose smut, seed decay and seedling blight, 1982 (*Ustilago nuda*, *Fusarium*, *Pythium* and *Helminthosporium* spp. on *Hordeum vulgare*). Pederson, V.D.FNETD. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 25-26. (NAL Call No.: 464.9 AM31R).

1015

Control of barley loose smut, seed decay and seedling blight, 1984. FNETD. Pederson, V.D. s.l. : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1985. v. 40. p. 168. (NAL Call No.: DNAL 464.9 AM31R).

1016

Control of barley loose smut, seed decay and seedling blight, 1985. FNETD. Pederson, V.D. s.l. : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1986. v. 41. p. 124. (NAL Call No.: DNAL 464.9 AM31R).

1017

Control of barley loose smut with seed treatments, 1978 (Barley (*Hordeum vulgare* 'Horsford'), loose smut; *Ustilago nuda*). Johnston, R.H. Metz, S.G.; Mathre, D.E. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1981. v. 36. p. 153. (NAL Call No.: 464.9 AM31R).

1018

Control of barley loose smut, 1979 (Barley (*Hordeum vulgare* 'Clipper'), loose smut; *Ustilago nuda*).

Barbetti, M.J. Trevenen, S. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1982. v. 37. p. 160-161. (NAL Call No.: 464.9 AM31R).

1019

Control of barley loose smut, 1980 (Barley (*Hordeum vulgare* 'Clipper'), loose smut; *Ustilago nuda*).

Barbetti, M.J. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1982. v. 37. p. 160. (NAL Call No.: 464.9 AM31R).

1020

Control of barley loose smut, 1981 (*Ustilago nuda* on *Hordeum vulgare*).

Barbetti, M.J.FNETD. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 22-23. (NAL Call No.: 464.9 AM31R).

1021

The control of cereal smuts by seed treatment /by F.D. Fromme.

Fromme, F. D. 1886-. Blacksburg, Va. : Virginia Polytechnic Institute. Virginia Agricultural Experiment Station, 1928. 16 p. : ill. ; 23 cm. (NAL Call No.: DNAL 100 V815 (1) no.262).

1022

Control of covered smut, seed decay and seedling blight, 1981 (Barley (*Hordeum vulgare* 'Dickson'), covered smut; *Ustilago hordei*, seed decay and seedling blight; *Fusarium*, *Pythium* and *Helminthosporium* spp.).

Pederson, V.D. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1982. v. 37. p. 162. (NAL Call No.: 464.9 AM31R).

1023

Control of covered smut, seed decay and seedling blight, 1981 (*Ustilago hordei*, *Fusarium*, *Pythium*, *Helminthosporium*, *Hordeum vulgare*).

Pederson, V.D.FNETD. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 25. (NAL Call No.: 464.9 AM31R).

1024

Control of loose and covered smut of barley by seed treatments, 1978 (Barley (*Hordeum vulgare* 'Zephyr' and 'Carlsberg'), loose smut; *Ustilago nuda*, covered smut; *Ustilago hordei*).

Sheridan, J.E. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 175. (NAL Call No.: 464.9 AM31R).

1025

Control of loose and covered smut of oats by seed treatments, 1978 (Oats (*Avena sativa* 'Onward'), loose smut; *Ustilago avenae*, covered smut; *Ustilago hordei*).

Sheridan, J.E. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 183. (NAL Call No.: 464.9 AM31R).

1026

Control of loose and covered smut of oats with seed treatments, 1980 (Oats (*Avena sativa* 'Onward' 'Mapua'), loose smut; *Ustilago avenae*, covered smut; *Ustilago hordei*).

Sheridan, J.E. Cromey, M.G. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1981. v. 36. p. 158. (NAL Call No.: 464.9 AM31R).

1027

Control of loose smut of oats and effect of seed treatments on plant establishment, 1979 (Oats (*Avena sativa* 'Miljee'), loose smut; *Ustilago avenae*).

Ballinger, D.J. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 181. (NAL Call No.: 464.9 AM31R).

1028

Control of loose smut of oats and effect of seed treatments on plant establishment, 1980 (Oats (*Avena sativa* 'Miljee'), loose smut; *Ustilago avenae*).

Ballinger, D.J. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1981. v. 36. p. 157. (NAL Call No.: 464.9 AM31R).

1029

Control of loose smut, seed decay and seedling blight, 1982 (*Ustilago avenae*, *Fusarium*, *Pythium* and *Helminthosporium* spp. on oats, *Avena sativa*).

Pederson, V.D.FNETD. (s.l.) : The Society. Fungicide and nematocide tests : results -

(PLANT DISEASES - FUNGAL)

American Phytopathological Society. 1983. v. 38. p. 29-30. (NAL Call No.: 464.9 AM31R).

1030

Control of net blotch of barley by seed treatments, 1978 (Barley (*Hordeum vulgare* 'Zephyr' and 'Carlsberg'), net blotch; *Drechslera teres*, fungicides).
Sheridan, J.E. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 175-176. (NAL Call No.: 464.9 AM31R).

1031

Control of net blotch of barley by seed treatments, 1979 (Barley (*Hordeum vulgare* 'Zephyr' and 'Carlsberg'), net blotch; *Drechslera teres*, field trials).
Sheridan, J.E. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 176. (NAL Call No.: 464.9 AM31R).

1032

Control of oat loose and covered smut with seed treatments, 1982 (*Ustilago avenae*, *Ustilago hordei*, *Avena sativa*).
Sheridan, J.E. FNETD. Grbavac, N. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 30. (NAL Call No.: 464.9 AM31R).

1033

Control of oat loose and covered smut, 1978 (Oats (*Avena sativa* 'Miljee' and 'Algerian'), loose smut; *Ustilago avenae*, covered smut; *Ustilago hordei*).
Alcock, K.T. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 179-180. (NAL Call No.: 464.9 AM31R).

1034

Control of oat loose smut, seed decay and seedling blight, 1984.
FNETD. Peterson, V.D. s.l. : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1985. v. 40. p. 171. (NAL Call No.: DNAL 464.9 AM31R).

1035

Control of oat loose smut, seed decay and seedling blight, 1985.
FNETD. Pederson, V.D. s.l. : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1986. v. 41. p. 132. (NAL Call No.: DNAL 464.9 AM31R).

1036

Control of oat smut by seed treatment / S.G. Lehman and G.W. Fant .
Lehman, Samuel George, 1887-. Fant, G. W. Raleigh, N.C. : The Agricultural Experiment Station of the North Carolina State College of Agriculture and Engineering, 1929. Cover title. 16 p. : ill. ; 23 cm. (NAL Call No.: DNAL 100 N81 (1) no.268).

1037

Control of oat smut with experimental seed dressings, 1980 (Oats (*Avena sativa* 'Compact'), smut; *Ustilago kolleri*).
Jones, J.P. Dolezal, W.E.; Collins, F.C. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1981. v. 36. p. 157-158. (NAL Call No.: 464.9 AM31R).

1038

Control of oat smut with seed treatment, 1979 (Oats (*Avena sativa* 'Compact'), smut; *Ustilago kolleri*).
Jones, J.P. Collins, F.C. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 181-182. (NAL Call No.: 464.9 AM31R).

1039

Control of oats loose smut, seed decay and seedling blight, 1979 (Oats (*Avena sativa* 'Froker'), loose smut; *Ustilago avenae*, seed decay and seedling blight; *Fusarium*, *Pythium* and *Helminthosporium* spp.).
Pederson, V.D. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 183. (NAL Call No.: 464.9 AM31R).

1040

Control of powdery mildew of barley with seed treatments, 1979 (Barley (*Hordeum vulgare* 'Clipper'), powdery mildew; *Erysiphe graminis hordei*).
Mayfield, A.H. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1981. v. 36. p. 153. (NAL Call No.: 464.9 AM31R).

1041

Control of Pyrenophora leaf stripe of barley with seed treatments, 1982 (Pyrenophora graminea on Hordeum vulgare).
Johnston, R.H.FNETD. Mathre, D.E. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 23-24. (NAL Call No.: 464.9 AM31R).

1042

Correlated inheritance in oats of reaction to diseases and other characters /David Clyde Smith.

Smith, D. C. 1906-. St. Paul : University Farm, 1934 . Cover title.~ Originally presented as: Thesis (Ph.D.)--University of Minnesota, 1934. 38 p. : ill. ; 23 cm. (NAL Call No.: DNAL 100 M66 (3) no.102).

1043

Cultivar differences in response to triadimenol seed treatment for control of barley net blotch caused by Pyrenophora teres.

PLDRA. Sheridan, J.E. Grbavac, N. St. Paul, Minn. : American Phytopathological Society. Plant disease. Jan 1985. v. 69 (1). p. 77-80. ill. Includes 6 references. (NAL Call No.: DNAL 1.9 P69P).

1044

Deposition and biological efficacy of a fungicide applied in charged and uncharged sprays in cereal crops.

Hislop, E.C.CRPTD6. Cooke, B.K.; Harman, J.M.P. Guildford, Eng. : Butterworths. Crop protection. Sept 1983. v. 2 (3). p. 305-316. ill. Includes references. (NAL Call No.: SB599.C8).

1045

Development of a sensitive enzyme-linked immunosorbent assay for the fungicide fenpropimorph.

JAFCAU. Jung, F. Meyer, H.H.D.; Hamm, R.T. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. July/Aug 1989. v. 37 (4). p. 1183-1187. Includes references. (NAL Call No.: DNAL 381 J8223).

1046

Development of crown rust epidemics in genetically diverse oat populations: effect of genotype unit area.

PHYTAJ. Mundt, C.C. Browning, J.A. St. Paul, Minn. : American Phytopathological Society. Phytopathology. May 1985. v. 75 (5). p.

607-610. Includes 31 references. (NAL Call No.: DNAL 464.8 P56).

1047

Disease control.

Drye, C.E. Kingsland, G.C. Clemson, S.C. : The Service. Circular - Clemson University, Cooperative Extension Service. In series analytic: Small grain production guidelines for South Carolina, 1988-89. Aug 1988. (463,rev.). p. 13-20. (NAL Call No.: DNAL 275.29 S08E).

1048

Disease severity and grain yield in barley multilines with resistance to powdery mildew.

CRPSAY. Kolster, P. Munk, L.; Stolen, O. Madison, Wis. : Crop Science Society of America. Superiority of yield in cultivar mixtures may be ascribed to specific and nonspecific resistance against leaf pathogens and to differential reactions of components to environment. This study was conducted to investigate the significance of specific resistance on powdery mildew (*Erysiphe graminis* DC.: Fr. f. sp. *hordei*) severity and grain yield in barley (*Hordeum vulgare* L.) multilines and isolines, each carrying different genes for specific resistance to powdery mildew. Multilines were composed of equal proportions of four or five isolines. In the multilines, disease levels were 50 to 80% lower than the mean level for the components grown in pure stands. The advantage of mixing increased with increasing mean level of resistance in the component isolines. Grain yield in both multilines and isolines increased with the mean level of resistance, with most resistant multilines producing higher grain yields than predicted by the mean level of the components in pure stands. The results indicate that components with higher levels of resistance gained the most benefit from utilizing them in a multiline. Crop science. Nov/Dec 1989. v. 29 (6). p. 1459-1463. Includes references. (NAL Call No.: DNAL 64.8 C883).

1049

Effect of a chemical mutagen on crown rust (*Puccinia coronata*) tolerance of diploid and tetraploid oats.

Simons, M.D. St. Paul, Minn., American Phytopathological Society. Phytopathology. Feb 1981. Abstract only. v. 71 (2). p. 255-256. (NAL Call No.: 464.8 P56).

1050

The effect of defence reactions on the energy balance and yield of resistant plants (Barley, *Erysiphe graminis hordei*).

Smedegaard-Petersen, V. New York, Plenum Press. NATO advanced study institutes series. Series A. Life sciences. 1982. v. 37. p. 299-315. ill.

(PLANT DISEASES - FUNGAL)

Includes 31 ref. (NAL Call No.: QH301.N32).

1051

Effect of energy-requiring defense reactions on yield and grain quality in a powdery mildew (*Erysiphe graminis*)-resistant barley cultivar.
Smedegaard-Petersen, V. Stolen, O. St. Paul, Minn., American Phytopathological Society. *Phytopathology*. Apr 1981. v. 71 (4). p. 396-399. ill. 21 ref. (NAL Call No.: 464.8 P56).

1052

Effect of environment and genotype and their interaction on pathogenicity of *Ustilago hordei* (causing the covered smut of barley). I. Parasite-environment effects.
Emara, Y.A. Freake, G.W. Washington, D.C., American Genetic Association. *The Journal of heredity*. July/Aug 1981. v. 72 (4). p. 261-263. 15 ref. (NAL Call No.: 442.8 AM3).

1053

Effect of environment on fungal leaf blights of small grains.
Shaner, G. Palo Alto, Annual Reviews. Annual review of phytopathology. 1981. Literature review. v. 19. p. 273-296. 108 ref. (NAL Call No.: 464.8 AN72).

1054

Effect of fungicide seed dressings on plant establishment and control of covered smut, 1982 (*Ustilago hordei* on barley, *Hordeum vulgare*).
Bollinger, D.J. FNETD. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 22. (NAL Call No.: 464.9 AM31R).

1055

Effect of fungicide seed treatment on plant establishment and disease control, 1980 (Barley (*Hordeum vulgare* 'Weeah'), covered smut; *Ustilago hordei*).
Ballinger, D.J. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1981. v. 36. p. 152. (NAL Call No.: 464.9 AM31R).

1056

Effect of fungicide seed treatment on plant establishment and disease control, 1982.
FNETD. Ballinger, D.J. s.l. : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1985. v. 40. p. 166. (NAL Call No.: DNAL 464.9 AM31R).

1057

Effect of fungicide seed treatments on control of barley loose smut; 1979 (Barley (*Hordeum vulgare* 'Beecher'), loose smut; *Ustilago nuda*).
Ballinger, D.J. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 172-173. (NAL Call No.: 464.9 AM31R).

1058

Effect of fungicides, applied at seeding on plant establishment and covered smut, 1983.
FNETD. Ballinger, D.J. s.l. : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1985. v. 40. p. 166. (NAL Call No.: DNAL 464.9 AM31R).

1059

Effect of host genotype unit area on epidemic development of crown rust following focal and general inoculations of mixtures of immune and susceptible oat plants.
PHYTA. Mundt, C.C. Leonard, K.J. St. Paul, Minn. : American Phytopathological Society. *Phytopathology*. Oct 1985. v. 75 (10). p. 1141-1145. Includes 26 references. (NAL Call No.: DNAL 464.8 P56).

1060

Effect of inoculations with *Cochliobolus sativus* at specific growth stages on grain yield and quality of malting barley.
CRPSAY. Nutter, F.W. Jr. Pederson, V.D.; Foster, A.E. Madison, Wis. : Crop Science Society of America. *Crop science*. Nov/Dec 1985. v. 25 (6). p. 933-938. Includes references. (NAL Call No.: DNAL 64.8 C883).

1061

Effect of potassium fertilizers on malting barley infected with common root rot.
AGJOAT. Timm, C.A. Goos, R.J.; Johnson, B.E.; Sobolik, F.J.; Stack, R.W. Madison, Wis. : American Society of Agronomy. *Agronomy journal*. Jan/Feb 1986. v. 78 (1). p. 197-200. Includes references. (NAL Call No.: DNAL 4 AM34P).

1062

The effect of powdery mildew on the response of barley to applied nitrogen.
Young, K.J. Khan, T.N. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 177-182. Includes references. (NAL Call No.: DNAL SB191.B2A9 1987).

1063

Effect of Puccinia coronata on straw yield and harvest index of oats.

Simons, M.D. AR-NC. St. Paul, Minn., American Phytopathological Society. Phytopathology. July 1980. v. 70 (7). p. 604-607. ill. 15 ref. (NAL Call No.: 464.8 P56).

1064

Effect of seed treatment fungicides on spot blotch of barley, 1981 (Cochliobolus sativus on Hordeum vulgare).

Luz, W.C. da.FNETD. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 23. (NAL Call No.: 464.9 AM31R).

1065

Effect of seed treatment on oat seedling emergence, 1979 (Oats (Avena sativa 'Bob'), seed decay and seedling blight).

Jones, J.P. Dolexal, W.E.; Collins, F.C. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1980. v. 35. p. 182. (NAL Call No.: 464.9 AM31R).

1066

Effect of seed treatments on barley loose smut, 1981 (Barley (Hordeum vulgare 'Horsford'), loose smut; Ustilago nuda).

Johnston, R.H. Mathre, D.E. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1982. v. 37. p. 161-162. (NAL Call No.: 464.9 AM31R).

1067

Effect of seed treatments on barley loose smut, 1982 (Ustilago nuda, Hordeum vulgare).

Johnston, R.H.FNETD. Mathre, D.E. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 24-25. (NAL Call No.: 464.9 AM31R).

1068

Effect of sexual and asexual reproduction on race abundance in cereal rust fungus populations (Puccinia graminis f. sp. tritici, Puccinia coronata, Puccinia recondita, genetic variation).

Groth, J.V.PHYTA. Roelfs, A.P. St. Paul : American Phytopathological Society. Phytopathology. Nov 1982. v. 72 (11). p. 1503-1507. 20 ref. (NAL Call No.: 464.8 P56).

1059

Effect of susceptibility and adult plant resistance on primary penetration of oats by Erysiphe graminis f. sp. avenae (Abstract only).

Douglas, S.M. Sherwood, R.T.; Lukezic, F.L. St. Paul, Minn., American Phytopathological Society. Phytopathology. Aug 1981. v. 71 (8). p. 871. (NAL Call No.: 464.8 P56).

1070

Effect of volatile compounds, nutrients, and source of sclerotia on eruptive sclerotial germination of Sclerotium rolfsii.

PHYTAJ. Punja, Z.K. Jenkins, S.F.; Grogan, R.G. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Nov 1984. v. 74 (11). p. 1290-1295. ill. Includes 29 references. (NAL Call No.: DNAL 464.8 P56).

1071

Effectiveness of seed treatments in controlling barley covered smut, 1981 (Barley (Hordeum vulgare 'Purcell'), covered smut; Ustilago hordei).

Johnston, R.H. Mathre, D.E. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1982. v. 37. p. 161. (NAL Call No.: 464.9 AM31R).

1072

Effects of chloride fertilizer and systemic fungicide seed treatments on common root rot of barley.

PLDRA. Shefelbine, P.A. Mathre, D.E.; Carlson, G. St. Paul, Minn. : American Phytopathological Society. Plant disease. July 1986. v. 70 (7). p. 639-642. Includes 22 references. (NAL Call No.: DNAL 1.9 P69P).

1073

The effects of heterogeneous oat populations on the epiphytotic development of Victoria blight / by Dafe Kpigere Gboriavhien Ayanru. -

Ayanru, Dafe Kpigere Gboriavhien, 1938-. 1970. Thesis (Ph.D.)--Iowa State University, 1970. Photocopy. Ann Arbor, Mich. : University Microfilms, 1971. iv, 193 leaves : ill. ; 21 cm. Bibliography: leaves 178-192. (NAL Call No.: DISS 71-7,243).

(PLANT DISEASES - FUNGAL)

1074

Effects of planting date on development of net blotch epidemics in winter barley in Pennsylvania.

PLDRA. Delserone, L.M. Cole, H. Jr. St. Paul, Minn. : American Phytopathological Society. Plant disease. May 1987. v. 71 (5). p. 438-441. Includes references. (NAL Call No.: DNAL 1.9 P69P).

1075

The effects of polyamine biosynthesis inhibitors on infection of *Hordeum vulgare* L. by *Erysiphe graminis* f.sp. *hordei* Marchal.

NEPHA. West, H.M. Walters, D.R. New York, N.Y. : Cambridge University Press. The New phytologist. Oct 1988. v. 110 (2). p. 193-200. Includes references. (NAL Call No.: DNAL 450 N42).

1076

Effects of powdery mildew (*Erysiphe graminis* f.sp. *hordei*) and water stress on CO₂ (carbon dioxide) exchange in uninfected leaves of barley.

Williams, G.M. Ayres, P.G. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1981. v. 68 (3). p. 527-530. 25 ref. (NAL Call No.: 450 P692).

1077

Effects of *Pyrenophora teres* and weeds on barley yield and yield components.

PHYTAJ. Burleigh, J.R. Tajani, M.; Seck, M. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Mar 1988. v. 78 (3). p. 295-299. Includes references. (NAL Call No.: DNAL 464.8 P56).

1078

Endophytic fungi in propiconazole-treated and untreated barley leaves.

MYCOAE. Riesen, T.K. Close, R.C. Bronx, N.Y. : The New York Botanical Garden. Mycologia. July/Aug 1987. v. 79 (4). p. 546-552. Includes references. (NAL Call No.: DNAL 450 M99).

1079

Ergots (Fungi, genus *Claviceps*, cereals, feed contamination, mycotoxicity).

Christensen, C.M. Springfield, Va., The Service. PB - U. S. National Technical Information Service. 1980. Literature review. 1980. (80-22-1773). p. 1-44. Bibliography p. 39-44. (NAL Call No.: 157.8 R29).

1080

***Erysiphe graminis* resistance in the *Hordeum murinum* complex (Powdery mildew, wild barley, genetic aspects).**

Giles, B. Barrett, J. Fort Collins : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1983. v. 13. p. 78-82. Includes references. (NAL Call No.: QK495.G74B34).

1081

Evaluation of foliar fungicides on net blotch of barley, 1982 (*Pyrenophora teres*, *Hordeum vulgare*).

Luz, W.C. da.FNETD. (s.l.) : The Society. Fungicide and nematicide tests : results - American Phytopathological Society. 1983. v. 38. p. 46. (NAL Call No.: 464.9 AM31R).

1082

Evaluation of foliar sprays for crown rust control in oats, 1981 (Oats (*Avena sativa* 'Mapua'), crown rust; *Puccinia coronata*).

Sheridan, J.E. (s.l.), The Society. Fungicide and nematicide tests; results - American Phytopathological Society. 1982. v. 37. p. 92. (NAL Call No.: 464.9 AM31R).

1083

Evaluation of foliar sprays for spot blotch control in barley, 1981 (Barley (*Hordeum vulgare* 'Zephyr'), spot blotch; *Drechslera sorokiniana*).

Sheridan, J.E. (s.l.), The Society. Fungicide and nematicide tests; results - American Phytopathological Society. 1982. v. 37. p. 91. (NAL Call No.: 464.9 AM31R).

1084

Evaluation of oat crown rust disease parameters and yield in moderately resistant cultivars (*Puccinia coronata* on *Avena sativa*).

Singleton, L.L. Moore, M.B.; Wilcoxson, R.D.; Kernkamp, M.F. St. Paul, Minn., American Phytopathological Society. Phytopathology. May 1982. v. 72 (5). p. 538-540. Includes 16 ref. (NAL Call No.: 464.8 P56).

1085

Evaluation of Rovral as a seed treatment for control of brown leaf spot in lupins, 1981 (*Pleiochaeta setosa*, *Avena sativa*, oats).

Woodcock, T.FNETD. (s.l.) : The Society. Fungicide and nematicide tests : results - American Phytopathological Society. 1983. v. 38. p. 28-29. (NAL Call No.: 464.9 AM31R).

(PLANT DISEASES - FUNGAL)

1086

Evaluation of seed treatment fungicides for control of net blotch of barley, 1983.
FNETD. Sheridan, J.E. Grbavac, N.; Sheridan, M.H. s.l. : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1985. v. 40. p. 167. (NAL Call No.: DNAL 464.9 AM31R).

1087

Evaluation of seed treatment fungicides for control of net blotch of barley, 1984.
FNETD. Sheridan, J.E. Grbavac, N.; Sheridan, M.H. s.l. : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1985. v. 40. p. 167. (NAL Call No.: DNAL 464.9 AM31R).

1088

Evaluation of seed treatments for barley stripe control, 1980 (Barley (*Hordeum vulgare* 'Summit'), barley stripe; *Pyrenophora graminea*).
Johnston, R.H. Metz, S.G.; Mathre, D.E. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1981. v. 36. p. 152. (NAL Call No.: 464.9 AM31R).

1089

Evaluation of seed treatments for control of barley covered smut, 1982 (*Ustilago hordei*, *Hordeum vulgare*).
Johnston, R.H. FNETD. Mathre, D.E. (s.l.) : The Society. Fungicide and nematocide tests : results - American Phytopathological Society. 1983. v. 38. p. 24. (NAL Call No.: 464.9 AM31R).

1090

Evolutionary response of barley composite cross II to *Rhynchosporium secalis* analyzed by pathogenic complexity and by gene-by-race relationships.
PHYTAJ. Webster, R.K. Saghai-Maroo, M.A.; Allard, R.W. St. Paul, Minn. : American Phytopathological Society. Phytopathology. July 1986. v. 76 (7). p. 661-668. Includes 24 references. (NAL Call No.: DNAL 464.8 P56).

1091

Factors affecting the development of loose smut in barley and its control by dust fungicides /by R.W. Leukel.
Leukel, R. W. 1888-. Washington : U.S. Dept. of Agriculture, 1932. Caption title. 20 p. ; 23 cm. Literature cited: p. 18-19. (NAL Call No.: DNAL 1 Ag84Te no.293).

1092

Factors related to partial resistance of barley to leaf rust.
PHYTAJ. Statler, G.D. Parlevliet, J.E. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Apr 1987. v. 77 (4). p. 549-551. Includes references. (NAL Call No.: DNAL 464.8 P56).

1093

Feasibility of selecting for resistance to kernel discoloration in barley (caused by *Bipolaris sorokiniana*).
Wilcoxson, R.D. Rasmusson, D.C.; Bantari, E.E.; Johnson, D.A. St. Paul, Minn., American Phytopathological Society. Plant disease. Oct 1980. v. 64 (10). p. 928-930. 2 ref. (NAL Call No.: 1.9 P69P).

1094

Ferrous complexes and chelating compounds in suppression of fungal diseases of cereals.
NASSD. Brown, A.E. New York, N.Y. : Plenum Press. NATO advanced science institutes series : Series A : Life sciences. In the series analytic: Iron, siderophores, and plant diseases / edited by T.R. Swinburne. Paper presented at the "NATO Advanced Research Workshop," July 1-5, 1985, Wye, Kent, England.~ Literature review. 1986. v. 117. p. 233-241. Includes references. (NAL Call No.: DNAL OH301.N32).

1095

Field and storage fungi.
Christensen, C.M. New York : Von Nostrand Reinhold, 1987. Food and beverage mycology / edited by Larry R. Beuchat. p. 211-232. ill. Includes references. (NAL Call No.: DNAL QR115.F59 1987).

1096

Field resistance of oats to *Puccinia graminis* f. sp. avenae measured via yield and seed weight reduction.
PLDIDE. Epstein, A.H. Simons, M.D.; Frey, K.J.; Rothman, P.G. St. Paul, Minn. : American Phytopathological Society. Plant disease. Feb 1988. v. 72 (2). p. 154-156. Includes references. (NAL Call No.: DNAL 1.9 P69P).

1097

Field studies on resistance of hybrid selections of oats to covered and loose smuts /by T.R. Stanton, F.A. Coffman, and V.F. Tapke.
Stanton, T. R. 1885-. Coffman, Franklin A. 1892-; Tapke, V. F. 1890-. Washington : U.S. Dept. of Agriculture, 1934. Caption title. 10

(PLANT DISEASES - FUNGAL)

p. ; 23 cm. Literature cited: p. 9. (NAL Call No.: DNAL 1 Ag84Te no.422).

1098

Field studies on resistance of hybrid selections of oats to covered and loose smuts by T.R. Stanton, F.A. Coffman, and V.F. Taper ; the Bureau of Plant Industry in cooperation with the Iowa, North Dakota, Montana, Idaho, and Oregon Agricultural Experiment Stations. Stanton, T. R. (Thomas Ray). Washington, D.C. U.S. Dept. of Agriculture 1934. 10 p. --. Bibliography: p. 9. (NAL Call No.: Fiche S-69 no.422).

1099

Field studies on the rust resistance of oat varieties by M.N. Levine, E.C. Starkman, and T.R. Stanton ; in cooperation with the Minnesota Agricultural Experiment Station. -. Levine, M. N. Washington, D.C. U.S. Dept. of Agriculture 1930. 36 p. : ill., map --. Bibliography: p. 34-35. (NAL Call No.: Fiche S-69 no.143).

1100

Field studies on the rust resistance of oat varieties /by M.N. Levine, E.C. Starkman, and T.R. Stanton. Levine, Mose N. Starkman, E. C.; Stanton, T. R. 1885-. Washington : U.S. Dept. of Agriculture, 1930. Caption title. 36 p. : ill. ; 23 cm. Literature cited: p. 34-35. (NAL Call No.: DNAL 1 Ag84Te no.143).

1101

Formaldehyde seed treatment for oat smuts / by V.F. Tapke . Tapke, V. F. 1890-. Washington, D.C. : U.S. Dept. of Agriculture, 1928. Cover title. 4 p. ; 21 x 10 cm. (NAL Call No.: DNAL 1 Ag84M no.21).

1102

Formalin as a preventive of oat smut / by William Stuart . Stuart, William, 1865-. Lafayette, Ind. : Purdue University Agricultural Experiment Station, 1901. Cover title. p. 38 ; 22 cm. (NAL Call No.: DNAL 100 In2P no.87).

1103

Formation of sodium bisulfite addition products with trichothecenes and alkaline hydrolysis of deoxynivalenol and its sulfonate.

JAFCAU. Young, J.C. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. Sept/Oct 1986. v. 34 (5). p. 919-923. Includes references. (NAL Call No.: DNAL 381 J8223).

1104

Fungicides evaluated for cereal and forage crop disease control.

PLDRA. Watkins, J.E. St. Paul, Minn. : American Phytopathological Society. Plant disease. Oct 1985. v. 69 (10). p. 911-912. (NAL Call No.: DNAL 1.9 P69P).

1105

Further experiments on the control of barley smuts /by R.W. Leukel.

Leukel, R. W. 1888-. Washington : U.S. Dept. of Agriculture, 1936. Caption title. 12 p. ; 23 cm. Literature cited: p. 10-11. (NAL Call No.: DNAL 1 Ag84Te no.513).

1106

Further experiments on the control of barley smuts by R.W. Leukel. -.

Leukel, R. W. Washington, D.C. U.S. Dept. of Agriculture 1936. 12 p. --. Bibliography: p. 10-11. (NAL Call No.: Fiche S-69 no.513).

1107

Genes conditioning resistance of *Hordeum spontaneum* (wild relative of barley) to *Erysiphe graminis* f. sp. *hordei* (barley powdery mildew).

Moseman, J.G. Baenziger, P.S.; Kilpatrick, R.A. Madison, Wis., Crop Science Society of America. Crop science. Mar/Apr 1981. v. 21 (2). p. 229-232. 24 ref. (NAL Call No.: 64.8 C883).

1108

Genetic analysis of changes in scald resistance in barley composite cross V (*Rhynchosporium secalis*, *Hordeum vulgare*).

Jackson, L.F. Webster, R.K.; Allard, R.W.; Kahler, A.L. St. Paul, Minn., American Phytopathological Society. Phytopathology. Aug 1982. v. 72 (8). p. 1069-1072. 16 ref. (NAL Call No.: 464.8 P56).

1109

Genetic dissection of a powdery mildew resistance gene in barley.

Jorgensen, J.H. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Aug 31, 1987. v. 17. p. 99-100. Includes references. (NAL Call No.: DNAL QK495.G74B34).

1110

Genetic factors governing resistance and susceptibility of oats to *Puccinia coronata* Corda var. avenae, F. and L. Race 57 /by Verne C. Finkner.

Finkner, Verne C. Ames, Iowa : Agricultural Experiment Station, Iowa State College, 1954. p. 1040-1063. 1 leaf of plates : ill. (1 col.) ; 22 cm. Bibliography: p. 1063. (NAL Call No.: DNAL 100 Io9 no.411).

1111

Genetic variability for pathogenicity, isozyme, ribosomal DNA and colony color variants in populations of *Rhynchosporium secalis*.

GENTA, McDermott, J.M. McDonald, B.A.; Allard, R.W.; Webster, R.K. Baltimore, Md. : Genetics Society of America. Samples of *Rhynchosporium secalis* were collected from two experimental barley populations known to carry a diverse array of alleles for resistance to this fungal pathogen. Classification of 163 isolates for four putative isozyme systems, a colony color dimorphism and 20 ribosomal DNA restriction fragment length variants revealed 49 different multilocus phenotypes (haplotypes). The six most common haplotypes differed significantly in pathogenicity. Genetic analyses of the data indicated that effective population sizes of the fungus were very large, that the effects of genetic drift were small, and that negligible recombination occurred in the populations studied. Frequency dependent selection was suggested as an explanation for the maintenance of variation in pathogenicity in the fungus. Genetics. July 1989. v. 122 (3). p. 561-565. ill. Includes references. (NAL Call No.: DNAL 442.8 G28).

1112

Geographical distribution and associations between resistance to four races of *Rhynchosporium secalis*.

PHYTAJ. Zhang, Q.F. Webster, R.K.; Allard, R.W. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Feb 1987. v. 77 (2). p. 352-357. Includes references. (NAL Call No.: DNAL 464.8 P56).

1113

Greenhouse experiments on the rust resistance of oat varieties /by John H. Parker.

Parker, John H. 1891-. Washington, D.C. : U.S. Dept. of Agriculture, 1918. Caption title.~ "February 11, 1918."~ "Professional paper.". 16 p., 3 p. of plates : ill. ; 24 cm. Bibliography: p. 16. (NAL Call No.: DNAL 1 Ag84B no.629).

1114

Growth and reproduction of *Drechslera sorokiniana* (leaf and root pathogen of cereals and grasses) as influenced by preemergence herbicides.

Hodges, C.F. Bronx, N.Y., New York Botanical Garden. Mycologia. Mar/Apr 1981. v. 73 (2). p. 244-251. ill. 25 ref. (NAL Call No.: 450 M99).

1115

Half-seed determination of hordeins associated with known M1-a alleles conferring race-specific resistance to barley powdery mildew (*Erysiphe graminis* f. sp. *hordei*).

Hash, C.T. Jr. Blake, T.K. Fort Collins, Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1981. v. 11. p. 74-76. ill. 4 ref. (NAL Call No.: QK495.G74B34).

1116

Histochemical changes in oat cell walls after victorin (Phytotoxic product of *Helminthosporium victoriae*) treatment.

Hanchey, P. St. Paul, Minn., American Phytopathological Society. Phytopathology. May 1980. v. 70 (5). p. 377-381. ill. 15 ref. (NAL Call No.: 464.8 P56).

1117

Histology of the relation between minor and major genes for resistance of barley to leaf rust (*Puccinia hordei* on *Hordeum vulgare*).

Niks, R.E. PHYTA. Kuiper, H.J. St. Paul : American Phytopathological Society. Phytopathology. Jan 1983. v. 73 (1). p. 55-59. 25 ref. (NAL Call No.: 464.8 P56).

1118

History of benzimidazole use and resistance.

Smith, C.M. St. Paul, Minn. : ARS Press, American Phytopathological Society, 1988. Fungicide resistance in North America / Charles J. Delp, editor. p. 22-24. (NAL Call No.: DNAL SB951.F88).

(PLANT DISEASES - FUNGAL)

1119

How to recognize and control ergot (*Claviceps purpurea*) of small grains and grasses.
Hart, L.P. Perry, S.K. East Lansing, Mich., The Service. Extension bulletin E - Michigan State University, Cooperative Extension Service. Aug 1980. Aug 1980. (1430). 2 p. (NAL Call No.: 275.29 M58B).

1120

Hybrid barley research: history and potential.
PMASA. Hockett, E.A. S.l. : The Academy. Proceedings of the Montana Academy of Sciences. Literature review. 1987. v. 47. p. 27-35. Includes references. (NAL Call No.: DNAL 500 M762).

1121

Hydroxypyrimidine fungicides inhibit adenosine deaminase in barley powdery mildew (*Erysiphe graminis*).
Hollomon, D.W. Chamberlain, K. New York, Academic Press. Pesticide biochemistry and physiology. Oct 1981. v. 16 (2). p. 158-169. ill. 27 ref. (NAL Call No.: SB951.P49).

1122

In vitro culture of barley: a method to study *Rhynchosporium secalis* disease and select plants resistant to the toxin, Rhynchosporoside (*Rhynchosporium secalis*, *Hordeum vulgare*).
Branchard, M. New York : Praeger, 1982. Variability in plants regenerated from tissue culture / edited by E.D. Earle, Y. Demarly. p. 343-350. ill. Includes references. (NAL Call No.: QK840.V37).

1123

In vitro selection for resistance.
Foroughi-Wehr, B. Friedt, W.; Schuchmann, R.; Kohler, F.; Wenzel, G. Hingham, Mass. : Martinus Nijhoff Publishers. Advances in agricultural biotechnology. 1986. (20). p. 35-44. ill. Includes references. (NAL Call No.: DNAL S494.5.B563A39).

1124

Induction of resistance to *Erysiphe graminis* f. sp. *hordei* in near-isogenic barley lines.
PHYTA. Cho, B.H. Smedegaard-Petersen, V. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Mar 1986. v. 76 (3). p. 301-305. ill. Includes 16 references. (NAL Call No.: DNAL 464.8 P56).

1125

Influence of genotype and environment on kernel discoloration of Midwestern malting barley.
PLDRA. Miles, M.R. Wilcoxson, R.D.; Rasmusson, D.C.; Wiersma, J.; Warnes, D. St. Paul, Minn. : American Phytopathological Society. Plant disease. June 1987. v. 71 (6). p. 500-504. Includes references. (NAL Call No.: DNAL 1.9 P69P).

1126

Inheritance and yield with particular reference to rust resistance and panicle type in oats /by R.J. Garber.
Garber, Ralph John, 1890-. St. Paul : University Farm, 1922 . Cover title. 62 p., 5 leaves of plates : ill. ; 23 cm. Bibliography: p. 41-43. (NAL Call No.: DNAL 100 M66 (3) no.7).

1127

Inheritance of blast resistance in two-rowed barley.
PLDIDE. Yaegashi, H. St. Paul, Minn. : American Phytopathological Society. Plant disease. July 1988. v. 72 (7). p. 608-610. ill. Includes references. (NAL Call No.: DNAL 1.9 P69P).

1128

Inheritance of resistance to barley leaf stripe.
Skou, J.P. Haahr, V. Fort Collins, Colo. : Department of Agronomy, Colorado State University. Barley genetics newsletter. 1988. v. 18. p. 50-52. Includes references. (NAL Call No.: DNAL QK495.G74B34).

1129

Inheritance of resistance to *Puccinia graminis* f. sp. *secalis* in barley (Breeding for disease resistance).
Steffenson, B.J. Wilcoxson, R.D.; Roelfs, A.P. St. Paul, Minn. : American Phytopathological Society. Plant disease. Sept 1984. v. 68 (9). p. 762-763. Includes 9 references. (NAL Call No.: 1.9 P69P).

1130

Inheritance of resistance to *Pyrenophora graminea* in barley.
PLDIDE. Boulif, M. Wilcoxson, R.D. St. Paul, Minn. : American Phytopathological Society. Plant disease. Mar 1988. v. 72 (3). p. 233-238. Includes references. (NAL Call No.: DNAL 1.9 P69P).

1131

Inheritance of resistance to specific races of crowns and stem rust, to helminthosporium blights, and of certain agronomic characters of oats /by S.C. Litzenberger.

Litzenberger, Samuel C. 1914-. Ames, Iowa : Agricultural Experiment Station, Iowa State College of Agriculture and Mechanic Arts, 1949. p. 454-496 : ill. ; 23 cm. Bibliography: p. 494-496. (NAL Call No.: DNAL 100 Io9 no.370).

1132

The inheritance of smut resistance in crosses of certain varieties of oats.

AGJQAT. Barney, A.F. Madison, Wis. : American Society of Agronomy. The data presented indicate that some varieties of oats contain one factor pair for resistance to loose smut, that other varieties contain two independent factor pairs, while in other varieties three factor pairs may be concerned with resistance. Selection within seventy families from two crosses indicated that resistant families can be isolated in the second generation which breed true to that characteristic thru the F3, F4 and F5 generations. Likewise, highly susceptible families can be isolated which breed true. Such results show the possibility of obtaining desirable types of smut resistant oats from crosses between resistant and susceptible varieties. The accompanying illustrations (Figures 1, 2, 3 and 4) are presented to indicate the response to smut exposure exhibited by two resistant and two susceptible families. Agronomy journal. Apr 1924. v. 16 (4). p. 283-291. (NAL Call No.: DNAL 4 AM34P).

1133

Initiation of induced nonhost resistance of oat leaves to rust infection (*Puccinia* sp.).

Tani, T. Yamashita, Y.; Yamamoto, H. St. Paul, Minn., American Phytopathological Society. Phytopathology. Jan 1980. v. 70 (1). p. 39-42. ill. 13 ref. (NAL Call No.: 464.8 P56).

1134

Localization of induced genes for powdery mildew resistance.

Robbelen, G. Heun, M. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Oct 10, 1985. v. 15. p. 4-7. Includes references. (NAL Call No.: DNAL QK495.G74B34).

1135

Measuring disease progress in pure and mixed stands of plant cultivars.

PHYTA. Gumpert, F.M. St. Paul, Minn. : American Phytopathological Society. A measure is presented that quantifies disease progress of fungal pathotypes during the exponential phase of the epidemic. This measure, referred to as rate of disease increase (p), represents the factor by which the number of infection units is multiplied from one day to another after the initial infection waves have damped. The rate of disease increase depends on four pathotype/cultivar-specific parameters (latent period, infectious period, infection efficiency, spore production rate) and two nonspecific parameters (deposition frequency and autodeposition frequency). Three applications are given. First, it is used to determine the levels of partial resistance of barley cultivars to leaf rust. The calculated p values were highly correlated both with total spore production per unit leaf area ($r = 0.97$) and with disease score in the field ($r = 0.81$). The rate of disease increase is particularly useful if the epidemiological parameters display compensatory effects. The second application deals with the long-term composition of pathotype mixtures. The pathotype with the largest p value will predominate in the long run. This is again exemplified by using barley leaf rust data. The third application concerns disease control strategies. A host stand should be composed so that the corresponding predominant pathotype has a p value smaller than the prevailing pathotype of any other composition. This concept does not presuppose selection against unnecessary genes for virulence. A condition is given under which a cultivar mixture may be more beneficial than each of its components grown in pure stands, and this condition is illustrated by a simple example of two pathotypes and two cultivars. Phytopathology. Sept 1989. v. 79 (9). p. 969-973. Includes references. (NAL Call No.: DNAL 464.8 P56).

1136

Method for screening bacteria and application thereof for field control of *Pythium* spp. on small grain crops.

Weller, D.M. Cook, R.J.; Becker, J.D. Washington, D.C.? : The Department. Abstract: A method for screening bacteria to select strains which will suppress *Pythium* spp. in small grain crops under field conditions and a method for applying field-suppressive bacteria to suppress *Pythium* spp. in a commercial setting are described. Four *Pseudomonas* strains which passed the screen test are disclosed. United States Department of Agriculture patents. Copies of USDA patents are available for a fee from the Commissioner of Patents and Trademarks, U.S. Patents and Trademarks Office, Washington, D.C. 20231. Mar 3, 1987. (4,647,533). 1 p. ill. Includes references. (NAL Call No.: DNAL aT223.V4A4).

(PLANT DISEASES - FUNGAL)

1137

Methods in breeding cereals for rust resistance.

AGJOAT. Johnson, E.C. Madison, Wis. : American Society of Agronomy. Agronomy journal. 1910. v. 2. p. 76-80. (NAL Call No.: DNAL 4 AM34P).

1138

Methods of eradicating buckthorn (*rhamnus*) susceptible to crown rust (*puccinia coronata*) of oats /by S.M. Dietz and L.D. Leach.

Dietz, S. M. 1893-. Leach, L. D. 1900-. Washington, D.C. : U.S. Dept. of Agriculture. 1930. Caption title.~ "Contribution from Bureau of Plant Industry.". 16 p. : ill., charts, maps ; 24 cm. Bibliography: p. 15. (NAL Call No.: DNAL 1 Ag84C no.133).

1139

Methods of seeding oats drilling and broadcasting /by Albert N. Hume, O.D. Center and Leonard Hegnauer.

Hume, Albert N. 1878-. Center, O. D. 1872-; Hegnauer, Leonard, 1872-. Urbana, Ill. : University of Illinois Agricultural Experiment Station, 1909. Cover title. p. 297 -312 : ill. ; 23 cm. (NAL Call No.: DNAL 100 I16S no.136).

1140

Microflora (Storage fungi, cereal grains and their products).

Christensen, C.M. Sauer, D.B. St. Paul, Minn. : American Association of Cereal Chemists, 1982. Storage of cereal grains and their products. 3rd ed. / edited by Clyde M. Christensen. p. 219-240. 1 p. ref. (NAL Call No.: SB190.C43 1982).

1141

Models explaining the specificity and durability of host resistance derived from the observations on the barley-*Puccinia hordei* system.

Parlevliet, J.E.NASSD. New York : Plenum Press. NATO advanced study institutes series. Series A. Life sciences. 1983. v. 55. p. 57-80. Includes references. (NAL Call No.: QH301.N32).

1142

Monitoring stem rust: a key to control (Cereal Rust Laboratory).

Pierce, R. Washington, D.C., The Administration. Agricultural research - United States Dept. of Agriculture, Science and Education Administration. Mar 1981. v. 29 (9). p. 9. ill. (NAL Call No.: 1.98 AG84).

1143

Morphologic and physiologic studies on stem-rust resistance in cereals /by Helen Hart. Hart, Helen, 1900-. Washington : U.S. Dept. of Agriculture, 1931. Cover title. 76 p. : ill. ; 23 cm. Literature cited: p. 72-75. (NAL Call No.: DNAL 1 Ag84Te no.266).

1144

Multiple inoculation technique for evaluating resistance of single barley seedlings to three fungi (*Rhynchosporium secalis*, *Puccinia hordei*, *Erysiphe graminis hordei*, breeding for disease resistance).

Kilpatrick, R.A. Baenziger, P.S.; Moseman, J.G. St. Paul, Minn., American Phytopathological Society. Plant disease. June 1981. v. 65 (6). p. 504-506. 8 ref. (NAL Call No.: 1.9 P69P).

1145

Near-isogenic barley lines with genes for resistance to powdery mildew.

CRPSAY. Kolster, P. Munk, L.; Stolen, O.; Lohde, J. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 903-907. Includes references. (NAL Call No.: DNAL 64.8 C883).

1146

New sources of resistance to *Puccinia hordei* in barley land race cultivars.

PHYTAJ. Yahyaoui, A.H. Sharp, E.L.; Reinhold, M. St. Paul, Minn. : American Phytopathological Society. Phytopathology. July 1988. v. 78 (7). p. 905-908. Includes references. (NAL Call No.: DNAL 464.8 P56).

1147

Partial characterization of fusicoccin (a phytotoxin isolated from the fungus *Fusicoccum amygdali*) binding to receptor sites on oat root membranes.

Stout, R.G. Cleland, R.E. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1980. v. 66 (3). p. 353-359. ill. 27 ref. (NAL Call No.: 450 P692).

1148

Pathogen fitness in cereal mildews (Plant breeding, disease resistance).

Wolfe, M.S.NASSD. Barrett, J.A.; Slater, S.E. New York : Plenum Press. NATO advanced study institutes series. Series A. Life sciences. 1983. v. 55. p. 81-100. Includes references. (NAL Call No.: QH301.N32).

1149

Pathogen population control in powdery mildew of barley (*Erysiphe graminis*, UK).
 Wolfe, M.S. Minneapolis, Minn. : Published for the Congress by Burgess Pub., c1981.
 Proceedings of symposia : IX International Congress of Plant Protection, Washington, D.C., U.S.A., August 5-11, 1979 / editor, Thor Kommedahl. p. 145-148. Includes 17 ref. (NAL Call No.: SB951.I5 1979).

1150

Pathogenesis and host-parasite interaction in cereal powdery mildew.
 Carver, T.L.W. New York : Gordon and Breach Science Pub., c1988. Experimental and conceptual plant pathology / edited by R.S. Singh ... et al. . Literature review. p. 351-381. ill. Includes references. (NAL Call No.: DNAL SB731.E97).

1151

Pathogenic variation in field populations of barley mildew.
 Chan, K. Boyd, W.J.R. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 242-245. (NAL Call No.: DNAL SB191.B2A9 1987).

1152

Performance of small grain varieties in Louisiana, 1987-88.
 Harrison, S.A. Boquet, D.J.; Colyer, P.D.; Groth, D.E.; Habetz, R.J.; Hallmark, W.B.; Hutchinson, R.L.; Moore, S.H.; Rabb, J.L. Baton Rouge, La. : The Department. Report of projects - Louisiana Agricultural Experiment Station, Department of Agronomy. 1988. p. 80-83. (NAL Call No.: DNAL 100 L936).

1153

Physiologic specialization in the oat smut fungi and its relation to breeding oats for smut resistance /by C.S. Holton and H.A. Rodenhiser.
 Holton, Charles Stewart, 1904-. Rodenhiser, H. A. 1899-. Washington : Dept. of Agriculture, 1948. Caption title. 16 p. ; 23 cm. Literature cited: p. 15-16. (NAL Call No.: DNAL 1 Ag84Te no.952).

1154

Physiologic specialization in the oat smut fungi and its relation to breeding oats for smut resistance by C.S. Holton and H.A. Rodenhiser. -.
 Holton, C. S. (Charles Stewart), 1904-. Washington, D.C. U.S. Dept. of Agriculture 1948. 16 p. --. Bibliography: p. 15-16. (NAL Call No.: Fiche S-69 no.952).

1155

Population dynamics of DMI fungicide sensitivity changes in barley powdery mildew.
 Heaney, S.P. St. Paul, Minn. : ARS Press, American Phytopathological Society, 1988. Fungicide resistance in North America / Charles J. Delp, editor. p. 89-92. (NAL Call No.: DNAL SB951.F88).

1156

A possible biological control agent (*Bacillus licheniformis*) for net blotch (*Pyrenophora teres*) of barley (Abstract only).
 Scharen, A.L. Bryan, M.D. St. Paul, Minn., American Phytopathological Society. Phytopathology. Aug 1981. v. 71 (8). p. 902-903. (NAL Call No.: 464.8 P56).

1157

Potato insecticides and fungicides in 1902 ; Oat smut and its prevention / Chas. D. Woods .
 Woods, Chas. D. 1856-1925. Orono : Maine Agricultural Experiment Station, 1902. Cover title. p. 198-212 ; 23 cm. (NAL Call No.: DNAL 100 M28S (1) no.87).

1158

The potato scab /by Roland Thaxter. The proteids or albuminoids of the oat-kernel /by Thomas B. Osborne.
 Thaxter, Roland, 1858-1932. Osborne, Thomas B. 1859-1929. New Haven : Connecticut Agricultural Experiment Station, 1890. Cover title. 8 p. ; 24 cm. (NAL Call No.: DNAL 100 C76St (1) no.105).

1159

Powdery mildew resistance gene M1-a8 (Reg1h8) in northwest European spring barley varieties (*Erysiphe graminis hordei*, *Hordeum*).
 Jorgensen, J.H. Jensen, H.P. Fort Collins : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1983. v. 13. p. 51-53. Includes references. (NAL Call No.: QK495.G74B34).

(PLANT DISEASES - FUNGAL)

1160

A preliminary bulletin on the prevention of smut on oats /E.F. Pernot.

Pernot, E. F. 1859-. Corvallis, Ore. : Oregon Agricultural Experiment Station, 1900. Cover title. 9 p. ; 22 cm. (NAL Call No.: DNAL 100 Or3 no.63).

1161

A preliminary note on the inheritance of rust resistance in oats.

AGJOAT. Garber, R.J. Madison, Wis. : American Society of Agronomy. Agronomy journal. Jan 1921. v. 13 (1). p. 41-43. ill. Includes references. (NAL Call No.: DNAL 4 AM34P).

1162

A preliminary study of the inheritance of rust resistance in oats.

AGJOAT. Parker, J.H. Madison, Wis. : American Society of Agronomy. Pedigree lines of two oat varieties, Burt and Sixty-Day, together with a large number of F2 generation hybrids between these varieties, were studied in relation to their rust resistance. Most of the inoculations were made on seedlings, but enough were made on plants at time of heading to show that the results were similar. The rusts used were the crown rust of oats, *Puccinia lolii avenae* McAlpine, and the stem rust of oats, *Puccinia graminis avenae* Erikss, and Henn. Burt and Sixty-Day and all the hybrids of these two varieties so far tested were found to be entirely susceptible to stem rust. All plants to Sixty-Day also were uniformly susceptible to crown rust. Of 223 inoculated plants of Burt, 48 were classified as resistant, 152 as intermediate, and 23 as susceptible. Each of the five hybrid families contained, in the F2 generation, some plants showing a high degree of resistance to crown rust and others which were as susceptible as plants of the Sixty-Day parent. In other words, there was definite segregation. There was, however a rather large number of plants which were classified as intermediate and which showed varying degrees of resistance. The numerical results of inoculations made in the F2 hybrids were as follows: seedlings classified as resistant no. 81 (17.3%); seedlings classified as intermediate no. 61 (13.0%); seedlings classified as susceptible no. 326 (69.7%); total no. 468 (100.0%). The fact that there were so many more susceptible than resistant plants indicates that susceptibility to crown rust in this cross is partially dominant, while resistance is recessive. These contrasted characters are not thought to be due to environmental conditions or to differences in the metabolism of the host plants, but to definite genetic factors. Nonhereditary factors may of course influence or modify their expression. Rust resistance and susceptibility hardly can be considered as simple characters or as being determined by a single factor difference. The F2 gene. Agronomy journal. Jan 1920. v. 12 (1). p. 23-38. plates. Includes

references. (NAL Call No.: DNAL 4 AM34P).

1163

Production of fungal inoculum using a substrate of perlite, cornmeal, and potato-dextrose agar (*Bipolaris sorokiniana*, *Drechslera oryzae*, *Alternaria alternata*, barley, production of contamination free inoculum).

Miles, M.R. Wilcoxson, R.D. St. Paul, Minn. : American Phytopathological Society. Plant disease. Apr 1984. v. 68 (4). p. 310. Includes references. (NAL Call No.: 1.9 P69P).

1164

Rapid removing of smut from seed oats / by J.C. Arthur .

Arthur, J. C. 1850-1942. Lafayette, Ind. : Purdue University Agricultural Experiment Station, 1905. Cover title. p. 255 -264 ; 22 cm. (NAL Call No.: DNAL 100 In2P no.103).

1165

A rapid simple liquid pickle applicator for small seed lots.

Ellis, S.E. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 262-263. (NAL Call No.: DNAL SB191.B2A9 1987).

1166

Rate-limiting resistance to *Pyrenophora* leaf blotch in spring oats.

PHYTAJ. Frank, J.A. Christ, B.J. St. Paul, Minn. : American Phytopathological Society. Phytopathology. July 1988. v. 78 (7). p. 957-960. Includes references. (NAL Call No.: DNAL 464.8 P56).

1167

Reaction of barley varieties to *helminthosporium sativum* /by H.K. Hayes ... et al. .

Hayes, H. K. 1884-. St. Paul : University Farm, 1923. Cover title. 47 p., 3 p. of plates : ill. ; 23 cm. Bibliography: p. 47. (NAL Call No.: DNAL 100 M66 (3) no.21).

1168

Reaction of *Hordeum bulbosum* L. to Japanese races of powdery mildew (*Erysiphe graminis* f. sp. *hordei*) (that attack *Hordeum vulgare*, barley, genetic transfer of resistance).

Prasad, G. Yasuda, S.; Konishi, T. Fort Collins : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1983. v.

13. p. 60-62. Includes references. (NAL Call No.: QK495.G74B34).

1169

Receptivity, incubation period, and lesion size as criteria for screening barley genotypes for resistance to pyrenophora teres.

PHYTAJ. Nutter, F.W. Jr. Pederson, V.D. St. Paul, Minn. : American Phytopathological Society. Phytopathology. May 1985. v. 75 (5). p. 603-606. Includes 12 references. (NAL Call No.: DNAL 464.8 P56).

1170

Recognizing and controlling Cephalosporium stripe, a disease of cereal grains.

Karow, R.S. Powelson, R.L.; Koepsell, P.A. Corvallis, Or. : The Service. FS, fact sheet - Oregon State University Extension Service. Feb 1985. (308). 2 p. (NAL Call No.: DNAL 275.29 OR36).

1171

Registration of 'Florida 401' rye.

CRPSAY. Pfahler, P.L. Barnett, R.D.; Luke, H.H. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 836. Includes references. (NAL Call No.: DNAL 64.8 C883).

1172

Registration of four stem rust and crown rust resistant oat germplasm lines.

CRPSAY. Rothman, P.G. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1984. v. 24 (6). p. 1217-1218. Includes 6 references. (NAL Call No.: DNAL 64.8 C883).

1173

Registration of 'Heartland' barley.

CRPSAY. Therrien, M.C. Irvine, R.B.; Campbell, K.W.; Wolfe, R.I. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1985. v. 25 (6). p. 1124. (NAL Call No.: DNAL 64.8 C883).

1174

Registration of 'Multiline E76' and 'Multiline E77' oats.

CRPSAY. Frey, K.J. Browning, J.A.; Simons, M.D. Madison, Wis. : Crop Science Society of America. Crop science. Nov/Dec 1985. v. 25 (6). p. 1125. Includes 1 references. (NAL Call No.: DNAL 64.8 C883).

1175

Registration of powdery mildew resistant barley composite cross XLII germplasm (Hordeum vulgare, Erysiphe graminis).

Bockelman, H.E. CRPSAY. Sharp, E.L.; Eslick, R.F.; Ramage, R.T. Madison : Crop Science Society of America. Crop science. Nov/Dec 1983. v. 23 (6). p. 1225. Includes references. (NAL Call No.: 64.8 C883).

1176

Registration of 'Webster' oat.

CRPSAY. Frey, K.J. Simons, M.D.; Michel, L.J.; Murphy, J.P.; Browning, J.A. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 374-375. Includes references. (NAL Call No.: DNAL 64.8 C883).

1177

Registration of Webster oat isolines as parental lines.

CRPSAY. Frey, K.J. Simons, M.D.; Michel, L.J.; Murphy, J.P.; Browning, J.A. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1988. v. 28 (2). p. 386-387. Includes references. (NAL Call No.: DNAL 64.8 C883).

1178

Resistance experiences in Europe.

Brent, K.J. St. Paul, Minn. : ARS Press, American Phytopathological Society, 1988. Fungicide resistance in North America / Charles J. Delp, editor. p. 19-21. (NAL Call No.: DNAL SB951.F88).

1179

Resistance of barley to Puccinia graminis f. sp. tritici and Puccinia graminis f. sp. secalis.

PHYTA. Steffenson, B.J. Wilcoxson, R.D.; Roelfs, A.P. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Oct 1985. v. 75 (10). p. 1108-1111. Includes 16 references. (NAL Call No.: DNAL 464.8 P56).

1180

Resistance of Hordeum spontaneum collected in Israel to infection with Erysiphe graminis hordei (Barley, powdery mildew, germplasm, disease).

Moseman, J.G. CRPSAY. Nevo, E.; Zohary, D. Madison : Crop Science Society of America. Crop science. Nov/Dec 1983. v. 23 (6). p. 1115-1119. Includes references. (NAL Call No.: 64.8 C883).

(PLANT DISEASES - FUNGAL)

1181

Resistance of wild barley accessions from Israel to leaf rust collected in the USA and Israel.

CRPSAY. Manisterski, J. Treeful, L.; Tomerlin, J.R.; Anikstest, Y.; Moseman, J.G.; Wahl, I.; Wilcoxson, R.D. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 727-730. Includes 22 references. (NAL Call No.: DNAL 64.8 C883).

1182

Resistance of winter barley to *Ustilago nuda* (Jens.) Rostr. /C.K. Cloninger and J.M. Poehlman.

Cloninger, C. K. Poehlman, John Milton, 1910-. Columbia, Mo. : University of Missouri, College of Agriculture, Agricultural Experiment Station, 1954. 35 p. : ill. ; 23 cm. Bibliography: p. 33-35. (NAL Call No.: DNAL 100 M693 (3) no.560).

1183

Resistance to covered smut in varieties and hybrids of oats.

AGJDAT. Gaines, E.F. Madison, Wis. : American Society of Agronomy. Of 210 varieties and selections of oats tested for resistance to covered smut, *Ustilago levis* (K. & S.) Magn., during the past eight years, 21 were smut free, the others ranging from slight infection to extreme susceptibility in which nine-tenths of the panicles were black masses of smut spores. Markton, one of the immune selections, is a thin hulled, white oat of sativa type, and is one of the best yielding varieties tested in recent years. It has been increased and grown commercially since 1924. The inheritance of the immunity of Red Rustproof has been studied in four crosses with susceptible varieties. The crosses with Black Tartarian and Abundance indicate that Red Rustproof carries three dominant factors for immunity, any one of which prevents the production of covered smut spores. In crosses with Large and Chinese Hulless, one factor apparently does not give complete dominance in hulless segregates, but otherwise the prepotency of the factors for immunity is similar in all four crosses. Of 56 F3 plants selected from smut-free rows, 45 produced only smut-free plants in the F4 generation. From the results already obtained in varietal testing and breeding for disease resistance in field crops, it is to be hoped that the major diseases will in time be successfully controlled by the development and introduction of resistant or immune varieties. Agronomy journal. Dec 1925. v. 17 (12). p. 775-789. (NAL Call No.: DNAL 4 AM34P).

1184

Resistance to disease and maximum potential yield.

Vanderplank, J.E. New York : Gordon and Breach Science Pub., c1988. Experimental and conceptual plant pathology / edited by R.S. Singh ... et al. . p. 593-599. Includes references. (NAL Call No.: DNAL SB731.E97).

1185

Response of plant pathogens to fungicides.

Wolfe, M.S. Barrett, J.A. Washington, D.C. : National Academy Press, 1986. Pesticide resistance : strategies and tactics for management / Committee on Strategies for the Management of Pesticide Resistant Pest Populations, Board on Agriculture, National Research Council. Paper presented at the "Symposium on Pesticide Resistance," November 27-29, 1984, Washington, D.C. p. 245-256. Includes references. (NAL Call No.: DNAL SB957.P48).

1186

Responses of two-, three-, and four-component barley mixtures to a variable pathogen population.

CRPSAY. McDonald, B.A. Allard, R.W.; Webster, R.K. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1988. v. 28 (3). p. 447-452. Includes references. (NAL Call No.: DNAL 64.8 C883).

1187

Rhizoctonia root rot of small grains favored by reduced tillage in the Pacific Northwest.

PLDRA. Weller, D.M. Cook, R.J.; MacNish, G.; Bassett, E.N.; Powelson, R.L.; Petersen, R.R. St. Paul, Minn. : American Phytopathological Society. Plant disease. Jan 1986. v. 70 (1). p. 70-73. ill. Includes 35 references. (NAL Call No.: DNAL 1.9 P69P).

1188

Seed treatment control of oat smut, and downy mildew, 1981 (Oats (*Avena sativa* 'Compact'), smut; *Ustilago kollerii*, downy mildew; *Sclerophthora macrospora*).

Jones, J.P. Bassi, A.; Collins, F.C. (s.l.), The Society. Fungicide and nematocide tests; results - American Phytopathological Society. 1982. v. 37. p. 166. (NAL Call No.: 464.9 AM31R).

1189

Seed treatment for control of Pyrenophora leaf stripe of barley.

PLDRA. Johnston, R.H. Metz, S.G.; Riesselman, J.H. St. Paul, Minn. : American Phytopathological Society. Plant disease. Dec 1982. v. 66 (12). p. 1122-1124. Includes references. (NAL Call No.: DNAL 1.9 P69P).

1190

Seed treatment for control of seed and soil-borne pathogens of barley, 1981 (Barley (*Hordeum vulgare* 'Maury'), seed decay, seedling blight; *Pythium* sp., *Rhizoctonia* sp., *Fusarium* sp., spot blotch; *Bipolaris sorokiniana*, net blotch; *Drechslera teres*).

Ayers, J.E. Frank, J.A.; Cole, H. Jr.; Watson, G.R.; Broschious, S.C.; Gregory, L.V. (s.l.), The Society. Fungicide and nematicide tests; results - American Phytopathological Society. 1982. v. 37. p. 159-160. (NAL Call No.: 464.9 AM31R).

1191

Seed treatment for controlling covered smut of barley /by R.W. Leukel.

Leukel, R. W. 1888-. Washington : U.S. Dept. of Agriculture, 1930. Caption title. 23 p. : ill. ; 23 cm. Literature cited: p. 20-22. (NAL Call No.: DNAL 1 Ag84Te no.207).

1192

Seed treatment for controlling covered smut of barley by R.W. Leukel. -.

Leukel, R. W. Washington, D.C. U.S. Dept. of Agriculture 1930. 23 p. : ill. --. Bibliography: p. 20-22. (NAL Call No.: Fiche S-69 no.207).

1193

Seed treatment trials for control of oat smut, 1980 (Oats (*Avena sativa* 'Compact'), smut; *Ustilago kollerii*).

Jones, J.P. Dolezal, W.E.; Collins, F.C. (s.l.), The Society. Fungicide and nematicide tests; results - American Phytopathological Society. 1981. v. 36. p. 158. (NAL Call No.: 464.9 AM31R).

1194

Seed treatment with systemic fungicides to control *Cochliobolus sativus* on barley.

Luz, W.C. Vieira, J.C. St. Paul, Minn., American Phytopathological Society. Plant disease. Feb 1982. v. 66 (2). p. 135-136. Includes 6 ref. (NAL Call No.: 1.9 P69P).

1195

Seedborne diseases of cereals.

Forster, R.L. Schaad, N.W. Moscow, Idaho : The Service. Current information series - Cooperative Extension Service, University of Idaho. Nov 1988. (833). 4 p. plates. (NAL Call No.: DNAL 275.29 ID13IDC).

1196

Selection of barley for slow rusting resistance to leaf rust in epidemics of different severity.

CRPSAY. Andres, M.W. Wilcoxson, R.D. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1986. v. 26 (3). p. 511-514. Includes 25 references. (NAL Call No.: DNAL 64.8 C883).

1197

Selective toxin of *Helminthosporium victoriae*: thermal relationships in effects on oat tissues and protoplasts.

Briggs, S.P. Scheffer, R.P.; Haug, A.R. St. Paul, Minn. : American Phytopathological Society. Phytopathology. July 1984. v. 74 (7). p. 768-773. ill. Includes references. (NAL Call No.: 464.8 P56).

1198

Selective toxin of *Helminthosporium victoriae*: thermal relationships in effects on oat tissues and protoplasts (Blight disease of oats).

Briggs, S.P. Scheffer, R.P.; Haug, A.R. St. Paul, Minn. : American Phytopathological Society. Phytopathology. July 1984. v. 74 (7). p. 768-773. ill. Includes references. (NAL Call No.: 464.8 P56).

1199

Smut in oats, insecticides and fertilizer analyses / W.W. Cooke .

Cooke, Wells Woodbridge, 1858-1916. Burlington : Vermont State Agricultural Experiment Station, 1888. Cover title. 5, 2 p. ; 22 cm. (NAL Call No.: DNAL 100 V59 no.9).

1200

Smuts of cereal and forage crops in Kansas and their control / L.E. Melchers .

Melchers, L. E. 1887-. Manhattan, Kan. : Agricultural Experiment Station, Kansas State College of Agriculture and Applied Science, 1948. "Bulletin 279 rev.--Cover.--"Contribution no. 475, serial no. 391, Dept. of Botany"--p. 5. 36 p. : ill. ; 22 cm. (NAL Call No.: DNAL 100 K13S (1) no.332).

(PLANT DISEASES - FUNGAL)

1201

Smuts of cereal and forage crops in Kansas and their control / L.E. Melchers .

Melchers, L. E. 1887-. Manhattan, Kan. : Agricultural Experiment Station, Kansas State College of Agriculture and Applied Science, 1938. "Contribution no. 362 from the Dept. of Botany"--p. 5. 37 p. : ill. ; 23 cm. (NAL Call No.: DNAL 100 K13S (1) no.279).

1202

Solid-phase radioimmunoassay of ochratoxin A in serum.

JAFCAU. Rousseau, D.M. Slegers, G.A.; Peteghem, C.H. van. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. Sept/Oct 1986. v. 34 (5). p. 862-865. Includes references. (NAL Call No.: DNAL 381 J8223).

1203

Some synthetic phytotoxins structurally related to rhynchosporoside (produced by Rhynchosporium secalis, cause of scald in barley).

Beltran, J.P. Strobel, G.A.; Beier, R.; Mundy, B.P. Bethesda, Md., American Society of Plant Physiologists. Plant physiology. Mar 1980. v. 65 (3). p. 554-556. ill. 15 ref. (NAL Call No.: 450 P692).

1204

Sources of genes resistant to Puccinia hordei in barley (leaf rust, Hordeum vulgare, cultivars and lines).

Sharp, E.L. Reinhold, M. St. Paul, American Phytopathological Society. Plant disease. Nov 1982. v. 66 (11). p. 1012-1013. 14 ref. (NAL Call No.: 1.9 P69P).

1205

Specific binding of victorin to a 100-kDa protein from oats.

PNASA. Wolpert, T.J. Macko, V. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. June 1989. v. 86 (11). p. 4092-4096. ill. Includes references. (NAL Call No.: DNAL 500 N21P).

1206

Stamp out oat smut! (Fungicides).

Grau, C.R. Arny, D.C. Madison, Wis., The Programs. Publication - Cooperative Extension Programs, University of Wisconsin Extension. Jan 1981. Jan 1981. (A2894). 2 p. ill. (NAL Call No.: S544.3.W6W53).

1207

Stem rust of barley.

Dill-Macky, R. Rees, R.G.; Johnston, R.P. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 240-241. (NAL Call No.: DNAL SB191.B2A9 1987).

1208

Studies on the toxicity of blighted barley to swine /J.J. Christensen, H.C.H. Kernkamp.

Christensen, Jonas Jergon, 1892-. Kernkamp, H. C. H. St. Paul : University Farm, 1935? . Cover title. 28 p. : ill. ; 23 cm. Bibliography: p. 27-28. (NAL Call No.: DNAL 100 M66 (3) no.113).

1209

Sugar composition and acid invertase activity in spring barley plants in relation to adult-plant resistance to powdery mildew.

PHYTA. Hwang, B.K. Heitefuss, R. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Mar 1986. v. 76 (3). p. 365-369. Includes 19 references. (NAL Call No.: DNAL 464.8 P56).

1210

Systemic seed treatment for control of barley net blotch, 1981 (Pyrenophora teres on Hordeum vulgare).

Luz, W.C. da.FNETD. (s.l.) : The Society. Fungicide and nematicide tests : results - American Phytopathological Society. 1983. v. 38. p. 23. (NAL Call No.: 464.9 AM31R).

1211

Transfer of crown rust (Puccinia cornata) field resistance from Avena sterilis to cultivated oats by backcrossing (Abstract only).

Simons, M.D. St. Paul, Minn., American Phytopathological Society. Phytopathology. Aug 1981. v. 71 (8). p. 904. (NAL Call No.: 464.8 P56).

1212

Transfer of field resistance to Puccinia coronata from Avena sterilis to cultivated oats by backcrossing.

PHYTAJ. Simons, M.D. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Mar 1985. v. 75 (3). p. 314-317. Includes 26 references. (NAL Call No.: DNAL 464.8 P56).

1213

Un8 allele for loose smut resistance associated with necrosis in embryos of infected barley.
PHYTAJ. Gabor, B.K. Thomas, P.L. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Apr 1987. v. 77 (4). p. 533-538. ill. Includes references. (NAL Call No.: DNAL 464.8 P56).

1214

Uptake and translocation of Carbon-14-labeled fungicides in cereals: macro- and microautoradiographic studies.
ACSMC. Fuhr, F. Washington, D.C. : The Society. ACS Symposium Series - American Chemical Society. 1986. (304). p. 53-71. ill. Includes 33 references. (NAL Call No.: DNAL QD1.A45).

1215

Use of phytotoxins in selection of disease resistant mutants in tissue culture.
ISJRA6. Gengenbach, B.G. Rines, H.W. Ames, Iowa : Iowa State University. Iowa state journal of research. May 1986. v. 60 (4). p. 449-476. ill. Includes references. (NAL Call No.: DNAL 470 I09).

1216

Variation of resistance to powdery mildew in wild barley, Hordeum spontaneum.
Fukuyama, T. Heta, H.; Takeda, K. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Nov 10, 1986. v. 16. p. 15-17. (NAL Call No.: DNAL QK495.G74B34).

1217

Varietal resistance and susceptibility of oats to powdery mildew, crown rust and smuts / George M. Reed .
Reed, George M. 1878-1956. Columbia, Mo. : University of Missouri, College of Agriculture, Agricultural Experiment Station, 1920. 41 p., 4 p. of plates : ill. ; 23 cm. Bibliography: p. 39-41. (NAL Call No.: DNAL 100 M693 (3) no.37).

1218

Varietal susceptibility of oats to loose and covered smuts /by George M. Reed and Marion A. Griffiths and Fred N. Briggs.
Reed, George M. 1878-1956. Zehner, Marion Alice Griffiths, 1895-; Briggs, Fred N., 1896-. Washington, D.C. : U.S. Dept. of Agriculture, 1925. Cover title.~ "April, 1925."~ Chiefly tables. 40 p., 2 p. of plates ; 24 cm. Bibliography: p. 38-39. (NAL Call No.: DNAL 1 Ag84B no.1275).

1219

Victorin as a tool for elucidating the molecular mechanism of disease specificity in victoria blight of oats.
Macko, V. Wolpert, T.J.; Acklin, W.; Arigoni, D. Columbia, Mo. : The Interdisciplinary Plant Biochemistry and Physiology Program. Current topics in plant biochemistry and physiology : Proceedings of the ... Plant Biochemistry and Physiology Symposium held at the University of Missouri, Columbia. 1987. v. 6. p. 55-58. Includes references. (NAL Call No.: DNAL QK861.P55).

1220

Victorin-induced changes in carbohydrate metabolism in oat leaves / by Carroll David Rawn. -.
Rawn, Carroll David. Ann Arbor, Mich. University Microfilms International 1978. Thesis--University of Kentucky, 1974. Facsimile produced by microfilm-xerography. ix, 79, 3 leaves. Bibliography: leaves 73-78. (NAL Call No.: DISS 75-18,508).

PLANT DISEASES - BACTERIAL

1221

Effect of plant species and environmental conditions on epiphytic population sizes of *Pseudomonas syringae* and other bacteria.
PHYTA. O'Brien, R.D. Lindow, S.E. St. Paul, Minn. : American Phytopathological Society. Selected biological and environmental effects influenced epiphytic colonization of plants by *Pseudomonas syringae*, *Escherichia coli*, *Salmonella typhimurium*, *Aeromonas hydrophila*, and *Rhizobium meliloti* when tested in a growth chamber at 24 C. Epiphytic population size varied with plant host, environmental conditions, and among strains of *P. syringae* tested. Strains of *P. syringae* achieved only slightly larger population sizes than strains from other genera when incubated on inoculated plants for 48 hr, and near 100% relative humidity (RH). However, the strains of *P. syringae* maintained populations at least 25 times higher after a subsequent 72 hr at 40% RH. Epiphytic population sizes of 15 different strains of *P. syringae* varied up to 10-fold on a given plant species, indicating epiphytic diversity within this bacterial species. Relative population sizes of three strains of *P. syringae* on plants under field conditions were predicted by growth chamber populations. Neither epiphytic strains, pathogenic strains, or toxin producing groups were associated with greater epiphytic population sizes. Different plant species varied up to 17-fold in the size of bacterial populations supported. Maceration of inoculated plant tissue increased bacterial population size estimates relative to cells removed by sonication, but only after low RH incubations. *Phytopathology*. May 1989. v. 79 (5). p. 619-627. Includes references. (NAL Call No.: DNAL 464.8 P56).

1222

Effects of tabtoxinine-beta-lactam on nitrogen metabolism in *Avena sativa* L. roots.
PLPHA. Knight, T.J. Durbin, R.D.; Langston-Unkefer, P.J. Rockville, Md. : American Society of Plant Physiologists. *Plant physiology*. Dec 1986. v. 82 (4). p. 1045-1050. Includes references. (NAL Call No.: DNAL 450 P692).

1223

Evaluation of sources of resistance in oats to *Puccinia coronata* by use of many fungus cultures.
Michel, L.J. PIAIA. Simons, M.D. Cedar Falls : The Academy. *The Proceedings of the Iowa Academy of Science*. Dec 1983. v. 90 (4). p. 144-146. Includes references. (NAL Call No.: 500 I093).

1224

Extension of localized freeze injury in barley by acute post-thaw bacterial disease (*Bacillus subtilis*).
Olien, C.R. Smith, M.N. New York, Academic Press. *Cryobiology*. Aug 1981. v. 18 (4). p. 404-409. ill. 9 ref. (NAL Call No.: QH324.C7).

1225

Genetic control of ethirimol resistance in a natural population of *Erysiphe graminis* f.sp. *hordei* (Barley powdery mildew).
Hollomon, D.W. Laramie, The Station. *Science monograph - University of Wyoming, Agricultural Experiment Station*. May 1981. v. 71 (5). p. 536-540. 20 ref. (NAL Call No.: S131.E2).

1226

Markton, an oat variety immune from covered smut /T.R. Stanton, D.E. Stephens and E.F. Gaines.
Stanton, T. R. 1885-. Stephens, David E. 1874-; Gaines, E. F. 1886-. Washington, D.C. : U.S. Dept. of Agriculture, 1924. 8 p. : ill. ; 23 cm. Includes bibliographical references. (NAL Call No.: DNAL 1 Ag84D no.324).

1227

Oats tolerant of *Pseudomonas syringae* pv. *tabaci* contain tabtoxinine-beta-lactam-insensitive leaf glutamine synthetases.
PLPHA. Knight, T.J. Bush, D.R.; Langston-Unkefer, P.J. Rockville, Md. : American Society of Plant Physiologists. *Plant physiology*. Oct 1988. v. 88 (2). p. 333-339. Includes references. (NAL Call No.: DNAL 450 P692).

1228

Performance of small grain varieties in Louisiana, 1987-88.
Harrison, S.A. Boquet, D.J.; Colyer, P.D.; Groth, D.E.; Habetz, R.J.; Hallmark, W.B.; Hutchinson, R.L.; Moore, S.H.; Rabb, J.L. Baton Rouge, La. : The Department. *Report of projects - Louisiana Agricultural Experiment Station, Department of Agronomy*. 1988. p. 80-83. (NAL Call No.: DNAL 100 L936).

1229

Registration of 'Lamont' barley.
CRPSAY. Wesenberg, D.M. Robbins, G.S. Madison, Wis. : Crop Science Society of America. *Crop science*. Mar/Apr 1988. v. 28 (2). p. 373. Includes references. (NAL Call No.: DNAL 64.8 C883).

1230

A relationship between ice-nucleation-active bacteria, freeze damage, and genotype in oats. PHYTAJ. Marshall, D. St. Paul, Minn. : American Phytopathological Society. Phytopathology. July 1988. v. 78 (7). p. 952-957. Includes references. (NAL Call No.: DNAL 464.8 P56).

1231

Seedborne diseases of cereals.

Forster, R.L. Schaad, N.W. Moscow, Idaho : The Service. Current information series - Cooperative Extension Service, University of Idaho. Nov 1988. (833). 4 p. plates. (NAL Call No.: DNAL 275.29 ID131DC).

1232

Survival of ice nucleation-active and genetically engineered non-ice-nucleating *Pseudomonas syringae* strains after freezing. APMBA. Buttner, M.P. Amy, P.S. Washington, D.C. : American Society for Microbiology. The survival after freezing of ice nucleation-active (INA) and genetically engineered non-INA strains of *Pseudomonas syringae* was compared. Each strain was applied to oat seedlings and allowed to colonize for 3 days, and the plants were subjected to various freezing temperatures. Plant leaves were harvested before and after freezing on two consecutive days, and bacterial populations were determined. Populations of the INA wild-type strain increased 15-fold in the 18 h after the oat plants incurred frost damage at -5 and -12 degrees C. Plants colonized by the non-INA strain were undamaged at -5 degrees C and exhibited no changes in population size after two freeze trials. As freezing temperatures were lowered (-7, -9, and -12 degrees C), oat plants colonized by the non-INA strain suffered increased frost damage concomitant with bacterial population increases following 18 h. At -12 degrees C, both strains behaved identically. The data show a relationship between frost damage to plants and increased bacterial population size during the following 18 h, indicating a potential competitive advantage of INA strains of *P. syringae* over non-INA strains in mild freezing environments. Applied and environmental microbiology. July 1989. v. 55 (7). p. 1690-1694. ill. Includes references. (NAL Call No.: DNAL 448.3 AP5).

1233

Use of aluminum-foil and oat-straw mulches for controlling aster leafhopper, *Macrosteles fascifrons* (Homoptera: Cicadellidae), and aster yellows in carrots.

GRLEA. Setiawan, D.P. Ragsdale, D.W. East Lansing, Mich. : Michigan Entomological Society. The Great Lakes entomologist. Summer 1987. v. 20 (2). p. 103-109. Includes references. (NAL Call No.: DNAL QL461.M5).

1234

Xanthomonas campestris pv. *translucens* strains active in ice nucleation.

PLDIDE. Kim, H.K. Orser, C.; Lindow, S.E.; Sands, D.C. St. Paul, Minn. : American Phytopathological Society. Plant disease. Nov 1987. v. 71 (11). p. 994-997. Includes references. (NAL Call No.: DNAL 1.9 P69P).

PLANT DISEASES – VIRAL

1235

Aberrant ratio revisited.

Nelson, D.E. New York : Alan R. Liss. UCLA symposia on molecular and cellular biology. Paper presented at the "Symposium on Molecular and Cellular Biology of Plant Stress," April 15-21, 1984, Keystone, Colorado. 1985. v. 22. p. 1-12. Includes references. (NAL Call No.: DNAL QH506.U34).

1236

Aphid feeding behavior: relationship to barley yellow dwarf virus resistance in *Agropyron* species.

PHYTAU. Shukle, R.H. Lampe, D.U.; Lister, R.M.; Foster, J.E. St. Paul, Minn. : American Phytopathological Society. Phytopathology. May 1987. v. 77 (5). p. 725-729. ill. Includes references. (NAL Call No.: DNAL 464.8 P56).

1237

Barley stripe mosaic virus: its economic importance and control in Montana.

Carroll, T.W. St. Paul, Minn. : American Phytopathological Society. Plant disease. Feb 1980. v. 64 (2). p. 136-140. ill. 11 ref. (NAL Call No.: 1.9 P69P).

1238

Barley yellow dwarf of California cereals.

CAGRA. Griesbach, J.A. Creamer, R.; Lorens, G.; Falk, B.W.; Qualset, C.O.; Jackson, L.F. Berkeley, Calif. : The Station. California agriculture - California Agricultural Experiment Station. Jan/Feb 1989. v. 43 (1). p. 23-24. ill. (NAL Call No.: DNAL 100 C12CAG).

1239

Barley yellow dwarf symptom severity in oat affected by plant growth stage at infection and plot type.

CRPSAY. Goulart, L.R. Ohm, H.W.; Foster, J.E. Madison, Wis. : Crop Science Society of America. Lack of a standard method for evaluation of barley yellow dwarf (BYD) symptoms may result in inconsistency for genetic evaluation. The BYD symptoms were determined in oat, *Avena sativa* L., subsequent to virus infection at three plant growth stages and with two types of field plots. Five treatments: noninoculated; and inoculated at the two- to three-leaf (transplanted), three-leaf, four- to five-leaf, and stem elongation stages (direct seeded) were whole plots. Five oat cultivars were subplots. Two types of plots, hill and space planted, were whole units across the five cultivars. Seven traits were measured: BYD symptoms, plant height, heading date, grain yield and yield components. The plot types were not significantly different and there was a high

correlation between the types of plot for these characters. Cultivar rank was similar at all growth stages, indicating that sensitive cultivars may be severely damaged by BYDV infection at any stage of development. Small differences among cultivars for tolerance to BYDV were detected, subsequent to infection at the stem elongation stage under conditions of moderate temperatures and well distributed rainfall during the growing season. The plants that were transplanted after inoculation at the two- to three-leaf stage were more severely affected by BYDV than plants that were seeded in the field and inoculated at the three-leaf stage. Differences among cultivars were most distinct for BYD symptom, kernel weight, grain yield, number of kernels per panicle and plant height. Inoculation at the three-leaf stage resulted in maximum differences among cultivars and hill plots involved minimal effort for BYDV evaluation. Crop science. Nov/Dec 1989. v. 29 (6). p. 1412-1416. Includes references. (NAL Call No.: DNAL 64.8 C883).

1240

Barley yellow dwarf virus in Pennsylvania: effect of the PAV isolate on yield components of Noble spring oats.

PLDIDE. Gildow, F.E. Frank, J.A. St. Paul, Minn. : American Phytopathological Society. Plant disease. Mar 1988. v. 72 (3). p. 254-256. Includes references. (NAL Call No.: DNAL 1.9 P69P).

1241

Comparative characterization of two luteoviruses: beet western yellows virus and barley yellow dwarf virus.

PHYTAU. Hewings, A.D. D'Arcy, C.J. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Nov 1986. v. 76 (11). p. 1270-1274. ill. Includes 32 references. (NAL Call No.: DNAL 464.8 P56).

1242

Development of a barley germ plasm resistant to the seed transmission of barley stripe mosaic virus (Abstract only).

Carroll, T.W. Zaske, S.K.; Hockett, E.A. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Aug 1981. v. 71 (8). p. 865. (NAL Call No.: 464.8 P56).

1243

The effect of barley stripe mosaic virus on isozyme patterns in barley seedlings.

Wu, F.C. Timian, R.G. Grand Forks, N.D. : The Academy. Proceedings of the North Dakota Academy of Science. Apr 1984. v. 38. p. 98. Includes 5 references. (NAL Call No.: 500 N813).

1244

Effect of barley yellow dwarf virus infection on winter survival and other agronomic traits in barley.

Grafton, K.F. Pehlman, J.M.; Sechler, D.T.; Sehgal, D.P. Madison, Wis., Crop Science Society of America. Crop science. May/June 1982. v. 22 (3). p. 596-600. ill. 2 p. ref. (NAL Call No.: 64.8 C883)..

1245

The effects of infections by Pyrenophora teres and barley yellow dwarf virus on the freezing hardiness of winter barley.

PHYTAU. Delserone, L.M. Cole, H. Jr.; Frank, J.A. St. Paul, Minn. : American Phytopathological Society. Phytopathology. Oct 1987. v. 77 (10). p. 1435-1437. Includes references. (NAL Call No.: DNAL 464.8 P56).

1246

Feeding responses of Rhopalosiphum padi (Homoptera: Aphidae) to barley yellow dwarf virus resistant and susceptible barley varieties.

EVETEX. Ullman, D.E. Qualset, C.O.; McLean, D.L. Lanham, Md. : Entomological Society of America. The probing and feeding behavior of an important vector of barley yellow dwarf virus (BYDV), the bird cherry-oat aphid, Rhopalosiphum padi was electronically monitored on barley varieties with and without the Yd2 gene, a gene that imparts BYDV resistance to certain cereal cultivars. The presence of the Yd2 gene was shown to have no significant effect on the time elapsed to the first sieve element contact, total number of sieve element contacts, or duration of phloem sieve element ingestion by R. padi. Furthermore, aphid probing and feeding was not greatly affected by BYDV infection in susceptible plants. These data demonstrate that the mechanisms underlying BYDV resistance in plants with the Yd2 gene do not involve factors significantly influencing plant-aphid interactions that invoke alterations in aphid probing or feeding behavior. In addition, the number of sieve element contacts and duration of ingestion from sieve element cells by R. padi indicate that this aphid should be highly efficient in acquiring and transmitting the phloem-delimited BYDV to 'California Mariout' barley, with or without the Yd2 gene. Environmental entomology. Dec 1988. v. 17 (6). p. 988-991. Includes references. (NAL Call No.: DNAL QL461.E532).

1247

Generation mean analysis of inheritance of resistance to barley yellow dwarf in crosses involving bates spring oats.

PLDRA. Gellner, J.L. Sechler, D.T. St. Paul, Minn. : American Phytopathological Society. Plant disease. Aug 1986. v. 70 (8). p. 795-797. Includes references. (NAL Call No.: DNAL 1.9

P69P).

1248

Genetic analysis of a mutant resistant to barley yellow mosaic virus.

Ukai, Y. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Apr 10, 1984. v. 14. p. 31-33. Includes references. (NAL Call No.: QK495.G74B34).

1249

Genetic analysis of resistance to barley yellow dwarf virus in hybrids between Avena sativa 'Lamar' and virus-resistant lines of Avena sterilis (Oats).

Landry, B. Comeau, A.; Minvielle, F.; St.Pierre, C.A. Madison, Wis. : Crop Science Society of America. Crop science. Mar/Apr 1984. v. 24 (2). p. 337-340. ill. Includes references. (NAL Call No.: 64.8 C883).

1250

Genetic studies of resistance to barley yellow mosaic virus of German winter barley cultivars.

Friedt, W. Fort Collins, Colo. : Dept. of Agronomy, Colorado State University. Barley genetics newsletter. Oct 10, 1985. v. 15. p. 58-61. Includes references. (NAL Call No.: DNAL QK495.G74B34).

1251

High efficiency T7 polymerase synthesis of infectious RNA from cloned brome mosaic virus cDNA and effects of 5' extensions on transcript infectivity.

VIRLA. Janda, M. French, R.; Ahlquist, P. Duluth, Minn. : Academic Press. Virology. May 1987. v. 158 (1). p. 259-262. ill. Includes references. (NAL Call No.: DNAL 448.8 V81).

1252

In vitro selection for resistance.

Foroughi-Wehr, B. Friedt, W.; Schuchmann, R.; Kohler, F.; Wenzel, G. Hingham, Mass. : Martinus Nijhoff Publishers. Advances in agricultural biotechnology. 1986. (20). p. 35-44. ill. Includes references. (NAL Call No.: DNAL S494.5.B563A39).

(PLANT DISEASES - VIRAL)

1253

Mass rearing the bird cherry oat aphid, *Rhopalosiphum padi* (L.).
PIACA. Kudagamage, C. Foster, J.E.
Indianapolis, Ind. : The Academy. Proceedings
of the Indiana Academy of Science. Includes
abstract. 1985. v. 94. p. 304. (NAL Call No.:
DNAL 500 IN2).

1259

Using cDNA probes to identify barley yellow
dwarf virus-resistant barley lines.
Heath, R. Van de Velde, R.; Holloway, P. New
South Wales : Dept. of Agriculture, 1987? .
Proceedings, Australian Barley Technical
Symposium : Wagga Wagga, 11-15th October 1987.
p. 247-248. (NAL Call No.: DNAL SB191.B2A9
1987).

1254

Mosaics of winter oats and their control in the
southeastern states /by H.H. McKinney ... et
al. .
McKinney, Harold Hall, 1889-. Washington, D.C.
: U.S. Dept. of Agriculture, 1949. Caption
title. 17 p. : ill. ; 23 cm. Includes
bibliographical references. (NAL Call No.: DNAL
1 Ag84C no.809).

1255

Recurrent selection for tolerance to barley
yellow dwarf virus in oat.
CRPSAY. Baltenberger, D.E. Ohm, H.W.; Foster,
J.E. Madison, Wis. : Crop Science Society of
America. Crop science. May/June 1988. v. 28
(3). p. 477-480. Includes references. (NAL Call
No.: DNAL 64.8 C883).

1256

Registration of 'Venus' barley.
CRPSAY. Brown, A.R. Johnson, J.W.; Rothrock,
C.S.; Bruckner, P.L. Madison, Wis. : Crop
Science Society of America. Crop science.
July/Aug 1988. v. 28 (4). p. 718-719. Includes
references. (NAL Call No.: DNAL 64.8 C883).

1257

Relationship between seed infection by barley
stripe mosaic virus and yield loss (*Hordeum
vulgare*).
Nutter, F.W. Jr. Pederson, V.D.; Timian, R.G.
St. Paul : American Phytopathological Society.
Phytopathology. Mar 1984. v. 74 (3). p.
363-366. Includes references. (NAL Call No.:
464.8 P56).

1258

Structural comparison of *Poa semilatent virus*
and barley stripe mosaic virus.
PHYTA. Hunter, B.G. Heaton, L.A.; Bracker,
C.E.; Jackson, A.D. St. Paul, Minn. : American
Phytopathological Society. Phytopathology. Mar
1986. v. 76 (3). p. 322-326. ill. Includes 27
references. (NAL Call No.: DNAL 464.8 P56).

PLANT DISEASES - PHYSIOLOGICAL

1260

Differential response of barley varieties to Fe (iron) stress (*Hordeum vulgare*).

Fleming, A.L. Foy, C.D. New York ; Basel : Marcel Dekker, 1982. Iron nutrition and interactions in plants : Brigham Young University, August 12-14, 1981 / edited by S.D. Nelson ... (et al.). p. 457-468. ill. 18 ref. (NAL Call No.: QK867.J67 v. 5, nos. 4-7).

1261

Distribution of potassium between vacuole and cytoplasm in response to potassium deficiency.
NASSD. Storey, R. Leigh, R.A. New York, N.Y. : Plenum Press. NATO advanced science institutes series : Series A : Life sciences. In the series analytic: Plant vacuoles: their importance in solute compartmentation in cells and their applications in plant biotechnology / edited by B. Marin. Proceedings of a Workshop, July 6-11, 1986, Sophia-Antipolis, France. 1987. 134. p. 95-98. ill. Includes references. (NAL Call No.: DNAL QH301.N32).

1262

Effect of nitrate reductase deficiency upon growth, yield, and protein in barley.
Oh, J.Y. Warner, R.L.; Kleinhofs, A. Madison, Wis., Crop Science Society of America. Crop science. July/Aug 1980. v. 20 (4). p. 487-490. ill. 27 ref. (NAL Call No.: 64.8 C883).

1263

Effect of zinc deficiency on the accumulation of boron and other mineral nutrients in barley.
SSSJD4. Graham, R.D. Welch, R.M.; Grunes, D.L.; Cary, E.E.; Norvell, W.A. Madison, Wis. : The Society. Soil Science Society of America journal. May/June 1987. v. 51 (3). p. 652-657. Includes references. (NAL Call No.: DNAL 56.9 S03).

1264

The effects of iron and light treatments on chloroplast composition and ultrastructure in iron-deficient barley leaves (*Hordeum vulgare*).
Pushnik, J.C. Miller, G.W. New York ; Basel : Marcel Dekker, 1982. Iron nutrition and interactions in plants : Brigham Young University, August 12-14, 1981 / edited by S.D. Nelson ... (et al.). p. 311-321. ill. 26 ref. (NAL Call No.: QK867.J67 v. 5, nos. 4-7).

1265

Effects of P and K (phosphorus and potassium) on cereal crop diseases (Mineral deficiencies).
Piening, L.J. Atlanta, Ga., Potash & Phosphate Institute. Better crops with plant food. Winter 1981/1982. v. 66. p. 36-37, 39. (NAL Call No.: 6 B46).

1266

Effects of sulfur supply on the yield, composition, and quality of grain from cereals, oilseeds, and legumes.
ACSTD. Randall, P.J. Wrigley, C.W. St. Paul, Minn. : American Association of Cereal Chemists. Advances in cereal science and technology. Literature review. 1986. v. 8. p. 171-206. ill. Includes references. (NAL Call No.: DNAL TS2120.A3).

1267

Induction of alcohol dehydrogenase and lactate dehydrogenase in hypoxically induced barley.
PLPHA. Good, A.G. Crosby, W.L. Rockville, Md. : American Society of Plant Physiologists. In barley (*Hordeum vulgare* L.), alcohol dehydrogenase (ADH) and lactate dehydrogenase (LDH) are induced by anaerobiosis in both aleurone layers and roots. Under aerobic conditions, developing seeds of cv Himalaya accumulate ADH activity, which survives seed drying and rehydration. This activity consists almost entirely of the ADH1 homodimer. Activity of LDH also increases during seed development, but the level of activity in dry or rehydrated seeds is very low, indicating that this enzyme may not be involved in anaerobic glycolysis during the initial stages of germination. In contrast to ADH, the LDH isozymes present in developing seeds are similar to those found in uninduced and induced roots. Developmental expression of ADH and LDH was monitored from 0 to 24 days postgermination. Neither activity was induced to any extent in the germinating seeds; however, both enzymes were highly induced by anoxia in root tissue during development. Based on gel electrophoresis, this increase in activity results from the differential expression of different Adh and Ldh genes in root tissue. The changes in ADH and LDH activity levels were matched by changes in the amount of these particular proteins, indicating that the increase in activity results from de novo synthesis of these two proteins. The level of inducible LDH activity in an ADH1- mutant was not found to differ from cv Himalaya. We suggest that although the ADH- plants are more susceptible to flooding, they are not capable of responding to the lack of ADH1 activity by increasing the amount of LDH activity in root tissue. Plant physiology. July 1989. v. 90 (3). p. 860-866. ill. Includes references. (NAL Call No.: DNAL 450 P692).

(PLANT DISEASES - PHYSIOLOGICAL)

1268

An iron chelating compound released by barley roots in response to Fe-deficiency stress.
JPNUDS. Jolley, V.D. Brown, J.C.; Blaylock, M.J. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Jan 1988. v. 11 (1). p. 77-91. Includes references. (NAL Call No.: DNAL QK867.J67).

1269

Nitrate utilization by nitrate reductase-deficient barley mutants (Nutritional deficiencies).
Warner, R.L. Kleinhofs, A. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Apr 1981. v. 67 (4). p. 740-743. 111. 22 ref. (NAL Call No.: 450 P692).

1270

Peptides related to phytosiderophore secretion by Fe-deficient barley roots.
JPNUDS. Mori, S. Hachisuka, M.; Kawai, S.; Takagi, S.; Kishi-Nishizawa, N. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Fourth International Symposium on Iron Nutrition and Interactions in Plants," July 6-9, 1987, University of New Mexico, Albuquerque. June/Nov 1988. v. 11 (6/11). p. 653-662. Includes references. (NAL Call No.: DNAL QK867.J67).

1271

Relative toxicities of inorganic aluminum complexes to barley.
SSSJD4. Cameron, R.S. Ritchie, G.S.P.; Robson, A.D. Madison, Wis. : The Society. Soil Science Society of America journal. Sept/Oct 1986. v. 50 (5). p. 1231-1236. Includes references. (NAL Call No.: DNAL 56.9 S03).

MISCELLANEOUS PLANT DISORDERS

1272

Absorption and translocation of CGA-82725 with additives.

WEESA6. Gillespie, G.R. Skrzypczak, G.A.; Nalewaja, J.D. Champaign, Ill. : Weed Science Society of America. Abstract: The influence of various additives on CGA-82725 2-propargyl-2,4-(3,5-dichloro-2-pyridyloxy)phenoxy propanoate absorption and translocation was determined in oats (*Avena sativa* L. 'Lyon'). The absorption and translocation of 14C was greater when 14C-CGA-82725 was applied with petroleum oil compared to soybean Glycine max (L.) Merr oil. The translocation of 14C was greater at 96 than 48 h after 14C-CGA-82725 application. The absorption of 14C was greater at 48 than 24 h but was similar at 48 and 96 h after 14C-CGA-82725 application with no additive, petroleum oil, or soybean oil. The absorbed and translocated 14C was greater when 14C-CGA-82725 was applied with oil at 1.2 compared to 0.6 L/ha. No additional increase in 14C absorption and translocation was obtained if the oil volume was increased to 2.3 L/ha. The addition of petroleum oil to 14C-CGA-82725 increased 14C absorption and translocation more than the addition of palm (*Eleais quineensis* Jalq.), safflower (*Carthamus tinctorius* L.), linseed (*Linum usitatissimum* L.), or soybean oil. The four seed oils and the emulsifier. At Plus 300F caused similar increases in 14C absorption and translocation over 14C-CGA-82725 applied alone. Ethylene glycol did not increase 14C absorption and translocation compared to 14C-CGA-82725 applied alone. Weed science. May 1988. v. 36 (3). p. 282-285. Includes references. (NAL Call No.: DNAL 79.8 W41).

1273

Absorption and translocation of fluazifop with additives.

WEESA6. Nalewaja, J.D. Skrzypczak, G.A. Champaign, Ill. : Weed Science Society of America. Weed science. July 1986. v. 34 (4). p. 572-576. Includes 9 references. (NAL Call No.: DNAL 79.8 W41).

1274

Action of selected herbicides and Tween 20 on oat (*Avena sativa*) membranes.

Watson, M.C. Bartels, P.G.; Hamilton, K.C. Champaign, Ill., Weed Science Society of America. Weed science. Jan 1980. v. 28 (1). p. 122-127. ill. 28 ref. (NAL Call No.: 79.8 W41).

1275

Alfalfa and oat yields as influenced by triazine herbicide residues in three tillage systems.

PNWSB. Lazowski, E.J. Hartwig, N.L.; Hall, J.K. Beltsville, Md. : The Society. Proceedings of the ... annual meeting - Northeastern Weed Science Society. 1986. v. 40. p. 10-15. Includes references. (NAL Call No.: DNAL 79.9

N814).

1276

Aluminum-induced changes in calmodulin.

NASSD. Haug, A. Weis, C. New York, N.Y. : Plenum Press. NATO advanced science institutes series : Series A : Life sciences. In the series analytic: Molecular and cellular aspects of calcium in plant development / edited by A. J. Trewavas. 1986. v. 104. p. 19-25. Includes references. (NAL Call No.: DNAL QH301.N32).

1277

Analysis of midwinter freezing stress (Barley, hardiness, injuries).

Olien, C.R. Boca Raton, Fla., CRC Press. Analysis and improvement of plant cold hardiness. 1981. p. 35-59. 65 ref. (NAL Call No.: SB781.A52).

1278

The antagonistic action of 2,4-D and bromoxynil on glyphosate phytotoxicity to barley (*Hordeum vulgare*) (Herbicide mixtures).

O'Donovan J.T. O'Sullivan, P.A. Champaign, Ill., Weed Science Society of America. Weed science. Jan 1982. v. 30 (1). p. 30-34. Includes 6 ref. (NAL Call No.: 79.8 W41).

1279

Antidote action of humic substances on atrazine inhibition of sulfate uptake in barley roots (Pesticide toxicity).

Dell'Agnola, G. Ferrari, G.; Nardi, S. New York, Academic Press. Pesticide biochemistry and physiology. Apr 1981. v. 15 (2). p. 101-104. ill. 7 ref. (NAL Call No.: SB951.P49).

1280

Barley (*Hordeum vulgare*) response to herbicides applied at three growth stages.

WETEE9. Martin, D.A. Miller, S.D.; Alley, H.P. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Jan 1988. v. 2 (1). p. 41-45. Includes references. (NAL Call No.: DNAL SB610.W39).

1281

Behavior and fate of ethoxylated alkylphenol nonionic surfactant in barley plants (*Hordeum vulgare*, uptake, phytotoxicity).

Stolzenberg, G.E. Olson, P.A.; Zaylskie, R.G.; Mansager, E.R. Washington, D.C., American Chemical Society. Journal of agricultural and food chemistry. July/Aug 1982. v. 30 (4). p. 637-644. ill. 2 p. ref. (NAL Call No.: 381

(MISCELLANEOUS PLANT DISORDERS)

J8223).

1282

Betaine accumulation: metabolic pathways and genetics.

Hanson, A.D. Grumet, R. New York : Alan R. Liss. UCLA symposia on molecular and cellular biology. Paper presented at the "Symposium on Molecular and Cellular Biology of Plant Stress," April 15-21, 1984, Keystone, Colorado. 1985. v. 22. p. 71-92. ill. Includes references. (NAL Call No.: DNAL QH506.U34).

1283

Boron toxicity in barley.

JPNUDS. Riley, M.M. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Tenth International Plant Nutrition Colloquium," August 4-9, 1986, Beltsville, Maryland. 1987. v. 10 (9/16). p. 2109-2115. Includes references. (NAL Call No.: DNAL QK867.J67).

1284

Breeding for preharvest sprouting tolerance.

Derera, N.F. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 111-128. ill. Includes references. (NAL Call No.: DNAL SB189.6.P73).

1285

Cadmium-induced accumulation of putrescine in oat and bean leaves.

PLPHA. Weinstein, L.H. Kaur-Sawhney, R.; Rajam, M.V.; Wettlaufer, S.H.; Galston, A.W. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Nov 1986. v. 82 (3). p. 641-645. Includes references. (NAL Call No.: DNAL 450 P692).

1286

Damage to barley spikes resulting from impact momentum similar to small sized hail.

AGUDAT. Ferguson, H. Jones, A.J.; Tsai, K.J. Madison, Wis. : American Society of Agronomy. Agronomy journal. Nov/Dec 1987. v. 79 (6). p. 1015-1018. Includes references. (NAL Call No.: DNAL 4 AM34P).

1287

Differential aluminum tolerances of two barley cultivars related to organic acids in their roots.

JPNUDS. Foy, C.D. Lee, E.H.; Wilding, S.B. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Tenth International Plant Nutrition Colloquium," August 4-9, 1986, Beltsville, Maryland. 1987. v. 10 (9/16). p. 1089-1101. Includes references. (NAL Call No.: DNAL QK867.J67).

1288

Differential inhibition of potassium ion absorption by difenzoquat in wild oat and cereals (Herbicide toxicity).

Cohen, A.S. Morrison, I.N. New York, Academic Press. Pesticide biochemistry and physiology. Oct 1982. v. 18 (2). p. 174-179. ill. 23 ref. (NAL Call No.: SB951.P49).

1289

Differential mRNA transcription during salinity stress in barley.

PNASA. Ramagopal, S. Washington, D.C. : The Academy. Proceedings of the National Academy of Sciences of the United States of America. Jan 1987. v. 84 (1). p. 94-98. ill. Includes references. (NAL Call No.: DNAL 500 N21P).

1290

Drought sensitivity at various growth stages of barley in relation to relative evapotranspiration and water stress.

Mogensen, V.O. Madison, Wis., American Society of Agronomy. Agronomy journal. Nov/Dec 1980. v. 72 (6). p. 1033-1038. ill. 12 ref. (NAL Call No.: 4 AM34P).

1291

Effect of application time on soil residue and efficacy of sulfonylureas.

SWSPBE. Foy, C.L. Mersie, W. Raleigh, N.C. : The Society. Proceedings - Southern Weed Science Society. 1986. (39th). p. 446-456. Includes references. (NAL Call No.: DNAL 79.9 S08 (P)).

1292

Effect of environment and adjuvants on asulam phytotoxicity.

WEESA6. Nalewaja, J.D. Woznica, Z. Champaign, Ill. : Weed Science Society of America. Abstract: Experiments were conducted to determine the influence of various factors on asulam methyl (4-amino-phenyl)sulfonyl carbamate toxicity to flax (*Linum usitatissimum* L.) and

(MISCELLANEOUS PLANT DISORDERS)

wild oats (*Avena fatua* L. ~ AVEFA). Asulam toxicity to both flax and wild oats generally increased as temperature, humidity, and soil moisture increased after treatment. Octoxynol (alpha- p-1,1,3,3-tetramethyl butyl phenyl -omega-hydroxypoly(oxyethylene) in the spray solution increased asulam toxicity to both species in all environments. Octoxynol and trimethylenonypolyethoxyethanol (WK) enhanced asulam toxicity more than other adjuvants evaluated. Asulam toxicity to both flax and wild oats increased as octoxynol concentration in the spray increased. Flax tolerance to asulam generally increased with flax height at treatment. 'Flor' flax was the most asulam susceptible of six cultivars evaluated. A 2-mm simulated rainfall within 3 or 6 h after asulam treatment reduced toxicity to wild oats and flax, respectively. Weed science. May 1988. v. 36 (3). p. 367-372. Includes references. (NAL Call No.: DNAL 79.8 W41).

1293

Effect of ethephon-bromoxynil and ethephon-DPXR9674 on spring barley yield. Lish, J.M. Thill, D.C. S.L. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 200. (NAL Call No.: DNAL 79.9 W52R).

1294

Effect of gaseous hydrogen chloride on seed germination and early development of seedlings (Tomatoes, barley, injury, air pollution). Granett, A.L. Taylor, O.C. Alexandria, Va., The Society. Journal of the American Society for Horticultural Science. July 1980. v. 105 (4). p. 548-550. 11 ref. (NAL Call No.: 81 S012).

1295

The effect of injury in imitation of hail damage on the development of small grain /by John C. Eldredge. Eldredge, John C. 1886-. Ames, Iowa : Agricultural Experiment Station, Iowa State College of Agriculture and Mechanic Arts, 1937. p. 286-302 : ill., charts ; 22 cm. Bibliography: p. 302. (NAL Call No.: DNAL 100 Io9 no.219).

1296

Effect of MCPA on 14C-Diclofop uptake and translocation (Wild oat, *Avena fatua*, phytotoxicity). Olson, W. Nalewaja, J.D. Champaign, Ill., Weed Science Society of America. Weed science. Jan 1982. v. 30 (1). p. 59-63. Includes 18 ref. (NAL Call No.: 79.8 W41).

1297

Effect of NaCl (sodium chloride) on some trace metals in barley (Salinity, interactions, cobalt, lithium, strontium). Wallace, A. Cha, J.W.; Mueller, R.T. New York, Marcel Dekker. Journal of plant nutrition. 1980. v. 2 (1/2). p. 115-117. ill. 5 ref. (NAL Call No.: QK867.J67).

1298

Effect of simulated hail damage on barley /E.L. Deckard and J.J. Hammond. Deckard, Edward Lee, 1943-. Hammond, James J. Fargo : Agricultural Experiment Station, North Dakota State University, 1978. Cover title. 8 p. : ill., charts, 1 map ; 28 cm. Bibliography: p. 8. (NAL Call No.: DNAL 100 N813 no.505).

1299

Effect of small-scale composting of sewage sludge on heavy metal availability to plants (*Lactuca sativa*, *Avena sativa*, cadmium, copper, zinc). Simeoni, L.A. Barbarick, K.A.; Sabey, B.R. Madison, Wis. : American Society of Agronomy. Journal of environmental quality. Apr/June 1984. v. 13 (2). p. 264-268. Includes references. (NAL Call No.: QH540.J6).

1300

Effects of aluminum on nutrient composition and yield of oats (Toxicity, chlorosis). Alam, S.M. Adams, W.A. New York, Marcel Dekker. Journal of plant nutrition. 1979. v. 1 (4). p. 365-375. ill. 31 ref. (NAL Call No.: QK867.J67).

1301

Effects of chlorsulfuron on meiosis and seed viability in rye (*Secale cereale* L.). WSWPA. Zollinger, R.K. Evans, J.G. Reno : The Society. Proceedings - Western Society of Weed Science. 1985. v. 38. p. 114-119. (NAL Call No.: DNAL 79.9 W52).

1302

Effects of haloxyfop and CGA-82725 on cell cycle and cell division of oat (*Avena sativa*) root tips. WEESA6. Kim, J.C. Bendixen, L.E. Champaign, Ill. : Weed Science Society of America. Weed science. Nov 1987. v. 35 (6). p. 769-774. Includes references. (NAL Call No.: DNAL 79.8 W41).

(MISCELLANEOUS PLANT DISORDERS)

1303

The effects of preharvest rain.

Derera, N.F. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 1-14. ill., maps. Includes references. (NAL Call No.: DNAL SB189.6.P73).

1304

Effects of variations in drop makeup on the phytotoxicity of glyphosate (Herbicides, controlled-drop application, barley).

Ambach, R.M. Ashford, R. Champaign, Ill., Weed Science Society of America. Weed science. May 1982. v. 30 (3). p. 221-224. Includes 6 ref. (NAL Call No.: 79.8 W41).

1305

Ethylene as an effector of wound-induced resistance to cellulase in oat leaves (*Avena sativa*).

Geballe, G.T. Galston, A.W. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1982. v. 70 (3). p. 788-790. 31 ref. (NAL Call No.: 450 P692).

1306

Evaluation of chlorophyll fluorescence parameters for an intact-plant herbicide bioassay.

CRPSAY, Shaw, D.R. Peeper, T.F.; Nofziger, D.L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 756-760. Includes references. (NAL Call No.: DNAL 64.8 C883).

1307

Evidence for allelopathy by residues and aqueous extracts of rye (*Secale cereale*).

WEESA6. Barnes, J.P. Putnam, A.R. Champaign, Ill. : Weed Science Society of America. Weed science. May 1986. v. 34 (3). p. 384-390. Includes references. (NAL Call No.: DNAL 79.8 W41).

1308

Extension of localized freeze injury in barley by acute post-thaw bacterial disease (*Bacillus subtilis*).

Olien, C.R. Smith, M.N. New York, Academic Press. Cryobiology. Aug 1981. v. 18 (4). p. 404-409. ill. 9 ref. (NAL Call No.: QH324.C7).

1309

Gas chromatography-mass spectrometry of acylalanine fungicides.

JAFCAU. Ripley, B.D. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. July/Aug 1985. v. 33 (4). p. 560-563. Includes references. (NAL Call No.: DNAL 381 J8223).

1310

Genetic variability for herbicide reaction in plant populations (Wild oat, *Avena barbata*, *Avena fatua*, *godetia*, *Clarkia williamsonii*, toxicity, Sierra Nevada in California).

Price, S.C. WEESA. Hill, J.E.; Allard, R.W. Champaign : Weed Science Society of America. Weed science. Sept 1983. v. 31 (5). p. 652-657. ill. Includes references. (NAL Call No.: 79.8 W41).

1311

Grass weed control in California cereals: update and review.

WSWPA. Mitich, L.W. Kyser, G.B. Reno : The Society. Proceedings - Western Society of Weed Science. Paper presented at the annual meeting of the Western Society of Weed Science, March 18-20, 1986, San Diego, California. 1986. v. 39. p. 169-172. (NAL Call No.: DNAL 79.9 W52).

1312

Growth and composition of alfalfa fertilized in greenhouse trials with deproteinized juice from low and high saponin alfalfa and from oat herbage (Phytotoxicity).

Weich, D.A. Smith, D.; Soberalske, R.M.; Ream, H.W. New York, Marcel Dekker. Journal of plant nutrition. 1979. v. 1 (2). p. 151-170. ill. 15 ref. (NAL Call No.: QK867.J67).

1313

Growth inhibition and disruption of mitosis by DCPA in oat (*Avena sativa*) roots.

WEESA6. Holmsen, J.D. Hess, F.D. Champaign, Ill. : Weed Science Society of America. Weed science. Nov 1984. v. 32 (6). p. 732-738. ill. Includes 28 references. (NAL Call No.: DNAL 79.8 W41).

1314

Herbicidal disruption of proton gradient development and maintenance by plasmalemma and tonoplast vesicles from oat root.

PCBPB. Ratterman, D.M. Balke, N.E. Duluth, Minn. : Academic Press. Pesticide biochemistry and physiology. Literature review. July 1988. v. 31 (3). p. 221-236. Includes references. (NAL Call No.: DNAL SB951.P49).

(MISCELLANEOUS PLANT DISORDERS)

1315

Ice nucleation temperature of individual leaves in relation to population sizes of ice nucleation active bacteria and frost injury.
PLPHA. Hirano, S.S. Baker, L.S.; Upper, C.D. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Feb 1985. v. 77 (2). p. 259-265. ill. Includes 25 references. (NAL Call No.: DNAL 450 P692).

1316

Induction of alpha-amylase inhibitor synthesis in barley embryos and young seedlings by abscisic acid and dehydration stress.
PLPHA. Robertson, M. Walker-Simmons, M.; Munro, D.; Hill, R.D. Rockville, Md. : American Society of Plant Physiologists. An endogenous alpha-amylase inhibitor was found to be synthesized in embryos of developing barley grain (*Hordeum vulgare* cv Bonanza). Accumulation of this protein occurred late in development (stage IV,) at the same time that endogenous abscisic acid (ABA) showed a large increase. The inhibitor could be induced up to 23-fold in isolated immature embryos (stage III) by culture in ABA. Precocious germination was also blocked in stage III embryos by ABA. Dehydration stress on the isolated immature embryos also induced higher levels of the inhibitor and ABA. An even greater response to dehydration stress was observed in young seedlings, where inhibitor content increased 20-fold and ABA increased 80-fold during water stress. The high degree of correlation between ABA and inhibitor contents in in situ embryos, dehydrated embryos and young seedlings, as well as the increase in inhibitor caused by exogenously applied ABA to isolated embryos, suggest that increased alpha-amylase inhibitor synthesis in response to dehydration stress is mediated by ABA. Plant physiology. Sept 1989. v. 91 (1). p. 415-420. Includes references. (NAL Call No.: DNAL 450 P692).

1317

Injury and grain yield of spring barley treated with diclofop and thiameturon.
Evans, R.M. Thill, D.C. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 193-195. (NAL Call No.: DNAL 79.9 W52R).

1318

Long term residual effects of applied selenium on the selenium uptake by plants (Cereals, forages, Canada).
Gupta, U.C. Winter, K.A. New York, Marcel Dekker. Journal of plant nutrition. 1981. v. 3 (1/4). p. 493-502. 19 ref. (NAL Call No.: QK867.J67).

1319

Loss of metribuzin and ethyl-metribuzin from glass and soil surfaces.
WETEE9. Peek, D.C. Appleby, A.P. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Jan/Mar 1989. v. 3 (1). p. 173-176. Includes references. (NAL Call No.: DNAL SB610.W39).

1320

Low temperature injury in winter cereals.
Pomeroy, M.K. Andrews, C.J. Boca Raton, FL : CRC Press, c1989. Low temperature stress physiology in crops / editor, Paul H. Li. Literature review. p. 107-122. ill. Includes references. (NAL Call No.: DNAL SB781.L68).

1321

Mechanical properties of the plasma membrane of isolated plant protoplasts. Mechanism of hyperosmotic and extracellular freezing injury (Secale cereale, rye seedlings, changes in area and membrane contraction in lysis).
Wolfe, J. PLPHA. Steponkus, P.L. Rockville : American Society of Plant Physiologists. Plant physiology. Feb 1983. v. 71 (2). p. 276-285. ill. 28 ref. (NAL Call No.: 450 P692).

1322

Metribuzin absorption and translocation in two barley (*Hordeum vulgare*) cultivars.
WEESA6. Gawronski, S. Haderlie, L.C.; Stark, J.C. Champaign, Ill. : Weed Science Society of America. Weed science. July 1986. v. 34 (4). p. 491-495. ill. Includes 13 references. (NAL Call No.: DNAL 79.8 W41).

1323

Mineral element concentration of two barley cultivars in relation to water deficit and aluminum toxicity.
JPNUDS. Krizek, D.T. Foy, C.D. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Apr 1988. v. 11 (4). p. 369-386. ill. Includes references. (NAL Call No.: DNAL QK867.J67).

1324

Mineral nutrition of plants exposed to salinity stress.
Lauchli, A. Davis : University of California, Davis?, 1981? . A Conference on biosalinity : the problem of salinity in agriculture : a joint conference of Egyptian, Israeli and American scientists, Univ. of California, Davis, September 1-4, 1981 / organized and. p. 63-69. ill. Includes 8 references. (NAL Call No.: DNAL S619.S24C6).

(MISCELLANEOUS PLANT DISORDERS)

1325

Mode of action studies on nitrodiphenyl ether herbicides. 1. Use of barley mutants to probe the role of photosynthetic electron transport. PLPHA. Bowyer, J.R. Smith, B.J.; Camilleri, P.; Lee, S.A. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Mar 1987. v. 83 (3). p. 613-620. ill. Includes references. (NAL Call No.: DNAL 450 P692).

1326

Molecular biology of salinity stress: preliminary studies and perspectives. Ramagopal, S. New York : Plenum Press, c1987. Tailoring genes for crop improvement : an agricultural perspective / edited by George Bruening ... et al. . p. 111-119. ill. Includes references. (NAL Call No.: DNAL SB123.57.C66 1986).

1327

Molybdenum enrichment of plants grown on fly ash-treated soils (Coal combustion residues, Alfalfa, Cynodon dactylon, white clover, barley, Swiss chard). Elseewi, A.A. Page, A.L. Madison, Wis. : American Society of Agronomy. Journal of environmental quality. July/Sept 1984. v. 13 (3). p. 394-398. ill. Includes references. (NAL Call No.: QH540.J6).

1328

Mutagenicity of selected chemicals in barley: repair and recovery (Storage of seeds treated with alkylating agents). Veleminsky, J. Gichner, T. New York, Plenum Press. Environmental science research. 1982. v. 24. p. 321-327. 58 ref. (NAL Call No.: TD172.E55).

1329

Mutagenicity of selected chemicals in barley test systems (Includes agricultural chemicals, environmental pollutants). Nilan, R.A. Veleminsky, J. New York, Plenum Press. Environmental science research. 1982. v. 24. p. 291-320. ill. 63 ref. (NAL Call No.: TD172.E55).

1330

Path coefficient analysis of correlation between stress and barley yield components. Campbell, W.F. Wagenet, R.U.; Bamatraf, A.M.; Turner, D.L. Madison, Wis., American Society of Agronomy journal. Nov/Dec 1980. v. 72 (6). p. 1012-1016. ill. 15 ref. (NAL Call No.: 4 AM34P).

1331

Phase transitions in liposomes formed from the polar lipids of mitochondria from chilling-sensitive plants. PLPHA. Raison, J.K. Orr, G.R. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. July 1986. v. 81 (3). p. 807-811. Includes 26 references. (NAL Call No.: DNAL 450 P692).

1332

Physiological and cytological aspects of manganese toxicity in barley seedlings. JPNUDS. Riedell, W.E. Schmid, W.E. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. 1986. v. 9 (1). p. 57-66. ill. Includes references. (NAL Call No.: DNAL QK867.J67).

1333

Physiology and biochemistry of Al³⁺ (aluminum ion) toxicity in barley roots (Hordeum vulgare). Siegel, N. Haug, A. East Lansing, Mich., The Laboratory. Annual report - Michigan State University, MSU/DOE Plant Research Laboratory. 1981. 1981 (16th). p. 75. (NAL Call No.: QK1.M5).

1334

Physiology of sprouting resistance. King, R.W. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 27-60. ill. Includes references. (NAL Call No.: DNAL SB189.6.P73).

1335

Phytotoxic interaction studies--Techniques for evaluation and presentation of results (Interactions of two or more pesticides in plants, tested on oats and cucumbers). Nash, R.G. Champaign, Ill., Weed Science Society of America. Weed science. Mar 1981. v. 29 (2). p. 147-155. ill. 22 ref. (NAL Call No.: 79.8 W41).

1336

Plasma membrane ATPase activity following reversible and irreversible freezing injury. PLPHA. Iswari, S. Palta, J.P. Rockville, Md. : American Society of Plant Physiologists. Plasma membrane ATPase has been proposed as a site of functional alteration during early stages of freezing injury. To test this, plasma membrane was purified from Solanum leaflets by a single step partitioning of microsomes in a dextran-polyethylene glycol two phase system.

(MISCELLANEOUS PLANT DISORDERS)

Addition of lysolecithin in the ATPase assay produced up to 10-fold increase in ATPase activity. ATPase activity was specific for ATP with a K_m around 0.4 millimolar. Presence of the ATPase enzyme was identified by immunoblotting with oat ATPase antibodies. Using the phase partitioning method, plasma membrane was isolated from *Solanum commersonii* leaflets which had four different degrees of freezing damage, namely, slight (reversible), partial (partially reversible), substantial and total (irreversible). With slight (reversible) damage the plasma membrane ATPase specific activity increased 1.5- to 2-fold and its K_m was decreased by about 3-fold, whereas the specific activity of cytochrome c reductase and cytochrome c oxidase in the microsomes were not different from the control. However, with substantial (lethal, irreversible) damage, there was a loss of membrane protein, decrease in plasma membrane ATPase specific activity and decrease in K_m while cytochrome c oxidase and cytochrome c reductase were unaffected. These results support the hypothesis that plasma membrane ATPase is altered by slight freeze-thaw stress. Plant physiology. July 1989. v. 90 (3). p. 1088-1095. ill. Includes references. (NAL Call No.: DNAL 450 P692).

1337

Polyamine metabolism and plant stress.

Flores, H.E. Young, N.D.; Galston, A.W. New York : Alan R. Liss. UCLA symposia on molecular and cellular biology. Paper presented at the "Symposium on Molecular and Cellular Biology of Plant Stress," April 15-21, 1984, Keystone, Colorado. 1985. v. 22. p. 93-114. Includes references. (NAL Call No.: DNAL OH506.U34).

1338

Preharvest sprouting damage and sprouting tolerance: assay methods and instrumentation. Mares, D.J. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 129-170. ill. Includes references. (NAL Call No.: DNAL SB189.6.P73).

1339

Preparation and use of mixed fumigant standards for multiresidue level determination by gas chromatography.

JAFCAU. Daft, J.L. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. July/Aug 1985. v. 33 (4). p. 563-566. Includes references. (NAL Call No.: DNAL 381 J8223).

1340

Pronamide phytotoxicity in ten Wisconsin soils (Oats).

Dutt, T.E. Harvey, R.G. Champaign, Ill., Weed Science Society of America. Weed science. July 1980. v. 28 (4). p. 429-432. ill. 15 ref. (NAL Call No.: 79.8 W41).

1341

Protection of grass crops from sulfonylurea and imidazolinone toxicity.

Barrett, M. San Diego : Academic Press, c1989. Crop safeners for herbicides : development, uses, and mechanisms of action / edited by Kriton K. Hatzios and Robert E. Hoagland. Literature review. p. 195-220. Includes references. (NAL Call No.: DNAL SB951.45.C76).

1342

A rapid method for the analysis of the mode of action of bleaching herbicides.

PCBPB. Young, A.J. Britton, G.; Musker, D. Duluth, Minn. : Academic Press. Pesticide biochemistry and physiology. Nov 1989. v. 35 (3). p. 244-250. Includes references. (NAL Call No.: DNAL SB951.P49).

1343

Recovery from winter injury (Barley).

Smith, M.N. Olien, C.R. Boca Raton, Fla., CRC Press. Analysis and improvement of plant cold hardiness. 1981. 1981. p. 117-138. 97 ref. (NAL Call No.: SB781.A52).

1344

Registration of 'Hercules' oat.

CRPSAY. Marshall, H.G. Kolb, F.L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1988. v. 28 (4). p. 719. Includes references. (NAL Call No.: DNAL 64.8 C883).

1345

A relationship between ice-nucleation-active bacteria, freeze damage, and genotype in oats.

PHYTAU. Marshall, D. St. Paul, Minn. : American Phytopathological Society. Phytopathology. July 1988. v. 78 (7). p. 952-957. Includes references. (NAL Call No.: DNAL 464.8 P56).

(MISCELLANEOUS PLANT DISORDERS)

1346

Residual effect of metsulfuron applied during the fallow year on barley and lentils.
Mashhadi, H.R. Evans, J.D. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 258-259. (NAL Call No.: DNAL 79.9 W52R).

1347

Response of oats to atrazine (Herbicides).
Brinkman, M.A. Langer, D.K.; Harvey, R.G.; Hardie, A.R. Madison, Wis., Crop Science Society of America. Crop science. Mar/Apr 1980. v. 20 (2). p. 185-189. ill. 18 ref. (NAL Call No.: 64.8 C883).

1348

Response to spring barley (*Hordeum vulgare*) to herbicides.
WETEE9. Clay, S.A. Thill, D.C.; Cochran, V.L. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Jan 1988. v. 2 (1). p. 68-71. Includes references. (NAL Call No.: DNAL SB610.W39).

1349

Role of benzoxazinones in allelopathy by rye (*Secale cereale* L.).
JCECD. Barnes, J.P. Putnam, A.R. New York, N.Y. : Plenum Press. Journal of chemical ecology. Apr 1987. v. 13 (4). p. 889-906. Includes references. (NAL Call No.: DNAL QD415.A1J6).

1350

Role of water stress in differential aluminum tolerance of two barley cultivars grown in an acid soil.
JPNUDS. Krizek, D.T. Foy, C.D. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Apr 1988. v. 11 (4). p. 351-367. ill. Includes references. (NAL Call No.: DNAL QK867.J67).

1351

Rye residues contribute weed suppression in no-tillage cropping systems (Agroecosystems, biomass).
Barnes, J.P. JCECD. Putnam, A.R. New York : Plenum Press. Journal of chemical ecology. Aug 1983. v. 9 (8). p. 1045-1057. ill. Includes references. (NAL Call No.: QD415.A1J6).

1352

Seedling injury and chromosome aberrations induced by Bladex, Dowpon, Princep and Tenoran (Herbicides, barley).
Kahlon, P.S. Nashville, The Academy. Journal of the Tennessee Academy of Science. Tennessee Academy of Science. Jan 1980. v. 55 (1). p. 17-19. ill. 9 ref. (NAL Call No.: 500 T25A).

1353

The site of the inhibition of the shikimate pathway by glyphosate (herbicide, buckwheat). II. Interference of glyphosate with chorismate formation in vivo and in vitro.
Amrhein, N. Deus, B.; Gehrke, P.; Steinrucken, H.C. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Nov 1980. v. 66 (5). p. 830-834. ill. 24 ref. (NAL Call No.: 450 P692).

1354

The site of the inhibition of the shikimate pathway by glyphosate (postemergence herbicide). I. Inhibition by glyphosate of phenylpropanoid synthesis in buckwheat (*Fagopyrum esculentum* Moench).
Hollander, H. Amrhein, N. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Nov 1980. v. 66 (5). p. 823-829. ill. 28 ref. (NAL Call No.: 450 P692).

1355

Soil bound residues of carbaryl and 1-naphthol: release and mineralization in soil, and uptake by plants.
JPFCD2. Murthy, N.B.K. Raghu, K. New York, N.Y. : Marcel Dekker. Journal of environmental science and health : Part B : Pesticides, food contaminants, and agricultural wastes. 1988. v. 23 (6). p. 575-585. Includes references. (NAL Call No.: DNAL TD172.J61).

1356

Some effects of arsenical vine killers on potatoes and oats /G.L. Terman, B.E. Plummer, Jr., and Donald Folsom.
Terman, G. L. Plummer, Bernie E.; Folsom, Donald, 1891-. Orono : Maine Agricultural Experiment Station, 1952. 11 p. : ill. ; 23 cm. Bibliography: p. 11. (NAL Call No.: DNAL 100 M28S (1) no.501).

(MISCELLANEOUS PLANT DISORDERS)

1357

Spring oats response to metsulfuron treatment during the fallow year.

Mashadi, H.R. Evans, J.D. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 260. (NAL Call No.: DNAL 79.9 W52R).

1358

Studies on the cytogenetic and genetic effects of fluoride on barley / by Shankarappa S. Bale.

Bale, Shankarappa S. (Shankarappa Siddaveerasetty), 1940-. 1971. Thesis (Ph.D.)--Texas A&M University, 1971. Photocopy. Ann Arbor, Mich. : University Microfilms, 1972. xiii, 90 leaves ; 21 cm. Bibliography: leaves 82-89. (NAL Call No.: DISS 72-5,635).

1359

Techniques for selection of cold hardiness in cereals (Oats).

Marshall, G.H. Olien, C.R.; Everson, E.H. Boca Raton, Fla., CRC Press. Analysis and improvement of plant cold hardiness. 1981. 1981. p. 139-159. 30 ref. (NAL Call No.: SB781.A52).

1360

Temperature and drought effects on blast and other characteristics in developing oats (Cultivars, environmental stresses, yield reductions).

Chinnici, M.F. AR-NC. Peterson, D.M. Madison, Wis., Crop Science Society of America. Crop science. Nov/Dec 1979. v. 19 (6). p. 893-897. ill. 29 ref. (NAL Call No.: 64.8 C883).

1361

Tolerance of spring-planted cereals to sulfonylurea herbicides.

Spinney, R.L. Appleby, A.P.; Brewster, B.D. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 257-259. (NAL Call No.: DNAL 79.9 W52R).

1362

Transpiration and soil water usage by oat as affected by postemergence herbicides.

AGJOAT. Beardmore, R.A. Linscott, D.L. Madison, Wis. : American Society of Agronomy. Changes in transpiration of plants as influenced by herbicides may or may not significantly affect soil water uptake in the field. We compared effects of the relatively new herbicides fluazifop (+/-)-butyl-2-(4- 5-(trifluoromethyl)-2-pyri-

dinyl oxy phenoxy)propanoate , haloxyfop methyl-2-(4- 3-chloro-5-(trifluoromethyl)-2--pyridinyl oxy phenoxy)propanoate , and sethoxydim (2- 1-(ethoxyimino)butyl -5-2 2-(ethylthio)propyl -3-hydroxy-2-cyclohexen-1-one applied at various rates to oat (*Avena sativa* L.) at the 5- to 6-cm stage on transpiration, soil water use, and plant development. In controlled environments, transpiration rates in oat declined about 6 d after herbicide treatment. Time for and the degree of transpiration reduction were herbicide and herbicide-rate dependent, as were the reductions in plant weight and leaf number. First reductions in soil moisture under laboratory conditions were found 10 d after treatment of oat with fluazifop and haloxyfop and 12 d after treatment with sethoxydim. Similarly, in the field fluazifop treatment required 9 d and haloxyfop and sethoxydim 11 d before significant differences in soil moisture occurred. The correlation coefficient of soil matric potential and dry weight of oat in the field was -0.89 and -0.90. respectively, for 2 yr, 1984 and 1985. Reduction of oat dry matter and soil water use in the field was also herbicide rate dependent. Agronomy journal. Nov/Dec 1988. v. 80 (6). p. 982-986. Includes references. (NAL Call No.: DNAL 4 AM34P).

1363

Triallate phytotoxicity and nitrogen fertilization (Oats).

McKercher, R.B. McGregor, W.R. Champaign, Ill., Weed Science Society of America. Weed science. Nov 1980. v. 28 (6). p. 740-744. ill. 10 ref. (NAL Call No.: 79.8 W41).

1364

Trifluralin and triallate retention by imbibed tame oat (*Avena sativa*) caryopses (Herbicides, phytotoxic effects, embryo culture, pericarp, testa).

Heath, M.C. Ashford, R.; McKercher, R.B. Champaign : Weed Science Society of America. Weed science. Mar 1984. v. 32 (2). p. 251-257. ill. Includes references. (NAL Call No.: 79.8 W41).

1365

Use of tonoplast and plasma membrane vesicles from oat root to investigate herbicidal disruption of proton gradients.

PCBPB. Ratterman, D.M. Balke, N.E. Duluth, Minn. : Academic Press. Pesticide biochemistry and physiology. May 1987. v. 28 (1). p. 17-28. Includes references. (NAL Call No.: DNAL SB951.P49).

(MISCELLANEOUS PLANT DISORDERS)

1366

Why cereals winterkill.

AGUDAT. Salmon, S.C. Madison, Wis. : American Society of Agronomy. Conclusions regarding the causes of winterkilling of cereal crops at this time would be decidedly premature. There can be scarcely any doubt that death occurs as a result of heaving of the soil, smothering, and direct effect of low temperature on the protoplasm. No doubt physiological drought causes injury and differences in resistance of certain cereals may perhaps be explained by their ability to absorb a larger quantity of water from the soil in proportion to the amount transpired. The duration and intensity of cold, rate of freezing, and in certain cases the rate of thawing and protection afforded by mulches, snow, and uneven surface of the ground are important factors. The moisture content of the tissue and its condition with respect to dormancy often have a determining influence. The following outline indicates the probable relation of the different factors. Causes of winterkilling: heaving; smothering; direct effect of low temperature (desiccation, chemical effect of cold, metabolism at low temperature); physiological drought. Conditions which modify the degree of injury: duration and intensity of cold; rate of freezing; rate of thawing; protection; kind of soil and moisture content of soil; habit of growth of plants; moisture content of tissue; dormancy; age of plants; concentration of sap; structure of tissue. Agronomy journal. Literature review. Nov 1917. v. 9 (8). p. 353-380. Includes references. (NAL Call No.: DNAL 4 AM34P).

1367

Wild mustard, *Sinapis arvensis*, control in common buckwheat, *Fagopyrum esculentum*, with desmedipham and fluorochloridone.

WETEE9. Friesen, G.H. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Apr 1988. v. 2 (2). p. 175-178. Includes references. (NAL Call No.: DNAL SB610.W39).

1368

Wound-induced lignin formation and resistance to cellulase in oat leaves (*Avena sativa*).

Geballe, G.T. PHYTA. Galston, A.W. St. Paul : American Phytopathological Society. Phytopathology. Apr 1983. v. 73 (4). p. 619-623. ill. Includes references. (NAL Call No.: 464.8 P56).

1369

Wound-induced resistance to cellulase in oat leaves (*Avena sativa*).

Geballe, G.T. Galston, A.W. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Sept 1982. v. 70 (3). p. 781-787. ill. 20 ref. (NAL Call No.: 450 P692).

1370

***Xanthomonas campestris* pv. *translucens* strains active in ice nucleation.**

PLDIDE. Kim, H.K. Orser, C.; Lindow, S.E.; Sands, D.C. St. Paul, Minn. : American Phytopathological Society. Plant disease. Nov 1987. v. 71 (11). p. 994-997. Includes references. (NAL Call No.: DNAL 1.9 P69P).

PROTECTION OF PLANT PRODUCTS - GENERAL AND MISC.

1371

Biochemical, functional, and nutritive changes during storage (Cereal grains).

Pomeranz, Y. St. Paul, Minn. : American Association of Cereal Chemists, 1982. Storage of cereal grains and their products, 3rd ed. / edited by Clyde M. Christensen. p. 145-217. 13 p. ref. (NAL Call No.: SB190.C43 1982).

1372

Buckwheat browning and color assessment.

CECHAF. Mazza, G. St. Paul, Minn. : American Association of Cereal Chemists. Cereal chemistry. July/Aug 1986. v. 63 (4). p. 361-364. Includes references. (NAL Call No.: DNAL 59.8 C33).

1373

Carbon-13 and proton nuclear magnetic resonance spectral assignments of deoxynivalenol and other mycotoxins from Fusarium graminearum (Contaminants of cereal crops).

Blackwell, B.A. Greenhalgh, R.; Bain, A.D. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. Sept/Oct 1984. v. 32 (6). p. 1078-1083. ill. Includes 35 references. (NAL Call No.: 381 J8223).

1374

The Cereal rusts /edited by William R. Bushnell and Alan P. Roelfs. --.

Bushnell, William R.; Roelfs, A. P. Orlando : Academic Press, c1984-1985. 2 v. : ill. ; 24 cm. Includes bibliographies and indexes. (NAL Call No.: DNAL SB741.R8C44).

1375

Field and storage fungi.

Christensen, C.M. New York : Von Nostrand Reinhold, 1987. Food and beverage mycology / edited by Larry R. Beuchat. p. 211-232. ill. Includes references. (NAL Call No.: DNAL QR115.F59 1987).

1376

Grain velocity measurement with optical sensors.

Chang, C.S. Martin, C.R.; Lai, F.S. St. Joseph, Mich. : The Society. American Society of Agricultural Engineers (Microfiche collection). Paper presented at the 1985 Winter Meeting of the American Society of Agricultural Engineers. Available for purchase from: The American Society of Agricultural Engineers, Order Dept., 2950 Niles Road, St. Joseph, Michigan 49085. Telephone the Order Dept. at (616) 429-0300 for information and prices. 1985. (fiche no.

85-3551). 22 p. ill. Includes references. (NAL Call No.: DNAL FICHE S-72).

1377

Measurement of thermally induced pressures in a model grain bin.

TAAEA. Manbeck, H.B. Muzzelo, L.M. St. Joseph, Mich. : The Society. Transactions of the ASAE - American Society of Agricultural Engineers. July/Aug 1985. v. 28 (4). p. 1253-1258, 1264. Includes references. (NAL Call No.: DNAL 290.9 AM32T).

1378

Microflora (Storage fungi, cereal grains and their products).

Christensen, C.M. Sauer, D.B. St. Paul, Minn. : American Association of Cereal Chemists, 1982. Storage of cereal grains and their products, 3rd ed. / edited by Clyde M. Christensen. p. 219-240. 1 p. ref. (NAL Call No.: SB190.C43 1982).

1379

Mutagenicity of selected chemicals in barley: repair and recovery (Storage of seeds treated with alkylating agents).

Veleminsky, J. Gichner, T. New York, Plenum Press. Environmental science research. 1982. v. 24. p. 321-327. 58 ref. (NAL Call No.: TD172.E55).

1380

Phosphine residue and its desorption from cereals.

JAFCAU. Rangaswamy, J.R. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. Nov/Dec 1985. v. 33 (6). p. 1102-1106. Includes references. (NAL Call No.: DNAL 381 J8223).

1381

Preservation and storage of grains, seeds, and their by-products cereals, oilseeds, pulses, and animal feed /edited by J.L. Multon ; preface by A.M. Reimbert ; translated from the French by D. Marsh ; reread by A.J. Eydt.

Multon, J. L. 1938-. New York, N.Y. : Lavoisier Pub., c1988. Translated from French.~ Translation of: Conservation et stockage des grains et graines et produits d'origine v'g'tale.~ Originally published: Lavoisier : Technique et documentation, 1982. xxiv, 1095 p. : ill. ; 25 cm. Includes bibliographies and index. (NAL Call No.: DNAL SB190.C6513).

(PROTECTION OF PLANT PRODUCTS - GENERAL AND MISC.)

1382

Programming and training for small farm grain storage /by Carl J. Lindblad.
Lindblad, Carl J. Washington, D.C. : Peace Corps, Information Collection and Exchange, 1981. "September 1981 ; September 1978.". iii, 100, 8, 1 p. : ill. ; 28 cm. Bibliography: p. 9 (3rd group). (NAL Call No.: DNAL SB190.L54).

1383

The rodent problem in Latin America.
Elias, D.J. Fall, M.W. Boca Raton, Fla. : CRC Press, c1988. Rodent pest management / editor, Ishwar Prakash. Literature review. p. 13-28. maps. Includes references. (NAL Call No.: DNAL SB994.R6R65).

1384

Rodents (Stored cereal grains).
Harris, K.L. Bauer, F.J. St. Paul, Minn. : American Association of Cereal Chemists, 1982. Storage of cereal grains and their products, 3rd ed. / edited by Clyde M. Christensen. p. 363-405. ill. 4 p. ref. (NAL Call No.: SB190.C43 1982).

1385

An uptake on mold inhibitors.
Hagler, W.M. Jr. College Park : The Conference. Proceedings - Maryland Nutrition Conference for Feed Manufacturers. 1983. p. 73-78. Includes references. (NAL Call No.: DNAL 389.9 UN342).

PROTECTION OF PLANT PRODUCTS - INSECTS

1386

Annotated list of the insects and mites associated with stored grain and cereal products, and of their arthropod parasites and predators /by R.T. Cotton and N.E. Good. Cotton, R. T. Good, Newell E._1905-. Washington, D.C. : U.S. Dept. of Agriculture, 1937. Caption title.~ Contribution from Bureau of Entomology and Plant Quarantine. 81 p. ; 23 cm. Bibliography: p. 63-74. (NAL Call No.: DNAL 1 Ag84M no.258).

1387

Another khapra beetle infestation confirmed by USDA in New Jersey (Pest of stored grains and cereal products). Washington, D.C., The Office. Major news releases and speeches - United States Department of Agriculture, Office of Governmental and Public Affairs. Apr 10/17, 1981. Apr 10/17, 1981. p. 20-21. (NAL Call No.: aS21.A8U51).

1388

Biological observations with *Zeteticontus* sp. (Hymenoptera: Encyrtidae) a parasite of *Carpophilus hemipterus* (L.) (Coleoptera: Nitidulidae) (Pest of dried fruits and stored cereals, natural and biological control, Israel). Gerling, D. Ben Mordechai, Y. Honolulu, The Society. Proceedings of the Hawaiian Entomological Society. Feb 1981. v. 23 (3). p. 351-354. 3 ref. (NAL Call No.: 420 H312).

1389

Cypermethrin and fenvalerate as grain protectants against *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Cyptolestes ferrugineus* (Coleoptera: Cucujidae) at different moisture levels and temperatures. JEENAI. Joia, B.S. Loschiavo, S.R.; Webster, G.R.B. College Park, Md. : Entomological Society of America. Journal of economic entomology. June 1985. v. 78 (3). p. 637-641. Includes references. (NAL Call No.: DNAL 421 J822).

1390

Defining the stored barley insect problem. Harein, P.K. Milwaukee, Wis.? : The Association?, 1985? . Proceedings, Barley Insect Conference : January 9, 1985, Minneapolis Plaza Hotel, Minneapolis, Minnesota / sponsored by American Malting Barley Association. p. 1-6. (NAL Call No.: DNAL SB608.B2B2 1985).

1391

Effect of oat constituents on aggregation behavior of *Oryzaephilus surinamensis* (L.) (Sawtoothed grain beetle). Mikolajczak, K.L.JAFCA. Freedman, B.; Zilkowski, B.W.; Smith, C.R. Jr.; Burkholder, W.E. Washington : American Chemical Society. Journal of agricultural and food chemistry. Jan/Feb 1983. v. 31 (1). p. 30-33. 1 p. ref. (NAL Call No.: 381 J8223).

1392

Effects of continuous and fractionated doses of gamma radiation on the survival and fertility of *Sitophilus granarius* (L.). Jefferies, D.J. New York : Pergamon Press, 1966. The Entomology of radiation disinfection of grain : a collection of original research papers / edited by P.B. Cornwell. p. 41-56. Includes references. (NAL Call No.: DNAL SB608.G6C6).

1393

Effects of 2-tridecanone and analogues on the reproduction and mortality of stored product insects. JKESA. Kramer, K.J. Beeman, R.W.; Speirs, W.E.; Burkholder, W.E.; McGovern, T.P. Lawrence, Kan. : The Society. Journal of the Kansas Entomological Society. Apr 1985. v. 58 (2). p. 254-260. Includes references. (NAL Call No.: DNAL 420 K13).

1394

Energy budget for *Oryzaephilus surinamensis* (Coleoptera: Cucujidae) feeding on rolled oats. White, N.D.G. Sinha, R.N. College Park, Md., Entomological Society of America. Environmental entomology. June 1981. v. 10 (3). p. 320-326. 111. 19 ref. (NAL Call No.: QL461.E532).

1395

Evaluation of *Bacillus thuringiensis* for controlling indianmeal moths (Lepidoptera: Pyralidae) in farm grain bins and elevator silos. JEENAI. McGaughey, W.H. College Park, Md. : Entomological Society of America. Journal of economic entomology. Oct 1985. v. 78 (5). p. 1089-1094. Includes references. (NAL Call No.: DNAL 421 J822).

(PROTECTION OF PLANT PRODUCTS - INSECTS)

1396

Evaluation of fenoxycarb, *Bacillus thuringiensis*, and malathion as grain protectants in small bins.

JEENAI. Kramer, K.J. Hendricks, L.H.; Wojciak, J.H.; Flyer, J. College Park, Md. : Entomological Society of America. Journal of economic entomology. June 1985. v. 78 (3). p. 632-636. ill. Includes references. (NAL Call No.: DNAL 421 J822).

1397

Factors affecting survival of the merchant grain beetle (Coleoptera: Cucujidae) and the confused flour beetle (Coleoptera: Tenebrionidae) exposed to silica aerogel.

JEENAI. White, N.D.G. Loschiavo, S.R. Lanham, Md. : Entomological Society of America. Mortality of *Oryzaephilus mercator* Fauvel exposed to 0.72 mg/cm² silica aerogel (SG-67) bound to paper was directly related to length of exposure of the insects and length of time between exposure and placement in food. Exposures of 15 s killed 97% of starved adults in 2 d, but mortality was low if adults had access to food within 2 h. As exposure times increased, insects survived only if food was available at shorter intervals. Type and quantity of food also affected mortality. Mortality of *O. mercator* adults exposed to SG-67 was greater on bread crumbs and whole rolled oats than on ground rolled oats. Starved *Tribolium confusum* Jacquelin du Val required exposures of 6 h to SG-67 for complete mortality; the presence of food following exposure, rather than type or quantity, affected survival, although mortality was slightly higher on bread crumbs. Early larval instars of *O. mercator* and *T. confusum* were more susceptible than late instars, which responded to SG-67 in the same way as adults. Adults of both species exposed to SG-67 and subsequently held in inert cellulose powder died within 1-3 d, but many insects in ground rolled oats survived, probably because they had ingested and metabolized food and produced metabolic water. Adult *O. mercator* immobilized at 2.5 degrees C during exposure to SG-67 for 4 h, then held at 25 degrees C without food, were unaffected by the treatment. Exposures of *O. mercator* for 15-60 s to silica aerogel on an adhesive surface will effectively control these insects if they cannot find food immediately. *T. confusum* cannot be controlled with exposure times of less than 6 h. Journal of economic entomology. June 1989. v. 82 (3). p. 960-969. ill. Includes references. (NAL Call No.: DNAL 421 J822).

1398

Factors influencing stored-oat insect populations in South Dakota.

JEENAI. Ingemansen, J.A. Reeves, D.L.; Walstrom, R.J. College Park, Md. : Entomological Society of America. Journal of economic entomology. Apr 1986. v. 79 (2). p. 518-522. Includes references. (NAL Call No.:

DNAL 421 J822).

1399

Grain pesticidesAmerican Association of Cereal Chemists. --.

Washington, D.C.? : The Association, 1985. Title from container. 4 videocassettes (362 min.) : sd., col. ; 1/2 in. (NAL Call No.: DNAL Videocassette no.21).

1400

Heat for control of cereal insects / W.H. Goodwin .

Goodwin, W. H. 1880-. Wooster, Ohio : Ohio Agricultural Experiment Station, 1922. Cover title. 18 p. ; 23 cm. (NAL Call No.: DNAL 100 OH3S (2) no.354).

1401

In-transit fumigation of truck-ship containers with hydrogen phosphide--a feasibility study (Insect control, stored product pests, cereals).

Jay, E. Davis, R.; Zehner, J.M. New Orleans : The Region. Advances in agricultural technology. AAT-S - United States, Dept. of Agriculture, Agricultural Research Service, Southern Region. Apr 1983. Apr 1983. (28). 13 p. Includes references. (NAL Call No.: aS21.A75U7).

1402

In vitro effect of bean amylase inhibitor on insect amylases (Cereals, insect pest control).

Powers, J.R. Culbertson, J.D. Ames, Iowa, International Association of Milk, Food, and Environmental Sanitarians. Journal of food protection. May 1982. v. 45 (7). p. 655-657. ill. Includes 14 ref. (NAL Call No.: 44.8 J824).

1403

The influence of starvation on mortality, development, and protein content in parasitized and unparasitized *Tribolium castaneum*.

JIVPA. Armstrong, E. Newton, P.B. New York, N.Y. : Academic Press. Journal of invertebrate pathology. July 1985. v. 46 (1). p. 103-108. Includes references. (NAL Call No.: DNAL 421 J826).

(PROTECTION OF PLANT PRODUCTS - INSECTS)

1404

Insect control in stored grain and peas and seed treatment for small grains.
WA. Pullman, Wash., The Service. EM - Cooperative Extension Service, Washington State University. Washington State University. Cooperative Extension Service. Feb 1980. Feb 1980. (3314). 7 p. (NAL Call No.: 275.29 W27MI).

1405

Insect infestation in grain loaded in railroad cars at primary elevators in Southern Manitoba, Canada.
JEENAI. Smith, L.B. College Park, Md. : Entomological Society of America. Journal of economic entomology. June 1985. v. 78 (3). p. 531-534. ill. Includes references. (NAL Call No.: DNAL 421 J822).

1406

Insects infesting barley stored on farms in Minnesota.
JEENAI. Subramanyam, B. Harein, P.K. Lanham, Md. : Entomological Society of America. Barley stored on farms in Minnesota in 12 storage facilities during 1985 and 17 facilities during 1986 was sampled for adults of insect species. Grain samples were removed with grain probing devices or a plastic scoop, and live insects were trapped with unbaited plastic probe traps. Analysis of grain and probe trap samples indicated that, on average, about 85.0 to 100.0% of the facilities holding barley were infested with more than one insect species. The insect species most commonly occurring in grain or probe trap samples were *Cryptolestes* spp.; red flour beetle, *Tribolium castaneum* (Herbst); sawtoothed grain beetle, *Oryzaephilus surinamensis* (L.); foreign grain beetle, *Ahasverus advena* (Waltl); lesser grain borer, *Rhyzopertha dominica* (F.); and hairy fungus beetle, *Typhaea stercorea* (L.). *Cryptolestes* spp. and *O. surinamensis* were the most abundant insect species. Indianmeal moth, *Plodia interpunctella* (Hubner), larvae or adults were rarely found in the grain or probe trap samples, although adult males were captured in pheromone traps that were placed above the grain. Temperature of the grain in July and August 1985 and grain samples between May and September 1986 was within the range (15.6 to 35.0 degrees C) conducive for insect survival and reproduction. Moisture content of the grain samples during 1985 ranged from 12.0 to 16.0% and from 11.0 to 20.0% in 1986. These moisture levels were adequate to support insect infestations. Moisture content of the grain samples during 1985 and 1986 decreased by approximately 0.1 and 0.2%, respectively, for every 1 degrees C rise in grain temperature between 7 and 35 degrees C. Based on test weight of the grain samples, 5 of 12 facilities during 1985 and 3 of 17 facilities during 1986 held poor quality barley (U.S. Sample Grade). The amount of dockage in the grain samples ranged from 0.0 to 4.6%. Comparison of probe

traps and grain sampling devices indicated that probe traps were more sensitive than a deep-bin cup probe. a. Journal of economic entomology. Dec 1989. v. 82 (6). p. 1817-1824. Includes references. (NAL Call No.: DNAL 421 J822).

1407

Interspecific sexual attraction between *Pyralis farinalis* L. and *Amyelois transitella* (Walker) (Lepidoptera: Pyralidae) (Minor stored grain pest, ecology).
Landolt, P.J. Curtis, C.E. Lawrence, Kan., The Society. Journal of the Kansas Entomological Society. Apr 1982. v. 55 (2). p. 248-252. ill. Includes 9 ref. (NAL Call No.: 420 K13).

1408

Irradiation of grain and grain products for insect control / prepared by Elvin M. Tilton and Stuart O. Nelson . --.
Tilton, Elvin W. Nelson, Stuart Owen, 1927-. Ames, Iowa : Council for Agricultural Science and Technology, 1984. "April 1984" - cover. 6 p. ; 28 cm. --. Bibliography: p. 5-6. (NAL Call No.: DNAL TP371.8.T55).

1409

Keep oats clean.
SDFHA. Ingemansen, J. Brookings, S.D. : The Station. South Dakota farm and home research - South Dakota, Agricultural Experiment Station. 1986. v. 37 (2). p. 3-5. ill. (NAL Call No.: DNAL 100 S0825).

1410

Laboratory studies on several plant materials as insect repellants for protection of cereal grains (Turmeric, *Curcuma longa*, neem, *Azadirachta indica*, fenugreek, *Trigonella foenum-graceum*, *Tribolium castaneum*, *Trogoderma granarium*, *Rhyzopertha dominica*).
Jilani, G. Su, H.C.F. College Park, Md. : Entomological Society of America. Journal of economic entomology. Feb 1983. v. 76 (1). p. 154-157. Includes references. (NAL Call No.: 421 J822).

1411

Management of stored barley in Minnesota: practices versus recommendations.
Gardner, R.D. Harein, P.K.; Subramanyam, B. College Park, Md. : The Society. Bulletin of the Entomological Society of America. Spring 1988. v. 34 (1). p. 22-26. Includes references. (NAL Call No.: DNAL 423.9 EN8).

(PROTECTION OF PLANT PRODUCTS - INSECTS)

1412

Modified atmosphere technology.

Storey, C.L. Milwaukee, Wis.? : The Association?, 1985? . Proceedings, Barley Insect Conference : January 9, 1985, Minneapolis Plaza Hotel, Minneapolis, Minnesota / sponsored by American Malting Barley Association. p. 14-21. (NAL Call No.: DNAL SB608.B2B2 1985).

1413

Moth control in stored grain and the role of *Bacillus thuringiensis*: an overview.

RREVA. Subramanyam, B. Cutkomp, L.K. New York, N.Y. : Springer. Residue reviews. Literature review. 1985. v. 94. p. 1-47. ill. Includes references. (NAL Call No.: DNAL 389.9 R314).

1414

Organophosphate resistance in adults of red flour beetle (Coleoptera: Tenebrionidae) and sawtoothed grain beetle (Coleoptera: Cucujidae) infesting barley stored on farms in Minnesota. JEENAI. Subramanyam, B. Harein, P.K.; Cutkomp, L.K. Lanham, Md. : Entomological Society of America. Four strains of adult red flour beetles, *Tribolium castaneum* (Herbst), and six strains of adult sawtoothed grain beetles, *Oryzaephilus surinamensis* (L.), infesting barley stored on farms in Minnesota were surveyed for resistance to malathion, pirimiphos-methyl, and chlorpyrifos-methyl. Adults were exposed to filter papers impregnated with each of the insecticides. Dose-response and discriminating-dose test procedures were used to evaluate pirimiphos-methyl and chlorpyrifos-methyl resistance. Dose-response and discriminating-dose tests indicated malathion resistance in all field strains of *T. castaneum*, but not in *O. surinamensis* field strains. *T. castaneum* strains did not show cross-resistance to pirimiphos-methyl or chlorpyrifos-methyl. All six *O. surinamensis* strains were susceptible to pirimiphos-methyl, but four strains showed reduced susceptibility (tolerance) to chlorpyrifos-methyl. The use of synergists in combination with malathion on *T. castaneum* strains and with chlorpyrifos-methyl on the four *O. surinamensis* strains indicated that carboxylesterase was involved in detoxifying malathion in resistant *T. castaneum* adults, whereas oxidases and esterases other than carboxylesterase were involved in detoxifying chlorpyrifos-methyl in tolerant *O. surinamensis* adults. Journal of economic entomology. Aug 1989. v. 82 (4). p. 989-995. Includes references. (NAL Call No.: DNAL 421 J822).

1415

Potential of ionizing radiation for insect control in the cereal food industry.

Watters, F.L. St. Paul, Minn. : American Association of Cereal Chemists, c1984. Insect management for food storage and processing / edited by Fred J. Baur. Literature review. p. 267-278. Includes references. (NAL Call No.: DNAL SB937.I49).

1416

Preliminary infectivity tests using six bacterial formulations against the red flour beetle, *Tribolium castaneum*.

JIVPA. Kumari, S.M. Neelgund, Y.F. New York, N.Y. : Academic Press. Journal of invertebrate pathology. Sept 1985. v. 46 (2). p. 198-199. (NAL Call No.: DNAL 421 J826).

1417

Presence of four species of stored-product moths in storage and field situations in north-central Florida as determined with sex pheromone-baited traps.

FETMA. Vick, K.W. Coffelt, J.A.; Weaver, W.A. Gainesville, Fla. : Florida Entomological Society. Florida entomologist. Dec 1987. v. 70 (4). p. 488-492. ill. Includes references. (NAL Call No.: DNAL 420 F662).

1418

Production dynamics of cucujolide pheromones and identification of 1-octen-3-ol as a new aggregation pheromone for *Oryzaephilus surinamensis* and *O. mercator* (Coleoptera: Cucujidae).

EVETEX. Pierce, A.M. Pierce, H.D. Jr.; Borden, J.H.; Oehlschlager, A.C. Lanham, Md. : Entomological Society of America. Aggregation pheromone production by *Oryzaephilus surinamensis* (L.) and *O. mercator* (Fauvel) was measured as a function of adult age and population density. Porapak Q-captured volatiles were monitored weekly from mixed-sex adults feeding on rolled oats at two different population densities. Macrolide aggregation pheromones (cucujolides) were produced by males of both species by the end of the first week after eclosion and maximum production rates were still maintained more than 3 mo later. For *O. mercator*, the previously identified cucujolides were 3(Z),11(R)-dodecen-11-olide and 3(Z),6(Z),11(R)-dodecadien-11-olide, and for *O. surinamensis*, 3(Z),6(Z),11(R)-dodecadien-11-olide, 3(Z),6(Z)-dodecadienolide, and 5(Z),8(Z),13(R)-tetradecadien-13-olide. *O. surinamensis* less than 1 mo after eclosion showed a two- to threefold increase in cucujolide production compared with adults maintained at a fivefold higher population density. In comparison, cucujolide production by young *O. mercator* was less sensitive to the presence of conspecifics. At the lower

(PROTECTION OF PLANT PRODUCTS - INSECTS)

population density, adults of both species more than 1 mo after eclosion produced almost optically pure (R)-(-)-1-octen-3-ol. Laboratory bioassays in a two-choice, pitfall olfactometer indicated that 1-octen-3-ol alone serves as an aggregation pheromone, and when added to mixtures of cucujolide pheromones, it extends the attractive range of the cucujolides for both *Dryzaephilus* species. *Environmental entomology*. Oct 1989. v. 18 (5). p. 747-755. Includes references. (NAL Call No.: DNAL QL461.E532).

1419

Purification, partial characterization, and postembryonic levels of amylases from *Sitophilus oryzae* and *Sitophilus granarius*.
Baker, J.E. Woo, S.M. New York, N.Y. : Alan R. Liss, Inc. Archives of insect biochemistry and physiology. 1985. v. 2 (4). p. 415-428. ill. Includes 37 references. (NAL Call No.: DNAL QL495.A7).

1420

Safe storage of small amounts of grains, edible seeds and processed foods.
Wallen, Stanley E. Kamble, Shripat T.; Gold, Roger E.; Roselle, Robert E.; Walker, Charles E. Document available from: University of Nebraska-Lincoln, Dept. of Agricultural Communications, Lincoln, Nebraska 68583 1983. Describes how to properly store these foods to ensure quality, including types of containers to use and how to destroy or repel insects. 4 p. (NAL Call No.: Document available from source.). (NAL Call No.: G83-642).

1421

A sealed paper carton to protect cereals from insect attack /by William B. Parker.
Parker, William B. 1885-. Washington, D.C. : U.S. Dept. of Agriculture, 1913. Caption title. ~ "October 16, 1913.". 8 p. : ill. ; 24 cm. (NAL Call No.: DNAL 1 Ag84B no.15).

1422

The search for a safe, effective spot fumigant for use in breweries.
Klimovitz, R.U. Milwaukee, Wis.? : The Association?, 1985? . Proceedings, Barley Insect Conference : January 9, 1985, Minneapolis Plaza Hotel, Minneapolis, Minnesota / sponsored by American Malting Barley Association. p. 11-13. (NAL Call No.: DNAL SB608.B2B2 1985).

1423

Stored-grain insect management in Montana.
Morrill, W.L. Lohr, K.; Jensen, G. Bozeman, Mont. : The Service. Montguide MT : Agriculture - Montana State University, Cooperative Extension Service. Oct 1985. (8528). 4 p. ill. Includes references. (NAL Call No.: DNAL S544.3.M9M65).

1424

Superheating as a control for cereal-mill insects /by J.H. Pepper and A.L. Strand.
Pepper, J. H. 1902-. Strand, A. L. 1894-. Bozeman, Mont. : Montana State College, Agricultural Experiment Station, 1935. 26 p. ; 23 cm. (NAL Call No.: DNAL 100 M76 (1) no.297).

1425

Testing for malathion resistance in field-collected populations of *Cryptolestes ferrugineus* (Stephens) and factors affecting reliability of the tests.
JEENAI. White, N.D.G. Loschiavo, S.R. College Park, Md. : Entomological Society of America. Journal of economic entomology. June 1985. v. 78 (3). p. 511-515. Includes references. (NAL Call No.: DNAL 421 J822).

1426

Tricalcium phosphate-soybean oil in fortified processed cereals to suppress insects, dusting, and separation.
JFDAZ. Bookwalter, G.N. Highland, H.A.; Warner, K. Chicago, Ill. : Institute of Food Technologists. Journal of food science. Jan/Feb 1985. v. 50 (1). p. 245-248. Includes references. (NAL Call No.: DNAL 389.8 F7322).

1427

Understanding some insect pests of cereal grains...Hessian fly.
Morrill, W.L. Bozeman : The Station. Capsule information series - Montana Agricultural Experiment Station. June 1983. June 1983. (31). 2 p. ill. (NAL Call No.: S83.M6).

1428

Update on the use and status of methyl bromide for barley insect control.
White, L.V. Milwaukee, Wis.? : The Association?, 1985? . Proceedings, Barley Insect Conference : January 9, 1985, Minneapolis Plaza Hotel, Minneapolis, Minnesota / sponsored by American Malting Barley Association. p. 7-8. (NAL Call No.: DNAL SB608.B2B2 1985).

(PROTECTION OF PLANT PRODUCTS - INSECTS)

1429

Update on the use and status of phosphine products for barley insect control.

Wilbur, D.A. Milwaukee, Wis.? : The Association?, 1985? . Proceedings, Barley Insect Conference : January 9, 1985, Minneapolis Plaza Hotel, Minneapolis, Minnesota / sponsored by American Malting Barley Association. p. 9-10. (NAL Call No.: DNAL SB608.B2B2 1985).

1430

Uptake of malathion from galvanized-steel surfaces by stored barley.

White, N.D.G. Abramson, D. College Park, Md. : Entomological Society of America. Journal of economic entomology. Apr 1984. v. 77 (2). p. 289-293. Includes references. (NAL Call No.: 421 J822).

WEEDS

1431

Annual Grass control in flax (*Linum usitatissimum*) with quizalofop. WETEE9. Friesen, G.H. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Apr 1988. v. 2 (2). p. 144-146. Includes references. (NAL Call No.: DNAL SB610.W39).

1432

The antagonistic action of 2,4-D and bromoxynil on glyphosate phytotoxicity to barley (*Hordeum vulgare*) (Herbicide mixtures). O'Donovan J.T. O'Sullivan, P.A. Champaign, Ill., Weed Science Society of America. Weed science. Jan 1982. v. 30 (1). p. 30-34. Includes 6 ref. (NAL Call No.: 79.8 W41).

1433

Antagonistic effects of MCPA on wild oat (*Avena fatua*) control with diclofop. Olson, W.A. Nalewaja, J.D. Champaign, Ill., Weed Science Society of America. Weed science. Sept 1981. v. 29 (5). p. 566-571. ill. 16 ref. (NAL Call No.: 79.8 W41).

1434

Application of herbicides in small grain stubble (with the boom mounted on a combine and incorporation with a rolling cultivator for weed control in soybeans). Barker, G.L.TAAEA. Buehring, N.W.; Whisler, F.D.; Colwick, R.F. St. Joseph : The Society. Transactions of the ASAE - American Society of Agricultural Engineers. Sept/Oct 1982. v. 25 (5). p. 1232-1236. ill. 11 ref. (NAL Call No.: 290.9 AM32T).

1435

Barban for control of wild oats in cereal grains including limited studies with Diallate / Chester L. Foy and David E. Bayer. Foy, Chester L. Bayer, David E. Berkeley, Cal. : California Agricultural Experiment Station, 1966. Cover title. 29 p. : ill. ; 23 cm. Bibliography: p. 27-29. (NAL Call No.: DNAL 100 C12S no.828).

1436

Barley no-till weed management trial Riverside county: 1986. Cudney, D.W. Baameur, A. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 280-282. (NAL Call No.: DNAL 79.9 W52R).

1437

The basis for synergism between barban and flamprop on wild oat (*Avena fatua*) (weed of cereal and oilseed crops in the prairie provinces of Canada, Herbicide combinations). Sharma, M.P. Qureshi, F.A.; Vandenborn, W.H. Champaign, Ill., Weed Science Society of America. Weed science. Mar 1982. v. 30 (2). p. 147-152. ill. Includes 10 ref. (NAL Call No.: 79.8 W41).

1438

The basis of the antagonistic effect of 2,4-D on diclofop-methyl toxicity to wild oat (*Avena fatua*). Todd, B.G. Stobbe, E.H. Champaign, Ill., Weed Science Society of America. Weed science. July 1980. v. 28 (4). p. 371-377. ill. 18 ref. (NAL Call No.: 79.8 W41).

1439

BAY FDE 3440 for selective wild oat control in spring barley. Rydrych, D.J. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 271. (NAL Call No.: DNAL 79.9 W52R).

1440

The biologic and economic assessment of Diallate; a report of the Diallate assessment team to the rebuttable presumption against registration of Diallate (Herbicide for the control of wild oats). USDA. Washington, D.C., The Department. Technical bulletin. United States. Dept. of Agriculture. Sept 12, 1977. Sept 12, 1977. (1620). 24 p. 28 ref. (NAL Call No.: 1 AG84TE).

1441

Breeding cereal grains for resistance to witchweed. Ramaiah, K.V. Boca Raton, Fla. : CRC Press, c1987-. Parasitic weeds in agriculture / editor, Lytton J. Musselman. Literature review. p. 227-242. Includes references. (NAL Call No.: DNAL SB611.P34).

1442

Broadleaf weed control barley. WAEB. Miller, S.D. Lauer, J. Laramie : The Station. Bulletin B - Wyoming, Agricultural Experiment Station. 1986. (885). p. 140-141. (NAL Call No.: DNAL 100 W99 (1)).

(WEEDS)

1443

Broadleaf weed control in barley.

Miller, S.D. Lauer, J. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 208-209. (NAL Call No.: DNAL 79.9 W52R).

1444

Broadleaf weed control in barley.

Miller, S.D. Lauer, J. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 264-265. (NAL Call No.: DNAL 79.9 W52R).

1445

Broadleaf weed control in barley with clopyralid and fluroxypyr alone or in combinations.

WAEBA. Miller, S.D. Krall, J.M. Laramie : The Station. Bulletin B - Wyoming, Agricultural Experiment Station. 1986. (885). p. 148-149. (NAL Call No.: DNAL 100 W99 (1)).

1446

Broadleaf weed control in barley with clopyralid and fluroxypyr alone or in combinations.

Miller, S.D. Krall, J.M. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 260-261. (NAL Call No.: DNAL 79.9 W52R).

1447

Broadleaf weed control in barley with postemergence herbicide treatments.

Miller, S.D. Krall, J.M. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 214-215. (NAL Call No.: DNAL 79.9 W52R).

1448

Broadleaf weed control in barley with sulfonyl urea herbicides.

Miller, S.D. Krall, J.M. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 213. (NAL Call No.: DNAL 79.9 W52R).

1449

Broadleaf weed control in barley with sulfonyl urea herbicides.

WAEBA. Miller, S.D. Krall, J.M. Laramie : The Station. Bulletin B - Wyoming, Agricultural Experiment Station. 1986. (885). p. 150-151. (NAL Call No.: DNAL 100 W99 (1)).

1450

Broadleaf weed control in barley with sulfonyl urea herbicides.

Miller, S.D. Krall, J.M. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 255-256. (NAL Call No.: DNAL 79.9 W52R).

1451

Broadleaf weed control in spring barley at Bonners Ferry, Idaho.

Zamora, D.L. Thill, D.C.; Callihan, R.H. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 194-195. (NAL Call No.: DNAL 79.9 W52R).

1452

Broadleaf weed control in spring barley at Pottlatch, Idaho.

Swensen, J.B. Thill, D.C.; Callihan, R.C. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 191-193. (NAL Call No.: DNAL 79.9 W52R).

1453

Broadleaf weed control in spring barley in Fremont county.

Dial, M.J. Thill, D.C. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 191-192. (NAL Call No.: DNAL 79.9 W52R).

1454

Broadleaf weed control in spring barley in Fremont County.

Dial, M.J. Lish, J.M.; Thill, D.C.; Herrman, T.J. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 268-269. (NAL Call No.: DNAL 79.9 W52R).

1455

Canada thistle control in barley.

Miller, S.D. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 270. (NAL Call No.: DNAL 79.9 W52R).

1456

Canada thistle control in spring barley.

Lish, J.M. Thill, D.C. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 189. (NAL Call No.: DNAL 79.9 W52R).

1457

Cell membrane permeability and ultrastructural effects of difenzoquat on wild oats (*Avena fatua*).

WEESA6. Thai, K.M. Jana, S.; Fowke, L.C. Champaign, Ill. : Weed Science Society of America. Effects of difenzoquat on wild oats grown under controlled environmental conditions were studied. Seedling height and fresh weight were significantly reduced 5 days after postemergence treatment. Dose-dependent increase in cell membrane permeability was detected after a 12-h exposure to the herbicide. Scanning electron micrography showed normal leaf hairs and cuticular wax but swollen guard cells 10 days after treatment. Ultrastructural changes occurred before the visible symptoms. The primary effect of difenzoquat appears to be the disruption of the tonoplast and plasmalemma. The tonoplast showed greater damage than the plasmalemma. Secondary effects included damage to mitochondria and chloroplasts. Mitochondria were swollen and often ruptured, but the effect did not increase with the duration of exposure. Chloroplasts became spherical in shape, and their contents were also affected. The changes included: accumulation and then disappearance of starch granules, swelling of frets, fusion of granal thylakoids, detachment and then rupture of the outer membrane of the envelope, and clumping of ribosomes. By contrast, natural senescence caused greater injury of the plasmalemma than the tonoplast, a marked increase in size of plastoglobuli, and loss of starch grains without early accumulation. Weed science. Jan 1989. v. 37 (1). p. 98-106. ill. Includes references. (NAL Call No.: DNAL 79.8 W41).

1458

Changes in composition of chlorophylls, carotenoids, and prenylquinones in green seedlings of *Hordeum* (barley) and *Raphanus* (radishes) induced by the herbicide San 6706--an effect possibly antagonistic to phytochrome action.

Kleudgen, H.K. New York, Academic Press. Pesticide biochemistry and physiology. Dec 1979. v. 12 (3). p. 231-238. ill. 36 ref. (NAL Call No.: SB951.P49).

1459

The characteristics of secondary dormant wild oat (*Avena fatua* L.) seed.

WSWPA. Fellows, G.M. Fay, P.K.; Foley, M.E. Reno : The Society. Proceedings - Western Society of Weed Science. 1985. v. 38. p. 105-107. (NAL Call No.: DNAL 79.9 W52).

1460

Chemical weed control in small grain and forages: 1985.

Wrage, L.J. Arnold, W.E.; Johnson, P.D. Brookings, S.D. : The Service. FS - South Dakota State University, Cooperative Extension Service. Jan 1985. (525A). 14 p. (NAL Call No.: DNAL 275.29 S085FS).

1461

Chemical weed control in small grains.

Bullock, F.D. Athens, Ga. : The Service. Circular - Cooperative Extension Service, University of Georgia. Oct 1985. (765,rev.). 8 p. ill. (NAL Call No.: DNAL 275.29 G29C).

1462

Chlorsulfuron reduced control of wild oat (*Avena fatua*) with diclofop, difenzoquat, and flamprop (Herbicide mixtures, antagonism).

O'Sullivan, P.A. Kirkland, K.J. Champaign, Ill. : Weed Science Society of America. Weed science. May 1984. v. 32 (3). p. 285-289. Includes references. (NAL Call No.: 79.8 W41).

1463

The combinations of chlorsulfuron and metasulfuron with AC 222,293 at various rates to determine an effective rate for broad spectrum weed control.

Stewart, V.R. Keener, T.K. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 216-218. (NAL Call No.: DNAL 79.9 W52R).

1464

Common buckwheat (*Fagopyrum esculentum*) tolerance to herbicides.

WEESA6. Friesen, G.H. Campbell, C.G. Champaign, Ill. : Weed Science Society of America. Weed science. May 1986. v. 34 (3). p. 435-439. Includes references. (NAL Call No.: DNAL 79.8 W41).

(WEEDS)

1465

Comparison of chlorophyll fluorescence and fresh weight as herbicide bioassay techniques. WEESA6. Shaw, D.R. Peeper, T.F.; Nofziger, D.L. Champaign, Ill. : Weed Science Society of America. Weed science. Jan 1985. v. 33 (1). p. 29-33. Includes 18 references. (NAL Call No.: DNAL 79.8 W41).

1466

Competitive effects of wild barley in seedling alfalfa. Cudney, D.W. Orloff, S.B. S.1. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 115. (NAL Call No.: DNAL 79.9 W52R).

1467

Control of foxtail barley (*Hordeum jubatum* L.) in perennial grass pastures. Ferrell, M.A. Whitson, T.D. S.1. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 31. (NAL Call No.: DNAL 79.9 W52R).

1468

Control of Jerusalem artichoke (*Helianthus tuberosus*) in Barley (*Hordeum vulgare*). WEESA6. Wall, D.A. Kiehn, F.A.; Friesen, G.H. Champaign, Ill. : Weed Science Society of America. Weed science. Sept 1986. v. 34 (5). p. 761-764. Includes references. (NAL Call No.: DNAL 79.8 W41).

1469

Control of volunteer barley in seedling alfalfa--1987. Orloff, S.B. Cudney, D. S.1. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 174-175. (NAL Call No.: DNAL 79.9 W52R).

1470

Control of weeds in an oat (*Avena sativa*)-soybean (*Glycine max*) ecofarming rotation (Herbicides and reduced tillage). Burnside, O.C. Wicks, G.A.; Carlson, D.R. Champaign, Ill., Weed Science Society of America. Weed science. Jan 1980. v. 28 (1). p. 46-50. ill. 15 ref. (NAL Call No.: 79.8 W41).

1471

Controlling wild oat on roadsides. SWSPB. McCully, W.G. Wiese, A.F. Champaign : The Society. Proceedings - Southern Weed Science Society. Jan 17-19, 1984. (37th). p. 287-289. Includes 4 references. (NAL Call No.: DNAL 79.9 S08).

1472

Days after seeding and application of diclofop for wild oat control. Dial, M.J. Thill, D.C.; Yenne, S.P. S.1. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 275-276. (NAL Call No.: DNAL 79.9 W52R).

1473

Diclofop (Hoelon) for wild oat (*Avena fatua*) and green foxtail (*Setaria viridis*) control. Miller, S.D. ND. Nalewaja, J.D. Fargo, N.D., The Station. North Dakota farm research - North Dakota, Agricultural Experiment Station. Sept/Oct 1980. v. 38 (2). p. 19-22. 5 ref. (NAL Call No.: 100 N813B).

1474

DPX-Y6202 for wild oats control in lentils. Curran, W.S. Whitesides, R.E.; Morrow, L.A. S.1. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 161-162. (NAL Call No.: DNAL 79.9 W52R).

1475

The effect of an oat companion crop on weeds in seedling alfalfa. Lanini, W.T. Orloff, S.B.; Vargas, R.N.; Marble, V.L.; Orr, J.; Grattan, S.R. Sacramento, Calif. : California Weed Conference Office. Proceedings - California Weed Conference. Paper presented at a conference on "Education and Communication--the Keys to the Future," January 18-21, 1988, Sacramento, California. 1988. (40). p. 79-89. Includes references. (NAL Call No.: DNAL 79.9 C122).

1476

Effect of application time on soil residue and efficacy of sulfonylureas. SWSPBE. Foy, C.L. Mersie, W. Raleigh, N.C. : The Society. Proceedings - Southern Weed Science Society. 1986. (39th). p. 446-456. Includes references. (NAL Call No.: DNAL 79.9 S08 (P)).

1477

Effect of environment and adjuvants on asulam phytotoxicity.

WEESA6. Nalewaja, J.D. Woznica, Z. Champaign, Ill. : Weed Science Society of America.

Abstract: Experiments were conducted to determine the influence of various factors on asulam

methyl (4-amino-phenyl)sulfonyl carbamate toxicity to flax (*Linum usitatissimum* L.) and wild oats (*Avena fatua* L. ~ AVEFA). Asulam toxicity to both flax and wild oats generally increased as temperature, humidity, and soil moisture increased after treatment. Octoxynol (alpha- p-1,1,3,3-tetramethyl butyl phenyl -omega-hydroxypoly(oxyethylene) in the spray solution increased asulam toxicity to both species in all environments. Octoxynol and trimethylenonypolyethoxyethanol (WK) enhanced asulam toxicity more than other adjuvants evaluated. Asulam toxicity to both flax and wild oats increased as octoxynol concentration in the spray increased. Flax tolerance to asulam generally increased with flax height at treatment. 'Flor' flax was the most asulam susceptible of six cultivars evaluated. A 2-mm simulated rainfall within 3 or 6 h after asulam treatment reduced toxicity to wild oats and flax, respectively. Weed science. May 1988. v. 36 (3). p. 367-372. Includes references. (NAL Call No.: DNAL 79.8 W41).

1478

Effect of MCPA on 14C-Diclofop uptake and translocation (Wild oat, *Avena fatua*, phytotoxicity).

Olson, W. Nalewaja, J.D. Champaign, Ill., Weed Science Society of America. Weed science. Jan 1982. v. 30 (1). p. 59-63. Includes 18 ref. (NAL Call No.: 79.8 W41).

1479

Effect of moisture stress on wild oat (*Avena fatua*) response to diclofop (Weeds).

Akey, W.C. WEESA. Morrison, I.N. Champaign : Weed Science Society of America. Weed science. Mar 1983. v. 31 (2). p. 247-253. ill. Includes references. (NAL Call No.: 79.8 W41).

1480

Effect of rate of glyphosate with and without frigate surfactant on weed control in orbit spring oats.

PNWSB. Morrow, L.S. Beltsville, Md. : The Society. Proceedings of the ... annual meeting - Northeastern Weed Science Society. 1985. v. 39. p. 112-113. (NAL Call No.: DNAL 79.9 N814).

1481

The effect of tank mixing sulfonyl urea herbicides with diclofop-methyl on wild oat control.

Downard, R.W. Evans, J.D. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 359-361. (NAL Call No.: DNAL 79.9 W52R).

1482

The effect of weed and invertebrate pest management on alfalfa establishment in oat stubble.

Bahler, C.C. Byers, R.A.; Stout, W.L. Ankeny, Iowa : Soil Conservation Society of America, c1987. The role of legumes in conservation tillage systems / J.F. Power, editor. Paper presented at the "National Conference on the Role of Legumes in Conservation Tillage Systems", April 27-29, 1987, University of Georgia, Athens, Georgia. p. 76-77. Includes references. (NAL Call No.: DNAL SB203.R6).

1483

The effect of wounding on primary dormancy in wild oat (*Avena fatua*) caryopses.

WEESA6. Foley, M.E. Champaign, Ill. : Weed Science Society of America. Weed science. Mar 1987. v. 35 (2). p. 180-184. Includes references. (NAL Call No.: DNAL 79.8 W41).

1484

Effect of 2,4-D (dichlorophenoxyacetic acid) on the hydrolysis of diclofop-methyl in wild oat (*Avena fatua*).

Hill, B.D. Todd, B.G.; Stobbe, E.H. Champaign, Ill., Weed Science Society of America. Weed science. Nov 1980. v. 28 (6). p. 725-729. ill. 19 ref. (NAL Call No.: 79.8 W41).

1485

Effects of chlorsulfuron or 2,4-D upon diclofop-methyl efficacy in oat (*Avena sativa*) (Herbicides, antagonism, antiauxin).

Hall, C. Edgington, L.V.; Switzer, C.M. Champaign : Weed Science Society of America. Weed science. Nov 1982. v. 30 (6). p. 672-676. ill. 12 ref. (NAL Call No.: 79.8 W41).

1486

Effects of fluazifop and barley windbreak competition on seeded onions in organic soil.

PNWSB. Lanterman, W.S. Warholc, D.T.; Ellerbrock, L.A.; Stachowski, P.J. Beltsville, Md. : The Society. Proceedings of the ... annual meeting - Northeastern Weed Science Society. 1985. v. 39. p. 184-187. (NAL Call No.: DNAL 79.9 N814).

(WEEDS)

1487

Effects of herbicides on the lipid composition of plant membranes (Barley, chloroplasts).
St John, J.B. Washington, D.C., The Society.
ACS symposium series - American Chemical
Society. 1982. 1982. (181). p. 97-109. ill.
Includes 23 ref. (NAL Call No.: QD1.A45).

1488

Effects of Pyrenophora teres and weeds on barley yield and yield components.
PHYTAJ. Burleigh, J.R. Tajani, M.; Seck, M. St.
Paul, Minn. : American Phytopathological
Society. Phytopathology. Mar 1988. v. 78 (3).
p. 295-299. Includes references. (NAL Call No.:
DNAL 464.8 P56).

1489

Effects of soil moisture on the vegetative growth of wild oat (Avena fatua) (Water stress, stomatal diffusion, Manitoba).
Akey, W.C. Morrison, I.N. Champaign, Ill. :
Weed Science Society of America. Weed science.
Sept 1984. v. 32 (5). p. 625-630. ill. Includes
14 references. (NAL Call No.: 79.8 W41).

1490

Effects of variations in drop makeup on the phytotoxicity of glyphosate (Herbicides, controlled-drop application, barley).
Ambach, R.M. Ashford, R. Champaign, Ill., Weed
Science Society of America. Weed science. May
1982. v. 30 (3). p. 221-224. Includes 6 ref.
(NAL Call No.: 79.8 W41).

1491

Efficacy of postharvest herbicides on Russian thistle (Salsola iberica) control and seed germination.
WEESA6. Young, F.L. Whitesides, R.E. Champaign,
Ill. : Weed Science Society of America. Weed
science. July 1987. v. 35 (4). p. 544-559.
Includes references. (NAL Call No.: DNAL 79.8
W41).

1492

Efficacy of two isomers of DPX-6202 on wild oat and spring peas.
Prather, T.S. Callihan, R.H.; Thill, D.C.
S.I. : Western Society of Weed Science.
Research progress report - Western Society of
Weed Science. 1987. p. 227. (NAL Call No.: DNAL
79.9 W52R).

1493

Established foxtail barley, Hordeum jubatum, control with glyphosate plus ammonium sulfate.
WETEE9. Donald, W.W. Champaign, Ill. : The
Society. Weed technology : a journal of the
Weed Science Society of America. July 1988. v.
2 (3). p. 364-368. ill. Includes references.
(NAL Call No.: DNAL SB610.W39).

1494

Establishment of legumes using no-till seedings with and without herbicides.
AAREEZ. Guillard, K. Allinson, D.W. New York :
Springer. Applied agricultural research. 1987.
v. 1 (5). p. 281-288. Includes references. (NAL
Call No.: DNAL S539.5.A77).

1495

Evaluation of bromoxynil tank mixes for weed control in spring barley.
Kidder, D.W. Drummond, D.P. S.I. : The
Society. Research progress report - Western
Society of Weed Science. 1988. p. 196-197. (NAL
Call No.: DNAL 79.9 W52R).

1496

Evaluation of early spring herbicide treatments in fallow.
Miller, S.D. S.I. : Western Society of Weed
Science. Research progress report - Western
Society of Weed Science. 1987. p. 345. (NAL
Call No.: DNAL 79.9 W52R).

1497

Evaluation of herbicides for broadleaf weed control in spring barley.
Miller, S.D. S.I. : Western Society of Weed
Science. Research progress report - Western
Society of Weed Science. 1986. p. 196-197. (NAL
Call No.: DNAL 79.9 W52R).

1498

Evaluation of herbicides for field bindweed (Convolvulus arvensis L.) control and crop tolerance.
Whitson, T.D. Tuck, B. S.I. : Western Society
of Weed Science. Research progress report -
Western Society of Weed Science. 1986. p. 190.
(NAL Call No.: DNAL 79.9 W52R).

1499

Evaluation of herbicides for weed control in barley.
WAEBA. Miller, S.D. Krall, J.M. Laramie : The Station. Bulletin B - Wyoming, Agricultural Experiment Station. 1986. (885). p. 152-153. (NAL Call No.: DNAL 100 W99 (1)).

1500

Evaluation of herbicides for weed control in barley.
Miller, S.D. Krall, J.M. S.l. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 262-263. (NAL Call No.: DNAL 79.9 W52R).

1501

Evaluation of herbicides for wild oat control in spring barley.
Miller, S.D. Gill, J.R. S.l. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 201. (NAL Call No.: DNAL 79.9 W52R).

1502

Evaluation of three wild oat herbicides in combination with broadleaf herbicides for the control of weeds in spring barley and the economic value of the crop.
WSWPA. Stewart, V.R. Keener, T.K. Reno, Nev. : The Society. Proceedings - Western Society of Weed Science. 1987. v. 40. p. 104-105. (NAL Call No.: DNAL 79.9 W52).

1503

The extent to which weeds modify the transpiration of cereals /by A.L. Bakke and H.H. Plagge.
Bakke, Arthur Laurence, b. 1886. Plagge, H. H. 1894-. Ames, Iowa : Agricultural Experiment Station, Iowa State College of Agriculture and Mechanic Arts. 1926. p. 212-239 : charts ; 22 cm. Bibliography: p. 238-239. (NAL Call No.: DNAL 100 Io9 no.96).

1504

Factors of wild oat (*Avena fatua*) interference on spring barley (*Hordeum vulgare*) growth and yield.
WEESA6. Morishita, D.W. Thill, D.C. Champaign, Ill. : Weed Science Society of America. Weed science. Jan 1988. v. 36 (1). p. 37-42. Includes references. (NAL Call No.: DNAL 79.8 W41).

1505

Field experiments with oats, 1890 / George E. Morrow, Thomas F. Hunt . Milk and butter tests / G.E. Morrow . Cream raising by dilution / G.E. Morrow, E.H. Farrington . The hessian fly / S.A. Forbes . Canada thistles : their extermination / T.J. Burrill .
Morrow, G. E. 1840-. Hunt, Thomas F. 1862-; Morrow, G. E. 1840-; Morrow, G. E. 1840-; Farrington, E. H. 1860-; Forbes, Stephen Alfred, 1844-1930.; Burrill, Thomas Jonathan, 1839-1916. Champaign, Ill. : University of Illinois Agricultural Experiment Station, 23 cm. Caption title. p. 353 -388 : ill. ; 23 cm. (NAL Call No.: DNAL 100 I165 no.12).

1506

Flurecol (as growth-retarding and suppressing agents to dicotyledonous plants, cereal herbicides).
Amadori, E. Heupt, W. New York, Academic Press, 1980. Updated general techniques and additional pesticides, edited by Gunter Zweig and Joseph Sherma. p. 319-329. ill. 5 ref. (NAL Call No.: 395).

1507

Fluroxypyr: a new environmentally compatible herbicide.
WSWPA. Schober, A.E. McMaster, S.A.; Gantz, R.L. Reno : The Society. Proceedings - Western Society of Weed Science. Paper presented at the annual meeting of the Western Society of Weed Science, March 18-20, 1986, San Diego, California. 1986. v. 39. p. 167-168. (NAL Call No.: DNAL 79.9 W52).

1508

Foxtail barley (*Hordeum jubatum* L.) control in perennial grass meadows.
Whitson, T.D. Langbehn, G. S.l. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 32. (NAL Call No.: DNAL 79.9 W52R).

1509

Glyphosate sled for alley cleanup in research plots.
CRPSAY. Nelson, L.R. Ward, S.; Collins, F. Madison, Wis. : Crop Science Society of America. A wooden sled with an attached sponge was developed and used to successfully apply glyphosate N(phosphonomethyl)glycine to alleyways between small grain plots. The objective for constructing this device was to improve the procedure for killing vegetation in alleyways of small grain experiments. The sponge was attached to the bottom of a hinged platform. The platform allowed to glyphosate-treated sponge to follow the contour

(WEEDS)

of the soil surface as the sled was pulled through the alleys. The sponge was wetted with a solution containing 150 mL L-1 of a glyphosate formulation with 356 g (acid equivalent) L-1 of glyphosate. Excellent vegetation control was accomplished with this device. Advantages over other procedures are (i) glyphosate can be applied when small grain is quite tall (20 cm), (ii) dead vegetation remains on the soil to reduce erosion, (iii) the method prevents spray drift, and (iv) construction costs are minimal. Crop science. Nov/Dec 1989. v. 29 (6). p. 1567-1568. ill. Includes references. (NAL Call No.: DNAL 64.8 C883).

1510

Graminicides for wild oat control in dry spring peas.

Huston, C.H. Callihan, R.H.; Thill, D.C. S.1. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 163-164. (NAL Call No.: DNAL 79.9 W52R).

1511

Grass weed control in California cereals: update and review.

WSWPA. Mitich, L.W. Kyser, G.B. Reno : The Society. Proceedings - Western Society of Weed Science. Paper presented at the annual meeting of the Western Society of Weed Science, March 18-20, 1986, San Diego, California. 1986. v. 39. p. 169-172. (NAL Call No.: DNAL 79.9 W52).

1512

Growth and control of wild oat (*Avena fatua*, Canada).

Banting, J.D. Champaign : Weeds Today, Inc. Weeds today. Winter 1982. v. 13 (4). p. 1-4. ill. (NAL Call No.: SB610.W4).

1513

Growth and reproduction of *Drechslera sorokiniana* (leaf and root pathogen of cereals and grasses) as influenced by preemergence herbicides.

Hodges, C.F. Bronx, N.Y., New York Botanical Garden. Mycologia. Mar/Apr 1981. v. 73 (2). p. 244-251. ill. 25 ref. (NAL Call No.: 450 M99).

1514

Influence of picloram alone or plus 2,4-D on control of wild oats (*Avena fatua*) with four postemergence herbicides (Antagonism, annual grass weed in western Canada).

O'Sullivan, P.A. WEESA. Champaign : Weed Science Society of America. Weed science. Nov 1983. v. 31 (6). p. 889-891. Includes references. (NAL

Call No.: 79.8 W41).

1515

Influence of thiameturon and DLX-L5300 on wild oats (*Avena fatua*) control with barban, diclofop, AC 222,293, and difenzoquat.

WEESA6. Eberlein, C.V. Miller, T.L.; Wiersma, J.V. Champaign, Ill. : Weed Science Society of America. In field studies, wild oat control with AC 222,293 and difenzoquat was not reduced when each herbicide was applied in combination with thiameturon or DPX-L5300. Mixtures of barban with thiameturon gave wild oat control similar to barban applied alone, but mixtures of barban with DPX-L5300 sometimes gave less wild oat control than barban applied alone. Control was reduced when diclofop was applied in combination with thiameturon in 1985 but not in 1986. When antagonism occurred, addition of crop oil concentrate (COC) at 1.2 L/ha to the diclofop-thiameturon spray mixture overcame the antagonism. When diclofop was applied in combination with DPX-L5300, control was reduced and antagonism could not be overcome by increasing the diclofop rate or by adding COC to the spray mixture. In greenhouse studies the inert ingredients in the thiameturon and DPX-L5300 formulations did not antagonize diclofop activity on wild oats. TLC analysis of ¹⁴C-diclofop solutions with and without thiameturon or DPX-L5300 revealed neither degradation products of diclofop nor evidence of complexing between diclofop and thiameturon or diclofop and DPX-L5300. Diclofop antagonism by thiameturon or DPX-L5300 was not due to a chemical or physical interaction between the herbicides in the spray mixture. Weed science. Nov 1988. v. 36 (6). p. 792-799. Includes references. (NAL Call No.: DNAL 79.8 W41).

1516

Influence of tillage on horseweed, *Conyza canadensis*.

WETEE9. Brown, S.M. Whitwell, T. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. July 1988. v. 2 (3). p. 269-270. Includes references. (NAL Call No.: DNAL SB610.W39).

1517

Influence of weed control treatments on soybean cultivars in an oat-soybean rotation.

AGJ0AT. Burnside, D.C. Moomaw, R.S. Madison, Wis. : American Society of Agronomy. Agronomy journal. Nov/Dec 1984. v. 76 (6). p. 887-890. Includes 13 references. (NAL Call No.: DNAL 4 AM34P).

1518

Influence of 2,4-D and MCPA formulations and oil on diclofop phytotoxicity.
 WEESA6. Gillespie, G.R. Nalewaja, J.D. Champaign, Ill. : Weed Science Society of America. Field experiments were conducted during 1984 and 1985 to determine the influence of 2,4-D and MCPA amine and ester formulations on control of foxtail millet with diclofop. Foxtail millet control increased as the diclofop rate increased from 0.6 to 1.7 kg ai/ha whether diclofop was applied alone or with 2,4-D or MCPA amine and ester formulations; however, foxtail millet control was lower when diclofop was applied with 2,4-D or MCPA amine and ester formulations compared to diclofop applied alone. MCPA ester, MCPA amine, 2,4-D ester, or 2,4-D amine at 0.14 kg ai/ha added to a diclofop spray mixture reduced foxtail millet control 8, 15, 20, and 30% compared to diclofop applied alone, averaged over diclofop rates of 0.6, 0.8, 1.1, 1.4, and 1.7 kg/ha. Adding a petroleum oil to diclofop increased foxtail millet control with diclofop at 0.6 kg/ha but not at 0.8 kg/ha. The petroleum oil additive did not overcome the antagonism of foxtail millet control when diclofop at 0.6 kg/ha was applied with 2,4-D or MCPA amine and ester formulations. The amount of 14C-diclofop absorbed by oats was greater when diclofop was applied with the ester formulations of 2,4-D or MCPA compared to the amine formulations. Either once-refined sunflower oil or sunflower oil methyl ester applied with 14C-diclofop increased the amount of 14C absorbed and translocated in oat plants compared to 14C-diclofop applied alone. Weed science. May 1989. v. 37 (3). p. 380-384. Includes references. (NAL Call No.: DNAL 79.8 W41).

1519

Inhibited mitotic entry is the cause of growth inhibition by cinmethylin.
 WEESA6. El-deek, M.H. Hess, F.D. Champaign, Ill. : Weed Science Society of America. Weed science. Sept 1986. v. 34 (5). p. 684-688. Includes references. (NAL Call No.: DNAL 79.8 W41).

1520

Integrated control of weeds in small grains in Nebraska : cooperative agreement between USDA/ARS and UN-L : final report, October 1, 1982. -.
 Wax, L. M. (Peoria, Ill. U.S. Dept. of Agriculture, Agricultural Research Service, North Central Region 1982). Caption title ~Cooperative agreement no. 58-519B-O-0891 ~Transmitted by L. M. Wax. 39 leaves ; 28 cm. Bibliography: leaf 39. (NAL Call No.: aSB612.N2I57).

1521

The interaction between the sulfonylurea herbicides and diclofop-methyl on control of wild oat (*Avena fatua* L.).
 WSWPA. Downard, R.W. Evans, J.O. Reno, Nev. : The Society. Proceedings - Western Society of Weed Science. 1987. v. 40. p. 26-31. Includes references. (NAL Call No.: DNAL 79.9 W52).

1522

Isolation and identification of some phytotoxic compounds from aqueous extracts of rye (*Secale cereale* L.).
 JAFCAU. Shilling, D.G. Jones, L.A.; Worsham, A.D.; Parker, C.E.; Wilson, R.F. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. July/Aug 1986. v. 34 (4). p. 633-638. Includes references. (NAL Call No.: DNAL 381 J8223).

1523

Key factors affecting weed-crop balance in agroecosystems.
 Sen, D.N. Boca Raton, Fla. : CRC Press, c1988. Weed management in agroecosystems : ecological approaches / editors, Miguel A. Altieri, Matt Liebman. Literature review. p. 157-182. ill. Includes references. (NAL Call No.: DNAL SB611.5.W43).

1524

Leaf wash techniques for estimation of foliar absorption of herbicides (*Fagopyrum tataricum*, *Hordeum vulgare*, *Cirsium arvense*, glyphosate).
 Devine, M.D. Bestman, H.D.; Hall, C.; Vanden Born, W.H. Champaign, Ill. : Weed Science Society of America. Weed science. May 1984. v. 32 (3). p. 418-425. ill. Includes references. (NAL Call No.: 79.8 W41).

1525

Low volume herbicide application for broadleaf weed control in barley.
 Miller, S.D. Krall, J.M. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 212. (NAL Call No.: DNAL 79.9 W52R).

1526

A management tool for small grain producers...the wild oat staging card (Weeds, herbicide application timing, Montana).
 Dyer, W. Fay, P.; Rardon, P.; Stewart, V. Bozeman : The Station. Capsule information series - Montana Agricultural Experiment Station. June 1983. June 1983. (30). 5 p. ill. (NAL Call No.: S83.M6).

(WEEDS)

1527

Mayweed chamomile and catchweed bedstraw control in winter barley in northern Idaho.
Lish, J.M. Thill, D.C. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 198-199. (NAL Call No.: DNAL 79.9 W52R).

1528

Metabolism of chlorsulfuron by plants: biological basis for selectivity of a new herbicide for cereals.
Sweetser, P.B. Schow, G.S.; Hutchison, J.M. New York, Academic Press. Pesticide biochemistry and physiology. Feb 1982. v. 17 (1). p. 18-23. ill. Includes 9 ref. (NAL Call No.: SB951.P49).

1529

Metsulfuron methyl--a new alternative for broadleaf weed control in cereals and reduced tillage fallow.
WSWPA. Warner, R.W. Kral, C.W.; Henson, M.A.; Saladini, J.L. Reno : The Society. Proceedings - Western Society of Weed Science. Paper presented at the annual meeting of the Western Society of Weed Science, March 18-20, 1986, San Diego, California. 1986. v. 39. p. 129-133. (NAL Call No.: DNAL 79.9 W52).

1530

The mode of action of chlorsulfuron: a new herbicide for cereals.
Ray, T.B. New York, Academic Press. Pesticide biochemistry and physiology. Feb 1982. v. 17 (1). p. 10-17. ill. Includes 18 ref. (NAL Call No.: SB951.P49).

1531

Modeling weed populations in cereals.
Cousens, R. Moss, S.R.; Cussans, G.W.; Wilson, E.J. Champaign, Ill. : Weed Science Society of America. Reviews of weed science. Literature review. 1987. v. 3. p. 93-112. ill. Includes references. (NAL Call No.: DNAL SB610.R47).

1532

Morphological and histological effects of three grass selective herbicides on developing wild oat (*Avena fatua*) stems.
WEESA6. Jain, R. Born, W.H. vanden. Champaign, Ill. : Weed Science Society of America. Three grass selective herbicides, sethoxydim, fluazifop, and haloxyfop, applied to wild oat plants at the five-leaf stage inhibited growth and induced chlorosis in leaves. Young and actively growing tissues were affected first. Stem elongation in wild oat was inhibited within 2 days of treatment with sethoxydim and

within 5 days of treatment with fluazifop or haloxyfop. At these same observation times, internodes that were elongating rapidly at the time of treatment were constricted at the base. These symptoms were followed by necrosis of the internode tissue. Histological examination of the affected internodes indicated that the herbicides inhibited cell division in very young internodes and inhibited both cell division and cell elongation in slightly older internodes. Initial injury occurred in the epidermal, cortical, and procambium cells of the peripheral regions of the stems located at the base of the affected internodes. Necrosis then progressed to the center of the stem tissue and all cells in the internodes were killed within 14 days of treatment. All three herbicides caused similar morphological and histological effects on developing wild oat stems. Weed science. July 1989. v. 37 (4). p. 575-584. ill. Includes references. (NAL Call No.: DNAL 79.8 W41).

1533

The morphological and physiological response of slender oat (*Avena barbata*) to the herbicides barban and difenzoquat.
WEESA6. Price, S.C. Hill, J.E.; Allard, R.W. Champaign, Ill. : Weed Science Society of America. Weed science. Jan 1988. v. 36 (1). p. 60-69. Includes references. (NAL Call No.: DNAL 79.8 W41).

1534

Morphological and physiological variation in wild oat (*Avena fatua*, dormancy, herbicide resistance, varieties, weeds).
Miller, S.D. Nalewaja, J.D.; Mulder, C.E.G. Madison, Wis., American Society of Agronomy. Agronomy journal. Sept/Oct 1982. v. 74 (5). p. 771-775. ill. 11 ref. (NAL Call No.: 4 AM34P).

1535

Mountain meadow (*Hordeum jubatum*, Weed control, Wyoming).
Alley, H.P. Vore, R.E.; Humburg, N.E. Laramie, Wyo., The Station. Research journal - University of Wyoming, Agricultural Experiment Station. Jan 1982. Jan 1982. (172). p. 19-27. (NAL Call No.: S131.E22).

1536

The number one weed problem in Montana: controlling wild oats (*Avena fatua*).
Fay, P.K. Stewart, V.R. Bozeman, The Station. Capsule information series - Montana Agricultural Experiment Station. May 1981. May 1981. (24). 3 p. ill. (NAL Call No.: S83.M6).

1537

An oats (*Avena sativa*)-soybean (*Glycine max*) rotation using ecofarming versus conventional tillage.

WEESA6. Moomaw, R.S. Champaign, Ill. : Weed Science Society of America. Weed science. July 1985. v. 33 (4). p. 544-550. Includes 29 references. (NAL Call No.: DNAL 79.8 W41).

1538

Oats research.

Reeves, D.L. Brookings, S.D. : The Station. Plant science pamphlet - Plant Science Dept., Agricultural Experiment Station, South Dakota State University. In the series analytic: 1986 Annual progress report, Northeast Research Station, Watertown, South Dakota. Jan 1987. (100). p. 7. (NAL Call No.: DNAL S541.5.S8P5).

1539

Oats research.

Reeves, D.L. Brookings, S.D. : The Station. Plant science pamphlet - Plant Science Dept., Agricultural Experiment Station, South Dakota State University. In the series analytic: 1987 Annual Progress Report--Northeast Research Station, Watertown, South Dakota. Jan 1988. (5). p. 37-38. (NAL Call No.: DNAL S541.5.S8P5).

1540

Pacific Northwest Weed Control Handbook, January 1989.

Burrill, L.C. Braunworth, W.S. Jr.; William, R.D.; Parker, R.; Swan, D.G.; Howard, S.W.; Kidder, D.W. Corvallis, Or. : The Services. Pacific Northwest weed control handbook - Extension Services of Oregon State University, Washington State University, and the University of Idaho. Jan 1989. 276 p. (NAL Call No.: DNAL SB612.A19P3).

1541

Performance of triallate as influenced by incorporation with various pieces of tillage equipment.

WSWPA. Ryerson, D.K. Reno : The Society. Proceedings - Western Society of Weed Science. Paper presented at the annual meeting of the Western Society of Weed Science, March 18-20, 1986, San Diego, California. 1986. v. 39. p. 136-138. (NAL Call No.: DNAL 79.9 W52).

1542

Postemergence control of an oat cover crop and broadleaf weeds in direct-seeded nursery beds. HUHSA. Warmund, M.R. Alexandria, Va. : American Society for Horticultural Science. HortScience. Aug 1987. v. 22 (4). p. 603-605. Includes references. (NAL Call No.: DNAL SB1.H6).

1543

Postemergence herbicide application on three accessions of wild oat.

Tapia, L.S. Dial, M.U.; Thill, D.C. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 364-365. (NAL Call No.: DNAL 79.9 W52R).

1544

Postemergence herbicide, wild oat control in irrigated, no-till spring barley.

Lish, J.M. Thill, D.C. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 200. (NAL Call No.: DNAL 79.9 W52R).

1545

The potential of scopoletin, a biotoxic root exudate of oats (*Avena sativa* L.), for biological weed control / by Peter K. Fay. -. Fay, Peter K., 1941-. Ann Arbor, Mich. University Microfilms 1976. Thesis--Cornell University, 1975. Facsimile produced by microfilm-xerography. viii, 96 leaves. Includes bibliographies. (NAL Call No.: DISS 75-18,127).

1546

A problem in small grains and alfalfa...controlling catchweed bedstraw (*Galium aparine*, weeds).

Trippet, B.U. Dyer, W.E.; Fay, P.K. Bozeman : The Station. Capsule information series - Montana Agricultural Experiment Station. May 1982. May 1982. (26). 2 p. ill. (NAL Call No.: S83.M6).

1547

A problem in winter and spring grains: controlling green foxtail and yellow foxtail (*pigeongrass*).

Nelson, J.E. Bozeman, Mont. : The Service. Montguide MT : Agriculture - Montana State University, Cooperative Extension Service. May 1985. (8517). 3 p. ill. (NAL Call No.: DNAL S544.3.M9M65).

(WEEDS)

1548

Racing to control wild oats.

Northcutt, G. MT. Bozeman, The Service. Focus on Montana agriculture - Cooperative Extension Service, Montana State University. Winter 1980. v. 2 (1). p. 10-11. ill. (NAL Call No.: S451.M9M9).

1549

Reduced tillage systems for Montana (Small grain production, includes herbicides and pesticides application guidelines).

Rardon, P. Bozeman : The Service. Bulletin - Cooperative Extension Service, Montana State University. Mar 1983. Mar 1983. (1286). 28 p. ill. (NAL Call No.: 275.29 M76C).

1550

Results of bindweed control experiments at the Fort Hays Branch Station, Hays, Kansas, 1935 to 1940 / F.L. Timmons .

Timmons, F. L. 1905-. Manhattan, Kan. : Agricultural Experiment Station, Kansas State College of Agriculture and Applied Science, 1941. "In cooperation with Division of Cereal Crops and Diseases, Bureau of Plant Industry, U.S. Dept. of Agriculture"--Cover. 50 p. : ill. ; 22 cm. (NAL Call No.: DNAL 100 K13S (1) no.296).

1551

Seedling competition between mountain rye, 'Hycrest' crested wheatgrass, and downy brome.

JRMGA. Buman, R.A. Monsen, S.E.; Abernethy, R.H. Denver, Colo. : Society for Range Management. Journal of range management. Jan 1988. v. 41 (1). p. 30-34. Includes references. (NAL Call No.: DNAL 60.18 J82).

1552

Selective control of Canada thistle in cereals with clopyralid.

Curtis, R.E. Haagsma, T. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 188. (NAL Call No.: DNAL 79.9 W52R).

1553

Selective control of Canada thistle in cereals with 3,6-dichloropicolinic acid (clopyralid).

WSWPA. Curtis, R.E. Haagsma, T. Reno : The Society. Proceedings - Western Society of Weed Science. Paper presented at the annual meeting of the Western Society of Weed Science, March 18-20, 1986, San Diego, California. 1986. v. 39. p. 159-166. Includes references. (NAL Call No.: DNAL 79.9 W52).

1554

Self-burial of wild oat florets.

AGJQAT. Somody, C.N. Nalewaja, J.D.; Miller, S.D. Madison, Wis. : American Society of Agronomy. Agronomy journal. May/June 1985. v. 77 (3). p. 359-362. ill. Includes 5 references. (NAL Call No.: DNAL 4 AM34P).

1555

Shortcut for bluetongue research.

AGREA. Senft, D. Washington, D.C. : The Administration. Agricultural research - U.S. Department of Agriculture, Agricultural Research Service. Mar 1989. v. 37 (3). p. 22. (NAL Call No.: DNAL 1.98 AG84).

1556

Small grains (Weed control, Wyoming).

Alley, H.P. Humburg, N.E. Laramie, Wyo., The Station. Research journal - University of Wyoming, Agricultural Experiment Station. Jan 1982. Jan 1982. (172). p. 47-59. (NAL Call No.: S131.E22).

1557

Soybean cultivar response to reduced tillage systems in northern dryland areas.

AGJQAT. Deibert, E.J. Madison, Wis. : American Society of Agronomy. Information on response of soybean Glycine max (L.) Merr. cultivars to reduced tillage systems in northern dryland areas is limited. A 4-yr field study (1984 to 1987) was conducted to evaluate the effect of tillage system, weed control method, and cultivar maturity on soybean seed yield variables. An early and a late-maturing soybean cultivar were grown on a Fargo clay (fine, montmorillonitic frigid Vertic Haplaquoll) on established tillage plots. Tillage systems included conventional (moldboard plow) and three reduced tillage systems (sweep, intertill, and no-till) with herbicides or herbicides plus cultivation for weed control. Climatic conditions resulted in differences among years in seed yield, seed weight, seed moisture, seed oil concentration, and seed oil yield. These seed variables were not significantly influenced by tillage system, weed control method, or cultivar maturity when grown in rotation with barley (*Hordeum vulgare* L.), but showed significant interactions. Cultivation for weed control depressed seed yield and weight of only the early cultivar. Early plant water stress (June and July) lowered yield of the early cultivar more than the late cultivar. Early cultivar no-till yields (1240 kg ha⁻¹) were greater than tilled system yields (average 1070 kg ha⁻¹), while late cultivar yields were similar among systems (average 1420 kg ha⁻¹). An early maturing cultivar performed similarly to a late-maturing cultivar irrespective of tillage system unless early plant water stress was encountered. Fall application of granular herbicide provided good

(WEEDS)

weed control, but cultivation for weed control was not beneficial for the yields parameters measured. *Agronomy journal*. July/Aug 1989. v. 81 (4). p. 672-676. Includes references. (NAL Call No.: DNAL 4 AM34P).

1558

Striga control under peasant farming conditions.

Ogborn, J.E.A. Boca Raton, Fla. : CRC Press, c1987-. Parasitic weeds in agriculture / editor, Lytton J. Musselman. p. 145-158. Includes references. (NAL Call No.: DNAL SB611.P34).

1559

Temperature requirements for mountain rye, Hycrest crested wheatgrass, and downy brome germination.

JRMGA. Buman, R.A. Abernethy, R.H. Denver, Colo. : Society for Range Management. *Journal of range management*. Jan 1988. v. 41 (1). p. 35-39. Includes references. (NAL Call No.: DNAL 60.18 J82).

1560

Tillage effects on spring barley production.

Flom, D.G. Thill, D.C.; Callihan, R.H. S.I. Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 198-199. (NAL Call No.: DNAL 79.9 W52R).

1561

Today's herbicide: glean (for cereals).

Palm, H.L. Champaign : Weeds Today, Inc. *Weeds today*. Winter 1982. v. 13 (4). p. 5-6. (NAL Call No.: SB610.W4).

1562

Translocation of different 2,4-D, bentazon, diclofop, or diclofop-methyl combinations in oat (*Avena sativa*) and soybean (*Glycine max*) (Herbicides, antagonism).

Hall, C. Edgington, L.V.; Switzer, C.M. Champaign : Weed Science Society of America. *Weed science*. Nov 1982. v. 30 (6). p. 676-682. 23 ref. (NAL Call No.: 79.8 W41).

1563

Treatment of currants and cherries to prevent spot diseases / L.H. Pammel, G.W. Carver. Squirrel-tail grass or wild barley, (*Hordeum jubatum* L.) / by L.H. Pammel. The chemical composition of Squirrel-tail grass / J.B. Weems, W.H. Heileman.

Pammel, L. H. 1862-1931. Carver, George Washington, 1864?-1943.; Weems, J. B.; Heileman, W. H. Ames, Iowa : Experiment Station, Iowa Agricultural College, 1895. Caption title. p. 289-321 : ill. ; 23 cm. Includes bibliographical references. (NAL Call No.: DNAL 100 Io9 no.30).

1564

Triallate phytotoxicity and nitrogen fertilization (Oats).

McKercher, R.B. McGregor, W.R. Champaign, Ill., Weed Science Society of America. *Weed science*. Nov 1980. v. 28 (6). p. 740-744. ill. 10 ref. (NAL Call No.: 79.8 W41).

1565

Use of a mathematical model to determine the fate of atrazine in barley (*Hordeum vulgare*).

WEESA6. Raynaud, S. Bastide, J.; Coste, C. Champaign, Ill. : Weed Science Society of America. *Weed science*. Nov 1985. v. 33 (6). p. 906-912. Includes 28 references. (NAL Call No.: DNAL 79.8 W41).

1566

Variability for response to herbicides in wild oat (*Avena fatua*) populations.

WEESA6. Thai, K.M. Jana, S.; Naylor, J.M. Champaign, Ill. : Weed Science Society of America. *Weed science*. Nov 1985. v. 33 (6). p. 829-835. Includes 10 references. (NAL Call No.: DNAL 79.8 W41).

1567

Variation within pure lines of wild oats (*Avena fatua*) in relation to temperature of development.

WEESA6. Adkins, S.W. Loewen, M.; Symons, S.J. Champaign, Ill. : Weed Science Society of America. *Weed science*. Mar 1987. v. 35 (2). p. 169-172. Includes references. (NAL Call No.: DNAL 79.8 W41).

1568

Volunteer barley (*Hordeum vulgare*) interference in canola (*Brassica campestris* and *B. napus*).

WEESA6. O'Donovan, J.T. Sharma, A.K.; Kirkland, K.J.; De St Remy, E.A. Champaign, Ill. : Weed Science Society of America. The yield potential and the effect on yield loss of canola of different densities of volunteer barley were investigated at three locations in western Canada. Field studies were conducted from 1982 to 1986. Rectangular hyperbolic models based on data pooled over years, locations, and canola cultivars, and incorporating different densities of volunteer barley and canola accurately portrayed field responses in most

(WEEDS)

instances. Results indicated that volunteer barley severely reduced canola yield. However, financial losses due to reduced canola yield were partly offset by the volunteer barley crop. Weed science. Nov 1988. v. 36 (6). p. 734-739. Includes references. (NAL Call No.: DNAL 79.8 W41).

1569

Volunteer Jerusalem artichoke (*Helianthus tuberosus*) interference and control in barley (*Hordeum vulgare*).
WETEE9. Wall, D.A. Friesen, G.H. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Jan/Mar 1989. v. 3 (1). p. 170-172. Includes references. (NAL Call No.: DNAL SB610.W39).

1570

Weed control.
Murdock, E.C. Clemson, S.C. : The Service. Circular - Clemson University, Cooperative Extension Service. In series analytic: Small grain production guidelines for South Carolina, 1988-89. Aug 1988. (463,rev.). p. 4-12. ill. (NAL Call No.: DNAL 275.29 S08E).

1571

Weed control and its impact on cereal production (Herbicides, yield increases, Tunisia).
Ettounsi, L. Corvallis : The Station. Special report - Agricultural Experiment Station, Oregon State University. 1982. French text p. 124-137. List of common weed species. 1982. (668). p. 321-331. maps. (NAL Call No.: 100 OR3M).

1572

Weed control by means of chemical sprays.
AGJ0AT. Bolley, H.L. Madison, Wis. : American Society of Agronomy. Agronomy journal. 1910. v. 1. p. 159-168. (NAL Call No.: DNAL 4 AM34P).

1573

Weed-control evaluations in no-till soybeans (*Glycine max*) double-cropped with rye (*Secale cereale*) (Georgia).
Banks, P.A.GARRA. Kvien, J.S. Athens : The Stations. Research report - University of Georgia, College of Agriculture, Experiment Stations. July 1983. July 1983. (431). 6 p. Includes references. (NAL Call No.: S51.E22).

1574

Weed control in a low-till oat (*Avena sativa*)-soybean (*Glycine max*) rotation.
Burnside, O.C.WEESA. Carlson, D.R. Champaign : Weed Science Society of America. Weed science. Nov 1983. v. 31 (6). p. 853-856. Includes references. (NAL Call No.: 79.8 W41).

1575

Weed control in barley.
Miller, S.D. Lauer, J. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 207. (NAL Call No.: DNAL 79.9 W52R).

1576

Weed control in conservation tillage systems--small grains.
Wicks, G.A. Champaign, Ill. : Weed Science Society of America. Monograph series of the Weed Science Society of America. Literature review. 1985. (2). p. 77-92. Includes references. (NAL Call No.: DNAL SB610.M65).

1577

Weed control in oats.
Mitich, L. Melvin, G. Berkeley, Calif., The Service. Leaflet - University of California, Cooperative Extension Service. Sept 1981. (21254). 2 p. (NAL Call No.: S544.3.C2C3).

1578

Weed control in small grain.
Wrage, Leon J. Arnold, W. E.; O'Neal, W. B. Document available from: South Dakota State Univ., Ag. Information Bulletin Room, Extension Bldg., Brookings, South Dakota 57007 19--?. Information in the publication discussed weed control for spring grains, winter grains, and flax. Spraying cost are included. 8 p. : ill. (NAL Call No.: Document available from source.). (NAL Call No.: FS 552).

1579

Weed control in small grains and flax.
Strand, Oliver E. Behrens, Richard. Document available from: University of Minnesota, Bulletin Room, 1420 Eckles Avenue, St. Paul, Minnesota 55108 1981. Summarizes herbicide treatments to control weeds in small grains and flax. 4 p. : ill. (NAL Call No.: Document available from source.). (NAL Call No.: Ext. Folder 493).

1580

Weed control in small grains (in Minnesota, chemicals).
Strand, O.E. Behrens, R. St. Paul, Minn., The Service. Extension folder - Minnesota University, Agricultural Extension Service. 1982. 1982. (493). 4 p. ill. (NAL Call No.: 275.29 M66EX).

1581

Weed control in small grains (Oregon as an example).
Appleby, A.P. Corvallis : The Station. Special report - Agricultural Experiment Station, Oregon State University. 1982. French text p. 138-143. 1982. (668). p. 333-340. 5 ref. (NAL Call No.: 100 OR3M).

1582

Weed control with clopyralid combinations in barley.
Miller, S.C. Krall, J.M. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 210-211. (NAL Call No.: DNAL 79.9 W52R).

1583

Weed management in dryland cereal production with special reference to the Near East.
PPRBA. Kukula, S.T. Rome : World Reporting Service on Plant Diseases and Pests, FAO. Plant protection bulletin. 1986. v. 34 (3). p. 133-138. Includes references. (NAL Call No.: DNAL 421 P692).

1584

Weed populations in conventional and no-tillage peanuts.
Costello, S.R. Gallaher, R.N. Gainesville, Fla. : The Station. Agronomy research report AY - Agricultural Experiment Stations, University of Florida. Includes statistical data. 1984? . (85-2). 9 p. Includes references. (NAL Call No.: DNAL S540.A2F62).

1585

Whipping weeds, naturally.
DeVault, G. Emmaus, Pa. : Regenerative Agriculture Association. The New farm. May/June 1987. v. 9 (4). p. 36-37. ill. (NAL Call No.: DNAL S1.N32).

1586

Wild barley control in seeding alfalfa.
Cudney, D.W. Orloff, S.B. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 116. (NAL Call No.: DNAL 79.9 W52R).

1587

Wild barley (foxtail) control in alfalfa.
Robinson, G.D. Gilbert, D.E. Reno, Nev. : College of Agriculture, University of Nevada-Reno, Nevada Cooperative Extension. Fact sheet - College of Agriculture, University of Nevada-Reno, Nevada Cooperative Extension. 1987? . (87-41). 4 p. ill. Includes references. (NAL Call No.: DNAL S544.3.N3C66).

1588

Wild mustard, *Sinapis arvensis*, control in common buckwheat, *Fagopyrum esculentum*, with desmedipham and fluorochloridone.
WETEE9. Friesen, G.H. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Apr 1988. v. 2 (2). p. 175-178. Includes references. (NAL Call No.: DNAL SB610.W39).

1589

Wild oat and wild buckwheat control in irrigated spring barley in southeast Idaho.
Dial, M.J. Thill, D.C. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 189-190. (NAL Call No.: DNAL 79.9 W52R).

1590

Wild oat (*Avena fatua*): competition and crop loss.
Lee, G.A. Coleman-Harrell, M.E.; Mundt, G.A. Moscow, Idaho, The Service. Current information series - Cooperative Extension Service, University of Idaho. Idaho. University. Cooperative Extension Service. Aug 1980. (541). 3 p. (NAL Call No.: 275.29 ID131DC).

1591

Wild oat (*Avena fatua*) and *Avena sterilis* morphological characteristics and response to herbicides.
Somody, C.N. Nalewaja, J.D.; Miller, S.D. Champaign, Ill. : Weed Science Society of America. Weed science. May 1984. v. 32 (3). p. 353-359. Includes references. (NAL Call No.: 79.8 W41).

(WEEDS)

1592

Wild oat (*Avena fatua*) and spring barley (*Hordeum vulgare*) growth and development in monoculture and mixed culture.
WEESA6. Morishita, D.W. Thill, D.C. Champaign, Ill. : Weed Science Society of America. Weed science. Jan 1988. v. 36 (1). p. 43-48. Includes references. (NAL Call No.: DNAL 79.8 W41).

1593

Wild oat (*Avena fatua*) control in Texas.
Lovelace, D.A. TX. Wiese, A.F. College Station, Tex., The Service. Leaflet L - Texas Agricultural Extension Service. Texas A and M University. Agricultural Extension Service. Dec 1978. Dec 1978. (1708). 2 p. (NAL Call No.: 275.29 T313).

1594

Wild oat (*Avena fatua*) herbicide studies. I. Physiological response of wild oat to five postemergence herbicides.
Chow, P.N.P. Champaign, Ill., Weed Science Society of America. Weed science. Jan 1982. v. 30 (1). p. 1-6. ill. Includes 31 ref. (NAL Call No.: 79.8 W41).

1595

Wild oat (*Avena fatua*) seed environment and germination (North Dakota).
Somody, C.N. Nalewaja, J.D.; Miller, S.D. Champaign, Ill. : Weed Science Society of America. Weed science. July 1984. v. 32 (4). p. 502-507. ill. Includes 11 references. (NAL Call No.: 79.8 W41).

1596

Wild oat control in lentils.
Callihan, R.H. Huston, C.H.; Thill, D.C. S.1. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 159-160. (NAL Call No.: DNAL 79.9 W52R).

1597

Wild oat control in no-till seeded spring barley.
Dial, M.J. Thill, D.C.; Lish, J.M. S.1. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 187-188. (NAL Call No.: DNAL 79.9 W52R).

1598

Wild oat control in no-tillage spring barley in southeastern Idaho.
Lish, J.M. Thill, D.C. S.1. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 277-279. (NAL Call No.: DNAL 79.9 W52R).

1599

Wild oat control in spring barley.
Mallory, C.A. Lish, J.M.; Thill, D.C. S.1. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 203. (NAL Call No.: DNAL 79.9 W52R).

1600

Wild oat control with imazamethabenz tank mixes.
Mallory, C.A. Dial, M.J.; Lish, J.M.; Thill, D.C. S.1. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 204-206. (NAL Call No.: DNAL 79.9 W52R).

1601

Wild oat control with PP604 plus vegetable crop oil.
Mallory, C.A. Lish, J.M.; Thill, D.C. S.1. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 201-202. (NAL Call No.: DNAL 79.9 W52R).

1602

Wild oat (Dryland barley, Weed control, Wyoming).
Alley, H.P. Humburg, N.E.; Costel, G.L. Laramie, Wyo., The Station. Research journal - University of Wyoming, Agricultural Experiment Station. Jan 1982. Jan 1982. (172). p. 39-41. (NAL Call No.: S131.E22).

1603

Wild oat identification and control.
Strand, Oliver E. 1981. This publication discusses the control of wild oat. Control measures are included. Document available from: Bulletin Room, 3 Coffey Hall, 1420 Eckles Avenue, University of Minnesota, St. Paul, Minnesota 55108. 1 sheet : ill. (NAL Call No.: Not available at NAL.). (NAL Call No.: Agricultural Chemicals Fact Sheet No. 9).

1604

Wild oats (*Avena fatua*) and IPM (Integrated Pest Management).
Drummond, W.D. Champaign, Ill., Weeds Today, Inc. Weeds today. Late Spring 1980. v. 11 (2). p. 16-17. ill. (NAL Call No.: SB610.W4).

1605

Wild oats control in barley.
WAEBA. Miller, S.D. Lauer, J. Laramie : The Station. Bulletin B - Wyoming, Agricultural Experiment Station. 1986. (885). p. 138-139. (NAL Call No.: DNAL 100 W99 (1)).

1606

Wild oats control in barley.
Miller, S.D. Lauer, J. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 272-273. (NAL Call No.: DNAL 79.9 W52R).

1607

Wild oats: one of the worst weeds.
CRSOA. Puy, D. van der. Madison, Wis. : American Society of Agronomy. Crops and soils magazine. Aug/Sept 1986. v. 38 (9). p. 13-14. ill. (NAL Call No.: DNAL 6 W55).

1608

Winter weed control in established alfalfa.
Orloff, S.B. Cudney, D.W. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 96-97. (NAL Call No.: DNAL 79.9 W52R).

1609

\$10 weed control in no-till beans.
Bruskc, M. Emmaus, Pa. : Regenerative Agriculture Association. The New farm. Feb 1987. v. 9 (2). p. 10-11. ill. (NAL Call No.: DNAL S1.N32).

1610

2-chloro-N-((4-methoxy-6-methyl-2,3,5-triazin-2-yl)aminocarbonyl)benzenesulfonamide, a new herbicide (Tested on cereal crop weeds and water hyacinth, *Eichorina crassipes*).
Levitt, G. Ploeg, H.L.; Weigel, R.C. Jr.; Fitzgerald, D.J. Washington, D.C., American Chemical Society. Journal of agricultural and food chemistry. Mar/Apr 1981. v. 29 (2). p. 416-417. 6 ref. (NAL Call No.: 381 J8223).

1611

2, 4-D for weed control in cereal crops /by C.I. Seely, Howard B. Roylance.
Seely, C. I. Roylance, Howard B. Moscow, Idaho : University of Idaho, College of Agriculture, 1954. 1 folded sheet (6 p.) : ill. ; 23 cm. (NAL Call No.: DNAL 100 ID14 no.205).

PESTICIDES - GENERAL

1612

Absorption and translocation of CGA-82725 with additives.

WEESA6. Gillespie, G.R. Skrzypczak, G.A.; Nalewaja, J.D. Champaign, Ill. : Weed Science Society of America. Abstract: The influence of various additives on CGA-82725 2-propenyl-2,4-(3,5-dichloro-2-pyridyloxy)phenoxy propanoate absorption and translocation was determined in oats (*Avena sativa* L. 'Lyon'). The absorption and translocation of 14C was greater when 14C-CGA-82725 was applied with petroleum oil compared to soybean Glycine max (L.) Merr oil. The translocation of 14C was greater at 96 than 48 h after 14C-CGA-82725 application. The absorption of 14C was greater at 48 than 24 h but was similar at 48 and 96 h after 14C-CGA-82725 application with no additive, petroleum oil, or soybean oil. The absorbed and translocated 14C was greater when 14C-CGA-82725 was applied with oil at 1.2 compared to 0.6 L/ha. No additional increase in 14C absorption and translocation was obtained if the oil volume was increased to 2.3 L/ha. The addition of petroleum oil to 14C-CGA-82725 increased 14C absorption and translocation more than the addition of palm (*Eleais quineensis* Jalq.), safflower (*Carthamus tinctorius* L.), linseed (*Linum usitatissimum* L.), or soybean oil. The four seed oils and the emulsifier. At Plus 300F caused similar increases in 14C absorption and translocation over 14C-CGA-82725 applied alone. Ethylene glycol did not increase 14C absorption and translocation compared to 14C-CGA-82725 applied alone. Weed science. May 1988. v. 36 (3). p. 282-285. Includes references. (NAL Call No.: DNAL 79.8 W41).

1613

Absorption and translocation of fluazifop with additives.

WEESA6. Nalewaja, J.D. Skrzypczak, G.A. Champaign, Ill. : Weed Science Society of America. Weed science. July 1986. v. 34 (4). p. 572-576. Includes 9 references. (NAL Call No.: DNAL 79.8 W41).

1614

Action of selected herbicides and Tween 20 on oat (*Avena sativa*) membranes.

Watson, M.C. Bartels, P.G.; Hamilton, K.C. Champaign, Ill., Weed Science Society of America. Weed science. Jan 1980. v. 28 (1). p. 122-127. ill. 28 ref. (NAL Call No.: 79.8 W41).

1615

Alfalfa and oat yields as influenced by triazine herbicide residues in three tillage systems.

PNWSB. Lazowski, E.J. Hartwig, N.L.; Hall, J.K. Beltsville, Md. : The Society. Proceedings of the ... annual meeting - Northeastern Weed Science Society. 1986. v. 40. p. 10-15. Includes references. (NAL Call No.: DNAL 79.9

N814).

1616

Application of 14C-labeled herbicides in lysimeter studies.

WEESA6. Fuhr, F. Champaign, Ill. : Weed Science Society of America. Weed science. Presented at the "Symposium on Assessment of Methodology for Field Evaluation of Herbicide Behavior in Soils," Weed Science Society of America, February 9, 1984, Miami, Florida. 1985. v. 33 (suppl.2). p. 11-17. ill. Includes 39 references. (NAL Call No.: DNAL 79.8 W41).

1617

Barley (*Hordeum vulgare*) response to herbicides applied at three growth stages.

WEETE9. Martin, D.A. Miller, S.D.; Alley, H.P. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Jan 1988. v. 2 (1). p. 41-45. Includes references. (NAL Call No.: DNAL SB610.W39).

1618

Biological properties of D-amino acid conjugates of 2,4-D (2,4-dichlorophenoxyacetic acid, Glycine max, soybeans, *Avena sativa*, oats, herbicides).

Davidonia, G.H. Hamilton, R.H.; Vallejo, R.P.; Bully, R.; Mumma, R.O. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Aug 1982. v. 70 (2). p. 357-360. 19 ref. (NAL Call No.: 450 P692).

1619

A comparative study of the persistence, movement, and metabolism of six carbon-14 insecticides (potential environmental pollutants) in soils and plants (Oats).

Fuhremann, T.W. Lichtenstein, E.P. Washington, D.C., American Chemical Society. Journal of agricultural and food chemistry. Mar/Apr 1980. v. 28 (2). p. 446-452. ill. 27 ref. (NAL Call No.: 381 J8223).

1620

Crop groupings: a survey of its possibilities for deltamethrin registrations.

AECTCV. Mestres, G. Mestres, R. New York, N.Y. : Springer-Verlag. Archives of environmental contamination and toxicology. May 1985. v. 14 (3). p. 321-324. Includes references. (NAL Call No.: DNAL TD172.A7).

(PESTICIDES - GENERAL)

1621

Degradation of the herbicide mecoprop (2-(2-methyl-4-chlorophenoxy)propionic acid) by a synergistic microbial community.

APMBA. Lappin, H.M. Greaves, M.P.; Slater, J.H. Washington, D.C. : American Society for Microbiology. Applied and environmental microbiology. Feb 1985. v. 49 (2). p. 429-433. Includes 26 references. (NAL Call No.: DNAL 448.3 AP5).

1622

Detection of pesticide residues.

Hascoet, M. New York, N.Y. : Lavoisier Pub., c1988. Preservation and storage of grains, seeds, and their by-products : cereals, oilseeds, pulses, and animal feed / edited by J.L. Multon ; preface by A.M. Reimbert ; translated from French by D. Marsh ; reread by A.J. Eydt. p. 516-526. Includes references. (NAL Call No.: DNAL SB190 C6513).

1623

Determination of chlorsulfuron residues in grain, straw, and green plants of cereals by high-performance liquid chromatography.

Slates, R.V.JAFCA. Washington : American Chemical Society. Journal of agricultural and food chemistry. Jan/Feb 1983. v. 31 (1). p. 113-117. ill. 1 p. ref. (NAL Call No.: 381 J8223).

1624

Developmental effects of (the herbicide) Sandoz 6706 on activities of enzymes of phenolic and general metabolism in barley shoots grown in the dark or under low or high intensity light.

Blume, D.E. McClure, J.W. Bethesda, Md., American Society of Plant Physiologists. Plant physiology. Feb 1980. v. 65 (2). p. 238-244. ill. 24 ref. (NAL Call No.: 450 P692).

1625

Diclofop and MCPA influence on (oat) coleoptile growth.

Olson, W.A. Nalewaja, J.D.; Schroeder, G.L.; Duysen, M.E. Champaign, Ill., Weed Science Society of America. Weed science. Sept 1981. v. 29 (5). p. 597-600. ill. 19 ref. (NAL Call No.: 79.8 W41).

1626

Differential inhibition of potassium ion absorption by difenzoquat in wild oat and cereals (Herbicide toxicity).

Cohen, A.S. Morrison, I.N. New York, Academic Press. Pesticide biochemistry and physiology. Oct 1982. v. 18 (2). p. 174-179. ill. 23 ref.

(NAL Call No.: SB951.P49).

1627

Effect of application time on soil residue and efficacy of sulfonylureas.

SWSPBE. Foy, C.L. Mersie, W. Raleigh, N.C. : The Society. Proceedings - Southern Weed Science Society. 1986. (39th). p. 446-456. Includes references. (NAL Call No.: DNAL 79.9 S08 (P)).

1628

Effect of atrazine carryover on malting quality of barley (Herbicide, residues).

Brinkman, M.A. Langer, D.K.; Harvey, R.G.; Burger, W.C. Madison, Wis., Crop Science Society of America. Crop science. Nov/Dec 1981. v. 21 (6). p. 973-976. 12 ref. (NAL Call No.: 64.8 C883).

1629

Effect of environment and adjuvants on asulam phytotoxicity.

WEESA6. Nalewaja, J.D. Woznica, Z. Champaign, Ill. : Weed Science Society of America.

Abstract: Experiments were conducted to determine the influence of various factors on asulam

methyl (4-amino-phenyl)sulfonyl carbamate toxicity to flax (*Linum usitatissimum* L.) and wild oats (*Avena fatua* L. ~ AVEFA). Asulam toxicity to both flax and wild oats generally increased as temperature, humidity, and soil moisture increased after treatment. Octoxynol (alpha- p-1,1,3,3-tetramethyl butyl phenyl -omega-hydroxypoly(oxyethylene) in the spray solution increased asulam toxicity to both species in all environments. Octoxynol and trimethylenonpolyethoxyethanol (WK) enhanced asulam toxicity more than other adjuvants evaluated. Asulam toxicity to both flax and wild oats increased as octoxynol concentration in the spray increased. Flax tolerance to asulam generally increased with flax height at treatment. 'Flor' flax was the most asulam susceptible of six cultivars evaluated. A 2-mm simulated rainfall within 3 or 6 h after asulam treatment reduced toxicity to wild oats and flax, respectively. Weed science. May 1988. v. 36 (3). p. 367-372. Includes references. (NAL Call No.: DNAL 79.8 W41).

1630

Effect of ethephon-bromoxynil and ethephon-DPXR9674 on spring barley yield.

Lish, J.M. Thill, D.C. S.l. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 200. (NAL Call No.: DNAL 79.9 W52R).

(PESTICIDES - GENERAL)

1631

Effect of glyphosate on indole-3-acetic acid metabolism in tolerant and susceptible plants. JPGRDI. Lee, T.T. Dumas, T. New York, N.Y. : Springer. Journal of plant growth regulation. 1985. v. 4 (1). p. 29-39. ill. Includes references. (NAL Call No.: DNAL QK745.J6).

1632

Effects of chlorsulfuron on meiosis and seed viability in rye (*Secale cereale* L.). WSWPA. Zollinger, R.K. Evans, J.O. Reno : The Society. Proceedings - Western Society of Weed Science. 1985. v. 38. p. 114-119. (NAL Call No.: DNAL 79.9 W52).

1633

Effects of haloxyfop and CGA-82725 on cell cycle and cell division of oat (*Avena sativa*) root tips. WEESA6. Kim, J.C. Bendixen, L.E. Champaign, Ill. : Weed Science Society of America. Weed science. Nov 1987. v. 35 (6). p. 769-774. Includes references. (NAL Call No.: DNAL 79.8 W41).

1634

Evaluation of chlorophyll fluorescence parameters for an intact-plant herbicide bioassay. CRPSAY. Shaw, D.R. Peeper, T.F.; Nofziger, D.L. Madison, Wis. : Crop Science Society of America. Crop science. July/Aug 1986. v. 26 (4). p. 756-760. Includes references. (NAL Call No.: DNAL 64.8 C883).

1635

Excretion and placental and mammary transfer of hexachlorobenzene in the European ferret (*Mustela putorius furo*) (Preemergence fungicides for cereal grain seeds). Bleavins, M.R. JTEHD. Breslin, W.J.; Aulerich, R.J.; Ringer, R.K. Washington : Hemisphere Publishing. Journal of toxicology and environmental health. Dec 1982. v. 10 (6). p. 929-940. Includes references. (NAL Call No.: RA565.A1J6).

1636

Flurecol (as growth-retarding and suppressing agents to dicotyledonous plants, cereal herbicides). Amadori, E. Heupt, W. New York, Academic Press, 1980. Updated general techniques and additional pesticides, edited by Gunter Zweig and Joseph Sherma. p. 319-329. ill. 5 ref. (NAL Call No.: 395).

1637

Fungicides evaluated for cereal and forage crop disease control. PLDRA. Watkins, J.E. St. Paul, Minn. : American Phytopathological Society. Plant disease. Oct 1985. v. 69 (10). p. 911-912. (NAL Call No.: DNAL 1.9 P69P).

1638

Gas chromatographic determination of flucythrinate synthetic pyrethroid residues in a range of crops. JAFCAU. Cordon, C. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. Nov/Dec 1986. v. 34 (6). p. 953-955. Includes references. (NAL Call No.: DNAL 381 J8223).

1639

Gas chromatography-mass spectrometry of acylalanine fungicides. JAFCAU. Ripley, B.D. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. July/Aug 1985. v. 33 (4). p. 560-563. Includes references. (NAL Call No.: DNAL 381 J8223).

1640

Genetic variability for herbicide reaction in plant populations (Wild oat, *Avena barbata*, *Avena fatua*, *godetia*, *Clarkia williamsonii*, toxicity, Sierra Nevada in California). Price, S.C. WEESA. Hill, J.E.; Allard, R.W. Champaign : Weed Science Society of America. Weed science. Sept 1983. v. 31 (5). p. 652-657. ill. Includes references. (NAL Call No.: 79.8 W41).

1641

Grain pesticides American Association of Cereal Chemists. -- Washington, D.C.? : The Association, 1985. Title from container. 4 videocassettes (362 min.) : sd., col. ; 1/2 in. (NAL Call No.: DNAL Videocassette no.21).

1642

Growth inhibition and disruption of mitosis by DCPA in oat (*Avena sativa*) roots. WEESA6. Holmsen, J.D. Hess, F.D. Champaign, Ill. : Weed Science Society of America. Weed science. Nov 1984. v. 32 (6). p. 732-738. ill. Includes 28 references. (NAL Call No.: DNAL 79.8 W41).

1643

Herbicidal disruption of proton gradient development and maintenance by plasmalemma and tonoplast vesicles from oat root.

PCBPB. Ratterman, D.M. Balke, N.E. Duluth, Minn. : Academic Press. Pesticide biochemistry and physiology. Literature review. July 1988. v. 31 (3). p. 221-236. Includes references. (NAL Call No.: DNAL SB951.P49).

1644

High-pressure liquid chromatographic determination of captan, captafol, and folpet residues in plant material (Fruits, grapes, cereals, fungicides).

Buttler, B. Hormann, W.D. Washington, D.C., American Chemical Society. Journal of agricultural and food chemistry. Mar/Apr 1981. v. 29 (2). p. 257-260. ill. 3 ref. (NAL Call No.: 381 J8223).

1645

HPLC (high performance liquid chromatography)--residue analysis of the herbicide pyridate in cereals.

Lindner, W. Ruckendorfer, H. New York, N.Y. : Gordon and Breach Science Publishers. International journal of environmental analytical chemistry. 1983. v. 16 (3). p. 205-218. Includes references. (NAL Call No.: QH540.I52).

1646

Injury and grain yield of spring barley treated with diclofop and thiameturon.

Evans, R.M. Thill, D.C. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 193-195. (NAL Call No.: DNAL 79.9 W52R).

1647

Interactions in the fate of chemicals in terrestrial systems.

EESAD. Scheunert, I. Korte, F. New York : Academic Press. Ecotoxicology and environmental safety. June 1985. v. 9 (3). p. 385-391. Includes references. (NAL Call No.: DNAL QH545.A1E29).

1648

Loss of metribuzin and ethyl-metribuzin from glass and soil surfaces.

WETEE9. Peek, D.C. Appleby, A.P. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Jan/Mar 1989. v. 3 (1). p. 173-176. Includes references. (NAL Call No.: DNAL SB610.W39).

1649

Metribuzin absorption and translocation in two barley (*Hordeum vulgare*) cultivars.

WEESA6. Gawronski, S. Haderlie, L.C.; Stark, J.C. Champaign, Ill. : Weed Science Society of America. Weed science. July 1986. v. 34 (4). p. 491-495. ill. Includes 13 references. (NAL Call No.: DNAL 79.8 W41).

1650

Mode of action studies on nitrodiphenyl ether herbicides. 1. Use of barley mutants to probe the role of photosynthetic electron transport.

PLPHA. Bowyer, J.R. Smith, B.J.; Camilleri, P.; Lee, S.A. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Mar 1987. v. 83 (3). p. 613-620. ill. Includes references. (NAL Call No.: DNAL 450 P692).

1651

Mutagenicity of selected chemicals in barley test systems (Includes agricultural chemicals, environmental pollutants).

Nilan, R.A. Veleminsky, J. New York, Plenum Press. Environmental science research. 1982. v. 24. p. 291-320. ill. 63 ref. (NAL Call No.: TD172.E55).

1652

Narrow row soybean production in untilled oat stubble.

AGJOAT. Burnside, D.C. Moomaw, R.S. Madison, Wis. : American Society of Agronomy. Agronomy journal. Jan/Feb 1985. v. 77 (1). p. 36-40. Includes 11 references. (NAL Call No.: DNAL 4 AM34P).

1653

Nature of photooxidative events in leaves treated with chlorosis-inducing herbicides (*Secale cereale*, winter rye seedlings).

Feierabend, J. Winkelhusener, T. Rockville : American Society of Plant Physiologists. Plant physiology. Nov 1982. v. 70 (5). p. 1277-1282. 30 ref. (NAL Call No.: 450 P692).

1654

Paraquat concentration and renal function in mice fed purified and cereal-based diets (Herbicide toxicity).

Evers, W.D. Hook, J.B.; Bond, J.T. New York : Alan R. Liss. Drug-nutrient interactions. 1983. v. 2 (2). p. 95-104. Includes references. (NAL Call No.: RM302.D76).

(PESTICIDES - GENERAL)

1655

Pest control in forages and small grains 1980 / J. D. Doll ... (et al.). -. WI. Doll, J. D. Madison University of Wisconsin, Cooperative Extension Programs 1980. Cover title. 37 p. : ill. 28 cm. (NAL Call No.: S544.3.W6W53 no. 1981).

1656

Phytotoxic interaction studies--Techniques for evaluation and presentation of results (Interactions of two or more pesticides in plants, tested on oats and cucumbers). Nash, R.G. Champaign, Ill., Weed Science Society of America. Weed science. Mar 1981. v. 29 (2). p. 147-155. ill. 22 ref. (NAL Call No.: 79.8 W41).

1657

Phytotoxicity, adsorption, and mobility of metribuzin and its ethylthio analog as influenced by soil properties. WEESA6. Peek, D.C. Appleby, A.P. Champaign, Ill. : Weed Science Society of America. The phytotoxicity, adsorption, and mobility of metribuzin and ethyl-metribuzin were studied in five soils to determine if differential soil behavior could explain: a) the greater activity of both chemicals in western Oregon than in eastern Oregon, and b) the greater activity of metribuzin than ethyl-metribuzin under similar conditions. Metribuzin had higher activity on oats than ethyl-metribuzin in all soils and in quartz sand. Metribuzin was absorbed less and moved more than ethyl-metribuzin in all soils. Activity of both herbicides decreased as sand content increased, and activity in quartz sand was lower than in soil. Activity of both herbicides increased in a bioassay in which leaching was prevented, indicating that leaching may be important in the loss of activity of metribuzin and ethyl-metribuzin. Lowest herbicide adsorption and greatest movement were observed in coarser textured soils. Weed science. May 1989. v. 37 (3). p. 419-423. Includes references. (NAL Call No.: DNAL 79.8 W41).

1658

The products of metabolism of (14C (carbon isotopes)) triadimefon in the grain and in the straw of ripe barley (Fungicides). Rouchaud, J. Moons, C.; Meyer, J.A. New York, Springer. Bulletin of environmental contamination and toxicology. Oct 1981. v. 27 (4). p. 543-550. Bibliography p. 550. (NAL Call No.: RA1270.P35A1).

1659

Pronamide phytotoxicity in ten Wisconsin soils (Oats). Dutt, T.E. Harvey, R.G. Champaign, Ill., Weed Science Society of America. Weed science. July 1980. v. 28 (4). p. 429-432. ill. 15 ref. (NAL Call No.: 79.8 W41).

1660

Protection of grass crops from sulfonylurea and imidazolinone toxicity. Barrett, M. San Diego : Academic Press, c1989. Crop safeners for herbicides : development, uses, and mechanisms of action / edited by Kriton K. Hatzios and Robert E. Hoagland. Literature review. p. 195-220. Includes references. (NAL Call No.: DNAL SB951.45.C76).

1661

Purge and trap method for determination of ethylene dibromide in whole grains, milled grain products, intermediate grain-based foods, and animal feeds. JANCA2. Heikes, D.L. Arlington, Va. : The Association. Journal of the Association of Official Analytical Chemists. Nov/Dec 1985. v. 68 (6). p. 1108-1111. Includes 13 references. (NAL Call No.: DNAL 381 AS7).

1662

A rapid method for the analysis of the mode of action of bleaching herbicides. PCBPB. Young, A.U. Britton, G.; Musker, D. Duluth, Minn. : Academic Press. Pesticide biochemistry and physiology. Nov 1989. v. 35 (3). p. 244-250. Includes references. (NAL Call No.: DNAL SB951.P49).

1663

Regulatory pest control. DeWitt, Jerald R. Document available from: Iowa State University, Publications Distribution, Printing & Publications Bldg., Ames, Iowa 50011 1977. This publication offers information on control principles, control methods, and pesticide useage on japanese beetles, gypsy moths, cereal leaf beetle, soybean nematode and barberry. 12 p. (NAL Call No.: Document available from source.). (NAL Call No.: CS-24).

1664

Requirements and developments in small grain seed treatments. PLDRA. Phipps, P.M. St. Paul, Minn. : American Phytopathological Society. Plant disease. Nov 1985. v. 69 (11). p. 1009-1010. (NAL Call No.: DNAL 1.9 P69P).

1665

Residual effect of metsulfuron applied during the fallow year on barley and lentils.
Mashhadi, H.R. Evans, J.O. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 258-259. (NAL Call No.: DNAL 79.9 W52R).

1666

Response of oats to atrazine (Herbicides).
Brinkman, M.A. Langer, D.K.; Harvey, R.G.; Hardie, A.R. Madison, Wis., Crop Science Society of America. Crop science. Mar/Apr 1980. v. 20 (2). p. 185-189. ill. 18 ref. (NAL Call No.: 64.8 C883).

1667

Response to spring barley (Hordeum vulgare) to herbicides.
WETEE9. Clay, S.A. Thill, D.C.; Cochran, V.L. Champaign, Ill. : The Society. Weed technology : a journal of the Weed Science Society of America. Jan 1988. v. 2 (1). p. 68-71. Includes references. (NAL Call No.: DNAL SB610.W39).

1668

Seedling injury and chromosome aberrations induced by Bladex, Dowpon, Princep and Tenoran (Herbicides, barley).
Kahlon, P.S. Nashville, The Academy. Journal of the Tennessee Academy of Science. Tennessee Academy of Science. Jan 1980. v. 55 (1). p. 17-19. ill. 9 ref. (NAL Call No.: 500 T25A).

1669

The site of the inhibition of the shikimate pathway by glyphosate (herbicide, buckwheat). II. Interference of glyphosate with chorismate formation in vivo and in vitro.
Amrhein, N. Deus, B.; Gehrke, P.; Steinrucken. H.C. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Nov 1980. v. 66 (5). p. 830-834. ill. 24 ref. (NAL Call No.: 450 P692).

1670

The site of the inhibition of the shikimate pathway by glyphosate (postemergence herbicide). I. Inhibition by glyphosate of phenylpropanoid synthesis in buckwheat (Fagopyrum esculentum Moench).
Hollander, H. Amrhein, N. Rockville, Md., American Society of Plant Physiologists. Plant physiology. Nov 1980. v. 66 (5). p. 823-829. ill. 28 ref. (NAL Call No.: 450 P692).

1671

Small grain insect management guidelines number 1 - aphids / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-O31).

1672

Small grain insect management guidelines number 2 - armyworms / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"July 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-O32).

1673

Small grain insect management guidelines, number 3 -- Cereal leaf beetles / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-O33).

1674

Small grain insect management guidelines number 4 - fall armyworms / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-O34).

1675

Small grain insect management guidelines, number 5 -- natural enemies / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-O35).

(PESTICIDES - GENERAL)

1676

Small grain insect pest management for Virginia / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1984. Pesticide Applicator Training collection ~"March 1984.". 9 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-015 1984).

1677

Smut in oats, insecticides and fertilizer analyses / W.W. Cooke .
Cooke, Wells Woodbridge, 1858-1916.
Burlington : Vermont State Agricultural Experiment Station, 1888. Cover title. 5, 2 p. ; 22 cm. (NAL Call No.: DNAL 100 V59 no.9).

1678

Soil bound residues of carbaryl and 1-naphthol: release and mineralization in soil, and uptake by plants.
JPFCD2. Murthy, N.B.K. Raghu, K. New York, N.Y. : Marcel Dekker. Journal of environmental science and health : Part B : Pesticides, food contaminants, and agricultural wastes. 1988. v. 23 (6). p. 575-585. Includes references. (NAL Call No.: DNAL TD172.J61).

1679

Spring oats response to metsulfuron treatment during the fallow year.
Mashhadi, H.R. Evans, J.O. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 260. (NAL Call No.: DNAL 79.9 W52R).

1680

Tolerance of spring-planted cereals to sulfonylurea herbicides.
Spinney, R.L. Appleby, A.P.; Brewster, B.D. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1987. p. 257-259. (NAL Call No.: DNAL 79.9 W52R).

1681

Translocation of different 2,4-D, bentazon, diclofop, or diclofop-methyl combinations in oat (Avena sativa) and soybean (Glycine max) (Herbicides, antagonism).
Hall, C. Edgington, L.V.; Switzer, C.M. Champaign : Weed Science Society of America. Weed science. Nov 1982. v. 30 (6). p. 676-682. 23 ref. (NAL Call No.: 79.8 W41).

1682

Transpiration and soil water usage by oat as affected by postemergence herbicides.
AGJ0AT. Beardmore, R.A. Linscott, D.L. Madison, Wis. : American Society of Agronomy. Changes in transpiration of plants as influenced by herbicides may or may not significantly affect soil water uptake in the field. We compared effects of the relatively new herbicides fluazifop (+/-)-butyl-2-(4- 5-(trifluoromethyl)-2-pyridinyl oxy phenoxy)propanoate, haloxyfop methyl-2-(4- 3-chloro-5-(trifluoromethyl)-2-pyridinyl oxy phenoxy)propanoate, and sethoxydim (2- 1-(ethoxyimino)butyl -5-2 2-(ethylthio)propyl -3-hydroxy-2-cyclohexen-1-one applied at various rates to oat (Avena sativa L.) at the 5- to 6-cm stage on transpiration, soil water use, and plant development. In controlled environments, transpiration rates in oat declined about 6 d after herbicide treatment. Time for and the degree of transpiration reduction were herbicide and herbicide-rate dependent, as were the reductions in plant weight and leaf number. First reductions in soil moisture under laboratory conditions were found 10 d after treatment of oat with fluazifop and haloxyfop and 12 d after treatment with sethoxydim. Similarly, in the field fluazifop treatment required 9 d and haloxyfop and sethoxydim 11 d before significant differences in soil moisture occurred. The correlation coefficient of soil matric potential and dry weight of oat in the field was -0.89 and -0.90, respectively, for 2 yr, 1984 and 1985. Reduction of oat dry matter and soil water use in the field was also herbicide rate dependent. Agronomy journal. Nov/Dec 1988. v. 80 (6). p. 982-986. Includes references. (NAL Call No.: DNAL 4 AM34P).

1683

Trifluralin and triallate retention by imbibed tame oat (Avena sativa) caryopses (Herbicides, phytotoxic effects, embryo culture, pericarp, testa).
Heath, M.C. Ashford, R.; McKercher, R.B. Champaign : Weed Science Society of America. Weed science. Mar 1984. v. 32 (2). p. 251-257. ill. Includes references. (NAL Call No.: 79.8 W41).

1684

The uptake, distribution and metabolism of four organic chemicals by soybean plants and barley roots.
ETOC DK. McFarlane, C. Nolt, C.; Wickliff, C.; Pfleeger, T.; Shimabuku, R.; McDowell, M. Elmsford : Pergamon Press. Environmental toxicology and chemistry. 1987. v. 6 (11). p. 847-856. ill. Includes references. (NAL Call No.: DNAL QH545.A1E58).

1685

Uptake of malathion from galvanized-steel surfaces by stored barley.

White, N.D.G. Abramson, D. College Park, Md. : Entomological Society of America. Journal of economic entomology. Apr 1984. v. 77 (2). p. 289-293. Includes references. (NAL Call No.: 421 J822).

1686

Use of tonoplast and plasma membrane vesicles from oat root to investigate herbicidal disruption of proton gradients.

PCBPB. Ratterman, D.M. Balke, N.E. Duluth, Minn. : Academic Press. Pesticide biochemistry and physiology. May 1987. v. 28 (1). p. 17-28. Includes references. (NAL Call No.: DNAL SB951.P49).

1687

Vitamins C and E: An antioxidative system against herbicide-induced lipid peroxidation in higher plants.

JAFCAU. Finckh, B.F. Kunert, K.J. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. July/Aug 1985. v. 33 (4). p. 574-577. Includes references. (NAL Call No.: DNAL 381 J8223).

1688

Wild oat herbicide studies. 3. Physiological and biochemical bases for interaction of barban and growth regulator herbicides in wild oat.

Chow, P.N.P.JAFCA. Taylor, H.F. Washington : American Chemical Society. Journal of agricultural and food chemistry. May 1983. v. 31 (3). p. 575-578. Includes references. (NAL Call No.: 381 J8223).

1689

2-chloro-N-((4-methoxy-6-methyl-2,3,5-triazin-2-yl)aminocarbonyl)benzenesulfonamide, a new herbicide (Tested on cereal crop weeds and water hyacinth, Eichorina crassipes).

Levitt, G. Ploeg, H.L.; Weigel, R.C. Jr.; Fitzgerald, D.J. Washington, D.C., American Chemical Society. Journal of agricultural and food chemistry. Mar/Apr 1981. v. 29 (2). p. 416-417. 6 ref. (NAL Call No.: 381 J8223).

SOIL BIOLOGY

1690

Degradation of the herbicide mecoprop (2-(2-methyl-4-chlorophenoxy)propionic acid) by a synergistic microbial community.

APMBA. Lappin, H.M. Greaves, M.P.; Slater, J.H. Washington, D.C. : American Society for Microbiology. Applied and environmental microbiology. Feb 1985. v. 49 (2). p. 429-433. Includes 26 references. (NAL Call No.: DNAL 448.3 AP5).

1691

Effect of burning cereal straw on soil properties and grain yields in Saskatchewan (Effects on physical, chemical and biological characteristics of the soil and on grain yields).

Biederbeck, V.O. Campbell, C.A.; Bowren, K.E.; Schnitzer, M.; McIver, R.N. Madison, Wis., The Society. Journal - Soil Science Society of America. Soil Science Society of America. Jan/Feb 1980. v. 44 (1). p. 103-111. ill. 42 ref. (NAL Call No.: 56.9 S03).

1692

Effects of different endomycorrhizal fungi on five host plants grown on calcined montmorillonite clay (Apple, asparagus, leek, strawberry, oats).

Plenchette, C. Furian, V.; Fortin J.A. Alexandria, Va., The Society. Journal of the American Society for Horticultural Science. July 1982. v. 107 (4). p. 535-538. 16 ref. (NAL Call No.: 81 S012).

1693

Inoculation effects on growth, nodulation, cytosol components and nitrogen fixation of narrowleaf and hairy vetch (*Rhizobium leguminosarum*, *Vici angustifolia*, *Vici villosa*, *Secale cereale*).

Lynd, J.Q. CSOSA. McNew, R.W. New York : Marcel Dekker. Communications in soil science and plant analysis. 1983. v. 14 (5). p. 411-426. ill. Includes references. (NAL Call No.: S590.C63).

SOIL CHEMISTRY AND PHYSICS

1694

Characteristics of cadmium and zinc in four soils treated with sewage sludge.
JEVQAA. Mullins, G.L. Sommers, L.E. Madison, Wis. : American Society of Agronomy. Journal of environmental quality. Oct/Dec 1986. v. 15 (4). p. 382-387. Includes references. (NAL Call No.: DNAL QH540.J6).

1695

Differential aluminum tolerances of two barley cultivars related to organic acids in their roots.
JPNUDS. Foy, C.D. Lee, E.H.; Wilding, S.B. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Tenth International Plant Nutrition Colloquium," August 4-9, 1986, Beltsville, Maryland. 1987. v. 10 (9116). p. 1089-1101. Includes references. (NAL Call No.: DNAL QK867.J67).

1696

Effect of burning cereal straw on soil properties and grain yields in Saskatchewan (Effects on physical, chemical and biological characteristics of the soil and on grain yields).
Biederbeck, V.O. Campbell, C.A.; Bowren, K.E.; Schnitzer, M.; McIver, R.N. Madison, Wis., The Society. Journal - Soil Science Society of America. Soil Science Society of America. Jan/Feb 1980. v. 44 (1). p. 103-111. ill. 42 ref. (NAL Call No.: 56.9 S03).

1697

Effect of calcium and phosphorus content of various soil series in western Washington upon the calcium and phosphorus composition of oats, red clover and white clover /by Henry F. Holtz.
Holtz, Henry F., 1880-1931. Pullman, Wash. : State College of Washington, Agricultural Experiment Station, 1930. Cover title. 45 p. ; 23 cm. Bibliography: p. 31-33. (NAL Call No.: DNAL 100 W27E no.243).

1698

Effect of legume cover crops and tillage on soil water, temperature, and organic matter.
Utomo, M. Blevins, R.L.; Frye, W.W. Ankeny, Iowa : Soil Conservation Society of America, c1987. The role of legumes in conservation tillage systems / J.F. Power, editor. Paper presented at the "National Conference on the Role of Legumes in Conservation Tillage Systems", April 27-29, 1987, University of Georgia, Athens, Georgia. p. 5-6. Includes references. (NAL Call No.: DNAL SB203.R6).

1699

Effect of sample pretreatment on extractable soil potassium.
CSOSA2. Haby, V.A. Sims, J.R.; Skogley, E.O.; Lund, R.E. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. 1988. v. 19 (1). p. 91-106. Includes references. (NAL Call No.: DNAL S590.C63).

1700

Fate of tagged urea N (nitrogen) in the field with different methods of N and organic matter placement (Broadcast and banded, barley).
Tomar, J.S. Soper, R.J. Madison, Wis., American Society of Agronomy. Agronomy journal. Nov/Dec 1981. v. 73 (6). p. 991-995. 18 ref. (NAL Call No.: 4 AM34P).

1701

Mutual influence of certain crops in relation to nitrogen.
AGJOAT. Kellerman, K.F. Wright, R.C. Madison, Wis. : American Society of Agronomy. Agronomy journal. July/Oct 1914. v. 6 (4/5). p. 204-210. Includes references. (NAL Call No.: DNAL 4 AM34P).

1702

Nitrification and total nitrogen as affected by crops, fertilizers, and copper sulfate.
AGJOAT. Jensen, C.A. Madison, Wis. : American Society of Agronomy. The accumulation of nitrates in the soils in the Arkansas River Valley in Colorado, which promised in 1910 and 1911 to become a serious factor in sugar beet growth, became less in 1912 and 1913. In the work in 1911, mustard appeared to have some effect in checking the accumulation of nitrate in the field. Copper sulfate at the rate of 100 pounds per acre on fallow was also effective in checking nitrification, reducing the average seasonal accumulation to about 60 percent of the amount found in the check plot. Molasses on fallow decreased nitrification about 25 percent as compared with the check, but the molasses-treated plot showed a little more nitrates than the plots cropped to cane and oats. Manure on fallow gave a slightly higher accumulation of nitrates than the fallow check. The waste lime on fallow caused strong nitrate accumulation, being more effective in this regard than any other treatment. In general, active nitrification did not set in until the first part of June. From then until the middle of July it was strongest and then suddenly decreased and became very feeble until the end of the experiment, August 17, regardless of the field treatments. The differences in the average seasonal accumulations of nitrates could not have been due entirely to the differences in beet yields from the plots. The mustard plot contained less total nitrogen than any of the others. The fallow plots receiving copper sulfate and molasses contained less

(SOIL CHEMISTRY AND PHYSICS)

total nitrogen than the other fallow plots. The fallow plots with waste lime and manure each contained less total nitrogen than the plot fallowed with nothing added. The reverse was true as regards nitrates. In general, there seemed to be an inverse relation between the amounts of total nitrogen. In the work in 1912, in the presence of a vigorously growing beet crop, the only treatment which showed decided increases in nitrification were cyanamid and manure plus ammonium sulfate. Aside from considerable weekly variations, there. *Agronomy journal*. Jan/Feb 1916. v. 8 (1). p. 10-22. Includes references. (NAL Call No.: DNAL 4 AM34P).

1703

Phytotoxicity, adsorption, and mobility of metribuzin and its ethylthio analog as influenced by soil properties.

WEESA6. Peek, D.C. Appleby, A.P. Champaign, Ill. : Weed Science Society of America. The phytotoxicity, adsorption, and mobility of metribuzin and ethyl-metribuzin were studied in five soils to determine if differential soil behavior could explain: a) the greater activity of both chemicals in western Oregon than in eastern Oregon, and b) the greater activity of metribuzin than ethyl-metribuzin under similar conditions. Metribuzin had higher activity on oats than ethyl-metribuzin in all soils and in quartz sand. Metribuzin was absorbed less and moved more than ethyl-metribuzin in all soils. Activity of both herbicides decreased as sand content increased, and activity in quartz sand was lower than in soil. Activity of both herbicides increased in a bioassay in which leaching was prevented, indicating that leaching may be important in the loss of activity of metribuzin and ethyl-metribuzin. Lowest herbicide adsorption and greatest movement were observed in coarser textured soils. *Weed science*. May 1989. v. 37 (3). p. 419-423. Includes references. (NAL Call No.: DNAL 79.8 W41).

1704

Soil bound residues of carbaryl and 1-naphthol: release and mineralization in soil, and uptake by plants.

JPFC2. Murthy, N.B.K. Raghu, K. New York, N.Y. : Marcel Dekker. *Journal of environmental science and health : Part B : Pesticides, food contaminants, and agricultural wastes*. 1988. v. 23 (6). p. 575-585. Includes references. (NAL Call No.: DNAL TD172.J61).

1705

Soil tests for copper, iron, manganese, and zinc in histosols: 4. Selection on the basis of soil chemical data and uptakes by oats, carrots, onions, and lettuce.

SOSCAK. Mathur, S.P. Levesque, M.P. Baltimore, Md. : Williams & Wilkins. *Soil science*. Dec 1989. v. 148 (6). p. 424-432. Includes

references. (NAL Call No.: DNAL 56.8 S03).

1706

Water stress: role in differential aluminum tolerance of barley genotypes.

AGABB. Krizek, D.T. Foy, C.D. Madison, Wis. : American Society of Agronomy. *Agronomy abstracts*. Includes abstract. 1981. (73rd). p. 181-182. (NAL Call No.: DNAL 241 AM39).

SOIL FERTILITY - FERTILIZERS

1707

Availability studies upon high potash nitrate.
AGJOAT. Allison, R.V. Madison, Wis. : American Society of Agronomy. Experimental studies upon the availability of the nitrogen and potash of High Potash Nitrate have been carried out under both green-house and field conditions. The results in pot culture indicated the value of the nitrogen and potash of this compound as being fully equivalent to that derived from sodium nitrate and potassium sulfate. Because of the fact that the field plots responded strongly to applications of nitrogen and not at all to potash, it may be assumed that the nitrogen of the High Potash Nitrate was as effective as that of sodium nitrate. Agronomy journal. Jan 1924. v. 16 (1). p. 26-30. (NAL Call No.: DNAL 4 AM34P).

1708

Boron toxicity in barley.
JPNUDS. Riley, M.M. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Paper presented at the "Tenth International Plant Nutrition Colloquium," August 4-9, 1986, Beltsville, Maryland. 1987. v. 10 (9/16). p. 2109-2115. Includes references. (NAL Call No.: DNAL QK867.J67).

1709

Characteristics of cadmium and zinc in four soils treated with sewage sludge.
JEVQAA. Mullins, G.L. Sommers, L.E. Madison, Wis. : American Society of Agronomy. Journal of environmental quality. Oct/Dec 1986. v. 15 (4). p. 382-387. Includes references. (NAL Call No.: DNAL QH540.J6).

1710

Comparison of selenium treatments of crops in the field.
Gissel-Nielsen, G. Passaic, N.J. : Humana Press. Biological trace element research. Sept 1986. v. 10 (3). p. 209-213. Includes references. (NAL Call No.: DNAL QP534.B56).

1711

Crop response to sludge loading rates.
BCYCDK. Day, A.D. Solomon, M.A.; Ottman, M.U.; Taylor, B.B. Emmaus, Pa. : J.G. Press. BioCycle. Aug 1989. v. 30 (8). p. 72-75. Includes references. (NAL Call No.: DNAL 57.8 C734).

1712

Crop rotations and manure versus agricultural chemicals in dryland grain production.
JSWCA3. Sahs, W.W. Lesoing, G. Ankeny, Iowa : Soil Conservation Society of America. Journal of soil and water conservation. Nov/Dec 1984. v. 40 (6). p. 511-516. Includes 27 references. (NAL Call No.: DNAL 56.8 J822).

1713

Differential aluminum tolerance of winter barley varieties and selections in associated greenhouse and field experiments.
AGJOAT. Reid, D.A. Jones, G.D.; Armiger, W.H.; Foy, C.D.; Koch, E.J.; Starling, T.M. Madison, Wis. : American Society of Agronomy. Agronomy journal. Mar/Apr 1969. v. 61 (2). p. 218-222. ill. Includes references. (NAL Call No.: DNAL 4 AM34P).

1714

Effect of ammonium and nitrate on growth and yield of barley on acid soils.
CSOSA2. Malhi, S.S. Nyborg, M.; Caldwell, C.D.; Hoyt, P.B.; Leitch, R.H. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. May/Sept 1988. v. 19 (7/12). p. 1049-1063. Includes references. (NAL Call No.: DNAL S590.C63).

1715

Effect of Basic-H on vegetable and agronomic crops and soil fertility at Pt. MacKenzie.
AGBOBO. Laughlin, W.M. Smith, G.R.; Peters, M.A. Fairbanks : The Station. Agroborealis - Alaska Agricultural and Forestry Experiment Station, University of Alaska-Fairbanks. July 1987. v. 19 (1). p. 31-33. Includes references. (NAL Call No.: DNAL S33.E2).

1716

Effect of Cu on the distribution of P, Ca, and Fe in barley plants.
SOSCAK. Brown, J.C. Foy, C.D. Baltimore, Md. : Williams & Wilkins. Soil science. Dec 1964. v. 98 (6). p. 362-370. ill. Includes references. (NAL Call No.: DNAL 56.8 S03).

1717

Effect of manganese and soil pH on the iron content of crops grown on Podzol soils (Hordeum vulgare, barley, Pisum sativum, peas).
Gupta, U.C. JPNUD. New York : Marcel Dekker. Journal of plant nutrition. 1982. v. 5 (10). p. 1229-1239. 14 ref. (NAL Call No.: QK867.J67).

(SOIL FERTILITY - FERTILIZERS)

1718

Effect of nitrogen nutrition on quality of agronomic crops.

Deckard, E.L. Tsai, C.Y.; Tucker, T.C. Madison, Wis. : American Society of Agronomy, 1984. Nitrogen in crop production : proceedings, symposium, 25-27 May, 1982, Sheffield, Alabama / spon. by National Fertilizer Development Center of Tennessee Valley Authority ... et al. ; Roland D. Hauck. Literature review. p. 601-615. Includes references. (NAL Call No.: DNAL S651.N57).

1719

The effect of powdery mildew on the response of barley to applied nitrogen.

Young, K.U. Khan, T.N. New South Wales : Dept. of Agriculture, 1987? . Proceedings, Australian Barley Technical Symposium : Wagga Wagga, 11-15th October 1987. p. 177-182. Includes references. (NAL Call No.: DNAL SB191.B2A9 1987).

1720

Effect of sample pretreatment on extractable soil potassium.

CSOSA2. Haby, V.A. Sims, J.R.; Skogley, E.O.; Lund, R.E. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. 1988. v. 19 (1). p. 91-106. Includes references. (NAL Call No.: DNAL S590.C63).

1721

Effect of seeding rate and time of N application on oat grain yield.

JPraen. Ahmadi, M. Wiebold, W.J.; Beuerlein, J.E. Madison, Wis. : American Society of Agronomy. Journal of production agriculture. July/Sept 1988. v. 1 (3). p. 242-244. Includes references. (NAL Call No.: DNAL S539.5.J68).

1722

The effect of surface applied soil amendments on barley root growth in an acid subsoil.

CSOSA2. Wright, R.U. Hern, J.L.; Baligar, V.C.; Bennett, D.L. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. Feb 1985. v. 16 (2). p. 179-192. ill. Includes 26 references. (NAL Call No.: DNAL S590.C63).

1723

The effect of thiosulfate on phosphorus availability and uptake by plants.

JPNUDS. Morden, G. Soper, R.; Huzel, V.; Swan, M. New York, N.Y. : Marcel Dekker. Journal of plant nutrition. Oct 1986. v. 9 (10). p. 1315-1321. Includes references. (NAL Call No.:

DNAL QK867.J67).

1724

Effects of by-product sulfuric acid on phyto availability of nutrients in irrigated calcareous, saline-sodic soils (Barley, *Hordeum vulgare*).

Cates, R.L. Jr. Haby, V.A.; Skogley, E.O.; Ferguson, H. Madison, Wis. : American Society of Agronomy. Journal of environmental quality. Apr/June 1984. v. 13 (2). p. 252-256. Includes references. (NAL Call No.: QH540.J6).

1725

Effects of chloride fertilizer and systemic fungicide seed treatments on common root rot of barley.

PLDRA. Shefelbine, P.A. Mathre, D.E.; Carlson, G. St. Paul, Minn. : American Phytopathological Society. Plant disease. July 1986. v. 70 (7). p. 639-642. Includes 22 references. (NAL Call No.: DNAL 1.9 P69P).

1726

Effects of dried sewage sludge on forage production from barley genotypes in the Sonoran Desert.

Day, A.D. Thompson, R.K.; Swingle, R.S. Superior : University of Arizona. Desert plants. 1986. v. 8 (1). p. 17-19. Includes references. (NAL Call No.: DNAL QK938.D4D4).

1727

Effects of forage legumes on yield and nitrogen uptake by a succeeding barley crop.

AAREEZ. Abernethy, R.H. Bohl, W.H. New York : Springer. Applied agricultural research. 1987. v. 2 (2). p. 97-102. Includes references. (NAL Call No.: DNAL S539.5.A77).

1728

Effects of N, P, and K (nitrogen, phosphorus, potassium) fertilization on barley grown in a newly cleared subarctic soil (Alaska).

Michaelson, G.J. Loynachan, T.E.; Woding, F.J.; Mitchell, G.A. Madison, Wis., American Society of Agronomy. Agronomy journal. July/Aug 1982. v. 74 (4). p. 694-699. ill., map. 16 ref. (NAL Call No.: 4 AM34P).

(SOIL FERTILITY - FERTILIZERS)

1729

Effects of nitrogen fertilization on manganese concentration and yield of barley and oats (*Hordeum vulgare*, *Avena sativa*).

Petrie, S.E. Jackson, T.L. Madison, Wis. : The Society. Journal - Soil Science Society of America. Mar/Apr 1984. v. 48 (2). p. 319-322. Includes references. (NAL Call No.: 56.9 S03).

1730

Effects of nitrogen fertilizer rate, seeding rate, and row spacing on semidwarf and conventional height spring oat.

CRPSAY. Marshall, H.G. Kolb, F.L.; Roth, G.W. Madison, Wis. : Crop Science Society of America. Crop science. May/June 1987. v. 27 (3). p. 572-575. Includes references. (NAL Call No.: DNAL 64.8 C883).

1731

Effects of various salts on barley growth /H.E. Dregne.

Dregne, H. E. University Park, N.M. : Agricultural Experiment Station, New Mexico State University, 1962. Caption title. 5, 1 p. : charts ; 26 cm. (NAL Call No.: DNAL 100 N465R no.62).

1732

Ethephon application and nitrogen fertilization of irrigated winter barley in an arid environment.

AGUOAT. Pearson, C.H. Golus, H.M.; Tindall, T.A. Madison, Wis. : American Society of Agronomy. Lodging in barley (*Hordeum vulgare* L.) can result in grain loss, lower seed quality, increased disease, high grain moisture, reduced harvest efficiency, and lower milling grade. Application of N fertilizer to achieve high barley yields often results in increased lodging. Consequently, N applications are often reduced in order to lessen the potential for lodging. Use of plant growth regulators may permit higher N application rates without increased lodging. Field studies were conducted in an irrigated arid environment during 1986 and 1987 on Fruita sandy clay loam (fine-loamy, mixed, mesic Typic Haplargid) soil. Ethephon (2-chloroethyl) phosphonic acid was applied at 0.42 kg a.i. ha⁻¹ to irrigated winter barley that received 56, 112, 168, and 224 kg ha⁻¹ of spring-applied N. The 1986 data indicated that, in the absence of lodging, ethephon did not affect barley grain yields. Ethephon increased grain yields in 1987 in association with reduced lodging. Plant height response to ethephon was similar across N rates. Ethephon shortened internode lengths on the upper portion of the culm and reduced lodging of winter barley grown at high N rates. Ethephon did not affect spikes m⁻², kernel mass, kernels spike⁻¹, or seed N content in either year. Application of higher N rates than traditionally used in irrigated winter barley

increased grain yield and seed N content. In years when lodging occurred, ethephon was effective as an antilodging agent. Agronomy journal. Sept/Oct 1989. v. 81 (5). p. 7171-719. Includes references. (NAL Call No.: DNAL 4 AM34P).

1733

Exploitation of soil potassium in layered profiles by root systems of cotton and barley.

SSSUD4. Gulick, S.H. Cassman, K.G.; Grattan, S.R. Madison, Wis. : The Society. On vermiculitic soils of the San Joaquin Valley, cotton (*Gossypium hirsutum* L.) often responds to K fertilization where other annual crops do not. These soils have markedly higher available K in surface soil than subsoil. A pot study was conducted to compare rooting patterns, K uptake, and dry matter production of cotton and barley (*Hordeum vulgare* L.) in soil profiles with layered K availability. Topsoil (0.15 cmolc K kg⁻¹) was layered to each of six depths above subsoil (0.09 cmolc K kg⁻¹) in 17-L pots with a total soil depth of 45 cm in all treatments. Two irrigation regimes were imposed and nutrients other than K were adequately supplied. For both crops, plant dry matter and K uptake increased linearly with increased topsoil depth but K uptake per unit increase in topsoil depth was 6.5- and 3.6-fold greater by barley than for cotton with frequent and infrequent irrigation, respectively. Increased K uptake per unit increase in topsoil depth reflected coincident root and K distribution: barley root length density (RLD) was 2.7 times greater than RLD of cotton in topsoil layers but little different in subsoil layers. Poor exploitation of topsoil layers by the cotton root system was attributed to greater sensitivity to low soil water potential. A root system with little compensatory root development in the surface soil when the subsoil is low in nutrients may limit K uptake and crop productivity on layered soils in the San Joaquin Valley and may require management systems designed to promote more congruent root and nutrient distribution. Soil Science Society of America journal. Jan/Feb 1989. v. 53 (1). p. 146-153. ill. Includes references. (NAL Call No.: DNAL 56.9 S03).

1734

Fate of tagged urea N (nitrogen) in the field with different methods of N and organic matter placement (Broadcast and banded, barley).

Tomar, U.S. Soper, R.J. Madison, Wis., American Society of Agronomy. Agronomy journal. Nov/Dec 1981. v. 73 (6). p. 991-995. 18 ref. (NAL Call No.: 4 AM34P).

(SOIL FERTILITY - FERTILIZERS)

1735

Induction of barley leaf urease.

PLPHA. Chen, Y.G. Ching, T.M. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. Mar 1988. v. 86 (3). p. 941-945. ill. Includes references. (NAL Call No.: DNAL 450 P692).

1736

Interactions between calcium amendments and phosphate on the response of alfalfa and barley growing on acid soils.

CSOSA2. Soon, Y.K. New York, N.Y. : Marcel Dekker. Communications in soil science and plant analysis. May/Sept 1988. v. 19 (7/12). p. 1343-1353. Includes references. (NAL Call No.: DNAL S590.C62).

1737

Modifications of plant growth and ash content as effected by acids added to soils.

AGUOAT. Carr, R.H. Havercamp, H.G. Madison, Wis. : American Society of Agronomy. A comparison has been made of the effect upon the growth of soybean and buckwheat plants of adding an organic (acetic) and an inorganic (sulfuric) acid to a silt and a clay-loam soil. It has been found that both acids seriously interfere with plant growth, but that acetic was more deleterious than sulfuric acid. This is believed to be due to the effect of acetic acid in supplying more toxic aluminum and iron, etc., to the plant and in decreasing calcium and magnesium to a greater extent than was done by the sulfuric acid. In addition two other series of acids (phosphoric and silicic) were used, because they make compounds with aluminum and iron which seem to be harmless to the growing plant, regardless of the fact that the root is surrounded by a soil which is highly acid. The good growth of both kinds of plants obtained, even when rather large amounts of these latter acids were added, indicates that relatively insoluble compounds of aluminum and iron, etc., were formed and that the acid by itself did no harm. Agronomy journal. Apr 1924. v. 16 (4). p. 278-283. (NAL Call No.: DNAL 4 AM34P).

1738

Physiological responses of barley leaves of foliar applied urea-ammonium nitrate.

CRPSAY. Turley, R.H. Ching, T.M. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 987-993. Includes references. (NAL Call No.: DNAL 64.8 C883).

1739

Plant response to topsoil thickness on an eroded loess soil.

JSWCA3. Mielke, L.N. Schepers, J.S. Ankeny, Iowa : Soil Conservation Society of America. Journal of soil and water conservation. Jan/Feb 1986. v. 41 (1). p. 59-63. Includes 15 references. (NAL Call No.: DNAL 56.8 J822).

1740

Profitable small grain production in the Texas Gulf Coast.

TAEBA. Miller, T.D. Livingston, S. College Station, Tex. : The Station. B - Texas Agricultural Experiment Station. Nov 1987. (1587). 8 p. (NAL Call No.: DNAL 100 T31S (1)).

1741

The rate of absorption of nitrate of soda by oats and cotton when applied at different stages of plant growth.

AGUOAT. Appleton, W.H. Helms, H.B. Madison, Wis. : American Society of Agronomy. Experiments were conducted in the greenhouse to study the rate of absorption of nitrate nitrogen by oats and cotton when applied as nitrate of soda at different stages of plant growth. The general plan of the experiment was to add nitrate to a series of cultures at a certain stage of growth. Then, at intervals of a few days, part of the cultures were harvested and the amount of nitrate absorbed was determined by plant analysis and also by determining the nitrate content of the soil. Cultures that did not receive nitrate were used as checks to indicate how much nitrogen the plant secured from the soil. With oats these studies were made at four stages of plant growth. With cotton the study was made at three stages of plant growth. The results may be summarized as follows: When sodium nitrate at the rate of 400 pounds per acre was applied to oats 14 days after planting, absorption of the nitrate was very slow for three weeks. After the third week absorption increased and all nitrate was absorbed by the close of the seventh week. When the nitrate was applied to oats at later stages of growth, the rate of absorption was more rapid. Nitrate applied 42, 70, and 92 days after planting was completely absorbed in 20, 14, and 10 days, respectively. With both oats and cotton there was a close correlation between the rate of growth and the rate of nitrate absorption. Sodium nitrate at the rate of 600 pounds per acre was applied to cotton 14, 40, and 61 days after planting. Absorption of the nitrate was complete in 36, 14, and 11 days, respectively. The results of both experiments indicate that the loss of soluble nitrogenous fertilizer by leaching may be reduced by delaying the application until the crop will absorb it rapidly. Agronomy journal. Oct 1925. v. 17 (10). p. 596-605. (NAL Call No.: DNAL 4 AM34P).

(SOIL FERTILITY - FERTILIZERS)

1742

Smut in oats, insecticides and fertilizer analyses / W.W. Cooke .

Cooke, Wells Woodbridge, 1858-1916.
Burlington : Vermont State Agricultural
Experiment Station, 1888. Cover title. 5, 2
p. ; 22 cm. (NAL Call No.: DNAL 100 V59 no.9).

1748

Triallate phytotoxicity and nitrogen fertilization (Oats).

McKercher, R.B. McGregor, W.R. Champaign, Ill.,
Weed Science Society of America. Weed science.
Nov 1980. v. 28 (6). p. 740-744. ill. 10 ref.
(NAL Call No.: 79.8 W41).

1743

Soil fertility and defoliation effects with arrowleaf clover and nitrogen fertilizer equivalence of crimson-arrowleaf clover combinations (Trifolium vesiculosum, Trifolium incarnatum, Secale cereale, rhizobium, nitrogen fixation by forage legumes, Oklahoma).

Lynd, J.O.AGJOAT. Hanlon, E.A. Jr.; Odell, G.V.
Jr. Madison : American Society of Agronomy.
Agronomy journal. Jan/Feb 1984. v. 76 (1). p.
13-16. ill. Includes references. (NAL Call No.:
4 AM34P).

1744

Storage protein accumulation in 'Scio' barley seed as affected by late foliar applications of nitrogen.

CRPSAY. Turley, R.H. Ching, T.M. Madison, Wis.
: Crop Science Society of America. Crop
science. July/Aug 1986. v. 26 (4). p. 778-782.
Includes references. (NAL Call No.: DNAL 64.8
C883).

1745

Studies on the effect of nitrogen applied to oats at different periods of growth.

AGJOAT. Gericke, W.F. Madison, Wis. : American
Society of Agronomy. Agronomy journal. Nov
1922. v. 14 (8). p. 312-320. (NAL Call No.:
DNAL 4 AM34P).

1746

Subsurface drip irrigation of cotton and small grains.

Tollefson, S. St. Joseph, Mich. : American
Society of Agricultural Engineers, c1985.
Drip/trickle irrigation in action : proceedings
of the Third International Drip/Trickle
Irrigation Congress, November 18-21, 1985,
Centre Plaza Holiday Inn, Fresno, California,
USA. p. 887-895. Includes 7 references. (NAL
Call No.: DNAL S612.I5 1985).

1747

Toxic factors in acid soils.

Foy, C.D. Burns, G.R. Washington, D.C. :
National Plant Food Institute. Plant food
review. Fall 1964. v. 10 (3). p. 2-3, 16. ill.
(NAL Call No.: DNAL 57.8 P694).

SOIL CULTIVATION

1749

Alfalfa and oat yields as influenced by triazine herbicide residues in three tillage systems.

PNWSB. Lazowski, E.J. Hartwig, N.L.; Hall, J.K. Beltsville, Md. : The Society. Proceedings of the ... annual meeting - Northeastern Weed Science Society. 1986. v. 40. p. 10-15. Includes references. (NAL Call No.: DNAL 79.9 N814).

1750

Barley vs. oat companion crops. I. Forage yield and quality response during alfalfa establishment.

CRPSAY. Brink, G.E. Marten, G.C. Madison, Wis. : Crop Science Society of America. Crop science. Sept/Oct 1986. v. 26 (5). p. 1060-1067. Includes references. (NAL Call No.: DNAL 64.8 C883).

1751

Conservation-tillage and residue-management systems for interior Alaska.

AGBOB. Siddoway, F.H. Lewis, C.E.; Cullum, R.F. Fairbanks : The Station. Agroborealis - Alaska Agricultural Experiment Station, Fairbanks. Includes lists of species. July 1984. v. 16 (2). p. 35-40. ill. Includes 5 references. (NAL Call No.: DNAL S33.E2).

1752

Crop rotations and manure versus agricultural chemicals in dryland grain production.

JSWCA3. Sahs, W.W. Lesoing, G. Ankeny, Iowa : Soil Conservation Society of America. Journal of soil and water conservation. Nov/Dec 1984. v. 40 (6). p. 511-516. Includes 27 references. (NAL Call No.: DNAL 56.8 J822).

1753

Drills and seeders for heavy residues and untilled soils (Small grain planting equipment, minimum tillage farming, Kansas).

Powell, G.M. Manhattan : The Service. L - Cooperative Extension Service, Kansas State University. June 1982. June 1982. (634). 8 p. ill. (NAL Call No.: 275.29 K13LE).

1754

The effect of different times of plowing small-grain stubble in eastern Colorado /by O.J. Grace.

Grace, Oliver J. 1880-. Washington, D.C. : U.S. Dept. of Agriculture, 1915. Caption title. 15 p. : charts ; 24 cm. (NAL Call No.: DNAL 1 Ag84B no.253).

1755

Influence of rye-cover crop management on soybean foliage arthropods.

EVETEX. Smith, A.W. Hammond, R.B.; Stinner, B.R. College Park, Md. : Entomological Society of America. Environmental entomology. Feb 1988. v. 17 (1). p. 109-114. Includes references. (NAL Call No.: DNAL QL461.E532).

1756

Metsulfuron methyl--a new alternative for broadleaf weed control in cereals and reduced tillage fallow.

WSWPA. Warner, R.W. Kral, C.W.; Henson, M.A.; Saladini, J.L. Reno : The Society. Proceedings - Western Society of Weed Science. Paper presented at the annual meeting of the Western Society of Weed Science, March 18-20, 1986, San Diego, California. 1986. v. 39. p. 129-133. (NAL Call No.: DNAL 79.9 W52).

1757

An oats (Avena sativa)-soybean (Glycine max) rotation using ecofarming versus conventional tillage.

WEESA6. Moomaw, R.S. Champaign, Ill. : Weed Science Society of America. Weed science. July 1985. v. 33 (4). p. 544-550. Includes 29 references. (NAL Call No.: DNAL 79.8 W41).

1758

Performance of triallate as influenced by incorporation with various pieces of tillage equipment.

WSWPA. Ryerson, D.K. Reno : The Society. Proceedings - Western Society of Weed Science. Paper presented at the annual meeting of the Western Society of Weed Science, March 18-20, 1986, San Diego, California. 1986. v. 39. p. 136-138. (NAL Call No.: DNAL 79.9 W52).

1759

Reduced tillage systems for Montana (Small grain production, includes herbicides and pesticides application guidelines).

Rardon, P. Bozeman : The Service. Bulletin - Cooperative Extension Service, Montana State University. Mar 1983. Mar 1983. (1286). 28 p. ill. (NAL Call No.: 275.29 M76C).

1760

Residual effect of metsulfuron applied during the fallow year on barley and lentils.

Mashhadi, H.R. Evans, J.O. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 258-259. (NAL Call No.: DNAL 79.9 W52R).

1761

Rhizoctonia root rot of small grains favored by reduced tillage in the Pacific Northwest.

PLDRA. Weller, D.M. Cook, R.J.; MacNish, G.; Bassett, E.N.; Powelson, R.L.; Petersen, R.R. St. Paul, Minn. : American Phytopathological Society. Plant disease. Jan 1986. v. 70 (1). p. 70-73. ill. Includes 35 references. (NAL Call No.: DNAL 1.9 P69P).

1762

Root-lesion nematodes, potassium deficiency, and prior cover crops as factors in apple replant disease.

JOSHB. Merwin, I.A. Stiles, W.C. Alexandria, Va. : The Society. Growth chamber evaluations of soil from an orchard replant site showed severe stunting of 'Northern Spy' apple (*Malus domestica* Borkh.) seedlings grown in field soil (FS) compared with pasteurized soil (PS) from the same site. The FS : PS seedling dry weight ratio of 0.44 indicated a serious replant problem. Leaf nutrient content was generally higher in PS than FS seedlings. Multiple-regression analysis indicated that leaf K and root-lesion nematode (*Pratylenchus penetrans* Filipjev) primary inoculum accounted for 75% of the variation in FS seedling dry weight. Apple seedling dry weight in FS was 97% of that in PS following a marigold (*Tagetes patula* L. cv. Sparky) cover crop, and 75% following oats (*Avena sativa* L. cv. Saia). Root-lesion nematodes were nearly eliminated from the plots with marigold. Other cover crops and weed-free fallow period were less effective in controlling apple replant disease and/or phytonematodes. Journal of the American Society for Horticultural Science. Sept 1989. v. 114 (5). p. 728-732. Includes references. (NAL Call No.: DNAL 81 S012).

1763

Rye residues contribute weed suppression in no-tillage cropping systems (Agroecosystems, biomass).

BARNES, J.P. JCECD. Putnam, A.R. New York : Plenum Press. Journal of chemical ecology. Aug 1983. v. 9 (8). p. 1045-1057. ill. Includes references. (NAL Call No.: QD415.A1U6).

1764

Seasonal establishment of bermudagrass using plastic and straw mulches.

AGUDAT. Sowers, R.S. Welterlen, M.S. Madison, Wis. : American Society of Agronomy. Bermudagrass *Cynodon dactylon* (L.) Pers. is normally established vegetatively during the early summer in the transition zone, to allow sufficient establishment time before the onset of freezing conditions in the fall. Clear polyethylene covers and straw mulches cause changes in the turfgrass microenvironment that may influence the rate of sprig establishment. The objective of this study was to evaluate the

effects of barley (*Hordeum vulgare* L.) straw and clear polyethylene plastic mulch (0.04-mm thickness) on the establishment of 'Midiron', 'Tufcote', and 'Vamont' bermudagrass from sprigs. Separate tests were conducted in the fall of 1983 and 1984 and the summer of 1984 and 1985. Summer spriggings were made in May, June, and July, months generally recommended for planting in the transition zone. Late-season spriggings were made in August, September, and October. Plantings were made on a Sassafras sandy loam (fine-loamy, siliceous, mesic Typic Hapludult). Summer bermudagrass establishment was reduced under straw and polyethylene covers. Injury under plastic occurred to plantings made in May and June, which were exposed to mean soil temperatures above 41 degrees C during the 8-wk cover period. In contrast, plastic mulch stimulated early fall bermudagrass growth and delayed dormancy. August and September plantings remaining under plastic throughout the winter exhibited high winter survival in comparison to unmulched or straw mulched turf, and were nearly 100% established by 1 July of the following year. Spriggings under plastic planted after 20 September exhibited poor establishment by 1 July the following year. These studies showed that plastic covers can be used to extend the establishment season of bermudagrass into the fall; however, plastic covers are detrimental to summer bermudagrass establishment. Agronomy journal. Jan/Feb 1988. v. 80 (1). p. 144-148. Includes references. (NAL Call No.: DNAL 4 AM34P).

1765

Soybean cultivar response to reduced tillage systems in northern dryland areas.

AGUDAT. Deibert, E.J. Madison, Wis. : American Society of Agronomy. Information on response of soybean *Glycine max* (L.) Merr. cultivars to reduced tillage systems in northern dryland areas is limited. A 4-yr field study (1984 to 1987) was conducted to evaluate the effect of tillage system, weed control method, and cultivar maturity on soybean seed yield variables. An early and a late-maturing soybean cultivar were grown on a Fargo clay (fine, montmorillonitic frigid Vertic Haplaquoll) on established tillage plots. Tillage systems included conventional (moldboard plow) and three reduced tillage systems (sweep, intertill, and no-till) with herbicides or herbicides plus cultivation for weed control. Climatic conditions resulted in differences among years in seed yield, seed weight, seed moisture, seed oil concentration, and seed oil yield. These seed variables were not significantly influenced by tillage system, weed control method, or cultivar maturity when grown in rotation with barley (*Hordeum vulgare* L.), but showed significant interactions. Cultivation for weed control depressed seed yield and weight of only the early cultivar. Early plant water stress (June and July) lowered yield of the early cultivar more than the late cultivar. Early cultivar no-till yields (1240 kg ha⁻¹) were greater than tilled system yields (average 1070 kg ha⁻¹), while late cultivar yields were similar among systems

(SOIL CULTIVATION)

(average 1420 kg ha⁻¹). An early maturing cultivar performed similarly to a late-maturing cultivar irrespective of tillage system unless early plant water stress was encountered. Fall application of granular herbicide provided good weed control, but cultivation for weed control was not beneficial for the yields parameters measured. *Agronomy journal*. July/Aug 1989. v. 81 (4). p. 672-676. Includes references. (NAL Call No.: DNAL 4 AM34P).

1766

Spring interseeding of winter rye with cover crops.

JSWCA3. Edwards, L.M. Ankeny, Iowa : Soil Conservation Society of America. *Journal of soil and water conservation*. May/June 1986. v. 41 (3). p. 190-191. Includes 5 references. (NAL Call No.: DNAL 56.8 J822).

1767

Spring oats response to metsulfuron treatment during the fallow year.

Mashhadi, H.R. Evans, J.O. S.I. : The Society. Research progress report - Western Society of Weed Science. 1988. p. 260. (NAL Call No.: DNAL 79.9 W52R).

1768

Tillage effects on spring barley production.

Flom, D.G. Thill, D.C.; Callihan, R.H. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 198-199. (NAL Call No.: DNAL 79.9 W52R).

1769

Using crop diversity to manage pest problems: some California examples.

Flint, M.L. Roberts, P.A. Greenbelt, Md. : Institute for Alternative Agriculture. Moderate to large scale California growers (as well as small scale ones) manipulate cropping patterns in a number of ways to reduce pest problems. Crop rotation, which can be defined as diversifying crops over time, is used to manage selected pests, primarily weeds, pathogens, and nematodes. As a substitute for pesticides, crop rotation has been most rewarding in the control of nematodes; sugarbeet cyst and root knot nematode examples are detailed. Some pests that invade fields from nearby areas can be managed by modifying adjacent cropping patterns or practices; Pierce's disease of grapes, sugarbeet yellows and border harvesting of alfalfa are given as examples. Finally, multiple crops can be grown within a single field or orchard. Although this approach is not widely practiced by many California growers, two examples of systems where intercropping has been shown to limit pest numbers without the use of pesticides are described: intercropping

of cotton with alfalfa and companion planting oats when seeding alfalfa. These examples show that using crop diversification to manage pests is feasible, but growers must be strongly motivated to make the necessary changes in cropping patterns. Most of the systems that have been widely adopted are those for which few other economically feasible methods were available. *American journal of alternative agriculture*. Fall 1988. v. 3 (4). p. 163-167. Includes references. (NAL Call No.: DNAL S605.5.A43).

1770

Weed-control evaluations in no-till soybeans (Glycine max) double-cropped with rye (Secale cereale) (Georgia).

Banks, P.A. GARRA, Kvien, J.S. Athens : The Stations. Research report - University of Georgia, College of Agriculture, Experiment Stations. July 1983. July 1983. (431). 6 p. Includes references. (NAL Call No.: S51.E22).

1771

Weed control in a low-till oat (Avena sativa)-soybean (Glycine max) rotation.

Burnside, O.C. WEESA, Carlson, D.R. Champaign : Weed Science Society of America. *Weed science*. Nov 1983. v. 31 (6). p. 853-856. Includes references. (NAL Call No.: 79.8 W41).

1772

Weed management in dryland cereal production with special reference to the Near East.

PPRBA. Kukula, S.T. Rome : World Reporting Service on Plant Diseases and Pests, FAO. *Plant protection bulletin*. 1986. v. 34 (3). p. 133-138. Includes references. (NAL Call No.: DNAL 421 P692).

1773

Whipping weeds, naturally.

DeVault, G. Emmaus, Pa. : Regenerative Agriculture Association. *The New farm*. May/June 1987. v. 9 (4). p. 36-37. ill. (NAL Call No.: DNAL S1.N32).

1774

Year-round forage production with sodseeded cool-season annuals and summer perennial grasses, 1982-1983.

Allen, M. Friesner, D.; Jodari, F.; Mason, L. Baton Rouge? , La. : The Station. Annual progress report - Southeast Research Station, Louisiana Agricultural Experiment Station. 1983. p. 75-82. (NAL Call No.: DNAL S67.E22).

1775

\$10 weed control in no-till beans.

Brusko, M. Emmaus, Pa. : Regenerative
Agriculture Association. The New farm. Feb
1987. v. 9 (2). p. 10-11. ill. (NAL Call No.:
DNAL S1.N32).

SOIL EROSION AND RECLAMATION

1776

Conservation-tillage and residue-management systems for interior Alaska.

AGB08B. Siddoway, F.H. Lewis, C.E.; Cullum, R.F. Fairbanks : The Station. Agroborealis - Alaska Agricultural Experiment Station, Fairbanks. Includes lists of species. July 1984. v. 16 (2). p. 35-40. ill. Includes 5 references. (NAL Call No.: DNAL S33.E2).

1777

Effect of legume cover crops and tillage on soil water, temperature, and organic matter.

Utomo, M. Blevins, R.L.; Frye, W.W. Ankeny, Iowa : Soil Conservation Society of America, c1987. The role of legumes in conservation tillage systems / J.F. Power, editor. Paper presented at the "National Conference on the Role of Legumes in Conservation Tillage Systems", April 27-29, 1987, University of Georgia, Athens, Georgia. p. 5-6. Includes references. (NAL Call No.: DNAL SB203.R6).

1778

Propagation of sea oats (*Uniola latifolia*).

Wilsey, M.H. Boulder, Colo. : International Plant Propagators' Society. The Plant propagator. June 1985. v. 31 (2). p. 15. (NAL Call No.: DNAL 81 P692).

1779

Spring interseeding of winter rye with cover crops.

JSWCA3. Edwards, L.M. Ankeny, Iowa : Soil Conservation Society of America. Journal of soil and water conservation. May/June 1986. v. 41 (3). p. 190-191. Includes 5 references. (NAL Call No.: DNAL 56.8 J822).

FORESTRY RELATED

1780

Buffer capacities of leaves, leaf cells, and leaf cell organelles in relation to fluxes of potentially acidic gases.

PLPHA. Pfanz, H. Heber, U. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. June 1986. v. 81 (2). p. 597-602. Includes 38 references. (NAL Call No.: DNAL 450 P692).

ENTOMOLOGY RELATED

1781

Annotated list of the insects and mites associated with stored grain and cereal products, and of their arthropod parasites and predators /by R.T. Cotton and N.E. Good. Cotton, R. T. Good, Newell E. 1905-. Washington, D.C. : U.S. Dept. of Agriculture, 1937. Caption title.- Contribution from Bureau of Entomology and Plant Quarantine. 81 p. ; 23 cm. Bibliography: p. 63-74. (NAL Call No.: DNAL 1 Ag84M no.258).

1782

Biology and control of cereal leaf beetle using natural enemies (authors: S.P. Briggs ... (et al.)). -. Briggs, S. P. (Blacksburg, Va.) Chemical, Drug and Pesticide Unit, Virginia Tech (1980?). Pesticide Applicator Training collection. 56 slides : col. + 1 sound cassette (17:50 min.) + 1 script. (NAL Call No.: Slide no.60).

1783

Feeding behavior, development, and damage by biotypes B, C, and E of *Schizaphis graminum* (Homoptera: Aphididae) on 'Wintermalt' and 'Post' barley. EVETEX. Peters, D.C. Kerns, D.; Puterka, G.J.; McNew, R. College Park, Md. : Entomological Society of America. Abstract: Greenbug, *Schizaphis graminum* (Rondani), biotypes B, C, and E, were monitored on susceptible 'Wintermalt' and resistant 'Post' barley to determine if there were behavioral differences among the biotypes when they fed on resistant and susceptible barley. All biotype-cultivar combinations were tested by observing feeding behavior, fecundity, and damage. Greenbug biotype B had shorter nonfeeding times on the host plant before probing than did biotypes C and E. Biotype B appeared to be less successful in feeding from the phloem of susceptible 'Wintermalt' but was relatively more successful than biotypes C and E on the resistant 'Post'. Development, reproduction, and particularly dry weight of biotype B were inferior to biotypes C and E on both resistant and susceptible plants. However, biotype B caused the greatest overall reduction in chlorophyll content of infested leaves and should be considered the most virulent but least biologically fit of the three biotypes in these comparisons. Environmental entomology. June 1988. v. 17 (3). p. 503-507. Includes references. (NAL Call No.: DNAL QL461.E532).

1784

Grasshoppers and their control /by J.R. Parker. Parker, J. R. 1884-. Washington, D.C. : U.S. Dept. of Agriculture, 1939. "Supersedes Farmers' bulletin 1691: How to control grasshoppers in cereal and forage crops.". 38 p. : ill., maps ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1828).

1785

How to control grasshoppers in cereal and forage crops /by J.R. Parker and W.R. Walton. Parker, J. R. 1884-. Walton, W. R. 1873-. Washington, D.C. : U.S. Dept. of Agriculture, 1938. Originally issued Apr. 1932. 17 p. : ill., plans ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1691 1938).

1786

How to control grasshoppers in cereal and forage crops /by J.R. Parker and W.R. Walton. Parker, J. R. 1884-. Walton, W. R. 1873-. Washington, D.C. : U.S. Dept. of Agriculture, 1932. "This bulletin is a revision of and supersedes Farmers' bulletin 747: Grasshopper control in Relation to Cereal and Forage Crops.". 14 p. : ill., plans ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1691).

1787

How to control grasshoppers in cereal and forage crops by J.R. Parker and W. R. Walton, and R.L. Shotwell. -. Parker, J. R. Washington, D.C. : U.S. Dept. of Agriculture, 1938. 17 p. : ill. --. (NAL Call No.: DNAL Fiche S-70 no.1691 1938).

1788

Pests not known to occur in the United States or of limited distribution. 80. A click beetle. Whittle, K. Hyattsville, Md. : The Service. APHIS 81 - U.S. Department of Agriculture. Animal and Plant Health Inspection Service. Sept 1987. (50). 5 p. ill., maps. Includes references. (NAL Call No.: DNAL aSB599.A3U5).

1789

Small grain insect management guidelines number 1 - aphids / Robert M. McPherson. -. McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-031).

1790

Small grain insect management guidelines number 2 - armyworms / Robert M. McPherson. -. McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"July 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-032).

1791

Small grain insect management guidelines, number 3 -- Cereal leaf beetles / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-033).

1792

Small grain insect management guidelines number 4 - fall armyworms / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-034).

1793

Small grain insect management guidelines, number 5 -- natural enemies / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1983. Pesticide Applicator Training collection ~Caption title ~"May 1983.". 2 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-035).

1794

Small grain insect pest management for Virginia / Robert M. McPherson. -.
McPherson, Robert M. Blacksburg Extension Division, Virginia Polytechnic Institute and State University 1984. Pesticide Applicator Training collection ~"March 1984.". 9 p. : ill. ; 28 cm. --. (NAL Call No.: S544.3.V8V52 no.444-015 1984).

1795

Webworms injurious to cereal and forage crops and their control /by Geo. G. Ainslie.
Ainslie, George G. 1886-1930. Washington, D.C. : U.S. Dept. of Agriculture, 1922. Cover title.~ "Contribution from the Bureau of Entomology.". 16 p. : ill., 1 map ; 23 cm. (NAL Call No.: DNAL 1 Ag84F no.1258).

1796

Webworms injurious to cereal and forage crops and their control by Geo. G. Ainslie. --.
Ainslie, George G. Washington, D.C. : U.S. Dept. of Agriculture, 1922. 16 p. : ill., map --. (NAL Call No.: DNAL Fiche S-70 no.1258).

1797

Feeding behavior, development, and damage by biotypes B, C, and E of *Schizaphis graminum* (Homoptera: Aphididae) on 'Wintermalt' and 'Post' barley.

EVETEX. Peters, D.C. Kerns, D.; Puterka, G.J.; McNew, R. College Park, Md. : Entomological Society of America. Abstract: Greenbug, *Schizaphis graminum* (Rondani), biotypes B, C, and E, were monitored on susceptible 'Wintermalt' and resistant 'Post' barley to determine if there were behavioral differences among the biotypes when they fed on resistant and susceptible barley. All biotype-cultivar combinations were tested by observing feeding behavior, fecundity, and damage. Greenbug biotype B had shorter nonfeeding times on the host plant before probing than did biotypes C and E. Biotype B appeared to be less successful in feeding from the phloem of susceptible 'Wintermalt' but was relatively more successful than biotypes C and E on the resistant 'Post'. Development, reproduction, and particularly dry weight of biotype B were inferior to biotypes C and E on both resistant and susceptible plants. However, biotype B caused the greatest overall reduction in chlorophyll content of infested leaves and should be considered the most virulent but least biologically fit of the three biotypes in these comparisons. Environmental entomology. June 1988. v. 17 (3). p. 503-507. Includes references. (NAL Call No.: DNAL QL461.E532).

1798

Interspecific sexual attraction between *Pyrallis farinalis* L. and *Amyelois transitella* (Walker) (Lepidoptera: Pyralidae) (Minor stored grain pest, ecology).

Landolt, P.J. Curtis, C.E. Lawrence, Kan., The Society. Journal of the Kansas Entomological Society. Apr 1982. v. 55 (2). p. 248-252. ill. Includes 9 ref. (NAL Call No.: 420 K13).

ANIMAL NUTRITION

1799

The effect of genotypes on the feed value of barley straws (North Dakota).

Erickson, D.O. JANS. Meyer, D.W.; Foster, A.E.
Champaign : American Society of Animal Science.
Journal of animal science. Nov 1982. v. 55 (5).
p. 1015-1026. 15 ref. (NAL Call No.: 49 J82).

1800

Effects of pasture management systems on cow-calf productivity on loessial soils in Northeast Louisiana.

LAXBA. Coombs, D.F. Bartleson, J.L.; Rogers, R.L.; Saxton, A.M.; Huffman, D.C.; Alison, M.W.
Baton Rouge, La. : The Station. Bulletin - Louisiana Agricultural Experiment Station. Oct 1989. (815). 19 p. Includes references. (NAL Call No.: DNAL 100 L93 (1)).

1801

Small grain production and utilization in Georgia.

Lee, R.D. Bullock, D.; Segars, W.I.; Bullock, F.; Suber, E.F.; Hudson, R.D.; Thompson, S.S.; Hammond, C.; Johnson, J.T.; Owsley, W.F.
Athens, Ga. : The Service. Bulletin - Cooperative Extension Service, University of Georgia, College of Agriculture. Feb 1986. (929). 18 p. (NAL Call No.: DNAL 275.29 G29B).

1802

Studies on the toxicity of blighted barley to swine /J.U. Christensen, H.C.H. Kernkamp.

Christensen, Jonas Jergon, 1892-. Kernkamp, H. C. H. St. Paul : University Farm, 1935? .
Cover title. 28 p. : ill. ; 23 cm.
Bibliography: p. 27-28. (NAL Call No.: DNAL 100 M66 (3) no.113).

ANIMAL TAXONOMY AND GEOGRAPHY

1803

Annotated list of the insects and mites associated with stored grain and cereal products, and of their arthropod parasites and predators /by R.T. Cotton and N.E. Good.
Cotton, R. T. Good, Newell E. 1905-.
Washington, D.C. : U.S. Dept. of Agriculture, 1937. Caption title.~ Contribution from Bureau of Entomology and Plant Quarantine. 81 p. ; 23 cm. Bibliography: p. 63-74. (NAL Call No.: DNAL 1 Ag84M no.258).

1804

Pests not known to occur in the United States or of limited distribution. 80. A click beetle.
Whittle, K. Hyattsville, Md. : The Service. APHIS 81 - U.S. Department of Agriculture, Animal and Plant Health Inspection Service. Sept 1987. (50). 5 p. ill., maps. Includes references. (NAL Call No.: DNAL aSB599.A3U5).

PESTS OF ANIMALS – GENERAL AND MISC.

1805

Compensation for vertebrate pest damage
(Cereals, forage, livestock, bees, Alberta,
Canada).
Gurba, J.B.PVPCB. Davis, Calif. : University of
California. Proceedings ... Vertebrate Pest
Conference. June 1982. June 1982. (10th). p.
90-94. ill. Includes references. (NAL Call No.:
SB950.A1V4).

ANIMAL DISEASES - FUNGAL

1806

Natural occurrence of the mycotoxin viomellein in barley and the associated quinone-producing penicillia.

Hald, B.APMBA. Christensen, D.H.; Krogh, P.
Washington : American Society for Microbiology.
Applied and environmental microbiology. Dec
1983. v. 46 (6). p. 1311-1317. ill. Includes
references. (NAL Call No.: 448.3 AP5).

ANIMAL DISORDERS – PHYSICAL TRAUMA

1807

Ergots (Fungi, genus *Claviceps*, cereals, feed contamination, mycotoxicity).
Christensen, C.M. Springfield, Va., The Service. PB - U. S. National Technical Information Service. 1980. Literature review. 1980. (80-22-1773). p. 1-44. Bibliography p. 39-44. (NAL Call No.: 157.8 R29).

AQUACULTURE RELATED

1808

Determining vigor of natural and planted stands of sea oats (*Uniola paniculata*) on the Texas Gulf Coast.

Baker, R.L. Dahl, B.E. Austin, Tex., Southwestern Association of Naturalists. The Southwestern naturalist. May 21, 1981. v. 26 (2). p. 117-123. ill. 17 ref. (NAL Call No.: 409.6 S08).

1809

2-chloro-N-((4-methoxy-6-methyl-2,3,5-triazin-2-yl)aminocarbonyl)benzenesulfonamide, a new herbicide (Tested on cereal crop weeds and water hyacinth, *Eichorina crassipes*).

Levitt, G. Ploeg, H.L.; Weigel, R.C. Jr.; Fitzgerald, D.J. Washington, D.C., American Chemical Society. Journal of agricultural and food chemistry. Mar/Apr 1981. v. 29 (2). p. 416-417. 6 ref. (NAL Call No.: 381 J8223).

AGRICULTURAL ENGINEERING

1810

Relation of dust fungicides to flow of small grains through drills and to drill injury /by R.W. Leukel.

Leukel, R. W. 1888-. Washington, D.C. : U.S. Dept. of Agriculture, 1930. Caption title.~ "Contribution from Bureau of Plant Industry.". 10 p. ; 23 cm. Includes bibliographical references. (NAL Call No.: DNAL 1 Ag84C no.119).

FARM EQUIPMENT

1811

Drills and seeders for heavy residues and untilled soils (Small grain planting equipment, minimum tillage farming, Kansas).

Powell, G.M. Manhattan : The Service. L - Cooperative Extension Service, Kansas State University. June 1982. June 1982. (634). 8 p. ill. (NAL Call No.: 275.29 K13LE).

1812

Grain velocity measurement with optical sensors.

Chang, C.S. Martin, C.R.; Lai, F.S. St. Joseph, Mich. : The Society. American Society of Agricultural Engineers (Microfiche collection). Paper presented at the 1985 Winter Meeting of the American Society of Agricultural Engineers. Available for purchase from: The American Society of Agricultural Engineers, Order Dept., 2950 Niles Road, St. Joseph, Michigan 49085. Telephone the Order Dept. at (616) 429-0300 for information and prices. 1985. (fiche no. 85-3551). 22 p. ill. Includes references. (NAL Call No.: DNAL FICHE S-72).

DRAINAGE AND IRRIGATION

1813

Performance of triallate as influenced by incorporation with various pieces of tillage equipment.

WSWPA. Ryerson, D.K. Reno : The Society. Proceedings - Western Society of Weed Science. Paper presented at the annual meeting of the Western Society of Weed Science, March 18-20, 1986, San Diego, California. 1986. v. 39. p. 136-138. (NAL Call No.: DNAL 79.9 W52).

1814

Postemergence herbicide, wild oat control in irrigated, no-till spring barley.

Lish, J.M. Thill, D.C. S.I. : Western Society of Weed Science. Research progress report - Western Society of Weed Science. 1986. p. 200. (NAL Call No.: DNAL 79.9 W52R).

1815

Root-lesion nematodes, potassium deficiency, and prior cover crops as factors in apple replant disease.

JOSHB. Merwin, I.A. Stiles, W.C. Alexandria, Va. : The Society. Growth chamber evaluations of soil from an orchard replant site showed severe stunting of 'Northern Spy' apple (*Malus domestica* Borkh.) seedlings grown in field soil (FS) compared with pasteurized soil (PS) from the same site. The FS : PS seedling dry weight ratio of 0.44 indicated a serious replant problem. Leaf nutrient content was generally higher in PS than FS seedlings. Multiple-regression analysis indicated that leaf K and root-lesion nematode (*Pratylenchus penetrans* Filipjev) primary inoculum accounted for 75% of the variation in FS seedling dry weight. Apple seedling dry weight in FS was 97% of that in PS following a marigold (*Tagetes patula* L. cv. Sparky) cover crop, and 75% following oats (*Avena sativa* L. cv. Saia). Root-lesion nematodes were nearly eliminated from the plots with marigold. Other cover crops and weed-free fallow period were less effective in controlling apple replant disease and/or phytonematodes. Journal of the American Society for Horticultural Science. Sept 1989. v. 114 (5). p. 728-732. Includes references. (NAL Call No.: DNAL 81 S012).

1816

Salinity effects on rye grain yield, quality, vegetative growth, and emergence.

AGUOAT. Francois, L.E. Donovan, T.J.; Lorenz, K.; Maas, E.V. Madison, Wis. : American Society of Agronomy. Although current rye (*Secale cereale* L.) grain production is concentrated mainly in the northern half of the USA and Canada, some rye grain is grown in the arid southwest. Soils in this area are, or have the potential to become, highly saline from the application of saline irrigation water. Since there is nearly a complete lack of information about the response of rye grown under saline

conditions, a 2-yr field plot study was conducted. Six salinity treatments were imposed on a Holtville silty clay (clayey over loamy, montmorillonitic calcareous, hyperthermic Typic Torrifluvent) by irrigating with Colorado River water artificially salinized with NaCl and CaCl₂ (1:1 by weight). Electrical conductivities of the irrigation waters were 1.1, 4.0, 8.0, 12.1, 16.0, and 20.1 dS m⁻¹ the first year, and 1.1, 3.9, 7.5, 11.6, 15.6, and 19.8 dS m⁻¹ the second year. Grain yield and vegetative growth were measured. Relative grain yield of two cultivars, Maton and Bonel, was unaffected up to a soil salinity of 11.4 dS m⁻¹ (electrical conductivity of the saturation extract; k(e)). Each unit increase in salinity above 11.4 dS m⁻¹ reduced yield by 10.8%. These results place rye in the salt-tolerant category. Yield reduction was attributed primarily to reduced spike weight and individual seed weight rather than spike number. Bread quality decreased slightly with increasing levels of salinity. Straw yield was more sensitive to salinity than was grain yield. Plant emergence was determined in greenhouse sand cultures. Both cultivars were slightly less salt tolerant during plant emergence than during subsequent stages of growth. Agronomy journal. Sept/Oct 1989. v. 81 (5). p. 707-712. Includes references. (NAL Call No.: DNAL 4 AM34P).

1817

Subsurface drip irrigation of cotton and small grains.

Tollefson, S. St. Joseph, Mich. : American Society of Agricultural Engineers, c1985. Drip/trickle irrigation in action : proceedings of the Third International Drip/Trickle Irrigation Congress, November 18-21, 1985, Centre Plaza Holiday Inn, Fresno, California, USA. p. 887-895. Includes 7 references. (NAL Call No.: DNAL S612.15 1985).

FOOD SCIENCE, FIELD CROP

1818

In-transit fumigation of truck-ship containers with hydrogen phosphide--a feasibility study (Insect control, stored product pests, cereals).

Jay, E. Davis, R.; Zehner, J.M. New Orleans : The Region. Advances in agricultural technology. AAT-S - United States, Dept. of Agriculture, Agricultural Research Service, Southern Region. Apr 1983. Apr 1983. (28). 13 p. Includes references. (NAL Call No.: aS21.A75U7).

1819

Third international symposium on pre-harvest sprouting in cereals / edited by James E. Kruger and Donald E. LaBerge. -.

Kruger, James E.; LaBerge, Donald E. Boulder, Colo. Westview Press 1983. xii, 312 p. : ill. ; 24 cm. Includes bibliographies and index. (NAL Call No.: SB608.G6T45).

FOOD PROCESSING

1820

Irradiation of grain and grain products for insect control / prepared by Elvin M. Tilton and Stuart O. Nelson . --.

Tilton, Elvin W. Nelson, Stuart Owen, _1927-. Ames, Iowa : Council for Agricultural Science and Technology, 1984. "April 1984" - cover. 6 p. ; 28 cm. --. Bibliography: p. 5-6. (NAL Call No.: DNAL TP371.8.T55).

FOOD STORAGE, FIELD CROP

1821

Biochemical, functional, and nutritive changes during storage (Cereal grains).

Pomeranz, Y. St. Paul, Minn. : American Association of Cereal Chemists, 1982. Storage of cereal grains and their products, 3rd ed. / edited by Clyde M. Christensen. p. 145-217. 13 p. ref. (NAL Call No.: SB190.C43 1982).

1826

Programming and training for small farm grain storage /by Carl J. Lindblad.

Lindblad, Carl J. Washington, D.C. : Peace Corps, Information Collection and Exchange, 1981. "September 1981 ; September 1978.". iii, 100, 8, 1 p. : ill. ; 28 cm. Bibliography: p. 9 (3rd group). (NAL Call No.: DNAL SB190.L54).

1822

Effects of continuous and fractionated doses of gamma radiation on the survival and fertility of Sitophilus granarius (L.).

Jefferies, D.J. New York : Pergamon Press, 1966. The Entomology of radiation disinfestation of grain : a collection of original research papers / edited by P.B. Cornwell. p. 41-56. Includes references. (NAL Call No.: DNAL SB608.G6C6).

1823

In vitro effect of bean amylase inhibitor on insect amylases (Cereals, insect pest control).

Powers, J.R. Culbertson, J.D. Ames, Iowa, International Association of Milk, Food, and Environmental Sanitarians. Journal of food protection. May 1982. v. 45 (7). p. 655-657. ill. Includes 14 ref. (NAL Call No.: 44.8 J824).

1824

The molecular biology of barley storage protein synthesis.

Cameron-Mills, V. Brandt, A.; Ingversen, J. New York, Academic Press, 1980. Cereals for food and beverages : recent progress in cereal chemistry and technology, edited by George E. Inglett, Lars Munck. International Conference on Cereals for Food and Beverages (1979 : Copenhagen). p. 339-364. ill. 33 ref. (NAL Call No.: TX557.I57 1979).

1825

Preservation and storage of grains, seeds, and their by-products cereals, oilseeds, pulses, and animal feed /edited by J.L. Multon ; preface by A.M. Reimbert ; translated from the French by D. Marsh ; reread by A.J. Eydt.

Multon, J. L. 1938-. New York, N.Y. : Lavoisier Pub., c1988. Translated from French.~ Translation of: Conservation et stockage des grains et graines et produits d'origine v'g'tale.~ Originally published: Lavoisier : Technique et documentation, 1982. xxiv, 1095 p. : ill. ; 25 cm. Includes bibliographies and index. (NAL Call No.: DNAL SB190.C6513).

FOOD CONTAMINATION AND TOXICOLOGY

1827

Crop groupings: a survey of its possibilities for deltamethrin registrations.
AECTCV. Mestres, G. Mestres, R. New York, N.Y. : Springer-Verlag. Archives of environmental contamination and toxicology. May 1985. v. 14 (3). p. 321-324. Includes references. (NAL Call No.: DNAL TD172.A7).

1828

HPLC (high performance liquid chromatography)--residue analysis of the herbicide pyridate in cereals.
Lindner, W. Ruckendorfer, H. New York, N.Y. : Gordon and Breach Science Publishers. International journal of environmental analytical chemistry. 1983. v. 16 (3). p. 205-218. Includes references. (NAL Call No.: QH540.I52).

1829

The products of metabolism of (¹⁴C (carbon isotopes)) triadimefon in the grain and in the straw of ripe barley (Fungicides).
Rouchaud, J. Moons, C.; Meyer, J.A. New York. Springer. Bulletin of environmental contamination and toxicology. Oct 1981. v. 27 (4). p. 543-550. Bibliography p. 550. (NAL Call No.: RA1270.P35A1).

1830

Purge and trap method for determination of ethylene dibromide in whole grains, milled grain products, intermediate grain-based foods, and animal feeds.
JANCA2. Heikes, D.L. Arlington, Va. : The Association. Journal of the Association of Official Analytical Chemists. Nov/Dec 1985. v. 68 (6). p. 1108-1111. Includes 13 references. (NAL Call No.: DNAL 381 AS7).

1831

Why the world doesn't want our grain.
McDonald, D. Philadelphia : The Journal. Farm journal. Oct 1985. v. 109 (12). p. 13-17. ill. (NAL Call No.: DNAL 6 F2212).

FOOD CONTAMINATION, FIELD CROP

1832

Aflatoxin decomposition in various soils.
JPFC2. Angle, J.S. New York, N.Y. : Marcel Dekker. Journal of environmental science and health. Part B. Pesticides, food contaminants, and agricultural wastes. 1986. v. 21 (4). p. 277-288. Includes references. (NAL Call No.: DNAL TD172.J61).

1833

Buckwheat browning and color assessment.
CECHAF. Mazza, G. St. Paul, Minn. : American Association of Cereal Chemists. Cereal chemistry. July/Aug 1986. v. 63 (4). p. 361-364. Includes references. (NAL Call No.: DNAL 59.8 C33).

1834

Detection of pesticide residues.
Hascoet, M. New York, N.Y. : Lavoisier Pub., c1988. Preservation and storage of grains, seeds, and their by-products : cereals, oilseeds, pulses, and animal feed / edited by J.L. Multon ; preface by A.M. Reimbert ; translated from French by D. Marsh ; reread by A.J. Eydt. p. 516-526. Includes references. (NAL Call No.: DNAL SB190.C6513).

1835

Effect of atrazine carryover on malting quality of barley (Herbicide, residues).
Brinkman, M.A. Langer, D.K.; Harvey, R.G.; Burger, W.C. Madison, Wis., Crop Science Society of America. Crop science. Nov/Dec 1981. v. 21 (6). p. 973-976. 12 ref. (NAL Call No.: 64.8 C883).

1836

Field and storage fungi.
Christensen, C.M. New York : Von Nostrand Reinhold, 1987. Food and beverage mycology / edited by Larry R. Beuchat. p. 211-232. ill. Includes references. (NAL Call No.: DNAL QR115.F59 1987).

1837

Formation of sodium bisulfite addition products with trichothecenes and alkaline hydrolysis of deoxynivalenol and its sulfonate.
JAFCAU. Young, J.C. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. Sept/Oct 1986. v. 34 (5). p. 919-923. Includes references. (NAL Call No.: DNAL 381 J8223).

1838

High-pressure liquid chromatographic determination of captan, captafol, and folpet residues in plant material (Fruits, grapes, cereals, fungicides).
Buttler, B. Hormann, W.D. Washington, D.C., American Chemical Society. Journal of agricultural and food chemistry. Mar/Apr 1981. v. 29 (2). p. 257-260. ill. 3 ref. (NAL Call No.: 381 J8223).

1839

Production of ochratoxin A in barley by Aspergillus ochraceus and Penicillium viridicatum: effect of fungal growth, time, temperature, and inoculum size (Mycotoxins).
Haggbloom, P. Washington, D.C., American Society for Microbiology. Applied and environmental microbiology. May 1982. v. 43 (5). p. 1205-1207. Includes 10 ref. (NAL Call No.: 448.3 AP5).

1840

Solid-phase radioimmunoassay of ochratoxin A in serum.
JAFCAU. Rousseau, D.M. Slegers, G.A.; Peteghem, C.H. van. Washington, D.C. : American Chemical Society. Journal of agricultural and food chemistry. Sept/Oct 1986. v. 34 (5). p. 862-865. Includes references. (NAL Call No.: DNAL 381 J8223).

1841

Tricalcium phosphate-soybean oil in fortified processed cereals to suppress insects, dusting, and separation.
JFDAZ. Bookwalter, G.N. Highland, H.A.; Warner, K. Chicago, Ill. : Institute of Food Technologists. Journal of food science. Jan/Feb 1985. v. 50 (1). p. 245-248. Includes references. (NAL Call No.: DNAL 389.8 F7322).

1842

Uptake of malathion from galvanized-steel surfaces by stored barley.
White, N.D.G. Abramson, D. College Park, Md. : Entomological Society of America. Journal of economic entomology. Apr 1984. v. 77 (2). p. 289-293. Includes references. (NAL Call No.: 421 J822).

FOOD CONTAMINATION, HORTICULTURAL CROP

1843

High-pressure liquid chromatographic determination of captan, captafol, and folpet residues in plant material (Fruits, grapes, cereals, fungicides).

Buttler, B. Hormann, W.D. Washington, D.C., American Chemical Society. Journal of agricultural and food chemistry. Mar/Apr 1981. v. 29 (2). p. 257-260. ill. 3 ref. (NAL Call No.: 381 J8223).

FOOD COMPOSITION, DAIRY

1844

Field experiments with oats, 1890 / George E. Morrow, Thomas F. Hunt . Milk and butter tests / G.E. Morrow . Cream raising by dilution / G.E. Morrow, E.H. Farrington . The hessian fly / S.A. Forbes . Canada thistles : their extermination / T.J Burrill .
Morrow, G. E. 1840-. Hunt, Thomas F. 1862-;
Morrow, G. E. 1840-; Morrow, G. E. 1840-;
Farrington, E. H. 1860-; Forbes, Stephen Alfred, 1844-1930.; Burrill, Thomas Jonathan, 1839-1916. Champaign, Ill. :
University of Illinois Agricultural Experiment Station, 23 cm. Caption title. p. 353 -388 :
ill. ; 23 cm. (NAL Call No.: DNAL 100 I16S no.12).

FOOD COMPOSITION, FIELD CROP

1845

Biochemical, functional, and nutritive changes during storage (Cereal grains).
Pomeranz, Y. St. Paul, Minn. : American Association of Cereal Chemists, 1982. Storage of cereal grains and their products, 3rd ed. / edited by Clyde M. Christensen. p. 145-217. 13 p. ref. (NAL Call No.: SB190.C43 1982).

1846

Breeding oats for food and feed: conventional and new techniques and materials.
Burrows, V.D. St. Paul, Minn. : American Association of Cereal Chemists, c1986. Oats : chemistry and technology / edited by Francis H. Webster. p. 13-46. ill. Includes references. (NAL Call No.: DNAL TP435.0502).

1847

The effects of preharvest rain.
Derera, N.F. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 1-14. ill., maps. Includes references. (NAL Call No.: DNAL SB189.6.P73).

1848

Modification of nutritional quality by environment and production practices.
AGRYA. Sander, D.H. Allaway, W.H.; Olson, R.A. Madison, Wis. : American Society of Agronomy. Agronomy. 1987. (28). p. 45-82. Includes references. (NAL Call No.: DNAL 4 AM392).

1849

Oat lipids and lipid-related enzymes.
Youngs, V.L. St. Paul, Minn. : American Association of Cereal Chemists, c1986. Oats : chemistry and technology / edited by Francis H. Webster. Literature review. p. 205-226. ill. Includes references. (NAL Call No.: DNAL TP435.0502).

1850

Preharvest sprouting damage and sprouting tolerance: assay methods and instrumentation.
Mares, D.J. Boca Raton, Fla. : CRC Press, 1988. Preharvest field sprouting in cereals / editor, N.F. Derera. Literature review. p. 129-170. ill. Includes references. (NAL Call No.: DNAL SB189.6.P73).

1851

Rapid electrophoresis of oat (Avena sativa L.) prolamins from single seeds for cultivar identification.
CECHAF. Hansen, A.E. Nassuth, A.; Altosaar, I. St. Paul, Minn. : American Association of Cereal Chemists. Cereal chemistry. Mar/Apr 1988. v. 65 (2). p. 153-154. ill. Includes references. (NAL Call No.: DNAL 59.8 C33).

FEED PROCESSING AND STORAGE

1852

Cereal vegetation as a source of forage.

AGRYA. Wedin, W.F. Hoveland, C. Madison, Wis. :
American Society of Agronomy. Agronomy. 1987.
(28). p. 83-99. Includes references. (NAL Call
No.: DNAL 4 AM392).

FEED CONTAMINATION TOXICOLOGY

1853

Ergots (Fungi, genus Claviceps, cereals, feed contamination, mycotoxicity).

Christensen, C.M. Springfield, Va., The Service. PB - U. S. National Technical Information Service. 1980. Literature review. 1980. (80-22-1773). p. 1-44. Bibliography p. 39-44. (NAL Call No.: 157.8 R29).

1854

Natural occurrence of the mycotoxin viomellein in barley and the associated quinone-producing penicillia.

Hald, B.APMBA. Christensen, D.H.; Krogh, P. Washington : American Society for Microbiology. Applied and environmental microbiology. Dec 1983. v. 46 (6). p. 1311-1317. ill. Includes references. (NAL Call No.: 448.3 AP5).

1855

Purge and trap method for determination of ethylene dibromide in whole grains, milled grain products, intermediate grain-based foods, and animal feeds.

JANCA2. Heikes, D.L. Arlington, Va. : The Association. Journal of the Association of Official Analytical Chemists. Nov/Dec 1985. v. 68 (6). p. 1108-1111. Includes 13 references. (NAL Call No.: DNAL 381 AS7).

1856

An uptake on mold inhibitors.

Hagler, W.M. Jr. College Park : The Conference. Proceedings - Maryland Nutrition Conference for Feed Manufacturers. 1983. p. 73-78. Includes references. (NAL Call No.: DNAL 389.9 UN342).

1857

Why the world doesn't want our grain.

McDonald, D. Philadelphia : The Journal. Farm journal. Oct 1985. v. 109 (12). p. 13-17. ill. (NAL Call No.: DNAL 6 F2212).

FEED COMPOSITION

1858

Breeding oats for food and feed: conventional and new techniques and materials.

Burrows, V.D. St. Paul, Minn. : American Association of Cereal Chemists, c1986. Oats : chemistry and technology / edited by Francis H. Webster. p. 13-46. ill. Includes references. (NAL Call No.: DNAL TP435.0502).

1859

The effect of genotypes on the feed value of barley straws (North Dakota).

Erickson, D.O. JANS. Meyer, D.W.; Foster, A.E. Champaign : American Society of Animal Science. Journal of animal science. Nov 1982. v. 55 (5). p. 1015-1026. 15 ref. (NAL Call No.: 49 J82).

1860

Oat lipids and lipid-related enzymes.

Youngs, V.L. St. Paul, Minn. : American Association of Cereal Chemists, c1986. Oats : chemistry and technology / edited by Francis H. Webster. Literature review. p. 205-226. ill. Includes references. (NAL Call No.: DNAL TP435.0502).

PHYSIOLOGY OF HUMAN NUTRITION

1861

Paraquat concentration and renal function in mice fed purified and cereal-based diets (Herbicide toxicity).

Evers, W.D. Hook, J.B.; Bond, J.T. New York : Alan R. Liss. Drug-nutrient interactions. 1983. v. 2 (2). p. 95-104. Includes references. (NAL Call No.: RM302.D76).

POLLUTION

1862

Aflatoxin decomposition in various soils.
JPFC2. Angle, J.S. New York, N.Y. : Marcel Dekker. Journal of environmental science and health. Part B. Pesticides, food contaminants, and agricultural wastes. 1986. v. 21 (4). p. 277-288. Includes references. (NAL Call No.: DNAL TD172.J61).

1863

Buffer capacities of leaves, leaf cells, and leaf cell organelles in relation to fluxes of potentially acidic gases.
PLPHA. Pfanz, H. Heber, U. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. June 1986. v. 81 (2). p. 597-602. Includes 38 references. (NAL Call No.: DNAL 450 P692).

1864

Control of nitrate leaching with winter annual cover crops.
McCracken, D. Lexington, Ky. : The Department. Soil science news & views - Cooperative Extension Service and University of Kentucky. College of Agriculture, Department of Agronomy. 1989. v. 10 (2). 2 p. (NAL Call No.: DNAL S591.55.K4S64).

1865

Effect of gaseous hydrogen chloride on seed germination and early development of seedlings (Tomatoes, barley, injury, air pollution).
Granett, A.L. Taylor, D.C. Alexandria, Va., The Society. Journal of the American Society for Horticultural Science. July 1980. v. 105 (4). p. 548-550. 11 ref. (NAL Call No.: 81 S012).

1866

Effect of small-scale composting of sewage sludge on heavy metal availability to plants (Lactuca sativa, Avena sativa, cadmium, copper, zinc).
Simeoni, L.A. Barbarick, K.A.; Sabey, B.R. Madison, Wis. : American Society of Agronomy. Journal of environmental quality. Apr/June 1984. v. 13 (2). p. 264-268. Includes references. (NAL Call No.: QH540.U6).

1867

Effects of chlorsulfuron on meiosis and seed viability in rye (Secale cereale L.).
WSWPA. Zollinger, R.K. Evans, J.O. Reno : The Society. Proceedings - Western Society of Weed Science. 1985. v. 38. p. 114-119. (NAL Call No.: DNAL 79.9 W52).

1868

Kinetics of the uptake of 14C-labelled chlorinated benzenes from soil by plants.
EESAD. Topp, E. Scheunert, I.; Korte, F. Duluth, Minn. : Academic Press. Ecotoxicology and environmental safety. Apr 1989. v. 17 (2). p. 157-166. Includes references. (NAL Call No.: DNAL QH545.A1E29).

1869

Mutagenicity of selected chemicals in barley test systems (Includes agricultural chemicals, environmental pollutants).
Nilan, R.A. Veleminsky, J. New York, Plenum Press. Environmental science research. 1982. v. 24. p. 291-320. ill. 63 ref. (NAL Call No.: TD172.E55).

1870

Pronamide phytotoxicity in ten Wisconsin soils (Oats).
Dutt, T.E. Harvey, R.G. Champaign, Ill., Weed Science Society of America. Weed science. July 1980. v. 28 (4). p. 429-432. ill. 15 ref. (NAL Call No.: 79.8 W41).

MATHEMATICS AND STATISTICS

1871

Leaf water potential, relative water content, and diffusive resistance as screening techniques for drought resistance in barley. AGJOAT. Matin, M.A. Brown, J.H.; Ferguson, H. Madison, Wis. : American Society of Agronomy. Rapid drought resistance screening techniques could accelerate selection of improved cultivars for semiarid areas. This study was conducted to determine if total leaf water potential, leaf relative water content, and leaf diffusive resistance could be used to differentiate between barley (*Hordeum vulgare* L.) cultivars differing in apparent drought resistance. Resistant and susceptible cultivars were selected by regressing individual cultivar yields against site mean yield for numerous yield trials. Individual plants were grown in cone-tainers (plastic tubes 25 mm in diameter and 123 mm long). The lower 30 mm of the cone-tainers were imbedded in sand so that roots emerging from the bottom of the cone-tainers were in the capillary fringe above a water table in the sand. Fifty plants were grown at a time in a sand bed, five replications each of five resistant and five susceptible barley cultivars of either two- or six-row head types. At the three-leaf growth stage total leaf water potential was measured on the third leaf of each plant under low stress conditions (predawn). Water was then drained from the sand bed to stress the plants. Leaf diffusive resistance (LDR) was measured at 1000 and 1300 h each day as stress developed. When the PM diffusive resistance reached about 1000 s m⁻¹, relative water content (RWC) of the second leaf was determined. All measurements were repeated in three different trials for each head type. Relative water content and total leaf water potential differentiated between drought-resistant and drought-susceptible groups of two- and six-rowed barley at the 0.01 probability level in every trial. Leaf diffusive resistance differentiated between resistant and susceptible groups at the 0.01 or 0.05 level under low and high stress conditions. Neither RWC nor LDR allowed separation of cultivars within resistant or susceptible groups, but total leaf water potential did show differences between cultivars within groups. These techniques provide an ea. Agronomy journal. Jan/Feb 1989. v. 81 (1). p. 100-105. Includes references. (NAL Call No.: DNAL 4 AM34P).

1872

A mathematical model for evaluating insect resistance in plants. JKESA. Webster, J.A. Inayatullah, C.; Fargo, W.S. Lawrence, Kan. : The Society. Journal of the Kansas Entomological Society. Includes abstract. July 1985. v. 58 (3). p. 564. (NAL Call No.: DNAL 420 K13).

1873

Measurement of thermally induced pressures in a model grain bin. TAAEA. Manbeck, H.B. Muzzello, L.M. St. Joseph, Mich. : The Society. Transactions of the ASAE - American Society of Agricultural Engineers. July/Aug 1985. v. 28 (4). p. 1253-1258, 1264. Includes references. (NAL Call No.: DNAL 290.9 AM32T).

1874

Measuring disease progress in pure and mixed stands of plant cultivars. PHYTA. Gumpert, F.M. St. Paul, Minn. : American Phytopathological Society. A measure is presented that quantifies disease progress of fungal pathotypes during the exponential phase of the epidemic. This measure, referred to as rate of disease increase (p), represents the factor by which the number of infection units is multiplied from one day to another after the initial infection waves have damped. The rate of disease increase depends on four pathotype/cultivar-specific parameters (latent period, infectious period, infection efficiency, spore production rate) and two nonspecific parameters (deposition frequency and autodeposition frequency). Three applications are given. First, it is used to determine the levels of partial resistance of barley cultivars to leaf rust. The calculated p values were highly correlated both with total spore production per unit leaf area ($r = 0.97$) and with disease score in the field ($r = 0.81$). The rate of disease increase is particularly useful if the epidemiological parameters display compensatory effects. The second application deals with the long-term composition of pathotype mixtures. The pathotype with the largest p value will predominate in the long run. This is again exemplified by using barley leaf rust data. The third application concerns disease control strategies. A host stand should be composed so that the corresponding predominant pathotype has a p value smaller than the prevailing pathotype of any other composition. This concept does not presuppose selection against unnecessary genes for virulence. A condition is given under which a cultivar mixture may be more beneficial than each of its components grown in pure stands, and this condition is illustrated by a simple example of two pathotypes and two cultivars. Phytopathology. Sept 1989. v. 79 (9). p. 969-973. Includes references. (NAL Call No.: DNAL 464.8 P56).

1875

Modeling weed populations in cereals. Cousens, R. Moss, S.R.; Cussans, G.W.; Wilson, B.J. Champaign, Ill. : Weed Science Society of America. Reviews of weed science. Literature review. 1987. v. 3. p. 93-112. ill. Includes references. (NAL Call No.: DNAL SB610.R47).

(MATHEMATICS AND STATISTICS)

1876

Models explaining the specificity and durability of host resistance derived from the observations on the barley-Puccinia hordei system.

Parlevliet, J.E.NASSD. New York : Plenum Press.
NATO advanced study institutes series. Series
A. Life sciences. 1983. v. 55. p. 57-80.
Includes references. (NAL Call No.: QH301.N32).

1877

The modification of the foliage structure and radiation absorption in a plant canopy by wind.

Hipps, L.E. Baiden, E. Boston : The Society,
1985. 17th Conference on Agricultural and
Forest Meteorology and seventh Conference on
Biometeorology and Aerobiology, May 21-24,
1985, Scottsdale, Ariz. : preprint volume /
sponsored by the American Meteorological
Society. p. 127-128. Includes references. (NAL
Call No.: DNAL S600.2.C6 1985).

1878

Student's method for interpreting paired experiments.

AGUOAT. Love, H.H. Brunson, A.M. Madison, Wis.
: American Society of Agronomy. Too close an
analogy should not be drawn between the
determination of exact constants and the
results of comparative agronomic experiments.
Bessel's and Peter's formulas are not adapted
to calculating the degree of significance of a
difference between two varieties or two
comparative treatments in experiments in which
the variability within each variety or
treatment is high because they extend through a
series of years or because of place effect. For
observations which naturally arrange themselves
in pairs, Student's method is a better method
with which to determine the probability of the
difference, and deserves a larger application
in the interpretation of agronomic
investigations. Agronomy journal. Jan 1924. v.
16 (1). p. 60-68. (NAL Call No.: DNAL 4 AM34P).

DOCUMENTATION

1879

A model for the regulation of K⁺ influx, and tissue potassium concentrations by negative feedback effects upon plasmalemma influx.
PLPHA. Siddiqi, M.Y. Glass, A.D.M. Rockville, Md. : American Society of Plant Physiologists. Plant physiology. May 1986. v. 81 (1). p. 1-7. Includes 32 references. (NAL Call No.: DNAL 450 P692).

1880

The rodent problem in Latin America.

Elias, D.J. Fall, M.W. Boca Raton, Fla. : CRC Press, c1988. Rodent pest management / editor, Ishwar Prakash. Literature review. p. 13-28. maps. Includes references. (NAL Call No.: DNAL SB994.R6R65).

AUTHOR INDEX

- . 383
AAREEZ. 88, 1727, 96, 1494
ABBIA. 604, 737
Abernethy, R.H. 797, 1559, 771, 1551, 88, 1727
Abramson, D. 1430, 1685, 1842
Acklin, W. 416, 1219
ACSMC. 420, 387, 919, 1214
ACSTD. 591, 1266
Adams, D. 805
Adams, P.A. 653
Adams, W.A. 1300
Adkins, S.W. 1567
AECTCV. 1620, 1827
AESAA. 241, 873
AGABB. 164, 1706
AGBOB. 57, 1751, 1776
AGBOBO. 33, 1715
AGJOAT. 97, 1732, 767, 1816, 803, 398, 1557, 1765, 219, 552, 1713, 51, 200, 290, 1132, 124, 678, 1737, 160, 402, 1878, 122, 44, 1707, 390, 1183, 490, 749, 1741, 117, 400, 783, 1366, 339, 1161, 399, 781, 34, 584, 161, 791, 1745, 340, 1162, 485, 1702, 484, 1701, 305, 1137, 81, 230, 36, 1572, 1362, 1682, 40, 1764, 663, 1871, 82, 61, 1286, 102, 261, 619, 62, 545, 459, 1061, 1554, 109, 283, 1517, 127, 1652, 133, 486, 695, 157, 491, 1743
AGREA. 2, 1555
AGRYA. 45, 503, 54, 1852, 123, 1848, 197, 838, 953, 191, 723, 704, 822
Ahlquist, P. 423, 1251
Ahmadi, M. 80, 1721
Ahokas, H. 215, 538
Ahrenst-Larsen, B. 341, 735
Ahring, R.M. 621
Ahuja, S.L. 180, 181, 182
Ainslie, George G. 936, 1796
Ainslie, George G. 1886-1930. 935, 1795
Aist, J.R. 510, 985, 988
AJBDA. 637, 444, 743, 446
AJBOAA. 529, 433, 521
Akey, W.C. 590, 1489, 1479
Akimoto, I. 589
Alam, S.M. 1300
Alberte, R.S. 741
Albrechtsen, R.S. 108, 267, 626
Alcock, K.T. 1033, 1008
Alexander, K. 583
Alfinito, S.H. 433, 521
Ali, Mohammed Abdul, 664
Alison, M.W. 27, 174, 1800
Allard, R.W. 259, 1111, 1000, 393, 1186, 311, 684, 1533, 1112, 1090, 258, 1310, 1640, 250, 1108
Allaway, W.H. 123, 1848
Allen, H.T. 359
Allen, M. 172, 1774
Alley, H.P. 46, 1280, 1617, 1556, 1602, 176, 1535
Allinson, D.W. 96, 1494, 116, 481
Allison, R.V. 44, 1707
Altosaar, I. 344, 824, 1851
Amadori, E. 1506, 1636
Amaral, A.C. 748
Ambach, R.M. 1304, 1490
Amrhein, N. 1353, 1669, 1354, 1670
Amy, P.S. 1232
Andersen, K. 269, 946
Andersen, S. 269, 946
Anderson, M.K. 191
Andres, M.W. 192, 983, 396, 1196
Andrews, C.J. 1320, 760
Angle, J.S. 975, 1832, 1862
Anikstesr, Y. 389, 1181
Aniol, A. 184, 499
APMBA. 1232, 1621, 1690, 425, 1806, 1854
Appleby, A.P. 1657, 1703, 1319, 1648, 1361, 1680, 1581
Appleton, W.H. 490, 749, 1741
Argandona, V.H. 387, 919
Arigoni, D. 416, 1219
Arioka, K. 573
Armiger, W.H. 219, 552, 1713
Armstrong, E. 1403
Arnold, R.N. 3, 76, 229
Arnold, W. E. 1578
Arnold, W.E. 1460
Arny, D.C. 7, 244, 958, 1206
Arthur, J. C. 1850-. 1164
Arvinte, T. 802
Ashford, R. 1364, 1683, 1304, 1490
Aslam, M. 597, 558, 649
Aspinall, D. 532, 648
Atsmon, D. 568
au, C. R. 910
Aulerich, R.J. 1635
Ayanru, Dafe Kpigere Gboriavhien, . 1073
Ayers, J.E. 1190
Ayres, P.G. 585, 1076
Ba-Angood, S.A. 931
Baameur, A. 1436
Baenziger, P.S. 237, 315, 1144, 248, 1107
Bahler, C.C. 85, 871, 1482
Baiden, E. 5, 439, 1877
Bain, A.D. 987, 1373
Baker, J.E. 1419
Baker, L.S. 1315
Baker, R.J. 240
Baker, R.L. 541, 1808
Bakke, Arthur Laurence, b. 1886. 605, 1503
Bale, Shankarappa S. 1358
Baligar, V.C. 463, 1722
Balke, N.E. 548, 420, 1314, 1643, 755, 1365, 1686
Ballesteros, A. 604
Ballinger, D.J. 1056, 1058, 1028, 1055, 1027, 1057
Balsamo, R.A. 802
Baltenberger, D.C.C. 102, 261, 619
Baltenberger, D.E. 347, 1255
Bamatraf, A.M. 141, 1330, 155
Banks, P.A. 166, 1573, 1770
Banting, J.D. 1512
Banttari, E.E. 1093
Barbarick, K.A. 1299, 1866
Barbetti, M.J. 1020, 1018, 1019
Barbour, N.W. 245, 609
Barker, G.L. 1434

AUTHOR INDEX

- Barnes, J.P. 419, 761, 1349, 1307, 429, 1351, 1763
 Barnett, R.D. 12, 353, 1171, 100
 Barney, A.F. 290, 1132
 Baron, O. 551
 Barrett, J. 242, 1080
 Barrett, J.A. 1185, 327, 1148, 201, 986
 Barrett, M. 1341, 1660
 Bartels, P.G. 1274, 1614
 Bartleson, J.L. 27, 174, 1800
 Bassett, E.N. 1187, 1761
 Bassi, A. 1188
 Bastide, J. 808, 1565
 Bauer, F.J. 1384
 Baumgartner, B.J. 337, 731
 Bayer, David E. 1435
 Baysdorfer, C. 752
 BBRCA. 748
 BCYCDK. 534, 1711
 Beachy, R.N. 756
 Beardmore, R.A. 1362, 1682
 Becker, J.O. 9, 1136
 Beeman, R.W. 1393
 Behrens, R. 1580
 Behrens, Richard. 1579
 Beier, R. 1203
 Belanger, A. 78, 461
 Belanger, F.C. 436, 629
 Belhadri, M. 292, 655
 Bell, M. A. 1897-. 83
 Beltran, J.P. 1203
 Ben Mordechai, Y. 1388
 Bendixen, L.E. 1302, 1633
 Bennett, O.L. 463, 1722
 Bent, A.F. 279, 644
 Benton, Curtis, 1898-. 55, 863
 Bernal-Lugo, I. 756
 Bernard, R.L. 392
 Bestman, H.D. 1524
 Beuerlein, J.E. 80, 1721
 Biederbeck, V.O. 68, 1691, 1696
 Bissonnette, H.L. 984
 Bjorn, S.E. 204, 999
 Blackwell, B.A. 987, 1373
 Blake, T.K. 295, 668, 268, 1115
 Blaylock, M.J. 480, 1268
 Bleavins, M.R. 1635
 Blevins, R.L. 455, 1698, 1777
 Bloom, A.J. 487, 708, 766, 281, 478, 646, 550
 Bloom, W.W. 747
 Blum, A. 705
 Blume, D.E. 546, 1624
 Boal, R. 703
 Bockelman, H.E. 366, 373, 369, 1175, 364, 297
 Boerboom, Marvin L. 233, 574
 BOGAA. 764
 Bohl, W.H. 88, 1727
 Bolley, H.L. 36, 1572
 Bollinger, D.J. 1054
 Bond, J.T. 1654, 1861
 Bonilla, C.A. 603
 Bookwalter, G.N. 1426, 1841
 Booth, Ernest Gordon, 1899-. 540
 Boozaya-Angoon, D. 287, 896
 Boquet, D.J. 333, 1152, 1228
 Borden, J.H. 1418
 Born, W.H. vanden. 441, 683, 1532
 Bothmer, R. von. 704, 822
 Boulif, M. 1130
 Bowman, D.T. 118, 300, 963, 119, 301, 964
 Bowren, K.E. 68, 1691, 1696
 Bowyer, J.R. 1325, 1650
 Boyd, W.J.R. 328, 1151
 Boyle, J.G. 747
 Bracken, A.F. 117
 Bracker, C.E. 1258
 Branchard, M. 275, 1122
 Brandenburg, R.L. 858
 Brandt, A. 681, 1824
 Brauen, S.E. 42, 185, 951
 Brauer, D. 516
 Braunworth, W.S. Jr. 1540
 Bregitzer, P. 217, 542
 Bregitzer, P.P. 313, 147, 335, 718
 Breidenbach, R.W. 467, 581
 Breman, J.W. 73
 Brent, K.J. 1178
 Brentzel, W. E. 1889-. 971
 Breslin, W.J. 1635
 Brewbaker, H. E. 1895-. 788
 Brewster, B.D. 1361, 1680
 Briggie, L.W. 409, 800
 Briggs, Fred N., 1896-. 1218
 Briggs, K.G. 240
 Briggs, S. P. 854, 1782
 Briggs, S.P. 1197, 1198
 Briggs, W.R. 721, 641, 421, 569
 Brink, G.E. 49, 505, 1750
 Brinkman, M.A. 1628, 1835, 1347, 1666
 Britton, G. 1342, 1662
 Brock, T.G. 438, 666, 529, 497, 637
 Brodl, M.R. 436, 629
 Broscious, S.C. 1190
 Brown, A.E. 1094
 Brown, A.H.D. 798
 Brown, A.R. 22, 378, 1256, 361, 114
 Brown, C.M. 137, 325, 701
 Brown, D.J. 665
 Brown, G.N. 787
 Brown, J.C. 453, 562, 1716, 492, 480, 1268
 Brown, J.H. 663, 1871
 Brown, K.C. 872
 Brown, S.M. 1516
 Browning, J.A. 25, 381, 1177, 19, 365, 1174, 1046
 Browning, J.A. 24, 380, 1176
 Bruckner, P.L. 22, 378, 1256
 Bruggers, R.L. 846
 Brunson, A.M. 160, 402, 1878
 Brusko, M. 1609, 1775
 Bryan, M.D. 1156
 Buchenauer, H. 570
 Buehring, N.W. 1434
 Buhl, M.B. 582
 Bukhov, N.G. 611
 Bullock, D. 156, 1801
 Bullock, F. 156, 1801
 Bullock, F.D. 1461
 Buly, R. 1618
 Buman, R.A. 797, 1559, 771, 1551
 Burger, W.C. 1628, 1835
 Burke, B.A. 419
 Burkholder, W.E. 1393, 1391
 Burleigh, J.R. 1077, 1488
 Burns, G.R. 801, 1747
 Burnside, O.C. 109, 283, 1517, 127, 1652, 1574, 1771, 1470
 Burrill, L.C. 1540
 Burrill, Thomas Jonathan, 1839-. 98, 878, 1505, 1844
 Burrows, V.D. 199, 1846, 1858
 Burton, R. 899, 110, 901, 900
 Bush, D.R. 702, 1227
 Bush, L. 216, 866
 Bushnell, William R. 53, 991, 1374
 Buttery, R.G. 697, 909

AUTHOR INDEX

- Buttler, B. 1644, 1838, 1843
 Buttner, M.P. 1232
 Byers, R.A. 85, 871, 1482
 Cadenas, R.A. 537
 CAGRA. 1238
 Caldwell, C.D. 67, 1714
 Caldwell, C.R. 630
 Caldwell, R.M. 487, 708, 766
 Callihan, R.C. 1452
 Callihan, R.H. 1560, 1768, 1451, 1510, 1596, 1492
 Cameron-Mills, V. 681, 1824
 Cameron, R.S. 1271
 Camilleri, P. 1325, 1650
 Camm, E.L. 494
 Cammaerts, D. 401
 Campbell, C.A. 68, 1691, 1696
 Campbell, C.G. 1464, 351
 Campbell, K.W. 23, 379, 14, 356, 1173
 Campbell, R.K. 853
 Campbell, W.F. 141, 1330, 155
 Camper, H.M. Jr. 383
 Carceller, M. 537
 Carlson, D.R. 1574, 1771, 1470
 Carlson, G. 87, 1072, 1725
 Carr, R.H. 124, 678, 1737
 Carroll, T.W. 11, 350, 218, 1242, 1237
 Carter, J.V. 553
 Cartwright, B.O. 853
 Carver, B.F. 13, 354, 917
 Carver, George Washington, 1864?- . 41, 1563
 Carver, T.L.W. 1150
 Cary, E.E. 465, 1263, 466
 Cassman, K.G. 473, 1733
 Cates, R.L. Jr. 86, 1724
 Cattoir-Reynaerts, A. 401
 CECHAF. 344, 824, 1851, 1372, 1833
 Center, O. D. 1872-. 121, 1139
 Cha, J.W. 457, 1297
 Chada, Harvey L. 884
 Chafai El Alaoui, A. 593
 Chamberlain, K. 1121
 Chamberlin, Thomas R. 1889-. 169, 937
 Chambers, G. R. 162
 Chan, K. 328, 1151
 Chandler, P.C. 418, 814
 Chang, C.S. 1376, 1812
 Chapin, F.S. III. 550
 Chapin, J.W. 898
 Chapko, L.B. 71, 225, 434
 Chedester, L.D. 875, 852
 Chen, K.H. 745
 Chen, Y.G. 643, 1735
 Cheng, C.L. 206, 522
 Cheng, H.C. 711
 Cherif, M. 366
 Ching, T.M. 643, 1735, 489, 722, 1738, 786, 1744
 Chinnici, M.F. 1360
 Cho, B.H. 280, 1124
 Choo, T.M. 128, 317, 688
 Chow, P.N.P. 1688, 1594
 Chowdhury, Ikbalur Rashid, . 422, 575
 Christ, B.J. 1166
 Christensen, C.M. 1095, 1375, 1836, 1140, 1378, 1079, 1807, 1853
 Christensen, D.H. 425, 1806, 1854
 Christensen, Jonas Jergon, 1892-. 1208, 1802
 Christianson, M.L. 433, 521
 Clay, S.A. 154, 1348, 1667
 Clayton, J.L. 830, 828, 829
 Cleland, R.E. 528, 662, 775, 748, 759, 709, 1147
 Cloninger, C. K. 1182
 Close, R.C. 1078
 Cloutier, Y. 557, 804
 Cochran, V.L. 154, 1348, 1667
 Coffelt, J.A. 1417
 Coffin, R.H. 72, 563
 Coffman, Franklin A. 1892-. 171, 63, 955, 106, 960, 424, 606, 1097
 Cohen, A.S. 1288, 1626
 Cohen, J.D. 745
 Cole, H. Jr. 1245, 93, 1074, 1190
 Coleman-Harrell, M.E. 1590
 Collins, F. 1509
 Collins, F.C. 1188, 1193, 1037, 1065, 1038
 Colville, D.C. 62, 545
 Colwick, R.F. 1434
 Colyer, P.D. 333, 1152, 1228
 Comeau, A. 251, 1249
 Conover, M.R. 839, 842
 Cook, R.J. 9, 1136, 1187, 1761
 Cooke, B.K. 1044
 Cooke, Wells Woodbridge, 1858-. 1199, 1677, 1742
 Coombs, D.F. 27, 174, 1800
 Copeland, L. O. 135, 50
 Copeland, L.O. 48
 Coppock, S. 899, 110, 901, 900
 Corcuera, L.J. 387, 919
 Cordon, C. 1638
 Corke, H. 568
 Cornish, P.S. 780
 Coste, C. 808, 1565
 Costel, G.L. 1602
 Costello, S.R. 1584
 Cotton, R. T. 1386, 1781, 1803
 Cousens, R. 1531, 1875
 Creamer, R. 1238
 Cribb, D.M. 583
 Criddle, R.S. 467, 581
 Crocker, R.L. 911
 Crome, M.G. 1026
 Crookston, R.K. 593
 Crosby, W.L. 278, 1267
 Crosslin, J.M. 11, 350
 Crowley, D.E. 810
 CRPSA. 238, 262, 620, 237, 240
 CRPSAY. 186, 978, 222, 1048, 1509, 193, 1239, 404, 793, 382, 376, 284, 895, 367, 313, 292, 655, 217, 542, 705, 202, 511, 13, 354, 917, 12, 353, 1171, 179, 392, 15, 357, 1344, 22, 378, 1256, 303, 347, 1255, 393, 1186, 25, 381, 1177, 24, 380, 1176, 23, 379, 18, 362, 1229, 593, 712, 366, 371, 383, 359, 108, 267, 626, 370, 91, 239, 1730, 128, 317, 688, 11, 350, 10, 349, 21, 374, 17, 360, 20, 372, 192, 983, 351, 71, 225, 434, 295, 668, 49, 505, 1750, 112, 658, 489, 722, 1738, 318, 1145, 226, 564, 221, 554, 786, 1744, 600, 1306, 1634, 147, 335, 718, 389, 1181, 134, 324, 696, 396, 1196, 19, 365, 1174, 14, 356, 1173, 16, 358, 1060, 368, 363, 361, 352, 386, 754, 254, 614, 355, 1172, 103, 264, 622, 373, 369, 1175, 364, 388, 1180
 CRPTD6. 1044
 CRSOA. 449, 1607
 CRYBA. 557
 CSOSA. 479, 1693
 CSOSA2. 111, 1736, 67, 1714, 462, 1699, 1720, 78, 461, 116, 481, 463, 1722
 Cudney, D. 1469
 Cudney, D.W. 1586, 1466, 1608, 1436
 Culbertson, J.D. 1402, 1823
 Cullum, R.F. 57, 1751, 1776
 Cunfer, B.M. 361, 114

AUTHOR INDEX

- Curran, W.S. 1474
 Curtis, C.E. 1407, 1798
 Curtis, R.E. 1552, 1553
 Cussans, G.W. 1531, 1875
 Cutkomp, L.K. 1414, 1413
 Cyr, K.L. 679
 Czaplewski, S.J. 103, 264, 622
 D'Arcy, C.U. 1241
 Daft, J.L. 1339
 Dahl, B.E. 541, 1808
 Dailey, J.E. 631
 Dalling, M.J. 334, 717
 Daniels, N.E. 852
 Davidonia, G.H. 1618
 Davis, M.P. 420
 Davis, R. 31, 1401, 1818
 Day, A.D. 534, 1711, 10, 349, 236, 1726
 De Boer, A.H. 775
 De Maggio, A.E. 737
 De St Remy, E.A. 30, 417, 1568
 Dean, J.G. 175
 Deane-Drummond, C.E. 693
 deBoer, A.H. 662
 Deckard, E.L. 74, 458, 1718
 Deckard, Edward Lee, 1943-. 1298
 Deibert, E.J. 398, 1557, 1765
 Dell'Agnola, G. 1279
 Delserone, L.M. 1245, 93, 1074
 Delucchi, G.M. 144, 331
 Denney, A. 474, 612
 Derera, N.F. 198, 1284, 4, 1303, 1847
 Detling, J.K. 603
 Deus, B. 1353, 1669
 DeVault, G. 1585, 1773
 Devine, M.D. 1524
 Dewdney, J. 206, 522
 DeWitt, Jerald R. 1663
 Dhindsa, R.S. 748
 Dial, M.J. 1543, 1600, 1453, 1589, 1597, 1472, 1454
 Dickson, D.W. 129, 948
 Dietz, S. M. 1893-. 120, 1138
 Dill-Macky, R. 1207
 Dirks, R. 401
 Ditz, K.M. 262, 620, 624, 623, 405, 794, 265, 625
 Doersch, R. E. 844
 Doersch, R.E. 833
 Dofing, S.M. 16, 358, 386, 754, 291, 437, 650
 Dolezal, W.E. 1065
 Dolezal, W.E. 1193, 1037
 Dolferus, R. 401
 Doll, J. D. 910, 844, 146, 1655
 Doll, J.D. 833
 Donald, W.W. 1493
 Donovan, T.J. 767, 1816
 Douglas, S.M. 1069
 Dowgert, M.F. 672, 671, 561
 Downard, R.W. 1521, 1481
 Dregne, H. E. 95, 1731
 Dreier, A.F. 16, 358
 Drummond, D.P. 1495
 Drummond, W.D. 1604
 Drury, R.E. 720
 Drye, C.E. 1047
 Dubbs, A.L. 238
 Duffus, C.M. 716
 Duffy, P.D. 26, 79
 Dumas, T. 1631
 DuPont, F.M. 665, 746, 587, 687
 Durbin, R.D. 592, 1222, 776
 Durrell, L. W. 1888-. 1002
 Dutt, T.E. 1340, 1659, 1870
 Duysen, M.E. 1625
 Dyer, M.I. 603
 Dyer, W. 29, 1526
 Dyer, W.E. 1546
 Dyson, D.H. 367, 20, 372
 Eberlein, C.V. 1515
 ECBDA. 427, 738, 823
 Edgington, L.V. 1562, 1681, 1485, 524
 Edwards, C.R. 856
 Edwards, L.H. 13, 354, 917, 287, 896
 Edwards, L.M. 158, 1766, 1779
 Edwards, Richard C. & Field crops insects. 857
 EESAD. 660, 1868, 1647
 Effron, D. 885
 Egli, D.B. 516
 Ehlers, W. 803
 Eikenbary, R.D. 209, 864, 853
 El-deek, M.H. 651, 1519
 Eldredge, John C. 1886-. 1295
 Elfman, B. 494
 Elias, D.J. 845, 1383, 1880
 Elich, T.D. 819
 Ellerbrock, L.A. 1486
 Elliott, C.C.H. 843
 Elliott, Charlotte, 1883-. 782
 Elliott, N.C. 865
 Ellis, S.E. 1165, 298, 947
 Elseewi, A.A. 1327
 Elseewi, Ahmed Abd Elhamid Mohamed, . 451, 517
 Emara, Y.A. 224, 1052
 Epel, B.L. 551
 Epstein, A.H. 1096
 Erdal, K. 341, 735
 Erickson, D.O. 227, 1799, 1859
 Eslick, R.F. 112, 658, 226, 564, 373, 369, 1175, 364, 238, 297
 Estes, P.M. 848
 ETOCDK. 806, 1684
 Ettounsi, L. 165, 1571
 Evans, D.A. 316
 Evans, J.O. 1357, 1679, 1767, 1346, 1665, 1760, 1521, 1481, 1301, 1632, 1867
 Evans, R.M. 1317, 1646
 Evans, W.F. 72, 563
 Evers, W.D. 1654, 1861
 Everson, E.H. 406, 796, 1359, 375
 EVETB. 872, 209, 864
 EVETEX. 1418, 877, 1246, 876, 1783, 1797, 893, 1755, 266, 883, 874
 Evins, Warren H., . 635
 Ewing, A.L. 757
 Falk, B.W. 1238
 Fall, M.W. 845, 1383, 1880
 Famelaer, Y. 401
 Fant, G. W. 1036
 Fargo, W.S. 906, 1872
 Farrington, E. H. 1860-. 98, 878, 1505, 1844
 Farris, T.N. 853
 Fay, P. 29, 1526
 Fay, P.K. 520, 1459, 1546, 1536
 Fay, Peter K., . 1545
 Feierabend, J. 689, 1653
 Fellows, G.M. 520, 1459
 Fenster, W. E. 162
 Ferguson, H. 663, 1871, 61, 1286, 86, 1724
 Fernandez, J.A. 869
 Fernandez, V.M. 604
 Fernando, M. 789
 Ferrari, G. 1279
 Ferrell, M.A. 1467
 FETMA. 1417
 Finazzo, J. 487, 708, 281, 478, 646
 Fincher, G.B. 308, 440, 680

AUTHOR INDEX

- Finckh, B.F. 1687
 Finkner, Verne C. 256, 1110
 Fischer, M.L. 621
 Fisher, G. 886
 Fitzgerald, D.J. 1610, 1689, 1809
 Fleming, A.L. 1260
 Fletcher, J.H. 73
 Flick, C.E. 316
 Flint, M.L. 837, 1769
 Flint, W. P. 1882-. 55, 863
 Flom, D.G. 1560, 1768
 Flores, H.E. 1337, 707, 706
 Fluke, C. L. 1891-. 169, 937
 Flyer, J. 1396
 FNETD. 1035, 1016, 1005, 1034, 1004, 1015, 1086, 1087, 1056, 1058, 996, 1081, 998, 1032, 1029, 1085, 1014, 1023, 1067, 1089, 1041, 1064, 1210, 1020, 1054
 Foley, M.E. 1483, 520, 1459
 Folsom, Donald, 1891-. 1356
 Forbes, Stephen Alfred, 1844-1930. 98, 878, 1505, 1844
 Fornari, C.S. 208, 527
 Foroughi-Wehr, B. 276, 1123, 1252
 Forster, R.L. 1195, 1231
 Fortin J.A. 579, 1592
 Foster, A.E. 1060, 227, 1799, 1859
 Foster, E. 45, 503
 Foster, J.E. 193, 1239, 299, 905, 1253, 347, 1255, 850, 1236
 Fowke, L.C. 431, 512, 1457
 Foy, C.D. 164, 1706, 453, 562, 1716, 219, 552, 1713, 801, 1747, 1323, 1350, 409, 800, 1287, 1695, 1260
 Foy, C.L. 1291, 1476, 1627
 Foy, Chester L. 1435
 Franckowiak, J.D. 274
 Francois, L.E. 767, 1816
 Frank, J.A. 1240, 1166, 1245, 1190
 Frantz, T.A. 776
 Freake, G.W. 224, 1052
 Freckleton, M.E. 6, 149
 Freed, R.D. 404, 793, 48
 Freedman, B. 1391
 French, R. 1251
 Frey, K.J. 376, 1096, 25, 381, 1177, 24, 380, 1176, 102, 261, 619, 62, 545, 134, 324, 696, 19, 365, 1174, 188, 980, 214, 314
 Friedman, W. 950, 949
 Friedrich, J.W. 713
 Friedt, W. 257, 1250, 276, 1123, 1252
 Friesen, G.H. 1367, 1588, 43, 1431, 163, 1569, 1464, 1468
 Friesner, D. 172, 1774
 Fromme, F. D. 1886-. 1021
 Frost, S. 145, 332
 Frye, W.W. 455, 1698, 1777
 Fuhr, F. 1214, 501, 1616
 Fuhremann, T.W. 1619
 Fujii, S. 573
 Fukuyama, T. 415, 812, 414, 1216
 Fuller, Wallace Hamilton. 493
 Fulton, Winston Cordell. 868
 Furian, V. 579, 1692
 Gabor, B.K. 411, 1213
 Gaines, E. F. 1886-. 1226
 Gaines, E.F. 390, 1183
 Gallagher, L.W. 292, 655
 Gallaher, R.N. 1584, 129, 948
 Galston, A.W. 1285, 1337, 719, 707, 706, 740, 1368, 599, 1305, 817, 1369
 Gamble, P.E. 639
 Gambrell, R.H. 21, 374
 Gantz, R.L. 1507
 Garbarino, J. 746, 687
 Garber, R.J. 339, 1161
 Garber, Ralph John, 1890-. 1126
 Garcia del Moral, L.F. 407, 799
 Garcia-Martinez, J.L. 805
 Gardner, R.D. 1411
 Garger, S.J. 433, 521
 GARRA. 114, 166, 1573, 1770
 Gassman, M.L. 792
 Gaur, B.K. 677
 Gawronski, S. 306, 1322, 1649
 Geballe, G.T. 1368, 599, 1305, 817, 1369
 Gehrke, P. 1353, 1669
 Gellner, J.L. 247, 1247
 Gengenbach, B.G. 412, 1215
 GENTA. 259, 1111
 Gerasimowicz, W.V. 809
 Gerhold, L.S. 673
 Gericke, W.F. 161, 791, 1745
 Gerling, D. 1388
 Ghosheh, N.S. 438, 666, 637, 432, 515
 Gichner, T. 1328, 1379
 Giddings, T.H. 502
 Giese, H. 183, 977
 Gilbert, D.E. 1587
 Gilbert, M.D. 466
 Gildow, F.E. 1240
 Giles, B. 242, 1080
 Giles, K.I. 533
 Gill, J.R. 1501
 Gillespie, G.R. 1518, 1272, 1612
 Gilmore, E.C. 195, 855
 Giridhar, G. 652
 Gissel-Nielsen, G. 475, 613, 526, 1710
 Glass, A.D.M. 789, 676, 1879, 452, 530, 693, 772, 533
 Gold, Roger E. 1420
 Golus, H.M. 97, 1732
 Good, A.G. 278, 1267
 Good, Newell E. 1905-. 1386, 1781, 1803
 Gooden, D.T. 178
 Goodman, H.M. 206, 522
 Goodwin, W. H. 1880-. 1400
 Goos, R.J. 459, 1061
 Gordon-Kamm, W.J. 661
 Gordon, T.R. 997
 Goudey, J.S. 638
 Goulart, L.R. 193, 1239
 Grace, Oliver J. 1880-. 70, 1754
 Gracen, Vernon Edward, 443, 714
 Grafius, J. E. 135
 Grafius, J. E. & A new spring barley for feed. 50
 Grafius, J.E. 377, 918
 Grafton, K.F. 1244
 Graham, R.D. 465, 1263
 Graham, W.D. 178
 Graham, W.D. Jr. 21, 374
 Granett, A.L. 1294, 1865
 Grattan, S.R. 473, 1733, 1475
 Grau, C. R. 844
 Grau, C.R. 833, 1206
 Grbavac, N. 1086, 1087, 1043, 1032
 Greaves, M.P. 1621, 1690
 Greenhalgh, R. 987, 1373
 Gregory, E.J. 133, 486, 695, 3, 76, 229
 Gregory, L.V. 1190
 Griesbach, J.A. 1238
 Griffith, M. 610, 494, 787
 GRLEA. 934, 1233
 Grogan, R.G. 572, 1070
 Groth, D.E. 333, 1152, 1228

AUTHOR INDEX

- Groth, J.V. 231, 1068
 Grument, R. 254
 Grumet, R. 108, 267, 626, 194, 506, 1282, 262, 620
 Grunes, D.L. 465, 1263, 466
 Grunwald, C. 496
 Guillard, K. 96, 1494, 116, 481
 Gulick, S.H. 473, 1733
 Gumpert, F.M. 302, 1135, 1874
 Gupta, U.C. 456, 1717, 482, 1318
 Gurba, J.B. 840, 1805
 Haagsma, T. 1552, 1553
 Haahr, V. 288, 1128
 Habetz, R.J. 333, 1152, 1228
 Haby, V.A. 462, 1699, 1720, 86, 1724
 Hachisuka, M. 488, 1270
 Haderlie, L.C. 306, 1322, 1649
 Haeder, H.E. 460
 Hagan, A.K. 47
 Haggblom, P. 342, 1839
 Hagler, W.M. Jr. 1385, 1856
 Hald, B. 425, 1806, 1854
 Hall, C. 1524, 1562, 1681, 1485
 Hall, D.H. 997
 Hall, J.K. 1275, 1615, 1749
 Hall, L. 17, 360, 363
 Hallmark, W.B. 333, 1152, 1228
 Halstead, R.P. 179
 Hamilton, K.C. 1274, 1614
 Hamilton, R.H. 1618
 Hamm, R.T. 1045
 Hammerton, R.W. 636
 Hammond, C. 156, 1801
 Hammond, James J. 1298
 Hammond, R.B. 893, 1755, 874
 Hanchey, P. 1116
 Hanes, J.A. 769
 Hanlon, E.A. Jr. 157, 491, 1743
 Hanscom, J.T. 144, 331, 145, 332, 143, 330
 Hansen, A.E. 344, 824, 1851
 Hansen, L.D. 467, 581
 Hanson, A.D. 108, 267, 626, 279, 644, 194, 506, 1282, 418, 814, 642, 254, 798, 262, 620, 624, 623, 502, 405, 794, 507, 265, 625, 744, 508
 Hanson, J.B. 476, 616
 HARAA. 848
 Hardie, A.R. 1347, 1666
 Harein, P.K. 1406, 1414, 1411, 1390
 Harman, J.M.P. 1044
 Harper, J.F. 310, 682
 Harper, J.N. 136
 Harrabi, M.M. 366
 Harris, K.L. 1384
 Harrison, M.A. 764, 598, 634
 Harrison, S.A. 333, 1152, 1228
 Hart, Helen, 1900-. 1143
 Hart, L. P. & MSU Ag Facts. 135
 Hart, L.P. 830, 828, 829, 1119
 Hart, R.H. 175
 Hartwig, N.L. 1275, 1615, 1749
 Harvey, R.G. 1628, 1835, 1340, 1659, 1870, 1347, 1666
 Hascoet, M. 1622, 1834
 Hash, C.T. Jr. 268, 1115
 Hatano, T. 820
 Hatfield, J.L. 82
 Haug, A. 498, 1276, 1333, 630
 Haug, A.R. 1197, 1198
 Havercamp, H.G. 124, 678, 1737
 Hayashi, J. 187, 979
 Hayes, H. K. 1884-. 403, 1167
 Haynes, D.L. 241, 873
 Heaney, S.P. 1155
 Heath, M.C. 1364, 1683
 Heath, R. 413, 1259
 Heaton, L.A. 1258
 Heber, U. 674, 509, 1780, 1863
 Hegnauer, Leonard, 1872-. 121, 1139
 Heid, Walter George, 1932-. 990
 Heikes, D.L. 1661, 1830, 1855
 Heileman, W. H. 41, 1563
 Heitefuss, R. 1209
 Hejgaard, J. 204, 999
 Heller, K. 432, 515
 Helm, J.H. 367, 20, 372
 Helms Jorgensen, J. 190, 982
 Helms, H.B. 490, 749, 1741
 Hendricks, L.H. 1396
 Henson, M.A. 1529, 1756
 Hern, J.L. 463, 1722
 Herrman, T.J. 1454
 Hess, F.D. 651, 1519, 1313, 1642
 Heta, H. 187, 979, 414, 1216
 Heun, M. 294, 1134
 Heupt, W. 1506, 1636
 Hewings, A.D. 1241
 Highland, H.A. 1426, 1841
 Hilbert, D.W. 603
 Hill, B.D. 1484
 Hill, J.E. 311, 684, 1533, 258, 1310, 1640
 Hill, R.D. 1316, 184, 499
 Hipps, L.E. 5, 439, 1877
 Hirano, S.S. 1315
 Hislop, E.C. 1044
 Hitz, W.D. 507, 744
 Hiura, U. 203, 994
 HJHSA. 38, 1542, 615
 Ho, T.H.D. 739, 436, 629, 636
 Hockett, E.A. 273, 1120, 320, 442, 690, 274, 11, 350, 112, 658, 226, 564, 238, 218, 1242
 Hodges, C.F. 1114, 1513
 Hodges, T.K. 773
 Hoffman, A.H. 122
 Hoffman, N.E. 279, 644
 Hollander, H. 1354, 1670
 Hollomon, D.W. 1121, 253, 1225
 Holloway, P. 413, 1259
 Holmsen, J.D. 1313, 1642
 Holton, C. S. 1154
 Holton, Charles Stewart, 1904-. 1153
 Holtz, Henry F., 1880-. 560, 1697
 Homeyer, U. 807
 Hook, J.B. 1654, 1861
 Hooley, R. 272, 633
 Hopkins, W.G. 732
 Hormann, W.D. 1644, 1838, 1843
 Hosoda, E. 778, 777
 Housley, T.L. 700, 426, 699
 Hoveland, C. 54, 1852
 Howard, S.W. 1540
 Hoxie, R.P. 861, 152, 916
 Hoyt, P.B. 67, 1714
 Hsiao, A.I. 769
 Huang, C.X. 296, 669, 667
 Huchzermeyer, B. 807
 Hudson, R.D. 156, 1801
 Huffaker, R.C. 467, 581, 519, 597, 558, 649, 713
 Huffman, D.C. 27, 174, 1800
 Humburg, N.E. 1556, 1602, 176, 1535
 Hume, Albert N. 1878-. 974, 121, 1139
 Humphrey, Harry Baker, 1873-. 992
 Huner, N.P.A. 543, 544, 610, 553, 787
 Hunt, Thomas F. 1862-. 98, 878, 1505, 1844
 Hunter, B.G. 1258
 Hurkman, W.J. 208, 527, 587, 586

AUTHOR INDEX

- Huston, C.H. 1510, 1596
Hutchinson, R.L. 333, 1152, 1228
Hutchison, J.M. 1528
Huzel, V. 464, 1723
Hwang, B.K. 1209
Hyslop, J. A. 939
Hyslop, J. A. 1884-. 938
Ikeda, K. 573
Inayatullah, C. 851, 906, 1872, 266, 883
Ingemansen, J. 1409
Ingemansen, J.A. 1398
Inglis, D. comp. 832
Ingversen, J. 341, 735, 681, 1824
Irvine, R.B. 23, 379, 14, 356, 1173
ISJRA6. 412, 1215
Isleib, T.G. 254
Israel, H.W. 988
Iswari, S. 1336
Jackson, A.O. 1258
Jackson, L.F. 1238, 997, 250, 1108
Jackson, P.C. 496, 809, 549, 594, 580
Jackson, T.L. 90, 1729
Jackson, W.B. 846
Jacobs, J.M. 737
Jacobs, M. 401
Jacobs, N.J. 737
Jacobsen, J.V. 418, 814
Jacobsen, N. 704, 822
JAFCA. 1688, 1, 1623, 1391
JAFCAU. 1045, 1638, 1103, 1837, 1202, 1840, 659, 1522, 589, 1687, 1339, 1309, 1639, 1380
Jain, R. 441, 683, 1532
James, D.R. 544
Jana, S. 431, 512, 1457, 1566
JANCA2. 1661, 1830, 1855
Janda, M. 1251
JANSA. 227, 1799, 1859
Jay, E. 31, 1401, 1818
JBCHA3. 711
JCECD. 761, 1349, 429, 1351, 1763
JCLBA3. 496
Jedel, P.E. 186, 978
JEENAI. 1406, 1414, 1397, 216, 866, 851, 865, 173, 941, 1396, 1395, 1389, 1396, 1405, 1425
Jefferies, D.J. 1392, 1822
Jende-Strid, B. 210, 531, 341, 735
Jenkins, S.F. 572, 1070
Jenner, C.F. 596
Jensen, C.A. 485, 1702
Jensen, E.H. 131, 323, 691
Jensen, G. 1423, 933
Jensen, H.P. 338, 1159, 190, 982, 183, 977
Jensen, J. 211, 212, 190, 982
Jensen, N.F. 382
Jensen, Ray Eugene. 742
JEVQAA. 518, 1694, 1709
JFDAZ. 1426, 1841
Jilani, G. 1410
JIVPA. 1416, 1403
JKESA. 906, 1872, 1393
JMBBBO. 671
Jodari, F. 172, 1774
Johnson, B.E. 459, 1061
Johnson, D.A. 1093
Johnson, E.C. 305, 1137
Johnson, H.G. 984
Johnson, J.T. 156, 1801
Johnson, J.W. 22, 378, 1256, 361, 114
Johnson, P.O. 1460
Johnson, W. 848
Johnston, R. H. 990
Johnston, R.H. 1189, 1067, 1089, 1041, 1066, 1071, 1017, 1007, 1088, 1006
Johnston, R.P. 1207
Joia, B.S. 1389
Jokinen, D.P. 241, 873
Jolley, V.D. 492, 480, 1268
JONEB. 513, 944, 942
Jones, A.J. 61, 1286
Jones, C. Allan. 60, 539
Jones, G.D. 219, 552, 1713
Jones, J.P. 1188, 1193, 1037, 1065, 1038
Jones, L.A. 659, 1522
Jordon, William R. 578
Jorgensen, J.H. 213, 841, 954, 293, 255, 1109, 338, 1159, 183, 977
JOSHB. 39, 1762, 1815
JOVIAM. 423
JPFCD2. 1355, 1678, 1704, 975, 1832, 1862
JPGRDI. 686, 537, 677, 1631, 673, 598
JPNUD. 647, 456, 1717
JPNUDS. 492, 488, 1270, 1323, 1350, 480, 1268, 475, 613, 1283, 1708, 409, 800, 1287, 1695, 1332, 464, 1723
JPRAN. 80, 1721
JRMGA. 797, 1559, 771, 1551, 175, 757
JSWCA3. 158, 1766, 1779, 148, 727, 1739, 59, 1712, 1752
JTEHD. 1635
Jung, F. 1045
Jung, J. 725
Jusaitis, M. 532, 648
JWMAA9. 842
KAEBB. 136
Kahler, A.L. 250, 1108
Kahlon, P.S. 1352, 1668
Kallsen, C.E. 133, 486, 695
Kamble, Shripat T. 1420
Kantack, B.H. 173, 941, 897
Karow, R.S. 1170
Karren, J.B. 860
Kasha, K.J. 235
Katayama, M. 820
Kaufman, P.B. 438, 665, 529, 497, 637, 764, 598, 432, 515, 634
Kaufmann, M.L. 359, 352
Kaur-Sawhney, R. 1285
Kawai, S. 488, 1270
Kee, D.D. 26, 79
Keener, T.K. 1463, 28, 1502
Keith, B. 703
Kellerman, K.F. 484, 1701
Kenefick, D.G. 295, 668
Kenney, J. F. 910
Kernkamp, H. C. H. 1208, 1802
Kernkamp, M.F. 1084
Kerns, D. 876, 1783, 1797
Kerns, D.L. 216, 866
Kessler, R.W. 673
Key, J.M. 395, 770, 825
Khan, T.N. 77, 1062, 1719
Kibite, S. 359, 352
Kidder, D.W. 1540, 1495
Kieckhefer, R.W. 865, 173, 941
Kiehn, F.A. 1468
Kilpatrick, R.A. 315, 1144, 248, 1107
Kim, H.K. 1234, 1370
Kim, J.C. 1302, 1633
King, R.W. 724, 1334
Kingsland, G.C. 1047, 21, 374
Kirby, E.J.M. 723
Kirkland, K.J. 30, 417, 1568, 1462
Kishi-Nishizawa, N. 488, 1270
Klein, R.R. 639
Kleinhofs, A. 206, 522, 795, 694, 1269, 1262
Kleudgen, H.K. 514, 1458

AUTHOR INDEX

- Klimovitz, R.J. 1422
 Klobus, G. 519
 Klostermeyer, E.C. 882
 Knight, T.J. 702, 1227, 592, 1222
 Koch, E.J. 219, 552, 1713
 Koehler, S. 739
 Koepsell, P.A. 1170
 Koga, H. 203, 994
 Kohler, F. 276, 1123, 1252
 Kolb, F.L. 15, 357, 1344, 91, 239, 1730, 277, 640
 Kolding, M.F. 343, 969
 Kolp, B. J. 1928-. 132, 692
 Kolster, P. 222, 1048, 318, 1145
 Konishi, T. 345, 1168
 Korte, F. 660, 1868, 1647
 Kral, C.W. 1529, 1756
 Krall, J.M. 1447, 1448, 1525, 1582, 1499, 52, 1449, 1445, 1500, 1446, 1450
 Kramer, K.J. 1393, 1396
 Kreckhefer, R. 897
 Kremer, D.F. 752
 Kristiansen, K.N. 341, 735
 Krizek, D.T. 164, 1706, 1323, 1350
 Krogh, P. 425, 1806, 1854
 Krol, M. 543
 Kruger, James E. 847, 1819
 Krupa, Z. 543, 544
 Kudagamage, C. 299, 905, 1253
 Kudrna, D.A. 348, 750
 Kuhna, A.S. 304, 675
 Kuiper, H.J. 270, 1117
 Kukula, S.T. 168, 1583, 1772
 Kumari, S.M. 1416
 Kunert, K.J. 1687
 Kunoh, H. 988
 Kuo, T.M. 795
 Kusano, T. 573
 Kvien, J.S. 166, 1573, 1770
 Kyle, D.J. 610
 Kyser, G.B. 105, 1311, 1511
 LaBerge, Donald E. 847, 1819
 LaCroix, J.D. 432, 515
 Ladyman, J.A.R. 262, 620, 405, 794, 507
 Lafever, H.N. 371
 Lagarias, J.C. 819, 711
 Lai, F.S. 1376, 1812
 Lampe, D.J. 850, 1236
 Lampert, E.P. 241, 873
 Landolt, P.J. 1407, 1798
 Landry, B. 251, 1249
 Langbehn, G. 1508
 Lange, O.L. 674
 Langer, D.K. 1628, 1835, 1347, 1666
 Langston-Unkefer, P.J. 702, 1227, 592, 1222
 Lanini, W.T. 1475
 Lanterman, W.S. 1486
 Lappin, H.M. 1621, 1690
 Laroche, A. 732
 Larsen, J. 341, 735
 Larter, E.N. 184, 499
 Lauchli, A. 483, 1324
 Lauer, J. 1443, 1575, 1442, 170, 1605, 1606, 1444
 Lauer, J.G. 202, 511, 712
 Laughlin, W.M. 33, 1715, 450, 504
 Lawton, J.H. 872
 LAXBA. 27, 174, 1800
 Lazowski, E.J. 1275, 1615, 1749
 Leach, L. D. 1900-. 120, 1138
 Leary, W.P. 75
 Lee, C.C. 420
 Lee, Chung, . 894
 Lee, E.H. 1287, 1695
 Lee, G.A. 1590
 Lee, R.D. 156, 1801
 Lee, S.A. 1325, 1650
 Lee, T.T. 1631
 Leep, R. H. 50
 Leep, R.H. 48
 Legge, R.L. 432, 515
 Leggett, J.E. 516
 Lehman, Samuel George, 1887-. 1036
 Leigh, R.A. 576, 555, 1261
 Leitch, R.H. 67, 1714
 Leith, B.D. 400, 783
 Leland, T. 623
 Leland, T.J. 642, 624
 Leonard, K.J. 228, 1059
 Leonard, M.E. 853
 Lerner, H.R. 446
 Leslie, J. 99, 246, 959
 Lesoing, G. 59, 1712, 1752
 Lester, G.E. 614
 Letham, D.S. 535
 Leukel, R. W. 952, 993, 1106, 1192
 Leukel, R. W. 1888-. 153, 1810, 1105, 1091, 1191
 Levesque, M.P. 774, 1705, 37, 94, 471
 Levesue, M. 78, 461
 Levine, M. N. 1099
 Levine, Mose N. 1100
 Levitt, G. 1610, 1689, 1809
 Lewis, C.E. 57, 1751, 1776
 Li, P.H. 615
 Lichtenstein, E.P. 1619
 Lindblad, Carl J. 1382, 1826
 Lindner, W. 1645, 1828
 Lindow, S.E. 1221, 1234, 1370
 Lindstrom, O.M. 553
 Ling, L.C. 697, 909
 Linscott, D.L. 1362, 1682
 Lish, J.M. 1600, 1599, 1601, 1293, 1630, 1527, 1597, 1544, 1814, 1456, 1598, 1454
 Lister, R.M. 850, 1236
 Litek, K. 807
 Little, J. 848
 Litzenberger, Samuel C. 1914-. 1131
 Livingston, D.P. III. 404, 793
 Livingston, S. 150, 835, 1740
 Lockley, P. 780
 Loewen, M. 1567
 Lohde, J. 318, 1145
 Lohr, K. 1423
 Lomax, T.L. 662, 775
 Lorens, G. 1238
 Lorenz, K. 767, 1816
 Loschiavo, S.R. 1397, 1389, 1425
 Love, H.H. 160, 402, 1878
 Lovelace, D.A. 1593
 Loynachan, T.E. 89, 1728
 Lu, C.R. 438, 666
 Ludeke, K.L. 10, 349
 Luk, T.M. 700
 Luke, H.H. 12, 353, 1171, 100
 Lukezic, F.L. 1069
 Lund, R.E. 462, 1699, 1720
 Luz, W.C. 1194
 Luz, W.C. da. 996, 1081, 998, 1064, 1210
 Lynch, D.V. 802, 730, 729
 Lynd, J.Q. 157, 491, 1743, 479, 1693
 Maas, E.V. 767, 1816
 MacGregor, John M. 84
 Mackie, W.W. 122
 Macko, V. 1205, 416, 1219
 MacNish, G. 1187, 1761

AUTHOR INDEX

- Maggioni, A. 657
 Maissan, E. 543, 544
 Makino, T. 232
 Malhi, S.S. 67, 1714
 Mallory, C.A. 1600, 1599, 1601
 Manbeck, H.B. 1377, 1873
 Manisterski, J. 389, 1181
 Mansager, E.R. 1281
 Marble, V.L. 1475
 Mares, D.J. 734, 1338, 1850
 Marks, G.C. 747
 Marschner, H. 472, 602
 Marshall, D. 385, 1230, 1345
 Marshall, G.H. 406, 796, 1359
 Marshall, H.G. 15, 357, 1344, 91, 239, 1730, 277, 640, 286
 Marten, G.C. 49, 505, 1750
 Martens, J.W. 186, 978
 Martin, C.R. 1376, 1812
 Martin, D.A. 46, 1280, 1617
 Martin, R.A. 524
 Martinoia, E. 674
 Marumo, S. 820
 Mashhadi, H.R. 1357, 1679, 1767, 1346, 1665, 1760
 Mask, P.L. 47
 Mason, C.T. Jr. 447, 932
 Mason, L. 172, 1774
 Mather, D.E. 128, 317, 688
 Mathre, D.E. 87, 1072, 1725, 1067, 1089, 1041, 1066, 1071, 1017, 1007, 1088, 1006
 Mathre, Don E., 1938-. 990, 831
 Mathur, S.P. 774, 1705, 78, 461, 37, 94, 471
 Matin, M.A. 663, 1871
 Matthew, D.L. Jr. 856
 Matthew, David L. Jr. 857
 Maxwell, D. 886
 Mayama, S. 203, 994
 Mayfield, A.H. 1040
 Mazuranich, P.C. 903
 Mazza, G. 1372, 1833
 Mc Gowan, Anthony Alexander, . 627
 McBeath, J.H. 142, 329, 144, 331, 145, 332, 143, 330
 McClure, J.W. 546, 1624
 McCracken, D. 58, 1864
 McCully, W.G. Wiese, A.F. 1471
 McDermott, J.M. 259, 1111, 1000
 McDonald, B.A. 259, 1111, 1000, 393, 1186
 McDonald, D. 32, 1831, 1857
 McDonald, S. 902
 McDowell, M. 806, 1684
 McFarlane, C. 806, 1684
 McGaughey, W.H. 1395
 McGovern, T.P. 1393
 McGraw, R.L. 313, 147, 335, 718, 260, 618
 McGregor, W.R. 1363, 1564, 1748
 McIntyre, G.I. 769
 McIver, R.N. 68, 1691, 1696
 McKercher, R.B. 1364, 1583, 1363, 1564, 1748
 McKinney, Harold Hall, 1889-. 126, 1254
 McLean, D.L. 877, 1246
 McLeod, A.R. 583
 McMaster, S.A. 1507
 McNew, R. 876, 1783, 1797
 McNew, R.W. 216, 866, 479, 1693
 McPherson, R.M. 928, 927, 929, 926, 925
 McPherson, Robert M. 922, 1673, 1791, 1676, 1794, 923, 1674, 1792, 921, 1672, 1790, 920, 1671, 1789, 924, 1675, 1793
 McProud, Wayne Lucas, . 617
 Melchers, L. E. 1887-. 1200, 1201
 Meller, E. 792
 Melvin, G. 1577
 Menke, J.W. 757
 Meredith, M.P. 466
 Merkle, O.G. 284, 895
 Mersie, W. 1291, 1476, 1627
 Merwin, I.A. 39, 1762, 1815
 Mestres, G. 1620, 1827
 Mestres, R. 1620, 1827
 Metcalfe, D.R. 186, 978
 Metz, S.G. 1189, 1017, 1007, 1088
 Metzger, D.D. 103, 264, 622
 Metzger, J.D. 763, 762
 Meyer, D.W. 227, 1799, 1859
 Meyer, H.H.D. 1045
 Meyer, J.A. 1658, 1829
 Michaelson, G.J. 89, 1728
 Michel, L.J. 376, 25, 381, 1177, 24, 380, 1176, 368, 243, 1223
 Michels, G.J. Jr. 875
 Mielke, L.N. 148, 727, 1739
 Mikolajczak, K.L. 1391
 Miles, M.R. 282, 1125, 1163
 Miller, A.N. 745
 Miller, C.O. 536
 Miller, G.W. 435, 1264, 474, 612
 Miller, S.C. 1582
 Miller, S.D. 46, 1280, 1617, 1447, 1448, 1525, 1443, 1575, 1499, 52, 1449, 1445, 1442, 170, 1605, 1501, 1497, 1496, 1606, 1455, 1444, 1500, 1446, 1450, 1554, 816, 1595, 1591, 445, 758, 312, 685, 1534, 1473
 Miller, T.D. 150, 835, 1740
 Miller, T.L. 1515
 Mills, D. 773
 Minvielle, F. 251, 1249
 Mishra, S.D. 677
 Mitchell, G.A. 89, 1728
 Mitich, L. 1577
 Mitich, L.W. 105, 1311, 1511
 Miyahara, T. 589
 Mogensen, H.L. 444, 743
 Mogensen, V.O. 1290
 Mohanty, P. 818
 Monsen, S.B. 771, 1551
 Moomaw, R.S. 1537, 1757, 109, 283, 1517, 127, 1652
 Moons, C. 1658, 1829
 Moore, D. 848
 Moore, M. B. 1905-. 403
 Moore, M.B. 1084
 Moore, R. A. 1861-. 140, 968
 Moore, S.H. 333, 1152, 1228
 Morden, G. 464, 1723
 Morey, D.D. 361, 114
 Morgan, G.H. 284, 895, 13, 354, 917
 Mori, S. 488, 1270
 Morishita, D.W. 815, 1592, 607, 1504
 Moriya, I. 384, 751
 Morrill, W.L. 1423, 1427, 933
 Morrison, I.N. 590, 1489, 1479, 1288, 1626
 Morrison, K.J. 42, 185, 951
 Morrow, G. E. 1840-. 98, 98, 98, 878, 878, 878, 1505, 1505, 1505, 1844, 1844, 1844
 Morrow, L.A. 1474
 Morrow, L.S. 1480
 Morton, B.C. Jr. 21, 374
 Moseman, J. G. 1921-. 601, 957
 Moseman, J.G. 389, 1181, 388, 1180, 315, 1144, 248, 1107
 Moss, S.R. 1531, 1875
 MUCBA. 830, 828, 829, 48, 914, 915
 Mueller, R.T. 457, 1297
 Muir, C.E. 42, 185, 951

AUTHOR INDEX

- Muir, R.M. 673
 Mulder, C.E.G. 312, 685, 1534
 Mullenax, Richard H. 65, 556
 Mullet, J.E. 639, 337, 731
 Mullins, G.L. 518, 1694, 1709
 Multon, J. L. 1938-. 834, 1381, 1825
 Mulvehill, John F. 84
 Mumma, R.O. 1618
 Mundt, C.C. 228, 1059, 1046
 Mundt, G.A. 1590
 Mundy, B.P. 1203
 Munk, L. 222, 1048, 318, 1145
 Munro, D. 1316
 Murata, M. 205
 Murdock, E.C. 1570
 Murphy, C.F. 245, 609, 286
 Murphy, H. C. 1902-. 966, 1003
 Murphy, J.P. 25, 381, 1177, 24, 380, 1176
 Murray, T.D. 989
 Murry, L.E. 433, 521
 Murthy, N.B.K. 1355, 1678, 1704
 Musker, D. 1342, 1662
 Muzzelo, L.M. 1377, 1873
 MXMRA. 92, 468
 MYCOA. 778
 MYCOAE. 1078
 Nalewaja, J.D. 1518, 1292, 1477, 1629, 1272, 1612, 1273, 1613, 1554, 816, 1595, 1591, 445, 758, 312, 685, 1534, 1296, 1478, 1625, 1433, 1473
 Nardi, S. 1279
 Nash, R.G. 1335, 1656
 Nash, R.L. 72, 563
 NASSD. 775, 576, 555, 1261, 1094, 498, 1276, 269, 946, 223, 945, 327, 1148, 307, 1141, 1876
 Nassuth, A. 344, 824, 1851
 Naylor, J.M. 1566
 NDFRA. 207, 1001
 Neelgund, Y.F. 1416
 Negrutiu, I. 401
 Nelson, J.E. 1547
 Nelson, L.R. 1509
 Nelson, D.E. 177, 1235
 Nelson, Stuart Owen, 1927-. 1408, 1820
 NEPHA. 667, 1075, 583
 Netsvetaev, V.P. 271, 632
 Nevo, E. 388, 1180
 Newton, P.B. 1403
 Nguyen, H.T. 266, 883
 Niks, R.E. 270, 1117
 Nilan, R.A. 341, 735, 42, 185, 951, 1329, 1651, 1869
 Nofziger, D.L. 600, 1306, 1634, 1465
 Noll, C.F. 34, 584
 Nolt, C. 806, 1684
 Noma, M. 432, 515
 Nomoto, K. 715
 Northcutt, G. 1548
 Norvell, W.A. 465, 1263, 466
 NOSCA. 428
 NUSRA. 118, 300, 963, 119, 301, 964
 Nutter, F.W. Jr. 1060, 346, 1169, 1257
 Nyborg, M. 67, 1714
 O'Brien, R.D. 1221
 O'Donovan J.T. 1278, 1432
 O'Donovan, J.T. 30, 417, 1568
 O'Neal, W. B. 1578
 O'Sullivan, P.A. 1462, 1514, 1278, 1432
 Oaks, A. 795
 Odell, G.V. Jr. 157, 491, 1743
 Oehlschlager, A.C. 1418
 Oelke, E. A. 162
 Ogborn, J.E.A. 1558
 Oh, J.Y. 1262
 Ohlrogge, J.B. 805
 Ohm, H.W. 193, 1239, 347, 1255
 Oji, Y. 566
 Okamoto, S. 566
 Oku, M. 573
 Olien, C.R. 404, 793, 614, 495, 406, 796, 1359, 1343, 500, 1277, 1224, 1308, 304, 675
 Olson, P.A. 1281
 Olson, R.A. 123, 1848
 Olson, W. 1296, 1478
 Olson, W.A. 1625, 1433
 Onsager, J.A. 903
 Oplinger, E.S. 75
 Orioli, G.A. 790
 Orloff, S.B. 1469, 1475, 1586, 1466, 1608
 Orr, G.R. 710, 1331
 Orr, J. 1475
 Orser, C. 1234, 1370
 Osborn, Herbert, 1856-. 115, 904
 Osborn, T.C. 631
 Osborne, Thomas B. 1859-. 733, 1158
 Ottman, M.J. 534, 1711, 10, 349
 Outtrup, H. 341, 735
 Owsley, W.F. 156, 1801
 Page, A.L. 1327
 Paleg, L.G. 532, 648
 Palm, H.L. 1561
 Palni, L.M.S. 535
 Palta, J.P. 1336
 Pammel, L. H. 1862-. 41, 1563
 Park, J. 686
 Parker, C.E. 659, 1522
 Parker, J. R. 892, 1787
 Parker, J. R. 1884-. 890, 1785, 881, 1784, 891, 1786
 Parker, J.H. 340, 1162
 Parker, John H. 1891-. 1002, 1113
 Parker, K.A. 914, 915
 Parker, R. 1540
 Parker, William B. 1885-. 1421
 Parlevliet, J.E. 1092, 307, 1141, 1876
 Pass, H. 287, 896
 Patel, J.D. 128, 317, 688
 Patterson, G.W. 745
 Payne, T.S. 313, 147, 335, 718
 PCBPB. 1342, 1662, 1314, 1643, 755, 1365, 1686
 Peahler, P.L. 101, 252
 Pearson, C.H. 97, 1732
 Pederson, D.V. 1010
 Pederson, V.D. 1035, 1016, 1005, 1004, 1015, 1060, 346, 1169, 1257, 1029, 1014, 1023, 1013, 1022, 1012, 1039, 1011, 1009
 Peek, D.C. 1657, 1703, 1319, 1648
 Peeper, T.F. 600, 1306, 1634, 1465
 Pepper, J. H. 1902-. 1424
 Pernot, E. F. 1859-. 1160
 Perring, T.M. 853
 Perry, S.K. 1119
 Person-Dedryver, F. 223, 945
 Peteghem, C.H. van. 1202, 1840
 Peter, A.M. 136
 Peters, D.C. 216, 886, 876, 1783, 1797
 Peters, M.A. 33, 1715, 450, 504
 Petersen, R.R. 1187, 1761
 Peterson, D.M. 631, 137, 325, 701, 700, 426, 699, 776, 1360
 Peterson, V. 882
 Peterson, V.D. 1034
 Petrie, S.E. 90, 1729
 Pfahler, P.L. 12, 353, 1171, 221, 554, 100
 Pfanz, H. 674, 509, 1780, 1863
 Pfeffer, P.E. 809

AUTHOR INDEX

- Pfleeger, T. 806, 1684
 Pharis, R.P. 432, 515
 Phipps, P.M. 35, 1664
 PHYTA. 302, 1135, 1874, 1221, 1209, 1258, 280, 1124, 228, 1059, 1179, 203, 994, 1368, 270, 1117, 231, 1068
 PHYTAJ. 1166, 385, 1230, 1345, 321, 1146, 1077, 1488, 1245, 850, 1236, 1092, 411, 1213, 1112, 1241, 1090, 1046, 346, 1169, 410, 1212, 572, 1070
 PIACA. 299, 905, 1253, 747
 PIAIA. 243, 1223
 Pickering, R.A. 189, 981
 Piening, L.J. 470, 1265
 Pierce, A.M. 1418
 Pierce, H.D. Jr. 1418
 Pierce, R. 1142
 Pike, K. 886
 Pike, K.S. 882
 Pike, P. 849
 Pitts, J.T. 899
 Plagge, H. H. 1894-. 605, 1503
 PLDIDE. 285, 1127, 1240, 1130, 1096, 1234, 1370
 PLDRA. 93, 1074, 1189, 282, 1125, 247, 1247, 87, 1072, 1725, 1187, 1761, 35, 1664, 188, 980, 1104, 1637, 997, 1043
 Plenchette, C. 579, 1692
 Ploeg, H.L. 1610, 1689, 1809
 PLPHA. 487, 708, 807, 1316, 438, 666, 208, 527, 296, 669, 348, 750, 548, 278, 1267, 528, 1336, 665, 789, 467, 581, 337, 731, 746, 523, 587, 510, 985, 819, 631, 702, 1227, 687, 516, 519, 766, 810, 568, 679, 497, 739, 643, 1735, 745, 674, 662, 732, 638, 752, 768, 544, 672, 729, 592, 1222, 1325, 1650, 586, 566, 234, 577, 279, 644, 1285, 509, 1780, 1863, 710, 1331, 809, 281, 478, 646, 676, 1879, 654, 636, 472, 602, 597, 272, 633, 418, 814, 784, 642, 721, 773, 753, 652, 1315, 610, 719, 558, 779, 551, 763, 653, 561, 795, 582, 741, 624, 740, 670, 1321
 Plummer, Bernie E. 1356
 PMASA. 273, 1120
 PNASA. 1205, 1000, 310, 682, 802, 436, 629, 220, 1289, 206, 522, 661
 PNWSB. 1275, 1615, 1749, 1486, 1480
 Poehlman, J.M. 1244
 Poehlman, John Milton. 1182, 976
 Poljakoff-Mayber, A. 446
 Pomeranz, Y. 1371, 1821, 1845
 Pomeroy, M.K. 1320, 760
 Pope, A.J. 576
 Post, T.J. 129, 948
 Potter, Alden A. 992
 Powell, G.M. 66, 1753, 1811
 Powelson, R.L. 1187, 1761, 1170
 Powers, J.R. 1402, 1823
 PPGGD. 820, 75, 72, 563
 PPRBA. 168, 1583, 1772
 Prasad, G. 345, 1168
 Prather, T.S. 1492
 Prentice, N. 45, 503
 Price, S.C. 311, 684, 1533, 258, 1310, 1640
 Price, T.P. 755
 Punja, Z.K. 572, 1070
 Pushnik, J. 474, 612
 Pushnik, J.C. 435, 1264
 Puterka, G.J. 216, 866, 876, 1783, 1797, 195, 855
 Putnam, A.R. 419, 761, 1349, 1307, 429, 1351, 1763
 Puy, D. van der. 449, 1607
 PVPCB. 840, 1805
 Qualset, C.O. 1238, 877, 1246, 238
 Qureshi, F.A. 1437
 Rabb, J.L. 333, 1152, 1228
 Radford, J. D. & Management of crops and soils in NC and NE Minnesota correspondence course. 162
 Raghu, K. 1355, 1678, 1704
 Ragsdale, D.W. 934, 1233
 Rai, B. 567
 Rains, D.W. 558, 649
 Raison, J.K. 710, 1331
 Rajagopal, R. 571
 Rajam, M.V. 1285
 Rajashekar, C. 615
 Ramage, R.T. 369, 1175
 Ramagopal, S. 309, 1326, 768, 220, 1289
 Ramaiah, K.V. 196, 1441
 Ramos, J.M. 407, 799
 Randall, P.J. 591, 1266
 Rangaswamy, J.R. 1380
 Rao, K.P. 72, 563, 649
 Rapp, J.C. 337, 731
 Rappaport, L. 805
 Rardon, P. 29, 1526, 151, 1549, 1759
 Rasmusson, D.C. 282, 1125, 71, 225, 434, 103, 264, 622, 408, 1093
 Ratterman, D.M. 548, 1314, 1643, 1365, 1686
 Rattunde, H.F. 134, 324, 696
 Rawn, Carroll David. 813, 1220
 Ray, T.B. 1530
 Rayapati, P.U. 234, 577
 Raymer, P.L. 392
 Raynaud, S. 808, 1565
 Ream, H.W. 477, 1312
 Recalde, L. 407, 799
 Reed, George M. 1878-1956. 1217, 1218
 Rees, R.G. 1207
 Reeves, D.L. 138, 1538, 139, 326, 1539, 17, 360, 1398, 363
 Reginato, R.J. 82
 Reicosky, D.A. 623, 265, 625
 Reid, C.P.P. 810
 Reid, D.A. 219, 552, 1713
 Reid, David A. 1914-. 56, 525
 Reinbergs, E. 128, 317, 688, 191
 Reinhold, M. 321, 1146, 364, 397, 1204
 Reisenauer, P. 42, 185, 951
 Retan, A. 849
 Retan, A.H. 882
 Rho, Y.B. 637
 Rhodes, D. 744
 Rickard, W.H. 428
 Rickman, R. 886
 Riedell, W.E. 1332, 559
 Riesen, T.K. 1078
 Riesselman, J.H. 1189
 Riley, M.M. 1283, 1708
 Rines, H.W. 217, 542, 179, 370, 412, 1215
 Ringer, R.K. 1635
 Ripley, B.D. 1309, 1639
 Ritchie, G.S.P. 1271
 Roane, C.W. 383
 Robbelen, G. 294, 1134
 Robbins, G.S. 18, 362, 1229
 Roberts, P.A. 837, 1769
 Roberts, T.M. 583
 Robertson, L.D. 188, 980, 214
 Robertson, M. 1316
 Robertson, R.L. 858, 859
 Robinson, G.D. 1587
 Robinson, K. 773
 Robson, A.D. 1271
 Rodenhiser, H. A. 1899-. 1153
 Roelfs, A. P. 53, 991, 1374

AUTHOR INDEX

- Roelfs, A.P. 1179, 289, 1129, 231, 1068
 Rogers, R. N. 1927-. 493
 Rogers, R.L. 27, 174, 1800
 Rohner, E. 570
 Romheld, V. 472, 602
 Roos, E.E. 205
 Roselle, Robert E. 1420
 Ross, C.W. 603
 Ross, M.A. 653
 Roth, G.W. 91, 239, 1730
 Rothman, P.G. 1096, 370, 355, 1172
 Rothrock, C.S. 22, 378, 1256
 Rouchaud, J. 1658, 1829
 Rousseau, D.M. 1202, 1840
 Roylance, Howard B. 1611
 Rozhkovskii, A.D. 611
 RREVA. 1413
 Rubinstein, B. 759
 Ruckendorfer, H. 1645, 1828
 Ruppel, R.F. 914, 915
 Ruth, T.J. 789
 Rydrych, D.J. 1439
 Ryerson, D.K. 1541, 1758, 1813
 Ryoma, Y. 566
 Sabey, B.R. 1299, 1866
 Sacher, R. 423
 Saghai-Maroo, M.A. 1090
 Sahs, W.W. 59, 1712, 1752
 Saladini, J.L. 1529, 1756
 Salmon, D.F. 367
 Salmon, S.C. 1366
 Salto, C.E. 209, 864
 Sammis, T.W. 133, 486, 695
 Sanchirico, C.A. 466
 Sander, D.H. 123, 1848
 Sands, D.C. 1234, 1370
 Sandstrom, R.P. 528, 662, 775
 Satler, S.O. 645
 Satterthwait, A. F. 889
 Satterthwait, A. F. 1879-. 887, 888
 Sauer, D.B. 1140, 1378
 Sawyer, A.J. 241, 873
 Saxe, H. 571
 Saxton, A.M. 27, 174, 1800
 Schaad, N.W. 1195, 1231
 Scharen, A.L. 1156
 Scheffer, R.P. 1197, 1198
 Schepers, J.S. 148, 727, 1739
 Scheunert, I. 660, 1868, 1647
 Schmid, W.E. 559
 Schmid, W.E. 1332
 Schmidt, J.W. 16, 358, 386, 754, 291, 437, 650
 Schnitzer, M. 68, 1691, 1696
 Schober, A.E. 1507
 Schow, G.S. 1528
 Schroeder, G.L. 1625
 Schuchmann, R. 276, 1123, 1252
 Schultz, G. 807
 Schwartz, S.M. 466
 Schwartz, T.K. 72, 563
 Scott, N.A. 508
 SDFHA. 1409, 99, 246, 959
 Sebesta, D.K. 762
 Sechler, D.T. 247, 1247, 1244
 Sechler, Dale Truman, 1926-. 394, 765
 Seck, M. 1077, 1488
 Seely, C. I. 1611
 Segars, W.I. 156, 1801
 Sehgal, O.P. 1244
 Sen, D.N. 1523
 Senft, D. 2, 1555
 SENTD. 875
 Sethi, G.S. 180, 181, 182
 Setiawan, D.P. 934, 1233
 Severson, R.K. 92, 468, 469
 Shamel, A. D. 1878-. 973, 962
 Shands, R. G. 1903-. 159, 970
 Shaner, G. 1053
 Sharma, A.K. 30, 417, 1568
 Sharma, M.P. 1437
 Sharp, E.I. 297
 Sharp, E.L. 321, 1146, 366, 197, 838, 953, 373, 369, 1175, 364, 397, 1204
 Shaw, D.R. 600, 1306, 1634, 1465
 Shefelbine, P.A. 87, 1072, 1725
 Sheridan, J.E. 1086, 1087, 1043, 1032, 1083, 1026, 1025, 1031, 1030, 1024
 Sheridan, M.H. 1086, 1087
 Sheridan, J.E. 1082
 Sherwood, R.T. 1069
 Shilling, D.G. 659, 1522
 Shimabuku, R. 806, 1684
 Shinkle, J.R. 721, 641
 Shires, S.W. 872
 Shishiyama, J. 203, 994
 Shivashankar, Gurubasavappa, . 588
 Shukle, R.H. 850, 1236
 Sicher, R.C. 752, 237
 Siddiqi, M.Y. 789, 676, 1879, 452, 530, 772, 533
 Siddoway, F.H. 57, 1751, 1776
 Siegel, N. 1333
 Simeoni, L.A. 1299, 1866
 Siminovitch, D. 557, 804
 Simmons, S.R. 202, 511, 593, 712, 723, 408
 Simons, M. D. 967
 Simons, M. D. 1925-. 966, 1003
 Simons, M.D. 376, 1096, 25, 381, 1177, 24, 380, 1176, 19, 365, 1174, 188, 980, 368, 410, 1212, 243, 1223, 1211, 1049, 1063
 Simpson, C.L. 911
 Sims, J.R. 462, 1699, 1720
 Singh, Bhisham Pal, . 995
 Singh, D.B. 567
 Singletary, G.W. 624, 623
 Singleton, L.L. 1084
 Sinha, R.N. 1394
 Skogley, E.O. 462, 1699, 1720, 86, 1724
 Skou, J.P. 288, 1128
 Skrzypczak, G.A. 1272, 1612, 1273, 1613
 Slater, J.H. 1621, 1690
 Slater, S.E. 327, 1148
 Slates, R.V. 1, 1623
 Slegers, G.A. 1202, 1840
 Slife, F. W. 1923-. 448
 Slosser, J.E. 216, 866, 195, 855
 Smail, V.W. 112, 658, 226, 564
 Smedegaard-Petersen, V. 280, 1124, 69, 1050, 1051
 Smith, A.W. 893, 1755
 Smith, B.J. 1325, 1650
 Smith, C.M. 8, 1118
 Smith, C.R. Jr. 1391
 Smith, D. 477, 1312
 Smith, D. C. 1906-. 1042
 Smith, D.H. 152, 916
 Smith, D.H. Jr. 409, 800, 377, 918, 375, 940
 Smith, David H. 867
 Smith, G.R. 33, 1715, 450, 504
 Smith, I.K. 784
 Smith, L.B. 1405
 Smith, M.N. 1343, 1224, 1308
 Smith, W.O. Jr. 679
 Soberalske, R.M. 477, 1312
 Sobolik, F.J. 459, 1061
 Sogaard, B. 293

AUTHOR INDEX

- Solomon, M.A. 534, 1711
 Somers, D.A. 217, 542, 795
 Sommers, L.E. 518, 1694, 1709
 Somody, C.N. 1554, 816, 1595, 1591, 445, 758
 Song, I. 637
 Soon, Y.K. 111, 1736
 Soper, R. 464, 1723
 Soper, R.J. 608, 1700, 1734
 SOPPAA. 611
 Sorrells, M.E. 382, 303
 SOSCA. 37, 94, 471
 SOSCAK. 774, 1705, 453, 562, 1716
 Souza, E.U. 303
 Sowers, R.S. 40, 1764
 Sozinov, A.A. 271, 632
 Spanswick, R.M. 510, 985
 Speirs, W.E. 1393
 Spencer, M.S. 638, 654
 Spinney, R.L. 1361, 1680
 Spragg, F.A. 81, 230
 Srivastava, L.M. 703
 SSSJD4. 473, 1733, 465, 1263, 466, 1271
 St John, J.B. 594, 1487, 580
 St. Pierre, C.A. 251, 1249
 Stachowski, P.J. 1486
 Stack, R.W. 207, 1001, 459, 1061
 Stakman, E. C. 1885-1979. 403
 Stanton, T. R. 64, 956, 107, 961, 1098
 Stanton, T. R. 1885-. 171, 130, 322, 1226, 63, 955, 106, 960, 1097, 1100
 Stanton, T.R. 51, 200
 Stark, J.C. 306, 1322, 1649
 Starkman, E. C. 1100
 Starks, K.J. 209, 864, 870, 287, 896
 Starling, T.M. 219, 552, 1713, 383
 Statler, G.D. 1092
 Steffenson, B.J. 1179, 289, 1129
 Steinrucken, H.C. 1353, 1669
 Stephens, David E. 1874-. 1226
 Steponkus, P.L. 802, 730, 672, 729, 671, 661, 561, 670, 1321
 Sterling, J.D.E. 128, 317, 688
 Stewart, C.R. 234, 577, 753, 779, 582
 Stewart, R.K. 931
 Stewart, V. 29, 1526
 Stewart, V.R. 1463, 28, 1502, 238, 1536
 Stewart, W.M. 367, 20, 372
 Stiles, W.C. 39, 1762, 1815
 Stinner, B.R. 893, 1755
 Stobbe, E.H. 1484, 1438
 Stolen, O. 222, 1048, 318, 1145, 1051
 Stolzenberg, G.E. 1281
 Stone, A.M. 447, 932
 Storey, C.L. 1412
 Storey, R. 555, 1261
 Stout, R.G. 709, 1147
 Stout, W.L. 85, 871, 1482
 Strand, A. L. 1894-. 1424
 Strand, O.E. 1580
 Strand, Oliver E. 827, 1579, 1603
 Stritzke, J.F. 621
 Strobel, G.A. 1203
 Stroman, G.N. 399, 781
 Stuart, William, 1865-. 1102
 Stuthman, D.D. 313, 370, 147, 335, 718, 260, 618
 Su, H.C.F. 1410
 Suber, E.F. 156, 1801
 Subramanyam, B. 1406, 1414, 1411, 1413
 Suhayda, C.G. 630
 Sullivan, B.P. 221, 554, 101, 252
 Sullivan, M.J. 898
 Summons, R.E. 535
 Sundqvist, C. 421, 569
 Surowy, T.K. 310, 682
 Sussman, M.R. 310, 682
 Swan, D.G. 1540
 Swan, M. 464, 1723
 Sweetser, P.B. 1528
 Swensen, J.B. 1452
 Swingle, R.S. 236, 1726
 Switzer, C.M. 1562, 1681, 1485
 SWSPB. 1471
 SWSPBE. 1291, 1476, 1627
 Symons, S.J. 1567
 Szaniszló, P.J. 810
 TAAEA. 1377, 1873, 1434
 TAEB. 150, 835, 1740
 Taira, H. 589
 Tajani, M. 1077, 1488
 Takagi, S. 488, 1270, 715
 Takeda, K. 263, 821, 415, 812, 414, 1216
 Takemoto, T. 715
 Tanaka, C.K. 208, 527, 587, 586
 Tani, T. 1133
 Tao, D. 615
 Tao, G.Q. 535
 Tapia, L.S. 1543
 Tapke, V. F. 1890-. 1101, 1097
 Taylor, B.B. 534, 1711
 Taylor, H.F. 1688
 Taylor, D.C. 1294, 1865
 Teng, P.S. 965
 Terman, G. L. 1356
 Thai, K.M. 431, 512, 1457, 1566
 Thaxter, Roland, 1858-. 733, 1158
 Therrien, M.C. 23, 379, 14, 356, 1173
 Thill, D.C. 154, 1348, 1667, 1543, 1600, 1599, 1601, 1293, 1630, 1527, 1317, 1646, 1453, 1589, 1597, 815, 1592, 607, 1504, 1544, 1814, 1560, 1768, 1451, 1452, 1456, 1510, 1596, 1598, 1472, 1454, 1492
 Thimann, K.V. 686, 652, 645
 Thomas, P.L. 411, 1213
 Thompson, R.K. 236, 1726
 Thompson, S.S. 156, 1801
 Tilton, Elvin W. 1408, 1820
 Timian, R.G. 1243, 1257
 Timm, C.A. 459, 1061
 Timmons, F. L. 1905-. 1550
 Tindall, T.A. 97, 1732
 TISAA. 559
 Tittle, F.L. 654
 Tittle, F.T. 638
 Todd, B.G. 1484, 1438
 Tollefson, S. 836, 1746, 1817
 Tomar, J.S. 608, 1700, 1734
 Tomerlin, J.R. 389, 1181
 Topp, E. 660, 1868
 Torres, V. 604
 Townshend, J.L. 942
 Toyoda, H. 203, 994
 TQMB. 341, 735
 Traynor, P.L. 265, 625
 Treeful, L. 389, 1181
 Trevenen, S. 1018
 Tripathy, B.C. 818
 Trippet, B.J. 1546
 Tsai, C.Y. 74, 458, 1718
 Tsai, K.J. 61, 1286
 Tsuchiya, t. 205
 Tuck, B. 1498
 Tucker, T.C. 74, 458, 1718
 Tungland, L. 71, 225, 434
 Turley, R.H. 489, 722, 1738, 786, 1744
 Turner, D.L. 141, 1330, 155

AUTHOR INDEX

- Uemura, M. 802, 656
 Ukai, Y. 249, 1248
 Ullman, D.E. 877, 1246
 Ullrich, S.E. 341, 735, 42, 185, 951
 Upper, C.D. 1315
 Utomo, M. 455, 1698, 1777
 Vallejo, R.P. 1618
 Van de Velde, R. 413, 1259
 Van Duyn, J.W. 858, 859
 Van Gundy, S.D. 943
 Van Steveninck, R.F.M. 296, 669, 667
 Van Veldhuizen, R.M. 144, 331, 145, 332, 143, 330
 Vanden Born, W.H. 1524
 Vandenborn, W.H. 1437
 Vanderplank, J.E. 391, 1184
 Varanini, Z. 657
 Vargas, R.N. 1475
 Varner, J.E. 756
 Varvel, G.E. 92, 468, 469
 Vasil, I.K. 336, 726
 Veierskov, B. 686
 Veleminsky, J. 1328, 1379, 1329, 1651, 1869
 Veltrup, W. 647, 454, 565
 Vick, K.W. 1417
 Vieira, J.C. 1194
 Vierling, E. 741
 VIRLA. 1251
 Voetberg, G. 234, 577, 753, 779
 Volkmar, K.M. 513, 944
 Vore, R.E. 176, 1535
 Voskresenskaya, N.P. 611
 WAEBA. 1499, 52, 1449, 1445, 1442, 170, 1605
 Wagenet, R.J. 141, 1330, 155
 Wagner, G.J. 811
 Wahl, I. 389, 1181
 Wakiuchi, N. 566
 Walgenbach, D.D. 865
 Walker-Simmons, M. 1316, 348, 750
 Walker, Charles E. 1420
 Wall, D.A. 163, 1569, 1468
 Wallace, A. 457, 1297
 Wallen, Stanley E. 1420
 Walmsley, M.H. 603
 Walsh, D.A. 711
 Walstrom, R.J. 1398
 Walters, D.R. 1075
 Walton, W. R. 879
 Walton, W. R. 1873-. 890, 1785, 891, 1786, 880
 Ward, C.Y. 26, 79
 Ward, M.R. 467, 581, 519, 597
 Ward, S. 1509
 Warholc, D.T. 1486
 Warmund, M.R. 38, 1542
 Warner, K. 1426, 1841
 Warner, R.L. 487, 708, 348, 750, 723, 795, 694, 1269, 1262
 Warner, R.W. 1529, 1756
 Warnes, D. 282, 1125
 Watkins, J.E. 1104, 1637
 Watson, G.R. 1190
 Watson, M.C. 1274, 1614
 Watters, F.L. 1415
 Wax, L. M. 1520
 Weaver, W.A. 1417
 Webster, G.R.B. 1389
 Webster, J.A. 284, 895, 13, 354, 917, 851, 906, 1872, 266, 883, 377, 918, 152, 916, 375, 940
 Webster, James A. 867
 Webster, R.K. 259, 1111, 1000, 393, 1186, 1112, 1090, 997, 250, 1108
 Wedberg, J. L. 910, 844
 Wedberg, J.L. 833
 Wedin, W.F. 54, 1852
 Weems, J. B. 41, 1563
 WEESA. 1514, 1574, 1771, 258, 1310, 1640, 1479
 WEESAG. 441, 683, 1532, 1518, 1657, 1703, 431, 512, 1457, 1515, 30, 417, 1568, 595, 1491, 1292, 1477, 1629, 1272, 1612, 311, 684, 1533, 815, 1592, 607, 1504, 1302, 1633, 1464, 1307, 1483, 1567, 1468, 651, 1519, 1273, 1613, 306, 1322, 1649, 501, 1616, 808, 1565, 1566, 1537, 1757, 1465, 1313, 1642
 Weigel, R.C. Jr. 1610, 1689, 1809
 Weimberg, R. 446
 Weinstein, L.H. 1285
 Weis, C. 498, 1276
 Weissbart, J. 487, 708
 Welch, D.A. 477, 1312
 Welch, R.M. 465, 1263, 466
 Weller, D.M. 9, 1136, 1187, 1761
 Wellso, S.G. 241, 873, 862, 697, 909, 861
 Welterlen, M.S. 40, 1764
 Wenzel, G. 276, 1123, 1252
 Werker, E. 446
 Wesenberg, D.M. 18, 362, 1229, 237
 West, H.M. 1075
 WETEE9. 1367, 1588, 43, 1431, 154, 1348, 1667, 46, 1280, 1617, 1493, 1516, 1319, 1648, 163, 1569
 Wettlaufer, S.H. 1285
 Wettstein, D. von. 341, 735
 Wheatley, W.G. 235
 Whisler, F.D. 1434
 White, L.V. 1428
 White, N.D.G. 1397, 1425, 1430, 1685, 1842, 1394
 Whitesides, R.E. 595, 1491, 1474
 Whitman, C.E. 82
 Whitson, T.D. 1508, 1467, 1498
 Whittle, K. 912, 1788, 1804
 Whitwell, T. 1516
 Wickliff, C. 806, 1684
 Wicks, G.A. 167, 826, 1576, 1470
 Wiebe, G. A. 56, 525
 Wiebold, W.J. 80, 1721
 Wiersma, J. 282, 1125
 Wiersma, J.V. 1515, 71, 225, 434, 408
 Wiese, A.F. 1593
 Wilbur, D.A. 1429
 Wilcoxson, R.D. 1130, 370, 282, 1125, 192, 983, 389, 1181, 396, 1196, 1179, 289, 1129, 1163, 1084, 1093
 Wilding, S.B. 1287, 1695
 William, R.D. 1540
 Williams, G.M. 585, 1076
 Williams, J.P. 543, 544
 Wilsey, M.H. 736, 1778
 Wilson, B.J. 1531, 1875
 Wilson, R.F. 659, 1522
 Wilson, R.G. 988
 Wilson, R.L. 870
 Windle, J.J. 523
 Winkelhusener, T. 689, 1653
 Winter, K.A. 482, 1318
 WLSBA. 839
 Woding, F.J. 89, 1728
 Wojciak, J.H. 1396
 Wolf, F.T. 430, 785
 Wolfe, D. 48
 Wolfe, J. 672, 671, 670, 1321
 Wolfe, M.S. 1185, 327, 1148, 1149, 201, 986
 Wolfe, R.I. 23, 379, 14, 356, 1173
 Wolpert, T.J. 1205, 416, 1219
 Wong, Y.S. 711
 Woo, S.M. 1419

AUTHOR INDEX

Woodcock, T. 1085
 Wooding, F.J. 142, 329, 144, 331, 145, 332,
 143, 330
 Woods, Chas. D. 1856-. 913, 1157
 Worrall, D.W. 216, 866
 Worsham, A.D. 659, 1522
 Woznica, Z. 1292, 1477, 1629
 Wrage, L.J. 1460
 Wrage, Leon J. 1578
 Wright, R.C. 484, 1701
 Wright, R.J. 463, 1722
 Wrigley, C.W. 591, 1266
 Wrona, A.F. 510, 985
 WSWPA. 28, 1502, 1521, 105, 1311, 1511, 1507,
 1553, 1541, 1758, 1813, 1529, 1756, 1301, 1632,
 1867, 520, 1459
 Wu, F.C. 1243
 WUEXA. 989
 Wych, R.D. 723, 260, 618
 Yaegashi, H. 285, 1127
 Yahyaoui, A.H. 321, 1146
 Yamamoto, H. 1133
 Yamashita, Y. 1133
 Yap, Thoo Chai, . 547
 Yasuda, S. 345, 1168, 384, 751
 Yenne, S.P. 1472
 Yoshida, S. 656
 Young, A.J. 1342, 1662
 Young, F.L. 595, 1491
 Young, J.C. 1103, 1837
 Young, J.O. 882
 Young, K.J. 77, 1062, 1719
 Young, N.D. 1337, 719, 740
 Young, Nevin Dale. 728
 Youngs, V.L. 698, 1849, 1860, 137, 325, 701
 Yu, M.H. 474, 612
 Zahour, A. 292, 655
 Zamora, D.L. 1451
 Zaske, S.K. 11, 350, 218, 1242
 Zaylskie, R.G. 1281
 Zehner, J.M. 31, 1401, 1818
 Zehner, Marion Alice Griffiths, _1895-. 1218
 Zhang, Q.F. 1112
 Zhi-Yi, T. 686
 Zilkowski, B.W. 1391
 Zohary, D. 427, 738, 823, 388, 1180
 Zollinger, R.K. 1301, 1632, 1867
 Zuniga, G.E. 387, 919
 Zwar, J.A. 272, 633
 1904-. 1154
 1932-. 627
 1934-. 588
 1937-. 451, 517
 1938-. 1073
 1939-. 422, 575
 1940-. 547, 1358
 1941-. 995, 664, 1545
 1943-. 894
 1944-. 578, 635
 1945-. 443, 714, 617

Cereals

1, 2, 4, 6, 7, 8, 9, 28, 31, 32, 33, 34, 35, 36, 53, 54, 55, 60, 66, 68, 70, 72, 74, 75, 83, 104, 105, 110, 115, 117, 118, 119, 122, 123, 125, 142, 143, 144, 145, 146, 149, 150, 151, 153, 156, 161, 165, 168, 169, 178, 195, 196, 198, 201, 216, 231, 244, 298, 300, 301, 305, 308, 309, 314, 327, 329, 330, 331, 332, 333, 334, 336, 343, 387, 392, 395, 427, 440, 448, 458, 460, 462, 470, 476, 482, 501, 536, 539, 563, 584, 591, 596, 605, 616, 628, 637, 680, 710, 716, 717, 724, 725, 726, 734, 738, 760, 770, 823, 825, 826, 832, 833, 834, 835, 836, 840, 843, 844, 845, 846, 847, 848, 849, 852, 853, 854, 855, 856, 857, 858, 859, 860, 863, 866, 867, 868, 870, 872, 879, 880, 881, 882, 884, 885, 886, 887, 888, 889, 890, 891, 892, 897, 898, 899, 900, 901, 902, 904, 907, 908, 910, 911, 912, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 933, 935, 936, 937, 938, 939, 943, 947, 949, 950, 958, 963, 964, 965, 969, 971, 975, 986, 987, 989, 990, 991, 992, 993, 1021, 1044, 1045, 1047, 1053, 1068, 1079, 1094, 1095, 1103, 1104, 1114, 1118, 1119, 1136, 1137, 1140, 1142, 1143, 1148, 1150, 1152, 1170, 1195, 1200, 1201, 1202, 1214, 1228, 1231, 1265, 1266, 1284, 1295, 1303, 1309, 1311, 1318, 1320, 1331, 1334, 1338, 1339, 1341, 1361, 1366, 1371, 1373, 1374, 1375, 1376, 1378, 1380, 1381, 1382, 1383, 1384, 1385, 1386, 1387, 1388, 1389, 1392, 1393, 1395, 1396, 1398, 1399, 1400, 1401, 1402, 1403, 1404, 1405, 1406, 1407, 1408, 1410, 1413, 1415, 1416, 1417, 1419, 1420, 1421, 1423, 1424, 1425, 1426, 1427, 1434, 1441, 1460, 1461, 1503, 1506, 1507, 1509, 1511, 1513, 1520, 1523, 1528, 1529, 1530, 1531, 1540, 1541, 1546, 1547, 1549, 1550, 1552, 1553, 1555, 1558, 1561, 1570, 1571, 1572, 1576, 1578, 1579, 1580, 1581, 1583, 1585, 1610, 1611, 1616, 1620, 1621, 1622, 1623, 1626, 1635, 1636, 1637, 1638, 1639, 1641, 1644, 1645, 1654, 1655, 1660, 1661, 1663, 1664, 1671, 1672, 1673, 1674, 1675, 1676, 1689, 1690, 1691, 1696, 1699, 1715, 1718, 1720, 1740, 1746, 1753, 1754, 1756, 1758, 1759, 1761, 1772, 1773, 1781, 1782, 1784, 1785, 1786, 1787, 1788, 1789, 1790, 1791, 1792, 1793, 1794, 1795, 1796, 1798, 1801, 1803, 1804, 1805, 1807, 1808, 1809, 1810, 1811, 1812, 1813, 1817, 1818, 1819, 1820, 1821, 1822, 1823, 1825, 1826, 1827, 1828, 1830, 1831, 1832, 1834, 1836, 1837, 1838, 1840, 1841, 1843, 1845, 1847, 1848, 1850, 1852, 1853, 1855, 1856, 1857, 1861, 1862, 1864, 1873, 1875, 1880

Buckwheat

124, 351, 428, 447, 573, 589, 678, 801, 932, 1353, 1354, 1367, 1372, 1464, 1524, 1588, 1589, 1631, 1669, 1670, 1687, 1737, 1747, 1833

Flax

43, 1292, 1431, 1477, 1579, 1629

Mustard

567, 1367, 1588, 1702

Sea Oats

541, 736, 1778, 1808

Rye

12, 26, 27, 73, 79, 129, 157, 158, 166, 184, 221, 252, 353, 399, 400, 419, 429, 455, 479, 491, 494, 495, 499, 523, 543, 544, 553, 554, 557, 561, 610, 614, 656, 659, 661, 670, 671, 672, 679, 689, 729, 730, 732, 761, 767, 771, 781, 783, 787, 788, 797, 802, 804, 830, 842, 874, 893, 903, 942, 948, 1171, 1301, 1307, 1321, 1349, 1351, 1494, 1516, 1522, 1551, 1559, 1573, 1584, 1609, 1632, 1653, 1693, 1698, 1707, 1743, 1755, 1763, 1766, 1770, 1774, 1775, 1777, 1779, 1800, 1816, 1867

Barley

3, 10, 11, 13, 14, 16, 18, 20, 22, 23, 28, 30, 40, 41, 42, 43, 45, 46, 47, 48, 49, 50, 52, 56, 57, 61, 65, 67, 69, 71, 76, 77, 82, 86, 87, 88, 89, 90, 92, 93, 95, 97, 103, 108, 111, 112, 114, 128, 133, 135, 141, 154, 155, 159, 163, 164, 170, 173, 175, 176, 177, 180, 181, 182, 183, 185, 186, 187, 189, 190, 191, 192, 194, 197, 202, 203, 204, 205, 206, 207, 208, 210, 211, 212, 213, 218, 219, 220, 222, 223, 224, 225, 226, 227, 229, 232, 233, 234, 235, 236, 237, 238, 240, 242, 247, 248, 249, 250, 253, 254, 255, 257, 259, 262, 263, 264, 265, 266, 267, 268, 270, 271, 273, 274, 275, 276, 278, 279, 280, 281, 282, 284, 285, 288, 289, 291, 292, 293, 294, 295, 296, 297, 302, 304, 306, 307, 315, 316, 317, 318, 319, 320, 321, 328, 337, 338, 341, 342, 345, 346, 348, 349, 350, 352, 354, 356, 358, 361, 362, 364, 366, 367, 369, 371, 372, 373, 377, 378, 379, 382, 383, 384, 386, 388, 389, 393, 396, 397, 398, 401, 404, 405, 407, 408, 411, 413, 414, 415, 417, 418, 421, 422, 423, 425, 433, 434, 435, 436, 437, 438, 442, 444, 446, 450, 451, 452, 453, 454, 456, 459, 463, 464, 465, 466, 467, 468, 469, 472, 473, 474, 475, 478, 480, 483, 484, 486, 487, 488, 489, 493, 495, 496, 498, 500, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 514, 516, 517, 518, 519, 521, 522, 524, 525, 526, 527, 530, 531, 533, 534, 537, 538, 546, 547, 549, 550, 552, 555, 556, 558, 559, 562, 564, 565, 566, 567, 568, 569, 570, 571, 574, 575, 577, 580, 581, 582, 583, 585, 586, 587, 588, 593, 594, 595, 597, 601, 602, 604, 608, 611, 612, 613, 617, 620, 621, 622, 623, 624, 625, 626, 627, 629, 630, 631, 632, 635, 636, 638, 639, 642, 643, 644, 646, 647, 649, 650, 654, 655, 658, 660, 663, 664, 665, 666, 667, 668, 669, 674, 675, 676, 677, 681, 687, 688, 690, 693, 694, 695, 703, 704, 705, 708, 712, 713, 715, 722, 723, 731, 735, 737, 739, 741, 742, 743, 744, 746, 747, 750, 751, 752, 753, 754, 756, 766, 768, 772, 779, 780, 784, 786, 789, 790, 792, 793, 794, 795, 798, 799, 801, 805, 806, 807, 808, 809, 811, 812, 814, 818, 821, 822, 828, 831, 838, 839, 841, 850, 851, 862, 865, 869, 876, 877, 883, 894, 895, 906, 914, 915, 917, 918, 931, 940, 941, 945, 951, 952, 953, 954, 957, 970, 976, 977, 978, 979, 981, 982, 983, 984, 985, 988, 994, 996, 997, 998, 999, 1000, 1001, 1004, 1005, 1006, 1007, 1008, 1009, 1010, 1011, 1012, 1013, 1014, 1015, 1016, 1017, 1018, 1019, 1020, 1022, 1023, 1024, 1030, 1031, 1040, 1041, 1043, 1048, 1050, 1051, 1052, 1054, 1055, 1056, 1057, 1058, 1060, 1061, 1062, 1064, 1066, 1067, 1071, 1072, 1074, 1075, 1076, 1077, 1078, 1080, 1081, 1083, 1086, 1087, 1088, 1089, 1090, 1091, 1092, 1093, 1105, 1106, 1107, 1108, 1109, 1111, 1112, 1115, 1117, 1120, 1121, 1122, 1123, 1124, 1125, 1127, 1128, 1129, 1130, 1134, 1135, 1141, 1144, 1145, 1146, 1149, 1151, 1155, 1156, 1159, 1163, 1165, 1167, 1168, 1169, 1173, 1175, 1178, 1179, 1180, 1181, 1182, 1185, 1186, 1187, 1189, 1190, 1191, 1192, 1194, 1196, 1203, 1204, 1207, 1208, 1209, 1210, 1213, 1216, 1224, 1225, 1229, 1234, 1235, 1236, 1237, 1238, 1239, 1240, 1241, 1242, 1243, 1244, 1245, 1246, 1250, 1251, 1252, 1256, 1257, 1258, 1259, 1260, 1261, 1262, 1263, 1264, 1267, 1268, 1269, 1270, 1271, 1276, 1277, 1278, 1279, 1280, 1281, 1282, 1283, 1286, 1287, 1289, 1290, 1291, 1293, 1294, 1297, 1298, 1304, 1308, 1316, 1317,

Barley continued:

1322,1323,1324,1325,1326,1327,1328,1329,1330,1331,1332,1333,1342,
1343,1346,1348,1350,1352,1355,1358,1370,1379,1390,1411,1412,1414,
1422,1428,1429,1430,1431,1432,1436,1442,1443,1444,1445,1446,1447,
1448,1449,1450,1451,1452,1453,1454,1455,1456,1458,1463,1466,1467,
1468,1469,1476,1486,1487,1488,1490,1491,1493,1495,1496,1497,1498,
1499,1500,1501,1502,1504,1508,1524,1525,1527,1535,1544,1557,1560,
1563,1565,1568,1569,1575,1582,1586,1587,1589,1592,1597,1598,1599,
1602,1605,1606,1608,1617,1624,1627,1628,1630,1646,1647,1649,1650,
1651,1658,1662,1665,1667,1668,1678,1684,1685,1694,1695,1700,1704,
1706,1708,1709,1710,1711,1713,1714,1716,1717,1719,1722,1723,1724,
1725,1726,1727,1728,1729,1731,1732,1733,1734,1735,1736,1738,1744,
1747,1750,1760,1764,1765,1768,1776,1780,1783,1797,1799,1802,1806,
1814,1824,1829,1835,1839,1842,1854,1859,1863,1865,1868,1870,1871,
1872,1874,1876,1879

Oats

5,15,17,19,21,24,25,37,38,39,51,58,62,63,64,78,80,81,84,85,90,91,
94,99,102,106,107,109,113,116,120,121,126,127,130,131,132,134,
136,137,138,139,140,147,148,152,160,171,172,179,188,193,199,200,
209,214,215,217,228,230,239,245,246,247,251,256,258,260,261,269,
272,277,283,286,287,290,299,303,310,311,312,313,322,323,324,325,
326,335,339,340,344,347,355,357,359,360,363,365,368,370,374,376,
380,381,385,390,391,394,402,403,406,409,410,412,416,420,424,426,
430,431,432,439,441,443,445,449,461,471,477,481,485,490,492,497,
500,512,513,515,520,528,529,532,535,540,542,545,548,557,560,572,
576,578,579,590,592,598,599,600,603,606,607,609,618,619,633,634,
640,641,645,648,651,652,653,657,662,673,682,683,684,685,686,691,
692,696,697,698,699,700,701,702,706,707,709,711,714,718,719,720,
721,727,728,733,740,745,748,749,755,757,758,759,762,763,764,765,
769,773,774,775,776,777,778,782,785,791,796,800,803,810,813,815,
816,817,819,820,824,827,829,837,864,871,875,878,896,905,909,913,
916,934,944,946,955,956,959,960,961,962,966,967,968,972,973,974,
975,980,995,1002,1003,1025,1026,1027,1028,1029,1032,1033,1034,
1035,1036,1037,1038,1039,1042,1046,1049,1059,1063,1065,1069,1070,
1073,1082,1084,1085,1096,1097,1098,1099,1100,1101,1102,1110,1113,
1116,1126,1131,1132,1133,1138,1139,1147,1153,1154,1157,1158,1160,
1161,1162,1164,1166,1172,1174,1176,1177,1183,1184,1188,1193,1197,
1198,1199,1205,1206,1211,1212,1215,1217,1218,1219,1220,1221,1222,
1223,1226,1227,1230,1232,1233,1247,1248,1249,1253,1254,1255,1272,
1273,1274,1275,1288,1292,1296,1299,1300,1302,1305,1306,1310,1312,
1313,1314,1315,1319,1335,1337,1340,1344,1345,1347,1356,1357,1359,
1360,1362,1363,1364,1365,1368,1369,1391,1394,1397,1409,1418,1431,
1435,1437,1438,1439,1440,1457,1459,1462,1465,1470,1471,1472,1473,
1474,1475,1477,1478,1479,1480,1481,1482,1483,1484,1485,1489,1492,
1501,1502,1504,1505,1510,1512,1514,1517,1519,1521,1526,1532,1533,
1534,1536,1537,1538,1539,1542,1543,1544,1545,1548,1554,1562,1564,
1566,1567,1574,1577,1589,1590,1591,1592,1593,1594,1595,1596,1597,
1598,1599,1600,1601,1602,1603,1604,1605,1606,1607,1612,1613,1614,
1615,1618,1619,1625,1626,1629,1633,1634,1640,1642,1643,1648,1652,
1656,1657,1659,1666,1677,1679,1680,1681,1682,1683,1686,1688,1692,

Oats continued:

1697,1703,1705,1712,1721,1730,1739,1741,1742,1745,1748,1749,1750,
1751,1752,1757,1762,1767,1769,1771,1774,1814,1815,1844,1846,1849,
1851,1858,1860,1866,1869,1877