Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

UNITED STATES DEPARTMENT OF AGRICULTURE

DEPARTMENT BULLETIN No. 1334

Washington, D. C.

November, 1925

TESTS OF BARLEY VARIETIES IN AMERICA

By

HARRY V. HARLAN, Agronomist in Charge, MARY L. MARTINI, Assistant Botanist, and
MERRITT N. POPE, Associate Agronomist, Barley Investigations, Office of Cereal Investigations
Bureau of Plant Industry

CONTENTS

	Page
Scope of the Bulletin	1
Barley Growing in America	2
Preparation of Material	13
Discussion of Station Yields	17
Regional Adaptation of Varieties	154
Well-Known or Promising Varieties	159
Accession Data of Cereal Investigations Numbers	175
Botanical Comparison of Prominent Varieties	205
Best Varieties of the Period from 1917 to 1921, Inclusive	207
Distribution of Varieties by Experiment Stations	212
Index	213

WASHINGTON
GOVERNMENT PRINTING OFFICE
1925

UNITED STATES DEPARTMENT OF AGRICULTURE

DEPARTMENT BULLETIN No. 1334

Washington, D. C.

November, 1925

TESTS OF BARLEY VARIETIES IN AMERICA

By HARRY V. HARLAN, *Agronomist in Charge*, MARY L. MARTINI, *Assistant Botanist*, and MERRITT N. POPE, *Associate Agronomist, Barley Investigations, Office of Cereal Investigations, Bureau of Plant Industry*

CONTENTS

	Page		Page
Scope of the bulletin.....	1	Botanical comparison of prominent varieties.....	205
Barley growing in America.....	2	Best varieties of the period from 1917 to 1921, inclusive.....	207
Preparation of material.....	13	Distribution of varieties by experiment stations.....	212
Discussion of station yields.....	17	Index.....	213
Regional adaptation of varieties.....	154		
Well-known or promising varieties.....	159		
Accession data of Cereal Investigations numbers.....	175		

SCOPE OF THE BULLETIN

Barley has been a minor crop in all but a few States of the United States. Many importations have been made, tried, and discarded by individuals and experiment stations. Often the same varieties have been reimported and retested at the same stations later. A few men have been particularly interested in barley. Until now no effort has been made to assemble the incidents and results of the early work. This task is becoming more difficult each year. Already the results of many tests have been lost. Cereál breeding is entering on a new basis, and if ever a summary is to be made of the pioneer work it is best made now.

The year 1921 has been chosen arbitrarily as the last one from which yields are included. This year divides fairly well a transition period. While the replication of plats is now the method in general use for the testing of varieties, the data for the year 1921 include about the last of the yields from unreplicated plats. Also, at about this time a large number of varieties produced by younger plant breeders made their appearance. The closer relationship existing between experiment stations resulted in these new varieties being grown promptly at a large number of places. To show the value of these new varieties, yields of several seasons after 1921 would need to be included.

This bulletin is the result of an attempt to gather together all the material available. The writers know that their work is far from complete. They feel sure that owing to the incompleteness of the records and the early date of many of the references the identification

of some varieties can not be certain, but they feel that they are submitting something tangible which will be of use to the agronomists who have assisted them.

As far as possible all the yields obtained in field plats in the United States have been included. In some instances plat yields have been rejected because of failure to establish the identity of the variety or because of accident of season that vitiated the results. Doubtless there are included many results that the station agronomists would desire omitted. However, as one of the main purposes of this bulletin is the defining of ecological boundaries of various types of barleys, it is thought that these yields are valuable, at least from that standpoint.

While the writers of this bulletin may point with some justice to many tedious months of compilation and to an endless effort of identification, the real authors of this bulletin are the agronomists of the various States. It is only by their unselfish donation of data that this compilation has been made possible. That all States are not represented in the compilations is due not to any unwillingness on the part of their officials, but to the fact that they have not grown barley. The agronomists of every barley-growing State but one have made available all their station results on barley. They have not only granted the use of these data, but they have extracted the results and histories, which has been no small task. At a few of the stations some of the results were secured by them in cooperation with the United States Department of Agriculture, but even in the cooperating States the material furnished was far more than that jointly obtained. The writers desire hereby to express their appreciation of these many courtesies received from State agronomists.

BARLEY GROWING IN AMERICA

The demarcation of areas of adaptation is obviously an ecological study. The usual method of studying variation in plants is by classifying them as to their morphological characters. Fortunately, groups of varieties which vary in their climatic requirements also vary in their taxonomic features as well. In either case the grouping is merely a convenient method of study. When large numbers of varieties are considered the group limits merge, whether the separations be made on an ecological or a morphological basis. Even so, the discussion is materially facilitated by the use of these distinctions, and in most cases there is no confusion.

VARIETAL GROUPS

There are several ecological groups of barley which are well marked when typical varieties are considered. The most common of these in the United States are the Manchurian, which embraces most of the 6-rowed barleys of the temperate plains of Europe and Asia, and the Coast group, which includes many 6-rowed barleys from North Africa and Spain. Other well-known groups include the Chevalier and Thorpe of England, the Hanna of Bavaria and Austria, the Smyrna group of Asia Minor, and the hull-less forms of Central Asia. The differences between Chevalier, Gold, Thorpe, Manchuria, and Coast varieties are very apparent, and there is no overlapping. These barleys are adapted also to definite sets of ecological condi-

tions. Other varieties adapted to these districts are often intermediate in character, especially where the districts merge into each other; that is, where sets of conditions overlap. The Chevalier variety is typical of England. Very similar barleys, however, are found in Denmark and Bohemia, where conditions similar to those of England exist. The Hanna barleys of Bavaria, Austria, and Bohemia differ somewhat from the 2-rowed sorts which are occasionally found in southern Russia.

The nodding Coast and more erect Peruvian types of North Africa are quite characteristic of that region. Stavropol, a variety brought from the dry Stavropol region of southern Russia, is similar to the nodding Coast type. In the areas north and east of the Black and Caspian Seas there are numerous 6-rowed barleys of local adaptation. These are grown on the border land between regions suited to the culture of barley of the North African type and those suited to that of the Manchurian type. Many of these varieties are intermediate between these groups. A variety partaking of the characters of two groups is defined to a certain extent by placing it in either one of them.

FOREIGN SOURCES

The original introductions of barley into America were entirely from Europe and North Africa. Later, barleys from Asia Minor were imported in a commercial way. The Nepal barley of India was introduced largely as a curiosity. In recent years numerous barleys have been brought into this country from Asia. Yields of many varieties from different parts of the world are reported in the tables of this bulletin. The commercial-barley acreage of the United States at the present time consists almost entirely of barleys from two ecological districts. These are (1) North Africa and (2) the central Eurasian Plain, which reaches through the level temperate countries from the North Sea, through Russia, and by a slender belt through Siberia to the Pacific Ocean.

The barleys of the western part of the United States are entirely of the North African group, with the exception of those grown in certain valleys, such as the Salinas of California, where unusual conditions occur. The North African barleys came into America with the Spanish invasion and spread with the settlements on the Pacific coast. These introductions were grown almost to the exclusion of all other varieties until the last few years, when two other North African barleys were introduced into general field agriculture. These were the Club Mariout and California Mariout from the district of Mariout, Egypt, the latter from nonirrigated lands. These varieties are now grown on an acreage which seems to assure their place in the western part of the United States, at least until others are brought in or developed. The 6-rowed barleys grown in the Mississippi Valley are of the type best adapted to the Eurasian Plain. The western European varieties of this group have been displaced in the last 50 years by introductions from Manchuria and Germany.

The two sources of barleys which are in most urgent need of exploration are North Africa, which has given us the dominant types of the Pacific coast, and Manchuria, which has furnished the most vigorous and most disease-resistant varieties of the Mississippi Valley.

In the Great Basin between the Rocky Mountains and the ranges along the Pacific coast there are many high plateaus and many mountain valleys, each differing from the neighboring ones. Barleys suited to the special valleys may best be found in southern and central Asia, where similar ecological conditions obtain.

HISTORY OF BARLEY GROWING

EARLY-GROWN BARLEYS

The first barleys grown in the various American settlements doubtless were those varieties from the homes of the settlers. The English probably brought with them the late 2-rowed barleys commonly grown in England, such as Chevalier and Thorpe. The Dutch brought the barleys of the mainland, and the Spanish the barleys of the Coast type. The Spanish settlements were largely in arid regions adapted to the culture of barleys of the Coast type. The barley of Spain was of North African origin, well suited to Mexico, western South America, and California. Although its history is not easily traced, it is likely that this cereal was an immediate success, that it was taken up widely in Mexico, and that it went into California with the missions. It is still the prevailing barley of the Western States.

Along the Atlantic seaboard conditions were not so favorable. All the settlements were in a region of humid summers, and most of them south of the latitudes where barley does well in humid growing seasons. The Chevalier and Thorpe types of England were suited to the eastern Provinces of Canada, but it was only when the settlements had penetrated western New York that an especially favorable area was found in the Eastern States. This area was better adapted to the culture of the Hanna and the 6-rowed varieties of the Continent than to the 2-rowed sorts of England. Barley production became important early in the history of New York.

After agricultural periodicals became common there are frequent notes on barley production, many of which indicate the nature of the varieties grown. The reports of the grain trade of Albany in the fifties show that 6-rowed barley was more common than 2-rowed in New York. A note by Chief Inspector Stevens, of Chicago, in the *Prairie Farmer* in 1860 stated that the 4-rowed barley was best suited for the Chicago market "because a majority of that grown in the Northwest is of that variety." In 1865 all correspondents of this paper seemed to be growing 6-rowed barley. The Wisconsin University Experiment Farm began distributing the Manchuria barley about 1873. This variety was widely disseminated, and its culture spread from farm to farm from the centers started by the distribution from the university farm. At the present time practically all barley that comes into the market in the northern Mississippi Valley is Manchuria or Oderbrucker.

The question naturally arises as to the identity of the 6-rowed barley widely grown before the introduction of Manchuria and Oderbrucker. There is little definite information on this point. In all probability it was very similar to Manchuria and Oderbrucker. It is inconceivable that it has now entirely disappeared. It probably is still grown by a small percentage of farmers. In all likelihood it differs little in appearance in the field or when threshed from

the more recent introductions. This early 6-rowed barley probably consisted of the mixed stocks of the 6-rowed barley under cultivation in England, Scotland, and western continental Europe at the time of its importation into this country. These mixtures seem to differ from the more vigorous strain of Manchuria only in vigor and in the smaller proportion of biotypes with blue aleurone layers. Some of the strains also are characterized by slightly larger kernels.

Barley was being bred by selection at an early date in western Europe. The efforts at improvement were expended on the 2-rowed varieties. Until very recently the 6-rowed barleys remained as they were developed by ecological conditions. The Oderbrucker barley of Germany has proved to be the best west European 6-rowed sort tried in this country. As originally introduced it was not widely different from the Manchuria form, but has exhibited somewhat less vigor and a narrower range of adaptation.

TREND OF BARLEY GROWING

As has been remarked before, barley was introduced into America by the early colonists. The colonies first founded were along the Atlantic seaboard. In recent times barley has not done well on the Atlantic Coastal Plain. It is not well adapted to the types of soil most common there. It is probable that it grew to better advantage on the virgin soils when the land was first cleared. It was introduced into all of the colonies, where it was in demand as a grain for brewing. Both in the early colonies and in the later settlements of the Mississippi Valley barley invariably was planted, and considerable acreage developed near all of the larger cities. This acreage was in response to a city demand, and barley often was grown in districts to which it was not suited.

In the annual report of the United States Patent Office (agricultural portion) for 1853 (pp. 156, 157) there are several notes on the early history of barley. According to this report the introduction of barley in the North American colonies may be traced back to the periods of their settlement. It was grown by Gosnold, together with other English grains, on Marthas Vineyard and the Elizabeth Islands in 1602 and by the colonists of the London Company in Virginia in 1611. By the year 1648 it was grown in abundance in that colony, but its culture was allowed to decline soon after, in consequence of the more profitable and increased production of tobacco.

Barley appears to have been cultivated in New Netherlands as early as the year 1626, as samples of the harvest of that year, grown by the colonists on Manhattan Island, were sent to Holland with other grains as evidence of their prosperous condition.

According to the records of the governor and of the Massachusetts Bay Company in New England, barley was introduced into that colony in 1629. In 1633 good crops were grown in Lynn.

In 1796 the chief agricultural product of Rhode Island was barley, considerable quantities of which were raised.

Barley grew better in the colonies of New England than in those farther south, with the possible exception of the southeastern part of Pennsylvania, where the soils were especially suited to its production. The combination of favorable soil and climatic conditions was not found in any settlement in the United States until these settle-

ments had reached central New York. Here both the soil conditions and the cool summers were favorable to the growing of barley.

A series of maps showing the production of barley in the census years from 1839 until 1919, inclusive, is given as Figures 1 to 8, inclusive. In 1839 some barley was still grown in the cooler sections of the Atlantic seaboard. Most of this was in Maine, although some was cultivated in New Hampshire and Massachusetts. The greater part of the American crop, however, was produced along the line of the Erie Canal, which traversed a section where the production of barley was profitable and gave ready access to the Albany, New York, and other city markets. The distribution of barley acreage along the canal is very apparent in Figure 1.

FIG. 1.—In 1839 most of the barley of the United States was produced in New York. The production was heaviest along the line of the Erie Canal. There also was a fringe of production near the coast of New England and a small acreage on the favorable soils of southeastern Pennsylvania. The production of barley in 1849 was still centered in New York State. In the Mississippi Valley a city demand for barley for brewing was reflected in the beginnings of production near Cincinnati, St. Louis, and Milwaukee.

There was but little change in the location of the acreage between 1839 and 1849. A little less was grown along the Atlantic seaboard. The New York acreage was more concentrated in the western two-thirds of the State. Barley cultivation had given way to the production of crops of higher money value on the soils of southeastern Pennsylvania. In response to the demands of growing cities, areas of production appeared near Pittsburgh, Cincinnati, St. Louis, and Milwaukee. This, of course, was a result of the demand for brewing barley.

By 1859 the New York acreage was centered in the western part of the State (fig. 2). The acreage around Cincinnati had expanded greatly, owing to the large demand for brewing in that city. This acreage disappeared in later years, when transportation was developed to the extent that barley could be brought from other sections more cheaply than it could be grown around Cincinnati.

The acreage about St. Louis, where the conditions are not favorable for the production of barley, never expanded as did that near Cincinnati. The cultivation of barley about Milwaukee, on the

other hand, was extended very rapidly. In southern Wisconsin, northern Illinois, and along the Mississippi in eastern Iowa conditions were particularly favorable for securing higher yields. The demand of the Milwaukee and Chicago markets was such as to encourage production. The river transportation not only gave a market in St. Louis, but probably was responsible for the fact that a large acreage was never developed near that city.

Barley was introduced into California with the early missions. It was grown wherever there were settlements, because the districts were very well suited to barley production and there was no other grain which could be grown so successfully for feed. To this demand was added the brewing demand on the west coast. In 1848 gold was discovered in California. The population in the area about San Francisco increased very rapidly. With this increase there was a

FIG. 2.—By 1859 New York was losing its dominant position in barley production. Production had increased in southwestern Ohio, about Cincinnati, and still more notably in northern Illinois and southern Wisconsin. In this section barley had spread away from the local city market and become a general farm crop. Only a slight increase took place near St. Louis. The settlement of central California, following the discovery of gold, resulted in a production in the State fully as great as that in New York.

great expansion of the barley acreage. By 1859 California was producing as much barley as New York.

In 1869 the barley acreage was in the same regions as in 1859, but the area was more extensive (fig. 3). Both California and New York were growing more barley. The malting industry of western New York had developed greatly, and there was a resultant increase in production in that section of the State. The acreage in the eastern portion of Michigan, which is especially favorable to the production of barley, had increased. The greatest expansion occurred in the districts west and north of Chicago and Milwaukee. Conditions in the upper Mississippi Valley are well suited to the raising of barley. The extensive cultivation of this grain followed the settlements as closely as transportation developed to take care of the surplus crop. Southeastern Minnesota is a district in which both soil and climate are quite favorable and the acre yields are large. There was a considerable acreage in this part of the State in 1869, as may be seen in Figure 3.

By the year 1879 a large quantity of barley was raised in eastern Oregon and Washington (fig. 4). The conditions here are favorable to the growing of barleys of the type found in California. Barley

FIG. 3.—In the decade from 1860 to 1869 barley became commonly cultivated in southeastern Minnesota and its culture was begun in eastern Oregon and Washington. There was a notable increase in southern Wisconsin and northern Illinois, and some increase in the southern section of the Great Plains area and in the Mountain States. California and New York, however, remained the most important producing States.

was grown over a wider acreage in California than in 1869. Figure 4 shows that centers of production had become prominent in western New York, eastern Wisconsin, southeastern Minnesota, and central California. These centers were maintained for many years. Barley was grown as a cash crop, and the production was far beyond the local

FIG. 4.—In the years from 1870 to 1879 the total production of barley in the United States increased almost 50 per cent. The culture of barley was extended westward into eastern Nebraska and southeastern Dakota. It also became more widely distributed in the Pacific Coast States, while a small production was developing in Utah, Nevada, and Arizona.

needs. It will be noted that there was a heavy increase in the production in southeastern Nebraska and a beginning of cultivation in the southeast corner of Dakota Territory. The acreage was not

maintained in southeastern Nebraska in later years, while that of Dakota Territory expanded along the Red River and later over the eastern half of both Dakotas.

In the 10 years from 1879 to 1889 the development of centers of production in barley was accentuated. As a cash crop in 1889 it was grown on large acreages in five different centers, as may be seen from Figure 5. These were western New York, eastern Wisconsin, southeastern Minnesota, western Iowa, and central California. A considerable acreage was grown in Michigan, the area extending into northwestern Ohio. Barley was not so widely grown in northern Illinois as it had been previously. There was a marked expansion of the acreage in the Red River Valley.

In 1890 a law was enacted which raised the tariff to 30 cents per bushel. Western New York up to this time had been a great malting center. The malt houses of this district had obtained large

FIG. 5.—By 1889 there was a marked concentration of the areas growing barley for the malting market. On the favorable soils of western New York, southeastern Wisconsin, southeastern Minnesota, and northwestern Iowa, and in the central valley of California barley was grown as a money crop. At the same time production was increasing in the Red River Valley of Minnesota and North Dakota. The production of barley about Cincinnati decreased in the face of competition from the northern Mississippi Valley.

quantities of barley from Canada. These houses had given a ready market to all the barley that could be produced in western New York. The tariff of 1890 was sufficient to make importation from Canada unprofitable. As malt weighs less per bushel than barley the malt houses of Wisconsin and Minnesota could ship to the eastern seaboard more cheaply than it was possible for the eastern malt houses to purchase barley for malting. The malting industry was gradually transferred from western New York to Wisconsin and neighboring States. This, in turn, decreased the market for barley in western New York, and this with the competition of other crops gradually reduced the acreage.

In 1899, as shown in Figure 6, western New York was no longer one of the principal centers of production. However, the loss of acreage in New York State was much more than made up by the expansion in the Red River Valley and in eastern Oregon and Washington. By 1899 a large acreage of barley was grown in western

Kansas. Both in Nebraska and Kansas there were extensive changes, not only in the acreage but in the sections in which barley was grown. The portions of these States best adapted to the production of barley,

FIG. 6.—In 1890 the tariff on barley was raised to 30 cents per bushel. The malt houses of western New York had been securing part of their barley from Canada, but this tariff made the importation of barley unprofitable. The near-by supplies were insufficient and the malting industry was transferred to Wisconsin and Minnesota. In New York the area devoted to barley decreased after 1890. Production increased notably in Minnesota and California, also in eastern Washington and Oregon, in the Red River Valley, and in the central section of the Great Plains area.

when economic as well as climatic conditions are taken into consideration, were not yet growing barley.

In Figure 7, showing the production of barley in 1909, the center of cultivation in Kansas is more nearly in the section in which it will

FIG. 7.—The decade ending in 1909 was marked by the rapid expansion of the acreage in the Dakotas and the definite beginning of a center of production in northwestern Kansas. Minnesota was now the leading State, producing, with the Dakotas, almost half of the national crop. California, however, was a close second to Minnesota, with an average production of 29 million bushels in the last five years of this decade.

be permanent. The climatic and soil conditions of northwestern Kansas are better suited to the production of barley than is the case either east or south. It is probable that for economic reasons

the cultivation will be expanded both east and south, but northwestern Kansas, as is shown in this figure, will probably remain the center of production. By 1909 the acreage of barley on either side of the Red River had been enormously expanded. North Dakota, South Dakota, Minnesota, and Wisconsin were producing a large proportion of the total crop of the United States. The acreages around Cincinnati and St. Louis had long since disappeared.

After 1909 the acreage in southeastern Minnesota gradually decreased. This area was one of very high fertility, and only the profits of barley production prevented its development for diversified farming at an earlier date. As general farming increased in southeastern Minnesota, the acreage of barley grown as a cash crop decreased. This movement was already beginning, in advance of the decrease of the demand from malt houses which came with the increase of statutory prohibition. The crop of 1919 (fig. 8) was not a normal

FIG. 8.—The production of barley in 1919 was abnormally distributed. The great increase in the spring-wheat acreage in the Dakotas, coupled with a low acre yield of barley, resulted in a lower production in these States in 1919 than in the years before or since. The decrease in southeastern Minnesota in 1919 was caused chiefly by the gradual drift away from barley as a money crop, a process which had been going on since 1910. The most remarkable development of production was in the central Great Plains area, especially Kansas. This probably is part of a permanent modification of the agricultural practice of the section

one as far as barley production was concerned. The war prices of wheat and the campaign to increase its acreage had resulted in a considerable expansion in the area sown to spring wheat in that year. The greatest increase in this crop came in the barley-producing sections of North Dakota, South Dakota, and Minnesota. The great difference which is apparent in the production of these two years, as shown in Figures 7 and 8, was not due entirely to the reduced acreage of barley. The acre yield of barley in the northern Mississippi Valley in 1919 was low, so that the production was reduced relatively much more than the acreage. The smaller production in southeastern Minnesota, on the other hand, was due in large part to the advent of diversified farming in that area, which had replaced the grain farming commonly practiced until about 1909.

The most interesting change of acreage shown in the 1919 map is in Nebraska, Kansas, and Oklahoma. A center of heavy production

is developing in northwestern Kansas. This area has extended southward into Oklahoma and Texas. The section is of especial interest, as there is an overlapping of conditions which allow types of barley adapted to quite different ecological regions to be grown in the same general area. In central Nebraska barleys of the Manchuria type grow very well. In northwestern Kansas those of the Coast and Stavropol types are most commonly grown. These varieties are cultivated farther south, and in Oklahoma their culture is found in the same district with winter barleys of the Tennessee Winter types. This is the only district in the United States where the extensive culture of both winter and spring varieties meets. In the Central States there is a margin of several hundred miles between the southern limit of profitable culture of spring barley and the northern limit of profitable culture of winter barley. It is true that in Ohio, Pennsylvania, and Maryland, and in the Palouse district of Washington, Oregon, and Idaho, both winter and spring barleys can be grown. Winter barleys in these States, however, are of only local importance, and the acreage is small.

The general distribution of barleys over the United States has followed the course of the settlements. It is a natural inquiry as to what influence agricultural experiment stations have had, especially on the varieties cultivated. Prior to 1870 the barleys under cultivation were doubtless those that had been introduced by the colonists, and agronomic experiments probably had little influence. Since then, however, the varieties grown on the farms have been greatly affected. As has been stated before, the experiment farm at Madison, Wis., began testing and distributing barley about 1873.

It is likely that most of the 6-rowed barley of the northern Mississippi Valley as now grown consists largely of varieties introduced by experiment stations. Recently in California these stations have introduced varieties which have been substituted on a large scale for others previously grown.

EXPERIMENTAL METHODS DEVELOPED

There is a feeling among many agronomists that the early experiments are without value, but a study of the data shows that this feeling is justifiable only in part and is not entirely warranted. The agronomic work of the older experiment stations has passed through three distinct phases of development. In the earliest phase the stations were undermanned and without suitable equipment for carrying on varietal tests. They worked with varieties obtained by chance and often with many of the desirable types absent from the tests. In the middle period both equipment and personnel were on a better basis, and larger collections of material were available for comparison. The tests were conducted, however, on the single-plat basis.

Recently the technic of field testing has been improved. Plats are more carefully laid out. Methods of testing have been developed, and a sufficient number of replications have been carried to form a basis for a statement of the probable mathematical significance of the yield. Unfortunately, one is not able to fix these periods by dates, as each station went through these phases independently. Some stations are not yet replicating plats.

With this better development of methods and with the better training of the younger generation of experimenters has come the doubt as to the value of the early tests. No student of agronomy fails to realize the progress that has been made in the methods of accurate valuation of cereal stocks. There is no reason to think that our present methods are fully accurate. On the other hand, there is no reason to believe that the early work was wholly inaccurate. Probably the early tests did not show the true rank of varieties of nearly equal value. They did afford an indication of these values, however, and most certainly did definitely and correctly establish the type of barley most valuable in many sections. The Wisconsin and Minnesota stations found the Manchuria-Oderbrucker barley to be the most promising type long before plats were replicated or varieties purified. In the typewritten report of the cooperative experiments at Davis, Calif., in 1909 the following statement occurs:

Hardly sufficient data are at hand to make any definite statement as to the most promising varieties. About all that can be said is that, so far, we have no new variety that promises nearly so well as the common barley. It is certain that none will withstand as much wind as the common barley.

The Coast barley is still the common barley of the interior valley of California. Recently two new varieties have been grown. The California Mariout has been widely distributed in California, and the Club Mariout has been grown in a limited area. Both of these are of North African origin, and it is unlikely that either will displace Coast as the dominant variety, although the California Mariout is probably more wind resistant than the Coast.

PREPARATION OF MATERIAL

IDENTIFICATION OF VARIETIES

The identification of the varieties appearing in the early experiments has been very difficult. The order in which the experimental results were received by the writers from the State stations also was unfortunate. If the work could have started with the results from the Central Experimental Farm at Ottawa, Canada, and the State experiment stations of Wisconsin, Minnesota, and North Dakota at hand, the final compilation of data from all stations would have been better. These four stations, with the Agricultural College at Guelph, Ontario, are the key stations in tracing the early varieties. Many of these varieties were produced and many others first were tested at one or the other of these five institutions. Varieties frequently were sent back and forth between stations, unfortunately acquiring a new number with each accession until their identity was much confused. The same varieties also were obtained by seedsmen and reacquired by the stations from them. There was continual opportunity for mechanical mixture and always a possibility of confusing varieties by accident in such a way that they might be carried under a name other than the one originally used.

The identifications were made largely by the records of the stations and checked in various ways. The accession records of the American stations were made available to the writers. These records usually showed the source from which the seed was obtained and stated whether the lot was 2 rowed or 6 rowed. The field records of yields

and growth often contained similar observations. The annual reports from the Canadian experimental farms frequently contained notes in addition to the yield and growth data. These notes were of great assistance. The senior writer visited Ottawa and Charlottetown for the purpose of obtaining further information as to the nature of the varieties grown in Canada and at both places examined specimens of many of the varieties grown in the earlier years at these stations. Among these were many type specimens of Doctor Saunders's hybrids. The senior writer has been identified with barley studies in the United States since 1909. Certain varieties grown at that time have since disappeared, and the memory of things seen even 10 or 12 years ago has been serviceable.

The writers know that the identifications are far from perfect. The records at many places were inadequate. Some stations had conducted varietal experiments for 15 years without an accession record of any kind. The writers, however, have spent nearly two years in an effort to make the presentation the best possible within the limits of reasonable expenditure of time and effort. For many stations it is thought that complete accuracy has been attained. For others the writers have done their best, but some uncertainty remains.

Probably some errors have been introduced by averaging varieties thought to be identical. The number of individual-plat yields was so large that to make the work as useful and available as possible yields were averaged where supposedly identical varieties were included.

From the standpoint of adaptation of barley types, minor errors and impurity of variety are not of dominant importance. There has been no attempt to make varietal recommendations from the results of the early experiments. Most of the barleys grown at that time were commercial varieties of what might be termed ecological races. Examples of these are Coast and Manchuria. Each of these contained a large number of biotypes which could be isolated as pure lines, but in each case these pure lines were adapted to the region of origin of the parent variety. These parent mixtures when grown for many years at a given experiment station in the United States under conditions which differed from those of the original habitat must alter decidedly in the proportions of the constituent strains. They must differ also from the resulting complex of the same mass variety grown at another station. They are comparable, however. The strains of each are similar, if not identical. They are as comparable with each other as with themselves from year to year. The yields of these varying mixtures of unpurified stocks must give a valuable indication of the types of barley adapted to the area where tested.

At experiment stations there also is mechanical mixing of varieties. This was doubtless greater in the earlier years than at present. Equipment and assistance are now more adequate. This mixing of varieties probably has not affected the identification and value of the results as here used. If a variety became mixed to the extent of 10 per cent with another that yielded only 80 per cent as much as the first, the yield would be reduced only 2 per cent. This is much less than is held to be a significant difference in this digest of results.

METHODS OF COMPILATION

The aim in this bulletin is to present the yields of barley varieties grown in field experiments in the United States and Canada. The Canadian yields were included because they have as much value for many American farmers as those of our own stations. Yields are reported of all varieties which are regarded as significant or which can be identified, at least as to group. A few varieties have been included without definite information even as to their group characters, but only when there was some hope of later identification or some particular significance to their yield. The number of varietal yields discarded because of lack of information has been very small, and such varieties usually have been tested but few years with mediocre results.

In many cases the individual yields of synonymous varieties are not reported. Unpurified Manchuria barleys with various station names and numbers are averaged under C. I. No. 244 when these trace directly to the Agricultural College of Guelph, Ontario. Likewise, those which trace directly to the original introductions of the Wisconsin Agricultural Experiment Station are averaged under C. I. No. 241. Averages have been made of commercial sorts only. Whenever a selection of any variety has been made, for example Manchuria, this selection has been given a number and retains its identity in all computations.

With the hybrid variety Horsford the averaging was perhaps more arbitrary, as the confusion was greater. Horsford (C. I. No. 507) here represents the hybrid type obtained by crossing Nepal with barleys of the Manchuria group. The resulting varieties are homozygous for only a few characters at best. While barleys of more than one mating may be listed under this number, they are of the same general type, and the procedure probably is justified. Not all varieties arising from such crosses are included under C. I. No. 507. In some cases the records of ancestry were clear enough to determine the history of the origin. Even in these cases the purpose of this bulletin would have been served by including these varieties also under C. I. No. 507. The greater number of the yields are presented without combining varieties or numbers. At most of the stations few, if any, averages were made.

For points where the data were sufficiently extensive, three tabular compilations have been made, although only the first and third are presented. The first of these shows the yields of the varieties by years. The varieties are listed in the chronological order of their first appearance in the experiments. This compilation embodies statements of the number of years grown; the average yield, which means little; and the average yield of each variety in terms of a percentage of the average of all varieties for the years grown. Suppose, for example, that Svanhals barley was grown during the years 1909, 1911, 1912, and 1913. The yields of all varieties for those years are then added together and divided by the number of yields. The average of the yield of Svanhals for these years is then divided by the average of all varieties for those same years; that is, the yield of Svanhals is computed as a percentage of the weighted mean yield of all varieties. Any basis of computing is open to objection. A comparison of the yearly average yield of Svanhals with the yearly average of all varieties gives undue emphasis to unusual yields in years

when few varieties were grown. On the other hand, the weighted mean does not properly care for the differences of season. The weighted mean, however, is a correct mathematical statement, and it probably is as good a basis of comparison as any which can be used.

At stations where the number of varieties tested has been large, they can be arranged in groups. At many places, for example, several varieties of the Hanna group and several of the Manchuria group have been grown. Where the results are not evident in the first compilation, a digest has been made by placing all varieties of the various types in groups and a comparison made of the resulting groups. This is not a very satisfactory method, in that the groups contain varying numbers of varieties, and these seldom can be compared for the same number of years. However, the adaptation of the groups is so different that the errors due to the difficulties of compilation usually are less than the variation in yield of the groups. The performance of barleys of the different types is much more easily determined in a group digest than in the tabulation of individual yields. The details of this digest are not presented in tabular form, as the relationship can be clearly and concisely stated in the text itself.

Most agronomists are interested in a few prominent well-known varieties. A second digest, therefore, has been made and the results reported in tabular form. In this second digest a well-known variety of each group has been taken and compared with well-known varieties from the other groups. In many cases the choice has been difficult. It is necessary for reasonable comparison that each pair of varieties shall have been in the experiment during several coincident years. This limits the choice of varieties. Those used have not always been the best of the group to which they belong. They are representative in most instances, and they have been chosen with a view both to their local importance and to their inclusion in experiments elsewhere. Usually varieties that have produced good yields over several years at any station have been sent to other stations. Occasionally a high-yielding variety has been grown at one or only a few stations, but this is the exception. For the most part the digest is intended to cover the period of experimentation as well as can be done. The treatment necessarily varies as the data from the stations vary.

Later in this bulletin varieties which have shown promise in recent years are described. In this discussion and the table accompanying it the varieties most worthy of distribution in the light of recent tests are given. This is the only place in the bulletin where it is felt that definite recommendations as to specific varieties should be made.

The earlier work forms an invaluable background for the later. The earlier results in a way are a check upon the behavior of types over a long period of time. If a selection of Hanna has yielded well for two or three years in an area where the Hanna group has been promising in only 6 of the last 20 years, these recent results are not so conclusive as if the Hanna group had been uniformly good.

USE OF CEREAL INVESTIGATIONS NUMBERS

In preparing the material for publication it was necessary to use a single number to designate the same variety at the various stations. It is probable that State officials would prefer that the identity of

the State number be preserved in this compilation. In practice this is not feasible, and it is especially impracticable with commercial varieties. At a given station most commercial varieties were obtained from other stations. If a single State number were to be given to a variety it should be that of the first State to number it. Often more than one State secured the variety from a common source. In many instances the variety was obtained by the State more than once either from the same or different sources. A serious study was made as to the best method of presenting the material. No other method seemed feasible, so a C. I. (Cereal Investigations) number was given to all varieties. Later in the bulletin under the C. I. number additional data are given which could not be repeated throughout the publication.

The part played by the various State stations in introducing varieties is emphasized in two ways. The table of C. I. numbers is intended to include all varieties listed in the tables, with a definite record of the origin of all pedigreed varieties. In the discussion of varieties all those of promise are treated individually and credited to the originators so far as the writers know them. If a variety is promising only at the original station it will be so shown in presenting the yields for that place. If it has a wider adaptation this is specifically considered elsewhere.

DISCUSSION OF STATION YIELDS

In discussing the varietal yields obtained in experiments at numerous stations covering most of the United States and Canada (fig. 9) it seems desirable to arrange them in three geographic groups, as

FIG. 9.—Map showing the location of agricultural experiment stations in the United States and Canada from which yields or observations are reported in this bulletin

follows: (1) Stations in Canada; (2) stations in the northern half of the United States, where spring varieties are grown from spring sowing in the north and fall sowing in the far West; and (3) stations in the southern half of the United States, where winter barleys are grown.

In each group the individual stations are taken in their order from east to west. This has the advantage of showing the areas of adaptation of different types or varietal groups of barley. For example, in the Canadian group of stations there becomes apparent (1) the area where the Chevalier barleys are adapted, (2) the area where the Manchuria and Hanna groups do better, and (3) the semiarid area, where still other varieties are best adapted. The name and location of each station included in each of these three groups follow.

Canadian stations

Charlottetown, Prince Edward Island.	Ottawa, Ontario.	St. Bernard Mission, Gr- ard, Alberta.
Nappan, Nova Scotia.	Brandon, Manitoba.	Beaver Lodge, Alberta.
Kentville, Nova Scotia.	Indian Head, Saskatche- wan.	Agassiz, British Colum- bia.
Fredericton, New Bruns- wick.	Rosthern, Saskatchewan.	Sidney, British Columbia.
Ste. Anne de la Pocatiere, Quebec.	Scott, Saskatchewan.	Invermere, British Co- lumbia.
Cap Rouge, Quebec.	Lethbridge, Alberta.	
	Lacombe, Alberta.	
	Fort Vermilion, Alberta.	

Northern United States stations

Orono, Me.	Edgeley, N. Dak.	Huntley, Mont.
Durham, N. H.	Langdon, N. Dak.	Bozeman, Mont.
Kingston, R. I.	Mandan, N. Dak.	Sheridan, Wyo.
Storrs, Conn.	Dickinson, N. Dak.	Archer, Wyo.
Ithaca, N. Y.	Williston, N. Dak.	Fort Collins, Colo.
State College, Pa.	Brookings, S. Dak.	Akron, Colo.
Wooster, Ohio.	Eureka, S. Dak.	Sandpoint, Idaho.
East Lansing, Mich.	Highmore, S. Dak.	Moscow, Idaho.
La Fayette, Ind.	Cottonwood, S. Dak.	Aberdeen, Idaho.
Urbana, Ill.	Newell, S. Dak.	Nephi, Utah.
De Kalb, Ill.	Lincoln, Nebr.	Pullman, Wash.
Madison, Wis.	North Platte, Nebr.	Lind, Wash.
St. Paul, Minn.	Mitchell, Nebr.	Moro, Oreg.
Waseca, Minn.	Manhattan, Kans.	Union, Oreg.
Duluth, Minn.	McPherson, Kans.	Burns, Oreg.
Grand Rapids, Minn.	Hays, Kans.	Newlands, Nev.
Crookston, Minn.	Colby, Kans.	Chico, Calif.
Morris, Minn.	Garden City, Kans.	Modesto, Calif.
Ames, Iowa.	Tribune, Kans.	Davis, Calif.
Columbia, Mo.	Havre, Mont.	
Fargo, N. Dak.	Moccasin, Mont.	

Southern United States stations

College Park, Md.	Ashburn, Ga.	Fayetteville, Ark.
Arlington Experiment Farm, Rosslyn, Va.	Quitman, Ga.	Stillwater, Okla.
Blacksburg, Va.	Morgantown, W. Va.	New Hope, Tex.
Clemson, S. C.	Lexington, Ky.	Amarillo, Tex.
Athens, Ga.	Knoxville, Tenn.	Channing, Tex.
	Jackson, Tenn.	State College, N. Mex.

CANADIAN STATIONS

As noted previously, these stations are taken in their order from east to west, beginning with Prince Edward Island and ending with those in British Columbia.

CHARLOTTETOWN, PRINCE EDWARD ISLAND

The climate of Prince Edward Island is analogous to that of England. The summers are mild, and the growing season of barley is long. As might be expected the dominant varieties are of the Chevalier type. This is due not only to their superior adaptation, but to the farming system on the island. Very few fields of pure barley are grown here. As in a large part of eastern Canada, barley and oats are grown together, and the resulting crop is used as feed without the separation of the two. The yield from these mixed fields is higher than from either barley or oats seeded alone. As the oat crop is later in maturing than barley, only the late-ripening varieties of barley are suited to mixing with oats at seeding time.

TABLE 1.—*Annual acre yields of varieties of barley grown at the experiment station for Prince Edward Island (at Charlottetown) in part or all of the six years from 1910 to 1915, inclusive*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)						Years grown	Average yield (bus.)	Percentage of weighted mean
		1910	1911	1912	1913	1914	1915			
Albert.....	1551	80.7	50.6	51.4	52.5	71.4	56.1	6	60.5	102.4
Claude.....	1557	79.5	51.9	—	—	—	—	2	65.7	121.4
Mensury.....	2557	79.2	37.4	—	—	—	—	2	58.3	107.8
Nugent.....	176	70.6	60.7	54.2	56.2	70.2	60.8	6	62.1	105.1
Mansfield.....	2241	66.0	35.7	—	—	—	—	2	50.9	94.1
Stella.....	2678	63.4	44.9	60.1	53.9	79.3	67.6	6	61.5	104.1
Odessa.....	182	61.6	45.7	65.3	51.8	65.3	63.7	6	58.9	99.7
O. A. C. 21.....	1470	60.1	53.5	66.7	53.1	70.9	64.5	6	61.5	104.1
Trooper.....	173	58.9	37.9	63.9	46.9	81.2	66.6	6	59.2	100.2
Yale.....	1592	55.2	39.7	—	—	—	—	2	47.5	87.8
Oderbrucker.....	2700	48.7	41.6	68.0	51.4	69.3	63.0	6	57.0	96.4
Invincible.....	590	53.3	45.3	53.0	58.7	65.4	61.0	6	61.1	103.4
Chevalier II.....	200	80.6	50.1	66.7	57.7	74.1	71.0	6	66.7	112.9
Hannchen.....	531	77.3	38.9	62.3	65.0	78.3	—	5	64.4	110.1
Standwell.....	584	59.6	45.7	57.4	49.8	64.5	54.9	6	55.3	93.6
Clifford.....	1910	59.4	41.7	49.0	48.7	65.2	—	5	52.8	90.3
French Chevalier.....	175	58.3	44.4	—	—	—	—	2	51.3	94.8
Beaver.....	1915	53.7	41.0	46.3	48.4	52.9	40.8	6	47.2	79.9
Canadian Thorpe.....	740	51.6	43.3	61.6	56.4	70.2	57.3	6	56.7	95.9
Jarvis.....	2672	50.7	42.1	—	—	—	—	2	46.4	85.8
Danish Chevalier.....	180	49.1	43.9	—	—	—	—	2	46.5	86.0
Manchurian.....	739	—	43.4	67.0	54.5	69.3	62.6	5	59.4	102.6
English Chevalier.....	2733	—	—	67.9	68.6	83.3	—	3	73.3	117.7
Gold.....	1145	—	—	—	63.2	82.3	73.5	3	73.0	115.9
Early Chevalier.....	2725	—	—	—	—	65.0	59.3	2	62.1	92.8
Charlottetown 80.....	2732	—	—	—	—	—	75.3	1	75.3	120.7

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF SIX REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages					
			Chevalier II	Invincible	Hannchen	Manchurian	Odessa	Stella
Chevalier II.....	200	Years comparable.....	—	6	5	5	6	6
		Percentage yield.....	—	109.1	102.3	107.7	113.2	108.4
Invincible.....	590	Years comparable.....	6	—	5	5	6	6
		Percentage yield.....	91.6	—	95.0	95.5	103.8	99.3
Hannchen.....	531	Years comparable.....	5	5	—	4	5	5
		Percentage yield.....	97.8	105.3	—	104.4	111.1	106.7
Manchurian.....	739	Years comparable.....	5	5	4	—	5	5
		Percentage yield.....	92.9	104.7	95.8	—	101.7	97.1
Odessa.....	182	Years comparable.....	6	6	5	5	—	6
		Percentage yield.....	88.3	96.4	90.0	98.3	—	95.7
Stella.....	2678	Years comparable.....	6	6	5	5	6	—
		Percentage yield.....	92.3	100.7	93.7	103.0	104.5	—

In Table 1 are given the yields of the varieties grown during all or part of the years 1910 to 1915, inclusive. Of those grown for the entire period of six years Chevalier II (C. I. No. 200) gave a yield of almost 113 per cent of the weighted mean. In the first group digest, the detailed statement of which does not appear here, the yields of the varieties falling in the Chevalier, Thorpe, Hanna, Manchuria, and hybrid groups were summarized. The Chevalier group gave the highest yield, Hanna the second highest, Manchuria third, and Thorpe fourth. Although nine hybrid varieties were tested the group yields of the hybrids were inferior to those of the other groups. In the summarized digest of Table 1 six important varieties are compared. It is obvious that the group yields are not always indicative of the relative value of types. The general average yield of a group with many varieties may be lower than that of a group which contains only a few, when in reality the best-yielding individual varieties are in the large group. This happens, of course, when all of those of the small group are much better than the average, while only a few of the varieties of the larger group are superior. The table shows that Chevalier II was the best variety during the six years compared. Hannchen (C. I. No. 531) was almost as good as Chevalier II, and Manchurian (C. I. No. 739) was fourth. Stella (C. I. No. 2678), one of the better hybrids produced by the Central Experimental Farm at Ottawa, was third among the six varieties compared.

NAPPAN, NOVA SCOTIA

At Nappan, Nova Scotia, 75 varieties of barley were tested during one or more of the 26 years from 1890 to 1915, inclusive. It will be noticed in the statement of percentages in Table 2 that varieties belonging to the Chevalier and Manchuria groups are almost uniformly good. Several varieties now well known were not introduced until late in the experiments, and it may be that the number of years they were tested was insufficient to give a proper estimate of their value. The yield obtained from Gold (C. I. No. 1145) was very good in the two years tested. Manchurian (C. I. No. 739), Hannchen (C. I. No. 531), and O. A. C. 21 (C. I. No. 1470) were not outstanding at Nappan. These have proved to be promising sorts at many other stations, and it is rather surprising that they do not yield better at Nappan. Others of the same groups did well at that place. In the group digest made from the yields of barley varieties at Nappan the Manchuria group was decidedly better than the others compared. The Chevalier was second, the Hybrid third, and the Hanna last in point of yield.

These group relationships show that although Nappan is on the mainland, only a few miles from Charlottetown, the section is not so favorable as Prince Edward Island for the growing of Chevalier barleys. This point is also brought out in the summarized digest in which 10 promising varieties of various groups are compared. Of these Mensury (C. I. No. 2657) is the best, French Chevalier (C. I. No. 175) is almost equal to it, and Stella (C. I. No. 2678), Odessa (C. I. No. 182), and Canadian Thorpe (C. I. No. 740) follow in the order named. The poorest of the 10 was Blue Long Head (C. I. No. 2685). This is of interest, as this variety belongs to the Coast group and is best adapted to arid and semiarid climates where the ripening season occurs during an almost rainless period. Success (C. I. No. 2707) was inferior to all others except Blue Long Head (C. I. No. 2685), Petschora (C. I. No. 2658), and Nepal (C. I. No. 595). Petschora is an early form received from a section of Russia much farther north than Nappan.

TABLE 2.—Annual acre yields of varieties of barley grown at the experimental farm for Nova Scotia (at Nappan) in part or all of the 23 years from 1890 to 1915, inclusive

[Data compiled from the Canadian Experimental Farms Reports⁴⁵]

Variety	C. I. No.	Acre yields (bushels)																							Years grown	Average yield (bus.)	Percentage of weight-mean		
		1890	1891	1892	1893	1894	1895	1896	1897	1898	1899	1900	1901	1902	1903	1904	1905	1906	1907	1908	1909	1910	1911	1912				1913	1914
Petschora.....	2658	35.0	—	—	—	—	31.7	37.9	17.9	45.4	20.2	45.0	52.5	46.7	42.5	—	—	—	—	—	—	—	—	—	—	—	—	38.4	96.0
Baku.....	333	14.5	41.1	135.0	15.8	19.0	638.7	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	25.8	71.9	
English Maltling.....	2654	21.5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	21.5	67.6	
Early Maltling.....	2661	19.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	19.0	76.7	
Peerless White.....	2647	32.0	47.5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	30.7	96.6	
Selected Chevalier.....	2662	44.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	44.0	155.5	
Goldthorpe.....	327	29.7	63.3	64.5	26.7	25.8	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	38.2	108.5	
Baxter.....	185	20.4	46.8	45.0	37.1	24.3	23.6	33.7	4.1	7.5	5.8	47.5	59.2	53.3	35.0	28.0	2.2	—	—	—	—	—	—	—	—	—	42.5	103.7	
Rennio Improved.....	181	40.0	57.5	50.0	19.0	26.3	32.2	9.1	9.4	8.3	4.4	2.5	5.6	7.5	1.7	45.8	45.0	34.1	6.6	—	—	—	—	—	—	39.9	97.8		
Odessa.....	182	40.0	56.0	80.0	20.0	31.3	32.1	12.2	9.6	3.0	8.4	5.8	5.1	7.6	0.0	8.3	3.1	3.4	5.0	35.5	8.4	3.8	3.5	2.1	3.9	2.4	44.2	106.8	
Rhegman Hills.....	2698	15.5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	20	97.8	
Prize Prolific.....	169	25.0	50.0	50.0	27.1	24.0	638.3	33.3	33.1	2.7	5.2	5.4	5.6	7.4	2.5	38.3	—	—	—	—	—	—	—	—	—	—	15.5	52.0	
Sharpe Chevalier.....	2650	31.0	44.6	—	26.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	33.6	94.8	
Saale.....	2645	51.7	40.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	45.9	96.6	
Golden Melon.....	411	52.1	55.0	32.2	34.2	32.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	41.1	112.9	
Danish Chevalier.....	180	44.2	45.0	—	26.7	42.5	26.7	30.3	33.3	3.0	2.3	33.8	3.6	8.6	5.0	37.5	42.9	51.7	5.6	7.5	4.0	2.4	4.5	8.8	—	—	40.8	112.9	
Thane.....	2652	45.4	45.0	22.1	20.4	43.5	4.3	2.9	2.4	2.1	7.5	0.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	36.1	83.3	
Kinver Chevalier.....	587	48.3	32.5	—	36.7	37.1	26.3	32.5	8.4	4.0	3.3	3.4	2.2	—	—	—	—	—	—	—	—	—	—	—	—	—	33.8	108.5	
Duckbill.....	1916	71.1	45.0	31.7	27.0	30.0	28.1	7.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	33.5	108.5	
Oderbrucker.....	2700	71.5	55.0	25.0	30.0	43.8	37.1	5.0	4.5	8.4	3.2	5.6	1.7	5.2	5.6	7.4	5.8	3.4	2.2	9.3	7.4	4.0	8.6	3.5	6.7	4.9	45.8	110.3	
Mensury.....	2657	65.6	50.0	19.0	63.4	8.4	6.6	1.1	7.5	1.3	7.5	5.5	8.0	0.0	5.6	7.5	0.0	4.9	2.8	3.3	4.1	7.2	5.9	5.3	2.3	1.4	43.7	118.4	
Golden Grain.....	588	24.0	60.0	18.6	30.0	44.4	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5	35.6	97.0
Himalaya.....	1984	19.8	45.0	24.4	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3	29.7	73.5
Canadian Thorpe.....	740	20.4	20.0	24.6	7.1	7.0	540.8	27.5	50.0	3.8	3.4	2.5	5.4	2.0	3.9	2.2	5.4	2.2	5.4	2.5	5.4	2.5	4.7	5.4	0.3	5.1	3	39.5	95.6
French Chevalier.....	175	32.3	18.3	33.6	74.7	9.1	33.5	8.0	2.6	5.5	0.5	2.9	2.8	3.5	8.4	6.4	7.5	3.7	8.5	0.3	2.9	2.6	2.6	5.3	8.5	4.8	23	113.0	
Common.....	184	40.0	16.2	14.2	12.5	8.2	5.3	3.7	3.4	2.3	3.4	2.6	3.3	3.4	2.6	3.3	3.4	2.6	3.3	3.4	2.6	3.3	3.4	2.6	3.3	3.4	14	108.9	
Newton.....	179	20.0	20.0	43.5	4.2	1.0	8.3	3.7	4.7	5.4	5.0	0.6	4.2	2.6	7.3	8.7	—	—	—	—	—	—	—	—	—	—	13	38.6	95.1
Royal.....	1252	37.4	45.4	49.4	29.4	33.3	33.6	7.6	7.1	7.5	2.9	6.8	3.9	2.3	2.3	3.8	—	—	—	—	—	—	—	—	—	—	13	44.1	107.8
Phoenix.....	2676	32.1	41.0	26.3	34.0	4.3	23.5	8.2	5.4	2.9	2.2	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	9	38.3	94.8
Summit.....	174	28.3	24.2	1.8	7.3	7.4	7.0	0.4	8.3	3.5	2.4	2.4	2.3	7.3	6.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	14	30.2	96.1
Trooper.....	173	26.3	34.3	37.4	9.0	0.4	43.8	3.7	5.4	1.7	5.4	6.5	0.0	3.4	6.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	14	30.2	96.1
Surprise.....	171	25.4	41.1	7.6	7.4	0.0	2.9	2.2	5.4	2.4	0.0	0.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	9	42.6	105.4
Beaver.....	1015	27.2	52.3	33.4	6.0	4.0	8.5	5.5	6.6	7.3	3.0	8.2	5.4	3.3	3.0	8.2	5.4	3.3	3.0	8.2	5.4	3.3	3.0	8.2	5.4	3.3	29	103.8	105.4
Sidney.....	178	40.2	27.6	3.8	3.2	8.3	3.2	5.8	3.4	8.3	3.2	5.8	3.4	8.3	3.2	5.8	3.4	8.3	3.2	5.8	3.4	8.3	3.2	5.8	3.4	8.3	15	36.3	88.8
Stella.....	2678	37.3	33.4	6.2	1.3	7.4	7.5	0.7	5.8	3.5	8.3	3.5	8.4	5.0	3.4	2.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3	22	107.2	105.4
Bolton.....	176	35.4	33.4	2.7	9.3	5.3	0.4	2.0	0.4	5.3	7.5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	9	38.5	95.3
Nugent.....	177	33.3	33.1	7.7	9.3	0.3	6.3	2.4	5.0	4.0	4.7	5.5	2.6	2.4	1.7	7.3	7.2	9.3	5.0	4.5	8.4	6.7	5.4	2.5	0.3	3.4	22	105.1	105.1

[illegible]

TABLE 2.—*Annual acre yields of varieties of barley grown at the experimental farm for Nova Scotia (at Nappan) in part or all of the 26 years from 1890 to 1915, inclusive—Continued*

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages									
			French Chevalier	Canadian Thorpe	Hannchen	Mensury	Petschora	Odessa	Blue Long Head	Nepal	Stella	Success
French Chevalier.	175.	Years comparable	-----	23	4	19	9	24	9	3	22	8
		Percentage yield	-----	120.3	124.2	94.4	115.4	107.1	132.8	131.8	109.0	111.2
Canadian Thorpe.	740.	Years comparable	23	-----	4	18	9	23	9	3	22	8
		Percentage yield	83.1	-----	99.9	80.8	104.9	91.8	111.6	115.8	90.1	102.3
Hannchen.	531.	Years comparable	4	4	-----	1	(*)	4	(*)	(*)	4	(*)
		Percentage yield	80.5	100.1	-----	50.0	-----	87.4	-----	-----	83.7	-----
Mensury.	2657.	Years comparable	19	18	1	-----	9	20	9	3	17	8
		Percentage yield	105.9	123.7	200.0	-----	129.9	108.0	130.0	129.1	109.0	127.1
Petschora.	2658.	Years comparable	9	9	(*)	9	-----	10	6	3	9	8
		Percentage yield	86.6	95.4	-----	77.0	-----	86.6	118.8	106.6	89.7	96.4
Odessa.	182.	Years comparable	24	23	4	20	10	-----	9	3	22	8
		Percentage yield	93.3	109.0	114.5	92.6	115.5	-----	124.2	125.1	98.4	112.8
Blue Long Head.	2685.	Years comparable	9	9	(*)	9	6	8	5	-----	3	9
		Percentage yield	75.3	89.6	-----	76.9	84.2	80.5	-----	94.6	86.3	92.1
Nepal.	595.	Years comparable	3	3	(*)	3	3	3	3	-----	3	3
		Percentage yield	75.9	86.4	-----	77.5	93.8	79.9	105.7	-----	88.9	93.8
Stella.	2678.	Years comparable	22	22	4	17	9	22	9	3	-----	-----
		Percentage yield	91.7	111.0	119.4	91.7	111.4	101.6	115.9	112.5	-----	106.7
Success.	2707.	Years comparable	8	8	(*)	8	8	8	6	3	8	-----
		Percentage yield	89.9	97.8	-----	78.7	103.8	88.6	108.6	106.6	93.7	-----

KENTVILLE, NOVA SCOTIA

At Kentville, Nova Scotia, only two varieties were reported tested before 1915. During the years 1914 and 1915, as shown in Table 3, Manchurian (C. I. No. 739) proved to be slightly better than Canadian Thorpe (C. I. No. 740). A test of only two years is of little significance; but here, as at Nappan, the Manchurian was superior to Canadian Thorpe.

TABLE 3.—*Annual acre yields of varieties of barley grown at the experimental station for Annapolis and Cornwallis Valleys (at Kentville, Nova Scotia) in 1914 and 1915*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)		Years grown	Average yield (bus.)	Percentage of weighted mean
		1914	1915			
Manchurian.	739	24.3	22.4	2	23.1	101.8
Canadian Thorpe.	740	22.2	21.7	2	21.9	96.5

FREDERICTON, NEW BRUNSWICK

Five barleys were grown at Fredericton, New Brunswick, in 1915. The yields are reported in Table 4. Four of the varieties were introduced into experiment-station tests in recent years. The best of the five was Gold (C. I. No. 1145). Second in yield came O. A. C. 21 (C. I. No. 1470), while Early Chevalier (C. I. No. 2725), Manchurian (C. I. No. 739), and Canadian Thorpe (C. I. No. 740), were third, fourth, and fifth, respectively.

TABLE 4.—*Annual acre yields of varieties of barley grown at the experimental station for New Brunswick (at Fredericton) in the year 1915*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yield (bus.)	Percent- age of weighted mean
Early Chevalier.....	2725	41.0	103.0
Gold.....	1145	46.0	115.6
O. A. C. 21.....	1470	43.0	108.0
Manchurian.....	739	40.0	100.5
Canadian Thorpe.....	740	29.0	72.9

STE. ANNE DE LA POCATIERE, QUEBEC

To 1921 the only yields published from Ste. Anne de la Pocatiere were those of Manchurian (C. I. No. 739) and Success (C. I. No. 2707) grown in 1915. The yield of Manchurian was 37 bushels and that of Success only 31 bushels.

TABLE 5.—*Annual acre yields of varieties of barley grown at the experimental station for central Quebec (at Cap Rouge) in one or more of the five years from 1911 to 1915, inclusive.*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percent- age of weighted mean
		1911	1912	1914	1915			
Escourgeon.....	2716	36.3	18.7	-----	-----	2	27.5	106.3
Odessa.....	182	29.4	25.0	-----	-----	2	27.2	107.1
Success.....	2707	27.2	22.5	14.2	-----	3	21.3	88.7
Manchurian.....	739	26.9	21.3	13.7	40.0	4	25.5	95.9
Stella.....	2678	24.4	25.0	-----	-----	2	24.7	97.2
Black Japan.....	2717	20.6	12.5	-----	-----	2	16.5	65.0
O. A. C. 21.....	1470	13.7	18.7	12.3	43.0	4	21.9	82.3
Chevalier II.....	200	41.3	-----	-----	-----	1	41.3	137.2
Early Chevalier.....	2725	40.0	28.7	-----	41.0	3	36.6	130.7
Hannchen.....	531	35.0	12.5	-----	-----	2	23.7	93.3
Beaver.....	1915	33.7	-----	-----	-----	1	33.7	112.0
Svanhals.....	187	33.1	8.7	-----	-----	2	20.9	82.3
Duckbill.....	1916	29.7	-----	-----	-----	1	29.7	98.7
Gold.....	1145	-----	-----	-----	46.0	1	46.0	115.6
Canadian Thorpe.....	740	-----	-----	-----	29.0	1	29.0	72.9

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF EIGHT REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages							
			Early Chevalier	Duckbill	Svanhals	Hannchen	Manchurian	Odessa	Stella	Success
Early Chevalier.....	2725	Years comparable.....	-----	1	2	2	3	2	2	2
		Percentage yield.....	-----	134.7	164.4	144.6	124.4	126.3	139.1	138.2
Duckbill.....	1916	Years comparable.....	-----	1	1	1	1	1	1	1
		Percentage yield.....	74.3	-----	89.7	84.9	110.4	101.0	121.7	109.2
Svanhals.....	187	Years comparable.....	2	1	-----	2	2	2	2	2
		Percentage yield.....	60.8	111.4	-----	88.0	86.7	76.8	84.6	84.1
Hannchen.....	531	Years comparable.....	2	1	2	-----	2	2	2	2
		Percentage yield.....	69.1	117.8	113.6	-----	98.5	87.3	96.2	95.6
Manchurian.....	739	Years comparable.....	3	1	2	2	-----	2	2	3
		Percentage yield.....	80.4	90.6	115.3	101.5	-----	88.6	97.6	96.9
Odessa.....	182	Years comparable.....	2	1	2	2	2	-----	2	2
		Percentage yield.....	79.2	99.0	130.1	114.5	112.9	-----	110.1	109.5
Stella.....	2678	Years comparable.....	2	1	2	2	2	2	-----	2
		Percentage yield.....	71.9	82.2	118.2	104.0	102.5	90.8	-----	99.4
Success.....	2707	Years comparable.....	2	1	2	3	2	2	2	-----
		Percentage yield.....	72.3	91.6	118.9	104.6	103.2	91.4	100.6	-----

CAP ROUGE, QUEBEC

During one or more of the five years from 1911 to 1915, inclusive, 15 varieties of barley were tested at Cap Rouge, Quebec. Only two of these varieties were grown for as many as four of the five years. The results are reported in Table 5. Early Chevalier (C. I. No. 2725) was undoubtedly the best variety for the three years in which it was tested. The indications are that the Chevalier barleys are well adapted to this section of Quebec. The summarized digest of Table 5 shows that Duckbill (C. I. No. 1916) and Odessa (C. I. No. 182) ranked high among the varieties. These two have been widely tested at Canadian stations and have given good yields at many places. Gold (C. I. No. 1145) grown only in 1915 produced the highest yield in that year.

OTTAWA, ONTARIO

The Central Experimental Farm of Canada is located at Ottawa, Ontario. The other Dominion experimental farms are under the direct supervision of this institution. Most of the barleys tested at these farms were received from the farm at Ottawa. The many hybrids tested at the various stations were produced by Dr. C. E. Saunders, of that farm.

The work at Ottawa began in 1889. Table 6 shows that 15 varieties were tested in that year. Barley was grown in all but one of the years from 1889 to 1915, inclusive. In all, nearly 150 varieties were tested. Although this large number makes inspection of the data difficult, it is obvious that barleys of the Manchuria and Odessa types are the highest yielding sorts at Ottawa. The Chevalier barleys are not particularly good, although Early Chevalier (C. I. No. 2725) has shown promise since its introduction in 1907. From the column of percentages it is evident that a number of hybrid sorts have given good yields. Among the best of these are Stella (C. I. No. 2678), Trooper (C. I. No. 173), Nugent (C. I. No. 176), and Albert (C. I. No. 1551). Among the widely tested varieties which do not yield well at Ottawa are Chevalier II (C. I. No. 200), Invincible (C. I. No. 590), and Standwell (C. I. No. 584).

Nine groups of barleys were compared. The number in the groups varied from only 2 in Coast and Nudum to 40 in the Hybrid. The groups ranked as follows: Manchuria, Hybrid, Hanna, Nudum, Coast, Thorpe, Polar, Chevalier, and Himalaya. The Manchuria was much better than any of the other groups compared. The high yields of the hybrids may have been due to the fact that they were developed at Ottawa and selected for yield at that place. They probably have done relatively better at this place than at the other stations. The Hanna barleys produced almost as much as the hybrids. The Nudum group was represented by only two varieties and therefore probably stands too high as a group. The high yield of the Nudum barleys is due to the unusual productivity of Caucasian (C. I. No. 2724). This is a 2-rowed naked sort, similar to the Baku grown at a number of western experiment stations in the United States. Its yield is especially interesting, inasmuch as the naked oats have yielded exceptionally well at Ottawa.

Ten well-known varieties are compared in the summarized digest of Table 6. Odessa (C. I. No. 182) is the best of the 10 compared in this table. Mensury (C. I. No. 2657), Hannchen (C. I. No. 531), Stella (C. I. No. 2678), and Svanhals (C. I. No. 187) follow in order. Blue Long Head (C. I. No. 2685) was relatively better here than at Nappan. The climatic conditions at Ottawa are not only more favorable to the growth of Manchuria varieties and less favorable to the Chevalier barleys than is the case in eastern Canada but also are such that barleys suited to the arid regions grow quite well. Neither the Chevalier nor the Thorpe groups were at their best, although the former were better than the latter. It is unfortunate that only a limited number of varieties can be compared in this way, as it resulted in eliminating several very promising sorts from the comparison. Manchurian (C. I. No. 739), Gold (C. I. No. 1145), and O. A. C. 21 (C. I. No. 1470) have been particularly promising since their introduction in the more recent years.

TABLE 6.—Annual acre yields of varieties of barley grown at the Central Experimental Farm, Ottawa, Ontario, in one or more of the 27 years from 1889 to 1915, inclusive—Continued

Variety	C. I. No.	Acre yields (bushels)																	Years grown	Average yield (bus.)	Percentage of weighted mean										
		1889	1890	1891	1892	1893	1894	1895	1896	1897	1898	1899	1900	1901	1902	1903	1904	1905				1906	1907	1908	1909	1910	1911	1912	1913	1914	1915
Laboul.	2690			25.1																								25.1	64.0		
Moulton.	2697			26.8																								26.8	68.4		
Mardan.	2696			30.5																								30.5	77.7		
Oderbrucker.	2700			51.7	27.1	25.8	31.0	47.3	48.1	46.9	49.2	52.2	52.6	52.6	52.5	42.9	36.7	68.3	42.5	42.5	48.7	60.5	57.1	53.1	30.0	18.7	46.3	44.5	104.7		
Palampur.	2691			38.9																								38.9	99.2		
Salakot.	2695			34.5	19.2																							26.9	74.3		
Simla.	2692			34.5																								34.5	88.0		
Soraj.	2693			34.5																								34.5	88.0		
French Chevalier.	175			33.3	32.7	32.6	53.4	45.5	52.4	63.6	9.47	5.56	7.55	2.08	3.44	6.44	2.60	0.57	9.27	5.28	7.48	1.45	0.48	1.1	60.0	55.0	44.3	103.7	87.0		
Duckbill.	1916			40.4	18.3	25.8	37.5	53.9	6.1					16.3	56.7													33.5	101.2		
Canadian Thorpe.	740			31.3	15.8	23.3	25.8	49.6	38.5	47.3	46.7	58.3	45.2	62.5	46.7	44.2	42.5	56.3	48.1	136.3	45.0						42.4	91.6	10.2		
Norway House.	2699			28.3																								28.3	91.2	91.2	
Victor.	179			30.2	23.1	33.5	46.7	73.4	63.9	244.6	49.2	221.9	48.3														28.3	36.2	107.2		
Newton.	172			19.2	27.1	29.4	51.0	41.2	36.7	30.8	50.0		50.0														19.2	46.0	96.9		
Summit.	174			35.0																								35.0	40.1	75.7	
Surprise.	171			28.3																								28.3	41.8	95.7	
Bolton.	177			30.8	35.5	51.0	31.5	54.3	1.38	3.52	5.25	2.53															30.8	40.1	96.9		
California Prolific.	2649			27.1	126.0	39.6																						27.1	75.7	30.9	
Sidney.	178			21.9	34.6	33.5	93.3	7.50	0.45	0.31	5.46	7.39	2.30	4.42	5.52	9.26	9.34										21.9	38.7	90.8		
Royal.	1252			36.0	51.3	36.2	1.48	1.52	7.50	0.58	2.83	5.47	5.41	3.34	2.54	6.41	7.1										36.0	47.0	109.0		
Vanguard.	2665			34.4	46.7	44.4	3.86	7.47	152.9	92.7	7.50																	34.4	103.4	96.4	
Phoenix.	2676			33.5	37.3	35.8	7.41	2.45	0.40	4.1	7.25	2.86															33.5	39.9	96.4		
Pioneer.	2677			33.5	54.7	56.7	7.50	8.57	1.50	8.60	0.77	7.00															33.5	48.8	117.9		
Stella.	2678			31.0	46.0	55.0	0.45	5.47	1.46	7.43	7.86	9.65	8.51	3.53	3.55	4.47	1.50	6.56	7.35	0.08	7.45	6.44	4.26	9.51	3.1	21.4	48.0	110.9	48.0		
Trooper.	173			30.8	46.9	59.0	0.48	9.88	7.49	2.47	1.25	2.65	8.52	1.27	1.69	2.69	8.31	9.49	4.40	6.59	4.1					30.8	47.7	110.4	48.0		
Nugent.	176			22.7	42.9	50.2	24.3	1.41	5.50	0.50	0.33	1.00	8.42	1.50	0.72	5.65	0.46	9.46	3.58	7.62	5.1	9.26	9.1			22.7	48.3	112.1	48.3		
Nepean.	183			37.4	45.6	63.1	6.37	7.45	4.49	2.41	0.46																37.4	41.9	99.3	41.9	
Pacer.	2667			36.3	48.3	132.9	33.7	1.41	9.43	3.26	9.48	3.1															36.3	38.9	92.2	38.9	
Beaver.	1915			35.0	44.4	23.4	2.55	4.49	2.54	2.46	0.52	5.42	5.31	4.7	5.57	5.38	1.36	9.50									35.0	45.0	104.2	45.0	
Prolific.	2738			30.8	40.8																							30.8	35.8	80.8	35.8
Suffolk Chevalier.	2701			30.2	44.8																							30.2	37.5	84.7	37.5
Success.	2707			51.3	34.0	14.4	3.37	5.41	7.50	8.16	3.50	8.1															51.3	42.1	99.8	42.1	
Monck.	2703			28.7	36.9	18.5	2.7	7.45	4.35	8.14	4.6	7.1															28.7	31.8	75.4	31.8	
Suffolk Chevalier.	2702			28.7	44.2																							28.7	36.5	82.4	36.5
Logan.	2668			47.9	98.8	6.35	4.45	4.3	3.39	4.4	2.37	1.20	5.0														47.9	40.3	94.6	40.3	
Leslie.	2669			46.6	65.5	6.45	8.45	8.0	8.11	0.39	2.2																46.6	37.8	88.9	37.8	
Douglas.	2704			46.3																								46.3	46.3	95.5	46.3
Dunham.	2670			46.3	237.4	46.0	45.0	49.2	30.6	53.3	37.9	15.0	40.8	33.7	34.4												46.3	40.8	93.4	40.8	
Kirby.	2671			45.6	63.8	4.31	2.47	1.1																			45.6	40.6	94.6	40.6	

[illegible]

TABLE 6.—Annual acre yields of varieties of barley grown at the Central Experimental Farm, Ottawa, Ontario, in one or more of the 27 years from 1889 to 1915, inclusive—Continued

Variety	C. I. No.	Acre yields (bushels)																	Years grown	Aver- age yield (bus.)	Per- centage of weight- ed mean											
		1889	1890	1891	1892	1893	1894	1895	1896	1897	1898	1899	1900	1901	1902	1903	1904	1905				1906	1907	1908	1909	1910	1911	1912	1913	1915		
Taganrog	2721																		69.2										6	58.4	117.3	
Manchurian	739																		63.3	48.1	53.7	62.5	68.1	53.1	40.6	42.5			8	54.0	119.5	
Small Blue Naked	2722																		34.6	33.1	42.5	84.4	58.1						5	40.5	86.7	
Erft White	1310																		66.7	36.9									2	51.8	107.0	
Old Irish	2723																		59.6	50.6	638.1	134.4							4	45.7	101.3	
Jewel	324																		58.7	42.5	25.6	33.7	37.5						5	39.6	84.8	
Hofbrau	338																		48.7	52.5	29.4	26.9	35.0						5	38.5	82.4	
Gambinus	330																		44.2	21.9									2	33.1	68.4	
Caucasian	2724																		50.0	50.6	52.5	46.3	49.4						2	45.3	102.3	
Black	2726																		39.4	51.9	45.6	31.3	62.5	45.6					6	46.1	101.5	
Early Chevalier	2725																		49.4	35.6	54.4	53.7	55.0						7	50.0	112.9	
Leuder	2727																		35.0	51.3	38.1								3	41.5	85.6	
O. A. C. 21	1470																		70.6	52.5	26.9	46.3	65.6						5	52.4	111.5	
Early Indian	1576																		43.7	33.1	14.4	14.4	40.6						5	33.4	71.1	
Kutais	2728																		60.6	58.1									4	50.9	101.9	
Reed Triumph	889																		26.3	45.0	63.1	58.7							4	48.3	108.5	
Gold	1145																		39.4											3	56.5	123.6
Himalaya	1934																		30.0	39.4	46.9								3	38.8	89.4	
Swedish Star	1701																		55.6											3	55.6	105.9
Blinder	1909																													1	40.6	77.3

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages									
			French Chevalier	Canadian Thorpe	Svanhals	Hann- chen	Mensury	Odessa	Blue Long Head	Nepal	Stella	Success
French Chevalier	175	Years comparable	-----	18	10	9	19	17	13	4	20	8
		Percentage yield	-----	100.5	96.9	92.0	82.9	81.8	96.2	164.4	94.9	109.7
Canadian Thorpe	740	Years comparable	18	-----	6	6	18	17	13	4	17	8
		Percentage yield	99.5	-----	94.8	86.8	84.5	81.5	106.1	153.6	92.8	111.0
Svanhals	187	Years comparable	10	6	-----	9	7	5	5	(*)	18	(*)
		Percentage yield	103.2	105.5	-----	92.7	88.3	85.5	111.0	(*)	98.8	(*)
Hannchen	531	Years comparable	9	6	9	-----	7	5	5	(*)	-----	(*)
		Percentage yield	108.7	115.3	107.9	105.7	94.6	86.8	128.8	-----	102.6	-----
Mensury	2657	Years comparable	19	18	7	7	-----	19	13	4	17	8
		Percentage yield	120.7	118.3	113.2	115.2	103.0	97.1	107.9	149.9	106.6	128.0
Odessa	182	Years comparable	17	17	5	5	19	-----	12	4	15	8
		Percentage yield	122.2	122.6	122.6	115.2	103.0	103.0	115.8	150.8	110.3	130.1
Blue Long Head	2685	Years comparable	13	13	5	5	13	12	-----	4	13	7
		Percentage yield	104.0	99.9	90.1	82.8	92.7	86.4	-----	141.3	94.1	117.6
Nepal	595	Years comparable	4	-----	(*)	(*)	4	-----	4	-----	4	4
		Percentage yield	60.8	65.1	-----	-----	66.7	66.3	70.8	-----	71.7	86.8
Stella	2678	Years comparable	20	16	10	9	17	15	13	4	-----	8
		Percentage yield	105.4	107.8	101.2	97.5	93.8	90.7	106.2	139.4	-----	114.9
Success	2707	Years comparable	8	8	(*)	(*)	8	8	7	-----	8	-----
		Percentage yield	91.1	90.1	-----	-----	78.1	76.9	83.0	115.2	87.1	-----

BRANDON, MANITOBA

One of the first of the Canadian experimental farms to be established was that at Brandon, Manitoba. The testing of barley varieties commenced in 1889 at the Ottawa station. Brandon is much farther north as well as farther west. The summers are cooler, and there is less humidity at ripening time. Table 7 shows that Brandon is undoubtedly in the territory of the Manchuria group. In the percentage valuation of the varieties Mensury (C. I. No. 2657) and Odessa (C. I. No. 182), which were grown for almost the entire period of years, ranked very high. From the average yields shown it is evident that barley is highly productive in Manitoba. Mensury (C. I. No. 2657), for example, produced an average yield of 56.3 bushels for a period of 24 years. Several of the hybrid varieties gave good yields at Brandon. Among the best were Yale (C. I. No. 1592), Mansfield (C. I. No. 2241), Argyle (C. I. No. 2681), and Nugent (C. I. No. 176). Of the varieties which were introduced so late that they were tested in only a few years Garton 68 (C. I. No. 2740), Hannchen (C. I. No. 531), Manchurian (C. I. No. 739), and Gold (C. I. No. 1145), were among the best. The Chevalier barleys gave only average yields at Brandon. Danish Chevalier (C. I. No. 180) yielded nearly 100 per cent, French Chevalier (C. I. No. 175) 102 per cent, and Chevalier II (C. I. No. 200) 99 per cent of the weighted mean. The Thorpe varieties were more irregular than the Chevalier sorts, but on the whole produced slightly better yields.

Seven groups of barleys were compared, which ranked as follows: Manchuria, Thorpe, Hanna, Polar, Himalaya, Hybrid, and Chevalier. The Manchuria is outstanding as compared with the other groups. It is significant that a group of Polar barleys which are 6-rowed sorts adapted to regions north of the Manchuria barleys yielded well at Brandon.

Ten varieties of different types are compared in the summarized digest of Table 7. Mensury (C. I. No. 2657) was the best of these. Odessa (C. I. No. 182) and Hannchen (C. I. No. 531) were almost as good. Hannchen was tested for only four years, being introduced in 1910. The comparative yield of Blue Long Head (C. I. No. 2685) was low, as is usually the case where Manchuria yields well. Likewise, Nepal and Manchuria seldom do well at the same place.

TABLE 7.—Annual acre yields of varieties of barley grown at the experimental farm for Manitoba (at Brandon) in one or more of the 27 years from 1889 to 1915, inclusive

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)																				Years grown	Average yield (bus.)	Per- centage of weight- ed mean					
		1889	1890	1891	1892	1894	1895	1896	1897	1898	1899	1900	1901	1902	1903	1904	1905	1906	1907	1908	1909				1910	1911	1912	1913	1914
Danish Chevalier	180	13.8	51.7	08.3	51.3	23.1	141.5	41.3	30.4	37.1	150.6	28.5	19.2	44.0	29.0	53.7	70.0	85.1	97.0	55.0	83.0	46.1	5	—	—	—	—	44.3	99.8
Prentice	917	12.3	46.8	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	29.5	104.7
Duckbill	1916	10.8	48.3	37.5	54.6	29.0	57.1	—	22.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	33.8	86.7
Peerless White	2047	8.7	49.8	—	42.9	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	40.3	94.6
Beardless	2046	8.7	48.4	48.5	45.2	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	39.9	90.5
Thaet	2052	9.4	48.2	—	49.6	29.2	44.0	85.6	42.1	156.7	74.4	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	32.3	82.8
English Maltling	2054	7.8	40.8	—	48.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	30.3	77.7
Golden Melon	411	7.3	47.7	—	36.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	38.0	97.4
Swedish	2055	7.1	49.6	—	57.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	33.1	84.9
New Zealand	2056	6.3	40.2	—	52.7	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	27.7	88.5
Baku	333	5.0	—	60.4	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	36.9	98.7
Petschora	2058	4.2	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	56.3	122.7
Mensury	2057	2.7	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	24	56.3
Polar	2053	1.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	1.0
Prize Prolific	169	48.8	75.7	52.9	27.1	142.3	36.7	84.8	36.6	52.3	72.4	20.4	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	12	40.8
Goldthorpe	327	56.5	65.4	67.5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3	63.1
Saale	2045	40.3	—	54.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	13	123.7
Odesa	182	68.5	66.3	45.0	33.3	160.2	22.5	40.6	53.7	52.3	342.7	26.7	29.2	66.3	57.5	508.3	60.8	71.3	361.3	64.6	50.2	64.1	80.0	1.4	6.7	61.7	25	54.6	116.4
Rennie Improved	181	54.5	56.8	51.9	31.8	51.7	39.4	42.7	59.2	25.4	0.96	0.82	1.36	3.54	0.43	3.66	7	—	—	—	—	—	—	—	—	—	—	15	46.7
Baxter	185	44.0	—	32.5	30.2	37.9	36.5	52.7	63.7	53.2	63.7	53.2	63.7	53.2	63.7	53.2	63.7	53.2	63.7	53.2	63.7	53.2	63.7	53.2	63.7	53.2	63.7	13	39.9
Kulu	2094	50.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	50.0
Palampur	2091	60.6	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	60.6
Seoraj	2093	47.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	47.0
Soti Valley	2064	39.4	—	27.9	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2	33.7
Bhagamany Hills	2098	39.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	39.3
California Prolific	2049	—	—	61.4	45.0	19.0	445.0	62.9	21.3	49.0	56.6	3.24	4.17	5.29	6	—	—	—	—	—	—	—	—	—	—	—	—	6	42.8
Kinver Chevalier	587	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	11	36.6
Italian	342	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2	43.5
Sharpe Chevalier	2050	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2	54.2
Phoenix Von Thalen	325	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	11	36.6
Bethorn	331	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2	43.5
Peacock	2060	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	52.5
Kangra Valley	2089	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	48.3
Fun	341	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	43.4
Selected Chevalier	2052	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	38.5
Excelsior	2081	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	9	38.5
Common	184	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3	47.5
Champion	2083	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	15	37.0

TABLE 7.—Annual acre yields of varieties of barley grown at the experimental farm for Manitoba (at Brandon) in one or more of the 27 years from 1889 to 1915, inclusive—Continued

Variety	C. I. No.	Acre yields (bushels)																	Years grown	Average yield (bus.)	Percentage of weight-mean									
		1889	1890	1891	1892	1894	1895	1896	1897	1898	1899	1900	1901	1902	1903	1904	1905	1906				1907	1908	1909	1910	1911	1912	1913	1914	1915
Oderbrucker	2700					27.5	49.0	34.6	63.4	6.2	155.8	40.4	42.9	6.17	9.63	3.41	7.05	8.47	9.81	3.45	4.62	1.42	7					17	47.1	105.6
Trooper	173					26.7	65.2	240.8	51.3	32.9	65.8	296.9	29.2	17.1	153.3	353.3	353.3	355.2	50.6	49.4	64.2	41.9	62.9					18	48.3	107.3
Success	2707					25.0	30.2	235.6	84.4	24.1	73.1	529.6	18.3															13	40.3	96.2
Summit	174					22.3	59.0	299.8	80.0	25.7	56.6	324.6	63.7	52.2	1.6	352.9	604.6	50.2	27.1	3								14	48.3	110.8
Royal	1252					21.3	65.1	61.9	25.5	45.6	4.9	19.6	533.7	9.6	10.7	8.46	350.0	50.4										13	40.7	95.8
Surprise	171					17.3	36.5	238.7	39.9	35.8	53.1	332.1	115.0															9	39.5	102.6
Beaver	1915					32.1	150.2	299.6	634.2	25.7	52.2	524.0	108.7	21.3	332.1	162.5	64.8	54.5	41.5	60.4	43.1	43.2	56.2	58.6	63.1	147.9		22	42.3	90.8
French Chevalier	175					31.5	62.2	341.0	43.3	57.9	60.0	829.0	23.3	72.6	327.7	742.9	62.1	61.0	67.9	94.2	53.0	41.5						17	45.4	101.8
Golden Grain	588					23.6	58.3	332.7	23.3	45.4	49.8	334.4	27.3	7.5	24.9	555.4	56.7	51.0	71.9	94.1	9.65	4.61	3.65	5.62	3.66	8.43	4.66	1	25.6	93.8
Canadian Thorpe	740					24.8	58.3	332.7	23.3	45.4	49.8	334.4	27.3	7.5	24.9	555.4	56.7	51.0	71.9	94.1	9.65	4.61	3.65	5.62	3.66	8.43	4.66	1	47.1	101.1
Newton	172					19.2	56.9	47.9	22.1	53.7	61.0	222.8	19.6	14.2	23.3	330.4	53.3											12	36.1	86.6
Pioneer	2677					33.1																						7	39.1	105.7
Bolton	177					30.2																						8	36.9	99.5
Sidney	178					24.4	60.2	237.7	46.7	53.3	124.0	25.8	14.2	23.9	45.8	34.8	34.8	34.8	34.8	34.8	34.8	34.8	34.8	34.8	34.8	34.8	15	42.2	96.3	
Victor	179					20.2																						8	36.1	97.3
Stella	2678					18.5	53.7	742.3	33.3	68.3	356.3	334.0	21.7	26.3	354.5	53.3	3.3	74.9	0.63	34.8	7.67	9	23.3				17	45.8	102.2	
Nugent	176					68.5	45.4	441.0	51.7	61.5	537.2	33.3	20.8	73.7	147.5	90.0	65.8	54.2	47.1	61.3	41.9	59.0					17	51.2	111.5	
Vanguard	2665					25.3	51.7	42.0	29.4	46.7	58.3	725.8	39.6	15.6														8	40.0	101.3
Phoenix	2676					61.7	36.7	40.4																				3	46.3	105.5
Emerson	2739					61.7	36.7	40.4																				3	46.3	108.4
Monk	2703					33.1	121.5	45.4	46.0																			2	32.8	77.5
Rigid	2682					33.1	121.5	45.4	46.0																			2	32.8	87.5
Pacer	2667					30.2	44.6	34.5	45.8	47.5																		4	36.5	90.2
Nepean	183					30.2	44.6	34.5	45.8	47.5																		4	42.5	90.2
Blue Long Head	2685					27.7	35.4	40.4	733.3	33.3	13.1																	7	40.4	106.0
Argyle	2681					22.5	63.7	115.2	243.7	24.2	55.4	6.67	1.51	63.5														10	44.5	102.1
Mansfield	2241					40.8	64.5	635.6	47.5	51.7	160.3	32.7	57.7	47.9	67.5	56.9	71.3	46.9	53.4	73.3	54.6	46.9					16	50.2	111.3	
Empire	2680					55.4	57.9	93.3	33.3	25.8	61.7	58.7	55.8	56.5	70.4	45.2	5											11	51.0	112.6
Kirby	2671					65.4	55.4		28.3																			3	49.7	105.7
Dunham	2670					62.5	63.7	17.9	94.1	7.27	9.40	6.50	7.33	7.49	8.50	6												10	47.4	105.1
Leslie	2669					57.1	55.0	20.0	35.4	28.7																		5	39.2	102.3
Logan	2668					54.2	68.1	119.8	835.4	26.3	37.7	52.5	52.9	47.7	50.2													10	45.4	100.7
Harvey	2675					64.1	61.7	042.5	53.7	93.5	55.8	63.3	53.5															8	40.2	108.7
Fulton	2706					62.1	116.9	396.3	32.1	33.3	59.6																	6	38.3	98.5
Clifford	1940					56.7	119.2	289.9	223.7	31.9	58.7	754.2	200.6	654.2	36.9	54.5	48.7	50.2	65.8	59.8							15	47.6	100.0	
Jarvis	2672					55.8	82.2	647.9	19.0	23.0	55.5	863.7	66.5	72.1	35.8	863.3	30.4	40.4										13	48.7	105.9
Claude	1557					67.1	130.3	388.7	71.9	4.55	45.5	898.3	57.5	45.2	35.7	9.39	2.62	2										13	52.0	113.0
Yale	1592					53.3	336.3	347.1	113.7	30.0	62.5	63.7	60.2	5.56	4.88	3.54	6.63	37.1	55.4									15	56.1	117.9
Garfield	2673					51.7	332.7	346.3	330.4	455.1	452.5																	6	44.8	115.2

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

Explanation.—The asterisk (*) indicates that the two varieties were not grown in the same years]

Variety	C. I. No.	Data shown	French Cheva- lier	Canadian Thorpe	Hann- chen	Mensury	Odessa	Blue Long Head	Nepal	Himalaya	Yale	Excelsior
French Chevalier	175	{ Years comparable. Percentage yield	----- ----- -----	17 105.5	1 64.1 4	17 82.4 22	17 88.6	10 106.7	4 131.7	{ (*)	{ 12 82.2 9	9 105.9
Canadian Thorpe	740	{ Years comparable. Percentage yield	17 ----- 94.8	----- ----- -----	----- ----- 90.7	79.3 4	87.4 4	91.9	114.6	{ 98.3 4	{ 88.2 4	85.5
Hannehen	531	{ Years comparable. Percentage yield	1 ----- -----	4 110.2	----- ----- -----	100.0	----- ----- -----	(*)	(*)	{ 126.0 4	{ 115.1 15	(*)
Mensury	2657	{ Years comparable. Percentage yield	17 121.3	22 126.1	4 100.0	----- ----- 23	----- 109.6	10 120.8	4 182.6	{ 121.2 4	{ 109.5 15	9 124.5
Odessa	182	{ Years comparable. Percentage yield	17 112.9	10 114.4	90.5	91.3	----- ----- -----	10 113.9	145.4	{ 102.7 4	{ 101.7 8	104.5
Blue Long Head	2685	{ Years comparable. Percentage yield	93.8 ----- -----	108.8 ----- -----	{ (*) ----- -----	82.8 4 4	57.8 4 4	----- ----- -----	121.8	{ (*)	{ 96.5 2	83.7
Nepal	595	{ Years comparable. Percentage yield	4 ----- -----	4 ----- -----	(*) ----- -----	54.8 4 4	68.8 4 4	82.1	(*)	{ (*)	{ 69.2 2	79.4
Himalaya	1934	{ Years comparable. Percentage yield	{ (*) ----- -----	87.2 ----- -----	2 79.4 4	4 82.5 15	4 97.4	(*)	(*)	{ 95.1 4	{ (*)	4
Yale	1592	{ Years comparable. Percentage yield	12 15 4	101.7 113.4	15 86.9	4 91.4	98.3	8 103.6	4 144.5	{ 104.1 4	{ 114.7 4	114.7
Excelsior	2684	{ Years comparable. Percentage yield	9 ----- -----	9 116.9	9 ----- -----	80.3	95.7	6 119.4	126.0	{ (*)	{ 87.2 -----	-----

INDIAN HEAD, SASKATCHEWAN

The experimental farm at Indian Head, Saskatchewan, was also one of the earlier ones to be established in Canada. Field tests of barley were begun in 1888, and by 1915 some varieties had been tested for as many as 25 years. The results are given in Table 8. The Manchuria barleys yielded well at Indian Head, but they were not relatively so good as at Brandon. While the Chevalier group did not rank high, the yield of Danish Chevalier (C. I. No. 182) was 109 per cent of the weighted mean over a period of 25 years. From the digest made of the various groups the Manchuria is shown to be distinctly superior to all the others. The Thorpe and Hanna groups were better than the Chevalier at Indian Head. The Polar varieties were promising, but the hybrid sorts did not average high in yield. Ten varieties of different types are compared in the summarized digest of Table 8. Blue Long Head (C. I. No. 2685), a barley of North African origin, was the best of these by a small margin. This is of interest because it is a definite indication of the change of ecological conditions, the more arid summers being suitable to the growing of barleys of the North African type. It appears that Indian Head is on the border line of an ecological area as far as the production of barley is concerned. Hannchen (C. I. No. 531), Odessa (C. I. No. 182), Danish Chevalier (C. I. No. 180), and Mensury (C. I. No. 2657) gave yields in the order named. A number of varieties given in Table 8 were not in the test a sufficient number of years to establish their value. The most promising of these were Hannchen (C. I. No. 531), O. A. C. 21 (C. I. No. 1470), Gold (C. I. No. 1145), Standwell (C. I. No. 584), and Invincible (C. I. No. 590). These last two are hybrid varieties of the Thorpe type.

TABLE 8.—Annual acre yields of varieties of barley grown at the experimental farm for southern Saskatchewan (at Indian Head) in one or more of the 28 years from 1888 to 1915, inclusive

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)																										Years grown	Aver- age yield (bus.)	Per- centage of weight- ed mean	
		1888	1889	1890	1891	1892	1893	1894	1895	1896	1897	1898	1899	1900	1901	1902	1903	1904	1905	1906	1907	1908	1909	1910	1911	1912	1913				1914
Golden Melon	411	34.0	17.7	70.6	42.2	48.7																						6	39.7	104.2	
Peerless White	2647	33.3	18.8	40.3					45.0																				3	30.8	96.0
Chevalier	2651	24.0	26.7	143.3																									3	31.3	97.5
Thane	2652	56.7	21.2	49.1	49.0	43.7	51.5	18.7	54.8	50.6	45.2	54.4	57.1															12	46.0	102.9	
Polar	2653	33.0																											1	33.0	91.2
Danish Chevalier	180		12.7	46.2	44.4																							25	57.2	108.7	
English Maltling	2654		23.4	37.5																								3	33.5	100.9	
Swedish	2655		24.6	33.2																								2	38.9	122.7	
Premative	917		13.9	37.2	40.0																							3	30.4	89.1	
New Zealand	2656		23.1	36.7	43.3																							4	32.6	93.7	
Monsieur	2657		12.4	47.4	43.0	36.3	36.3	19.7	43.7	71.9	66.3	47.3	30.2	1.5	4.2	67.1	151.3	363.7	53.5	50.8	57.5	61.3	45.4	54.2	64.3	80.8	23	52.7	105.4		
Petschorn	2658		7.8																									11	43.6	93.4	
Prince Edward	2660		13.4																									3	37.6	113.3	
Beardless	2646		27.3	45.0		40.6																						2	36.6	115.5	
Scall	2645		19.6	33.6																								2	23.9	75.4	
Early Mintage	2661		13.7	34.0																								4	30.8	88.5	
Selected Chevalier	2662		12.0	25.0	30.7	35.4																						4	32.7	94.0	
Peerless	2663		16.3	39.0	26.4	38.7																						4	30.8	88.5	
Prize Prolific	169		49.1	45.0	48.4	44.0	41.7	79.0	52.1	50.0	53.7	56.7															4	48.3	101.5		
Duckbill	1916		55.4	60.0	31.1	50.4	41.1	74.5	60.0																			4	44.9	105.2	
Goldthorpe	327		45.5																									4	38.5	110.2	
Sharpe Chevalier	2650		46.9	47.2	49.2	42.1	115.6	45.2																				13	47.3	95.6	
Baxter	185		41.3																									10	57.7	112.3	
Odessa	182		38.1																									16	52.3	106.1	
Rennie Improved	181		39.3	46.7	41.7	49.4	118.7	62.3	55.4	468.1	56.7	60.6	46.3	52.5	46.7	57.3	62.2	63.3									20	50.9	50.9		
Stalkot	2665		20.0																									1	25.5	64.9	
Scoraj	2663		25.5																									1	38.2	97.2	
Kulu	2664		28.0																									1	24.0	61.1	
Simla	2662		24.0																									1	30.6	77.9	
Palampur	2661		30.6																									1	31.9	81.2	
Bhagamany Hills	2668		31.9																									1	28.0	71.2	
Lahoul	2660		28.0																									3	27.4	69.5	
Splai Valley	2664		37.2	24.7	20.4																							3	25.9	65.7	
Baku	333		28.0	26.7	22.9																							10	48.1	90.2	
Kinner Chevalier	587		41.7	54.7	19.4	56.5	54.2	55.1	55.3	55.0																		3	41.7	99.5	
California Prolific	2649		30.8	50.3	15.6	48.0	49.4																					5	35.6	95.4	
Golden Grain	588		35.6																									1	48.4	93.8	
Newton	172		39.3	15.6	64.8	108.7	75.1	55.3	34.3																			12	48.4	93.8	

TABLE 8.—Annual acre yields of varieties of barley grown at the experimental farm for southern Saskatchewan (at Indian Head) in one or more of the 28 years from 1888 to 1915, inclusive—Continued

C. I. No.	Variety	Acre yields (bushels)																		Years grown	Average yield (bus.)	Per- centage of weight- mean							
		1888	1889	1890	1891	1892	1893	1894	1895	1896	1897	1898	1899	1900	1901	1902	1903	1904	1905				1906	1907	1908	1909	1910	1911	1912
184	Common	36.9	16.3	41.4	408.	771.	3.40	0.58.	3.50.	8.53.	7.50.	0.00.	4.64.	6.65.	4.00.	8.	---	---	---	---	---	---	---	---	---	---	---	---	53.4
184	Himalaya	41.0	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	41.0
740	Canadian Thorpe	43.7	12.5	50.	165.	0.63.	1.50.	2.58.	7.15.	0.50.	48.7.	271.	0.55.	8.55.	0.48.	351.	7.40.	0.45.	8.51.	7.75.	0.	87.5	0.8.	3.52.	6.00.	2.	23	53.1	91.9
171	Surprise	39.0	12.5	38.	552.	3.55.	6.37.	5.56.	3.88.	7.56.	3.33.	7.46.	3.53.	7.	---	---	---	---	---	---	---	---	---	---	---	---	---	43.0	
174	Summit	40.7	15.6	40.	659.	4.55.	6.40.	0.62.	5.36.	7.56.	3.60.	8.01.	0.61.	9.62.	0.48.	7.42.	9.	---	---	---	---	---	---	---	---	---	---	50.1	
2700	Oderbrucker	22.9	40.	0.65.	271.	0.44.	7.58.	7.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	58.7	
2682	Rigid	20.0	48.	3.60.	4.50.	2.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	44.7	
1252	Royal	19.2	41.	4.61.	8.03.	7.45.	6.60.	0.41.	7.63.	3.58.	7.00.	0.38.	7.60.	3.50.	8.	---	---	---	---	---	---	---	---	---	---	---	---	52.8	
2676	Phoenix	17.7	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	42.8	
173	Trooper	16.9	45.	8.67.	3.37.	9.51.	5.69.	2.40.	0.50.	6.48.	3.64.	2.57.	0.62.	5.53.	7.58.	9.50.	2.	49.	2.67.	1.78.	3.	72.5	80.	8.	---	---	58.0		
1915	Beaver	15.7	52.	10.6.	7.92.	5.37.	1.55.	4.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	47.5	
177	Bolton	15.0	46.	2.55.	8.47.	3.41.	3.58.	3.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	48.2	
179	Victor	12.5	52.	6.0.	8.45.	6.37.	5.53.	3.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	86.3	
178	Sidney	12.5	52.	9.61.	9.44.	8.50.	6.37.	7.34.	2.61.	7.59.	1.62.	5.49.	0.51.	3.54.	0.50.	6.46.	3.	---	---	---	---	---	---	---	---	---	---	42.9	
2677	Pioneer	6.7	38.	2.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	49.6	
2707	Success	45.	0.50	0.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	41.6	
176	Nugent	42.	55.	4.	56.	3.47.	5.55.	0.27.	5.55.	8.53.	7.54.	6.07.	5.78.	7.49.	6.55.	6.37.	1.55.	2.62.	9.00.	8.	---	---	---	---	---	---	44.1		
2678	Stella	41.	7.55.	6.	38.	3.7.	3.61.	3.33.	5.1.	3.48.	7.54.	1.65.	4.82.	9.47.	9.52.	1.55.	8.45.	0.08.	3.85.	8.	90.0	30.0	57.	1.	83.	3.	107	55.6	
2684	Excelsior	41.	7.54.	3.7.	7.46.	3.55.	0.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	40.9	
2665	Vanguard	37.	5.55.	2.06.	7.30.	4.55.	0.33.	3.3.	45.8	40.4.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	45.4	
2683	Champion	37.	5.53.	3.54.	4.46.	3.46.	3.	47.1	36.7	45.4.	41.1	36.3	3.41.	7.47.	5.35.	8.	---	---	---	---	---	---	---	---	---	---	---	46.9	
2703	Monck	57.	9.37.	5.36.	5.48.	7.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	43.8	
175	French Chevalier	54.	57.3	3.53.	3.53.	1.65.	8.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	43.8	
183	Nepean	57.	5.47.	7.41.	7.50.	0.28.	3.38.	3.7.	9.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	45.1	
2667	Pacer	46.	0.	43.	5.35.	6.45.	4.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	45.9	
2685	Blue Long Head	63.	8.92.	5.65.	8.97.	1.	57.	1.	62.	5.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	42.6	
2681	Argyle	42.	4.45.	4.96.	7.64.	6.62.	1.47.	9.65.	6.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	60.0	
2241	Mansfield	43.	3.66.	7.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	55.9	
2689	Empire	37.	5.61.	3.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	61.8	
2671	Kirby	44.	6.58.	7.	37.	5.39.	2.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	56.0	
2670	Dunham	43.	7.57.	1.	20.	4.45.	4.	42.	1.32.	3.46.	6.41.	7.39.	2.48.	3.	---	---	---	---	---	---	---	---	---	---	---	---	---	47.5	
2669	Leslie	40.	0.55.	8.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	43.7	
2668	Logan	39.	0.49.	2.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	50.8	
1557	Claude	64.	6.	12.	5.62.	3.47.	6.47.	9.46.	3.56.	3.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	49.6	
1592	Yale	57.	3.0.	8.7.	3.49.	6.03.	5.00.	8.	61.	3.44.	6.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	63.3	
2679	Brome	57.	5.30.	8.58.	3.49.	6.03.	5.00.	8.	61.	3.44.	6.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	112.8	
2673	Gartfield	55.	8.25.	4.	53.	3.52.	1.55.	6.63.	8.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	53.3	
1551	Albert	50.	8.	45.	8.42.	1.53.	6.63.	7.	57.	5.42.	5.	51.	3.45.	1.	47.	1.	55.	4.	---	---	---	---	---	---	---	---	---	49.6	
596	Black Hull-less	44.	8.	45.	8.47.	5.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	46.0	
595	Nepal	40.	4.31.	7.37.	1.35.	0.	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	36.1	

Acre yields (bushels)

C. I.
No.

Variety

Years
grownAverage
yield
(bush.)Per-
centage
of
weight-
ed
mean

Jarvis.....	2672	51.315	0.57	1.52	5.61	5.46	6.50	0.62	9.61	0.47	148.7	54.2	61.7	---	---	---	---	13	50.7	92.7
Fulton.....	2705	50.0	20.9	8.42	9.59	0.90	9	---	---	---	---	---	---	---	---	---	---	6	45.6	86.4
Clifton.....	1910	43.6	20.8	5.4	5.5	5.67	3.43	0.51	7.32	5	53.7	54.5	58.7	60.6	---	---	---	15	54.2	92.4
Harvey.....	2675	49.6	57.5	4.8	7.88	3.43	5.47	5	46.7	---	---	---	---	---	---	---	---	17	51.1	92.4
Standwell.....	584	---	57.9	5.5	8.80	8.67	2.62	1.32	3.59	2.54	2	40.5	59.7	73.7	---	---	---	15	62.1	107.3
Invinible.....	590	---	55.8	5.7	3.77	3.67	1.74	6.32	3.54	6.41	7	45.4	63.8	74.2	---	---	---	15	61.9	106.9
Gordon.....	2674	---	52.5	4.5	8.64	6.43	5.04	6.52	5.54	6.96	3	52.3	---	---	---	---	---	9	54.7	101.3
Princess.....	552	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	1	62.5	109.6
Hannchen.....	529	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	1	60.4	106.0
Chevalier H.....	531	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	5	80.3	118.4
Black Japan.....	200	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	12	61.3	102.0
O. A. C. 21.....	2717	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	2	60.4	127.2
Manchurian.....	1470	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	6	83.2	117.3
Early Chevalier.....	739	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	6	75.7	106.8
Gold.....	2725	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	4	63.8	92.9
---	1145	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	3	75.9	114.0

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages									
			Danish Chevalier	Canadian Thorpe	Hannchen	Mon-sury	Odessa	Blue Long Head	Nepal	Baku	Stella	Success
Danish Chevalier.....	180	{ Years comparable.....	---	22	5	21	19	9	3	2	20	10
---	---	{ Percentage yield.....	---	110.2	91.5	101.4	98.3	92.6	157.8	165.6	101.6	139.2
Canadian Thorpe.....	740	{ Years comparable.....	22	---	5	19	19	10	4	(*)	21	10
---	---	{ Percentage yield.....	90.8	---	84.2	87.8	84.5	78.7	120.5	(*)	92.0	128.5
Hannchen.....	531	{ Years comparable.....	5	5	---	3	3	(*)	(*)	(*)	5	1
---	---	{ Percentage yield.....	100.3	118.7	---	111.3	102.7	(*)	(*)	(*)	103.1	137.6
Mensury.....	2657	{ Years comparable.....	21	19	3	---	---	10	4	3	17	7
---	---	{ Percentage yield.....	95.6	113.8	89.8	---	97.1	97.1	162.8	163.3	106.5	129.2
Odessa.....	182	{ Years comparable.....	19	19	3	20	---	10	4	1	17	7
---	---	{ Percentage yield.....	101.7	118.3	97.4	103.0	---	97.3	173.4	136.1	109.1	136.4
Blue Long Head.....	2685	{ Years comparable.....	9	10	(*)	10	---	---	---	(*)	10	5
---	---	{ Percentage yield.....	108.0	127.1	(*)	103.0	102.8	---	154.3	(*)	111.3	131.9
Nepal.....	595	{ Years comparable.....	3	4	(*)	---	---	4	---	(*)	4	3
---	---	{ Percentage yield.....	64.3	83.0	(*)	61.4	57.7	64.8	(*)	(*)	73.9	87.3
Baku.....	333	{ Years comparable.....	2	(*)	(*)	3	73.5	(*)	(*)	(*)	(*)	(*)
---	---	{ Percentage yield.....	60.4	---	---	61.2	---	---	---	---	---	---
Stella.....	2678	{ Years comparable.....	20	21	5	17	17	10	4	(*)	---	10
---	---	{ Percentage yield.....	98.4	108.7	97.0	93.9	91.7	89.9	135.2	(*)	10	132.5
Success.....	2707	{ Years comparable.....	10	7	1	7	7	5	3	(*)	75.5	---
---	---	{ Percentage yield.....	71.8	77.8	72.7	77.4	73.3	75.8	114.5	(*)	---	---

ROSTHERN, SASKATCHEWAN

Yields of barley are first reported in 1911. In the years 1911 to 1915 the tests included 15 varieties. The relative value of these varieties is more than usually apparent in Table 9, since 11 of them were tested for the full five years. Those of the Manchuria type gave very good yields. The best of these were O. A. C. 21 (C. I. No. 1470) and Manchurian (C. I. No. 739). Unfortunately, no variety of the Coast type was included in the test. Neither are the results quite comparable with those of the older stations, as many of the varieties tested on them were not grown at Rosthern. In the summarized digest of Table 9 a more direct comparison is made with eight varieties. The best of these again was O. A. C. 21, although Svanhals (C. I. No. 187) was practically as good. Duckbill (C. I. No. 1916), a barley of the Thorpe group, ranked third among the varieties compared. Barleys of this group also yielded well at Indian Head. Stella (C. I. No. 2678) was fourth in the list of eight varieties compared. This is one of the better hybrids produced by Doctor Saunders. Odessa (C. I. No. 182) yielded slightly less than Stella.

TABLE 9.—*Annual acre yields of varieties of barley grown at the experimental station for central Saskatchewan (at Rosthern) in part or all of the five years from 1911 to 1915, inclusive*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)					Years grown	Average yield (bus.)	Percentage of weighted mean
		1911	1912	1913	1914	1915			
Odessa.....	182	100.8	44.2	46.7	67.4	62.5	5	64.3	103.5
Manchurian.....	739	96.7	55.8	---	67.5	74.6	4	73.7	115.2
O. A. C. 21.....	1470	94.2	57.1	73.3	75.0	67.5	5	73.4	118.2
Black Japan.....	2717	93.3	70.8	58.3	67.5	74.6	5	72.9	117.4
Stella.....	2678	83.3	49.2	50.8	66.3	82.9	5	66.5	107.1
Mensury.....	2657	81.7	58.3	---	---	---	2	70.0	104.5
Taganrog.....	2721	81.7	59.2	59.2	67.1	57.5	5	64.9	104.5
Success.....	2707	58.3	49.2	31.7	32.7	43.7	5	43.1	69.4
Early Indian.....	1576	19.2	---	11.7	45.8	32.4	4	27.3	42.7
Duckbill.....	1916	85.0	61.7	67.5	66.9	53.3	5	66.9	107.7
Hannchen.....	531	81.7	47.5	---	---	---	2	64.6	96.4
Early Chevalier.....	2725	79.2	54.6	65.0	54.6	56.3	5	61.9	99.7
Svanhals.....	187	78.3	66.7	74.2	77.1	70.0	5	73.3	118.0
Chevalier II.....	200	71.7	55.8	56.7	53.3	72.9	5	62.1	100.0
Beaver.....	1915	70.8	38.3	46.7	45.4	40.8	5	48.4	77.9

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF EIGHT REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages							
			Chevalier II	Duckbill	Svanhals	Hannchen	O. A. C. 21	Odessa	Stella	Success
Chevalier II..	200	Years comparable.....	---	5	5	2	5	5	5	5
		Percentage yield.....	---	92.8	84.7	98.7	84.6	96.5	93.4	144.0
Duckbill.....	1916	Years comparable.....	5	---	5	2	5	5	5	5
		Percentage yield.....	107.7	---	91.3	113.5	91.1	104.0	100.6	155.1
Svanhals.....	187	Years comparable.....	5	5	---	2	5	5	5	5
		Percentage yield.....	118.0	109.5	---	112.2	99.8	113.9	110.2	169.9
Hannchen.....	531	Years comparable.....	2	2	2	---	2	2	2	2
		Percentage yield.....	101.3	88.1	89.1	---	85.4	89.1	97.5	120.2
O. A. C. 21.....	1470	Years comparable.....	5	5	5	2	---	5	5	5
		Percentage yield.....	118.3	109.8	100.2	117.1	---	114.1	110.4	170.3
Odessa.....	182	Years comparable.....	5	5	5	2	5	---	5	5
		Percentage yield.....	103.6	96.2	87.8	112.2	87.6	---	96.7	149.2
Stella.....	2678	Years comparable.....	5	5	5	2	5	5	---	5
		Percentage yield.....	107.1	99.4	90.8	102.6	90.6	103.4	---	154.2
Success.....	2707	Years comparable.....	5	5	5	2	5	5	5	---
		Percentage yield.....	69.5	64.5	58.9	83.2	58.7	67.0	64.8	---

SCOTT, SASKATCHEWAN

Yields from field plats of barley grown at Scott, Saskatchewan, in the years 1912, 1913, 1914, and 1915 are reported in Table 10. Five of the nine varieties are compared in detail in the summarized digest. Duckbill was the best of the varieties tested for the full four years. O. A. C. 21 (C. I. No. 1470), while second in point of yield, was distinctly inferior to Duckbill (C. I. No. 1916) for these four years. Black Japan (C. I. No. 2717) was third, and Early Chevalier (C. I. No. 2725) was fourth for productivity. There was very little difference in yield between O. A. C. 21, Black Japan, Early Chevalier, and Manchurian (C. I. No. 739). At Scott, as at the other points in Saskatchewan, barleys of the Thorpe type showed much greater promise than in eastern Canada, while barleys of the Manchuria type still yielded well.

TABLE 10.—*Annual acre yields of varieties of barley grown at the experimental station for northwestern Saskatchewan (at Scott) in part or all of the four years from 1912 to 1915, inclusive*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1912	1913	1914	1915			
Manchurian.....	739	65.0	25.8	10.0	61.9	4	40.7	101.0
O. A. C. 21.....	1470	64.4	35.8	14.6	65.2	4	45.0	111.7
Black Japan.....	2717	37.7	37.5	20.8	73.9	4	42.5	105.5
Success.....	2707	10.8	16.3	15.6	31.4	4	18.5	45.9
Early Indian.....	1576	2.1	-----	-----	-----	1	2.1	4.3
Duckbill.....	1916	88.0	29.6	25.8	69.2	4	53.1	131.8
Hannchen.....	531	75.3	-----	-----	-----	1	75.3	152.4
Early Chevalier.....	2725	51.8	37.1	17.1	60.2	4	41.5	103.0
Brewers Favorite.....	586	-----	-----	28.3	57.7	2	43.0	109.1

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF FIVE REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages				
			Early Chevalier	Duckbill	O. A. C. 21	Black Japan	Success
Early Chevalier.....	2725	Years comparable.....	-----	4	4	4	4
		Percentage yield.....	-----	78.2	92.3	97.8	221.3
Duckbill.....	1916	Years comparable.....	4	-----	4	4	4
		Percentage yield.....	127.9	-----	118.1	125.1	286.9
O. A. C. 21.....	1470	Years comparable.....	4	4	-----	4	4
		Percentage yield.....	108.3	84.7	-----	105.9	242.9
Black Japan.....	2717	Years comparable.....	4	4	4	-----	4
		Percentage yield.....	102.2	79.9	94.4	-----	220.3
Success.....	2707	Years comparable.....	4	4	4	4	-----
		Percentage yield.....	44.6	34.9	41.2	43.6	-----

LETHBRIDGE, ALBERTA

A part of the farm at Lethbridge, Alberta, is under irrigation. Barley varieties were grown both on dry land and under irrigation. With the exception of 1911 varietal experiments were conducted on the dry land at Lethbridge from 1908 to 1915, inclusive. During this period 32 varieties were tested for one or more years (Table 11). Of five groups compared at this station the Thorpe was distinctly the best. In order of yield the Hanna, Chevalier, Manchuria, and Hybrid follow the Thorpe. It is unfortunate that more groups were not represented. The only variety of the Coast group tested was Blue Long Head (C. I. No. 2685), and it was grown only for the year 1909. It was the best of the 24 varieties grown that year. It was also unfortunate that Hannchen (C. I. No. 531) was not continued in the experiment after 1912. For the two years in which it was grown, Hannchen was superior to the six other varieties with which it is compared in Table 11. Following Hannchen were Chevalier II (C. I. No. 200), Invincible (C. I. No. 590), and Odessa (C. I. No. 182), in the order named. It is apparent that Lethbridge is in a region better adapted to the cultivation of

the 2-rowed barleys than to those of the Manchuria type. The data at this point do not show which are superior, the 2-rowed barleys or those of the Coast type. The hybrid varieties were not, for the most part, as promising as they were at the stations farther east. Gold (C. I. No. 1145), a pedigreed barley of the Hanna type, was grown during the years 1914 and 1915. It showed promise during these years, but was not so good as several other varieties.

TABLE 11.—*Annual acre yields of varieties of barley grown on dry land at the experimental station for southern Alberta (at Lethbridge) in one or more of the eight years from 1908 to 1915, inclusive*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)							Years grown	Average yield (bus.)	Percentage of weighted mean
		1908	1909	1910	1912	1913	1914	1915			
Blue Long Head	2685	---	56.9	---	---	---	---	---	1	56.9	137.8
Claude	1557	41.3	55.4	11.9	28.7	40.0	30.0	79.1	7	40.9	114.6
Empire	2680	---	48.2	---	---	---	---	---	1	48.2	116.7
Mansfield	2241	48.8	42.3	10.0	27.5	35.0	32.5	96.9	7	41.9	117.4
Albert	1551	33.4	39.2	7.5	---	---	---	---	3	26.7	91.1
Odessa	182	41.1	37.9	12.5	34.4	40.0	42.5	96.9	7	43.6	122.1
Mensury	2657	31.9	37.9	10.6	---	---	---	---	3	26.8	91.5
Stella	2678	34.8	37.2	8.1	---	---	---	---	3	26.7	91.1
Nugent	176	33.2	36.4	6.3	---	---	---	---	3	25.3	86.3
Yale	1592	33.0	32.1	8.7	---	---	---	---	3	24.6	84.0
Oderbrucker	2700	33.9	30.6	7.5	---	---	---	---	3	24.0	81.9
Trooper	173	33.7	29.2	9.4	---	---	---	---	3	24.1	82.3
Champion	2683	---	20.4	---	---	---	---	---	1	20.4	49.4
Chevalier II	200	43.7	55.4	11.9	41.3	50.0	25.6	86.6	7	44.9	125.8
Invincible	590	39.8	54.0	10.6	42.5	47.5	27.8	83.4	7	43.7	122.4
Sidney	178	---	51.0	---	---	---	---	---	1	51.0	123.5
Standwell	584	34.9	49.6	9.4	---	---	---	---	3	31.3	106.8
Danish Chevalier	180	27.6	45.2	10.0	---	---	---	---	3	27.6	94.2
French Chevalier	175	28.8	43.7	6.3	---	---	---	---	3	26.3	89.8
Gordon	2674	---	41.6	---	---	---	---	---	1	41.6	100.7
Clifford	1910	31.3	41.6	15.6	30.0	35.0	23.7	70.0	7	35.3	98.9
Canadian Thorpe	740	46.4	40.8	10.6	---	---	---	---	3	32.6	111.3
Jarvis	2672	28.9	35.0	8.7	---	---	---	---	3	24.2	82.6
Beaver	1915	25.6	30.1	9.4	---	---	---	---	3	21.7	74.1
Hannchen	531	---	16.3	38.7	---	---	---	---	2	27.5	153.6
O. A. C. 21	1470	---	11.9	---	28.1	35.0	32.2	92.5	5	39.9	118.0
Cole	2731	---	30.6	40.0	---	---	---	---	2	35.3	101.4
Himalaya	1934	---	28.1	28.7	24.1	16.4	---	---	4	24.3	54.9
Manchurian	739	---	23.1	32.5	25.0	86.6	---	---	4	41.8	94.4
Early Chevalier	2725	---	23.7	39.4	24.7	67.2	---	---	4	38.7	87.4
Gold	1145	---	---	---	---	30.3	82.5	---	2	56.4	105.6
Duckbill	1916	---	---	---	---	20.0	85.6	---	2	52.8	98.9

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF SEVEN REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages						
			Chevalier II	Invincible	Hannchen	O. A. C. 21	Odessa	Himalaya	Mansfield
Chevalier II	200	Years comparable	---	7	2	5	7	4	7
		Percentage yield	---	102.9	96.7	107.9	103.0	209.1	107.3
Invincible	590	Years comparable	7	---	2	5	7	4	7
		Percentage yield	97.2	---	96.5	106.1	100.1	206.8	104.3
Hannchen	531	Years comparable	2	2	---	---	---	1	2
		Percentage yield	103.4	103.6	---	137.5	117.3	137.7	146.7
O. A. C. 21	1470	Years comparable	5	5	2	---	5	4	5
		Percentage yield	92.7	94.3	72.7	---	88.2	193.0	98.9
Odessa	182	Years comparable	7	7	2	5	---	4	7
		Percentage yield	97.1	99.9	85.3	113.3	---	219.7	104.2
Himalaya	1934	Years comparable	4	4	1	4	4	---	4
		Percentage yield	47.8	48.4	72.6	51.8	45.5	---	50.7
Mansfield	2241	Years comparable	7	7	2	5	7	4	---
		Percentage yield	93.2	95.9	68.2	101.1	96.0	197.2	---

Similar experiments were carried on under irrigation from 1908 to 1915, inclusive. Varieties were grown under irrigation in 1911, for which year no yields were reported on dry land. The yields under irrigation are reported in Table 12. Most of the varieties were in the tests for a very short period, which makes difficult a direct inspection of this table. In the unreported group digest the Chevalier group produced the highest yields, while Thorpe, Manchuria, Hanna, and Hybrid ranked second, third, fourth, and fifth, respectively. Although the Hybrid group as a group was quite poor, it will be seen in the summarized digest of Table 12 that Claude (C. I. No. 1557) was the best variety compared. Chevalier II was almost the equal of Claude and Invincible (C. I. No. 590) somewhat inferior, while O. A. C. 21 (C. I. No. 1470) was nearly equal to Invincible. Gold was grown during the years 1914 and 1915, for which years it was third in point of yield, being inferior to both Claude and Chevalier II.

TABLE 12.—*Annual acre yields of varieties of barley grown on irrigated land at the experimental station for southern Alberta (at Lethbridge) in part or all of the eight years from 1908 to 1915, inclusive*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)								Years grown	Average yield (bus.)	Percentage of weighted mean
		1908	1909	1910	1911	1912	1913	1914	1915			
Claude	1557	63.6	59.8	42.5	81.3	81.6	93.7	97.5	80.0	8	75.0	122.0
Mansfield	2241	58.7	49.6	37.5	61.3	61.9	84.1	62.5	83.7	8	62.4	101.5
Blue Long Head	2085		45.2							1	45.2	108.1
Yale	1592		45.2							1	45.2	108.1
Odessa	182	61.4	44.5	30.0	65.0	80.6	91.0	66.9	75.0	8	64.3	104.6
Empire	2680		37.9							1	37.9	90.7
Mensury	2657	53.6	37.2	25.0	51.3					4	41.8	87.6
Nugent	176		36.5							1	36.5	87.3
Oderbrueker	2700		36.5							1	36.5	87.3
Albert	1551		30.6							1	30.6	73.2
Stella	2678		30.6							1	30.6	73.2
Champion	2683		27.7							1	27.7	66.3
Trooper	173		24.8							1	24.8	59.3
Standwell	584	64.3	70.0	43.3	63.7					4	60.3	126.4
Sidney	178		62.7							1	62.7	150.0
Chevalier II	200	68.9	61.3	53.7	70.0	77.5	93.1	90.6	80.6	8	74.5	121.1
Danish Chevalier	180		45.2							1	45.2	108.1
Canadian Thorpe	740		43.7							1	43.7	104.5
Gordon	2674		42.3							1	42.3	101.2
French Chevalier	175		40.1							1	40.1	95.9
Clifford	1910		36.5	27.5	41.3	55.4	65.0	53.7	67.5	7	49.6	80.9
Invincible	590		35.2	43.3	62.5	69.4	94.4	82.8	91.3	7	68.4	111.2
Jarvis	2672		32.8							1	32.8	78.5
Beaver	1915		28.4							1	28.4	67.9
O. A. C. 21	1470				73.7	76.9	88.7	67.8	90.0	5	79.4	108.0
Manchurian	739				41.9	78.9	73.9	63.1	72.5	5	66.1	89.9
Hannehen	531					66.9				1	66.9	95.3
Early Chevalier	2725					52.8	68.6	54.4	80.0	4	63.9	83.9
Cole	2731						100.6			1	100.6	120.2
Himalaya	1934						68.1	75.0	66.9	3	70.0	89.7
Gold	1145							87.2	81.3	2	84.3	112.0
Duekhill	1916							63.4	73.7	2	68.5	91.0

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF SIX REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages					
			Chevalier II	Invincible	O. A. C. 21	Odessa	Himalaya	Claude
Chevalier II	200	Years comparable	-----	7	5	8	3	8
		Percentage yield	-----	110.0	103.7	115.8	125.9	99.3
Invincible	590	Years comparable	7	-----	5	7	3	7
		Percentage yield	90.9	-----	100.8	105.7	127.9	89.3
O. A. C. 21	1470	Years comparable	5	5	-----	5	3	5
		Percentage yield	96.4	99.2	-----	104.9	117.4	91.5
Odessa	182	Years comparable	8	7	5	-----	3	8
		Percentage yield	86.3	94.6	95.3	-----	110.9	85.7
Himalaya	1934	Years comparable	3	3	3	3	-----	3
		Percentage yield	79.5	78.2	85.2	90.2	-----	77.4
Claude	1557	Years comparable	8	7	5	8	3	-----
		Percentage yield	100.7	112.0	109.3	116.6	129.1	-----

TABLE 13.—*Annual acre yields of varieties of barley grown at the experimental station for central Alberta (at Lacombe) in part or all of the nine years from 1907 to 1915, inclusive*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)									Years grown	Average yield (bus.)	Percentage of weight-mean
		1907	1908	1909	1910	1911	1912	1913	1914	1915			
Nugent	176	72.5	40.0	18.1	53.7	80.8	-----	-----	-----	-----	5	53.0	105.6
Mensury	2657	72.5	47.5	40.0	63.7	-----	-----	-----	-----	-----	4	55.9	122.3
Oderbrucker	2700	70.0	29.2	20.0	58.1	77.5	-----	-----	-----	-----	5	51.0	101.6
Mansfield	2241	68.7	62.5	28.7	63.1	83.3	79.6	68.3	72.9	48.8	9	64.0	125.0
Stella	2678	65.0	43.7	18.1	80.0	70.5	62.5	60.4	59.0	41.7	9	59.3	115.8
Claude	1557	60.0	41.3	33.1	79.1	92.3	-----	-----	-----	-----	5	61.2	121.9
Seamit	174	57.5	-----	-----	-----	-----	-----	-----	-----	-----	1	57.5	119.3
Odessa	182	26.9	45.0	27.5	77.5	77.5	75.4	55.0	71.9	50.0	9	56.3	110.0
Blue Long Head	2685	53.7	60.0	-----	-----	-----	-----	-----	-----	-----	2	56.9	127.6
Empire	2680	51.3	31.3	-----	-----	-----	-----	-----	-----	-----	2	41.3	92.6
Albert	1551	48.7	43.7	19.4	73.1	87.7	-----	-----	-----	-----	5	54.5	108.6
Champion	2683	48.7	33.7	-----	-----	-----	-----	-----	-----	-----	2	41.2	92.4
Yale	1592	47.5	33.7	19.4	77.5	-----	-----	-----	-----	-----	4	44.5	97.4
Argyle	2681	47.5	-----	-----	-----	-----	-----	-----	-----	-----	1	47.5	98.5
Trooper	173	42.5	23.7	16.9	71.3	74.0	-----	-----	-----	-----	5	45.7	91.0
Clifford	1910	50.0	27.5	18.7	47.5	55.4	-----	-----	-----	-----	5	39.8	79.3
French Chevalier	175	46.3	37.5	23.4	47.2	-----	-----	-----	-----	-----	4	38.6	84.5
Gordon	2674	45.0	43.7	21.3	-----	-----	-----	-----	-----	-----	3	36.7	94.3
Canadian Thorpe	740	45.0	33.7	28.7	68.7	54.6	-----	-----	-----	-----	5	46.1	91.8
Dunham	2670	43.7	-----	-----	-----	-----	-----	-----	-----	-----	1	43.7	90.7
Logan	2668	42.5	-----	-----	-----	-----	-----	-----	-----	-----	1	42.5	88.2
Standwell	584	40.0	53.7	32.5	80.9	56.0	-----	-----	-----	-----	5	52.6	104.8
Sidney	178	38.3	55.0	18.7	-----	-----	-----	-----	-----	-----	3	37.3	95.9
Invincible	590	37.5	56.9	43.7	75.0	69.8	65.2	61.3	71.5	31.7	9	55.8	109.0
Beaver	1915	36.9	18.7	19.7	44.4	49.2	-----	-----	-----	-----	5	33.8	67.3
Danish Chevalier	180	32.5	32.5	18.4	49.4	57.7	-----	-----	-----	-----	5	38.1	75.9
Jarvis	2672	31.3	21.3	16.0	45.3	-----	-----	-----	-----	-----	4	28.5	62.4
Chevalier II	200	28.7	52.5	28.1	85.0	60.8	67.1	64.0	85.8	46.3	9	57.6	112.5
Baku	333	-----	-----	52.3	67.5	-----	-----	-----	-----	-----	2	59.9	127.7
O. A. C. 21	1470	-----	-----	77.8	80.8	79.2	58.3	67.3	45.4	-----	6	68.1	112.0
Himalaya	1934	-----	-----	80.0	56.3	50.0	41.0	34.6	31.7	-----	6	48.9	80.4
Hannchen	531	-----	-----	83.7	73.4	76.3	68.7	-----	-----	-----	4	75.5	116.0
Early Chevalier	2725	-----	-----	-----	57.7	56.5	50.4	46.3	11.5	-----	5	44.5	75.9
Manchurian	739	-----	-----	-----	88.3	69.2	54.4	69.2	41.3	-----	5	64.5	110.1
Princess	529	-----	-----	-----	66.9	-----	-----	-----	-----	-----	1	66.9	96.5
Primus	532	-----	-----	-----	66.4	-----	-----	-----	-----	-----	1	66.4	95.8
Svanhals	187	-----	-----	-----	60.0	75.4	67.1	45.4	9.2	-----	5	51.4	87.7
Success	2707	-----	-----	-----	-----	36.5	26.9	10.0	32.5	-----	4	26.5	49.3
Early Indian	1576	-----	-----	-----	-----	-----	12.1	-----	-----	-----	1	12.1	22.9
Gold	1145	-----	-----	-----	-----	-----	-----	71.3	53.1	-----	2	62.2	129.9

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages									
			Chevalier II	Invincible	Svanhals	Hannchen	O. A. C. 21	Odessa	Blue Long Head	Himalaya	Mansfield	Success
Chevalier II	200	Years comparable	-----	9	5	4	6	9	2	6	9	4
		Percentage yield	-----	103.1	126.0	91.7	100.0	102.3	71.4	139.3	90.0	248.5
Invincible	590	Years comparable	9	-----	5	4	6	9	2	6	9	4
		Percentage yield	97.0	-----	112.6	86.5	89.2	99.2	83.0	124.1	87.3	216.9
Svanhals	187	Years comparable	5	5	-----	3	5	5	(*)	5	5	4
		Percentage yield	79.4	88.8	-----	92.7	77.7	78.0	-----	120.4	72.9	186.1
Hannchen	531	Years comparable	4	4	3	-----	4	4	(*)	4	4	2
		Percentage yield	109.1	115.6	107.9	-----	102.0	105.9	-----	132.9	102.7	228.7
O. A. C. 21	1470	Years comparable	6	6	5	4	-----	6	(*)	6	6	4
		Percentage yield	100.0	112.2	128.7	98.0	-----	100.4	-----	139.2	98.3	236.3
Odessa	182	Years comparable	9	9	5	4	6	-----	2	6	9	4
		Percentage yield	97.8	100.8	128.3	94.5	99.6	-----	63.2	138.7	88.0	238.2
Blue Long Head	2685	Years comparable	2	2	(*)	(*)	(*)	2	-----	(*)	2	(*)
		Percentage yield	140.0	120.4	-----	-----	158.1	-----	-----	-----	86.7	-----
Himalaya	1934	Years comparable	6	6	5	4	6	6	(*)	-----	6	4
		Percentage yield	71.8	80.5	83.1	75.2	71.8	72.1	-----	-----	70.6	148.5
Mansfield	2241	Years comparable	9	9	5	4	6	9	2	6	-----	4
		Percentage yield	111.1	114.6	137.3	97.4	101.8	113.7	115.4	141.7	-----	254.6
Success	2707	Years comparable	4	4	4	2	4	4	(*)	4	4	-----
		Percentage yield	40.2	46.1	53.7	43.7	42.3	42.0	-----	67.3	39.3	-----

LACOMBE, ALBERTA

Barleys were grown at Lacombe, Alberta, from 1907 to 1915, inclusive, and the results are given in Table 13. Here, again, the groups are not so well represented as would be desirable. The Coast group is represented by one variety, Blue Long Head (C. I. No. 2685), which was grown in 1907 and 1908. For these years its yield was 128 per cent of the weighted mean of all varieties. It produced second highest yield of all the sorts tested in 1908, the last year it was in the test. It is also unfortunate that Hannchen (C. I. No. 531) was not carried longer. It gave the highest yield of all varieties tested in 1913, the year in which it was discarded. There were only four groups that could be compared. Of these, Manchuria ranked highest. The Thorpe, Hybrid, and Chevalier II followed in the order named.

In the summarized digest of Table 13 10 varieties are compared with each other. Owing to the fact that these varieties were seldom comparable for many years, the results are not very conclusive. The varietal comparison in this case is more indicative of the type values than the group comparison. Hannchen was the best of the varieties compared. Mansfield (C. I. No. 2241), one of the hybrid varieties, was slightly inferior to Hannchen. O. A. C. 21 (C. I. No. 1470) and Chevalier II (C. I. No. 200) were practically equal in yield. Blue Long Head was synchronous with only 4 of the 10 varieties compared. In the two years in which comparisons can be made, it was inferior to Mansfield but superior to Chevalier II, Invincible (C. I. No. 590), and Odessa (C. I. No. 182). Success (C. I. No. 2707) in all but two instances gave a yield of less than half that of other varieties. Barleys of this type have been included in the digest of results from most stations, because they are widely sold by seedsmen and because of the fact that if they possessed high yielding power they would be preferred by farmers. Gold (C. I. No. 1145) was grown during the years 1914 and 1915. It was the highest yielding variety in 1915, but only fifth in yield in 1914.

TABLE 14.—*Annual acre yields of varieties of barley grown at the experimental station at Fort Vermilion, Alberta, in part or all of the four years from 1912 to 1915, inclusive*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1912	1913	1914	1915			
Mensury.....	2657	75.6	61.3	55.0	66.3	4	64.5	111.0
Claude.....	1557	63.3	60.0	57.5	71.8	4	63.1	108.6
Canadian Thorpe.....	740	58.7	48.7	62.5	46.6	4	54.1	93.1
Sidney.....	178	55.6	30.4	61.3	51.7	4	49.7	85.5
Champion.....	2683		65.0		67.2	2	66.1	116.0
Success.....	2707		60.0	51.3	60.5	3	57.3	100.5
Nepal.....	595			53.7	51.9	2	52.8	90.6

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF FIVE REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages				
			Canadian Thorpe	Mensury	Nepal	Claude	Success
Canadian Thorpe.....	740	{Years comparable.....	-----	4	2	4	3
		{Percentage yield.....	-----	83.8	103.3	85.7	91.9
Mensury.....	2657	{Years comparable.....	4	-----	2	4	3
		{Percentage yield.....	119.3	-----	114.1	102.2	106.3
Nepal.....	595	{Years comparable.....	2	2	-----	2	2
		{Percentage yield.....	96.8	87.1	-----	81.7	94.5
Claude.....	1557	{Years comparable.....	4	4	2	-----	3
		{Percentage yield.....	116.7	97.8	122.4	-----	110.2
Success.....	2707	{Years comparable.....	3	3	2	3	-----
		{Percentage yield.....	108.9	94.1	105.9	90.8	-----

FORT VERMILION, ALBERTA

The yields reported in Table 14 were from barleys grown at Fort Vermilion, Alberta. Seven varieties were tested in the four years from 1912 to 1915, inclusive. In the summarized digest of the table five of these are compared. The best is Mensury (C. I. No. 2657). Claude was second highest in yield, while Success (C. I. No. 2707) was better than Canadian Thorpe (C. I. No. 740). Nepal (C. I. No. 595), as usual, was inferior to Success. The range of varieties in this test is not sufficient to draw any conclusions as to the types best adapted to this region. It does appear, however, that Manchuria does very well in the Fort Vermilion section.

ST. BERNARD MISSION SUBSTATION, GROUARD, ALBERTA

Two varieties of barley, Odessa (C. I. No. 182) and Manchurian (C. I. No. 739), were grown at St. Bernard Mission substation, Grouard, Alberta, in 1913.

Their yields were 35.2 bushels and 34 bushels, respectively. Manchurian was grown again in 1914, when it produced 40 bushels to the acre. It would seem that barleys give good yields as far north as Grouard.

BEAVER LODGE, ALBERTA

Three varieties of barley were grown at Beaver Lodge, Alberta, in 1915. Owing to an accident the yield of Success (C. I. No. 2707) was not determined, although it appeared to be the poorest of the three. Manchurian (C. I. No. 739) yielded 42.5 bushels and French Chevalier (C. I. No. 175) 38.7 bushels.

EXPERIMENTS IN THE NORTHWEST TERRITORY

In 1915 barley was grown at three points in the Northwest Territory, but the yields were not reported. At Fort Resolution, in the Mackenzie district, barley and wheat did not mature successfully. They were seeded rather late and did not ripen before frost. At Fort Providence, also in the Mackenzie district, Manchuria and Success were grown. They were seeded on April 23 and ripened about August 1. At Fort Smith the results were considered promising, although the plats were injured by heavy frosts in June.

AGASSIZ, BRITISH COLUMBIA

Barleys were first grown at Agassiz in 1891. More than 70 varieties were tested in one or more of the years from 1891 to 1915, inclusive. The annual yields are reported in Table 15.

A comparison was made of six groups of barley. The Thorpe was the best of the six. The Manchuria was almost as good as the Thorpe, while the Hybrid, Chevalier, Hanna, and Himalaya groups follow in the order named. There were not sufficient varieties of the Coast type to include a Coast group. Ten well-known varieties of various types are compared in the summarized digest of Table 15. Hannehen (C. I. No. 531) was grown only four years, but was superior to the other nine varieties for the years in which the yields were comparable. Mensury (C. I. No. 2657) exceeded all varieties except Hannehen. Since Blue Long Head (C. I. No. 2685), Danish Chevalier (C. I. No. 180), Canadian Thorpe (C. I. No. 740), and Odessa (C. I. No. 182) also produced good yields, it is evident that barleys of many types grow well in the valley in which Agassiz is situated. This is not true in the barley-growing districts of California, eastern Oregon, and Washington, where varieties of the Coast type are outstanding.

TABLE 15.—Annual acre yields of barley grown at the Agassiz (British Columbia) Experimental Farm in part or all of the 25 years from 1891 to 1915, inclusive

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C.I. No.	Acre yields (bushels)															Years age grown	Average yield (bus.)	Per-centage of weighted mean										
		1891	1892	1893	1894	1895	1896	1897	1898	1899	1900	1901	1902	1903	1904	1905				1906	1907	1908	1909	1910	1911	1912	1913	1914	1915
Golden Melon	411	36.0	21.9	29.0																						3	29.0	105.1	
Sealo	2645	33.6	27.9																								2	30.7	105.5
Pitso Prolific	169	32.7	16.9	18.6	33.3	32.9	23.2	53.6	33.0	8.36	3.59	2.43	3.													11	33.6	97.8	
Thano	2652	24.0	20.6	12.2	52.5	8.3	31.2	22.2	93.3	7.																8	25.4	83.8	
Duckbill	1916	24.0	23.0	6.24	23.0	4.4	0.28	3.																		8	30.3	111.4	
Kniver Chevalier	587	20.8	23.3	22.2	18.3	28.3	33.0	64.0	84.0	0.29	2.82	5.50	8.48	3.												12	32.0	95.0	
Peerless White	2647	20.0	26.9	9.																						2	27.9	93.9	
Sharpe Chevalier	2650	32.9	28.3	120.3																						4	25.7	98.7	
Danish Chevalier	180	32.7	27.9	30.3	30.6	40.0	33.1	33.3	33.3	3.7	1.47	1.37	5.65	4.	56.7	35.8	4.1	3.3	3.	58.3	5.1	3.6	1.53	7.41	33.7	41.3	43.5	108.7	
Goldthorn	327	42.1	18.0	13.0	29.9	9.14	20.0																			6	34.2	123.7	
Golden Grain	388	47.7	18.0	13.0	123.8	82.7	28.2																			6	30.5	102.7	
Bennie Improved	183	37.5	27.9	23.3	15.3	3.	28.3	32.2	83.4	6	40.8	38.9	53.	735.	6.64	6.										12	33.7	99.7	
Odesa	182	31.7	27.9	23.3	15.3	3.	28.3	19.6	37.2	27.9	30.0	54.	5.62	3.65	4.	49.2										14	36.3	93.8	
Oderbrucker	2700	22.8	11.0	0.28	20.0	33.3	28.3	30.4	42.0	43.8	3.51	3.90	8.71	3.88	3.48	3.47	1.42	1.47	5.54	2.50	2.36	9.88	7.28	3.46	3.33	1	40.1	101.5	
Common	184	37.1	19.4	43.3	16.1	128.1	23.2	23.8	3.54	6.30	8.35	9.00	5.80	4.67	1.36	3.47	5.36	7.								16	33.2	102.1	
Himalaya	1934	41.9	22.2	120.3																						3	28.1	101.8	
Mensury	2657	35.8	28.3	25.1																						20	43.2	108.3	
Spiti Valley	2664	20.5	19.2																							2	19.9	68.4	
French Chevalier	175		22.9	93.8	3.4	4.2	23.7	131	730	0.34	51.9	33.3	370	8.82	1.46	5.41	9.41	7.	51.7	40	3.41	5.				18	40.9	102.0	
Canadian Thorpe	740	24.8	27.3	7.41	23.7	127	936	7.31	50.4	44.6	7.69	5.66	51	7.32	9.48	7.42	9.55	0.61	3.33	7	46	33.3	338	7	50.0	23	42.9	107.3	
Newton	172	16.2	28.8	8.34	218	43.1	33.3	33.9	23.1	5.51	9.40	5.62	9.0	2	55.0											13	38.0	97.4	
Petchora	2658	23.	7.16	73.3	17.9	33.8	3.28	7.32	33.9	8.55	4.35	2.														10	32.1	92.5	
California Prolific	2649		37.1	24.6	25.4																					3	24.0	107.8	
Beaver	1915		23.8	36.0	37.1	128.	739	23.1	33.4	2.61	9.33	175.	0.50	4.51	337.5	42.7	50.0	53.7	743.	135.0	42	534	243	7	86.6	22	44.1	108.9	
Royal	1252		17.1	129	825	536	538	334	632	67	539	608	7	46	344	231.7										13	39.4	99.7	
Phoenix	2676		15.0	82	8.15	0.33	340	0.85	3.66	5.53	0.35	8.														9	32.5	92.1	
Monck	2703		38.3	616	3.28	325	830	8.																		5	28.0	88.9	
Bolton	177		30.5	37.1	128.	131	333	33.1	9.50	829.6																8	32.8	91.	
Victor	179		27.7	24	230	429	234	634	645	831.1																8	32.2	89.4	
Sidney	178		27.7	32.2	127.	530	833	7.30	4	37	939	273	3.56	4	39.2	41.0	52	52.7								14	40.3	98.8	
Rigid	2082		26	30	0.27																					3	24.5	81.4	
Trooper	173		34.0	22.2	3.65	0.92	938	7.50	8	36	65.5	4	50.1	33	339	248	5	49	2	43.1	9	31	3	33.7	30	46	3	68.7	
Stella	2678		32.1	119	634	234	235	429	647	337.1	172	5	53	6.50	2	37	589	2	40	0	50.8					21	41.1	100.2	
Vanguard	2665		31.3	16	737	531	531	336	947	339.4																16	34.0	94.0	
Summit	174		28.4	20	33.3	535	822	929	847	155	0.90	4	50	8	39	2	33	342	5						8	34	94	90	
Champion	2683		26.7	20	0	33.3	335	436	4	37	751	138	1	61	7	53	1	45	8	35	841	9	30	8		13	39.5	97.3	
																										14	39.1	95.	

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C.I. No.	Data shown	Varieties and percentages									
			Danish Chevalier	Canadian Thorpe	Hann- chen	Mensury	Pet- schora	Odessa	Blue Long Head	Nepal	Beaver	Success
Danish Chevalier	180	{ Years comparable. Percentage yield.	22 102.3	22 102.3	4 95.6	20 98.4	9 108.5	24 107.8	10 97.7	4 139.4	21 100.5	12 130.3
Canadian Thorpe	740	{ Years comparable. Percentage yield.	22 97.7	22 97.7	4 95.9	18 99.0	10 114.2	23 104.5	10 97.2	4 135.8	22 99.2	12 127.5
Hannchen	531	{ Years comparable. Percentage yield.	4 104.6	4 104.2	— —	2 133.4	(*) —	4 118.1	(*) —	(*) —	4 117.6	2 123.3
Mensury	2657	{ Years comparable. Percentage yield.	20 101.7	18 101.0	2 74.9	— —	9 111.0	20 105.1	10 107.9	3 135.4	17 104.7	8 125.2
Petschora	2658	{ Years comparable. Percentage yield.	9 92.2	10 87.6	(*) —	9 90.1	— —	10 90.3	6 100.0	3 121.8	9 91.7	8 112.4
Odessa	182	{ Years comparable. Percentage yield.	24 92.7	23 95.7	4 84.7	20 95.1	10 110.7	— —	10 101.1	4 120.3	22 95.0	12 118.2
Blue Long Head	2885	{ Years comparable. Percentage yield.	10 102.4	10 102.9	(*) —	10 92.6	6 100.0	10 99.0	— —	3 121.4	10 102.3	6 109.2
Nepal	595	{ Years comparable. Percentage yield.	4 71.7	4 73.6	(*) —	3 73.9	3 82.1	4 83.2	3 82.4	— —	— —	4 89.8
Beaver	1915	{ Years comparable. Percentage yield.	21 99.5	22 100.8	4 85.1	17 95.5	9 109.0	22 105.3	10 97.7	4 156.7	— —	134.0
Success	2707	{ Years comparable. Percentage yield.	12 76.7	12 78.4	2 81.1	8 79.9	8 89.0	12 84.6	6 91.6	4 111.3	12 74.6	— —

SIDNEY, BRITISH COLUMBIA

At Sidney, British Columbia, two varieties of spring barley were grown in 1914 and 1915 and two varieties of winter barley in 1915. The yields are reported in Table 16. The number of varieties tested was not sufficient to determine the general type of barley best suited to the region. The spring varieties, however, were superior to the winter varieties. The yields were not so good as those obtained at Agassiz in the same years.

TABLE 16.—*Annual acre yields of varieties of barley grown at the Sidney (British Columbia) Experimental Station in one or both of the years 1914 and 1915*

[Data compiled from the Canadian Experimental Farms Reports]

Variety	C. I. No.	Acre yields (bushels)		Years grown	Average yield (bus.)	Percentage of weighted mean
		1914	1915			
Spring barley:						
Manchurian.....	739	30.0	35.2	2	32.6	112.8
Canadian Thorpe.....	740	23.1	34.9	2	29.0	100.3
Winter barley:						
Arlington Awnless.....	702		30.0	1	30.0	99.7
Tapp Winter.....	1707		20.5	1	20.5	68.1

INVERMERE, BRITISH COLUMBIA

Barley yields were reported from Invermere, British Columbia, only for the year 1915. Four varieties were tested during that year. The yields were as follows: Manchurian (C. I. No. 739), 25 bushels; Gold (C. I. No. 1145), 24.2 bushels; Early Chevalier (C. I. No. 2725), 20 bushels; Success (C. I. No. 2707), 16.9 bushels. In this year at Agassiz and Sidney barleys of the Manchuria type gave very good yields.

DATA FROM NORTHERN AND WESTERN STATIONS IN THE UNITED STATES

The data from the agricultural experiment stations in the United States are presented in two sections, namely, (1) those from northern and western stations and (2) those from southern stations. As noted previously, the stations are arranged in sequence from east to west in each section.

The yields of barley at the Canadian stations which have been followed across Canada from east to west can be used as a background for interpreting the results of experiments at the stations in the northern United States. The barley area in the United States with the exception of the Southwest lies mostly along the Canadian border. The tests at stations in Canada have been more extensive and less subject to interruption than those at experiment stations in the United States. The results at points just north of our northern border are fully applicable to our northern tier of States. In many cases the Canadian stations are closer to the northern counties of our States than are our own stations.

ORONO, ME.

In the annual reports of the Maine Agricultural Experiment Station from 1886 to 1889, yields are given for seven varieties of barley. These yields are reported in Table 17. It will be seen that the best two varieties were Chevalier (C. I. No. 2802) and Champion of Vermont (C. I. No. 1892). The latter is a barley of the Hanna type. The Golden Melon (C. I. No. 411) and Imperial (C. I. No. 2805) are barleys of the Thorpe group, but Mensury (C. I. No. 2657) belongs to the Manchuria group. It is of special interest that the Hanna type gave good

yields in Maine, for it will be remembered that the Chevalier barleys were best adapted to Prince Edward Island and Manchuria to the mainland of eastern Canada, while Hanna was not particularly promising. But Hanna has long been known to be a good barley in New England and New York State. In other words, the Hanna barleys are adapted to a somewhat warmer district than those of the Chevalier type.

TABLE 17.—*Annual acre yields of varieties of barley grown at the Maine Agricultural Experiment Station (at Orono) in some or all of the four years from 1886 to 1889, inclusive*

[Data obtained through the courtesy of the Maine Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1886	1887	1888	1889			
Sibley Imperial.....	2805	28.7	17.1	20.8	-----	3	22.2	80.7
Mensury.....	2657	28.7	16.7	33.1	-----	3	26.2	95.3
Chevalier.....	2802	23.2	19.6	50.4	33.9	4	31.8	110.8
Nepal.....	595	17.1	32.5	31.2	-----	3	26.9	97.8
Golden Melon.....	411	26.5	18.8	34.2	-----	3	26.5	96.4
Black Hull-less.....	596	21.0	24.6	25.0	-----	3	23.5	85.5
Champion of Vermont.....	1892	-----	34.8	46.7	46.3	3	42.6	140.1

DURHAM, N. H.

Through the courtesy of the New Hampshire Agricultural Experiment Station, barley yields for the years 1908 and 1909 were supplied for this bulletin (Table 18). The varieties were not tested for a sufficient number of years to show which types were superior. In 1908 the best variety was a 2-rowed brewing barley which has not been identified. Seedsmen of the northeast district often sell seed barley as Two-Row Brewing. Hanna is more often sold under this name than any other variety, and it is probable that the barley grown in 1908 was of this sort. In 1909 Horsford gave the highest yield. This is not consistent with the behavior of Horsford at other points, and it is likely that it would not maintain this high relative position if it were grown for a period of years.

TABLE 18.—*Annual acre yields of varieties of barley grown at the New Hampshire Agricultural Experiment Station (at Durham) in one or both of the years 1908 and 1909*

[Data obtained through the courtesy of the New Hampshire Agricultural Experiment Station]

Variety	C. I. No.	Acre yield (bushels)		Years grown	Average yield (bus.)	Percentage of weighted mean
		1908	1909			
Two-Row Brewing.....	3178	22.9	18.7	2	20.8	110.1
Manchuria.....	244	18.7	18.9	2	18.8	99.5
Oderbrucker.....	537	12.1	-----	1	12.1	81.7
Nepal.....	595	5.4	-----	1	5.4	36.5
Horsford.....	507	-----	29.1	1	29.1	131.1
Caswell.....	3170	-----	26.1	1	26.1	117.6
Black Hull-less.....	596	-----	18.4	1	18.4	82.9

KINGSTON, R. I.

Barleys have not been grown in varietal tests at the Rhode Island Agricultural Experiment Station. Through the courtesy of the station a few yields have been furnished from barley plats utilized in fertilizer and other experiments. In 1910 a plat of Manchuria (C. I. No. 241) yielded 10.5 bushels of grain. The same variety grown in 1913 produced 24.2 bushels and in 1914 12.8 bushels. In 1910 a second plat of Manchuria barley produced 28 bushels. These results show nothing of the relative value of types of barley, but they give an idea of the range of yields which might be expected in Rhode Island.

STORRS, CONN.

The results at Storrs, Conn., were furnished through the courtesy of the Connecticut Agricultural Experiment Station (Table 19). Few varieties were tested for more than one or two years. Barley is not an important crop in Connecticut, and the varietal tests have not been extensive. Two of the best sorts were Heil Hanna (C. I. No. 681) and Franconian (C. I. No. 680). These are both of the Hanna group, and the results agree with those in Maine. French Chevalier (C. I. No. 175) was grown for two years but was inferior to the two Hanna varieties. Oderbrucker (C. I. No. 1273) yielded slightly more than either Hanna. It would seem that varieties of the Hanna and the Manchuria types are best adapted to Connecticut, while Chevalier and Duckbill (C. I. No. 1916) are but slightly inferior.

TABLE 19.—*Annual acre yields of varieties of barley grown at the Connecticut Agricultural Experiment Station (at Storrs) in one or more of the eight years from 1914 to 1921, inclusive*

[Data obtained through the courtesy of the Connecticut Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)							Years grown	Average yield (bus.)	Percentage of weighted mean
		1914	1915	1916	1917	1919	1920	1921			
Heil Hanna 1.....	681	35.4	46.0	42.6	42.6	-----	-----	-----	4	41.7	117.5
Manchuria.....	244	30.4	-----	-----	-----	-----	-----	-----	1	30.4	105.2
French Chevalier.....	175	29.2	44.1	-----	-----	-----	-----	-----	2	36.7	106.4
Himalaya.....	620	21.7	-----	-----	-----	-----	-----	-----	1	21.7	75.1
Duckbill.....	1916	32.9	46.9	24.0	-----	-----	-----	-----	3	34.6	100.9
O. A. C. 21.....	1470	29.6	-----	-----	-----	32.0	35.4	-----	3	32.3	104.5
Success Beardless.....	1808	25.0	34.7	-----	-----	-----	-----	-----	2	29.9	86.7
Franconian.....	680	34.4	47.9	31.7	42.6	-----	-----	-----	4	39.1	110.1
Heil Hanna 4.....	677	21.2	-----	-----	-----	-----	-----	-----	1	21.2	73.4
Champion of Vermont.....	1892	24.8	-----	-----	-----	-----	-----	-----	1	24.8	85.8
Oderbrucker.....	1273	33.1	49.7	-----	46.0	-----	37.8	26.6	5	38.6	109.3
Wisconsin Pedigree.....	835	-----	-----	-----	-----	33.7	38.2	26.9	3	32.9	100.0

ITHACA, N. Y.

The results of varietal tests of barley are available from Cornell University, Ithaca, N. Y., from 1913 to 1921, inclusive. The annual yields are reported in Table 20. It will be seen that most of the varieties are those which have come into experiment stations for testing during recent years. A comparison was made of six groups of barley, of which the Manchuria was the best. The Hybrid group gave yields next to Manchuria, the Hanna was third, and the Chevalier fourth. Neither the Thorpe nor the Horsford group was promising. In the summarized digest of the table a comparison of 12 varieties is shown. The best five of these are Featherston (C. I. No. 1120), Alpha (C. I. No. 959), Hanchamont (C. I. No. 1121), Champion of Vermont (C. I. No. 1892), and Trebi (C. I. No. 936). There is very little difference in the yield of these varieties. Although the Manchuria group ranks highest as a group, there are three barleys of the Hanna type among the five just named. These are Champion of Vermont, Alpha, and Hanchamont, the two latter being of hybrid origin. The barleys of the Hanna type are relatively better adapted to New York conditions than they are to the sections north and east. The Chevalier sorts, which were superior in Prince Edward Island and promising in New England, did not produce high yields here. The Manchuria barleys, on the other hand, grew exceptionally well. The summers in New York are not so hot as in other parts of the humid-spring region. In consequence some varieties succeed which do not seem primarily suited to New York conditions. Trebi (C. I. No. 936), which has done so well under irrigation in southern Idaho, has also yielded well in New York State. New York is probably at the western edge of that section of the North-eastern States where 2-rowed barley can be grown successfully. Because of the warmer summers of the valley lands farther west and in the region of Ottawa to the north, the Manchuria barleys grow best.

TABLE 20.—Annual acre yields of varieties of barley grown at the New York Agricultural Experiment Station at Cornell University (at Ithaca) in one or more of the nine years from 1913 to 1921, inclusive

[Data obtained in cooperation with the New York Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)								Years grown	Average yield (bus.)	Percentage of weighted mean
		1913	1914	1915	1916	1918	1919	1920	1921			
Oderbrucker	836	42.6	47.3	44.9		28.6	32.5	33.7	48.9	7	39.8	109.0
Do	2700	44.9	46.1							2	45.5	112.6
Wisconsin Pedigree	835	46.3	50.6			27.5				3	41.5	116.2
Silver King	890	46.2	46.4	44.9	34.8	26.7	33.4	34.3	38.6	8	38.2	107.6
Manchuria	2949	43.9	47.1	48.2	33.8	25.2				5	39.7	109.4
Do	241	40.2	46.2	42.5	34.8	29.7	33.7			6	37.9	106.8
Do	2944	44.0	44.7	45.7						3	44.8	109.0
Do	2950	40.2	48.1	45.0						3	44.4	108.0
Do	2952	41.3	45.8	43.7						3	43.6	106.1
Do	2946	42.2	47.3	43.0						3	44.2	107.5
Do	2945	41.7	46.8	47.1	33.9	25.8				5	39.1	107.7
Do	244	44.2	48.0	44.5	31.9	28.5				5	40.1	108.5
Do	2951	40.0	47.8	47.8	40.7	24.0				5	40.1	110.5
Baxter	185	43.3	45.9	46.1						3	45.1	109.7
Manchuria	882	42.7	48.5	47.3	35.0	27.0	32.3			6	38.8	109.3
Canadian	2635	32.9	36.6	45.5						3	38.3	93.2
Hanna	319	38.0	35.2	36.3		34.9	39.3	31.5		6	36.1	99.2
Hannchen	531	45.8	47.1	41.5	26.9	31.8	23.2			6	36.1	101.7
Silver King	905	36.2	47.2	45.0						3	42.8	104.1
Hanna	1854	35.7	46.7	47.8	34.8	32.3	36.3	33.1		7	38.1	107.9
New Zealand	2656	35.7	43.1	35.8						3	38.2	92.9
Champion of Vermont	1892	34.0	49.3	44.9	37.2	33.3	37.2	34.7		7	38.7	109.6
Horsford	507	30.8	31.6	38.2						3	33.5	81.5
Evans	892	29.4	30.4	29.9						3	29.9	72.7
Himalaya	620	27.5	38.0	25.3	12.5	10.9	16.7			6	21.8	61.4
Nepal	595	19.8	24.8	18.7		17.5	7.7			5	17.7	48.1
California Prolific	2049	23.1	32.2	27.7						3	27.7	67.4
White Smyrna	910	26.9	34.8	39.4	24.2					4	31.3	82.2
Bohemian	933	28.7	41.3	31.6	18.1					4	29.9	78.5
Chevalier II	200		40.0	34.0	22.1					3	32.0	83.8
Primus	532		31.9	38.0	17.8					3	29.2	76.4
Princess	529		40.9	33.7						2	37.3	88.6
Gold	1145		46.5	43.9	28.0					3	39.5	103.4
Svanhals	187		47.6	38.0	23.9					3	36.5	95.5
Featherston	1120		51.7	57.0	41.2	27.1	34.6	29.8	42.6	7	40.6	115.3
Do	954		31.4	36.4	29.6					3	32.5	85.1
Trebl	936		43.4	51.3	35.3	34.9	36.2	32.3	39.1	7	38.9	110.5
Servian	915		44.4	41.2	32.7					3	39.4	103.1
Sandrel	937		32.9	40.0	31.5					3	34.8	92.1
Manchuria	1178			39.0	29.6					2	34.3	93.2
Alpha	959			52.6	41.0	33.6	38.9	34.3	25.5	6	37.7	110.6
Hanchamont	1121			45.4	40.0	32.6	39.8	40.1	25.7	6	37.3	109.4
Hanna	1122			43.8	26.5					2	35.1	95.4
Frankish	953			45.7	28.6	32.5	27.8			4	33.7	99.4
Manchuria	956			42.1	30.4					2	36.3	98.6
Oderbrucker	957			48.5	32.6	23.8	32.7			4	34.4	101.5
Cortile	1123			38.8	26.7					2	32.7	88.9
Featherston	1118			43.8	31.4					2	37.6	102.2
Luth	972			51.0	33.5	30.0	34.7	34.5		5	36.7	108.6
Red River	973			47.7	35.2	30.7	33.1			4	36.7	108.3
Featherston	1119			52.6	34.2	29.1				3	38.6	111.6
Lion	923			34.2	28.1					2	31.2	84.8
Hanna	906			45.3	28.9	33.9	30.0			4	34.5	101.8
Steigum	907			41.5	32.6	34.7	34.1			4	35.7	105.3
Goldfoil	925			41.4	33.8	30.3				3	35.2	101.7
Australian White	925			37.2	24.0					2	30.6	83.2
Mährische	912			45.4	34.5	33.8	30.6			4	36.1	106.5
Italian	914			34.5	25.1					2	29.8	81.0
Odessa	916			40.0	31.8					2	35.9	97.6
Do	927			47.3	32.4					2	39.9	108.4
Club Mariout	932			38.2	33.6					2	35.9	97.6
Eagle	913			48.0	30.8	28.0				3	35.6	102.9
Peruvian	935			41.0	29.3					2	35.1	95.4
Featherston	911			44.7	31.4	28.0				3	34.7	100.3
Luth	908			41.9	28.2					2	35.1	95.4
Summit	929			39.2	26.2					2	32.7	88.9
Proskowetz	893			40.4	29.1	32.8	31.7	31.8		5	33.2	98.2
Scholey	962			44.9	25.9					2	35.4	96.2
Bolton	922			37.4	27.7					2	32.5	88.3
Horn	926			37.9	25.9					2	31.9	86.7
Oderbrucker	1272					27.0	33.5	38.8	46.0	4	36.3	116.3
Coast	690					31.5	32.4			2	31.9	108.1

TABLE 20.—*Annual acre yields of varieties of barley grown at the New York Agricultural Experiment Station at Cornell University (at Ithaca) in one or more of the nine years from 1913 to 1921, inclusive—Continued*

Variety	C. I. No.	Acre yields (bushels)								Years grown	Average yield (bus.)	Percentage of weighted mean
		1913	1914	1915	1916	1918	1919	1920	1921			
Manchuria.....	2823	---	---	---	---	29.9	33.5	37.5	43.4	4	36.1	115.7
O. A. C. 21.....	1470	---	---	---	---	29.2	31.9	29.6	39.8	4	32.6	104.5
Oderbrucker.....	1526	---	---	---	---	28.0	25.8	---	---	2	26.9	91.2
Wing Pedigree.....	1177	---	---	---	---	22.8	28.3	19.5	---	3	23.5	77.6
Black Hull-less.....	596	---	---	---	---	22.0	20.9	---	---	2	21.5	72.9
Beardless.....	1797	---	---	---	---	14.4	6.8	---	---	2	10.6	35.9
Hybrid.....	2636	---	---	---	---	36.2	28.2	33.9	23.4	4	30.4	97.4
Saul.....	1884	---	---	---	---	31.8	35.5	31.8	28.9	4	32.0	102.6
Charlottetown 80.....	2732	---	---	---	---	31.5	32.9	32.4	---	3	32.2	106.3
Hanna.....	1554	---	---	---	---	30.8	32.1	35.2	---	3	32.0	105.6
Hybrid.....	2637	---	---	---	---	29.5	30.5	29.8	---	3	29.9	98.7
Manchuria.....	2638	---	---	---	---	25.2	33.3	32.0	41.9	4	33.1	106.1
Selection.....	2639	---	---	---	---	---	---	36.4	47.7	2	42.1	120.3
Do.....	2640	---	---	---	---	---	---	35.6	43.4	2	39.5	112.9

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 12 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same year. Under "Data shown" P=Percentage yield and Y=Years comparable]

Variety	C. I. No.	Data shown	Varieties and percentages											
			Charlottetown 80	Svanhals	Champion of Vermont	Hannchen	Featherston	O. A. C. 21	White Smyrna	Nepal	Himalaya	Trebi	Alpha	Hanchamont
Charlottetown 80.....	2732	Y----- P-----	----- -----	(*) -----	3 91.9	2 116.9	3 105.7	3 106.6	(*) -----	2 255.2	2 233.0	3 93.5	3 90.5	3 86.0
Svanhals.....	187	Y----- P-----	(*) -----	(*) -----	3 83.3	3 94.8	3 73.0	3 111.3	3 196.8	2 144.5	3 84.2	3 66.1	2 66.1	2 72.5
Champion of Ver- mont.	1892	Y----- P-----	3 108.8	3 120.0	----- -----	6 109.1	3 98.0	4 116.0	5 132.0	5 224.5	5 180.2	5 101.4	5 93.5	5 94.6
Hannchen.....	531	Y----- P-----	2 85.5	3 105.5	3 91.7	6 109.1	5 80.6	3 90.0	4 128.7	5 214.0	5 165.2	5 84.8	5 74.3	5 78.2
Featherston.....	1120	Y----- P-----	3 94.6	3 136.9	3 102.0	6 124.1	5 102.8	4 152.3	3 248.0	3 204.6	4 104.2	4 102.8	4 103.9	4 103.9
O. A. C. 21.....	1470	Y----- P-----	3 93.8	3 -----	3 86.2	2 111.1	4 97.3	3 -----	3 242.5	4 221.4	4 91.6	3 98.6	2 98.6	2 91.4
White Smyrna.....	910	Y----- P-----	(*) -----	3 89	3 75.8	4 77.7	3 65.6	3 -----	3 159.7	4 121.3	4 75.7	3 67.9	2 74.5	2 74.5
Nepal.....	595	Y----- P-----	2 39.2	2 50.8	5 44.5	4 46.7	4 40.3	2 41.2	3 62.6	5 -----	5 74.7	5 41.4	3 35.1	3 37.3
Himalaya.....	620	Y----- P-----	2 42.9	3 69.2	6 55.5	5 60.5	5 48.9	2 45.2	4 82.4	5 133.8	5 -----	5 51.4	4 39.4	4 41.4
Trebi.....	936	Y----- P-----	3 106.9	3 118.7	3 98.6	5 117.9	7 96.0	4 109.2	3 132.1	4 241.3	5 194.5	5 -----	6 101.4	6 102.5
Alpha.....	959	Y----- P-----	3 110.4	2 151.2	5 107.0	4 134.6	4 97.2	2 101.4	2 147.2	3 285.0	4 254.0	6 98.6	6 -----	6 101.0
Hanchamont.....	1121	Y----- P-----	3 116.3	2 138.0	5 105.7	4 127.9	4 96.3	2 105.9	3 134.3	4 268.3	4 241.3	6 97.6	6 99.0	6 -----

STATE COLLEGE, PA.

The results from Pennsylvania were furnished through the courtesy of the Pennsylvania Agricultural Experiment Station. Some of these were published in the annual reports of the station. Experiments were carried at three points in Pennsylvania. From the central farm, yields are available for the years 1869, 1870, 1872, 1873, 1888, 1889, 1911, and from 1917 to 1921, inclusive. Results from the eastern farm are at hand for only the five years, 1869 to 1873, inclusive, and from the western farm for the years 1870, 1872, and 1873. The annual yields are given in Table 21. While it is very difficult to compare a few varieties grown scatteringly over so many years, it is obvious that both the 2-rowed and Manchuria barleys did well. Good yields were obtained from Charlottetown

80 (C. I. No. 2732) and Michigan (C. I. No. 2782) in recent years. Wisconsin Pedigree (C. I. No. 835) did not yield so high as might be expected in view of the fact that Manchuria (C. I. No. 241) yielded well the two years in which it was grown. At the eastern farm the 2-rowed varieties were promising. The best yielding barley was a winter sort which was grown for two years. The winter barleys usually show up relatively better than their actual value, owing to the fact that they are soon discontinued on account of winterkilling and the final year of failure is not reported. Probstier (C. I. No. 122), a barley of the Hanna group, gave a relatively high yield for the entire five years. Two North African varieties, Coast (C. I. No. 690) and Cape (C. I. No. 557), were entirely out of their region of adaptation and produced very low yields.

The results at the western farm are not consistent. The yields are low except in 1873, when only a few varieties were grown. The Common barley, which was probably of the Manchuria type, was better than the average. While the tests in Pennsylvania are neither extensive nor conclusive, they indicate that Pennsylvania belongs in the same district as New York, that is, the Hanna and Manchuria barleys probably are the ones best adapted. In southeastern Pennsylvania it is probable that the winter barleys would give higher returns and would not winterkill with such frequency as to make their culture unprofitable.

TABLE 21.—*Annual acre yields of varieties of barley grown at the Pennsylvania Agricultural Experiment Station (at State College) in one or more of the 13 stated years in the 53-year period from 1869 to 1921, inclusive*

[Data obtained through the courtesy of the Pennsylvania Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)													Years grown	Aver- age yield (bus.)	Per- centage of weight- ed mean
		1869	1870	1871	1872	1873	1888	1889	1911	1917	1918	1919	1920	1921			
Central farm:																	
Saxonian	2803	32.2	10.7		11.4										3	18.1	120.7
Probstier	122	28.3	11.7												2	20.0	109.3
Common	3112		20.4		15.4	7.3									3	14.4	123.1
Rottingham	3113		8.2												1	8.2	55.0
Golden Melon	411		16.0		6.7										2	11.3	89.0
Chevalier	2802		18.2		8.7										2	13.5	106.3
Brewers De- light	3114		18.9		7.3										2	13.1	103.1
Thanet	2652				11.1	3.3									2	7.2	80.9
Nepal	595						20.0	11.4	0						3	10.5	48.2
Manchuria	241						30.6	31.5							2	31.1	125.9
Black	256							23.1							1	23.1	94.3
Scotch	848							31.9							1	31.9	130.2
Charlottetown 80	2732								50.1	38.1	22.9	38.6	34.8		5	36.9	112.1
Horsford	1271								21.4	40.0	16.7	16.9			4	23.7	88.4
Wing Pedigree	1177									46.1	14.4	28.7			3	29.7	97.7
Michigan	2782												37.4	40.9	2	39.1	109.8
Wisconsin Pedigree	835								26.5				22.7	28.6	3	25.9	85.8
Lion	923												33.7	47.9	2	40.8	114.6
Eastern farm:																	
Coast	690	6.5	10.6	4.9											3	7.3	64.0
Probstier	122	12.2	10.6	9.2	18.0	16.7									5	13.3	126.7
Saxonian	2803	8.3	14.0	9.8	8.1										4	10.1	99.0
Winter (fall sown)	3115	45.0	11.5												2	28.3	193.8
Common																	
Two-Row	3116	18.5													1	18.5	110.8
Nonpareil	3117	9.6		9.7											2	9.7	93.3
Rottingham	3113		14.0	5.0											2	9.5	97.9
Golden Melon	411		12.6	4.9	3.6										3	7.0	81.4
Chevalier	2802		10.6	5.0	1.3										3	5.6	65.1
Brewers De- light	3114		13.0	4.5	1.2										3	6.2	72.1
Common	3112		21.5		4.3	11.0									3	12.3	117.1
Cape	557			2.5											1	2.5	40.3
Thanet	2652				5.9	11.3									2	8.6	106.2
Western farm:																	
Saxonian	2803		12.3		17.4										2	14.9	108.8
Common	3112		10.2		16.0	32.5									3	19.6	117.4
Chevalier	2802		13.3		3.5										2	8.4	61.3
Thanet	2652				18.0	27.5									2	22.7	123.4
Golden Melon	411				17.0										1	17.0	116.4
Brewers De- light	3114				15.5										1	15.5	106.2

WOOSTER, OHIO

Results of barley tests at Wooster, Ohio, were furnished through the courtesy of the Ohio Agricultural Experiment Station from 1904 to 1921. Both winter and spring varieties were grown. As may be seen in Table 22, the winter varieties produced more grain per acre than the spring sorts. None of the Hanna type was included among the spring varieties, probably because they had not done well at the lower altitudes in the Mississippi Valley. Two 2-rowed barleys—Primus (C. I. No. 532) and Princess (C. I. No. 529)—were grown. These gave only fair yields and are probably not so well adapted to the section as even Hanna would be. Two varieties of the Manchuria type—Oderbrucker (C. I. No. 836) and Wisconsin Pedigree (C. I. No. 835)—were grown in most of the years since 1912. Wisconsin Pedigree gave higher yields in six of the nine years in which both of the varieties were grown. The barleys of the Tennessee Winter group were the best of the winter varieties grown. Ohio Winter (C. I. No. 2033), which is a locally adapted form of the Tennessee Winter group, was the best variety. Orel (C. I. No. 351), a 2-rowed barley, gave a very good yield but was not equal to the Ohio Winter in the only year in which it was grown. It is probable that in sections of Ohio the winter barleys are the better commercial group. In northwestern Ohio spring barleys are likely to be more dependable. Those of the Manchuria group are the best adapted of the spring varieties. They endure hot, moist weather at ripening time better than most other sorts and are therefore well adapted to culture in the upper Mississippi Valley.

TABLE 22.—*Annual acre yields of varieties of barley grown at the Ohio Agricultural Experiment Station (at Wooster) in one or more of the 18 years from 1904 to 1921, inclusive*

[Data obtained through the courtesy of the Ohio Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)									
		1904	1905	1906	1908	1909	1910	1911	1912	1913	1914
Manchuria.....	2778	38.5	39.3	41.7	-----	-----	-----	-----	-----	-----	-----
Ohio Beardless.....	231	25.9	18.3	32.7	-----	-----	-----	-----	-----	-----	-----
Black Hull-less.....	596	41.9	22.1	30.1	-----	-----	-----	-----	-----	-----	-----
Oderbrucker.....	836	-----	-----	-----	44.1	37.8	44.7	14.2	34.4	14.4	27.4
Primus.....	532	-----	-----	-----	36.6	9.0	34.2	-----	-----	-----	-----
Princess.....	529	-----	-----	-----	34.9	24.6	34.1	-----	-----	-----	-----
Wisconsin Pedigree.....	835	-----	-----	-----	-----	-----	-----	-----	28.0	13.9	28.7
Winter barleys:											
Texas Winter.....	554	-----	-----	-----	-----	-----	51.0	-----	-----	-----	53.6
Ohio Winter.....	2033	-----	-----	-----	-----	-----	63.0	-----	-----	-----	46.9
Maryland Winter.....	518	-----	-----	-----	-----	-----	46.3	-----	-----	-----	43.7
Tennessee Winter.....	257	-----	-----	-----	-----	-----	57.3	-----	-----	-----	40.1
Indiana Winter.....	2039	-----	-----	-----	-----	-----	-----	-----	-----	-----	37.5
Wakamatsu.....	579	-----	-----	-----	-----	-----	-----	-----	-----	-----	34.4
Orel.....	351	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Michigan Winter.....	2036	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Variety	C. I. No.	Acre yields (bushels)							Years grown	Average yield (bus.)	Percentage of weighted mean
		1915	1916	1917	1918	1919	1920	1921			
Manchuria.....	2778	-----	-----	-----	-----	-----	-----	-----	3	39.8	123.2
Ohio Beardless.....	231	-----	-----	-----	-----	-----	-----	-----	3	25.6	79.3
Black Hull-less.....	596	-----	-----	-----	-----	-----	-----	-----	3	31.4	97.2
Oderbrucker.....	836	-----	33.7	42.2	59.4	21.8	30.0	24.2	13	32.9	83.9
Primus.....	532	-----	-----	-----	-----	-----	-----	-----	3	26.6	79.9
Princess.....	529	-----	-----	-----	-----	-----	-----	-----	3	31.2	93.7
Wisconsin Pedigree.....	835	-----	37.7	43.4	59.0	23.9	30.7	27.0	9	32.5	81.9
Winter barleys:											
Texas Winter.....	554	61.2	58.3	63.5	42.5	54.9	13.5	-----	8	49.8	108.5
Ohio Winter.....	2033	67.7	54.2	74.0	45.9	47.3	16.1	-----	8	51.9	113.1
Maryland Winter.....	518	62.0	46.9	74.0	-----	-----	-----	-----	5	54.6	109.4
Tennessee Winter.....	257	62.0	-----	-----	-----	-----	-----	-----	3	53.1	109.5
Indiana Winter.....	2039	56.8	-----	-----	-----	-----	-----	-----	2	47.1	95.7
Wakamatsu.....	579	54.7	-----	-----	-----	-----	-----	-----	2	44.5	90.4
Orel.....	351	62.0	-----	-----	-----	-----	-----	-----	1	62.0	101.8
Michigan Winter.....	2036	-----	-----	-----	39.1	55.4	26.6	-----	3	40.4	107.2

EAST LANSING, MICH.

The yields from East Lansing, Mich., given in Table 23, were furnished through the courtesy of the Michigan Agricultural Experiment Station. They cover the six years from 1916 to 1921, inclusive. The results are best considered in two periods. During the three years, 1916 to 1918, inclusive, the yields of 13 varieties are reported. Hannchen (C. I. No. 531) was the best of the varieties grown for the three years, Chevalier II (C. I. No. 200) and Princess (C. I. No. 529) were equal in yield, and Svanhals (C. I. No. 187) ranked fourth. Gold (C. I. No. 1145) was highly productive for the two years in which it was grown. Oderbrucker (C. I. No. 1273), which was grown in 1917 and 1918, was the only variety of the Manchuria type included. In 1917 it was the second highest variety. In 1918 it was relatively inferior. During the years from 1919 to 1921, inclusive, a number of selections were grown. Since Gold is a Swedish variety, originating from a single plant, it is likely that the three selections of this variety were identical. Oderbrucker (C. I. No. 1273) was supposed to be pedigreed. The selections from Crawford, a local Michigan barley, were uniformly good. The selections from Gold were slightly inferior to Crawford, while Charlottetown 80 (C. I. No. 2732) was slightly less promising than Gold. The Oderbrucker and Manchuria varieties were slightly less than average in yield. In that section of Michigan in which East Lansing is located it is apparent that the Manchuria barleys are not so dominant as they were in Ohio, Indiana, or Ontario. Conditions in this section of Michigan are perhaps more like those in New York State, where barleys of the Hanna type are about on a par with those of the Manchuria type. However, there are doubtless sections in Michigan in which the Manchuria barleys would give higher yields than the Hanna barleys.

TABLE 23.—*Annual acre yields of varieties of barley grown at the Michigan Agricultural Experiment Station (at East Lansing) in two or three of the six years from 1916 to 1921, inclusive*

[Data furnished through the courtesy of the Michigan Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)						Years grown	Average yield (bus.)	Percentage of weighted mean
		1916	1917	1918	1919	1920	1921			
Franconian	680	26.9	21.8	31.5	-----	-----	-----	3	26.7	93.4
Svanhals	187	31.0	31.7	33.5	-----	-----	-----	3	32.1	112.2
Chevalier II	200	33.5	30.6	38.6	-----	-----	-----	3	34.2	119.6
Princess	529	35.4	29.1	38.1	-----	-----	-----	3	34.2	119.6
Hannchen	531	44.6	18.3	35.4	-----	-----	-----	3	32.8	114.7
Do	531	42.9	27.6	41.3	-----	-----	-----	3	37.3	130.4
Lion	923	23.3	29.0	30.8	-----	-----	-----	3	27.7	96.9
Semibarbless	2809	21.7	21.3	16.3	-----	-----	-----	3	19.8	69.2
Arlington Awnless	702	18.4	11.3	14.5	-----	-----	-----	3	14.7	51.4
Arizona	2808	30.1	20.7	25.9	-----	-----	-----	3	25.6	89.5
Gold	1145	-----	38.1	33.3	-----	-----	-----	2	35.7	128.9
Charlottetown 80	2732	-----	10.1	28.8	-----	-----	-----	2	19.5	70.4
Oderbrucker	1273	-----	33.4	29.1	-----	-----	-----	2	31.3	113.0
Gold	1145	-----	-----	38.0	38.5	29.9	-----	3	35.5	107.6
Do	1145	-----	-----	31.0	38.2	33.8	-----	3	34.3	103.9
Do	1145	-----	-----	27.3	38.1	29.9	-----	3	31.8	96.4
Crawford	2810	-----	-----	27.9	38.9	32.4	-----	3	33.1	100.3
Do	3136	-----	-----	34.3	38.7	31.8	-----	3	34.9	105.8
Do	3137	-----	-----	29.4	37.5	41.3	-----	3	36.1	109.4
Do	3138	-----	-----	39.0	37.9	40.9	-----	3	39.3	119.1
Do	3139	-----	-----	35.2	36.7	33.5	-----	3	35.1	106.4
Charlottetown 80	2732	-----	-----	27.6	37.3	36.1	-----	3	33.7	102.1
Oderbrucker	1273	-----	-----	27.3	38.3	32.4	-----	3	32.7	99.1
Do	1273	-----	-----	31.5	30.9	32.5	-----	3	31.6	95.8
Do	1273	-----	-----	31.8	32.1	33.0	-----	3	32.3	97.9
Do	1273	-----	-----	25.3	30.7	36.1	-----	3	30.7	93.0
Do	1273	-----	-----	28.9	34.7	31.1	-----	3	31.6	95.8
Manchuria	3140	-----	-----	27.6	30.8	29.7	-----	3	29.4	89.1
Do	3141	-----	-----	37.3	34.0	30.8	-----	3	34.0	103.0
Do	3142	-----	-----	30.2	34.2	28.3	-----	3	30.9	93.6
Do	3143	-----	-----	24.9	29.9	28.0	-----	3	27.6	83.6

TABLE 24.—*Annual acre yields of varieties of barley grown at the Indiana Agricultural Experiment Station (at La Fayette) in some or all of the 17 years from 1905 to 1921, inclusive*

[Data obtained through the courtesy of the Indiana Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)									
		1905	1906	1907	1908	1909	1910	1911	1912	1913	1914
Success Beardless.....	1808	20.6	21.7	16.2	21.3	30.5	45.7	20.9	37.8	7.3	6.4
Hanna.....	203	19.6	24.8	8.4	17.7	33.1	-----	-----	-----	-----	-----
Silver Beardless.....	641	17.5	24.0	11.8	14.4	32.7	-----	-----	-----	-----	-----
Silver King.....	890	22.3	33.6	21.2	19.0	46.3	-----	-----	-----	-----	-----
Manchuria.....	241	20.4	30.3	10.0	13.4	47.1	-----	-----	-----	-----	-----
Highland Chief.....	883	20.0	20.9	9.0	16.3	37.6	-----	-----	-----	-----	-----
Black Hull-less.....	596	16.8	18.6	15.0	14.4	31.3	-----	-----	-----	-----	-----
Nepal.....	595	16.4	-----	9.3	13.2	35.4	-----	-----	-----	-----	-----
Great Beardless.....	2779	20.6	23.6	17.3	15.7	32.2	-----	-----	-----	-----	-----
Oderbrucker.....	2780	-----	-----	-----	21.1	41.3	42.0	18.6	37.5	9.0	3.2
Manchuria.....	244	-----	-----	-----	19.7	43.7	43.2	16.3	34.4	5.2	5.2
O. A. C. 21.....	1470	-----	-----	-----	18.8	48.0	41.8	19.9	55.0	7.1	5.9
Garton 986.....	645	-----	-----	-----	-----	31.3	-----	1	4.6	2.7	4.9
Reed Triumph.....	889	-----	-----	-----	-----	32.3	-----	5.1	38.4	5.8	6.2
Sexradigt.....	607	-----	-----	-----	-----	-----	-----	17.6	46.2	8.7	6.3
Hannchen.....	531	-----	-----	-----	-----	-----	-----	17.8	55.5	8.7	8.1
Primus.....	532	-----	-----	-----	-----	-----	-----	3.8	-----	-----	0
Princess.....	529	-----	-----	-----	-----	-----	-----	4.9	56.1	3.7	6.3
Chevalier II.....	200	-----	-----	-----	-----	-----	-----	9.1	42.0	5.4	2.7
Stoeckinger.....	2781	-----	-----	-----	-----	-----	-----	-----	-----	8.1	5.1
Svanhals.....	187	-----	-----	-----	-----	-----	-----	-----	-----	-----	1.9
Gold.....	1145	-----	-----	-----	-----	-----	-----	-----	-----	-----	2.3
Wing Pedigree.....	1177	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Horsford.....	1271	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Charlottetown 80.....	2732	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Wisconsin Pedigree.....	835	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Manchuria.....	2330	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Lion.....	923	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Michigan.....	2782	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Oderbrucker.....	1529	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Gatami.....	575	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Featherston.....	1120	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Variety	C. I. No.	Acre yields (bushels)							Years grown	Average yield (bus.)	Percentage of weighted mean
		1915	1916	1917	1918	1919	1920	1921			
Success Beardless.....	1808	20.5	28.7	48.8	42.5	17.5	16.2	25.0	17	25.1	104.1
Hanna.....	203	-----	-----	-----	-----	-----	-----	-----	5	20.7	89.2
Silver Beardless.....	641	-----	-----	-----	-----	-----	-----	-----	5	20.1	86.6
Silver King.....	890	-----	-----	-----	-----	-----	-----	-----	5	28.5	122.8
Manchuria.....	241	-----	-----	-----	-----	-----	-----	-----	5	24.2	104.3
Highland Chief.....	883	-----	-----	-----	-----	-----	-----	-----	5	20.8	89.7
Black Hull-less.....	596	-----	-----	-----	-----	-----	-----	-----	5	19.2	82.8
Nepal.....	595	-----	-----	-----	-----	-----	-----	-----	4	18.6	80.9
Great Beardless.....	2779	-----	-----	-----	-----	-----	-----	-----	5	21.9	94.4
Oderbrucker.....	2780	-----	-----	-----	-----	-----	-----	-----	7	24.7	118.2
Manchuria.....	244	-----	-----	-----	-----	-----	-----	-----	7	24.0	114.8
O. A. C. 21.....	1470	-----	-----	-----	-----	-----	-----	-----	7	28.1	134.4
Garton 986.....	645	-----	-----	-----	-----	-----	-----	-----	6	12.7	58.8
Reed Triumph.....	889	14.2	29.2	65.6	-----	-----	-----	-----	8	24.6	103.8
Sexradigt.....	607	18.1	33.9	51.8	-----	-----	-----	-----	7	26.1	116.0
Hannchen.....	531	17.9	35.2	50.5	-----	-----	-----	-----	7	27.7	123.1
Primus.....	532	12.8	23.4	45.7	-----	-----	-----	-----	5	17.1	76.3
Princess.....	529	24.8	33.6	44.9	-----	-----	-----	-----	7	24.9	110.7
Chevalier II.....	200	15.7	25.3	38.8	-----	-----	-----	-----	7	19.9	88.4
Stoeckinger.....	2781	16.0	31.3	38.3	-----	-----	-----	-----	5	19.8	93.0
Svanhals.....	187	16.8	30.7	45.8	-----	-----	-----	-----	4	23.8	96.0
Gold.....	1145	16.0	32.7	48.8	-----	-----	-----	-----	4	24.9	100.4
Wing Pedigree.....	1177	-----	-----	46.3	42.7	24.3	13.4	23.3	5	30.0	92.0
Horsford.....	1271	-----	-----	45.8	32.5	20.1	15.0	21.0	5	26.9	82.5
Charlottetown 80.....	2732	-----	-----	53.3	45.5	14.6	22.6	22.1	5	31.6	96.9
Wisconsin Pedigree.....	835	-----	-----	-----	38.0	18.6	21.0	28.7	4	26.6	101.1
Manchuria.....	2330	-----	-----	-----	-----	21.4	25.7	32.2	3	26.4	110.9
Lion.....	923	-----	-----	-----	-----	-----	26.7	32.2	2	29.5	118.0
Michigan.....	2782	-----	-----	-----	-----	-----	29.2	21.9	2	25.5	102.0
Oderbrucker.....	1529	-----	-----	-----	-----	-----	24.1	33.2	2	28.7	114.8
Gatami.....	575	-----	-----	-----	-----	-----	13.3	32.5	2	22.9	91.6
Featherston.....	1120	-----	-----	-----	-----	-----	-----	46.2	1	46.2	159.9

LA FAYETTE, IND.

The annual yields of the barley varieties grown at Purdue University, La Fayette, Ind., were furnished through the courtesy of the Indiana Agricultural Experiment Station. These yields, which extend over a period of 17 years, are reported in Table 24. Success Beardless (C. I. No. 1808) was the only variety carried in the test for the entire period. Its yield was 104 per cent of that of the weighted mean of all varieties. Owing to the fact that many varieties were discarded after the harvests in 1909, 1914, and 1917, it is easier to discuss their relative value by periods. In the years 1905 to 1909, inclusive, Silver King (C. I. No. 890) and Manchuria (C. I. No. 241) were the best of the nine varieties grown. From 1908 until 1914 only three varieties, all of which were fully comparable, belong to the Manchuria group. Of these, O. A. C. 21 (C. I. No. 1470) was the best. Hannchen (C. I. No. 531) produced the highest yield of the nine varieties discarded after 1917. Unfortunately there were no barleys of the Manchuria group in the test for the years 1915, 1916, and 1917. Hannchen was in the test for four of the years in which O. A. C. 21 was grown and was better than O. A. C. 21 in three of these years. The difference in yield, however, was very small.

Hannchen was not grown in 1919, 1920, and 1921, when varieties of the Manchuria group were included in the experiment. During the years 1917 to 1921, inclusive, 10 barleys were tested one or more years. Charlottetown 80 (C. I. No. 2732) was the most promising of those grown for the full five years. Manchuria (C. I. No. 2330), a selection made at the Minnesota Agricultural Experiment Station, was the best of the five varieties that were carried for three years. Lion (C. I. No. 923), introduced in 1920, gave relatively high yields for two years, while Featherston (C. I. No. 1120) was by far the best variety in 1921, the only year in which it was grown. These results indicate that the Manchuria barleys are the ones best adapted to this section of Indiana. This agrees with the results in the neighboring States and with what is known of the ecological adaptation of varieties. The Manchuria barleys are adapted to climates where the ripening season is warm and humid without excessive heat. Barleys of the Hanna and Chevalier groups grow best in cooler regions. That the Charlottetown 80 has done well not only at La Fayette but at many other points in the Mississippi Valley is proof that this selection differs materially from the original Chevalier of England. Primus (C. I. No. 532), a barley of the Thorpe type, was very low in productivity. This variety must have a cool ripening season before good yields are secured. It is of especial interest that high yields were obtained from Success Beardless (C. I. No. 1808), a barley of the Horsford group. It was the only variety carried for the entire 17 years. In each of the four periods it was one of the better sorts.

TABLE 25.—*Annual acre yields of varieties of barley grown at the Illinois Agricultural Experiment Station (at Urbana) in part or all of the seven years from 1915 to 1921, inclusive*

[Data obtained through the courtesy of the Illinois Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)							Years grown	Average yield (bus.)	Percentage of weighted mean
		1915	1916	1917	1918	1919	1920	1921			
Oderbrucker.....	836	53.3	57.1	72.7	43.8	34.2	31.8	48.9	7	48.8	122.6
Horsford.....	507	44.6	43.7	46.7	39.0	26.9	30.1	-----	6	38.5	99.0
Nepal.....	595	17.7	26.8	24.4	14.4	-----	-----	-----	4	20.8	50.6
Montana.....	3118	-----	28.0	39.7	13.8	-----	-----	-----	3	27.2	65.4
Wisconsin Pedigree.....	835	-----	55.4	71.7	46.2	36.3	35.4	53.0	6	49.7	124.6
Lion.....	923	-----	-----	-----	-----	-----	-----	39.0	1	39.0	83.0

URBANA, ILL.

The Illinois Agricultural Experiment Station at Urbana furnished data as to the yields of the barley varieties tested there for the seven years from 1915 to 1921, inclusive. These are reported in Table 25. Six varieties were tested during this period. Of these, Oderbrucker (C. I. No. 836) was in the test for the full seven years. Unfortunately few types of barley were included. The only 2-rowed barley was Montana (C. I. No. 3118), which was probably a Cheva-

lier. The Chevalier barleys, perhaps, are the least suited of the 2-rowed forms for Illinois conditions, except possibly those of the Thorpe group. Of all the varieties in the test those of the Manchuria group were by far the best. Oderbrucker (C. I. No. 836) and Wisconsin Pedigree (C. I. No. 835) were the only two varieties which yielded more than the weighted mean of all varieties for the years grown. Wisconsin Pedigree was somewhat better than Oderbrucker. Horsford (C. I. No. 507) gave a higher relative yield than at most places. With the exception of the two Oderbrucker strains, however, the competing varieties were not well adapted.

DE KALB, ILL.

The reports of yields at De Kalb, Ill., shown in Table 26 were furnished through the courtesy of the Illinois Agricultural Experiment Station. Varieties were grown at this point during the same period as at Urbana. Eight varieties were tested at De Kalb. As was the case at Urbana, barleys of the Manchuria group were markedly superior. In three of the five years in which both varieties were grown at De Kalb Oderbrucker (C. I. No. 836) was better than Wisconsin Pedigree (C. I. No. 835). Lion (C. I. No. 923) gave very good yields for the two years grown. This is a black, 6-rowed, smooth-awned variety distributed by the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul. The strain grown in Illinois was from a plant selection made by the Michigan Agricultural Experiment Station from Lion (C. I. No. 923). It was sent out by the Michigan station as Black Barbless. The Lion sent to Michigan apparently had become accidentally mixed or hybridized in Minnesota. The plant selection made in Michigan seems identical with the Lion selection made in Minnesota; at least it differs so little from the original variety that, for the purpose of this description, it is carried under the original name. The Montana barley (C. I. No. 3118) produced higher yields at De Kalb than at Urbana. Its yields were as high as those of Wisconsin Pedigree in 1915 only. In the three following years it was much inferior.

From the results at Urbana and De Kalb it is apparent that northern Illinois is definitely in the district where barleys of the Manchuria type are especially adapted. This agrees with the results in the neighboring States, with the exception of parts of Michigan. In Ohio, Indiana, Illinois, and Wisconsin the barleys of the Manchuria type unquestionably are superior as far as the varieties now known have been compared. Hooded barleys of the Horsford type have given fair yields but are undoubtedly inferior to the Manchuria and Oderbrucker barleys in yielding capacity. Lion has shown some promise in Illinois and in neighboring States. Hybrids of this variety with varieties of the Manchuria group have been made at the Minnesota, Wisconsin, and other stations, and it is possible that smooth-awned barleys which are white in color may prove valuable in this section.

TABLE 26.—*Annual acre yields of varieties of barley grown at the Soil Experiment Field, De Kalb, Ill., in part or all of the seven years from 1915 to 1921, inclusive*

[Data obtained through the courtesy of the Illinois Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)							Years grown	Average yield (bus.)	Percentage of weighted mean
		1915	1916	1917	1918	1919	1920	1921			
Michigan	2782	24.9	-----	-----	-----	-----	-----	-----	1	24.9	70.5
Montana	3118	50.0	34.6	32.9	30.3	-----	-----	-----	4	36.9	86.0
Nepal	595	21.4	28.2	12.5	32.9	-----	-----	-----	4	23.7	55.2
Wisconsin Pedigree	835	44.7	47.2	61.6	77.1	27.2	51.3	37.2	7	49.5	118.7
Horsford	507	-----	-----	42.0	52.0	23.5	47.8	-----	4	41.3	91.2
Oderbrucker	836	-----	-----	63.8	73.3	34.3	51.8	36.0	5	51.8	118.8
Lion	923	-----	-----	-----	-----	55.7	36.7	36.7	2	46.2	104.8
Silver King	890	-----	-----	-----	-----	-----	36.7	36.7	1	36.7	100.0

MADISON, WIS.

Wisconsin is one of the more important barley-producing States of the United States. The testing of varieties at the Wisconsin Agricultural Experiment Station was begun at an early date. The yields shown in Table 27 were furnished through the courtesy of this station, in whose reports some of the earlier yields

were published. An organization of farmers interested in the testing and growing of good barley in Wisconsin assisted in establishing farm culture of pedigreed varieties in the State, and this acted as an incentive to experiment-station workers to produce better varieties. Not only has very important work been done here, but the Wisconsin station has been a distributing center of varieties, especially those of the Manchuria group. Many of these have been sent out under the names of Oderbrucker, Wisconsin Pedigree, and Silver King. In the earlier years of the station a Manchuria barley was widely distributed, often under the spelling "Manshury." This has been changed to Manchuria, principally to make it conform with varieties which have been widely tested at other stations and which also came from Manchuria. The history of this variety is given under C. I. No. 241 in a special paragraph elsewhere in this bulletin.

Six groups of barley were compared, of which the Manchuria group was outstanding. The Hanna group, which was next in point of yield, was much inferior to the Manchuria. The Thorpe, Chevalier, Horsford, and Hybrid groups follow in the order named.

It will be seen in the summarized digest of Table 27, in which eight varieties are compared, that barleys of the Manchuria group are as outstanding in their performance as was the group itself. The best variety of the eight compared was Oderbrucker (C. I. No. 2700), though Manchuria (C. I. No. 241) was practically equal in point of yield. The best of the other types compared was Svanhals (C. I. No. 187). Its yield was only 81 per cent of that of Oderbrucker. Hannchen (C. I. No. 531), which usually gives good yields where barleys of the Hanna type are adapted, yielded only 61 per cent of that of Oderbrucker during the five years in which both were in the test.

It is obvious that among so many varieties it is impossible to compare in detail all of the important ones. In recent years several varieties have been tested which have shown promise elsewhere. Charlottetown 80 (C. I. No. 2732) was grown in the years 1917, 1918, and 1920. For these years its average yield was 50.6 bushels, which was 102 per cent of the weighted mean of all varieties. Manchurian (C. I. No. 739) was grown in 1907 and in the seven years from 1915 to 1921, inclusive. Its yield was 115 per cent of the weighted mean. Gold (C. I. No. 1145) was grown in the six years from 1915 to 1920, inclusive. Its yield was 91 per cent of the weighted mean. For the seven years from 1914 to 1920, inclusive, O. A. C. 21 produced an average yield of 49.2 bushels, which was 107 per cent of the weighted mean. Of the varieties introduced into the test since 1912 O. A. C. 21 (C. I. No. 1470) seems to be especially promising, while Charlottetown 80 deserves consideration when it is compared with 2-rowed barleys.

The results at Madison show that barleys of the Manchuria-Oderbrucker type are especially suited to growing in Wisconsin. Although varieties of other types have been tested as extensively, they have not given sufficient promise to justify carrying many of them. Several varieties from the Wisconsin station, such as Wisconsin Pedigree (C. I. No. 835), Oderbrucker (C. I. No. 836), and Silver King (C. I. No. 890), have been widely tested over the United States and at many stations have given very high yields. These varieties are discussed elsewhere.

[illegible]

TABLE 27.—Annual acre yields of varieties of barley grown at the Wisconsin Agricultural Experiment Station (at Madison) in one or more of the years from 1871 to 1921, inclusive—Continued

Variety	C. I. No.	Acre yields (bushels)																	Years grown	Average yield (bus.)	Percent- age of weighted mean
		1906	1907	1908	1909	1910	1911	1912	1913	1914	1915	1916	1917	1918	1919	1920	1921				
		2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1				
Garton.....	263	13.0																1	13.0	51.0	
Do.....	264	10.8																1	10.8	42.4	
Do.....	266	16.1	3.7	26.2														3	13.3	58.4	
Odessa.....	182	19.4	10.8	38.9														3	23.0	87.8	
Royal.....	1252	19.0	14.1	31.2														3	21.4	81.7	
Claude.....	1557	17.2																1	17.2	67.5	
Mensury.....	2657	35.7	27.5	36.2														3	33.1	126.3	
Serradigt.....	607		13.7															1	13.7	72.5	
Gutckorn.....	606		12.5	57.0														2	34.7	130.5	
Gotland.....	2749		14.7															1	14.7	77.8	
Manchurian.....	739		26.5									66.7	43.5	45.6	60.6	25.4	46.2	8	42.3	115.3	
Hanna.....	319		8.9															1	8.9	47.1	
Canadian Lake Shore.....	2750		20.2															1	20.2	154.5	
Coast.....	680		26.5															1	26.5	140.2	
Standwell.....	584		19.5															1	19.5	103.2	
Mansfield.....	2241		20.0															1	20.0	105.8	
Invincible.....	500		8.3															1	8.3	43.9	
Sidney.....	178		20.0															1	20.0	105.8	
Chevalier.....	156		7.9															1	7.9	41.8	
Garton 936.....	2752		42.9		22.9	54.1	35.4	52.0				39.3	59.3	42.8	53.5	60.0	23.5	2	32.9	137.1	
Manchuria.....	1271			42.5	40.4	40.8	41.2	41.2					48.3					13	44.4	107.5	
Oderbrucker.....	1274			38.6	46.7		29.5	34.7	45.0									6	42.8	103.1	
Do.....	1273			31.1	40.8		32.0	35.4	49.1									5	36.2	92.3	
Silver King.....	1276			37.6			32.0	35.4	49.1									4	38.5	96.3	
Do.....	905			34.0	37.5	35.8	35.4	49.5				43.0	62.3	36.3	47.5	65.0		13	44.6	103.5	
Oderbrucker.....	1529			35.3	37.5	45.0	35.4	47.0				38.4	58.3	38.3	47.1	65.4	28.3	10	42.6	103.1	
Golden Queen.....	2753			27.2	17.9	42.3	35.4	50.8										5	34.8	88.8	
Silver King.....	2754			26.8	32.1	44.4	40.8	47.9										5	38.4	98.0	
Oderbrucker.....	1272			45.0	42.5	41.6	44.1					50.4	38.8	50.4	57.3	23.7	50.2	11	43.1	102.1	
Manchuria.....	2750			45.8	43.5	40.0	46.6					61.7	39.7					6	46.2	108.5	
Golden Queen.....	1511			29.4		42.5	41.6	49.1										1	29.4	84.2	
Manchuria.....	2758			33.3		35.4	45.8	51.2										4	41.6	103.5	
Do.....	2760			31.3		45.0	35.4	48.3										4	40.9	101.7	
Do.....	2762					31.2	45.8	43.7				58.0	42.7	53.7	55.3			8	45.6	102.7	
Wisconsin Pedigree 20.....	2761					50.0	31.6	18.7										3	40.2	96.2	
Manchuria.....	2764																	3	33.4	79.9	

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF EIGHT REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages							
			Chevalier II	Primus	Svan-hals	Hanna	Hann-chen	Oder-brucker	Man-churia	Nepal
Wisconsin Pedigree 18.....	2763	-----	-----	59.3	39.6	-----	-----	26.6	4	47.8
Wisconsin Pedigree IV.....	2765	-----	-----	50.0	39.6	-----	-----	-----	4	43.5
O. A. C. 21.....	1470	-----	41.5	58.1	42.1	48.4	63.1	44.6	7	49.2
Gold.....	1145	-----	-----	40.3	32.8	52.5	72.9	30.8	6	42.2
Wisconsin Pedigree 35.....	2766	-----	-----	61.4	41.1	-----	-----	-----	2	51.3
Arlington Awnless.....	702	-----	-----	24.1	-----	-----	-----	-----	2	43.7
Charlotteown 80.....	2732	-----	-----	-----	57.5	50.0	44.4	-----	3	43.7
Wing Pedigree.....	1177	-----	-----	-----	35.3	38.9	31.8	21.6	3	42.4
Success.....	2807	-----	-----	-----	-----	-----	29.2	-----	1	31.2
Korsdyg.....	918	-----	-----	-----	-----	-----	-----	-----	1	21.2
Reed Triumph.....	889	-----	-----	-----	-----	-----	-----	20.2	1	20.2

Variety	C. I. No.	Data shown	Varieties and percentages							
			Chevalier II	Primus	Svan-hals	Hanna	Hann-chen	Oder-brucker	Man-churia	Nepal
Chevalier II.....	200	{ Years comparable	-----	4	4	2	4	4	3	1
		{ Percentage yield	-----	98.1	67.5	103.8	86.2	53.3	54.3	170.5
Primus.....	532	{ Years comparable	4	-----	5	3	5	5	4	1
		{ Percentage yield	101.9	-----	85.0	104.6	97.3	59.5	58.3	141.5
Svanhals.....	187	{ Years comparable	4	5	-----	6	5	8	7	3
		{ Percentage yield	148.1	117.6	-----	101.1	114.5	80.8	82.4	192.7
Hanna.....	203	{ Years comparable	2	3	6	-----	3	6	5	2
		{ Percentage yield	96.3	95.6	98.9	-----	122.8	83.4	79.6	172.3
Hannchen.....	531	{ Years comparable	4	5	5	3	-----	5	4	1
		{ Percentage yield	116.1	102.7	87.4	81.4	-----	61.2	65.7	183.0
Oderbrucker.....	2700	{ Years comparable	4	5	8	6	5	-----	9	4
		{ Percentage yield	187.7	168.0	123.8	119.9	163.5	-----	102.0	150.2
Manchuria.....	241	{ Years comparable	3	4	7	5	4	9	-----	8
		{ Percentage yield	184.1	171.4	121.3	125.6	152.3	98.0	-----	149.1
Nepal.....	595	{ Years comparable	1	3	2	2	1	4	8	-----
		{ Percentage yield	58.7	70.7	51.9	58.0	54.6	66.6	67.0	-----

TABLE 28.—Annual acre yields of varieties of barley grown at the Minnesota Agricultural Experiment Station (at St. Paul) in one or more of the 29 years from 1893 to 1921, inclusive—Continued

Variety	C. I. No.	Acre yields (bushels)																				Years grown	Average yield (bus.)	Per- centage of weight- ed mean		
		1893	1894	1895	1896	1897	1898	1906	1907	1908	1909	1910	1911	1912	1913	1914	1915	1916	1917	1918	1919				1920	1921
Manchuria	2867																							2	46.3	117.8
Do	2868																							3	43.8	110.9
Do	2869																							3	45.7	115.7
Do	2870																							3	41.7	105.6
Do	2871																							3	43.7	110.9
Do	2872																							14	48.5	123.4
Do	2873																							2	48.5	123.4
Do	2874																							2	44.7	113.7
Do	2875																							2	42.1	107.1
Do	2876																							2	40.9	104.1
Do	2877																							2	33.1	84.7
Do	2878																							2	33.9	86.3
Do	2879																							3	38.9	99.5
Do	2880																							2	39.7	102.6
Do	2881																							1	15.4	40.3
Do	2882																							6	36.0	86.7
Do	2883																							4	40.9	114.9
Do	2884																							4	47.1	132.3
Do	2885																							4	43.0	120.8
Do	2886																							4	42.0	118.0
Do	2887																							4	45.2	127.0
Do	2888																							2	36.5	95.5
Do	2889																							1	26.7	69.9
Do	2890																							2	33.4	87.4
Do	2891																							5	43.5	105.8
Do	2892																							3	32.5	92.6
Do	2893																							3	32.0	91.2
Do	2894																							3	34.9	99.4
Do	2895																							3	32.4	92.3
Do	2896																							3	27.4	78.1
Do	2897																							3	28.6	81.5
Do	2898																							3	26.0	74.1
Do	2899																							2	35.8	93.7
Do	2900																							1	23.3	61.0
Do	2901																							5	34.3	83.5
Do	2902																							13	42.6	101.2
Do	2903																							3	34.2	97.4
Do	2904																							1	30.5	79.8

Chevalier	156	28.8	11.7	33.4	36.8	35.0	20.0	23.3	50.1	28.8	11.7	75.5
Tennessee Winter	257	33.4	36.8	35.0	20.0	23.3	50.1	28.8	11.7	75.5	30.0	87.5
Canadian Lake	2750	36.8	35.0	20.0	23.3	50.1	28.8	11.7	75.5	30.0	87.5	87.5
Shore	2750	36.8	35.0	20.0	23.3	50.1	28.8	11.7	75.5	30.0	87.5	87.5
Gottland	739	37.0	36.7	34.3	21.9	23.3	50.1	28.8	11.7	75.5	30.0	87.5
Manchurian	266	32.0	43.0	24.1	9	23.3	50.1	28.8	11.7	75.5	30.0	87.5
Garton	2853	39.3	38.3	35.0	20.0	23.3	50.1	28.8	11.7	75.5	30.0	87.5
Manchuria	2853	39.3	38.3	35.0	20.0	23.3	50.1	28.8	11.7	75.5	30.0	87.5
Do	2900	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0
Do	2862	46.8	36.7	47.5	36.7	40.8	42.0	42.2	42.2	42.2	42.2	42.2
Do	2864	47.5	36.7	40.8	42.0	42.2	42.2	42.2	42.2	42.2	42.2	42.2
Daniels	2880	43.3	36.7	40.8	42.0	42.2	42.2	42.2	42.2	42.2	42.2	42.2
Mensury	2657	40.8	42.0	42.2	42.2	42.2	42.2	42.2	42.2	42.2	42.2	42.2
Matting	2909	38.3	32.0	33.4	32.0	33.4	32.0	33.4	32.0	33.4	32.0	33.4
Silver King	890	43.3	33.1	33.8	3	69.3	47.5	46.0	107.7	98.1	98.1	98.1
Chevalier	2910	40.8	42.0	42.2	42.2	42.2	42.2	42.2	42.2	42.2	42.2	42.2
Nepal	595	25.8	29.0	22.5	25.8	29.0	22.5	25.8	29.0	22.5	25.8	29.0
Reed Triumph	889	15.3	30.4	67.0	56.0	64.2	52.1	64.2	52.1	64.2	52.1	64.2
Hybrid	2911	28.8	33.8	8	34.7	30.7	56.0	34.7	30.7	56.0	34.7	30.7
Do	2912	26.5	33.5	0	36.2	22.7	25.0	22.7	25.0	22.7	25.0	22.7
Do	2913	22.7	25.0	22.7	25.0	22.7	25.0	22.7	25.0	22.7	25.0	22.7
Do	2914	26.5	33.5	3	61.6	26.5	33.5	3	61.6	26.5	33.5	3
Manchuria	2903	25.0	24.4	2	69.9	25.0	24.4	2	69.9	25.0	24.4	2
Do	2904	23.9	36.3	2	73.9	23.9	36.3	2	73.9	23.9	36.3	2
Do	2905	28.8	33.8	3	68.2	28.8	33.8	3	68.2	28.8	33.8	3
Do	2906	23.1	33.8	3	68.2	23.1	33.8	3	68.2	23.1	33.8	3
Do	2907	24.7	33.5	0	65.3	24.7	33.5	0	65.3	24.7	33.5	0
Do	2908	28.8	33.8	3	68.2	28.8	33.8	3	68.2	28.8	33.8	3
Do	2915	31.2	29.6	62.5	70.5	31.2	29.6	62.5	70.5	31.2	29.6	62.5
Do	2916	26.2	24.0	8	70.5	26.2	24.0	8	70.5	26.2	24.0	8
Do	2917	24.7	33.5	0	65.3	24.7	33.5	0	65.3	24.7	33.5	0
Do	2918	38.3	35.0	2	51.7	38.3	35.0	2	51.7	38.3	35.0	2
Hanna	2919	10.0	21.5	14.3	3	15.3	30.4	15.3	30.4	15.3	30.4	15.3
Gisborne	1030	45.8	36.7	22.9	45.8	36.7	22.9	45.8	36.7	22.9	45.8	36.7
Malster	585	67.0	60.1	17.5	6	67.0	60.1	17.5	6	67.0	60.1	17.5
Proskowetz	893	40.8	42.0	42.2	42.2	42.2	42.2	42.2	42.2	42.2	42.2	42.2
Black	2726	15.8	43.1	14.1	72.3	15.8	43.1	14.1	72.3	15.8	43.1	14.1
Chevalier	1170	36.7	22.9	45.8	36.7	22.9	45.8	36.7	22.9	45.8	36.7	22.9
Baku	2920	45.8	36.7	22.9	45.8	36.7	22.9	45.8	36.7	22.9	45.8	36.7
Saxradig	607	63.3	33.5	5	63.3	33.5	5	63.3	33.5	5	63.3	33.5
Coast	2921	67.0	60.1	17.5	6	67.0	60.1	17.5	6	67.0	60.1	17.5
Scotch	2922	75.1	56.1	17.0	83.8	75.1	56.1	17.0	83.8	75.1	56.1	17.0
Manshield	2923	63.6	33.5	5	63.6	33.5	5	63.6	33.5	5	63.6	33.5
Mensury	2924	64.7	51.5	74.5	64.7	51.5	74.5	64.7	51.5	74.5	64.7	51.5
Manchuria	2925	79.6	63.6	67.4	9	79.6	63.6	67.4	9	79.6	63.6	67.4
Do	2926	77.9	54.4	76.2	77.9	54.4	76.2	77.9	54.4	76.2	77.9	54.4
Do	2926	68.8	60.2	72.4	68.8	60.2	72.4	68.8	60.2	72.4	68.8	60.2
Do	2823	76.0	61.3	76.1	137.9	42.8	45.2	36.0	13.9	76.0	61.3	76.1
Do	2927	77.1	53.6	66.4	77.1	53.6	66.4	77.1	53.6	66.4	77.1	53.6
Do	2928	81.9	63.0	73.6	86.6	33.8	5.52	3	81.9	63.0	73.6	86.6
Do	2929	70.5	61.3	2	65.9	124.3	65.9	124.3	65.9	124.3	65.9	124.3
Do	2930	79.1	57.3	80.3	79.1	57.3	80.3	79.1	57.3	80.3	79.1	57.3

TABLE 28.—Annual acre yields of varieties of barley grown at the Minnesota Agricultural Experiment Station (at St. Paul) in one or more of the 29 years from 1893 to 1921, inclusive—Continued

Variety	C. I. No.	Acre yields (bushels)																			Years grown	Aver- age yield (bus.)	Per- centage of weight- ed mean										
		1893	1894	1895	1896	1897	1898	1899	1900	1901	1902	1903	1904	1905	1906	1907	1908	1909	1910	1911				1912	1913	1914	1915	1916	1917	1918	1919	1920	1921
White Smyrna	195	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	48.7	33.8	---	---	---	---	---	---	---	2	41.3	77.9
Composite	1147	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	55.0	48.7	69.2	---	---	---	---	---	---	3	57.6	102.7
Gatami	575	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	69.3	48.8	---	---	---	---	---	---	---	2	59.1	111.5
Black Hull-less	1106	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	41.8	13.8	---	---	---	---	---	---	---	2	27.8	52.5
Hybrid	2931	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	73.9	52.1	75.4	---	---	---	---	---	---	3	67.1	119.6
Do	2932	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	70.2	41.3	81.0	---	---	---	---	---	---	3	64.2	114.4
Luth	908	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	52.7	74.7	---	---	---	---	---	---	---	2	63.7	120.6
Hanna	906	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	48.1	51.5	---	---	---	---	---	---	---	2	49.8	94.3
Featherston	911	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	45.8	52.5	---	---	---	---	---	---	---	2	49.1	93.0
Steigum	907	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	45.8	52.5	---	---	---	---	---	---	---	2	49.7	94.1
Eagle	913	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	41.3	63.4	---	---	---	---	---	---	---	2	52.3	99.1
Servian	915	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	66.5	85.6	---	---	---	---	---	---	---	2	45.9	103.8
Mährische	912	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	43.5	50.3	---	---	---	---	---	---	---	5	43.5	99.5
Black	2936	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	1	50.3	74.4
Hybrid	2935	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	4	33.1	106.4
Minsturd	1556	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	53.8	24.2	29.1	43.3	---	---	---	---	---	4	37.6	120.9
Svanota	1907	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	56.5	24.0	29.3	32.5	---	---	---	---	---	4	35.6	114.5
Aker	1577	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	34.0	29.9	0.7	5.5	---	---	---	---	---	3	30.2	113.5
Samola	1211	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	25.7	26.9	98.0	---	---	---	---	---	---	3	26.9	101.1
Bohrman	2933	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	26.9	27.7	23.0	---	---	---	---	---	---	3	25.9	97.4
Lion	923	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	21.6	24.4	33.1	2	---	---	---	---	---	3	25.7	96.6
Gold	1145	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	2	17.1	65.0
Manchuria	1478	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	29.4	23.0	---	---	---	---	---	---	---	2	26.2	99.6
Do	1189	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	2	24.3	92.4
Barb	2793	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	1	22.0	80.9

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 12 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages											
			Chevalier II	Highland Chief	Svanhals	Hanna	Hannchen	Manchuria	Odessa	Coast	Nepal	Black Hull-less	Hybrid	Horsford
Chevalier II	200	Years comparable	-----	4	5	5	5	8	4	2	1	1	5	(*)
	-----	Percentage yield	-----	96.1	97.7	91.2	102.4	71.7	69.7	93.7	126.0	90.8	91.0	-----
	-----	Years comparable	4	-----	1	4	4	5	2	(*)	(*)	3	4	3
Highland Chief	883	Percentage yield	104.1	-----	110.3	101.0	111.7	82.7	94.2	-----	-----	109.7	103.2	105.3
	-----	Years comparable	5	1	-----	6	6	9	6	3	4	4	6	2
	-----	Percentage yield	102.3	90.6	-----	97.7	110.5	76.0	84.5	85.9	167.6	121.1	96.6	118.6
Svanhals	187	Percentage yield	109.7	99.0	102.4	-----	111.8	90.9	102.7	76.2	159.4	115.2	99.9	108.3
	-----	Years comparable	5	4	6	-----	9	9	3	1	4	4	9	2
	-----	Percentage yield	97.6	89.6	90.5	89.5	-----	81.3	80.4	107.6	145.9	105.5	89.4	99.5
Hanna	531	Percentage yield	139.4	120.9	131.6	110.0	123.0	-----	112.3	131.5	195.7	141.9	109.1	132.7
	-----	Years comparable	4	2	6	3	3	10	-----	5	4	7	3	4
	-----	Percentage yield	143.6	106.2	118.3	97.4	124.3	89.1	-----	115.2	163.7	120.6	95.9	123.7
Manchuria	690	Percentage yield	106.8	-----	116.4	131.2	92.9	76.1	86.8	-----	168.3	123.5	86.2	98.3
	-----	Years comparable	1	(*)	4	4	4	6	4	2	-----	6	5	3
	-----	Percentage yield	79.4	-----	59.7	62.7	68.5	51.1	61.1	59.4	-----	72.5	55.5	59.4
Nepal	595	Percentage yield	110	91.2	82.6	86.8	94.8	70.5	82.9	81.0	137.9	-----	77.9	92.0
	-----	Years comparable	5	4	6	9	9	10	3	2	5	5	5	7
	-----	Percentage yield	109.9	96.9	103.5	100.1	111.8	91.7	104.3	116.0	180.2	128.4	-----	118.2
Hybrid	2838	Percentage yield	(*)	3	2	2	2	4	3	3	3	7	3	-----
	-----	Years comparable	-----	94.9	84.3	92.3	100.5	75.4	80.9	101.8	168.4	108.7	84.6	-----
	-----	Percentage yield	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Horsford	507	Percentage yield	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

ST. PAUL, MINN.

Barley varieties have been tested at University Farm, St. Paul, Minn., in all of the 29 years from 1893 to 1921, inclusive. The annual yields given in Table 28 were furnished through the courtesy of the Minnesota Agricultural Experiment Station. During a part of the time the work was carried on in loose cooperation with the United States Department of Agriculture. The annual yields were not filed with the department but have been supplied by the Minnesota station for the years in which there was no connection with the Department of Agriculture as well as for the years of cooperation.

The history of the test at the Minnesota station has been like that at all the other stations of the United States. During the earlier years most of the varieties were not pedigreed, and the identification of many of them is not adequate. It is thought, however, that the chance of their being incorrectly identified is small. The facilities for keeping varieties pure were not so good as at present, and the barleys were grown on unreplicated plats. For these reasons the officials of the Minnesota station have expressed a desire that conclusions be drawn from the yields only when the limitations of the earlier experiments are taken into full consideration. The use made of the yields in this bulletin is but little affected by these conditions of experimentation.

The conditions at Minnesota were no different from those at any of the other stations in the same period, and the work was more carefully carried out and results more carefully recorded than at most places. When it is considered that this bulletin is intended to be partially historical and partially a comparison of commercial varieties and that one of the principal aims is to delimit all regions adapted to types of barley, it can be readily seen that the Minnesota results are most valuable. The large number of barleys of each type included is in a way equivalent to a replication of varieties. In many of the important sorts there was actual replication. Varieties were sent from Minnesota to other stations, reaccessioned at those places, and again brought back to Minnesota, so that the same variety was sometimes carried under two or several numbers. This happened at many stations. Where they could be definitely identified the yields of such duplications have been combined and the average yield reported as the yield of the variety.

The Minnesota station has been one of the most important distributing centers of barley varieties in the United States. From a historical standpoint Minnesota is an invaluable link in tracing the distribution of the early varieties. The Wisconsin Agricultural Experiment Station was responsible for the distribution of many of the varieties grown at the experiment stations under the name of Oderbrucker. The Minnesota station is responsible for a great many of the varieties grown under the name of Manchuria. In a similar way, the Central Experimental Farm at Ottawa, Canada, has furnished a large percentage of our named hybrids. These three stations have been the three most important points distributing to the experiment stations.

In Table 28 it will be seen that the average yields of the barley varieties at St. Paul have been very good. In the column of percentages, where the percentage yield of each variety is given, the rank of the Manchuria group must be very high, as most of the percentages over 100 are those of barleys of this group. Since 200 varieties were tested in one or more of the 29 years, an inspection of Table 28 is difficult. To analyze the results more readily a digest was made by arranging most of the varieties in eight groups. The Manchuria group, which included about 80 varieties, was the best, with the same large percentage superiority as in Wisconsin. The Coast group was second in point of yield, but contained only two varieties, which were grown but a few years. It is not thought that this group is well adapted to Minnesota, and the acre yields were far less than those of the Manchuria. The Hybrid and the Chevalier groups, which follow, were about equal, as were Hanna and Polar, which are next in rank. The Hull-less and Thorpe groups gave low yields.

Table 28 presents a summarized digest in which 12 varieties are compared. These were not selected because of their value, but because they were good varieties, representative of the different groups and were grown long enough to make comparisons possible. Manchuria (C. I. No. 244) was by far the best variety. A hybrid barley (C. I. No. 2838) gave very good yields, as did Highland Chief (C. I. No. 883), Odessa (C. I. No. 182), and Coast (C. I. No. 690).

Several conclusions can be definitely drawn from the earlier experiments. It is unquestionably demonstrated that Minnesota is in a region preeminently adapted to the production of barleys of the Manchuria group. This conclusion is supported by the experiments in the neighboring States as well as at the sub-

stations in Minnesota. The behavior of the hybrid barleys indicates that high-yielding 2-rowed sorts can be produced if desired. The Chevalier and Hanna barleys are not well adapted at St. Paul. This probably is caused by a disease factor, which might be overcome by crossing them with the more resistant Manchuria types. Manchuria (C. I. No. 244), which was widely distributed by the Minnesota station, was a very vigorous stock of barley. It produced high yields, not only at St. Paul but also at many stations to which it was sent. This barley is discussed elsewhere in this bulletin, as are several other varieties which were produced at Minnesota. In recent years the plats have been replicated and the varieties purified. Among the most promising of the new sorts are Manchuria (Minnesota No. 184; C. I. No. 2330), Minsturdi (C. I. No. 1556), Svansota (C. I. No. 1907), Aker (C. I. No. 1577), and Samofa (C. I. No. 1211).

WASECA, MINN.

Barleys were grown on the Southeastern Demonstration Farm and Substation at Waseca, Minn., from 1918 to 1921, inclusive. The annual yields, which are given in Table 29, were furnished through the courtesy of the Minnesota Agricultural Experiment Station. One of the best varieties at Waseca was the hybrid C. I. No. 2935 (Minnesota No. 438). It was developed in the breeding experiments cooperative between the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul. This hybrid is a selection from a cross of Lion $\times$ Manchuria. It is of especial interest because it is smooth awned. If Manchuria barleys with smooth awns could be developed they would be of great importance. The yield of this variety at Waseca is encouraging. Minsturdi (C. I. No. 1556), another hybrid, also yielded well. Manchuria (C. I. No. 2330), a selection made at the Minnesota station, was the most productive for the entire period. A selection of a French Chevalier gave good yields for this type of barley.

TABLE 29.—*Annual acre yields of varieties of barley grown at the Southeast Demonstration Farm and Substation at Waseca, Minn., in part or all of the four years from 1918 to 1921, inclusive*

[Data obtained through the courtesy of the Minnesota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1918	1919	1920	1921			
Manchuria.....	2330	66.0	34.8	33.5	47.8	4	45.5	120.4
French Chevalier.....	2900	56.1	29.8	37.4	34.3	4	39.4	104.2
Manchuria.....	244	58.5				1	58.5	97.2
Hybrid.....	2935		35.6	35.1	48.5	3	39.7	113.8
Minsturdi.....	1556		37.2	33.2	42.6	3	37.7	108.0
Svansota.....	1907		33.0	35.1	29.0	3	32.4	92.8
Manchuria.....	1478			31.2	35.8	2	23.5	67.0
Do.....	1189			31.7	28.2	2	30.0	85.5
Samofa.....	1211			33.4	38.3	2	35.9	102.3
Aker.....	1577				34.4	1	34.4	95.0
Bohman.....	2933				22.8	1	22.8	63.0

DULUTH, MINN.

Barley varieties were grown on the Northeast Demonstration Farm and Substation at Duluth, Minn., during the years 1919, 1920, and 1921. The yields from these tests were furnished by the Minnesota Agricultural Experiment Station and are reported in Table 30. Of the nine varieties included four were grown for all of the three years. Svansota (C. I. No. 1907) was the best of these. This hybrid seemed to be particularly well adapted to the district about Duluth, having produced the highest individual yield in each of the three years. Manchuria (C. I. No. 2330) was second in point of yield, while French Chevalier (C. I. No. 2900) was third. Two other Manchuria selections (C. I. Nos. 1478 and 1189) yielded well in 1920 and 1921.

The results at Duluth indicate that while Duluth is in a district that is best suited to barleys of the Manchuria type the summers are sufficiently cool to allow a normal development of varieties which do not grow normally in the higher temperatures to the south.

TABLE 30.—*Annual acre yields of varieties of barley grown at the Northeast Demonstration Farm and Substation at Duluth, Minn., in part or all of the three years from 1919 to 1921, inclusive*

[Data obtained through the courtesy of the Minnesota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)			Years grown	Average yield (bus.)	Percentage of weighted mean
		1919	1920	1921			
Manchuria.....	2330	47.1	45.1	25.5	3	39.2	107.4
French Chevalier.....	2900	41.3	33.5	30.4	3	35.1	96.2
Minsturdi.....	1556	42.6	40.1	29.0	3	37.2	101.9
Svansota.....	1907	53.7	50.1	33.5	3	45.8	125.5
Manchuria.....	1478	-----	42.3	28.9	2	35.6	104.7
Do.....	1189	-----	39.3	32.2	2	35.7	105.3
Aker.....	1577	-----	-----	28.3	1	28.3	98.3
Samofa.....	1211	-----	-----	25.7	1	25.7	89.2
Bohman.....	2933	-----	-----	25.4	1	25.4	88.2

GRAND RAPIDS, MINN.

The yields of barley varieties grown at the North-Central Experiment Farm at Grand Rapids, Minn., were furnished through the courtesy of the Minnesota Agricultural Experiment Station. These yields are reported in Table 31. Barley was grown in all of the years from 1918 to 1921, inclusive. Only 4 of the 21 varieties were grown for the full four years. Manchuria (C. I. No. 2330) produced the highest yield, as it did at Waseca. French Chevalier (C. I. No. 2900), Hanna (C. I. No. 319), and Nepal (C. I. No. 595) followed in the order named. For the three years in which Minsturdi (C. I. No. 1556) was grown it was slightly superior to Manchuria (C. I. No. 2330). Svansota (C. I. No. 1907), grown for the same years, gave higher yields than Hanna, but did not yield so well as Manchuria (C. I. No. 2330), French Chevalier (C. I. No. 2900), or Minsturdi. Samofa (C. I. No. 1211) is a hybrid between South African and Manchuria, produced at the Minnesota Agricultural Experiment Station. It was tested only in the year 1921, when it gave the highest yield. Although several hybrid varieties were very promising at Grand Rapids, barleys of the Manchuria type are unquestionably well adapted. It is probable that the hybrids are valuable in proportion to their resemblance to Manchuria, especially in the matter of disease resistance.

TABLE 31.—*Annual acre yields of varieties of barley grown at the North-Central Experiment Farm at Grand Rapids, Minn., in part or all of the four years from 1918 to 1921, inclusive*

[Data obtained through the courtesy of the Minnesota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1918	1919	1920	1921			
Manchuria.....	2330	51.5	37.5	34.1	14.5	4	34.4	115.1
French Chevalier.....	2900	50.3	34.8	30.5	9.9	4	31.4	105.0
Odesa.....	182	52.8	38.3	-----	10.8	3	34.0	112.6
Hanna.....	319	55.7	29.7	24.7	12.0	4	30.5	102.0
Nepal.....	595	24.8	18.3	18.8	4.5	4	16.6	55.5
Manchuria.....	244	51.1	-----	-----	-----	1	51.1	109.4
Blue Ribbon.....	611	46.7	-----	-----	-----	1	46.7	100.0
Oderbrucker.....	2700	39.0	-----	-----	-----	1	39.0	83.5
O. A. C. 21.....	1470	47.8	-----	-----	-----	1	47.8	102.4
Golden Queen.....	558	47.7	-----	-----	-----	1	47.7	102.1
Manchuria.....	241	50.3	-----	-----	-----	1	50.3	107.7
Champion of Vermont.....	2899	40.7	-----	-----	-----	1	40.7	87.2
Svanhals.....	187	49.0	-----	-----	-----	1	49.0	104.9
Hybrid.....	2935	-----	32.4	27.6	9.1	3	23.0	104.1
Svansota.....	1907	-----	32.0	31.6	9.4	3	24.3	110.0
Minsturdi.....	1556	-----	40.3	33.1	14.9	3	29.4	133.0
Aker.....	1577	-----	-----	-----	11.8	1	11.8	98.3
Manchuria.....	1478	-----	-----	-----	12.7	1	12.7	105.8
Do.....	1189	-----	-----	-----	14.9	1	14.9	124.2
Samofa.....	1211	-----	-----	-----	16.5	1	16.5	137.5
Bohman.....	2933	-----	-----	-----	15.1	1	15.1	125.8

CROOKSTON, MINN.

The annual yields of barley varieties from the Northwest Experiment Farm at Crookston, Minn., were furnished through the courtesy of the Minnesota Agricultural Experiment Station. The yields for the years 1919 to 1921, inclusive, are reported in Table 32. Of the 21 varieties 13 were tested for the three years. Minsturdi (C. I. No. 1556) produced the highest yield for this period, while Manchuria (C. I. No. 2330), Beardless (C. I. No. 3144), O. A. C. 21 (C. I. No. 1470), and Hybrid (C. I. No. 2935) were second, third, fourth, and fifth. Of the varieties grown but two years Samofa (C. I. No. 1211) and Manchuria (C. I. No. 1189) were the best. Several of the varieties tested were produced at the Minnesota Agricultural Experiment Station at St. Paul. The origin, identity, and general value of the better ones are discussed elsewhere in this bulletin. At Crookston barleys of the Manchuria group are the superior ones. French Chevalier (C. I. No. 2900), Hannchen (C. I. No. 531), and Svanhals (C. I. No. 187), which represent three 2-rowed groups, did not compare favorably in yielding capacity with the Manchuria barleys.

TABLE 32.—*Annual acre yields of varieties of barley grown at the Northwest Experiment Farm at Crookston, Minn., in part or all of the three years 1919 to 1921, inclusive*

[Data obtained through the courtesy of the Minnesota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)			Years grown	Average yield (bus.)	Percentage of weighted mean
		1919	1920	1921			
Manchuria	2330	25.9	30.7	24.4	3	27.0	120.0
Do	2823	22.2	22.3	17.8	3	20.8	92.4
French Chevalier	2900	14.4	30.8	17.0	3	20.7	92.0
Hybrid	2935	15.8	31.0	26.4	3	24.4	108.4
Minsturdi	1556	25.2	34.3	25.2	3	28.2	125.3
Svansota	1907	15.5	28.2	23.1	3	22.3	99.1
Aker	1577	22.4	—	—	1	22.4	119.8
Hanna	319	16.6	26.6	21.4	3	21.5	95.6
O. A. C. 21	1470	26.9	25.8	21.6	3	24.8	110.2
Beardless	3144	27.2	32.7	14.7	3	24.9	110.7
Hannchen	531	9.3	31.3	16.0	3	18.9	84.0
Svanhals	187	12.8	31.6	14.9	3	19.8	88.0
Oderbrucker	2700	20.2	26.4	20.0	3	22.2	98.7
Princess	529	13.4	26.7	18.1	3	19.4	86.2
Nepal	595	12.0	—	—	1	12.0	64.2
Manchuria	2823	—	22.3	14.9	2	18.6	77.5
Do	1189	—	34.7	17.6	2	26.1	108.7
Samofa	1211	—	36.7	16.1	2	26.4	110.0
Bohman	2933	—	25.7	19.9	2	22.8	95.0
Mahrtsche	912	—	26.9	15.2	2	21.1	87.9
Manchuria	1478	—	29.7	14.9	2	22.3	92.9

TABLE 33.—*Annual acre yields of varieties of barley grown at the West-Central Experiment Farm at Morris, Minn., in part or all of the four years from 1918 to 1921, inclusive*

[Data obtained through the courtesy of the Minnesota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1918	1919	1920	1921			
Manchuria	2330	38.1	29.4	32.1	28.8	4	32.1	117.2
French Chevalier	2900	35.2	32.7	—	21.0	3	29.6	102.8
Manchuria	2823	41.1	—	23.0	29.5	3	31.2	116.0
Wisconsin Pedigree	835	35.7	25.7	24.9	—	3	28.8	98.0
Manchuria	244	30.4	—	—	—	1	30.4	81.7
Oderbrucker	2700	42.7	—	—	—	1	42.7	114.8
Hybrid	2935	—	24.4	17.0	11.6	3	17.7	70.2
Minsturdi	1556	—	34.3	19.4	21.3	3	25.0	99.2
Svansota	1907	—	31.5	28.7	21.5	3	27.2	107.9
Manchuria	1478	—	—	24.4	27.0	2	25.7	108.0
Do	1189	—	—	30.5	30.6	2	30.5	128.2
Samofa	1211	—	—	16.4	23.7	2	20.1	84.5
Aker	1577	—	—	—	23.4	1	23.4	98.7
Bohman	2933	—	—	—	22.0	1	22.0	92.8

MORRIS, MINN.

Barley varieties were tested on the West-Central Experiment Farm at Morris, Minn., from 1918 to 1921, inclusive. The yields reported in Table 33 were furnished by the Minnesota Agricultural Experiment Station. The highest yields were obtained from varieties of the Manchuria type, of which Manchuria (C. I. No. 2330) and Manchuria (C. I. No. 1189) were probably the best. Svansota (C. I. No. 1907), the best of the hybrids, produced an average yield slightly less than that of Manchuria (C. I. No. 2330). The selection of French Chevalier (C. I. No. 2900) was better than the average at Morris. Although the varietal tests have been run only four years it is quite apparent that barleys of the Manchuria type are particularly well adapted to cultivation in this section.

AMES, IOWA

Varietal tests of barley were conducted at Ames, Iowa, in cooperation with the Iowa Agricultural Experiment Station from 1913 to 1921, inclusive. Ten varieties were tested, and all of them were carried in the experiment for the entire period. The annual yields are reported in Table 34. Through 1917 the yields were figured to check. From 1918 to 1921, inclusive, actual yields are given. The best two varieties were Oderbrucker (C. I. No. 1272) and Oderbrucker (C. I. No. 2700). Oderbrucker (C. I. No. 1272) is a pedigreed variety developed by the Wisconsin Agricultural Experiment Station under the pedigree No. 5. O. A. C. 21 (C. I. 1470) was third in point of yield. Like the Oderbrucker barleys it belongs to the Manchuria group. Following O. A. C. 21 is Manchuria (C. I. No. 241).

The 2-rowed varieties tested, Frankish (C. I. No. 295) and Hanna (C. I. No. 203), are barleys belonging to the Hanna group. Frankish gave an average yield of 111 per cent of the weighted mean, but was only fifth in point of yield. Hanna was quite inferior to Frankish, exceeding only the low-yielding hull-less and hooded varieties. Caucasian (C. I. No. 90), a Russian 6-rowed bearded sort, differing greatly from Caucasian (C. I. No. 2724) grown in Canada, gave fairly good yields, but was also surpassed by the Manchuria barleys. There is no question as to the types of barleys best suited to this district. The barleys of the Manchuria group are outstanding in yield at Ames, as they were in Wisconsin and Minnesota. For many years Oderbrucker has been the predominating variety grown on the farms in Iowa.

TABLE 34.—*Annual acre yields of varieties of barley grown at the Iowa Agricultural Experiment Station (at Ames) in the nine years from 1913 to 1921, inclusive*

[Data obtained in cooperation with the Iowa Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)									Years grown	Average yield (bus.)	Per centage of weighted mean
		1913	1914	1915	1916	1917	1918	1919	1920	1921			
Caucasian.....	90	41.5	32.5	11.0	40.5	42.1	26.5	8.0	26.3	24.3	9	28.1	110.6
Oderbrucker.....	2700	34.0	43.9	12.7	36.5	49.2	35.4	15.4	27.5	20.8	9	30.6	120.5
Do.....	1272	41.3	35.0	11.4	35.8	49.2	35.4	20.8	31.3	20.8	9	31.2	122.8
Manchuria.....	241	33.9	35.8	9.5	32.4	52.9	30.4	18.3	31.7	20.8	9	29.5	116.1
O. A. C. 21.....	1470	35.5	34.2	16.5	32.0	50.8	27.5	16.3	33.8	22.9	9	29.9	117.7
Frankish.....	295	40.7	31.7	12.6	23.2	54.6	27.5	12.9	27.5	22.9	9	28.2	111.0
Hanna.....	203	36.8	16.9	10.0	18.7	42.0	25.4	2.5	14.6	20.8	9	20.9	82.3
Black Hull-less.....	596	32.2	22.6	9.1	28.3	31.2	21.3	7.9	18.8	14.6	9	20.7	81.5
Nepal.....	595	27.0	18.9	4.0	21.4	21.3	26.3	6.3	20.0	12.5	9	17.5	68.9
Horsford.....	507	26.3	19.6	5.2	26.0	24.2	18.8	5.4	16.7	14.6	9	17.4	68.5

TABLE 35.—*Annual acre yields of varieties of barley grown at the Missouri Agricultural Experiment Station (at Columbia) and at the soil-experiment farms at Lewistown, Carthage, Maryville, and Warrensburg, Mo., in one or more of the 12 years from 1910 to 1921, inclusive*

[Data obtained through the courtesy of the Missouri Agricultural Experiment Station]

Location and variety	C. I. No.	Acre yields (bushels)										Years grown	Average yield (bus.)	Per- centage of weight- ed mean	
		1910	1911	1912	1914	1915	1916	1917	1918	1919	1920				1921
Columbia:															
Hanna	906								15.8	6.9		2	11.3	59.5	
Steigum	907								26.1	11.6		2	18.9	99.5	
Luth	908								32.2	17.5	21.2	3	23.6	112.4	
Eagle	913								28.6	12.2		2	20.4	107.4	
Italian	914								9.5	5.5		2	7.5	39.5	
Servian	915								30.0	11.8		2	20.9	110.0	
Odessa	916								20.3	7.7		2	14.0	73.7	
Lion	923								32.8	18.9	32.3	3	28.0	133.3	
Australian White	925								2.1			1	2.1	7.4	
Horn	926								32.1	8.2		2	20.1	105.8	
Odessa	927								29.5	14.8	31.9	3	25.4	121.0	
Summit	929								31.9	13.6	28.0	3	24.5	116.7	
Club Mariout	932								26.5			1	26.5	93.0	
Odessa	934								14.4	3.5		2	8.9	46.8	
Peruvian	935								31.9	10.3		2	21.1	111.1	
Trebi	936								43.5	14.9	39.6	3	32.7	155.7	
Sandrel	937								47.1	17.8	39.6	3	34.8	165.7	
Oderbrucker	940								31.5	12.6		2	22.1	116.3	
Frankish	953								29.8	13.6	33.9	3	25.8	122.9	
Manchuria	956								41.7	12.1	32.6	3	28.8	137.1	
Oderbrucker	957								34.5	8.6		2	21.5	113.2	
Alpha	959								21.5	11.4		2	16.5	86.8	
Luth	972								30.5	4.5		2	17.5	92.1	
Red River	973								33.3	6.6		2	19.9	104.7	
Featherston	1118								35.3	6.1		2	20.7	108.9	
Do	1119								33.1	3.7		2	18.4	96.8	
Do	1120								43.8	13.9	35.0	3	30.9	147.1	
Hanchamont	1121								18.1	15.6		2	16.9	88.9	
Manchuria	1125								26.5	6.2		2	16.3	85.8	
Malting	1129								22.2	9.3		2	15.7	82.6	
Manchuria	244									8.6		1	8.6	83.5	
Oderbrucker	537									12.2	38.7	2	25.5	160.4	
Nepal	595									2.5		1	2.5	24.3	
Black Hull-less	596									2.5		1	2.5	24.3	
Horsford	507									14.6	27.5	2	21.1	132.7	
Lewistown:															
Horsford	507		2.8	6.1								2	4.5	62.5	
Manchuria	244		3.7	10.8								2	7.3	101.4	
Oderbrucker	537			10.4								1	10.4	114.3	
Nepal	595			9.2								1	9.2	101.1	
Carthage:															
Horsford	507	25.7	6.1									2	15.9	102.6	
Manchuria	244	28.5	4.2									2	16.3	105.2	
Nepal	595	24.8										1	24.8	94.3	
Black Hull-less	596		3.7									1	3.7	78.7	
Maryville:															
Manchuria	244				15.2			29.7	36.8	18.5	31.6	5	26.4	94.6	
Horsford	507				15.4		22.7	29.3		30.3	36.3	26.3	6	26.7	
Oderbrucker	537				14.4			36.5	37.3		46.3	29.9	5	32.9	
Black Hull-less	596										13.1		1	13.1	
Warrensburg:															
Manchuria	244				3.4	25.3	25.7				5.5		4	15.0	
Oderbrucker	537				2.7	8.2	14.8				6.7	12.7	5	9.0	
Nepal	595				.7								1	.7	
Horsford	507				4.0	25.1	19.2					8.5	4	14.2	

COLUMBIA, MO.

The yields of barley varieties grown at Columbia, Mo., were furnished through the courtesy of the Missouri Agricultural Experiment Station. The yields for the years 1910 to 1921, inclusive, are reported in Table 35. In 1919 30 varieties were grown, 28 of which were continued in the experiment the following year. Five other varieties were introduced in 1920. This was a year of very low yields, and only a few of the better sorts were continued in 1921. Of these Sandrel (C. I. No. 937) and Trebi (C. I. No. 936) were undoubtedly the best. Two varieties of the Man-

churia group, Featherston (C. I. No. 1120) and Manchuria (C. I. No. 956), were third and fourth, respectively. Lion (C. I. No. 923), a 6-rowed smooth-awned barley, also gave a high average yield. Oderbrucker (C. I. No. 537) produced the highest yield of the five varieties introduced in 1920. It was, however, surpassed by Trebi, Sandrel, and Lion. The 2-rowed sorts were quite inferior. Frankish (C. I. No. 953) and Horn (C. I. No. 926) were the only ones whose averages exceeded that of the weighted mean. While a 3-year test is not of sufficient length to make any definite statements regarding the value of the individual varieties, it would seem that there is an overlapping of growing conditions here. Barleys of the Manchuria group and such varieties as Sandrel and Trebi, usually grown under the more arid conditions of the Western States, gave relatively high yields in this locality. The barley commonly grown in this region, however, is of the Manchuria-Oderbrucker type.

The Missouri station has conducted barley experiments at four of its soil-experiment fields. These are located at Lewistown, Warrensburg, Carthage, and Maryville, Mo. At each of these places four varieties were tested, two of which, Manchuria (C. I. No. 244) and Horsford (C. I. No. 507), were carried at all of them.

At Lewistown Manchuria (C. I. No. 244) and Oderbrucker (C. I. No. 537) produced the best yields, the former having been grown in the years 1911 and 1912 and the latter in 1912 only. Horsford (C. I. No. 507), grown in 1911 and 1912, and Nepal (C. I. No. 595), grown in 1911, did not compare favorably with the other two.

At Warrensburg these same four varieties were tested during the years from 1914 to 1916, inclusive, and in 1920 and 1921. Manchuria (C. I. No. 244) was again the leading variety, but Horsford was superior to Oderbrucker at this station. Nepal (C. I. No. 595) gave such a low yield in 1914 that it was not continued further in the experiment.

The results at Maryville are quite comparable to those at Warrensburg. Oderbrucker (C. I. No. 537) was first in point of yield, Horsford (C. I. No. 507) second, Manchuria (C. I. No. 244) third, and Black Hull-less fourth. The Manchuria and Oderbrucker varieties belong to the same varietal group and are almost identical. At Warrensburg, however, Manchuria gave the higher yields.

Varieties were tested at Carthage during the years 1910 and 1911. Oderbrucker was not grown at this station, but both Nepal and Black Hull-less were included. Manchuria (C. I. No. 244) was again the leading variety, with Horsford following. The Hull-less varieties Nepal (C. I. No. 595) and Black Hull-less (C. I. No. 596) were grown for only one year, Nepal in 1910 and Black Hull-less in 1911, and were the lowest in point of yield in these years.

Results at these four stations indicate that Manchuria (C. I. No. 244) and Oderbrucker (C. I. No. 537) were the best of the varieties tested. At both Maryville and Warrensburg, however, Horsford gave relatively higher yields than is usually the case.

FARGO, N. DAK.

Varietal tests of barley have been made at the North Dakota Agricultural Experiment Station at Fargo since 1892. There were developed at this station a number of strains which have been tested at other places. The North Dakota station, like those of Minnesota, Wisconsin, and Ontario, was a center for the distribution of varieties to other stations. It is an important link in the tracing of varieties. At Fargo, as at many other places, the same variety was introduced more than once and in some cases was carried under different numbers at the same time. Manchuria (C. I. No. 244) was grown under North Dakota Nos. 788 and 792 for eight years. The average yield of the variety under each of these two numbers was exactly the same at the end of this period. The average deviation from the mean was only 0.9 bushel, while the maximum deviation from the mean was only 1.6 bushels. This is of special interest, since many agronomists feel that the earlier tests, which were conducted without any replication of plats, are not of significant value. It is unlikely that the slight variations of yield shown by these two varieties are typical of what might be expected. Probably the actual accuracy is far less than is indicated by this comparison. The writers feel, however, that the value of this early work has been underestimated. Certainly, the deviation from the mean was very slight when compared with an average yield of more than 38 bushels.

The data reported in Table 36 were obtained through the courtesy of the North Dakota Agricultural Experiment Station. A group digest of the varieties was made, which showed the Manchuria group to be the best of six compared.

The Thorpe group was much inferior to the Manchuria, while the Chevalier was slightly inferior to the Thorpe. Higher yields were obtained from the Hanna than from the Hull-less or the Horsford groups. In the summarized digest of Table 36 eight well-known varieties belonging to different groups are compared. Manchuria (C. I. No. 244) is the best variety. Chevalier II (C. I. No. 200) is superior to all except Manchuria. No comparison between Chevalier II and Odessa (C. I. No. 182) was possible, as they were not grown in the same years. Hannchen (C. I. No. 531) produced yields almost as high as those of Chevalier II. Mansury (C. I. No. 617) is a representative of the Thorpe group. Mansury was superior to Odessa, Evans (C. I. No. 892), Horsford (C. I. No. 507), and Nepal (C. I. No. 595).

In recent years several promising selections of Manchuria have been made at the North Dakota station. Manchuria (C. I. No. 2945; N. Dak. 2119), Manchuria (C. I. No. 2947; N. Dak. 2121), Manchuria (C. I. No. 2949; N. Dak. 2123), and Manchuria (C. I. No. 2951; N. Dak. 2125) are among the best of these. The Michigan selection of Lion (C. I. No. 923) gave high yields in 1920 and 1921, the only years in which it was tested. The results here demonstrate that Fargo is in the section where Manchuria barleys are particularly adapted. Barleys of the Chevalier, Thorpe, and Hanna groups yielded relatively higher than at St. Paul or Madison. Although the Manchuria barleys are the ones best adapted, it is evident that the conditions are increasingly favorable to the production of 2-rowed barleys in the districts north and west of Madison and St. Paul.

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF EIGHT REPRESENTATIVE VARIETIES

[*Explanation.*—The asterisk (*) indicates that the two varieties were not grown in the same years]

[illegible]

33610°—25†——6

Variety	C. I. No.	Data shown	Varieties and percentages							
			Cheva- lier II	Man- sury	Hann- chen	Man- churia	Odessa	Evans	Nepal	Horsford
Chevalier II	200	{ Years comparable { Percentage yield	9	9	9	9	(*)	9	5	8
Man-sury	617	{ Years comparable { Percentage yield	111.6	101.8	94.9	94.9	3	117.9	211.3	147.9
Hannchen	531	{ Years comparable { Percentage yield	89.6	91.2	91.2	13	(*)	14	11	13
Manchuria	244	{ Years comparable { Percentage yield	98.3	109.6	15	15	2	108.2	178.3	128.6
Odessa	182	{ Years comparable { Percentage yield	105.4	113.6	117.3	85.3	2	115.9	208.1	142.1
Evans	892	{ Years comparable { Percentage yield	96.2	91.6	3	2	3	123.4	205.0	136.6
Nepal	595	{ Years comparable { Percentage yield	84.8	86.3	10	15	3	111.8	160.1	130.6
Horsford	507	{ Years comparable { Percentage yield	47.3	56.1	5	9	2	159.4	110.9	8
			8	13	8	13	3	62.7	64.9	
			67.6	77.8	70.4	73.2	76.6	90.2	154.0	

TABLE 37.—Annual acre yields of barley grown at the Edgeley (N. Dak.) substation in two or more of the years from 1903 to 1921, inclusive

Variety	C. I. No.	Acre yields (bushels)										Years grown	Aver- age yield (bus.)	Percent- age of weighted mean
		1903	1904	1905	1906	1907	1908	1909	1910	1911	1914	1920	1921	
Evans.....	892	21.1	42.1	50.6	25.3	10.8	19.0	32.4	1.7	1.5	---	---	---	34.8
Mansury.....	617	22.5	57.8	60.8	42.7	10.8	19.0	32.4	1.7	1.5	---	---	---	86.6
Manchuria.....	244	24.6	40.3	48.1	39.5	10.1	21.1	34.3	2.1	1.0	38.0	28.6	12.7	131.1
Drum.....	241	22.1	49.9	53.5	39.5	8.1	21.8	28.7	7.1	5.5	---	---	---	117.4
Horsford.....	507	---	48.3	37.6	24.3	---	15.0	---	---	---	---	---	---	121.2
Club Maricot.....	261	---	44.0	48.8	---	18.9	30.3	43.2	8.8	14.6	---	---	---	101.6
Baldi Dwarf.....	100	---	43.5	40.2	---	18.9	30.3	43.2	8.8	14.6	---	---	---	143.3
Talli.....	194	---	28.8	50.8	---	27.0	14.8	---	---	---	---	---	---	105.7
Nopal.....	595	---	---	37.6	---	14.1	22.0	30.5	3.0	1.6	---	---	---	31.5
Blue Ribbon.....	611	---	---	---	---	10.8	33.8	41.7	6.7	7.0	21.0	---	---	111.4
New Moroccan.....	2706	---	---	---	---	8.9	27.4	31.2	1.1	2.5	---	---	---	17.3
Oatbrucker.....	2706	---	---	---	---	16.2	22.2	34.5	2.8	2.5	---	---	---	20.0
Bohemian.....	27	---	---	---	---	20.7	26.8	35.1	4.4	2.5	---	---	---	14.2
Do.....	32	---	---	---	---	18.9	20.4	31.9	5.0	1.0	---	---	---	15.4
Mansury.....	170	---	---	---	---	18.9	31.6	33.9	3.4	2.5	---	---	---	18.1
Princess.....	520	---	---	---	---	18.2	6.3	---	---	---	---	---	---	12.3
Hansley.....	524	---	---	---	---	36.5	18.3	41.4	2.3	3.0	---	---	---	20.3
Chevalier II.....	209	---	---	---	---	20.3	16.9	30.1	6.6	1.5	---	---	---	13.9
Primus.....	539	---	---	---	---	10.2	23.2	31.5	5.9	2.0	26.8	---	---	17.6
Hanna.....	203	---	---	---	---	33.7	28.5	36.3	3.1	1.7	---	---	---	20.7
Hanna.....	31	---	---	---	---	13.9	21.1	---	---	---	---	---	---	19.1
Stratum.....	47	---	---	---	---	27.0	26.1	37.1	2.7	2.5	---	---	---	104.4
Goldfoil.....	162	---	---	---	---	18.9	17.6	---	---	---	---	---	---	80.6
Kitzing.....	107	---	---	---	---	13.5	6.3	---	---	---	---	---	---	79.2
Surprise.....	171	---	---	---	---	8.1	19.0	---	---	---	---	---	---	9.9
Summit.....	174	---	---	---	---	29.3	28.2	---	---	---	---	---	---	13.5
Summit.....	184	---	---	---	---	32.3	13.4	35.2	5.9	1.5	---	---	---	19.0
Scotchman.....	187	---	---	---	---	27.3	28.2	---	---	---	---	---	---	103.8
Black Smyrna.....	197	---	---	---	---	27.0	28.2	30.5	5.0	2.5	---	---	---	18.6
White Smyrna.....	191	---	---	---	---	35.4	46.0	40.0	14.3	12.0	---	---	---	28.7
Frankish.....	207	---	---	---	---	32.4	38.7	31.0	6.5	1.0	---	---	---	15.7
Highland Chief.....	883	---	---	---	---	24.3	12.6	36.8	3.8	2.4	---	---	---	14.4
Giant.....	575	---	---	---	---	16.2	31.7	42.8	7.4	13.6	---	---	---	24.5
Kitzing.....	189	---	---	---	---	27.9	23.2	32.6	1.0	1.0	---	---	---	19.1
Oden.....	182	---	---	---	---	37.8	41.2	---	---	---	---	---	---	41.0
Golden Queen.....	558	---	---	---	---	27.8	35.0	30.0	2.1	1.0	---	---	---	17.5
Robertson.....	559	---	---	---	---	43.2	20.4	38.6	4.8	2.0	---	---	---	104.9
Hannchen.....	531	---	---	---	---	31.1	31.3	33.0	5.8	2.0	29.3	---	---	21.8

[Data for 1904 to 1909, inclusive, obtained in cooperation with the North Dakota Agricultural Experiment Station, through whose courtesy the data for other years were obtained]

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages									
			Chevalier II	Man- sury	Svan- hals	Hanna	Hann- chen	Man- churia	Club Mariout	White Smyrna	Nepal	Summit
Bavarian.....	158	-----	-----	24.3	30.3	35.9	6.9	1.0	-----	-----	5	19.7
Do.....	159	-----	-----	18.9	15.0	30.1	7.1	3.0	-----	-----	5	14.8
Black Hull-less.....	506	-----	-----	-----	-----	36.7	6.1	3.6	-----	-----	4	16.9
Red Triumph.....	889	-----	-----	-----	21.4	30.9	1.4	-----	-----	-----	3	10.9
Oderbrucker.....	836	-----	-----	-----	-----	-----	-----	-----	-----	-----	3	25.0
Do.....	1272	-----	-----	-----	-----	-----	-----	33.0	30.4	11.7	2	76.2
									29.7	11.3		106.2
Chevalier II.....	200	{ Years comparable.....	-----	5	5	5	5	5	5	5	5	5
		{ Percentage yield.....	-----	106.1	74.5	67.2	67.2	101.2	59.9	53.5	5	73.0
Mansury.....	617	{ Years comparable.....	5	-----	5	5	5	9	7	5	6	5
		{ Percentage yield.....	94.2	-----	70.2	63.3	63.4	113.2	88.7	50.5	116.0	68.8
Svanhals.....	187	{ Years comparable.....	5	5	-----	5	5	5	5	5	5	5
		{ Percentage yield.....	134.3	142.5	-----	90.2	90.3	135.9	80.5	71.9	130.9	98.0
Hanna.....	203	{ Years comparable.....	5	5	-----	-----	-----	5	5	5	5	5
		{ Percentage yield.....	148.8	158.0	110.8	-----	100.1	150.6	89.2	79.7	145.1	108.6
Hannchen.....	531	{ Years comparable.....	5	5	5	5	5	6	5	5	5	5
		{ Percentage yield.....	148.7	157.8	110.7	99.9	-----	124.3	89.1	79.6	144.9	108.5
Manchuria.....	244	{ Years comparable.....	5	9	5	5	6	-----	7	5	8	5
		{ Percentage yield.....	98.8	88.4	73.6	66.4	80.5	-----	75.3	52.9	113.8	72.1
Club Mariout.....	261	{ Years comparable.....	5	5	5	5	5	7	-----	5	6	5
		{ Percentage yield.....	166.9	112.8	124.3	112.1	112.2	132.9	-----	89.4	151.3	121.8
White Smyrna.....	195	{ Years comparable.....	5	5	5	5	5	5	5	-----	5	5
		{ Percentage yield.....	186.7	198.2	139.1	125.5	125.6	188.9	111.9	-----	182.0	136.3
Nepal.....	595	{ Years comparable.....	5	6	5	5	5	8	6	5	5	5
		{ Percentage yield.....	102.6	86.2	76.4	68.9	69.0	57.8	66.1	54.9	-----	74.9
Summit.....	174	{ Years comparable.....	5	5	5	5	5	5	5	5	5	5
		{ Percentage yield.....	137.0	145.4	102.0	92.1	92.2	138.6	82.1	73.4	133.6	-----

EDGELEY, N. DAK.

Varietal tests of barley were made at the Edgeley substation in North Dakota in 12 of the 19 years from 1903 to 1921, inclusive. During the years 1904 to 1909, inclusive, they were in cooperation with the United States Department of Agriculture. The yields before and after these dates were obtained through the courtesy of the North Dakota Agricultural Experiment Station. The annual yields are given in Table 37. A group digest not reported here showed the Chevalier and Hanna groups to exceed the Manchuria in yield. The Hybrid, Hull-less, and Thorpe groups follow in the order named.

In the summarized digest of Table 37, in which 10 individual varieties representing different groups are compared, it is still more apparent that the Manchuria barleys do not hold here the position of prominence that they maintain throughout most of the Mississippi Valley. White Smyrna (C. I. No. 195), Club Mariout (C. I. No. 261), Hanna (C. I. No. 203), and Hannchen (C. I. No. 531) produced yields in the order named. The two highest yielding sorts were varieties which do best where the ripening season comes during a period of dry weather. The Hanna and Hannchen also are varieties that produce their best yields in regions in which the diseases that flourish in hot humid climates do not occur. It will be remembered that at the Canadian stations, the results of which have already been presented, the 2-rowed barleys began to yield relatively well in about the same longitude in Canada. The conditions favorable to the production of 2-rowed barley which exist in New York State probably obtain far to the north of the Mississippi Valley in Canada, but extend south again into the northern Great Plains of the United States.

LANGDON, N. DAK.

The results of barley varietal experiments at the Langdon substation in North Dakota (Table 38) were furnished through the courtesy of the North Dakota Agricultural Experiment Station. Only three varieties, all of the Manchuria type, Manchuria (C. I. Nos. 244 and 241) and Common (C. I. No. 184), were tested for the full seven years of the experiment, 1909 to 1915, inclusive. Of these three, Common produced the highest average yield. Manchuria (C. I. Nos. 241 and 244) followed, being almost the equal of Common. Silver King (C. I. No. 890), a pedigreed Manchuria barley, was grown from 1909 to 1914, inclusive. It produced comparatively high yields, as did Mansury (C. I. No. 617), a barley of the Thorpe group, which was grown for the same period. Two hull-less varieties, Evans (C. I. No. 892) and Nepal (C. I. No. 595), were low yielders, as was also Horsford (C. I. No. 507). Three new sorts, Oderbrucker (C. I. No. 836), Hannchen (C. I. No. 531), and Primus (C. I. No. 532), were grown in 1914 and 1915. Hannchen produced an average yield slightly higher than that of Oderbrucker and Primus. The results obtained indicate that, while Langdon is still in the Manchuria district, the 2-rowed barleys can be grown with almost equal success. These results correspond with those found at Edgeley.

TABLE 38.—*Annual acre yields of varieties of barley grown at the Langdon (N. Dak.) substation in part or all of the seven years from 1909 to 1915, inclusive*

[Data obtained through the courtesy of the North Dakota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)							Years grown	Average yield (bus.)	Percentage of weighted mean
		1909	1910	1911	1912	1913	1914	1915			
Manchuria.....	244	38.5	15.1	38.7	37.3	29.2	12.5	41.5	7	30.4	105.6
Do.....	241	41.6	17.3	40.1	37.0	28.3	15.8	40.1	7	31.5	109.4
Silver King.....	890	44.5	15.7	39.0	34.7	26.7	12.1	-----	6	28.8	107.1
Common.....	184	42.7	13.6	39.6	39.3	35.0	16.3	36.7	7	31.9	110.8
Evans.....	892	51.4	10.1	25.0	26.6	22.0	16.3	-----	6	25.2	93.7
Mansury.....	617	37.0	10.0	30.7	53.0	26.7	12.9	-----	6	28.4	105.6
Horsford.....	507	-----	14.8	34.4	31.7	20.3	15.0	31.0	6	24.5	90.4
Nepal.....	595	-----	9.1	36.6	30.5	22.0	-----	-----	4	24.5	88.1
Oderbrucker.....	836	-----	-----	-----	-----	-----	17.1	44.2	2	30.7	119.5
Hannchen.....	531	-----	-----	-----	-----	-----	15.3	48.1	2	31.9	124.1
Primus.....	532	-----	-----	-----	-----	-----	13.3	48.1	2	30.7	119.5

MANDAN, N. DAK.

Six varieties of barley were grown at the United States Northern Great Plains Field Station at Mandan, N. Dak., from 1916 to 1921, inclusive. This work was carried on in cooperation with the Office of Dry-Land Agriculture Investigations, United States Department of Agriculture. The barley varieties, although few in number, represent several groups. White Smyrna (C. I. No. 195) gave the highest yields, as will be seen in Table 39. Hannchen (C. I. No. 531) was somewhat less productive than White Smyrna, and Svanhals (C. I. No. 187) was third in point of yield. Coast (C. I. No. 690) and Club Mariout (C. I. No. 261) ranked fourth and fifth, while Manchuria (C. I. No. 244) was last. Here, again, it is evident that the Manchuria barleys are not better adapted than those of other types. The six years reported, however, contain no years of high yields. It is likely that the Manchuria would produce relatively higher yields in years of greater rainfall. The best three varieties were 2-rowed sorts.

TABLE 39.—*Annual acre yields of varieties of barley grown at the United States Northern Great Plains Field Station (Mandan, N. Dak.) in the six years from 1916 to 1921, inclusive*

[Data obtained in cooperation with the Office of Dry-Land Agriculture Investigations]

Variety	C. I. No.	Acre yields (bushels)						Years grown	Average yield (bus.)	Percentage of weighted mean
		1916	1917	1918	1919	1920	1921			
White Smyrna.....	195	29.0	23.1	25.0	21.8	11.3	7.7	6	19.7	117.3
Club Mariout.....	261	25.8	14.5	22.6	13.6	10.8	7.3	6	15.5	94.0
Coast.....	690	25.8	16.8	24.5	17.5	11.4	2.9	6	16.5	98.2
Manchuria.....	244	20.8	8.1	17.6	13.7	11.4	.6	6	12.0	71.4
Hannchen.....	531	31.0	20.4	25.3	20.4	16.0	0	6	18.9	112.5
Svanhals.....	187	26.2	20.3	26.2	18.7	18.1	0	6	18.3	108.9

DICKINSON, N. DAK.

The yields of barley varieties grown at the Dickinson (N. Dak.) substation were obtained in cooperation with the North Dakota Agricultural Experiment Station. Results furnished for all of the 15 years from 1907 to 1921, inclusive, are reported in Table 40. Only one variety, Hanna (C. I. No. 203), was grown for the entire period. Five varieties were included for 14 years. A comparison is made between 10 well-known varieties in the summarized digest of the table. Hannchen (C. I. No. 531) gave the highest yields of the 10 varieties compared. Svanhals (C. I. No. 187) was almost as good as Hannchen. The average yield of Hanna (C. I. No. 203) was 97 per cent of that of Svanhals and Hannchen. Gatami (C. I. No. 575) was the best 6-rowed variety, but it was not so good as White Smyrna (C. I. No. 658), the lowest yielding 2-rowed sort in the comparison. Manchuria (C. I. No. 244) was not higher than sixth in yield. Since the tests at Dickinson cover a period of 15 years, the results are much more conclusive than those at Edgeley or Mandan. The best varieties at Dickinson are unquestionably 2-rowed sorts. At Lethbridge, Alberta, they were also superior. It would seem that the ecological conditions that favored this type of barley in southern Alberta obtain in western North Dakota.

TABLE 40.—Annual acre yields of varieties of barley grown at the Dickinson (N. Dak.) substation in part or all of the 15 years from 1907 to 1921, inclusive

[Data obtained in cooperation with the North Dakota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)										Years grown	Average yield (bus.)	Percentage of weighted mean			
		1907	1908	1909	1910	1911	1912	1913	1914	1915	1916				1917	1918	1919
Nepal.....	595	40.9	22.1	24.1	23.0	8.4	---	20.9	12.5	41.5	29.6	11.1	9.7	0.2	38.0	5.7	83.0
Hanna.....	203	46.8	28.5	39.8	34.8	17.0	46.0	31.4	16.6	62.0	39.8	15.5	16.5	1.0	47.5	7.5	120.0
Gatani.....	575	36.7	31.4	27.9	34.5	11.9	---	27.5	24.3	39.7	29.6	10.7	22.2	4.9	39.6	7.3	100.8
Manchuria.....	244	16.5	27.9	30.2	28.0	7.8	---	30.6	16.3	49.3	32.5	9.7	14.9	3.2	41.5	6.8	91.1
Imperial.....	289	34.4	28.3	40.4	---	---	---	---	---	---	---	---	---	---	---	---	101.2
Oderbrucker.....	2700	---	22.1	30.0	23.5	---	---	---	---	---	---	---	---	---	---	---	78.7
Chevalier II.....	200	---	36.1	45.8	---	---	---	---	---	---	---	---	---	---	---	---	87.7
Primus.....	532	---	29.3	36.7	36.7	12.3	41.3	42.7	22.9	66.0	30.8	14.2	---	---	---	---	121.4
Hannchen.....	531	---	37.5	48.7	35.4	15.2	39.2	35.0	17.8	62.8	39.8	16.2	14.8	1.9	46.1	5.8	117.7
Princess.....	529	---	32.7	38.5	---	---	---	---	---	---	---	---	---	5.1	48.0	7.7	120.2
Svarthals.....	187	---	39.4	44.9	---	---	---	---	---	---	---	---	---	1.2	38.4	5.9	111.9
Coast.....	626	---	---	---	34.8	8.2	41.1	26.7	18.5	61.0	44.8	17.0	13.3	---	---	---	116.5
Farnesh.....	616	---	---	---	32.9	12.0	---	---	---	---	---	---	---	---	---	---	95.7
Horsford.....	507	---	---	---	14.7	3.5	---	---	---	---	---	---	---	---	---	---	43.3
White Gatani.....	920	---	---	---	19.5	---	---	---	---	---	---	---	---	6.8	40.6	11.9	94.4
Proskowetz.....	883	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	83.4
White Smyrna.....	658	---	---	---	---	---	---	---	13.5	61.7	39.4	14.5	---	---	---	---	117.9
Oderbrucker.....	888	---	---	---	---	---	---	---	22.1	41.3	27.6	14.8	20.4	3.7	48.1	9.8	104.4
Chile Common.....	663	---	---	---	---	---	---	---	9.6	---	---	---	---	---	---	---	56.8
Manchuria.....	241	---	---	---	---	---	---	---	20.8	36.8	14.8	11.6	15.9	3.7	29.4	3.7	76.0
Steigum.....	907	---	---	---	---	---	---	---	7.3	---	---	---	---	---	---	---	43.2
Club Mariout.....	932	---	---	---	---	---	---	---	---	30.1	15.9	15.9	23.6	3.1	47.8	9.5	109.6
Wing Pedigree.....	1177	---	---	---	---	---	---	---	---	16.7	10.1	10.1	37.2	5.5	37.2	8.5	84.8
Scholey.....	962	---	---	---	---	---	---	---	---	---	---	---	13.8	1.5	47.1	5.9	90.5
Heil Hanna 4.....	677	---	---	---	---	---	---	---	---	---	---	---	---	6.3	51.4	8.6	115.1
White Smyrna.....	2169	---	---	---	---	---	---	---	---	---	---	---	---	6.1	45.6	7.2	102.1
Odessa.....	182	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	118.2
White Smyrna.....	2170	---	---	---	---	---	---	---	---	---	---	---	---	---	44.7	10.9	102.7

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages									
			Svanhals	Hanna	Hann- chen	Man- churia	Gatami	Chile Common	Club Mariout	White Smyrna	Nepal	Wing Pedigree
Svanhals	187	{ Years comparable	13	13	13	13	13	8	6	8	13	4
		{ Percentage yield	103.0	97.7	123.4	118.3	146.4	119.6	106.5	106.5	149.3	86.1
Hanna	203	{ Years comparable	13	14	14	14	14	8	6	106.5	14	4
		{ Percentage yield	97.1	97.0	128.4	116.2	151.0	126.8	106.9	106.9	140.7	100.1
Hannchen	531	{ Years comparable	13	13	13	13	13	8	6	8	13	4
		{ Percentage yield	102.3	126.2	126.2	121.0	150.1	123.6	106.3	106.3	152.8	100.4
Manchuria	244	{ Years comparable	13	13	13	13	13	8	6	8	14	4
		{ Percentage yield	103.0	79.2	110.5	90.5	127.4	107.7	92.8	92.8	109.6	97.2
Gatami	575	{ Years comparable	13	13	13	13	13	8	6	94.9	14	4
		{ Percentage yield	81.1	82.6	110.5	130.4	127.4	113.4	94.9	94.9	121.0	108.3
Chile Common	663	{ Years comparable	8	8	8	8	8	6	6	72.8	8	4
		{ Percentage yield	68.3	66.6	78.5	76.7	127.4	78.5	72.8	72.8	92.2	77.2
Club Mariout	932	{ Years comparable	6	6	6	6	6	6	6	81.0	6	4
		{ Percentage yield	83.6	80.9	92.8	88.2	127.4	123.4	81.0	81.0	106.9	108.3
White Smyrna	658	{ Years comparable	8	8	8	8	8	8	6	126.6	8	4
		{ Percentage yield	93.9	91.5	107.8	105.3	137.4	126.6	126.6	126.6	120.1	120.1
Nepal	595	{ Years comparable	13	13	13	13	13	8	6	8	126.6	4
		{ Percentage yield	67.0	65.5	91.3	82.6	108.5	93.6	79.0	79.0	4	78.5
Wing Pedigree	1177	{ Years comparable	4	4	4	4	4	4	4	4	4	4
		{ Percentage yield	116.2	94.2	102.9	92.3	129.6	92.3	83.3	83.3	127.4	127.4

TABLE 41.—Annual acre yields of varieties of barley grown at the Williston (N. Dak.) substation in one or more of the 14 years from 1908 to 1921, inclusive

[Data for the years 1908 to 1918, inclusive, obtained in cooperation with the North Dakota Agricultural Experiment Station, through whose courtesy data for 1920 and 1921 were furnished]

Variety	C. I. No.	Acre yields (bushels)												Years grown	Average yield (bus.)	Percent- age of weighted mean
		1908	1909	1910	1911	1912	1913	1914	1915	1916	1917	1918	1920	1921		
Success	891	11.3	46.0	1.2	8.5	68.3	26.9	48.5	73.8	53.7	---	---	---	---	37.6	91.9
Munsey	617	6.4	56.0	2.0	8.1	51.2	---	---	---	---	---	---	---	---	24.7	86.4
Russian	1855	11.3	57.8	3.9	4.3	60.6	---	---	---	---	---	---	---	---	5	96.5
Baxter	1855	9.8	51.3	2.6	8.5	56.9	---	---	---	---	---	---	---	---	5	96.5
Hanna	203	10.0	56.0	3.9	10.6	65.8	19.6	66.7	80.2	61.7	23.5	10.2	13.8	39.2	13	103.5
Manchuria	241	12.6	54.5	6.6	8.2	68.5	44.8	53.2	69.2	56.5	---	---	---	---	9	101.7
Silver King	890	12.0	57.2	4.7	7.7	63.1	35.4	---	---	---	---	---	---	---	6	100.7
Bernard	881	17.1	52.1	5.7	9.8	62.9	40.0	53.9	64.2	56.3	26.0	4.2	10.0	---	8	97.7
Manchuria	244	13.7	51.1	4.7	8.9	64.0	37.1	58.7	72.1	---	---	---	---	---	12	98.5
Highland Chief	883	10.6	47.8	2.4	5.5	66.6	24.0	---	---	---	---	---	---	---	6	87.6
Manchuria	882	18.2	53.7	6.1	12.3	65.5	51.3	71.3	70.2	54.8	25.0	6.3	12.5	32.5	13	107.6
Nopal	595	---	---	1.7	8.1	52.1	---	---	---	---	---	---	---	---	4	82.2
Reed Triumph	889	---	46.8	2.2	5.1	61.5	51.8	---	---	---	---	---	---	---	5	103.7
Oderbrucker	888	---	55.6	3.7	7.7	59.2	49.8	52.9	75.8	58.3	25.4	3.7	10.0	45.0	12	104.8
Gatani	575	---	---	7.3	19.5	45.0	---	53.2	72.6	45.2	25.4	3.0	10.8	20.8	10	93.0
Evans	892	---	---	4.5	7.2	45.0	---	---	---	---	---	---	---	---	3	73.5
Canadian Thorpo	740	---	---	---	---	50.0	---	---	---	---	---	---	---	---	1	83.3
Manchurian	739	---	---	---	---	58.3	46.3	76.6	66.7	58.8	24.4	3.1	10.0	27.5	9	100.2
Himalaya	619	---	---	---	---	27.2	27.2	---	---	---	---	---	---	---	1	75.1
Proskowetz	893	---	---	---	---	25.8	25.8	54.7	80.0	63.4	29.8	10.2	10.8	---	7	101.0
Hannchen	531	---	---	---	---	26.2	26.2	72.9	83.4	65.9	24.1	9.0	10.8	34.2	8	106.8
Svanhals	187	---	---	---	---	---	---	53.7	65.5	52.5	16.7	8.0	5.6	31.7	7	86.8
Primus	582	---	---	---	---	---	---	63.6	75.0	40.4	20.0	4.6	5.8	20.8	7	85.5
Arnold	2741	---	---	---	---	---	---	---	67.5	45.4	15.4	---	---	---	3	83.4
Flaxton	2742	---	---	---	---	---	---	---	75.5	---	---	---	---	---	1	104.4
Kropshadt	2743	---	---	---	---	---	---	---	50.2	36.3	---	---	---	---	2	68.5
Tartar	1116	---	---	---	---	---	---	---	80.0	68.3	26.9	8.4	---	---	4	112.0
Triega	2744	---	---	---	---	---	---	---	---	45.9	21.3	---	---	---	2	84.8
White Smyrna	668	---	---	---	---	---	---	---	---	---	20.0	8.3	12.9	---	3	103.0
Wing Pedigree	1177	---	---	---	---	---	---	---	---	---	---	5.2	---	---	1	81.3
Horsford	1271	---	---	---	---	---	---	---	---	---	---	5.6	9.6	36.7	3	117.7
Oderbrucker	886	---	---	---	---	---	---	---	---	---	---	---	10.8	---	1	104.9
Manchuria	2947	---	---	---	---	---	---	---	---	---	---	---	33.3	---	1	103.4

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF EIGHT REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages							
			Primus	Svanhals	Hanna	Hann- chen	Man- churia	Gatami	Tartar	Nepal
Primus	532	1/4 ears comparable.	7	7	7	7	7	7	4	(*)
		Percentage yield	98.5	78.0	76.7	84.4	99.7	76.3	76.3	(*)
Svanhals	187	1/4 ears comparable.	7	7	7	7	7	7	4	(*)
		Percentage yield	101.5	79.1	77.8	85.7	101.2	77.7	77.7	(*)
Hanna	203	1/4 ears comparable.	7	7	8	13	10	10	4	4
		Percentage yield	128.3	126.4	96.4	96.1	118.5	96.6	96.6	131.9
Hannchen	531	1/4 ears comparable.	7	8	8	8	7	7	4	(*)
		Percentage yield	130.5	103.7	100.8	100.8	130.0	99.3	99.3	(*)
Manchuria	882	1/4 ears comparable.	7	18	8	8	10	10	4	4
		Percentage yield	118.4	104.0	99.2	99.2	112.5	85.1	85.1	133.2
Gatami	575	1/4 ears comparable.	7	10	7	7	10	4	4	3
		Percentage yield	100.3	84.4	76.9	76.9	88.9	79.6	79.6	138.9
Tartar	1116	1/4 ears comparable.	4	4	4	4	4	4	4	(*)
		Percentage yield	131.1	128.7	100.7	100.7	117.5	125.6	125.6	(*)
Nepal	595	1/4 ears comparable.	(*)	(*)	(*)	(*)	4	3	(*)	(*)
		Percentage yield	75.8	75.8	75.1	75.1	72.0	72.0	72.0	(*)

WILLISTON, N. DAK.

Varietal tests of barley were conducted at the Williston (N. Dak.) substation in cooperation with the North Dakota Agricultural Experiment Station from 1908 to 1918, inclusive. Records of yields of 1920 and 1921 were obtained through the courtesy of the North Dakota Agricultural Experiment Station. The annual yields are shown in Table 41. Four groups of barley were compared, of which the Manchuria and Hanna were practically equal in rank, while the Thorpe ranked higher than the Hull-less. In the summarized digest of Table 41 the yields of eight well-known varieties are compared. Tartar (C. I. No. 1116) produced slightly higher yields than Hannchen (C. I. No. 531). Yields practically equal to those of Tartar and Hannchen were obtained from Manchuria (C. I. No. 882). Hanna (C. I. No. 203) was fourth and Svanhals (C. I. No. 187) fifth in point of yield. Tartar (C. I. 1116) is a selection made from the same stock as Trebi (C. I. No. 936). Under irrigation in southern Idaho Trebi has proved to be a highly productive variety. Tartar has not been widely tested, and its value is not known. Disregarding the Tartar, it would seem that the Manchuria barleys were probably slightly superior to the 2-rowed sorts at Williston. This means that the line separating the districts where Manchuria is superior from that in which the 2-rowed barleys grow best runs in a north-westerly direction through North Dakota.

BROOKINGS, S. DAK.

Yields are reported in this bulletin from several stations in South Dakota. These data are available only for those years in which the work was carried on in cooperation at the various South Dakota stations or, in some cases, where the yields in other years have been published in bulletins of the State station. It is thought that the conclusions which are drawn from the results are not affected by the missing years.

In Table 42 yields are reported from a large number of varieties grown at Brookings in one or more of the years from 1902 to 1919, inclusive. To facilitate the study of the results six different groups of varieties were compared. The Manchuria was much superior to the other five groups. The Hanna, Chevalier, and Thorpe, in the order given, ranked next. The Coast group was not represented in this comparison, but probably would have been inferior to the four named.

In Table 42 the yields of 11 varieties representative of several different groups are compared. These are not necessarily the highest yielding ones of their groups. They are, however, well-known varieties which are of interest both because they are representative of a group and are widely grown. Of the 11 varieties compared the highest yield was obtained from Odessa (C. I. No. 182), which seems particularly adapted to eastern South Dakota. Manchuria (C. I. No. 241) was almost the equal of Odessa. The third variety in point of yield was Gatami (C. I. No. 575), which was tested in only a few years and was comparable with some of the varieties in only one year. Its place, therefore, is not conclusively determined. Following Gatami are three 2-rowed varieties of almost equal value. These are Gold (C. I. No. 1145), Svanhals (C. I. No. 187), and White Smyrna (C. I. No. 195). In recent years three selections have shown promise. These are Manchuria (C. I. No. 2156), Manchuria (C. I. No. 2155), and Odessa (C. I. No. 2154). Oderbrucker (C. I. No. 1146) also yielded well.

It is apparent from the original data (Table 42) and the summarized digest that barleys of the Manchuria type have produced the highest yields in eastern South Dakota. Odessa has given somewhat better yields than the best strains of either Manchuria or Oderbrucker. This variety belongs to the same general group as the Manchuria. Selections have been made from Odessa, however, which differ very much from the Manchuria barleys. Odessa (C. I. No. 182) as tested in South Dakota, while containing a few of these variant types, was of the same general character as Manchuria. The advantage of early ripening, which is so important in western South Dakota, was seen to have some influence even at Brookings, for both Gatami and White Smyrna gave fairly good yields. Nepal (C. I. No. 595) produced very low yields at Brookings but not much lower than at most points in the Mississippi Valley.

TABLE 42.—Annual acre yields of varieties of barley grown at the South Dakota Agricultural Experiment Station (at Brookings) in one or more of the 18 years from 1902 to 1919, inclusive

[Data obtained in cooperation with the South Dakota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)													Years grown	Average age of weighted mean (bus.)	Percentage of weighted mean				
		Acre yields (bushels)																			
		1902	1904	1905	1906	1907	1908	1909	1910	1911	1912	1913	1914	1915				1916	1917	1918	1919
Chevalier	23	33.0	34.5																2	33.7	87.1
Hanna	24	34.7	47.0	60.6	46.4	19.2	30.1	14.6	19.2	0	25.2	35.4	38.5						12	30.9	96.0
Scale	25	37.8	44.5	39.0															3	40.4	104.9
Utano	26	38.0	42.9	54.8	46.4	21.3	15.2												6	36.4	111.0
Bokenan	27	35.6	45.4	42.7	49.5	22.1	18.7				52.0	32.3	30.2						9	36.5	106.3
Utano	28	40.4	46.6	46.2															3	44.4	115.3
Beshorn Imperial	29	29.0	38.3																2	32.1	82.9
Hanna	30	33.8	45.0	43.5	44.0	20.2	22.0												6	34.7	105.8
Utano	31	28.6	43.7	31.0	41.4														4	36.4	93.8
Bokenan	32	39.0	47.7	35.4	44.4														4	41.6	107.2
Hanna	33	33.2	43.5	37.0	36.8														4	38.3	98.7
Do	34	28.6	47.5	38.5	41.7														4	39.1	100.8
Chevalier	35	31.3	42.9	64.2	50.3	22.5	27.8	15.4	11.0	.8	32.0	41.7							11	30.9	96.6
Nishima	45	10.4																	1	10.4	35.4
Strigum	47	22.3	42.5	30.0	43.1	20.8	13.9												6	28.8	87.8
Golden Melon	48	26.0	45.8	46.3	37.0	25.6	16.0												6	32.8	100.0
Nita	49	11.7																	1	11.7	38.8
Tunkaze	50	15.6																	1	15.6	53.1
Santaku	52	23.5	37.7	23.4															3	28.2	73.2
Shiro-Nishiki	54	20.9	36.6	39.3															3	35.3	91.7
Uyetake	56	52.0																	1	52.0	176.9
Nijo	60	36.5	34.6	35.4															3	35.5	92.2
Doitsu	62	39.0	56.6	26.9															3	40.8	106.0
Baitori	64	17.4																	1	17.4	59.2
Sangatsuka	78	22.0	27.9	32.9	31.2	21.3	13.3												6	24.8	75.6
Mocki-Mugi	80	20.8																	1	20.8	70.7
Beldi Dwarf	190	8.6	50.7	26.0							41.7	25.0	52.0						6	34.0	84.2
Black Smyrna	191	35.0	59.1	35.2															3	43.1	111.9
Telli	194	18.8	27.0																2	22.9	59.2
Hanna	203	45.0	47.0	48.3	49.0	23.3	14.7												6	37.9	115.5
Manchuria	244	29.7	46.8								47.2	64.8	53.2	61.4	43.7	45.8	56.3	21.9	10	47.1	119.2
Do	241	31.5	56.6	33.3	41.0	34.6	27.4	24.6	24.2	0	39.3	64.6	64.5	64.5	33.2	57.3	14.6	12	36.7	111.9	
White Smyrna	195	31.5	49.3												31.6	33.2		10	38.2	97.9	
Coreth	36	31.5	56.6	33.3											2	26.1		2	64.1	64.1	
Nepal	22		29.0	29.4											2	23.3		5	23.3	70.4	
Do	158		29.0	41.3	25.3	12.0	8.8											5	37.0	111.8	
Bavarian	159		54.6	40.5	40.2	21.0	19.5											5	34.5	104.2	
Do	170		59.9	48.9	36.8	14.6	12.5											5	37.7	113.9	
Mensury	175		48.5	27.1	31.8													5	37.7	113.9	
Medney	178		50.7	29.9	36.5	20.8												4	34.0	96.9	

TABLE 42.—Annual acre yields of barley grown at the South Dakota Agricultural Experiment Station (at Brookings) in one or more of the 18 years from 1902 to 1919, inclusive—Continued

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 11 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages										
			Cheval- lier	Primus	Svanhals	Bohe- mian	Hann- chen	Man- churia	Odessa	Gatami	Gold	White Smyrna	Nepal
Chevalier	35	Years comparable	-----	5	10	7	10	11	10	1	1	4	5
Primus	532	Percentage yield	-----	130.8	97.8	105.5	102.3	90.4	79.7	87.1	85.8	108.0	204.1
Svanhals	187	Years comparable	-----	5	5	5	5	5	5	(*)	(*)	2	4
Bohemian	27	Percentage yield	-----	76.5	75.1	89.0	81.2	82.8	74.8	-----	-----	97.7	166.9
Hannchen	531	Years comparable	-----	10	6	6	10	10	10	1	1	3	5
Manchuria	241	Percentage yield	-----	102.2	-----	106.0	104.5	91.6	81.4	68.7	67.7	108.6	173.2
Odessa	182	Years comparable	-----	5	6	-----	8	8	8	3	3	6	7
Gatami	575	Percentage yield	-----	94.8	94.3	106.4	94.0	84.1	77.1	67.9	75.5	96.9	148.1
Gold	1145	Years comparable	-----	10	10	8	-----	11	16	61.0	81.1	88.8	137.3
White Smyrna	195	Percentage yield	-----	97.8	95.7	118.9	119.9	-----	93.1	2	2	5	6
Nepal	595	Years comparable	-----	110.6	109.2	118.9	116	11	69.4	107.8	130.9	124.6	211.1
		Percentage yield	-----	125.5	122.8	129.7	144.1	107.5	-----	103.8	137.9	128.4	195.6
		Years comparable	-----	114.9	145.6	147.3	163.8	92.8	96.4	-----	-----	-----	181.0
		Percentage yield	-----	1	1	3	7	2	7	7	132.9	128.0	7
		Years comparable	-----	116.5	147.7	132.5	123.2	76.4	72.5	75.2	-----	96.3	136.2
		Percentage yield	-----	4	3	6	9	5	11	7	7	-----	8
		Years comparable	-----	92.6	92.1	103.1	112.6	80.3	77.9	78.1	103.8	137.1	137.1
		Percentage yield	-----	5	5	7	11	6	11	7	7	8	-----
		Years comparable	-----	49.0	57.7	67.5	72.8	47.4	51.1	55.2	73.4	73.0	-----

EUREKA, S. DAK.

Tests of barley varieties were carried on at the substation at Eureka, S. Dak., in cooperation with the South Dakota Agricultural Experiment Station for 11 years, 1909 to 1919, inclusive. In at least two of the years 1909, 1910, and 1911 17 varieties were tested (Table 43). The yields for 1909 were very good. The crop of 1911 was a total failure. Only three varieties were tested for more than five years. In four of the seven years in which Gatami (C. I. No. 575) and Hannchen (C. I. No. 531) were both grown Hannchen was superior to Gatami. Gatami excelled Manchuria (C. I. No. 244) in four of the seven comparable years. Under favorable seasonal conditions, however, Manchuria produced much more than Gatami, so that the average production of Manchuria for the seven years was greater than that of Gatami. White Smyrna (C. I. No. 195) was better than Hannchen in two out of four comparable years and exceeded Manchuria in all of them, but these were not favorable years. Odessa (C. I. No. 182), which was also grown four years, was better than Manchuria in two of these years and superior to Hannchen in three of them. White Smyrna, comparable with Odessa for only three years, exceeded it in two of the three.

Eureka is in a district where there is a high premium on early maturity. In years of adequate rainfall barleys of the Manchuria type produced good yields. The conditions which were favorable to the 2-rowed barleys in western North Dakota obviously extended into South Dakota. This is reflected in the high yield of Hannchen. In years of low yield, that is, in years of low rainfall, such varieties as White Smyrna and Gatami are most likely to produce grain. It is entirely possible that even a small return in these years may be more valuable than a higher return from other varieties in the good years. The early-season varieties are more likely to produce feed in all years.

TABLE 43.—*Annual acre yields of varieties of barley grown at the Eureka (S. Dak.) substation in some of the 11 years from 1909 to 1919, inclusive*

[Data obtained in cooperation with the South Dakota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)											Years grown	Average yield (bus.)	Percent-age of weighted mean
		1909	1910	1911	1912	1913	1914	1915	1916	1917	1918	1919			
Manchuria.....	241	30.9	5.1	0	-----	-----	-----	-----	-----	-----	-----	-----	3	12.0	122.4
Do.....	244	31.2	6.0	0	2.6	6.4	22.8	60.1	43.2	27.0	0.6	14.3	11	19.5	139.3
Oderbrucker.....	1146	33.3	2.7	0	-----	-----	-----	-----	-----	-----	-----	-----	3	12.0	122.4
Odessa.....	182	29.5	5.2	0	-----	-----	-----	-----	-----	-----	1.0	18.9	5	10.9	112.4
Bohemian.....	27	32.1	8.0	0	-----	-----	-----	-----	-----	-----	-----	-----	3	13.4	136.7
Chevalier II.....	200	31.0	.3	0	-----	-----	-----	-----	-----	-----	-----	-----	3	10.4	106.1
Golden Melon.....	48	26.2	1.3	0	-----	-----	-----	-----	-----	-----	-----	-----	3	9.2	93.9
Hanna.....	24	18.7	4.3	0	-----	-----	-----	-----	-----	-----	-----	-----	3	7.7	78.6
Do.....	34	23.1	3.3	0	-----	-----	-----	-----	-----	-----	-----	-----	3	8.8	89.8
Do.....	203	18.7	3.1	0	-----	-----	-----	-----	-----	-----	-----	-----	3	7.3	74.5
Hannchen.....	531	29.6	3.8	0	21.4	16.5	30.5	67.5	33.0	32.2	.7	17.8	11	23.0	164.3
Horn.....	31	27.5	12.2	0	-----	-----	-----	-----	-----	-----	-----	-----	3	13.2	134.7
Primus.....	532	18.7	1.7	0	-----	-----	-----	-----	-----	-----	-----	-----	3	6.8	69.4
Princess.....	529	28.3	0	0	-----	-----	-----	-----	-----	-----	-----	-----	3	9.4	95.9
Striegum.....	47	27.1	3.9	0	-----	-----	-----	-----	-----	-----	-----	-----	3	10.3	105.1
Gatami.....	575	7.3	0	23.4	11.4	30.9	39.2	16.8	19.6	-----	-----	-----	8	18.6	167.6
White Smyrna.....	195	7.2	0	7.8	-----	-----	-----	-----	-----	-----	2.9	16.3	5	6.8	154.5

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 11 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages												
			Golden Melon	Bohemian	Hannchen	Manchuria	Odessa	Gatami	Coast	Club Marlot	White Smyrna	Beldi Dwarf	Horsford		
White Gatami		920											4	31.6	102.7
Caucasian		714											1	43.7	105.3
Iranian		679											4	23.1	75.0
Iranian		314											4	17.8	57.8
Poppenheim		507											4	22.6	73.4
Horsford		1177											4	22.3	83.8
Living Pedigree		2169											3	34.7	131.4
White Smyrna		2170											2	37.1	140.5
Do		2171											2	35.3	133.7
Do		2172											2	33.7	127.7
SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 11 REPRESENTATIVE VARIETIES															
[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]															
Golden Melon	48	Years comparable													
		Percentage yield	7	82.1	5	3	3	3	5	5	5	7	89.4	4	(*)
Bohemian	27	Years comparable	121.7		10	8	8	101.1	92.7	121.1	12	107.3	125.4	5	(*)
		Percentage yield	5		95.6	139.9	101.4	10	10	6	15	125.4	5		
Hannchen	531	Years comparable	122.6	104.6		111.6	13	95.1	91.4	123.5	104.5	147.4	5		
		Percentage yield	3				13	10	10	6	13	104.5	5		
Manchuria	244	Years comparable	112.8	83.4	13		13	84.2	80.9	101.8	88.4	90.8	5		
		Percentage yield	3		88.6		13	10	10	6	13	88.4	5		
Odessa	182	Years comparable	122.5	98.7	116.7	130.2	10	115.1	110.6	127.8	115.1	128.3	5		
		Percentage yield	(*)	5		10	10		10	6	10	128.3	5		
Gatami	575	Years comparable		98.9	105.1	118.7	10		96.0	121.6	102.5	140.4	5		
		Percentage yield	(*)	5		10	10		6	6	10	140.4	5		
Coast	626	Years comparable		107.9	103.3	123.6	90.4	104.1		127.7	106.7	128.1	5		
		Percentage yield	(*)	5		6	6	6	6	6	6	128.1	5		
Club Marlot	261	Years comparable		82.6	81.0	98.2	78.2	82.3	78.3		85.3	124.1	5		
		Percentage yield	7		15	13	13	10	10	6	6	124.1	5		
White Smyrna	195	Years comparable	111.8	93.2	95.7	113.2	86.9	97.6	93.7	117.3					
		Percentage yield	(*)	4		5	5	5	5	5	5	142.0			
Beldi Dwarf	190	Years comparable		79.7	67.8	110.1	78.0	71.2	78.1	80.6	70.4				
		Percentage yield	(*)	4		4	4	4	4	(*)	4				
Horsford	507	Years comparable			81.4	84.6	56.9	71.7	69.5		72.6				
		Percentage yield													

HIGHMORE, S. DAK.

The yields of barley varieties grown at Highmore, S. Dak., in cooperation with the South Dakota Agricultural Experiment Station, from 1903 to 1919, inclusive, are reported in Table 44. Eleven representative varieties are compared in the summarized digest of the same table. Odessa (C. I. No. 182) was the highest yielding sort. It was also the best of the varieties compared at Brookings. Second in point of yield was Coast (C. I. No. 626). Bohemian (C. I. No. 27), a representative of the Hanna group, was third in rank. This variety, however, was dropped from the experiment in 1915, thus making it comparable with a number of the varieties for only the last four or five years it was grown. The crops in two of these years were almost complete failures owing to drought. The rank of Bohemian is not so conclusive as that of the other varieties. Following it is Gatami (C. I. No. 575). Although this barley came from the same general region in which Manchuria originated it ripens somewhat earlier, thus making it more drought escaping.

While the 2-rowed varieties Hannchen (C. I. No. 531) and White Smyrna (C. I. No. 195) do not rank relatively so high as those just mentioned, they were included in the experiment for a longer period than any of the other varieties tested and have produced uniformly good yields with the exception of the three years of drought, 1911, 1912, and 1913. In 1911 no yields were obtained from any of the varieties with the exception of Manchuria (C. I. No. 241), which produced only 0.1 bushel. In 1912 and 1913 White Smyrna exceeded all of the other sorts in yield. Manchuria (C. I. No. 244) was very low in rank, being superior only to Golden Melon (C. I. No. 48), Club Mariout (C. I. No. 261), and Horsford (C. I. No. 507). In 1918 and 1919 four promising selections of White Smyrna were grown. They were exceeded only by Odessa (C. I. No. 182).

It is apparent that in moving westward in South Dakota, while Odessa holds its position of prominence, barleys of the Manchuria group are no longer among the better yielding varieties. They are superseded by Coast, Gatami, and the 2-rowed types.

COTTONWOOD, S. DAK.

Barleys were grown at the Cottonwood (S. Dak.) substation, in cooperation with the South Dakota Agricultural Experiment Station from 1909 to 1919, inclusive. The yields during this period are reported in Table 45. The relative values of the varieties are difficult to determine. Many of them were tested for only one or two years, and these were years of drought. Four varieties are comparable for the years 1916 to 1919, inclusive. Of these, Odessa (C. I. No. 182) produced the highest yield, White Smyrna (C. I. No. 195) was second, while Gatami (C. I. No. 575) and Hannchen (C. I. No. 531) were third and fourth, respectively. Here, again, it will be noted that two varieties characterized by early ripening yielded well. Odessa, which was the best sort at Brookings for many years, was slightly better than the two early-season varieties. White Smyrna and Hannchen have given good results from Highmore north and west over a considerable area.

TABLE 45.—Annual acre yields of varieties of barley grown at the Cottonwood (S. Dak.) substation in one or more of the 11 years from 1909 to 1919, inclusive

[Data obtained in cooperation with the South Dakota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)										Years grown	Average yield (bus.)	Percentage of weighted mean
		1909	1910	1912	1913	1914	1915	1916	1917	1918	1919			
Manchuria.....	241	1.0	2.0									2	1.5	53.6
Do.....	244	2.0	3.1	2.6	0.4	1.3		19.9	0.3			7	4.2	68.9
Oderbrucker.....	1146	2.2	3.1									2	2.7	96.4
Bohemian.....	32	3.9	1.6									2	2.7	96.4
Chevalier II.....	200	3.4	2.9									2	3.1	110.7
Hanna.....	34	2.8										1	2.8	93.3
Do.....	24	2.4	3.5									2	2.9	103.6
Do.....	203	1.6										1	1.6	53.3
Hannchen.....	531	6.1	3.5			3.1	8.6	18.5	10.0	17.7	24.4	8	11.5	111.7
Horn.....	31	4.9	2.2									2	3.5	125.0
Golden Melon.....	48		1.2									1	1.2	46.1
Gatami.....	575					6.6		19.9	5.4	25.0	23.9	5	16.2	94.7
Odessa.....	182							25.8	11.6	27.8	37.3	4	25.6	132.6
White Smyrna.....	195							19.4	10.5	25.2	25.4	4	29.1	104.1

NEWELL, S. DAK.

Dry-land conditions.—The yields of barley varieties grown on dry land at the Belle Fourche Experiment Farm, Newell, S. Dak., were obtained in cooperation with the Office of Western Irrigation Agriculture. These yields for the years 1908 to 1919, inclusive, are shown in Table 46. Ten varieties representing different types and groups are compared in the summarized digest of Table 46. The highest yields were obtained from White Smyrna (C. I. No. 658) and Hannchen (C. I. No. 531). Coast (C. I. No. 690), which was grown only during the last five years of the experiment, was not comparable with Chevalier II (C. I. No. 200) or Hanna (C. I. No. 24). It was superior, however, to the other five of the ten varieties. Odessa (C. I. No. 182), Gatami (C. I. No. 575), and Manchuria (C. I. No. 241) follow in the order named. Here again the results confirm the conclusions that in the western two-thirds of South Dakota the 2-rowed barleys are equal or superior to the 6-rowed sorts. At Newell under dry-land conditions all other varieties are surpassed by the 2-rowed varieties Hannchen and White Smyrna.

TABLE 46.—*Annual acre yields of varieties of barley grown on dry land at the Belle Fourche (Newell, S. Dak.) Experiment Farm in part or all of the 12 years from 1908 to 1919, inclusive*

[Data obtained in cooperation with the South Dakota Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)												Years grown	Average yield (bus.)	Percentage of weighted mean
		1908	1909	1910	1911	1912	1913	1914	1915	1916	1917	1918	1919			
Manchuria.....	244	26.0	17.3	4.3	0	7.4	6.6	—	—	—	—	—	—	6	10.3	115.7
Nepal.....	262	16.3	9.6	2.5	0	—	—	—	—	—	—	—	—	4	7.1	74.7
Do.....	595	12.0	9.0	1.7	0	10.7	8.9	4.9	50.8	13.1	9.5	8.7	2.0	12	10.9	67.3
Hanna.....	24	29.0	23.8	1.0	0	10.7	2.7	—	—	—	—	—	—	7	9.6	114.3
Do.....	203	27.9	21.4	1.4	0	—	—	—	—	—	—	—	—	5	10.1	114.8
Manchuria.....	241	—	19.8	5.2	0	8.1	8.0	4.2	62.1	20.2	12.9	18.3	9.0	11	15.3	96.8
Odessa.....	182	—	22.1	8.1	0	9.8	8.7	3.5	69.4	26.1	18.3	20.0	8.0	11	17.6	111.4
Chevalier II.....	200	—	22.1	0	0	0	6.8	1.1	—	—	—	—	—	6	5.0	70.4
Hannchen.....	531	—	19.2	3.1	0	0	12.7	6.7	85.9	23.5	17.5	23.6	6.1	11	18.0	113.9
White Smyrna.....	658	—	—	—	—	10.7	14.3	14.4	76.2	21.6	23.3	18.4	9.2	8	23.9	123.8
Gatami.....	575	—	—	—	—	18.3	8.2	6.9	50.6	28.2	26.0	14.4	7.1	8	20.0	103.6
Coast.....	690	—	—	—	—	—	—	—	82.8	18.3	22.1	18.4	9.8	5	30.3	111.0
Himalaya.....	620	—	—	—	—	—	—	—	53.4	18.3	12.4	12.7	5.4	5	21.4	78.4
White Smyrna.....	195	—	—	—	—	—	—	—	77.9	—	—	—	—	1	77.9	114.2

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages									
			Chevalier II	Hanna	Hannchen	Manchuria	Odessa	Gatami	Coast	White Smyrna	Nepal	Himalaya
Chevalier II.....	200	Years comparable.....	—	6	6	6	6	3	(*)	3	6	(*)
		Percentage yield.....	—	78.5	71.9	66.2	57.5	23.7	—	20.1	85.2	—
Hanna.....	24	Years comparable.....	6	—	6	6	6	3	(*)	3	7	(*)
		Percentage yield.....	127.3	—	91.6	84.3	73.2	40.1	—	34.0	142.4	—
Hannchen.....	531	Years comparable.....	6	6	—	11	11	8	5	8	11	5
		Percentage yield.....	139.0	109.2	—	118.2	102.2	110.2	103.4	92.1	166.2	146.1
Manchuria.....	241	Years comparable.....	6	6	11	—	11	8	5	8	11	5
		Percentage yield.....	151.0	118.6	84.6	—	86.5	89.4	80.9	74.7	149.7	114.3
Odessa.....	182	Years comparable.....	6	6	11	11	—	8	5	8	11	5
		Percentage yield.....	174.0	136.6	97.8	115.6	—	102.6	93.7	85.7	162.6	132.3
Gatami.....	575	Years comparable.....	3	3	8	8	8	—	5	8	8	5
		Percentage yield.....	422.8	249.3	90.7	111.8	97.5	—	83.4	83.6	147.1	117.8
Coast.....	690	Years comparable.....	(*)	(*)	5	5	5	5	—	5	5	—
		Percentage yield.....	—	—	96.7	123.6	106.8	119.9	—	99.8	180.0	141.2
White Smyrna.....	658	Years comparable.....	3	3	8	8	8	8	5	—	8	—
		Percentage yield.....	498.7	294.0	108.6	133.8	113.7	119.7	100.2	—	176.0	141.5
Nepal.....	595	Years comparable.....	6	7	11	11	11	8	5	8	—	—
		Percentage yield.....	117.3	70.2	60.2	71.1	61.5	68.0	55.5	56.8	—	78.5
Himalaya.....	620	Years comparable.....	(*)	(*)	5	5	5	5	5	5	5	—
		Percentage yield.....	—	—	68.5	87.5	75.6	84.9	70.8	70.7	127.5	—

Irrigated land.—Barleys were grown under irrigation at the Belle Fourche Experiment Farm, Newell, S. Dak., in cooperation with the Office of Western Irrigation Agriculture from 1912 to 1921, inclusive. The yields of the 12 varieties tested during this period are presented in Table 47. Seven of these were grown for more than three years. This table shows that the 2-rowed varieties are high-yielding sorts under irrigation at Newell, as they also were under dry-land conditions there. Hannchen (C. I. No. 531) was surpassed by Chevalier II (C. I. No. 200) for the seven comparable years in which they were grown. Coast (C. I. No. 690), which was tested for 6 of the 10 years of the experiment, was excelled by both Chevalier II and Hannchen. Trebi (C. I. No. 936) produced the highest average yield of all varieties during the period from 1917 to 1921, inclusive. This has been a very productive variety at rather high altitudes under irrigation in the West. White Smyrna (C. I. No. 195), which was the highest yielding sort on dry land here, was tested only in 1921, when it gave but a fair return.

TABLE 47.—*Annual acre yield of varieties of barley grown on irrigated land at the Belle Fourche (Newell, S. Dak.) Experiment Farm in one or more of the 10 years from 1912 to 1921, inclusive*

[Data obtained in cooperation with the South Dakota Agricultural Experiment Station from 1912 to 1919, inclusive. Data for 1920 and 1921 used by courtesy of the Office of Western Irrigation Agriculture]

Variety	C. I. No.	Acre yields (bushels)										Years grown	Average yield (bus.)	Percentage of weighted mean
		1912	1913	1914	1915	1916	1917	1918	1919	1920	1921			
Manchuria	241	16.9										1	16.9	111.2
Odessa	182	17.4										1	17.4	114.5
Chevalier II	200	13.0		23.0	39.2	23.9	38.5	74.3	32.0	25.0	15.4	9	31.6	111.7
Hannchen	531	19.8	32.9	15.0		25.1	40.7	66.9	27.8		10.4	8	29.8	105.3
Nepal	595	9.1		19.2	20.4	14.9	22.4	51.6	18.4			7	22.3	74.1
Silver King	905		25.8	21.8	20.9	14.7	23.9	66.7	17.7			7	27.3	86.4
Manchuria	244		23.5	17.8								2	20.7	83.1
Chevalier	1142			26.8	37.2	23.2						3	29.1	128.8
Himalaya	620			29.6	23.4	17.5	23.7	62.7	17.5			6	29.1	91.2
Coast	690				23.0	15.2	37.9	65.6	25.4		7.1	6	29.0	93.2
Trebi	936						38.0	75.9	28.8	34.6	12.8	5	38.0	109.2
White Smyrna	195										11.7	1	11.7	101.7

LINCOLN, NEBR.

Results of the varietal experiments at Lincoln, Nebr., were obtained through the courtesy of the Nebraska Agricultural Experiment Station and are presented in Table 48. Barley experiments were conducted at three distinct periods, in 1900 and 1901, from 1905 to 1912, inclusive, and from 1917 to 1921, inclusive. Three varieties were tested for both years in the first period. Of these, Black Hull-less (C. I. No. 596) produced the highest average, with Russian Naked (C. I. No. 2815) second and Sisolsk (C. I. No. 89) third. This latter variety was originally grown in the extremely cold regions of northern Russia, so it is not surprising that it was surpassed by these hull-less sorts, even though they are low-yielding barleys.

Eight varieties were grown in the second period, among which Black Hull-less and Russian Naked again appear. These two barleys were grown in the years 1908 to 1910, inclusive. Black Hull-less was still the better of the two, although they were both surpassed by Oderbrucker (C. I. No. 1517). This latter variety was grown for four years, 1907 to 1910, inclusive, producing an average yield of 135 per cent of the weighted mean. Tennessee Winter (C. I. No. 257) was the only variety grown in all the eight years of the second period. It produced very high yields for the first four years but was a complete failure during the last four. The winters are usually too severe for the successful cultivation of winter barley in Nebraska. Four varieties from an English seed firm were introduced in 1909. None of these seemed promising, and with the exception of Garton (C. I. No. 2816), which was grown again in 1910, they were dropped from the experiment.

TABLE 48.—Annual acre yields of barley grown at the Nebraska Agricultural Experiment Station (at Lincoln) in one or more of the 22 years from 1900 to 1921, inclusive

[Data obtained through the courtesy of the Nebraska Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)												Years grown	Aver- age of yield (bus.)	Percent- age of weighted mean
		1900	1901	1905	1906	1907	1908	1909	1910	1911	1912	1917	1918			
Sisolsk	89	18.9	3.5													54.8
Black Hull-less	596	25.4	4.0				28.9	34.9	44.5							11.2
Russian Naked	2815	23.8	3.7				14.9	24.0	33.7							27.5
Tennessee Winter	257			58.5	65.3	68.0	49.4	0	0	0	0					100.3
Oderbrucker	1517					41.1	39.6	44.3	46.6							30.1
Garton 5004	2816							27.5	44.5							93.6
Garton 5003	2817							23.5								135.3
Garton 5002	2818							17.6								36.0
Garton 5001	2819															130.0
Manchuria	2820															107.1
Do.	2821															73.9
Gold	1145							16.8								70.6
Manchuria	2330										57.7	0	19.8	47.9	18.6	82.3
Club Marabout	261										33.1	0	13.2	41.5	19.5	93.0
Acc.	1653										61.2	0				25.9
O. A. C. 21	1470												18.5	49.5	19.8	23.6
Odessa	182													60.7	23.9	128.6
Gatami	575													49.8	24.3	37.1
Oderbrucker	1329													49.4	22.3	112.8
Svanhals	187													48.5	19.3	35.9
White Smyrna	195													45.5	16.2	109.1
Chevalier	1886													44.5	10.2	93.9
Wing Pedigree	1177													43.7	26.7	97.0
Horsford	507													43.7	26.7	31.9
Hannchen	531													41.7	29.5	35.2
Manchuria	1562													40.0	20.4	108.2
														35.3	21.5	35.6
														33.3	21.5	91.8
														33.5	20.0	86.3
														29.2	20.0	28.4
														28.3	21.5	66.0
														27.5	20.0	86.0
														42.9	20.0	95.7

Two varieties, Manchuria (C. I. Nos. 1580 and 2820), were grown for the full five years of the third period. In 1917 they were both surpassed by Gold (C. I. No. 1145). All three of these were complete failures in 1918, and Gold was not grown again. In 1919 Manchuria (C. I. No. 2330), a pedigreed sort, was introduced and was continued through 1921. These three Manchuria barleys, however, were surpassed by a number of varieties grown only in 1920 and 1921. Club Mariout (C. I. No. 261), a North African barley, produced the highest average yield, 42.3 bushels, for those years. Ace (C. I. No. 1853) was second in point of yield. This is a 2-rowed variety of the Smyrna type. Following Ace is O. A. C. 21 (C. I. No. 1470), another Manchuria barley. Fourth and fifth in rank are two 2-rowed sorts, White Smyrna (C. I. No. 195) and Svanhals (C. I. No. 187).

From the results of these three periods it is rather difficult to say which types are best adapted here. From the yields in recent years Lincoln appears to be on the border between the districts where the barleys of the Manchuria group and those of the North African and Asia Minor groups can be successfully grown.

TABLE 49.—*Annual acre yields of varieties of barley grown at the North Platte (Nebr.) substation in one or more of the 13 years from 1909 to 1921, inclusive*

[Data obtained through the courtesy of the Nebraska Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)												Years grown	Average yield (bus.)	Percentage of weighted mean
		1909	1910	1912	1913	1914	1915	1916	1917	1918	1919	1920	1921			
Coast	690	19.0	7.9	14.7	6.3	11.4	29.7	58.6	25.5	12.0	25.5	40.3	11.3	12	21.9	107.4
Manchuria	244	19.2	4.4	—	—	—	—	—	—	—	—	—	—	2	11.8	106.3
Black Hull-less	596	18.6	4.6	14.0	—	—	—	—	—	—	—	—	—	3	12.4	108.8
Hanna	24	14.8	—	—	—	8.3	3.5	—	—	—	—	—	—	3	6.4	61.5
Beardless	639	15.2	5.4	8.5	1.0	8.4	30.0	—	—	—	—	—	—	6	11.4	88.4
Nepal	595	9.2	4.3	—	—	—	—	—	—	—	—	—	—	2	6.7	60.4
Stavropol	2103	—	—	—	5.2	15.0	27.8	58.2	29.1	12.0	23.4	39.0	10.3	9	24.4	106.5
Wisconsin Pedigree	835	—	—	—	—	8.7	30.6	54.2	24.4	5.5	21.1	127.3	10.3	8	22.8	93.1
Horsford	877	—	—	—	—	—	22.7	38.9	24.7	10.1	120.7	36.7	9.2	7	23.3	88.6
White Smyrna	658	—	—	—	—	—	66.6	28.5	12.2	23.1	136.4	7.7	—	6	29.1	111.5
Gatani	575	—	—	—	—	—	—	—	—	—	18.4	24.9	—	2	21.7	71.9
Trebi	936	—	—	—	—	—	—	—	—	—	45.2	10.9	—	2	28.1	120.6
McClymont	2126	—	—	—	—	—	—	—	—	—	41.3	11.2	—	2	26.3	112.9
Manchuria	2330	—	—	—	—	—	—	—	—	—	—	10.2	—	1	10.2	101.0

NORTH PLATTE, NEBR.

Through the courtesy of the Nebraska Agricultural Experiment Station the records of barley yields for the years 1909, 1910, and 1912 to 1921, inclusive, for the North Platte substation are presented in Table 49. As only a few of the varieties were grown during the same years or over a long period, a varietal digest would mean little or nothing. Only one variety, Coast (C. I. No. 690), was grown for the entire period. The five others grown in 1909 were all dropped from the experiment by 1916. Two of these, Manchuria (C. I. No. 244) and Hanna (C. I. No. 24), are humid-area spring varieties and, as might be expected, produced very poor results. The other three, Black Hull-less (C. I. No. 596), Beardless (C. I. No. 639), and Nepal (C. I. No. 595), were inferior, as is usually the case wherever they are grown. Stavropol (C. I. No. 2103) was introduced in 1913 and was grown through 1921. During these nine years it produced nearly the same average yield as did Coast (C. I. No. 690). While somewhat different in appearance, these barleys belong to the same ecological group and are almost equally well suited to growing conditions here. Wisconsin Pedigree (C. I. No. 835), another Manchuria barley, was grown from 1914 to 1921, inclusive. It was surpassed by both Coast and Stavropol for the period. Horsford (C. I. No. 877) was grown for seven years, 1915 to 1921, inclusive, but was inferior to the three varieties first mentioned. White Smyrna (C. I. No. 658) produced the highest average yield of all the barleys grown for the six years, 1916 to 1921, inclusive. It will be remembered that in the western part of North Dakota and South Dakota the 2-rowed varieties White Smyrna and Hannchen are very productive. Hannchen, however, usually ripens a little later than White Smyrna and, although not tested at North Platte, probably would not be so successful. Gatani (C. I. No. 575) was the poorest sort in both 1919 and 1920 and was dropped from the experiment. Two varieties, Trebi (C. I. No. 936)

and McClymont (C. I. No. 2126), were grown in 1920 and 1921. Trebi produced the highest average of all varieties for these two years and McClymont, a local barley, was second. Manchuria (C. I. No. 2330), grown in 1921, was superior only to Horsford and White Smyrna. Although it exceeded White Smyrna in this one year, it is apparent that when grown for several years the barleys of the Manchuria group can not compete with such varieties as Coast, Stavropol, or White Smyrna which are adapted to more arid regions.

There is no doubt as to the types of barley best suited to this section. White Smyrna is a high-yielding variety, but both Coast and Stavropol yield almost equally well and are preferable for general cultivation. They are the common barleys grown here.

MITCHELL, NEBR.

The records of yields of varieties of barley grown at the Scottsbluff Experiment Farm, Mitchell, Nebr., during the years 1914 to 1919, inclusive, were received through the courtesy of the Office of Western Irrigation Agriculture and are reported in Table 50. In 1914 12 varieties were tested, all of which were continued through 1916 with the exception of White Smyrna (C. I. No. 195). In 1915 this variety produced a relatively poor yield, 53.2 bushels, when all the other varieties yielded from 77.6 to 103.1 bushels. Of the 11 varieties grown during the years 1914 to 1916, inclusive, White Moravian (C. I. No. 977) ranks first. This is a 2-rowed barley of the Hanna group. Barbary (C. I. No. 695), a Coast barley, ranks next. It was continued through 1919, with an average yield of 68.1 bushels for the six years of the experiment. Barbary was the only variety grown for this entire period. Han River (C. I. No. 206) was third in rank for the three years 1914, 1915, and 1916. Svanhals (C. I. No. 187) was also a high-yielding sort for these years, but was exceeded in 1917 and 1918 by both Barbary and Trebi. The latter was introduced in 1917 and was continued through 1919. It produced the highest yield of any variety grown in the comparable years. Trebi, both in these and other tests at Scottsbluff, was so evidently a superior sort that it was distributed to farmers. On the farms it has given even better relative yields, and according to James A. Holden, superintendent of the Scottsbluff farm, 85 per cent of the acreage in that vicinity is of Trebi. Yields of 75 to 80 bushels to the acre are not uncommon.

Both the Coast and 2-rowed types have yielded well here, but the 6-rowed barleys are the ones usually grown.

TABLE 50.—*Annual acre yields of varieties of barley grown at the Scottsbluff Experiment Farm, Mitchell, Nebr., in part or all of the six years from 1914 to 1919, inclusive*

[Data furnished through the courtesy of the Office of Western Irrigation Agriculture]

Variety	C. I. No.	Acre yields (bushels)						Years grown	Average yield (bus.)	Percentage of weighted mean
		1914	1915	1916	1917	1918	1919			
Hannchen	531	51.8	103.1	40.5	-----	-----	-----	3	65.1	98.8
White Moravian	977	91.6	102.2	37.0	-----	-----	-----	3	76.9	116.7
Franconian	680	61.0	90.1	47.1	56.2	-----	-----	4	63.6	97.4
Svanhals	187	64.0	77.6	67.8	60.4	56.9	-----	5	65.3	99.5
White Smyrna	195	65.1	53.2	-----	-----	-----	-----	2	59.1	78.0
Han River	206	72.9	102.0	47.4	-----	-----	-----	3	74.1	112.4
Barbary	695	72.2	101.0	49.3	63.2	71.2	51.7	6	68.1	104.8
Caucasian	90	29.1	100.5	57.4	-----	-----	-----	3	62.3	94.5
Scotch	848	62.8	99.8	36.8	-----	-----	-----	3	66.5	100.9
Coast	690	49.7	97.6	35.3	-----	-----	-----	3	60.9	92.4
Club Mariout	261	53.9	93.6	40.7	-----	-----	-----	3	62.7	95.1
Manchuria	244	37.0	88.5	47.9	-----	-----	-----	3	57.8	87.7
Thomas	2501	-----	-----	44.4	-----	-----	-----	1	44.4	96.5
Trebi	936	-----	-----	-----	79.4	79.5	53.2	3	70.7	114.4
Nepal	595	-----	-----	-----	46.5	-----	-----	1	46.5	76.1

MANHATTAN, KANS.

The barley yields at Manhattan, Kans., were obtained in cooperation with the Kansas Agricultural Experiment Station. Several varieties were grown in plat tests at this station in periods of varying length during the years 1903 to 1912, inclusive. The varieties were not sufficient in number and were not grown in plats for a period of sufficient length to be of value in this discussion. Ten-

nessee Winter (C. I. No. 257) produced very high yields in 1904, 1905, and 1906, but it has never attained commercial importance in this section, because of the danger of winterkilling.

During the 10-year period from 1912 to 1921, inclusive, numerous varieties were grown in replicated nursery rows. Here, as at Cornell University Agricultural Experiment Station, it is thought that replicated row tests are adequate for the purposes of this discussion. From 1918 to 1921, inclusive, however, some of the varieties reported in Table 51 were replicated only three times or grown in only a single row. These yields are believed to be of sufficient importance to warrant including them, but the conclusions drawn from the varietal comparison may not be so accurate as in the case of some of the other stations.

The summarized digest of Table 51 shows a comparison of 10 representative varieties. Coast (C. I. No. 690) is the best of these, but as it was grown only three years its rank is not so definite as that of the others compared. O. A. C. 21 (C. I. No. 1470), a pedigreed Manchuria barley, was exceeded only by Coast. Third in point of yield is Odessa (C. I. No. 182), followed by Manchuria (C. I. No. 1472). Horsford (C. I. No. 1474), which is usually one of the lowest yielding sorts, is better than the remaining four in this group. White Smyrna (C. I. No. 195) is quite inferior at Manhattan.

TABLE 51.—*Annual acre yields of varieties of barley grown at the Kansas Agricultural Experiment Station (at Manhattan) in one or more of the 10 years from 1912 to 1921, inclusive*

[Data obtained in cooperation with the Kansas Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)										Years grown	Average yield (bus.)	Percentage of weighted mean
		1912	1914	1915	1916	1917	1918	1919	1920	1921				
Manchuria	1473	13.5	20.8	31.1	70.2	57.9	30.5	42.3	15.7	10.7	9	32.5	110.9	
Black Hull-less	596	9.4	11.8	8.3	42.0	25.7	12.0	27.4	11.7	4.6	9	17.0	58.0	
Black Smyrna	191	28.6	44.8	31.7	57.5	35.0					5	39.5	108.8	
Horsford	1807	8.8	22.2	14.1	40.2	28.1	18.0	31.5	18.4	7.8	9	21.0	71.7	
Do	1778	19.0	22.9	9.3	45.1	43.7	21.1	33.3	14.8	5.5	9	23.9	81.6	
Wisconsin Winter	519	17.4	33.1	14.8	56.9	55.2	25.8	49.1	16.1	10.1	9	30.9	105.5	
Bonanza	1800	9.7	22.8	19.2	53.8	42.7	21.0	47.3	14.1	6.4	9	26.3	89.8	
Horsford	3171	6.7	19.5	10.1							3	12.1	55.8	
Manchuria	3172		20.0	20.7	62.0	48.2					4	37.7	98.9	
Odessa	182		31.3	42.7	57.5	65.3	23.3	38.9	15.4	8.7	8	35.4	118.0	
Manchuria	1472		31.3	26.3	60.0	48.2	28.5	37.9	16.1	13.2	8	32.7	109.0	
Do	3173		18.0	11.4							2	14.7	63.1	
Do	2181		29.9	28.1	59.0						3	39.0	105.7	
Do	3174		24.2	18.4							2	21.3	91.4	
Do	2182		22.2	41.4	55.1	57.6					4	44.1	115.7	
Do	3175		27.3	17.7							2	22.5	96.6	
Bonanza	3176		25.0	13.9							2	19.5	83.7	
Success	2183		43.0	9.0	42.3						3	31.4	85.1	
Horsford	1474		22.2	18.0	56.9	45.5	24.8	41.4	17.6	11.4	8	29.7	99.0	
Bonanza	3177		25.4	27.0	59.7						3	37.4	101.4	
Swedish Star	1701				58.2	36.8	10.6				3	35.2	83.0	
Gold	1145				54.8	40.6	20.9	31.3	9.8		5	31.5	88.0	
Oderbrucker	888				78.4						1	78.4	146.8	
Manchuria	244				61.1			45.9	17.9	14.2	4	34.8	109.4	
Hannchen	531				41.6						1	41.6	77.9	
Chevalier II	200				55.2						1	55.2	103.4	
White Smyrna	658				36.7	37.8					2	37.3	76.0	
Gatami	575				51.7	48.9	17.9	34.4	23.2	14.0	6	31.7	98.8	
Nepal	595				31.7	17.7	10.0	19.3	7.9		5	17.3	48.3	
Himalaya	620				39.9						1	39.9	74.7	
Mensury	2657				43.0						1	43.0	80.5	
White Smyrna	195				70.0	28.1	23.8	32.3	10.0	8.7	6	28.8	89.7	
Chevalier	2178				51.3						1	51.3	96.1	
Silver King	890				57.1						1	57.1	106.9	
Manchuria	1275				62.1	50.7	19.1	38.2	15.7	10.4	6	32.7	101.9	
Garton 68	2740				62.9	18.4					2	40.7	82.9	
O. A. C. 21	1470				68.7	60.7	26.8	32.5	15.1	10.8	6	35.8	111.5	
Svanhals	187						19.5	24.2	3.8		4	16.5	48.1	
Coast	690							45.9	14.6	12.5	3	24.3	119.1	
Horsford	507							27.0	13.8	8.9	3	16.6	81.4	
Himalaya	1585							40.0	8.6	5.6	3	18.1	88.7	
Jet	1928							40.0	14.2	14.5	3	22.9	112.3	
Club Mariout	261								19.8	19.9	2	19.9	159.2	
California Mariout	1455									14.7	1	14.7	138.7	

TABLE 51.—*Annual acre yields of varieties of barley grown at the Kansas Agricultural Experiment Station (at Manhattan) in one or more of the 10 years from 1912 to 1921, inclusive—Continued*

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages									
			Gold	Manchuria	O. A. C. 21	Gatami	Odessa	Coast	White Smyrna	Nepal	Black Hull-less	Horsford
Gold-----	1145	Years comparable-----	-----	5	5	5	5	2	5	5	5	5
		Percentage yield-----	-----	82.5	77.2	89.4	78.5	67.9	95.9	181.8	132.5	84.5
Manchuria----	1472	Years comparable-----	-----	5	6	6	8	3	6	5	8	8
		Percentage yield-----	-----	121.2	95.0	107.3	92.4	92.1	117.9	220.2	182.2	110.0
O. A. C. 21----	1470	Years comparable-----	-----	5	6	6	6	3	6	5	6	6
		Percentage yield-----	-----	129.5	105.2	112.9	102.6	80.0	124.1	235.3	173.9	108.6
Gatami-----	575	Years comparable-----	-----	5	6	6	6	3	6	5	6	6
		Percentage yield-----	-----	111.9	93.2	88.6	90.9	98.1	109.9	203.3	154.1	96.2
Odessa-----	182	Years comparable-----	-----	5	8	6	6	3	6	5	8	8
		Percentage yield-----	-----	127.3	108.3	97.4	110.0	86.3	120.9	231.4	197.3	119.0
Coast-----	690	Years comparable-----	-----	2	3	3	3	3	3	2	3	3
		Percentage yield-----	-----	147.2	108.6	125.0	102.0	115.9	143.1	222.4	167.0	103.7
White Smyrna	195	Years comparable-----	-----	5	6	6	6	3	3	5	6	6
		Percentage yield-----	-----	104.3	84.8	80.6	91.0	82.7	69.9	189.6	140.1	87.5
Nepal-----	595	Years comparable-----	-----	5	5	5	5	2	5	5	5	5
		Percentage yield-----	-----	55.0	45.4	42.5	49.2	42.3	45.0	52.7	72.9	46.5
Black Hull-less	596	Years comparable-----	-----	5	8	6	8	3	6	5	5	8
		Percentage yield-----	-----	75.5	54.9	57.5	64.9	59.9	72.4	137.2	100.3	60.3
Horsford-----	1474	Years comparable-----	-----	5	8	6	8	3	6	5	8	8
		Percentage yield-----	-----	118.3	90.9	92.1	103.9	84.0	96.4	215.0	165.7	-----

As shown in Table 51 four varieties were added in 1919. Coast was the best of these. Jet (C. I. No. 1928), a black hull-less sort, was second. Club Mariout (C. I. No. 261), grown in 1920 and 1921, produced the highest average yield of any barley grown in those years. Most of the varieties having an average yield of more than 100 per cent of the weighted mean, however, are of the Manchuria group. Although barley is not of commercial importance in this section of the State, probably Odessa and Manchuria are the types best adapted here.

McPHERSON, KANS.

Barley was grown at the McPherson Experiment Farm in cooperation with the Kansas Agricultural Experiment Station from 1906 to 1909, inclusive. The results are shown in Table 52. Of the varieties grown for the entire period Caucasian (C. I. No. 90) was the highest yielding sort. It is probably identical with Stavropol. Yenidje (C. I. No. 205), second in yield, is a barley which resembles the Coast type, although it originated in Asiatic Turkey. Odessa (C. I. No. 182), which also produced high yields at Manhattan, ranks third at this station. White Smyrna (C. I. No. 195) and Black Smyrna (C. I. No. 191) follow in the order named. Coast (C. I. No. 549) also yielded well. At Manhattan the barleys of the Manchuria and Odessa types produced the best yields of any tested over a period of years. At McPherson barleys from southern Russia, Asia Minor, and North Africa are the superior sorts, although several of the Manchuria barleys, such as Common (C. I. No. 184) and Baxter (C. I. No. 185), still do well.

TABLE 52.—*Annual acre yields of varieties of barley grown at the McPherson (Kans.) Experiment Farm in some or all of the four years from 1906 to 1909, inclusive*

[Data obtained in cooperation with the Kansas Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1906	1907	1908	1909			
Perm.....	20	30.2	4.0	25.1	20.9	4	20.1	102.0
Caucasian.....	90	51.5	10.4	33.9	22.5	4	29.6	150.3
Rennie Improved.....	181	28.7	3.5	23.9	23.1	4	19.8	100.5
Odessa.....	182	44.5	9.9	30.5	27.1	4	28.0	142.1
Nepean.....	183	37.8	4.0	26.2	-----	3	22.7	115.2
Common.....	184	36.2	5.8	28.2	21.7	4	23.0	116.8
Baxter.....	185	33.1	4.1	26.8	24.2	4	22.1	112.2
Beldi Dwarf.....	190	23.5	2.1	15.3	-----	3	13.6	69.0
Black Smyrna.....	191	48.9	10.5	17.1	19.4	4	24.0	121.8
White Smyrna.....	195	50.6	3.3	22.3	22.8	4	24.7	125.4
Albacete.....	199	45.4	9.1	16.8	16.4	4	21.9	111.2
Yenidje.....	205	51.8	4.8	32.5	25.3	4	28.6	145.2
Baku.....	210	28.2	.6	15.6	-----	3	14.8	75.1
Black Hull-less.....	596	30.4	2.1	15.7	11.9	4	15.0	76.1
Success.....	212	27.3	2.0	18.3	18.0	4	16.4	83.2
Hanna.....	203	40.7	.7	16.4	19.4	4	19.3	98.0
Manchuria.....	235	31.1	5.6	14.1	25.2	4	19.0	96.4
Do.....	237	35.2	4.7	15.3	23.6	4	19.7	100.0
Do.....	240	30.5	4.9	14.5	23.6	4	18.4	93.4
Club Mariout.....	261	28.3	3.4	25.4	7.9	4	16.3	82.7
Garton.....	263	32.2	2.5	4.9	-----	3	13.2	67.0
Do.....	264	29.7	.8	4.1	-----	3	11.5	58.4
Do.....	265	31.3	-----	-----	-----	1	31.3	87.4
Horsford.....	507	26.1	3.9	12.7	20.8	4	15.9	80.7
Coast.....	549	40.9	4.1	25.4	15.8	4	21.5	109.1
Gatani.....	575	-----	-----	12.8	23.1	2	17.9	94.7
Svanhals.....	187	-----	-----	8.1	17.3	2	12.7	67.2
Primus.....	532	-----	-----	2.2	4.4	2	3.3	17.5
Princess.....	529	-----	-----	13.5	13.7	2	13.6	72.0

HAYS, KANS.

In cooperation with the Kansas Agricultural Experiment Station, barley yields were obtained at the Fort Hays branch station in 12 of the 19 years from 1903 to 1921, inclusive. These are presented in Table 53, and a comparison of the yields of 10 representative varieties is also shown. The best of these was Ellis (C. I. No. 2107), a local barley. It is a mixture of several strains, with Coast types predominant. Club Mariout (C. I. No. 261), Stavropol (C. I. No. 2105), and Coast (C. I. No. 690), followed in the order named. Although Stavropol came from southern Russia, it is of the same general group as Coast and Club Mariout. The latter was very badly mixed at this station and its yields may mean little. Fifth in yield is Beldi Dwarf (C. I. No. 190), a North African barley. While White Smyrna (C. I. No. 195) ranks only sixth in this varietal comparison, it produced an average yield of 117 per cent of the weighted mean for the 11 years in which it was grown. The Manchuria barleys and Hannchen (C. I. No. 531) are not adapted to western Kansas, the climate there being too dry for the successful growing of spring barleys from humid climates.

Table 53 shows that several winter varieties were tested during the period from 1913 to 1916, inclusive. Tennessee Winter (C. I. No. 257) was the best of these. However, the winters here are usually too severe for the successful growing of these sorts. In 1920 five additional spring varieties were introduced. Two of these, Manchuria (C. I. Nos. 1275 and 1473), produced very poor yields and were not included in 1921. Meloy (C. I. No. 1176) averaged 27.5 bushels for the two years. This is a fairly good yield for a hooded barley, but Meloy has ranked relatively high at a number of the other stations. It is the only hooded spring barley that has proved at all successful in any of the experiments reported in the drier sections of the West. Odessa (C. I. No. 182) was not promising in 1920 and 1921, the two years in which it was grown. Of those tested, the Coast-Stavropol barleys are unmistakably the types best adapted to this region.

TABLE 53.—*Annual acre yields of varieties of barley grown at the Fort Hays (Kans.) branch station in one or more of the years from 1903 to 1921, inclusive—Continued.*

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

Variety	C. I. No.	Data shown	Varieties and percentages									
			Austrian	Hann-chen	Man-churia	Gatami	Coast	Stav-ropol	Beldi Dwarf	Club Mariout	White Smyrna	Ellis
Austrian.....	2185	{ Years comparable { Percentage yield.....	2	297.0	247.5	4	4	4	4	4	3	4
Hannchen.....	531	{ Years comparable { Percentage yield.....	2	33.7	118.3	94.5	86.4	82.7	96.6	80.7	79.2	80.5
Manchuria.....	244	{ Years comparable { Percentage yield.....	2	4	84.5	77.7	58.9	66.5	71.9	63.4	85.9	3
Gatami.....	575	{ Years comparable { Percentage yield.....	4	105.8	130.8	76.4	53.1	56.5	61.8	55.0	62.1	48.6
Coast.....	690	{ Years comparable { Percentage yield.....	4	128.8	188.4	9	80.2	80.3	86.5	78.1	87.0	47.7
Stavropol.....	2105	{ Years comparable { Percentage yield.....	4	169.8	177.1	124.8	9	99.8	108.4	97.9	103.2	8
Beldi Dwarf.....	190	{ Years comparable { Percentage yield.....	4	150.4	161.9	115.6	100.2	10	107.8	97.3	107.2	89.9
Club Mariout.....	261	{ Years comparable { Percentage yield.....	4	139.1	181.9	128.0	92.3	10	107.8	90.3	101.5	8
White Smyrna.....	195	{ Years comparable { Percentage yield.....	8	157.7	161.1	114.9	102.2	10	110.8	9	109.3	80.6
Ellis.....	2107	{ Years comparable { Percentage yield.....	4	124.3	202.7	131.6	111.2	8	93.3	91.5	7	93.7
			124.3	203.9				8	8	106.7	108.5	92.2
								103.3	111.6			

COLBY, KANS.

Tests of barley varieties were carried on at the Colby substation during the four-year period from 1918 to 1921, inclusive. The data found in Table 54 were furnished through the courtesy of the Kansas Agricultural Experiment Station. Six varieties were grown for the entire period, of which Odessa (C. I. No. 182) was the best and Ellis (C. I. No. 2107) almost its equal. Coast (C. I. No. 690) and White Smyrna (C. I. No. 195) also produced good yields. It is evident from their yields that Manchuria (C. I. No. 244) and Gatami (C. I. No. 575) are not adapted to this section. Horsford (C. I. No. 507) was grown only in 1918 and 1919 and was then dropped, doubtless because it is a low-yielding sort. In 1921 seven other barleys were tested. Of these Meloy (C. I. No. 1176) was the best, being exceeded only by Odessa (C. I. No. 182). Stavropol (C. I. No. 2103) produced a yield slightly lower than Meloy, while Flynn (C. I. No. 1311) and Hannchen (C. I. No. 531) followed in the order named. Flynn is a smooth-awned hybrid and is one of the high-yielding smooth-awned barleys. Club Mariout (C. I. No. 261), Blackhull (C. I. No. 878), and California Mariout (C. I. No. 1455) were not outstanding in yield. It is again apparent that the barleys best adapted to this region are of the southern Russian, Asia Minor, and North African types.

TABLE 54.—*Annual acre yields of varieties of barley grown at the Colby (Kans.) substation in part or all of the four years from 1918 to 1921, inclusive*

[Data obtained through the courtesy of the Kansas Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1918	1919	1920	1921			
Ellis.....	2107	20.2	53.6	20.9	18.4	4	28.3	125.8
White Smyrna.....	195	12.7	42.6	24.9	17.7	4	24.5	108.9
Gatami.....	575	8.6	43.6	7.1	19.0	4	19.6	87.1
Odessa.....	182	9.6	53.9	19.7	30.3	4	28.4	126.2
Manchuria.....	244	1.5	49.2	9.7	19.6	4	20.0	88.9
Coast.....	690	7.1	47.1	26.1	19.6	4	25.0	111.1
Horsford.....	507	9.6	16.4	-----	-----	2	13.0	48.5
California Mariout.....	1455	-----	-----	-----	10.2	1	10.2	51.3
Meloy.....	1176	-----	-----	-----	23.4	1	23.4	117.6
Stavropol.....	2103	-----	-----	-----	23.3	1	23.3	117.1
Hannchen.....	531	-----	-----	-----	20.0	1	20.0	100.5
Flynn.....	1311	-----	-----	-----	22.0	1	22.0	110.6
Blackhull.....	878	-----	-----	-----	17.1	1	17.1	85.9
Club Mariout.....	261	-----	-----	-----	17.6	1	17.6	88.4

TABLE 55.—*Annual acre yields of varieties of barley grown at the Garden City (Kans.) substation in part or all of the three years 1919, 1920, and 1921*

[Data furnished through the courtesy of the Kansas Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)			Years grown	Average yield (bus.)	Percentage of weighted mean
		1919	1920	1921			
Coast.....	690	8.1	13.6	28.9	3	16.9	109.7
Odessa.....	182	7.5	22.6	20.6	3	16.9	109.7
Manchuria.....	244	6.3	24.6	8.3	3	13.1	85.1
Gatami.....	575	1.3	13.3	-----	2	7.3	57.0
White Smyrna.....	195	8.1	14.1	33.0	3	18.4	119.5
Club Mariout.....	261	-----	21.4	-----	1	21.4	116.9

GARDEN CITY, KANS.

The annual yields of barley varieties at the Garden City substation were furnished through the courtesy of the Kansas Agricultural Experiment Station. These are reported in Table 55 for the years 1919, 1920, and 1921. Only four varieties were tested in all three years. Of these White Smyrna (C. I. No. 195) produced the highest average yield. Coast (C. I. No. 690) and Odessa (C. I.

No. 182) follow, each having an average yield of 109.7 per cent of the weighted mean. While Manchuria (C. I. No. 244) was the best in 1920 it was very low in 1919 and 1921, making the average yield quite low as compared with the other three varieties. Gatami (C. I. No. 575) was grown in 1919 and 1920, but was inferior to all others in both years. Club Mariout (C. I. No. 261), with a yield of 21.4 bushels, ranked third of the six varieties tested in 1920, the only year it was grown. White Smyrna has yielded well at nearly all of the Kansas stations, but because of its very short straw it probably is not so good for commercial purposes as either Odessa or Coast.

TRIBUNE, KANS.

The annual yields of barley varieties grown at the Tribune (Kans.) substation were furnished through the courtesy of the Kansas Agricultural Experiment Station. Barley was grown here for only the four years, 1918 to 1921, inclusive (Table 56). In 1921 the crop was a complete failure. Coast (C. I. No. 690) was the best in the four-year period. Next in order was Ellis (C. I. No. 2107), a local barley of the Coast type, with Odessa (C. I. No. 182) third. White Smyrna (C. I. No. 195) was the highest yielding variety in 1920, but produced very low yields in the two previous years. Club Mariout (C. I. No. 261) was grown only in 1920, but ranked second among the varieties grown that year. At practically all the stations in western Kansas these five, together with Stavropol, have been the barleys outstanding in yield. At Tribune, as at the other stations in western Kansas, Manchuria (C. I. No. 244) and Gatami (C. I. No. 575) are in a region unsuited to their culture. While several barleys showed up well at these stations, only Coast and Stavropol are grown commercially.

TABLE 56.—*Annual acre yields of varieties of barley grown at the Tribune (Kans.) substation in one or more of the four years from 1918 to 1921, inclusive*

[Data furnished through the courtesy of the Kansas Agricultural Experiment Station]

Variety	C. I. No.	Acre yield (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1918	1919	1920	1921			
Odessa.....	182	14.1	21.0	38.8	-----	3	24.6	102.1
Gatami.....	575	13.1	16.2	25.1	-----	3	18.1	75.1
White Smyrna.....	195	2.4	17.7	49.4	0	4	17.4	87.0
Manchuria.....	244	2.3	20.2	18.8	0	4	10.3	51.5
Coast.....	690	23.6	25.3	47.6	0	4	24.1	120.5
Ellis.....	2107	27.1	14.9	43.6	0	4	21.4	107.0
Horsford.....	507	11.9	-----	-----	-----	1	11.9	88.1
Club Mariout.....	261	-----	-----	47.9	-----	1	47.9	123.8

HAVRE, MONT.

Barley varieties have been tested at the northern Montana substation at Havre during the 6-year period from 1916 to 1921, inclusive. The yields, presented in Table 57, were obtained in cooperation with the Montana Agricultural Experiment Station. Four varieties were first grown in 1916, two of which, White Smyrna (C. I. No. 195) and Hannchen (C. I. No. 531), were continued through the six years of the experiment. White Smyrna produced the higher average yield. Coast (C. I. No. 690) was grown from 1916 to 1920, inclusive, but also was surpassed by White Smyrna. Nepal (C. I. No. 595) was inferior to the other three varieties during the five years from 1916 to 1919, inclusive, and in 1921.

In 1917 eight varieties were introduced, all of which were continued in the experiment through 1921. Horn (C. I. No. 926), a 2-rowed variety of the Chevalier type, was superior to all the other sorts. Following Horn are three other lax 2-rowed barleys, Bohemian (C. I. No. 27), Goldfoil (C. I. No. 928), and Holland (C. I. No. 952). These four in this period excelled both White Smyrna and Hannchen as well as all of the 6-rowed sorts. Han River (C. I. No. 206), which was the best of the 6-rowed barleys, surpassed the North African varieties, Beldi Giant (C. I. No. 2777) and Sandrel (C. I. No. 937), by only a small margin. Trebi (C. I. No. 936) was not at all suited to the conditions which prevailed during the years it was grown.

Without doubt, this is a district favorable to the cultivation of 2-rowed barleys. The Chevalier and Hanna varieties, which seem to be superior types here, differ somewhat from Hannchen and White Smyrna in that they require a somewhat longer but cooler growing season. At Lethbridge, Alberta, Canada, the second highest yielding variety grown for a period of years also was a barley of the Chevalier group, namely, Chevalier II (C. I. No. 200).

TABLE 57.—*Annual acre yields of varieties of barley grown at the northern Montana substation (at Havre) in part or all of the six years from 1916 to 1921, inclusive*

[Data obtained in cooperation with the Montana Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)						Years grown	Average yield (bus.)	Percentage of weighted mean
		1916	1917	1918	1919	1920	1921			
White Smyrna.....	195	61.7	16.7	12.0	1.5	8.3	12.7	6	18.8	142.4
Coast.....	690	63.8	8.6	9.6	3.4	10.7	---	5	19.2	140.1
Hannchen.....	531	55.4	13.0	15.4	3.5	6.9	9.7	6	17.3	131.1
Nepal.....	595	43.7	9.0	11.9	3.5	---	8.5	5	15.3	115.0
Trebi.....	936	---	8.3	14.2	1.7	8.6	9.4	5	8.4	81.6
Han River.....	206	---	12.0	12.7	3.0	12.8	6.6	5	9.4	91.3
Horn.....	926	---	13.0	17.7	3.0	22.9	15.3	5	14.4	139.8
Goldfoil.....	928	---	11.0	15.4	3.6	13.1	15.4	5	11.7	113.6
Beldi Giant.....	2777	---	12.5	12.7	2.8	10.4	7.1	5	9.1	88.3
Bohemian.....	27	---	13.0	14.8	3.5	15.9	12.1	5	11.9	115.5
Sandrel.....	937	---	11.0	10.2	2.6	12.5	10.4	5	9.3	90.3
Holland.....	952	---	11.9	15.9	3.3	14.6	13.3	5	11.6	112.6
Flynn.....	1311	---	---	---	---	---	11.5	1	11.5	104.5

MOCCASIN, MONT.

The barley yields at the Judith Basin (Moccasin, Mont.) substation were obtained in cooperation with the Montana Agricultural Experiment Station. Barley was harvested in 13 of the years from 1908 to 1921, inclusive. The crop of 1912 was destroyed by hail. Annual and average yields for the entire period are shown in Table 58.

TABLE 58.—*Annual acre yields of varieties of barley grown at the Judith Basin substation (Moccasin, Mont.) in one or more of the 14 years from 1908 to 1921, inclusive*

[Data obtained in cooperation with the Montana Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)														Years grown	Aver- age yield (bus.)	Per- centage of weight- ed mean
		1908	1909	1910	1911	1913	1914	1915	1916	1917	1918	1919	1920	1921				
Mansury	617	20.0	45.7	13.0	38.3	48.7	38.0	70.8	40.0	11.2	45.0	3.5			11	34.0	104.0	
Thorpe	921	10.0	53.3	11.1	28.3	52.6	39.2	66.7	36.0	9.0					9	34.0	97.7	
Evans	892	5.5	32.6	10.3	19.3										4	16.9	65.3	
Black Hull-less	596	10.0	43.8	13.6	25.3										4	23.2	89.6	
Himalaya	620	5.5	33.3	12.3	25.0	44.3	30.3	48.8	18.7	10.3	33.9	4.8	29.6	43.4	13	26.3	79.0	
Nepal	595	20.0	42.4	16.4	30.0	37.0	25.7	41.8	22.1	8.1	30.0	2.3	21.9	39.6	13	25.9	77.8	
Bohemian	27	48.5	11.2												2	29.9	127.2	
Manchuria	244	45.2	9.8	40.8	39.2	36.0	52.2	53.5	9.6	43.3	2.5	15.6	39.6		12	30.6	89.7	
Hannchen	531	14.4	45.8	54.3	46.0	78.8	46.0	13.3	33.3	5.2	29.2	52.4			11	39.9	118.7	
White Smyrna	195	12.3	52.0	65.7	50.0	80.3	46.0	16.0	42.8	5.0	29.2	39.2			11	39.9	118.7	
Beldi Giant	2777	8.0	43.7	53.7	42.2	73.6	27.9	13.1	42.3	6.5					9	34.6	104.5	
Coast	690	15.0	43.7	55.0	43.1	83.1	52.0	14.6	39.0	8.3	27.5	44.5			11	36.9	109.8	
Gatari	575	8.2	34.4												2	21.3	89.1	
Club Mariout	261	16.2	47.0	47.9	51.3	72.0	32.8	15.2	35.1	6.9	24.4	45.8			11	35.9	106.8	
Canadian Thorpe	740	50.0	36.0	65.1											3	50.4	98.1	
Svanhals	187	48.9	43.5	68.1	39.6	11.9	47.3	3.1	29.2	50.2					9	38.0	106.4	
Clyde	909	37.5	22.0												2	29.7	69.4	
Franconian	680	31.2	80.0	40.0	11.9	50.0	4.4	27.3	48.5						8	36.7	107.3	
Heil Hanna 2	678	29.5	78.2	37.5	9.8										4	38.7	102.4	
White Smyrna	910	33.7	20.2	46.1	6.9	27.5	48.7								6	30.5	110.9	
Meloy	1176	33.7	15.8	44.8	5.0	34.0	44.8								6	29.7	108.0	
July	1563													30.2	1	30.2	67.9	
Horn	926													52.1	1	52.1	117.1	

TABLE 58.—*Annual acre yields of varieties of barley grown at the Judith Basin substation (Moccasin, Mont.) in one or more of the 14 years from 1903 to 1921, inclusive—Continued.*

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

Variety	C.I. No.	Data shown	Varieties and percentages									
			Mansury	Svanhals	Franconian	Hannchen	Manchuria	Coast	Club Mari- out	White Smyr- na	Meloy	Nepal
Mansury-----	617	Years comparable-----	-----	7	6	9	10	9	9	4	4	11
		Percentage yield-----	-----	98.0	95.9	86.4	113.5	92.4	95.1	93.3	100.4	135.7
Svanhals-----	187	Years comparable-----	7	-----	8	9	9	9	9	6	6	9
		Percentage yield-----	102.0	-----	99.9	90.3	125.9	98.5	103.1	99.0	101.8	149.6
Franconian-----	680	Years comparable-----	6	8	-----	8	8	8	8	6	6	8
		Percentage yield-----	104.3	100.1	-----	90.5	126.3	100.4	103.5	99.5	102.2	153.2
Hannchen-----	531	Years comparable-----	9	9	8	-----	11	11	11	6	6	11
		Percentage yield-----	115.8	110.7	110.5	-----	136.2	108.1	111.2	108.9	112.0	159.6
Manchuria-----	244	Years comparable-----	10	9	8	11	-----	11	11	6	6	12
		Percentage yield-----	88.1	79.4	79.2	73.4	-----	79.4	81.6	78.7	80.9	115.8
Coast-----	690	Years comparable-----	9	9	8	11	11	-----	11	6	6	11
		Percentage yield-----	108.2	101.6	99.6	92.5	126.0	-----	102.8	90.6	93.1	147.6
Club Mariout-----	261	Years comparable-----	9	9	8	11	11	11	-----	6	6	11
		Percentage yield-----	105.2	97.0	96.7	89.9	122.5	97.2	-----	87.5	89.9	143.5
White Smyrna-----	910	Years comparable-----	4	6	6	6	6	6	6	-----	6	6
		Percentage yield-----	107.2	101.0	100.5	91.8	127.1	110.4	114.3	-----	102.8	147.7
Meloy-----	1176	Years comparable-----	4	6	6	6	6	6	6	6	-----	6
		Percentage yield-----	99.6	98.2	98.7	89.3	123.6	107.4	111.2	97.3	-----	143.6
Nepal-----	595	Years comparable-----	11	9	8	11	12	11	11	6	6	-----
		Percentage yield-----	73.7	66.9	65.3	62.7	86.4	67.7	69.7	67.7	69.6	-----

Ten well-known varieties representative of the different types and groups were chosen for the purposes of the varietal comparison. The highest yields were obtained from Hannchen (C. I. No. 531) and White Smyrna (C. I. No. 910), while Franconian (C. I. No. 680), another 2-rowed barley representative of the Hanna group, ranked third. Coast (C. I. No. 690) and Svanhals (C. I. No. 187) are of almost equal importance. While Meloy (C. I. No. 1176) does not show up well in this comparison, it produced an average yield of 108 per cent of the weighted mean for the six years in which it was grown. The results from this hooded variety have been quite encouraging at a number of the stations in this same general region. Manchuria (C. I. No. 244) exceeded only Nepal (C. I. No. 595), the poorest of the varieties compared. Both of the North African barleys, Coast (C. I. No. 690) and Club Mariout (C. I. No. 261), yielded well during the 11 years they were grown but were inferior to the 2-rowed varieties, Hannchen and White Smyrna.

In 1921 varieties named July (C. I. No. 1563) and Horn (C. I. No. 926) were introduced. July, a 6-rowed sort, produced the lowest yield, but Horn, a Chevalier barley, was exceeded only by Hannchen in that one year. At Havre, Mont., the Chevalier and Hanna barleys were superior to Hannchen and White Smyrna for the years in which they were comparable.

It is evident, however, that to the south of Havre Hannchen and White Smyrna are the most widely adapted 2-rowed varieties. This is due in part to the fact that they mature somewhat earlier than the barleys of the Chevalier and Hanna types.

HUNTLEY, MONT.

The barley data from Huntley, Mont., as presented in Table 59, were obtained through the courtesy of the Office of Western Irrigation Agriculture. Only one variety, Trebi (C. I. No. 936), was grown for the five years from 1917 to 1921, inclusive. The other varieties tested were grown only in 1918. Trebi, with a yield of 70.5 bushels, ranked highest of the five tested that year. The other varieties were all 2-rowed sorts. White Smyrna (C. I. No. 195) was the best of these, with Chevalier II (C. I. No. 200) almost its equal. Hannchen (C. I. No. 531) and Svanhals (C. I. No. 187) were fourth and fifth, respectively, but were quite low in yield as compared with Trebi. Although Huntley is in a district favorable to the production of 2-rowed barleys, Trebi seems to be a superior sort there under irrigation, and in drier, hotter years it would doubtless yield much more than the 2-rowed varieties.

TABLE 59.—*Annual acre yields of varieties of barley grown at the Huntley (Mont.) Experiment Farm in part or all of the five years from 1917 to 1921, inclusive*

[Data obtained through the courtesy of the Office of Western Irrigation Agriculture]

Variety	C. I. No.	Acre yields (bushels)					Years grown	Average yield (bus.)	Percentage of weighted mean
		1917	1918	1919	1920	1921			
Trebi.....	936	49.7	70.5	70.7	56.5	47.0	5	58.9	101.4
White Smyrna.....	195	-----	62.0	-----	-----	-----	1	62.0	103.7
Chevalier II.....	200	-----	61.5	-----	-----	-----	1	61.5	102.8
Hannchen.....	531	-----	53.9	-----	-----	-----	1	53.9	90.1
Svanhals.....	187	-----	50.9	-----	-----	-----	1	50.9	85.1

BOZEMAN, MONT.

The data on yields of barley varieties grown at Bozeman, Mont., were received through the courtesy of the Montana Agricultural Experiment Station. From 1905 to 1912, inclusive, barley was grown in fortieth-acre plats. From 1913 to 1921, inclusive, it was grown only in replicated rows. It is thought that the yields for the period 1905 to 1912, inclusive, are best suited for the purposes of this bulletin. These yields are shown in Table 60.

TABLE 60.—*Annual acre yields of varieties of barley grown at the Montana Agricultural Experiment Station (at Bozeman) in some or all of the eight years from 1905 to 1912, inclusive*

[Data obtained through the courtesy of the Montana Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)								Years grown	Average yield (bus.)	Percentage of weighted mean
		1905	1906	1907	1908	1909	1910	1911	1912			
New Zealand.....	2656	90.8	86.7	64.2	75.0	78.3	39.4	85.2	64.2	8	73.0	117.9
Kitzing.....	189	101.7	76.7	58.3	57.5	74.2	47.9	88.3	76.7	8	72.7	117.4
Scotch Chevalier.....	2767	82.9	70.8	69.6	66.7	82.5	47.5	92.5	68.3	8	72.6	117.3
Chevalier.....	2768	64.2	48.3	72.9	54.2	80.0	50.0	100.0	79.2	8	68.6	110.8
Princess.....	529	60.0	60.0	72.2	48.1	60.0	34.8	93.3	71.3	8	62.5	101.0
Hannchen.....	531	45.8	68.3	50.6	53.4	76.3	56.6	83.7	86.4	8	65.1	105.2
Beldi Dwarf.....	190	81.0	58.3	58.8	56.7	72.5	35.4	93.3	67.5	8	65.4	105.7
California Prolific.....	2649	91.8	66.7	49.6	60.8	85.0	31.7	75.8	56.7	8	64.8	104.7
Himalaya.....	620	86.7	62.5	52.5	50.0	48.7	42.7	83.3	70.7	8	63.4	102.4
Manchuria.....	244	43.9	56.3	56.1	62.5	68.7	34.8	77.9	81.3	8	60.2	97.3
Frankish.....	207	62.5	62.5	66.7	63.7	64.2	31.6	81.7	50.8	8	60.5	97.7
Odesa.....	182	66.3	60.8	50.4	51.7	58.3	34.2	69.2	85.0	8	59.5	96.1
Berkley.....	2769	55.8	58.3	60.0	38.7	67.5	47.5	86.7	85.0	8	58.9	95.2
Nepal.....	630	70.4	46.7	50.0	52.1	77.1	37.3	68.7	62.0	8	58.0	93.7
Nepbant.....	2770	73.3	54.2	54.6	45.0	45.0	39.3	64.7	47.3	8	52.9	85.5
Evans.....	892	45.0	46.7	59.1	58.3	-----	35.3	67.3	62.0	7	53.4	87.1
Nepal.....	595	58.3	55.8	57.9	28.8	56.7	33.1	67.6	55.7	8	51.7	83.5
Black Hull-less.....	596	39.6	38.3	58.8	62.1	-----	38.7	64.7	61.3	7	51.9	84.7
Bavarian.....	208	63.3	57.5	60.0	63.7	-----	-----	-----	-----	4	61.1	105.9
Canadian Thorpe.....	740	59.2	37.5	-----	-----	-----	-----	-----	-----	2	48.3	80.0
Kulu.....	2694	54.2	47.5	-----	-----	-----	-----	-----	-----	2	50.9	84.3
Manchuria.....	2774	43.3	34.2	49.6	45.9	-----	-----	-----	-----	4	43.3	75.0
O. A. C. 21.....	1470	41.7	44.2	54.6	50.0	-----	-----	-----	-----	4	47.6	82.5
Chevalier II.....	200	-----	-----	62.1	70.8	89.2	58.3	84.2	77.5	6	73.7	118.3
Svanhals.....	187	-----	-----	47.9	67.5	76.2	69.6	78.3	77.5	6	69.5	111.6
Gutekorn.....	606	-----	-----	44.6	55.8	65.8	49.2	80.8	76.7	6	62.1	99.7
Sextradigt.....	607	-----	-----	45.0	55.0	58.3	50.8	80.8	85.0	6	62.5	100.3
Primus.....	532	-----	-----	40.0	48.5	71.4	67.3	86.6	75.8	6	64.9	104.2
Spear.....	2776	-----	-----	59.6	55.0	84.2	64.2	103.7	91.7	6	76.4	122.6
Hanna.....	2773	-----	-----	-----	35.4	55.0	62.5	86.7	67.5	5	61.4	97.3
Manhattan Malting.....	157	-----	-----	-----	51.1	51.7	64.2	67.5	69.2	5	60.7	96.2
Garton 986.....	645	-----	-----	-----	46.7	76.7	47.5	95.0	71.7	5	67.5	107.0
White Smyrna.....	658	-----	-----	-----	-----	35.0	104.2	85.0	-----	3	74.7	118.8
Shansi.....	2775	-----	-----	-----	-----	58.3	40.0	58.3	62.0	4	54.7	84.3
Chevalier.....	2771	-----	-----	-----	-----	-----	19.2	95.4	69.2	3	61.3	95.2
Chile Chevalier.....	659	-----	-----	-----	-----	-----	18.8	82.5	60.8	3	54.0	83.9
Chile Common.....	663	-----	-----	-----	-----	-----	27.1	99.2	83.3	3	69.9	108.5
Chile Brewing.....	2772	-----	-----	-----	-----	-----	20.4	95.8	76.7	3	64.3	99.8
Tatien.....	665	-----	-----	-----	-----	-----	19.7	62.0	54.0	3	45.2	70.2
Wyoming Hull-less.....	629	-----	-----	-----	-----	-----	27.7	67.3	58.7	3	51.2	79.5
Kashgar Hull-less.....	528	-----	-----	-----	-----	-----	35.0	56.7	56.7	3	49.5	76.9
Poda.....	652	-----	-----	-----	-----	-----	2.7	96.7	89.8	3	63.1	98.0
Peru.....	653	-----	-----	-----	-----	-----	18.3	95.8	98.3	3	70.8	109.9
Silver King.....	1276	-----	-----	-----	-----	-----	26.3	77.5	75.0	3	59.6	92.5

TABLE 60.—*Annual acre yields of varieties of barley grown at the Montana Agricultural Experiment Station (at Bozeman) in some or all of the eight years from 1905 to 1912, inclusive—Continued*

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 12 REPRESENTATIVE VARIETIES

[Under "Data shown" P=Percentage yield and Y=Years comparable]

Variety	C. I. No.	Data shown	Varieties and percentages										
			New Zea- land	California Prolific	Svanhals	Kitzing	Hannchen	Manchuria	Odessa	Peru	White Smyr- na	Beldi Dwarf	Nepal
New Zealand.....	2656	Y----- P-----	8 ----- 112.7	6 ----- 97.4	8 ----- 100.4	8 ----- 112.0	8 ----- 121.2	8 ----- 122.7	3 ----- 88.9	3 ----- 90.5	8 ----- 111.5	8 ----- 125.7	8 ----- 115.1
California Prolific.....	2649	Y----- P-----	8 ----- 88.7	6 ----- 86.2	8 ----- 89.1	8 ----- 99.4	8 ----- 107.6	8 ----- 108.9	3 ----- 77.3	3 ----- 85.9	8 ----- 99.0	8 ----- 111.6	8 ----- 102.2
Svanhals.....	187	Y----- P-----	6 ----- 102.6	6 ----- 116.0	6 ----- 103.5	6 ----- 109.4	6 ----- 119.6	6 ----- 106.1	3 ----- 100.0	3 ----- 108.5	6 ----- 120.1	6 ----- 116.5	6 ----- 116.5
Kitzing.....	189	Y----- P-----	8 ----- 99.6	6 ----- 112.2	6 ----- 96.6	8 ----- 111.6	8 ----- 120.7	8 ----- 122.1	3 ----- 100.2	3 ----- 93.8	8 ----- 111.0	8 ----- 125.2	8 ----- 114.6
Hannchen.....	531	Y----- P-----	8 ----- 89.3	6 ----- 100.6	6 ----- 97.6	8 ----- 89.6	8 ----- 108.2	8 ----- 109.5	3 ----- 106.7	3 ----- 96.6	8 ----- 99.5	8 ----- 112.2	8 ----- 102.8
Manchuria.....	244	Y----- P-----	8 ----- 82.5	6 ----- 92.9	6 ----- 91.4	8 ----- 82.8	8 ----- 92.4	8 ----- 101.2	3 ----- 91.3	3 ----- 80.9	8 ----- 92.0	8 ----- 103.7	8 ----- 95.0
Odessa.....	182	Y----- P-----	8 ----- 81.5	6 ----- 91.9	6 ----- 83.6	8 ----- 81.9	8 ----- 91.3	8 ----- 98.8	3 ----- 88.7	3 ----- 72.1	8 ----- 90.9	8 ----- 102.5	8 ----- 93.8
Peru.....	653	Y----- P-----	3 ----- 112.5	3 ----- 129.4	3 ----- 94.2	3 ----- 99.8	3 ----- 93.7	3 ----- 109.5	3 ----- 112.7	2 ----- 60.3	3 ----- 108.3	3 ----- 126.4	3 ----- 108.0
White Smyrna.....	658	Y----- P-----	3 ----- 110.5	3 ----- 116.5	3 ----- 100.0	3 ----- 106.0	3 ----- 103.5	3 ----- 123.6	3 ----- 138.7	2 ----- 165.8	3 ----- 111.4	3 ----- 122.4	3 ----- 128.3
Beldi Dwarf.....	190	Y----- P-----	8 ----- 89.7	6 ----- 101.0	6 ----- 92.1	8 ----- 90.1	8 ----- 100.5	8 ----- 108.7	3 ----- 110.0	3 ----- 92.4	3 ----- 89.7	8 ----- 112.8	8 ----- 103.2
Nepal.....	630	Y----- P-----	8 ----- 79.5	6 ----- 89.6	6 ----- 83.3	8 ----- 79.9	8 ----- 89.1	8 ----- 96.4	3 ----- 97.6	3 ----- 79.1	8 ----- 81.7	8 ----- 88.7	8 ----- 91.6
Himalaya.....	620	Y----- P-----	8 ----- 86.9	6 ----- 97.9	6 ----- 85.8	8 ----- 87.2	8 ----- 97.3	8 ----- 105.3	3 ----- 106.6	3 ----- 92.6	8 ----- 77.9	8 ----- 96.9	8 ----- 109.2

A comparison of six varietal groups was made, of which Coast was probably the best. As only three varieties were included and as this group is comparable with other groups for only three years, its relative position is not positive. The Chevalier group is second, with Hanna and Thorpe following, successively. These three consist only of 2-rowed sorts. The Manchuria barleys are entirely out of their ecological district here. The hull-less group was the poorest, as is the case at most stations.

In the summarized digest of Table 60, 12 important and representative varieties are compared. White Smyrna (C. I. No. 658) ranked first. It is to be regretted that it can be compared with the other varieties for only three years, as it produced an exceptionally high yield in 1910 but relatively low yields in 1909 and 1911. Svanhals (C. I. No. 187) exceeded all the others compared in this table except White Smyrna and was its equal in the three years it was grown. New Zealand (C. I. No. 2656), a representative of the Chevalier group, was third in point of yield, with Kitzing (C. I. No. 189), of the Hanna group, following. These barleys are also 2-rowed sorts.

Peru (C. I. No. 653), a variety of the Coast group, is fifth. Beldi Dwarf (C. I. No. 190), a 6-rowed North African barley, gave an average yield of 106 per cent of the weighted mean for the eight years in which it was tested, but it was exceeded by Peru and four 2-rowed varieties. As demonstrated at the other stations in Montana this section is undoubtedly in the region where the 2-rowed barleys are well adapted. At the present time a number of varieties are being tested which seem quite promising. Among these are the 6-rowed varieties Trebi (C. I. No. 936), Flynn (C. I. No. 1311), and Meloy (C. I. No. 1176).

SHERIDAN, WYO.

The barley yields at the Sheridan Experiment Farm were obtained in cooperation with the Office of Dry-Land Agriculture Investigations. Table 61 shows the yields for the varieties grown from 1917 to 1921, inclusive. Of these, White Smyrna (C. I. No. 195) was the best, Trebi (C. I. No. 936) second, and Svanhals (C. I. No. 187) third. Coast (C. I. No. 690) was almost the equal of Svanhals, with Hann-

chen (C. I. No. 531) following. Club Mariout (C. I. No. 261) and Manchuria (C. I. No. 244) produced only fair yields. In 1921 Flynn (C. I. No. 1311) and Horn (C. I. No. 926) were introduced into the experiment. As all the yields for this year were very low, owing to scant rainfall, a fair comparison can not be made. It is interesting to note that Trebi, a 6-rowed barley that has given high yields under irrigation, is second here to White Smyrna, a 2-rowed sort, which is one of the best varieties for dry-land conditions. As far as tested, it appears that, in general, the 2-rowed barleys are the most successful types in northern Wyoming.

TABLE 61.—*Annual acre yields of varieties of barley grown at the Sheridan (Wyo.) Experiment Farm in part or all of the five years from 1917 to 1921, inclusive*

[Data obtained in cooperation with the Office of Dry-Land Agriculture Investigations]

Variety	C. I. No.	Acre yields (bushels)					Years grown	Average yield (bus.)	Percentage of weighted mean
		1917	1918	1919	1920	1921			
Svanhals	187	11.7	35.1	0	21.3	4.6	5	14.5	109.8
White Smyrna	195	24.6	26.5	.8	29.4	3.9	5	17.0	128.8
Hannchen	531	13.3	30.9	.1	21.1	2.7	5	13.6	103.0
Club Mariout	261	20.8	17.9	.4	19.2	2.5	5	12.2	92.4
Manchuria	244	12.1	28.5	.3	22.9	1.6	5	13.1	99.2
Coast	690	18.3	26.7	.5	24.0	2.1	5	14.3	108.3
Trebi	936	18.3	33.1	.3	23.9	2.5	5	15.6	118.2
Nepal	595	10.3	13.0	0	10.4		4	8.4	52.2
Flynn	1311					3.6	1	3.6	120.0
Horn	926					3.3	1	3.3	110.0

ARCHER, WYO.

Varietal tests of barley have been conducted at the Cheyenne Experiment Farm, Archer, Wyo., since 1913. The data presented in Table 62 were obtained in cooperation with the Wyoming State Board of Farm Commissioners.

TABLE 62.—*Annual acre yields of varieties of barley grown at the Cheyenne (Archer, Wyo.) Experiment Farm in part or all of the nine years from 1913 to 1921, inclusive*

[Data obtained in cooperation with the Wyoming State Board of Farm Commissioners]

Variety	C. I. No.	Acre yields (bushels)									Years grown	Average yield (bus.)	Percentage of weighted mean
		1913	1914	1915	1916	1917	1918	1919	1920	1921			
Hannchen	531	9.6	16.0	34.1	7.9	28.9	56.3	3.9	23.3	24.6	9	22.7	111.8
Primus	532	8.9	8.3	33.9	4.3	22.9					5	15.7	98.1
Hanna	24	8.8	10.8	32.0							3	17.2	102.4
Manchuria	241	6.6	11.8	33.5	5.6	25.9	47.1	2.7	10.0	16.7	9	17.8	87.7
Nepal	595	6.4	10.0	29.7	2.4	16.3					5	13.0	81.3
Manchuria	244	7.2	9.6	30.2							3	15.7	93.5
Baku	709	6.5	10.4	29.1							3	15.3	91.1
Blackhull	878	3.5	14.8	22.7	7.8	12.9	41.4	1.7	12.4	20.4	9	15.3	75.4
Horsford	877	6.3	14.5	26.6	6.2	19.7	41.7	1.9	11.5		8	16.1	80.1
Gatami	203	7.5									1	7.5	102.7
Gatami	575	4.4	15.5	23.7	7.0	21.6					5	14.4	90.0
Coast	690	6.6	14.4	41.2	8.0	28.4	59.4	3.8	24.3	21.8	9	23.1	113.8
White Smyrna	658	10.0	16.3	38.3	8.3	27.2	64.3	3.8	20.6	24.4	9	23.7	116.7
Black Hull-less	1106	12.3	9.4	34.7	2.4	23.0	55.9	6.0	17.1	18.4	9	19.8	97.5
Yerli	660	5.7	9.6	30.6	4.4						4	12.6	87.5
Svanhals	187	5.9	11.9	31.6	7.1	27.2	62.4	2.8	18.4	20.2	9	20.8	102.5
Himalaya	620				7.3		62.3	3.4	16.7	22.3	6	21.9	98.2
Horn	926					30.2	68.1	3.6	25.0	23.7	5	30.1	119.4
Italian	914						66.6	2.9	22.1	21.4	4	28.3	110.5
Sandrel	937						73.1	4.2	23.3	23.4	4	31.0	121.1
Odessa	927						52.4	4.8	21.5	26.6	4	26.3	102.7
Trebi	936						57.3	5.0	26.9	24.3	4	28.4	110.9
Meloy	1176								23.2	22.0	2	22.6	108.1
Flynn	1311									22.0	1	22.0	99.5

TABLE 62.—*Annual acre yields of varieties of barley grown at the Cheyenne (Archer, Wyo.) Experiment Farm in part or all of the nine years from 1913 to 1921, inclusive—Continued*SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 12 REPRESENTATIVE VARIETIES
[Under "Data shown" P=Percentage yield and Y=Years comparable]

Variety	C. I. No.	Data shown	Varieties and percentages										
			Horn	Svanhals	Italian	Hannchen	Manchuria	Odessa	Coast	Sandrel	Trebi	White Smyr- na	Black Hull- less
Horn.....	926	(Y)----- (P)-----	5 115.0	4 106.5	5 109.9	5 147.1	4 114.3	5 109.4	4 97.1	4 106.1	5 107.3	5 125.1	4 169.7
Svanhals.....	187	(Y)----- (P)-----	5 87.0	4 91.9	9 91.6	9 117.3	9 98.6	9 90.2	4 83.7	4 91.5	9 87.9	4 104.6	3 130.3
Italian.....	914	(Y)----- (P)-----	4 93.9	4 108.9	4 104.5	4 147.7	4 107.3	4 103.4	4 91.1	4 99.6	9 99.9	4 116.0	3 166.2
Hannchen.....	531	(Y)----- (P)-----	5 91.0	9 109.1	9 95.7	9 128.0	9 102.7	9 98.4	4 87.2	4 95.2	9 96.0	9 114.2	8 140.2
Manchuria.....	241	(Y)----- (P)-----	5 68.0	9 85.3	9 67.7	9 78.2	9 72.6	9 76.9	4 61.7	4 67.4	9 75.0	9 89.2	8 111.5
Odessa.....	927	(Y)----- (P)-----	5 87.5	4 101.4	4 93.2	4 97.4	4 137.6	4 96.3	4 84.9	4 92.8	9 93.1	4 108.1	8 142.8
Coast.....	690	(Y)----- (P)-----	5 91.4	9 110.9	9 93.7	9 101.6	9 130.0	4 103.8	4 88.1	4 96.3	9 97.5	4 116.0	9 144.9
Sandrel.....	937	(Y)----- (P)-----	4 103.0	4 119.5	4 109.7	4 114.7	4 162.1	4 117.8	4 113.4	4 109.3	4 109.6	4 127.3	3 182.6
Trebi.....	936	(Y)----- (P)-----	4 94.3	4 109.3	4 100.4	4 105.0	4 148.4	4 107.8	4 103.8	4 91.5	4 100.4	4 116.5	3 161.9
White Smyrna.....	658	(Y)----- (P)-----	5 93.2	9 113.7	9 100.1	9 104.2	9 133.3	9 107.4	4 102.5	4 91.2	4 99.6	4 119.0	8 147.0
Black Hull-less.....	1106	(Y)----- (P)-----	5 79.9	9 95.6	9 86.2	9 87.6	9 112.1	9 92.5	4 86.2	4 78.5	4 85.8	4 84.1	3 125.2
Horsford.....	877	(Y)----- (P)-----	4 58.9	8 76.7	3 60.2	3 71.3	3 89.7	3 70.0	3 69.0	3 54.8	3 61.8	3 68.0	3 79.9

Among the varieties compared in Table 62 there are four, Sandrel (C. I. No. 937), Trebi (C. I. No. 936), Italian (C. I. No. 914), and Odessa (C. I. No. 927), which were grown for only four years. Although Sandrel is the highest yielding variety for these years, its rank, as well as that of the other three grown for this period, is not so definite as it would be if the results for a greater number of years had been included. Following Sandrel is Horn (C. I. No. 926), a 2-rowed sort which is promising at several of the Montana stations. Trebi (C. I. No. 936) ranks third. White Smyrna (C. I. No. 658), which is next in order, was grown for the entire period of nine years, its average yield being 116.7 per cent of the weighted mean. Another high-yielding variety was Italian (C. I. No. 914), a barley of the Hanna type. Although Coast (C. I. No. 690) and Hannchen (C. I. No. 531) do not make so good a showing in this table, they produced relatively good yields during the nine years of the experiment. As usual, Horsford (C. I. No. 877) and Black Hull-less (C. I. No. 1106) were poor-yielding sorts. Manchuria (C. I. No. 241) and Odessa (C. I. No. 927) are evidently unsuited to growing conditions here. The table of yields shows that Meloy (C. I. No. 1176) gave fair returns for two years, 1920 and 1921. Flynn (C. I. No. 1311), which was grown in 1921, was the equal of Meloy.

White Smyrna in general should give better results than other 2-rowed sorts of the Coast types. Because of its short straw, however, it is probably not so desirable for field culture as some of the other varieties tested. It seems quite evident that Archer is near the line where the Coast is equal to the 2-rowed varieties in yield.

FORT COLLINS, COLO.

The yields of barley varieties at the Colorado Agricultural Experiment Station, Fort Collins, for the period from 1918 to 1921, inclusive, are shown in Table 63. Records of these yields were obtained through the courtesy of the Colorado station. Twenty-two varieties were grown for the entire four years. Of these Coast (C. I. No. 2785) produced the highest average yield, 76.9 bushels. Two 2-rowed barleys belonging to the Hanna group, Hanna (C. I. No. 2784) and Gold

(C. I. No. 1145), were second and third. Following Gold are Coast (C. I. No. 2790) and Colless (C. I. No. 2792), the latter a hybrid variety which was produced at this station. Coast (C. I. No. 2789) is almost the equal of Colless. As shown in Table 63 the varieties grown during the entire four years which had an average of more than 100 per cent of the weighted mean were either Coast or Hanna types with the exception of Colless and Charlottetown 80 (C. I. No. 2732). Coast (C. I. No. 2791) was introduced in 1919, but was found inferior to several of the other pedigreed Hanna and Coast barleys. Golden Drop (C. I. No. 2135), a variety of the Chevalier group, was grown in 1920 and 1921, but was not a superior sort in either of those years.

Twelve varieties were introduced in 1921. With the exception of Bark (C. I. No. 2793) all of these produced relatively high yields. Manchuria (C. I. No. 2783) and O. A. C. 21 (C. I. No. 1470), both pedigreed strains of Manchuria barley, were the highest yielders that year. This is rather surprising, as the Manchuria types would appear to be entirely out of their ecological region in Colorado.

Considering the whole period of the experiment, however, there is no doubt that the barleys of the Coast and Hanna types were the highest yielding sorts. Fort Collins, like Archer, Wyo., is apparently near the line separating the district where the 2-rowed varieties grow best from that in which Coast is a superior type. In the 2-rowed district east and north of Fort Collins, Hannchen and White Smyrna were usually the best of the 2-rowed barleys. At this station Hannchen yielded well in 1921, the only year it was grown, but White Smyrna was not tested.

TABLE 63.—*Annual acre yields of varieties of barley grown at the Colorado Agricultural Experiment Station (Fort Collins) in part or all of the four years from 1918 to 1921, inclusive*

[Data obtained through the courtesy of the Colorado Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1918	1919	1920	1921			
Hanna.....	2784	73.1	59.0	76.5	83.1	4	72.9	115.3
Coast.....	2785	78.8	67.5	81.8	79.7	4	76.9	121.7
Gold.....	1145	65.8	59.6	81.8	76.9	4	71.0	112.3
Hanna.....	2786	61.6	58.3	82.0	76.0	4	69.5	110.0
Do.....	2787	68.7	50.5	74.6	75.6	4	67.3	106.5
Coast.....	2789	70.0	58.0	78.9	73.6	4	70.1	110.9
Colless.....	2792	70.4	53.1	84.8	73.1	4	70.3	111.2
Hanna.....	2788	66.0	53.8	75.2	72.4	4	66.9	105.9
Coast.....	2790	69.8	62.9	78.0	72.4	4	70.8	112.0
Charlottetown 80.....	2732	67.8	50.0	69.6	70.8	4	64.5	102.1
Black Hull-less.....	596	33.0	49.7	62.7	65.9	4	52.8	83.5
Silver King.....	890	53.7	33.7	60.0	64.5	4	53.0	83.9
Coast.....	690	69.2	64.2	77.9	63.8	4	68.8	108.9
Telli.....	194	53.1	49.5	71.3	63.6	4	59.4	94.0
Garton 986.....	645	59.2	51.5	73.9	60.5	4	61.3	97.0
Chevalier II.....	200	59.7	55.9	76.2	59.2	4	62.7	99.2
Horsford.....	507	32.2	28.2	50.5	56.3	4	41.8	66.1
Wing Pedigree.....	1177	34.8	27.3	57.5	55.6	4	43.8	69.3
Horsford.....	1271	35.9	33.7	55.2	50.3	4	43.8	69.3
Princess.....	529	61.7	45.7	66.2	54.9	4	57.1	90.3
Nepal.....	595	32.0	44.7	63.3	47.7	4	46.9	74.2
Blackhull.....	878	53.2	46.9	51.3	49.0	4	50.1	79.3
Coast.....	2791		60.0	78.1	65.5	3	67.9	105.1
Golden Drop.....	2135			73.3	68.5	2	70.9	101.3
O. A. C. 21.....	1470				88.6	1	88.6	127.5
Manchuria.....	2783				88.4	1	88.4	127.2
Moister.....	2799				83.8	1	83.8	120.6
Elfy.....	2800				81.3	1	81.3	117.0
Canadian Thorpe.....	740				80.1	1	80.1	115.3
Samofa.....	1211				80.1	1	80.1	115.3
Manchuria.....	2330				77.3	1	77.3	111.2
Minstardi.....	1556				76.9	1	76.9	110.6
Svansota.....	1907				73.0	1	73.0	105.0
Lion.....	923				69.8	1	69.8	100.4
Bark.....	2793				47.0	1	47.0	67.6
Hannchen.....	531				76.2	1	76.2	109.6

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages									
			Primus	Hanna	Hannehen	Manehuria	Coast	Smyrna	Baku	Blackhull	Orel	Horsford
Primus.....	532	{ Years comparable.....	5	5	5	5	5	(*)	2	5	3	5
		{ Percentage yield.....	66.4	61.1	75.7	62.3	62.3	---	71.0	81.3	50.5	101.6
Hanna.....	203	{ Years comparable.....	5	7	6	12	12	5	5	12	4	11
		{ Percentage yield.....	150.7	89.2	112.6	87.1	87.1	75.1	148.4	103.8	88.2	135.7
Hannehen.....	531	{ Years comparable.....	5	7	6	6	7	(*)	2	7	3	7 *
		{ Percentage yield.....	163.6	112.1	120.1	100.1	100.1	---	215.4	137.7	96.4	152.2
Manehuria.....	244	{ Years comparable.....	5	6	6	6	6	(*)	2	6	3	6
		{ Percentage yield.....	132.0	88.8	83.3	79.4	79.4	---	173.9	107.3	75.9	127.2
Coast.....	690	{ Years comparable.....	5	12	7	6	---	7	7	14	4	13
		{ Percentage yield.....	160.6	114.8	99.9	125.9	---	93.5	154.4	122.7	97.2	154.9
Smyrna.....	2642	{ Years comparable.....	(*)	5	(*)	(*)	7	---	5	7	1	6
		{ Percentage yield.....	---	133.2	---	---	107.0	---	146.3	115.4	117.3	170.5
Baku.....	709	{ Years comparable.....	2	5	2	2	7	5	---	7	2	7
		{ Percentage yield.....	140.8	67.4	46.4	57.5	64.8	68.3	---	74.7	38.6	91.6
Blackhull.....	878	{ Years comparable.....	5	12	7	6	14	7	7	---	4	13
		{ Percentage yield.....	123.0	96.4	72.6	93.2	81.5	86.6	135.9	---	83.9	120.5
Orel.....	2190	{ Years comparable.....	3	4	3	3	4	1	2	4	---	3
		{ Percentage yield.....	197.9	113.4	103.8	131.8	102.9	85.2	---	119.2	---	187.0
Horsford.....	877	{ Years comparable.....	5	11	7	6	13	6	7	13	3	---
		{ Percentage yield.....	98.4	73.7	65.7	78.6	64.6	58.6	106.2	83.0	53.5	---

AKRON, COLO.

The barley yields at the Akron field station (Table 64) were obtained in cooperation with the Office of Dry-Land Agriculture Investigations. Ten important varieties are compared in the summarized digest of the table. Smyrna (C. I. No. 2642) is the best of the varieties included in the comparison. It was not grown in the same years as Hannchen (C. I. No. 531), Primus (C. I. No. 532), or Manchuria (C. I. No. 244), so that a comparison with these varieties can not be made. Following Smyrna is Hannchen (C. I. No. 531). In 1915 this variety became confused with another. As a consequence, no comparison can be made after 1914. Orel (C. I. No. 2190) is third in point of yield. This strain is thought to be identical with Orel (C. I. No. 351), which was grown from 1917 to 1921, inclusive. However, Orel (C. I. No. 351) was surpassed by Coast (C. I. No. 690) as well as by the two Smyrna varieties in this later period. Orel is usually grown as a winter barley, and it is of interest that it is a high-yielding sort when sown in the spring at Akron. Coast (C. I. No. 690) was almost the equal of both Hannchen and Orel. For the 14 years of the experiment Coast produced an average yield of 113 per cent of the weighted mean. Hanna (C. I. No. 203) and Manchuria (C. I. No. 244) follow in the order named. Blackhull (C. I. No. 878) was superior to Primus (C. I. No. 532), Baku (C. I. No. 709), and Horsford (C. I. No. 877). (See Pl. III, fig. 2, A and B.) Grown for the full period of the experiment, 1908 to 1921, inclusive, its average yield was 92 per cent of the weighted mean. Blackhull has yielded better at Akron than at most of the other stations, and those who have been in charge of the testing think that the uniformity of its yields and other excellent qualities make it a desirable variety. As can be seen in Table 64, the winter barleys, Tennessee Winter (C. I. No. 257), Arlington Awnless (C. I. No. 702), and Nakano Wase (C. I. No. 754), which were grown during different periods, were not high-yielding sorts as compared with the spring varieties.

While the 2-rowed barleys are undoubtedly the highest yielding sorts at Akron, it is evident that Coast is also well adapted to this district. It appears that Akron, Archer, and Fort Collins are all near the border line between the 2-rowed and Coast districts. Where the yield is equal, farmers usually prefer 6-rowed varieties, as the straw of the 2-rowed ones is often weak and in White Smyrna is very short.

SANDPOINT, IDAHO

The yields of barley varieties at the Sandpoint substation for the years 1920 and 1921 were obtained through the courtesy of the Idaho Agricultural Experiment Station. These yields are found in Table 65. As the crop of 1921 was almost a failure, the yields for this year are of little value in determining the relative importance of the varieties grown. Han River (C. I. No. 206) produced the highest average yield for the two years of the experiment, while White Smyrna (C. I. No. 910) ranked second. Baker (C. I. No. 975), Trebi (C. I. No. 936), and Peruvian (C. I. No. 935) follow in the order named. Winter Club (C. I. No. 488) was grown only in 1921, but was exceeded by both White Smyrna and Peruvian.

TABLE 65.—*Annual acre yields of varieties of barley grown at the Sandpoint (Idaho) substation in 1920 and 1921*

[Data obtained through the courtesy of the Idaho Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)		Years grown	Average yield (bus.)	Percentage of weighted mean
		1920	1921			
Han River.....	206	24.5	3.2	2	13.9	149.4
Trebi.....	936	14.5	2.3	2	8.4	90.3
White Smyrna.....	910	17.2	4.0	2	10.6	114.0
Peruvian.....	935	9.1	3.7	2	6.4	68.8
Baker.....	975	17.2	3.0	2	10.1	108.6
Winter Club.....	488		3.2	1	3.2	100.0

MOSCOW, IDAHO

Table 66 shows the yields of varieties of barley grown from 1910 to 1921, inclusive, at Moscow, Idaho. Records of these yields were obtained through the courtesy of the Idaho Agricultural Experiment Station. The high yield of the winter varieties shown here is most striking. Moscow is situated in the Palouse district, which lies just to the west of the western Rocky Mountain Ranges. It has a mild winter climate, and in the better part of the Palouse district there is an average annual rainfall of 20 to 25 inches, with an average of only a few inches in the months of April to July, inclusive. The soil, which is unusually fertile, is a wind-formed deposit. These conditions are favorable to the growth of winter barleys, and it is likely that only the time needed for wheat seeding prevents their extensive culture.

Of the spring-sown varieties introduced into the experiment before 1919 only Winter Club (C. I. No. 488) was grown through 1921, the others being dropped in 1920. Of these varieties White Smyrna (C. I. No. 195) was perhaps the best. However, a selection of this variety (C. I. No. 910) which was grown in 1920 and 1921 did not prove to be as high yielding as the others grown in those two years. Winter Club (C. I. No. 488), when spring sown, was second, and Hannehen (C. I. No. 531) was third in point of yield. Following Hannehen are Chile Common (C. I. No. 663) and California Centgener (C. I. No. 3147), both North African barleys of the Coast group. Svanhals (C. I. No. 187) also gave a fair average yield, but was surpassed by the varieties mentioned above in the years for which it was comparable. Of the five spring barleys grown only in 1920 and 1921 Peruvian (C. I. No. 935) was the best. This is also of North African origin and belongs to the Coast group. Trebi (C. I. No. 936) and Baker (C. I. No. 975) were second and third in rank, respectively. Han River (C. I. No. 206) was better than either Winter Club or White Smyrna in the two years.

TABLE 66.—*Annual acre yields of varieties of barley grown at the Idaho Agricultural Experiment Station (at Moscow) in one or more of the 12 years from 1910 to 1921, inclusive*

[Data obtained through the courtesy of the Idaho Agricultural Experiment Station]

Variety	C.I. No.	Acre yields (bushels)												Years grown	Average yield (bus.)	Percentage of weighted mean
		1910	1911	1912	1913	1914	1915	1916	1917	1918	1919	1920	1921			
White Smyrna.....	195	38.0	47.9	82.0	48.5	52.5	84.0	70.3	32.8	14.5	44.3	-----	-----	10	51.5	126.5
Nepal.....	595	33.8	27.5	537.6	-----	-----	-----	33.3	14.8	6.3	22.4	-----	-----	7	25.1	65.7
Hannehen.....	531	43.3	34.6	53.3	60.2	42.7	-----	76.4	22.5	12.0	37.1	-----	-----	9	42.5	107.9
Winter Club.....	488	-----	-----	97.3	60.2	60.0	48.5	54.8	31.8	20.5	37.6	77.6	54.2	10	54.3	115.8
Oderbrucker.....	3146	-----	-----	13.2	48.1	49.7	56.2	51.2	214.8	37.1	-----	-----	-----	7	38.6	97.5
Svanhals.....	187	-----	-----	38.1	58.7	65.1	69.4	17.8	7.9	39.8	-----	-----	-----	7	42.4	107.1
White Moravian.....	1877	-----	-----	33.3	51.4	29.8	51.1	22.2	7.0	39.7	-----	-----	-----	7	33.5	84.6
California Centgener.....	3147	-----	-----	-----	-----	70.3	59.8	22.4	20.0	33.2	-----	-----	-----	5	41.1	107.0
Beldi Dwarf.....	190	-----	-----	-----	-----	43.1	70.1	23.9	4.7	42.7	-----	-----	-----	5	36.9	96.1
Arlington Awnless.....	702	-----	-----	-----	-----	9.9	26.6	22.2	6.0	35.5	-----	-----	-----	5	20.0	52.1
Chile Common.....	663	-----	-----	-----	-----	71.7	57.1	29.0	21.6	35.9	-----	-----	-----	5	43.1	112.2
Wing Beardless.....	2147	-----	-----	-----	-----	-----	27.1	13.6	5.5	21.7	-----	-----	-----	4	17.0	46.7
Abed.....	1389	-----	-----	-----	-----	-----	52.9	21.5	11.6	40.9	-----	-----	-----	4	31.7	87.1
Beardless.....	3145	-----	-----	-----	-----	-----	60.8	40.7	9.7	39.3	-----	-----	-----	4	37.6	103.3
White Smyrna.....	910	-----	-----	-----	-----	-----	-----	-----	-----	-----	76.9	47.2	-----	2	62.1	84.4
Han River.....	206	-----	-----	-----	-----	-----	-----	-----	-----	-----	77.5	56.5	-----	2	67.0	91.0
Trebi.....	936	-----	-----	-----	-----	-----	-----	-----	-----	-----	89.6	62.7	-----	2	76.1	103.4
Peruvian.....	935	-----	-----	-----	-----	-----	-----	-----	-----	-----	87.1	72.9	-----	2	80.0	108.7
Baker.....	975	-----	-----	-----	-----	-----	-----	-----	-----	-----	57.1	62.5	-----	2	74.8	101.6
Winter barley:																
Winter Club.....	488	-----	-----	-----	-----	-----	-----	-----	41.0	102.9	83.7	81.2	-----	4	77.2	161.8
Wisconsin Winter.....	519	-----	-----	-----	-----	-----	-----	-----	40.5	101.3	74.3	77.3	-----	4	73.3	153.7
Tennessee Winter.....	257	-----	-----	-----	-----	-----	-----	-----	26.7	87.9	60.8	75.0	-----	4	62.6	131.2
Michigan Winter.....	2036	-----	-----	-----	-----	-----	-----	-----	37.9	88.6	72.9	95.8	-----	4	73.8	154.7

The winter barleys mentioned were first seeded in the fall of 1917, being harvested in the summers of 1918 to 1921, inclusive. Winter Club (C. I. No. 488) produced the highest average yield, 77.2 bushels, in these years. This variety is a superior one here when either spring or fall sown. Michigan Winter (C. I. No. 2036) and Wisconsin Winter (C. I. No. 519) follow in the order named. Tennessee Winter (C. I. No. 257) was the lowest in rank of the winter sorts. Except in 1920 all of the winter barleys were superior to the spring varieties.

It is evident that the climatic conditions here are such that several distinct types of barley may be successfully grown. Of the spring varieties both the 2-rowed sorts and the barleys of the Coast group are outstanding in yield. Trebi and Baker, representative of still another group, also seem promising. There is no doubt, however, that the winter barleys are superior in yield so far as they have been tested at Moscow. Even in field culture, despite occasional winter-killing, which might be expected, the winter barleys may be more profitable than the spring varieties.

ABERDEEN, IDAHO

Irrigated land.—The Aberdeen substation is most advantageously situated for testing barleys under irrigation, particularly those of the types adapted to arid and semiarid conditions. It is located in the Snake River basin of southern Idaho at an altitude of 4,400 feet. The growing season is short, and the river makes available a water supply for irrigation purposes. Extensive nursery and plat tests of barley varieties have been made at the Aberdeen substation in recent years. Most of the varieties tested seem to produce maximum yields here.

TABLE 67.—*Annual acre yields of varieties of barley grown on irrigated land at the Aberdeen (Idaho) substation in part or all of the nine years from 1913 to 1921, inclusive*

[Data obtained in cooperation with the Idaho Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)										Years grown	Average yield (bus.)	Percent- age of weight- ed mean
		1913	1914	1915	1916	1917	1918	1919	1920	1921				
Silver King	905	26.0									1	26.0	46.2	
Svanhals	187	60.8	58.2	70.5							3	63.2	95.6	
Hanna	906	54.1	59.5	65.0							3	59.5	90.0	
Manchuria	244	43.3	63.0	73.9							3	60.1	90.9	
Club Mariout	261	63.8	39.0								2	51.4	83.2	
Princess	529	46.6									1	46.6	82.8	
Hannchen	531	61.6	73.3	80.6	89.5	84.1	68.7	20.8	55.4	86.2	9	68.9	102.7	
White Smyrna	195	69.5									1	69.5	123.4	
Han River	206	78.3	68.7	81.7	67.4	92.9	57.0	28.0	55.7	97.9	9	69.7	103.9	
Woodrow	986	71.5									1	71.5	127.0	
White Moravian	977	64.1	57.8	72.8	85.8						4	70.1	103.9	
Composite	1147	56.8	47.0	89.6							3	64.5	97.6	
Wisconsin Winter	519	63.3	77.9	78.4							3	73.2	110.7	
Coast	690	51.6	67.6	53.7							3	57.6	87.1	
Gatami	575	49.5									1	49.5	87.9	
Beldi Giant	2777	56.0	88.2	80.6	62.4	96.6	70.7	30.5	69.9	96.0	9	72.3	107.7	
Coast	2235	65.0	67.6								2	66.3	107.3	
Chevalier	278	50.4									1	50.4	89.5	
Union	2198	49.0									1	49.0	87.0	
Do	2199	48.0									1	48.0	85.3	
Nepal	230	40.3									1	40.3	71.6	
Bavarian	160	70.0	59.5								2	64.7	104.7	
Odessa	927		81.3	79.5	66.6						3	75.8	106.5	
White Smyrna	910		70.4	63.8	90.4	87.5	53.7	29.3	53.7	93.7	8	67.8	98.0	
Chevalier II	200		58.6	71.6							2	65.1	92.7	
Club Mariout	932		59.7								1	59.7	90.7	
Trebi	936		85.9	84.0	69.6	105.0	92.8	25.3	78.7	96.7	8	79.7	115.2	
Servian	915		76.7	81.7	66.2						3	74.9	105.3	
Eagle	913		68.7	76.1	54.1						3	66.3	93.2	
Sandrel	937		84.7	80.5	67.4	98.7	64.9	29.9	72.5	86.0	8	73.1	105.6	
Peruvian	935		74.4	84.0	74.9	87.9	54.5	30.4	67.5	94.2	8	71.0	102.6	
Featherston	911		69.8								1	69.8	106.1	
Luth	908		66.4	73.9							2	70.1	99.9	
Manchuria	1275		34.9								1	34.9	53.0	
Steigum	907		43.3								1	43.3	65.8	
Goldfoil	928		67.6	71.7	83.4						3	74.2	104.4	
Horn	926		67.0	78.4	83.3	80.0					4	77.2	105.2	
Australian White	925		66.4	70.5							2	68.5	97.7	
Italian	914		71.0	81.7	88.4	75.4					4	79.1	107.8	
Baker	975						43.3	30.1	63.7	85.2	4	55.6	91.9	
Rasput	996						34.9				1	34.9	60.8	
Petro	994						33.3				1	33.3	58.0	
Algerian	1179								57.5	70.8	2	64.1	83.6	

TABLE 67.—*Annual acre yields of varieties of barley grown on irrigated land at the Aberdeen (Idaho) substation in part or all of the nine years from 1913 to 1921, inclusive—Continued*

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 12 REPRESENTATIVE VARIETIES

[Under "Data shown" P=Percentage yield and Y=Years comparable]

Variety	C. I. No.	Data shown	Varieties and percentages										
			Horn	Svanhals	Italian	Hannchen	Manchuria	Odessa	Sandrel	Beldi Giant	White Smyrna	Trebi	Han River
Horn-----	926	{Y----- P-----	----- 113.0	2 97.5	4 94.3	4 106.2	2 100.6	3 93.2	4 94.2	4 98.9	4 89.6	4 99.4	4 93.0
Svanhals-----	187	{Y----- P-----	----- 88.5	2 84.3	3 87.9	3 105.2	2 80.0	2 77.9	3 84.3	2 95.9	2 75.8	3 82.9	3 86.3
Italian-----	914	{Y----- P-----	----- 102.5	2 118.6	4 96.6	2 111.5	2 106.0	4 95.5	4 96.6	4 101.4	4 91.9	4 101.9	2 97.7
Hannchen-----	531	{Y----- P-----	----- 106.1	3 113.7	4 103.5	----- 119.6	3 107.0	3 95.6	3 95.3	3 103.0	3 87.6	3 98.8	3 98.1
Manchuria-----	244	{Y----- P-----	----- 94.2	2 95.1	3 89.7	3 83.6	----- 85.1	2 82.9	2 80.2	2 102.0	2 80.6	3 78.8	2 82.1
Odessa-----	927	{Y----- P-----	----- 99.4	2 124.9	4 94.3	8 93.4	3 117.5	----- 97.8	3 98.4	3 101.2	3 94.9	3 104.4	2 102.9
Sandrel-----	937	{Y----- P-----	----- 107.3	2 128.4	4 104.7	9 104.7	3 120.7	3 102.3	----- 98.3	3 107.8	3 91.6	3 16.4	3 105.7
Beldi Giant-----	2777	{Y----- P-----	----- 106.2	4 118.6	8 103.6	8 105.0	2 124.8	3 101.7	----- 101.8	----- 109.7	8 93.2	8 103.7	2 102.4
White Smyrna-----	910	{Y----- P-----	----- 101.1	4 104.3	8 98.6	8 97.1	2 98.0	3 98.8	3 92.8	3 91.2	----- 85.0	8 98.8	2 85.9
Trebi-----	936	{Y----- P-----	----- 111.6	4 132.0	8 108.0	9 114.2	3 124.1	3 105.3	3 109.1	3 107.2	3 117.6	----- 116.1	2 108.7
Han River-----	206	{Y----- P-----	----- 100.6	4 120.7	8 98.2	9 101.2	3 126.9	3 95.8	3 94.0	3 96.4	3 101.3	----- 86.1	2 104.1
Wisconsin Winter	519	{Y----- P-----	----- 107.5	2 115.9	2 102.4	3 101.9	3 121.9	2 97.2	2 94.6	2 97.7	2 116.5	2 92.0	----- 96.0

Barley is an important crop in southern Idaho as feed for stock. Several varieties that have been grown successfully at this substation have been distributed to farmers in the vicinity. The cultivation of one of these, Trebi (C. I. No. 936), has spread over a considerable area in the Snake River basin. The annual yields of the barley varieties grown in plats under irrigation from 1913 to 1921, inclusive, at Aberdeen, are shown in Table 67. These yields were obtained in cooperation with the Idaho Agricultural Experiment Station. While a number of ecological groups have been represented in these experiments, the varieties of the humid groups were not grown for a sufficient number of years to justify a group digest.

In the summarized digest of Table 67 12 important or representative varieties are compared. Trebi (C. I. No. 936) is undoubtedly the best of these. During the eight years in which it was grown it produced an average acre yield more than 5 bushels higher than that of Beldi Giant (C. I. No. 2777), its nearest competitor in those years. Trebi is a selection made from a bulk importation of barley from Samsun, Asiatic Turkey. From its behavior in southern Idaho it seems probable that it was originally grown under irrigation in one of the small valleys south of the Black Sea. Following Trebi are two North African barleys of the Coast group, Beldi Giant (C. I. No. 2777) and Sandrel (C. I. No. 937). Both of these have proved to be high-yielding sorts at a number of the western stations.

Hannchen (C. I. No. 531), a 2-rowed sort, is fourth in point of yield. Hannchen is one of the few 2-rowed spring varieties that has produced high yields in the Western States, particularly in the sections where the growing season is short and the moisture supply is fairly abundant either from rainfall or irrigation. Han River (C. I. No. 206) also produced comparatively high yields. White Smyrna (C. I. No. 910), which was the highest yielding variety at this station under dry-land conditions, was the lowest of the varieties grown under irrigation in the years 1914 to 1921, inclusive. Hannchen and White Smyrna usually are

high-yielding sorts in the same general regions, Hannchen yielding better in years when there is more than the average rainfall and White Smyrna when there is less.

Odesa (C. I. No. 927) and the barleys of the Manchuria type were not tested for more than three years, as none of them is adapted to growing conditions in these arid regions. Horn (C. I. No. 926), a Chevalier variety with an average yield of 105 per cent of the weighted mean for the years 1914 to 1917, inclusive, was exceeded by the other 2-rowed varieties grown those years. Baker (C. I. No. 975) was introduced into the experiment in 1918 and was continued through 1921. This is also a pedigreed variety which was originally selected from the same bulk importation as Trebi. It has not proved so good at this station, however, as Trebi.

While the irrigated lands in southern Idaho are well adapted to the growing of barleys of the Coast type, such as Beldi Giant and Sandrel, and to the 2-rowed variety Hannchen, in nearly all of the places where Trebi has been grown it has proved superior to them.

Dry-land conditions.—Barley varieties were grown on dry land at the Aberdeen (Idaho) substation during the period from 1913 to 1918, inclusive. These yields, shown in Table 68, were obtained in cooperation with the Idaho Agricultural Experiment Station. Only two varieties were grown during the full period of the experiments. They were White Smyrna (C. I. No. 195) and Club Mariout (C. I. No. 261). These produced an average yield of 20.7 and 20.5 bushels, respectively, for the six years, so that they are essentially equal. Winter Club (C. I. No. 592) was grown from 1913 to 1916, inclusive, but was inferior to most of the spring barleys grown in those years. Although Hannchen (C. I. No. 531) was not grown in 1914, in the five years in which it was comparable it did not prove so good as White Smyrna or Club Mariout. Coast (C. I. No. 690) produced fairly good yields in 1913 and 1914, but was very poor in 1915 and consequently was dropped from the experiment the following year. Svanhals (C. I. No. 187) was grown for this same period, but is evidently not adapted to dry-farming conditions in southern Idaho.

TABLE 68.—*Annual acre yields of varieties of barley grown on dry land at the Aberdeen (Idaho) substation in part or all of the six years from 1913 to 1918, inclusive*

[Data obtained in cooperation with the Idaho Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)						Years grown	Average yield (bus.)	Percentage of weighted mean
		1913	1914	1915	1916	1917	1918			
Winter Club.....	592	13.7	26.3	8.4	1.5	-----	-----	4	12.5	60.4
Nepal.....	230	16.0	-----	-----	-----	-----	-----	1	16.0	68.7
Bavarian.....	160	14.1	-----	-----	-----	-----	-----	1	14.1	60.5
Do.....	2198	15.3	-----	-----	-----	-----	-----	1	15.3	65.7
Do.....	2199	14.3	-----	-----	-----	-----	-----	1	14.3	61.4
Gatami.....	575	27.5	-----	-----	-----	-----	-----	1	27.5	118.0
Coast.....	690	28.3	38.1	10.1	-----	-----	-----	3	25.5	111.8
White Smyrna.....	195	30.4	35.4	19.1	24.1	10.0	5.4	6	20.7	121.8
Hannchen.....	531	27.3	-----	18.5	10.4	2.5	7.2	5	13.2	96.3
Hanna.....	906	21.4	-----	-----	-----	-----	-----	1	21.4	91.8
Club Mariout.....	261	43.8	35.5	13.5	17.0	7.9	5.2	6	20.5	120.6
Princess.....	529	15.8	-----	-----	-----	-----	-----	1	15.8	67.8
Manchuria.....	244	22.2	-----	-----	-----	-----	-----	1	22.2	95.3
Svanhals.....	187	35.4	28.6	11.2	-----	-----	-----	3	25.1	110.1
Winter Club.....	488	-----	27.0	-----	-----	-----	-----	1	27.0	80.4
Maryland Winter.....	518	-----	47.0	-----	-----	-----	-----	1	47.0	139.9
Tennessee Winter.....	257	-----	43.2	15.7	3.0	2.2	-----	4	16.0	95.8
Turkestan.....	711	-----	-----	3.9	-----	-----	-----	1	3.9	29.3
Moravian.....	343	-----	13.1	-----	-----	-----	-----	1	13.1	39.0
Lion.....	923	-----	42.0	14.6	13.9	6.6	4.9	5	16.4	108.6
Bulgarian.....	521	-----	-----	1.6	-----	-----	-----	1	1.6	120.3
Beldi Giant.....	2777	-----	-----	22.5	15.8	6.2	7.4	4	13.0	131.3
Sandrel.....	937	-----	-----	20.3	11.2	4.1	5.8	4	10.3	104.0
Trebi.....	936	-----	-----	-----	13.7	5.4	6.8	3	8.6	103.6

In 1914 three winter barleys, Winter Club (C. I. No. 488), Maryland Winter (C. I. No. 518), and Tennessee Winter (C. I. No. 257), were grown. Tennessee Winter was continued in the experiment through 1917. Turkestan (C. I. No. 711) and Bulgarian (C. I. No. 521), two other winter barleys, were grown in

1915. None of these winter sorts proved successful, owing to the severity of the winters at Aberdeen. Spring barley is commonly grown in southern Idaho, although at lower altitudes winter barley is sometimes grown successfully.

Lion (C. I. No. 923) produced an average yield of 108.6 per cent of the weighted mean for the years 1914 to 1918, inclusive. Beldi Giant (C. I. No. 2777) and Sandrel (C. I. No. 937) were grown from 1915 to 1918, inclusive. Beldi Giant was exceeded in yield by only White Smyrna for those years. While Sandrel ranked second in 1915, its yield was comparatively low the other three years. Trebi (C. I. No. 936) yielded fairly well in the three years it was grown, considering the fact that it is best adapted to irrigated conditions in this section.

From Table 68 it is evident that White Smyrna, Club Mariout, and Beldi Giant are about equally well adapted to culture under dry-land conditions. White Smyrna is perhaps the better sort in years of very little rainfall.

NEPHI, UTAH

Barley varieties were tested at Nephi, Utah, in the period from 1908 to 1921, inclusive. The yields shown in Table 69 were obtained in cooperation with the Utah Agricultural Experiment Station during the years from 1908 to 1920, inclusive, and through the courtesy of this station in 1921.

In 1908 three spring barleys, Hull-less (C. I. No. 649), California Prolific (C. I. No. 651), and Coast (C. I. No. 650), were grown. Although retained for several years none of these produced high yields.

Two winter barleys, Winter Club (C. I. No. 592) and Tennessee Winter (C. I. No. 257), were sown in the fall of 1908 and were continued in the experiment through 1921. In 1909 and 1910 they produced yields much higher than those of the three spring sorts just mentioned. A number of spring barleys were sown along with the winter varieties in 1911. As can be seen from the yields in 1912, the crop was practically a failure. However, Gatami (C. I. No. 575), a spring barley, produced the highest yield that year. It was the only spring variety tested further as a winter sort, having been grown in 1914, 1915, and 1916, when it proved inferior to the true winter sorts. With the exception of Turkestan (C. I. No. 711) the other winter barleys sown in the fall of 1911 were also dropped from the experiment after that year. Turkestan was grown through 1921. Bulgarian (C. I. No. 521) was tested for the years 1914 to 1921, inclusive, making possible a comparison of the four winter barleys, Winter Club, Tennessee Winter, Turkestan, and Bulgarian, for those eight years. Bulgarian, with an average yield of 33.7 bushels, was the best of these four. Turkestan was second, with Tennessee Winter and Winter Club third and fourth, respectively.

TABLE 69.—*Annual acre yields of varieties of barley grown at the Nephi (Utah) substation in one or more of the 14 years from 1908 to 1921, inclusive*

[Data for 1908 to 1920, inclusive, obtained in cooperation with the Utah Agricultural Experiment Station, through whose courtesy the data for 1921 are used. The varieties marked with an asterisk (*) were sown in the fall]

Variety	C. I. No.	Acre yields (bushels)														Years grown	Average yield (bush.)	Percentage of weighted mean
		1908	1909	1910	1911	1912	1913	1914	1915	1916	1917	1918	1919	1920	1921			
Hull-less	649	16.0	7.7	---	---	---	---	---	---	---	---	---	---	---	---	2	11.9	86.9
California Prolific	651	20.2	7.7	1.2	---	---	---	---	---	---	---	---	---	---	---	3	9.7	81.3
Coast	650	13.3	4.2	1.4	---	---	---	---	---	---	---	---	---	---	---	3	6.9	51.8
*Tennessee Winter	257	---	18.1	16.5	24.6	4.1	1.0	50.9	43.5	8.1	21.7	26.0	13.6	16.7	41.1	13	22.0	108.4
*Winter Club	592	---	22.0	9.8	32.7	4.6	0	39.8	53.5	12.7	16.7	20.1	13.6	16.7	36.2	13	21.4	105.4
*Nima	2196	---	---	---	22.5	---	---	---	---	---	---	---	---	---	---	1	22.5	84.6
*Virginia Hooded	648	---	---	---	---	1.4	---	---	---	---	---	---	---	---	---	1	1.4	41.2
*Black Hull-less	596	---	---	---	---	1.6	---	---	---	---	---	---	---	---	---	1	.6	17.6
*Chevalier	156	---	---	---	---	3.9	---	---	---	---	---	---	---	---	---	1	3.9	114.7
*Orel	351	---	---	---	---	3.2	---	---	---	---	---	---	---	---	---	1	3.2	94.1
*Hannchen	531	---	---	---	---	4.0	---	---	---	---	---	---	---	---	---	1	4.0	117.6
*Turkestan	711	---	---	---	---	2.9	0	61.7	50.2	7.9	25.1	30.4	14.3	12.5	61.3	10	26.6	123.7
*Gatami	575	---	---	---	---	5.7	---	43.5	38.0	3.0	---	---	---	---	---	4	22.5	96.6
*Bulgarian	521	---	---	---	---	---	---	59.0	49.6	13.7	24.3	36.5	18.6	19.4	48.8	8	33.7	127.7
Coast	690	---	---	---	---	---	---	27.3	13.7	19.0	16.7	6.2	10.7	---	---	6	15.6	74.3
White Smyrna	195	---	---	---	---	---	---	29.5	16.8	23.3	11.9	7.3	9.1	---	---	6	16.3	77.6

Both Coast (C. I. No. 690) and White Smyrna (C. I. No. 195) were spring sown for the years 1915 to 1920, inclusive, but were surpassed by the four winter barleys for that period.

Nephi, which is located in an intermountain valley, is evidently better suited to the growing of winter than of spring varieties. The winter sorts are somewhat protected from extremes of temperature and ripen about 10 days before the spring varieties, thus escaping the drought which usually begins about June 1.

PULLMAN, WASH.

The results of barley tests at Pullman, Wash., for the period from 1914 to 1921, inclusive, were obtained through the courtesy of the Washington Agricultural Experiment Station. These data are found in Table 70. Ten spring and four winter varieties were tested.

Of the spring sorts Coast (C. I. No. 1249) produced the highest average yield. Beldi Giant (C. I. No. 2777) was almost the equal of Coast. Following Beldi Giant is Blue (C. I. No. 1247), which was grown for the seven years from 1915 to 1921, inclusive. Club Mariout (C. I. No. 261) ranked fourth in point of yield for the four years in which it was grown. All of these four varieties are of North African origin. Next in order is Horsford (C. I. No. 1775), which is noteworthy only because it produced a higher average yield than White Smyrna. Horsford, like Nepal, usually is one of the poorest sorts. Both Eureka (C. I. No. 1250) and Nepal (C. I. No. 595), 6-rowed, hooded, hull-less varieties, produced comparatively low yields.

Of the winter barleys grown for the eight years of the experiments Winter Club (C. I. No. 592) ranks first. It exceeded all the spring varieties for this period. When sown in spring, however, Winter Club did not compare favorably with the spring sorts. Following Winter Club are Wisconsin Winter (C. I. No. 519) and Maryland Winter (C. I. No. 518). Neither gave as high average yields as Coast or Beldi Giant. Wisconsin Winter (C. I. No. 1894) was first tested in 1919. It was inferior to Winter Club, but exceeded all other winter or spring varieties in the three years from 1919 to 1921, inclusive.

While the spring barleys undoubtedly are of more commercial importance, it is of especial interest that the winter barleys produce such high yields in the same locality. It is seldom that both types can be successfully grown in the same district. At Moscow, Idaho, only a few miles distant, the relative yield of the winter sorts was even higher. It is unlikely that their culture will be greatly increased in the near future, as the seeding time is the same as that of winter wheat, and labor is not often available for the extension of the fall-seeded acreage. The yields following severe winters drop very low.

TABLE 70.—*Annual acre yields of varieties of barley grown at the Washington Agricultural Experiment Station (at Pullman) in some or all of the eight years from 1914 to 1921, inclusive*

[Data obtained through the courtesy of the Washington Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)								Years grown	Average yield (bus.)	Per- centage of weight- ed mean
		1914	1915	1916	1917	1918	1919	1920	1921			
Spring barleys:												
Beldi Giant.....	2777	43.2	60.2	85.4	37.5	25.8	69.1	79.8	52.3	8	56.7	108.6
Coast.....	1249	49.2	57.8	89.6	37.1	28.4	65.0	72.5	55.2	8	56.9	109.0
Eureka.....	1250	36.7	45.9	65.0	20.3	16.9	50.9	57.9	33.5	8	40.9	78.4
Nepal.....	595	42.6	39.5	73.5	16.6	12.6	43.4	59.1	-----	7	41.0	79.0
Excelsior.....	1218	40.7	47.7	68.7	-----	-----	-----	-----	-----	3	52.4	91.8
Blue.....	1247	-----	58.7	87.5	30.8	23.7	60.1	72.2	53.5	7	55.2	104.7
Winter Club.....	592	-----	41.8	89.8	23.5	30.0	55.4	61.0	48.9	7	50.1	95.1
Horsford.....	1775	-----	-----	-----	39.1	24.1	55.1	70.6	38.3	5	45.4	90.8
Club Mariout.....	261	-----	-----	-----	-----	25.0	56.0	65.4	63.3	4	52.4	98.1
White Smyrna.....	658	-----	-----	-----	-----	25.2	46.0	61.0	48.3	4	45.1	84.5
Winter barleys:												
Maryland Winter.....	518	50.6	76.3	40.8	38.6	24.5	61.8	70.2	53.5	8	52.0	99.6
Winter Club.....	592	68.9	59.4	24.6	47.1	16.7	88.2	91.2	65.6	8	57.7	110.5
Wisconsin Winter.....	519	49.9	63.4	40.4	36.8	24.1	88.8	72.4	65.0	8	55.1	105.6
Do.....	1894	-----	-----	-----	-----	-----	93.7	73.1	75.8	3	80.9	128.4

LIND, WASH.

The yields of barley at the Adams Branch Experiment Station at Lind, Wash., were obtained in cooperation with the Washington Agricultural Experiment Station until the end of 1920 and through the courtesy of that station in 1921. These yields for the 5-year period from 1917 to 1921, inclusive, are shown in Table 71. Only two varieties, Coast (C. I. No. 1249) and White Smyrna (C. I. No. 910), were grown for the entire period. Coast was only slightly superior, with no significant difference, as its average yield was 13.8 bushels, as compared to 13.3 bushels for White Smyrna. The 2-rowed varieties, Hannchen (C. I. No. 531) and White Smyrna, are usually successful sorts in the same general localities, but Hannchen was surpassed by the other four varieties grown at this station from 1917 to 1919, inclusive. The two hooded barleys Eureka (C. I. No. 1250) and Excelsior (C. I. No. 1248) produced the lowest yields in 1917 and were discarded after that year. Meloy (C. I. No. 1176) and Himalaya (C. I. No. 620) were grown for the four years from 1918 to 1921, inclusive. Meloy was exceeded only by Coast in those years. This is a hooded hulled variety that has yielded well in recent years at a number of the western stations. Trebi (C. I. No. 936), which was grown in 1918 and 1919, was exceeded by both Coast and Meloy. Horsford (C. I. No. 1775) yielded 20.1 bushels in 1919, the highest that year, but was superior only to Himalaya the two following years.

The three winter barleys, Winter Club (C. I. No. 592), Hybrid (C. I. No. 3168), and Wisconsin Winter (C. I. No. 519), grown in 1918 and 1919, were very low in productivity. At Pullman both Winter Club and Wisconsin Winter were superior in yield, but it is evident that in this section of Washington the winters are too severe for the successful cultivation of these varieties.

Of the barleys tested at Lind three varieties of entirely different types did almost equally well. Coast, the 6-rowed bearded sort, is the one commonly grown in this section. White Smyrna, a 2-rowed bearded variety, has produced high yields at a number of western stations, but it has not gained a wide commercial reputation, because of its short straw and general appearance which are very much against it. Meloy, a 6-rowed hooded hull-less variety, has been only recently tested, but from its behavior at the various stations where it is grown it is likely that it will prove valuable in this section.

TABLE 71.—*Annual acre yields of varieties of barley grown at the Adams Branch Experiment Station, Lind, Wash., in part or all of the five years from 1917 to 1921, inclusive*

[Data obtained in cooperation with or through the courtesy of the Washington Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)					Years grown	Average yield (bus.)	Percentage of weighted mean
		1917	1918	1919	1920	1921			
Club Mariout.....	932	13.9	7.0	13.1	-----	-----	3	11.3	104.6
Coast.....	1249	15.0	6.4	18.6	12.1	16.9	5	13.8	121.1
Blue.....	1247	13.3	6.4	16.7	-----	-----	3	12.1	112.0
Hannchen.....	531	12.5	6.0	15.6	-----	-----	3	11.4	105.6
White Smyrna.....	910	16.2	7.4	15.0	11.2	16.7	5	13.3	116.7
Eureka.....	1250	8.2	-----	-----	-----	-----	1	8.2	66.1
Excelsior.....	1248	8.0	-----	-----	-----	-----	1	8.0	64.5
Trebi.....	936	-----	6.5	17.1	-----	-----	2	11.8	114.6
Meloy.....	1176	-----	5.8	17.9	11.2	17.0	4	13.0	117.1
Himalaya.....	620	-----	6.4	13.4	9.2	11.5	4	10.1	91.0
Winter Club (fall sown).....	592	-----	2.8	10.2	-----	-----	2	6.5	63.1
Hybrid (fall sown).....	3168	-----	1.3	-----	-----	-----	1	1.3	24.5
Wisconsin Winter (fall sown).....	519	-----	2.2	10.6	-----	-----	2	6.4	62.1
Horsford.....	1775	-----	-----	20.1	10.8	13.4	3	14.8	104.2

MORO, OREG.

Varietal experiments with barley were conducted at the Sherman County Branch Station, Moro, Oreg., in cooperation with the Oregon Agricultural Experiment Station from 1910 to 1921, inclusive. The annual yields are reported in Table 72. Of the five groups compared, the Coast was far superior to the Hanna, which ranked second. The Tennessee Winter group was third, Manchuria fourth, and Hull-less fifth. No yields were recorded for the Tennessee Winter varieties in 1913. It is probable that they winterkilled that year. If so, this would lower the place of rank of the group.

TABLE 72.—*Annual acre yields of varieties of barley grown at the Sherman County Branch Station, Moro, Oreg., in part or all of the 12 years from 1910 to 1921, inclusive*

[Data obtained in cooperation with the Oregon Agricultural Experiment Station. The varieties in the table proper marked with an asterisk (*) were sown in the fall]

Variety	C. I. No.	Acre yields (bushels)												Years grown	Average yield (bus.)	Percentage of weighted mean
		1910	1911	1912	1913	1914	1915	1916	1917	1918	1919	1920	1921			
Himalaya	620	14.7	2.9	6.0	---	---	---	---	---	---	---	---	---	3	7.9	53.4
Coast	626	14.1	2.6	23.3	38.3	32.5	51.2	72.5	36.5	20.4	---	---	---	9	32.4	102.5
Bohemian	188	14.4	---	---	---	---	---	---	---	---	---	---	---	1	14.4	137.1
Beldi Dwarf	190	14.3	3.5	15.5	37.5	36.2	48.8	59.9	35.8	21.5	38.1	40.6	42.1	12	32.8	101.5
Nepal	595	14.2	3.0	---	---	---	---	---	---	---	---	---	---	2	8.6	106.8
White Smyrna	658	14.1	9.3	27.0	42.0	36.7	49.8	71.8	38.0	22.8	35.0	30.4	37.7	12	34.5	108.9
Club Mariout	261	13.9	6.3	28.0	40.6	42.1	54.5	69.3	45.4	29.0	40.2	39.2	41.7	12	37.5	116.1
Hannchen	531	11.3	8.4	25.2	48.3	32.8	46.4	68.3	36.8	25.0	39.8	28.5	33.4	12	33.7	104.3
Oderbrucker	537	13.4	7.0	23.0	53.0	63.4	51.5	61.9	39.2	22.9	---	---	---	9	30.5	96.5
Horsford	610	13.4	5.4	24.5	14.4	20.8	22.5	---	---	---	---	---	---	6	16.8	63.9
Mansury	617	13.4	2.7	21.2	23.1	24.2	---	---	---	---	---	---	---	5	18.5	80.1
Princess	529	9.5	3.3	20.3	20.5	19.1	---	---	---	---	---	---	---	5	14.5	62.8
Evans	892	9.7	2.3	---	---	---	---	---	---	---	---	---	---	2	6.0	75.9
Hanna	24	5.6	10.0	26.0	29.5	26.9	40.0	---	---	---	---	---	---	6	23.0	87.5
Gatami	575	7.7	5.9	16.8	36.4	41.2	36.0	45.2	---	---	---	---	---	7	27.0	84.9
Svanbals	187	7.3	13.8	25.1	34.1	30.6	38.7	71.5	31.9	18.4	34.0	25.6	32.7	12	30.3	93.8
Black Hull-less	596	5.0	3.7	17.8	37.9	25.2	36.6	---	---	---	---	---	---	6	21.0	79.8
Chevalier II	200	3.1	6.4	26.2	31.0	21.8	32.5	---	---	---	---	---	---	6	20.2	76.8
Imperial	289	9	4.3	25.0	17.3	---	---	---	---	---	---	---	---	4	11.9	64.3
Manchuria	244	---	3.2	3.9	33.0	34.6	27.0	32.2	---	---	---	---	---	6	25.2	74.1
*Tennessee Winter	257	---	5.0	24.9	---	44.9	44.7	53.9	---	---	---	---	---	5	34.7	101.5
*Baku	709	---	8	2.5	---	---	---	---	---	---	---	---	---	2	1.7	10.6
*Maryland Winter	518	---	8.3	26.0	---	51.7	46.1	56.1	---	---	---	---	---	5	37.6	109.9
*Wisconsin Winter	519	---	12.2	37.7	---	44.6	---	---	---	---	---	---	---	3	31.5	134.0
*Texas Winter	554	---	11.0	22.5	---	59.4	46.5	67.5	---	---	---	---	---	5	41.4	121.1
*Winter Club	592	---	11.0	21.1	---	35.0	30.4	60.9	---	---	---	---	---	5	31.7	92.7
*Chevalier	612	---	7.7	28.3	---	51.0	39.0	51.4	---	---	---	---	---	5	35.5	103.8
*Nesbian	647	---	---	34.8	---	42.3	---	---	---	---	---	---	---	2	38.5	131.4
*Turkestan	711	---	---	22.7	---	---	---	---	---	---	---	---	---	1	22.7	97.8
*Chevalier	156	---	---	25.4	---	34.2	---	---	---	---	---	---	---	2	29.8	101.7
*Bavarian	159	---	---	22.0	---	33.9	---	---	---	---	---	---	---	2	27.9	95.2
*Orel	351	---	---	21.0	---	40.2	---	---	---	---	---	---	---	2	30.6	104.4
*Hannchen	531	---	---	32.0	---	50.6	---	---	---	---	---	---	---	2	41.3	141.0
*Phoson	516	---	---	31.1	---	---	---	---	---	---	---	---	---	1	34.1	147.0
*Gatami	575	---	---	29.0	---	48.7	---	---	---	---	---	---	---	2	38.9	132.8
*Hanse Hull-less	703	---	---	20.7	---	---	---	---	---	---	---	---	---	1	20.7	89.2
*Black Hull-less	596	---	---	21.2	---	39.2	---	---	---	---	---	---	---	2	30.2	103.1
*Virginia Hooded	648	---	---	13.1	---	30.8	---	---	---	---	---	---	---	2	21.9	74.7
Bevede	2295	---	---	20.7	33.1	23.3	27.0	---	---	---	---	---	---	4	26.0	79.8
Manchuria	1275	---	---	22.3	25.4	---	---	---	---	---	---	---	---	2	23.9	89.8
O. A. C. 21	1470	---	---	---	34.0	23.4	---	---	---	---	---	---	---	2	28.7	84.2
Manchuria	241	---	---	---	34.3	25.4	---	---	---	---	---	---	---	2	29.9	87.7
Manchurian	739	---	---	---	38.7	29.2	27.2	---	---	---	---	---	---	3	31.7	87.3
Black Algerian	2296	---	---	---	---	46.3	46.4	37.5	---	---	---	---	---	3	43.4	95.8
Chile Brewing	2297	---	---	---	---	23.7	47.5	69.2	34.4	---	---	---	---	4	43.7	99.1
Himalaya	2298	---	---	---	---	33.3	39.3	---	---	---	---	---	---	2	36.3	97.1
Do	2299	---	---	---	---	25.0	42.2	66.8	37.9	24.8	40.1	29.8	36.0	8	37.8	94.7
Black Abyssinian	2300	---	---	---	---	24.3	---	---	---	---	---	---	---	1	24.3	69.8
Franconian	679	---	---	---	---	29.2	47.5	66.1	29.0	---	---	---	---	4	42.9	97.3
Coast	2301	---	---	---	---	37.9	42.6	67.1	40.6	18.8	43.1	31.0	38.4	8	39.9	100.0
Peru	2302	---	---	---	---	40.4	48.3	60.9	45.0	22.3	46.9	38.3	41.7	8	43.0	107.8
Mahan	1144	---	---	---	---	36.0	---	---	---	---	---	---	---	1	36.0	88.2
Withycombe	2303	---	---	---	---	36.5	59.4	---	---	---	---	---	---	2	47.9	91.4
Meloy	1176	---	---	---	---	51.1	63.9	47.5	18.6	52.5	36.2	34.8	---	7	43.5	105.1
*Dufur	2304	---	---	---	---	40.8	56.4	---	---	---	---	---	---	2	48.6	92.7
Servian	915	---	---	---	---	---	55.0	---	---	---	---	---	---	1	55.0	85.4
Horn	926	---	---	---	---	---	79.6	34.6	---	---	---	---	---	2	57.1	105.9
Odessa	927	---	---	---	---	---	81.7	37.5	19.0	44.2	18.8	36.3	---	6	39.6	95.4
Peruvian	935	---	---	---	---	---	77.5	43.7	24.1	44.4	39.3	43.4	---	6	45.4	109.4
Trebi	936	---	---	---	---	---	95.8	27.5	24.2	47.3	28.5	53.4	---	6	42.9	103.4
Sandrel	937	---	---	---	---	---	84.2	34.6	21.7	37.9	23.5	---	---	5	40.4	95.7
Hanna	906	---	---	---	---	---	69.6	32.8	19.0	37.5	26.9	26.4	---	6	35.4	85.3
Flynn	1311	---	---	---	---	---	---	---	21.7	44.6	37.9	38.8	---	4	35.7	110.5
White Smyrna	2169	---	---	---	---	---	---	---	---	---	27.5	40.6	---	2	34.1	102.4
Ace	1853	---	---	---	---	---	---	---	---	---	17.5	---	---	1	17.5	58.1
Beldi Giant	2777	---	---	---	---	---	---	---	---	---	---	21.6	---	1	21.6	71.8
California Mariout	1455	---	---	---	---	---	---	---	---	---	---	33.3	---	1	33.3	90.0

TABLE 72.—*Annual acre yields of varieties of barley grown at the Sherman County Branch Station, Moro, Oreg., in part or all of the 12 years from 1910 to 1921, inclusive—Continued*

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 13 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years. Under "Data shown" P=Percentage yield and Y=Years comparable]

Variety	C. I. No.	Data shown	Varieties and percentages											
			Chevalier II	Svanhals	Hannchen	Coast	Peruvian	Club Mariout	White Smyrna	Beldi Dwarf	Trebi	Himalaya	Meloy	Flynn
Chevalier II	200	Y----- P-----	6 80.9	6 70.2	6 74.7	(*)	6 65.3	6 67.6	6 77.7	(*)	2 80.8	1 63.6	(*)	4 62.3
Svanhals	187	Y----- P-----	6 123.6	12 90.0	9 93.1	6 78.6	6 80.8	6 87.7	6 92.4	6 83.1	6 93.7	6 83.0	7 77.4	5 86.9
Hannchen	531	Y----- P-----	6 142.5	12 111.1	9 103.8	6 85.1	6 89.8	6 97.5	6 102.6	6 89.9	6 102.8	6 91.3	7 88.6	5 87.5
Coast	626	Y----- P-----	6 133.9	9 107.4	9 96.3	6 89.1	6 88.5	6 93.5	6 106.7	6 87.7	6 108.3	6 99.7	4 94.0	1 88.0
Peruvian	935	Y----- P-----	(*) 127.2	6 117.5	6 112.3	6 102.9	6 115.6	6 114.5	6 105.7	6 115.7	6 107.5	6 105.7	6 114.8	
Club Mariout	261	Y----- P-----	6 153.2	12 123.8	9 111.4	6 112.9	6 97.2	6 108.6	6 114.3	6 102.8	6 119.4	6 104.8	8 105.0	5 96.8
White Smyrna	658	Y----- P-----	6 147.9	12 114.0	9 102.6	6 106.9	6 86.5	6 92.1	6 105.3	6 91.5	6 106.5	6 93.7	8 88.0	5 94.1
Beldi Dwarf	190	Y----- P-----	6 128.8	12 108.3	9 97.4	6 93.7	6 87.4	6 87.5	6 95.0	6 92.4	6 106.7	6 94.2	7 99.5	4 79.2
Trebi	936	Y----- P-----	(*) 120.4	6 111.2	6 114.0	6 94.6	6 97.3	6 109.3	6 108.3	6 109.5	6 101.7	6 94.0	6 141.9	
Himalaya	2299	Y----- P-----	2 123.8	8 106.8	5 97.3	6 92.3	6 86.4	6 83.7	6 93.9	6 93.7	6 91.3	7 91.1	4 91.4	5 77.3
Meloy	1176	Y----- P-----	1 157.2	7 120.5	4 109.5	6 100.3	6 93.1	6 95.4	6 106.7	6 98.4	6 109.7	6 99.4	4 100.9	
Flynn	1311	Y----- P-----	(*) 129.2	4 112.9	1 106.4	4 94.6	4 95.3	4 113.6	4 100.5	4 106.4	4 109.4	4 100.6	4 -----	(*)
Texas Winter	554	Y----- P-----	4 160.4	5 115.1	5 114.2	5 113.6	1 87.1	5 103.3	5 106.3	5 126.2	1 70.5	3 129.4	2 99.1	-----

Thirteen representative varieties are compared in the summarized digest of Table 72. Of these, Peruvian (C. I. No. 935) produced the highest yields. It is a North African barley of the Coast group. Second in point of yield was Club Mariout (C. I. No. 261), also a North African barley but not of the Coast type. Trebi (C. I. No. 936) produced very high yields. Considering the low precipitation in this part of the Columbia Basin it would hardly be expected that Trebi would be so successful here. However, the soil in this region absorbs water quite readily and has a high capacity for retaining it. Following Trebi are two hybrid sorts, Flynn (C. I. No. 1311) and Meloy (C. I. No. 1176), both of which have been distributed from the Moro station. Flynn is a smooth-awned barley and Meloy a hooded one. The stiff awns of barley cause many farmers to object to its cultivation. The smooth-awned and hooded sorts are not open to this objection. Flynn came from a cross of a smooth-awned barley on Club Mariout, which parent it has almost equaled in yield. Meloy is by far the best of the hooded barleys widely tested in the West, producing higher yields and shattering less than any other hooded hybrid.

In the earlier years a number of winter barleys were grown, but they were all dropped from the experiments after 1916. Texas Winter (C. I. No. 554) produced the highest average yield of those tested for five years. While a small acreage of winter barley is grown successfully in this basin, the possibilities of winter-killing are too great for the crop to become of much commercial importance.

White Smyrna (C. I. No. 658) was grown for the full 12 years of the experiments, producing an average yield of 107 per cent of the weighted mean. Hannchen (C. I. No. 531) was also grown during the entire period, giving an average yield of 104 per cent of the weighted mean. Beldi Dwarf (C. I. No. 190) produced fair yields, but was not so good as either of the two just mentioned for the 12 years.

As can be seen from Table 72, the humid-area spring varieties, such as the 6-rowed Manchuria, Oderbrucker, and Odessa and the 2-rowed sorts, Princess, Hanna, and Chevalier, are entirely out of their ecological range. From the results at Moro it may be seen that the North African barleys, particularly those of the Coast group, and Club Mariout, are the ones best adapted to this region. Coast has been the common barley grown in this section for many years.

UNION, OREG.

Table 73 shows the varietal yields of barley grown at the Eastern Oregon Branch Station (at Union) in seven of the years from 1910 to 1921, inclusive. Because of the varying periods in which some of the barleys were grown, a varietal comparison is almost impossible. The plats also varied greatly in size, thus making conclusions less dependable than if they had been uniform. In 1910 all the higher yielding varieties except Beardless (C. I. No. 639) were selections made at this station. Two hybrids of the Nepal type, Union (C. I. Nos. 2198 and 2199), produced high average yields in the four years they were grown. Blue (C. I. No. 1247), a barley of the Coast type, is the only variety tested for as many as six years. It produced an average yield of 102 per cent of the weighted mean. Of the varieties grown during the more recent period, 1919 to 1921, inclusive, Trebi (C. I. No. 936) is perhaps the best. Hannchen (C. I. No. 531) produced an average yield of 98 per cent of the weighted mean. The 2-rowed varieties, Horn (C. I. No. 926) and White Smyrna (C. I. No. 910), yielded well in 1919 and 1920. Club Mariout (C. I. No. 261) and Flynn (C. I. No. 1311), which were introduced in 1921, were the best varieties that year.

TABLE 73.—*Annual acre yields of varieties of barley grown at the Eastern Oregon Branch Station (at Union) in one or more of the years from 1910 to 1921, inclusive*

[Data furnished through the courtesy of the Oregon Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)							Years grown	Average yield (bus.)	Percentage of weighted mean
		1910	1911	1912	1914	1919	1920	1921			
Beardless	639	56.5	42.6	44.6	49.2			28.5	5	44.3	101.8
Gutekorn	606	41.5			54.4				2	47.9	103.9
Marsbury	3121	40.8			47.3	26.3	55.1		4	42.4	98.1
Blue	1247	38.2	32.8	51.1	62.0	27.4	46.3		6	43.0	101.9
Primus	532	36.5							1	36.5	79.0
Highland Chief	883	34.9							1	34.9	75.5
Princess	529	32.8			52.7	20.0			3	35.2	84.8
Chevalier II	200	32.6			55.1	17.8			3	35.2	84.8
Svanhals	187	32.5							1	32.5	70.3
Sexradigt	607	25.9							1	25.9	56.1
Black Hull-less	596	20.3							1	20.3	43.9
Union	3122	57.1	45.6	26.7					3	43.1	99.8
Do	3123	58.8	44.1						2	51.5	113.7
Do	3179	50.8	38.2						2	44.5	98.2
Do	2198	63.7	65.2	28.1	46.7				4	50.9	116.5
Do	3124	63.9	58.3						2	61.1	134.9
Do	3127	54.6	42.0						2	48.3	106.6
Do	3125	53.7	36.4						2	45.1	99.6
Do	3126	46.9	28.6						2	37.7	83.2
Do	3128	50.6	45.8						2	48.2	106.4
Do	3129	51.5	43.3						2	47.4	104.6
Do	3130	50.7	33.6	35.6	36.4				4	39.1	89.5
Do	2199	58.7	68.4	34.5	39.1				4	50.2	114.9
Do	3131	50.9			23.1				2	37.0	80.3
Do	3132	41.8			49.8				2	45.8	99.3
Do	3134	56.0	33.7	22.1					3	37.3	86.3
Do	3133		39.8	21.2					2	30.5	76.1
New Hull-less	3135			39.7	35.6				2	37.7	92.6
Odessa	927					23.6	52.3	31.0	3	35.6	98.2
Sandrel	937					25.7	51.7		2	38.7	102.4
White Smyrna	910					26.7	56.4		2	41.5	109.8
Hannchen	531					26.0	48.1		2	37.1	98.1
Trebi	936					32.6	64.4	41.1	3	46.0	120.4
Horn	926					31.9	52.1		2	42.0	111.1
Peruvian	935					21.0	45.5		2	33.3	88.1
Nepal	595					42.2			1	42.2	157.5
Flynn	1311							45.8	1	45.8	114.2
Club Mariout	261							54.2	1	54.2	135.2

Although the barleys grown in recent years at Union have not been tested sufficiently to show their relative value, the behavior of Trebi, Club Mariout, and Flynn here and in other sections of Oregon indicates that they would be desirable sorts to grow.

BURNS, OREG.

Dry-land conditions.—The yields of barley varieties grown on dry land at the Harney Valley Branch Station, Burns, Oreg., were obtained in cooperation with the Oregon Agricultural Experiment Station for the seven years from 1913 to 1919, inclusive, and through the courtesy of that station for the years 1920 and 1921. Burns is situated in an old lake bed on a high plateau between the mountain ranges. The annual precipitation is low, and although there are no such extreme climatic changes as in the humid States frosts may occur in any month of the year. Because of this fact and the low rainfall, there is only one good crop in about every three on the dry land. The yields of varieties tested on dry land are shown in Table 74.

In the summarized digest of Table 74 a comparison of 12 well-known varieties is shown. The 2-rowed sorts, Hannchen (C. I. No. 531) and White Smyrna (C. I. No. 195), produced the highest yields. The data in Table 74 indicate that Hannchen was usually the better of the two in the years of heavier rainfall and that in the drier years the reverse was true. Following White Smyrna was Coast (C. I. No. 690), which was the best of the 6-rowed barleys compared. Svanhals (C. I. No. 187) was fourth in point of yield. Winter Club (C. I. No. 592) grown as a spring variety gave fair results. It was the poorest of the varieties compared, however, when fall sown. As can be seen from this table, the barleys of the Manchuria group are entirely out of their region of adaptation. The hooded and hull-less varieties, Horsford (C. I. No. 610), Nepal (C. I. No. 595), and Black Hull-less (C. I. No. 596), also produced very low yields.

In the fall of 1912 two spring and four winter varieties were sown. They produced low yields as compared with the spring-sown varieties, probably owing to both lack of precipitation and partial winterkilling. Winter Club (C. I. No. 592) and Tennessee Winter (C. I. No. 257) were the only fall-sown varieties tested further. Neither of these could compete at all with those which were spring sown. Trebi (C. I. No. 936) was grown in 1919 and 1920, but it is evident that it can not succeed here without irrigation.

Hannchen and White Smyrna are unquestionably the superior varieties on the dry lands. At the stations in eastern Oregon the 2-rowed varieties have yielded relatively higher than elsewhere in the region west of the Rocky Mountains although they have been promising at many other places.

Irrigated land.—Barley varieties have been tested on irrigated land at the Harney Valley Branch Station, Burns, Oreg., for the five years from 1917 to 1921, inclusive. The data were obtained in cooperation with the Oregon Agricultural Experiment Station for the years 1917 to 1919, inclusive, and through the courtesy of that station for the years 1920 and 1921. The annual yields of these barleys are found in Table 75. While it is difficult to compare the varieties when grown for such varying periods of years, it is obvious that Hannchen (C. I. No. 531), White Smyrna (C. I. No. 658), and Trebi (C. I. No. 936) were superior sorts at this station when tested under irrigation. Coast (C. I. No. 690) was inferior to these varieties in all the years for which it was comparable with them with the exception of 1920, when it gave the unusually high yield of 81.3 bushels, the highest in that year. Svanhals (C. I. No. 187) was fourth in point of yield. Under irrigation, Winter Club (C. I. No. 592) was quite successful when spring sown. It gave the highest average yield of all the varieties grown during the years 1917, 1920, and 1921. Trebi (C. I. No. 936) was tested only in the years 1919 and 1920 but gave the highest average for those years. The hooded barleys Nepal (C. I. No. 595) and Horsford (C. I. No. 610) were inferior in yield.

While both Trebi and Winter Club are promising sorts, their values are not so well established as those of Hannchen and White Smyrna.

TABLE 74.—Annual acre yields of varieties of barley grown on dry land at the Harney Valley Branch Station, Burns, Oreg., in one or more of the nine years from 1913 to 1921, inclusive

TABLE 75.—*Annual acre yields of varieties of barley grown on irrigated land at the Harney Valley Branch Station, Burns, Oreg., in part or all of the five years from 1917 to 1921, inclusive*

[Data for years 1917, 1918, and 1919 were obtained in cooperation with the Oregon Agricultural Experiment Station, through whose courtesy the data for 1920 and 1921 were furnished]

Variety	C. I. No.	Acre yields (bushels)					Years grown	Average yield (bus.)	Percentage of weighted mean
		1917	1918	1919	1920	1921			
White Smyrna.....	658	17.5	43.0	35.6	75.0	-----	4	42.8	106.5
Svanhals.....	187	11.0	38.5	-----	61.4	50.0	4	40.2	95.9
Hannchen.....	531	17.9	40.0	49.1	63.0	55.0	5	45.0	107.7
Coast.....	690	12.5	35.7	27.9	81.3	43.1	5	40.1	95.9
Winter Club.....	592	7.9	-----	-----	80.9	70.3	3	53.0	121.8
Club Mariout.....	261	13.9	-----	-----	-----	-----	1	13.9	110.3
Horsford.....	610	7.5	-----	-----	35.5	-----	2	21.5	51.8
Coast.....	2305	-----	32.0	-----	-----	-----	1	32.0	88.2
Nepal.....	595	-----	28.5	-----	58.3	30.7	3	39.2	75.2
Trebi.....	936	-----	-----	52.0	67.3	-----	2	59.7	104.7
Blackhull.....	878	-----	-----	-----	53.1	-----	1	53.1	83.0

NEWLANDS, NEV.

The results of varietal experiments at the Newlands Experiment Farm are used through the courtesy of the Office of Western Irrigation Agriculture. The yields of these varieties for the period from 1915 to 1919, inclusive, are shown in Table 76. Four of the seven varieties tested were grown during the entire period. Coast (C. I. No. 626) was the best of these. It produced an average yield of 34.6 bushels, which was 7.6 bushels higher than that of Hannchen (C. I. No. 531), its nearest competitor. Kent (C. I. No. 2644), a local barley of the Coast type, also was superior to Hannchen for the four years from 1915 to 1918, inclusive. It was exceeded only by Coast in that period. Besides Hannchen, two other 2-rowed varieties, Svanhals (C. I. No. 187) and Chevalier (C. I. No. 1142), were grown. They do not compare favorably with the two Coast barleys. Nepal (C. I. No. 595) was the lowest yielding sort. Trebi (C. I. No. 936), grown in the years from 1917 to 1919, inclusive, excelled all other varieties for those years with the exception of Coast.

It is quite obvious that the 2-rowed varieties are not superior here. The barleys of the Coast type are outstanding in yield at Newlands.

TABLE 76.—*Annual acre yields of varieties of barley grown at the Newlands (Nev.) Experiment Farm in some or all of the five years from 1915 to 1919, inclusive*

[Data used through the courtesy of the Office of Western Irrigation Agriculture]

Variety	C. I. No.	Acre yields (bushels)					Years grown	Average yield (bus.)	Percentage of weighted mean
		1915	1916	1917	1918	1919			
Coast.....	626	42.5	36.8	20.4	50.7	22.4	5	34.6	126.3
Svanhals.....	187	29.9	29.3	17.5	34.8	16.8	5	25.7	93.8
Hannchen.....	531	32.9	30.3	21.7	34.3	16.0	5	27.0	98.5
Nepal.....	595	25.4	26.9	10.6	32.7	9.6	5	21.0	76.6
Kent.....	2644	29.9	39.9	20.5	32.0	-----	4	30.6	102.3
Chevalier.....	1142	-----	31.3	17.4	35.9	17.6	4	25.5	96.2
Trebi.....	936	-----	-----	23.6	40.1	18.9	3	27.5	111.3

CHICO, CALIF.

Varietal tests of barley were conducted at Chico, Calif., from 1910 to 1921, inclusive. This work was done in cooperation with the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture. The annual yields from the varietal tests are given in Table 77. Although in recent years plats have been replicated, the results at Chico were not entirely satisfactory, as the soil used was not sufficiently uniform to give wholly satisfactory results.

TABLE 77.—*Annual acre yields of varieties of barley grown at the United States Plant Introduction Garden, Chico, Calif., in part or all of the 12 years from 1910 to 1921, inclusive*

[Data obtained in cooperation with the Office of Foreign Seed and Plant Introduction]

Variety	C. I. No.	Acre yields (bushels)												Years grown	Average yield (bus.)	Percent- age of weighted mean
		1910	1911	1912	1913	1914	1915	1916	1917	1918	1919	1920	1921			
Beldi Dwarf.....	190	13.8	58.4	50.0	41.2	47.0	26.6	49.2	52.0	58.5	47.7	65.7	28.4	12	44.9	110.6
Telli.....	194	18.0	60.8	60.0	41.2	40.3	23.0							6	40.5	102.8
White Smyrna.....	195	25.3	59.2	56.7	54.0	43.5	21.5	55.7	46.4	48.6	48.6	49.4	20.8	12	44.1	108.6
Club Mariout.....	261	13.0	32.4	66.7	37.5			48.2	50.6	61.8	58.6	63.2	30.2	10	46.2	102.4
Coast.....	626	13.3	66.3	60.0	47.3	47.9	24.3	64.8	54.1	63.4	36.5	62.5	35.2	12	48.0	118.2
Tennessee Winter.....	257		60.0						38.7	59.1	31.8	48.8	26.3	6	44.1	102.1
Maryland Winter.....	518		83.2	90.0	66.7									3	80.0	152.7
Wisconsin Winter.....	519		52.8	83.3										2	68.1	125.6
Nepal.....	22		41.2	53.3										2	47.3	87.3
Chevalier.....	278		44.8	60.0	32.3			57.7	45.7	46.3	28.9			7	45.1	95.3
Poda.....	652		43.6	60.0	45.0	31.1	29.3	64.6	55.5	59.2	60.4	52.6	32.8	11	48.6	118.2
Peru.....	653		68.0	60.0	51.0									3	59.7	113.9
Coast.....	691		94.4	50.0										2	72.2	133.2
Bavarian.....	159			33.3	45.0									2	39.1	77.1
Black Algerian.....	708			56.7	33.3									2	45.0	88.8
Abyssinia.....	361			60.0	64.5	68.7	25.0	47.4	25.3	30.0	24.0			8	42.5	105.5
Popeline.....	704			43.3	51.7									2	47.5	93.7
Princess.....	529			63.3	57.7									2	60.5	119.3
Hannchen.....	531			73.3	57.0		27.7	33.7	33.0	35.1	36.2	54.4		8	43.8	105.3
Orel.....	351			53.3	45.7									2	49.5	97.6
Chevalier.....	156			60.0	38.3									2	49.1	96.8
Hankow.....	197			26.7										1	26.7	51.1
Nepal.....	234			46.7				41.7						2	44.2	90.8
Abyssinia.....	355			36.7										1	36.7	70.2
Gatami.....	575			60.0										1	60.0	114.7
Golden Grain.....	588			60.0										1	60.0	114.7
Black Hull-less.....	596			50.0										1	50.0	95.6
Nesbian.....	647			36.7										1	36.7	70.2
Virginia Hooded.....	648			16.7										1	16.7	31.9
Hanse Hull-less.....	703			33.3										1	33.3	63.7
Black Russian.....	705			40.0				38.0						2	39.0	80.1
Turkestan.....	711			46.7										1	46.7	89.3
Black Hull-less.....	735			26.7										1	26.7	51.1
Nepal.....	2315					35.4	24.0							2	29.7	103.1
Abyssinian.....	938			41.5										1	41.5	111.9
Palestine.....	939			23.2										1	23.2	62.5
Svanhals.....	187			28.5	14.5									2	21.5	74.7
White Smyrna.....	910			36.4	18.7									2	27.5	95.5
Steigum.....	931			33.1	14.3	40.1								3	29.2	83.7
Goldroll.....	928			34.5	21.0	36.3								3	30.6	87.7
Italian.....	914			28.2	12.0									2	20.1	69.8
Proskowetz.....	893			41.1	19.5	50.9								3	37.2	106.6
Summit.....	929			39.3	14.0									2	26.7	92.7
Oderbrucker.....	940			36.4	18.0									2	27.2	94.4
Servian.....	915			26.5										1	26.5	71.4
Odessa.....	916			32.3	23.3	43.5	56.9	48.0	40.1	44.5				7	41.2	108.7
Do.....	934			35.4	33.3									2	34.3	119.1
Abyssinian.....	941			48.5	16.5									2	32.5	112.8
Hanna.....	942			32.3	17.0									2	24.7	85.8
Coast.....	2306			51.8										1	51.8	139.6
Do.....	2307			46.5										1	46.5	125.3
Abina.....	2308			49.7	17.5									2	33.6	116.7
Hanna.....	2309			22.5	21.5									2	22.0	76.4
Invincible.....	590			29.8	18.0	46.7								3	31.5	90.3
Peru.....	2311			25.5	25.5	59.0								3	36.7	105.2
Heil Hanna 2.....	678			31.6	19.2	58.3	40.8	44.9	34.9					6	38.3	104.1
Franconian.....	679			31.3	21.5	37.9								3	30.2	86.5
Heil Hanna 1.....	681			30.5	22.0									2	26.3	91.3
Coast.....	2312			42.9										1	42.9	115.6
Mährische.....	912					16.3	38.1							2	27.2	80.2
Luth.....	908					17.7	63.6							2	40.7	120.1
Abyssinian.....	2313					19.7								1	19.7	94.7
Manchuria.....	244					17.5								1	17.5	84.1
Hanna.....	2316					18.3								1	18.3	88.0
Abyssinian.....	2314					27.0	46.7							2	36.9	108.8
Hanna.....	416						43.1							1	43.1	94.5
Cortney.....	976						22.9							1	22.9	50.2
Hanchamont.....	1121						41.2							1	41.2	90.4
Sandrel.....	937						52.3							1	52.3	114.7
Abyssinian.....	950						36.5							1	36.5	80.0
Australian White.....	925						43.2							1	43.2	94.7
Abyssinian.....	943						51.8							1	51.8	113.6
Coast.....	276						53.9							1	53.9	118.2
Golden Melon.....	958						36.7							1	36.7	80.5

Of the varieties given a detailed comparison of 13 is shown in Table 77. The 2 North African barleys, Coast (C. I. No. 626) and Poda (C. I. No. 652), were the best of the 13. Poda is of the Peruvian type. The North African barleys as now grown in California contain two quite different types, although both are obviously North African in origin and character. The predominating type is a nodding barley with kernels which, though they are long, are shorter than those of most North African varieties. Barleys of the Poda, or Peruvian, type have more erect spikes and longer kernels. They are also more likely to have blue color in the endosperm.

Beldi Dwarf (C. I. No. 190), which is similar to the California Mariout (C. I. No. 1455), was third in point of yield at Chico. White Smyrna (C. I. No. 195), ranking fourth, gave relatively higher yields at Chico than at Davis. Club Mariout (C. I. No. 261) was fifth. This barley came from the same region in Egypt from which the California Mariout was imported. The former has given very good yields at Chico, and the quality of the grain when well grown is not surpassed by that of any barley in America. These varieties are discussed in detail in the section of this bulletin devoted to descriptions of the better varieties.

In the summarized digest of Table 77 a comparison of one of the Abyssinian barleys with 12 other varieties is shown. While it was superior only to Oderbrucker (C. I. No. 940) and Invincible (C. I. No. 590) its yield was more than 90 per cent of that of five other varieties. Few Abyssinian barleys have been imported into the United States. The varieties from Abyssinia grow well in California, and from their appearance it is probable that they would have value in hybridizing.

Tennessee Winter (C. I. No. 257) has been grown at Chico but has not proved equal to the barleys of North Africa and Asia Minor. It has been a very good variety, however. Chico unquestionably is in the district well suited to the production of barleys of the North African type. It is also possible that barleys from Asia Minor may be found which would be very high yielding.

MODESTO, CALIF.

The barleys grown at Modesto, Calif., from 1906 to 1909, inclusive, were tested in cooperation with the California Agricultural Experiment Station. The conditions under which these tests were made were not altogether favorable, and no definite conclusions can be drawn.

It is quite evident from Table 78 that White Smyrna (C. I. No. 195) and Coast (C. I. No. 626) were the best yielding sorts. In 1907 Coast gave the better yield, in 1908 White Smyrna exceeded all varieties grown, and in 1909 White Smyrna and Coast were almost equal. Club Mariout (C. I. No. 261) produced the highest yield of the spring varieties in 1909. In that year Tennessee Winter (C. I. No. 257) gave the unusually high yield of 67.6 bushels. The spring barleys were seeded later than the optimum date in the tests made.

TABLE 78.—*Annual acre yields of varieties of barley grown at Modesto, Calif., in two or more of the four years from 1906 to 1909, inclusive*

[Data obtained in cooperation with the California Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)				Years grown	Average yield (bus.)	Percentage of weighted mean
		1906	1907	1908	1909			
Beldi Dwarf.....	190	5.7	-----	39.7	-----	2	22.7	102.3
Black Smyrna.....	191	9.8	-----	12.4	-----	2	11.0	50.0
White Smyrna.....	195	10.9	-----	65.0	13.6	4	24.5	111.4
Telli.....	194	8.8	-----	12.2	13.6	3	11.5	49.1
Tennessee Winter.....	257	9.4	-----	-----	67.6	2	38.5	230.5
Club Mariout.....	261	8.4	-----	28.8	23.4	3	20.2	86.3
Coast.....	626	-----	12.2	55.0	13.0	3	26.7	95.0

DAVIS, CALIF.

Davis, Calif., is situated in the Sacramento Valley in a section of heavy barley production. The Sacramento and San Joaquin Valleys are the most important centers of production of barleys of the Coast type in the United States. Barley was brought into California with the early missions. Its production extended rapidly in the Sacramento and San Joaquin Valleys after the discovery of gold in 1848.

The University Farm at Davis has been an important station in the testing and production of varieties and is very favorably situated for the continuation of experiments in barley. Yields from 1908 to 1920 have been furnished by the California Agricultural Experiment Station. These are reported in Table 79. Three or four varieties which could not be definitely identified were not included in this table. Ten varieties are compared in detail in the summarized digest of Table 79. Of these, the two highest yielding were Coast (C. I. Nos. 2821 and 626). Coast (C. I. No. 2821) is a selection developed by the California Agricultural Experiment Station and distributed as California No. 4000. The name "Coast" is given here not with the intention of changing the varietal name as far as its use in California is concerned, but simply to identify its type more definitely in this bulletin. Coast (C. I. No. 626) is the ordinary commercial barley of California. California Mariout (C. I. No. 1455), which was third in point of yield, is a variety which has been distributed by the California station. The history of this and of California No. 4000 as well as the adaptation of these varieties is considered elsewhere in this bulletin. Of the 10 varieties compared, Beldl Giant (C. I. No. 2777) was fourth. All four of these varieties originated in North Africa.

Fifth in point of yield was White Smyrna (C. I. No. 195). This variety originated in Asia Minor under conditions somewhat similar to those found in North Africa. Hannchen was the highest yielding sort of any of those which do not trace back to either North Africa or Asia Minor. Hannchen (C. I. No. 531) produces very good results in California, but is obviously not so well adapted as the barleys of the North African group. Perhaps the most interesting variety from the standpoint of yield is Oderbrucker (C. I. No. 537). This barley was not grown many years, as it was obviously inferior. Although it is extremely well adapted to the northern Mississippi Valley, it is almost worthless in California. It was superior to none of the nine varieties with which it was compared, being inferior even to Nepal (C. I. No. 595), which is usually the poorest type grown. There is no question as to types of barleys best suited to the district around Davis. Barleys of North African origin are outstanding in their yielding capacity.

Recently two barleys have been distributed in California which have quickly attained a commercial status. These are Club Mariout (C. I. No. 261) and California Mariout (C. I. No. 1455). Both came originally from Egypt. The acreage of the former is increasing rapidly in the Sacramento and upper San Joaquin Valleys, where it is meeting with a ready market in the export trade. California Mariout has been found to be superior under low rainfall conditions in the San Joaquin Valley, although its acreage is not limited to this locality.

[illegible]

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 10 REPRESENTATIVE VARIETIES

[*Explanation.*—The asterisk (*) indicates that the two varieties were not grown in the same years]

Variety	C. I. No.	Data shown	Che- valier	Svan- hals	Hann- chen	Oder- brucker	Cali- fornia Marlout	Coast	Coast	Beldi Giant	White Smyrna	Nepal
Chevalier	290	{ Years comparable	---	99.1	9	4	6	8	5	6	9	7
		{ Percentage yield	---	99.1	97.1	170.0	83.5	78.1	74.5	81.7	86.8	151.4
Svanhals	187	{ Years comparable	9	---	11	4	8	10	6	5	11	7
		{ Percentage yield	101.0	---	95.5	167.9	82.8	78.1	74.1	82.8	88.3	148.7
Hannchen	531	{ Years comparable	9	---	---	4	8	10	6	8	11	7
		{ Percentage yield	103.0	104.7	---	178.6	85.8	81.6	73.7	85.8	92.5	147.9
Oderbrucker	537	{ Years comparable	4	4	4	---	1	3	(*)	1	3	3
		{ Percentage yield	98.8	99.6	56.0	---	38.5	48.3	---	48.0	55.9	85.9
California Marlout	1455	{ Years comparable	6	---	8	1	---	8	6	8	8	5
		{ Percentage yield	119.7	120.8	116.6	289.4	---	95.4	84.8	100.0	101.7	168.7
Coast	626	{ Years comparable	8	10	10	3	8	6	6	9	10	6
		{ Percentage yield	128.0	128.0	122.5	207.2	104.8	---	92.7	103.5	110.7	178.9
Do.	2821	{ Years comparable	5	6	6	(*)	6	6	4	6	6	4
		{ Percentage yield	134.1	134.9	132.1	---	117.9	107.8	---	109.4	119.2	209.3
Beldi Giant	2777	{ Years comparable	6	8	8	1	8	9	6	---	8	5
		{ Percentage yield	122.4	120.8	116.6	203.2	100.0	96.6	91.4	---	101.7	172.9
White Smyrna	195	{ Years comparable	9	11	11	4	8	10	6	8	---	7
		{ Percentage yield	115.2	113.2	108.1	178.9	98.4	90.3	83.9	98.4	---	175.1
Nepal	595	{ Years comparable	7	7	7	3	5	6	4	5	7	---
		{ Percentage yield	66.0	67.2	67.6	116.3	59.3	55.9	47.8	57.8	57.1	---

DATA FROM SOUTHERN STATIONS IN THE UNITED STATES

Two ecological regions have already been described. The first of these is the region with the growing and ripening seasons both humid. This extends from eastern Canada and the northern seaboard of the United States across the Northern States to the Great Plains and through the prairie Provinces of Canada. The second comprises the region where the rainfall is much less and the ripening season comes during a part of the year when the humidity is low. This region extends from the northern Great Plains west and south, including all the Mountain and Western States. The remaining district, a relatively unimportant one in barley production, includes the Southeastern States and extends westward, including Texas and Oklahoma. The summers here are too hot and humid for the growing of spring barley. The barley is fall sown. The varieties used are those more than usually winter hardy and adapted to humid conditions. As heretofore stated, the stations are arranged in order from east to west.

COLLEGE PARK, MD.

Barley varieties were grown at the Maryland Agricultural Experiment Station (at College Park) in the years 1908 to 1916, inclusive. The yields of the varieties shown in Table 80 were obtained in cooperation with the Maryland station. All of the 15 barleys grown were true winter sorts and were sown in the fall. Only four of these were grown for a longer period than three years. Maryland Winter (C. I. No. 518) was obviously the best of the four. Mammoth Winter (C. I. No. 410) and Tennessee Winter (C. I. No. 257) produced nearly the same average yields for the comparable years, 1908 to 1914, inclusive. Winter Six-Rowed (C. I. No. 328) was surpassed by the three mentioned above. All four varieties are quite similar and belong to the Tennessee Winter group.

Arlington Awnless (C. I. No. 702), which was grown from 1913 to 1915, inclusive, was much inferior in yield to the varieties of the Tennessee Winter group. While Arlington Awnless is a very desirable type of barley, it shatters quite easily, thus causing the loss of much grain. The only 2-rowed barley grown in this experiment was Nesbian (C. I. No. 647), which was tested in 1915. It produced a very low yield, as did Osaka (C. I. No. 2798) and Virginia Hooded (C. I. No. 648), which were also grown in that year. The results at College Park indicate that the barleys of the Tennessee Winter type are the ones best adapted to the growing conditions of this section of Maryland.

TABLE 80.—*Annual acre yields of varieties of barley grown at the Maryland Agricultural Experiment Station (at College Park) in part or all of the nine years from 1908 to 1916, inclusive*

[Data obtained in cooperation with the Maryland Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)									Years grown	Average yield (bus.)	Percentage of weighted mean
		1908	1909	1910	1911	1912	1913	1914	1915	1916			
Maryland Winter.....	518	32.9	36.7	24.7	31.9	42.0	32.1	37.5	22.7	48.2	9	34.3	121.6
Mammoth Winter.....	410	23.5	34.7	20.1	33.0	47.4	35.2	39.6	-----	-----	7	33.4	110.2
Tennessee Winter.....	257	39.8	29.8	20.7	32.8	42.6	31.8	35.8	15.7	-----	8	31.1	111.9
Winter Six-Rowed.....	328	24.7	27.6	22.8	36.4	33.9	24.9	35.4	-----	-----	7	29.4	97.0
Arlington Awnless.....	702	-----	-----	-----	-----	-----	16.4	17.6	15.9	-----	3	16.6	71.2
Squarehead Winter.....	252	29.7	35.2	-----	-----	-----	-----	-----	-----	-----	2	32.5	111.7
Tenarab.....	662	24.1	30.4	-----	-----	-----	-----	-----	-----	-----	2	27.3	93.8
Dutrow Winter.....	2797	-----	32.8	22.6	-----	-----	-----	-----	-----	-----	2	27.7	98.2
Han River.....	205	28.5	-----	-----	-----	-----	-----	-----	-----	-----	1	28.5	106.3
Black Winter.....	337	25.4	-----	-----	-----	-----	-----	-----	-----	-----	1	25.4	94.8
Albaette.....	199	21.6	-----	-----	-----	-----	-----	-----	-----	-----	1	21.6	80.6
Greece.....	221	17.8	-----	-----	-----	-----	-----	-----	-----	-----	1	17.8	66.4
Nesbian.....	647	-----	-----	-----	-----	-----	-----	6.9	-----	-----	1	6.9	61.6
Osaka.....	2798	-----	-----	-----	-----	-----	-----	3.5	-----	-----	1	3.5	31.3
Virginia Hooded.....	648	-----	-----	-----	-----	-----	-----	2.3	-----	-----	1	2.3	20.5

ROSSLYN, VA.

At Arlington Experiment Farm 72 varieties and strains of barley have been tested in periods of varying length during the years 1909 to 1921, inclusive. This farm, which is at Rosslyn, Va., across the Potomac River from Washington, D. C., is owned and operated by the United States Department of Agriculture. Table 81 shows the yields of the 72 varieties, all of which were sown in the fall. The 11 prominent winter and spring sorts are compared in the summarized digest of the table. Orel (C. I. No. 351), a 2-rowed variety, perhaps is one of the best of those compared. Although it was surpassed in yield by both Chevalier (C. I. No. 156) and Hanna (C. I. No. 287) in 1911 and 1912, the years in which they were comparable, neither of these varieties produced high yields. Orel entirely winterkilled in 1912, making its average yield for the two years very low. Han River (C. I. No. 2163), a 6-rowed variety, also produced very high yields. It was not grown the same years as Chevalier and Hanna, so no comparison can be made with these varieties.

Wisconsin Winter (C. I. No. 2159), which was equal to Orel and Han River in point of yield, seems quite winter hardy at Arlington. It makes a fall growth heavier than that of the Tennessee Winter, so that the chances of winterkilling would be greater at most stations. In recent years Han River has been surpassed by both Wisconsin Winter and Tennessee Winter. Following Wisconsin Winter is Scottish Pearl (C. I. No. 277), a barley of the Tennessee Winter group, and Pidor (C. I. No. 901), a hybrid resulting from the cross Tennessee Winter $\times$ Hankow. Although Tennessee Winter (C. I. No. 257) was surpassed by the varieties mentioned above for the years in which it was comparable, it averaged 30.3 bushels for the 13 years of the experiments, as compared with 31.2 bushels for Wisconsin Winter (C. I. No. 2159) for this same period.

The three spring varieties compared in Table 81—Chevalier (C. I. No. 156), Hanna (C. I. No. 287), and Svanhals (C. I. No. 187)—all produced very low yields. Neither the spring nor the hull-less varieties have given satisfactory results at Arlington. When the spring varieties are sown in the fall their growth is so rapid that unless the winter months are very mild they will not survive. When sown in the spring they do not reach full development because of the early advent of high temperatures.

Orel produced unusually high yields during the four years from 1918 to 1921, inclusive (Table 81). Texas Winter (C. I. No. 554), a locally adapted variety of the Tennessee Winter group, produced an average yield of 118 per cent of the weighted mean for the nine years it was tested. Two hybrids of a Tennessee Winter cross, Cusado (C. I. No. 895) and Tenkow (C. I. No. 646), also produced fair yields.

From the results at Arlington Experiment Farm it is evident that several types, such as Orel, Wisconsin Winter, and varieties of the Tennessee Winter group, are successful here.

Wisconsin Winter and Orel appear to be better varieties than Tennessee Winter. The value of Tennessee Winter may not be fully indicated by the yields at Arlington farm, which is protected by the hills and is doubtless warmer than most of the grain-growing sections of Maryland and Virginia. Tennessee Winter is more winter hardy than the varieties which gave higher yields at Arlington. In the Piedmont area it probably would be more dependable.

Orel obviously has unusual qualities. It yields well. It is winter hardy and at least at Arlington it is less affected by smut than Tennessee Winter. As it succeeds as a spring variety at Akron, Colo., its winter dormancy is likely to be less than that of Tennessee Winter. In the warmer sections of the South this lack of dormancy may prove to be an asset. The experimental results with barley in the South are too meager as yet to form the basis of an intelligent opinion, but there are indications that on the Coastal Plains the highest yields will be obtained from varieties which take advantage of the first growing weather of the spring. In the latitude of Washington, D. C., Tennessee Winter does this, but farther south on the Coastal Plains its dormancy may be a handicap.

Another desirable quality in winter barleys either in the South or in the latitude of Washington, D. C., is the habit of early growth in the spring, which is of great advantage in checking the development of weeds. Wisconsin Winter is one of the better varieties in this respect.

TABLE 81.—Annual acre yields of varieties of barley grown at the Arlington Experiment Farm, Rosslyn, Va., in part or all of the 13 years from 1909 to 1921, inclusive—Continued

SUMMARIZED DIGEST, SHOWING THE RELATIVE PERFORMANCE OF 11 REPRESENTATIVE VARIETIES

[Explanation.—The asterisk (*) indicates that the two varieties to which it relates were not grown in the same years]

Variety	C. I. No.	Data shown	Varieties and percentages										
			Cheva-lier	Hanna	Svan-hals	Tennes-see Winter	Han-kliver	Scottish Pearl	Wisconsin Winter	Hansoe Hull-less	Nakano Wase	Orel	Pidor
Chevalier	156	{ Years comparable { Percentage yield.....	-----	4 101.7	4 104.6	4 76.8	(*)	1 40.6	4 84.6	3 84.1	(*)	2 329.1	(*)
Hanna	287	{ Years comparable { Percentage yield.....	98.4	4 102.9	4 102.9	4 75.6	(*)	1 52.2	4 83.3	3 65.1	(*)	2 198.5	(*)
Svanhals	187	{ Years comparable { Percentage yield.....	4	4 97.2	-----	12 62.5	7	9 61.5	12 61.0	11 80.2	6	0 50.5	8
Tennessee Winter	257	{ Years comparable { Percentage yield.....	95.6	4 132.3	12 159.9	-----	49.8	3 91.5	13 97.1	11 132.2	6	0 97.1	8
Han River	2163	{ Years comparable { Percentage yield.....	130.2	(*)	7	7	90.5	97.5	7	7	128.8	97.1	7
Scottish Pearl	277	{ Years comparable { Percentage yield.....	-----	1 200.9	0 200.9	9 110.5	7	115.0	103.7	172.6	137.8	70.7	109.9
Wisconsin Winter	2159	{ Years comparable { Percentage yield.....	246.6	191.5	162.7	102.5	87.0	9	93.4	148.0	131.4	76.2	96.0
Hansoe Hull-less	703	{ Years comparable { Percentage yield.....	118.1	120.1	103.8	103.0	96.4	107.0	-----	11	6	100.7	105.5
Nakano Wase	2166	{ Years comparable { Percentage yield.....	118.9	153.7	124.7	75.6	57.9	9	71.7	-----	6	6	68.8
Orel	351	{ Years comparable { Percentage yield.....	-----	2 50.4	6 197.8	6 103.0	72.6	76.1	6	6	-----	43.4	72.1
Pidor	901	{ Years comparable { Percentage yield.....	30.4	(*)	8 173.1	8 101.2	7 91.0	8 104.1	8 99.3	8 142.0	6 230.3	4 79.3	126.0

BLACKSBURG, VA.

The yields of barley varieties grown at Blacksburg, Va. (Table 82), were obtained through the courtesy of the Virginia Agricultural Experiment Station. During the years from 1914 to 1921, inclusive, 20 winter barleys were tested, and 17 spring varieties from 1917 to 1919, inclusive. In 1919 the spring varieties were destroyed by extremely wet weather.

All of the winter varieties grown in 1914 were selections made at the Virginia station. Five of these were grown during the entire eight years of the experiments, all of them producing comparatively high average yields with the exception of C. I. No. 3153. Because of crop failure no yields were recorded for the years 1917 and 1921. In 1918 eight winter barleys were introduced and continued through 1921. Union Winter (C. I. No. 583), a barley of the Tennessee Winter group, produced the highest average yield of all varieties for these four years. C. I. No. 3153 was second in point of yield for this period. Its average for the eight years tested was quite low. Following C. I. No. 3153 are two other selections, C. I. Nos. 3164 and 3163. Both of these are selections from C. I. No. 3154, the first being 6-rowed and the second 2-rowed. It is to be regretted that there is no record of the source of the stock from which the early selections were made. Both Arlington Awnless (C. I. No. 702) and Tennessee Winter (C. I. No. 257) produced comparatively high yields during this 4-year period. Arlington Awnless is not usually a high-yielding sort and it shatters very badly.

TABLE 82.—*Annual acre yields of varieties of barley grown at the Virginia Agricultural Experiment Station (at Blacksburg) in part or all of the eight years from 1914 to 1921, inclusive*

[Data obtained through the courtesy of the Virginia Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)								Years grown	Average yield (bus.)	Percentage of weighted mean
		1914	1915	1916	1917	1918	1919	1920	1921			
Winter barley:												
Virginia Selection	3148	12.5	25.4	0	—	—	—	—	—	3	12.6	66.0
Do.	3149	20.3	44.0	26.6	0	10.3	17.5	34.4	0	8	19.1	134.5
Do.	3150	9.9	57.0	18.1	0	11.2	20.3	32.7	0	8	18.7	131.7
Do.	3151	6.8	39.4	30.2	0	9.6	17.5	25.8	0	8	16.2	114.1
Do.	3152	13.5	28.1	0	—	—	—	—	—	3	13.9	72.8
Do.	3153	3.6	7.2	10.8	0	13.4	21.7	36.1	0	8	11.6	81.7
Do.	3154	9.4	40.5	32.7	0	—	—	—	—	4	20.7	120.3
Do.	3155	8.9	21.6	9.6	0	—	—	—	—	4	10.0	58.1
Do.	3156	12.5	31.6	0	—	—	—	—	—	3	14.7	77.0
Do.	3157	6.8	44.0	45.9	0	10.3	16.5	30.9	0	8	19.3	135.9
Hull-less	3158	5.2	26.6	0	—	—	—	—	—	3	10.6	55.5
Virginia Selection	3159	1.3	17.8	—	—	—	—	—	—	2	9.5	46.1
Arlington Awnless	702	—	—	—	—	9.7	22.3	29.2	0	4	15.3	127.5
Winter	3160	—	—	—	—	7.2	20.3	27.5	0	4	13.7	114.2
Tennessee Winter	257	—	—	—	—	11.3	19.9	27.5	0	4	14.7	122.5
Union Winter	583	—	—	—	—	13.2	21.7	37.8	0	4	18.2	151.7
Winter Beardless	3161	—	—	—	—	16.2	11.0	15.5	0	4	10.7	89.2
Tennessee	3162	—	—	—	—	6.9	17.9	8.6	0	4	8.3	69.2
Virginia Selection	3163	—	—	—	—	14.9	25.1	24.1	0	4	16.0	133.3
Do.	3164	—	—	—	—	17.2	26.5	20.6	0	4	16.1	134.2
Spring barley:												
Manchuria	3166	—	—	—	19.2	15.5	0	—	—	3	11.6	94.3
Gatami	575	—	—	—	20.3	14.5	0	—	—	3	11.6	94.3
Eagle	913	—	—	—	20.3	16.3	0	—	—	3	12.2	99.2
Black Hull-less	596	—	—	—	15.3	—	—	—	—	1	15.3	105.5
Featherston	911	—	—	—	26.5	22.2	0	—	—	3	16.2	131.7
Chevalier	3165	—	—	—	20.6	11.0	0	—	—	3	10.5	85.4
Black Egyptian	1246	—	—	—	17.2	10.7	0	—	—	3	9.3	75.6
Luth	908	—	—	—	22.9	17.1	0	—	—	3	13.3	108.1
Proskowetz	893	—	—	—	19.6	14.6	0	—	—	3	11.4	92.7
Reed Triumph	839	—	—	—	18.9	14.1	0	—	—	3	11.0	89.4
Hanna	319	—	—	—	27.1	10.6	0	—	—	3	12.6	102.4
Svanhals	187	—	—	—	15.9	—	—	—	—	1	15.9	109.7
Nepal	595	—	—	—	19.7	16.2	0	—	—	3	12.0	97.6
Servian	915	—	—	—	26.0	9.0	0	—	—	3	11.7	95.1
Odessa	916	—	—	—	25.6	7.7	0	—	—	3	11.1	90.2
Manchuria	3167	—	—	—	20.8	20.4	0	—	—	3	13.7	111.4
Two-Row Hull-less	3180	—	—	—	12.9	—	—	—	—	1	12.9	89.0

Of the spring varieties tested, Featherston (C. I. No. 911) was the best. Manchuria (C. I. No. 3167) and Luth (C. I. No. 908) were second and third, respectively. These three belong to the Manchuria group. Hanna (C. I. No. 319) was the only 2-rowed variety that compared at all favorably with the 6-rowed sorts. The barleys of the Manchuria type are undoubtedly the best adapted spring varieties for this section. Although fairly high yields were obtained from some of these barleys, they were usually much diseased and had a great tendency to lodge. The use of spring barley is not generally advocated in this State. It is rather difficult to draw any definite conclusions from the table of yields as to the winter types best adapted to this section because of lack of information regarding the identity of the winter selections. In recent years, however, Union Winter has proved a superior sort. As Union Winter and Tennessee Winter are the most frost resistant of the winter varieties, they are likely to be the most successful in this humid region where heavy winterkilling is likely to occur every three or four years.

CLEMSON, S. C.

The annual yields of barley grown in 1912, 1914, and 1920 at Clemson, S. C. (Table 83), were furnished through the courtesy of the South Carolina Agricultural Experiment Station. Barley is a crop of little or no importance in South Carolina. All of the varieties tested were winter barleys and were fall sown. Tennessee Winter (C. I. No. 257) is the only variety tested for more than one year. Of the eight varieties grown in 1912 Tennessee Winter was outstanding, with a yield of 40.3 bushels. Alaska (C. I. No. 534) was second. There is little information about this variety, as it has not been generally distributed to the various stations. Both Han River (C. I. No. 206) and Greece (C. I. No. 221) produced yields of 32 bushels. The other varieties grown this year did not seem promising, yielding less than 30 bushels per acre. In 1914 four varieties were grown, but since only Tennessee Winter can be identified it is the only one included in the table. It yielded 21.1 bushels, which did not compare favorably with its yield in 1912. Two varieties, Virginia Hooded (C. I. No. 648) and Arlington Awnless (C. I. No. 702), were grown in 1920. Virginia Hooded produced a yield of 9.1 bushels, but Arlington Awnless failed entirely.

While the varieties in this experiment were not tested for a sufficient period to show their relative value, the results in 1914 and in 1920 were not promising. In 1912, on the other hand, they were quite good. It is the belief of the State officials that barley may be grown with success, but owing to the competition with other crops it is unlikely to be seeded in any considerable acreage.

TABLE 83.—*Annual yields of varieties of barley grown at the South Carolina Agricultural Experiment Station (at Clemson) in 1912, 1914, and 1920*

[Data furnished through the courtesy of the South Carolina Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)		
		1912	1914	1920
Han River.....	206	32.0		
Greece.....	221	32.0		
Besthorn Giant.....	414	27.9		
Tennessee Winter.....	257	40.3	21.1	
Winter Club.....	488	29.9		
Wisconsin Winter.....	519	25.5		
Alaska.....	534	38.2		
Texas Winter.....	554	28.9		
Virginia Hooded.....	648			9.1
Arlington Awnless.....	702			0

ATHENS, GA.

The yields of barley varieties shown in Table 84 were obtained in cooperation with the Georgia State College of Agriculture. These varieties were grown during the years 1914 to 1921, inclusive. Unfortunately the barns were burned in 1921, destroying the entire crop for that year. The college farm at Athens is one of the few points in the South where barleys of the humid-winter type have been tested in sufficient numbers and for an adequate period of years to make definite

conclusions possible. Nineteen varieties were tested in two or more of the seven years. Six were grown for the full period.

During the earlier years of the experiments five spring barleys, Bavarian (C. I. No. 159), Chevalier (C. I. No. 156), Hannchen (C. I. No. 531), Hanna (C. I. No. 287), and Princess (C. I. No. 529), were tested along with the winter varieties. Both Bavarian and Hannchen were continued for six years, but it is obvious that none of the spring varieties can compete successfully with the winter sorts. While Hannchen produced the third highest yield of all varieties grown in 1920, it was surpassed by a number of the winter sorts in all the other years.

Of the varieties grown for the seven years, Turkestan (C. I. No. 711) produced the highest average yield, 44.7 bushels. Tennessee Winter (C. I. No. 257) was almost its equal, with an average yield of 44.1 bushels. Wisconsin Winter (C. I. No. 519), which was third, was also a superior variety at Arlington Experiment Farm, near Washington, D. C. Argentine (C. I. No. 223), Mammoth Winter (C. I. No. 220), and Greece (C. I. No. 221) follow in the order named. None of the other winter varieties seems at all promising. Orel (C. I. No. 351), which has produced high yields at several of the stations as a spring variety and at others as a winter sort, does not appear to be hardy enough to withstand the winters here. Because of the absence of rough awns, Virginia Hooded (C. I. No. 648) and Arlington Awnless (C. I. No. 702) are desirable sorts, but they were very poor yielders, as was Tanbash (C. I. No. 578), a hull-less variety. These varieties have not been highly productive at any of the stations where they have been tested. Black Russian (C. I. No. 705), which was grown during the years 1916 to 1920, inclusive, produced only a fair average yield.

The results show conclusively that spring barleys are not at all dependable when fall sown at Athens. Turkestan, Tennessee Winter, and Wisconsin Winter were all satisfactory varieties well adapted to conditions here. The small acreage of barley grown in Georgia is largely of the Tennessee Winter type.

TABLE 84.—*Annual acre yields of varieties of barley grown at the Georgia State College (at Athens) in some or all of the seven years from 1914 to 1920, inclusive*

[Data obtained in cooperation with the Georgia State College of Agriculture. Yields in 1921 were destroyed by fire]

Variety	C. I. No.	Acre yields (bushels)							Years grown	Average yield (bus.)	Percentage of weighted mean
		1914	1915	1916	1917	1918	1919	1920			
Turkestan.....	711	43.0	46.6	54.2	14.1	55.2	48.9	51.2	7	44.7	154.7
Tennessee Winter.....	257	40.6	43.8	61.5	8.6	70.8	43.2	40.4	7	44.1	152.6
Wisconsin Winter.....	519	35.1	44.0	57.9	11.7	56.3	42.1	37.7	7	40.7	140.8
Mammoth Winter.....	220	33.4	42.7	56.3	6.8	7.3	44.7	50.0	7	34.5	119.4
Argentine.....	223	33.0	41.9	55.3	0	39.8	50.0	44.5	7	37.8	130.8
Greece.....	221	30.2	39.8	39.6	4.2	20.0	39.0	44.4	7	31.0	107.3
Maryland Winter.....	518	23.9	34.1	-----	-----	-----	-----	-----	2	29.0	93.9
Virginia Hooded.....	648	21.9	25.5	32.8	0	-----	-----	-----	4	20.1	74.7
Arlington Awnless.....	702	-----	22.9	37.0	0	4.2	-----	26.3	5	18.1	63.9
Nesbian.....	647	31.5	32.7	36.0	0	0	-----	-----	5	20.0	76.6
Orel.....	351	20.8	33.5	26.6	0	11.9	-----	41.1	6	22.3	79.6
Bavarian.....	159	16.9	36.7	35.4	0	4.7	29.3	-----	6	20.5	75.1
Chevalier.....	156	16.6	35.7	-----	-----	-----	-----	-----	2	26.1	84.5
Hannchen.....	531	21.3	29.7	27.1	0	6.3	-----	7.7	6	22.0	78.6
Hanna.....	287	21.3	28.9	-----	-----	-----	-----	-----	2	25.1	81.2
Princess.....	529	11.5	30.0	-----	-----	-----	-----	-----	2	20.7	67.0
Tanbash.....	578	26.8	22.9	19.1	0	1.0	-----	-----	5	14.0	53.6
Black Russian.....	705	-----	-----	48.0	0	30.7	18.7	43.7	5	28.2	101.1
Nakano Wase (See Pl. III, fig. 1, D).....	754	-----	-----	46.4	0	2.1	-----	-----	3	16.2	72.0

ASHBURN, GA.

The results at the substation at Ashburn, Ga., were obtained in cooperation with the Georgia State College of Agriculture. The yields of the barley varieties grown in the years 1915 and 1916 are shown in Table 85. With the exception of Nesbian (C. I. No. 647), all of the barleys were grown both years. Greece (C. I. No. 221) produced the highest average yield, with Orel (C. I. No. 351) second and Tennessee Winter (C. I. No. 257) third. Following Tennessee Winter were Argentine (C. I. No. 223) and Turkestan (C. I. No. 711). Arlington

Awnless (C. I. No. 702) and Tanbash (C. I. No. 578), a hull-less variety, produced the lowest yields in both years. Nesbian (C. I. No. 647), which was grown in 1916, was the second in rank that year. While a test in only two years does not show definitely the value of any particular variety, it does give a general idea of what varieties might be expected to yield well in this section.

TABLE 85.—*Annual acre yields of varieties of barley grown at the Ashburn (Ga.) substation in 1915 and 1916*

[Data obtained in cooperation with the Georgia State College of Agriculture]

Variety	C. I. No.	Acre yields (bushels)		Years grown.	Average yield (bus.)	Percentage of weighted mean
		1915	1916			
Greece.....	221	19.3	14.5	2	16.9	128.0
Argentine.....	223	18.7	9.3	2	14.0	106.1
Tennessee Winter.....	257	15.6	14.0	2	14.8	112.1
Orel.....	351	15.6	16.1	2	15.9	120.5
Turkestan.....	711	13.0	12.0	2	12.5	94.7
Arlington Awnless.....	702	9.4	8.8	2	9.1	68.9
Tanbash.....	578	8.3	8.0	2	8.1	61.4
Nesbian.....	647		15.6	1	15.6	126.8

QUITMAN, GA.

The yields of barley varieties grown at the Quitman substation during the year 1915 are found in Table 86. These yields were obtained in cooperation with the Georgia State College of Agriculture. Of the seven winter barleys grown in this year, Greece (C. I. No. 221) was the highest yielder, with Mammoth Winter (C. I. No. 220) and Argentine (C. I. No. 223) following. Orel (C. I. No. 351) and Tennessee Winter (C. I. No. 257) were fourth and fifth in point of yield. Tennessee Winter is usually relatively higher in rank at stations where humid-winter barleys are grown than it is here. A test of only one year is not sufficient to show which types are superior, but it is likely that the less dominant winter types are better adapted this far south.

TABLE 86.—*Annual acre yields of varieties of barley grown at the Quitman (Ga.) substation in 1915*

[Data obtained in cooperation with the Georgia State College of Agriculture]

Variety	C. I. No.	Acre yields (bushels)	Variety	C. I. No.	Acre yields (bushels)
Greece.....	221	8.1	Tennessee Winter.....	257	4.4
Mammoth Winter.....	220	7.8	Turkestan.....	711	4.3
Argentine.....	223	7.3	Wisconsin Winter.....	519	4.2
Orel.....	351	5.5			

MORGANTOWN, W. VA.

The yields of the barley varieties grown in 1921 at Morgantown were furnished through the courtesy of the West Virginia Agricultural Experiment Station. As very little barley is grown in this State, the experiments with this cereal have been very limited. The four varieties tested here were grown in rod rows replicated seven times. Manchuria (C. I. No. 244) produced the highest yield, 34.4 bushels. A 2-rowed barley, Chevalier (C. I. No. 278), was second in rank, with a yield of 32.7 bushels. Neither of the hull-less sorts, Himalaya (C. I. No. 620) and Nepal (C. I. No. 595), compared favorably with the two hulled varieties, Himalaya producing a yield of only 18 bushels and Nepal 13.6 bushels.

No data are available on which to base an opinion as to the value of winter varieties. At the higher elevations the spring varieties would likely be the better sorts, but at the lower elevations they would probably become badly diseased. This was the case in Kentucky and Virginia where spring barleys were tested. There the winter sorts proved to be superior.

LEXINGTON, KY.

The results at the Kentucky Agricultural Experiment Station at Lexington are shown in Table 87. These data, furnished through the courtesy of that station, include yields for seven of the years from 1910 to 1917, inclusive. In 1910 both winter and spring varieties were tested. The spring-sown barleys produced yields much higher than those sown in the fall, 1910 being one of the few favorable years for spring-sown varieties in this section. In 1911 the spring-sown varieties, with the exception of Franconian (C. I. No. 679), were grown in rod rows instead of in plots. The straw was unusually short and the yields very low. Franconian, which produced the high yield of 85.5 bushels in 1910, was again grown in 1911 and 1912, but was quite inferior in both these years. No further tests were made with spring-sown barleys.

It was soon found that the spring varieties when fall sown could not survive the winters. In 1910 both Hanna (C. I. No. 287) and Horsford (C. I. No. 610) were entirely winterkilled, and Odessa (C. I. No. 182) and Telli (C. I. No. 194) were excelled by the winter sorts. None of the spring barleys survived in 1911 and 1912.

Mammoth Winter (C. I. No. 220) and Tennessee Winter (C. I. No. 257) are the only winter varieties so far tested that have produced uniformly good yields. These varieties are almost identical. In 1912 a number of the winter barleys winterkilled, and none of these was given further trial. Beardless 20 (C. I. No. 2745), a hybrid variety, produced a yield of 40.8 bushels in 1917, which was less than that of Tennessee Winter.

Barley is a crop of minor importance in Kentucky. The spring varieties are very unreliable. Tennessee Winter and Mammoth Winter seem to be the most productive of the winter sorts, and what little barley is grown is of this type.

TABLE 87.—*Annual acre yields of varieties of barley grown at the Kentucky Agricultural Experiment Station (at Lexington) in one or more of the eight years from 1910 to 1917, inclusive*

[Data obtained through the courtesy of the Kentucky Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)						Years grown	Average yield (bus.)	Percentage of weighted mean
		1910	1911	1912	1913	1914	1915			
Winter barley:										
Odessa	182	16.3	—	—	—	—	—	1	16.3	50.0
Telli	194	20.4	—	—	—	—	—	1	20.4	62.6
Mammoth Winter	220	40.0	25.0	31.0	69.0	28.7	51.2	6	40.8	176.6
Tennessee Winter	257	39.0	24.5	29.0	65.5	23.3	50.5	7	39.8	167.2
Hanna	287	0	—	—	—	—	—	1	0	—
Maryland Winter	518	33.0	14.2	4.5	0	—	—	4	12.9	58.1
Texas Winter	554	42.5	17.1	11.0	52.0	13.3	—	5	27.2	123.1
Horsford	610	0	—	—	—	—	—	1	0	—
Wisconsin Winter	519	30.5	22.9	26.7	49.0	—	—	4	32.3	145.5
Svanhals	187	0	—	—	—	—	—	1	0	—
Princess	529	0	0	—	—	—	—	2	0	—
Large-Grained Winter	408	—	22.8	30.0	47.5	16.4	—	4	29.2	170.8
Nesbian	647	0	0	—	—	—	—	2	0	—
Arlington Awnless	702	0	0	—	—	—	—	2	0	—
Hanse Hull-less	703	—	—	—	—	—	—	1	0	—
Chevalier	156	—	—	0	—	—	—	1	0	—
Virginia Hooded	648	—	—	0	—	—	—	1	0	—
Orel	351	—	—	0	—	—	—	1	0	—
Bavarian	159	—	—	0	—	—	—	1	0	—
Pontius	731	—	—	0	—	—	—	1	0	—
Turkestan	711	—	—	0	—	—	—	1	0	—
Beardless 20	2745	—	—	—	—	—	40.8	1	40.8	93.2
Spring barley:										
Heil Hanna 3	682	40.5	—	—	—	—	—	1	40.5	124.2
Barbary	695	19.0	—	—	—	—	—	1	19.0	58.3
Kashmir	685	44.5	—	—	—	—	—	1	44.5	136.5
White Smyrna	655	47.5	—	—	—	—	—	1	47.5	145.7
Chile Common	663	28.5	—	—	—	—	—	1	28.5	87.4
Abyssinia	669	44.5	—	—	—	—	—	1	44.5	136.5
Do	672	25.5	—	—	—	—	—	1	25.5	78.2
Franconian	679	85.5	17.0	9.0	—	—	—	3	37.2	195.8
Heil Hanna 1	681	30.0	—	—	—	—	—	1	30.0	92.0

KNOXVILLE, TENN.

The annual yields of barley in varietal experiments at the Tennessee Agricultural Experiment Station, Knoxville, furnished through the courtesy of this station, are presented in Table 88. While the plat experiments have not been so extensive here as at some of the other southern stations, much work has been done in breeding and selecting winter varieties for general cultivation in the warm and humid regions of the South. Tennessee undoubtedly has the largest winter-barley acreage of any humid State. The parent stock of Tennessee Winter (C. I. No. 257) has been the common barley in this region. This variety, as improved by mass selection at this station, was first distributed about 1900 as "Tennessee Winter." Several hooded hybrids which seem promising as winter barleys have been developed and distributed to other stations and to farmers.

Table 88 shows that Union Winter (C. I. No. 583), a barley of the Tennessee Winter group, is the only variety grown during the full period of the experiment. Tennessee Winter (C. I. No. 257), however, was grown for 11 of the 14 years, so that a fair comparison of the yields of the two varieties can be made. Union Winter undoubtedly is the higher yielding of the two at this station for this period. It produced an average acre yield of 42.8 bushels, as compared with 39.9 bushels for Tennessee Winter. In general appearance these varieties are quite similar, although Union Winter is a few inches shorter and several days earlier than Tennessee Winter.

Four varieties were introduced in 1908. Of these, Tenarab (C. I. No. 662) was the only one that was further tested, being grown in 1910 and 1911. In all of the three years it was inferior to both Union Winter and Tennessee Winter. Maryland Winter (C. I. No. 518) yielded 65.2 bushels in 1908, which was the highest that year. Bestehorn Giant (C. I. No. 414) and Black Winter (C. I. No. 337) were both quite inferior to the other four varieties. Beardless 20 (C. I. No. 2745), grown in 1913, 1914, 1916, and 1920, and Beardless 6 (C. I. No. 2746), grown in 1920 and 1921, are hooded selections made from a number of hybrids between Tennessee Winter $\times$ Horsford and Union Winter $\times$ Horsford. According to C. A. Mooers, of this station, Beardless 20 has been distributed to some of the farmers in the vicinity and is being more extensively cultivated. On poor or average land it produces nearly as much grain as Union Winter, with a larger yield of hay of superior quality. It does not stand wet-land conditions well, however. Beardless 6, although inferior to Beardless 20 in 1920, according to Professor Mooers, is "more robust and has greater capacity to produce both grain and straw. It also stands wet-land conditions well." Neither of these varieties was the equal of Union Winter or Tennessee Winter at this station in the years when comparable yields were obtained.

TABLE 88.—*Annual acre yields of varieties of barley grown at the Tennessee Agricultural Experiment Station (at Knoxville) in one or more of the 22 years from 1900 to 1921, inclusive*

[Data obtained through the courtesy of the Tennessee Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)															Years grown	Aver- age yield (bus.)	Per- centage of weight- ed mean
		1900	1901	1902	1903	1904	1905	1910	1911	1912	1913	1914	1916	1920	1921				
Tennessee Winter.....	257	58.9	30.0	54.5	21.9	28.1	63.3	33.9	44.5	30.2	-----	-----	29.5	44.2	-----	11	39.9	93.5	
Union Winter.....	583	71.1	27.1	54.2	41.7	35.4	57.5	34.6	44.5	27.6	43.7	54.2	33.0	50.9	23.7	14	42.8	106.5	
Tenarab.....	662	-----	-----	-----	-----	-----	53.2	32.8	39.1	-----	-----	-----	-----	-----	-----	3	41.7	96.7	
Bestehorn Giant.....	414	-----	-----	-----	-----	29.0	-----	-----	-----	-----	-----	-----	-----	-----	-----	1	29.0	60.3	
Maryland Winter.....	518	-----	-----	-----	-----	65.2	-----	-----	-----	-----	-----	-----	-----	-----	-----	1	65.2	135.6	
Black Winter.....	337	-----	-----	-----	-----	20.2	-----	-----	-----	-----	-----	-----	-----	-----	-----	1	20.2	42.0	
Beardless 20.....	2745	-----	-----	-----	-----	-----	-----	-----	-----	-----	37.5	49.0	27.5	37.4	-----	4	37.9	94.3	
Beardless 6.....	2746	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	34.9	21.9	-----	2	28.4	80.0	

It is evident from these results and from the types in general cultivation that the Tennessee Winter barleys are well adapted to growing conditions in Tennessee. While the hooded hybrids developed here are recommended especially for hay

and on poor lands, none has yet been found that will produce as much grain on fertile soil as varieties of the Tennessee Winter group.

JACKSON, TENN.

Two varieties of barley, Union Winter (C. I. No. 583) and Beardless 20 (C. I. No. 2745), were tested at the Jackson substation during the years 1915, 1916, 1917, 1919, 1920, and 1921. The yields of these varieties (Table 89) are used through the courtesy of the Tennessee Agricultural Experiment Station. Union Winter produced an average yield of 19.5 bushels for the six years, as compared with 16.5 bushels for Beardless 20. At Knoxville Union Winter was superior to all other varieties tested. It is a barley of the Tennessee Winter group which seems to be particularly well adapted to this region. Beardless 20, a hooded variety, produced good yields in all of the years it was grown with the exception of 1920. The hooded winter barleys are not, however, as high yielding as those of the Tennessee Winter type.

TABLE 89.—*Annual acre yields of varieties of barley grown at the Jackson (Tenn.) substation in six of the seven years from 1915 to 1921, inclusive*

[Data obtained through the courtesy of the Tennessee Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)						Years grown	Average yield (bus.)	Percentage of weighted mean
		1915	1916	1917	1919	1920	1921			
Union Winter-----	583	19.2	22.0	11.5	22.5	19.2	22.9	6	19.5	108.3
Beardless 20-----	2745	20.0	13.3	14.2	25.6	7.9	18.0	6	16.5	91.7

FAYETTEVILLE, ARK.

The data from the Arkansas Agricultural Experiment Station at Fayetteville were furnished through the courtesy of that station. Only winter barley was grown, usually in connection with experiments with winter wheat and oats. Comparative tests of these three cereals were conducted during the period from 1915 to 1921, inclusive. According to information received from this station, good stands were established each season. No winterkilling due to low temperature or to heaving was observed. The crop proved more hardy than Turf oats, but is less hardy than either winter wheat or rye. Arkansas undoubtedly is in the humid winter-barley district where the varieties of the Tennessee Winter type are probably the ones generally cultivated. As this was not a varietal experiment, no varietal names were mentioned in the data received.

STILLWATER, OKLA.

The data on barley yields at Stillwater, Okla., were furnished through the courtesy of the Oklahoma Agricultural Experiment Station. While these data were not adequate for tabulation, they are of enough significance to warrant consideration. Barley is becoming much more important as a feed crop in this section, but the experiments with this cereal so far have been very limited. For this reason any information as to types grown is of value.

In 1920 the best results were obtained from the fall-sown varieties. A yield of 53.3 bushels was procured from a 6-rowed club barley, which is probably the variety known as Winter Club. It has been grown for many years in small localized areas in the Western States, particularly in Utah, where it is known as Utah Winter. Another 6-rowed variety, thought to be a barley of the Tennessee Winter group, is commonly grown in the vicinity of Stillwater. It produced the relatively low yield of 26 bushels, but an improved selection of this variety yielded 32.2 bushels. The barleys sown in the spring of 1920 were practically failures. Scarcely more than enough seed for 1921 was obtained from these plats.

The fall-sown barleys also were the higher yielding sorts in 1921. The common 6-rowed barley, which is thought to be of the Tennessee Winter type, yielded 36.6 bushels. It was again surpassed by the improved strain just mentioned, which this year produced 50.1 bushels. Results with the spring varieties

were somewhat better than in 1920, although they did not compare favorably with the winter sorts. The acre yields varied from 3.2 to 17.1 bushels.

It is evident that the fall-sown barleys are more successful than the spring-sown sorts in the vicinity of Stillwater. According to Adrian Daane, of the Oklahoma station, relatively good yields have been obtained from the spring varieties when fall sown in the eastern part of the State, where they furnish winter pasture and later a crop of grain. In western Oklahoma the spring sorts, Coast and Stavropol, are probably grown along with the winter barleys of the Tennessee Winter type. Both Coast and Stavropol are extensively grown in western Kansas and have spread southward through Oklahoma into northern Texas. Barley as feed is considered an excellent substitute for kafir and corn where these crops have failed.

TEXAS

Experimental data on barley production in Texas are very limited. Most of the varietal records were obtained 15 or 20 years ago. It would appear that barley is not an important crop in Texas. According to A. H. Leidigh, agronomist, College Station, Tex., "From a practical standpoint, everything centers around Tennessee Winter barley. Possibly, however, there are indications of the importance of other things."

Most of the barley acreage is in northwestern Texas adjacent to Oklahoma, where the Stavropol and Coast varieties of western Kansas and Oklahoma probably might be grown. On the remainder of the acreage Tennessee Winter doubtless is the prevailing type.

NEW HOPE, TEX.

The barley yields recorded at New Hope, near Dallas, in 1903 and 1904 are shown in Table 90. These yields were obtained in cooperation with the Texas Agricultural Experiment Station. Nine varieties were grown for the two years, but the period is too short to justify the drawing of reliable conclusions. The two varieties which produced the best average yields were Svanhals (C. I. No. 187) and Hanna (C. I. No. 203), both of which are 2-rowed humid-spring sorts usually grown in cooler regions of the North. Telli (C. I. No. 194), a barley of the Coast type, was third, and Bohemian (C. I. No. 188), another barley of the Hanna group, was fourth in point of yield. None of the varieties of the Tennessee Winter group were included in the test. While the spring barleys appear to have produced high yields in the two years they were grown, the period was not of sufficient length to establish their value. These same barleys have not generally shown themselves adapted to cultivation under climatic conditions such as exist here.

TABLE 90.—*Annual acre yields of varieties of barley grown at New Hope, Tex., in 1903 and 1904*

[Data obtained in cooperation with the Texas Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)		Years grown	Average yield (bus.)	Percentage of weighted mean
		1903	1904			
Beldi Dwarf.....	190	63.6	20.0	2	41.8	87.1
Telli.....	194	74.5	34.0	2	54.3	113.1
Chevalier II.....	200	59.3	45.0	2	52.1	108.5
Svanhals.....	187	74.5	40.0	2	57.3	119.4
Kitzing.....	189	60.0	35.0	2	47.5	99.0
Hanna.....	203	70.3	40.0	2	55.1	114.8
Bohemian.....	188	58.1	46.0	2	52.1	108.5
Hankow.....	197	36.1	20.0	2	28.1	58.5
Do.....	192	51.1	37.0	2	44.1	91.9

MARILLO, TEX.

The data from the Amarillo Cereal Field Station found in Table 91 were obtained independently by the Office of Cereal Investigations. In 1906 and 1907 six varieties were grown, and all except Tennessee Winter were spring sorts. Tennessee Winter (C. I. No. 257) was the only variety tested for the entire period from 1906 to 1919, inclusive. In 1906 the spring barleys did not com-

pare at all favorably with Tennessee Winter, but in the following year the latter entirely winterkilled. While it produced high yields in some of the years, it gave little or no return in others. Winterkilling occurs too often here for the safe cultivation of winter varieties. The spring sorts tested all produced poor yields. Barleys of the Coast and Stavropol types should do better than those tried. These varieties, however, were not included in the experiments.

TABLE 91.—*Annual acre yields of varieties of barley grown at the Amarillo (Tex.) Cereal Field Station in some or all of the 14 years from 1906 to 1919, inclusive*

[Data obtained in cooperation with the Texas Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)														Years grown	Average yield (bus.)	Percentage of weighted mean
		1906	1907	1908	1909	1910	1911	1912	1913	1914	1915	1916	1917	1918	1919			
Tennessee Winter.....	257	23.9	0	22.5	0	3.1	14.1	10.6	6.3	35.1	24.6	6.6	16.3	0	25.7	14	13.5	143.6
Horsford.....	507	6.3	3.4	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	2	4.9	96.1
Hanna.....	203	3.5	2.2	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	2	2.9	56.9
Manchuria.....	244	3.3	2.9	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	2	3.1	60.8
White Smyrna.....	195	1.8	6.2	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	2	4.0	78.4
Club Mariout.....	261	2.3	5.0	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	2	3.7	72.5

CHANNING, TEX.

The data on the varietal experiments at Channing, Tex., were obtained in cooperation with the Capitol Syndicate Freehold Land & Investment Co. on the X. I. T. Ranch. The yields of these varieties for the years 1905 and 1906 are shown in Table 92. In 1905 one winter and four spring barleys were grown. Club Mariout (C. I. No. 261) and White Smyrna (C. I. No. 195) produced the highest yields. Tennessee Winter (C. I. No. 257) was third in rank. The humid-area spring varieties, Manchuria (C. I. No. 244) and Hanna (C. I. No. 203), did not compare favorably with these other sorts. Horsford (C. I. No. 507) was introduced in 1906. It yielded only 8.5 bushels, as compared with 25.1 bushels for Tennessee Winter. As the climatic conditions at Channing and Amarillo are quite similar, it is likely that winter barley would be too frequently winterkilled for successful cultivation. The Coast and Stavropol barleys might be successful here.

TABLE 92.—*Annual acre yields of varieties of barley grown at Channing, Tex., in one or both of the years 1905 and 1906*

[Data obtained in cooperation with the Capitol Syndicate Freehold Land & Investment Company]

Variety	C. I. No.	Acre yields (bushels)		Variety	C. I. No.	Acre yields (bushels)	
		1905	1906			1905	1906
Tennessee Winter.....	257	11.4	25.1	Manchuria.....	244	8.5	-----
White Smyrna.....	195	12.5	-----	Hanna.....	203	5.7	-----
Club Mariout.....	261	12.7	-----	Horsford.....	507	-----	8.5

STATE COLLEGE, N. MEX.

The annual yields of barley varieties grown at the New Mexico Agricultural Experiment Station were obtained through the courtesy of that station. These yields for five years, 1913, 1916, 1917, 1920, and 1921, are shown in Table 93. All of the varieties grown during the years 1913, 1916, and 1917 were spring sorts, although they were fall sown. Of these Scotch (C. I. No. 848) produced the highest average yield for the three years. Frankish (C. I. No. 207) was second, and Six-Row (C. I. No. 2811), a barley of unknown identity, was third in point of yield. With the exception of Coast and Telli, however, all these

varieties are humid-area spring sorts, and when fall sown or winter sown they produce an excessive growth, thus making their chances of winterkilling much greater. This was the cause of the very low yields in 1916.

The varieties tested in 1920 and 1921 with the exception of Coast were all winter sorts. Tennessee Winter (C. I. No. 257) far exceeded all others with an average yield of 74.5 bushels, as compared with 58.5 bushels for O. A. C. Selection 7 (C. I. No. 2814), its nearest competitor. Coast (C. I. No. 690) was third in rank. In 1913, however, the two Coast barleys, Telli and Coast, were the lowest yielding sorts.

Barley as an agricultural crop in New Mexico is of only minor importance. It is grown largely for winter pasturage. Although Tennessee Winter was not sufficiently tested at this station to determine its relative importance, it is both high yielding and winter hardy and is probably unequaled for the purpose for which barley is usually grown in this State. For this reason State College is considered with the stations in the winter-barley region, although ecologically it belongs in the Coast region.

TABLE 93.—*Annual acre yields of varieties of barley grown at the New Mexico Agricultural Experiment Station (at State College) in one or more of the nine years from 1913 to 1921, inclusive*

[Data obtained through the courtesy of the New Mexico Agricultural Experiment Station]

Variety	C. I. No.	Acre yields (bushels)					Years grown	Average yield (bus.)	Percentage of weighted mean
		1913	1916	1917	1920	1921			
Frankish	207	51.1	9.5	40.4			3	33.7	111.6
Coast	690	28.1			55.6	56.6	3	46.8	95.1
Six-Row	2811	61.6	9.7	25.7			3	32.3	107.0
Bavarian	208	43.1					1	43.1	102.4
Scotch	848	59.9	8.3	34.6			3	34.3	113.6
Hull-less	2812	29.7	6.9	22.5			3	19.7	65.2
Telli	194	21.5					1	21.5	51.1
Maryland Winter	518				51.6	46.2	2	48.9	92.3
Nakano Wase (See Pl. III, fig. 1, D)	754				14.6		1	14.6	26.6
Wisconsin Winter	519				51.5	37.1	2	44.3	83.6
Tennessee Winter	257				83.1	66.0	2	74.5	140.6
O. A. C. Selection 8	2813				61.5	47.7	2	54.6	103.0
O. A. C. Selection 7	2814				66.0	51.1	2	58.5	110.4

REGIONAL ADAPTATION OF VARIETIES

In order to make the discussion of results at each station fairly complete, statements of the adaptation of varieties and types have been repeated throughout the bulletin. When the results of all the stations are brought together they should enable one to map the various regions of barley culture. There are many difficulties, however, in mapping. The boundary lines are intangible and fluctuate from year to year according to the season. The data available are often fragmentary or for so few years as to be inadequate. The boundaries themselves do not separate the regions of adaptation of one type from that of another. There are lines where two well-adapted types are equal in value and on each side of which either produces good yields. Often types of probable high yield have not been tested or have been inadequately tested. It is possible that in many cases the most desirable sorts have never been introduced into America. For these reasons the history of barley cultivation and of varieties is a very desirable background for defining present limits.

The regions are most easily separated on a basis of rainfall and temperature. In States east of the Rocky Mountains rains occur during the growing season. In this district of summer rainfall there

are three distinct subdivisions: (1) Those States north of the Ohio River extending from the Atlantic seaboard to the northern Great Plains area, (2) the States of the northern Great Plains, and (3) the Southern States.

The Northeastern States are characterized by moderate summer temperature and an ample summer rainfall. This area includes the most important barley-producing center east of the Rocky Mountains. On the northern Great Plains there is precipitation during the summer, but the quantity is considerably less than in the Northeastern States, and droughts are frequent. In the Southern States the summers are hot, and the winters are sufficiently mild for the growing of winter varieties.

West of the Rocky Mountains the rainfall occurs during the winter season, and the crop ripens during hot dry weather.

SPRING BARLEYS FOR HUMID AREAS

In the earlier years in New England the 2-rowed barleys of the Thorpe and Chevalier types predominated. The barley acreage now has almost disappeared. New York was for many years an important center of production. Both soil and climate are suited to the growing of barley, which was in great demand for brewing and malting purposes. As has been previously stated, the types grown in these earlier years were doubtless the 2-rowed Hanna and Chevalier and a 6-rowed barley of unknown identity, probably brought from western Europe and northern England. At the present time types adapted to general farming, such as Manchuria and Hanna, usually are grown. To supply the demand for feed in the dairy industry the acreage sown to these varieties has been considerably increased in recent years. In Pennsylvania, where barley cultivation is of only local importance, the varieties are of the same general types as in New York.

In the upper Mississippi Valley the Manchuria-Oderbrucker varieties have been grown almost exclusively for many years. The earlier agricultural records show that a 6-rowed barley which was probably quite similar to the varieties now grown commercially was in cultivation there.

Numerous experiments at the stations in the various States have been conducted for a sufficient number of years to prove beyond doubt that the Manchuria-Oderbrucker types are best adapted to the region. The pioneer stations in barley experimentation were those in Wisconsin, Minnesota, and North Dakota. The first experiments at Madison, Wis., were conducted in 1870; at Fargo, N. Dak., in 1892; and at St. Paul, Minn., in 1893. At these stations the Manchuria-Oderbrucker varieties were superior to all others tested. From the types grown in the various localities and from the results at the various stations, boundaries can be defined.

Spring barleys adapted to humid climates are grown in the northeastern part of the region of summer rainfall. This area (fig. 10) is arbitrarily bounded by a line starting near Indian Head, Saskatchewan, crossing the northern boundary of North Dakota near the Montana line, running diagonally to southeastern South Dakota, and continuing to southeastern Iowa. Then crossing Illinois and Indiana it divides these two States in halves. It then curves northward to

the more favorable soils of the upper counties of Ohio, thence southward to central Pennsylvania and east to the coast. The accompanying map shows that this humid spring-barley region extends far into Canada. In most of this region the barleys are of the Manchuria type, although in a small area in Maine and southeastern Canada Chevalier is grown and there is a considerable acreage of Hanna and other 2-rowed barleys in New York and Pennsylvania.

Just west of this humid region is an area of summer rainfall where the precipitation is quite light. Here a number of 2-rowed varieties do particularly well. This is the largest area in the United States suited to the growth of the 2-rowed sorts. It includes southwestern Iowa, northwestern Missouri, all of Nebraska, extensive areas in the

FIG. 10.—Map of the United States and Canada showing the areas and sections to which groups of barleys are adapted

western parts of the Dakotas, the northeastern half of Wyoming, and all of Montana with the exception of a western fringe. It extends far into Canada, embracing all the territory west of Indian Head, Saskatchewan, to the mountains, and theoretically should connect with the Chevalier district of the extreme eastern Provinces by crossing above the central Manchuria area. All of the 2-rowed barleys grown here except White Smyrna are spring sorts adapted to humid climates. In Montana, where it is cool and the days are long, the varieties of the Chevalier group are grown quite successfully. They are even more successful in Canada, where these conditions are intensified. In those portions of Wyoming, North Dakota, South Dakota, and Nebraska adapted to 2-rowed barleys Hannchen and White Smyrna are superior sorts, Hannchen producing higher yields where there is more moisture. The Thorpe barleys have been tested in the

northern part of this region, but have not proved superior to Hannchen or the Chevalier varieties.

In addition to the areas mentioned, 2-rowed barleys do well in two districts in the West. In Washington and Oregon there is a district including Moro, Union, and Burns, Oreg., where Smyrna and Hannchen produce high yields, while in a part of the Salinas Valley of California practically the only variety grown is a Chevalier barley.

There are thus three distinct centers where different types of barleys adapted to humid climates seem to do particularly well. These are (1) the Northeastern States, where Hanna and Manchuria do equally well; (2) the upper Mississippi Valley, where the Manchuria-Oderbrucker barleys predominate; and (3) the large area west of the upper Mississippi Valley, where the 2-rowed barleys are well adapted, an area included with these districts for two reasons. The rainfall, although too light to characterize the region as really humid, is a summer rainfall, nevertheless. Many of the varieties grown are those originating in humid climates.

The difficulty of defining either the regions or the range of varieties is apparent. The Hanna and Chevalier barleys succeed along the Atlantic seaboard from New England to Prince Edward Island, because the summers are somewhat similar to the cool rainy summers of northern Europe, whence they came. These conditions are partially fulfilled in the northern Great Plains, where there are sections in which they do well despite the lower rainfall and lesser humidity. In part of this latter area White Smyrna from the dry coast of Asia Minor grows well in the same district as Hannchen from Sweden.

In the same way the growth of plants on soils of low water content, as found in the West where the rainfall comes in winter, is duplicated east of the Rocky Mountains on the western Plains, where the rainfall is light and comes in summer. The coast district overlaps the western 2-rowed area. Farther south, in western Kansas and adjoining sections, it similarly overlaps the border of the humid-spring region. In this section Coast and Stavropol, allied varieties tolerant of more humid conditions, are largely grown, and this section is pertinently considered with the western region.

SPRING BARLEYS FOR SEMIARID AREAS

West of the Rocky Mountains, Coast is the dominant commercial variety, constituting most of the acreage. East of the mountains in the Great Plains area its acreage exceeds that of all other semiarid barleys, if Stavropol, which is of Russian origin, is excepted. Because of the extreme variation in the topography and of the precipitation of the region, numerous varieties other than Coast and Stavropol are cultivated. Many of these are 6-rowed barleys of North African origin. In western Kansas and southern Nebraska the similar varieties Stavropol and Coast are widely cultivated. A small part of the Colorado acreage is devoted to the growing of hull-less varieties. In Colorado, Nebraska, Wyoming, Montana, and the western Dakotas the 2-rowed humid-spring types already discussed and a number of other types occur. Chief among these are Coast, White Smyrna from Asia Minor, Club Mariout of the North African group, and Trebi from Asiatic Turkey, the latter usually grown under

irrigation. In Idaho, Oregon, and Washington Coast is the dominant variety, although Trebi, White Smyrna, Club Mariout, Beldi Giant, and other varieties of the Coast type are promising. In California the barley acreage and production exceed those of any other State. By far the most important variety is Coast. California Mariout is grown in parts of the interior valleys where warmer and drier conditions prevail. There is an increasing acreage of Club Mariout in the Sacramento Valley and in the northern portion of the San Joaquin Valley, where it brings good prices for export.

WINTER BARLEYS FOR HUMID AREAS

The humid winter-barley region extends from the Atlantic seaboard to western Texas and as far north as central Ohio, including several very different areas: The Piedmont, the Coastal Plain, the southeastern mountain section, the Ohio Valley, and the southern area west of the Mississippi. Winter barleys are grown in only a few sections of the South, the center of production being in Tennessee.

Experiments in this region have been few. Occasional satisfactory yields have been obtained, but the available information is so meager that conclusions are quite untrustworthy. The fact that barley is not grown is an indication of its lack of suitability.

The potential economic importance of barley in the Coastal Plain section is great. Most of the hogs of this plain are fattened on peanuts, resulting in soft pork. Barley is one of the best of all feeds for hardening pork. Where it can be grown here it will assist greatly in establishing the hog industry on a satisfactory basis. Unfortunately barley is not very promising when grown under Coastal Plain conditions.

Much higher yields have been obtained in the Piedmont region. This section is much better suited to production than the Coastal Plain, but even here barley does not grow particularly well.

In eastern Tennessee barley is an important crop. Here both climatic and soil conditions are favorable. The dominating variety is Tennessee Winter, the most frost-resistant variety known in humid climates. Recently Professor Mooers, of the Tennessee Agricultural Experiment Station, has developed some hooded hybrids which seem to be true winter barleys and have done well in Tennessee.

The northern limit of safe cultivation of Tennessee Winter barley is rather indefinite. Besides the area under cultivation in Tennessee, some winter barley is grown commercially in Ohio, and there is even a small acreage in Michigan. Barley winterkills, however, in Kentucky and is grown but rarely in northern Ohio. The southern limit of winter-barley production is equally indefinite. The true winter sorts are unquestionably the ones best adapted to Ohio, Tennessee, and northwestern Georgia. South of this area it is probable that fall-sown spring varieties are better adapted, although information on this point is very limited. Experiments were conducted at Tifton, Ga., in 1921 in cooperation with the experiment station at that point. Spring barleys responded favorably to the early season which occurred that year, while the winter forms remained dormant until the days were of sufficient length to induce flowering. Spring barleys under these conditions might be caught by late frosts and in the long run prove unsuited. With the information available,

however, spring sorts appear to be worthy of extensive trials on the southern Coastal Plain. Yields equal to those obtained in Minnesota and Wisconsin can hardly be expected, as the soil type is not one to which barley is well adapted, and the climatic conditions are unquestionably more adverse.

East of the Mississippi River there is a conspicuous gap between the winter and spring areas. Spring barley is at present grown extensively in northern Illinois, Indiana, and Ohio. South of this section little barley, either spring or winter, is grown until the winter-barley district of Tennessee is reached. On the Great Plains the spring-sown semiarid forms and the fall-sown humid forms of the Tennessee Winter type overlap. In western Oklahoma and southwestern Kansas both types are grown. Winter barley also is cultivated on the Plains in Texas.

WELL-KNOWN OR PROMISING VARIETIES

At various times varieties of barley have been introduced into the United States or developed at the agricultural experiment stations. Some of these disappeared quickly, while others were tested for long periods at many stations. The number of varieties which had been tested up to 1921 was so large that no attempt has been made to describe them individually. From this number, however, a few of the more important and better known varieties were chosen for the purpose of amplifying the meager details mentioned in describing the results at the several stations.

The varieties are not presented in the order of their Cereal Investigations (C. I.) numbers. This arrangement, perhaps, would have been the most desirable in checking references arising within the text of the bulletin. For any question arising outside this discussion the varieties are more easily found when arranged in related groups. Thus, in the following descriptions the Chevalier barleys are grouped together, as are the Manchuria varieties, etc. Where it is desired to check varieties by the Cereal Investigations numbers they will be found in numerical order in Table 99, in which also is inserted a reference to the varietal description under this heading.

TWO-ROWED HULLED AWNED BARLEYS

CANADIAN THORPE (C. I. NO. 740)

A variety known as Canadian Thorpe has been widely tested at stations in Canada and has been grown on a few experiment stations in the United States. It is a barley of the Thorpe type. According to Dr. C. E. Saunders, "Canadian Thorpe was introduced into Canada many years ago by a Canadian seedsman and is essentially identical with Duckbill." The original Canadian Thorpe produced good yields in the western prairie Provinces. Selections were made from the Canadian Thorpe, which Seager Wheeler, of Rosthern, Saskatchewan, considered to be superior to the original variety. This barley was grown at Moccasin, Mont., for three years. During these years its yield was 98 per cent of that of the weighted mean of all varieties.

INVINCIBLE (C. I. NO. 590)

Invincible (C. I. No. 590) was originated by the Garton Brothers, seedsmen, of England. It came from a compound cross, Chevalier × Golden Melon, crossed on Standwell. Invincible is a late 2-rowed variety of the Goldthorpe type. It has large straw and a dense spike. It produced relatively higher yields in Canada than in the United States.

STANDWELL (C. I. NO. 584)

Standwell is a hybrid barley which has been classed in the Thorpe group in this bulletin. It was produced by Garton Brothers, of England. It came from a cross, Fan × Golden Melon. It may not be entirely consistent to place Standwell and Invincible in the Thorpe group, as they might more consistently be placed with the hybrids. It is entirely possible, however, that the entire group was of hybrid origin. Those standard varieties of England with wide spikes may very easily have come from accidental hybrids. While this type is a desirable one in England there is no area in the United States to which it is well adapted.

SVANHALS (C. I. NO. 187)

Originated by the Svalof Plant-Breeding Association, Svalof, Sweden, Svanhals (C. I. No. 187) was first introduced into the United States by the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture in March, 1901. It was reintroduced in 1906 as C. I. No. 605. In Sweden this variety was considered suited to cold wet land. It was not there believed to be the equal of Princess (C. I. No. 529) or of Chevalier II (C. I. No. 200).

In the United States it has been superior to both these varieties at almost all places. It has proved to be adapted to an unexpectedly wide range, producing relatively good yields in the northern Great Plains and also under irrigation farther west. Svanhals has been grown to some extent in field culture, as shown in Plate I, D. It is a 2-rowed barley with a dense spike, similar to Thorpe, but ripens earlier than the Thorpe barleys. The density of Svanhals was about 2.6 millimeters in an ordinary year at Aberdeen, Idaho.

EARLY CHEVALIER (C. I. NO. 2725)

According to the Canadian Experimental Farms Report of 1908 Early Chevalier (C. I. No. 2725) is a selection from French Chevalier made in 1904 on account of earliness and strength of straw. While it has many of the characters of Chevalier, it is hardly typical of that variety. The spikes are somewhat wider at the base than is the case with most varieties of the Chevalier group. Early Chevalier is one of the highest yielding sorts at Cap Rouge. It has not been as widely tested in Canada as many of the other varieties.

FRENCH CHEVALIER (C. I. NO. 175)

A barley known as French Chevalier was widely tested at stations in Canada. It is mentioned in the Canadian Experimental Farms Report of 1892. This variety also was grown at a number of stations

HEADS OF TWO-ROWED BARLEY VARIETIES

A, Chevalier II (C. I. No. 200), an English brewing barley adapted to sections with cool summer temperatures in Canada and the United States; B, Hammett (C. I. No. 531), a Swedish brewing barley adapted to sections with summer temperatures slightly higher than those suited to Chevalier; C, Alpha (C. I. No. 953), a barley of recent hybrid origin not yet widely distributed, promising in New York State; D, Svanhals (C. I. No. 187), a Swedish variety adapted to much the same sections as Hammett; E, White Shyryna (C. I. No. 165), from Asia Minor, widely adapted to the drier sections of the United States.

HEADS OF BARLEY VARIETIES.—I

A, Manchuria (C. I. No. 24), a barley of the Manchuria group of varieties grown on most of the northern Mississippi Valley where the summers are hot and humid; B, Coast (C. I. No. 690), a head from varieties grown on most of the barley acreage of the Western States where the grain ripens during a season of low rainfall; C, Club Marston (C. I. No. 261), an Egyptian variety with requirements similar to those of Coast, commercially grown in California; D, California Marston (C. I. No. 1455), a dry-land variety from the Egyptian sea-coast suited to conditions of very low rainfall, especially in California.

in the United States. Both the original variety and selections from it are still being grown. The French Chevalier itself, while a very good variety, has not been outstanding at any one point. At Nappan, Nova Scotia, it produced good yields, but not so good as Mensury (C. I. No. 2657). This is not surprising, as the number of stations where the Chevalier group is the highest producing one is few.

CHEVALIER II (C. I. NO. 200)

Originated by the Svalof Plant-Breeding Association, Svalof, Sweden, Chevalier II (C. I. No. 200) was imported into the United States by the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture. Several importations were made, the first of which was C. I. No. 200 (Seed and Plant Introduction No. 5473), received March 11, 1901. Another importation (S. P. I. No. 10584; C. I. No. 530) was made in 1904, and a third (S. P. I. No. 19781; C. I. No. 604) was made in 1907. These later numbers have been discarded in favor of C. I. No. 200. The original selection was made by Doctor Nilsson from Horsford Chevalier and was a pedigreed barley of the Chevalier type. It is about as dense as Charlottetown 80; in other words, more dense than the usual Chevalier. A spike of Chevalier II is shown in Plate I, A.

HORN (C. I. NO. 926)

Horn is a pedigreed selection from C. I. No. 31 of the same name, which originally came from Gassen, Austria. This selection, a 2-rowed bearded barley of the Chevalier group, was made by the United States Department of Agriculture in breeding experiments in cooperation with the Minnesota Agricultural Experiment Station at St. Paul, Minn. Small quantities of seed were sent from St. Paul to a number of places, where it was grown in nursery rows. At Archer, Wyo., it showed such promise that it was increased and included in the field plats. The yields on the larger area continued to be large, and seed was sent to a number of other stations for plat testing. While it has not been widely tested it has given high yields not only at Archer but also at Havre, Mont., where it produced the highest average yield of all varieties grown for the five years from 1917 to 1921, inclusive.

CHARLOTTETOWN 80 (C. I. NO. 2732)

Charlottetown 80 was selected from Old Island Two-Rowed by J. A. Clark, superintendent of the Experimental Farm, Charlottetown, Prince Edward Island, Canada.

The Old Island Two-Rowed is a variety that has been grown on Prince Edward Island for many years. It consists principally of Chevalier types but also contains a much taller barley with a dense spike. It was doubtless brought there from England. Mr. Clark selected one of the Chevalier types which had a tendency to drop the awns at maturity. While this barley belongs to the Chevalier group it would seem that some modification has occurred which favors its culture in areas south of those normally best suited to that group. In recent years it has given very good results not only

at Charlottetown but at many of the stations in western Canada and in the United States. It is unquestionably one of the best Chevalier selections which have been made so far as its culture in America is concerned. Besides the tendency to drop the awns this selection often has more red on the nerves of the lemma just before maturity than is common with the Chevalier barleys.

PRINCESS (C. I. NO. 529)

Princess (C. I. No. 529) is a pedigreed variety originated by the Svalof Plant-Breeding Association, Svalof, Sweden. It was first imported into the United States by the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture in April, 1904, under S. P. I. No. 10583. Several other importations of Princess were made under later numbers, but because it is a pedigreed variety these numbers have been discarded in favor of C. I. No. 529. It is a 2-rowed bearded barley. Like the Chevalier it requires a long growing season, but it is usually several days later in ripening than Chevalier. It has not proved a high-yielding sort in the United States.

HANNA (C. I. NO. 203)

Introduced into the United States by the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture in January, 1901, from Kwassitz, Moravia, Austria, Hanna (C. I. No. 203) has been grown for many years in Germany and Austria for brewing purposes. It is a 2-rowed bearded barley with lax head like that of Chevalier. It is earlier than Chevalier and has a long-haired rachilla. While it has been widely tested at a number of experiment stations throughout the United States, it has not been outstanding in point of yield at any one of them. This type of barley seems best adapted to the New England States, New York, Pennsylvania, and western North Dakota. At the Cornell Agricultural Experiment Station several crosses have been made with Hanna which seem very promising.

GOLD (C. I. NO. 1145)

Produced by the Svalof Plant-Breeding Station of Svalof, Sweden, several importations of Gold (C. I. No. 1145) have been made both by the United States Department of Agriculture and by experiment stations in the United States and Canada. As it originated as a plant selection the various importations may be considered identical. Gold is a barley of the Hanna type. Its appearance is not at all striking. It was only after it had been tested for a number of years at the Svalof station that its yielding ability was sufficiently demonstrated to attract attention. This variety has been widely tested in the United States and Canada in recent years. It has been in the test only a few years at most places. In recent years it has produced high yields at Ottawa, Ontario, and Indian Head, Saskatchewan, but was inferior to Manchurian (C. I. No. 739) at Ottawa and to O. A. C. 21 (C. I. No. 1470) at Indian Head. It was not so good as Garton 68 (C. I. No. 2740), Hannchen (C. I. No. 531), or Manchurian (C. I. No. 739) at Brandon, Manitoba. In the United States Gold has been grown at a number of stations. It has shown promise over a

wide range of territory but has not been outstanding at any place. At some points it has produced very high yields, but the number of years it has been in the tests has been too few to determine its value.

HANNCHEN (C. I. NO. 531)

Hannchen (C. I. No. 531) was introduced into the United States by the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture in April, 1904. It was originated by the Svalof Plant-Breeding Association of Svalof, Sweden, and is a 2-rowed barley of the Hanna type. (Pl. I, B.) In 1921 at Aberdeen, Idaho, the average length of the internodes of 25 spikes was 3.4 millimeters.

Hannchen has given by far the best yields and has exhibited a much wider range than any of the other barleys imported from Svalof. Although coming from the very different climate of northern Europe it has proved to be of especial value in regions of low rainfall in the United States. In the past five years it has been among the best five varieties at 13 stations located between Eureka, S. Dak., and Burns, Oreg.

ALPHA (C. I. NO. 959)

Alpha (C. I. No. 959), shown in Plate I, C, is a 2-rowed selection from a cross of Manchuria $\times$ Champion of Vermont. It was produced in the cooperative breeding experiments conducted by the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul, Minn. It was later tested and distributed by the agricultural experiment station at Cornell University, Ithaca, N. Y. It has produced high yields on the farms in New York State and its acreage is increasing.

WHITE SMYRNA (C. I. NO. 195)

Received by the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture from Smyrna, Asia Minor, on November 15, 1901, White Smyrna (C. I. No. 195) is a 2-rowed bearded barley with an average density of 3.6 millimeters at Aberdeen, Idaho, in 1921. (Pl. I, E.) It has been widely tested at the experiment stations in the western half of the United States and is the most drought resistant of any of the varieties so far tested. The straw is very short, and the head when fully emerged usually remains about half inclosed in the boot. In point of yield it has done particularly well in sections of Montana, Wyoming, western South Dakota, northern Colorado, Idaho, and eastern Oregon.

White Smyrna (C. I. No. 910) is a selection of this variety which had done particularly well at Moccasin, Mont.

White Smyrna under numbers C. I. 195, 658, or 910 has been one of the five highest yielding varieties at 22 stations in the past five years. These cover an area from Eureka, S. Dak., to Chico, Calif., and from Havre, Mont., to Tribune, Kans.

WHITE SMYRNA (C. I. NO. 658)

White Smyrna (C. I. No. 658) differs only in a few minor details from White Smyrna (C. I. No. 195). It has been grown under the name "Ouchac," or "Ouchac Smyrna," at a number of the western stations. The most characteristic difference in the two White

Smyrna strains as grown at Aberdeen, Idaho, in 1921 was the longer straw length of C. I. No. 658. At Bozeman, Mont., Burns, Oreg., Dickinson, N. Dak., and Archer, Wyo., it has yielded well.

SIX-ROWED HULLED AWNED AND AWNLESS BARLEYS

MANCHURIA (C. I. NO. 244)

All the unpedigreed Manchuria barleys originally sent out from the Ontario Agricultural College, Guelph, Ontario, have been given the number C. I. 244. According to Ontario Agricultural College Bulletin No. 71, page 6, this barley, under the name "Mandscheuri," was "imported from Russia in 1881 and is probably the same as Mensury." The writers do not know the full history of this variety. It was grown and distributed under the name "Mandscheuri," which is German for Manchuria. Other variations of the spelling occurred later. It is not stated in the report whether the barley was imported direct from Russia or through Germany. All varieties tracing to Guelph have been kept separate from those tracing to Madison, Wis. It may be that it is from the same source and the same stock as C. I. No. 241. As both varieties are merely mixtures of an indefinite number of strains, their culture at Madison and Guelph would eventually result in a different residual complex, whether they were the same or different to start with. The variety resulting at Guelph seems to have been the better of the two for most of the Manchuria district.

In 1894 this variety was sent from Guelph, Ontario, to St. Paul, Minn., from which place it was received at Fargo, N. Dak. It was soon grown by many seedsmen and was rapidly distributed to the various experiment stations and farmers in the upper Mississippi Valley. At many of the stations it is still grown as Minnesota No. 32 or Minnesota No. 105. Its immediate origin is thought to be the humid plains of eastern Manchuria, though descendants from the same common stock doubtless spread to the plains of central Europe. In the United States its area of best adaptation is undoubtedly the humid States of the upper Mississippi Valley, where soil and climatic conditions are quite comparable to those of central Eurasia. Manchuria is a 6-rowed bearded variety with lax head and long straw. (Pl. II, A.) Since it is a mixture of a number of strains the aleurone layer may be either blue or white. This variety has been widely tested for many years in the United States. It has produced exceptionally high yields at St. Paul, Minn., and Fargo, N. Dak., and is the common barley grown commercially in these States. A number of selections have been made from C. I. No. 244 which have also been high yielders at the various stations.

MANCHURIA (C. I. NO. 241)

Manchuria (C. I. No. 241) was the first introduction of the Manchuria barley into the United States. About 1859 a traveler brought to Germany a sample of barley from the Amur River district of Manchuria. About 1861 Dr. Herman Grunow, of Mifflin, Iowa County, Wis., while on a visit to Germany, secured some of this seed, which he brought to America. The Wisconsin Agricultural Experiment Station received this barley from Doctor Grunow and grew it

in 1872. For many years this was the best variety at the Wisconsin station. It was widely distributed to farmers in Wisconsin and at one time constituted a large part of the acreage of barley in that State. It is probable that there is much of the original "Manshury" stock or selections of it grown in Wisconsin and neighboring States. This Wisconsin Manchuria was sent to other stations and is the foundation of many selections made elsewhere. It proved to be generally better than the 6-rowed barleys from western Eurasia more commonly grown before its introduction. This variety was grown and distributed under various names and State numbers, such as Minnesota No. 6, North Dakota No. 787, Manshury, Mansury, and Manchuria.

MANCHURIA (C. I. NO. 2330)

Manchuria (C. I. No. 2330) is a selection made at the Minnesota Agricultural Experiment Station, where it is grown as Minnesota No. 184. This barley has been tested at comparatively few stations outside of Minnesota but produced high yields for the years it was grown at La Fayette, Ind. At all of the Minnesota stations where barley has been grown Manchuria (C. I. No. 2330) has been one of the best varieties. It seems to be a high-yielding Manchuria strain that is particularly well adapted to Minnesota conditions.

SILVER KING (C. I. NO. 890)

Silver King (C. I. No. 890) is a barley of the Manchuria type which has been widely distributed over the United States by the Wisconsin Agricultural Experiment Station. It has proved a high-yielding variety at many places. Numerous high-yielding selections of Silver King were made at Madison.

FEATHERSTON (C. I. NO. 1120)

Featherston (C. I. No. 1120) was first grown in the cooperative breeding experiments conducted by the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul, Minn. It came from a head selection made on a farm near Red Wing, Minn. This barley belongs to the Manchuria group. Since 1913 it has been grown at the Cornell Agricultural Experiment Station at Ithaca, N. Y., where it has yielded exceptionally well. During these years its yield was 115.3 per cent of that of the weighted mean of all varieties. Featherston has been distributed from Ithaca to farmers and to other stations.

O. A. C. 21 (C. I. NO. 1470)

A selection of Manchuria (C. I. No. 244) made by Prof. C. A. Zavitz at the Ontario Agricultural College at Guelph, Ontario, proved to be of unusual merit and was given the number O. A. C. 21 (C. I. No. 1470). It is probably the best Manchuria selection tested in Canada. It has been grown at a number of points in the United States in recent years. The yields have not been relatively so high as in Canada, but the variety should be more thoroughly tested in the States. It produced the best yields at Indian Head, Saskatchewan, and yielded well at Lacombe and Rosthern.

WISCONSIN PEDIGREE (C. I. NO. 835)

Wisconsin Pedigree is a selection made from a barley of the Manchuria type at Madison, Wis. It has been distributed to a number of stations where it has given high yields.

MENSURY (C. I. NO. 2657)

A barley of the Manchuria group very widely tested at the Canadian stations under the name "Mensury" has been grown at a few stations in the United States under this name. The history of this variety is not clear. It may be that a Manchuria barley was received with this spelling, and it may differ a little in origin from C. I. Nos. 241 and 244, which were the common stocks of Manchuria barley found at American experiment stations. Mensury was among the highest yielding barleys at Nappan, Nova Scotia; Ottawa, Ontario; Brandon, Manitoba; and Fort Vermilion, Alberta, Canada. It was mostly grown in the earlier years. A selection from Mensury called Manchurian (C. I. No. 739) has been tested recently at many stations in the United States as well as in Canada.

MANCHURIAN (C. I. NO. 739)

Manchurian (C. I. No. 739) is a selection which was made from Canadian Mensury by Dr. C. E. Saunders, of Ottawa, Canada. It is a barley of the Manchuria type which has a blue color in the aleurone layer. It has been more widely tested in Canada than in the United States.

ODERBRUCKER (C. I. NO. 836)

Oderbrucker (C. I. No. 836) is a pedigreed selection of the Manchuria type made at the Wisconsin Agricultural Experiment Station. This barley has been widely distributed over the United States, giving high yields at many of the stations. While it has been promising over a wide area it is suited to the southern fringe of the Manchuria district relatively better than many other strains of that variety.

ODERBRUCKER (C. I. NO. 1529)

This variety is a pedigreed selection of Oderbrucker barley made at the Wisconsin Agricultural Experiment Station, where it was known as Wisconsin Pedigree No. 6. The Oderbrucker barley belongs to the Manchuria group and seems particularly well adapted to Wisconsin. This selection has produced very good results at Madison, Wis., and at La Fayette, Ind.

ODERBRUCKER (C. I. NO. 2700)

There were several independent introductions of the parent stock of the Oderbrucker variety into the United States and Canada. A reference in the report of the United States Department of Agriculture for 1865 (p. 28) states that this variety "is grown very extensively on the low, formerly swamp lands of the valley of the Oder, but which were drained during the reign of Frederick the Great and has since produced the very best barley known on the continent of Europe. It is a very favorite variety for the porter brewers of Eng-

land and for which they pay a very high price. The sample experimented with, of this variety, was very small, and the season was too far advanced before it was received to test it properly. It, however, came up vigorously, and appears to be of robust growth, but had not time to mature."

In 1889 the Ontario Agricultural College at Guelph received this barley from Germany and later sent it to the Wisconsin Agricultural Experiment Station at Madison, where it was first tested in 1899 as Wisconsin No. 55. From Madison it was widely distributed, and it and later selections from it made by the Wisconsin station are the barleys commonly grown by the farmers of Wisconsin. In its appearance and general botanical description it closely resembles Manchuria. However, it is not so widely adapted to growing conditions of the northern Mississippi Valley as is Manchuria.

STELLA (C. I. NO. 2678)

Dr. C. E. Saunders produced a great many hybrid varieties of barley at Ottawa, Canada. These were distributed to the Dominion experiment farms for testing. Stella is one of the best of these hybrids. It is a selection from a cross of Swedish $\times$ Baxter. The Swedish parent was a 2-rowed barley received from Harris & Co., of London, England. Baxter is a 6-rowed selection with a peculiarly short kernel with thin glumes. The Stella hybrid looks much like a Manchuria barley and has a short-haired rachilla. This variety produced good yields at Charlottetown, Prince Edward Island; Nappan, Nova Scotia; Ottawa, Ontario; and Rosthern, Saskatchewan.

YALE (C. I. NO. 1592)

Yale is one of the barley hybrids which were produced by Dr. C. E. Saunders at Ottawa, Canada. It was a selection from a cross of Duckbill $\times$ Rennie Improved. The strain of Yale now being grown at Ottawa is a Manchuria type of barley with a short-haired rachilla. Yale was the best of the hybrid varieties tested at Brandon, Manitoba. At most places it was not the equal of Stella, but it is unquestionably among the better hybrid varieties produced by Doctor Saunders.

HYBRID (C. I. NO. 2935)

This barley is a 6-rowed smooth-awned hybrid selected from a cross of Lion $\times$ Manchuria. It was developed in the cooperative breeding experiments conducted by the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul, Minn. This hybrid is of special interest, as very few smooth-awned barleys have produced high yields. At Waseca, Minn., it was one of the highest yielding sorts for the three years in which it was grown. It also gave good yields at Grand Rapids, Minn. It is hoped that it will be more widely tested in the Manchuria regions.

MINSTURDI (C. I. NO. 1556)

Minsturdi (C. I. No. 1556) was developed in the cooperative breeding experiments conducted by the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul, Minn. It is a 6-rowed bearded hybrid variety selected

from the cross South African $\times$ Manchuria. At Aberdeen, Idaho, in 1921 it had an average spike density of about 2.8 mm., which is considerably greater than that of the average Manchuria barley. In recent years it has been outstanding in yield at St. Paul, Grand Rapids, and Crookston, Minn.

ODESSA (C. I. NO. 182)

Odessa (C. I. No. 182) is a barley belonging to the same general group as Manchuria. Since it is adapted to conditions in southern Russia, one would expect the region of its best adaptation in the United States to be south and west of the section where Manchuria grows best. This is only partially true. Odessa has produced very high yields in South Dakota and seems particularly well adapted to culture in that State. On the other hand, it has done very well in the Canadian districts where Manchuria is also at its best. For many years Odessa was the best barley at Ottawa, Ontario. It gave good yields at Brandon, Manitoba, and Indian Head, Saskatchewan, but was not so good as Manchuria in Minnesota and North Dakota. Odessa contains several types, and numerous selections were made from it in the cooperative studies at St. Paul, Minn. Some of these have been grown in field plats in recent years. They are not of the same type as the barleys grown in Canada and North Dakota.

STAVROPOL (C. I. NO. 2103)

Stavropol (C. I. No. 2103) belongs to the same general group as the Coast barley and probably originated in southern Russia near the Caucasus Mountains. Being adapted to conditions of this region it is not surprising that this type of barley does well in the western half of Kansas and the southwestern part of Nebraska, where the altitude varies from 2,000 to 4,000 feet and the climatic conditions are comparable. It has produced exceptionally good yields at Hays and Colby, Kans. Stavropol is not so good as Trebi or Coast at North Platte, Nebr., but it is one of the types now grown extensively in western Kansas.

TREBI (C. I. NO. 936)

Trebi barley is a pure-line selection made in 1907 in the cooperative breeding experiments conducted by the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul, Minn. The original importation was brought into the United States by the Office of Foreign Seed and Plant Introduction from Samsun, Asiatic Turkey, in 1905, being designated as S. P. I. No. 15821. Mr. Papathopoulos, from whom the seed was received, obtained it from Trebizond. This importation contained a number of strains, and several other selections were made. It is a 6-rowed bearded barley with heads very similar to those of Coast. At Aberdeen, Idaho, the plants are usually several days earlier and the straw is a little shorter than those of Coast. In threshing, the awns break off from the kernel more easily than those of the Coast variety.

At Aberdeen, Idaho (irrigated), Trebi has produced the highest average yield of any of the barleys grown. It also produced unusually high yields at Scottsbluff (irrigated) and North Platte, Nebr., Newlands, Nev. (irrigated), and Moro, Burns (irrigated), and Union,

Oreg. On the dry lands Trebi has yielded well in years when there was a fair amount of rainfall, but it seems to be especially adapted to irrigated lands. On a farm near Aberdeen, Idaho, a field of 28 acres under irrigation produced a yield of more than 90 bushels to the acre. Trebi appears to do best under such climatic conditions as are found in southern Idaho.

Baker (C. I. No. 975), another selection from the same parent stock, has produced high yields at a few stations.

COAST (C. I. NO. 690)

Coast (C. I. No. 690) was selected for this discussion, but it includes the characteristics of Coast (C. I. No. 626), commonly grown in California. For many years it has been the most important of all varieties grown commercially in the western half of the United States. It has been known as Bay Brewing, Common, California Feed, and Blue. It is of North African origin. It is a 6-rowed bearded variety with short-haired rachilla (Pl. II, B). The head is lax but not so lax as that of Manchuria, and the straw is much shorter. The awns usually emerge earlier and are considerably heavier and rougher than those of Manchuria. The awns contain much less ash than those of Manchuria, which makes them less brittle. The tissues of the lemma are quite tender, and short pieces of awn are left on many kernels in threshing. Often where this occurs the lemma is torn for a short distance below the tip. Such kernels do not pack closely, and the bushel weight of Coast may be low even when the individual kernels are plump and heavy. Coast barley requires a semiarid climate and for that reason is particularly well adapted to the barley-growing sections of the Western States. It is the common feed grain of the West, and varying quantities are exported from California, where the production is far above the feed requirements. The better grades of this barley are in high favor with European brewers. This variety has been widely tested at the experiment stations throughout the West and has yielded exceptionally well at Burns, Oreg. (dry land), and (under C. I. No. 626) at Davis and Chico, Calif. Varieties of this type have been among the better sorts at many of the western stations.

COAST (CALIFORNIA NO. 4000, C. I. NO. 2821)

Coast (California No. 4000) barley is a pedigreed variety developed from Common (Coast) barley at Davis, Calif., about 12 years ago. Its general appearance is the same as that of Coast (C. I. No. 626). From Davis it has been widely distributed in the California barley districts. California No. 4000, according to Prof. G. W. Hendry, University Farm, Davis, Calif., is about 10 per cent more productive than the ordinary commercial Coast barley.

SANDREL (C. I. NO. 937)

Sandrel is a white 6-rowed hulled bearded selection from a bulk importation from Moravia. Since it is a barley of the Coast type, it was probably an accidental mixture. It differs from most of the North African selections in that the aleurone layer is white and not blue. At Archer, Wyo., Aberdeen, Idaho (irrigated), and Columbia, Mo., Sandrel produced very high yields.

BLUE LONG HEAD (C. I. NO. 2685)

The Blue Long Head variety has been widely tested at Canadian experiment farms. It is mentioned in the Canadian Experimental Farms Report of 1896 (p. 17). A specimen seen at Ottawa in 1922 indicated that Blue Long Head belongs to the Coast group of barleys. It is not quite typical of the Coast group, but is of importance in marking the limits of the ecological districts, since it was the only barley grown in Canada which belonged to the Coast group. It was one of the highest yielding varieties at Indian Head, Saskatchewan, while in eastern Canada it produced very low yields. Yields of Blue Long Head at points in the West other than Indian Head were not so high as might be expected.

PERUVIAN (C. I. NO. 935)

Peruvian (C. I. No. 935) is a pedigreed selection made in the cooperative breeding experiments conducted by the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul, Minn. This barley also belongs to the Coast group, but it differs from Coast (C. I. No. 690) in several respects. At Aberdeen, Idaho, in 1921 it had a culm length of 97 cm., which was the longest of all the barleys of the Coast type. The spike, while more erect, was not so dense as Coast or Stavropol, the density being 3.7 mm. The seed length was 11.2 mm., the longest of all the barleys tested at Aberdeen in that year. The length of kernel is one of the conspicuous characters which differentiate it from Coast. Peruvian produced exceptionally high yields at Moro, Oreg., in the years in which it was tested.

BELDI GIANT (C. I. NO. 2777)

This barley undoubtedly belongs to the Coast group and is a variety of the Peruvian type. It was apparently substituted for the original Beldi (C. I. No. 190) at Davis, Calif., and distributed to the stations at Gooding, Aberdeen (Idaho), Bozeman (Mont.), Moccasin (Mont.), and Pullman (Wash.). At Pullman, Davis, and Aberdeen (irrigated) it was the second highest yielding spring barley for the years in which it was grown.

BELDI DWARF (C. I. NO. 190)

The Beldi Dwarf variety was first introduced into the United States from Algeria by the Office of Foreign Seed and Plant Introduction on September 26, 1901. It is a 6-rowed bearded barley with short straw. In Algeria it was grown on the saline soils in the vicinity of the Sahara Desert. It has been grown at most of the western stations, where its region of best adaptation would be expected. Although a good variety it has not proved to be superior at any of the stations where tested. It seems to have done best at Chico, Calif., where it ranked third in point of yield compared with the other varieties grown in the same years.

CALIFORNIA MARIOUT (C. I. NO. 1455)

The California Mariout barley, according to Prof. G. W. Hendry, of the California Agricultural Experiment Station, was secured from the E. Clements Horst Co., of San Francisco, from whose London

office it had been obtained directly in 1905. This variety undoubtedly came from the dry-hill region west of Lake Mariout, Egypt, where many variations were found and brought to America in 1923 by the senior writer. Like Coast (C. I. No. 626) and Club Mariout (C. I. No. 261), this is a 6-rowed bearded barley which originated in northern Africa. It is easily distinguished from these types by its long-haired rachilla, shorter straw, etc.

In a comparison of the California and Club Mariout at Aberdeen, Idaho, in 1921 it was found that the awns of the former emerged three days earlier than those of the latter. The density of California Mariout was 3.7 mm., as compared with 2.6 mm. for Club Mariout. (Pl. II, C and D.) The straw length of the first was only 55 cm., as compared with 76 cm. for the second. Both in appearance and field requirements these barleys are quite different. The California Mariout is drought resistant, early, and well adapted to the warmer, drier interior areas of California. It will outyield the Coast barleys in these limited areas, but the Coast types remain by far the more important commercially.

CLUB MARIOUT (C. I. NO. 261)

Club Mariout (C. I. No. 261) was imported into the United States from Cairo, Egypt, by the Office of Foreign Seed and Plant Introduction, United States Department of Agriculture. In a letter from the Traders and Growers Union, Cairo, Egypt, March 3, 1919, it is stated that the Mariout barley comes from the district west of Alexandria, along the Mediterranean, where it was cultivated in Roman times. In 1923 the senior writer collected a large number of barleys in Egypt. From these samples it would appear that the Club Mariout came from east of the lake, while the California Mariout was cultivated on the dry hills to the west of it. Club Mariout is a 6-rowed bearded variety having straw of medium length and a short compact spike. At Aberdeen, Idaho, in 1921 the average density of Club Mariout was 2.6 mm. This variety is adapted in general to the same conditions as the Coast barleys. It is outstanding in yield at Chico, Calif., Moro, Oreg., and Hays, Kans. At these places it produced the highest average yield of all barleys grown for the five years from 1917 to 1921, inclusive. It is grown commercially in California, where the acreage is increasing, a large part of the crop being exported to Europe for brewing. There is also a limited acreage in Oregon. In most other localities in the West it is surpassed by the Coast varieties.

Club Mariout (C. I. No. 932) is a pedigreed selection of C. I. No. 261 made in the cooperative breeding experiments conducted by the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul, Minn.

FLYNN (C. I. NO. 1311)

Flynn (C. I. No. 1311) is a 6-rowed smooth-awned hybrid made in the cooperative breeding experiments conducted by the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul, Minn. It came from a cross, Club Mariout $\times$ Lion, and was first tested in the nursery at Moro, Oreg., in 1917, where it was selected from several hybrids of the same cross

because of its high yield and good appearance. Of the varieties grown in the field plats at Moro from 1918 to 1921, inclusive, Flynn was exceeded in yield only by Club Mariout and Peruvian. This barley is now being tested at a number of experiment stations where Club Mariout has done well. It now seems to be a very promising sort in these localities. The potential importance of this type of barley is great. The rough awns of the varieties now grown are objectionable, and if smooth-awned sorts which yield well can be produced they will be greatly preferred by farmers.

WINTER CLUB (C. I. NO. 592)

Winter Club (C. I. No. 592) is a bearded 6-rowed barley with a short compact head. (Pl. III, fig. 2, D.) It has been grown commercially for a number of years in Utah, usually under the name Utah Winter or White Winter. It is one of the oldest types of barley known, having been cultivated in western Europe before historical times. It is plausible that the dense 2-rowed barleys came from crosses of this 6-rowed and a lax 2-rowed sort. This variety does not have so pronounced a winter habit as most other winter barleys. At Burns, Oreg., it yielded very well when spring sown. At Pullman, Wash., it also produced fair yields as a spring barley. As a winter barley at Pullman it made the highest average yield of all varieties grown in the same years.

TENNESSEE WINTER (C. I. NO. 257)

Tennessee Winter (C. I. No. 257) is the dominant variety among the winter barleys. Its origin is not known. The parent stock has long been grown in Tennessee. Somewhat similar varieties have been received from Switzerland and the Balkans. It has been tested at a number of the experiment stations and is found to be the most frost resistant of all winter barleys. It is a 6-rowed bearded sort. While this barley seems particularly well adapted to the growing conditions of Tennessee, it has also produced high yields at Arlington Experiment Farm in Virginia, at Athens, Ga., and at State College, N. Mex. At Wooster, Ohio, and Blacksburg, Va., the highest yielding winter barleys were of the Tennessee Winter group.

ARLINGTON AWNLESS (C. I. NO. 702)

Arlington Awnless (C. I. No. 702) was developed by selection from a hybrid made between Tennessee Winter (C. I. No. 257) and Black Arabian (C. I. No. 202). This variety was originated by H. B. Derr, formerly of the Office of Cereal Investigations of the United States Department of Agriculture. It is a 6-rowed awnless hulled barley with a lax spike. Because of its earliness and awnless character it was thought it would become an important winter barley. The rachis, however, is very brittle, which causes much loss of grain by shattering. It is inferior to Nakano Wase (C. I. No. 754), another awnless variety. The latter is a Japanese barley which is quite winter hardy, but does not yield well enough to warrant extensive culture.

FIG. 1.—A, Nepal (C. I. No. 595), the original hooded variety, parent of all hooded forms, much inferior to hulled awned varieties except in a few mountain sections of the West; B, Horsford (C. I. No. 507), a variety of hybrid origin typical of those (usually Nepal) crossed on varieties of the Manchuria group; C, Mcloy (C. I. No. 1176), a hooded hulled variety from Oregon, probably from a cross of Nepal on Coast, shattering less than Horsford; D, Nakano Wase (C. I. No. 754), an awnless winter form from Japan

FIG. 2.—A, Baku (C. I. No. 709), a naked variety, common in dry mountain plateaus of southern Asia, not commercially important in America; B, Blackhull (C. I. No. 878), a deficient 2-rowed variety, promising in Colorado; C, Himalaya (C. I. No. 620), a naked variety from the same region as Baku, the best-yielding naked variety in America; D, Winter Club (C. I. No. 592), one of the old winter varieties once common in Europe, grown occasionally in the Great Basin

SIX-ROWED HULLED HOODED BARLEYS

MELOY (C. I. NO. 1176)

The Meloy (C. I. No. 1176) barley is evidently a natural hybrid. It is a 6-rowed hooded hulled variety which was found in the field of George Meloy, Moro, Oreg. In general appearance it is like the ordinary Horsford except that the hoods are more elevated. (Pl. III, fig. 1, C.) At Aberdeen, Idaho, in 1921 Meloy had a culm length of 97 cm., as compared with 114 cm. for Horsford (C. I. No. 507). The average density was 3.3 mm., which was almost the same as that of Horsford. As the latter barley has been a low yielder in practically all localities, it is of interest to note that Meloy has produced very high yields at Moro, Oreg.

The Horsford varieties have all exhibited the defect of shattering badly. The awn of barley contains as high as 32 per cent of ash. When the awn is removed by crossing, the awnless hybrids of the progeny shatter easily, probably owing to an excess of ash in the rachis. The rachis of the Coast barleys contains much less ash than that of the Manchuria types, and it is probable that less ash is absorbed by the Coast plants. Crosses with Nepal on Coast should give hooded hybrids less brittle than when Nepal is crossed on Manchuria. Meloy is thought to have resulted from a hybrid of Nepal and Coast and shatters less than the progeny of Manchuria-Nepal crosses.

HORSFORD (C. I. NO. 507)

The Horsford variety was originated by F. H. Horsford, of Charlotte, Vt., for whom it was named. In 1879 or 1880 Mr. Horsford crossed a Nepal on a 6-rowed bearded hulled barley and obtained a 6-rowed hooded hulled variety which has proved constant for these characters. C. I. No. 507 (Pl. III, fig. 1, B), however, has been given to a number of the barleys of this description that are not selections or pedigreed strains of this type. A number of these undoubtedly did not come from Mr. Horsford's cross, but they are of the same general type, having been produced by crossing Nepal with barleys of the Manchuria type. This variety is grown extensively in the United States and has been tested at most of the experiment stations throughout the country. Like Nepal, however, it is a low-yielding sort, as compared with the better varieties of a region, and has the additional defect of shattering when ripe. It is grown commercially under such names as Beardless, Great Beardless, and Success. Although it is not hull-less, in general appearance it is quite like the Nepal barley, but the head is not so dense, the hoods are not usually so sessile, and the straw is longer. The beardless character of both of these barleys makes them desirable, because of the greater ease with which they may be handled. Nearly all seedsmen carry a Horsford barley under some commercial name, and farmers are led by their descriptions to grow these barleys in entirely unsuitable localities.

SIX-ROWED HULLED AWNED BLACK BARLEYS**LION (C. I. NO. 923)**

Lion (C. I. No. 923) is a pedigreed selection from S. P. I. No. 32767 made in the cooperative breeding experiments conducted by the United States Department of Agriculture and the Minnesota Agricultural Experiment Station at St. Paul, Minn. The original barley from which this selection was made was first found at the agricultural experiment station in Taganrog, southern Russia. It was obtained from the experiment station near Rostov on Don through the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture, in December, 1911. Lion is a 6-rowed black smooth-awned variety. It has been used as one of the parents in a number of crosses in the hope of producing a high-yielding smooth-awned hybrid. While it has produced good yields at several of the experiment stations, it has no present commercial importance. At the Michigan Agricultural Experiment Station this barley was called Michigan Black Barbless and was distributed under that name. Apparently the Lion sent to the Michigan station from Minnesota had become mixed, as more than one type was selected at Michigan. It is the opinion of the writers that the strain finally chosen for distribution was identical with Lion, as they have been able to find no differences between the two.

GATAMI (C. I. NO. 575)

Gatami (C. I. No. 575) barley was imported into the United States from Mukden, Manchuria, by the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture in April, 1907. At many of the experiment stations in earlier years it was grown under the Chinese name Gai ta mi or some variation of this spelling. It is a black-bearded 6-rowed barley having the same general growth requirements as Manchuria and Odessa. While it has been widely tested in the United States, it is usually surpassed in yield by Manchuria and Odessa when grown at the same stations. It is a very early variety, ripening several days before Manchuria or Odessa at most points. For this reason it has proved to be of value in localities in South Dakota where hot dry weather occurs early in the summer.

SIX-ROWED NAKED BARLEYS**NEPAL (C. I. NO. 595)**

Nepal is one of the older types of barley grown in the United States. It is thought to have originated in Nepal, India, where it was first noticed by Europeans in 1839. The number C. I. No. 595 (Pl. III, fig. 1, A) has been given to most of the 6-rowed naked hooded barleys that are not selections or pedigreed strains. This variety has been tested at one period or another at nearly all the experiment stations in the United States. It has also been grown under many names at these stations, the more common ones being White Hull-less, Giant White Hull-less, Ideal Hull-less, and Colossal Hull-less. Because it has no awns and is hull-less it is not unpleasant to handle and is very easily threshed. These qualities make it a desirable type of

barley. At none of the stations, however, has Nepal produced high yields. At most of them it was among the lowest in yield. Many crosses have been made with other types of barley with the hope of producing high-yielding hooded varieties, but with few exceptions these have not proved worthy of note. Crosses with barleys of the Manchuria type have resulted in progeny in which the rachis is so brittle that large losses occur from shattering.

HIMALAYA (C. I. NO. 620)

Several importations of barleys of the Himalaya type have been tested on experimental farms. Others have most likely been grown by farmers without having found their way to the stations. In all cases the type is quite similar. (Pl. III, fig. 2, C.) This blue naked 6-rowed bearded barley is quite common in central Asia and occurs at the higher altitudes in northern India. It has given higher yields than other naked sorts and is found in occasional fields throughout the barley-producing sections of the western mountain region of the United States.

BLACK HULL-LESS (C. I. NO. 596)

The Black Hull-less variety is widely distributed over central Asia. Its kernels when grown under semiarid conditions are brilliant purple at harvest time. As they dry, this purple darkens until the kernels are almost black in appearance. It is a 6-rowed bearded barley, like Himalaya, and has been tested and grown in the same general regions, but it has not proved to be the equal of that variety in yield.

ACCESSION DATA OF CEREAL INVESTIGATION NUMBERS

The barley varieties appearing in this bulletin have been accessioned under Cereal Investigations (C. I.) numbers. In a great many cases these varieties were received from foreign countries through the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture. In these cases the C. I. number is the one under which they were first grown by the Office of Cereal Investigations and distributed to the experiment stations. In many other cases the varieties were received from State experiment stations and accessioned. Many of the varieties have not been received by the department, especially those of Canadian stations. As they often were grown at several stations under different station numbers, it was necessary to accession these as well.

In Table 94 the data occurring under the Cereal Investigations numbers are abridged in tabular form. The list of station numbers and synonyms must necessarily be incomplete. An attempt is made to indicate the character of the barley by stating that it belongs in a certain group. Such grouping is necessarily approximate. For example, the Chevalier barleys merge into the Hanna types by way of the Bohemian group, which has some of the characters of each. There is some confusion as to whether certain varieties originated in Austria or Germany. Conditions in these countries are so similar, however, that it is immaterial which is credited as the place of origin.

TABLE 94.—Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of caryopsis					
3	Plumage	Sterile	Hulled	Awed	White		Thorpe	England	Various		Plumage from Nor- way.
17	Pern	do	do	do	do			Russia	Russia		
20	do	Fertile	Naked	Hooded	do			India	New York		
22	Nepal	Sterile	Hulled	Awed	do		Nepal	England	Russia		
23	Chevalier	Sterile	do	do	do		Chevalier	Germany	Various		
24	Hanna	do	do	do	do		Hanna	do	do		
25	Scale	do	do	do	do		do	do	do		
26	Hanna	do	do	do	do		do	Austria	do		
27	Bohemian	do	do	do	do		do	Bohemia	do		
28	Hanna	do	do	do	do		do	Germany	do		
29	Basthorn	Im- perial	do	do	do		Thorpe	do	do		
30	Hanna	do	do	do	do		Hanna	Germany	do		
31	Hanna	do	do	do	do		Chevalier	England	do		
32	Bohemian	do	do	do	do		Hanna	Bohemia	do		
33	Hanna	do	do	do	do		do	Germany	do		
34	do	do	do	do	do		do	Plant selection	do		
35	Chevalier	do	do	do	do		do	England	do		
36	Corinth	Fertile	Naked	do	do		Chevalier	Central Asia	Mexico		
39	Rokaku Cheva- lier	Sterile	Hulled	do	do		Chevalier	England	Japan		
45	Mishima	Fertile	Naked	do	do			Japan	do		
46	Sonbatu	Sterile	Hulled	do	do			do	do		
47	Strigum	Sterile	do	do	do		Hanna	Germany	do		
48	Golden Melon	do	do	do	do		Thorpe	England	do		
49	Nifo	Fertile	Naked	do	do			Japan	do		
50	Tunkaze	do	Hulled	do	do			do	do		
52	Sontake	do	do	do	do			do	do		
54	Shiro-Nishiki	do	do	do	do			do	do		
55	Tsutsuzaki	do	Naked	do	do			do	do		
56	Nyotake	do	do	do	do			do	do		
60	Nyotake	Sterile	do	do	do		Baku	do	do		
62	Daisen	Fertile	Hulled	do	do			do	do		
64	Baifu	do	do	do	do			do	do		
78	Sonotsuka	do	Naked	do	do			do	do		
80	Mochi-Mugi	do	do	do	do			do	do		
84	Kostrona	Sterile	Hulled	do	do			Russia	Russia	North Dakota No. 327.	

TABLE 94.—Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of caryopsis					
200	Chevalier II (Pl. I, A).	Sterile..	Hulled..	Awned..	White..	Lax..	Chevalier..	Plant selection..	Sweden..	North Dakota No. 1398.	
201	Kizing..	do..	do..	do..	do..	do..	Hanna..	Germany..	Germany..		
202	Black Arabian..	do..	do..	do..	Black..	do..	do..	Asia Minor..	Arabia..		
203	Hanna..	do..	do..	do..	White..	Lax..	Hanna..	Moravia..	do..	North Dakota No. 649.	Moravian.
204	Bohemian..	do..	do..	do..	do..	do..	Bohemian..	Bohemia..	Austria..		
205	Yenidje..	Fertile..	do..	do..	do..	do..	do..	Asia Minor..	Asia Minor..		
206	Han River..	do..	do..	do..	do..	do..	do..	China..	Han River, China..		
207	Frankish..	Sterile..	do..	do..	do..	Lax..	Hanna..	Germany..	Bavaria..		
208	Bavarian..	do..	do..	do..	do..	do..	do..	do..	do..		Hanna, Lower Frankish.
209	Highland Chief..	do..	do..	do..	do..	do..	Thorpe..	England..	Various..		
210	Baku..	do..	Naked..	do..	do..	Dense..	Baku..	Central Asia..	McPherson, Kans..		
211	Success..	Fertile..	Hulled..	Hooded..	do..	do..	Hybrid..	Nepal X..	Utah..		
212	Norway 8..	do..	do..	Awned..	do..	do..	do..	Norway..	Various..		
213	Nerchinsk..	do..	Naked..	do..	do..	do..	do..	Central Asia..	Siberia..		
214	Mammoth Win- ter.	do..	Hulled..	do..	do..	do..	Tennessee Winter.	Europe..	Russia..		
221	Greece..	do..	do..	do..	do..	do..	Greece..	Greece..	Greece..		
223	Argentine..	do..	do..	do..	do..	Blue..	do..	Argentina..	Argentina..		Ideal Hull-less.
220	Nepal..	do..	Naked..	Hooded..	White..	do..	Nepal..	India..	Iowa..		
231	Onto Beardless..	do..	Hulled..	do..	do..	do..	Hybrid..	Nepal X..	Ohio..		Golossal Hull-less.
230	Nepal..	do..	do..	do..	do..	do..	Nepal..	India..	Michigan..		Manshury.
233	Manchuria..	do..	Hulled..	Awned..	do..	do..	Manchuria..	Manchuria..	Massachusetts..		Manchschuri.
232	do..	do..	do..	do..	do..	do..	do..	do..	Various..		Manshury.
237	do..	do..	do..	do..	do..	do..	do..	do..	Kansas..		Do.
240	do..	do..	do..	do..	do..	do..	do..	do..	Various..		
241	do..	do..	do..	do..	do..	do..	do..	do..	do..	Minnesota No. 6, Wisconsin No. 62, North Dakota No. 787.	
244	Manchuria (Pl. II, A).	do..	do..	do..	do..	do..	do..	do..	do..	Minnesota Nos. 32 and 105, North Da- kota No. 788.	Manshury, Man- shury, Mand- scheuri.
248	Bavarian..	Sterile..	do..	do..	do..	do..	Hanna..	Germany..	Bavaria..		
252	Squarehead Win- ter.	Fertile..	do..	do..	do..	Lax..	do..	Europe..	France..		

253	Baku	Sterile	Naked	do	do	Blue	do	Baku	Central Asia	do	do	Two-Rowed Hull-less, Himalaya Hull-less.
254	Himalaya	Fertile	do	do	do	do	do	Himalaya	do	do	do	Himalaya Hull-less.
255	Fan	Sterile	Hulled	do	do	White	do	Zeeriton	Asia	Various	do	Rice, Fischer, Peacock.
256	Black	Fertile	do	do	do	Black	do	Tennessee Winter	Europe	France	do	Maryland Winter, Texas Winter.
257	Tennessee Winter	do	do	do	do	White	do	Arctic	Various	do	do	
259	Early Lapland	do	do	do	do	do	do	Club Marabout	Finland	France	do	
261	(Pl. II, C).	do	do	do	do	do	do	Club Marabout	Egypt	Egypt	do	
262	Nepal	do	Naked	Hooded	do	do	do	Nepal	India	Colorado	do	Myer Hull-less.
263	Garton	do	Hulled	Awne	do	do	do	do	England	England	do	
264	do	do	do	do	do	do	do	do	do	do	do	
265	do	Sterile	do	do	do	do	do	do	do	do	do	
266	do	do	do	do	do	do	do	Thorp	do	do	do	
267	Chevalier	do	do	do	do	do	do	Chevalier	do	Bavaria	do	
268	Webs	do	do	do	do	do	do	Hanna	Germany	Bavaria	do	
269	Frankish	do	do	do	do	do	do	Bohemian	Czechoslovakia	do	do	
270	Bohemian	do	do	do	do	do	do	Hanna	Germany	do	do	
271	Gold Medal	do	do	do	do	do	do	do	do	France	do	
272	Kitzing	do	do	do	do	do	do	do	do	do	do	
274	Langer	do	do	do	do	do	do	do	do	do	do	
275	Kwasitzer	do	do	do	do	do	do	do	do	do	do	
276	Coast	Fertile	do	do	do	do	do	Coast	Plant selection	U. S. D. A., Chico, Calif.	do	
277	Scottish Pearl	do	do	do	do	do	do	Tennessee Winter	Europe	U. S. D. A., Arlington Farm, Va.	do	Tennessee Winter.
278	Chevalier	Sterile	do	do	do	do	do	Chevalier	England	France	do	
279	Fuinettener	do	do	do	do	do	do	Hanna	Germany	do	do	
282	Frankish	do	do	do	do	do	do	do	do	do	do	
284	Bohemian	do	do	do	do	do	do	Bohemian	Czechoslovakia	do	do	
285	do	do	do	do	do	do	do	do	do	do	do	
287	Hanna	do	do	do	do	do	do	Hanna	Germany	do	do	
289	Imperial	do	do	do	do	do	do	Thorp	do	do	do	
290	Chevalier	do	do	do	do	do	do	Chevalier	England	do	do	
293	Bohemian	do	do	do	do	do	do	Bohemian	Czechoslovakia	do	do	
295	Frankish	do	do	do	do	do	do	Hanna	Germany	do	do	
301	Hanna	do	do	do	do	do	do	do	do	do	do	
302	Langer	do	do	do	do	do	do	do	do	do	do	
314	Poppenhelm	do	Naked	do	do	do	do	do	do	do	do	
317	Prague	do	Hulled	do	do	do	do	Baku	Central Asia	Czechoslovakia	do	Austrian Hanna.
319	Hanna	do	do	do	do	do	do	Hanna	do	do	do	
320	Crimean Hull-less.	do	Naked	do	do	do	do	Baku	North Dakota No. 1326.	do	do	
322	Himalaya	Fertile	do	do	do	Blue	do	Baku	Central Asia	Germany	do	Baku.
323	Victoria	Sterile	Hulled	do	do	White	do	Himalaya	Germany	do	do	
324	Jewel	do	do	do	do	do	do	Thorp	Western Europe	Various	do	

TABLE 94.—Accession data of varieties of barley arranged in the order of their *Cereal Investigations numbers*—Continued

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of carypsids					
325	Phoenix Von Thalen.	Sterile..	Hulled..	Awed..	White..	Lax..	Hanna Thorpe..	Germany..	Various..		
326	Cheney..	do..	do..	do..	do..	do..	do..	do..	Germany..		
327	Goldthorpe..	do..	do..	do..	do..	Dense..	do..	do..	Canada..		
328	Winter Six- Rowed.	Fertile..	do..	do..	do..			England..	Maryland..		
329	Manchuria..	do..	do..	do..	do..		Manchuria..		Germany..		
330	Gambinus..	Sterile..	do..	do..	do..	Lax..	Hanna..	Germany..	Various..		
331	Beschnorn..	do..	do..	do..	do..	do..	Chevalier..	England..	Canada..		
332	Kaiser..	do..	do..	do..	do..	do..	Hanna..	Germany..	Germany..		
333	Baku..	do..	Naked..	do..	do..	do..	Baku..	Central Asia..	Canada..		
334	Trifurcate Hull- less.	do..	do..	Hooded..	do..		Hybrid..	Nepal × Distichon	Germany..		Large Two-Rowed Naked.
335	Coelste..	Fertile..	do..	Awed..	do..			Central Asia..	do..		
336	Greek..	do..	Hulled..	do..	do..			Greece..	Various..		
337	Black Winter..	do..	do..	do..	Black..			Germany..	Germany..		Grecian Six-Rowed.
338	Rotbraun..	Sterile..	do..	do..	White..	Lax..	Hanna..	Germany..	Various..		
339	Sholey Cheva- lier.	do..	do..	do..	do..	do..	Chevalier..	England..	Germany..		
341	Fan..	do..	do..	do..	do..	Dense..	Zecriton..	Asia..	Various..		Rice, Fächer, Pea- cock. Italian Compact.
342	Italian..	do..	do..	do..	do..		Thorpe..	England..	do..		
343	Moravian..	do..	do..	do..	do..	Lax..	Hanna..	Czechoslovakia..	Germany..		
344	Bohemian..	do..	do..	do..	do..	do..	Bohemian..		France..		
345	Schwarzenberg I..	do..	do..	do..	do..	do..	Hanna..	Germany..	do..		
346	Schwarzenberg III..	do..	do..	do..	do..	do..	do..	do..	do..		
347	Voronezh..	do..	do..	do..	do..	do..	do..	do..	do..		
348	Orel..	do..	do..	do..	do..	do..	Orel..	Russia..	Russia..		
349	Nepal..	Fertile..	Naked..	Hooded..	do..	do..	Nepal..	Orel, Russia..	Washington..		
350	Abyssinia..	do..	do..	Awed..	do..	do..	Abyssinia..	Abyssinia..	Abyssinia..		Mexican Beardless.
351	Centre..	do..	Hulled..	do..	do..	Lax..	Abyssinia..	do..	do..		
352	Brasserie..	Sterile..	do..	do..	do..		Hanna..	Germany..	France..		
353	Large-Grained	do..	do..	do..	do..			Hungary..	Hungary..		
354	Winter.	Fertile..	do..	do..	do..			Europe..	Germany..		
409	Goldthorpe	Sterile..	do..	do..	do..	Dense..	Thorpe..	England..	St. Louis, Mo..		
410	Mammoth Win- ter.	Fertile..	do..	do..	do..		Tennessee Winter.	Europe..	Germany..		

411	Golden Melon...	Sterile	do	do	do	Lax	Chevalier	England	Various	North Dakota
414	Besthorn Giant...	Fertile	do	do	do	Lax	Hanna	Europe	Germany	Nos. 147, 378,
416	Hanna	Sterile	do	do	do	do	do	Germany	do	and 720.
434	Catania	do	do	do	do	do	do	do	Sicily	
442	Aquila	Fertile	do	do	do	do	do	do	Italy	
455	Caveda	do	do	do	do	do	do	do	Portugal	
468	Quebec	do	do	do	do	do	do	do	Mexico	
488	Winter Club	do	do	do	do	do	Winter Club, Hybrid	North Africa	Lewiston, Idaho	
507	Horsford (Pl. III, fig. 1, B).	do	do	Hooded	do	do	do	Nepal X	Various	
516	Phoson	do	Naked	Awne	do	do	Tennessee	India	India	
518	Maryland Winter	do	Hulled	do	do	do	Winter	Europe	Maryland	
519	Wisconsin Winter	do	do	do	do	do	Wisconsin	do	Various	
521	Bulgarian	do	do	do	do	do	do	Bulgaria	Bulgaria	
528	Kashgar Hull-less	do	Naked	do	do	do	do	Asia	Kashgar	
529	Princess	Sterile	Hulled	do	do	Lax	Plant selection	Plant selection	Svalof, Sweden	
531	Hannchen	do	do	do	do	do	Hanna	do	do	North Dakota
532	Primus	do	do	do	do	Dense	Thorpe	do	do	No. 1397.
533	Nepal	Fertile	Naked	Hooded	do	do	Nepal	India	Geneva, Idaho	Minnesota No.
534	Alaska	do	Hulled	Awne	do	do	Manchuria	Plant selection	Fairbanks, Alaska	150, North Da-
537	Oderbrucker	do	do	do	do	do	Tennessee	Europe	Madison, Wis.	kota No. 1399.
539	Terwin	do	do	do	do	do	Coast	North Africa	Channing, Tex.	Minnesota No.
549	Coast	do	do	do	do	do	Winter	Europe	Utah	151, North
554	Texas Winter	do	do	do	do	do	Tennessee	do	Texas	Dakota No.
557	Cape	do	do	do	do	do	Winter	do	State College, Pa.	1400.
558	Golden Queen	do	do	do	do	do	Manchuria	Manchuria	Various	Boehme's Hull-less.
559	Robertson	Sterile	do	do	do	Lax	Chevalier	England	Scotland	
575	Gatami	Fertile	do	do	Black	do	do	Manchuria	Manchuria	North Dakota
577	Kamamugi	do	Naked	do	White	Dense	do	Japan	Japan	Nos. 258 and
578	Tanbashi	do	do	do	do	do	do	do	do	791.
579	Wakamatsu	do	do	do	do	do	Tennessee	do	do	Minnesota No.
583	Union Winter	do	Hulled	do	do	do	Winter	Europe	Knoxville, Tenn.	401.
584	Standwell	Sterile	do	do	do	Dense	Hybrid	Fan X Melon	Garton Bros., Eng-land.	Tanbashi Hull-less.

TABLE 94.—Accession data of varieties of barley arranged in the order of their *Cereal Investigations numbers*—Continued

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of earcopsis					
585	Maltster.	Sterile	Hulled.	Awned	White		Hybrid	(Jewel X Italian) X (Chevalier X Irishable).	Garton Bros., England.	Minnesota No. 364.	
586	Brewers Favor- ite.	do	do	do	do		Thorpe	England	do		
587	Kinver Chevalier	do	do	do	do		Chevalier	do	Various		
588	Golden Grain	do	do	do	do		do	do	do		
590	Invineible.	do	do	do	do		Hybrid	(Chevalier X Gol- den Moh) X Standwell.	Garton Bros., Eng- land.		New Golden Grains.
592	Winter Club (Pl. III, fig. 2, D).	Fertile	do	do	do		Winter Club.	Europe	Various		
595	Nepal (Pl. III, fig. 1, A).	do	Naked.	Hooded	do		Nepal	India	do		Utah Winter-Tapp Winter. White
596	Black Hull-less	do	do	Awned		Purple	Black Hull-less.	Central Asia	do	Minnesota No. 332, North Dakota Nos. 392 and 848.	Giant White Hull- less, White Hull- less.
606	Gutekorn	Sterile	Hulled	do	do			Plant selection	Svalof, Sweden	Minnesota No. 207.	
607	Sexradigt	Fertile	do	do	do			do	do	Minnesota No. 380.	Yatte Sexradigt.
610	Horsford	do	do	Hooded	do			Nepal X	Dallas, Tex.		Texas Beardless.
611	Blue Ribbon	do	do	Awned	do		Hybrid	Manchuria	Various	Minnesota No. 196.	
612	Chevalier	Sterile	do	do	do		Chevalier	England	Montana		
613	Manchuria	Fertile	do	do	do		Manchuria	Manchuria	do		
616	Fancsh- date.	Inferme	do	Awless	do		Awless	China	China		
617	Mansury	Sterile	do	Awned	do			Central Asia	Various	North Dakota No. 172	
619	Himalaya	Fertile	Naked	do		Blue	Himalaya	Central Asia	do	Wyoming No. 46	Guy Mayle.
620	Himalaya (Pl. III, fig. 2, C).	do	do	do		do	do	do	do	Minnesota No. 201.	
626	Coast	do	Hulled	do	White		Coast	North Africa	do		California Feed, Bay Brewing.

TABLE 94.—Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adherence	Appendages to central florets	Color of lemma	Color of caryopsis					
702	Arlington Awnless.	Intermediate.	Hulled.	Awnless.	White.		Hybrid.	Tennessee Winter X Black Arabian.	U. S. D. A. Arlington Farm, Va.		
703	Hanseer Hull-less.	Fertile.	Naked.	Awed.	do.		do.	Horsford X Hanseer.	do.		
704	Popeline.	do.	Hulled.	do.	Black.			Caucasus.	Caucasus.		
705	Black Russian.	do.	do.	do.	do.			do.	do.		
708	Black Algerian.	do.	do.	Smooth awed.	do.			do.	Algeria.		
709	Baku (Pl. III, fig. 2, A).	Sterile.	Naked.	Awed.	White.		Baku.	Central Asia.	Caucasia.		
711	Turkestan.	Fertile.	Hulled.	do.	do.			Turkestan.	Turkestan.		
712	do.	do.	do.	do.	do.			do.	do.		
713	Canada Winter.	do.	do.	do.	do.		Tennesseer.	Europe.	Guelph, Ontario.		
714	Caucasian.	Sterile.	do.	do.	do.			Caucasia.	Caucasia.		
731	Pontius.	Mixed.	do.	do.	do.			China.	China.		
735	Black Hull-less.	Fertile.	Naked.	Awed.		Purple.		Central Asia.	Chinese Turkestan.		
737	Nilver.	Fertile.	Hulled.	Awed.	White.		Manchuria.	Plant selection.	Ottawa, Canada.	Minnesota No. 303.	
739	Manchurian.	Fertile.	do.	do.	do.		Thorpe.	England.	do.		
740	Canadian Thorpe.	Sterile.	do.	do.	do.		Awless.	Japan.	Japan.		
754	Nakano Wase (Pl. III, fig. 1, D).	Intermediate.	do.	Awless.	do.						
835	Wisconsin Pedigree.	Fertile.	do.	Awed.	do.		Manchuria.	Manchuria.	Wisconsin.	North Dakota No. 2406.	Manchuria.
836	Oderbrucker.	do.	do.	do.	do.		do.	do.	Various.	North Dakota No. 2406.	Oderbrucker.
848	Scotch.	do.	do.	do.	do.		do.	do.	do.		
869	Kopeck.	do.	do.	do.	do.		do.	do.	do.		
877	Horsford.	do.	do.	Hooded.	do.		do.	North Africa.	Bolivia.		
878	Blackhull (Pl. III, fig. 2, B).	Deficient.	do.	Awed.	Black.		Hybrid.	Nepal X Abyssinia.	Various.		

881	Bernard	Fertile	do	do	White	Manchu- ria.	Manchuria	do	Minnesota No. 28, Wisconsin No. 63, North Dakota No. 789
882	Manchuria	do	do	do	do	do	do	do	Common.
883	Highland Chief	Sterile	do	do	do	Thorpe	England	do	North Dakota No. 966, Williston Nos. 61 and 170, North Dakota No. 847, Williston No. 63
888	Oderbrucker	Fertile	do	do	do	Manchu- ria, Hybrid	Manchuria	Minneapolis, Minn.	Triumph.
889	Reed Triumph	do	do	do	do	Manchu- ria.	Highland Chief X Manchuria.	Various	
890	Silver King	do	do	do	do	Manchu- ria.	Manchuria	do	Minnesota No. 361, Wisconsin No. 98
891	Success	Sterile	do	do	do			do	Williston No. 50, North Dakota No. 171, Williston No. 171, North Dakota No. 149
892	Evans	do	Naked	do	do	Baku	Central Asia	North Dakota	Salzer's Silver King.
893	Proskowetz	do	Itulled	do	do	Hybrid	Czechoslovakia	Kassowitz (Aus- tria)	McEvan, McEwan, Baku.
895	Cusado	Fertile	do	do	do	Lax	Tennessee Win- ter X Black Arabian.	U. S. D. A., Col- lege Park, Md.	
896	Arabel	do	do	do	do	do	do	do	
898	Omar	Sterile	do	do	do	do	do	do	
901	Pidor	Fertile	do	do	do	do	Tennessee Win- ter X Hankow.	do	
902	Valkow	do	do	do	do	do	do	do	
905	Silver King	do	do	do	do	Manchu- ria.	Plant selection	do	
906	Hanna	Sterile	do	do	do	Hanna	do	Madison, Wis.	Manchuria.
907	Steigum	do	do	do	do	Lax	do	do	Wisconsin Pedigree 13.
908	Luth	Fertile	do	do	do	do	do	do	Minnesota No. 422
909	Clyde	Sterile	Naked	do	do	Manchu- ria.	Central Asia	U. S. D. A., St. Paul, Minn.	Minnesota No. 424
910	White Smyrna	do	Itulled	do	do	Lax	Plant selection	do	Minnesota No. 421
911	Featherston	Fertile	do	do	do	Manchu- ria.	do	do	
							Moccasin, Mont. U. S. D. A., Aber- deen, Idaho.	U. S. D. A., St. Paul, Minn.	Minnesota No. 423

TABLE 94.—Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of caryopsis					
912	Mahrliche	Sterile	Hulled	Awned	White	Lax	Hanna	Plant selection	U. S. D. A., St. Paul, Minn.	Minnesota No. 427	
913	Eagle	Fertile	do	do	do		Manchu- ria	do	do	Minnesota No. 425	
914	Italian	Sterile	do	do	do	Lax	Hanna	do	do	Minnesota No. 426	
915	Servian	Fertile	do	do	do			do	do		
916	Odessa	do	do	do	do	Lax	Odessa	do	do		
917	Prentice	Sterile	do	do	do		Chevalier	England	Ottawa, Canada		Danish Prentice Chevalier.
918	Korsbyg	Fertile	do	do	do			Plant selection	Denmark	Wisconsin No. 97	
920	White Gatami	do	do	do	do			do			
921	Thorpe	Sterile	do	do	do	Dense	Thorpe	Plant selection	U. S. D. A., Dick- inson, N. Dak.		
922	Bolton	do	do	do	do	do	Hybrid	do	Moessin, Mont.		
923	Lion	Fertile	do	Smooth awned	Black			do	U. S. D. A., St. Paul, Minn.	North Dakota No. 31001	Michigan Black Barbless.
925	Australian White.	Sterile	do	Awned	White	Lax	Hanna	do	do		
926	Horn	do	do	do	do			do	do		
927	Odessa	Fertile	do	do	do	do	Chevalier	do	do		
928	Goldfoil	Sterile	do	do	do	Lax	Odessa	do	do		
929	Summit	Fertile	do	do	do		Hanna	do	do		
931	Steinrum	Sterile	do	do	do	Lax	Hanna	do	do		
932	Club Marout	Fertile	do	do	do		Club Ma- rout	do	do		
933	Bohemian	Sterile	do	do	do	Lax	Hanna	do	do		
934	Odessa	Fertile	do	do	do		Odessa	do	do		
935	Peruvian	do	do	do	do	Blue	Coast	do	do		
936	Trebi	do	do	do	do	do		do	do		
937	Sandral	do	do	do	do	do	Coast	do	do		
938	Abyssinian	Deficient	do	do	White		Deficients.	do	U. S. D. A., Chi- co, Calif.		
939	Palestine	Sterile	do	do	do			do	U. S. D. A., St. Paul, Minn.		
940	Oderbrueker	Fertile	do	do	do		Manchu- ria	do	do		

941	Abyssinian	Sterile	Hulled	Awned	White		Lax	Hanna	Plant selection	U. S. D. A., Chi- co, Calif.
942	Hanna	Fertile	do	do	do			Abyssinia	do	U. S. D. A., St. Paul, Minn.
943	Abyssinian	do	do	do	do			do	do	U. S. D. A., Chi- co, Calif.
946	do	do	do	do	do			do	do	U. S. D. A., St. Paul, Minn.
950	do	do	do	do	do			Hanna	do	U. S. D. A., Chi- co, Calif.
952	Holland	Sterile	do	do	do		Lax	do	do	U. S. D. A., St. Paul, Minn.
953	Frankish	do	do	do	do		do	do	do	do
954	Featherston	Fertile	do	do	do			Manchu- ria	do	do
956	Manchuria	do	do	do	do			do	do	do
957	Oderbrucker	do	do	do	do			Thorpe	do	Japan
958	Golden Melon	Sterile	do	do	do		Dense	Hybrid	do	U. S. D. A., St. Paul, Minn.
959	Alpha (Pl. I, C)	do	do	do	do			do	Champion of Vermont	do
962	Scholey	do	do	do	do		Lax	Chevalier	Plant selection	do
972	Luth	Fertile	do	do	do			Manchu- ria	do	do
973	Red River	do	do	do	do			Trebi	do	do
975	Baker	do	do	do	do	Blue		Hybrid	do	do
976	Cortney	do	do	do	White			Hanna	6-row X 2-row	St. Paul, Minn.
977	White Moravian	Sterile	do	do	do		Lax	do	Plant selection	U. S. D. A., St. Paul, Minn.
982	Hodge	Fertile	do	do	do			do	Turkestan	Mexico
986	Woodrow	do	do	do	do			do	Russia	Russia
994	Petro	do	do	do	do			do	do	do
996	Raspit	Fertile	Hulled	Awned	White			do	do	do
1030	Gisborne	Sterile	do	do	do		Lax	Hanna	do	do
1106	Black Hull-less	Fertile	Naked	do	do	Purple		do	do	Minnesota No. 302
1107	Cartouch	do	Hulled	do	do	Blue		do	do	Minnesota No. 402
1116	Tartar	do	do	do	White			Trebi	do	do
1117	Khayyam	Sterile	do	do	do			Hybrid	Tennessee Winter X Black Ara- bian	U. S. D. A., Ross- lyn, Va.
1118	Featherston	Fertile	do	do	do			do	Plant selection	U. S. D. A., St. Paul, Minn.
1119	do	do	do	do	do			Manchu- ria	do	do
1120	do	do	do	do	do			do	do	do
1121	Hanchamont	Sterile	do	do	do			Hybrid	Hanna X Chan- pion of Vermont	do
1122	Hanna	do	do	do	do		Lax	Hanna	Plant selection	do
1123	Cortile	Fertile	do	do	do			Hybrid	6-row smooth X 2-row Indian Head	do

Mexican.

TABLE 94.—*Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued*

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adherence	Appendages to central florets	Color of lemma	Color of caryopsis					
1125	Manchuria	Fertile	Hulled	Awned	White		Manchuria	Plant selection	U. S. D. A. St. Paul, Minn.		
1129	Making	do	do	do	do		do	do	do		
1142	Chevalier	Sterile	do	do	do	Lax	Chevalier	England	Newell, S. Dak		
1144	Mahan	do	do	do	do	do	Hama	Plant selection	North Dakota		
1145	Gold	do	do	do	do	do	do	do	Svalof, Sweden	Minnesota No. 435	
1146	Oderbrucker	Fertile	do	do	do		Manchuria	Manchuria	Various	Wisconsin No. 6	
1147	Composite						Hybrid	Mixture of hybrids	U. S. D. A., St. Paul, Minn.	Minnesota No. 400	
1170	Chevalier	Sterile	Hulled	Awned	White	Lax	Chevalier	England	New Zealand	Minnesota No. 306	Chevalier Prolific.
1174	Oderbrucker	Fertile	do	do	do		Manchuria	Manchuria	Various	Wisconsin No. 6, South Dakota No. 178	
1176	Mealy (Pl. III, fig. 1, C)	do	do	Hooded	do		Hybrid	Nepal X	Moro, Oreg		
1177	Wing Podigree	do	do	do	do		do	do	Chas. B. Wing, Ohio	Colorado No. 30, North Dakota No. 3101	Horsford.
1178	Manchuria	do	do	Awned	do		Manchuria	Plant selection	U. S. D. A., St. Paul, Minn.		
1179	Algerian	do	do	do	do		Coast	do	do		
1189	Manchuria	do	do	do	do		Manchuria	do	do		
1211	Samofa	do	do	do	do		Hybrid	South African X Manchuria	do	Minnesota No. 1-16-06	
1236	Abyssinian	Deficient	do	do	do		Abyssinia	Plant selection	U. S. D. A., Chico, Calif.	Minnesota No. 11-16-77	Mensury.
1246	Black Egyptian	Sterile	do	do	do		do	do	do		
1247	Blue	Fertile	do	do	Black	Blue	Coast	North Africa	Various	Washington No. 973	
1248	Excelsior	do	Naked	do		Purple	do	Central Asia	Pullman, Wash	Washington No. 989	Black Hull-less.
1249	Coast	do	Hulled	do	White		Coast	North Africa	Various	Washington No. 970	California.
1250	Eureka	do	Naked	Hooded	do		Nepal	India	do	Washington No. 988	

1252	Royal	do	Hulled	do	Awne	do	Blue	Hybrid	Swedish X Baxter	Ottawa, Canada	Wing's Ordinary.
1256	Arquipa	do	do	do	do	White	do	Coast	North Africa	Peru	
1263	Fichte	do	do	do	do	do	do	Coast	South Russia	Crimea	
1269	Sagina	do	do	do	do	do	do	Hybrid	North Africa	Sudan	
1271	Horsford	do	do	do	Hooded	do	do	do	Nepal X	Chas. B. Wing, Ohio.	
1272	Oderbrucker	do	do	do	Awne	do	do	Manchuria	Plant selection	Madison, Wis	Manchuria.
1273	do	do	do	do	do	do	do	do	do	do	Do.
1274	do	do	do	do	do	do	do	do	do	do	Manshury.
1275	Manchuria	do	do	do	do	do	do	do	do	do	Manchuria.
1276	Silver King	do	do	do	do	do	do	do	do	do	
1286	Hero	do	do	do	Smooth awne	do	do	Hybrid	Club Mariout X Lion	U. S. D. A., St. Paul, Minn.	
1303	Blarney	do	do	do	do	do	do	do	Lion X Manchuria	do	
1304	Hurst	do	do	do	do	do	do	do	Featherston X Lion	do	
1308	Louden	do	do	do	do	do	do	do	Bay Brewing X Lion	do	
1310	Erfurt White	do	do	do	Awne	do	do	Hanna	Germany	Various	
1311	Flynn	do	do	do	do	do	do	Smooth- awne hybrid.	Club Mariout X Lion	U. S. D. A., Moro, Oreg.	
1312	Himalaya	do	Naked	do	do	White	Blue	do	Coolest X Coast	Lompoc, Calif.	Zvolanek.
1389	Abel	do	Hulled	do	do	do	do	do	North Africa	Moscow, Idaho	
1455	California Mar- out (P. II, D).	do	do	do	do	do	do	California	do	Davis, Calif.	
1470	O. A. C. 21	do	do	do	do	do	do	Manchuria	Plant selection	Guelph, Ontario	Manchuria.
1472	Manchuria	do	do	do	do	do	do	do	do	Manhattan, Kans.	Manchuria.
1473	do	do	do	do	do	do	do	do	do	do	Do.
1474	Horsford	do	do	do	Hooded	do	do	Hybrid	Nepal X	do	
1478	Manchuria	do	do	do	Awne	do	do	Manchuria	Plant selection	St. Paul, Minn.	
1511	Golden Queen	do	do	do	do	do	do	do	do	Madison, Wis	Manchuria.
1517	Oderbrucker	do	do	do	do	do	do	do	Manchuria	Lincoln, Nebr	
1526	do	do	do	do	do	do	do	do	Plant selection	Ithaca, N. Y.	
1529	do	do	do	do	do	do	do	do	do	Madison, Wis	
1551	Albert	do	do	do	do	do	do	Hybrid	4-rowed X Cana- dian Thorpe	Ottawa, Canada	Oderbrucker- Manchuria.
1554	Hanna	do	do	do	do	do	do	Hanna	Germany	Ithaca, N. Y.	
1556	Minstirdi	do	do	do	do	do	do	Hybrid	South African X Manchuria	U. S. D. A., St. Paul, Minn.	Mensury.
1557	Claude	do	do	do	do	do	do	Hybrid	Duckbill X Com- mon.	Ottawa, Canada	

TABLE 94.—Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of caryopsis					
1562	Manchuria	Fertile	Hulled	Awned	White		Manchuria	Manchuria	Various		
1563	July	do	do	do	do		do	Plant selection	Sweden		
1576	Early Indian	do	do	do	do		do	India	Canada		
1577	Aker	do	do	do	do		Manchuria	Russia	St. Paul, Minn.	Minnesota No. 442	
1580	Manchuria	do	do	do	do		do	Manchuria	Lincoln, Nebr.		
1585	Himalaya	do	Naked	do	do	Blue	Himalaya	Central Asia	Manhattan, Kans.	Kansas No. 7128	
1592	Yale	do	Hulled	do	White		Hybrid	Duckbill X Ren- due Improved	Ottawa, Canada		
1701	Swedish Star	do	do	do	do	Dense	Winter Club	Europe	Canada		Common six-row. Abyssinian.
1707	Tupp Winter	do	do	do	do	do	do	Sweden	Sidney, British Columbia		Star. Utah Winter, Win- ter Club, White Winter.
1775	Horsford	do	do	Hooded	do		Hybrid	Nepal X	Pullman, Wash.	Washington No. 873	
1778	do	do	do	do	do		do	do	Manhattan, Kans.	Kansas No. 7023	
1787	Beardless	do	do	do	do		do	Nepal X Coast	Various		
1800	Bonanza	do	do	Awned	do		Manchuria	Manchuria	Kansas	Kansas No. 7025	
1807	Horsford	do	do	Hooded	do		Hybrid	Nepal X	do	Kansas No. 7020	
1808	Success Beardless	do	do	do	do		do	do	Indiana		
1853	Acc	Sterile	do	Awned	do	Lax	Smyrna	Plant selection	U. S. D. A., High- more, S. Dak.		Horsford.
1854	Hanna	do	do	do	do	do	Hanna	Germany	Ithaca, N. Y.		Two-rowed.
1877	White Moravian	do	do	do	do	do	Chevalier	England	Moscow, Idaho.	Puyallup No. 61	
1884	Saul	do	do	do	do	do	Hanna	Germany	Ithaca, N. Y.		
1886	Chevalier	do	do	do	do	do	Chevalier	England	Various	North Dakota Nos. 169 and 253.	
1892	Champion of Vermont	do	do	do	do	do	Hanna	Germany	do		
1894	Wisconsin Win- ter	Fertile	do	do	do		Wisconsin Winter	Europe	Pullman, Wash.	Washington No. 971	Chevalier Winter.
1907	Swansota	Sterile	do	do	do		Hybrid	No. 456 X Svanhals	U. S. D. A., St. Paul, Minn.	Minnesota Nos. 11-16-37 and 440.	
1909	Binder	do	do	do	do	Lax	Chevalier	England	Canada		
1910	Clifford	do	do	do	do		do	do	do		
1915	Beaver	do	do	do	do		Hybrid	Swedish X Baxter	Ottawa, Canada		
1916	Duckbill	do	do	do	do	Dense	Thorpe	England	Various		
1928	Jet	do	Naked	do	Black		Jet	Abyssinia	Manhattan, Kans.	Kansas No. 7129	

1934	Himalaya	Fertile	do	do	White	Blue	Himalaya Tennessee Winter	South-central Asia Europe	Various	Guy Mayle. Tennessee Winter.
2033	Ohio Winter	do	Hulled	do	do		do	do	Ohio	
2036	Michigan Winter	do	do	do	do		do	do	Indiana	
2039	Indiana Winter	do	do	do	do		do	do	do	
2080	White Smyrna	Sterile	do	do	do	Lax	Smyrna	Plant selection	U. S. D. A., Akron, Colo.	Do. Ouehae.
2087	Blackhull	Deficient	do	do	Black		Coast	do	do	
2103	Stavropol	Fertile	do	do	White		do	do	Various	
2107	Ellis	do	do	do	do		do	do	Hays, Kans.	
2126	McClymont	do	do	do	do		do	do	North Platte, Nebr.	
2135	Golden Drop	Sterile	do	do	do	Lax	Chevalier	England	Various	
2147	Wing Beardless	Fertile	do	Hooded	do		Hybrid	Nepal X	Moscow, Idaho	
2154	Odessa	do	do	Awned	do		Odessa	Plant selection	South Dakota	
2155	Manchuria	do	do	do	do		Manchuria	do	do	
2156	do	do	do	do	do		do	do	do	
2157	do	do	do	do	do		do	do	do	
2158	Odessa	do	do	do	do		Odessa	do	do	
2159	Wisconsin Winter	do	do	do	do		do	do	U. S. D. A., Ross- lyn, Va.	
2160	Arlington Awn- less	Inter- mediate	do	Awnless	do		Hybrid	Tennessee Winter X Black Arabian	do	
2161	do	do	do	do	do		do	do	do	
2162	do	do	do	do	do		do	do	do	
2163	Han River	Fertile	do	Awned	do		do	do	do	
2164	Nakano Wase	Inter- mediate	do	Awnless	do		do	Plant selection	do	
2165	do	do	do	do	do		do	do	do	
2166	do	do	do	do	do		do	do	do	
2167	Wisconsin Win- ter	Fertile	do	Awned	do		do	do	do	
2168	Odessa	do	do	do	do		Odessa	do	Higmore, S. Dak.	South Dakota No. 252.
2169	White Smyrna	Sterile	do	do	do	Lax	Smyrna	do	do	
2170	do	do	do	do	do	do	do	do	do	
2171	do	do	do	do	do	do	do	do	do	
2172	do	do	do	do	do	do	do	do	do	
2178	Chevalier	do	do	do	do	Lax	Chevalier	England	Manhattan, Kans.	
2181	Manchuria	Fertile	do	do	do		Manchuria	Plant selection	do	Kansas No. 7115
2182	do	do	do	do	do		do	do	do	Kansas No. 8025
2183	Success	do	do	do	do		Hybrid	Nepal X	do	Kansas No. 8029
2184	Oderbrueker	do	do	Hooded	do		Manchuria	do	do	Kansas No. 8034
2185	Austrian	do	do	Awned	do		do	do	Hays, Kans.	
2186	Chile Common	Fertile	Hulled	Awned	White		Coast	Plant selection	do	
2187	Odessa	do	do	do	do		Odessa	do	Akron, Colo.	
2189	Yendje	do	do	do	do		do	do	do	
2190	Orel	Sterile	do	do	do		do	do	do	
2191	Oderbrueker	Fertile	do	do	do	Lax	Manchuria	do	do	
2198	Union	do	Naked	Hooded	do		Hybrid	Horsford X Black Hull-less	Union, Oreg	
2199	do	do	do	do	do		do	do	do	

TABLE 94.—Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued

C. I. No.	Name	Description				Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of caryopsis				
2235	Coast.....	Fertile	Hulled.	Awned.	White.		Coast.....	Davis, Calif.		California Cent- gener No. 3318.
2241	Mansfeld.....	do.	do.	do.	do.		Hybrid	Ottawa, Canada.		
2295	Bevede.....	do.	Naked	Hooded.	do.		do.	Moro, Oreg.		
2296	Black Algerian.....	do.	Hulled.	Smooth awned.	Black		Lion.....	U. S. D. A., Moro, Oreg.		
2297	Chile Brewing.....	do.	do.	do.	do.		Coast.....	do.	Moro No. 708-1.	
2298	Himalaya.....	do.	Naked	do.	Blue		Himalaya.....	do.	Moro No. 657-1.	
2299	do.....	do.	do.	do.	do.		do.	do.		
2300	Black Abyssinian.....	do.	Hulled.	do.	White		Coast.....	do.		
2301	Coast.....	do.	do.	do.	do.		Coast.....	do.		
2302	Peru.....	do.	do.	do.	do.		Hybrid	Union, Oreg.		
2303	Withycombe.....	do.	Naked	Hooded.	do.		Hybrid	do.		
2304	Dufur.....	do.	Hulled.	Awned.	do.		Tennessee Winter.	Moro, Oreg.		Withycombe Hull- less.
2305	Coast.....	do.	do.	do.	do.		Coast.....	Burns, Oreg.		Tennessee Winter.
2306	do.....	do.	do.	do.	do.		do.	U. S. D. A., Chico, Calif.		
2307	do.....	do.	do.	do.	do.		do.	do.		
2308	Abina.....	do.	Naked	do.	do.		Hanna.....	do.		
2309	Hanna.....	Sterile.	Hulled.	do.	do.	Lax.	do.	do.		
2311	Peru.....	Fertile	do.	do.	do.		Coast.....	do.		
2312	Coast.....	do.	do.	do.	do.		do.	do.		
2313	Abyssinian.....	do.	do.	do.	do.		do.	do.		
2314	do.....	do.	do.	do.	do.		do.	do.		
2315	Nepal.....	Fertile	Naked	Hooded.	White		Nepal.....	do.		
2316	Hanna.....	Sterile.	Hulled.	Awned.	do.	Lax.	Hanna.....	do.		
2330	Manchuria.....	Fertile	do.	do.	do.		Manchuria.....	St. Paul, Minn.	Minnesota No. 184.	Man- Improved churia.
2617	do.....	do.	do.	do.	do.		do.	Various		
2635	Canadian.....	do.	do.	do.	do.		do.	Ithaca, N. Y.		
2636	Hybrid.....	Sterile.	do.	do.	do.	Lax.	Hybrid	do.		
2637	do.....	do.	do.	do.	do.		do.	do.		
2638	Manchuria.....	Fertile.	Hulled.	Awned.	White.		do.	do.		

[illegible]

[illegible]

TABLE 94.—Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of ears/opsis					
2761	Wisconsin Pedit- gree 20.	Fertile	Hulled	Awnecl.	White		Manchuria	Plant selection	Madison, Wis.	Wisconsin Pedit- gree 19.	Mansbury.
2762	Manchuria	do	do	do	do		do	do	do		
2763	Wisconsin Pedit- gree 18.	do	do	do	do		do	do	do		Do.
2764	Manchuria	do	do	do	do		do	do	do		
2765	Wisconsin Pedit- gree 17.	do	do	do	do		do	do	do		
2766	Wisconsin Pedit- gree 35.	do	do	do	do		do	do	do		
2767	Scotch Chevalier	Sterile	do	do	do		Chevalier	England	Bozeman, Mont.		Chevalier. Do.
2768	Chevalier	do	do	do	do		do	do	do		Probably Nepal.
2769	Berkley	do	do	do	do		do	do	do		
2770	Nepbant	Sterile	Hulled	Awnecl.	White		Chevalier	England	do		
2771	Chevalier	do	do	do	do		Hanna	Germany	do		
2772	Chile Brewing	Sterile	Hulled	Awnecl.	White		Manchuria	Manchuria	do		
2773	Hanna	do	do	do	do		do	China	do		
2774	Manchuria	Fertile	do	do	do		do	do	do		
2775	Shansi	do	Naked	do	do		do	do	do		
2776	Spear	Fertile	Hulled	Awnecl.	White	Blue	Coast	Plant selection	Davis, Calif.		Oderbrucker. Shansi Hull-less.
2777	Beldi Giant	do	do	do	do		Manchuria	Manchuria	Wooster, Ohio.		Spear and Mason Beldi, Coast.
2778	Manchuria	do	do	do	do		do	do	La Fayette, Ind.		
2779	Great Beardless	Fertile	Hulled	Awnecl.	White		Manchuria	Manchuria	do		Manchuria.
2780	Oderbrucker	do	do	do	do		do	do	do		
2781	Stoeckinger	Sterile	Hulled	Awnecl.	White		Manchuria	Manchuria	do		
2782	Michigan	Fertile	do	do	do		Manchuria	Manchuria	Colorado		
2783	Manchuria	Fertile	do	do	do		Hanna	Germany	do		
2784	Hanna	Fertile	do	do	do		Coast	Plant selection	do		
2785	Coast	Fertile	do	do	do		Hanna	do	do		
2786	Hanna	Sterile	do	do	do		do	do	do		
2787	do	do	do	do	do		do	do	do		
2788	do	do	do	do	do		do	do	do		
2789	Coast	Fertile	do	do	do		Coast	do	do		
2790	do	do	do	do	do		do	do	do		
2791	do	do	do	do	do		do	do	do		
2792	Colless	do	do	do	do		do	do	do		
2793	Bark	do	Hooded	Awnecl.	White	Blue	Hybrid	do	do	Minnesota No. 444.	

2794	Giant	Fertile	Hulled	Awmed	White				Russia	South Dakota, U.S.D.A., Fargo, N. Dak.	North Dakota No. 107.
2795	Russian	Sterile	do	do	do	Lax	Hanna Tennessee Winter.		Czechoslovakia Europe	Edgeley, N. Dak. Maryland.	
2796	New Moravian	Fertile	do	do	do				Plant selection	U. S. D. A., Arlington Farm, Va.	
2797	Dutrow winter	do	do	do	do					U. S. D. A., St. Paul, Minn.	Minnesota No. 11-17-2.
2798	Osaka	do	do	do	do					do	Minnesota No. 11-16-91.
2799	Moister	do	do	Smooth awmed	do	Hybrid			Lion X Manchuria	do	
2800	Elfy	do	do	do	do	do			Lion X Featherston.	do	
2801	Thomas Chevalier	do	do	Awmed	do	Lax	Chevalier		England	Scottsbluff, Nebr. Madison, Wis.	North Dakota No. 170.
2802	Saxonian	Fertile	Hulled	Awmed	White		Hybrid		6-Row X 2-Row	do	
2803	Sibley Imperial	Fertile	Hulled	Awmed	White					Charlottesville, Va. Madison, Wis.	
2804	Hungarian Hulled	Fertile	Hulled	Hooded	White	Hybrid			Nepal X	do	
2805	Success	Fertile	Hulled	Smooth awmed	Black				Plant selection	East Lansing, Mich.	
2806	Arizona	Fertile	Hulled	Awmed	White					do	
2807	Semibarless	Fertile	Hulled	Awmed	White					State College, N. Mex.	
2808	Crawford	Fertile	Hulled	Awmed	White					do	
2809	Six-Row	Fertile	Hulled	Awmed	White					do	
2810	Hull-less	O. A. C. Selection 8.								do	
2811	O. A. C. Selection 7.									do	
2812	Russian Naked									do	
2813	Garton 5004									Lincoln, Nebr.	
2814	Garton 5006									do	
2815	Garton 5007									do	
2816	Garton 5008									do	
2817	Garton 5009									do	
2818	Garton 5010									do	
2819	Garton 5011									do	
2820	Manchuria	Fertile	Hulled	Awmed	White		Manchuria Coast.		Manchuria	do	
2821	Coast	do	do	do	do				Plant selection	Davis, Calif.	California No. 2285.
2822	Manchuria	do	do	do	do		Manchuria		Manchuria X Manchuria	St. Paul, Minn.	Minnesota No. 388.
2823	Empress	Sterile	do	do	do				Manchuria	Canada	
2824	Culver	do	do	do	do					St. Paul, Minn., Fargo, N. Dak.	North Dakota No. 257, Minnesota No. 86.
2825	South African	Fertile	Hulled	Awmed	White	Dense	Winter Club.			St. Paul, Minn.	
2826	Manchuria	do	do	do	do		Manchuria		Plant selection	do	Minnesota No. 158.

TABLE 94.—Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued

C. I. No.	Name	Description						Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of caryopsis	Density					
2828	Duckbill Bald.							Hybrid		St. Paul, Minn.	Minnesota No. 125.	
2829	Brush.	Sterile	Hulled	Awed.	White.					do.	Minnesota No. 126.	
2830	Ross.	Fertile	do.	do.	do.					do.	Minnesota No. 127.	
2831	Manchuria.	do.	do.	do.	do.			Manchuria	Plant selection	do.	Minnesota No. 129.	
2832	Crop Nursery.	do.	do.	do.	do.				do.	do.	Minnesota No. 130.	
2833	Hybrid	Sterile	do.	do.	do.			Hybrid	2-rowed X branch- ing Duckbill.	do.	Minnesota No. 131.	
2834	do.	do.	do.	do.	do.			do.	Black X long 2- rowed.	do.	Minnesota No. 132.	
2835	Manchuria.	Fertile	do.	do.	do.			Manchuria	Plant selection	do.	Minnesota No. 134.	
2836	do.	do.	do.	do.	do.			do.	do.	do.	Minnesota No. 135.	
2837	do.	do.	do.	do.	do.			do.	do.	do.	Minnesota No. 136.	
2838	Hybrid	Sterile	do.	do.	do.			Hybrid	2-rowed X branch- ing Duckbill.	do.	Minnesota No. 137.	
2839	do.	do.	do.	do.	do.			do.	Black X long 2- rowed.	do.	Minnesota No. 138.	
2840	do.	do.	do.	do.	do.			do.	do.	do.	Minnesota No. 139.	
2841	do.	do.	do.	do.	do.			do.	6-row smooth X 2- row Indian Head.	do.	Minnesota No. 140.	
2842	do.	Fertile	do.	do.	do.			do.	do.	do.	Minnesota No. 141.	
2843	do.	do.	do.	do.	do.			do.	do.	do.	Minnesota No. 142.	
2844	do.	Sterile	do.	do.	do.			do.	do.	do.	Minnesota No. 143.	
2845	do.	do.	do.	do.	do.			do.	do.	do.	Minnesota No. 144.	
2846	do.	Mixed	do.	do.	do.			do.	do.	do.	Minnesota No. 145.	
2847	do.	Sterile	Hulled	Awed.	White.			do.	do.	do.	Minnesota No. 146.	

[illegible]

TABLE 94.—Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of caryopsis					
2874	Manchuria	Fertile	Hulled	Awned	White		Manchuria	Plant selection	St. Paul, Minn.	Minnesota No. 189.	
2875	Hybrid	do	do	do	do		Hybrid	6-row smooth X 2- row Indian Head.	do	Minnesota No. 190.	
2876	do						do	do	do	Minnesota No. 191.	
2877	Chevalier	Sterile	Hulled	Awned	White		Chevalier	England	do	Minnesota No. 192.	
2878	Frankish Brew- ing	do	do	do	do		Hanna	Germany	do	Minnesota No. 197.	
2879	Manchuria	Fertile	do	do	do		Manchuria	Plant selection	do	Minnesota No. 197.	
2880	Daniels	do	do	do	do		do	Manchuria	do	Minnesota No. 197.	
2881	Manchuria	do	do	do	do		do	Manchuria	do	Minnesota No. 196.	
2882	do	do	do	do	do		do	Plant selection	do	Minnesota No. 196.	
2883	do	do	do	do	do		do	do	do	Minnesota No. 198.	
2884	do	do	do	do	do		do	do	do	Minnesota No. 199.	
2885	do	do	do	do	do		do	do	do	Minnesota No. 210.	
2886	Hanna	Sterile	do	do	do		do	do	do	Minnesota No. 211.	
2887	Silver Beardless	do	do	do	do		Hanna	do	do	Minnesota No. 212.	
2888	Hybrid	do	do	Hooded	do		Hybrid	do	do	Minnesota No. 213.	
2889	do	do	do	Awned	do		do	do	do	Minnesota No. 214.	
2890	do	Fertile	do	do	do		do	Manchuria X Champion of Vermont.	do	Minnesota No. 218.	
2891	do	do	do	do	do		do	do	do	Minnesota No. 219.	
2892	do	Sterile	do	do	do		do	Manchuria X Chevalier.	do	Minnesota No. 220.	
							do	do	do	Minnesota No. 221.	
							do	Manchuria X Champion of Vermont.	do	Minnesota No. 222.	

[illegible]

TABLE 94.—*Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued*

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of caryopsis					
2018	Manchuria	Fertile	Hulled	Awned	White		Manchuria	Manchuria × Manchuria	St. Paul, Minn.	Minnesota No. 324	
2019	Hanna	Sterile	do	do	do		Hanna	Plant selection	do	Minnesota No. 356	
2020	Baku	do	Naked	do	do		Baku	do	do	Minnesota No. 370	
2021	Coast	Fertile	Hulled	do	do		Coast	do	do	Minnesota No. 382	
2022	Scotch	do	do	do	do		Manchuria	do	do	Minnesota No. 383	
2023	Mansfield	do	do	do	do		Hybrid	Duckbill × Ren- nie improved	do	Minnesota No. 384	
2024	Mensury	do	do	do	do		Manchuria	Plant selection	do	Minnesota No. 385	
2025	Manchuria	do	do	do	do		do	Golden Queen × Manchuria	do	Minnesota No. 386	
2026	do	do	do	do	do		do	Manchuria × Manchuria	do	Minnesota No. 387	
2027	do	do	do	do	do		do	do	do	Minnesota No. 389	
2028	do	do	do	do	do		do	do	do	Minnesota No. 390	
2029	do	do	do	do	do		do	Golden Queen × Manchuria	do	Minnesota No. 391	
2030	do	do	do	do	do		do	do	do	Minnesota No. 392	
2031	Hybrid	do	do	do	do		Hybrid	Golden Queen × Blue	do	Minnesota No. 410	
2032	do	do	do	do	do		do	Blue × Golden Queen	do	Minnesota No. 414	
2033	Bohman	Sterile	do	do	do		do	Böhemian × Manchuria	U. S. D. A., St. Paul, Minn.	Minnesota No. 11-16-78	
2035	Hybrid	Fertile	do	Smooth awned	do		do	Lion × Manchu- ria	do	Minnesota No. 438	
2035	Black	Sterile	do	do	Black		do	Asia Minor	St. Paul, Minn.	Minnesota No. 417	
2037	Senca Purple	Fertile	do	do	do		do	do	Fargo, N. Dak.	North Dakota No. 681	
2038	Hoover Hull-less	Fertile	do	do	do		do	do	do	North Dakota No. 234	
2039	Hull-less	Fertile	do	do	do		do	do	do	North Dakota No. 168	
2041	Highland Scotch	Sterile	Hulled	do	White		do	do	do		
2042	Carter	Fertile	do	do	do		do	do	do		
2043	Salzer	Sterile	do	do	do		Chevalier	do	do		

Row	Montana	Rowed.
2944	Manchuria	do
2945	do	do
2946	do	do
2947	do	do
2948	Hull-less	do
2949	Manchuria	do
2950	do	do
2951	do	do
2952	do	do
3112	Common	do
3113	Rottingham	do
3114	Brewers Delight	do
3115	Winter	do
3116	Common Two-Row	do
3117	Nonpareil	do
3118	Montana	do
3121	Marsbury	do
3122	Union	do
3123	do	do
3124	do	do
3125	do	do
3126	do	do
3127	do	do
3128	do	do
3129	do	do
3130	do	do
3131	do	do
3132	do	do
3133	do	do
3134	do	do
3135	New Hull-less	do
3136	Crawford	do
3137	do	do
3138	do	do
3139	do	do
3140	Manchuria	do
3141	do	do
3142	do	do
3143	do	do
3144	Beardless	do
3145	do	do
3146	Oderbrucker	do
3147	California Cent- gener.	do
3148	Virginia Selection	do
3149	do	do

Montana
T w o -
Rowed.

TABLE 94.—Accession data of varieties of barley arranged in the order of their Cereal Investigations numbers—Continued

C. I. No.	Name	Description					Group	Probable origin	Immediate source	Designations	Synonyms
		Lateral florets	Adher- ence	Append- ages to central florets	Color of lemma	Color of caryopsis					
3150	Virginia Selection								Blacksburg, Va.	Blacksburg 3	
3151	do.								do	Blacksburg 4	
3152	do.								do	Blacksburg 5	
3153	do.	Fertile	Hulled	Awned	White				do	Blacksburg 6	
3154	do.								do	Blacksburg 7	
3155	do.								do	Blacksburg 8	
3156	do.								do	Blacksburg 9	
3157	do.								do	Blacksburg 10	
3158	Hull-less								do	Blacksburg 11	
3159	Virginia Selection								do	Blacksburg 12	
3160	Winter								do	Blacksburg 13	
3161	Winter Beardless								do	Blacksburg 14	
3162	Tennessee	Fertile	Hulled	Awned	White		Tennessee Winter	Europe	do	Blacksburg 45	
3163	Virginia Selection							Plant selection	do	Blacksburg 46	
3164	do.							do	do	Two-rowed from Selection 7	
3165	Chevalier	Sterile	Hulled	Awned	White		Chevalier	England	do	Blacksburg 30	Manchury.
3166	Manchuria	Fertile	do	do	do		Manchuria	Manchuria	do	Blacksburg 20	
3167	do.	do	do	do	do		do	do	do		
3168	Hybrid						Hybrid	Rice X Beardless	Land, Wash		
3170	Caswell								Durham, N. H		
3171	Horsford	Fertile	Hulled	Hooded	White		Hybrid	Nepal X	Manhattan, Kans.	Kansas No. 8001	
3172	Manchuria	do	do	Awned	do		Manchuria	Plant selection	do	Kansas No. 7015	
3173	do.	do	do	do	do		do	do	do	Kansas No. 8022	
3174	do.	do	do	do	do		do	do	do	Kansas No. 8030	
3175	do.	do	do	do	do		do	do	do	Kansas No. 8032	
3176	Bonanza	do	do	do	do		do	do	do	Kansas No. 8049	
3177	do.	do	do	do	do		do	do	do		
3178	Two-Row Brew- ing	Sterile	do	do	do		do	do	Durham, N. H		
3179	Union							Plant selection	Union, Oreg	Union No. 28-2	
3180	Two-Row Hull- less								Blacksburg, Va.		
3181	O. A. C. Winter								Burns, Oreg		
3182	Manchuria	Fertile	Hulled	Awned	White		Manchuria	Manchuria	Hays, Kans.	Hays No. 57	

In Table 94 the terms relating to the fertility of lateral florets must necessarily be definite. The central floret is always fully fertile. In the 6-rowed barleys the lateral florets also are fertile. In the 2-rowed varieties they are present but sterile. Besides these two classes, which include all our common varieties, there are two other groups, the homozygous intermediate and the deficient. In the intermediate barleys the lateral florets are fertile, but the resulting kernels are usually smaller than in the 6-rowed varieties, and the percentage of fertility is lower. The most conspicuous characteristic of the intermediate barley is in the lack of appendages on the lateral floret. These florets are characterized by rounded lemmas, while the central ones may bear long awns or hoods.

The fourth variation, the deficient barleys, concerns a still further reduction of the lateral florets than is found in the common 2-rowed varieties. These barleys are 2-rowed, of course, but the lateral spikelets are reduced to the outer glumes and lemma. The palet is sometimes present, but sexual organs are never found.

BOTANICAL COMPARISON OF PROMINENT VARIETIES

It is quite obvious that the abridged description given in Table 94 affords only a general conception of the nature of the varieties. The separation of similar strains from the same parent stock is very difficult. It is the opinion of the writers that the nomenclature used in taxonomic work should not be carried beyond the major groups. The variations of barley are so numerous that the varieties can be arranged in a great number of ways.

If an attempt is made to gather these under named taxonomic groups, it is impossible to avoid placing forms of totally different and unrelated adaptation in a group where the type variety is ecologically unrelated to them. Thus, if the Svalof characters of minor variation, such as the toothing of the nerves and the nature of the basal bristle, are appended to the usual classifications it is perfectly possible, even in the ultimate divisions of morphological groupings, to find a winter variety suited to the arid conditions of North Africa occurring as a variation of a spring variety from Manchuria suited to humid conditions.

TABLE 95.—*Botanical comparison of prominent varieties of barley*

[Many of these characters are not comparable unless grown under identical conditions. *Explanation.*—In column 3: D=deficient, I=intermediate, 2=2-rowed, 6=6-rowed. In column 4: H=hulled, N=naked. In column 5: A=awned, Al=awnless, H=hooded, Sa=smooth awned. In column 6: * = 1 head only, † = variable. In column 7: L=long, S=short, LS=mixed (both long and short). In column 8: B=blue, Bk=black, BW=blue and white, V=violet, W=white. In column 10: A=Aberdeen, Idaho; Ag=Arlington Farm, Va. (greenhouse); Ak=Akron, Colo.; C=Chico, Calif.; D=Davis, Calif.; Di=Dickinson, N. Dak.; S=St. Paul, Minn.; W=Madison, Wis.

Variety and station No.	C. I. No.	Fertility	Caryopsis	Lemma	Spike density (mm.)	Ra-chilla	Color	Seed length (mm.)	Place and year grown	At Aberdeen, Idaho, in 1921	
										Awns emerged	Culm length (cm.)
1	2	3	4	5	6	7	8	9	10	11	12
Charlottetown 80.....	2732	2	II	A	3.5	S	W	9.1	A 1921	July 30	94
Chevalier II.....	200	2	II	A	3.6	S	W	9.2	A 1921	do.....	86
French Chevalier.....	175	2	II	A	3.5	S	W	7.9	Ag 1923	-----	-----
Horn.....	926	2	II	A	3.6	S	W	9.0	A 1921	July 28	101

TABLE 95.—*Botanical comparison of prominent varieties of barley*—Continued

Variety and station No.	C. I. No.	Fer- tility	Cary- opsis	Lem- ma	Spike den- sity (mm.)	Ra- chilla	Color	Seed length (mm.)	Place and year grown	At Aberdeen, Idaho, in 1921	
										Awns emerged	Culm length (cm.)
1	2	3	4	5	6	7	8	9	10	11	12
Canadian Thorpe.....	740	2	H	A	2.2	L	W	8.1	C 1921	-----	-----
Primus.....	532	2	H	A	2.4	S	W	10.3	D 1922	-----	-----
Manchuria (Minn. 6).....	241	6	H	A	3.3	L	W	8.5	A 1921	July 27	100
Manchuria (Minn. 105).....	244	6	H	A	3.4	LS	B	8.6	A 1921	July 24	113
Manchuria (Minn. 184).....	2330	6	H	A	3.4	S	BW	8.8	A 1921	July 25	111
O. A. C. 21.....	1470	6	H	A	3.5	S	B	8.8	A 1921	July 26	102
Manchurian (no sample).....	739	6	H	A	3.4	S	B	10.5	D 1923	-----	-----
Mensury (no sample).....	2657	6	H	A	-----	-----	-----	-----	-----	-----	-----
Featherston.....	1120	6	H	A	3.6	S	B	9.3	A 1921	July 23	100
Oderbrucker.....	836	6	H	A	3.9	LS	BW	8.8	A 1921	July 24	113
Oderbrucker (S. Dak. 178).....	1174	6	H	A	3.9	S	W	8.8	A 1920	-----	-----
Oderbrucker (Wis. Ped. 5).....	1272	6	H	A	3.7	S	W	8.8	A 1921	July 24	103
Oderbrucker (Wis. Ped. 6).....	1529	6	H	A	4.1	S	W	9.0	A 1921	July 26	111
Hanna.....	203	2	H	A	3.8	L	W	9.4	A 1921	do.....	88
Gold.....	1145	2	H	A	3.4	L	W	8.8	A 1921	July 30	90
Hannchen.....	531	2	H	A	3.4	L	W	9.9	A 1921	July 29	93
Odessa.....	182	6	H	A	3.5	S	W	9.2	A 1921	July 20	92
Svanhals.....	187	2	H	A	2.5	L	W	9.5	A 1921	July 30	92
Gatami.....	575	6	H	A	3.5	L	Bk	9.0	A 1921	July 20	91
Princess.....	529	2	H	A	3.4	L	W	9.5	C 1917	-----	-----
Lion.....	923	6	H	A	4.0	L	Bk	9.6	A 1921	July 23	94
Mansury.....	617	2	H	A	2.5	L	W	-----	{Di1907 Ak1908}	-----	-----
Coast.....	626	6	H	A	3.4	S	B	10.4	A 1921	July 19	87
Do.....	690	6	H	A	3.1	S	B	10.1	A 1921	July 17	90
Beldi Giant.....	2777	6	H	A	3.4	S	B	10.5	A 1921	July 23	83
Sandrel.....	937	6	H	A	3.2	S	W	9.1	C 1919	-----	-----
Stavropol.....	2103	6	H	A	3.3	S	B	10.2	A 1921	July 23	87
Peruvian.....	935	6	H	A	3.7	S	B	11.2	A 1921	July 21	97
Telli.....	194	6	H	A	4.2	LS	B	10.6	A 1921	July 24	79
Poda.....	652	6	H	A	3.7	S	B	10.5	A 1921	July 23	96
Club Mariout.....	261	6	H	A	2.6	S	W	10.6	A 1921	July 16	76
California Mariout.....	1455	6	H	A	3.7	L	B	10.2	A 1921	July 13	55
White Smyrna.....	195	2	H	A	3.6	L	W	10.2	A 1921	July 20	62
Do.....	658	2	H	A	4.0	L	W	10.2	A 1921	do.....	67
Beldi Dwarf.....	190	6	H	A	2.6	L	BW	10.3	A 1921	July 16	66
Nepal.....	595	6	N	H	2.7	L	W	8.0	A 1921	July 25	92
Himalaya.....	620	6	N	A	3.4	L	B	8.5	A 1921	July 24	85
Trebi.....	936	6	H	A	3.5	S	B	10.6	A 1921	do.....	93
Baker.....	975	6	H	A	3.5	S	W	10.6	A 1921	July 22	90
Black Hull-less.....	596	6	N	A	3.1	L	V	7.9	A 1921	July 23	96
Baku.....	709	2	N	A	4.4	L	W	9.5	A 1921	July 22	82
Olonets.....	198	6	H	A	3.1	S	W	9.4	D 1922	-----	-----
Tennessee Winter.....	257	6	H	A	3.0	S	W	8.6	C 1918	-----	-----
Ohio Winter.....	2033	6	H	A	*3.8	S	W	10.2	S 1921	-----	-----
Winter Club.....	592	6	H	A	1.7	L	W	9.5	A 1921	Aug. 8	72
Wisconsin Winter.....	519	6	H	A	3.7	S	B	9.9	A 1921	July 21	75
Orel.....	351	2	H	A	3.8	LS	W	9.2	A 1921	Aug. 1	77
Turkestan.....	711	6	H	A	3.1	L	B	9.0	-----	-----	-----
Blackhull.....	878	D	H	A	3.5	S	Bk	10.1	A 1921	July 16	78
Han River.....	206	6	H	A	3.9	S	B	9.9	A 1921	July 18	89
Nakano Wase (Pl. III, fig. 1, D).....	754	I	H	Al	2.5	L	W	8.0	Ag 1920	-----	-----
Meloy.....	1176	6	H	H	†3.2	S	B	9.5	A 1921	July 24	97
Flynn.....	1311	6	H	Sa	3.9	L	W	9.9	A 1921	July 17	93
Horsford.....	507	6	H	H	3.4	S	W	8.8	A 1921	July 29	114
Stella.....	2678	6	H	A	4.0	S	W	8.1	Ag 1923	-----	-----
Alpha.....	959	2	H	A	3.3	S	W	9.0	A 1921	July 26	96
Minsturdi (Minn. 437).....	1556	6	H	A	2.4	S	W	8.8	A 1921	July 22	96
Svansota (Minn. 1140).....	1907	2	H	A	2.9	L	W	9.4	A 1921	July 30	90
Hanchamont.....	1121	2	H	A	2.7	L	W	8.0	C 1921	-----	-----
Yale.....	1592	6	H	A	3.3	S	W	9.5	A 1921	July 25	85
Standwell (no sample).....	584	2	H	A	-----	L	W	9.3	W 1907	-----	-----
Early Chevalier.....	2725	2	H	A	3.2	L	W	8.4	Ag 1923	-----	-----
Invincible.....	590	2	H	A	2.0	L	W	8.1	C 1921	-----	-----
Blue Long Head (no sample).....	2685	6	H	A	-----	-----	B	-----	-----	-----	-----

As far as the agronomic classification is concerned the ecological features with their relation to season and rainfall are of more importance than the taxonomic ones. Some of the physiological differences which are expressed in height, time of flowering, or even in yield are the ones on which the final distinctions between varieties are likely to be made. In 1921 a large collection of barleys from various experiment stations was grown at Aberdeen, Idaho. The observations made on a few of these are shown in Table 95. One of the most useful observations made was the time of emergence of awns, which is shown in column 11 in this table. The date the awns emerged from the leaf sheath was noted. This is a much more accurate statement of the stage of development than the time of ripening. It can be much more accurately taken, and the plants at this time are not affected in their rate of development by disease and usually not by diminution of water supply. The great difference in appearance which is represented by only one or two days in the time of emergence of the awns can be appreciated only in the field.

It is certain that as our selected varieties are increased in number by the many agencies now working in barley improvement, their separation will become more and more difficult. Most certainly varieties will be put out which are indistinguishable. It is important that their relationships be studied, and some such system as was used at Aberdeen in 1921 is likely to be useful. The occasional assembling of the varieties at a single place would serve to detect the confusion which is inevitable when so many are handled by a large number of stations.

BEST VARIETIES OF THE PERIOD FROM 1917 TO 1921, INCLUSIVE

Emphasis throughout this bulletin has been placed on group yields and on the ecological and regional adaptation of types. Specific varieties have been mentioned from time to time as having done well at the stations. Since the methods and equipment of testing in the earlier years were obviously imperfect the performances of individual sorts were not stressed. In the last five years covered, however, replications of plats and other improvements have been common, and most of the varieties tested were pure-line selections.

In Table 96 the varieties giving the best yield in some or all of the five years from 1917 to 1921, inclusive, at each station, are listed in the order of productiveness. It will be noticed that very few non-pedigreed varieties are included. The few that are found are for the most part very definite types, such as White Smyrna (C. I. No. 195) and Club Mariout (C. I. No. 261). But very little can be added in the way of amplifying the facts brought out in the table. The material contained should be of use to agronomists in obtaining varieties for testing which have done well at near-by stations, but which for some reason have not been included in their plats.

TABLE 96.—*Varities of barley giving the best yields at different stations in part or all of the five years from 1917 to 1921, inclusive, listed in order of yield*

[A figure in parentheses following the name of a variety indicates that such variety was grown only for the number of years thus shown, not for the entire five years. In like manner F = sown in fall; S = sown in spring]

Place	Num-ber of varie-ties com-pared	First variety		Second variety		Third variety		Fourth variety		Fifth variety	
		Name	C. I. No.	Name	C. I. No.	Name	C. I. No.	Name	C. I. No.	Name	C. I. No.
Storrs, Conn.	3	Wisconsin Pedigree	835	Oderbrucker	1273	O. A. C. 21	1470	Trebi (4)	936	Hanchmont (4)	1121
Utica, N. Y.	15	Oderbrucker (4)	1272	Manchuria (4)	2823	Oderbrucker (4)	836	Wisconsin Pedigree	835	Wing Pedigree (3)	1177
State College, Pa.	922	Charlottetown 80	2732	Lion (2)	923	Michigan (2)	2782	(2)			
Wooster, Ohio	2633	Ohio Winter (4)	554	Texas Winter (4)	554	Gold (3)	2036	Wisconsin Pedigree	835	Oderbrucker	836
East Lansing, Mich.	18	Crawford (3)	3133	Crawford (3)	3137	Michigan (3)	1145	Crawford (3)	3139	Crawford (3)	3136
La Fayette, Ind.	9	Lion (2)	923	Manchuria (3)	2330	Oderbrucker (2)	1529	Michigan (2)	2782	Wisconsin Pedigree (4)	835
Urbana, Ill.	5	Wisconsin Pedigree	835	Oderbrucker	835	Horsford	507	Montana	3118	Nepal	595
De Kalb, Ill.	6	Oderbrucker	836	Wisconsin Pedigree	835	Lion	923	Horsford	507	Montana	3118
Madison, Wis.	10	O. A. C. 21 (4)	1470	Oderbrucker	1529	Oderbrucker	1272	Manchuria	1272	Manchuria	739
St. Paul, Minn.	10	Minsturd (4)	1556	Manchuria	2330	Aker (3)	1577	Svansota (4)	1907	French Chevalier	2900
Waseca, Minn.	8	Manchuria (4)	2330	Hybrid (3)	2635	Minsturd (3)	1556	French Chevalier (4)	2900	Samola (2)	1211
Duluth, Minn.	6	Svansota (3)	1907	French Chevalier (3)	2900	Manchuria (3)	2330	Manchuria (2)	1189	Manchuria (2)	1478
Grand Rapids, Minn.	8	Minsturd (3)	1556	Manchuria (4)	2330	French Chevalier (4)	2900	Svansota (3)	1907	Hybrid (3)	2635
Crookston, Minn.	19	Minsturd (3)	1556	Manchuria (3)	2330	Samola (2)	1211	Manchuria (2)	1189	O. A. C. 21 (3)	1470
Morris, Minn.	14	Manchuria (2)	1189	Manchuria (4)	2330	Manchuria (3)	2823	do	1478	Svansota (3)	1907
Ames, Iowa	35	Oderbrucker	1272	Oderbrucker	2700	O. A. C. 21	1470	Manchuria	241	Frankish	295
Columbia, Mo.	3	Sandral (3)	937	Trebi (3)	496	Featherston (3)	1120	Manchuria (3)	923	Lion (3)	923
Fargo, N. Dak.	2945	Manchuria (4)	2945	Manchuria (4)	2947	Lion (2)	923	Manchuria (4)	244	Wisconsin Pedigree (4)	835
Edgeley, N. Dak.	4	Oderbrucker (2)	836	Manchuria (2)	214	Oderbrucker (2)	1273	Nepal (2)	595	Club Marlout	261
Maritan, N. Dak.	6	White Smyrna	195	Svansals	187	Hannchen	531	Coast	690	Scholey (3)	962
Dickinson, N. Dak.	18	Stelgun	997	White Smyrna	658	Hanna	203	Hannchen	531	Manchuria (4)	882
Williston, N. Dak.	12	Hanna (4)	203	Oderbrucker (4)	888	Hannchen (4)	531	Wing Pedigree (3)	1177	Manchuria (3)	244
Brookings, S. Dak.	14	Odessa (3)	182	Manchuria (3)	2156	Gatami (3)	575	Odessa (3)	2154	Manchuria (4)	575
Eureka, S. Dak.	5	Odessa (2)	182	White Smyrna (2)	195	Hannchen (3)	531	Manchuria (3)	244	Gatami (1)	626
Higmore, S. Dak.	15	Odessa (3)	182	do	2170	White Smyrna (2)	2171	White Smyrna (2)	2169	Coast (3)	626
Cottonwood, S. Dak.	8	do	182	White Smyrna (3)	195	Gatami (3)	575	Hannchen (3)	531	Manchuria (1)	244
Newell, S. Dak. (dry land)	6	White Smyrna (3)	658	Coast (3)	690	do	575	do	531	Odessa (3)	182
Newell, S. Dak. (irrigated)	16	Trebi	935	Chevalier II	200	Hannchen (4)	531	Coast (4)	690	Silver King (3)	905
Lincoln, Nebr.	7	Club Marlout (2)	936	Ace (2)	1353	O. A. C. 21 (2)	1470	White Smyrna (2)	195	Svansals (2)	187
North Platte, Nebr.	6	Trebi (2)	936	McClymont (2)	2126	Coast	690	Stavropol	2103	White Smyrna	658
Mitchell, Nebr.	5	Trebi (3)	936	Babary (3)	695	Svansals (2)	187	Franchon (1)	680	Nepul (1)	595
Manhattan, Kans.	28	Manchuria	1473	Manchuria (3)	244	Wisconsin Winter (F)	519	Coast (3)	690	Odessa	182
Hays, Kans.	15	Club Marlout	261	Coast (4)	690	Gatami	575	Beldi Dwarf	190	White Smyrna	195

TABLE 97.—Five-year average performance of varieties of barley, 1917 to 1921, inclusive, shown by stations

Varieties and stations	Rank	Number compared	Varieties and stations	Rank	Number compared
Svanhals (C. I. No. 187):			Manchuria (C. I. No. 244)—Con.		
Mandan, N. Dak.	2	6	Eureka, S. Dak.	4	5
Mitchell, Nebr.	3	5	Garden City, Kans.	4	6
Sheridan, Wyo.	3	8	Cottonwood, S. Dak.	5	5
Lincoln, Nebr.	5	16	Brookings, S. Dak.	5	14
Moccasin, Mont.	5	13	Manchuria (C. I. No. 2330):		
Huntley, Mont.	5	5	La Fayette, Ind.	2	9
Newlands, Nev.	5	7	Waseca, Minn.	1	8
French Chevalier (C. I. No. 2900):			St. Paul, Minn.	2	10
St. Paul, Minn.	5	10	Grand Rapids, Minn.	2	8
Waseca, Minn.	4	8	Crookston, Minn.	2	19
Duluth, Minn.	2	6	Morris, Minn.	2	14
Chevalier II (C. I. No. 200):			Duluth, Minn.	3	6
Newell, S. Dak. (irrigated)	2	7	O. A. C. 21 (C. I. No. 1470):		
Huntley, Mont.	3	5	Storrs, Conn.	3	3
Horn (C. I. No. 926):			Ames, Iowa	3	10
Havre, Mont.	1	12	Lincoln, Nebr.	3	16
Archer, Wyo.	2	13	Crookston, Minn.	5	19
Union, Oreg.	2	10	Madison, Wis.	1	10
Charlottetown 80 (C. I. No. 2732):			Wisconsin Pedigree (C. I. No. 835):		
State College, Pa.	1	6	Storrs, Conn.	1	3
Hanna (C. I. No. 203):			Urbana, Ill.	1	5
Dickinson, N. Dak.	3	18	De Kalb, Ill.	2	6
Williston, N. Dak.	1	12	State College, Pa.	4	6
Gold (C. I. No. 1145):			Wooster, Ohio	4	5
East Lansing, Mich.	3	18	La Fayette, Ind.	5	9
Fort Collins, Colo.	3	24	Fargo, N. Dak.	5	8
Hannchen (C. I. No. 531):			Oderbrucker (C. I. No. 1272):		
Moccasin, Mont.	1	13	Ithaca, N. Y.	1	15
Burns, Oreg. (irrigated)	1	7	Ames, Iowa	1	10
Burns, Oreg. (dry land)	2	7	Madison, Wis.	3	19
Mandan, N. Dak.	3	6	Oderbrucker (C. I. No. 836):		
Williston, N. Dak.	3	12	De Kalb, Ill.	1	6
Eureka, S. Dak.	3	5	Edgeley, N. Dak.	1	4
Newell, S. Dak. (irrigated)	3	7	Ithaca, N. Y.	3	15
Newlands, Nev.	3	7	Urbana, Ill.	2	5
Dickinson, N. Dak.	4	18	Minsturd (C. I. No. 1556):		
Cottonwood, S. Dak.	4	5	St. Paul, Minn.	1	10
Newell, S. Dak. (dry land)	4	8	Grand Rapids, Minn.	1	8
Huntley, Mont.	4	5	Crookston, Minn.	1	19
Sheridan, Wyo.	5	8	Waseca, Minn.	3	8
White Smyrna (C. I. No. 195):			Odessa (C. I. No. 182):		
Mandan, N. Dak.	1	6	Brookings, S. Dak.	1	14
Garden City, Kans.	1	6	Eureka, S. Dak.	1	5
Sheridan, Wyo.	1	8	Highmore, S. Dak.	1	15
Aberdeen, Idaho (dry land)	1	7	Cottonwood, S. Dak.	1	5
Eureka, S. Dak.	2	5	Colby, Kans.	1	14
Cottonwood, S. Dak.	2	5	Garden City, Kans.	2	6
Huntley, Mont.	2	5	Tribune, Kans.	3	8
Akron, Colo.	2	10	Newell, S. Dak. (dry land)	5	8
Lincoln, Nebr.	4	16	Manhattan, Kans.	5	28
Tribune, Kans.	4	8	Trebi (C. I. No. 936):		
Hayes, Kans.	5	15	Newell, S. Dak. (irrigated)	1	7
Havre, Mont.	5	12	North Platte, Nebr.	1	6
Chico, Calif.	5	18	Mitchell, Nebr.	1	5
White Smyrna (C. I. No. 658):			Huntley, Mont.	1	5
Newell, S. Dak. (dry land)	1	8	Aberdeen, Idaho (irrigated)	1	9
Burns, Oreg. (dry land)	1	7	Union, Oreg.	1	10
Dickinson, N. Dak.	2	18	Columbia, Mo.	2	35
Burns, Oreg. (irrigated)	2	7	Sheridan, Wyo.	2	8
Archer, Wyo.	4	13	Newlands, Nev.	2	7
North Platte, Nebr.	5	6	Archer, Wyo.	3	13
White Smyrna (C. I. No. 910):			Ithaca, N. Y.	4	15
Moccasin, Mont.	2	15	Sandpoint, Idaho	4	6
Sandpoint, Idaho	2	6	Moscow, Idaho	4	23
Lind, Wash.	4	5	Aberdeen, Idaho (dry land)	4	7
Svansota (C. I. No. 1907):			Burns, Oreg. (irrigated)	4	7
Duluth, Minn.	1	6	Coast (C. I. No. 626):		
Grand Rapids, Minn.	4	8	Newlands, Nev.	1	7
Morris, Minn.	5	14	Davis, Calif.	1	11
Manchuria (C. I. No. 241):			Chico, Calif.	4	18
Ames, Iowa	4	10	Highmore, S. Dak.	5	15
Manchuria (C. I. No. 244):			Sandrel (C. I. No. 937):		
Edgeley, N. Dak.	2	4	Columbia, Mo.	1	35
Manhattan, Kans.	2	28	Archer, Wyo.	1	13
Fargo, N. Dak.	4	8	Aberdeen, Idaho (irrigated)	3	9

TABLE 97.—Five-year average performance of varieties of barley, 1917 to 1921, inclusive, shown by stations—Continued

Varieties and stations	Rank	Number compared	Varieties and stations	Rank	Number compared
Coast (C. I. No. 690):			Winter Club (C. I. No. 592):		
Tribune, Kans.	1	4	Moscow, Idaho.	1	23
Newell, S. Dak. (dry land)	2	8	Pullman, Wash.	1	13
Hays, Kans.	2	15	Burns, Oreg. (irrigated)	3	7
North Platte, Nebr.	3	6	Nephi, Utah	4	6
Colby, Kans.	3	14	Burns, Oreg. (dry land)	5	7
Garden City, Kans.	3	6	Tennessee Winter (C. I. No. 257):		
Akron, Colo.	3	10	State College, N. Mex.	1	6
Burns, Oreg. (dry land)	3	7	Athens, Ga.	2	14
State College, N. Mex.	3	6	Knoxville, Tenn.	2	4
Mandan, N. Dak.	4	6	Nephi, Utah	3	6
Newell, S. Dak. (irrigated)	4	7	Union Winter (C. I. No. 583):		
Manhattan, Kans.	4	28	Knoxville, Tenn.	1	4
Sheridan, Wyo.	4	8	Jackson, Tenn.	1	2
Archer, Wyo.	5	13	Wisconsin Winter (C. I. No. 519):		
Nephi, Utah	5	6	Pullman, Wash.	3	13
Burns, Oreg. (irrigated)	5	7	Roslyn, Va.	2	19
Beldi Giant (C. I. No. 2777):			Manhattan, Kans.	3	28
Aberdeen, Idaho (irrigated)	2	9	Athens, Ga.	3	14
Aberdeen, Idaho (dry land)	2	7	Moscow, Idaho	5	23
Davis, Calif.	2	11	Meloy (C. I. No. 1176):		
Pullman, Wash.	4	13	Lind, Wash.	2	5
Beldi Dwarf (C. I. No. 190):			Moccasin, Mont.	3	13
Chico, Calif.	3	18	Colby, Kans.	4	14
Hays, Kans.	4	15	Moro, Oreg.	5	15
Club Mariout (C. I. No. 261):			Lion (C. I. No. 923):		
Lincoln, Nebr.	1	16	La Fayette, Ind.	1	9
Hays, Kans.	1	15	State College, Pa.	2	6
Moro, Oreg.	1	15	De Kalb, Ill.	3	6
Chico, Calif.	1	18	Fargo, N. Dak.	3	8
Aberdeen, Idaho (dry land)	3	7	Columbia, Mo.	5	35
Mandan, N. Dak.	5	6	Aberdeen, Idaho (dry land)	5	7
Garden City, Kans.	5	6	Gatami (C. I. No. 575):		
Tribune, Kans.	5	8	Brookings, S. Dak.	3	14
Flynn (C. I. No. 1311):			Cottonwood, S. Dak.	3	5
Moro, Oreg.	4	15	Newell, S. Dak. (dry land)	3	8
			Hays, Kans.	3	15
			Eureka, S. Dak.	5	5

It will be noted that the names of some varieties occur a large number of times. Such varieties must possess high yielding capacity. Their range of adaptation perhaps is better shown in Table 97. In this table many of the prominent varieties are included in such a manner as to show the stations at which they have given high yields for this period. The rank at each place also is shown. The wide range of adaptation of White Smyrna already has been pointed out. The Coast varieties (C. I. Nos. 690 and 626) are shown to be good at Tribune, Kans., at Davis, Calif., at State College, N. Mex., and at Mandan, N. Dak. The range of Winter Club (C. I. No. 592) is limited to the Great Basin. The Oderbrucker and Manchuria varieties have done well at stations from Ithaca, N. Y., to Edgeley, N. Dak., and from Ames, Iowa, to Crookston, Minn. The good yields of Gatami (C. I. No. 575) have been limited to that section of the Great Plains from Hays, Kans., north to Eureka, S. Dak. It is interesting to note that the high yields from Hannchen (C. I. No. 531) and Svanhals (C. I. No. 187) were obtained at stations from Eureka, S. Dak., and Mandan, N. Dak., west to the Pacific coast. To judge from the origin of these varieties, it would be expected that their best adaptation would be farther east.

DISTRIBUTION OF VARIETIES BY EXPERIMENT STATIONS

It is obvious from the volume of the results reported that there has been a large expenditure of funds in the introduction, breeding, and testing of barley varieties in the United States and Canada. The question naturally arises as to how profitable this investment has been and to what extent the varieties so produced and so recommended are in cultivation to-day. A review of the situation leads to the conclusion that the investment has already been many times repaid and will continue to be repaid as long as barley is cultivated.

Four basic types—(1) Coast, (2) a barley closely allied to the Manchuria, (3) Tennessee Winter, and (4) Stavropol—found their way into cultivation without the assistance of national or State investigations. Coast was so well adapted to the Western States that for most of the period of experimentation it has been the variety recommended by experiment stations. It is only recently that superior sorts have been found for local conditions in the West. The introduction of Manchuria and Oderbrucker, which were tested and distributed by the stations, resulted in displacing the related form mentioned above as basic type No. 2. These varieties now constitute the greater part of the acreage of the upper Mississippi Valley and adjoining sections of Canada.

Stavropol probably was imported by the Office of Foreign Seed and Plant Introduction of the United States Department of Agriculture and was distributed by the Kansas Agricultural Experiment Station. It is more than plausible to assume, however, that this same barley was brought repeatedly from southern Russia by Russian immigrants and that there was a large acreage of this type of barley in northwestern Kansas before any was sent out by the State station.

In comparing the expense with the returns in barley studies it must be realized that the great bulk of the expenditure has been in the last few years. There has not been time to secure returns on those varieties recently produced. Club Mariout was brought to America in 1904. After years of testing it is just now becoming of commercial importance. There are many varieties still with a very limited acreage which have great potential importance. Trebi, now extensively grown in southern Idaho and the irrigated section of western Nebraska, is meeting with popular favor and is giving yields far greater than those of the varieties displaced. California Mariout is advantageously cultivated in parts of the San Joaquin Valley which are too dry for high yields of Coast barleys. Many superior selections of Oderbrucker and Manchuria have been distributed in Wisconsin, Minnesota, and North Dakota. Alpha, Horn, O. A. C. 21, Minsturdi, Svansota, Featherston, and many more such new productions have yielded extremely well in recent tests and are now appearing in field agriculture. The potential value of such varieties in the next 10 years is very great. Back of these, on the field plats and breeding nurseries, are many others with which much of the work of breeding and testing has already been accomplished.

INDEX

Experiments in Canada, by Provinces

Page	Page	Page			
Alberta.....	41-46	Northwest Territory.....	46	Prince Edward Island.....	19-20
British Columbia.....	46-50	Nova Scotia.....	20-24	Quebec.....	25-26
Manitoba.....	32-35	Ontario.....	26-31	Saskatchewan.....	36-41
New Brunswick.....	24-25				

Experiments in the United States, by States

	Page		Page		Page
Arkansas	151	Michigan	57	Pennsylvania	54-55
California	133-139	Minnesota	66-76	Rhode Island	51
Colorado	116-120	Missouri	77-78	South Carolina	146
Connecticut	52	Montana	110-114	South Dakota	90-100
Georgia	146-148	Nebraska	100-103	Tennessee	150-151
Idaho	120-125	Nevada	133	Texas	132-153
Illinois	59-60	New Hampshire	51	Utah	125-126
Indiana	58-59	New Mexico	153-154	Virginia	141-146
Iowa	76	New York	52-54	Washington	126-127
Kansas	103-110	North Dakota	75-90	West Virginia	148
Kentucky	149	Ohio	56	Wisconsin	60-65
Maine	50-51	Oklahoma	151-152	Wyoming	114-116
Maryland	140	Oregon	127-133		

Experiments at Agricultural Experiment Stations in the United States and Canada

Page	Page	Page
Abdeen, Idaho. 122-125, 209-211	Fort Providence, North-west Territory. 46	Mitchell, Nebr. 103, 208, 210
Agassiz, British Columbia. 46-49	Fort Resolution, North-west Territory. 46	Moccasin, Mont. 111-112, 209-211
Akron, Colo. 118-120, 209-211	Fort Smith, Northwest Territory. 46	Modesto, Calif. 136
Amarillo, Tex. 152-153	Fort Vermilion, Alberta. 45-46	Morgantown, W. Va. 148
Ames, Iowa. 76, 208, 210	Frederickton, New Brunswick. 24-25	Moro, Oreg. 127-130, 209, 211
Archer, Wyo. 115-116, 209-211	Garden City, Kans. 109-110, 210, 211	Morris, Minn. 75-76, 208, 210
Arlington Experiment Farm. (See Rosslyn, Va.)	Grand Rapids, Minn. 74, 208, 210	Moscow, Idaho. 121-122, 209-211
Ashburn, Ga. 147-148	Grouard, Alberta. 46	Nappan, Nova Scotia. 20-24
Athens, Ga. 146-147, 209-211	Havre, Mont. 110-111, 209, 210	Nephi, Utah. 125-126, 209, 211
Beaver Lodge, Alberta. 46	Hays, Kans. 106-108, 208, 210, 111	Newell, S. Dak. 208, 210, 211
Blacksburg, Va. 145-146, 209	Highmore, S. Dak. 96-98, 208, 210	New Hope, Tex. 152
Bozeman, Mont. 113-114	Huntley, Mont. 112-113, 209, 210	Newlands, Nev. 133, 209, 210
Brandon, Manitoba 32-35	Indian Head, Saskatchewan. 36-39	North Platte, Nebr. 102-103, 208, 210, 211
Brookings, S. Dak. 90-94, 208, 210, 211	Invermere, British Columbia. 50	Orono, Me. 50-51
Burns, Oreg. 131-133, 209-211	Ithaca, N. Y. 52-54, 208, 210	Ottawa, Ontario. 26-31
Cap Rouge, Quebec. 25-26	Jackson, Tenn. 151, 209, 211	Pullman, Wash. 126, 209, 211
Carthage, Mo. 77-78	Kentville, Nova Scotia. 24	Quitman, Ga. 148
Channing, Tex. 153	Kington, R. I. 51	Rosslyn, Va. 141-144, 209, 211
Charlottetown, Prince Edward Island. 19-20	Knoxville, Tenn. 150-151, 209, 211	Rosthern, Saskatchewan. 40
Chico, Calif. 133-136, 209-211	Lacombe, Alberta. 44-45	Saint Bernard, Alberta. 46
Clemson, S. C. 146	La Fayette, Ind. 58-59, 208, 210, 211	Sainte Anne de la Pocati- tierre, Quebec. 25
Colby, Kans. 109, 209-211	Langdon, N. Dak. 84	Saint Paul, Minn. 66-73, 208, 210
College Park, Md. 140	Lethbridge, Alberta. 41-43	Sandpoint, Idaho. 120, 209, 210
Columbia, Mo. 77-78, 208, 210, 211	Lewistown, Mo. 77-78	Scott, Saskatchewan. 41
Cottonwood, S. Dak. 98, 208, 210, 211	Lexington, Ky. 149	Sheridan, Wyo. 114-115, 209-211
Crookston, Minn. 75, 208, 210	Lincoln, Nebr. 100-101, 208, 210, 211	Sidney, British Columbia. 50
David, Calif. 137-139, 209-211	Lind, Wash. 127, 209-211	State College, N. Mex. 153-154, 209, 211
De Kalb, Ill. 60, 208, 210, 211	McPherson, Kans. 105-106	State College, Pa. 64-55, 208, 210, 211
Dickinson, N. Dak. 85-87, 208, 210	Madison, Wis. 60-65, 208, 210	Stillwater, Okla. 151-152
Duluth, Minn. 73-74, 208, 210	Mandan, N. Dak. 85, 208, 210, 211	Storrs, Conn. 52, 208, 210
Durham, N. H. 51	Manhattan, Kans. 103-105, 208, 210, 211	Tribune, Kans. 110, 209-211
East Lansing, Mich. 57, 208	Maryville, Mo. 77-78	Urbana, Ill. 59-60, 208, 210
Edgeley, N. Dak. 82-84, 208, 210		Union, Oreg. 130-131, 209, 210
Eureka, S. Dak. 95, 208, 210, 211		Warrensburg, Mo. 77-78
Fargo, N. Dak. 78-81, 208, 210, 211		Waseca, Minn. 73, 208, 210
Fayetteville, Ark. 151		Williston, N. Dak. 88-90, 208, 210
Fort Collins, Colo. 116-117, 209, 210		Wooster, Ohio. 56, 208, 210

Varieties included in Botanical and Yield Tables

	Page		Page		Page
Abed.....	121, 189	Beldi Giant.....	111,	Canadian Lake Shore.....	64, 69, 195
Abina.....	134, 192	122-124, 126, 128, 139, 196,		Canadian Thorpe.....	19,
Abyssinia:		206, 209, 211.		22, 24, 25, 28, 31, 34, 35, 38, 39,	
C. I. No. 355.....	134, 180	Benghazi.....	139, 183	42-45, 47, 49, 50, 66, 88, 111,	
361.....	134, 185, 180	Bere.....	29, 195	113, 117, 184, 206.	
669.....	149, 183	Berkley.....	113, 196	Cape.....	53, 181
672.....	149, 183	Bernard.....	62, 63, 66, 80, 88, 185	Carter.....	80, 202
Abyssinian:		Bestehorn.....	27, 33, 93, 180	Cartouch.....	143, 187
C. I. No. 673.....	135, 183	Bestehorn Giant.....	142, 146, 150, 181	Caswell.....	51, 204
938.....	134, 186	Bestehorn Imperial.....	91, 96, 176	Catania.....	93, 181
941.....	134, 187	Bestehorn Kaiser.....	29, 194	Caucasian:	
943.....	134, 187	Bevede.....	128, 192	C. I. No. 90.....	76, 103, 106, 177
946.....	135, 187	Bhagamany Hills.....	22, 33, 37, 194	714.....	97, 184
950.....	134, 187	Binder.....	30, 190	2724.....	30, 195
1236.....	135, 188	Black:		Caveda.....	143, 181
2313.....	134, 192	C. I. No. 256.....	55, 62, 63, 92, 179	Centre.....	93, 180
2314.....	134, 192	2726.....	30, 69, 195	Champion.....	23,
Ace.....	101, 128, 190, 208	2936.....	70, 202	29, 33, 38, 42-45, 47, 194	
Aker.....	70, 73-75, 190, 208	Black Abyssinian.....	128, 192	Champion of Vermont:	
Alaska.....	143, 146, 181	Black Algerian:		C. I. No. 1892.....	51-54,
Albacete.....	63, 106, 107, 140, 142, 177	C. I. No. 708.....	134, 184	2899.....	62, 63, 66, 80, 190
Albert.....	19,	2296.....	128, 192	19, 54, 55, 57,	
23, 29, 35, 38, 42-44, 48, 189		Black Arabian.....	107, 142, 178	Charlottetown 80.....	58, 65, 117, 195, 205, 208, 210
Algerian.....	122, 188	Black Egyptian.....	145, 188	Cheney:	
Alpha.....	53, 54, 77, 135, 187, 206	Blackhall:		Chevalier:	
Altamura.....	62, 63, 177	C. I. No. 878.....	109, 115,	C. I. No. 23.....	91, 96, 176
Aquila.....	93, 181	117-119, 133, 184, 206, 209		35.....	91, 94, 96, 176
Arabel.....	142, 185	2087.....	118, 191	156.....	64, 69, 92, 125, 128,
Archer Chevalier.....	29, 195	Black Hull-less:		134, 142, 144, 147, 149, 177	
Arequipa.....	135, 189	C. I. No. 596.....	23,	267.....	138, 179
Argentine.....	143, 147, 148, 178, 209	27, 35, 38, 48, 51, 54, 56,		278.....	122, 134, 135, 179
Argyle.....	23, 29, 34, 38, 44, 48, 193	58, 62, 63, 66, 71, 76, 77,		290.....	138, 139, 179
Arizona.....	57, 197	80, 83, 101, 102, 104-107,		612.....	128, 142, 182
Arlington Awnless:		111, 113, 117, 125, 128, 130,		1142.....	100, 133, 188, 209
C. I. No. 702.....	50,	132, 134, 142, 145, 182, 206		1170.....	69, 188
57, 65, 96, 118, 121, 140,		735.....	132, 134, 184	1886.....	101, 190
143, 145-149, 184.		1106.....	70, 115, 116, 187	2178.....	104, 191
2160.....	143, 191	Black Japan.....	25, 29, 39-41, 195	2651.....	37, 193
2161.....	143, 191	Black Russian.....	134, 143, 147, 184	2768.....	113, 196
2162.....	143, 191	Black Smyrna.....	63, 82, 91, 96,	2771.....	113, 196
Arnt.....	88, 195	104, 106, 107, 136, 142, 177		2802.....	51,
Australian White:		Black Winter.....	140, 150, 180	55, 62, 63, 66, 80, 197	
C. I. No. 925.....	53,	Blarney.....	135, 189	2877.....	68, 200
77, 122, 134, 186		Blue.....	126, 127, 130, 188	2910.....	69, 201
2898.....	68, 201	Blue Long Head.....	23,	3165.....	145, 204
Austrian.....	107, 108, 191	24, 29, 31, 34, 35, 38, 39,		Chevalier II.....	19,
Baitori.....	91, 176	42-44, 48, 49, 194, 206.		23, 25, 29, 35, 39, 40, 42-44, 48,	
Baku.....	120-122, 187, 206, 209	Blue Ribbon.....	67, 74, 82, 143, 182	53, 57, 58, 62, 63, 65, 67, 71	
C. I. No. 210.....	106, 178	Blue Short Head.....	29, 194	81-83, 86, 92, 95, 96, 98-100,	
253.....	92, 179	Bohemian:		104, 113, 117, 122, 128-130, 132,	
333.....	22,	C. I. No. 27.....	82,	138, 142, 152, 178, 205, 208-210	
27, 33, 37, 39, 44, 93, 180		91, 94-97, 111, 176, 209		Cheyney.....	66, 93, 180
709.....	115,	32.....	82, 91, 96, 98, 176	Chili Brewing:	
118, 119, 128, 184, 206		188.....	92, 128, 152, 177	C. I. No. 2297.....	128, 192
2920.....	69, 202	204.....	92, 138, 178	2772.....	113, 196
Barbary.....	103, 149, 183, 208	270.....	138, 179	Chili Chevalier.....	113, 183
Bark.....	70, 117, 196	284.....	138, 179	Chili Common:	
Bavarian:		285.....	138, 179	C. I. No. 663.....	86,
C. I. No. 158.....	83, 91, 177	293.....	138, 179	87, 113, 121, 149, 183	
159.....	83,	344.....	138, 180	2186.....	118, 191
91, 107, 128, 134, 142, 147,		933.....	53, 186	Chinese Hull-less.....	29, 195
149, 177.		Bohman.....	70, 73-75, 202	Chousk.....	29, 195
160.....	92, 122, 124, 142, 177	Boiton:		Claude.....	19,
208.....	92, 113, 154, 178	C. I. No. 177.....	22,	23, 29, 34, 38, 42-45, 48, 64, 189	
248.....	107, 178	28, 34, 38, 47, 92, 177		Clifford.....	19,
Baxter.....	22,	922.....	53, 186	23, 29, 34, 39, 42-44, 48, 190	
27, 33, 37, 47, 53, 66, 80, 88,		Bonanza:		Club Mariout:	
92, 106, 177.		C. I. No. 1800.....	104, 190	C. I. No. 261.....	63,
Beardless:		3176.....	104, 204	82, 83, 85, 87, 93, 96, 97,	
C. I. No. 639.....	102, 130, 183	3177.....	104, 204	101, 103, 104, 106-112,	
1797.....	54, 190	Brasserie.....	93, 180	115, 122, 124, 126, 128-	
2646.....	27, 33, 37, 193	Brewers Delight.....	55, 203	130, 132-136, 153, 179,	
3144.....	75, 203	Brewers Favorite.....	29, 35, 41, 182	206, 208, 209, 211.	
3145.....	121, 203	Brome.....	23, 29, 35, 38, 48, 193	932.....	53,
Beardless 6.....	150, 195, 209	Brush.....	67, 198	77, 86, 122, 127, 186	
Beardless 20.....	149, 150, 151, 195, 209	Bulgarian.....	124, 125, 181, 209	Clyde.....	111, 185
Beaver.....	19,	California Centgener.....	121, 203	Coast:	
22, 25, 28, 34, 38, 40, 42-44,		California Mariout.....	104, 109,	C. I. No. 276.....	134, 179
47, 49, 190.		128, 135, 139, 189, 206, 209		549.....	106, 181
Beldi Dwarf.....	62,	California Prolific:		626.....	86,
63, 82, 91, 96, 97, 106-108, 113,		C. I. No. 651.....	125, 183	93, 96, 97, 128, 129, 133-	
114, 121, 128, 129, 132, 134,		2649.....	23, 28,	136, 138, 139, 182, 206,	
135, 136, 138, 152, 177, 206,		33, 37, 47, 53, 113, 114, 193		208-210.	
208, 209, 211.		Canada Winter.....	142, 184	650.....	125, 183
		Canadian.....	53, 192		

	Page		Page		Page
Coast—Continued		Featherston—Continued		Hanchamont	53,
C. I. No. 690	53,	C. I. No. 1118	53, 77, 187	C. I. No. 192	54, 77, 134, 187, 206, 208
55, 64, 67, 71, 85, 99, 100,		1119	53, 77, 187	Hankow:	
102-105, 107-112, 115-		1120	53,	C. I. No. 192	107, 152, 177
119, 122, 124, 125, 132,		54, 58, 77, 187, 206, 208		197	107, 134, 142, 152, 177
133, 154, 183, 206, 208,		Fichtel Mountain	29, 194	Hanna:	
209, 211.		Flaxton	88, 195	C. I. No. 24	82, 91,
691	96, 134, 183	Fleche	143, 189	95, 96, 98, 99, 102, 115	
1249	126, 127, 188, 209	Flynn	109, 111,	118, 128, 132, 176,	
2235	122, 192	115, 128-130, 189, 206, 209, 211		26	91, 96, 176
2301	128, 192	Francian:		28	91, 96, 176
2305	132, 133, 192	C. I. No. 679	97,	30	91, 96, 176
2306	134, 192	680	128, 134, 149, 183	33	91, 96, 176
2307	134, 192	103, 111, 112, 183, 208, 209		34	91, 95, 96, 98, 176
2312	134, 192	Frankish:		203	58, 62,
2785	117, 196, 209	C. I. No. 207	67,	63, 65, 67, 76, 80, 82, 83,	
2789	117, 196	82, 92, 113, 154, 178		86-89, 91, 95, 96, 98, 99,	
2790	117, 196, 209	269	138, 179	106, 107, 115, 118, 119,	
2791	117, 196	282	62, 63, 138, 179	138, 142, 152, 153, 178,	
2821	139, 197, 209	295	76, 138, 179, 208	206, 208, 210.	
2921	69, 202	953	53, 77, 135, 187	287	138,
Coeleste	93, 180	Frankish Brewing	68, 200	142, 144, 147, 149, 179	
Cole	42, 43, 195	French Chevalier:		301	138, 179
Colless	117, 196, 209	C. I. No. 175	19,	319	53, 64,
Common:		22, 24, 28, 31, 34, 35, 38, 42-		67, 71, 74, 75, 80, 145, 179	
C. I. No. 184	22,	44, 47, 52, 62, 63, 66, 80,		416	93, 134, 181
27, 33, 38, 47, 66, 82, 84,		177, 205		906	53, 70,
92, 106, 177.		2900	68,	77, 122, 124, 128, 185	
3112	55, 203	73-75, 201, 208, 210		942	134, 187
Common Two-Row	55, 203	Fulton	23, 29, 34, 39, 48, 194	1122	53, 187
Composite	70,	Fünfstettener	138, 179	1554	54, 189
93, 96, 118, 122, 188		Gambirinus	30, 93, 180	1854	53, 190
Coreth	91, 176	Garfield	23, 29, 34, 38, 48, 193	2309	134, 192
Cortile	53, 187	Garton:		2316	134, 192
Cortney	134, 187	C. I. No. 263	64, 106, 179	2773	113, 196
Crawford:		264	64, 106, 179	2784	117, 196, 209
C. I. No. 2810	57, 197	265	106, 179	2786	117, 196
3136	57, 203, 208	266	64, 69, 179	2787	117, 196
3137	57, 203, 208	Garton 68	35, 104, 195	2788	117, 196
3138	57, 203, 208	Garton 936	64, 195	2886	68, 200
3139	57, 203, 208	Garton 986	58, 113, 117, 183	2902	68, 201
Crimean Hull-less	92, 179	Garton 5001	101, 197	2919	69, 202
Crop Nursery	67, 198	Garton 5002	101, 197	Hannchen	19,
Culver	66, 80, 197	Garton 5003	101, 197	23-25, 29, 31, 35, 39-44, 48, 49,	
Cusado	142, 185	Garton 5004	101, 197	53, 54, 57, 58, 62, 63, 65, 67, 71,	
Daniels	69, 200	Gatami	58,	75, 81, 82-89, 92, 94-101, 103,	
Danish Chevalier	19,	70, 82, 86-89, 93-99, 101, 102,		104, 107-109, 111-119, 121-	
22, 27, 33, 37, 39, 42-44, 47,		104-111, 115, 118, 122, 124,		125, 127-130, 132-135, 138,	
49, 66, 92, 177.		125, 128, 132, 134, 142, 145,		139, 142, 147, 181, 206, 208-210.	
Doitsu	91, 96, 176	181, 206, 208, 211		Han River:	
Douglas	28, 194	Giant	93, 197	C. I. No. 206	103,
Duckbill	22,	Gisborne	69, 187	111, 120-123, 140, 143,	
25, 28, 33, 37, 40-43, 47, 52, 190		Gold	19,	146, 178, 206, 209.	
Duckbill Bald	67, 198	23, 25, 30, 35, 39, 42-44, 48, 53,		2163	143, 144, 191
Dufur	128, 192	57, 58, 65, 70, 93, 94, 96, 101,		Hanseel Hull-less	128,
Dunham	23, 28, 34, 38, 44, 48, 193	104, 105, 117, 188, 206, 208,		134, 142, 144, 149, 184	
Dutch	27, 194	209, 210		23, 29, 34, 39, 48, 193	
Dutrow Winter	140, 197	Golden Drop	66, 117, 191	Heil Hanna 1	52, 134, 149, 183
Eagle	53, 70, 77, 122, 145, 186	Golden Grain	22,	111, 134, 135, 183	
Early Chevalier	19,	27, 34, 37, 47, 66, 134, 182		Heil Hanna 3	149, 183
25, 30, 39-44, 135, 206		Golden Melon:		Heil Hanna 4	52, 86, 183
Early Indian	30, 40, 41, 44, 190	C. I. No. 48	91, 95-98, 176	Hero	135, 189
Early Lapland	92, 179	411	22, 27,	Highland Chief:	
Early Mintage	22, 27, 37, 193	33, 37, 47, 51, 55, 62, 63, 181		C. I. No. 209	62, 63, 92, 178
Elipse	29, 195	958	134, 187	883	58,
Elfy	117, 197	Golden Queen:		66, 71, 80, 82, 88, 130, 155	
Ellis	107-110, 191, 209	C. I. No. 538	62	80, 202	
Emerson	34, 193	63, 66, 74, 80, 82, 181		Highland Scotch	80, 202
Empire	23, 29, 34, 38, 42-44, 48,	1511	64, 189	Himalaya:	
Empress	66, 197	2753	64, 195	C. I. No. 254	92, 179
English Chevalier	19, 195	2901	68, 201	322	92, 179
English Malting	22, 27, 33, 37, 193	Goldfoil:		619	88, 182
Erfurt White	30, 189	C. I. No. 162	82, 92, 177	620	52,
Escourgeon	25, 29, 195	928	53,	53, 54, 62, 63, 68, 99, 100,	
Eureka	126, 127, 188	111, 122, 134, 186, 209		104, 107, 111, 113-115,	
Evans	53, 62, 63, 80,	Gold Medal	138, 179	118, 127, 128, 182, 206,	
81, 82, 84, 88, 111, 113, 128, 185		Goldthorpe:		209.	
Excelsior:		C. I. No. 327	22,	1312	135, 189
C. I. No. 1248	126, 127, 188	27, 33, 37, 47, 66, 80, 93, 180		1585	104, 190
2684	23,	409	138, 180	1934	22, 27,
29, 33, 35, 38, 48, 194		Gordon	23,	30, 35, 38, 42-44, 47, 191	
Famesh	86, 139, 182	29, 35, 39, 42-44, 48, 193		2298	128, 192
Fan:		Gotland	64, 69, 195	2299	128, 129, 192
C. I. No. 255	92, 179	Great Beardless	58, 196	Hodge	139, 187, 209
341	27, 33, 93, 180	Greece	140, 142, 146-148, 178	Hofbrau	30, 93, 180
Featherston:		Greek	27, 93,	Holland	111, 135, 187, 209
C. I. No. 911	53,	Gutekorn	64, 68, 113, 130, 132, 182	Hoover Hull-less	80, 202
70, 122, 145, 185, 209				Horn:	
954	53, 135, 187			C. I. No. 31	82,
				91, 95, 96, 98, 176	

	Page		Page		Page	
Horn—Continued.		Kitzing:		Manchuria—Continued.		
C. I. No. 926.....	53, 77, 111, 115, 116, 122, 123, 128, 130, 186, 205, 209, 210.	C. I. No. 167.....	82, 93, 177 189..... 67, 82 92, 113, 114, 138, 152, 177 201..... 67, 92, 107, 138, 178 272..... 138, 179	C. I. No. 2820.....	101, 197 2823..... 54, 69, 75, 197, 208 2827..... 67, 197 2831..... 67, 198 2835..... 67, 198 2836..... 67, 198 2837..... 67, 198 2848..... 67, 199 2849..... 67, 199 2850..... 67, 199 2851..... 67, 199 2852..... 67, 199 2853..... 67, 199 2854..... 67, 199 2855..... 67, 199 2856..... 67, 199 2857..... 67, 199 2858..... 67, 199 2859..... 69, 199 2860..... 69, 199 2861..... 67, 199 2862..... 69, 199 2863..... 67, 199 2864..... 69, 199 2865..... 67, 199 2866..... 68, 199 2867..... 68, 199 2868..... 68, 199 2869..... 68, 199 2870..... 68, 199 2871..... 68, 199 2872..... 68, 199 2873..... 68, 199 2874..... 68, 200 2875..... 68, 200 2876..... 68, 200 2877..... 68, 200 2878..... 68, 200 2879..... 68, 200 2880..... 68, 200 2881..... 68, 200 2882..... 68, 200 2883..... 68, 200 2884..... 68, 200 2885..... 68, 200 2886..... 68, 200 2887..... 68, 200 2888..... 68, 200 2889..... 68, 200 2890..... 68, 200 2891..... 68, 200 2892..... 68, 200 2893..... 68, 201 2894..... 68, 201 2895..... 68, 201 2896..... 68, 201 2897..... 68, 201 2898..... 68, 201 2899..... 68, 201 2900..... 68, 201 2901..... 68, 201 2902..... 68, 201 2903..... 68, 201 2904..... 68, 201 2905..... 68, 201 2906..... 68, 201 2907..... 68, 201 2908..... 68, 201 2909..... 68, 201 2910..... 68, 201 2911..... 68, 201 2912..... 68, 201 2913..... 68, 201 2914..... 68, 201 2915..... 68, 201 2916..... 68, 201 2917..... 68, 201 2918..... 68, 201 2919..... 68, 201 2920..... 68, 201 2921..... 68, 201 2922..... 68, 201 2923..... 68, 201 2924..... 68, 201 2925..... 68, 201 2926..... 68, 201 2927..... 68, 201 2928..... 68, 201 2929..... 68, 201 2930..... 68, 201 2931..... 68, 201 2932..... 68, 201 2933..... 68, 201 2934..... 68, 201 2935..... 68, 201 2936..... 68, 201 2937..... 68, 201 2938..... 68, 201 2939..... 68, 201 2940..... 68, 201 2941..... 68, 201 2942..... 68, 201 2943..... 68, 201 2944..... 68, 201 2945..... 68, 201 2946..... 68, 201 2947..... 68, 201 2948..... 68, 201 2949..... 68, 201 2950..... 68, 201 2951..... 68, 201 2952..... 68, 201 2953..... 68, 201 2954..... 68, 201 2955..... 68, 201 2956..... 68, 201 2957..... 68, 201 2958..... 68, 201 2959..... 68, 201 2960..... 68, 201 2961..... 68, 201 2962..... 68, 201 2963..... 68, 201 2964..... 68, 201 2965..... 68, 201 2966..... 68, 201 2967..... 68, 201 2968..... 68, 201 2969..... 68, 201 2970..... 68, 201 2971..... 68, 201 2972..... 68, 201 2973..... 68, 201 2974..... 68, 201 2975..... 68, 201 2976..... 68, 201 2977..... 68, 201 2978..... 68, 201 2979..... 68, 201 2980..... 68, 201 2981..... 68, 201 2982..... 68, 201 2983..... 68, 201 2984..... 68, 201 2985..... 68, 201 2986..... 68, 201 2987..... 68, 201 2988..... 68, 201 2989..... 68, 201 2990..... 68, 201 2991..... 68, 201 2992..... 68, 201 2993..... 68, 201 2994..... 68, 201 2995..... 68, 201 2996..... 68, 201 2997..... 68, 201 2998..... 68, 201 2999..... 68, 201 3000..... 68, 201	
Horsford:		Large-Grained Winter.....	142, 149, 180	Leader.....	30, 195	
C. I. No. 507.....	51, 53, 59, 60, 63, 66, 71, 76, 77, 80-82, 84, 86, 93, 97, 101, 104, 106, 107, 109, 110, 117, 153, 181, 206, 208.	Leslie.....	23, 28, 34, 38, 48, 193	Lion.....	53, 55, 57-60, 70, 77, 81, 117, 124, 186, 206, 208, 209, 211.	
610.....	128, 132, 133, 149, 182	Logan.....	23, 28, 34, 38, 44, 48, 193	Louden.....	135, 189	
877.....	102, 115, 116, 118, 119, 184	Luth:		C. I. No. 908.....	53, 70	
1271.....	55, 58, 81, 88, 117, 189	Lytton.....	77, 122, 134, 145, 185	972.....	33, 77, 187	
1474.....	104, 105, 189	Mahan.....	29, 194	Lytton.....	29, 194	
1775.....	126, 127, 190, 209	Mahrische.....	128, 188	Mahan.....	128, 188	
1778.....	104, 190	Malting:		Mahrische.....	53, 70, 75, 134, 186	
1807.....	104, 190	C. I. No. 1129.....	77, 188	Malting:		
3171.....	104, 204	2909.....	69, 201	C. I. No. 1129.....	77, 188	
Hull-less:		Maltster.....	29, 69, 182	2909.....	69, 201	
C. I. No. 649.....	125, 183	Mammoth Winter:		C. I. No. 220.....	143, 147- 149, 178, 209	
2812.....	154, 197	C. I. No. 220.....	143, 147- 149, 178, 209	410.....	140, 180	
2939.....	80, 202	Manchuria:		410.....	140, 180	
2948.....	81, 203	C. I. No. 235.....	106, 178	Manchuria:		
3158.....	145, 204	237.....	106, 178	C. I. No. 235.....	106, 178	
Hungarian Hull-less.....	62, 63, 197	240.....	106, 178	237.....	106, 178	
Hurst.....	135, 189	241.....	53, 55, 58, 62, 63, 65, 66, 74, 76, 80, 82, 84, 86, 88, 91, 94- 96, 98-100, 115, 116, 128, 178, 206, 208, 210.	241.....	53, 55, 58, 62, 63, 65, 66, 74, 76, 80, 82, 84, 86, 88, 91, 94- 96, 98-100, 115, 116, 128, 178, 206, 208, 210.	
Hybrid:		244.....	29, 51- 53, 58, 62, 63, 66, 71, 73- 75, 77, 80-88, 91, 95-100, 102-104, 107-115, 118, 119, 122-124, 128, 132, 134, 153, 178, 206, 208-210.	244.....	29, 51- 53, 58, 62, 63, 66, 71, 73- 75, 77, 80-88, 91, 95-100, 102-104, 107-115, 118, 119, 122-124, 128, 132, 134, 153, 178, 206, 208-210.	
C. I. No. 2636.....	54, 192	329.....	93, 180	329.....	93, 180	
2637.....	54, 192	613.....	142, 182	613.....	142, 182	
2833.....	67, 198	643.....	183	643.....	183	
2834.....	67, 198	882.....	53, 80, 88, 89, 185, 208	882.....	53, 80, 88, 89, 185, 208	
2838.....	67, 71, 198	956.....	53, 77, 187, 208	956.....	53, 77, 187, 208	
2839.....	67, 198	1125.....	77, 188	1125.....	77, 188	
2840.....	67, 198	1178.....	53, 188	1178.....	53, 188	
2841.....	67, 198	1189.....	70, 73-75, 188, 208	1189.....	70, 73-75, 188, 208	
2842.....	67, 198	1275.....	64, 104, 107, 122, 128, 132, 189, 208	1275.....	64, 104, 107, 122, 128, 132, 189, 208	
2843.....	67, 198	1472.....	104, 105, 189	1472.....	104, 105, 189	
2844.....	67, 198	1473.....	104, 107, 189, 208	1473.....	104, 107, 189, 208	
2845.....	67, 198	1478.....	70, 73-75, 189, 208	1478.....	70, 73-75, 189, 208	
2846.....	67, 198	1562.....	101, 190	1562.....	101, 190	
2847.....	67, 198	1580.....	101, 190	1580.....	101, 190	
2875.....	68, 200	2155.....	93, 191	2155.....	93, 191	
2876.....	68, 200	2156.....	93, 191, 208	2156.....	93, 191, 208	
2888.....	68, 200	2157.....	93, 191	2157.....	93, 191	
2889.....	68, 200	2181.....	104, 191	2181.....	104, 191	
2890.....	68, 200	2182.....	104, 191	2182.....	104, 191	
2891.....	68, 200	2330.....	58, 68, 73-75, 101, 102, 117, 192, 206, 208, 210	2330.....	58, 68, 73-75, 101, 102, 117, 192, 206, 208, 210	
2892.....	68, 200	2517.....	93, 96, 192	2517.....	93, 96, 192	
2893.....	68, 201	2638.....	54, 192	2638.....	54, 192	
2894.....	68, 201	2756.....	64, 195	2756.....	64, 195	
2895.....	68, 201	2758.....	64, 195	2758.....	64, 195	
2896.....	68, 201	2760.....	64, 195	2760.....	64, 195	
2897.....	68, 201	2762.....	64, 196	2762.....	64, 196	
2898.....	68, 201	2764.....	64, 196	2764.....	64, 196	
2899.....	68, 201	2774.....	113, 196	2774.....	113, 196	
2900.....	68, 201	2778.....	56, 196	2778.....	56, 196	
2901.....	68, 201	2783.....	117, 196	2783.....	117, 196	
2902.....	68, 201					
2903.....	68, 201					
2904.....	68, 201					
2905.....	68, 201					
2906.....	68, 201					
2907.....	68, 201					
2908.....	68, 201					
2909.....	68, 201					
2910.....	68, 201					
2911.....	68, 201					
2912.....	68, 201					
2913.....	68, 201					
2914.....	68, 201					
2915.....	68, 201					
2916.....	68, 201					
2917.....	68, 201					
2918.....	68, 201					
2919.....	68, 201					
2920.....	68, 201					
2921.....	68, 201					
2922.....	68, 201					
2923.....	68, 201					
2924.....	68, 201					
2925.....	68, 201					
2926.....	68, 201					
2927.....	68, 201					
2928.....	68, 201					
2929.....	68, 201					
2930.....	68, 201					
2931.....	68, 201					
2932.....	68, 201					
2933.....	68, 201					
2934.....	68, 201					
2935.....	68, 201					
2936.....	68, 201					
2937.....	68, 201					
2938.....	68, 201					
2939.....	68, 201					
2940.....	68, 201					
2941.....	68, 201					
2942.....	68, 201					
2943.....	68, 201					
2944.....	68, 201					
2945.....	68, 201					
2946.....	68, 201					
2947.....	68, 201					
2948.....	68, 201					
2949.....	68, 201					
2950.....	68, 201					
2951.....	68, 201					
2952.....	68, 201					
2953.....	68, 201					
2954.....	68, 201					
2955.....	68, 201					
2956.....	68, 201					
2957.....	68, 201					
2958.....	68, 201					
2959.....	68, 201					
2960.....	68, 201					
2961.....	68, 201					
2962.....	68, 201					
2963.....	68, 201					
2964.....	68, 201					
2965.....	68, 201					
2966.....	68, 201					
2967.....	68, 201					
2968.....	68, 201					
2969.....	68, 201					
2970.....	68, 201					
2971.....	68, 201					
2972.....	68, 201					
2973.....	68, 201					
2974.....	68, 201					
2975.....	68, 201					
2976.....	68, 201					
2977.....	68, 201					
2978.....	68, 201					
2979.....	68, 201					
2980.....	68, 201					
2981.....	68, 201					
2982.....	68, 201					
2983.....	68, 201					
2984.....	68, 201					
2985.....	68, 201					
2986.....	68, 201					
2987.....	68, 201					
2988.....	68, 201					
2989.....	68, 201					
2990.....	68, 201					
2991.....	68, 201					
2992.....	68, 201					
2993.....	68, 201					
2994.....	68, 201					
2995.....	68, 201					
2996.....	68, 201					
2997.....	68, 201					
2998.....	68, 201					
2999.....	68, 201					
3000.....	68, 201					

	Page		Page		Page
Maryland Winter.....	56,	Oderbrucker—Continued.		Primus.....	29,
124, 126, 128, 134, 140, 142,		C. I. No. 1146.....	93,	39, 44, 53, 56, 58, 62, 63, 65, 67,	
147, 149, 150, 154, 181, 209.		95, 96, 98, 132, 188.		81, 82, 84, 86, 88, 89, 92, 94-96,	
McClymont.....	102, 191, 208	1174.....	93, 188, 206	106, 113, 115, 118, 119, 130,	
Meloy.....	107, 109, 111, 112, 115,	1272.....	53,	132, 181, 206.	
127-129, 135, 188, 206, 209, 211		64, 76, 83, 189, 206, 208,		Prince Edward.....	37, 193
Mensury:		210.		Princess.....	29,
C. I. No. 170.....	66, 82, 91, 177	1273.....	52, 57, 64, 189, 208	39, 44, 53, 56-58, 62, 63, 67, 75,	
2657.....	19,	1274.....	64, 189	80, 82, 86, 92, 95, 96, 106,	
22, 24, 27, 31, 33, 35, 37,		1517.....	101, 189	113, 117, 122, 124, 128, 130,	
39, 40, 42-45, 47, 49, 51,		1526.....	54, 189	134, 138, 142, 147, 149, 181,	
64, 69, 104, 138, 193, 206.		1529.....	58,	206.	
2924.....	69, 202	64, 101, 189, 206, 208		Prize Prolific.....	22, 27,
Michigan.....	55, 58, 60, 196, 208	2184.....	107, 191	33, 37, 47, 66, 80, 93, 142, 177	
Michigan Winter.....	56,	2191.....	118, 191	Probstier.....	55, 62, 63, 177
121, 191, 208, 209		2700.....	19,	Prolific.....	28, 195
Minsturd.....	70,	22, 28, 34, 38, 42-44, 47,		Proskowetz.....	53,
73-75, 117, 189, 206, 208, 210		33, 62, 63, 65, 66, 74-76,		69, 86, 88, 134, 145, 185	
Mishima.....	91, 176	82, 86, 194, 208.		Purposco.....	93, 177
Mocki-Mugi.....	91, 176	2780.....	58, 196	Quitepec.....	93, 181
Moister.....	117, 197	3146.....	121, 203	Rasput.....	122, 187
Monck.....	23, 28, 34, 38, 47, 194	Odessa:		Red River.....	53, 77, 187
Monde Hull-less.....	29, 194	C. I. No. 182.....	19,	Reed Triumph.....	30,
Monro.....	29, 194	22, 24, 25, 27, 31, 33, 35,		58, 65, 69, 83, 88, 145, 185	
Montana.....	59, 60, 193, 208	37, 39, 40, 42-44, 47, 49,		Rennie Improved.....	22,
Moravian.....	93, 124, 180	64, 66, 71, 74, 80-82, 86,		27, 33, 37, 47, 106, 177	
Moulton.....	28, 194	92, 94-101, 104-107, 109,		Rigid.....	23, 29, 34, 38, 47, 194
Nakano Wase:		110, 113, 114, 142, 149,		Robertson.....	82, 181
C. I. No. 754.....	118,	177, 206, 208-210.		Rokaku Chevalier.....	96, 176
143, 147, 154, 184, 206		916.....	53,	Ross.....	67, 198
2164.....	143, 191	77, 134, 135, 145, 186		Rottingham.....	55, 203
2165.....	143, 191	927.....	53,	Royal.....	22, 28, 34, 38, 47, 64, 66, 189
2166.....	143, 144, 191	77, 115, 116, 122, 123,		Russian.....	80, 88, 197
Nepal:		128, 130, 186.		Russian Naked.....	101, 197
C. I. No. 22.....	91, 134, 176	934.....	77, 134, 186	Saale:	
230.....	122, 124, 178	2154.....	93, 96, 191, 208	C. I. No. 25.....	91, 96, 176
234.....	134, 178	2158.....	93, 191	2645.....	22, 27, 33, 37, 47, 193
262.....	99, 179	2168.....	96, 191	Sagina.....	135, 189
352.....	93, 180	2187.....	118, 191	Salzer.....	80, 202
633.....	63, 181	2737.....	27, 195	Salzer Beardless.....	29, 62, 63, 183
595.....	23,	Ohio Beardless.....	56, 178	Samofa.....	70, 73-75, 117, 188, 208
24, 29, 31, 35, 38, 39, 45,		Ohio Winter.....	56, 191, 206, 208	Sandrel.....	53,
48, 49, 51, 53-55, 58-60,		Old Irish.....	30, 195	77, 111, 115, 116, 122-124, 128,	
62, 63, 65, 69, 71, 74-77,		Olonets.....	92, 177, 206	130, 134, 186, 206, 208-210.	
80-84, 86-89, 92, 94, 99,		Omar.....	107, 143, 185	Sangatsuka.....	91, 142, 176
100, 102-105, 107, 111-		Oregon.....	29, 195	Santaku.....	91, 176
113, 115, 117, 118, 121,		Orel:		Saul.....	54, 190
126, 128, 130, 132, 133,		C. I. No. 351.....	56,	Saxonian.....	55, 62, 63, 197
139, 145, 182, 206, 208.		93, 118, 119, 125, 128, 134,		Scholey.....	53, 86, 187, 208
630.....	113, 114, 183	142, 144, 147-149, 180,		Schwarzenberg I.....	138, 180
2315.....	134, 192	206, 209.		Schwarzenberg III.....	138, 180
Nepbant.....	113, 196	2190.....	118, 191	Scotch:	
Nepean.....	23,	Osaka.....	140, 143, 197	C. I. No. 848.....	55,
28, 34, 38, 48, 92, 106, 177		Pacer.....	23, 28, 34, 38, 48, 193	62, 63, 66, 103, 154, 184,	
Nerchinsk.....	135, 178	Palampur.....	28, 33, 37, 194	2922.....	69, 202
Nesbian.....	128, 132,	Palestine.....	134, 186	Scotch Chevalier.....	113, 196
134, 140, 142, 147-149, 183		Parkin.....	29, 194	Scottish Pearl.....	107,
New Hull-less.....	130, 203	Peacock.....	27, 33, 193	143, 144, 179, 209	
New Moravian.....	82, 197	Peerless.....	37, 193	Selected Chevalier.....	22,
Newton.....	22, 28, 34, 37, 47, 92, 177	Peerless White.....	22,	27, 33, 37, 193	
New Zealand.....	27, 33,	27, 33, 37, 47, 193		Selection:	
37, 53, 66, 80, 113, 114, 193		Pelham.....	29, 194	C. I. No. 2639.....	54, 193
Nijo.....	91, 176	Perm:		2640.....	54, 193
Nina.....	125	C. I. No. 17.....	92, 176	Semibarless.....	57, 197
Nita.....	91, 176	20.....	92, 106, 176	Senbaku.....	92, 176
Niver.....	143, 184	Peru:		Senca Purple.....	80, 202
Nonpareil.....	55, 203	C. I. No. 653. 113, 114, 134, 183		Seoraj.....	28, 33, 37, 194
Norway 8.....	178	2302.....	128, 192, 209	Servian.....	53,
Norway House.....	28, 194	2311.....	134, 192	70, 77, 122, 128, 134, 143, 145,	
Norwegian.....	69, 195	Peruvian.....	53, 77,	186.	
Nugent.....	19, 22	120-122, 128-130, 186, 206, 209		Sextradit.....	58, 64, 69, 113, 130, 182
28, 34, 38, 42-44, 48, 92, 177		Petro.....	122, 187	Shansi.....	113, 196
O. A. C. 21.....	19,	Petschora.....	22,	Sharpe Chevalier.....	22,
23, 25, 30, 35, 39, 40-44, 48,		24, 27, 33, 37, 47, 49, 66, 193		27, 33, 37, 47, 66, 193	
52, 54, 58, 65, 74-76, 101, 104,		Phoenix.....	22, 28, 34, 38, 47, 193	Shiro-Nishiki.....	91, 176
105, 113, 117, 128, 132, 189,		Phoenix Von Thalen 27, 33, 93, 180		Shiale Chevalier.....	93, 180
206, 208, 210.		Phoson.....	128, 181	Shalkot.....	28, 37, 194
O. A. C. Selection 7. 154, 197, 209		Pidor.....	143, 144, 185, 209	Sibley Imperial.....	51,
O. A. C. Selection 8.....	154, 197, 209	Pilsen.....	67, 177	62, 63, 66, 80, 197	
O. A. C. Winter.....	132, 204	Pioneer.....	23, 28, 34, 38, 48, 193	Sidney.....	22,
Oderbrucker:		Plumage.....	29, 176	28, 34, 38, 42-45, 47, 64, 91, 177	
C. I. No. 537.....	51,	Pod.....	113, 134, 135, 183, 206, 209	Silver Beardless:	
77, 128, 132, 138, 139, 181		Polar.....	27, 33, 37, 193	C. I. No. 641.....	58, 183
836.....	53,	Pontius.....	143, 149, 184	2887.....	68, 200
56, 59, 60, 81, 83, 84, 88,		Popeline.....	134, 184	Silver King:	
184, 206, 208, 210.		Poppenheim.....	93, 97, 179	C. I. No. 890.....	23,
888.....	86, 88, 104, 185, 208	Portuguese.....	139, 193	29, 35, 53, 58, 60, 62,	
940.....	77, 134, 135, 186	Prague.....	138, 179	63, 69, 80, 84, 88, 104,	
957.....	53, 77, 187	Prentice.....	27, 33, 37, 48, 186	117, 185.	

	Page		Page		Page
Silver King—Continued.		Tennessee	145, 204	Virginia Selection—Con.	
C. I. No. 905	53,	Tennessee Winter	56,	C. I. No. 3159	145, 204
64, 100, 122, 132, 185, 208		62, 63, 69, 101, 107, 118, 121,		3163	145, 204, 209
1276	64, 113, 189	124, 125, 128, 132, 134-136,		3164	145, 204, 209
2754	64, 195	138, 140, 142, 144-150, 153,		Voronezh	93, 180
Simla	28, 37, 194	154, 179, 206, 209, 211.		Wakamatsu	56, 142, 181
Sisolsk	29, 62, 63, 67, 80, 101, 177	Terwin	143, 181	Warren	29, 194
Six-Row	154, 197	Tetcherit	107, 177	Webbs	138, 179
Small Blue Naked	30, 195	Texas Winter	56, 128,	White Gatami	86, 93, 97, 186
Smyrna	118, 119, 193, 209	129, 142, 146, 149, 181, 208		White Moravian:	
South African	67, 197	Thanet	22, 192	C. I. No. 977	103, 122, 187
Spears	113, 196	27, 33, 37, 47, 55, 66, 193		1877	121, 190
Spiti Valley	27, 33, 37, 47, 193	Thomas	103, 197	White Smyrna	
Squarehead Winter	140, 142, 178	Thorpe	111, 186	C. I. No. 195	63, 70,
Standwell	19, 23, 29, 35,	Tioga	88, 195	82, 83, 85, 91, 94-101,	
39, 42-44, 48, 64, 67, 181, 206		Trebi	53,	103, 104-111, 113, 115,	
Stavropol	102,	54, 77, 100, 102, 103, 111, 113,		118, 121, 122, 124, 125,	
107-109, 191, 206, 208, 209		113, 116, 120-124, 127-130, 132,		134-136, 138, 139, 143,	
Steigum:		133, 135, 186, 206, 208-210		153, 177, 206, 208-210.	
C. I. No. 907	53,	Trifurcate Hull-less	93, 180	655	149, 183
70, 77, 86, 122, 185, 208		Trooper	19, 22,	658	86-88,
931	134, 186	28, 34, 38, 42-44, 47, 66, 92, 177		99, 102, 104, 107, 113-116,	
Stella	19, 22, 24, 25, 28, 31, 34,	Turkestan:		118, 128, 129, 129, 132,	
38-40, 42-44, 47, 193, 206		C. I. No. 711	124, 125, 128,	133, 183, 206, 208-210.	
Stoeckinger	58, 196	134, 147-149, 184, 206, 209		910	53, 54,
Strigum	82, 91, 95, 96, 176	712	142, 184	111, 112, 120-123,	
Success:		2715	29, 80, 194	130, 134, 185, 209, 210,	
C. I. No. 212	106, 178	Two-Row Brewing	51, 204	2080	118, 191
891	80, 88, 185	Two-Row Hull-less	145, 204	2169	86, 97, 128, 191, 208
2183	104, 191	Union:		2170	86, 97, 191, 208
2707	23-25, 28, 31, 34,	C. I. No. 2198	122,	2171	97, 191, 208
38-41, 44, 45, 48, 49, 194		124, 130, 191		2172	97, 191
2807	65, 197	2199	122, 124, 130, 191	Wing Beardless	121, 191
Success Beardless	52, 58, 190	3122	130, 203	Wing Pedigree	54, 55, 58, 65, 81,
Suffolk Chevalier:		3123	130, 203	86-88, 93, 97, 101, 117, 183, 208	
C. I. No. 2701	28, 194	3124	130, 203	Winter:	
2702	28, 194	3125	130, 203	C. I. No. 3115	55, 203
Summit:		3126	130, 203	3160	145, 204
C. I. No. 174	22, 28, 34, 38,	3127	130, 203	Winter Beardless	145, 204
44, 47, 66, 82, 83, 92, 177		3128	130, 203	Winter Club:	
929	53, 77, 134, 186	3129	130, 203	C. I. No. 488	120,
Surprise	22, 28,	3130	130, 203	121, 124, 146, 181, 209	
34, 38, 48, 66, 82, 93, 177		3131	130, 203	592	124-128,
Svanhals	25,	3132	130, 203	132, 133, 182, 206, 209, 211	
29, 31, 40, 44, 53, 54, 57, 58,		3133	130, 203	Winter Six-Rowed	140, 180
62, 63, 65, 67, 71, 74, 75, 82,		3134	130, 203	Wisconsin Pedigree	52, 53, 55, 56,
83, 85, 86-89, 92, 94, 101, 103,		3179	130, 204	58-60, 75, 81, 102, 184, 208, 210	
104, 106, 107, 111-116, 118,		Union Winter	143,	Wisconsin Pedigree 20	64, 196
121, 122-124, 128-130, 132-		145, 150, 151, 181, 209, 211		Wisconsin Pedigree 17	65, 196
134, 138, 139, 142, 144, 145,		Uyetake	91, 176	Wisconsin Pedigree 18	65, 196
149, 152, 177, 206, 208-210.		Yalkow	143, 185	Wisconsin Pedigree 35	65, 196
Svansota	70, 73-75,	Vanguard	23, 28, 34, 38, 47, 193	Wisconsin Winter:	
117, 190, 206, 208, 210		Victor	23,	C. I. No. 519	104,
Swedish	27, 33, 37, 123	28, 34, 38, 47, 66, 92, 177		107, 121-123, 126-128, 134,	
Swedish Star	30, 104, 193	Victoria	93, 179	144, 146-149, 154, 181, 206,	
Szechwan	143, 183	Virginia Hooded	125, 183, 134,	208, 209, 211	
Toganrog	30, 40, 195	140, 142, 146, 147, 149, 183.		1894	126, 190, 209
Tanbush	147, 148, 181	Virginia Selection:		2159	142, 191, 209
Tanikaze	91, 176	C. I. No. 3148	145, 203	2167	143, 191, 209
Tapp Winter	50, 190	3149	145, 203	Withycombe	128, 192
Tartar	88, 89, 187	3150	145, 204	Woodrow	122, 187
Tarumizu	92, 176	3151	145, 204	Wyoming Hull-less	113, 183
Tatien	113, 183	3152	145, 204	Yale	19, 23, 29, 34, 35,
Telli	82, 91, 107, 117, 134, 136, 138,	3153	145, 204, 209	38, 42-44, 48, 190, 206.	
142, 149, 152, 154, 177, 206		3154	145, 204	Yendide:	
Tenarab	140, 150, 183	3155	145, 204	C. I. No. 205	93, 106, 178
Tenkow	142, 183	3156	145, 204	2189	118, 191
		3157	145, 204	Yerli	115, 183

Varieties of Interest or Promise

	Page		Page		Page
Alpha	163	Gatami	174	Oderbrucker	166-167
Arlington Awnless	172	Gold	162-163	Odessa	168
Beldi Dwarf	170	Hanna	162	Peruvian	170
Beldi Giant	170	Hannchen	163	Princess	162
Black Hull-less	175	Himalaya	175	Sandel	169
Blue Long Head	170	Horn	161	Silver King	165
California Marliout	170-171	Horsford	173	Standwell	160
Canadian Thorpe	159	Hybrid	167	Stavropol	168
Charlottetown 80	161-162	Invincible	160	Stella	167
Chevalier II	161	Lion	174	Svanhals	160
Club Marliout	171	Manchuria	164-165	Tennessee Winter	172
Coast	169	Manchurian	166	Trebi	168-169
Coast (Calif., 4000)	169	Meloy	173	White Smyrna	163-164
Early Chevalier	160	Mensury	166	Winter Club	172
Featherston	165	Minstard	167-168	Wisconsin Pedigree	166
Flynn	171-172	Nepal	174-175	Yale	167
French Chevalier	160-161	O. A. C. 21	165		

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE

September 30, 1925

<i>Secretary of Agriculture</i>	W. M. JARDINE.
<i>Assistant Secretary</i>	R. W. DUNLAP.
<i>Director of Scientific Work</i>	
<i>Director of Regulatory Work</i>	WALTER G. CAMPBELL.
<i>Director of Extension Work</i>	C. W. WARBURTON.
<i>Director of Information</i>	NELSON ANTRIM CRAWFORD.
<i>Director of Personnel and Business Administration</i>	W. W. STOCKBERGER.
<i>Solicitor</i>	R. W. WILLIAMS.
<i>Weather Bureau</i>	CHARLES F. MARVIN, <i>Chief</i> .
<i>Bureau of Agricultural Economics</i>	THOMAS P. COOPER, <i>Chief</i> .
<i>Bureau of Animal Industry</i>	JOHN R. MOHLER, <i>Chief</i> .
<i>Bureau of Plant Industry</i>	WILLIAM A. TAYLOR, <i>Chief</i> .
<i>Forest Service</i>	W. B. GREELEY, <i>Chief</i> .
<i>Bureau of Chemistry</i>	C. A. BROWNE, <i>Chief</i> .
<i>Bureau of Soils</i>	MILTON WHITNEY, <i>Chief</i> .
<i>Bureau of Entomology</i>	L. O. HOWARD, <i>Chief</i> .
<i>Bureau of Biological Survey</i>	E. W. NELSON, <i>Chief</i> .
<i>Bureau of Public Roads</i>	THOMAS H. MACDONALD, <i>Chief</i> .
<i>Bureau of Home Economics</i>	LOUISE STANLEY, <i>Chief</i> .
<i>Bureau of Dairying</i>	C. W. LARSON, <i>Chief</i> .
<i>Fixed Nitrogen Research Laboratory</i>	F. G. COTTRELL, <i>Director</i> .
<i>Office of Experiment Stations</i>	E. W. ALLEN, <i>Chief</i> .
<i>Office of Cooperative Extension Work</i>	C. B. SMITH, <i>Chief</i> .
<i>Library</i>	CLARIBEL R. BARNETT, <i>Librarian</i> .
<i>Federal Horticultural Board</i>	C. L. MARLATT, <i>Chairman</i> .
<i>Insecticide and Fungicide Board</i>	J. K. HAYWOOD, <i>Chairman</i> .
<i>Packers and Stockyards Administration</i>	JOHN T. CAINE, <i>in Charge</i> .
<i>Grain Futures Administration</i>	J. W. T. DUVEL, <i>In Charge</i> .

This bulletin is a contribution from

<i>Bureau of Plant Industry</i>	WILLIAM A. TAYLOR, <i>Chief</i> .
<i>Office of Cereal Investigations</i>	CARLETON R. BALL, <i>Senior Agronomist in Charge</i> .

219

ADDITIONAL COPIES
OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.

AT
30 CENTS PER COPY