Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

UNITED STATES DEPARTMENT OF AGRICULTURE

DEPARTMENT BULLETIN No. 1210

Washington, D. C.

May 1, 1924

SUMMARY OF LITERATURE ON BUNT, OR STINKING SMUT, OF WHEAT

By

HORACE M. WOOLMAN, Field Assistant in Cereal Disease Investigations, and HARRY B. HUMPHREY,
Pathologist in Charge of Cereal Disease Investigations, Office of Cereal Investigations,
Bureau of Plant Industry

CONTENTS

	Page
Introduction	1
General historical retrospect	2
The pre-Tillet period	2
The Tillet-to-Kühn period	4
The Kühn or modern period	7
Physiologic factors affecting the development of bunt	9
History	9
Physiologic response of spores of <i>Tilletia</i> to temperature	13
Control of bunt	13
History of seed treatment for the prevention of bunt	13
Varietal resistance to bunt	18
Summary	20
Bibliography	22

WASHINGTON
GOVERNMENT PRINTING OFFICE
1924

UNITED STATES DEPARTMENT OF AGRICULTURE

DEPARTMENT BULLETIN No. 1210

Washington, D. C.

May 1, 1924

SUMMARY OF LITERATURE ON BUNT, OR STINKING SMUT, OF WHEAT.¹

By HORACE M. WOOLMAN, *Field Assistant in Cereal Disease Investigations*,
and HARRY B. HUMPHREY, *Pathologist in Charge of Cereal Disease Investi-*
*gations, Office of Cereal Investigations, Bureau of Plant Industry.*²

CONTENTS.

	Page.
Introduction.....	1
General historical retrospect.....	2
The pre-Tillet period.....	2
The Tillet-to-Kühn period.....	4
The Kühn or modern period.....	7
Physiologic factors affecting the development of bunt.....	9
History.....	9
Physiologic response of spores of <i>Tilletia</i> to temperature.....	13
Control of bunt.....	13
History of seed treatment for the prevention of bunt.....	13
Varietal resistance to bunt.....	18
Summary.....	20
Bibliography.....	22

INTRODUCTION.

The world-wide importance of bunt, or stinking smut, as a factor influencing wheat production, its great antiquity as a cereal disease, and the vast amount of published matter relating to it, much of it not generally accessible, are facts seeming to justify the publication of this bulletin.

In the preparation of the subject matter it has been considered advantageous to present the data in the following historical periods:

- (1) The pre-Tillet period (500 B. C. to 1755 A. D.).
- (2) The Tillet-to-Kühn period (1755-1870).
- (3) The Kühn or modern period (1870 to date).

The first of these periods is characterized by the almost complete absence of information based on an experimental investigation of the problem and by the preponderance of inferences and conclusions derived from hearsay and speculation. The second, or Tillet-to-Kühn period, marks the beginning of bunt investigations by the experimental method. During this period the causal relation-

¹The manuscript for this bulletin was submitted for publication August 6, 1923.

²The investigations on the physiology and control of bunt, or stinking smut, of wheat, were conducted by the Office of Cereal Investigations, Bureau of Plant Industry, United States Department of Agriculture, in cooperation with the Washington Agricultural Experiment Station.

ship between the bunt fungus and the disease was established, and the essential facts pertaining to the life history of the organism were definitely ascertained. The Kühn or modern period is characterized by a searching and detailed consideration of the more strictly pathologic aspects of the problem and an experimental study of various methods of controlling bunt. More recently, attention has been given to the study of varietal resistance to bunt and to the relation of certain growth factors to infection and the subsequent development of the disease within the host.

GENERAL HISTORICAL RETROSPECT.

THE PRE-TILLET PERIOD.

There is no doubt that bunt, or stinking smut, of wheat, due either to *Tilletia tritici* (Bjerk.) Wint.³ or *T. laevis* Kühn, has been known to wheat growers for many centuries, probably ever since wheat became a cultivated crop. Without question it was known to Theophrastus, as well as to Vergil, Pliny, and other ancient Romans. It seems, however, that the same term was made to apply to both rust and smut by these early writers; in fact, these diseases certainly were thus confused by some writers as late as the middle of the eighteenth century. In the early English writings on the subject, mildew was used as synonymous with smut. In like manner it is found that "Mehltau" and "Brand" were used interchangeably by early German writers.

R. Bradley (56),⁴ professor of botany in the University of Cambridge, London, 1725, says:

We find here a fresh Scene of Mischief, from the Mildew in the first Places, which is here call'd *Rubigo*, and by some *Rubigo*, is the Smut in Corn, which often spoils whole Fields, or Crops of Grain; for where this happens in any Quantity to distinguish itself, it spoils the whole Parcel of Corn for Sale, and renders it unwholesome to be used in Flower or Meal; or to be sown, we must certainly expect our Crop to produce a great deal of it. *Scaliger*, on *Theophrastus*, tells us, that Smut, or Mildew, is a certain Putrefaction, when the Dew, or Rain, that is lodg'd in the Ears of Corn, is not shaken off, but by the adventitious Heat of the Sun corrupts and putrefies. It is supposed, that this Disease happens most in enclosed Grounds, where the Air is more close than in Hills or Champain. The Remedy of this, says *Pliny*, is by sticking Laurel Branches in the Ground (that is, what we call the Bay) which will draw the blasting Vapour to them; and for this End were the Feasts, call'd *Rubigalia*, instituted by *Numa*, in the Eleventh Year of his Reign. But I differ in Opinion from *Scaliger*, that this Smut in Corn is the Effect of Putrefaction: My Opinion is, that all Blights proceed from Insects, for besides the Observations I have made with Microscopes, upon the Black in Corn, which plainly discovers it to be the Eggs of Insects; there is a certain Remedy found against it, by Colonel *Plummer*, late of *Hertfordshire*, which consists in steeping of Corn before it is sown, by which these pestiferous Eggs are destroy'd; so that the Corn steep'd, as he directs, will not produce any smutty Corn. The Receipt for making this Brine, for steeping of Corn, is in my Monthly Writings. [The writers were unable to obtain access to Mr. Bradley's Monthly Writings to which reference is here made.]

³ *Tilletia tritici* (Bjerk.) Wint. Syn. *Lycoperdon tritici* (in Svensk. Vetensk. Akad. Handl. 1775, p. 326, Abhdlg. d. Schwed. Akademie, Deutsch von Kastner).

Uredo caries DC., 1815, *Flora franc.*, 5(6), p. 78.

Caeoma segetum Nees, 1816, *Syst. Pilze*, p. 14, pl. I, fig. 7.

Uredo sitophila Ditmar, 1813, in *Sturm Deutschl. Flora*, Abt. III, Bd. 1, p. 69, pl. 34.

Caeoma sitophilum Link, 1833, in *Linne, Spec. Plant*, ed. 6, v. 2, p. 2.

Uredo foetida Bauer, 1821, in *Ann. Sci. Nat. Ser. 1*, Bd. 2, p. 167.

Erysibe foetida Wallr., 1833, *Flora Crypt Germ.*, v. 2, p. 213.

Tilletia caries Tul., 1847, *Mem. Ustil.*, p. 113.

⁴ The serial numbers (italic) in parentheses refer to the "Bibliography" at the end of this bulletin.

Camerarius (73) quotes one Merodanus as follows:

In the year 1550 the Gauls first raised wheat which they called *roratum* or *mellitum* (honeydew) referring to the odor when the head is green: but when mature there was nothing but a black, foetid dust.

Hieronymus Bock (47, p. 665-666), who lived between 1498 and 1554, described bunt, but spoke of it as something already well known. Camerarius shows that the question had been raised in his day whether the causes of bunt were internal or external. He makes it clear that he regards bunt as being essentially different from the loose smuts of wheats, oats, and barley. It had occurred to him that the smuts might be fungi because they produced a black powder like the *Lycoperdons*. He finally abandoned this theory because he was unable to find with the microscope any such fungous threads as had been described for *Mucor* by Malpighi.

According to Richard Remnant (321), bunt had become a source of serious loss in England prior to 1637. He mentions having experimented with certain steepes as bunt preventives, but withholds any information as to their ingredients until satisfied as to their value. This is the earliest record of combating bunt by seed treatment that the writers have discovered.

The subject of smuts, in general, occupies considerable space in the agricultural and so-called scientific journals of the eighteenth century, but little of this was available to the writers, since it could not be found in the principal libraries of the United States. Enough of this early literature has been examined, however, to show the trend of thought for that period as to causes and means of prevention. It can be said that the ideas on the subject, particularly in the first half of the century, were largely empirical and charged with an admixture of superstition.

Several smuts were found listed by Dr. Caspar Bauhin (36), in *Phytopinax*, among which was *Ustilago secalina* (*Tilletia secalina*).

Three anonymous articles, probably by the same author, two of which are mentioned in Benevenuti's (42) notes, appeared in the *Sammlung von Nature und Medicin*: Breslau (1, 2, 3). A few abstracts from these articles follow:

A large amount of "*brand*" was found in the wheat in the year 1717; also some in the rye but not so much, rye being less subject to the trouble. The *brand* occurred mostly under certain conditions: (1) In low damp soil. (2) When the plants came up too slowly. (3) Where the soil had become contaminated. (4) When planted in double light, i. e., "Wenn früh oder Abends Sonn und Mond zugleich am Himmel zu sehen." (5) When new seed was planted damp. (6) In case of injurious dew.

He relates that honeydew and damp July weather had a bad influence. He quotes an unnamed writer to the effect that the sowing of damp new wheat, i. e., new wheat not fully cured, favors the production of *brand* and that it is safest to sow seed from the previous year, but that this should be held over, unthreshed, until just before sowing. The same writer informs us that many means of prevention were practiced, mostly superstitions, and cites a few, such as mixing powdered lime with the seed, sowing in the dark of the moon "in God's name," and soaking the seed in salt water for 24 hours. To prevent smut in barley, running the seed through burning straw (flaming) before sowing was sometimes practiced. He notes that smut can be produced experimentally in the window of a room as well as in the open field. He had observed

plants that produced both smutted and sound heads and that the smut heads were diseased before they emerged from the boot. He suggests that interference with sap flow may be a factor.

According to Jethro Tull (378), the use of salt brine for treating seed originated about 1650. A shipload of wheat had been beached near Bristol at high tide. The wheat was salvaged and, being unfit for grinding, but viable, was used for seed. This resulted in a clean crop, while fields in which home-grown seed was sown were bunted. He states, however, that this was only partially remedial, proving quite ineffectual in certain years. He seems to have had some idea that the disease was infective, as he recommends changing seed and suggests that results obtained by sowing the salvaged wheat might have been due to the fact that the wheat came from a locality where there was no smut.

Abbé Pluche (309) applies the term "smut" to what is unquestionably rust. This disease he attributes to the burning effect of the sun's rays which are concentrated by the small drops of dew or rain. Bunt, which he describes as a "black meal," results, he believes, from a failure of fertilization of the ovaries of the affected heads.

C. F. Meyer (269) believes "bunt dust" to be infective and thinks it acts by producing a fermentation of the sap. He advises treating with lime.

Aucante (27) maintains that treatment of the seed with lime and arsenic, as well as with corrosive sublimate, is useless and that it is injurious to germination. He believes smut to be due to atmospheric conditions. It is uncertain whether Aucante refers to loose smut or bunt, or whether he regards them as one. The article is interesting chiefly because it contains the first mention of either arsenic or corrosive sublimate as a treating medium.

J. H. Ritter (330), in response to Aucante's article, holds that the bunt dust is infective, but that bunt production is influenced by growth conditions under which the seed grain matures. He advises (1) sowing well-ripened and well-dried seed; (2) treating with a mixture of salt, saltpeter, lime, and wood ashes; and (3) in the long months sowing in the forenoon, in the short months in the afternoon.

THE TILLET-TO-KÜHN PERIOD.

In 1755, Abbé Tillet (370) proved by experiment the infective character of the bunt dust, thus establishing the first real step toward the discovery of the true cause of bunt. Tillet sowed seed artificially blackened with bunt spores alongside of rows sown to clean seed and obtained very much bunt in the former and perfectly sound wheat in the latter. He repeated his experiments at Versailles in 1759 and obtained the same results. The writers have been unable to find a copy of Tillet's original paper in the United States; consequently, they have been compelled to rely on such commentaries as are available. Comments pro and con regarding his conclusions furnished a large part of the literature on the subject from 1760 to 1800. Tillet's work and conclusions were imperfect in two respects: (1) He did not recognize *Tilletia* as a parasite or even as a fungus, notwithstanding the fact that he discovered the contents of bunt balls to be of uniform size and shape. (2) He regarded bunt and loose smut as differing only in degree, or merely as different stages of develop-

ment. However, he rejected the theory that rusts were related to the smuts.

Schreber (339) undertook to refute Tillet's conclusions in toto. He denied that bunt was infectious, basing his conclusion on the fact that a clean crop could sometimes be produced from smutty seed. He argued that bunt was due to a disordered sap produced by atmospheric conditions. He agreed with Tillet that rust was in no way related to the smuts. Even at this date there were persons who held the opposite view.

In 1763 Benevenuti (42) asserted that smut or bunt was an advanced stage of rust, or at least resulted from it, and that rust was caused by concentration of the sun's rays on drops of dew or rain water.

In marked contrast to Schreber's article is one by J. L. Sturler (359), who strongly supported Tillet. After presenting evidence of the infective nature of the disease he proceeds to a somewhat vigorous expression of his opinion of such farmers as refuse to treat their seed. Referring to the evidence he had presented, he says:

In spite of this clear demonstration few farmers make use of this easy and cheap means of help. Deplorable blindness and unaccountable laziness! Were they ignorant and prejudicial peasants my surprise would be less. But shall not one grieve that people of another sort, some of whom belong to the first rank, should persist in their old humdrum ways and stiffneckness? I know such without having made a single experiment maintain that smut is an accidental disease of the grain which man can not anticipate or prevent and that the powder can not infect sound grain. Can an error be more plain? However, we will hope that time and the light which the society [Economic Society of Bern] is distributing will help even such impaired understandings!

Schulthess (342) makes what is probably the first mention of the use of copper sulphate as a treatment medium.

Tschanner (371) supports Tillet, but thinks bunt can be avoided by sowing old seed. Tschanner states that red spelt is less liable to bunt than white spelt, a fact observed by him in fields containing both varieties. So far as the writers can determine, this is the first mention of varietal difference in resistance to bunt.

In 1774 Johann Christian Fabricius (105) published a paper on phytopathology, in which he took the ground that fungi found in plant lesions were distinct organisms and not morbid plant tissues. How much influence this had on the final discovery of parasitism in its relation to plant diseases in general and smuts in particular is hard to say.

In 1783 and 1786 Tessier (368, 369) repeated Tillet's experiments, confirmed his conclusions, and performed some further experiments of his own. He says:

According to the experiments of Mr. Tillet and myself, the stinking smut spreads itself by contact with great facility and multiplies rapidly.

From his own experiments he came to the conclusion that the original cause is not to be found in the proximity of bushes, kind of fertilizer used, or the nature of the soil, but thinks the disease must originate from some inherent peculiarity of individual plants. He observed that the presence of smut "powder" on the grain retarded germination. He gives a very accurate morphological description of the affected heads at different stages of their development, from just before their emergence from the boot until maturity, and of the microscopic appearance of the spores, from which one would conclude

that he was dealing with *Tilletia laevis*. He mentions several methods of prevention by means of seed treatment.

Volkart (384) quotes Prévost to the effect that sulphate of copper was also recommended by Tessier.

Bryant (69) states that the treatment of seed wheat with salt brine, soap lees, and lime had been practiced in England for generations prior to 1784. He believes it to be useless for prevention of bunt and injurious to the seed. He also states that the farmers knew it to be ineffectual, but that they persisted in it because it was a time-honored habit. He made a careful examination of the bunted heads and gave a correct morphological description of them, including the aborted stamens. He was convinced that atmospheric or other external factors caused the shrinking or hardening of the ligule at the base of the flag leaf, thus causing the young head to be compressed in emerging from the boot. This supposed interference with the flow of the sap he believed resulted in the abortion of the stamens and consequent failure of fertilization, in which case bunt was produced. Where the compression was more severe, both pistils and stamens were aborted, and loose smut resulted. This was an ingenious theory. Bryant clearly failed to discover the presence of bunt in the young head before it was subjected to this pressure. The ideas here expressed were common, in a more or less modified form, during the eighteenth century and, together with the idea of injurious or poisonous dew and disordered sap, dominated the pseudoscientific thought, at least down to the time of Tillet's discoveries. Meyen (268) was absolutely certain that smuts of cereals were not infectious diseases but were due to stagnation of the sap, caused by overfertilization or improper fertilization of the soil.

The work of Prévost (312) was a distinct contribution to our knowledge of the smuts. He observed the germination of the spores of *Tilletia* and the production of the primary and secondary sporidia. He also tried to demonstrate the entrance of the fungus into the seedling.

Tulasne (377) carefully followed up and confirmed Prévost's work and established the genus, naming it *Tilletia* in honor of Tillet. In his second memoir, 1854, he figures the germination and development of *Tilletia* as far as the formation of the secondary sporidia. There was now no logical ground to doubt the parasitism of this smut, although no entrance to the host plant had been demonstrated.

Berkeley (43) says that it has been known for many years that the principal diseases of cereal plants, such as rusts, bunt, mildew, etc., are of vegetable origin. Further, he states that, in spite of Unger's (380) attempt to overthrow this theory, the work of Cordu, Léveillé (239), and others had completely established the fact that the peculiar structures found in plant lesions were not modifications of the host tissue, but that they sprang from a distinct mycelium and were as certainly vegetable as other fungi. In the same article, Berkeley reports bunt studies carried on by him during the year 1846 for the purpose of demonstrating the parasitism of the organism as well as the method of infection and the physiologic relationship of the fungus to the host. He observed the germination of the spores and the production of sporidia, though he did not recognize the latter as belonging to *Tilletia*, but thought it a *Fusarium* attacking the bunt. Farther on he states that in one case he observed on a bunted plant a streak

of bunt upon the stem "in which the fetid smell and peculiar structure were not to be mistaken." Evidently he was ignorant of Prévost's work, for he says (44) that he believes himself to have been the first to observe the fusiform sporidia, although he misinterpreted them. In 1857 (45) he correctly figures the germination and development as far as the formation of secondary sporidia.

The controversy between those who recognized fungi as causes of plant diseases and those who regarded them merely as symptoms or effects of plant disorders raged from 1800 to 1860, and the briefest review of the writings of this period would extend this bulletin beyond reasonable limits. Of those who, by their investigations and writings, have contributed to the establishment of the parasitism theory, the following deserve special notice: De Candolle (74, 75, p. 1435-1438), Banks (31), Prévost (312), Rudolphi (331), Tulasne (377), Lévillé (239), Berkeley (43, 44), and De Bary (34, 35). The leaders of the opposition were Unger (379, 380, 381), Meyen (268), and Fries (126). In so far as bunt and other covered smuts were concerned, the controversy ended with the work of Kühn (220, 223), and Wolff (398), who demonstrated the entrance of the mycelium at the root node or between it and the point of the leaf sheath. The work of Maddox (255, 256), Hecke (144), and Brefeld (61, 62) in proving the blossom infection of the loose smuts put them in a class apart from the bunt and covered smuts and facilitated in great measure the further study of both classes.

THE KÜHN OR MODERN PERIOD.

Kühn (220) observed what he conceived to be the entrance of the fungus threads into the host plant through the leaf sheath and stated that it grew across, through the leaf layers, to the growing point.

The next successful experiments were those conducted by Wolff (398). He investigated several smuts, but principally the stem smut of rye, *Urocystis occulta* (Wallr.) Rabenh. His conclusions were that the rye-smut organism entered the host, not necessarily at the root node, but within 8 or 10 millimeters above it. Wolff observed that the mycelium, while passing through the epidermal cell, was inclosed in a cellulose sheath, and that after this passage its growth within the host became intercellular, pushing haustoria into the cells. With the smuts of wheat, oats, and barley, Wolff claimed that the mycelium was, in general, intracellular. In 1874 Kühn (223) announced that infection could take place at almost any point in the seedling.

After Kühn, Brefeld carried on an extensive series of inoculation experiments, using sporidia produced in pure culture (see Ward, 390). He maintained that the fungus could enter the plant at any point where it could be brought in contact with meristematic tissue. He asserts also that he found the mycelium in all parts of the young growing plant, from the root hairs and roots upward, and that he obtained infection by passing the inoculum downward to the growing point through the rolled leaves by means of a capillary tube. In this latter case he says that many points of entrance could be discovered when the material was killed soon after the inoculation, but that the openings soon closed and became invisible with the growth of the plant. Brefeld used oats and barley and their smuts in his inoculation experiments, so his results may not be entirely applicable to *Tilletia*.

Jensen (184) suggested the possibility of blossom infection in case of the loose smut of barley. His deductions, based on certain experiments and observations, were briefly as follows:

- (1) He was unable to produce infected plants by applying spores to the seed.
- (2) Seed taken from a field in which loose smut had appeared produced infected plants in spite of treatment with copper sulphate, while seed from a clean field produced a clean crop.
- (3) Seed treated with copper sulphate produced a smutty crop, while the same seed treated with hot water produced a clean crop. From these facts, Jensen reasoned that either an internal infection had obtained or that the spores were lodged between the grain and its adhering glumes.

Maddox (255) demonstrated the blossom infection of the loose smut of wheat. This was fully confirmed by Brefeld (61) in 1903. Hecke (144) in 1906, and Hori (170) in 1907.

After this date the history of the investigation of the bunt and loose smut of wheat becomes for the most part dissociated. Before taking leave of the subject of the loose smuts, a few words should be said about the contributions of Maddox to the discovery of blossom infection, inasmuch as he has never been accorded the complete recognition he deserves. His methods of investigation are best expressed in his own words, quoted in 1896 in the *Agricultural Gazette of Tasmania*:

I will now give the conclusions I have arrived at with smut [of wheat] from the results of my experiments. I have never been able to cause infection and reproduce the disease with spores on the grain or in the ground, which I can so easily do with bunt spores to reproduce bunt. The only way I have been able to infect grain and reproduce smut (which seldom ever fails) is by putting the spores on the ovary of the plant at flowering time, about the same time as the pollen grains are being shed. The grain will mature without the slightest signs of being diseased. I have hit the time so well now that I may say I never have a failure. I think this accounts for when I did fail, viz, the ovary was not forward enough for the spore to get its seed bed, or possibly sometimes the spores were not matured enough. The comparison of bunt and smut spores finding their seed bed are the very opposite. It is really wonderful to me how the smut spores do, as the ovary is well protected by the glumes or chaff, and there is no doubt that here we have a practical demonstration of the fact that infection by the loose smut of wheat occurs during the flowering period and that this is the first record of it.

McAlpine (252) in his "Smuts of Australia," 1910, accords due recognition to Maddox by quoting the foregoing paragraph. On the other hand, Kirk (211), in the Annual Reports of the New Zealand Department of Agriculture, gives the entire credit to Brefeld and Hecke, with no mention of Maddox.

Lang (231) published his conclusions from somewhat extensive investigations of the physiologic relationships of the smuts to their hosts and of their modes of infection. His studies were confined to *Ustilago tritici* (Pers.) Jens. and to the smuts of oats and barley, but it is of interest to note that he found them all to be intercellular parasites and that they obtained their nutriment through their high osmotic power. So far as the seedling infection in case of oats and barley is concerned, he is of the opinion that the entrance of the parasite is promoted by the degeneration of the cells of the leaf sheath, if not entirely dependent on it.

In the period following the final determination of the parasitism and the mode of infection of bunt and extending to the present time, much investigation has been carried on and a great volume of literature published. This work and publication can be grouped under

three general heads: (1) Physiologic and ecologic factors, (2) varietal resistance or immunity, and (3) seed treatment.

PHYSIOLOGIC FACTORS AFFECTING THE DEVELOPMENT OF BUNT.

HISTORY.

Throughout the pre-Tillet period, from Theophrastus down, the cause of bunt, as well as that of other smuts and the rusts, was sought almost entirely in physiologic and environmental factors, Bradley's supposed discovery that the bunt spores were insect eggs being the only notable exception. Theophrastus, and evidently Pliny, believed this smut to be due to moisture entering the young head, causing putrefaction under the action of the sun's heat. In all the literature of the early part of the eighteenth century we find the cause attributed to such things as dew, honeydew, fog, rains followed by hot sunshine, damp soil, interference with flow of sap, and failure of fertilization.

The experiments of Jethro Tull (378) furnished strong evidence that certain climatic and soil conditions were contributory factors. He cites the fact that a large amount of bunt appeared in certain years. He believed that excessive soil moisture was conducive to bunt production and finds strong support for this belief in the results of what seems to have been the earliest controlled physiologic experiment on record dealing with the bunt problem. He took up several seedling wheat plants from a field and transferred them to the window of his chamber, where they finished their growth in excessively wet soil, a part of their roots being at all times in water. These plants were entirely bunted, while bunt heads were very rare in the field from which he took up his experimental plants. Others observed that late sowings were more subject to bunt than crops sown earlier.

While Tillet's experiments proved bunt to be an infective disease, evidence has steadily accumulated to show that the occurrence of bunt in the crop is largely dependent on ecologic factors, not only at the time of sowing but to some extent during subsequent growth.

Ferle (116) says that dampness, protected locations, impermeable soil, and fresh stall manure favor bunt production. Ivanoff (178), working with *Tilletia laevis* in Tiflis, Transcaucasia, found an apparent relation between bunt prevalence and either altitude or moisture.

The first recorded observations on the relation of temperature to spore germination in *Tilletia* were those of Prévost (312) about 1807. He states that in water they germinated at 17° to 18° C. after 2½ to 3 days, at 14° to 15° after 3 to 4 days, at 11° to 12° after 5 to 6 days, at 7° to 8° after 9 to 10 days, and at 4° to 5° after 11 to 12 days. Hoffmann (164) and Schindler (336) investigated the thermolethal points of spores.

Tubeuf, in his comprehensive investigations of all phases of the bunt problem (372, 374), concludes that at a temperature of 3° to 4° C. the spores will not germinate, and that even at temperatures of 5° to 11° C. they may remain ungerminated in the soil and, consequently, a source of danger. He finds the optimum temperature for their germination to be between 16° and 18° C. He is in agreement with Appel that high temperatures promote bunt production.

Volkart (384) found that the optimum temperature for spore germination was between 16° and 18° C. Between 18° and 20° C. the germination began just as quickly, but was less vigorous. Above 25° C. there was no germination, but the spores remained uninjured. Volkart apparently did not undertake to fix the minimum. The figures in Table 1 are from a series of Volkart's experiments to determine the relative response of wheat and bunt spores to various temperatures.

TABLE 1.—*Data on germination of wheat and of bunt spores, at various temperatures according to Volkart.*

Temperature.	Wheat germination.	Spore germination.
5.0° to 6.1° C.....	4 per cent on the 8th day. 93 per cent on the 12th day.	A few on the 12th day. Abundant on the 15th day.
12.0° to 12.9° C.....	61 per cent on the 5th day. 97 per cent on the 6th day.	Moderate on the 5th day. Abundant on the 6th day.
17.8° to 21.4° C.....	3 per cent on the 2d day. 99 per cent on the 6th day.	Abundant on the 5th day.
28.3° to 33.6° C.....	9 per cent on the 2d day. 98 per cent on the 8th day.	None.
33.1° to 37.2° C.....	15 per cent on the 3d day. 63 per cent on the 12th day.	Do.

Volkart (384), contrary to the view held by Tubeuf (372, 374), concludes that the lower the temperature of the soil the greater the liability to infection.

Hecke (147) conducted careful experiments with spring sowings in 1907 and 1908, extending from March 3 to May 21, during which time he kept a continuous record of soil temperatures. He states his conclusions as follows: "In case of bunt, the view that the degree of infection is in inverse proportion to the temperature is without doubt correct." He also notes that this would indicate early fall sowing, which is generally known to be less infected than late. Hecke, however, does not believe that the temperature influence is limited to the germination period. He further states that there are three ways in which temperature may exert an influence: (1) It may affect the germination of the spores and the seed; (2) it may lengthen or shorten the duration of susceptibility to infection; and (3) it may limit or promote the ability of the fungus to reach the growing point of the host plant.

The field experiments of Munerati (283, 284, 285) furnished further support for the views of Volkart and Hecke. All his seedings showed a reduction of bunt with rising soil temperature, but in his case, however, the soil temperature can be inferred only, as he depends on the air-temperature records taken at a neighboring weather station.

Müller and Molz (275) also investigated this question (see table 2 for their results). Their temperature observations are open to the same source of error as those of Munerati, i. e., they are air temperatures from a weather station, the distance of which from the experimental plat is not stated.

TABLE 2.—Data on influence of temperature on bunt production, according to Müller and Molz.

Winter wheat.			Spring wheat.			
Date sown.	Bunt heads per 30 square meters.	Average temperature for 7 days following.	Date sown.	Bunt heads per square meter.		Average temperature for 7 days following.
				Red Schlanstedter.	Red Bordeaux.	
October 5.....	8,040	3.31	March 14.....	465	165	5.6
October 26.....	2,970	8.40	April 2.....	12	13	9.7
November 23.....	174	2.13	April 17.....	17	1	7.34

While the results of these spring sowings are in harmony with Hecke's conclusions, the fall sowings appear somewhat paradoxical. At that time they challenged explanation, and to some extent they still remain unexplained. Müller and Molz's conclusion that lower temperatures promote bunt production is true only with limitations.

All agree that temperature at and subsequent to sowing is an important factor, but are not in entire accord as to what is the optimum temperature. Hecke is positive that lower temperatures favor infection. He is supported by the field experiments of Muncrati, cited above.

All these investigations seem to have failed to establish the important fact that there is an optimum temperature for bunt infection and production above and below which the susceptibility or liability of wheat to bunt falls off. The authors of this paper believe that their investigations, while not entirely solving the question of temperature relations to bunt, have contributed to its solution.

Brefeld (63) demonstrated that on a suitable nutritive substratum *Tilletia* would grow saprophytically for an indefinite period with a great increase of secondary conidia and eventually would produce a spore closely resembling the chlamydospore. This had been fully confirmed by the writers, but his further conclusions, i. e., that the spores are favorably influenced and their germination promoted by passing through the digestive tract of animals, have not been confirmed by later investigation. Nor has it been shown that Brefeld was correct in assuming that infection was spread by wind dissemination of the conidia, or that the application of infested stall manure to the fields would establish a permanent and increasing saprophytic growth of the bunt organism.

Tubeuf (372, 374), after a comprehensive investigation of the bunt problem, states his conclusion in part as follows:

One must assume that a distribution of the bunt of wheat from plot to plot by means of conidia in general does not take place, that spores passing through the digestive tract of animals are destroyed or injuriously affected, and that fresh stall manure is toxic to spores.

In fact, Tubeuf believes that normal germination of spores does not take place in nutritive solutions. His experiments seem to indicate the possibility of a crop being infected through bunt-infested manure if applied fresh and very shortly before sowing, particularly if the spores come from infested straw used for litter, but that danger

from this source would disappear in a short time after applying the manure.

Steglich (356) agrees that danger from spores introduced with stall manure lasts but a few days, because, as he asserts, manure acts as a stimulant to germination and thus causes the spores quickly to become innocuous in the absence of a host plant, as continuous reproduction of conidia does not take place in the soil. A microscopic examination of spores evacuated by swine showed 75 per cent loss of viability. Spores placed in a small sack and buried in a manure pile for 193 days had practically all germinated and become innocuous. When these were mixed with water and sprayed on seed wheat they apparently caused the appearance of three bunt heads in 15 square meters of field area.

Appel and Riehlm (20) are confident that in no case do bunt spores in the soil survive the winters of northern Europe. At that time they believed that infection from spores in the soil seldom, if ever, occurs.

Honecamp, Zimmerman, and Schneider (167) conducted some experiments to determine: (1) The toxic effect of bunt fed to animals, (2) the effect on the spores produced by passing through the animal, and (3) the probability of wheat infection from manure-borne spores or from spores reaching the manure after its evacuation. They found that spores passing through the animal had in great part lost their viability, but that this loss was less pronounced in the case of swine than with other animals.

Appel and Riehlm (20) conclude that infection of wheat by way of stall manure containing spores which have passed through the digestive tract of animals is so slight as to be disregarded in practice. Thus their results in general agree with those of Tubeuf, but are not in entire accord with the conclusions of Steglich. Although infection from spores mixed with the manure after evacuation does obtain, it is, nevertheless, slight in comparison with that resulting from spores adhering to the seed. Regarding the duration of bunt viability in the soil, Appel and Riehlm maintain that in uninjured bunt balls the spores can lie in damp soil for some time without germination. Thus infection from such balls might result, provided they are able to withstand external influences and remain intact until broken by tillage or seeding operations. The spores resist cold, and in dry storage they lose but little viability in two years. Appel and Riehlm further state that the conidia are resistant to weather influences so long as the soil contains sufficient moisture, but that they are destroyed by dryness. An infection due to such soil infestation would, therefore, be dependent on the soil-moisture factor.

The above conclusions regarding the life of the conidia in the soil would seem to support Brefeld's theory of the continuous saprophytic life subject only to moisture relations. It is the opinion of the writers that Honecamp and others are mistaken in this conclusion and that they have overlooked a most important factor, i. e., the organic flora of the soil. The studies of the writers have consistently shown that *Tilletia* grows readily enough in soil or on nutrient media as long as conditions are such as to prevent its being overrun with other organisms.

Concerning the possibility of a crop becoming infected by means of wind-borne bunt spores from threshing machines, there is little to be found in the literature. It was first suggested by Saunders.

Bedford, and Mackay (332), who recommended treating bunt-free seed with a strong solution of copper sulphate as a protection against such wind-distributed spores. The idea is next found in the same paper in which Maddox (255) presented his evidence of blossom infection of the loose smuts in 1896. Maddox here notes that bunt is very liable to appear in a fallow field lying to the lee of threshing operations.

Appel (14) relates a specific instance of such infection. A machine threshing a bunt-free lot of wheat was so situated that the dust was blown across a part of a field that was sown soon afterwards, and the crop was about half bunt in this part and bunt free in another part that was not exposed to the wind-blown dust.

PHYSIOLOGIC RESPONSE OF SPORES OF *TILLETIA* TO TEMPERATURE.

Hoffmann (164) investigated the thermolethal points of several of the Ustilagineae, but did not include *Tilletia* in his experiments.

Schindler (336), on the contrary, confined his work to *Tilletia caries* Tul. He found that spores were somewhat injured by an exposure to 65° C. of dry heat for a period of two hours, but that the absolute death point for a 2-hour exposure was between 95° and 100° C.; that when placed in water for two hours at 40° C., they were somewhat injured; and that the death point lay between 45° and 50° C. He states that an exposure of -16° C. of dry cold for two hours was not injurious and that the spores were not killed by outdoor exposure in the soil from December 5 to January 16. During this period the average daily minimum temperature was -8.5° C. and the average maximum -1.3° C. According to Jensen, a 5-minute exposure to damp heat ranging from 52.5° to 60° C. is fatal to the spores of *Tilletia*. Kühn states that most of the spores are killed by a 5-minute bath at 52.5° C. According to Kirchner (201), they are all killed within five minutes at 55° and 56° C.

Tubeuf (372, 374) germinated spores in water that had been subjected to a temperature of 37.5° to 38° C. for 24 hours. There was, however, a tendency to abnormality in the germination. He obtained no germination from those exposed for three days to a temperature of 39° to 40.5° C.

Volkart (384) found that spores of *Tilletia* would not germinate at and above a temperature of 30° C. but were uninjured at that temperature and germinated perfectly in four days when transferred to suitable temperatures.

CONTROL OF BUNT.

HISTORY OF SEED TREATMENT FOR THE PREVENTION OF BUNT.

Seed treatment for the prevention of bunt, by one method or another, has been practiced nearly three centuries. Before the true nature of the disease was known many of the more intelligent farmers conceived methods of seed treatment on the theory that the seed needed a stimulus to enable it to resist the disease, hence the use of fermenting animal manures. With the idea of the infective nature of smut came the use of toxins. The origin of the use of copper sulphate has been generally credited to Kühn, but he makes no claim to originality and credits it to Prévost, who in turn passes it back to Tessier. The fact is, it is of still earlier origin, for it was recommended

by Schulthess (342), and Taylor (367) of England recommended its use at the rate of half a pound per bushel of seed. So far as the writers have been able to ascertain, Kühn (221) was the first to investigate carefully the effect of copper sulphate on bunt spores as well as upon the seed and to recommend a definite formula consisting of immersion for 12 to 14 hours in a 0.5 per cent solution. His further experiments in 1873 (222) resulted in the conclusion that soaking for 12 hours was sufficient and that injury to seed was negligible. He found that when compared with seed soaked 12 hours in water, the treatment caused a slight reduction in the average number of roots but increased by about 30 per cent the sum total of root length. He does not state the time interval between the beginning of the germination experiment and measuring the roots, nor does he say how the grain was threshed.

Dreisch (101) conducted like investigations and came to a different conclusion regarding seed injury caused by the copper sulphate. He found that soaking in a solution as weak as 0.1 per cent for one hour resulted in root injury and inhibited germination by hardening the testa of the grain. He further found that such seed showed a decided difference in behavior when germinated in soil instead of in a moist chamber and that when sown in the ordinary way it suffered but little loss. He attributes this to lime in the soil and states that an aftertreatment with milk of lime counteracts the injury from copper sulphate. The only available abstract of Dreisch's article says nothing about the kind of seed used. It is entirely probable that the discrepancies between his results and those of Kühn are due to seed injury.

Nebbe (287) had compared the effect of copper-sulphate treatment on hand-threshed and machine-threshed grain and found the latter to be much more susceptible to injury. He found, however, that grain with abraded seed coats germinated more promptly than sound seed, which fact he attributes to the more rapid absorption of water. Like Dreisch, he found that the soil neutralized the injurious effect of copper sulphate on the seed. He believed this was due to chemical reaction between soil substances and the copper sulphate in which the toxic property of the copper was destroyed. He found support for this belief in the results of an experiment in which 152.02 grams of air-dry soil was wet with 150 cubic centimeters of a 2 per cent solution of copper salt. This was filtered off and followed by 12 washings. The total amount of filtrate was 2,240 cubic centimeters, and contained only 7 per cent of the copper salt originally added. Grassmann (137) investigated the effect of copper sulphate on machine-threshed wheat, immersing the grain in solutions varying in strength from 0.1 to 2 per cent. He found that the 0.1 per cent solution caused but little injury, but that the injury increased with the strength of the solution, resulting in total loss of viability at 2 per cent strength. He found also that seed injury increased with the lapse of time between treatment and sowing for a period of 10 days. Seed treated with a 0.5 per cent solution 16 hours and tested for viability showed increasing injury as follows: 24 hours after treatment, 34 per cent; 48 hours, 43.07 per cent; 72 hours, 47.64 per cent; 6 days after treatment, 51.33 per cent; 10 days after, 55.73 per cent.

Jensen (184) announced the results of experiments involving the use of hot water for the prevention of smut. Up to this time his studies were confined chiefly to oats and barley. This method, as

well as dry-heat sterilization, held an important place in subsequent investigations.

Experiments by Kellerman and Swingle in 1889 (196) and Kellerman in 1890 to 1896 (191-194) led them to conclude that of the many methods of seed treatment tested by them, the hot-water method was best for general use. Their experiments also established the practical value of potassium sulphid as a preventive of oat smut and the covered smuts of wheat and barley. The results of these experiments indicated that potassium sulphid reacted favorably on the germination of the seed and on the early growth and vigor of the plants. It was found that the promptest germination resulted from seed soaked 24 hours in a 0.5 per cent solution of potassium sulphid where this test was made immediately following treatment. The prompt germination and stimulation of seedling vigor were considered as factors favoring rapid growth and hence possible escape from infection.

Arthur in 1890 (25) concluded that seed wheat was practically uninjured by hot-water at 54° C. for 5 minutes. Experiments by Hickman (151) supported Kellerman's findings as to increased yield following treatment with hot water. Harwood and Holden (143) found that a 10-minute treatment at a temperature of 57.7° C. or less produced no change in germinative energy.

Kirchner (201) reviews the results of all experimental study of the Jensen method and gives the data resulting from his own experiments, which, while not extensive, were carried out with such careful regard for all factors and details as to give the results high value. His conclusions favor the Jensen method.

Geuther (133) suggested the use of formaldehyde as a sterilizing agent in combating smuts in general and supported his recommendation by experimental evidence. Bolley (56) gives the results of laboratory and field experiments in the treatment of seed wheat for the years 1893 to 1896, inclusive. In these experiments he used copper sulphate, mercuric chlorid, hot water, and formaldehyde. His is the first recorded use of formaldehyde in the United States as a fungicide for the prevention of bunt. As to the efficacy of a bunt preventive, he considered corrosive sublimate superior to all other fungicides, with the possible exception of formaldehyde, which at that time had received only preliminary tests.

Kittlauss (213) reported on extensive experiments regarding the injury to seed grain from the use of copper sulphate. His opinion was that the degree of damage was to some extent dependent on injury to the seed coat.

Falke (106) found that an application of 0.5 per cent copper sulphate for a period of 16 hours produced no injury when the seed coat was uninjured and notes that injury was greater to machine-threshed than to flail-threshed grain.

From 1900 to 1903 Farrer (109, 110, 113, 114) carried on extensive seed-treatment experiments, from which he concluded: (1) That bunt seed wheat could not be sufficiently cleaned by repeated washings in water to make its use safe. (2) The spores within unbroken bunt balls could not be killed by any treatment that the wheat could survive. (3) Copper sulphate was fairly efficient in killing the spores on the grain, and, although it was more injurious to the seed than either formaldehyde or mercuric chlorid, it furnished greater protection against reinfestation. (4) Formaldehyde was cheap.

effectual, and safe; and compared with copper sulphate was noninjurious to the seed, but afforded no protection against reinfestation.

Tubeuf (372, 374) states that laboratory experiments seemed to show that spores of *Tilletia tritici* could be killed with an application of formaldehyde gas that would not injure seed wheat, but in field experiments as high as 15 per cent of smut occurred in the crop grown from seed thus treated. Damp grain was severely injured by the gas treatment. The use of gas, therefore, was not recommended.

According to Tubeuf, a 12-hour treatment with a solution of 0.5 per cent copper sulphate reduced the yield of both plants and heads. The injury caused by copper sulphate he reduced by an afterbath in milk of lime. Machine-threshed seed he found to be more susceptible to copper-sulphate injury than flail threshed. All grain treated with copper sulphate germinated better in earth or sand than in filter paper. Disinfection with hot water at 55° to 56° C. for 15 minutes resulted in complete freedom from bunt and no injury to seed, but he considered its use inconvenient. Seed which had been washed in water and subsequently immersed in a 0.1 per cent solution of formaldehyde for a period of 4 hours, followed by quick drying, showed no resultant injury. This treatment he found thoroughly effective in killing the smut spores, and he also proved Bordeaux mixture to be entirely effectual but likely to result in seed injury if the action is too prolonged.

Tubeuf (374) gave attention to the effect of incrusting the seed with material containing (1) nutritive substances, such as Chile saltpeter, potassium salts, kainit, and superphosphate; and (2) toxic substances.

Peacock (295), using copper sulphate at the rate of 1 pound to 4 gallons of water in an open tank from 5 to 10 minutes, obtained an average of 86.3 per cent germination and a bunt-free crop. A lot of seed resmutted after treating produced 4.5 per cent of bunted heads. Another lot, washed after treating and resmutted, produced 10 per cent of bunted heads, while an untreated check plat gave 35 per cent of bunt.

Falke (108) found that injury from formaldehyde treatment is entirely prevented by rapid drying at 51° to 53° C. Volkart (384) published the results of a careful and comprehensive inquiry into the effects of the various treatments upon both the spores of bunt and those of oat smut, as well as upon the seed and the resulting harvest yield. Regarding the treatments with copper sulphate, the work showed conclusively that *Tilletia* spores were not killed in the ordinary methods of its use, but only delayed or at best prevented from germinating, and that the viability of the spores could be restored by washing in either dilute hydrochloric or acetic acid or by placing the spores in the soil. He found for *Tilletia*, as Hecke had found for covered smut of barley, *Ustilago hordei*, that a reaction took place between the spores and the copper sulphate in which the copper ion united with some element of the epispore to form a compound insoluble in water, while the sulphate ion forms a soluble compound with some element or compound drawn from the spore. It follows from this that when a solution of copper sulphate is being used in farm practice for treating wheat that contains a large amount of bunt it will lose effectiveness much more rapidly than when the wheat is comparatively clean. It was found that the lethal effect of copper sulphate was increased by raising the temperature of the treating bath from

20° to 33° C. With formaldehyde, no germination was observed in spores treated 2 hours in a 0.1 per cent solution. Those treated 1 hour and washed gave complete germination in 14 days, while untreated spores gave complete germination on the fifth day.

Sutton and Pridham (362), Ditzell and Downing (99), Sutton and Downing (361), and Soutter (348, 349) performed some experimental work to determine what fungicidal treatments gave the most effectual protection against infection through reinfestation of the seed. They all agree that copper sulphate gave the most satisfactory results.

In 1918 Darnell-Smith and Ross (94) reported on three years' experiments in the use of dry copper carbonate as a seed treatment for bunt prevention. They found it effectual as a fungicide and noninjurious to the seed. In fact, they believed it stimulated germination and increased the yield. These conclusions have been fully confirmed in experimental results obtained by Mackie and Briggs (254).

Störmer (357, 358) studied the relative merits of a number of treating processes, in both laboratory and field, in 1909 and 1910. He asserts that a 16-hour immersion in a 0.5 per cent solution of copper sulphate kills the spores in unbroken balls. This statement is open to question and is not confirmed by results obtained by other investigators. He failed to restore viability to spores treated with copper sulphate by washing them with 0.5 per cent solution of hydrochloric acid.

Hiltner (155) and Hiltner and Gentner (161) urge the use of mercuric chlorid as a treatment for the seed of winter grains, chiefly as a protection against *Fusarium* and other soil fungi. Their experimental evidence of a resultant increase in plant vigor and in the prevention of winterkilling is so convincing as to warrant our consideration of its more general adoption as a bunt preventive.

Riehm (326) conducted some experiments on the lethal effect of certain synthetic dyes on *Tilletia* spores and found that none of them killed the spores in the unbroken ball. Spores in balls, treated 10 minutes in a 1 per cent solution of Antiavit green and Antiavit blue, were delayed in germination. Free spores were killed by Antiavit blue, Antiavit green, Antiavit, Victoria blue, and acid violet, but chlorophenate of mercury, 0.05 per cent solution, killed spores in one hour with negligible injury to the seed grain.

Lehn and Vasters (237) found that the amount of spores carried on the seed grain was approximately proportional to the percentage of bunt in the resulting crop. Gentner (132) found treatment with mercuric chlorid a perfect protective against *Penicillium* and *Fusarium*, while formaldehyde promoted the attack of these fungi. He says this fungicide is much used in Denmark, Sweden, and Russia in the treatment of seed wheat and rye.

Remy and Vasters (322) experimented with a proprietary fungicide known as Uspulun, the active principle of which is chlorophenate of mercury, which is reported to afford perfect control of bunt without injury to the seed when used in a strength of 0.25 per cent for an immersion period of one hour.

Weck (391) records the results of experiments with Uspulun, which he carried on for three years. He found that the action of this fungicide as a bunt preventive was perfect, that the viability of the seed was unaffected, and subsequent growth of the seedling was stimulated.

Müller and Molz (275, 276) report on additional methods of seed treatment. Germination tests were made in moist chambers and in the open field involving the sowing of 8,000 grains in each trial. Entire bunt prevention was obtained in but one case, viz, when paraformaldehyde was used, but this fungicide showed the highest degree of seed injury.

Reynolds (323) performed extensive experiments with many fungicides to determine their protective properties against reinfestation. His conclusions were that any treatment involving the use of copper sulphate, except immersion in Bordeaux compounds, was highly protective, and that aftertreatment with lime reduced the protection. Dry copper carbonate mixed with the seed gave the best results.

Saunders, Bedford, and Mackay (332) in 1893 recommended copper sulphate in preference to other media for treatment because of its protective action against infection from bunt-infested soil. But so far as the authors have been able to ascertain no experiments of this kind had been undertaken in highly infested soil. It was thought that a sufficient amount of soluble toxin might be made to adhere to the seed to establish an immunizing zone around the grain which would prevent or reduce the infection from infested soil. Obviously it could not be expected that protection by means of seed-borne fungicides could be made to extend to the surface of the soil, but it was hoped that it would be effectual in the region where the seedling is most subject to infection, i. e., immediately above the seed.

VARIETAL RESISTANCE TO BUNT.

In 1764 N. E. Tschärner (371) stated that for several years he had sown a mixture of red and white spelt and always found much less bunt in the former. It was not until 130 years later, however, that a scientific search was begun for resistant varieties.

In 1900 William Farrer (110), the noted wheat breeder of New South Wales, reported on a number of varieties which he had studied. His investigations must have been made under severe conditions, for with two exceptions the average percentage of bunt was 83.

Kühn (225) noted that certain varieties were less susceptible to bunt than others and suggested that this fact might be related to moisture content of the plant, i. e., that low moisture content might enable the plant to resist entrance of the mycelium or to check its spread within the host.

Tubeuf (372, 374) gives in great detail the results of extensive varietal resistance experiments. He noted the number and percentage of plants partly bunted, as well as the number of sound and bunted heads appearing in each of such plants. Tubeuf is of the opinion that resistance is correlated with the varietal characters of prompt germination and rapid growth up to the point where infection is inhibited, i. e., the appearance of the first leaf. He found Ohio,⁵ Ontario, and Besthorn's Ueberfluss highly resistant. He suggests the possibility of increasing the resistance of a variety by selection through a long period of years. He found no uniform improvement as a result of selection for only one year.

Appel and Gassner (15) mention Ohio and Strubes Grannenweizen as being constantly resistant. Kirchner (204) published the results

⁵ So far as the writers have been able to determine, this wheat is unknown in the United States.

of studies begun in 1903. He tested 208 varieties, including both winter and spring wheats, of *Triticum vulgare*, *T. compactum*, *T. turgidum*, *T. durum*, *T. spelta*, *T. dicoccum*, *T. polonicum*, and *T. monococcum*. The results varied from 0 to 85.92 per cent bunt. Ohio was bunt free for 1905 and gave 0.7 per cent bunt in 1906. In most winter varieties that were tested in both years, 1903 and 1904, the percentage of bunt was much larger in the crop of the second year. The 1903 sowing was made on September 9 and that of 1904 on October 7. One variety of einkorn was tested for three years and in all three was bunt free. In his article published in 1908, Kirchner (206) expresses doubt as to resistance being dependent on quick germination and growth as a varietal character, though he concedes that such characters would lessen liability by shortening the period of susceptibility. He says it is necessary to investigate closely to what extent these traits are inherent transmissible characters rather than changeable phenomena due to external factors. Kirchner continued his investigations up to 1916. Important reports were published by him in 1913 (207) and 1916 (208). He (209) states that his investigations have been carried on since 1903, during which time he has tested for resistance to bunt 241 varieties of winter wheat and 119 of spring wheat. In winter wheats he found Cimbal's Fürst Hatzfeld and Hohenheimer No. 77 very resistant. The last named was attacked in his experimental plats but four times in 10 years. Some of the Galician wheats gave favorable, though somewhat inconstant, results. Beardless Odessa was tested six times, four tests being smut free, while one gave 2.5 per cent of bunt and another 0.56 per cent. These results agree with those obtained by Hecke (144, 146). Miracle, a smooth, red rivet wheat, was bunt free four years in succession. Kirchner found the durum wheats generally hard to infect, and two varieties proved very resistant. He described them as white, hard, smooth, the awns of one variety black, the other white. The Polish wheats were hard to infect. Of the spelts, two blue ones never showed infection. He notes that the same botanical group will contain varieties of high and low susceptibility. He recognizes external factors as having a great influence on infection, and he agrees with Hecke that degree of resistance is a constant hereditary varietal character. He was unable to establish any relation between germinative power and resistance. His observations led him to the conclusion that resistance is dependent on chemical composition and that the acid content of resistant varieties is comparatively higher than that of susceptible ones possessing the same morphological characters. In 1909 Pye, of Australia (315), stated that for several years he had been trying to obtain resistant varieties possessing other desirable qualities by crossing resistant varieties with other high-typed wheats which possessed some resistance. In these crosses he made considerable use of Medeah, a bearded durum. In his varietal tests, the highest percentage of bunt obtained was 58. The well-known wheats shown in Table 3 are selected from his results.

McAlpine (253) reports the following percentages of bunt infection: Federation, 88 to 100; Ohio, 4.3; Genoa, 4.5.

Hecke, in his admirable paper, "Der Einfluss von Sorte und Temperatur auf den Steinbrandbefall" (147), in 1909, gives the results

of exact and comprehensive experiments which are of high scientific value in that he takes into consideration the temperature factor in tests of varietal resistance. Ohio wheat, previously almost immune, in his experiments yielded as high as 61.9 per cent of bunted heads.

TABLE 3.—*Comparative resistance to bunt shown by well-known varieties of wheat in Pye's experiments.*

Variety.	Percentage of bunt.	Variety.	Percentage of bunt.
Queens Jubilee.....	58.00	Mudgee.....	6.00
Federation.....	49.00	Cedar.....	14.00
Bobs.....	43.00	Florence.....	3.03
Medeah.....	0	Genoa.....	2.01

Hecke's conclusions in part are that resistance is a varietal character and is transmissible to the progeny in crosses, but that it is modified by other factors, one of which is temperature, and that varieties react differently to external factors, temperature in particular. He is very skeptical as to resistance being constantly related to the characters of prompt germination and growth, though he recognizes the fact that the shorter the time from the beginning of germination to the appearance of the first leaf, all other factors being equal, the less the liability to infection.

Darnell-Smith (91) finds no evidence that rapidity of growth is related to resistance, but notes that Florence, Cedar, and Medeah, all resistant, have but little brush as compared with susceptible varieties and suggests that this lack of capacity to hold a larger amount of spores may be a factor. Cobb's (84) observations on the amount and character of the brush of the variety Medeah confirm those of Darnell-Smith.

SUMMARY.

Bunt of wheat has been known from very early times and is referred to by Theophrastus and other early Greek and Roman writers.

Down to the time of Tillet (1755) bunt was believed to be caused by environmental or providential influences.

In 1755 Tillet proved by experiment its infective character.

Prévost (1807) demonstrated the fungous origin and nature of bunt.

Kühn (1858) established the fact that bunt arises from infection of the wheat seedling.

The use of copper sulphate as a bunt preventive was first suggested by Schulthess in 1761, and Kühn (1866) first evolved a formula for its use.

The hot-water treatment of seed grain for the prevention of smuts was recommended by Jensen in 1888.

The use of formaldehyde as a general seed disinfectant was suggested by Geuther in 1895.

Maddox (1896) demonstrated that loose smut (*Ustilago tritici*) is caused by a fungous infection of the blossom, thus establishing the fundamental difference between bunt and loose smut.

Appel (1913) recorded the observation that a bunt crop was caused by spores blown from a threshing machine.

BIBLIOGRAPHY.

[Owing to the fact that many of the published contributions to our knowledge of bunt, or stinking smut, of wheat, were not accessible to the writers the bibliography here presented is not exhaustive. The originals of the references marked with an asterisk (*) were not available and were not examined. Many titles bearing on the subject appear in the bibliography which are not cited in the text.]

(1) ANONYMOUS.

1719. Experimentirter Nutzen der Invention das Getrayde, vermittelst der Asche, vor Brand zu bewahren. *In Samml. Nat. u. Med. Gesch.*, 1718, Sommer-Quartal, p. 1409–1412.

(2) 1719. Neuentdeckte Ursache des Brandtes in dem Getreide. *In Samml. Nat. u. Med. Gesch.*, 1718, Winter-Quartal, p. 594–596.

(3) 1720. Von einem neuen Experiment das Korn zu vermehren, und solches vor Brand zu bewahren. *In Samml. Nat. u. Med. Gesch.*, 1719, Frühlings-Quartal, p. 490–493.

(4) 1756. Beytrag zu den Mitteln den Brand in dem Weizen zu verhindern. *In Hannoversche Anz.*, 1855, Nützliche Saml., Stück 100, p. 1594–1598.

(5) 1764. Anzeige einer leichten Zubereitung des Getreides um 1764 die Saat. vordem Mehltau und dem Brande zu verwahren. . . . *In Abhandl. u. Beobacht. Oekonom. Gesell. Bern*, Jahrg. 5, Stück 2, p. 43–59.

(6) 1765. Expériences de M. Tillet; épi de froment ergoté. *In Hist. Acad. Roy. Sci. [Paris]*, ann. 1759, p. 116–118.

(7) 1768. Vom Brand im Getreide. *In Abhandl. u. Beobacht. Oekonom. Gesell. Bern*, 1768, Stück 2, p. 138–141.

(8) 1854. Versuche über die Wirkung verschiedener Beizen bei Saatfrucht. *In Centbl. Gesam. Landeskultur*, Jahr. 5, No. 50, p. 398–399.

(9) ALBERT, FRIEDRICH.

1898. Zur Bekämpfung des Steinbrandes beim Weizen. *In Deut. Landw. Presse*, Jahrg. 25, No. 87, p. 920–921, fig. 679–680.

(10)* APPEL, OTTO.

1904. Der Steinbrand des Weizens und seine Bekämpfung. *In Landbote*, Prenzlau, 25, 1904, p. 1073–1074; *Landw. Ann. Rostock*, N. F. 43, 1904, p. 325–327; *Westpr. Landw. Mitt. Danzig*, 9, 1904, p. 231–232. (Not seen. Reference from *Internat. Cat. Sci. Lit.*, M, Bot., v. 5, 1905/06, p. 47. 1908.)

(11) 1904. Der Steinbrand des Weizens und seine Bekämpfung. K. Gsmdtsamt. Biol. Abt. Land- u. Forstw. Flugbl. 26, 4 p., 8 fig.

(12) 1909. Aussprache über die bei den verschiedenen Methoden der Brandverteilung bei Kulturpflanzen gemachten Erfahrungen. *In Jahrb. Deut. Landw. Gesell.*, Bd. 24, Lfg. 2, p. 319–333.

(13) 1909. Theorie und Praxis der Bekämpfung von *Ustilago tritici* und *Ustilago nuda*. *In Ber. Deut. Bot. Gesell.*, Bd. 27, Heft 10, p. 606–610.

(14) 1913. Das Auftreten und die Bekämpfung des Steinbrandes. *In Mitt. Deut. Landw. Gesell. Beil. (Flugbl. 16)*, 3 p., 3 fig.

(15) ——— and GASSNER, GUSTAV.

1907. Untersuchungen über den Brand, ins besondere den Flugbrand des Getreides. *In Mitt. K. Biol. Anst. Land- u. Forstw.*, Heft 4, p. 9–12, illus.

- (16) APPEL, OTTO and RIEHM, EDUARD.
1910. Untersuchungen über die Brandkrankheiten des Getreides. *In* Mitt. K. Biol. Anst. Land- u. Forstw., Heft 10, p. 7-11.
- (17) 1911. Die Bekämpfung der Flugbrandes von Weizen und Gerste. *In* Arb. K. Biol. Anst. Land- u. Forstw., Bd. 8, Heft 3, p. 343-426, 2 fig., pl. 3 (col.).
- (18) 1911. Versuche über die Keimfähigkeit verfütterter Steinbrandsporen. *In* Mitt. K. Biol. Anst. Land- u. Forstw., Heft 11, p. 12.
- (19) 1912. Untersuchungen über die Brandkrankheiten des Getreides. *In* Mitt. K. Biol. Anst. Land- u. Forstw., Heft 12, p. 9-14.
- (20) 1914. Zur Frage der Ueberwinterung des Steinbrandes im Boden. *In* Mitt. K. Biol. Anst. Land- u. Forstw., Heft 15, p. 6.
- (21) APPL, JOHANN.
1915. Saatzeit und Steinbrandbefall des Weizens. *In* Ztschr. Landw. Versuchsw. Oesterr., Jahrg. 18, Heft. 3, p. 45-54.
- (22)* ARDUINO, PIETRO.
1770. Modi di preparare la semenza per preservare il frumento dal carbone. Venezia. 1770. Milocco. 8. (Not seen. Reference from Re, Filippo. Dizionario ragionato di libri d'agricoltura . . . t. 1, p. 229. Venezia. 1808.)
- (23) ARIETI, GIUSEPPE.
1900. I trattamenti preventivi dei cereali contro la carie ed il carbone. *In* Staz. Sper. Agr. Ital., v. 33, fasc. 5, p. 405-429. Bibliografia, p. 424-429. Abstract in Jahresber. Pflanzenschutz [Hollrung], Bd. 3, 1900, p. 41, 204. 1902.
- (24) ARTHUR, JOSEPH CHARLES.
1899. Formalin for grain and potatoes. *In* Ind. Agr. Exp. Sta. Bul. 77, p. 38-44.
- (25) 1890. Treatment for smut in wheat. *In* Ind. Agr. Exp. Sta. Bul. 32, p. 3-9.
- (26) ——— and JOHNSON, A. G.
1910. The loose smut of oats and stinking smut of wheat and their prevention. *In* Ind. Agr. Exp. Sta. Circ. 22, 15 p., 9 fig.
- (27) AUCANTE.
1756. Schreiben, von schädlicher Zubereitung der Getraidesaat. *In* Hannoversche Anz., 1755, Nützliche Saml., Stück 75, p. 1194-1198.
- (28) AUMÜLLER, FRITZ.
1920. Steinbrand und Streifenkrankheit als Sorten- und Linienkrankheit. *In* Deut. Landw. Presse, Jahrg. 47, No. 54, p. 383-384.
- (29) AYMEN.
1763. Von der Natur, dem Size und den Ursachen des Brandes im Getreide. *In* Abhandl. u. Beobacht. Oekon. Gesell. Bern, Jahrg. 4, Stük 1, p. 217-223. Translated from the English.
- (30)* BALSAMO-CRIVELLI, GIUSEPPE.
1857. Di alcune crittogame che furono riscontrate sul frumento nei contorni di Sesto di Monza ed altre località del Milanese. *Gaz. della Prov. di Pavia*, Juli, 1857. (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 1, p. 81, no. 1619. 1908.)
- (31) BANKS, JOSEPH.
1805. A short account of the cause of the disease in corn, called by farmers the blight, the mildew, and the rust. 31 p., fold. pl. London.
- (32) BARBECK, WILLIAM.
1879. Microscopical fungi infesting our cereals. *In* Amer. Nat., v. 13, no. 10, p. 612-620.
- (33) BARRUS, MORTIER FRANKLIN.
1916. Observations on the pathological morphology of stinking smut of wheat. *In* Phytopathology, v. 6, no. 1, p. 21-28, 3 fig.

- (34) BARY, ANTON DE.
1853. Untersuchungen über die Brandpilze und die durch sie verursachten Krankheiten der Pflanzen, mit Rücksicht auf das Getreide und andere Nutzpflanzen. 144 p., 8 pl. Berlin.
- (35) 1887. Comparative morphology and biology of the fungi, mycetoza and bacteria . . . Authorized English translation by Henry E. F. Garnsey . . . rev. by Isaac Bayley Balfour . . . xviii, 525 p., illus. Oxford. "Literature" at the end of each chapter.
- (36) BAUHIN, CASPAR.
1596. Phytopanax; seu. Enumeratio plantarum ad herbariis nostro seculo descriptarum, cum earum differentiis . . . 22 p. l., 669, 10 p., port. Basileae.
- (37) BEDFORD, S. A.
1893. Preventives of smut in wheat. *In Canada Exp. Farms Rpts.*, 1892, p. 196-197.
- (38) 1894. Test of bluestone as a smut preventive. *In Canada Exp. Farms Rpts.*, 1893, p. 237-238.
- (39) 1906. Preventives of smut in wheat. *In Canada Exp. Farms Rpts.*, 1905, p. 324-325.
- (40) BELLINI, MICHELE.
1827. Memoria sul modo di preservare la sementa del grano dal carbone o golpe. *In Atti R. Accad. Econ. Agr. Georg. Firenze*, t. 5, pt. 1, p. 268-272. Also in *Giorn. Agr. Tosc.*, v. 1, p. 526-530.
- (41)* BENDANDI, N.
1910. Contro il carbone dei cereali. I trattamenti cogli anticrittogamici sono inefficaci. *In Italia Agricola Piacenza*, 47, 1910, p. 490-492. (Not seen. Reference from *Internat. Cat. Sci. Lit., M, Bot.*, v. 10, 1910, p. 48. 1913.)
- (42) BENEVENUTI, GIUSEPPE.
1763. Abhandlung von den Ursachen des Brandes (Rostes) im Getreide und den Mitteln dagegen. *In Hamburg. Mag.*, Bd. 26, Stück 6, p. 553-569. [Bibliographical footnote], p. 553-561.
- (43) BERKELEY, MILES JOSEPH.
1847. Observations on the propagation of bunt (*Uredo caries*, D. C.) made with an especial reference to the potato disease. *In Jour. Hort. Soc. London*, v. 2, p. 107-114, illus.
- (44) 1856. Vegetable pathology.—No. CXLIV. Parasitae (*Tilletia*, bunt). *In Gard. Chron.*, 1856, no. 46, p. 757.
- (45) 1857. Introduction to cryptogamic botany. 604 p., illus. London, New York, Paris. "A list of some of the most useful works and memoirs," p. 569-579.
- (46)✓ BESSEY, CHARLES EDWIN.
1884. The wheat smuts. *In Iowa Agr. Coll. Bul.*, Nov. 1884, p. 118-126, fig. 3-4.
- (47) BOCK, HIERONYMUS.
1552. De stirpium, maxime earum, quae in Germania nostra nascuntur . . . 33 p. l., 1200, 64 p., illus., port. Argentorati.
- (48) BOLLEY, HENRY LUKE.
1891. Grain smuts. *N. Dak. Agr. Exp. Sta. Bul.* 1, 28 p., 7 fig.
- (49) ✓ 1895. Stinking smut in wheat. *In N. Dak. Agr. Exp. Sta. Bul.* 19, p. 125-130, fig. 1.
- (50) 1897. New studies upon the smut of wheat, oats, and barley, with a résumé of treatment experiments for the last three years. *N. Dak. Agr. Exp. Sta. Bul.* 27, p. 109-164, 13 fig.
- (51) 1899. Prevention of the smuts of cereal grains, and prevention of potato scab. *N. Dak. Agr. Exp. Sta. Bul.* 37, p. 363-379, illus.
- (52) 1900. Studies upon wheat and other cereals. *In N. Dak. Agr. Exp. Sta. 10th Ann. Rpt.* [1899/1900], p. 19-25, fig. 1.

- BOLLEY, HENRY LUKE—Continued.
- (53) 1900. The course of the hyphal filaments of *Tilletia* in the body of the wheat plant: A correction of an error. *In Proc. 21st Ann. Meeting Soc. Prom. Agr. Sci.*, 1900, p. 147-148.
- (54) 1902. The use of the centrifuge in diagnosing plant diseases. *In Proc. 23d Ann. Meeting Soc. Prom. Agr. Sci.*, 1902, p. 82-85.
- (55) 1913. Wheat: . . . Causes of soil sickness in wheat lands . . . N. Dak. *Agr. Exp. Sta. Bul.* 107, 96 p., 45 fig.
- (56) BRADLEY, RICHARD.
1725. A survey of the ancient husbandry and gardening, collected from Cato, Varro, Columella, Virgil, and others, the most eminent writers among the Greeks and Romans. 373 p., 4 pl. London.
- (57) BREFELD, OSKAR.
1883-1905. Die Brandpilze I-IV. *His Untersuchungen aus dem Gesamtgebiete der Mykologie*, Heft 5, 220 p., 13 pl., 1883; Heft 11, p. 1-98, 5 col. pl., 1895; Heft 12, p. 99-236, pl. 6-12, 1895; Heft 13, 75 p., 2 pl., 1905. Münster i. W.
- (58) 1888. Neue Untersuchungen über die Brandpilze und die Brandkrankheiten. II. *In Nachr. Klub Landw. Berlin*, No. 220, p. 1577-1584; No. 221, p. 1588-1594; No. 222, p. 1597-1601. (For English translation by Erwin F. Smith, see no. 60.)
- (59) 1889. Basidiomyceten III. *His Untersuchungen aus dem Gesamtgebiete der Mykologie*, Heft 8, 305 p., 12 pl. (partly col.). Leipzig.
- (60) 1890. Recent investigations of smut fungi and smut diseases. *In Jour. Mycol.*, v. 6, no. 1, p. 1-8; no. 2, p. 59-71; no. 4, p. 153-164. (Translation by Erwin F. Smith from *Nachr. Klub Landw. Berlin*, No. 220-222. See no. 58.)
- (61) 1903. Neue Untersuchungen und Ergebnisse über die natürliche Infektion und Verbreitung der Brandkrankheiten des Getreides. *In Nachr. Klub Landw. Berlin*, No. 466, p. 4224-4232. Discussion, p. 4232-4234.
- (62) 1907. Über die Brandkrankheiten des Getreides ihre natürliche Verbreitung und ihre Verhütung. *In Jahrb. Deut. Landw. Gesell.*, Bd. 22, Lfg. 1, p. 75-89. Discussion, p. 89-91.
- (63) 1908. Die Kultur der Pilze . . . *His Untersuchungen aus dem Gesamtgebiete der Mykologie*, Heft 14, 256 p. Münster i. W.
- (64) BRITTLERANK, CHARLES C.
1913. Effect of formalin and bluestone pickle on the germination of wheat. *In Jour. Dept. Agr. Victoria*, v. 11, pt. 8, p. 473-476, 2 fig.
- (65) BROILI, JOSEF.
1911. Über Versuche mit Brandinfektion zur Erzielung brandfreier Gerstenstämme. *In Fühlings Landw. Ztg.*, Jahrg. 60, Heft 3, p. 105-107.
- (66)* BROSLAVSKII, P.
1909. Novyi metod bor'by sel'skokhozyaistvennymi vreditel'ami. [Eine neue Methode zur Bekämpfung der Pflanzenkrankheiten.] *In Sel'skii Khoziaïn*, 1909, no. 23, 24, 25. (Not seen. Abstract in *Russ. Jour. Exp. Landw. god* 10, kniga 5, p. 683-684. 1909. Listed under German title in *Just's Bot. Jahresber.*, Jahrg. 38, 1910, Abt. 1, Heft 1, p. 219. 1911.)
- (67) BROWN, ADRIAN JOHN.
1907. On the existence of a semi-permeable membrane enclosing the seeds of some of the Gramineae. *In Ann. Bot.*, v. 21, no. 81, p. 79-87.
- (68) BROŽ, OTTO.
1911. Der Getreidebrand und seine Bekämpfung. *In Monatsh. Landw.*, Jahrg. 4, Heft 10, p. 289-293, 9 fig.

- (69) BRYANT, HENRY.
1784. A particular enquiry into the causes of that disease in wheat, commonly called brand; being an attempt to shew that the general practice of drenching the seed in salt and water, soap lees, quicklime, &c. are not only ineffectual for preventing the disorder, but are destructive to the seed. 58 p. Norwich.
- (70) BUCHANAN, J.
1907. Treatment of grain for smut. *In* 32d Ann. Rpt. Ontario Agr. Col. and Exp. Farm, 1906, p. 176-178.
- (71) BUESGEN, MORITZ.
1893. Ueber einige Eigenschaften der Keimlinge parasitischer Pilze. *In* Bot. Ztg., Jahrg. 51, Abt. 1, Heft 3/4, p. 53-72, pl. 3.
- (72) BURMESTER, HERMANN.
1908. Vergleichende Untersuchungen über den Einfluss der verschiedenen Samenbeizmethoden auf die Keimfähigkeit gebeizten Saatgutes und über ihre pilztötende Wirkung. *In* Ztschr. Pflanzenkrankh., Bd. 18, Heft 3, p. 154-187, pl. 6.
- (73) CAMERARIUS, RUDOLPH JACOB.
1709. De Ustilagine frumenti. 16 p. Tubingae.
- (74) CANDOLLE, AUGUSTIN PYRAMUS DE.
1807. Sur les champignons parasites. *In* Ann. Mus. Hist. Nat. [Paris], t. 9, p. 56-74.
- (75) 1832. Physiologie végétale . . . t. 3. Paris.
- (76) CARON-ELDINGEN, VON.
1920. Steinbrand und physiologische Spaltungen. *In* Deut. Landw. Presse, Jahrg. 47, No. 75, p. 514, fig. 492.
- (77) ——— and HEUSER.
1920. Nochmals: "Steinbrand und physiologische Spaltungen." *In* Deut. Landw. Presse, Jahrg. 47, No. 87, p. 598-599.
- (78)* CHLÁDEK, C.
1909. Der jetzige Stand unserer Kenntnisse über die Verbreitung der Getreidebrandpilze. *In* Zeitschr. der Lehrer der Landw. Anst. in Böhmen, Tabor, IX, 1909, p. 8-12. (Böhmisch.) (Not seen. Reference from Lindau and Sydow, Thesaurus, Litt. Mycol., v. 3, p. 155, no. 32057. 1913.)
- (79) CLINTON, G. P.
1902. North American Ustilagineae. *In* Jour. Mycol., v. 8, p. 128-156.
- (80) 1904. North American Ustilagineae. *In* Proc. Boston Soc. Nat. Hist., v. 31, no. 9, p. 329-529. Literature, p. 505-524.
- (81) 1905. The Ustilagineae, or smuts, of Connecticut. Conn. State Geol. and Nat. Hist. Survey Bul. 5, 43 p., 55 fig. on 7 pl.
- (82) 1906. Ustilaginales. *In* N. Y. Bot. Gard., No. Amer. Flora, v. 7, pt. 1, p. 1-82.
- (83) COBB, N. A.
1896. The hot-air treatment of bunt or stinking smut. *In* Agr. Gaz. N. S. Wales, v. 7, pt. 2, p. 82-83.
- (84) 1902. Comparative observations on the brush of about fifty varieties of wheat. *In* Agr. Gaz. N. S. Wales, v. 13, pt. 6, p. 647-649, 3 fig.
- (85) 1905. Quantitative estimation of bunt in seed-wheat. *In* Agr. Gaz. N. S. Wales, v. 16, pt. 11, p. 1113-1117.
- (86) COOK, MELVILLE THURSTON.
[1913?]. Grain smuts: Their causes and treatments. N. J. Agr. Exp. Sta. Circ. 36, 4 p.
- (87) COUSTON, F.
1911. Le formol contre la carie et le charbon des céréales. *In* Jour. Agr. Prat., ann. 75, sem. 1 (n. s. t. 21), no. 1, p. 14-16.

- (88) CRAMER, KARL EDUARD.
1890. Die Brandkrankheiten der Getreidearten, nach dem neuesten Stand der Frage. *In* Landw. Jahrb. Schweiz. Bd. 4, p. 9-16, pl. 4.
- (89) DAMSEAUX, ADOLPHE, and LAURENT, ÉMILE.
1899. Enquête sur la carie du froment en Belgique en 1898. *In* Bul. Agr. [Brussels], t. 15, livr. 3, p. 107-114.
- (90)* ————
1899. Contre la carie. Union, 1899, p. 453-454. (Not seen. Reference from Just's Bot. Jahresber., Jahrg. 28, 1900, Abt. 2, p. 468. 1902.)
- (91) DARNELL-SMITH, G. P.
1910. Observations upon the disease of wheat known as "bunt." *In* Rpt. Govt. Bur. Microbiol. [N. S. Wales], 1909, p. 64-69, 9 fig.
- (92) ————
1910. Some observations on bunt and fungicides. *In* Agr. Gaz. N. S. Wales, v. 21, pt. 9, p. 751-756, 4 fig.
- (93) ————
1915. Fungicide experiments, 1914. *In* Agr. Gaz. N. S. Wales, v. 26, pt. 6, p. 494-495.
- (94) ———— and ROSS, H.
1919. A dry method of treating seed wheat for bunt. *In* Agr. Gaz. N. S. Wales, v. 30, pt. 10, p. 685-692, 7 fig.
- (95) DAVID, ST.
1900. Zur Frage über die Wirkung des Formaldehyds auf Getreidesamen und Brandsporen. *In* Sitzber. Naturf. Gesell. Univ. Jurjeff (Dorpat), Bd. 12, 1899, Heft 2, p. 222-247. *In* Russian. German résumé, p. 202-204. Review in Jahresber. Pflanzenschutzes [Hollrung], Bd. 3, 1900, p. 42, 204. 1902.
- (96) DAVIS, HORACE.
1868. Wheat in California. *In* Overland Monthly, v. 1, no. 5, p. 442-452.
- (97) DEGRULLY, LÉON.
1904. Maladies des céréales. *In* Prog. Agr. et. Vit., ann. 21, no. 40, p. 371-374; no. 44, p. 502-504.
- (98) DEMOUSSY, ÉMILE.
1901. La germination des grains de blé traités au sulfate de cuivre. *In* Ann. Agron., t. 27, no. 6, p. 257-261.
- (99) DITZELL, F., and DOWNING, R. G.
1911. Some experiments with fungicides used for the prevention of "stinking smut" (Bunt), Cowra, 1910. *In* Agr. Gaz. N. S. Wales, v. 22, pt. 4, p. 341-357.
- (100) DOBENECK, VON.
1899. Umfrage im Königreich Bayern über Vorkommen und Bekämpfung des Weizenbrandes im Jahre 1898. *In* Vrtljschr. Bayer. Landw.-Rat., Jahrg. 4, Heft 3, p. 255-282.
- (101) DREISCH, FRANZ EMIL.
1873. Untersuchungen über die Einwirkung verdünnter Kupferlösungen auf den Keimproceß des Weizens. 50 p. Dresden. Inaug. Diss., Rostock.
- (102)* ÉGERT, K. L.
1914. [Potassium permanganate treatment for seed grains.] *In* Selsk. Khoz., 1914, p. 1343-1346. (Russian original not seen. Abstract in Mat. Mikol. i Fitopatol. Ross. [Jour. Mycol. and Phytopath. Russia], v. 1, no. 2, p. 66. 1915. English abstract in Exp. Sta. Rec. v. 34, no. 9, p. 844. 1916.)
- (103) EHRENBURG, PAUL.
1918. Zur Frage der Beizung des Winterweizens gegen Steinbrand. *In* Fühlings Landw. Ztg., Jahrg. 67, Heft 23/24, p. 425-432.
- (104) ERIKSSON, JAKOB.
1908. Om sot i hvete. *In* K. Landtbr. Akad. Handl. och Tidskr., årg. 47, no. 4, p. 274-279, 2 fig.

- (105) **FABRICIUS, JOHANN CHRISTIAN.**
1774. Forsøg til en Afhandling om Planternes Sygdomme. *In* K. Norske Vidensk. Selsk. Skr., deel 5, p. 431-492.
- (106) **FALKE, FR.**
1900. Die Dehne'sche Desinfektionsmaschine für Saatgetreide. *In* Landw. Wehnschr. Prov. Sachsen, Jahrg. 2, No. 41, p. 365-367, illus.; No. 42, p. 374-375.
- (107) 1902. Versuche zur Gewinnung brandfreien Saatgutes. *In* Deut. Landw. Presse, Jahrg. 29, No. 62, p. 518.
- (108) 1902. Ueber in Eckendorf angestellte Versuche zur Gewinnung von brandfreiem Saatgetreide. Erster Bericht. *In* Deut. Landw. Presse, Jahrg. 29, No. 70, p. 575-576; No. 71, p. 584-585, illus.
- (109) **FARRER, WILLIAM.**
1900. Some experiments in dealing with bunt or the stinking smut of wheat. *In* Agr. Gaz. N. S. Wales, v. 11, pt. 5, p. 335-344.
- (110) 1901. Results of the Lambrigg bunt experiments of 1900. *In* Agr. Gaz. N. S. Wales, v. 12, pt. 4, p. 419-430.
- (111) 1901. Bunt in self-sown wheat. *In* Agr. Gaz. N. S. Wales, v. 12, pt. 9, p. 1182-1183.
- (112) 1902. The effect on the milling quality and nutritive value of the resulting crop of wheat when bunt-infected seed is sown. *In* Agr. Gaz. N. S. Wales, v. 13, pt. 11, p. 1094-1097.
- (113) 1903. Bunt experiments of 1901. *In* Agr. Gaz. N. S. Wales, v. 14, pt. 3, p. 206-216.
- (114) 1905. The effect, in actual farm practice, of treatment with bluestone on the germination of wheat. *In* Agr. Gaz. N. S. Wales, v. 16, pt. 12, p. 1246-1247.
- (115) ——— and **SUTTON, GEORGE L.**
1905. The effects of some solutions of formalin and bluestone, which are in common use, on the germination of wheat seeds. *In* Agr. Gaz. N. S. Wales, v. 16, pt. 12, p. 1248-1255.
- (116) **FERLE, FR.**
1905. Beizversuche, ausgeführt an Weizen. *In* Fühlings Landw. Ztg., Jahrg. 54, Heft 18, p. 601-604.
- (117) **FISCHER VON WALDHEIM, ALEKSANDR ALEKSANDROVICH.**
1867. Sur la structure des spores des Ustilaginées. *In* Bul. Soc. Imp. Nat. Moscou, t. 40, pt. 1, no. 1, p. 242-261, pl. 3 (col.)
- (118) 1869. Beiträge zur Biologie und Entwicklungsgeschichte der Ustilagineen. *In* Jahrb. Wiss. Bot. [Pringsheim], Bd. 7, p. 61-144, pl. 7-12. Translation in Trans. N. Y. State Agr. Soc., 1870, p. 280-354, 6 pl. 1871.
- (119) 1876. Les Ustilaginées et leurs plantes nourricières. *In* Ann. Sci. Nat. Bot., s. 6, t. 4, p. 190-276.
- (120)* **FOURCROY, A. F. DE.**
1810. Ueber die chemische Beschaffenheit des brandigen Weizens. *In* Arch. Agr.-Chem. Denkende Landwirthe Herbstädt, IV, p. 296-301. (Not seen. Reference from Cat. Sci. Papers., Roy. Soc. Lond., v. 7, p. 694. 1877.)
- (121) **FRANK, ALBERT BERNHARD.**
1895-96. Die Krankheiten der Pflanzen. Ein Handbuch für Land- und Forstwirte, Gärtner, Gartenfreunde und Botaniker . . . Aufl. 2, 3 v. Breslau.
- (122) ——— and **SORAUER, PAUL.**
1898. Jahresbericht des Sonderausschusses für Pflanzenschutz 1897. *In* Jahrb. Deut. Landw. Gesell., Bd. 13, p. 466-467.

- (123) FRASSI, ALFREDO.

1913. Azione di alcuni disinfettanti sul potere germinativo delle cariossidi di frumento. *In Staz. Sper. Agr. Ital.*, v. 46, fasc. 1, p. 25-56. Bibliografia, p. 53-56.

- (124) FREEMAN, E. M., and JOHNSON, EDWARD C.

1909. The loose smuts of barley and wheat. U. S. Dept. Agr., Bur. Plant Indus. Bul. 152, 48 p., 2 fig., 6 pl.

- (125) ——— and STAKMAN, E. C.

1911. The smuts of grain crops. *Minn. Agr. Exp. Sta. Bul.* 122, p. 35-64, 11 fig.

- (126)* FRIES, ELIAS MAGNUS.

1821. Om brand och rost på växter, jemte fullständig underrättelse om deras kännetecken, orsaker, skada samt medel till dess förekommande. *In "Lösning för allmogen i Kronobergs län,"* No. 3, Lundae 1821, p. 1-51. (Not seen. Reference from Lindau and Sydow, *Thesaurus, Litt. Mycol.*, v. 1, p. 495, no. 9213. 1908.)

- (127) FUSCHINI, CARLO.

1912. Dei mezzi più idonei a combattere la "carie" ed il "carbone" del frumento. *In Staz. Sper. Agr. Ital.*, v. 45, fasc. 8, p. 549-586.

- (128) [GAILLOT, L.]

1903. Emploi du formol pour le traitement des graines de céréales contre le charbon et la carie. *In Sta. Agron. Aisne Bul.*, s. 2, ann. 13, 1902, p. 79-81.

- (129) GALLOWAY, B. T.

1889. A partial list of the parasitic fungi of Missouri, with remarks on the species of economic importance. *In U. S. Dept. Agr., Bot. Div. Bul.* 8, p. 54-59.

- (130) 1894. A new method of treating grain by the Jensen process for the prevention of smut.
- In Jour. Mycol.*
- , v. 7, no. 4, p. 372-373.

- (131) GAUL.

1920. Die Bekämpfung des Weizensteinbrandes in der Praxis. *In Deut. Landw. Presse, Jahrg.* 47, No. 78, p. 536-537.

- (132) GENTNER, GEORG.

1913. Kann Sublimat als Beizmittel gegen Pilzbefall des Getreides durch Chinosol und andere Mittel ersetzt werden? *In Prakt. Bl. Pflanzenbau u. Schutz, Jahrg.* 16 (n. F. 11), p. 6-12, illus.

- (133) GEUTHER, TH.

1895. Ueber die Einwirkung von Formaldehydlösungen auf Getreidebrand. *In Ber. Pharm. Gesell.*, Jahrg. 5, Heft 12, p. 325-329. Diskussion, p. 329-330.

- (134)* GOMILEVSKIJ, VASILII.

1906. Die Sterilisation des Saatkornes des Weizens von den Sporen des Stein- und des Flugbrandes. *In Selisk. Choziain, St. Petersburg*, 21, 1906, p. 763-764. *In Russian.* (Not seen. Reference from *Internat. Cat. Sci. Lit.*, M, Bot., v. 8, 1908/09, p. 124. 1910.)

- (135) GRADENWITZ, A.

1912. Der Widerstand der Getreidesorten gegen Pflanzenkrankheiten. *In Prometheus, Jahrg.* 23, No. 1147, p. 39-40, illus. Apparatus devised by Fr. Straňak.

- (136) GRÄF, K.

1913. Roggenbeizung mit Sublimat. *In Prakt. Bl. Pflanzenbau u. Schutz, Jahrg.* 16 (n. F. 11), Heft 8, p. 97-101, 4 fig.

- (137) GRASSMANN, P.

1886. Die Verluste beim Weizenbau infolge unzweckmässiger Anwendung des Kupfervitriols als Schutzmittel gegen den Schmierbrand. *In Landw. Jahrb.*, Bd. 15, p. 293-307.

- (138) GRÓH, JULIUS.

1912. Ueber die Bestimmung des Brandsporengehaltes in Kleien. *In Arch. Chem. u. Mikros.*, Jahrg. 5, Heft 4, p. 177-187, 2 fig.

- (139) GROSSER.
1913. Wissenswerte über das als Saatschutzzmittel angepriesene Präparat "Corbin." *In* Ztschr. Landw. Kammer Prov. Schlesien, Jahrg. 17, Heft 33, p. 1170-1171.
- (140) GÜSSOW, HANS THEODOR.
1913. Smut diseases of cultivated plants, their cause and control. Canada Exp. Farms Bul. 73, 57 p., illus.
- (141)* HABERLANDT, FRIEDRICH.
1874. Kupfervitriollösungen in ihrem Einflusse auf die Keimfähigkeit des Weizens. *In* Landwirtsch. Centralbl., 22, 1874, p. 353-356. (Not seen. Reference from Cat. Sci. Papers, Roy. Soc. Lond., v. 10, p. 104. 1894.)
- (142) HARPER, ROBERT ALMER.
1900. Nuclear phenomena in certain stages in the development of the smuts. *In* Trans. Wis. Acad. Sci., v. 12, pt. 2, 1899, p. 475-498, pl. 8-9.
- (143) HARWOOD, P. M., and HOLDEN, P. G.
1892. Smut in oats and wheat. Jensen or hot water treatment. Mich. Agr. Exp. Sta. Bul. 87, 10 p., illus.
- (144) HECKE, LUDWIG.
1906. Die Blüteninfektion des Getreides durch Flugbrand. *In* Jahresb. Angew. Bot., Jahrg. 3, 1904/05, p. 63-64.
- (145) 1906. Die Brandkrankheiten des Getreides und ihre Bekämpfung. *In* Wiener Landw. Ztg., Jahrg. 56, No. 33, p. 318-319, fig. 137-139.
- (146) 1907. Die Triebinfektion bei Brandpilzen. *In* Ztschr. Landw. Versuchsw. Oesterr., Jahrg. 10, Heft 6, p. 572-574.
- (147) 1909. Der Einfluss von Sorte und Temperatur auf den Steinbrandbefall. *In* Ztschr. Landw. Versuchsw. Oesterr., Jahrg. 12, Heft 2, p. 49-66, pl. 3.
- (148) HENDERSON, LOUIS FOURNIQUET.
1898. Smuts and rusts of grains in Idaho, and the most approved methods of dealing with them. Idaho Agr. Exp. Sta. Bul. 11, 34 p., 15 fig.
- (149) 1906. Experiments with wheat and oats for smut. Idaho Agr. Exp. Sta. Bul. 53, 15 p.
- (150) HENKEMEYER, A.
1920. Zur Bekämpfung des Weizensteinbrandes. *In* Deut. Landw. Presse, Jahrg. 47, No. 71, p. 489.
- (151) HICKMAN, J. FREMONT.
1891. Experiments in wheat seeding and treatment of seed for smut. *In* Ohio Agr. Exp. Sta. Bul., v. 4, no. 4, p. 77-89.
- (152) 1899. Experiments with oats. Seeding on different soil to exterminate smut. *In* Ohio Agr. Exp. Sta. Bul. 101, p. 179-181.
- (153) HILTNER, LORENZ.
1908. Ueber die Abhängigkeit der Brandanfälligkeit des Getreides von dessen Keimungsenergie und Entwicklungsgeschwindigkeit. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 11 (n. F. 6), Heft 6, p. 67-69.
- (154) 1910. Anweisung zur Beizung des Weizens gegen Steinbrand mit Formalinlösung. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 13 (n. F. 8), Heft 9, p. 110-111.
- (155) 1910. Über die Beizung des Wintergetreides mit Sublimatlösung. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 13 (n. F. 8), Heft 10, p. 114.
- (156) 1912. Über die Beizung des Saatgutes von Wintergetreide. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 15 (n. F. 10), Heft 9, p. 97-98.
- (157) 1913. Über die Wirkung der Sublimatbeizung des Winterroggens und des Winterweizens im Jahre 1912/13. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 16 (n. F. 11), Heft 8, p. 101-104.

HILTNER, LORENZ—Continued.

- (158) 1914. Über die Beizung des Saatgutes von Wintergetreide mit sublimat-haltigen Mitteln. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 17 (n. F. 12), Heft 8/9, p. 85-89.
- (159) 1915. Über die Erfolge der im Herbst 1914 in Bayern durchgeführten Beizung des Saatgutes von Winterroggen und Winterweizen mit sublimathaltigen Mitteln. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 18 (n. F. 13), Heft 6/7, p. 65-90.
- (160) 1915. Über die Beizung des Winterweizens gegen *Fusarium*, *Penicillium* und Steinbrand. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 18 (n. F. 13), Heft 8, p. 97-109, illus.; Heft 9, p. 113-124.
- (161) ——— and GENTNER, GEORG.
1912. Über die schützende Wirkung der Sublimatbeizung des Roggens gegen den Befall durch Bodenfusarien. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 15 (n. F. 10), Heft 11, p. 129-131.
- (162) HÜSTERMANN.
1913. Brandbekämpfungsversuch. *In* Landw. Jahrb., Bd. 45, Ergänzungsbd. 1, p. 107-111.
- (163) HOFFMANN, HERMANN.
1860. Untersuchungen über die Keimung der Pilzsporen. *In* Jahrb. Wiss. Bot. [Pringsheim], Bd. 2, p. 267-337, pl. 26-32.
- (164) HOLLRUNG, MAX.
1898. Handbuch der chemischen Mittel gegen Pflanzenkrankheiten. xii, 178 p. Berlin.
- (165) 1903. Zur Würdigung der sog. Kulturabeize (Cerespulver). *In* Landw. Wchnschr. Sachsen, Jahrg. 5, No. 36, p. 300-301.
- (166) HONCAMP, FRANZ.
1911. Untersuchungen über die Wirkung der Brandsporen im Futter und im Dünger. *In* Landw. Vers. Stat., Bd. 74, Heft 3/5, p. 364-376.
- (167) ——— and ZIMMERMANN, H.
1910. Untersuchungen über das Verhalten von Brandsporen im Tierkörper und im Stalldünger. Unter Mitwirkung von G. Schneider. *In* Centbl. Bakt. [etc.], Abt. 2, Bd. 28, No. 22/24, p. 590-607.
- (168) HORI, SHOTARO.
1901. Komungi no Namagusakurobo Yobo Shiken (Experiments for the prevention of bunt in wheat). *In* Nishigahara Imp. Agr. Exp. Sta. Tech. Rpt. 18, p. 1-5.
- (169) 1907. Seed infection by smut fungi of cereals. *In* Bul. Imp. Cent. Agr. Exp. Sta. Japan, v. 1, no. 2, p. 163-176.
- (170) HOWARD, ALBERT, and HOWARD, GABRIELLE L. C.
1907. Note on immune wheats. *In* Jour. Agr. Sci., v. 2, pt. 3, p. 278-280.
- (171) HUMPHREY, HARRY B., and POTTER, ALDEN A.
1918. Cereal smuts and the disinfection of seed grain. U. S. Dept. Agr., Farmers' Bul. 939, 28 p., 16 fig.
- (172) HURD, ANNIE MAY.
1921. Seed-coat injury and viability of seeds of wheat and barley as factors in susceptibility to molds and fungicides. *In* Jour. Agr. Research, v. 21, no. 2, p. 99-122, pl. 13-23. Literature cited, p. 120-122.
- (173) HURST, ROBERT J.
1911. Bunt and germination experiments. *In* Agr. Gaz. N. S. Wales, v. 22, pt. 9, p. 749-752.
- (174) ISOLANI, ALAMANNO.
1844. [Memoria intorno alla golpe del frumento.] (Abstract 6a, Sessione ordinaria, 7 Marzo 1812.) *In* Nuovi Ann. Sci. Nat. Rend. Soc. Agr. Bologna, s. 2, t. 1, p. 384.

- (175) ISSATSCHENKO, BORIS.
1909. Über die Bestimmung des Verunreinigungsgrades durch Brandsporen. *In* Bul. Bur. Angew. Bot., Jahrg. 1, 1908, No. 11/12, p. 273-276. *In* Russian, German summary, p. 276-277.
- (176)* ISTVÁNYFI, GYULA.
1891(?) Újabb vizsgálatok az ősök gombákról. (Neuere Untersuchungen über die Brandpilze. Supplementhefte zum T. K. Budapest, 1891. 16. Heft. p. 152-165. Mit Abb. Magyarisch.) (Not seen. Reference from Just's Bot. Jahresber., Jahrg. 19, 1891, Abt. 1, p. 210. 1894. Also given in Lindau and Sydow, Thesaurus, v. 1, p. 730, no. 13346, 1903, as follows: Természettud. Közlöny, XVI, Pótfüzet, 1893, p. 152-165. Ungarisch.)
- (177)* 1896. A gabonafélék ősökbetegségéről. (Ueber die Brandpilzkrankheiten der Getreidearten.) *In* Pótfüzet. Természettud. Közlöny, XXXIX, 1896, p. 157-171. (Magyarisch.) (Not seen. Reference from Just's Bot. Jahresber., Jahrg. 24, 1896, Abt. 1, p. 280. 1899.)
- (178) IVANOFF, K. S.
1903. Phytopathologisches aus Transkaukasien. *In* Ztschr. Pflanzenkrank., Bd. 13, Heft. 4, p. 221-222.
- (179)* JACZEWSKI, ARTHUR.
1901. Die parasitären Pilze der Kulturpflanzen. V. Der Brand des Getreides. St. Petersburg, Ackerbau Minist. u. Departm., 1901. (Russisch.) (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 3, p. 331, no. 34898. 1913.)
- (180) 1902. Golovnia psheinitsy i rzhi. (Die Brandkrankheit des Weizens und Roggens.) *In* Listok Boriby Boliezn. Rastenij Imp. Bot. Sad. St. Petersburg, god 1, no. 7, p. 49-52, fig. 7-8. *In* Russian. (German title from Internat. Cat. Sci. Lit., M, Bot., v. 2, 1902, p. 165. 1904.)
- (181) 1904. Obezrazhanie formalinom siemian khliebnykh zlakov i drugikh kul'turnykh rastenii. (Sterilisieren der Getreide- und anderer Kulturpflanzensamen mit Formalin.) *In* Listok Bor. Boliezn. Rastenij Imp. Bot. Sad, St. Petersburg, god 3, no. 4, p. 33-39. *In* Russian. (German title from Internat. Cat. Sci. Lit., M, Bot., v. 5, 1905/06, p. 172. 1908.)
- (182)* 1907. Der Steinbrand und seine Bekämpfung. *In* Věst. Selisk.-Choz., Moskva, 8, 1907, N. 46, p. 11-12. *In* Russian. (Not seen. Reference from Internat. Cat. Sci. Lit., M, Bot., v. 7, 1907/08, p. 161-162. 1910.)
- (183) 1910. Über die Beizung der Samen unserer Kulturgewächse mit Formalin. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 13 (n. F. 8), Heft 11, p. 130-132.
- (184) JENSEN, JENS LUDWIG.
1888. The propagation and prevention of smut in oats and barley. *In* Jour. Roy. Agr. Soc. England, s. 2, v. 24, p. 397-415.
- (185) 1889. Le charbon des céréales. 8 p. Copenhagen.
- (186)* 1891. Et par Meddelelser om Bortskaffelsen af Vaarsaeds Brand. *In* Markfrøkontorets Aarsberetninger 1890, p. 22-24. (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 3, p. 732, no. 41322. 1913.)
- (187)* 1892. Varmvandsmetoden mod Brand i Vaarsaeden. Kjöbenhavn, Marts 1892, 8 p. (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 3, p. 732, no. 41324. 1913.)
- (188)* 1895. Tilbageblik paa de senere Aars Forsøg med Varmvandsmetoden. *In* Markfrøkontorets Aarsberet. 1894, p. 17-20. (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 3, p. 732, no. 41329. 1913.)
- (189) JOHNSON, EDWARD C.
1912. The smuts of wheat, oats, barley, and corn. U. S. Dept. Agr., Farmers' Bul. 507, 32 p., 11 fig.

- (190) JORDI, ERNST.
[1907]. Versuch zur Bekämpfung des Steinbrandes des Weizens. *In* Jahresber. Landw. Schule Rätti, 1906/07, p. 103-105.
- (191) KELLERMAN, W. A.
1890. Prevention of smut in cereals. *In* Agr. Sci., v. 4, no. 4, p. 99-101.
- (192) ———
1891. Second report on fungicides for stinking smut of wheat. *Kans. Agr. Exp. Sta. Bul.* 21, p. 47-72, pl. 1.
- (193) ———
1896. Ceres-pulver; Jensen's new fungicide for the treatment of smuts. (Abstract.) *In* Science, n. s. v. 4, no. 91, p. 438.
- (194) ———
[1896]. New experiments with fungicides for smut of wheat and oats. *In* Proc. 17th Ann. Meeting Soc. Prom. Agr. Sci., 1896, p. 60-70.
- (195) ——— and JENNINGS, O. E.
1902. Smut infection experiments. *In* Ohio Nat., v. 2, no. 6, p. 258-261, 1 fig.
- (196) ——— and SWINGLE, W. T.
1889. Preliminary report on smut in oats. *Kans. Agr. Exp. Sta. Bul.* 8, p. 91-104, 4 pl.
- (197) ———
1890. Preliminary experiments with fungicides for stinking smut of wheat. *Kans. Agr. Exp. Sta. Bul.* 12, p. 27-51, pl. 1.
- (198) KIESSLING, LUDWIG.
1918. Ueber schädliche Nebenwirkungen der Formalinbeizung des Saatgutes auf die Keimung. *In* Jour. Landw., Jahrg. 66, Heft 1, p. 7-51.
- (199) KILLER, J.
1918. Versuche über die Eignung des essigsauren Kupfers zur Bekämpfung des Steinbrandes. *In* Ztschr. Pflanzenkrank., Bd. 28, Heft 3/4, p. 106-109.
- (200)* KIRCH, JOHANN.
1783. Bewährte Methode den geschlossenen Brand im Waizen abzuhalten. *In* Bemerk. d. Kurpfälz, Phys.-Oek. Ges., 1783, p. 269-302. (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 1, p. 766, no. 13893. 1908.)
- (201) KIRCHNER, OSKAR.
1893. Ueber die Behandlung des Saatgetreides mit warmem Wasser als Mittel gegen den Flug- und Steinbrand. *In* Ztschr. Pflanzenkrank., Bd. 3, Heft 1, p. 2-15.
- (202)* ———
1903. Die Getreidefeinde, ihre Erkennung und Bekämpfung. Gemeinverständlich dargestellt. iv, 33 p., 4 fig., 2 tab. Stuttgart. E. Ulmer. 8°. (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 1, p. 767, no. 13911. 1908.)
- (203) ———
1903. Versuche zur Bekämpfung der Getriede-Brandkrankheiten. *In* Naturw. Ztschr. Land- u. Forstw., Jahrg. 1, Heft 12, p. 465-470.
- (204) ———
1906. Ueber die Empfänglichkeit verschiedener Weizensorten für die Steinbrand-Krankheit. *In* Fühlings Landw. Ztg., Jahrg. 55, Heft 23, p. 781-794.
- (205) ———
1906. Die Krankheiten und Beschädigungen unserer landwirtschaftlichen Kulturpflanzen. . . Aufl. 2, 675 p. Stuttgart.
- (206) ———
1908. Neue Beobachtungen über die Empfänglichkeit verschiedener Weizensorten für die Steinbrandkrankheit. *In* Fühlings Landw. Ztg., Jahrg. 57, Heft 5, p. 161-170.
- (207) ———
1913. Bericht über die Tätigkeit der K. Anstalt für Pflanzenschutz in Hohenheim im Jahre 1912. *In* Württemb. Wchnbl. Landw., Jahrg. 36, No. 29, p. 439-442; No. 30, p. 455-457.
- (208) ———
1916. Ueber die verschiedene Empfänglichkeit der Weizensorten für die Steinbrandkrankheit. *In* Ztschr. Pflanzenkrank., Bd. 26, Heft 1, p. 17-25.

KIRCHNER, OSKAR—Continued.

- (209) 1916. Untersuchungen über die Empfanglichkeit unserer Getreide für Brand- und Rostkrankheiten. *In* Fühlings Landw. Ztg., Jahrg. 65, Heft 1, p. 1-27; Heft 2, p. 41-72; Heft 3/4, p. 92-137. Literatur, p. 109-111.
- (210) ——— and BOLTSHAUSER, H.
1896. Atlas der Krankheiten und Beschädigungen unserer landwirtschaftlichen Kulturpflanzen. Ser. I, Getreidearten. 20 col. pl. with descriptive letterpress. Stuttgart.
- (211) KIRK, T. W.
1907. Diseases of field crops and forage plants. *In* New Zealand Dept. Agr. 15th Rpt. [1906]07, p. 159-163.
- (212) KITTLAUSS, K.
1899. Ueber die Einwirkung der Kupfervitriol-Beize auf die Keimkraft des Saatgetreides bei verschiedener Zeitdauer und Stärke der Lösung. *In* Fühlings Landw. Ztg., Jahrg. 48, Heft 15, p. 572-586; Heft 16, p. 605-616.
- (213) 1900. Mittel gegen Gerstenbrand und Schutz des Getreides gegen Stein- und Flugbrand. *In* Deut. Landw. Presse, Jahrg. 27, No. 100, p. 1200.
- (214) KLEBAHN, HEINRICH.
1893. Einige Versuche, betreffend den Einfluss der Behandlung des Saatguts gegen Brandpilze auf die Keimfähigkeit und den Ertrag des Getreides. *In* Ztschr. Pflanzenkrank., Bd. 3, Heft 2, p. 65-69.
- (215) KLEBS, GEORG.
1900. Zur Physiologie der Fortpflanzung einiger Pilze. III. Allgemeine Betrachtungen. *In* Jahrb. Wiss. Bot. [Pringsheim], Bd. 35, Heft 1, p. 80-203. Literatur-Verzeichniss, p. 199-203.
- (216)* KNEUTTINGER.
1861. Ueber die Verbreitung und Verhütung des Brandes der Getreidepflanzen. *In* Landw. Centbl. Deutschl. (Wilda), Jahrg. 9, Bd. 1, p. 350-359. (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 1, p. 778, no. 14100. 1908.)
- (217)* KÖCK, GUSTAV.
1905. Die wichtigsten Brandkrankheiten des Getreides und ihre Bekämpfung. *In* Mitt. K. K. Pflanzenschutzstat. Wien, 2. Flugbl. Oesterr. Landw. Wchnbl., 1905, No. 9, p. 72. (Not seen. Reference from Just's Bot. Jahresber., Jahrg. 33, 1905, Abt. 2, p. 218, no. 214. 1907.)
- (218)* 1907. Praktische Erfahrungen mit Formaldehyd als Getreidebrandbekämpfungsmittel. *In* Österr. Landw. Wchnbl. xxxiii, 1907, No. 13, p. 99. (Not seen. Abstract in Jahresber. Pflanzenkrank. [Hollrung], Bd. 10, 1907, p. 102, 115. 1909.)
- (219) KÖRNICKE, FRIEDRICH AUGUST.
1890. Ueber den Steinbrand des Weizens. *In* Verhandl. Naturhist. Ver. Preuss. Rheinlande, Jahrg. 47 (F. 5, Jahrg. 7), Sitzber., p. 92-93.
- (220) KÜHN, JULIUS GOTTHELF.
1859. Die Krankheiten der Kulturgewächse, ihre Ursachen und ihre Verhütung. Aufl. 2, xxii, 312 p., 7 pl. Berlin.
- (221) 1866. Einbeizen des Weizens. *In* Ztschr. Landw. Cent. Ver. Prov. Sach., Jahrg. 23, No. 4, p. 86-87.
- (222) 1873. Die Anwendung des Kupfervitrioles als Schutzmittel gegen den Steinbrand des Weizens. *In* Bot. Ztg., Jahrg. 31, No. 32, p. 502-505.
- (223) 1874. Mittheilungen über die Entwicklungsformen des Getreidebrandes und besprach eingehender die Art des Eindringens der Keimfäden in die Nährpflanze. *In* Ber. Naturf. Gesell. Halle, 1874, p. 5-7.
- (224) 1874. Ueber die Entwicklungsformen des Getreidebrandes und die Art des Eindringens der Keimfäden in die Nährpflanze. *In* Centbl. Agr. Chem., Bd. 5, Feb., p. 150-153.

KÜHN, JULIUS GOTTHELF—Continued.

- (225)* 1880. Beobachtungen über den Steinbrand des Weizens. Oesterr. Landw. Wehnbl., VI, 1880, No. 1 u. 2. (Not seen. Abstract in Bot. Centbl., Bd. 1, No. 6, p. 175-176, 1880.)
- (226)* KUZMA.
1878. Beobachtungen über den Weizensteinbrand. In Oesterr. Landw. Wehnbl., 1878, p. 511. (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 1, p. 813, no. 14667. 1908.)
- (227) LANG, HANS.
1908. Pilztötung mit dem Getreidetrockenapparat. In Illus. Landw. Ztg., Jahrg. 28, No. 70, p. 603-604.
- (228)* 1910. Das Beizen von Weizen und Dinkelsaatgut gegen Steinbrand. In Wehnbl. Landw. Ver. Baden, Karlsruhe 1910, p. 797-798. (Not seen. Reference from Internat. Cat. Sci. Lit., M, Bot., v. 10, 1910/11, p. 188. 1913.)
- (229) ——— and LANG, WILHELM.
1907. Das Beizen von Getreide-Elitekörnern. In Deut. Landw. Presse, Jahrg. 34, No. 20, p. 162-163.
- (230) LANG, WILHELM.
1912. De Getreideernte von 1911 und das Beizen. In Württemb. Wehnbl. Landw., Jahrg. 35, No. 15, p. 243-245.
- (231) 1913. Zum Parasitismus der Brandpilze. In Jahresber. Angew. Bot., Jahrg. 10, 1912, p. 172-180.
- (232)* 1917. Ueber die Beeinflussung der Wirtspflanze durch *Tilletia tritici*. In Ztschr. Pflanzenkrank., Bd. 27, Heft 2/3, p. 80-99. Zitierte Literatur, p. 99.
- (233)* LANGENBECK, ERNST.
1904. Die Pilzkrankungen der Getreidearten im Sommer 1903 in ihrem Zusammenhang mit abnormen Witterungserscheinungen. In Königsberger Land- u. Forstw. Ztg., 39, No. 51, 1903, p. 381-382. (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 1, p. 827, no. 14873a. 1908.)
- (234) LATTI, W. C.
1892. Field experiments with wheat. Ind. Agr. Exp. Sta. Bul. 41, p. 84-100.
- (235) LAUPERT, BRUNO.
1920. Einige Worte über Steinbrandbekämpfung. In Deut. Landw. Presse, Jahrg. 47, No. 62, p. 435-436.
- (236) LAWSON, JOHN.
1838. On the structure of smut-balls in wheat. In Quart. Jour. Agr. [Edinburgh], v. 9, no. 42, p. 269-272, illus.
- (237) LEHN, D., and VASTERS, J.
1913. Feinde und Krankheiten an Feldgewächsen. In Veröffentl. Landw.-Kammer Rheinprovinz, 1913, No. 1, p. 5-48.
- (238) LESAGE, PIERRE.
1912. Sur l'attitude de quelques semences soumises à l'action de solutions diverses de sulfate de cuivre. In Bul. Soc. Sci. et Méd. Ouest, t. 21, no. 3, p. 129-132.
- (239) LÉVEILLÉ, JOSEPH HENRI.
1851. Lettre sur la carie des grains, à M. Decaisne. In Rev. Hort., s. 3, t. 5, p. 385-391.
- (240)* LIEBENBERG, VON.
1879. Über die Dauer der Keimkraft der Sporen einiger Brandpilze. Österr. Landw. Wehnbl., III, 1879, No. 43, 44. (Not seen. Reference from Lindau and Sydow, Thesaurus Litt. Mycol., v. 1, p. 859, no. 15389. 1908.)

- (241)* LIEBENBERG, VON—Continued.
1880. Die schweflige Säure als Mittel gegen den Steinbrand des Weizens. *In Oesterr. Landw. Wehnbl.*, IV, 1880, p. 13. (Not seen. Reference from Lindau and Sydow. *Thesaurus Litt. Mycol.*, v. 1, p. 859, no. 15390. 1908.)
- (242)* 1880. Mittel gegen den Steinbrand des Weizens. *In Fühlings Landw. Ztg.*, 1880, p. 567. *Nach Oesterr. Landw. Wehnbl.* (Not seen. Reference from Just's *Bot. Jahresber.*, Jahrg. 9, 1881. Abt. 1. p. 260. 1884.)
- (243)* LONAY, H.
1911. Les blés belges (Belgian wheats). *In Jour. Soc. Roy. Agr. de l'Est de la Belgique*, 62. an., No. 30, p. 147, Liège, 24 juillet 1911. (Not seen. Abstract in *Internat. Inst. Agr. Bul. Bur. Agr. Intell. and Plant Diseases*, year 2, no. 7, p. 1631. 1911.)
- (244) LOVÉN, CHRISTIAN.
1884. Om sot på vårsäden och medlen däremot. *In K. Landtbr. Akad. Handl. och Tidskr.*, årg. 23, no. 5, p. 309-316.
- (245) LOVERDO, JEAN.
1892. Les maladies cryptogamiques des céréales. 312 p., illus. Paris.
- (246) LUSOR.
✓1844. Smut in wheat. *In Gard. Chron.*, 1844, no. 27, p. 452.
- (247) LUTMAN, BENJAMIN FRANKLIN.
1910. Some contributions to the life history and cytology of the smuts. *In Trans. Wis. Acad. Sci.*, v. 16, pt. 2, p. 1191-1244, pl. 88-95. Literature cited. p. 1225-1228.
- (248) M., P. D.
1776. Exposition des principales maladies des grains. *In Observ. Phys. Hist. Nat.*, t. 7, p. 435-437, pl. 1.
- (249) McALPINE, DANIEL.
1902. Experiments in the prevention of stinking smut or bunt of wheat. *In Jour. Dept. Agr. Victoria*, v. 1, pt. 4, p. 413-414.
- (250) 1903. Experiments in rust and stinking smut in wheat during 1901. (Abstract.) *In Rpt. 9th Meeting Aust. Assoc. Adv. Sci.*, 1902, p. 610-611.
- (251) 1906. Effect of formalin and bluestone on the germination of seed wheat. *In Agr. Gaz. N. S. Wales*, v. 17, pt. 5, p. 423-439. Also published as *Dept. Agr. So. Aust. Bul.* 12, 21 p. 1906.
- (252) 1910. The smuts of Australia, their structure, life history, treatment, and classification. 288 p., illus., 56 pl. Melbourne.
- (253) 1910. Rust and smut resistance in wheat and smut experiments with oats and maize. (Wheat improvement committee. II.) *In Jour. Dept. Agr. Victoria*, v. 8, pt. 5, p. 284-289.
- (254) MACKIE, W. W., and BRIGGS, FRED N.
1920. Fungicidal dusts for control of smut. *In Science*, n. s., v. 52, no. 1353, p. 540-541.
- (255) MADDOX, FRANK.
✓ 1896. Smut and bunt. *In Agr. Gaz. Tasmania*, v. 4, no. 6, p. 92-95. Reprinted in Maddox, Frank, Notes and results on agricultural experiments carried on under the auspices of the Council of Agriculture of Tasmania at Eastfield, Newnham. p. 72-84. Launceston, 1897.
- (256) ✓ 1910. Smut in wheat. *In Agr. Gaz. N. S. Wales*, v. 21, pt. 1, p. 58-59. Extracts from article by Maddox in *Agr. Gaz. Tasmania*, v. 4, no. 6, p. 92-95, with additional notes.
- (257) MAIRE, R.
1902. Sur la coexistence de la nielle et de la carie dans les grains de blé. *In Bul. Soc. Mycol. France*, t. 18, p. 130.

- (258) MALKOFF, KONSTANTIN.
1904-07. Versuche zur Bekämpfung des Steinbrandes des Weizens. In Jahresber. Staatl. Landw. Vers. Sta. Sadovo, Bulgarien. Jahrg. 1, 1903, p. 185-200; 2, 1904, p. 139-149; 3, 1905, p. 140-160 incl. 8 pl.; 4, 1906, p. 147-150. In Bulgarian. German résumé, Jahrg. 2, Sup. p. 12-14; 3, Sup. p. 9-11; 4, Sup. p. 5.
- (259) 1907. Untersuchungen über den Steinbrand des Weizens (*Tilletia laevis* Kühn). In Arb. Staatl. Landw. Vers. Sta. Sadovo, Bulgarien, No. 2, p. 1-23. In Bulgarian.
- (260) MALPEAUX, LÉOPOLD.
1916. Traitement des semences de céréales. In Jour. Agr. Prat., ann. 80 (n. s., t. 29), no. 5, p. 98-99.
- (261) MALZEW, A.
1908. Die Gewichtsbestimmung der Verunreinigung des Kornes durch der Brand. In Bul. Bur. Angew. Bot., Jahrg. 1, No. 5/6, p. 141-144. In Russian, German abstract, p. 145.
- (262)* MARAZZI, GIUSEPPE.
1788. Osservazioni sopra il carbonchio del formento. In Memoria Società Agraria Torino, vol. 2, p. 120. 1788. (Not seen. Reference from Ann. R. Accad. Agr. Torino, v. 28, 1885, Appendice, p. 146. 1886.)
- (263)* MARESCALCHI, A.
1899. Il carbone dei cereali. Bollett. di Entomol. agraria e Patol. veget. Padova. VI, 1899, p. 196-198. (Not seen. Reference from Just's Bot. Jahresber., Jahrg. 27, 1899, Abt. 2, p. 411. 1902.)
- (264) MARTINET, GUSTAVE.
1906. Traitement des blés de semence contre la carie et le charbon. In Chron. Agr. Vaud., ann. 19, no. 18, p. 477-480.
- (265) MASSEE, GEORGE EDWARD.
1891. British fungi. Phycomycetes and Ustilagineae. xv, 232 p., 3 pl. London. Literature quoted, p. 207-211.
- (266) ——— and MASSEE, IVY.
1913. Mildews, rusts and smuts. 229 p., col. front., 4 pl. London.
- (267) MAZÉ, PIERRE MARIE.
1916. Chlorose toxique du maïs: la sécrétion interne et la résistance naturelle des végétaux supérieurs aux intoxications et aux maladies parasitaires. In Compt. Rend. Soc. Biol. [Paris], t. 79, no. 19, p. 1059-1066.
- (268) MEYEN, FRANZ JULIUS FERDINAND.
1841. Pflanzen-Pathologie . . . xi, 330 p. Berlin. (Handbuch der Pflanzen-Pathologie und Pflanzen-Teratologie. Hrsg. von Chr. Gottfr. Nees von Esenbeck . . . Bd. I)
- (269) MEYER, C. F.
1753. Abhandlung von dem Brande im Getreide, und den Mitteln solchen zu wehren. In Hannoversche Anz., 1752, Gelehrte Anz., Stück 78, p. 1022-1026.
- (270) MICZYŃSKI, KÁZMÉR.
1911. Der Einfluss des Steinbrandes auf die Form der Weizenähren. In Ztschr. Landw. Versuchsw. Oesterr., Jahrg. 14, Heft 3, p. 232-234, illus.
- (271) MÖLLER, ALFRED.
1904. Neue Forschungsergebnisse über Brandkrankheiten des Getreides. In Ztschr. Forst- u. Jagdw., Jahrg. 36, Heft 4, p. 254-257.
- (272) MORITZ, JULIUS.
1902. Ueber die Einwirkung von Pictolin auf die Keimfähigkeit von Getreide. In Arb. K. Gsndhtsamt., Biol. Abt., Bd. 2, Heft 4, p. 512-513.

- (273) MORTENSEN, MORTEN LARSEN.
[1910.] Foreløbig meddelelse om forsøg anstillende af "De samvirkende danske landboforeninger's plantepatologiske forsøgsvirksomhed, [4] p. [n. p., n. d.]. (Abstract in Jahresber. Pflanzenkrank. [Holl-rung], Bd. 13, 1910, p. 130-131, 147. 1912.)
- (274) MÜLLER, H. C., and MOLZ, EMIL.
1913. Beizempfindlichkeit des Getreides der Ernte 1912 und Vorschläge zu dessen Beizung. *In* Deut. Landw. Presse, Jahrg. 40, No. 16, p. 190-192, fig. 211-212.
- (275) 1914. Über den Steinbrand des Weizens. *In* Fühlings Landw. Ztg., Jahrg. 63, Heft 6, p. 204-214.
- (276) 1914. Versuche zur Bekämpfung des Steinbrandes bei dem Winterweizen mittels des Formaldehyd-Verfahrens. *In* Fühlings Landw. Ztg., Jahrg. 63, Heft 24, p. 742-752.
- (277) 1918. Ergebnisse unserer letztjährigen Beizversuche mit Uspulun gegen den Steinbrand des Winterweizens. *In* Deut. Landw. Presse, Bd. 45, No. 71, p. 435.
- (278) 1918. Versuche mit Saatschutzmitteln. *In* Landw. Jahrb., Bd. 52, Heft 1, p. 67-130, pl. 5-6.
- (279) ——— and MORGENTHALER, OTTO.
1912. Beizempfindlichkeit des Getreides von der Ernte 1911. *In* Landw. Wchnschr. Sachsen, Jahrg. 14, No. 8, p. 58-59.
- (280) 1913. Über Brandbekämpfung und den Einfluss der Bestellzeit beim Sommerweizen auf dessen Ertrag und Gesundheit. *In* Landw. Vers. Sta., Bd. 83, Heft 3/4, p. 211-220.
- (281) ——— and MORGENTHALER, O.
1913. Versuche über die Bekämpfung des Steinbrandes bei Winterweizen. *In* Fühlings Landw. Ztg., Jahrg. 62, Heft 14, p. 481-487.
- (282) MÜLLER, L.
1912. Die Bekämpfung des Getreidebrandes. *In* Hess. Landw. Ztschr., Jahrg., 82, No. 34, p. 646-649, 7 fig.
- (283) MUNERATI, OTTAVI.
1911. La recettività del frumento per la carie in rapporto col tempo di semina. *In* Atti R. Accad. Lincei, Rend. Cl. Sci. Fis., Mat. e Nat., v. 20, sem. 1, fasc. 11, p. 835-840.
- (284) 1912. Sulla recettività del frumento per la carie in rapporto al tempo di semina. *In* Atti R. Accad. Lincei, Rend. Cl. Sci. Fis., Mat. e Nat., s. 5, v. 21, sem. 2, fasc. 12, p. 875-878.
- (285) ——— and HITIER, HENRI.
1912. Sur l'attaque du blé par la carie. Influence de l'époque de la semaille. *In* Jour. Agr. Prat., ann. 76 (n. s., t. 24), no. 42, p. 494-496.
- (286) NEGER, FRANZ WILHELM.
1898. Über Desinfektion von Saatgut mittels Formaldehyd-Dämpfe. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 1, Heft 11, p. 84-85.
- (287) NOBBE, FRIEDRICH.
1872. Ueber die Wirkungen des Maschinendrusches auf die Keimfähigkeit des Getreides. *In* Landw. Vers. Stat., Bd. 15, p. 252-275.
- (288) NORTON, J. B. S.
1896. A study of the Kansas Ustilagineae, especially with regard to their germination. *In* Trans. Acad. Sci. St. Louis, v. 7, no. 10, p. 229-241, pl. 25-29.
- (289) ØRSTED, ANDERS SANDØE.
1863. Om Sygdomme hos Planterne, som forårsages af Snyltesvampe, navnlig om Rust og Brand, og om Midlerne til deres Forebyggelse. 146 p., 3 col. pl. Kjøbenhavn.
- (290) OETKEN, W.
1912. Einige Beobachtungen über Steinbrand im Weizen. *In* Deut. Landw. Presse, Jahrg. 39, No. 70, p. 803-804.

(291)* OLSEN, ABRAHAM.

1791. Tilforladelig Underretning, hvorledes man paa bedste Maade skal behandle Hveden førend den saaes, at den skal blive fri for den skadelige Brand. Kjöbenhavn. (Not seen. Reference from Katalog K. Vet.- og Landb. Bibliotek [Denmark], 1894, p. 711. 1898.)

(292) OPITZ, KURT, and OBERSTEIN, OTTO.

1918. Neue Versuche zur Steinbrandbekämpfung mit Uspulun und Weizenfusariol. *In* Deut. Landw. Presse, Bd. 45, No. 86, p. 532.

(293) ORTH, A.

1881. [Über eine eigentümliche Einwirkung von *Tilletia caries* Tul. auf *Triticum spelta* L.] *In* Verhandl. Bot. Ver. Prov. Brandenb., Jahrg. 22, p. xvii-xviii.

(294) PAMMEL, LOUIS HERMANN. WEEMS, J. B., and SCRIBNER, FRANK LAMSON-.

1901. Fungus diseases of grasses. *In* Iowa Geol. Survey Bul. 1, p. 185-292, fig. 80-137, 3 col. pl.

(295) PEACOCK, ROBERT WILLIAM.

1902. Bunt experiments at Coolabah farm. *In* Agr. Gaz. N. S. Wales, v. 13, pt. 10, p. 1013-1014.

(296) 1905. Bunt preventives and their effects upon the germination of the grain. *In* Agr. Gaz. N. S. Wales, v. 16, pt. 3, p. 251-252.(297) 1906. Treatment of seed wheat—formalin. *In* Agr. Gaz. N. S. Wales, v. 17, pt. 9, p. 911-912.

(298)* PEGLION, VITTORIO.

1908. Intorno alla difesa del grano del carbone. *L'Agricoltore Ferrarese*, Ferrara, 13, 1908, N. 19. (Not seen. Reference from *Internat. Cat. Sci. Lit., M, Bot.*, v. 9, 1909/10, p. 219. 1911.)

(299) 1908. Contributo allo studio del carbone dei cereali. *In* Atti R. Accad. Econ. Agr. Georg. Firenze, s. 5, v. 5 (v. 86), p. 482-486, 2 fig.(300) 1908. Sulla immunità dei semi di frumento provenienti da piante colpito de infezione diffusa. *In* Atti Accad. Sci. Med. e Nat. Ferrara, ann. 82, p. 137-138.(301) 1910. Intorno alla carie del frumento. *In* Atti R. Accad. Lincei, Rend. Cl. Sci. Fis., Mat. e Nat., s. 5, v. 19, sem. 2, fasc. 4, p. 216-220.

(302) PHILIPPART, FRANÇOIS AKEN.

1837. Traité organographique et physiologico-agricole sur la carie, le charbon, l'ergot, la rouille et autres maladies du même genre qui ravagent les céréales. 220 p., 9 pl. Versailles.

(303) PIEPER, H.

1920. Steinbrand und physiologische Spaltungen. *In* Deut. Landw. Presse, Jahrg. 47, No. 79, p. 543-544.

(304) PIPER, CHARLES V.

1894. Common fungus diseases and methods of prevention. Dodder. Wash. Agr. Exp. Sta. Bul. 8, p. 131-144, 2 fig.

(305)* PIRET, E.

1898. Le carie des céréales et le chaulage et le sulfatage de grains de semences dans la province de Namur. *Agronome* 1848 [1898] Nr. 40. (Not seen. Reference from Jahresber. Pflanzenkrank. [Hollrung], Bd. 2, 1899, p. 228. 1900. Also in *Exp. Sta. Rec.*, v. 10, p. 764. 1899.)

(306) PLOWRIGHT, CHARLES BAGGE.

1889. A monograph of the British Uredineae and Ustilagineae . . . 347 p., illus., 8 pl. London. "Authors quoted," p. 309-315.

(307) 1892. The diseases of the reproductive organs of plants caused by parasitic fungi. *In* Gard. Chron. s. 3, v. 11, p. 301-302, 331-332, 373, 397, 472-473, 490-491, 524-525.(308) [1899.] On the recent additions to our knowledge of the Uredineae and Ustilagineae with special reference to the British species. *In* Brit. Mycol. Soc. Trans. 1898/99, p. 90-105.

- (309) **PLUCHE**, [NOËL ANTOINE].
1740. Part of a letter from the Abbé Pluche to Dr. Mortimer, Sec. R. S. concerning the smut of corn. Translated from the French. *In* Phil. Trans. [Roy. Soc. London], v. 41, pt. 1, no. 456, p. 357-358.
- (310) **POGGI**, TITO.
1899. Alcuni mali del frumento . . . ed. 3, 72 p. Casale. (On cover: Biblioteca agraria Ottavi, v. 12.)
- (311)* **POSPÉLOV**, V.
1905. Die neuesten Beiträge zur Frage über die Bekämpfung des Brandes. Védom. Sel'sk. Choz. i Promysl. Kiev, 9, 1905, p. 268-270, 287-289. *In* Russian. (Not seen. Reference from Internat. Cat. Sci. Lit., M, Bot., v. 6, 1906/07, p. 240. 1908.)
- (312) **PRÉVOST**, ISAAC BÉNÉDICT.
1807. Mémoire sur la cause immédiate de la carie ou charbon des blés et de plusieurs autres maladies des plantes. 80 p., 3 pl. Paris.
- (313)* **PRIANISHNIKOV**, D. N.
1902. [Experiments for smut prevention.] *In* Khozyain. 1902, No. 20, p. 647-652. *In* Russian. (Not seen. English abstract in Exp. Sta. Rec., v. 15, no. 3, p. 267. 1903. Russian abstract in Zhur. Opuitn. Agron. (Jour. Exp. Landw.), god 4, kniga 1, p. 112. 1903.)
- (314) **PUSCH**, G.
1912. Über die Schädlichkeit der Tilletia im Futter unserer Haustiere. *In* Ztschr. Infektionskrank. Par. Krank. u. Hyg. Haust., Bd. 11, Heft 1, p. 1-14.
- (315) **PYE**, H.
1909. Diseases and pests of cereals. (Wheat improvement committee.) *In* Jour. Dept. Agr. Victoria, v. 7, pt. 6, p. 368-373.
- (316) **QUANJER**, HENDRIK MARIUS.
1914. Ontsmetting van zaaigranen met heet water. Tweede, herziene druk. 18 p., pl. 's Gravenhage. (At head of title: Directie van den Landbouw.)
- (317) ——— and **BOTJES**, J. OORTWIJN.
1915. Nederlandsche onderzoekingen over de bestrijding van graanen grasbrand en van strepenziekte. *In* Meded. R. Hoog. Land-, Tuin- en Boschbouwsch., deel 8, afl. 3, p. 129-156, pl. 3-5.
- (318) **REED**, GEORGE MATTHEW.
1912. The smuts of cereals. *In* 44th Ann. Rpt. Mo. State Bd. Agr., 1911, p. 253-265, illus.
- (319) ——— **MUNDY**, EMMA B., and **GIBBS**, N. M.
1916. Grain smut investigation and control. *In* Mo. Agr. Exp. Sta. Bul. 141, p. 26.
- (320) **REED**, JOSEPH.
1903. Treatment of stinking smut in wheat. Colo. Agr. Exp. Sta. Bul. 79, 8 p.
- (321) **REMNANT**, RICHARD.
1637. A discourse or historie of bees . . . Whereunto is added the causes, and cure of blasted wheat. And some remedies for blasted hops, and rie, and fruit. Together with the causes of smutty wheat; all of which are very usefull for this later age. 47 p. London. (Wheat, p. 40-47.)
- (322) **REMY**, THEODOR, and **VASTERS**, J.
1914. Beobachtungen über Chlorphenol-Quecksilber als Pflanzenschutzmittel. *In* Illus. Landw. Ztg., Jahrg. 34, No. 91, p. 769-770; No. 92, p. 776-778, illus.
- (323) **REYNOLDS**, M. H.
1913. Prevention of bunt in wheat. *In* Agr. Gaz. N. S. Wales, v. 24, pt. 6, p. 461-476.
- (324) **RICHARDSON**, A. E. V.
1910. Bunt tests, 1909. *In* Jour. Dept. Agr. So. Aust., v. 13, no. 6, p. 491-494.

- (325) RIEHM, EDUARD.
1911-15. Getreidekrankheiten und Getreideschädlinge. Eine Zusammenstellung der wichtigeren, im Jahre 1910, 1911, 1912, 1913, 1914 veröffentlichten Arbeiten. In *Centbl. Bakt. etc.*, Abt. 2, Bd. 30, p. 465-492, 1911; Bd. 34, p. 434-472, 1912; Bd. 39, p. 81-107, 1913; Bd. 43, p. 177-218, 1915; Bd. 44, p. 385-407, 1915. Literatur, Bd. 34, p. 466-472; Bd. 39, p. 103-107; Bd. 43, p. 210-218; Bd. 44, p. 402-407.
- (326) 1913. Prüfung einiger Mittel zur Bekämpfung des Steinbrandes. In *Mitt. K. Biol. Anst. Land- u. Forstw.*, Heft 14, p. 8-9.
- (327) 1913. Ueber Apparate zur Brandbekämpfung. In *Deut. Landw. Presse*, Jahrg. 40, No. 10, p. 107-108.
- (328) 1914. Prüfung einiger neuerer Beizmittel. In *Mitt. K. Biol. Anst. Land- u. Forstw.*, Heft 15, p. 7-8.
- (329) 1915. Beizversuche zur Bekämpfung einiger Getreidekrankheiten. In *Illus. Landw. Ztg.*, Jahrg. 35, No. 24, p. 161-162.
- (330) RITTER, J. H.
1756. Nützliche Anmerkungen über das communicirte Schreiben des Herrn Aucante, darinnen die Zubereitung des Getreydes mit Kalk und Arsenik als höchst schädlich verworfen, und im 75ten Stücke dieser nützlichen Samlungen dem Publico mitgetheilet worden. In *Hannoversche Mag.* 1755, Nützl. Samml., Stück 96, p. 1534-1535.
- (331) RUDOLPHI, FR.
1829. Plantarum vel novarum vel minus cognitarum descriptiones. In *Linnaea*, Bd. 4, Quartalheft 1, p. 114-120.
- (332) SAUNDERS, WILLIAM, BEDFORD, S. A., and MACKAY, ANGUS.
1894. Smut in wheat. In *Canada Exp. Farms Rpts.*, 1893, p. 41-42.
- (333)* SAVASTANO, LUIGI.
1881. Malattie delle Graminacee. La carie del frumento. *L'Agricolt. Meridionale Portici*, IV, 1881, No. 1, 2, 4. (Not seen. Reference from *Just's Bot. Jahresber.*, Jahrg. 10, 1882, Abt. 1, p. 141. 1884.)
- (334) SCHELLENBERG, HANS CONRAD.
1911. Die Brandpilze der Schweiz. xlv, 180 p., illus. Bern. (Beiträge zur Kryptogamenflora der Schweiz, Bd. 3, Heft 2.) Literaturverzeichnis, p. 164-171.
- (335) SCHEUNERT, ARTHUR, and LÖTSCH, ERNST.
1911. Fütterungsversuche mit *Tilletia*. Ein Beitrag zur Hygiene der Ernährung. In *Ztschr. Infektionskrank. Par. Krank. u. Hyg. Haust.*, Bd. 9, Heft 3/4, p. 177-187.
- (336) SCHINDLER, FRANZ.
1880. Ueber den Einfluss verschiedener Temperaturen auf die Keimfähigkeit der Steinbrandsporen. In *Forsch. Geb. Agr. Phys.*, Bd. 3, p. 288-293.
- (337) SCHMITZ, N.
1910. Wheat—variety tests and diseases. *Md. Agr. Exp. Sta. Bul.* 147; p. 33-45.
- (338)* SCHÖLER, N. P.
1807. Om at opdage Brand i Hvede. In *Ökonomske Annaler*, X, 1807, p. 346. (Not seen. Reference from *Lindau and Sydow, Thesaurus Litt. Mycol.*, v. 2, p. 478, no. 24123. 1909.)
- (339) SCHREBER, D.
1758. Prüfung der Meynung vom Brande im Getreide, welche Herr Tillet in einer französischen Preisschrift vorgetragen hat. In *Hannoversche Anz.* 1757, Nützliche Saml., Stück 25, p. 386-399.
- (340) SCHRIBAU, ÉMILE.
1892. Le germinateur quarante et le sulfatage des céréales. In *Jour. Agr. Prat.*, ann. 56, t. 2, p. 454-458.

- (341) SCHROETER, JOSEPH.
1877. Bemerkungen und Beobachtungen über einige Ustilagineen. *In* Beitr. Biol. Pflanz., Bd. 2, Heft 3, p. 349-383, pl. 12.
- (342) SCHULTHESS, HEINRICH.
1761. Vorschlag einiger durch die Erfahrung bewährter Hilfsmittel gegen den Brand im Korn. *In* Abhandl. Naturf. Gesell. Zürich, Bd. 1, p. 498-506.
- (343) SELBY, AUGUSTINE DAWSON.
1898. Some diseases of wheat and oats. Ohio Agr. Exp. Sta. Bul. 97, p. 31-61, 4 fig.
- (344) SESSOUS.
1920. Beobachtungen über Steinbrandbefall. *In* Deut. Landw. Presse, Jahrg. 47, No. 81, p. 556.
- (345) SORAUER, PAUL.
1874. Handbuch der Pflanzenkrankheiten . . . 406 p., illus., 16 pl. (partly col.) Berlin.
- (346) 1893. Populäre Anleitung für den Landwirt zur Unterscheidung der im Getreide vorkommenden Stein- und Staubbbrandarten. *In* Ztschr. Pflanzenkrank., Bd. 3, p. 271-277, pl. 5.
- (347) 1895. Die Schutzbehandlung des Getreides gegen die Brandkrankheiten. *In* Ztschr. Pflanzenkrank., Bd. 5, Heft 3, p. 187-190.
- (348) SOUTTER, R. E.
1912. Experiments with smut preventives. *In* Queensland Agr. Jour., v. 28, pt. 1, p. 1-5.
- (349) 1913. Smut experiments at the state farm, Bungeworgorai. *In* Queensland Agr. Jour., v. 30, pt. 3, p. 162-163.
- (350) STAES, G.
1898. De invloed van het gebruik van molenstof op den brand der graangewassen. *In* Tijdschr. Plantenziekten, jaarg. 4, afl. 3, p. 72-77.
- (351) 1898. Noordamerikaansche middelen tot het voorkomen van den brand der graangewassen. *In* Tijdschr. Plantenziekten, jaarg. 4, afl. 3, p. 78-83.
- (352) 1899. Een onderzoek over den stink- of steenbrand der tarwe in België in 1898. *In* Tijdschr. Plantenziekten, jaarg. 5, afl. 5/6, p. 170-176.
- (353) 1902. De brand der graangewassen. *In* Tijdschr. Plantenziekten, jaarg. 8, afl. 4/5, p. 122-141, 2 fig.
- (354) STAKMAN, E. C.
1913. Spore germination of cereal smuts. Minn. Agr. Exp. Sta. Bul. 133, 52 p., 25 pl. Bibliography, p. 46-48.
- (355) STEELE, W. EDWARD.
1850. Chloride of lime a prevention for smut in wheat. *In* Gard. Chron., 1850, no. 41, p. 651.
- (356) STEGLICH, B.
1909-11. Die Uebertragung des Weizensteinbrandes auf den Pflanzenbestand der Weizenfelder durch infizierten Stalldünger, Samen und Ackerboden. *In* Fühlings Landw. Ztg., Jahrg. 58, Heft 20, p. 738-742, 1909: 59, Heft 5, p. 162-164, 1910: 60, Heft 2, p. 54-55, 1911.
- (357) STÖRMER, KURT.
1904. Bericht über die Ergebnisse von Versuchen zur Bekämpfung des Weizensteinbrandes. *In* Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 2, Heft 11, p. 141-146; Heft 12, p. 149-152.
- (358) 1911. Ueber die Bekämpfung des Steinbrandes beim Winterweizen. *In* Deut. Landw. Presse, Jahrg. 38, No. 80, p. 917-918; No. 81, p. 929.
- (359) STURLER, J. L.
1763. Von den Mitteln dem Brande des Getreides (Carie) durch eine Lauge Vorzubiegen. *In* Abhandl. u. Beobacht. Oekonom. Gesell. Bern, Jahrg. 4, Stük 1, p. 209-216.

- (360) SUTTON, GEORGE L.
1910. Experiments with fungicides for the prevention of "stinking smut" or "bunt" in wheat. [Preliminary note.] *In Agr. Gaz. N. S. Wales*, v. 21, pt. 4, p. 289-291.
- (361) ——— and DOWNING, R. G.
1910. Some experiments with fungicides used for the prevention of "stinking smut" (bunt), Cowra. 1909. *In Agr. Gaz. N. S. Wales*, v. 21, pt. 5, p. 382-397.
- (362) ——— and PRIDHAM, J. T.
1907. The effects of some fungicides recommended for the prevention of "stinking smut" (bunt) on the germination of wheat seeds. *In Agr. Gaz. N. S. Wales*, v. 18, pt. 3, p. 235-253, illus.
- (363) SWINGLE, W. T.
1892. Treatment of smut of oats and wheat. U. S. Dept. Agr., Farmers' Bul. 5, 8 p., pl. 1.
- (364) ———
1895. The grain smuts: Their causes and prevention. *In U. S. Dept. Agr. Yearbook 1894*, p. 409-420, fig. 92-99.
- (365) ———
1898. The grain smuts: How they are caused and how to prevent them. U. S. Dept. Agr., Farmers' Bul. 75, 19 p., 8 fig.
- (366) ———
1906. The prevention of stinking smut of wheat and loose smut of oats. U. S. Dept. Agr., Farmers' Bul. 250, 16 p., 5 fig.
- (367) TAYLOR, SAMUEL.
1846. Smut in wheat. *In Gard. Chron.*, 1846, no. 15, p. 242-243.
- (368) TESSIER, ALEXANDRE HENRI.
1783. *Traité des maladies des grains*. 351 p., 7 fold. pl. Paris.
- (369)* 1786. *Moyens éprouvés pour préserver les fromens de la carie*. 15 p. Avignon. (Not seen. Reference from Pritzel, *Thesaurus Lit. Bot.*, p. 291. 1851.)
- (370)* TILLET, MATHIEU.
1755. *Dissertation sur la cause qui corrompt et noircit les grains de bled dans les épis, et sur les moyens de prévenir ces accidents*. 150 p. Bordeaux. (Not seen. Reference from Pritzel, *Thesaurus Lit. Bot.*, p. 296. 1851.)
- (371) TSCHARNER, NIKLAUS EMANUEL.
1764. Von dem Brand und von dem Rost im Getreide. *In Abhandl. u. Beobacht. Oekonom. Gesell. Bern*, Jahrg. 5, Stük 2, p. 27-31.
- (372) TUBEUF, CARL VON.
1901. Studien über die Brandkrankheiten des Getreides und ihre Bekämpfung. *In Arb. K. Gsndtsamt., Biol. Abt.*, Bd. 2, Heft 2, p. 179-372, 19 fig., 1 col. pl.
- (373) ———
1901. Vorschläge zur Bekämpfung des Weizen-Steinbrandes. *In Mitt. Deut. Landw. Gesell.*, Jahrg. 16, Stück 34, p. 201-202.
- (374) ———
1902. Weitere Beiträge zur Kenntnis der Brandkrankheiten des Getreides und ihrer Bekämpfung. *In Arb. K. Gsndtsamt., Biol. Abt.*, Bd. 2, Heft 3, p. 437-467, illus.
- (375)* 1910. *Pflanzenpathologische Wandtafeln*. Tafel 7-8: Brandkrankheiten des Getreides. 51 p., 2 col. pl. Stuttgart. (Not seen. Reference from *Internat. Cat. Sci. Lit., M, Bot.*, v. 10, 1910/11, p. 327. 1913.)
- (376) ———
1910. Die Uebertragung des Weizensteinbrandes auf den Pflanzenbestand der Weizenfelder durch infizierten Stalldünger, Samen und Ackerboden. *In Fühlings Landw. Ztg.*, Jahrg. 59, Heft 5, p. 161-162. (Criticism of article by Steglich, no. 356.)
- (377) TULASNE, LOUIS RÉNE, and TULASNE, CHARLES.
1847-54. *Mémoire sur les Ustilaginées comparées aux Urédinées*. *In Ann. Sci. Nat. Bot.*, s. 3. t. 7. p. 12-127. pl. 2-7. 1847; s. 4. t. 2, p. 77-196, pl. 7-12, 1854.

- (378) TULL, JETHRO.
1733. Horse-hoeing husbandry: or, an essay on the principles of tillage and vegetation . . . x, 269 p., 6 pl. London.
- (379) UNGER, F. J. A. N.
1833. Die Exantheme der Pflanzen . . . 422 p., 7 col. pl. Wien.
- (380)* 1836. Ueber den Einfluss des Bodens auf die Vertheilung der Gewächse, nachgewiesen in der Vegetation des nordöstlichen Tirols. xxiv, 367 p., 9 maps. Wien. (Not seen. Reference from Pritzel, Thesaurus Lit. Bot., p. 304. 1851.)
- (381) 1840. Beiträge zur vergleichenden Pathologie. vi, 42 p., 1 col. pl. Wien.
- (382) VAVILOV, N. I.
1913. Beiträge zur Frage über die verschiedene Widerstandsfähigkeit der Getreide gegen parasitische Pilze. In Arb. Vers. Stat. Pflanzenzücht. Moskauer Landw. Inst., F. 1, p. 1-110, 3 col. pl. In Russian. German résumé, p. 90-108.
- (383)* VOLKART, ALBERT.
1906. Krankheiten und Schädlinge des Getreides und ihre Bekämpfung. Mitt. Gesell. Schweiz. Landwirte. 1906 (II, 1905, p. 3-20). (Not seen. Reference from Just's Bot. Jahresber., Jahrg. 35, 1907, Abt. 2, p. 494. 1910. Reference in curves from Lindau and Sydow, Thesaurus Litt. Mycol., v. 2, p. 704, no. 27974. 1909.)
- (384) 1906 Die Bekämpfung des Steinbrandes des Weizens und des Kornes. In Landw. Jahrb. Schweiz. Jahrg. 20. Heft 8, p. 445-490, 3 fig.
- (385) WAGNER, R. J.
1910 Wassertoffionenkonzentration und natürliche Immunität der Pflanzen. In Centbl. Bakt. [etc.], Abt. 2, Bd. 44, No. 24/25, p. 708-719.
- (386)* WAHL, BRUNO.
1907. Die wichtigsten Krankheiten und Beschädigungen unseres Getreides. Landes-Amtsblatt d. Erzherzogtums Österreich unter der Enns, 1907, Nos. 3, 5, 6, 7, 8. 48 p., fig. (Not seen. Reference from Just's Bot. Jahresber., Jahrg. 35, 1907, Abt. 1, p. 513. 1909.)
- (387)* WAHRlich, W. K.
1897. [Krankheiten des Getreides.] iv, 37 p., 19 fig. St. Petersburg. (Russian original not seen. Abstract in Russian in Selsk. Khoz. i Lycsov., t. 187, no. 10, p. 449. 1897. German title from Lindau and Sydow, Thesaurus. Litt. Mycol., v. 3, p. 668. 1913.)
- (388) WALLDÉN, J. N.
1912. Höstvetets betning mot brand. In Sveriges Utsädesför. Tidskr., årg. 22, häfte 4, p. 242-252. (Abstract in Bot. Centbl. v. 122, p. 149. 1913.)
- (389) 1916. Tröskskada å hvete och råg samt dess inflytande på känsligheten för betning och lagring. (Threshing injuries of wheat and rye and their influence on the germination and keeping qualities of the grain.) In Sveriges Utsädesför. Tidskr., årg. 26, häfte 1, p. 24-47, 1 fig., 1 pl.
- (390) WARD, HARRY MARSHALL.
1889. Smut-fungi. In Gard. Chron., s. 3, v. 5, no. 113, p. 233-235; no. 114, p. 267-268.
- (391) WECK, L.
1916. Untersuchungen über Uspulun als Beizmittel. In Illus. Landw. Ztg., Jahrg. 36, No. 82, p. 552.
- (392) WEISS, J. E.
1900. Die rationelle Bekämpfung der Getreidebrandarten. In Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 3, Heft 9, p. 68-69, illus.
- (393) 1901. Das richtige Beizen des Saatgutes gegen die Brandpilze, besonders gegen den Steinbrand des Weizens. In Prakt. Bl. Pflanzenbau u. Schutz, Jahrg. 4, Heft 9, p. 65-67.

- (394) WHEELER, W. A.
1904. Preliminary experiments with vapor treatments for the prevention of the stinking smut of wheat. S. Dak. Agr. Exp. Sta. Bul. 89, 19 p.
- (395) WILCOX, EDWIN MEAD.
1912. Smuts of Nebraska cereals. Nebr. Agr. Exp. Sta. Bul. 131 (v. 25, art. 1), 16 p., 13 fig.
- (396) WINDISCH, RICHARD.
1897. Ueber die Einwirkung des Formaldehyds auf die Keimung. In Landw. Vers. Sta., Bd. 49, Heft 3, p. 223-226.
- (397) WINTER, GEORG.
1884. Die Pilze Deutschlands, Oesterreichs und der Schweiz. Abt. 1. Leipzig. In Rabenhorst, Ludwig. Kryptogamen-Flora, Aufl. 2, Bd. 1, Abt. 1.
- (398) WOLFF, REINHOLD.
1873. Beitrag zur Kenntniss der Ustilagineen. In Bot. Ztg., Jahrg. 31, No. 42, p. 657-661; No. 43, p. 673-677; No. 44, p. 689-694, pl. 7.
- (399) 1873. Der Brand des Getreides . . . 37 p. Halle. Inaug.-Diss.
- (400) ZADE, ADOLF.
1920. Ein neues Verfahren zur Bekämpfung des Weizensteinbrandes. In Deut. Landw. Presse, Jahrg. 47, No. 27, p. 204-205; No. 28/29, p. 210-211.
- (401) ZAVITZ, C. A.
1914. Loose smut in oats, and stinking smut in wheat. In 39th Ann. Rpt. Ontario Agr. Col. and Exp. Farm, 1913, p. 132-134.
- (402) ZELLNER, JULIUS.
1910. Chemistry of the higher fungi. VI. Relations of the higher parasitic fungi and their substrata. (Abstract.) In Jour. Chem. Soc. [London], v. 98, pt. 2, p. 886-887.
- 403) 1910. Zur Chemie der höheren Pilze. VI. Mitteilung: Chemische Beziehungen zwischen höheren parasitischen Pilzen und ihrem Substrat. In Monatsh. Chem., Bd. 31, Heft 6, p. 635-641.
- (404) 1911. Zur Chemie der höheren Pilze. VIII. Mitteilung: Ueber den Weizenbrand (*Tilletia levis* Kühn und *tritici* Winter). In Monatsh. Chem., Bd. 32, Heft 10, p. 1065-1074.

ADDITIONAL COPIES

OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.

AT
10 CENTS PER COPY