2 1/2

3 1/2

Digitized by the Internet Archive
in 2015

<https://archive.org/details/manualofbritishv11jeny>

Cardley Hall.

A
M A N U A L
OF
BRITISH VERTEBRATE ANIMALS:
OR
DESCRIPTIONS
OF
ALL THE ANIMALS BELONGING TO THE CLASSES,
MAMMALIA, AVES, REPTILIA, AMPHIBIA,
AND PISCES,
WHICH HAVE BEEN
HITHERTO OBSERVED IN THE BRITISH ISLANDS:
INCLUDING THE
DOMESTICATED, NATURALIZED, AND EXTIRPATED SPECIES:
THE WHOLE SYSTEMATICALLY ARRANGED.

BY THE
REV. LEONARD JENYNS, M.A.

FELLOW OF THE LINNEAN, ZOOLOGICAL, AND ENTOMOLOGICAL SOCIETIES
OF LONDON; AND OF THE CAMBRIDGE PHILOSOPHICAL SOCIETY.

CAMBRIDGE:

PRINTED AT THE PITT PRESS, BY JOHN SMITH,
PRINTER TO THE UNIVERSITY.

SOLD BY J. & J. J. DEIGHTON; AND T. STEVENSON, CAMBRIDGE;
AND LONGMAN & CO., LONDON.

M.DCCC.XXXV.

E
#23136
v. 1
sp coll

PREFACE.

THE present Work contains descriptions of all the Vertebrate Animals, including the domesticated, naturalized, and extirpated species, which have been hitherto observed in the British Islands. The object of its author is to present Naturalists with a Manual in this department of our Fauna, adapted to the existing state of our knowledge, and such as shall be calculated to meet the wants of the Science in that advanced stage to which it has attained since the publication of former works of this nature. In furtherance of this end two points appeared necessary to be attended to. One was to ascertain, as far as practicable, the additions which had been made of late years to our lists of British Animals, to inquire into the respective claims of those which had been admitted into these lists previously, and carefully to distinguish between such species as have unquestionably occurred within the limits of our own Islands, or in the adjoining seas, and others of which reasonable doubts might be entertained in reference to this matter. The other important point, as it appeared to the author, was to take care that the descriptions should as far as possible be obtained from the animals themselves, and nothing inserted upon the credit of other writers which was capable of being verified by personal examination. The day is for ever gone by in which mere compilations will be thought to be of any service to the

science of Zoology. So far from advancing its progress, it may be said unhesitatingly that they tend only to retard it. It is through such channels that errors already of long standing become more widely circulated, at the same time that new ones, to a greater or less extent, are infallibly introduced. The author who is too indolent to examine and describe for himself, has often spared himself the trouble of even investigating the nature of the materials which he has obtained from others. The consequence is, that he has perhaps blended together the descriptions of two or more perfectly distinct species, or out of one made several, or, led away by the identity of mere names, has transferred to our native animals the descriptions of exotic, nearly allied species, with which he has confounded them.

After the above enunciation of the two leading points which have been borne in mind whilst preparing this work for publication, the author hopes that it will not be thought uncalled for by those for whose use and guidance it is principally intended. The latest previous work upon the same subject and conducted upon the same general plan is the "History of British Animals" by Dr. Fleming. This was completed in 1827 and published the year following, since which period a great variety of species have been added to the Fauna of this country, more particularly in the Class of Fish, though many occur in the Classes of Quadrupeds and Birds. As these additions have been already indicated in the "Systematic Catalogue" lately published by the author of the present work, it is unnecessary to dwell upon them individually. It may, however, afford interest to present a comparative view of the aggregate numbers of species in each Class, as they appear in the "British Animals" of Dr. Fleming and in the Manual now offered to the public.

The total number of Mammalia noticed by that Naturalist, excluding the domesticated and naturalized species, as well as all those not met with at the present day, is fifty-three, from which deducting three as of rather doubtful character, there remain fifty. The number of Birds, excluding in like manner the domesticated and naturalized kinds, but including Stragglers, which Dr. Fleming has only briefly alluded to and not allowed to form part of the regular numbering, amounts to two hundred and eighty-one, from which deducting seventeen as either decidedly not British or of doubtful character, there remain two hundred and sixty-four. The number of Reptiles and Amphibious Animals, which are united in one Class, is fourteen, from which deducting one as a doubtful native, and two others as very doubtful species, there remain eleven. The number of indigenous Fish amounts to one hundred and seventy, from which deducting eleven not distinct from others, one not British, and three of doubtful character, there remain one hundred and fifty-five, which added to about seven noticed only as stragglers*, gives one hundred and sixty-two. Hence it appears that the total number of truly British *Vertebrata*, restricting the expression in the manner above indicated, described or mentioned by Dr. Fleming, amounts to only four hundred and eighty-seven. In the present work, excluding in a similar way the domesticated, naturalized, and extirpated species, as well as all those of doubtful character, the number of described Mammalia amounts to sixty-one; that of Birds to two hundred and ninety-seven; that of Reptiles and Amphibious Animals together to thirteen;

* Dr. Fleming has alluded briefly to several other species which have been included by some authors in the British Fauna, but the claims of these to be considered as true natives being extremely doubtful, they are not here taken into the account.

that of Fish to two hundred and ten. These added together give five hundred and eighty-one as the total number, exceeding that above stated by ninety-four. But independently of these additions which have been made of late years to our native Animals, Dr. Fleming's work professing to be in a great measure only a compilation, it was thought that one which partook more of an original character, would not prove unacceptable at the present day. It must not be forgotten that two other works connected with the Vertebrate Division of the British Fauna have appeared since the one just spoken of. I allude to the excellent "Illustrations of British Ornithology" by Mr. Selby, published in 1833, in which the number of Birds is raised to two hundred and eighty*, and the valuable "British Fishes" of Mr. Yarrell now in course of publication. But not to dwell upon the circumstance, that these works are upon totally different plans from that of the present one, the author conceives that a single volume comprising all the Vertebrate Animals which have been observed hitherto in these Islands will form a useful addition to the library of the British Naturalist, as well as prove a convenient travelling companion.

A few remarks may now be made upon the particular plan which is adopted in the present work. And first, it must be stated why a deviation has been made from Dr. Fleming's method of separating from the other species, and partially excluding from the British Fauna, those animals which have occurred as Stragglers only in a few rare instances. The reason is founded in the impossibility of drawing any marked line where such separation shall com-

* The exact number of species given by Mr. Selby is two hundred and eighty-seven, but six, if not seven, of these cannot be considered otherwise than as doubtful natives.

mence. However just it may be in general terms to speak of the animals of any country under the three heads of Residents, Periodical Visitants, and Irregular Visitants or Stragglers, it must be obvious to every one that between the last two there exists so little boundary, that there are many species of which it is difficult to say whether they belong to the former, or to the latter of these divisions*. Every intermediate degree of frequency, as regards the visits of different species of Birds to this country, may be found between what constitutes an annual migration and a solitary appearance. There are some which occur most years in greater or less numbers, though not with that unerring punctuality which attends the movements of our more regular migrants; there are others which only shew themselves at uncertain and more distant intervals; there are others, again, which have not been observed in more than a few instances; and of these last a series might easily be made out in which such instances became continually fewer, till the number was reduced ultimately to one. Now under such circumstances it is clearly impossible to draw the line which Dr. Fleming has attempted to point out. For this reason, in the present work, the species above alluded to are noticed exactly in the same manner as those which are permanently resident in this country, or which visit it at fixed intervals. In so doing there is no wish to raise them to a higher rank in the Fauna than they really deserve. They are still considered only as Stragglers to a greater or less extent. But if it be the object of a Fauna to present naturalists with an account

* Dr. Fleming himself must have experienced some difficulty in determining what species were to be "degraded to the rank of Stragglers," judging from the circumstance, that some which he has only briefly noticed as of this character, have in fact occurred in this country oftener than, many quite as often as, others described at length in the body of his work.

of such animals as are to be met with in their own country, it must surely include all those which have been known to occur in it hitherto. Neither is the reason obvious why the characters, or even a detailed description, of such species, should be suppressed. If it be quite certain that they have appeared in one instance, it cannot be deemed improbable, clearly not impossible, that they may occur in other instances*, and should this prove to be the case, it is very desirable that the student should have the means of identifying them.

A line of quite as much importance, in the opinion of the author, as that on which Dr. Fleming has insisted, and one more easily drawn, though never yet attempted to be drawn with accuracy†, is that between species, the occurrence of which in this country at one time or another there is no ground for questioning, and others whose claims to be considered as British have not yet been made substantially good. In distinguishing between these two classes, the author has generally been guided, at least in the case of Birds, by the fact of the existence or not of British-killed specimens in any known collection, or by the comparative recency of the occurrence of any species, and the circumstances under which that occurrence has been announced. Species which are not to be found in any of our Museums, for which no authority is known, or whose claims rest on statements made many years back, at a time when specific differences were but little attended to, he has no hesitation in saying, ought to have a separate place

* Several cases might be mentioned of species, which only a few years back had occurred but in a single instance, and which have since been met with more than once.

† This is said with reference only to the Vertebrate Division of the British Fauna. Mr. Stephens has taken great pains to draw the line spoken of in the case of our native Insects.

allotted to them as *doubtful natives*, and not to be mixed up with others which labour under no uncertainty of this nature.

Another line which the author has attempted to draw, is that between what may be termed *good* or *genuine* species, and such as are probably not distinct from others, or which are involved in some obscurity from the circumstance of their true characters not being well understood. In this latter division he has placed such animals as the *Catodon Sibbaldi* of Fleming, the *Red Lark* of Lewin, the *Lacerta ædura* and *L. anguiformis* of Sheppard, the *Comber Wrasse* of Pennant, the *Lesser Fork-beard* of Jago (*Raniceps Jago*, Flem.), and a vast many others, especially amongst the Fish, all which he would designate as *doubtful species*; but he has *not* placed in it animals, such as the *Exocætus* and *Hippocampus*, which are perfectly good species, as well as undoubtedly British, but of which the *exact* species met with in our seas or islands remains to be determined*. These are often cases in which it was not supposed that more than *one* species existed, at the time of their being enrolled in the British Fauna.

It must now be explained in what way these and other divisions, under which the Animals of this country may be parcelled, are distinguished in the present work. For this purpose recourse has been had to different types, and to two distinct sets of numbers. Firstly, all those well-ascertained, or at least *genuine* species, which are met with at the present day, or which have undoubtedly occurred at one time or another within the record of history, have one continuous numbering in each Class respectively, and form

* If there are cases which appear exceptions to this rule, such as those of the *Manatus borealis* of Fleming and the *Albacore* of Couch, they are instances in which not only the *species*, but even the *genus*, is as yet unascertained.

what may be termed the body of the work *. These are then divided into, firstly, such as are found now existing in a *truly wild* state; secondly, such as are *domesticated*, or, when at large, are supposed to have been originally *introduced* into these Islands; and thirdly, such as were indigenous in former times, but are now *extirpated*. In the case of the *truly wild* and *now existing* species, which of course constitute the great bulk of the Fauna, and which may be subdivided by those who please into *Residents*, *Periodical Visitants*, and *Stragglers*, the names and specific characters are printed in Small Pica; the synonyms and descriptions in Bourgeois. In the case of the *domesticated*, *naturalized*, and *extirpated* species, these types are exchanged for Bourgeois and Minion respectively; the first two are, however, particularly distinguished by having an asterisk (*) prefixed to their names, the last by a dagger (†) in like manner. The second principal division comprises the *doubtful natives* and *doubtful species*, which terms are employed in the sense in which they have been already explained. These, although inserted in their proper places in the system, are cut off, as it were, from all those above mentioned, and marked by a distinct numbering enclosed in brackets. The types resorted to are the same as those adopted in the case of the naturalized and extirpated species, but these types are here set a little way in from the margin. By such an arrangement the attention is more readily drawn to these animals which stand in such particular need of further investigation by the naturalist.

* It may be here observed, that a single species has been inserted in the body of the work as British, which perhaps is hardly entitled to be considered in that light. The author alludes to the *Balaena Physalus*, Linn., of the occurrence of which in our seas, he can find no certain instance on record. He also entertains some doubts respecting the *Sparus Aurata*, from the circumstance of its having been so often confounded with the *S. centrodonatus*.

Some other matters remain now to be spoken of. It has been already intimated that the descriptions are as far as possible original. It may be added, that in a large number of instances they are derived from recent specimens. It must be obvious, however, that in preparing a work of this nature, there will always be many species, especially among the large marine animals, which it is impossible for any individual to describe from his own observation. In such cases, then, recourse has been had to what have appeared the best authorities, more especially to such persons as have published any thing original on the species in question; and where under such circumstances the accounts of authors differ, the discrepancies are pointed out. It is believed that in almost all the above cases the name of the author, from whom any thing is borrowed, is sub-joined*, who of course is responsible for the accuracy of what is stated. Desmarest and Scoresby (the last for the Cetaceous Animals) are the authorities mostly resorted to in the Class *Mammalia*; Temminck in that of *Birds*; Cuvier and Valenciennes, as well as Bloch, and occasionally Pennant and Donovan, in that of *Fish*. By some it may be thought that the descriptions are too long, and run needlessly into detail. But when it is considered how many species have been overlooked from their supposed identity with others; how many, some even of the most common occurrence, have been misunderstood, and referred

* A few instances occur in the Class *Mammalia*, in which this authority was omitted to be annexed. The cases in question are those of the *Extirpated Quad-rupeds*, the *Common and Great Seals*, the *Walrus*, the *Red Deer*, the *Roe*, and two or three of the *Cetacea*, the descriptions of which have been borrowed, partly from Desmarest, and partly from other sources. There are also three species of *Birds* similarly circumstanced; the *Shore Lark*, the *Rock Ptarmigan*, and the *Virginian Partridge*. The description of the first is taken from Temminck and Wilson; that of the second from Sabine, and the "*Fauna Boreali-Ame-ricana*;" that of the third from Temminck.

to others which perhaps are not inhabitants of this country ; and that these and similar errors have arisen not merely from the imperfect, but, it must be added, careless descriptions which have been given of such animals, it is hoped that the pains which have been taken to render this portion of the work as complete and accurate as possible, will not be thought entirely thrown away. The plan which the author has adopted in most instances, in the case of the Fish especially, which of all our British *Vertebrata* have till lately been the least attended to, and, in regard to the distinction of species, the worst understood, has been to describe as minutely as possible the first species in each genus, or sub-genus. if it present a marked modification of form, and then, when there were other nearly allied species, to restrict himself principally to the *differences* which were observable in these last, with reference to the first and to one another. By this method, which is adopted from MM. Cuvier and Valenciennes, the enquirer is more readily enabled to identify the particular species he may have before him. In the Class of Birds, the different variations of plumage, arising from age and season, have been pointed out and characterized so far as they are known ; and this plan has been pursued even in the case of those species which have occurred in this country as yet only in the immature state, for the more complete guidance of our own naturalists, in the event of their being met with in the adult plumage.

What has been said of the descriptions applies also to the measurements. They are to be considered as original, excepting where the name of any author stands attached to them. They have been taken with much care, in by far the larger number of instances, from recently killed specimens, and many of them are the mean results obtained from mea-

asuring a large number of individuals. In the Class of Birds, they are generally those of the *male*, unless stated to the contrary. It may be observed that in this Class the *entire length* is measured from the tip of the bill to the extremity of the tail; the length of the bill is estimated in two directions, firstly from the frontal feathers, following the curvature of the ridge, secondly from the gape or angle of the mouth; the length of the tail is measured from the extremity of the longest quill to its insertion in the coccyx.

Fish being very variable in size, and having no very well marked limit of growth, it was thought that the *absolute* dimensions of any species, would, if given, prove of little value. Hence in this Class, the author has generally restricted himself to noting the average length to which the species attains; the *relative proportions*, which are often of great importance, being introduced into the body of the description. With reference to these proportions, it may be stated, that the *entire length*, unless mentioned to the contrary, is always measured to the extremity of the caudal fin. The length of the head is measured from the end of the snout to the posterior margin of the gill-cover. The *depth* (termed the *breadth* in the case of the *Pleuronectidæ*) is reckoned vertically from the most elevated point in the line of the back, termed the *dorsal line*, to the corresponding point in the line of the abdomen, or *ventral line*, the dorsal and anal fins, unless mentioned to the contrary, not being included.

It may be added that in computing the fin-ray formula, which is similar to that adopted by MM. Cuvier and Valenciennes, all the rays are included which it was possible to distinguish, on the ground that if the short rays be omitted, which often pass insensibly into the longer ones,

it were difficult to know where to begin the reckoning. The mention of this circumstance will serve to explain why the number of fin-rays as stated in this work will be often found to exceed the number given by former authors, who appear in general to have made their computation without much attempt at accuracy. The above remark, however, does not apply to the caudal fin, in which, generally, though not always, there is a tolerably well marked line of separation between what may be termed the *principal* rays, and the *accessory* or shorter ones. In many instances, in which this distinction is evident, these two kinds of rays are reckoned separately.

Appended to the description of each species, are a few general remarks illustrative of its habits; more especially those connected with locality, food, and propagation. It was thought that these would render the work more generally useful. Many of them are the result of the author's own observation, though some are confessedly obtained from other sources.

On the subject of Classification, it must be remarked that the system of no one individual author has been rigidly adhered to. Regard has been paid to what has been written on this subject by the most recent writers in each department, and all the larger groups, as well as, in most instances, their mode of collocation, have been derived from such sources. The arrangement of the Mammalia has been drawn up from a combined view of the system of Cuvier, and the systems of Gray* and MacLeay†. That of the Birds from a similar view of the system of Vigors‡, and the modifications of that system as adopted by Swainson and

* *Ann. of Phil.* vol. xxvi. p. 337. † *Linn. Trans.* vol. xvi. p. 1, &c.

‡ *Linn. Trans.* vol. xiv. p. 395. See also some Articles by the same author in the first and second volumes of the "*Zoological Journal*."

Selby; assistance has been also gained, in regard to the situation of a few genera, from the system of Lesson*. The arrangement of the Reptiles and Amphibious Animals is for the most part in accordance with that in the "Regne Animal" of Cuvier. The author has, however, followed Latreille and many modern naturalists in considering these groups as two distinct classes. The arrangement of the Fish is likewise similar to that in the *second edition* of the "Regne Animal," excepting a slight alteration in the value of the larger groups, adopted from the Prince of Musignano†. In no case must it be imagined that the order of affinities is exactly the same as the order of arrangement given, since, if it be true that all natural groups be circular, an opinion now generally prevalent, and one in which the author is strongly disposed to join, it is clearly impossible to preserve such a coincidence under circumstances which necessarily entail the appearance of a linear series.

The author has also exercised his own judgment in the adoption of certain genera and sub-genera. With respect to these last, having expressed his views elsewhere‡, it is unnecessary to repeat them in this place. He may simply state that he has endeavoured to acknowledge, and to act up to, the principle which he has there advocated; although, from the ignorance in which we are upon the subject of the real value of many groups, it cannot be doubted that numerous instances occur, in which he has failed making a correct application of it. Being aware of

* *Traité d'Ornithologie; ou Tableau méthodique des Ordres, Sous-ordres, Familles, Tribus, Genres, Sous-genres et Races d'Oiseaux.* Par. 1831. 8vo.

† *Saggio di una Distribuzione metodica degli Animali Vertebrati.* Rom. 1831. 8vo. The alteration above alluded to has been adopted by the author since the publication of his "Systematic Catalogue."

‡ *Lond. Mag. of Nat. Hist.* vol. VI. p. 385. and vol. VII. p. 97.

the dislike which many of our own naturalists have to the adoption of sub-genera, he begs to state that they are inserted in such a manner, that any one who chooses may place them upon the same footing with the genera, or, if he be not a friend to the subdivision of the old-established groups, take no notice of them at all.

Instead of prefixing the generic characters to the several genera respectively as they occur in order, they are presented in a synoptic form at the head of each of the Classes, by which means a better view is obtained of the relative collocation and affinities of the larger groups. Moreover, the same kind of arrangement with respect to types prevails here, as that which occurs in the other part of the work. The characters of all those genera and sub-genera which contain *truly wild*, as well as *genuine* and *now existing* species, are printed in Small Pica and Bourgeois respectively, the names of the genera standing in LARGE CAPITALS, those of the sub-genera in SMALL CAPITALS. In the case of the *domesticated*, *naturalized*, and *extirpated* animals, these types are exchanged respectively, as before, for Bourgeois and Minion; the first two, being, also in like manner, particularly distinguished by an asterisk, the last by a dagger. The same types are employed for the characters of those genera and sub-genera which contain only *doubtful natives* or *doubtful species*, but these may be readily distinguished from the last by the circumstance of the names standing in ITALIC Capitals, which are large or small, as in the former instances, according as the group in question is either a genus or sub-genus. Moreover, in this division, the genera have a distinct numbering enclosed in brackets: the sub-genera no numbering at all. It may be further observed, that the characters of the orders and families are printed in Pica, and that

here likewise the type is exchanged for one of subordinate size in the case of two families (*Phasiagnidæ* and *Siluridæ*), one containing no true natives, the other only a doubtful native. In accordance, also, with the rule before given, the former is distinguished by an asterisk; the latter, by the name standing in *ITALIC CAPITALS*, and its being, as in the case of sub-genera similarly circumstanced, without a number.

The specific characters, which are for the most part entirely new, have been drawn up, as far as practicable, with a view to the exact differences of species; but the difficulty of doing this in the instance of the aquatic Birds, which are subject to such great changes of plumage from age and other circumstances, it must be confessed is very great, and has not in all cases been satisfactorily got over. In the Class of Fish, specific characters have seldom been added in the case of genera or sub-genera containing not more than one *British* species, or when given, bear reference solely to other, nearly allied, *European* species; the essential characters of those met with in other countries, in many groups, not having been hitherto determined with precision.

The terms employed in characterizing the larger groups, or in drawing up the descriptions of species, have for the most part been derived from the works of Illiger*, Desmarest†, and Cuvier. The “*Histoire Naturelle des Poissons*” of this last author has been exclusively resorted to in the Class of Fish. To these works, the reader, to whom the terms are not familiar, is accordingly referred.

* *Prodromus Systematis Mammalium et Avium*. Berol. 1811. 8vo. This work will be found to contain a very complete terminology with reference to the two classes of Mammalia and Birds.

† *Terminologie des Mammifères*; prefixed to his “*Mammalogie*.”

On the subject of Nomenclature it may be remarked that the oldest names have been adopted in most instances, unless a different one has been sanctioned by general use. As the author is of opinion that in common parlance there is no occasion to name the sub-genus, when speaking of any particular species, he has not thought it necessary to change the names of species which are the same as those of the sub-genera to which they belong.

The synonyms will be found to embrace references to some of the principal writers, more especially those in our own country, who have treated of the different Classes, or of any subordinate group, or any particular species. These references have, in every instance, been examined personally by the author. To facilitate the inquiries of such persons, as may wish to examine for themselves on this subject, a complete list has been annexed of all those works, with their several editions, which are quoted, either for the synonyms, or for any other purpose.

It has been thought proper to annex two distinct indexes, one containing the Latin, the other the English names. No pains have been spared to render them as complete as possible. It is believed that every name and synonym, in the above two languages, occurring in the body of the work, will be found in its proper place.

It now only remains for the author to express his acknowledgments to those friends who have assisted him in this undertaking. To Mr. Yarrell in particular, he begs publicly to return his sincere thanks for the able help which he has experienced at his hands, and such as alone has enabled him to complete the work upon the plan first contemplated. This help has been especially felt upon the subject of the British Fishes. Had it not been for the very liberal manner in which that gentleman offered him

the almost unlimited use of his Manuscripts and rich collections, the author has no hesitation in saying that he could never have extended the Manual to that department, or presumed to enter upon a field, to which he was previously almost an entire stranger. Assistance, however, has been not the less afforded him in the other Classes. Mr. Yarrell's well-known practical acquaintance with our British Birds has enabled the author to detail more at length the changes of plumage to which some species are liable, and to correct a few errors into which previous writers had fallen on this subject. The same gentleman kindly volunteered an accurate description, accompanied by measurements, of the egg of every species, of which his extensive collection afforded specimens; thus enhancing the utility of the work by an addition, which, but for this circumstance, the author would have been unable to supply. He begs it may be distinctly understood that this portion of it is from the pen of Mr. Yarrell.

To Mr. Gray, he desires to make an acknowledgment of the readiness with which he has at all times allowed him to consult the specimens in the British Museum. The same return must be made to the officers of the Zoological Society for similar liberty to examine a few specimens contained in their collection.

In conclusion, the author may state that he has no intention of extending his work to the Invertebrate Division of the British Fauna. In the present advanced state of the Science, a complete Manual of all the Animals occurring in these Islands can only be accomplished by the united labours of many individuals. With the view, however, of continuing his researches into that portion of it which is here treated of, he begs to solicit such observations, notices of new or rare species, and, where it may

not be inconvenient, specimens, as it may be in the power of any of his readers to supply. Even the Vertebrate Animals of our own country are far yet from being thoroughly understood. Much confusion prevails in several groups, and, without doubt, many additional species remain to be detected. The *Shrews* and *Bats* amongst Quadrupeds; the *Cetaceous* Animals; the *Wrasses*, *Gobies*, *Blennies*, the *Salmon* Tribe, the *Sharks* and *Rays*, amongst Fish; these and other families might be pointed out, with respect to which we want more information, and which particularly invite the attention of the British Naturalist. Should any one be disposed to honour him with specimens in these or other instances, the author begs to state that they will not be reserved for his own use exclusively, but will, with the permission of the donor, be deposited in the Museum of the Cambridge Philosophical Society, which already possesses an extensive collection of British Animals, and which the author is anxious to render as perfect as possible in that department of the Fauna, to the advancement of which the Work now offered to the Public is directed.

SWAFFHAM BULBECK,
Oct. 24, 1835.

THE Author takes this opportunity of expressing his grateful acknowledgments to the SYNDICS of the University Press, for their liberality in taking upon themselves the expense of printing this Work.

ERRATA.

PAGE	LINE	ERROR.	CORRECTION.
52	16	LANIIDÆ.	LANIADÆ.
53	11	from the bottom ; SYLVIIDÆ.	SYLVIADÆ.
59	12	 CERTHIIDÆ.	CERTHIADÆ.
64	8	 CHARADRIIDÆ.	CHARADRIADÆ.
126	7	 vol. 11.	vol. 1. part ii.
137	2	 dele &.	
160	13	<i>Europæan.</i>	<i>European.</i>
252	14	<i>C. Urinator.</i>	<i>Colymbus Urinator.</i>
360	14	from the bottom, } and elsewhere } <i>Mackarel.</i>	<i>Mackerel.</i>
434	2	CLUPEA, <i>Linn.</i>	CLUPEA, <i>Cuv.</i>
445	7	<i>peritonæum.</i>	peritoneum.

ALPHABETICAL LIST

OF

WORKS QUOTED.

- Ann. des Sci.*—ANNALES des Sciences Naturelles. Paris, 1824, &c. 8vo.
- Ann. du Mus.*—Annales du Muséum d'Histoire Naturelle. Paris, 1802–1813. 4to. 20 vols.
- Ann. of Phil.*—The Annals of Philosophy. New Series. Lond. 1821–1826. 8vo. (*Vol. 6, or 22 from the commencement, quoted.*)
- Berkenh. Syn.*—Berkenhout (J.). Synopsis of the Natural History of Great Britain and Ireland; being a Second Edition of the Outlines, &c. Lond. 1789. 8vo. 2 vols.
- Bew. Brit. Birds.*—Bewick (T.). A History of British Birds. Sixth Edition. Newcastle, 1826. 8vo. 2 vols.
- *Quad.*—Bewick (T.). A General History of Quadrupeds. The Figures engraved on Wood. Eighth Edition. Newcastle-upon-Tyne, 1824. 8vo.
- Bloch, Ichth.*—Bloch (M. E.). Ichthyologie; ou Histoire Naturelle des Poissons. Berlin, 1785–1795. fol. 12 parts.
- Bon. Am. Orn.*—Bonaparte (C. L.). American Ornithology; or the Natural History of Birds inhabiting the United States, not given by Wilson. Philadelph. 1825, &c. 4to. 4 vols.
- *Faun. Ital.*—Bonaparte (C. L.). Iconographia della Fauna Italica. Rom. 1832, &c. fol. (*In course of publication.*)
- *Syn.*—Bonaparte (C. L.). The Genera of North American Birds, and a Synopsis of the Species found within the Territory of the United States. New York, 1828. 8vo. (*From the Annals of the Lyceum of Natural History of New York.*)
- Borl. Cornw.*—Borlase (W.). The Natural History of Cornwall. Oxford, 1758. fol.
- Bowd. Brit. fr. wat. Fish.*—Bowdich (Mrs. T. E.). The Fresh-water Fishes of Great Britain; drawn and described. Lond. 1828, &c. 4to. (*In course of publication.*)

- Briss. Orn.*—Brisson. Ornithologia, sive Synopsis Methodica Avium, &c. Lug. Bat. 1763. 8vo. 2 vols.
- *Reg. An.*—Brisson. Regnum Animale in classes ix. distributum, cum duarum primarum classium, Quadrupedum scilicet et Cetaceorum particulari divisione. Paris, 1756. 4to.
- Brown, Illust.*—Brown (P.). New Illustrations of Zoology. Lond. 1776. 4to.
- Buff. Hist. Nat.*—Buffon. Histoire Naturelle générale et particulière, avec la description du Cabinet du Roi. Paris, 1750–1767. 4to. 15 vols.
- Bull. des Sci. Nat.*—Bulletin des Sciences Naturelles et de Géologie. Paris, 1824–1831. 8vo. (*Vol. 4, for the year 1825, quoted.*)
- Camb. Phil. Trans.*—Transactions of the Cambridge Philosophical Society. Cambridge, 1822, &c. 4to.
- Child. Add. to Zool. Cl.*—Children (J. G.). An Address delivered at the Anniversary Meeting of the Zoological Club of the Linnean Society, held Nov. 29, 1827. Lond. 1827. 8vo.
- Cuv. Oss. Foss.*—Cuvier (G.). Recherches sur les Ossemens Fossiles. Paris, 1821–1824. 4to. 5 vols.
- *Reg. An.*—Cuvier, (G.) Le Règne Animal distribué d'après son organisation. Paris, 1817. 8vo. 4 vols. The same. Paris, 1829. 8vo. 5 vols. (*The second edition is the one quoted, unless stated to the contrary.*)
- & *Val. Poiss.*—Cuvier et Valenciennes. Histoire Naturelle des Poissons. Paris, 1828–1833. 4to. 9 vols. (*In course of publication.*)
- Dan. Rur. Sports.*—Daniel (W. B.). Rural Sports. Lond. 1801–1802. 4to. 2 vols.
- Daud. Hist. Nat. des Rept.*—Daudin (F. M.). Histoire Naturelle des Reptiles. Paris, 1805. 8vo. 8 vols.
- Desm. Mammal.*—Desmarest (A. G.). Mammalogie; ou description des espèces de Mammifères. Paris, 1820–1822. 4to. 2 parts.
- Don. Brit. Birds.*—Donovan (E.). The Natural History of British Birds. Lond. 1794–1819. 8vo. 10 vols.
- *Brit. Fish.*—Donovan (E.). The Natural History of British Fishes. Lond. 1808. 8vo. 5 vols.
- *Brit. Quad.*—Donovan (E.). The Natural History of British Quadrupeds. Lond. 1820. 8vo. 3 vols.
- Edinb. Journ. of Nat. and Geog. Sci.*—The Edinburgh Journal of Natural and Geographical Science. Edinb. 1830, &c. Directed by W. Ainsworth and H. Cheek. 8vo. 3 vols.
- Edinb. New Phil. Journ.*—The Edinburgh New Philosophical Journal. Conducted by R. Jameson. Edinb. 1826, &c. 8vo.

- Edinb. Phil. Journ.*—The Edinburgh Philosophical Journal. Conducted by R. Jameson. Edinb. 1819–1826. 8vo. 14 vols.
- Edw. Glean.*—Edwards (G.). Gleanings of Natural History. Lond. 1758–1764. 4to. 3 parts.
- *Nat. Hist.*—Edwards (G.). A Natural History of uncommon Birds, and of some other rare and undescribed Animals. Lond. 1743–1751. 4to. 4 parts.
- Encycl. Brit.*—Encyclopædia Britannica. Seventh Edition. Edited by Professor Napier. Edinb. 1834, &c. 4to. (*In course of publication.*)
- Faun. Bor. Am.*—Fauna Boreali-Americana; or the Zoology of the Northern parts of British America. Part second, The Birds. By W. Swainson, and J. Richardson. Lond. 1831. 4to.
- *Franç.*—Faune Française, ou Histoire Naturelle, générale et particulière, des Animaux qui se trouvent en France, &c. Par MM. Vieillot, Desmarest, De Blainville, Audinet-Serville, St. Fargeau, et Walkenäer. Paris, 1824, &c. 8vo. (*Incomplete.*)
- Flem. Brit. An.*—Fleming (J.). A History of British Animals, &c. Edinb. 1828. 8vo.
- *Phil. Zool.*—Fleming (J.). The Philosophy of Zoology; or a general view of the structure, functions, and classification of Animals. Edinb. 1822. 8vo. 2 vols.
- Gmel. Linn.*—Gmelin (J. F.). Caroli a Linné, Systema Naturæ, &c. Lips. 1788, &c. 8vo. vol. I., Animal Kingdom.
- Gould, Europ. Birds.*—Gould (J.). The Birds of Europe. Lond. 1832, &c. fol. (*In course of publication.*)
- Gray, Syn. Rept.*—Gray (J. E.). Synopsis Reptilium; or short descriptions of the Species of Reptiles. Part i. Cataphracta. Lond. 1831. 8vo. (*No more has been hitherto published.*)
- Grew, Rar.*—Grew (N.). Musæum Regalis Societatis; or a Catalogue and Description of the natural and artificial Rarities belonging to the Royal Society and preserved at Gresham College. Lond. 1681. fol.
- Hast. Worcest.*—Hastings (C.). Illustrations of the Natural History of Worcestershire, &c. Lond. 1834. 8vo.
- Heysh. Cumb. An.*—Heysham (J.). A Catalogue of Cumberland Animals. (*Prefixed to the 1st volume of Hutchinson's History of the County of Cumberland.* Carlisle, 1794. 4to.)
- Horsf. Zool. Research.*—Horsfield (T.). Zoological Researches in Java, and the neighbouring Islands. Lond. 1824. 4to.
- Jago.*—Catalogus quorundam Piscium rariorum, quos oris Cornubiæ maritimis nuper observavit, et delineavit, &c. G. Jago, &c. (*In Ray's Synopsis Piscium.* p. 162–166.)

- Jard. and Selb. Orn.*—Jardine (W.) and Selby (P. J.). Illustrations of Ornithology. Edinb. 1827, &c. 4to. (*In course of publication.*)
- Jen. Cat. of Brit. Vert. An.*—Jenyns (L.). A Systematic Catalogue of British Vertebrate Animals. Cambridge, 1835. 8vo.
- Jesse, Glean.*—Jesse (E.). Gleanings in Natural History. Second Series. Lond. 1834. 8vo.
- Journ. de Phys.*—Journal de Physique, de Chimie, d'Histoire Naturelle, et des Arts. Paris, 1773–1823. 4to. (*Vol. 71, the only one quoted.*)
- Lacép Cétac.*—La Cépède. Histoire Naturelle des Cétacées. Paris, 1803. 4to.
- *Poiss.*—La Cépède. Histoire Naturelle des Poissons. Paris, 1798–1803. 4to. 5 vols.
- Lath. Ind. Orn.*—Latham (J.). Index Ornithologicus; sive Systema Ornithologiæ; &c. Lond. 1790. 4to. 2 vols.
- *Syn.*—Latham (J.). A General Synopsis of Birds. Lond. 1781–1785. 4to. 3 vols. in 6 parts.
- *Supp.*—Supplement to the General Synopsis of Birds. Lond. 1787. 4to. Supplement II. Lond. 1801. 4to.
- Latr. Salam. de France.*—Latreille. Histoire Naturelle des Salamandres de France, précédé d'un Tableau méthodique des autres Reptiles indigènes. Paris, 1800. 8vo.
- Leach, Zool. Misc.*—Leach (W. E.). The Zoological Miscellany; being descriptions of new, or interesting Animals. Lond. 1814–1817. 8vo. 3 vols.
- Leigh, Lancash.*—Leigh (C.). The Natural History of Lancashire, Cheshire, and the Peak, in Derbyshire, &c. Oxford, 1700. fol.
- Lew. Brit. Birds.*—Lewin (W.). The Birds of Great Britain, with their Eggs, accurately figured. Lond. 1789–1794. 4to. 7 vols.
- Linn. Syst. Nat.*—Linnaeus (C.). Systema Naturæ, &c. Edit. 12. Holm. 1766–1768. 8vo. 3 vols. Vol. 1, Animal Kingdom.
- Linn. Trans.*—Transactions of the Linnean Society of London. Lond. 1791, &c. 4to.
- L'Institut.*—L'Institut; Journal des Académies et Sociétés scientifiques de la France et de l'Etranger. Paris, 1833, &c. 4to.
- Lond. & Edinb. Phil. Mag.*—The London and Edinburgh Philosophical Magazine and Journal of Science. By Sir D. Brewster, R. Taylor, and R. Philips. Lond. 1832, &c. 8vo.
- Lond. Quart. Journ. of Sci.*—The Quarterly Journal of Science, Literature, and the Arts. Lond. 1816–1830. 8vo. 29 vols.
- Lond. Mag. of Nat. Hist.*—The Magazine of Natural History, and Journal of Zoology, Botany, Mineralogy, Geology, and Meteorology. Conducted by J. C. Loudon. Lond. 1829, &c. 8vo.

- Low, Faun. Orc.*—Low (G.). Fauna Orcadensis; or the Natural History of the Quadrupeds, Birds, Reptiles, and Fishes, of Orkney and Shetland. Edinb. 1813. 4to.
- Lyell, Geol.*—Lyell (C.). Principles of Geology, &c. Lond. 1830–1833. 8vo. 3 vols.
- Mac Cull. West. Isles.*—Mac Culloch (J.). A Description of the Western Islands of Scotland, including the Isle of Man. Lond. 1819. 2 vols. 8vo. and 4to atlas of plates.
- Mém. de la Soc. de Gen.*—Mémoires de la Société de Physique, et d'Histoire Naturelle de Genève. Genév. 1821, &c. 4to. (*Vol. 3 quoted.*)
- Mém. du Mus.*—Mémoires du Muséum d'Histoire Naturelle. Paris, 1815–1832. 4to. 20 vols.
- Merr. Pin.*—Merrett (C.). Pinax Rerum Naturalium Britannicarum. Lond. 1666. 8vo.
- Mont. Orn. Dict.*—Montagu (G.). Ornithological Dictionary; or Alphabetical Synopsis of British Birds. Lond. 1802. 8vo. 2 vols.
- *Supp.*—Montagu (G.). Supplement to the Ornithological Dictionary. Exeter, 1813. 8vo.
- Mull. Zool. Dan.*—Muller (O. F.). Zoologia Danica. Haun. 1788–1806. fol. 4 vols.
- Nash, Worcest.*—Nash (T.). Collections for the History of Worcestershire. Lond. 1781. fol. 2 vols.
- Newcast. Nat. Hist. Trans.*—Transactions of the Natural History Society of Northumberland, Durham, and Newcastle. Newcastle. 1831, &c. 4to.
- Nichol. Journ.*—A Journal of Natural Philosophy, Chemistry, and the Arts. Lond. 1802–1813. 8vo. 36 vols. (*Vol. 22, for the year 1809, quoted.*)
- Nilss. Prod. Ichth. Scand.*—Nilsson (S.). Prodrömus Ichthyologiæ Scandinaviæ. Lund. 1832. 8vo.
- Nutt. Orn. of Un. St.*—Nuttall (T.). A Manual of the Ornithology of the United States and of Canada. Boston, 1832–1834. 8vo. 2 vols.
- Osbeck, Voy. to China.*—Osbeck (P.). A Voyage to China and the East Indies. Lond. 1771. 8vo. 2 vols. (*Translated from the German by J. R. Forster.*)
- Paget, Nat. Hist. of Yarm.*—Paget (C. J. & J.). Sketch of the Natural History of Yarmouth and its Neighbourhood, containing Catalogues of the Species of Animals, Birds, Reptiles, Fish, Insects, and Plants, at present known. Yarmouth, 1834. 8vo.
- Penn. Arct. Zool.*—Pennant (T.). Arctic Zoology. Lond. 1792. 4to. 2 vols.

- Penn. Brit. Zool.*—Pennant (T.). British Zoology. Fourth Edition. Lond. 1776-1777. 8vo. 4 vols. The same. New Edition. Lond. 1812. 8vo. 4 vols. (*The Fourth Edition is the one quoted, unless stated to the contrary.*)
- Phil. Trans.*—Philosophical Transactions of the Royal Society of London. Lond. 1665, &c. 4to.
- Plot, Oxfordsh.*—Plot (R.). The Natural History of Oxfordshire. Oxford, 1705. fol. (*Second Edition.*)
- Proceed. of Berw. Nat. Club.*—Proceedings of the Berwickshire Naturalist's Club. Edinb. 1834, &c. 8vo.
- *Zool. Soc.*—Proceedings of the Zoological Society of London. Lond. 1830, &c. 8vo.
- Pult. Cat. Dors.*—Pulteney (R.). Catalogue of the Birds, Shells, and some of the more rare Plants of Dorsetshire. From the new and enlarged Edition of Mr. Hutchin's History of that county. Lond. 1813. fol.
- Quart. Rev.*—The Quarterly Review. vol. 47. Lond. 1832. 8vo.
- Ray, Syn. Av.*—Ray (J.). Synopsis methodica Avium. Lond. 1713. 8vo.
- *Syn. Pisc.*—Ray (J.). Synopsis methodica Piscium. Lond. 1713. 8vo.
- *Syn. Quad.*—Ray, (J.). Synopsis methodica Animalium Quadrupedum et Serpentinum Generis. Lond. 1693. 8vo.
- Richards. App. Parr. Voy.*—Richardson (J.). Zoological Appendix to Parry's Second Voyage. 1824. 4to.
- Riss. Hist. Nat.*—Risso (A.). Histoire Naturelle des principales productions de l'Europe Méridionale, et particulièrement de celles des environs de Nice et des Alpes maritimes. Par. 1826. 8vo. 5 vols. (*Vol. 3, quoted.*)
- Ros. Ran.*—Ræsel (A. J.). Historia Naturalis Ranarum nostratium. Nurnberg. 1758. fol.
- Rond. Pisc.*—Rondeletius (G.). De Piscibus marinis. Lug. Bat. 1554. fol.
- Ross, Voy. App.*—Appendix to Ross's Voyage of Discovery. Lond. 1819. 4to.
- Sab. Supp. Parr. Voy.*—A Supplement to the Appendix of Captain Parry's Voyage for the discovery of a N.W. passage in the years 1819, 1820, containing an account of the subjects of Natural History. Lond. 1824. 4to. The Birds by Capt. Sabine.
- Salmon.*—Salmonia; or Days of Fly-fishing. By an Angler. Second Edition. Lond. 1829. 8vo.
- Scoresb. Arct. Reg.*—Scoresby (W.). An Account of the Arctic Regions, with a history and description of the Northern Whale-fishery. Edinb. 1829. 8vo. 2 vols.

- Selb. Illust.*—Selby (P. J.). Illustrations of British Ornithology. Edinb. 1833. 8vo. 2 vols. Plates to the same. Eleph. fol.
- Shaw, Gen. Zool.*—Shaw (G.). General Zoology. Lond. 1800–1826. 8vo. 14 vols. (*The last six volumes are by Mr. Stephens.*)
- *Nat. Misc.*—Shaw (G.). The Naturalists Miscellany. Lond. 1790, &c. 8vo. 24 vols.
- Sibb. Phalain.*—Sibbald (R.). Phalainologia Nova; sive observationes de rarioribus quibusdam Bakenis in Scotiæ Littus nuper ejectis. Edinb. 1773. 8vo.
- *Scot. Illust.*—Sibbald (R.). Scotia Illustrata; sive Prodrômus Historiæ Naturalis, &c. Edinb. 1684. fol.
- Sloane, Jam.*—Sloane (Hans). A Voyage to the Islands Madeira, Barbados, Nieves, St. Christopher's and Jamaica, with the Natural History of the last of those Islands. Lond. 1707–1725. fol.
- Sow. Brit. Misc.*—Sowerby (J.). The British Miscellany; or coloured figures of new, rare, or little known Animal subjects; many not before ascertained to be inhabitants of the British Isles. vol. 1. Lond. 1806. 8vo. (*The work was not continued beyond the first volume.*)
- Stew. El. of Nat. Hist.*—Stewart (C.). Elements of the Natural History of the Animal Kingdom. Second Edition. Edinb. 1817. 8vo. 2 vols.
- Swains. Zool. Ill.*—Swainson (W.). Zoological Illustrations; or original figures and descriptions of new, rare, or interesting Animals, &c. Lond. 1820–1823. 8vo. 3 vols.
- Temm. Man. d'Orn.*—Temminck (C. J.). Manuel d'Ornithologie, ou Tableau Systématique des Oiseaux qui se trouvent en Europe; &c. Second Edition. Parts 1 & 2. Paris, 1820. Part 3. Par. 1835. 8vo.
- *Monog. de Mamm.*—Temminck (C. J.). Monographies de Mammalogie. Paris, 1827. 4to.
- *Pig. et Gall.*—Temminck (C. J.). Histoire Naturelle Générale des Pigeons et des Gallinacés. Amsterdam, 1813–1815. 8vo. 3 vols.
- Turt. Brit. Faun.*—Turton (W.). British Fauna; containing a Compendium of the Zoology of the British Islands. Swansea, 1807. 12mo.
- *Linn.*—Turton (W.). A General System of Nature, &c. Translated from Gmelin's last edition of the Systema Naturæ, by Sir Charles Linné. Swansea, 1800. 8vo. Vols. 1–4, Animal Kingdom.
- Vieill. Gal. Ois.*—Vieillot (L. P.). La Galerie des Oiseaux. Paris, 1825. 4to. 2 vols.
- Wagl. Syst. Av.*—Wagler (J.). Systema Avium. Pars prima. Stuttgart, 1827. 8vo. (*Incomplete.*)

- Wern. Mem.*—Memoirs of the Wernerian Natural History Society.
Edinb. 1811, &c. 8vo.
- White, Selb.*—White (G.). The Natural History of Selborne, to which
are added The Naturalist's Calendar, Miscellaneous Observations,
and Poems. Lond. Edit. 1813. 8vo. 2 vols.
- Will. Hist. Pisc.*—Willughby (F.). De Historia Piscium Libri Quatuor,
&c. Totum opus recognovit, coaptavit, supplevit, Librum etiam
primum et secundum integros adjecit, Johannes Raius. Curâ
Cromwelli Mortimeri, M.D. Lond. 1743. fol.
- Wils. Amer. Orn.*—Wilson (A.). American Ornithology; or the Natural
History of the Birds of the United States. Philadelph. 1808—1814.
4to. 8 vols.
- Yarr. Brit. Fish.*—Yarrell (W.). A History of British Fishes. Illustrated
by Wood-cuts of all the species. vol. 1. Lond. 1835. 8vo. (*In course
of publication.*)
- Zool. Journ.*—The Zoological Journal. Lond. 1825, &c. 8vo.
-

VERTEBRATA.

VERTEBRATE animals are characterized by having the brain and principal trunk of the nervous system included in a bony articulated case, composing the skull and vertebral column. They have all red blood, and a muscular heart. The mouth is furnished with two jaws moving vertically. They have distinct organs of sight, hearing, smell, and taste. The limbs are never more than four in number. The sexes are in all cases separate.

They are divisible into the following five classes :

* *With warm blood : heart with two auricles and two ventricles.*

I. MAMMALIA.—Viviparous animals, suckling their young : breathing by lungs. Body generally covered with hair, and furnished in most cases with four feet.

II. AVES.—Oviparous animals ; breathing by lungs. Body covered with feathers ; furnished with two wings and two feet.

****** *With cold blood: heart with one or two auricles,
and one ventricle.*

III. REPTILIA.—Oviparous animals; breathing by lungs. Body covered with scales or shelly plates. Impregnation effected by sexual union: the young undergoing no metamorphosis.

IV. AMPHIBIA.—Oviparous animals; breathing by lungs and gills. Body covered with a soft naked skin: feet without claws. Eggs impregnated after exclusion: the young undergoing metamorphosis.

V. PISCES.—Oviparous animals; breathing by gills only. Body covered with scales, and furnished with fins instead of feet.

.

CLASS I. MAMMALIA.

§ 1. *Incisor, canine, and molar teeth all present, forming a continuous series.*

ORDER I. FERÆ.

Feet armed with strong claws: mammæ abdominal: penis sheathed.

* *Six incisors in each jaw: grinders of three sorts.*

† *Walk on the soles of the feet.*

† 1. URSUS.—Lower incisors set in the same line: grinders varying in number, the three last large and tubercular: no glandular pouch under the tail.

2. MELES.—Second incisor on each side in the lower jaw placed behind the others: grinders five above and six below on each side; the first very small, the last tubercular: body low on the legs: a glandular pouch or scent-bag under the tail.

†† *Walk on the extremities of the toes.*

3. MUSTELA.—Grinders varying in number: body very much elongated: feet short; the toes separate: ears short and rounded.

(1. MUSTELA.) Grinders five above and six below on each side: tongue smooth.

(2. PUTORIUS.) Grinders four above and five below on each side: tongue rough.

4. LUTRA.—Grinders five above and five or six below: body long: feet short, with five palmated toes: head depressed: ears very short: tongue slightly rough: tail flattened horizontally.

5. CANIS.—Grinders six above and seven below on each side; the two last in each jaw tubercular: ears moderately large: tongue smooth: claws not retractile.

(1. CANIS.) Pupil circular.

(2. VULPES.) Pupil linear: tail bushy.

6. FELIS.—Grinders four above and three below on each side; no tubercular grinder in the lower jaw: tongue armed with prickly papillæ pointing backwards: claws retractile.

*** Teeth various: feet short and fin-shaped, adapted to swimming; those behind horizontal.*

7. PHOCA.—Incisors and tusks in both jaws: grinders uniform; each of three lobes; the central lobe triangular, large and cutting, the anterior and posterior lobes small.

8. TRICHECHUS.—No incisors or tusks in the lower jaw: tusks in the upper jaw greatly produced, directed downwards: grinders cylindrical, short, and obliquely truncated.

**** Incisors varying in number: summits of the grinders with conical points: soles of the hinder feet applied to the ground.*

9. TALPA.—Incisors six above and eight below, nearly equal: tusks large: grinders seven above and six below on each side: body very thick: fore feet short and broad, formed for digging: tail short: no external ears: eyes very minute.

10. SOREX.—Two middle incisors produced beyond the others; those above bent and notched at the base: lateral incisors or false tusks very small: grinders four above and three below on each side: body covered with hair: snout attenuated: tail long.

11. ERINACEUS.—Two middle incisors produced beyond the others, longer than the tusks; those above cylindrical, with a space between: grinders five above and four below on each side: body covered with spines: tail very short.

ORDER II. PRIMATES.

Claws flat and small: anterior extremities with a distinct thumb: mammæ pectoral: penis free and pendulous.

VESPERTILIONIDÆ.—*Incisors varying in number; summits of the grinders armed with sharp points: anterior and posterior extremities connected by a naked expansion of the skin, adapted to flight: fingers very long.*

12. RHINOLOPHUS.—Incisors two above and four below: grinders five on each side above and below: nostrils with two foliaceous appendages, the posterior one erect: ears free; tragus wanting.

13. VESPERTILIO.—Incisors four above and six below: grinders varying in number: nostrils without foliaceous appendages: ears sometimes free, sometimes united at their base; tragus always present.

- (1. VESPERTILIO.) Ears moderately large, lateral, separate: grinders from four to six above, and from three to six below on each side.

- (2. *PLECOTUS*.) Ears very large, much longer than the head, with their inner edges united at the base above the eyes: grinders five above and six below on each side.
- (3. *BARBASTELLUS*.) Ears moderate, united at the base above the eyes: a flat naked space on the forehead surrounded by a membranous edge: grinders four on each side above and below.
-

§ 11. *Incisor, canine, and molar teeth not all present, or not forming a continuous series.*

ORDER III. GLIRES.

Incisors two in each jaw, large, and strong, remote from the grinders: tusks none: toes distinct, with small conical claws.

** Furnished with strong clavicles.*

14. *SCIURUS*.—Grinders simple, with tubereular summits, five above and four below on each side: upper incisors chisel-shaped; lower ones pointed, compressed laterally: toes long, armed with sharp claws: tail long and bushy.

15. *MYOXUS*.—Grinders simple, their summits marked with transverse ridges of enamel, four on each side above and below: hair very soft and fine: tail very long, somewhat bushy.

16. *MUS*.—Grinders simple, with tubereular summits, three on each side above and below: upper incisors wedge-shaped, lower ones compressed and pointed: tail nearly naked, annulated with scales.

17. *ARVICOLA*.—Grinders compound, with flat summits, the enamel appearing in ridges on the surface: upper

incisors chisel-shaped, lower ones pointed: tail round and hairy, shorter than the body.

† 18. CASTOR.—Grinders compound, with flat summits, four on each side above and below: toes of the hinder feet palmated: tail oval, depressed, and covered with scales.

** *Clavicles rudimentary.*

19. LEPUS.—Two subsidiary incisors in the upper jaw immediately behind the others: grinders six above and five below on each side, the summits flat, with transverse plates of enamel: inside of the cheeks hairy: ears very large: toes five before and four behind: tail short, and turned up.

* 20. CAVIA.—Grinders compound; each with one simple and one fork-shaped plate of enamel; four on each side above and below: toes four before and three behind: tail none.

ORDER IV. UNGULATA.

Teeth various: incisors and tusks often wanting in one or both jaws: grinders of one sort: toes large, covered with hoofs.

* *No incisors in the upper jaw; eight in the lower: two middle toes separate: frontal bones furnished with horns: ruminant.*

† *Horns hollow, growing on a bony core; persistent.*

* 21. BOS.—Horns smooth; directed laterally at first, afterwards recurved: body thick and heavy: limbs strong: tail moderately long, terminated by a tuft of hair: four inguinal mammae.

* 22. OVIS.—Horns rough and angular; directed backwards and laterally, more or less spirally twisted: chin without a beard: limbs rather slender: tail moderately short: two inguinal mammae.

* 23. CAPRA.—Horns rough and angular; directed upwards and backwards: chin bearded: tail short: mammae two.

†† *Horns bony and solid: deciduous.*

24. CERVUS.—Horns branched or palmated; while growing covered with a soft velvety skin: body and limbs slender: four inguinal mammæ.

** *Six incisors in each jaw: two middle toes soldered in one: no horns: do not ruminare.*

* 25. EQUUS.

*** *Six incisors in each jaw; lower ones projecting forwards: four toes on each foot; two middle ones large and hoofed, lateral ones much smaller and not touching the ground.*

†* 26. SUS.—Tusks exerted, inclining both upwards and to one side: snout elongated, and truncated at the extremity: body covered with stiff bristles.

ORDER V. CETACEA.

Teeth various; sometimes entirely wanting: body fish-shaped: anterior extremities in the form of fins; posterior united, forming a flat horizontal tail.

* *Grinders with flat summits: nostrils at the extremity of the snout, not acting as blow-holes: whiskers very distinct: mammæ pectoral.*

(1.) MANATUS.—Grinders eight or nine on each side above and below; with two transverse ridges on the summits: no incisors or tusks in the adult state: vestiges of nails on the margins of the pectoral fins.

(2.) STELLERUS.—A single grinder on each side above and below; summit flat with projecting plates of enamel: fins without vestiges of nails.

*** Teeth, when present, conical: nostrils opening on the crown of the head, acting as blow-holes: whiskers none: mammae near the anus.*

† Head small: blow-holes united.

27. DELPHINUS.—Both jaws with numerous teeth, all simple and equal.

(1. DELPHINUS.) Snout produced into a beak, broad at its base, rounded at its extremity, and separated from the forehead by a kind of furrow: a dorsal fin.

(2. PHOCÆNA.) Snout short and blunt; not beaked: a dorsal fin.

(3. DELPHINAPTERA.) Head blunt; snout not produced: no dorsal fin.

28. MONODON.—Head blunt: no true teeth; upper jaw with one, rarely two, very long straight tusks projecting forwards in a line with the body: no dorsal fin.

29. HYPEROODON.—Snout produced: teeth generally but two in number, placed in front of the lower jaw: palate studded with tubercles: a dorsal fin.

†† Head very large; half or one third of the entire length.

30. PHYSETER.—Head enormously large, truncated in front: upper jaw without whalebone or visible teeth; teeth in the lower jaw numerous: blow-holes united.

(1. CATODON.) No dorsal fin.

(2. PHYSETER.) An elevated dorsal fin.

31. BALÆNA.—Head very large: palate furnished with whalebone: no teeth: blow-holes separate.

(1. BALÆNA.) No dorsal fin.

(2. BALÆNOPTERA.) A dorsal fin.

ORDER I. FERÆ.

† GEN. 1. URSUS, Linn.

† 1. U. *Aretos*, Linn. (*Brown Bear*).—Blackish brown: forehead convex above the eyes: snout suddenly tapering: soles of the hinder feet moderately long.

U. *Aretos*, *Desmar. Mammal.* p. 163. Common Bear, *Shaw, Gen. Zool.* vol. 1. p. 450. pl. 102.

DIMENS. Length of the head and body three feet seven inches; of the head one foot; of the fore foot seven inches seven lines; of the hinder foot eight inches ten lines.

DESCRIPT. Body entirely covered with thick shaggy hair, varying in colour from chestnut brown to black: soles of the fore feet with their anterior half naked; those behind naked throughout: ears short and rounded: eyes small: tail very short.

Formerly an inhabitant of Great Britain; but extirpated many centuries back. Infested Scotland (according to Pennant) so late as the year 1057. Still common on many parts of the European continent.

GEN. 2. MELES, Cuv.

2. M. *Taxus*, Flem. (*Badger*).—Gray above, black beneath; a black band on each side of the head extending from the nose over the eyes to behind the ears.

M. *Taxus*, *Flem. Brit. An.* p. 9. M. *vulgaris*, *Desm. Mammal.* p. 173. Badger, *Penn. Brit. Zool.* vol. 1. p. 85. pl. 8. no. 13. *Shaw, Gen. Zool.* vol. 1. p. 467. pl. 106.

DIMENS. Length of the head and body two feet six inches; of the head six inches eight lines; of the ears one inch four lines; of the tail seven inches.

DESCRIPT. Body thick and clumsy: hair rigid and very long; gray on the upper parts, black on the throat, breast, belly, and legs: head above white, with a longitudinal black spot on each side, which takes its origin between the extremity of the nose and the eye, and terminates behind the ear: toes five on each foot: claws long and bent: eyes very small: ears short and rounded, almost concealed in the hair: a transverse glandular pouch between the tail and the anus, secreting a fœtid substance.

Found in several parts of the kingdom, but not of very general occurrence. Burrows in the ground, concealing itself during the day, and coming abroad at night. Feeds indiscriminately on animal and vegetable substances.

GEN. 3. MUSTELA, Linn.

(1. MUSTELA, Cuv.)

3. *M. Foina*, Linn. (*Common Martin*.)—Brown : throat and breast white.

M. Foina, *Desm. Mammal.* p. 182. *Martes Fagorum*, *Flem. Brit. An.* p. 14. *Martin*, *Penn. Brit. Zool.* vol. i. p. 92. pl. 6. no. 15.

DIMENS. Length of the head and body eighteen inches; of the head four inches three lines; of the tail nine inches six lines.

DESCRIPT. Hair of two sorts; the shorter very fine and soft, and of a pale ash-colour; the longer somewhat rigid, less abundant than the last, ash-coloured at the roots, dusky brown towards the extremity, with a tinge of chestnut red: legs and tail dusky: under parts somewhat paler than the upper: a white spot upon the throat extending itself over the under surface of the neck and anterior portion of the breast.

More generally diffused than the next species, and said to prefer the vicinity of houses. Preys on poultry, game, rats, moles, &c. Breeds frequently in hollow trees, producing from three to seven young at a time.

4. *M. Martes*, Linn. (*Pine Martin*.)—Brown: throat and breast yellow.

M. Martes, *Desm. Mammal.* p. 181. *Martes Abietum*, *Flem. Brit. An.* p. 14. *Pine Martin*, *Penn. Brit. Zool.* vol. i. p. 94. *Don. Brit. Quad.* pl. 13.

DIMENS. Length of the head and body one foot seven inches; of the head four inches; of the tail ten inches.

DESCRIPT. Differs from the last species in having the throat, under surface of the neck, and anterior portion of the breast, yellow: the head also shorter, the hair rather darker, and the legs a little longer.

Inhabits the fir woods of Scotland: occurs also sparingly in the West of England. Builds its nest on the tops of trees, and produces from two to three young. Frequents wild situations. Preys on the smaller quadrupeds and birds.

(2. PUTORIUS, Cuv.)

5. *M. Putorius*, Linn. (*Polecat*.)—Dusky brown with a tinge of yellow; beneath paler: some white spots about the ears and mouth.

M. Putorius, *Desm. Mammal.* p. 177. *Flem. Brit. An.* p. 14. *Fitchet*, *Penn. Brit. Zool.* vol. i. p. 89. pl. 6. no. 14. *Polecat*, *Shaw, Gen. Zool.* vol. i. p. 415. pl. 98.

DIMENS. Length of the head and body one foot six inches; of the head two inches ten lines; of the ears six lines; of the tail five inches six lines.

DESCRIPT. Tail proportionably shorter than in the last sub-genus: space round the mouth, and edge of the ears, white: hair on the body of two kinds; the longer sort somewhat harsh, shining, and of a dusky brown colour: the shorter more woolly, and of a yellowish or tawny white; from the mixing of these two the general tint appears brown with a slight cast of tawny yellow: the legs and tail are of a uniform dusky brown.

A common inhabitant of woods and plantations in all parts of the country. Preys on game, poultry, eggs, and the smaller quadrupeds: is particularly fond of blood. Produces in the Spring from five to six young.

* 6. *M. Furo*, Linn. (*Ferret*).—Yellowish white with the eyes red.

M. Furo, *Desm. Mammal.* p. 178. *Ferret*, *Shaw, Gen. Zool.* vol. 1. p. 418. pl. 98.

DIMENS. Length of the head and body one foot two inches; of the head two inches six lines; of the ears six lines; of the tail five inches six lines.

DESCRIPT. In general somewhat smaller than the polecat, and of a more slender shape; the snout is also proportionably longer: colour of the fur bright yellow, here and there tinged with white; sometimes a mixture of white, black, and tawny, with the tail entirely black.

Originally a native of Africa, from whence imported into Europe. Known in this country only in a domesticated state, where it is much used in the destruction of rabbits and rats. Supposed by some to be a mere variety of the polecat; but independently of the differences above mentioned, the circumstance of its being a native of a warmer climate seems to militate against that idea. Habits similar to those of that species.

7. *M. vulgaris*, Gmel. (*Weasel*).—Reddish brown above; beneath white: tail of the same colour with the body.

M. vulgaris, *Desm. Mammal.* p. 179. *Flem. Brit. An.* p. 13. Common Weasel, *Penn. Brit. Zool.* vol. 1. p. 95. pl. 7. no. 17. *Shaw, Gen. Zool.* vol. 1. p. 420. pl. 98.

DIMENS. Length of the head and body (*male*) eight inches three lines, (*female*) seven inches; of the head (*male*) one inch nine lines, (*female*) one inch six lines; of the ears (*male*) four lines, (*female*) three lines; of the tail (*male*) two inches four lines, (*female*) two inches.

DESCRIPT. Upper part of the head, neck and body, shoulders, external and anterior portions of the fore legs, and hind legs wholly, reddish brown tinged with yellowish; the under parts, from the extremity of the lower jaw to the vent, white: a brown spot below each corner of the mouth.

Var. β . White, with a few black hairs at the extremity of the tail.

Common everywhere in the vicinity of barns and outhouses. Devours young birds, eggs, rats, mice, moles, &c. Breeds twice or thrice in the year, and produces from four to six young at a birth. The white variety is rare.—*Obs.* The female of this species is constantly much smaller than the male, and is probably the animal alluded to in White's *Nat. Hist. of Selb.* (vol. 1. p. 73.) under the name of *Cane*.

8. *M. Erminea*, Linn. (*Stoat or Ermine*.)—Reddish brown above, beneath white; or wholly white: extremity of the tail always black.

M. Erminea, *Desm. Mammal.* p. 180. *Flem. Brit. An.* p. 13. *Stoat*, *Penn. Brit. Zool.* vol. 1. p. 89. pl. 17. no. 18. *Shaw, Gen. Zool.* vol. 1. p. 426. pl. 99.

DIMENS. Length of the head and body ten inches; of the head one inch eleven lines; of the ears six lines; of the tail five inches.

DESCRIPT. Larger than the last species, to which it is closely allied.—(*Summer dress*.) Upper part of the head, neck, and back, as well as a considerable portion of the tail, reddish brown; under parts white tinged with yellow; tail terminating in a tuft of black hair.—(*Winter dress*.) Wholly white, or white with a slight tinge of yellow, the extremity of the tail excepted, which remains black.—*Obs.* In spring and autumn these two liveries are found intermixed.

Equally common with the weasel. Habits similar.

GEN. 4. LUTRA, Cuv.

9. *L. vulgaris*, Desm. (*Otter*.)—Deep brown: sides of the head, throat and breast, cinereous.

L. vulgaris, *Desm. Mammal.* p. 188. *Flem. Brit. An.* p. 16. *Otter*, *Penn. Brit. Zool.* vol. 1. p. 92. pl. 8. no. 19. *Shaw, Gen. Zool.* vol. 1. p. 437. pl. 100.

DIMENS. Length of the head and body two feet three inches; of the head five inches; of the ears eight lines; of the tail one foot four inches and a half: girth one foot four inches.

DESCRIPT. Head broad and flattened: muzzle obtuse: upper lip very thick and muscular, projecting over the lower: mouth rather small: whiskers strong, nearly three inches in length: eyes small, situate one inch behind the nostrils: ears short and rounded, almost hid in the fur: hair on the body of two kinds; the finer sort grayish white; the longer and coarser grayish white at the roots, deep brown at the extremity, the latter colour alone appearing externally: sides of the head, throat, under surface of the neck, and breast, cinereous: hair on the feet short, brown with a reddish tinge: tail dusky brown.

Inhabits the banks of rivers, lakes, and marshes. Swims and dives with great facility, and is destructive to fish, on which it preys. Breeds in March: goes with young nine weeks, and produces from four to five at a birth.

GEN. 5. CANIS, Linn.

(1. CANIS, *Flem.*)

* 10. *C. familiaris*, Linn. (*Dog*.)—Tail recurved.

C. familiaris, *Desm. Mammal.* p. 190. *Flem. Brit. An.* p. 10. *Dog*, *Penn. Brit. Zool.* vol. 1. p. 59. *Shaw, Gen. Zool.* vol. 1. p. 273. pl. 75.

DESCRIPT. Known only in the domesticated state. Varieties extremely numerous, offering every gradation of size, form, colour, and quality of the hair.

Primitive stock supposed to have been allied to the *Shepherd's Dog*, a race chiefly characterized by straight ears, long hair, and a bushy tail.

Feeds principally on animal substances, and is particularly attached to carrion. Female goes with young sixty-three days, and produces from six to twelve at a time. The puppies are born blind, and continue so for the first ten days.

† 11. *C. Lupus*, Linn. (*Wolf*).—Tail straight: eyes oblique.

C. Lupus, *Desm. Mammal.* p. 197. *Wolf*, *Penn. Brit. Zool.* vol. i. p. 75. pl. 5. *Shaw, Gen. Zool.* vol. i. p. 290. pl. 75.

DIMENS. Length of the head and body three feet nine inches; of the head ten inches six lines; of the ears four inches nine lines; of the tail one foot five inches.

DESCRIPT. Larger than the dog; the limbs stronger, and the body more muscular: muzzle sharp and pointed: eyes set obliquely: ears erect: tail straight and pendent; somewhat long and bushy: fur tawny or yellowish gray, with a black streak on the fore legs in the adult state.

Formerly abundant throughout Great Britain, but long since extirpated. Continued in Scotland till the year 1680, and in Ireland so late as 1710. Still found in most parts of the continent.

(2. VULPES, *Flem.*)

12. *C. Vulpes*, Linn. (*Fox*).—Tawny brown above; white beneath: ears externally black: tail thick and bushy, tipped with white.

C. Vulpes, *Desm. Mammal.* p. 201. *Vulpes vulgaris*, *Flem. Brit. An.* p. 13. *Fox*, *Penn. Brit. Zool.* vol. i. p. 71.

DIMENS. Length of the head and body two feet three inches; of the head six inches; of the ears four inches; of the tail one foot four inches: girth one foot one inch.

DESCRIPT. Muzzle sharp: head rather large, and somewhat flattened on the forehead: tusks relatively longer than in the dog, and more slender: ears erect and pointed: eyes oblique: tail very thick and bushy: fur thick and long, of a tawny or reddish brown colour: lips, lower jaw, forepart of the neck, abdomen, and inside of the thighs, white: back of the ears blackish brown; a streak of the same colour passing from the corner of each eye to the nose: extreme tip of the tail white. Varies occasionally in size and colour, as well as in the quality of the hair.

Frequents woods in the vicinity of farmyards and villages. Prowls about during the night, preying on poultry, small birds, rabbits, &c. Brings forth its young under ground, and produces from three to six at a birth.

GEN. 6. FELIS, *Linn.*

13. *F. Catus*, Linn. (*Wild Cat*).—Yellowish gray, with longitudinal and transverse bars of black: tail annulated with black; of equal thickness throughout.

F. Catus, *Desm. Mammal.* p. 232. *Flem. Brit. An.* p. 15. *Wild Cat*, *Penn. Brit. Zool.* vol. i. p. 80.

DIMENS. Length of the head and body (*male*) twenty-seven inches nine lines, (*female*) twenty-one inches three lines; of the tail (*male*) thirteen inches six lines, (*female*) twelve inches four lines.

DESCRIPT. Hair long and bushy; gray or brownish gray above, more or less tinged with yellow; beneath paler: a longitudinal black line down the middle of the back, with transverse bars of the same colour branching out laterally in parallel waves and extending over the flanks, shoulders, and thighs; several narrow parallel black lines on the top of the head between the ears: lips black: feet yellowish; the soles black: tail annulated with black; the tip entirely of that colour; of equal thickness throughout its whole length, and as it were truncated at the extremity.

Confined entirely to the northern parts of the island. Resides in woods: and climbs trees with great facility. Preys on birds and small quadrupeds. Goes with young fifty-six days, and produces from four to five at a birth.

* 14. *F. maniculata*, Rüpp. (*Domestic Cat*.)—Tail relatively longer than in the last species, and gradually tapering to a point.

F. maniculata, Temm. *Monog. de Mammal.* tom. i. p. 123. *Domestic Cat*, Penn. *Brit. Zool.* vol. i. p. 82.

DESCRIPT. Invariably smaller than the wild cat: the head larger in proportion to the body; the tail longer, more slender, and gradually terminating in a point: colours extremely variable.

I have followed Rüppel and Temminck in referring the domestic Cat to this species, which in a wild state inhabits the North of Africa, where it was discovered by the former of the two naturalists above mentioned. He supposes it to have been first reclaimed by the Egyptians.

GEN. 7. PHOCA, Linn.

15. *P. vitulina*, Linn. (*Common Seal*.)—Fur yellowish gray more or less marked and spotted with brown; whiskers with the bristles undulated.

P. vitulina, Desm. *Mammal.* p. 244. *Flem. Brit. An.* p. 17. *Common Seal*, Penn. *Brit. Zool.* vol. i. p. 137. pl. 12. *Shaw, Gen. Zool.* vol. i. p. 250. pl. 70.

DIMENS. Entire length from five to six feet, sometimes more.

DESCRIPT. Body elongated and somewhat conical, tapering gradually from the breast to the tail: neck very short: head large and round: muzzle broad, flat, and as it were truncated: upper lip furnished with long whiskers, the bristles of which have alternate contractions and dilatations so as to appear of unequal thickness: incisors six above and four below, the lower ones separated in pairs by an intermediate space; grinders all of the false kind, on each side five above and below: no external ears: feet very short; the toes, which are five in number, enveloped in a membrane; claws very strong, larger on the hind than on the fore feet: hair fine and close-set, abundant on the young: colour variable; the general tint yellowish gray with different shades of brown, usually deeper on the head and back than on the sides and abdomen: tail shorter than the hind feet.

Common on many parts of the coast, but prefers rocky shores. Swims and dives readily. Preys on fish, which it devours under the water. Breeds about Midsummer, and produces two young.—*Obs.* The *Pied Seal* of Pennant is probably only a variety of this species.

16. *P. barbata*, Mull. (*Great Seal*).—Fur blackish: middle toes on the forefeet longer than the lateral ones.

P. barbata, *Desm. Mammal.* p. 246. *Flem. Brit. An.* p. 18. *Great Seal*, *Penn. Brit. Zool.* vol. 1. p. 136. *Don. Brit. Quad.* pl. 11.

DIMENS. Entire length from ten to twelve feet.

DESCRIPT. Teeth as in the last species: head somewhat elongated: muzzle broad: upper lip rounded, fleshy, divided into two lobes by a deep furrow; each lobe furnished with numerous strong white bristles, semitransparent, and curled at the ends: eyes large; irides dark hazel: auricular apertures larger than in the last species: fore feet rather long, with the middle toe more developed than the lateral ones; claws of these black, horny, and curved; those of the hind feet long and straight: body elongated; when young, clothed with a long woolly fur, which after the space of fourteen or fifteen days is cast and superseded by a new covering of close short hair*: general colour dusky gray, in very old individuals black.

Inhabits the coast of Scotland, and (according to Selby) the Farn Islands, but is much less common than the last species. Breeds in November.

GEN. 8. TRICHECHUS, Linn.

17. *T. Rosmarus*, Linn. (*Walrus*.)

T. Rosmarus, *Desm. Mammal.* p. 253. *Flem. Brit. An.* p. 18. *Arctic Walrus*, *Shaw, Gen. Zool.* vol. 1. p. 234. pl. 68. *Nat. Miscell.* vol. viii. pl. 276.

DIMENS. Entire length from eleven to fifteen feet.

DESCRIPT. Head small, rounded, obtuse: lips very thick and swollen; upper lip divided into two large rounded lobes, over which are scattered numerous semitransparent bristles, somewhat flattened towards their roots, and slightly pointed at their extremities: eyes small and brilliant: auricular apertures situate very much behind: mouth small, armed with two enormous tusks bent downwards, and attaining in some individuals the length of two feet: neck short: body very thick and heavy: tail relatively longer than in the last genus: skin of a dusky hue, with a very few short, scattered, reddish hairs: hind feet very broad.

Very rare in the British seas. A solitary individual shot on the east coast of Harris in December 1817: a second killed in Orkney in June 1825. Habits resembling those of the Seals.

* See Selby's observations on this species in *Zool. Journ.* vol. 11. p. 465.

GEN. 9. TALPA, *Linn.*18. *T. Europæa*, *Linn. (Mole.)*

T. Europæa, *Desm. Mammal.* p. 160. *Flem. Brit. An.* p. 9. Mole, *Penn. Brit. Zool.* vol. i. p. 128. *Shaw, Gen. Zool.* vol. i. p. 515. pl. 117.

DIMENS. Length of the head and body five inches three lines; of the head one inch seven lines; of the tail one inch two lines.

DESCRIPT. Body thick, oblong, almost cylindrical: snout sharp and slender: eyes extremely small; entirely concealed in the fur: no external ears: feet extremely short; the anterior pair larger, very robust, and inclining sideways: claws strong, as long as the toes themselves. Fur very soft and silky, shining, black or deep ash-colour, according to the direction in which it is viewed, sometimes white, or yellowish white.

Common in England and Scotland, but said to be unknown in Ireland. Habits subterraneous: constructs galleries beneath the surface of the soil, which it throws up at intervals in hillocks. Feeds on worms and insects, but principally on the former. Breeds twice in the year, producing from four to six at each birth.

GEN. 10. SOREX, *Linn.*

19. *S. Araneus*, *Linn. (Common Shrew.)*—Reddish brown above; paler beneath: tail shorter than the body, somewhat square, not ciliated on its under surface.

S. Araneus, *Desm. Mammal.* p. 149. *Flem. Brit. An.* p. 8. Fetid Shrew, *Penn. Brit. Zool.* vol. i. p. 125. Common Shrew, *Shaw, Gen. Zool.* vol. i. p. 527. pl. 118.

DIMENS. Length of the head and body two inches five lines; of the head one inch; of the ears two lines; of the tail one inch nine lines.

DESCRIPT. Somewhat variable in size and colours. Upper parts generally dusky brown, more or less deep, with a tinge of red; under parts grayish white with a tinge of yellow: in some specimens a triangular whitish patch upon the throat: ears small, hardly showing themselves above the fur, furnished internally with two lobes or duplicatures of the skin placed one above the other and fringed with hair: incisors deep ferruginous brown*: tail varying in length, always shorter than the body, roundish approaching to square, rather stout, of equal thickness throughout and blunt at its extremity, uniformly clothed with short dusky hairs, but having no fringe along its under surface: feet much smaller than in the other species of this genus, the toes scarcely ciliated.

A very common species inhabiting gardens and hedge rows. Feeds on insects, and also on vegetable substances. Possesses a strong musky odour.

* In the *Ann. du Mus.* (tom. xvii. p. 176.) M. Geoffroy St Hilaire describes this species as having the incisors *entirely white*. This circumstance, together with one or two others, induces me to suspect that the *S. Araneus* of the continental authors may be distinct from ours.

20. *S. fodiens*, Gmel. (*Water Shrew*).—Blackish brown above; grayish white beneath: tail two-thirds the length of the body: feet and tail strongly ciliated with white hairs.

S. fodiens, *Flem. Brit. An.* p. 8. *S. Daubentonii*, *Desm. Mammal.* p. 150. *Water Shrew*, *Penn. Brit. Zool.* vol. i. p. 126. pl. 11. no. 33. *Don. Brit. Quad.* pl. 6.

DIMENS. Length of the head and body three inches two lines; of the head one inch; of the ears two lines; of the tail two inches.

DESCRIPT. Upper parts deep brown, approaching to black; under parts pale ash gray, in some individuals silvery white; in general the two colours separated by a well-defined line; a triangular dusky spot on the anus: snout long, somewhat depressed, emarginated at the extremity: whiskers long: eyes small, almost concealed: ears very short, not projecting above the fur, furnished internally with three valves or lobes, on one of which is a tuft of white hair, giving the effect of a white spot upon the auricle: intermediate incisor teeth with the tips ferruginous: tail more slender than in the last species, quadrangular throughout the greater part of its extent, the extreme tip being flattened; dusky, with a cilium of white hairs along its under surface: feet dusky; the toes fringed with white hairs. *Weight* three drachms.—*Obs.* English authors do not quite agree in their descriptions of this animal, from which circumstance it would seem that it is either subject to much variation, or that there is some other indigenous species with which it has been confounded.

Not uncommon in many parts of the country, inhabiting marshy districts. Swims and dives with great facility. Preys on insects: is also said to attack frogs. Produces in the Spring from six to eight young.

21. *S. remifer*, Geoff.? (*Oared Shrew*).—Black above; scarcely paler beneath; throat brownish ash: tail quadrangular at the base, flattened at the extremity, and, as well as the feet, ciliated.

S. remifer, *Geoff. Ann. du Mus.* tom. xvii. p. 182. pl. 2. f. 1? *Desm. Mammal.* p. 152? *Yarr. in Loud. Mag. of Nat. Hist.* vol. v. p. 598. *S. ciliatus*, *Sow. Brit. Misc.* pl. 49.

DIMENS. Length of the head and body three inches two lines; of the head one inch; of the ears two lines; of the tail two inches one line; of the fore foot five lines; of the hind foot eight lines and a half.

DESCRIPT. Body more thick and bulky for its length than in the last species: snout broad, and rather obtuse: feet and tail ciliated; the latter distinctly quadrangular for two-thirds of its length, compressed towards the tip, where the hairs of the cilium become longer. Colour darker, and more uniform than that of the *S. fodiens*; all the upper parts, sides of the abdomen, and region of the pubes, black; throat, breast, and middle of the abdomen, dusky ash, with a faint tinge of yellowish; sometimes, but not always, a white spot on the ears; incisors ferruginous at their extremities; feet and tail dusky gray, the latter with the cilium of hairs underneath of the same colour as above. *Weight* three drachms fifty-one grains.

This species, which has been taken in Norfolk, Cambridgeshire, and Surrey, as well as in Scotland, appears to be identical with a foreign one in the British Museum, ticketed *S. remifer*: nevertheless I feel some

Common in the Lake of Wight.

doubts as to its being the species originally described by Geoffroy under that name. It however approaches more nearly to it than to any other I am acquainted with. My specimen was taken in a corn-field at some distance from any water. The others have occurred in ditches. The ciliated feet and tail evidently mark it to be of aquatic habits.

GEN. 11. ERINACEUS, Linn.

22. *E. Europæus*, Linn. (*Hedgehog*.)—Ears short : spines moderately long.

E. Europæus, *Desm. Mammal.* p. 147. *Flem. Brit. An.* p. 7. Common Urchin, *Penn. Brit. Zool.* vol. i. p. 133. *Hedgehog*, *Shaw, Gen. Zool.* vol. i. p. 542. pl. 121.

DIMENS. Length of the head and body nine inches six lines; of the head two inches eleven lines; of the ears one inch one line; of the tail nine lines.

DESCRIPT. Body oblong, convex above, very low on the legs: head produced into a snout which is truncated at the extremity: nostrils narrow: ears short, broad, and rounded: neck thick and short: upper part of the body clothed with round sharp-pointed spines, barely an inch in length, which cross and interlace one another in all directions; the colour of them is whitish with a black ring a little higher up than the middle: snout, forehead, sides of the head, under part of the neck, breast, and legs, covered with coarse stiff hairs of a yellowish white colour.

Resides in hedges, thickets, &c. Habits nocturnal. Omnivorous; devouring roots, insects, worms, flesh, and even snakes. Becomes torpid during the Winter. Produces in the early part of the Summer from two to four young. Spines soft at birth, and all inclining backwards; become hard and sharp in twenty-four hours. In the adult state, has the power of rolling itself into a ball to avoid danger.

ORDER II. PRIMATES.

GEN. 12. RHINOLOPHUS, Geoff.

23. *R. Ferrum-equinum*, Gmel. (*Greater Horse-shoe Bat*.)—Posterior foliaceous appendage lanceolate, expanding laterally at the base: ears notched on their external margins.

R. Ferrum-equinum, *Flem. Brit. An.* p. 5. *R. unihastatus*, *Desm. Mammal.* p. 125. *Horse-shoe Bat*, *Penn. Brit. Zool.* vol. i. p. 147. pl. 14.

DIMENS. Length of the head and body two inches five lines; of the head eleven lines and a half; of the tail one inch two lines and a half; of the ears nine lines; breadth of the ears six lines; length of the thumb two lines and a half: extent of wing thirteen inches.

DESCRIPT. Upper incisors very small, separated from each other by a space; lower incisors each with three lobes: ears nearly as long as the head, somewhat triangular, broad at the base, terminating upwards in an

acute point; the external margin notched at the base, from which point it becomes inflexed and rises into an elevated round lobe that guards the orifice of the ear and appears to act the part of the tragus, which is wanting: nostrils placed at the bottom of a cavity close to each other, surrounded by a naked membrane in the form of a horse-shoe arising from the upper lip; anterior foliaceous appendage rising vertically immediately behind the nostrils, of a somewhat pyramidal form, sinuous at the margins and at the apex, which last is obliquely truncated; the posterior one situate on the forehead, placed transversely with respect to the first and standing more erect, lanceolate, expanding laterally at the base, in front of which are two small cup-shaped cavities formed by a duplicature of the skin. Colour of the fur reddish ash, inclining to gray beneath: membranes dusky: ears within and without slightly hairy.

A local species, inhabiting caves and buildings. Found in Bristol and Rochester Cathedrals, Dartford Powder Mills, and in some other parts of the country.

24. *R. Hipposideros*, Bechst. (*Lesser Horse-shoe Bat.*)
—Posterior foliaceous appendage lanceolate, without any lateral expansions: ears deeply notched on their external margins.

R. Hipposideros, Leach, *Zool. Misc.* vol. 111. p. 2. pl. 121. *Flem. Brit. An.* p. 5. *R. bihastatus*, Desm. *Mammal.* p. 125. *Vespertilio minutus*, Mont. in *Linn. Trans.* vol. ix. p. 163. pl. 8. f. 5.

DIMENS. Length of the head and body one inch four lines; of the head eight lines; of the tail nine lines; of the ears five lines; breadth of the ears four lines and a half; length of the thumb two lines: extent of wing eight inches four lines.

DESCRIPT. Principally distinguished from the last by its very inferior size. General appearance of the ears, nose, and foliaceous appendages similar; the anterior appendage is however less obliquely truncated at the apex, and the posterior one narrower at the base, and without the lateral expansions: the ears likewise are more deeply notched and the external margin altogether more sinuous. Fur soft and rather long; pale rufous brown above, grayish ash beneath with a tinge of yellow.

Found in Wiltshire, Dorsetshire, Devonshire, Somersetshire, and also in Wales. Is often met with in company with the last species, but is considered as less common.

GEN. 13. VESPERTILIO, Linn.

(1. VESPERTILIO, Geoff.)

25. *V. murinus*, Desm.—Ears oval, as long as the head; tragus falciform, half the length of the auricle: fur reddish brown above; dirty white beneath.

V. murinus, Desm. *Mammal.* p. 134. *La Chauve-souris*, Buff. *Hist. Nat.* tom. viii. pl. 16.

DIMENS. Length of the head and body three inches five lines; of the head eleven lines; of the tail one inch eight lines; of the ears eleven lines and a half; of the tragus five lines; of the thumb five lines: extent of wing fifteen inches.

DESCRIPT. Face almost naked: forehead very hairy: eyes rather large, with a few dusky hairs immediately above them: ears inclining backwards, as long as the head, oval, naked, grayish ash-colour externally, somewhat yellowish within: tragus falciform, about half the length of the auricle. Fur above pale reddish brown; beneath dirty white, inclining to yellowish: flying and interfemoral membranes brownish.

This must not be confounded with the *V. murinus* of English authors, from which it may be readily distinguished by its very superior size. It is a common species on the Continent, but apparently rare in Britain. The only indigenous specimens known were taken in the gardens of the British Museum, and are in that collection. Said to inhabit churches and buildings: never found in trees.

26. *V. Bechsteinii*, Leisl.—Ears oval, a little longer than the head; tragus falciform, bending outwards at the extremity, not half the length of the auricle: fur reddish gray above; white beneath.

V. Bechsteinii, *Desm. Mammal.* p. 135.

DIMENS. Length of the head and body two inches one line; of the head nine lines; of the tail one inch three lines; of the ears ten lines; of the tragus four lines; of the thumb four lines: extent of wing eleven inches.

DESCRIPT. Allied to the last species, but distinguished by its smaller size, relatively larger ears, and remarkably slender thumb. Face almost naked: muzzle long, and conical: ears oval, somewhat longer than the head, rounded at the extremity; tragus lanceolate, pointed, a little bent outwards towards the extremity. Fur reddish gray on the upper parts, whitish on the under.

A rare species, of which specimens are in the British Museum from the New Forest. Resides constantly amongst trees, and is never found in buildings.

27. *V. Nattereri*, Kuhl.—Ears oblong-oval, about as long as the head; tragus lanceolate, nearly two-thirds the length of the auricle: fur reddish gray above, white beneath: interfemoral ample, the margin crenate on each side towards the tip, and fringed with short bristly hairs.

V. Nattereri, *Desm. Mammal.* p. 135.

DIMENS. Length of the head and body one inch eleven lines; of the head eight lines and a half; of the tail one inch seven lines; of the ears eight lines and a half; of the tragus five lines; breadth of the ears three lines and three quarters; of the tragus at the base one line; length of the fore-arm one inch six lines; of the thumb two lines and three quarters: extent of wing ten inches eight lines.

DESCRIPT. Head rather small; snout attenuated: nose a line in breadth at the extremity, slightly emarginated between the nostrils, convex above: all the face, excepting immediately above the nose, hairy, but the hair very thinly scattered about the eyes and chin, with a few bristly hairs longer than the others intermixed: gape extending as far as the posterior angle of the eye: a row of longish hairs on the upper lip

forming a moustache: a large sebaceous gland above the eye advancing towards the nose, of a yellowish white colour; another gland on the upper lip immediately above the canine tooth, of the same colour with the rest of the face, causing a protuberant swelling at that spot, and furnished with long bristles: ears oblong-oval, rather broad, about the length of the head, the inner margin bending outwards, the external margin nearly straight, with a shallow notch about midway, nearly naked, excepting towards the base on the outside; tragus longer than in the last species, lanceolate, curving slightly outwards towards the tip, naked: flying and interfemoral membranes naked; the latter ample, furnished with a long spur*, between which and the end of the tail the margin has a crenate or puckered appearance, and is set with short bristly hairs; tip of the tail somewhat blunt and compressed, scarcely exerted for more than half a line: hind claws very strong, furnished with long hairs: thumb much smaller than in the last species. Fur long and silky, particularly about the upper part of the head and neck, of a light rufous brown, approaching to reddish gray above, the tips of the hairs being of this colour, the roots dusky brown; beneath silvery gray at the tips, black towards the roots: ears yellowish gray, the yellow tinge being most obvious within towards the base; tragus almost entirely yellowish: flying and interfemoral membranes dusky, but the latter paler than the former. In the *female*, the colour of the fur on the upper parts has a more reddish tinge than in the *male*.

This species occurs in hollow trees in the neighbourhood of buildings at Swaffham Prior in Cambridgeshire. Mr Yarrell has received it from Colchester and Norwich. There are also specimens in the British Museum taken near London. It appears however to be very locally distributed.

28. *V. Serotinus*, Gmel. (*Serotine*).—Ears oval-triangular, shorter than the head; tragus semicordate: muzzle short, and obtuse: fur chestnut brown above; pale beneath.

V. Serotinus, *Desm. Mammal.* p. 137. *La Sérotine*, *Buff. Hist. Nat.* tom. viii. pl. 18. f. 2.

DIMENS. Length of the head and body two inches seven lines; of the head ten lines; of the tail one inch ten lines; of the ears eight lines; of the tragus three lines; of the thumb three lines: extent of wing twelve inches six lines.

DESCRIPT. Face almost naked: muzzle remarkably short, broad, and obtuse; extremity of the nose one line and a half across: ears oval, approaching triangular, shorter than the head, with the inner margin bending outwards in an arcuate form, externally hairy on their basal half, naked above; tragus semicordate, somewhat elongated, pointed at the extremity. Fur in the *male* of a deep chestnut brown on the upper parts, passing beneath into a yellowish gray; that of the *female* said to be much brighter.

Apparently a rare species in Great Britain: has hitherto only occurred in the neighbourhood of London. Is said to frequent trees, and also houses occasionally. Habits somewhat solitary.

* By the *spur* I mean a long tendinous process from the heel of the foot which runs along the margin of the interfemoral membrane and serves to stretch it. It in fact represents the *os calcis*. It will be found of very different length in different species, varying from three to seven lines or more.

Abundant at Dover, (W Boner) and in the
Isle of Wight. Sandown &c. Feeds on a small
kind of Chafer. (Rev. C Bury).

In the Isle of Wight I have seen it in great numbers.

mononucleosis

29. *V. Noctula*, Gmel. (*Noctule*).—Ears oval-triangular, shorter than the head; tragus small, arcuate, terminating in a broad round head: fur short, of a uniform reddish brown above and below.

V. Noctula, *Desm. Mammal.* p. 136. *Flem. Brit. An.* p. 6. *La Noctule*, *Buff. Hist. Nat.* tom. VIII. pl. 18. f. 1. *Great Bat*, *Penn. Brit. Zool.* vol. 1. p. 146. pl. 13. no. 38.

DIMENS. Length of the head and body two inches eleven lines; of the head ten lines; of the tail one inch eight lines; of the ears seven lines and a half; of the tragus two lines and a half; breadth of the ears six lines; of the tragus one line and a half; length of the fore-arm two inches; of the thumb two lines and a half: extent of wing fourteen inches.

DESCRIPT. Head very broad; muzzle short and thick in the adult state, though less so than in the last species, somewhat elongated when young; nostrils tumid at the edges, slightly bilobated; forehead very hairy; rest of the face almost naked: ears shorter than the head, somewhat triangular, rounded at the extremity; the posterior margin folded back, with a projecting ridge internally, and a small protuberance at the base, which extends round nearly to the corners of the mouth; tragus very small, somewhat arcuate with the bend directed inwards, terminating above in a broad round head. Fur rather short, but soft and thick, of a uniform reddish brown colour above and below: membranes dusky, with a ridge of hair along the bones of the arm. Tail shorter than the fore-arm, protruding from the interfemoral membrane to the extent of a line and a quarter.

Common in many parts of the country, but only to be seen on wing during the summer months. Flight high and rapid. Habits gregarious. Retires early in the Autumn into hollow trees, and beneath the roofs of large buildings. Has a strong disagreeable smell.

30. *V. Leisleri*, Kuhl.—Ears oval-triangular, shorter than the head; tragus terminating in a round head: fur long, bright chestnut above, darker beneath: under surface of the flying membrane with a broad band of hair along the fore-arm.

V. Leisleri, *Desm. Mammal.* p. 138.

DIMENS. Length of the head and body two inches two lines; of the head seven lines and a half; of the tail one inch eight lines; of the ears five lines; of the tragus two lines and a half; breadth of the ears four lines; of the tragus one line and a quarter; length of the fore-arm one inch six lines and a half; of the thumb one line and three quarters: extent of wing eleven inches.

DESCRIPT. Muzzle rather more elongated than in the *Noctule*; nose depressed, naked, as is also the region of the eyes: ears oval-triangular, shorter than the head, broad, the outer basal margin advancing forwards to nearly the corners of the mouth; tragus terminating in a round head, bulging out about the middle of the external margin, slightly bending inwards at the extremity, considered relatively, somewhat larger and longer than in the last species. Fur on the upper parts bright chestnut, this

colour occupying the tips of the hairs, the roots of which are deep brown; under parts darker than the upper, the hair being grayish brown at the tips, dusky at bottom: flying membranes dusky; the portion contiguous to the body very hairy; a band of scattered hair also extends along and beneath the fore arm, about five lines in breadth: upper half of the interfemoral membrane both above and below likewise very hairy. Thumb short: bones of the arm and fore-arm much slenderer than in the *Noctule*.

The only indigenous specimen which I have seen of this species is in the British Museum. It is not known where it was taken. According to Desmarest it resides constantly in hollow trees in large companies, and is attached to the neighbourhood of stagnant waters.

31. *V. discolor*, Natt.—Ears shorter than the head, oval, bending outwards, with a projecting lobe on the inner margin; tragus of equal breadth throughout: fur above reddish brown, with the tips of the hairs white; beneath dirty white.

V. discolor, *Desm. Mammal.* p. 139.

DIMENS. Length of the head and body two inches four lines; of the head nine lines; of the tail one inch five lines; of the ears six lines and a half; of the tragus two lines and a quarter; of the thumb three lines: extent of wing ten inches six lines.

DESCRIPT. Forehead broad and hairy; muzzle long, and very broad; nose thick and obtuse, measuring one line and two-thirds across the extremity: eyes very small: ears shorter than the head, rounded, oval, bending outwards and reaching almost to the corners of the mouth, with a projecting lobe near the base of the internal margin, clothed externally on their lower half with thick woolly hair; tragus short, and nearly of equal breadth throughout. Fur on the back reddish brown, with the extreme tips of the hairs white, presenting a marbled appearance; underneath dirty white, with a large patch of a somewhat darker tint covering the breast and abdomen; throat pure white. Tail protruding from the interfemoral membrane to the extent of three lines.

A single individual of this species, taken at Plymouth by Dr Leach, is now in the British Museum. Said to be peculiar to houses, and never to be found in trees.

32. *V. Pipistrellus*, Gmel. (*Pipistrelle*).—Ears shorter than the head, oval-triangular, deeply notched on their external margins; tragus nearly straight, terminating in a blunt rounded head: fur reddish brown above; somewhat paler beneath.

V. Pipistrellus, *Desm. Mammal.* p. 139. *Jen. in Linn. Trans.* vol. xvi. p. 163. *Le Pipistrelle*, *Buff. Hist. Nat.* tom. viii. pl. 19. f. 1. Common Bat, *Penn. Brit. Zool.* vol. i. p. 148.

DIMENS. Length of the head and body one inch seven lines; of the head six lines; of the tail one inch two lines; of the ears four lines; of the tragus two lines; breadth of the ears three lines; of the tragus three-

quarters of a line; length of the fore-arm one inch two lines; of the thumb one line and three quarters: extent of wing eight inches four lines.

DESCRIPT. Resembling the *Noctule* in many of its characters but much smaller: head depressed in front, convex behind; muzzle short in the adult state, somewhat elongated when young; nose obtuse at the extremity, and slightly emarginated between the nostrils; a protuberant swelling on each side of the face above the upper lip formed by a congeries of sebaceous glands: eyes very small; above each an elevated wart furnished with a few black hairs: ears broad, oval-triangular, rather more than half as long as the head, with their external margins deeply notched about midway down: tragus half the length of the auricle, nearly straight, oblong, with a blunt rounded head: tail as long as the fore-arm. Fur rather long and silky, yellowish red on the forehead and at the base of the ears, on the rest of the upper parts reddish brown, with the lower half of each hair dusky; on the under parts wholly dusky, except the extreme tips of the hairs which are of the same colour as above, but paler. Young specimens generally brownish gray, sometimes black, without any tinge of red. Nose, lips, ears, and membranes dusky.

The most common species in this country, although for a long time confounded with the *V. murinus* described above. Congregates in large numbers in the crevices of old walls, decayed door-frames, &c. Resists cold more than the other species, continuing on wing till near the end of the year. Is first seen about the beginning of March.

33. *V. pygmæus*, Leach. (*Pygmy Bat*).—Forehead with an impressed longitudinal furrow: ears a little shorter than the head, broad at the base, obtuse and rounded at the extremity; tragus linear: fur above brown, darkest on the head and dorsal line; paler beneath.

V. pygmæus, Leach, in *Zool. Journ.* vol. i. p. 560. pl. 22.

DIMENS. Length of the head and body one inch two lines and a half; of the head five lines; of the tail nine lines; of the ears four lines: extent of wing five inches four lines.

DESCRIPT. Differs from the last species in its smaller dimensions, as well as in its relative proportions. Head high; the forehead marked with a longitudinal furrow; muzzle short and obtuse, nearly of equal breadth throughout; nostrils small, opening laterally: ears somewhat shorter than the head, broad at the base, obtuse and rounded at the extremity; anterior margin nearly straight, posterior slightly concave and convolute: tragus about half the length of the auricle, regularly linear, simple and rounded at the extremity: tail as long as the body exclusively of the head; the tip naked, protruding one line beyond the interfemoral membrane. Fur short and delicate, dark brown on the upper parts, the colour being deepest on the head and on the highest part of the back along the spine; inclining to gray underneath: flying membrane dark brown.—LEACH.

A new species discovered by Dr Leach at Spitchweek, near the Forest of Dartmoor, where it is said to be extremely common. Has not hitherto occurred elsewhere.

34. *V. emarginatus*, Geoff.—Ears oblong, nearly as long as the head, deeply notched on their external margins; tragus half the length of the auricle, subulate, bending slightly inwards towards the tip: fur reddish gray above, ash-colour beneath.

V. emarginatus, Geoff. in *Ann. du Mus.* tom. viii. p. 198. pl. 46.
Desm. Mammal. p. 140.

DIMENS. Length of the head and body two inches; of the head eight lines; of the tail one inch five lines; of the ears six lines; of the tragus three lines; breadth of the ears three lines and three quarters; of the tragus at the base three quarters of a line; length of the fore-arm one inch four lines; of the thumb two lines and three quarters: extent of wing nine inches six lines.

DESCRIPT. Bearing some relation to the *Pipistrelle* in its physiognomy, but distinguished from that species by its somewhat superior size. Muzzle rather obtuse; head flattish; face hairy, but the hair on the nose and chin thinly scattered and longer than on the other parts; a moustache of soft longish hair on the upper lip, above which is a congeries of glands extending from the eyes to the nostrils: ears moderate, a little shorter than the head, oblong approaching triangular, rounded at the extremity, the inner margin bending outwards, the outer one sinuate and rather deeply notched about half way down; tragus half the length of the auricle, linear approaching to subulate, nearly straight but having a slight bend inwards: thumb longer and stouter than in the *Pipistrelle*: hind feet remarkably large and strong, much more so than in either the *Pipistrelle* or the next species, the toes set with bristly hairs: interfemoral moderately ample; the spur rather short: tail a little longer than the fore-arm, exerted for about one line. Fur long; the hair on the upper parts with the basal half dusky, the tips reddish brown, this last colour alone appearing externally; beneath black at the roots, ash-gray at the tips: ears and flying membrane dusky; under surface of the interfemoral whitish.

Stated to have been found near Dover by M. A. Brongniart, and in Fifeshire by Dr Fleming. The description given above is from a specimen in my possession taken at Milton Park in Northamptonshire. It may possibly be distinct from the true *V. emarginatus* of the continental authors, but it appears to approach nearer to that species than to any other I am acquainted with. Mr Yarrell has also a specimen which was taken at Islington.

35. *V. mystacinus*, Leisl.—Ears oblong, shorter than the head, bending outwards and notched on their external margins; tragus half the length of the auricle, lanceolate, straight: a moustache of fine close hairs on the upper lip: fur blackish chestnut above, dusky ash beneath.

V. mystacinus, *Desm. Mammal.* p. 140.

DIMENS. Length of the head and body one inch eight lines; of the head seven lines and a half; of the tail one inch five lines; of the ears five lines and a half; of the tragus three lines; breadth of the ears three lines and a half; of the tragus at the base one line; length of the fore-

I have taken one here (Keymer).

arm one inch three lines; of the thumb two lines and a half: extent of wing eight inches six lines.

DESCRIPT. Head small and flattish; muzzle short; nose swollen, with a shallow cleft in the middle; face much more hairy than in the last species, with a few scattered hairs on the nose and chin, longer than the rest, intermixed; a row of fine, soft, close-set hairs on the upper lip forming a conspicuous moustache; a similar row crossing the forehead: ears shorter than the head, moderately broad, oblong, rounded at the extremities, curving outwards and rather deeply notched on their external margins; tragus half or rather more than half the length of the auricle, lanceolate, perfectly straight, narrowing regularly from the base upwards to the tip which terminates in a sharp point: thumb moderate: hind feet much smaller than in the last species: interfemoral with the spur of about the same length: tail longer than the fore-arm, exserted for about one line. Fur long, thick and woolly; hair dusky, approaching to black, throughout the greater part of its length, the extreme tips being reddish brown on the upper parts and ash-gray beneath: ears, flying and interfemoral membranes dusky; this last sometimes transversely marked on its under surface with numerous white ciliated lines.

This species occurs, though rarely, in houses in Cambridgeshire. I have also received specimens from Milton Park in Northamptonshire. Mr Yarrell has others which were taken at Colchester in the caverns under the castle. Said to frequent the neighbourhood of water, and to retire into hollow trees and houses.

(2. PLECOTUS, Geoff.)

36. *V. auritus*, Linn. (*Greater Long-eared Bat*).—

Ears more than double the length of the head; tragus oval-lanceolate: fur brownish gray on the upper parts; paler beneath.

V. auritus, *Desm. Mammal.* p. 144. *Long-eared Bat*, *Penn. Brit. Zool.* vol. 1. p. 147. pl. 13. no. 40. *Shaw, Gen. Zool.* vol. 1. p. 123. pl. 40.

DIMENS. Length of the head and body one inch ten lines; of the head eight lines; of the tail one inch eight lines; of the ears one inch five lines; of the tragus seven lines; breadth of the ears nine lines; of the tragus two lines and a half; length of the fore-arm one inch five lines; of the thumb two lines and three quarters: extent of wing ten inches two lines.

DESCRIPT. Head and face flattened; muzzle somewhat swollen about the nose; nostrils with their anterior and inner edges tumid, elongated posteriorly into a sort of *cul de sac*: eyes small: ears extremely large, more than twice the length of the head, oblong-oval, thin, and semi-transparent; the inner margin presenting a broad longitudinal fold, which doubles back nearly at right angles to the rest of the auricle and is ciliated with hair along its external and internal edges; near the base of this fold is a small projecting lobe, also ciliated; tragus long, oval-lanceolate, with the outer margin somewhat sinuous, the inner one straight: ears united over the head to the height of one line and a half; extending round at the base to the corners of the mouth: flying and interfemoral membranes broad and ample: tail longer than the fore-arm, the tip obtuse, protruding to the extent of three quarters of a line: forehead, and anterior surface of the connecting membrane of the ears, hairy; posterior

surface of the same membrane naked. Fur long and silky, brownish gray on the upper parts, paler beneath; the hair every where dusky at the roots.

A common and generally diffused species, resorting principally to the roofs of houses and churches. Flight swifter than that of the Pipistrelle. In the living animal the ears are generally curled, the bend being directed outwards. when at rest they are sometimes wholly concealed beneath the fore-arm, the tragus alone remaining erect.

37. *V. brevimanus*, Jenyns. (*Lesser Long-eared Bat*.)
—Ears not double the length of the head; tragus oval-lanceolate: fur reddish brown on the upper parts; yellowish white beneath.

V. brevimanus, *Linn. Trans.* vol. xvi. p. 55. pl. 1. f. 2.

DIMENS. Length of the head and body one inch six lines; of the head seven lines; of the tail one inch two lines; of the ears one inch; of the tragus five lines and a half; breadth of the ears five lines; of the tragus two lines; length of the fore-arm one inch two lines; of the thumb three lines: extent of wing six inches six lines.

DESCRIPT. Similar to the last in general appearance, but much smaller, and differing in its relative proportions: ears shorter with respect to the head, and rather narrower at the extremity; tragus relatively longer: bones of the hands much shorter: thumb somewhat longer: tail the length of the fore-arm; the tip more exerted from the interfemoral membrane, and terminating in a fine point. Fur on the upper parts reddish brown, presenting a marked contrast with that on the under which is yellowish white; the hair everywhere of the same colour throughout its whole length, and not dusky at the roots as in the last species: ears and membranes dusky with a tinge of red.

A new species first described by myself in the Linnæan Transactions as above quoted. Only one individual has as yet occurred, which is a female. It was found adhering to the bark of a pollard-willow, in Grunten Fen in the Isle of Ely.

(3. BARBASTELLUS, Gray.)

38. *V. Barbastellus*, Gmel. (*Barbastelle*.)—Ears shorter than the head, broad, triangular, notched on their external margins; tragus semicordate: fur black, with the tips of the hairs grayish white.

V. Barbastellus, *Desm. Mammal.* p. 145. *Sow. Brit. Misc.* pl. 5. *Mont. Linn. Trans.* vol. ix. p. 171.

DIMENS. Length of the head and body two inches; of the head seven lines and a half; of the tail one inch nine lines; of the ears five lines; of the tragus three lines and a half; breadth of the ears five lines; of the tragus one line and a half; length of the fore-arm one inch four lines and a half; of the thumb two lines and a half: extent of wing ten inches three lines.

DESCRIPT. Muzzle short and obtuse, somewhat swollen at the extremity; a naked space above the nose extending up to the ears, sunk and hollowed out in front, with the nostrils placed in the cavity; on each

I have taken 3 specimens from a low summer house thatched with heath, in my grandfather's garden at Henfield Sussex. Several others are in the collection of my Uncle W. Borser all taken from the house or out-houses in the same garden.

side of the face above the upper lip a large protuberant swelling covered with black hair: eyes very small, placed within the auricle and almost concealed: ears united over the head, oval-triangular, as broad as long, externally hairy, and notched on their outer margins; tragus semicordate, very protuberant near the base of the external margin, above which it suddenly bends inward and terminates in a point; inner margin straight. Fur very long and silky, particularly on the top of the head behind the ears, blackish brown, with the tips of the hairs silvery gray on the upper parts, and ash-coloured beneath; flying and interfemoral membranes dusky.

A rare species. Has occurred in Devonshire, Kent, Northamptonshire, and Cambridgeshire. Habits more diurnal than usual with this family. Flight slow and near the ground. Resorts to buildings for retirement.

ORDER III. GLIRES.

GEN. 14. SCIURUS, *Linn.*

39. *S. vulgaris*, *Linn.* (*Common Squirrel.*) — Fur brownish red above; white beneath: ears with a pencil of long hairs at the extremity.

S. vulgaris, *Desm. Mammal.* p. 330. *Flem. Brit. An.* p. 20.
Common Squirrel, *Penn. Brit. Zool.* vol. i. p. 107.

DIMENS. Length of the head and body nine inches; of the head two inches one line; of the ears nine lines and a half; of the tail (to the end of the bone) six inches six lines, (to the end of the hair) eight inches six lines.

DESCRIPT. Head thick, with the cheeks rather flattened; nose prominent; the upper lip projecting considerably beyond the lower; the first grinder in the upper jaw extremely small, consisting of a single tubercle, and disappearing altogether at a certain age; eyes black, large, and round; ears erect, moderately large, ornamented at the extremity with a tuft or pencil of long hairs: neck short; legs strong and muscular, much longer behind than before; fore feet with four long, deeply divided toes, and a claw in the place of a thumb; hind feet with five toes; claws sharp and strong: tail long and bushy, with the hair spreading out laterally. Colour of the upper parts reddish, or bright chestnut brown, the red tint being deepest on the sides of the head and neck, the shoulders, and the external surface of the legs; lower parts, including the under portion of the neck, breast, abdomen, and inside of the legs, white: tail the colour of the back.

Common in extensive woods, residing and building in trees. Climbs and leaps with great agility. Feeds on buds, acorns, nuts, and other fruits. When at rest, often sits erect, using its fore feet as hands, with its tail turned back over the body. Produces in the Spring from three to four young.

GEN. 15. MYOXUS, *Gmel.*

40. *M. avellanarius*, *Desm. (Dormouse.)*—Fur tawny red above; white beneath: tail bushy, the length of the body.

M. avellanarius, *Desm. Mammal.* p. 295. *Flem. Brit. An.* p. 22. *Dormouse, Penn. Brit. Zool.* vol. i. p. 110. *Shaw, Gen. Zool.* vol. ii. p. 167. pl. 154.

DIMENS. Length of the head and body two inches ten lines; of the head eleven lines and a half; of the ears four lines; of the tail two inches eight lines.

DESCRIPT. Head broad; nose blunt; eyes black, large, and prominent; ears broad, oval, rounded at the extremity: body plump: hind legs not disproportionably longer than the fore; toes four in front with the rudiment of a thumb, five behind: tail very long, somewhat flattened horizontally, thickly clothed with hair on every side. Colour of the upper parts bright tawny red; under parts white, with here and there a yellow tinge: tail red throughout.

Not uncommon in some parts of the country, frequenting woods and thick hedges. Less active than the squirrel: food similar. Builds in the hollows of trees, and produces from three to four young. Becomes torpid in the Winter.

GEN. 16. MUS, *Linn.*

41. *M. sylvaticus*, *Linn. (Field Mouse.)*—Fur yellowish brown above; whitish beneath, with a ferruginous spot on the breast: ears more than half the length of the head.

M. sylvaticus, *Desm. Mammal.* p. 301. *Flem. Brit. An.* p. 19. *Field Mouse, Penn. Brit. Zool.* vol. i. p. 120. *Wood Mouse, Shaw, Gen. Zool.* vol. ii. p. 58. pl. 132.

DIMENS. Length of the head and body three inches nine lines; of the head one inch one line; of the ears seven lines; of the tail three inches seven lines.

DESCRIPT. Forehead somewhat convex; nose blunt; eyes black, large and prominent; ears oblong-oval, with the anterior margin doubled in at the base, and a projecting lobe opposite to it, this last arising within the auricle near the base of the posterior margin; whiskers very long, measuring one inch two lines. Upper part of the head and body, cheeks, sides, and external portion of the legs, of a tawny gray or yellowish brown colour, each hair being dusky ash from the root upwards throughout two-thirds of its length, then tawny yellow, and lastly (more particularly some longer ones than the others) black at the extremity; under part of the head and body, and inside of the legs, whitish, with here and there a tinge of dusky ash, the hair on these parts being dusky from the root upwards the same as above, but the extremities white; the line of separation between the above colours is tolerably well defined, and the red tinge is most prevalent just at that part; there is also a faint tawny spot upon the breast: tail a little shorter than the body, more slender and tapering

than in the *M. Musculus*; dusky on the upper surface, whitish beneath: toes furnished above with long white hairs extending beyond the claws.

Common in gardens; and is occasionally, though very rarely, found in houses. A larger variety, measuring four inches and a half in length, exclusively of the tail, which is four inches, is sometimes met with in woods. Resides in holes under ground, where it amasses seeds, roots, &c. on which it feeds. Produces nine or ten young at a litter.

42. *M. messorius*, Shaw. (*Harvest Mouse*.)—Fur tawny brown on the upper parts; beneath white: ears one-third the length of the head.

M. messorius, *Desm. Mammal.* p. 302. *Flem. Brit. An.* p. 19.
Harvest Mouse, *Shaw, Gen. Zool.* vol. II. p. 62. *Fig. on Frontisp.*

DIMENS. Length of the head and body two inches six lines; of the head ten lines; of the ears three lines; of the tail two inches six lines.

DESCRIPT. Much smaller than the last species, and of a more elongated form: head narrower in proportion: eyes and ears smaller, the latter barely one-third the length of the head, with the projecting lobe at the base of the posterior margin relatively larger: tail varying in length, equalling the body, or somewhat shorter. Fur above of a bright tawny red colour, with a tinge of brown, the basal half of each hair being dusky, and the upper portion red; beneath almost pure white, the hair being of one tint throughout its whole length: a distinct line of separation between the above colours running parallel with the sides.

First found in Hampshire by White. Since met with in Wiltshire, Devonshire, Cambridgeshire, and other parts of the country. Frequents cornfields during the harvest. Nest round, composed of dry straws and grasses, and suspended at a small height from the ground. Produces seven or eight at a litter.

43. *M. Musculus*, Linn. (*House Mouse*.)—Fur dusky gray above with a tinge of yellow; beneath cinereous: ears about half the length of the head.

M. Musculus, *Desm. Mammal.* p. 301. *Flem. Brit. An.* p. 19.
Common Mouse, *Penn. Brit. Zool.* vol. I. p. 122. pl. 11. no. 30.
Shaw, Gen. Zool. vol. II. p. 56. pl. 131.

DIMENS. Length of the head and body three inches two lines; of the head eleven lines; of the ears five lines; of the tail two inches eleven lines.

DESCRIPT. Head of a somewhat triangular form: muzzle pointed, sharper than in the *M. sylvaticus*: eyes and ears, especially the former, smaller than in that species; the latter oval, rounded at the extremity, and narrower in proportion to their length: whiskers shorter: tail stouter: feet and toes, particularly those behind, shorter. All the upper parts of the body and sides, dusky gray with a faint tinge of yellow, this last colour occupying a small portion of each hair near the summit, the basal half and extreme tip being dusky; under parts cinereous, paler than those above, likewise tinged with yellow: ears, feet, and tail, clothed with a very short fine hair; the last dusky, of one colour above and

below.—*Obs.* The colours vary a little in different specimens, and sometimes individuals occur wholly white; or white with dark spots.

Found in houses and stacks. Those which occur in the latter situation are remarkable for the brightness of their colour, which has a more decided yellow tinge than in house specimens. They also sometimes attain a larger size, measuring nearly four inches in length. Feeds on various animal and vegetable substances. Breeds several times in the year, and is extremely prolific.

44. *M. Rattus*, Linn. (*Black Rat*).—Fur grayish-black above; ash-colour beneath: tail a little longer than the body.

M. Rattus, *Desm. Mammal.* p. 300. *Flem. Brit. An.* p. 20. Black Rat, *Penn. Brit. Zool.* vol. i. p. 113. *Shaw, Gen. Zool.* vol. 11. p. 52. pl. 130.

DIMENS. Length of the head and body seven inches four lines; of the head one inch ten lines; of the ears eleven lines and a half; of the tail seven inches eleven lines.

DESCRIPT. Head long; muzzle sharp-pointed; lower jaw very short; eyes large and prominent; ears oval, broad, and naked; whiskers long; fore feet with four toes, and a claw in the place of a thumb; hind feet with five; tail longer than the body, almost entirely naked, and covered with small scales disposed in rings. Colour of the upper parts deep iron-gray, or grayish black; those beneath dull ash-colour: feet and tail dusky.

Frequents houses, and is truly indigenous; but is now much less general than the next species. Food various. Breeds frequently in the year, producing from six to eleven at a time.

* 45. *M. decumanus*, Pall. (*Brown Rat*).—Fur above grayish brown, tinged with yellow; beneath whitish: tail scarcely so long as the body.

M. decumanus, *Desm. Mammal.* p. 299. *Flem. Brit. An.* p. 20. Norway Rat, *Penn. Brit. Zool.* vol. i. p. 115. *Shaw, Gen. Zool.* vol. 11. p. 51. pl. 130.

DIMENS. Length of the head and body eleven inches; of the head two inches four lines and a half; of the ears eight lines and a half; of the tail eight inches four lines.

DESCRIPT. Larger than the last species; thicker and of a stronger make: muzzle not quite so sharp at the extremity: eyes large and prominent: ears as broad as long, rounded at the extremity, and almost naked: tail naked and scaly, with about one hundred and eighty rings, generally not quite so long as the body, sometimes equalling it in length. Hair on the upper parts dusky ash at the roots, and reddish yellow at the extremity, with longer hairs intermixed of a deep brown colour throughout their whole length; general resulting tint grayish brown with a cast of tawny yellow; under parts dirty white, inclining here and there to ash-colour.

Introduced originally by shipping from Asia: unknown in England before the year 1730. Abundant now throughout the country, inhabiting houses, barns, granaries, drains, and other situations. Habits similar to those of the last species. Takes occasionally to the water, and swims readily.

GEN. 17. ARVICOLA, *Lacép.*

46. *A. amphibia*, Desm. (*Water Campagnol.*)—Fur above dusky gray tinged with yellow; beneath paler: tail more than half the length of the body.

A. amphibia, Desm. *Mammal.* p. 280. *A. aquatica*, *Flem. Brit. An.* p. 23. *Water Rat*, *Penn. Brit. Zool.* vol. i. p. 118. *Shaw, Gen. Zool.* vol. ii. p. 73. pl. 129.

DIMENS. Length of the head and body eight inches four lines; of the head one inch ten lines; of the ears five lines and a half; of the tail four inches nine lines.

DESCRIPT. Head thick and blunt; eyes small; ears short, scarcely projecting beyond the fur; incisors of a deep yellow colour, very strong, and measuring in the lower jaw half an inch in length: fore feet with four toes and the rudiment of a thumb; hind feet with five; these last connected for a little way at their base, but not regularly webbed: tail more than half as long as the body, covered with short scattered hairs. Fur very thick, grayish black above, or reddish brown with scattered black hairs; beneath iron gray.

Var. β. *A. ater*, *Macgillivray in Wern. Mem.* vol. vi. p. 429. Fur of a deep black colour above and below.

Frequents the banks of rivers and ditches, in which it burrows. Swims and dives readily. Feeds principally on roots and tender plants. Produces in June from six to eight young. The black variety is not uncommon in the fens of Cambridgeshire: it differs in no respect but that of colour.

47. *A. agrestis*, *Flem.* (*Field Campagnol.*)—Fur reddish brown above; pale gray beneath: tail one-third the length of the body.

A. agrestis, *Flem. Brit. An.* p. 23. *A. vulgaris*, Desm. *Mammal.* p. 282. Short-tailed Mouse, *Penn. Brit. Zool.* vol. i. p. 123. Meadow Mouse, *Shaw, Gen. Zool.* vol. ii. p. 81. pl. 136.

DIMENS. Length of the head and body four inches one line: of the head one inch two lines; of the ears five lines; of the tail one inch three lines and a half.—*Obs.* These dimensions are rather larger than usual.

DESCRIPT. Resembles the last species, but is much smaller. Head large; muzzle blunt at the extremity; ears rounded, moderately projecting, relatively longer than in the *A. amphibia*; incisors white: tail not more than one-third the length of the body, sometimes less. Fur soft and silky, reddish brown mixed with dusky on the upper parts, grayish ash beneath.

Common in meadows, preferring moist situations. Food seeds and roots. Is very prolific; and in certain seasons multiplies to a great extent. Nest constructed of dried grass.

48. *A. riparia*, Yarr. (*Bank Campagnol.*)—Fur bright chestnut-red above, ash-colour beneath: tail half the length of the body; the hairs at the tip a little elongated.

A. riparia, *Yarrell in Proceed. of Zool. Soc.* (1832) p. 109. *Id. in Loud. Mag. of Nat. Hist.* vol. v. p. 598.

DIMENS. Length of the head and body three inches four lines; of the head one inch half a line; of the ears five lines; of the tail one inch eight lines.

DESCRIPT. Distinguished from the *A. agrestis* by its brighter and more rufous colour; tail longer, with the hairs at the tip extending beyond the bone; ears rather larger, and more prominent: under parts cinereous, with a faint yellow tinge along the mesial line of the abdomen and on each side of it: tail blackish above, white beneath, the two colours separated by a well-defined line.

First discovered by Mr Yarrell at Birchanger in Essex. Has since occurred in Hertfordshire, Middlesex, Berkshire, and Cambridgeshire, but is not so plentiful as the *A. agrestis*. Frequents hedge-bottoms and ditch-banks, also occasionally stacks of corn. Is said to construct its nest of wool.—*Obs.* Independently of the above external differences between this and the last species, there are others connected with their anatomy, which will be found detailed at length in *Loud. Mag.* l. c.

† GEN. 18. CASTOR, Linn.

† 49. *C. Fiber*, Linn. (*Beaver.*)—Fur of a deep chestnut-brown; glossy above, dull beneath.

C. Fiber, *Desm. Mammal.* p. 277. *Common Beaver*, *Shaw, Gen. Zool.* vol. 11. p. 30. pl. 123.

DIMENS. Length of the head and body two feet six lines; of the head five inches; of the tail one foot; breadth of the tail four inches two lines.—**DESM.**

DESCRIPT. Head short and thick, somewhat flattened at the top; muzzle obtuse: eyes small and black: ears short, and rounded at the extremity: neck short: body thick, very convex on the back: tail depressed, broad, and of an oval form; the surface naked and scaly. Fur of a deep chestnut-brown colour, smooth and glossy on the upper parts, more dull underneath.

Formerly an inhabitant of Wales and Scotland. Observed in the former country by Giraldus de Barri in the year 1188. Not exactly known when the species was extirpated. Has occurred in a fossil state in Cambridgeshire.

GEN. 19. LEPUS, Linn.

50. *L. timidus*, Linn. (*Common Hare.*)—Tawny gray, shaded with brown: ears longer than the head; black at the tips: tail black above, white beneath.

L. timidus, *Desm. Mammal.* p. 347. *Flem. Brit. An.* p. 21. *Hare*, *Penn. Brit. Zool.* vol. 1. p. 98. *Shaw, Gen. Zool.* vol. 11. p. 197. pl. 162.

DIMENS. Length of the head and body twenty-one inches nine lines; of the head four inches; of the ears four inches ten lines; of the tail three inches six lines.

DESCRIPT. Head thick and large: inside of the cheeks hairy; eyes placed laterally, large and prominent; ears longer than the head: limbs slender, much longer behind than before: soles of the feet hairy. Fur composed of a fine down with longer hairs intermixed, of a tawny gray or rusty brown colour, the red tint prevailing in certain parts more than in others: each individual hair gray at the roots, black in the middle, and tawny at the tip; abdomen, inside of the thighs, and a transverse patch beneath the lower jaw, white; ears externally cinereous towards the base of the outer margin, above that colour a black spot reaching to the extremity of the auricle: tail black above, white underneath.

Var. β. Irish Hare. *Yarr. in Proceed. of Zool. Soc.* (1833) p. 88. Head shorter and more rounded than in the *Common Hare*; ears shorter, not equalling the head in length; limbs less lengthened; fur composed of only one sort of hair, the long dark hairs, observable in the English Hare, being wanting.

Frequents chiefly open fields. Feeds entirely on vegetables, coming abroad in the evening for that purpose. Breeds frequently in the year. Goes with young thirty days, and produces from one to four, rarely five, at a time. Young born with their eyes open, and the body clothed with fur. When full grown, has been known to weigh thirteen pounds one ounce and a half.

Var. β, which is the only Hare found in Ireland, might almost deserve to be considered as a distinct species. From the shortness and inferior quality of the hair, its fur is useless in trade.

51. *L. Cuniculus*, Linn. (*Rabbit*.)—Brownish gray, mixed with tawny: ears scarcely longer than the head: tail brown above, white beneath.

L. Cuniculus, *Desm. Mammal.* p. 348. *Flem. Brit. An.* p. 21.
Rabbit, Penn. Brit. Zool. vol. i. p. 104. pl. 10. no. 22. *Shaw*,
Gen. Zool. vol. ii. p. 204. pl. 162.

DIMENS. Length of the head and body sixteen inches; of the head three inches four lines: of the ears three inches six lines; of the tail three inches.

DESCRIPT. Approaching the last species in general appearance, but with the ears and hind legs proportionably shorter: fur of a less ferruginous colour; the general tint brownish gray, or dusky brown, with the nape reddish; throat and abdomen white; ears gray, without the black spot at the extremity; tail dusky brown above, white beneath. In the domesticated state the colours vary extremely.

Resides in holes under ground. Like the Hare, comes abroad in the evening to feed. Still more prolific than that species, breeding repeatedly in the year, and producing from four to eight at a litter. Young blind and naked at birth.

52. *L. albus*, Briss. (*Alpine Hare*.)—Dusky gray in summer, with a tinge of tawny; white in winter: ears shorter than the head, always black at the extremity.

L. albus, *Briss. Reg. An.* p. 139. *L. variabilis*, *Flem. Brit. An.*
 p. 22. *Alpine Hare, Penn. Brit. Zool.* vol. i. p. 102. pl. 10.
 no. 21. *Varying Hare, Shaw, Gen. Zool.* vol. ii. p. 201.

DIMENS. Length of the head and body twenty-one inches six lines; of the head four inches six lines; of the ears three inches three lines

of the tail two inches six lines; of the fore leg, from the olecranon to the end of the toe-nails, seven inches three lines; of the hind foot, from heel to toe, six inches.

DESCRIPT. In general somewhat smaller than the *Common Hare*, with shorter ears and more slender legs: tail shorter. Fur in summer grayish tawny mixed with black, here and there inclining to ash-colour; in winter wholly white, with the exception of the tips of the ears, which remain always black.

Inhabits the Scotch mountains: keeps near the summits, and never descends into the valleys, or mixes with the common species. Change of dress effected in the months of September and April; caused by an actual change of colour in the hair itself, without the shedding of the fur.—*Obs.* The *L. variabilis* of Pallas is probably a distinct species.

* GEN. 20. CAVIA, *Gmel.*

* 53. *C. Cobaya*, *Gmel.* (*Guinea Pig*).—Reddish yellow, variegated with black and white.

C. Cobaya, *Desm. Mammal.* p. 356. Variegated Cavy, *Shaw, Gen. Zool.* vol. 11. p. 17. pl. 126.

DIMENS. Length of the head and body twelve inches; of the head three inches one line; of the ears ten lines and a half.

DESCRIPT. Body thick and short; the neck scarcely to be distinguished from it: ears broader than long, straight, naked, transparent, partly concealed by the hair: eyes round, large, and prominent. Hair sleek and smooth; varying in colour in different individuals; entirely white; or variegated with black, white, and tawny yellow.

A native of Brazil; but domesticated in Europe. Is very prolific in confinement, breeding repeatedly in the year, and producing from four to twelve at a litter. Feeds on different vegetable substances.

ORDER IV. UNGULATA.

* GEN. 21. BOS, *Linn.*

* 54. *B. Taurus*, *Linn.* (*Common Ox*).—Forehead flat, longer than broad: horns taking their origin from the extremities of the occipital ridge.

B. Taurus, *Desm. Mammal.* p. 499. *Flem. Brit. An.* p. 24. *Ox, Penn. Brit. Zool.* vol. 1. p. 18. pl. 2. *Shaw, Gen. Zool.* vol. 11. pl. 208.

DESCRIPT. Varying extremely in the domesticated state in size and colour, as well as in the form and direction of the horns, which last are sometimes wholly wanting. A wild breed, (*Bewick, Quad.* p. 38.) formerly met with in Scotland, but now extinct, said to have been characterized by their white colour, with the muzzle and ears black.

Is capable of breeding at the age of two years. Goes nine months with young, and produces one, rarely two, at a time. The two central incisors are shed in the tenth month; the adjoining ones in the sixteenth; by the end of the third year the change is wholly completed.

* GEN. 22. OVIS, Linn.

* 55. O. *Aries*, Linn. (*Sheep*).—Horns compressed, and lunated.

O. *Aries*, *Desm. Mammol.* p. 488. *Flem. Brit. An.* p. 25. *Sheep, Penn. Brit. Zool.* vol. i. p. 27. *Shaw, Gen. Zool.* vol. ii. p. 385.

DESCRIPT. Varying much in size and colour: with or without horns: wool coarse or fine, long or short, according to the breed, sometimes approaching the quality of hair: tail short, or reaching below the knees.

Probably not originally indigenous. Primitive stock supposed to be the Argali* (*Ovis Ammon*, Gmel.), which is found in a wild state in the mountainous parts of Asia. Period of gestation about five months. Produces one or two at a birth, rarely more. The two middle incisors fall at the end of the first year, and are replaced by others; the next in succession at about the age of two and a half: by the end of the third year, or soon after, all have been renewed, and the individual is said to be *full-mouthed*.

* GEN. 23. CAPRA, Linn.

* 56. C. *Hircus*, Linn. (*Goat*).—Horns edged in front, rounded posteriorly.

C. *Hircus*, *Desm. Mammal.* p. 482. *Flem. Brit. An.* p. 25. Domestic Goat, *Penn. Brit. Zool.* vol. i. p. 35. pl. 3. *Shaw, Gen. Zool.* vol. ii. p. 369. pl. 199.

DESCRIPT. Subject to less variation than the *Sheep*. Usual colour black, white, or pied; occasionally brown, approaching more or less to a tawny red: horns in the *male*, in some instances upwards of three feet long; in the *female* much smaller, or wanting altogether: tail about seven inches in length; often black.

Ranges in a state of liberty on the mountains of Wales, Scotland, and Ireland, and is often domesticated; but it is doubtful whether it be indigenous. Supposed to be derived either from the C. *Iber*, Linn. or the C. *Ægagrus*, Gmel., the former of which species is found wild on the Alps of Europe, the latter on the mountains of Persia. Goes with young five months.

GEN. 24. CERVUS, Linn.

57. C. *Elaphus*, Linn. (*Stag, or Red Deer*).—Horns branched; round; diverging at the base, somewhat converging at the extremity.

C. *Elaphus*, *Desm. Mammal.* p. 434. *Flem. Brit. An.* p. 26. Stag, *Penn. Brit. Zool.* vol. i. p. 41. *Shaw, Gen. Zool.* vol. ii. p. 276. pl. 177.

DIMENS. Length of the body about six feet six inches; of the horns about two feet; of the tail seven inches: height about three feet eight inches.

DESCRIPT.—Varying in size and colour: usually reddish brown in summer, with a dusky line along the spine; in winter brownish gray; under parts whitish: horns at first simple, afterwards branched; the number of antlers increasing with age till they amount to ten or twelve; three of them being always directed forwards: eyes large, with a distinct lachrymal furrow: ears long and pointed: tail of moderate length. Female or *Hind* smaller, and without horns. The young or *Calf* is generally spotted with white, or as it is termed *menilled*, on the upper parts: the first indication of horns takes place during the latter part of the first year, when it is called a *Knobber*.

* *Shaw, Gen. Zool.* vol. ii. p. 379. pl. 201.

Formerly abundant throughout the kingdom, but now chiefly confined to the Highlands of Scotland. The horns are shed in March, and reappear in the course of the summer. Rutting season from Michaelmas to the end of November. Period of gestation rather more than eight months. Usually but one at a birth.

* 58. *C. Dama*, Linn. (*Buck, or Fallow Deer*.)—Horns branched; compressed; palmated at the top, diverging.

C. Dama, *Desm. Mammal.* p. 438. *Flem. Brit. An.* p. 26. *Fallow Deer*, *Penn. Brit. Zool.* vol. i. p. 41. *Shaw, Gen. Zool.* vol. ii. p. 282. pl. 178-9.

DIMENS. Length of the body about five feet; of the tail seven inches five lines: height about two feet ten inches.

DESCRIPT. Body much smaller than in the last species; tawny brown, with the back, flanks, shoulders, and thighs, more or less spotted with white; a dusky line down the middle of the back; buttocks white, bounded on each side by a descending black line; tail longer than in the *Stag*, blackish brown above, white beneath; abdomen and inside of the thighs whitish: horns round at bottom, with two antlers directed forwards; their upper portion compressed, and dilated into a broad palm, with tooth-like processes along the outer margin. Female or *Doe* without horns, and likewise the *Fawns* during the first year.

Var. β. Entirely white.

Var. γ. Deep brown, approaching to black.

Abundant in parks and forests in a half-reclaimed state, but doubtful whether indigenous. The black variety said to have been introduced from Sweden by King James the First; the others supposed to have come from Asia. Congregate in small herds. Rutting season in the Autumn. Female goes with young eight months, and produces one or two, rarely three, at a birth. The first two central incisors shed at the age of a year and a half; the change completed by the end of the fourth year.

59. *C. Capreolus*, Linn. (*Roe-Buck*.)—Horns branched; cylindrical; small and erect, with furcate summits.

C. Capreolus, *Desm. Mammal.* p. 439. *Flem. Brit. An.* p. 26. *Roe*, *Penn. Brit. Zool.* vol. i. p. 49. pl. 4. *Shaw, Gen. Zool.* vol. ii. p. 291.

DIMENS. Length of the body three feet nine inches; of the horns eight to nine inches; of the tail one inch: height (in front) two feet three inches, (behind) two feet seven inches.

DESCRIPT. Smaller than either of the preceding species, but of similar form. Colour variable; the general tint yellowish gray or reddish brown, more or less deep in different individuals, sometimes dusky; each hair being ash-coloured at the roots, then black, with the extreme tip tawny yellow; lower part of the neck, abdomen, and inside of the thighs, grayish white; contour of the anus pure white: ears long, furnished internally with long whitish hairs; extremity of the nose dusky, with a white spot on each side of the upper lip; chin white: horns very rugged, sulcated longitudinally, about as long as the head, with only two antlers, the first arising about the middle, directed forwards,—the second higher up directed backwards: no lachrymal furrow: tail very short. Summer coat much shorter and finer, and of a redder tint, than the winter one. *Doe* without horns. In the *young Buck*, the horns are plain and unbranched during the second year; in the third year furnished with a single antler, and in the fourth year complete.

Common formerly in Wales, in the North of England, and in Scotland, but at present almost confined to the Scottish Highlands. Loses its horns

at the latter end of Autumn. Rutting season during the first half of November. Doe goes with young five months and a half, and produces in April. Two at a birth, which are always male and female.

* GEN. 25. EQUUS, Linn.

* 60. E. *Caballus*, Linn. (*Horse*).—Tail uniformly covered with long hair; mane long and flowing; ears of moderate size: no dorsal line or transverse band.

E. *Caballus*, Desm. *Mammal.* p. 416. *Flem. Brit. An.* p. 27. *Horse*, Penn. *Brit. Zool.* vol. i. p. 1. pl. 1. Common Horse, Shaw, *Gen. Zool.* vol. ii. p. 419. pl. 214.

DESCRIPT. Offering every variety of size and colour: generally bay, or chestnut-brown, more or less deep; black, or grayish white: head long and tapering: teeth, incis. $\frac{6}{6}$, can. $\frac{1-1}{1-1}$ (seldom present in the *mare*), mol. $\frac{6-6}{6-6}$, = 40: ears erect and pointed, much smaller than in the next species: a naked callosity on the inside of the fore legs above the knee; another on the hind legs just under the knee.

Probably brought originally from Asia, where the species still exists in a truly wild state. A small variety occurs in the Highlands of Scotland, and in the Shetland Islands, half-reclaimed, but can scarcely be considered as indigenous. Period of gestation eleven months. Seldom more than one at a birth. Central incisors cast at the age of two years and a half, and replaced by permanent ones; the adjoining pair at three and a half; the remaining ones at four and a half; these last replaced more slowly than the others.

* 61. E. *Asinus*, Linn. (*Ass*).—Gray, inclining to reddish; with the dorsal line, and a transverse band across the shoulders, black: ears very large: tail terminating in a tuft of long hair.

E. *Asinus*, Desm. *Mammal.* p. 414. *Ass*, Penn. *Brit. Zool.* vol. i. p. 13. Shaw, *Gen. Zool.* vol. ii. p. 429. pl. 216.

DESCRIPT. Smaller than the *Horse*, and subject to less variation than that species. Colour generally gray, more or less dark; sometimes approaching to silvery white, obscurely spotted with stains of a reddish cast, at other times dark brown or dusky: a transverse black stripe upon the shoulders, crossing another of the same colour down the middle of the back; these marks always more or less obvious. Head shorter and thicker than that of the horse; ears long and slouching: tail tipped with long hair: no naked callosities on the hind legs.

A native of the East. Introduced into this country towards the close of the tenth century. Goes eleven months with young, and produces one at a birth. Breeds occasionally with the Horse: the hybrid production termed a *Mule* or a *Hinny*, according as the Ass is the *male* or *female* parent.

GEN. 26. SUS, Linn.

† 62. S. *Scrofa*, Linn. (*Boar*).—Tusks strong, triangular, of moderate length, directed to one side.

S. *Scrofa*, Desm. *Mammal.* p. 389. *Flem. Brit. An.* p. 28. *Hog*, Penn. *Brit. Zool.* vol. i. p. 55. Wild Boar, and Common Hog, Shaw, *Gen. Zool.* vol. ii. p. 459. pl. 221 and 222.

DIMENS. (*Wild Boar*). Length of the head and body three feet three inches; of the head one foot; of the tail seven inches six lines: height about one foot nine inches.

DESCRIPT. Head elongated: neck short: body thick and muscular: legs short and strong: ears rather short: eyes small: mouth large, with the upper lip pushed

up by the tusks; body covered with long stiff bristles, intermixed at the roots with a soft woolly hair; the longest and strongest bristles on the back. General colour dusky gray.—DESM.

**Var. β. domestica.* (*Domestic Hog.*)—In this variety the ears are longer and more or less pendulous; the bristles more thinly scattered, and of one sort; the tusks comparatively short; the tail more or less twisted; the size and colour very variable.

Formerly abundant in a wild state throughout the country. Continued to inhabit the forests about London so late as the reign of Henry the Second, but it is not exactly known how soon after that period they were extirpated. In the domestic state the Sow goes with young about four months, and is very prolific, producing sometimes as many as twenty at a litter. Food extremely various.

ORDER V. CETACEA.

GEN (1.) *MANATUS*, Cuv.?

GEN. (2.) *STELLERUS*, Cuv.?

- (1.) ———? *Manatus borealis*, Flem. Brit. An. p. 29. *Mermaid of the Shetland Seas*, Edinb. New Phil. Journ. (1829) vol. vi. p. 57.

Two instances at least are on record in which specimens of the *Hervivorous Cetacea* have been observed in the British seas, but there is no satisfactory evidence by which the species can be determined, or even the genus, to which they belonged*. In the *Elements of Nat. Hist.* by Mr Stewart, mention is made (vol. i. p. 125.) of the carcase of one of these animals which was thrown ashore near Leith; and in the *Edinb. New Philos. Journ.* as quoted above, there will be found some account of another individual, which occurred a few years ago off the Shetland Islands. See also Dr Fleming's *Brit. An.* p. 30.

GEN. 27. DELPHINUS, Linn.

(1. DELPHINUS, Cuv.)

63. *D. Delphis*, Linn. (*Common Dolphin.*)—Jaws moderately produced; nearly of equal length: teeth more than forty on each side above and below; slender, slightly bent, pointed.

D. Delphis, *Desm. Mammal.* p. 514. *Flem. Brit. An.* p. 35. *Dolphin*, *Penn. Brit. Zool.* vol. III. p. 65. *Shaw, Gen. Zool.* vol. II. p. 507. pl. 229.

DIMENS. Entire length from six to seven feet.

DESCRIPT. Body slender, thickest in the middle, gradually tapering towards the head and tail: muzzle or beak, measured from the forehead, equalling in length the rest of the head; much depressed, narrow, and

* On this account I have introduced, in the Synoptic Arrangement of Genera, the characters of both *Manatus* and *Stellerus* Cuv., which are perhaps equally likely to occur in the British seas. At any rate they will prove useful in determining the *true* genus of any of these animals which may be met with hereafter.

somewhat pointed at the extremity: jaws nearly equal, the upper being slightly shorter than the lower: teeth very numerous, $\frac{42-42}{42-42}$ to $\frac{47-47}{47-47}$ = 168 to 188, slender, somewhat bent inwards, sharp-pointed; placed at equal distances from each other, and locking in between each other when the jaws are closed: eyes small, almost in a line with the mouth: blow-hole situate on the top of the head a little above the eyes: pectoral fins placed very low, in form somewhat falcate: dorsal fin pointed, taking its origin from a little beyond the middle of the back; rather elevated, and when measured along the line of flexure, equalling one-sixth of the entire length of the body: tail crescent-shaped. Skin smooth; of a dusky colour above, white beneath, and grayish on the sides.

Met with occasionally on the British shores, but not of very frequent occurrence. Feeds on animal substances. Period of gestation said to be ten months. Produces one or two at a birth.

64. *D. Tursio*, Fab. (*Blunt-toothed Dolphin*).—Jaws moderately produced; the lower longer than the upper: teeth rather more than twenty on each side above and below; straight, with obtuse summits.

D. Tursio, *Desm. Mammal.* p. 514. *Flem. Brit. An.* p. 37. *D. truncatus*, *Mont. in Wern. Mem.* vol. III. p. 75. pl. 3. Bottle-nose Whale, *Hunter in Phil. Trans.* for 1787. pl. 18.

DIMENS. Entire length eleven feet: girth seven feet four inches: length of the mouth fourteen inches: from the snout to the eye sixteen inches: from the same to the pectoral fin two feet: from the same to the dorsal four feet eight inches: from the same to the vent seven feet three inches: length of the dorsal twenty-three inches: height of the same ten inches.

DESCRIPT. Larger than the last species: lower jaw projecting further beyond the upper: teeth less numerous, $\frac{21-21}{21-21}$ to $\frac{23-23}{23-23}$ = 84 to 92, straight, somewhat conical, but blunt and truncated at the extremity.

But little known as a British species. Hunter's specimen was caught upon the sea-coast near Berkeley, and is now in the Museum of the College of Surgeons. In the same collection is a second individual from the Thames. The one described by Montagu, which appears to have been large and aged, with the summits of the teeth more than usually truncated, was taken in the river Dart, in 1814. The measurements given above are those of a fourth individual which occurred a few years since in the river at Preston. They were obligingly sent me by Mr Gilbertson of that place.

(2. *PHOCÆNA*, Cuv.)

65. *D. Phocæna*, Linn. (*Porpesse*).—Under jaw slightly projecting beyond the upper: teeth twenty-two to twenty-five on each side above and below; straight, compressed, and rounded at the summits.

D. Phocæna, *Desm. Mammal.* p. 516. *Flem. Brit. An.* p. 33. *Porpesse*, *Penn. Brit. Zool.* vol. III. p. 69. *Shaw, Gen. Zool.* vol. II. p. 504. pl. 229.

DIMENS. Entire length from four feet to five feet and a half.

DESCRIPT. Body elongated, gradually tapering towards the tail: snout short, rather obtuse at the extremity: under jaw somewhat longer

than the upper: teeth numerous; from twenty-two to twenty-five in each jaw on each side, compressed, and nearly straight: eyes small, almost in a line with the mouth: blow-hole crescent-shaped, with the concavity directed forwards: pectoral fins placed low down, oval, and somewhat pointed: dorsal fin straight, triangular, rather beyond the middle of the body. Skin smooth; dusky on the back, whitish on the belly, the two colours meeting on the sides.

A constant inhabitant of the British seas, often entering the mouths of rivers. Preys on mackerel, herrings, and other fish. A specimen which occurred in the London market in May 1833 was found to contain a full-formed fœtus; it is probable, therefore, that they produce their young at about that period of the year.

66. *D. Orca*, Fab. (*Grampus*).—Upper jaw projecting a little beyond the lower: teeth about eleven on each side above and below; conical, bent at the summits: pectoral fins broad and oval.

D. Orca, *Flem. Brit. An.* p. 34. *D. Grampus*, *Desm. Mammal.* p. 517. *Grampus*, *Hunter in Phil. Trans.* 1787, pl. 16.

DIMENS. Entire length from twenty to twenty-five feet.

DESCRIPT. Much larger than the last species; the body deeper and thicker in proportion to its length. Snout very short and obtuse: upper jaw somewhat longer than the lower, but this last broader than the upper: teeth unequal, conical, a little bent at the summits; varying in number according to the age of the individual, generally about twenty two in each jaw: eyes almost in the same line with the mouth: dorsal fin nearly in the middle, very much elevated, and pointed at the extremity: pectorals very broad, of an oval form: tail crescent-shaped. Skin smooth; glossy black above, white beneath, the two colours meeting on the sides but separated by a well-defined line; an oval white spot behind each eye.

Inhabits the British seas in large herds, and occasionally enters rivers. Is of a fierce and voracious disposition, preying upon the larger species of fish.—*Obs.* The *Delphinus Gladiator* and the *D. ventricosus* of Lacépède, two species constituted by that author from individuals taken in the Thames, are considered by Cuvier as not really distinct from the above.

67. *D. melas*, Traill. (*Caving Whale*).—Top of the head very convex: teeth conical, varying in number: pectoral fins long and narrow.

D. melas, *Traill in Nichol. Journ.* vol. XXII. p. 81. *Flem. Brit. An.* p. 34. *D. globiceps*, *Cuv. Ann. du Mus.* tom. XIX. p. 14. pl. 1. f. 2, 3. *D. Deductor*, *Scoresby, Arct. Rég.* vol. I. p. 496. pl. 13. f. 1.

DIMENS. Entire length from twenty to twenty-four feet.

DESCRIPT. Equalling the *Grampus* in size, but differing essentially from that species in the form and character of the fins; the dorsal much shorter, the pectorals longer, narrower, and more pointed: head short and round, with the forehead remarkably convex and prominent: upper jaw projecting a little beyond the lower: teeth conical, sharp, and a little bent; varying much in number in different individuals, not visible in

very young specimens, and falling in advanced life; average number from twenty-two to twenty-four in all. Skin smooth, of a deep bluish black colour, with the exception of a whitish band beneath the body extending from the throat to the anus.

Common in large herds off the Orkney and Shetland Islands.

(3. DELPHINAPTERA, *Lacép.*)

68. *D. albicans*, Fab. (*Beluga*).—Head blunt: teeth about nine on each side above and below; short, with obtuse summits.

D. leucas, *Desm. Mammal.* p. 519. *Delphinaptera albicans*, *Flem. Brit. An.* p. 36. *Beluga*, *Barclay and Neill in Wern. Mem.* vol. III. p. 371. pl. 17. *Scoresby, Arct. Reg.* vol. I. p. 500. pl. 14.

DIMENS. Entire length from twelve to eighteen feet.

DESCRIPT. Body thick in the middle, somewhat tapering towards each extremity, but more especially towards the tail: head small, blunt, and round: the forehead rising abruptly: eyes and mouth small: jaws equal: teeth in the adult state about nine on each side above and below, short, straight, slightly compressed, with obtuse summits; those in the upper jaw falling in advanced life: a longitudinal ridge on the back supplying the place of the dorsal fin: pectorals short, broad, and oval. Skin smooth: colour wholly white; sometimes with a tinge of yellow.

A native of high Northern latitudes, where it is found in herds of thirty or forty together. Must be considered as a very rare visitant of the British seas. The individual described by Mr. Neill l.c. was taken in the Frith of Forth in June 1815.

GEN. 28. MONODON, *Linn.*

69. *M. Monoceros*, *Linn.* (*Narwhal*).—Body sub-conical, with a dorsal and ventral ridge: head obtuse, about one-seventh of the entire length.

M. Monoceros, *Scoresby, Arct. Reg.* vol. I. p. 486. pl. 15. *Flem. Brit. An.* p. 37. Small-headed Narwhal, *Flem. in Wern. Mem.* vol. I. p. 131. pl. 6.

DIMENS. Entire length, exclusive of the tusk, thirteen to sixteen feet; circumference (at the thickest part) eight to nine feet; length of the tusk about five feet*.

DESCRIPT. Anterior half of the body nearly cylindrical; posterior half conical: this latter portion furnished with a dorsal and ventral ridge, which take their origin about three feet from the extremity, and extend half way across the tail; the edges of the tail run in like manner six or eight inches along the body, forming ridges on the sides of the rump: head about one-seventh of the entire length; small, blunt, and round; the forehead very prominent, rising suddenly from the snout: mouth small; intermaxillary bones furnished each with one tooth directed forwards; in the *female* these teeth generally remain through life concealed in the sockets, not appearing externally; in the *male*, that on the left side is exerted, growing to the length of several feet;

* The above measurements are from Scoresby, from whose excellent work on the Arctic Regions much assistance has been derived in drawing up the characters of this species, as well as of some others of the *Cetacea*.

it is spirally striated from right to left, nearly straight, and tapering to a round blunt point; very rarely, in this last sex both teeth are equally developed, and both exerted: blow-hole semicircular, situate directly over the eyes: pectoral fins short: no dorsal fin, but instead of it an irregular sharpish fatty ridge, two inches in height, extending two feet and a half along the back, nearly midway between the snout and the tail: tail divided by a notch into two lobes, which project laterally and are somewhat pointed. Prevailing colour white, or yellowish white, with dark gray or blackish spots of different degrees of intensity.

Has only occurred hitherto in two or three instances on the British shores. In the Northern seas is said to be gregarious; each sex herding separately. Feeds on *sepia* and other molluscous animals.

GEN. 29. HYPEROODON, *Lacép.*

70. *H. bidens*, *Flem.* (*Bottle-head.*)—Teeth two only in the fore part of the lower jaw.

H. bidens, *Flem. Brit. An.* p. 36. *H. Butskopf*, *Lacép. Cétac.* p. 319. Bottle-nose Whale, with two teeth, *Hunter in Phil. Trans.* 1787. pl. 19. Beaked Whale, *Penn. Brit. Zool.* vol. 111. p. 59. pl. 5. f. 1. Two-toothed Cachalot, *Sow. Brit. Misc.* pl. 1.

DIMENS. Entire length from twenty to twenty-five feet.

DESCRIPT. Body elongated; greatest circumference in the region of the pectoral fins: forehead high, very convex, rising suddenly from the snout; this last short and depressed, terminating in a kind of beak somewhat similar to that of the genus *Delphinus*: lower jaw rather longer and larger than the upper: teeth conical, and pointed; only two, situate in the fore part of the lower jaw; sometimes altogether wanting, or not appearing above the gums: palate studded with little horny eminences, considered by Cuvier as rudimentary vestiges of *whalebone*: eyes large, a little above the line of the lips: blow-hole crescent-shaped, with the horns of the crescent directed towards the tail: dorsal fin placed considerably beyond the middle of the body, but little elevated, lanceolate, pointed, inclining backwards: pectorals small, oval, in the same horizontal line with that of the mouth. Skin smooth and glossy, blackish lead-colour above, whitish underneath, the two colours mixing on the sides.

Occasionally met with on the British shores. Hunter's specimen was taken in the Thames above London Bridge in 1783.—*Obs.* This species has been unnecessarily split into several by many authors: it probably embraces all the following of Desmarest's "Mammalogie;" *Delphinus Chemnitzianus*, *D. Hunteri*, *D. edentulus*, *D. Hyperoodon*, and *D. Sowerbyi*.

GEN. 30. PHYSETER, *Linn.*

(1. CATODON, *Lacép.*)

71. *P. macrocephalus*, *Shaw.* (*Blunt-headed Cachalot.*)—Teeth in the lower jaw from twenty to twenty-four on each side; mostly conical with obtuse summits.

Catodon macrocephalus and *C. Trumbo*, *Lacép. Cétac.* pp. 165 and 212. *C. macroceph.* *Flem. Brit. An.* p. 39. Blunt-headed Cachalot, *Penn. Brit. Zool.* vol. 111. p. 61. pl. 6. Spermaceti Whale, *Alderson in Camb. Phil. Trans.* vol. 11. p. 253. pl. 12—14.

DIMENS. Entire length from fifty to sixty-three feet.

DESCRIPT. Head enormously large, forming more than one-third of the entire bulk; the body gradually tapering from the posterior part of it towards the tail: upper part of the snout very thick and swollen, as it were truncated in front, and overhanging considerably the lower jaw; this last of a narrow elongated form, fitting, when the mouth is closed, into a grooved cavity above: upper jaw without whalebone or visible teeth, although a few teeth may be found concealed within the gums on cutting through the integuments; in the lower jaw from forty to forty-nine teeth (there being occasionally an odd one), entering likewise when the mouth is shut into corresponding cavities above; the last three or four on each side smaller than the others, and somewhat hooked; the rest projecting above the gums about two inches, cylindrical or conical, with bluntish summits; those in front inclining *backwards*, those situate more behind *forwards*, the middle tooth on each side being nearly vertical: blow-hole single, near the extremity of the snout, and placed rather to the left of the median line: no dorsal fin; instead of it a callous ridge commencing gradually, and terminating behind abruptly in a sort of hook-like process: pectorals small. General colour black or dusky, somewhat paler beneath*.

Occasionally stranded on different parts of the coast, but not of frequent occurrence in the British seas. The upper part of the head in this species consists of large cavities, separated from each other by a cartilaginous substance, and filled with an oily fluid, which, in its congealed state, forms the *spermaceti* of commerce. These cavities are quite distinct from that of the cranium which is situate beneath.

(2.) *P. Catodon*, Linn.—*Catodon Sibbaldi*, Flem. Brit. An. p. 39.

This supposed species is too imperfectly characterized, and rests on too doubtful authority, to rank as distinct. In the opinion of Cuvier, (*Oss. Foss. tom. v. p. 335.*) the herd stated by Sibbald (*Phalainolog. Nov. p. 24.*) as having occurred at Kairston in Orkney, consisting of an hundred individuals, were probably Belugas (*Delphinus albicans*), some of which had lost the teeth in the upper jaw through age. Moreover Sibbald appears only to have had his account from others, and not to have seen any of the individuals himself.

(2. PHYSETER, *Lacép.*)

72. *P. Tursio*, Linn. (*High-finned Cachalot.*)—Teeth very slightly bent, with flat summits.

P. Tursio, *Flem. Brit. An.* p. 38. *P. Mular*, *Lacép. Cétac.* p. 239. *Desm. Mammal.* p. 526. High-finned Cachalot, *Penn. Brit. Zool.* vol. III. p. 64.

DESCRIPT. Said to be characterized by an erect dorsal fin, very high, and pointed at the extremity: blow-hole in front: teeth very slightly bent, with flat or obtuse summits: in other respects similar to the *P. macrocephalus*.

But very little is known of this species. The few particulars on record respecting it are principally taken from Sibbald, who briefly describes a Cachalot with the above characters, which came on shore on the Orkney Isles in 1687.

* Some parts of this description are borrowed from Alderson's account of this species, l. c.

- (3.) *P. microps*, Linn. *Flem. Brit. An.* p. 38.

This species, which entirely owes its existence to Sibbald's *Phalainologia*, rests upon very vague and uncertain authority. It is said to resemble the *P. Tursio*, excepting in having the teeth more pointed. Cuvier does not admit that there is any well-founded distinction between the two. Indeed, in his *Ossements Fossiles*, (tom. v. p. 328.) he would seem to entertain some doubts with respect to the existence of either.

GEN. 31. BALÆNA, Linn.

(1. BALÆNA, *Lactp.*)

73. *B. Mysticetus*, Linn. (*Common Whale*).—Gape of the mouth arched: upper jaw with about six hundred and fifty laminae of whalebone.

B. Mysticetus, Scoresby, *Arctic Reg.* vol. i. p. 449. pl. 12. *Flem. Brit. An.* p. 33. *Common Whale*, Penn. *Brit. Zool.* vol. III. p. 50.

DIMENS. Entire length averaging from fifty to sixty-five feet: greatest circumference from thirty to forty feet.

DESCRIPT. One of the most bulky, but not in general the longest of the Cetaceous tribe. Body thickest in the middle, a little behind the fins, from which point it gradually tapers, in a conical form, towards the tail, and slightly towards the head: this last very large, of a somewhat triangular form; "the under part, the arched outline of which is given by the jawbones, flat, and measuring sixteen to twenty feet in length, and ten to twelve in breadth: the lips, which are five or six feet high, and form the cavity of the mouth, are attached to the under jaw, and rise from the jaw-bones, at an angle of about eighty degrees, having the appearance, when viewed in front, of the letter U: the upper jaw, including the crown-bone or skull, bent down at the extremity, so as to shut the front and upper parts of the cavity of the mouth, and overlapped by the lips in a squamous manner at the sides*:" no teeth; but the palate furnished with two extensive rows of whalebone, generally curved longitudinally, and giving an arched form to the roof of the mouth; each series consists of upwards of three hundred laminae, the interior edges of which are covered with a fringe of hair: eyes remarkably small: pectoral fins situate about two feet beyond the angle of the mouth: tail horizontal, of great breadth, and of a semilunar form; the lateral lobes somewhat pointed, and turned a little backward. Colour black, or blackish gray, with the exception of the fore part of the under jaw and a portion of the belly, which are white.

Appears to have been formerly of not unfrequent occurrence in the British seas, but must be considered in these days as an extremely rare visitant. Sibbald mentions one which came ashore near Peterhead in 1682. A small one is stated to have been taken near Yarmouth, July 8, 1784†. The food of this species is said to consist principally of shrimps and molluscous animals.

* Scoresby.

† C. and J. Paget's *Nat. Hist. of Yarmouth and its Neighbourhood*.

(2. BALÆNOPTERA, *Lacép.*)

74. *B. Physalus*, Linn. (*Fin-Fish.*)—Pectoral skin without longitudinal folds.

Balænoptera Gibbar, *Scoresb. Arct. Reg.* vol. i. p. 478. *Physalis vulgaris*, *Flem. Brit. An.* p. 32. *Fin-Fish*, *Penn. Brit. Zool.* vol. iii. p. 57. *Fin-backed Mysticete*, *Shaw, Gen. Zool.* vol. ii. p. 490. pl. 227.

DIMENS. Entire length about one hundred feet: greatest circumference from thirty to thirty-five feet.

DESCRIPT. The longest of the Cetaceous tribe. Body more slender, and less cylindrical than that of the *B. Mysticetus*; considerably compressed at the sides, and angular on the back: head smaller than in that species; the snout more pointed, with the jaws nearly equal; the whalebone shorter, the longest lamina measuring about four feet: a small horny protuberance, or dorsal fin, near the extremity of the back: pectorals long and narrow. Colour pale bluish black, or dark bluish gray.

Apparently of equal rarity in the British seas with the species last described. It is included by Pennant in his *British Zoology*, but it is not said on what authority.

75. *B. Boops*, Linn. (*Sharp-lipped Whale.*)—Pectoral skin with longitudinal folds admitting of dilatation: jaws pointed.

Balænoptera Jubartes, *Scoresb. Arct. Reg.* vol. i. p. 484. *B. Boops*, *Flem. Brit. An.* p. 31. *Pike-headed Mysticete*, *Shaw, Gen. Zool.* vol. ii. p. 492. pl. 227. *Fin-Whale*, *Neill in Wern. Mem.* vol. i. p. 201.

DIMENS. Entire length about forty-six feet: greatest circumference about twenty feet.

DESCRIPT. Body very thick, and somewhat elevated, immediately over the pectoral fins; gradually tapering from that point towards the tail: head moderately large, becoming narrower towards the extremity of the snout, which terminates however in a somewhat broadish tip: lower jaw one-third of the entire length: palate with about three hundred laminæ of whalebone on each side, the longest measuring about eighteen inches in length: dorsal protuberance or fin placed far down the back; two feet and a half high: pectorals four or five feet long, scarcely a foot broad. Colour black above, whitish on the belly, inclining to red between the pectoral folds.

Represented as being of not unfrequent occurrence in the Scotch seas; and occasionally stranded on different parts of the English coast. There is some doubt whether the whale described and figured by Dr Johnston in the *Trans. of Newcastle Nat. Hist. Soc.* (vol. i. p. 6.) be referable to this species or not, as it possessed pectorals nine feet in length, being one-fourth of the length of the body. It was thrown on shore about two miles north of Berwick, in Sept. 1829.

76. *B. Musculus*, Linn. (*Round-lipped Whale.*)—Pectoral skin with longitudinal folds: margin of the under lip semicircular.

Balænoptera Rorqual, *Scoresb. Arct. Reg.* vol. i. p. 482. *B. Musculus*, *Flem. Brit. An.* p. 30. *Round-lipped Whale*, *Penn. Brit. Zool.* vol. iii. p. 58.

DIMENS. Entire length from seventy to eighty-six feet.

DESCRIPT. Strongly resembling the last species, but said to grow to a much larger size: differs also in having the lower lip much broader than the upper, and semicircularly turned at its extremity, while the upper is somewhat sharp or pointed at the tip: gape large; longest lamina of whalebone three feet in length.

Found in the Scotch seas with the preceding. Said to feed principally on herrings.

Obs. The characters of the species in this sub-genus are not sufficiently well understood to admit of being laid down with any degree of certainty, and there can be little doubt that the species themselves have been unnecessarily multiplied. Cuvier appears to be of opinion (*Regne An.* tom. 1. p. 298.) that even the *B. Physalus* may eventually turn out to be the same with the *B. Boops*, and that the supposed difference between them may be the result of hasty and imperfect examination. There is likewise but little dissimilarity between the *B. Boops* and the *B. Musculus*, and it is not as yet satisfactorily ascertained that even that little is to be depended upon as constant in all cases. With respect to the *Balæna rostrata* of Fabricius and Hunter (*Phil. Trans.* 1787. pl. 20.) which occurs not unfrequently on the British coasts, it is probably nothing more than the young state of one of the two last mentioned; in the opinion of Cuvier*, of the *B. Boops*.

* See his *Ossemens Fossiles* (tom. v. p. 365.) which contains a valuable dissertation on the different species of the *Cetacea*.

CLASS II. AVES.

§ 1. *Feet formed for grasping.*

ORDER I. RAPTORES.

Bill strong, covered at the base with a cere, hooked towards the extremity: legs strong and muscular, short, or of moderate length: toes four in number, three directed forwards and one behind; rough underneath; armed with powerful, sharp, curved, retractile talons.

I. VULTURIDÆ.—*Head and neck more or less divested of feathers: nostrils lateral, placed in the cere, oval or elongated: feet generally naked: claws moderately curved.*

1. NEOPHRON.—Bill long, slender, straight, hooked at the extremity; lower mandible bending downward; no gonys: nostrils longitudinal, lateral, situate near the ridge: fore part of the head naked; neck feathered: feet strong, moderate, naked: wings long; third quill the longest: tail of fourteen feathers.

II. FALCONIDÆ.—*Head feathered: nostrils lateral, placed in the cere, rounded or oval: legs feathered to the toes, or naked: claws strongly curved, very sharp.*

2. AQUILA.—Bill long, very robust, convex or slightly angular above, straight at the base, much hooked

at the tip: cere hispid: nostrils rounded, or lunulate: wings long.

- (1. *AQUILA*.) Tarsi feathered to the toes: claws unequal, grooved beneath: fourth quill longest.
- (2. *HALIÆTUS*.) Tarsi half-feathered; acrotarsia scutellated; claws unequal, grooved beneath: fourth quill longest: nostrils transverse.
- (3. *PANDION*.) Tarsi naked; acrotarsia reticulated with small rounded scales: outer toe reversible: claws equal, rounded beneath: second quill longest.

3. *FALCO*.—Bill short, strong, bending from the base; upper mandible strongly toothed, lower one emarginated: nostrils round: tarsi short and strong; acrotarsia reticulated: wings long; second quill longest; first and second with the inner webs deeply notched near the extremity.

4. *ACCIPITER*.—Bill short, rather strong, bending from the base: nostrils oval: tarsi slender; acrotarsia scutellated: middle toe much longer than the lateral ones: wings short, not reaching beyond two-thirds of the length of the tail: fourth quill longest.

- (1. *ASTUR*.) Tarsi moderate; rather robust: acrotarsia scutellated.
- (2. *ACCIPITER*.) Tarsi long and slender: acrotarsia scutellated; the sutures of the scales obsolete.

5. *MILVUS*.—Bill moderate, rather weak, bending slightly from the base, somewhat angular above: nostrils oblique, oval or elliptic: tarsi short: toes and claws weak: wings very long: tail long, and forked.

- (1. *MILVUS*.) Tarsi feathered a little below the knee; acrotarsia scutellated: fourth quill longest.
- (*ELANUS*.) Tarsi very short, half-feathered; acrotarsia reticulated: second quill generally the longest.

6. *BUTEO*.—Bill moderate, bending from the base: nostrils oval or roundish: tarsi partly naked, or clothed with feathers: wings long; third and fourth quills generally the longest; the four first emarginated on their inner webs: tail even.

- (1. *BUTEO*.) Lore without feathers: tarsi short and strong; naked, with the acrotarsia scutellated; or feathered to the toes.

(2. **PERNIS.**) Lore closely covered with small scaly feathers: tarsi moderate, half-feathered; acrotarsia reticulated.

(3. **CIRCUS.**) Sides of the head furnished with a circle of feathers approaching that of the owls: tarsi elongated, feathered a little below the joint; acrotarsia scutellated: tail long; somewhat rounded.

III. STRIGIDÆ.—*Head large, feathered; base of the bill clothed with stiff bristly feathers directed forwards, concealing the cere and nostrils: eyes surrounded by a circle of radiating feathers: legs feathered to the toes: outer toe reversible.*

7. **BUBO.**—Bill strong, bending from the base: nostrils oval or rounded: facial disk small and incomplete: auditory aperture small, oval, without an operculum: head furnished with large conspicuous egrets: wings moderate; third quill generally the longest.

(1. **BUBO.**) Legs robust, feathered to the claws.

(2. **SCOPS.**) Tarsi feathered: toes naked.

8. **OTUS.**—Bill bending from the base: nostrils oval, oblique: facial disk complete, of moderate size: auditory conch large, the aperture covered by an operculum: egrets more or less conspicuous: wings long; second quill longest: legs feathered to the claws.

9. **STRIX.**—Bill somewhat elongated, bending at the tip only: nostrils oval: facial disk complete and full-sized: auditory aperture large, furnished with an operculum: head without egrets: wings long; second quill longest: tarsi feathered: toes hairy; middle claw serrated beneath.

10. **SYRNIUM.**—Bill bending from the base: nostrils round: facial disk large and complete: auditory aperture large, furnished with an operculum: head without egrets: wings short and rounded; fourth and fifth quills longest: legs feathered to the claws.

11. **NOCTUA.**—Bill bending from the base: nostrils oval: facial disk small, and generally very incom-

plete: auditory aperture small, oval: head without egrets: wings moderate; third quill longest.

(1. SURNIA.) Legs and toes thickly feathered.

(2. NOCTUA.) Tarsi feathered: toes either feathered or hairy.

ORDER II. INSESSORES.

Bill various: legs short, or of moderate length: feet adapted for perching: toes four, varying in position, flat underneath; hind toe articulated on the same plane with the fore toes: claws slender, somewhat retractile, curved and acute.

I. DENTIROSTRES.—*Bill moderate, more or less weak, furnished with a notch or tooth on each side towards the tip: rictus armed with bristles: feet generally slender; toes three before and one behind.*

(LANIIDÆ.)

12. LANIUS.—Bill robust, convex above, much compressed at the sides, furnished at the base with hairy feathers directed forwards; upper mandible hooked at the extremity, and strongly notched: nostrils basal, lateral, nearly round, partly closed by a membrane: wings with the first quill short; the second shorter than the third and fourth, which are longest.

(MUSCICAPIDÆ.)

13. MUSCICAPA.—Bill moderate, somewhat triangular, depressed at the base, compressed towards the tip, which is deflected; the upper mandible emarginated: rictus armed with long stiff bristles: tarsus a little longer

than the middle toe: side toes of equal length: wings with the first quill very short; third and fourth longest.

(MERULIDÆ.)

14. CINCLUS.—Bill rather slender, straight, rounded, compressed at the sides; upper mandible emarginated at the tip, slightly bending over the lower one: nostrils longitudinally cleft, and partly covered by a membrane: tarsus longer than the middle toe: wings and tail short; the former with the first quill very short, the third and fourth longest.

15. TURDUS.—Bill moderate, convex above, slightly bending towards the point, which is rather compressed; upper mandible emarginated: rictus furnished with a few bristles: nostrils oval; partly covered by a naked membrane: tarsus longer than the middle toe: wings and tail moderate; the former with the first quill extremely short, almost abortive; the second somewhat shorter than the third and fourth, which are longest.

16. ORIOLUS.—Bill subconic, depressed at the base; upper mandible carinated above, slightly emarginated, inclining over the lower one: nostrils opening longitudinally in a large membrane: tarsus not longer than the middle toe: wings moderate; first quill very short; third longest

(SYLVIIDÆ.)

17. ACCENTOR.—Bill rather strong, subconic, straight, acute, broader than high at the base, very much compressed in front; upper mandible emarginated at the tip; the tomia of both mandibles inflected: nostrils basal, naked, pierced in a large membrane: tarsi strong: wings moderate; first quill very short; third and fourth longest.

18. SYLVIA.—Bill slender, depressed at the base, compressed in front; upper mandible bending over the lower one, and slightly emarginated at the tip: lower man-

dible straight: nostrils basal, pierced in a membrane, more or less exposed: wings short, or moderate; first quill very short, sometimes almost abortive; third, or third and fourth, generally the longest.

- (1. ERITHACA.) Bill rather strong, as broad as high at the base; gonys slightly ascending: rictus with long weak diverging bristles: tarsi slender, somewhat elongated: wings moderate; fourth and fifth quills longest: tail divaricated; the tips of the feathers pointed.
- (2. PHOENICURA.) Bill slender, as broad as high at the base: rictus nearly smooth: tarsi moderate: wings rather long; third and fourth quills longest: tail even; the feathers obtuse, and generally rufous.
- (3. SALICARIA.) Bill very slender, broader than high at the base: rictus with long diverging bristles: tarsi moderate: wings short, and somewhat rounded; first quill nearly abortive; third longest: tail cuneiform; the tips of the feathers rounded.
- (4. PHILOMELA.) Bill rather strong, as broad as high at the base; gonys slightly ascending: rictus with a few short bristles: tarsi elongated: wings moderate; third and fourth quills longest: tail with the two middle feathers longest; the tips of all rounded.
- (5. CURRUCA.) Bill rather strong, broader than high at the base: gonys slightly ascending: rictus nearly smooth: tarsi moderate: wings moderate; first quill nearly abortive; third longest: tail slightly forked.
- (6. SYLVIA.) Bill very slender, broader than high at the base: rictus with a few longish bristles: tarsi rather elongated: wings moderate; third and fourth quills longest: tail slightly forked; the tips of the feathers rounded.

19. MELIZOPHILUS.—Bill very slender, short, slightly arched from the base, compressed, with the tip finely emarginated; tomia of both mandibles inflected towards the middle: rictus with a few bristles: tarsi strong, longer than the middle toe: wings short; first quill very small; third, fourth, and fifth, equal and longest: tail much elongated, cuneiform.

20. REGULUS.—Bill very slender, subulate, straight, compressed throughout its whole length; the tomia inflected inwards: nostrils concealed by small bristly feathers directed forwards: tarsi rather long: wings with the first quill very short; second much shorter than the third; fourth and fifth longest: tail moderate; divaricated, the tips of the feathers pointed.

21. *MOTACILLA*.—Bill slender, subulate, straight, carinated, angulated between the nostrils, emarginated at the tip; the tomia of both mandibles slightly compressed inwards: tarsus considerably longer than the middle toe: wings with the first quill extremely short; second and third longest; one of the scapulars as long as the quills: tail elongated, even.

22. *ANTHUS*.—Bill slender, straight, subulate towards the extremity, with the tomia compressed inwards about the middle; upper mandible carinated at the base, bending downwards at the tip, and slightly emarginated: tarsus generally longer than the middle toe; hind claw more or less produced: wings with the first quill abortive; second shorter than the third and fourth, which are longest: two of the scapulars as long as the quills.

23. *SAXICOLA*.—Bill rather strong, straight, carinated above, dilated at the base, advancing on the forehead; upper mandible emarginated, slightly bending at the tip: rictus with some stiff longish bristles: tarsi elevated; hind claw shorter than the toe: wings with the first quill not half the length of the second; third and fourth longest: coverts and scapulars short.

24. *PARUS*.—Bill strong, short, straight, subconical, slightly compressed, sharp-pointed, without any notch: nostrils basal, round, concealed by some short bristly feathers reflected over them: tarsi robust: hind claw strongest, and most hooked: wings with the first quill of moderate length; second shorter than the third; fourth and fifth longest.

(1. *PARUS*.) Bill moderate: tail slightly forked.

(2. *MECISTURA*.) Bill very short: tail elongated, cuneiform.

25. *CALAMOPHILUS*.—Bill short, subconical, very slightly compressed; the upper mandible convex above, curved at the extremity, and projecting over the lower one, which is nearly straight: nostrils covered by reflected bristles: tarsi slender: wings with the first quill very short, almost abortive; fourth and fifth longest: tail elongated, cuneiform.

(AMPELIDÆ.)

26. BOMBYCILLA.—Bill strong, short, and straight, broad at the base; upper mandible convex above, slightly bending at the tip, and emarginated: nostrils oval, covered by small hairy feathers directed forwards: tarsus shorter than the middle toe: wings long, and pointed; first and second quills longest; secondaries tipped with wax-like appendages.

II. CONIROSTRES.—*Bill moderate or elongated, strong, more or less conic, entire or slightly emarginated: feet robust; toes three before and one behind.*

(FRINGILLIDÆ.)

27. ALAUDA.—Bill short, subconic; mandibles of equal length; the upper one convex, and slightly curved: nostrils basal, oval, covered by small bristly feathers directed forwards: hind claw nearly straight, longer than the toe: wings long; first quill almost abortive; second a little shorter than the third, which is longest: wing-coverts shorter than the quills.

28. EMBERIZA.—Bill short, strong, conic, sharp-pointed; the mandibles a little distant from each other, and forming an angle at the gape, compressed in front, with the tomia bending inwards; upper mandible smaller and narrower than the lower; palate furnished with a hard bony knob.

- (1. PLECTROPHANES.) Wings long, and acuminate; first and second quills longest, and nearly equal: hind claw produced, and nearly straight.
- (2. EMBERIZA.) Wings moderate; first quill shorter than the second and third, which are longest: hind claw short and hooked.

29. FRINGILLA.—Bill thick, strong, more or less perfectly conic: nostrils basal, lateral, partly concealed by short bristly feathers directed forwards: tarsi short, or

moderate: wings with the first three quills nearly equal; second and third generally a little the longest: tail more or less forked.

- (1. FRINGILLA.) Bill straight, perfectly conic, the cone rather elongated, sharp-pointed; mandibles nearly equal, with the tomia of the lower one a little inflected: first quill a little shorter than the second and third, which are longest and equal.
- (2. PYRGITA.) Bill strong, conic; the culmen somewhat arched, and the tip a little obtuse; lower mandible rather smaller than the upper: three first quills nearly equal, longer than the fourth.
- (3. COCCOTHAUSTES.) Bill conic, very thick and strong; the sides bulging, and the culmen much rounded: lower mandible nearly equal to the upper, with the tomia inflected: wings rather long, acuminate; first and fourth quills equal, shorter than the second and third, which are longest.
- (4. CARDUELIS.) Bill straight, perfectly conic, the cone rather elongated, much compressed at the tip, and sharp-pointed; upper mandible angulated at the base, and slightly sinuated: wings moderate; first quill a little shorter than the second and third, which are nearly equal and longest.
- (5. LINARIA.) Bill straight, perfectly conic, short, compressed anteriorly, sharp-pointed; commissure straight; wings long, and acuminate; first three quills equal, longer than the fourth.

30. PYRRHULA.—Bill short and thick, the sides inflated and bulging; upper mandible convex above, advancing on the forehead at the base, deflected at the tip and overhanging the lower one: nostrils concealed by hairy feathers directed forwards: wings rather short; first quill shorter than the second, which is longest; third and fourth nearly equal to the second.

31. LOXIA.—Bill moderate, strong, thick at the base, much compressed anteriorly; both mandibles equally convex, hooked at the tips, and crossing each other when at rest; the tomia bending inwards: nostrils round, concealed by bristly feathers directed forwards: wings with the first and second quills equal, shorter than the third, which is longest.

(STURNIDÆ.)

32. STURNUS.—Bill moderate, straight, subconic; depressed, especially at the tip, which is rather blunt;

upper mandible advancing on the forehead, the edges rather dilated and extending beyond those of the lower: nostrils basal, half closed by an arched membrane: tarsus longer than the middle toe: wings long; first quill very short; second longest; the rest gradually decreasing.

33. PASTOR.—Bill subconic, compressed, slightly arched, the tip somewhat emarginated: nostrils basal, oval, partly covered by a plumose membrane: feet robust; tarsus much longer than the middle toe: wings with the first quill very short, almost abortive; second and third longest.

(CORVIDÆ.)

34. FREGILUS.—Bill long, rather slender, arched from the base, a little compressed, subulate and pointed at the extremity: nostrils basal, oval, concealed by reflected bristles: feet robust; tarsus longer than the middle toe: wings long; the fourth and fifth quills longest: tail square.

35. CORVUS.—Bill strong, thick, convex above, compressed at the sides, with the tomia sharp and cutting; upper mandible slightly bending towards the tip; lower one nearly straight: nostrils basal, oval, generally concealed by setaceous feathers directed forwards: wings acuminate; first quill much shorter than the second and third; fourth longest.

(1. CORVUS.) Tail moderate, rounded.

(2. PICA.) Tail long, cuneiform.

36. GARRULUS.—Bill rather short, strong, compressed, clothed at the base with feathers directed forwards; both mandibles inclining equally towards the tip, the upper one slightly emarginated: crown feathers long, and capable of erection: wings rounded; fifth and sixth quills longest: tail moderate, square, or slightly rounded.

37. NUCIFRAGA.—Bill long, and straight; upper mandible rounded, and longer than the lower one; both of them terminating in a slightly obtuse and depressed point:

nostrils basal, round, covered by reflected feathers: wings rather acuminate; first quill shorter than the second and third; fourth longest: tail rounded.

III. SCANSORES.—*Bill various: feet short; adapted for climbing; toes four, two each way, or three in front and one behind.*

* *Bill straight, robust: feet zygodactyle.*

(PICIDÆ.)

38. PICUS.—Bill robust, long, straight, angular, compressed, cuneate at the tip: tongue long and extensible, lumbriciform, the tip barbed: nostrils concealed by reflected bristly feathers: wings with the first quill very short; fourth and fifth longest: feet robust: tail feathers stiff, and pointed at the extremity.

39. YUNX.—Bill short, straight, somewhat conical, depressed, rounded above, the tip sharp and pointed: tongue long and extensible, lumbriciform: nostrils naked, partly closed by a membrane: wings moderate; first quill a little shorter than the second, which is longest: tail feathers soft and flexible.

** *Bill generally slender: feet anisodactyle.*

(CERTHIIDÆ.)

40. CERTHIA.—Bill moderate, slender, arcuate, compressed, angulated above, sharp-pointed: tongue short: nostrils naked, partly covered by an arched membrane: wings with the first quill very short; fourth longest: hind toe strong, and longer than the others: tail feathers stiff and pointed at the extremity, deflected.

41. TROGLODYTES.—Bill slender, slightly compressed, a little arcuate: nostrils basal, oval, partly covered by an arched naked membrane: wings short, and rounded; first quill very short; second shorter than the third; fourth and fifth equal and longest: tail short, rounded, carried erect.

42. UPUPA.—Bill long and slender, arcuate, compressed, convex above, sharp-pointed: nostrils oval, open: crown furnished with a crest: wings rather long; the first quill very short; fourth and fifth longest: claws short, and not much hooked: tail square at the extremity.

43. SITTA.—Bill moderate, rather strong, straight, compressed; both mandibles equally inclining to the tip which is somewhat cuneated: nostrils oval, open, covered by setaceous feathers directed forwards: wings rather short; third and fourth quills longest: hind toe strong, and lengthened: tail short, nearly even.

*** *Bill more or less curved: feet zygodactyle.*

(CUCULIDÆ.)

44. CUCULUS.—Bill as long as the head, rather compressed, moderately curved, and a little hooked at the tip: gape wide: nostrils round, surrounded by a naked and prominent membrane: wings long, acuminate: first quill short; third longest; tarsi short, feathered a little below the knee: outer hind toe partly reversible: tail long, more or less cuneated.

45. COCCYZUS.—Bill strong, arched, the culmen convex, triangular at the base, compressed at the sides and tip: nostrils basal, longitudinally cleft in a membrane: orbits naked: wings short, and concave: tarsi very long: tail long, and cuneated.

IV. FISSIROSTRES.—*Bill broad at the base; gape wide: wings long: feet short, and weak.*

* *Bill strong, more or less elongated.*

(MEROPIDÆ.)

46. CORACIAS.—Bill moderate, robust, higher than broad, compressed, straight; upper mandible bending at the tip: nostrils basal, linear, pierced diagonally, half closed by a plumose membrane: toes three before and one

behind, entirely divided: wings long, and acuminate; first quill shorter than the second, which is longest.

47. MEROPS.—Bill longer than the head, triangular at the base, slightly curved, carinated above, sharp-pointed: nostrils basal, oval, open: toes three before and one behind; the outer toe connected with the middle one as far as the second joint, the middle with the inner one as far as the first: wings long, and pointed; first quill very short, the second longest.

(HALCYONIDÆ.)

48. ALCEDO.—Bill very long, straight, quadrangular, thick and pointed: nostrils basal, pierced obliquely, linear, almost closed by a naked membrane: toes three before and one behind; the outer toe connected with the middle one as far as the second joint, the middle with the inner one as far as the first: wings moderate; the second and third quills equal and longest: tail short.

*** Bill short and weak: gape extremely large.*

(HIRUNDINIDÆ.)

49. HIRUNDO.—Bill short, and much depressed; upper mandible carinated, deflected at the tip; gape extending nearly to the eyes: toes three before and one behind; the outer toe connected with the middle one as far as the first joint: wings very long; first quill longest: tail forked.

50. CYPSELUS.—Bill very short, triangular, depressed at the base; the upper mandible deflected at the tip; gape extending to the posterior angle of the eye: toes divided to their origin; all directed forwards: wings extremely long; first quill a little shorter than the second: tail forked.

(CAPRIMULGIDÆ.)

51. CAPRIMULGUS.—Bill very short, rather curved, broad and depressed at the base; the upper mandible de-

flected at the tip, furnished at the basal edge with strong vibrissæ directed forwards; gape extending beyond the eyes: anterior toes connected by a membrane as far as the first joint; hind toe reversible: wings long; first quill shorter than the second, which is longest: tail rounded, or forked.

§ II. *Feet not formed for grasping.*

ORDER III. RASORES.

Bill short, or moderate, convex, often furnished with a cere at the base; upper mandible arched: wings short: legs strong and muscular, adapted for walking: hind toe sometimes wanting; when present, generally articulated high on the tarsus: claws robust, short, slightly curved, somewhat blunt.

I. COLUMBIDÆ.—*Hind toe present, articulated nearly on the same plane with the others: tarsi without spurs: tail of twelve feathers.*

52. COLUMBA.—Bill moderate, straight, compressed, deflected at the tip; upper mandible covered at the base with a soft tumid membrane, in which the nostrils are situate: tarsi short, reticulated; toes entirely divided.

(1. COLUMBA.) Wings with the first quill shorter than the second, which is longest: tail even: side toes equal.

(2. ECTOPISTES.) Wings pointed; two first quills equal, longer than the third: tail long, and cuneated: side toes unequal; the inner one longest.

* II. PHASIANIDÆ.—*Hind toe present, elevated above the others: tarsi generally armed with spurs: tail of more than twelve feathers: head more or less naked.*

*53. MELEAGRIS.—Bill short and thick, furnished at the base with an elongated, pendulous, fleshy appendage: head and neck naked;

throat with a pendulous carunculated wattle: tarsi of the male armed with spurs: tail broad, expansile, consisting of from fourteen to eighteen feathers.

*54. PAVO.—Bill naked at the base, thick, convex above, deflected at the tip: cheeks partially naked: head ornamented with a crest: tarsi of the male spurred: tail of eighteen feathers: upper tail-coverts longer than the tail, broad and expansile, ocellated.

*55. GALLUS.—Bill smooth at the base, thick, slightly curved: nostrils covered by an arched scale: generally an erect fleshy crest on the head; throat with fleshy wattles on each side of the lower mandible: ears naked: tarsi with strong spurs: anterior toes united by a membrane as far as the first joint: tail of fourteen feathers, compressed, more or less arched, ascending.

*56. PHASIANUS.—Bill short and thick, naked at the base; upper mandible very convex, with the tip deflected: nostrils basal, lateral, covered by an arched membrane: cheeks naked, adorned with scarlet papillæ: ears covered: tarsi of the male spurred: anterior toes united by a membrane as far as the first joint: wings short; fourth and fifth quills longest: tail very long, cuneated, of eighteen feathers.

*57. NUMIDA.—Bill thick, covered at the base with a warty membrane, in which the nostrils are placed: head naked, the crown with a callous horny protuberance; beneath the cheeks pendulous carunculated wattles: tarsi without spurs: anterior toes united by a membrane as far as the first joint: tail short, bent down, of fourteen or sixteen feathers.

III. TETRAONIDÆ.—*Hind toe short, and weak; sometimes altogether wanting: tarsi generally armed with spurs: tail short.*

58. TETRAO.—Bill short and strong, convex above, bending towards the tip: nostrils partly closed by an arched scale, concealed by the frontal feathers: eyebrows naked, adorned with scarlet papillæ: tarsi feathered; without spurs: wings with the first quill much shorter than the second; third and fourth longest: tail of sixteen or eighteen feathers.

(1. TETRAO.) Anterior toes naked, with pectinated margins: hind toe longer than the nail: tail broad, and slightly rounded, or lyrate.

(2. LAGOPUS.) Anterior toes feathered, the margins not pectinated: hind toe shorter than the nail: tail nearly even.

59. PERDIX.—Bill short, strong, naked at the base, convex above, deflected towards the tip: nostrils basal, half closed by an arched naked scale: tarsi naked: feet with

four toes, naked: wings short, and concave; the fourth and fifth quills generally longest: tail short, of from twelve to eighteen feathers.

- (1. PERDIX.) Bill short, and strong: orbits naked: tarsus in the male armed with a blunt tubercle: tail short, and bent down.
- *(2. ORTYX.) Bill thick, strong, higher than broad: orbits feathered: tarsi unarmed: wings with the first quill very short; fifth longest: tail short or moderate, of twelve feathers.
- (3. COTURNIX.) Bill short, and slender: orbits feathered: tarsi unarmed: wings with the first quill longest: tail very short, almost concealed by the upper coverts.

IV. STRUTHIONIDÆ.—*Hind toe wanting: tarsi without spurs: wings very short, often unfit for flight.*

60. OTIS.—Bill moderate, subconic, nearly straight, compressed; the upper mandible arched towards the tip: nostrils oval, open, a little remote from the base: legs long, naked above the knee; toes connected by a membrane at the base: wings moderate; third quill longest.

ORDER IV. GRALLATORES.

Bill various: legs moderate or elongated, slender, with the lower part of the tibia generally naked, adapted for wading: toes long, three or four in number, more or less connected by a membrane at the base, sometimes lobated.

I. CHARADRIIDÆ.—*Bill short or moderate, rarely elongated; robust or slender: legs moderate, or elongated: toes three, all directed forwards; rarely the rudiment of a fourth; the outer and middle toes generally united at the base by a membrane.*

61. CURSORIUS.—Bill shorter than the head, depressed at the base, somewhat arched and bent down to-

wards the tip, pointed: nostrils basal, oval, covered above by a protuberant membrane: legs long and slender: toes three, very short; almost divided to their origin: claws very small, the middle one serrated: wings moderate; second quill longest.

62. **ÆDICNEMUS**.—Bill longer than the head, straight, very robust, a little depressed at the base, compressed laterally; culmen elevated towards the tip; lower mandible angulated beneath: nostrils medial, longitudinally cleft: tarsi long and slender: toes three, united by a membrane at the base: wings moderate; second quill longest.

63. **CHARADRIUS**.—Bill shorter than the head, straight, slender, compressed, somewhat tumid and enlarged towards the tip: nasal channel extending two-thirds of its length, covered by a large membrane; nostrils basal, linear, pierced in the membrane: tarsi moderate, slender: toes three; outer and middle ones connected by a short membrane; inner toe free: wings moderate; first quill longest.

64. **VANELLUS**.—Bill moderate, straight, compressed, slender at the base, tumid towards the extremity; upper mandible slightly bending at the tip: nasal channel large, covered by a membrane; nostrils linear, pierced in the membrane: tarsi moderate, slender: toes four; outer and middle toes connected at the base; hind toe small, and elevated: wings long and acuminate.

(1. **SQUATAROLA**.) Nasal channel extending half the length of the bill: tarsi reticulated: hind toe rudimentary: first quill longest.

(2. **VANELLUS**.) Nasal channel two-thirds the length of the bill: acrotarsia scutellated: hind toe moderately developed: fourth and fifth quills longest.

65. **STREPSILAS**.—Bill moderate, strong, nearly straight, forming a lengthened cone, with the culmen a little depressed, and the apex subtruncated; the commissure slightly ascending: nasal channel half the length of the bill: toes four; the anterior ones almost entirely divided; hind toe moderately developed, touching the ground: wings long; first quill longest.

66. CALIDRIS. — Bill moderate, straight, slender and flexible throughout, compressed at the base, depressed and dilated towards the tip: nasal groove extending nearly the whole length; nostrils lateral, linear: tarsi moderate, slender: toes three, divided nearly to their origin: wings moderate; first quill longest.

67. HÆMATOPUS. — Bill long, robust, straight, compressed, the tip cuneated: nostrils lateral, nearly basal, oblong-linear, placed in a groove: tarsi moderate, robust: toes three; the outer and middle ones connected as far as the first joint; all the toes bordered by a narrow membrane: wings long; first quill longest.

II. GRUIDÆ.—*Bill strong, moderate or elongated: legs long: toes four in number; the outer and middle ones united at the base by a small membrane; hind toe short, elevated on the tarsus.*

68. GRUS.—Bill a little longer than the head, straight, compressed, deeply channelled at the base of the upper mandible on each side, the tip forming a lengthened cone: nostrils medial, placed in the lateral groove, closed behind by a membrane: head generally more or less naked: a considerable part of the tibia naked: wings moderate; third quill longest.

III. ARDEIDÆ.—*Bill strong, elongated: legs long: feet with four toes; anterior ones more or less united at the base; hind toe long, resting on the ground.*

69. ARDEA.—Bill as long as, or a little longer than the head, strong, straight or very slightly inclined, thick at the base, gradually tapering towards the tip; upper mandible channelled for about two-thirds of its length: nostrils nearly basal, placed in the groove, and partly closed by a membrane: orbits and lore naked:

lower part of the tibia more or less naked: middle claw with the inner margin pectinated: wings ample; second and third quills longest.

- (1. ARDEA.) Bill longer than the head, the upper mandible nearly straight: occiput usually ornamented with a pendent crest: neck slender, elongated.
- (2. BOTAURUS.) Bill scarcely longer than the head, higher than broad at the base, very much compressed; upper mandible slightly curved: neck thick, rather short; the feathers on the fore part loose and elongated, capable of erection; those behind short and downy.
- (3. NYCTICORAX.) Bill scarcely longer than the head, higher than broad, much compressed; upper mandible slightly curved: occiput with three very long subulate feathers: neck thickish, rather short.

70. CICONIA. — Bill much longer than the head, straight, robust, cylindrical, acute, forming a lengthened cone; upper mandible convex above; under mandible inclining a little upwards at the tip: nostrils nearly basal, open, longitudinally pierced in the horny substance of the bill: orbits naked: tarsi very long: a considerable part of the tibia naked: claws short, depressed, not pectinated: wings moderate; third, fourth, and fifth quills longest.

71. PLATALEA. — Bill very long, robust, much depressed, dilated and rounded at the extremity in the form of a spatula: nostrils dorsal, oblong, open, bordered by a membrane: head and face more or less naked: tibiæ semiplumed: tarsi long: the three anterior toes united as far as the second joint by a deeply cut membrane: wings moderate; second quill longest.

72. IBIS. — Bill very much elongated, slender, arcuate, broad at the base, somewhat quadrangular, the tip depressed, obtuse, and rounded; upper mandible deeply channelled throughout its whole length: nostrils nearly basal, placed in the groove, oblong, narrow, surrounded by a membrane: face and throat more or less naked: legs long, and slender; lower part of the tibia naked: wings moderate; second and third quills longest.

IV. SCOLOPACIDÆ.—*Bill long and slender; legs moderate, or elongated: toes four in number; anterior ones entirely divided, or united by a small basal membrane; hind toe weak and elevated, very rarely altogether wanting.*

73. NUMENIUS.—Bill much longer than the head, slender, arcuate, rounded, the tip rather obtuse; upper mandible projecting beyond the lower, channelled throughout three-fourths of its length: nostrils basal, oblong-linear, placed in the lateral groove: lore, and rest of the face feathered: legs slender; lower part of the tibia naked: anterior toes connected as far as the first joint: wings moderate; first quill longest.

74. TOTANUS.—Bill long or moderate, slender, generally straight, rarely inclining a little upwards, soft at the base, solid and sharp-pointed at the extremity; upper mandible channelled through half its length or more, slightly bending over the lower one at the tip: nostrils lateral, linear, placed in the groove: legs long, slender, with the lower part of the tibia naked: anterior toes, or the outer ones only, connected by a small membrane.

75. RECURVIROSTRA.—Bill very long, slender, weak, depressed, pointed, flexible at the tip, and considerably recurved; upper mandible channelled on each side: nostrils long, linear, placed at the base of the groove: legs long and slender; lower half of the tibia naked: anterior toes connected by a membrane as far as the second joint; hind toe very small.

76. HIMANTOPUS.—Bill very long, slender, attenuated, nearly straight, rounded, a little compressed at the tip; mandibles channelled laterally for half their length: nostrils linear, elongated: legs very long and slender: nearly the whole of the tibia naked: outer and middle toes connected by a broad membrane; hind toe wanting: wings very long; first quill longest.

77. LIMOSA.—Bill very long, slender, soft and flexible throughout, slightly recurved, depressed, the tip obtuse and somewhat dilated; upper mandible channelled for nearly its whole length: nostrils linear, pierced in the membrane of the groove: legs long and slender; lower part of the tibia naked: outer and middle toes connected by a membrane as far as the first joint: wings moderate: first quill longest.

78. SCOLOPAX.—Bill very long, straight, slender, soft, the tip obtuse and dilated; upper mandible a little longer than the lower; both mandibles channelled for nearly two-thirds of their length: nostrils lateral, basal, linear, covered by a membrane: legs moderate, slender; naked space on the tibia very small: wings with the first quill longest; second nearly equal to it.

- (1. *RUSTICOLA*.) Tibia feathered to the joint: tarsi short: anterior toes entirely divided: hind claw shorter than the toe.
- (2. *SCOLOPAX*.) Lower part of the tibia naked: tarsi more elongated: toes entirely divided, or united at the base by an extremely short membrane: hind claw longer than the toe.
- (3. *MACRORAMPHUS*.) Lower part of the tibia naked: tarsi rather elongated: outer toe united to the middle one as far as the first joint: hind claw longer than the toe.

79. TRINGA.—Bill as long as, or a little longer than the head; straight, or slightly curved, soft and flexible throughout, compressed at the base, depressed and dilated at the extremity; upper mandible longer than the lower, with the nasal channel extending nearly to the tip: nostrils lateral, short, pierced in the membrane of the groove: tarsi moderate, or elongated: wings with the first quill longest.

- (1. *MACHETES*.) Tarsi elongated: outer and middle toes united by a membrane as far as the first joint: neck (in the male) with an ornamental ruff during the breeding season.
- (2. *TRINGA*.) Tarsi moderate: anterior toes divided to their origin: hind toe very small,

80. LOBIPES.—Bill long, slender, weak, a little depressed at the base, subulate at the tip; upper mandible slightly curved, and channelled throughout: tongue slender,

and pointed: nostrils basal, oblong-linear, prominent, surrounded by a membrane: tarsi somewhat elongated, and compressed: anterior toes rather long, bordered by narrow lobated membranes: wings with the first quill longest.

81. PHALAROPUS.—Bill as long as the head, straight, strong, broad and depressed at the base, a little compressed at the extremity; upper mandible inclining over the lower, with the nasal channel extending the whole length: tongue short, broad, and rounded at the tip: nostrils basal, oval, prominent, surrounded by a membrane: tarsi short and strong, scarcely compressed: anterior toes rather short, united as far as the first joint; bordered beyond with broad and deeply scalloped membranes: wings with the first and second quills longest.

V. RALLIDÆ.—*Bill short or moderate, rarely elongated; rather strong: body generally compressed: legs moderate: toes four; anterior ones elongated, entirely divided, or united by a membrane at the base, sometimes pinnated.*

82. GLAREOLA.—Bill short, convex, compressed towards the tip; the upper mandible curved for half its length: nostrils basal, lateral, obliquely cleft: a small part of the tibia naked: tarsi moderate, slender: outer and middle toes connected by a short membrane; hind toe small, and elevated: claws long, and pointed: wings very long; first quill considerably the longest.

83. RALLUS.—Bill longer than the head, rather slender, very slightly arched, compressed at the base, somewhat cylindrical at the tip; upper mandible channelled for two-thirds of its length: nostrils submedial, oblong-linear, placed in the groove, partly covered by a membrane lower: part of the tibia naked: toes long, and slender; anterior ones entirely divided: wings short, rounded; first quill shorter than the two next, which are longest: bastard winglet with a short spur.

84. CREX.—Bill shorter than the head, somewhat thick, straight, convex above, higher than broad at the base, attenuated and compressed at the tip; upper mandible a little deflected; gonys of the lower mandible ascending to meet it: nostrils lateral, medial, longitudinally cleft, half closed by the membrane of the nasal channel: anterior toes long, entirely divided: wings short; first quill shorter than the second and third, which are longest.

85. GALLINULA.—Bill shorter than the head, straight, robust, convex above, much compressed; upper mandible expanding at the base on the forehead, forming a naked disk: nostrils lateral, medial, oblong-linear: anterior toes very long, divided to their origin, bordered by a narrow, entire membrane: wings with the second and third quills longest.

86. FULICA.—Bill shorter than the head, straight, robust, somewhat convex above, compressed, thick at the base: a naked disk on the forehead: nostrils lateral, medial, longitudinally cleft: anterior toes very long, united at the base, bordered by a scalloped membrane: wings moderate: second and third quills longest.

ORDER V. NATATORES.

Bill various: legs short, often placed far behind, adapted for swimming: lower part of the tibia more or less naked: tarsi compressed: toes generally four; the three anterior ones palmated, semi-palmated, or fissopalmated; or all four united: hind toe sometimes wanting.

I. ANATIDÆ.—*Bill stout, straight, covered with a thin membranous skin; the tomia armed with lamellæ, or with small denticulations; more or*

less depressed; the tip rounded and obtuse, furnished with a nail: wings moderate: legs placed in or near the equilibrium: feet four-toed, palmed; hind toe free, placed high on the tarsus.

* (ANSERINÆ.)

87. ANSER.—Bill as long as, or shorter than the head, higher than broad at the base, subconic; the upper mandible deflected at the tip; the lower one flat and straight, narrower: nostrils a little behind the middle: neck of moderate length: wings tuberculated, or spurred: legs in the equilibrium: tarsi slightly elongated.

- (1. ANSER.) Bill as long as the head; the tips of the lamellæ exerted, appearing like sharp teeth.
- (2. BERNICLA.) Bill shorter than the head, somewhat slender; the edges of the mandibles concealing the tips of the lamellæ.
- (3. PLECTROPTERUS.) Bill moderate; furnished with a fleshy tubercle at its base: wings armed with spurs.

88. CYGNUS.—Bill moderate, of nearly equal breadth throughout, furnished at its base with a knob or fleshy tumour, convex above; the upper mandible deflected at the tip; the lower one flat: nostrils medial: neck very long: legs a little beyond the equilibrium: tarsi short.

** (ANATINÆ.) — *Hind toe simple: tarsi somewhat round.*

89. TADORNA.—Bill shorter than the head, with an elevated tubercle at the base, depressed or concave in the middle, of nearly equal breadth throughout; the upper mandible laterally grooved near the tip, which is slightly recurved: face and lore feathered: nostrils large, oval, a little before the base: tail even.

* 90. CAIRINA.—Bill furnished with an elevated tubercle at the base; the edges of the mandibles sinuated: face and lore covered with a naked warty skin: nostrils round, basal: wings tuberculated.

91. ANAS.—Bill as long as, or rather longer than the head, simple at the base, broad and depressed generally throughout its whole length, sometimes widening ante-

riorly: nostrils small, oval, a little before the base: tail more or less acute, sometimes elongated.

- (1. ANAS.) Bill longer than the head, semicylindric at its origin, depressed and very much dilated towards the tip; tomia furnished with long pectinated lamellæ, projecting far beyond the margin: tail a little longer than the wings, of fourteen feathers, moderately acute.
- (2. CHAULIOPDUS.) Bill the length of the head, of nearly equal breadth throughout; pectinated lamellæ distinct, appearing below the edge of the upper mandible: tail slightly lengthened: the two middle feathers pointed, and rather longer than the others.
- (3. DAFILA.) Bill full as long as the head; the upper mandible of equal breadth throughout; lamellæ not projecting beyond the margin: tail of sixteen feathers, elongated, acute.
- (4. BOSCHAS.) Bill as long as the head, of nearly equal breadth throughout; the denticulations of the upper mandible scarcely projecting beyond the margin: tail of sixteen feathers, moderate, rather acute.

92. MARECA.—Bill shorter than the head, higher than broad at the base, depressed and narrowing towards the tip; lamellæ slightly projecting: nostrils small, oval, a little before the base: tail of fourteen feathers, short, acute.

* 93. DENDRONESSA.—Bill shorter than the head, as broad as high at the base, towards the tip narrow and contracted: nostrils large, pervious, submedial: head with an occipital crest: tertials ornamented: tail of sixteen feathers, rounded.

*** (FULIGULINÆ.)—*Hind toe with a lobated membrane: tarsi compressed.*

94. SOMATERIA.—Bill swollen and elevated at the base, nearly straight, extending on the forehead, where it is divided by an angular projection of feathers, narrow and obtuse anteriorly, the nail strong and hooked; tomia set with coarse distant lamellæ: nostrils small, oval, medial: tail of fourteen feathers.

95. OIDEMIA.—Bill short, broad, with an elevated tumour at the base, towards the tip much depressed and flattened; the nail obtuse and rounded, slightly deflected; lamellæ coarse, widely set, scarcely projecting: nostrils oval, submedial, elevated: tail short, graduated, acute.

96. **FULIGULA.**—Bill as long as the head, simple at the base or very slightly elevated, broad and very much depressed anteriorly, a little dilated towards the tip; upper lamellæ not projecting beyond the edges of the mandible: nostrils rather small, oblong-oval, a little before the base: wings and tail short, the latter generally of fourteen acute feathers.

97. **CLANGULA.**—Bill shorter than the head, elevated at the base, much more contracted than in the last genus, and becoming continually narrower towards the tip; lamellæ not projecting: nostrils roundish-oval, medial: tail of moderate length, graduated, the feathers semi-acute.

98. **HARELDA.**—Bill much shorter than the head, rather high at the base, slender, suddenly contracting towards the tip; nail strong and arched; upper lamellæ projecting considerably below the edge of the mandible: nostrils large, oblong-linear, a little before the base: forehead high: tail of fourteen feathers; the central pair acute, and very much elongated.

**** (MERGANINÆ.)

99. **MERGUS.**—Bill short or moderate, straight, slender, narrow, approaching to cylindrical; upper mandible high at the base, tapering towards the tip, which is armed with a strong hooked nail; tomia of both mandibles set with acute teeth directed backwards: nostrils oblong-oval, pervious, medial: legs short, placed far behind: tarsi compressed: hind toe lobated: tail of twelve feathers, rounded.

II. **COLYMBIDÆ.**—*Bill slightly compressed, smooth, not covered by a membranous skin; the tomia simple, or only slightly denticulated: wings short: legs placed at the extremity of the body: tarsi very much compressed: feet four-toed, palmated, or fissopalmated: hind toe free.*

100. **PODICEPS.**—Bill moderate, straight, robust, forming a lengthened cone, sharp-pointed; upper mandible

slightly inclining at the tip ; lower mandible with the gonys ascending : feet short ; anterior toes fissopalmed ; hind toe lobated : claws broad, and much depressed : wings with the three first quills nearly equal, and longest : tail wanting.

101. COLYMBUS.—Bill rather longer than the head, straight, robust, nearly cylindrical, sharp-pointed ; the tomia of both mandibles inflected ; gonys ascending : feet short ; anterior toes long, entirely palmed ; hind toe short, and lobated : claws depressed : wings with the first or second quill longest : tail very short, rounded.

III. ALCIDÆ.—*Bill very much compressed, often transversely grooved at the sides : wings short, sometimes wholly unfit for flight : legs placed at the extremity of the body : feet three-toed, entirely palmed.*

102. URIA.—Bill moderate, strong, straight, convex above, much compressed at the sides, sharp-pointed ; upper mandible slightly deflected at the tip ; gonys of the lower mandible ascending ; tomia curving inwards from the base to the middle : nostrils basal, lateral, pervious, elliptic-linear, half concealed by a plumose membrane : tarsi very much compressed : wings very short ; first quill longest : tail short, of twelve feathers.

103. MERGULUS.—Bill shorter than the head, rather thick, convex, conical, slightly curved, covered at the base with downy feathers ; both mandibles notched at the tip : nostrils basal, lateral, round, partly concealed by a plumose membrane : wings not reaching to the tail ; first and second quills equal, and longest : tail short, of twelve feathers.

104. FRATERCULA.—Bill shorter than the head, thick at the base, very much compressed anteriorly, higher than long, the culmen more elevated than the crown of the head ; both mandibles arched, transversely furrowed, and notched at the tip : nostrils basal, marginal, oblong-linear,

almost entirely closed by a large naked membrane: wings short, narrow, and acuminate; first and second quills equal, and longest: tail short, of sixteen feathers.

105. ALCA.—Bill a little shorter than the head, rather longer than high, much compressed, cultrated; both mandibles laterally suleated; upper one hooked at the extremity; lower one with the gonys ascending, the mental angle prominent and rounded: nostrils marginal, nearly concealed by the advancing frontal feathers: wings short, and narrow: tail short, of twelve to sixteen feathers.

IV. PELECANIDÆ.—*Bill strong, sometimes compressed; the tomia denticulated: wings long, and powerful: legs very short, in or near the equilibrium: feet with four toes, all united by one continuous membrane.*

106. PHALACROCORAX.—Bill a little longer than the head, straight, compressed, culmen rounded; upper mandible with a longitudinal furrow on each side, the dertrum distinct and strongly hooked; lower one with the tip truncate: nostrils concealed: face and throat naked: tibiæ feathered: middle claw with the inner margin denticulated: wings moderate; first quill shorter than the second, which is longest: tail rounded.

107. SULA.—Bill a little longer than the head, robust, straight, large at the base, compressed and attenuated towards the tip; culmen rounded; upper mandible laterally suleated, the tip slightly hooked: gape extending beyond the eyes: face naked: nostrils concealed: tibiæ naked at bottom: middle claw with the inner margin denticulated: wings long, and acuminate; first quill longest, or equal to the second: tail graduated.

V. LARIDÆ.—*Bill without denticulations: wings very long: legs in or near the equilibrium:*

three anterior toes united ; hind toe free, sometimes represented by a simple nail, or altogether wanting.

108. STERNA.—Bill as long as, or a little longer than the head, nearly straight, rather compressed, slender, attenuated towards the tip, sharp-pointed ; mandibles of equal length, the upper one slightly bending at the tip, lower one with the mental angle a little projecting : nostrils longitudinal, oblong-linear, pervious : feet weak, small, four-toed ; membranes deeply notched : wings long, acuminate ; first quill longest : tail generally forked.

(1. STERNA.) Wings very long : tail deeply forked.

(2. ANOUS.) Wings moderate : tail even.

109. LARUS.—Bill moderate, strong, cultrated, much compressed ; the culmen continuous, bending downwards at the tip ; gonys obliquely ascending, straight ; mental angle strongly projecting ; tomia sharp : nostrils longitudinal, medial, oblong-linear, pervious : legs moderate : anterior toes with the membranes entire ; hind toe small, high on the tarsus, rarely wanting : wings long ; first or second quill longest : tail generally even.

(1. XEMA.) Tail slightly forked.

(2. LARUS.) Tail even.

110. LESTRIS.—Bill moderate, strong, thick, rounded above, compressed towards the tip ; culmen and dertrum distinct, the latter convex and strongly hooked ; gonys straight, obliquely ascending ; mental angle projecting : nostrils immediately behind the dertrum, diagonal, narrow, closed behind, pervious : hind toe very small, nearly on a level with the others : wings with the first quill longest : two middle tail feathers elongated.

111. PROCELLARIA.—Bill more or less elongated, straight, subcylindrical, depressed and dilated at the base ; culmen and dertrum distinct ; the latter convex, much compressed, and strongly hooked ; lower mandible with a longi-

tudinal furrow on each side: gonys concave, more or less ascending: nostrils at the extremity of an open truncated tube extending along the culmen: a simple claw in the place of a hind toe: wings long; first or second quill longest.

- (1. PROCELLARIA.) Bill moderate, robust; lower mandible straight, somewhat truncated at the tip; mental angle projecting; nostrils opening by a single orifice, the tube extending half the length of the bill: tail slightly rounded.
 - (2. PUFFINUS.) Bill longer than the head, slender; lower mandible pointed, deflected at the tip, following the curvature of the upper; mental angle obsolete: nostrils in a double tube extending one-fourth of the length of the bill: tarsi moderate: tail rounded.
 - (3. THALASSIDROMA.) Bill shorter than the head, slender; lower mandible pointed, deflected at the tip, following the curvature of the upper: mental angle obsolete: nostrils opening by a single orifice, the tube extending half the length of the bill: tarsi slender, elevated: tail square, or slightly forked.
-

ORDER I. RAPTORES.

GEN. 1. NEOPHRON, *Sav.*

1. *N. Percnopterus*, *Sav.* (*Egyptian Neophron*).—Plumage above and below white; quills alone black: occipital feathers long, and filiform: tail cuneated.

N. Percnopterus, *Jard. and Selb. Orn.* pl. 33. *Cathartes Percnopterus*, *Temm. Man. d'Orn.* tom. i. p. 8. *Egyptian Neophron*, *Selb. Illust.* vol. i. p. 4. pl. A.

DIMENS. Entire length two feet seven inches: length of the bill (from the forehead) two inches six lines; of the tarsus three inches; of the middle toe, claw included, three inches: breadth, wings extended, five feet nine inches. **SELBY.**

DESCRIPT. (*Immature plumage.*) "Bill brownish black, or horn-coloured: cere somewhat bulging at the base, and occupying half the length of the bill, wine yellow: nostrils in the middle of the cere, large and open: crown of the head, cheeks, and throat, covered with a naked skin, of a livid flesh-coloured red, with a few straggling bristly feathers between the bill and the eyes, and upon the margins of the mandibles: ears round, open, and large: occiput and nape covered with a close, thick-set white down, with small black feathers intermixed: neck clothed with long, arched, and acuminate feathers, forming a kind of ruff of a deep umber brown, tipped with cream yellow: back and scapulars cream white, the latter intermixed and varied with umber brown: lesser wing-coverts nearest the body deep umber brown, margined with a paler shade; these are succeeded by two rows of cream-coloured sharp-pointed feathers: greater coverts umber brown, varied with cream white: secondaries pale umber brown, their tips and margins yellowish white: quills black: tail cuneiform, umber brown at the base, the tip yellowish white: under parts mixed with umber brown: legs strong and fleshy, of a pale yellowish gray: tarsi covered with a rough reticulated skin: middle toe with four entire scales upon the last phalanx; the exterior and interior each with three; hinder toe short and strong: claws blackish brown, strong, but not greatly arched." **SELBY.** In the *adult state*, the entire plumage is white; the greater quills alone excepted, which are black.

Common in Egypt, and in some parts of the Continent, but not known as a visitant in this country till 1825, in the Autumn of which year, two specimens were observed on the shores of the Bristol Channel near Kilve in Somersetshire, and one killed. From this individual, which proved to be in immature plumage, the above description was taken by Mr Selby. This species is said to feed principally upon carrion, but occasionally on lizards and other reptiles; more rarely attacks living birds and the smaller quadrupeds. Builds in the crevices and hollows of rocks.

GEN. 2. AQUILA, *Briss.*(1. AQUILA, *Cuv.*)

2. A. *Chrysaëtos*, Vigors. (*Golden Eagle*.) — Tail longer than the wings; rounded at the extremity: the last phalanges of all the toes with only three scales.

Faleo fulvus, *Temm. Man. d'Orn.* tom. 1. p. 38. Golden Eagle, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. 1. p. 12. pls. 1, 1*, and 2. *Bew. Brit. Birds*, vol. 1. p. 5.

DIMENS. Entire length three feet and a half: breadth eight feet. MONT.

DESCRIPT. (*Adult.*) Top of the head, and nape of the neck, bright rust-colour; the feathers long and acuminate: rest of the plumage dull brown, approaching to dusky: inside of the thighs, and feathers on the tarsi, light brown: tail deep gray, barred and tipped with blackish brown: bill bluish at the base, black at the extremity: irides brown: cere and feet yellow. (*Young.*) Plumage throughout of a uniform reddish brown: vent and under tail-coverts whitish: inside of the thighs, and feathers on the tarsi, white: tail white for two-thirds of its length from the base; the remainder dark brown. Perfect plumage not attained till the fourth year. (*Egg.*) Dirty white, mottled all over with pale reddish brown: longitudinal diameter three inches; transverse diameter two inches five lines.

Found principally in the mountainous parts of Scotland and Ireland: rare in England, but has been killed as far south as in Sussex. Preys on lambs, fawns, &c. as well as on the larger birds. Builds on rocks and tall trees, and lays two, rarely three, eggs. *Obs.* The Ring-tail Eagle (*F. fulvus*, Linn.) is the young of this species.

(2. HALIÆTUS, *Sav.*)

3. A. *Albicilla*, Briss. (*Cinereous Eagle*.) — Plumage brown; with spots of a darker tint in the immature state: tail not extending beyond the wings: all the phalanges of the toes scaled.

Faleo Albicilla, *Temm. Man. d'Orn.* tom. 1. p. 49. Cinereous Eagle, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. 1. p. 18. pls. 3, and 3*. *Bew. Brit. Birds*, vol. 1. p. 9.

DIMENS. Entire length two feet four to ten inches.

DESCRIPT. (*Adult.*) Head, and upper part of the neck, brownish ash: rest of the plumage on the body, including the wings, brown clouded with cinereous: tail white: bill whitish: cere and feet pale yellow. (*Young.*) Head and neck deep brown, with the tips of the feathers somewhat paler: upper parts in general reddish brown, each feather being pale at the root and dark along the shaft: quills dusky: under parts brown, with deeper coloured spots, and here and there a few white feathers: tail whitish at the base, irregularly spotted with brown on the outer webs, the tip also being of this colour: bill bluish black, paler at the base. (*Egg.*) Dirty white, with a few pale red marks: long. diam. three inches; trans. diam. two inches four lines.

More generally diffused than the last species. Most plentiful in Scotland, Ireland, and in the Orkney and Shetland Isles, but is occasionally met with, particularly in the immature state, in various parts of England. Feeds principally on fish, but attacks also birds and quadrupeds. Breeds generally on the most inaccessible cliffs, and lays two eggs. *Obs.* The Sea Eagle (*F. Ossifragus*, Linn.) is the young of this species.

(3. PANDION, Sav.)

4. *A. Haliaetus*, Meyer. (*Osprey*.) — Wings longer than the tail.

Falco Haliaetus, *Temm. Man. d'Orn.* tom. i. p. 47. *Osprey*, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. i. p. 24. pl. 4. *Bew. Brit. Birds*, vol. i. p. 13.

DIMENS. Entire length one foot eleven inches: breadth five feet four inches. **FLEM.**

DESCRIPT. (*Adult.*) Crown of the head and nape of the neck deep brown, the feathers, which are long and narrow, edged with white: a streak of blackish brown on each side of the neck, proceeding from the posterior angle of the eye, and reaching almost to the shoulders: upper parts in general deep umber-brown: under parts white, with some faint indications of a darker patch on the breast: tail-feathers transversely barred with white on their inner webs, the two middle ones excepted, which are wholly brown: bill blackish: irides yellow: cere and feet grayish blue. (*Young.*) Chiefly characterized by having the under parts more or less mottled; a large patch especially of tawny yellow on the breast, spotted with brown: feathers on the upper parts edged with reddish yellow. (*Egg.*) Reddish white: the larger end blotched with dark red brown, the smaller end spotted with the same colour: long. diam. two inches four lines; trans. diam. one inch ten lines.

Not a common species. Met with in Scotland, and in the West of England. Has occurred also in Hampshire, Hertfordshire, Norfolk, and Suffolk. Resides chiefly in the neighbourhood of large pieces of water, preying on fish. Builds on trees and rocks, and lays from three to four eggs.

GEN. 3. FALCO, Linn.

5. *F. Islandicus*, Lath. (*Jer-Falcon*.) — Plumage white, with dusky lines and spots.

F. Islandicus, *Temm. Man. d'Orn.* tom. i. p. 17. *Jer-Falcon*, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. i. p. 36. pl. 14. *Bew. Brit. Birds*, vol. i. p. 15.

DIMENS. Entire length (*male*) one foot ten inches, (*female*) two feet. **TEMM.**

DESCRIPT. (*Old male.*) Ground of the plumage white: upper parts marked with narrow streaks of brown; under parts with spots of the same colour, which become larger and more numerous on the sides: bill yellowish: cere and orbits livid yellow: irides brown: feet bright yellow. (*Old female, and immature male.*) Spots on the under parts more numerous; confluent on the sides, forming transverse bars: the markings above more extended, and occupying a larger portion of the entire plumage. (*Young of the year.*) Upper parts of a uniform brownish

ash, the white appearing only in small spots at the tips of the feathers: under parts with large brown spots disposed longitudinally upon a white ground: cere and orbits bluish: feet lead-colour, with a tinge of yellow. (*Egg*). "Spotted, of the size of a Ptarmigan's." FLEM.

Occurs in the northern parts of Scotland, particularly in the Orkney and Shetland Isles: but is of very rare occurrence in England. Sheppard mentions one that was shot on Bungay Common in Suffolk. Preys on birds and the smaller quadrupeds. *Obs.* The *Spotted Falcon* of Pennant is the young of this species.

6. *F. peregrinus*, Gmel. (*Peregrine Falcon*.)—Above bluish ash, with darker fasciæ; beneath yellowish white, with brown transverse bars; a broad black moustache: wings reaching to the extremity of the tail.

F. peregrinus, *Temm. Man. d'Orn.* tom. i. p. 22. *Peregrine Falcon*, *Mont. Orn. Dict. and Supp.* *Selb. Illust.* vol. i. p. 39. pls. 15 and 15*. *Bew. Brit. Birds*, vol. i. p. 17.

DIMENS. Entire length (*male*) fifteen inches, (*female*) seventeen inches; from the carpus to the end of the wing fourteen inches; tarsus two inches.

DESCRIPT. (*Adult male*.) Head, upper part of the neck, and a broad moustache descending from the corners of the mouth, blackish blue: rest of the upper parts bluish ash, with shades of a darker tint: throat and breast white, with a few fine longitudinal streaks; belly, vent, and thighs, grayish white, with brown transverse bars: quills dusky, the inner webs with a series of reddish white spots: tail alternately barred with black and gray: bill bluish at the base, black at the tip: cere, space surrounding the eyes, irides, and feet, yellow. (*Adult female*.) Colour of the upper plumage more dull: under parts with a faint reddish tinge. (*Young*.) Upper parts ash-brown, the feathers with pale reddish edges; region of the eyes, and moustaches, dusky, the latter ill-defined: throat white; rest of the under parts whitish, with large brown longitudinal spots of an oblong heart-shaped form: tail with irregular reddish bars; the tip white: irides brown. (*Egg*.) Mottled all over with two shades of red brown: long. diam. two inches one line; trans. diam. one inch eight lines.

Not very abundant in England: more plentiful in the northern parts of Scotland. Preys on birds. Builds on rocks, or in trees, and lays from three to four eggs. *Obs.* The *Lanner* of Pennant is the young of this species: the *Falco Lanarius* of Temminck is a distinct species, which has not hitherto been found in this country.

7. *F. Subbuteo*, Linn. (*Hobby*.)—Above bluish black; beneath reddish yellow, with longitudinal streaks of brown: a broad black moustache: wings reaching beyond the extremity of the tail.

F. Subbuteo, *Temm. Man. d'Orn.* tom. i. p. 25. *Hobby*, *Mont. Orn. Dict.* *Selb. Illust.* vol. i. p. 43. pl. 16. *Bew. Brit. Birds*, vol. i. p. 42.

DIMENS. Entire length twelve to fourteen inches: breadth two feet to two feet three inches.

DESCRIPT. (*Adult male*.) Upper parts bluish black: under parts whitish, passing into reddish yellow, with longitudinal brown streaks: throat white: a black patch or moustache proceeding from the corners of the mouth down each side of the neck: rump, thighs, and under tail-coverts, rust-red: quills, and outer tail-feathers, barred with reddish brown on their inner webs: bill bluish black: irides reddish brown: cere, eyelids, and feet, yellow. (*Female*.) Colours generally more obscure; upper parts approaching to dusky brown; the red on the rump and thighs not so bright; the spots underneath deeper. (*Young*.) Plumage above dusky, all the feathers edged with reddish yellow; this last colour most prevalent on the head: some white spots on the occiput: under parts yellowish white, passing into reddish yellow, with long streaks of brown: tip of the tail reddish: cere and feet greenish yellow. (*Egg*.) Yellowish white, speckled all over with reddish brown: long. diam. one inch eight lines; trans. diam. one inch four lines.

A summer visitant; arriving in April, and departing in October. Preys on larks, and other small birds. Builds in tall trees, and lays from three to four eggs.

8. *F. Æsalon*, Gmel. (*Merlin*.)—Above bluish ash, beneath reddish yellow, with longitudinal dark spots: no moustache: wings reaching to two-thirds the length of the tail.

F. Æsalon, *Temm. Man. d'Orn.* tom. i. p. 27. *Merlin*, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. i. pp. 46 and 48. *Selb. Illust.* vol. i. p. 51. pls. 18 and 18*.

DIMENS. Entire length eleven to twelve inches: breadth two feet.

DESCRIPT. (*Adult male*.) Upper parts, including the tail, bluish ash, with the shafts of the feathers black: throat white: under parts reddish yellow, with dark brown, oblong-oval spots: quills with the inner webs barred with white: a broad black bar near the extremity of the tail; the tip itself white: bill bluish: irides brown: cere, orbits, and feet, yellow. (*Female*.) Crown of the head dusky brown, streaked with black; nape white, with reddish brown spots: back and scapulars grayish brown, with reddish brown spots and black shafts: under parts yellowish white, with spots larger and more numerous than in the male bird: tail brown, with six or seven yellowish white bars. (*Young*.) Upper plumage brown, the feathers edged with red: tail dusky, with several narrow bars of reddish brown; the tip of this last colour: quills brownish black, spotted or barred with reddish brown: under parts as in the female. (*Egg*.) Mottled all over with two shades of dark red brown: long. diam. one inch seven lines; trans. diam. one inch three lines.

Supposed to be migratory in the South of England, which it visits during the winter months. Breeds in the counties of Northumberland, Cumberland, and Westmoreland, making its nest on the ground. Preys on small birds. *Obs.* This is a very variable species. The male in adult plumage is the *Stone-Falcon* of Latham, and other authors.

9. *F. rufipes*, Bechst. (*Red-legged Falcon*)—General plumage dark bluish gray, without spots: cere and feet

red: claws yellow: wings reaching to the extremity of the tail.

F. rufipes, *Temm. Man. d'Orn.* tom. i. p. 33. Orange-legged Hobby, *Lath. Syn. Supp.* II. p. 46. *Selb. Illust.* vol. i. p. 45. pl. B. Red-footed Falcon, *Gould, Europ. Birds.*

DIMENS. Entire length ten inches six lines. **TEMM.**

DESCRIPT. (*Adult male.*) Head, neck, breast, belly, and all the upper parts, dark lead-gray, without spots; thighs, vent, and under tail-coverts, deep ferruginous: cere, orbits, and feet, bright red: claws yellow, the tips brown. (*Female.*) Head and neck of a uniform reddish colour, without streaks; rest of the upper parts dusky blue; all the feathers, except the quills, edged with bluish black; under parts deep reddish, streaked with dusky brown: thighs red: tail bluish gray, with six or seven dusky bars, the tip also of this colour: cere, orbits, and feet, orange-red. (*Young.*) Top of the head brown, with dusky streaks; throat and ear-coverts white; eyes encircled with black; also a small black moustache extending from beneath the eye downwards: upper parts brown, the feathers with reddish brown edges; under parts yellowish white, with longitudinal brown streaks on the breast and abdomen: thighs without spots: tail with numerous alternate bars of brown and reddish white; the tip white. (*Egg.*) Unknown.

Three specimens of this species, unknown before as a native of this country, occurred at Horning in Norfolk, in May 1830. They consisted of an adult male, a female, and a young male in immature plumage. A fourth specimen was killed afterwards in Holkham Park, in the same county: and a fifth in Yorkshire. Said to feed on insects. Nidification unknown.

10. *F. Tinnunculus*, Linn. (*Kestrel*).—Upper plumage reddish brown, spotted with black: cere and feet yellow: claws black: wings reaching to three-fourths the length of the tail.

F. Tinnunculus, *Temm. Man. d'Orn.* tom. i. p. 29. *Kestrel*, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. i. pp. 39 and 41. *Selb. Illust.* vol. i. p. 47. pls. 17 and 17*.

DIMENS. Entire length fifteen inches: length of the bill (from the forehead) nine lines and a half, (from the gape) ten lines and a half; of the tarsus one inch eight lines; of the tail seven inches seven lines; from the carpus to the end of the wing nine inches ten lines: breadth, wings extended, two feet six inches.

DESCRIPT. (*Male.*) Crown of the head and nape bluish gray: upper parts reddish brown, with scattered black spots of an angular form: breast, belly, and thighs, white, inclining to reddish, with oblong brown spots: rump bluish gray: tail cinereous, with one broad black bar near the extremity: the tip white: bill bluish: cere, orbits, irides, and feet, yellow. (*Female.*) All the upper parts reddish orange, with transverse bars of dusky brown: tail reddish, with numerous black bars; one broad bar near the extremity, which last is reddish white. (*Young.*) Upper part of the head, and nape, reddish brown with dark streaks; rest of the plumage above with numerous arrow-shaped spots: a blackish streak under each eye, arising from the corners of the mouth, and extending to the neck: tail reddish, with transverse dusky bars in both sexes, as in

the adult female : irides brown : cere greenish yellow. (*Egg.*) Mottled all over with dark red brown : in some specimens blotches of red brown upon a pale reddish white ground : long. diam. one inch seven lines ; trans. diam. one inch three lines.

A common species ; preying principally on mice and insects. Builds in tall trees, and lays from three to five eggs. Hatches towards the end of April or beginning of May.

GEN. 4. ACCIPITER, Willugh.

(1. ASTUR, *Vig.*)

11. *A. palumbarius*, Will. (*Goshawk.*)—A broad white streak over the eye.

Falco palumbarius, *Temm. Man. d'Orn.* tom. i. p. 55. Goshawk, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 29. pls. 12 and 12*. *Bew. Brit. Birds*, vol. i. p. 28.

DIMENS. Entire length (*female*) two feet, (*male*) one-third less. TEMM.

DESCRIPT. (*Adult male.*) Upper parts deep bluish ash-colour : a broad white streak over the eyes : under parts white, with transverse bars and fine longitudinal streaks of dark brown : tail cinereous, with four or five bars of blackish brown ; the tip white : bill bluish gray : cere greenish yellow : irides and feet bright yellow. (*Adult female.*) Upper parts with a brownish tinge : spots on the throat more numerous. (*Young.*) Top of the head and upper part of the neck reddish brown, the feathers edged with white : under parts reddish white, with long lanceolate streaks of deep brown : tail brown, with four broad bars of a darker colour ; the tip white : irides gray : cere and feet livid yellow. (*Egg.*) "Skim-milk white, marked with streaks and spots of reddish brown." SELBY.

Nearly confined to the mountainous parts of Scotland, where it breeds, and is said to be a great destroyer of game. Very rare in England. *Obs.* The Gentil Falcon (*Falco gentilis*, Gmel.) is the young of this species.

(2. ACCIPITER, *Vig.*)

12. *A. fringillarius*, Will. (*Sparrow-Hawk.*)—A white spot on the nape.

Falco Nisus, *Temm. Man. d'Orn.* tom. i. p. 56. Sparrow-Hawk, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 32. pls. 13 and 13*. *Bew. Brit. Birds*, vol. i. p. 44.

DIMENS. Entire length (*male*) twelve inches, (*female*) fifteen inches : length of the bill (from the forehead) eight lines, (from the gape) ten lines and a half ; of the tarsus two inches five lines ; of the middle toe, claw included, one inch ten lines ; of the tail seven inches six lines ; from the carpus to the end of the wing nine inches three lines : breadth, wings extended, (*male*) two feet, (*female*) two feet four inches.

DESCRIPT. (*Adult male.*) Upper parts bluish ash-colour : above the eye an indistinct pale line, conducting to a white spot on the nape : under parts white, with transverse waves of deep brown ; on the throat a few longitudinal streaks : tail grayish ash, with broad brownish black bars : bill bluish gray, the tip black : cere greenish yellow : feet and irides bright gamboge-yellow. (*Female.*) All the upper parts blackish brown,

passing into blackish gray: throat whitish, with fine longitudinal dusky streaks; rest of the under parts reddish white, undulated with transverse bars of dusky brown. (*Young.*) Upper parts brown, the feathers with pale reddish edges: some whitish spots on the shoulders: under parts yellowish white, with transverse reddish bars; or whitish, with brown bars: irides grayish: feet livid yellow. (*Egg.*) Pale bluish white, blotched and spotted with dark red brown: long. diam. one inch seven lines; trans. diam. one inch four lines.

Common throughout the country. Preys upon the smaller quadrupeds and birds, and is very destructive to partridges. Builds in trees, forming a shallow nest of slender twigs. Lays from three to six eggs. This species is remarkable for the great difference of size between the male and female.

GEN. 5. MILVUS, *Bechst.*

(1. MILVUS, *Vig.*)

13. *M. Ictinus*, Sav. (*Kite.*)—Reddish brown above; beneath ferruginous, with dark longitudinal streaks.

Falco Milvus, *Temm. Man. d'Orn.* tom. I. p. 59. *Kite*, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. I. p. 74. pl. 5. *Bew. Brit. Birds*, vol. I. p. 32.

DIMENS. Entire length twenty-five inches: length of the bill (from the forehead) one inch eight lines, (from the gape) one inch eleven lines; of the tarsus two inches two lines; of the tail twelve inches eight lines; from the carpus to the end of the wing twenty inches: breadth, wings extended, five feet two inches.

DESCRIPT. (*Male.*) Head and neck grayish white, with fine streaks of dusky brown; the feathers on these parts long, and acuminate: rest of the upper parts reddish brown; the feathers with pale edges: under parts ferruginous, with longitudinal brown streaks: tail long and deeply forked, reddish orange, with obsolete brown bars: bill yellowish brown at the base, towards the tip dusky: cere and irides yellow. (*Female.*) Upper plumage of a deeper brown, with less of the ferruginous tinge; the edges of the feathers paler, approaching to white. (*Young of the year.*) Feathers on the head shorter, and less acuminate; bright red without streaks, tipped with white: upper parts redder than in the adult; feathers on the back and wings dusky in the centre, reddish yellow at the edges: on the lower part of the neck some large white spots. (*Egg.*) Dirty white; the larger end spotted with red brown: long. diam. two inches two lines; trans. diam. one inch nine lines.

Common in some parts of the country, but not generally diffused. Frequents wooded districts, and builds in trees. Food young game, and the smaller quadrupeds.

(*ELANUS*, Sav.)

- (1.) *M. furcatus*, Nob. *Falco furcatus*, Linn. Syst. Nat. tom. I. p. 129. *Nauclerus furcatus*, Vig. in Zool. Journ. vol. II. p. 387. *Swallow-tailed Falcon*, Shaw, Nat. Misc. vol. VI. pl. 204. *Swallow-tailed Elanus*, Selb. Illust. vol. I. p. 77.

This species, which is a native of North America, is stated by Dr Fleming (*Brit. An.* p. 52) as having occurred to the late Dr Walker in Argyleshire, in 1772. A second individual is said to have been taken alive in Yorkshire, in Sept. 1805. (*Linn. Trans.* vol. XIV. p. 583.) No British-killed specimen, however, is known to exist in any of our museums.

GEN. 6. BUTEO, *Nob.*(1. BUTEO, *Bechst.*)

14. *B. vulgaris*, Will. (*Common Buzzard*).—Tarsi naked: tail barred transversely throughout its whole length.

Falco Buteo, *Temm. Man. d'Orn.* tom. i. p. 63. Common Buzzard, *Mont. Orn. Dict.* *Selb. Illust.* vol. i. p. 55. pl. 6. *Bew. Brit. Birds*, vol. i. p. 22.

DIMENS. Entire length twenty inches six lines: length of the bill (from the forehead) one inch four lines, (from the gape) one inch six lines; of the tarsus three inches one line; of the tail eight inches six lines; from the carpus to the end of the wing fifteen inches three lines: breadth, wings extended, three feet eleven inches.

DESCRIPT. (*Adult.*) All the upper parts, as well as the neck and breast, deep brown: throat whitish, with a few brown streaks on the shafts of the feathers; abdomen grayish white, with longitudinal spots of deep brown: tail with about twelve transverse dusky bars, the extremity slightly rounded: bill lead-colour: cere, irides, and feet, yellow. (*Young.*) Upper plumage brown, much varied with yellowish white and brownish yellow: throat pure white, with longitudinal brown spots; breast and belly yellowish white, with large oval stains and blotches of deep brown, prevailing most upon the breast: irides pale brownish yellow. *Obs.* The plumage of this species varies considerably, even in individuals of the same age: the brown above is more or less uniform, and the markings beneath more or less regular: occasionally the brown extends nearly over the whole body, and the white appears only on the throat, and in transverse bars on the middle of the abdomen. (*Egg.*) Dirty white, spotted with pale brown: long. diam. two inches three lines; trans. diam. one inch ten lines.

Of frequent occurrence throughout the country. Habits sluggish and inactive. Builds in old trees, and lays from two to four eggs. Feeds on rats, mice, moles, leverets, small birds, and occasionally on reptiles.

15. *B. Lagopus*, Vig. (*Rough-legged Buzzard*).—Tarsi feathered nearly to the toes: basal portion of the tail white, the remainder brown.

Falco Lagopus, *Temm. Man. d'Orn.* tom. i. p. 65. Rough-legged Falcon, *Mont. Orn. Dict.* *Rough-legged Buzzard*, *Selb. Illust.* vol. i. p. 58. pl. 7. *Bew. Brit. Birds*, vol. i. p. 20.

DIMENS. Entire length twenty inches six lines: from the carpus to the end of the wing sixteen inches six lines.

DESCRIPT. (*Adult.*) Head, upper part of the neck, throat, breast, and thighs, yellowish white, spotted and streaked with brown: back and wing-coverts deep brown, the feathers edged with reddish yellow: quills dusky: with one half of the inner webs white, spotted with deep brown: lower portion of the abdomen deep umber-brown: rump and under tail-coverts yellowish white: tail white at the base, the remaining part brown: the extreme tip dirty white: bill bluish black, darkest at the tip: cere yellow: irides brown. (*Young.*) Upper parts more or less varied with white: a considerable portion of the abdomen whitish, spotted with

brown; the sides with larger spots of the same colour: tail with two or three transverse bars towards the extremity: irides yellowish brown. (*Egg.*) Pale brownish white, blotched with darker brown: long. diam. two inches two lines; trans. diam. one inch eight lines.

A rare and only occasional visitant in this country. Has been twice killed in Cambridgeshire; been met with also in Norfolk, Suffolk, Kent, the Isle of Wight, Gloucestershire, Northumberland, and other parts of England. Food and habits much resembling those of the last species.

(2. *PERNIS*, *Cuv.*)

16. *B. apivorus*, Ray. (*Honey Buzzard*.)—Upper plumage brown, with cinereous bars on the wings: under parts white, with triangular brown spots.

Falco apivorus, *Temm. Man. d'Orn.* tom. 1. p. 67. *Honey Buzzard*, *Mont. Orn. Dict. Selb. Illust.* vol. 1. p. 62. pl. 8. *Bew. Brit. Birds*, vol. 1. p. 24.

DIMENS. Entire length one foot ten inches: length of the bill (from the forehead) eleven lines, (from the gape) one inch six lines; of the tarsus two inches three lines; of the tail eight inches; from the carpus to the end of the wing one foot four inches: breadth, wings extended, four feet one inch.

DESCRIPT. (*Adult male.*) Top of the head bluish ash; upper parts of the body brown, inclining to cinereous: secondaries barred alternately with dusky brown and bluish gray: under parts whitish, with brown spots, which assume a triangular form on the breast and abdomen: tail with three transverse dusky bars: cere greenish grey: inside of the mouth, irides, and feet, yellow. (*Female and young.*) Forehead alone bluish ash; rest of the upper parts pale reddish brown, with a large dusky spot in the centre of each feather: throat nearly white: fore part of the neck, breast and belly, yellowish red, with large brown spots; sometimes light brown, with reddish brown bars. (*Young of the year.*) Upper parts like those of the adult female, but the feathers on the head, neck, and greater wing-coverts, tipped with white: quills dusky, tipped with white: throat whitish; rest of the under parts reddish white with large brown spots, or reddish brown with a few partially white feathers intermixed, occasionally of a uniform pale brown with fine longitudinal dusky streaks: cere yellow: irides brown. *Obs.* This species appears to vary considerably in plumage, especially in the immature state. (*Egg.*) Mottled nearly all over with two shades of dark red brown, the white ground only here and there visible: long. diam. two inches one line; trans. diam. one inch nine lines.

Of rare occurrence in this country, especially in the adult state. Has been met with in Hampshire, Berkshire, Norfolk, Northumberland, and several times in Cambridgeshire. Feeds principally upon the larvæ and pupæ of wasps during the season in which they are to be obtained; at other times, preys on small quadrupeds and birds. Is said to build in tall trees.

(3. *CIRCUS*, *Bechst.*)

17. *B. rufus*, Nob. (*Marsh Harrier*.)—General plumage deep reddish brown: crown of the head yellowish white: wings a little shorter than the tail.

Falco rufus, *Temm. Man. d'Orn.* tom. i. p. 69. Marsh Harrier, *Selb. Illust.* vol. i. p. 66. pl. 9. Moor-Buzzard, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. i. p. 26.

DIMENS. Entire length one foot seven to nine inches : breadth, wings extended, three feet eleven inches to four feet two inches.

DESCRIPT. (*Adult in the fourth year.*) Head and neck yellowish brown, with longitudinal darker streaks: scapulars and wing-coverts reddish brown: first five primaries dusky lead-colour; all the secondaries and tail-feathers cinereous: rest of the plumage bright ferruginous: bill black: irides reddish yellow: cere and feet yellow. (*Second year.*) Crown of the head, throat, and nape, straw-yellow, with fine longitudinal dusky streaks: rest of the body dark umber-brown, tinged beneath with rust-red: a few yellowish spots upon the wings: irides brown. (*Young of the year.*) Whole plumage deep chocolate-brown, inclining to yellowish on the throat, crown, and back of the head: quills, wing-coverts, and tail-feathers, tipped with yellowish brown: occasionally a few white patches on the lower part of the abdomen: irides dusky brown. (*Egg.*) Rather pointed: white: long. diam. two inches one line; trans. diam. one inch six lines.

Common in marshy districts. Preys on water-fowl, small quadrupeds, and reptiles. Nest on the ground amongst rushes: eggs three to five in number. *Obs.* The *Falco ceruginosus* of Linnæus is the young of this species.

18. *B. cyaneus*, Nob. (*Hen-Harrier.*)—Upper plumage (*male*) bluish gray, (*female*) reddish brown: quills dusky, without transverse bars: third and fourth primaries of equal length: wings reaching to three-fourths the length of the tail.

Falco cyaneus, *Temm. Man. d'Orn.* tom. i. p. 72. Hen-Harrier and Ring-tail, *Mont. Orn. Dict. and Supp.* *Bew. Brit. Birds*, vol. i. pp. 34 and 36. *Selb. Illust.* vol. i. p. 68. pl. 10.

DIMENS. Entire length (*male*) one foot six or seven inches, (*female*) one foot eight or nine inches: length of the tarsus three inches: breadth, wings extended, three feet two to four inches.

DESCRIPT. (*Adult male.*) Head, neck, back, scapulars, and wing-coverts, bluish gray; rump white: quills black, whitish at the base: breast, belly, sides, under wing and tail-coverts, pure white, without spots of any kind: upper part of the tail ash-gray, whitish at the tip: bill bluish black: irides, cere, and feet, yellow. (*Adult female.*) Space surrounding the eyes white: upper parts of the plumage deep brown: feathers on the head, neck, top of the back, and wing-coverts, edged with rust-red: under parts pale reddish yellow, with deep orange-brown longitudinal streaks and spots: quills dusky, barred underneath with white: tail with alternate broad bars of deep brown and pale yellowish rust. *Obs.* The *young* of both sexes resemble the old female: *after the first autumnal moult*, the male begins to assume the adult plumage, and exhibits a mixture of ash-gray and reddish brown: the transverse bars on the tail also gradually disappear. (*Egg.*) White: long. diam. one inch eight lines; trans. diam. one inch four lines.

Pretty generally distributed throughout the country, but seemingly most partial to fenny districts. Preys on small birds and quadrupeds. Makes its nest on the ground; lays from three to five eggs, and hatches

in the beginning or middle of June. *Obs.* The *Ring-tail* of English authors is the female of this species.

19. *B. cineraceus*, Flem. (*Ash-coloured Harrier*).—Upper plumage (*male*) bluish ash, (*female*) reddish brown: third primary much longer than the others: secondaries with three black transverse bars: wings reaching to the extremity of the tail.

Faleo cineraceus, *Mont. in Linn. Trans.* vol. ix. p. 188. *Temm. Man. d'Orn.* tom. i. p. 76. *Ash-coloured Harrier*, *Selb. Illust.* vol. i. p. 70. pl. 11. *Ash-coloured Faleon*, *Mont. Orn. Dict. and Supp. with fig.* *Bew. Brit. Birds*, vol. i. p. 37.

DIMENS. Entire length seventeen inches: length of the bill (from the forehead) ten lines, (from the gape) eleven lines and a half; of the tarsus two inches three lines; of the tail eight inches two lines; from the carpus to the end of the wing thirteen inches nine lines. *Female*.

DESCRIPT. (*Adult male*.) Head, throat, breast, and all the upper parts of the body, deep bluish ash; belly, sides, and thighs, white, with longitudinal rust-coloured streaks: primaries black; secondaries ash-gray above, paler beneath, with three transverse dusky bars, one of which is visible externally: under wing-coverts barred with reddish brown: two middle tail-feathers brownish gray; the rest cinereous, their inner webs barred with reddish brown: bill bluish black: irides and feet yellow. (*Adult female*.) Crown of the head reddish brown, with dusky spots: nape yellowish red, sometimes approaching to white: above and below the eye a pale fascia: upper parts of the body deep brown, the feathers with reddish edges: lower part of the rump, and tail-coverts, white, streaked with pale orange-brown: all the under parts bright ferruginous, with the shafts of the feathers somewhat darker, appearing like fine slender streaks: tail with the two middle feathers of a uniform brown; the rest with brown and ferruginous bars. (*Young of the year*.) Upper plumage much resembling that of the adult female: under parts of an uniform rust-red colour, without any spots or streaks: irides brown. (*Egg*.) White: long. diam. one inch seven lines; trans. diam. one inch four lines.

First discovered by Montagu in Devonshire: has since been occasionally met with in the North of England, and likewise in the fens of Cambridgeshire. Nest placed on the ground; often amongst furze: the eggs, which are generally four in number, are hatched about the second week in June.

GEN. 7. BUBO, *Isid. Geoff.*

(1. BUBO, *Cuv.*)

20. *B. maximus*, Flem. (*Eagle Owl*).—Upper plumage variegated with black and ochre: under parts ochre-yellow, with longitudinal black spots.

Strix Bubo, *Temm. Man. d'Orn.* tom. i. p. 100. *Eagle Owl*, *Selb. Illust.* vol. i. p. 82. pl. 19. *Bew. Brit. Birds*, vol. i. p. 52. *Great-eared Owl*, *Mont. Orn. Dict.*

DIMENS. Entire length two feet. TEMM.

DESCRIPT. All the upper parts elegantly varied and spotted with black, ochre-yellow, and yellowish gray: under parts ochre-yellow, with longitudinal black spots and streaks: throat white: egrets composed of six or eight elongated feathers, coloured like the rest of the plumage: tarsal feathers reddish yellow: bill and claws horn-colour: irides bright orange. (*Female*.) Rather larger than the male, and without the white throat. (*Egg*.) White: long. diam. two inches five lines; trans. diam. one inch ten lines.

An occasional visitant of extreme rarity in this country. According to Montagu has been shot in Yorkshire and in Sussex, as well as in Scotland. More recently four specimens are said to have been observed on the northern coast of Donegal, Ireland. Inhabits mountainous and rocky situations, where it breeds. Food small quadrupeds, and reptiles.

(2. SCOPS, Sav.)

21. *B. Scops*, Nob. (*Scops-eared Owl*.) — Plumage variegated with brown, gray, black, and rufous.

Strix Scops, *Temm. Man. d'Orn.* tom. i. p. 103. *Scops-eared Owl*, *Selb. Illust.* vol. i. p. 92. pl. 22. *Bew. Brit. Birds*, vol. i. p. 67. Little-horned Owl, *Mont. Orn. Dict. Supp. App.*

DIMENS. Entire length seven inches and a half. FLEM.

DESCRIPT. Head, neck, and egrets, (the latter consisting of six or eight feathers,) brownish gray, speckled with black; rest of the upper parts reddish ash, with spots and zigzag streaks of black and brown: under parts ash-gray, streaked and speckled with black and reddish brown: quills barred with white: tail variegated with black, brown, and white: bill black: irides yellow: toes bluish gray. (*Egg*.) Of a short oval form: white, with a few faint dusky specks: long. diam. one inch three lines; trans. diam. one inch and half a line.

Very rare. Has been killed near York, and in one or two other parts of the country. Is probably migratory. Preys on mice and insects. *Obs.* The *Strix pulchella* of Donovan (*Brit. Birds*, vol. vii. pl. 165.) appears to be the same as this species.

GEN. 8. OTUS, Cuv.

22. *O. vulgaris*, Flem. (*Long-eared Owl*.) — Plumage reddish yellow, variegated with gray and dusky brown: egrets elongated, composed of eight or ten feathers.

Strix Otus, *Temm. Man. d'Orn.* tom. i. p. 102. *Long-eared Owl*, *Mont. Orn. Dict.* *Selb. Illust.* vol. i. p. 85. pl. 20. *Bew. Brit. Birds*, vol. i. p. 56.

DIMENS. Entire length fourteen inches: length of the tail six inches; of the egrets one inch three lines; from the carpus to the end of the wing eleven inches six lines.

DESCRIPT. Upper parts of a yellowish rust-colour, irregularly marked and spotted with deep brown, gray and cinereous: egrets of eight or ten black feathers, edged with yellow and white: under parts orange-yellow, with oblong and arrow-shaped streaks of dusky brown: quills and tail barred with brownish black: tarsal feathers yellow, without spots: bill

dusky: irides rich orange-yellow. (*Female*.) Throat and face white: rest of the plumage like that of the male, but much mottled in places with grayish white. (*Egg*.) Oval; smooth: white: long. diam. one inch eight lines and a half; trans. diam. one inch three lines and a half.

Found in wooded districts not very uncommonly. Partial to fir-plantations and old ivy, where it breeds early in the Spring, laying from four to five eggs: these are hatched towards the end of April. Remains with us the whole year.

23. *O. Brachyotos*, Flem. (*Short-eared Owl*.)—Upper plumage dusky brown, edged with ochre-yellow: egrets small and inconspicuous, composed of three or four feathers.

Strix Brachyotos, *Temm. Man. d'Orn.* tom. i. p. 99. Short-eared Owl, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. i. p. 88. pl. 21. *Bew. Brit. Birds*, vol. i. pp. 58 and 60.

DIMENS. Entire length fifteen inches: length of the bill (along the ridge) one inch three lines; of the tarsus one inch; of the tail six inches; of the egrets six lines; from the carpus to the end of the wing eleven inches eight lines: breadth, wings extended, three feet one inch.

DESCRIPT. Head small: facial circle dirty white, with dark streaks radiating outwards; immediate contour of the eyes black: egrets black, with tawny edges: head, neck, back, and wing-coverts, variegated with dusky brown and tawny yellow, the feathers being mostly edged with this last colour, with a dark spot in the centre of each: breast and belly pale orange-yellow, with brownish black streaks on the shafts of the feathers; these streaks most abundant on the breast: vent and under tail-coverts yellowish white: quills with alternate bars of dark brown and ochre-yellow: tail barred like the quills; the two outermost feathers much paler than the others; the four central ones with dusky spots in the middle of the yellow bars: feet and toes thickly coated with downy ochreous feathers: bill dusky: irides rich golden yellow: claws black. (*Egg*.) Smooth: white, with a slight blush of red: long. diam. one inch eight lines; trans. diam. one inch three lines and a half.

A migratory species, visiting England in October, and departing in April. Has been known, however, to breed in Norfolk, as well as in some parts of Scotland. Habits somewhat diurnal. Met with chiefly on moors and in open fields, often in flocks consisting of several individuals. Resorts to plantations in the evening to roost. Food principally the field-campagnol. Nest placed on the ground.

GEN. 9. *STRIX*, Linn.

24. *S. flammea*, Linn. (*White Owl*.)—Upper plumage tawny yellow, variegated with gray and brown: underneath white.

S. flammea, *Temm. Man. d'Orn.* tom. i. p. 91. White Owl, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 99. pl. 24. *Bew. Brit. Birds*, vol. i. p. 61.

DIMENS. Entire length fourteen inches six lines: length of the bill (from the forehead) one inch four lines, (from the gape) one inch seven

lines; of the tarsus two inches six lines; of the tail four inches eight lines; from the carpus to the end of the wing eleven inches: breadth, wings extended, two feet eleven inches six lines.

DESCRIPT. Facial circle white: all the upper parts bright tawny yellow, variegated with fine zigzag lines of brown and gray, and sprinkled with numerous white dots: under parts pure white, or with a faint tinge of reddish yellow, occasionally speckled with a few brownish points: bill and claws yellowish white: irides bluish black. (*Egg.*) Dull white: long. diam. one inch six lines; trans. diam. one inch three lines.

Common in every part of the kingdom. Frequents churches and the eaves of old buildings, where it breeds, laying from three to five eggs. Habits nocturnal: comes abroad about sunset. Feeds principally on mice, but will occasionally devour other of the smaller quadrupeds: rejects the shrew. Screams in its flight, but does not hoot.

GEN. 10. SYRNIUM, *Sav.*

25. *S. Aluco*, Nob. (*Tawny Owl.*) — Upper plumage reddish brown, variegated with black and ash-gray: scapulars and wing-coverts spotted with white: under parts white, with reddish bars, and longitudinal dusky streaks.

Strix Aluco, *Temm. Man. d'Orn.* tom. 1. p. 89. *Tawny Owl*, *Mont. Orn. Dict. Selb. Illust.* vol. 1. p. 102. pl. 25. *Bew. Brit. Birds*, vol. 1. p. 63.

DIMENS. Entire length sixteen inches: length of the bill (from the forehead) one inch four lines, (from the gape) one inch six lines; of the tarsus two inches; of the tail six inches eight lines; from the carpus to the end of the wing ten inches ten lines: breadth, wings extended, three feet.

DESCRIPT. Head large: facial circle white, with a tinge of reddish brown: upper parts of the plumage ferruginous brown, variously marked and spotted with dark brown, black, and ash-gray: several large white spots upon the scapulars and wing-coverts, disposed in rows: under parts white, with transverse bars of reddish brown, and longitudinal dusky streaks on the shafts of the feathers: quills barred alternately with dusky and tawny brown: two middle tail-feathers of a uniform tawny brown; the others barred like the quills: bill yellowish white: eyes very large: irides bluish black. (*Female.*) General colour more ferruginous than in the male, with less white: the transverse bars on the wings and tail alternately red and brown. (*Egg.*) Smooth: dull white: long. diam. one inch ten lines; trans. diam. one inch six lines.

Equally common with the last species, but found only in woods. Builds in the hollows of old trees, or amongst ivy, and lays four eggs, which are hatched in the beginning of April. Preys upon various small quadrupeds and birds. Comes abroad only during the night, and has a clamorous hooting note.

GEN. 11. NOCTUA, *Sav.*

(1. *SURNIA*, *Selby.*)

26. *N. nyctea*, Nob. (*Snowy Owl.*) — Plumage snow-white, more or less marked with transverse brown bars.

Strix nyctea, *Temm. Man. d'Orn.* tom. i. p. 82. Snowy Owl, *Mont. Orn. Dict. Supp. App. Selb. Illust.* vol. i. p. 95. pl. 23. *Bew. Brit. Birds*, vol. i. p. 54.

DIMENS. Entire length (*female*) two feet one inch: breadth, wings extended, five feet. *Male* of inferior size. **SELBY.**

DESCRIPT. Head small: plumage above and below snow-white, more or less variegated with brown spots and transverse bars, which are largest and most conspicuous in *immature* birds; in very old individuals (*males* more especially) these spots wholly disappear, leaving the entire plumage pure white: bill black, almost concealed by the bristly feathers at its base: irides bright orange-yellow: tail rounded, rather short, extending nearly two inches beyond the folded wings: legs and toes very thickly clothed with hair-like feathers partly concealing the claws. (*Egg.*) Pure white.

Found in the Orkney and Shetland Isles, where it remains the whole year. Very rare in England. Has been killed in Northumberland, and even as far south as in Norfolk. In the latter county it has occurred twice. Habits said to be diurnal. Food hares, rats, mice, grouse, and other birds. Builds in rocky situations, and lays two eggs.

(2. NOCTUA, Selby.)

27. *N. passerina*, Selby. (*Little Owl.*)—Tarsi feathered; toes thinly covered with a few white hairs.

Strix passerina, *Temm. Man. d'Orn.* tom. i. p. 92. Little Owl, *Mont. Orn. Dict.* Little Night-Owl, *Selb. Illust.* vol. i. p. 107. pl. 27.

DIMENS. Entire length nine inches. **TEMM.**

DESCRIPT. Facial circle mostly white: upper plumage grayish brown; head and nape rather thickly spotted with white; wing-coverts and scapulars with large irregular white spots of an oval-oblong form; middle of the back nearly free from spots; sides of the nape much variegated with white; quills with a regular series of yellowish white spots on their outer webs; tail with alternate bars of brown and yellowish white: under parts white, with oblong longitudinal spots of grayish brown; chin and under tail-coverts free from spots: tarsi clothed with short downy white feathers; longer (according to Selby) than in the next species; toes thinly covered with a few white hairs. (*Egg.*) Of a short oval form: white: long. diam. one inch four lines; trans. diam. one inch one line.

A very rare and occasional visitant in this country. Has been killed in a few instances, but from having been formerly confounded with the next species, such instances can hardly be particularized with certainty. Said to frequent towers and old buildings, in which situations it breeds, laying from two to four eggs. Food, according to Temminck, mice, bats, small birds, and insects.

28. *N. Tengmalmi*, Selby. (*Tengmalm's Owl.*)—Tarsi and toes thickly clothed with downy feathers.

Strix Tengmalmi, *Temm. Man. d'Orn.* tom. i. p. 94. Tengmalm's Night-Owl, *Selb. Illust.* vol. i. p. 105. pl. 26. Little Owl, *Bew. Brit. Birds*, vol. i. p. 65.

DIMENS. Entire length eight inches four lines. **TEMM.**

DESCRIPT. Facial circle more developed than in the last species; blackish round the eyes, and at the base of the bill; the rest of it white,

bordered externally with dark brown and small white spots intermixed: upper parts nearly similar: liver-brown, spotted with white, but the spots on the head and nape not so numerous; the white spots on the scapulars and wing-coverts disposed more in lines: wings and tail relatively longer; the latter with interrupted white bars: under parts white; the dark spots much as in *N. passerina*, but having a tendency to unite to form transverse bars: bill yellowish white: legs and toes thickly clothed with soft downy feathers. (*Egg.*) Elongated; pointed: white: long. diam. one inch four lines; trans. diam. eleven lines.

Of equal rarity with the last in this country. Has been killed near Morpeth in Northumberland, and probably in other instances. Said to feed on mice and insects, and to breed in the hollows of firs.

ORDER II. INSESSORES.

GEN. 12. LANIUS, *Linn.*

* *Tail long; graduated.*

29. *L. Excubitor*, *Linn.* (*Cinereous Shrike.*)—Head, neck, and back, cinereous; a black band beneath the eyes: under parts white.

L. Excubitor, *Temm. Man. d'Orn.* tom. i. p. 142. *Cinereous Shrike*, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. i. p. 148. pl. 43. f. 1. *Bew. Brit. Birds*, vol. i. p. 71.

DIMENS. Entire length ten inches: length of the bill (from the forehead) eight lines, (from the gape) one inch; of the tarsus one inch; of the tail four inches three lines; from the carpus to the end of the wing four inches six lines: breadth, wings extended, fourteen inches.

DESCRIPT. (*Male.*) Upper plumage, including the head neck and back, ash gray; scapulars, rump, and upper tail-coverts, somewhat paler: a black band beneath the eye, commencing at the base of the upper mandible, and extending beyond the orifice of the ear: under parts pure white: wings black; primaries white at the base; secondaries tipped with white: tail cuneiform, of twelve feathers; the two middle ones black; the two next tipped with white; the others with the white portion gradually increasing to the outermost feather, which is almost wholly white: bill and feet black: irides blackish brown. (*Female.*) Upper plumage resembling that of the male, but the colours more obscure: under parts dirty white, with numerous crescent-shaped dusky lines. (*Egg.*) Light bluish white; the larger end nearly covered with spots of two shades of light brown and ash: long. diam. one inch one line; trans. diam. nine lines and a half.

A migratory, and not abundant species: visits this country sparingly towards the end of Autumn, departing in the Spring. Feeds on small birds, mice, insects, &c. Transfixes its prey when killed upon a thorn before devouring it. Builds in trees, and lays from five to seven eggs.

*** Tail short; even, or slightly rounded.*

30. *L. rufus*, Briss. (*Wood-Chat*).—Upper plumage variegated with black, white, and ferruginous: under parts white: second quill equal to the fifth: tail slightly rounded.

L. rufus, *Temm. Man. d'Orn.* tom. i. p. 146. *Wood-Chat*, *Bew. Brit. Birds*, vol. i. p. 377. *Selb. Illust.* vol. i. p. 153. pl. 100. f. 1. *Wood-Shrike*, *Mont. Orn. Dict. Supp.*

DIMENS. Entire length seven inches five lines.

DESCRIPT. Forehead, region of the eyes and ears, and upper part of the back, black: occiput and nape ferruginous: scapulars white: wings black: quills white at the base; some of them also edged and tipped with white: rump gray: under parts pure white: tail with the two outermost feathers wholly white, excepting a black spot at the base of the inner web: the two next with the black spot of larger size, and on both webs; the rest black, tipped with white; the two middle ones wholly black. (*Female*.) Colours more obscure: occiput and nape streaked with brown: the black on the back approaching to that colour: under parts dirty white; the breast with transverse brown lines. (*Egg*.) Pale bluish white: the larger end spotted with hair-brown and ash-gray: long. diam. eleven lines; trans. diam. eight lines.

Extremely rare in this country. Has been killed in the neighbourhood of Canterbury, and near Swaffham in Norfolk. In the latter locality, it has been observed to breed more than once. Feeds on insects. Places its nest in the forked branches of bushes.

31. *L. Collurio*, Linn. (*Red-backed Shrike*).—Back, scapulars, and wing-coverts, ferruginous: breast and flanks rose-red: second quill longer than the fifth: tail even at the extremity.

L. Collurio, *Temm. Man. d'Orn.* tom. i. p. 147. *Red-backed Shrike*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 150. pl. 43. f. 2. and pl. 43. f. 1, 2. *Bew. Brit. Birds*, vol. i. p. 73.

DIMENS. Entire length seven inches: length of the bill (from the forehead) six lines, (from the gape) nine lines and a half; of the tarsus eleven lines; of the tail three inches one line and a half; from the carpus to the end of the wing three inches seven lines: breadth, wings extended, eleven inches four lines.

DESCRIPT. (*Male*.) Crown of the head and nape ash-gray: a black band through the eyes, commencing at the base of the upper mandible, and reaching half-way down the neck: back, scapulars, and wing-coverts, ferruginous brown: rump ash-gray: quills dusky, edged with reddish brown: throat and vent white: breast, belly, and flanks, rose-red: tail nearly even at the extremity: the two middle feathers wholly black; the rest white from the base through two-thirds of their length, the remaining portion black, except the extreme tip which is also white: bill and feet black: irides chestnut-brown. (*Female*.) All the upper parts dull ferruginous brown: nape and rump tinged with ash-gray: under parts grayish white; sides of the neck, breast, and flanks, barred transversely with narrow semicircular dusky lines: tail with the outer webs of the two external feathers edged with white: the four middle ones of a uni-

form reddish brown. (*Young*.) General plumage resembling that of the adult female, but with less of the grayish tinge on the nape and rump; upper parts here and there barred with blackish brown. (*Egg*.) Pointed: ground colour reddish white, spotted with darker red and ash-gray, the spots forming a band near the large end: long. diam. eleven lines; trans. diam. eight lines. *Obs.* The eggs of this species vary occasionally, and are sometimes greenish white, with spots of light brown and ash-gray intermixed.

A regular summer visitant, arriving in the Spring, and departing in the early part of the Autumn. Not uncommon in some parts of the country. Haunts copses, tall hedges, and furzy commons, where it builds. Nest formed of moss and wool, intermixed with grass, and lined with hair: eggs five or six. Food principally coleopterous insects. *Obs.* The *Wood-Chat* figured at p. 75 (vol. i.) of Bewick's *British Birds*, is only the female of this species.

GEN. 13. MUSCICAPA, Linn.

32. *M. grisola*, Linn. (*Spotted Flycatcher*.)—Upper plumage brownish ash: under parts white: head and breast spotted with dusky.

M. grisola, *Temm. Man. d'Orn.* tom. i. p. 152. Spotted Flycatcher, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 141. pl. 43*. f. 1. *Bew. Brit. Birds*, vol. i. p. 210.

DIMENS. Entire length six inches one line: length of the bill (from the forehead) five lines and a half, (from the gape) nine lines; of the tarsus seven lines; of the tail two inches eight lines and a half: from the carpus to the end of the wing three inches four lines: breadth, wings extended, ten inches three lines.

DESCRIPT. All the upper parts of the plumage brownish ash; crown of the head obscurely spotted with dusky: throat and middle of the belly white; sides of the neck and breast with longitudinal streaks of brown; flanks tinged with pale orange-red: legs dusky brown. (*Egg*.) White tinged with blue, and mottled all over with pale reddish yellow brown: long. diam. nine lines; trans. diam. seven lines.

A migratory, and common species. Arrives in this country about the middle, or towards the latter end of May: departs in the Autumn. Frequents gardens. Builds in holes of trees and old walls; often in the branches of trees nailed against walls. Nest formed of bents, moss, &c. interwoven with spider's webs and a little wool; lined with dry grass and feathers. Eggs five in number; hatched about the second week in June. Food insects, taken on the wing. Has no song.

33. *M. luctuosa*, Temm. (*Pied Flycatcher*.)—Upper plumage black; forehead, and under parts of the body, white.

M. luctuosa, *Temm. Man. d'Orn.* tom. i. p. 155. Pied Flycatcher, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. i. p. 143. pl. 43*. f. 2, 3. *Bew. Brit. Birds*, vol. i. p. 206.

DIMENS. Entire length five inches two lines.

DESCRIPT. (*Adult male in summer plumage*.) All the upper parts, including the tail, deep black; forehead and under parts pure white:

wings black; the middle and greater coverts white. (*Female*.) Upper parts grayish brown: the white on the forehead and wings less conspicuous: the three outermost tail-feathers edged with white. (*Young male*.) Upper parts grayish brown: wings black; coverts and scapulars broadly edged with white: tail black; the three outermost feathers edged with white on the exterior web, the white portion gradually decreasing: under parts yellowish white, tinged with brown. (*Egg*.) Uniform pale blue: long, diam. eight lines and a half; trans. diam. six lines and a half.

Found in Yorkshire, and in some other parts of the North during the summer months; but is very local. Is probably a bird of passage. Habits similar to those of the last species.

- (2.) *M. albicollis*, Temm. Man. d'Orn. tom. i. p. 153. *White-collared Flycatcher*, Gould, Europ. Birds, part 7.

Mr Gould, in his valuable work above referred to, states that he "has seen this species in a collection of British birds, and was informed that it was supposed to have been killed in England." Its claims, however, to a place in our Fauna require to be confirmed by some further authority. The general character of its plumage is very similar to that of the *M. luctuosa*.

GEN. 14. CINCLUS, *Bechst.*

34. *C. aquaticus*, *Bechst.* (*Water-Ouzel*.)—Upper parts brown: throat and breast white: abdomen brownish red.

- C. aquaticus*, *Temm. Man. d'Orn.* tom. i. p. 177. *Water-Ouzel*, *Mont. Orn. Dict. and Supp.* *Bew. Brit. Birds*, vol. ii. p. 118. *European Dipper*, *Selb. Illust.* vol. i. p. 172. pl. 45*.

DIMENS. Entire length seven inches six lines.

DESCRIPT. (*Male*.) Upper parts deep brown, passing into black; the feathers on the back and wings edged with ash-gray: throat, eyelids, sides and fore part of the neck, and upper part of the breast, white: lower part of the breast, and belly, brownish red: vent and under tail-coverts dusky brown: bill blackish: irides yellowish brown: legs yellowish. (*Female*.) Upper part of the head and neck cinereous brown: the red and white on the under parts less pure. (*Young of the year*.) Head and upper part of the neck gray: some whitish spots on the wings: the white on the under parts extending almost to the vent, and marked with transverse streaks of grayish brown. (*Egg*.) Pointed: white: long, diam. one inch; trans. diam. nine lines.

Inhabits rocky streams, and the banks of rapid rivers, chiefly in the mountainous parts of the country, where it resides all the year. Common in the North of England, and in Scotland: found also in Wales, Devonshire, and occasionally in other parts. Feeds on aquatic insects; and is capable of diving. Nest on the ground, composed of moss and leaves, and arched over. Eggs four to six; hatched in May.

GEN. 15. TURDUS, *Linn.*

* *Plumage brown, and spotted.*

35. *T. viscivorus*, *Linn.* (*Missel-Thrush*.)—Lore grayish white: upper plumage cinereous brown; wing-coverts edged with white: all the under parts spotted.

T. viscivorus, *Temm. Man. d'Orn.* tom. i. p. 161. Missel-Thrush, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 158. pl. 44. f. 1. *Bew. Brit. Birds*, vol. i. p. 112.

DIMENS. Entire length eleven inches: length of the bill (from the forehead) nine lines, (from the gape) one inch two lines; of the tarsus one inch three lines; of the tail four inches; from the carpus to the end of the wing six inches: breadth, wings extended, eighteen inches.

DESCRIPT. Upper parts brown, with a tinge of ash-colour: space between the bill and the eye grayish white: all the under parts white, lightly shaded here and there with reddish yellow, and marked with deep brown spots, which are triangular on the throat and fore part of the neck, oval on the abdomen: upper wing-coverts edged and tipped with whitish: the three outermost tail feathers tipped with white: bill dusky: feet pale brown. (*Young of the year.*) Upper parts cinereous brown: head, back, and scapulars, distinctly spotted with yellow and dusky, the former colour occupying the central portion, the latter the extreme tip of each feather: wing-coverts broadly tipped and edged with yellow; quills brown, with a narrow edging of yellow: tail brown, all the feathers, except the two middle ones, tipped with white: under parts spotted much as in the adult bird: legs very pale yellowish brown. *Obs.* This species is subject to considerable variation of plumage. (*Egg.*) Greenish white, spotted with red brown; sometimes reddish white, spotted with dark red brown: long. diam. one inch three lines; trans. diam. eleven lines.

Not uncommon in some parts of the country; in others less frequent. Haunts woods and orchards. Commences its song very early in the year; continues it till the end of May. Nest placed in the forked branches of trees; formed of moss, wool, and coarse grass, and lined with the finer grasses. Eggs from three to five in number, incubation commencing early in April. Food insects, and berries, particularly those of the misseltoe.

36. *T. pilaris*, Linn. (*Fieldfare.*) — Lore black: back and wing-coverts brown; head, neck, and rump, ash-gray: abdomen without spots.

T. pilaris, *Temm. Man. d'Orn.* tom. i. p. 163. Fieldfare, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 160. pl. 45. f. 1. *Bew. Brit. Birds*, vol. i. p. 116.

DIMENS. Entire length nine inches nine lines: breadth, wings extended, seventeen inches two lines.

DESCRIPT. Head, nape, and lower part of the back (including the rump and upper tail-coverts), deep ash-colour: upper part of the back and wing-coverts chestnut-brown: space between the eye and bill black; above the eyes a white streak: throat, sides of the neck, and breast, yellowish, with oblong dusky spots; feathers on the flanks spotted with black and edged with white; abdomen without spots: quills dusky with pale edges: tail black, the two outermost feathers tipped with whitish on their inner webs: bill yellow; the tips of the mandibles black: irides brown: feet dusky. (*Female.*) Head ash-colour, obscurely spotted with brown: throat whitish: feet paler than in the male bird. (*Egg.*) Light blue, mottled all over with spots of dark red brown.

A winter visitant; first appearing in November: remains with us till very late in the Spring, but does not breed in this country. During its stay keeps in flocks. Feeds on haws and other berries; also on insects and worms.

37. *T. musicus*, Linn. (*Song-Thrush*.)—Lore yellowish: all the upper parts brown tinged with olive: wing-coverts edged with reddish yellow: throat without spots.

T. musicus, *Temm. Man. d'Orn.* tom. i. p. 164. *Song-Thrush*, *Selb. Illust.* vol. i. p. 162. pl. 45. f. 2. *Bew. Brit. Birds*, vol. i. p. 114. *Throstle*, *Mont. Orn. Dict.*

DIMENS. Entire length eight inches eleven lines: length of the bill (from the forehead) nine lines, (from the gape) one inch one line; of the tarsus one inch five lines; of the tail three inches four lines; from the carpus to the end of the wing four inches four lines: breadth, wings extended, thirteen inches eight lines.

DESCRIPT. Head, and all the upper parts, brown with a slight tinge of olive: greater wing-coverts edged with orange-yellow: space between the eye and the bill yellowish: throat pure white, without spots; sides of the neck, and upper part of the breast, reddish yellow, with triangular dusky spots; belly and flanks white, with oval spots: under wing-coverts pale orange-yellow: bill dusky; the under mandible yellowish at the base: feet light brown. (*Egg*.) Light blue, with a few black spots at the larger end: long. diam. one inch one line; trans. diam. ten lines.

Common throughout the country; remaining with us the whole year. Feeds on berries, insects, and snails. In song from the beginning of February till the end of July. Has frequently two broods in a season, the first of which is hatched about the beginning of April, sometimes earlier. Nest placed in shrubs or low trees; composed of moss and grass externally, plastered over within with clay and cow-dung. Eggs three to five in number. This species does not collect in flocks during the winter.

38. *T. iliacus*, Linn. (*Redwing*.)—A broad white streak above the eyes: upper parts olive-brown: under wing-coverts and flanks deep orange-red: middle of the abdomen without spots.

T. iliacus, *Temm. Man. d'Orn.* tom. i. p. 165. *Redwing*, *Mont. Orn. Dict.* *Selb. Illust.* vol. i. p. 165. pl. 45. f. 3. *Bew. Brit. Birds*, vol. i. p. 118.

DIMENS. Entire length eight inches four lines: length of the bill (from the forehead) eight lines and a half, (from the gape) one inch and half a line; of the tarsus one inch and half a line; of the tail three inches three lines; from the carpus to the end of the wing four inches seven lines: breadth, wings extended, thirteen inches.

DESCRIPT. All the upper parts olive-brown: a broad whitish streak above the eye, extending from the base of the bill backwards for an inch or more: space between the eye and the bill black mixed with yellow: sides of the neck, breast, and sides of the abdomen, white, with numerous large oblong dusky spots; middle of the abdomen pure white, without spots: under wing-coverts and flanks deep orange-red: feet pale brown: bill dusky; base of the under mandible yellowish. (*Egg*.) Resembling that of the *Fieldfare*, but smaller, with the spots less numerous.

A migratory species, arriving in large flocks about the beginning of October, or rather later, but generally before the *Fieldfare*. Haunts pastures and hedges. Feeds principally on insects, and the smaller

helices; or, when these cannot be obtained, on berries. Suffers more from very severe weather than the Fieldfare. Retires northward in the Spring to breed.

- (3.) *T. varius*, Horsf. Zoolog. Research. in Java. Linn. Trans. vol. XIII. p. 149.

A bird, supposed to be of this species, was shot by Lord Malmesbury in a small gorze-covert, in the parish of Heron Court in Hampshire, Jan. 24, 1828. It being a native of Java, it is not easy to conjecture by what accident it could have reached this country. The plumage, however, was in very fine condition, and did not exhibit any marks of confinement.

** *Plumage uniform; the ground colour black.*

39. *T. Merula*, Linn. (*Blackbird*.)—Plumage wholly black.

T. Merula, Temm. *Man. d'Orn.* tom. I. p. 168. Blackbird, *Mont. Orn. Dict. Selb. Illust.* vol. I. p. 167. pl. 45. f. 4. and pl. 43. f. 4. *Bew. Brit. Birds*, vol. I. p. 120.

DIMENS. Entire length ten inches eleven lines; length of the bill (from the forehead) eleven lines, (from the gape) one inch three lines; of the tarsus one inch six lines; of the tail four inches eleven lines; from the carpus to the end of the wing five inches: breadth, wings extended, sixteen inches.

DESCRIPT. (*Male*.) The whole of the plumage deep black: orbits, bill, and inside of the mouth, bright yellow: irides and feet dusky brown. (*Female*.) Upper parts dusky brown: throat pale brown, with spots of a darker tint; breast reddish brown: abdomen and under tail-coverts dark cinereous brown: bill and legs dusky brown. (*Young newly fledged*.) Upper parts dusky brown, as in the female, but with reddish streaks on the head, nape, upper part of the back, and lesser wing-coverts, occupying the shafts of the feathers: throat and breast reddish brown, obscurely spotted with dusky. (*Egg*.) Light blue, speckled and spotted with light red brown; occasionally uniform blue without spots: long. diam. one inch two lines; trans. diam. ten lines.

Equally common with the Thrush, and like that bird a constant resident in this country. Food the same. Commences its song in the Spring, a little later than that species, and ceases a little earlier. Habits solitary. Nest formed of moss, twigs, and roots, plastered internally, and afterwards lined with the finer grasses. Eggs four or five in number; hatched the beginning of April.

40. *T. torquatus*, Linn. (*Ring-Ouzel*.) — Plumage black, edged with gray: a crescent-shaped white spot on the breast.

T. torquatus, Temm. *Man. d'Orn.* tom. I. p. 166. Ring-Ouzel, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. I. p. 169. pl. 44. f. 2. *Bew. Brit. Birds*, vol. I. p. 122.

DIMENS. Entire length eleven inches.

DESCRIPT. (*Male*.) The whole plumage black: all the feathers edged with ash-gray: on the upper part of the breast a large crescent-shaped gorget of pure white: bill and legs dusky. (*Female*.) Plumage more

clouded with gray: the gorget on the breast smaller, not so well defined, and tinged with reddish brown and gray. In the *young* female the gorget is scarcely visible: in the *young male* it is reddish white. (*Egg*.) Light blue, mottled with spots of ash-gray and light brown: long. diam. one inch two lines; trans. diam. ten lines.

Not a common species. Occasionally observed in small flocks about Spring and Autumn, on its passage to and from the northern and mountainous parts of the kingdom where it breeds. Does not winter in England. Nest placed on the ground; in form and texture resembling that of the Blackbird. Food insects and berries.

GEN. 16. ORIOIUS, Linn.

41. *O. Galbula*, Linn. (*Golden Oriole*.) — Golden yellow; lore, wings, and tail, black.

O. Galbula, *Temm. Man. d'Orn.* tom. 1. p. 129. Golden Oriole, *Mont. Orn. Dict. Selb. Illust.* vol. 1. p. 176. pl. 35. f. 1 and 2. *Bew. Brit. Birds*, vol. 1. p. 103.

DIMENS. Entire length nine inches six lines: length of the bill (from the forehead) eleven lines and a half, (from the gape) one inch one line and a half; of the tarsus ten lines; of the tail three inches one line; from the carpus to the end of the wing six inches.

DESCRIPT. (*Male*.) Bright golden yellow: space between the eye and the bill black: wings black; the outer webs of the quills edged with white; primary coverts tipped with yellow: two middle tail-feathers wholly black; the others black, tipped with yellow: bill reddish: irides red: feet black. (*Female and young*.) Upper parts olive-green: throat, breast, and abdomen, grayish white, tinged with yellowish, with dusky streaks on the shafts of the feathers: wings brownish black, the quills edged with pale olive-gray: tail olive with a tinge of dusky, the tips of all the feathers, except the two middle ones, yellowish white. (*Egg*.) White, tinged with pink, and sparingly spotted with ash-colour and liver-brown: long. diam. one inch two lines; trans. diam. ten lines.

An occasional visitant in this country, but not often met with. Has been killed in Suffolk, Norfolk, Essex, Hampshire, Devonshire, Cheshire, Lancashire, Northumberland, as well as in Scotland and Ireland. Many of the individuals occurred in the Spring. Frequents wooded districts: feeding on berries and coleopterous insects. Nest purse-shaped, suspended from the forked branches of tall trees. Eggs four or five in number.

GEN. 17. ACCENTOR, Bechst.

42. *A. alpinus*, Bechst. (*Alpine Accentor*.) — Cinereous gray, with large brown spots on the back: throat white with brown scales: * abdomen grayish white mixed with reddish.

A. alpinus, *Temm. Man. d'Orn.* tom. 1. p. 248. Alpine Accentor, *Selb. Illust.* vol. 1. p. 247. pl. D. f. 3.

DIMENS. Entire length six inches eight lines. **TEMM.**

DESCRIPT. Head, breast, neck and back, cinereous gray, with large brown spots between the shoulders: wings and tail dusky brown, varie-

gated with ash-colour; middle and lesser wing-coverts tipped with white: throat white with brown scales; abdomen and sides grayish white, mixed with reddish: bill yellow at the base, black at the tip: feet yellowish. (*Egg.*) Of a uniform bluish green, like that of the next species, but rather larger.

A native of the European Alps, but has twice occurred in this country. In the first instance, a pair were observed at Cambridge by Dr Thackeray, Nov. 23, 1822, and one, a female, killed, which is at present in his collection. In the second, a single individual was shot (a few years since) on the borders of Epping Forest. Said to build in the clefts of rocks, and under the eaves of houses. Food insects and seeds.

43. *A. modularis*, Cuv. (*Hedge Accentor.*)—Crown of the head cinereous, with brown streaks: back and wing-coverts with large brown spots: throat and breast bluish gray: middle of the abdomen whitish.

A. modularis, *Temm. Man. d'Orn.* tom. i. p. 249. *Hedge Accentor*, *Selb. Illust.* vol. i. p. 248. pl. 43*. f. 4. *Hedge Warbler*, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. i. p. 244.

DIMENS. Entire length five inches eight lines: length of the bill (from the forehead) five lines, (from the gape) seven lines; of the tarsus ten lines and a half; of the tail two inches six lines; from the carpus to the end of the wing two inches seven lines: breadth, wings extended, eight inches five lines.

DESCRIPT. Top of the head cinereous, with brown streaks: back and wing-coverts yellowish brown, with a large darker-coloured spot in the centre of each feather; middle wing-coverts tipped with whitish: sides of the neck, throat, and breast, bluish gray: rump and flanks yellowish brown; middle of the abdomen whitish: under tail-coverts brown, edged with white: bill dusky: irides hazel: feet yellowish brown. (*Egg.*) Of a uniform bluish green: long. diam. nine lines and a half; trans. diam. six lines and a half.

Common in hedges throughout the country; and in song the greater part of the year. Feeds principally on insects. Nest formed of moss and wool, and lined with hair; placed in shrubs and low bushes. Eggs four to six in number; hatched early in April.

GEN. 18. SYLVIA, Lath.

(1. ERITHACA, *Swains.*)

44. *S. Rubecula*, Lath. (*Redbreast.*)—Upper parts brownish gray, tinged with olive: throat and breast bright ferruginous.

S. Rubecula, *Temm. Man. d'Orn.* tom. i. p. 215. *Redbreast*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 188. pl. 46. f. 2. *Bew. Brit. Birds*, vol. i. p. 235.

DIMENS. Entire length five inches nine lines: length of the bill (from the forehead) six lines, (from the gape) eight lines and a half; of the tarsus one inch; from the carpus to the end of the wing two inches eleven lines: breadth, wings extended, nine inches two lines.

DESCRIPT. (*Male.*) Head, and all the upper parts of the body, grayish brown with a tinge of olive: forehead, cheeks, throat, and breast, bright ferruginous red, edged round with ash-gray: belly white: greater and middle wing-coverts sometimes tipped with pale reddish orange: irides black: feet yellowish brown. (*Female.*) Upper parts cinereous brown; the red on the breast not so bright, with the surrounding margin of ash-gray less conspicuous. (*Young of the year.*) Upper parts olive-gray, speckled with pale reddish spots: the breast faintly tinged with reddish yellow, and marked with fine transverse streaks of olive-brown. (*Egg.*) Ground-colour white, spotted with pale yellow brown: long. diam. nine lines and a half; trans. diam. seven lines and a half.

A common and generally diffused species. Frequents gardens and woods. Feeds on insects and worms. In song nearly all the year. Nest formed of moss, dead leaves, and stalks of plants, and lined with hair; placed on the ground, occasionally in the holes of trees and old buildings. Has often two broods in the season, the first of which is sometimes hatched as early as the end of March. Eggs five to seven in number.

(2. PHENICURA, Swains.)

45. *S. Suecica*, Lath. (*Blue-throated Warbler.*)—Upper parts ash-brown tinged with olive: throat, and fore part of the neck blue, with a central white spot.

S. Suecica, Temm. *Man. d'Orn.* tom. i. p. 216. Blue-throated Warbler, *Lath. Syn.* vol. ii. p. 444. Blue-throated or Swedish Redbreast, *Shaw, Nat. Misc.* vol. xvi. pl. 661. Blue-throated Redstart, *Sebb. Illust.* vol. i. p. 195. pl. 160. f. 2, 3.

DIMENS. Entire length five inches ten lines: length of the bill (from the forehead) five lines and a half, (from the gape) nine lines: of the tarsus one inch and half a line; of the tail two inches two lines; from the carpus to the end of the wing two inches eleven lines.

DESCRIPT. (*Adult male.*) Upper parts brownish ash, tinged with olive; a whitish streak above the eye; throat, and fore part of the neck, bright azure, in the centre of which is a pure white spot, disappearing in advanced age; beneath the blue a border of black, and beyond this a broader one of red: belly, thighs, and vent, white: basal half of the tail rust-red, the remainder black: the two middle feathers similar to the back, and of one colour throughout. (*Female.*) Upper parts resembling those of the male: throat, and fore part of the neck, white; upper part of the breast dusky, tinged with cinereous and azure, beneath which is a faint indication of the transverse red band; the rest of the under parts dirty white. (*Young.*) Plumage brown, spotted with whitish; a large white space on the throat. (*Egg.*) Uniform greenish blue: long. diam. eight lines; trans. diam. five lines and a half.

Extremely rare in this country. A single individual killed on Newcastle Town Moor, in May 1826, is now in the Museum at that place. Said to reside in forests, and to breed in the holes of decayed trees.

46. *S. Phenicurus*, Lath. (*Redstart.*)—Bluish gray above: throat black; breast, rump, and under tail-coverts, red: second quill equal to the sixth.

S. Phœniceurus, *Temm. Man. d'Orn.* tom. i. p. 220. Redstart, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 191. pl. 46. f. 3. *Bew. Brit. Birds*, vol. i. p. 239.

DIMENS. Entire length five inches seven lines: length of the bill (from the forehead) five lines and a half, (from the gape) eight lines and a half; of the tarsus eleven lines; of the tail two inches five lines; from the carpus to the end of the wing three inches: breadth, wings extended, nine inches four lines.

DESCRIPT. (*Adult male.*) Forehead white: base of the bill, lore, cheeks, throat, and a portion of the under surface of the neck, black: head, upper part of the neck, and back, deep bluish gray: breast, rump, flanks, and under tail-coverts, bright rust-red; tail the same, excepting the two middle feathers, which are brown: abdomen whitish: feet black. (*Adult female.*) Upper parts reddish gray: quills and coverts edged with reddish yellow: throat white; in very old individuals dusky, mixed with reddish: breast, flanks, and under tail-coverts pale reddish orange: abdomen whitish: rump rust-red: tail the same, the two middle feathers excepted, which are brown edged with red. (*Young males of the year.*) Upper parts reddish ash; no white on the forehead; the black on the throat, and the red on the breast, variegated with white. (*Egg.*) Uniform grayish blue: long. diam. eight lines and a half; trans. diam. six lines and a half.

A summer visitant, first appearing about the second week in April. Common in most parts of the kingdom, but said to be rare in some of the western counties. Frequents gardens and woods, and builds in the holes of trees and old walls. Nest composed of moss, and lined with hair and feathers. Eggs five to seven in number. Young broods fledged about the second week in June.

47. S. *Tithys*, Scop. (*Black Red-tail.*)—Upper parts bluish ash: throat and breast black: rump and under tail-coverts red: second quill equal to the seventh.

S. *Tithys*, *Temm. Man. d'Orn.* tom. i. p. 218. Phœniceura *Tithys*, *Jard. and Selb. Orn.* pl. 86. figs. 1 and 2. Black Red-tail, *Lath. Syn.* vol. ii. p. 426. *Tithys* Redstart, *Selb. Illust.* vol. i. p. 193. pl. D. f. 1, 2.

DIMENS. Entire length five inches three lines. **TEMM.**

DESCRIPT. (*Male.*) Upper parts bluish ash: space between the eye and the bill, cheeks, throat and breast, deep black, passing into bluish ash on the belly and sides: rump and under tail-coverts rust-red: the two middle tail feathers brown; the others red: lower part of the abdomen whitish: greater wing-coverts edged with pure white. (*Female.*) Upper and under parts cinereous, dull above, brighter beneath, passing into whitish on the lower part of the abdomen: quills and wing-coverts dusky, edged with cinereous gray: rump, tail, and under tail-coverts reddish, not so bright as in the male. (*Egg.*) White: long. diam. ten lines; trans. diam. seven lines.

First added to the British Fauna by Mr Gould, who obtained a specimen which was shot at Kilburn, near London, Oct. 25, 1829. (See *Zool. Journ.* vol. v. p. 102.) Two other individuals were procured during the Summer of 1830, one near Bristol, the other at Brighton. Common on some parts of the Continent. Habits said to resemble those of the last species.

(3. SALICARIA, *Selby*.)

48. *S. Locustella*, Lath. (*Grasshopper Warbler*).—Upper plumage olivaceous brown, spotted with dusky: tail of one colour, long, and very much cuneated.

S. Locustella, *Temm. Man. d'Orn.* tom. i. p. 184. Grasshopper Warbler, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. i. p. 199. pl. 45**. f. 1. *Bew. Brit. Birds*, vol. i. p. 247.

DIMENS. Entire length five inches six lines: length of the bill (from the forehead) five lines, (from the gape) seven lines and a half; of the tarsus nine lines and a half; of the tail two inches three lines; from the carpus to the end of the wing two inches four lines and a half.

DESCRIPT. All the upper parts olivaceous brown, with a dusky spot in the centre of each feather, most conspicuous on the head and back: throat white, bounded by a circle of small oval brown spots; breast and belly yellowish white; under tail-coverts pale yellowish brown, with dusky streaks occupying the shafts of the feathers: quills and tail dusky brown, with pale olive edges: feet yellowish brown: hind claw shorter than the toe. (*Egg*.) Pale reddish white, speckled all over with darker red brown: long. diam. eight lines; trans. diam. six lines.

Visits this country in April, but is not generally distributed, and no where very plentiful. Haunts thickets and furzy commons, principally in damp situations, and is of shy habits, seldom exposing itself to view. Note resembling that of the *gryllidæ*. Nest artfully concealed, placed on the ground, or in thick bushes of furze and bramble; composed of dried stalks and goose-grass, and lined with fibrous roots. Eggs six in number. Food principally small coleopterous insects.

49. *S. Phragmitis*, Bechst. (*Sedge Warbler*).—Upper plumage olivaceous brown, tinged with yellow, and spotted with dusky: above the eye a broad white streak: tail moderate.

S. Phragmitis, *Temm. Man. d'Orn.* tom. i. p. 189. Sedge Warbler, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 201. pl. 45**. f. 2. Reed Warbler, *Bew. Brit. Birds*, vol. i. p. 246.

DIMENS. Entire length five inches five lines: length of the bill (from the forehead) five lines, (from the gape) seven lines and a half; of the tarsus nine lines and a half; of the tail two inches one line; from the carpus to the end of the wing two inches four lines: breadth, wings extended, seven inches five lines.

DESCRIPT. Top of the head, back, scapulars and wing-coverts, olivaceous brown, with a dark spot in the centre of each feather: neck, lower part of the back, rump and upper tail-coverts, plain yellowish brown: a broad and very distinct white streak above the eye, arising from the base of the bill, and extending the length of the head: all the under parts yellowish white, with a reddish tinge on the breast, sides, and under tail-coverts: quills brown with pale edges: tail deep yellowish brown: bill dusky above, whitish beneath: irides dark hazel: feet yellowish brown: soles yellow. (*Egg*.) Greenish white; mottled all over with yellow brown: long. diam. eight lines; trans. diam. six lines.

Very abundant in marshy districts, and by the sides of rivers, visiting this country about the third week in April. Song much varied, imitative

of that of other birds; kept up unceasingly during the breeding season, and often heard in the night. Nest generally suspended between the stems of reeds at a little distance from the ground; composed of dried stalks, with the addition of a little moss, and lined with hair and the finer grasses. Eggs five or six in number; hatched towards the end of May, or beginning of June.

50. *S. arundinacea*, Lath. (*Reed-Wren*.) — Upper parts plain olivaceous brown, without spots: between the bill and the eye a white streak.

S. arundinacea, Temm. *Man. d'Orn.* tom. 1. p. 191. *Reed-Wren*, *Mont. Orn. Dict. Selb. Illust.* vol. 1. p. 203. pl. 45**. f. 3.

DIMENS. Entire length five inches four lines: length of the bill (from the forehead) five lines and a half, (from the gape) eight lines; of the tarsus ten lines; of the tail two inches three lines; from the carpus to the end of the wing two inches four lines: breadth, wings extended, seven inches five lines.

DESCRIPT. Upper parts of a plain uniform olive brown without spots; rump and tail faintly tinged with reddish: quills brown with cinereous edges: between the bill and the eye a narrow white streak, but not continued *above* the eye as in the last species: under parts yellowish white, lightest on the throat and down the middle of the belly; sides inclining to rufous: bill broader throughout, but especially at the base, and also somewhat longer, than in the *S. Phragmitis*; in colour dusky above, yellowish beneath, and along the margin of the upper mandible: irides light hazel: feet pale brown; soles yellow. (*Egg*.) Greenish white, spotted and speckled with ash-green and light brown; the markings darkest, and most numerous at the larger end: long. diam. nine lines; trans. diam. six lines and a half.

Found in the same situations with the last species, but is much less plentiful, and not so generally diffused. Habits similar. Food the smaller species of *libellulæ*, and other insects. Nest of an oblong panier-shaped form, suspended like that of the *S. Phragmitis*; composed of grasses and the seed-branches of the reed. Eggs four or five in number.

(4.) *S. Cetti*, Temm. *Man. d'Orn.* tom. 1. p. 194.

According to Temminck, this species has been killed in England. It does not, however, appear to be known to our own naturalists, nor am I aware that it is to be found in any of our British collections.

(4. PHILOMELA, *Swains*.)

51. *S. Luscinia*, Lath. (*Nightingale*.) — Upper plumage reddish brown; beneath, cinereous white.

S. Luscinia, Temm. *Man. d'Orn.* tom. 1. p. 195. *Nightingale*, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. 1. p. 206. pl. 46. f. 1. *Bew. Brit. Birds*, vol. 1. p. 231.

DIMENS. Entire length six inches two lines: length of the bill (from the forehead) six lines, (from the gape) nine lines and a half: of the tarsus one inch one line; of the tail two inches eight lines; from the carpus to the end of the wing three inches two lines: breadth, wings extended, nine inches seven lines.

DESCRIPT. All the upper parts reddish brown: tail rust-red: breast and sides pale ash, passing into yellowish white: throat and belly whitish: quills pale dusky, edged with reddish brown: bill brown: irides hazel: feet yellowish brown. (*Egg.*) Uniform olive-brown: long, diam. ten lines; trans. diam. eight lines and a half.

Common in the southern, midland, and eastern counties; rare in the western and northern. Has not been observed in Cornwall or in Devonshire, excepting on the eastern borders. First heard about the middle or end of April: song continued till the first or second week in June. Frequents woods, copses, and tall hedges. Nest placed on or near the ground, sometimes suspended between the stems of herbaceous plants; of a very deep, oval form; composed principally of withered leaves, and lined with fine grasses. Eggs four to six in number; hatched towards the end of May, or beginning of June.

(5. CURRUCA, *Bechst.*)

52. *S. atricapilla*, Lath. (*Black-cap Warbler*).—Top of the head and occiput deep black (*male*), or reddish brown (*female*); rest of the upper plumage greenish ash.

S. atricapilla, *Temm. Man. d'Orn.* tom. i. p. 201. *Black-cap*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 209. pl. 46. f. 5. *Bew. Brit. Birds*, vol. i. p. 249.

DIMENS. Entire length five inches eight lines: length of the bill (from the forehead) four lines and a half; (from the gape) seven lines and a half: of the tarsus nine lines and a half; of the tail two inches four lines; from the carpus to the end of the wing three inches: breadth, wings extended, eight inches six lines.

DESCRIPT. (*Male*.) Forehead, crown, and occiput, deep black: rest of the upper parts cinereous, with a slight tinge of olive-green: neck and breast grayish ash; belly and vent grayish white: bill and feet dusky. (*Female*.) Crown of the head reddish brown; rest of the plumage resembling that of the male but rather darker. (*Egg.*) Pale greenish white, mottled with light brown and ash-colour, with a few spots and streaks of dark brown: long, diam. nine lines; trans. diam. seven lines.

A common and widely dispersed species, visiting this country about the same time as the Nightingale. Song melodious, and rather powerful, heard till the middle, or occasionally the end of July. Is partial to orchards and gardens. Nest placed in a low bush; loosely put together; constructed of bents and dried stalks, patched on the outside with threads and cobwebs, and lined with fibrous roots and a few long hairs. Eggs five in number, on which the male sits occasionally, as well as the female.

53. *S. hortensis*, Lath. (*Greater Pettychaps*).—Upper parts grayish brown, tinged with olive: orbits white: tail of one colour.

S. hortensis, *Temm. Man. d'Orn.* tom. i. p. 206. *Greater Pettychaps*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 211. pl. 46. f. 4. *Passerine Warbler*, *Bew. Brit. Birds*, vol. i. p. 243.

DIMENS. Entire length five inches ten lines: length of the bill (from the forehead) five lines, (from the gape) eight lines; of the tarsus ten lines; of the tail two inches two lines; from the carpus to the end of the

wing two inches eleven lines: breadth, wings extended, eight inches six lines.

DESCRIPT. All the upper parts cinereous brown, tinged with olive: orbits white: below the ear on each side of the neck a patch of ash-gray: throat whitish: breast and sides yellowish gray inclining to brown: belly, vent and under tail-coverts, nearly pure white: quills and tail dusky, with pale edges: feet bluish gray. (*Egg.*) Greenish white ground, speckled with two shades of ash-green: long, diam. nine lines; trans. diam. six lines and a half.

Not uncommon in many parts of the country, making its first appearance about the beginning of May. Frequents gardens, copses, and thick hedges. Song resembling that of the Blackcap, but less powerful; continued till the middle of July. Nest of flimsy structure; formed of the decayed stems of goose-grass, and other fibrous plants. Eggs four or five in number, very like those of the last species.—*Obs.* The *Garden Warbler* and the *Passerine Warbler* of Bewick are probably both referable to the present species. Montagu, however, thought that the *Passerine Warbler* of the author just mentioned was the same as the *Reed Wren*.

54. *S. cinerea*, Lath. (*White-Throat*).—Top of the head cinereous; back and wing-coverts grayish brown inclining to rufous: under parts white; the breast tinged with rose-red: tail particoloured.

S. cinerea, *Temm. Man. d'Orn.* tom. 1. p. 207. *White-Throat*, *Mont. Orn. Dict.* *Selb. Illust.* vol. 1. p. 213. pl. 46. f. 7. *Bew. Brit. Birds*, vol. 1. p. 251.

DIMENS. Entire length five inches ten lines: length of the bill (from the forehead) four lines and a half, (from the gape) seven lines; of the tarsus nine lines and a half; of the tail two inches eight lines; from the carpus to the end of the wing two inches eight lines: breadth, wings extended, eight inches seven lines.

DESCRIPT. (*Male*.) Crown of the head, space between the eye and the bill, cheeks, and ear-coverts, cinereous; rest of the upper parts cinereous brown, inclining to rufous on the back: quills dusky; secondaries and greater coverts broadly edged with brownish red: throat and middle of the abdomen white; breast faintly tinged with rose-red; flanks and thighs reddish gray: tail deep brown; outer feather white, except at the base of the inner web; the next tipped with white: bill dusky above, paler beneath: irides yellowish: legs yellowish brown. (*Female and young.*) Upper parts more inclining to reddish brown; breast white, without the rosy tinge; the outer tail-feather reddish, instead of white. (*Egg.*) Greenish white ground, spotted and speckled with ash-colour and two shades of ash-green: long. diam. nine lines; trans. diam. six lines and a half.

Common in hedges and thickets throughout the kingdom. First seen about the third week in April. Song often exerted on wing, accompanied by peculiar jerks and gesticulations of the body. Nest similar to that of the last species; placed in low bushes, where there is thick covert. Eggs four or five in number. Young broods fledged about the end of May.

55. *S. Curruca*, Lath. (*Lesser White-Throat*).—Crown of the head cinereous; back and wing-coverts

grayish ash, tinged with brown: under parts silvery white: tail particoloured.

S. Curruca, *Temm. Man. d'Orn.* tom. i. p. 209. Lesser White-Throat, *Lath. Syn. Supp.* p. 185. pl. 113. *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. i. p. 253. *Selb. Illust.* vol. i. p. 215. pl. 100. f. 4.

DIMENS. Entire length five inches three lines.

DESCRIPT. Crown of the head cinereous; space between the eye and the bill, and ear-coverts, the same, but darker-coloured: rest of the upper parts grayish ash, inclining to brown: quills dusky brown, edged with cinereous: all the under parts, from the throat to the vent, silvery white: tail dusky; outer feather white, except at the base of the inner web; the two next faintly tipped with white: bill dusky; the lower mandible pale at the base: irides yellowish: feet dusky lead-colour. (*Egg.*) White, sparingly spotted and speckled, principally at the larger end, with ash-colour and light brown: long. diam. eight lines; trans. diam. six lines.

Common in the eastern, and in some of the midland counties, making its first appearance about the same time as the last species, but apparently rare, or altogether unknown, in many parts of the kingdom. Song soft and inward; generally ending in a shrill shivering cry, which last note is continued at intervals till near the middle of July. Chiefly frequents gardens and copses, concealing itself in the thickest covert. Nest placed in a low bush; of very flimsy structure, composed of dry bents mixed with wool, and lined with finer grasses and a few hairs. Eggs five in number; incubation commencing about the 20th of May.

(6. SYLVIA, *Selb.*)

56. *S. sibilatrix*, Bechst. (*Wood-Wren.*) — Upper parts bright yellowish green: belly and under tail-coverts pure white: wings with the second quill equal to the fourth; third and fourth with their outer webs sloped off at the extremity.

S. sibilatrix, *Temm. Man. d'Orn.* tom. i. p. 223. Wood-Wren, *Lamb in Linn. Trans.* vol. ii. p. 245. pl. 24. *Mont. in Linn. Trans.* vol. iv. p. 35. *Mont. Orn. Dict.* *Selb. Illust.* vol. i. p. 224. pl. 47. f. 2. Yellow Wren, *Bew. Brit. Birds*, vol. i. p. 255.

DIMENS. Entire length five inches three lines: length of the bill (from the forehead) four lines and a half, (from the gape) six lines and a half; of the tarsus nine lines; of the tail two inches one line; from the carpus to the end of the wing three inches: breadth, wings extended, eight inches eight lines.

DESCRIPT. All the upper parts bright yellow green: above the eye a broad streak of bright sulphur-yellow, arising on the forehead at the base of the upper mandible and terminating on the temples: sides of the head, throat, fore part of the neck, axillæ, and thighs, pale primrose-yellow; rest of the under parts pure white: quills dusky, edged with green: tail slightly forked, extending eight lines and a half beyond the tips of the folded wings; dusky, with each feather, except the outermost, likewise edged with bright yellow-green: bill and legs pale brown. (*Egg.*) White, spotted and speckled all over with dark purple, red, and ash-colour: long. diam. eight lines; trans. diam. six lines.

Common in the southern, western, midland, and some of the northern counties, but not generally diffused. In Cambridgeshire very rare. Haunts woods and tall trees; being first heard towards the end of April or beginning of May. Note very peculiar, resembling the word *tree* repeated several times in succession, at first slowly, but afterwards in a hurried manner, and accompanied by a singular shake of the wings. Nest placed on the ground, of an oval form, constructed of dry grass, dead leaves, and a little moss, lined with finer grass and a few hairs. Eggs six in number.

57. *S. Trochilus*, Lath. (*Willow-Wren*.) — Upper parts pale olive-green, tinged with yellow: under tail-coverts yellowish white: wings with the second quill equal to the sixth; the third, fourth, and fifth, with their outer webs sloped off at the extremity.

S. Trochilus, *Temm. Man. d'Orn.* tom. i. p. 224. Yellow Wren, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 226. pl. 47. f. 3. Willow-Wren, *Bew. Brit. Birds*, vol. i. p. 257.

DIMENS. Entire length four inches eleven lines: length of the bill (from the forehead) four lines, (from the gape) six lines; of the tarsus nine lines; of the tail two inches three lines; from the carpus to the end of the wing two inches six lines and a half: breadth, wings extended, eight inches.

DESCRIPT. Distinguished from the last principally by its paler colour, and shorter wings; the difference in size not very obvious. Upper parts pale olive-green, with a yellowish tinge, inclining in some specimens to cinereous: from the base of the bill a bright streak of primrose-yellow extending over each eye to the temples, but narrower than in the *S. sibilatrix*: all the under parts yellowish white; the yellow tinge darkest on the breast, sides, axillæ, and ridges of the wings; palest on the middle of the belly: quills and tail brownish ash, faintly edged with olive-green; the latter extending eleven lines beyond the tips of the folded wings, with the two middle feathers shorter than the others: base of the under mandible yellowish: feet light brown. *Obs.* The colour of this species, especially the intensity of the yellow tinge, varies much in different individuals. The variation, however, results more from a difference of age than of sex. The male and female are equally pale in the adult state. The yellowest birds, which are also generally the smallest, occur in the early part of the Spring, and appear to be the young males of the preceding year. (*Egg.*) White, with numerous small specks of pale red: long. diam. seven lines and a half; trans. diam. six lines.

Of frequent occurrence in gardens, plantations, hedges, willow-grounds, and a variety of other situations. On the whole a much more generally diffused species than the last, but, according to Montagu, does not extend so far westward. Song pleasing, but simple, consisting of several plaintive notes in a regularly descending scale; heard from the middle of April to the beginning of July. Nest placed on the ground, nearly spherical, with a small opening at the side near the top; composed of dry grass, stalks, and other herbage, and profusely lined with feathers. Eggs five to seven; hatched the end of May or beginning of June.

58. *S. Hippolais*, Lath. (*Chiff-Chaff*.) — Upper parts olive-green, tinged with yellow; beneath yellowish white:

wings with the second quill equal to the seventh; third, fourth, fifth, and sixth, with their outer webs sloped off at the extremity.

S. Hippolais, *Lath. Ind. Orn.* vol. 11. p. 507. *S. rufa*, *Temm. Man. d'Orn.* tom. 1. p. 225. Lesser Pettychaps, *Mout. Orn. Dict. and Supp. Selb. Illust.* vol. 1. p. 222. pl. 47. f. 1. Chiff-Chaff, *Bew. Brit. Birds*, vol. 1. p. 258.

DIMENS. Entire length four inches six lines: length of the bill (from the forehead) four lines, (from the gape) six lines and a half; of the tarsus nine lines; of the tail one inch seven lines; from the carpus to the end of the wing two inches three lines: breadth, wings extended, seven inches three lines.

DESCRIPT. Usually somewhat smaller than the *S. Trochilus*, with the wings and tail, *relatively* considered, still shorter than in that species; also to be distinguished from it by the characters of the quills above pointed out, but in colour and general appearance almost absolutely the same. Upper parts olive-green, tinged with yellow and ash-gray: between the bill and the eye, and over each eye, a narrow, faint yellowish white streak: quills cinereous brown, the outer webs edged with yellowish green: all the under parts, including the under tail-coverts, whitish tinged with yellow; the yellow having a tendency on the breast to appear in streaks: axillæ and under wing-coverts bright primrose-yellow: tail extending an inch beyond the tips of the folded wings: feet rather darker than in the last species. (*Egg.*) White, with a few specks of dark purplish red: long. diam. seven lines; trans. diam. five lines and a half.

Very abundant in some parts of England, but in others much less plentiful than the last species. Is one of the earliest of our summer visitants, making its first appearance about the middle, or towards the end of March. Chiefly frequents woods and tall trees, and is of restless habits, being always in motion in search of insects. Song consisting of only two, rather loud, hollow notes, resembling the words *chip-chop*, or *chiff-chaff*, which are occasionally heard till near the end of September. Nest similar to that of the last species, placed on the ground, or in very low bushes. *Obs.* Having compared our English *S. Hippolais* with the *S. rufa* of Temminck*, and found them agreeing closely in all their characters, I have no hesitation in considering them as the same species.

GEN. 19. MELIZOPHILUS, *Leach*.

59. *M. provincialis*, *Leach*. (*Dartford Warbler*).—Upper plumage deep grayish brown: throat, neck, and breast, ferruginous red.

Sylvia provincialis, *Temm. Man. d'Orn.* tom. 1. p. 211. *S. Dartfordiensis*, *Mont. in Linn. Trans.* vol. 1x. p. 191. Dartford Warbler, *Mout. Orn. Dict. and Supp. Selb. Illust.* vol. 1. p. 219. pl. 46. f. 6. *Bew. Brit. Birds*, vol. 1. p. 234.

DIMENS. Entire length five inches six lines. **MONT.**

DESCRIPT. (*Male.*) All the upper parts dark grayish brown: cheeks cinereous: throat, neck, and breast, deep ferruginous, inclining to purple;

* The comparison was made with a specimen of *S. rufa* named by Temminck himself, and sent by him to Mr Gould, to whose kindness I am indebted for it.

sides the same, but not so bright; middle of the belly white: quills dusky; the outer webs edged with dark cinereous, the inner with dark reddish brown: tail blackish brown: the outer feather tipped, and edged externally, with white: bill yellowish white at the base, black at the tip: feet yellowish. (*Female and young.*) Colours paler: throat more or less speckled and streaked with white. (*Egg.*) Greenish white ground, speckled all over with olivaceous brown and cinereous; the markings most numerous at the larger end: long. diam. eight lines; trans. diam. five lines and a half.

A local species; found in Kent, Middlesex, Hampshire, Sussex, Surrey, Devonshire, Cornwall, and a few other parts of England. Remains in this country throughout the year. Haunts chiefly downs and furzy commons. Feeds on insects. Nest somewhat similar to that of the Whitethroat; artfully concealed in the thickest bushes. Young broods fledged in May.

GEN. 20. REGULUS, *Cuv.*

60. *R. aurocapillus*, Selby. (*Gold-crested Regulus.*)—Upper parts greenish yellow: cheeks and region of the eyes cinereous, without any white bands: crown-feathers elongated, forming a bright yellow orange crest.

Sylvia Regulus, *Temm. Man. d'Orn.* tom. i. p. 229. Gold-crested *Regulus*, *Selb. Illust.* vol. i. p. 229. pl. 47. f. 4 & 5. Golden-Crested Wren, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. i. p. 260.

DIMENS. Entire length three inches six lines: length of the bill (from the forehead) three lines and a half, (from the gape) five lines; of the tarsus seven lines and a half; of the tail one inch five lines; from the carpus to the end of the wing one inch eleven lines: breadth, wings extended, five inches six lines.

DESCRIPT. Upper parts of the plumage olive-green, tinged with yellow: feathers on the top of the head long and narrow, forming a crest, of a rich orange-yellow, the sides bordered with black: base of the bill, cheeks, throat, and region of the eyes, cinereous; the rest of the under parts the same, but with a tinge of yellow: quills dusky, edged with yellowish green: secondaries crossed by a black bar; wing-coverts dusky brown, edged with yellow and tipped with white: tail blackish brown; the feathers edged outwardly with yellowish green, inwardly with whitish: bill black: irides hazel: feet brown. (*Female.*) Colours generally paler: crest on the head smaller, and of a pale lemon-yellow; the lateral edging of black narrower, and less conspicuous. (*Egg.*) Dull pale reddish white: long. diam. six lines: trans. diam. five lines.

Not uncommon, and generally dispersed throughout the kingdom. Does not migrate. Frequents woods and plantations, especially those of fir. Song weak and inward; heard as early as February. Nest generally suspended from the under surface of the branches of spruce firs; formed principally of moss, and lined with feathers. Eggs seven to ten in number. Breeds early in the year.

61. *R. ignicapillus*, Nob. (*Fire-crested Regulus.*)—Upper parts yellowish green, passing into pure yellow on the sides of the neck; cheeks with three longitudinal streaks, one black and two white: crown-feathers elongated, of a brilliant flame-red.

Sylvia ignicapilla, Temm. *Man. d'Orn.* tom. i. p. 231. Fire-crested Wren, Gould, *Europ. Birds*, part 3.

DIMENS. Entire length four inches: length of the bill (from the forehead) four lines and a half, (from the gape) five lines and a half; of the tarsus seven lines and a half; of the tail one inch nine lines; from the carpus to the end of the wing two inches and half a line.

DESCRIPT. Closely resembling the last species, but somewhat larger, with the bill longer and broader at the base. (*Adult male.*) Upper parts yellowish tinged with green, passing into nearly pure yellow on the sides of the neck; crest of a brilliant flame-red, bordered in front and at the sides of the crown with deep black; cheeks particularly distinguished by three longitudinal streaks, of which there is no indication in the *R. aurocapillus*; one of these, which is black, passes directly through the eyes in a line from the base of the upper mandible to the ear-coverts; the other two are white, and are situate one above, the other beneath the eyes; forehead cinereous, with a pale reddish tinge: wings and tail the same as in the last species: under parts more approaching to white: bill black: irides deep brown: tarsi brown; toes and claws yellowish. (*Female.*) The longitudinal streaks on the sides of the face the same as in the male, but the colours more obscure: crest of a dull orange: sides of the neck olivaceous green, instead of yellow: in other respects the plumage of the two sexes is similar. (*The young of the year before the first moult.*) Only to be distinguished from those of the last species, by the longer and broader bill: cheeks cinereous, without any appearance of the longitudinal streaks: crest of a pale lemon-yellow, scarcely developed: forehead, and sides of the neck, cinereous: upper parts not so bright as in the adult; under parts cinereous, tinged with yellow.

A single individual of this species (a young bird of the year) was killed in a garden at Swaffham Bulbeck in Cambridgeshire, in August 1832. Since then others have been observed at Brighton by Mr J. E. Gray. Is probably to be met with in other parts of the country, though from its general resemblance to the *R. aurocapillus*, it is easily overlooked. According to Temminck, this last species resides chiefly at the tops of trees; the *R. ignicapillus*, more on the lower branches and in small bushes. The habits of the two in other respects, and nidification, are similar.

GEN. 21. MOTACILLA, Linn.

* *Hind claw moderate; much curved.*

62. *M. alba*, Linn. (*Pied Wagtail*).—Plumage variegated with black and white.

M. alba, Temm. *Man. d'Orn.* tom. i. p. 255. Pied Wagtail, *Selb. Illust.* vol. i. p. 251. pl. 49. f. 1. *Bew. Brit. Birds*, vol. i. p. 226. White Wagtail, *Mont. Orn. Dict.*

DIMENS. Entire length seven inches five lines: length of the bill (from the forehead) six lines, (from the gape) eight lines; of the tarsus eleven lines; of the hind toe, claw included, seven lines; of the tail three inches four lines and a half: from the carpus to the end of the wing three inches four lines: breadth, wings extended, ten inches nine lines.

DESCRIPT. (*Adult in summer plumage.*) Forehead, cheeks, and sides of the neck, white; crown of the head, nape, throat, breast, and upper part of the body, black; belly and vent white: quills and greater wing-coverts edged with white: the two outer tail-feathers nearly all white;

the others black : bill black : irides dusky. (*Winter plumage.*) Throat and fore part of the neck white : a crescent-shaped black patch on the upper part of the breast : all the upper parts inclining to blackish gray. (*Young of the year.*) Crown of the head, and all the upper parts, ash-gray, inclining to bluish gray ; wing-coverts broadly edged and tipped with white : on the upper part of the breast a crescent-shaped patch of dusky ash, the horns of the crescent extending upwards to the ears in a narrow line on each side of the neck : throat, cheeks, under part of the neck, and lower part of the breast, dirty white : abdomen nearly pure white : quills blackish, with an oblong white spot on their inner webs. (*Egg.*) White, speckled with ash-colour : long. diam. nine lines ; trans. diam. seven lines.

A common and well-known species, frequenting pastures and newly ploughed lands, more especially those in the vicinity of water. Runs with great celerity, and occasionally wades in shallow streams in pursuit of aquatic insects. Migrates in the North of England, but is stationary in the southern counties. Nest on the ground, or in the holes of trees and old buildings ; composed of moss, fibres, and wool, and lined with hair. Eggs four or five.

63. *M. Boarula*, Linn. (*Gray Wagtail.*)—Head and back bluish gray ; rump and under parts bright yellow : tail very long.

M. Boarula, *Temm. Man. d'Orn.* tom. i. p. 257. *Gray Wagtail*, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. i. p. 253. pl. 49. f. 2. *Bew. Brit. Birds*, vol. i. p. 228.

DIMENS. Entire length seven inches nine lines : length of the bill (from the forehead) five lines, (from the gape) seven lines and a half ; of the tarsus ten lines ; of the hind toe, claw included, six lines ; of the tail three inches ten lines ; from the carpus to the end of the wing three inches one line and a half.

DESCRIPT. (*Summer plumage.*) Head, neck, back, and scapulars, bluish gray : above the eyes a pale streak : throat black : rump, and under parts, bright yellow, the former tinged with green : quills and six middle tail-feathers grayish black, more or less edged with yellowish white ; outer tail-feather entirely white ; the two next white, with a part of the outer web black : feet dusky brown. (*Winter plumage.*) Throat yellowish white ; under parts of a paler yellow. *Obs.* Female at all times paler than the male. (*Egg.*) White, nearly covered with reddish yellow brown, varying in intensity : long. diam. eight lines and a half ; trans. diam. seven lines.

A regular *winter* visitant in the southern parts of the kingdom, but in the northern a *summer* visitant. Habits and nidification very similar to those of the last species. Is said to have two broods in the year, the first of which is generally fledged about the end of May.

**** Hind claw elongated ; but slightly curved.**

64. *M. flava*, Ray. (*Yellow Wagtail.*)—Head, and all the upper parts of the body, olive-green : streak above the eye, and under parts, bright yellow : tail moderate.

M. flava, Ray, *Syn. Av.* p. 75. Yellow Wagtail, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. 1. p. 255. pl. 49. f. 3. *Bew. Brit. Birds*, vol. 1. p. 229.

DIMENS. Entire length six inches eight lines: length of the bill (from the forehead) five lines and a half, (from the gape) eight lines; of the tarsus eleven lines; of the hind toe, claw included, eight lines and a half; of the tail two inches eleven lines; from the carpus to the end of the wing three inches: breadth, wings extended, nine inches nine lines.

DESCRIPT. (*Male*.) Crown of the head, nape, and ear-coverts, pale olive-green; back, rump, and scapulars, of the same colour but darker: over the eyes a bright yellow streak: all the under parts bright yellow: quills and coverts blackish brown, edged and tipped with yellowish white: tail dusky; the two middle feathers edged with pale olive; the two outer ones with the whole of their external, and a large part of their inner webs white: bill and feet black. (*Female*.) Upper plumage inclining more to cinereous: throat yellowish white; the rest of the under parts yellow, but not so bright as in the male. (*Young of the year*.) Resembling the female, but still paler: upper parts dull cinereous, very faintly tinged with olivaceous: all the under parts yellowish white: greater and lesser coverts broadly tipped with white, forming two bars across the wings: tail as in the adult. (*Egg*.) White, mottled nearly all over with yellow and ash-brown: long. diam. eight lines and a half; trans. diam. six lines and a half.

A migratory species, visiting this country about the end of March, or beginning of April. Frequents downs, and arable lands, and is rather less attached to water than either of the foregoing species. Nest placed on the ground; composed of dried stalks and fibres, and lined with hair. Eggs four or five in number.

65. *M. neglecta*, Gould. (*Blue-headed Wagtail*.)—Head bluish lead-colour; streak above the eye white: body above olivaceous green, beneath yellow.

M. neglecta, Gould in *Proceed. of Zool. Soc.* 1832, p. 129. *Id. Europ. Birds*, part 3. *M. flava*, Temm. *Man. d'Orn.* tom. 1. p. 260.

DIMENS. Entire length six inches. GOULD.

DESCRIPT. Closely resembling the last species, from which it scarcely differs excepting in the colour of the head and streak above the eye: the former, instead of being olivaceous like the rest of the upper parts, is of a fine lead gray, approaching to blue; the latter, as well as a second streak beneath the eyes, is white. The female has the colours paler, and the eye-streaks indistinctly marked. (*Egg*.) "Greenish olive, with light flesh-coloured blotches." GOULD.

Mr Gould has the merit of having first distinguished this from the *M. flava*, with which it had been previously confounded. The differences between them, he observes, do not depend upon the season, he having obtained specimens of each killed in the same month. The present species, which appears to be the only one known to continental authors, in our own country has hitherto occurred but once. This individual, which was an old bird, was killed near Colchester by Mr H. Doubleday, in the Autumn of 1834. A pair were seen. Food and nidification similar to those of the *M. flava*.

GEN. 22. ANTHUS, *Bechst.*

* *Hind claw longer than the toe; very little curved.*

66. *A. Richardi*, Vieill. (*Richard's Pipit.*)—Upper plumage deep brown, with lighter edges: breast spotted: tarsi very long.

A. Richardi, *Temm. Man. d'Orn.* tom. i. p. 263. *Vigors in Zool. Journ.* vol. i. p. 411. pl. 14. *Bew. Brit. Birds*, vol. i. p. 379. *Richard's Pipit*, *Selb. Illust.* vol. i. p. 264. pl. 100. f. 5.

DIMENS. Entire length six inches seven lines: length of the hind toe, claw included, one inch. TEMM.

DESCRIPT. Plumage on the crown of the head, back, and scapulars, deep olive-brown; the edges of the feathers somewhat paler: a broad streak above the eye, temples, throat, belly and vent, pure white; breast and sides reddish yellow, the former marked with lanceolate dusky spots: wings blackish brown; quills and coverts broadly edged with yellowish white: tail dusky, edged in the same manner; outer feather entirely white; the next with the greater portion white: bill somewhat stronger, and rather more elongated, than in the other species of this genus; upper mandible brown; lower mandible yellow: irides brown: feet yellowish: hind claw very long, almost straight. (*Egg.*) Reddish white ground, speckled with darker red and light brown: long, diam. ten lines and a half; trans. diam. seven lines and a half.

Very rare in this country. Has hitherto only occurred in two instances, both times in the neighbourhood of London. Habits unknown.

67. *A. pratensis*, Bechst. (*Meadow Pipit.*)—Upper plumage olivaceous green, with dusky spots: neck, breast, and sides, spotted: tarsi moderate.

A. pratensis, *Temm. Man. d'Orn.* tom. i. p. 269. *Meadow Pipit*, *Selb. Illust.* vol. i. p. 260. pl. 49. f. 4. *Titlark*, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. i. p. 220.

DIMENS. Entire length five inches nine lines: length of the bill (from the forehead) five lines and a half, (from the gape) eight lines; of the tarsus ten lines and a half; of the hind toe, claw included, ten lines and a half; of the tail two inches four lines and a half; from the carpus to the end of the wing three inches: breadth, wings extended, nine inches ten lines.

DESCRIPT. (*Summer plumage.*) Upper parts olivaceous green tinged with ash-gray; feathers on the head, back, and scapulars, spotted with dusky: under parts yellowish white; throat, in the *male*, during the breeding season only, with a deep reddish tinge; sides of the neck, breast, and upper part of the abdomen, marked with blackish brown spots; flanks with long dusky streaks: tail dusky; outer feather edged externally, and broadly tipped, with white; the next with a small white spot at the extremity: bill slender, dusky; base of the under mandible yellowish: irides hazel: legs yellowish brown: hind claw not so long as in the last species, slightly curved. (*Winter plumage.*) Upper parts of a brighter and more decided olive-green, with the dusky spots not occupying so large a portion of the feather: under parts ferruginous yellow, and not so much spotted. (*Egg.*) Reddish white ground, mottled all over with darker reddish brown: long, diam. nine lines; trans. diam. seven lines.

Very abundant on moors, barren heaths, and extensive fenny districts. Stays the whole year. Sings in its descent, after rising to a considerable height in the air, always returning to the ground, or to some low bush, with motionless wings and expanded tail. Nest on the ground; composed of dry grass, bents, and stalks of plants, patched on the outside with a small quantity of green moss, and lined with fine grasses and long horse-hair. Eggs five or six, hatched towards the end of May. *Obs.* The *Pipit Lark* of Montagu is this species in its winter plumage, after the autumnal moult.

** *Hind claw about equal to the toe in length, or shorter; much curved.*

68. *A. arboreus*, Bechst. (*Tree Pipit*).—Upper parts olivaceous green, with dusky spots: breast and sides spotted: wings with two transverse bars of yellowish white.

A. arboreus, *Temm. Man. d'Orn.* tom. i. p. 271. *Tree Pipit*, *Selb. Illust.* vol. i. p. 262. pl. 49. f. 5. *Field Lark*, *Mont. Orn. Dict.* *Tree Lark*, *Bew. Brit. Birds*, vol. i. p. 218.

DIMENS. Entire length six inches: length of the bill (from the forehead) five lines and a half, (from the gape) eight lines; of the tarsus nine lines; of the tail two inches six lines; from the carpus to the end of the wing three inches six lines.

DESCRIPT. Strongly resembling the last species, but always to be distinguished by the hind claw, which is shorter than the toe, and so much curved as to form the quadrant of a circle; the bill is also somewhat stronger, and more dilated at the base. Upper parts olivaceous green, tinged with cinereous; the centre of each feather, more especially those on the head and back, dusky brown: lesser and middle wing-coverts tipped with yellowish white, so as to shew a double transverse bar on the wings when closed: throat almost white: fore part of the neck, breast, and flanks, ochre-yellow, marked with dusky spots, large and oval on the breast, but assuming the form of long narrow streaks on the sides; middle of the abdomen pure white; under tail-coverts tinged with yellow, but free from spots: tail dusky brown; the outer feather with the greater portion white; the next only tipped with white: upper mandible wholly dusky, under mandible dusky at the tip, flesh-coloured at the base: irides hazel: legs flesh-coloured, inclining to yellowish brown. (*Egg.*) Grayish white ground, spotted and streaked with ash-colour and dark brown; sometimes pale purple, spotted with darker purple and dark red brown: long. diam. nine lines; trans. diam. seven lines.

A migratory species, visiting this country about the third week in April. Not uncommon in wooded districts, but rarely or never to be found in open country. Sings in its descent like the last species, but always rises from the top of some tall tree, to which it returns gradually, with expanded wings and tail. Song heard till the middle of July. Nest placed on the ground; formed of dry grass, patched externally with moss, and lined with the finer grasses. Eggs four or five.

69. *A. petrosus*, Flem. (*Rock Pipit*).—Upper parts greenish brown with darker spots: breast and sides streaked with brown: above the eye a yellowish white streak.

A. aquaticus, *Temm. Man. d'Orn.* tom. i. p. 265. *Alauda petrosa*, *Mont. in Linn. Trans.* vol. iv. p. 41. Rock Pipit, *Selb. Illust.* vol. i. p. 258. pl. 49. f. 6. Rock Lark, *Mont. Orn. Dict.* Field Lark, *Bew. Brit. Birds*, vol. i. p. 216.

DIMENS. Entire length six inches eight lines: length of the bill (from the forehead) six lines and a half, (from the gape) nine lines; of the tarsus ten lines and a half; of the hind toe, claw included, ten lines; of the tail two inches nine lines; from the carpus to the end of the wing three inches five lines: breadth, wings extended, eleven inches three lines.

DESCRIPT. (*Adult.*) Upper parts dark brown, with a tinge of olive-green; the feathers on the back, and scapulars, obscurely marked with dusky streaks: quills and coverts dusky, edged with greenish white: above the eye a pale streak: all the under parts whitish, with longitudinal brown streaks on the sides of the neck, breast, and flanks: the two middle tail-feathers cinereous brown; the others dusky; the outer one being white in its greater portion, the next tipped with white: bill dusky: irides hazel: feet brown: hind claw curved, about equal to the toe in length. (*Young.*) The olivaceous tinge of the upper parts more conspicuous: under parts yellowish, spotted as in the adult bird: wing-coverts more broadly edged with pale cinereous brown: the white on the outer tail-feathers less pure, inclining to pale dusky. (*Egg.*) Greenish white ground, speckled with ash-brown, darkest at the larger end: long. diam. nine lines and a half; trans. diam. seven lines and a half.

Common on rocky shores, in many parts of the kingdom, where it resides the whole year, but is never found inland. Feeds on marine insects. Song and habits somewhat similar to those of the two last species. Breeds in the clefts of rocks, sometimes at a considerable height from the ground. Nest composed of marine plants, with the addition of a little moss externally, lined with fine grass and a few long hairs. Eggs four or five; hatched early in the Spring.

GEN. 23. SAXICOLA, *Bechst.*

70. S. *Ænanthe*, *Bechst.* (*Wheat-Ear.*)—Upper parts ash-gray: forehead, throat, streak above the eyes, and basal portion of the tail, white.

S. Ænanthe, *Temm. Mon. d'Orn.* tom. i. p. 237. *Wheat-Ear*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 181. pl. 48. f. 1. *Bew. Brit. Birds*, vol. i. p. 264.

DIMENS. Entire length six inches six lines: length of the bill (from the forehead) six lines, (from the gape) ten lines; of the tarsus eleven lines and a half; of the tail two inches six lines; from the carpus to the end of the wing four inches: breadth, wings extended, twelve inches.

DESCRIPT. (*Male in summer.*) Upper parts of the body bluish ash: from the corners of the bill a black band, passing across the eyes, and extending to the ear-coverts; above the black, a white streak: forehead and throat white: wings brownish black: fore part of the neck, and upper part of the breast, reddish yellow; the rest of the under parts white: tail, with the exception of the two middle feathers which are wholly black, white for two thirds of its length from the base; the remaining portion black: bill and feet black. (*Female.*) Upper parts cinereous brown, tinged with yellow: the band across the eyes brown: quills with

pale edges: less white on the tail. (*Male, after the autumnal moult.*) All those parts above, which in summer are bluish gray, tinged with rufous: rump, and streak above the eye, pure white: lore pure black; but the black on the ear-coverts tinged with rufous: wing-coverts, and secondary quills, broadly edged with rufous; primaries tipped with reddish white: chin white; throat, breast, and sides of the neck, decided rufous; rest of the under parts paler, but still tinged with rufous. The *female*, at this season, resembles the male, but is more rufous, with the colours not so well-defined: the black streak through the eye indistinct; the white one above it dull and inclining to rufous. (*Young of the year.*) General plumage closely resembling that of the female in autumn: the band across the eyes very indistinct. (*Egg.*) Uniform delicate pale grey blue: long, diam. ten lines and a half; trans. diam. seven lines and a half.

A migratory species, making its first appearance about the end of March, and remaining till September. Common on open downs, and in uninclosed districts. Builds on the ground, under stones, or in old rabbit-burrows. Nest composed of moss and grass, and lined with wool or hair. *Obs.* Previously to quitting this country, it assumes the autumnal plumage, in which state it does not appear to have been noticed by Ornithologists.

71. *S. Rubetra*, Bechst. (*Whin-Chat.*)—Upper plumage dusky brown, edged with reddish yellow: streak above the eyes, a spot on the wings, and base of the tail, white.

S. Rubetra, *Temm. Man. d'Orn.* tom. 1. p. 244. *Whin-Chat*, *Mont. Orn. Dict. Selb. Illust.* vol. 1. p. 183. pl. 48. f. 2. *Bew. Brit. Birds*, vol. 1. p. 266.

DIMENS. Entire length five inches: length of the bill (from the forehead) five lines, (from the gape) eight lines; of the tarsus ten lines; of the tail two inches; from the carpus to the end of the wing two inches ten lines and a half; breadth, wings extended, eight inches eleven lines.

DESCRIPT. (*Male.*) Crown of the head, nape, back, and scapulars, dusky brown, the feathers broadly edged with reddish yellow: from the base of the upper mandible a white streak, passing above the eye, and extending considerably beyond it: cheeks and ear-coverts blackish brown: throat and breast pale yellowish red, edged laterally with white; rest of the under parts white, with a faint tinge of yellow: wings brown, with a large oblong white patch near the shoulder, and another smaller one of the same colour on the greater coverts of the primaries: rump and upper tail-coverts yellowish brown streaked with dusky: tail short; the basal half white, the remainder dusky brown: the two middle feathers entirely dusky: bill and feet black. (*Female.*) The rufous edging of the upper plumage broader and more conspicuous: less white on the wing: all the under parts yellowish white, inclining to rust-red on the breast: upper tail-coverts rust-red. (*Egg.*) Uniform bluish green, with specks (some very minute) of dull reddish brown: long, diam. nine lines; trans. diam. six lines and a half.

Migratory like the last species: seldom appears before the middle of April. Haunts moors and commons. Nest placed on the ground, artfully concealed, constructed almost wholly of dried grasses and stalks, the coarser parts being on the outside, and the finer blades forming a lining within. Eggs five or six in number; hatched towards the end of May.

72. *S. Rubicola*, Bechst. (*Stone-Chat*).—Head and throat black; back black, the feathers with reddish edges: sides of the neck, a spot on the wings, and rump, white.

S. Rubicola, *Temm. Man. d'Orn.* tom. i. p. 246. *Stone-Chat*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 185. pl. 48. f. 3 & 4. *Bew. Brit. Birds*, vol. i. p. 268.

DIMENS. Entire length five inches one line: length of the bill (from the forehead) five lines, (from the gape) eight lines; of the tarsus ten lines; of the tail two inches; from the carpus to the end of the wing two inches six lines: breadth, wings extended, eight inches five lines.

DESCRIPT. (*Male in Summer*.) Head and throat black; on each side of the neck a large white spot; back and nape black, some of the feathers on these parts faintly edged with reddish; quills dusky with pale edges; a large white spot on the greater coverts next the body: breast rust-red; the rest of the under parts white, tinged with yellow: rump white: tail black: bill and feet black. In the *winter plumage*, the black on the head and throat is mixed with cinereous brown. (*Female*.) Head and upper parts dusky brown; the feathers edged with yellowish red: quills and tail deep brown, with paler edges: throat dusky, with whitish and reddish spots: the white on the sides of the neck and on the wings not so conspicuous as in the male; the red on the breast paler. (*Young of the year before the first moult*.) Crown of the head dusky brown, with fine yellowish white streaks on the shafts of the feathers; rest of the upper parts as in the adult female, but with a few whitish spots on the nape and top of the back: wing-coverts and scapulars broadly edged with reddish yellow; upper tail-coverts pale rust-colour: throat whitish, speckled with dusky brown. (*Egg*.) Pale gray blue; the larger end speckled with dull reddish brown: long. diam. eight lines and a half; trans. diam. seven lines.

Found in the same places with the preceding species, but remains throughout the year. Habits similar. Nest on the ground; composed of moss and dry bents, and lined with hair or feathers. Eggs five or six; hatched early in the season.

GEN. 24. PARUS, Linn.

(1. PARUS, *Leach*.)

73. *P. major*, Linn. (*Great Titmouse*).—Head, throat, lower part of the neck, and mesial line of the abdomen, black; cheeks, and a spot on the nape, white.

P. major, *Temm. Man. d'Orn.* tom. i. p. 287. *Great Titmouse*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 233. pl. 51. f. 1. *Bew. Brit. Birds*, vol. i. p. 272.

DIMENS. Entire length five inches eleven lines: length of the bill (from the forehead) five lines and a half, (from the gape) seven lines; of the tarsus nine lines and a half; of the tail two inches nine lines; from the carpus to the end of the wing three inches one line; breadth, wings extended, nine inches three lines.

DESCRIPT. Head, throat, fore part of the neck, and a longitudinal streak down the middle of the abdomen, black: cheeks and ear-coverts, and a spot on the nape, white: back olive-green: rump bluish ash:

breast and belly, on each side of the black line, yellow: quills dusky; wing-coverts bluish ash, the larger tipped with white: tail dusky ash; the outer feather with the exterior web white; the next tipped with white: under tail-coverts white: bill black: legs lead-colour. In the *female*, the colours are not so vivid, and the abdominal black streak does not extend so low down. (*Egg*.) White, spotted and speckled with pale red: long. diam. nine lines and a half; trans. diam. seven lines.

Common in woods and gardens throughout the country. Has a great variety of notes, which are heard very early in the year. Feeds principally on insects, but will also devour grain, as well as flesh. Breeds in the hollows of decayed trees, or in old walls. Nest constructed chiefly of moss, and lined with feathers. Eggs very numerous.

74. *P. cæruleus*, Linn. (*Blue Titmouse*.)—Crown blue, encircled with white: cheeks white, bordered with black, and dark blue.

P. cæruleus, *Temm. Man. d'Orn.* tom. i. p. 289. *Blue Titmouse*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 235. pl. 51. f. 2. *Bew. Brit. Birds*, vol. i. p. 276.

DIMENS. Entire length four inches four lines: length of the bill four lines; of the tarsus eight lines; of the tail one inch eleven lines; from the carpus to the end of the wing two inches four lines; breadth, wings extended, seven inches eight lines.

DESCRIPT. Forehead white, the white extending backwards in the form of a narrow band above the eyes, and reaching to the occiput; crown of the head bright blue: cheeks white, bounded above by a narrow black line commencing at the bill and passing through the eyes, beneath by another line of dark blue extending from the chin to the hinder part of the neck where it becomes broader: back yellowish green: throat, and longitudinal streak on the middle of the abdomen, deep blue; breast and sides bright yellow: wings and tail blue; secondaries and greater coverts tipped with white: bill and legs bluish gray. In the *female*, the blue colours are more obscure, and the abdominal streak not so well defined. (*Egg*.) White, spotted with pale red: long. diam. seven lines and a half; trans. diam. six lines.

Equally common with the last species. Habits and food similar. Nest placed in the holes of trees; formed of moss, and lined with hair and feathers.

75. *P. cristatus*, Linn. (*Crested Titmouse*.)—Crown-feathers elongated, forming a crest, black, edged with white: cheeks and sides of the neck dull white: throat, a streak on the temples, and collar, black.

P. cristatus, *Temm. Man. d'Orn.* tom. i. p. 290. *Crested Titmouse*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 243. pl. 43. f. 5. *Bew. Brit. Birds*, vol. i. p. 274.

DIMENS. Entire length four inches eight lines: length of the bill (from the forehead) three lines and a half, (from the gape) five lines; of the tarsus nine lines; of the tail one inch eight lines; from the carpus to the end of the wing two inches six lines.

DESCRIPT. Feathers on the crown and forehead long and pointed, and capable of erection; of a black colour, edged with white: cheeks and sides of the neck dull white, speckled with black; beneath the eye, and across the ear-coverts, a black streak; throat, upper part of the neck, and a narrow line bounding the white on the sides, deep black: back, and other upper parts, greenish brown: breast and belly white, tinged with buff: bill black: feet bluish gray. The *female* has less black on the throat, and the crown-feathers not so much elongated. (*Egg.*) White, with a few spots and specks of pale red: long. diam. seven lines; trans. diam. six lines.

Confined to Scotland; but stated to be not uncommon amongst the large tracts of pines in the northern parts of that country, particularly in the forest of Glenmoor. Is said to be of solitary habits, and to build in the holes of trees.

76. *P. palustris*, Linn. (*Marsh Titmouse*).—Crown, throat, and nape, deep black: cheeks yellowish white.

P. palustris, *Temm. Man. d'Orn.* tom. i. p. 291. *Marsh Titmouse*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 237. pl. 51. f. 4. *Bew. Brit. Birds*, vol. i. p. 282.

DIMENS. Entire length four inches six lines; length of the bill (from the forehead) four lines, (from the gape) four lines and a half; of the tarsus eight lines; of the tail two inches two lines: from the carpus to the end of the wing two inches four lines and a half: breadth, wings extended, seven inches ten lines.

DESCRIPT. Crown of the head, nape, and throat, deep black; cheeks yellowish white: upper parts of the body gray tinged with brown: wings and tail bluish gray, the edges of the feathers somewhat paler: breast and belly white, faintly tinged with grayish brown: bill black, stronger and more conical than in the next species: feet bluish gray. (*Egg.*) White, with a few dark red spots: long. diam. seven lines and a half; trans. diam. six lines.

Not an uncommon species in woods and thickets. Has a loud shrill note, repeated quickly several times in succession, heard very early in the year. Food insects and seeds, occasionally flesh. Builds in the holes of decayed trees: nest composed of moss and the seed-down of the willow, and lined with the last mentioned material. Eggs from six to eight in number.

77. *P. ater*, Linn. (*Cole Titmouse*).—Head, throat, and fore part of the neck, black; cheeks, and a spot on the nape, white: two transverse white bars on the wings.

P. ater, *Temm. Man. d'Orn.* tom. i. p. 288. *Cole Titmouse*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 239. pl. 51. f. 3. *Bew. Brit. Birds*, vol. i. p. 278.

DIMENS. Entire length four inches three lines: length of the bill (from the forehead) three lines and a half, (from the gape) four lines and a half; of the tarsus seven lines and three quarters: of the tail one inch seven lines and a half; from the carpus to the end of the wing two inches three lines.

DESCRIPT. Crown of the head, and nape of the neck, black; the latter with a central white spot: cheeks and sides of the neck white:

throat and fore part of the neck black: upper parts of the body cinereous, inclining to yellowish on the rump: greater and lesser coverts tipped with white, forming two transverse bars across the wings when closed: belly and abdomen grayish white: tail relatively shorter than in the last species, and slightly forked. The *female* has less white on the sides of the neck, and less black on the throat. (*Egg.*) White, with numerous pale red spots: long. diam. seven lines and a half; trans. diam. six lines.

Not so abundant as the last species. Confined to woods and extensive plantations. Feeds principally on insects. Builds in holes. Nest constructed of moss and wool, and lined with hair.

(2. MECISTURA, *Leach.*)

78. *P. caudatus*, Linn. (*Long-tailed Titmouse.*)—Forehead, crown, cheeks, and throat, white: a broad streak across the eye, nape, and back, black; scapulars reddish.

P. caudatus, *Temm. Man. d'Orn.* tom. 1. p. 296. Long-tailed Titmouse, *Mont. Orn. Dict. Selb. Illust.* vol. 1. p. 241. pl. 51. f. 5. *Bew. Brit. Birds*, vol. 1. p. 279.

DIMENS. Entire length five inches nine lines: length of the bill (from the forehead) two lines and a half, (from the gape) three lines and a half; of the tarsus eight lines; of the tail three inches six lines; from the carpus to the end of the wing two inches four lines: breadth, wings extended, six inches eight lines.

DESCRIPT. Forehead, crown, cheeks and throat, pure white; over each eye a broad black streak, which extends backward to the nape, where it unites with its fellow to form a broader band of the same colour down the middle of the back: sides of the back, and scapulars, rose-red: quills black; greater coverts dusky ash; these last, as well as the secondary quills, edged with white: belly, sides, and abdomen, whitish, tinged with red: tail cuneated, and very long; the six middle feathers black; the others tipped, and obliquely edged on their outer webs, with white: bill black, very short: irides brown: feet dusky. *Obs.* According to Temminck, it is only the *female* which possesses the black streak over the eyes, but as far as my observation goes it is common to both sexes, nor did I ever see a specimen in which it was wanting. *Young* birds have the white parts more or less mottled with brown and dusky; and the black on the back not so deep and well-defined. (*Egg.*) White, with a few pale red specks; frequently quite plain: long. diam. seven lines; trans. diam. five lines.

Far from uncommon in woods and shrubberies, in most parts of the kingdom. Feeds entirely on insects. Nest placed generally in thick bushes; of a very elegant form; long and oval, covered at top, with a small aperture at the side; constructed principally of moss and wool, studded externally with lichens, and lined with a profusion of soft feathers. Eggs ten to twelve in number. The young, when fledged, follow the parent birds till the ensuing spring.

GEN. 25. CALAMOPHILUS, Leach.

79. *C. biarmicus*, Leach. (*Bearded Titmouse*.)—Crown and occiput bluish ash; throat white; beneath the eye a black pointed moustache.

Parus biarmicus, *Temm. Man. d'Orn.* tom. i. p. 298. *Bearded Titmouse*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 244. pl. 51. f. 6. *Bew. Brit. Birds*, vol. i. p. 283.

DIMENS. Entire length six inches four lines: length of the bill (from the forehead) three lines and a half, (from the gape) five lines; of the tarsus nine lines and a half; of the tail three inches one line; from the carpus to the end of the wing two inches three lines.

DESCRIPT. (*Male*.) Head and occiput bluish gray: between the bill and the eye a tuft of loose pendent black feathers, forming a pointed moustache on each side of the face; throat and fore part of the neck white; breast and belly tinged with rose-red: nape, back, and rump, light rufous orange; scapulars paler: quills dusky, the primaries edged with white; greater coverts deep black, edged with red on the outer webs, and with reddish white on the inner: under tail-coverts deep black: tail long and cuneated, orange-brown; the outer feathers tipped, and edged externally, with grayish white: bill orange: irides yellow: feet black. (*Female*.) No black moustache on the sides of the neck; crown, and upper parts of the body, yellowish brown, spotted with dusky: under tail-coverts of the same colour with the abdomen. (*Egg*.) White, with a few minute specks and streaks of dark red: long. diam. eight lines and a half; trans. diam. six lines and a half.

A local species confined to fenny districts, and extensive marshes, where there is abundance of reeds. Found near Winchelsea in Sussex, between Erith and London, and also in some parts of Gloucestershire, Lancashire, Cambridgeshire, and Norfolk. Nest placed near the ground, fixed to the stems of coarse grasses; composed entirely of dried bents, the finer ones forming the lining. Eggs four to six in number. Said to feed on land mollusca, especially the *Succinea amphibia*, as well as insects.

GEN. 26. BOMBYCILLA, Briss.

80. *B. garrula*, Bonap. (*Bohemian Wax-Wing*.)—Head and upper part of the body purplish red; the feathers on the crown elongated; throat black: secondaries and greater coverts tipped with white.

Bombycivora garrula, *Temm. Man. d'Orn.* tom. i. p. 124. *Bohemian Wax-Wing*, *Selb. Illust.* vol. i. p. 268. pl. 34*. *Bohemian Chatterer*, *Mont. Orn. Dict.* *Chatterer*, *Bew. Brit. Birds*, vol. i. p. 98.

DIMENS. Entire length seven inches eleven lines.

DESCRIPT. Crown-feathers elongated, forming a pendent crest, of a pale brown purplish red colour: nostrils covered with small black feathers: throat and region of the eyes deep black: all the upper and under parts of the body purplish red, with a tinge of ash-gray, being darkest above: greater wing-coverts black, tipped with white: quills black; some of the

primaries with a *yellow*, others, and all the secondaries, with a *white* spot at the extremity of the outer web: six or eight of the secondaries with the shafts terminating in a small, oblong, flat, cartilaginous process, of a bright red colour, and having the appearance of wax: vent and under tail-coverts reddish orange: tail black, tipped with yellow: bill and feet black: irides purplish red. In the *female*, there is less black on the throat, and the wax-like appendages are not so numerous: the yellow on the wings, and extremity of the tail, is also not so bright. In *young* birds the waxen tags do not appear at all till after the first moult; from that time they gradually increase in number as the age of the individual advances. (*Egg.*) Pale blue, with a few specks of ash-gray and black: long. diam. eleven lines; trans. diam. seven lines.

A very uncommon visitant in this country; making its appearance at irregular intervals. Occasionally observed in small flocks during the winter season. Feeds on heps, and the berries of the mountain-ash.

GEN. 27. ALAUDA, Linn.

81. *A. alpestris*, Linn. (*Shore Lark.*)—Upper parts reddish brown; streak above the eye, moustache, and a broad transverse bar on the breast, black.

A. alpestris, Temm. *Man. d'Orn.* tom. i. p. 279. *Shore Lark*, Wils. *Amer. Orn.* vol. i. p. 85. pl. 5. f. 4. *Penn. Arct. Zool.* vol. n. p. 84.

DIMENS. Entire length six inches ten lines. TEMM.

DESCRIPT. (*Male.*) Forehead, throat, and sides of the head behind the eyes, pale yellow: a streak above the eye, continued over the forehead and bounding the yellow on that part, a moustache from the corners of the bill across the cheeks, and a broad transverse bar on the upper part of the breast, deep black: upper parts of the body, wing-coverts, and sides of the breast, reddish brown: quills dusky, edged internally with white: tail with the two middle feathers brown; the rest black; the outer one edged externally with white: lower part of the breast and flanks pale red; belly and abdomen white: bill and feet black. (*Female.*) Forehead yellowish; crown of the head varied with black and brown; the black parts with fine yellowish streaks: the transverse bar on the breast smaller than in the other sex; the black feathers in the tail tipped with whitish. (*Egg.*) White, spotted with black.

A single specimen of this Lark, the only one that has hitherto occurred in this country, was killed on the beach near Sherringham in Norfolk, in March 1830. The species inhabits the northern parts of Europe, Asia, and America. Said to frequent extensive plains, and to feed on insects and seeds. Nest placed on the ground: eggs four or five in number.

(5.) *A. rubra*, Gmel. *Syst.* vol. n. p. 794. *Red Lark*, Lewin, *Brit. Birds*, vol. iii. pl. 93. *Mont. Orn. Diet. and Supp. App.*

An obscure species, if indeed it be a species, of which very little is known. Montagu says, "Taken in the Winter of 1812, near Woolwich, in a net with other larks." The specimen, however, is not in his collection in the British Museum, nor does one exist in any other with which I am acquainted.

82. *A. arvensis*, Linn. (*Sky-Lark*.) — Upper parts reddish brown; neck and breast reddish white with dusky spots.

A. arvensis, *Temm. Man. d'Orn.* tom. i. p. 281. *Sky-Lark*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 273. pl. 50. f. 1. *Bew. Brit. Birds*, vol. i. p. 213.

DIMENS. Entire length seven inches three lines: length of the bill (from the forehead) five lines and a half, (from the gape) eight lines; of the tarsus eleven lines and a half; of the hind toe, claw included, one inch one line and a half; of the tail two inches eight lines; from the carpus to the end of the wing four inches four lines: breadth, wings extended, thirteen inches six lines.

DESCRIPT. All the upper parts reddish brown, with a dusky spot in the middle of each feather; these colours palest on the nape, darkest on the head and upper part of the back: a whitish streak above the eyes; cheeks pale yellowish brown: throat yellowish white; fore part of the neck, and breast, tinged with reddish yellow, and spotted with blackish brown; sides with dark streaks of this last colour; middle of the abdomen white: quills edged externally with reddish; the secondaries notched at their extremities: outer tail-feathers with the tip and exterior web white; the two next simply edged with white: bill dusky; the base of the lower mandible yellowish: feet yellowish: hind claw nearly straight, longer than the toe. (*Egg*.) Greenish white ground, spotted all over with darker green and ash-brown: long. diam. eleven lines; trans. diam. eight lines and a half.

An abundant, and widely dispersed species. Frequents open country, more especially arable and cultivated lands. Sings in its ascent, and also whilst suspended at a height in the air. Commences its song, in the middle of Summer, as early as two A.M. Breeds in May. Nest placed on the ground; composed of stalks and dry grasses, and lined with the finer blades. Eggs four or five in number. Occasionally two broods in the season. Congregates in large flocks during the Winter.

83. *A. arborea*, Linn. (*Wood-Lark*.) — Upper parts reddish brown: neck and breast yellowish white, with dusky spots: a distinct yellowish white band above the eyes, reaching from the bill to the occiput.

A. arborea, *Temm. Man. d'Orn.* tom. i. p. 282. *Wood-Lark*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 276. pl. 50. f. 2. *Bew. Brit. Birds*, vol. i. p. 222.

DIMENS. Entire length six inches five lines: breadth, wings extended, twelve inches six lines.

DESCRIPT. Distinguished from the last species, by its somewhat smaller size, shorter tail, and very distinct yellowish white streak passing from the bill above the eyes and reaching nearly to the occiput. Upper plumage light reddish brown, with dusky spots: crown-feathers long, and capable of erection: ear-coverts pale brown: cheeks yellowish white: neck and breast yellowish white, with longitudinal dusky spots; belly dirty white: quills dusky, slightly edged with brown; wing-coverts tipped with white: the two middle tail-feathers brown: the next two

dusky; the others black, with white tips; the outermost of all white on the exterior web: bill brownish black; the base of the lower mandible yellowish white: feet flesh-colour, tinged with yellowish brown; hind claw very long, and nearly straight. (*Egg.*) Light greenish brown ground, mottled with darker brown, with a few dusky streaks at the larger end: long. diam. ten lines; trans. diam. seven lines.

Much less generally distributed than the last species: not uncommon in the southern and western parts of England, but rare in the eastern. Habits similar to those of the Sky-Lark. Sings at a great height in the air, flying round and round in large irregular circles. In song nearly the whole year. Nest on the ground, constructed principally of dry grasses, and lined with a few hairs. Breeds very early in the season. Does not congregate in Winter.

GEN. 28. EMBERIZA, *Linn.*

(I. PLECTROPHANES, *Meyer.*)

84. *E. Lapponica*, Nilss. (*Lapland Bunting.*)—Crown of the head, and upper part of the breast, black; above the eye a white band, prolonged on the sides of the neck: nape, back, and scapulars, varied with brown and red.

E. calcarata, *Temm. Man. d'Orn.* tom. i. p. 322. *Plectrophanes Lapponica*, *Selb. in Linn. Trans.* vol. xv. p. 156. pl. 1. (Young.) *Faun. Bor. Amer.* part 2. p. 248. pl. 48. (Adult.) Lapland Finch, *Lath. Syn.* vol. ii. p. 263. Lapland Lark-Bunting, *Selb. Illust.* vol. i. p. 283. pl. 100. f. 6.

DIMENS. Entire length six inches eight lines: length of the bill (from the forehead) six lines, (from the gape) six lines and a quarter; of the tarsus ten lines; of the hind toe, claw included, nine lines and a quarter; of the tail two inches nine lines; of the folded wing three inches nine lines. *Faun. Bor. Amer.*

DESCRIPT. (*Young.*) "Bill yellowish brown, palest towards the base of the under mandible: head, and all the upper parts of the body, pale wood-brown, tinged with yellowish gray; the shafts of the feathers blackish brown: greater wing-coverts, and secondary quills, blackish brown, deeply margined with chestnut or orange-brown; the tips white: quills dusky brown, paler at the edge: above the eyes a broad streak of pale wood-brown: cheeks and ear-coverts wood-brown, the latter mixed with black: from the corners of the under mandible, on each side of the throat, a streak of blackish brown: throat yellowish white: lower part of the neck and breast dirty white, with numerous dusky spots: belly and vent white: flanks with oblong dusky streaks: tail dusky, the outer feather with the exterior web, and half of the interior, dirty white; the second with a small wedge-shaped white spot near the tip: legs and toes brown: claws not much hooked, the posterior nearly straight, and longer than the toe." **SELBY.** In the *adult male*, which has not yet occurred in this country, the head, chin, throat, and upper part of the breast, are velvet black, with a broad whitish band down the sides of the neck, commencing above the eyes and dilating into an open space in about the middle of its course: nape bright chestnut; rest of the upper plumage pale reddish brown, with a blackish streak in the middle of each feather: wing-coverts with two obsolete white bands; primaries hair-brown, their

exterior edges whitish: belly and under tail-coverts dusky white; sides of the breast, and flanks, spotted with black: bill bright lemon-yellow, tipped with black: legs pitch-black. The *female* differs in having the chin grayish, the black plumage of the head and breast edged with pale brown and gray, and the chestnut feathers of the nape fringed with white: the white band duller. (*Egg.*) Pale ochre-yellow, spotted with brown. *Faun. Bor. Amer.*

Of this species, which is a native of high northern latitudes, only four individuals have hitherto occurred in this country: the first of these was taken in Cambridgeshire; the second in the neighbourhood of Brighton; the third near London, in Sept. 1828; the fourth near Preston in Lancashire, in Oct. 1833. All the specimens were in immature plumage. Habits said to resemble those of the Larks. Food principally seeds. Nest composed externally of dry stems of grass, and lined with hair. Eggs usually seven in number.

85. *E. nivalis*, Linn. (*Snow-Bunting.*)—Head, neck, a large space on the wings, and all the under parts, white.

E. nivalis, *Temm. Man. d'Orn.* tom. i. p. 319. *Snow-Bunting*, *Mont. Orn. Dict. and Supp.* *Selb. Illust.* vol. i. p. 279. pl. 52. f. 5. *Bew. Brit. Birds*, vol. i. p. 178.

DIMENS. Entire length six inches eight lines: length of the bill (from the forehead) four lines and a half, (from the gape) five lines and a half; of the tarsus ten lines; of the hind toe, claw included, eight lines; of the tail two inches eight lines; from the carpus to the end of the wing four inches four lines and a half: breadth, wings extended, twelve inches four lines.

DESCRIPT. (*Male in Summer.*) Head, neck, rump, wing-coverts, upper half of the quills, and all the under parts, pure white: back, scapulars, bastard wing, the lower portion of the quills, and the three secondaries nearest the body, black: the three outer tail-feathers white, with a black spot near the tip; the next with the upper portion of the outer web white; the rest black: bill yellow, dusky at the tip: irides deep brown: feet black. (*Female, and male in Winter.*) Head and neck white, but deeply tinged on the occiput, nape, and ear-coverts, with brownish red; feathers on the back and scapulars, and three secondaries nearest the body, black, all broadly edged and tipped with reddish ash: greater wing-coverts white, stained with dusky towards the tips of the feathers: breast, rump, and upper tail-coverts, tinged more or less deeply with brownish red: the middle tail-feathers edged, chiefly towards their extremities, with reddish white: belly and abdomen white. (*Young of the year.*) Crown of the head yellowish brown: forehead, ear-coverts, and a transverse band on the breast, deep reddish brown; flanks the same, but paler; nape, throat, and fore part of the neck, cinereous, with a tinge of reddish; the feathers on the back, rump, and scapulars, dusky, with a broad edging of yellowish red; middle of the wing, and under parts, whitish; quills and middle tail-feathers dusky, edged with reddish; the three outer tail-feathers whitish, with a large black spot towards the tip: bill yellowish: feet black. (*Egg.*) Pale reddish white, slightly speckled and spotted with pale red and purple brown: long. diam. eleven lines; trans. diam. seven lines and a half.

A regular winter visitant in the northern parts of the island, first appearing about the middle of October; but very rare, and only of accidental occurrence, in the southern and midland counties. Breeds

within the arctic circle: nest said to be placed within the fissures of rocks. Habits terrestrial: does not perch on trees. Food seeds and insects. *Obs.* The *Emberiza mustelina* and *E. montana* of authors are this species, either in its winter plumage, or in an immature state.

(2. EMBERIZA, *Meyer.*)

86. *E. Miliaria*, Linn. (*Common Bunting.*)—Upper parts yellowish brown, with darker spots: beneath yellowish white, spotted and streaked with black.

E. Miliaria, *Temm. Man. d'Orn.* tom. i. p. 306. *Common Bunting*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 286. pl. 52. f. 1. *Bew. Brit. Birds*, vol. i. p. 168.

DIMENS. Entire length seven inches six lines: length of the bill (from the forehead) six lines and one-third, (from the gape) seven lines and one-third; of the tarsus one inch one line; of the tail two inches eleven lines; from the carpus to the end of the wing three inches nine lines: breadth, wings extended, twelve inches three lines.

DESCRIPT. Upper parts yellowish brown, inclining to olive, the shafts of the feathers with longitudinal dusky streaks: under parts yellowish white, spotted on the breast, and streaked on the sides, with dark brown: quills and tail-feathers dusky, with pale edges; wing-coverts tipped with yellowish white: upper mandible dusky, under one pale: irides dark hazel: feet pale brown. *Obs.* This species is occasionally found white, or pied. (*Egg.*) Reddish white, or purple red ground, streaked and spotted with dark purple brown: long. diam. one inch; trans. diam. eight lines and a half.

Common throughout the kingdom, particularly in open cultivated lands. Note harsh and inharmonious, heard at intervals during a great part of the year. Food principally seeds. Nest placed on the ground, formed of straw and fibrous roots, and lined with fine grass and hairs. Congregates in the winter season.

87. *E. Schœniclus*, Linn. (*Reed-Bunting.*) — Head, throat, and fore part of the neck, black: back and scapulars reddish brown, with dusky streaks.

E. Schœniculus, *Temm. Man. d'Orn.* tom. i. p. 307. *Reed-Bunting*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 290. pl. 52. f. 3. *Bew. Brit. Birds*, vol. i. p. 176.

DIMENS. Entire length six inches: length of the bill (from the forehead) four lines and a half, (from the gape) five lines and a half; of the tarsus ten lines; of the tail two inches seven lines; from the carpus to the end of the wing three inches one line: breadth, wings extended, nine inches eleven lines.

DESCRIPT. (*Male in Summer.*) Crown of the head, occiput, cheeks, throat, and upper part of the breast, deep black: a white streak from the corner of the bill downwards joining a collar of the same colour encircling the nape and sides of the neck: back, scapulars, and wing-coverts, reddish brown, with a dusky spot in the middle of each feather: rump and upper tail-coverts bluish ash, tinged with brown: sides of the breast, belly, and under tail-coverts, white; flanks with a few longitudinal dusky streaks: quills dusky, edged with reddish: tail black; the two middle

feathers broadly edged with rufous; the two outer ones on each side obliquely marked with white towards their extremities, the shafts and tips black: bill black; feet dusky. In *Winter* the black feathers on the head, throat, and breast, are edged with reddish brown. (*Female*.) Crown of the head reddish, with dusky spots: above the eye a pale streak of yellowish brown: throat whitish, bordered on each side by a black line: breast and flanks tinged with reddish, and spotted with dusky: nape and sides of the neck brownish ash: the rest much as in the male bird. The *young* of both sexes are very similar to the adult female. (*Egg*.) Purple white ground, sparingly streaked with dark purple brown: long. diam. nine lines and a half; trans. diam. seven lines.

Common in marshy districts, and by the sides of rivers, in most parts of the kingdom. Feeds on the seeds of aquatic plants, and occasionally on aquatic insects. Song simple and inharmonious. Nest generally placed on the ground, and concealed amongst rushes; constructed of stalks, and other dry vegetable substances, and lined with fine grass, and a scanty supply of long hairs. Eggs four or five in number; incubation commencing about the first week in May. Occasionally a second brood in July.

88. *E. Citrinella*, Linn. (*Yellow Bunting*.)—Head, neck, and upper part of the breast, yellow: back and scapulars reddish brown, with darker spots.

E. Citrinella, *Temm. Man. d'Orn.* tom. i. p. 304. *Yellow Bunting*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 288. pl. 52. f. 2. *Bew. Brit. Birds*, vol. i. p. 172.

DIMENS. Entire length six inches four lines: length of the bill (from the forehead) five lines, (from the gape) seven lines; of the tarsus nine lines and a half; of the tail two inches eleven lines; from the carpus to the end of the wing three inches three lines: breadth, wings extended, ten inches three lines.

DESCRIPT. (*Male*.) Head, cheeks, fore part of the neck, abdomen, and under tail-coverts, gamboge-yellow: breast and sides streaked with brownish orange: back and scapulars reddish brown, tinged with olive, with a dusky spot in the centre of each feather; rump bright chestnut, the feathers edged with grayish white: tail dusky: the two outer feathers with a large white conical spot on their inner webs: feet yellowish brown. (*Female*.) The yellow colours much less vivid; that on the head, neck, and throat, marked with olivaceous brown spots; the under parts more clouded and streaked than in the male bird. (*Egg*.) Pale purplish white, streaked and speckled with dark red brown: long. diam. ten lines and a half; trans. diam. eight lines.

Abundant, and generally distributed. Feeds principally on grain. Song scarcely more varied than that of the last species; heard from the first week in February to the beginning or middle of August. Nest placed on or near the ground; composed of straw and dried herbage, and lined with fine grass and long hair. Eggs three to five in number. Breeds late; incubation seldom commencing before the beginning or middle of May. Young fledged about the second week in June. Congregates in Winter.

89. *E. Cirlus*, Linn. (*Cirl Bunting*.)—Head and nape olivaceous green, with dusky spots: throat, and a narrow band across the eye, black: a yellow gorget on the neck.

E. Cirlus, *Temm. Man. d'Orn.* tom. i. p. 313. Cirl Bunting, *Mont. Orn. Dict. and Supp. with fig.* *Id. in Linn. Trans.* vol. vii. p. 276. *Selb. Illust.* vol. i. p. 292. pl. 52. f. 4. *Bew. Brit. Birds*, vol. i. p. 174.

DIMENS. Entire length six inches six lines. **MONT.**

DESCRIPT. (*Male in Summer.*) Crown of the head, and nape of the neck, olivaceous green, with a dusky streak in the centre of each feather; throat, and a narrow band through the eye reaching from the bill to the ear-coverts, black; streak above the eye, and another below it, bright primrose-yellow; a gorget of the same colour encircling the lower part of the neck: breast, immediately below the gorget, yellowish gray, inclining to olive green: back and scapulars fine chestnut brown, the former streaked with dusky; rump olivaceous brown: belly and vent bright yellow; sides of the breast and abdomen brownish red: tail with the two middle feathers chestnut brown; the rest black, the two outer ones obliquely marked with white; bill bluish gray: feet brown, with a tinge of flesh red. In *Winter*, the black on the throat is more dull, the feathers edged with pale yellow. (*Female.*) Head and nape olivaceous brown, with numerous spots of a darker colour; above the eye a dull yellow streak, passing down the sides of the head: chin whitish; the rest of the under parts pale yellow; the breast spotted with reddish; the sides, and under tail-coverts, with large dusky streaks: upper parts as in the male bird, but the colours not so bright, and the spots on the back larger. (*Egg.*) Dirty white, streaked and speckled with dark liver-brown: long. diam. ten lines; trans. diam. eight lines.

Not uncommon in Devonshire, and in one or two of the adjoining counties, where it was first observed by Montagu. Is also found in the Isle of Wight, but is scarcely known in other parts of England. Song and habits somewhat similar to those of the last species. Nest placed in a low bush; composed of dry stalks, roots, and a little moss, and lined with long hair and fibrous roots. Eggs four or five; laid early in May. Congregates in the winter season with Chaffinches, and Yellow Buntings. Said to feed on the berries of the *Solanum*.

90. *E. Hortulana*, Linn. (*Ortolan Bunting*).—Head and neck olive-gray, obscurely spotted with dusky brown; throat, orbits, and a narrow streak from the corner of the bill downwards, greenish yellow.

E. Hortulana, *Temm. Man. d'Orn.* tom. i. p. 311. Green-headed Bunting, *Brown, Illust. of Zool.* p. 74. pl. 30. *Lewin, Brit. Birds*, vol. ii. pl. 76. *Bew. Brit. Birds*, vol. i. p. 170. *Ortolan Bunting*, *Selb. Illust.* vol. i. p. 294. pl. 100. f. 7.

DIMENS. Entire length six inches six lines: length of the bill (from the forehead) four lines and three-quarters, (from the gape) five lines and a half; of the tarsus eight lines and a half; of the tail two inches eight lines; from the carpus to the end of the wing three inches five lines.

DESCRIPT. (*Male.*) Head and neck olivaceous green, with a tinge of ash-gray, obscurely spotted with dusky brown; throat, circle round the eyes, and a broadish streak passing downwards from the corner of the bill, greenish yellow: feathers on the back, rump, scapulars, and wing-coverts, deep brown in the middle, reddish at the edges; those on the breast, belly, and abdomen, chestnut red, edged with ash-gray: quills and tail dusky brown; the former with pale edges; the latter with the

two outer feathers white on their inner webs: irides brown: bill and feet flesh red. (*Female*.) Smaller; the yellow on the throat, and round the eyes, paler; breast spotted with brown; the rest of the under parts reddish white: above, as in the male bird, but the colours not so deep, and the spots on the head and nape larger, and more numerous. (*Egg*.) Pale reddish white, streaked and speckled with purple brown.

Only two examples of this species have hitherto occurred in this country: the first taken near London many years ago, and described by Brown, Lewin, and others, under the name of the *Green-headed Bunting**; the second killed near Manchester, in November 1827. Said to build in bushes, hedges, or among corn, and to lay from four to five eggs. Food grain and insects.

- (6.) *E. Ciris*, Linn. Syst. Nat. tom. i. p. 313. *Painted Finch*, Edwards, Nat. Hist. pl. 130. Gleanings, pl. 273. *Painted Bunting*, Lath. Syn. vol. ii. p. 206.

A single individual of this species is recorded by Montagu (*Orn. Dict. Supp. Art. Grosbeak-White-Winged*.) as having been taken alive on Portland Island, in the year 1802; but considerable doubts are entertained, whether it had not escaped from some vessel going up channel. It is a native of some parts both of N. and S. America.

GEN. 29. FRINGILLA, Linn.

(1. FRINGILLA, Cuv.)

91. *F. Cælebs*, Linn. (*Chaffinch*.)—Head and nape grayish blue: back and scapulars chestnut brown, tinged with olive; rump greenish.

F. Cælebs, *Temm. Man. d'Orn.* tom. i. p. 357. *Chaffinch*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 303. pl. 54. f. 4. & 4*. *Bew. Brit. Birds*, vol. i. p. 188.

DIMENS. Entire length five inches eleven lines: length of the bill (from the forehead) five lines, (from the gape) six lines and a half; of the tarsus nine lines; of the tail two inches seven lines; from the carpus to the end of the wing three inches three lines: breadth, wings extended, ten inches.

DESCRIPT. (*Male in Spring*.) Forehead black; crown of the head, and nape, deep grayish blue: back and scapulars chestnut brown, with a tinge of olive-green; rump greenish yellow: cheeks, throat, and neck, reddish brown; breast and flanks the same, but somewhat paler: lower part of the abdomen white: lesser wing-coverts white; those of the primary quills, and the bastard wing, entirely black; secondary coverts black, tipped with primrose-yellow: the three first quills black, edged externally with yellowish white; the rest of the primaries, and all the secondaries, with a white spot at the base, with part of the inner web white, and with a portion of the outer web edged with pale yellow: the two middle tail-feathers ash-gray; the others black; the two outer ones on each side having a large white spot on the inner web: bill bluish; the tip black: feet brown. In *Winter*, the colours are paler; and the feathers on all the upper and under parts tipped with ash-gray. (*Female*.) Head, nape, back, and scapulars, pale olive-green, tinged with cinereous brown: cheeks, and under parts, cinereous white: the transverse bars on the wings not so distinct as in the male bird: bill yellowish

* See some remarks on this subject by Mr Yarrell, in *Zool. Journ.* vol. iii. p. 498.

gray. *Obs.* This species is subject to much variation of plumage, especially about the wings, which are occasionally almost entirely white. (*Egg.*) Pale purplish white, sparingly streaked and spotted with red brown: long. diam. nine lines and a half; trans. diam. seven lines.

Common throughout the country. In song from the first week in February, to the end of June, or beginning of July. Nest placed against the side of a tree, or in the forked branch of a bush; constructed principally of moss, elegantly studded on the outside with lichens and wool, and lined with feathers and hair. Eggs four or five in number; laid about the third week in April. Collects in flocks, at the approach of Winter.

92. *F. Montifringilla*, Linn. (*Mountain Finch*).—Head, cheeks, nape, and upper part of the back, black: rump white: axillæ bright gamboge-yellow.

F. Montifringilla, *Temm. Man. d'Orn.* tom. i. p. 360. *Mountain Finch*, *Selb. Illust.* vol. i. p. 306. pl. 54. f. 5. & 5*. *Bew. Brit. Birds*, vol. i. p. 190. *Brambling*, *Mont. Orn. Dict.*

DIMENS. Entire length six inches five lines: length of the bill (from the forehead) five lines, (from the gape) six lines and a half; of the tarsus ten lines; of the tail two inches seven lines; from the carpus to the end of the wing three inches eight lines: breadth, wings extended, ten inches seven lines.

DESCRIPT. (*Male in Winter.*) Head, cheeks, nape, sides of the neck, and upper part of the back, black; the feathers edged and tipped with reddish brown and ash-gray: throat, fore part of the neck, breast, scapulars, and lesser wing-coverts, pale orange-brown; greater coverts black, tipped with orange-red; quills black, edged with yellow on their outer webs; some of the primaries with a white spot at the base, forming an oblique bar of that colour when the wing is closed: axillæ bright gamboge-yellow: rump, belly, and under tail-coverts, white: sides inclining to reddish, with a few dusky spots: tail black; the two middle feathers edged with reddish ash; the outer one white on the exterior web: bill lemon-yellow at the base, black at the tip: irides dusky: feet grayish brown. In *Summer*, the plumage on the head, neck, and back, is deep black, without the edging of reddish brown: the bill is also bluish. (*Female.*) Colours not so bright: crown of the head reddish brown, the feathers edged with ash-gray; a black streak above the eye; cheeks, nape, and sides of the neck, cinereous; back dusky brown, the feathers broadly margined with reddish ash: the rufous tinge on the breast and wings much fainter than in the male bird. In *young* birds the throat is white. (*Egg.*) "Spotted with yellowish." *TEMM.*

A native of the northern parts of Europe. In this country only a winter visitant, making its appearance towards the end of Autumn, and departing early in the Spring. Is more plentiful, and more generally distributed, some seasons than others, according to the state of the weather. Habits very similar to those of the last species. Said to build in fir and pine-forests, and to construct a nest of moss and wool, lined with hair and feathers: eggs five in number.

(2. PYRGITA, *Cuv.*)

93. *F. domestica*, Linn. (*House Sparrow*).—Crown and occiput bluish ash; cheeks, and sides of the neck, grayish white: a transverse whitish bar on the wing.

F. domestica, *Temm. Man. d'Orn.* tom. i. p. 350. House Sparrow, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 298. pl. 54. f. 2. & 2*. *Bew. Brit. Birds*, vol. i. p. 184.

DIMENS. Entire length six inches two lines: length of the bill (from the forehead) five lines and a half, (from the gape) seven lines; of the tarsus nine lines; of the tail two inches four lines; from the carpus to the end of the wing two inches eleven lines: breadth, wings extended, nine inches four lines.

DESCRIPT. (*Male*.) Crown of the head, and occiput, bluish ash; space between the bill and the eye, throat, and fore part of the neck, deep black; above the eyes, and behind the ears, a band of reddish brown: cheeks, sides of the neck, breast, and abdomen, grayish white: plumage on the back and wings dusky brown, edged with reddish; the latter with one transverse whitish bar: tail brown, edged with yellowish gray: bill bluish black: irides hazel: feet brown. (*Female*.) Head and nape cinereous brown: above and behind the eye a yellowish streak: upper parts plain brown, darkest in the middle of the feathers: under parts grayish white, without any black on the throat and neck: bill much paler than in the male. *Obs.* White, and other varieties of this species are not unfrequent. (*Egg*.) White; spotted and streaked with ash-colour and dusky brown: varies considerably in the number and intensity of the markings: long. diam. ten lines; trans. diam. seven lines.

Plentiful in all parts of the kingdom. Attached to the habitations of man. Food, grain and insects. Nest generally placed under the eaves of buildings, or in the holes of old walls; composed of hay and straw in large quantities and loosely put together, lined principally with feathers. Eggs five or six.

94. *F. montana*, Linn. (*Tree Sparrow*.)—Crown and occiput chestnut-brown; sides of the neck, and a collar on the nape, white: wings with two transverse white bars.

F. montana, *Temm. Man. d'Orn.* tom. i. p. 354. Tree Sparrow, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 300. pl. 55. f. 2. Mountain Sparrow, *Bew. Brit. Birds*, vol. i. p. 187.

DIMENS. Entire length five inches six lines. **MONT.**

DESCRIPT. Crown of the head, and nape of the neck, deep chestnut-brown, tinged with cinereous: space between the eye and the bill, ear-coverts, and throat, black: cheeks, sides of the neck, and an interrupted collar on the nape, white: feathers on the back, and scapulars, dusky in the middle, rufous brown at the edges; wing-coverts chestnut-brown, tipped with white, forming two distinct bars of that colour across the wings: quills black, with reddish edges: tail rufous brown: bill black: feet pale brown. In the *female*, the colours are more obscure; the black on the throat and ear-coverts hardly visible. (*Egg*.) Dirty white, speckled all over with light ash-brown: long. diam. eight lines and a half; trans. diam. six lines.

Much less abundant, and more partially distributed, than the last species. Met with sparingly in Yorkshire, Lancashire, Lincolnshire, and some other of the northern and eastern counties. Does not frequent buildings, but is partial to old trees, in the holes of which it builds. Nest formed of the same materials as that of the House-Sparrow. Number of eggs the same.

(3. COCCOTHAUSTES, *Briss.*)

95. *F. Coccothraustes*, Temm. (*Common Grosbeak*).
—Base of the bill, lore, and throat, black : crown, cheeks, and rump, reddish brown.

F. Coccothraustes, *Temm. Man. d'Orn.* tom. i. p. 344. *Haw Grosbeak*, *Mont. Orn. Dict.* *Haw-Finch*, *Selb. Illust.* vol. i. p. 324. pl. 55. f. 1. *Grosbeak*, *Bew. Brit. Birds*, vol. i. p. 159.

DIMENS. Entire length seven inches : length of the bill (from the forehead) nine lines, (from the gape) ten lines ; of the tarsus ten lines and a half ; of the tail one inch ten lines ; from the carpus to the end of the wing three inches ten lines.

DESCRIPT. Base of the bill, space between the bill and the eye, chin and throat, deep black : a broad collar of ash-gray on the nape : crown of the head, cheeks, rump, and upper tail-coverts, reddish, or chestnut-brown ; back, scapulars, and lesser wing-coverts, the same, but of a deeper tint ; greater coverts grayish white, forming a broad bar of that colour across the wings ; secondary quills, and some of the primaries, glossy black, with an oblong white spot on the middle of their inner webs, the former truncated at their extremities ; the rest of the quills entirely black : breast and belly vinous red ; vent and under tail-coverts white : tail with the two middle feathers like the back ; the others dusky brown with a large oblong white spot at the extremities of their inner webs : bill blue in Summer, whitish in Winter. In the *female*, the colours are paler, but similarly disposed to those of the other sex. (*Young of the year before the first moult.*) Throat yellow ; crown of the head, cheeks, and all the upper parts of the body, yellowish brown : under parts yellowish white ; the breast, belly and flanks, spotted with brown, this last colour occupying the tip of each feather. (*Egg.*) White tinged with blue, spotted and streaked with ash-gray and dark brown : long. diam. eleven lines ; trans. diam. eight lines and a half.

Only an occasional visitant in this country during the winter months. Principally observed in the southern counties. In a few instances, has been known to remain and breed. Feeds on haws, and other stone fruits. Builds in hedges and tall trees : nest composed of twigs, lichens, and vegetable fibres ; lined with feathers or horse-hair, and other soft materials. Eggs three to five in number.

96. *F. Chloris*, Temm. (*Green Grosbeak*).—All the upper and under parts of the body yellowish green.

F. Chloris, *Temm. Man. d'Orn.* tom. i. p. 346. *Green Grosbeak*, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. i. p. 163. *Selb. Illust.* vol. i. p. 326. pl. 54. f. 3.

DIMENS. Entire length six inches : length of the bill (from the forehead) six lines, (from the gape) seven lines ; of the tarsus nine lines ; of the tail two inches five lines ; from the carpus to the end of the wing three inches four lines : breadth, wings extended, ten inches four lines.

DESCRIPT. (*Male.*) All the upper and under parts of the body, scapulars, and lesser wing-coverts, yellowish green ; greater coverts and secondary quills ash-gray ; primaries dusky, edged on the outer web with gamboge-yellow : tail a little forked ; the two middle feathers blackish gray, edged with yellowish gray ; the four outer feathers on each side

with the greater part of the exterior web yellow : bill and feet pale flesh-red : irides dark hazel. (*Female*.) Upper parts cinereous, tinged with green : under parts much paler than in the male bird : the yellow edging on the outer webs of the primaries and tail-feathers not so bright. (*Egg*.) White, tinged with light blue ; the larger end speckled and spotted with purplish gray and dark brown : long. diam. nine lines and a half ; trans. diam. six lines and a half.

Common in all parts of the country, and stationary throughout the year. Feeds on seeds and grain. Has a harsh monotonous note, heard from the end of February to the middle of August or September. Nest generally placed in a thick bush ; composed of slender twigs, bents, and moss, interwoven with wool, and lined with hair and feathers. Eggs four or five. Is a late breeder, and seldom hatches before the middle of May. Collects into flocks in Winter.

(4. CARDUELIS, *Briss.*)

97. *F. Carduelis*, Linn. (*Goldfinch*.)—Base of the bill, occiput, and nape, black : forehead and throat, blood-red.

F. Carduelis, *Temm. Man. d'Orn.* tom. 1. p. 376. *Goldfinch*, *Mont. Orn. Dict. Selb. Illust.* vol. 1. p. 312. pl. 55. f. 8, 9. *Bew. Brit. Birds*, vol. 1. p. 192.

DIMENS. Entire length five inches : length of the bill (from the forehead) six lines, (from the gape) six lines and a half ; of the tarsus seven lines ; of the tail two inches one line ; from the carpus to the end of the wing three inches one line : breadth, wings extended, nine inches three lines.

DESCRIPT. Base of the bill, lore, occiput and nape, deep black : forehead and throat arterial blood-red : cheeks, ear-coverts, fore part of the neck, belly and abdomen, white : back, scapulars, and sides of the breast, deep yellowish brown : lesser wing-coverts black ; greater coverts, and basal half of the quills, bright yellow ; the remaining portion of the quills black, with a white spot at the tips : tail black : the two outer feathers on each side with a large oblong white spot on the middle of the inner web ; the others tipped with white : bill yellowish white ; the tip dusky : feet pale flesh-red. In the *female*, there is less red on the forehead and throat ; the lesser wing-coverts are brown, and the colours in general not so bright as in the male bird. (*Egg*.) Pale bluish white, with a few spots and lines of pale purple and brown : long. diam. eight lines and a half ; trans. diam. six lines.

An abundant and generally diffused species. Feeds on seeds, especially those of the thistle. Song pleasing and varied ; heard from April till towards the end of the Summer. Builds in orchards, gardens, and plantations : nest placed in trees or shrubs, composed of moss and wool with the addition of a few bents or lichens, and lined with wool, hair, and thistle-down. Eggs four or five in number. Is rather a late breeder.

98. *F. Spinus*, Linn. (*Siskin*.)—Crown of the head, and throat, black : nape dusky green : above and behind the eye a broad yellow streak.

F. Spinus, *Temm. Man. d'Orn.* tom. 1. p. 371. *Siskin*, *Mont. Orn. Dict. & Supp. Selb. Illust.* vol. 1. p. 309. pl. 55. f. 6, 7. *Bew. Brit. Birds*, vol. 1. p. 194.

DIMENS. Entire length four inches seven lines: length of the bill (from the forehead) four lines and a half, (from the gape) five lines and a half; of the tarsus six lines and a half; of the tail one inch nine lines; from the carpus to the end of the wing two inches nine lines.

DESCRIPT. (*Male.*) Throat, crown, and occiput, deep black: nape variegated with dusky and greenish yellow: above and rather behind the eye a broad streak of sulphur-yellow: neck, breast, and belly, yellow: back, scapulars, and lesser wing-coverts, yellowish green, inclining to ash-gray, with a longitudinal dusky spot in the centre of each feather: greater coverts black, tipped with yellow: quills dusky with yellow edges: tail dusky, all the feathers except the two middle ones with a large yellow spot near the base extending over both webs: sides and lower part of the abdomen whitish, with longitudinal dusky streaks: bill whitish, the tip black: feet pale brown. (*Female.*) All the upper parts, cheeks, and sides of the neck, cinereous, with longitudinal dusky spots: under parts yellowish white: breast, flanks, and under tail-coverts, with dusky green streaks: the transverse bars on the wings, and the yellow edging of the quills, paler than in the other sex. (*Egg.*) Pale bluish white, spotted with purple brown.

A winter visitant, but not making its appearance regularly. Occasionally observed in large flocks in Cambridgeshire about the commencement of the new year, during its stay in which parts, it appears to feed almost entirely on the seeds of the alder. Is known also in Norfolk, Suffolk, Sussex, Middlesex, and Northumberland. Breeds in the North of Europe, and is said to build in the highest branches of the pine. Lays five eggs.

(5. LINARIA, *Steph.*)

99. *F. Linaria*, Linn. (*Lesser Redpole.*)—Forehead, lore, and throat, black: bill much compressed towards the tip, yellowish: feet brown.

F. Linaria, *Temm. Man. d'Orn.* tom. 1. p. 373. *Lesser Redpole*, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. 1. p. 200. *Selb. Illust.* vol. 1. p. 320. pl. 54. f. 6.

DIMENS. Entire length five inches three lines: length of the bill (from the forehead) three lines, (from the gape) four lines and a quarter; of the tarsus six lines and a half; of the tail two inches one line; from the carpus to the end of the wing two inches eight lines.

DESCRIPT. (*Male in Spring.*) Forehead, space between the bill and the eye, chin and throat, black: crown of the head deep crimson: occiput, nape, and upper part of the back, blackish brown, the feathers edged with pale reddish brown: sides of the throat, fore part of the neck, breast, sides of the abdomen, and rump, earmine-red, the latter tinged with grayish; middle of the belly, vent, and under tail-coverts, white tinged with rose-colour: wings and tail dusky, the feathers edged with pale reddish brown; wing-coverts tipped with pale yellowish brown, so as to show two transverse bars: bill more compressed than in the next species; yellow, the culmen and tip dusky: feet brown. (*Female.*) Less red on the crown of the head; forehead brown, mixed with yellowish white: rump and under parts only slightly tinged with rose-red: throat dusky; breast and middle of the belly whitish; flanks, vent, and under tail-coverts, marked with large longitudinal spots of blackish brown. (*Egg.*) Pale blue, speckled and streaked with pale purplish red, and reddish brown: long. diam. seven lines and a half; trans. diam. five lines and a half.

Var. β. *Linaria borealis*, *Selb. in Newcast. Nat. Hist. Trans.* vol. i. p. 263. Lesser Redpole, (a large variety,) *Selb. Illust.* vol. i. pl. 53^{**}. f. 2. Mealy Redpole, (*Lin. canescens*,) *Gould, Europ. Birds*, part xi. Larger than the more common variety: general plumage paler: rump grayish white: tail rather longer: the two cross-bars on the wings broader and more strongly marked.

Resident in Scotland and the North of England throughout the year. In the southern counties only a winter visitant, appearing at that season in large flocks, and frequently associating with other species. Nest, according to Selby, placed in a bush or low tree; constructed of moss and the stalks of dry grass, intermixed with down from the catkin of the willow, which also forms the lining. Eggs four or five in number. Young seldom fledged before the end of June or beginning of July. Feeds on the catkins of the birch and alder, as well as on the seeds and buds of other trees. The *Var. β.* is considered by some as a distinct species, but this point requires further investigation. It is known to the bird-catchers in the neighbourhood of London, by the name of *Mealy-backed Redpole*, the more common variety being called by them *Stone Redpole*.

100. *F. cannabina*, Linn. (*Common Linnet*.)—Throat yellowish white, with dusky streaks: scapulars and wing-coverts, chestnut-brown: bill thick at the base; bluish gray: feet brown.

F. cannabina, *Temm. Man. d'Orn.* tom. i. p. 364. Common or Brown Linnet, *Selb. Illust.* vol. i. p. 315. pl. 55. f. 3, 4. Greater Redpole, and Linnet, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. i. pp. 198 & 202.

DIMENS. Entire length five inches five lines: length of the bill (from the forehead) five lines, (from the gape) five lines and a half; of the tarsus seven lines and a half; of the tail two inches three lines; from the carpus to the end of the wing three inches one line: breadth, wings extended, nine inches six lines.

DESCRIPT. (*Male in Spring*.) Forehead and breast bright carmine-red: throat and fore part of the neck yellowish white, with longitudinal dusky streaks: crown, nape, and sides of the neck, cinereous: back, scapulars, and wing-coverts, chestnut-brown: middle of the belly and abdomen grayish white; flanks pale reddish brown: quills black, edged with white, so as to exhibit a longitudinal bar of the latter colour when the wing is closed: tail forked; the two middle feathers wholly black, and pointed; the rest black, narrowly edged with white on the outer web, and more broadly so on the inner: bill bluish gray; not so much compressed as in the last species: feet brown. In *young* birds, the red on the breast and forehead is not so intense, or so widely extended. (*Male in Winter*.) Crown of the head with large dusky spots: back and scapulars chestnut-brown, broadly edged with pale yellowish brown: breast cinereous brown, faintly tinged with red, the tips of the feathers yellowish white: flanks with large oblong brown streaks. (*Female*.) Smaller in size. All the upper parts pale cinereous brown, with dusky spots; wing-coverts reddish brown: under parts pale reddish, inclining to white on the belly: breast and flanks with streaks of dusky brown. (*Egg*.) Pale bluish white, speckled with pale purple and red-brown: long. diam. nine lines; trans. diam. six lines and a half.

Common throughout the country. Song commenced in March or April, and continued during the greater part of the Summer. Nest

placed in hedges and low bushes; formed of moss and bents interwoven with wool, and lined with wool and hair. Eggs four or five in number; hatched the beginning of May. Occasionally a second brood. In Winter, collects into large flocks. *Obs.* The *Fringilla Linota* of authors is this species in its winter plumage.

101. *F. Montium*, Gmel. (*Mountain Linnet*).—Throat reddish brown, without spots; greater wing-coverts edged with white; rump purplish red: bill thick and short; yellow: feet black.

F. Montium, *Temm. Man. d'Orn.* tom. 1. p. 368. *Mountain Linnet* or Twite, *Selb. Illust.* vol. 1. p. 318. pl. 55. f. 5. *Bew. Brit. Birds*, vol. 1. p. 204. *Twite*, *Mont. Orn. Dict.*

DIMENS. Entire length five inches three lines: length of the bill (from the forehead) three lines, (from the gape) four lines; of the tarsus seven lines and a half; of the tail two inches three lines; from the carpus to the end of the wing two inches eleven lines: breadth, wings extended, eight inches nine lines.

DESCRIPT. (*Male in Spring*.) Throat, fore part of the neck, and sides of the head, pale reddish brown: the feathers on the crown, nape, and back, deep black in the middle, reddish at the edges: rump fine purplish red: greater wing-coverts edged and tipped with whitish: quills dusky; the primaries edged with pale brown, the secondaries with white, on their outer webs: breast and sides, yellowish brown, with large dusky spots; middle of the belly and vent whitish: tail forked; deep brown, edged with dirty white: bill smaller than in the last species, wax-yellow: irides brown: feet black. In *Winter*, the black on the upper parts assumes more of a brownish tinge. (*Female*.) All the upper parts brown, the feathers edged with pale yellowish red; rump the same as the back, without the tinge of purplish red: bill yellowish, tipped with brown. (*Egg*.) Pale bluish white, speckled with pale purple-red: long. diam. eight lines; trans. diam. six lines and a half.

Not uncommon in the neighbourhood of London, and in many of the eastern counties, during the winter season, flocking with the last species. At the approach of Spring, retires to the northern and mountainous parts of England and Scotland, where it breeds. Nest placed amongst heath, and formed of that plant with the addition of dry grass and wool, lined with this latter material, fibres of roots, and the finer parts of the heath. Eggs four or five in number.

GEN. 30. PYRRHULA, *Briss.*

102. *P. vulgaris*, Temm. (*Common Bullfinch*).—Base of the bill, crown, throat, wings and tail, black: cheeks, and under parts, red.

P. vulgaris, *Temm. Man. d'Orn.* tom. 1. p. 338. *Bullfinch*, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. 1. p. 165. *Selb. Illust.* vol. 1. p. 336. pl. 54. f. 1. & 1*.

DIMENS. Entire length six inches four lines: length of the bill (from the forehead) four lines and a half, (from the gape) five lines; of the tarsus nine lines; of the tail two inches six lines; from the carpus to the

end of the wing three inches two lines: breadth, wings extended, nine inches nine lines.

DESCRIPT. (*Male*.) Crown and occiput, base of the bill, throat, wings, and tail, velvet black tinged with purple: nape and back bluish ash: cheeks, sides and under part of the neck, breast, belly, and flanks, bright tile-red: rump, vent, and under tail-coverts, white: greater wing-coverts tipped with grayish white, shewing a transverse bar across the wing: bill and feet dusky brown. (*Female*.) Upper parts bluish ash, with a tinge of yellowish brown: under parts reddish brown: less white on the rump. (*Egg*.) Pale blue, speckled and streaked with purplish gray and dark purple: long. diam. nine lines and a half: trans. diam. seven lines.

Common in most parts of the country. Frequents woods, gardens, and orchards. Food seeds and the buds of trees. Nest generally placed in thick bushes; composed of dry twigs, and lined with fibrous roots. Eggs four or five; hatched towards the end of May.

103. *P. Enucleator*, Temm. (*Pine Bullfinch*.)—General plumage reddish orange, spotted on the back and scapulars with dusky brown.

P. Enucleator, Temm. *Man. d'Orn.* tom. i. p. 333. Pine Grosbeak, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. i. p. 161. Pine Bullfinch, *Selb. Illust.* vol. i. p. 334. pl. 53*. f. 1, 2.

DIMENS. Entire length seven inches four or five lines. TEMM.

DESCRIPT. (*Adult male*.) Head, throat, neck, and under parts, fine reddish orange, inclining to yellowish orange on the breast and belly: the feathers on the back, scapulars, and rump, deep brown, broadly bordered with yellowish orange: wings and tail black; the former with two transverse white bars: secondaries edged with white: primaries and tail-feathers, with orange. (*Young male*.) Head, neck, throat, breast, and rump, bright crimson; feathers on the back and scapulars broadly bordered with the same colour: the transverse bars on the wings, and the edging of the secondary quills, likewise crimson: flanks, belly, and vent, cinereous, tinged with red. (*Female*.) Head, nape, and rump, yellowish brown, more or less inclining to orange: back and scapulars cinereous brown: under parts ash-gray very faintly tinged with yellowish orange: the transverse bars on the wings grayish white, and not so broad as in the other sex; quills edged with greenish orange. (*Egg*.) White: long. diam. one inch one line; trans. diam. ten lines.

An occasional visitant in the northern districts of Scotland, but very rare. According to Messrs. C. & J. Paget (*Nat. Hist. of Yarm.* p. 6.) a flight was once observed near Yarmouth. Partial to pine-forests. Food seeds and berries. Breeds within the Arctic Circle. Nest placed in trees, not far from the ground, formed of twigs and sticks, and lined with feathers. Eggs four in number.

GEN. 31. LOXIA, *Briss.*

104. *L. curvirostra*, Linn. (*Common Cross-Bill*.)—Bill as long as the middle toe, moderately hooked; the crossing point of the lower mandible passing beyond the ridge of the upper.

L. curvirostra, *Temm. Man. d'Orn.* tom. i. p. 328. Common Cross-Bill, *Selb. Illust.* vol. i. p. 329. pl. 53. Cross-Bill, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. i. p. 153.

DIMENS. Entire length six inches four lines.

DESCRIPT. (*Adult male.*) General colour of the plumage cinereous, deeply tinged with greenish yellow: rump yellow: lower part of the abdomen ash-gray with dusky spots: quills and tail-feathers dusky, with greenish edges; great and middle wing-coverts edged with yellowish white: bill yellowish brown: irides and feet brown. (*Male after the first moult.*) All the upper and under parts brick-red, tinged with yellowish gray: quills and tail-feathers dusky, edged with yellowish green: under tail-coverts white, spotted with dusky. (*Female, and young of the year.*) Brownish gray, more or less tinged with greenish: rump yellow: under parts whitish, with longitudinal dusky streaks. (*Egg.*) Pale bluish white, speckled with red-brown: long. diam. ten lines and a half; trans. diam. eight lines.

An occasional visitant in this country, at irregular intervals. Generally observed during the summer and autumnal months, in larger or smaller flocks. Breeds in the northern parts of Europe during the Winter, or very early in the Spring. According to Sheppard has been known to breed in Suffolk in one or two instances. Nest placed in the forked branches of pines, composed of moss and lichens, and lined with feathers. Eggs four or five in number. Food principally the seeds of the pine and other firs.

105. *L. Pytiopsittacus*, Bechst. (*Parrot Cross-Bill.*) —Bill shorter than the middle toe, very strong and broad at the base, much hooked; the crossing point of the lower mandible not reaching so high as the ridge of the upper.

L. Pytiopsittacus, *Temm. Man. d'Orn.* tom. i. p. 325. Parrot Cross-Bill, *Selb. Illust.* vol. i. p. 332. pl. 53*. f. 1. *Bew. Brit. Birds*, vol. i. p. 157.

DIMENS. Entire length seven inches. **TEMM.**

DESCRIPT. (*Adult male.*) All the upper and under parts of the body olive-ash: cheeks, throat, and sides of the neck, cinereous: head spotted with brown: rump greenish yellow; breast and belly the same, tinged with gray; sides streaked with dusky: quills and tail-feathers dusky brown, edged with olive-ash: bill dusky horn-colour: irides, and feet, brown. (*Male after the first moult.*) Upper and under parts bright red: wings and tail blackish brown; the quills edged with reddish. (*Female.*) "Upper parts greenish ash, with large spots of cinereous brown: throat and neck grayish, tinged with brown; rest of the under parts cinereous, slightly tinged with greenish yellow: rump yellowish: abdomen whitish; under tail-coverts spotted with brown." **TEMM.** The *young of the year* very much resemble the female. (*Egg.*) Similar to that of the last species.

This species, which inhabits the northern parts of Europe and America, has been occasionally met with in this country, but is much more rare than the last. Habits similar. Breeds, according to Temminck, in some climates, during the Winter; in others, in May. Eggs four or five in number.

106. *L. leucoptera*, Gmel. (*White-winged Cross-Bill*.)
—Bill longer than the middle toe: wings with two broad transverse white bars.

L. leucoptera, Bonap. *Amer. Orn.* vol. 11. pl. 15. f. 3. (Female.) *Faun. Bor. Am.* vol. 11. p. 263. *L. falcistrostra*, Lath. *Ind. Orn.* p. 371. White-winged Cross-bill, Lath. *Syn.* vol. 11. p. 108. *Wilson, Amer. Orn.* vol. 1v. pl. 31. f. 3.

DIMENS. Entire length six inches three lines: length of the bill (from the forehead) seven lines, (from the gape) seven lines and a half; of the tarsus seven lines and a half; of the tail two inches seven lines; of the wing three inches six lines. *Faun. Bor. Am.*

DESCRIPT. (*Immature male*.) Head, nape, back, rump, and all the under parts, crimson-red, partially spotted with dusky: wings and tail black, the former with two broad, white, transverse bars: region of the bill brown. "As the bird acquires its *mature* plumage, the red parts change to greenish yellow, the rump assuming a purer yellow. The *female and young*, before their first moult, are greenish, with yellowish rumps; their bellies whitish, streaked with blackish brown." *Faun. Bor. Am.* (*Egg*.) White, marked with yellowish spots.

A native of North America, but has been killed in Ireland, in one instance, within two miles of Belfast. Partial to thick forests of white spruce, feeding on the seeds of the cones. Nest generally fixed about half way up a pine-tree, composed of grass, mud, and feathers. Eggs five in number.

GEN. 32. STURNUS, Linn.

107. *S. vulgaris*, Linn. (*Common Starling*.)—Plumage black, with purple and green reflections; the feathers tipped with yellowish white.

S. vulgaris, Temm. *Man. d'Orn.* tom. 1. p. 132. Common Starling, *Selb. Illust.* vol. 1. p. 340. pl. 36. f. 1. Starling, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. 1. p. 105.

DIMENS. Entire length eight inches four lines: length of the bill (from the forehead) one inch, (from the gape) one inch four lines; of the tarsus one inch one line; of the tail two inches seven lines; from the carpus to the end of the wing four inches ten lines: breadth, wings extended, fifteen inches.

DESCRIPT. (*Male*.) General colour of the plumage black, with brilliant purple and golden-green reflections; the feathers on the upper parts tipped with small triangular yellowish white spots: quills and tail-feathers dusky brown, with pale reddish edges: under tail-coverts edged with white: bill yellow: feet flesh-red, inclining to brown. In the *Female*, the spots are more numerous, and diffused over the under as well as the upper parts. In the *young of the year previously to the autumnal moult*, the plumage is of a uniform cinereous brown without any spots; the throat and lower part of the abdomen whitish. In this state it is the *Solitary Thrush* of Montagu. The perfect plumage is probably not attained till the third year. (*Egg*.) Of a uniform delicate pale blue: long. diam. one inch two lines; trans. diam. ten lines.

A plentiful and widely dispersed species. Partial to old trees, church steeples, and ruinous buildings, in the holes of which it breeds. Nest formed of dry grass. Eggs four or five in number. In the Autumn, congregates in immense flocks. Food insects and worms, occasionally grain.

GEN. 33. PASTOR, Temm.

108. *P. roseus*, Temm. (*Rose-coloured Pastor*.)—Head and neck violet-black, the feathers on the crown elongated, forming a crest: back and belly rose-red.

P. roseus, Temm. *Man. d'Orn.* tom. i. p. 136. Rose-coloured Pastor, *Selb. Illust.* vol. i. p. 343. pl. 36. f. 2, 3. Rose-coloured Ouzel, *Mont. Orn. Dict.* Rose-coloured Starling, *Bew. Brit. Birds*, vol. i. p. 110.

DIMENS. Entire length eight inches six lines: breadth, wings extended, fourteen inches six lines.

DESCRIPT. (*Male*.) Head and neck velvet-black, with violet and green reflections; the feathers on the crown very much elongated, forming a pendent crest: back, rump, breast, and belly, rose-red: wings and tail blackish brown, with violet reflections: under tail-coverts, and thighs, black: upper mandible, and tip of the lower one, yellowish red, the remaining portion black: irides deep brown: feet yellowish red. (*Female*.) General colours of the plumage much paler; the red tinged with brown: the feathers on the crown shorter. (*Young of the year*.) "No crest: all the upper parts of the body isabella-brown: wings and tail brown, all the feathers fringed with white and ash-colour: throat and middle of the abdomen pure white; the rest of the under parts cinereous brown: bill yellow at the base, brown at the tip: feet brown." TEMM. (*Egg*.) Colour unknown.

An occasional, but very rare, visitant in this country. Has been observed in Sussex, Suffolk, Norfolk, Cambridgeshire, Oxfordshire, Dorsetshire, Northumberland, Lancashire, and Ireland. Habits similar to those of the Starling. Food insects and their larvæ; also seeds, and occasionally cherries. Said to build in the holes of trees and old walls, and to lay six eggs.

GEN. 34. FREGILUS, Cuv.

109. *Graculus*, Selb. (*Cornish Chough*.)—Black: bill and feet coral-red.

Pyrrhocorax Graculus, Temm. *Man. d'Orn.* tom. i. p. 122. Cornish Chough, *Selb. Illust.* vol. i. p. 365. pl. 33. Chough, *Bew. Brit. Birds*, vol. i. p. 90. Red-legged Crow, *Mont. Orn. Dict.*

DIMENS. Entire length sixteen inches: length of the bill (from the forehead) two inches two lines and a half, (from the gape) two inches two lines and a half; of the tarsus two inches; of the tail five inches seven lines; from the carpus to the end of the wing ten inches eight lines: breadth, wings extended, two feet eight inches.

DESCRIPT. The whole plumage black, with purple and green reflections: bill and feet bright coral-red: claws black. (*Egg*.) Yellowish white, spotted with ash-gray and light brown: long. diam. one inch eight lines; trans. diam. one inch one line.

A local species; principally met with on the rocky coasts of Cornwall, Devonshire, Dorsetshire, and Wales. Found also in some of the Scotch Islands. Never observed inland, but breeds on the cliffs, or in old buildings near the sea. Nest formed of sticks, and lined with wool and hair. Eggs four or five in number. Food insects, grain, and berries.

GEN. 35. CORVUS, *Linn.*

(1. CORVUS, *Cuv.*)

110. *C. Corax*, *Linn.* (*Raven*.)—Plumage black, glossed with blue: nostrils covered by bristly feathers half the length of the bill: tail considerably rounded, extending two inches beyond the folded wings.

C. Corax, *Temm. Man. d'Orn.* tom. i. p. 107. *Id.* tom. III. p. 56.
Raven, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 346. pl. 27*.
Bew. Brit. Birds, vol. i. p. 79.

DIMENS. Entire length two feet one inch: length of the bill two inches nine lines: breadth, wings extended, four feet.

DESCRIPT. The whole plumage black; the upper parts with purple and blue reflections, beneath less glossy: throat feathers long, loose, and acuminate: bill very strong, black: irides with two circles, the outer one brown, the inner gray: tail very much rounded, much longer than in the next species, reaching about two inches beyond the tips of the folded wings: feet black. According to *Low*, *white varieties* have been met with in the Orkneys. (*Egg*.) Pale green ground, spotted and speckled with darker greenish brown: long. diam. two inches; trans. diam. one inch four lines.

The largest species in the genus. Widely dispersed over the country, but not very plentiful. Breeds very early in the year, on steep cliffs, or in large and lofty trees. Nest formed of sticks, and lined with wool, hair, and other substances. Eggs five or six in number. Food small quadrupeds, poultry, game, as well as carrion, and other animal substances. The male and female pair for life, and generally haunt the same spot every year for the purpose of nidification.

111. *C. Corone*, *Linn.* (*Carrion Crow*.)—Plumage bluish black: nostrils covered by bristly feathers one-third the length of the bill: tail moderately rounded, extending an inch and a quarter beyond the wings.

C. Corone, *Temm. Man. d'Orn.* tom. i. p. 108. *Id.* tom. III. p. 58.
Carrion Crow, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 349. pl. 28.
The Crow, *Bew. Brit. Birds*, vol. i. p. 81.

DIMENS. Entire length nineteen inches: length of the bill (from the forehead) one inch ten lines, (from the gape) two inches one line; of the tarsus two inches one line; of the tail seven inches six lines; from the carpus to the end of the wing twelve inches: breadth, wings extended, three feet.

DESCRIPT. Much smaller than the last species, but very similar in general appearance: plumage entirely black; the upper parts reflecting green and violet: tail relatively shorter than in the *C. Corax*, and not so

much rounded at the extremity: bill and feet black: irides dark hazel. (*Egg.*) Ground colour pale bluish green, spotted and speckled with two shades of ash-colour and clove-brown: long. diam. one inch eight lines; trans. diam. one inch two lines.

Common throughout the kingdom, but most abundant in wooded districts. Food and habits similar to those of the Raven. Like that species found in pairs all the year. Breeds as early as February. Nest generally placed in the forked branch of a lofty tree, composed of sticks, and lined with wool, hair, and other soft materials. Eggs four or five in number.

112. *C. Cornix*, Linn. (*Hooded Crow.*)—Head, throat, wings and tail, black, glossed with blue; the rest of the plumage ash-gray.

C. Cornix, *Temm. Man. d'Orn.* tom. i. p. 109. *Id.* tom. III. p. 59.
Hooded Crow, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 351. pl. 29.
Bew. Brit. Birds, vol. i. p. 83.

DIMENS. Entire length nineteen inches six lines: length of the bill (from the forehead) two inches two lines, (from the gape) two inches two lines and a half; of the tarsus two inches three lines; of the tail seven inches six lines; from the carpus to the end of the wing twelve inches ten lines: breadth, wings extended, three feet two inches.

DESCRIPT. Head, throat, wings, and tail, black, with blue and green reflections: all the rest of the plumage cinereous gray: bill strong, shaped like that of the *Raven*, black: nostrils covered by reflected bristly feathers, as in the last species: irides brown: feet black. (*Egg.*) Mottled all over with dark greenish brown on a lighter green ground: long. diam. one inch ten lines; trans. diam. one inch three lines.

Resident all the year in the western and northern parts of Scotland, and in some parts of Ireland, but only a winter visitant in England, where it arrives about the end of October, and remains till the approach of Spring. During its stay chiefly frequents the sea-coast, or open and extensive downs. Builds in trees and rocky cliffs, constructing a nest similar to that of the Carrion Crow. Breeds early in the year, and lays four or five eggs. Feeds indiscriminately on all kinds of animal substances.

113. *C. frugilegus*, Linn. (*Rook.*)—Plumage black, glossed with blue: base of the bill (in the *adult* bird) denuded of feathers, and covered with a white scurf.

C. frugilegus, *Temm. Man. d'Orn.* tom. i. p. 110. *Id.* tom. III. p. 59.
Rook, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 353. pl. 30. *Bew. Brit. Birds*, vol. i. p. 85.

DIMENS. Entire length eighteen inches: length of the bill (from the forehead) two inches two lines, (from the gape) two inches four lines; of the tarsus two inches; of the tail seven inches five lines; from the carpus to the end of the wing twelve inches six lines: breadth, wings extended, three feet.

DESCRIPT. Strongly resembling the *C. Corone*, but may always be distinguished in the adult state by the naked, scabrous, whitish skin surrounding the base of the bill, and by the entire absence of nasal fea-

thers: plumage wholly black as in that species, but rather more glossy, reflecting rich tints of blue and violet-purple: bill not quite so strong, and of not so deep a black: colour of the feet similar. *Young* birds, on leaving the nest, have the base of the bill feathered as in the *C. Corone*, and the nostrils likewise covered by reflected bristles: in the course of the autumn these feathers fall off, and are not replaced by others. *Varieties* are occasionally met with entirely white, or pied, or with the tips of all the feathers whitish. (*Egg*.) Ground colour pale green, nearly covered with blotches of dark greenish brown: long. diam. one inch eight lines; trans. diam. one inch two lines.

Common throughout the country, especially in cultivated districts. Of gregarious habits, breeding together, and likewise seeking food in large companies. Subsists principally on the grub of the cockchafer, wire-worm, and other insects; but will occasionally devour corn, and, during the winter season, is very destructive to turnips. Builds early in March, and hatches in April. Nests crowded together at the tops of the tallest trees, composed principally of fresh twigs forcibly detached from the branches of the neighbouring trees, and lined with grass and fibrous roots. Eggs four or five in number.

114. *C. Monedula*, Linn. (*Jackdaw*.)—Crown of the head, and upper parts of the body, bluish black; occiput and nape ash-gray.

C. Monedula, *Temm. Man. d'Orn.* tom. i. p. 111. *Id.* tom. iii. p. 60. *Jackdaw*, *Mont. Orn. Dict.* *Selb. Illust.* vol. i. p. 356. pl. 31. f. 1. *Bew. Brit. Birds*, vol. i. p. 88.

DIMENS. Entire length twelve inches nine lines: length of the bill (from the forehead) one inch two lines, (from the gape) one inch five lines; of the tarsus one inch eight lines; of the tail four inches eleven lines; from the carpus to the end of the wing nine inches: breadth, wings extended, two feet three inches three lines.

DESCRIPT. Much smaller than any of the foregoing species: crown of the head, and all the upper parts of the body, black, glossed with violet-blue; occiput and nape ash-gray: wings and tail the same as the back: under parts deep black, not so much glossed as the upper: bill and feet black: irides grayish white. *Varies* occasionally like the last species. (*Egg*.) Pale bluish white, spotted with ash-colour and clove-brown: long. diam. one inch seven lines; trans. diam. one inch and half a line.

Common in all parts of the country. Much attached to churches and other buildings, especially such as are in a ruinous and deserted state. Builds in such situations, as well as in the holes of decayed trees; occasionally in chimnies, and even rabbit-burrows. Nest composed of sticks mixed up occasionally with horse-dung, and lined with wool and other soft substances. Eggs four to six in number. Gregarious like the last species, with which it often associates. Feeds on a great variety of animal and vegetable substances.

(2. *PICA*, *Cuv.*)

115. *C. Pica*, Linn. (*Magpie*.)—Head, throat, back, and breast, black; scapulars and belly white.

C. Pica, *Temm. Man. d'Orn.* tom. i. p. 113. Garrulus Picus, *Id.* tom. iii. p. 63. Magpie, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 358. pl. 31. f. 2. *Bew. Brit. Birds*, vol. i. p. 92.

DIMENS. Entire length eighteen inches: length of the bill (from the forehead) one inch four lines, (from the gape) one inch eight lines; of the tarsus two inches one line; of the tail nine inches ten lines; from the carpus to the end of the wing seven inches eight lines: breadth, wings extended, twenty-three inches.

DESCRIPT. Head, throat, neck, upper part of the breast, back, vent, under tail-coverts, and thighs, deep velvet-black; the feathers on the throat rather loose and open, each tipped with a short bristle: scapulars, lower part of the breast, and belly, pure white: wings and tail greenish black, reflecting rich tints of purple, bronze, and blue, according to the position of the eye; the inner webs of the primaries with a large oblong white spot: tail very much cuneated, the length of the feathers rapidly decreasing: irides dark brown: bill and feet black. (*Egg.*) Pale bluish white, spotted all over with ash-colour and two shades of greenish brown: long. diam. one inch four lines and a half; trans. diam. one inch.

A common and generally dispersed species. Frequents woods and thickets, but prefers those near adjoining to villages. Generally observed in pairs. Nest placed in a thick bush, or on the top of a lofty tree; of an oval shape, constructed of sticks closely interwoven with each other, the bottom plastered with clay, and lined with fibrous roots. Eggs six or seven in number, laid early in the Spring. Omnivorous.

GEN. 36. GARRULUS, *Briss.*

116. *G. glandarius*, *Flem. (Jay.)*—A black moustache descending from the corners of the bill: greater wing-coverts barred with black and blue.

Corvus glandarius, *Temm. Man. d'Orn.* tom. i. p. 114. Garrulus gland. *Id.* tom. iii. p. 65. Jay, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 362. pl. 32. *Bew. Brit. Birds*, vol. i. p. 94.

DIMENS. Entire length thirteen inches eight lines: length of the bill (from the forehead) one inch two lines, (from the gape) one inch five lines and a half; of the tarsus one inch eight lines and a half; of the tail five inches nine lines; from the carpus to the end of the wing seven inches three lines: breadth, wings extended, twenty-two inches.

DESCRIPT. The feathers on the forehead and crown narrow and elongated, forming a crest capable of erection, of a whitish colour streaked with black; chin whitish: a broad black moustache, about an inch in length, descending from the corners of the lower mandible: general colour of the plumage above and below reddish or purple ash; darkest on the nape and upper part of the back; rump, and lower part of the abdomen, whitish: quills and tail dusky, the former edged with white; bastard wing and greater coverts bright blue, with several narrow transverse bars of black; secondaries with an oblong white spot on their outer webs: bill black: irides bluish white: feet livid brown. White and other varieties sometimes occur. (*Egg.*) Yellowish white ground, minutely and thickly speckled all over with light brown, presenting the appearance of a uniform yellow gray-brown: long. diam. one inch four lines; trans. diam. one inch.

Found in extensive woods in most parts of the kingdom. Like the last species omnivorous. Has a harsh disagreeable cry. Nest placed in close thickets, formed of sticks and lined with fibrous roots. Lays five or six eggs.

GEN. 37. NUCIFRAGA, *Briss.*

117. *N. Caryocatactes*, Temm. (*Nutcracker*).—General plumage brown, spotted on the back and under parts with white.

N. Caryocatactes, Temm. *Man. d'Orn.* tom. i. p. 117. *Id.* tom. III. p. 67. *Nutcracker*, Mont. *Orn. Dict. Selb. Illust.* vol. i. p. 368. pl. 33*. *Bew. Brit. Birds*, vol. i. p. 97.

DIMENS. Entire length thirteen inches: length of the bill two inches. MONT.

DESCRIPT. Crown of the head and nape of the neck blackish brown: the rest of the plumage above and below the body rusty brown, spotted with white; the spots on the back guttiform; those beneath disposed longitudinally on each feather, and occupying a larger portion of the ground-colour: vent white: quills and tail black; the latter with a broad white bar at the extremity: irides brown: bill and feet dusky. (*Egg*.) Pale buff-coloured white: long. diam. one inch three lines; trans. diam. ten lines and a half.

Very rare in this country. Has been only observed in three or four instances. One killed near Moyston in Flintshire in October 1753; a second in Kent; a third in Northumberland in the autumn of 1819. Common in some of the mountainous parts of the North of Europe, residing principally in woods. Feeds on seeds and nuts; also on insects. Said to build in the holes of trees, and to lay five or six eggs.

GEN. 38. PICUS, *Linn.*

118. *P. viridis*, Linn. (*Green Woodpecker*).—Plumage above, green; beneath, greenish ash; crown and occiput bright red.

P. viridis, Temm. *Man. d'Orn.* tom. i. p. 391. *Id.* tom. III. p. 280. *Green Woodpecker*, Mont. *Orn. Dict. Selb. Illust.* vol. i. p. 372. pl. 38. f. 2. *Bew. Brit. Birds*, vol. i. p. 136.

DIMENS. Entire length thirteen inches three lines: length of the bill (from the forehead) one inch seven lines, (from the gape) one inch nine lines and a half; of the tarsus one inch two lines; of the tail three inches ten lines and a half; from the carpus to the end of the wing six inches four lines; breadth, wings extended, twenty-one inches.

DESCRIPT. Crown and occiput bright crimson-red; base of the bill, and region of the eyes, black; a broad crimson moustache edged with black descending from the corner of the bill: hind part of the neck, back, and wing-coverts, bright green; rump gamboge-yellow: under parts

grayish white, with a tinge of yellowish green: primaries dusky, with a regular series of pale yellow spots on the outer webs: tail-feathers presenting alternate bars of green and dusky-brown; the extreme tips black: bill dusky; the base of the lower mandible yellowish: irides white: feet pale brown with a tinge of green. In the *female*, there is less red on the head, and less black about the eyes; the moustache is entirely black. In *very young* birds, the red on the head is mixed with yellowish gray; the green tint on the upper parts paler, and irregularly spotted with ash-gray; the moustache incomplete; and the under parts marked with transverse brown bars. (*Egg*.) Smooth and shining: pure white: long. diam. one inch two lines and a half; trans. diam. ten lines and a half.

Common in wooded districts throughout the country. Feeds on ants, and other insects; more especially the larvæ of the timber-eating species, which it extracts by means of its long tongue, after having perforated the wood with its bill. Breeds in the holes of trees. Eggs four or five in number, deposited on the rotten wood.

119. *P. major*, Linn. (*Great Spotted Woodpecker*.)
—Crown, and upper part of the body, black; scapulars white; a crimson patch on the occiput.

P. major, *Temm. Man. d'Orn.* tom. i. p. 395. *Id.* tom. iii. p. 281.
Greater Spotted Woodpecker, *Mont. Orn. Dict. Selb. Illust.*
vol. i. p. 376. pl. 38. f. 3. Pied Woodpecker, *Bew. Brit. Birds*,
vol. i. p. 138.

DIMENS. Entire length nine inches six lines: length of the bill (from the forehead) one inch, (from the gape) one inch three lines; of the tarsus eleven lines; of the tail three inches seven lines; from the carpus to the end of the wing five inches: breadth, wings extended, fourteen inches.

DESCRIPT. A transverse band of dirty white on the forehead; crown of the head black; occiput crimson-red: cheeks and ear-coverts white, bounded beneath by a black line, proceeding from the corner of the mouth towards the nape, from whence arises another line of the same colour that passes down the side of the neck and terminates on the breast: a white spot on each side of the back part of the neck: back and lesser wing-coverts velvet-black; scapulars, and some of the greater coverts nearest the body, pure white: quills black, with a series of white spots on each web: throat, fore part of the neck, breast and belly, dirty white; abdomen and under tail-coverts rich crimson: tail black; the three lateral feathers on each side spotted and tipped with dirty white: irides red: bill and feet dusky lead-colour. In the *female*, there is no red on the occiput. In *young* birds, until the first moult, the crown of the head is red, and the patch on the occiput black; the black on the upper parts is also tinged with brown. (*Egg*.) Smooth; shining white: long. diam. one inch; trans. diam. nine lines.

Not so abundant as the last species, but hardly to be esteemed rare. Haunts, food, and habits, very similar. Makes a loud jarring noise in the Spring, (probably a call-note to the other sex) by striking its bill very rapidly and many times in succession against the branch of a tree. Makes no nest, but deposits its eggs, which are four or five in number, in the holes of decayed trees. *Obs.* The *Middle Spotted Woodpecker* of Montagu and other English authors is only the young of this species: the *Picus medius* of Temminck is distinct; but not hitherto found in this country.

120. *P. minor*, Linn. (*Lesser Spotted Woodpecker*.)
—Occiput and nape black: middle of the back, and scapulars, with white and black bars: crown of the head red.

P. minor, *Temm. Man. d'Orn.* tom. i. p. 399. *Id.* tom. iii. p. 283.
Lesser Spotted Woodpecker, *Mont. Orn. Dict. Selb. Illust.* vol. i.
p. 379. pl. 38. f. 4. Barred Woodpecker, *Bew. Brit. Birds*, vol. i.
p. 140.

DIMENS. Entire length five inches six lines: length of the bill seven lines and a half: breadth, wings extended, twelve inches.

DESCRIPT. Forehead dirty white; crown of the head bright red; streak over the eyes, occiput and nape, black; cheeks and sides of the neck white; from the corners of the lower mandible a black streak directed towards the shoulders: upper part of the back, rump, and lesser wing-coverts, glossy black; the rest of the upper parts, including the middle region of the back, scapulars, and quills, transversely barred with black and white: all the under parts of a dirty brownish white, with a few fine longitudinal dark streaks on the breast and sides: the four middle tail-feathers glossy black; the three outer ones on each side tipped with white and barred with black: bill and feet dusky lead-colour: irides reddish brown. In the *female*, the crown is dirty white instead of red. (*Egg*.) Smooth, delicate white: long. diam. nine lines and a half; trans. diam. seven lines.

Much less frequent than either of the preceding species, and only partially distributed. Met with in the counties of Gloucester, Wilts, Hereford, Cambridge, Norfolk, and Suffolk. Said to be very rare in the North of England. Habits and nidification similar to those of the last.

(7.) *P. martius*, Linn. (*Great Black Woodpecker*.) *Temm. Man. d'Orn.* tom. i. p. 390. Gould, *Europ. Birds*, part i. *Selb. Illust.* vol. i. p. 375. pl. D. f. 4.

This species, which is not uncommon in the northern parts of Europe, has been included in the British lists principally on the authority of Drs. Latham and Pulteney. The former author states (*Syn. Supp.* p. 104.) that it has been sometimes seen in Devonshire; the latter (*Cat. Dors.* p. 6.) that it has been more than once shot in Dorsetshire. No specimen, however, known to have been certainly killed in this country, exists in any of our museums, and there is strong reason to doubt the reality of its claims to a place in the British Fauna.

(8.) *P. villosus*, Linn. (*Hairy Woodpecker*.) Lewin, *Brit. Birds*, vol. II. pl. 50. *Mont. Orn. Dict.*

A pair of this species, in the collection of the late Dowager Duchess of Portland, were said to have been shot near Halifax in Yorkshire. It is supposed, however, that this was an error, and that the above locality had been confounded with Halifax in North America, where the species is not uncommon.

(9.) *P. tridactylus*, Linn. (*Three-toed Woodpecker*.) *Temm. Man. d'Orn.* tom. i. p. 401. Don, *Brit. Birds*, vol. vi. pl. 143.

A native of the northern parts of Europe and America. According to Donovan, has been shot in the North of Scotland. This, however, is probably a mistake.

GEN. 39. YUNX, *Linn.*

121. *Y. Torquilla*, *Linn.* (*Wryneck*.)—Upper parts yellowish gray, irregularly spotted and lined with brown and black; a broad black mesial stripe on the back of the neck.

Y. Torquilla, *Temm. Man. d'Orn.* tom. I. p. 403. *Id.* tom. III. p. 284. *Wryneck*, *Mont. Orn. Dict. Selb. Illust.* vol. I. p. 381. pl. 38. f. 1. *Bew. Brit. Birds*, vol. I. p. 129.

DIMENS. Entire length seven inches four lines: length of the bill (from the forehead) six lines, (from the gape) nine lines and a half; of the tarsus eight lines and a half; of the tail two inches eight lines: from the carpus to the end of the wing three inches six lines: breadth, wings extended, eleven inches.

DESCRIPT. General colour of the upper plumage reddish or yellowish gray, irregularly spotted and marked with various shades of brown and black; more particularly a broad black mesial streak extending from the occiput to the upper part of the back: throat and fore part of the neck reddish yellow, with transverse undulating dusky lines; rest of the under parts whitish, with arrow-shaped black spots: quills marked on the outer webs with oblong red spots: tail-feathers rounded at the end, mottled like the back, with four transverse black bars: irides yellowish brown: bill and feet brown. (*Egg*.) Smooth, delicate white: long. diam. nine lines and a half; trans. diam. seven lines.

A summer visitant, first appearing about the second week in April. Not very uncommon in wooded districts throughout the southern, midland, and eastern counties: said to be more rare in the West of England. Utters a loud and oft-repeated cry during the breeding season, somewhat resembling that of the Kestrel Hawk. Food principally ants. Eggs six to ten in number, deposited in the holes of trees on the rotten wood, without a nest.

GEN. 40. CERTHIA, *Linn.*

122. *C. familiaris*, *Linn.* (*Common Creeper*.)—Upper parts yellowish brown; variegated with dusky, and white spots: rump reddish yellow.

C. familiaris, *Temm. Man. d'Orn.* tom. I. p. 410. *Id.* tom. III. p. 288. *Common Creeper*, *Mont. Orn. Dict. Selb. Illust.* vol. I. p. 388. pl. 39. f. 2. *Bew. Brit. Birds*, vol. I. p. 148.

DIMENS. Entire length five inches: length of the bill (from the forehead) five lines and three quarters, (from the gape) seven lines; of the tarsus seven lines; of the tail two inches one line and a half; from the carpus to the end of the wing two inches four lines: breadth, wings extended, seven inches three lines.

DESCRIPT. Head, and all the upper parts, yellowish brown, mixed with dusky, with an oblong whitish spot in the centre of each feather; rump rust-red; above the eyes a whitish streak: all the under parts white, tinged with reddish on the lower part of the abdomen: quills tipped with white; the four first dusky, the rest with a broad transverse reddish white bar about the middle: tail-feathers long, stiff, and acumi-

nated; brownish gray, tinged with red: upper mandible brown; lower mandible yellowish: feet yellowish brown. (*Egg.*) White, with pale red spots often confined to the larger end: long. diam. eight lines; trans. diam. five lines and a half.

Common and generally dispersed throughout the country. Climbs trees with great facility. Feeds entirely on insects. Has a peculiar, but rather monotonous song, heard early in the Spring, and continued during the breeding season. Nest placed in the holes, or under the loose bark, of decayed trees: formed of small sticks, wool, and mosses packed rudely together, and lined with feathers and fine shreds of wood. Eggs six or eight in number, laid towards the end of April.

GEN. 41. TROGLODYTES, *Cuv.*

123. *T. Europæus*, *Selb.* (*Common Wren.*)—Upper plumage reddish brown, with transverse dusky lines: over the eye a narrow white streak.

Sylvia Troglodytes, *Temm. Man. d'Orn.* tom. i. p. 233. *Troglodytes vulgaris*, *Id.* tom. iii. p. 160. *Common Wren*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 390. pl. 47. f. 6. *Bew. Brit. Birds*, vol. i. p. 262.

DIMENS. Entire length three inches nine lines: length of the bill (from the forehead) three lines and a half, (from the gape) five lines and a half; of the tarsus nine lines; of the tail one inch four lines; from the carpus to the end of the wing one inch nine lines and a half: breadth, wings extended, six inches three lines.

DESCRIPT. Upper parts of the body deep reddish brown, faintly marked with transverse dusky lines: over each eye a pale narrow streak: quills barred alternately on their outer webs with blackish brown and reddish: tail dusky, with transverse black bars: under parts light rufous-brown; the sides and thighs streaked with darker lines; under tail-coverts obscurely spotted with white: bill brown: irides hazel: feet yellowish brown. (*Egg.*) White, with a few specks of pale red: long. diam. seven lines and a half; trans. diam. six lines.

Abundant in all parts of the country, remaining with us the whole year. Frequents gardens and out-houses. Song shrill and loud. Nest often fixed against the thatch of buildings, or placed in the holes of trees, more rarely on the ground; of an oval form, covered over at top, with the entrance on one side; formed of moss, hay, leaves, and other materials, and lined with feathers occasionally mixed with hair. Eggs five to eight in number.

GEN. 42. UPUPA, *Linn.*

124. *U. Epops*, *Linn.* (*Hoopoe.*)—Head, neck, and breast, purplish red; wings black, barred with white.

U. Epops, *Temm. Man. d'Orn.* tom. i. p. 415. *Id.* tom. iii. p. 291. *Hoopoe*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 393. pl. 40. f. 2. *Bew. Brit. Birds*, vol. i. p. 144.

DIMENS. Entire length twelve inches: length of the bill two inches six lines: breadth, wings extended, nineteen inches.

DESCRIPT. Crest composed of a double row of elongated feathers, orange-red tipped with black; a small white patch intervening between these two colours: head, neck, and breast, purplish red: upper part of

the back pale grayish brown; middle and lower regions barred with black and white: wings black, with five transverse bars of yellowish white: tail black, with one broad, white, crescent-shaped bar: abdomen white; flanks with a few longitudinal brown streaks: bill flesh-red at the base, black at the tip: irides and feet brown. In the *female*, the colours are paler, and the crest not so long. (*Egg*.) Uniform pale lavender-gray: long. diam. one inch and half a line; trans. diam. eight lines.

Visits this country most years, but in very small numbers. Occasionally met with in different parts of England: rarely breeds with us. Habits somewhat terrestrial. Food coleopterous, and other insects. Builds in the holes of trees. Nest formed of bents, and lined with feathers and other soft materials. Eggs four or five in number.

GEN. 43. SITTA, Linn.

125. *S. Europæa*, Linn. (*Nuthatch*.)—Upper parts bluish ash; breast and belly reddish yellow; a black streak across the eye and ear-coverts.

S. Europæa, *Temm. Man. d'Orn.* tom. i. p. 407. *Id.* tom. iii. p. 285.

Nuthatch, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 385. pl. 39. f. 1. *Bew. Brit. Birds*, vol. i. p. 142.

DIMENS. Entire length six inches ten lines.

DESCRIPT. Crown of the head, and all the upper parts of the plumage bluish ash: a black streak from the corner of the bill across each eye, and extending down the sides of the neck as far as the shoulder; cheeks and throat white; breast and belly dull orange-red; sides and thighs ferruginous chestnut; under tail-coverts white, edged with ferruginous: tail consisting of twelve short flexible feathers; the two middle ones gray; the four outer ones black, with a white spot near the extremity, the tips themselves gray: bill blackish above, white at the base of the lower mandible: irides hazel: feet yellowish gray. (*Egg*.) White, spotted and speckled with pale red: long. diam. nine lines; trans. diam. seven lines.

Common in wooded districts in many parts of the country, but not in all. Rare in some of the northern and western counties. Remains with us the whole year. Climbs trees with great facility, and equally well in all directions. Feeds principally on nuts, which it breaks with its bill after having firmly fixed them in the crevices of old trees; occasionally on insects. Nest placed in the holes of trees, and formed of dead leaves. Eggs six or seven in number.

GEN. 44. CUCULUS, Linn.

126. *C. canorus*, Linn. (*Common Cuckow*.)—Head, neck, breast, and upper parts, bluish ash: abdomen whitish, with transverse dusky bars.

C. canorus, *Temm. Man. d'Orn.* tom. i. p. 381. *Id.* tom. iii. p. 272.

Common Cuckow, *Mont. Orn. Dict. and Supp. Selb. Illust.* vol. i. p. 397. pl. 37, and pl. 43. f. 3. *Bew. Brit. Birds*, vol. i. p. 124.

DIMENS. Entire length thirteen inches six lines: length of the bill (from the forehead) ten lines, (from the gape) one inch two lines; of the

tarsus ten lines and a half; of the tail six inches ten lines; from the carpus to the end of the wing eight inches nine lines: breadth, wings extended, twenty-three inches four lines.

DESCRIPT. (*Adult.*) Head, and all the upper parts, bluish ash; throat, fore part of the neck, and breast, the same, but rather paler; belly, thighs, and under tail-coverts, whitish, with transverse streaks of dusky brown: quills dusky, barred on the inner webs with oval white spots: tail black, with a series of small white spots on the shafts of the feathers; the tips also white: bill dusky, yellowish at the base and edges; inside of the mouth, and orbits, orange-yellow: irides and feet yellow. (*Young of the year.*) All the upper parts of a deep clove-brown, with transverse bars of pale ferruginous brown, the feathers tipped with whitish; a patch of white on the occiput: throat, and under parts, yellowish white, with transverse black bars: quills spotted with reddish brown on their inner webs: tail with alternate oblique bars of red and brown, the brown bar nearest the extremity broader than the others; the shafts of the feathers with a series of white spots; the tips white: irides liver-brown. (*Egg.*) White, speckled all over with ash-brown; or reddish white, speckled with nutmeg-brown: long. diam. eleven lines; trans. diam. eight lines and a half.

Visits this country early in April, and leaves it again about the beginning of July; the young of the year remaining till September. Feeds principally on caterpillars, and other insects. Makes no nest, but commits its eggs (five or six in number) to the nests of other birds, generally selecting those of the Hedge Sparrow, Wagtail, or Tit-Pipit.

GEN. 45. COCCYZUS, Vieill.

127. *C. Americanus*, Bonap. (*Carolina Cuckow*.)

Cuculus Americanus, Linn. *Syst. Nat.* tom. i. p. 170. *Lath. Ind. Orn.* vol. i. p. 219. *Cuc. Carolinensis*, Wils. *Amer. Orn.* vol. iv. p. 13. pl. 28. f. 1. *Cuc. cinerosus*, Temm. *Man. d'Orn.* tom. iii. p. 277. *Carolina Cuckow*, *Lath. Syn.* vol. ii. p. 537.

DIMENS. Entire length eleven inches four lines: length of the bill (from the forehead) one inch, (from the gape) one inch four lines; of the tarsus eleven lines and a half; of the tail five inches seven lines; from the carpus to the end of the wing five inches nine lines.

DESCRIPT. All the upper parts of the head and body, wings, and two middle tail-feathers, cinereous brown, with a slight tinge of olivaceous; the other tail-feathers black, with a broad white space at the extremity of each of the three outermost; the fourth just tipped with white; primaries and wing-coverts bright rufous: throat, sides of the neck, and all the under parts white: upper mandible black, edged with yellow at the base; lower mandible yellow, tipped with black: legs black: tarsi long, naked. (*Egg.*) "Of a uniform greenish blue colour". WILS.

The above description of this species is taken from a specimen in the collection of the Zoological Society, which was killed in the preserves of Lord Cawdor in Wales, in the autumn of 1832. Three other individuals have occurred in this country, two in Ireland, and one in Cornwall. Inhabits the northern parts of America. Habits said to be essentially different from those of the *Common Cuckow*. Constructs its own nest, and rears its own young. Eggs three or four in number. Food, according to Wilson, principally caterpillars.

GEN. 46. CORACIAS, Linn.

128. *C. garrula*, Linn. (*Garrulous Roller*).—Head, neck, and under parts, light bluish green: back and scapulars reddish brown.

C. garrula, Temm. *Man. d'Orn.* tom. i. p. 127. *Id.* tom. iii. p. 72.
Garrulous Roller, Mont. *Orn. Dict. Selb. Illust.* vol. i. p. 117.
pl. 34. Roller, Bew. *Brit. Birds*, vol. i. p. 100.

DIMENS. Entire length twelve inches six lines. MONT.

DESCRIPT. Head, neck, breast, and belly, verditer blue, inclining to sea-green: back and scapulars reddish brown: lesser wing-coverts rich violet-blue; greater coverts pale green: rump, and basal portion of the quills, purplish blue; the tips of the quills dusky: tail somewhat forked; the outer feather pale ultra-marine blue, tipped with black; the others dusky green, tipped with pale bluish green, the two middle ones excepted, which are dusky throughout: bill black at the tip, yellowish brown at the base: feet yellowish. (*Egg*.) Smooth, shining white: long. diam. one inch five lines; trans. diam. one inch one line.

Very rare in this country, and only an accidental visitant. First noticed as a British bird by Pennant, who describes one killed in Cornwall. Solitary individuals have been since met with in Sussex, Suffolk, Norfolk, Scotland, Northumberland, and Yorkshire. Said to be common in Germany and Sweden, inhabiting large forests. Feeds principally on insects. Builds in the holes of decayed trees, and lays from four to seven eggs.

GEN. 47. MEROPS, Linn.

129. *M. Apiaster*, Linn. (*Common Bee-eater*).—Forehead greenish white; nape, and upper part of the back, deep chestnut: throat yellow, bounded by a black line.

M. Apiaster, Temm. *Man. d'Orn.* tom. i. p. 420. *Id.* tom. iii. p. 293.
Sow. Brit. Misc. pl. 69. Common Bee-eater, Mont. *Orn. Dict. Selb. Illust.* vol. i. p. 114. pl. 41. *Bew. Brit. Birds*, vol. i. p. 146.

DIMENS. Entire length eleven inches. TEMM.

DESCRIPT. Forehead white, passing into bluish green; crown of the head, nape, and upper part of the back, deep chestnut: middle and lower regions of the back of a paler chestnut, passing into reddish yellow: from the corner of the bill a black streak passing over the eyes and across the ear-coverts, where it meets another line of black encircling the lower portion of the neck; throat golden yellow; the rest of the under parts bluish green: lesser wing-coverts grass-green; middle portion of the wing brownish red: quills dusky, passing into bluish green on their outer webs: tail greenish blue; the two middle feathers somewhat darker, pointed, and longer than the others by more than an inch: bill black: irides red: feet brown. In the *female*, the colours are for the most part similar, only paler: the two middle tail-feathers are relatively shorter. (*Egg*.) Smooth, and white; nearly round: long. diam. one inch and half a line; trans. diam. eleven lines.

Very rarely seen in this country. A flight of about twenty was observed near Mattishall in Norfolk, and one killed, in June 1794; since

which period, other specimens have been met with on different occasions in Cornwall, Devonshire, and Ireland. Common in some parts of the Continent. Feeds principally on Hymenopterous insects. Nest placed in deep holes excavated in the banks of rivers for that purpose. Eggs five to seven in number.

GEN. 48. ALCEDO, Linn.

130. *A. Ispida*, Linn. (*Common King-Fisher*.)—Crown of the head, and wing-coverts, deep green, spotted with azure-blue: behind the eye a patch of orange-brown passing into white.

A. Ispida, *Temm. Man. d'Orn.* tom. i. p. 423. *Id.* tom. iii. p. 296. *Common King-Fisher*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 136. pl. 40. f. 1. *Bew. Brit. Birds*, vol. ii. p. 121.

DIMENS. Entire length seven inches three lines: length of the bill (from the forehead) one inch six lines, (from the gape) two inches; of the tarsus four lines and a half; from the carpus to the end of the wing three inches: breadth, wings extended, ten inches.

DESCRIPT. Crown of the head deep olive-green, the feathers tipped with bright azure-blue: from the upper mandible to the eye a dusky streak: behind the eye a band of orange-brown, passing on the sides of the neck into a white patch; below this, extending from the base of the lower mandible to near the insertion of the wing, a broad streak of azure-green: middle of the back, rump, and upper tail-coverts, fine bright azure: throat, and fore part of the neck, yellowish white: rest of the under parts ferruginous orange: wing-coverts and quills deep greenish blue, spotted like the crown of the head, but more sparingly: tail greenish blue; the shafts of the feathers black: bill dusky brown, reddish at the base: irides hazel: feet orange-red. (*Egg*.) Smooth, and white; nearly round: long. diam. ten lines and a half; trans. diam. nine lines.

Generally diffused over the country, and resident all the year. Haunts the banks of rivers and clear streams, feeding on fish and aquatic insects. Is rapid on wing, and frequently utters a shrill cry in its flight. Nest placed in holes in the ground near the water's edge, consisting simply of fish bones and other indigestible parts of the food rejected from the stomach after eating. Eggs six or seven in number.

GEN. 49. HIRUNDO, Linn.

131. *H. rustica*, Linn. (*Chimney Swallow*.)—Upper parts, and a transverse bar on the breast, bluish black; forehead and throat chestnut-red.

H. rustica, *Temm. Man. d'Orn.* tom. i. p. 427. *Id.* tom. iii. p. 297. *Chimney Swallow*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 120. pl. 42. f. 1. *Swallow*, *Bew. Brit. Birds*, vol. i. p. 287.

DIMENS. Entire length seven inches: length of the bill (from the forehead) three lines and a half, (from the gape) eight lines; of the tarsus five lines; of the tail four inches; the same, excluding the long

lateral feathers, two inches six lines; from the carpus to the end of the wing four inches eleven lines: breadth, wings extended, thirteen inches four lines.

DESCRIPT. Forehead and throat chestnut-red: all the upper parts, sides of the neck, and a broad transverse bar on the breast, black, with purple and blue reflections: belly and vent reddish white: tail long, and very much forked; the two middle feathers plain; the rest with a large white spot on the inner web: bill and feet black. The *female* has less red on the forehead, and less black on the breast: the under parts are also whiter, and the outer tail-feathers not so long as in the male bird. In *young* birds, the long tail-feathers do not appear till after the first moult. White *varieties* are sometimes met with. (*Egg.*) White, spotted and speckled with ash-colour and dark red-brown: long. diam. nine lines and a half; trans. diam. six lines and a half.

A summer visitant, making its first appearance about the second or third week in April, and staying till towards the middle or end of October. Feeds entirely on insects, taken on the wing. Builds generally in chimnies: nest formed of mud plastered together, and lined with feathers. Eggs four or five in number: two broods in the season. Previous to migration, congregates in large flocks.

132. *H. urbica*, Linn. (*House Martin.*)—Head, nape, and back, bluish black; rump white: tarsi and toes feathered.

H. urbica, *Temm. Man. d'Orn.* tom. i. p. 428. *Id.* tom. III. p. 300.
Martin, Mont. Orn. Dict. Selb. Illust. vol. i. p. 123. pl. 42. f. 2.
Bew. Brit. Birds, vol. i. p. 293.

DIMENS. Entire length five inches six lines: length of the bill (from the forehead) three lines, (from the gape) six lines; of the tarsus six lines; of the tail two inches six lines and a half; from the carpus to the end of the wing four inches four lines.

DESCRIPT. Head, nape, and all the upper region of the back, glossy bluish black: wings and tail dusky brown; the latter forked, but not so long as in the last species: rump, and all the under parts from the chin to the vent, pure white: tarsi and toes covered with a white down. (*Egg.*) Smooth, delicate white: long. diam. nine lines and a half; trans. diam. six lines.

Rather later in its arrival than the last species, being seldom observed in abundance before the end of April or beginning of May. Departs in October; but a few stragglers may occasionally be seen on to November. Food and habits similar to those of the Swallow: flight not so rapid. Nest usually fixed under the eaves of houses or in the angles of windows; formed entirely of mud externally, and lined with feathers. Lays four or five eggs, and has frequently two broods in the Summer. Congregates at the approach of Autumn.

133. *H. riparia*, Linn. (*Bank Martin.*)—All the upper parts, and a transverse band on the breast, cinereous brown: tarsi and toes naked.

H. riparia, *Temm. Man. d'Orn.* tom. i. p. 429. *Id.* tom. III. p. 300.
Sand Martin, Mont. Orn. Dict. Selb. Illust. vol. i. p. 125. pl. 42.
 f. 3. *Bew. Brit. Birds*, vol. i. p. 295.

DIMENS. Entire length five inches two lines: length of the bill (from the forehead) three lines, (from the gape) six lines; of the tarsus five lines and a half; of the tail two inches one line; from the carpus to the end of the wing four inches.

DESCRIPT. All the upper parts, cheeks, and a transverse band on the breast, cinereous or mouse-coloured brown: wings and tail inclining to dusky brown: throat, fore part of the neck, belly, and under tail-coverts, white: tarsi and toes naked, with the exception of a few small feathers near the origin of the hind toe. In *young* birds, all the upper parts of the plumage are edged with pale reddish brown; the tail-feathers with yellowish white. (*Egg.*) White: long. diam. eight lines; trans. diam. six lines.

First seen about the beginning of April. More locally distributed than either of the preceding species. Found only in the neighbourhood of sand-pits and the high banks of rivers, in which situations it builds, excavating a horizontal hole in the loose soil to the depth of two or three feet. Nest placed at the extremity of the hole, formed of dry grass and straw, and lined with feathers. Eggs four or five in number.

GEN. 50. CYPSELUS, *Illig.*

134. *C. Apus*, Flem. (*Common Swift.*)—Chin white; all the rest of the plumage black.

C. murarius, *Temm. Man. d'Orn.* tom. i. p. 434. *Id.* tom. iii. p. 303.
Common Swift, *Selb. Illust.* vol. i. p. 127. pl. 42. f. 4. Swift,
Mont. Orn. Dict. Bew. Brit. Birds, vol. i. p. 296.

DIMENS. Entire length eight inches: length of the bill (from the forehead) three lines and a half, (from the gape) eight lines and a half; of the tarsus six lines; of the tail three inches four lines; from the carpus to the end of the wing five inches ten lines and a half: breadth, wings extended, seventeen inches.

DESCRIPT. Throat grayish white: all the rest of the plumage above and below sooty black, with greenish reflections: wings and tail extremely long; the latter much forked: tarsi covered with small feathers: irides deep brown: bill and feet black. (*Egg.*) White: long. diam. one inch; trans. diam. eight lines.

Late in its arrival, seldom shewing itself before the beginning, and in some parts of the country not till near the end, of May. Departs also much sooner than the other species of this family; generally about the middle of August. Haunts churches, towers, and other lofty buildings, in the holes of which it breeds. Nest formed of dried grass, and lined with feathers. Eggs two in number. Only one brood in the season. Feeds entirely on insects, taken on the wing. Flight high, and extremely rapid.

135. *C. alpinus*, Temm. (*Alpine Swift.*)—General colour of the plumage grayish brown: throat, and middle of the abdomen, white.

C. alpinus, *Temm. Man. d'Orn.* tom. i. p. 433. *Id.* tom. iii. p. 303.
Greatest Martin, *Edw. Nat. Hist.* vol. i. pl. 27. White-bellied
Swift, *Lath. Syn.* vol. ii. p. 586.

DIMENS. Entire length nearly nine inches. **TEMM.**

DESCRIPT. All the upper parts of a uniform grayish brown; breast, sides of the abdomen, and under tail-coverts the same: throat, and middle of the belly, pure white: tarsi covered with brown feathers. (*Egg.*) White: of the same size as that of the last species.

Shot within eight or ten miles of the south coast of Ireland in 1829, about Midsummer. A second individual has been since killed at Kingsgate, in the Isle of Thanet, and a third in Norfolk. Common in alpine and rocky districts in the southern parts of Europe. Habits somewhat similar to those of the Common Swift. Said to build in the clefts of rocks and old buildings, and to lay three or four eggs.

GEN. 51. CAPRIMULGUS, *Linn.*

136. *C. Europæus*, *Linn.* (*European Goatsucker.*)—

General colour of the plumage ash-gray, variegated with black, brown, and ferruginous.

C. Europæus, *Temm. Man. d'Orn.* tom. i. p. 436. *Id.* tom. iii. p. 304. *European Goat-sucker*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 131. pl. 42^f. *Night-Jar*, *Bew. Brit. Birds*, vol. i. p. 302.

DIMENS. Entire length ten inches four lines: length of the bill (from the forehead) four lines and a half, (from the gape) one inch two lines; of the tarsus eight lines; of the tail four inches eight lines; from the carpus to the end of the wing seven inches nine lines: breadth, wings extended, twenty-two inches.

DESCRIPT. General colour of the plumage ash-gray, beautifully variegated with black, brown, ferruginous, and white, disposed in bars, spots and specks, of different shades and sizes: on the head, and down the middle of the back, some longitudinal black streaks: a little white on the throat, as well as beneath each eye: under parts in general yellowish brown, with transverse undulating lines of black: quills dusky, with ferruginous spots on their outer webs; the three first with also a large white patch on the inner web about midway: tail nearly square, yellowish gray, with transverse zigzag bars of black: the two outer feathers on each side tipped with white: bill and irides dusky: feet yellowish brown. The *female* wants the white spots on the tips of the quills and two outer tail-feathers. (*Egg.*) Of an oval form: ground colour white, mottled with cinereous; this last colour including spots of two shades of brown: long. diam. one inch two lines; trans. diam. ten lines and a half.

One of our latest summer visitants, not appearing before the middle or end of May. Stays till September or October. Generally distributed over the kingdom, but not equally plentiful in all parts. Found principally in wild uncultivated districts, where there is wood. Habits crepuscular. Food insects; particularly the *Melolontha* tribe, and the larger *Lepidoptera*. Makes no nest, but lays two eggs on the bare ground amongst fern, heath, or long grass. During the season of incubation the male utters a peculiar noise somewhat resembling that of a spinning-wheel.

ORDER III. RASORES.

GEN. 52. COLUMBA, Linn.

(1. COLUMBA, Swains.)

137. *C. Palumbus*, Linn. (*Ring-Dove*).—Bluish ash; a white space on the sides of the neck, and on the edge of the wing; extremity of the tail black.

C. Palumbus, *Temm. Man. d'Orn.* tom. II. p. 444. *Id. Pig. et Gall.* tom. I. p. 78. *Ring-Dove*, *Mont. Orn. Dict. Selb. Illust.* vol. I. p. 406. pl. 56. f. 1. *Bew. Brit. Birds*, vol. I. p. 307.

DIMENS. Entire length sixteen inches six lines: length of the bill (from the forehead) nine lines and a half, (from the gape) one inch two lines; of the tarsus one inch four lines; of the tail six inches six lines; from the carpus to the end of the wing ten inches: breadth, wings extended, twenty-nine inches four lines.

DESCRIPT. Head, cheeks, throat, rump, and basal portion of the tail, bluish ash; on each side of the neck a patch of white; a narrow line of the same colour running longitudinally down the wing near the edge; breast and upper part of the belly vinaceous red, with glossy green reflections: back and wing-coverts deep bluish gray; quills dusky, edged with white: a broad black bar at the extremity of the tail; vent, thighs, and under tail-coverts, grayish white: bill orange; the soft portion at the base covered with a white mealy substance: irides light yellow: feet red. (*Egg*.) White: long. diam. one inch eight lines; trans. diam. one inch two lines.

Common in wooded districts throughout the country, remaining the whole year. Utters a cooing note, heard from February to the beginning of October. Builds in April: nest placed in trees, formed of twigs loosely put together, and very shallow. Eggs always two in number. Frequently a second, or even a third brood. Food, grain and seeds of various kinds; during severe weather, the leaves of turnips, and other vegetables. Collects in large flocks at the approach of Winter.

138. *C. Œnas*, Linn. (*Stock-Dove*).—Bluish ash; sides of the neck with green and purple reflections; some of the wing-coverts spotted with black, forming an irregular transverse bar; rump bluish gray; tip of the tail black.

C. Œnas, *Temm. Man. d'Orn.* tom. II. p. 445. *Id. Pig. et Gall.* tom. I. p. 118. *Stock-Dove*, *Selb. Illust.* vol. I. p. 408. pl. 56*. f. 1.

DIMENS. Entire length twelve inches six lines: length of the bill (from the forehead) eight lines and a half, (from the gape) eleven lines

and a half; of the tarsus one inch; of the tail four inches; from the carpus to the end of the wing eight inches six lines.

DESCRIPT. Head and throat deep bluish gray; sides of the neck glossed with metallic hues of green and purple; back brownish gray; rump and upper tail-coverts pale bluish ash; breast pale vinaceous red; belly, thighs, and under tail-coverts, bluish gray; quills dusky, passing into bluish gray at the base of the feathers; wing-coverts of the same colour as the back, but rather paler; some of the greater ones, as well as the three last secondary quills, spotted with black, forming an irregular bar across the wings: tail bluish ash passing into black at the extremity; the outer web of the external feather with an oblong white spot towards the base: irides deep reddish brown: feet red. (*Egg*.) White: oval: long. diam. one inch six lines and a half; trans. diam. one inch two lines.

Inhabits woods with the preceding species, but is less plentiful and more local. Not uncommon in some of the midland and eastern counties, where it remains the whole year. Builds in the hollows of old pollard trees, and lays two eggs. Does not coo like the Ring-Dove, but utters a hollow rumbling note, heard at intervals throughout the spring and summer months. Flocks with that species in the Winter, and supports itself in the same manner.

139. *C. Livia*, Briss. (*Rock-Dove*.) — Bluish ash; sides of the neck glossed with green reflections; wings with two distinct transverse bars; rump white; tip of the tail black.

C. Livia, *Temm. Man. d'Orn.* tom. 11. p. 446. *Id. Pig. et Gall.* tom. 1. p. 125. *Rock-Dove*, *Selb. Illust.* vol. 1. p. 410. pl. 56*. f. 2. *Wild Pigeon*, *Bew. Brit. Birds*, vol. 1. p. 309.

DIMENS. Entire length twelve inches eight lines.

DESCRIPT. Head, throat, upper part of the back, wing-coverts, and under parts, bluish ash; sides of the neck, and upper part of the breast, glossed with shades of green and purple-red; rump, and lower part of the back, white: wings with two distinct black transverse bars; quills tipped with black: tail deep bluish gray at the base, black at the extremity; outer feather with the external web white: irides pale reddish orange: feet red. (*Egg*.) White: of a sub-oval form, and rather pointed: long. diam. one inch five lines; trans. diam. one inch two lines and a half.

Found on rocky cliffs, principally those in the neighbourhood of the sea. Met with in various parts of England, but is most abundant on the eastern coast. Breeds in caverns, and the recesses of rocks, and lays two eggs. Has two or three broods in the year. Feeds on grain, seeds, and some of the smaller *Helices*. *Obs.* The *Domestic Pigeon*, with its numerous varieties, is descended from this species.

140. *C. Turtur*, Linn. (*Turtle-Dove*.) — Back and rump cinereous brown; a black space on the sides of the neck; tip of the tail white.

C. Turtur, *Temm. Man. d'Orn.* tom. 11. p. 448. *Id. Pig. et Gall.* tom. 1. p. 305. *Turtle-Dove*, *Mont. Orn. Dict.* *Selb. Illust.* vol. 1. p. 413. pl. 56. f. 2. *Bew. Brit. Birds*, vol. 1. p. 312.

DIMENS. Entire length eleven inches three lines: length of the bill (from the forehead) eight lines, (from the gape) ten lines; of the tarsus ten lines; of the tail four inches four lines; from the carpus to the end of the wing six inches nine lines: breadth, wings extended, twenty inches.

DESCRIPT. Head and nape cinereous, with a tinge of vinaceous red; on each side of the neck a patch of black feathers tipped with grayish white; back and rump cinereous brown; inside and edge of the wings bluish ash; greater quills brownish black; secondaries bluish gray; scapulars and wing-coverts ferruginous brown inclining to rust-red, with a black spot in the middle of each feather: fore part of the neck and breast pale vinaceous; belly and under tail-coverts pure white: the two middle tail-feathers wholly brown; the rest bluish black, tipped with white; the outer one also white on the external web: bill brown: irides reddish orange: orbits and feet red. (*Egg.*) White: rather more pointed than in any of the former species: long. diam. one inch two lines and a half; trans. diam. ten lines.

Var. β. Spotted-necked Turtle. *Lath. Syn.* vol. 11. p. 645. *Don. Brit. Birds.* vol. vii. pl. 149. Characterized principally by having the whole side of the neck black, each feather having a round white spot near the extremity, instead of being tipped with white.

A migratory species, visiting this country in May, and departing at the approach of Autumn. Not generally diffused. Said to be most abundant in Kent and Buckinghamshire. Occurs sparingly in some of the eastern and western counties, but is rare northward. Frequents thick woods, and builds in trees, constructing a nest like that of the Ring-Dove, but smaller. Eggs two in number, laid about the middle of June. Utters a peculiar plaintive note during the breeding season, which is sometimes continued at intervals till near the middle of August. Food all kinds of grain and seeds. The *Spotted-necked* variety, first noticed by Dr Latham, appears to have been only met with in Buckinghamshire.

(2. ECTOPISTES, *Swains.*)

141. *C. migratoria*, Linn. (*Passenger Pigeon.*)—
Cinereous; sides of the neck glossed with green and purple; wing-coverts spotted with black.

C. migratoria, *Temm. Pig. et Gall.* tom. 1. p. 346. *Passenger Pigeon*, *Lath. Syn.* vol. 11. p. 661. (Male.) *Canada Turtle*, *Id.* vol. 11. p. 658. (Female.) *Pass. Pig., Wilson, Amer. Orn.* vol. v. p. 102. pl. 44. f. 1.

DIMENS. Entire length sixteen inches nine lines and a half: length of the bill one inch; of the tarsus one inch one line and a half: breadth, wings extended, twenty-four inches six lines. **FLEM.**

DESCRIPT. "Chin, cheeks, head, back, and rump, bluish gray; shoulders with a tinge of yellowish brown: side of the neck, and behind, rich reddish purple, iridescent: fore-neck deep chestnut, becoming paler on the breast, or rather salmon-coloured, and passing to white on the belly and vent: thighs like the breast: quills brownish black, the gray colour of the margin of the outer web increasing at the base of the secondaries, and towards the ends of the inner ones: bastard wing, and greater coverts of the primaries, brownish black; greater coverts of the secondaries gray: lesser coverts and outer scapulars tinged with yellowish brown, with black spots: tail of twelve feathers, the two middle ones

produced, the rest decreasing to the exterior: the two middle ones dusky black, the next gray, the inner margin white towards the extremity, with a black and brown spot near the base; the fourth and third gray, with the black spot; the second gray, with the black and brown spot; the outer web and tip of the first white, lower half of the inner web gray, with a black and brown spot: upper tail-coverts long, produced; lower ones white: bill black: bare space round the eyes livid: irides reddish orange: feet reddish, paler behind than before; claws black." FLEM. The *female*, according to Wilson, is somewhat smaller, with the colours in general less vivid and more tinged with brown: the gold spot on the sides of the neck smaller, and less brilliant.

A native of North America. The only individual which has hitherto occurred in this country is recorded by Dr Fleming (*Brit. An.* p. 145.) to have been "shot in the neighbourhood of a pigeon-house at Westhall, in the parish of Monymenal, Fifeshire, Dec. 31, 1825. The feathers were quite fresh and entire, like those of a wild bird."

* GEN. 53. MELEAGRIS, *Linn.*

* 142. *M. Gallopavo*, *Linn.* (*Turkey*).—Head, and upper part of the neck, almost naked, with a bluish papillose skin: a tuft of black hairs on the breast.

M. Gallopavo, *Temm. Pig. et Gall.* tom. II. p. 374. and tom. III. p. 677. *Turkey*, *Bew. Brit. Birds*, vol. I. p. 325.

DIMENS. Entire length three feet six inches: breadth, wings extended, about four feet.

DESCRIPT. Head, and upper portion of the neck, bare of feathers; skin bluish, rough with numerous fleshy papillæ of various hues and sizes; throat furnished with a pendulous carunculated wattle, of a bright scarlet colour, increasing in intensity when the bird is under excitement: also an elongated fleshy appendage arising from the base of the upper mandible, in its contracted state about an inch long, but when relaxed hanging down considerably below the bill: a tuft of long pendent hair from the middle of the breast: colour of the plumage very variable; generally dark gray, inclining to black, or black with transverse whitish bars; occasionally pure white, or pied; more rarely of a fine deep copper-colour, with the greater quills pure white, and the tail dirty white. In the *female* the pectoral tuft is wanting; the frontal caruncle is smaller, and remains always contracted; there is also no power of erecting and expanding the tail, as in the male bird. (*Egg*.) Yellowish white, spotted and speckled all over with reddish yellow: long. diam. three inches; trans. diam. one inch eleven lines.

Found in a wild state in North America. Generally supposed to have been introduced into England about the year 1524. Lays early in the Spring, and produces from fifteen to seventeen in a brood. Seldom hatches more than once in the season in this climate.

* GEN. 54. PAVO, *Linn.*

* 143. *P. cristatus*, *Linn.* (*Crested Peacock*).—Crest on the head compressed.

P. cristatus, domesticus, *Temm. Pig. et Gall.* tom. II. p. 35. and tom. III. p. 651. *Peacock*, *Bew. Brit. Birds*, vol. I. p. 328.

DIMENS. Entire length three feet eight inches: length of the bill one inch ten lines; of the train four feet five inches; of the tail one foot seven inches.

DESCRIPT. (*Male*.) Crest of twenty-four feathers, the shafts slender, and scarcely webbed except at the tips, which are of a golden green colour: head, throat, neck, and breast, of a rich blue, glossed with green and gold; above and

beneath the eye a white streak; back and rump golden green, glossed with copper, the feathers edged and tipped with velvet black; train, or upper tail-coverts, very much elongated, reaching considerably beyond the tail, capable of being erected and expanded at will; each of the feathers composing it having a conspicuous ocellated spot at the extremity; the true tail concealed beneath the train, consisting of eighteen feathers, of a grayish brown colour: scapulars and lesser wing-coverts variegated with black, and reddish cream-colour; middle coverts deep blue, glossed with golden green; greater coverts, bastard wing, and primary quills, rufous; the other quills dusky, some of them being variegated with red, and tinged with golden green: belly and sides greenish black; thighs tawny yellow: irides yellow: bill whitish: feet grayish brown. (*Female.*) Smaller; the train shorter than the tail, and without the ocellated spots; sides of the head with more white; throat of that colour; neck green; general colour of the body, and wings, cinereous brown, the feathers on the breast being tipped with white: irides lead gray. White and pied varieties sometimes occur in both sexes. (*Egg.*) Yellowish white, sparingly speckled with reddish yellow: long. diam. two inches eight lines; trans. diam. two inches one line.

A native of India, where it is still met with in an unreclaimed state. The period of its first introduction into this country, not very well ascertained. Lays from five to eight eggs. Time of incubation from twenty-seven to thirty days. The approach of the breeding season announced by the loud discordant screams of the male bird, first heard towards the end of March, and continued at intervals through the Summer. The two sexes are similar in plumage during the first year, and the train of the male does not appear till the third.

* GEN. 55. GALLUS, *Briss.*

* 144. *G. domesticus*, *Briss.* (*Domestic Cock.*)—Caruncle on the head compressed, denticulated; throat with two pendulous wattles: neck-feathers linear and elongated.

G. domesticus, *Temm. Pig. et Gall.* tom. II. p. 92. and tom. III. p. 654. The Cock, *Bew. Brit. Birds*, vol. I. p. 316.

DIMENS. Extremely variable.

DESCRIPT. Crest or comb with eight or nine serratures, of a bright coral-red; two long wattles beneath the lower mandible of the same colour; throat, and space round the eyes, naked, the skin on these parts red; beneath the ears a naked white spot: neck-feathers very much elongated, and linear throughout their whole length: tail ascending, compressed, forming two planes inclined to one another at an acute angle; of fourteen feathers, the two middle ones considerably the longest, and bending gracefully over the others: thighs strong and muscular: tarsi with long bent spurs: colours of the plumage very variable; more commonly, the head, neck, back, and wing-coverts, orange-red; under parts whitish, or velvet black; the sickle-shaped feathers of the tail blackish blue: sometimes the whole plumage pure white. The *Hen* is always smaller, with the comb and wattles less developed; the colours are also less brilliant, and the tail wants the long pendent feathers. (*Egg.*) White; varying in size and shape according to the breed.

Var. β. cristatus. (Crested Cock.) *Temm. Pig. et Gall.* tom. II. p. 239. Distinguished by having a tuft of feathers on the head instead of a comb.

Var. γ. pusillus. (Bantam Cock.) *Temm. Pig. et Gall.* tom. II. p. 242. A small variety, with the feet and toes feathered.

Var. δ. pumilio. (Dwarf Cock.) *Temm. Pig. et Gall.* tom. II. p. 244. Very small, scarcely larger than a pigeon: feet short, generally feathered.

Var. ε. pentadactylus. (Dorking Cock.) *Temm. Pig. et Gall.* tom. III. p. 658. Differs from the others in having five toes on each foot, three in front, and two behind. Generally considered as most abundant in the neighbourhood of Dorking, in Surrey.

Known in a state of domestication from the earliest times. The original stock very uncertain; referred by Temminck to the *G. Bankiva*, *Temm. (Pig. et Gall.* tom. II. p. 87.) a species met with at the present day in a wild state in the Island of Java. In the domestic varieties, the cock is polygamous. The hen is very prolific, and continues to lay during a great part of the year. Period of incubation about three weeks.

Besides the above, the following species are domesticated in some parts of England.

- * (10.) *G. Morio*, Temm. (*Negro Cock.*) Pig. et Gall. tom. 11. p. 253.

Originally from India. Remarkable for having the comb, wattles, skin, and periosteum black.

- * (11.) *G. lanatus*, Temm. (*Silk Cock.*) Pig. et Gall. tom. 11. p. 256.

A native of Japan. Has the webs of the feathers disunited, and of a very silky texture. Plumage white.

- * (12.) *G. crispus*, Briss. (*Frizzled Cock.*) Temm. Pig. et Gall. tom. 11. p. 259.

From Asia. All the feathers reflexed, and as it were curled. Smaller than the common poultry.

- * (13.) *G. ecaudatus*, Temm. (*Rumpless Cock.*) Pig. et Gall. tom. 11. p. 267.

Inhabits Ceylon. No tail or tail-coverts: wants the last dorsal vertebra.

* GEN. 56. PHASIANUS, Linn.

* 145. *P. Colchicus*, Linn. (*Common Pheasant.*)—Head and neck metallic green, glossed with blue; breast and flanks reddish orange with purple reflections, the feathers edged and tipped with violet black.

P. Colchicus, Temm. *Man. d'Orn.* tom. 11. p. 453. *Id.* Pig. et Gall. tom. 11. p. 289. *Common Pheasant*, Mont. *Orn. Dict. Selb. Illust.* vol. 1. p. 417. pl. 57.

DIMENS. Entire length three feet: length of the bill (from the forehead) one inch two lines, (from the gape) one inch four lines; of the tarsus three inches; of the tail one foot nine inches; from the carpus to the end of the wing ten inches: breadth, wings extended, two feet seven inches six lines. (*Male.*)

DESCRIPT. (*Male.*) Head and neck of a rich metallic green, passing beneath into blue and violet; on each side of the occiput a tuft of dark golden green feathers capable of being erected at will, and most conspicuous in the breeding season: a large naked space on the cheeks, thickly studded with scarlet papillæ intermixed with minute black specks: lower part of the neck, breast, belly, and sides, of a brilliant orange red, with a faint tinge of purple, each feather being edged and tipped with violet black; lower part of the abdomen, and thighs, blackish brown: scapulars, and feathers on the back, dusky brown in the middle, broadly edged with purplish orange, within which is a yellowish white band; lower part of the back, tail-coverts, and saddle-hackle feathers, exhibiting different shades of green, intermixed with orange and purple: primary and secondary quills dusky brown, with yellowish white bars: tertials and wing-coverts reddish yellow, stained and spotted with dark purple-red: tail long, and very much unequaled, of an olive-gray, or grayish brown colour, with transverse black bars, each feather being fringed with purplish red: bill pale horn-colour: irides yellowish orange: feet grayish black: spurs sharp and pointed, half an inch or more in length. (*Female.*) Smaller: general colour of the plumage yellowish brown, variegated with gray and rufous; head, neck, and upper part of the body, with the central portion of each feather black: region of the eyes feathered: tail much shorter than in the male bird, but barred in a similar manner. (*Young of the year, till after the first moult.*) Of a uniform gray colour, somewhat resembling the adult female: spur of the male short and blunt. *Varieties*, white and pied, are not unfrequent. (*Egg.*) Uniform olive-brown: long. diam. one inch ten lines; trans. diam. one inch five lines.

Var. β. Ring Pheasant. *Bew. Brit. Birds*, vol. 1. p. 321. With a collar of white round the neck. This variety, which is not uncommon, has originally proceeded from a cross between this and the following species, with which last it must not be confounded.

Var. γ. Hybrid Pheasant. (Male.) Head, neck, and breast, deep brownish black, with a slight gloss of bottle-green; the rest of the plumage above and below the body, scapulars and wing-coverts, deep red brown, the feathers edged with glossy black, and many of them streaked with yellow down the shafts: tail shorter than in the common Pheasant, compressed, and slightly arched, of a deep brown, approaching to black, variegated with specks and transverse undulating lines of ochre-yellow. (*Female.*) Head, neck, and breast, dull brown without the gloss of green; a greater portion of the upper plumage reddish or yellowish brown; many of the feathers without the black border: tail brown, barred with reddish yellow; more cuneated than in the male bird. This variety, which is occasionally met with, is an hybrid production between the *Common Pheasant* and the *Domestic Fowl*. The plumage is of course subject to considerable variation. The above descriptions were taken from a pair in the museum of the Cambridge Philosophical Society. Is sometimes called a *Pero*. The *female* has been known to breed again with the *male Game-Fowl*.

Supposed to have been brought originally from the banks of the Phasis, a river of ancient Colchis. Common at present throughout the greater part of Asia and Europe. Period of its first introduction into Britain uncertain. As a naturalized species, is generally distributed throughout England, but considered rare in Scotland. Frequents woods. Is polygamous, like the *Domestic Cock*. Commences laying in April, and hatches towards the end of May. Number of eggs from ten to fourteen, deposited on the ground, amongst long grass, or in cornfields; sometimes in thick copses. Barren hens, which have partially assumed the spurs and plumage of the cock, are not unfrequently met with, and are termed by sportsmen *Mule-Birds*.

* 146. *P. torquatus*, Temm. (*Ring-necked Pheasant.*)—Head and neck bluish green, the latter encircled by a white collar; a line of white over each eye; lower part of the neck and breast copper-red, the feathers deeply divided by a black line.

P. torquatus, Temm. *Pig. et Gall.* tom. II. p. 326. and tom. III. p. 670. *Ring Pheasant*, Leach, *Zool. Misc.* vol. II. p. 14. pl. 66. *Shaw, Gen. Zool.* vol. XI. p. 228.

DIMENS. Rather less than those of the last species.

DESCRIPT. (*Male.*) Head and upper part of the neck deep green, with a tinge of violet blue; below this colour, a collar of pure white, broadest on the sides of the neck, narrow behind, entirely interrupted in front: over each eye a narrow white line, not always present: plumage on the upper part of the back, between the shoulders, of a paler colour than in the *P. Colchicus*, approaching to orange-yellow, the feathers of a more pointed form, and deeply divided by a blackish blue line; all the lower portion of the back varied with different shades of green, passing off at the sides into bluish ash; saddle-hackle feathers pale rust-colour; upper tail-coverts the same, passing into greenish yellow; feathers on the fore part of the neck below the white collar, and those on the breast, of a fine bright copper-red, with a much narrower edging of black than in the last species, but deeply indented at the tip by a lanceolate black line: wings and tail paler than in the *Common Pheasant*. (*Female.*) General colour similar to that of the female of *C. Colchicus*; breast not so much spotted; transverse bars on the tail more distinctly marked. (*Egg.*) Bluish green, with small spots of a darker tint. TEMM.

A native of China, from whence it has been introduced into England, and naturalized in many parts of the country. In consequence, however, of its breeding freely with the *Common Pheasant**, it has become so intermixed with that species, as very rarely to occur at present exhibiting the pure plumage which characterizes it in its wild state. The above description of the cock bird was taken from a fine specimen in the possession of Mr Leadbeater.

* (14.) *Bohemian Pheasant.*

This name is employed in many parts of England to distinguish a species or variety of the Pheasant, which is met with in several preserves, but which does not appear to have received the notice of ornithologists. The

* This does not happen in its native country, where both species are equally plentiful, but, according to Temminck, keep perfectly separate.

head and neck are coloured much the same as in the *P. colchicus*, but all the rest of the plumage is of a uniform pale brownish yellow, the feathers being edged with black, and indented at the tips, as in the species last described: tail rather darker than the body, but paler than in the common Pheasant. The history of this peculiar breed, together with the origin of its name, does not appear to be well ascertained.

* GEN. 57. NUMIDA, Linn.

* 147. *N. Meleagris*, Linn. (*Guinea Pintado*).—Head and upper part of the neck naked, the skin bluish: plumage on the body bluish gray, with white spots.

N. Meleagris, Temm. *Pig. et Gall.* tom. II. p. 431. and tom. III. p. 680. *Guinea Pintado*, Lath. *Syn.* vol. II. p. 685. *Bew. Brit. Birds*, vol. I. p. 332.

DIMENS. Entire length twenty-two inches: length of the bill one inch three lines.

DESCRIPT. Head bare of feathers, and covered with a naked bluish skin; on the crown a callous conical protuberance, compressed at the sides and directed backwards, of a bluish red colour; at the base of the upper mandible, on each side, a loose pendulous wattle, bluish in the male, red in the female: upper part of the neck nearly naked, being thinly furnished with hair-like feathers, which on the nape are directed upwards; the skin bluish ash; lower part of the neck feathered, inclining to purple: general colour of the plumage on the other parts, dusky, or dark bluish gray, sprinkled all over, the breast alone excepted, with round white spots of various sizes: back very much rounded: tail short, and bent down: feet brownish red. (*Egg.*) Yellowish white ground, mottled all over with reddish yellow: long. diam. two inches; trans. diam. one inch seven lines.

Brought originally from Africa, but has long been domesticated in Britain. Is very prolific; but the young are difficult to rear. Of a restless disposition, and very clamorous.

GEN. 58. TETRAO, Linn.

(I. TETRAO, Steph.)

† 148. *T. Urogallus*, Linn. (*Wood Grouse*).—Chin and throat-feathers elongated: breast glossed with dark green: bill white: tail rounded at the extremity.

T. Urogallus, Temm. *Man. d'Orn.* tom. II. p. 457. *Id. Pig. et Gall.* tom. III. pp. 114, & 696. *Wood Grouse*, Lewin, *Brit. Birds*, vol. IV. pl. 132. *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. I. p. 335. (*Trachea*) Linn. *Trans.* vol. XVI. pl. 21. f. 1.

DIMENS. Entire length (*male*) two feet ten inches, (*female*) two feet one inch: breadth, wings extended, three feet six inches.

DESCRIPT. (*Male*.) Head and neck dusky ash, passing into black on the chin and throat where the feathers are elongated; a bare red skin above the eye, and beneath it a small spot of white feathers; breast of a fine dark glossy green; rest of the under parts black, with white spots: wings and scapulars chestnut brown, finely speckled with dusky; lower part of the back, rump, sides, and upper tail-coverts, marked with numerous small undulating lines and specks of black, upon an ash-coloured ground: tail very much rounded, black, some of the feathers having a white spot on each side near the extremity: tarsi thickly clothed with brown hair-like feathers: bill yellowish white: irides hazel. (*Female*.) Very much smaller: head, neck, and back, barred transversely with black and tawny red; throat tawny red, without spots; breast deep red, with a few white spots; belly barred like the back, some of the feathers tipped with white: quills dusky, mottled on their outer webs with light rufous brown: tail dark red with black bars; the tip white: bill dusky. (*Young male after the first moult.*) Upper plumage not so deep coloured as in the adult, inclining to cinereous gray; green on the breast more dull; a few red feathers shewing themselves in different parts of the body; tail often tipped with

white. *Before the first moult*, both sexes resemble the adult female. (*Egg.*) Light reddish yellow brown, spotted all over with two shades of darker brown: long, diam. two inches three lines; trans. diam. one inch eight lines.

Formerly abundant in the mountainous forests of Scotland and Ireland, but now extirpated. The last specimen killed in the former country is said to have been shot near Inverness, about fifty years ago. Ceased to exist in Ireland at a considerably earlier period. Occurs plentifully at the present day in many parts of the Continent, and is much attached to pines, birch, and juniper. Said to feed on the berries of the latter, and on the buds and tender sprays of the two former. The males are polygamous. The females build on the ground, and lay from six to sixteen eggs.

- † (15.) *T. medius*, Meyer. (*Hybrid Grouse.*) Temm. Man. d'Orn. tom. 11. p. 459. Id. Pig. et Gall. tom. 111. pp. 129, & 698. *Spurious Grouse*, Lath. Syn. Supp. p. 214.

Said to have been formerly found in Scotland, on the authority of Mr Tunstall; but there is no existing evidence by which the truth of this assertion can be proved. Is principally distinguished from the last species by having the neck and breast of a rich bronzed purple hue, the bill black, and the tail slightly forked, with the outermost feathers the longest. The *egg*, which is figured by Klein, is very similar in colour to that of the *Wood Grouse*, but in size a little smaller.

149. *T. Tetrix*, Linn. (*Black Grouse.*) — Throat-feathers not elongated: the general plumage violet-black, with a white bar on the wings: tail very much forked; the lateral feathers bending outwards.

T. Tetrix, Temm. Man. d'Orn. tom. 11. p. 460. Id. Pig. et Gall. tom. 111. pp. 140, & 699. *Black Grouse*, Mont. Orn. Dict. Selb. Illust. vol. 1. p. 423. pls. 58, & 58*. Bew. Brit. Birds, vol. 1. p. 338.

DIMENS. Entire length twenty-three inches: breadth, wings extended, three feet.

DESCRIPT. (*Male.*) Head, neck, breast, back, and rump, black, with purple and blue reflections: eyebrows naked, vermilion-red: beneath the eye a white spot: abdomen, wing-coverts, and tail, deep black. secondary quills tipped with white, forming, with the adjacent coverts, a broad bar of that colour across each wing: under tail-coverts pure white: the lateral feathers of the tail much longer than the middle ones, curling outwards: tarsi clothed with blackish gray hair-like feathers: bill black: irides bluish. (*Female.*) Smaller; general colour of the upper plumage ferruginous yellow, barred and mottled with black: greater wing-coverts tipped with white: breast orange-brown, with black bars: belly dusky brown, with whitish and red bars: tail very slightly forked, variegated with ferruginous and black; the tip grayish white. *The young of the year, till after the first moult*, resemble the adult female. (*Egg.*) Yellowish white, spotted and speckled with orange-brown: long. diam. two inches; trans. diam. one inch five lines.

Most abundant in Scotland, and the northern parts of England. Occurs more or less sparingly in some parts of North Wales, as well as in the counties of Stafford, Somerset, Devon, Sussex, and Hants. Partial to woody, heathy, and mountainous situations. Feeds on berries, and on the tops of heath and birch. Is polygamous. Nest placed on the ground, generally under the shelter of a low bush, composed of a few dried stems of grass. Eggs from six to ten in number, laid in May. *Obs.* The *Hybrid Bird* figured in White's *Nat. Hist. of Selborne*, is probably a young male of this species, having nearly completed the first moult.

(2. LAGOPUS, *Vieill.*)

150. *T. Scoticus*, Temm. (*Red Ptarmigan.*)—The entire plumage chestnut-brown, variegated with black: tail of sixteen feathers; the six outer ones on each side dusky.

T. Seotieus, Temm. *Man. d'Orn.* tom. 11. p. 465. *Red Grouse*, Mont. *Orn. Dict. Selb. Illust.* vol. 1. p. 427. pl. 59. f. 1. *Bew. Brit. Birds*, vol. 1. p. 341.

DIMENS. Entire length sixteen inches: length of the bill (from the forehead) nine lines and a half, (from the gape) eleven lines; of the tarsus one inch five lines; of the tail four inches; from the carpus to the end of the wing eight inches three lines.

DESCRIPT. (*Male.*) The whole plumage of a deep chestnut-brown, nearly plain on the head and neck, but marked on the back and wing-coverts with black spots of different sizes, and beneath the body with numerous undulating black lines: orbits, and a small patch at the base of the lower mandible, white: naked space above the eyes fringed, of a bright scarlet colour: frequently some of the feathers on the abdomen tipped with white: tail of sixteen feathers; the four middle ones reddish brown, with transverse black lines; the rest of a uniform dusky brown: bill black, half concealed by the nasal feathers: irides chestnut-brown: tarsi and toes thickly clothed with grayish white feathers; claws light horn-colour. (*Female.*) Colours not so dark as in the male; the brown varied with reddish yellow, and marked with a greater number of black spots and lines: naked space above the eyes less conspicuous. (*Egg.*) Reddish white, nearly covered with blotches and spots of umber-brown: long, diam. one inch nine lines; trans. diam. one inch three lines.

Peculiar to the British Islands. Found plentifully in Scotland, as well as in some of the mountainous parts of England and Wales. Frequents moors, heaths, and extensive uncultivated wastes: never resorts to woods. Feeds on berries, and the tender tops of heaths. Is monogamous, and pairs in January. Commences laying in March or April. Eggs eight to twelve in number, deposited on the ground.

151. *T. Lagopus*, Sab. (*Common Ptarmigan.*)—Cinereous, with transverse undulating lines and spots of black, (*summer*); or wholly white, (*winter*); shafts of the quills, and lateral tail-feathers, always black.

T. Lagopus, Sab. *Supp. Parry's First Voy.* p. exxvii. *Richards. App. Parry's Second Voy.* p. 350. *Ptarmigan*, Mont. *Orn. Dict. Selb. Illust.* vol. 1. p. 430. pl. 59. f. 2. & pl. 60. *Shaw, Gen. Zool.* vol. xi. p. 287. pl. 21. *Bew. Brit. Birds*, vol. 1. p. 343.

DIMENS. Entire length fifteen inches: length of the bill (from the forehead) eight lines, (from the gape) ten lines: of the tarsus one inch four lines; of the tail four inches three lines: of the wing seven inches six lines: breadth, wings extended, twenty-four inches six lines.

DESCRIPT. (*Male in winter plumage.*) A streak from the corner of the bill across the eyes, lateral tail-feathers, and shafts of the quills, black: the rest of the plumage pure white: above the eyes a scarlet fringed membrane: tarsi and toes thickly clothed with woolly feathers:

bill and claws black. In the *female*, the naked membrane above the eyes is less conspicuous, and the black on the lore altogether wanting. (*Male in summer plumage*.) All the upper parts of the body, scapulars, tertiaries, neck, breast, and sides under the wings, cinereous brown, with transverse undulating black lines, and minute dusky spots: a few reddish orange bars on the head and neck only: primary and secondary quills, (with the exception of the black shafts), greater part of the wing-coverts, belly, under tail-coverts, and legs, white: two middle tail-feathers nearly all white; the others black, some of them slightly tipped with white. The *female in summer* does not show so much of the rufous tint on the head and neck. (*Egg*.) Yellowish white, sparingly blotched and spotted with black brown: long. diam. one inch eight lines; trans. diam. one inch two lines.

Found only in the Highlands of Scotland, and the adjacent isles. Was formerly met with in some of the mountainous parts of Cumberland and Westmoreland, but is supposed now to be extinct in England. Frequents the summits of the loftiest hills, from whence it rarely descends into the plains. Feeds on the berries and tender shoots of alpine plants. Pairs early in the Spring, and lays its eggs on the bare ground, from eight to fourteen in number. *Obs.* This species is also met with in North America; but it is not found on the Continent, where it has been always confounded with the following.

152. *T. rupestris*, Sab. (*Rock Ptarmigan*.)—Brownish yellow, with transverse black bars, (*summer*); or pure white, (*winter*); shafts of the quills, and lateral tail-feathers, always black.

T. rupestris, Sab. *Supp. Parry's First Voy.* p. cxv. *Richards. App. Parry's Second Voy.* p. 348. *Faun. Bor. Amer.* pt. 2. p. 354. pl. 64. *T. Lagopus*, *Temm. Man. d'Orn.* tom. II. p. 468. *Rock Ptarmigan*, *Shaw, Gen. Zool.* vol. XI. p. 290.

DIMENS. Entire length fourteen inches: length of the bill (from the forehead) seven lines, (from the gape) one inch; of the tarsus one inch four lines and a half; of the tail four inches; of the wing seven inches. *Faun. Bor. Am.*

DESCRIPT. Smaller than the last species, with the bill longer and narrower. *Winter plumage* exactly similar. *Summer plumage* characterized by its brownish yellow colour, with rather broad blackish brown bars, exhibiting none of the cinereous tint which predominates in the *Common Ptarmigan*: on the *upper* parts the black markings prevail over the yellow; on the *under* the yellow ground is most conspicuous: primary and secondary quills, with some of the coverts, white; the shafts of the quills black: tail with the two middle feathers barred like the back; the rest black, faintly tipped with white. The *male* is at all seasons distinguished from the *female* by a black band across the eye, as in the last species: in its *summer plumage*, it is furthermore characterized by having the middle of the belly white. (*Egg*.) Buffly white, nearly covered with spots of two shades of dark red brown: long. diam. one inch nine lines; trans. diam. one inch two lines.

This species, originally described by Pennant, (*Arct. Zool.* vol. II. p. 364. no. 184.) occurs in North America, and very plentifully in some parts of the Continent, but has been confounded by Temminck and others

with the Common Ptarmigan. In Great Britain, it has hitherto only occurred once. This specimen was killed in Perthshire in Scotland, and is now in the collection of Lord Derby. Is said to frequent dry rocky grounds, and to feed on the tops of small birch. Hatches in June.

GEN. 59. PERDIX, *Briss.*

(1. PERDIX, *Steph.*)

153. *P. cinerea*, *Briss.* (*Common Partridge.*)—Upper parts cinereous, variegated with brown and black; a deep chestnut crescent-shaped spot on the breast.

P. cinerea, *Temm. Man. d'Orn.* tom. ii. p. 488. *Id. Pig. et Gall.* tom. iii. pp. 373, & 728. *Common Partridge*, *Mont. Orn. Dict. Selb. Illust.* vol. i. p. 433. pl. 61. *Bew. Brit. Birds*, vol. i. p. 348.

DIMENS. Entire length thirteen inches: length of the bill (from the forehead) seven lines and a half, (from the gape) eleven lines; of the tarsus one inch nine lines; of the tail three inches four lines; from the carpus to the end of the wing six inches two lines: breadth, wings extended, twenty inches six lines.

DESCRIPT. (*Male.*) Sides of the face, throat, and eyebrows, bright rust-colour: behind the eye a naked red skin: neck and breast bluish gray with fine zigzag black lines; on the lower part of the breast a large patch of deep chestnut-brown in the shape of a horse-shoe; flanks cinereous, with undulating black lines, and a large rust-coloured bar towards the tip of each feather: back, rump, and upper tail-coverts, cinereous brown, with transverse zigzag black lines and a few narrow bars of reddish brown; scapulars and wing-coverts of a deeper hue than the back, with the shafts of the feathers yellowish white; quills blackish gray, spotted and barred with pale yellowish red: tail of eighteen feathers; the four middle ones marked like the back; the others bright rust-colour: bill and feet bluish gray: irides hazel. (*Female.*) Less of the rust-coloured tinge on the head and throat; feathers on the crown of the head spotted with white: upper plumage generally darker, with a greater number of black bars and spots: the horse-shoe mark on the breast very indistinct, or altogether wanting. White and pied *varieties* are occasionally met with. (*Egg.*) Of a uniform olive-brown: long. diam. one inch five lines; trans. diam. one inch and half a line.

Abundant in all the cultivated parts of Great Britain, but seldom found at a distance from arable land, which is its favourite haunt. Feeds on seeds and insects, and especially on the pupæ of ants. Pairs in February, and commences laying about the middle or end of May. Eggs twelve to twenty in number, deposited on the ground, amongst brush-wood and long grass, or in fields of clover and standing corn. Period of incubation three weeks. After the young are hatched, they flock together in *coveys* till the following Spring.

* 154. *P. rubra*, *Briss.* (*Red-legged Partridge.*)—Upper parts reddish ash: throat and cheeks white; bounded by a collar of black, expanding on the breast, and spotted with white.

P. rubra, *Temm. Man. d'Orn.* tom. ii. p. 435. *Id. Pig. et Gall.* tom. iii. pp. 361, and 726. *Guernsey Partridge*, *Mont. Orn. Dict.* *Lewin, Brit. Birds*, vol. iv. pl. 137. *Red-legged Partridge*, *Bew. Brit. Birds*, vol. i. p. 345.

DIMENS. Entire length thirteen inches six lines: length of the bill (from the forehead) eight lines, (from the gape) eleven lines; of the tarsus one inch seven lines; of the tail four inches one line; from the carpus to the end of the wing six inches six lines: breadth, wings extended, twenty inches ten lines.

DESCRIPT. Forehead cinereous; crown and occiput reddish brown; throat, cheeks, and a broad band above the eyes reaching nearly to the nape, white; the white space on the throat bounded by a black collar, which spreads itself out on the upper part of the breast and sides of the neck, in the form of black spots on a light reddish ground; intermixed with the black spots are a few white ones; lower part of the breast bluish ash; belly and abdomen rust-red; feathers on the sides cinereous, each with two transverse bars across the middle, the first white and the second black, beyond which last is a large semilunar rust-coloured spot, occupying the tip: all the upper parts of the body reddish ash: quills grayish brown, with the outer webs ochre-yellow: the four middle tail-feathers like the back; the others rust-colour: orbits, bill, and feet, bright red. In the *female* the colours are paler, but in other respects similar to those of the male. (*Egg.*) Reddish yellow white, spotted and speckled with reddish brown: long. diam. one inch seven lines and a half; trans. diam. one inch three lines.

A native of France, Italy, and the Islands of Guernsey and Jersey, from whence it has been imported into England, and naturalized in some parts of the country*. Is very common in Suffolk, frequenting waste heaths and extensive barren lands, more than cornfields. Occasionally met with in other counties. On the Continent, is said to lay from fifteen to eighteen eggs.

* (2. ORTYX, *Steph.*)

* 155. *P. Virginiana*, Lath. (*Virginian Partridge*).—Body above brownish chestnut, variegated with black and rufous; beneath whitish, with transverse undulating black lines: throat, and a broad band above the eye, white.

Tetrao Virginianus, *Wils. Amer. Orn.* vol. VI. p. 21. pl. 47. f. 2. *Perdix borealis*, *Temm. Pig. et Gall.* tom. III. pp. 436, and 735. *Virginia and Maryland Partridge*, *Lath. Syn.* vol. II. pp. 777, and 778. *Northern Colin*, *Shaw, Gen. Zool.* vol. XI. p. 377.

DIMENS. Entire length nine inches: length of the bill six lines and a half; of the tarsus one inch three lines.

DESCRIPT. (*Male.*) Forehead black; a broad white band above the eye, reaching to the nape, above which is another narrow one of black; throat white, bounded by a black collar, which expands over the front and sides of the neck in the form of black spots, mixed with others white and red: crown of the head chestnut-red, spotted with black; upper part of the body brownish chestnut; the edges of the feathers cinereous, with fine black streaks; in the middle of the back some large black spots edged with rufous: scapulars and greater wing-coverts spotted with black and red on their inner webs, with cinereous and red on their outer; lesser coverts red with fine undulating black lines: breast reddish white, with transverse black lines; belly pure white, with black lines of a semicircular form; feathers on the sides rufous, with an edging of white spots surrounded with black; under tail-coverts red, spotted with black; tail bluish ash; the middle feathers rufous towards their extremities, and marked with brown undulating lines: bill black, reddish at the base: feet brownish red. (*Female.*) Throat, and band above the eyes, pale red, the latter without the black streak above; the red on the throat surrounded by spots of black, brown, and white: nape, and upper part of the head, spotted with red: feathers on the back more deeply edged with cinereous; rest of the upper parts paler than in the male bird: breast bright red, with two little white spots towards the extremity of each feather: tail bluish ash, with fine undulating lines of brown and whitish towards the tips of all the feathers: bill with more red at the base than in the male. The *young of the year* very much resemble the adult female, but the transverse undulating lines on the back and tail-feathers are more numerous. (*Egg.*) White: of a pointed form: long. diam. one inch two lines and a half; trans. diam. one inch.

* Pulteney observes that this species has been killed near Weymouth in Dorsetshire, and suggests the probability of its sometimes reaching this country from the Islands of Jersey and Guernsey. If this be ever the case, which is rather doubtful, it has a claim to be considered as one of our *native* species.

A native of North America, from whence it has been recently introduced into this country. Is now naturalized in Suffolk. Said to frequent woody situations, and to perch in trees. Lays from twenty to twenty-five eggs, and has two broods in the year.

(3. COTURNIX, *Briss.*)

156. *P. Coturnix*, Lath. (*Common Quail*).—Upper parts cinereous brown, variegated with black: over each eye, and on the crown of the head, a longitudinal whitish streak.

P. Coturnix, *Temm. Man. d'Orn.* tom. II. p. 491. *Coturnix dactylisonans*, *Temm. Pig. et Gall.* tom. III. pp. 478, & 740. *Common Quail*, *Selb. Illust.* vol. 1. p. 437. pl. 62. *Quail*, *Mont. Orn. Dict. and Supp.* *Bew. Brit. Birds*, vol. 1. p. 351.

DIMENS. Entire length eight inches: length of the bill (from the forehead) five lines, (from the gape) seven lines and a half; of the tarsus one inch two lines; of the tail one inch six lines; from the carpus to the end of the wing four inches six lines: breadth, wings extended, fourteen inches four lines.

DESCRIPT. (*Male*.) Crown and occiput black, the feathers edged with rufous brown; over each eye a yellowish white streak extending to a considerable distance down the neck; a similar streak along the top of the head running parallel with the above; throat rufous, bounded by a double crescent of dusky brown: upper part of the neck, back, scapulars, and wing-coverts, cinereous brown, variegated with black; the shafts and central portions of the feathers yellowish white: lower part of the neck, breast, and sides, pale rufous brown with longitudinal white streaks; belly yellowish white: tail blackish brown; the shafts and tips of the feathers whitish: bill and feet yellowish brown. (*Female*.) Throat white, without the double crescent-shaped markings: breast yellowish white, spotted with blackish brown: general colours of the plumage paler. *Obs.* This species is subject to some variation of colours: in *very old males*, the cheeks and throat are dusky brown, and the whole plumage of a more vivid hue. Occasionally found pure white; or of a uniform deep brown. (*Egg*.) Yellowish white, blotched and spotted with umber-brown: long. diam. one inch one line; trans. diam. eleven lines.

A migratory species visiting this country in May, and usually departing towards the end of the Autumn. Occasionally remains through the Winter. Chiefly frequents corn-fields and open lands. Feeds on seeds, grain, and insects. The males are polygamous, and utter during the breeding season a peculiar whistling note. Eggs eight or ten in number, deposited on the bare ground.

GEN. 60. OTIS, *Linn.*

157. *O. Tarda*, Linn. (*Great Bustard*).—Upper part of the body reddish orange, with transverse black bars; beneath whitish: a tuft of long filiform feathers (*male*) from the corners of the lower mandible.

O. Tarda, *Temm. Man. d'Orn.* tom. II. p. 506. Great Bustard,
Mont. Orn. Dict. and Supp. Selb. Illust. vol. I. p. 442. pls. 64,
 & 64*. *Bew. Brit. Birds*, vol. I. p. 355.

DIMENS. Entire length nearly four feet: breadth nine feet. PENN.

DESCRIPT. (*Male*.) Head and neck cinereous; a longitudinal black streak on the crown reaching to the occiput; feathers on the chin loose, and somewhat elongated, with the barbs disunited; a long moustache of similar feathers on each side of the lower mandible;* back, scapulars, lesser wing-coverts, rump, and tail-coverts, yellowish or reddish orange, barred and variegated with black; greater wing-coverts, and some of the secondaries, pale cinereous; primaries black, slightly tipped with white; upper part of the breast reddish orange; rest of the under parts white; tail brownish orange, some of the lateral feathers with the basal half and tip white; the whole crossed by two or three black bars near the extremity: bill dusky, the under mandible somewhat paler than the upper: irides light hazel; feet dusky brown. (*Female*.) Much smaller, and without the tuft of long feathers on each side of the lower mandible; the longitudinal streak on the crown not so obvious; fore part of the neck of a deeper gray: in other respects similar to the male. (*Egg*.) Olive-brown, sparingly and indistinctly blotched with greenish brocoli-brown: long. diam. two inches eleven lines; trans. diam. two inches two lines.

Formerly met with in great plenty on the plains of Wiltshire and Dorsetshire, the Wolds of Yorkshire, as well as in many other parts of England, and also in Scotland. Is now become extremely rare, and almost confined to the open parts of Norfolk and Suffolk, where the species still continues to breed in small quantities. Single individuals are occasionally observed in Cambridgeshire, and a fine male specimen was killed near Ickleton in that county in January 1831. Where plentiful, usually found in flocks. Food green corn, seeds, insects, as well as various other vegetable and animal substances. Lays two eggs early in the Spring, which are deposited on the bare ground; rarely, however, hatches more than one in this country. Period of incubation about four weeks. The male of this species possesses a capacious pouch, situate along the fore part of the neck, said to be capable of holding several quarts of water.

158. O. *Tetrax*, Linn. (*Little Bustard*.) — Upper part of the body yellowish orange, with fine zigzag dusky lines; neck black, encircled by a double collar of white.

O. *Tetrax*, *Temm. Man. d'Orn.* tom. II. p. 507. Little Bustard,
Mont. Orn. Dict. and Supp. Selb. Illust. vol. I. p. 447. pl. 65.
Bew. Brit. Birds, vol. I. p. 359.

DIMENS. Entire length seventeen inches and a half: breadth, wings extended, two feet ten inches and a half. SELBY.

DESCRIPT. (*Adult male*.) Crown and occiput yellowish orange, with brown spots: sides of the head and throat deep cinereous, bounded by a white collar encircling the upper part of the neck: lower part of the neck black; on the breast another, and somewhat broader, white collar, below which is a narrow one of black: upper parts yellowish orange, crossed by numerous fine zigzag dusky lines, and mottled with large spots more

* Bewick's figure represents this species with a longitudinal black streak or band on the sides of the neck. It is possible that such a character may exist during the short period of the breeding season only, but it is not usually present.

sparingly distributed: the edge of the wing, belly, and upper tail-coverts, pure white: irides orange: bill and feet gray. (*Female and young male.*) Throat white; sides of the head, and upper part of the neck, yellowish orange, with longitudinal dusky streaks; lower part of the neck and upper part of the breast the same, with transverse crescent-shaped dusky bars; lower breast and flanks whitish, spotted with dusky, the latter with a few fine longitudinal streaks besides on the shafts of the feathers: belly and abdomen white and unspotted: upper parts of the body much as in the male. *Obs.* It is not certain whether the *adult male* does not lose the black neck and double white collar in Winter, and resemble the female during that season. (*Egg.*) Of a uniform olive-brown: long. diam. two inches; trans. diam. one inch six lines.

A very rare, and only occasional, visitant in this country. Has been taken alive on the edge of Newmarket Heath, and more recently near Caxton in Cambridgeshire. Other specimens have been killed in Sussex, Kent, Devonshire, Northumberland, and Suffolk. Frequents plains, and large tracts of open country. Said to be graminivorous. Eggs three to five in number, deposited on the ground.

ORDER IV. GRALLATORES.

GEN. 61. CURSORIUS, *Lath.*

159. *C. isabellinus*, Meyer. (*Cream-coloured Courser.*)—General plumage reddish cream-colour: behind the eyes a double black streak.

C. isabellinus, *Temm. Man. d'Orn.* tom. II. p. 513. Cream-coloured Plover, *Lath. Syn.* vol. III. p. 217. *Id. Supp.* p. 254. pl. 116. *Mont. Orn. Dict.* Cream-coloured Swiftfoot, *Selb. Illust.* vol. II. p. 217. pl. 33**.

DIMENS. Entire length ten inches: length of the bill nine lines.
MONT.

DESCRIPT. General colour of the plumage reddish cream or buff orange, the feathers in some places with pale edges; throat whitish; behind the eye a double black streak directed towards the occiput: quills black: lateral tail-feathers black towards their extremities, the black including a small white spot: abdomen whitish: bill black: legs yellowish brown. In *immature birds*, the scapulars and wing-coverts are crossed by numerous, fine, zigzag bars of a darker tint, more particularly towards the tips of the feathers: the black streak behind the eyes very faint. (*Egg.*) Unknown.

A native of Africa, very rarely occurring in Europe. In England only three specimens have been hitherto observed. The first of these was shot near St Albans, in Kent, many years back; the second was killed in October 1827, at Charnwood Forest, Leicestershire; the third in the year following, at Freston, near Aldborough, in Suffolk. Very little is known of its habits. Said to inhabit dry plains, and to run very swiftly.

GEN. 62. *ŒDICNEMUS*, Temm.

160. *Œ. crepitans*, Temm. (*Common Thick-knee*).—Reddish ash, with longitudinal dusky spots: quills black; the first two primaries with a broad white bar across each web.

Œ. crepitans, Temm. *Man. d'Orn.* tom. II. p. 521. Common Thick-knee, *Selb. Illust.* vol. II. p. 250. pl. 40. Great Plover, *Bew. Brit. Birds*, vol. I. p. 362. Thick-kneed Bustard, *Mont. Orn. Dict.*

DIMENS. Entire length eighteen inches.

DESCRIPT. Upper parts reddish ash, with a longitudinal dusky spot down the middle of each feather: space between the eye and the bill, cheeks, throat, belly, and thighs, white: neck and breast tinged with reddish, and marked with fine longitudinal dusky streaks: a pale bar across the wing-coverts: quills black; the first with a large and conspicuous white spot near the middle; the second with one somewhat smaller: under tail-coverts red: all the tail-feathers, the two middle ones excepted, tipped with black: bill yellowish at the base, black at the tip: irides, orbits, and feet, yellow. In *young birds*, the markings are less distinct. (*Egg*.) Stone-colour, blotched, spotted, and streaked, with ash-blue and dark brown: long. diam. two inches two lines; trans. diam. one inch seven lines.

A migratory species visiting this country about the latter end of April or beginning of May, and departing in the Autumn. Is most abundant in the southern, midland, and eastern counties. Frequents heaths, extensive corn-lands, and other open districts. Feeds on insects, worms, and reptiles. Makes no nest, but lays its eggs, which are two in number, on the bare ground. During the breeding season the male utters a loud shrill cry, heard more particularly in the dusk of the evening. The name of this species is derived from a peculiar enlargement of the upper part of the tarsus, and of the joint immediately above it, most conspicuous in the young birds of the year.

GEN. 63. *CHARADRIUS*, Linn.

161. *C. pluvialis*, Linn. (*Golden Plover*).—Upper parts dusky brown, spotted with yellow; cheeks, neck, and breast, variegated with brown and ash-colour; throat and abdomen white.

C. pluvialis, Temm. *Man. d'Orn.* tom. II. p. 535. Golden Plover, *Mont. Orn. Dict.* *Selb. Illust.* vol. II. p. 231. pl. 37. *Bew. Brit. Birds*, vol. I. p. 367.

DIMENS. Entire length eleven inches: length of the bill (from the forehead) ten lines and a half, (from the gape) one inch one line; of the tarsus one inch seven lines; of the tail three inches; from the carpus to the end of the wing seven inches six lines: breadth, wings extended, twenty-three inches.

DESCRIPT. (*Winter plumage*.) Crown of the head, and all the upper parts, dusky brown, thickly spotted with king's yellow, the spots being disposed along the edges and at the tips of the feathers; cheeks, neck,

and breast, somewhat variegated with streaks and spots of ash-gray and yellowish brown; throat, belly, and abdomen, white; quills, and greater coverts, dusky; the former white along the shafts; the latter tipped with white: bill dusky: irides dark brown: legs deep ash-colour. (*Summer plumage.*) Upper parts deep black; the edges of the feathers spotted with bright yellow: forehead, and space above the eyes, white: sides of the neck white, with large black and yellow spots: throat, fore part of the neck, and all the under parts, black. (*Young of the year.*) Upper parts cinereous brown, with spots of yellowish ash. (*Egg.*) Yellowish stone-colour, blotched and spotted with brownish black: long. diam. two inches; trans. diam. one inch four lines.

Not uncommon; migrating from the southern to the northern parts of the kingdom at the approach of the breeding season. Haunts moors, heaths, and other open districts; occasionally, during severe weather, the sea-coast. Food worms and insects. Eggs four in number; laid in May.

162. *C. Morinellus*, Linn. (*Dotterel Plover.*)—Upper plumage brownish ash, edged with reddish: breast reddish ash: a broad streak above the eyes, and a gorget on the breast, white.

C. Morinellus, *Temm. Man. d'Orn.* tom. 11. p. 537. *Dotterel*, *Mont. Orn. Dict. Selb. Illust.* vol. 11. p. 236. pl. 39. *Bew. Brit. Birds*, vol. 1. p. 369.

DIMENS. Entire length nine inches five lines: length of the bill (from the forehead) seven lines and a half, (from the gape) one inch; of the tarsus one inch six lines; from the carpus to the end of the wing six inches.

DESCRIPT. (*Winter plumage.*) Forehead, throat, and sides of the face, whitish, dotted and streaked with black; above each eye a broad band of reddish white, reaching to the nape: crown of the head, and occiput, dusky ash; rest of the upper parts cinereous, tinged with olivaceous brown; the feathers on the back, wing-coverts, and scapulars, edged with rust-red: fore part of the neck, breast, and flanks, reddish ash; a gorget on the upper part of the breast extending on each side to the bend of the wing, and middle of the belly, pure white: shaft of the first primary white; the rest dusky: tail cinereous, passing into dusky towards the end; the extreme tip white: bill black: irides brown: feet greenish ash. (*Male in summer plumage.*) Cheeks, throat, and band above the eyes, pure white; crown of the head, and occiput, deep brown; back, wing-coverts, and scapulars, olivaceous ash, the feathers edged with deep red; nape, also the sides and lower part of the neck in front, cinereous; on the upper part of the breast a white gorget bounded above by a blackish line; lower part of the breast, and flanks, bright rust-colour; middle of the belly deep black; vent and under tail-coverts reddish white. In the *female*, the white band above the eyes is less conspicuous; the gorget on the breast has a reddish tinge; the black patch on the abdomen is smaller, and much intermixed with white. In *young birds*, the crown of the head is reddish, with longitudinal dusky spots; the upper parts are more cinereous; and the tail is tipped with pale red. (*Egg.*) Yellowish olive, blotched and spotted with dark brownish black: long. diam. one inch seven lines and a half; trans. diam. one inch two lines and a half.

Common in some parts of England, appearing twice in the year, Spring and Autumn, in its passage to and from more northern latitudes, in which it breeds. Winters in the South of Europe. With us generally observed in small flocks on heaths, moors, and other open districts. Feeds on insects and worms. This and the preceding species are in high estimation for the table.

163. *C. Hiaticula*, Linn. (*Ringed Plover*.) — Upper parts cinereous brown: a broad gorget of black on the breast: bill orange and black: feet orange.

C. Hiaticula, *Temm. Man. d'Orn.* tom. II. p. 539. Ringed Plover, *Mont. Orn. Dict. Selb. Illust.* vol. II. p. 240. pl. 38. f. 1, 2. *Bew. Brit. Birds*, vol. I. p. 371.

DIMENS. Entire length seven inches six lines: length of the bill (from the forehead) seven lines, (from the gape) seven lines and a half; of the tarsus eleven lines and a half; of the tail two inches five lines; from the carpus to the end of the wing five inches: breadth, wings extended, sixteen inches.

DESCRIPT. (*Adult in Summer and Winter*.) Forehead, space between the eye and the bill, and sides of the face, black; across the forehead, and through the eyes, a white band, passing backwards to the occiput: on the upper part of the breast a broad gorget of black, the ends of which unite on the nape becoming narrower: throat, a collar round the neck immediately above the black gorget, and the rest of the under parts, pure white: crown of the head, and all the upper parts, cinereous brown: quills dusky, with an oval white spot about the middle of each feather; shafts partly white: outermost tail-feather wholly white; the next white, with a small brown spot on the inner web; the rest dusky, tipped with white, the two middle ones excepted, which are dusky throughout: bill orange at the base, black at the tip: irides hazel: feet yellowish orange. (*Adult female*.) Lore and cheeks cinereous brown; forehead white, surmounted by a narrow transverse band of dusky ash; streak through the eye faint; gorget dusky: in other respects similar to the male. (*Young of the year*.) Between the bill and the eye a dusky streak; forehead dirty white, without the coronal black band: upper parts cinereous brown, the feathers edged with yellowish white: gorget on the breast cinereous brown: bill dusky: feet yellowish. (*Egg*.) Stone-colour, spotted and streaked with ash-blue and black: long. diam. one inch five lines; trans. diam. one inch and half a line.

Common on most parts of the sea-coast, resorting occasionally to inland marshes and the banks of rivers. Remains with us the whole year. Food insects and worms. Pairs in May, and towards the end of that month deposits its eggs, which are four in number, in a small cavity in the sand, just above high-water mark. In the Autumn becomes gregarious, and keeps in small flocks throughout the Winter.

164. *C. minor*, Meyer. (*Little Ringed Plover*.) — Upper parts cinereous brown: a gorget of black on the breast: bill entirely black: feet flesh-colour.

C. minor, *Temm. Man. d'Orn.* tom. II. p. 542. Little Ring-Dotterel, *Gould, Europ. Birds*, part XI. pl. 2.

DIMENS. Entire length five inches: length of the bill (from the forehead) five lines and a half; of the tarsus eleven lines and a half; of the tail two inches four lines; from the carpus to the end of the wing four inches four lines.

DESCRIPT. (*Adult male and female.*) "Bill black; a band of the same colour passing from the bill to the eye, and extending over the ear-coverts; the forehead pure white, above which on the crown a black band passes from eye to eye; the occiput gray, beneath which a white circle spreads from the throat round the neck; this is succeeded by a black band, broad on the chest, but narrowing until it meets at the back of the neck; the whole of the upper plumage, with the exception of the rump, which is white, of a fine brownish gray; under surface white; feet and legs flesh-colour; irides hazel. (*Young.*) Wants the black collar and facial markings, the crown of the head and face being brownish gray; in every other respect resembles the adult, except that a brownish tint pervades the whole of the upper plumage, and that every feather is edged with a lighter margin. (*Egg.*) Yellowish white, marked with blotches of black and brown." GOULD.

A single individual of this species, a young bird of the year, has been recently killed at Shoreham in Sussex. It is now in the possession of Mr Henry Doubleday of Epping. No other has hitherto occurred in Britain, although from its close resemblance to the *C. Hiaticula*, it is possible that the species may have been overlooked. Common on some parts of the Continent. According to Temminck, resorts by preference more to the banks of rivers than to the shores of the sea. Food and nidification similar to those of the Ringed Plover.

165. *C. Cantianus*, Lath. (*Kentish Plover.*)—Upper parts cinereous brown: a large patch of black on each side of the breast: bill and feet black.

C. Cantianus, Temm. *Man. d'Orn.* tom. II. p. 544. *Kentish Plover*, Lewin, *Brit. Birds*, vol. v. pl. 85. *Lath. Syn. Supp.* II. p. 316. *Mont. Orn. Dict. Supp. Selb. Illust.* vol. II. p. 243. pl. 38. f. 3.

DIMENS. Entire length six inches ten lines: length of the bill (from the forehead) seven lines and a half, (from the gape) nine lines; of the tarsus one inch and half a line; of the tail two inches one line; from the carpus to the end of the wing four inches three lines: breadth, wings extended, thirteen inches four lines.

DESCRIPT. (*Male.*) Forehead, a broad streak over each eye, a ring round the neck, and all the under parts, pure white: space between the eye and the bill, and ear-coverts, black; an angular black patch immediately behind the forehead, advancing towards the crown; a larger patch of the same colour on each side of the breast: crown of the head, and nape of the neck, light brownish red; rest of the upper parts cinereous brown: shafts of the primaries white: the two outermost tail-feathers white; the next whitish; the rest cinereous brown: bill and feet black. (*Female.*) The white on the forehead of less extent; behind it a small transverse black streak instead of the large angular spot; space between the eye and the bill, ear-coverts, and the patch on each side of the breast, cinereous brown: the red on the head and nape tinged with ash-gray. (*Young before the first moult.*) The white on the forehead, above the eyes, and across the nape of the neck, very indistinct: no black any

where; the large spot on the sides of the breast light brown, and very faintly indicated: all the upper parts light cinereous brown. (*Egg*.) Stone-colour, spotted with black: long. diam. one inch three lines; trans. diam. eleven lines.

A rare, and only occasional visitant in this country. Observed many years ago near Sandwich in Kent, by the late Mr Boys. Has more recently occurred in two or three instances on the coasts of Norfolk and Sussex. Habits similar to those of the last two species. Feeds on marine insects and worms. Lays from three to five eggs on the bare sand.

GEN. 64. VANELLUS, *Briss.*

(I. SQUATAROLA, *Cuv.*)

166. *V. griseus*, *Briss.* (*Gray Plover*.)—Upper plumage dusky brown, variegated with white and ash-colour; breast and belly white with dark spots (*winter*), or of a uniform deep black (*summer*).

V. melanogaster, *Temm. Man. d'Orn.* tom. II. p. 547. *Gray Plover*, *Selb. Illust.* vol. II. p. 227. pl. 35. *Bew. Brit. Birds*, vol. II. p. 83. *Gray Sandpiper*, *Mont. Orn. Dict.*

DIMENS. Entire length eleven inches six lines: length of the bill (from the forehead) one inch and half a line, (from the gape) one inch three lines and a half; of the tarsus one inch eleven lines; of the naked part of the tibia ten lines; of the tail three inches: from the carpus to the end of the wing seven inches eleven lines: breadth, wings extended, twenty-five inches six lines.

DESCRPT. (*Winter plumage*.) Forehead and chin white; streak over the eyes, fore part of the neck, sides of the breast, and flanks, white, variegated with spots of brown and ash-colour: head, and all the upper parts of the body, dusky brown, the feathers edged and tipped with grayish white: belly, abdomen, thighs, and upper tail-coverts, pure white: beneath the wing some long black feathers arising from the axilla: tail white, towards the tip reddish, with transverse brown bars which become paler and less numerous on the lateral feathers: bill black: irides dusky: feet blackish gray. (*Summer plumage*.) Space between the eye and the bill, throat, sides and fore part of the neck, middle of the breast, belly and flanks, deep black; forehead, a broad streak above the eyes continued down the sides of the neck and breast and bounding the black on those parts, thighs and abdomen, pure white: crown and occiput cinereous brown, with the shafts of the feathers black; nape variegated with brown, black, and white; back, scapulars, and wing-coverts, black, all the feathers broadly tipped and barred with white. (*Young of the year*.) Somewhat similar to the adult bird in winter plumage: the spots on the sides of the breast and flanks larger, but of a paler tint: upper parts of a uniform light gray, variegated with whitish spots; the transverse bars on the tail gray. (*Egg*.) Light clay-brown, spotted and streaked all over with dark brown: long. diam. one inch ten lines; trans. diam. one inch four lines.

Found on many parts of the British coast during the autumn and winter months, but not in very great abundance. Retires northward to breed. Feeds on worms and marine insects. Lays four eggs.

(2. VANELLUS, *Cur.*)

167. *V. cristatus*, Meyer. (*Crested Lapwing*.)—Crown, fore part of the neck and breast, greenish black; occipital feathers very much elongated, slightly recurved: back and scapulars olive-green.

V. cristatus, *Temm. Man. d'Orn.* tom. II. p. 550. Crested or Green Lapwing, *Selb. Illust.* vol. II. p. 221. pl. 34. Lapwing, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. II. p. 79.

DIMENS. Entire length twelve inches six lines: length of the bill (from the forehead) ten lines and a half, (from the gape) one inch and half a line; of the tarsus one inch ten lines; of the tail four inches six lines: from the carpus to the end of the wing eight inches seven lines: breadth, wings extended, twenty-eight inches five lines.

DESCRIPT. (*Winter plumage*.) Occipital feathers very much elongated, filiform, and turned upwards at the extremities: forehead, crown, crest, fore part of the neck, and upper part of the breast, of a shining greenish black; throat, region of the eyes, and sides of the neck, white: back, scapulars, and wing-coverts, olive-green, glossed with blue and purplish red: quills black; the first four primaries tipped with white: lower part of the breast, and all the belly, white; upper and under tail-coverts pale ferruginous: basal half of the tail white; the remaining portion black, tipped with white; outer feather almost entirely white: bill dusky: feet brownish red. (*Summer plumage*.) The black on the neck of a deeper tint, and extending over the throat; the occipital crest somewhat longer; the green and blue reflections on the upper parts more brilliant. (*Young of the year*.) The crest shorter; throat variegated with white and cinereous brown; beneath the eyes a dusky streak; all the feathers, as well on the upper as under parts, tipped with reddish yellow: feet olivaceous ash. (*Egg*.) Olive ground, blotched and spotted nearly all over with blackish brown: long. diam. one inch eleven lines; trans. diam. one inch four lines.

Common in most parts of the kingdom, frequenting fens and moist fields, as well as heaths, warrens, and upland situations. Has a peculiar note resembling the word *pee-wit*. Feeds on insects, worms, and snails. Breeds early in the Spring, depositing its eggs, which are four in number, on the bare ground. Collects in vast flocks at the approach of Autumn.

GEN. 65. STREPSILAS, *Illig.*

168. *S. Interpres*, Leach. (*Common Turnstone*.)—Upper parts variegated with white, black, and ferruginous; breast and abdomen white.

S. collaris, *Temm. Man. d'Orn.* tom. II. p. 553. Common Turnstone, *Selb. Illust.* vol. II. p. 204. pl. 33*. Turnstone, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. II. pp. 108, and 110.

DIMENS. Entire length nine inches three lines: length of the bill (from the forehead) ten lines, (from the gape) eleven lines and a half; of the tarsus eleven lines and a half; of the tail two inches eight lines;

from the carpus to the end of the wing five inches ten lines: breadth, wings extended, eighteen inches eight lines.

DESCRIPT. (*Adult male.*) Forehead, space between the eye and the bill, throat, and a broad collar on the nape of the neck, white: across the forehead a narrow black streak, which, passing over the eyes, becomes somewhat enlarged on the sides of the face; from thence it branches off on one side to the base of the lower mandible, on the other to the sides of the neck, where it again enlarges before uniting with a large gorget of the same colour on the lower part of the neck and breast: rest of the under parts pure white: crown of the head reddish white, with longitudinal black streaks; upper part of the back, scapulars, and wing-coverts, ferruginous brown, with large irregularly distributed black spots; lower back and upper tail-coverts white; a broad black bar on the rump: quills dusky; the shafts of the primaries, and tips of the secondaries, white: tail black, tipped with white; the outer feather entirely white: bill black: feet orange-yellow. The *female* differs only in having the colours not so distinct; the white on the head and neck less pure; the black not so deep. (*Young of the year.*) Head and nape cinereous brown, with darker variegations; some white spots on the cheeks and sides of the neck; throat and fore part of the neck whitish; sides of the breast deep brown, the feathers tipped with whitish; rest of the under parts, and a considerable portion of the back, pure white; upper part of the back, scapulars, and wing-coverts, deep brown, all the feathers broadly edged with yellowish; the transverse bar on the rump deep brown, bordered with red: feet yellowish red. (*Egg.*) Reddish white, blotched and spotted with dark chestnut-brown: long. diam. one inch six lines; trans. diam. one inch.

A regular winter visitant on many parts of the coast, appearing in August and departing in the Spring. Goes northward to breed. Derives its name from its habit of turning over the small stones on the shore in search of marine insects and worms on which it feeds. Is said to lay three or four eggs, in a small hollow in the sand. *Obs.* The *Tringa Morinella* of Linnæus is this bird in immature plumage.

GEN. 66. CALIDRIS, *Illig.*

169. *C. arenaria*, *Illig.* (*Sanderling.*) — Cinereous above, with darker streaks; all the under parts white (*winter*): or ferruginous and white spotted with black; belly and abdomen white (*summer*).

C. arenaria, *Temm. Man. d'Orn.* tom. II. p. 524. *Sanderling*, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. I. p. 375. *Common Sanderling*, *Selb. Illust.* vol. II. p. 208. pl. 36.

DIMENS. Entire length eight inches: length of the bill (from the forehead) one inch and half a line, (from the gape) one inch one line; of the tarsus eleven lines and a half; of the middle toe, claw included, eight lines and a half; from the carpus to the end of the wing four inches nine lines.

DESCRIPT. (*Winter plumage.*) All the upper parts cinereous, with the shafts of the feathers blackish brown: forehead, cheeks, throat, sides of the neck, and all the under parts, pure white: the bend and edge of the wing blackish gray; primaries dusky, the edges and tips inclining to brown; wing-coverts broadly edged with white: tail deep gray, the

feathers edged with white; the two middle ones darkest: bill, irides, and feet, black. (*Summer plumage*.) Cheeks and upper part of the head black, the feathers edged and variegated with ferruginous and white: throat, neck, breast, and upper part of the flanks, reddish ash; the central portion of each feather black, the tip whitish: the rest of the under parts pure white: back and scapulars deep ferruginous, with large irregular patches and spots of black; all the feathers edged and tipped with whitish: wing-coverts dusky brown, margined and tipped with white, forming a transverse bar: the two middle tail-feathers blackish brown with ferruginous edges. (*Young of the year before the first moult*.) "Feathers on the crown of the head, back, scapulars, and wing-coverts, black, edged and spotted with yellowish; between the bill and the eye a cinereous brown streak; nape, sides of the neck, and sides of the breast, pale gray, with fine undulating streaks; forehead, throat, fore part of the neck, and all the under parts, pure white: wings and tail as in the adult." TEMM. (*Egg*.) Olive ground, spotted and speckled with black: long. diam. one inch four lines; trans. diam. one inch.

Not uncommon in small flocks on many parts of the coast, but rarely observed inland. Is partial to sandy shores, feeding upon marine insects. Breeds in high northern latitudes. Said to construct a rude nest of grass in the marshes, and to lay four eggs; incubation commencing in the middle of June.

GEN. 67. HÆMATOPUS, Linn.

170. *H. ostralegus*, Linn. (*Pied Oyster-Catcher*.)—

Head, neck, and upper parts black: breast and abdomen white: bill and feet red.

H. ostralegus, Temm. *Man. d'Orn.* tom. II. p. 531. *Pied Oyster-Catcher*, Mont. *Orn. Dict. and Supp.* *Oyster-Catcher*, Bew. *Brit. Birds*, vol. II. p. 113. *Common Oyster-Catcher*, Selb. *Illustr.* vol. II. p. 200. pl. 33.

DIMENS. Entire length sixteen inches ten lines: length of the bill (from the forehead) two inches nine lines, (from the gape) two inches ten lines and a half; of the tarsus one inch eleven lines; of the tail four inches two lines; from the carpus to the end of the wing ten inches nine lines: breadth, wings extended, thirty-two inches.

DESCRIPT. (*Winter plumage*.) Head, neck, upper part of the breast, back, wings, and extremity of the tail, deep black; a collar on the throat, and a small spot beneath the eye, white: under parts, rump, basal portion of the tail and quill feathers, and a transverse bar on the wings, pure white: bill, and orbits, bright orange-red: irides crimson: feet purplish red. The *summer plumage* is distinguished by the absence of the white collar; in other respects similar. (*Young of the year*.) "The black parts of the plumage shaded with brown, the feathers being edged with this last colour: the white dull and soiled: bill and orbits dusky brown: irides brown: feet livid gray." TEMM. (*Egg*.) Yellowish stone-colour, spotted with ash-gray and dark brown: long. diam. two inches two lines; trans. diam. one inch six lines.

Common on the coast, assembling in small flocks during the winter season. Is never found inland. Feeds principally on marine insects and the bivalve mollusca. Makes no nest, but deposits its eggs, two to four in number, on the bare ground above high-water mark. The male bird has a loud screaming note during the season of incubation.

- (16.) *Psophia crepitans*, Linn. (*Gold-breasted Trumpeter*.) Lath. Syn. vol. II. p. 793. pl. 68.

An individual of this species is recorded by Montagu, in the Supplement to his Ornithological Dictionary, as having been taken in Surrey; it had, however, probably escaped from confinement, being a native of South America.

GEN. 68. GRUS, *Pall.*

171. *G. cinerea*, Bechst. (*Common Crane*.) — Cinereous; occiput, throat, and fore part of the neck, dusky gray.

G. cinerea, *Temm. Man. d'Orn.* tom. II. p. 557. Common Crane, *Selb. Illust.* vol. II. p. 4. pl. 1. *Bew. Brit. Birds*, vol. II. p. 3. Crane, *Mont. Orn. Dict.* (Trachea,) *Linn. Trans.* vol. IV. pl. 12. f. 4.

DIMENS. Entire length five feet: length of the bill above four inches. MONT.

DESCRIPT. Forehead, and space between the bill and the eyes, covered with black bristly hairs; crown of the head naked, the skin of an orange-red colour: occiput, throat, and fore part of the neck, of a deep blackish gray; all the upper and under parts of the body dark ash-colour: quills, and greater coverts, black; secondaries and tertials elongated, and of an arched form, with the barbs of the feathers disunited, forming an elegant tuft of floating plumes capable of being erected or depressed at will: bill greenish black, the tip horn-colour: feet black. "*Young birds, previously to the second autumnal moult*, are without the naked patch on the crown of the head, or have it very small; the deep blackish tint on the occiput and fore part of the neck is simply indicated by a few longitudinal spots of that colour. In *old individuals*, a large whitish space is found behind the eyes, and on each side of the upper part of the neck." TEMM. (*Egg*.) Pale greenish olive ground, blotched and spotted with darker green and olive-brown: long. diam. four inches; trans. diam. two inches six lines.

According to Ray, this species was formerly met with in Cambridge-shire, in large flocks, during the winter months. In still earlier times it is stated by Turner to have bred in some of our fens. It must now be considered as an extremely rare and accidental visitant. Pennant makes mention of a single specimen which was killed in his time near Cambridge. One or two others are said to have occurred in Kent; and so recently as in the year 1828, an individual is recorded to have been killed in Cornwall. This last is now in the collection of Mr Drew, Devonport. (*Loud. Mag.* vol. III. p. 177.) Is a common inhabitant of marshy plains in the eastern parts of Europe. Feeds on grain, aquatic plants, worms, and small reptiles. Said to build amongst rushes and other thick herbage, sometimes on the roofs of solitary houses. Lays two eggs.

GEN. 69. ARDEA, Linn.

(1. ARDEA, Steph.)

* *A large portion of the tibia naked; tarsi long.*

172. *A. cinerea*, Lath. (*Common Heron*).—General plumage bluish ash: middle toe, claw included, much shorter than the tarsus.

A. cinerea, Temm. *Man. d'Orn.* tom. II. p. 567. *Common Heron*, Mont. *Orn. Dict. and Supp.* Selb. *Illust.* vol. II. p. 11. pl. 2. Heron, Bew. *Brit. Birds*, vol. II. p. 9.

DIMENS. Entire length three feet two inches nine lines; length of the bill (from the forehead) four inches seven lines, (from the gape) five inches eleven lines; of the tarsus five inches eight lines; of the naked part of the tibia two inches eight lines; of the middle toe three inches ten lines; of the tail eight inches; from the carpus to the end of the wing one foot five inches seven lines: breadth, wings extended, five feet six inches.

DESCRIPT. (*Adult, after the age of three years.*) A pendent crest of narrow, elongated, black feathers at the back of the head: forehead, crown, neck, middle of the breast, belly, edge of the wing, thighs, and under tail-coverts, pure white; fore part of the neck with large longitudinal black spots; the feathers towards the bottom loose and very much elongated, falling over the breast in an elegant manner: occiput, sides of the breast, and flanks, deep black: back and wings of a fine bluish gray: quills black: scapulars long and pointed, forming graceful plumes of a silvery gray colour: tail bluish ash: bill deep yellow: irides bright gamboge-yellow: lore, and naked skin round the eyes, greenish: feet brown; the naked part of the tibia red. (*Young, and until the age of three years.*) Occipital crest very short, almost wanting: without the long plumes on the scapulars and lower part of the neck: forehead and crown cinereous; throat white; neck pale ash-colour, inclining to whitish underneath, with large longitudinal dusky spots: rest of the upper parts bluish ash; ridge of the wing spotted with white and reddish brown; quills of a deep bluish black: breast, belly, and vent, white, spotted with black: upper mandible dusky brown, variegated with yellowish spots; lower mandible wholly yellow: irides yellow: lore and orbits greenish yellow: feet dark ash-colour; the lower part of the tibia yellowish. (*Egg.*) Of a uniform sea-green: long. diam. two inches three lines; trans. diam. one inch nine lines.

A common and generally distributed species. Congregates in large societies during the breeding season, building on the loftiest trees. At other times of the year, of rather solitary habits, residing principally in marshy districts, or in the neighbourhood of streams and rivers. Feeds principally on fish and small reptiles. Lays three or four eggs.

173. *A. purpurea*, Linn. (*Purple Heron*).—General plumage reddish ash, inclining to purple: middle toe, claw included, as long as, or longer than the tarsus.

A. purpurea, Temm. *Man. d'Orn.* tom. II. p. 570. *Crested Purple Heron*, Lath. *Syn.* vol. III. p. 95. *Shaw*, *Zool.* vol. XI. p. 556. *Selb. Illust.* vol. II. p. 15. pl. 3.

DIMENS. Entire length two feet eleven inches: length of the bill (from the forehead) four inches eight lines, (from the gape) five inches seven lines; of the tarsus four inches seven lines; of the naked part of the tibia two inches seven lines; of the middle toe, claw included, four inches nine lines; from the carpus to the end of the wing thirteen inches eight lines.

DESCRIPT. (*Adult in perfect plumage.*) Crown, and occipital crest, black, with green reflections; throat white; cheeks, and sides of the neck, reddish brown, with three longitudinal narrow black bands, two lateral reaching from the eyes to the breast, the third commencing at the nape and running down the back of the neck for two-thirds of its length: front of the neck variegated with red, black, and purple; the feathers on the lower part long and acuminate, of a purplish white colour: back, wings, and tail, reddish ash; scapulars long and subulate as in the last species, of a rich brilliant purple red: breast and flanks of a deep brownish red, tinged with purple: belly and thighs red: bill and orbits bright yellow: irides yellowish orange: soles of the feet, posterior part of the tarsus, and naked space on the tibia, yellow; fore part of the tarsus, and upper surface of the toes, greenish brown. (*Immature plumage.*) No occipital crest, or simply a few red feathers somewhat longer than the others: without the long subulate plumes on the lower part of the neck and scapulars: forehead black: nape and cheeks pale red; throat white; fore part of the neck, and sides of the breast, yellowish white, with numerous longitudinal black spots: back, wings, and tail, dusky ash, all the feathers edged with reddish ash: belly and thighs whitish: a large portion of the upper mandible dusky; lower mandible, orbits, and irides, pale yellow. (*Egg.*) Pale asparagus-green: long. diam. two inches four lines; trans. diam. one inch seven lines.

A very rare, and only occasional visitant in this country. Has been killed in Berkshire, Cambridgeshire, Norfolk, and in one or two other parts of the kingdom. A specimen is also said to have flown on board a fishing boat off the coast of Cornwall in May 1822, and to have been taken. Frequents the same kind of situations as the last species. Food similar. Is said to breed amongst reeds and thick underwood, rarely in trees. Lays three eggs. *Obs.* The *African Heron* of Montagu's *Ornithological Dictionary* (A. Caspica, Gmel.) is this species in immature plumage.

- (17.) *A. alba*, Linn. Syst. Nat. tom. i. p. 239. *Great White Heron*, Mont. Orn. Dict. & Supp. Selb. Illust. vol. II. p. 18. *A. Egretta*, Temm. Man. d'Orn. tom. II. p. 572. *White Heron*, Penn. Brit. Zool. vol. II. p. 427. pl. 62.

Recorded as a British species by Ray. Asserted by Latham, on the authority of Dr Heysham, to have been shot in Cumberland: supposed also to have been seen in Devonshire and Suffolk. There is, however, no well authenticated instance of its having been met with in this country of late years, nor any British-killed specimen in existence. It is distinguished by its pure white plumage, greenish yellow bill, and long slender legs: the *adult*, in the breeding season, possesses an occipital crest and long dorsal plumes like the next species. Is common in some of the eastern parts of Europe. Food and habits resembling those of its congeners.

174. *A. Garzetta*, Linn. (*Little Egret Heron.*)—
Whole plumage white: bill black.

- A. Garzetta*, Temm. Man. d'Orn. tom. II. p. 574. *Egret*, Penn. Brit. Zool. vol. II. Append. p. 631. pl. 7. *Mont. Orn. Dict.* *Little Egret*, Lath. Syn. vol. III. p. 90. *Bew. Brit. Birds*, vol. II. p. 17. *Little Egret Heron*, Selb. Illust. vol. II. p. 21. pl. 5.

DIMENS. Entire length one foot ten or eleven inches. TEMM.

DESCRIPT. (*Mature plumage in Summer.*) The whole plumage pure white: a pendent crest of long narrow feathers on the occiput: a tuft of similar feathers, the webs silky and disunited, on the lower part of the neck: dorsal plumes likewise of the same character, being of a soft silky texture, with the shafts very much elongated, and floating loosely over the back and tail from between the shoulders: bill black: lore and orbits greenish: irides bright yellow: feet dusky green: toes greenish yellow. (*Immature plumage, and adult in Winter.*) The long silky plumes on the occiput, back, and lower part of the neck, wanting. In *very young* birds, the white is of a dull tint; the bill, lore, orbits, irides, and feet, black. (*Egg.*) White. TEMM.

An extremely rare and accidental visitant in this country, though supposed to have been more plentiful formerly. Said to have been shot in Anglesey, and also in Ireland, many years ago. More recently, in April 1824, two specimens are recorded to have been killed near Penzance in Cornwall, and one of them to have been preserved. Common on some parts of the Continent. Said to breed in marshes, and to lay four or five eggs.

175. *A. russata*, Wagler. (*Buff-backed Heron.*) — Occiput, nape, and hind-neck, saffron-yellow: wings, tail, and under parts, pure white.

A. russata, Wagler, *Syst. Av.* part i. sp. 12. *A. æquinoctialis*, Mont. in *Linn. Trans.* vol. ix. p. 197. Little White Heron, Mont. *Orn. Dict. Supp. with fig.* Buff-backed Heron, *Selb. Illust.* vol. 11. p. 24.

DIMENS.* Entire length twenty inches six lines: length of the bill (from the forehead) two inches two lines, (from the gape) two inches ten lines; of the tarsus three inches; of the naked part of the tibia one inch one line; of the middle toe, claw included, two inches nine lines; of the tail four inches; from the carpus to the end of the wing ten inches.

DESCRIPT. (*Mature plumage.*) "Occiput, nape, and hinder part of the neck, clothed with rigid open feathers of a saffron yellow: throat, front part of the neck and breast, white, tinged with sienna-yellow: long flowing plumes of the back ochre-yellow: wings, tail, and under parts of the body, pure white." SELBY. (*Immature plumage.*) "The whole plumage snow white, excepting the crown of the head, and the upper part of the neck before, which are buff: the skin of a very dark colour, almost black, so that on the cheeks and sides of the neck, where the feathers are thin, it is partly seen, giving a dingy shade to the white plumage of those parts: feathers on the back of the head somewhat elongated, but scarcely forming a crest: those on the lower part of the neck before, rather more elongated, hanging loosely over the upper part of the breast: tail slightly forked, and so short as to be entirely covered by the wings when folded: bill, lore, and orbits, orange-yellow: irides pale yellow: legs nearly black, with a tinge of green." MONT.

A single individual of this species, in immature plumage, and probably a young bird of the year, is recorded by Montagu to have been shot near Kingsbridge in Devonshire, the latter end of October 1805. This specimen, which is now in the British Museum, is the only one that has

* These measurements were taken from Montagu's original specimen.

hitherto occurred in this country. It is a native of the warmer parts of Europe, as well as of Asia and Africa, but is not found in America, "the *Ardea æquinoctialis*, with which it has been confounded, being a distinct species, and confined to that continent." Habits unknown.

** *A small portion of the tibia naked; tarsi short.*

176. *A. Ralloides*, Scop. (*Squacco Heron*.) — Occipital feathers very long, white edged with black: neck, back, and scapulars, pale buff-orange.

A. Ralloides, *Temm. Man. d'Orn.* tom. 11. p. 581. *Squacco Heron*, *Selb. Illust.* vol. 11. p. 25. pl. 6. *Mont. Orn. Dict. Supp.* *Shaw, Zool.* vol. x1. p. 574.

DIMENS. Entire length seventeen inches: length of the bill (from the forehead) two inches six lines, (from the gape) three inches one line; of the tarsus two inches; of the naked part of the tibia eleven lines and a half: of the middle toe, claw included, two inches seven lines; of the tail three inches one line; from the carpus to the end of the wing eight inches eleven lines.

DESCRIPT. (*Mature plumage*.) Forehead and crown yellowish, streaked with black; a pendent crest from the occiput, consisting of eight or ten very long narrow white feathers edged at the sides with black: throat white; neck, breast, upper part of the back, and scapulars, pale buff orange; middle and lower regions of the back inclining to ferruginous chestnut, the feathers on those parts very long, with disunited webs, and floating loosely over the tail: all the rest of the plumage pure white: bill bluish green at the base, the tip black: lore and orbits gray tinged with green: irides yellow: feet yellow tinged with green. (*Plumage during the first and second year*.) "Without the long occipital feathers; head, neck, and wing-coverts, ferruginous brown, with large longitudinal spots of a darker tint; throat, rump, and tail, pure white; upper part of the back and scapulars of a more or less deep brown: upper mandible greenish brown; lower mandible greenish yellow: orbits green: irides very pale yellow: feet greenish ash." *TEM.* (*Egg*.) Unknown.

A very rare, and only occasional visitant. Shot at Boyton in Wiltshire, in 1775. A second specimen taken at Ormsby in Norfolk, in 1820; a third killed in Cambridgeshire; and a fourth, a female, at Bridgewater, in the Summer of 1825. Common in some parts of the Continent. Frequents marshes and the borders of rivers, &c. Food small fish, mollusca, and insects. Said to build in trees. *Obs.* This species is the *A. comata* of Pallas and some other authors.

*** *Tibia entirely feathered; tarsi short.*

177. *A. minuta*, Linn. (*Little Heron*.) — Crown, back, scapulars, and secondaries, black, glossed with green; neck, wing-coverts, and under parts, reddish yellow.

A. minuta, *Temm. Man. d'Orn.* tom. 11. p. 584. *Little Bittern*, *Penn. Brit. Zool.* vol. 11. *App.* p. 633. pl. 8. *Mont. Orn. Dict. and Supp.* *Selb. Illust.* vol. 11. p. 36. pl. 6*. *Bew. Brit. Birds*, vol. 11. pp. 25, & 27.

DIMENS. Entire length fourteen inches three lines: length of the bill two inches; of the tarsus one inch nine lines; of the middle toe, claw included, two inches; from the carpus to the end of the wing five inches six lines.

DESCRIPT. (*Adult.*) Crown of the head, occiput, back, scapulars, secondary quills, and tail, black, with green reflections; cheeks, neck, wing-coverts, and all the under parts, reddish yellow; flanks, and sides of the breast, with a few brownish streaks; primary quills dusky: bill yellow; the tip brown: orbits and irides yellow: feet greenish yellow. (*Young of the year.*) Crown of the head brown; fore part of the neck whitish with longitudinal dark streaks; cheeks, nape, breast, back, and wing-coverts, chestnut-brown more or less deep, some of the feathers with pale rufous edges: greater quills and tail-feathers deep brown. (*Egg.*) White: long. diam. one inch six lines; trans. diam. eleven lines.

Occasionally met with, but not common in this country. Montagu records that three specimens were shot in Devonshire during the Spring and Summer of 1808. Others have been killed at different times in the Orkneys, in Somersetshire, Shropshire, Northumberland, Hampshire, Suffolk, and, more recently, on Uxbridge Moor, and on the banks of the Thames near Windsor. Frequents marshes, and rushy places on the borders of lakes and rivers. Breeds in such situations, constructing a large nest of leaves and rushes. Lays five or six eggs. Food small reptiles, fish, worms, &c.

(2. *BOTAURUS*, *Steph.*)

178. *A. stellaris*, Linn. (*Common Bittern.*)—Upper parts ochre-yellow, variegated with black; beneath paler, with oblong dusky streaks.

A. stellaris, *Temm. Man. d'Orn.* tom. II. p. 580. *Bittern*, *Mont. Orn. Dict. Selb. Illust.* vol. II. p. 30. pl. 8. *Bew. Brit. Birds*, vol. II. p. 22.

DIMENS. Entire length two feet six inches six lines: length of the bill (from the forehead) three inches, (from the gape) four inches; of the tarsus three inches eleven lines; of the middle toe four inches six lines; of the tail four inches six lines; from the carpus to the end of the wing one foot: breadth, wings extended, three feet nine inches.

DESCRIPT. Crown of the head, and a broad streak or moustache from the corners of the mouth, black; neck-feathers loose and elongated, capable of being raised and depressed at will, ochre-yellow, with brown zigzag transverse lines on the sides, and long streaks and spots of reddish brown in front: all the upper parts of the body ochre-yellow tinged with orange-red, with a large dusky spot in the middle of each feather; primary and secondary quills, primary coverts, and spurious winglet, ferruginous, with transverse bars of blackish brown; rest of the coverts and scapulars, mottled like the back with ochre-yellow and zigzag dusky lines: under parts of the same colour as above but paler, with large oblong longitudinal dusky streaks: upper mandible brown, passing into yellow at the edges; lower mandible, lore, orbits, and feet, greenish yellow; irides bright gamboge-yellow. (*Egg.*) Of a uniform light olive-brown: long. diam. two inches two lines; trans. diam. one inch six lines.

Occasionally met with in extensive marshes, as well as on the borders of rivers and lakes, but not so abundant as formerly. Is partial to sedge

and reedy situations. Breeds in such spots, constructing a nest of rushes and other coarse plants. Lays four or five eggs. During the Spring the male utters a singular bellowing note which may be heard to a considerable distance. Feeds on small fish and reptiles, as well as on aquatic worms and insects.

179. *A. lentiginosa*, Mont. (*American Bittern*.)—Back and scapulars chocolate-brown glossed with purple, the edges of the feathers paler: under parts ochre-yellow, marked on the neck, breast, and belly, with broad chestnut streaks.

A. lentiginosa, *Faun. Amer. Bor.* vol. II. p. 374. Freckled Heron, *Mont. Orn. Dict. Supp. with fig.* American Bittern, *Selb. Illust.* vol. II. p. 34.

DIMENS. Entire length about twenty-three inches: length of the bill (from the forehead) two inches nine lines: of the tarsus three inches nine lines. **MONT.**

DESCRIPT. Crown of the head chocolate brown, passing into dull yellow at the nape, where the feathers are much elongated; chin and throat white, with a row of brown feathers down the middle; from the base of the lower mandible a black streak increasing on the upper part of the neck on each side; cheeks yellowish, with an obscure dusky line at the corner of the eye; the feathers on the neck long and broad, with their webs partly unconnected; those in front pale dull yellow with broad chestnut streaks; hind neck bare; the feathers on the breast long, of a fine chocolate brown, glossed with purple, and edged with dull yellow; belly and sides the same, but not so bright, the brown marks becoming speckled; vent and under tail-coverts yellowish white: back and scapulars chocolate-brown, minutely speckled and glossed with purple, the edges of the feathers paler: wing-coverts dull yellow, darkest in the middle of each feather, the margins prettily speckled; first and second order of quills, their greater coverts, and the spurious wing, dusky lead-colour, with a tinge of cinereous; primaries very slightly tipped with brown; secondaries and the greater coverts tipped more deeply with the same, and prettily speckled on the light part: the closed wings not reaching to the end of the tail: upper mandible dusky above, greenish yellow at the sides; lower mandible greenish yellow: legs greenish. **MONT.** (*Egg.*) Cinereous green. *Faun. Am. Bor.*

A single individual of this species, which is common in North America, was shot by Mr Cunningham, in the parish of Piddletown, in Dorsetshire, in the Autumn of 1804. This specimen was procured by Montagu, and, with the rest of his collection, is now in the British Museum. No other has occurred since in this country. Has a loud booming note similar to that of the last species. Said to frequent marshes and willow thickets, and to lay four eggs.

(3. NYCTICORAX, Steph.)

180. *A. Nycticorax*, Linn. (*Common Night-Heron*.)—Crown, back, and scapulars, black, glossed with green: under parts white.

A. Nyctieorax, *Temm. Man. d'Orn.* tom. 11. p. 577. Night-Heron, *Mont. Orn. Dict. Selb. Illust.* vol. 11. p. 39. pls. 7, & 7*. *Bew. Brit. Birds*, vol. 11. p. 14.

DIMENS. Entire length twenty-one inches: length of the bill (from the forehead) two inches eight lines, (from the gape) three inches eight lines; of the tarsus two inches eleven lines; of the naked part of the tibia ten lines; of the middle toe, claw included, three inches one line and a half; from the carpus to the end of the wing twelve inches.

DESCRIPT. (*Adult.*) Crown of the head, nape, upper part of the back, and scapulars, black glossed with green: occiput ornamented with three very narrow white feathers, measuring six or seven inches in length: lower part of the back, wings, and tail, of a fine pearl-gray: forehead, a narrow streak above the eye, throat, and all the under parts, pure white; bill black, yellowish at the base: lore and orbits green: irides deep orange: feet yellowish green. (*Young of the year.*) Without the long subulate feathers on the occiput: upper part of the head, nape, back, and scapulars, dull brown, with longitudinal streaks of yellowish white, one in the middle of each feather: throat white, spotted with brown; feathers on the sides and fore part of the neck, yellowish, broadly edged with brown; rest of the under parts variegated with brown, white, and ash-colour; middle of the belly whitish: wing-coverts, and primary quills, cinereous brown, with large pisciform yellowish white spots at the tips of the feathers: culmen and tip of the bill dusky brown, the remaining portion greenish yellow: irides brown: feet olivaceous brown. (*Egg.*) Pale blue: long. diam. two inches three lines; trans. diam. one inch nine lines.

Very rare in this country. Has been killed in Norfolk, Suffolk, Bedfordshire, Oxfordshire, Buckinghamshire, and, in more than one instance, near London. Common in the South of Europe; inhabiting the borders of rivers and lakes where there is covert. Feeds on fish, reptiles, worms, &c. Said to build on the ground, and to lay three or four eggs. *Obs.* The *Ardea Gardani* of Gmelin and Latham is the young of the year of this species. The supposed *A. Cayennensis*, recorded by Mr Youell to have been taken near Yarmouth, May 23, 1824, (*Linn. Trans.* vol. xiv. p. 588.) is probably only a variety of the common Night-Heron, having double the usual number of long occipital feathers.

GEN. 70. CICONIA, *Briss.*

181. *C. alba*, Ray. (*White Stork.*)—White; scapulars and wings black.

C. alba, *Temm. Man. d'Orn.* tom. 11. p. 560. Stork, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 6. White Stork, *Selb. Illust.* vol. 11. p. 45. pl. 11.

DIMENS. Entire length three feet eight inches: length of the bill (from the forehead) seven inches six lines; of the tarsus twelve inches; of the naked part of the tibia six inches.

DESCRIPT. Head, neck, and all the body, pure white; scapulars and wings black: bill and legs red: naked skin round the eyes black: irides brown. In *young birds*, the black on the wings inclines to brown, and the bill is dusky tinged with reddish. (*Egg.*) White: long. diam. two inches ten lines; trans. diam. one inch eleven lines.

Very common in Holland, France, and other parts of the Continent, but in this country a rare, and only accidental visitant. Montagu mentions one which was killed near Salisbury, in February 1790; a second shot at Sandwich in Kent, in 1805; and a third in Hampshire, in the Autumn of 1808. Has occurred in Suffolk more recently, and in three or four instances. Food, reptiles and insects, as well as small quadrupeds and birds. Builds on the tops of houses, or in old trees. Lays generally three eggs.

182. *C. nigra*, Ray. (*Black Stork*.)—Black; lower part of the breast, and belly, white.

C. nigra, Temm. *Man. d'Orn.* tom. II. p. 561. *Black Stork*, Mont. in *Linn. Trans.* vol. XII. p. 19. *Selb. Illust.* vol. II. p. 48. pl. 11*.

DIMENS. Entire length nearly three feet. TEMM.

DESCRIPT. (*Adult.*) Head, neck, all the upper parts of the body, wings, and tail, black, with purple and green reflections; lower part of the breast and belly dingy white: bill and orbits bright orange: irides hazel: legs and toes deep red. (*Young.*) Head, and upper part of the neck, pale reddish brown, with the central portion of the feathers dusky; back, scapulars, wings, and tail, dusky brown, very slightly glossed with greenish hues: bill, orbits, and legs, olivaceous green. (*Egg.*) Of a uniform greenish buff: long. diam. two inches seven lines; trans. diam. one inch eleven lines.

A single individual of this species is recorded by Montagu as having been shot in West Sedge-Moor, in Somersetshire, May 13, 1814. The specimen is now in the British Museum. A second is stated to have been shot in October 1832, in the parish of Otley, about eight miles from Ipswich. (*Loud. Mag. of Nat. Hist.* vol. VII. p. 53.) Inhabits various parts of Europe, but is less plentiful than the last species. Said to frequent wooded swamps and extensive forests: Food, small fish, reptiles, and insects. Builds in lofty trees, and lays two or three eggs.

GEN. 71. PLATALEA, Linn.

183. *P. Leucorodia*, Linn. (*White Spoonbill*.)—Whole plumage white: a pendent crest of long subulate feathers on the occiput.

P. Leucorodia, Temm. *Man. d'Orn.* tom. II. p. 595. *Spoonbill*, Penn. *Brit. Zool.* vol. II. *App.* p. 634. pl. 9. *Bew. Brit. Birds*, vol. II. p. 29. *White Spoonbill*, Mont. *Orn. Dict. & Supp. Selb. Illust.* vol. II. p. 51. pl. 10. (*Trachea*.) *Linn. Trans.* vol. XVI. pl. 19.

DIMENS. Entire length thirty-one inches: length of the bill (from the forehead) seven inches three lines; breadth of the spoon one inch ten lines; length of the tarsus four inches eleven lines; of the naked part of the tibia two inches eight lines; from the carpus to the end of the wing fourteen inches six lines.

DESCRIPT. (*Old male.*) The whole plumage pure white, with the exception of a large patch of buff yellow on the upper part of the breast, from whence a narrow band of the same colour ascends on each side towards the top of the back: lore, orbits, and naked space on the throat, orange-yellow: bill black; the tip ochre-yellow: irides red: legs black,

(*Old female*.) Somewhat smaller; the occipital crest shorter; the buff-coloured patch on the breast not so distinct. (*Young of the year*.) No occipital crest; the feathers on the head short and rounded; shafts of all the quills, and tips of the primaries, deep black; the rest of the plumage white: bill deep ash-colour, soft, and very flexible: irides cinereous: lore and orbits dingy white. "The patch of buff yellow on the breast does not shew itself till the second or third year." TEMM. (*Egg*.) White, spotted with pale reddish brown: long, diam. two inches nine lines; trans. diam. one inch nine lines.

A rare visitant in England. Pennant mentions a flock which migrated into the marshes near Yarmouth in Norfolk, in April 1774. Since then, specimens have been killed in Somersetshire, Devonshire, Dorsetshire, Lincolnshire, and Suffolk. In the latter county, three were shot out of a flight of seven, which appeared at Thorpe, in the Autumn of 1828. Common in Holland, frequenting the banks of rivers near their junction with the sea. Food, small fish, mollusca, and aquatic insects. Builds in trees or amongst rushes. Eggs two or three in number.

GEN. 72. IBIS, *Lacép.*

184. I. *Falcinellus*, Temm. (*Glossy Ibis*.)—Neck, breast, and under parts, bright chestnut-red; lower back, rump, wings and tail, glossy green with purple reflections.

I. *Falcinellus*, Temm. *Man. d'Orn.* tom. II. p. 598. *Glossy Ibis*, *Mont. Orn. Dict. & Suppl. Selb. Illust.* vol. II. p. 56. pl. 12. *Bew. Brit. Birds*, vol. II. p. 33. *Bay Ibis*, *Sow. Brit. Misc.* pl. 17. *Brazilian Curlew*, *Shaw, Nat. Misc.* vol. XVII. pl. 705.

DIMENS. Entire length twenty-two inches: length of the bill (from the forehead) four inches, (from the gape) four inches three lines; of the tarsus three inches six lines; of the naked part of the tibia one inch ten lines; from the carpus to the end of the wing ten inches eight lines.

DESCRIPT. (*Old bird in perfect plumage*.) Head chestnut-red passing into dusky brown; neck, breast, upper part of the back, and all the under parts, bright chestnut-red; lower part of the back, rump, wing-coverts, primary quills, and tail-feathers, deep dusky green, with purple and bronze reflections: bill and legs greenish black: lore and orbits green: irides brown. (*Young till the age of three years*.) Head, throat, and upper part of the neck, dusky brown, the feathers with whitish edges; occasionally a few large irregular spots and transverse bars of this last colour: lower part of the neck, breast, belly, and thighs, cinereous black, with a few greenish reflections on the breast: back and scapulars greenish brown; wings and tail as in the perfect plumage, but with the gloss of green and purple much less brilliant. The *young of the year* have the whole plumage inclining more to cinereous brown; and the feathers on the head and neck more broadly edged with white. (*Egg*.) Unknown.

An occasional but rare visitant in this country. Has been killed in the counties of Cornwall, Devon., Kent, Berks., Norfolk, and Northumberland; also in Anglesea and Ireland. Most of the specimens have occurred during the autumnal and winter months. Common in the eastern parts of Europe, inhabiting the borders of lakes and rivers. Feeds on insects, worms, shell-fish, and aquatic plants. Nidification unknown. *Obs.* The *Green Ibis* of Latham and other authors is this species in immature plumage.

GEN. 73. NUMENIUS, *Briss.*

185. *N. arquata*, Lath. (*Common Curlew*).—Head, neck, breast, and all the upper parts, rufous ash; belly and abdomen white; the whole variegated with dusky spots.

N. arquata, *Temm. Man. d'Orn.* tom. II. p. 603. Common Curlew, *Mont. Orn. Dict. & Supp. Selb. Illust.* vol. II. p. 62. pl. 13. *Bew. Brit. Birds*, vol. II. p. 36.

DIMENS. Entire length twenty-two to twenty-five inches: length of the bill from four to six inches and a half; of the tarsus three inches three lines; of the naked part of the tibia one inch two lines; of the tail four inches six lines; from the carpus to the end of the wing from eleven to eleven inches and a half: breadth, wings extended, from thirty-eight to forty-two inches.

DESCRIPT. Head, neck, and breast, pale ash tinged with rufous, the shafts and central portion of the feathers dusky; upper part of the back and scapulars blackish brown, the feathers broadly edged with pale rufous brown; lower part of the back inclining to white, spotted with black; tail yellowish white, with transverse brown bars; belly and abdomen white, with longitudinal dusky spots: upper mandible blackish brown; lower one flesh-colour: irides brown: legs and toes pale bluish gray. The *female* has less of the rufous, and more of the cinereous, tinge. The *young of the year* have the bill considerably shorter, and nearly straight. (*Egg*.) Olive green; blotched and spotted with darker green and dark brown: long. diam. two inches seven lines; trans. diam. one inch eleven lines.

Common on most parts of the coast during the winter season. Retires inland, and to more elevated spots, to breed. Deposits its eggs amongst heath, rushes, or long grass, to the number of four or five. Feeds on worms, insects, and molluscous animals.

186. *N. Phæopus*, Lath. (*Whimbrel*).—Head, neck, breast, and upper parts, pale ash-colour, spotted with dusky: lower back, belly and abdomen, white.

N. Phæopus, *Temm. Man. d'Orn.* tom. II. p. 604. Whimbrel, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. II. p. 38. Whimbrel Curlew, *Selb. Illust.* vol. II. p. 65. pl. 14.

DIMENS. Entire length seventeen inches: length of the bill three inches three lines: breadth, wings extended, twenty-nine inches.

DESCRIPT. Forehead and crown cinereous brown, the latter divided in the middle by a longitudinal pale streak; over each eye, another but broader streak of white variegated with brown; sides of the head, neck, and breast, pale ash, with longitudinal brown streaks: upper part of the back, scapulars, and wing-coverts, deep brown, the feathers with pale edges; lower part of the back white; rump barred with cinereous brown; tail cinereous brown, with oblique bars of a darker shade: belly and abdomen white; sides barred with brown: bill dusky, reddish towards the base: irides brown: legs deep bluish gray. In *young birds*, the bill is shorter, and not so much curved. (*Egg*.) Dark olive-brown, blotched with darker brown: long. diam. two inches five lines; trans. diam. one inch eight lines.

Found on many parts of the coast during the autumnal and winter months, but is less common than the last species. Generally observed in small flocks. Retires to the northern parts of the country to breed. Nest said to be placed on open and exposed heaths. Food, worms and insects.

GEN. 74. TOTANUS, *Bechst.*

* *Bill slender, straight.*

187. *T. fuscus*, Leisl. (*Dusky Sandpiper*.)—Upper plumage ash-gray; rump, and all the under parts, pure white; legs, and base of the lower mandible, red.

T. fuscus, *Temm. Man. d'Orn.* tom. II. p. 639. Cambridge Godwit, *Penn. Brit. Zool.* vol. II. p. 447. Dusky Sandpiper, *Shaw, Zool.* vol. XII. p. 132. *Selb. Illust.* vol. II. p. 69. pl. 15. f. 1, & 2.

DIMENS. Entire length twelve inches: length of the bill two inches five lines; of the tarsus two inches three lines; of the naked part of the tibia one inch three lines and a half; of the tail two inches seven lines; from the carpus to the end of the wing six inches ten lines.

DESCRIPT. (*Adult in winter plumage*.) Crown of the head, nape, and back, cinereous gray, with fine dusky streaks on the shafts of the feathers; wing-coverts and scapulars cinereous gray edged with white*; between the bill and the eye a blackish patch, above which is a white streak; cheeks, sides and fore part of the neck, variegated with white and ash-colour; throat, breast, rump, belly and abdomen, pure white; flanks whitish, passing into ash-gray: upper tail-coverts, and three or four outermost tail-feathers, with alternate transverse bars of white and dusky brown; central tail-feathers of a uniform ash-gray narrowly edged with white: bill black; base of the lower mandible red: legs bright orange-red. (*Adult in summer plumage*.) All the upper parts deep blackish brown, the feathers on the back, scapulars, and wing-coverts, marked on the edges with white spots mostly of a triangular form; head, neck, and under parts, dusky gray; the neck without spots, but the breast and belly sometimes with a narrow crescent-shaped white edging at the tip of each feather: vent, and under tail-coverts, barred transversely with white and dusky; tail dusky ash, with little transverse white streaks at the edges of the feathers, not reaching to the shafts: legs reddish brown. (*Young of the year*.) Upper parts olivaceous brown, the feathers on the back with a narrow edging of white; wing-coverts and scapulars with triangular white spots: all the under parts whitish, with indistinct spots and transverse undulating bars of cinereous brown: legs and toes orange-red.

A rare species in Great Britain, but has been killed at different times in various parts of the country. Frequents marshes, and the borders of lakes and rivers. Food, principally molluscous animals. Retires to high northern latitudes to breed. *Obs.* The *Cambridge Godwit* of Pennant is this species in the adult winter plumage: the *Spotted Redshank* of the same author (the *Spotted Snipe* of Latham and Montagu) is the young bird of the year.

188. *T. Calidris*, *Bechst.* (*Redshank Sandpiper*.)—Upper parts cinereous brown; rump, belly, and greater portion of the secondary quills, white: legs, and basal half of both mandibles, red.

* Probably, in very old birds, this white edging disappears.

T. *Calidris*, *Temm. Man. d'Orn.* tom. II. p. 643. Redshank Sandpiper, *Selb. Illust.* vol. II. p. 72. pl. 16. f. 1. Redshank, *Mont. Orn. Dict. & Supp.* *Bew. Brit. Birds*, vol. II. p. 71.

DIMENS. Entire length twelve inches two lines: length of the bill (from the forehead) one inch seven lines and a half, (from the gape) one inch ten lines; of the tarsus one inch ten lines; of the naked part of the tibia nine lines; of the middle toe one inch four lines and a half; of the tail two inches ten lines; from the carpus to the end of the wing six inches eleven lines: breadth, wings extended, twenty-one inches nine lines.

DESCRIPT. (*Winter plumage.*) Crown of the head, space between the eye and the bill, back of the neck, upper part of the back, scapulars, and wing-coverts, cinereous brown, darkest on the shafts of the feathers; throat, and streak above the eye, white; sides of the head, fore part of the neck, and breast, grayish white, the shafts of the feathers brown; lower part of the back, belly and abdomen, white: greater quills with their coverts dusky brown; secondaries white for a considerable portion of their length: tail and upper tail-coverts with transverse zigzag bars of white and deep brown: bill red; the tip black: irides brown: feet pale red. (*Summer plumage.*) A white streak from the base of the upper mandible to the eye: all the upper parts cinereous inclining to olivaceous brown, with a broad longitudinal dusky streak in the middle of each feather; lower part of the back white: sides of the head, throat, front of the neck, breast, and belly, white, spotted and streaked with blackish brown: tail barred with black and white, and tipped with pure white, the white passing into ash-colour on the four middle feathers: legs, and basal half of the bill, bright vermilion-red. (*Young, till after the first moult.*) Upper parts dark cinereous brown, all the feathers edged with yellowish white; on the back and scapulars this edging assumes the form of angular spots; wing-coverts dusky brown, edged and tipped with yellowish white: throat whitish, dotted with brown; neck and breast cinereous, with longitudinal narrow streaks; belly white; flanks, abdomen, and under tail-coverts, spotted with brown: tip of the tail reddish: bill livid flesh-colour; the tip brown: legs orange-yellow. (*Egg.*) Pale reddish white, tinged with green; blotched, spotted, and speckled, with dark red brown: long. diam. one inch six lines and a half; trans. diam. one inch two lines.

Not uncommon on the coast during the autumnal and winter months. Retires inland to breed. Nest placed in marshes and wet pastures. Eggs four in number. Food, insects and worms, and bivalve mollusca. *Obs.* The *Gambet* of Pennant is either this species, or a *Ruff* in winter plumage.

189. T. *Ochropus*, *Temm.* (*Green Sandpiper.*)—Upper parts greenish brown: tail barred with brown and white; the two outer feathers on each side almost entirely white: bill and legs greenish.

T. *Ochropus*, *Temm. Man. d'Orn.* tom. II. p. 651. T. *Glareola*, *Markwick in Linn. Trans.* vol. I. p. 128. tab. 11. Green Sandpiper, *Mont. Orn. Dict. & Supp.* *Bew. Brit. Birds*, vol. II. p. 85. *Selb. Illust.* vol. II. p. 75. pl. 16. f. 2.

DIMENS. Entire length nine inches eight lines: length of the bill (from the forehead) one inch three lines and a half, (from the gape) one

inch six lines; of the naked part of the tibia nine lines; of the tarsus one inch four lines; of the middle toe one inch two lines; of the tail two inches five lines; from the carpus to the end of the wing five inches six lines.

DESCRIPT. (*Winter plumage.*) Crown of the head, nape, sides of the neck, and all the upper parts, cinereous brown, tinged with olivaceous green; the feathers on the back, scapulars, and wing-coverts, with an edging of small whitish spots: from the bill to the eye a brown streak, above which is another one of white: upper tail-coverts, and all the under parts, pure white; the breast, and front of the neck, variegated with numerous longitudinal dusky streaks: quills deep brown: under wing-coverts marked with transverse undulating white bars: tail white for one-third of its length from the base; the remaining portion barred with brown and white; the two middle feathers with five broad bars of the former colour; the two next with four; the two next with three; the two next with two; the two next with only one bar, but with a small spot higher up on the exterior web; the two outer feathers almost entirely white, having only a single small spot on the exterior web: bill dusky green passing into black at the tip: irides deep brown: legs and toes greenish gray. The *summer plumage* is distinguished by the deeper tint of the upper parts, which are also more glossed with green, and more spotted on the edges of the feathers; the streaks on the fore part of the neck are likewise more distinct. (*Young of the year.*) "All the upper parts paler; the spots fewer, and of a yellowish colour; nape tinged with cinereous: sides of the breast of the same colour as the back, and marked with white spots; all the fore part of the neck, and middle of the breast, with lanceolate brown streaks: less white at the base of the tail; the bars of brown on the middle feathers broader." TEMM. (*Egg.*) Yellowish clay-colour, blotched and spotted with ash-gray and dark brown: long. diam. one inch four lines and a half; trans. diam. one inch.

An occasional visitant in this country, principally during the autumnal and winter months, but scarcely to be called common. Frequents the borders of streams, rivers, and other pieces of fresh water, but rarely resorts to the coast. Said to breed in the central parts of Europe, constructing its nest in the sand or grass by the water-side. Lays from three to five eggs. Food, insects and worms.

190. *T. Glareola*, Temm. (*Wood Sandpiper.*)—Upper parts deep brown, spotted with white: all the tail-feathers barred with brown and white; the two outer feathers on each side with the inner web entirely white: bill and legs greenish.

T. Glareola, Temm. *Man. d'Orn.* tom. II. p. 654. *Wood Sandpiper*, *Mont. Orn. Dict. & Supp. with fig. Selb. Illust.* vol. II. p. 77. pl. 16. f. 3. Long-legged Sandpiper, *Mont. Orn. Dict. Supp. App.*

DIMENS. Entire length nine inches: length of the bill not quite one inch three lines. MONT.

DESCRIPT. (*Winter plumage.*) Between the bill and the eye a dusky streak, above which is a broader one of white continued over the eye: forehead, crown, back, and wings, deep brown; the edges of the feathers on the back, scapulars, and wing-coverts, marked with white and grayish white spots: ear-coverts dusky; cheeks, nape, fore-neck, and breast, dirty

white, irregularly variegated with spots and streaks of cinereous brown; flanks with transverse undulating bars of the same colour: throat, middle of the abdomen, upper and under tail-coverts, pure white; some of the tail-coverts with fine dusky streaks on the shafts of the feathers: tail with alternate black and white bars, narrower and more numerous than in the last species; the two outer feathers with their inner webs pure white: legs greenish; much longer than in the *T. Ochropus*, with a greater portion of the tibia naked: bill black, towards the base greenish. (*Summer plumage*.) "Crown of the head, and nape, streaked longitudinally with brown and whitish; cheeks, fore part of the neck, breast, and flanks, nearly pure white, streaked longitudinally with deep brown: all the feathers of the back with a very large black spot in the centre, and with two whitish spots on each side of the webs; scapulars, when raised, appearing barred with broad dusky bands: the rest as in winter." TEMM. (*Young of the year*.) "All the deep brown part of the plumage covered with small, closely approximating, red spots: the whole of the breast waved with cinereous, with irregular brown spots: legs, and base of the bill, dirty yellowish green: tail-feathers irregularly barred." TEMM. (*Egg*.) Reddish white; the larger end nearly covered with dark red brown: long. diam. one inch five lines; trans. diam. one inch one line.

First noticed as British by Montagu, who has accurately pointed out the distinctions between this and the last species. The individual described by him was shot on the coast of South Devon early in August. He mentions another which was shot at Woolwich, at the same period of the year. Since his time individuals have been killed in the counties of Durham, Northumberland, Norfolk, and Cambridge. The species, however, must be considered as a rare and accidental visitant in this country. Common on many parts of the Continent. Breeds, according to Temminck, within the arctic circle. Constructs its nest in marshes, and lays four eggs. Food, insects and worms.

- (18.) *T. Macularia*, Temm. *Man. d'Orn.* tom. II. p. 656. Wils. *Amer. Orn.* vol. VII. p. 60. pl. 59. f. 1. *Spotted Sandpiper*, Mont. *Orn. Dict. & Supp. Selb. Illust.* vol. II. p. 84. pl. 17. *Spotted Tringa*, Edw. Glean. pl. 277.

There is no well-authenticated instance in which this species has occurred in Great Britain, although by all our authors it has been included in the British lists. The bird shot in Essex described by Edwards, and supposed by him to be the same as his *Spotted Tringa* from America, is evidently nothing more (as Mr Selby has suggested) than the common *Totanus Hypoleucos*. The same may be said of the *Spotted Sandpiper* of Bewick. (*Brit. Birds*, vol. II. p. 97.) The true *T. Macularia* is abundant in the United States, where it represents the Common Sandpiper of this country. Nevertheless, as (according to Temminck) it has certainly been killed in two or three instances on the Continent, it is still possible that it may occasionally occur in this country.

191. *T. Hypoleucos*, Temm. (*Common Sandpiper*.)

—All the upper parts cinereous brown glossed with olive; under parts pure white, without spots.

T. Hypoleucos, Temm. *Man. d'Orn.* tom. II. p. 657. Common Sandpiper, Mont. *Orn. Dict. Bew. Brit. Birds*, vol. II. p. 93. *Selb. Illust.* vol. II. p. 81. pl. 15. f. 3, & 4.

DIMENS. Entire length seven inches nine lines: length of the bill (from the forehead) one inch, (from the gape) one inch one line and

a half; of the tarsus eleven lines; of the naked part of the tibia four lines; of the tail two inches four lines; from the carpus to the end of the wing four inches three lines: breadth, wings extended, thirteen inches three lines.

DESCRIPT. (*Summer and winter plumage.*) All the upper parts cinereous brown, glossed with olivaceous green; the shafts of the feathers being of a darker tint; back and wing-coverts marked with fine transverse undulating lines of dusky brown: over the eye a whitish streak: under parts pure white, streaked on the breast and sides of the neck with cinereous brown: quills brown, with a large white spot on their inner webs, the two first excepted: the four middle tail-feathers like the back; the two next on each side tipped with white; the outer one tipped with white, and barred on the exterior web with white and brown: bill and legs grayish brown, the latter tinged with green. (*Young of the year.*) Throat, and fore part of the neck, pure white, spotted only on the sides; the white streak above the eyes broader and more distinct; the feathers on the back edged with reddish and dusky; wing-coverts tipped with red and black bars. (*Egg.*) Reddish white, spotted and speckled with umber brown: long. diam. one inch four lines; trans. diam. one inch.

A regular summer visitant, making its appearance in the Spring and departing in the Autumn. Chiefly frequents the banks of rivers and lakes. Nest constructed of moss and dry leaves, and placed in the grass by the water side. Eggs four or five in number. Utters a clear piping note in its flight. Food, worms and insects.

*** Bill rather strong, slightly recurved.*

192. *T. Glottis*, Bechst. (*Greenshank.*)—Back, scapulars, and wing-coverts, dusky brown, the feathers edged with yellowish white; lower back, and under parts, white: legs greenish.

T. Glottis, *Temm. Man. d'Orn.* tom. II. p. 659. *Greenshank*, *Mont. Orn. Dict. & App. to Supp. Bew. Brit. Birds*, vol. II. p. 67. *Selb. Illust.* vol. II. p. 86. pl. 19.

DIMENS. Entire length fourteen inches. MONT.

DESCRIPT. (*Winter plumage.*) From the upper mandible to the eye a white streak; head, cheeks, sides and back part of the neck, and sides of the breast, cinereous white, with longitudinal brown streaks; upper part of the back, scapulars, and wing-coverts, dusky brown, all the feathers broadly edged with yellowish white: lower part of the back, upper tail-coverts, throat, fore part of the neck, middle of the breast, belly, and other under parts, pure white: quills dusky; some of them spotted with white on their inner webs: tail white, with transverse bars of brown; the outer feathers entirely white, with the exception of a narrow longitudinal streak on the exterior web: bill cinereous brown: legs yellowish green; long and slender. (*Summer plumage.*) Crown, nape, cheeks, and sides of the neck, with dusky streaks; throat white; fore-neck, breast, upper part of the abdomen, and flanks, white, with oval black spots; rest of the under parts pure white: upper part of the back and scapulars deep black, the feathers on the back edged with white, those on the scapulars with reddish white spots; greater coverts reddish ash with black shafts, edged with white: the two middle tail-feathers cinereous, with transverse brown zigzag bars. (*Egg.*) Olive-brown, covered all over with dusky spots.

Occasionally observed in small flocks on the coast during the winter months, as well as in marshes and on the banks of rivers, but not a very common species. Doubtful whether it ever remains through the Summer. According to Temminck, breeds within the arctie circle. Food, small fish, worms, and molluscous animals. *Obs.* The *Cinereous Godwit* of Pennant is this species in its winter plumage.

GEN. 75. RECURVIROSTRA, Linn.

193. *R. Avocetta*, Linn. (*Scooping Avocet*).—Crown, nape, scapulars, and lesser wing-coverts, black : cheeks, and under parts, pure white.

R. Avocetta, *Temm. Man. d'Orn.* tom. II. p. 590. *Scooping Avocet*, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. II. p. 116. *Selb. Illust.* vol. II. p. 90. pl. 20.

DIMENS. Entire length eighteen inches : length of the bill (from the forehead) three inches, (from the gape) three inches two lines; of the tarsus two inches nine lines; of the tail two inches ten lines; from the carpus to the end of the wing eight inches six lines.

DESCRIPT. (*Adult male and female.*) Crown of the head, nape, two-thirds of the hind part of the neck, scapulars, lesser wing-coverts, and primary quills, deep black; all the rest of the plumage pure white: bill black: irides reddish brown: legs bluish gray. (*Young of the year, before the first moult.*) "All the black parts of the plumage tinged with brown; the brownish black on the head not extending below the occiput; scapulars edged with red, and tipped with reddish ash: legs cinereous." *TEM.* (*Egg.*) Olive-brown, blotched and spotted with brownish black: long. diam. two inches three lines; trans. diam. one inch five lines.

Common in small flocks on some parts of the coast during Winter. Breeds in the fens of Lincolnshire, and in other extensive marshy districts. Lays two eggs. Food, worms and marine insects.

GEN. 76. HIMANTOPUS, Briss.

194. *H. melanopterus*, Temm. (*Black-winged Long-shanks*).—Occiput, nape, back, and wings, black: rest of the plumage white.

H. melanopterus, *Temm. Man. d'Orn.* tom. II. p. 528. *Long-legged Plover*, *Penn. Brit. Zool.* vol. II. p. 476. *White*, *Nat. Hist. Selb.* vol. II. p. 84. *with fig.* *Mont. Orn. Dict.* *Shaw, Nat. Misc.* vol. VI. pl. 195. *Bew. Brit. Birds*, vol. I. p. 365. *Black-winged Stilt*, *Selb. Illust.* vol. II. p. 247. pl. 39*.

DIMENS. Entire length thirteen inches: length of the bill two inches six lines; of the naked part of the tibia three inches six lines; of the tarsus four inches six lines: breadth, wings extended, two feet five inches.

DESCRIPT. Occiput and nape black, the latter sometimes whitish with dusky streaks; back and wings glossy black, with greenish reflections: forehead, sides of the face, neck, and all the under parts, pure white, slightly tinged on the breast and belly with rose-red: rump white: tail cinereous gray; the outer feathers white: bill black: irides crimson:

legs vermillion red. In *very old males*, according to Temminck, the nape, and sometimes the occiput also, is perfectly white. (*Female*.) Smaller; the black on the back and wings inclining more to brown, and without the greenish gloss. (*Young*.) Crown, occiput, and nape, dusky ash; back and wings brown; all the feathers with whitish edges: legs orange-red. (*Egg*.) Pale blue, blotched and streaked with ash green and red brown: long. diam. one inch nine lines; trans. diam. one inch three lines.

A very rare and accidental visitant in this country. Sibbald mentions a pair which were killed in Scotland. Pennant has recorded another which was shot near Oxford; and White, in his *Nat. Hist. of Selborne*, speaks of five which were shot out of a flock of six on the verge of Frinsham pond, near Farnham in Surrey, in April 1779. Has also been killed in Anglesea, and more recently in Lincolnshire and Norfolk. Said to be not uncommon in the eastern parts of Europe, frequenting the borders of rivers and lakes. Breeds in the salt marshes of Hungary and Russia. Food, tadpoles, aquatic and other insects, &c.

GEN. 77. LIMOSA, Briss.

195. *L. rufa*, Briss. (*Bar-tailed Godwit*.)—Tail with eight or nine transverse dusky bars: bill curving considerably upwards: middle claw short, and without serratures: tarsi moderate.

L. rufa, *Temm. Man. d'Orn.* tom. II. p. 668. Godwit, and Red Godwit, *Bew. Brit. Birds*, vol. II. pp. 59, & 61. *Shaw, Gen. Zool.* vol. XII. p. 77. Red Godwit, *Selb. Illust.* vol. II. p. 98. pl. 22.

DIMENS. Entire length sixteen inches: length of the bill (from the forehead) three inches, (from the gape) three inches one line; of the tarsus one inch eleven lines and a half; of the tail two inches ten lines; from the carpus to the end of the wing eight inches.

DESCRIPT. (*Male and female in winter plumage*.) Crown of the head, space between the bill and the eyes, cheeks, and neck, pale ash-colour, with longitudinal streaks of brown: a broad line above the eyes, throat, breast, belly, and abdomen, white; flanks, and some of the under tail-coverts, streaked with dusky brown: upper part of the back, and scapulars, cinereous gray, the shafts of the feathers, together with a small portion of the webs on each side, brownish black; wing-coverts white, the middle of each feather deep brown: quills dusky: lower part of the back, and rump, white with dusky spots: tail with alternate transverse bars of white and blackish brown: basal half of the bill purplish red; the tip black: irides brown: legs and toes black. (*Male in summer plumage*.) Crown and nape reddish orange, with brown streaks; line above the eyes, throat, sides of the neck, and all the under parts, bright ferruginous; the feathers on the sides of the breast, flanks, and under tail-coverts, longitudinally streaked with dusky brown: back and scapulars deep black, with oval spots of pale orange-red at the edges of the feathers; wing-coverts dusky ash, edged with white: quills black with white shafts: rump white spotted with black: tail with alternate bars of brown and reddish white. In the *female*, the ferruginous tint is not so bright: the upper parts are deep brown, with cinereous undulating lines, and yellowish spots on the edges of the feathers: under parts pale reddish yellow, with the exception of the middle of the belly, which is pure white; sides of the breast, flanks, and under tail-coverts, streaked with black. Dimensions always larger than those of the male.

(*Young of the year.*) "Head, nape, back, and scapulars, deep brown, the feathers with an interrupted border of yellowish white; wing-coverts broadly edged with white: neck, breast, and flanks, reddish ash, with fine longitudinal streaks of brown; band above the eyes, throat, and belly, pure white; rump and under tail-coverts white, with large lanceolate dusky spots: tail with broad zigzag bars of brown and reddish white; the tip white: legs dusky ash." TEMM. (*Egg.*) Unknown.

Not uncommon on our coasts during the winter months, and occasionally observed in marshes inland. Said to resort to higher latitudes to breed. Food, worms, insects, and the smaller bivalve mollusca. *Obs.* The *Common Godwit* of Montagu and Bewick is this species in its winter plumage. The *Red Godwit* of this last author (but not of Montagu) is the same in its summer plumage.

196. *L. melanura*, Leisl. (*Black-tailed Godwit.*)—Tail of a uniform black, the base white: wings with a transverse white bar: bill very slightly curved: middle claw long and serrated: tarsi very long.

L. melanura, Temm. *Man. d'Orn.* tom. II. p. 664. Black-tailed Godwit, *Selb. Illust.* vol. II. p. 94. pl. 21. Jadreka Snipe, *Mont. Orn. Dict. Supp. with fig.*

DIMENS. Entire length seventeen inches six lines: length of the bill (from the forehead) four inches seven lines, (from the gape) four inches ten lines; of the tarsus three inches four lines: of the naked part of the tibia one inch eleven lines; of the tail three inches eight lines: from the carpus to the end of the wing nine inches six lines.

DESCRIPT. (*Winter plumage.*) All the upper parts of a uniform cinereous brown, with the shafts of the feathers of a somewhat deeper tint: streak over the eyes, throat, fore part of the neck, breast, and flanks, grayish white; belly, abdomen, rump, and under tail-coverts, white: quills dusky; the basal portion of some of the primaries white: greater wing-coverts ash-gray, broadly edged with white: tail white for one-third of its length from the base; the remaining portion black; the two middle feathers tipped with white: bill orange-red at the base, the tip black: legs dusky gray; the tarsi much longer than in the last species. (*Summer plumage.*) Crown of the head black, the feathers edged with bright red: eye-streak reddish white; space between the eye and the bill brown; throat and neck bright ferruginous, with brownish specks; breast and flanks the same, with transverse crescent-shaped bars of brownish black: upper part of the back, and scapulars, deep black, all the feathers tipped with reddish brown, and edged at the sides with small spots of the same colour; wing-coverts much as in winter: middle of the belly, abdomen, and base of the tail, white; lower part of the back, and remainder of the tail, deep black: base of the bill bright orange: legs black. (*Young of the year, before the first moult.*) Eye-streak, throat, base of the tail, belly, and abdomen, white: upper part of the head brown, the feathers with reddish edges; neck and breast pale red, tinged with cinereous; feathers on the back and scapulars dusky brown, surrounded by a red border; wing-coverts cinereous, broadly edged and tipped with reddish white; tips of the tail-feathers edged with white: extremity of the bill brown. (*Egg.*) Light olive-brown, blotched and spotted with darker brown: long. diam. two inches two lines; trans. diam. one inch six lines.

Much less frequent than the last species. Haunts marshes and low pastures, and is not often observed on the sea-coast. Is occasionally known to breed in the fenny districts of Cambridgeshire and Norfolk. Lays four eggs. Food, insects and their larvæ, worms, &c.

GEN. 78. SCOLOPAX, *Linn.*

(1. *RUSTICOLA*, *Vieill.*)

197. *S. Rusticola*, *Linn.* (*Woodcock*.)—Plumage, above, variegated with black, gray, and ferruginous; beneath, yellowish white, with transverse undulating bars: occiput barred transversely.

S. Rusticola, *Temm. Man. d'Orn.* tom. II. p. 673. *Woodcock*, *Mont. Orn. Dict. & Suppl. Bew. Brit. Birds*, vol. II. p. 43. *Selb. Illust.* vol. II. p. 107. pl. 23. f. 1.

DIMENS. Entire length thirteen inches nine lines: length of the bill (from the forehead) two inches nine lines, (from the gape) two inches seven lines and a half; of the tarsus one inch four lines and a half; of the middle toe one inch eight lines and a half; of the tail three inches three lines; from the carpus to the end of the wing seven inches ten lines: breadth, wings extended, twenty-four inches six lines.

SCRIPT. Forehead and crown ash-gray, tinged with rufous; from the corner of the bill to the eyes a dusky streak; occiput with four broad transverse bars of blackish brown; rest of the upper parts variegated with chestnut-brown, ochre-yellow, and ash-gray, with zigzag lines and large irregular spots of black: throat plain white; rest of the under parts yellowish white, passing into rufous on the breast and fore part of the neck, with transverse undulating bars of dusky brown: quills barred with ferruginous and black on their outer webs: tail of twelve feathers, black, the outer webs edged with rufous; the tips ash-gray above, silvery white beneath: bill flesh-colour, tinged with gray: legs livid. The *female* is somewhat larger, with the colours more dull. (*Egg*.) Pale yellowish white; the larger end blotched and spotted with ash-gray and two shades of reddish yellow brown: long. diam. one inch nine lines; trans. diam. one inch four lines.

A winter visitant, appearing regularly about the beginning of October, and departing in February or March. Frequents moist woods, and is of common occurrence in most parts of the kingdom. Occasionally known to breed with us. Nest placed on the ground, in a small hollow. Eggs four in number. Food, insects and worms.

(2. *SCOLOPAX*, *Vieill.*)

198. *S. Sabini*, *Vig.* (*Sabine's Snipe*.) — Whole plumage variegated with black and chestnut; paler beneath: tail of twelve feathers.

S. Sabini, *Vigors in Linn. Trans.* vol. XIV. p. 556. pl. 21. *Jard. and Selb. Orn.* vol. I. pl. 27. *Sabine's Snipe*, *Selb. Illust.* vol. II. p. 118. pl. 24. f. 1. *Bew. Brit. Birds*, vol. II. p. 416.

DIMENS. Entire length nine inches and three-tenths: length of the bill two inches and seven-tenths; of the tarsus one inch three lines; from the carpus to the end of the wing five inches and one-tenth. **VIG.**

DESCRIPT. Cheeks, throat, neck, and breast, brownish black, speckled with chestnut-brown; belly and abdomen grayish black, with transverse bars of chestnut-brown: back and scapulars deep black, with chestnut-brown bars and spots; under wing-coverts blackish gray: tail of twelve feathers, black at the base, ferruginous at the tip, with transverse black bars: bill brownish black, with a tinge of chestnut at the base of the upper mandible: legs brownish black. (*Egg.*) Unknown.

A rare and little known species first described by Mr Vigors from a specimen shot in the Queen's County in Ireland, in August 1822. Since then, two other individuals have occurred in Kent, a third in another part of Ireland, and a fourth near Morpeth in Northumberland. This last is in the collection of Mr Selby. Has not hitherto been met with in any other part of the world.

199. *S. major*, Gmel. (*Great Snipe*).—All the under parts spotted and barred with black: tail of sixteen feathers.

S. major, *Temm. Man. d'Orn.* tom. 11. p. 675. *Great Snipe*, *Mont. Orn. Dict. & Suppl. Selb. Illust.* vol. 11. p. 115. pl. 23. f. 2. *Bew. Brit. Birds*, vol. 11. p. 415.

DIMENS. Entire length twelve inches six lines: length of the bill two inches nine lines: breadth, wings extended, nineteen inches.

DESCRIPT. Crown of the head black, divided in the middle by a longitudinal streak of yellowish white; between the bill and the eye a dusky brown streak: cheeks, eyelids, and throat, yellowish white, speckled with brown: hind part of the neck pale yellowish brown, spotted with black: back and scapulars variegated with black and chestnut-brown, many of the feathers edged on their outer webs with yellowish white; greater coverts tipped with white: under parts white, tinged with yellowish brown, spotted and barred all over with blackish brown: quills dusky: tail of sixteen feathers, black for two-thirds of its length from the base, the remaining portion bright chestnut with black bars, the tip yellowish white: bill pale brown, inclining to flesh-colour at the base: legs greenish gray. (*Egg.*) Yellow olive-brown, spotted with two shades of reddish brown: long. diam. one inch nine lines; trans. diam. one inch two lines.

A rare and only occasional visitant in this country. Inhabits extensive marshes in the North of Europe. Nest said to be placed in such situations, amongst rushes and other herbage. Eggs three or four in number. Food, worms, insects, and small shells.

200. *S. Gallinago*, Linn. (*Common Snipe*).—Neck and breast mottled with black and pale ferruginous; belly and abdomen pure unspotted white: tail of fourteen feathers.

S. Gallinago, *Temm. Man. d'Orn.* tom. 11. p. 676. *Common Snipe*, *Mont. Orn. Dict. Selb. Illust.* vol. 11. p. 121. pl. 23. f. 3. *Snipe*, *Bew. Brit. Birds*, vol. 11. p. 50.

DIMENS. Entire length eleven inches five lines: length of the bill (from the forehead) two inches ten lines and a half, (from the gape) two inches ten lines; of the tarsus one inch two lines; of the tail two inches

three lines; from the carpus to the end of the wing five inches four lines : breadth, wings extended, seventeen inches five lines.

DESCRIPT. Upper parts very similar to those of the last species : crown black, divided by a yellowish white line; a similar streak from the base of the upper mandible over each eye; between the bill and the eye a dusky line : back and scapulars velvet-black, with transverse bars of chestnut-brown, and longitudinal streaks of ochre-yellow; wing-coverts dusky brown, edged with reddish white; quills black: chin and throat white; cheeks, neck, and upper part of the breast, mottled with black and pale ferruginous brown; flanks with white and dusky transverse bars; lower part of the breast, belly and abdomen, pure white without spots: tail of fourteen feathers, black for two-thirds of its length from the base, the remaining portion reddish brown, with black bars, the tip reddish white: bill brown, paler at the base: legs dusky gray, tinged with green. (*Egg.*) Pale yellowish white; the larger end spotted with three shades of brown: long. diam. one inch six lines; trans. diam. one inch one line.

A common inhabitant of marshes and low meadows throughout the kingdom. Is generally considered as migratory, appearing early in the Autumn and departing in the Spring. In certain districts, however, many remain annually through the breeding season. Nest placed on the ground, concealed amongst rushes and coarse grass. Eggs four or five in number. Food, worms, insects, &c.

201. *S. Gallinula*, Linn. (*Jack Snipe.*) — Crown divided longitudinally by a black band reaching to the nape: neck and breast spotted; belly and abdomen pure white: tail of twelve feathers.

S. Gallinula, *Temm. Man. d'Orn.* tom. II. p. 678. *Jack Snipe*, *Mont. Orn. Dict. Selb. Illust.* vol. II. p. 125. pl. 23. f. 5. *Judcock*, *Bew. Brit. Birds*, vol. II. p. 54.

DIMENS. Entire length eight inches six lines: length of the bill (from the forehead) one inch eight lines and a half, (from the gape) one inch seven lines: of the tarsus eleven lines; of the tail one inch eleven lines: from the carpus to the end of the wing four inches three lines.

DESCRIPT. Crown divided by a black band slightly edged with reddish brown, extending from the forehead to the nape; beneath this, and parallel with it, are two streaks of yellowish white, separated by another of black; between the bill and the eye a dusky line; throat white: front of the neck, and upper part of the breast, pale yellowish brown tinged with ash, with longitudinal spots of a deeper tint: back and scapulars black, glossed with green and purple reflections; the latter with the outer webs cream-yellow, forming two conspicuous longitudinal bands extending from the shoulders to the tail; quills dusky; wing-coverts black, edged with pale brown and white: lower part of the breast, belly and abdomen, pure white: tail of twelve feathers, dusky, edged with pale ferruginous brown: bill bluish at the base, black towards the tip: legs greenish gray. (*Egg.*) Yellowish olive; the larger end spotted with two shades of brown: long. diam. one inch three lines; trans. diam. ten lines.

Found with the last species, but not quite so plentiful. Arrives early in October and departs in March. Very rarely remains with us to breed. Nest said to resemble that of the Common Snipe. Eggs four or five. Food similar.

(3. MACRORAMPHUS, *Leach.*)

202. *S. grisea*, Gmel. (*Brown Snipe.*) — Cinereous brown (*winter*), or reddish brown variegated with black (*summer*); rump and tail white, with numerous dusky transverse bars.

S. grisea, *Temm. Man. d'Orn.* tom. 11. p. 679. *S. Noveboracensis*, *Wils. Amer. Orn.* vol. VII. p. 45. pl. 58. f. 1. (*Summer.*) Brown Snipe, *Mont. Orn. Dict.* vol. 11. *with fig. in Supp.* Brown Long-beak, *Selb. Illust.* vol. 11. p. 103. pl. 24. f. 2. *Shaw, Zool.* vol. XII. p. 61. pl. 9.

DIMENS. Entire length eleven inches: length of the bill two inches six lines. **MONT.**

DESCRIPT. (*Winter plumage.*) Between the bill and the eye a dusky streak; above that, passing over the eye, a white one; cheeks and throat white, the former with a few brown streaks; crown of the head, neck, and upper part of the breast, cinereous brown, with the shafts of the feathers somewhat darker; back and scapulars dark brown, the feathers edged with cinereous and rufous brown; primary quills dusky, the inner ones and largest coverts immediately over them, slightly tipped with white; smaller coverts above and below the bastard winglet dusky and white; the rest of the coverts cinereous brown, darkest in the middle of the feathers; secondary quills dusky brown, tipped and edged with white: lower part of the breast, belly and thighs, pure white; flanks whitish with dusky bars; lower part of the back white; rump and under tail-coverts white, with transverse crescent-shaped dusky spots; upper tail-coverts and tail thickly barred with black and white: tail of twelve feathers: bill dusky, lightest at the base: legs yellow olivaceous green.

MONT. (*Summer plumage.*) Crown of the head, nape, back and scapulars, reddish brown, variegated with black and yellowish; cheeks, and streak over the eye, pale reddish; fore part of the neck and breast reddish brown; wing-coverts cinereous, with whitish edges; belly, rump and tail, as in winter. **TEMM.** (*Young of the year.*) All the upper parts blackish brown, the feathers broadly edged with bright ferruginous; eyestreak and under parts dirty white tinged with reddish, the ferruginous tint most distinct on the breast; breast and flanks spotted with dusky brown; the middle tail-feathers tipped with red. **TEMM.** (*Egg.*) Unknown.

A native of North America. In this country an extremely rare and accidental visitant. The specimen described by Montagu was shot in the beginning of October, on the coast of Devonshire. Since then, a second, in summer plumage, has been killed at Yarmouth. Wilson states that it differs in its habits from the Common Snipe, keeping on the sea-coast, and being seldom or never seen inland. Food, worms and small bivalve mollusca. Nidification unknown.

GEN. 79. TRINGA, *Briss.*(1. MACHETES, *Cuv.*)

203. *T. pugnax*, Linn. (*Ruff.*)—Tail rounded; the two middle feathers barred; the three outer ones on each side always of one colour.

T. pugnax, *Temm. Man. d'Orn.* tom. II. p. 631. *Ruff, Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. II. p. 75. *Selb. Illust.* vol. II. p. 130. pl. 25.

DIMENS. Entire length twelve inches five lines: length of the bill (from the forehead) one inch six lines, (from the gape) one inch six lines; of the tarsus one inch ten lines; of the tail two inches eight lines; from the carpus to the end of the wing seven inches five lines.

DESCRIPT. (*Male, during the breeding season.*) Face naked, covered with warty pimples of a reddish yellow colour: sides of the occiput adorned with two tufts of elongated feathers: beneath the throat a large frill or ruff of similar feathers standing out in a very conspicuous manner: colours of the ruff extremely variable, changing occasionally even in the same individual in different seasons: yellowish white, barred with black; or entirely black, glossed with purple; in some, of a uniform yellowish brown; in others, pure white; or varied with black, white, and yellow: upper parts of the body likewise variable, in general presenting a mixture of cinereous brown, yellowish white, reddish brown, and black: sides of the breast and flanks pale reddish brown, with transverse black bars, sometimes entirely black; middle of the belly, abdomen, and under tail-coverts, white: quills dusky: the four middle tail-feathers barred with black; the rest of a uniform colour: bill yellowish orange: legs yellow. (*Male in autumn and winter.*) Face feathered: no elongated tufts on the occiput, or frill on the neck: throat, fore part of the neck, belly, and other under parts, pure white; or stained and mottled with black: sometimes black, with transverse undulating white lines: breast reddish brown, with spots of a deeper tint: upper plumage variable, generally brown with black spots, the feathers edged with reddish: greater coverts, and middle tail-feathers, barred with black and reddish brown: bill brownish: legs yellowish brown. (*Female or Reeve.*) Smaller: at all times without the ruff and occipital tufts: upper parts of the body cinereous brown, mixed with black, the black glossed with steel blue: neck and breast the same, but paler: belly and abdomen white: bill black: legs yellowish. (*Young of the year.*) Very much resembling the female in winter plumage, but with the fore part of the neck and breast of a dull reddish ash: the feathers on the head, back, scapulars, and greater wing-coverts, dusky brown, broadly edged with reddish yellow; lesser coverts edged with reddish white: throat, belly, and abdomen, pure white: bill black: feet greenish. (*Egg.*) Olive, blotched and spotted with clove and liver-brown: long. diam. one inch seven lines; trans. diam. one inch one line and a half.

A migratory species, arriving early in the Spring, and departing in September. Rather locally distributed. Principally confined to the marshes of Lincolnshire, Norfolk, and the Isle of Ely. Males very pugnacious during the breeding season. Nest usually placed on a hassock of grass, in the most swampy situations. Eggs four in number, laid in the first or second week of May. Towards the end of June, or beginning of July, the ruff on the neck of the male bird begins to fall. Food, insects and worms. *Obs.* The *Shore Sandpiper*, *Greenwich Sandpiper*, and *Equestrian Sandpiper* of Latham, are all referable to this species in different states of plumage.

(2. TRINGA, *Selb.*)

204. *T. subarquata*, *Temm.* (*Pigmy Curlew.*)—Bill slightly bent down, much longer than the head: the two

middle tail-feathers longer than the others: a considerable part of the tibia naked: tarsus fourteen lines in length.

T. subarquata, *Temm. Man. d'Orn.* tom. II. p. 609. Pigmy Curlew, *Mont. Orn. Dict. & Supp. with fig. Bew. Brit. Birds*, vol. II. p. 40. Curlew *Tringa*, *Selb. Illust.* vol. II. p. 157. pl. 26. f. 4, & 5.

DIMENS. Entire length eight inches: length of the bill one inch seven lines; of the tarsus one inch two lines: of the naked part of the tibia six lines; of the middle toe, claw included, eleven lines; from the carpus to the end of the wing five inches two lines.

DESCRIPT. (*Winter plumage.*) Face, streak above the eye, throat, belly, and other under parts, pure white; from the bill to the eye a brown streak; upper part of the head, back, scapulars, and wing-coverts, cinereous brown, with the shafts of the feathers somewhat darker; feathers on the nape streaked longitudinally with brown, and edged with whitish; fore part of the neck and breast the same, but paler: upper tail-coverts white: tail cinereous gray, the feathers edged with white: bill black: irides brown: legs dusky gray. (*Summer plumage.*) Face, streak over the eye, and throat, white, speckled with brown: crown of the head black, the feathers edged with pale reddish brown; hind part of the neck reddish brown, with longitudinal black streaks; fore part of the neck, breast, belly, and abdomen, chestnut red, with a few white feathers and brown spots sometimes intermixed, according to the period of the season: back, scapulars, and greater coverts, deep black, the edges of the feathers marked with a series of reddish brown spots, the tips yellowish gray; rest of the coverts deep brown, margined with grayish white, or (during the season of incubation) with reddish yellow: upper and under tail-coverts white, with transverse black bars. (*Young of the year, before the first moult.*) Back, scapulars, and wing-coverts, dusky ash, all the feathers broadly edged and tipped with yellowish white: breast tinged with white and pale yellowish brown, faintly streaked, but without any distinct spots: feet brown. (*Egg.*) Yellowish, with brown spots. **TEMM.**

A rare visitant in this country. Occasionally met with on the eastern and southern coasts, but not often observed inland. Common in Holland in Spring and Autumn at the periods of its migration. Sometimes breeds in that country. Nest placed on the sea-shore, or on the borders of extensive waters. Eggs four or five in number. Food, insects and worms. *Obs.* The *Red Sandpiper* of Latham and Montagu is this species in its summer plumage.

205. *T. variabilis*, Meyer. (*Dunlin.*)—Bill nearly straight, a little longer than the head: the two middle tail-feathers longer than the others, and pointed: a small part of the tibia naked: tarsus eleven lines and a half.

T. variabilis, *Temm. Man. d'Orn.* tom. II. p. 612. Dunlin and Purre, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. II. pp. 102, & 104. Dunlin, *Selb. Illust.* vol. II. p. 153. pl. 26.

DIMENS. Entire length seven inches nine lines: length of the bill (from the forehead) one inch three lines, (from the gape) one inch two lines and a half: of the tarsus eleven lines and a half; of the naked part of the tibia three lines; of the tail two inches one line; from the carpus

to the end of the wing four inches eight lines: breadth, wings extended, fourteen inches seven lines.

DESCRIPT. (*Winter plumage*.) Throat, and a streak from the base of the upper mandible to the eye, pure white: head, back of the neck, and all the upper parts, cinereous brown, the shafts of the feathers being of a darker tint; quills dusky; greater wing-coverts tipped with white: lower part of the neck and breast grayish white, mottled with pale cinereous brown; belly, vent, and under tail-coverts, pure white: rump and two middle tail-feathers dusky brown; the other tail-feathers brownish ash, edged with white: bill black: irides and legs dusky brown. (*Summer plumage*.) Throat white; cheeks, sides and fore part of the neck, and breast, reddish white, with fine longitudinal streaks of black; belly and abdomen, during the period of incubation, deep black, but afterwards mixed with white: crown of the head black, the feathers edged with reddish brown; back, scapulars, and lesser wing-coverts, deep black, all the feathers broadly edged with reddish brown, and tipped with grayish white; quills and greater coverts dusky brown, the latter tipped with white: two middle tail-feathers blackish brown; the rest cinereous edged with white. (*Spring and autumn, at the time of change*.) Throat and eye-streak white; upper parts dusky brown, the feathers edged with reddish yellow; amongst them several cinereous brown, as in winter: wing-coverts wholly cinereous brown, without any red: neck and breast reddish yellow, spotted with brown; belly white, more or less spotted and blotched with dusky brown; vent and under tail-coverts pure white. *Obs.* The *female* is somewhat larger than the *male*, and has a longer bill. In *young* birds, the bill is nearly straight. (*Egg*.) Greenish white, blotched and spotted with two shades of dark red brown: long. diam. one inch four lines and a half; trans. diam. eleven lines and a half.

A common species on all parts of the coast during the greater part of the year. At the approach of the breeding season, retires to inland marshes and the banks of rivers. Lays four eggs. Food, worms and insects. *Obs.* The *Purra* of authors is this species in its winter plumage.

206. *T. pectoralis*, Bon. (*Pectoral Sandpiper*.)—"Bill shorter than the head, compressed and reddish yellow at the base: rump black: middle tail-feathers longest: feet greenish yellow: tarsus one inch." BON.

T. pectoralis, Bon. *Syn.* p. 318. *Id. Amer. Orn.* vol. iv. p. 43. pl. 23. f. 2.* *Pectoral Sandpiper*, Nuttall, *Orn. of Unit. States*, vol. II. p. 111.

DIMENS. Entire length about ten inches: length of the bill (from the rictus) rather more than one inch; of the tarsus one inch. NUTT.

DESCRIPT. "*Summer plumage*, varied with black and rufous, beneath white; breast cinereous, strongly lined with blackish. *Winter plumage*, cinereous brown, beneath white." BON.

Mr Yarrell informs me that a single individual of this species, which is not uncommon in the United States, has been killed at Yarmouth. It is at present in the possession of Mr Hoy of that place. Its identity with the *T. pectoralis* of America was confirmed by Mr Audubon, to whom the specimen was submitted for examination.

* The above reference to Bonaparte's *American Ornithology* is on the authority of Nuttall. I have not been able to get sight of the fourth volume of that work myself.

207. *T. maritima*, Brunn. (*Purple Sandpiper*.)—Bill slightly curved at the tip, longer than the head: tibia almost entirely feathered: feet, and base of the bill, coloured.

T. maritima, Temm. *Man. d'Orn.* tom. II. p. 619. *T. nigricans*, Mont. in Linn. *Trans.* vol. IV. p. 40. pl. 2. Purple Sandpiper, Mont. *Orn. Dict. & Supp.* Purple or Rock Tringa, Selb. *Illustr.* vol. II. p. 150. pl. 26. f. 6.

DIMENS. Entire length eight inches three lines: length of the bill one inch three lines; of the tarsus ten lines and a half; of the tail two inches three lines; from the carpus to the end of the wing five inches two lines.

DESCRIPT. (*Winter plumage*.) Crown of the head, cheeks, sides and nape of the neck, dusky brown, with a tinge of ash-gray; throat white; orbits, and a spot near the anterior angle of the eye, grayish white: back and scapulars black, with purple and violet reflections, the feathers edged with deep ash-colour; wing-coverts black, edged with white: breast gray, the feathers fringed with white; rest of the under parts white, streaked on the flanks with dark ash-colour; under tail-coverts with lanceolate dusky spots: rump, and two middle tail-feathers, deep black; the other tail-feathers dusky ash, fringed with white: bill reddish orange at the base, the tip dusky: irides dusky: legs ochre-yellow. (*Summer plumage*.) "Crown of the head, back, and scapulars, violet black, the feathers with a broad edging of white mixed with a little ferruginous: fore part of the neck, breast, and belly, cinereous white, with lanceolate dusky spots; flanks, and sides of the neck, with oval spots; under tail-coverts with longitudinal streaks; middle of the belly pure white." TEMM. (*Young of the year*.) "Crown, back, scapulars, secondary quills, and middle tail-feathers, dull black, all the feathers edged and tipped with pale red; wing-coverts broadly tipped with white: sides and fore part of the neck with longitudinal streaks, the feathers edged with cinereous; some large spots disposed longitudinally on the flanks and under tail-coverts: legs and base of the bill pale yellowish." TEMM. (*Egg*.) Yellowish olive, spotted and speckled with reddish brown: long. diam. one inch three lines; trans. diam. eleven lines.

Inhabits rocky shores, and is found on many parts of the British coast during the winter months. Retires to higher latitudes to breed. A few have been observed by Mr Selby to remain for that purpose on the Fern Islands. Food, marine insects and small bivalve mollusca. *Obs.* The *Knot* of Bewick is the young of the year of this species.

208. *T. Temminckii*, Leisl. (*Temminck's Stint*.)—Bill very slightly curved at the tip, shorter than the head: tail cuneiform, the outer feathers pure white: tarsus eight lines and a half.

T. Temminckii, Temm. *Man. d'Orn.* tom. II. p. 622. Little Sandpiper, Mont. *Orn. Dict. App.* Temminck's Tringa, Selb. *Illustr.* vol. II. p. 144. pl. 27*. f. 1, & 2.

DIMENS. Entire length six inches: length of the bill (from the forehead) eight lines and a half, (from the gape) eight lines; of the tarsus

eight lines and a half or nine lines; of the naked part of the tibia four lines and a half; of the tail one inch nine lines; from the carpus to the end of the wing three inches nine lines: breadth, wings extended, eleven inches eight lines.

DESCRIPT. (*Winter plumage.*) All the upper parts deep brown, with dusky streaks on the shafts of the feathers: throat white; breast and front of the neck cinereous brown; belly, vent, and under tail-coverts, pure white: tail cuneated; the four middle feathers cinereous brown; the others whitish; the two outer ones on each side nearly pure white: bill dusky: legs olivaceous brown. (*Summer plumage.*) "Upper parts black, all the feathers broadly edged with deep ferruginous: throat white; forehead, breast, and fore part of the neck, reddish ash, with very small longitudinal black spots; the rest of the under parts, and outer tail-feathers, pure white; the two middle tail-feathers dusky brown, edged with deep ferruginous." **TEMM.** (*Young of the year, before the first moult.*) Upper parts cinereous brown, palest on the nape; all the feathers, those on the nape excepted, with a double edging of dusky brown and yellowish white: throat white; over the eyes a whitish streak; breast and sides of the neck cinereous gray, with a faint tinge of reddish; rest of the under parts white: tail-feathers, the outer one excepted, tipped with reddish: bill and legs somewhat paler than in the adult. (*Egg.*) Unknown.

A rare, and only an occasional, visitant in this country. Montagu's specimen was killed on the south coast of Devonshire. Others have been since met with in Sussex and Norfolk. Frequents the borders of lakes and rivers. Probably breeds in high northern latitudes. Food, insects and worms.

209. *T. minuta*, Leisl. (*Little Stint.*)—Bill straight, shorter than the head: tail doubly forked, the outer feathers cinereous brown edged with whitish: tarsus ten lines and a half.

T. minuta, *Temm. Man. d'Orn.* tom. II. p. 624. *Little Stint*, *Bew. Brit. Birds*, vol. II. p. 107. *Minute Tringa*, *Selb. Illust.* vol. II. p. 147. pl. 27*. f. 3, & 4.

DIMENS. Entire length six inches: length of the bill eight lines and a half; of the tarsus ten lines and a half.

DESCRIPT. (*Winter plumage.*) All the upper parts cinereous brown, with the shafts of the feathers dusky: a brown streak between the bill and the eye; above the eyes another indistinct one of white; sides of the breast cinereous, tinged with brown; throat, fore part of the neck, middle of the breast, and all the other under parts, pure white: the outer tail-feathers cinereous brown, edged with whitish; the two middle ones brown: bill and legs black. (*Summer plumage.*) Crown black, the feathers edged with reddish brown; forehead and eye-streak somewhat paler than the rest of the head: throat white; cheeks, sides of the neck, and sides of the breast, pale reddish spotted with brown; middle of the breast, and all the under parts, pure white: feathers on the back, scapulars, wing-coverts, rump, and two middle tail-feathers, deep black, all broadly edged and tipped with bright red brown; the outer tail-feathers cinereous brown, edged lighter. (*Young of the year, before the first moult.*) "Crown of the head dusky, the feathers edged with yellowish red; forehead, eye-streak, throat, front of the neck, middle of the breast, and other under

parts, pure white; between the bill and the eye a brown streak; sides of the breast reddish, mixed with cinereous brown; nape and sides of the neck cinereous and brown: back, scapulars, and wing-coverts, dusky brown: the feathers on the upper part of the back broadly edged with red, the scapulars with yellowish white; wing-coverts with a narrow edging of yellowish red: two middle tail-feathers dusky, edged with reddish ash: the others edged with white." TEMM. (*Egg.*) Reddish white ground, spotted and speckled with dark red brown: long. diam. one inch one line and a half; trans. diam. nine lines.

Like the last, a rare and occasional visitant. Habits similar. Supposed to retire northward to breed.

210. *T. Canutus*, Linn. (*Knot.*)—Bill straight, a little longer than the head, very much dilated at the tip: tail even at the extremity: a small part of the tibia naked.

T. cinerea, Temm. *Man. d'Orn.* tom. 11. p. 627. Knot and Ash-coloured Sandpiper, *Mont. Orn. Dict. & Supp.* Knot, *Selb. Illust.* vol. 11. p. 138. pl. 27. f. 2, 3, & 4.

DIMENS. Entire length ten inches: length of the bill (from the forehead) one inch four lines: of the tarsus one inch two lines; of the tail two inches six lines; from the carpus to the end of the wing six inches three lines.

DESCRIPT. (*Winter plumage.*) Crown of the head, hind part of the neck, back, and scapulars, light ash-gray, with the shafts of the feathers darker; wing-coverts the same, but tipped with white, forming a transverse bar: from the bill to the eye a dusky streak; over the eye a white one: forehead, throat, and all the under parts, white; sides and front of the neck longitudinally streaked with brown; breast and flanks with transverse bars of cinereous brown: rump and upper tail-coverts white, with transverse crescent-shaped black bars: tail-feathers cinereous, edged with white: bill and legs blackish gray: irides hazel. (*Summer plumage.*) Throat, eye-streak, sides and fore part of the neck, breast, belly, and flanks, ferruginous brown: crown of the head, and nape, ferruginous, with small longitudinal black streaks; back and scapulars deep black, all the feathers edged with bright ferruginous; on the scapulars some large spots of the same colour: abdomen white, mixed with ferruginous: upper tail-coverts white, barred with black and spotted with ferruginous: tail dusky ash, the feathers edged with whitish. (*Young of the year, before the first moult.*) Crown of the head, and nape, ash-gray, with large brown spots longitudinally disposed; back and scapulars deep cinereous gray, all the feathers with a double edging of black and yellowish white, the latter colour being outermost: a brown streak between the bill and the eye; breast with a faint tinge of reddish gray: bill greenish gray: legs greenish yellow: in other respects like the adult in winter. (*Egg.*) Pale yellowish olive, blotched and spotted with two shades of dark red brown: long. diam. one inch eight lines; trans. diam. one inch one line and a half.

A winter visitant, arriving in September and departing in April. Frequents the coast in large flocks; and, during certain parts of the year, inland fens and marshy districts. Retires northward to breed. Said to lay four eggs on a tuft of withered grass. Food, worms, marine insects, and small bivalve mollusca.

211. *T. rufescens*, Vieill. (*Buff-breasted Sandpiper*.)—Bill slender, very slightly curved, not longer than the head: tail cuneated: the outer feathers light brown, edged with white: a large portion of the tibia naked.

T. rufescens, Vieill. *Gal. Ois.* tom. 11. p. 105. pl. 238. *Varr. in Linn. Trans.* vol. xvi. p. 109. pl. 11. *Buff-breasted Tringa*, *Selb. Illust.* vol. 11. p. 142. pl. 27. f. 1.

DIMENS. Entire length eight inches: length of the bill (from the forehead) nine lines, (from the gape) one inch; of the naked part of the tibia six lines; of the tarsus one inch three lines. **YARR.**

DESCRIPT. Feathers on the top of the head dark brown, approaching to black, edged with very light brown, giving a mottled appearance; hind part of the neck light brown, with a minute dark spot in the centre of each feather; back, scapulars, and tertials, blackish brown, the edges of the feathers paler; primaries nearly black, tipped with white; the shafts white; tail-coverts brown, with lighter-coloured edges; tail cuneiform, the middle feathers black, the shafts and edges lighter; the feathers on each side light brown, inclosed by a zone of black, and edged with white: chin, sides of the neck, throat, and breast, light brown tinged with buff; abdomen, flanks, and under tail-coverts, white, but pervaded also by the buff colour of the higher parts; sides of the neck with darker coloured spots: anterior portion of the under surface of the wing rufous brown; the outer portion spotted; under coverts pure white; shafts of the primaries on their under surface pearl-white, the outer web dusky, the inner web also dusky, and plain on the part nearest the shaft, the other inner half of the web beautifully mottled with dark specks; secondary quills also mottled at their bases, and ending in sabre-shaped points, presenting a regular series of lines formed by alternating shades of white, black, and dusky bands, well defined in the adult bird, and presenting a beautifully variegated appearance, peculiar to the species: bill black: legs brown. **YARR.** (*Egg.*) Unknown.

A single individual of this species, which is a native of Louisiana in America, was shot early in September 1826, in the parish of Melbourne in Cambridgeshire, in company with some dotterel. It is now in the possession of Mr Yarrell. A second was killed at Sheringham in Norfolk, July 28, 1832. This last, which proved to be a female, is in the Norwich Museum. It appears to be a very rare species even in its native country. Habits and nidification unknown.

GEN. 80. LOBIPES, Cuv.

212. *L. hyperboreus*, Steph. (*Red Lobefoot*.)—Crown, nape, and sides of the breast, deep ash; sides of the neck ferruginous: legs greenish gray.

Phalaropus hyperboreus, Temm. *Man. d'Orn.* tom. 11. p. 709. *P. Williamsii*, Simm. in *Linn. Trans.* vol. viii. p. 264. *Red Phalarope*, Mont. *Orn. Dict. & Supp.* *Red-necked Phalarope*, Sow. *Brit. Misc.* pl. 10. *Bew. Brit. Birds*, vol. 11. p. 149. *Red Lobefoot*, *Selb. Illust.* vol. 11. p. 166. pl. 28.

DIMENS. Entire length seven inches three lines: length of the bill (from the forehead) ten lines and a half, (from the gape) eleven lines; of

the tarsus nine lines and a quarter; of the naked part of the tibia three lines; of the tail one inch eleven lines; from the carpus to the end of the wing four inches two lines.

DESCRIPT. (*Adult male in spring plumage.*) Crown of the head, nape, sides of the breast, space between the eye and the bill, as well as a little streak behind the eye, deep ash-gray; sides and front of the neck bright ferruginous; throat, middle of the breast, and all the other under parts, pure white; flanks marked with large ash-coloured spots: back, and scapulars, deep black, the feathers with broad ferruginous edges; wing-coverts black, tipped with white, forming a transverse bar: two middle tail-feathers deep black; the lateral feathers ash-gray bordered by a narrow white band: bill black: irides brown: legs greenish gray. (*Adult female.*) The ash-gray about the eyes tinged with reddish; the red on the front of the neck of less extent, and mixed with cinereous: the spots on the flanks larger and more numerous. (*Young of the year.*) "Crown of the head, occiput, nape, and a spot behind the eyes, dusky brown; back, scapulars, and two middle tail-feathers, the same, but all the feathers broadly edged with pale red: primaries and wing-coverts dusky, edged and tipped with whitish: the transverse bar on the wing not so broad as in the adult: forehead, throat, front of the neck, breast, and other under parts, white, passing into pale ash on the sides of the breast and on the flanks; a faint yellowish tinge on the sides of the neck: inside of the tarsi yellow: the outer portion, and toes, yellowish green." **TEMM.** The *adult in winter* is nearly similar to the young. (*Egg.*) Olive ground, blotched and streaked all over with dark red-brown: long. diam. one inch two lines; trans. diam. ten lines.

Abundant in the Orkneys, Hebrides, and northern parts of Scotland, but very rarely met with in England. Has been shot in Yorkshire, and in Norfolk. Frequents fresh-water lakes, on the banks of which it breeds. Lays three eggs. Food, worms and insects.

GEN. 81. PHALAROPUS, *Briss.*

213. *P. lobatus*, *Flem.* (*Gray Phalarope.*) — Gray above; beneath white (*winter*): or brown above, the feathers edged with rufous; beneath red (*summer*): a transverse white bar on the wings.

P. platyrhynchus, *Temm. Man. d'Orn.* tom. II. p. 712. *Gray Phalarope*, *Mont. Orn. Dict. Selb. Illust.* vol. II. p. 162. pl. 28. Red and Gray Phalaropes, *Bew. Brit. Birds*, vol. II. pp. 146, & 147.

DIMENS. Entire length eight inches six or nine lines. **TEMM.**

DESCRIPT. (*Winter plumage.*) Forehead and crown white; occiput, ear-coverts, and a streak down the nape of the neck, dusky gray; sides of the breast, back, scapulars, and rump, bluish ash, with the shafts of the feathers dusky; some of the scapulars tipped with white; wing-coverts the same, forming a transverse bar on the wing: throat, sides of the neck, middle of the breast, and all the other under parts, pure white: tail dusky gray, the feathers edged with cinereous: bill yellowish red at the base, dusky brown at the tip: irides reddish yellow: legs greenish gray. (*Summer plumage.*) Forehead, throat, crown, and occiput, deep grayish ash; region of the eyes, and sides of the occiput, white: sides and fore part of the neck, breast, belly, and abdomen, brick-red: nape, back, scapulars, and upper tail-coverts, dusky brown, all the feathers

broadly edged with reddish yellow; wing-coverts black, tipped with white; rump white, spotted with black. (*Young of the year, before the first moult.*) "Occiput, and a streak over the eyes, dusky; nape, back, scapulars, and upper tail-coverts, cinereous brown; the feathers on the back and scapulars broadly edged with yellowish; rump white, variegated with brown; wing-coverts edged and tipped with yellowish white; tail cinereous brown, the middle feathers with a broad yellowish edging: forehead, throat, sides and front of the neck, breast, and other under parts, pure white: bill cinereous brown: feet greenish yellow." TEMM. (*Egg.*) Stone-colour, tinged with green; blotched and spotted all over with blackish brown: long. diam. one inch two lines; trans. diam. ten lines and a half.

A rare and occasional visitant in this country, but has been killed in several instances and in widely separated localities. Said to be not uncommon in some seasons on the coast of Cornwall. Very abundant in the arctic regions, and in the eastern parts of Europe and Asia, where it is supposed to breed. Inhabits the borders of rivers and large lakes. Swims with great facility, and preys upon marine and other aquatic insects. *Obs.* The *Red Phalarope* of Bewick is this species in summer plumage.

GEN. 82. GLAREOLA, *Briss.*

214. *G. Pratincola*, Leach. (*Collared Pratincole*).—Above brown; throat and front of the neck reddish white, bounded by a black line: tail very much forked.

G. Pratincola, *Leach in Linn. Trans.* vol. XII. p. 131. pl. 12. *G. torquata*, *Temm. Man. d'Orn.* tom. II. p. 500. *Collared Pratincole*, *Selb. Illust.* vol. II. p. 213. part i. pl. 63. *Austrian Pratincole*, *Mont. Orn. Dict. Supp. with fig.*

DIMENS. Entire length ten inches. MONT.

DESCRIPT. (*Adult.*) All the upper parts brown, more or less inclining to gray, occasionally with a reddish tinge on the crown and nape: back and scapulars with faint reflections of greenish bronze: throat and front of the neck reddish or yellowish white, bounded by a black line passing upwards to the corners of the bill; space between the bill and the eye black; breast clouded with brown: under wing-coverts bright ferruginous; belly, abdomen, upper and under tail-coverts, white, the two former sometimes tinged with reddish: quills dusky brown: tail very much forked, dusky, with more or less of white at the base: bill black, the base reddish: irides reddish brown: eyes surrounded by a naked red circle: legs rufous brown. (*Young.*) Upper parts cinereous brown, shaded with brown of a darker tint; the feathers edged with reddish white: throat whitish, the surrounding black line simply indicated by a few spots: breast and belly deep gray, occasionally spotted with brown: tail but little forked; the outer feathers much shorter than in the adult bird. *Obs.* This species is subject to considerable variation of plumage: the colours are more or less intense even in individuals of the same age. (*Egg.*) White: long. diam. one inch two lines; trans. diam. ten lines.

This species, which is a native of some parts of Europe and Asia, must in this country be regarded as an extremely rare and accidental visitant. The first recorded British-killed specimen was shot near Liverpool on the 18th of May 1804, and is now in the collection of Lord Derby. A second

individual was killed by Mr Bullock in the Isle of Unst, about three miles from the northern extremity of Britain, on the 16th of August 1812. More recently (May 1827) a pair are recorded to have been shot on the Breydon-wall, Yarmouth*. Said to frequent the banks of rivers, lakes, inland seas, and extensive marshes. Builds in the reeds and other thick herbage growing in such situations. Lays three or four eggs. Flight extremely rapid. Food, principally insects, which are taken on the wing.

GEN. 83. RALLUS, Linn.

215. *R. aquaticus*, Linn. (*Water-Rail*.) — Above, olive-brown spotted with dusky; beneath, bluish gray; flanks with black and white bars.

R. aquaticus, Temm. *Man. d'Orn.* tom. II. p. 683. Rail, Mont. *Orn. Dict. & Supp. Selb. Illust.* vol. II. p. 172. pl. 29. Water-Rail, Bew. *Brit. Birds*, vol. II. p. 126.

DIMENS. Entire length eleven inches five lines: length of the bill (from the forehead) one inch seven lines, (from the gape) one inch eight lines and a half: of the tarsus one inch eight lines: of the naked part of the tibia six lines; of the middle toe, claw included, two inches; of the tail two inches three lines; from the carpus to the end of the wing four inches eight lines: breadth, wings extended, sixteen inches.

DESCRIPT. Crown of the head, and all the upper parts, olive-brown, with a dusky spot in the middle of each feather: chin whitish; sides of the head, sides and fore part of the neck, breast and belly, deep ash or lead-gray; flanks with alternate transverse bars of black and white; under tail-coverts white: quills dusky brown: tail blackish; the feathers edged with olive-brown: bill red; the culmen, and tips of both mandibles, brown: irides red: legs and toes brownish flesh colour. A white variety is said to have occurred not long since in Berkshire†. (*Egg*.) Buff white; the larger end speckled with ash-gray and orange-brown: long. diam. one inch four lines; trans. diam. one inch.

Not uncommon in most parts of the country, frequenting marshes, streams, and the banks of rivers. Remains with us the whole year. Nest placed amongst rushes and other aquatic herbage. Eggs six to ten in number. Food, insects, and molluscous animals; also vegetables.

GEN. 84. CREX, Bechst.

216. *C. pratensis*, Bechst. (*Corn-Crake*.) — Above, rufous brown, the centres of the feathers dusky; beneath, yellowish white: wing-coverts ferruginous.

Gallinula Crex, Temm. *Man. d'Orn.* tom. II. p. 686. Crake Gallinule, Mont. *Orn. Dict.* Land-Rail or Corn-Crake, Bew. *Brit. Birds*, vol. II. p. 130. Meadow or Corn-Crake, *Selb. Illust.* vol. II. p. 176. pl. 30.

DIMENS. Entire length ten inches: length of the bill (from the forehead) nine lines and a half, (from the gape) one inch one line; of the tarsus one inch five lines and a half; of the tail two inches and half a line; from the carpus to the end of the wing five inches four lines.

* Paget's *Nat. Hist. of Yarm.* p. 10.

† Loudon's *Mag. of Nat. Hist.* vol. v. p. 384.

DESCRIPT. All the upper parts dusky brown, the feathers broadly margined with reddish ash; over each eye a broad cinereous streak, prolonged down the sides of the head and neck; wing-coverts ferruginous brown; quills reddish brown, tinged with gray; under parts yellowish white, tinged with cinereous, passing into pale yellowish brown on the breast; flanks and under tail-coverts with transverse bars of pale orange-brown: upper mandible of the bill brown; lower mandible whitish: irides pale brown: legs yellowish brown, tinged with gray. (*Egg.*) Pale reddish white, spotted and speckled with ash-gray and pale red brown: long. diam. one inch six lines; trans. diam. one inch one line.

A regular summer visitant, appearing about the latter end of April, and departing in October. Is pretty generally distributed throughout the kingdom, though said to be most plentiful in the northern parts of it, and in Ireland. Frequents cornfields and meadows, particularly such as are in the neighbourhood of water. Nest on the ground, generally in high grass, rudely constructed of moss and a few dry plants. Eggs from seven to twelve in number, laid the middle of June. During the breeding season, the male is heard to utter a singular noise resembling the word *Crex*, frequently repeated. Food, insects, worms, and snails; also seeds and aquatic vegetables.

217. *C. Porzana*, Selby. (*Spotted Crake.*)—Above olive-brown, with dusky streaks and white spots; beneath cinereous olive, spotted with white.

Gallinula Porzana, Temm. *Man. d'Orn.* tom. II. p. 688. Spotted Crake, *Selb. Illust.* vol. II. p. 179. pl. 30*. f. 1, & 2. Spotted Gallinule, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. II. p. 132.

DIMENS. Entire length nine inches: length of the bill (from the forehead) nine lines and a quarter, (from the gape) ten lines and a half; of the tarsus one inch four lines and a half; of the naked part of the tibia six lines; of the middle toe, claw included, one inch eight lines; of the tail two inches; from the carpus to the end of the wing four inches eight lines; breadth, wings extended, fifteen inches two lines.

DESCRIPT. (*Adult male.*) Crown of the head, back, scapulars and rump, olive-brown, with a dusky spot in the middle of each feather, all except the first elegantly spotted and streaked with pure white: forehead, throat, and a broad streak above the eyes, ash-gray, the latter speckled with white; nape thickly spotted with black and white; cheeks cinereous, speckled with black; fore part of the neck and breast pale olivaceous, tinged with ash-gray, and spotted with white; belly and vent cinereous white; flanks marked with transverse bars of white, black, and olivaceous brown: quills brown: wing-coverts olivaceous brown, sparingly spotted with white: bill greenish yellow, passing into orange-yellow at the base: irides reddish hazel: legs and toes yellowish green. In the *female*, the cheeks have a reddish tinge, and are speckled with brown. (*Egg.*) Pale reddish white, spotted and speckled with dark red brown: long. diam. one inch three lines; trans. diam. eleven lines.

Sparingly distributed over many parts of the kingdom, frequenting marshes, the banks of rivers, and other watery places. Is usually considered as a migratory species, retiring at the approach of Winter, but it is highly probable that many individuals remain with us throughout the year. It is certainly to be met with from early in March to the middle of November. Is fond of concealing itself in the thickest covert,

amongst reeds and rushes, in which situations it is not easily roused. Nest rudely constructed of aquatic plants, and said to float upon the water. Eggs eight to twelve in number. Food, similar to that of the last species.

218. *C. Baillonii*, Selby. (*Baillon's Crake*).—Upper parts olivaceous red, with numerous white spots on the back and wings: these last reaching to half the length of the tail: bill green: legs flesh-colour.

Gallinula Baillonii, *Temm. Man. d'Orn.* tom. II. p. 692. *Jard. and Selb. Orn.* pl. 15. *Baillon's Crake*, *Selb. Illust.* vol. I. p. 182. pl. 30*. f. 3. *Shaw, Gen. Zool.* vol. XII. p. 228. pl. 27.

DIMENS. Entire length seven inches: length of the bill (from the forehead) eight lines, (from the gape) eight lines and a half; of the naked part of the tibia five lines; of the tarsus one inch and half a line; of the middle toe, claw included, one inch six lines; of the tail one inch eight lines; from the carpus to the end of the wing three inches five lines.

DESCRIPT. (*Adult male*.) Crown of the head, and back of the neck, wood-brown; throat, cheeks, front and sides of the neck, breast and belly, bluish gray: back, scapulars, and wing-coverts, yellowish brown, tinged with olivaceous, and marked with numerous irregular white spots, each surrounded by a black border: flanks, abdomen, and under tail-coverts, black, with narrow transverse white bars: bill dark green, thicker and shorter than in the next species: irides reddish: legs and toes flesh-colour. In the *female* the colours are the same, but of a paler tint. (*Young*.) Throat, and middle of the belly, white, with transverse undulations of cinereous and olivaceous brown; flanks olivaceous, spotted with white: upper parts as in the adult, but the white spots fewer in number: bill greenish brown. (*Egg*.) Light olive-brown, spotted with darker brown: long. diam. one inch one line; trans. diam. nine lines and a half.

A rare and accidental visitant in this country. A specimen, caught alive at Melbourne in Cambridgeshire, in January 1823, is now in the collection of Dr Thackeray. According to Sheppard, it has also occurred in Suffolk. Common in the southern and eastern parts of Europe. Frequent the same situations as the last species. Food similar. Nest always placed in the vicinity of water. Eggs seven or eight in number.

219. *C. pusilla*, Selby. (*Little Crake*).—Upper parts olivaceous with a few white streaks; wings without spots, reaching to the extremity of the tail: bill and legs sap-green.

Gallinula pusilla, *Temm. Man. d'Orn.* tom. II. p. 690. *Little Crake*, *Selb. Illust.* vol. II. p. 185. pl. 30*. f. 4. *Little and Olivaceous Gallinules*, *Mont. Orn. Dict. Supp. with figs.* *Little Gallinule*, *Bew. Brit. Birds*, vol. II. p. 134.

DIMENS. Entire length seven inches nine lines: length of the bill (from the forehead) eight lines and a half, (from the gape) nine lines; of the naked part of the tibia five lines; of the tarsus one inch one line and a half; of the middle toe, claw included, one inch seven lines; of the tail

one inch seven lines and a half; from the carpus to the end of the wing three inches ten lines and a half.

DESCRIPT. (*Adult male.*) Throat, sides of the head and neck, breast and belly, deep bluish gray, without spots: crown of the head, and all the upper parts, olivaceous green, with the centres of the feathers dusky; on the back a large blackish patch, marked with a few longitudinal white streaks: abdomen and flanks with indistinct transverse bars of white and brown; under tail-coverts black, barred with white: bill longer and more slender than in the last species, of a fine sap-green, somewhat reddish at the base: irides red: legs and toes sap-green. (*Adult female.*) Sides of the head pale cinereous; throat whitish; front of the neck, breast and belly, reddish ash; thighs and abdomen cinereous; under tail-coverts tipped with white; all the upper parts reddish brown; on the back a large dusky patch, sparingly spotted with white; wing-coverts olivaceous ash. (*Young.*) Colours paler: the whole of the throat whitish; fore-neck and upper breast light buff; the white marks on the upper part of the back very few in number; flanks brown, with transverse white bars. (*Egg.*) Light olive-brown, spotted with darker brown: long. diam. one inch two lines; trans. diam. nine lines and a half.

Like the last, a very rare and only occasional visitant in this country. Montagu's specimen was shot near Ashburton, in Devonshire, in 1809. Others have been since killed in Yorkshire, Derbyshire, Cambridgeshire, and Norfolk. Common in the eastern parts of Europe. Haunts and food similar to those of the last species. Nest said to be placed amongst reeds and other aquatic herbage. Eggs seven or eight in number. *Obs.* The *Olivaceous Gallinule* of Montagu is supposed to be the adult state, and the *Little Gallinule* of the same author the young, of this species.

GEN. 35. GALLINULA, *Lath.*

220. *G. chloropus*, *Lath.* (*Common Gallinule.*)—Plumage above deep olive-brown; beneath blackish gray; ridge of the wing, and under tail-coverts, white.

G. chloropus, *Temm. Man. d'Orn.* tom. II. p. 693. *Common Gallinule*, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. II. p. 137. *Selb. Illust.* vol. II. p. 188. pl. 31.

DIMENS. Entire length thirteen inches: length of the bill (from the base of the frontal disk) one inch five lines, (from the gape) one inch one line and a half; of the tarsus one inch eleven lines; of the middle toe two inches eight lines and a half; of the tail two inches eleven lines; from the carpus to the end of the wing six inches nine lines: breadth, wings extended, twenty inches ten lines.

DESCRIPT. Head, throat, neck, and upper part of the breast, dusky gray; rest of the under parts deep bluish gray, the feathers on the belly and abdomen edged with grayish white: upper parts dark olive: flanks with large longitudinal streaks of white: ridge of the wing, and under tail-coverts, pure white, the latter with a few black feathers intermixed: frontal disk, and base of the bill, red; tip of the bill yellow: irides red: legs and toes olive-green: on the naked part of the tibia a red circle. (*Young, till after the second autumnal moult.*) Crown of the head and nape, as well as the rest of the upper parts, olivaceous brown: throat, front of the neck, and a spot beneath the eye, whitish; breast, belly,

and abdomen, pale gray; flanks tinged with olivaceous: tip of the bill greenish, passing into olivaceous brown at the base: the frontal disk of small size, of a deep olivaceous brown: irides brown: legs olivaceous; the naked part of the tibia yellowish. (*Egg.*) Reddish white, sparingly spotted and speckled with orange-brown: long. diam. one inch eight lines and a half; trans. diam. one inch three lines and a half.

A common inhabitant of marshy places and the banks of rivers throughout the country. Runs swiftly, and is also a good swimmer. Nest constructed of rushes and other dry herbage; generally placed on the ground near the water's edge, but occasionally in trees. Eggs from five to eight in number, laid early in April. Food, insects, seeds, and aquatic vegetables.

GEN. 86. FULICA, *Linn.*

221. *F. atra*, *Linn.* (*Common Coot.*)—Head and neck black; back black, tinged with cinereous: beneath paler.

F. atra, *Temm. Man. d'Orn.* tom. II. p. 706. *Common Coot*, *Mont. Orn. Dict. & Suppl. Selb. Illust.* vol. II. p. 193. pl. 32. *Coot*, *Bew. Brit. Birds*, vol. II. p. 141.

DIMENS. Entire length sixteen inches: length of the bill (from the base of the frontal disk) one inch ten lines, (from the gape) one inch five lines; of the tarsus two inches two lines; of the middle toe three inches three lines; of the tail two inches; from the carpus to the end of the wing seven inches nine lines.

DESCRIPT. Head and neck deep black; back, wings, and all the upper parts, black, with a tinge of ash-gray: breast, belly, flanks, thighs, and under tail-coverts, deep bluish gray: frontal disk large, of a pure white: bill white, with a tinge of rose-red: irides crimson red: legs and toes ash-gray, tinged with greenish; naked part of the tibia orange-yellow. In *young birds*, the frontal disk is very small; the under parts of the plumage pale gray. (*Egg.*) Stone-colour, speckled with nutmeg-brown: long. diam. two inches one line; trans. diam. one inch six lines.

Common in most parts of the country, frequenting lakes, rivers, and extensive ponds. Makes a large floating nest of flags, and other aquatic plants, and lays from eight to twelve eggs. Feeds on aquatic insects and vegetables. *Obs.* The *Greater Coot* (*F. aterrima*, *Linn.*) is not distinct from the present species.

ORDER V. NATATORES.

GEN. 87. ANSER, *Briss.*(1. ANSER, *Steph.*)

222. *A. ferus*, *Steph.* (*Wild Goose*).—Bill strong and elevated; orange, the nail whitish: legs flesh-colour: wings not reaching to the extremity of the tail.

Anas Anser ferus, *Temm. Man. d'Orn.* tom. II. p. 818. Gray-Lag Goose, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. II. p. 282. Gray-Lag Wild Goose, *Selb. Illust.* vol. II. p. 261. pl. 41.

DIMENS. Entire length two feet ten inches.

DESCRIPT. Head and neck clove-brown, tinged with gray, the feathers on this last part loose and furrowed: back, scapulars, greater and middle wing-coverts, clove-brown, the feathers deeply edged with grayish white; lesser wing-coverts, and base of the primary quills, bluish gray: breast and belly grayish-white, undulated with bars of a deeper tint: rump, abdomen, and under tail-coverts, pure white: tail deep clove-brown; the middle feathers edged with white; the outer one on each side almost wholly white: bill large and elevated; orange-red, the nail whitish: irides deep brown: legs flesh-colour. (*Egg*.) Ivory white: smooth and shining: long. diam. three inches one line; trans. diam. two inches one line.

Said to have been formerly very plentiful in this country; and resident in the fens of Lincolnshire all the year. Now only met with in small flocks during the winter months, and that not very frequently. Breeds in marshes, and lays from five to ten eggs. Food, aquatic vegetables, and grain of all kinds. *Obs.* The *Domestic Goose* is usually considered as having been derived from this species, but such a circumstance is rendered highly improbable from the well known fact that the Common Gander after attaining a certain age is invariably white. Montagu also observes* that a specimen of the *Anser ferus*, which was shot in the wing by a farmer in Wiltshire, and kept alive many years, would *never associate* with the tame geese. In fact the origin of these last is unknown.

223. *A. Segetum*, *Steph.* (*Bean Goose*).—Bill long, and somewhat depressed; orange, the base and nail black: legs orange: wings extending beyond the extremity of the tail.

Anas Segetum, *Temm. Man. d'Orn.* tom. II. p. 820. Bean Goose, *Mont. Orn. Dict. & Supp. Selb. Illust.* vol. II. p. 263. pl. 42.

DIMENS. Entire length two feet six inches: length of the tarsus three inches three lines; from the carpus to the end of the wing eighteen inches six lines.

* *Orn. Dict.* vol. I. Art. *Goose-Gray-Lag*.

DESCRIPT. Head, and upper part of the neck, dark cinereous brown, the feathers of the latter having a furrowed appearance; lower part of the neck, breast and belly, ash-gray, clouded with shades of a deeper tint: back, scapulars and wing-coverts, deep clove-brown, the feathers edged with whitish: quills dark gray, passing into black at the tips; the shafts white: rump blackish brown: tail dusky, tipped with white: vent, and under tail-coverts, pure white: bill black at the base and on the nail, the middle part orange-red: irides deep brown: legs orange. *Obs.* This species is subject to some little variation of plumage: the brown tint on the upper parts prevails over the ash-colour, or the ash-colour over the brown: the secondary quills are also sometimes edged and tipped with white: occasionally, though very rarely, the nail of the bill is white, in which case it becomes extremely difficult to distinguish it from the last species, which at all times it closely resembles; it is however smaller than the *A. ferus*, and the wings when closed extend nearly half an inch beyond the tip of the tail. (*Egg.*) Dull yellowish white: long. diam. three inches five lines: trans. diam. two inches five lines.

Much more plentiful than the last species. Visits this country in large flocks at the approach of Winter, resorting to fens, and also to upland corn-fields for the sake of the green wheat. Clamorous in its flight, which, when the flock is numerous, generally assumes a figured form. Retires in May to more northern latitudes to breed. Said to make a nest in marshes, amongst the coarse herbage, and to lay from ten to twelve eggs.

224. *A. albifrons*, Steph. (*White-fronted Goose*.)—
Bill and legs orange; the former with the nail white: wings reaching a little beyond the tail: a large white space on the forehead.

Anas albifrons, Temm. *Man. d'Orn.* tom. II. p. 821. White-fronted Goose, Mont. *Orn. Dict.* Bew. *Brit. Birds*, vol. II. p. 294. *Selb. Illust.* vol. II. p. 266. pl. 43.

DIMENS. Entire length two feet three inches: length of the bill (from the gape) two inches; of the tarsus three inches: from the carpus to the end of the wing sixteen inches.

DESCRIPT. Region of the bill, and a large space on the forehead, pure white, the latter bordered with deep brown; a small white spot also under each eye: rest of the head, and neck, brownish ash: back, scapulars, and flanks, clove-brown, the feathers edged with reddish ash: quills black; greater wing-coverts tipped with white: breast, belly and abdomen, dirty white, irregularly marked with patches and transverse bars of black: tail blackish gray; the middle feathers edged with white, the outer one on each side almost wholly white: upper and under tail-coverts white: bill flesh-red, tinged with orange: the nail white: irides brown: legs yellowish orange. In *young birds*, there is no white patch on the forehead, but occasionally two or three small white spots at the base of the bill: the head and neck are tinged with reddish; the ash-coloured tints on the rest of the plumage paler than in the adult. (*Egg.*) White, tinged with buff: long. diam. two inches ten lines; trans. diam. one inch eleven lines.

Appears in small flocks during the winter months. Not uncommon in the fenny districts in the eastern parts of England. Said to breed in the Arctic Regions. Food, grain, seeds and vegetables.

(2. BERNICLA, Steph.)

225. *A. Leucopsis*, Bechst. (*Common Bernicle*.)—Forehead, cheeks, and throat, white: crown, occiput, neck and breast, black.

Anas Leucopsis, *Temm. Man. d'Orn.* tom. II. p. 823. Bernicle Goose, *Mont. Orn. Dict. & Supp. Selb. Illust.* vol. II. p. 268. pl. 44. Bernacle, *Bew. Brit. Birds*, vol. II. p. 302.

DIMENS. Entire length two feet one inch.

DESCRIPT. Forehead, cheeks and throat, white; between the bill and the eyes a dark streak; crown, occiput, neck, and upper part of the breast, black; back, scapulars and wing-coverts, undulated with transverse bars of ash-gray, black, and grayish white, this last colour occupying the tips of the feathers; quills blackish gray: under parts of the body pure white, passing into ash-gray on the flanks: tail black: irides dusky brown: bill and legs black. The *young* have the white on the forehead mottled with dusky; the feathers on the back and wings tipped with reddish; the flanks of a darker colour than in the adult: legs dusky brown. (*Egg*.) Greenish white: long. diam. three inches; trans. diam. one inch eleven lines.

Common as a winter visitant on the western coast of England, and in some parts of Ireland. Rarely seen in the southern and eastern districts. Retires early in the year to high latitudes, where it breeds. Food, seeds, grain, and aquatic and other vegetables.

226. *A. torquatus*, Frisch. (*Brent Bernicle*.)—Head, neck, and breast, black; a white patch on each side of the neck.

Anas Bernicla, *Temm. Man. d'Orn.* tom. II. p. 824. Brent Goose, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. II. p. 300. *Selb. Illust.* vol. II. p. 271. pl. 45.

DIMENS. Entire length twenty-nine inches.

DESCRIPT. (*Adult*.) Head, neck, and upper part of the breast, black; on each side of the neck about half-way down, a patch of white, mottled with black: back, scapulars, and wing-coverts, clove-brown, the feathers edged with ash-gray: middle of the belly dark gray, tinged with brown; flanks barred transversely with gray and white; vent, upper and under tail-coverts, pure white: rump, quills, and tail, black: irides dusky brown: bill and legs black. (*Young of the year*.) Head and neck dusky ash, the latter without the white patch on the sides; the feathers on the back and breast tipped with reddish brown; flanks gray, with transverse bars of reddish white. (*Egg*.) Dull white, slightly tinged with green: long. diam. two inches eleven lines; trans. diam. one inch nine lines.

Like the other species of this family, a regular winter visitant. Most abundant in the eastern and southern parts of the kingdom. Frequents the sea coast, and also inland marshes. Retires northward in February and March to breed. Said to lay from eight to ten eggs. Food, similar to that of the last.

227. *A. ruficollis*, Pall. (*Red-breasted Bernicle*).—Crown and throat black; fore-neck and breast ferruginous; lore, and a double longitudinal line on the sides of the neck, white.

Anas ruficollis, Temm. *Man. d'Orn.* tom. II. p. 826. Red-breasted Goose, Mont. *Orn. Dict.* Bew. *Brit. Birds*, vol. II. p. 280. *Selb. Illust.* vol. II. p. 275. pl. 46.

DIMENS. Entire length twenty-one inches. MONT.

DESCRIPT. Forehead, crown, list down the back of the neck, throat, and a band extending upwards to the eyes, black: between the bill and the eye, a large oval white spot; behind the eye, and surrounding a large patch of orange-brown on the side of the neck, a list of white, which is further extended, and forms a line of division between the orange-brown and black of the lower neck: fore part of the neck, and breast, fine orange-brown; the latter margined by a list of black, and another of white; a second bar of white immediately before the shoulders: back, belly, quills and tail, black: abdomen, vent, thighs, upper and under tail-coverts, pure white: greater wing-coverts black, edged with white: bill reddish brown; the nail black: legs blackish brown, tinged with red. SELB. (*Egg*.) Unknown.

A very rare and accidental visitant in this country. According to Montagu, one was shot near London in 1766, and another taken alive in Yorkshire about the same time. Other specimens are stated to have been killed in Cambridgeshire during the severe winter of 1813. It has also occurred in Norfolk. Inhabits the North of Asia, about the shores of the Frozen Sea. Appears in Russia as a regular bird of passage. Said to breed in the northern parts of that country.

* 228. *A. Ægyptiacus*, Briss. (*Egyptian Goose*).—Forehead, crown, and throat, white; a large patch encircling the eyes, and another on the breast, chestnut-red.

Chenalopex Ægyptiaca, Steph. in Shaw's *Gen. Zool.* vol. XII. part 2. p. 43. pl. 42. *Egyptian Goose*, Lath. *Syn.* vol. III. p. 453. Shaw, *Nat. Misc.* vol. xv. pl. 603. Bew. *Brit. Birds*, vol. II. p. 298.

DIMENS. Entire length two feet three inches: length of the bill two inches. LATH.

DESCRIPT. (*Male*.) Forehead, crown, and throat, white; this last somewhat spotted with chestnut: on the sides of the head, surrounding the eyes, a large patch of chestnut-red: upper part of the neck pale chestnut, becoming deeper at bottom where it unites with the back: this last, and scapulars, brownish red, with numerous transverse undulating dusky lines: wing-coverts white; the greater ones barred with black near their tips: primary quills black, the first five entirely so, the rest edged with glossy green: secondaries tinged with reddish bay, and edged with chestnut: lower back, rump and tail, black: middle of the belly white; all the rest of the under parts pale rufous ash, with narrow undulating dusky lines; on the breast a large deep chestnut-coloured spot: bill red, the tip black: eyelids red: irides yellowish: legs red. (*Female*.) "Chestnut patch round the eye smaller: chin white: the chestnut patch on the breast smaller, if not wholly wanting: lesser wing-coverts white; the others pale ash-colour, with darker edges; the lower order fringed with white, forming a bar on the wing: scapulars and secondary quills much inclined to chestnut: the rest as in the male." LATH. (*Egg*.) Dull white, tinged with buff: long. diam. two inches nine lines; trans. diam. two inches.

A native of Egypt, and other parts of Africa. Has been introduced into this country, and become partly naturalized in some places. A small flock is recorded

to have visited the banks of the Tweed, at Carham, in February 1832, and two to have been killed; but according to the conjectures of Mr Selby, they had probably made their escape from Gosforth, the seat of the Earl of Wemyss, upon the Firth of Forth, where great numbers of these birds are kept in the artificial pieces of water.

(3. PLECTROPTERUS, *Leach.*)

229. *A. Gambensis*, Briss. (*Spur-winged Goose.*)— Neck, and all the upper plumage, black, glossed with purple on the back; cheeks, throat, and under parts, white: bill and legs red.

Plectropterus Gambensis, *Steph. in Shaw's Gen. Zool.* vol. XII. part ii. p. 6. pl. 36. *Spur-winged Goose*, *Lath. Syn.* vol. III. p. 452. pl. 102. *Bew. Brit. Birds*, vol. II. p. 296.

DIMENS. Size of the *Common Goose*, but standing higher on its legs: length of the bill more than two inches. **LATH.**

DESCRIPT. Upper part of the head and neck dingy brown; ear-coverts, and sides of the throat, white, spotted with brown; the lower part of the neck, sides of the breast, and all the upper plumage, appear black, but this colour is lost, particularly in the scapulars and tertials, which are most resplendently bronzed and glossed with brilliant green, and most of the outer webs of the other feathers partake of the same hue: on the bend of the wing or wrist a strong white horny spur, turning upwards, about five-eighths of an inch (Latham says an inch and a half) in length, and pointing rather inwards: the whole of the edges of the wing, from the alula spuria to the elbow and shoulder, white: all the under parts the same: bill reddish yellow: legs and toes somewhat longer than those of the *Wild Goose*; of a red or orange-yellow. **BEW.** (*Egg.*) Unknown.

A single individual of this rare species, shot near St Germain's, Cornwall, June 20th, 1821, is now in the Newcastle Museum. No other has hitherto occurred in this country. Inhabits Gambia, and other parts of Africa. Habits and nidification unknown.

GEN. 88. CYGNUS, *Meyer.*

* 230. *C. Guineensis*, Nob. (*Guinea Swan.*)—An elevated knob at the base of the upper mandible: upper parts of the body brownish gray, the edges of the feathers paler: a black list down the nape of the neck.

Anser Guineensis, *Briss. Orn.* vol. II. p. 435. *Chinese Goose*, *Lath. Syn.* vol. III. p. 447. *Swan Goose*, *Bew. Brit. Birds*, vol. II. p. 274.

DIMENS. Entire length three feet four inches.

DESCRIPT. (*Male.*) A white streak from the corners of the mouth surrounding the base of the upper mandible: down the nape a longitudinal black stripe reaching from the occiput to the back; this last, and all the rest of the upper parts, brownish gray, with the edges of the feathers somewhat paler: fore part of the neck and breast yellowish brown; belly, abdomen, and under tail-coverts, white; flanks brownish gray, the feathers edged with white: bill with a large protuberance at the base of the upper mandible; black, the base and protuberance sometimes orange: under the throat a loose skin, forming a kind of pouch and almost bare of feathers: legs black. (*Female.*) Smaller: the frontal protuberance not so much developed. (*Egg.*) White: long. diam. three inches three lines; trans. diam. two inches two lines.

A domesticated species, common in many parts of England on artificial pieces of water. Native country somewhat doubtful. Said by some to have been brought

originally from Guinea. Of restless habits, and very clamorous. Spends much of its time on land, seldom taking to the water except for safety in times of danger. Feeds on grain and vegetables. Lays in March.

* 231. *C. Canadensis*, Steph. (*Canada Swan*).—No protuberance at the base of the bill: head and neck black; under the throat a crescent-shaped white patch.

C. Canadensis, Steph. in Shaw's *Gen. Zool.* vol. xii. part 2. p. 19. Anser Canadensis, *Faun. Bor. Am.* part ii. p. 468. Canada Goose, *Lath. Syn.* vol. iii. p. 450. Wils. *Amer. Orn.* vol. viii. p. 53. pl. 67. f. 4. Cravat Goose, *Bew. Brit. Birds*, vol. ii. p. 276.

DIMENS. Entire length forty-one inches: length of the bill (above) two inches two lines, (from the gape) two inches six lines; of the tarsus three inches seven lines; of the tail nine inches; of the wing nineteen inches six lines. *Faun. Bor. Am.*

DESCRIPT. Head, and greater part of the neck, black; under the throat a crescent-shaped white patch, passing upwards behind the eyes, and reaching nearly to the occiput: back, scapulars, wings and flanks, grayish brown, the tips of the feathers paler: lower part of the neck in front, breast, belly, and under parts, pure white: primary quills, rump and tail, black: bill and legs black. (*Egg.*) Dull dirty white: long. diam. three inches four lines; trans. diam. two inches four lines.

A native of North America, from which country it has been introduced into Europe. Is not uncommon in some parts, and may be considered as in a great measure naturalized. Small flocks are occasionally observed in England in a state of liberty and independence.

232. *C. ferus*, Ray. (*Whistling Swan*).—Bill black; the base and sides, these last to beyond the nostrils, bright yellow: plumage white: tail of twenty feathers.

Anas Cygnus, *Temm. Man. d'Orn.* tom. ii. p. 828. Whistling Swan, *Mont. Orn. Dict. & Supp. Selb. Illust.* vol. ii. p. 278. pl. 47. Wild Swan, *Bew. Brit. Birds*, vol. ii. p. 265. (Trachea,) *Linn. Trans.* vol. iv. pl. 12. f. 1.

DIMENS. Entire length five feet: length of the bill (to the forehead) four inches four lines and a half, (to the eye) five inches three lines; of the tarsus four inches; of the middle toe six inches six lines; from the carpus to the end of the wing twenty-five inches six lines: breadth, wings extended, seven feet ten inches. YARR.

DESCRIPT. The whole plumage pure white, with the exception of the head and nape, which have a faint tinge of orange-yellow: bill black; cere at the base, and a portion of the sides extending beyond the line of the nostrils, lemon-yellow; the same cere passes backward, and forms a yellow space round the eyes: irides brown: legs black. In the *female*, the bill is of a paler yellow than in the male. In *young birds*, the whole plumage is pale gray; the cere, and naked skin surrounding the eyes, pale flesh-colour; the legs reddish gray. The plumage is probably not perfected till the fourth or fifth year. (*Egg.*) Dull white, faintly tinged with greenish: long. diam. four inches one line; trans. diam. two inches eight lines.

A periodical winter visitant in the northern parts of Britain. Seldom observed southward except in very severe seasons. According to Low, some few pairs remain and breed in the Orkneys; but the greater

number retire to the Arctic Regions at the approach of Spring. Builds on the ground near water, and lays from five to seven eggs. Has a loud harsh note. Food, aquatic plants and insects.

233. *C. Bewickii*, Yarr. (*Bewick's Swan*).—Bill black; the base orange-yellow: plumage white: tail of eighteen feathers.

C. Bewickii, *Yarrell in Linn. Trans.* vol. xvi. p. 445. *Selby in Newcastle Nat. Hist. Trans.* vol. i. p. 17. *Faun. Bor. Amer.* part ii. p. 465. *Bewick's Swan*, *Selb. Illust.* vol. ii. p. 284. pl. 47*. (Trachea,) *Linn. Trans.* vol. xvi. pls. 24, & 25.

DIMENS. Entire length three feet nine inches: length of the bill (to the forehead) three inches six lines, (to the eye) four inches four lines and a half; of the tarsus three inches nine lines; of the middle toe five inches three lines; from the carpus to the end of the wing twenty inches six lines: breadth, wings extended, six feet one inch. YARR.

DESCRIPT. Closely resembling the last species, but one third smaller. The whole plumage pure white, tinged, in immature specimens, on the crown, nape, and belly, with reddish orange: bill black; the cere at the base and sides orange-yellow; but no part of this last colour advancing beyond the nostrils, (as in the *C. ferus*): irides orange-yellow: legs black: tail cuneiform, of only eighteen feathers*. The *young* are gray, and the adult plumage is not perfected till the third or fourth year. (*Egg*.) Brownish white, slightly clouded with a darker tint: long. diam. three inches seven lines; trans. diam. two inches six lines.

Like the last, a winter visitant. First distinguished as a peculiar species by Mr Yarrell, and, about the same time, by Mr Richard Wingate of Newcastle. Breeds in the Arctic Regions. Said to possess a weaker voice than that of the *C. ferus*. Habits similar.

* 234. *C. Olor*, Steph. (*Mute Swan*).—Bill red; the edges of the mandibles, nail, and a protuberance at the base, black: plumage white: tail of twenty-four feathers.

C. Olor, *Steph. in Shaw's Gen. Zool.* vol. xii. part ii. p. 15. pl. 38. *Anas Olor*, *Temm. Man. d'Orn.* tom. ii. p. 830. *Mute Swan*, *Lewin, Brit. Birds*, vol. vii. pl. 237. *Bew. Brit. Birds*, vol. ii. p. 270.

DIMENS. Entire length five feet or upwards.

DESCRIPT. The whole plumage, without exception, pure white: bill red; the edges of the mandibles, nail at the tip, nostrils, and a large protuberance on the forehead, as well as the space round the eyes, deep black: irides brown: legs black; sometimes tinged with reddish. The *female* is smaller, with the frontal protuberance not so much developed. The *young of the year* are ash-brown, with the bill and legs dusky gray. In the *second year* the bill assumes a yellowish tint, and white feathers appear intermixed with the gray ones. In the *third year*, the plumage is perfected. (*Egg*.) Dull greenish white: long. diam. four inches; trans. diam. two inches nine lines.

Well known in a domesticated or half-reclaimed state on many of our rivers and artificial pieces of water. Found wild in the eastern parts of Europe. Makes a nest amongst aquatic herbage in February or March, and lays from six to eight eggs. Food, aquatic vegetables and insects; occasionally small fish. Is destitute of the shrill voice which distinguishes the two preceding species.

* According to Mr Thompson there are twenty tail-feathers. See *Lond. & Edinb. Phil. Mag.* Oct. 1834. p. 299.

GEN. 89. TADORNA, *Leach*.

235. *T. rutila*, Steph. (*Ruddy Sheldrake*.) — Back, scapulars, and under parts, ferruginous yellow; a black collar on the neck; a large white space on the wings: bill and legs black.

Anas rutila, *Temm. Man. d'Orn.* tom. II. p. 832. Ruddy Goose, *Lath. Syn.* vol. III. p. 456. Ruddy or Casarka Sheldrake, *Selb. Illust.* vol. II. p. 293. pl. 48**. Ferruginous Duck, *Bew. Brit. Birds*, vol. II. p. 313.

DIMENS. Entire length about twenty-three inches. **SELB.**

DESCRIPT. Forehead, cheeks and chin, pale ochreous yellow; region of the eyes, crown and nape, grayish white; neck, as far as the collar, ochreous yellow, tinged with orange; collar about half an inch in width, black, glossed with green: breast, back, scapulars, and under parts of the body, deep gallstone-yellow, tinged with orange, deepest upon the breast; the feathers on the upper parts with paler edges, and the ends of the long tertials passing into sienna-yellow: lesser and middle wing-coverts white: secondary quills, and greater coverts, green, glossed with purple, forming a very large speculum: greater quills black: tail-coverts black, glossed with duck-green: bill, legs and feet, black. **SELB.** The *female*, according to Temminck, wants the black collar, and has more white about the head and neck. (*Egg*.) White. **TEMM.**

An extremely rare and accidental visitant in this country. One killed at Bryanston in Dorsetshire, in the Winter of 1776, is now in the New-castle Museum. Mr Selby has figured a second individual which was killed in the South of England, and which is in his own collection. Inhabits the eastern parts of Europe. Said to make a nest in the banks of rivers, or in burrows excavated for the purpose in deserted hillocks; occasionally in hollow trees. Lays eight or nine eggs. Food, aquatic plants and insects.

236. *T. Bellonii*, Steph. (*Common Sheldrake*.) — Back, wing-coverts, and flanks, white; a broad ferruginous band on the breast: bill, frontal protuberance, and legs, red.

Anas Tadorna, *Temm. Man. d'Orn.* tom. II. p. 833. Sheldrake, *Mont. Orn. Dict. & Supp.* *Bew. Brit. Birds*, vol. II. p. 341. Common Sheldrake, *Selb. Illust.* vol. II. p. 289. pl. 48. (Trachea.) *Linn. Trans.* vol. IV. pl. 15. f. 8, & 9.

DIMENS. Entire length twenty-three inches six lines: length of the bill (from the forehead) two inches, (from the gape) two inches three lines: of the tarsus one inch eleven lines; of the tail three inches eight lines; from the carpus to the end of the wing twelve inches three lines.

DESCRIPT. (*Male*.) Head, and upper part of the neck, deep blackish green with glossy reflections; lower part of the neck, back, wing-coverts, flanks, rump, and base of the tail, pure white: across the breast a broad band of ferruginous brown, the ends passing upwards and uniting between the shoulders: scapulars, a mesial abdominal list dilating at the vent, and tip of the tail, black; under tail-coverts pale reddish brown: primaries black: speculum rich bronzed-green: three or four of the

secondaries next the body with their outer webs rich orange-brown: bill, and a fleshy protuberance on the forehead, blood-red: irides brown: legs flesh-red, inclining to crimson. (*Female*.) Smaller: no frontal protuberance, but a whitish spot instead: the colours in general not so bright as in the male: the pectoral band, and abdominal list, narrower; the latter often interrupted by large white spots. (*Young of the year*.) Forehead, cheeks, front of the neck, back, and all the under parts, white: crown and back of the neck dusky brown, with whitish specks: the pectoral band faintly indicated by a very pale tinge of reddish on that part: scapulars dusky gray, edged with pale ash: lesser wing-coverts white, with dusky tips, having a mottled appearance: tail tipped with cinereous brown: bill reddish brown: legs livid gray. (*Egg*.) Smooth; shining white: long. diam. two inches nine lines; trans. diam. one inch eleven lines.

A common species on many parts of the British coast, where it resides all the year. Prefers flat and sandy shores. Is rarely observed inland. Breeds in rabbit-burrows, or in holes in sand-banks excavated for the purpose. Eggs eight to ten in number. Food, sea-weed, marine insects, and small bivalve mollusca.

* GEN. 90. CAIRINA, *Flem.*

* 237. *C. moschata*, *Flem.* (*Musk Duck*.)—Bill, legs, and naked skin surrounding the eyes, red: plumage various.

Anas moschata, *Linn. Syst. Nat.* tom. i. p. 199. Muscovy Duck, *Lath. Syn.* vol. III. p. 476. Musk Duck, *Bew. Brit. Birds*, vol. II. p. 317. (*Trachea*.) *Linn. Trans.* vol. xv. pl. 15. f. a.

DIMENS. Entire length twenty-four inches six lines.

DESCRIPT. Crown-feathers slightly elongated, forming a kind of tuft capable of being erected: colours of the plumage extremely variable in the domesticated state; often wholly white, the tuft on the crown excepted, which is black; or slate-gray, with more or less white on the under parts: tail of twenty feathers; the outer one on each side always white: bill, a fleshy protuberance at its base, and a naked papillose skin surrounding the eyes, red: legs short and thick; red, or reddish yellow. In the *female*, the naked papillose skin is of a paler hue, and does not cover so large a part of the face: she is also smaller than the male. (*Egg*.) Yellowish white: long. diam. two inches six lines; trans. diam. one inch nine lines.

Common in a domesticated state, throughout Europe, but not indigenous. Native country said to be America, where they are still found wild. A prolific species, laying often, and producing from eight to twelve in a brood. Nest occasionally placed in the holes of trees. Has a musky smell arising from the gland on the rump. *Obs.* The *Anas bicolor* of Donovan* is a hybrid between this species and the common *Domestic Duck*, which not unfrequently breed together.

GEN. 91. ANAS, *Linn.*

(1. ANAS, *Swains.*)

238. *A. clypeata*, *Linn.* (*Common Shoveller*.)—Head and neck glossy green; back brown; belly and abdomen brownish red; lesser wing-coverts pale blue.

A. clypeata, *Temm. Man. d'Orn.* tom. II. p. 842. Shoveller, *Mont. Orn. Dict. & Supp.* *Bew. Brit. Birds*, vol. II. p. 345. Common Shoveller, *Selb. Illust.* vol. II. p. 297. pl. 48*. (*Trachea*.) *Linn. Trans.* vol. IV. pl. 13. f. 4, & 5.

* *Brit. Birds*, vol. IX. pl. 212.

DIMENS. Entire length twenty inches: length of the bill (from the forehead) two inches eight lines, (from the gape) two inches eleven lines; of the tarsus one inch five lines; of the tail four inches; from the carpus to the end of the wing nine inches: breadth, wings extended, two feet six inches.

DESCRIPT. (*Male.*) Head, and upper half of the neck, deep green, with glossy reflections; lower part of the neck, breast, and scapulars, white; belly, abdomen, and flanks, brownish red: back, and primary quills, umber-brown: lesser wing-coverts pale blue; greater ones tipped with white, forming an oblique bar across the wings and an upper border to the speculum, which last is of a brilliant grass-green: rump, upper and under tail-coverts, brown, glossed with blackish green; sides of the rump white: tail brown, the feathers edged with white; the outer one wholly white: bill black: irides yellow: legs orange. (*Female.*) Head pale reddish brown, with fine dusky streaks; rest of the upper parts dusky brown, the feathers edged with reddish white: under parts reddish, with large brown spots: lesser wing-coverts slightly glossed with pale blue: speculum not so bright as in the male. *Obs.* During the breeding season the *male* has a red breast, in which state it is the *Red-breasted Shoveller* of Pennant: after the expiration of that season, he partially assumes the *female* plumage, which is retained till the period of the autumnal moult. (*Egg.*) White, tinged with green: long. diam. two inches two lines; trans. diam. one inch six lines.

Not a very abundant species, but met with occasionally during the winter months, principally in the eastern parts of the country. A few pairs are said to remain and breed in the marshes in Norfolk. Nest placed near the water's edge amongst aquatic herbage. Eggs twelve to fourteen in number. Food, worms and aquatic insects.

(2. CHAULIODUS, *Swains.*)

239. A. *Strepera*, Linn. (*Gadwall.*)—Back, breast, scapulars and flanks, dusky brown, marked with undulating white lines: speculum white.

A. *Strepera*, *Temm. Man. d'Orn.* tom. 11. p. 837. *Gadwall*, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. 11. p. 348. Common *Gadwall*, *Selb. Illust.* vol. 11. p. 301. pl. 51, & pl. 49*. f. 1. *Gould, Europ. Birds*, part viii. (Trachea,) *Linn. Trans.* vol. 1v. pl. 13. f. 7, & 8.

DIMENS. Entire length twenty inches: length of the bill (from the forehead) one inch nine lines, (from the gape) two inches; of the tarsus one inch six lines; of the tail three inches five lines; from the carpus to the end of the wing ten inches seven lines.

DESCRIPT. (*Male.*) Head and neck grayish white, speckled with brown; lower part of the neck, breast, and back, clove-brown, marked with crescent-shaped white lines; scapulars and flanks undulated with white and blackish brown; middle wing-coverts chestnut-brown; greater coverts, rump, and upper and under tail-coverts, black, glossed with purplish blue: speculum white: belly and abdomen white, minutely speckled with grayish brown: tail cinereous, edged with white: bill brownish black, pale beneath: legs orange. (*Female.*) Not very dissimilar to the male, but with the undulating lines and crescent-shaped bars less distinctly marked. (*Young of the year.*) "Of a uniform rusty brown above, each feather having a central mark of dusky black; the under surface white." *GOULD.* (*Egg.*) "Greenish ash-colour." *TEMM.*

A winter visitant, but not of very common occurrence. According to Temminck, is very abundant in Holland, frequenting the same situations as the common Wild Duck. Breeds in marshes, and lays eight or nine eggs. Food, fish, and aquatic insects and vegetables.

(3. *DAFILA*, *Leach*.)

240. *A. acuta*, Linn. (*Pintail*.)—A longitudinal white line on each side of the occiput; back and flanks undulated with black and grayish white; two central elongated tail-feathers black.

A. acuta, *Temm. Man. d'Orn.* tom. II. p. 838. Pintail Duck, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. II. p. 356. Common Pintail, *Selb. Illust.* vol. II. p. 311. pl. 49, & pl. 49*. f. 2. (Trachea,) *Linn. Trans.* vol. IV. pl. 13. f. 6.

DIMENS. Entire length twenty-six inches; the same, central tail-feathers excluded, twenty-four inches: length of the bill (from the forehead) two inches, (from the gape) two inches three lines; of the tarsus one inch seven lines; of the middle toe, nail included, two inches four lines; of the tail six inches; from the carpus to the end of the wing ten inches six lines.

DESCRIPT. (*Male*.) Forehead and crown umber-brown, the feathers with paler edges; rest of the head, chin, and throat, dark hair-brown, slightly glossed behind the ears with purplish green: fore part of the neck, and two lateral streaks passing upwards to the occiput, white; neck above deep blackish brown: the whole of the back, flanks, and sides of the breast, beautifully marked with transverse undulating lines of black and grayish white: scapulars black; tertials long and acuminate, velvet black, with a broadish edging of grayish white: wing-coverts and primaries hair-brown: speculum blackish green, glossed with purple, bordered above by a pale ferruginous bar, below by a white one: breast, belly, and abdomen, white; the latter minutely speckled with gray towards the vent: tail, and upper coverts, dark cinereous brown, the edges of the feathers paler; two central elongated feathers, and under coverts, black: bill black, the sides of the upper mandible bluish gray: legs blackish gray. (*Female*.) Smaller than the male: head and neck reddish brown, speckled and streaked with dusky: all the upper plumage blackish brown, the feathers edged with reddish white: under parts reddish yellow, obscurely spotted with brown: speculum dull, without the green gloss; bordered above with yellowish, beneath with whitish: tail conical, but the two middle feathers scarcely longer than the others. *Obs.* The *male* of this species (as in the case of the *Shoveller*) partially assumes the *female* plumage after the expiration of the breeding season: it is, however, not retained beyond the autumnal moult. (*Egg*.) Greenish white, tinged with buff: long. diam. two inches three lines; trans. diam. one inch seven lines and a half.

Not of unfrequent occurrence during the winter months. Breeds in higher latitudes. Said to lay eight or nine eggs. Food, aquatic insects and vegetables, fish, and molluscous animals.

(4. *BOSCHAS*, *Swains*.)

241. *A. glocitans*, Pall. (*Bimaculated Duck*.)—Before and behind the eyes an irregular patch of chestnut-brown;

back and flanks undulated with black: two middle tail-feathers somewhat elongated, straight.

Querquedula gloecitans, *Vigors in Linn. Trans.* vol. XIV. p. 559.
Bimaculated Duck, *Penn. Brit. Zool.* vol. II. p. 602. pl. 100.
 no. 287. *Bew. Brit. Birds*, vol. II. p. 362. *Bimaculated Teal*,
Selb. Illust. vol. II. p. 321. pls. 55, & 55*.

DIMENS. Length of the body, bill included, fifteen inches nine lines; of the bill (to the forehead) one inch and nine-tenths, (to the gape) two inches and one-tenth; of the tarsus one inch and six lines; from the carpus to the end of the wing eight inches and two-fifths. **VIG.**

DESCRIPT. (*Male.*) Forehead, crown, and occiput, very deep reddish brown, glossed with purplish black, passing, on the hinder part of the neck, into deep violet purple: between the bill and the eye, and behind the ear-coverts, two large irregular patches of chestnut-brown, margined and varied with white: cheeks, and sides of the neck, glossy duck-green, the rest of the upper neck, and throat, greenish black: front of the lower neck, and sides of the breast, reddish brown, with oval black spots; middle part of the breast pale reddish brown, spotted with black: ground-colour of the back pale sienna-yellow, undulated with black lines; scapulars the same, tipped with glossy Scotch blue; wing-coverts hair-brown, the lower row tipped with pale wood-brown; speculum dark green, glossed with purple: upper and under tail-coverts greenish black, glossed with purple: tail cuneiform, the two middle feathers black, narrow and acuminate, considerably longer than the others, which are hair-brown, edged with white: belly and abdomen yellowish white, with undulating black lines, darkest and most distinct upon the flanks: bill blackish gray, passing towards the base and margins, into dirty orange-yellow: legs pale orange. **SELB.** (*Female.*) Chin and throat pale buff; head and neck the same, but spotted and streaked with black, the spots largest and most distinct upon the crown; lower neck, and sides of the breast, pale yellowish brown, with blackish brown spots; middle of the breast, and belly, white; abdomen white, with faint hair-brown spots; flanks variegated with yellowish brown and blackish brown: upper parts blackish brown, the feathers deeply edged with reddish white and yellowish brown; lesser wing-coverts hair-brown, the lower tier deeply tipped with pale reddish brown: upper half of the speculum green, with purple reflections; the lower half velvet-black, with white tips to the feathers: quills and tail hair-brown, the latter edged with white and reddish white: legs orange-yellow. **SELB.**

Very little is known of this extremely rare species, which was first described by Pennant from a male specimen taken in a decoy in 1771. Two other individuals, a male and female, occurred near Maldon in Essex, in the Winter of 1812-13. These last, which are described in the Linnæan Transactions by Mr Vigors, are now in the Museum of the Zoological Society. Habits and nidification unknown. Stated by Pallas to be a native of Siberia, frequenting Lake Baikal and the River Lena.

242. *A. Boschas*, Linn. (*Mallard.*)—Head, and upper part of the neck, deep green; below the green a white collar: four middle tail-feathers (in the *male*) recurved.

- A. Boschas, *Temm. Man. d'Orn.* tom. 11. p. 835. Common Duck, *Mont. Orn. Dict. & Supp.* Common Wild Duck, *Selb. Illust.* vol. 11. p. 305. pls. 50 & 50*. Mallard, *Bew. Brit. Birds*, vol. 11. p. 325. (Trachea,) *Linn. Trans.* vol. 1v. pl. 13. f. 10.

DIMENS. Entire length twenty-four inches: length of the bill (from the forehead) two inches three lines, (from the gape) two inches seven lines; of the tarsus one inch eleven lines; of the middle toe, nail included, two inches five lines; of the tail three inches five lines; from the carpus to the end of the wing ten inches nine lines: breadth, wings extended, three feet.

DESCRIPT. (*Male.*) Head, and upper half of the neck, deep emerald-green, approaching to black on the cheeks and forehead; beneath the green a white collar; rest of the neck, and breast, dark chestnut; upper part of the back, wing-coverts, and primary quills, cinereous brown of different tints, the first with fine transverse lines of gray; rump, upper and under tail-coverts, blackish green; scapulars, flanks, abdomen, and sides of the rump, grayish white, with fine transverse undulating lines of clove-brown; some of the outer scapulars chestnut, with the transverse lines darker; speculum deep Prussian-blue, with purple and green reflections; bounded above and below by a double border, the inner one velvet-black, the outer one white: tail grayish brown, all the feathers bordered with white: bill greenish yellow: irides reddish brown: legs orange. (*Female.*) Smaller: all the upper parts umber-brown, of different shades, the feathers edged with pale reddish brown: head and neck with dusky streaks; throat whitish; breast and under parts yellowish gray, obscurely spotted and streaked with brown: speculum as in the male: four central tail-feathers straight: bill greenish gray. The *young males, till after the first moult*, resemble the females. (*Egg.*) Smooth; greenish white: long. diam. two inches three lines and a half; trans. diam. one inch seven lines.

A common species in most parts of the country during the winter months, some few remaining to breed. Frequents lakes, marshes, and rivers. Nest generally placed on the ground amongst aquatic herbage; sometimes in trees. Eggs ten or twelve in number. The young are called *Flappers*. Food, insects, slugs, grain, and aquatic vegetables. *Obs.* The *Domestic Duck* derives its origin from this species.

- * (19.) *A. adunca*, Linn. (*Hook-billed Duck.*) Lath. Syn. vol. 111. p. 495. Don. *Brit. Birds*, vol. 1x. pl. 218.

Differs from the *Common Mallard* in the bill being broader, longer, and inclined more downwards at the tip. In other respects similar. It does not seem to be satisfactorily determined, whether it be a peculiar species, or only a variety of that last described; probably, however, the latter. Not uncommon in the domestic state.

243. *A. Querquedula*, Linn. (*Garganey.*) — Crown, occiput, and list on the nape, dark brown; a white stripe from the eye down each side of the neck: speculum grayish green.

- A. Querquedula*, *Temm. Man. d'Orn.* tom. 11. p. 844. Garganey, *Mont. Orn. Dict. & Supp.* *Bew. Brit. Birds*, vol. 11. p. 372. Garganey Teal, *Selb. Illust.* vol. 11. p. 318. pl. 53. (Trachea,) *Linn. Trans.* vol. 1v. pl. 13. f. 2, & 3.

DIMENS. Entire length sixteen inches six lines.

DESCRIPT. (*Male*.) Crown, occiput, and a list down the back part of the neck, dark amber-brown; throat black; over each eye a band of pure white, prolonged down the sides of the neck; cheeks, and upper part of the neck, chestnut-brown, with fine longitudinal streaks of white; lower part of the neck, and breast, pale yellowish brown, beautifully marked with crescent-shaped black bars: back grayish black, the feathers edged with ash-colour and yellowish brown: scapulars long and narrow, black, with a broad central white streak; wing-coverts bluish ash: speculum grayish green, bordered above and below by a white bar: belly white; flanks marked with transverse undulating black lines: vent, upper and under tail-coverts, yellowish white, spotted with black: tail dusky gray, the edges of the feathers lighter: irides light hazel: bill and legs blackish gray. (*Female*.) Smaller: throat white: head, neck, and upper parts of the body, brown, approaching to dusky, the feathers with pale whitish edges: the streak behind the eye very faint and ill-defined: lower part of the breast, and belly, white; flanks and abdomen spotted with brown: wing-coverts dark ash-gray; speculum dull, the green tinge almost wanting. The *young males* resemble the females: as they advance to maturity, the plumage assumes a mixed character. (*Egg*.) Buff-colour: long. diam. one inch nine lines; trans. diam. one inch three lines.

Met with in small numbers principally during the winter and spring months, and probably remains to breed in some parts of the country, but this last point does not appear to have been fully ascertained. Frequents marshes and the reedy banks of rivers. Nest placed on the ground in situations of the above nature. Eggs ten or twelve. Food, slugs, insects, seeds, and aquatic plants.

244. *A. Crecca*, Linn. (*Teal*.) — Crown, cheeks, and neck, ferruginous brown; from the eyes to the nape a broad longitudinal band of green: speculum, half green and half black.

A. Crecca, *Temm. Man. d'Orn.* tom. II. p. 846. *Teal*, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. II. p. 374. Common Teal, *Selb. Illust.* vol. II. p. 315. pl. 54. (*Trachea*.) *Linn. Trans.* vol. IV. pl. 13. f. 1.

DIMENS. Entire length fourteen inches six lines: length of the bill (from the forehead) one inch five lines, (from the gape) one inch eight lines; of the tarsus one inch two lines; of the tail three inches two lines; from the carpus to the end of the wing seven inches five lines.

DESCRIPT. (*Male*.) Crown of the head, cheeks, front and sides of the neck, ferruginous brown; on the sides of the head, inclosing the eye, a large patch of deep green, passing off backwards to the nape in the form of a broad band; sides of the lower part of the neck, back, scapulars, and flanks, beautifully marked with transverse undulating lines of black and white; some of the longer scapulars cream-yellow, with a portion of their outer webs velvet-black: lower part of the neck in front, and breast, reddish white, with round black spots: wing-coverts brown, tinged with gray; speculum deep green in the middle, velvet-black at the sides, bordered above by a broad white bar: belly and abdomen white: under tail-coverts blackish brown, bordered at the sides with yellowish white: tail cuneiform, brown, the feathers edged with white: bill dusky gray: irides light hazel: legs brown, with a tinge of ash-gray. (*Female*.) Extremely similar to that of the last species, but may always be distinguished by the speculum, which is of the same dark black and green

colours as in the male: head, neck, and all the upper parts, dusky brown, the feathers more or less broadly edged with pale reddish brown; throat, cheeks, and a band behind the eyes, yellowish white, spotted with black: under parts yellowish white. The *young males* resemble the female. (*Egg.*) White, tinged with buff: long. diam. one inch nine lines; trans. diam. one inch four lines.

A common species during the winter months, appearing in small flocks, and frequenting fresh waters. Remains to breed in some parts of the country. Nest (according to Selby) composed of rushes and other aquatic grasses, and lined with down. Eggs ten to twelve in number. Food, similar to that of the Garganey. *Obs.* The name of *Summer Teal* appears to be applied in some places indiscriminately to this and the last species, when met with during the Spring of the year.

GEN. 92. MARECA, Steph.

245. *M. Penelope*, Selb. (*Wigeon*.) — Forehead yellowish white; rest of the head, and neck, chestnut-red: back and flanks undulated with black and white.

Anas Penelope, *Temm. Man. d'Orn.* tom. II. p. 840. *Wigeon*, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. II. p. 350. Common *Wigeon*, *Selb. Illust.* vol. II. p. 324. pl. 52. (*Trachea*), *Linn. Trans.* vol. IV. pl. 13. f. 9.

DIMENS. Entire length twenty inches: length of the bill (from the forehead) one inch six lines, (from the gape) one inch nine lines; of the tarsus one inch six lines; of the tail three inches ten lines; from the carpus to the end of the wing ten inches three lines.

DESCRIPT. (*Male.*) Forehead and crown cream-yellow; rest of the head, and upper part of the neck, chestnut-red, the cheeks speckled with black: throat black; lower part of the neck, and breast, vinaceous red: back and flanks marked with transverse undulating lines of black and white; scapulars black, edged with white; wing-coverts white, some of the lesser ones nearest the body pale grayish brown: quills cinereous brown: speculum composed of three bars, the middle one glossy green, the upper and under ones black: belly and abdomen white: tail cuneiform; the two middle feathers pointed, and considerably longer than the others; of a blackish gray colour: under tail-coverts black: bill bluish gray, the tip black: irides hazel: legs dusky lead-colour. (*Female.*) Head and neck rufous brown, speckled with dusky; back and scapulars dusky brown, the feathers with reddish edges; wing-coverts brown, edged with whitish; speculum without the gloss of dark green: breast, belly, and abdomen, much as in the male: flanks rufous brown, the tips of the feathers inclining to ash-gray: bill and legs dusky gray. *Obs.* This species appears to be subject to considerable variation of plumage. In *very old males*, according to Temminck, the forehead alone is yellowish white, this colour not extending over the crown. *Young males of the year* resemble the female. (*Egg.*) Brownish white: long. diam. two inches; trans. diam. one inch seven lines.

A winter visitant of common occurrence in most parts of the country. Frequents rivers and marshes, as well as inlets of the sea. Keeps in flocks, and has a peculiar whistling note. Breeds in the north of Europe, and is said to lay eight or nine eggs. Food, aquatic insects and vegetables, mollusca, and small fish. In confinement, has been known to breed with the Pintail and the Common Duck.

* GEN. 93. DENDRONESSA, *Swains.*

* 246. *D. Sponsa*, Swains. (*Summer Duck*.)—A pendent occipital crest, variegated with green, purple, and white.

D. Sponsa, *Faun. Bor. Amer.* part ii. p. 446. *Summer Duck*, *Lath. Syn.* vol. iii. p. 546. *Wils. Amer. Orn.* vol. viii. p. 97. pl. 70. f. 3. (Trachea,) *Mont. Orn. Dict. Supp. App.* last plate, f. 5.

DIMENS. Entire length twenty-one inches: length of the bill (above) one inch four lines and a half, (to gape) one inch ten lines; of tarsus one inch six lines; of middle toe and claw two inches; of tail four inches six lines; of folded wing nine inches. *Faun. Bor. Am.*

DESCRIPT. (*Male*.) Head above, and space between the eye and the bill, glossy dark green; cheeks, and a large patch on the sides of the throat, purple, with blue reflections; pendent occipital crest of green and auricle purple, marked with two narrow white lines, one of them terminating behind the eye, the other extending over the eye to the bill; sides of the neck purplish red, changing on the front of the neck and sides of the breast to brown, and there spotted with white: scapulars, wings and tail, exhibiting a play of duck-green, purple, blue, and velvet-black, colours; interscapulars, posterior part of the back, rump, and upper tail-coverts, blackish green and purple; several of the lateral coverts reddish orange; a hair-like, splendid, reddish purple tuft on each side of the rump; the under coverts brown: chin, throat, a collar round the neck, a crescentic bar on the ears, the middle of the breast, and whole of the abdomen, white: flanks yellowish gray, finely undulated with black; the tips of the long feathers, and also of those on the shoulder, broadly barred with white and black: inner wing-coverts white, barred with brown: almost all the coloured plumage shows a play of colours with metallic lustre: bill red; a space between the nostrils, its tip, margins, and lower mandible, black: legs orange. *Faun. Bor. Am.* (*Female*.) Without the fine lines on the flanks, and the hair-like tufts on the sides of the rump: crest shorter: the plumage less vivid, particularly about the head, where it is mostly brown. *Faun. Bor. Am.* (*Egg*.) Smooth; yellowish white: long. diam. one inch ten lines; trans. diam. one inch six lines.

A pair of these birds, male and female, were shot about two or three years since near Dorking in Surrey, but as the species is strictly a native of America, and is often kept in aviaries, its claims to a place in the British Fauna must be considered as extremely doubtful. It is probable that the above individuals had escaped from confinement. Common in Mexico, and other parts of North America; breeding in hollow trees. Said to lay thirteen eggs. Food, according to Wilson, principally acorns, seeds of the wild oats, and insects.

GEN. 94. SOMATERIA, *Leach.*

247. *S. mollissima*, Leach. (*Eider Duck*.)—Lateral prolongations at the base of the bill in the form of two narrow flat lamellæ: bill and legs greenish gray.

Anas mollissima, *Temm. Man. d'Orn.* tom. ii. p. 848. *Eider Duck*, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. ii. p. 305. Common Eider, *Selb. Illust.* vol. ii. p. 338. pls. 70, & 70*. (Trachea,) *Linn. Trans.* vol. xii. pl. 30. f. 1, & 2.

DIMENS. Entire length twenty-four inches: length of the bill (from the forehead) two inches three lines, (from the gape) two inches ten lines; of the tarsus two inches; of the tail four inches; from the carpus to the end of the wing eleven inches six lines.

DESCRIPT. (*Male*.) Contour of the frontal lamellæ, forehead and crown, taking in the eyes, black, glossed with violet; a longitudinal band on the top of the head, cheeks, throat, and front of the neck, whitish; occiput, nape, and a large patch beneath the temples, greenish white; lower part of the neck, back, scapulars, lesser wing-coverts,

curved tertiaries, and sides of the rump, pure white; breast white, with a tinge of vinaceous red: belly, abdomen, and rump, deep black: quills and tail-feathers dark ash-gray: bill dull green: irides brown: legs dull greenish gray. (*Female*.) The whole plumage reddish brown, with transverse black bars: head and back part of the neck marked with dusky streaks: wing-coverts black, edged with ferruginous: across the wing two rather indistinct white bars: belly and abdomen dark reddish brown, with obscure transverse black bars. *Young males* are said not to attain to maturity till the *fourth year*. According to Montagu, during the *first year*, the back is white, and the usual parts, except the crown, black: but the rest of the body variegated with black and white. In the *second year*, the crown of the head is black, and the neck and breast spotted with black and white. (*Egg*.) Smooth, shining, olive-green: long, diam. three inches; trans. diam. two inches one line.

Common in the northern parts of Britain, particularly in the Scotch Islands, where it breeds. Rarely observed southwards; but has been killed in a single instance on the coast of Devon. Nest placed on the ground, near the edge of the sea, or on projecting rocks, formed of seaweed, and copiously lined with down, which the female plucks from her body. Eggs five or six in number. Food, marine plants and insects, small fish, and bivalve mollusca.

248. *S. spectabilis*, Leach. (*King Duck*.) — Lateral prolongations at the base of the bill in the form of two elevated, compressed tubercles: bill red: legs ochre-yellow.

Anas spectabilis, Temm. *Man. d'Orn.* tom. II. p. 851. *King Duck*, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. II. p. 310. *King Eider*, *Selb. Illust.* vol. II. p. 343. pl. 71. (Trachea.) *Linn. Trans.* vol. XV. pl. 15. f. c & d.

DIMENS. Entire length twenty-four inches six lines: length of the bill (from the forehead) one inch two lines, (from the gape) two inches seven lines; of the tarsus one inch ten lines and a half; of the tail four inches three lines; of the wing eleven inches six lines: height of the frontal tubercles one inch six lines; breadth of ditto one inch. *Faun. Bor. Am.*

DESCRIPT. (*Male*.) Contour of the frontal tubercles, under eyelid, edge of the upper one, and two converging bands on the throat, meeting on the chin, rich velvet-black: crown, occiput, and nape, fine bluish gray: cheeks and ear-coverts glossy pea-green, passing into dull white towards the chin and neck; a whitish line from the corners of the eye down the sides of the occiput reaching to the nape; lower part of the neck, and breast, white tinged with buff; upper part of the back, wing-coverts, and a large space on each side of the rump, pure white: scapulars, secondaries, curved tertiaries, lower part of the back, rump, tail-coverts, belly, and abdomen, deep black: greater quills and tail blackish brown: bill vermilion-red: frontal tubercles, and base of the lower mandible, orange: legs ochre-yellow. (*Female*.) "Exactly resembling the female *Eider*, except that the frontal plates of the upper mandible, instead of being almost horizontal, are more nearly vertical. The bill is also shorter than that of the *Eider*." *Faun. Bor. Am.* (*Young male*.) "Head and neck dusky yellowish gray, crowded with blackish spots: upper plumage mostly pitch-black, with yellowish brown edgings: breast and flanks yellowish brown, spotted and barred with black; belly the

same colours intimately mixed: bill as in the female." *Faun. Bor. Am. (Egg.)* Greenish yellow white: long. diam. two inches nine lines; trans. diam. one inch nine lines.

A native of high northern latitudes, but met with occasionally in small numbers in some of the Scotch Islands. Mr Bullock is stated by Montagu to have found it breeding in Papa Westra, one of the Orkneys, in the latter end of June. The nest was on a rock impending the sea. Eggs six in number, covered, as in the case of the last species, with the down of the female. Very rare on the English coast, but has been killed at Aldborough in Suffolk.

GEN. 95. OIDEMIA, *Flem.*

249. *O. fusca*, *Flem.* (*Velvet Scoter.*)—Black; a white speculum on the wings: legs red.

Anas fusca, *Temm. Man. d'Orn.* tom. II. p. 854. *Velvet Duck*, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. II. p. 320. *Velvet Scoter*, *Selb. Illust.* vol. II. p. 333. pl. 67. (Trachea,) *Linn. Trans.* vol. XVI. pl. 21. f. 2, & 3.

DIMENS. Entire length twenty-three inches: length of the bill (from the forehead) one inch eight lines, (from the gape) two inches seven lines; of the tarsus one inch nine lines; of the tail three inches ten lines; from the carpus to the end of the wing ten inches ten lines.

DESCRIPT. (*Male.*) A small crescent-shaped spot beneath the eyes, and speculum on the wings, white; all the rest of the plumage deep velvet-black: bill orange; a protuberance at the base, nostrils, and margins of both mandibles, black: legs and toes scarlet-red; membranes black. (*Female.*) Similar to the male, but smaller, with the upper plumage more inclining to brown; under parts gray, spotted and streaked with dusky brown; between the bill and the eye, and on the orifice of the ears, a grayish white spot: bill blackish brown; slightly inflated at the base: legs and toes dull red. (*Egg.*) Buff-colour: long. diam. two inches ten lines; trans. diam. one inch ten lines.

A periodical winter visitant, but not very numerous in the southern parts of the country. Frequents the sea-coast, and is very rarely observed inland. Breeds, according to Temminck, in the Arctic Regions, and lays from eight to ten eggs. Dives well in search of its food, which consists principally of bivalve mollusca.

250. *O. nigra*, *Flem.* (*Black Scoter.*)—The whole plumage black; no speculum on the wings: legs dusky gray.

Anas nigra, *Temm. Man. d'Orn.* tom. II. p. 856. *Scoter*, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. II. p. 322. *Black Scoter*, *Selb. Illust.* vol. II. p. 329. pl. 68.

DIMENS. Entire length eighteen inches: length of the bill (from the forehead) one inch eight lines, (from the gape) two inches; of the tarsus one inch nine lines; of the tail three inches four lines; from the carpus to the end of the wing eight inches nine lines: breadth, wings extended, two feet nine inches. (*Female.*)

DESCRIPT. (*Male.*) The whole plumage, without exception, deep velvet-black, with glossy reflections: bill, and a globular protuberance at the base, black; the latter divided longitudinally by a mesial band of

orange yellow, which, passing onwards, spreads over half the bill, but does not reach the tip by half an inch: irides brown: orbits yellow: tarsi and toes dusky gray; membranes black. (*Female*.) Crown, occiput and nape, dusky brown; cheeks, throat and breast, dull brown, with a mixture of ash-gray: back, wings, and under parts of the body, deep brown, the feathers edged at the tips with whitish brown: bill somewhat elevated at the base, but without the globular protuberance; of a blackish colour; the nostrils, and a spot towards the tip, yellowish: orbits brown. (*Young male, during the first year*.) "Resembles the adult female, but the colours are somewhat paler: space between the eye and the bill, crown, occiput, nape and breast, deep brown; beneath the eyes, sides and fore part of the neck, pure white; the rest of the plumage sooty-brown: bill elevated at the base; livid brown; the nostrils flesh-colour: legs dirty yellowish green; membranes dusky." TEMM. (*Egg*.) Pale buff-colour: long. diam. two inches six lines; trans. diam. one inch nine lines.

Like the last a winter visitant. Not uncommon on most parts of the British coast, but seldom observed inland. Breeds in high northern latitudes. Dives well, and feeds principally on shell-fish.

- (20.) *O. perspicillata*, Steph. (*Surf Scoter*.) Faun. Bor. Amer. part ii. p. 449. Selb. Illust. vol. 11. p. 335. pl. 69. *Anas persp.*, Temm. Man. d'Orn. tom. 11. p. 853. Wils. Amer. Orn. vol. viii. p. 49. pl. 67. f. 1.

According to Temminck, this species, which is plentiful in North America, has occurred occasionally in the Orkneys. It does not, however, appear to have been met with by any of our own naturalists, nor am I aware that any native specimens exist in our collections. It is distinguished from the *O. nigra*, which it most nearly resembles, by a frontal band, and a large patch on the nape, of pure white: rest of the plumage black: no speculum.

- (21.) *O. leucocephala*, Steph. (*White-headed Scoter*.) Flem. Brit. An. p. 119. *Anas leuc.*, Temm. Man. d'Orn. tom. 11. p. 859.

Dr Fleming is of opinion that the *female Scoter* described by Montagu in the *Supp.* to the *Orn. Dict.* and the *White-throated Duck* of Pennant (*Brit. Zool.* vol. 11. pl. 98.) are referable to this species, to which he has accordingly given a place in the British Fauna. It is, however, very probable, that the individuals above alluded to may have been only immature specimens of the *O. nigra*, which has generally more or less white on the head and neck during the first year. There is no good authority for considering the *O. leucocephala* as British. It inhabits the eastern parts of Europe, and, according to Temminck, is never found in Holland.

GEN. 96. FULIGULA, Ray.

251. *F. rufina*, Steph. (*Red-crested Pochard*.)—Head, throat, and crest, reddish brown; lower part of the neck, breast, belly and abdomen, deep black; flanks, a large spot on the shoulders, and speculum on the wing, white: bill and legs red.

- F. rufina*, Steph. in Shaw's *Gen. Zool.* vol. xii. part ii. p. 188. pl. 54. *Anas rufina*, Temm. Man. d'Orn. tom. 11. p. 864. Yarrell in *Zool. Journ.* vol. 11. p. 492. Red-crested Duck, Lath. *Syn.* vol. 111. p. 544. Red-crested Pochard, Selb. Illust. vol. 11. p. 350. (Trachea,) Linn. *Trans.* vol. xv. pl. 15. f. c.

DIMENS. Entire length twenty-one inches: length of the bill (from the forehead) two inches two lines, (from the gape) two inches three lines; of the tarsus one inch four lines and a half: from the carpus to the end of the wing ten inches.

DESCRIPT. (*Male.*) Head, cheeks, throat, and upper part of the neck, reddish-brown or bay; the feathers on the crown elongated, and of a silky texture, forming a crest, somewhat paler than the rest of the head; lower part of the neck, breast, belly, and abdomen, deep black: back, wings, and tail, pale brown: flanks, bend of the wing, a large spot on the sides of the back, speculum, and basal part of the primary quills, white: bill, tarsi, and toes, bright red: nail of the bill white: membranes of the feet black: irides bright red. (*Female.*) "Crown, occiput and nape, deep brown; the crest less tufted; cheeks, throat, and sides of the neck, ash-brown; breast and flanks yellowish brown; belly and abdomen gray: back, wings, and tail, brown, with a slight tinge of ochre; no white spot on the sides of the back; speculum half grayish white, the other half pale brown; base of the quills white tinged with brown: bill, tarsi, and toes, reddish brown." **TEMM.** (*Immature male.*) Nape, fore part of the neck, and breast, dark brown; abdomen of a lighter brown: flanks white, tinged with pink: tail ash-brown; upper and under tail-coverts dark brown: legs and toes orange: the rest as in the adult. (*Egg.*) Uniform olive-brown: long. diam. two inches two lines; trans. diam. one inch six lines.

An immature male of this species, supposed to be in the second year, was shot near Boston, in January 1826, while feeding on fresh water in company with some Wigeons. It is now in the collection of Mr Yarrell. Since then a second individual has occurred in the London markets; a third has been shot at Yarmouth; and a fourth killed near Colchester. This last is in the Museum of the Cambridge Philosophical Society. Said to inhabit the north-eastern parts of Europe. Food, according to Temminck, shell-fish and aquatic vegetables.

252. *F. ferina*, Steph. (*Common Pochard.*)—Head and neck bright chestnut; breast black; flanks and scapulars with undulating lines of black and grayish white: no speculum: bill black, with a broad transverse blue band.

Anas ferina, Temm. *Man. d'Orn.* tom. 11. p. 868. Pochard, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 353. Red-headed Pochard, *Selb. Illust.* vol. 11. p. 347. pl. 63. f. 1. (*Trachea*), *Linn. Trans.* vol. 14. pl. 14. f. 5.

DIMENS. Entire length nineteen inches: length of the bill (from the forehead) two inches, (from the gape) two inches three lines; of the tarsus one inch nine lines; of the tail two inches six lines; from the carpus to the end of the wing eight inches three lines: breadth, wings extended, twenty-nine inches seven lines.

DESCRIPT. (*Male.*) Head and neck bright chestnut-red: upper part of the back, breast, and rump, black; middle and lower regions of the back, scapulars, wing-coverts, flanks, and thighs, grayish white, with numerous fine undulating black lines; belly and abdomen whitish, very faintly undulated in the same manner, the lines becoming darker towards

the vent: secondary quills bluish gray: primaries, and tail, dark cinereous brown, approaching to dusky: upper and under tail-coverts black: bill black, with a broad transverse band in the middle of deep bluish gray: irides orange-yellow: legs bluish gray; membranes black. (*Female*.) Crown, nape, sides of the neck, and upper part of the back, reddish brown; throat, and fore part of the neck, white, mixed with reddish; breast reddish brown, the feathers edged and mottled with reddish white; flanks with large brown spots; middle of the belly grayish white: back and wings much as in the male, but the undulating lines less distinct. *Young males of the year* resemble the female. In those of *one and two years*, the breast is dusky brown, and the head and neck not so bright as in the adult. (*Egg*.) Greenish white: long. diam. two inches three lines and a half; trans. diam. one inch seven lines and a half.

Not uncommon on the British coasts during the winter season. Has been known to breed in Norfolk, but the greater part retire northwards in the Spring for that purpose. Eggs twelve to fourteen in number. Food, small fish, marine plants, and insects.

253. F. Nyroca, Steph. (*Nyroca Pochard*.)—Head, neck, breast, and flanks, ferruginous: irides, and a spot on the chin, white: speculum white, edged with black beneath: bill and legs bluish gray.

F. Nyroca, *Steph. in Shaw's Gen. Zool.* vol. xii. part ii. p. 201. pl. 55. *Anas leucophthalmos*, *Temm. Man. d'Orn.* tom. ii. p. 876. Olive-tufted Duck, *Sow. Brit. Misc.* pl. 21. Ferruginous Duck, *Penn. Brit. Zool.* vol. ii. p. 601. pl. 99? *Mont. Orn. Dict. Supp. with fig.* Castaneous Duck, *Mont. Orn. Dict. Supp. App. Bew. Brit. Birds*, vol. ii. p. 315. *Nyroca Pochard*, *Selb. Illust.* vol. ii. p. 352. pl. 63. f. 2. (*Trachca*.) *Mont. Orn. Dict. Supp. App.* last pl. f. 1.

DIMENS. Entire length (*male*) sixteen inches six lines, (*female*) eighteen inches. **MONT.**

DESCRIPT. (*Male*.) Head and neck dark ferruginous; on the lower part of the neck a narrow collar of blackish brown; back, scapulars, and wing-coverts, dusky brown, somewhat glossed with green and purple reflections, the whole finely powdered with pale reddish brown: a whitish spot on the chin; lower part of the neck, breast, and flanks, bright ferruginous; belly and under tail-coverts pure white; abdomen and vent blackish gray, finely speckled with yellowish white: speculum white, edged with black at the lower part: primaries dusky: tail, and upper tail-coverts, dusky brown, with a slight dash of ferruginous: bill and legs bluish gray; the nail of the bill, and the membranes of the toes, black: irides white. (*Female*.) Head, neck, breast, and flanks, brown, all the feathers tipped with reddish; the collar on the lower part of the neck indistinct: upper parts dusky, the feathers tipped with pale brown: the rest as in the male. (*Young of the year*.) "Crown of the head dusky brown: all the feathers on the upper parts edged and tipped with reddish brown: the white on the belly clouded with pale brown." **TEMM.** (*Egg*.) "White, tinged with greenish." **TEMM.**

A rare species in this country, but met with occasionally during the winter months. Frequents rivers, extensive lakes, and other inland waters. Said to breed in marshes and on the rushy banks of rivers, laying from eight to ten eggs. Food, insects, aquatic vegetables, &c.

Obs. The *Collared Duck* of Donovan*, is either a female or a young male of this species, with a slight variation of plumage. The *Ferruginous Duck* of Pennant is also probably referable to this species in some one of its different states.

254. *F. dispar*†, Steph. (*Western Pochard*).—White; beneath ferruginous: forehead, and occipital band, green; orbits, throat, collar and back, black.

F. dispar, Steph. in *Shaw's Gen. Zool.* vol. XII. part ii. p. 206. *Anas dispar*, Gmel. *Syst.* tom. i. part ii. p. 535. *Western Duck*, Penn. *Arct. Zool.* vol. II. p. 289. pl. 23. *Lath. Syn.* vol. III. p. 532. *Id. Supp.* p. 275. *Shaw, Nat. Misc.* vol. i. pl. 32. *Western Pochard*, Shaw, *Gen. Zool.* l. c. *Selb. Illust.* vol. II. p. 360. pl. 66**.

DIMENS. Entire length seventeen inches. LATH.

DESCRIPT. (*Male*.) Crown, sides of the head and neck, and hind part of this last for half-way, white; across the forehead a band of pea-green; a transverse fascia on the nape of the same colour, but deeper, at the lower corner of which, on each side, is a round black spot of the size of a pea; behind the eye, another spot of the same colour, but irregular in shape: chin, throat, and fore part of the neck, black, communicating with a collar of the same which surrounds the neck about the middle; from the hind part of this the black passes down over the back, quite to the tail: breast and sides pale ferruginous, deepening into chestnut at the middle, growing still deeper as it passes on towards the vent, where the colour is black: wing-coverts white: primary quills dusky black: secondaries six inches long, and curving downwards, partly white, partly black, the colours divided obliquely on each feather; scapulars also long, and curving elegantly downwards over the greater coverts, as in the *Garganey*; each of these has the web next the body scarcely broader than the shaft itself, and both of them white; the other web very broad, and black: tail pointed, brown: bill and legs black. LATH. (*Female*.) The whole plumage mixed brown and ferruginous, not unlike that of the *Woodcock*: the quills all straight, and of a dusky colour; some of the secondaries with white tips, forming a spot on the wing; some of the wing-coverts also tipped with white: legs black. LATH. (*Egg*.) Unknown.

Only one individual of this species has hitherto occurred in Britain. This, which was a male specimen, is recorded to have been shot at Caistor near Yarmouth, in February 1830. It is now in the Norwich Museum. Said to inhabit Kamtschatka, and the western coast of America, breeding amongst rocks. Pennant states that it flies in flocks, and never enters the mouths of rivers.

255. *F. Marila*, Steph. (*Scaup Pochard*).—Head and neck black, glossed with green; back and scapulars whitish, with undulating black lines; belly, flanks, and speculum on the wing, white; bill blue; legs ash-colour.

* *Brit. Birds*, vol. VI. pl. 147.

† I have followed Stephens and Selby in referring this species to the present genus, until its true situation be determined. It may be questioned, however, whether it be not more properly an *Oidemia*.

Anas Marila, *Temm. Man. d'Orn.* tom. 11. p. 865. Scaup Duck, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 339. Scaup Pochard, *Selb. Illust.* vol. 11. p. 354. pls. 66, & 66*. (Trachea,) *Linn. Trans.* vol. 1v. pl. 14. f. 3, & 4.

DIMENS. Entire length twenty inches: length of the bill (from the forehead) one inch ten lines, (from the gape) two inches three lines; of the tarsus one inch five lines and a half; of the tail two inches seven lines; from the carpus to the end of the wing eight inches six lines.

DESCRIPT. (*Male.*) Head and upper part of the neck greenish black, with glossy reflections; lower part of the neck, and breast, pitch-black: upper part of the back, and scapulars, grayish white, with fine, distant, transverse, undulating, black lines: first three primaries, and greater wing-coverts, dusky; lesser coverts the same, but finely powdered with white; the rest of the primaries, and all the secondaries, white, with dusky tips, forming a bar across the wings when closed: rump, tail, and upper tail-coverts, deep brown, approaching to black: belly and flanks pure white, the latter slightly undulated with black: vent, and under tail-coverts, spotted with brown: bill pale grayish blue; the nail black: irides bright yellow: legs ash-gray; the membranes dusky. (*Female.*) A broad, transverse band of yellowish white surrounding the base of the bill; rest of the head, and neck, chocolate-brown; lower part of the neck, breast, and rump, deep brown; back and scapulars dusky brown, with minute specks and close undulating lines of grayish white; flanks spotted and undulated with brown: irides dull yellow. (*Young male.*) More or less similar to the adult female: some white feathers at the base of the bill; the black on the head and neck without gloss, and mixed with feathers of a dusky brown colour: the white on the back variegated with brown spots; and the undulating lines closer together than in the *adult male*: belly dirty white, spotted with gray; flanks spotted with dusky brown. (*Egg.*) Greenish yellow white: long, diam. two inches five lines; trans. diam. one inch eight lines and a half.

A regular winter visitant, and not uncommon at that season on most parts of the British coast. Occasionally met with inland on fresh waters. Retires in the Spring to high northern latitudes to breed. Food, marine plants and insects, fish, and molluscous animals. *Obs.* The *Anas fraenata* of Sparmann* is the female of this species.

256. *F. cristata*, Steph. (*Tufted Pochard.*)—Head, neck, crest, and upper parts, black, with glossy reflections; belly, flanks, and speculum on the wing, white: bill and legs bluish gray.

Anas Fuligula, *Temm. Man. d'Orn.* tom. 11. p. 873. Tufted Duck, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. 11. p. 370. Tufted Pochard, *Selb. Illust.* vol. 11. p. 357. pl. 65.

DIMENS. Entire length seventeen inches: length of the bill (from the forehead) one inch seven lines, (from the gape) one inch nine lines and a half; of the tarsus one inch four lines and a half; of the tail two inches four lines; from the carpus to the end of the wing eight inches two lines: breadth, wings extended, twenty-eight inches two lines.

DESCRIPT. (*Male.*) Head, neck, breast, and a pendent crest of elongated feathers on the occiput, black, with green and purple reflections: back, wings, and rump, brownish black, slightly glossed with green, and

* *Sow. Brit. Misc.* pl. 62.

finely powdered with grayish white: belly, flanks, and a transverse bar on the wings, pure white; vent black: bill pale bluish gray; the nail black: irides gamboge-yellow: legs bluish gray; the membranes dusky. (*Female*.) Smaller than the male; the crest shorter, and of not so bright a colour: head, neck, breast, and upper part of the back, dull black, clouded with deep brown; back and wings dull blackish brown, speckled with grayish brown; breast and flanks with large spots of reddish brown; belly whitish, tinged with reddish brown: speculum smaller than in the male: bill and legs of a deeper colour: irides pale yellow. (*Young of the year of both sexes*.) "No indication of a crest: a large whitish spot on the sides of the bill; some white also on the forehead and occasionally behind the eyes: head, neck, and breast, dull brown, this last variegated with reddish brown: back and wings dusky brown, the feathers edged with paler brown: flanks reddish brown: the bar on the wings small and whitish: abdomen variegated with brown and ash-colour: irides dirty yellow. *Young males* have the belly of a purer white than *young females*. After the first moult, the white at the base of the bill disappears; the crest begins to shew itself; and the plumage becomes deeper coloured." TEMM. (*Egg*.) Greenish white, spotted with light brown: long. diam. two inches one line; trans. diam. one inch five lines.

Like the last species, a winter visitant. Met with on the coast, and also not unfrequently on fresh waters. Said to breed within the Arctic Circle. Food, similar to that of the preceding.

GEN. 97. CLANGULA, *Flem.*

257. *C. chrysophthalmos*, Steph. (*Golden-Eye Garrot*.)

—A large patch before the eye, lower neck, all the under parts, and speculum on the wing, white: legs orange-yellow.

Anas Clangula, Temm. *Man. d'Orn.* tom. 11. p. 870. *Golden-Eye*, Mont. *Orn. Dict. & Supp.* Bew. *Brit. Birds*, vol. 11. p. 365. *Common Golden-Eye Garrot*, Selb. *Illust.* vol. 11. p. 367. pl. 62. (*Trachea*.) Linn. *Trans.* vol. 1v. pl. 15. f. 1, & 2.

DIMENS. Entire length eighteen inches six lines: length of the bill (from the forehead) one inch three lines, (from the gape) one inch seven lines; of the tarsus one inch six lines: of the tail three inches seven lines; from the carpus to the end of the wing eight inches three lines.

DESCRIPT. (*Male*.) A large white space at the corner of the bill, immediately beneath the lore; rest of the head, and upper half of the neck, glossy black, with green and violet reflections; lower part of the neck, breast, belly, abdomen, and flanks, pure white: back, rump, and upper tail-coverts, deep glossy black: scapulars partly white and partly black; primary quills, four outer secondaries, and coverts bordering the wing, brownish black; rest of the secondaries, middle and greater coverts, white: tail blackish gray: bill black: irides golden yellow: legs orange; the membranes, and joints of the toes, dusky. (*Female*.) No white space at the corner of the bill; the whole head, and upper part of the neck, deep umber-brown: lower part of the neck, belly, and abdomen, pure white; breast and flanks deep ash-colour, margined with grayish white: all the upper parts grayish black, the feathers edged and tipped with ash: greater coverts white, tipped with black; lesser coverts blackish gray, tipped with white: bill black: the tip yellowish: irides and legs pale

yellow. *Young males of the year* resemble the adult female. *At the end of the first year*, the white patch at the base of the bill begins to shew itself, and the head becomes black, but without any glossy reflections. (*Egg.*) Buff-coloured white: long. diam. two inches four lines; trans. diam. one inch seven lines.

Met with in small flocks during the winter season, but not very plentiful. Frequents principally rivers, lakes, and other inland waters. Retires northward in the Spring to breed. Said to lay as many as fourteen eggs. Food, aquatic insects and vegetables, small fish, and mollusca. *Obs.* The *Anas Glaucion* of Linnæus may be referred either to the female, or to the young, of this species.

258. *C. histrionica*, Steph. (*Harlequin Garrot.*)—A large patch on the lore, spot on the ears, longitudinal band on the sides of the neck, collar, and pectoral fascia on each side, white: speculum blue: legs dusky.

C. histrionica, Steph. in *Shaw's Gen. Zool.* vol. xii. part ii. p. 180. pl. 57. *Anas histrionica*, Temm. *Man. d'Orn.* tom. ii. p. 878. *Sow. Brit. Misc.* pl. 6. *Harlequin Duck*, Mont. *Orn. Dict.* *Harlequin Garrot*, Selb. *Illust.* vol. ii. p. 371. pl. 60.

DIMENS. Entire length seventeen inches: length of the bill one inch six lines. MONT.

DESCRIPT. (*Male.*) Head and neck black, glossed with violet: a large patch between the bill and the eye, a spot behind the eyes, a longitudinal band on the sides of the neck, collar lower down, a broad crescent-shaped fascia on the sides of the breast, and part of the scapulars, all pure white: lower part of the neck, and breast, bluish ash; flanks ferruginous; belly brown: back, wings, and rump, glossy black, with blue and violet reflections: speculum on the wing very deep violet: bill black: irides brown: legs, and membranes of the toes, dusky blue. TEMM. (*Female.*) Smaller: all the upper plumage deep brown, clouded with ash-colour: towards the forehead, and a little before the eyes, a small white spot; a large patch of the same colour on the orifice of the ears: throat whitish; breast and belly whitish, clouded and spotted with brown; flanks reddish brown. TEMM. (*Egg.*) White, tinged with buff-colour: long. diam. two inches two lines; trans. diam. one inch seven lines.

A very rare occasional winter visitant. A pair, male and female, killed some years back in the north of Scotland, were in the collection of the late Mr James Sowerby. A young female has also been shot in one of the Orkney Islands. According to Messrs. C. and J. Paget*, it has also been obtained in one instance in the market at Yarmouth. Common in the northern parts of Europe and America. Said to breed on the borders of waters, amongst shrubs and herbage, and to lay from ten to twelve eggs. Food, shell-fish and insects. *Obs.* The *Anas minuta* of Linnæus is the female of this species.

(22.) *C. albeola*, Steph. (*Buffel-headed Garrot.*) Faun. Bor. Amer. part ii. p. 458. *Anas albeola*, Wils. Amer. Orn. vol. viii. p. 51. pl. 67. f. 2 & 3. Don. Brit. Birds, vol. x. pl. 226.

Inserted as British by Donovan in his "British Birds," but without any authority being given. A common species in North America.

GEN. 98. HARELDA, *Leach*.

259. *H. glacialis*, Steph. (*Long-tailed Hareld.*)—Crown and nape white; a large chestnut patch on the sides of the neck: bill black, with a transverse orange band: legs yellow.

Anas glacialis, *Temm. Man. d'Orn.* tom. II. p. 860. *Sabine in Linn. Trans.* vol. XII. p. 555. Long-tailed Duck, *Mont. Orn. Dict. & Suppl. Bew. Brit. Birds*, vol. II. p. 359. Long-tailed Hareld, *Selb. Illust.* vol. II. p. 363. pl. 61. (*Trachca*,) *Linn. Trans.* vol. XII. pl. 30. f. 3, & 4.

DIMENS. Entire length, central tail-feathers included, twenty-two inches. **MONT.**

DESCRIPT. (*Adult male in winter.*) Crown of the head, nape, front and lower part of the neck, pure white; cheeks and chin ash-gray; on each side of the upper part of the neck a large patch of chestnut-brown: breast, back, wings, rump, and the two elongated central tail-feathers, brownish black: scapulars, and lateral tail-feathers, white; the former long, narrow, and acuminate: flanks cinereous; belly and abdomen white: bill black, with a broad transverse band of deep orange before the nostrils: irides reddish yellow: legs yellowish; the membranes of the toes dusky. (*Adult female.*) "Forehead, chin, and eyebrows, whitish ash; crown of the head, and the large patch on the sides of the neck, dusky ash: nape, front and lower part of the neck, as well as the belly and abdomen, pure white: breast variegated with brown and dark ash-colour: back, scapulars, and wing-coverts, blackish brown, the feathers edged and tipped with reddish ash: the rest of the upper parts sooty brown: tail short, the central feathers not elongated; blackish brown, edged with white: bill bluish; the transverse band yellowish: irides light brown: legs bluish gray." **TEMM.** (*Male in summer.*) "The whole under part of the neck, and breast, black; sides of the head, and a little beyond the eye, brownish white; round the eye some white feathers; from the bill a black line runs on the top of the head to the crown, which is black; back of the neck chiefly black, but at a small distance below the crown with a few white feathers intermixed; the black of the back of the neck extends down the back, but in the centre of the upper part of the back near the neck is a patch of black feathers edged with ferruginous; scapulars long and narrow, black in the centre, and edged with ferruginous white, the longer ones having more white; wings brownish black, the quills being palest; lower belly and sides to the rump, and under tail-coverts, white, a line of black descending between the white from the back to the tail." **SAB.** (*Young of the year.*) Not very different from the adult female: face, crown, occiput, and nape, grayish white, mottled with brown and ash-colour; on the sides of the neck a large dusky patch; throat and upper part of the neck in front, belly and abdomen, white; lower part of the neck in front, and upper part of the breast, spotted and variegated with brownish ash; sides of the breast with transverse crescent-shaped bars of the same colour; back, and rest of the upper parts, deep brown; the scapulars edged with ash-colour, and some of the shorter ones with white feathers intermixed; tail short, the two middle feathers not elongated. (*Egg.*) Greenish yellow white: long. diam. two inches two lines; trans. diam. one inch six lines and a half.

A regular winter visitant in the Orkneys, and on the northern coasts of England and Scotland. Seldom met with in the southern parts of

Britain; such individuals generally young birds. Breeds in the Arctic Regions. Nest said to be placed amongst grass, near the edge of the sea. Eggs, according to Temminck, five in number; according to others, from ten to fourteen. Food, marine insects, and bivalve mollusca. *Obs.* The *Anas hyemalis* of Linnæus is the male of this species in summer plumage.

GEN. 99. MERGUS, Linn.

260. *M. Merganser*, Linn. (*Goosander*.) — Breast, and speculum on the wing, white; the latter without transverse bars: bill and legs red: crest (in the *adult male*) short and bushy.

M. Merganser, *Temm. Man. d'Orn.* tom. 11. p. 881. *Goosander*, *Mont. Orn. Dict. & Suppl. Selb. Illust.* vol. 11. p. 375. pl. 57. *Goosander and Dun-Diver*, *Bew. Brit. Birds*, vol. 11. pp. 250, & 253. (Trachea,) *Linn. Trans.* vol. xv. pl. 15. f. h.

DIMENS. Entire length twenty-four inches: length of the bill (from the forehead) two inches four lines, (from the gape) three inches one line; of the tarsus two inches one line; from the carpus to the end of the wing ten inches six lines. (*Young male*.) The *adult* sometimes attains a length of twenty-nine inches.

DESCRIPT. (*Adult male*.) Head and upper part of the neck glossy greenish black; the feathers on the crown and occiput elongated, forming a short crest: lower part of the neck, breast, belly, and abdomen, cream-yellow, fading after death to pure white: upper part of the back, inner scapulars, humeral wing-coverts, bastard winglet, basal halves of the greater coverts, and fourteen outer quills, black; outer scapulars, all the lesser coverts except the humeral ones, six of the secondary quills, and tips of the greater coverts immediately above them, white: lower back, rump, and tail, deep ash-gray: bill vermilion red; the ridge and nail black: irides red: legs vermilion. (*Adult female*.) Crown, and occipital crest (the latter longer and more slender than in the male), ferruginous brown; rest of the head, and upper part of the neck, bright ferruginous; chin and throat pure white; lower part of the neck before, sides of the breast, flanks, and thighs, white and ash-colour mixed; belly and abdomen yellowish white: back, scapulars, wing-coverts, and tail, deep ash-colour: primaries dusky; six of the secondaries, and tips of the impending coverts, white, forming a large speculum: bill dull red; the ridge and nail blackish: irides brown: legs yellowish red. *The young males of the year* resemble the adult female. *At the end of the first year*, the white on the throat begins to be spotted with black; the crown becomes dusky, and the ferruginous brown on the neck is edged at bottom with black; the white also appears on the wing-coverts. (*Egg*.) Pale olive white, tinged with buff: long. diam. two inches six lines; trans. diam. one inch eight lines.

Found in the Orkneys, and some other of the Scotch Islands, throughout the year. In England, only a winter visitant, and seldom seen at all in the more southern districts except in severe seasons. Frequents lakes, large rivers, and estuaries. Nest said to be placed amongst loose stones near the edge of the water, occasionally in bushes, or in hollow trees.

Eggs twelve to fourteen in number. Food, fish and amphibious reptiles. *Obs.* The *Dun-Diver* of English authors (*M. Castor*, Linn.) is the female of this species.

261. *M. Serrator*, Linn. (*Red-breasted Merganser.*)

—Breast (in the *male*) reddish brown: speculum white, divided by one or two transverse bars: bill and legs red: crest (in the *adult male*) long, slender, pendent.

M. Serrator, *Temm. Man. d'Orn.* tom. 11. p. 884. Red-breasted Merganser, *Mont. Orn. Dict. Bew. Brit. Birds*, vol. 11. p. 257. *Selb. Illust.* vol. 11. p. 379. pls. 58, & 58*. (Trachea.) *Linn. Trans.* vol. 1V. pl. 16. f. 1, & 2.

DIMENS. Entire length twenty-two inches.

DESCRIPT. (*Adult male.*) Head, a pendent crest on the occiput, and upper part of the neck, greenish black, with glossy reflections; lower part of the neck white, the mesial line behind black; breast reddish brown, spotted and variegated with black; near the insertion of the wing, several large white spots bordered with black; upper part of the back, scapulars, edge of the wing and adjoining coverts, and primary quills, black; the rest of the coverts, and secondary quills, white, but these last, as well as the greater coverts, black at the base, the whole together forming a large speculum divided by two transverse bars; some of the secondaries nearest the body white edged with black: lower part of the back, rump, and flanks, grayish white, with fine transverse undulating black lines; belly and abdomen white; tail brown: bill orange-red; the ridge and nail dusky: irides red: legs orange. (*Adult female.*) Head, crest, and neck, dull ferruginous; throat white; fore part of the neck and breast variegated with white and ash-gray; lower part of the neck behind, back, rump, wings, and scapulars, dark ash-colour; speculum white, with one transverse bar, by which it may always be distinguished from the female of the last species, which in other respects it greatly resembles; belly and abdomen white; flanks dark ash-colour: bill and legs dull orange: irides brown. (*Young male of the year.*) Head deep brown; throat grayish white; bill pale red; irides yellowish: in other respects like the adult female. (*Egg.*) Smooth and shining; buff-colour: long. diam. two inches six lines; trans. diam. one inch nine lines.

A winter visitant in England, and not often met with in the southern counties. More abundant northward. Breeds in some of the Scotch lakes. Nest, according to Selby, composed of small sticks and grass, lined with the down of the female; built in the thick herbage on the edge of the main land, or on small islets in the lake. Eggs nine in number. Food and habits similar to those of the last species.

262. *M. cucullatus*, Linn. (*Hooded Merganser.*)—

Head, neck, and upper parts, dusky brown; beneath white: bill and legs reddish brown: crest (in the *adult male*) semicircular, compressed.

M. cucullatus, *Faun. Bor. Amer.* part ii. p. 463. Hooded Merganser, *Lath. Syn.* vol. 111. p. 426. pl. 101. *Wils. Amer. Orn.* vol. VII. p. 79. pl. 69. f. 1. *Selb. in Trans. of Nat. Hist. Soc. of Newcast.* vol. I. p. 292. *Id. Illust.* vol. 11. p. 383. pl. 58*.

DIMENS. Entire length eighteen inches: length of the bill (from the forehead) one inch eight lines, (from the gape) two inches; of the tarsus one inch three lines; of the tail three inches ten lines; from the carpus to the end of the wing seven inches six lines.

DESCRIPT. (*Adult male.*) Head and neck brownish black; the former furnished with a large, erect, compressed, semicircular crest, extending from the forehead to the occiput; frontal portion of the crest, space above the eyes, and a narrow edging all round, black; the remaining part of the crest, and a space on the sides of the occiput commencing five lines behind the eyes, white: back, rump, tail, primaries, and lesser wing-coverts, dark cinereous brown, inclining in some places to dusky: greater coverts, and secondary quills, black at the base, the tips of the former and outer borders of the latter white; the whole together forming a speculum consisting of four bars, black and white alternately: scapulars long and pointed, greenish black, streaked longitudinally down the middle with white: sides of the breast with two or three crescent-shaped transverse black bars; flanks, and sides of the body near the bend of the wing, finely undulated with black and yellowish brown: rest of the under parts pure white: bill and legs reddish brown. (*Adult female.*) Forehead, sides of the head, and all the neck, dark ash-colour; crest shorter than in the male, reddish ash, without any white: all the upper plumage dark brown, approaching to dusky, with a tinge of cinereous: upper part of the breast, and flanks, lead-gray, the feathers edged with white: the rest much as in the male. (*Young female?*) "Chin grayish white, speckled with grayish brown; the whole of the face, cheeks, and neck, of an uniform grayish brown, or mouse-colour; crown of the head darker; occipital crest pale reddish brown, tinged with gray; breast gray, the margins of the feathers paler; upper back and wing-coverts grayish black, the feathers margined with obscure grayish brown; scapulars and lower back black; margins of four or five of the secondary quills white, forming a small spot in the middle of the wings; quills and tail grayish black: lower part of the breast, belly, and abdomen, pure white, with a silken lustre; sides and flanks deep grayish brown; bill reddish brown at the base, the tip black; legs reddish brown." SELB. (*Egg.*) White. LATH.

A single individual of this species, supposed to be a young female, of which the above is a description borrowed from Selby, was killed near Yarmouth in Norfolk, in the Winter of 1829. It is a native of North America, and had never before been seen in Europe. According to Latham, appears at Hudson's Bay as a summer visitant at the end of May, and builds close to the lakes. Nest composed of grass, lined with feathers from the breast. Eggs four to six in number. Frequents fresh waters.

263. *M. albellus*, Linn. (*Smew.*)—Head, neck, and all the under parts, white; cheeks and occiput greenish black: bill and legs bluish gray: crest (in the *adult male*) bushy, moderately elongated.

M. albellus, *Temm. Man. d'Orn.* tom. 11. p. 887. *Smew*, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 260. *Selb. Illust.* vol. 11. p. 385. pl. 59. (*Trachea.*) *Linn. Trans.* vol. 1v. pl. 16. f. 3, & 4.

DIMENS. Entire length seventeen inches: length of the bill (from the forehead) one inch three lines, (from the gape) one inch eight lines; of the tarsus one inch six lines.

DESCRIPT. (*Adult male.*) A large patch on each side of the base of the bill enclosing the eyes, and another longitudinal one on the occiput, black glossed with green; rest of the head, occipital crest, neck, some of the lesser wing-coverts, and all the under parts, pure white: two crescent-shaped fasciæ advancing forwards from the shoulders on each side, and partly encircling, one the lower part of the neck, the other the upper part of the breast, black; back, lesser coverts bordering the wing, and primary quills, black; scapulars white, edged on the outer webs with black; secondary quills and greater coverts black, tipped with white: tail and upper tail-coverts bluish gray: flanks and thighs with transverse undulating lines of black: bill and legs deep bluish gray; the membranes of the toes dusky. (*Female.*) Crown, cheeks, and occiput, reddish brown; crest shorter than in the male; throat, sides and front of the upper part of the neck, belly, and abdomen, white; lower part of the neck, breast, and flanks, clouded with ash-colour: back, tail, and upper tail-coverts, deep ash-gray: wings much as in the male, only the dark parts gray instead of black. *The young of the year* resemble the adult female. *After the second moult*, the male begins to shew a few black feathers on the sides of the face, the first indication of the large patch which characterizes that part in the adult; some white appears on the crown and occiput; the back becomes partially black; and the two crescent-shaped fasciæ are faintly traced out on the sides of the breast. (*Egg.*) "Whitish."

TEMM.

A winter visitant, but not of very frequent occurrence. Principally met with in the neighbourhood of fresh waters. Breeds in high latitudes on the borders of lakes and rivers. Eggs, according to Temminck, from eight to twelve in number. Food, fish, and aquatic vegetables. *Obs.* The *Minute Merganser* of Montagu and other English authors (*M. minutus*, Linn.) is referable to the female and young of this species.

GEN. 100. PODICEPS, Lath.

264. *P. cristatus*, Lath. (*Great Crested Grebe.*) — Bill longer than the head, reddish, the tip white; distance from the nostrils to the tip about eighteen lines.

P. cristatus, *Temm. Man. d'Orn.* tom. II. p. 717. Crested Grebe, *Mont. Orn. Dict. Selb. Illust.* vol. II. p. 394. pl. 73. Great Crested Grebe, *Bew. Brit. Birds*, vol. II. p. 153.

DIMENS. Entire length twenty-one inches: length of the bill, from the forehead, two inches.

DESCRIPT. (*Adult in perfect plumage.*) Streak above the eye, cheeks, and throat, white: crown of the head, occipital crest, and a large ruff of feathers standing out round the neck, glossy black, passing into reddish yellow on the sides of the head: fore part of the neck, and all the under parts of the body, silvery white: hind part of the neck, back, scapulars, and wing-coverts, dusky brown: secondaries pure white, forming an oblique bar across the wings: naked space between the bill and the eye red: bill reddish, dusky brown on the ridge, the tip grayish white: irides crimson red: legs dusky; internally yellowish white. (*From two to three years of age.*) "Occipital crest very short, edged with white: face white, without the reddish tinge; the feathers of the ruff very short: a dusky streak passing from the bill beneath the eyes, and reaching to

the occiput. (*Young before the age of two years.*) No occipital crest or ruff round the neck: forehead and face white; these parts as well as the upper part of the neck with zigzag streaks of dusky brown disposed in every direction: irides pale yellow. *The young of the year before the first moult* have the head and upper part of the neck deep brown." TEMM. (*Egg.*) White: long. diam. two inches two lines; trans. diam. one inch six lines.

Not an uncommon species, remaining with us the whole year. Frequents lakes and extensive tracts of fen; during severe weather in Winter, the sea-coast and the estuaries of large rivers. Is said to breed in the meres of Shropshire and Cheshire, and also in Lincolnshire. Makes a large floating nest of aquatic plants. Eggs four in number. Food, principally fish. Dives well, and is rarely seen on land. *Obs.* The *Tippet Grebe* of Pennant (*C. Urinator*, Linn.) is this species in its immature state.

265. *P. rubricollis*, Lath. (*Red-necked Grebe.*)—Bill of the length of the head, black, towards the base yellow; distance from the nostrils to the tip thirteen or fourteen lines: no ruff; occipital crest very short.

P. rubricollis, Temm. *Man. d Orn.* tom. II. p. 720. *Red-necked Grebe*, Mont. *Orn. Dict. & Suppl.* Bew. *Brit. Birds*, vol. II. p. 161. *Selb. Illust.* vol. II. p. 392. pl. 72.

DIMENS. Entire length seventeen inches six lines: length of the bill (from the forehead) one inch nine lines, (from the gape) two inches two lines; of the tarsus two inches two lines; from the carpus to the end of the wing seven inches two lines: breadth, wings extended, twenty-nine inches three lines.

DESCRIPT. (*Adult.*) Forehead, crown, and occipital tuft, blackish brown: cheeks and throat ash-gray: back part of the neck, and all the upper parts of the body, deep blackish gray: front of the neck, sides and upper part of the breast, bright orange-brown: rest of the under parts white; flanks and thighs with a few dusky streaks: lower half of the secondary quills white: bill black, with the base of both mandibles bright gamboge-yellow: irides reddish brown: legs externally black; internally yellowish green. (*Young, at the age of two years.*) Throat and cheeks white: forehead, crown, and rest of the upper parts, dusky brown, without any elongated feathers on the occiput: sides of the neck, and upper part of the breast, pale reddish, mixed with dusky and brown; rest of the under parts white, faintly spotted with dusky on the flanks and towards the vent: base of the bill livid yellow. (*Egg.*) Dirty white: long. diam. one inch ten lines; trans. diam. one inch four lines.

Of rare occurrence in this country, particularly in the adult state. Abundant in the eastern parts of Europe. Frequents rivers and lakes, and also the sea-coast. Food, small fish, aquatic insects and vegetables. Nest said to resemble that of the last species. Eggs three or four in number.

266. *P. cornutus*, Lath. (*Sclavonian Grebe.*)—Bill strong, shorter than the head, compressed throughout its length; black, the tip red: eyes with a double iris: distance from the nostrils to the tip of the bill seven lines.

P. cornutus, *Temm. Man. d'Orn.* tom. II. p. 721. Slavonian Grebe, *Mont. Orn. Dict. & Supp. with fig.* Horned Grebe, *Selb. Illust.* vol. II. p. 397. pl. 74.

DIMENS. Entire length thirteen inches six lines: length of the bill (from the forehead) ten lines and a half, (from the gape) one inch four lines; of the tarsus one inch eight lines: breadth, wings extended, twenty-two inches.

DESCRIPT. (*Adult.*) Forehead, crown, and an ample ruff surrounding the upper part of the neck, deep glossy black; above and behind the eyes two large spreading tufts of feathers of an orange-red colour, resembling horns: nape, and all the upper parts of the body, dusky brown: secondaries white: front of the neck, and breast, bright ferruginous chestnut; rest of the under parts pure white, with the exception of the flanks, which are tinged with ferruginous: base of the bill, and naked space between it and the eyes, rose-red; rest of the bill black, the extreme tip red: irides with a double circle; the inner one yellow; the outer and broader one bright red: legs externally black, internally gray. (*Young, during the first year.*) Head, nape, and all the upper parts, dusky ash, deepest on the head, where it approaches to black, with a slight gloss of green; no indications of a ruff or tufts above the eyes: throat and cheeks pure white; a narrow line of the same colour extends from behind the ears on each side to the back of the head: middle of the fore part of the neck ash-colour; sides of the breast, and flanks, tinged with dusky; rest of the under parts, as well as the secondary quills, pure white: bill dusky, with the sides of the upper, and base of the under, mandible, of a livid flesh-colour, the tip pale brown: the inner circle of the iris white; the outer one red: legs externally dusky; internally bluish ash. (*Egg.*) Dirty greenish white: long. diam. one inch nine lines; trans. diam. one inch three lines.

More common than the last species, but not very plentiful, at least in the adult state. Found on fresh waters, as well as on the sea-coast. Food, similar to that of the two preceding species. Constructs a floating nest amongst reeds and rushes, and lays three or four eggs. *Obs.* The second of the two states of plumage described above is that of the *Dusky Grebe* of Pennant, (*P. obscurus*, Lath.) which Temminck considers as the young of this species. Not being prepared to prove to the contrary, I have followed him in this last instance. Yet the correctness of such an opinion may reasonably be called in question on the ground of anatomical differences observed by Mr Yarrell. This last gentleman remarked that in an adult male specimen of the *P. cornutus*, the stomach was membranous-muscular, and the cæcal appendages only an inch and a half in length; whereas in a specimen of the *P. obscurus* dissected by him, the stomach was more decidedly muscular, and the cæcal appendages nearly five inches in length. These facts appear to indicate a difference of species.

267. *P. auritus*, Lath. (*Eared Grebe.*)—Bill shorter than the head, depressed at the base, slightly recurved at the tip, black: iris single: distance from the nostrils to the tip of the bill six lines and a half.

P. auritus, *Temm. Man. d'Orn.* tom. II. p. 725. Eared Grebe, *Mont. Orn. Dict. & Supp. with fig.* *Bew. Brit. Birds*, vol. II. p. 157. *Selb. Illust.* vol. II. p. 399. pl. 74.

DIMENS. Entire length twelve inches: length of the bill (from the forehead) eleven lines, (from the gape) one inch one line and a half; of the tarsus one inch six lines and a half; from the carpus to the end of the wing five inches.

DESCRIPT. (*Adult.*) Cheeks, forehead, crown, a short occipital crest, and upper part of the neck, deep black: sides of the head ornamented with two tufts of long slender feathers, of a pale yellow passing into deep orange; these tufts arise from behind the upper part of the eye, and passing backwards cover the orifice of the ears: throat, lower part of the neck, sides of the breast, and all the upper parts, dull black: secondary quills white: flanks and thighs deep chestnut-red, mixed with dusky; the rest of the under parts pure white: bill black, reddish at the base: irides vermilion: legs externally dusky ash, internally greenish ash. (*Young of the year.*) Closely resembling the young of the last species, but may always be distinguished by the peculiar shape of the bill, which is depressed at the base, and slightly recurved at the tip: the iris is also of one colour, and the white on the cheeks more extended, reaching down the sides of the neck. (*Egg.*) Dirty yellowish white: long. diam. one inch nine lines; trans. diam. one inch three lines.

A rare species in this country, particularly in the adult state. Met with principally in the winter months, but, according to Montagu, breeds in the fens of Lincolnshire. Chiefly frequents lakes and inland waters, occasionally the estuaries of rivers. Nest placed amongst reeds and other aquatic herbage. Eggs three or four in number.

268. *P. minor*, Lath. (*Little Grebe.*) — Bill very short, compressed; distance from the nostrils to the tip six lines: no ruff or occipital crest: tarsi furnished posteriorly with a double row of serratures.

P. minor, *Temm. Man. d'Orn.* tom. 11. p. 727. *Colymbus Hebridicus*, *Sow. Brit. Misc.* pl. 70. Little and Black-chin Grebes, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 163. Little Grebe, *Selb. Illust.* vol. 11. p. 401. pl. 75.

DIMENS. Entire length nine inches ten lines: length of the bill one inch: of the tarsus one inch three lines; from the carpus to the end of the wing four inches three lines.

DESCRIPT. (*Adult in summer.*) Throat, crown of the head, and nape, deep black; cheeks, sides and fore part of the neck, bright chestnut: upper parts of the body dark brown, tinged with olivaceous green; rump inclining to reddish brown: primary quills brownish ash; secondaries white at the base and on their inner webs: under parts dusky ash, darkest on the breast, vent, and flanks; thighs tinged with reddish: bill black; the extreme tip, and base of the lower mandible, together with the lore, yellowish white: irides reddish brown: legs externally dusky olive, internally livid flesh-colour. (*Young of the year.*) Crown of the head, nape, and all the upper parts of the body, a mixture of brown and ash-colour, with a slight tinge of rufous: throat pure white: sides of the neck reddish ash; fore part of the neck, upper part of the breast, and flanks, the same but paler; belly and abdomen white: bill shorter than in the adult; the lower mandible, and edges of the upper mandible, yellowish white; rest of the bill deep brown: irides brown. (*Egg.*) Smooth; white: long. diam. one inch seven lines; trans. diam. one inch one line.

The most abundant species in the genus. A frequent inhabitant of ponds, marshes, lakes, and the banks of rivers. Rarely observed on the sea-coast. Feeds on *Dyticidæ* and other aquatic insects. Is an excellent diver. Constructs a large floating nest, which it attaches to the stems of rushes. Eggs four to six in number.

GEN. 101. COLYMBUS, *Lath.*

269. *C. glacialis*, Linn. (*Northern Diver*.)—Head and neck violet-black, the latter with a double interrupted white collar: bill upwards of four inches in length; the upper mandible nearly straight.

C. glacialis, *Temm. Man. d'Orn.* tom. II. p. 910. *Northern Diver*, *Mont. Orn. Dict. Selb. Illust.* vol. II. p. 406. pl. 76. *Great Northern Diver*, *Bew. Brit. Birds*, vol. II. p. 174.

DIMENS. Entire length thirty-three inches: length of the bill (from the forehead) two inches eleven lines, (from the gape) four inches one line; of the tarsus three inches; of the middle toe, nail included, four inches two lines; of the tail three inches six lines; from the carpus to the end of the wing thirteen inches six lines.

DESCRIPT. (*Adult*.) Head and neck velvet black, with green and purple reflections; beneath the throat a transverse semilunar white band with black streaks; on the lower part of the neck an interrupted broad collar, streaked longitudinally with black and white: all the upper parts black, elegantly spotted with white; the spots on the back and rump small and round; those on the scapulars larger and of an oblong-square form, disposed in rows, two at the extremity of each feather: quills and tail without spots: breast and other under parts white; flanks and sides of the breast with a few black streaks: bill black: the ridge of the upper mandible very slightly arched above; lower mandible channelled beneath, appearing deepest in the middle; the gonys sloping upwards to the tip: tomia of both mandibles, but particularly of the lower one, inflected: irides brown: feet externally dusky brown; the inner portions, and membranes between the toes, whitish. (*Young of the year*.) Head, occiput, and all the back part of the neck, cinereous brown; cheeks speckled with white and ash-colour; throat, front of the neck, and all the other under parts, pure white; the feathers on the back, wings, rump, and flanks, deep brown in the middle, the edges and tips ash-colour: upper mandible cinereous gray; lower mandible whitish. *At the age of one year*, a transverse bar of dusky brown appears on the middle of the neck, forming a sort of collar; the back assumes a dusky tint, and the small white spots begin for the first time to shew themselves. *At the end of the second year*, the dusky bar on the neck extends further; the brown on the head and nape becomes mixed with greenish black; the spots on the back and wings increase in number, and the band on the throat, and collar on the lower part of the neck, are more distinctly indicated by longitudinal streaks of brown and white. *At the age of three years*, the plumage is perfect. (*Egg*.) Dark olive-brown, with a few spots ofumber-brown: long. diam. three inches six lines; trans. diam. two inches three lines.

A native of high northern latitudes. Abundant in the Orkneys, Hebrides, and other Scotch islands, but very rare in the southern parts

of Britain. Has been taken in Cornwall, Bedfordshire, and occasionally, in the immature state, on different parts of the English coast. Seldom flies, but dives well. Feeds on fish, particularly herrings. Nest said to be placed on the borders of fresh-water lakes, or in small islands. Eggs two in number. *Obs.* In its immature state, this species is the *C. Immer* of authors, but it is probable that in some cases this name has been also applied to the young of the next species.

270. *C. arcticus*, Linn. (*Black-throated Diver*).— Occiput and nape cinereous brown; front of the neck violet-black: bill upwards of three inches in length; the upper mandible slightly curved.

C. arcticus, *Temm. Man. d'Orn.* tom. 11. p. 913. Black-throated Diver, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. 11. p. 181. *Selb. Illust.* vol. 11. p. 411. pl. 77.

DIMENS. Entire length from twenty-four to twenty-six inches.
TEMM.

DESCRIPT. (*Adult plumage.*) Head and nape cinereous brown, passing into black on the forehead; throat and fore part of the neck black, with purple and green reflections; immediately beneath the throat a narrow transverse white band longitudinally streaked with black; on the sides of the neck a broader band extending downwards from the ears, streaked longitudinally with black and white; lower part of the neck, and sides of the breast, streaked in a similar manner: back, rump, and flanks, deep black, without spots; scapulars with twelve or thirteen transverse bars of pure white; wing-coverts black, speckled with white: breast and under parts of the body pure white: quills dusky: tail grayish black; under tail-coverts barred with black: bill dusky; the upper mandible more curved than in the last species; the lower mandible not thickened in the middle, and without the groove: irides brown: feet externally brown; internally whitish; membranes whitish. (*Young of the year.*) Only to be distinguished from those of the last species by their inferior size, and characters of the bill as above pointed out: plumage almost entirely similar: there is generally, however, a dusky band running longitudinally down the sides of the neck, which is not present in the young of *C. glacialis*. *At the age of one year*, a few black feathers begin to appear on the throat, as well as on the back, rump, and flanks; there is also a slight indication of the broad band of black and white streaks on the sides of the neck. *At the end of the second year*, the colour of the forehead begins to deepen; the violet-black extends itself over the throat and fore part of the neck, still however mixed with a few white feathers; the band of longitudinal streaks appears more distinct; and the black and white on the upper parts becomes tolerably well-defined. *At the age of three years*, the plumage is generally matured. (*Egg.*) Dark olive-brown, thinly spotted with umber-brown: long. diam. two inches nine lines; trans. diam. one inch ten lines.

Inhabits the Arctic Regions, but appears occasionally as a winter visitant on different parts of the British coast, as well as on lakes and rivers inland. Feeds on fish, frogs, insects, and aquatic vegetables. Nest placed amongst reeds and rushes in marshy situations, or on the banks of rivers. Eggs two in number. *Obs.* The *Lesser Imber* of Bewick is probably the young of this species.

271. *C. septentrionalis*, Linn. (*Red-throated Diver*.)

—Occiput and nape with black and white streaks; front of the neck deep orange-brown: bill scarcely, or not exceeding, three inches; slightly recurved; tomia very much inflexed.

C. septentrionalis, *Temm. Man. d'Orn.* tom. II. p. 916. Red-throated Diver, *Mont. Orn. Dict. & Suppl. Bew. Brit. Birds*, vol. II. p. 177. *Selb. Illust.* vol. II. p. 414. pls. 78, & 78*.

DIMENS. Entire length twenty-six inches three lines: length of the bill (from the forehead) one inch eleven lines, (from the gape) two inches eleven lines and a half; of the tarsus two inches ten lines; of the tail three inches three lines; from the carpus to the end of the wing eleven inches three lines: breadth, wings extended, three feet four inches seven lines.

DESCRPT. (*Adult.*) Sides of the head, throat, and sides of the neck, mouse-gray; crown spotted with black; occiput, nape and lower part of the neck, marked with longitudinal streaks of black and white; on the front of the neck a large patch of deep orange-brown; breast and belly silvery white; flanks and thighs streaked with dusky; back, and all the other upper parts dusky brown, without spots in very old individuals, but in birds of three and four years of age, with small indistinct whitish spots: bill black; the commissure quite straight, but the upper mandible being considerably depressed above the nostrils, the ascending gonys of the lower one causes the bill, when closed, to appear slightly recurved: tomia of both mandibles greatly inflexed: irides orange-brown: feet externally greenish gray; internally livid white; membranes the same. (*Young of the year.*) Forehead, crown and nape, dusky ash, finely streaked with grayish white; back, scapulars and rump, dusky brown, copiously sprinkled with small oval white spots arranged along the edges of the feathers; wing-coverts bordered with white towards their extremities: throat, sides and fore part of the neck, breast and belly, white; flanks streaked with dusky: quills and tail blackish gray: bill grayish white; the ridge of the upper mandible dusky: irides brown: legs externally dark greenish gray; internally, as well as the webs, greenish white. *At the end of the first year*, a few red feathers begin to show themselves on the fore part of the neck; and the white spots on the upper parts become smaller and less distinct. *After the second moult*, the patch of orange-brown is nearly complete, but still mixed with a little white: the spots on the upper parts gradually disappear as the bird advances to maturity. (*Egg.*) Chestnut-brown, rather thickly spotted with dark umber-brown: long. diam. two inches eight lines; trans. diam. one inch ten lines.

A winter visitant on the English coast of not unfrequent occurrence, particularly in the immature state. Remains all the year in the Orkney and Shetland Islands, where it breeds. Nest placed on the extreme borders of lakes, composed of rushes and dry grass. Eggs two in number. Food similar to that of the last species. Flies well, and is also an excellent diver. *Obs.* The *Speckled Diver* of Pennant and Montagu (*C. stellatus*, Gmel.) is the young of the year of this species. The *Second Speckled Diver* of Bewick is perhaps the same bird a little nearer advanced to maturity.

GEN. 102. URIA, *Briss.*

272. *U. Troile*, Lath. (*Foolish Guillemot*).—Above brownish black; secondaries tipped with white: bill as long as, or somewhat longer than, the head; very much compressed throughout: feet dusky.

U. Troile, *Temm. Man. d'Orn.* tom. II. p. 921. *Foolish Guillemot*, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. II. p. 166. *Selb. Illust.* vol. II. p. 420. pl. 79.

DIMENS. Entire length seventeen inches nine lines: length of the bill (from the forehead) one inch eleven lines, (from the gape) two inches eleven lines; of the tarsus one inch seven lines; of the tail two inches; from the carpus to the end of the wing six inches two lines: breadth, wings extended, two feet.

DESCRIPT. (*Adult in winter.*) Forehead, crown, space between the eye and the bill, a longitudinal streak running backwards from the eyes to the distance of an inch and a half, and all the upper parts, including the wings, brownish black with a slight tinge of cinereous: occiput grayish white, opening on each side into a white patch above the longitudinal streak behind the eyes: secondaries tipped with white: all the under plumage pure white; the black on the sides of the neck advancing in front towards the lower part, and forming a faint collar of grayish ash: bill black; very much compressed throughout its whole length: inside of the mouth dull yellow: irides brown: feet and toes yellowish brown; posterior part of the tarsus and membranes black. (*Summer plumage*.) The whole of the head, including the region of the eyes, throat, and all the upper part of the neck, pitchy brown: inside of the mouth bright yellow: the rest as in winter. (*Young of the year.*) Resemble the adult in winter, but distinguished by the shorter bill: the longitudinal streak behind the eyes is also indistinct; and the white on the under parts less pure, with more of cinereous brown on the lower part of the neck: tarsi and toes dull yellowish; membranes brown. (*Egg*.) Very variable in colour, scarcely two being found precisely alike; generally bluish green, more or less blotched and streaked with black: long. diam. three inches two lines; trans. diam. one inch eleven lines.

Met with in great abundance on various parts of the British coast throughout the year. Breeds on cliffs and rocky islands, generally in large companies. Lays a single egg. Food, fish, marine insects, and small bivalve mollusca. *Obs.* The *Lesser Guillemot* of English authors is the young of the year of this species.

(23.) *U. Brunnichii*, Sab. in Linn. Trans. vol. XII. p. 538. *Temm. Man. d'Orn.* tom. II. p. 924. *Franks' Guillemot*, Shaw, Gen. Zool. vol. XII. p. 243. pl. 62.

According to the statement of a writer in the *Quarterly Review*, (vol. XLVII. p. 354.) this species has been killed off one of the Shetland Isles. It is distinguished from the last, principally by the form of the bill, which is shorter and more dilated at the base.

273. *U. Grylle*, Lath. (*Black Guillemot*).—Above black; a large white patch on the middle of the wings: legs red.

U. Grylle, *Temm. Man. d'Orn.* tom. 11. p. 925. Black Guillemot, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 170. *Selb. Illust.* vol. 11. p. 426. pl. 80.

DIMENS. Entire length thirteen inches six lines: length of the bill (from the forehead) one inch two lines, (from the gape) one inch five lines; of the tarsus one inch two lines; of the tail one inch nine lines; from the carpus to the end of the wing six inches seven lines.

DESCRIPT. (*Adult in winter.*) Crown, nape, and all the rest of the upper plumage, sooty black, with the exception of the middle and greater coverts, which are white, forming a large patch of this last colour on the middle of the wing: sides of the face, and all the under parts, white: bill black: irides brown: inside of the mouth, and legs, reddish. (*Summer plumage.*) The whole plumage above and below, with the exception of the large white space in the middle of the wing, deep sooty black: bill black: inside of the mouth, and legs, bright red. (*Young of the year.*) Crown of the head and nape blackish, the feathers edged with white; lower part of the neck, and sides of the breast, gray and white mixed: throat, breast, and rest of the under parts, white: upper back and scapulars black, many of the feathers edged with white; lower back black: rump white mixed with black: wings as in the adult, but the white speculum marked with cinereous and dusky spots: irides dusky brown: inside of the mouth, and legs, reddish. (*Egg.*) White, tinged with green: blotched, spotted and speckled, with ash-gray, red brown, and very dark brown: long. diam. two inches three lines; trans. diam. one inch six lines.

Met with principally in the northern parts of the kingdom. Common in the Hebrides and Orkney Islands, where they remain all the year, breeding in the crevices of rocks. Like the last species, lays but one egg*. Food, small fish, and marine crustacea. *Obs.* The *Spotted Guillemot* of authors is this species in its immature state.

GEN. 103. MERGULUS, Ray.

274. M. *Alle*, Selby. (*Common Rotche.*) — Black; breast, belly, a spot above the eyes, and tips of the secondaries, white.

Uria Alle, *Temm. Man. d'Orn.* tom. 11. p. 928. Little Auk, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 408. Common Rotche, *Selb. Illust.* vol. 11. p. 430. pl. 81.

DIMENS. Entire length eight inches ten lines: length of the bill (from the forehead) seven lines, (from the gape) eleven lines; of the tarsus nine lines and a half; of the tail one inch seven lines; from the carpus to the end of the wing four inches nine lines.

DESCRIPT. (*Winter plumage.*) Crown, region of the eyes, nape, sides of the breast, and all the upper parts of the body, deep black: wings pitchy brown; secondaries tipped with white; tertials broadly edged with the same colour: throat, front and sides of the neck, and all the under parts, pure white: sides of the head also white, variegated with dusky streaks, the white passing off towards the occiput in the form of an indistinct narrow band: bill black: irides dusky brown: tarsi and toes

* A writer in *Loudon's Magazine of Natural History*, (vol. v. p. 413.) asserts that in a large number of instances he has invariably found the eggs *two* in number.

yellowish brown; membranes greenish brown. (*Summer plumage.*) Head, cheeks, throat, and neck, deep sooty black; a small white spot above the eyes: the rest as in winter. (*Young of the year.*) Distinguished by their shorter and smaller bill: plumage similar to that of the adult in winter. (*Egg.*) Of a uniform pale blue: long. diam. one inch seven lines; trans. diam. one inch one line.

Very abundant in the Arctic Regions, where it breeds. Met with occasionally on the British coasts during the winter season. Lays two eggs, which are deposited in the holes and crevices of the steepest rocks. Food, marine insects and crustacea.

GEN. 104. FRATERCULA, *Briss.*

275. *F. arctica*, Steph. (*Puffin.*)—Crown, collar and back, black; cheeks, throat, breast and belly, white.

Mormon *Fratercula*, *Temm. Man. d'Orn.* tom. II. p. 933. *Puffin*, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. II. p. 404. *Selb. Illust.* vol. II. p. 439. pl. 83*.

DIMENS. Entire length thirteen inches six lines: length of the bill (from the forehead) one inch eleven lines, (from the gape) one inch six lines; height of the bill at the base one inch seven lines; length of the tarsus one inch one line; of the tail two inches two lines; from the carpus to the end of the wing six inches six lines.

DESCRIPT. (*Adult at all seasons.*) Crown of the head, a broad collar surrounding the neck, and all the upper parts of the body, deep glossy black: quills dusky brown: cheeks, throat, and a broad band above the eyes, grayish white; breast, belly, and rest of the under parts, pure white: bill bluish gray at the base, orange-yellow in the middle, bright red at the tip; upper mandible with three obliquely transverse furrows; lower mandible with two: irides, and horny appendages to the eyelids, grayish white: orbits red: legs orange-red; claws black. (*Young of the year.*) Bill much smaller and narrower: the sides smooth, with the transverse furrows very indistinctly marked; colour yellowish brown: space between the eye and the bill dusky ash; cheeks and throat of a deeper gray than in the adult bird; collar inclining to dusky gray: legs dull red. (*Egg.*) White; indistinctly spotted with ash-gray: long. diam. two inches three lines; trans. diam. one inch seven lines.

Found in great abundance on many parts of the British coast during the spring and summer months. First seen about the middle of April. Breeds in the crevices of rocks, and, in some places, in holes in the ground burrowed out for the purpose. Deposits a single egg about the middle of May. Towards the end of August or beginning of September, leaves the British kingdom, and retires southward for the Winter. Food, small fish, particularly sprats, marine crustacea and insects, occasionally sea-weed.

GEN. 105. ALCA, *Linn.*

276. *A. Torda*, Linn. (*Razor-Bill.*)—Wings reaching to the rump: bill black, with a transverse white band: a narrow white line in advance of the eye.

A. Torda, *Temm. Man. d'Orn.* tom. II. p. 936. Razor-Bill, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. II. p. 399. Razor-Bill Auk, *Selb. Illust.* vol. II. p. 435. pl. 83.

DIMENS. Entire length seventeen inches: length of the bill (from the forehead) one inch six lines, (from the gape) one inch eleven lines; of the tarsus one inch three lines; of the tail three inches four lines; from the carpus to the end of the wing seven inches six lines.

DESCRIPT. (*Adult in winter.*) Crown of the head, nape, sides of the neck, and all the upper parts, deep black: from the eye to the middle of the bill a narrow longitudinal interrupted white line: secondaries tipped with white, forming a transverse bar on the wings: throat, front of the neck, breast, and all the under parts, pure white; sides of the occiput white mixed with cinereous, bounded beneath by a narrow black band reaching backwards from the eyes: bill black, with three or four transverse furrows, the middle one pure white: irides hazel: legs dusky. (*Adult in summer.*) The narrow streak from the eye to the middle of the bill pure uninterrupted white: head, throat, and all the upper and fore part of the neck, deep brownish black: the rest as in winter. (*Young of the year.*) Similar to the adult in winter plumage, but easily distinguished by the smaller and narrower bill, without the transverse white furrow: the white line from the bill to the eye obscure and ill-defined: crown and nape cinereous black; sides of the neck, and towards the occiput, white, tinged with ash-colour: under parts pure white. (*Egg.*) White; blotched and spotted with dark red brown and blackish brown: long. diam. two inches nine lines; trans. diam. one inch ten lines.

Like the Puffin, a regular summer visitant. Common on all the rocky parts of the coast during the breeding season, and generally found in large companies. Deposits a single egg, about the beginning of May, on the tops and projecting shelves of the highest cliffs. Food, sprats and other small fish, and marine crustacea. *Obs.* The *Black-billed Auk* of English authors (*A. Pica*, Gmel.) is this species in its immature state.

277. *A. impennis*, Linn. (*Great Auk*).—Wings very short, not reaching to the rump: bill black: an oval white patch in advance of the eye.

A. impennis, *Temm. Man. d'Orn.* tom. II. p. 939. Great Auk, *Mont. Orn. Dict.* *Bew. Brit. Birds*, vol. II. p. 397. *Selb. Illust.* vol. II. p. 433. pl. 82.

DIMENS. Entire length thirty-two inches: length of the bill (from the forehead) three inches six lines, (from the gape) four inches two lines; of the tarsus two inches five lines; of the tail three inches three lines; from the carpus to the end of the wing seven inches.

DESCRIPT. (*Summer plumage.*) Head, nape, back, wings, and tail, deep glossy black; front and sides of the neck black tinged with brown: immediately before the eye a large oval white patch: flanks dark ash-colour; rest of the under parts pure white: wings very short, incapacitated for flight; secondary quills tipped with white: bill black; crossed obliquely by several transverse furrows and ridges: irides and legs black. (*Winter plumage.*) "Unknown." TEMM. (*Egg.*) Dirty white, tinged with yellow; blotched and streaked, principally at the larger end, with black: long. diam. four inches ten lines: trans. diam. two inches nine lines.

A native of high northern latitudes, very rarely visiting the British coast. Occasionally seen in the Orkneys, and off the island of St. Kilda. Resides entirely in the open sea, never coming on land except for the purpose of breeding. Swims and dives with the greatest rapidity. Lays one egg in the clefts of rocks close to the water's edge. Food, fish and marine vegetables.

GEN. 106. PHALACROCORAX, *Briss.*

278. *P. Carbo*, Steph. (*Common Cormorant*.) — A white gorget under the throat: tail of fourteen feathers.

Carbo Cormoranus, *Temm. Man. d'Orn.* tom. ii. p. 894. *Corvorant*, *Mont. Orn. Dict.* *Cormorant*, *Bew. Brit. Birds*, vol. ii. p. 379. *Leach, Zool. Misc.* vol. iii. pl. 123. *Common Cormorant*, *Selb. Illust.* vol. ii. p. 446. pl. 84.

DIMENS. Entire length three feet: length of the bill (from the forehead) two inches ten lines, (from the gape) four inches.

DESCRIPT. (*Adult in winter.*) Crown of the head, neck, breast, and all the under parts, black, with blue and green reflections; throat dusky, mixed with whitish; on the neck a few indistinct whitish streaks: upper part of the back, scapulars, and wing-coverts, cinereous brown, glossed with purple and bronze, all the feathers broadly edged with deep black: lower back and rump bluish black: quills and tail dusky: bill dark horn-colour: naked skin encircling the eyes greenish yellow: irides green: guttural pouch yellowish: legs black. (*Summer plumage.*) The feathers on the occiput and nape slender and elongated, forming a crest of a bluish black colour, glossed with green: a broad gorget of pure white on the throat: crown and upper part of the neck variegated with numerous long slender white feathers of a silky texture: upon the thighs a large patch of pure white: the rest as in winter. (*Young of the year.*) "Crown, nape, and back, deep brown, with a slight gloss of green: gorget whitish gray; front of the neck, and all the under parts, grayish brown, with whitish spots, more particularly on the breast and middle of the belly, intermixed: upper part of the back, scapulars, and wing-coverts, ash-gray, the feathers broadly edged with deep brown: bill pale brown: irides brown." **TEMM.** (*Egg.*) White, tinged with pale blue; the outer surface of a rough calcareous texture: long. diam. two inches nine lines; trans. diam. one inch seven lines.

Common on most parts of the British coast throughout the year. Occasionally observed inland on the banks of rivers. Swims and dives with great ease. Breeds on the summits of lofty rocks, sometimes in trees. Constructs a nest of sea-weed, and lays three or four eggs. Feeds entirely on fish. *Obs.* The *Crested Cormorant* of Bewick is this species in its summer plumage.

279. *P. cristatus*, Steph. (*Crested Shag*.) — The whole of the plumage, throat included, deep green: tail short, of twelve feathers.

Carbo cristatus, *Temm. Man. d'Orn.* tom. ii. p. 900. *Shag*, and *Crested Shag*, *Mont. Orn. Dict. & Supp.* *Shag*, *Bew. Brit.*

Birds, vol. 11. p. 387. Crested Shag, or Green Cormorant, *Selb. Illust.* vol. 11. p. 450. pl. 86. *Gould, Europ. Birds*, part x.

DIMENS. Entire length twenty-eight inches: length of the bill (from the forehead) two inches six lines, (from the gape) three inches six lines; of the tarsus two inches two lines; of the tail five inches two lines; from the carpus to the end of the wing ten inches nine lines.

DESCRIPT. (*Adult in winter.*) The whole plumage of a rich deep glossy green: upper part of the back, scapulars, and wing-coverts, with bronze and purple reflections, each feather surrounded with a narrow edging of velvet-black: tail short and rounded; black: base of the bill and gular pouch gamboge-yellow: bill brown: irides green: legs black. (*Summer plumage.*) Remarkable for an elegant crest of long spreading feathers rising from the forehead between the eyes, capable of erection: no white feathers on the neck and thighs as in the last species. (*Young of the year.*) "Distinguished by the whole of the upper part of the plumage being brown, slightly tinted with green; the under surface brownish ash, more or less inclining to white." *GOULD. (Egg.)* White, tinged with pale greenish blue in patches; the outer surface rough and calcareous: long. diam. two inches five lines; trans. diam. one inch five lines.

Found principally in the rocky islands of the North of England and Scotland. Common in the Orkneys. Breeds on the shelves of steep rocks. Nest formed of sea-weed. Eggs from three to five in number. Feeds on fish. Is never seen inland.

(24.) *P. Graculus*, Steph. in Shaw's Gen. Zool. vol. xiiii. part i. p. 82. *Carbo Graculus*, Temm. Man. d'Orn. tom. 11. p. 897. *Black Cormorant*, Gould, *Europ. Birds*, part x.

This species has been considered as British by nearly all our own Ornithologists, though upon very insufficient authority. It appears to have been confounded with the *P. cristatus*, from which it differs in the darker colour of its plumage, as well as in other characters which will be found pointed out in Mr Gould's work above referred to.

GEN. 107. SULA, *Briss.*

280. *S. Bassana*, *Briss. (Common Gannet.)*—Crown and occiput pale buff; quills black; the rest of the plumage white.

S. alba, *Temm. Man. d'Orn.* tom. 11. p. 905. *Gannet*, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 390. *Solan Gannet*, *Selb. Illust.* vol. 11. p. 455. pls. 86*, & 87.

DIMENS. Entire length three feet.

DESCRIPT. (*Adult in perfect plumage.*) Crown of the head, and occiput, pale buff-yellow; all the rest of the plumage milk-white, with the exception of the quills and bastard winglet, which are black: bill bluish gray at the base, the tip white: naked skin surrounding the eyes pale blue: at the corners of the mouth a dusky membrane having the appearance of being a prolongation of the gape: chin destitute of feathers, and of a dusky colour; capable of great distention, forming a kind of pouch: irides yellow: acrotarsia and acrodactyla with longitudinal streaks of pale green; membranes dusky; claws white: tail-feathers strong and pointed; twelve in number; the middle ones longest. (*Young*

during the first year.) All the upper plumage dusky brown; under parts brown, varied with ash-gray: bill, naked membranes, and irides, brown: tail simply rounded. *After the second moult*, the head, neck and breast, are grayish black, with a small triangular white spot at the tip of each feather: back, rump, and wings, the same, but the spots on these parts larger and more scattered: abdomen whitish, the feathers margined with cinereous brown: quills and tail grayish black; the latter more conical, with the shafts of the feathers white: bill grayish brown, whitish towards the tip: acetarsia and acrodactyla greenish brown; the longitudinal streaks grayish white: membranes cinereous brown: claws whitish. *At each succeeding moult* the plumage becomes whiter, till the commencement of the fourth year, when it is fully matured. (*Egg.*) Dull white, tinged with green: long. diam. three inches three lines; trans. diam. one inch ten lines.

Plentiful during the breeding season on some parts of the northern coasts, particularly in the Isle of Bass in the Frith of Forth, St. Kilda, and some other of the Scotch Islands. Migrates southward in the Autumn, and, during Winter, may be observed in most parts of the British channel. At such times generally keeps far out at sea, but has been killed inland in a few rare instances. Feeds on herrings, pilchards, and other fish, on which it darts with great force and velocity. Nest placed on steep inaccessible rocks. Lays a single egg.

(25.) *Pelecanus Onocrotalus*, Temm. Man. d'Orn. tom. 11. p. 891. *Pelican*, Mont. Orn. Dict. Supp. Gould, Europ. Birds, part xii.

An individual of this species, which is a native of Eastern Europe, is recorded (according to Montagu) to have been shot in England, in May 1663, at Horsey Fen. It is conjectured, however, that it might have escaped from confinement.

GEN. 108. STERNA, Linn.

(1. STERNA, Steph.)

281. *S. Caspia*, Pall. (*Caspian Tern.*)—Bill thick and strong, bright red: tarsus one inch six lines in length: tail short, forked.

S. Caspia, Temm. Man. d'Orn. tom. 11. p. 733. *Caspian Tern*, Lath. Syn. vol. 111. p. 350. *Selb. Illust.* vol. 11. p. 463.

DIMENS. Entire length twenty-one inches: length of the bill (from the forehead) two inches nine lines, (from the gape) three inches two lines; of the tarsus one inch six lines; of the tail six inches; from the carpus to the end of the wing sixteen inches six lines.

DESCRIPT. (*Summer plumage.*) Forehead, crown, and long occipital feathers, deep black: nape, back, scapulars and wing-coverts, bluish ash: quills brown, tinged with ash-gray: sides of the head, fore part of the neck, and all the under parts of the plumage, pure white: tail pale ash-gray: bill coral-red: legs black. Wings reaching about four inches and a half beyond the tail. (*Winter plumage.*) Forehead, and a part of the crown, pure white: occiput variegated with black and white: the rest as in summer. (*Young of the year.*) "Forehead and crown as in the adult in winter plumage; rest of the upper parts ash-coloured brown, with large dusky spots and transverse bars: tail with a large dusky space at

the extremity: quills almost entirely of the same colour: under parts white: bill dull red; the tip dusky." TEMM. (*Egg*.) Yellowish stone-colour, spotted with ash-gray and dark red-brown; long. diam. two inches six lines; trans. diam. one inch eight lines.

Inhabits the borders of the Baltic, the Caspian Sea, and Archipelago. Is also occasionally seen on the coasts of France and Holland. In England, a few specimens have occurred at Yarmouth and Aldborough. One killed at this last place is in the Museum of the Cambridge Philosophical Society. Feeds on fish. Eggs two or three in number; deposited in a hole in the sand, or on the rocks by the edge of the sea.

282. *S. Cantiaea*, Gmel. (*Sandwich Tern*.)—Bill long; black, the tip yellowish: legs short; black: tarsus one inch: tail long, very much forked, shorter than the wings.

S. Cantiaea, Temm. *Man. d'Orn.* tom. 11. p. 735. *Sandwich Tern*, Mont. *Orn. Dict. & Supp. with fig.* Bew. *Brit. Birds*, vol. 11. p. 189. *Selb. Illust.* vol. 11. p. 464. pl. 88. f. 3.

DIMENS. Entire length eighteen inches: length of the bill two inches: breadth, wings extended, two feet nine inches. MONT.

DESCRIPT. (*Summer plumage*.) Forehead, crown, and long occipital feathers, deep black; nape and upper part of the back pure white; middle and lower regions of the back, scapulars and wing-coverts, pale bluish ash: all the under parts pure white, tinged on the breast and fore part of the neck, with rose-red: quills deep ash-gray, having a velvety appearance, bordered on their inner webs with a broad band of pure white; some of the primaries tipped with black: tail white; considerably forked: bill black for two-thirds of its length from the base, the tip straw-yellow: legs black; the under surface of the toes and membranes yellow: claws black. (*Winter plumage*.) Forehead and crown white, spotted with black towards the occiput; the long occipital feathers deep black, but fringed with white: under parts pure white, without the rosy tinge on the neck and breast: the rest as in winter. (*Young of the year*.) Crown and occiput a mixture of black, white, and pale reddish brown: all the under parts pure white: upper part of the back and scapulars reddish, with transverse bars of dusky brown; some of the larger scapulars broadly bordered with brown: tips of the wing-coverts with a crescent-shaped edging of this colour: primary and secondary quills dusky ash, edged and tipped with white: tail ash-coloured at the base; the remaining portion dusky, with the tips of the feathers white: bill livid black; the extreme tip yellowish. (*Egg*.) Yellowish stone-colour; thickly spotted with ash-gray, orange-brown, and deep red-brown: long. diam. two inches; trans. diam. one inch five lines.

A summer visitant. Frequents the coasts of Kent, Sussex, and Suffolk. Occasionally met with in other parts of the country, but not a generally diffused species. Breeds in large societies. Eggs two or three, laid on sandy shores or on rocks. Feeds on fish.

283. *S. Dougallii*, Mont. (*Roseate Tern*.)—Bill slender; black, the base reddish: legs orange: tarsus nine lines: tail greatly forked, extending far beyond the wings.

S. Dougalli, *Temm. Man. d'Orn.* tom. II. p. 738. Roseate Tern, *Mont. Orn. Dict. & Supp. with fig. Bew. Brit. Birds*, vol. II. p. 192. *Selb. Illust.* vol. II. p. 470. pl. 89. f. 1, & 2.

DIMENS. Entire length fifteen inches six lines: length of the bill (from the forehead) one inch seven lines and a half; of the tarsus nine lines; of the tail seven inches; from the carpus to the end of the wing nine inches three lines.

DESCRIPT. (*Summer plumage.*) Forehead, crown, and long occipital feathers, deep black; cheeks, throat, neck, and all the under parts, pure white, tinged on the fore part of the neck, breast, and belly, with rose-red: back, scapulars, and wing-coverts, pale ash-gray: first quill with the outer web hoary black; the rest gray; all of them with a deep border of white on their inner webs: tail white, greatly forked; the outer feathers very long and subulate, extending upwards of two inches beyond the wings: bill long and slender, black, passing into orange-red at the base: legs bright orange: claws small and black. (*Young.*) "Bill brownish black; the base orange-yellow: forehead and crown cream-yellow, tinged with gray: region of the eyes, ear-coverts, and nape, grayish black, mixed with yellowish white: throat, sides of the neck, and under parts, white: ridge of the wings blackish gray, margined paler: back and wing-coverts bluish gray, marbled with grayish black and yellowish white: tail-feathers with the exterior webs gray, the interior and tips white: quills gray, margined with white: legs pale gallstone-yellow." **SELB.** (*Egg.*) Yellowish stone-colour; spotted and speckled with ash-gray and dark brown: long. diam. one inch nine lines and a half; trans. diam. one inch two lines and a half.

First noticed on the Cumbrey Islands in the Frith of Clyde, by Dr Macdougall, who communicated the species to Montagu. Since observed on other parts of the Scotch coast, and also in the Fern Islands, to which last locality, according to Selby, they resort annually to breed. Eggs two or three in number. Food, small fish.

284. S. *Hirundo*, Linn. (*Common Tern.*) — Bill moderate; red, the tip black: legs red: tarsus nine lines and a half: tail very much forked; as long as, or a little shorter than, the wings.

S. *Hirundo*, *Temm. Man. d'Orn.* tom. II. p. 740. Common Tern, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. II. p. 185. *Selb. Illust.* vol. II. p. 468. pl. 90. f. 1.

DIMENS. Entire length thirteen inches six lines: length of the bill (from the forehead) one inch seven lines and a half, (from the gape) two inches two lines; of the tarsus nine lines and a half; of the tail four inches six lines; from the carpus to the end of the wing ten inches four lines: breadth, wings extended, two feet six inches eight lines.

DESCRIPT. (*Summer plumage.*) Forehead, crown, and long occipital feathers, deep black; hind part of the neck, back, scapulars, and wing-coverts, bluish ash: all the under parts white, the breast and belly faintly tinged with ash-colour: quills ash-gray passing into dusky brown at the tips, with a large oblong white space on their inner webs: tail whitish; the outer webs of the two exterior feathers dusky ash: bill red, passing into dusky at the tip: feet bright red. The *winter plumage* differs simply in having the forehead and crown of a more dull black. (*Young of the year.*) Forehead and anterior part of the crown dirty white; rest

of the crown towards the occiput marked with large dusky spots; the long occipital feathers brownish black with whitish tips: rest of the upper parts dull bluish ash, irregularly spotted with pale reddish brown, all the feathers whitish at the tips: under parts whitish: many of the quills tipped with white: lesser wing-coverts near the top of the wing dusky gray with whitish edges: tail ash-gray, passing into white on the inner webs, and towards the tips of the feathers: bill much shorter than in the adult bird, dusky at the tip, dull orange at the base: legs orange. (*Egg.*) Yellowish stone-colour, blotched and spotted with ash-gray and dark red-brown: long. diam. one inch eight lines; trans. diam. one inch two lines.

Common on most parts of the British coast during the spring and summer months. Retires in the Autumn. Occasionally observed inland on fresh waters. Breeds on flat shingly shores, and lays from two to four eggs. Of noisy and restless habits: constantly on wing. Food, insects and small fish. *Obs.* The *Brown Tern* of English authors is probably this species in immature plumage.

235. *S. arctica*, Temm. (*Arctic Tern.*)—Bill slender; red throughout its whole length: legs red: tarsus seven lines: tail very much forked; longer than the wings.

S. arctica, Temm. *Man. d'Orn.* tom. II. p. 742. *Arctic Tern*, Shaw, *Gen. Zool.* vol. XIII. p. 152. *Selb. Illust.* vol. II. p. 473. pl. 90. f. 2.

DIMENS. Entire length fifteen inches: length of the bill (from the forehead) one inch four lines, (from the gape) one inch ten lines; of the tarsus seven lines; of the tail seven inches nine lines; from the carpus to the end of the wing eleven inches.

DESCRIPT. Very similar to the last species, but may always be distinguished by the much shorter tarsus, and the bill which is also shorter and generally of one colour throughout. (*Summer plumage.*) Forehead, crown, and long occipital feathers, deep black; rest of the upper parts bluish ash: throat, front of the neck, breast and belly, of the same colour as the back; vent, under tail-coverts, and a band beneath the eyes, pure white: tail very much forked; always longer than in the last species, extending (in the *adult male*) one inch two lines beyond the wings: tarsi and toes very short; of a deep crimson-red: bill crimson-red. (*Egg.*) Colour the same as in the last species: long. diam. one inch seven lines; trans. diam. one inch one line.

Inhabits the Arctic Regions; resorting to the northern coasts of Britain during the summer months. According to Selby, very abundant in the Fern Islands, where they arrive about the middle of May and stay till August. Eggs laid on the gravelly beach or bare ground; three or four in number. Food, small fish, particularly the fry of the sand-eel (*Ammodytes Tobianus*.)

236. *S. minuta*, Linn. (*Lesser Tern.*)—Bill orange, the tip black: legs orange: tarsus seven lines and a half: tail very much forked, shorter than the wings: forehead white.

S. minuta, *Temm. Man. d'Orn.* tom. 11. p. 752. Lesser Tern, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 187. *Selb. Illust.* vol. 11. p. 475. pl. 89. f. 3, & 4.

DIMENS. Entire length nine inches six lines: length of the bill (from the forehead) one inch two lines, (from the gape) one inch eight lines; of the tarsus seven lines and a half; of the middle toe, claw included, nine lines; of the tail three inches three lines; from the carpus to the end of the wing seven inches one line: breadth, wings extended, twenty inches ten lines.

DESCRIPT. (*Summer and winter plumage.*) Forehead, and a streak above the eyes, pure white; between the bill and the eye a broad black streak; crown, occiput, and nape, deep black; back, scapulars, and wing-coverts, pale bluish ash: first three quills blackish gray, their inner webs broadly edged with white: cheeks, all the under parts, and tail, pure white: bill yellowish orange; the tip black: irides black: legs and toes orange-yellow. (*Young of the year.*) Forehead yellowish white; crown, occiput and nape, brownish, with dusky streaks; before and behind the eyes a black spot; back, scapulars and wing-coverts, yellowish white, with a crescent-shaped dusky bar near the extremity of each feather: quills cinereous gray, edged and tipped with yellowish white: tail nearly even at the end, white tinged with cinereous, the tips of the feathers yellowish: throat, and all the under parts, white: bill shorter than in the adult bird, pale yellow, the tip blackish brown: legs dull yellow. (*Egg.*) Stone-colour, spotted and speckled with ash-gray and dark chestnut-brown: long. diam. one inch four lines; trans. diam. eleven lines.

Common on some parts of the British coast during the summer months, but not so generally diffused as the *S. Hirundo*. Habits similar to those of that species. Eggs two or three in number, deposited on the shingly beach a little above high water-mark. Food, marine worms and winged insects.

287. *S. nigra*, Linn. (*Black Tern.*) — Bill black: legs reddish brown: membranes of the toes deeply emarginated: tarsus seven lines and a half: tail not much forked, shorter than the wings.

S. nigra, *Temm. Man. d'Orn.* tom. 11. p. 749. Black Tern, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 195. *Selb. Illust.* vol. 11. p. 477. pl. 91.

DIMENS. Entire length ten inches three lines: length of the bill (from the forehead) one inch two lines, (from the gape) one inch seven lines and a half; of the tarsus seven lines and a half; of the tail three inches three lines; from the carpus to the end of the wing eight inches eight lines: breadth, wings extended, twenty-three inches six lines.

DESCRIPT. (*Summer plumage.*) Head and neck deep black: breast and abdomen dark ash-colour: all the upper parts, including the rump and tail, deep bluish gray: vent and under tail-coverts pure white: wings, when closed, extending one inch seven lines beyond the tail: bill black: irides brown: legs purplish brown. (*Winter plumage.*) Forehead, space between the bill and the eyes, throat and all the fore part of the neck, white: the rest as in summer. *In the intermediate seasons,*

these parts are often found mottled with black upon a white ground. (*Young of the year.*) "Forehead, space between the eye and the bill, sides and front of the neck, as well as all the under parts, pure white: on the sides of the breast a large patch of dusky ash: before the eyes a crescent-shaped spot of the same colour: crown, occiput and nape, black: back and scapulars brown, the feathers edged and tipped with reddish white: wings, rump and tail, ash-colour; the wing-coverts tipped with reddish white: bill brown at the base: irides brown: legs livid brown." TEMM. (*Egg.*) Dark olive-brown, blotched and spotted with black, principally at the larger end: long. diam. one inch five lines: trans. diam. one inch.

A more inland species than any of the foregoing. Found principally in marshes, and on the banks of rivers and lakes. First seen towards the latter end of April: departs in October. Common in Romney Marsh in Kent, and also in the fens of Cambridgeshire and Lincolnshire, where they breed. Eggs two to four in number, deposited on the bare grass in swampy and sedgy spots. Young hatched in the beginning of July. Food, worms and aquatic insects. *Obs.* The *S. naevia* of Gmelin is the young of this species.

288. *S. Anglica*, Mont. (*Gull-billed Tern.*) — Bill short and thick; entirely black: legs long; black: claws nearly straight: tarsus one inch three or four lines: tail not much forked, considerably shorter than the wings.

S. Anglica, Temm. *Man. d'Orn.* tom. II. p. 744. Gull-billed Tern, *Mont. Orn. Dict. Supp. with fig. Selb. Illust.* vol. II. p. 480. pl. 88. f. 1.

DIMENS. Entire length thirteen inches six lines: length of the bill (from the forehead) one inch six lines, (from the gape) one inch ten lines and a half; of the tarsus one inch two lines and a half to four lines and a half*.

DESCRIPT. (*Summer plumage.*) Forehead, crown, occiput and nape, deep black; the feathers on this last part long and silky: all the rest of the upper plumage pale bluish ash: quills dusky gray, having a hoary appearance: the tips of the first five black: under parts white: wings extending three inches beyond the extremity of the tail: bill deep black; remarkably thick and strong; gonys of the lower mandible ascending; mental angle very prominent, as in the genus *Larus*: irides brown: legs long; black, with a reddish tinge: toes long; the claws unusually straight; membranes deeply emarginated. (*Winter plumage.*) Forehead and crown white; anterior angle of the eye, and spot behind the ears, grayish black: the rest as in summer, with the exception of the primary quills, which are not quite so deeply coloured at the tips. (*Young of the year.*) "Crown white, with small longitudinal brownish streaks: back and wings bluish ash, mixed with gray and yellowish brown: all the under parts pure white: quills brownish ash: tail very little forked, ash-colour, the tips of the feathers white: base of the bill yellowish; the rest, towards the tip, dusky brown: legs brown." TEMM. (*Egg.*) Dark olive-brown, spotted with ash-colour and two shades of dark red-brown: long. diam. one inch eleven lines; trans. diam. one inch four lines.

* A difference of two lines was found in the length of the tarsus in two British-killed specimens in the British Museum.

A rare species in this country, first described by Montagu, who obtained specimens from the coast of Sussex. According to Temminck, very abundant in Hungary. Said to breed on the marshy borders of salt-water lakes, and to lay three or four eggs. Food, winged insects.

(2. *ANOUS*, *Steph.*)

289. *S. stolidus*, Linn. (*Black Noddy*).—Forehead and crown whitish; the rest of the plumage, as well as the bill and legs, black.

S. stolidus, *Lath. Ind. Orn.* vol. II. p. 805. *Noddy*, *Lath. Syn.* vol. III. p. 354. *Black Noddy*, *Steph. in Shaw's Gen. Zool.* vol. XIII. p. 140. pl. 17.

DIMENS. Entire length fifteen inches: length of the bill (from the forehead) one inch nine lines, (from the gape) two inches; of the tarsus eleven lines; of the tail five inches; from the carpus to the end of the wing ten inches.

DESCRIPT. Forehead and crown whitish, passing into ash-gray on the occiput and nape: all the rest of the plumage brownish black, deepest on the quills and tail-feathers: wings just reaching to the extremity of the tail, which is even, or somewhat rounded, at its tip: bill and legs black: the membranes of the toes entire, without any indication of a notch.

Two adult individuals of this species were taken a few years since during Summer on the Wexford coast, between the Tasker Lighthouse and Dublin Bay. From one of these, which was obligingly submitted to my examination by Mr W. Thompson of Belfast, the above measurements and description were taken. Not previously known as British. Frequently met with at sea, but generally within the Tropics. Stated by Latham to breed in the Bahama Islands, laying its eggs on the bare rocks.

GEN. 109. *LARUS*, Linn.

(1. *XEMA*, *Leach*.)

290. *L. Sabini*, Sabine. (*Sabine's Gull*.)

Xema Sabini, *Leach in Ross's Voy. App.* p. LVII. with fig. *Larus Sabini*, *Sab. in Linn. Trans.* vol. XII. p. 520. pl. 29. *Sabine's Xeme*, *Steph. in Shaw's Gen. Zool.* vol. XIII. part I. p. 177. pl. 20.

DIMENS. Entire length from twelve inches six lines to fourteen inches: length of the bill one inch; of the tarsus one inch six lines; of the tail five inches: breadth, wings extended, about thirty-three inches.

SAB.

DESCRIPT. (*Adult male and female in summer.*) "The whole of the head, and upper part of the neck, a very dark ash or lead-colour; the remainder of the neck behind and before, as well as the breast and belly, pure white; a narrow black collar surrounding the neck at the meeting of the ash-colour and white: back, scapulars, and wing-coverts, ash-coloured, very much lighter than the head, but darker than the corresponding parts of the *Larus ridibundus*; the lower ends of the scapulars tipped with white: the first five primary quills with black shafts, the whole outer webs of these black, the edge of their upper webs white to within an inch and a half of the tips, the white sometimes continued to the tip; the tips of the first and second of these quills in some white, in

others black; the tips of the third, fourth, and fifth, white, giving the wing when closed a spotted appearance; the sixth primary quill with a white shaft, having the web more or less black, but principally white, with sometimes a black spot near the end; the other primaries, the secondaries, and tertials, white; the whole under parts of the wings white: wings extending an inch or more beyond the longest feather of the tail: upper and under tail-coverts white: the tail feathers twelve, the outer narrower than the central ones; the depth of the fork nearly an inch: bill with the base of both mandibles black as far as the angular projection of the lower mandible, the remainder yellow; inside of the mouth bright vermilion: irides dark, surrounded by a naked circle of the same colour as the inside of the mouth; a small white speck beneath the eye, scarcely perceptible: legs black." SAB. In the *immature* state and in the *adult winter* plumage, the black head is wanting. (Egg.) Of a regular shape, and not much pointed: colour olive, much blotched with brown: long. diam. one inch six lines. SAB.

Two individuals of this species, which was first observed in Greenland by Captain Sabine, are recorded by Mr Thompson* to have been killed in Ireland. The specimens, which are both in the plumage of the first year, are now in the Museums of the Natural History Society of Belfast and the Royal Society of Dublin. Not much is known respecting its habits. Said to lay two eggs on the bare ground, which are hatched in the last week in July. Food, marine insects.

(2. LARUS, *Steph.*)

* *Small; less than twenty inches in length.*

291. *L. minutus*, Pall. (*Little Gull.*)—Bill brownish red: legs bright vermilion: tarsus eleven or twelve lines: hind toe very small; the claw straight: all the quills tipped with white; the shafts of the primaries brown.

L. minutus, Temm. *Man. d'Orn.* tom. 11. p. 787. *Little Gull*, Mont. *Orn. Dict. App. to Supp. with fig.* Lath. *Syn.* vol. III. p. 391. *Selb. Illust.* vol. 11. p. 484. pl. 92. Gould, *Europ. Birds*, part xi.

DIMENS. Entire length rather exceeding ten inches: length of the bill (from the forehead) rather more than nine lines: of the tarsus rather more than one inch; from the carpus to the end of the wing eight inches six lines. MONT.

DESCRIPT. (*Adult in winter.*) Forehead, cheeks, and a space behind the eyes, throat, and all the under parts, pure white: occiput, nape, a spot at the anterior angle of the eye, and ear-coverts, deep blackish gray: upper parts of the body and wings pale bluish ash: primary and secondary quills broadly tipped with white: inside of the wings deep blackish gray: tail white: bill and irides dusky brown: legs bright vermilion-red. (*Adult in summer.*) "All the head, and upper part of the neck, black: a white crescent-shaped spot behind the eyes; lower part of the neck, and all the under parts, white, with a faint blush of rose-red: back, scapulars, and wings, very pale bluish ash; rump and tail pure white: primary quills cinereous; all of them, as well as the secondaries, tipped with white: bill deep lake-red: irides dark brown: legs crimson." TEMM. (*Young of the year.*) "Forehead and crown white, tinged with gray: nape, and back part of the neck, and upper parts of the body, blackish

gray, tinged with clove-brown, the margins of the feathers fringed with grayish white: scapulars edged with white: greater coverts pale bluish gray, deeply edged with white: first four quills with the outer webs and tips black, the inner webs grayish white: under plumage white: tail white for two-thirds of its length, terminated by a broad black bar: legs livid or flesh-red." SELB. (*Egg*.) Unknown.

A very rare occasional visitant in this country. First noticed by Montagu, who describes a specimen shot on the Thames near Chelsea. Has since occurred at Yarmouth, and on one or two other parts of the eastern coast, as well as in Cornwall and Scotland. Said to be common in the East of Europe, frequenting rivers and lakes. Food, insects and worms. Nidification unknown.

292. *L. capistratus*, Temm. (*Brown-headed Gull*).—Bill and legs reddish brown: tarsus one inch seven lines: shafts of the outer primaries white: *in summer*, a brown hood reaching to the occiput.

L. capistratus, Temm. *Man. d'Orn.* tom. II. p. 785. *Flem. Brit. An.* p. 142. Yarr. in *Proceed. of Zool. Soc.* 1831. p. 151. Masked Gull, Shaw, *Gen. Zool.* vol. XIII. p. 204.

DIMENS. Entire length fifteen inches: length of the bill (above, from the first feathers) one inch and half a line, (from the gape) one inch ten lines; of the tarsus one inch seven lines; of the middle toe, nail included, one inch six lines; from the carpus to the end of the wing eleven inches eight lines. YARR.

DESCRIPT. (*Winter plumage*.) According to Temminck, exactly similar to that of the next species, from which however the present may always be distinguished by its smaller size, its more slender as well as shorter bill, shorter tarsi, and smaller feet: the colour of the feet is also reddish brown. (*Summer plumage*.) "Head, and upper part of the neck, brocoli-brown, bounded by blackish brown, descending lowest at the fore part, some of the dark feathers at the margin in front tipped with white; the remaining portion of the neck, breast, abdomen, vent and tail, pure white: upper surface of the wings pale ash-gray; under surface grayish white: primaries white, edged and tipped with black, broadest on the inner web; shafts white: bill, legs, and toes, brownish red; membranes chocolate-brown." YARR. (*Egg*.) Unknown.

But little is known of the habits of this species, which was first characterized as distinct from the two following, by Temminck. Inhabits the Shetland and Orkney Islands. Has been also shot in Ireland, in the neighbourhood of Belfast, by Mr Thompson. Food and nidification unknown.

293. *L. ridibundus*, Linn. (*Black-headed Gull*).—Bill and legs bright red: tarsus one inch ten lines: a large white space in the middle of the primaries: *in summer*, a deep brown hood reaching below the occiput.

L. ridibundus, Temm. *Man. d'Orn.* tom. II. p. 780. Black-headed Gull, *Mont. Orn. Dict. & Supp. Selb. Illust.* vol. II. p. 486. pl. 92. Black-headed and Red-legged Gulls, Bew. *Brit. Birds*, vol. II. pp. 222, & 225.

DIMENS. Entire length seventeen inches: length of the bill (from the forehead) one inch four lines, (from the gape) two inches one line; of the tarsus one inch ten lines; of the middle toe, nail included, one inch eight lines; of the tail four inches eight lines; from the carpus to the end of the wing twelve inches nine lines: breadth, wings extended, three feet three inches five lines.

DESCRIPT. (*Winter plumage.*) Head, neck, and tail, pure white, with the exception of a small black spot at the anterior angle of the eye, and a dusky patch on the ear-coverts: back, scapulars, secondary quills, and greater portion of the wing-coverts, pale bluish ash; some of the long coverts annexed to the primaries, as well as the spurious winglet, white: primaries themselves white, passing into dusky ash towards the margins of the inner webs, and broadly tipped with black; (according to Temminck, *in very old birds*, the extreme tips of the primaries are *white*;) first primary with the outer web black from the base nearly to the tip, the inner edging of dusky ash very narrow; under surface of the wings dark ash-gray: breast, belly, and abdomen, white, in some individuals faintly tinged with flesh-red: irides deep brown: bill and legs bright vermilion-red. (*Summer plumage.*) Head, and upper part of the neck, deep brown: at the posterior angle of the eye a white spot; lower part of the neck, and all the under plumage, fine white, tinged with rose-red: bill and legs deep purplish red: the rest as in winter. (*Young of the year.*) Head and occiput very pale brown; a large white spot behind the eyes: under parts, and a collar on the nape, white; the white on the front of the neck with a faint reddish tinge: flanks with brown crescent-shaped marks: back, scapulars, and middle coverts, deep brown, the feathers with yellowish edges; upper edge of the wing, rump, and greater part of the tail-feathers, white, these last tipped with dusky brown: primaries white at the base and on the inner webs, black on the outer, and at the tips; greater coverts bluish ash: bill livid at the base; the tip black: legs yellowish. *After the first autumnal moult*, the back and wings are bluish ash, with a few brown feathers intermixed, some of those on the wings still with yellowish edges: forehead, and all the under parts, pure white; head white, spotted with pale ash-colour: ear-coverts, and a spot at the anterior angle of the eye, brown: tail still with a dark bar at its tip: bill reddish at the base: the tip brown. *In this state they remain till the following spring, when they assume the perfect summer plumage.* (*Egg.*) Yellowish olive-brown, spotted with two shades of darker brown: long. diam. two inches one line and a half; trans. diam. one inch six lines.

A common species; frequenting inland marshes and the banks of rivers. Resorts to the sea-coast in Winter. Breeds in fens and low swampy meadows, laying three eggs on tufts of grass previously trodden down. Food, insects, worms, and small fish. *Obs.* The *Red-legged Gull* (*L. cinerarius*, *Gmel.*) is this species in its winter plumage. The *Brown-headed Gull* (*L. erythropus*, *Gmel.*) is the young of the year after the first autumnal moult.

294. *L. Atracilla*, Linn. (*Laughing Gull*).—Bill and legs deep brownish red: tarsus one inch ten lines: primaries entirely black; extending considerably beyond the tail: *in summer* a lead-coloured hood reaching below the occiput.

L. Atracilla, *Temm. Man. d'Orn.* tom. II. p. 779. *L. ridibundus*, *Wils. Amer. Orn.* vol. IX. pl. 74. f. 4. *Laughing Gull*, *Mont. Orn. Dict.*

DIMENS. Entire length eighteen inches. MONT.

DESCRIPT. (*Winter plumage*.) According to Temminck, unknown. (*Summer plumage*.) "Head, and upper part of the neck, deep lead-gray, the hood extending lower in front than behind; a white spot above and beneath the eyes; lower part of the neck, breast, belly, and tail, pure white: back, wings, and secondary quills, lead-gray; tips of the secondaries white; the whole of the primaries, which extend two inches beyond the tail, deep black: bill and legs very deep lake-red." TEMM. (*Egg*.) Dun clay-colour, thinly marked with small irregular touches of pale purple and pale brown: long. diam. two inches and a quarter; trans. diam. one inch and a half. WILS.

But little known as a British species. Montagu records having seen five individuals together feeding in a pool upon the Shingly Flats near Winchelsea in Sussex, and two others near Hastings. One was shot. According to Temminck, not uncommon in the South of Europe. Food, the remains of fish and crustacea, and also insects. Said by Wilson to breed in marshes, and to lay three eggs.

295. *L. tridactylus*, Lath. (*Kittiwake Gull*.) — Bill greenish yellow: legs dusky: tarsus one inch four lines: a small knob instead of a back toe: primaries bluish ash, the outer one edged externally with black.

L. tridactylus, Temm. *Man. d'Orn.* tom. II. p. 774. Kittiwake, Mont. *Orn. Dict. & Supp. Bew. Brit. Birds*, vol. II. pp. 218, & 220. *Scb. Illust.* vol. II. p. 493. pl. 94.

DIMENS. Entire length fifteen inches six lines: length of the bill (from the forehead) one inch four lines, (from the gape) two inches one line; of the tarsus one inch four lines; of the tail five inches five lines; from the carpus to the end of the wing twelve inches: breadth, wings extended, three feet one inch four lines.

DESCRIPT. (*Adult in winter*.) Crown, occiput, nape, and sides of the neck, pale bluish ash, with a few fine dusky streaks before the eyes; forehead, region of the eyes, and all the under parts, pure white: back, scapulars, and wing-coverts, bluish ash: quills the same; the first with the whole of the outer web black; first four primaries tipped with black, two or three of them with a small white spot at the extreme point; the fifth white at the extremity, with a black bar near the tip: rump and tail pure white: bill greenish yellow: inside of the mouth, and eye-lids, orange-red: irides brown: legs dusky brown, tinged with olivaceous. (*Summer plumage*.) The whole head and neck pure white, without any bluish ash on the nape, or dusky streaks before the eyes: the rest as in winter. (*Young of the year*.) Head, neck, and all the under parts, white, with the exception of a small spot at the anterior angle of the eye, a patch on the ear-coverts, and a larger crescent-shaped patch on the nape, which are dusky ash: back and wings deep bluish ash, the feathers tipped with dusky brown; the bend, and upper edge, of the wing, black; scapulars and secondary quills with large dusky spots; primaries black: tail black at the end, the extreme tips of the feathers whitish; the outer feather entirely white: bill, irides, and eye-lids, black. *After the first moult*, the back assumes the bluish ash, with still a few feathers intermixed spotted with brown; the spots on the head, and crescent-shaped patch on the nape, appear of a dark bluish ash; the wing-coverts also still spotted with dark brown; tail black at the extremity: bill greenish

yellow spotted with dusky: forehead, and all the under parts, white. *The plumage is not matured till after the second autumnal moult.* (Egg.) Stone-colour, thickly spotted with ash-gray and two shades of light brown: long. diam. two inches two lines and a half; trans. diam. one inch seven lines.

Not uncommon during the summer months on rocky coasts in the northern parts of the island. More rarely observed southwards. Breeds on the narrow ledges of steep cliffs, and lays three eggs. Food, small fish and marine insects. *Obs.* The *Tarrock Gull* of English authors is this species in its immature plumage.

296. *L. canus*, Linn. (*Common Gull*).—Bill greenish, yellow at the tip: legs greenish gray: tarsus two inches one line: wings reaching beyond the tail; first two primaries black, with a large white spot near the tips.

L. canus, Temm. *Man. d'Orn.* tom. II. p. 771. Common Gull, *Mont. Orn. Dict. & Supp.* Bew. *Brit. Birds*, vol. II. p. 216. *Selb. Illust.* vol. II. p. 490. pl. 93.

DIMENS. Entire length seventeen inches three lines: length of the bill (from the first feathers) one inch three lines, (from the gape) two inches two lines; of the tarsus two inches one line; of the naked part of the tibia seven lines; of the middle toe, nail included, one inch nine lines; of the tail five inches four lines; from the carpus to the end of the wing thirteen inches ten lines.

DESCRIPT. (*Adult in winter.*) Head, occiput, nape, and sides of the neck, white, spotted with blackish gray: all the under parts, rump and tail, pure white: back, scapulars and wing-coverts, fine bluish ash: first two primaries black, with a large white space near the tips; the rest black towards the extremities, the tips themselves, as well as those of the scapulars and secondary quills, white: bill greenish at the base, the tip ochre-yellow: inside of the mouth orange: irides brown: naked circle round the eyes reddish brown: legs greenish gray; the webs blotched with yellowish. (*Summer plumage.*) Head, occiput, nape and sides of the neck, pure white, without the dusky spots: bill ochre-yellow: naked circle round the eyes bright vermilion: legs pale ochre-yellow, spotted with bluish ash: the rest as in winter. (*Young of the year.*) A black spot before the eyes: all the upper plumage grayish brown, the feathers on the back and wings edged and tipped with yellowish white: those on the upper part of the back with fine streaks of this colour: forehead, and all the under parts, whitish, the breast and flanks spotted with ash-gray; throat, and middle of the belly, pure white: quills of a uniform dusky brown, neither tipped nor spotted with white: tail white, with a broad black bar near the extremity: base of the bill livid dirty white: the tip black: naked circle round the eyes brown: legs livid white. *In the second year*, the head, neck, rump, and under parts, are white, the two former streaked with dusky brown: back and scapulars bluish ash; wing-coverts still mottled with brown and white feathers: quills dusky brown, sometimes tipped with white: tail white, with a black bar towards the tip. *The plumage is not matured till after the second spring moult.* (Egg.) Dark olive-brown, spotted with two shades of darker brown: long. diam. two inches three lines; trans. diam. one inch six lines.

Common on all parts of the coast, and, during the winter months, occasionally observed inland at a considerable distance from the sea.

Breeds on the ledges of rocks. Nest constructed of sea-weed. Eggs two or three in number. Food, worms, marine insects, and small fish. *Obs.* The *Winter Gull* of English authors (*L. hybernus*, Gmel.) is this species in its immature plumage.

297. *L. eburneus*, Gmel. (*Ivory Gull*).—Bill strong; bluish gray, the tip yellow: legs black: naked part of the tibia very small: tarsus one inch six lines: membranes of the toes somewhat abbreviated: *adult* plumage pure white.

L. eburneus, *Temm. Man. d'Orn.* tom. II. p. 769. *Sabine in Linn. Trans.* vol. XII. p. 548. *Edmondst. in Wern. Mem.* vol. IV. p. 501. *Ivory Gull*, *Bew. Brit. Birds*, vol. II. p. 214. *Selb. Illust.* vol. II. p. 497. pl. 94*.

DIMENS. Entire length sixteen inches, (according to *Temminck*, nineteen inches): breadth, wings extended, three feet three inches and a half. **EDMONDST.**

DESCRIPT. (*Adult plumage*.) The whole plumage pure white, without spots of any kind: bill thick and strong; deep bluish gray at the base, the rest ochre-yellow: irides deep brown: wings extending an inch and a half beyond the tail: legs black; covered with a rough skin: tibia feathered nearly to the tarsal joint: membranes of the toes short, and slightly hollowed out: claws much hooked. In *young birds*, the plumage is pale grayish white, more or less spotted and barred with blackish brown: the tips of the quills, and a transverse bar at the extremity of the tail, of this last colour: bill blackish, the tip yellow: legs blackish gray. (*Egg*.) Unknown.

Inhabits the Arctic Regions. A rare and accidental visitant in this country. A solitary individual in its second year's plumage is recorded by Mr. Edmondston to have been killed in Balta Sound, Zetland, Dec. 13, 1822. It is now in the Edinburgh Museum. According to Selby, it has been also killed, in an immature state, in the Frith of Clyde. Food, according to Captain Sabine, blubber and the flesh of whales. Nidification unknown.

** *Large; exceeding twenty inches in length.*

298. *L. argentatus*, Brunn. (*Herring Gull*).—Bill yellow: legs livid flesh-colour: tarsus two inches six lines: wings reaching very little beyond the tail: shafts of the primaries black: mantle (in the *adult*) bluish ash.

L. argentatus, *Temm. Man. d'Orn.* tom. II. p. 764. *Herring Gull*, *Mont. Orn. Dict. & Supp.* *Bew. Brit. Birds*, vol. II. p. 207. *Selb. Illust.* vol. II. p. 504. pls. 96, & 96*.

DIMENS. Entire length twenty-three inches: length of the bill (from the forehead) two inches three lines, (from the gape) two inches nine lines; of the tarsus two inches six lines; from the carpus to the end of the wing seventeen inches three lines.

DESCRIPT. (*Adult in winter*.) Crown, region of the eyes, occiput, nape and sides of the neck, white, with longitudinal streaks of pale brown on the shafts of the feathers: back, scapulars and wing-coverts, fine bluish ash: primaries dusky, passing into deep black towards their extre-

mities, all terminated by a large white spot; scapulars and secondaries edged and tipped with white: forehead, throat, and all the other under parts, as well as the tail, pure white: bill ochre-yellow; the angle of the lower mandible orange-red: orbits orange: irides pale yellow: legs livid flesh-colour. (*Summer plumage.*) The whole head and neck pure white, without any brown streaks: the rest as in winter. (*Young of the year.*) Head, neck, and all the under parts, ash-gray, spotted and variegated with light brown: upper parts brown and ash-colour, all the feathers edged with reddish: quills dusky brown, without any white at the tips: tail whitish at the base, becoming browner towards the extremity, the tips of the feathers reddish yellow: bill grayish black: irides and orbits brown: legs livid brown. *In the second year*, the colours are similar, but somewhat paler. *After the second autumnal moult*, the bluish ash begins to appear upon the back mottled with brown, and the irides get lighter, inclining to yellow. *The following spring*, the colour of the bill changes; the bluish ash extends further and assumes a purer tint; and the bar on the tail gradually disappears. *At the age of three years, or after the third autumnal moult*, the plumage is matured. (*Egg.*) Light olive-brown, spotted with two shades of dark brown: long. diam. two inches six lines; trans. diam. one inch nine lines.

Common on all parts of the coast throughout the year. In the young or immature state, occasionally observed inland in the vicinity of rivers and fresh-water lakes. Breeds on rocky cliffs, constructing a nest of long dry grass. Eggs two or three in number. Feeds on fish, worms, and marine *rejectionenta*. *Obs.* The name of *Wagel Gull* has been applied indiscriminately to the young of this, and the two following species.

299. *L. fuscus*, Linn. (*Lesser Black-backed Gull.*)—Bill and legs yellow: tarsus two inches three lines: wings reaching two inches beyond the tail: shafts of the primaries black: mantle (in the *adult*) grayish black.

L. fuscus, *Temm. Man. d'Orn.* tom. 11. p. 767. Lesser Black-backed Gull, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 205. *Selb. Illust.* vol. 11. p. 509. pl. 95.

DIMENS. Entire length twenty-four inches. **MONT.**

DESCRIPT. (*Adult in winter.*) Crown, region of the eyes, occiput, nape and sides of the neck, white, with longitudinal streaks of pale brown on the shafts of the feathers: upper part of the back, scapulars and wing-coverts, deep blackish gray: primaries black, deepening in tint towards their extremities; the first two with an oval white spot near their tips; the rest finely tipped with white; secondaries, and some of the longer scapulars, likewise tipped with white: forehead, throat, and all the under parts, lower part of the back, rump and tail, pure white: bill ochre-yellow; the angle of the lower mandible orange-red: irides gamboge-yellow: orbits bright red: legs yellow. (*Summer plumage.*) The whole head and neck pure white, without any brown streaks: the rest as in winter. (*Young of the year.*) Scarcely to be distinguished from that of the last species, excepting by the shorter tarsus and somewhat longer wings: plumage extremely similar: throat, and front of the neck, whitish, with longitudinal streaks of pale brown: rest of the under parts grayish white, almost entirely covered with large deep brown spots: upper plumage dusky brown, all the feathers edged with a narrow yel-

lowish band: primaries black, without any white at the tips: tail grayish white at the base, variegated with black; the remaining portion deep blackish brown, the feathers tipped with white: bill blackish, brown at the base: legs dull yellow. (*Egg.*) Yellow stone-colour, thickly spotted with ash-gray and two shades of brown: long. diam. two inches ten lines; trans. diam. one inch eleven lines.

Not uncommon on the northern coasts of Britain: less abundant southward, though generally diffused. Like the last species, occasionally observed inland on the banks of rivers. Breeds on rocks and steep shores, and lays from two to four eggs. Food, fish, worms, and marine *rejectamenta*.

300. *L. marinus*, Linn. (*Great Black-backed Gull*).—Bill yellow: legs pale flesh-colour: tarsus three inches: wings reaching very little beyond the tail: shafts of the primaries black: mantle (in the *adult*) grayish black.

L. marinus, *Temm. Man. d'Orn.* tom. 11. p. 760. *Great Black-backed Gull*, *Mont. Orn. Dict. Selb. Illust.* vol. 11. p. 507. pl. 97. *Black-backed Gull*, *Bew. Brit. Birds*, vol. 11. p. 201.

DIMENS. Entire length near thirty inches: length of the bill three inches six lines: breadth, wings extended, five feet nine or ten inches. **MONT.**

DESCRIPT. Distinguished from the last species, which it closely resembles, by its superior size, longer tarsus, and by the colour of the legs. (*Adult in winter.*) Crown, region of the eyes, occiput and nape, white, with longitudinal streaks of pale brown on the shafts of the feathers: back, scapulars and wing-coverts, deep grayish black: forehead, throat, neck, all the under parts, rump and tail, pure white: primaries black towards their extremities, all terminated by a large white space: secondaries and scapulars tipped with white: bill thick and strong: bright yellow; the angle of the lower mandible blood-red, enclosing a dusky spot not found in the last species: irides gamboge-yellow: orbits red: legs livid white, or pale flesh-colour. (*Summer plumage.*) The whole head and neck pure white, without any streaks of brown: orbits orange: the rest as in winter. (*Young of the year.*) Similar to that of the two last species, but always larger, with the bill stronger. Head, and fore part of the neck, grayish white, with numerous brown spots largest on the neck: upper plumage dusky brown, all the feathers edged and tipped with reddish white; wing-coverts marked with transverse bars of this last colour: under parts dirty gray, with broad zig-zag streaks and spots of brown: primaries dusky, sometimes finely tipped with white: middle tail-feathers almost wholly dusky; lateral feathers black towards their extremities; all edged and tipped with whitish: bill black: irides and orbits brown: legs livid gray. *During the second year*, the colours of the plumage undergo no material change: only the gray on the head and under parts gradually gives place to a purer white, while the brown spots decrease in size and number; the base and tip of the bill also assume a livid tinge. *After the second autumnal moult*, the mantle of grayish black begins to show itself, though still variegated with large irregular brown spots; the white on the under parts is now nearly perfected: the tail white, variegated in parts with black; the bill becomes livid yellow spotted with black: and the red spot, enclosing a black one, appears at the corner of the lower

mandible. *At the age of three years, or after the third autumnal moult, the plumage is matured. (Egg.)* Yellow-brown tinged with green, sparingly spotted with slate-colour and dark brown: long. diam. three inches two lines; trans. diam. two inches four lines.

Not so plentiful as the two last species, but occasionally met with on most parts of the British coast. Observed by Montagu in considerable abundance on the extensive sandy flats of the coast of Caermarthenshire. In the immature state, it is sometimes seen far inland. Breeds in the Orkneys, and on the northern shores of Scotland. Nest placed on the shelves of insulated rocks. Eggs three or four in number. Food, fish, carrion, and any animal matter.

301. *L. Islandicus*, Edmondst. (*Iceland Gull.*)—Bill yellow: legs livid: tarsus two inches three lines: wings reaching a little beyond the tail: shafts of the primaries white: mantle (in the *adult*) pale bluish ash.

L. Islandicus, Edmondst. in *Wern. Mem.* vol. iv. p. 506. *Flem. Brit. An.* p. 139. *L. argentatus*, *Sab. in Linn. Trans.* vol. xii. p. 546. ? *Iceland Gull*, *Selb. Illust.* vol. ii. p. 501. pl. 98.

DIMENS. Entire length twenty-two inches, (EDMONDST.): length of the bill (from the forehead) two inches: of the tarsus two inches three lines, (SELB.): breadth, wings extended, four feet four inches, (EDMONDST.)

DESCRIPT. Distinguished from the next species by its smaller size, and from the *L. argentatus*, which it also closely resembles, by the entire absence of black on the primary quills. (*Adult in winter.*) "Back, and upper wing-coverts, very pale blue; all the rest of the plumage white, except the head and upper part of the neck, which are streaked with gray, as occurs in the winter dress of the other large gulls: wing-feathers and scapulars tipped with a more brilliant and pure white tinge than that which occurs on the rest of the plumage: irides pale yellow: bill smaller and more slender than in the *Herring Gull*: feet deep flesh-colour." EDMONDST. (*Immature Bird.*) "Ground-colour of the entire plumage pale yellowish gray; the feathers being barred and mottled with pale broccoli-brown: quills grayish white, with a slight tinge of broccoli-brown: tail pale broccoli-brown, marbled with white: bill pale flesh-red, or livid, at the base, with the tip blackish, or dark horn-colour: irides pale yellowish gray: legs and toes pale livid flesh-red." SELB.

A winter visitant in the Shetland Isles, and the northern parts of Scotland, where it was first observed by Mr. Edmondston, who has pointed out the distinctions between this and the next species. Mr. Selby has also obtained a few specimens on the coast of Northumberland, though all in immature plumage. By this last gentleman it is considered to be the same as the *L. argentatus* of Sabine, *l. c.* (the *L. arcticus* of Macgillivray, *Wern. Mem.* vol. v. p. 268), which is described as being plentiful in Baffin's Bay, Davis's Straits, and Melville Island. Mr. Selby observes that it is also common upon the Iceland coast, and thinks it probable that many of those which winter with us retire there to breed. Feeds on fish, the flesh of whales, and other carrion.

302. *L. glaucus*, Brunn. (*Glaucous Gull.*)—Bill yellow: legs livid: tarsus three inches: wings barely

reaching to the end of the tail: primaries terminated by a large white space; the shafts entirely white: mantle (in the *adult*) pale bluish ash.

L. glaucus, *Temm. Man. d'Orn.* tom. II. p. 757. *Sab. in Linn. Trans.* vol. XI. p. 543. Glaucous Gull, *Bew. Brit. Birds*, vol. II. pp. 209, & 212. *Selb. Illust.* vol. II. p. 498, pl. 99. Iceland Gull, *Edmondst. in Wern. Mem.* vol. IV. pp. 176, & 182.

DIMENS. Entire length twenty-nine inches: length of the bill (above) three inches, (from the gape) four inches; of the tarsus three inches; of the tail ten inches; of the wing nineteen inches: breadth, wings extended, five feet two inches. *Faun. Bor. Am.*

DESCRIPT. Similar to the last, but larger, with the bill more robust, and the tarsus longer. (*Adult in winter.*) Head and neck white, streaked and mottled with gray: back, scapulars, and wing-coverts, pale bluish ash, not so dark as in the *L. argentatus*: primaries terminated by a large white space, extending upwards more than two inches; the shafts entirely white: tips of the secondaries, tail, and all the under parts, also pure white: bill very strong; wax-yellow; the angular projection of the lower mandible bright arterial blood-red: orbits red: irides yellow: legs livid flesh-colour. (*Summer plumage.*) Head and neck pure white, without the gray streaks: the rest as in winter. (*Young of the year.*) Whole plumage mottled throughout with white, gray, and light brown, as in the young of the *L. marinus* and *L. argentatus*, but the colours are always paler than in those species, especially those of the quills, which are brown instead of dusky, with the shafts entirely white: bill horn-coloured at the base, the tip brownish black; the angular projection of the lower mandible not so strongly defined as in the adult: legs flesh-colour. From the young of the last species, they are only to be distinguished by their greater size. (*Egg.*) Stone-colour, spotted with ash-gray, and two shades of red-brown: long. diam. two inches nine lines; trans. diam. one inch eleven lines.

A regular winter visitant in the Shetland Islands, where it arrives about the middle of Autumn. In the southern parts of Britain rare, and only occasionally met with: such individuals generally young birds in immature plumage. A bold and voracious species, preying on fish and other birds, as well as on carrion. Retires to the Arctic Regions to breed.

GEN. 110. LESTRIS, *Illig.*

303. *L. Cataractes*, *Temm.* (*Common Skua.*)—Central tail-feathers but slightly elongated; of equal breadth throughout; square at the extremities: tarsus two inches eight lines.

L. Cataractes, *Temm. Man. d'Orn.* tom. II. p. 792. *Cataractes vulgaris*, *Flem. in Edinb. Phil. Journ.* vol. I. p. 99. Skua Gull, *Mont. Orn. Dict. & Supp.* *Bew. Brit. Birds*, vol. II. p. 229. Common Skua, *Selb. Illust.* vol. II. p. 514. pl. 100.

DIMENS. Entire length twenty-five inches; the same, central tail-feathers excluded, twenty-two inches: length of the bill one inch nine lines; of the tarsus two inches eight lines.

DESCRIPT. Head, and region of the eyes, deep brown: neck, and all the under parts, reddish ash, tinged with brown: back and scapulars dark ferruginous brown, the feathers edged at the sides with dusky brown: wing-coverts, secondary quills, and tail-feathers, brown: basal half of the primaries white; the remaining portion deep brown; the first with the whole of the outer web brown; shafts of the quills, as well as those of the tail-feathers, white: bill black, the base brownish: irides brown: orbits black: legs covered with large black scales; back part of the tarsus with very little indication of the projecting asperities which characterize the next species: claws strong and black. *Obs.* The plumage of this species is not subject to any important variation, either from age or season. One, which lived ten years in confinement, exhibited no tendency towards the light colour which characterizes the adult under plumage of the other species of the genus. (*Egg.*) Olive-brown, blotched with darker brown: long. diam. two inches nine lines: trans. diam. two inches.

Met with in small numbers in the Shetland Islands, particularly in those of Foulah and Unst, where they remain the whole year. Of very rare and accidental occurrence in the southern parts of Britain. Montagu mentions one that was shot at Sandwich in Kent, in the Winter of 1800. Another has been since killed in Somersetshire. A bold and rapacious species, obtaining its food principally by pursuing the larger kinds of Gulls, and compelling them to disgorge the fish which they have obtained. Flight very impetuous. Breeds in large companies on high hills and unfrequented moors. Nest constructed of a few dried weeds. Eggs two in number.

304. *L. pomarinus*, Temm. (*Pomarine Skua.*) — Central tail-feathers projecting three inches; of equal breadth throughout; rounded at the tips: tarsus two inches and half a line; covered posteriorly with rough angular scales.

L. pomarinus, Temm. *Man. d'Orn.* tom. 11. p. 793. *Faun. Bor. Amer.* part ii. p. 429. *Pomarine Skua*, Shaw, *Gen. Zool.* vol. XIII. p. 216. pl. 24. *Selb. Illust.* vol. 11. p. 517. pl. 101**.

DIMENS. Entire length twenty-one inches; the same, central tail-feathers excluded, eighteen inches: length of the bill (above) one inch seven lines, (from the gape) one inch ten lines; of the tarsus two inches and half a line; of the central tail-feathers nine inches six lines; of the wing fifteen inches. *Faun. Bor. Am.*

DESCRIPT. (*Male and female in perfect plumage.*) Face, crown of the head, occiput, back, wings, and tail, of a uniform deep brown: neck straw-yellow; the feathers on the nape long and subulate: ear-coverts, throat, breast and belly, white; vent, and under tail-coverts, blackish brown; flanks, and sides of the breast, blotched with the same: shafts of the quills and tail-feathers white, except at their tips: bill dark brown, the tip black: legs black: tail slightly rounded, excluding the central pair of feathers; these last of equal breadth throughout, and rounded at the tips: tarsus posteriorly very rugose, clothed with angular projecting scales. (*Middle age.*) "The whole plumage above and below very deep brown; the feathers on the neck and nape slightly elongated, tinged with yellow: central tail-feathers shorter than in the adult in

perfect plumage, but always of the same breadth throughout, and rounded at the tips: bill and legs the same as described above." TEMM. (*Young of the year.*) Head, neck, and all the upper parts, very dark cinereous brown, approaching to dusky; the feathers on the back, scapulars, and wing-coverts, edged with reddish yellow: quills and tail dusky: under parts yellowish ash, with dusky spots and indistinct transverse undulating bars; upper and under tail-coverts with more distinct bands of brown and yellowish white: base of the bill greenish blue; the tip black: legs bluish ash; base of the toes, and membranes, whitish: central tail-feathers not more than half an inch longer than the others; the tips broad and rounded. (*Egg.*) Pale green; the larger end spotted with two shades of red brown: long. diam. two inches three lines; trans. diam. one inch six lines and a half.

Inhabits the Arctic Regions. In this country, at least in the adult state, a rare and accidental visitant. Immature birds have been killed at Dover, Brighton, near London, and in several instances on the eastern coast of England. Food and habits similar to those of the last species. Said to breed on low hills in marshy districts, or on rocks. Constructs a nest of grass and moss coarsely put together, and lays two or three eggs.

305. *L. Richardsonii*, Swains. (*Richardson's Skua*.)
—Central tail-feathers projecting three inches; gradually tapering from the base; the tips acute: tarsus one inch nine lines; slightly rugose posteriorly.

L. Richardsonii, *Faun. Bor. Amer.* part ii. p. 433. pl. 73. *Cataractes parasiticus*, *Flem. in Edinb. Phil. Journ.* vol. 1. p. 101. *Id. Brit. An.* p. 138. *Larus parasiticus*, *Edmondst. in Edinb. Phil. Journ.* vol. v. p. 169, & vol. vii. p. 91. Arctic and Black-toed Gulls, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. ii. pp. 232, & 235. Arctic Skua, *Selb. Illust.* vol. ii. p. 520. pls. 101, & 101*. *Richardson's Lestrís*, *Gould, Europ. Birds*, part iv.

DIMENS. Entire length twenty-one inches; the same, central tail-feathers excluded, eighteen inches: length of the bill (from the forehead) one inch two lines and a half, (from the gape) one inch nine lines and a half; of the tarsus one inch nine lines: of the middle toe, nail included, one inch eight lines and a half; of the central tail-feathers nine inches; from the carpus to the end of the wing thirteen inches.

DESCRIPT. (*Adult male and female.*) Forehead, crown, occiput, wings and tail, deep black; the shafts of the quills and tail-feathers white to near their tips: rest of the upper plumage deep dusky brown tinged with cinereous: nape, and ear-coverts, straw-yellow: throat, breast, and belly, yellowish white: flanks, vent, and under tail-coverts, deep dusky gray: bill and legs black. (*Young of the year.*) Extremely similar to those of the last species, from which they are scarcely to be distinguished except by their inferior size, less robust bill, shorter and smoother tarsi, and the sharp-pointed central tail-feathers, which in length, however, do not at this age exceed the others by more than half an inch: legs, and basal portion of the webs as well as toes, flesh-colour; the remaining portion of the membranes black, in which state it is the *Black-toed Gull* of English authors. *Obs.* In the *adult state* this species is subject to considerable variation of plumage; in some instances all the under parts being of a uniform dark brownish gray without any of the yellow and

yellowish white tints, which are probably not assumed till after a certain number of years. This variety is common to both sexes, and individuals may be found shewing every intermediate shade of colouring. (*Egg*.) Olive-brown, spotted with two shades of darker brown: long. diam. two inches four lines; trans. diam. one inch eight lines.

Of much more frequent occurrence than the next species, with which it appears to have been confounded by many authors. It is, however, principally confined to the Orkney and Shetland Islands, where it breeds in considerable abundance. Immature birds are occasionally met with on different parts of the English coast. Habits predatory, like those of the Common Skua. Nest constructed of dry grass, on unfrequented heaths. Eggs two in number. Is gregarious during the breeding season.

306. *L. parasiticus*, Temm. (*Arctic Skua*).—Central tail-feathers projecting six inches or more; gradually tapering from the base; the tips acute: tarsus one inch six lines; slightly rugose.

L. parasiticus, Temm. *Man. d'Orn.* tom. II. p. 796. *Faun. Bor. Amer.* vol. II. p. 430. Parasitic Gull, Gould, *Europ. Birds*, part iv.

DIMENS. Entire length twenty-one inches: length of the bill one inch and six-eighths; of the tarsus one inch six lines; of the central tail-feathers twelve inches; of the wing eleven inches nine lines. GOULD*.

DESCRIPT. Extremely similar to the last species, from which it is principally distinguished by its smaller size, longer central tail-feathers, and shorter tarsi. (*Adult male and female*.) "Top of the head, and the space between the bill and the eyes, of a deep blackish brown, terminating at the occiput; the whole of the upper surface of a clear brownish gray; quills and tail-feathers much darker; the throat, neck, and under surface, of a pure white, with the exception of the cheeks and sides of the neck, which are tinged with a delicate straw-yellow; legs and feet black. The *young* resemble in colouring the other species of the genus of the same age." GOULD.

Apparently of rare occurrence in this country; nor is it known whether it ever breeds with us like the last species. A British-killed specimen is in the British Museum, but I am ignorant from what locality it was obtained. According to Mr. Thompson, it has repeatedly occurred in the Bays of Dublin and Belfast. It is possible that the *Arctic Gull* noticed by Montagu in the *Supplement* to his "Ornithological Dictionary," may have been of this species, but in all other cases the bird described under that name by English authors is referable to the *L. Richardsonii*. According to Mr. Gould, the present species inhabits the shores of the Baltic, the rugged coasts of Norway, and the Polar Regions. Habits similar to those of the rest of the genus.

* The following are the dimensions of a *British specimen*, in the British Museum, as given in the *Faun. Bor. Amer.* by Dr. Richardson. "Length, total, twenty inches; the same, excluding central tail-feathers, fourteen inches; of central pair eleven inches; of wing eleven inches six lines; of bill above one inch; of bill to rectus one inch seven lines and a half; of tarsus one inch seven lines; of middle toe one inch three lines; of hind toe two lines; of hind nail two lines; transverse diameter of the bill at the front four lines."

GEN. 111. PROCELLARIA, Linn.

(1. PROCELLARIA, Vig.)

307. *P. glacialis*, Linn. (*Northern Fulmar*.)—White; back and wings hoary ash: bill and legs yellow.

P. glacialis, *Temm. Man. d'Orn.* tom. 11. p. 802. *Fulmar*, *Mont. Orn. Dict. & Supp. Bew. Brit. Birds*, vol. 11. p. 239. *Fulmar Petrel*, *Selb. Illust.* vol. 11. p. 525. pl. 102.

DIMENS. Entire length seventeen inches: length of the bill (from the forehead) one inch seven lines, (from the gape) one inch nine lines and a half; of the tarsus one inch ten lines; of the tail five inches; from the carpus to the end of the wing twelve inches.

DESCRIPT. (*Adult.*) Head, neck, rump, tail, and all the under parts, pure white: back, scapulars, wing-coverts, and secondary quills, bluish ash; primaries dusky gray: tail rounded at the tip: bill yellow; the nasal tube tinged with orange: irides and legs yellowish. (*Young of the year.*) "All the parts of the body pale gray tinged with brown; the feathers on the back and wings tipped with deeper brown: primary quills and tail-feathers of one uniform tint of brownish gray: before the eyes an angular dusky spot: bill and legs yellowish ash." *TEM.* (*Egg.*) Dirty yellowish white: long. diam. two inches ten lines; trans. diam. two inches.

Not uncommon in some of the islands off the North of Scotland, but very rarely seen in the southern parts of Britain. Has been killed on the coast of Durham, as well as, in a single instance, in South Wales. Breeds in St. Kilda, and is said to lay one egg in the holes of rocks. Habits very oceanic: seldom resorts to land, except for the purpose of nidification. Food, fish, the flesh of whales, and other animal matter.

(2. PUFFINUS, Ray.)

308. *P. Puffinus*, Linn. (*Cinereous Shearwater*.)—Bill two inches long: tarsus one inch ten lines: tail conical. *TEM.*

P. Puffinus, *Temm. Man. d'Orn.* tom. 11. p. 805. *Cinereous Shearwater*, *Shaw, Gen. Zool.* vol. XIII. part i. p. 227. *Selb. Illust.* vol. 11. p. 528. pl. 102*.

DIMENS. Length eighteen inches. *TEM.*

DESCRIPT. "Head, back part of the neck, and the upper plumage, blackish brown, with the margins and tips of the feathers of the scapulars lighter: throat, lower part of the neck, and the whole of the under plumage, deep ash-gray, with a tinge of broccoli-brown: quills and tail brownish black: legs having the outer part of the tarsus deep gray; the inner part, and webs, yellowish." *SELB.*

Mr. Selby informs us that he has lately obtained an individual of this species on the coast of Northumberland. It has not been noticed by any other British author. Possibly, however, it may have been confounded with the *P. Anglorum*, which it is said closely to resemble. According to Temminck, it is common in the Mediterranean and on the coast of Spain. Habits unknown.

309. *P. Anglorum*, Temm. (*Manks Shearwater*).— Bill very slender; one inch seven or eight lines in length; tail rounded; wings extending a little beyond its extremity; tarsus one inch nine lines.

P. Anglorum, Temm. *Man. d'Orn.* tom. 11. p. 806. Shearwater, *Mont. Orn. Dict.* Bew. *Brit. Birds*, vol. 11. p. 241. Manks Shearwater, *Selb. Illust.* vol. 11. p. 529. pl. 102.

DIMENS. Entire length fourteen inches: length of the bill (from the forehead) one inch six lines, (from the gape) one inch nine lines; of the tarsus one inch nine lines; of the tail three inches seven lines; from the carpus to the end of the wing nine inches six lines.

DESCRIPT. Forehead, crown, nape, and all the upper parts, including the wings and tail, deep grayish black, with a slight gloss: sides of the neck mottled with indistinct transverse bars of gray and white: chin, throat, and all the other under parts, pure white: some of the under tail-coverts black on their outer webs: bill yellowish brown at the base, dusky towards the tip: legs gray; the front of the tarsus, and part of the webs, yellowish white, tinged with flesh-red. (*Egg*.) White: long. diam. two inches five lines; trans. diam. one inch nine lines.

Formerly very abundant during the breeding season in the Calf of Man, but less so of late years. Common in the Orkneys, and some other of the Scotch Islands, to which they resort in February or March. Builds in the interstices of rocks, sometimes in rabbit-burrows, laying but one egg. Food, fish, and marine insects and worms. Habits crepuscular; seldom appearing abroad during the bright sunshine.

310. *P. fuliginosa*, Strickland.

Puffinus fuliginosus, *Proceed. of Zool. Soc.* 1832. p. 129.

DIMENS. Entire length eighteen inches: length of the bill (from the forehead) one inch and seven-eighths, (from the gape) two inches and a half; of the tarsus two inches and a quarter; of the middle toe two inches and a half; of the wing twelve inches. STRICKL.

DESCRIPT. Brown above and below; the wings of a deeper tint: throat faintly tinged with gray: bill of the same colour: tarsi externally, and the outer toes, brown: tarsi internally, and membranes, ochraceous brown. STRICKL.

A single individual of this species, supposed to be the *Procellaria fuliginosa* of Kuhl, was shot in the middle of August 1828, at the mouth of the Tees, and is now in the collection of Mr. Arthur Strickland, of Boynton in Yorkshire. It is the only specimen hitherto obtained. Native country and habits unknown.

(3. THALASSIDROMA, Vig.)

311. *P. pelagica*, Linn. (*Stormy Petrel*).— Black: tail even; the wings extending a little beyond its tip: tarsus ten lines and a half.

P. pelagica, Temm. *Man. d'Orn.* tom. 11. p. 810. Stormy Petrel, *Mont. Orn. Dict. & Supp.* Bew. *Brit. Birds*, vol. 11. p. 246. *Selb. Illust.* vol. 11. p. 533. pl. 103. f. 2.

DIMENS. Entire length five inches ten lines.

DESCRIPT. Head, back, quills, and tail, ink-black: wing-coverts brownish black; a broad transverse bar of white on the rump; some of the scapulars and secondary quills tipped with white: all the under parts brownish black: bill and legs black: irides brown. (*Egg.*) White; of a roundish-oval form: long. diam. one inch one line; trans. diam. ten lines.

A common species on some parts of the British coast, especially northwards. Very abundant in the Orkneys and Hebrides, where it breeds in holes in the ground or in the clefts of rocks, laying one, sometimes two eggs. Generally keeps far out at sea, though occasionally driven inland by storms. Individuals under such circumstances, have been met with in various parts of England. Skims the surface of the ocean, feeding on the marine insects which float on the waves. When approached, ejects from its mouth an oily substance of a very rancid smell.

312. *P. Leachii*, Temm. (*Leach's Petrel.*) — Black: tail forked; the wings not reaching beyond its tip: tarsus eleven lines and a half.

P. Leachii, Temm. *Man. d'Orn.* tom. II. p. 812. Leach's Petrel, *Steph. in Shaw's Gen. Zool.* vol. XIII. part I. p. 219. pl. 25. Fork-tailed Petrel, *Bew. Brit. Birds*, vol. II. p. 244. *Selb. Illust.* vol. II. p. 537. pl. 103. f. 1.

DIMENS. Entire length seven inches six lines.

DESCRIPT. Head, neck, and under parts, grayish black: back and scapulars pitch-black: quills black; the coverts lighter, forming a bar of dusky brown: upper and lateral under tail-coverts white, the shafts and tips black: tail black, of twelve feathers, forked; the depth of the fork half an inch: bill and legs black. (*Egg.*) White; roundish-oval: long. diam. one inch four lines; trans. diam. eleven lines.

First discovered by Mr. Bullock in St. Kilda. Since ascertained to be not uncommon in that island, though rare elsewhere. Solitary individuals have been occasionally met with in different parts of England. Breeds in the clefts of rocks, and lays one egg. Food and habits similar to those of the last species.

(26.) *P. Wilsoni*, Bonap. — Vigors in *Zool. Journ.* vol. I. p. 425. *P. pelagica*, Wils. *Amer. Orn.* vol. VII. p. 90. pl. 60. f. 6.

I am informed by Mr. Yarrell that this species has been killed in the British Channel, though at some distance from land. It inhabits the western shores of the Atlantic, and is principally distinguished by a large oblong yellow spot on the membranes of the toes. Length of the tarsus nearly an inch and a half.