580.5

FB

v.2

1900-1916

cop.2

Inc.

FIELD COLUMBIAN MUSEUM.

PUBLICATION 106.

BOTANICAL SERIES.

VOL. II, No. 3.

PRÆNUNCIÆ BAHAMENSES-I.

CONTRIBUTIONS TO A FLORA OF THE
BAHAMIAN ARCHIPELAGO.

BY

CHARLES FREDERICK MILLSPAUGH, M.D.

Curator Department of Botany.

CHICAGO, U. S. A.

February, 1906.

BOTANICAL SERIES.

ACTUAL DATES OF ISSUE.

<i>Publication Number.</i>	<i>Bot. Ser.</i>	<i>Title</i>	<i>Date of Issue</i>
4	Vol. I No. 1	Flora of Yucatan.	Sept. 20, 1895
9	Vol. I No. 2	Flora of West Virginia.	Mar. 21, 1896
15	Vol. I No. 3	Contrib. Fl. Yucatan II.	Mar. 1, 1897
25	Vol. I No. 4	Contrib. Fl. Yucatan III.	Mar. 7, 1898
39	Vol. I No. 5	Higinbothamia, etc.	Sept. 9, 1899
43	Vol. II No. 1	Plantae Utowanæ.	Apr. 20, 1900
48	Vol. I No. 6	Pl. Ins. Ananas.	Sept. 7, 1900
50	Vol. II No. 2	Pl. Uto. Suppl.	Sept. 7, 1900
68	Vol. I No. 7	Fl. St. Croix.	Dec. 16, 1902
69	Vol. III No. 1	Pl. Yuc. Fasc. I.	Apr. 6, 1903
92	Vol. III No. 2	Pl. Yuc. Fasc. II.	Apr. 26, 1904

Volume I Is closed and indexed.

Volume II Is still open. Reserved for papers on Antillean Botany.

Volume III Is still open. It is reserved for the sequence of fascicles of Plantae Yucatanæ until that Flora is completed.

FIELD COLUMBIAN MUSEUM.

PUBLICATION 106.

BOTANICAL SERIES.

VOL. II, No. 3.

PRÆNUNCIÆ BAHAMENSES-I.
CONTRIBUTIONS TO A FLORA OF THE
BAHAMIAN ARCHIPELAGO.

BY

CHARLES FREDERICK MILLSPAUGH, M.D.

Curator Department of Botany.

CHICAGO, U. S. A.

February, 1906.

ISSUED Feb. 17, 1906

PRÆNUNCIÆ BAHAMENSES—I.

CONTRIBUTIONS TO A FLORA OF THE BAHAMIAN ARCHIPELAGO.

CHARLES F. MILLSPAUGH, M.D.

During field and herbarium studies of the flora of the Greater Antillean Islands, conducted by Dr. N. L. Britton, Director of the New York Botanical Garden, and the Author, each came to the conclusion that a thorough investigation of the Bahamian Archipelago would throw much desired light upon many problems of plant distribution and mutation, and offer a probable solution of the question of the age of these interesting islands.

A joint investigation of such collections as had been previously made on the archipelago proved them to be in many cases meagre in material, scrappy in character, or from islands too widely separated to afford even a partial solution of the problems presented. They therefore decided to solicit the consent of those botanists representing the Government of Great Britain, to which these islands belong, and the support of the institutions they themselves represent, in view of organizing a thorough exploration of the islands included in the archipelago. This being in both instances unhesitatingly given, three expeditions were equipped for immediate field work and others planned, to cover within the next two years, all the islands included.

Up to the present writing, the survey thus organized has covered the following groups:—Great Bahama, the Abacos, the Isaacs, the Berry Islands, the Bimini Islands, the New Providence area, the Exuma Chain, the Inaguas and Grand Turk. In these, a large amount of information and material has been accumulated by Dr. and Mrs. Britton, Mr. Nash, Mr. Brace, and the Author. In accomplishing this, not only have the usual lines of communication been utilized, but special vessels have been equipped and commissioned.

The following material (over 7,000 sheets), resulting from this survey, or secured by exchange, gift, loan, or purchase, is deposited in duplicate in both institutions, where it is being studied by the co-workers, who are also taking occasion at various intervals to consult the collections of Catesby, Eggers, Hjalmarsson, Brace, Daniell, Herrick, Allen, Barbour & Bryant, and others, deposited in European and American herbaria. In addition, a complete series of the orchids, cacti, tillandsias, and other difficult groups is being collected alive and studied in the conservatories of the Garden.

COLLECTORS.

Brace — Mr. Louis J. K. Brace began his collections in the Bahamas in 1875. He worked principally on New Providence until 1880, but also secured material on Fortune Island or Long Cay, and Andros. His work resulted in about 525 numbers, which were sent to Kew Gardens. A duplicate set is still in his possession. He began field work under the present survey in September, 1904 (see Britton & Brace), and later in the same year was commissioned by the Garden to make an exploration of the Abaco Islands, where he collected as follows: Green Turtle Cay (1475-1515), Allen's Cay (1526-54), Spanish Cay (1555-62), Man-o'-War Cay (1563-89), Pigeon Cay (1662-73), Elbow Cay (1688-1712), and on the island of Abaco at Butler Bay (1516-25), Marsh Harbor (1590-1661, 1713-53, 1787-1846), Great Cistern (1674-87, 1754-86), Cherokee Sound (1847-9, 1898-1980, 1999, 2080-84, 2086), Eight Mile Bay (1850-97), opposite Cherokee Settlement (1981-98, 2064-79), Old Kerr's Point (2000-2036), and California road (2037-63), a total of 611 numbers. Following this expedition he continued work on New Providence in localities from which, from time to time, it became necessary to have more material of previously secured species.

In June, 1905, a commission was given him by both the Garden and this Museum to continue, on Great Bahama, the work already undertaken there. (See Britton & Millspaugh.) On this exploration he collected at Eight Mile Rocks (3633-5, 3671-3738), Deadman's Reef (3631-2), and the western extremity of the island (3484-3630, 3636-57). He also collected Garden Cay (3658-70), and visited (during his sailings to and from Great Bahama) North Bimini (3461-75, 3479-83), South Bimini (3476-8), North Cat Cay (3739-54), and Andros Island (3755-6), and after returning to New Providence collected Nos. 3893-3918.

Britton — Dr. N. L. Britton collected on New Providence in April, 1904, in the neighborhood of Nassau, South Side Beach, Blue Hills, Far-

ringdon Road, West Bay Street Road, Cunningham Road, Lake Cunningham, and Old Fort. In this preliminary investigation, in company with the Author (see Millsp.), he collected 158 numbers, (1-158). An account of the field work is published in the Journal of the New York Botanical Garden 5: 129-136.

See also Britton & Brace, and Britton & Millspaugh.

B. & Br. — Britton & Brace — Dr. N. L. Britton, Mrs. E. G. Britton, and Mr. Louis J. K. Brace collected together on New Providence (Nos. 159-328, 361-872), and Hog Island (329-360), in September and October, 1904. They ranged New Providence with great thoroughness, penetrating to the coast region north, east, south, and west, and to the coppices, pine barrens, and openings in the neighborhood of Nassau, Fort Montague, Farringdon Road, Swingate, Maidenhead, Tea House, Race Course, Delaport, Fox Hills, Village Road, Mt. Vernon, Rifle Range, Harold Road, Carmichael, Adelaide, Southwest Landing, Miller's, Blue Hills, Soldiers Road, Lake Cunningham, Winton, Old Fort, Waterloo, Clifton, Mt. Pleasant, Fort Charlotte, Gambier, Killarney, Bonefish Pond, and Seven Hills. The commoner things were not taken, the attempt being made to secure only such species as were previously unfamiliar or those which had only been gathered when out of character. This survey resulted in 714 particularly interesting plants embracing the series of numbers indicated above.

An account of this expedition is published in the Journal of the New York Botanical Garden 5: 201-209.

B. & M. — Britton & Millspaugh — Drs. N. L. Britton and C. F. Millspaugh, accompanied by Mr. M. A. Howe, began an exploration of the Berry Islands, Great Bahama, and the Exuma Chain, in January, 1905. A schooner was chartered and equipped at Nassau, where, after a day's delay on account of heavy weather (during which collecting was done in the scrublands and coppices south of Nassau (2085-2110), the expedition began at Rose Island (2111-2166), and Hog Island (2167-2176). The first island of the Berry Group visited was Whale Cay (2177-2201), followed by Frozen Cay (2202-2224), Little Harbor Cay (2225-2254), Great Sturup Cay (2255-2285), Goat Cay (2286-2301), Lignum Vitæ Cay (2302-2332), and Great Harbor Cay (2333-2354). The exploration began on Great Bahama Island at Eight Mile Rocks (2355-2596), and was continued at Barnett's Point (2597-2716), and Golden Grove (2717-2741). Returning to Nassau, a fresh start was made to the Exuma Chain, of which the following islands were diligently searched: Ship Channel

Cay (2742-2767), Cay north of Wide Opening (2768-2804), Cave Cay (2805-2836), Little Galiot (2837-2850), Great Galiot (2851-2860), Great Guana (2861-2922), Great Exuma (2923-3051, 3073-3141), and Stocking Island (3052-3072).

On Great Exuma the regions worked were the scrub lands and coppices lying west of Georgetown; Hayne's Road from the east to the west shores across the island; and the Rolletown scrub lands and coppices. The total series of this exploration is 1057 sheets.

An account of this expedition may be found in the *Journal of the New York Botanical Garden* 6: 77-85.

Coker — Dr. William C. Coker, botanist of the Bahamian expedition of the Geographical Society of Baltimore, assisted by Messrs. C. A. Shore and F. M. Hanes, collected in the summer of 1903 on the following islands: New Providence (1-193, 247-305, 551), Andros Island and Little and Great Mangrove Cays (194-234), Green Cay (235-246), George's Island (306-317), Eleuthera (318-421), Cat Island (422-437), Rum Cay (438-457), Watlings Island (458-491, 521, 528), Long Island (492-520, 524), Water Cay (523, 525), Abaco (558-568, 575-6), and Elbow Cay (569-574). This collection was deposited in the herbarium of the New York Botanical Garden, and forms the basis of his "Vegetation of the Bahama Islands," in Shattuck's "The Bahama Islands." Owing to the insufficiency of the material secured many phanerogams therein published are based upon provisional determination only. The cryptogams exist under a separate series of numbers.

Cooper — Dr. William Cooper paid a visit to New Providence in 1859 for the purpose of making dredgings for zoölogical material. While there he collected in the neighborhood of Nassau about 100 plants for his friend, Dr. John Torrey. The prime set of these plants is now in the herbarium of the New York Botanical Garden. See *Bull. Torrey Club* 17:190.

Curtiss — Mr. A. H. Curtiss collected his first series of "West Indian Plants" in April, 1903, in the neighborhood of Nassau, New Providence. This series comprises numbers 1-211, of which the prime set, with the unnumbered unicates, is deposited in the herbarium of the New York Botanical Garden, and the first distributed set in the herbarium of this Museum.

Earle — Prof. F. S. Earle collected during one day only, on New Providence (Nos. 1-79), while on his way to Cuba, in March, 1903. His plants, largely from the vicinity of Nassau, are in the herbarium of the Garden.

E. G. B. — Mrs. N. L. Britton, often accompanied by Mr. Brace, visited, in April and May, 1905, all those stations on New Providence previously collected by Britton & Brace, at which it had become desirable to secure additional material of interesting plants or characters missing on previous notable species. In addition to achieving marked success in this undertaking, she secured many other plants of particular interest. Her survey resulted in 310 sheets (Nos. 3141-3460), and is mentioned in the Journal of the New York Botanical Garden 5:129-136. See also Britton & Brace.

Hitchcock — Dr. J. T. Rothrock and Prof. Albert S. Hitchcock collected in the Bahamas in November and December, 1890. The collections were largely made by Prof. Hitchcock on New Providence, Eleuthera, Cat, Watlings, Crooked, Fortune, and Inagua, and form the basis of his "Plants collected in the Bahamas," etc. The material collected was deposited in the herbarium of the Missouri Botanical Garden, from which, through the kindness of the director, Dr. William Trelease, the author has been allowed to reassemble the sheets. The collection, embracing about 600 sheets, is at this writing a loan to the Field Columbian Museum.

Howe — Dr. Marshall A. Howe, algologist of the New York Botanical Garden, has collected extensively the sea vegetation of the Bahamas, accompanying Dr. Britton and the author on their first trip, and later, Britton & Millspaugh. He also accompanied the author upon the Bimini explorations. See Journal New York Botanical Garden 5:129-136 and 6:77-85.

Millsp. — C. F. Millspaugh—the author began his collections among the islands on New Providence, in April, 1904, in company with Dr. Britton (see Britton), visiting substantially the same stations, where, as his collecting was to extend to other islands, each kept his individual series of numerals. He collected at the following localities: Blue Hills (2048-2100), South Shore (2101-2151 and 2262-2277), West Bay Street road (2152-2178), pine barrens and hammock lands along Lake Cunningham road (2179-2217), region of Lake Cunningham (2218-2244), Farrington Road (2245-7), Swingate (2248-51), Old Fort (2252-4), and Waterloo (2255-61). Then hiring and commissioning a sloop, he sailed in company with Dr. M. A. Howe (see Howe) for an investigation of the Bimini Islands, collecting at the following stations: West End Bight, New Providence (2278-2293), Joulter's Cays (2294-2304), Gun Cay (2305-2328), North Cat Cay (2329-2347, 2415), South Bimini (2348-2368, 2379-85, 2387-98,

2406-2414), North Bimini (2369-78, 2386, 2399-2405), South Cat Cay, (2416-2441).^{*} Again in January, 1905, he collected in the following localities while awaiting the arrival of Dr. Britton (see Britton & Millspaugh): Blue Hills road (2442-4, 2461-81), Silver Cay (2445-2460), and South Side, Soldiers Road, Fort Montague and Fort Fincastle (2482-2502). Following immediately upon this, see Britton & Millspaugh.

N. & T. — Nash & Taylor—Messrs. Geo. V. Nash and Norman Taylor, of the New York Botanical Garden, were commissioned by the Garden, in October, 1904, to investigate the Inaguas. They made an exhaustive survey of the islands, collecting on Inagua 482 numbers (874-1138 and 1258-1474), on Sheep Cay 23 numbers (1139-1161), and on Little Inagua 96 numbers (1162-1257). In September, 1905, on their return from an exploration in Haiti, they spent a week on Grand Turk Island, collecting all such plants as were noted by them at that season, 136 numbers (3757-3892.) Accounts of these expeditions may be found in the Journal of the New York Botanical Garden 6:1-19 and 189-191.

Northrop — Mrs. Alice (Rich) Northrop, accompanied by her husband, Prof. John I. Northrop, made, in 1890, a large collection of the plants of New Providence (Nos. 1-239, 280-331), Rose Island (251), Salt Cay (240-244, 271-279), Hog Island (245-8, 252-70), and Andros (332-758). These plants formed the basis of Mrs. Northrop's "Flora of New Providence and Andros." They are now deposited in equal sets in the herbaria of this Museum and the Garden.

Wight — Mr. Alex. E. Wight, under the patronage of the Gray Herbarium, of Cambridge, Mass., made a Bahamian collection of 275 numbers from March to May, 1905. He collected on New Providence and Hog Island (Nos. 1 226 and 271-275), and on Andros (227-270). The resulting plants are being determined at the Gray Herbarium, where the initial set will be retained and the duplicate sets distributed to this Museum, the New York Botanical Garden, and others. Prof. B. L. Robinson has kindly communicated, in advance, to the author all those numbers falling within the natural orders treated in this Prænuncia.

^{*}An account of this expedition may be found in the Journal of the New York Botanical Garden 5:129-136.

Islands Represented in the Collections.

- Abaco — *Brace, Coker.*
Allen's Cay (Abaco) — *Brace.*
Andros — *Brace, Coker, Northrop, Wight.*
Bimini, North — *Brace, Howe, Millspaugh.*
Bimini, South — *Brace, Howe, Millspaugh.*
Cat Cay, North (Biminis) — *Brace, Howe, Millspaugh.*
Cat Cay, South (Biminis) — *Howe, Millspaugh.*
Cat Island — *Hitchcock.*
Cave Cay (Exuma Chain) — *Britton & Millspaugh, Howe.*
Crooked Island — *Hitchcock.*
Eleuthera — *Coker, Hitchcock.*
Elbow Cay (Abaco) — *Brace, Coker.*
Exuma — *Britton & Millspaugh, Howe.*
Fortune Island — *Brace, Hitchcock.*
Frozen Cay (Berry Is.) — *Britton & Millspaugh, Howe.*
Galiot Cay, Great (Exuma Chain) — *Britton & Millspaugh, Howe.*
Galiot Cay, Little (Exuma Chain) — *Britton & Millspaugh, Howe.*
Garden Cay (Gt. Bahama) — *Brace.*
George's Isl. (Eleuthera) — *Coker.*
Goat Cay (Berry Is.) — *Britton & Millspaugh.*
Great Bahama Island — *Brace, Britton & Millspaugh, Howe.*
Green Cay — *Coker.*
Green Turtle Cay (Abaco) — *Brace.*
Guana, Great (Exuma Chain) — *Britton & Millspaugh, Howe.*
Gun Cay (Biminis) — *Howe, Millspaugh.*
Harbor Cay, Great (Berry Isl.) — *Britton & Millspaugh, Howe.*
Harbor Cay, Little (Berry Isl.) — *Britton & Millspaugh, Howe.*
Hog Island (N. Providence) — *Brace, Britton, N. L., Britton, E. G.,
Britton & Brace, Britton & Millspaugh, Northrop, Wight.*
Inagua — *Hitchcock, Nash & Taylor.*
Inagua, Little — *Nash & Taylor.*
Isaac, Great — *Brace.*
Joulter's Cays (Andros) — *Howe, Millspaugh.*
Lignum Vitæ Cay (Berry Is.) — *Britton & Millspaugh, Howe.*
Long Cay see Fortune Island.
Long Island — *Coker.*
Mangrove Cay, Little (Andros) — *Coker.*
Mangrove Cay (Andros) — *Coker.*

- Man o'War Cay (Abaco) — *Brace*.
 New Providence — *Brace, Britton, N. L., Britton, E. G., Britton & Brace, Britton & Millspaugh, Coker, Cooper, Curtiss, Earle, Hitchcock, Howe, Millspaugh, Northrop, Wight*.
 No Harbor Cay (Rose Isl.) — *Britton & Millspaugh*.
 Pigeon Cay (Abaco) — *Brace*.
 Rose Island (N. Providence) — *Britton & Millspaugh, Howe, Northrop*.
 Rum Cay — *Coker*.
 Salt Cay (N. Providence) — *Northrop*.
 Sheep Cay (Inagua) — *Nash & Taylor*.
 Ship Channel Cay (Exuma Chain) — *Britton & Millspaugh, Howe*.
 Silver Cay (N. Providence) — *Howe, Millspaugh*.
 Spanish Cay (Abaco) — *Brace*.
 Stocking Island (Exuma Chain) — *Britton & Millspaugh, Howe*.
 Sturup Cay, Great (Berry Isl.) — *Britton & Millspaugh, Howe*.
 Turk Island, Grand — *Nash & Taylor*.
 Water Cay (Cay Sal Bank) — *Coker*.
 Watlings Island — *Coker, Hitchcock*.
 Whale Cay (Berry Islands) — *Britton & Millspaugh, Howe*.
 Wide Opening, Cay N. of (Exuma Chain) — *Britton & Millspaugh, Howe*.
-

Critical investigation of the above collections has so far resulted in the following publications:

- "A New *Waltheria* from the Bahamas" N. L. Britton in *Torrey* 3: 105 (July 25, 1903).
 "Notes on Bahaman Algæ" Marshall A. Howe in *Bull. Torr. Club*, 31: 93-100 (Feb. 11, 1904).
 "*Savia Bahamensis*" N. L. Britton in *Torrey* 4: 104 (Nov. 21, 1904).
 "A new Bahaman *Euphorbia*" C. F. Millspaugh in *Torrey* 4: 172 (Nov. 21, 1904).
 "Notes on the Flora of the Bahamas" N. L. Britton before the Torrey Botanical Club. Digest in *Torrey* 4: 190 (Dec. 30, 1904).
 "On *Pisonia obtusata* and its Allies" N. L. Britton in *Bull. Torr. Club* 31: 611-615 (Jan. 9, 1905).
 "Contributions to the Flora of the Bahama Islands - I" N. L. Britton in *Bull. New York Bot. Gard.*, 3: 441-453 (Feb. 7, 1905).
 "The Polyporaceæ of North America - X." W. A. Murrill in *Bull. Torr. Club* 32: 91. *Agaricus deplanatus* (Fr.) Murr. (March 22, 1905).

- "Phycological Studies - I" Marshall A. Howe in Bull. Torr. Club 32: 241-252 (May 6, 1905).
- "Contributions to the Flora of the Bahama Islands - II" N. L. Britton in Bull. New York Bot. Gard. 4: 115-127 (Aug. 24, 1905).
- "Phycological Studies - II" Marshall A. Howe in Bull. Torr. Club 31: 563-586 (Dec. 6, 1905).

AMARANTHACEÆ, EUPHORBIACEÆ, RUBIACEÆ,
VERBENACEÆ, AND SOLANUM DIDYMACANTHUM.

AMARANTHACEÆ.

Amaranthus crassipes Schlecht.

Waste grounds infrequent:

New Providence — Fort Charlotte, *B. & Br.* 778.

Amaranthus emarginatus Salzm.

In cultivated grounds:

New Providence — Nassau, *B. & Br.* 794.

Not before known from the Bahamas. There appears to be no doubt as to the validity of this species. It comes up from Brazil through Guadeloupe (Duss 4067) to the Bahamas.

Amaranthus hybridus Linn.

Waste grounds about dwellings:

Fortune Island — *Hitchcock (A paniculatus)*.

Amaranthus polygonoides Linn.

Sandy waste places:

Eleuthera — Governor's Harbor, *Hitchcock*.

Inagua — *N. & T.* 1079.

Amaranthus spinosus Linn.

Waste grounds:

New Providence — Nassau, *Hitchcock; Brace* 204.

Amaranthus tristis Linn.

Waste grounds near habitations:

Abaco — Cherokee Sound, *Brace* 1922.

New Providence — Grants Town, *B. & Br.* 792; Soldiers Road, *Coker* 264.

Great Guana Cay — *B. & M.* 2910.

Not before known from the Bahamas.

Amaranthus viridis Linn.

Waste grounds near dwellings:

Great Bahama — Eight Mile Rocks Settlement, *B. & M.* 2429.

New Providence — Nassau, *B. & Br.* 785; *Brace* 9, 206; *Curtiss sine num.*

Not before known from the Bahamas.

Achyranthes indica (L.) Mill.

A. aspera obtusifolia (Lam) Griseb. A weed in waste places and cultivated soils:

Abaco — Cherokee Sound, *Brace* 1921.

New Providence — Nassau, *Earle* 23; *B. & M.* 2086; *Hitchcock*.

Exuma — Georgetown, *B. & M.* 2991.

Alternanthera maritima St. Hil.

In maritime sands on the upper beach line, often within wash of the waves:

Great Bahama — Barnett's Point, *B. & M.* 2620; near Dead Man's Reef, *Brace* 3632.

Frozen Cay — *B. & M.* 2205.

South Bimini — *Millsp.* 2353, 2410.

New Providence — Southwest Bay, *B. & Br.* 478.

Alternanthera paronychioides St. Hil.

In sand, especially where trodden down:

Abaco — Cherokee Sound, *Brace* 1925.

Great Bahama — Eight Mile Rocks, *B. & M.* 2367, 2412, 2419; Road to Dead Man's Reef, *Brace* 3623.

New Providence — near Nassau, *Hitchcock* (*A repens*); *Northrop* 197; *B. & Br.* 263; *Curtiss* 161; *Wight* 121.

St. George's Cay — *Coker* 313.

Inagua — *N. & T.* 979.

Lithophila muscoides Sw.

Usually in trodden sands of roadside paths:

Great Bahama — Eight Mile Rocks, *B. & M.* 2573.

New Providence — interior of Old Fort, *Millsp.* 2253; Nassau, *Curtiss* 69; *Northrop* 150.

Inagua — *Hitchcock* (*Alternanthera muscoides*); *N. & T.* 892.

Grand Turk Island — *N. & T.* 3786.

Lithophila vermicularis (L.) Uline.

On maritime rocks and beach dunes:

Abaco — Butler Bay, *Brace* 1514.

Great Bahama — Barnett's Point, *B. & M.*

Great Sturru Cay — *B. & M.*

Frozen Cay — *B. & M.*

South Bimini — *Mills* sp. 2409.

Little Mangrove Cay — *Coker* 201.

Rose Island — *B. & M.*

New Providence — Southwest Bay, *B. & Br.* 459; Swingate, *Mills* sp. 2251;

Nassau neighborhood, *Coker* 37; *Cooper sine num.*; *Northrop* 147;

Wight 10.

Eleuthera — Governor's Harbor, *Hitchcock* (*Philoxerus vermicularis*).

Exuma — rocks of West Beach, *B. & M.*

Watlings Island — *Coker* 476.

***Iresine keyensis* sp. nov.**

Caule lignoso erecto ramoso glabro, foliis petiolatis anguste lanceolatis vel oblongo-lanceolatis rotundo-obtusis submucronatis basi ad petiolam utrobique decurrentibus, margine integris subrevolutis pagina utrinque minute pustulosis, paniculis elongatis tenuiter ramosis laxis subaphyllis, rachi glabris, floribus albis paleaceis, calyce bracteis laterales medio superante, sepalis obtusiusculis enerviis glabris. Semen obeso-lenticularis.

Plants strictly erect 4.5–6.0 dm. Leaves 4 cm. x 6 mm., 9 x 1.5 cm. to 6 x 2 cm. petiole about one-quarter the length of the blade, panicles 1–2 dm. the ultimate branchlets often up to 4 cm. long, very delicate, almost capillary. Seeds dark-red, smooth and polished .5 x .5 mm.

On maritime rocks:

Great Bahama — Eight Mile Rocks, *B. & M.* 2485.

Great Sturru Cay — *B. & M.*

Little Harbor Cay — *B. & M.*

Lignum Vitae Cay — where it forms a strong character plant, *B. & M.*

Goat Cay — *B. & M.* 2289.

Frozen Cay — *B. & M.*

Whale Cay — *B. & M.* 2179 type.

South Bimini — *Mills* sp. 2390.

South Cat Cay — *Mills* sp. 2437.

Andros — Calabash Cay, Stafford Creek, *Northrop* 362a (*I. paniculata*).

Here called "NEWBURN WEED."

Mangrove Cay — *Bryant* 11.

Silver Cay — *Mills* sp. 2450.

New Providence — Southwest Bay, *B. & Br.* 480; neighborhood of Nassau, *Brace* 424; *Hitchcock* (*I. celosioides*)

Eleuthera — Governor's Harbor, *Hitchcock* (*I. celosioides*).

Great Galiot Cay — *B. & M.* 2854.

Cave Cay — *B. & M.* 2820.

Exuma — West Beach, *B. & M.*

Cat Cay — Port Howe, *Hitchcock (I. celosioides)*.

Watlings Island — *Hitchcock (I. celosioides)*.

***Iresine inaguensis* sp. nov.**

Caule ad basin lignoso erecto ramosissimo glabro. Foliis sessilis oppositis vel opposito-fasciculatis, glabris, anguste linearis obtusis margine integris; paniculis elongatis, terminalis densus et tenuiter ramosissimis subaphyllis, ramis densis, spiculis oppositis vel alternis, sessilibus, tenuis, rachi glabris, floribus sessilibus paleaceis, calyce bracteas laterales paulo superante, sepalis acutiusculis deltoideis in sicco carinatis; utriculus glaber, semen reniforme, obesum, nitidum, fusco-nigrum.

Plants erect 6–9 dm. compactly myriad branched, branches tenuous, internodes 5–8 cm. long, leaves 3–6 cm. long, 1 mm. wide or slightly larger. Seeds .6 x .5 mm.

Sheep Cay — Inagua, Oct. 18, 1904, *Nash & Taylor 1139 type*.

Inagua — Salt Pond Hill, *N. & T. 929*. This with almost capilliform leaves.

EUPHORBIACEÆ

***Savia bahamensis* Britton.**

Rocky places in coppices and pine barrens :

Abaco — Marsh Harbor, *Brace 1644, 1723, 1609, 1638*, and Cherokee Sound, *Brace 1965, 1970*.

Great Bahama — Eight Mile Rocks, *B. & M. 2407*; West End, *Brace 3504, 3501, 3640*.

Andros — Fresh Creek, *Northrop 610 (S. erythroxyloides)*, where it is known as "MAIDEN BUSH."

New Providence — *Brace 345, 3906*; *B. & Br. 670*; *Coker 157, 160*; Coppice on West Bay Street Road, *Britton 84 type*; and *Mills. 2163* from the same bush.

Hog Island — *B. & Br. 438*.

Wide Opening, Cay north of — *B. & M. 2772*.

Exuma — near Georgetown, *B. & M. 2959*.

Inagua — *N. & T. 1439*.

***Securinega acidothamnus* (Griseb.) Muell.**

In maritime coppices and scrub lands :

New Providence — at Southwest Landing, *B. & Br. 475*; *E. G. B. 3332*.

Mangrove Cay — *Coker 225*.

Exuma — near Georgetown. *B. & M. 2958*.

(*PHYLLANTHUS ANGUSTIFOLIUS* Gard. & Br.

This species of Swartz, reported by Gardner & Brace in Proc. Acad. Nat. Sci. Phila. 1889: 405, has not been collected by Mr. Brace, nor seen as yet from the islands.)

Phyllanthus bahamensis Urb.

In coppices:

Abaco — California Road, *Brace* 2037.Andros — at Red Bays, *Northrop* 488.New Providence — near Nassau, *Curtiss* 183; Waterloo, *B. & Br.* 741; *E. G. B.* 3435; Farringdon Road, *E. G. B.* 3403.Eleuthera — *Coker* 328.**Phyllanthus distichus** L.

Escaped from cultivation:

Andros — Fresh Creek, *Northrop* 653, where it is known as "GOOSE-BERRY TREE."**Phyllanthus epiphyllanthus** L.(*P. falcatus* Sw.) In various situations throughout the islands from Andros southward:Andros — Morgan's Bluff, where it is known as "HARDHEAD," *Northrop* 146a.North Bimini — *Mills* 2375.South Bimini — *Mills* 2411.South Cat Cay — *Mills* 2426.Silver Cay — *Mills* 2453.New Providence — *B. & Br.* 223, 520; *Britton* 14, 57; *Coker* 31; *Northrop* 325; *Cooper* 48; *Earle* 57; *Curtiss* 4; *Brace* 58, 230; *Hitchcock*; *Mills* 2068, 2081, 2132; *Wight* 18.Eleuthera — Governor's Harbor and Palmetto Sound, *Hitchcock*.Cave Cay — *B. & M.*Little Galiot Cay — *B. & M.* 2837.Great Guana Cay — *B. & M.* 2866, 2919.Exuma — *B. & M.* 3034, 3080.Cat Island — *Hitchcock*.Fortune Island — *Hitchcock*.Water Cay — *Coker* 524.Watlings Island — *Hitchcock*.Inagua — *N. & T.* 875; *Hitchcock*.Grand Turk Island — *N. & T.* 3766.

Varies from very long, slender leaves, through broader forms to short, broad, nearly ovate phyllodes.

Phyllanthus niruri L.

Open sandy places general:

Abaco — Cherokee Sound, *Brace* 2083.Little Harbor Cay — *B. & M.* 2252.

Great Harbor Cay — *B. & M.* 2338.

Andros — Nichols Town, *Northrop* 338; *Wight* 265.

North Bimini — *Millsp.* 2400.

New Providence — *E. G. B.* 3296 known as "GALE OF WIND"; *B. & Br.* 736; *Curtiss* 68; *Brace* 260; *Wight* 93.

Mangrove Cay — *Coker* 227.

Great Guana Cay — *B. & M.*

Exuma — *B. & M.*

Fortune Island — *Hitchcock*.

Inagua — *N. & T.* 1376; *Hitchcock*.

Grand Turk Island — *N. & T.* 3866.

Phyllanthus radicans (Muell Arg.) Small.

In pine barrens :

Great Bahama — Eight Mile Rocks, *B. & M.* 2381.

Phyllanthus saxicola Small.

On coral rock or sands in open places. The **type** is from South Florida between Coconut Grove and Cutler, Small & Carter 775 in herb. N. Y. Bot. Garden.

Great Bahama — Eight Mile Rocks and Barnett's Point, *B. & M.* 2489, 2623 (simple stemmed).

Great Sturup Cay — *B. & M.* 2273 (simple stemmed).

Little Harbor Cay — *B. & M.* 2250.

Great Harbor Cay — *B. & M.* 2333.

New Providence — at Tea House, *E. G. B.* 3447; near Nassau, *Curtiss* 79.

Great Guana Cay — *B. & M.* 2861.

Drypetes sp.

Specimen in leaf only :

Exuma — in a rocky coppice near Georgetown, *B. & M.* 2984. Apparently quite a different species from the following two. Leaves ovate narrowing to a prolonged sharp point, reticulate on both surfaces, upper surface dark, shining, lower surface pale, smooth. 4-6' x 2-1.5 cm., petioles about .5 cm.

Drypetes diversifolia Kr. & Urb.

D. keyensis Kr. & Urb. See remarks upon this species by Dr. Britton in Bull. N. Y. Bot. Gard. 3: 444, since which the following new localities have been determined :

Great Bahama — West End, *Brace* 3618.

Lignum Vitae Cay — *B. & M.* 2319.

Little Harbor Cay — *B. & M.*, where it is known as "WHITE WOOD."

Whale Cay — *B. & M.*

New Providence — at Winton, *E. G. B.* 3290.

Wide Opening, Cay north of — *B. & M.* 2769.

Little Galiot Cay — *B. & M.*

Stocking Island — *B. & M.* 3060.

Exuma — *B. & M.*

***Drypetes lateriflora* (Sw.) Kr. & Urb.**

Maritime sandy coppices:

Elbow Cay — *Brace* 1688.

Great Harbor Cay — *B. & M.* 2350.

New Providence — coppice on the north side of Blue Hills, *E. G. B.* 3452., the Race Course 3421, and at Winton 3293; *Eggers* 4331.

***Croton cascarilla* (L.) Linn.**

This species, first depicted by Catesby vol. 2 plate 46, has only been collected once since, then on New Providence by Mr. W. F. Daniell, whose specimen, as seen by Dr. Britton, proves Catesby's drawing to be a good representation of the species. All other reports of the species from the Bahamas or Antillean Islands are in error and are principally based upon forms of *Croton linearis*. Prof. Hitchcock's specimens are not this species.

***Croton discolor* Willd.**

So far known from only one station:

Rum Cay — *Coker* 453 at Port Nelson.

***Croton eluteria* (L.) Sw.**

In coppices and scrub lands, where it is known as "CASCARILLA" or "SWEETWOOD BARK." This is the *C. niveus* of Gardner & Brace:

Great Harbor Cay — *B. & M.* 2339.*

Hog Island — *Eggers* 4151.

New Providence — *Daniell*; *Brace* 56, 289; *Curtiss* 185; *Coker* 3a; *E. G. B.* 3143.

Eleuthera — *Eggers* 4425; *Hitchcock*, at Governor's Harbor.

Exuma — rocky coppices near Georgetown, *B. & M.* 2985.

***Croton flocculosus* Geis.**

So far known from only one island:

New Providence — near Nassau, *Curtiss* 61; *Brace* 516.

*Dr. Britton finds that Wright's type of *Croton homolepidus* in herb. British Museum is the same as *B. & M.* 2339, and identical with Catesby's specimen of *C. eluteria*.

Croton Hjalmarssoni Griseb.

In sandy opens and whitelands:

Great Guana Cay — *B. & M.* 2891.

Fortune Island — *Hitchcock*.

Inagua — *N. & T.* 874, 882 ("compares with the type in Herb. Kew" — Britton); *Hitchcock*.

Grand Turk Island — *N. & T.* 3782, 3802.

Croton humilis Linn.

The only specimens that we have seen from the Bahamas are:

Cat Island — at Port Howe, *Hitchcock*.

Croton linearis Jacq.

Maritime sand plains and dunes from the Berry Islands southward, where it is known generally as "GRANNY BUSH":

Great Sturup Cay — *B. & M.*

Little Harbor Cay — *B. & M.* 2244.

Great Harbor Cay — *B. & M.*

Whale Cay — *B. & M.* 2184.

South Bimini — *Millsp.* 2407.

Gun Cay — *Millsp.* 2315.

Andros — Fresh Creek, *Northrop* 615.

Rose Island — *B. & M.* 2113.

New Providence — *Northrop* 113; *Cooper sine num.*; *Coker* 23; *B. & Br.* 304; *Britton* 85; *Millsp.* 2154; *Wight* 65.

Eleuthera — Governor's Harbor, *Hitchcock*. (*C. cascarilla linearis*); *Coker* 387.

Great Guana Cay — *B. & M.*

Little Galiot Cay — where it occupies the southwest plateau almost to the exclusion of all other vegetation, *B. & M.* 2844.

Exuma — in scrublands north of Georgetown, *B. & M.* 2942.

Cat Island — at Port Howe, *Hitchcock*.

Watlings Island — *Hitchcock* (*C. cascarilla L.*). This specimen is a counterpart of a sheet of *C. linearis* from Palm Beach, Florida, Curtiss 5360, in Herb. Field Col. Mus.

Inagua — *N. & T.* 906, these specimens have very long and narrow leaves, and smaller seeds than the usual form.

Croton lobatus Linn.

In cultivated soils. This species can be unhesitatingly placed among the Antillean weeds that have been more or less lately introduced in the Bahamas:

New Providence — *Earle* 32; *B. & Br.* 783; *Coker* 553; *Brace* 353.

Croton lucidus Linn.

Sandy scrub and white-lands near the sea :

Great Sturrupey — *B. & M.* 2283.

Great Harbor Cay — *B. & M.*

New Providence — near Clifton, *E. G. B.* 3323; *Cooper* 22; *Britton* 21, 82; *Brace* 68, 288, 3896; *Curtiss* 108; *Mills* 2077, 2156; *Wight* 109, 173.

Eleuthera — *Coker* 326.

Exuma — *B. & M.* 2930.

Watlings — *Coker* 471.

Croton rosmarinifolius Griseb.

One locality only is known up to this date :

Exuma — abundant in one situation, an open, rocky field just north of Georgetown, where it forms small, bushy trees up to 3.5 m. high, *B. & M.* 2978.

Our specimens compare exactly with Wright's, Cuba, 1968 in Hb. Kew.

Argythamnia argentea sp. nov.

Fruticulus ad basi ramosissimus, rami virgatis subdense foliosis, ramuli canescentibus. Foliis lanceolatis utraque attenuatis breve petiolatis margine superne paucodentatis utraque pagina argenteo-sericeis; stipulis crassis cuspidatis. Racemulis depauperatis supra axillaribus. Flores masc. 3, sepalis 4 ovato-lanceolatis, petalis 4 crassis, sepalis similiter sed brevissimis, bracteola 1 scaphoideis; flores foem. 3, sepalis elongato-lanceolatis, petalis minute ligulatis glandulae alternantibus; stylis 3, prox basi bifurcatis ad apice bilobatis, ovario canescentibus.

Near *A. sericea* from which, however, its characters plainly separate it. The **type** and only plants so far seen are from :

Grand Turk Island — in scrubland Aug. 27–Sept. 1, 1905, *Nash & Taylor* 3830.

Argythamnia candicans Sw.

So far only known in the Bahamas through the following specimens that are clearly this species :

Cat Island — Port Howe, *Hitchcock*.

Inagua — *Hitchcock*.

Argythamnia lucayana sp. nov.

Fruticulus ramosis prostratis vel ascendentibus. Rami minute sericeis sparse foliosis, foliis sessilibus elliptico-velobovato-lanceolatis apice acute vel brevissime acuminatis margine integris vel praeter admodum paucos minute et remote glanduloso-denticulatis, matura glabris imma-

tura subtus sparse et minute strigose pilosis. Racemuli 4-6 floris. Sepalis masc. deltoideo-lanceolatis petalem aequantibus, petalis spathulatis, pilosis, staminis 4, laevis, calycis foem. laciniis foliaceis lanceolatis acuminatis, petalis minutis, angustis-lanceolatis obtusis paleaceis calyce pluries brevioribus, ovario villosa-pubescentis, stylis villosis bis dichotome divisis, dichotomiis terminis brevis. Semen ad maturitas globosis, apiculatis, anastomoso-reticulatis.

A small shrubby plant with branches 3-5 dm. long. Leaves bluish-green 2.5-3.5 cm. long, 1-1.5 cm. broad. Racemes 6-10 mm. long, 4-6 flowered. Seeds 1.5 mm. in both diameters, dark brown, the surface coated with a hydnum-like porous integument.

This is the common *Argythamnia* of the Bahamian islands, where it is to be found on the floor of the coppices or at times in more open situations:

Great Sturrupey — *B. & M.* 2280.

Little Harbor Cay — *B. & M.* 2232.

Eleuthera — Tarpum Bay, *Coker* 409 (*A. candidans*).

Ship Channel Cay — *B. & M.* 2760.

Wide Opening, Cay north of — *B. & M.* 2789.

Cave Cay — *B. & M.* 2834.

Great Guana Cay — in rocky coppice, *B. & M.* 2913 type.

Little Galiot Cay — *B. & M.* 2839.

Exuma — *B. & M.* 3601.

***Argythamnia sericea* Griseb.**

The only island upon which this species has as yet been found in typical form is:

Inagua — in sand on whitelands at Tenados, Oct. 14, 1904, *N. & T.* 1030.

A striking form with flabellate leaves and minor character differences is returned from:

Fortune Island — *Brace* 462.

***Mercurialis annua* L.**

Cultivated ground:

New Providence — *Wight* 118.

This is the first return of this species from the Bahamas.

***Bernardia bernardia* (L.) Britton.**

Coker in Shattuck's "Bahama Islands" p. 257. *Adelia Bernardia* L.

Bernardia carpinifolia Griseb., *Bernardia dichotoma* Mill.

Eleuthera — at Tarpum Bay, *Coker* 405; Governor's Harbor, *Hitchcock* (*B. mexicana*).

Cat Island — Port Howe, *Hitchcock* (*B. mexicana*).

Fortune Island — *Brace* 458.

Lasiocroton macrophyllus Griseb.

So far known only from Mrs. Northrop's specimens which, while not absolutely agreeing with Marsh's Jamaican specimen in Hb. N. Y. Bot. Garden, are undoubtedly the species. The sheet in Hb. Field Col. Mus. is in good flowering condition.

Andros — at Deep Creek, *Northrop 689*, where it is known as "WILD OAK," "LIGHTWOOD" or "BITTERS".

Acalypha alopecuroidea Jacq.

Doubtless an introduced weed in waste and cultivated grounds :

Abaco — at Marsh Harbor, *Brace 1751*.

New Providence — near Nassau, *Northrop 32; Curtiss 22; Hitchcock; Wight 39; Brace 132*, and at Southwest Landing, *B. & Br. 474*.

Eleuthera — at Governor's Harbor, *Hitchcock*.

Inagua — at Mathew Town, *N. & T. 1080*.

Acalypha ostryaefolia Ridd.

Known only by the following plant, probably introduced from Florida:

New Providence — at Nassau, *Brace 385 (300?)*

Acalypha setosa A. Rich:

Known only by the following plant, doubtless an introduced weed.

New Providence — at Nassau, *Brace 299 (267?)*.

Pera bumeliaefolia Griseb.

In coppices; known as "BLACK EBONY":

Abaco — at Marsh Harbor, *Brace 1603*.

Andros — *Brace 411*.

New Providence — coppice at Seven Hills, *E. G. B. 3360* the *forma parvifolia* of Urban in Hb. Kew.; Fox Hills, *B. & Br. 376*; near Nassau, *B. & Br. 267; Coker 537; Curtiss sine num., Brace 226; Wight 166*.

Ricinus communis Linn.

Escaped from gardens on :

New Providence — *Wight 169*.

Fortune Island — *Hitchcock*.

Jatropha curcas Linn.

Known only by the following specimen, probably an escape from a garden :

New Providence — near Nassau, *Brace 305*.

Jatropha gossypifolia Linn.

The only specimens seen are from waste lands near Nassau :
New Providence — *Hitchcock*; *Brace*.

Manihot manihot (Linn.) Cockl.

Manihot apii Pohl. Known only as cultivated at :
Andros — Nichols Town, *Northrop* 363.
Eleuthera — Governor's Harbor, *Hitchcock*.

Excoecaria Sagraei Muell.

In coppices. The plant has a strong mephitic odor :
Andros — Stafford Creek, *Northrop* 539.
New Providence — *Curtiss* 190; *B. & Br.* 665; Fox Hills Road, *E. G. B.* 3388; *Britton* 78; *Millsp.* 2160, 2486.
Exuma — *B. & M.* 3079.
Watlings Island — *Coker* 478.

Bonania cubana A. Rich.

In coppices and scrub lands :
Andros — at Fresh Creek, *Northrop* 628 (*B. emarginata*); *Wight* 254.
New Providence — in a coppice at Winton, *E. G. B.* 3294; *Brace* 141, 340.
Cat Island — in a low, dry coppice at Arthur's Town, *Coker* 428.
Exuma — near Rolletown, *B. & M.* 3074.

Hippomane mancinella Linn.

In coppices and scrub lands. Known generally as "MANCHINEEL" or "POISON BUSH."
Abaco — Cherokee Sound, *Brace* 1920.
Andros — Fresh Creek and Conch Sound, *Northrop* 622, 565 (misprinted 556 in Fl. N. P. & Andros).
Great Guana Cay — *B. & M.* 2873.
Watlings Island — *Coker* 485.
Inagua — at Lantern Head and Tenados, *N. & T.* 1392, 1048; *Hitchcock*.
Grand Turk Island — in scrublands, *N. & T.* 3826.

Gymnanthes lucida Sw.

Coppices and scrublands. Known generally as "CRABWOOD":
Allen's Cay — *Brace* 1547.†
Green Turtle Cay — *Brace* 1512.†
Great Bahama — at Barnett's Point, *B. & M.* 2642.†
Great Harbor Cay — *B. & M.*
Lignum Vitae Cay — *B. & M.*
Great Sturup Cay — *B. & M.*

Goat Cay — *B. & M.*, where it forms character thickets.

Whale Cay — *B. & M.*

South Bimini — *Brace* 3476.†

Gun Cay — *Mills* 2327.°

Cat Cay — *Brace* 3751.†

Andros — at Fresh Creek and Nichols Town, *Northrop* 375;° *Wight* 248.°

Rose Island — *B. & M.*

New Providence — near Clifton, *E. G. B.* 3326;° coppice at Lake

Cunningham, *B. & Br.* 606,† 607;§ Fox Hills, *B. & Br.* 862;°

E. G. B. 3393;° *Britton* 103;§ *Brace* 299,† 335,† *Curtiss* 190;

Mills 2183.§

George's Island — Spanish Wells, *Coker* 314.†

Ship Channel Cay — *B. & M.*

Wide Opening, Cay north of — *B. & M.* 2804.†

Great Guana Cay — *B. & M.*

Exuma — *B. & M.*

Watlings Island — *Coker* 467.†

Inagua — on Salt Pond Hill, *N. & T.* 927,* 992.*

Sheep Cay — *N. & T.* 1145.*

Hura crepitans Linn.

Cultivated, only at Nassau.

Euphorbia Berteriana Balbis.

So far seen only in the redlands of:

Exuma — *B. & M.* 2938, 3031.

Euphorbia Blodgettii Engelm.

On rocks, sand dunes, whitelands and in pot holes common. Not yet found on the islands of the Abaco group.

Great Bahama — *B. & M.* 2383, 2470, 2619.

Great Sturru Cay — *B. & M.* 2257, 2272, 2274.

Frozen Cay — *B. & M.* 2216.

Whale Cay — *B. & M.* 2182.

North Bimini — *Mills* 2404; *Brace* 3462.

South Bimini — *Mills* 2364.

Gun Cay — *Mills* 2310.

Andros — Nichols Town, *Northrop* 379.

Salt Cay — *Northrop* 277 (*E. serpens*).

Hog Island — *B. & M.* 2169, 2174.

Rose Island — *B. & M.* 2146.

*Leaves ovate 3 x 1.5 cm. †Leaves ovate-spatulate 5 x 1.7 cm. ‡Leaves ovate-lanceolate 8 x 2.7 cm. °Leaves ovate-lanceolate 6 x 1.8 cm. §Leaves willow-like 6-8 x 1 cm.

New Providence — *Cooper sine num.*; *Millsp.* 2100, 2149, 2150, 2167, 2271; *Curtiss* 70 (*E. Nashii Small*); *Britton* 72, 74, 90; *B. & Br.* 195, 279, 309, 313, 340, 400, 477, *Northrop* 41; *Hitchcock*.

Eleuthera — Governor's Harbor, *Hitchcock* (3 sheets).

Ship Channel Cay — *B. & M.* 2746, 2755,

Great Guana Cay — *B. & M.* 2863,

Cave Cay — *B. & M.* 2805, 2818.

Great Galiot — *B. & M.* 2859.

Exuma — *B. & M.* 2961, 2966, 3107, 3112.

Fortune Island — *Hitchcock*.

Crooked Island — *Hitchcock*.

Inagua — *N. & T.* 954, 1090, 1435; *Hitchcock*.

Sheep Cay — *N. & T.* 1159.

Grand Turk Island — *N. & T.* 3851, 3869.

Euphorbia Bracei sp. nov;

§ Chamaesyceae. Caulibus prostratis ad basin ramosis, rami lignosis canescentis, foliis petiolatis oppositis oblongis vel ovatis ad basin oblique-deltoides apice rotundo-obtusis margine integris revolutis utrinque pagina hirsutis et resinosis, stipulis triangulo-lanceolatis acutis, involucris axillaribus, cupuliformis, pedicelis tubo aequantibus, extus canescentis intus ciliatis, mure crassis interiore in columnae alaribusque 5 productis, lobis triangularis acutis, glandulis minutis transverse-ovatis appendice angustissima vel obsoleta crenato-3 dentatis, stigmatis ad basin bifurcatis. Capsulae aciculo-hirsutis, semen triangulo-ovoides leviter anastomo-rugosis.

A low, grayish plant near *E. cayensis* Millsp. from which it differs in many characters. Leaves 8 x 5 to 5 x 3 mm. dotted with minute amber-like resin globules; petiole one-fifth the length of the blade; internodes 1.3-2 cm.; fifth gland replaced by a broad fifth tooth equalling the breadth of a gland; seed .9 x .7 mm., the angles prominent.

Abaco — in sand at Eight Mile Bay, *Brace* 1856 type.

Man o'War Cay — on whitelands, *Brace* 1571.

Euphorbia brasiliensis Lam.

In open places and grassy coverts:

New Providence — at Lake Cunningham, *Britton* 137; *Millsp.* 2227; cultivated ground, Farringdon Road, *E. G. B.* 3199.

Eleuthera — at Gregory Town, *Coker* 371.

Euphorbia Brittonii sp. nov.

§ Chamaesyceae. Pilosa praegracilis erectis ad basin ramosis, rami filiosis ascendentibus, foliis oppositis breviter petiolatis rotundatis vel oblongo-ellipticis apice rotundatis basi leviter et oblique cordatis margine integerrimis vel acute subdentatis revolutis obsolete venosis nervo

medio subtus prominentis, crassiusculis, pagina subtus pilosis glauco-viridis; stipulis triangulatis plus minusve aequaliter fimbriatis; involucris solitariis axillaribus, campanulatis, pedicellatis, extus glabris intus ad faucam dense tomentosis, lobis triangularis inflexis praeciliatis, glandulis ovatis albo-viridis, appendice magnis orbiculatis albis. Stamina insertis pilosis (paleis linearibus glabris intermixtis), stigmatis supra bifurcatis; capsulae glabris; semina roseo-cinereis ovatis valide triangulatis leviter transverse rugosis.

A small, erect, wiry plant branching from the base, 5–7 cm. high; leaves 4 x 3 mm. glands 4 the fifth replaced by a sharp fissure flanked by 2 larger involucral lobes.

So far known only from sandy whitelands, as follows:

New Providence,—at the Race Course, Aug. 26, 1904, *B. & Br.* 839 type and 279.

Euphorbia buxifolia Lam.

Maritime beaches from Great Bahama southward.

Green Turtle Cay — *Brace* 1508.

Abaco — Cherokee Sound, *Brace* 1902, and Butler Bay, 1520.

Great Bahama — infrequent, low and undeveloped. Eight Mile Rocks, in a clearing near the sea, *B. & M.* 2781.

Great Sturup Cay — *B. & M.* 2256.

Great Harbor Cay — *B. & M.*

Lignum Vitae Cay — *B. & M.*

Frozen Cay — *B. & M.*

Whale Cay — *B. & M.*

Gun Cay — *Millsp.* 2317.

South Cat Cay — *Millsp.* 2417, 2418.

Andros — at Red Bays and Big Cabbage Creek, *Northrop* 457, 672.

Silver Cay — *Millsp.* 2454.

Rose Island — *B. & M.*

New Providence — South shore, *Millsp.* 2109; near Nassau, *Cooper sine num.*; *E. G. B.* 3185; *Northrop* 87; *Coker* 114; *Hitchcock*; *Wight* 55.

Eleuthera — Governor's Harbor, *Hitchcock*.

Ship Channel Cay — *B. & M.*

Wide Opening, Cay north of — *B. & M.* 2781.

Great Guana Cay — *B. & M.*

Great Galiot Cay — *B. & M.*

Little Galiot Cay — *B. & M.*

Cave Cay — *B. & M.*

Exuma — *B. & M.*

Stocking Island — *B. & M.*

Cat Island — Port Howe, *Hitchcock*.

Watlings Island — *Hitchcock*.

Crooked Island — *Hitchcock*.

Fortune Island — *Hitchcock*.

Inagua — *N. & T.* 1038.

Sheep Cay — *N. & T.* 1160.

Grand Turk Island — *N. & T.* 3793, 3889.

Euphorbia cassythoides Boiss.

Known so far only from the following specimens:

Andros — Deep Creek, *Northrop* 702.

Euphorbia cayensis Millsp.

Maritime sands. So far found only on the smaller cays from Joulter's Cays northward.

Man o'War Cay — *Brace* 1571.

Great Sturup Cay — *B. & M.* 2266.

Great Harbor Cay — *B. & M.* 2334 (prostrate).

Little Harbor Cay — *B. & M.*

Frozen Cay — *B. & M.*

Whale Cay — *B. & M.* 2178, where it is the character plant of the maritime whitelands, and is known as "MILK BUSH."

Joulter's Cays — *Millsp.* 2295 **type**.

Euphorbia flexuosa Kth.

E. buxifolia flexuosa (Kth.) Boiss. Anyone seeing this species in the field would at once recognize it as clearly distinct from *E. buxifolia*. Its purplish coloration, cinereous glaucescence, rounded flat leaves and spreading habit cause it to resemble *E. Cayensis* far more closely than *E. buxifolia*.

Leaves glauco-cinereous, ovate, rounded at the apex and unequal base .3-1 cm. x .2-.6 cm. Involucres turbinate, short pediceled, smooth without, densely bearded within, margin entire; glands 4, large, cordate, nearly equalling the appendages; stamens 5, the filaments bearded and tufted below; seeds triangular, pink-ashen, farinose, the angles rounded and prominent, the facets slightly anastomose-rugose transversely.

Maritime sands and sand-pocketed rocks. Not seen north of the Exuma Chain.

Ship Channel Cay — *B. & M.* 2747.

Wide Opening, Cay north of — *B. & M.* 2777, 2779, On the open sea-side of this cay the plants are strongly characteristic and to be found in great quantity both erect and prostrate.

Little Galiot Cay — *B. & M.*

Stocking Island — *B. & M.* 3071.

Rum Cay — *Coker* 452 (*E. cayensis*). Mr. Coker mentions seeing this plant also on Abaco, but doubtless mistook the Abacan *E. Bracei* for this species.

***Euphorbia havanensis* Willd.**

E. heterophylla graminifolia Eng. This species as it occurs in South Florida and the Keys, Cuba and the Bahamas, is plainly distinct from *E. heterophylla*.

Green Turtle Cay — *Brace* 1479.

Great Sturrupey — *B. & M.* 2255.

New Providence — *Northrop* 92; *Millsp.* 2146; *Britton* 69; *Coker* 45; *Wight* 14.

Eleuthera — Governor's Harbor, *Hitchcock* (*E. heterophylla graminifolia*).

Exuma — near Georgetown, *B. & M.* 3133.

Cat Island — Port Howe, *Hitchcock* (*E. heterophylla graminifolia*).

***Euphorbia heterophylla* Linn.**

This species appears to exist in all its forms in the Bahama Islands where it is principally found in open waste grounds.

Green Turtle Cay — *Brace* 1503, 1479.

Great Bahama — Eight Mile Rocks in pine barren, *B. & M.* 2461; West End, *Brace* 3499.

Little Harbor Cay — *B. & M.*

South Bimini — *Millsp.* 2365.

South Cat Cay — *Millsp.* 2435.

New Providence — near Nassau, *Northrop* 96; *Millsp.* 2204; *Hitchcock*; *Wight* 29.

Eleuthera — Governor's Harbor, *Hitchcock*.

Great Guana Cay — *B. & M.*

Little Guana Cay — *B. & M.*

Exuma — in a cornfield, *B. & M.*

The form with the lower (and often all) leaves broadly ovate-lanceolate, entire or obsoletely repand, *E. prunifolia* Jacq. This latter form plainly belongs to *E. heterophylla* rather than to *E. geniculata* Ortega. Under this form the following specimens may be grouped:

Green Turtle Cay — *Brace* 1539.

South Cat Cay — *Millsp.* 2426.

New Providence — near Nassau, *Earle* 33; *Curtiss* 73; *Hitchcock* (*E. geniculata*); *Wight* 5.

Eleuthera — Governor's Harbor, *Hitchcock* (*E. geniculata*).

Cat Island — Port Howe, *Hitchcock* (*E. geniculata*).

Fortune Island — *Hitchcock* (*E. geniculata*).

Inagua — Mathew Town, *N. & T.* 1086; *Hitchcock* (*E. geniculata*).

Abaco — Eight Mile Bay, *Brace* 1870.

Euphorbia hypericifolia Linn.

In grassy open lands :

Great Bahama — Eight Mile Rocks, *B. & M.* 2476.

North Bimini — *Mills* sp. 2370, 2399.

Andros — Nichols Town, *Northrop* 380 (*E. nutans*).

New Providence — at Lake Cunningham, *Mills* sp. 2227; near Nassau, *Earle* 46^a; *B. & Br.* 470, 535; *Hitchcock*; *Wight* 46.

Eleuthera — Governor's Harbor, *Hitchcock*.

Great Guana Cay — *B. & M.* 2906.

Exuma — near Georgetown, *B. & M.* 2945.

Inagua — Smith's Thatch Pond, *N. & T.* 1401, 1406; *Hitchcock*.

Euphorbia lactea Haw.

Widely escaped from cultivation and forming dense thickets on:

New Providence — along the West Bay Street Road, *Northrop* (*E. antiquorum*).

Inagua — Mathew Town, *Hitchcock* (*E. antiquorum*).

Euphorbia lecheoides sp. nov.

Tota griseo furfureo-farinoso caulibus e rhizomate lignoso, ramosissimo. Rami longis, diffusis rigidis dichotomis, internodiis inferioribus prox. 1 cm. longis, superne tenuiter virgatis rigidis. Foliis breviter petiolatis, deltoideo-ovatis vel triangulari-ovatis abrupte attenuatis ad margine revoluti-intumescens, pagina utroque furfureo-farinoso, stipulis late ovatis blepharociliatis, involucris solitariis turbinatis brevi pedicellatis, extus glabris intus ad faucem barbatis, lobis triangularibus acutis ciliatis, glandulis nigris complano-orbiculatis, facie notis variorum insignis, appendices minutis carnosus vel obsoletis, stigmata ad mediam bifurcatis, filamentae barbatis. Capsulae glabris. Semen fuscis triangulo-ovatis fascies minute papillatis.

Plants rigidly erect or diffusely prostrate 20–30 cm. Leaves 2.5 to 5 mm. long, 1 to 2.5 mm. wide. Seeds .8 x .5 mm. The specimens representing the extremes of this species are so widely different in general appearance as to almost convince one that they represent two strongly marked species. This difference, however, lies entirely in the habit of the plants and the size of the leaves. The tallest erect form with small, ovate leaves (3796) passes through the type (3888) to a more diffuse, larger leaved form (1305) and a fully prostrate plant with much larger, sickle-shaped leaves (1 cm. x 3 mm.) with a somewhat less intumescent margin.

Inagua — in scrubland on the trail from Canfield Bay to Cabbage Pond, Oct. 22, 1904, *Nash & Taylor* 1305.

Grand Turk Island — in sandy scrublands, *Nash & Taylor 3888 type* and on the strand, *3796*.

Euphorbia obliterata Jacq.

This species, considered a synonym of *E. pilulifera procumbens* by Boissier, is distinct both in characters and in habitat.

Great Bahama — at Eight Mile Rocks, *B. & M. 2471*.

Little Harbor Cay — *B. & M. 2247*.

New Providence — on coral limestone (the usual habitat) along Farringdon Road, *B. & Br. 199*, and Soldiers Road *807*; *Wight 179*.

Cave Cay — *B. & M. 2817*.

Exuma — *B. & M. 2938, 3031*.

Inagua — *Hitchcock (E. pilulifera obliterata.)*

Euphorbia pilulifera Linn.

Frequent in cultivated grounds and along roads and paths:

New Providence — *B. & Br. 757, 805*; *Mills. 2216*; *Hitchcock*.

Eleuthera — Governor's Harbor, *Hitchcock*.

Cat Island — Port Howe, *Hitchcock*.

The prostrate form with the same characteristic jointed amber colored hairs (*E. pilulifera procumbens* Boiss.) from:

New Providence — *B. & Br. 757*.

Inagua — *N. & T. 1088*.

Euphorbia prostrata Aiton.

So far known only from two localities:

New Providence — near Nassau, *Curtiss sine num.*

Grand Turk Island — *N. & T. 3871*.

Euphorbia punicea Sw.

On the southern islands only:

Watlings Island — on a small cay in the lake, *Coker 487*.

Fortune Island — *Hitchcock*.

Inagua — *N. & T. 1263, 1446*.

Little Inagua — *N. & T. 1216*.

Euphorbia trichotoma Lam.

This species, common on the maritime sands of the Florida Keys and the Antillean Islands, is so far only known in the Bahamas by specimens collected on:

Allen's Cay — *Brace 1551*.

Great Bahama — at Eight Mile Rocks, *B. & M. 2364*; West End, *Brace 3493*.

Euphōrbia vaginulata Griseb.

The following specimens represent the important rediscovery of this endemic species.

Little Inagua — Moujean Harbor, *N. & T.* 1222.

Inagua — *N. & T.* 1170, 1304.

Grand Turk Island — *Hjalmarsson (type)* *N. & T.* 3764.

The plants grow in great profusion on Grand Turk Island. The Inagua specimens, while apparently differing from those of Grand Turk Island and Little Inagua on account of their longer leaves (twice or thrice the length) and darker bark, have the same characters as the type.

Pedilanthus tithymaloides (L.) Poit. ?

In coppices, rare:

Andros — at Deep Creek, *Northrop* 693 (*P. angustifolius* ?)

Inagua — at Mathew Town, *N. & T.* 1363.

Grand Turk Island — *N. & T.* 3761.

Plants from Grand Turk Island are now growing in the conservatory of the Garden which give promise of settling upon the specific character of this form.

RUBIACEÆ.

Rachicallis maritima (Jacq.) K. Sch.

R. rupestris (Sw.) D. C. On maritime rocks throughout the Islands. Often the only vegetation on many of the sea washed islets. Observed by *B. & M.* and *Mills* on each island visited. The collections contain the following sheets:

Spanish Cay — *Brace* 1562.

Great Bahama — *Allen* 36.

South Cat Cay — *Mills* 2425.

North Cat Cay — *Mills* 2415.

Andros — Red Bays and Fresh Creek, *Northrop* 458 (*R. Americana*) where it is known as "SAND-FLY BUSH," "SALT WATER BUSH," "SEAWEED" and "WILD THYME."

New Providence — *Cooper* 2; *B. & Br.* 293; *Curtiss* 32; *Coker* 19; *Wight* 190; *Mills* 2249; *Brace* 161, 165, 173; *Hitchcock* (*R. Americana*.)

Crooked Island — *Hitchcock* (*R. Americana*).

Inagua — *N. & T.* 904, 1366, 1391.

Grand Turk Island. — *N. & T.* 3812.

Exostema caribaeum (Jacq.) Willd.

In shady coppices. Known as "PRINCE WOOD."

Elbow Cay — *Brace* 1698.

Abaco — Cherokee Sound, *Brace* 1930.

Great Bahama — Eight Mile Rocks and Golden Grove, *B. & M.* 2433, 2733; West End, *Brace* 3516.

Andros — Kemp Sound and Deep Creek. *Northrop*, 685.

Rose Island — *B. & M.* 2117.

New Providence — Fort Montague, *B. & Br.* 316; *Brace* 187; Fox Hills, *E. G. B.* 3392.

Wide Opening, Cay north of — *B. & M.* 2787.

Exuma — *B. & M.*

Watlings Island — *Coker* 465.

Crooked Island — *Hitchcock*.

Inagua — *N. & T.* 952, 1431.

Genipa clusiaefolia (Jacq.) Griseb.

Coastal rocks of all islands visited (*B. & M., Millsp.*). Known generally as "SEVEN YEAR APPLE."

Andros — Mastic Point, *Northrop* 299.

North Cat Cay — *Millsp.* 2346.

New Providence — *Brace* 41; *Millsp.* 2113; *Hitchcock*.

Watlings Island — *Hitchcock*.

Cat Island — *Hitchcock*.

Crooked Island — *Hitchcock*.

Fortune Island — *Hitchcock*.

Inagua — *N. & T.* 937, 1345.

Little Inagua — *N. & T.* 1183.

Randia aculeata Linn.

In thickets, coppices and scrublands. The farther south in the Archipelago the plants are found the smaller the leaves appear. The spinous or aspinous character is not, however, so constant. Not seen on the Bimini Group.

Elbow Cay — *Brace* 1703.

Abaco — Cherokee Sound, *Brace* 1918.

Great Bahama — Eight Mile Rocks, *B. & M.* 2432, West End, *Brace* 3540.

Berry Islands — on all islands visited except Frozen and Goat Cays, *B. & M.*

Little Harbor Cay — *B. & M.* 2225.

Cat Cay — *Brace* 3743.

Andros — Nichols Town, *Northrop* 383; *Wight* 252.

Rose Island — *B. & M.*

New Providence — *B. & Br.* 323; *Coker* 56; *Hitchcock*.

Exuma Chain — noted on all islands visited except Great Guana Cay, *B. & M.*

Eleuthera — Palmetto Sound, *Hitchcock*.

Cat Island — Port Howe, *Hitchcock*.

Crooked Island — *Hitchcock*.

Inagua — *N. & T.* 1060, 1062 with very small leaves; *Hitchcock*.*

Grand Turk Island — *N. & T.* 3759.

***Hamelia patens* Jacq.**

Waste grounds near habitations. This habitat, and the absence of the species from other islands, place it as a doubtless escape from cultivation.

New Providence — Lake Cunningham, *B. & Br.* 635; Bluebeard's Tower, *Coker* 127; Nassau, *Northrop* 40; *Wight* 147.

***Catesbaea campanulata* La Sagra.**

C. parviflora septentrionalis Urb. In coppices and scrub lands:

New Providence — *Brace* 167.

Crooked Island — *Hitchcock* (*C. parviflora*). See remarks on Grand Turk Specimen below.

Inagua — *N. & T.* 1272.

Grand Turk Island — *N. & T.* 3771. This is a low, scrubby specimen having no spines developed upon the short new shoots. It appears to be a depauperate mutant with narrower and more revolute leaves and smaller seeds than the typical form.

***Catesbaea fasciculata* Northrop.**

In sandy coppices and whitelands:

Abaco — Eight Mile Bay, *Brace* 1852.

Little Harbor Cay — *B. & M.* 2226.

Great Harbor Cay — *B. & M.*

Andros — Fresh Creek, *Northrop* 627 **type**.

New Providence — scrublands along West Bay Street Road, *Britton* 83; *Mills* 2162; coppice at Winton, *E. G. B.* 3292; whitelands near Race Course and Miller's, *B. & Br.* 284, 529.

Green Cay — maritime beach, *Coker* 247.

Ship Channel Cay — *B. & M.* 2750.

Wide Opening, Cay north of — *B. & M.*

Great Guana Cay — *B. & M.*

***Catesbaea spinosa* Linn.**

Sandy open lands:

Andros — at Fresh Creek, *Northrop* 624. Known as "PRICKLY APPLE."

*Prof. Hitchcock observes on p 93 Pl. Bah.: "The Inagua specimens are spineless and have small leaves about 7 mm. long." The Nash & Taylor specimens have such leaves and also spines.

Eleuthera — on a side hill near Gregory Town, *Coker* 367.

New Providence — So far only seen in the Governor's Garden (*Britton* 153; *Mills* 2247) where it is planted from Harbor Island. Mr. Brace remarks that he once knew a bush of this species in the scrubland back of Nassau but that it has long since disappeared.

Guettarda elliptica, Sw.

In coppices and scrub lands:

Abaco — Cherokee Sound, *Brace* 1935.

Great Bahama — Eight Mile Rocks, *B. & M.* 2583; West End, *Brace* 3554.

Andros — Lisbon Creek, *Northrop* 677.

Hog Island — *B. & Br.* 349.

New Providence — near Nassau, *Brace* 166; *Hitchcock*; Fort Montague, *B. & Br.* 173.

Eleuthera — Governor's Harbor, *Hitchcock*.

Great Guana Cay — *B. & M.*

Exuma — *B. & M.* 2979.

Cat Island — Arthur's Town, *Coker* 425.

Long Island — Clarence Harbor, *Coker* 515.

Fortune Island — *Hitchcock*.

Inagua — *N. & T.* 1266; *Hitchcock*.

Guettarda Krugii Urban.

In scrub lands and rocky coppices:

Man o'War Cay — *Brace* 1582.

Great Guana Cay — *B. & M.* 2870.

Exuma — scrub lands along Hayne's Road, *B. & M.* 3022.

Little Inagua — *N. & T.* 1309.

Inagua — Mathew Town, *N. & T.* 918, 919, 1369; *Hitchcock* (*G. calyptrata*).

Grand Turk Island — *N. & T.* 3800.

Guettarda scabra Lam.

In scrub lands and coppices:

Abaco — Cherokee Sound, *Brace* 1909.

Great Bahama — Eight Mile Rocks, *B. & M.* 2399.

Andros — at Fresh Creek, *Northrop*, 535, 730.

New Providence — Near Nassau, *Curtiss sine num.*; *Hitchcock*; *Brace* 186, 197; Village Road and Farringdon Road, *B. & Br.* 369, 259.

Eleuthera — opposite Spanish Wells, *Coker* 318.

Inagua — *Hitchcock*.

Antirrhoea.

This genus is being considered by Dr. Britton,

Laugeria densiflora (Griseb.) Benth. & Hook.

A viscid shrub frequent in coppices:

Abaco — Marsh Harbor, *Brace* 1604, 1614, 1714.

New Providence — *Eggers* 4460; *Curtiss sine num.*; *Brace* 389; Blue Hills Road, *Coker* 551; Soldiers Road, *B. & Br.* 587; Fox Hills, *E. G. B.* 3389; *Hitchcock*.

Erithalis fruticosa Linn.

Everywhere. In low, sandy coppices, sandy thickets, scrub lands, coastal thickets and sand dunes, where it is generally known as "BLACK TORCH."

Man o'War Cay — *Brace* 1569.

Abaco — Cherokee Sound, *Brace* 1899.

Great Bahama — Eight Mile Rocks, *B. & M.* 2371; West End, *Brace* 3511.

Great Sturup Cay — *B. & M.*

Little Harbor Cay — *B. & M.* 2229.

Great Harbor Cay — *B. & M.*

Frozen Cay — *B. & M.*

Whale Cay — *B. & M.*

South Bimini — *Mills* 2413

Gun Cay — *Mills* 2311.

Andros — Deep Creek and Red Bays, *Northrop* 691, 365.

Silver Cay — *Mills* 2449.

Rose Island — *B. & M.* 2126.

New Providence — near Nassau, *Cooper* 4; *Brace* 27; *Coker* 73, 161, 192; *Wight* 138; South Shore, *Britton* 43, 52; *Mills* 2115, 2123; *Hitchcock*.

Eleuthera — Governor's Harbor, *Hitchcock*.

Exuma Chain — on all cays visited and on Exuma Island, *B. & M.*

Cat Island — Port Howe, *Hitchcock*.

Watlings Island — *Hitchcock*; *Coker* 463.

Crooked Island — *Hitchcock*.

Fortune Island — *Hitchcock*.

Inagua — *Hitchcock*; *N. & T.* 949, 1029, 1054, 1443.

Sheep Cay — *N. & T.* 1149.

Grand Turk Island — *N. & T.* 3768, 3809, 3816, 3817.

The form of the exposed sea beaches has leaves nearly orbicular, in that of the denser coppices the leaves are quite narrowly lanceolate.

Erithalis odorifera Jacq.

"Common in coppices on New Providence, and distinct from the more abundant *E. fruticosa* L., being much taller, the corolla twice as

large as in that species, the fruit and persistent calyx-limb larger."—

N. L. Britton.

Andros — Deep Creek, *Northrop* 739 (*E. rotundata*).

New Providence — low coppices, *Coker* 43 (*E. fruticosa*); near Nassau,

B. & Br. 234, Blue Hills road, 580, Lake Cunningham 605; *Britton* 147; *Millsp.* 2242; Nassau, *Curtiss sine num.* :

Phialanthus myrtilloides Griseb.

Coppices and scrublands. Known generally as "CANDLE WOOD."

Andros — Fresh and Stafford Creeks, *Northrop* 698, 541.

New Providence — near Nassau, *B. & Br.* 677; *Curtiss* 189.

Watlings Island — *Coker* 469.

Exuma — near Rolltown, *B. & M.* 3081.

Fortune Island — *Hitchcock*.

Little Inagua — *N. & T.* 1196.

Inagua — *N. & T.* 1271, 1379,

Chiococca alba (L.) Rusby.

C. racemosa Jacq. Pine barrens, coppices and scrub lands common :
Green Turtle Cay — *Brace* 1484.

Abaco — Marsh Harbor, *Brace* 1713, 1789, Butler Bay, 1522, and Eight
Mile Bay, 1881.

Great Bahama — Barnett's Point, *B. & M.* 2713.

Great Harbor Cay — *B. & M.*

Lignum Vitae Cay — *B. & M.*

Goat Cay — *B. & M.*

Mangrove Cay — *Bryant* 7.

Silver Cay — *Millsp.* 2447.

Rose Island — *B. & M.*

New Providence — near Nassau, *Cooper* 9; *Northrop* 28; *B. & Br.* 230,
434, 656; *Wight* 149.

Eleuthera — Governor's Harbor, *Hitchcock*; Gregory Town, *Coker* 363.

Wide Opening, Cay north of — *B. & M.*

Exuma — *B. & M.*

Cat Island — Port Howe, *Hitchcock*.

Crooked Island — *Hitchcock* (*C. parvifolia*).

Inagua — *N. & T.* 1298.

Chiococca parvifolia Wulls.

In coppices, infrequently collected :

Abaco — Marsh Harbor, *Brace* 1839.

Andros — Deep Creek, *Northrop* 688.

New Providence — near Winton, *B. & Br.* 729; along Soldiers Road, *Coker* 557; near Nassau, *Brace* 80.

Chiococca pinetorum Britton sp. nov.

A vine, creeping or low-climbing sometimes 1.5 m. long, usually shorter. Branches short; leaves ovate, lanceolate or elliptic, small, 2–4.5 cm. wide, varying from acute to obtuse at the apex, narrowed at the base, coriaceous, dark green and shining above, paler and dull beneath, the midvein impressed on the under side, the lateral veins few and obscure, the petiole slender, 2–3 mm. long; racemes few-flowered, shorter than the leaves or about equalling them; pedicels about as long as the calyx, or sometimes a little longer; calyx campanulate 2 mm. long; corolla funnel-form-campanulate, about 5 mm. long, white or cream color, veined with purple; berry white, somewhat compressed, 4–6 mm. in diameter.

Differs from *C. parvifolia* Wulsch. by its smaller darker green leaves, smaller pale corolla and less compressed berries. *C. pinetorum* occurs only, so far as known, on islands inhabited by *Pinus bahamensis* Griseb., and in association with it.

Abaco — opposite Cherokee Settlement, *Brace* 1990.

Great Bahama — Eight Mile Rocks, *B. & M.* 2408; West End, *Brace* 3563.

Andros — near Red Bays, *Northrop* 477.

New Providence — near Nassau, *Northrop* 138 (*C. parvifolia*); *Hitchcock* (*C. parvifolia*); *B. & Br.* 430 **type**; *Cooper* 19, 25; *Britton* 20; *Curtiss* 159 (*C. parvifolia*); *B. & M.* 2103; *Millsp.* 2076, 2238; *Coker* 18, 80, 163, 185.

Scolosanthus bahamensis Britton.

Occasional in coppices:

Andros — at Fresh Creek, *Northrop* 646 (*Scolosanthus* sp.).

New Providence — coppice on East Road near Nassau, *Coker* 138; coppice on Soldiers Road, *Millsp.* 2488; coppice on Village Road, *B. & Br.* 367 **type**.

Strumpfia maritima Jacq.

Borders of brackish swamps and on coastal rocks:

Abaco — opposite Cherokee Settlement, *Brace* 1981.

Great Bahama — *Allen* 33.

Whale Cay — *B. & M.*

Joulter's Cays — *Millsp.* 2300.

Rose Island — *B. & M.* 2138.

New Providence — *Hitchcock*; *Cooper* sine num.; *Coker* 59; *Curtiss* 188; *Brace* 159, 550; southeast shore near Fox Hills, *B. & Br.* 550; Dix Point, *Northrop* 151; Swingate on the shore rocks, *Millsp.* 2248; *Wight* 116.

Eleuthera — Governor's Harbor, *Hitchcock*.

Wide Opening, Cay north of — *B. & M.*

Great Guana Cay — *B. & M.*

Watlings Island — *Hitchcock*.

Crooked Island — *Hitchcock*.

Little Inagua — *N. & T.* 1186, 1250.

Inagua — *N. & T.* 986.

Grand Turk Island — *N. & T.* 3813; *Hjalmarsson*.

***Psychotria bahamensis* Millsp.**

Bull. N. Y. Bot. Gard. 3: 451. In coppices and thickets. Fruits globular, bright orange; flowers white.

Great Bahama — Eight Mile Rocks, *B. & M.* 2529; *Brace* 3675; Barnett's Point, *B. & M.* 2665.

Great Sturrupe Cay — *B. & M.* 3269.

Lignum Vitae Cay — *B. & M.*

Great Harbor Cay — *B. & M.*

South Bimini — *Millsp.* 3478.

Rose Island — *B. & M.* 2134.

New Providence — *B. & Br.* 206 **type**, 208, 662; *Earle* 43, 52; *Britton* 17; *Millsp.* 2483; *E. G. B.* 3359.

Fortune Island — *Hitchcock* (*Myrstiphyllum undatum*). (Reported on Eleuthera in Pl. Baham p. 94, label says Fortune Is.)

(*PSYCHOTRIA LANCEOLATA* Nutt.)

This species reported by Grisebach as having been seen in the so-called Swainson collection, has not yet been re-collected in the Bahamas.)

***Psychotria ligustrifolia* (Northrop)**

Myrstiphyllum ligustrifolium, Northrop, Flora New Prov. & Andros p. 68.

New Providence — near Nassau, *Northrop* 206 **type**; Blue Hills, *Millsp.* 2071; *Wight* 205.

North Cat Cay — *Millsp.* 2342.

***Psychotria pubescens* Sw.**

In coppices and pine barrens:

Abaco — Marsh Harbor, *Brace* 1836.

Great Bahama — Eight Mile Rocks, *Brace* 3700.

Andros — Conch Sound, *Northrop* 585 (*Myrstiphyllum pubescens*).

New Providence — Farringdon Road, *B. & Br.* 204; near Nassau, *Cooper* 78; *Brace* 168.

Cat Island — Port Howe, *Hitchcock* (*Myrstiphyllum undatum*).

Psychotria undata Jacq.

In coppices and pine barrens. The common form in the Bahamas.

Known as "WILD COFFEE."

Abaco — Marsh Harbor, *Brace* 1596, 1613.

Great Bahama — West End, *Brace* 3547.

South Bimini — *Millsp.* 2381.

Andros — Nichols Town, *Northrop* 361 (*Myrstiphyllum undatum*).

Hog Island — *B. & Br.* 347.

Rose Island — *B. & M.* 2116.

New Providence — *Britton* 26, 100, 117, 142; *Coker* 28, 29; *B. & Br.* 179, 347, 437; *Millsp.* 2051, 2083, 2180, 2197, 2236, 2484; *E. G. B.* 3149, 3147; *Brace* 29; *Cooper* 59; *Hitchcock* (*Myrstiphyllum undatum*).

Eleuthera — opposite Spanish Wells and at Tarpum Bay, *Coker* 330, 412.

Great Guana Cay — *B. & M.* 2868.

Exuma — near Georgetown, *B. & M.* 3138.

Long Island — Clarence Harbor, *Coker* 507.

Morinda royoc L.

Scrublands and pine barrens. Known as "RHUBARB" and "WILD MULBERRY."

Abaco — Marsh Harbor, Great Cistern, and Butler Bay, *Brace* 1590, 1763, 1517.

Great Bahama — Eight Mile Rocks, *B. & M.* 2427; West End, *Brace* 3487.

Andros — Conch Sound, *Northrop* 425.

Eleuthera — Gregory Town, *Coker* 379.

Ernodea

This genus is being considered by Dr. Britton.

Hemidiodia ocimifolia (Willd) Schum.

In water holes. Not before collected on the Islands:

New Providence — water hole on Harold Road, *B. & Br.* 532.

Borreria

This genus is being considered by Dr. Britton.

Spermacoce aspera Aubl.

On waste grounds and in sink holes:

Abaco — Eight Mile Bay, *Brace* 1861.

Great Bahama — Eight Mile Rocks, *B. & M.* 2473.

New Providence — Farringdon Road and Fort Charlotte, *B. & Br.* 182, 779; *Brace* 490; *Hitchcock*; *Wight* 32.

Eleuthera — Governor's Harbor, *Hitchcock*.

Cat Island — Port Howe, *Hitchcock*.

Inagua — *N. & T.* 1332, 1377; *Hitchcock*,

Spermacoe tenuior (L.) Lam.

Only to be found in pot or banana holes:

Abaco — Marsh Harbor, *Brace* 1640.

Great Bahama — Eight Mile Rocks, *B. & M.* 2475.

New Providence — South West Point, *Northrop* 319; near Nassau, *Eggers* 1441; *Earle* 18 b; *Curtiss* 18; Blue Hills road, *Millsp.* 2471.

Eleuthera — Governor's Harbor, *Hitchcock*.

Great Guana Cay — *B. & M.* 2898.

Inagua — *N. & T.* 1414; *Hitchcock*.

Galium hispidulum Mx.

Pinelands infrequent. Identical with the species as found in the southeastern United States.

Elbow Cay — *Brace* 1694.

Abaco — Marsh Harbor, *Brace* 1621; Which Point, *Coker* 563.

Great Bahama — Eight Mile Rocks, *B. & M.* 2403.

Andros — Conch Sound, *Northrop* 523 (*Relbunium hypocarpium*, this is a high mountain species of Jamaica).

New Providence — near Harold Road, *B. & Br.* 429; near Nassau, *Brace* 330; Blue Hills, *Millsp.* 2062.

VERBENACEÆ.

Ghinia curassavica (L.)

Ischnia verbenacea D. C. *Verbena curassavica* Linn. *Ghinia verbenacea* Sw.

Waste grounds, doubtless an introduced weed:

New Providence — Farringdon Road, *B. & Br.* 183; near Nassau, *Curtiss* *sine num.*; *Brace* 192, 432.

Lantana bahamensis Britton.

In coppices and scrub lands:

New Providence — near Ft. Montague, *B. & Br.* 174 **type**, 328; near Nassau, *Curtiss* 109; *Coker* 126 (*L. crocea*); *Northrop* 129 (*L. camara*); *E. G. B.* 3441; *Hitchcock* (*L. crocea*); *Millsp.* 2089, 2257; *Wight* 87, 122.

Andros — Conch Sound and Nichols Town, *Northrop* 561, 352 (*L. crocea*); Red Bays, 490 (*L. camara*).

Eleuthera — Governor's Harbor, *Hitchcock* (*L. crocea*).

Exuma — near Georgetown, *B. & M.* 2939.

Cat Island — Port Howe, *Hitchcock* (*L. crocea*).

Lantana balsamifera Britton.

The range of this striking species is as yet rather intermittent. It is found in sand and on whitelands, and is known on Inagua as "MOUJEAN TEA."

Whale Cay — *B. & M.* 2187.

Eleuthera — Tarpum Bay, *Coker* 410 (*L. involucrata*).

Little Inagua — *N. & T.* 1211 **type**, 1198, 1203.

Lantana camara L.

So far known only from the following Islands :

Watlings — *Hitchcock*.

Inagua — Mathew Town, *N. & T.* 1468.

Long Island — Clarence Harbor, *Coker* 513 (*L. crocea*).

(*LANTANA CROCEA* Jacq.

This species, though reported by all writers on Bahamian plants, has not yet been seen from the islands).

Lantana demutata sp. nov.

Frutex glabro valido ramoso cortice rugoso. Rami virgatis ascendentis, foliis petiolatis oblongis vel oblanceolatis ad basin attenuatis, obtusis margine revolutis crenato-dentatis pagina utraque minute et strigose pilosis, supra reticulato-rugosis infra pilis densis nervis crassis prominentis. Inflorescentia axillaribus longe pedunculatis pedunculæ filosis ad apicem inflorescentis capitulæ brevis globosis 5-8-floris bractæ foliosis ovato-rotundatis crassis. calyce paleaceis coronatis irregulariter dentatis margine ciliatis, corolla tubo longis infundibuliformis aliquid curvatis limbo irregulariter lobatis, lobis superiora erectis rotundatis integris, lateralis parvis irregulariter fisso-dentatis inferiora transverse-ovatis reflexis integris ad basi auriculatis. Fructus globosis carnis, pyrena profunde et similiter tuber cibariæ foveolatis.

A spreading, globose-headed, odorless shrub 1.5 to 2 m. high. Leaves 1-1.5 cm. x 6-8 mm., petioles 2 mm., peduncles 2-4 cm. long. Calyx 1 mm., corolla 3.5 mm. The species approximates *L. involucrata*, from which it differs, however, in many characters. No counterpart of the specimens could be found in Hb. Kew or British Museum.

In scrublands:

Exuma — near Georgetown, Feb. 26, 1905, *B. & M.* 2940 **type**.

Lantana involucrata Linn.

Common in scrub and open lands. Known as "WILD SAGE."

Allen's Cay — *Brace* 1534.

Abaco — Cherokee Sound, *Brace* 1940.

Great Bahama — Eight Mile Rocks, *B. & M.* 2444.

Great Sturup Cay — *B. & M.*

Lignum Vitae Cay — *B. & M.*

Little and Great Harbor Cays — *B. & M.*

Frozen Cay — *B. & M.*

Whale Cay — *B. & M.* 2188.

Gun Cay — *Mills* 2306.

South Cat Cay — *Mills* 2422.

Mangrove Cay — *Bryant* 15.

Rose Island — *B. & M.* 2127.

New Providence — near Ft. Montague, *Northrop* 56; *Curtiss sine num.*; *Britton* 51; *Earle* 45; *Coker* 21; *Brace* 70; *Hitchcock*; *Wight* 45.

Eleuthera — Governor's Harbor, *Hitchcock*.

Exuma Chain — on all islands visited except Exuma Island, *B. & M.*;

Cave Cay, *B. & M.* 2821.

Cat Island — Port Howe, *Hitchcock*.

Watlings Island — *Hitchcock*.

Crooked Island — *Hitchcock*.

Fortune Island — *Hitchcock*.

Inagua — Mathew Town, *N. & T.* 890, 933; *Hitchcock*.

Grand Turk Island — *N. & T.* 3763; *Hjalmarsson*.

***Lantana odorata* Linn.**

Doubtless rare on the Bahamian Islands. So far only known by the following specimen :

Green Turtle Cay — *Brace* 1478.

***Lantana ovatifolia* Britton.**

A peculiar large and coarse-leaved species, sprawling among shrubs and Pteridium in the pine lands of :

Great Bahama — at Eight Mile Rocks, *B. & M.* 2430 **type**; *Brace* 3686.

NASHIA gen. nov.

Frutices. Truncus 4-costatus. Cortex roseo-cinereus in laciniae tenuiter exfolientibus. Rami opposito robustis, ramuli ad tuberculae reductus. Foliis ad apicem tuberculae fasciculato vel ad rami oppositi dispositis. Inflorescentia capitellata, rhachi crasso, calyce in annulo reducto corolla tubulo-campanulatis in aestivo valvato lobis 4 aequaliter magnitudo stamina 4 didymamo. Fructus drupaceus, pyrenae in nuculis osseis coherentibus.

I take great pleasure in dedicating this striking genus to Mr. George Valentine Nash, who, in his energetic field work in the Bahamas, South Florida and Haiti, has rediscovered many little known species and amassed very valuable collections toward a flora of these regions.

Nashia inaguensis sp. nov.

Frutex citronellae odoratis. Rami crassis oppositis tereto-quadrangularis cortex roseo-cinereis in laminii longitudinale exfoliatis. Folia brevissime petiolatis ad ramuli, baculatum et lanatum in fasciculi dispositis, ovatis vel obovatis vel spathulatis vel ligulatis, obtusis ad basin attenuatis pagina supra valde reticulo-rugosa strigosis infra strigose-tomentosis margine ad siccam revolutis integris. Inflorescentia sessilis, ad media ramulorum dispositis; flores 6–8 albis cum tubo aurentio-luteus, bractae rutelliformis apiculatis ciliatis floribus longioribus. Calyce minutis annulatis dentatis ciliatis ad rachi persistentibus, corolla campanulatis lobis aequantis duo exterior magnis ad basin auriculatis apice emarginatis duo interior integris, stigma peltatis post anthesis sagittatis, stylis indusis, filmenta abbreviata tubo tertia demittere insertis. Drupa pyriformis extus carnosus, putamen duris laevis.

A stout, straggling, endemic bush 1.5 to 2 m. high, with strong, stout branches issuing at right angles. Trunk 4–5 dm. long 3–4 cm. in diameter, leaves 5–8 x 3–5 mm. ; bracts 2.5 x 2 mm. Drupes fleshy 3 x 4 mm. ; nutlets 2.5 x 3.5 mm. So far only known from:

Inagua — in scrublands back of Mathew Town, Oct. 13, 1904, *Nash & Taylor 1006 type, 1454*. A sheet also found in the *Hitchcock* collection marked "Lantana sp. from Inagua Dec. 3, 1890". Mr. *Brace* sends a small specimen of the species from a garden in Nassau "where it was planted from Inagua." It is well known to the natives, who use the leaves in decoction as a febrifuge known as "MOUJEAN TEA."

Lippia nodiflora (L.) Michx.

Open sandy places :

Great Bahama — clearing at Eight Mile Rocks, *B. & M. 2368*; West End, *Brace 3575*.

North Cat Cay — *Millsp. 2337*

Andros — Red Bays, *Northrop 481*.

Hog Island — *Northrop 250*.

New Providence — near Nassau, *Brace 265*; near Grants Town, *Coker 280*; at Lake Cunningham, *Britton 105, Millsp. 2185; Wight 162*.

Eleuthera — Governor's Harbor, *Hitchcock*.

Exuma — near Georgetown, *B. & M.*

Cat Island — Port Howe, *Hitchcock*.

Inagua — *N. & T. 1091, 1276, 1465*.

Lippia reptans H. B. K.

Shores of the sea or brackish marshes :

New Providence — near Nassau, *Northrop 114, (L. canescens); Coker 98, (L. canescens); Hitchcock (L. nodiflora)*.

Inagua — Lower Savannah near Blakeville, *N. & T.* 1131; *Hitchcock* (*L. nodiflora*).

Lippia stoechadifolia (L.) H. B. K.

Borders of fresh water pools or marshes and in water holes :

New Providence — near Clifton, *B. & Br.* 745.

Exuma — in a water hole in palmetto lands back of Georgetown, *B. & M.* 2964.

Cat Island — near Arthur's Town, *Coker* 430.

Bouchea prismatica (L.) Ktze.

B. Ehrenbergii Cham. In waste grounds :

New Providence — at Fort Charlotte, *B & Br.* 782; *Brace* 371.

Valerianodes fruticosa sp. nov.

Stachytarpheta et *Stachytarpha* et *Abena*. § Pachyurae: Frutex. Caule erectis duro-lignosis superne ramosis, rami teretis glabriusculis, ramuli subtetragonis; foliis ovato-lanceolatis basi contracta in petiolum spurium decurrentibus, apice acutis margine plus minusve patentidentatus, pagina supra nitidis subtus pallidulo-viridis glabriusculis, nervi medius alboviridis, utrinque minute nigro-punctatis. Spices terminalibus cylindraceus brevis tortis, glabris, floribus arrectis demum rhachi-incrassatae immersis bracteis ovato-lanceolatis calyce subcarinatis margine integris non scariosis, calyce compressis bifidis aequantibus. Fructus obpyriformis manifeste apiculatis, pyrenae jugis dorsalisibus et citra apicem rugis evidentis anastomosantibus.

A shrub 1-2 m. high with a strong, erect stem one-half to three-fourths of a meter long, with white bark and dense, pinkish wood (in our specimen showing 12 annular rings). Leaves 4-7 cm. long by 1-2 cm. broad. Spikes 5-10 cm. long, 4 mm. thick. Pyrenae 6 mm. long, by 2 mm. broad, full twice the dimensions of those of *V. jamaicensis* to which the species appears nearest related. From that species it also strongly differs in its shrub character, short spikes, and punctate glabrous leaves. In scrublands, infrequent :

Great Bahama — West End, *Brace* 3574.

Rose Island — Jan. 27, 1905, *Britton & Millspaugh* 2125 type.

Wide Opening, Cay north of — from crevices in rocks, *B. & M.* 2771.

Valerianodes jamaicensis (L.) Medic.

In scrublands and open waste grounds:

Great Bahama — Eight Mile Rocks, *B. & M.* 2479.

Great Sturup Cay — *B. & M.*

Goat Cay — *B. & M.* 2300.

Lignum Vitae Cay — *B. & M.*

Mangrove Cay — *Bryant* 18.

Rose Island — *B. & M.*

New Providence—near Nassau, *Earle* 8; *Brace* 320; *Northrop* 19 (*Abena*); *Coker* 113. (*Abena*); *Millsp.* 2465; *Hitchcock* (*Abena*); *Wight* 85.

Eleuthera — Governor's Harbor, *Hitchcock* (*Abena*).

Great Guana Cay — *B. & M.*

Exuma — *B. & M.*

Fortune Island — *Hitchcock* (*Abena*).

Inagua — *N. & T.* 1068; *Hitchcock* (*Abena*).

Grand Turk Island — *N. & T.* 3870.

***Priva lappulacea* (L.) Pers.**

P. echinata Juss. Waste places near habitations, doubtless an introduced weed.

Green Turtle Cay — *Brace* 1482.

New Providence — near Nassau, *Curtiss* 24; *B. & Br.* 534, 655; *Earle* 12; *Coker* 268; *Wight* 40.

Eleuthera — Governor's Harbor, *Hitchcock*.

Inagua — *Hitchcock*.

Citharexylon** *sp.

A shrub with obovate short-acuminate acute or emarginate glabrous leaves and paniculate inflorescence.

Inagua — Salt Pond Hill, *N. & T.* 935.

The specimens are too incomplete for satisfactory specific determination.

***Citharexylon Berterii* Spreng.(?)**

In coppices:

Andros — Fresh Creek, *Northrop* 608.

The specimen under this number in herb. Field Col. Museum possesses either teratological or very immature flowers, hence this determination is rendered somewhat doubtful.

***Citharexylon caudatum* L.**

In a coastal coppice:

Andros — Couch Sound, *Northrop* 571 (*C. lucidum*) Cham. & Schlecht.

(*CITHAREXYLON QUADRANGULARE* Jacq.

Prof. Coker's specimens referred to this species in his "Vegetation. The Bahama Islands" have not been seen by the writer.)

*This consideration of the genus *Citharexylon* is by Dr. J. M. Greenman.

Citharexylon villosum Jacq.

In coastal coppices:

Abaco — Eight Mile Bay, *Brace* 1882.

New Providence — at Southwest Bay and along Farringdon Road, *B. & Br.* 499, 233; near Nassau, *Hitchcock*.

Cat Island — Port Howe, *Hitchcock*.

This species shows considerable variation in leaf outline, degree of pubescence and size and color of the pyrenae. Mr. Combs' No. 468 from Cieneguita, Cuba; *Mr. Brace's* Abaco No. 1882; and Mr. Nash's 991 from Hayti are broad leaved forms. These pass through *Britton & Brace's* New Providence 499 to narrower leaved forms representing *C. bahamense* Millsp. (Bull. N. Y. Bot. Gard. 3. 450.) To this form are referable, *Britton & Brace's* New Providence 233 type; *Hitchcock's* New Providence and Cat Island specimens; and Paulsen's St. Thomas No. 156, the last specimen cited was distributed as *C. cinereum* L.

Duranta repens Linn.

D. Plumieri Jacq. Plentiful where found at all. Generally in the pine barrens and open scrublands. Apparently native though possibly introduced and now spreading.

Abaco — Marsh Harbor, *Brace* 1740.

Great Bahama — pine barrens at Eight Mile Rocks, *B. & M.* 2434; *Brace* 3687.

New Providence — near Nassau, *Northrop* 39; *Cooper* 33; *Curtiss* 1; *Coker* 61; *Brace* 22; *Millsp.* 2050; *Hitchcock*; *Wight* 4, 49.

Eleuthera — Governor's Harbor, *Hitchcock*.

Callicarpa fulva Rich.

In scrub land and coppice:

Cat Island — *Hitchcock*.

Mr. Hitchcock's plant, while having (on account of its greatly reduced leaves) a quite distinct general appearance of difference from *C. fulva* (as well represented by *C. Wright's* 1357 Monte Verde, Cuba, May 30, 1859,) nevertheless has no other characters of differentiation. It is fairly well connected in leaf size and form with the Wright plant through Mr. Wright's other 1357 collected at the base of Farallones, Sept. 29, 1860, which is in the same fruiting stage as the Hitchcock plant. Of this Farallones plant Mr. Wright says, "A slender bush 6 to 10 feet high: in thick woods."

Sauvalle (Fl. Cub. 113) considers this species synonymous with the Jamaican *C. ferruginea* Swartz, this however is not the case, the differences are broad and evident.

Grisebach in his *Catalogus Plantarum Cubensium* includes under *C. fulva* Rich. Mr. Wright's Majanabajo, Cuba 3173 with the remark "forma foliis lanceolatis." The sheet of this number in Herb. Gray, Cambridge (one of Mr. Wright's original series), proves to be an entirely different species. This may be designated as follows:

CALLICARPA LANCIFOLIA sp. nov.

Ramulis cum cymis petiolis foliorumque tomento stellari ferrugineo canescentibus, foliis crassis lanceolatis apice acutis ad basin attenuatis brevipetiolatis, supra nigra, ad sicca valde reticulato-rugosis et minute resinosis, infra lutea dense stellato-canescens, margine leviter et irregulariter crenato-denticulosis. Cymis axillaribus folio medio brevioribus ramoso-dichotomis, calyce campanulatis dense resinosis granulatis cum quidam pilae stellatis et perpauci hirtis, limbo in lobis triangularis valde divisus.

Eastern Cuba, C. Wright No. 3173 in Herb. Gray, Cambridge, U. S. A. **type.** Leaves 5 cm. long, 1 cm. broad, strongly bicolor; they differ from those of *C. fulva* Rich. in their shape, the far less prominence of the veins beneath, and in not having the pubescence thereon so thick and strongly tufted.

Petitia domingensis Jacq.

P. Poeppigii Schl. In pine barrens and scrublands:

Abaco — Marsh Harbor, *Brace* 1630, 1721.

Great Bahama — Eight Mile Rocks, *B. & M.* 2593.

Andros — Nichols Town, *Northrop* 358.

Mangrove Cay — *Coker* 224.

New Providence — near Nassau, *Cooper* 34; *Eggers* 4201; *Britton* 27; *Millsp.* 2090; *Wight* 206; *Coker* 162, 556; *Curtiss* 136; *Brace* 55; near

Lake Cunningham, *B. & Br.* 604; Lake Killarney, *E. G. B.* 3288.

Eleuthera — Governor's Harbor, *Hitchcock* (*P. Poeppigii*).

Cat Island — Arthur's Town, *Coker* 423 (*P. Poeppigii*).

PSEUDOCARPIDIUM gen nov.

Arbusculae aut frutices folia simplicia opposita integra aut crenato-dentata, flores paniculata. Calyces campanulata 5-dentata aequalia. Corolla tubulosa 5-fida. Stamina 4, didynama exserta, stigma bifida. Drupa 2-pyrenensis, pyrene osseis 2-loculus monospermis in drupa centralia dispositis.

The name from *ψευδής* false, *καρπίδιον* carpid, in reference to the four carpid-like prominences on the fruit; these apparent cocci are simply fleshy masses, Richard's fig. 3, pl. 64 in *La Sagra* to the contrary, notwithstanding. The known species are as follows:

(PSEUDOCARPIDIUM ILICIFOLIUM (Rich.)

Vitex ilicifolia Rich. in La Sagra Hist. Cuba Bot. 2; 148).

(PSEUDOCARPIDIUM AVICENNIOIDES (Rich.)

Vitex avicennioides Rich. in La Sagra Hist. Cuba Bot. 2; 149).

Pseudocarpidium Wrightii sp. nov.

Frutex a basi ramosissimis, rami glabris crassis, cortice argenteo-albis. Foliis coriaceis late obovatis vel orbiculatis vel ovato-lanceolatis apice obtusis, basi subcordatis, margine superante subsinuoso-dentatis spinosis vel tantummodo spinosis, pagina supra nitidis valde reticulovenosis, infra cum grana resinosa pulverulentibus ad venae minutissime pubescentibus. Inflorescentia paniculatis, paniculae laxis longe et filisae pedunculatis folio plus longioribus supra dichotomis, tota minutissime resinoso-pulverulentibus, bractae linearis. Calyce campanulatis obtuse 5-deltoideo-dentatis, corolla bilabiatis limbo explanatis, labia superiore erectis integris apice deltoideis, labia inferiore in dentibus quatuor productis, dentes lateralia ovatis acutis, dentes inferiore deltoideis. Drupa depressa producte in lobis quatuor ad centro bipyrenis, pyrenae osseis hemisphaeris dorso bi-costatis ventro profunde sulcatis.

A tall, lax and woody shrub with opposite, ascending branches and shining holly-like leaves. Leaves 2.5–3.5 x 1.3–2 cm., petioles about 4 mm., the spines confined to the upper half of the leaf. Peduncles 3.5–5 cm.

In coppices and scrublands:

Mangrove Cay — *Bryant* 2.

Andros — at Fresh Creek, *Northrop* June 6, 1890, no. 625, type; *Wight* 261.

The specimens agree with those of Combs, Calicita, Cuba 239, and Shafer, Madruga, Cuba 74. There are two plants of Wright, Cuba orientali bearing his number 421, one being this species and the other *P. ilicifolium*, evidently a case of attempted matching of plants before distribution, unfortunately therefore I am compelled to take the later gathered Bahamian plant for the type.

The foliage form and shape is very variable, never however even approaching that of *P. ilicifolium*, in which the margins are crenate and spinous throughout, while those of *P. Wrightii* are simply spinous and that only on the upper half of the leaf. Both sides of the leaf in *P. ilicifolium* are pubescent, while in *P. Wrightii* the upper side is shining and strongly reticulate, while the lower surface is resinous pulverulent.

Clerodendron aculeatum (L.) Griseb.

Rare in collections. Only known by:

New Providence — "Escaped" near Nassau, *Northrop* 296 (*Ovieda*).

Inagua — *N. & T.* 1471.

Ovieda fragrans (Willd.) Hitch.

Escaped from cultivation and becoming fully naturalized along roadways on:

New Providence — Blue Hills Road, *Northrop* 328; roadside at Lake Cunningham, *Mills* 2221; Nassau, cult. *Hitchcock*.

Avicennia nitida Jacq.

In salt and brackish borders, nowhere common:

Abaco — Cherokee Sound, *Brace* 1849.

Great Bahama — Barnett's Point, *B.* & *M.*

Great Harbor Cay — In the large savannah, *B.* & *M.*

Frozen Cay — *B.* & *M.*

South Bimini — at the south end, *Mills* 2396.

New Providence — collected by M. A. Howe far out from the south side beach, *Mills* 2482.

Andros — at Mastic Point, *Northrop* 593.

Ship Channel Cay — *B.* & *M.*

Great Guana Cay — *B.* & *M.*

Exuma — in the tide pond at Georgetown, *B.* & *M.*

Fortune Island — *Hitchcock*.

Inagua — *N.* & *T.* 1326.

Grand Turk Island — *N.* & *T.* 3823.

SOLANACEÆ.

Solanum didymacanthum sp. nov.

§ *Leptostemonum* — *Graciliflora*. Rami pilis stellatis pilosi, aculeatissimis, foliis minutis integris utrinque stellati-pilosis. Caulis fruticosus ramosissimus 1–1.5 m. alt. Rami nigro-viridibus subtomentosis, pilis stellaris, aculeatissimis, aculeis acicularibus ad nodi geminibus foliis bis vel tres longioribus, prae-gracilis leviter recurvis divaricato-horizontalibus rubro-brunneus basi incrassatis, medio centim. longis. Foliis fasciculatis omnes minutis, petiolatis, oblongis vel obovatis integris 2–2.5 mm. longis utrinque stellati-pilosis, basi acutis, apice acutis vel obtusis. Inflorescentia solitaria terminalia. Pedunculi 3 mm. longi, uniflori, stellato-pilosi inermi; calyce stellato-pilosis inermis 5-dentatis, laciniis lanceolato-triangularis acutis, 1 mm. longis; corolla alba (?) 5-partita, laciniis liguliformis stellato-pilosis, 8 mm. longis. Stamina 5 corolla nonnihil breviora. Ovarium globosum minimum; stylus filiformis staminibus longior. Bacca ignota.

A densely virgate-branching, spiny shrub, with minute leaves, the uppermost of which often bear a single spine central upon the midrib of the same length and character as those of the nodes; slightly recurved, needle-like spines full twice or thrice the length of the

leaves, issuing in twins from each node; and single, comparatively large flowers terminal on each branchlet.

Exuma — in the open scrubland near the west end of Hayne's road, *B.* & *M.* 3037 **type**.

Cat Island — at Port Howe, *Hitchcock*. This specimen has ovate-spatulate leaves as long as or nearly the length of the spines or sometimes exceeding them.