STAND DYNAMICS SYSTEM FIELD REMEASUREMENT MANUAL

MARCH 2005

**Public Lands and Forests Division
Forest Management Branch
8th Fl. 9920 – 108 Street
Edmonton, AB
T5K 2M4**

Phone: (780) 427 – 8474
Or visit the website: <http://www3.gov.ab.ca/srd/forests/psp>

Digitized by the Internet Archive
in 2016

TABLE OF CONTENTS

1.0	INTRODUCTION.....	1
2.0	FIELD PROCEDURES.....	1
2.1	PLOT ESTABLISHMENT	1
2.2	PLOT MAINTENANCE	1
2.3	MAINTENANCE REPORT	2
3.0	REMEASUREMENT PROCEDURE.....	3
3.1	REGEN PLOTS (RECORD TYPE 6 – (REFER TO FIGURE 4.23)	5
3.2	REGENERATION HEIGHT CLASS RECORD (CSTM 100)	7
3.3	SAPLING PLOT (RECORD TYPE 7)	8
3.4	TREE PLOT (RECORD TYPE 8)	9
3.4A	CROWN WIDTH MEASUREMENT.....	9
3.5	PLOT PHOTOGRAPHY	10
3.6	VEGETATION DESCRIPTION	10
3.7	PLOT VEGETATION ASSESSMENT.....	10
3.8	PLOT RETREATMENT REPORT (RECORD TYPE 9).....	11
3.9	PLOT AGING	12
4.0	ALLOWABLE ERRORS.....	14
4.1	REGENERATION PLOT MEASUREMENTS	14
4.2	SAPLING PLOT MEASUREMENTS	14
4.3	VEGETATION MEASUREMENTS	15
4.3A	PLOT RETREATMENT MEASUREMENTS.....	15
4.4	PLOT REJECTION	15
4.5	DATA ENTRY ON HEADER AND TALLY SHEETS.....	16
4.6	STAND DYNAMICS GENERAL INFORAMTION (CSTM 100 AND CSTM 101) – SEE APPENDIX # 17.....	16
4.7	SUGGESTED EQUIPMENT AND SUPPLIES - SEE APPENDIX # 1.....	16
4.8	TREE SPECIES CODE – SEE APPENDIX #12.	16
4.9	TREE HEIGHT MEASUREMENT USING A CLINOMETER – SEE APPENDIX #14.	16
4.10	HEIGHT TO LIVE CROWN.....	16
4.11	SLOPE CORRECTION FACTORS AND TABLES – SEE APPENDIX #5.	17
4.12	CONDITION CODES – SEE MASTER CONDITION CODE LIST.....	17
4.13	MISTLETOE RATING SYSTME – SEE FIGURES 11 AND 12.	17
4.14	LEGAL SURVEY OF ALBERTA – SEE APPENDIX # 8	17
4.15	DETERMINING BREAST HEIGHT – SEE APPENDIX #14.....	17
4.16	ROUNDING OFF – SEE APPENDIX 9-H.....	17
4.17	PROBLEMS WITH RECORDING DATA – SEE APPENDIX #15 AND 16.	17
4.18	CROWN CLASS.....	17
4.19	STEM MAPPING.....	17
4.20	SEED VS. SUCKER – ORIGIN ON POPULUS SP. – SEE FIGURE 4.32.	17

1.0 INTRODUCTION

As Alberta's forest industry continues to grow and expand so does the requirement for timber, and the demand for better forest management. There is a need for an increased awareness and understanding of this resource. Two gaps in our knowledge centre around the growth and dynamics of stands younger than 20 years of age, and the dynamics of stands established after harvesting.

To fill these gaps, the Public Lands and Forests Division, established a series of permanent sample plots throughout areas of Alberta where harvesting occurs. This is a long-term project monitoring stand growth and dynamics from seedling to mature tree. Due to the various phases of growth that will be monitored, it was necessary to establish nested sample plots. At the regeneration stage, the four regen plots will fall within a larger area that is monitored in the sapling stage. This plot area falls within the bounds of a standardized 1/10 ha.

Three other areas also under study are:

- 1) The regeneration, growth, and development of new stands after pine beetle infestation, estimating the rate of deterioration and breakdown of standing dead trees infested with the mountain pine beetle.
- 2) The effect of herbicide spraying in the MOF (Maintain our Forests) cut blocks and
- 3) stand growth and regeneration on the Lodgepole well blowout site.

2.0 FIELD PROCEDURES

The field procedures outlined here are compatible with procedures used for remeasuring the standard Forest Management mature PSP's.

2.1 PLOT ESTABLISHMENT

The plot establishment procedure is not included in this manual. The measurements of the plot (Figure 2.1) can be used to replace any missing posts. For more information see the STAND DYNAMICS FOLLOWING HARVESTING FIELD MANUAL for establishment procedure.

2.2 PLOT MAINTENANCE

The following items are checked:

- 1) Evaluate the access (see Master Appendix 7)
- 2) Rate the overall condition of the plot and buffer, noting any damages and their location on the plot maintenance sheet (Figure 2.10).

If the trees are too small to paint, a 200 m x 200 m buffer should be clearly marked with blue PSP flagging. Contact Forest Management Branch, Resource Analysis Centre for this flagging.

Do not flag into standing mature timber or across seismic lines, roads, etc. that are less than 75 m away from the tree plot. Use them as a boundary and record the appropriate changes on the maintenance report, Figure 4.22.

- 3) Check the condition and tags of all the posts, centre and corners. Replace and retag where necessary. Paint all corner, plot centre and regen posts blue. Mark corner posts with orange and blue geo-flagging. Plot centre is flagged blue, orange and yellow. Sapling aluminum angles are flagged orange.
- 4) Rate the overall condition of the plot and buffer noting any damages and their location on the plot maintenance sheet (Figure 4.22) and Plot Retreatment Report (Figure 4.29).
- 5) Measure all regen, sapling, and tree plot sides. Record sizes on maintenance tally sheet. Remeasure only if you had to re-establish posts, otherwise record previous measurements.
- 6) The tie-point for each plot must be confirmed. This includes checking the distance and azimuth from plot centre to the tie-point. The previous azimuth and distance was to northwest corner, redo to plot centre, if necessary. Keep in mind the distances are horizontal and must be adjusted for slopes exceeding 10%. Whenever possible, a second tie-point should be established in the event that the original tie-point is destroyed. Suitable tie-points include definite bends in the road, stream crossings such as bridges and culverts, or any other permanent land features. A topofil or survey chain must be used to measure the distance between tie points. A vehicle odometer is not accurate by our standards therefore it is not acceptable for measuring distance.
- 7) Ensure the North arrow is correctly located on the maintenance tally sheet.
- 8) Tie-points will be taken to plot centre from now on. Redo maintenance map if they were taken to northwest corner.
- 9) Complete check off list.

Planted trees in the NE, SW and SE corners have their main stem painted blue. Re-paint at time of remeasurement.

2.3 MAINTENANCE REPORT

The information recorded on the maintenance report is documented on the Header Sheet (TM 267). Some of this information is subjective, and requires good judgement, as budget requirements, planning of field work, and other office decisions are based on this information. The PSP maintenance sheet (Figure 4.22) is located on the reverse side of the PSP Header Sheet, TM 267.

The legal location of the group is confirmed once the tie-point information and associated map work is completed. An SDS plot can conceivably occupy up to four sections and legal subdivisions.

Access is extremely important for planning purposes for both remeasurement and maintenance crews. Budget estimates rely heavily on the amount of time it takes to travel to a plot and what type of transportation is required (i.e. 4-wheel drive vehicle, all terrain vehicle, or helicopter). Access is assessed using the following codes (see Master Appendix 7 for description).

- 1 - All weather road
- 2 - Dry weather road
- 3 - Deteriorating road
- 4 - All terrain vehicles only
- 5 - Helicopter access only
- 6 - Unknown (office use only)

Plot and buffer damage to the plot is assessed using the following codes:

- 1 - No damage
- 2 - Natural damage
- 3 - Manmade damage
- 4 - Natural and manmade damage
- 5 - Closed
- 6 - Over Damage inside buffer greater than 20 m from the plot

Code 6 is used to indicate buffer damage only.

- 7 - If there is seismic damage to the plot or buffer that was not recorded on the previous maintenance report, attempt to determine the year the damage was done, and indicate on the maintenance report exactly where the damage is.
- 8 - The lengths and azimuths of the buffer must be recorded on the map.
- 9 - Make general comments on planted trees, ingress etc. on maintenance sheet. Eg. Planted still in bare soil and chlorotic - probably from nutrient deficiency.

3.0 REMEASUREMENT PROCEDURE

The following procedures are for remeasurement and are to be used whether data is collected on tally sheets or on field computer program (produced February, 1998 - see field computer instructions).

If trees decrease in height a condition code must be given. If trees over 1.3m in height, the DBH is required.

If a tree height grows only 2 to 3 cm in 2 or 3 years then something is wrong. Give trees an appropriate condition code.

Make sure you $\sqrt{\sqrt{}}$ HT and $\sqrt{\sqrt{}}$ DBH (in comments) if they shrink or stay the same size.

DBH's have to remain the same or increase otherwise use appropriate condition code.

Each Regen plot should be separated by a line and/or on a new sheet. Checklist must be completed **BEFORE** leaving plot.

All trees that are added in need to **HAVE** an Age and ORIGIN; if Birth year is > than harvest year Origin = "I", if Birth year < harvest year then origin = "A", planted (use guideline sheet provided) "B" or "C" is for. Write birth years in Age column. Place all other comments in the comments column. When trees die (Code 25) you **MUST** give a cause. If a tree dies it must be given code 25 and cause of death. Look for previous damage codes in comment or examine dead tree for clues. If the tree is >9.1cm DBH and tree is dead and down (CC1=61), can have a CC2 if necessary to say what put the tree down on the ground.

Write proper code in condition codes to verify that the tree has died. A missing (15) code can only be used following a dead code (25). Look carefully and try to match previous height to find the tree. All trees over 1.1cm DBH and >1.3m tall require stem mapping.

√ azimuth and distance that are obviously outside the proper range. Saplings and planted trees should only be found between 270° and 360°. Trees noted as out of range on conflict sheet, verify Distance and Bearing. Visually verify "IN" or "OUT" if these trees are okay and tree is in plot **TAKE THE BEARING ON THE CLOSEST LEG'S** (need 2 bearings) and record in comments section.

Be careful when using codes higher than 51. Many have lower size limits.

Use caution before changing any species. This is particularly important with Sw and Sb. If in doubt **DO NOT** change the species.

Cmp Hgt-Type (Competition Height and Type) - This is a two-digit code describing the height and type of competition affecting individual seedlings/saplings.

Code - Cmp Hgt

- 0 - Little or no competing vegetation present.
- 1 - Competing vegetation below seedling/sapling height.
- 2 - Competing vegetation equal to seedling/sapling height.
- 3 - Competing vegetation above seedling/sapling height.
- 4 - Microsite shading (by dead or inorganic materials).
- 5 - Crushing - seeding in close contact with organic/inorganic debris such that physical damage may occur. (Remember to replace debris to minimize the remeasurement disturbance).

Code - Cmp Typ

- 0 - No competition
- 1 - Herbs, forbes, etc.
- 2 - Grasses, sedges, etc.
- 3 - Shrubs (non-acceptable species eg. willows, roses)
- 4 - Trees (acceptable species eg. Fir, Birch, Aspen)

Competition should be tallied only if the canopy area of competing vegetation overlaps approximately 40% of the seedling/sapling canopy area. The canopy area refers to the area directly above and below the leaves of the vegetation. If there is a number of different types of competition, the type that occupies or intercepts the greatest canopy area of the seedling/sapling should be tallied. This dominant competition type may occupy less than 40% of a seedling/sapling canopy area (approximately 30%) if combined with other vegetation type. The height of the competition is evaluated by distribution of competing leaf surface area relative to the height of the seedling/sapling. The two factors of height and density of competing leaves must be weighted together. It was decided that height would have more weight than density of leaf surface area when overtopping occurs. Significant overtopping is roughly defined as having at least 20% of the competing leaf surface area above the height of the seedling/sapling.

Competition will be tallied as above if the seedling/sapling is significantly overtopped by competing vegetation. Competition is to be tallied as equal to if the competition is approximately the same height as the seedling/sapling. There can be some overtopping but not a significant amount (less than 20% leaf surface area). The densest competing leaf surface area should be generally the same height as the seedling/sapling leaf surface area. Competition will be tallied as below if the height of the competing vegetation is less than the seedling/sapling. There must be an insignificant amount of overtopping and the densest competing leaf surface area should be below the seedling/sapling's densest leaf surface area.

There is no room on SDS tally sheets for this 2 digit code, so record it in comments area.

Field computer program requires these 2 variables.

3.1 REGEN PLOTS (RECORD TYPE 6 – (refer to Figure 4.23))

There are 4 circular regen plots located in the northwest corner of the tree plot. All regen is tagged up to a maximum of 50 per plot. If there are more than 50 regen, a height class table is used (see section 3.2). Be careful to number new trees up to 50 with a new number. If 50 trees have been tagged in past and a new tree that used to be counted on the tree tally form grows into 1.3m (doesn't need DBH >1.1cm), number this tree to the next number eg. Plot #1, Tree # 1051. It is very important that you write the correct regen plot number next to these trees.

All pinned and tagged seedlings must be numbered. Numbering starts in plot one at the point closest to the NW corner and continues sequentially in a clockwise direction. Numbering in plot two starts at the point closest to plot one and continues in a clockwise direction, numbering in plot three starts at the point closest to plot two and numbering in plot four starts at the point closest to plot three. When numbering ingress the numbering continues where it was left off from the last measurement. The numbering system is as follows:

Plot one	1001-1050
Plot two	1051-1100
Plot three	1101-1150
Plot four	1151-1200
Plot five (planted)	1201-

On all planted trees greater than 60 cm leave pin in place and put new tag on lateral branch and paint lateral branch blue.

All planted trees must be stem mapped and given crown classes, even those in regeneration plots.

At establishment all seedlings are pinned and numbered to a maximum of fifty. If there are less than fifty seedlings in the plot at establishment then only the amount present is pinned and tagged then when the plot is remeasured all ingress is tagged and numbered until a maximum of fifty in total is tagged and numbered, for example:

Establishment in 1987 - 27 seedlings are pinned and tagged numbers used are 1001-1027

Remeasurement in 1989 - 2 seedlings have died and 14 ingress are found. All of the ingress is pinned and tagged bringing the total to 39 live seedlings and 2 dead = 41 total (last #1041).

Remeasurement in 1991 - 4 more seedlings have died and 17 ingress are found. 9 of the ingress are pinned and tagged bringing the total number tagged to 50. The remaining 8 ingress are measured and recorded on the height class tally sheet. The total number of seedlings is now 44 live tagged seedlings, and 6 dead = 50 tagged, and 8 more recorded on the height class tally sheet (Figure 4.27). Numbers used 1001-1050.

Now that all of the numbers available in regen plot one have been used, any more ingress that occurs in future remeasurements will be recorded into the height class tallies. (CSTM 100). Refer to Section 4.24. **DO NOT DUPLICATE OR RE-USE PREVIOUSLY ASSIGNED NUMBERS.**

There should be no trees with 2000 numbers (sapling) in the 4 regen plots. Call supervisor if found. **DO NOT** change numbers to sapling numbers if a tree grows into a sapling. Keep original number.

The height of all pinned, tagged, and numbered seedlings is measured to the nearest cm (excluding current years growth) and recorded on the tally sheet (CSTM 101). Species and any condition codes (refer to for the Master Tree Condition Codes) present are also recorded, crown classes are not recorded until tree's DBH becomes > 1.1cm. If code 15 is recorded for the previous measurement, look for tree again and if found but dead, try to decide what killed it. i.e. 25 18. If unable to find it or tell what killed the tree use code 25(dead), 37(unknown). If tree is found alive, record previous missed height in comments section.

If a seedling grows to a sapling (greater than 1.3 m in height and 1.1 cm DBH) the pin is pulled and the tag and number is put on the sapling. This sapling has the same number and retains its place in the regen plot tally. It is stem mapped from the main plot centre (azimuth and distance recorded) and the height, diameter, crown class and condition codes are recorded. Diameters are recorded to the nearest millimetre. Mark DBH with blue paint. It is still recorded as record type 6.

If the area has been planted (paint lateral branch blue), then all planted seedlings within the sapling plot were pinned, tagged, and numbered at establishment. At remeasurement, current height and any condition codes for the planted seedlings are recorded on the tally sheet (CSTM

100) under regen plot five. Planted trees are stem mapped regardless of size and crown class recorded. All planted trees in the NE, SW and SE corners have had their main stem painted blue. Repaint these trees at time of measurement. If establishing a plot on a new cutblock that has been planted, pin the planted trees in the NE, SW and SE with "blue" painted wire pigtailed. At each remeasurement, these pins should be repainted "blue". Maintain this procedure, until the planted trees DBH >9.1 cm and then they will be tagged as a 3000 tree number.

Paint all around the main stem, up high with "blue" paint to make them more visible.

Aspen and pine are shade intolerant and may die if they were overtopped by other trees. Spruce are shade tolerant. Be aware of this concept when damage codes are considered.

If a seedling dies, the pin is pulled and the condition code 25 (dead) is entered as the 1st code and you have to use the 2nd condition code to tell what the tree died from (3rd code can be used if necessary). You can pull the tree out of the ground to look for armillaria, flooding, etc. Species **HAS** to be recorded. If a pin is missing and the seedling to which it was assigned cannot be located, using historical height and the numbering sequence, then the condition code 25 for dead is used with a condition code of 15 for missing as the second code. These seedling numbers are not reused.

If a pin will not stay firmly in the ground due to bedrock, etc., loosely tie the tag to a lateral branch of the tree. At **NO** time should a tree be tagged on a main stem.

Starting in the 1999 field season, root collar diameter (RCD) was collected on regeneration or planted trees that were less than 130cm in height. If the plot is measured on paper, this variable (RCD) will have to be recorded in the comments section. The office will enter it into the electronic data file later. Field computer programs automatically prompts the surveyor for the RCD if the tree is less than 130cm in height.

3.2 REGENERATION HEIGHT CLASS RECORD (CSTM 100)

This is a record of untagged live seedlings, filled out if the number of tagged seedlings on a regeneration plot exceeds 50. The number of seedlings by species, in each height class, for each regeneration plot is recorded on form CSTM 100 (Figure 4.27). In the header line, **Record type is 61**, plot type is left blank and each species within each plot is recorded on a separate line.

Each untagged seedling is to have a pin placed beside it. The reason for pinning all non-tagged seedlings during remeasurement (when seedlings total more than fifty) is to ensure that all seedlings are counted. When you measure and record the height class and species you can pull the pin, measure the next seedling and pull the pin, and continue in this manner until all of the seedlings are measured and all the pins without tags are pulled. This avoids missing or double measuring any seedlings.

This form **MUST** be filled out for each Regen Plot. Record No Tally in first column (0-10 cm) and first row if no trees found on regen plots with 50 tagged trees or if plot has less than 50 trees.

Dot tally initially and then write number of trees once completed. Right justify and zero fill to left.

DOT TALLY

Regen Plot 5

Planted trees must have azimuth and distance recorded even if they are less than 1.1cm DBH and/or less than 1.3m in height.

ALL planted trees should get a crown class.

If previous species is wrong put $\sqrt{\vee}$ species in comments. If DBH or HT shrink $\sqrt{\vee}$ check and put in comments.

3.3 SAPLING PLOT (RECORD TYPE 7)

Any stems within the sapling plot over 1.3 m in height and 1.1 cm DBH, (excluding current year's growth), that are not yet tagged must be tagged and an azimuth and distance recorded from plot centre. Place tags on the north side of the tree and mark DBH with a blue painted stripe. Any stem that is forked below breast height is counted as two stems and both stems are tagged. All new saplings are numbered continuing from the last number previously assigned. **DO NOT USE DUPLICATE TREE NUMBERS.** The numbering system for saplings starts at 2000 and continues until all saplings are numbered (up to 2999). The original numbering order started at the SW corner and proceeded in a fan pattern to the NE corner. Ingrowth or new, additional saplings should be tagged in a similar pattern. Stem map all new saplings (**note that, for analysis purposes, 2 saplings cannot have identical azimuths and distances**) and record height, diameter, conditions codes and crown class of all saplings. All of this data is recorded on the sapling tally sheets (CSTM 101) under record type 7 (see Figure 4.25).

In extremely dense stands, stems that are on the border of the tree plot sides, but not actually within the plot, can be painted with an orange or yellow vertical stripe facing into the tree plot. This will assist future crews in determining the plot boundary.

Age and get origin on all **NEW** sapling - using guideline provided if necessary.

Ensure proper swathing techniques are followed. See Figure 2.3.

If trees cannot be tagged on a lateral branch, tie the tag around the tree with a long wire and leave it loose so that the tree is not girdled before the next measurement.

Trees that have a DBH > 9.1cm needs height to live crown measurements.

3.4 TREE PLOT (RECORD TYPE 8)

The tree plot is the full 1/10 ha plot. Once a stem reaches 9.1 cm in diameter at breast height, a tag is tied to a lateral brach or around main stem with a long piece of wire and a number assigned. Refer to the Permanent Sample Plot field procedures manual for proper "swathing" methods. The numbering system for the tree plot starts at 3001 and continues until all the trees are numbered. The height, height to live crown and diameter of all trees and azimuth and distance from the tree plot centre are measured and recorded along with any condition codes present and crown class. This information is recorded on the stand dynamics tally sheet (CSTM 101) under record type 8 (see Figure 4.26 Use condition codes in Master Condition Code List to assess the trees health. DBH is painted blue, just like in the sapling plot.

Trees from record plot 6 or 7 that grow over 9.1cm will **KEEP** their original number, i.e.; #1051 will stay as #1051 but you must also record height to live crown. New trees that grow over 9.1cm are recorded from 3001 and up.

Age and get origin on all trees - use guideline provided if necessary. Also use other trees similar in height in NW corner to get a good estimate of Birth year.

Use codes => 51 for this record. May use other codes if these are in appropriate.

Planted trees in NE, SW and SE quarters have had the main stem painted blue. If these trees become 3000 trees, we don't need Birth year or origin, just write planted in comments.

3.4A CROWN WIDTH MEASUREMENT

Starting with the year 2000 field season the crown width of a sample of trees/saplings/regen by species will have 4 crown widths measured. The crown width to the North, West, South and East of the selected tree/sapling/regen will be recorded to the closest decimeter and recorded in the comments section. The criteria for selecting trees/saplings/regen to have crown width measured **by species** is as follows for new trees/saplings/regen (previous measured trees/saplings/regen have already been selected).

- a) If < 10 trees/saplings/regen select all new trees/saplings/regen for crown measurements.
- b) If > 10 but <=20 trees/saplings/regen select every second tree for crown measurements.
- c) If > 20 but <=40 trees/saplings/regen select every fifth tree for crown measurements.
- d) If > 40 but <=100 trees/sapling/regen select every eighth tree for crown measurements.
- e) If > 100 trees/saplings/regen select every tenth tree for crown measurements.

This measurement is to be taken in the 4 cardinal point directions and is measured from the stem out. A reading of 2.5 metres is to be recorded as 25 (decimeters). This measurement is not to be done on dead trees/saplings/regen. If a tree has died that has been selected to have its crown measured, do not measure dead branches. Select the next tree of the same species as the new crown width tree and record the 4 measurements on the tally sheets.

The crown measurement is estimated at the widest portion of the foliage looking up from the base of the tree (Figure 2.8).

3.5 PLOT PHOTOGRAPHY

35 mm colour photos of each plot are taken once the measurements are complete. These photos serve as visual documentation of the overstorey and understorey of the plot and cover type. Numbered cards must be used in the photographs to identify the plot, as well as a card showing the month/ year the photo was taken. Two photographs are taken from plot centre; one facing east and the other facing west, with the photographer standing at plot centre.

3.6 VEGETATION DESCRIPTION

Figure 4.28 shows the format used for plot vegetation. The record type is #5 (Parent Vegetation is only measured at establishment).

3.7 PLOT VEGETATION ASSESSMENT

The vegetation plot is a 10m x 10m square plot located in the southwest corner of the tree plot (see Diagram 2.1). The vegetation plot can be moved to the northeast corner of the tree plot only if the southwest corner is not representative of the entire tree plot. If the vegetation plot is moved to the northeast corner, be sure to indicate this on the vegetation description form under the comments section. All this information is recorded on Stand Dynamic Vegetation Sheet (CSTM 102). Coverage can be over 100% as a result of one species growing beneath another. Assess the vigour of the species using a three part scale:

- 1 - Poor
- 2 - Average
- 3 - Outstanding

- a) **Field Name** - The field name is a seven letter code based on the scientific name of the species (see Master Appendix 12 & 13). The first four letters of the genus are recorded, followed by the first three letters of the species. For example, white spruce (*Picea glauca*) is tallied as PICEGLA. In some instances it is acceptable to record only the genus (i.e. *Salix*). When this is done the species part of the field name is filled in as "SP".

When you do not know what exact species the plant is write down the 4 letters of the genus and 'SP'. **Do not** use a period or an extra P. (eg. PYROSPP. Should be written as PYROSP). If in doubt about what to record, please check Appendix 13. The only exceptions are the Poa species. They require a space after POA. Eg. POA GLA, POA INT, POA SP.

- b) **Code** - The next range of columns divides the plant cover into eight groups, based on height and vegetation type. Within each group there are three columns labelled "%" and "vigour". The first two of these columns are reserved for recording the percentage of the area which is covered by each coded species in each group, and the last column (vigour) is used to record a subjective evaluation of the plant species vigour within the plot.
- c) **Trees** - All vegetation taller than 5.0 metres, regardless of species, is recorded in this column.
- d) **Tall Shrubs** - Includes all vegetation between 2.5 and 4.99 metres tall.
- e) **Low Shrubs** - Includes all vegetation between 0.3 and 2.49 metres tall.
- f) **Ground Shrubs** - All woody species that are less than .29 metres tall.
- g) **Herbs** - Record all herbaceous species in this column.
- h) **Grass** - Record all grass species within the plot.
- i) **Moss/Lichen** - Record all moss species within the plot.
- j) For trace amounts of any species, record the percent cover as 0%. In cases where there is more than or equal to 0.5%, record the percent cover as 1%. Note that 1% cover on 1/10 ha is ten metres squared (1 m x 1 m).

3.8 PLOT RETREATMENT REPORT (RECORD TYPE 9)

The plot re-treatment report is filled out at each remeasurement. The header information (boxes 1-24) must be filled out regardless of the plot has been disturbed or not. See figure 4.29

The report consists of 5 parts:

1. **Stand Tending:** The year of treatment refers to the last year the plot was tended e.g. 1989 - record as 89, type of treatment. Indicate Y(yes) or N(no) in each box that describes whether the plot was thinned, cleaned or deciduous species only were removed. More than one box can be filled in. Record the % of the tree plot to the nearest 10 that has been cleaned, etc. If the plot has been fertilized, obtain the fertilizer type and application rate from the appropriate district office, or forest headquarters.
2. **Replant:** Thee plot and/or buffer was replanted indicate this in columns 46 with an x. Obtain stock type from the district office and the date it was replanted.
3. **Weather/Natural Damage:** If there is any evidence of flooding, hail damage, browsing, snow or frost damage, indicate with an 'x' beside the appropriate description. Record the percent damage to the tree plot to the nearest 10%.
4. **Physical Description:** Indicate true or false in the box if there is a windthrow within the tree plot. If there is exposed mineral soil and no vegetation growing on it record the

percentage of the tree plot that it covers to the nearest 10%. If the buffer is a standard 200m x 200m record 'Y'. If it is smaller or larger, record 'N'.

5. **Plot Sizes:** Record the north and west plot sides for both the sapling and tree plots. Measure to nearest 0.1m. Write above each leg, the leg you are measuring, e.g. NW to NE, NW to SW etc.

Azimuth and distance to each Regen Plot from plot centre is to be recorded in the comments section. (This only has to be done once - check with supervisor if needed).

3.9 PLOT AGING

1. Ages in the four regeneration plots.

Ages on tally sheets should not be used to age the trees. This is an approximation and may change based on your age that you get in the field.

When aging trees, it is important to use the height increment printout along with counting increments. If a tree was tagged in 1986, it cannot have a birth year of 1989.

- a) Fill in the plot number and date on the age sheet (see Figure 4.30).

In Regen Plot# column record appropriate regen plot number. Use form for sapling and trees with appropriate Record number.

- b) These forms can be used for more than one regen plot **BUT** separate plots by a blank line.
- c) Select three of the tallest, tagged, healthiest trees for both tree types, Advanced and Ingress for each species. These trees should have a high probability of being there at least 25 years from today. **DO NOT** select trees that are obvious of poor vigour (dying).

Advanced tree type are trees that were left during harvesting and their age is older than the remeasurement date minus harvest date.

Ingress tree type are trees that germinated after harvest date or scarification date (see SDS checklist form). (Record "A" in tree type column for Advanced and "I" for Ingress).

- d) If less than 3 trees per species on a plot select all trees and record appropriate condition codes (on next line) if necessary (see lines 3 and 4, Figure 4.30).
- e) Planted trees do not need aging.
- f) Be AWARE that you may encounter a previous dead top or some other damage that the tree recovered from and you should include another year or two on top of the current age since the tree recovered. Pine have lammas branches which should not be counted as a whorl. This is common on pine but rare on spruce. Past condition codes (One

Only) can be recorded on the next line for that specific year. Mark the next lines Tree # column with CC (see Line#4 on tallysheet). If more HT number columns are needed use the next line (after CC line if necessary) and write in Year HT, etc. and record total height immediately below the new heading (see lines 6 and 7 on tallysheet).

- g) Deciduous trees over 8 metres should not be taken as age trees. Tall deciduous trees yearly increments are hard to discern and the guidelines provided are to be used (see line #7).
- h) On conifers greater than 5 metres in height, yearly Total height is taken up to 5 metres and then Total height is taken in 5 year increments (see line 10 and 11 on tallysheet).
- i) An additional five trees per tree type, per species in each plot are selected systematically for Total age and height ONLY (see lines 15 and 16 on tallysheet).
 - Count the total number of trees alive for each species and subtract the three trees already aged. Divide this number by five to decide which trees to measure for height and age only i.e. 20 Pl on regen plot minus 3 trees already measured = 17 divided by 5 = 3.4. Round off to 3. Take tree #3, 6, 9, 12 and 15 depending on point #2 below.
 - These trees are taken whether they are damaged or healthy. If the tree has already been used as one of the initial three tallest trees, take the tree (same species) before or after it.
 - If less than five trees for that species on the plot, record them all.
- j) DO NOT fill in any height measurements if you can't tell the increment on that particular year, eg. Tree #2009 in 1989 was 464 and in 1992 it was 381, not sure for 1990 and 1991. DO NOT fill 1990 and 1991 with 381.

2. Ages in Sapling Plots.

DO NOT choose trees over 8 metres in height for age trees (see lines 18 to 21 on tallysheet).

4.0 ALLOWABLE ERRORS

STANDARDS FOR STAND DYNAMICS PLOT MEASUREMENT

4.1 REGENERATION PLOT MEASUREMENTS

<u>ITEM</u>	<u>STANDARD</u>	<u>ALLOWABLE ERROR</u>
Tagged Seedlings (Type 6 data)	All seedlings must be located. Condition code (if any) filled - i.e. for dead, damaged or diseased seedlings.	None Any impacted seedlings must have appropriate condition code recorded
Species	All seedlings must be correctly identified by species. Special consideration should be given to Sb versus Sw, Pb versus Aw.	None
Height	Using Regeneration height measuring stick, measure to base of the candle or branch whorl on stem (exclude current years growth).	The allowable error for measured seedling heights is ± 4.0 cm
Seedling growth to Sapling height	All stems located and stem mapped (azimuth & distance)	Azimuth recorded within 1 degrees and distance within 10 cm.
Number of non-tagged seedling tally (type 61)	All seedlings not tagged must be accounted for and species and height class recorded on height class tally sheet	Total number of seedlings recorded within 2 seedlings.

4.2 SAPLING PLOT MEASUREMENTS

Saplings already tagged	All saplings located and condition code (if any) filled in	Any impacted sapling must have appropriate condition code recorded
New Saplings (ingrowth)	Saplings must be located, correctly numbered and condition code, if any, filled in	Numbering continues from last tagged sapling. Any impacted sapling must have appropriate condition

		code recorded
Species	All saplings must be correctly identified by species	None
Height	Measured with sapling height pole	Measured within 20 cm
Diameter	Measure at breast height (1.3 m)	Diameter measured within 1 cm. Breast height $\pm$ 1.0cm.
Number of Trees	All new trees must be recorded and stem mapped (azimuth & distance). Take care with 2 stems from the same stump, both tagged and measured if fork below 1.3 m	Allowable Error $\pm$ 1% of all trees tallied Azimuth recorded within 1 degree, distance within 10 cm. Record declination used

4.3 VEGETATION MEASUREMENTS

Vegetation Tally Sheets	All major species must be identified and recorded. Identify as much vegetation as possible and record an estimate of vigour.	Major species are those > 2% cover must be identified. Total plot vegetation cover to be measured within 20%.
-------------------------	--	---

4.3A PLOT RETREATMENT MEASUREMENTS

	All disturbances must be marked in the appropriate box	Percent damage recorded must be within $\pm$ 10% of the check cruise.
--	--	---

4.4 PLOT REJECTION

The plot may be required to be remeasured if the entire plot has:

- more than 5 percent of regeneration height measurements are outside specified allowable error - or;
- more than 5 percent of sapling height measurements are outside specified allowable error - or;
- species is incorrect on more than 5 percent of regeneration and sapling measurements - or;

- the number of seedlings by species is incorrect in more than 2 height classes on a plot
- the condition codes are incorrect on more than 10 percent of the regeneration, saplings and trees - or;
- azimuth and distance measurements are incorrect on more than 10 percent of the saplings or trees - or;
- vegetation tally sheets are incomplete, completely missing more than 3 major species (which cover more than 2 percent of plot area).
- plot retreatment report is incomplete.

4.5 DATA ENTRY ON HEADER AND TALLY SHEETS

Measurements taken on permanent sample plots are recorded on Permanent Sample Plot Header Sheets (TM267), Permanent Sample Plot Tally Sheets (TM249) and on Regeneration Tally Sheets (TM261). Data recorded on the front of each tally sheet in the white columns is keypunched in the same format to keep the tally sheets and computer files compatible. It is important that all letters and numbers are legible to avoid keypunch errors resulting in costly computer errors. Quality measurements are of no value if they are not legible. See Appendix 14 for common problems.

4.6 STAND DYNAMICS GENERAL INFORMATION (CSTM 100 AND CSTM 101) – See Appendix # 17.

4.7 SUGGESTED EQUIPMENT AND SUPPLIES - See Appendix # 1.

4.8 TREE SPECIES CODE – See Appendix #12.

4.9 TREE HEIGHT MEASUREMENT USING A CLINOMETER – See Appendix #14.

4.10 HEIGHT TO LIVE CROWN

The height from the ground to the base of the live crown (see Figure 2.5) is measured on all trees that have been measured for total height. The base of the live crown is the point that separates the continuously branched portion of the tree and the part that has sporadic or no branching. Live crowns on deciduous species start at the leaves, not at the branches. Live crowns on coniferous species start at the tip of the live branch, not at the base of the branch. The height to live crown is quite variable depending on stand maturity and density with young, open stands having low live crowns and mature, stocked stands having higher, live crowns. Height to live crown is measured on age trees when they are on the ground and being prepared for sectioning.

- 4.11 SLOPE CORRECTION FACTORS AND TABLES – See Appendix #5.**
- 4.12 CONDITION CODES – See Master Condition Code List.**
- 4.13 MISTLETOE RATING SYSTEM – See Figures 11 and 12.**
- 4.14 LEGAL SURVEY OF ALBERTA – See Appendix # 8**
- 4.15 DETERMINING BREAST HEIGHT – See Appendix #14.**
- 4.16 ROUNDING OFF – See Appendix 9-h.**
- 4.17 PROBLEMS WITH RECORDING DATA – See Appendix #15 and 16.**
- 4.18 CROWN CLASS**

Crown class (CC) refers to the position of an individual tree within the canopy of the stand inside the plot. Crown class is assessed on a plot-by-plot basis, not on the stand as a whole. For example, an intermediate tree in one PSP plot may be codominant in the next. The following figure shows the types of crown class in a single layer stand. Crown classes are recorded for all trees with the exception of those with a broken top/system, are dead, cut down, missing, or have a severe lean (see Figure 2.6).

4.19 STEM MAPPING

All tagged stems within the tree sapling plot are stem mapped. Stem mapping is used to identify the position of each tree or sapling with respect to other surrounding trees or sapling and can be used in distance dependent growth models and is used in plot remeasurement to locate missing trees or sapling.

A staff compass and a metric tape is used to determine the azimuth and distance to the centre of each tree or sapling, at breast height, from the plot centre. Azimuth are recorded from 1°-360° and distances are measured to the nearest 0.1 m. Standing dead trees are to be stem mapped.

Regeneration are not stem mapped.

- 4.20 SEED VS. SUCKER – ORIGIN ON POPULUS SP. – See Figure 4.32.**

References for Condition Code Assessments for Wildlife

- Conner, R.N. 1979 Seasonal changes in woodpecker foraging methods: strategies for winter survival. *In* Dickson, J.G., Conner, R.N., Fleet, R.R. Jackson, J.A. and J.C. Kroll (eds), *The Role of Insectivorous Birds in Forest Ecosystems*. Academic Press. New York. pp 95-105.
- Harrison, H. 1979 *Western Bird Nests*. Peterson Field Guide Series. 280 pp.
- Hiratsuka, Y. 1987 *Forest Tree Diseases of the Prairie Provinces*. Information Report NOR-X-286.
- Murie, O.J. 1975 *A Field Guide to Animal Tracks*. Houghton Mifflin. 375 pp.
- Rangen, S.A. and L.D. Roy. 1997
 A Field Guide to Animal Damage of Alberta's Native Trees. Alberta Research Council. Begreville, Alberta. ARCV97-R1. 58 pp.
- Sousa, P.J. 1987 *Habitat Suitability Models: Hairy Woodpecker*. Biological Report 82(10.146). U.S. Department of the Interior. Fish and Wildlife Service. 19 pp.

PSP MANUALS MASTER CONDITION CODE LIST

MARCH 2005

**Public Lands and Forests Division
Forest Management Branch
8th Floor, 9920 – 108 Street
Edmonton, AB
T5K 2M4**

Phone: (780) 427 - 8474

Or visit the website: <http://www3.gov.ab.ca/srd/forests/psp>

1974

CLARKSON, W. A.
117 1112
171, 1712 17121212
1712 17121212

1712 17121212

1712 17121212

CONDITION CODES

CODE	DESCRIPTIONS	CODE	DESCRIPTIONS
00	Healthy	47	Witche's Broom
01	Insects	48	Frost Crack
02	Disease	49	Dying
03	Rabbit Browsing	51	Conks/Blind Conks
04	Shepherd's Crook	52	Open Scars
05	Browsing (Other animal)	53	Burls and Galls
06	Fire	54	Fork
07	Mechanical	55	Pronounced Crook
08	Windthrow	56	Broken Top (DBH > 9.1) (No CC) (see "24" for DBH <9.1)
09	Climate	57	Limby
10	Flooding	58	Severe Lean (No CC) – see code #35
11	Poor Planting	59	Broken Stem (>=10cm DIB at Break DBH >9.1) (No CC)
12	Suppression	60	Generic woodpecker feeding (often smaller species)
13	Frost Heaving	61	Dead and Down (No CC)
14	Erosion	62	Stem Insects (Bark + Sawyer Beetles)
15	Missing	63	Stem Disease (Cankers)
16	Dead Top/Dieback	64	Foliar Insects
17	Poor Seedbed	65	Foliar Disease (Needle blights + rusts)
18	Herbicide	66	Stem Form Defect (>=7.0cm DIB at point where stem form begins)
19	Western Gall Rust (only on Pine)	67	Closed Scars
20	Armillaria Root Rot	68	Atropellis canker
21	Moldy Planting Stock	69	Comandra Blister Rust
22	Multiple Leader	70	Elytroderma needle cast of pine
23	Poor Form	71	Hypoxylon Canker
24	Broken Top (DBH <9.1) (see "56" for DBH > 9.1)	72	Spruce cone Rust
25	Dead Tree Standing (No CC)	73	Stalactiform Blister Rust
26	Snow Press (No CC)	74	Tomentosus Root Rot
27	Dead Top Dieback with NEW Leader	75	Spruce Spanworm
28	Sucker(s) (from OLD Stump)	76	Spruce Cone Maggot
29	Cut down	77	Spruce Cone worm
30	Terminal Weevil	78	Eastern Spruce Budworm
31	SW Gall Adelgid	79	Mountain Pine Beetle
32	Tent Caterpillar	80	Spruce Beetle
33	Root Collar Weevil	81	Spruce Needle Rust
34	J-Root	82	Yellow Headed Spruce Sawfly
35	Leaning (No CC) – see code # 58	83	Large Aspen Tortrix
36	Same Stump	84	Excavations by woodpeckers
37	Unknown	85	Yellow-bellied sapsucker feeding
38	Pitch Moth	86	Small mammal feeding on tree bole
39	DBH Taken on New Leader	87	Small Cavity
40	Nutrient Deficiency	88	Large Cavity
41	Mouse (feeding)	89	Hollow tree or hollow bole section
42	Ungulate feeding/rubbing	90	Beaver (feeding/harvesting)
43	Domestic livestock (rubbing)	91-96	Hawksworth Mistletoe Rating System
44	Nest	97	Available for future consideration
45	Other mammalian/avian evidence	98	Data changed by office
46	Sweep/Bow/Bend	99	Do not look for tree

Note: No CC means no crown class.

CONDITION CODES DESCRIPTION

Condition Codes	Description
00 Healthy	No Defect.
01 Insects	Damage or mortality due to destruction of plant parts or tissue by insects. Look for evidence of eggs, egg cases, nests, chewed plant parts, etc. Similar signs on plants located off site may aid in identification of insect mortality.
02 Disease	Damage or mortality caused by disease or fungi. Cankers, discoloration, rust spotting, fungal coverings, etc. help to identify mortality under this code.
03 Rabbit Browsing	Trees killed or damaged by rabbits can be identified by clean, sharp cut marks along the branches and stems (approximately 45° angles). Chewed bark and needles also indicate rabbit damage.
04 Shepherd's Crook	Damage results in blackening and wilting of young shoots and leaves. Tips of the blackened shoots often bend back. On older leaves brownish black, irregularly shaped spots appear.
05 Browsing (other animals)	Mortality or damage due to browsing by ungulates or other animals (e.g. moose, cattle, beavers). Look for chewed tops with rough cuts or breaks.
06 Fire	Mortality or damage due to actual burning of the seedling or scorching by nearby flames. Not to be used when seedlings are killed by sun scald.
07 Mechanical	Trees killed or damaged by mechanical or physical means such as scarification machinery, trampling or crushing by animals, etc. Stem scars and rough breakage help to identify mortality under this code.
08 WindThrow	Damage or mortality due to crushing by fallen or displaced logs, snags, branches, uprooted trees, etc.
09 Climate	Trees damaged or killed solely by climatic factors. These include death by freezing, sun scald, severe desiccation, ice accumulation, red belt, etc.
10 Flooding	Trees damaged or killed by drowning alone. Look for evidence of high water marks on the seedling, or in the immediate area. Pull tree out of ground and check roots to see if the root outer coverings is falling off and is blackened.
11 Poor Planting	Damage or mortality due to improper placement of nursery stock (hand or mechanical planting). Trees may have been planted too deep, too shallow, too loosely, or at an acute angle.
12 Suppression	Trees which have been suppressed by the surrounding vegetation for a period of time long enough to damage or kill them. Mortality may be due to severe lack of light, water, nutrients (removed by the competition) or by physical smothering (i.e. heavy grasses). Reference to the previous year's damage tally may help in determining this mortality call. A tree that is over topped by grass or shrubs is not necessarily suppressed. Look for a spindly main stem with very few long needles spaced wide apart or evaluate the last five increments. If the tree has only grown 1cm a year, it is probably suppressed.
13 Frost Heaving	This code is used only when mechanical frost action can be clearly identified as the direct cause of damage or mortality. Usually upheaval and separation of the seedling's root system from the soil occurs as a result of ice lens formation. This is most commonly associated with containerized seedlings planted in silty soil.
14 Erosion	Damage or mortality due to the removal of the seedling's seedbed, by the forces of water, wind or soil slumping. Trees killed by partial or total burial (deposited soil or organic matter) would also be tallied using this code.

15	Missing	This code is to be used when a seedling from the previous year's measurement cannot be located. It can also be used where the seedling was removed from the site and probably died (i.e. tag found, no morphological signs of live seedling remaining). Using in conjunction with code 25 ONLY .
16	Dead top/Dieback	Top is dead (die back) without any indication of insect or climate (frost) damage. See Figure 4.31
17	Poor Seedbed	This code is to be used only when the cause of death or damage for a seedling can be traced to the type of seedbed on which it is growing. In most cases the seedling will show signs of desiccation due to the poor moisture holding capacity of the seedbed material (e.g. rotten logs, dry clay).
18	Herbicide	Should only be used when the cutblock (or parts of the cutblock) has received a recent herbicide treatment; either before or after the stock was in place. Spruce seedlings exhibit needle loss and/or reddish brown coloration of stems and foliage. Deciduous species exhibit yellowish/brown leaf mottling and dieback of terminal growth. Hexazinone causes reddish brown coloration of conifer foliage and needle loss. Deciduous foliage turns red to black. Glyphosate causes chlorosis especially in new growing shoots. 2,4-D causes rapid growth and spiralling and twisting. If applied during conifer flush bad dieback similar to frost damage may occur. Often chemical damage will also be indicated by phytotoxicity spotting on exposed foliage.
19	Western Gall Rust (only in Pine)	This code is used when Lodgepole pine damage or death can be attributed to Western Gall Rust. This is usually clearly identifiable due to swelling of succulent tissue (and subsequent formation of a gall) and the bright orange spores produced in that affected area. This gall can be on the main stem or a lateral branch.
20	Armillaria Root Rot	This code is utilized when a seedling is damaged or killed by Armillaria Root Rot. Identification of the disease is in recognizing mycelial fans of the cambium of damaged and dead trees. Pull tree out of ground and examine root collar.
21	Moldy Planting Stock	This code is usually used on Bareroot Planting Stock. Grey mold will usually be found around the root collar and lower branches.
22	Multiple Leader	When a tree has two or more leaders, but is otherwise healthy this code should be entered. The tree is considered multiple leader if all leaders are within 5cm (height) of each other. This code also applies to saplings and regeneration that appear forked. Be aware of normal branching of deciduous trees.
23	Poor Form	This code is used on trees, which exhibit a general poor form, due to previous damage. DBH < 9.1 See code 66 for >9.1 DBH
24	Broken Top (DBH <9.1) (see "56" for DBH > 9.1)	It should be used as long as the broken top is noticeable and has some effect on the growth of the tree.
25	Dead Tree Standing (No CC)	Tree has no signs of being alive. A standing dead tree is one that is dead but still standing. No green foliage or buds present. The tree must be able to withstand a firm push. Record a diameter and species but do not record height. Pound nail into tree. No crown class.
26	Snow Press (No CC)	This code is normally used for trees that show signs of being pressed down to the ground for a few years after germinating. May Happen to bigger trees. See Figure 4.8
27	Dead Top Dieback with New Leader	This refers to stems that have had previous leader damage and a new leader has formed.
28	Sucker(s) (From Old stump)	Refers to stems that have been cut-down through thinning and have started to sucker. Do not re-use the previous stem number, but assign a new number to each sucker.
29	Cutdown	Self explanatory.
30	Terminal Weevil	Terminal leaders of Pine or Spruce bend over and die. Two or more years growth are affected. Bore Holes which are exit holes for the larvae MUST be present to

		use this code.
31	SW Gall Adelgid	Adelgid galls on spruce located at the end of new growth and may persist for many years.
32	Tent Caterpillar	A tent of silk forms on the tree and the caterpillars defoliate the tree.
33	Root Collar Weevil	This weevil feeds mainly on Sw, Pj and Pl. They feed in the bark and cambial area of the host tree at or below the duff surface, causing copious flows of resin. The tunnels often girdle small trees. This insect allows root rots to enter the tree.
34	J-Root	This code is used after the tree has had a poor planting code in the previous measurement.
35	Leaning (No CC) see code 58	Tree leaning more than 20% off of vertical axis.
36	Same Stump	Used when 2 or more trees can be distinguished above ground level but below DBH. Used a lot on Deciduous that have been cutdown and resprouted at stump.
37	Unknown	This condition code is to be used only when there appears to be something affecting the tree but the other condition codes do not describe the situation. This would include burnt trees etc. A description of what is affecting the tree should be included as well in the comments column. In the event that this code is used for more than 5% of the tallies, it is up to the crew leader or a forester to decide on the cause of the condition.
38	Pitch Moth	Primary host is Lodgepole Pine. May weaken or kill the terminal leader, resulting in stem deformities and height growth reduction. Blisters are mainly on main stem and are characteristic resin coated up to 20mm in diameter.
39	DBH Taken on New Leader	
40	Nutrient Deficiency	This may occur on blocks that have had the humus layer removed by scarification (i.e.; Blade). Trees are chlorotic and usually in bare mineral soil. Usually noted on spruce. May be confused with flooding damage.
41	Mouse Feeding	Mice and voles can girdle seedlings and consume seeds. See Rangen and Roy (1997) for more detail.
42	Ungulate feeding/rubbing	Ungulate feeding on twigs is generally recognized by the ragged appearance of twig terminals. Rubbing of trees as antler rubs and feeding on bark also occurs; these conditions are further described in Rangen and Roy (1997). Antler rubs can also be associated with "scrapes" (smaller patches of scraped ground) and small tufts of hair on twigs. If the bark on aspen trees has been consumed ensure that ungulates (as opposed to other mammals) are responsible. The extent of the bitten area, track identity and grooves that indicate tooth size and pattern should all be inspected in order to differentiate ungulate bark feeding from similar feeding by small mammals (i.e. see code number 86 and applicable photograph).
43	Domestic livestock (rubbing)	Rangen and Roy (1997) describe rubbing of trees by livestock; rubbed trees are occasionally seen in areas where cattle grazing occurs. If this code is used, ensure that other signs in general area (i.e. presence of cattle droppings, cow trails and grazed vegetation) also supports this.
44	Nest	This code indicates the presence of a nest on a given tree in the PSP. It refers only to an "open" nest; cavity nests are excluded from this category, as it is difficult to ascertain if a given cavity is indeed used as a nest site. Field guides that assist with the identification of "open" nests are available (see Harrison 1979). Of particular importance are colonial complexes of large nests on islands in lakes. Mammalian nests also exist and should be indicated as such if this is known. To do this, use the comments section which applies to a given tree and indicate as required. If the occupants of the nest can be identified, the identity can also be entered in the comments section.

45	Other mammalian/avian evidence	Other agents (i.e. bears, grouse, shrew, pocket gophers) which leave evidence on trees or leave evidence closely associated with trees are described in Rangen and Roy (1997). Pocket gophers leave soil mounds (Rangen and Roy - (1997). Bears can leave a characteristic series of claw marks on aspen trees, indicating that the tree was scaled, and rotted stumps/logs are also occasionally ripped apart. In addition, it has been suggested that bark on live trees is occasionally consumed (see Hiratsuka 1987 for a depiction). Ensure that ripped up stumps/logs, etc. are accompanied by other evidence of bear.
46	Sweep/Bow/Bend	Is a gradual bowing or curving of the main tree system. It has no decay significance but may cause a loss of volume in a sawlog.
47	Witches Broom	Yellow witches broom is the most conspicuous disease of spruce in the province. Can be recognized from a distance. See Figure 4.10
48	Frost Crack	A frost crack is a deep radial splitting of a trunk caused by an uneven shrinkage of the wood after a sudden drop in temperature. The cracks usually start at the base and extend up the trunk. They may re-opened repeatedly by wind stresses or a low temperature.
49	Dying	Tree is in distress and will die before next measurement.
51	Conk/Blind Conks	Conks appear most frequently on the underside of dead branch stubs or on the underside of live branches in the crown. Conks, by definition, are woody, shelflike basidiocarps (fruiting bodies) of wood-rotting fungi. See Figure 4.3
52	Open Scars	Open scars are wounds which have been penetrated through to the cambium. These wounds must not be healed over and may be caused by a variety of reasons such as fire, lightning, old blazing, machinery, animals, etc. Scars are considered to be entry points for decay fungi. Open scars are illustrated in Figure 4.4. Animal damage usually penetrates the cambium therefore code as an open scar. A common mistake is to call stem disease such as atopellis canker an open scar. See Figure 4.4
53	Burls and Galls	Burls are abnormal swelling of the main stem or branches resulting from abnormal wood cell development following disturbance to the cambial layer. A burl is illustrated in Figure 4.5. Galls are localized trunk and branch swelling of mainly tissue. There is little or no damage to the underlying wood.
54	Fork	Forks usually develop when there is malformation, injury or death of the terminal leader. Forks tend to be V-shaped and will only be recorded when above 1.3 m (DBH level). Forks below this point are recorded as same stump (condition code 36). Natural branching on deciduous trees is not to be recorded. Figure 4.6 demonstrates the difference between forks and natural branching.
55	Pronounced Crook	This condition develops from the death of the terminal leader or the breaking off of a forked leader. When this occurs, a lateral branch takes over apical dominance as shown in Figure 4.7.
56	Broken Top (No CC) (see "24" for DBH <9.1)	Broken tops occurs usually in the top third of the tree. No Crown Class.
57	Limby	A tree is recorded as limby if more than 75% of the tree has live, low sweeping branches. Usually branches on coniferous >2.0m on any part of tree could be considered limby.
58	Severe Lean (No CC) see code 35	A tree is considered leaning if it is standing greater than 20° off of vertical (see Figure 4.8). If the angle is greater than 45° to the ground, the tree has a severe lean. No crown class if severe.

59	Broken Stem (No CC)	A broken stem is recorded if the tree bole broken. Usually found in bottom 2/3 of the tree. No crown class.
60	Generic woodpecker feeding (often smaller species)	Figure 4.13 indicates feeding by woodpeckers. Species such as the Black-backed woodpecker and Three-toed woodpeckers will often leave signs like this on old coniferous trees, and Hairy and Downy woodpeckers typically peel off scales ("scale") and "peck" the bark as do Pileated woodpeckers in summer months (Conner 1979). Note the evidence of very small holes (arthropods) and holes made by the woodpeckers themselves. The appearance of tree trunks fed on in this manner is often reddish from a distance.
61	Dead and Down (No CC)	A dead and down tree is one that was previously tagged and measured in a PSP plot but at the present time is now dead and no longer standing. The cause of death must be by natural causes (i.e. windfall, beavers, insect or disease, etc.). No crown class.
62	Stem Insects (Bark + Sawyer Beetles)	This code is recorded when there is evidence of an insect infestation attacking the bole of the tree. Bark beetles are the most prevalent stem insects but sawyer beetles and others are included. Bark beetles, <u>Dendroctonus spp.</u> , are a very serious problem in Alberta. The adult female enters the bark in early summer and lays eggs in the tree's cambium. The eggs overwinter and hatch as larvae in the early spring. Damage to the tree is done by the larvae eating the cambium and usually results in death. The tree will not turn red until the next summer. Other symptoms of attack are piles of "sawdust" (frass) at the base of the tree, entry holes in the bark, and pitch tubes (the tree tries to push the beetles out with resin). The beetles also carry a blue stain that causes further deterioration of wood quality. Beetles attack all species of pines, spruce, and Douglas fir. Sawyer beetle infestations are common in burned timber.
63	Stem Disease (Cankers)	<p>All diseases that infect the main stem are documented with this code. Included in this code are cankers, rusts, rotten branches and root rot.</p> <p><u>Stem cankers</u> are caused by fungi that invade stems and branches resulting in localized areas of infection in the bark and underlying wood tissue. Cankers may be annual or perennial. In perennial cankers the infected area may be eventually exposed to the underlying wood when the deadbark sloughs off. A common stem canker on lodgepole pine is <u>Atropellis piniphila</u> (Figure 4.9). Exudation of resin from the bark surface is the first external symptom. They are sunken elongated on one side of the trunk and indicate resin flow. This can cause a distortion in growth and a blue-black stain on the wood.</p> <p><u>Stem rusts</u> are also included in this condition code. Rusts are host specific parasitic fungi usually requiring two alternating living hosts. Stems and branches may be girdled resulting in large malformations or even death. In particular, <u>Endroconartium harknessii</u> on young pines is a serious problem in Alberta. Spruce broom rust, <u>Chrysomyxa arctostaphi</u> (see Figure 4.10), can also be noted but only if the broom is no longer green (i.e. red or missing needles).</p> <p>Large rotten branches typically appear on overmature, decadent trees and can be indicative of decay. Large rotten branches are those well below the base of the live Crown and are > 5 cm in diameter, are unweathered, appear punky, and are weeping. Often a black ring appears on the stem surrounding the branch.</p> <p>Some of the typical symptoms of Armillaria root rot are reddish brown or yellowish foliage; mycelial fans form between the bark and wood around the base; fungal (shoestring) strands in the soil surrounding the diseased roots and honey mushrooms growing around the base of the diseased tree.</p>

64	Foliar Insects	<p>This condition code pertains to all insects that infest parts of the tree off the main stem. Included in this category are the tent caterpillar, spruce budworm, jack pine budworm, spruce gall aphid, etc.</p> <p>The forest tent caterpillar, <u>Malacasoma disstria</u>, causes severe defoliation in hardwood stands in Alberta resulting in a significant reduction in annual growth.</p> <p>The spruce budworm, <u>Choristoneura fumiferana</u>, infests mature white and black spruce, and balsam fir stands. This insect attacks the buds and new needles. Their feeding spreads to old needles and eventually kills the tree.</p> <p>The jack pine budworm, <u>Choristoneura pinus</u>, attacks stands of jack and lodgepole pine and is a relatively new forest pest in Alberta. This insect feeds and spreads in the same manner as the spruce budworm.</p>
65	Foliar Disease (Needle blights + rusts)	<p>This code is used for all diseases that infect parts of the tree off the main stem. Needle casts and blights, and needle rusts are included in this condition code.</p>
66	Stem Form Defects	<p>This condition code is used when there is damage or a distortion resulting in a loss of volume. Used for trees >9.1 DBH. See code 23 for <9.1cm DBH.</p> <p>A sweep or bend is the gradual bowing or curving of the main tree stem. If has no decay significance, but may cause a loss of volume in a sawlog.</p> <p>Spiral grain is the twisting of the grain seen in exposed wood or in the direction of the bark fissures. Spiralling frost cracks and scars also indicate the presence of spiral grain.</p> <p>Windshake is a splitting in the wood along the grain or less frequently within an annual growth layer. It is caused by wind or snow stresses and is also known as ringshake.</p>
67	Closed Scars	<p>Wounds that had penetrated the cambium but have now healed over are considered closed scars. A closed scar is characterized by an irregular indentation in the bole of the tree that would result in loss of volume due to poor wood quality. Before healing over, the scar provided an entry point for disease. Frost crack is not included in this code.</p>
68	Atropellis Canker	<p>Widespread on pine, from small to large trees. Symptoms are elongated, sunken, cankers on the stem with copious yellowish resin flow. Wood is discoloured blue/black. Figure 4.9</p>
69	Comandra Blister Rust	<p>Pl and Pj are hosts. Local occurrence only. Infected stems are spindle-shaped with conspicuous swelling of the bark. Fungus is orange-yellow in early summer. Cankers are circular and grow laterally as quickly as longitudinally. They thus girdle the stem faster than stalactiform. It should not be confused with western gall rust, which is mainly a swelling of the wood. Alternate host is Indian Paint Brush.</p>
70	Elytroderma Needle of Pine	<p>Mostly on Pl. Current years needles turn red in fall. In severe cases only current needles remain, giving branches a "lion's tail" appearance.</p>
71	Hypoxylon Canker	<p>Hosts are aspen and balsam poplar. Canker starts as a slightly sunken orange-yellowish area on stem. Eventually girdles the stem and has an orange/black appearance. A mycelial fan on the cambium is a reliable field symptom.</p>
72	Spruce Cone Rust	<p>Rust is <u>only</u> on spruce cones. Cones become prematurely brown then orange-yellow. When spores are abundant, the forest floor has an orange colour.</p>
73	Stalactiform Blister Rust	<p>Pl and Pj are hosts. Local occurrence. Causes slight swelling of bark. Orange-yellow in summer. Cankers are elongated and grow faster longitudinally compared</p>

		to Comandra. Alternate host is Bastard Toad Flax.
74	Tomentosus Root Rot	Most important on Sw and Sb. Symptoms are excessive branch mortality, thinning of crown and openings in the stand. Disease develops slowly (over 15-20 years) so is not so obvious in regenerating stands.
75	Spruce Spanworm	Chiefly affects aspen. Damage shows mostly as holes in the leaves. Resembles forest ten caterpillar but no pupal cases or egg masses on the foliage. Caterpillars are typically light green and have one prominent and two indistinct yellowish lines along each side of the body. The head is dark-brown.
76	Spruce Cone Maggot	No external symptoms. Dissected cone shows frass-filled spiral tunnel around the central axis.
77	Spruce Cone Worm	Feeding larvae expel frass, which adheres to silken webbing on cone surface.
78	Eastern Spruce Budworm	First symptoms are webbing and frass in buds or on previous year's needles. Later, webbing is spun on branch tips. By late June tree crowns appears rust brown.
79	Mountain Pine Beetle	Main host is Pl. Symptoms are standing dead trees with beetle exit holes about eye-level. Accumulations of pitch or sawdust are conspicuous around entrance holes bored into the bark of trees by adult beetles from mid-July to mid-August.
80	Spruce Beetle	Host are Sw and Se. Symptoms are standing dead trees with beetle exit holes about eye-level. Conspicuous boring dust accumulates on bark below holes until the wind blows it away.
81	Spruce Needle Rust	Feed on needles in the upper crown of the tree. Partly chewed needles and needle stubs impart a brownish color and ragged appearance to the foliage. No webbing present. Found on all spruce.
82	Yellow-Headed Spruce Sawfly	Discoloration of needles. May find dotlike sexual fruiting structures on needles. Infected needles drop prematurely.
83	Large Aspen Tortrix	Affected foliage has a clumped, irregular appearance and leaves do not move as freely in the wind as uninfested leaves. Larval instars feed within rolled leaves or within 2 or more leaves pulled together and secured with silken webbing.
84	Excavations by woodpeckers (likely Pileated woodpecker)	Feeding by Pileated woodpecker can occur on dead or senescent deciduous and coniferous trees, and feeding holes (as indicated in the figures below) are thought to occur towards the base of the tree (Rangen and Roy 1997). Excavated holes indicate subcambial penetration (holes <u>penetrate beneath the bark and into the sapwood</u>) and large wood chips can be associated with excavations. Excavated feeding holes can be large (Figure 4.16). In such excavations, evidence of carpenter ants (burrows, sawdust) or other boring arthropods might also be found in the sapwood. In living trees with a sound bole, initial feeding holes might be more restricted such as that indicated in Figure 4.12. Elsewhere in North America, the Pileated woodpecker has been found to excavate holes extensively in winter and to a greater extent than other woodpeckers (Conner 1979). The Hairy woodpecker might also create deeper holes in trees, however, it is considered an opportunistic feeder (Sousa 1987) and spends a smaller portion of its time "excavating" during winter months (Conner 1979). In Iowa, it has also been found to generally feed at higher locations in trees (5-7m) (Sousa 1987). If this feeding evidence exists on a given tree, indicate in comments its extent (i.e. restricted, such as in Figure 4.16).
85	Yellow-bellied sapsucker feeding	Figure 4.14 illustrates the characteristic pattern of regularly spaced small holes left by Yellow-bellied sapsucker (also see Hiratsuka 1987 for another depiction of sapsucker feeding). These are often found on birch, however they also have been observed on willows, and have been reported on aspen and pine (Rangen and Roy 1997, Hiratsuka 1987).
86	Small mammal feeding on tree bole (hare, porcupine, squirrel, bushy-tailed)	Figure 4.15 is an example of feeding by hare on small saplings. In this case the bark was bitten off. When hares feed on twigs, it is generally thought that twigs are clipped off in a characteristic razored fashion (Figure 106, Rangen and Roy, 1997). Small mammals such as porcupine, woodrat and squirrel might also feed on bark

woodrat)	however, if such feeding evidence occurs high in trees, one could probably rule out hare because hare do not climb trees (also see Hiratsuka 1987 for a depiction of porcupine feeding on pine). Ensure other evidence (i.e. tracks, pellets, etc.) Supports a specific determination of the agent involved. Also refer to Rangen and Roy (1997) for more information on how to identify the specific causes of girdling and refer to Murie (1975) for assistance on identifying tracks if this is required. Evidence of squirrel feeding is common and could also be indicated, however, the value of this information is probably less valuable.
87 Small Cavity	Small woodpeckers create small cavities (approximately 5 cm in diameter) in snags and stubs (Figure 4.17), however, height of the cavity above ground probably varies. Among the species which might use such cavities are smaller woodpeckers, kestrel, chickadee, nuthatch, swallow, wren, flycatchers, and small mammals (etc). One could explore whether such cavities are occupied by rubbing the bark with a stick. Should a cavity be occupied the occupant (if known) should be identified in the comments section.
88 Large Cavity	A large cavity is a round/excavated opening greater than or equal to 10 cm in diameter (see Figure 4.18). The cavity in the figure was approximately 15 m high. Pileated woodpeckers have been known to excavate such cavities, however, a variety of species (birds as well as mammals) may use them as nest sites, roosting sites or dens. As in the case of smaller cavities, one could investigate the identity of the occupant by rubbing/tapping the bark of such trees with a stick. It might be possible to ascertain the identity of the tracks which are associated with the cavity, during winter, by checking surrounding snow cover and identifying tracks that appear to lead towards the cavity in the tree (see Murie 1975).
89 Hollow tree or hollow bole section	Hollow trees can be used as denning sites by bats and other birds and mammals. This condition code should be used to identify these sites.
90 Beaver (feeding-/harvesting)	Beaver girdle large trees in a characteristic fashion and evidence of their harvesting activities (i.e. cone shaped stumps) are well known to many. Refer to Rangen and Roy (1997) and Hiratsuka (1987) for more details.
91-96 Hawksworth Mistletoe Rating System	<p>Dwarf mistletoes are parasitic flowering plants requiring living hosts. Mistletoe is usually recognized by swellings on branches and stems or by witches brooms. Heavy infestation makes trees susceptible to secondary attack (such as bark beetles), lower wood quality and growth losses (can be from 30-60%). The major tree hosts in Alberta are: lodgepole pine, Douglas fir and larch. Figure 4.11 illustrates the effect resulting from mistletoe infestations and the individual flowering plant.</p> <p>The Hawksworth Rating System for mistletoe is used to determine the severity of mistletoe infestation on individual trees. Figure 4.12 outlines instructions and gives an example of the use of the 6-class mistletoe rating systems (Hawksworth 1961, 1977). If a tree has mistletoe, record only the 90 series code, do not record 33 unless there is a second distinct foliar disease.</p>
97 Available for future consideration	
98 Data changed by office	
99 Do not look for Tree

PSP MANUALS MASTER APPENDICES

MARCH 2005

**Public Lands and Forests Division
Forest Management Branch
8th Fl. 9920 – 108 Street
Edmonton, AB
T5K 2M4**

Phone: (780) 427 – 8474

Or visit the website: <http://www3.gov.ab.ca/srd/forests/psp>

LEAFHOPPER
NOTES
FEBRUARY 1

THEY ARE

1-2-1

1-2-1

TABLE OF CONTENTS

1.	SUGGESTED EQUIPMENT AND SUPPLIES	1
2.	PLOT TOPOGRAPHY	1
a.	Location	1
3.	SOIL CODES	1
a.	Erosion potential	1
b.	Drainage Codes	2
4.	TREE HEIGHT MEASUREMENT	3
a.	Determining Tree Height with a Laser Clinometer or Haglof Vertex	5
5.	SLOPE CORRECTION FACTORS AND TABLES	5
6.	USE AND CONSTRUCTION OF A BILTMORE STICK	6
7.	ACCESS EVALUATION CODES	7
8.	LEGAL SURVEY SYSTEM OF ALBERTA	8
9.	FIELD SHEET HANDBOOK	8
a.	Tally Sheet Instructions	8
b.	General Information (Columns 1 – 37)	8
c.	Record 01 – Header Information (Columns 38-212)	11
d.	Record 02 – Tree Information – header Sheet (Columns 1-37) – Tally Sheet (Columns 38-124)	17
e.	Record 03 – Regen Information (Columns 38-85)	20
f.	Summary of Allowable Errors	21
g.	Plot Measurement Summary	22
h.	Rounding-Off	23
10.	MISTLETOE AND IMMATURE PLOT TYPES	24
11.	ASCII FORMAT OF PSP TREE DATA, RECORDS 1, 2 AND 3	25
a.	RECORD TYPE “01” – STANDARD SAMPLE PLOT HEADER RECORDS	27
b.	RECORD TYPE “02” – STANDARD TREE DESCRIPTION RECORD	29
c.	RECORD TYPE “03” – STANDARD REGENERATION RECORD	31
d.	RECORD TYPE “04” – STANDARD GPS RECORDS	32
e.	DETAIL OF PLOT TREATMENT (TYPE 05) - In progress	32
12.	TREE SPECIES CODE	33
13.	PLANT SPECIES CODE	34
14.	COMMON PROBLEMS	48
15.	RECORDING DATA	48
16.	GENERAL INFORMATION (CSTM 101)	49
17.	GENERAL INFORMATION (CSTM 100)	50
18.	EXAMPLE OF A TIE PLATE	51

1. SUGGESTED EQUIPMENT AND SUPPLIES

Accountable – Individual	Accountable - Crew	Expendable
Clinometer	Staff compass	Spiral nails (6.5 cm)
Surveyors hand compass	Axe	Blue tree marking paint
Metal diameter tape	Cloth tape(s) (50m)	Plot posts-metal (corner, regen and sapling)
Loggers tape	Photo holder	Pencils 2H and grease
Topofil	Camera	Geo-flagging tape (orange, blue and yellow)
Biltmore stick	Calipers (tree and seedling)	Aluminum tags-plot corner and tree
Claw hammer	Chainsaw, tool box, stretcher and necessary safety equipment	String
2 lb hammer	Canthook	Loggers crayon
Hammer holder	Whistle	Field notebook
Pocket calculator	Tally Sheets (TM 267, TM 249, Regen and Sectioning)	Paint pens/markers
Douglas protractor	Cruisers Vest	Tie plates
Hand lens	Hard Hat/with shield	
Tallyboard (metal 8x12)	ATV/snowmobile helmet	
Pocket First Aid Kit	Trapper Nelson backpack	
Flare gun with flares and bear bangers	Crayon holder	
Bear repellent	Seedling calipers	
Engineers scale (15 cm)		

2. PLOT TOPOGRAPHY

a. Location

- 1 – Hollow For plots located in local topographic depressions – collecting water
- 2 – Flat For plots located on flat terrain – receiving water
- 3 – Slope For plots on mid and upper slopes – shedding water
- 4 – Hilltop For plots located on ridge crests – shedding water

3. SOIL CODES

a. Erosion potential

- 1 – Slight When 0 to 25% of the area is, or could be eroded in a flood situation.
- 2 – Moderate When 26% to 75% of the area is, or could be eroded in a flood situation.
- 3 – High When more than 75% of the area is, or could be eroded in a flood situation.

b. Drainage Codes

1 – Very rapidly drained	Water is removed from the soil very rapidly relation to supply. Excess water flows downward very rapidly if underlying material is permeable. There may be very rapid subsurface flow during heavy rainfall provided there is a steep gradient. Soils have very low available water storage capacity (usually less than 2.5 cm) within the control section and are usually coarse textured, or shallow, or both. Water source is precipitation.
2 – Rapidly drained	Water is removed from the soil rapidly in relation to supply. Excess water flows downward if underlying material is permeable. Subsurface flow may occur on steep gradients during heavy rainfall. Soils have low available water storage capacity (2.5-4 cm) within the control section, and are usually coarse textured, or shallow, or both. Water source is precipitation.
3 – Well Drained	Water is removed from the soil readily but not rapidly. Excess water flows downward readily into underlying permeable material or laterally as subsurface flow. Soils have intermediate available water storage capacity (4-5m) within the control section, and are generally intermediate in texture and depth. Water source is precipitation. On slopes subsurface flow may occur for short durations but additions are equalled by losses.
4 – Moderately Well	Drained Water is removed from the soil somewhat slowly in relation to supply. Excess water is removed somewhat slowly due to low permeability, shallow water table, lack of gradient, or some combination of these. Soils have intermediate to high water storage capacity (5-6 cm) within the control section and are usually medium to fine textured. Precipitation is the dominant water source in medium to fine textured soils, precipitation and significant additions by subsurface flow are necessary in coarse textured soils.
5 – Imperfectly Supply	Drained Water is removed from the soil sufficiently slowly in relation to keep the soil wet for a significant part of the growing season. Excess water moves slowly downward if precipitation is major supply. If subsurface water or ground water, or both, is main source, flow rate may vary but the soil remains wet for a significant part of the growing season. Precipitation is main source if available water storage capacity decreases. Soils have a wide range in available water supply, texture, and depth, and are gleyed phases of well drained subgroups.
6 – Poorly drained	Water is removed so slowly in relation to supply that the soil remains wet for a comparatively large part of the time the soil is not frozen. Excess water is evident in the soil for a large part of the time. Subsurface flow or ground water flow, or both, in addition to precipitation are main water sources; there may also be a perched water table, with precipitation exceeding evapotranspiration. Soils have a wide range in available water storage capacity, texture, and depth and are gleyed subgroups, Gleysols and Organic soils.
7 – Very poor drained	Water is removed from the soil so slowly that the water table remains drained at or on the surface for the greater part of the time the soil is not frozen. Excess water is present in the soil for the greater part of the time. Ground water flow and subsurface is less important except where there is a perched water table with precipitation exceeding evapotranspiration. Soils have a wide range in available water storage capacity, texture, and depth, and are either Gleysolic or Organic.

4. TREE HEIGHT MEASUREMENT

The height of a tree is defined as the length between the point of germination and the tip of the terminal leader. Heights are measured using a clinometer (with the percent scale) and a 30 or 50m measuring tape. Tree height calculations must be completed on the reverse side of the PSP Tally Sheet (TM 249) (see Figure 4.2A). All data fields are required to be filled in except:

- a) Only Slope distance and slope % or horizontal distance is used.
- b) Correction Factor is only used when the bottom % reading is recorded at a different reference point other than the germination point. ie. DBH height = 1.3m
- c) Check cruise height columns are not to be used by field crew members. This space is only filled in if there has been an actual check cruise completed.

It is very important that field crews understand the process of measuring height so that data is calculated correctly.

At a distance far enough away from the tree to keep the clinometer scale below 100%, take readings for the top % of the tree (tip of the terminal leader), top % to live crown, and the bottom % of the tree (germination point) as shown in Figures 4.1. This may be difficult for deciduous trees as the top of the tree may not be visible through the crown. The slope of the ground must also be measured and recorded if it is greater than 10%.

During remeasurement if the present height is shorter than the past height then a second height must be taken and recorded on the tally sheet directly below the previous measurement. It is advised that the horizontal distance be increased 5 to 10 meters before taking the second reading. It will be up to the cruisers discretion to decide which is the correct height data to be transferred to the front of the tally sheet. Put a line through the height information that is not used **-never erase height information**. On the front of the tally sheet record in comments **HT** to indicate that the height was double-checked in the field.

If live crown height is measured using a metric tape instead of the clinometer, record on the front of the PSP tally sheet (TM 249) in the comments section "height to L.C. measured directly" (Figure 2.14, tree #10).

You should be approximately 25% further than previous height; ie. previous height is 20.2m, you should be 25.2 m, 36.0m should be 43.0m

Also record the slope and slope distance or horizontal distance to the tree. To calculate the tree height, to the nearest 0.1 m, use the following formula:

$$\begin{array}{l} \text{Slope Distance x Slope Correction Factor x} \\ \text{Top reading \% - Bottom reading \%} \\ \text{= Tree Height (m)} \quad \text{_____ x 100\%} \end{array}$$

or

$$\begin{array}{l} \text{Horizontal Distance x} \\ \text{Top reading \% - Bottom reading \%} \\ \text{= Tree Height (m)} \\ \text{_____ 100\%} \end{array}$$

For example, is 22.8 m away on a slope of 15%. The Suunto readings are +80% and +12%. Therefore, the tree is:

$$22.8 \text{ m} \times 0.989 \times \frac{+80 - (+12)}{100} = 22.55 \times 0.68 = 15.33 = 15.3 \text{ m}$$

*Obtained from table in Section 4.5

$$22.8 \times \frac{+80 - (+12)}{100} = 22.8 \times 0.68 = 15.5 \text{ m}$$

There are times when the germination point cannot be seen. In this situation, breast height is often used for the bottom % reading and a correction factor of 1.3 m is added on the calculated total height.

Trees with a lean that require height measurement should have the slope readings taken from a location perpendicular to the lean as shown in Figure 4.2. This will prevent an erroneous measurement that could result in a shorter or taller tree because of the lean.

If at all possible, all height measurements should be taken perpendicular to the slope.

Common errors made during tree height calculations are:

- 1) Misreading $\pm$ signs
 - bottom % readings may either read as positive or negative numbers. Bottom % reading are always subtracted from the top reading regardless on the $\pm$ sign of the number.

ie: if the top % is +90 and bottom % is -3 then:

$$+90 - (-3) = 93$$

if the bottom % reading was +3 then:

$$+90 - (+3) = 87$$

- 2) Not adding the appropriate correction factor, usually 1.3m (breast height). Zero fill if the bottom reading is taken at the germination point, or put line through column.

- a) adding the correction factor at the wrong time:

ie: top %	bottom %	Horizontal distance	Correction factor
+90	-3	20	+1.3

$$\text{Incorrect } ([90 - (-3)] + 1.3) \times .20 = 18.86 = 18.9$$

$$\text{Correct } ([90 - (-3)] \times .20) + 1.3 = 19.90 = 19.9$$

- 3) Miscalculating total height through standard arithmetic errors. Refer to Appendix 4.10.4 for rounding off procedures.

4) Total height information recorded with no calculations.

All tree height calculations must be recorded for each sample tree in the space provided otherwise, the data will be considered invalid and deleted.

TM 249 (Rev. 08/03)

Tree Number	Species	Top %	Top % to Live Crown	Bottom %	Not to Top of Tree	Not to Top of Live Crown	Slope Distance	% Slope	Horizontal Distance	Correction	Total Height	Height to Live Crown	Check Cruise Height	Tree Number
1	SW	92	33	+3	89	30	-	-	20.0	+1.3	19.1	7.3		1
4	SW	84	60	-1	85	61	22.0	13	21.8	-	18.4	13.3		4
7	SW	98	28	0	98	28	-	-	24.0	+1.3	24.8	8.0		7
10	SW	64	10	-2	66	12	-	-	20.0	+1.3	14.5	3.7		10
13	AW	73	63	+1	72	62	-	-	21.0	-	15.1	13.0		13
16	PB	99	80	0	99	80	-	-	25.0	+1.3	26.0	21.3		16
24	SW	No	Height	Forked	Top									

a. Determining Tree Height with a Laser Clinometer or Haglof Vertex

Before using the electronic instrument, be sure that you are familiar with the operators manual and the operation of the tool.

Only total tree height and height to live crown is recorded on the tree tally sheet. Indicate on the back of the sheet which tool was used.

The operator must have the blue direction dot in sight and double check the height of the tree if it does not grow or the height remains the same. Record both sets of numbers on the back of the tally sheets.

5. SLOPE CORRECTION FACTORS AND TABLES

To convert slope distance (S.D.) horizontal distance (H.D.)

$$H.D. = S. D. \times \text{slope distance factor}$$

To convert horizontal distance (H.D. to slope distance (S. D.)

$$S. D. = \frac{H. D.}{\text{slope distance factor}}$$

SLOPE DISTANCE FACTORS

% Slope		% Slope		% Slope	
10	0.995	40	0.928	71	0.819
11	0.994	41	0.925	71	0.815
12	0.993	42	0.922	72	0.812
13	0.992	43	0.919	73	0.808
14	0.990	44	0.915	74	0.804
15	0.989	45	0.912	75	0.800
16	0.987	46	0.908	76	0.796
17	0.986	47	0.905	77	0.792
18	0.984	48	0.902	78	0.789
19	0.982	49	0.898	79	0.785
20	0.980	50	0.894	80	0.781
21	0.979	51	0.891	81	0.777
22	0.977	52	0.887	82	0.773
23	0.974	53	0.883	83	0.769
24	0.972	54	0.880	84	0.766
25	0.970	55	0.876	85	0.762
26	0.968	56	0.872	86	0.758
27	0.965	57	0.869	87	0.754
28	0.963	58	0.865	88	0.751
29	0.960	59	0.861	89	0.747
30	0.958	60	0.857	90	0.743
31	0.955	61	0.854	91	0.740
32	0.952	62	0.850	92	0.736
33	0.950	63	0.846	93	0.732
34	0.947	64	0.842	94	0.729
35	0.944	65	0.838	95	0.725
36	0.941	66	0.835	96	0.721
37	0.938	67	0.831	97	0.718
38	0.935	68	0.827	98	0.714
39	0.932	69	0.823	99	0.711

6. USE AND CONSTRUCTION OF A BILTMORE STICK

Biltmore Sticks are used to measure diameter of saplings or tagged stems <9.1 cm as measuring a stem with a small diameter often results in a broken diameter tape. To use a Biltmore Stick, align "0" on one side of the stem, at breast height, and read the diameter on the other keeping the focal length at 64cm (metric equivalent of 25 in ¹). The following table shows the straight line distance from "0", to be marked on the stick, associated with the corresponding diameter (in centimeters).

¹ Avery, T.E. and H. E. Burkharl. 1983. Forest Measurements, 3rd ed. McGraw-Hill Inc., New York, N.Y. pp. 68-69

DBH	Biltmore	DBH	Biltmore
1.0	0.99	5.2	5.00
1.2	1.19	5.4	5.19
1.4	1.38	5.6	5.37
1.6	1.58	5.8	5.55
1.8	1.77	6.0	5.74
2.0	1.97	6.2	5.92
2.2	2.16	6.4	6.10
2.4	2.36	6.6	6.28
2.6	2.55	6.8	6.46
2.8	2.74	7.0	6.65
3.0	2.93	7.2	6.83
3.2	3.12	7.4	7.01
3.4	3.31	7.6	7.18
3.6	3.50	7.8	7.36
3.8	3.69	8.0	7.54
4.0	3.88	8.2	7.72
4.2	4.07	8.4	7.90
4.4	4.26	8.6	8.07
4.6	4.44	8.8	8.29
4.8	4.63	9.0	8.43
5.0	4.81		

7. ACCESS EVALUATION CODES

1. All Weather Road	All roads in this category are paved or are well traveled gravel roads. These roads are well drained with little possibility of washing out or flooding in heavy rain situations. In the winter, these roads are plowed on a regular basis.
2. Dry Weather Road	This type of road tends to be quite slippery in the spring and fall and becomes heavily rutted when wet. The shoulder on these roads are generally quite soft most of the year. Slopes on these roads should not exceed 10% as they are difficult to drive up or down when wet, even in a four wheel drive vehicle. Minor flooding or washouts can occur but the roads can still be traveled in a four wheel drive vehicle as the roads have solid bottoms.
3. Deteriorating Road	These roads are not used very often and are starting to grow over with grass, small shrubs, or small trees. During heavy rains they can be easily washed out or heavily rutted. It may be very difficult to travel on these roads even with a four wheel drive and the use of an all terrain vehicle should be considered.
4. All Terrain Vehicles only	Included in this category are seismic lines, old trails, and any roads inaccessible using a four wheel drive vehicle. If a plot is more than 1000m along a seismic line or trail, this access is to be indicated. If the distance is less than this, the remeasurement crew can walk to the plot.
5. Helicopter Access.	This access codes should be used only when there is no other way into the plot (i.e. cannot cross river, too far off roads to feasibly drive all terrain vehicle to etc.) It is important to remember to have a suitable location for a helicopter to land and take off from. Keep in mind that openings used for a landing may grow over within 10 years presenting a helicopter to land in the future
6. Unknown	This code is for office use only and is used when access has not been verified and maps do not provide any assistance.

8. LEGAL SURVEY SYSTEM OF ALBERTA

The province of Alberta has been surveyed using a system based on a grid framework. The largest divisions in this system are called meridians. In Alberta there are meridians numbered 4, 5 and 6. The fourth meridian, as shown in Figure 4.19 corresponds to the Alberta Saskatchewan border.

Each meridian has been divided into parcels of land, called townships, 36 square miles (93.2 square kilometers) in size. At six mile (9.654 kilometres) intervals, in a north-south direction, are divisions also called townships and are numbered 1 to 126 starting from the United States border and extending to the Northwest Territories border. The east-west six mile intervals are called "ranges" and are numbered westward from each meridian. The numbering of townships begin in the southeast corner of the province. An example of locating any given township is shown in Figure 4.19.

The grid system is further refined by taking each township and dividing it into 36-one square mile (1.604 square kilometers) parcels of land called "sections". The numbering scheme for each township is shown in Figure 4.19.

The last division in the survey system takes each section and divides it into 16 equal parts called "legal subdivisions" (LSD's). The numbering scheme of each section is demonstrated in Figure 4.19.

If a PSP was located using the example in Figure 4.19, the legal land description would be recorded as 13-1-87-18-4. This translates to Legal Subdivision 13 of Section 1 in Township 87, Range 18, West of the Fourth Meridian. If a PSP is located in two or more sections/legal subs, all legal descriptions are tallied and the location of plot enter is noted, i.e. if a PSP was located at the cross section of legal subs 11, 12, 13, and 14 then the following information would be tallied.

Recorded on Maintenance Sheet as:

11-1-87-18-4

12-1-87-18-4

13-1-87-18-4.1 (this would be recorded as the correct legal descriptions)

14-1-87-18-4

The center of the PSP falls in L.S. 13-recorded on the PSP Header Sheet (TM 267) during establishment remeasurement.

9. FIELD SHEET HANDBOOK

a. Tally Sheet Instructions

The PSP field handbook contains instructions pertaining to data entry on the tally sheet, a summary of allowable errors, and a plot measurement summary.

b. General Information (Columns 1 – 37)

This information is recorded on the Header Sheet and every Tally Sheet.

Column 1-2

Agency: numeric, right justified, zero filled. As listed on the Agency list.

Column 3-12

Group Number: numeric, right justified, zero filled.

e.g.

0	2	0
---	---	---

 = Group 20

Column 13

Sub Plot Number: numeric (1,2,3,4)

e.g.

1

 = sub plot 1

Column 14-15

Measurement Number: numeric, '00' for an establishment plot, subsequent measurements would be 01, 02, 03, etc.

e.g.

0	3
---	---

 = 3rd measurement

Column 16-18

Year: numeric; 1961

e.g.

1	9	6	1
---	---	---	---

Column 20-21

Month: numeric, right justified, May=05, June=06, etc., zero filled.

e.g.

0	6
---	---

 = June

Column 22-23

Day: numeric, right justified, zero filled.

e.g.

0	3
---	---

 = 3rd day

Column 24-25

L.S.: (Legal Subdivision), numeric, right justified, zero filled, only use if known.

e.g.

1	3
---	---

 = Legal subdivision 13

Column 26-27

Section: numeric, right justified, zero filled.

e.g.

0	2
---	---

 = Section 2

Column 28-30

Twp.: (Township), numeric, right justified, zero filled.

e.g.

0	3	2
---	---	---

 = Township 32

Column 31-32

Rge.: (Range), numeric, right justified, zero filled.

e.g.

0	3
---	---

 = Range 3

Column 33

M.: (Meridian), numeric.

e.g.

5

 = West of the 5th meridian

Column 34-36

Plot Type: Numeric, and left justified. First column is primary description. The remaining two columns are not defined yet; zero fill until further notice.

e.g.

3	0	0
---	---	---

 = Thinned PSP

e.g.

0	0	0
---	---	---

 = An untreated fire origin plot

Code

- 0 = No treatment
- 1 = Growth Intercept Plot
- 2 = Fertilized
- 3 = Thinned
- 4 = Drained
- 5 = Burned
- 6 = Mistletoe
- 7 = Immature (16-55 years)
- 8 = Paired Plot
- 9 = University of Alberta Plot

Column 37

Imp.: (Imperial), "X" if information is collected in imperial units; blank if collected in metric.

e.g.

X

 = Indicates measurements done in imperial units.

c. Record 01 – Header Information (Columns 38-212)

Column 38-39

Record Type: previously filled as 01

Column 40-60

Plot Sizes (Tree, Sapling and Regen): numeric, recorded in square metres right justified. Plot sizes are determined in the office using the horizontal distances recorded on the plot maintenance form.

e.g.

		1	0	0	0	.0
--	--	---	---	---	---	----

 = 1,000.0 m²

Column 61-71

Phase 1, 2 or 3 Overstorey - See PSP Header information for description.

Column 72-93

AVI Interpreted Overstorey: numeric and alpha using AVI ecological moisture regime, crown closure class, height and tree species composition.

Ecological moisture regime can be numeric or character. Small cap letter goes in column 72. If using numeric label this goes in column 73. Do not input both. Only one entry needed.

Ecological Moisture Regime Labels

Moisture Regime	Database Numeric Label	Database Character Label
Very xeric	0	d - dry
Xeric	1	d - dry
Subzeric	2	d - dry
Submesic	3	m – mesic (default)
Mesic	4	m – mesic (default)
Subhygric	5	m – mesic (default)
Hygric	6	w - wet
Subhygric	7	w - wet
Hygric	8	a - aquatic

Crown closure class can be numeric or character. Capital letter goes in column 74. If using numeric, label goes in column 75. Do not use both. Only one entry needed.

Crown Closure Class Labels

Crown Closure	Database	Database
---------------	----------	----------

Class (%)	Numeric Label	Character Label
01-05		V
06-10	0	A
11-20	1	A
21-30	2	A
31-40	3	B
41-50	4	B
51-60	5	C
61-70	6	C
71-80	7	D
81-90	8	D
91-100	9	D

Height is recorded to the closest metre and is numeric and goes in columns 76 and 77. Input as 2 digits i.e.: 9m = 09.

Tree species composition is recorded in 10% classes and must add up to 100%. A maximum of 5 species. First letter is capitalized and second letter is small.

Species 1 (Alpha) in Columns 78-79

Species 1 % in Columns 80-81

Species 2 (Alpha) in Columns 82-83

Species 2 % in Columns 84

Species 3 (Alpha) in Columns 85-86

Species 3 % in Columns 87

Species 4 (Alpha) in Columns 88-89

Species 4 % in Columns 90

Species 5 (Alpha) in Columns 91-92

Species 5 % in Columns 93

e.g.

d	1	A	2	0	9	S	w	0	9	P	I	1		
---	---	---	---	---	---	---	---	---	---	---	---	---	--	--

Column 37

Imp: (Imperial), "X" if information is collected in imperial units; blank if collected in metric.

e.g.

☐ X = indicates measurement done in imperial units.

Column 94-108

Phase 1, 2 or 3 Understorey – See PSP Header information for description.

Column 109-128

AVI Interpreted Understorey: numeric and alpha using AVI crown closure class, height and tree species compositions.

Crown closure class can be numeric or a character. Capital letter goes in column 109. If using numeric, label goes in column 110. Do not use both. Only one entry needed.

Height is recorded to the closest metre and is numeric and goes in columns 111 and 112. Input as 2 digits i.e.: 9m=09.

Tree species composition is recorded in 10% classes and must add up to 100%. A maximum of 5 species. First letter is capitalized and second letter is small.

Species 1 (Alpha) in Columns 113-114

Species 1 % in Columns 115-116

Species 2 (Alpha) in Columns 117-118

Species 2 % in Columns 119

Species 3 (Alpha) in Columns 120-121

Species 3 % in Columns 122

Species 4 (Alpha) in Columns 123-124

Species 4 % in Columns 125

Species 5 (Alpha) in Columns 126-127

Species 5 % in Columns 128

e.g.

A	2	0	9	S	W	0	9	P	I	1
---	---	---	---	---	---	---	---	---	---	---

Column 129

Location: numeric

Code

- 1 = hollow
- 2 = flat
- 3 = slope
- 4 = hilltop

e.g.

3

 = plot location on a slope

Column 130-132

% slope: average percent slope for the plot, numeric characters, right justified.

e.g.

0	1	5
---	---	---

 = 15% slope

Column 133-134

Aspect: alpha characters, right justified

Code

N = North

S = South
 E = East
 W = West
 NE = Northeast
 NW = Northwest
 SE = Southeast
 SW = Southwest
 NA = Non Applicable (Slope = "0")

e.g.

--

 N = North

Column 135-138

Elevation: elevations in metres, numeric characters, right justified (convert feet to metres by multiplying by .3048).

e.g.

1	0	9	7
---	---	---	---

 = 1097 m

Column 139

Erosion Potential: numeric. This describes the chance of water eroding down to or into the mineral soil layer and is based upon water flow, slope and soil type.

Code

- 1 = Slight – when 0 to 25% of the area is, or could be eroded in a flood situation.
- 2 = Moderate – when 25% to 75% of the area is, or could be eroded in a flood situation.
- 3 = High – when more than 75% of the area is, or could be eroded in a flood situation.

e.g.

2

 = moderate erosion potential

Column 140

Drainage: numeric, for definitions of codes see Appendix 4.3 of the PSP Manual

Code

- 1 = Very rapidly drained
- 2 = Rapidly drained
- 3 = Well drained
- 4 = Moderately well drained
- 5 = Imperfectly drained
- 6 = Poorly drained
- 7 = Very poorly drained

e.g. 6 = poorly drained soil

Column 141-142

Depth to Mineral Soil: numeric, right justified, measured in cm or inches.

If greater than 99 cm, fill with 99. This will indicate a depth to mineral soil of 99 or greater.

e.g.

1	0
---	---

 = 10cm

Column 143

Surface Vegetation-Type: numeric, nine possible types or combinations of grass, lichen/=moss, herbs (flowers and non-woody stems) and shrubs (woody stems). The dominant type is recorded.

Code

- 1 = grass
- 2 = grass + herbs
- 3 = grass + shrubs
- 4 = lichen/moss
- 5 = lichen/moss + herbs
- 6 = lichen/moss + shrubs
- 7 = herbs
- 8 = herbs + shrubs
- 9 = shrubs

e.g.

4

 = lichen/moss

Column 144 -146

Surface Vegetation - % Ground Cover: numeric, right justified. Tally the percent of ground that is covered by surface vegetation type.

e.g.

	9	5
--	---	---

 = 95% of ground covered by surface vegetation type

Column 147

Maintenance Report – Access: numeric (for definitions of Access codes see Appendix 4.8).

Code

- 1 = all weather road
- 2 = dry weather road
- 3 = deteriorating road
- 4 = all terrain vehicle
- 5 = helicopter
- 6 = unknown

Column 148

Plot Damage: numeric

- 1 = none
- 2 = natural damage
- 3 = manmade damage
- 4 = natural and manmade damage
- 5 = closed
- 6 = (this code is not used)
- 7 = plot was previously closed but has been reopened (office use only)

Column 149

Buffer Damage: numeric

- 1 = none
- 2 = natural damage
- 3 = manmade damage
- 4 = natural and manmade damage
- 5 = closed
- 6 = damage inside buffer greater than 20m from subplot
- 7 = plot was previously closed but has been reopened (office use only)

e.g. Comments: 2. (access) 1. (plot damage)
3. (buffer damage)

= dry weather road, none, manmade damage

Column 150-153

Establishment Year: numeric

e.g.

1	9	6	1
---	---	---	---

 = 1961
Column 164

Plot Configuration

Column 180-183

Stand Origin

Column 184

Site Index/TPR: character

Column 185-186

Site Index/TPR: numeric

Column 187-190

Photo year

Column 191-198

Ecosite

Column 199-201

Aspect in Degrees

Column 202

Nutrient Regime

Column 203-204

Natural Region

Column 203-208

Minimum diameter Measured (mm)

Column 209-210

Stump Height (cm)

Column 211

Surface Expression

Column 212

Slope Position

d. Record 02 – Tree Information – header Sheet (Columns 1-37) – Tally Sheet (Columns 38-124)**Column 38-39**

Record type: previously filled as 02

Column 40-43

Tree Number: numeric, right justified.

Tree numbers will be:

1. Consecutive i.e. 1 through 9997 (saplings included).
2. 0000 – representing a tree felled outside the plot to determine stump age and increments (pre-filled on Header Sheet).

e.g.

0	0	9	8
---	---	---	---

 = tree #98
Column 44-45

Species: tree species, alpha characters

Code

FA = Alpine fir

FB = Balsam fir
 FD = Douglas fir
 LA = Alpine larch
 LT = Tamarack
 LW = Western larch
 PF = Limber pine
 PJ = Jack pine
 PL = Lodgepole pine
 PW = Whitebark pine
 SB = Black spruce
 SE = Englemann spruce
 SW = White spruce
 AW = Aspen
 PB = Balsam poplar
 BW = White birch
 NO = Indicates No trees to tally in regeneration plot

e.g.

A	W
---	---

 = Aspen

Column 46-49

DBH: (Diameter at Breast Height), one decimal place, numeric, right justified. In the computer the format is in millimeters (No Decimal).

e.g.

	3	2	.1
--	---	---	----

 = 32.1 cm dbh

Column 50-53

Height: (Tree Height), one decimal place, numeric, right justified. In the computer the format is in decimeters (No Decimal).

e.g.

	3	0	.6
--	---	---	----

 = 30.6 m tree height

Column 54-57

Height to Live Crown: (Tree Height from base to point of first branch of the crown), one decimal place, numeric, right justified. In the computer the format is in decimeters (No Decimal).

e.g.

	1	0	.5
--	---	---	----

 = 10.5 m from tree base to live crown

Column 58

C.C. (Crown Class), alpha character

Code

D	Dominant	-crowns extend above the general level of the canopy
---	----------	--

C	Codominant	-crowns form the general level of the canopy
I	Intermediate	-crowns below but extending into the bottom of the general level of the canopy
S	Suppressed	- crowns entirely below the general level of the canopy
O	Open-grown	-if the trees branches does not interact with another trees branches.

e.g. S = suppressed tree

Column 59-64

Condition Codes: (Tree condition codes) numeric. If the tree has no defect, record 00. Zero filling is not required. **See Master Condition Code List**

e.g.

5	1	5	4		
---	---	---	---	--	--

 = tree with conks and a fork

e.g.

2	9				
---	---	--	--	--	--

 = tree cut down

e.g.

6	3	5	6	6	6
---	---	---	---	---	---

 = tree with a stem disease, broken and a stem form defect

If there is not visible defect, a condition code 00 is recorded.

The following columns 65-76 are on Header Sheet only (TM 267)

Column 65-67

DBH Age: (Tree DBH age), numeric characters, right justified, zero filled

e.g.

1	1	6
---	---	---

 = 116 yrs of age at DBH (1.3 m)

Column 68-70

Stump Age: (Tree stump age), numeric characters, right justified, zero filled.

e.g.

1	2	1
---	---	---

 = 121 yrs of age at stump (.3 m)

Column 71-76

Stump Increment Width: increment width for the last 20 yrs. growth done in 2 measurements: 0 – 10 yrs, 11-20 yrs, numeric characters, right justified, one decimal place, zero filled. Entered as mm in digital file.

e.g.

0	2	3	0	1	5
---	---	---	---	---	---

= 2.3 cm increment for 0 – 10 yrs
 = 1.5 cm increment for 11 – 20 yrs

Column 77-79

Azimuth: azimuth of tree from plot center, numeric, right justified, 1 to 360°

e.g.

0	2	0
---	---	---

 = 20⁰ from the plot centre

Column 80-82

Distance: distance of tree from plot center, numeric, right justified one decimal place. In the computer, the format is in decimeters (No Decimal).

e.g.

0	6	.3
---	---	----

 = 6.3 m from plot center

Column 83-89

Tree Plot size (m²): numeric, right justified, go to one decimal place. Zero filled.

Tree Plot Size (m²): numeric, right justified, go to one decimal place. Zero filled.

Column 97-103

Regen Plot Size (m²)

Column 104-107

Establishment Year: numeric, right justified.

Column 108

Crown Status: "Y" indicates tree needs crown measurements. "Blank" indicates no crown measurement required.

Column 109-112

Crown Width North: recorded in decimeters (dm). Right justified.

Column 113-116

Crown Width West: recorded in decimeters (dm). Right justified.

Column 117-120

Crown Width West: recorded in decimeters (dm). Right justified.

Column 121-124

Crown Width East: recorded in decimeters (dm). Right justified.

e. *Record 03 – Regen Information (Columns 38-85)*

Column 38-39

Record Type: previously recorded as 03.

Column 40-43

Tree number: previously recorded as 9998.

Column 44-45

Species: tree species, alpha characters.

Column 40-48, 49-51, 52-54, 55-57, 58-60

Total (1-5): numeric, right justified, zero filled. Add in 3 plot sizes and establishment year from Page 41: numeric right justified, zero filled.

e.g.

0	0	2
---	---	---

 = 2 regeneration of that species in that height class
Column 61-67

Tree Plot Size (m²): right justified.

e.g.

		1	0	0	0	.0
--	--	---	---	---	---	----

 = 1,000.0 m²
Column 68-74

Sapling Plot Size (m²): right justified.

e.g.

			2	5	0	.0
--	--	--	---	---	---	----

 = 250.0 m²
Column 75-81

Regen Plot Size (m²): right justified.

e.g.

				6	2	.0
--	--	--	--	---	---	----

 = 62.0 m²
Column 81-85

Establishment Year: right justified.

e.g.

1	9	6	0
---	---	---	---

 = 1960
f. Summary of Allowable Errors

Item	Allowable Error
Location of Plot Centre (Tie Point and Tie Line)	- 2% of the cruise line horizontal distance (e.g. 3m of a 150m tie line H.D.)
Plot Size (Sides) – Tree Plot	- 31.62 ± .25m (31.37 – 31.87m)
	38.73 ± .31m (38.72 – 39.04m)
	44.72 ± 0.36m (44.36 – 45.08m)
– Sapling/Regen Plot	- 7.90 ± .06m (7.84 – 7.96m)
	9.69 ± .08m (9.61 – 9.77m)
	11.09 ± .09m (11.09 – 11.27m)

Plot Size (bearings)	$\pm 2^\circ$
No. of Trees Talled and Species Identified	Tree plot - none Sapling plot - $\pm 5\%$ of total (e.g. 2- saplings ± 1) – no allowable error in identification Regen plot - $\pm 10\%$ (e.g. 20 regen ± 2)
DBH	Diameter Breast height – 1.3m $\pm$ 6.5cm ($\pm 5\%$) diameter – ± 1.0 cm
Height and Height to Live Crown	$\pm 3\%$ (e.g. 16.2m $\pm$.5m)
Crown Class and Condition Codes	10% of stems tallied may have incorrect crown class or condition codes (e.g. 10 trees of 100 tallied)
Stem Mapping	Azimuth - $\pm 2^\circ$ Distance - ± 0.5 m
Section Lengths	± 5 cm after “cookie” is cut
Perpendicular Cuts	$\pm 10^\circ$ from perpendicular
DHB and Stump Age	Conifers - ± 1 year Deciduous - ± 5 years
Increment Width	$\pm 5\%$ for each set of years

g. Plot Measurement Summary

Plot	Length of Side (m)	Area (ha)	Area (m ²)	Buffer Side (m)	Buffer Area ((m ²))
Tree	31.62	0.100	1,000	200	40,000
	38.73	0.150	1,500	237	56,250
	44.72	0.200	2,000	300	90,000
Sapling/Regen	7.90	0.006	62		
	9.69	0.009	94		
	11.18	0.012	125		

Plot	Stem Size	Tagged	Number Recorded
Tree	≥ 9.1 cm DBH	Yes	0001
			to 7999
Sapling	1.3 m in height	Yes	8001
Regen	≥ 0.10 height up to a height of ≤ 1.29 m	No	9998
Outside of Plot	Representatives of stand age	No	0000

h. Rounding-Off

The following rules are to be followed for rounding off height measurements on sample trees.

- 1) Delay rounding off numbers as long as possible, preferably to the last stage, as rounding can significantly affect the accuracy of the final answer.

- 2) If the digit to be rounded is followed by a digit greater than 5, round up.

e.g. 10.66 10.7
16.394 16.4 (6, 9 & 7 are greater than 5)
21.17 21.2

- 3) If the digit to be rounded is followed by a digit less than 5, leave as is.

e.g. 10.64 10.6
10.339 16.3 (4, 3 & 1 are less than 5)
21.11 21.1

- 4) If the digit to be rounded is followed by a 5 then:

- a) If the digits following the 5 are greater than 0, round up.

e.g. 10.651 10.7
16.359 16.4
21.154 21.2

- b) If the digit being rounded is followed by a 5 and no other digits then:

- i) If the digit being rounded is odd, round up.

10.55 10.6
16.35 16.4
21.15 21.2

- ii) If the digit being rounded is even, leave as is.

10.65 10.6
16.45 16.4
21.25 21.2
25.5 25.0

NOTE: The number 0 is considered to be an even number.

10. MISTLETOE AND IMMATURE PLOT TYPES

In September 1989 two new plot types were introduced:

- i) 600 – mistletoe plot type
- ii) 700 - immature plot type (16 - 55 years old)
- iii) The mistletoe plot type is established and re-measured according to the procedures outlines in Section 2.0. The severity of the mistletoe on the infected stem(s) is recorded using the 6 digit Hawksworth Mistletoe Rating System (see Figure 4.12 for rating system). Trees are sectioned in accordance with the A.F.S. Tree Sectioning Procedures Manual (see also Section 2.1.4.7).

Starting in 2000 field season in the immature plot, we do not have a minimum DBH requirement. All trees $\geq 1.3\text{m}$ in height will be tagged and measured in the 1/16 sapling/regen plot.

- iv) In the remainder of the plot, all trees $\geq 9.1\text{cm}$ at DBH are numbered and measured. Diameters, crown class and condition codes are assigned to each tree stem. Heights are measured according to procedures outlined in Section 2.1.4.4. All tagged trees are stem mapped.

NOTE: On the stems that are too small to withstand a nail, DBH is marked with a blue painted band so that future measurements are taken at the same point. A numbered tree tag is then placed loosely around the stem or a branch so as to prevent girdling of the stem. Seedling calipers or a baltimore stick is used to take sapling diameters.

When sectioning the immature plots three representative trees within the buffer are cut. When bucking, use 1.0m section lengths.

In very dense stands, trees on the boundary of the tree plot, but not in the plot can have a yellow or orange vertical strip painted on them, facing into the tree plot. This will assist future re-measurements crews in determining border line trees.

11. ASCII FORMAT OF PSP TREE DATA, RECORDS 1, 2 AND 3

STANDARD SAMPLE PLOT FILE FORMATS KEYPUNCH AND MASTER FILES (Format as of February, 2000)

There are 4 record types found in these standard sample plot files:

- a) Plot Header records (Type 01)
- b) Tree description Records (Type 02)
- c) Regeneration Records (Type 03)
- d) GPS Records (Type 04)
- e) Detail of Plot Treatment (Type 05)

All numeric variables are right justified and character variables are left justified.

Variable's abbreviated name is located before the variable description.

Record types are identified by a 2 digit code found in columns 38-39. A list of variables found in each record type is described below:

There are 15 different plot types and are as follows: Please advise FMD if plot type is not on list and put plot type in next available 2 digit position. Zero filled:

Ptype: plot type

- 1.) PSP Regular
- 2.) PSP Stand Dynamics Survey
- 3.) PSP Reforestation Monitor Plot
- 4.) TSP Operational Cruise
- 5.) TSP Inventory
- 6.) TSP Large Scale Photography
- 7.) Mistletoe
- 8.) Thinned
- 9.) Fertilized
- 10.) Herbicide
- 11.) Pruned
- 12.) Spruce Budworm
- 13.) Cleaned
- 14.) Thinned and Fertilized
- 15.) Herbicide and Thinned

There are 3 different plot configuration types and are as follows: Please advise FMD if plot configuration is not on list and put plot configuration in next available position.

Pconfig: plot configuration

- Fixed area (Any shape) - 1
- Fixed area subplot - 2
- Variable radius (Prism) plot - 3

There are 34 different Agency codes but this can be expanded. Please advise FMD if agency is not on list and put agency in next available 2-digit position. Zero filled. List is as follows:

Agency

01 – Alberta Forest Service	02 – B.C. Forest Service
03 – Saskatchewan Forest Service	04 – Pedology Consulting
05 – B.C. Forest Products.	06 – Proctor and Gambler
07 – Champion Forest Products	08 – Canfor (N.C.F.I)
09 – Ziedler Plywood Ltd.	10 – Makin Consulting
11 – Montreal Engineering	12 – Blue Ridge Lumber Co.
13 – Pelican – Spruce Sawmill	14– Special Projects
15 – University of Alberta	16 – Canadian Forest Service
17 – Alberta Newsprint	18 – Alpac
19 – Canadian Forest Products	20 – Daishowa-Marubeni International
21 – Manning Diversified Forest Products	22 – Millar Western
23 – Slave Lake Pulp Corp	24 – Spray Lake Sawmill
25 – Sundance	26 – Sunpine
27 – Tolko – High Prairie	28 – Tolko – High Level
29 – Vanderwell	30 – Weldwood
31 – Weyerhaeuser – Drayton Valley	32 – Weyerhaeuser - Edson
33 - Weyerhaeuser – Grande Prairie	34 – Weyerhaeuser – Saskatchewan
35 – Weyerhaeuser – Slave Lake	

There are 3 different measurement types (number) and are as follows:

Measure: measurement number

TSP Always = 0

PSP Establishment = 0

PSP Re-measurement = 1, 2, 3, etc.

In Record Type “01”, subplot variable indicates if there is subplots within the plot. If “No” subplots (1 in record Type 1) then Subplot variable in other records left blank. Otherwise enter appropriate subplot number.

a. RECORD TYPE "01" – STANDARD SAMPLE PLOT HEADER RECORDS

Column	Description	ASCII Format
1-2	Ag: Agency	I2
3-12	Plot: Plot number	I10
13	Subplot: Subplot number (1 = No subplots and 2 = Subplots)	I1
14-15	Measure: Measurement number	I2
16-19	Year: Year of measurement	I4
20-21	Month: Month of measurement	I2
22-23	Day: Day of measurement	I2
24-25	LSD: Legal subdivision	I2
26-27	Sec: Section	I2
28-30	Twp: Township	I3
31-21	Rge: Range	I2
33	Mer: Meridian	I1
34-36	Ptreat: Plot treatment	I3
37	Imp: Imperial	A1
38-39	Recty: Record Type (Always '01')	I2
40-46	Plotsize: Plot size (m ²)	F7.1
47-53	Sapsize: Sapling Plot size (m ²)	F7.1
54-60	Regsize: Regen Plot size (m ²)	F7.1
61-71	Phase 1, 2 or 3 Overstorey	A11
61	Dense: Overstorey Density class	A1
62	Htphase: Overstorey Height class	I1
63-64	Spph1: Overstorey Species 1	A2
65-66	Spph1: Overstorey Species 2	A2
37-38	Spph1: Overstorey Species 3	A2
69-70	Brack: Overstorey Bracketed Species	A2
71	Com: Overstorey Stand Commercialism	A1
72-93	AVI Interpreted Overstorey	A22
72	Moistch: Moisture Regime (Alpha-small letter)	A1
73	Moistnu: Moisture Regime (Numeric)	I1
74	Crcch: Crown Closure Class (Alpha-cap letter)	A1
75	Crcnu: Crown Closure Class (Numeric)	I1
76-77	Htc: Height to closest metre	I2

78-79	Sp1: Species 1 (Capital + small)	A2
80-81	Per1: Species 1 Percent (10% classes)	I2
82-83	Sp2: Species 2 (Capital + small)	A2
84	Per2: Species 2 Percent (10% classes)	I1
85-86	Sp3: Species 3 (Capital + small)	A2
87	Per3: Species 3 Percent (10% classes)	I1
88-89	Sp4: Species 4 (Capital + small)	A2
90	Per4: Species 4 Percent (10% classes)	I1
91-92	Sp5: Species 5 (Capital + small)	A2
93	Per5: Species 5 Percent (10% classes)	I1
94-108	Phase 1, 2 or 3 Understorey	A15
94	Denseu: Understorey Density class	A1
95	Htphaseu: Understorey Height class	I1
96-97	Spph1u: Understorey Species 1	A2
98-99	Spph2u: Understorey Species 2	A2
100-101	Spph3u: Understorey Species 3	A2
102-103	Bracku: Understorey Bracketed	A2
104	Commu: Understorey Stand Commercialism	A1
105-108	Soringinu: Understorey Stand Origin	I4
109-128	AVI Interpreted Understorey	A20
109	Crcchu: Crown Closure Class (Alpha-cap letter)	A1
110	Crcchu: Crown Closure Class (Numeric)	I1
111-112	Hrcu: Height to closest metre	I2
113-114	Spu1: Species 1 (Capital + small)	A2
115-116	Peru1: Species 1 Percent (Capital + small)	I2
117-118	Spu2: Species 2 (Capital + small)	A2
119	Peru2: Species 2 Percent (Capital + small)	I1
120-121	Spu3: Species 3 (Capital + small)	A2
122	Peru3: Species 3 Percent (Capital + small)	I1
123-124	Spu4: Species 4 (Capital + small)	A2
125	Peru4: Species 4 Percent (Capital + small)	I1
126-127	Spu5: Species 5 (Capital + small)	A2
128	Peru5: Species 5 Percent (Capital + small)	I1
129	Locate: Location	I1
130-132	Slope: Slope percent	I3

133-134	Aspect: N, W, NE, NW, etc. NA= No Aspect	A2
135-138	Elev: Elevation (m)	I4
139	Eros: Erosion potential	I1
140	Drain: Drainage	I1
141-142	DMS: Depth to mineral soil (cm)	I2
143	SVT: Surface vegetation type	I1
144-146	Gnd: Ground cover percent	I3
147	Access	I1
148	Pldam: Plot damage	I1
149	Budam: Buffer damage	I1
150-153	Estyr: Establishment Year	I4
162-163	Ptype: Plot Type	I2
164	Pconfig: Plot Configuration	I1
180-183	Storigin: Stand origin	I4
184	SIA: Site Index/TPR	A1
185-186	SII: Site Index/TPR	I2
187-190	Photo: Photo year	I4
191-193	Struct: Stand Structure	A3
194-198	Ecosite	A5
199-201	AspDeg: Aspect in Degrees	I3
202	Nutri: Nutrient Regime	A1
203-204	Nregion: Natural region	I2
205-208	Mindbh: Minimum Diameter measured (mm)	I4
209-210	Stumpht: Stump Height (cm)	I2
211	Surf: Surface Expression	A1
212	Slopepos: Slope Position	A1

b. RECORD TYPE "02" – STANDARD TREE DESCRIPTION RECORD

<u>Column</u>	<u>Description</u>	<u>ASCII Format</u>
1-2	Ag: Agency	I2
3-12	Plot: Plot number	I10
13	LPSPSub: LFS PSP Subplot number	I1
14-15	Measure: Measurement number	I2
16-19	Year: Year of measurement	I4

20-21		Month: Month of measurement	I2
22-23		Day: Day of measurement	I2
24-25		LSD: Legal subdivision	I2
26-27		Sec: Section	I2
28-30		Twp: Township	I23
31-32		Rge: Range	I2
33		Mer: Meridian	I1
34-36		Ptreat: Plot treatment	I3
37		Imp: Imperial	A1
38-39		Recty: Record Type (Always '02')	I
40-43		Tree: Tree number	I
44-45		Sp: Species (Caps)	A2
<u>On No Tally Plots Enter Tree = 0001 and Sp= NO</u>			
46-49		Dbh: Diameter Breast Height (mm)	I4
50-53		Ht: Height (dm)	I4
54-57		Htlcm: Height to live crown (dm)	I4
58		Cclass: Crown Class	A1
59-60		Cond1: Condition Code 1	I2
61-62		Cond2:Condition Code 2	I2
63-64		Cond3:Condition Code 3	I2
65-67		Dbhage: Diameter Breast Age	I3
68-70		Stumpage: Stump Age	I3
71-73		Inc10: Increment Prev. 10 yrs (mm)	I4
74-76		Inc20: Increment Prev. 11-20 yrs (mm)	I4
77-79		Azimuth	I3
80-82		Distance (dm)	I3
83-89		Plotsize: Plot/subplot size (m ²)	F7.1
90-96		Sapsize: Sapling Plot size (m ²)	F7.1
97-103		Regsize: Regen Plot size (m ²)	F7.1
104-107		Estyr: Establishment year	I4
108		Crstat: Crown status	A1
109-112		Crn: Crown Width North (dm)	I4

113-116		Crw: Crown Width West (dm)	I4
117-120		Crs: Crown Width South (dm)	I4
121-124		Cre: Crown Width East (dm)	I4
125-126		Subplot: Subplot number	I2
127-141		Stand: Stand/Polygon number	I15
142-146		Stump: Stump Diameter (mm)	I5

c. RECORD TYPE "03" – STANDARD REGENERATION RECORD

<u>Column</u>		<u>Description</u>	<u>ASCII Format</u>
1-2		Ag: Agency	I2
3-12		Plot: Plot number	I10
13		LPSPSub: LFS PSP Subplot number	I1
14-15		Measure: Measurement number	I2
16-19		Year: Year of measurement	I4
20-21		Month: Month of measurement	I2
22-23		Day: Day of measurement	I2
24-25		LSD: Legal subdivision	I2
26-27		Sec: Section	I2
28-30		Twp: Township	I3
31-32		Rge: Range	I4
33		Mer: Meridian	I1
34-36		Ptreat: Plot treatment	I3
37		Imp: Imperial	A1
38-39		Recty: Record Type (Always '03')	I2
40-43		Tree: Tree number	I4
44-45		Sp: Species (Caps)	A2
(On No Tally Plots Enter Sp=NO)			
46-48		Regen 1: Regeneration in height class 1	I3
49-51		Regen 2: Regeneration in height class 2	I3
52-54		Regen 3: Regeneration in height class 3	I3
55-57		Regen 4: Regeneration in height class 4	I3
58-60		Regen 5: Regeneration in height class 5	I3

61-67		Plotsize: Plot size (m ²)		F7.1
68-74		Sapsize: Sapling Plot size (m ²)		F7.1
75-81		Regsiz: Regen Plot size (m ²)		F7.1
82-85		Estyr: Establishment Year		I4
86-87		Subplot: Subplot number		I2

d. RECORD TYPE '04' – STANDARD GPS RECORDS

<u>Column</u>		<u>Description</u>		<u>ASCII Format</u>
1-2		Ag; Agency		I2
3-12		Plot: Plot number		I10
13-14		Sec: Section		I2
15-17		Twp: Township		I3
18-19		Rge: Range		I2
20		Mer: Meridian		I1
21-22		Subplot: Subplot number		I2
23-37		Stand: Stand/Polygon number		I15
38-39		Reacty: Record Type (Always '04')		I2
40-44		Declong: Longitude Dec. Degrees		F15.10
45-59		Declat: Latitude Dec. Degrees		F15.10
60-70		Easting		F11.5
71-81		Northing		F11.5
82-85		UTM		I4
86-87		NAD (Preferable 83)		I2

e. DETAIL OF PLOT TREATMENT (TYPE 05) - On progress

12. TREE SPECIES CODE

Only the following species will be measured:

Common Name Genus/Species	Scientific Name Code	Species
<u>Fir</u>	<u>Abies</u>	
Alpine fir	A. lasiocarpa	FA
Balsam fir	A. balsamea	FB
<u>Birch</u>	<u>Betula</u>	
White birch	B. papyrifera	BW
<u>Douglas-fir</u>	<u>Pseudotsuga</u>	
Douglas-fir	P. menziesii	FD
<u>Larch</u>	<u>Larix</u>	
Alpine larch	L. lyalii	LA
Tamarack	L. laricina	LT
Western larch	L. occidentalis	LW
<u>Pine</u>	<u>Pinus</u>	
Limber pine	P. flexilis	PF
Jack pine	P. banksiana	PJ
Lodgepole pine	P. contorta	PL
Whitebark pine	P. albicaulis	PW
<u>Poplar</u>	<u>Populus</u>	
Aspen (White Poplar)	P. tremuloides	AW
Balsam poplar (Black poplar)	P. balsamifera	PB
<u>Spruce</u>	<u>Picea</u>	
Black spruce	P. mariana	SB
Englemann spruce	P. englemannii	SE
White spruce	P. glauca	SW

13. PLANT SPECIES CODE

TREE LAYER

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
ABIE BAL	<i>Abies balsamifera</i>	Balsam Fir
ABIE LAS	<i>Abies lasiocarpa</i>	Alpine Fir
BETU PAP	<i>Betula papyrifera</i>	White Birch
DC		Dead conifer
DD		Dead Deciduous
LARI OCC	<i>Larix occidentalis</i>	Western Larch
LARI LAR	<i>Larix laricina</i>	Larch
LARI LYA	<i>Larix lyallii</i>	Alpine Larch
PICE MAR	<i>Picea mariana</i>	Black Spruce
PICE GLA	<i>Picea glauca</i>	White Spruce
PICE ENG	<i>Picea engelmannii</i>	Engelman Spruce
PINU CON	<i>Pinus contorta</i>	Lodgepole Pine
PINU FLE	<i>Pinus flexilis</i>	Limber Pine
PINU ALB	<i>Pinus albicaulis</i>	Whitebark Pine
POPU BAL	<i>Populus balsamifera</i>	Balsam Poplar
POPU TRE	<i>Populus tremuloides</i>	Aspen
PSEU MEN	<i>Pseudotsuga menziesii</i>	Douglas Fir

SHRUB LAYER

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
ARCT UVA	<i>Arctostaphylos uvaursi</i>	Bearberry, Kinnickinnick
ALNU CRI	<i>Alnus crispa</i>	Green Alder
ALNU TEN	<i>Alnus tenuifolia</i>	River Alder
AMEL ALN	<i>Amelanchier alnifolia</i>	Saskatoon Berry
BERB REP	<i>Berberis repens</i>	Creeping Mahonia
BETU GLA	<i>Betula glandulosa</i>	Dwarf Birch
BETU PUM	<i>Betula pumila</i> v. <i>glandulifera</i>	Swamp Birch
BETU OCC	<i>Betula occidentalis</i>	Water Birch
CHIM UMP	<i>Chimaphila umbellata</i>	Prince's Pine
CLEM OCC	<i>Clematis occidentalis</i>	Purple clematis
CORN STO	<i>Cornus stolonifera</i>	Red Osier Dogwood
CORY COR	<i>Corylus cornuta</i>	Beaked Hazelnut
GAUL HIS	<i>Gaultheria hispidula</i>	Creeping Snowberry
JUNI COM	<i>Juniperus communis</i>	Ground Juniper
JUNI HOR	<i>Juniperus horizontalis</i>	Creeping Juniper
LEDU GRO	<i>Ledum groenlandicum</i>	Labrador Tea
LINN BOR	<i>Linnaea borealis</i>	Twinflower
LONI DIO	<i>Lonicera dioica</i> v. <i>glaucescens</i>	Twining honeysuckle

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
LONI INV	<i>Lonicera involucrata</i>	Bracted Honeysuckle
LONI UTA	<i>Lonicera utahensis</i>	Red Twinberry
MENZ FER	<i>Menziesia ferruginea</i>	MenziesiaPLO
HOR	<i>Oplopanax horridum</i>	Devil's Club
OXYC MIC	<i>Oxycoccus microcarpus</i>	Small Bog Cranberry
POTE FRU	<i>Potentilla fruticosa</i>	Shrubby Cinquefoil
POTE NOR	<i>Potentilla norvegica</i>	Rough Cinquefoil
PRUN PEN	<i>Prunus pensylvanica</i>	Pin Cherry
PRUN VIR	<i>Prunus virginiana</i>	Choke Cherry
PRUN SP	<i>Prunus species</i>	Cherry
RHOD ALB	<i>Rhododendron albiflorum</i>	White-flowered Rhododendron
RIBE GLA	<i>Ribes glandulosum</i>	Skunk Currant
RIBE HIR	<i>Ribes hirtellum</i>	Wild Gooseberry
RIBE HUD	<i>Ribes hudsoniaum</i>	Northern Black Currant
RIBE LAC	<i>Ribes lacustre</i>	Bristly Black Currant
RIBE OXY	<i>Ribes oxycanthoides</i>	Wild Gooseberry
RIBE TRI	<i>Ribes triste</i>	Wild Red Currant
RIBES SP	<i>Ribes species</i>	
ROSA ACI	<i>Rosa acicularis</i>	Prickly Rose
ROSA SP	<i>Rosa species</i>	Rose
ROSA WOO	<i>Rosa woodsii</i>	Common Wild Rose
RUBU IDA	<i>Rubus idaeus</i>	Wild Red Raspberry
RUBU PAR	<i>Rubus parviflorus</i>	Thimble Berry
RUBU SP	<i>Rubus species</i>	Raspberry Species
SALI ATH	<i>Salix athabascensis</i>	Willow
SALI BAR	<i>Salix barklayi</i>	Barclay's Willow
SALI BEB	<i>Salix bebbiana</i>	Beaked Willow
SALI GLA	<i>Salix glauca</i>	Smooth Willow
SALI MYR	<i>Salix myrtillifolia</i>	Myrtle-leaved willow
SALI PED	<i>Salix pedicellaris</i>	Bog Willow
SALI PET	<i>Salix petiolaris</i>	Basket Willow
SALI PYR	<i>Salix pyrifolia</i>	Balsam Willow
SALI SCO	<i>Salix scouleriana</i>	Willow
SALI SP	<i>Salix species</i>	Willow
SAMB RAC	<i>Sambucus Racemosa</i>	Red Elderberry
SHEP CAN	<i>Shepherdia canadensis</i>	Canadian Buffaloberry
SORB SCO	<i>Sorbus scopulina</i>	Mountain Ash

Species Code	Latin Name	Common Name
SPIR BET	<i>Spiraea betulifolia</i>	Birch-leaved Spirea
SPIR DEN	<i>Spiraea densiflora</i>	Pink Meadowsweet
SPIR SP	<i>Spiraea species</i>	Meadowsweet
SYMP ALB	<i>Symphoricarpos albus</i>	Snowberry
SYMP OCC	<i>Symphoricarpos occidentalis</i>	Wolfberry
VACC CAE	<i>Vaccinium caespitosum</i>	Dwarf Bilberry
VACC MEM	<i>Vaccinium membranaceum</i>	Tall Bilberry
VACC MYR	<i>Vaccinium myrtilloides</i>	Blueberry
VACC MYT	<i>Vaccinium myrtillus</i>	Low Bilberry
VACC SCO	<i>Vaccinium scoparium</i>	Grouse-berry
VACC ULI	<i>Vaccinium uliginosum</i>	Bog Bilberry
VACC VIT	<i>Vaccinium vitis-idaea v. minus</i>	Bog Cranberry
VIBU EDU	<i>Viburnum edule</i>	Lowbush Cranberry
VIBU OPU	<i>Viburnum opulus</i>	High-bush Cranberry

GRASS LAYER

Species Code	Latin Name	Common Name
(POA species - Record 1st 3 letters of genus and 1st 4 letters of species)		
AGRO RIP	<i>Agropyron riparium</i>	Steambank Wheatgrass
AGRO SMI	<i>Agropyron smithii</i>	Western Wheatgrass
AGRO SUB	<i>Agropyron subsecundum</i>	Bearded Wheatgrass
AGRO TRA	<i>Agropyron trachycaulum</i>	Slender Wheatgrass
ALOP AEQ	<i>Alopecurus aequalis</i>	Water Foxtail
AGRO SCA	<i>Agrostis Scabra</i>	Hair Bentgrass
AGRO SP	<i>Agropyron species</i>	
AVEN FAT	<i>Avena fatua</i>	Wild Oat
BROM CAR	<i>Bromus carinatus</i>	Brome
BROM INE	<i>Bromus inermis</i>	Awnless Brome
BROM CIL	<i>Bromus ciliatus</i>	Fringed Brome
BROM VUL	<i>Bromus vulgaris</i>	Columbia Brome
CALA CAN	<i>Calamagrostis canadensis</i>	Bluejoint Marsh Reed
CALA INE	<i>Calamagrostis inexpansa</i>	Northern Reed Grass
CALA NEG	<i>Calamagrostis neglecta</i>	Narrow, Plains Reed Grass
CALA PUR	<i>Calamagrostis purpurascens</i>	Purple Reed Grass
CALA RUM	<i>Calamagrostis rubescens</i>	Pinegrass
CARE AQU	<i>Carex aquatilis</i>	Water Sedge
CARE BEB	<i>Carex bebbii</i>	Sedge

Species Code	Latin Name	Common Name
CARE BRU	<i>Carex brunnescens</i>	Brownish Sedge
CARE CON	<i>Carex concinna</i>	Beautiful Sedge
CARE DIS	<i>Carex disperma</i>	Two-seeded Sedge
CARE FOE	<i>Carex foenea</i>	Sedge
CARE GYN	<i>Carex gynocrates</i>	Northern Bog Sedge
CARE HOU	<i>Carex houghtoniana</i>	Carex
CARE LAS	<i>Carex lasiocarpa</i>	Hairy-fruited Sedge
CARE LEP	<i>Carex leptales</i>	Bristle Stalked Sedge
CARE OBT	<i>Carex obtusata</i>	Blunt Sedge
CARE PRA	<i>Carex praticola</i>	Sedge
CARE RIC	<i>Carex richardsonii</i>	Richardson's Sedge
CARE ROS	<i>Carex rostrata</i>	Beaked Sedge
CARE SP	<i>Carex species</i>	Carex Species
CARE UMB	<i>Carex umbellata</i>	Umbellate Sedge
CARE VAG	<i>Carex vaginata</i>	Sheathed Sedge
CINN LAT	<i>Cinna latifolia</i>	<i>Drooping Wood Reed</i>
DANT PAR	<i>Danthonid parryi</i>	Parry's Oatgrass
DESC CAE	<i>Deschampsia caespitosa</i>	Tufted Hair Grass
DIST STR	<i>Distichlis stricta</i>	Salt Grass
ELYM INN	<i>Elymus innovatus</i>	Hairy Wild Rye
ELYM SP	<i>Elymus species</i>	Wild Rye
ERIO POL	<i>Eriophorum polystachion</i>	Cotton Grass
ERIO SP	<i>Eriophorum species</i>	Cotton Grass
FEST OCC	<i>Festuco occidentalis</i>	Western Fescue
FEST PRA	<i>Festuca pratensis</i>	Meadow Fescue
FEST RUB	<i>Festuca rubra</i>	Red Fescue
FEST SAX	<i>Festuca saximontana</i>	Sheet Fescue
FEST SCA	<i>Festuca scabrella</i>	Rough Fescue
FEST SP	<i>Festuca species</i>	Fescue Species
GYLYC STR	<i>Glyceria striata</i>	Fowl Manna Grass
GRASS SP		<i>Grass species</i>
HIER ALP	<i>Hierochloe Alpina</i>	Alpine Sweetgrass
HIER ODO	<i>Hierochloe odorata</i>	Common Sweetgrass
HORD JUB	<i>Hordeum jubatum</i>	Foxtale Barley
JUNC BAL	<i>Juncus balticus</i>	Wire Rush
KOEL CRI	<i>Koeleria cristata</i>	June Grass
LUZU PAR	<i>Luzula parviflora</i>	Small-flowered Woodrush
ORYZ ASP	<i>Oryzopsis asperifolia</i>	Mountain Rice Grass
ORYZ PUN	<i>Oryzopsis pungens</i>	Short-awned Rice Grass
PANI SP	<i>Panicum species</i>	Panic Grass
PHLE PRA	<i>Phleum pratense</i>	Timothy Grass

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
PHRA AUS	<i>Phragmites australis</i>	Reed
POA GLAU	<i>Poa glauca</i>	Bluegrass
POA INTE	<i>Poa interior</i>	Bluegrass
POA PALU	<i>Poa palustris</i>	Fowl Bluegrass
POA PRAT	<i>Poa pratensis</i>	Kentucky Bluegrass
POA SP	<i>Poa species</i>	
SCHI PUR	<i>Schizachne purpurascens</i>	False Melic
TRIS SPI	<i>Trisetum spicatum</i>	Spike Trisetum
XERO TEN	<i>Xerophyllum tenax</i>	Bear Grass

HERB LAYER

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
ANAP MAR	<i>Anaphalis margaritacea</i>	Pearly Everlasting
ACHI MIL	<i>Achillea millefolium</i>	Common Yarrow
ACHI SIB	<i>Achillea sibirica</i>	Yarrow
ACHI SP	<i>Achillea species</i>	Yarrow Species
ACTA RUB	<i>Actaea rubra</i>	Red Baneberry
AGOS GLA	<i>Agoseris glauca</i>	Pale False Dandelion
AGOS SP	<i>Agoseris species</i>	False Dandelion
ALLI CER	<i>Allium cernuum</i>	Nodding Onion
ANEM MUL	<i>Anemone multifida</i>	Windflower, Cutleaf Anemone
ANTE MIC	<i>Antennaria microphylla</i>	Rosy Pussytoes
ANTE SP	<i>Antennaria species</i>	
ANTE NEG	<i>Antennaria neglecta</i>	
ANTE PUL	<i>Antennaria pulcherrima</i>	Showy Everlasting
ANTE RAC	<i>Antennaria racemosa</i>	Racemose Everlasting
ANTE ROS	<i>Antennaria rosea</i>	Rosy Everlasting
APOC AND	<i>Apocynum androsaemifolium</i>	Spreading Dogbane
AQUI BRE	<i>Aquilegia brevistyla</i>	Blue Columbine
AQUI FLA	<i>Aquilegia flavescens</i>	Yellow Columbine
ARAL NUD	<i>Aralia nudicaulis</i>	Wild Sarasparilla
ARCE AME	<i>Arceuthobium americanum</i>	Dwarf Mistletoe
ARNI CHA	<i>Arnica chamissonis</i>	Leafy Arnica
ARNI COR	<i>Arnica cordifolia</i>	Heart-leaved Arnica
ARNI LAT	<i>Arnica latifolia</i>	Mountain Arnica
ARNI LON	<i>Arnica longifolia</i>	Long-leaved Arnica
ARNI SP	<i>Arnica species</i>	Arnica
ASTR SP	<i>Astragalus species</i>	Milk Vetch
ARTE CAM	<i>Artemisia campestris</i>	Plains Wormwood
ARTE FRI	<i>Artemisia frigida</i> v. <i>americanus</i>	Pasture Sage

Species Code	Latin Name	Common Name
ASTE CIL	<i>Aster ciliolatus</i>	Lindley's Aster
ASTE CON	<i>Aster conspicuous</i>	Showy Aster
ASTE FOL	<i>Aster folicaeus</i>	Aster
ASTE JUN	<i>Aster junciformis</i>	Rush Aster
ASTE LAE	<i>Aster laevis</i>	Smooth Aster
ASTE SP	<i>Aster species</i>	Aster species
ASTR ALP	<i>Astragalus alpinus</i>	Alpine Milk Vetch
ASTR EUC	<i>Astragalus eucosmus</i>	Milk Vetch
ASTR FRI	<i>Astragalus frigidus</i>	American Milk Vetch
ASTR STR	<i>Astragalus striatus</i>	Ascending Purple Milk Vetch
BOTR VIR	<i>Botrychium virginianum</i>	Grape Fern
CALY BUL	<i>Calypso bulbosa</i>	Venus' Slipper
CAMP ROT	<i>Campanula rotundifolia</i>	Bluebell, Harebell
CAST MIN	<i>Castilleja miniata</i>	Red Indian Paintbrush
CAST PAR	<i>Castilleja parviflora</i>	Small Flowered Paintbrush
CAST SP	<i>Castilleja species</i>	Paintbrush
CERA ARV	<i>Cerastium arvense</i>	Field Chickweed
CERA SP	<i>Cerastium species</i>	Chickweed species
CHRY IOW	<i>Chrysplenium iowense</i>	Golden Saxifrage
CHRY LEU	<i>Chrysanthemum leucanthemum</i>	Ox-eye Daisy
CICU MAC	<i>Cicuta maculata</i>	Water Hemlock
CIRS ARV	<i>Cirsium arvense</i>	Canada Thistle
CIRS HOO	<i>Cirsium hookerianum</i>	White Thistle
CIRS SP	<i>Cirsium species</i>	Thistle Species
CIRS VUL	<i>Cirsium vulgare</i>	Bull Thistle
CORA MAC	<i>Corallorhiza maculata</i>	Spotted Coralroot
CORA TRI	<i>Corallorhiza trifida</i>	Pale Coralroot
CORN CAN	<i>Corn canadensis</i>	Bunch Berry
CORY AUR	<i>Corydalis aurea</i>	Yellow Corydalis
CORY SEM	<i>Corydalis sempervirens</i>	Pink/Pale Corydalis
CORY SP	<i>Corydalis species</i>	Corydalis
CREP SP	<i>Crepis species</i>	Hawksbeard species
CREP TEC	<i>Crepis tectorum</i>	Annual Hawksbeard
CYST FRA	<i>Cystopteris fragilis</i>	Bladder Fern
DELP GLA	<i>Delphinium glaucum</i>	Tall Larkspur
DISP TRA	<i>Disporum trachyarpum</i>	Fairy-bells
DODE RAD	<i>Dodecatheon radiculatum</i>	Shooting Star
DRAB AUR	<i>Draba aurea</i>	Golden Whitlow-Grass
DRYO CRI	<i>Dryopteris cristata</i>	Fern
DRYO EXP	<i>Dryopteris expansa</i>	Spiny Woodfern
DRYO SP	<i>Dryopteris species</i>	Fern
DRYO SPI	<i>Dryopteris spinulosa</i>	Narrow Spinulose Shield Fern
ERIG GLA	<i>Erigeron glabellus</i>	Wild Daisy

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
EPIL ANG	<i>Epilobium angustifolium</i>	Fireweed
EQUI ARV	<i>Equisetum arvense</i>	Field Horsetail
EQUI PRA	<i>Equisetum pratense</i>	Meadow Horsetail
EQUI SCI	<i>Equisetum scirpoides</i>	Dwarf Scouring Rush
EQUI SP	<i>Equisetum species</i>	Horsetail Species
EQUI SYL	<i>Equisetum sylvaticum</i>	Woodland Horsetail
EQUI VAR	<i>Equisetum variegatum</i>	Variegated Horsetail
ERIG PER	<i>Erigeron peregrinus v. callianthemus</i>	Wondering Daisy
ERIG SP	<i>Erigeron species</i>	Fleabane
FRAG VIR	<i>Fragaria virginiana</i>	Wild Strawberry
GAIL ARI	<i>Gaillardia aristata</i>	Gaillardia
GALE TET	<i>Galeopsis tetrahit</i>	Hemp Nettle
GALI BOR	<i>Galium boreale</i>	Northern Bedstraw
GALI TRI	<i>Galium triflorum</i>	Sweet-Scented Bedstraw
GENT AM'A	<i>Gentianella amarella</i>	Felwort, Northern Gentian
GENT SP	<i>Gentianella species</i>	Gentian species
GEOC LIV	<i>Geocaulon lividum</i>	Bastard Toadflax
GERA RIC	<i>Geranium richardsonii</i>	Richardson Geranium
GERA SP	<i>Geranium species</i>	Geranium
GERA VIS	<i>Geranium viscosissimum</i>	Sticky purple geranium
GEUM ALE	<i>Geum aleppicum</i>	Yellow Avens
GEUM MAC	<i>Geum macrophyllum</i>	Yellow Avens
GEUM RIV	<i>Geum rivale</i>	Purple or Water Avens
GEUM TRI	<i>Geum triflorum</i>	Old Man's Whiskers
GOOD REP	<i>Goodyera repens</i>	Rattlesnake Plantain
GYMN DRY	<i>Gymnocarpium dryopteris</i>	Oak Fern
HABE HYP	<i>Habenaria hyperborea</i>	Northern Green Orchid
HABE OBT	<i>Habenaria obtusata</i>	Blunt-leaved Orchid
HABE ORB	<i>Habenaria orbiculata</i>	Round-leaved Orchid
HABE VIR	<i>Habenaria viridis v. bracteata</i>	Bracted Orchid
HALE DEF	<i>Halenia deflexa</i>	Spurred Gentian
HEDY ALP	<i>Hedysarum alpinum v. americanum</i>	American Hedysarum
HEDY SP	<i>Hedysarum species</i>	Hedysarum
HERA LAN	<i>Heracleum lanatum</i>	Cow Parsnip
HIER ALB	<i>Hieracium albiflorum</i>	White hawkweed
HIER CAN	<i>Hieracium canadense</i>	Canada Hawkweed

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
HIER SP	<i>Hieracium species</i>	Hawkweed species
HIER UMB	<i>Hieracium umbellatum</i>	Narrow-leaved Hawkweed
LATH OCH	<i>Lathyrus ochroleucus</i>	Cream-coloured Vetchlin
LATH VEN	<i>Lathyrus venosus</i>	Peavine
LILI PHI	<i>Lilium philadelphicum</i>	Western Wood Lily
LINU LEW	<i>Linum lewisii</i>	Wild Blue Flax
LIST BOR	<i>Listera borealis</i>	Western Twayblade
LIST COR	<i>Listera cordata</i>	Heart-leaved Twayblade
LYCH DRU	<i>Lychnis drummondii</i>	Drummond's Cogle
LYCO ANN	<i>Lycopodium annotinum</i>	Stiff Club-Moss
LYCO COM	<i>Lycopodium complanatum</i>	Ground Cedar
LYCO SP	<i>Lycopodium species</i>	Club-moss
LUPI ARG	<i>Lupinus argenteus</i>	Perennia lupine
MAIA CAN	<i>Maianthemum canadense</i>	Wild Lily-of-the-Valley
MELI ALB	<i>Melilotus alba</i>	White sweet clover
MELI OFF	<i>Melilotus officinalis</i>	Yellow sweet clover
MERT PAN	<i>Mertensia paniculata</i>	Tall Mertensia, Lungwort
MITE NUD	<i>Mitella nuda</i>	Bishop's Cap, Mitrewort
ORCH ROT	<i>Orchis rotundifolia</i>	Round-leaved Orchid
ORTH SEC	<i>Orthilia secunda</i>	One-Sided Wintergreen
OSMO CHI	<i>Osmorhiza chilensis</i>	Blunt-fruited Sweet Cicely
OSMO DEP	<i>Osmorhiza depauperata</i>	Sweet Cicely
OXYT CAM	<i>Oxytropis campestris</i>	Late Yellow Locoweed
OXYT SER	<i>Oxytropis sericeus</i> v. <i>spicata</i>	Early Yellow Locoweed
OXYT SPL	<i>Oxytropis splendens</i>	showy loco-weed
PARN FIM	<i>Parnassia fimbriata</i>	Fringed Grass-of Parnas
PARN PAL	<i>Parnassia palustris</i>	Northern grass-of parnassus
PEDI BRA	<i>Pedicularis bracteosa</i>	Bracted lousewort
PEDI CAP	<i>Pedicularis capitata</i>	Few-flowered lousewort
PEDI GRO	<i>Pedicularis groenlandicam</i>	Elephant Head
PEDI LAB	<i>Pedicularis labradoricam</i>	Labrador Lousewort
PENS PRO	<i>Penstemon procerus</i>	Slender Blue Beardtongue
PETA PAL	<i>Petasites palmatus</i>	Palmate-leaved Coltsfoot
PETA SAG	<i>Petasites sagittatus</i>	Arrow-leaved Coltsfoot

Species Code	Latin Name	Common Name
PICR ECH	<i>Picris echioides</i>	Bristly ox-tongue
PLAN MAJ	<i>Plantago major</i>	common plantain
POLY BIS	<i>Polygonum bistortoides</i>	Western Bistort
POLY CAE	<i>Polygonum caeruleum</i> spp. <i>occidentale</i>	Jacob's Ladder
POLY VIV	<i>Polygonum viviparum</i>	Alpine Bistort
POTE ARG	<i>Potentilla arguta</i>	White Cinquefoil
POTE GLA	<i>Potentilla glandulosa</i> spp. <i>pseudorupestris</i>	Cinquefoil
POTE GRA	<i>Potentilla gracilis</i>	Graceful Cinquefoil
PYRO ASA	<i>Pyrola asarifolia</i>	Common Pink Wintergreen
PYRO BRA	<i>Pyrola bracteata</i>	Large wintergreen
PYRO CHL	<i>Pyrola chlorantha</i>	Green Wintergreen
PYRO ORB		
PYRO SP	<i>Pyrola species</i>	Wintergreen
RANU ACR	<i>Ranunculus acris</i>	Tall buttercup
RHIN CRI	<i>Rhinanthus cristagalli</i>	Yellow Rattle
RHIN MIN	<i>Rhinanthus minor</i>	Yellow rattle
RUBU ARC	<i>Rubus arcticus</i>	Dwarf Raspberry
RUBU PED	<i>Rubus pedatus</i>	Five-leaved bramble
RUBU PUB	<i>Rubus pubescens</i>	Dewberry
RUBU SP	<i>Rubus species</i>	
RUME ACE	<i>Rumex acetosa</i>	Green sorrel
RUME OCC	<i>Rumex occidentalis</i> v. <i>fenestratus</i>	Western Dock
SAXI BRO	<i>Saxifraga bronchialis</i>	Prickly Saxifrage
SCIR VAL	<i>Scirpus validus</i>	Common great bulrush
SCUT GAL	<i>Scutellaria galericulata</i>	Skullcap
SEDU STE	<i>Sedum stenopetalum</i>	Common Stonecap
SELA DEN	<i>Selaginella densa</i>	Prairie Selaginella
SENE CAN	<i>Senecio canus</i>	Prairie Groundsel
SENE PAU	<i>Senecio pauperculus</i>	Balsam Groundsel
SENE PSE	<i>Senecio pseud aureus</i>	Stream bank butterweed
SENE SP	<i>Senecio species</i>	Groundsel
SENE TRI	<i>Senecio triangularis</i>	Arrow-leaved groundsel
SMIL RAC	<i>Smilacina racemosa</i>	False Solomon's Seal
SMIL STE	<i>Smilacina stellata</i>	Star-flowered Solomon Seal
SMIL TRI	<i>Smilacina trifolia</i>	Three-leaved Solomon Seal
SOLI CAN	<i>Solidago canadensis</i>	Canada goldenrod
SOLI GIG	<i>Solidago gigantea</i>	Goldenrod
SOLI MUL	<i>Solidago multiradiata</i>	Alpine Goldenrod

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
SOLI SP	<i>Solidago species</i>	Goldenrod species
SOLI SPA	<i>Solidago spathulata</i>	Spike Like Golden Rod
SONC ARV	<i>Sonchus arvensis</i>	Perennial sow thistle
SONC SP	<i>Sonchus species</i>	Sow thistle
SPIR ROM	<i>Spiranthes romanzoffiana</i>	Ladies' Tresses
STEL CAL	<i>Stellaria calycantha</i>	Northern starwort
STEL LON	<i>Stellaria longifolia</i>	Long-leaved Chickweed
STEL SP	<i>Stellaria species</i>	Starwort species
STEN OCC	<i>Stenanthium occidentale</i>	Bronze Bells
STRE AMP	<i>Streptopus amplexifolius</i>	Twisted Stalk
TANA VUL	<i>Tanacetum vulgare</i>	Common tansy
TARA CER	<i>Taraxacum ceratophrum</i>	Horned Dandelion
TARA OFF	<i>Taraxacum officinale</i>	Common Dandelion
TARA SP	<i>Taraxacum species</i>	
THAL VEN	<i>Thalictrum venulosum</i>	Veiny Meadow Rue
THER RHO	<i>Thermopsis rhombifolia</i>	Golden bean
TRIF AUR	<i>Trifolium aureum</i>	Hop clover
TRIF PRA	<i>Trifolium pratense</i>	Red Clover
TRIF REP	<i>Trifolium repens</i>	White Clover
TRIF SP	<i>Trifolium species</i>	Clover Species
THPH LAT	<i>Typha latifolia</i>	Cattail
URTI DIO	<i>Urtica dioica</i>	Common nettler
VERA ESC	<i>Veratrum eschscholtzii</i>	False hellebore
VERO SP	<i>Veronica species</i>	
VERO WOR	<i>Veronica wormskioldii</i>	Alpine Speedwell
VICI AME	<i>Vicia americana</i>	Wild Vetch
VIOL ADU	<i>Viola adunca</i>	Early Blue Violet
VIOL CAN	<i>Viola canadensis</i>	Canada Violet
VIOL REN	<i>Viola renifolia</i>	Kidney-leaved Violet
VIOL RUG	<i>Viola rugulosa</i>	Western Canada Violet
VIOL SP	<i>Viola species</i>	Violet
ZIZI APT	<i>Zizia aptera</i>	Meadow Parsnip
ZYGA ELE	<i>Zygadenus elegans</i>	Death Camas

MOSESSES AND LIVERWORTS

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
AMBL SER	<i>Amblystegium serpens</i>	
ANAS HEL	<i>Anastrophyllum helleranum</i>	
AULA PAL	<i>Aulacomnium palustre</i>	Glow Moss
AULA SP	<i>Aulacomnium species</i>	
BARB HAT	<i>Barbilophozia hatcheri</i>	Liverwort
BRAC OED	<i>Brachythecium Oedipum</i>	Short-leaved Ragged Moss
BRAC SAL	<i>Brachythecium salebrosum</i>	
BRAC STA	<i>Brachythecium starkei</i>	
BRAC TUR	<i>Brachythecium turgidum</i>	
BRYU CAE	<i>Bryum caespitium</i>	
BRYU PSE	<i>Bryum pseudotriquetrum</i>	
CALL GIG	<i>Calliergon giganteum</i>	
CALY SPH	<i>Calypogeja sphagnicola</i>	
CAMP CHR	<i>Campylium chrysophyllum</i>	
CAMP HIS	<i>Campylium hispidulum</i>	
CAMP STE	<i>Campylium stellatum</i>	
CEPH LUN	<i>Cephalozia lunifolia</i>	
CERA PUR	<i>Ceratodon purpureus</i>	Fire Moss
CLAD BAC	<i>Cladonia bacillaris</i>	
CLIM DEN	<i>Climacium dendroides</i>	
DICR ACU	<i>Dicranum acutifolium</i>	
DICR CON	<i>Dicranum condensatum</i>	
DICR ELO	<i>Dicranum elongatum</i>	
DICR FLA	<i>Dicranum flagellare</i>	
DICR FRA	<i>Dicranum fragilifolium</i>	
DICR FUS	<i>Dicranum fuscenscens</i>	
DICR GRO	<i>Dicranum groenlandicum</i>	
DICR MUE	<i>Dicranum muehlenbeckii v. cirratum</i>	
DICR POL	<i>Dicranum polysetum</i>	
DICR SCO	<i>Dicranum scoparium</i>	
DICR SP	<i>Dicranum species</i>	
DICR UND	<i>Dicranum undulatum</i>	
DIST CAP	<i>Distichium capillaceum</i>	
DREP ADU	<i>Drepanocladus aduncus</i>	
DREP REV	<i>Drepanocladus revolvens</i>	
DREP UNC	<i>Drepanocladus uncinatus</i>	
DREP VER	<i>Drepanocladus vernicosus</i>	
EURH PUL	<i>Eurhynchium pulchellum</i>	
FUNA HYG	<i>Funaria hygrometrica</i>	
GEOC GRA	<i>Geocalyz graveolans</i>	
HEDW CIL	<i>Hedwigia ciliate</i>	

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
HYLO SP	<i>Hylocomium species</i>	
HYLO SPL	<i>Hylocomium splendens</i>	Stair Step Moss
HYPN PRA	<i>Hypnum pratense</i>	
HYPN REV	<i>Hypnum revolutum</i>	
JAME AUT	<i>Jamesoniella autumnalis</i>	
LEPI REP	<i>Lepidozia reptans</i>	
LEPT PYR	<i>Leptobryum pyriforme</i>	
LOPH GUT	<i>Lophozia guttulata</i>	
LOPH INC	<i>Lophozia incisa</i>	
LOPH LON	<i>Lophozia longidens</i>	
LOPH VEN	<i>Lophozia ventricosa</i>	
MNIU SP	<i>Mnium species</i>	
MNIU SPI	<i>Mnium spinulosum</i>	
MYLI ANO	<i>Mylia anomala</i>	
MYUR JUL	<i>Myurella julacea</i>	
ONCO WAH	<i>Oncophorus wahlenbergii</i>	
ORTH RUP	<i>Orthotrichum rupestre</i>	
ORTH SPE	<i>Orthotrichum speciosum</i>	
PLAG ASP	<i>Plagiochila asplenoides</i>	
PLAG CUS	<i>Plagiomnium cuspidatum</i>	
PLAG DRU	<i>Plagiomnium drummondii</i>	
PLAG MED	<i>Plagiomnium medium</i>	
PLEU SCH	<i>Pleurozium schreberi</i>	
POHL NUT	<i>Pohlia nutans</i>	
POLY	<i>ALPPolytrichum Alpinum</i>	Stiff-leaved Polytrichum
POLY COM	<i>Polytrichum commune</i>	
POLY JUN	<i>Polytrichum juniperinum</i>	
POLY SP	<i>Polytrichum Species</i>	
POLY STR	<i>Polytrichum strictum</i>	
PTIL CIL	<i>Ptilium ciliare</i>	
PTIL CRI	<i>Ptilium crista-castrensis</i>	Knight's Plume
PTIL PUL	<i>Ptilium pulcherrimum</i>	
PYLA POL	<i>Pylaisiella polyantha</i>	
RHIZ PSE	<i>Rhizomnium pseudopunctatum</i>	
RHYT RUG	<i>Rhytidium rugosum</i>	
SPHA ANG	<i>Sphagnum angustifolium</i>	
SPHA CAP	<i>Sphagnum capillaceum</i>	Common red sphagnum
SPHA FUS	<i>Sphagnum fuscum</i>	
SPHA SP	<i>Sphagnum Species</i>	
SPHA WAR	<i>Sphagnum warnstorffii</i>	
SPLA AMP	<i>Splachnum ampullaceum</i>	
APLA SPH	<i>Splachnum sphaericum</i>	
TAYL SER	<i>Tayloria serrata</i>	

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
TETR ANG	<i>Tetraplodon angustatus</i>	
TETR MIN	<i>Tetraplodon minoides</i>	
TETR PEL	<i>Tetraphis pellucida</i>	
THUI ABI	<i>Thuidium abietinum</i>	
THUI REC	<i>Thuidium recognitum</i>	
TIMM AUS	<i>Timmia austriaca</i>	
TOME NIT	<i>Tomenthypnum nitens</i>	
TORT MUC	<i>Tortula mucronifolia</i>	
TORT RUR	<i>Tortula ruralis</i>	
TRIT EXS	<i>Tritomaria exsecta</i>	

LICHENS

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
BACI SPH	<i>Bacidian sphaeroides</i>	
BRYO FRE	<i>Bryoria fremontii</i>	
BRYO FUS	<i>Bryoria fuscescens</i>	
CETR CUC	<i>Cetraria cucullata</i>	
CETR ERI	<i>Cetraria ericetorum</i>	
CETR HAL	<i>Cetraria halei</i>	
CETR ISL	<i>Cetraria islandica</i>	
CETR NIV	<i>Cetraria nivalis</i>	
CETR PIN	<i>Cetraria pinastri</i>	
CLADI SP	<i>Cladina Species</i>	
CLAD BOT	<i>Cladonia botrytes</i>	
CLAD CAR	<i>Cladonia carneola</i>	
CLAD CEN	<i>Cladonia cenotea</i>	
CLAD CHL	<i>Cladonia chlorophaea</i>	
CLAD COC	<i>Cladonia coccifera</i>	
CLAD CON	<i>Cladonia coniocraea</i>	
CLAD COR	<i>Cladonia cornuta</i>	
CLAD DEF	<i>Cladonia deformis</i>	
CLAD ECM	<i>Cladonia ecmocyna</i>	
CLAD FIM	<i>Cladonia fimbriata</i>	
CLAD GRA	<i>Cladonia gracilis</i>	
CLAD MIT	<i>Cladonia multiformis</i>	
CLAD PLE	<i>Cladonia pleurota</i>	
CLAD PYX	<i>Cladonia pyxidata</i>	
CLAD RAN	<i>Cladonia rangiferina</i>	
CLAD SP	<i>Cladonia Species</i>	
EVER MES	<i>Evernia mesomorpha</i>	
HYPO BIT	<i>Hypogymnia bitteri</i>	

<u>Species Code</u>	<u>Latin Name</u>	<u>Common Name</u>
HYPO PHY	Hypogymnia physodes	
ICMA ERI	Imadophila ericetorum	Spraypaint Lichen
LETH VUL	Letharia vulpina	
LOBA PUL	<i>Lobaria pulmonaria</i>	Lungwort
PARM ALE	Parmeliopsis aleurites	
PARM AMB	Parmeliopsis ambigua	
PARM HYP	Parmeliopsis hyperopta	
PARM SUL	Parmelia sulleata	
PELT APH	Peltigera aphthosa	Green Dog Lichen
PELT CAN	Peltigera canina	
PELT MAL	Peltigera malacea	
PELT POL	Peltigera polydactyla	
PELT SP	<i>Peltigera species</i>	
PHYS ADS	Physcia adscendens	
PLAG SP	<i>Plagiomnium species</i>	Leafy moss
PLAT GLA	Platismatia glauca	
RAMA FAS	Ramalina fastigiata	
RAMA POL	Ramalina pollinaria	
STER TOM	Stereocaulon tomentosum	
USNE GLA	Usnea glabrescens	
USNE HIR	Usnea hirta	
USNE SOR	Usnea sorediifera	
USNE SUB	Usnea subfloridana	
USNE SP	Usnea species	
XANT POL	Xanthoria polycarpa	
XANT STR	Xanthoria	

14. COMMON PROBLEMS

Duplicate Tree Numbers

Try to determine the "oldest" tree and assume that is the initial tagged tree. If this can't be done, assume the tree with the largest DBH and/or height is the original tree. The condition and age of the tree tag may give a clue as to what tree was originally tagged as well. Number the other tree with the next available sequential number.

Past crews would also assign a sapling number to a seedling that has grown to 1.1cm DBH. If it is certain that 2 different numbers were used on the same stem, assume the tree is the one with the lower number (regen number).

In all cases fully document this in the comments column beside the appropriate tree(s).

Establishment crews would also rarely follow the proper initial numbering sequence in the regen plots. Use extreme care when measuring these plots.

15. RECORDING DATA

All measurements taken on these plots must be recorded on the appropriate tally sheets. Data entered on the sheets is later entered into a computer so legibility is very important. All header lines must be completed.

- 1) All letters must be capitalized.
- 2) Use only those species codes listed in Appendix 6.1 and 6.2.
- 3) Asterisks, numeric characters in alphabetic fields (e.g. B3 in the species columns) and alphabetic characters in numeric fields (e.g. H in DBH columns) are not acceptable.
- 4) Comments are written in the shaded areas only. Comments written elsewhere on the tally sheets are not acceptable.
- 5) Alphabetic characters that are commonly illegible are:

N that looks like W
C that looks like L or O
D that looks like P or O
I that looks like T or L

- 6) Numeric characters that are commonly illegible are:

2 that is 'looped' and looks like 0
6 and 9 that looks like 0 or 4
0 incompletely closed and looks like 6
5 that looks like S
7 and 1 mistaken for each other
Scientific (European) 7 is not acceptable

The number is written as open (i.e. four not 4).

16. GENERAL INFORMATION (CSTM 101)

<u>Column</u>	<u>Name</u>	<u>Data Entry</u>
1	Record Type	03 - cruise tally 04 - cruise vegetation 05 - plot vegetation 06 - regeneration (planted & natural) 61 - regen height class tallies 07 - sapling 08 - tree plot 09 - plot retreatment 1 record type per page.
3	Group Number	2 digits, right justified.
5	Plot Number	5 digits - zero fill, right justified.
8	Plot Type	Leave blank.
10	Year	Last 2 digits of year are recorded e.g. 1996 recorded as 96.
12	Month	Right justified, zero filled e.g. June is recorded as 06.
14	Day	Right justified zero filled. Note the date is the same on all tally sheets even if the plot took several days to remeasure with the exception of the vegetation tally sheet CSTM 102, which the actual date is recorded.
15	Crew Identification	Full crew names, initials are not acceptable e.g. John Smith, Mary Jones.
16	Regen Plot	Fill in the number of regeneration plot in which pinned seedlings are found. For planted seedlings outside the regen plots but within the pre 1993 sapling plot, code regen # as 5.

<u>Column</u>	<u>Name</u>	<u>Data Entry</u>
17	Tree Number	Fill in as required.
21	Species	Recorded as 2 capitalized letters as given in Appendix 6.1.
23	Diameter	Numeric, right justified, tree must be > 1.3 m tall to have diameter recorded. Measurements to nearest millimeter.
27	Height	Numeric right justified. Regen & sapling exclude current years growth and measure to nearest centimeter. Trees (>9.1cm DBH) - measure and record total height for all trees. Refer to Appendix 6.3
35	Crown Class	Crown class is recorded on trees and saplings as one letter (D, C, I, S or O). Refer to Appendix 6.9
36	Condition Codes	Left justified, recorded as a 2 digit number. Do not zero fill e.g. a leaning tree with conks is recorded as 51,58, blank. If a tree has no defect it is recorded as '00'. Refer to Appendix 6.5
42	Azimuth	Right justified, recorded as 1°-360°. Only 360° can be used, not 0°.
45	Distance	Right justified, measured to nearest decimeter on all saplings and trees (standing dead as well). Record declination in comments. Note that 2 stems cannot have identical azimuths and distances.

17. GENERAL INFORMATION (CSTM 100)

Columns	Refer to section 4.1
1-16	Refer to section 4.1
17	Species - recorded as 2 capitalized letters as given in Appendix 6.1
19-54	Height classes - recorded to nearest centimeter, excluding current years growth. Right justified, zero filled.

18. EXAMPLE OF A TIE PLATE

1. FOREST MANAGEMENT PROJECT

PERMANENT SAMPLE PLOT NO 456

TIE POINT 'A' 75 M C 180° TO P.C.

LS. SEC. TWP. RGE. W. MER.

1 36 101 19 4

FOR MORE INFORMATION CONTACT ALBERTA FOREST SERVICE

RANGER STATION AT *Fort McMurray*

OR FOR DISTANCE MEASUREMENT AT 427-8474

PSP MANUALS MASTER TABLE OF FIGURES

March 2005

**Public Lands and Forests Division
Forest Management Branch
8th Fl. 9920-108 Street
Edmonton, AB
T5K 2M4**

Phone: (780) 427 – 8474

Or visit the website: <http://www3.gov.ab.ca/srd/forests/psp>

12/1/18

12/1/18

12/1/18

12/1/18

12/1/18

TABLE OF FIGURES

FIGURE 2.1	PLOT LAYOUT	1
FIGURE 2.2	PLOT BOUNDARIES.....	1
FIGURE 2.3	A SWATHS AND TAGGING ORDER	2
FIGURE 2.3B	AN EXAMPLE OF NUMBERING ON TREE TAG	2
FIGURE 2.4	DETERMINING POINT OF GERMINATION AND BREAST HEIGHT	3
FIGURE 2.5	BASE OF LIVE CROWN.....	4
FIGURE 2.6	CROWN CLASS	4
FIGURE 2.7	INCREMENT WIDTHS	5
FIGURE 2.8	CROWN WIDTH MEASUREMENT	6
FIGURE 2.9	PRE-APRIL 1981 PLOT LAYOUT	7
FIGURE 2.10	GROUP PLOT MAINTENANCE FORM.....	8
FIGURE 2.11	BUFFER DAMAGES AREA	9
FIGURE 2.12	EXAMPLES OF BUFFER DAMAGE	10
FIGURE 2.13	DATA ENTRY ON PSP HEADER SHEET	11
FIGURE 2.14	DATA ENTRY ON PSP TALLY SHEET	12
FIGURE 2.15	DATA ENTRY ON REGENERATION TALLY SHEET.....	13
FIGURE 3.1	PSP TREE TALLY CHECK SHEET.....	14
FIGURE 4.1	TREE HEIGHT MEASUREMENT	15
FIGURE 4.2	HEIGHT MEASUREMENT OF LEANING TREES.....	16
FIGURE 4.3	CONK AND BLIND CONKS	17
FIGURE 4.4	OPEN SCARS.....	17
FIGURE 4.5	LARGE BURL ON MAIN STEM	18
FIGURE 4.6	FORKS.....	18
FIGURE 4.7	PRONOUNCED CROOK	19
FIGURE 4.8	LEANING TREE.....	19

FIGURE 4.9	ATROPELLIS CANKER	20
FIGURE 4.10	WITCHES ON SPRUCE.....	20
FIGURE 4.11	MISTLETOE.....	21
FIGURE 4.12	INSTRUCTIONS AND EXAMPLE OF THE USE OF THE 6.....	22
	CLASS MISTLETOE RATING SYSTEM (HAWKSWORTH 1961, 1977).....	22
FIGURE 4.13	GENERIC WOODPECKER.....	23
FIGURE 4.14	YELLOWBELLIED SAPSUCKER FEEDING	23
FIGURE 4.15	SMALL MAMMA L FEEDING ON TREE BOLE	24
FIGURE 4.16	EXCAVATIONS BY WOODPECKERS.....	24
FIGURE 4.17	SMALL CAVITY	25
FIGURE 4.18	LARGE CAVITY	25
FIGURE 4.19	ALBERTA SURVEY SYSTEM	26
FIGURE 4.20	PLOT BOUNDARIES.....	27
FIGURE 4.21	STAND DYNAMICS PLOT LAYOUT	28
FIGURE 4.22	PLOT MAINTENANCE REPORT	29
FIGURE 4.23	SEEDLING NUMBERING	30
FIGURE 4.24	STAND DYNAMICS TALLY SHEET (CSTM 101) RECORD TYPE 6	31
FIGURE 4.25	STAND DYNAMICS TALLY SHEET (CSTM 101) RECORD TYPE 7 (SAPLINGS).....	32
FIGURE 4.26	STAND DYNAMICS TALLY SHEET (CSTM 101) RECORD TYPE 8 (TREE PLOT).....	33
FIGURE 4.27	STAND DYNAMICS REGENERATION HEIGHT CLASS RECORD (CSTM 100) RECORD TYPE	
61		34
FIGURE 4.28	VEGETATION DESCRIPTION FORM	35
FIGURE 4.29	PLOT RETREATMENT REPORT (RECORD TYPE 9).....	36
FIGURE 4.30	STAND DYNAMICS TREE AGE TALLY SHEET	37
FIGURE 4.31	DIEB ACK	38

FIGURE 2.1 PLOT LAYOUT

FIGURE 2.2 PLOT BOUNDARIES

FIGURE 2.3 A SWATHS AND TAGGING ORDER

FIGURE 2.3B AN EXAMPLE OF NUMBERING ON TREE TAG

Trees forked below 1.3m are treated as two separate stems and are tagged and tallied as such.

FIGURE 2.4 DETERMINING POINT OF GERMINATION AND BREAST HEIGHT

FIGURE 2.5 BASE OF LIVE CROWN

FIGURE 2.6 CROWN CLASS

Note: a crown class is recorded for dieback/dead top (code 16). The top of the live foliage is used to determine crown class in this case.

D – Dominant	-crowns extend above the general level of the canopy.
C – Codominant	-crowns form the general level of canopy.
I – Intermediate	- crowns below but extending into the bottom of the general level of the canopy.
S – Suppressed	-crowns entirely below the general level of the canopy.
O – Open grown	-if the trees branches does not interact with another trees branches.

FIGURE 2.7 INCREMENT WIDTHS

Picture A no longer apply

FIGURE 2.8 CROWN WIDTH MEASUREMENT

PERMANENT SAMPLE PLOT MAINTENANCE SHEET

Agency	Ground Number	Year	Month	Day	V	Twp	Rge	Section	Section	Section	LS	LS	LS	LS
01		679	90	02	12	5	08	31	16	09		01	08	

Field Overlay	Plot 1	Field Overlay	Plot 2	Field Overlay	Plot 3	Field Overlay	Plot 4
	C,2,P,L,S,W		C,2,P,L,S,W		C,2,P,L,S,W		C,2,P,L,S,W

Access A.W.R.

Plot Damage	Insect Damage
1	3
2	6
3	6
4	6

Comments From seismic line (310') (TPA) go 250m @ 90° to middle of road (TPB). At TPB go 12m @ 0° to tieplate. From tieplate go 100m @ 0° to group plot centre.

NOTE: T.P.A. refers to tie-point A
"X" marks the location of the sectioned trees

FIGURE 2.10

GROUP PLOT MAINTENANCE FORM

FIGURE 2.11 BUFFER DAMAGES AREA

The seismic line, in this example does not come within 20m of any of the subplots. Therefore, all subplots would have a code 6 recorded for buffer damages.

The seismic line, in this example is within 20 m of subplots 1 and 4. For these two subplots, a code 3 (manmade damage) is recorded in the buffer damage space. Code 6 is recorded for subplots 2 and 3 as damage is not within those buffer damage zones.

318 m

FIGURE 2.12 EXAMPLES OF BUFFER DAMAGE

Agency (1)	Group Number (2)	Plot No. (3)	Measure (m ²) (4)	Year (5)	Month (6)	Day (7)	Season (8)	Temp (9)	Age (10)	Alt (11)	Plot Type (12)	Imp (13)
		1	0390	06	03	04	02	03.2	03	5	000	

Standard Tree Type (14)	Tree Hgt. (m) (15)	Shading Plot Size (16)	Region Plot Size (17)	Field Intersected Quarter (18)	Field Intersected Understory (19)
0	10.00	63	63		

Plot Type (20)	Age Class (21)	Soils (22)	Depth (m) (23)	Surface Vegetation (24)	Management Agent (25)	AVI (26)	Field Overlay Type (27)	Field Understory Type (28)
1	75		3	095	122	4126	SN10	

AGE TREES ONLY

Record Type (29)	Tree Diameter (30)	Species (31)	DBH (32)	Height (33)	Height to Live Crown (34)	Co. (35)	Circulation Codes (36)	DBH Age (37)	Stem Age (38)	Group Intersected Width (39)
0	2	0	0	0	0	0	0	09.1	09.5	01.1013
		0	0	0	0	0	0	08.7	09.2	01.0012
		0	0	0	0	0	0	09.5	09.9	00.7009
		0	0	0	0	0	0	09.9	10.6	01.0009
		0	0	0	0	0	0	10.7	11.6	01.1010
		0	0	0	0	0	0			00.8009

Legend	Tree Type
A	Arctostaphylos
ED	Redwood
CC	Crown Class
D	Drainage
EP	Eastern Ponderosa
L	Location
PD	Plot Diagram
T	Tree

FIGURE 2.13 DATA ENTRY ON PSP HEADER SHEET

PERMANENT SAMPLE PLOT TALLY SHEET

FORESTRY, LANDS AND WILDLIFE
Forest Service

FORESTRY, LANDS AND WILDLIFE
Forest Service

Agency		Group Number		Plot No.		Measurements		Year		Month		Day		U.S.		Section		Twp		Rge		M		Plot Type		Info	
Agency		Group Number		Plot No.		Measurements		Year		Month		Day		U.S.		Section		Twp		Rge		M		Plot Type		Info	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2		0.3		0.4		0.5		0.6		0.7		0.8		0.9		1.0		1.1		1.2		1.3		1.4	
0.1		0.2																									

FIGURE 2.14 DATA ENTRY ON PSP TALLY SHEET

ENVIRONMENTAL PROTECTION

REGENERATION TALLY SHEET

Height Class 5	120 m
Height Class 4	90 m
Height Class 3	60 m
Height Class 2	30 m
Height Class 1	10 m
No Tally	

Page	3	of	3
Crew	Zagor, Marky Gubrank		

Agency	Occurrence Number	Plot ID	Measure (m ²)	Year	Month	Day	L.S.	Species	Tree	Range	ID	Tree Type	Imp.
01		020	1039005	110402	032	03	5000						

Record Type	Tree Number	Species	Height Class	Total (1)	Height Class 1 (10m - 20m)	Height Class 2 (20m - 30m)	Total (2)	Height Class 3 (30m - 40m)	Total (3)	Height Class 4 (40m - 50m)	Total (4)	Height Class 5 (50m - 60m)	Total (5)
02	93958	S.W.	1	003									
03	93988	S.B.											
03	93988	F.B.											
03	93988	P.L.											
03	93988	A.W.											
03	93988	S.W.											
03	93988	L.T.											
03	93988	F.D.											
03	93988	P.B.											
03	93988												

FIGURE 2.15 DATA ENTRY ON REGENERATION TALLY SHEET

FORESTRY, LANDS AND WILDLIFE
Forest Service

PERMANENT SAMPLE PLOT TALLY SHEET

[illegible]**FIGURE 3.1 PSP TREE TALLY CHECK SHEET**

FIGURE 4.1 TREE HEIGHT MEASUREMENT

Incorrect

Correct

FIGURE 4.2 HEIGHT MEASUREMENT OF LEANING TREES

FIGURE 4.3 CONK AND BLIND CONKS

FIGURE 4.4 OPEN SCARS

FIGURE 4.5 **LARGE BURL ON MAIN STEM**

This tree would be considered a fork if fork occurred above the DBH

FIGURE 4.6 **FORKS**

FIGURE 4.7 PRONOUNCED CROOK

FIGURE 4.8 LEANING TREE

FIGURE 4.9 ATROPELLIS CANKER

FIGURE 4.10 WITCHES ON SPRUCE

FIGURE 4.11 MISTLETOE

FIGURE 4.12 INSTRUCTIONS AND EXAMPLE OF THE USE OF THE 6 CLASS MISTLETOE RATING SYSTEM (HAWKSWORTH 1961, 1977)

CODES

- 91 One of the 1/3 sections has light infection (1) + other 2/3 have no visible infections.
- 92 Two of the 1/3 sections has light infection (10) + or one 1/3 section has a heavy infection only.
- 93 All three of the 1/3 sections has light infection (1) + or one 1/3 section has a heavy infection, one has a light infection + last 1/3 has no infection.
- 94 If total ratings = 4 then this code is used.
- 95 If total ratings = 5 then this code is used.
- 96 If total ratings = 6 then this code is used.

FIGURE 4.13 GENERIC WOODPECKER

FIGURE 4.14 YELLOWBELLIED SAPSUCKER FEEDING

FIGURE 4.15 SMALL MAMMAL FEEDING ON TREE BOLE

FIGURE 4.16 EXCAVATIONS BY WOODPECKERS

FIGURE 4.17 SMALL CAVITY

FIGURE 4.18 LARGE CAVITY

FIGURE 4.19 ALBERTA SURVEY SYSTEM²

² Alberta Bureau of Surveying and Mapping. 1986 Maps of Alberta Catalogue 1986-87. Government of Alberta ENR No. Ref 11 86 pp.iv.

FIGURE 4.20 PLOT BOUNDARIES

- - T bar and plot plaque at corner.
- L - Aluminum right-angle post at the exact corner.
- x ~ Galvanized metal post approximately 1 - 1 1/2 m. long.

FIGURE 4.21 STAND DYNAMICS PLOT LAYOUT

PERMANENT SAMPLE PLOT HEADER SHEET

Handwritten: 22

FIGURE 4.22 PLOT MAINTENANCE REPORT

FIGURE 4.23 SEEDLING NUMBERING

STAND DYNAMICS TALLY SHEET

Page 1 of 1

Record Type	Group No.	Plot No.	Plot Type	Year	Month	Day
C6	04	0210		910	08	22

Record Type 3 - Cruise Tally
Record Type 6 - Regeneration
Record Type 7 - Sapling
Record Type 8 - Tree Plot

Crew: LOGAN & KELSEY

Regen. Plot No.	Tree Number	Species	Species	DBH	DBH	Height	Height	Height to Live Crown	CC	Condition Codes	Azimuth	Distance	Comments
	11001	AW	AW	1.1	1.1	11.2	11.2	1.1					
	11002	AW	PL	1.1	1.1	11.4	11.4	1.1			312.0	16.9	SPECIES - VV
	11003	PL	PL	1.1	1.1	12.0	12.0	1.1			311.0	17.2	
	11004	PL	PL	1.1	1.1	12.0	12.0	1.1			311.0	17.6	
	11005	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11006	PL	PL	1.1	1.1	11.0	11.0	1.1					
	11007	PL	PL	1.1	1.1	11.1	11.1	1.1			321.1	16.9	
	11008	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11009	PL	PL	1.1	1.1	11.0	11.0	1.1					
	11010	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11011	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11012	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11013	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11014	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11015	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11016	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11017	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11018	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11019	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11020	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11021	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11022	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11023	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11024	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11025	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11026	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11027	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11028	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11029	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11030	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11031	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11032	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11033	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11034	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11035	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11036	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11037	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11038	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11039	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11040	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11041	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11042	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11043	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11044	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11045	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11046	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11047	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11048	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11049	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11050	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11051	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11052	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11053	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11054	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11055	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11056	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11057	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11058	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11059	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11060	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11061	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11062	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11063	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11064	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11065	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11066	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11067	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11068	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11069	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11070	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11071	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11072	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11073	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11074	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11075	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11076	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11077	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11078	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11079	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11080	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11081	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11082	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11083	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11084	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11085	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11086	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11087	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11088	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11089	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11090	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11091	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11092	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11093	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11094	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11095	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11096	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11097	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11098	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11099	PL	PL	1.1	1.1	11.1	11.1	1.1					
	11100	PL	PL	1.1	1.1	11.1	11.1	1.1					

STM 101

FIGURE 4.24 STAND DYNAMICS TALLY SHEET (CSTM 101) RECORD TYPE 6

Crew: SMITH & JONES

[illegible]

NOTES: 1. 20

Record Type	Group No.	Plot No.	Plot Type	Year	Month	Day
00	014	0210	1	90	08	22

Record Type 3 - Cruise Tally
Record Type 6 - Regeneration
Record Type 7 - Sapling
Record Type 8 - Tree Plot

Crew: Smith & Smith

Region: Plot No.	Tree Number	Species	Species	DBH	DBH	Height	Height	Height to Live Crown	CC	Condition Codes	Azimuth	Distance	Comments
	310101	PL	PL	110.2	111.9	110.6	111.2	112.0	C	02	1	271	16.4
	310102	PL	PL	110.1	112.1	10.1	110.6	111.0	C	00	1	143	14.9
	310103	PL	PL	111.1	111.1	111.1	110.0	112.9	C	00	1	151	10.0
	310104	PL	PL	111.1	111.6	111.1	111.9	117.9	C	00	1	310	11.9
	310105	PL	PL	111.1	110.0	111.1	111.9	117.1	C	00	1	318	16.0
	310106	PL	PL	111.1	111.1	111.1	117.6	116.2	C	00	1	216	17.1
	310107	PL	PL	111.1	112.3	111.1	113.2	111.0	C	01	1	1212	19.6
	310108	PL	PL	111.1	112.2	111.1	111.1	110.0	C	01	1	125	19.4
	310109	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310110	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310111	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310112	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310113	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310114	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310115	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310116	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310117	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310118	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310119	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310120	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310121	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310122	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310123	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310124	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310125	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310126	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310127	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310128	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310129	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1
	310130	PL	PL	111.1	111.1	111.1	111.1	111.1	C	01	1	111	11.1

FIGURE 4.26 STAND DYNAMICS TALLY SHEET (CSTM 101) RECORD TYPE 8 (TREE PLOT)

STAND DYNAMICS
REGENERATION HEIGHT CLASS RECORD

Page 1 of 1

Drawn: Smith + Jones

Record Type	Group No.	Plot No.	Plot Type	Year	Month	Day
61	04	020	9	91	06	06

Species	Height Classes (cm)											
	0-10	11-20	21-30	31-40	41-50	51-60	61-70	71-80	81-90	91-100	101-120	121 +
REGEN. PLOT NO.	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
1PL	006				002							
2Aw									003			005

FIGURE 4.27 STAND DYNAMICS REGENERATION HEIGHT CLASS RECORD (CSTM 100)
RECORD TYPE 61

FIELD NAME	TREES		TALL SHRUBS		LOW SHRUBS		GROUND SHRUBS		TOTAL SHRUBS		HERBS		GRASS		MOSS LICHEN		TOTAL ALL	
	%	VIGOR	%	VIGOR	%	VIGOR	%	VIGOR	%	VIGOR	%	VIGOR	%	VIGOR	%	VIGOR	%	VIGOR
PINNICON	5	2																
POPV TRE			4	2														
POPU BAL			2	2														
BETULAP			1	2														
ALNUCR			1	2														
SALISPP			1	2														
RIBELAC					0	2												
LEDUGRO					1	2												
LIINBDR							1	2										
VACC MYR							1	2										
VACC VIT							1	2										
MAIACAN											0	2						
EDILANG											4	2						
PVROSDP											0	2						
FRAGVIR											0	2						
ARALNUD											0	2						
DIKROSDP															5	2		
PIHEUSCH															1	2		
POLYJUN															1	2		
GAKEISD														8	2			

Record Type	Comments
C BROWSING & GRAZING -	
C DISEASE > 20%	
C OTHER -	

Master Table of Figures – PSP Manuals

ENVIRONMENTAL PROTECTION

(1) Plot Number	(2) Measure ment (3) No	(4) Year	(5) Month	(6) Day	(7) Plot Type	CREW:

NO CHANGES FROM LAST REMEAS. ☐ (8)

LEAVE BLANK IF RETREATMENT HAS NOT OCCURRED.

1. STAND TENDING YEAR OF RETREATMENT (9) (10)

TYPE OF RETREATMENT: THINNED ☐ (11) % (12) CLEANED ☐ (13) % (14)

DEC. ONLY ☐ (15)

FERTILIZED ☐ (16) TYPE (17)

RATE (18) kg/ha

2. REPLANTED SPECIES (19) STOCKING (20) DATE (21) (22)

MO YR

3. WEATHER / NATURAL DAMAGE:

FLOODING ☐ (23) % (24)

BROWSING ☐ (25) % (26)

SNOW PRESS ☐ (27) % (28)

FROST DAMAGE ☐ (29) % (30)

4. PHYSICAL DESCRIPTION:

WINDROW IN TREE PLOT ☐ (31)

EXPOSED MINERAL SOIL ☐ (32) % (33)

STANDARD BUFFER (Y / N) ☐ (34)

5. PLOT SIZES:

SAPLING (35) m X (36) m

TREE (37) m X (38) m

COMMENTS

FIGURE 4.29 PLOT RETREATMENT REPORT (RECORD TYPE 9)

Record Type	Grp.	Plot #	Plot Type	Year	Month	Day	Meas. #	Crew	Page	1 of	3	1986 HT	1987 HT	1988 HT	1989 HT	1990 HT	1991 HT	1992 HT	1993 HT	1994 HT	1995 HT	1996 HT
6	4	3		95	05	01			Gilday + Yusep													
Regen. Plot #	Tree #	Sp.	Tree Type	Byear	HT1	Byear + Year	1983 HT	1984 HT	1985 HT	1986 HT	1987 HT	1988 HT	1989 HT	1990 HT	1991 HT	1992 HT	1993 HT	1994 HT	1995 HT	1996 HT		
1	1001	PL	A	85	30	90								42	55	62	75	91	105	121		
1	1008	PL	I	87	2	-							4	10	18	25	40	58	74	102		
1	1010	SW	A	71	54	78	64	72	70	91	104	121	132	145	161	180	191	205	218	234		
	CC																					
							1994 HT	1995 HT	1996 HT													
1	1015	AW	I	80			252	286	295				457									
2	1051	PL	A	79	25	83		50	101	155	205	280	370	450	490	550	720	840	910	1996 HT		
2	1058	PL	I	89	20	92														980		
2	1082	PL	I	90	18	92														35		
2	1074	PL	I	90	25	93														24		
2	1082	PL	I	80	-	-												32	38	45		
2	1081	PL	I	90	-	-														62		
																				42		
7	2095	PL	I	84	15	87						25	35	45	62	88	101	150	240	355		
7	2110	PL	I	85	18	88							30	42	55	65	91	121	201	305		
7	2004	PL	I	85	22	88					40	52	65	81	111	155	210	310	410	490		
7	2010	PL	I	85																590		

FIGURE 4.30 STAND DYNAMICS TREE AGE TALLY SHEET

FIGURE 4.31 DIE BACK

3 3286 53216566 5