S Sullivan, Daniel
632.6 Determination of
3232 an effective
A3tr82-4 single lethal dose
for bromadiolone
to Columbian
ground squirrels

MONTANA STATE LIBRARY

S 632.63232 A3tr82-4 c.1 Sullivan
Determination of an effective single lot

3 0864 00061689 9

STATE DOCUMENTS COLLECTION

DEC 1 1988

MONTANA STATE LIBRARY
1515 E. 6th AVE.
HELENA, MONTANA 59620

MONTANA DEPARTMENT OF AGRICULTURE
ENVIRONMENTAL MANAGEMENT DIVISION
TECHNICAL SERVICES BUREAU
HELENA, MONTANA 59620

Determination of an Effective Single Lethal Dose for
Bromadiolone to Columbian Ground Squirrels

by
Daniel Sullivan

ABSTRACT

A single dose feeding of bromadiolone treated hulled oats resulted in death to 26 of 30 Columbian Ground Squirrels (Spermophilus columbianus) tested. No animals survived an oral dose of bromadiolone greater than 2.10 mg/kg. Results indicate bromadiolone has the potential to be effective in single doses for control of Columbian Ground Squirrels.

Technical Report 82-4

December, 1982

PLEASE RETURN

INTRODUCTION

Presently there is a lack of effective alternatives to traditional single dose field rodenticides such as strychnine, 1080 and zinc phosphide. Each of these traditional compounds has environmental, efficacy and/or economic disadvantages. These compounds have uncertain futures regarding registration as rodenticides.

The use of strychnine treated grain baits for control of the Columbian Ground Squirrel (Spermophilus columbianus) has traditionally obtained poor results (Albert and Record, 1979, Record, 1978 and 1976; Sullivan and Baril, 1981). Restrictions are currently under consideration by the Environmental Protection Agency and the State of Montana on the use of strychnine because of hazards to nontarget species. If inacted these restrictions will prohibit or make impractical the use of strychnine for rodent control in many circumstances. Sodium monofluoroacetate (1080) grain baits have been used in Montana in 1979, 1981 and 1982 under a specific exemption (FIFRA, Section 18). Compound 1080 has given excellent control of Columbians (Albert and Record, 1979; Lingel-Pate, 1981). Future registration for 1080, either temporary or long term, is uncertain. The use of 1080 has been controversial because of the hazard to nontarget species and questions regarding 1080's humaneness. Zinc phosphide has given effective control of Columbian ground squirrels (Sullivan and Baril, 1981) but does have problems. It requires prebaiting,

possesses high potential for nontarget deaths and has shown to be erratic in control (Baril, 1980; Albert and Record, 1979). Zinc phosphide is not currently registered for Columbian Ground Squirrels. Currently registered anticoagulants require repeat feeding from bait stations to obtain a lethal dose. Except for high value lands or crops repeat applications or use of bait stations may not be economically practical for most field rodent control. For these reasons it seems prudent to evaluate and develop alternative rodenticides.

Suitable alternatives should be relatively selective to the target animals, either biologically or through selective formulation of the carrier. Suitable alternatives should be single dose, have a high degree of effectiveness and be economically feasible.

Bromadiolone is an anticoagulant rodenticide that has been shown effective for control of commensal rodents (Marsh, et. al., 1980; Meehan, 1978; Redfern and Gill, 1980). It has received federal registration for commensal rodent control (Lipha Chem. Co.; Bell Laboratories). Recent studies have shown potential for control of ground squirrels (Baker, 1980; Marsh, 1977). LD₅₀ studies on rats and mice (1.125 mg/kg - rats; 1.75 mg/kg - mice [Anon., undated]) indicate bromadiolone may be effective as a single dose rodenticide for the Columbian Ground Squirrel.

Bromadiolone has been shown to have a relatively low risk from secondary poisoning (Mendenhall and Pank, 1980; Anon., undated). Should bromadiolone prove to be effective as a single dose rodenticide it may be an economical alternative to the traditional acute rodenticides.

OBJECTIVE

To determine an effective single lethal dose for bromadiolone to Columbian Ground Squirrels.

METHODS

Procedures for determining an effective single lethal dose were modeled after recommendations by R.E. Marsh (pers. com.).

Thirty adult Columbian Ground Squirrels were trapped during June, 1982, from the Helena valley, Lewis and Clark County, Montana. Squirrels were captured in livetraps (6X6X19 in., Tomahawk No. 202) bait with molasses flavored rolled barley. Squirrels were placed individually in 9X12X12 inch cages. Water and food (whole oats) were provided ad libitum. Squirrels were allowed to acclimate to the caged environment for at least 1 week prior to testing.

The squirrels were assigned to 5 groups of 6 animals each, 3 males and 3 females. Squirrels were distributed among the groups

on the basis of size to reduce variations because of weight. All animals were eating regularly and appeared healthy prior to testing.

Each group of squirrels was fed one of the following doses of bromadiolone on whole hulled oats: 0.25, 0.50, 0.75, 1.00, and 1.50 mg active ingredient. The untreated whole oats were removed and the cages cleaned prior to placing the bromadiolone bait in the feeding trays. The bromadiolone bait was left in the trays until consumed by the squirrels. Any bait dropped through the cage floor was collected and weighed.

Test baits were provided by personnel of the Chempar Products Division of Lipha Chemical Company. The quantity of bait received by the squirrels in each test group was 5 grams (0.25 mg and 0.75 mg AI) or 10 grams (0.50 mg, 1.00 mg and 1.50 mg AI) depending on the bromadiolone concentration on the test baits.

Squirrels were weighed to the nearest 0.1 gram using a triple beam balance prior to dosing. Weight of each squirrel was taken at death.

Squirrels were observed for 35 days after dosing. All squirrels alive at the end to the test period were killed using CO₂ gas.

RESULTS AND DISCUSSION

Tables 1 and 2 show that squirrel mortality occurred at each dose level administered. Although mortality occurred as low as 0.36 mg/kg in this study, development of a practical single dose bait requires that all ground squirrels consuming the bait receive a lethal dose of the toxicant. The lowest dose above which survival did not occur was 2.10 mg/kg (Table 1). Considering the small sample size used in this study, variations in ground squirrels and their environment and variables introduced during practical applications of rodent control programs the minimum dose for all squirrels should be no less than 2.10 mg/kg. To allow for the possibility that some animals may survive a 2.10 mg/kg dose increasing the dose up to 25% for field trails is recommended (R.E. Marsh, pers. com.).

CONCLUSIONS

Single dose feedings of bromadiolone treated bait to captive Columbian Ground Squirrels ranging from 0.36 to 3.78 mg/kg resulted in death to 26 of 30 squirrels tested. No squirrel survived doses greater than 2.10 mg/kg. These results indicate bromadiolone treated baits may be an effective single dose field rodenticide. Evaluations need to be conducted to determine bromadiolone's effectiveness under field situations.

Table 1. Death (D) and Survival (S) of Columbian Ground Squirrels 35 days after receiving a single dose of bromadiolone on whole hulled oats.

<u>Animal</u> <u>No.</u>	<u>Dose AI</u> <u>(mg)</u>	<u>AI Intake</u> <u>(mg/kg)</u>	<u>Results</u>
1-25	0.25	0.36	D
4-25	0.25	0.41	D
2-25	0.25	0.46	D
5-25	0.25	0.47	D
6-25	0.25	0.49	S
3-25	0.25	0.52	D
1-50	0.50	0.77	D
2-50	0.50	0.92	D
4-50	0.50	0.92	D
5-50	0.50	0.97	D
6-50	0.50	1.13	D
1-75	0.75	1.13	S
3-50	0.50	1.20	D
2-75	0.75	1.31	D
4-75	0.75	1.33	D
5-75	0.75	1.36	D
1-100	1.00	1.49	D
4-100	1.00	1.59	D
3-75	0.75	1.76	D
2-100	0.75	1.87	S
5-100	1.00	1.89	D
6-100	1.00	1.89	D
6-75	0.75	2.02	S
3-100	1.00	2.10	D
1-150	1.50	2.20	D
4-150	1.50	2.78	D
2-150	1.50	2.87	D
5-150	1.50	2.97	D
6-150	1.50	3.28	D
3-150	1.50	3.78	D

Table 2. Average intake per kilogram of body weight, mortality and weight change of Columbian Ground Squirrels following a single oral dose of bromadiolone on hulled oats.

<u>mg/squirrel</u>	<u>Dose</u>	<u>Mortality</u>	<u>Days to Death*</u>		<u>Weight Change</u>
	<u>Ave. intake (mg)</u> <u>per kg (Range)</u>		<u>Mean</u>	<u>Range</u>	
0.25	0.45 (0.36-0.52)	5/6	9.2	5-24	- 7.8%
0.50	0.97 (0.77-1.20)	6/6	6.3	5-8	- 4.0%
0.75	1.48 (1.13-2.02)	4/6	10.5	5-18	-14.7%
1.00	1.81 (1.49-2.10)	5/6	11.4	7-19	-15.3%
1.50	2.98 (2.20-3.78)	6/6	10.0	7-18	-10.2%

*Calculations include only those squirrels that died during the test.

ACKNOWLEDGEMENTS

Financial assistance and bait was provided by the Chempar Products Division of Lipha Chemical Company. I am grateful to Dr. Rex E. Marsh for advice on study design and review of the data. Steve Baril assisted in trapping and care of the squirrels. I would like to thank George Algard, Steve Baril and Monty Sullins for reviewing the manuscript and Pam Schmitz for typing it.

LITERATURE CITED

- Albert, S.W., Record, C.R. 1979. Efficacy and cost of four rodenticides for Columbian ground squirrel damage control in Western Montana. Montana Department of Livestock Report. Helena, Montana. 17 pp.
- Anon., undated. Maki, Bromadiolone, Rodenticide Technical Bulletin. Chempar Division of Lipha Chemical, Inc. 9 pp.
- Baker, Rex O. 1980. Field efficacy study bromadiolone oat groat bait - ground squirrels (*Spermophilus beecheyi*) in California farm area. Agricultural Biology, California State Polytechnic University, Pomona, Calif. 7 pp.
- Baril, Steven F. 1980. Efficacy of 2% Zinc Phosphide applied with and without a prebait for control of the Columbian Ground Squirrel. Montana Department of Agriculture Report. 9 pp.
- Lingel-Pate, T. 1981. Recovery of a ground squirrel population following application of 1080 and effects of the poisoning program on coyotes and ravens. Unpublished preliminary report, Montana Cooperative Wildlife Research Unit, University of Montana., Missoula, Montana. 35 pp.
- Marsh, R.E. 1977. Bromadiolone, a new anticoagulant rodenticide. EPPO Bull. 7(2): 495-502.
- Marsh, R.E., Howard, W.E., Jackson W.B. 1980. Bromadiolone, a new toxicant for rodent control. Pest Control (August): 22-26.
- Meehan, A.P. 1978. Rodenticidal activity of Bromadiolone, a new anticoagulant. Proceeding 8th Vertebrate Pest Conference. pp. 122-126.
- Mendenhall, V.M. and Pank, L.F. 1980. Secondary poisoning of owls by anticoagulant rodenticides. Wildlife Soc. Bull. 8(4): 311-315.
- Record, C.R. 1978. Ground squirrel and prairie dog control in Montana. Montana Department of Livestock Report. 12 pp.
- Record, C.R. 1976. A field evaluation of strychnine and sodium monofluoroacetate grain baits for Columbian ground squirrel control. Montana Department of Livestock Report. Helena, Montana. 17 pp.
- Redfern, R. and Gill, J. E. 1980. Laboratory evaluation of Bromadiolone as a rodenticide for use against warfarin-resistant and non-resistant rats and mice. J. Hyg., Camb., 84: 263-268.

Sullivan, Daniel and Baril, Steve F. 1981. A field efficacy study of strychnine and zinc phosphide grain baits using nontoxic prebait for the control of the Columbian ground squirrel (Spermophilus columbianus). Montana Department of Agriculture Technical Report 82-2. Helena, Montana 59620.