S Montana Parks
632.954 Division
F2dnwev Noxious weed and
1994 exotic vegetation
management plan,
Montana Fish,
Wildlife and

DRAFT

DECEMBER • 1994

NOXIOUS WEED AND EXOTIC VEGETATION

MANAGEMENT PLAN AND ENVIRONMENTAL ASSESSMENT

**Montana Fish,
Wildlife & Parks**

Region 1

STATE DOCUMENTS COLLECTION

MAY 30 1995

MONTANA STATE LIBRARY
1515 E. 6th AVE.
HELENA, MONTANA 59620

PLEASE RETURN

MONTANA STATE LIBRARY

S 632.954 F2dnwev 1994 c.1

Noxious weed and exotic vegetation manag

3 0864 00093897 0

NOXIOUS WEED AND EXOTIC VEGETATION MANAGEMENT PLAN

**Montana Fish, Wildlife and Parks
Region One**

**Prepared by
State Parks Division
490 North Meridian Road
Kalispell, Montana 59901**

DECEMBER 1994

TABLE OF CONTENTS

	Page
LIST OF TABLES	iii
LIST OF FIGURES	iv
1.0 INTRODUCTION	1
1.1 Program Guidance and Objectives	1
1.2 Existing Statutory Requirements	2
1.3 Overview and Setting	2
1.3.1 Property Groups	4
2.0 MANAGEMENT STRATEGIES	15
2.1 General Approach	15
2.2 Conceptual Framework to Prioritize Activities	15
2.2.1 Species Specific Directives	15
2.2.2 Location and Population Size	16
2.2.3 Special Management Zones	17
2.3 Treatment Methods and Techniques	21
2.3.1 Manual	21
2.3.2 Cultural	21
2.3.3 Biological	22
2.3.4 Chemical	23
2.4 Inventory and Monitoring Procedures	26
2.5 Other Subjects of Concern	26
2.5.1 Oversight of Plan Implementation	26
2.5.2 Threatened, Endangered, and Sensitive Species	26
2.5.3 Applicator Licenses and Record Keeping	27
2.5.4 Herbicide Labels	27
2.5.5 Contracted Services	27
3.0 PROPERTY GROUP TREATMENT PLANS	29
3.1 Wildlife Management Areas and Affiliated Lands	29
3.1.1 Introduction	29
3.1.2 Kootenai/DeRozier-Woods Ranch	32
3.1.3 Kootenai Falls	32
3.1.4 West Kootenai	34
3.1.5 Mount Silcox	34
3.1.6 Ninepipe	35
3.1.7 Pablo	35
3.1.8 Ray Kuhns	35
3.1.9 Roundhorn	36
3.1.10 Buffalo Head Park, Flathead Lake, Flathead River, Lower Stillwater Lake, Swan Lake, Thompson Falls WHPA's . . .	37

TABLE OF CONTENTS, continued

	Page
3.2 Fishing Access, Affiliated Lands and Administrative Sites	37
3.2.1 Introduction	37
3.2.2 Clark Fork River Drainage	41
3.2.3 Flathead River Drainage	43
3.2.4 Kootenai River Drainage	45
3.2.5 Fish Hatcheries and Administrative sites	46
3.3 State Parks and Affiliated Lands	49
3.3.1 Introduction	49
3.3.2 Flathead Lake	52
3.3.3 Flathead Lake - Wild Horse Island	53
3.3.4 Lake Mary Ronan	55
3.3.5 Logan	55
3.3.6 Lone Pine	56
3.3.7 Thompson Falls	57
3.3.8 Whitefish Lake	58
4.0 REFERENCES	60
5.0 APPENDICES	
Appendix A County Noxious Weed Control Act of 1979	
Appendix B Synopsis of Noxious Weed and Exotic Plant Species	
Appendix C Laws and Policies Relating to the Management of Noxious Weeds	
Appendix D Biological Control Agents Available in Montana and Those Released on Fish, Wildlife and Parks Properties	
Appendix E Herbicide Emergency Response Plan	
Appendix F Sensitive Species Occurring in the Vicinity of Fish, Wildlife and Parks Region One Properties	
Appendix G Herbicides Proposed for Use on Fish, Wildlife and Parks Region One Properties	
Appendix H Herbicide Label Instructions and Safety Precautions	

LIST OF TABLES

	Page
Table 1. Noxious weed species present and status on Department of Fish, Wildlife and Parks Wildlife Management Areas in Region One	7
Table 2. Noxious weed species present and status on Department of Fish, Wildlife and Parks Fishing Access in Region One by drainage	9
Table 3. Noxious weed species present and status on Montana State Parks in Region One	12
Table 4. Soil index values, risk classes, and soil descriptions used to identify groundwater areas sensitive to chemical contamination	19

Digitized by the Internet Archive
in 2016

<https://archive.org/details/noxiousweedexoti1994mont>

LIST OF FIGURES

	Page
Figure 1	Properties owned by or leased to Montana Fish, Wildlife & Parks in Region One 5
Figure 2.	Wildlife Management Areas (WMA's) administered by Montana Department of Fish, Wildlife and Parks in Region One 31
Figure 3.	Fishing Access administered by Montana Fish, Wildlife & Parks in Region One 39
Figure 4.	State Parks administered by Montana Fish, Wildlife & Parks in Region One 51

1.0 INTRODUCTION

1.1 Program Guidance and Objectives

Montana Fish, Wildlife & Parks (FWP) recently adopted its Vision document which describes the direction in which FWP is heading and how the goals articulated within that document will be reached (FWP 1992). The basic Mission of the agency is:

"to provide for the stewardship of the fish, wildlife, parks, and recreational resources of Montana, while contributing to the quality of life for present and future generations."

A key element of all Montana Fish, Wildlife & Parks programs is to consider how policies and programs affect adjacent lands, and in turn, how adjacent lands affect FWP properties. This perspective is rooted in a strong commitment to public involvement in management decisions. In 1983, FWP adopted a Noxious Weed Control Program for lands managed by the agency (FWP 1983). The objective of the program is:

"to prevent, to the greatest extent possible, the reproduction, proliferation, and distribution of agriculturally undesirable plant species throughout department lands or from department lands onto adjacent lands."

Overall direction for regional implementation of the Noxious Weed Management Program is provided by the Vision document and statutory requirements based in state law and local ordinances. The purpose of FWP Region One's Noxious Weed and Exotic Vegetation Management Plan is to describe the strategies and management actions necessary for the prevention, stabilization, control, or eradication of noxious weed and some categories of exotic plant species on FWP lands.

Management actions must be appropriate to land classification and use, and must satisfy FWP's legal mandate to serve as steward of specific natural and recreational lands within northwestern Montana. This Plan outlines an integrated approach to weed management based on site conditions by employing a variety of treatment methods and techniques including manual, cultural, biological, and chemical. More specifically, the objectives of the Plan are to:

- Meet legal requirements (including the County Noxious Weed Control Act and the Montana Environmental Policy Act) to manage plants declared as noxious weeds by the State and by the Flathead, Lake, Lincoln, and Sanders county weed boards.
- Develop and implement an integrated approach to noxious weed management and a strategy to prioritize treatment areas.
- Attempt to contain, reduce or eradicate weed populations, depending on the location and status of the population.

- Prevent the introduction and establishment of new noxious weed species, and exotic plant species in Natural Zones of State Parks, to the extent feasible.
- Work cooperatively with adjacent landowners and the County Weed Boards to enable consistent and sound weed management efforts.
- Conduct treatments in an ecologically responsible manner to maintain fish, wildlife, and vegetation resources and protect human health and safety.
- Involve the public in the planning process.

1.2 Existing Statutory Requirements

The State of Montana has recognized the damaging effects of noxious weeds in laws and regulations. The County Noxious Weed Control Act of 1979 (MCA 7-22-21-1 *et seq.*, as amended 1991, Appendix A) defines "noxious weeds" as "any exotic plant species established or that may be introduced in the state which may render land unfit for agriculture, forestry, livestock, wildlife, or other beneficial uses or that may harm native plant communities..." Administered at the county level, this law requires counties to develop weed control districts to plan and implement a weed control program. County commissioners appoint a district weed board which is responsible for the administration and enforcement of the district's noxious weed control efforts. The law provides for the promulgation of rules to list statewide noxious weeds and, it allows for county-wide listing of additional species. In addition, this law also gives specific direction to counties for cooperation with state and federal-aid programs. Furthermore, the law states that "it is unlawful for any person to permit any noxious weed to propagate or go to seed on his [or her] land" (7-22-2115) and, it encourages landowners to file weed control plans (7-22-2116).

The weed species declared as noxious by the State of Montana (through the Department of Agriculture) and by the individual county weed boards within the administrative boundaries of FWP Region One are listed in Appendix B. The treatment of noxious weeds, particularly through the use of herbicides, is also subject to other state and federal laws as well as the policies and programs of other agencies. These are summarized in Appendix C.

1.3 Overview and Setting

Exotic (non-native) "weedy" plants have been noted in Montana throughout this century. A number of these species are increasing in quantity, area and density. This presents a problem for the perpetuation of native plant communities and ecological processes on FWP properties. It also impacts the quality of recreation and increases the potential for spread to lands adjacent to FWP properties.

"Native" plant species occur and evolve naturally without human intervention or manipulation. Species that move into an area without the aid of humans are considered native. A goal of FWP is to perpetuate native plants and animals as part of natural ecosystems.

"Exotic" plant species are those that occur in a given place as a result of direct or indirect, deliberate or accidental actions by humans (not including deliberate reintroduction). The exotic species introduced because of such human action have not evolved with species native to the place in question, and therefore are not a natural component of the ecological system characteristic of that place.

A "weed" is any plant that is out of place, is undesirable and is growing where it is not wanted. "Noxious weeds" are plants in the wrong place that are harmful or destructive to man or other organisms (Holmes, 1979; Webster, 1981). Noxious weeds are prohibited or restricted by Montana law. Transporting seed or parts of these plants, or allowing them to seed on one's property is prohibited.

A "pest" is a population of plants or animals that interfere with the purpose of a state park.

FWP draws a distinction between "weedy" species and "noxious" species. FWP recognizes the value of some weed species in Wildlife Management Areas as food and/or cover. For example, some "weedy" species in the composite family (e.g. dandelion, prickly lettuce, and goat's beard) are important components of upland game bird summer diets. As declared in law, "noxious" weeds are those non-native species which are very aggressive and successful competitors for nutrients, moisture, and space. In the past few decades, some noxious weed species have become significant components of local vegetation communities, actually dominating some sites. Noxious weeds may reduce vegetation biodiversity and may limit the diversity of wildlife. Native vegetation species are also more desirable for providing erosion control and visual aesthetics.

Noxious weeds occur on many sites currently administered by FWP, although most invasions occurred prior to FWP acquisition. The number of species and the total coverage vary considerably by site and the nature of land use. Important factors influencing noxious weed establishment include proximity to water, previous overgrazing practices, human activities, site disturbance and seed dispersal -- the latter two of which can be aggravated by road access.

This Plan considers the treatment and management of noxious weeds on 74 separate properties totalling 18,178 acres owned by or leased to FWP Region One within Flathead, Lake, Lincoln and Sanders counties (FWP 1994). Figure 1 illustrates the location of these properties. Several land parcels owned by FWP are leased to other individuals or municipalities. This Plan will note the location of such properties but not address noxious weed management as individual leases may specifically address the topic. Chapter 2.0 identifies strategies to prioritize treatment activities and reviews methods and techniques of noxious weed treatments. Chapter 3.0 outlines specific weed management programs for the groups of properties outlined below.

1.3.1 Property Groups

Lands administered by FWP are classified according to the Department Land Classification Manual (FWP 1988) based on management, uses and funding sources. The Wildlife Division manages Wildlife Management Areas (WMA's) and Wildlife Habitat Protection Areas (WHPA's). These areas are set aside primarily to preserve wildlife habitat and secondarily to provide opportunities for both consumptive and non-consumptive outdoor recreation. There are 8 WMA's and 6 WHPA's totalling 9,774 acres in Region One.

The Fisheries Division oversees the management of Fishing Access Areas, Sites, Greenways, Hatcheries and Administrative Sites. Maintenance activities on Fishing Access Areas and Sites (FAS) is the responsibility of the Parks Division. Fishing Accesses provide the public with opportunities to access river corridors and lakes to fish, boat, hunt, picnic, watch wildlife, camp, etc. There are 33 Fishing Accesses and 9 affiliated lands totalling 5,411 acres in Region One.

The Parks Division manages State Parks classified as either: natural, historic, or recreation. State Parks are established for the purpose of conserving the scenic, historic, archaeologic, scientific, and recreational resources of the state and providing for their use and enjoyment, thereby contributing to the cultural, recreational, and economic life of the people and their health (23-1-101 MCA). The Parks division also oversees the maintenance of all FAS's and the Region One Headquarters complex. There are 6 State Parks (SP's) and 2 affiliated lands totalling 2,993 acres in Region One.

One administrative site for the Region One Headquarters complex, totalling 3.27 acres, is owned by FWP. The Parks Division oversees the maintenance of this administrative site.

Because of differences in land use, site disturbance, and the degree of noxious weed establishment, these lands have different opportunities and constraints for the management of noxious weeds. Therefore, this Management Plan will examine groups of properties where similarities merit and will direct appropriate management strategies accordingly. Site specific management strategies will be outlined for property groups, while recognizing that some deviations will occur. These differences will be highlighted and addressed more specifically in Part 3.0 of this document. The following discussion describes the groupings in general, whereas Tables 1, 2, and 3 outline the property groups, list discrete properties by name, and report the findings of the most recent noxious weed inventory conducted on each property during summer, 1993.

Figure 1.

Legend

- Fishing Access Site
State Park
Wildlife Management Area
Wildlife Habitat Protection Area

Key to the Fishing Access Sites

CLARK FORK RIVER DRAINAGE

8. Flatiron Ridge	30. Thompson Chain of Lakes

FLATHEAD RIVER DRAINAGE

1. Ashley Creek	7. Ducharme	19. Old Steel Bridge	26. Spring Creek
2. Ashley Lake	11. Horseshoe Lake/Ferndale	20. Pressentine	27. Swan River
3. Beaver Lake	12. Kokanee Bend	21. Skyles Lake	28. Teakettle
4. Bigfork	13. Kuhns	22. Smith Lake	31. Walstad
5. Blanchard Lake	15. Loon Lake/Ferndale	23. Somers	32. Whitefish River
6. Bootjack Lake	17. Marlowe Springs	25. Sportsmans Bridge	33. Woods Bay

KOOTENAI RIVER DRAINAGE

9. Frank Lake	14. Loon Lake/Eureka	18. Moran Lake	29. Tetrault Lake
10. Glen Lake	16. Marl Lake	24. Sophie Lake	

Table 1. Noxious weed species present and status on Department of Fish, Wildlife and Parks (FWP) Wildlife Management Areas in Region One.

FWP PROPERTY NAME and ACREAGE	WEED SPECIES PRESENT	WEED STATUS		
		SIZE ¹	DENSITY ²	LOCATION
Kootenai/DeRozier- Woods Ranch WMA (1,357)	spotted knapweed	25	MD	roadsides, meadows in section 2, fencelines
Kootenai Falls WMA (344.23)	spotted knapweed	15	LDA	roadsides, hay meadows, riverbank
West Kootenai WMA (920.12)	spotted knapweed	< 1	VLD	Has not been detected
Mount Silcox WMA (1,552.3)	spotted knapweed	189	LBD	mostly in foothills
	St. Johnswort	3	VLD	very spotty
Ninepipe WMA (3,142.36)	spotted knapweed	unk	LDB	roadsides,
	whitetop	unk	MD	roadsides,
	Russian knapweed	unk	LDB	old fields
	Canada thistle	unk	LDA	roadsides
Pablo WMA (415.92)	spotted knapweed	unk	LDB	roadsides
	Canada thistle	unk	LDB	roadsides
Ray Kuhns WMA (1,609.3)	spotted knapweed	unk	LDA	roadsides
RoundHorn WMA (27)	spotted knapweed	5	MD	pastureland

¹ SIZE: total acres, including roadsides and scattered patches away from roads

² DENSITY: number of weed stems in the population

VLD:	very low density	--	<i>one or a few individual plants, accounting for a very small proportion of the vegetative ground cover; native species well represented</i>
LDA:	low density - A	--	<i>individual <u>plants clumped in small areas</u>, accounting for a small proportion of the vegetative ground cover; native species well represented</i>
LDB:	low density - B	--	<i>individual <u>plants scattered throughout site</u>, accounting for a small proportion of the vegetative ground cover; native species well represented</i>
MD:	moderate density	--	<i>individual <u>plants numerous</u>, accounting for a moderate proportion of the vegetative ground cover; native species present</i>
HD:	high density	--	<i>individual <u>plants abundant</u>, accounting for a high proportion of the ground cover; native species present but not well represented</i>
VHD:	very high density	--	<i>individual <u>plants very abundant</u>, accounting for almost all of the vegetative ground cover; native species almost absent</i>

Table 2. Noxious weed species present and status on Department of Fish, Wildlife and Parks (FWP) Fishing Access Sites in Region One, by drainage.

FWP PROPERTY NAME and ACREAGE	WEED SPECIES PRESENT ¹	WEED STATUS SIZE ²	DENSITY ³
CLARK FORK RIVER			
Flatiron Ridge (24.9)	SK	20	MD
Thompson Chain of Lakes (3,965.07)	SK, CT, SJ	1500, 900, 10	MD, MD, VLD
FLATHEAD RIVER			
Ashley Creek (18.25)	unknown		
Ashley Lake (31.56)	unknown		
Beaver Lake (2)	SK	1/4	VLD
Bigfork (0.69)	NONE		
Blanchard Lake (9.31)	SK,CT	1,4	LDA,LDB
Bootjack Lake (0)	unknown		
Ducharme (51.25)	SK	1/4	LDA
Horseshoe Lake/Ferndale (22.92)	unknown		
Kokanee Bend (172.34)	SK, LS, CT	15, 3, 1	MD, LDA, VLD
Kuhns (82.59)	unknown		
Loon Lake/Ferndale (4.36)	unknown		
Marlowe Spgs (13.53)	unknown		
Old Steel Bridge/Kiwanis (122.3)	SK, CT	25, 2	MD, LDA
Pressentine (7.02)	SK, CT	4, 1/8	LDB, VLD
Skyles Lake (0.75)	SK	1/2	LBD
Smith Lake (3.16)	SK,CT	1/2, 1 1/2	LDA,LDB
Somers (2)	SK	1/64	LDA
Sportsmans Bridge (5.77)	SK	1/16	LDA
Spring Creek (240)	unknown		
Swan River (58.52)	SK	1/16	VLD
Teakettle (2.5)	SK	1/64	MD
Walstad (4.57)	CT	1/64	MD
Whitefish River (3.29)	NONE		
Woods Bay (11.7)	SK	1/2	VLD

FWP PROPERTY NAME and ACREAGE	WEED SPECIES PRESENT ¹	WEED STATUS SIZE ²	DENSITY ³
KOOTENAI RIVER			
Frank Lake (88.88)	unknown		
Glen Lake (1.4)	NONE		
Loon Lake/Eureka (0.56)	unknown		
Marl Lake (3.02)	unknown		
Moran Lake (0)	unknown		
Sophie Lake (17.82)	SK	6	LDB
Tetrault Lake (7.88)	SK	5	HD
FISH HATCHERIES and ADMINISTRATIVE SITES			
Flathead Lake Salmon Hatchery (25.47)	SK	1/2	VLD
Jocko River Trout Hatchery (59.71)	SK	6	LDA
Lake Mary Ronan Spawn Station (40.27)	unknown		
Lore Lake Spawn Station (0)	unknown		
Libby Field Station (67.57)	SK,CT	11	LDA
Lone Lake Hatchery (4.33)	unknown		
Region One Headquarters Complex (3.27)	NONE		
Rogers Lake Spawn Station (0.63)	unknown		
Rose Creek Hatchery Site (20.2)	CT	10	LDA

¹ Weed species abbreviations:

SK = spotted knapweed	LS = leafy spurge	ST = sow thistle
CT = Canada thistle	MT = musk thistle	BT = bull thistle
MU = mullein	DT = dalmatian toadflax	SJ = St. Johnswort
OW = other weed-like vegetation		

² SIZE: total acres or partial acres for each species, respectively:

1 acre = 43,560 ft ² (200' x 200')	1/16 acre = 2,722 ft ² (50' x 50')
1/2 acre = 21,780 ft ²	1/32 acre = 1,361 ft ²
1/4 acre = 10,890 ft ² (100' x 100')	1/64 acre = 680 ft ² (25' x 25')
1/8 acre = 5,445 ft ²	< 1/64 acre estimation in square feet

³ DENSITY: number of weed stems in the population, respectively:

VLD: very low density	--	<i>one or a few <u>individual plants</u>, accounting for a very small proportion of the vegetative ground cover; native species well represented</i>
LDA: low density - A	--	<i>individual <u>plants clumped in small areas</u>, accounting for a small proportion of the vegetative ground cover; native species well represented</i>
LDB: low density - B	--	<i>individual <u>plants scattered throughout site</u>, accounting for a small proportion of the vegetative ground cover; native species well represented</i>
MD: moderate density	--	<i>individual <u>plants numerous</u>, accounting for a moderate proportion of the vegetative ground cover; native species present</i>
HD: high density	--	<i>individual <u>plants abundant</u>, accounting for a high proportion of the ground cover; native species present but not well represented</i>
VHD: very high density	--	<i>individual <u>plants very abundant</u>, accounting for almost all of the vegetative ground cover; native species almost absent</i>

Table 3. Noxious weed species present and status on Montana State Parks in Region One.

FWP PROPERTY NAME and ACREAGE	WEED SPECIES PRESENT ¹	WEED SPECIES SIZE ²	WEED SPECIES DENSITY ³
Flathead Lake/Big Arm (217.49)	SK, CT,SC	150, 1, 1/64	LDB, HD, VLD
Flathead Lk/Finley Point (27.83)	SK, CT	2, 1/64	LDA, MD
Flathead Lake/Wayfarers (68.86)	SK, CT	10, 1/2	LDA, MD
Flathead Lk/West Shore (129.25)	SK	16	LDB
Flathead Lk/Wild Horse Is (2,163.92)	SK, MU	10, 10	LDA, LDA
	LS,OW	1/8, 1/64	MD, MD
	CT	1/64, 1/64, 1/64	MD, MD, MD
Flathead Lk/Yellow Bay (15)	SK	1/4	VLD
Lake Mary Ronan (76.08)	SK, CT	4, 1/16	LDB, LDA
Logan (17.58)	SK, CT	2, 10	LDA, HD
Lone Pine (182.35)	SK, CT, SJ	80, 10, 40	LDA, LDA, VLD
Thompson Falls (36.4)	SK, LS	12, 1/16	LDB, MD
Whitefish Lake (10.35)	NONE		

¹ Weed species abbreviations

SK = spotted knapweed	LS = leafy spurge	CT = Canada thistle
MU = mullein	DT = dalmatian toadflax	SJ = St. Johnswort
SC = sulphur cinquefoil	OW = other weed-like vegetation	

² SIZE: total acres or partial acres for each species, respectively:

1 acre = 43,560 ft ² (200' x 200')	1/16 acre = 2,722 ft ² (50' x 50')
1/2 acre = 21,780 ft ²	1/32 acre = 1,361 ft ²
1/4 acre = 10,890 ft ² (100' x 100')	1/64 acre = 680 ft ² (25' x 25')
1/8 acre = 5,445 ft ²	< 1/64 acre estimation in square feet

³ DENSITY: number of weed stems in the population

VLD: very low density	--	one or a few <u>individual plants</u> , accounting for a very small proportion of the vegetative ground cover; native species well represented
LDA: low density - A	--	individual <u>plants clumped in small areas</u> , accounting for a small proportion of the vegetative ground cover; native species well represented
LDB: low density - B	--	individual <u>plants scattered throughout site</u> , accounting for a small proportion of the vegetative ground cover; native species well represented
MD: moderate density	--	individual <u>plants numerous</u> , accounting for a moderate proportion of the vegetative ground cover; native species present
HD: high density	--	individual <u>plants abundant</u> , accounting for a high proportion of the ground cover; native species present but not well represented
VHD: very high density	--	individual <u>plants very abundant</u> , accounting for almost all of the vegetative ground cover; native species almost absent

Wildlife Management Areas

FWP manages the Wildlife Management Areas with the needs of wildlife and the maintenance of habitat as primary objectives. Secondary objectives include providing recreational opportunities for the public. Each WMA is a unique portrait of landscape features, vegetation communities, and the environmental forces which shape them. With varying elevations, habitat types, precipitation zones, and management programs, the WMA's will be reviewed individually. These include: Kootenai/DeRosier-Woods Ranch, Kootenai Falls, West Kootenai, Mount Silcox, Ninepipe, Pablo, Ray Kuhns and Roundhorn. Part 3.0 of this Plan will highlight these differences and propose weed treatments which suit each property. The smaller WHPA's without an active management program include: Buffalo Head Park, Flathead Lake, Flathead River, Lower Stillwater Lake, Swan Lake, and Thompson Falls.

Fishing Access

FWP manages Fishing Access to provide public access to river corridors or lakes as well as to provide reasonable accommodations for vehicle parking, boat launching, and general recreation. As noted earlier, proximity to water and seed dispersal influence the degree and success of noxious weed establishment. Water courses disperse and transport seeds quite effectively. A small, localized weed population will be a seed source for downstream lands if seeds are introduced into a waterway. For that reason, all FAS's for a particular drainage will constitute a property group. For example, spotted knapweed is a problem common to most access sites on the Flathead River; therefore, management strategies should address those sites comprehensively if long-term reduction and/or eradication is to be successful.

In Region One, many FAS's access ponds or lakes. These will be treated under the river property group for the appropriate river drainage. However, fish hatcheries and administrative sites will be treated as a separate property group. The Region One Headquarters complex will be grouped and considered with fish hatcheries and administrative sites.

State Parks

State Parks attract visitors because of the natural, cultural, recreational, and historical resources. Picnicking and overnight camping opportunities also attract many visitors. Although there may be similar opportunities and constraints for the treatment and management of noxious weeds in some State Parks, each Park was established for its unique features. Thus groups are based on broad natural, historic, or recreation resource similarities. Five of the six units of Flathead Lake State Park, because they are all situated on the shores of Flathead Lake, will be considered together. Flathead Lake State Park-Wild Horse Island, Lake Mary Ronan, Logan, Lone Pine, Thompson Falls, and Whitefish Lake will each be discussed individually due to the uniqueness of the settings and the resources involved. Differences in facility development and human use will be discussed in greater detail in Part 3.0.

2.0 MANAGEMENT STRATEGIES

2.1 General Approach

It is the goal of FWP to be in compliance with state laws and county ordinances concerning noxious weeds where they are established. Furthermore, FWP hopes to discourage the establishment of noxious weeds through the application of good land management principles and to reduce noxious weed populations over time with as few negative impacts to the State's resources as possible. To achieve these goals, FWP crafted this Noxious Weed and Exotic Vegetation Management Plan based on the concept of Integrated Noxious Weed Management (INWM). INWM is a multifaceted approach which tailors weed management activities to the treatment needs and site characteristics of a particular weed population. First, weed management priorities are established. Then, management methods and techniques are selected, including natural predators and parasites, environmental modifications, and chemical herbicides when necessary and appropriate. INWM allows for the most flexibility in managing noxious weeds as site conditions change, treatment methods improve, and funding opportunities arise. This approach is consistent with state law and the noxious weed plans developed by the counties. FWP has been practicing some form of INWM informally since the early 1980's. This Plan formally adopts this approach.

2.2 Conceptual Framework to Prioritize Activities

Within the context of INWM, there are several ways to prioritize noxious weed management activities. These approaches include species-specific directives based in state law, treatment objectives based on location or size of weed populations, special management zones which require modification of general (area-wide) techniques, and property groups based on watershed or political boundaries. In using INWM, FWP has adopted an approach which utilizes all of these concepts.

2.2.1 Species Specific Directives

As a state law, the County Noxious Weed Control Act mandates the management of certain species declared as "noxious." Thus, FWP must attempt to treat and manage those species declared by the state as well as those species declared by the individual counties. On FWP sites, spotted knapweed and leafy spurge are the most prevalent, but other species are treated as well. Thus, efforts which have primarily been focused on these species will continue, but additional species may need to be added as the state and county noxious weed lists are amended.

Categories of Exotic Species in State Parks

Given the differing characteristics of exotic species and the different mission of state parks, it is helpful to rank species into categories for management action within state parks. Individual exotics are not all equal from biological as well as management perspectives. An Exotic Species Ranking System (Hiebert, 1986) was used to provide objectivity to the placement of species into categories. Emphasis is on significance of impact and feasibility of control. Each exotic species has been tentatively classified into one of five categories (see Appendix B). As new exotics are identified in the parks, they also will be classified.

Within this scheme, Category 1 species have the highest priority for management actions, while Category 5 species have the lowest priority. Control actions are justified for exotics in Categories 1 and 2. Occasionally, control actions may be justified for category 3 species, but are unlikely for Categories 4 and 5.

2.2.2 Location and Population Size

Noxious weed treatment priorities depend on the location and size of a weed population. Factors to consider are: the potential impact on site values in the absence of weed treatment, potential of weed spread, and the difficulty of satisfactory weed control. These same factors are considered by the state and counties during the process of listing weeds as "noxious." Priorities based on location and population size described below (Lolo National Forest 1989).

Prevention

Prevention describes the implementation of policies, programs, or activities that will minimize the possibility of introduction of noxious weeds or their seeds to FWP sites from lands nearby. This approach will have a high priority for FWP since the long term cost effectiveness is high. Attempts will be made to prevent the establishment of a new noxious weed species or new populations on Region One properties. Where newly discovered populations are small, the objective is immediate treatment to prevent any increase in size beyond manageable limits.

Preventative measures under consideration include but are not limited to the following:

- Regulate or limit the use of vehicles that could spread and carry noxious weed seeds into an area.
- Require or promote the use of weed seed free materials for livestock feed, landscaping, revegetation, and construction activities, including topsoil, gravel, hay mulch, and seed mixes.
- Require the thorough cleaning of all heavy equipment used on FWP lands, where feasible.

- Cooperate in education programming initiated by other organizations, where feasible, concerning the problems caused by noxious weeds, how weeds spread, and how the spread can be curtailed.

Eradication

Eradication is an attempt to totally eliminate a noxious weed species or a particular weed patch from FWP properties. Eradication is preferred where weed populations are small but have the potential to spread rapidly. However, this approach has limitations. Long term efforts may be required where the seed source is established; current technologies may not be adequate or practical; or total eradication efforts may result in unacceptable environmental consequences. When appropriate, eradication efforts would continue as long as detectable levels of the weed were present. For small populations, the cost effectiveness of eradication efforts is high.

Reduction or Suppression

Reduction or suppression is an attempt to prevent seed production and distribution throughout the treatment area and reduce the coverage of the weed. Reduction will prevent the weed from dominating the vegetation of the area but will require the public and land managers to accept noxious weed presence at low levels since complete elimination would not be feasible. Reduction is preferred in most instances when light to moderate noxious weed populations exist in accessible areas.

Containment

Containment is an attempt to prevent (through eradication or die off) the spread of the weed beyond the perimeter of a weed patch or treatment area. Managers and the public must tolerate weeds within established areas, but not beyond current boundaries. Containment is the preferred approach when weed populations have grown so large or dense that reduction is not practical or possible. In some situations, weeds may exist in settings where all methods of treatment are either unacceptable or ineffective so that containment is the only practical course of action.

Tolerance

Tolerance is to accept the continued presence of established weed populations and the probable spread to ecological limits for a particular species. Tolerance should only be applied in the context of containment. Tolerance is only acceptable if noxious weeds are contained within an area and not allowed to spread beyond the ecosystem. Like containment, tolerance situations usually exist where noxious weeds are beyond control or are in environments where all methods of treatment are either unacceptable or ineffective. The manager must identify these situations and try to discourage weed encroachment through preventive practices.

2.2.3 Special Management Zones

Although weed treatment needs may be similar from site to site or within the boundaries of one particular site, each site is a unique juxtaposition of environmental features, visitor facility development, and level of human use. For these reasons, individual sites or groups of sites may be distinguished into management zones based on the following: presence of surface water or shallow groundwater, threatened, endangered, or sensitive species, intensity of human use, weed treatment objectives, park management zones, or other factors. The framework of management zones permits the identification of weed treatment methods and techniques amenable to a particular weed patch should it require the modification of general area-wide weed treatments. Three zones have been identified: surface and groundwater, sensitive species, and high human use. Because of the sensitive nature of these special management zones, herbicide use will be somewhat limited. A discussion of each zone and guidelines for their protection follow.

Surface and Groundwater Zone

Sensitive surface and groundwater zones constitute locations of aquatic vegetation, surface water, shallow groundwater, and groundwater recharge. Examples include wetlands or marshes, riparian areas, lakes or ponds, springs, rivers, flood plain, and pathways of overland runoff. Because of their high moisture content and abundant nutrients, these environments are highly productive, yielding valuable fishery and wildlife habitats, recreation areas, and livestock foraging areas. Such zones occur on a significant portion of FWP properties in Region One. Noxious weeds exist within many sensitive surface and groundwater areas. The weed management goal is to attempt to protect riparian areas and wetlands from encroachment by noxious weeds while preventing chemical contamination of surface and groundwater if weeds are treated in these zones.

Surface water zones can be determined by studying topographic features of a site. After surveying the presence and location of surface water, high water marks, and associated overland runoff contours, a chemical applicator can discover and avoid avenues of potential surface water contamination. In addition, a preliminary soil analysis will determine soil textures. Course, gravelly soils tend to have higher infiltration rates, which imply the potential to transport chemicals. The opposite is true of fine textured, clayey soils.

Sensitive groundwater zones could be identified on a soils map using a procedure developed for the Lewis & Clark County Conservation District (Lacey 1991). Risk of chemical contamination of water resources is determined for a site based on depth to the seasonal or permanent water table, soil permeability, soil organic matter content, flooding hazard, and percent slope. A risk index based on these variables was calculated on a scale of 1 to 100 and divided into 4 categories or classes of risk. Risk factors and classes were then indicated on a map of soil units (Table 4).

The U.S. Department of Agriculture (USDA) Soil Conservation Service completed the Flathead County soil map in the 1960's using data from the 1940's and 1950's and is scheduled to update it beginning 1995. The Lake County soil map is unpublished but is available on Geographic Information System (GIS) computer maps from the SCS Polson office. The SCS is still in the process of mapping soils in the other two counties in Region One. Until detailed soil maps become available and some of the other variables are quantified, the following guidelines are suggested to protect surface and groundwater from chemical contamination:

- Target noxious weeds encroaching upon sensitive surface and groundwater environments (e.g. purple loosestrife) so weeds don't become well established.
- Avoid applying herbicides directly to standing or running water; maintain a 50 foot distance from surface waters when applying chemicals; when applying chemicals within 50 feet of a surface water, use a wick applicator; because manufacturer label restrictions are not always restrictive enough, more restrictive guidelines could be established as site conditions merit.
- Mix and load herbicides off-site as much as possible; mixing, if done on site, should be at least 500 feet from open water or shallow groundwater; water for tank mixing should be obtained off-site.
- Avoid applying picloram on the immediate banks of a water body, below the high water line, or in high groundwater risk class areas.
- Encourage manual, cultural, and biological methods; if chemical methods are used, design and schedule herbicide treatments to have the least impact on fish, wildlife, non-target vegetation, and water quality.
- Avoid using boom sprayers where noxious weed plants and seeds are mixed with desirable woody vegetation; biological, cultural, or manual methods which target individual plants (e.g. wick applicator) are preferred in these settings.
- Avoid broadscale applications with high herbicide rates on areas where fractured bedrock lies close to the soil surface.

Table 4. Soil index values, risk classes, and soil descriptions used to identify groundwater areas sensitive to chemical contamination (Lacey 1991).

Index	Class	Soils Descriptions
100+	1-Very High	Soils with water tables within 3 feet and moderately rapid to very rapid permeability; herbicide use not recommended on these sites.

80-99	2-High	Soils with water tables within 3-6 feet, moderate to rapid permeability, no flooding risk; <u>or</u> soils with water tables within 15 feet, moderately rapid to very rapid permeability, high flooding risk, and low organic matter; spot treatments with herbicides may be utilized in some areas; reduce herbicide rates and application frequency to decrease contamination risk; low leach potential and short residual herbicides recommended for any large scale applications.
60-79	3-Moderate	Soils with water tables within 6-10 feet and moderately slow to very slow permeability; <u>or</u> soils with water tables within 15 feet and rapid to very rapid permeability; these soils often occur in association with high water table soils; herbicides can be used for spot treatments on these sites; large broadcast applications with high herbicide rates not recommended; decrease application frequency to reduce risk.
40-59	4-Slight	Soils with water tables within 15-25 feet and moderate to moderately rapid permeability; soils occur where landscape features resemble those with high water table soils; may be located on benches and terraces adjacent to perennial or intermittent streams; few restrictions for herbicide use.

Sensitive Species Zone

Sensitive species zones possess physical and biological features essential to the conservation or survival of plants or animals. Typically, these zones support animal or plant populations which may be adversely affected by the presence of noxious weeds or efforts to treat them. More specifically, they might provide vital habitat for species designated threatened or endangered under the Endangered Species Act or as sensitive by the Montana Natural Heritage Program. Noxious weeds occur in many sensitive species zones. The weed management goal within this zone is to protect existing, diverse habitats from encroachment by weeds while ensuring that threatened, endangered, rare or unique species are not adversely affected by noxious weed treatments. The following guidelines are offered:

- Determine sensitive species zones or identify and locate sensitive species populations for each property or group of properties and establish treatment buffers if necessary.
- Direct management towards weed populations in the vicinity of sensitive species zones before weeds become established in the sensitive zone.
- Encourage manual treatments, although some cultural and spot herbicide treatments may be considered where appropriate.
- Design and schedule herbicide treatments to have the least impact on wildlife, fisheries, and non-target vegetation.
- Avoid using boom sprayers in sensitive species zones where weeds are closely mixed with desirable woody vegetation.
- Consider revegetation or rehabilitation of treated wildlife habitat areas as part of the treatment plan to ensure that desirable ground cover is established without affecting sensitive plant species.

High Human Use Zone

High human use zones are areas which have concentrated and frequent public or FWP employee use. Examples include picnic table areas, tent sites, restrooms, drinking water well sites, interpretive sites, and boat launch areas. Because of the high degree of ground disturbance and the intense activity level, these areas are very susceptible to the encroachment and rapid spread of noxious weeds. Once invaded, these areas require aggressive treatment actions. The weed management goal is to reduce noxious weed populations in areas which are continually or heavily used by people, while simultaneously ensuring public health and safety. In the majority of situations, a buffer zone of at least 50 feet is maintained from such zones during chemical application. Weed management guidelines for this zone follow:

- Emphasize prevention as the key component of management efforts, primarily by cooperating with educational programming initiated by other organizations to increase public awareness of noxious weeds.
- Encourage manual, cultural, or biological treatments; consider the plausibility of spot herbicide treatments on a site by site basis.
- Post signs in high human use zones prior to and for 24 hours after herbicide applications to notify the public that spraying activities will/have occurred.
- Wash and rinse residue from surfaces used by the public for food handling if herbicides are applied within 50 feet.
- Use only those chemicals with label designations specifying permission for use in human occupation areas (e.g. glyphosate, clopyralid, and 2,4-D).

2.3 Treatment Methods and Techniques

Within the framework of INWM, there are four general weed treatment methods, each with several specific techniques or agents. Manual, cultural, biological, and chemical methods for the treatment and management of noxious weeds are described below, with specific techniques or agents identified. Additional methods and techniques will be considered in the future as they become available.

2.3.1 Manual

Manual techniques involve grubbing with hand tools (shovel, hoe, pulaski etc...), topping (shears, clippers, and "weed-whackers"), and hand-pulling to treat noxious weeds. These techniques can effectively reduce small populations of annual or biennial noxious weed species, primarily by reducing seed production. Periodic re-treatment is required. Because these

techniques are labor intensive, their application would be mostly limited to environmentally or culturally sensitive areas.

2.3.2 Cultural

Cultural techniques include burning, mowing, tilling, reseeding, and grazing. These techniques effectively reduce some noxious weed species, although implementation is often limited by topography and access. These techniques can also be costly and time consuming. Furthermore, some techniques disturb the soil which can contribute to noxious weed proliferation if revegetation does not occur rapidly.

Burning

As a treatment technique, burning alone is generally of limited effectiveness. It may be considered in the future in conjunction with other treatment techniques such as reseeding.

Mowing and Tilling

Mowing and tilling can prevent noxious weed plants from producing seeds when treated in the bud stage or earlier. Repeated efforts during a single growing season can deplete the underground food supply of some perennial species. These treatments would need to be repeated over at least a five-year period to attain satisfactory results since they do not eliminate weed plants themselves. These methods may also weaken non-target species. Mowing and tilling opportunities are often limited because of steep slopes and rugged terrain. Mowing would be considered on level or nearly level slopes, where suitable soils exist and where a small percentage of the vegetation consists of shrubs. Mowing is also an alternative in environmentally sensitive areas where other treatment methods are unsuitable. Tilling would be considered only for areas that had previously been or are currently under cultivation because ground disturbance may increase the likelihood of noxious weed establishment beyond the potential benefits of tilling. However, tilling combined with a reseeding program would be considered when economically feasible. Other treatment methods would also be considered in conjunction with tilling such as alternate cropping systems, grazing systems, or chemical applications.

Reseeding

Reseeding is required where plant communities are completely disturbed by facility construction. It may also be advisable where vegetation communities are dominated entirely by noxious weeds. Such revegetation projects require considerable effort to eliminate existing weeds, prepare a seedbed, and re-plant. In radically disturbed areas, it is imperative to reseed as soon as possible. Topsoil should be salvaged and replaced on the disturbed site immediately to

encourage native species to sprout from seed, rhizomes, or other plant parts in the salvaged soil. Despite the significant expense, dramatic results can be achieved.

Grazing

Grazing by domestic sheep and goats have effectively controlled the spread of leafy spurge in certain situations in Flathead County and other parts of Montana, as well as in North Dakota. Other weed species were often selected early in the growing season since they tend to green up earlier than native plants. Controlled livestock grazing incorporating adequate vegetative rest is utilized on several of the WMA's to maintain and/or improve range condition thereby discouraging weed proliferation. Many leafy spurge populations on Region One properties occur where careful herding and/or installation of temporary fences would be necessary. When considering this treatment technique, the effect of grazing on soils, water, and riparian areas will be considered.

2.3.3 Biological

Biological control methods are those which introduce natural enemies (insect pathogens) to a noxious weed population. Biological control agents stress noxious weeds, severely reducing seed production and/or plant viability. Biological agents will not eliminate the species, but can reduce the population to a tolerable level, particularly when used in combination with other treatment techniques. Biological control methods may be the only treatment possible in areas where manual, cultural, or chemical methods are not feasible.

Biocontrol agents are currently being developed and screened by universities and the USDA Animal, Plant and Health Inspection Service (APHIS). There are also commercial sources for a number of agents. Several biological agents have been released in the counties of Region One. Release of biological control agents onto FWP properties will be in cooperation with the counties, the Montana Agricultural and Experiment Station at Montana State University, and APHIS. FWP personnel in Region One have been working with APHIS to establish insectaries on certain properties which meet selected criteria. Over the long term, biological control agents are cost effective because expenses are limited to the collection and release of new agents and periodic monitoring to insure spread and infection. However, it remains to be seen whether biocontrol agents can keep pace with the rate of spread of certain noxious weed species, whether on FWP properties or elsewhere.

With any biocontrol program, it is necessary to maintain a viable host population for the biological control organism. Areas selected for release of these agents will be screened with respect to location and plant community. As such, these areas preclude weed eradication objectives and the use of herbicides. Appendix D summarizes biological control agents currently available in Montana, and those which have been released on FWP properties in Region One.

2.3.4 Chemical

Chemical methods involve the use of herbicides to kill weeds directly or the application of soil sterilants which prevent the germination of noxious weed seeds. Over the past half-century, chemical herbicides have been developed to treat a variety of weed species. While the benefits of chemical techniques include efficacy and low cost, there are valid concerns about any application of chemicals to the landscape. In addition to concerns about the effects of chemical exposure on human health and the environment, attention has also focused on chemical storage, handling, use, and disposal procedures. Herbicide manufacturers have responded to these issues with attempts to develop products that are: (a) more pest specific to reduce effects on non-target species, (b) less toxic to humans, birds, fish and other animals, and (c) less likely to impact the environment, particularly water quality. The following discussion highlights these points in greater detail.

Efficacy and Cost

Properly used chemical herbicides are usually effective against their targets. Variation in effectiveness is attributable to environmental conditions, skill of the applicator, condition of the equipment, application rates, and the target species. For example, leafy spurge is difficult to eradicate with chemicals whereas spotted knapweed can be successfully eradicated, in the short-term. Herbicides are relatively inexpensive to obtain and apply.

Human Health Effects

A definitive evaluation of human health risks from exposure to herbicides is beyond the scope of this Plan. The long-term health effects of chemical herbicides continue to be researched. Short-term health effects can be serious for sensitive individuals. Health effects are most commonly reported among herbicide applicators. Due to uncertainty over health and environmental effects, FWP hopes to reduce the use of chemical herbicides over time.

Environmental Concerns

Environmental concerns over the storage, handling, use, and disposal of chemicals are based on four issues: persistence, potential for movement, contamination of surface and groundwater, and the potential for chemical spills.

Persistence refers to the longevity of a chemical's integrity. It is usually expressed as half-life or the length of time required for half of the substance to become inert. Herbicides are degraded by physical, chemical and biological processes. Many herbicides degrade to very low levels in soil after several weeks or months, whereas others persist for several months to years. Most herbicides are broken down and adsorbed in the uppermost 12 inches of the soil profile.

Potential for movement describes the transport of chemicals through the environment as solids, liquids, and gases. Once applied, herbicides are absorbed, adsorbed, accumulated, degraded, diluted, inactivated, and/or mobilized. Solubility in water gives a preliminary indication of chemical mobility. The ultimate fate is heavily influenced by the type of herbicide, the application rate, environmental conditions at the time of application, and site characteristics. Individual properties or characteristics of herbicides also affect persistence and potential for movement due to the interaction of active ingredients with soil organic matter and clay particles. Research has been conducted on most common herbicides to establish environmental pathways and final fate. Research findings are incorporated into herbicide label instructions distributed with chemicals at purchase. These instructions are an important component of herbicide use and safety and will be discussed later.

Contamination of surface and groundwater results from improper use and application of chemical herbicides. Sandy and gravelly soils, having high infiltration and hydraulic conductivity rates, may transport chemicals directly to surface or groundwater. Transport may also occur where fractured bedrock lies close to the surface. Drainage paths, whether natural or artificially created, could also transport chemicals. These mechanisms of transport involve relatively low chemical concentrations, though some level of contamination may still result if appropriate applications guidelines are not followed.

In contrast, the potential for high chemical concentration levels exists in the event of a spill, with the extent of contamination depending of the nature of the spill. Many spills occur during transport, resulting from auto accidents. However, spills may also occur during application or storage. For these reasons, this Plan also outlines emergency response procedures in the event of an accidental spill. The Herbicide Emergency Response Plan is in Appendix E.

Because of the aforementioned concerns, FWP intends to minimize the reliance on chemical methods as much as possible, while still meeting its legal responsibility to manage and treat noxious weeds on its properties. However, because the proliferation of noxious weeds does not appear to be decelerating, chemical methods remain a necessary component of this Plan. Therefore, it will be under the following guidelines that FWP applies chemical herbicides to its properties:

- Consider the treatment objective, season of application, stage of plant growth, topography, species present, expected cost, equipment limitations, and potential environmental impacts during site specific planning.
- Determine rates of application from label instructions, taking into account target species, other vegetation in the area including desirable vegetation species, soil type, human occupation, wildlife, and the presence of surface waters or shallow groundwater.
- Conduct and supervise all applications with an applicator licensed by the State of Montana, whether that individual is a Department employee or contractor.

- Use vehicle-mounted spray applicators (hand-gun, boom, injector etc...) in open areas that are readily accessible and where vehicle operation will not damage soil or vegetation; the boom will only be used for concentrated weed patches.
- Use hand applicators (backpacks or wipers) on small, scattered patches in rough terrain or environmentally sensitive areas.
- Dismiss the possibility of aerial herbicide applications for the present time; reconsider in the future as warranted.
- Monitor weather conditions during application periods to ensure effectiveness of the application and to reduce drift; spraying would cease when rainfall is expected within 24 hours, air temperatures exceed 85° F, or wind speeds exceed 8 mph.
- Design and adjust spray equipment to deliver a median drop diameter of 200-800 microns to reduce excessive drift.
- Conduct calibration checks at the beginning of the spraying season and periodically to ensure that equipment is functioning properly and that the desired applications rates are achieved, when feasible.
- Attempt to apply herbicides during the optimum stage of plant growth to achieve maximum effect using the least amount of herbicide; this is usually in the spring and/or fall when adequate soil moisture is available for plant growth and herbicide uptake.
- Mix a dye, when feasible, with the herbicide solution to aid in identifying where spraying has occurred in order to prevent "double-spraying" and to provide a means of checking coverage for monitoring purposes.
- Avoid applying picloram to dense stands of noxious weeds in sequential years to prevent overuse of a persistent chemical capable of leaching into groundwater.
- Follow all herbicide label instructions, observing restrictions; additional precautions may also be taken.

2.4 Inventory and Monitoring Procedures

Prior to the preparation of this Plan, inventories were conducted during summer, 1994 and during previous years. Inventories consisted of field reconnaissance by FWP personnel or FWP contractors who noted the weed species present, location, approximate acreage, and its density. See Tables 1, 2, and 3 for a summary. Based on these reports, weed treatment activities are prioritized using the conceptual framework presented previously. Subsequently, work plans for

the following field season are outlined. Monitoring on a yearly basis will be conducted through similar informal field reconnaissance methods. Priorities and treatment methods can be adjusted as necessary, due to the dynamic nature of noxious weed seed germination and plant growth.

2.5 Other Subjects of Concern

2.5.1 Oversight of Plan Implementation

The ultimate responsibility for implementation of this Plan and compliance with its provisions rests with the Regional Supervisor. However, the most direct oversight of field personnel conducting weed treatments is the responsibility of the Regional Managers of the Division lands under their jurisdiction, respectively. For example, the Regional Wildlife Manager would oversee weed management activities on Wildlife Management Areas, the Regional Fisheries Manager would oversee weed management activities on Fishing Access, and the Regional Parks Manager would oversee weed management activities on State Parks.

2.5.2 Threatened, Endangered, and Sensitive Species

One goal of this Plan is the preservation of fisheries, wildlife, and vegetation resources under the stewardship of FWP. Land management agencies and conservation organizations have developed lists of threatened, endangered, or sensitive plant and animal species. Those species listed in Montana are of concern due to relatively small population sizes and/or limited distributions. Appendix F summarizes reported occurrences of such species on or near FWP properties in Region One (Montana Natural Heritage Program 1993). Properties in this Region include aquatic, moist forest types, dry forest types, and shrub grasslands -- some with similar environmental and site conditions as habitats supporting listed species. FWP personnel involved in weed management activities should be aware of this list. In addition, they should periodically consult with the FWP Plant Ecologist and the Endangered Species Program Coordinator.

2.5.3 Applicator Licenses and Record Keeping

Applicators of restricted use herbicides must be licensed by the State of Montana. To obtain a license, the applicator must take training courses and pass an exam. To retain that license, the applicator must earn re-certification credits in government-approved courses. State law also requires that applicators record their use of restricted herbicides. Records must indicate the location's legal description, the chemical applied, rate of mixing, the applicators, the weather conditions at the time of application, and the number of acres treated. These records must be submitted to the State Department of Agriculture one out of every five years. In addition, the records must be made available to state investigative officers at any time following an herbicide application. All aspects of applicator licensing, re-certification training, and record keeping

regulations administered by the State are supervised by the United States Environmental Protection Agency. Appendix G outlines herbicides used in Region One.

In addition to the State reporting requirements, FWP policy also requires the annual reporting of all measurable weed treatment activities conducted on any Department property, regardless of the methods and techniques employed. All cultural, manual, biological, and chemical treatments are recorded according to county and site name, type of action, species treated, number of acres treated, application rate, frequency of treatment, and total cost. Periodically, FWP produces summary reports of all noxious weed management activities statewide. Region One reports are filed at the Regional Headquarters Office in conjunction with yearly workplans.

2.5.4 Herbicide Labels

Herbicide labels are an important component of chemical use and safety. Manufacturer labels are in booklet format and supplied with each container of product. The booklet contains detailed information to meet four important goals of herbicide regulation, including: chemical identification, protection of human health and the environment, special practices, and legal requirements. Herbicide labels have the force of law and should be an integral part of any chemical weed management program. Appendix H contains manufacturer labels for those herbicides proposed for use in Region One.

2.5.5 Contracted Services

Some weed management activities in Region One, especially herbicide applications, have and will continue to be contracted to private individuals. Contractors must be licensed by the State and knowledgeable concerning the specific weeds to be treated and chemicals to be used. Contractors must also sign and comply with a Personal Services Contract, in addition to carrying a liability insurance policy of \$750,000. Contractors should be familiar with all components of this Plan, especially those regarding safety and emergency response. Contractors must be provided with clear information and directions and be supervised to insure compliance with contract specifications and this Plan.

Contractors should keep detailed records concerning applications, including: time and date of applications, chemicals used, amounts and mixing notes, public contacts, weather, and other pertinent information. This record-keeping can be coordinated with other herbicide use and reporting programs.

Contractors utilizing non-chemical methods of weed treatment would also be provided with the necessary information and direction to meet the specifications of the contract and this Plan. Their activities will also be recorded and filed.

3.0 PROPERTY GROUP TREATMENT PLANS

In most cases, existing information is sufficient to define the conditions of FWP Region One properties with respect to noxious weeds. Property group treatment plans were prepared based on these data. Plan preparation was also guided by the following factors, each of which stems from the conceptual framework to prioritize weed management activities discussed in Part 2.0 of this Plan:

- weed species present
- size and density of weed populations
- management objectives
- potential for successful treatment
- potential for spread on FWP property
- potential for spread beyond FWP property
- public concern over weeds or weed management efforts
- existing local weed management projects
- budget and personnel constraints

3.1 Wildlife Management Areas and Affiliated Lands

3.1.1 Introduction

Wildlife Management Areas (WMAs) are managed to meet the needs of wildlife, preserve wildlife habitat, and provide recreational opportunities to hunt, fish, or view wildlife in their natural habitats. Site specific, integrated resource management plans are in various stages of development and/or revision and will be filed at the Regional Headquarters Office as they are finalized. These management plans describe overall management direction and specify certain activities to be undertaken to reach some desired future condition. Common to every WMA management plan is the application of proper land management practices to maintain or enhance healthy, vigorous vegetation communities. To the degree that vegetation communities are healthy and vigorous, they will be able to resist noxious weed encroachment. Furthermore, the quality of wildlife habitat is directly linked to the health and stability of vegetation communities. Thus, the prevention of noxious weed encroachment is inherent in the administration of these properties.

On WMAs, good land management begins with protecting the soil and vegetation resources. All WMAs have vehicle restrictions intended to prevent excessive soil disturbance and/or erosion. Motorized vehicles are restricted to established roads or trails and designated parking areas. Vegetation and soil resources are also protected by requiring the use of certified weed seed free hay to prevent the introduction of noxious weed seeds to WMAs.

WMAs are a small part of a larger landscape in which, unfortunately, noxious weeds occur, and noxious weeds have been documented on the WMAs in Region One. FWP is required by law to attempt to manage noxious weeds on its properties. As such, any noxious weed management should be approached on the landscape level, considering neighboring landowners, land use practices, and mechanisms of noxious weed seed dispersal. To that end, noxious weed management on the WMAs will center around maintaining healthy vegetation communities, deterring seed dispersal, and the detection and treatment of noxious weed outbreaks before they become well established.

Quick detection and treatment of small patches is the most efficient approach to managing noxious weed outbreaks. As populations increase, the number of potential treatment methods dwindles as other considerations come into play. For example, chemicals could be applied to a 1/10 acre patch with some risk to other vegetation, but they could not be applied broadscale to a 10 acre patch without potential impacts on non-target vegetation or other resource values. In addition, seed production influences the long term success of weed treatments. The volume of seeds produced on a 1/10 acre patch is much less than that of a 10 acre patch, thus making it easier to confine or eradicate the smaller one.

Budget and personnel constraints necessitate the implementation (when necessary) of treatment methods which maximize the ratio of expedient, successful noxious weed treatment to cost. Of the methods currently available, chemical treatments offer the most cost effective means of weed management. However, there are circumstances where chemical methods are not feasible, such as near surface waters or where weeds cover a wide area. Manual methods are not practical in most cases because of budget and personnel limitations. Cultural methods, though effective in certain situations, sometimes result in additional soil and vegetation disturbances, two factors contributing to the spread of noxious weeds. Biocontrol methods offer a cost effective alternative though they are not appropriate in all settings. Furthermore, biocontrol agents are not yet readily available. Only a few sources dispense agents commercially, at a premium price; otherwise, agents must be obtained from APHIS.

Each WMA is discussed individually with respect to the status of noxious weeds and management strategies. Recall that Table 1 summarizes the noxious weed species present and status for WMAs in Region One. It should be pointed out that the estimated area (in acres) presented in Table 1 is cumulative for an entire site, since weeds are not continuously distributed throughout a WMA. That is, estimated acres of weeds along roadways are added to estimated acres in remote locations to obtain a total amount of acres. Figure 2 illustrates the location of the WMAs in Region One.

Figure 2. Wildlife Management Areas (WMAs) administered by FWP in Region One.

3.1.2 Kootenai/DeRoziert-Woods Ranch - Weed control on this property is currently being accomplished on a site-specific basis. There is no weed management plan for this property at this time.

3.1.3 Kootenai Falls

Existing Environment

Location: The Kootenai Falls WMA consists of 172 acres situated as a narrow, linear strip of land running east and west through the middle of Sections 15, 16 and 17, T31N R32W. Legal description is as follows: N S Sec. 15, N S Sec. 16, S N Sec. 17. T31N R32W. The WMA occupies approximately three miles of north shore Kootenai River frontage six miles downstream from the city of Libby.

Land use: The Kootenai Falls WMA was purchased by FWP for bighorn sheep and other big game grazing purposes. Public recreational use of the area is permitted and consists of hunting, fishing, wildlife viewing, walking, hiking and mountain biking. No overnight camping is allowed. No unauthorized motor vehicle traffic is permitted.

Fish and Wildlife: The Kootenai River, which forms the southern boundary of the WMA for three miles, is a designated Blue Ribbon trout stream in Montana. No fish supporting waters exist on the WMA.

The WMA is critical bighorn sheep range. Other big game species that utilize the property extensively include the white-tailed deer, mule deer and black bear. Moose and elk are occasional visitors to the area as are mountain lions. Other species of interest include ruffed grouse and a variety of waterfowl that nest along the Kootenai River shoreline. A host of neotropical migrant bird species utilize the riparian, as well as, upland sites on the property. Osprey and bald eagles are conspicuous raptors using shoreline trees for perching and roosting. No nests for either species exist on the WMA. Aquatic furbearers on or adjacent to the property include beaver, muskrat, mink and otter. Other medium sized mammals inhabiting the property are coyotes, bobcats and badgers.

Vegetation: Coniferous forest types common to the Pacific Northwest occur on the WMA. Douglas fir/ponderosa pine dominate the overstory. Western red cedar and grand fir occur on isolated sites associated with seeps or moist shaded areas. Common understory shrubs include ninebark, snowberry, oceanspray, white spirea, serviceberry, mock orange, Oregon grape, chokecherry and rose. The common habitat type is designated Douglas Fir/Ninebark. Cottonwood, birch and other deciduous species occupy riparian zones and moist microsites. Most flat tillable land has been converted to hay fields where alfalfa/grass hay is grown.

Water Resources: Three ephemeral streams, Hunter Gulch, Dad Creek and Burrell Creek dissect the property. The Kootenai River forms the entire southern boundary of the WMA. A few wet seep areas occur on the property.

Soils: Soils are not well developed on the area. Alluvial deposits from the Kootenai River mix with colluvial deposits from up-slope to form sandy-loam and silty-loam soils over most of the WMA.

Sensitive Species: Bald eagles use cottonwood snags and live trees along the Kootenai River for perching and roosting. No bald eagle nest sites occur on the property. Grizzly bears may occasionally use the area as a travel corridor.

Weed Status

Weed Infestation: Spotted knapweed infestations, primarily associated with disturbed sites, are delineated on a weed map.

Site Constraints

Aquatic chemicals have to be used for stream bank weed control along the Kootenai River. Use continues of target-specific chemicals to kill knapweed. Herbicide application should occur at opportune plant phenology in May or June.

Weed Management Strategies and Methods

Aside from disturbed sites, approximately 80 percent of the WMA remain unaffected by spotted knapweed. Past control measures have helped prevent further spread of this noxious weed onto other portions of the WMA. Prohibition of unauthorized motor vehicle use on the area will continue to prevent spread of seeds and soil disturbance. However, heavy recreational traffic via foot, bicycle and horseback will continue to disperse seed from the existing primitive roadway traversing the length of the property until the seed source is removed. Spotted knapweed encroachment onto the small meadows and hay fields will continue to reduce forage production for big game ungulates, primarily bighorn sheep.

Prevention: Continue prohibition of motor vehicle use on the area.

Physical: Possible application of hand pulling along Kootenai River banks.

Chemical: The proposed action is to treat the three miles of roadway (Management Area 1) with an application of Stinger during spring or early summer, monitor results and reapply Stinger in spot application as needed.

Meadows and portions of hay fields (Management Area 2) would also be treated with Stinger in spring/summer with retreatment the following year on a spot application basis.

Kootenai River banks (Management Area 3) could be treated with Aquatic Label 2,4-D during spring/summer and in subsequent years due to the short-term effect of 2,4-D on knapweed. As an alternative, labor and cost intensive hand pulling of spotted knapweed could be applied along the river bank each year.

3.1.4 West Kootenai - Weed control on this property is currently being accomplished on a site-specific basis. There is no weed management plan for this property at this time.

3.1.5 Mount Silcox

Existing Environment

Location: Mount Silcox WMA is located about 1 miles east of Thompson Falls, Montana. The area is bordered by private property to the east and west, State Highway 200 to the south, and the Lolo National Forest to the north. Total of 1,552 acres.

Land Use: The area is used primarily for winter and spring wildlife habitat. Cattle grazing is allowed from May to September. The WMA is used for recreational purposes such as hunting, hiking, and horseback riding. The area is accessible to the general public by horse, mountain bike, or walk-in. Roads are used for administrative purposes only.

Fish and Wildlife: The forested areas are used by elk, mule deer, whitetail deer, black bear, and mountain lion during part or all of the year. The grassland portion of the WMA is used by whitetail deer, elk, and bighorn sheep during the winter and spring seasons. Turkeys and mountain grouse are found occasionally on the WMA.

Cutthroat trout are found in Ashley Creek in the northwestern portion of the WMA.

Vegetation: The southern area contains a mixed species of pasture grass. The foothills to the north are mostly a mixture of Douglas fir and Ponderosa pine with numerous shrub species for an understory. Numerous rock outcrops are also present.

Water Resources: Ashley Creek flows through the western end of the WMA eventually going subsurface on adjacent private property. The Clark Fork River about 1 mile south represents the only other surface water in the area.

Soils: Soil characteristics are mostly sandy with some areas of gravel. Slopes vary from 0 to 60 percent.

Sensitive Species: Bald Eagles use cottonwood snags and live trees along the Clark Fork River for nesting and wintering. The rare Twin Clover and Rhombic-petaled Clarkia have been found in the vicinity.

Weed Status

Weed Infestation: Noxious weeds are predominately spotted knapweed with sporadic infestations of St. Johnswort. Several hundred acres of spotted knapweed have already been chemically treated to date.

Site Constraints

Herbicide application should occur at opportune plant phenology in May or June.

Weed Management Strategies and Methods

The flat grassland area will be controlled to reduce the spread of noxious weeds through chemical application of herbicide. The foothills and forested area will have limited chemical application of herbicide. Control will be attempted through various biological methods. Present weed infestations will be closely evaluated to see if they can be reduced. Herbicide application will be on a 2-5 year rotation.

Biological: Infestations of spotted knapweed will be possibly controlled with several insect species. Two seed-head flies Urophora affinis and U. quadrifasciata currently exist on the WMA. Other species of insect which could be introduced are a moth Metzneria paucipunctella, a root moth Agapeta zoezana, and a root weevil Cyphocleonus achates.

Chemical: Spotted knapweed and St. Johnswort can be controlled with an application of 1-2 pints per acre of picloram (Tordon 22K).

Priority: Area will be inspected annually to monitor noxious weed infestations and possibly initiate management efforts before weeds become well established. Spotted knapweed will receive top priority because of its ability to spread rapidly.

3.1.6 Ninepipe - Weed control on this property is currently being accomplished on a site-specific basis. There is no weed management plan for this property at this time. Efforts have concentrated on chemical and mechanical control of spotted knapweed, whitetop and purple loosestrife, if and when found.

3.1.7 Pablo - Weed control on this property is currently being accomplished on a site-specific basis. There is no weed management plan for this property at this time. Efforts have concentrated on chemical and mechanical control of spotted knapweed, whitetop and purple loosestrife, if and when found.

3.1.8 Ray Kuhns - Weed control on this property is currently being accomplished on a site-specific basis. There is no weed management plan for this property at this time.

3.1.9 Roundhorn

Existing Environment

Location: Roundhorn WMA is located about 15 miles east of Thompson Falls, Montana. The area is bordered by private property to the east and west, State Highway 200 to the south, and the Lolo National Forest to the north. Total of 27 acres.

Land Use: The area is used primarily for winter and spring bighorn sheep habitat. Light livestock grazing is permitted. Most recreational activity is done by walking to gain access to the Lolo National Forest. No roads are currently on the WMA. Vehicle access is allowed for administrative use only.

Fish and Wildlife: The area is used by bighorn sheep from November through April.

Vegetation: The area consists of flat pasture land of various grass species and forested areas dominated by Douglas fir and Ponderosa pine. Numerous rock outcrops and scree slopes are also present.

Water Resources: No surface water is located on the WMA. The Clark Fork River, about half-mile south, represents the only surface water in the area.

Soils: Soils are mostly sandy and rocky. Slopes vary from 0 to 80 percent.

Sensitive Species: Bald Eagles use cottonwood snags and live trees along the Clark Fork River for nesting and wintering.

Weed Status

Weed Infestation: Noxious weeds are predominately spotted knapweed which is sporadic throughout the WMA.

Site Constraints

Herbicide application should occur in May or June to correspond with opportune plant phenology.

Weed Management Strategies and Methods

Weed Management Strategies and Methods

Noxious weeds in the pasture area have been controlled with chemical application of herbicide to reduce their spread. Herbicide application has been on a 2-5 year rotation. Weed infestations will be monitored to see if reduction is possible. The foothills and forested areas are not planned for herbicide application, but will be controlled through natural and biological methods.

Biological: Infestations of spotted knapweed will be possibly controlled with two existing seed-head flies: Urophora affinis and U. quadrifasciata. No introduction of currently known controlling insects are planned for the WMA.

Chemical: Spotted knapweed can be controlled with an application of 1-2 pints per acre of picloram (Tordon 22K).

Priority: Area will be inspected annually to monitor noxious weed infestations and possibly initiate management efforts before weeds become well established. Spotted knapweed will receive top priority because of its ability to spread rapidly.

3.1.10 Buffalo Head Park, Flathead Lake, Flathead River, Lower Stillwater Lake, Swan Lake, Thompson Falls - There are no active management programs for these Wildlife Habitat Protection Areas at this time.

3.2 Fishing Access, Affiliated Lands and Administrative Sites

3.2.1 Introduction

Approximately twenty (20) percent of the statewide total hours of fishing per year (angling pressure) is spent in FWP Region One. Region One contains 2,700 miles of fishing streams. Region One also contains 600 lakes of which 450 have fish populations. These lakes sustain about one-fourth of the statewide lake angling pressure. Public enjoyment of such resources would not be possible without access.

Fishing Access properties are owned by or leased to FWP along surface watercourses or lakes. Through this ownership, FASes provide access to river corridors for the general public to fish, float, watch wildlife, picnic, hunt, camp, etc. Wetlands and watercourses with their associated vegetation communities (riparian zones) are some of the most diverse, productive habitats available to wildlife. For example, riparian woodlands in the west support the greatest diversity and abundance of neotropical migrant songbirds both during the breeding season and during migration (Dobkin 1992). These species are particularly dependent on the complexity and density of vegetation structure, especially in the shrub and herbaceous layers. Furthermore, riparian zones add tremendous diversity to the surrounding landscape. In addition to their significant value to wildlife, FASes are also valued by outdoor recreationists. Public use occurs

during all seasons, although it is more concentrated during the angling season. The setting and visitor use facilities at a typical FAS include the following: access road, parking area, surface water body (stream or lake), toilet, and a boat launch. Additional facilities may include picnic tables, cooking grills, and camp sites.

It is the goal of FWP to administer FASes by enacting good land management policies which protect vegetative and soil sources. To the degree that vegetation communities are healthy and vigorous, they will resist noxious weed encroachment. However, FASes are a small part of a larger landscape in which, unfortunately, noxious weeds occur, and noxious weeds have been documented on FASes in Region One. Furthermore, lake frontage and river corridors are increasing becoming subdivided into "ranchettes." This contributes to noxious weed problems for everyone if those landowners are not sensitive to the issue and make no attempt to manage weeds on their properties. Therefore, any noxious weed management should be approached on the landscape level, considering neighboring landowners, land use practices, and mechanisms of noxious weed seed dispersal. As well, noxious weed management strategies must take into account the natural resource and human use values of these sites.

To that end, noxious weed management on FASes will center around maintaining healthy vegetation communities, deterring seed dispersal, and the detection and treatment of noxious weed outbreaks before they become well established. Healthy vegetation communities can be promoted by limiting excessive soil disturbance which facilitates weed establishment. Soil disturbance can be prevented by confining the use of motorized vehicles to established roads and trails and by not disturbing vegetation unnecessarily during the development of visitor use facilities. After the development of visitor use facilities, the site will be reseeded. By restricting vehicular traffic to established routes, seed dispersal is also confined to areas where weeds can be easily detected and treated. Quick detection and treatment of small patches is the most efficient approach to manage noxious weed outbreaks. As populations increase, potential treatment methods and techniques dwindle as other considerations come into play. Given the number of FASes in Region One, budget and personnel constraints require the implementation of methods which maximize the ratio of successful weed management to cost. Therefore, the management of noxious weeds on FASes will be accomplished using the concept of Integrated Noxious Weed Management (INWM). Manual, cultural, chemical, and biological methods will all be used. Specific techniques will vary as individual circumstances require.

FAS treatment plans, organized by major river drainage, follow. All access sites in that particular drainage will be discussed in common. Unique circumstances will be pointed out as necessary. Access sites are located on the following rivers: Clark Fork, Flathead, and Kootenai. Ponds and Lakes with access sites will be discussed under the drainage they occur in. Fish hatcheries and administrative sites will be discussed as a separate category. Figure 3 illustrates the locations of FASes in Region One.

Figure 3. Fishing Access Sites administered by FWP in Region One. Key to the Fishing Access Sites appears on the facing page.

Key to the Fishing Access Sites

CLARK FORK RIVER DRAINAGE

8. Flatiron Ridge	30. Thompson Chain of Lakes

FLATHEAD RIVER DRAINAGE

1. Ashley Creek	7. Ducharme	19. Old Steel Bridge	26. Spring Creek
2. Ashley Lake	11. Horseshoe Lake/Ferndale	20. Pressentine	27. Swan River
3. Beaver Lake	12. Kokanee Bend	21. Skyles Lake	28. Teakettle
4. Bigfork	13. Kuhns	22. Smith Lake	31. Walstad
5. Blanchard Lake	15. Loon Lake/Ferndale	23. Somers	32. Whitefish River
6. Bootjack Lake	17. Marlowe Springs	25. Sportsmans Bridge	33. Woods Bay

KOOTENAI RIVER DRAINAGE

9. Frank Lake	14. Loon Lake/Eureka	18. Moran Lake	29. Tetrault Lake
10. Glen Lake	16. Marl Lake	24. Sophie Lake	

3.2.2

Clark Fork River Drainage

Existing Environment

FWP maintains one FAS on the Clark Fork River and one 4,200 acre FAS that encompasses 21 lakes at the upper end of the Thompson River.

Clark Fork Flatiron Ridge

Lakes Thompson Chain of Lakes

The Clark Fork enters Region One at the confluence with the Flathead River and flows 94 miles (including Noxon Rapids and Cabinet Gorge Reservoirs) to the Idaho state line. The Clark Fork has the largest annual discharge of any river in Montana. The river flows through a steep, deep, rocky canyon. The river corridor was impacted geologically by the repeated creation and flushing of glacial Lake Missoula so there are deposits of mixed sand, gravel, and rock, and soils are generally thin. Vegetation is primarily Douglas-fir and Ponderosa pine with a grass/brush understory. Cottonwood trees are also common.

Fish populations are generally low density due to warm water temperatures and poor water quality. Fishing and floating are relatively low (1,212 days of fishing in 1991). Noxon rapids and Cabinet Gorge reservoirs have been the subject of intensive fish enhancement programs by FWP and Washington Water Power the past 12 years. The bass fishery is now rated as one of the best in the state and reservoirs sustained 4,445 days of fishing in 1991.

The Thompson River is one of the major tributaries to the lower Clark Fork. The river originates in a complex of 21 lakes known as the Thompson Chain of Lakes. Champion International Corporation donated 4,200 acres of its landholdings in the Chain of Lakes to the state for recreation access. The area consists of rolling terrain with glacially mixed soils. The area also contains numerous streams and wetlands. The lakes range from 3 to 1,328 acres in surface area and contain a mix of cold water and warm water fish species including rainbow, westslope cutthroat, lake, brown, and brook trout, kokanee, largemouth and smallmouth bass, northern pike, pumpkinseeds, yellow perch, mountain whitefish, squawfish, and suckers. The lakes produced about 15,000 days of fishing in 1991.

The area has been used for logging and for cattle grazing. Some of the lakes have residential development. There are numerous roads and undeveloped campsites scattered throughout the area.

Weed Status

The most common noxious weed species on these FASes are spotted knapweed and Canada thistle, although other species do occur in small numbers (Table 2). The largest populations occur on Thompson Chain of Lakes and Flatiron Ridge FAS. The proximity of Clark Fork

River FASes to well-travelled highways serving as a primary transportation routes increases the potential for noxious weed introduction to these FASes.

Site Constraints

Chemical treatment of noxious weeds at these FASes is constrained by the presence of surface waters and by bedrock or gravelly soil types which easily transport chemicals to shallow groundwater tables or to surface waters. Furthermore, high levels of human use, particularly during the summer months, limit the use of chemicals even in those areas where they can be safely applied without risk to water resources. Riparian vegetation consisting of cottonwood/willow bottoms with a shrub and herbaceous understory component may also preclude certain chemical treatments. The wetlands habitat type at Thompson Chain of Lakes FAS would require special consideration when applying chemicals so as to avoid impacting non-target vegetation species.

Weed Management Strategies and Methods

Manual and cultural methods have been applied on roughly half of the FASes. Hand-pulling and cutting treatments were usually administered at least once during any one growing season. Many of these access sites receive annual treatments of this nature. Target weed species are spotted knapweed and thistle.

Some weed management on Clark Fork River FASes has been by chemical means, although FWP has begun to move away from a total reliance on chemical treatment methods on its FASes. Chemical methods will be reserved for those sites (or weed patches on a site) where application will not impair site values and resources. Manual (hand pulling), cultural (cutting or mowing), and biological methods will be emphasized.

In addition to treating established weeds, FWP will attempt to curtail noxious weed spread by restricting vehicles to established roads through traffic control devices such as rock barriers. Disturbance to soil and vegetation communities during facility development and maintenance will be minimized. Sites disturbed during development will be reseeded.

3.2.3 Flathead River Drainage

Existing Environment

FWP maintains 24 fishing access sites within the Flathead River drainage in Region One. They are:

Flathead River

Kokanee Bend

Kuhns

Old Steel Bridge/Kiwanis Lane

Pressentine

Sportsmans Bridge

Teakettle

Flathead Lake

Bigfork

Ducharme

Somers

Walstad

Woods Bay

Rivers

Ashley Creek

Marlowe Springs

Spring Creek

Swan River

Whitefish River

Lakes

Ashley Lake

Beaver Lake

Blanchard Lake

Bootjack Lake

Horseshoe Lake/Ferndale

Loon Lake/Ferndale

Skyles Lake

Smith Lake

The Flathead River and its forks flow from the Great Bear and Bob Marshall Wilderness; Glacier National Park and the province of British Columbia. The 295 miles of river provided 20,578 days of fishing in 1991. The forks of the river are bounded mostly by forested federal land. The mainstem river flows through a broad glaciated valley. The primary land use is agricultural, although much of the land along the river is being converted to rural homesites. The river channel shows the effects of several large floods with large cut banks and large bedload deposits. The river channel is braided downstream from Pressentine Bar. The river is greatly affected by hydropower dams with varying flows above Kalispell. The lower 20 miles of river is affected by Kerr Dam with reduced velocity and varying water levels. Some parts of the riparian zone have mixed cottonwood/douglas fir/spruce overstory while other parts are cleared and converted to cropland. Old oxbows provide wetland and warmwater fish habitat.

Angling is popular on the river with migratory runs of westslope cutthroat from Flathead Lake providing spring and summer fishing and lake whitefish providing fishing near Kalispell during the fall spawning run. The Flathead drainage is popular with floaters. The forks provide whitewater rafting while the mainstem provides uncongested floating with few hazards.

Waterfowl hunting and bird watching are also popular. Bald eagles, osprey, great blue herons and a number of other birds are common along the river corridor.

Flathead Lake is a 125,000 acre glaciated trough lake. It averages 100 feet deep with a maximum depth of 386'. The lake fluctuates 10 feet annually as regulated by Kerr Dam. Most of the shoreline is rocky with thin soils. Most of the lakeshore is in private ownership. There are only about 10 public access points on the entire lake, but many private commercial and homeowner association marinas. Flathead Lake provides about 73,000 days of fishing annually for lake trout, lake whitefish, yellow perch and westslope cutthroat. The lake is also popular for swimming, sailing and power boating. About two-thirds of the boat use is for angling.

Weed Status

Spotted knapweed is the most frequent noxious weed species encountered on FASes along the Flathead River (Table 2). The largest populations occur on the Pressentine and Kokanee Bend sites. The proximity of the Flathead FASes to the well-travelled U.S. Highways 2 and 93 increases the potential for noxious weed introduction to the FASes in this drainage. Not only are Highways 2 and 93 primary routes to Glacier National Park, they also serve as a shortcut between Canada and Idaho, and Commercial traffic along these highways is heavy.

Site Constraints

Chemical treatment of noxious weeds on Flathead River FASes is constrained by the presence of surface waters and by gravelly soil types which easily transport chemicals to shallow groundwater tables or to surface waters. Furthermore, high levels of human use, particularly during the summer months, limit the use of chemicals even in those areas where they can be safely applied without risk to water resources. Riparian vegetation consisting of cottonwood/willow bottoms with a shrub and herbaceous understory component may also preclude certain chemical treatments.

Weed Management Strategies and Methods

Manual and cultural methods have been applied on these sites intermittently in the past. Hand-pulling and cutting treatments were usually administered at least once during any one growing season. Many of these access sites receive annual treatments of this nature. Target weed species are spotted knapweed and leafy spurge.

Biological control agents (*Hyles euphorbiae*, *Aphthona nigriscutis*, *Aphthona flava*, and *Oberea erythrocephala*) were released on Kokanee Bend and *Urophora affinis* on Old Steel Bridge FASes respectively by the Flathead County Weed Board and Montana State University Extension Service in an attempt to treat leafy spurge. Initial releases of the first agent were made on Kokanee Bend on July 21, 1985. Releases of the latter three agents occurred at Kokanee Bend on July 10, 1992. Sites are inspected for agent survival, distribution, and effectiveness at least annually. Periodically, agents will be redistributed to new or different weed patches once

they've been introduced to a site. This redistribution sustains the life cycles of host-specific agents and provides weed treatment benefits to a broader area. Thus far, these agents have survived and appear to be expanding their territories. Biocontrol techniques will be relied upon more in the future as they become more readily available.

Some weed management on Flathead River FASes has been by chemical means, although FWP has begun to move away from a total reliance on chemical treatment methods on its FASes. Chemical methods will be reserved for those sites (or weed patches on a site) where application will not impair site values and resources. Manual (hand pulling), cultural (mowing or cutting), and biological control methods will be emphasized.

In addition to treating established weeds, FWP will attempt to curtail noxious weed spread by restricting vehicles to established roads through traffic control devices such as rock barriers. Disturbance to soil and vegetation communities during facility development and maintenance will be minimized. Sites disturbed during facility development will be reseeded.

3.2.4 Kootenai River Drainage

Existing Environment

FWP maintains the following 7 fishing access sites within the Kootenai drainage:

Frank Lake	Moran Lake
Glen Lake	Sophie Lake
Loon Lake/Eureka	Tetrault Lake
Marl Lake	

The Kootenai flows approximately 140 miles from the British Columbia border to the Idaho state line. Ninety miles of the river is impounded by Lake Koocanusa and 50 miles is flowing river from Libby Dam to the Idaho border. The Kootenai River at the Idaho border is the lowest elevation point in the state.

The west side of Lake Koocanusa is mountainous with coniferous forests. There is heavy logging activity with some agriculture. The east side of Lake Koocanusa, where the fishing access sites are located, is in a rainshadow with low precipitation. The terrain is low, rolling, glaciated hills with pothole lakes. Vegetation is Douglas-fir, Ponderosa pine, grassland and even some sagebrush. Primary land use is agricultural and residential with some logging. Lake Koocanusa supported 47,300 days of fishing and the Kootenai River supported 16,800 days of fishing in 1991, making them some of the most heavily waters in the state. The small lakes in the area each provide from 300 to 1,000 days of fishing annually. The lakes contain rainbow and cutthroat trout, mountain whitefish, kokanee, pumpkinseeds and bluegill, yellow perch, largemouth bass and northern pike.

Weed Status

Kootenai River drainage FASes are accessible via well-travelled U.S. highway 2 serving both as a route to Glacier National Park and the northernmost route connecting Montana with Idaho. and U.S. Highway 93 which connects Montana with British Columbia. This proximity increases the potential of noxious weed introduction to these FASes. Also, subdivided "ranchettes" in the Libby and Eureka areas lacking adequate noxious weed management efforts serve as potential sources of noxious weed seeds for the entire drainage downstream. In the upper Kootenai River valley (near the town of Eureka) spotted knapweed is a problem for many landowners, not just FWP. All of the FASes within the Kootenai River drainage are located on Ponds or Lakes.

Site Constraints

Chemical treatment of noxious weeds on Kootenai River FASes is constrained by the presence of surface waters and by bedrock or gravelly soil types which easily transport chemicals to shallow groundwater tables or to surface waters. Furthermore, high levels of human use, particularly during the summer months, limit the use of chemicals even in those areas where they can be safely applied without risk to water resources. Riparian vegetation consisting of cottonwood/willow bottoms with a shrub and herbaceous understory component may also preclude certain chemical treatments.

Weed Management Strategies and Methods

Manual and cultural methods have been applied on some of these sites intermittently in the past. Hand-pulling and cutting treatments were usually administered at least once during any one growing season. Many of these access sites received annual treatments of this nature. Target weed species were spotted knapweed.

Some past weed management on Kootenai River FASes was by chemical means, although FWP has begun to move away from a total reliance on chemical treatment methods on its FASes. Chemical methods will be reserved for those sites (or weed patches on a site) where application will not impair site values and resources. Manual (hand pulling), cultural (mowing or cutting), and biological methods will be emphasized.

In addition to treating established weeds, FWP will attempt to curtail noxious weed spread by restricting vehicles to established roads through traffic control devices such as rock barriers. Disturbance to soil and vegetation communities during facility development and maintenance will be minimized. Sites disturbed during facility development will be reseeded.

3.2.5 Fish Hatcheries and Administrative Sites

Existing Environment

Flathead Lake Salmon Hatchery-This site consists of a hatchery building, a residence and an upper spring site. The lower site is largely in lawn, and weeds are controlled by cutting. The proximity to Flathead Lake makes chemical control difficult.

Jocko River Trout Hatchery-The Jocko River Trout Hatchery consists of the hatchery site, residences on an upper bench and a riparian area leading to the spring site. Much of the administrative site is in lawns, and weeds are controlled by cutting. Some weeds are controlled by spot chemical applications. The riparian area/spring site are difficult to treat due to the coarse porous soils and high groundwater.

Lake Mary Ronan Spawn Station-This station consists of 0.27 acres of fee title land and 40 acres of leased land. The site has not been in use for many years and is under the management of the private landowner.

Lore Lake Spawn Station-The Lore Lake spawn station is no longer in use and is under the management of a private landowner.

Libby Field Station-This 67 acre site consists of administrative offices and parking, outbuildings, an extensive spring area and riparian zone. Some of the site is in lawn, and cutting is primary weed control method. Chemical treatment is complicated by coarse, porous soils and shallow groundwater.

Lone Lake Hatchery-This site is a 4.33 acre lease which is no longer in use. The site is under management of the private landowner.

Rogers Lake Spawn Station-This site is 0.63 acre of fee title land with an access easement through Plum Creek timberlands. The site consists of a small stream and associated riparian zone. The site is used occasionally and generally is weed free. There is currently no weed control program. Chemical treatment would be complicated by shallow groundwater.

Region One Headquarters Complex - Administrative sites are those reserved for use by FWP. The Regional Headquarters complex is located in Kalispell, totalling 3.27 acres. This complex serves as the public office of FWP and as a parking/storage facility for vehicles, trailers, and supplies and equipment.

Rose Creek Hatchery Site-This site is 20.2 acres of fee title land. The site consists of a small stream and associated riparian zone. There is currently no weed control program. Chemical treatment would be complicated by shallow groundwater.

Weed Status

Because vehicles are one agent of weed seed dispersal, occasionally a noxious weed will root in the parking lot or storage compound of the Regional Headquarters. Spotted knapweed and some thistle species (Canada, bull, and sow) can occur from time to time on these sites.

Site Constraints

On all sites, chemical treatment of noxious weeds is constrained by the presence of surface waters and by gravelly soil types which easily transport chemicals to shallow groundwater tables or to surface waters. Furthermore, high levels of human use limit the use of chemicals even in those areas where they can be safely applied without risk to water resources.

Weed Management Strategies

Manual and cultural methods have been applied intermittently when necessary. Hand-pulling and cutting treatments were usually administered at least once during any one growing season. Many of these sites receive annual treatments. Target weed species were spotted knapweed and thistle.

Some weed management on these sites was by chemical means, although FWP has begun to move away from the use of chemical treatment methods on its administrative sites. Chemical methods will be reserved for those sites (or weed patches on a site) where application will not impair site values and resources. Manual (hand pulling), cultural (mowing or cutting), biological and chemical methods will all be used, although the first three will be emphasized.

In addition to treating established weeds, FWP will attempt to curtail noxious weed spread by landscaping and irrigation where feasible, and restricting vehicles to established roads through traffic control devices. Disturbance to soil and vegetation communities during facility development and maintenance will be minimized. Sites disturbed during facility development will be reseeded.

3.3 State Parks and Affiliated Lands

3.3.1 Introduction

Preserved for their natural, historic, and recreation features, Montana's State Parks (SP's) add a great dimension to recreational opportunities in Montana. Protecting some of the best natural and cultural resources of Montana, State Parks offer visitors a glimpse into Montana's past, the discovery of unique natural features, and the enjoyment of leisure time spent outdoors.

The Montana State Park System is managed through a number of administrative regions. The region one park operation is comprised of two districts responsible for all management functions on 6 state parks totalling 2,993 acres. They are: Flathead Lake, Lake Mary Ronan, Logan, Lone Pine, Thompson Falls, and Whitefish Lake. This includes such complex areas as 2100-acre Wild Horse Island unit of Flathead Lake State Park with rare and endangered plants, and bald eagle and bighorn sheep populations; and Logan State Park, the centerpiece of the Thompson Chain of Lakes with 21 fragile recreational lakes and wetlands in checkerboard ownership with timberland, private cabin sites and livestock grazing.

Responsibilities also include the 260-mile groomed snowmobile trail program in four counties with the highest avalanche hazards in the state, and recreational trails and off-highway vehicle programs.

Visitation for all region one parks responsibilities exceeds 700,000 recreation days annually including the highest percentage of non-resident visitors in the state. Over 60 personnel carry out a variety of tasks, including park managers, park rangers, maintenance supervisors, crew chiefs, park and visitor station attendants, administrative aids, interns, youth crews, and park hosts. This includes a high percentage of seasonal employees and volunteers with 14.39 funded FTE responsible for operations, maintenance and visitor services of day use parks and 256 primitive to full-service campsites and boat slips in and adjacent to more urban areas than any other region of the state.

Combined region one state park operations and capital expenditures exceed \$1 million annually from over a dozen earmarked special state and federal revenue sources. Earned revenues from entrance and camping permits, facility rentals, concessions, sales of publications and supplies, tours, grazing and timber leases, access easements, and commercial use permits exceed \$200,000 annually.

These properties provide natural settings to camp, hike, boat, picnic, view wildlife, take pictures, etc. The setting and visitor use facilities at a typical state park include the following: access roads, parking areas, interpretive signs, flush toilets, tent and RV campsites with or without utility hookups, drinking water faucets, picnic tables, firegrills, boat ramp, day use and overnight boat docks with and without utility hookups, swimming areas, and picnic areas. Additional facilities may include day use and overnight group use areas, hiking trails, a meeting room and a visitor center.

The goal of FWP is to administer state parks by enacting good land management policies which protect vegetation and soil resources. To the degree that vegetation communities are healthy and vigorous, they will resist noxious weed encroachment. However, state parks are a small part of a larger landscape in which, unfortunately, noxious weeds occur, and noxious weeds have been documented on most state parks in Region One. FWP is required by law to treat noxious weeds. As such, any noxious weed management program undertaken should be approached at the landscape level, considering neighboring landowners, land use practices, and mechanisms of seed dispersal. As well, noxious weed management strategies must take into account the high natural resource and human use values of these sites. To that end, weed management in state parks will center around maintaining healthy vegetation communities, deterring seed dispersal, and the detection and treatment of noxious weed outbreaks before they become well established.

Healthy vegetation communities can be promoted by limiting excessive soil disturbance which facilitates noxious weed establishment. Soil disturbance can be prevented by confining the use of motorized vehicles to established roads and trails and by not disturbing vegetation unnecessarily during the development or maintenance of visitor use facilities. By restricting vehicular traffic to designated roadways, seed dispersal is also confined to areas where weeds can be easily detected and treated.

Preventing noxious weed dispersal is the ultimate goal. However, the mobility of our society makes that a difficult goal to attain. Quick detection and treatment of noxious weed outbreaks is the most efficient approach to manage weeds, keeping their dispersal rate at a manageable level. As weed patches increase in size, potential treatment methods and techniques dwindle as other considerations come into play. Given the amount of acreage contained within state parks in Region One, budget and personnel constraints require the implementation of methods which maximize the successful weed management to cost ratio. Therefore, the management of noxious weeds in state parks will be accomplished using the concept of Integrated Noxious Weed Management (INWM). Where appropriate, chemicals will be applied. However, other methods and techniques such as hand-pulling, mowing, and biological control agents will be emphasized.

Noxious weed treatment plans for state parks follow. The Flathead Lake units except for Wild Horse Island will be discussed concurrently, whereas Flathead Lake-Wild Horse Island, Lake Mary Ronan, Logan, Lone Pine, Thompson Falls, and Whitefish Lake will be discussed individually. Figure 4 illustrates the locations of these parks in Region One.

Figure 4. State Parks administered by FWP in Region One.

3.3.2 Flathead Lake

Existing Environment

Flathead Lake State Park is located between Kalispell and Polson on Highways 93 and 35. It is renowned for boating, sailing, fishing, camping and swimming. The park's six units, totalling 2606 acres acquired over the past 40 years, provide public access to the shores of Flathead Lake, which are privately owned, and provide boat access to the park's 2164 acre Wild Horse Island. The 217 acre Big Arm unit itself is leased from the Montana Department of State Lands. The current lease cost is \$14,000 per year.

The lake itself, the largest freshwater lake west of the Mississippi, was dredged out by receding glaciers during the Wisconsin period of glaciation which ended 10,000 years ago. It is 28 miles long and from 5 to 15 miles wide. Encompassing 188 square miles at an elevation of 2,892 feet, the lake's maximum depth is 386 feet and it has 128 miles of shoreline.

There are five campgrounds and 128 campsites with picnic tables and firegrills. Other facilities include drinking water faucets, hiking and vista trails, flush and vault toilets, and all areas have boat ramps and access to the lake for swimming and fishing. A trailer sewage dump station is located at Wayfarers, and a boat sewage dump station is located at Finley Point. Finley Point has RV hookups and overnight boat mooring docks with hookups. Group picnicking and group camping areas are available at Big Arm.

Weed Status

Noxious weeds occurring at Flathead Lake include: spotted knapweed, Canada thistle, leafy spurge, Sulphur cinquefoil and mullein. Other exotic plant species include Russian olive at Big Arm and apple and pear orchards associated with homesteading at Wayfarers and Wild Horse Island.

Spotted knapweed is the most serious of the noxious weed species documented at this park. Where it occurs near the lakeshore and traditional camping areas it is difficult to treat. Leafy spurge is not a problem at this park currently. However, small patches have been found on Wild Horse Island where access is difficult, and seed sources exist upstream along the Flathead River.

Flathead Lake lies along two major State Highways, so the potential for noxious weed introduction from such a source is great. Recent increases in tourism and tourism advertizing is also adding to the volume of traffic, bringing visitors to the park from all over the nation and Canada as well. This increased volume of traffic increases the potential for noxious weed introduction to Flathead Lake.

Site Constraints

Chemical methods of noxious weed treatment at Flathead Lake is constrained by the presence of surface water and by high levels of human use, particularly during the spring and summer seasons.

Weed Management Strategies and Methods

In the past, weed management at Flathead Lake was by a combination of pulling, mowing, and chemical means. As FWP has begun to move away from a total reliance on chemical treatment methods on its state parks, chemical methods will be reserved for those weed patches where application of chemicals will not impair site values and resources. Should weeds spread to areas where chemical methods are not appropriate, cultural, manual, and biocontrol methods will be used. The biological control agents, *Urophora affinis* and *Urophora quadrifasciata*, were introduced by the Flathead County Extension Agent in July, 1992 and earlier to attempt to treat spotted knapweed.

In addition to treating established weeds, FWP will attempt to curtail noxious weed spread by restricting vehicles to established roads through traffic control devices and on-site patrols. In natural zones, the introduction of exotic plant species will be prevented, or where already present, controlled or eradicated. Disturbance to soil and vegetation communities during facility development and maintenance will be minimized. Sites disturbed during facility development will be reseeded.

Bald eagles utilize areas on the lakeshore, on Wild Horse Island, and near the mouth of the Flathead River for feeding, nesting and flight paths during their migration. Weed management activities should not adversely affect this rare species of bird.

There is no record of any other threatened or endangered species in the immediate area. Species considered include: black-footed ferret, gray wolf, grizzly bear, eskimo curlew, interior least tern, piping plover, whooping crane, bald eagle and peregrine falcon. Among Montana fishes, the White Sturgeon is listed as endangered, and the Bull Trout is currently listed as warranted but precluded (see Appendix F).

3.3.3 Flathead Lake-Wild Horse Island

Existing Environment

One of the largest island in the inland U.S., this natural park, acquired in 1978, is noted for its endangered Spalding Campion palouse prairie plant species and incredible wildlife. A herd of about 200 rocky Mountain bighorn sheep, 100 mule deer, several whitetail deer, songbirds, waterfowl, bald eagles, and five wild horses make this 2,163 acre island their home. Boat access via Big Arm is carefully regulated. The island is maintained as a state primitive park.

There are no facilities or marked trails. There are 56 private lots along the shoreline of the island of which 27 have houses, docks or other developments on them to date.

Weed Status

The proximity of the lake will continue to be a source of weed seeds since they can disperse via water. All weed populations on the shores of Flathead Lake are potential sources of noxious weed seeds to Wild Horse Island, as well as human dispersion from boats, clothing, etc.

Noxious weeds occurring at Wild Horse Island include: Canada thistle, spotted knapweed, and leafy spurge (Sawyer 1994). Spotted knapweed is the most serious of the noxious weed species documented on this site. Canada thistle is near riparian areas and is difficult to treat. Two patches of leafy spurge have been found on the island and wildlife could spread seed if it is not eradicated soon.

Other exotic plant species include mullein and cheatgrass on the drier infertile hilltops, and an old homestead orchard (pre 1940) containing apple and pear trees.

Site Constraints

Chemical methods of noxious weed treatment on Wild Horse Island may be constrained along the shoreline by the presence of nearby surface water. However, low human use levels and lack of wetlands favor chemical use on inland sites. Management of the area as a primitive park precludes mechanical treatment although pulling and biological control may be feasible.

Weed Management Strategies and Methods

In the past, weed management on Wild Horse Island was by pulling. In 1993 a weed management survey was completed by Dr. Pete Fay of Montana State University (Fay 1993). Recommendations included an aggressive weed prevention/eradication program, public education program, and range monitoring plan be funded soon by FWP. As FWP moves away from total reliance on chemical treatment methods on state parks, chemical methods will be reserved for those weed patches where application of chemicals will not impair site values and resources. Should weeds spread to areas where chemical methods are not appropriate, cultural, manual, and biocontrol methods will be used. The biological control agent, *Ceutorhynchus litura*, will be introduced to treat Canada thistle patches if it proves feasible.

FWP will attempt to eradicate noxious weeds on Wild Horse Island using chemical, manual, cultural, and biological methods while protecting and encouraging endangered plant and animal species. Exotic plants such as mullein and cheatgrass will be controlled, and the specific varieties of apple and pear in the orchard maintained as a "living history" resource. Additionally, Ponderosa pine encroachment into the Palouse Prairie grassland will be controlled to protect this important habitat type.

3.3.4 Lake Mary Ronan

Existing Environment

Lake Mary Ronan is managed as a state primitive park. It was acquired in 1965 to provide access to Lake Mary Ronan for recreational activities and totals 76 acres. There are private resorts within walking distance with restaurants and lounges. Attractions include lake fishing for trout, bass, and kokanee salmon, and bird watching. Visitor facilities include: drinking water faucets, vault toilets, a 26 unit campground, a boat ramp, parking area, and gravel access road.

Weed Status

Noxious weeds occurring at Lake Mary Ronan include: Canada thistle and spotted knapweed. Spotted knapweed is the most commonly encountered species. Currently, its distribution is associated with adjacent recently logged areas and established roadways, but knapweed and thistle both are expanding from adjacent recently logged areas.

Site Constraints

Chemical treatment of noxious weeds on this site is constrained by the high levels of human use, especially during the spring when fishing is best in the lake. Chemical applications should not occur in the vicinity of the lake.

Weed Management Strategies and Methods

In the past, weed management at Lake Mary Ronan was by a combination of pulling, mowing, and chemical means. As FWP begins to move away from a total reliance on chemical treatments on its state parks, chemical methods will be reserved for those weed patches occurring in open areas where application of chemicals will not impair other site values and resources. Other methods such as cultural, manual, and biocontrol will be applied elsewhere.

In addition to treating established weeds, FWP will attempt to curtail noxious weed spread by restricting vehicles to established roads through traffic control devices and on-site patrols. Disturbance to soil and vegetation communities during facility repair and maintenance will be minimized. Sites disturbed during facility repairs will be reseeded.

3.3.5 Logan

Existing Environment

Located 45 miles west of Kalispell in Lincoln County, this 17.58 acre state park was acquired in 1967. With frontage on the north shore of Middle Thompson Lake, Logan is heavily forested with western larch, Douglas-fir, and Ponderosa pine. A channel connects Upper, Middle, and

Lower Thompson lakes. Attractions include swimming, boating, camping, water skiing, and fishing for rainbow trout, largemouth bass, kokanee salmon, and yellow perch. Visitor facilities include: drinking water faucets, flush and vault toilets, a 41 unit campground, a boat launch, parking area, and gravel access road.

Weed Status

The proximity to U.S. highway 2 and spotted knapweed populations in nearby lands disturbed by recent logging increases the potential for noxious weed introduction to Logan. Spotted knapweed beginning to invade this site. There are also dense patches of Canada thistle near the lakeshore and highway.

Site Constraints

Chemical treatment of noxious weeds at Logan is constrained in developed areas by a high water table and high levels of human use, particularly during the summer season.

Weed Management Strategies and Methods

Past weed management at Logan was by pulling and mowing. Thus far, annual treatments do not appear to be effective. In addition to treating established weeds, FWP will attempt to curtail noxious weed spread by restricting vehicles to established roads through traffic control devices and on-site patrols. Disturbance to soil and vegetation communities during facility development and maintenance will be minimized. Sites disturbed during development will be reseeded.

3.3.6 Lone Pine

Existing Environment

Lone Pine, five miles southwest of Kalispell, was acquired by donation in 1941. Its 182 acres offers several hiking trails, as well as horse trails and an archery range. There are three scenic overlooks that provide views from Flathead Lake to Big Mountain Ski Area. One of the overlook trails is designed for the mobility impaired visitor. The visitor center has nature and interpretive programs and includes a 100-person capacity meeting room that can be reserved. A group picnic shelter is also available on a reservation basis.

Weed Status

Noxious weeds occurring at Lone Pine include spotted knapweed, Canada thistle, and St. Johnswort (Goatweed). Spotted knapweed is the most commonly encountered species. Currently, its distribution is associated with adjacent recently logged areas, private roadway easements, trails and dry hillsides, but it is expanding despite previous treatment efforts. Presently St. Johnswort is of low density and restricted to the dry south-facing hillsides.

Biological control agents (*Urophora affinis*, *Chrysolina quadrigemini*) were released at Lone Pine by the Flathead County Weed Board in an attempt to treat spotted knapweed and St. Johnswort, respectively. One release of the first agent was made in about 1987, and two releases of the latter were made on July 1 and July 10, 1993.

Site Constraints

Chemical treatment of noxious weeds on this site is constrained by adjacent residential subdivisions and high levels of human use, especially during the spring and summer.

Weed Management Strategies and Methods

In the past, weed management at Lone Pine was by a combination of pulling, mowing, and chemical means. As FWP begins to move away from a total reliance on chemical treatments on its state parks, chemical methods will be reserved for those weed patches occurring in open areas where application of chemicals will not impair other site values and resources. Other methods such as cultural, manual, and biocontrol will be applied elsewhere.

In addition to treating established weeds, FWP will attempt to curtail noxious weed spread by restricting vehicles to established roads through traffic control devices and on-site patrols. Disturbance to soil and vegetation communities during facility repair and maintenance will be minimized. Sites disturbed during facility repairs will be reseeded.

3.3.7 Thompson Falls

Existing Environment

Located one mile northwest of the town of Thompson Falls on old Montana highway 200, this 36 acre park is leased from the Montana Department of State Lands for \$2,000 per year. It is managed as a state primitive park. Attractions include bird watching, fishing for bass, trout, and ling, nature walks, and boating on Noxon Rapids Reservoir. Visitor facilities include: drinking water faucets, 3 vault toilets, a 17 unit campground, a boat ramp, parking area, and paved access road.

Weed Status

The proximity to highway 200 has introduced spotted knapweed to a good portion of the park. The Clark Fork river adjacent to the park also increases the potential for additional noxious weed introduction to Thompson Falls. Thus far, spotted knapweed, Canada thistle, and two 25 foot by 50 foot patches of leafy spurge have been inventoried at this park.

Site Constraints

Chemical treatment of noxious weeds at Thompson Falls is constrained in developed areas by proximity to the river and high levels of human use, particularly during the summer season.

Weed Management Strategies and Methods

Past weed management at Thompson Falls was by pulling and mowing. Annual treatments do not appear to be effective. In addition to treating established weeds, FWP will attempt to curtail noxious weed spread by restricting vehicles to established roads through traffic control devices and on-site patrols. Disturbance to soil and vegetation communities during facility repairs and maintenance will be minimized. Sites disturbed during repair will be reseeded.

3.3.8 Whitefish Lake

Existing Environment

Whitefish Lake State Park is located one mile west of the town of Whitefish and north of U.S. highway 93. It was acquired in 1960 from the Whitefish Junior Chamber of Commerce. Attractions at this small 10.35 acre park include boating, swimming, and fishing. Whitefish Lake is rarely windy, often providing glassy conditions for waterskiing. Looking north across the lake, you can see the ski runs of Big Mountain. The nearby city of Whitefish is a year-round resort, with the Whitefish Golf Course within walking distance of the park. Good restaurants and motel accommodations are also available in the city. Visitor facilities include: drinking water faucets, flush and vault toilets, a 33 unit campground, a boat ramp, parking area, and access road.

Weed Status

The proximity to U.S. highway 93, the lakeshore, and nearby lands disturbed by recent residential development increases the potential for noxious weed introduction to Whitefish Lake state park. Thus far, no noxious weeds have been inventoried at this park.

Site Constraints

Chemical treatment of noxious weeds at Whitefish Lake is constrained in developed areas by a high water table and high levels of human use, particularly during the summer season.

Weed Management Strategies and Methods

FWP will attempt to prevent noxious weed introductions by restricting vehicles to established roads through traffic control devices and on-site patrols. Disturbance to soil and vegetation communities during facility development and maintenance will be minimized. Sites disturbed during development will be reseeded.

There is no weed management program at the present time. However, noxious weeds may become a problem sometime in the future. FWP will continue informal inspections to monitor weed status. Should weed management be necessary in the future, any of the four treatment methods (manual, cultural, chemical, or biological) may be used, depending on the circumstances. Any weed management actions will be conducted subject to the guidelines presented in this Plan.

4.0 REFERENCES

- Dobkin, D. S. 1992. Neotropical migrant landbirds in the Northern Rockies and Great Plains. U.S. Department of Agriculture Forest Service Northern Region. Publication no. R1-93-34. Missoula, Montana.
- Fay, P. 1993. Weed Management Suggestions for Wild Horse Island on Flathead Lake. Extension Weed Scientist, Department of Plant and Soil Science, Montana State University, Bozeman, Montana.
- Flath, D. 1993. Non-Game Coordinator for the Montana Department of Fish, Wildlife and Parks. Personal communication with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Flathead County Weed District. March, 1994. Flathead County Weed Management Plan. Kalispell, Montana.
- Flathead National Forest. 1991. Final Environmental Impact Statement Noxious Weed Management Projects, Spotted Bear and Hungry Horse Ranger Districts. U.S. Department of Agriculture, Flathead National Forest. Hungry Horse, Montana.
- Glacier National Park. 1991. Exotic Vegetation Management Plan. Division of Resource Management and Visitor Protection, Glacier National Park, West Glacier, Montana.
- Heidel, B. 1993. Botanist with the Montana Natural Heritage Program. Personal communication with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Hiebert, Ronald A. 1986. Indiana Dunes National Lakeshore.
- Konizeski, D. 1982. The Montanan's Fishing Guide, Volume II. Revised and Updated by J. A. Derleth. Mountain Press Publishing Company. Missoula, Montana. 307pp.
- Johnson, J. L. and R. D. Pfister. 1981. Survey of potential ecological natural landmarks of the northern rocky mountains. IFRES U.S. Department of Agriculture Forest Service. MT-25.
- Jorgenson, H. 1993. Plant Ecologist for the Montana Department of Fish, Wildlife and Parks. Personal communication with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Lacey, C. 1991. Groundwater and agrichemical protection pilot project. Unpubl. report and maps prepared for Lewis and Clark County Conservation District.
- Lake County Weed Control District. 1993. Proposed Noxious Weed Management Plan for Lake County. Polson, Montana.

- Lessica, P. 1993. Botanist. Missoula, Montana. Personal communication with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Lincoln County Weed and Rodent Board. April, 1991. Lincoln County Integrated Noxious Weed Management Plan. Libby, Montana.
- Lolo National Forest. 1989. Noxious weed management draft environmental impact statement, amendment to the Lolo National Forest Plan. U.S. Department of Agriculture, Lolo National Forest. Missoula, Montana.
- McKenna, D. 1993. Associate Research Hydrogeologist, Montana Groundwater Assessment Program, Montana College of Mineral Science and Technology. Personal communication with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Montana Department of Agriculture. 1991. Draft noxious weed trust fund programmatic environmental impact statement. Helena, Montana.
- Montana Fish and Game Department. 1975. Inventory and plans: Game Management Areas. Montana Department of Fish, Wildlife and Parks files.
- Montana Department of Fish, Wildlife and Parks. 1983. Weed control program for lands managed by the Montana Department of Fish, Wildlife and Parks. Montana Department of Fish, Wildlife and Parks files. 4pp.
- Montana Department of Fish, Wildlife and Parks. 1989. Montana statewide angling pressure, 1989. Montana Department of Fish, Wildlife and Parks files.
- Montana Department of Fish, Wildlife and Parks. 19 . Kootenai Wildlife Management Area draft management plan. Montana Department of Fish, Wildlife and Parks files. pp.
- Montana Department of Fish, Wildlife and Parks. 1991. Montana statewide angling pressure, 1991. Montana Department of Fish, Wildlife and Parks files.
- Montana Department of Fish, Wildlife and Parks. 1992. Creating a Vision for the future. Montana Department of Fish, Wildlife and Parks files.
- Montana Department of Fish, Wildlife and Parks. 1992. Lands inventory. Montana Department of Fish, Wildlife and Parks Lands Section files, Helena.
- Montana Department of Fish, Wildlife and Parks. 1993. Draft integrated noxious weed management plan for Region 8. Prepared by OEA Research Inc. Helena.
- Montana Department of Fish, Wildlife and Parks. 1993. Draft weed management plan for Region 2. Prepared by Land and Water Consulting Inc. Missoula.

- Montana Natural Heritage Program. 1993. Results of a Heritage Program database search. Montana Natural Heritage Program, Helena.
- Newby, R. 1988. Great Escapes: Montana State Parks. Falcon Press Publishing Co., Inc. 74pp.
- Phillips, G. 1993. Pollution Control Biologist and Water Quality Specialist for the Montana Department of Fish, Wildlife and Parks. Personal communication with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Sanders County Weed Control District. 1994. Sanders County Weed Management Program. Thompson Falls, Montana.
- Sawyer, J. 1994. Flathead Lake State Park-Wild Horse Island Management Plan Update. Parks Division-Region One, Montana Department of Fish, Wildlife and Parks. Kalispell, Montana.

5.0 APPENDICES

- APPENDIX A:** County Noxious Weed Control Act of 1979
- APPENDIX B:** Synopsis of Noxious Weed and Exotic Plant Species
- APPENDIX C:** Laws and Policies Relating to the Management of Noxious Weeds
- APPENDIX D:** Biological Control Agents Available in Montana and Those Released on Fish, Wildlife & Parks Properties
- APPENDIX E:** Herbicide Emergency Response Plan
- APPENDIX F:** Sensitive Species Occurring in the Vicinity of Fish, Wildlife & Parks Region One Properties
- APPENDIX G:** Herbicides Proposed for Use on Fish, Wildlife & Parks Region One Properties
- APPENDIX H:** Herbicide Label Instructions and Safety Precautions

APPENDIX A

COUNTY NOXIOUS WEED CONTROL ACT OF 1979

**COUNTY NOXIOUS WEED
CONTROL ACT
AND
ADMINISTRATIVE RULES**

**STATE OF MONTANA
DEPARTMENT OF AGRICULTURE
HELENA, MONTANA**

JULY, 1991

COUNTY NOXIOUS WEED CONTROL ACT

Title 7, Chapter 22

Sections

7-22-2101 through 7-22-2153

M C A

Amended 1991

AND RULES

Rules 4.5.201 through 4.5.203

State of Montana
Department of Agriculture
Agricultural and Biological Sciences Division
Capitol Station
Helena, MT 59620-0205
(406) 444-2944

TABLE OF CONTENTS

<u>Section</u>	<u>Title</u>	<u>Page</u>
7-22-2101	Definitions	1
7-22-2102	Weed Management Districts Established	2
7-22-2103	District Weed Board -- Appointment and Term .	2
7-22-21-4	Term of Office.	2
7-22-2105	Organization of District Weed Board and Compensation	2
7-22-2106	Renumbered 7-22-2115 by Code Commission . . .	3
7-22-2107	Renumbered 7-22-2116 by Code Commission . . .	3
7-22-2108	Renumbered 7-22-2117 by Code Commission . . .	3
7-22-2109	Powers and Duties of Board.	3
7-22-2110	Administrative Hearing -- Appeals	3
7-22-2111	(Temporary) Liability Restrictions.	4
7-22-2112	(Temporary) Information on Herbicide Use. . .	4
7-22-2113 and 7-22-2114	Reserved.	4
7-22-2115	Noxious Weeds and Seeds Declared Nuisance . .	4
7-22-2116	Unlawful to Permit Noxious Weeds to Propagate.	4
7-22-2117	Violations.	4
7-22-2118 through 7-22-2120	Reserved.	5
7-22-2121	Weed Management Program.	5
7-22-2122	Repealed.	5
7-22-2123	Procedure in a Case of Noncompliance.	5
7-22-2124	Destruction of Weeds by Board	6
7-22-2125	Repealed.	7
7-22-2126	Embargo	7
7-22-2127	Repealed.	7
7-22-2128 and 7-22-2129	Reserved.	7
7-22-2130	Weed District Supervisor Training	7
7-22-2131 through 7-22-2140	Reserved.	7
7-22-2141	Noxious Weed Fund Authorized.	7
7-22-2142	Sources of Money for Noxious Weed Fund. . . .	7
7-22-2143	Determination of Cost of Weed Control Program.	8
7-22-2144	Payment of Cost of Weed Control Program . . .	8
7-22-2145	Expenditures From Noxious Weed Fund	8
7-22-2146	Financial Assistance to Persons Responsible for Weed Control	8
7-22-2147	Repealed.	9
7-22-2148	Tax Liability for Payment of Weed Control Expenses.	9
7-22-2149	Responsibility for Assessments and Taxes for Weed Control Levied on Leased State Lands.	9
7-22-2150	Cooperation With State and Federal-Aid Programs	9
7-22-2151	Cooperative Agreements.	10
7-22-2152	Revegetation of Rights-of-Way and Disturbed Areas.	10
7-22-2153	Voluntary Agreements for Control of Noxious Weeds Along Roads	11

COUNTY NOXIOUS WEED CONTROL RULES

4.5.201	Designation of Noxious Weeds.	12
4.5.202	Category 1.	12
4.5.203	Category 2.	12
4.5.204	Category 3.	13
Part Cross References		14

COUNTY NOXIOUS WEED CONTROL ACT

7-22-2101. Definitions. As used in this part, unless the context indicates otherwise, the following definitions apply:

- (1) "Board" means a district weed board created under 7-22-2103.
- (2) "Commissioners" means the board of county commissioners.
- (3) "Department" means the department of agriculture provided for in 2-15-3001.
- (4) "District" means a weed management district organized under 7-22-2102.
- (5) "Native plant" means a plant endemic to the state of Montana.
- (6) "Native plant community" means an assemblage of native plants occurring in a natural habitat.
- (7) (a) "Noxious weeds" or "weeds" means any exotic plant species established or that may be introduced in the state which may render land unfit for agriculture, forestry, livestock, wildlife, or other beneficial uses or that may harm native plant communities and that is designated:
 - (i) as a statewide noxious weed by rule of the department; or
 - (ii) as a district noxious weed by a board, following public notice of intent and a public hearing.(b) A weed designated by rule of the department as a statewide noxious weed must be considered noxious in every district of the state.
- (8) "Person" means an individual, partnership, corporation, association, or state or local government agency or subdivision owning, occupying, or controlling any land, easement, or right-of-way, including any county, state, or federally owned and controlled highway, drainage or irrigation ditch, spoil bank, borrow pit, or right-of-way for a canal or lateral.
- (9) "Supervisor" means the person employed by the board to conduct the district noxious weed management program and supervise other district employees.
- (10) "Weed management" or "control" means the planning and implementation of a coordinated program for the containment, suppression, and, where possible, eradication of noxious weeds.

7-22-2102. Weed Management Districts Established. A weed management district shall be formed in every county of this state and shall include all the land within the boundaries of the county, except that a weed management district may include more than one county through agreement of the commissioners of the affected counties.

7-22-2103. District Weed Board -- Appointment and Term. (1) The commissioners shall appoint a district weed board.

(2) The commissioners shall, at a public meeting, pass a resolution establishing the number of members of the district weed board and the terms of the appointments. The board must consist of at least three members and no more than nine members, and the members of the board must be residents of the district. A majority of the board members must be rural agricultural land owners.

(3) The county extension agent in each county and other interested individuals may be appointed to serve as nonvoting members of that district's weed board.

(4) The board members are public officers.

(5) The board may call upon the county attorney for legal advice and services as it may require.

7-22-2104. Term of Office. (1) Except as provided in subsection (2), a member of a district weed board serves a term of 3 years and until the qualification of his successor. The term of office begins January 1.

(2) When a three-member weed board is established, the initial board members serve terms of 1, 2, and 3 years, respectively, as designated by the commissioners. When a five-member weed board is established, two of the initial members serve terms of 1 year, two serve terms of 2 years, and one serves a term of 3 years. After expiration of an initial term of office, the successor serves a 3-year term as provided in subsection (1).

7-22-2105. Organization of District Weed Board and Compensation.

(1) The board shall organize by choosing a chairman and a secretary. The secretary may or may not be a member of the board.

(2) Salary, per diem, and mileage of such board members shall be set by resolution of the commissioners.

(3) A majority of the board constitutes a quorum for the conduct of business.

7-22-2106. Renumbered 7-22-2115 by Code Commissioner, 1985.

7-22-2107. Renumbered 7-22-2116 by Code Commissioner, 1985.

7-22-2108. Renumbered 7-22-2117 by Code Commissioner, 1985.

7-22-2109. Powers and Duties of Board. (1) The board may:

(a) employ a supervisor and other employees as necessary and provide for their compensation;

(b) purchase such chemicals, materials, and equipment and pay other operational costs as it determines necessary for implementing an effective weed management program. Such costs must be paid from the noxious weed fund.

(c) determine what chemicals, materials, or equipment may be made available to persons controlling weeds on their own land. The cost for such chemicals, materials, or equipment must be paid by such person and collected as provided in this part.

(d) enter into agreements with the department for the control and eradication of any new exotic plant species not previously established in the state which may render land unfit for agriculture, forestry, livestock, wildlife, or other beneficial use if such plant species spreads or threatens to spread into the state; and

(e) perform other activities relating to weed management.

(2) The board shall:

(a) administer the district's noxious weed program;

(b) establish management criteria for noxious weeds on all land within the district;

(c) make all reasonable efforts to develop and implement a noxious weed program covering all land within the district owned or administered by a federal agency.

7-22-2110. Administrative Hearing -- Appeals. (1) A person adversely affected by any notice, action, or order of the board may request an administrative hearing before the board. The board shall hold a hearing within 30 days of the request. Participants may be represented by legal counsel. The board shall make a record of the proceeding and enter its order and findings within 7 days after the hearing.

(2) An order of the board may be appealed to the commissioners within 30 days from the time the order is entered.

The commissioners shall hear such appeal within 30 days after the notice of appeal and shall render their order and findings within 7 days after such hearing. Participants may be represented by legal counsel.

(3) Within 30 days after the commissioners render their order and findings, the person adversely affected may file a petition in district court requesting that the order and findings of the commissioners be set aside or modified. The court may affirm, modify, or set aside the order complained of, in whole or in part.

7-22-2111. (Temporary) Liability Restrictions. A district, as defined in 7-22-2101, is liable for damages caused by its use of herbicides only for an act or omission that constitutes gross negligence. The provisions of 2-9-305 apply to board members, supervisors, and employees of a district. (Terminates July 1, 1995 -- sec. 7, Ch. 530, L. 1991.)

7-22-2112. (Temporary) Information on Herbicide Use. The district must provide information on protective clothing, health hazards, and proper application techniques to mixers, loaders, and applicators of herbicides and make information available for review by the public at the district office. (Terminates July 1, 1995 -- sec. 7, Ch. 530, L. 1991.)

7-22-2113 and 7-22-2114 reserved.

7-22-2115. Noxious Weeds and Seeds Declared Nuisance. Noxious weeds and the seed of any noxious weed are hereby declared a common nuisance.

7-22-2116. Unlawful to Permit Noxious Weeds to Propagate. It is unlawful for any person to permit any noxious weed to propagate or go to seed on his land, except that any person who adheres to the noxious weed management program of his district or who has entered into and is in compliance with a noxious weed management agreement is considered to be in compliance with this section.

7-22-2117. Violations. (1) Any person who in any manner interferes with the board or its authorized agent in carrying out the provisions of this part or who refuses to obey an order or notice of the board is guilty of a misdemeanor, and upon conviction thereof, he shall be fined not to exceed \$100 for the first offense and not less than \$100 or more than \$200 for each subsequent offense.

(2) All fines, bonds, and penalties collected under the provisions of this part, except those collected by a justice's court, shall be paid to the county treasurer of each county and placed by him to the credit of a fund to be known as the noxious weed fund.

7-22-2118 through 7-22-2120 reserved.

7-22-2121. **Weed Management Program.** (1) The noxious weed management program must be based on a plan approved by the board.

(2) The noxious weed management plan must:

(a) specify the goals and priorities of the program;

(b) review the distribution and abundance of each noxious weed species known to occur within the district and specify the locations of new infestations and areas particularly susceptible to new infestations;

(c) specify pesticide management goals and procedures, including but not limited to water quality protection, public and worker safety, equipment selection and maintenance, and pesticide selection, application, mixing, loading, storage, and disposal; and

(d) estimate the personnel, operations, and equipment costs of the proposed program.

(3) The board shall provide for the management of noxious weeds on all land or rights-of-way owned or controlled by a county or municipality within the confines of the district. It shall take particular precautions while managing the noxious weeds to preserve beneficial vegetation and wildlife habitat. Where at all possible, methods for such control shall include cultural, chemical, and biological methods.

(4) The board may establish special management zones within the district. The management criteria in such zones may be more or less stringent than the general management criteria for the district.

7-22-2122. **Repealed.** Sec. 32, Ch. 607, L. 1985.

7-22-2123. **Procedure in a Case of Noncompliance.** (1) Where complaint has been made or the board has reason to believe that noxious weeds described in this part are present upon a person's land within the district in violation of the law, that person must be notified by mail or telephone of the complaint and the board may request inspection of such land. The board or its authorized agent and the landowner or his representative shall inspect the land at an agreeable time, within 10 days of notification of the landowner. If after reasonable effort the board is unable to gain cooperation of the person, the board or its authorized agent may enter and inspect the land to determine if the complaint is valid.

(2) If noxious weeds are found, the board or supervisor shall notify the person or his representative and seek voluntary compliance with the district weed control program. If voluntary compliance is not possible, notice of noncompliance must be sent to the person by certified mail.

(3) The notice must specify:

(a) the basis for the determination of noncompliance;

(b) the geographic location of the area of noncompliance, by legal description or other reasonably identifiable description;

(c) measures to be undertaken in order to comply with the district's management criteria; '

(d) a reasonable period of time, not less than 10 days, in which compliance measures must be initiated; and

(e) the right of the person to request, within the time specified in subsection (3) (d), an administrative hearing as provided by 7-22-2110.

(4) A person is considered in compliance if he submits and the board accepts a proposal to undertake specified control measures and is in compliance for so long as he performs according to the terms of the proposal. If the measures proposed to be taken extend beyond the current growing season, the proposal and acceptance must be in writing.

(5) In accepting or rejecting a proposal, the board shall consider the economic impact on the person and his neighbors, practical biological and environmental limitations, and alternative control methods to be used.

7-22-2124. Destruction of Weeds by Board. (1) If corrective action is not taken and no proposal is made and accepted or no request for an administrative hearing is made within the time specified in the notice, the board may forthwith enter upon the person's land and institute appropriate control measures.

In such case the board shall submit a bill to the person, itemizing man-hours of labor, material, and equipment time, together with a penalty not exceeding 10% of the total cost incurred. Labor and equipment must be valued at the current rate paid for commercial management operations in the district. The bill must specify and order a payment due date of 30 days from the date the bill is sent.

(2) A copy of the bill must also be submitted by the board to the county clerk and recorder.

(3) If a person receiving an order to take corrective action requests an administrative hearing, the board may not institute control measures until the matter is finally resolved, except in case of an emergency. In such a case, the person is liable for costs as provided in subsection (1) only to the extent determined appropriate by the board, commissioners, or court that finally resolves the matter.

7-22-2125. Repealed. Sec. 32, Ch. 607, L. 1985.

7-22-2126. Embargo. The board may establish voluntary embargo programs to reduce the spread of noxious weeds within the district or the introduction of noxious weeds into the district.

7-22-2127. Repealed. Sec. 32, Ch. 607, L. 1985.

7-22-2128 and 7-22-2129 reserved.

7-22-2130. Weed District Supervisor Training. Within the limitations of available funds, the board shall ensure that the weed district supervisor obtains training to properly implement the noxious weed management program described in 7-22-2121. The department shall specify through rulemaking the level and type of training necessary to fulfill this requirement.

7-22-2131 through 7-22-2140 reserved.

7-22-2141. Noxious Weed Fund Authorized. (1) The commissioners of each county in this state shall create a noxious weed management fund, to be designated the "noxious weed fund".

(2) This fund shall be kept separate and distinct by the county treasurer.

7-22-2142. Sources of Money For Noxious Weed Fund. (1) The commissioners may create the noxious weed fund and provide sufficient money in the fund for the board to fulfill its duties, as specified in 7-22-2109, by:

(a) appropriating money from the general fund of the county;

(b) at any time fixed by law for levy and assessment of taxes, levying a tax not exceeding 2 mills on the dollar of total taxable valuation in the county. The tax levied under this subsection must be identified on the assessment as the tax that will be used for noxious weed control; and

(c) levying a tax in excess of 2 mills if authorized by a majority of the qualified electors voting in an election held for this purpose pursuant to 7-6-2531 through 7-6-2536.

(2) The proceeds of the noxious weed control tax must be used solely for the purpose of managing noxious weeds in the county and must be designated to the noxious weed fund.

(3) Any proceeds from work or chemical sales must revert to the noxious weed fund and must be available for reuse within that fiscal year or any subsequent year.

(4) The commissioners may accept any private, state, or federal gifts, grants, contracts, or other funds to aid in the management of noxious weeds within the district. These funds must be placed in the noxious weed fund.

7-22-2143. Determination of Cost of Weed Control Program. Based on the board's recommendations, the commissioners shall determine and fix the cost of the control of noxious weeds in the district, whether the same be performed by the individual landowners or by the board.

7-22-2144. Payment of Cost of Weed Control Program. The total cost of such control shall be paid from the noxious weed fund. The cost of controlling such weeds growing along the right-of-way of a state or federal highway shall, upon the presentation by the board of a verified account of the expenses incurred, be paid from the state highway fund in compliance with 7-14-2132 and any agreement between the board and the department of highways. Costs attributed to other lands within the district shall be assessed to and collected from the responsible person as set forth in 7-22-2116.

7-22-2145. Expenditures From Noxious Weed Fund. (1) The noxious weed fund must be expended by the commissioners at the time and in the manner as is recommended by the board to secure the control of noxious weeds.

(2) Warrants upon the fund must be drawn by the board. Warrants may not be drawn except upon claims duly itemized by the claimant, except payroll claims that must be itemized and certified by the board, and each claim must be presented to the commissioners for approval before the warrant is countersigned by the commissioners.

7-22-2146. Financial Assistance to Persons Responsible For Weed Control. (1) The commissioners, upon recommendation of the board, may establish cost-share programs with any person, specifying costs that may be paid from the noxious weed fund and costs that must be paid by the person. Cost-share programs may be established for special projects and for established management zones.

(2) (a) When under the terms of any voluntary agreement, whether entered into pursuant to 7-22-2123 or otherwise, or under

any cost-share program entered pursuant to this section a person incurs any obligation for materials or services provided by the board, the board shall submit a bill to the person, itemizing man-hours of labor, material, and equipment time. The bill must specify and order a payment due date not less than 30 days from the date the bill is sent.

(b) A copy of the bill must be submitted by the board to the county clerk and recorder. If the sum to be repaid by the person billed is not repaid on or before the date due, the county clerk and recorder shall certify the amount thereof, with the description of the land to be charged, and shall enter the sum on the assessment list as a special tax on the land, to be collected in the manner provided in 7-22-2148.

7-22-2147. Repealed. Sec. 32, Ch. 607, L. 1985.

7-22-2148. Tax Liability For Payment of Weed Control Expenses.

(1) The expenses referred to in 7-22-2124 shall be paid by the county out of the noxious weed fund, and unless the sum to be repaid by the person billed under 7-22-2124 is repaid on or before the date due, the county clerk shall certify the amount thereof, with the description of the land to be charged, and shall enter the same on the assessment list of the county as a special tax on the land. If the land for any reason is exempt from general taxation, the amount of such charge may be recovered by direct claim against the lessee and collected in the same manner as personal taxes. When such charges are collected, they shall be credited to the noxious weed fund.

(2) In determining what lands are included as land covered by the special tax and are described in the certificate of the county clerk, it is presumed that all work done upon any of the land of any one landowner is for the benefit of all of the land within the district belonging to the owner, together with the parcel upon which the work was done, and the amount certified becomes a tax upon the whole thereof.

7-22-2149. Responsibility For Assessments And Taxes For Weed Control Levied on Leased State Lands. The lessee of agricultural state land is responsible for assessments and taxes levied by the board of county commissioners for the district as provided in 77-6-114.

7-22-2150. Cooperation With State And Federal-Aid Programs. The board is empowered to cooperate with any state or federal-aid program that becomes available. Under such a plan of cooperation, the direction of the program shall be under the direct supervision of the board of the district in which the program operates.

7-22-2151. Cooperative Agreements. (1) Any state agency controlling land within a district, including the department of highways; the department of state lands; the department of fish, wildlife, and parks; the department of institutions; the department of natural resources and conservation; and the university system, shall enter into a written agreement with the board. The agreement must specify mutual responsibilities for noxious weed management on state-owned or state-controlled land within the district.

(2) The board and the governing body of each incorporated municipality within the district shall enter into a written agreement and shall cooperatively plan for the management of noxious weeds within the boundaries of the municipality. The board may implement management procedures described in the plan within the boundaries of the municipality for noxious weeds only. Control of nuisance weeds within the municipality remains the responsibility of the governing body of the municipality, as specified in 7-22-4101.

(3) A board may develop and carry out its noxious weed management program in cooperation with boards of other districts, with state and federal governments and their agencies, or with any person within the district. The board may enter into cooperative agreements with any of these parties.

7-22-2152. Revegetation of Rights-Of-Way And Disturbed Areas.

(1) Any state agency or local government unit approving a mine, major facility, transmission line, solid waste facility, highway, subdivision, or any other development resulting in significant disturbance of land within a district shall notify the board.

(2) Whenever any person or agency disturbs vegetation on an easement or right-of-way within a district by construction of a road, irrigation or drainage ditch, pipeline, transmission line, or other development, the board shall require that the disturbed areas be seeded, planted, or otherwise managed to reestablish a cover of beneficial plants.

(3) (a) The person or agency disturbing the land shall submit to the board a written plan specifying the methods to be used to accomplish revegetation. The plan must describe the time and method of seeding, fertilization practices, recommended plant species, use of weed-free seed, and the weed management procedures to be used.

(b) The plan is subject to approval by the board, which may require revisions to bring the revegetation plan into compliance with the district weed management plan. Upon approval by the board, the revegetation plan must be signed by the chairman of the board and the person or agency responsible for the disturbance and constitutes a binding agreement between the

board and such person or agency.

7-22-2153. Voluntary Agreements For Control of Noxious Weeds Along Roads. (1) Any person may voluntarily seek to enter into an agreement for the management of noxious weeds along a state or county highway or road bordering or running through his land. The supervisor may draft such an agreement upon the request of and in cooperation with the person; however, the agreement must, in the board's judgment, provide for effective weed management. The weed management agreement must be signed by the person and, upon approval of the board, by the chairman. An agreement involving a state highway right-of-way must also be signed by a representative of the department of highways.

(2) The agreement must contain a statement disclaiming any liability of the board and, if applicable, the department of highways for any injuries or losses suffered by the person in managing noxious weeds on the state or county highway right-of-way. The signed agreement transfers responsibility for managing noxious weeds on the specified section of right-of-way from the board to the person signing the agreement. If the board later finds that the person has failed to adhere to the agreement, the board shall issue an order informing the person that the agreement will be void and that responsibility for the management of noxious weeds on the right-of-way will revert to the board unless the person complies with the provisions of the agreement within a specified time period.

RULES
COUNTY NOXIOUS WEED LIST
Sub-Chapter 2
Designation of Noxious Weeds

4.5.201. Designation of Noxious Weeds. (1) The department designates certain exotic plants listed in these rules as statewide noxious weeds under the County Weed Control Act 7-22-2101 (5), MCA. All counties must implement management standards for these noxious weeds consistent with weed management criteria developed under 7-22-2109 (2)(b) of the Act. The department established three categories of the noxious weeds. (History: Sec. 7-22-2101 MCA; IMP, Sec. 7-22-2101 MCA; NEW 1986, p. 337, Eff. 3/14/86; AND, 1991 MAR p. 511, Eff. 4/26/91.)

4.5.202. Category 1. (1) Category 1 noxious weeds are weeds that are currently established and generally widespread in many counties of the state. Management criteria includes awareness and education, containment and suppression of existing infestations and prevention of new infestations. These weeds are capable of rapid spread and render land unfit or greatly limit beneficial uses.

(2) The following are designated as category 1 noxious weeds:

- (a) Canada Thistle (Cirsium arvense)
- (b) Field Bindweed (Convolvulus arvensis)
- (c) Whitetop or Hoary Cress (Cardaria draba)
- (d) Leafy Spurge (Euphorbia esula)
- (e) Russian Knapweed (Centaurea repens)
- (f) Spotted Knapweed (Centaurea maculosa)
- (g) Diffuse Knapweed (Centaurea diffusa)
- (h) Dalmatian Toadflax (Linaria dalmatica)
- (i) St. Johnswort (Hypericum perforatum).

(History: Sec. 7-22-2101 MCA; IMP, Sec. 7-22-2101 MCA; NEW 1986 MAR p. 337, Eff. 3/14/86; AND, 1991 MAR p. 511, Eff. 4/26/91.)

4.5.203. Category 2. (1) Category 2 noxious weeds have recently been introduced into the state or are rapidly spreading from their current infestation sites. These weeds are capable of rapid spread and invasion of lands, rendering lands unfit for beneficial uses. Management criteria includes awareness and education, monitoring and containment of known infestations and eradication where possible.

(2) The following are designated as category 2 noxious weeds:

(a) Dyers Woad (Isatis tinctoria)

(b) Purple Loosestrife or Lythrum (Lythrum salicaria, L. virgatum, and any hybrid crosses thereof).

(c) Sulfur (Erect) Cinquefoil (Potentilla recta)

(History: Sec. 7-22-2101 and 80-7-802 MCA; MP Sec. 7-22-2101 MCA: EW 1986 MAR p. 337, Eff. 3/14/86; AND, 1989 MAR p. 899, Eff. 7/14/89; AND 1991 MAR p. 511, Eff. 4/26/91.)

4.5.204 Category 3. (1) Category 3 noxious weeds have not been detected in the state or may be found only in small, scattered, localized infestations. Management criteria includes awareness and education, early detection and immediate action to eradicate infestations. These weeds are know pests in nearby states and are capable of rapid spread and render land unfit for beneficial uses.

(2) The following are designated as category 3 noxious weeds:

(a) Yellow Starthistle (Centaurea solstitialis)

(b) Common Crupina (Crupina vulgaris)

(c) Rush Skeletonweed (Chondrilla juncea)

(History: Sec. 80-7-802 MCA; IMP, Sec. 7-22-2102 MCA; NEW 1991 MAR p.511, Eff. 4/26/91.)

Part Cross References:

Weed Control - Department of Agriculture, Title 80, Chapter 7, Part 7.
Municipal Weed Control, 7-22-4101.
Noxious Weed Management Funding, Title 80, Chapter 7, Part 8.
Embargo against introduction of noxious weed seed from other state, 80-7-701.
General authority of county commissioners, 7-5-2101.
County officers - term of office, 7-4-2205.
Nuisance, Title 27, Chapter 30.
Classification of offenses, 45-1-201.
Department of State Lands, general powers and duties, Title 77, Chapter 1, Part 3.
Mining on State Lands, Title 77, Chapter 3.
Department of Fish, Wildlife and Parks, general powers and duties, Title 87, Chapter 1, Part 2.
Department of Highways, general powers and duties, Title 60, Chapter 2, Part 7.
Highways, acquisition and disposition of property, Title 60, Chapter 4.
Highway maintenance agreements with local government, 60-2-204.
Montana Environmental Protection Act, Title 75, Chapter 1.
Montana Solid Waste Management Act, Title 75, Chapter 10, Part 2.
County Taxation, Title 7, Chapter 6, Part 25.
Department of Institutions, general powers and duties, Title 53, Chapter 1, Part 2.
University system, Title 20, Chapter 5.
Department of Natural Resources and Conservation established 2-15-3301.
Major Facility Siting Act, Title 75, Chapter 20.
Subdivisions, Title 76, Chapter 2 and 3.
Coal mining, Title 82, Chapter 3.
Oil and gas conservation, Title 82, Chapter 11.
Hard rock mining impact, Title 90, Chapter 6, Part 3.
Role and duties of county clerk, 7-4-2611.
Employment of personnel by county commissioners. 7-5-2107.

5.1\weed\county1.wp

APPENDIX B

SYNOPSIS OF NOXIOUS WEED AND EXOTIC PLANT SPECIES

State Declared Noxious Weed Species

CATEGORY 1 noxious weeds are weeds that are currently established in many counties of the state. Management criteria for control of these weeds is necessary in all counties to contain or suppress existing infestations or to prevent, through eradication or other appropriate measures, new infestations of these weeds. All of these weeds render land unfit or greatly limit the beneficial uses (rule 4.5.202).

Canada thistle (*Cirsium arvense*)

Canada thistle is a perennial forb that reproduces by horizontal roots and seed. The stems are hollow and branch near the top. The wavy leaves are deeply cut with spiny #to smooth margins. The flower heads are numerous, small, compact, and vary from light lavender to rose-purple. Canada thistle grows in cultivated fields, meadows, pastures, and waste areas.

Dalmatian toadflax (*Linaria dalmatica*)

Dalmatian toadflax is a perennial forb that spreads by creeping rootstalk and seed. The plants are pale green and have very showy yellow flowers. The "spurred" flowers are tinged with orange and are about 1 inch long. The leaves are broad, heart-shaped, and clasp the stem. Dalmatian toadflax is an escaped ornamental that invades rangeland, and is difficult to control.

Field bindweed (*Convolvulus arvensis*)

Field bindweed, a perennial forb, has an extensive, deep root system. Stems are prostrate, 1-4 feet long with numerous arrowhead-shaped stem leaves which form dense tangled mats. The white to pinkish flowers are trumpet-shaped and bloom from late June to frost. The plant propagates by seed and rhizome. Seeds remain viable for up to 50 years making this persistent weed very difficult to control. It is largely a problem in cultivated fields and waste areas.

Leafy spurge (*Euphorbia esula*)

Leafy spurge is a perennial forb which reproduces by vigorous rootstalk and seeds. The small flowers are enclosed by a pair of yellowish-green, heart shaped bracts which have the appearance of flowers. Stems, leaves and flowers contain a milky sap called latex, which may cause severe rashes in humans. The weed forms dense patches and may be

the most persistent noxious weed in Montana. It has wide habitat suitability, prolific reproductive capabilities, strong competitive ability and is difficult to control, especially along water bodies. Although it is unpalatable to cattle, leafy spurge will be grazed by sheep and goats when they are confined to an infested area.

Russian knapweed (*Centaurea repens*)

Russian knapweed is a perennial forb that spreads by creeping rootstalk and seed. The plant has numerous branches that are tipped with a single, lavender thistle-like flower. The leaves are small and narrow with broken edges. The roots are dark brown and have a scaly appearance. Once, established, it will completely crowd out other vegetation. Livestock tend to avoid the weed because of its bitter, quinine-like taste. Horses, if forced to graze Russian knapweed, will develop nervous disorders.

Spotted knapweed (*Centaurea maculosa*)

Spotted knapweed is a short-lived perennial forb that reproduces by seed. The seeds germinate in the spring and fall, whenever growing conditions are favorable. The showy, purple flowers are held in spotted bracts. The alternate leaves have deep, narrow divisions and a rough, hairy appearance. It is very nutrient competitive, accelerating the decline of native vegetation.

Diffuse knapweed (*Centaurea diffusa*)

Diffuse knapweed is a biennial or short lived perennial forb. The tip of each branch has a single flower head. The flowers are usually white or sometimes pinkish. Bracts are yellowish green with a light brown margin. The upper part of each bract narrows into a stiff spine. It is an excellent competitor on dry sites.

St. Johnswort (*Hypericum perforatum*)

St. Johnswort is most commonly known as "goatweed". A perennial forb, it reproduces by seed and rootstock. The stems are smooth, branched and woody at the base. The opposite leaves have small glandular dots. Flowers are orange-yellow with 5 petals. Goatweed is found in meadows, dry pastures, rangelands, neglected fields, and along the roadside. Goatweed causes photo-sensitive reactions in livestock and should be regarded as a poisonous plant.

Sulfur cinquefoil (*Potentilla recta*)

Sulfur cinquefoil is also referred to as "upright" or "rough fruited" cinquefoil. It is a perennial forb and reproduces by seed. The stems are hairy and rigid. The leaves are alternate and palmately divided with 5 to 7 coarsely toothed leaflets. The clustered flowers are a sulfur yellow. Sulfur cinquefoil is difficult to distinguish from several other native cinquefoils.

Whitetop (*Cardaria draba*)

Whitetop, also known as "hoary cress", is a perennial forb which reproduces by seeds and creeping roots. The dense flowers give the plant a white, flat-topped appearance. The leaves are grayish-green. The upper leaves clasp the stem.

CATEGORY 2 noxious weeds are weeds that have not been detected in or have recently been introduced into the State of Montana. These weeds have the potential for rapid spread and invasion of lands, thereby rendering them unfit for beneficial uses. County planning to prevent the spread or introduction of these weeds is necessary. Management criteria for detection and immediate action to eradicate or contain these weeds is necessary in all counties (rule 4.5.203).

Dyers woad (*Isatis tinctoria*)

Dyers woad is a perennial, biennial or annual forb reproducing by seeds and from roots. The plants have a smooth, bluish-green color. The lower leaves clasp the stem with ear-like projections. The yellow flowers are very small and form a flat-topped inflorescence. The seed pods are winged like a maple seed and turn black when mature. The purplish-black seed pods have been used for their dye extract, thus the name dyers woad.

Purple loosestrife (*Lythrum salicaria*)

Purple loosestrife is an aggressive perennial plant that is capable of invading our wetlands and replacing valuable wetland plants. Its dense root mats choke waterways, eliminating food and shelter for wildlife. It grows from underground roots, sprouts from broken off plant parts, and is a prolific seed producer. The flowers have a purple-magenta color. The leaves are linear shaped with smooth edges and are attached directly to a four-sided stem. Purple loosestrife is often confused with Blazing Star, Fireweed, or Blue Vervain.

CATEGORY 3 noxious weeds have not been detected in the state or may be found only in small, scattered localized infestations. Management criteria includes awareness and education, early detection and immediate action to eradicate infestations. These weeds are known pests in nearby states and are capable of rapid spread and render land unfit for beneficial uses.

Common crupina (*Crupina vulgaris*)

Common crupina is a annual that reproduces by seed. It is closely related to the knapweed species. The leaves are large, thick, and dark green with stiff hairs that feel sticky. The plant produces 5 to 100 flower heads with a lavender to purple color, which from a distance could be confused with spotted knapweed. It is generally found on well-drained, rocky to silt loam soils in pastures or rangeland. Cattle will not feed on common crupina plants. The species is competitive and forms solid stands reducing forage production and range carrying capacity. Infestations start in disturbed sites with sparse vegetation. Common crupina is not believed to be in Montana at this time.

Rush skeletonweed (*Chondrilla juncea*)

Rush skeletonweed, a member of the sunflower family, is a perennial 1-4 feet tall with a deep, extensive root system. It branches off of several main stems which have downward bent, coarse hairs on the lower 4-6 inches. Leaves occur at the base and resemble those of dandelions. Yellow flowering heads are small and scattered on branches. Flowering and seed production occurs from late summer through frost. This species generally inhabits well-drained, light-textures soils along roadsides, in rangelands, grain fields and pastures. Rush skeletonweed occupies large acreage in Oregon, Washington, Idaho and California. This species should not be confused with the rangeland native, skeletonweed, (*Lygodesmia juncea*) which has pink flowers.

Yellow star-thistle (*Centaurea solstitialis*)

Yellow star-thistle is an annual or, occasionally, biennial that reproduces by seed. The leaves are alternate and have slender, pointed lobes. The upper leaves are smaller and narrow with sharp, spiny tips. Leaf bases extend down the plant stem giving a winged effect. The flowers heads have sharp stiff spines on either side with one flower per branch. The flowers are yellow, long and tubular. It is toxic to horses and when eaten in quantity, results in a neurological disorder which over time causes death. The chewing disease associated with it comes from the spikes on the flower.

Other Watch Species

Eurasian watermilfoil (*Myriophyllum spicatum*)

Eurasian watermilfoil is an aggressive aquatic weed which is flourishing in Washington and Idaho. It resembles native milfoils with its lacy leaves and reproduces by seed and vegetative propagation. Water-milfoil roots in shallow streams and lakes in up to 15 feet of water forming extremely dense mats. It chokes waterways and irrigation ditches and depletes oxygen and therefore is a serious problem for recreationists, fisheries and landowners. The species appears to have a wide tolerance to severe environmental conditions and can resprout from seemingly dried out stems and roots. Thus it is easily spread by boats and trailers that carry fragments from infested waters to clean waters.

County Declared Noxious Weed Species

In addition to the state-declared noxious weed species specified in the County Noxious Weed Control Act, individual counties may also declare additional species as noxious and mandate control actions through their County Noxious Weed Control Management Plans. The following list summarizes these additional species for the counties within the administrative boundaries of FWP Region One.

Weed Species	Scientific Name	County (-ies)
Common Toadflax	<i>Linaria vulgaris</i>	Lake
Hound's Tongue	<i>Cynoglossum officinale</i>	Lake, Lincoln
Kochia	<i>Kochia scoparia</i>	Lincoln
Meadow Hawkweed	<i>Hieracium pratense</i>	Lincoln
Meadow Salsify	<i>Tragopogon pratensis</i>	Lake
Musk Thistle	<i>Carduus nutans</i>	Lake, Lincoln
Orange Hawkweed	<i>Hieracium aurantiacum</i>	Flathead, Lincoln
Oxeye Daisy	<i>Heliopsis helianthoides</i>	Lake
Poison Hemlock	<i>Conium maculatum</i>	Lincoln
Purple Mustard	<i>Chorispora tenella</i>	Lake
Scentless Chamomile	<i>Anthemis arvensis</i>	Sanders
Sheep Sorrel	<i>Rumex acetosella</i>	Lake

State Park Exotic Plant Categories

Category 1: Invasive and aggressive; capable of displacing native species; typically noxious to native plant communities and to neighboring agricultural lands; spread can be rapid from localized populations; elimination unlikely once populations are well established, but control can be effective in limiting spread and in elimination of localized or recent infestations; monitoring of these species should be frequent and regular.

<i>Cardaria draba</i>	whitetop
<i>Centaurea diffusa</i>	diffuse knapweed
<i>Centaurea maculosa</i>	spotted knapweed
<i>Centaurea repens</i>	Russian knapweed
<i>Centaurea solstitialis</i>	yellow star-thistle
<i>Chondrilla juncea</i>	rush skeletonweed
<i>Chrysanthemum leucanthemum</i>	oxeye daisy
<i>Crupina vulgaris</i>	common crupina
<i>Euphorbia esula</i>	leafy spurge
<i>Hypericum perforatum</i>	St. Johnswort
<i>Isatis tinctoria</i>	Dyars Woad
<i>Linaria dalmatica</i>	Dalmation toadflax
<i>Senecio jacobaea</i>	tansy ragwort

Category 2: Invasive but less aggressive; less capable of displacing native species; undesirable but not usually noxious; populations typically localized with gradual spread; control actions have a high probability of eliminating or reducing treated populations; monitoring should document effects of control or inaction.

<i>Acer platanoides</i>	Norway maple
<i>Aegopodium podagraria</i>	bishops weed
<i>Caragana arborescens</i>	siberian pea
<i>Centaurea Montana</i>	mountain bluet
<i>Convolvulus arvensis</i>	field bindweed
<i>Cynoglossum officinale</i>	hounds tongue
<i>Dianthus barbatus</i>	sweet william
<i>Digitalis purpurea</i>	foxglove
<i>Hieracium aurantiacum</i>	orange hawkweed
<i>Tanacetum vulgare</i>	common tansy
<i>Verbascum thapsus</i>	common mullein

Category 3: Invasive and aggressive; capable of displacing native species; undesirable; widespread distribution; control actions have low probability of reducing, containing or eliminating treated populations, but actions may be taken to discourage dispersal into uncolonized areas; monitoring may be beneficial but will have lower priority than Category 1 or 2 species.

<i>Agropyron repens</i>	Quackgrass
<i>Agrostis alba</i> var. <i>alba</i>	redtop
<i>Bromus inermis</i> var. <i>inermis</i>	smooth brome
<i>Bromus tectorum</i>	cheatgrass
<i>Cirsium arvense</i>	Canada thistle
<i>Cirsium vulgare</i>	bull thistle
<i>Linaria vulgaris</i>	butter and eggs
<i>Medicago lupulina</i>	black medic
<i>Medicago sativa</i>	alfalfa
<i>Melilotus alba</i>	white sweet clover
<i>Melilotus officinalis</i>	yellow sweet clover
<i>Phleum pratense</i>	common timothy
<i>Poa compressa</i>	Canada bluegrass*
<i>Poa pratensis</i>	Kentucky bluegrass*
<i>Sonchus arvensis</i>	field sow-thistle
<i>Sonchus asper</i>	prickly sow-thistle

*May have utilitarian value in Development Zones

Category 4: Less invasive and less aggressive; less capable of displacing native species; distribution variable but typically localized; relatively innocuous exotics for which limited control efforts are foreseen and only if the control actions would be less deleterious to the environment than continued presence of the exotic species; limited monitoring actions may be undertaken.

<i>Agropyron cristatum</i>	crested wheatgrass
<i>Alyssum alyssoides</i>	pale alyssum
<i>Amaranthus graecizans</i>	prostate pigweed
<i>Amaranthus retroflexus</i>	rough pigweed
<i>Ambrosia artemisiifolia</i>	annual ragweed
<i>Anthemis tinctoria</i>	golden marguerite
<i>Arabis glabra</i>	tower mustard
<i>Aremaroa serpyllifolia</i>	sandwort
<i>Artemisia absinthium</i>	absinth wormwood
<i>Asparagus officinalis</i>	asparagus

<i>Bellis perennis</i>	English daisy
<i>Brassica juncea</i>	Chinese mustard
<i>Brassica kaber</i>	wild mustard
<i>Bromus secalinus</i>	rye-brome
<i>Camelina microcarpa</i>	little-pod false flax
<i>Campanula glomerata</i>	clustered bellflower
<i>Campanula rapunculoides</i>	purple bell
<i>Capsella bursa-pastoris</i>	shepherds purse
<i>Carum carvi</i>	caraway
<i>Centaurea pratensis</i>	meadow knapweed
<i>Cerastium vulgatum</i>	big chickweed
<i>Chenopodium album</i>	lambsquarter
<i>Chenopodium botrys</i>	Jerusalem oak
<i>Crepis tectorum</i>	hawksbeard
<i>Dactylis glomerata</i>	orchard grass
<i>Diploaxis muralis</i>	wall rocket
<i>Echinocloa crusgalli</i>	barnyard grass
<i>Elaeagnus angustifolia</i>	Russian olive
<i>Erodium cicutarium</i>	storks-bill
<i>Erysimum cheiranthoides</i>	treacle mustard
<i>Filago arvensis</i>	filago
<i>Galium verum</i>	yellow bedstraw
<i>Geranium carolinianum</i>	geranium
<i>Lactuca serriola</i>	prickly lettuce
<i>Lappula echinata</i>	bristly stickseed
<i>Lathyrus latifolius</i>	perennial peavine
<i>Lepidium densiflorum</i>	common pepperwood
<i>Linum usitatissimum</i>	flax
<i>Lupinus polyphyllus</i> var. <i>regallis</i>	russell lupine
<i>Lychnis alba</i>	white campion
<i>Lysimachia nummularia</i>	creeping jenny
<i>Matricaria matricarioides</i>	pineapple weed
<i>Mentha spicata</i>	spearmint
<i>Nepeta cataria</i>	catnip
<i>Panicum capillare</i>	common witchgrass

<i>Pastinaca sativa</i>	parsnip
<i>Phalaris arundinacea</i>	reed canarygrass
<i>Picea pungens</i> Engelm.	Blue Spruce
<i>Pilea nummularifolia</i>	creeping charley
<i>Pinus sylvestris</i> L.	Scotch (or Scots) pine
<i>Plantago lanceolata</i>	English plantain
<i>Plantago major</i>	common plantain
<i>Poa annua</i>	annual bluegrass
<i>Polygonum aviculare</i>	dooryard knotweed
<i>Polygonum cuspidatum</i>	knotweed
<i>Potentilla argentea</i>	silvery cinquefoil
<i>Potentilla norvegica</i>	Norwegian cinquefoil
<i>Potentilla recta</i>	cinquefoil
<i>Prunella vulgaris</i> var. <i>vulgaris</i>	self-heal
<i>Ranunculus repens</i>	creeping buttercup
<i>Rubus laciniatus</i>	evergreen blackberry
<i>Rumex acetosella</i>	sheep sorrel
<i>Rumex crispus</i>	curly dock
<i>Salsola kali</i>	Russian thistle
<i>Saponaria officinalis</i>	bouncing bet
<i>Senecio vulgaris</i>	old-man-in-the-spring
<i>Silene cserei</i>	campion
<i>Silene cucubalis</i>	bladder campion
<i>Sisymbrium altissimum</i>	tumble mustard
<i>Solanum dulcamara</i>	bittersweet
<i>Solanum sarrachoides</i>	hairy nightshade
<i>Spergularia rubra</i>	red sandspurry
<i>Stellaria media</i>	chickweed
<i>Taraxacum laevigatum</i>	red-fruited dandelion
<i>Taraxacum officinale</i>	common dandelion
<i>Thlaspi arvense</i>	pennycress
<i>Tragopogon dubius</i>	yellow salsify
<i>Tragopogon pratensis</i>	meadow salsify
<i>Trifolium agrarium</i>	yellow clover
<i>Trifolium hybridum</i>	alsike clover
<i>Trifolium pratense</i>	red clover

Trifolium procumbens
Trifolium repens
Triticum aestivum
Verbascum blattaria
Verbena bracteata

hop clover
Dutch clover
wheat
moth mullein
bracted vervain

Veronica chamaedrys
Veronica officinalis
Vinca minor

germander speedwell
Pauls betony
periwinkle

Category 5: Least invasive and least aggressive; not likely to displace native species; distribution may be localized to historic landscapes or developed areas; species may be retained for cultural value at specific historic sites identified in plans; sites should be monitored for indication of spread beyond designated areas.

Aquilegia x hybrida
Chrysanthemum coccineum
Chrysanthemum maximum
Convallaria majalis
Iris sp.

columbine hybrids
painted daisy
shasta daisy
lily-of-the-valley
ornamental iris

Malus sp.
Narcissus sp.
Papavera orientalis
Prunus sp.

apple
daffodil
oriental poppy
cherry

Pyrus sp.
Sorbus aucuparia
Syringa vulgaris

pear
European mountain ash
lilac

APPENDIX C

LAWS AND POLICIES RELATING TO THE MANAGEMENT OF NOXIOUS WEEDS

Laws and Policies Relating to the Management of Noxious Weeds

State and federal laws, policies, and programs that affect FWP Region One's activities include:

- Creating a VISION for the Future of Montana's Department of Fish, Wildlife & Parks (1992)
- Weed Control Program for Lands Managed by the Montana Department of Fish, Wildlife and Parks (1983)
- Montana Pesticides Act (80-8-801 *et seq.*, MCA)
- Montana Weed Control Act (80-7-701 *et seq.*, MCA)
- Montana Noxious Weed Trust Fund Act of 1985 (80-7-801 *et seq.*, MCA) as amended 1991 and Rules
- Montana Water Quality Act (75-5-101 *et seq.*, MCA)
- Montana Agricultural Chemical Ground Water Protection Act of 1989 (80-15-100 *et seq.*, MCA)
- Montana Environmental Policy Act (MEPA, 75-1-101 *et seq.*, MCA)
- County Weed Boards for Flathead, Lake, Lincoln, and Sanders counties; 5-Year Plans
- Montana Department of Agriculture Chemical Use Reporting Procedures
- Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) P.L. 92-516; CFR 40-171
- USDA Forest Service - Flathead, Kootenai, and Lolo National Forests
- USDI Bureau of Land Management - Missoula Division
- US Environmental Protection Agency

APPENDIX D

BIOLOGICAL CONTROL AGENTS AVAILABLE IN MONTANA

Biological Control Agents Available in Montana

Host	Control Agent	Action
Knapweeds	<i>Agapeta zoegana</i>	root mining moth
	<i>Chaetorellia acrolophi</i>	seed head fly
	<i>Cyphocleonus achates</i>	root weevil
	<i>Larinus minutus</i>	seed head weevil
	<i>Larinus obtusus</i>	seed head beetle
	<i>Metzneria paucipunctella</i>	flowerette and seed moth
	<i>Pelochrista medullana</i>	root mining moth
	<i>Pterolonche inspersa</i>	root moth
	<i>Sclerotinia spp.</i>	fungus
	<i>Sphenoptera jugoslavica</i>	root mining beetle
	<i>Subanguinia picridis</i>	leaf gall nematode
	<i>Terellia virens</i>	seed head fly
	<i>Urophora affinis</i>	seed head gallfly larvae
	<i>Urophora quadrifasciata</i>	seed head fly
Leafy Spurge	<i>Aphthona nigriscutis</i>	root and leaf flea beetle
	<i>Aphthona cyparissiae</i>	root and leaf flea beetle
	<i>Aphthona flava</i>	root and leaf flea beetle
	<i>Aphthona czwalinae</i>	root and leaf flea beetle
	<i>Bayeria spp.</i>	shoot tip gall midge
	<i>Hyles euphorbiae</i>	defoliating hawk moth larvae
	<i>Oberea erythrocephala</i>	stem and root beetle larvae/adults
Canada Thistle	<i>Spurgia esulae</i>	shoot tip gall midge
	<i>Ceutorhynchus litura</i>	stem mining weevil
	<i>Urophora cardui</i>	stem and shoot gallfly
Dalmatian Toadflax	<i>Larinus planus</i>	seedhead weevil
	<i>Calophasia lunula</i>	defoliating larvae/moth
Musk Thistle	<i>Rhinocyllus conicus</i>	seed head weevil
	<i>Trichosiracalus horridus</i>	rosette weevil
St. Johnswort	<i>Chrysolina quadrigemini</i>	defoliating beetle

Biological Control Agents Released on Department of Fish, Wildlife and Parks Properties in Region One

Host	Control Agent	Release Site
Knapweeds	<i>Agapeta zoegana</i>	Ninepipe WMA*, Flatiron Ridge FAS
	<i>Cyphocieonus achates</i>	Ninepipe WMA*
	<i>Urophora affinis</i>	Mt. Silcox WMA, Ninepipe WMA*, Flatiron Ridge FAS, Flathead Lk, Lone Pine, Thompsn Fls ST PKS, Old Steel Bridge FAS
	<i>Urophora quadrifasciata</i>	Mt. Silcox WMA, Ninepipe WMA*, Flatiron Ridge FAS, Flathead Lk, Thompson Falls State Parks
Leafy Spurge	<i>Aphthona nigriscutis</i>	Kokanee Bend FAS
	<i>Aphthona flava</i>	Kokanee Bend FAS
	<i>Hyles euphorbiae</i>	Kokanee Bend FAS
	<i>Oberea erythrocephala</i>	Kokanee Bend FAS
Canada Thistle	<i>Ceutorhynchus litura</i>	Ninepipe WMA*
	<i>Urophora cardui</i>	Ninepipe WMA*
	<i>Larinus planus</i>	Ninepipe WMA*
	<i>Rhinocyllus conicus</i>	Ninepipe WMA*
	<i>Trichosirocalus horridus</i>	Ninepipe WMA*
Dalmatian Toadflax	<i>Calophasia lunula</i>	Ninepipe WMA*
	<i>Brachyteronius palicarius</i>	Ninepipe WMA*
Musk Thistle	<i>Rhinocyllus conicus</i>	Ninepipe WMA*
	<i>Trichosiracalus horridus</i>	Ninepipe WMA*
Purple Loosestrife	<i>Galerucella californiensis</i>	Ninepipe WMA*
St. Johnswort	<i>Agrilus hyperici</i>	Ninepipe WMA*
	<i>Aplocera plagiata</i>	Ninepipe WMA*
	<i>Chrysolina hyperici</i>	Ninepipe WMA*
	<i>Chrysolina quadrigemini</i>	Ninepipe WMA*, Lone Pine State Park

*Released at National Bison Range, but established on Ninepipe WMA

APPENDIX E

HERBICIDE EMERGENCY RESPONSE PLAN

Herbicide Emergency Response Plan

The improper use or accidental release of an herbicide may pose serious health or environmental hazards. Accidental spills and releases must be managed quickly and efficiently to protect human health and the environment. Additional information on the handling and use of herbicides can be found in the Montana Pesticide Act (80.8.101 MCA) and its Administrative Rules (ARM 4.10.101).

During herbicide emergencies, the first priority is the safety of personnel involved in the accident. The next priority is spill containment and clean-up to minimize environmental contamination. This plan outlines field response and reporting procedures to follow in the event of an herbicide spill. This plan applies to all FWP personnel and contractors when an accident occurs involving the transport, application, use or handling of an herbicide.

IN THE EVENT OF A SPILL:

- 1. Administer First Aid to Injured or Contaminated Persons**
- 2. Identify Type of Pesticide Released**
- 3. Notify the Appropriate Authorities**
- 4. Quarantine the Area**
- 5. Contain the Spill if Possible**
- 6. Complete Pesticide Emergency Response Record**
- 7. Develop a Clean-Up Plan Where Appropriate**

1. Administer First Aid to Injured Persons

Of greatest importance is the immediate threat to human health and the aid of someone injured during an herbicide spill. Any injured parties should be removed from the contaminated area immediately. Contaminated clothing should be removed. All persons should avoid direct contact with the spilled material until it is positively identified.

If the released chemical is known, labels and Material Safety Data Sheets can be consulted for the appropriate procedures for decontamination and administering first aid. For most herbicides, washing with water and detergent is the best method. Initial first aid for eye contact usually includes rinsing with eye-safe solutions (e.g. contact lens saline solution) or water. Injured parties, should be transported to medical facilities as soon as possible, with the name of the released material, label, and material safety data sheet accompanying the injured person.

2. Identify Spilled Material

To administer the most appropriate emergency response, the spilled herbicide must be accurately identified. Herbicide label information and Material Safety Data Sheets (MSDS) should accompany all DFWP personnel and their contractors if an herbicide is handled, transported, or applied. In addition, MSDS's should also be kept on file at any location where herbicides are stored. MSDS's describe appropriate response procedures for administering emergency aid to affected parties and appropriate protective clothing for containing a spill. The transporter and applicator of the herbicide must carry records describing mixtures of all herbicides applied.

3. Notify the Authorities

Herbicides can pose serious threats to human health and the environment and should be handled with extreme care. All accidental herbicide releases must be reported to the Montana Department of Agriculture within 48 hours. The written report must include the time of the incident, its location, herbicide name, type of formulation, method of application, and the FWP contact for the project. The report should also name all parties involved in the incident, including the names and addresses of individuals who may be affected by the spill. If the spilled herbicide is classified as either extremely toxic or highly toxic, the Department of Agriculture must be notified immediately. The following authorities may be contacted:

FWP: Regional Parks Maintenance Supervisor	752-5501
Local Emergency Response:	911
County Sheriff's Office:	Flathead 756-5668
	Lake 883-2121
	Lincoln 293-4112
	Sanders 827-3584
Montana Disaster & Emergency Services:	(406) 444-6911
National Response Center:	(800) 424-8802
Montana Department of Agriculture:	(406) 444-3144
Montana Poison Control Center:	(800) 525-5042
Chemtrec:	(800) 424-9300

The Chemical Manufacturer's Association maintains this 24 hour hotline to provide information about chemicals and their related health and environmental hazards for fire and police crews responding to chemical accidents and spills.

4. Quarantine the Area

In the event of a spill, restrict access to the site, using physical barriers such as emergency tape, flagging, or signs. The area should be secured to prevent entry of unauthorized personnel into the spill area. Only authorized persons wearing the appropriate protective clothing should be permitted into the area.

5. Contain the Spill

Once the spilled material is positively identified, it must be contained as soon as possible. Technical labels and MSDS's provide information on protective clothing requirements for clean-up personnel. Generally, it will be clothing similar to the protective clothing required for application.

Put on all necessary protective clothing, including respirators, before approaching the spill from an upwind direction. Avoid inhaling fumes, vapors, and dust from the spill. Smoking is not allowed in a spill area. Examine the area and determine an effective method to contain the spill. Any person attempting to contain a spilled herbicide should also follow these guidelines:

- minimize human contact with the spill, using mechanized equipment if possible
- avoid raising dust
- avoid diluting material with water (except for misting dry substances)
- treat all absorbent material used during containment as a hazardous waste
- remove and dispose of all contaminated soil as a hazardous waste
- wash and properly decontaminate hands and face prior to consuming any food

Potential spill containment methods include:

Liquid Spills

- create small collection pools for runoff
- create dikes to impound runoff
- cover spill material with approximately double its volume in absorbent material (hydrated lime, saw dust, kitty litter)
- transfer absorbent material onto an impermeable barrier (e.g. tarp)
- divert spilled material away from open waters
- monitor and plug leaks in containment structures

Dry Spills

- cover with a plastic tarp and secure edges of the tarp
- spray fine mist to minimize dust
- shovel material into clearly marked plastic bags or drums and seal

6. Record Herbicide Spill

An herbicide spill or accident can have long lasting effects beyond immediate health and environmental threats. It is important to document all circumstances and events pertaining to the accident and/or spill. Important information includes the name and type of herbicide (include MSDS sheets), the names of any injured or contaminated persons, amount spilled or released, spill location, and whether chemicals were discharged into a water body. Below is a form to record all appropriate information.

HERBICIDE SPILL EMERGENCY RESPONSE FORM

Complete all appropriate items and add additional notes to document special conditions or unusual circumstances. Take photos when possible to supplement notes.

1. Immediately contact the appropriate response agencies.
2. Were there injuries related to the spill? ☐yes ☐no
Were any individuals contaminated with herbicide? ☐yes ☐no
Was aid provided to affected individuals? ☐yes ☐no
Was transportation to a medical facility required? ☐yes ☐no
3. Was a vehicle accident involved? ☐yes ☐no
4. What materials/herbicides were involved? (include MSDS)

5. Estimate amount of herbicide or herbicide mixture spilled in gallons.
Herbicide: _____ gals Mixture: _____ gals
Herbicide: _____ gals Mixture: _____ gals
Herbicide: _____ gals Mixture: _____ gals
6. Location of Incident (name of site): _____
____ 1/4 ____ 1/4 ____ Sec ____ T ____ R Highway/Road: _____
Time of Incident: _____ Date: _____

Provide a description of the site (water bodies, vegetation, landmarks)

Distance to picnic/play campground area: _____
nearest well: _____
surface water: _____
7. Is there a danger of surface water contamination? ☐yes ☐no
Has an herbicide already entered the water body? ☐yes ☐no
Name of Stream/Lake _____
8. Is there a danger of ground water contamination? ☐yes ☐no
Depth to groundwater: _____
Elevation above nearest surface water: _____

HERBICIDE SPILL EMERGENCY RESPONSE FORM, page 2

9. General Soil Description (check one for surface and subsurface soil)

- Surface ☐ Dark Colored, Organic, Rock Free
 ☐ Medium Textured
 ☐ Light Colored, Rocky
- Subsoil ☐ Dark Colored, Organic, Rock Free
 ☐ Medium Textured
 ☐ Light Colored, Rocky

10. List Agencies Notified

FWP Authority _____	When _____
Local Authority _____	When _____
State Authority _____	When _____
Federal Authority _____	When _____
Other Authority _____	When _____

11. Site Sketch

12. Describe Incident: _____

13. Public Contact/Comment During Incident: _____

14. Name: _____ Date: _____

CHEMICAL EMERGENCY SPILL KIT

All FWP personnel and contractors applying or transporting herbicides must carry the following safety items in the vehicle.

- at least one pair of neoprene or plastic gloves for each crew member present
- at least one pair of rubber, neoprene, or plastic material boots
- at least one pair of unvented goggles for each crew member present
- at least one approved respirator for applied pesticide
- at least one pair of coveralls or disposable coveralls (Tyvek) for each crew member
- absorbent material (kitty litter, floor dry)
- heavy duty plastic garbage sacks
- plastic tarp
- shovel
- dustpan and shop brush
- one pint liquid detergent
- portable eye wash kit
- five-pound ABC-type fire extinguisher
- first aid kit
- safety tape/flagging
- Material Safety Data Sheets and technical label information for each herbicide handled and used
- Herbicide Spill Response Plan

APPENDIX F

SENSITIVE SPECIES OCCURRING IN THE VICINITY OF MONTANA FISH, WILDLIFE & PARKS PROPERTIES IN REGION ONE

Sensitive Species Occurring in the Vicinity of Department of Fish, Wildlife and Parks Region One Properties¹

Site Name	Species Occurrence and Status
Flthd Lk WHPA/Bird I	Bald Eagle (<i>Haliaeetus leucocephalus</i>); nesting site and territory; listed as endangered under the Endangered Species Act.
Kootenai Falls WMA ²	Bald Eagle (<i>Haliaeetus leucocephalus</i>); listed as endangered under ESA. No nesting on WMA. Use trees along Kootenai River shoreline for perching and roosting. No critical habitat exists on the WMA.
Kootenai Falls WMA ²	Grizzly Bear (<i>Ursus arctos</i>); listed as threatened under the ESA. The WMA may be in a corridor of travel between the Cabinet and Yaak portions of the Cabinet/Yaak Ecosystem. No critical habitat exists on the WMA.
Mount Silcox WMA ²	Bald Eagle (<i>Haliaeetus leucocephalus</i>); nesting and wintering territory; listed as endangered under the Endangered Species Act.
Mount Silcox WMA ²	Twin Clover (<i>Trifolium latifolium</i>); possibly first record in Montana, not globally endangered.
Mount Silcox WMA ²	Rhombic-petaled Clarkia (<i>Clarkia rhomboidea</i>); known from less than five sites in Montana, not globally endangered.
Ninepipe WMA ²	Bald Eagle (<i>Haliaeetus leucocephalus</i>); nesting and wintering territory; listed as endangered under the Endangered Species Act.
Ninepipe WMA ²	Columbia Water-meal (<i>Wolffia columbiana</i>); Post canal ponds; sensitive species.
Roundhorn WMA ²	Bald Eagle (<i>Haliaeetus leucocephalus</i>); nesting and wintering territory; listed as endangered under the Endangered Species Act.
Thompson Ch of Lakes	Bald Eagle (<i>Haliaeetus leucocephalus</i>); nesting site and territory; listed as endangered under the Endangered Species Act.
Thompson Ch of Lakes	Common Loon (<i>Gavia immer</i>); nesting site and territory; sensitive species of restricted range in Montana.
Thompson Ch of Lakes	Great Blue Heron Rookery; nesting site and territory (30 nests active in 1986, exact location unknown; sensitive species.
Flathead Lake SP ³	Bald Eagle (<i>Haliaeetus leucocephalus</i>); nesting site and territory; listed as endangered under the Endangered Species Act.
Flathead Lake SP ³	Bull Trout (<i>Salvelinus confluentus</i>); sensitive species.
Flathead Lake SP ³	Slender Sedge (<i>Carex tinctoria</i>); sensitive species observed at Yellow Bay.
Flathead Lake SP ³	Giant Helleborine (<i>Epipactis gigantea</i>); sensitive species observed at Yellow Bay, possibly extirpated by 1984.

Flathead Lake SP ³	Adder's Tongue (<i>Ophioglossum vulgatum</i>); sensitive species observed at Yellow Bay, possibly extirpated by 1984.
Flthd Lk SP/W Hrse Is	Spalding Campion (<i>Silene spaldingii</i>); restricted range or rare throughout its range.
Flthd Lk SP/W Hrse Is	Bald Eagle (<i>Haliaeetus leucocephalus</i>); nesting site and territory; listed as endangered under the Endangered Species Act.
Lake Mary Ronan SP ³	Bald Eagle (<i>Haliaeetus leucocephalus</i>); territory; listed as endangered under the Endangered Species Act.
Whitefish Lake SP ³	Small yellow Lady's Slipper (<i>Cypripedium calceolus var parviflorum</i>); sensitive species.
Flathead Lake SP ³	Bull trout (<i>Salvelinus confluentus</i>); listed as warranted but precluded.
Flathead Lake SP ³	Westslope cutthroat (<i>Oncorhynchus clarki lewisi</i>); Montana species of special concern.
Flathead River	Bull trout (<i>Salvelinus confluentus</i>); listed as warranted but precluded.
Flathead River	Westslope cutthroat (<i>Oncorhynchus clarki lewisi</i>); Montana species of special concern.
Clk Fk, Flatiron Ridge	Bull trout (<i>Salvelinus confluentus</i>); listed as warranted but precluded.
Swan River	Westslope cutthroat (<i>Oncorhynchus clarki lewisi</i>); Montana species of special concern.
Swan River	Bull trout (<i>Salvelinus confluentus</i>); listed as warranted but precluded.
Whitefish River	Westslope cutthroat (<i>Oncorhynchus clarki lewisi</i>); Montana species of special concern.
Whitefish River	Bull trout (<i>Salvelinus confluentus</i>); listed as warranted but precluded.
Ashley Lake	Westslope cutthroat (<i>Oncorhynchus clarki lewisi</i>); Montana species of special concern.
Thompson Ch of Lakes	Westslope cutthroat (<i>Oncorhynchus clarki lewisi</i>); Montana species of special concern.
Flatiron	Westslope cutthroat (<i>Oncorhynchus clarki lewisi</i>); Montana species of special concern.
Kootenai Drainage	Redband trout (<i>Oncorhynchus mykiss</i>); petitioned for listing.
Kootenai Drainage	White Sturgeon (<i>Acipenser transmontanus</i>); listed as endangered.
Kootenai WMA ²	Redband trout (<i>Oncorhynchus mykiss</i>); petitioned for listing.
Kootenai WMA ²	White Sturgeon (<i>Acipenser transmontanus</i>); listed as endangered.

¹ Information provided by the Montana Natural Heritage Program.

² Wildlife Management Area

³ State Park

APPENDIX G

HERBICIDES PROPOSED FOR USE ON DEPARTMENT OF FISH, WILDLIFE AND PARKS REGION ONE PROPERTIES

Herbicides Proposed for Use on Fish, Wildlife & Parks

Region One Properties

The following herbicides have been applied on properties within FWP Region One. Individual chemicals have different properties and thus different circumstances for their proper application. This list also represents chemicals from which FWP will select in the future, so as to use the appropriate herbicide in any given setting. All chemical applications are subject to the guidelines presented in this Plan. Herbicide label instructions will follow in APPENDIX H.

Trade Name	Chemical Ingredient
Tordon	piclorum
Roundup/Ruler	glyphosate
Stinger	clopyralid
Transline	clopyralid
Curtail	clopyralid + 2,4-D
Weedar 64	2,4-D Amine
Hi-Dep	2,4-D Ester

APPENDIX H

HERBICIDE LABEL INSTRUCTIONS AND SAFETY PRECAUTIONS

Specimen Label

Restricted Use Pesticide

May Injure (Phytotoxic) Susceptible, Non-Target Plants. For retail sale to and use only by Certified Applicators or person under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

For use in areas west of the Mississippi River for the control of susceptible broadleaf weeds and woody plants on rangeland and permanent grass pastures, fallow cropland, wheat, barley and oats not underseeded with a legume on grainland (which is not flood or sub-irrigated and not rotated to broadleaf crops), non-cropland, and on Conservation Reserve Program (CRP) acres and wildlife openings in forest and non-crop areas

Active Ingredient:

picloram: 4-amino-3,5,6-trichloropicolinic acid, potassium salt 24.4%

Inert Ingredients 75.6%

Acid Equivalent

picloram: 4-amino-3,5,6-trichloropicolinic acid - 21.1% - 2 lb per gal

EPA Reg. No. 62719-6

EPA Est. 464-MI-1

Net Contents 2.5 gal

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out Of Reach Of Children

WARNING AVISO:

Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Causes Substantial But Temporary Eye Injury • Harmful If Inhaled Or Absorbed Through Skin

Do not get in eyes or on clothing. Wear goggles, face shield or safety glasses when handling. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid breathing spray mist.

First Aid

If in eyes: Flush with plenty of water for at least 15 minutes. Get medical attention.

Environmental Hazards

Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes by cleaning of equipment or disposal of wastes. Do not allow run-off or spray to contaminate wells, irrigation ditches or any body of water used for irrigation or domestic purposes. Do not make application when circumstances favor movement from treatment site.

Picloram is a chemical which can travel (seep or leach) through soil and under certain conditions has the potential to contaminate groundwater which may be used for irrigation and drinking purposes. Users are advised not to apply picloram where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

An aquifer is defined as "an underground, saturated, permeable, geologic formation capable of producing significant quantities of water to a well or spring. It is the ability of the saturated zone, or portion of that zone, to yield water which makes it an aquifer" (American Chemical Society, 1983).

Note: Use In Hawaii limited exclusively to Supplemental Labeling. See "General Use Precautions" for details.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" sections elsewhere on this label.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Tordon* 22K

Table of Contents	Page
Precautionary Statements	1
Hazards to Humans and Domestic Animals	1
First Aid	1
Environmental Hazards	1
Directions for Use	2
Storage and Disposal	2
General Information	2
General Use Precautions	2
Precautions for Avoiding Spray Drift	3
Ground Equipment	3
Aerial Application	3
Woody Plants and Broadleaf Weeds	
Controlled by Tordon 22K	3
Mixing and Application Directions	3
Use with Surfactants	3
Use with Sprayable Liquid Fertilizer Solutions	4
Spot Treatment	4
Wick Application	4
Approved Uses	
Non-cropland Areas	4
Rangeland and Permanent Grass Pastures	4
Grazing Restrictions	4
Broadcast Treatment (Ground and	
Aerial Applications)	4
Weed Control Guidelines for Tordon	
22K in Non-cropland,	
Rangeland and Pasture	5
Spot Treatment	5
Spot Concentrate Application	5
Wick Application	5
Barley, Oats, and Wheat Not Underseeded	
With a Legume	5
Broadcast Treatment (Ground	
and Aerial Applications)	5
Spring Wheat, Barley and Oats	5
Durum Wheat	5
Winter Wheat and Barley	5
Use Rates for Wheat, Barley and Oats	6
Fallow Cropland (Not Rotated to Broadleaf Crops)	6
Spot Treatment	6
Conservation Reserve Program (CRP)	
for Seeding to Permanent	
Grasses Only	6
Broadcast Treatment (Ground and	
Aerial Applications)	7
Spot Treatment	7
Wick Application	7
Warranty Limitations and Disclaimer	7
Inherent Risks of Use	7
Limitation of Remedies	7

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

STORAGE AND DISPOSAL

Do not contaminate water, food, feed or fertilizer by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Metal Container Disposal: Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Plastic Container Disposal: Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Sprayer Clean-Out: To avoid injury to desirable plants, equipment used to apply Tordon 22K should be thoroughly cleaned before reusing to apply any other chemicals.

1. Rinse and flush application equipment thoroughly after use. Dispose of rinse water in non-cropland area away from water supplies.
 2. Rinse a second time, adding 1 quart of household ammonia for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 minutes). Let the solution stand for several hours, preferably overnight.
 3. Flush the solution out the spray tank through the boom.
 4. Rinse the system twice with clean water, recirculating and draining each time.
 5. Nozzles and screens should be removed and cleaned separately.
- General:** Consult federal, state or local disposal authorities for approved alternative procedures.

GENERAL INFORMATION

In areas west of the Mississippi River use Tordon 22K herbicide to control susceptible broadleaf weeds and woody plants on rangeland and permanent grass pastures, fallow cropland, wheat, barley and oats not underseeded with a legume on grainland (which is not flood or sub-irrigated and not rotated to broadleaf crops), non-cropland, and on Conservation Reserve Program (CRP) acres, and wildlife openings in forest and non-crop areas. This product is NOT for sale or use in the San Luis Valley of Colorado.

General Use Precautions

Use this product only as specified on this label. Observe any special use and application restrictions and limitations, including method of application and permissible areas of use as promulgated by state or local authorities.

To prevent damage to crops and other desirable plants, read and follow all directions and precautions on this label and container before using.

Do not use for manufacturing or formulating.

Do Not Mix With Dry Fertilizer.

Chemigation: Do not apply this product through any type of irrigation system.

In Hawaii, approved uses of Tordon 22K are limited to those described in Supplemental Labeling. This Supplemental Labeling may be obtained from your DowElanco representative or chemical dealer. Refer to this Supplemental Labeling for specific use directions and precautions.

Do not make application when circumstances favor movement from treatment site.

Do not apply or otherwise permit Tordon 22K or sprays containing Tordon 22K to contact crops or other desirable broadleaf plants, including but not limited to alfalfa, beans, grapes, melons, peas, potatoes, safflower, soybeans, sugar beets, sunflower, tomatoes, and other vegetable crops, flowers, fruit plants, ornamentals or shade trees or the soil containing roots of nearby valuable plants.

Precautions for Avoiding Injurious Spray Drift

Applications should be made to avoid spray drift since very small quantities of spray, which may not be visible, may seriously injure susceptible crops during both growing and dormant periods. To minimize spray drift, use low nozzle pressure; apply as a coarse spray; and use nozzles designed for herbicide application that do not produce a fine droplet spray. To aid in further reducing spray drift, a drift control and deposition aid such as Nalco-Trol may be used with this product. If such a drift control aid is used, follow all use recommendations and precautions on the product label.

Ground Equipment: With ground equipment spray drift can be lessened by keeping the spray boom as low as possible; by applying 20 gallons or more of spray per acre; by keeping the operating spray pressures at the manufacturers recommended minimum pressures for the specific nozzle type used (low pressure nozzles are available from spray equipment manufacturers); by spraying when the wind velocity is low (follow state regulations). Avoid calm conditions which may be conducive to air inversions. In hand-gun applications, select the minimum spray pressure that will provide adequate plant coverage (without forming a mist).

Aerial Application: With aerial applications, drift may be lessened by using a coarse spray; by using a drift control system; or by using Nalco-Trol drift control additive or equivalent. Adjust spray pressure to provide coarse spray droplets and by using nozzles which do not create fine droplets. Spray boom should be no longer than 3/4 of the wingspan or rotor length. Do not use a thickening agent with the Microfoil or the Thru-Valve booms, or other systems that cannot accommodate thick sprays. Spray only when the wind velocity is low (follow state regulations). Avoid calm conditions which may be conducive to air inversions.

Determine Air Movement and Direction Before Foliar Application. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. It is suggested that a continuous smoke column at or near the spray site or a smoke generator on the spray equipment be used to detect air movements, lapse conditions, or temperature inversions (stable air). If the smoke layers or indicates a potential of hazardous spray drift, do not spray.

Do not contaminate water intended for irrigation or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes.

Do not use on flood or sub-irrigated land.

Do not spray if the loss of forage legumes cannot be tolerated. Tordon 22K may injure or kill legumes. New legume seedlings may not grow for several years following application of this herbicide.

Do not use manure from animals grazing treated areas on land used for growing broadleaf crops, ornamentals, orchards or other susceptible, desirable plants. Manure may contain enough picloram to cause injury to susceptible plants.

Do not use grass or hay from treated areas for composting or mulching of susceptible broadleaf crops.

Do not transfer livestock from treated grazing areas onto sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated grass pasture. Otherwise, urine may contain enough picloram to cause injury to sensitive broadleaf plants.

Do not apply to snow or frozen ground. Application during very cold (near freezing) weather is not advisable.

Tordon 22K should not be applied on residential or commercial lawns or near ornamental trees and shrubs. Untreated trees can occasionally be affected by root uptake of herbicide through movement into the top soil or by excretion of the product from the roots of nearby treated trees. Do not apply Tordon 22K within the root zone of desirable trees unless such injury can be tolerated.

Do not rotate food or feed crops on treated land if they are not registered for use with picloram until an adequately sensitive bioassay or chemical test shows that no detectable picloram is present in the soil.

Do not move treated soil to areas other than those treatment sites for which Tordon 22K is registered for use. Also, do not use treated soil to grow plants for which use of Tordon 22K is not registered until an adequately sensitive bioassay or chemical test shows that no detectable residue of picloram is present in the soil.

Woody Plants and Broadleaf Weeds Controlled by Tordon 22K

Woody Plants:

absinth wormwood	junipers/cedars
aspen	locust
blackberries	multiflora rose
catclaw acacia	pinyon pine
chaparral sp.	rabbitbrush
fringed sagebrush	Scotch broom
gorse	

Annual and Perennial Broadleaf Weeds:

brackenfern	larkspurs	starthistles
buckwheat, wild	geyer	Iberian
buffalobur	plains	purple
bur sage	tall	yellow
burweed	lambquarters	St. Johnswort
camelthorn	leafy spurge	sulfur cinquefoil
clover	licorice, wild	sunflower
crupina, common	locoweeds	tansy ragwort
dock	lupines	tasajillo
field bindweed	milkweed	toadflaxes
goldenrod	ox-eye daisy	thistles
henbane, black	pigweed	artichoke
horsenettle	pricklypear cactus	beaumont
Carolina	ragweeds	bull
white	common	Canada
horseweed	bur	distaff
ironweed	lanceleaf	golden
knapweed	western	Italian
diffuse	rush skeletonweed	musk
Russian	Russian thistle	plumeless
spotted	snakeweeds	Scotch
squarrose	sowthistle	wavy leaf

Mixing and Application Methods

Mix the required amount of Tordon 22K in water and apply as a coarse low pressure spray using ground equipment or aircraft. Use enough spray volume to provide uniform coverage of the weeds. For best results treat when the weeds are growing actively in the spring before full bloom or late summer into fall. Treatments during full bloom or seed stage of some weeds may not give good control.

To prepare the spray, add about half the desired amount of water in the spray tank. Then with agitation, add the recommended amount of Tordon 22K and other registered tank mix herbicides. Finally, with continued agitation, add the rest of the water and additives such as surfactants or drift control and deposition aids.

Use with surfactants

Addition of wetting or penetration agents is not usually necessary when using Tordon 22K. Under extreme conditions, such as drought, addi-

tion of a surfactant may improve efficacy. However, if foliar burn occurs too rapidly, translocation of Tordon 22K will not occur and control of perennial weeds, such as field bindweed, may be reduced.

Use With Sprayable Liquid Fertilizer Solutions

Tordon 22K is compatible with most non-pressurized liquid fertilizer solutions; however, if these solutions are to be sprayed with Tordon 22K, a compatibility test (jar test) should be made prior to mixing. Jar tests are particularly important when a new batch of fertilizer or pesticide is used, when water sources change, or when tank mixture ingredients or concentrations are changed. Compatibility may be determined by mixing the spray components in the desired order and proportions in a clear glass jar before large scale mixing of spray components in the spray tank. Use of a compatibility aid such as Unite or Compex may help obtain and maintain a uniform spray solution during mixing and application. Agitation in the spray tank must be vigorous to be comparable with jar test agitation. For best results, liquid fertilizer rates should not exceed 50% of the total spray volume. Premix Tordon 22K with water and add to the liquid fertilizer/water mixture while agitating contents of the spray tank. Apply the spray the same day it is prepared while maintaining continuous agitation. Rinse spray tank thoroughly after use.

Note: Foliar applied liquid fertilizers can cause yellowing or leaf burn of crop foliage.

Local conditions may affect the use of herbicides. State agricultural experiment stations or extension service weed specialists in many states issue recommendations to fit local conditions. Be sure that use of this product conforms to all applicable regulations.

Spot Treatment

Use application rates as suggested in the "Approved Uses" section of this label or recommended by your area weed control specialist. Apply in a total spray volume of 20 to 100 gallons per acre. Make sure equipment is properly calibrated and that the amount of Tordon 22K added to the spray mixture corresponds to the desired rate and spray volume.

To Calibrate:

1. Measure an area 18.5 ft by 18.5 ft in the target application area.
2. Spray the measured area uniformly with water only and record the number of seconds required to cover the area.
3. Measure the amount of water delivered to the test area by spraying into a container for this amount of time.
4. The amount of water collected in fl oz equals spray volume in gallons per acre.
5. Refer to the chart below for the amount of Tordon 22K to mix at the spray volume indicated by the calibration procedure. This chart contains the amount of Tordon 22K to mix when the application rate is 1 quart per acre. For a rate of 1/2 quart per acre (1 pint), divide the amount in the table by 2. For an application rate of 2 quarts per acre, multiply the table value by 2.

To Apply the Equivalent of 1 Quart of Tordon 22K per Acre at the Spray Volume Indicated, Mix the Following:		
Spray Volume (gallons per acre)	Amount of Tordon 22K per 1 gallon of water	Amount of Tordon 22K per 100 gallons of water
20	10 tsp	5 quarts
40	4 3/4 tsp	2 quarts
60	3 1/4 tsp	1 2/3 quarts
80	2 1/3 tsp	1 1/4 quarts
100	2 tsp	1 quart

Note: tsp = teaspoon
6 tsp = 1 fluid ounce

Do not exceed 4 quarts per acre in any one year as a spot treatment.

Tank Mixture for Spot or Broadcast Treatment of Susceptible Weeds

Tordon 22K may also be tank mixed with 2,4-D products or other registered herbicides for use on areas having mixed species including those which respond well to 2,4-D. Read and follow all directions and use precautions on other product labels.

Wick Application

Mix 1 part of Tordon 22K with 2 parts of water to prepare a 33% solution. Apply when weeds are actively growing and are above most desirable plants. For ironweed and goldenrod, best results are obtained with applications made prior to early bud stage. Wick applicator should be drained and cleaned after each use. Ropes should be changed when flow is reduced from wear, extended use, poor cleaning or intermittent use.

APPROVED USES

NON-CROPLAND AREAS

Use Tordon 22K to control susceptible broadleaf weeds and woody plants on non-cropland areas such as on roadsides or other rights-of-way, along fence rows, and around farm buildings. Use up to 2 quarts of Tordon 22K per acre as a broadcast treatment and up to 4 quarts per acre as a spot treatment. Wick Application may be used on non-cropland. See "Wick Application" in "Mixing and Application Methods" section for directions.

RANGELAND AND PERMANENT GRASS PASTURES

Use Tordon 22K on rangeland and permanent grass pastures to control susceptible broadleaf weeds and woody plants such as (but not limited to) those shown in the table.

Grazing Restrictions: When applying more than 1 quart of Tordon 22K per acre, do not cut grass for feed within two weeks after treatment. Meat animals grazing for up to two weeks after treatment should be removed from treated areas three days prior to slaughter. Do not graze lactating dairy animals on treated areas within two weeks after treatment.

Broadcast Treatment (Ground and Aerial Applications)

Tordon 22K can be applied as a broadcast treatment by ground or aerially to control several broadleaf weeds and woody plants. Apply Tordon 22K at the suggested rates in 2 or more gallons of water per acre by air or in 10 or more gallons of water per acre by ground. Re-treat as necessary but do not exceed 1 quart of Tordon 22K per acre per season. For control of actively growing susceptible annual broadleaf weeds, including Russian thistle, apply 1/4 to 1/2 pint per acre of Tordon 22K. Tordon 22K can also be tank mixed with 1/2 to 1 pound per acre 2,4-D where species present are sensitive to 2,4-D.

Tordon 22K at rates over 1 quart may suppress certain established grasses, such as brome grass, blue gramma, and buffalograss. However, subsequent grass growth should be improved by release from weed competition.

Weed Control Guidelines for Tordon 22K in Non-cropland, Rangeland and Pasture*

Weed Species**	Rate per Acre	Comments
biennial thistles bull musk plumeless Scotch	Fall: 1/2 pint Spring: 6 - 8 fl oz with 1.0 lb ae 2,4-D	Apply when thistles are in the rosette stage before bolting in the spring or in the fall prior to soil freeze up.
bolton musk thistle	1/2 - 1 pint + 1 lb ae 2,4-D per acre	Apply before flowering.
broom snakeweed geyer larkspur locoweeds multiflora rose plains larkspur prickly pear cactus sulfur cinquefoil	1 pint	For pricklypear cactus, use of a diesel oil-water emulsion spray mixture may improve control.
black henbane crupina diffuse knapweed spotted knapweed yellow starthistle	1 - 2 pints	
absinth wormwood bursage Douglas rabbitbrush goldenrod ox-eye daisy	1 - 2 pints	Tank mix the lower rate with 1.0 lb ae per acre 2,4-D. Lower rates may require annual spot treatments.
wild lonicice	1 quart	
Canada thistle field bindweed gorse lupines rush skeletonweed St. Johnswort tansy ragwort	1 - 2 quarts	Tank mix the lower rate with 1.0 lb ae per acre 2,4-D. Lower rates may require annual spot treatments.
dalmation toadflax juniper perennial sowthistle Russian knapweed	2 - 3 quarts	For Russian knapweed, apply at bud stage or in the fall
tall larkspur yellow toadflax	3 - 4 quarts	
leafy spurge	1 - 4 quarts	Lower rates will require annual retreatment for several years. Retreat when control drops below 80%

*For additional species or more specific rates consult your area's current Weed Control Guide and/or your local DowElanco representative.

**Many seedling annual weeds can be controlled using 1 pt per acre.

For rates exceeding 1 quart per acre, apply only as a spot treatment and the total area treated in a single season should not exceed 25% of a landowner's acreage found in any particular watershed.

Spot Treatment

See "Spot Treatment" in "Mixing and Application Methods" section for directions for calibration, spray volume determination and mixing.

Spot Concentrate Application

Eastern red cedar can be controlled with spot concentrate applications of Tordon 22K in either the spring (April-May) or fall (September-October). For best results, use 3 ml to 4 ml of Tordon 22K (undiluted) per 3 feet of plant height. Application should precede periods of expected rainfall. Apply directly to soil within the dripline and on the upslope side of the tree. Application to trees taller than 15 feet is not recommended. Do not use more than 2 pints of Tordon 22K per acre in any one year.

Wick Application

See "Wick Application" in "Mixing and Application Methods" section for directions. Apply when weeds are actively growing and are above most desirable plants. For ironweed and goldenrod, best results are obtained with applications made prior to early bud stage.

Barley, Oats, and Wheat Not Underseeded With a Legume (Which is Not Flood or Sub-Irrigated and Not Rotated to Broadleaf Crops)

Use Tordon 22K for the control of susceptible annual broadleaf weeds such as (but not limited to) volunteer sunflower, wild buckwheat, lamb-squarters, pigweed, Russian thistle, and sowthistle.

Use Precautions

Do not apply Tordon 22K within 50 days before harvest.

Spray mixtures may cause shorter straw on some varieties of cereals but grain yields are usually not affected.

Do not graze or feed forage from treated areas for 2 weeks after treatment. Do not harvest hay from treated grain fields.

Use only on land that will be planted the following year to grass, barley, oats, wheat or fallowed. Do not apply more than 1 1/2 fluid ounces of Tordon 22K per acre during the small grain growing season.

Broadcast Treatment (Ground and Aerial Applications)

Tordon 22K can be applied as a single broadcast treatment by ground or aerially to control several broadleaf weeds by itself or as a tank mix with 2,4-D, MCPA, or sulfonyleurea herbicides such as Ally. Apply Tordon 22K at the rates suggested in the following table in 2 to 5 gallons of water per acre by air or in 5 to 20 gallons of water per acre by ground. The addition of surfactants may aid control under dry conditions, but may cause injury to grain if used over the top. Read and follow directions and precautions on other product labels when tank mixing.

Spring Wheat, Barley and Oats

Apply from the 3 to 5 leaf stage to the early jointing stage of growth as indicated in the table below. Applications at the 3 to 5 leaf stage occasionally cause slight head malformations and straw shortening but normally do not affect yield.

Durum Wheat

Do not treat durum wheat since some varieties of durum wheat may be injured.

Winter Wheat and Barley

Apply after resumption of active growth in the spring until the early jointing stage.

Use Rates for Wheat, Barley and Oats†					
Weeds	Weed Growth Stage††	Grain Growth Stage	Amounts of Each Product Per Acre†††		
			Tordon 22K	4 lb ae/gal 2,4-D or MCPA	6 lb ae/ga 2,4-D or MCPA
More susceptible species, such as: lambsquarters pennycress wild mustard mayweed	3 inches	3 to 5 leaf to early tillering	1 fl oz	1/2 pint	1/3 pint
	3 to 6 inches	Tillering to early jointing	1 1/2 fl oz	3/4 pint	1/2 pint
Less susceptible species such as: volunteer sunflower wild buckwheat Russian thistle pigweed Canada thistle, top growth suppression	1 to 6 inches	Tillering to early jointing	1 1/2 fl oz	3/4 to 1 pint	1/2 to 2/3 pint

† For oats, do not tank mix with 2,4-D herbicides.

†† For best results, treat when weeds have 2 to 4 leaves and are actively growing.

††† When measuring small amounts of Tordon 22K, special care should be taken not to exceed suggested rates.

FALLOW CROPLAND (NOT ROTATED TO BROADLEAF CROPS)

Apply Tordon 22K as a post harvest or fallow treatment in continuous grain or during the fallow period. Tordon 22K may be applied alone or in tank mix combination with 2,4-D or other herbicides registered for this use. Apply in 2 or more gallons of water per acre by air or 5 or more gallons per acre by ground. Spray only under conditions that will prevent injury to nearby susceptible crops or ornamentals. Refer to "General Use Precautions" section for information on preventing drift to off-target areas.

Application Rates

Note: Do not apply more than 1 pint per acre as a broadcast treatment in any calendar year.

Annual Weeds: To control annual weeds such as Russian thistle and wild buckwheat, apply 1/4 to 1/2 pint per acre of Tordon 22K in tank mix combination with 1/2 to 1 lb ae of 2,4-D or other herbicides registered for use on fallow land. Apply when weeds are actively growing.

Field Bindweed: Apply 1/2 to 1 pint per acre of Tordon 22K plus 1/2 to 1 lb ae per acre of 2,4-D when bindweed is actively growing. Optimum time for treatment is when plant runners reach 8 to 12 inches. Use 1/2 pint per acre to control light to moderate infestations under good growing conditions or to reduce the potential for crop injury. Use higher rates for heavy infestations and longer term control. Some regrowth will occur the following season and a re-treatment program for successive years is recommended.

Canada thistle: Apply 1 pint per acre of Tordon 22K plus 1 lb ae per acre of 2,4-D when the majority of thistle plants are emerged but prior to bud stage.

Crop Rotation

Use only on land to be planted the following year to grass, barley, oats, wheat or fallowed. Many broadleaf crops are extremely sensitive to soil residues of Tordon 22K. Do not plant sensitive broadleaf crops for 36 months after treatment or until soil residues have declined to a safe level as indicated by an adequately sensitive bioassay using the intended broadleaf crop. A bioassay is recommended following treatment prior to planting any sensitive broadleaf crop.

Preplant Interval

A preplant interval following Tordon 22K application prior to planting small grains is recommended to reduce or eliminate potential crop injury and/or yield reduction. The possibility for crop injury or yield reduction to occur depends on application rate, soil organic matter, rainfall, temperature and incidence of cereal diseases. Adequate soil moisture and soil temperature during the preplant interval is important in reducing, but may not eliminate, the risk of crop injury. When considering use of Tordon 22K on fallow land, growers should consider the benefit of weed control against the risk of crop damage and treat only if the risk of injury to small grains can be tolerated. The following preplant intervals are recommended:

For applications up to 1/2 pint per acre, allow a minimum of 45 days of soil temperatures above 40°F between application and planting.

For applications of greater than 1/2 pint and up to 1 pint per acre, allow a minimum of 60 days of soil temperatures above 40°F between application and planting, except in the states of Idaho, North Dakota, Nebraska, Montana, Oregon, South Dakota, Washington and Wyoming, where the minimum preplant interval is 90 days.

Spot Treatment

See "Spot Treatment" in "Mixing and Application Methods" section for directions for calibration, spray volume determination and mixing.

Spot treatments of Tordon 22K at rates over 1 pint per acre can be made on fallow, non-irrigated cropland if the treated areas comprise less than 10% of the immediate field in any one year. Tordon 22K should not be applied to cropland at rates exceeding 2 quarts per acre. When Tordon 22K at rates above 1 pint per acre are applied, injury to small grains may result for periods up to two years after treatment.

CONSERVATION RESERVE PROGRAM (CRP) FOR SEEDING TO PERMANENT GRASSES ONLY

Do not use Tordon 22K if legumes are a desired cover during CRP.

Conditions that stress grasses, such as drought, will increase potential for injury to the grass at all stages of growth.

To reduce potential damage to subsequent small grain crops, use the lower rate or discontinue the use of Tordon 22K at least 2 years prior to the seeding of small grain crops. After CRP, do not plant broadleaf crops in treated acres until an adequately sensitive bioassay shows that no detectable picloram is present in the soil.

Broadcast Treatment (Ground and Aerial Applications)

Applications of Tordon 22K should be made after perennial grasses are well established (have developed a good secondary root system and show good vigor). Most perennial grasses show improved tolerance to the herbicide at this stage of development.

For control of actively growing perennial weeds, use up to 1 quart per acre of Tordon 22K after the grass is established. For best results, use in 2 or more gallons of water per acre by air or in 5 or more gallons of water per acre by ground. Increasing the rate of application can increase the risk of injury.

For control of actively growing susceptible annual broadleaf weeds, (including Russian thistle) apply 1/4 to 1/2 pint per acre of Tordon 22K. Tordon 22K can also be tank mixed with 1/2 to 1 pound per acre of 2,4-D where 2,4-D sensitive species present. Read and follow all directions for use and use precautions on other product labels.

Spot Treatment

See "Spot Treatment" in "Mixing and Application Methods" section for directions for calibration, spray volume determination and mixing.

For spot applications when perennial grasses are established, use 1 to 8 pints per acre of Tordon 22K. Rates of 2 quarts per acre or more should only be used for control of deep-rooted perennial broadleaf weeds.

Tordon 22K at rates over 2 pints per acre may suppress certain established grasses such as bromegrass, bluegrass and buffalograss. However, subsequent grass growth should be improved by release from weed competition.

Wick Application

See "Wick Application" in "Mixing and Application Methods" section for directions. Apply when weeds are actively growing and are above most desirable plants. For ironweed and goldenrod, best results are obtained with applications made prior to early bud stage.

Warranty Limitations and Disclaimer

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268, U.S.A.

LABEL CODE 112-42-002 DATE CODE 292
EPA APPROVAL 12/20/91 REPLACES 112-42-001

Revisions Include:

- 1) Label reformatted and edited for clarity.
- 2) Updated woody plants and broadleaf weeds controlled listed.

Specimen Label

Transline*

Herbicide

For Selective Control of Broadleaf Weeds in Non-Cropland Areas, Industrial Manufacturing and Storage Sites and Rights-of-way

Active Ingredient(s):

Clopyralid (3,6-dichloro-2-pyridinecarboxylic acid) as the monoethanolamine salt40.9%

Inert Ingredient(s):.....59.1%

ACID EQUIVALENT:

3,6-dichloro-2-pyridinecarboxylic acid, 31% - 3 lb/gal

E.P.A. Registration No. 62719-73

E.P.A. Est. 464-MI-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUCION:

PRECAUCION AL USUARIO:

Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

**CAUSES EYE INJURY • HARMFUL IF
INHALED OR ABSORBED THROUGH
SKIN**

Avoid Contact With Eyes, Skin Or Clothing

• Avoid Breathing Spray Mist • Wash
Thoroughly With Soap And Water After
Handling • Remove Contaminated Clothing
And Wash Before Reuse

STATEMENTS OF PRACTICAL TREATMENT: If in eyes,
flush with plenty of water. Get medical attention if irritation
persists. If on skin, wash with plenty of soap and water.
Get medical attention.

Physical or Chemical Hazards

**COMBUSTIBLE • Do Not Use or Store Near
Heat or Open Flame. Do Not Cut or Weld
Container.**

Environmental Hazards

Do not contaminate water when disposing of equipment
washwaters. Do not contaminate water used for irrigation
or domestic purposes. Do not apply directly to any body of
water or wetlands.

See additional precautionary statements
elsewhere on this label.

NOTICE

Read and understand the entire label before using. Use
only according to label directions.

Before buying or using this product, read "WARRANTY
LIMITATIONS AND DISCLAIMER" elsewhere on this label.
If terms are not acceptable, return unopened package at
once to seller for full refund of purchase price paid.
Otherwise, use by the buyer or any other user constitutes
acceptance of the terms under WARRANTY LIMITATIONS
AND DISCLAIMER.

IN CASE OF AN EMERGENCY

endangering life or property involving this
product, call collect 517-636-4400

AGRICULTURAL CHEMICAL

**Do Not Ship or Store with Food, Feeds,
Drugs, or Clothing**

*
Transline

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

TRANSLINE Herbicide is recommended for selective, postemergence control of the following broadleaf weeds in non-cropland areas including equipment pathways, industrial manufacturing and storage sites and rights-of-way such as along roadsides, electrical power lines, communication lines, pipelines and railroads.

acacias	ladysthumb ¹
artichoke, Jerusalem	lettuce, prickly
buckwheat, wild	locoweed, white
buffalobur ¹	locoweed, Lambert
burdock, common	marshelder
Canada thistle	mesquite
(rosette to bud)	musk thistle (rosette to bud)
chamomile, false	nightshade, Eastern black
(scintless)	nightshade, cutleaf
chamomile, mayweed	nightshade, hairy
(dogfennel)	oxeye daisy
clover, sweet	pineappleweed
clover, red	ragweed, common
cocklebur, common	ragweed, giant
coffoewood	salsify, meadow (goatsbeard)
cornflower	sicklepod
(bachelor button)	smartweed, green ¹
dandelion	sorrel, red
dock, curly	sowthistle, annual
groundsel, common	sowthistle, perennial ¹
hawkbeard, narrowleaf	starthistle, yellow
horsetweed	sunflower (common and wild)
jimsonweed	thistle, Canada
knapweed, diffuse	thistle, musk
knapweed, Russian ¹	vetch
knapweed, spotted	

¹ These weeds may only be suppressed. Suppression is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment.

For perennial weeds, TRANSLINE Herbicide will control the initial topgrowth and inhibit regrowth during the season of application (season-long control). At higher use rates shown on this label, TRANSLINE Herbicide may cause a reduction in shoot regrowth in the season following application; however, plant response may be inconsistent due to inherent variability in shoot regrowth from perennial root systems.

Timing: Apply to actively growing weeds. Extreme growing conditions such as drought or near freezing temperatures prior to, at, and following time of application may reduce weed control. Only weeds which are emerged at the time of application will be affected. Wet foliage at the time of application may decrease control. The treatment with TRANSLINE Herbicide will be rainfast within 6-8 hours after application.

Rate: Generally, lower labeled application rates will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species, perennials, and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands, and/or larger weeds), the higher rates will be needed.

Coverage: Adequate spray coverage and drift control are important. Obtaining a balance between spray coverage and drift control may sometimes be difficult but can be achieved provided the applicator understands the factors affecting coverage and drift. Factors affecting spray coverage include spray volume, and weed density. As weed density increases, spray volume should be increased to obtain equivalent weed control. Refer to manufacturer's recommendations for information on the relationship between gallons per acre,

spray pressure, sprayer speed, nozzle type and arrangement, nozzle height above the target canopy, droplet size, and drift potential for respective application equipment. Use equipment and nozzle types which are designed for herbicide application. Reducing total spray volume may result in decreased coverage and weed control. Use enough total spray volume and a delivery system to provide thorough coverage and a uniform spray pattern. Do not apply where spray drift may be a problem due to proximity of susceptible crops or other desirable plants. **Use of Adjuvants:** Addition of surfactants, crop oils, or other adjuvants is not usually necessary when using TRANSLINE Herbicide. Adding a surfactant to the spray mixture may increase effectiveness on weeds. If an adjuvant is added to the spray solution, follow all manufacturer use guidelines.

Tank Mixes: When tank mixing, read and follow the label of each tank-mix product used for precautionary statements, directions for use, weeds controlled, and geographic and other restrictions. Use in accordance with the most restrictive of label limitations and precautions. No label dosages should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Non-Cropland

For use on non-cropland areas such as industrial manufacturing and storage sites and rights-of-way such as along roadsides, electrical power lines, communication lines, pipelines and railroads.

Broadcast Application (Ground): For control of broadleaf weeds, apply 1/4 to 1 1/2 pints of TRANSLINE Herbicide per acre (equivalent to 0.09 to 0.5 lbs ae per acre). Non-ionic surfactant should be used in spray mixtures at 1 to 2 quarts per 100 gallons of spray mixture. The lower rate of 1/4 pint per acre provides acceptable control of weeds only under highly favorable plant growing conditions and when plants are no larger than 3 to 6 inches tall. Where Canada thistle or knapwoods are the primary pest, best results are obtained by applying 2/3 to 1 1/2 pints of TRANSLINE Herbicide per acre after basal leaves are produced. For roadside applications, spray volumes of 25 to 50 gallons per acre will ensure adequate coverage. Addition of 1/4 to 1 ounce Telar per acre may enhance thistle control when low rates of TRANSLINE are used. TRANSLINE Herbicide can be tank mixed with diesel oil or equivalent inert agent approved for use on agricultural crops. Established grasses are tolerant but new grass seedlings may be injured to varying degrees until the grass has become well established.

High-Volume Leaf Stem Treatment (Woody Plants): For control of brush, use 1 to 3 quarts of TRANSLINE Herbicide per 100 gallons of total spray solution. Thorough coverage is necessary for good results, therefore, apply as a complete spray-to-wet foliar application, including all leaves, stems, and root collars but do not exceed application of more than 1 1/2 pints of TRANSLINE Herbicide per acre. To minimize drift, use a maximum spray pressure of 50 psi and keep sprays no higher than the tree crowns. Trees taller than 8 feet in height may be hard to treat efficiently and to obtain necessary coverage.

Do not apply by air.

Unsatisfactory control may result if application is made when brush and weeds are under severe drought stress or other conditions that inhibit good growth. Environmental conditions may influence results considerably. For best results on mesquite, apply in the spring or early summer during the period 40 to 90 days after the first green growth appears and when soil moisture is adequate for good growth. Soil temperatures of 75° to 83°F at a 12 to 18 inch depth are optimal for good plant kills. Soil temperature of less than 75°F at this depth will reduce the ultimate root kill of mesquite.

USE PRECAUTIONS

This product can affect susceptible broadleaf plants directly through foliage and indirectly by root uptake from treated soil. Therefore, do not apply TRANSLINE Herbicide directly to or allow spray drift to come in contact with vegetables, flowers, grapes, tomatoes, potatoes, beans, lentils, peas, alfalfa, sunflowers, soybeans, safflower, or other desirable broadleaf crops and ornamental plants or soil where these sensitive crops will be planted the same season.

Do not contaminate irrigation ditches or water used for irrigation or domestic purposes.

Avoid Spray Drift: Applications should be made to avoid spray drift since very small quantities of the spray, which may not be visible, may severely injure susceptible crops during both growing and dormant periods. Use coarse sprays to minimize drift since, under adverse weather conditions, fine spray droplets may drift a mile or more. A drift control or deposition agent such as Nalco-Trol may be used with this product to aid in reducing spray drift. If used, follow all use recommendations and precautions on the product label.

Ground Application: To minimize spray drift, apply TRANSLINE Herbicide in a total spray volume of 10 or more gallons per acre as large-droplet, low-pressure spray. Refer to manufacturer's recommendations for additional information on gallons per acre, spray pressure, sprayer speed, nozzle types and arrangements, nozzle heights above the target canopy, etc., for respective application equipment. With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible; by using no more than 30 pounds spraying pressure with large droplet-producing nozzle tips; by spraying when wind velocity is low; and by stopping all spraying when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone-type insecticide or other nozzles that produce a fine-droplet spray.

Do not apply by aircraft.

Do Not Transfer Livestock from treated grazing areas onto sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated pasture. Otherwise, urine may contain enough clopyralid to cause injury to sensitive broadleaf plants.

Straw from treated areas, or manure from animals that have grazed treated areas, cannot be used for composting or mulching on ground where susceptible crops may be grown the following season. To promote herbicide decomposition, plant material should be evenly incorporated or burned. Adequate moisture is also required to promote breakdown of plant residues which contain clopyralid.

Do not use in a greenhouse. Excessive amounts of this herbicide in the soil may temporarily inhibit seed germination or plant growth.

Spray Equipment - Cleaning Instructions

To avoid injury to desirable plants, equipment used to apply TRANSLINE Herbicide should be thoroughly cleaned before reusing to apply any other chemicals.

1. Rinse and flush application equipment thoroughly after use at least three times with water, and dispose of rinse water in non-cropland area away from water supplies.
2. During the second rinse, add 1 qt of household ammonia for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15-20 min.). Let the solution stand for several hours, preferably overnight.

3. Flush the solution out the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Nozzles and screens should be removed and cleaned separately.

PRECAUTIONARY STATEMENTS

Environmental Hazards

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store above 28°F or warm to 40°F and agitate before use.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal (Metal): Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Container Disposal (Plastic): Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY LIMITATIONS AND DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions therein under normal conditions of use. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL.

Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.

When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify in writing DowElanco of any claims to be eligible to receive other remedy given below. The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of DowElanco or any other seller will be one of the following, at the election of DowElanco:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

The seller will not be liable for consequential or incidental damages or losses.

The terms of this Warranty Limitations And Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations And Disclaimer in any manner.

11388-L1

290

DowElanco
Indianapolis, Indiana 46268 U.S.A.
*Trademark of DowElanco

FORM # 113-20-001 (1-92)
SPECIMEN LABEL 86-1936 DATE CODE 290
THIS IS AN INITIAL DOWELANCO PRINTING

NO OTHER POST-EMERGENT DOES MORE, SO WHY PAY MORE?

- Canada thistle, field bindweed and other tough broadleaf perennials controlled at last — including multiflora rose, buckbrush, and other woody species.
- Also very effective on annual broadleaf weeds, including velvetleaf, cocklebur, morningglory, mustards, and ragweed. Over 120 species listed!
- Apply as a low-volume spray, preferably 5 to 10 gals. total solution per acre for most ground applications. Aerially applied, HI-DEP needs only enough water for a total spray volume of 2 quarts minimum per acre.
- Droplets resist evaporation between nozzle and target, thus remaining moist on the leaf surface for improved absorption.

Unlike ordinary 2,4-D, more HI-DEP translocates into the horizontal roots, killing the root buds.

Read label before use.

G pbi / Gordon CORPORATION

1217 W. 12th St. • P.O. BOX 4090
KANSAS CITY, MO 64101 • 816/421-4070

1-800-821-7925

FAX: 816-474-0462

The literature contained herein is not intended to be used as a substitute for the information contained on the label of the product container. Specimen labels and other literature are subject to revision. Before using this product, read and follow all label instructions on the container/package.

CONVENTIONAL 2,4-D DROPLET

LITTLE HERBICIDE PENETRATES INTO THE LEAF; MUCH EVAPORATION LOSS.

VS. HI-DEP SPRAY DROPLET

HI-DEP IS RAPIDLY AND MORE COMPLETELY ABSORBED; VERY LITTLE LOSS.

HI-DEP[®] BROADLEAF HERBICIDE

CONTROLS MANY TOUGH PERENNIAL WEEDS THAT ESCAPE ORDINARY 2,4-D; ANNUALS, TOO.

Acme HI-DEP[®] High Deposition, Low Volume Broadleaf Herbicide provides outstanding post-emergence control of weeds such as Canada thistle, field bindweed, and hemp dogbane which, until now, have proven very difficult if not extremely costly to control. HI-DEP overcomes the problem because it penetrates deeply enough to get into the extensive root system and kills the root buds where regrowth occurs.

Contrast this to conventional 2,4-D treatments. While they often appear to kill the weeds and may even eliminate their competition with the crop, all too often only the tops are burned back. The same weeds — and more — reappear the next growing season, and year after year.

HI-DEP is a unique formulation of the mixed amine salts, dimethylamine and diethanolamine, of 2,4-D, combined with a special additive which enhances the product's non-evaporative qualities. Unlike ordinary 2,4-D, when applied, HI-DEP droplets tend to resist evaporation in flight and do not crystallize. Therefore, HI-DEP is absorbed more rapidly and completely by the leaf.

A radioisotope tracer study conducted by a major midwestern university, shows that, within four hours, *over twice as much* active ingredient will move into the roots of field bindweed than will standard 2,4-D.

Because of HI-DEP's unique characteristics, the finished spray won't burn the leaf, even when applied straight or in low volume solutions. Penetration is increased, even where thick or waxy cuticles or hairy leaf surfaces tend to leave other herbicides high and dry. HI-DEP gets to the root of the problem — and at a cost that is little more than for ordinary 2,4-D!

HI-DEP[®] is a registered trademark of PBI/Gordon Corporation.

RECOMMENDED USES:

Broadleaf weed control in corn, pasture, wheat and other small grains, stubble, fallow, and grain sorghum and other crops, orchards and non-crop areas as indicated on the label.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid 33.2%
Diethanolamine salt of 2,4-Dichlorophenoxyacetic acid 16.3%

PACKAGING:

2 x 2½ gallon plastic containers per case, shipping 55 lbs.
30 gallon drum, shipping weight 323 lbs.

© 1991, PBI/GORDON CORPORATION

ACME®

AGRICULTURAL PRODUCTS

HI-DEP®

HIGH DEPOSITION BROADLEAF HERBICIDE

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 33.2%

Diethanolamine Salt of 2,4-Dichlorophenoxyacetic acid 16.3%

INERT INGREDIENTS 50.5%

TOTAL 100.0%

This Product Contains:

3.8 lbs. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 38.6% isomer Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN DANGER — PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

See below for additional Precautionary Statements and Statement of Practical Treatment.

U.S. Patent No. 4,971,630

READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS:

DANGER: Corrosive, causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling. When handling this product, wear chemical resistant gloves. Wash thoroughly with soap and water after handling. Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT:

IF IN EYES: In case of eye contact, immediately flush eyes with plenty of water for 15 minutes. Call a physician at once.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly. If irritation develops, get medical attention.

IF SWALLOWED: Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol. Call a physician at once.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

IF INHALED: Remove victim to fresh air and apply respiration if indicated.

ENVIRONMENTAL HAZARDS: Under no circumstances should this herbicide product or any 2,4-D weed killer be used in the vicinity of cotton, tomatoes, garden crops, grapes, ornamentals or other susceptible crops or severe damage may result. Do not apply on windy days. Do not use equipment used in applying this product or any 2,4-D weed killer to apply insecticides, fungicides, or other materials to susceptible crops. Do not use this product through any type of irrigation system. Avoid contamination of water supplies that may be used to irrigate or water susceptible crops or to be used for domestic purposes. This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water except as specified on this label. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas. Do not apply in any manner not specified on this label.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing, loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing, loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application and until sprays have dried. Regular long-sleeved work clothing should be worn when working in treated fields. See Precautionary Statements, Statement of Practical Treatment and Note to Physician for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER — PELIGRO, area treated with HI-DEP Broadleaf Herbicide, date of application, appropriate protective clothing, and reentry interval (i.e. until sprays have dried).

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing. To prevent cross-contamination, do not store near other herbicides, fertilizers, insecticides, fungicides, or near seeds.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For plastic container, triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed, by state and local authorities, by burning. If burned, stay out of smoke. For metal drums, triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

AUTOMOBILE FINISH PRECAUTION

Undiluted spray droplets may damage automobile finishes. Cars should not be sprayed. If accidental exposure does occur, the car should be washed before product dries.

NOTICE TO USER: This product must be applied in compliance with the pesticide regulations of the state in which application is made. Check with local authorities regarding regulations which may affect the application of this product.

— USE INSTRUCTIONS —

AERIAL APPLICATION: Ready-To-Use, not necessary to dilute for application rates of 1/2 gallon (2 quarts) per acre or higher. For rates lower than 1/2 gallon, dilute with water for a total solution per acre of not less than 1/2 gallon.

AIRCRAFT SPECIFICATIONS (FIXED WING OR ROTARY WING): Boom width should not exceed 3/4 the length of the aircraft wingspan. Do not exceed 25 psi nozzle pressure. Number of nozzles required to obtain desired volume per acre is dependent on swath width and speed of aircraft. Nozzles should be positioned between 135 and 175 degrees from direction of flight for fixed wing. **DO NOT APPLY THROUGH BECOMIST NOZZLE SYSTEMS.** Maintain aircraft altitude of 10 to 12 feet during application. See manufacturer's technical bulletin regarding nozzling and method of application specifications.

GROUND APPLICATION: Apply in water, 1 to 10 gallons total solution per acre minimum with conventional equipment. Use nozzle systems capable of spraying correct gallonage; 25 psi is recommended.

— WEEDS CONTROLLED LIST —

Use HI-DEP herbicide to control many broadleaf weeds including:

PERENNIAL WEEDS

Artichoke
Aster
Austrian fieldcress
Birdweed
Black-eyed susan
Blue lettuce
Canada thistle
Catsnip
Chicory
Clover (many types)
Coffee-weed
Dandelion

Docks
Dogbane
Goldenrod
Ground ivy
Hemlock
Ironweed
Leafy spurge
Knapweed
(spotted)
(Russian)
(diffuse)

Locoweed
Mugwort
Nettles
Orange hawkweed
Povertyweed
Rushes
Sowthistle
St. Johnswort
Singing nettles
Strawberry (wild)
Tall buttercup
Tanweed

Toad flax
Vervains
Whinnetop
(Hoary cress)
Wild garlic
Wild onion
Wild sweet potato
Yellow rocket

ANNUAL AND BIENNIAL WEEDS

Beggar-ticks
Broomrape
Black medic
Broomweed
Bull thistle
Burdock
Caperweed
Catchweed bedstraw
Chickweed
Cinquefoil
Cockle
Cocklebur
Croton
Devilsclaw
Falseflax
Fleabane (daisy)
Flaxweed
Franchweed

Gainsage
Goatsbeard
Gooseloot
Groundsel
Gumweed
Hamlet
Jewelweed
Jimsonweed
Jim Hill mustard
(Tumble mustard)
Knotweed
Lambsquarters
Lettuce (wild)
Mallow
Marshelder
Marshall
(Horseweed)
Manzanita

Mediterranean sage
Miners lettuce
Morningglory (annual)
Musk thistle
Mustard
Parsnip
Penny-cress
Pepperweed
Pigweed (redroot)
Plantains
Prickly lettuce
Primrose
Puncturevine
Ragwort (wild)
Ragweed
Russian thistle
Scotch thistle
Shepherdspurse

Sneezeweed
Sow thistle
(common)
Spanshneedles
Sunflower
Tansy mustard
Tansy ragwort
Tumbleweed
Tumble pigweed
Velvetleaf
Vetch
Wild carrot
Wild parsnip
Wild turnip
Wildweed
Wormwood
Yellow starthistle

ALSO CERTAIN 2,4-D SUSCEPTIBLE WOODY PLANTS SUCH AS:

Big sagebrush
Buckbrush
Chamise
Coastal sage

Elderberry
Hazel
Locust
Manzanita

Poison ivy
Poison oak
Rabbitbrush
Sagebrush

Sand shinnery oak
Sumac
Tules (bulrush)
Willow

TO CONVERT LOCAL RECOMMENDATIONS INTO TERMS OF HI-DEP USE THE FOLLOWING TABLE:

2,4-D Acid Equivalent	1 lb	3/4 lb	1/2 lb	3/4 pt	1/2 pt	3/4 qt	1/2 qt
HI-DEP	2 pt.	1 1/2 pt.	1 pt.	3/4 pt.	1/2 pt.	3/4 qt.	1/2 qt.

TIMING OF APPLICATION MAY VARY — Your State Cooperative Extension Service may have information on the correct application time for your area.

WHEAT, BARLEY, OATS, RYE

See Table I for recommended use rates. Spray after grain begins heading and before the boot stage (usually 4 to 8 inches tall) and weeds are small. Do not apply before the tiller stage nor from early boot through the milk stage. To control large weeds, preharvest treatment can be applied when the grain is in the dough stage. Best results will be obtained when soil moisture is adequate for plant growth and weeds are growing well. Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within two weeks of treatment.

WHEAT

Perennial broadleaf weeds — Apply 2 pints per acre when weeds are in bud stage, but do not spray grain in the boot to dough stage. The 2 pint (1 pound acid equivalent) per acre application of any 2,4-D product can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury.

CORN

See Table I for recommended use rates.

PREEMERGENCE — Apply to soil anytime after planting but before corn emerges. Do not use on very light, sandy soil.

EMERGENCE — Apply just as corn plants are breaking ground.

POSTEMERGENCE — Apply to emerged corn. When corn is over 8 inches tall, use drop nozzles and direct spray from corn plant. Do not apply from tasseling to dough stage. Injury to corn is most likely to occur if applied when corn is growing rapidly under high temperatures and high soil moisture conditions. In such situations, use the low rate of 1/2 pint per acre. After application, delay cultivation for 10 to 10 days to allow the corn to overcome any temporary brittleness.

PREHARVEST: After the hard dough or denting stage, apply by air or ground equipment to suppress perennial weeds, decrease weed seed production and control tall weeds such as bindweed, cocklebur, dogbane, jimsonweed, ragweed, sunflower, velvetleaf, and vines that interfere with harvesting.

NOTE: Do not forage or feed corn or fodder for 7 days following application.

NOTE: Hybrids vary in tolerance to 2,4-D. Some are easily injured. Spray only hybrids known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service Weed Specialist for this information.

WITH LIQUID NITROGEN SOLUTIONS

The compatibility of HI-DEP, water and the liquid nitrogen solutions should be determined before combining in the spray tank. The testing can be conducted by mixing all the components in a small container in proportional quantities. If the mixture separates after standing but can be mixed readily by shaking then the mixture can be used as long as good agitation is maintained. If large flakes, sludges, gels or other precipitates form, or if a separate oily layer or oil globules appear, then the herbicide and the liquid fertilizer should not be used in the same spray tank. For late season control of young smartweeds, cocklebur, annual morningglory and other broadleaf weeds less than 1 inch high. Field should be as clean as possible and corn 20 to 30 inches tall. Apply 1 pint with 80 to 120 lbs. nitrogen per acre. The spray MUST be prepared by first adding required amount of liquid nitrogen solution to spray tank. Next dilute 1 pint with 2 quarts of clean water for each acre to be treated with one tankful. Start the tank agitator and SLOWLY add the diluted 2,4-D solution. Spray immediately, maintaining continuous agitation until spray tank is empty. Direct the spray to lower 3' to 4' of corn stalks. Use spray equipment designed to handle corrosive liquid nitrogen solutions. After spraying remove any remaining solution and rinse spray rig thoroughly with water. Mix only one tank at a time. Do not spray during or immediately following cold weather.

SORGHUM (MILO)

See Table 1 for recommended use rates. Treat only after sorghum is 6 inches high and preferably before it is 15 inches high. Do not treat during the boot, tasseling, or early dough stages. If crop is taller than 8 inches, use drop nozzles to keep the spray off the leaves. Temporary crop injury can be expected under conditions of high soil moisture and high air temperatures. If it is necessary to apply under these conditions, use no more than 1/2 pints per acre. NOTE: Hybrids vary in tolerance to 2,4-D. Some are easily injured. Spray only hybrids known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service Weed Specialist for this information.

RICE

See Table 1 for recommended use rates. Apply in the late tillering stage of rice development, at the time of first joint development (first to second green ring), usually 6 to 9 weeks after emergence. Do not apply after panicle initiation, after rice internodes exceed 1/2 inch, at early seeding, early panicle boot, flowering, or early heading growth stages.

NOTE: Some rice varieties under certain conditions can be injured by 2,4-D. Therefore, before spraying, consult local Extension Service or University Specialists for appropriate rates and timing of 2,4-D sprays.

SUGARCANE

See Table 1 for recommended use rates. Use up to 4 applications per year in accordance with State recommendations.

TABLE 1: AMOUNT OF HERBICIDE TO USE IN CROPS

SMALL GRAINS		
Spring postemergence wheat, barley, rye	1/2-1 1/2 pt	2-3 pts
Spring postemergence oats	1/2-1 pt.	1 1/2-2 pts
Preharvest*** (dough stage) wheat, barley, oats, rye	1-2 pts.	2-3 pts
CORN*		
Preemergence	2-4 pts.	
Emergence*	1 pt.	1 1/2 pt.
Postemergence* up to 8 inches tall	1/2-1 pt.	
8 inches to tasseling (use only directed spray)	1 pt.	1 1/2-2 1/2 pts
Preharvest***	1-2 pts.	1 1/2-2 1/2 pts
SORGHUM (MILO)*		
Postemergence 6 to 8 inches tall	1/2-1 pt.	
8 to 15 inches tall (use only directed spray)	1 pt.	1 1/2-2 pts.
RICE		
	1-2 1/2 pts	2-3 pts.
SUGARCANE		
Fall, after harvesting or planting	2-4 pts.	
Spring, once or twice before close-in	2-4 pts.	
Summer	2 1/2 pts	

*Corn and sorghum hybrids vary in tolerance to 2,4-D; some are easily injured. Before spraying, get information on 2,4-D tolerance of specific hybrids and spray only those known to be resistant to 2,4-D injury. If plants are more than 8 inches tall, use directed spray and keep off corn and sorghum foliage.

**These higher rates may be needed to handle difficult weed problems in certain areas such as dry conditions, especially in areas west of the Mississippi River. However, do not use unless possible crop injury will be acceptable. Consult State Agricultural Experiment Station or Extension Service Weed Specialists for recommendations or suggestions to fit local conditions.

***Apply after the hard dough (corn) or dough stage (wheat) by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control tall weeds such as bindweed, cocklebur, dogbane, Jimsonweed, ragweed, sunflower, valerian and vines that interfere with harvesting.

FALLOW LAND AND STUBBLE

Annual Weeds — Use 1 to 2 qts. acre. Apply when weeds are actively growing.

Perennial Weeds — Use 2 to 3 qts. acre on weeds such as Canada thistle (apply in late bud or early bloom), field bindweed (50% or greater bloom) and other perennial weeds listed. Do not make application within 90 days of planting or until chemical has disappeared from soil.

PASTURE AND RANGELAND

Annual Weeds — Use 1 to 2 qts. acre. Apply when weeds are actively growing.

Perennial Weeds — Use 2 to 4 qts. acre when perennial weeds are translocating carbohydrates, i.e. Canada thistle (late bud to early bloom), bull thistle (bud stage), musk thistle (spring or fall in rosette or early bud stage), leafy spurge (4 qts.) (early to late bloom). Field bindweed (80% or greater bloom). Higher rates may cause temporary yellowing of grasses.

Do not use on bermgrass, alfalfa, clover, or other legumes. Do not use on newly seeded areas until grass is well established. Do not use from early boot to milk stage where grass seed production is desired.

OBSERVE THESE INTERVALS

1. A 7-day pregrazing interval for dairy cattle.
2. A 30-day preharvest interval for grass cut for hay.
3. A pre-slaughter interval for meat animals of 3 days. Withdrawal is not needed if two weeks or more have elapsed since application.

CONTROL OF SOUTHERN WILD ROSE

On rangelands, roadsides, and fence rows, use 1 gallon and spray thoroughly as soon as foliage is well developed. Two or more treatments may be required. On rangeland, apply a maximum of 6 quarts per acre per season. See grazing restrictions in pasture and rangeland section above.

GRASS SEED CROPS

Use 1 to 4 pints per acre in spring or fall to control broadleaf weeds in grass being grown for seed. Do not apply from early boot to the milk stage. Spray seeding grass only after the five-leaf stage, using 1/4 to 1 pint per acre to control small seedling weeds. After the grass is well established, higher rates of up to 4 pints can be used to control hard-to-kill annual or perennial weeds. For best results, apply when soil moisture is adequate for good growth.

NOTE: Do not use on bermgrass unless grass injury can be tolerated. See grazing restrictions in pasture and rangeland section above.

GENERAL WEED CONTROL

Roadsides, Vacant Lots, Fence Rows and Drainage Ditchbanks — Use 1/2% solution.

Turf — 1/2 to 1 gallon 100 gallons of water (4 to 8 tsp. per 1 gallon water).

Woody Plants — 1 1/2 to 2 fluid ounces per 1 gallon water (1 1/2% solution). Wet thoroughly all parts of the plant and stems to point of run-off.

TANK MIXTURES FOR SMALL GRAINS, FALLOW AND RANGE/NONCROPLAND

HI-DEP can be applied as a tank-mixture with Banvel*, Glean*, Lexone*, Sencor*, Roundup*, or Tordon* to broaden the spectrum of weed control. In order to assure maximum safety and weed control follow all precautions and limitations on this label and the labels of products used in tank mixtures with HI-DEP. Where a rate range is given, the rate should be varied depending on the weeds present.

SMALL GRAINS

HI-DEP - Glean*	1 pint A - 1/2-1/2 oz. A
HI-DEP - Lexone* or Sencor*	1 1/2 pints A - 1/2 lb. ai A

Feb. 1991: Glean has been withdrawn from CO, MT, ND, SD, MN, WY and NE Pannhandle. Still available in South Central Plains and Pacific Northwest.

FALLOW

HI-DEP - Banvel*	3 pints A - 1 pint A
HI-DEP - Roundup*	1-2 pints A - 1/2-1 pint A

RANGE AND NON-CROP AREA

HI-DEP - Banvel*	1-3 quarts A - 1-2 pints A
HI-DEP - Tordon*	1-2 quarts A - 1/2-2 pints A

STONEFRUIT, NUT AND PISTACHIO ORCHARDS

Broadleaf weeds. Use 1 1/2 qts. in 20 to 50 gallons of water per acre of ground sprayed. For band or spot treatment, calculate rates according to the actual portion of an acre treated. Apply as a directed spray onto the weeds to point of run-off when weeds are young and actively growing (pre-bud to early bud stage). Make up to two applications through the growing season as needed. Do not harvest stonefruits within 40 days of application. Do not harvest nuts and pistachios within 60 days of application. Do not graze or feed cover crops from treated orchards to livestock.

FILBERTS

Sucker Control. Mix 1 quart in 100 gallons of water plus 8 fl. ozs. of non-ionic spreader sticker (such as Ortho X-77 or R-11). Spray to run-off when suckers are 6 to 9 inches tall. Spray when needed from April through August. Use large orifice nozzles (.04 nozzle) and low tank pressure (30-35 psi) to produce large droplet size. Apply no more than four times per year. Do not harvest filberts within 45 days of last application. Do not allow livestock to graze in treated areas or the feeding of cover crops grown in treated orchards.

PRECAUTIONS IN APPLYING 2,4-D IN ORCHARDS

Apply only after irrigation and allow maximum time before the next irrigation. Do not apply around fruit trees or vines with hand gun. Use only flat, fan-type nozzles and low pressures — 20-30 psi. Use a fixed boom applicator which can be calibrated and which will deposit the spray uniformly. Avoid contact with fruit, foliage, stems or lower limbs of trees or vines. DO NOT spray bare ground. Apply precisely and uniformly to prevent damage to the trees or vines and to obtain satisfactory weed control. Do not apply during windy periods or extremely high temperatures. In California — not for use in desert valleys or on shallow or sandy soils. Allow maximum time after application and before next irrigation. Late fall applications after harvest and before frost preferred.

BROADLEAF WEED CONTROL IN NON-CROPLAND GRASS AREAS SUCH AS LAWNS, GOLF COURSES, CEMETERIES AND PARKS, AIRFIELDS, ROADSIDES, VACANT LOTS, DRAINAGE DITCHBANKS

Use 1 to 3 quarts per acre. Treat when weeds are young and growing well. Usually 2 quarts per acre will provide adequate weed control. Do not use on dichondra or other herbaceous ground covers. Do not use on creeping grasses such as bermgrass except for spot treating nor on freshly seeded turf until grass is well established. Reseeding of lawns should be delayed following treatment. With spring application, reseed in the fall; with fall application, reseed in the spring. Legumes are usually damaged or killed. Deep rooted perennial weeds such as bindweed and Canada thistle may require repeated applications.

SPOT TREATMENT/NON-CROP

Hand-held and high volume equipment. For control of weeds listed using knapsack sprayers or high volume equipment utilizing hand guns or other nozzle arrangements — Unless otherwise specified, make a 3/4% solution in water and apply to foliage as a coarse spray for general vegetation control. For hard-to-kill woody plants use a 1 1/4% solution. Applications should be made on a spray-to-wet basis and coverage uniform. Do not spray to point of run-off. Prepare the spray solution by mixing in water as per the following table.

DESIRED VOLUME	3/4%	3/4%	1%	1 1/4%
1 gallon	3/4 fl. oz. (4 tsp.)	1 fl. oz. (2 Tbs.)	1 1/4 fl. oz. (8 tsp.)	2 fl. oz. (4 Tbs.)
25 gallon	1 pt.	1 1/4 pt.	1 ct.	1 1/4 qt.
100 gallon	3 gal.	4 gal.	1 gal.	1 1/4 gal.

2 Tablespoons = 1 fluid ounce

1 Teaspoon = 1/2 Tablespoon = .17 fluid ounce

When using in knapsack sprayers, insure mixture is complete by shaking or inverting sprayer several times.

CONTROL OF WOODY PLANTS OR BRUSH AND BROADLEAF WEEDS ON ROADSIDES, DRAINAGE DITCHBANKS, RIGHTS-OF-WAY, RAILROADS, FIREBREAKS, FORESTS (Forest Site Prep), FENCEROWS, INDUSTRIAL SITES & OTHER SIMILAR NON-CROP AREAS

GROUND APPLICATIONS

HIGH VOLUME: Mix at the rate of 1 to 2 gallons per 100 gallons of water (1-2% solution). Rate per acre depends on the density of brush and/or weeds. For small broadleaf weeds, use the lower rate. Heavy dense stands of brush require the high rate with higher water volume. For small applications with small tank sprayers use at the rate of 1.25-2.5 fluid ounces per gallon of water.

To effectively control brush, all leaves, stems and suckers should be thoroughly wetted to the ground. Apply when plants come into full leaf (spring) to the time plants begin to go dormant. Best results are obtained when brush and broadleaf weeds are young and actively growing. Do not cut brush until the herbicide has translocated throughout the plant causing root death. DO NOT APPLY as a stand release or cover spray to established conifers as injury may result.

AERIAL APPLICATIONS

Forestry Site Preparation — For use in vegetation controlled burning programs, use 1/2 to 2 gallons of HI-DEP in tank mixes with other herbicides labeled for forestry site preparation (e.g. Garlon®, Tordon®, ARSENAL® Herbicide Applicators Concentrate). Use sufficient water to achieve uniform wetting of target brush species. Do not exceed 25 gallon total spray per acre.

Utility & Pipeline Rights-of-Way — Use 1/2 to 2 gallons of HI-DEP in tank mix combination with other herbicides labeled for rights-of-way sites and apply in a total spray volume of 5 to 30 gallons per acre.

TANK MIXTURES FOR INDUSTRIAL/NON-CROPLAND AREAS

HI-DEP can be applied as a tank mixture with other recommended herbicides such as Garlon®, Tordon®, and Banvel® to broaden the spectrum of control. In order to assure maximum safety and weed control, follow all precautions and limitations on this label and the labels of products used in tank mixtures with HI-DEP. Where a rate range is given, the rate should be varied according to the density and target species.

PRODUCTS	RATES
HI-DEP - Garlon® 3A	1/2-2 ga. A - 1/2-1 gal. A
HI-DEP - Garlon® 4E	1/2-2 ga. A - 2-4 qts. A
HI-DEP - Tordon® K	1/2-2 ga. A - 1/2-4 qts. A
HI-DEP - Banvel®	1/2-2 ga. A - 1 qt.-2 ga. A

LEAFY SPURGE CONTROL IN COLORADO, IDAHO, MINNESOTA, MONTANA, NEBRASKA, NORTH DAKOTA, SOUTH DAKOTA, WASHINGTON AND WYOMING

HI-DEP is recommended for use in combination with Tordon® or Banvel® for the suppression and/or control of leafy spurge on industrial, non-cropland sites in Colorado, Idaho, Minnesota, Montana, Nebraska, North Dakota, South Dakota, Washington and Wyoming.

HOW TO USE

Apply 1 to 2 quarts of HI-DEP in combination with 1 quart of Tordon®, or 2 quarts of HI-DEP plus 2 quarts of Banvel®, or 2 quarts of HI-DEP plus 1 pint of Tordon® plus 1 quart of Banvel®.

Rates are on a per acre basis. Mix with water, 1 to 10 gallons per acre minimum with conventional equipment. Use nozzle systems capable of spraying correct gallonage. Add a non-ionic surfactant such as Gordon's Staypart® at 0.25% by volume (1 quart per 100 gallons of solution).

IMPORTANT: BEFORE USING HI-DEP, TORDON® AND OR BANVEL® IN THESE COMBINATIONS, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

FOREST-TREE INJECTION

To control unwanted hardwood trees make injections as near the root collar as possible using one injection per inch of trunk diameter at breast height. For resistant species such as hickory, injections should overlap. For best results injections should be made during the growing season — May 15 to October 1.

For Concentrate Injection — Use 1 to 2 ml. of concentrate per injection. The injector bit must penetrate the inner bark.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

HI-DEP® and STAYPART® are registered trademarks of PBI GORDON CORPORATION.

Garlon® and Tordon® are registered trademarks of DowElanco.

Banvel® is a registered trademark of Sandoz Crop Protection.

Arsenal® is a registered trademark of American Cyanamid Company.

Glean® and Lexone® are registered trademarks of E. I. DuPont de Nemours and Company, Inc.

Sencor® is a registered trademark of Bayer AG, Germany.

Roundup® is a registered trademark of Monsanto Agricultural Products Company.

808 992 AP090991
EPA REG NO 2217-703

MANUFACTURED BY

G pbi / Gordon
corporation
KANSAS CITY, MISSOURI 64101

SPECIMEN LABEL: READ CURRENT LABEL ON PRODUCT CONTAINER BEFORE USE.

Specimen Label

Curtail*

Herbicide

For Selective Control of Broadleaf Weeds in Wheat and Barley Not Underseeded with a Legume, Fallow Cropland, Grasses Grown for Seed and Conservation Reserve Program (CRP) Acres

Active Ingredient(s):

Clopyralid (3,6-Dichloro-2-pyridinecarboxylic acid) as alkanolamine salts (of the ethanol and isopropanol series)	7.5%
2,4-Dichlorophenoxyacetic Acid as alkanolamine salts (of the ethanol and isopropanol series)	38.4%

Inert Ingredient(s):54.1%

Acid Equivalents:

3,6-Dichloro-2-pyridinecarboxylic acid 3.9% - 0.38 lb/gal

2,4-Dichlorophenoxyacetic acid 20.9% - 2.0 lb/gal

E.P.A. Registration No. 62719-48

E.P.A. Est. 464-MI-1

KEEP OUT OF REACH OF CHILDREN

DANGER

PELIGRO:

PRECAUCION AL USUARIO:

Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUSES EYE DAMAGE AND SKIN

IRRITATION • HARMFUL IF

SWALLOWED

Do Not Get In Eyes, On Skin Or On Clothing

• **Wear Goggles Or Face Shield When Handling**

• **When Handling This Product, Wear Chemical**

Resistant Gloves • Wash Thoroughly With

Soap And Water After Handling • Remove

Contaminated Clothing And Wash Before

Reuse

STATEMENTS OF PRACTICAL TREATMENT: If in eyes, flush with plenty of water for at least 15 minutes. Get medical attention. If on skin, wash with plenty of soap and water. Get medical attention if irritation persists. If swallowed, induce vomiting immediately by giving two glasses of water and sticking finger down throat. Call a physician. Do not induce vomiting or give anything by mouth to an unconscious person.

Environmental Hazards

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters. See additional precautionary statements elsewhere on this label.

NOTICE

Read the entire label. Use only according to label directions.

Before buying or using this product, read Warranty Disclaimer and Limitation of Remedies sections elsewhere on this label.

IN CASE OF AN EMERGENCY

endangering life or property involving this product, call collect 517-636-4400

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing

*Curtail

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

CURTAIL Herbicide is recommended for selective, postemergence control of the following broadleaf weeds in wheat and barley not underseeded with a legume, fallow cropland (including summer fallow, post-harvest, and set-aside acres) and land in the Conservation Reserve Program (CRP).

alfalfa	mustard, tumble (Jim Hill)
artichoke, Jerusalem	mustard, wild
buckwheat, wild	nightshade, cutleaf
buffaloburn	nightshade, hairy
burdock, common	pennycress, field (fanweed)
chamomile, false (scentless)	pigweed, redroot
chamomile, mayweed	pineappleweed
(dogfennel)	plantain
clover, sweet	radish, wild
clover, red	ragweed, common
cocklebur, common	ragweed, giant
coffeweed	salsify, meadow (goatsbeard)
cornflower (bachelor button)	shepherdspurse
dandelion	sicklepod
dock, curly	smartweed, Pennsylvania
flixweed ¹	sorrel, red
groundsel, common	sowthistle, annual
hawksbeard, narrowleaf	sowthistle, perennial ¹
horseweed	starthistle, yellow
jimsonweed	sunflower, common
knapweed, diffuse	tansymustard, pinnate ¹
knapweed, Russian ¹	thistle, Canada
knapweed, spotted	thistle, musk
kochia (2-4 leaf) ¹	thistle, Russian (1-3 leaf)
ladysthumb	velvetleaf
lambsquarters, common	vetch
lettuce, prickly	volunteer beans
locoweed, white	volunteer lentils
locoweed, lambert	volunteer peas
marshelder	

¹ These weeds may only be suppressed. Suppression is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during and after the time of treatment. For perennial weeds, CURTAIL Herbicide will control the initial topgrowth and inhibit regrowth during the season of application (season-long control). At higher use rates shown on this label, CURTAIL Herbicide may cause a reduction in shoot regrowth in the season following application, however, plant response may be inconsistent due to inherent variability in shoot regrowth from perennial root systems.

Timing: Apply to actively growing weeds. Extreme growing conditions such as drought or near freezing temperatures prior to, at and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Only weeds which are emerged at the time of application will be affected. Wet foliage at the time of application may decrease control. The CURTAIL Herbicide treatment will be rainfast within 6-8 hours after application.

Rate: Generally, lower labeled application rates will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species, perennials and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands and/or larger weeds) the higher rates will be needed. Weeds in fallow or other areas where crop competition is not a factor will generally require higher rates to obtain control or suppression.

Coverage: Adequate spray coverage and drift control are important. Obtaining a balance between spray coverage and drift control may sometimes be difficult but can be achieved, provided the applicator understands the factors affecting coverage and drift. Factors affecting spray coverage include spray volume, crop canopy and weed density. As crop canopy

and weed density increase, spray volume should be increased to obtain equivalent weed control. Refer to manufacturer's recommendations for information on the relationship between gallons per acre, spray pressure, sprayer speed, nozzle type and arrangement, nozzle height above the target canopy, droplet size and drift potential for respective application equipment. Do not apply less than 2 and not more than 40 gallons per acre total spray volume. For best results, apply 5 or more gallons per acre by air and 10 or more gallons per acre by ground. Reducing total spray volume may result in decreased coverage and weed control. Use enough total spray volume and a delivery system to provide thorough coverage and a uniform spray pattern. Do not apply where spray drift may be a problem due to proximity of susceptible crops or other desirable plants.

Use of Surfactants: Addition of wetting and/or penetration agents is not usually necessary when using CURTAIL Herbicide; however, if a surfactant will be added to the spray solution, use a nonionic surfactant approved for use on growing crops, such as X-77, of at least 80% active ingredient and do not exceed 4 pints per 100 gallons of spray solution (0.5% v/v). Adding a surfactant to the spray mixture may increase effectiveness on weeds but may reduce selectivity to the crop, particularly under conditions which promote plant stress.

Use With Sprayable Liquid Fertilizer Solutions: CURTAIL Herbicide is compatible with most non-pressurized liquid fertilizer solutions, however, if these solutions are to be sprayed with CURTAIL Herbicide, a compatibility test (jar test) should be made prior to mixing. Jar tests are particularly important when a new batch of fertilizer or pesticide is used, when water source changes, or when tank mixture ingredients or concentrations are changed. A compatibility test is performed by pouring the spray components (in the desired order and proportions) into a clear glass jar before mixing in the spray tank. Use of a compatibility aid such as Unite or Compex may help obtain and maintain a uniform spray solution during mixing and application. Agitation in the spray tank must be vigorous to compare with jar test agitation. For best results, liquid fertilizer rates should not exceed 50% of the total spray volume. Premix CURTAIL Herbicide with water and add to the liquid fertilizer/water mixture while agitating contents of the spray tank. Apply the spray the same day it is prepared while maintaining continuous agitation.

NOTE: Foliar-applied liquid fertilizers can cause yellowing or leaf burn of crop foliage.

Wheat and Barley

Timing: Apply CURTAIL Herbicide in the spring to actively growing wheat or barley once 4 leaves have unfolded on the main stem and tillering has begun up to the jointing stage (first node of main stem detectable). To control or suppress weeds listed above, make application after maximum emergence of the target weeds but before they exceed 3 inches in height or diameter (for rosettes). To obtain season-long control of perennial weeds such as Canada thistle, apply after the majority of the weed's basal leaves have emerged from the soil, but before bud stage. A later timing of application (when the grain is between the jointing and boot stages) may be used to treat later-emerging weeds, however, do not apply unless the risk of injury is acceptable. Do not apply after the boot stage.

NOTE: Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 1 week after treatment. Do not harvest hay from treated grain fields.

Rate: Apply 2 to 2²/₃ pints of CURTAIL Herbicide per acre. The higher rate may be used when the condition of the weeds and/or crop at the time of treatment may prevent optimum control.

NOTE: Higher rates of CURTAIL Herbicide or CURTAIL Herbicide applications following spring postemergence treatments with 2,4-D or MCPA may increase the risk of crop injury.

Tank Mixtures for Wheat and Barley: This product may be tank mixed with the following herbicides for control of additional weeds in wheat and barley. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions. When tank mixing, use in accordance with the most restrictive of label limitations and precautions. No label dosages should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Active Ingredient	Product	Formulation	Amount of Product Per Acre
bromoxynil ¹	Buctril	2 lb/gal	3/4 to 1 pt
	Buctril 4EC	4 lb/gal	3/8 to 1/2 pt
	Buctril ME4		
chlorsulfuron dicamba ^{1,2} difenzoquat diuron	Glean	75% DF	1/8 to 1/4 wt oz
	Barvel	4 lb/gal	1/8 to 1/4 pt
	Avenge	2 lb/gal	2 1/2 to 4 pt
	Direx 4L	4 lb/gal	1 to 1 1/2 pt
	Diuron 4L		
MCPA or 2,4-D ² metribuzin	Diuron 80 WDG	80% DF	1/2 to 1 lb
	Diuron DF	80% WP	Up to 1/2 pt
	Lexone DF	75% DF	2 1/2 to 4 wt oz
metsulfuron methyl terbutryn thifensulfuron methyl DPX L-5300	Sencor DF		
	Ally	60% DF	1/10 wt oz
	Igran 80 WP	80% WP	6 to 10 wt oz
	Harmony	75% DF	1/3 to 1/2 wt oz
	Express	75% DF	1/6 to 1/4 wt oz

¹ Buctril or Barvel tank mixes with CURTAIL Herbicide may be useful in broadening the annual weed control spectrum but may reduce control of perennials, such as Canada thistle.

² Note: Do not tank mix CURTAIL Herbicide with 2,4-D or dicamba unless the risk of crop injury is acceptable.

Fallow Cropland

Timing: CURTAIL Herbicide can be applied either postharvest, in the spring/summer (during fallow period) or to set-aside acres to control or suppress weeds listed above (refer to rotation restrictions). Apply to young, emerged weeds under conditions that promote active growth. For best results on tough perennial weeds such as Canada thistle, apply after the majority of the basal leaves have emerged, but before bud stage. Later applications may result in less consistent control. Extreme growing conditions (such as drought or near freezing temperatures) prior to, at, and following the time of application may reduce weed control.

For best results, wait 14 to 20 days after application before cultivating or fertilizing with shank-type applicators to allow for thorough translocation. To avoid potential phytotoxicity, allow at least 20 days after application before seeding to wheat, barley or grasses.

Rate: Apply 4 pints of CURTAIL Herbicide per acre. Use the higher rate on perennial weeds or when the condition of the weeds at the time of treatment may prevent optimum control.

Tank Mixtures For Fallow Cropland: To improve control of certain broadleaf weeds, CURTAIL Herbicide at 2 pints per acre may be applied with up to 1.5 lb ae per acre additional 2,4-D. When mixing, read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions. When tank mixing, use in accordance with the most restrictive of label limitations and precautions. No label doses should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Grasses Grown for Seed

Timing: Apply only to established grasses before the boot stage. Applications in the boot stage and beyond will result in increased injury. Do not apply to bentgrass unless injury can be tolerated.

Rate: Use 2 to 4 pints of CURTAIL Herbicide per acre for control of annual weeds and Canada Thistle. The potential for crop injury exists due to the 2,4-D component of this product and must be balanced against the benefits of improved weed control. Potential for crop injury increases with higher rates. Re-treat as necessary, but do not exceed 4 pints per acre of CURTAIL per season.

NOTE: Do not graze (dairy) cattle in treated areas for 14 days after application. Remove meat animals from freshly treated areas for 7 days before slaughter. Withdrawal is not needed if 2 weeks or more have elapsed since application. Do not cut treated grass for hay within 30 days after application.

Conservation Reserve Program (CRP) For Seeding To Permanent Grasses Only

Do not use CURTAIL Herbicide if legumes or bentgrass are a desired cover during CRP.

Grass grown on CRP acres cannot be grazed or used for hay.

Conditions that stress grasses, such as drought, will increase potential for injury to the grass at all stages of growth. Do not use in newly seeded areas until grass is established.

Timing: CURTAIL Herbicide can be applied when perennial grasses have become established (tillering, with at least 1.5 inches secondary roots, and good vigor). For control of weeds such as musk thistle, Canada thistle and knapweed (diffuse, spotted and Russian), apply to actively growing weeds after the majority of the basal leaves have emerged, but before bud stage. Later applications may result in less consistent control.

Rate: Apply 2 to 5 quarts of CURTAIL Herbicide per acre.

USE PRECAUTIONS

Apply only once per 12-month period, except for grass grown for seed (see Directions For Use). A fallow treatment that precedes or follows a small grain application is also allowed.

This product can affect susceptible broadleaf plants directly through foliage and indirectly by root uptake from treated soil. Therefore, do not apply CURTAIL Herbicide directly to or allow spray drift to come in contact with vegetables, flowers, grapes, tomatoes, potatoes, beans, lentils, peas, alfalfa, sunflowers, soybeans, safflower, or other desirable broadleaf crops and ornamental plants or soil where these sensitive crops will be planted the same season.

Do not contaminate irrigation ditches or water used for irrigation or domestic purposes.

Wheat, barley, oats, grasses, or sugar beets (including beets grown for seed) may be planted less than 12 months after treatment.

Alfalfa, asparagus, canola, cole crops, mint, onions, safflower, and strawberries may be planted 12 months after treatment. Dry beans, soybeans and sunflowers may also be planted at 12 months after treatment, except in soils with less than 2% organic matter receiving less than 15 inches of natural precipitation in the 12 month period following treatment. For these areas see SPECIAL CONDITIONS below.

To avoid potential phytotoxicity all other crops including peas, lentils, potatoes, and broadleaf crops grown for seed should not be planted for 18 months after treatment unless the risk of injury is acceptable.

SPECIAL CONDITIONS: CURTAIL residues in plant tissues which have not completely decayed may affect succeeding susceptible crops. In areas defined previously as low in organic matter and precipitation, sensitive crops such as dry beans, soybeans, and sunflowers may be injured when planted 12 months after treatment. Unless the risk of injury is acceptable, these crops should not be planted until 18 months after treatment. The potential for injury may be reduced by burning, removal, or incorporation of treated crop residues with a minimum of 2 supplemental Fall irrigations.

Avoid Spray Drift: Applications should be made to avoid spray drift since very small quantities of the spray, which may not be visible, may severely injure susceptible crops during both growing and dormant periods. Use coarse sprays to minimize drift since, under adverse weather conditions, fine spray droplets may drift a mile or more. Fine droplets can present a drift hazard. A drift control or deposition agent such as Nalco-Trol may be used with this product to aid in reducing spray drift. If used, follow all use recommendations and precautions on the product label.

Ground Application: To minimize spray drift, apply CURTAIL Herbicide in a total spray volume of 10 or more gallons per acre as a large-droplet, low pressure spray. Refer to manufacturer's recommendations for additional information on gallons per acre, spray pressure, sprayer speed, nozzle types and arrangements, nozzle heights above the target canopy, etc., for respective application equipment. Spot treatments should only be applied with a calibrated boom to prevent misapplication. With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible; by applying no more than 20 gallons of spray per acre; by using no more than 30 pounds spraying pressure with large droplet-producing nozzle tips; by spraying when wind velocity is low; and by stopping all spraying when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone-type insecticide or other nozzles that produce a fine-droplet spray.

Aerial Application: With aircraft, drift can be lessened by applying a coarse spray; by using no more than 30 pounds spray pressure at the nozzles; by using straight-stream nozzles directed straight back; by using a spray boom no longer than 3/4 the wing span of the aircraft; and by spraying only when wind velocity is less than 6 mph.

Do Not Apply By Aircraft When An Air Temperature Inversion Exists. Such a condition is characterized by little or no wind and with lower air temperature near the ground than at higher levels. The use of a smoke device on the aircraft or continuous smoke column at or near site of application will indicate air direction and velocity, and whether a temperature inversion is present, which is shown by layering of the smoke.

Do not move treated soil and avoid situations where treated soil particles may blow into areas where susceptible crops are grown. Violent windstorms may move soil particles. If this product is on soil particles and they are blown onto susceptible plants, visible symptoms may appear. Serious injury is unlikely. The hazard of movement of this product on dust is reduced if treated fields are irrigated or if rain occurs shortly after application.

Straw from treated areas, or manure from animals which have grazed treated areas, cannot be used for composting or mulching on ground where susceptible crops may be grown the following season. To promote herbicide decomposition, plant material should be evenly incorporated or burned.

Do not use in a greenhouse. Excessive amounts of this herbicide in the soil may temporarily inhibit seed germination or plant growth.

Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

Do not transfer livestock from treated grazing areas onto sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated pasture. Otherwise, urine may contain enough clopyralid to cause injury to sensitive broadleaf plants.

Mixing and Loading: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

PRECAUTIONARY STATEMENTS

Environmental Hazards

Clpyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clpyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store above 10°F or warm and agitate before use.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal (Metal): Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Container Disposal (Plastic): Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General: Consult federal, state, or local disposal authorities for approved alternative procedures.

Be sure that use of this product conforms to all application regulations.

WARRANTY DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below.

DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

08103-L1

A790

DowElanco

Indianapolis, Indiana 46268 U.S.A.

*Trademark of DowElanco

Specimen Label

Stinger*

Herbicide

For Selective Postemergence Control of Broadleaf Weeds in Sugar Beets, Field Corn, Wheat, Barley, and Oats Not Underseeded With a Legume, Christmas Tree Plantations and Nurseries, Grasses Grown for Seed, Fallow Cropland, Rangeland and Permanent Grass Pastures, Non-Cropland Areas, and Conservation Reserve Program (CRP) Acres

Active Ingredient(s):

Clopyralid (3,6-dichloro-2-pyridinecarboxylic acid) as the monoethanolamine salt 40.9%
Inert Ingredient(s): 59.1%

ACID EQUIVALENT: 3,6-dichloro-2-pyridinecarboxylic acid, 31% - 3 lb/gal

E.P.A. Registration No. 62719-73

E.P.A. Est. 464-MI-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTION:

PRECAUCION AL USUARIO:

Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

**CAUSES EYE INJURY • HARMFUL IF
INHALED OR ABSORBED THROUGH
SKIN**

Avoid Contact With Eyes, Skin Or Clothing

• Avoid Breathing Spray Mist • Wash
Thoroughly With Soap And Water After
Handling • Remove Contaminated
Clothing And Wash Before Reuse

STATEMENTS OF PRACTICAL TREATMENT: If in eyes,
flush with plenty of water. Get medical attention if irritation
persists. If on skin, wash with plenty of soap and water.
Get medical attention.

Physical or Chemical Hazards

**COMBUSTIBLE • Do Not Use or Store Near
Heat or Open Flame. Do Not Cut or Weld
Container.**

Environmental Hazards

Do not contaminate water when disposing of equipment
washwaters. Do not contaminate water used for irrigation or
domestic purposes. Do not apply directly to any body of
water or wetlands.

See additional precautionary statements elsewhere on
this label.

NOTICE

Read and understand the entire label before using. Use only
according to label directions.

Before buying or using this product, read "WARRANTY
LIMITATIONS AND DISCLAIMER" elsewhere on this label. If
terms are not acceptable, return unopened package at
once to seller for full refund of purchase price paid.
Otherwise, use by the buyer or any other user constitutes
acceptance of the terms under WARRANTY LIMITATIONS
AND DISCLAIMER.

IN CASE OF AN EMERGENCY

endangering life or property involving this
product, call collect 517-636-4400

AGRICULTURAL CHEMICAL

**Do Not Ship or Store with Food, Feeds,
Drugs, or Clothing**

Stinger*

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

STINGER Herbicide is recommended for selective, postemergence control of the following broadleaf weeds in sugar beets, field corn, wheat, barley, and oats not underseeded with a legume, fallow cropland, rangeland and permanent grass pastures, grasses grown for seed, Christmas trees, conservation reserve program (CRP) acres, and non-cropland areas including fence rows, around farm buildings, and equipment pathways.

artichoke, Jerusalem	marshelder
buckwheat, wild	nightshade, Eastern black
buffalobur ¹	nightshade, cutleaf
burdock, common	nightshade, hairy
chamomile, false (scentless)	oxeye daisy
chamomile, mayweed	pineappleweed
(dogfennel)	ragweed, common
clover, sweet	ragweed, giant
clover, red	salsify, meadow
cocklebur, common	(goatsbeard)
coffeeweed	sicklepod
cornflower (bachelor button)	smartweed, green ¹
dandelion	sorrel, red
dock, curly	sowthistle, annual
groundsel, common	sowthistle, perennial ¹
hawksbeard, narrowleaf	starthistle, yellow
horseweed	sunflower
jimsonweed	thistle, Canada
knapweed, diffuse	thistle, musk
knapweed, Russian ¹	vetch
knapweed, spotted	volunteer alfalfa
ladythumb ¹	volunteer beans
lettuce, prickly	volunteer lentils
locoweed, white	volunteer peas
locoweed, Lambert	

¹ These weeds may only be suppressed. Suppression is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment. For perennial weeds, STINGER Herbicide will control the initial topgrowth and inhibit regrowth during the season of application (season-long control). At higher use rates shown on this label, STINGER Herbicide may cause a reduction in shoot regrowth in the season following application; however, plant response may be inconsistent due to inherent variability in shoot regrowth from perennial root systems.

Timing: Apply to actively growing weeds. Extreme growing conditions such as drought or near freezing temperatures prior to, at, and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Only weeds which are emerged at the time of application will be affected. Wet foliage at the time of application may decrease control. The treatment with STINGER Herbicide will be rainfast within 6-8 hours after application.

Rate: Generally, lower labeled application rates will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species, perennials, and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands, and/or larger weeds), the higher rates will be needed. Weeds in fallow or other areas where crop competition is not a factor will generally require higher rates to obtain control or suppression.

Coverage: Adequate spray coverage and drift control are important. Obtaining a balance between spray coverage and drift control may sometimes be difficult but can be achieved provided the applicator understands the factors affecting coverage and drift. Factors affecting spray coverage include spray volume, crop canopy, and weed density. As crop canopy and weed density increase, spray volume should be increased to obtain equivalent weed control. Refer to manufacturer's recommendations for information on the relationship between gallons per acre, spray pressure, sprayer speed, nozzle type and arrangement, nozzle height above the target canopy, droplet size, and drift potential for respective application equipment. Use equipment and nozzle types which are designed for herbicide application. Do not apply less than 2 and not more than 40 gallons per acre total spray volume. For best results, apply 5 or more gallons per acre by air and 10 or more gallons per acre by ground. Reducing total spray volume may result in decreased coverage and weed control. Use enough total spray volume and a delivery system to provide thorough coverage and a uniform spray pattern. Do not apply where spray drift may be a problem due to proximity of susceptible crops or other desirable plants.

Use of Adjuvants: Addition of surfactants, crop oils, or other adjuvants is not usually necessary when using STINGER Herbicide. Adding a surfactant to the spray mixture may increase effectiveness on weeds but may reduce selectivity to the crop, particularly under conditions which promote plant stress. If an adjuvant is added to the spray solution, follow all manufacturer use guidelines.

Tank Mixes: When tank mixing, read and follow the label of each tank-mix product used for precautionary statements, directions for use, weeds controlled, and geographic and other restrictions. Use in accordance with the most restrictive of label limitations and precautions. No label dosages should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

WEED CONTROL GUIDELINES ¹ Amount of STINGER Per Acre X Use Site ²

Weed Species	Growth Stage	Sugar Beet, Christmas Trees	Wheat, Barley, Oats	Grasses For Seed	Fallow, Non-Crop	Range & Pasture, CRP	Field Corn
clover cocklebur sunflower ragweeds jimsonweed volunteer soybean vetch marshelder	Up to 5 leaf	¼ - ½ pt	¼ - ½ pt	¼ - ½ pt	¼ - ½ pt	½ - ¾ pt	¼ - ½ pt
wild buckwheat nightshade sp. buffalobur smartweeds (suppression)	1-3 leaf stage, but before vining 2-4 leaf 2-3 leaf	½ pt					
Canada thistle sowthistle (suppression) knapweeds, spotted/diffuse knapweeds, Russian (suppression)	rosette to prebud up to bud stage	½ - ¾ pt ¾ pt	¼ - ½ pt —	½ - ¾ pt ¾ - 1 pt	½ pt —	¾ - 1 pt ¾ - 1 pt 1 - 1½ pt	½ - ¾ pt — —

¹ This table is intended as a reference only. For complete instructions see the body of the text.

² Use the lower rate for light to moderate infestations and good growing conditions and the higher rate for dense infestations or under poor growing conditions such as drought.

Sugar Beets

STINGER Herbicide is recommended for the control of various annual and perennial broadleaf weeds infesting sugar beets. Apply ¼ to ¾ pint of STINGER Herbicide per acre with ground equipment as a broadcast foliar spray. Apply in 10 or more gallons total spray volume per acre when the sugar beets are in the cotyledon to 8 leaf stage of growth and the weeds are young and actively growing. Re-treat as necessary but do not exceed ¾ pint of STINGER Herbicide per acre per season. Do not apply within 105 days before harvest of beet roots and tops. Do not apply to sugar beets grown for seed.

STINGER Herbicide may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast rate per treated acre}}{\text{per treated acre}} = \frac{\text{Band rate per treated acre}}{\text{per treated acre}}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast volume per treated acre}}{\text{per treated acre}} = \frac{\text{Band volume per treated acre}}{\text{per treated acre}}$$

For annual weed control spray ¼ - ½ pt of STINGER Herbicide per acre on weeds up to the 5 leaf growth stage. Wild buckwheat applications should be made at the 1-3 leaf stage, before vining begins.

For the most effective control of perennials such as Canada thistle and sowthistle, apply ½ - ¾ pint of STINGER Herbicide per acre as a broadcast treatment to the entire infested area. Apply when the majority of basal leaves have emerged, but before the bud stage. Cultivation can disrupt translocation to the roots of perennials such as Canada thistle. For best results do not cultivate thistle patches.

To promote herbicide efficacy wait a minimum of 7 days after application before flood or furrow irrigation.

Tank Mixes: To control additional broadleaf weeds and provide consistent control of difficult weeds such as wild buckwheat, tank mix ¼ - ½ pint of STINGER Herbicide per acre with 2-6.5 pints of Betamix or Betanex. For best results, apply ¼ pint of STINGER tank mixed with 2-6.5 pints of Betamix or Betanex followed 1-2 weeks later by a second application of ¼ - ½ pint of STINGER per acre tank mixed with Betamix or Betanex. Note: Do not add additional adjuvants when employing a Betamix or Betanex tank mix with STINGER Herbicide due to increased potential for crop injury. (See tank mix section under Directions for Use.)

STINGER Herbicide may be tank mixed with grass herbicides such as Poast for grassy weed control. Be sure to include crop oil or Dash surfactant to optimize grass weed control. (See tank mix section under Directions for Use.)

Field Corn

STINGER Herbicide is recommended for postemergence control of Canada thistle, Jerusalem artichoke, annual sowthistle, common sunflower, common cocklebur, giant and common ragweed, jimsonweed and other broadleaf weeds infesting field corn. Apply STINGER Herbicide at suggested timing and rates for field corn as indicated below.

Apply STINGER Herbicide to actively growing broadleaf weeds any time after corn emergence through 24 inch tall corn. Apply with ground equipment as a postemergence broadcast or directed spray in 10 or more gallons of spray volume per acre to ensure uniform and thorough spray coverage of the weed foliage. Use only spray nozzles designed for herbicide application. The use of flat fan nozzles provides the best coverage and distribution of chemical on the plant foliage. Use spray pressures (at the boom) which nozzle manufacturers recommend to obtain desired spray volume. Use higher spray pressures and volumes when weed foliage is dense.

For the most effective control of Canada thistle, apply $\frac{1}{2}$ to $\frac{3}{4}$ pint of STINGER Herbicide per acre as a broadcast treatment to the entire infested area. Apply when the majority of the basal leaves have emerged, thistles are at least 4 inches in diameter or height, but before bud stage. Use the higher rate listed for dense patches or when greater residual control is desired. Cultivation can disrupt translocation to the roots of Canada thistle. For best results, do not cultivate prior to application, and wait 14 to 20 days after application before cultivating to allow for thorough translocation.

For control of common cocklebur, giant ragweed, common ragweed, sunflower, other annual weeds and Jerusalem artichoke, apply $\frac{1}{4}$ to $\frac{1}{2}$ pint of STINGER Herbicide on weeds up to the 5 leaf stage.

Hand-Held Sprayers - Applications should be made on a spray-to-wet basis with spray coverage uniform and complete. Do not spray to the point of runoff. Prepare the desired volume of spray solution by mixing the amount of STINGER Herbicide with water as shown in the following table.

Desired Volume Spray Solution	Amount of STINGER
1 gallon	$\frac{1}{4}$ fl. ounce
25 gallons	$\frac{1}{2}$ pint
100 gallons	1 $\frac{1}{2}$ pints

Restrictions: Re-treat as necessary, but do not apply more than $\frac{3}{4}$ pint of STINGER Herbicide per acre per year. Do not apply to field corn greater than 24 inches tall. Do not allow livestock to graze treated areas or harvest treated corn silage as feed within 40 days after last treatment.

Christmas Tree Plantations And Nurseries

Timing: STINGER Herbicide can be safely applied over the top of actively growing: balsam fir, blue spruce, Douglas fir, Fraser fir, grand fir, lodgepole pine, noble fir, ponderosa pine, Scotch pine, and white pine. For the Pacific Northwest: do not apply in the first year of transplanting. Apply only to trees transplanted at least one year. (Some needle curling has been observed on 1st year transplants.) Apply to actively growing weeds. For control of annual weeds apply STINGER Herbicide up to the 5 leaf growth stage (for wild buckwheat application at 3-5 leaf, but before vining, is recommended). For control of weeds such as Canada thistle and knapweeds, apply after the majority of the basal leaves have emerged, but before bud stage. Later application may result in less consistent control.

Rate: Apply $\frac{1}{4}$ - $\frac{1}{2}$ pint of STINGER Herbicide per acre for control of annual weeds. Apply $\frac{1}{2}$ - $\frac{3}{4}$ pint of STINGER Herbicide per acre for difficult to control weeds such as Canada thistle and knapweeds. Apply as a broadcast or band application in a minimum of 10 gallons per acre by ground application. Use the formula under "sugar beets" to determine the appropriate rate and volume per treated acre. Apply as often as needed, but do not exceed $\frac{3}{4}$ pint per acre. Do not exceed $\frac{1}{2}$ pint per acre for blue spruce. Tree injury may occur with the addition of a surfactant or crop oil with STINGER Herbicide. Do not use unless previous experience shows injury is tolerable.

Grasses Grown For Seed

Timing: Apply only to established grasses before the boot stage. Applications in the boot stage and beyond can result in increased injury. Do not apply to bentgrass unless injury can be tolerated.

Rate: Use $\frac{1}{4}$ to $\frac{3}{4}$ pint of STINGER Herbicide per acre for control of annual weeds and Canada thistle. Re-treat as necessary, but do not exceed $\frac{3}{4}$ pint of STINGER per acre per season.

Fallow Cropland

Timing: STINGER Herbicide can be applied either postharvest, in the spring/summer (during fallow period), or to set-aside acres to control or suppress weeds listed above (refer to rotation restrictions). Apply to young, emerged weeds under conditions that promote active growth. For best results on perennial weeds such as Canada thistle, apply after the majority of the basal leaves have emerged, but before bud stages. Later applications may result in less consistent control. Extreme growing conditions (such as drought or near freezing temperatures) prior to, at, and following the time of application may reduce weed control.

For best results, wait 14 to 20 days after application before cultivating or fertilizing with shank-type applicators to allow for thorough translocation.

Rate: Apply $\frac{1}{4}$ to $\frac{1}{2}$ pint of STINGER Herbicide per acre. Use the higher rate on perennial weeds or when the condition of the weeds at the time of treatment may prevent optimum control.

Tank Mixtures for Fallow Cropland: To improve control of certain broadleaf weeds, STINGER Herbicide may be applied with 0.5-2.0 lb ae per acre 2,4-D.

Non-Cropland

For use on non-cropland areas such as fencerows, around farm buildings and equipment pathways. For control of broadleaf weeds, apply $\frac{1}{4}$ to $\frac{1}{2}$ pints of STINGER Herbicide per acre. The lower rate of $\frac{1}{4}$ pint per acre provides acceptable control of weeds only under highly favorable growing conditions and when plants are 1-3 inches tall. Apply $\frac{1}{2}$ pint per acre when weeds are 3 to 6 inches tall or under dry conditions. Where Canada thistle or knapweeds are the primary pest, best results are obtained by applying $\frac{3}{4}$ to 1 $\frac{1}{2}$ pints of STINGER Herbicide per acre. To improve spectrum of activity or to increase activity against taller weeds, STINGER may be tank mixed with 0.5 to 2.0 lbs. ae per acre of 2, 4-D amine or low volatile ester.

Wheat, Barley And Oats

Apply $\frac{1}{4}$ to $\frac{1}{2}$ pint of STINGER Herbicide per acre from the 3 leaf stage up to early boot stage of growth. For control of perennial weeds such as Canada thistle, $\frac{1}{2}$ pint of STINGER Herbicide per acre should be used. Russian knapweed will only be suppressed at this rate.

NOTE: Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 1 week after treatment. Do not harvest hay from treated grain fields.

APPLICATION PERIOD FOR STINGER HERBICIDE

WHEAT, BARLEY AND OATS

Tank Mixtures for Wheat, Barley and Oats: Tank mix $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre of STINGER Herbicide with the herbicides listed below for the control of additional weeds.

Active Ingredient	Product	Formulation	Amount of Product Per Acre
bromoxynil	Buctril	2 lb/gal	$\frac{3}{4}$ to 1 pt
	Buctril 4EC	4 lb/gal	$\frac{3}{4}$ to $\frac{1}{2}$ pt
	Buctril ME4		
chlorsulfuron	Glean	75% DF	$\frac{1}{4}$ to $\frac{1}{4}$ wt oz
dicamba	Banvel	4 lb/gal	$\frac{1}{4}$ to $\frac{1}{4}$ pt
diuron	Direx 4L	4 lb/gal	$\frac{3}{4}$ to $1\frac{1}{4}$ pt
	Diuron 4L		
	Diuron 80 WDG	80% DF	$\frac{1}{2}$ to 1 lb
MCPA or 2,4-D ¹	Diuron DF	80% WP	
		4 lb/gal	$\frac{1}{2}$ to 1 pt
	Lexone DF	75% DG	$2\frac{1}{2}$ to 4 wt oz
metribuzin	Sencor DF		
metsulfuron	Ally	60% DF	$\frac{1}{10}$ wt oz
methyl	Igran 80WP	80% WP	7.5 to 12.5 wt oz
terbutryn	Harmony	75% DF	$\frac{1}{2}$ to $\frac{1}{2}$ wt oz
thiameturon	Express	75% DF	$\frac{1}{4}$ to $\frac{1}{4}$ wt oz

¹Tank mix with 2,4-D for application on wheat and barley only.

Rangeland and Permanent Grass Pastures

Apply $\frac{3}{4}$ to $1\frac{1}{2}$ pints of STINGER Herbicide per acre when weeds are young and actively growing. Grasses are tolerant, but new grass seedlings may be injured to varying degrees until the grass has become well established.

NOTE: Some forbs are susceptible to STINGER Herbicide. Do not spray pastures containing desirable forbs, especially legumes, unless injury can be tolerated. However, the stand and growth of established perennial grasses is usually improved after spraying, especially when rainfall is adequate and grazing is deferred.

Do not use hay or straw from treated areas for composting or mulching on susceptible broadleaf crops.

Conservation Reserve Program (CRP) For Seeding To Permanent Grasses Only

Do not use STINGER Herbicide if legumes or bentgrass are a desired cover during CRP.

Grass grown on CRP acres cannot be grazed or used for hay.

Conditions that stress grasses, such as drought, will increase potential for injury to the grass at all stages of growth. Do not use in newly seeded areas until grass is established.

After CRP, do not plant broadleaf crops in treated areas until an adequately sensitive bioassay shows that no detectable clopyralid is present in the soil.

Broadcast Applications (Ground): Applications of STINGER Herbicide should be made when perennial grasses have become established (has tillered, developed a good secondary root system and shows good vigor) since most perennial grasses have shown better tolerance to the herbicide at that stage.

For control of actively growing weeds such as musk thistle, Canada thistle, and knapweed (spotted, diffuse and Russian), use $\frac{3}{4}$ to $1\frac{1}{2}$ pints per acre of STINGER Herbicide after the majority of basal leaves have emerged, but before bud stage. For the control of wild buckwheat, volunteer sunflower and musk thistle rosettes, apply $\frac{3}{4}$ pint per acre of STINGER Herbicide. STINGER Herbicide can also be tank mixed with $\frac{1}{4}$ to 1 lb/acre of 2,4-D where species present are sensitive to 2,4-D. For best results, use in 10 or more gallons of water per acre by ground. Increasing the rate of application can increase the risk of injury. Application prior to the flowering stage is recommended (still in the bud stage).

USE PRECAUTIONS

Apply only once per 12 month period, except for sugar beets, field corn and grasses grown for seed. A fallow treatment that precedes or follows a small grain application is also allowed. (See Directions for Use.)

Wheat, barley, oats, grasses, field corn or sugar beets (including beets grown for seed) may be planted at anytime following treatment.

Alfalfa, asparagus, canola, cole crops, mint, onions, safflower, and strawberries may be planted 12 months after treatment. Dry beans, soybeans, and sunflowers may also be planted at 12 months after treatment, except in soils with less than 2% organic matter receiving less than 15 inches of natural precipitation in the 12 month period following treatment. For these areas see SPECIAL CONDITIONS below.

To avoid potential phytotoxicity, all other crops, including peas, lentils, potatoes, and broadleaf crops grown for seed, should not be planted for 18 months after treatment unless the risk of injury is acceptable.

SPECIAL CONDITIONS: Residues of STINGER in plant tissues which have not completely decayed may affect succeeding susceptible crops. In areas defined previously as low in organic matter and precipitation, sensitive crops such as dry beans, soybeans, and sunflowers may be injured when planted 12 months after treatment. Unless the risk of injury is acceptable, these crops should not be planted until 18 months after treatment. The potential for injury may be reduced by burning, removal, or incorporation of treated crop residues with a minimum of 2 supplemental Fall irrigations.

This product can affect susceptible broadleaf plants directly through foliage and indirectly by root uptake from treated soil. Therefore, do not apply STINGER Herbicide directly to or allow spray drift to come in contact with vegetables, flowers, grapes, tomatoes, potatoes, beans, lentils, peas, alfalfa, sunflowers, soybeans, safflower, or other desirable broadleaf crops and ornamental plants or soil where these sensitive crops will be planted the same season.

Do not contaminate irrigation ditches or water used for irrigation or domestic purposes.

Avoid Spray Drift: Applications should be made to avoid spray drift since very small quantities of the spray, which may not be visible, may severely injure susceptible crops during both growing and dormant periods. Use coarse sprays to minimize drift since, under adverse weather conditions, fine spray droplets may drift a mile or more. A drift control or deposition agent such as Nalco-Trol may be used with this product to aid in reducing spray drift. If used, follow all use recommendations and precautions on the product label.

Ground Application: To minimize spray drift, apply STINGER Herbicide in a total spray volume of 10 or more gallons per acre as large-droplet, low-pressure spray. Refer to manufacturer's recommendations for additional information on gallons per acre, spray pressure, sprayer speed, nozzle types and arrangements, nozzle heights above the target canopy, etc., for respective application equipment. Spot treatments should only be applied with a calibrated boom to prevent misapplication. With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible; by applying no more than 20 gallons of spray per acre; by using no more than 30 pounds spraying pressure with large droplet-producing nozzle tips; by spraying when wind velocity is low; and by stopping all spraying when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone-type insecticide or other nozzles that produce a fine-droplet spray.

Do not apply by aircraft.

Do Not Transfer Livestock from treated grazing areas onto sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated pasture. Otherwise, urine may contain enough clopyralid to cause injury to sensitive broadleaf plants.

Do not move treated soil and avoid situations where treated soil particles may blow into area where susceptible crops are grown. Violent windstorms may move soil particles. If this product is on soil particles and they are blown onto susceptible plants, visible symptoms may appear. Serious injury is unlikely. The hazard of movement of this product on dust is reduced if treated fields are irrigated or if rain occurs shortly after application.

Straw from treated areas, or manure from animals that have grazed treated areas, cannot be used for composting or mulching on ground where susceptible crops may be grown the following season. To promote herbicide decomposition, plant material should be evenly incorporated or burned. Adequate moisture is also required to promote breakdown of plant residues which contain clopyralid.

Do not use in a greenhouse. Excessive amounts of this herbicide in the soil may temporarily inhibit seed germination or plant growth.

Spray Equipment - Cleaning Instructions

To avoid injury to desirable plants, equipment used to apply STINGER Herbicide should be thoroughly cleaned before reusing to apply any other chemicals.

1. Rinse and flush application equipment thoroughly after use at least three times with water, and dispose of rinse water in non-cropland area away from water supplies.
2. During the second rinse, add 1 qt of household ammonia for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15-20 min.). Let the solution stand for several hours, preferably overnight.
3. Flush the solution out the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Nozzles and screens should be removed and cleaned separately.

PRECAUTIONARY STATEMENTS

Environmental Hazards

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store above 28°F or warm to 40°F and agitate before use.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal (Metal): Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Container Disposal (Plastic): Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY LIMITATIONS AND DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions therein under normal conditions of use. **THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL.** Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.

When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify in writing DowElanco of any claims to be eligible to receive either remedy given below. The **EXCLUSIVE REMEDY OF THE BUYER OR USER** and the **LIMIT OF LIABILITY** of DowElanco or any other seller will be one of the following, at the election of DowElanco:

- (1) **Refund of purchase price** paid by buyer or user for product bought, or
- (2) **Replacement** of amount of product used.

The seller will not be liable for consequential or incidental damages or losses.

The terms of this Warranty Limitations And Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations And Disclaimer in any manner.

08505-L6

A390

DowElanco

Indianapolis, Indiana 46268 U.S.A.

*Trademark of DowElanco

This sample label is current as of January 1, 1993. The product descriptions and recommendations provided in this sample label are for background information only. Always refer to the label on the product container before using Monsanto or any other agrichemical product.

Wilbur-Ellis

Ideas to grow with®

HERBICIDE

Complete Directions for Use

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

®Ruler is a registered trademark used under license.

®Trademark of Wilbur-Ellis

EPA Reg. No. 524-326-2935

1992-1

GPL-1100.04/CG

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION IS PROHIBITED. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED ON CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF

THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

PRECAUTIONARY STATEMENTS

Hazards to
Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

MAY CAUSE EYE IRRITATION.

Avoid contact with eyes, skin or clothing.

Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Get medical attention.

IF ON SKIN, immediately flush with plenty of water. Wash clothing before reuse.

In case of a human health emergency involving this product. Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT USING GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl) glycine,
in the form of its isopropylamine salt . . . 41.5%
INERT INGREDIENTS: 58.5%
100.0%

*Contains 480 grams per litre or 4 pounds per U.S. gallon of the active ingredient glyphosate in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

STORE ABOVE 10° F (-12° C) TO KEEP PRODUCT FROM CRYSTALLIZING.

Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68° F (20° C) for several days to redissolve and shake or roll to mix well before using.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all label safeguards until container is destroyed.

Do not reuse container. Triple rinse container. Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

This product, a water soluble liquid, mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of aboveground growth and deterioration of underground plant parts.

Unless otherwise specified on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "Weeds Controlled" section of this label. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per acre within the recommended range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (noncultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of Ruler with herbicides or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation, since minute quantities of this product can cause severe damage or destruction to the crop, plants, or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. DO NOT APPLY WHEN WIND OR OTHER CONDITIONS FAVOR DRIFT. HAND GUN APPLICATIONS SHOULD BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS. **NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.**

MIXING

This product requires use with a nonionic surfactant. When using this product, mix 4 to 8 quarts (1 to 2 percent solution) of a nonionic surfactant per 100 gallons of spray solution. Use the 8-quart rate for any surfactant containing less than 50 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

Wilbur-Ellis recommends the use of at least an 80% active ingredient surfactant such as R-11® or similar type agricultural surfactant.

• R-11 is a registered trademark of Wilbur-Ellis Company.

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the recommended amount of this product (see "Directions for Use" and "Weeds Controlled" sections of this label) and nonionic surfactant near the end of the filling process and mix well. Remove hose from tank immediately after filling to avoid siphoning back into the carrier source. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

TANK MIXTURES

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Mix labeled tank mixtures of this product with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it SLOWLY through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
6. Continue filling the sprayer tank with water and add the required amount of this product near the end of the filling process.
7. Add nonionic surfactant to the spray tank before completing the filling process.
8. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water soluble liquid followed by surfactant.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles.

CLEAN SPRAYER AND PARTS IMMEDIATELY AFTER USING THIS PRODUCT BY THOROUGHLY FLUSHING WITH WATER AND DISPOSE OF RINSATE ACCORDING TO LABELED USE OR DISPOSAL INSTRUCTIONS.

ADDITIVES

AMMONIUM SULFATE

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product and this product plus 2,4-D, Banvel™ or residual herbicide tank mixtures on annual and perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Low-quality ammonium sulfate may contain material that will not readily dissolve which could result in nozzle tip plugging. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet lines. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides or surfactant. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: The use of ammonium sulfate as an additive does not preclude the need for surfactant. Do not use herbicide rates lower than recommended in this label.

™ Banvel is a trademark of Sandoz, Inc.

COLORANTS OR DYES

Agriculturally approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to manufacturer's recommendations.

APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

Aerial — Fixed Wing and Helicopter

Broadcast Spray

Controlled Droplet Application (CDA) — Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

Hand-held and High-Volume Spray Equipment — Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, lances and other hand-held spray equipment used to direct the spray onto weed foliage and vehicle-mounted, high-volume spray equipment for spray-to-wet applications.

Selective Equipment — Recirculating sprayers, shielded sprayers and wiper applicators.

See the appropriate part of this section for specific rates of application and instructions.

AERIAL EQUIPMENT

Use the recommended rates of this product in 3 to 15 gallons of water per acre unless otherwise specified on this label. See the "Weeds Controlled" section of this label for specific rates. Unless otherwise specified, do

not exceed one quart per acre. (Aerial applications of this product may be made in annual cropping conventional tillage systems, fallow and reduced tillage systems, and preharvest. Refer to the individual use area sections of this label for recommended volumes and application rates.)

Avoid direct application to any body of water.

AVOID DRIFT — DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application — To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

BROADCAST EQUIPMENT

For control of annual or perennial weeds listed on this label using broadcast equipment — Use the recommended rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label. See the "Weeds Controlled" section of this label for specific rates. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

CONTROLLED DROPLET APPLICATION (CDA)

The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre.

For the control of labeled annual weeds with hand-held CDA units, apply a 20 percent solution of this product plus a 5 percent solution of nonionic surfactant at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 MPH (1 quart per acre). For the control of labeled perennial weeds, apply a 20 to 40 percent solution of this product plus a 5 to 10 percent solution of nonionic surfactant at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 MPH (2 to 4 quarts per acre).

Controlled droplet application equipment produces a spray pattern which is not easily visible. Extreme care

must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation as damage or destruction may result.

HAND-HELD and HIGH-VOLUME EQUIPMENT

Use coarse sprays only.

Mix this product in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff.

For control of annual weeds listed on this label, apply 0.5 percent solution of this product plus nonionic surfactant to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grasses or bud formation in broadleaf weeds. Allow three or more days before tillage or mowing.

For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 percent solution. For best results, use a 2 percent solution on harder-to-control perennials, such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

Less than complete coverage of weeds may result from the use of spray equipment designed for motorized spot treatments. Where less than complete coverage of annual weeds occurs, use a 5 percent solution. Do not reduce recommended concentrations of this product when adding surfactant.

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

Spray Solution						
AMOUNT OF RULER®						
DESIRED VOLUME	1/2%	1%	1-1/2%	2%	5%	
1 gallon	2/3 oz	1-1/3 oz	2 oz	2-2/3 oz	6-1/2 oz	
25 gallons	1 pt	1 qt	1-1/2 qt	2 qt	5 qt	
100 gallons	2 qt	1 gal	1-1/2 gal	2 gal	5 gal	
2 tablespoons = 1 fluid ounce						
Add surfactant at a rate of 1 to 2 oz. per gallon of spray mix.						

For use in knapsack sprayers, it is suggested that the recommended amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution and add the correct amount of surfactant.

SELECTIVE EQUIPMENT

This product may be applied through a recirculating spray system, a shielded applicator, or a wiper applicator after dilution and thorough mixing with water to listed weeds growing in any noncrop site specified on this label and only when specifically recommended in cropping systems.

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

A shielded applicator directs the herbicide solution onto weeds while shielding desirable vegetation from the herbicide.

A wiper applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution.

AVOID CONTACT WITH DESIRABLE VEGETATION.

Contact of the herbicide solution with the desirable vege-

tation may result in damage or destruction. Applicators used above desired vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least two inches above the desirable vegetation. Droplets, mist, foam, or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting, or destruction.

Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

See the "Weeds Controlled" section of this label for recommended stage of growth for specific weeds.

NOTE

- Maintain equipment in good operating condition. Avoid leakage or dripping onto desirable vegetation.
- Adjust height of applicator to ensure adequate contact with weeds.
- Keep nozzle tips and wiping surfaces clean.
- Keep spray patterns aligned into recovery chamber of the recirculating sprayer.
- Keep shields on shielded applicators adjusted to protect desirable vegetation.
- Maintain recommended roller RPM on roller applicators while in use.
- Keep wiper material at proper degree of saturation with herbicide solution.
- DO NOT use wiper equipment when weeds are wet.
- DO NOT operate equipment at ground speeds greater than 5 mph. Weed control may be affected by speed of application equipment. As weed density increases, reduce equipment ground speed to ensure good coverage of weeds.
- Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying on the upper end of a wiper applicator.
- Variation in equipment design may affect weed control. With wiper applicators, the wiping material and its orientation must allow delivery of sufficient quantities of the recommended herbicide solution directly to the weed.
- Care must be taken with all types of wipers to ensure that the absorbent material does not become oversaturated, causing the herbicide to drip on desirable vegetation.
- Mix only the amount of solution to be used during a one-day period, as reduced activity may result from use of leftover solutions. With all equipment, drain and clean sprayer and wiper parts immediately after using this product by thoroughly flushing with water.

RECIRCULATING SPRAYERS

Recirculating sprayer calibration is made on the basis of ground speed and delivery volume. Two procedures can be used to calibrate: (1) determining the discharge being delivered per minute, then operate at the designated ground speed, or (2) select the desired ground speed and then adjust the sprayer to deliver the recommended volume per minute (this may require nozzle changes). Use the appropriate time table below.

Do not operate at nozzle pressure above 20 PSI.

Table 1. Use this table when calibrating box or row-type recirculating sprayers. Box or row-type sprayer calibration is based on the total discharge collected per row.

Use only straight stream or 15° fan-type nozzles.

***VOLUME PER MINUTE PER ROW**

MPH	Fluid Ounces
2	26 to 35
3	38 to 51
4	51 to 68
5	65 to 86

*NOTE: Be certain the amount collected is for all spray streams treating one row.

Table 2. Use this table when calibrating broadcast type recirculating sprayers. Broadcast recirculating sprayer calibration is based on the discharge collected per minute from one nozzle on a 20-inch spacing.

VOLUME PER MINUTE PER NOZZLE

MPH	Fluid Ounces
2	7 to 9
3	10 to 13
4	13 to 18
5	16 to 22

When applied as recommended under the conditions described for recirculating sprayers, this product will control the following weeds growing a minimum of 6 inches above desirable vegetation.

Perennial Broadleaf Weeds — To SUPPRESS the following weeds, mix in a ratio of 4 quarts of this product in 20 gallons of water and apply as directed.

Dogbane, hemp	Milkweed
<i>Apocynum cannabinum</i>	<i>Asclepias syriaca</i>

Perennial Grasses and Annual Broadleaf Weeds — To control the following weeds, mix in a ratio of 3 quarts of this product in 20 gallons of water and apply as directed:

Cocklebur	Pigweed, redroot
<i>Xanthium strumarium</i>	<i>Amaranthus retroflexus</i>
Johnsongrass	Sunflower
<i>Sorghum halepense</i>	<i>Helianthus annuus</i>

Annual Grasses — To CONTROL the following weeds, mix in a ratio of 2 quarts of this product in 20 gallons of water and apply as directed:

Corn	Shattercane
<i>Zea mays</i>	<i>Sorghum bicolor</i>

SHIELDED APPLICATORS

When applied as directed under conditions described for shielded applicators, this product will control those weeds listed in the "Weeds Controlled" section of this label.

Shielded applicators which apply the herbicide solution as a spray band should be calibrated on a broadcast equivalent rate and volume basis. To determine these:

Band width in inches	x	Herbicide broadcast RATE per acre	=	Herbicide band RATE per acre
Row width in inches				

Band width in inches	x	Broadcast VOLUME of solution per acre	=	Band VOLUME of solution per acre
Row width in inches				

Use nozzles that provide uniform coverage within the treated area. EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT WITH DESIRABLE VEGETATION.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

WIPER APPLICATORS

Wiper applicators include either roller or wick devices which physically wipe appropriate concentrations or amounts of this product directly onto the weed. Equipment must be designed, maintained, and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if two applications are made in opposite directions.

For Roller Applicators — Mix 1 gallon of this product in enough water to prepare 10 gallons of herbicide solution (10 percent solution). Apply this solution to perennial weeds or annual broadleaf weeds listed in this "Wiper Applicators" section.

Mix 1 gallon of this product in enough water to provide 20 gallons of herbicide solution (5 percent solution). Apply this solution to annual grasses listed in this "Wiper Applicators" section.

Roller speed should be maintained at 40 to 60 RPM.

For Wick or Wiper Applicators — Mix 1 gallon of this product in 2 gallons of water to prepare a 33 percent solution. Apply this solution to weeds listed in this "Wiper Applicators" section.

In severe infestations, reduce equipment ground speed to ensure that adequate amounts of this product are wiped on the weeds. A second treatment in the opposite direction may be beneficial.

Do not permit herbicide solution to contact desirable vegetation.

When applied as recommended under the conditions described for "Wiper Applicators", this product CONTROLS the following weeds:

ANNUAL GRASSES

Corn	Rye, common
<i>Zea mays</i>	<i>Secale cereale</i>
Panicum, Texas	Shattercane
<i>Panicum texanum</i>	<i>Sorghum bicolor</i>

ANNUAL BROADLEAVES

Sicklepod	Starbur, bristly
<i>Cassia obtusifolia</i>	<i>Acanthospermum hispidum</i>
Spanishneedles	
<i>Bidens bipinnata</i>	

When applied as recommended under the conditions described for "Wiper Applicators", this product SUPPRESSES the following weeds:

ANNUAL BROADLEAVES

Beggarweed, Florida	Ragweed, giant
<i>Desmodium tortuosum</i>	<i>Ambrosia trifida</i>
Dogtunnel	Sunflower
<i>Eupatorium capilliflorum</i>	<i>Helianthus annuus</i>
	Thistle, musk
	<i>Carduus nutans</i>

Pigweed, redroot	Velvetleaf
<i>Amaranthus retroflexus</i>	<i>Abutilon theophrasti</i>

Ragweed, common	
<i>Ambrosia artemisiifolia</i>	

PERENNIAL GRASSES

Bermudagrass	Smutgrass
<i>Cynodon dactylon</i>	<i>Sporobolus poiretii</i>
Guineagrass	Vaseygrass
<i>Panicum maximum</i>	<i>Paspalum urvillei</i>
Johnsongrass	
<i>Sorghum halepense</i>	

PERENNIAL BROADLEAVES

Dogbane, hemp	Nightshade, silverleaf
<i>Apocynum cannabinum</i>	<i>Solanum elaeagnifolium</i>
Milkweed	Thistle, Canada
<i>Asclepias syriaca</i>	<i>Cirsium arvense</i>

WEEDS CONTROLLED

This herbicide controls many annual and perennial grasses and broadleaf weeds.

ANNUAL WEEDS

- Apply to actively growing grass and broadleaf weeds.
- Allow at least 3 days after treatment before tillage.
- For maximum agronomic benefit, apply when weeds are 6 inches or less in height.
- To prevent seed production, applications should be made prior to seedhead formation.
- This product does not provide residual control; therefore, delay application until maximum weed emergence. Repeat treatments may be necessary to control later germinating weeds.

**LOW-VOLUME BROADCAST APPLICATION
(LOW-RATE TECHNOLOGY)**

When applied as directed under the conditions described, this product will control the weeds listed below when water carrier volumes of 3 to 10 gallons per acre for ground applications and 3 to 5 gallons per acre for aerial applications are recommended. (See the "Aerial Equipment" section of this label for approved sites.)

NOTE

- The addition of 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water may increase the performance of this product on annual weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the "Mixing, Additives and Application Instructions" section of this label.
- Do not tank-mix with soil-residual herbicides when using these rates unless otherwise specified.
- For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.
- Refer to the "Tank Mixtures" portion of this section for control of additional broadleaf weeds.

WEED SPECIES	MAXIMUM HEIGHT/ LENGTH	RATE PER ACRE ²⁰ (FLUID OUNCES)
Foxtail		
<i>Setaria spp.</i>	12"	8 oz

WEED SPECIES	MAXIMUM HEIGHT/LENGTH	RATE PER ACRE ^{1,2} (FLUID OUNCES)
Barnyardgrass <i>Echinochloa crus-galli</i>	6" (0 to 4" 4 to 6")	12 oz. 16 oz. ¹ 24 oz. ¹
Bluegrass, annual <i>Poa annua</i>		
Brome, downy ^o <i>Bromus tectorum</i>		
Mustard, blue <i>Chorispora tenella</i>		
Mustard, tansy <i>Descurainia pinnata</i>		
Mustard, tumble <i>Sisymbrium altissimum</i>		
Mustard, wild <i>Sinapis arvensis</i>		
Spurry, umbrella <i>Holosteum umbellatum</i>		
Barley <i>Hordeum vulgare</i>	12"	
Rye <i>Secale cereale</i>		
Sandbur, field <i>Cenchrus spp.</i>		
Shattercane <i>Sorghum bicolor</i>		
Stinkgrass <i>Eragrostis cilianensis</i>		
Wheat <i>Triticum aestivum</i>	18"	
Morningglory <i>Ipomoea spp.</i>	2"	16 oz.
Sicklepod <i>Cassia obtusifolia</i>		
Bluegrass, bulbous <i>Poa bulbosa</i>	6"	
Cheat <i>Bromus secalinus</i>		
Chickweed, common <i>Stellaria media</i>		
Chickweed, mouseear <i>Cerastium vulgatum</i>		
Corn <i>Zea mays</i>		
Goatgrass, jointed <i>Aegilops cylindrica</i>		
Groundsel, common <i>Senecio vulgaris</i>		
Horseweed/Marestail <i>Conyza canadensis</i>		
Lambsquarters, common <i>Chenopodium album</i>		
Pennycress, field Fanweed <i>Thlaspi arvense</i>		
Rocket, London <i>Sisymbrium irio</i>		

WEED SPECIES	MAXIMUM HEIGHT/LENGTH	RATE PER ACRE ^{1,2} (FLUID OUNCES)
Ryegrass, Italian <i>Lolium multiflorum</i>	6"	16 oz.
Shepherdspurse <i>Capsella bursa-pastoris</i>		
Spurge, annual <i>Euphorbia spp.</i>		
Buttercup <i>Ranunculus spp.</i>	12"	
Cocklebur <i>Xanthium strumarium</i>		
Crabgrass <i>Digitaria spp.</i>		
Dwarf dandelion <i>Krigia cespitosa</i>		
Falseflax, smallseed <i>Camelina microcarpa</i>		
Foxtail, Carolina <i>Alopecurus carolinianus</i>		
Johnsongrass, seedling <i>Sorghum halepense</i>		
Oats, wild <i>Avena fatua</i>		
Panicum, fall <i>Panicum dichotomiflorum</i>		
Panicum, Texas <i>Panicum texanum</i>		
Pigweed, redroot <i>Amaranthus retroflexus</i>		
Pigweed, smooth <i>Amaranthus hybridus</i>		
Witchgrass <i>Panicum capillare</i>		
Sicklepod <i>Cassia obtusifolia</i>	3 to 4"	24 oz.
Signalgrass, broadleaf <i>Bracharia platyphylla</i>	4"	
Horseweed/Marestail <i>Conyza canadensis</i>	7 to 12"	
Lambsquarters, common <i>Chenopodium album</i>		
Spurge, annual <i>Euphorbia spp.</i>		
Rice, red <i>Oryza sativa</i>	4"	32 oz.
Teaweed <i>Sida spinosa</i>		
Sprangletop <i>Leptochloa spp.</i>	6"	
Geranium, Carolina <i>Geranium carolinianum</i>	12"	
Goosegrass <i>Eleusine indica</i>		

WEED SPECIES	MAXIMUM HEIGHT/LENGTH	RATE PER ACRE ^{1,2} (FLUID OUNCES)
Primrose, cutleaf evening <i>Oenothera laciniate</i>	12"	32 oz.
Pusley, Florida <i>Richardia scabra</i>		
Sicklepod <i>Cassia obtusifolia</i>	5 to 12"	
Spanishneedles <i>Bidens bipinnata</i>		
Filaree <i>Erodium spp.</i>	12"	48 oz.
Sprangletop <i>Leptochloa spp.</i>		

¹Use these rates to control barnyardgrass in Alabama, Arkansas, Mississippi, Louisiana, Missouri and Texas for preplant treatments.

²For control in no-till systems, use 16 fluid ounces per acre.

³For those rates less than 32 fluid ounces per acre, this product at rates up to 32 fluid ounces per acre may be used where heavy weed densities exist.

■ TANK MIXTURES ■
RULER® plus BANVEL
 ■ plus NONIONIC SURFACTANT ■
RULER plus 2,4-D
 ■ plus NONIONIC SURFACTANT ■

These tank mixtures are recommended for use in fallow and reduced tillage areas only. Follow use directions as given in this "Low-Volume Broadcast Application" section.

This product plus Banvel or 2,4-D will control the annual grasses and broadleaf weeds previously listed for this product alone at the indicated heights (except 8 fluid ounces per acre applications), plus the following broadleaf weeds. For those weeds previously listed at 8 fluid ounces of this product alone per acre, use 12 fluid ounces in these tank mixtures.

NOTE: Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if Banvel is applied within 45 days of planting. The addition of Banvel in a mixture with this product may provide short-term residual control of selected weed species.

Apply 12 to 16 fluid ounces of this product plus 0.25 lb. a.i. of Banvel or 0.5 pound a.i. of 2,4-D per acre to control dense populations of the following annual broadleaf weeds when less than the height indicated:

Cocklebur (12") <i>Xanthium strumarium</i>	Morningglory (6") <i>Ipomoea spp.</i>
Kochia* (6") <i>Kochia scoparia</i>	Pigweed, redroot (12") <i>Amaranthus retroflexus</i>
Lambsquarters (12") <i>Chenopodium album</i>	Pigweed, smooth (12") <i>Amaranthus hybridus</i>
Lettuce, prickly (6") <i>Lactuca serriola</i>	Thistle, Russian (12") <i>Salsola kali</i>
Marestail/Horseweed (6") <i>Conyza canadensis</i>	

*Controlled with Banvel tank mixture only.

Apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D per acre to control the following annual broad-leaf weeds when less than 6 inches in height.

Ragweed, common <i>Ambrosia artemisiifolia</i>	Smartweed, Pennsylvania <i>Polygonum pennsylvanicum</i>
Ragweed, giant <i>Ambrosia trifida</i>	Velvetleaf <i>Abutilon theophrasti</i>

HIGH-VOLUME BROADCAST APPLICATIONS

When applied as directed under the conditions described, this product will control the weeds listed below when water carrier volumes are 10 to 40 gallons per acre for ground applications.

Use 1 quart of this product per acre if weeds are less than 6 inches tall. If weeds are over 6 inches tall, use 1.5 quarts of this product per acre. If weeds have been mowed, grazed, or cut, allow adequate time for new growth to recommended stages prior to treatment. These rates will also provide control of weeds listed in the "Low-Volume Broadcast Application" section.

WEED SPECIES

Balsamapple* <i>Momordica charantia</i>	Panicum <i>Panicum spp.</i>
Bassia, fivehook <i>Bassia hyssopifolia</i>	Ragweed, common <i>Ambrosia artemisiifolia</i>
Brome <i>Bromus spp.</i>	Ragweed, giant <i>Ambrosia trifida</i>
Fiddleneck <i>Amsinckia spp.</i>	Smartweed, Pennsylvania <i>Polygonum pennsylvanicum</i>
Flaxleaf, hairy <i>Conyza bonariensis</i>	Sowthistle, annual <i>Sonchus oleraceus</i>
Fleabane <i>Erigeron spp.</i>	
Kochia <i>Kochia scoparia</i>	Sunflower <i>Helianthus annuus</i>
Lettuce, prickly <i>Lactuca serriola</i>	Thistle, Russian <i>Salsola kali</i>
	Velvetleaf <i>Abutilon theophrasti</i>

*Apply with hand-held equipment only.

PERENNIAL WEEDS

Apply this product as follows to control or destroy most perennial weeds:

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages of growth.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product on perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the "Mixing, Additives and Application Instructions" section of this label.

When applied as recommended under the conditions described, this product **WILL CONTROL** the following **PERENNIAL WEEDS**:

Alfalfa <i>Medicago sativa</i>	Johnsongrass <i>Sorghum halepense</i>
Alligatorweed* <i>Alternanthera philoxeroides</i>	Kikuyugrass <i>Pennisetum clandestinum</i>

Artichoke, Jerusalem <i>Helianthus tuberosus</i>	Knapweed <i>Centaurea repens</i>
Bahiagrass <i>Paspalum notatum</i>	Lantana <i>Lantana camara</i>
Bentgrass <i>Agrostis spp.</i>	Milkweed <i>Asclepias spp.</i>
Bermudagrass <i>Cynodon dactylon</i>	Muhly, wirestem <i>Muhlenbergia frondosa</i>
Bermudagrass, water (knotgrass) <i>Paspalum distichum</i>	Mullein, common <i>Verbascum thapsus</i>
Bindweed, field <i>Convolvulus arvensis</i>	Napiergrass <i>Pennisetum purpureum</i>
Bluegrass, Kentucky <i>Poa spp.</i>	Nightshade, silverleaf <i>Solanum elaeagnifolium</i>
Blueweed, Texas <i>Helianthus ciliaris</i>	Nutsedge: purple, yellow <i>Cyperus rotundus</i> <i>Cyperus esculentus</i>
Brackenfern <i>Pteridium aquilinum</i>	Orchardgrass <i>Dactylis glomerata</i>
Bromegrass, smooth <i>Bromus inermis</i>	Pampas <i>Cortaderia jubata</i>
Bursage, woollyleaf <i>Franseria tomentosa</i>	Paragrass <i>Brachiaria mutica</i>
Canarygrass, reed <i>Phalaris arundinacea</i>	Phragmites* <i>Phragmites spp.</i>
Cattail <i>Typha spp.</i>	Quackgrass <i>Agropyron repens</i>
Clover, red <i>Trifolium pratense</i>	Redvine* <i>Brunnichia ovata</i>
Clover, white <i>Trifolium repens</i>	Reed, giant <i>Arundo donax</i>
Cogongrass <i>Imperata cylindrica</i>	Ryegrass, perennial <i>Lolium perenne</i>
Dallisgrass <i>Paspalum dilatatum</i>	Smartweed, swamp <i>Polygonum coccineum</i>
Dandelion <i>Taraxacum officinale</i>	Spurge, leafy* <i>Euphorbia esula</i>
Dock, curly <i>Rumex crispus</i>	Sweet potato, wild* <i>Ipomoea pandurata</i>
Dogbane, hemp <i>Apocynum cannabinum</i>	Thistle, Canada <i>Cirsium arvense</i>
Fescues <i>Festuca spp.</i>	Timothy <i>Phleum pratense</i>
Fescue, tall <i>Festuca arundinacea</i>	Torpedograss* <i>Panicum repens</i>
Guineagrass <i>Panicum maximum</i>	Trumpetcreeper* <i>Campsis radicans</i>
Horsenettle <i>Solanum carolinense</i>	Vaseygrass <i>Paspalum urvillei</i>
Horseradish <i>Armoracia rusticana</i>	Wheatgrass, western <i>Agropyron smithii</i>

*Partial control

Alfalfa — Apply 1 quart of this product in 3 to 10 gallons of water per acre. Make application after last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.

Alligatorweed — Apply 4 quarts of this product per acre or apply a 1-1/2 percent solution with hand-held equipment to provide partial control. Apply when most of the

plants are in bloom. Repeat applications will be required to maintain such control.

Bentgrass — For suppression in grass seed production areas. For ground applications only, apply 1.5 quarts of this product in 10 to 20 gallons of water per acre. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is recommended for best results. Failure to use tillage after treatment may result in unacceptable control.

Bermudagrass — For control, apply 5 quarts of this product per acre and, for partial control, apply 3 quarts per acre. Treat when bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control. Allow 7 or more days after application before tillage.

Bermudagrass, water (knotgrass) — Apply 1.5 quarts of this product in 5 to 10 gallons of water per acre. Apply when water bermudagrass is actively growing and 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field.

Fall applications only — Apply 1 quart of this product in 5 to 10 gallons of water per acre. Fallow fields should be tilled prior to application. Apply prior to frost on water bermudagrass that is actively growing and 12 to 18 inches in length. Allow 7 or more days before tillage.

Bindweed, field — For control, apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts east of the Mississippi River. Apply when the weeds are actively growing and are at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.

Also for control, apply 2 quarts of this product plus 0.5 pound a.i. of Banvel in 10 to 20 gallons of water per acre. At these rates, apply using ground application only.

The following tank mixtures with 2,4-D may be applied using aerial application equipment in fallow and reduced tillage systems only.

For suppression on irrigated agricultural land apply 1 to 2 quarts of this product plus 1 pound a.i. of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications should be made following harvest or on fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth.

For suppression, apply 16 fluid ounces of this product plus 0.5 pounds a.i. of 2,4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

Bluegrass, Kentucky / Bromegrass, smooth / Orchardgrass — Apply 2 quarts of this product in 10 to 40 gallons of water per acre when the grasses are actively growing and most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1-1/2 quarts of this product in 5 to 10 gallons of water per acre.

Apply to actively growing plants when most have reached 4 to 12 inches in height. Allow 7 or more days after application before tillage.

Blueweed, Texas—Apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when weed is actively growing and is at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.

Brackenfern—Apply 3 to 4 quarts of this product per acre as a broadcast spray or as a 1 to 1-1/2 percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 18 inches long.

Bursage, woollyleaf—For control, apply 2 quarts of this product plus 1 pint of Banvel per acre. For partial control, apply 1 quart of this product plus 1 pint of Banvel per acre. Apply 3 to 20 gallons of water per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.

Canarygrass, reed / Timothy / Wheatgrass, western—Apply 2 to 3 quarts of this product per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage.

Cogongrass—Apply 3 to 5 quarts of this product in 10 to 40 gallons of water per acre. Apply when cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

Dandelion/Dock, curly—Apply 3 to 5 quarts of this product per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.

Also for control, apply 16 fluid ounces of this product plus 0.5 pounds a.i. 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre.

Dogbane, hemp—Apply 4 quarts of this product per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.

For suppression, apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.

Fescue, tall—Apply 3 quarts of this product in 10 to 40 gallons of water per acre to actively growing plants when most have reached boot-to-early seedhead stage of development.

Fall applications only—Apply 1 quart of this product in 3 to 10 gallons of water per acre. Apply to fescue in the fall when actively growing and plants have 6 to 12 inches of new growth. Allow 7 or more days after application before tillage. A sequential application of 1 pint per acre of this product will improve long-term control and con-

trol seedlings germinating after fall treatments or the following spring.

Guineagrass—Apply 3 quarts of this product per acre or use a 1 percent solution with hand-held equipment. Apply to actively growing guineagrass when most have reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. Allow 7 or more days after application before tillage.

Johnsongrass / Ryegrass, perennial—Apply 1 to 3 quarts of this product per acre. In annual cropping systems, apply 1 to 2 quarts of this product per acre. Apply 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. In noncrop or areas where annual tillage (no-till) is not performed, apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage. Do not tank-mix with residual herbicides when using the 1 quart per acre rate.

For burndown of johnsongrass, apply 1 pint per acre in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.

Spot Treatment (partial control or suppression)—Apply a 1 percent solution of this product when johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.

Kikuyugrass—Apply 2 to 3 quarts of this product per acre. Spray when most kikuyugrass is at least 8 inches in height (3 or 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.

Knapweed/Horseradish—Apply 4 quarts of this product per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.

Lantana—Apply this product as a 1 to 1-1/4 percent solution using hand-held equipment only. Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth. Allow 7 or more days after application before tillage.

Milkweed, common—Apply 3 quarts of this product per acre. Apply when actively growing and most of the milkweed has reached the late bud to flower stage of growth. Following small grain harvest or mowing, allow milkweed to regrow to a mature stage prior to treatment. Allow 7 or more days after application before tillage.

Muhly, wirestem—Apply 1 to 2 quarts of this product per acre. Use 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre or in pasture, sod, or noncrop areas. Spray when the wirestem muhly is 8 inches or more in height and actively growing. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage. This product will not provide residual control of wirestem muhly from seeds which germinate after a Ruler herbicide application. Do not tank mix with residual herbicides when using the 1 quart per acre rate.

Nightshade, silverleaf—For control, apply 2 quarts of this product in 3 to 10 gallons of water per acre. Applications should be made when at least 60 percent of plants have berries. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage. Do not treat when weed is under drought stress, as good soil moisture is necessary for active growth.

Nutsedge: purple, yellow—Apply 3 quarts of this product per acre as a broadcast spray, or apply a 2 percent solution from hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control of ungerminated tubers. Sequential applications of 1 to 2 quarts of this product in 3 to 10 gallons of water per acre will provide control. Make applications when a majority of the plants are in the 3 to 5 leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3 to 5 leaf stage. Subsequent applications will be necessary for long-term control.

For suppression of existing plants, apply 1 pint to 2 quarts of this product in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants. Wait 7 days after treatment before tillage or mowing.

Pampas—Apply this product as a 1-1/2 to 2 percent solution using hand-held equipment. Apply to plants that are actively growing at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

Phragmites—For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 5 quarts per acre as a broadcast spray or apply a 2 percent solution from hand-held equipment. In other areas of the U.S., apply 3 quarts per acre as a broadcast spray or apply a 1 percent solution from hand-held equipment for partial control. For best results, treat during late summer or fall months or when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptom will be slow to develop.

Quackgrass—In Annual Cropping Systems, or in Pastures and Sods Followed by Deep Tillage: Apply 1 to 2 quarts of this product per acre. For the one quart rate, apply in 3 to 10 gallons of water per acre. For the 2 quart rate, apply in 10 to 40 gallons of water per acre. Do not tank mix with residual herbicides when using the 1 quart rate. Spray when quackgrass is 8 to 12 inches in height and actively growing. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, for best results use a mold-board plow.

Quackgrass—Pasture or Sod or Other Noncrop Areas Where Deep Tillage is Not Planned Following Application: Apply 2 to 3 quarts in 10 to 40 gallons of water per acre. Spray when the quackgrass is greater than 8 inches tall and actively growing. Do not till between harvest and fall application or in fall or spring prior to spring application. Allow 3 or more days after application before tillage.

Redvine—For suppression, apply 24 fluid ounces of this product per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply recommended rates in 5 to 10 gallons of water per acre. Apply to actively growing plants in late September or early October, which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least one week before a killing frost.

Reed, giant — For control of giant reed, apply a 2 percent solution of this product when plants are actively growing. Best results are obtained when applications are made in late summer to fall.

Smartweed, swamp—Apply 3 to 5 quarts of this product per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.

Also for control, apply 16 fluid ounces of this product plus 0.5 pound active ingredient of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.

Spurge, leafy—For suppression, apply 16 fluid ounces of this product plus 0.5 pound active ingredient 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall. Allow 7 or more days after application before tillage.

Sweet Potato, wild — Apply this product as a 2 percent solution using hand-held equipment. Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications will be required. Allow the plant to reach the recommended stage of growth before retreatment. Allow 7 or more days before tillage.

Thistle, Canada — Apply 2 to 3 quarts of this product per acre. Apply to actively growing thistles when most are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

For suppression of Canada thistle, apply 1 quart of this product, or 1 pint of this product plus 0.5 pound a.i. 2,4-D, in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.

Torpedograss — Apply 4 to 5 quarts of this product per acre to provide partial control of torpedograss. Apply to actively growing torpedograss when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost. Allow 7 or more days after application before tillage.

Trumpet creeper—For control, apply 2 quarts of this product in 5 to 10 gallons of water per acre. Apply in late September or October to actively growing plants which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Other perennials listed on this label — Apply 3 to 5 quarts

of this product per acre. Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.

WOODY BRUSH AND TREES

When applied as recommended under the conditions described, this product CONTROLS or PARTIALLY CONTROLS the following woody brush, plants and trees:

Alder	Maple:
<i>Alnus</i> spp.	Red**
Ash*	<i>Acer rubrum</i>
<i>Fraxinus</i> spp.	Sugar
Aspen, quaking	<i>Acer saccharum</i>
<i>Populus tremuloides</i>	
Bearmat (Bearclover)	Vine*
<i>Chamaebatia foliolosa</i>	<i>Acer circinatum</i>
Birch	Monkey Flower*
<i>Betula</i> spp.	<i>Mimulus guttatus</i>
Blackberry	Oak:
<i>Rubus</i> spp.	Black*
Broom:	<i>Quercus velutina</i>
French	Northern Pin
<i>Cytisus monspessulanus</i>	<i>Quercus palustris</i>
Scotch	Post
<i>Cytisus scoparius</i>	<i>Quercus stellata</i>
Buckwheat, California*	Red
<i>Eriogonum fasciculatum</i>	<i>Quercus rubra</i>
Cascara*	Southern Red
<i>Rhamnus purshiana</i>	<i>Quercus falcata</i>
Catsclaw*	White*
<i>Acacia greggii</i>	<i>Quercus alba</i>
Ceanothus*	Persimmon*
<i>Ceanothus</i> spp.	<i>Diospyros</i> spp.
Chamise	Poison Ivy
<i>Adenostoma fasciculatum</i>	<i>Rhus radicans</i>
Cherry:	Poison Oak
Bitter	<i>Rhus toxicodendron</i>
<i>Prunus emarginata</i>	Poplar*, yellow
Black	<i>Liriodendron tulipifera</i>
<i>Prunus serotina</i>	Raspberry
Pin	<i>Rubus</i> spp.
<i>Prunus pensylvanica</i>	Rose, multiflora
Coyote brush	<i>Rosa multiflora</i>
<i>Baccharis consanguinea</i>	Russian-olive
Creep, Virginia*	<i>Eleagnus angustifolia</i>
<i>Parthenocissus quinquefolia</i>	Sage, black
Dewberry	<i>Salvia mellifera</i>
<i>Rubus trivialis</i>	Sagebrush, California
Elderberry	<i>Artemisia californica</i>
<i>Sambucus</i> spp.	Salmonberry
Elm*	<i>Rubus spectabilis</i>
<i>Ulmus</i> spp.	Sassafras
Eucalyptus, bluegum	<i>Sassafras albidum</i>
<i>Eucalyptus glotulus</i>	Sourwood
Hasardia*	<i>Oxydendrum arboreum</i>
<i>Haplopappus squamosus</i>	Sumac:
Hawthorn	Poison*
<i>Crataegus</i> spp.	<i>Rhus vernix</i>
	Smooth*
	<i>Rhus glabra</i>
	Winged*
	<i>Rhus copallina</i>

Hazel	Sweetgum
<i>Corylus</i> spp.	<i>Liquidambar styraciflua</i>
Honeysuckle	Swordfern*
<i>Lonicera</i> spp.	<i>Polystichum munitum</i>
Kudzu	Tallowtree, Chinese
<i>Pueraria lobata</i>	<i>Sapium sebiferum</i>
Locust, black*	Tan oak
<i>Robinia pseudoacacia</i>	<i>Lithocarpus densiflorus</i>
Madrone	Thimbleberry
<i>Arbutus menziesii</i>	<i>Rubus parviflorus</i>
Manzanita	Tobacco, tree*
<i>Arctostaphylos</i> spp.	<i>Nicotiana glauca</i>
	Trumpet creeper
	<i>Campsis radicans</i>
	Willow
	<i>Salix</i> spp.

*Partial control

**See below for control or partial control instructions.

NOTE: If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stages of growth.

Apply this product when plants are actively growing, and unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring to early summer when brush species are at high moisture content and are flowering. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

Apply this product as follows to control or partially control the following woody brush and trees.

Alder / Dewberry / Honeysuckle / Post Oak / Raspberry — For control, apply 3 to 4 quarts per acre of this product as a broadcast spray or as a 1 to 1-1/2 percent solution with hand-held equipment.

Aspen, quaking / Cherry, bitter, black, pin / Hawthorn / Oak, southern red / Sweetgum / Trumpet creeper — For control, apply 2 to 3 quarts of this product per acre as a broadcast spray or as a 1 to 1-1/2 percent solution with hand-held equipment.

Birch / Elderberry / Hazel / Salmonberry / Thimbleberry — For control, apply 2 quarts per acre of this product as a broadcast spray or as a 1 percent solution with hand-held equipment.

Blackberry — For control, apply 3 to 4 quarts per acre of this product as a broadcast spray, or 1 to 1-1/2 percent solution with hand-held equipment. Make application after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. After berries have set or dropped in late fall, blackberry can be controlled by applying a 3/4

percent solution of this product with hand-held equipment. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of this product in 10 to 40 gallons of water per acre.

Broom: French, Scotch — For control, apply a 1-1/2 to 2 percent solution with hand-held equipment.

Buckwheat, California / Hasardia / Monkey Flower / Tobacco, tree — For partial control of these species, apply a 1 to 2 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Catsclaw — For partial control, apply as a 1 to 1-1/2 percent solution with hand-held equipment.

Coyote Brush — For control, apply a 1-1/2 to 2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Eucalyptus, bluegum — For control of eucalyptus resprouts, apply a 2 percent solution of this product with hand-held equipment when resprouts are 6 to 12 feet tall. Ensure complete coverage. Apply when plants are growing actively. Avoid application to drought-stressed plants. For control of eucalyptus trees 2 inches or less in diameter, cut trees as close to the soil surface as desired. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delay in applying this product may result in reduced performance.

Kudzu — For control, apply 4 quarts of this product per acre as a broadcast spray or as a 2 percent solution with hand-held equipment. Repeat applications will be required to maintain control.

Madrone resprouts — For suppression or partial control, apply a 2 percent solution of this product to resprouts less than 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.

Maple, red — For control, apply as a 1 to 1-1/2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of this product per acre as a broadcast spray.

Maple, sugar / Oak, northern pin / Oak, red — For control, apply as a 1 to 1-1/2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Poison ivy / Poison Oak — For control, apply 4 to 5 quarts of this product per acre as a broadcast spray or as a 2 percent solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.

Rose, multiflora — For control, apply 2 quarts of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Treatments should be made prior to leaf deterioration by leaf-feeding insects.

Sage, black / Sagebrush, California / Chamise / Tallow-tree, Chinese — For control of these species, apply a 1 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Tan oak resprouts — For suppression or partial control, apply a 2 percent solution of this product to resprouts less than 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.

Willow — For control, apply 3 quarts of this product per

acre as a broadcast spray or as a 1 percent solution with hand-held equipment.

Other Woody Brush and Trees listed on this label — For partial control, apply 2 to 4 quarts of this product per acre as a broadcast spray or as a 1 to 1-1/2 percent solution with hand-held equipment.

NONCROP USES

See "General Information" and "Mixing, Additives and Application Instructions" sections of this label for essential product performance information and the following "Noncrop" sections for specific recommended uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OF DESIRABLE TURFGRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

This product does not provide residual weed control. For subsequent weed control, follow a label-approved herbicide program.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

FARMSTEAD WEED CONTROL

When applied as directed for "Noncrop Uses", under conditions described, this product controls undesirable vegetation listed on this label around farmstead building foundations, along and in fences, shelterbelts, and for general nonselective farmstead weed control.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

GRASSES FOR SEED PRODUCTION

■ PREPLANT AND RENOVATION ■

When applied as directed for "Noncrop" uses, under conditions described, this product controls most existing vegetation prior to the planting or renovation of grass seed production areas.

For specific rates of application and instructions for control of various annual and perennial weeds, and woody brush and trees, see the "Weeds Controlled" section of this label.

For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as bermudagrass, summer or fall applications provide best control. DO NOT DISTURB SOIL OR UNDERGROUND PLANT PARTS BEFORE TREATMENT. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.

Apply this product to actively growing weeds at the stages of growth given in the "Weeds Controlled" section of this

label prior to planting or renovation of turf or forage grass areas grown for seed production.

DO NOT feed or graze treated areas within 8 weeks after application.

CROPPING SYSTEMS

When applied as directed for "Cropping Systems", under the conditions described, this product controls annual and perennial weeds listed on this label, prior to the emergence of direct seeded crops or prior to transplanting of crops listed on this label.

See the "General Information" and "Mixing, Additives and Application Instructions" sections of this label for essential product performance information.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE, GREEN STEMS OR FRUIT OF DESIRABLE CROPS, PLANTS, TREES OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Except as otherwise specified on this label, repeat treatments must be made before the crop emerges in accordance with the instructions of this label.

Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts per acre of this product per year.

Do not plant subsequent crops other than those on the label for 30 days following application.

Do not harvest or feed treated vegetation for 8 weeks following application. Following spot treatment or selective equipment use, allow 14 days before grazing domestic livestock or harvesting forage grasses and legumes.

ALFALFA*	LOGANBERRY
ARTICHOKE	LONGAN
(Jerusalem)	LYCHEE
ASPARAGUS*	MELONS***
ATEMOYA	MUSTARD GREENS
BARLEY*	OATS*
BEANS (All)	OKRA
BEET GREENS	OLALLIEBERRY
BEETS (Red, Sugar)	ONION
BLACKBERRY	PARSNIPS
BLUEBERRY	PASSION FRUIT
BOYSENBERRY	PEANUTS
BREADFRUIT	PEAS (All)
BROCCOLI	PEPPER***
CABBAGE	PERSIMMONS
CANISTEL	PINEAPPLE††
CARAMBOLA	POTATO (Irish, Sweet)
CARROT	PUMPKIN***
CAULIFLOWER	RADISH
CELERY	RASPBERRY
CHICORY	(Black, Red)
CORN (All)*	RICE**
COTTON*	RUTABAGA
CRANBERRY	SAPODILLA
CUCUMBER***	SAPOTE (Black, Mamey, White)
CURRENT	SORGHUM (Milo)*
DATES	SOURSOP
DEWBERRY	

EGGPLANT***
 ELDERBERRY
 FORAGE GRASSES*
 FORAGE LEGUMES*
 GARLIC***
 GOOSEBERRY
 GOURDS***
 HORSERADISH
 HUCKLEBERRY
 JABOTICABA
 JACKFRUIT
 KALE
 LENTILS
 LETTUCE

SOYBEANS*
 SPINACH
 SQUASH***
 (Summer, Winter)
 SUGAR APPLE
 SUGARCANE
 TAMARIND
 TOMATILLO***
 TOMATOES***†
 TURNIPS
 WATERCRESS***
 WATERMELON***
 WHEAT*
 YAMS

*Spot treatments may be applied in these crops.

**Do not treat rice fields or levees when the fields contain flood water.

***Apply only prior to planting. Allow at least 3 days between application and planting.

†Use is restricted to direct seeded crops only.

††Do not feed or graze treated pineapple forage following application.

When applying this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product from the plastic prior to transplanting. Residues can be removed by ½ inch natural rainfall or by applying water via a sprinkler irrigation system.

Spot Treatment (Only those crops with "*" can be spot treated) — Applications in growing crops must be made prior to heading of small grains and milo, initial pod set in soybeans, silking of corn, or boll opening on cotton.

For forage grasses and forage legumes, see the "Spot Treatment" in the "Pastures" section of "Cropping Systems" in this label.

For dilution and rates of application using boom or hand-held equipment, see the "Mixing, Additives and Application Instructions" and "Weeds Controlled" sections of this label.

NOTE: FOR FORAGE GRASSES AND FORAGE LEGUMES, NO MORE THAN ONE-TENTH OF ANY ACRE SHOULD BE TREATED AT ONE TIME. FOR ALL OTHER CROPS, DO NOT TREAT MORE THAN 10 PERCENT OF THE TOTAL FIELD AREA TO BE HARVESTED.

THE CROP RECEIVING SPRAY IN TREATED AREA WILL BE KILLED. TAKE CARE TO AVOID DRIFT OR SPRAY OUTSIDE TARGET AREA FOR THE SAME REASON.

Selective Equipment — This product may be applied through recirculating sprayers, shielded applicators, or wiper applicators in cotton and soybeans. Shielded and wiper applicators may also be used in tree crops and grapes. Wiper applicators may be used in rutabagas, forage grasses and forage legumes, including pasture sites and grain sorghum (milo).

See the "Selective Equipment" part of the "Application Equipment and Techniques" section of this label for information on proper use and calibration of this equipment.

Allow at least the following time intervals between application and harvest:

Cotton, Soybeans 7 days
 Apples, Atemoya, Avocado, Breadfruit, Canistel,

Carambola, Cherry, Citrus, Dates, Grapes,
 Jaboticaba, Jackfruit, Longan, Lychee, Passion
 Fruit, Pear, Persimmons, Rutabagas, Sapodilla,
 Sapote, Soursop, Sugar apple, Tamarind 14 days
 Stone Fruit 17 days
 Nut Crops 21 days
 Sorghum (milo)* 40 days

*Do not use roller applicators. Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

ASPARAGUS

When applied as directed for "Cropping Systems", under the conditions described, this product controls weeds listed on this label in asparagus.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

Prior to Crop Emergence — Apply this product prior to planting or emergence of spears in an established crop for the control of emerged labeled annual and perennial weeds. **DO NOT APPLY WITHIN A WEEK BEFORE THE FIRST SPEARS EMERGE.**

Spot Treatment — Apply this product immediately after cutting, but prior to the emergence of new spears. Do not treat more than 10 percent of the total field to be harvested. Do not harvest within 5 days of treatment.

Postharvest — Apply this product after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears. Direct contact of the spray with the asparagus may result in serious crop injury.

NOTE: Select and use recommended types of spray equipment for postemergence postharvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of the spray with the crop.

BERRIES AND SMALL FRUIT

For cranberries, apply after fruit set and no later than 30 days before harvest.

Wiper applicators may be used in cranberries in accordance with instructions in this section.

For other berries, apply as a preplant broadcast application, or as a directed spray or wiper application post-planting.

See the "General Information" and "Mixing, Additives and Application Instructions" sections of this label for essential product performance information.

See the "Selective Equipment" part of the "Application Equipment and Techniques" section of this label for information on recommended use and calibration of this equipment.

For Wick or other Wiper Applicators — Mix 1 gallon of this product in 4 gallons of water to prepare a 20 percent solution. Apply the solution to emerged weeds. Apply after cranberry fruit set and no later than 30 days before harvest.

In severe infestations, reduce equipment ground speed

to insure that adequate amounts of this product are wiped on the weeds. A second treatment in the opposite direction may be beneficial.

Do not permit herbicide solution to contact desirable vegetation, including green shoots, canes, or foliage.

FALLOW AND REDUCED TILLAGE SYSTEMS

Use this product in fallow and reduced tillage systems for control of annual weeds prior to emergence of crops listed in this label. Refer to the "Weeds Controlled" section of this label for specific rates and instructions. This product may be applied using ground or aerial spray equipment. See the "Application Equipment and Techniques" section of this label for instructions.

■	TANK MIXTURES	■
■	RULER® plus BANVEL	■
■	RULER plus 2,4-D	■
■	RULER plus GOAL™	■

Applications of 2,4-D or Banvel must be made at least 7 days prior to planting corn. Applications of 2,4-D must be made at least 30 days prior to planting soybeans.

The addition of Banvel in a mixture with this product may provide short-term residual control of selected weed species. Some crop injury may occur if Banvel is applied within 45 days of planting. Refer to the Banvel and 2,4-D labels for cropping restrictions and other use instructions.

Ruler® plus Goal Tank Mixtures

This product alone or in tank mixtures with Goal will provide control of those weeds listed below.

Make applications when weeds are actively growing and at the recommended stages of growth. Avoid spraying when weeds are subject to moisture stress, when dust is on the foliage or when straw canopy covers the weeds.

RULER® 12 fluid oz/acre		RULER® 16 fluid oz/acre	
Wheat	18"*	Annual grasses at	
Barley	12"	left plus:	
Bluegrass,		Ryegrass,	
annual	6"	annual	6"
Barnyardgrass	6"	Chickweed	6"
Rye	6"	Groundsel	6"
		Marestail	6"
		Rocket, London	6"
		Shepherdspurse	6"
		Crabgrass	12"
		Johnsongrass,	
		seedling	12"
		Lambsquarters	12"
		Oats, wild	12"
		Pigweed, redroot	12"
		Mustards	12"

RULER 12 fluid oz/acre + GOAL** 2 to 4 fluid oz/acre		RULER 16 fluid oz/acre + GOAL** 2 to 4 fluid oz/acre	
Annual grasses above		Annual weeds above	
plus:		plus:	
Cheeseweed	3"	Cheeseweed	6"

RULER 12 fluid oz/acre + GOAL**	RULER 16 fluid oz/acre + GOAL**
2 to 4 fluid oz/acre	2 to 4 fluid oz/acre
Chickweed 3"	Groundsel 6"
Groundsel 3"	Chickweed 6"
Rocket, London 6"	Rocket, London 12"
Shepherdspurse 6"	Shepherdspurse 12"

*Maximum height or length in inches.

**Use the higher rate when weeds approach maximum recommended height or stands are dense.

™Goal is a trademark of Rohm and Haas Company.

ECOFARMING SYSTEMS

The Ecofarming System consists of the following rotation: winter wheat, corn/sorghum, ecofallow.

Use the following tank mixtures for control of emerged annual weeds before planting corn or sorghum in the Ecofarming System.

RULER® at 16 to 20 fluid ounces per acre plus

2,4-D at 0.375 to 0.5 pound a.i. per acre plus

ATRAZINE at 0.75 to 1 pound a.i. per acre plus

LASSO® at 2.5 to 3 quarts per acre

The above tank mixture should be applied in 28-0-0 or 32-0-0 liquid fertilizer carrier at 20 to 30 gallons per acre.

The liquid fertilizer may be diluted with water to achieve the required carrier volume.

WEEDS CONTROLLED—The following weeds, up to a maximum height of 4 inches, will be controlled:

Brome, downy <i>Bromus tectorum</i>	Lettuce, prickly <i>Lactuca serriola</i>
Cheat <i>Bromus secalinus</i>	Pigweed, redroot <i>Amaranthus retroflexus</i>
Foxtail, green <i>Setaria viridis</i>	Thistle, Russian <i>Salsola kali</i>
Foxtail, yellow <i>Setaria lutescens</i>	Wheat, volunteer <i>Triticum aestivum</i>
Kochia* <i>Kochia scoparia</i>	

*For improved control of kochia, add 4 fluid ounces per acre (0.125 pound a.i. per acre) of Banvel to the above tank mixture.

Risk of crop injury from 2,4-D or Banvel can be reduced by applying this treatment 7 to 14 days before planting.

Refer to the label booklet for Lasso herbicide for pre-emergence weed control achieved by this tank mixture.

Refer to the specific product labels for crop rotation restrictions and cautionary statements for all products used in these tank mixtures.

*Lasso is a registered trademark of Monsanto Company.

AID TO TILLAGE

This product, when used in conjunction with preplant tillage practices, will provide control of downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 8 fluid ounces of this product in 3 to 10 gallons of water per acre. Make applications when weeds are actively growing and before they are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and

before regrowth occurs. Allow at least 1 day after application before tillage. Tank mixtures with residual herbicides may result in reduced performance.

PASTURES

Apply this product prior to planting forage grasses and legumes.

Pasture or Hay Crop Renovation—When applied as a broadcast spray, this product controls the annual and perennial weeds listed in this label prior to planting forage grasses or legumes. Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Spot Treatment—When applied as a spot treatment as recommended, this product controls annual and perennial weeds listed in this label which are growing in pastures, forage grasses and forage legumes composed of bahiagrass, bermudagrass, bluegrass, brome, fescue, orchardgrass, ryegrass, timothy, wheatgrass, alfalfa or clover.

Wiper Application—When applied as directed, this product controls or suppresses the weeds listed under "Wiper Applicators" in the "Selective Equipment" section of this label.

For spot treatment and wiper application, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre should be treated at one time. Further applications may be made in the same area at 30-day intervals. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

SUGARCANE

When applied as directed for "Cropping Systems", under the condition described, this product controls those emerged annual and perennial weeds listed on this label growing in or around sugarcane or in fields to be planted to sugarcane. This product will also control undesirable sugarcane.

NOTE: Where repeat treatments are necessary, do not exceed a total of 10.6 quarts of this product per acre per year. Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

Broadcast Treatment—Apply this product in 10 to 40 gallons of water per acre on emerged weeds growing in fields to be planted to sugarcane.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

For removal of last stubble or ratoon cane, apply 4 to 5 quarts of this product in 10 to 40 gallons of water per acre to new growth having at least 7 or more new leaves. Allow 7 or more days after application before tillage.

Spot Treatment In or Around Sugarcane Fields—For dilution and rates of application using hand-held equipment, see "Mixing, Additives and Application Instructions" and "Weeds Controlled" sections of this label.

For control of volunteer or diseased sugarcane, make a 1 percent solution of this product in water and spray to wet the foliage of vegetation to be controlled.

NOTE: When spraying volunteer or diseased sugarcane, the plants should have at least 7 new leaves.

Avoid spray contact with healthy cane plants since severe damage or destruction may result.

Do not feed or graze treated sugarcane forage following application.

CONSERVATION TILLAGE, MINIMUM TILLAGE AND NO-TILL SYSTEMS CORN AND SOYBEANS Tank Mixtures

When applied as recommended under the conditions described, the tank mixtures listed in this section control many emerged weeds, and give preemergence control of many annual weeds where corn or soybeans will be planted directly into a cover crop, established sod, or in previous crop residues.

Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures. For mixing instructions, see the "Mixing, Additives and Application Instructions" section of this label.

Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solutions per acre before, during or after planting. Do not apply these mixtures after crop emergence.

The addition of 1 to 2 percent dry ammonium sulfate by weight may increase the performance of this product.

NOTE: When using these tank mixtures, do not exceed 4 quarts of this product per acre.

CORN

For residual control, this product may be tank-mixed with the following herbicides or combination of herbicides:

LASSO®/ALACHLOR	ATRAZINE
LARIAT®	CYANAZINE
BULLET®	SIMAZINE
DUAL™	PROWL™
BICEP™	

For improved burndown, this product may be tank-mixed with 2,4-D or dicamba. Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn. See the "Weeds Controlled" section for specific rate information.

SOYBEANS

For residual control, this product may be tank-mixed with the following herbicides or combination of herbicides:

CANOPY™	LOROX™ PLUS
COMMAND™	PREVIEW™
DUAL™	PROWL™
GEMINI™	TURBO™
LASSO®/ALACHLOR	SCEPTER™
LEXONE™	SENCOR™
LINURON	SQUADRON™

For improved burndown, this product may be tank-mixed with the following herbicides:

2,4-DB
2,4-D*

*Applications of 2,4-D must be made at least 30 days prior to planting soybeans. See the "Weeds Controlled" section for specific rate information.

CORN AND SOYBEANS

Annual Weeds—For difficult to control weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other

labeled annual weeds, apply 1 to 1.5 pints of this product per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall. For a complete list of annual weeds controlled, see the "Weeds Controlled" section of this label.

Perennial Weeds—At normal application times in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. See the "Weeds Controlled" section of this label for the proper stage of growth for perennial weeds.

Use of 2 to 4 quarts of this product per acre in the tank mixtures mentioned above, under these conditions provides top kill and reduces competition from many emerged perennial grass and broadleaf weeds. For emerged perennial weeds controlled, see the "Weeds Controlled" section of this label.

To obtain the desired stage of growth, it may be necessary to apply this product alone in the late summer or fall and then follow with a label-approved, seedling weed control program at planting.

USE OF THESE TANK MIXTURES FOR BERMUDAGRASS OR JOHNSONGRASS CONTROL IN MINIMUM TILLAGE SYSTEMS IS NOT RECOMMENDED. For bermudagrass control, follow the instructions under "Control of Perennial Weeds" section of this label and then use a label-approved, seedling weed-control program in a minimum tillage or conventional tillage system. For johnsongrass control, follow instructions under "Control of Perennial Weeds" section of this label, and then use a label approved seedling weed control program with conventional tillage.

*Bullet and Lariat are registered trademarks of Monsanto Company.

**Canopy, Bladex, Gemini, Lexone, Lorox Plus and Preview are trademarks of E. I. duPont de Nemours and Company.

**Dual is a trademark of Ciba-Geigy Corporation.

**Sencor and Turbo are trademarks of Bayer AG.

**Prowl, Scepter and Squadron are trademarks of American Cyanamid Company

**Command is a trademark of FMC Corporation.

PREHARVEST APPLICATIONS

COTTON

When applied as directed under the conditions described, this product controls annual and perennial weeds listed on this label prior to the harvest of cotton.

Broadcast Applications—This product may be applied using either aerial or ground spray equipment. For ground applications with broadcast equipment, apply this product in 10 to 20 gallons of water per acre. For aerial applications, apply this product in 3 to 10 gallons of water per acre.

FOR AERIAL APPLICATIONS, REFER TO THE "APPLICATION EQUIPMENT AND TECHNIQUES" AND "AERIAL EQUIPMENT" SECTIONS OF THIS LABEL.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR.

Weed Control—For specific rates of application and instructions for control of various annual and perennial weeds for this product used alone or in the following tank mixtures, see the "Weeds Controlled" section of this label.

To control johnsongrass using multiple directed or broadcast over-the-top spray equipment, apply 1 quart of this

product in 10 to 20 gallons of water per acre. Ensure complete coverage.

For partial control of field bindweed, apply 1 quart of this product in 3 to 20 gallons of water per acre. Apply when bindweed is actively growing and 12 inches or greater in length. Reduced performance may result if bindweed is under drought stress.

Tank Mixtures

RULER® plus DEF™ 6

RULER plus FOLEX™

RULER plus PREP™

RULER plus PREP plus DEF 6 or FOLEX

When applied as recommended under the conditions described, these tank mixtures control annual and perennial weeds listed on this label prior to the harvest of cotton. For application guidelines, precautions and use-rates, refer to the Def, Folex and Prep labels.

This product when tank mixed with Def 6 or Folex defoliant may provide enhancement of cotton leaf drop and regrowth inhibition.

Timing of Application — Apply this product or these tank mixtures for preharvest weed control after 60 percent of the cotton bolls have opened.

NOTE: DO NOT APPLY TO CROPS GROWN FOR SEED.

Allow a minimum of 7 days between application and harvest. Do not feed or graze treated cotton forage or hay following preharvest applications.

*Def is a trademark of Mobay Chemical Company

**Folex and Prep are trademarks of Rhone-Poulenc, Inc.

TREE AND VINE CROPS

This product is recommended for weed control in established groves, vineyards, or orchards, or for site preparation prior to transplanting crops listed in this section. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, or orchard guns, or with wiper applicator equipment, except as directed in this section. See the "Application Equipment and Techniques" section of this label for specific information on use of equipment.

When applying this product, refer to the "Weeds Controlled" section of this label and to specific recommendations in this section for rates to be used.

NOTE

Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. This product does not provide residual weed control. For subsequent weed control, use repeated applications of this product. Do not apply more than 10.6 quarts of this product per acre per year.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

AVOID PAINTING OUT STUMPS WITH THIS PRODUCT AS INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut and have not been allowed to regrow to the recommended stage for treatment.

For specific rates of applications and instructions, see the "Weeds Controlled" section of this label, and to specific recommendations which follow.

MIDDLES MANAGEMENT

FOR ANNUAL WEEDS IN MIDDLES BETWEEN ROWS OF TREE AND VINE CROPS

For citrus crops treat uniformly between trees.

■ **RULER®** ■
■ **RULER plus GOAL** ■

This product alone or in mixtures with Goal will control or suppress the annual weeds listed below.

Apply the recommended rates of this product, either alone or in mixtures with Goal, in 3 to 10 gallons of water per acre. Apply when weeds are actively growing and less than 6 inches in height or rosette diameter. If weeds are under drought stress, irrigate prior to application. Reduced control may occur if weeds have been mowed prior to application. Up to 48 fluid ounces per acre of this product may be used to control weeds which have been mowed, are stressed, or are growing in dense populations.

WEED SPECIES	MAXIMUM HEIGHT/ DIAMETER (INCHES)	Rate Per Acre	
		RULER (FLUID OUNCES)	GOAL (FLUID OUNCES)
Barley <i>Hordeum vulgare</i>	6	8	•
Bluegrass, annual <i>Poa annua</i>			
Barnyardgrass <i>Echinochloa crus-galli</i>	6	12	•
Chickweed, common <i>Stellaria media</i>			
Red Maids <i>Calandrinia ciliata</i>			
Crabgrass <i>Digitaria spp.</i>	6	16	•
Fleabane, flaxleaf <i>Conyza bonariensis</i>			OR
Groundsel, common <i>Senecio vulgaris</i>		16 to 32 + 4 to 16**	
Junglerice <i>Echinochloa colonum</i>			
Lambsquarters, common <i>Chenopodium album</i>			
Pigweed, redroot <i>Amaranthus retroflexus</i>			
Rocket, London <i>Sisymbrium irio</i>			
Ryegrass, common <i>Lolium multiflorum</i>			
Shepherdspurse <i>Capsella bursa-pastoris</i>			
Sowthistle, annual <i>Sonchus oleraceus</i>			
Cheeseweed, common <i>Malva spp.</i>	3	12 to 32 + 4 to 16	
Cheeseweed, common <i>Malva spp.</i>	6	16 to 32 + 4 to 16	
Filaree* <i>Erodium spp.</i>			

Horseweed/Marestail

Conyza canadensis

Nettle, stinging

Urtica dioica

Purslane, common*

Portulaca oleracea

*Suppression only

**The mixture of this product plus Goal is recommended when weeds are stressed or growing in dense populations.

STRIPS

FOR ANNUAL AND PERENNIAL WEEDS IN STRIPS OF TREE AND VINE CROPS

TANK MIXTURES

WITH RESIDUAL HERBICIDES

When applied as a tank mixture, this product provides control of the emerged annual weeds and control or suppression of emerged perennial weeds listed in this label. The following residual herbicides will provide pre-emergence control of those weeds listed in the individual product labels.

■	RULER® plus GOAL 1.6E	■
■	RULER plus KARMEX™ DF	■
■	RULER plus KROVAR I	■
■	RULER plus KROVAR II	■
■	RULER plus SIMAZINE, PRINCEP CALIBER™ 90	■
■	RULER plus SIMAZINE 4L	■
■	RULER plus SIMAZINE 80W	■
■	RULER plus SOLICAM™ 80DF	■
■	RULER plus SURFLAN™ AS	■
■	RULER plus SURFLAN 75W	■
■	RULER plus SIMAZINE (80W, or 4L, or PRINCEP CALIBER 90) plus SURFLAN (AS or 75W)	■
■	RULER plus GOAL (1.6E) plus SURFLAN (AS or 75W)	■
■	RULER plus GOAL (1.6E) plus SIMAZINE (80W, or 4L, or PRINCEP CALIBER 90)	■
■	RULER plus GOAL (1.6E) plus SURFLAN (AS or 75W) plus SIMAZINE (80W, 4L, or PRINCEP CALIBER 90)	■

Do not apply these tank mixtures in Puerto Rico.

Refer to the individual product labels for specific crops, rates, geographical restrictions and precautionary statements.

Read and carefully observe the label claims, cautionary statements, rates and all other information on the labels of all products.

*Karmex is a trademark of E.I. duPont de Nemours and Company.

RECOMMENDED RATES

Annual Weeds—Apply 1 to 5 quarts per acre of this product in these tank mixtures. Use rates at the higher end of the recommended range when weeds are stressed, growing in dense populations or are greater than 12 inches tall.

Perennial Weeds—Apply 1 pint to 5 quarts per acre of this product in these tank mixtures to control or suppress

perennial weeds. Follow the recommendations in the "Weeds Controlled" section of this label for stage of growth and application rates for specific perennial weeds.

RULER® plus GOAL plus SIMAZINE/SURFLAN

This product plus low rates of Goal in three-way or four-way mixtures with simazine and/or Surflan will provide postemergence control of the weeds listed below.

Refer to the individual simazine and Surflan labels for preemergence rates, weeds controlled, precautionary statements and other important information.

Apply 1 to 5 quarts per acre of this product plus 4 to 48 fluid ounces per acre of Goal plus labeled rates of simazine and/or Surflan in 3 to 40 gallons of water per acre to control the following weeds:

Barley, wild	Horseweed/Marestail
<i>Hordeum leporinum</i>	<i>Conyza canadensis</i>
Bluegrass, annual	Nettle, stinging
<i>Poa annua</i>	<i>Urtica dioica</i>
Cheeseweed, common	Pineappleweed
<i>Malva spp.</i>	<i>Matricaria matricariodes</i>
Chickweed, common	Rocket, London
<i>Stellaria media</i>	<i>Sisymbrium irio</i>
Filaree*	Shepherdspurse
<i>Erodium spp.</i>	<i>Capsella bursa-pastoris</i>
Fleabane, flaxleaf	Sowthistle, annual
<i>Conyza bonariensis</i>	<i>Sonchus oleraceus</i>
Groundsel, common	
<i>Senecio vulgaris</i>	

*Use a minimum of 1.5 quarts of this product in these mixtures.

NOTE: This recommendation does not preclude the use of Goal in these mixtures at higher, labeled rates for preemergence weed control.

PERENNIAL GRASS SUPPRESSION ORCHARD FLOORS

When applied as directed, this product will suppress vegetative growth as indicated below.

Bahiagrass

This product will provide significant inhibition of seed-head emergence and will suppress vegetative growth for a period of approximately 45 days with a single application and approximately 120 days with sequential applications. Apply this product 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 6 fluid ounces of this product in 10 to 25 gallons of water per acre.

Sequential applications of this product may be made at approximately 45-day intervals to extend the period of seedhead and vegetative growth suppression. For continued seedhead suppression, sequential applications must be made prior to seedhead emergence. Apply no more than 2 sequential applications per year. As a first sequential application, apply 4 fluid ounces of this product per acre. A second sequential application of 2 to 4 fluid ounces per acre may be made approximately 45 days after the last application.

Bermudagrass

For burndown, apply 1 to 2 quarts of this product in 3 to 20 gallons of water per acre. Use 1 quart in 3 to 20 gallons of water per acre east of the Rocky Mountains. Use 1 to 2 quarts of this product in 3 to 10 gallons of water per acre west of the Rocky Mountains. Use this treatment only if reduction of the bermudagrass stand

can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

Suppression only — (east of the Rocky Mountains). Apply 6 to 16 fluid ounces of this product in 3 to 20 gallons of water per acre no sooner than 1 to 2 weeks after full green-up. Mowing prior to application may occur provided a minimum height of 3 inches is maintained. Rates of 6 to 10 fluid ounces should be used where a lesser degree of suppression is desired. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated.

Suppression only — (west of the Rocky Mountains). Apply 16 fluid ounces of this product in 3 to 10 gallons of water per acre to bermudagrass up to 6 inches in height and no sooner than 1 to 2 weeks after full green-up. Mowing prior to application may occur provided a minimum height of 3 inches is maintained. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated.

Cool-Season Grass Covers

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of this product in 10 to 20 gallons of water per acre. For best suppression, add ammonium sulfate to the spray solution at a rate of 2 percent by weight or 17 pounds per 100 gallons of spray solution.

For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of this product per acre. Do not add ammonium sulfate.

For best results, mow cool-season grass cover in the spring to even the height and apply the recommended rate of this product 3 to 4 days after mowing. Avoid treating cool-season grass covers under poor growing conditions, such as drought stress (drip irrigation), disease or insect damage.

LOW-VOLUME APPLICATION (Florida and Texas)

For burndown or control of the weeds listed, apply the recommended rate of this product in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

Annual Weeds

Goatweed—Apply 2 to 3 quarts per acre of this product plus 17 pounds of ammonium sulfate per 100 gallons of water. Apply in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are less than 8 inches tall and 3 quarts per acre when plants are greater than 8 inches. If goatweed is greater than 8 inches, the addition of Krovar™ II or Karmex™ may improve control. Use labeled rates for these residual products.

Read and carefully observe the label claims, cautionary statements, rates and all other information on the Krovar II and Karmex labels.

Perennial Weeds

Apply when weeds are actively growing and at the growth states listed in the "Perennial Weeds Controlled" section of this label. If perennial weeds are mowed, allow weeds to regrow to the recommended stage of growth.

S = Suppression, B = Burndown, C = Control,
PC = Partial control

WEED SPECIES	RULER® RATE PER ACRE			
	1 qt	2 qt	3 qt	5 qt
Bermudagrass	B	•	PC	C
Guineagrass				
Texas and Florida Ridge	B	C	C	C
Florida Flatwoods	•	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	•	PC	C

™Caliber is a trademark of Ciba-Geigy Corporation.

™Goal is a trademark of Rohm and Haas Company.

™Karmex and Krovar are trademarks of E. I. du Pont de Nemours and Company.

™Solcam is a trademark of Sandoz, Inc.

™Surflan is a trademark of Elanco Products Company.

TREE CROPS

Citrus®: citron, grapefruit, kumquat, lemon, lime, orange, pummelo, tangelo, tangerine, tangors.

Nuts®: almond, chestnuts, filbert, macadamia, pecan, pistachio, walnut.

Pome Fruit®: apple, pear.

Stone Fruit®: apricots, cherries, nectarines, olives, peaches, plums/prunes.

For cherries, any application equipment listed in this section may be used in all states.

For citron and olives, apply as a directed spray only.

Any application equipment listed in this section may be used in apricots, nectarines, peaches, and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah, and Washington, except for peaches grown in the states specified in the following paragraph. In all other states use wiper equipment only.

For PEACHES grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, and Tennessee only, apply with a shielded boom sprayer or shielded wiper applicator which prevents any contact of this product with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees which have been planted in the orchard for 2 or more years. **EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED.**

Tropical Fruit: acerola, atemoya®, avocado®, banana (plantains), breadfruit®, canistel®, carambola®, coffee®, dates®, figs®, guava, jaboticaba®, jackfruit®, longan®, lychee®, mango®, papaya, passion fruit®, persimmons®, sapodilla, sapote, soursop, sugar apple®, tamarind, tea. Allow a minimum of 1 day between last application and harvest of guava and papaya. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

NOTE:

*Allow a minimum of 14 days between last application and harvest.

**Allow a minimum of 21 days between last application and harvest.

***Allow a minimum of 17 days between last application and harvest.

****Allow a minimum of 28 days between last application and harvest.

VINE CROPS

Kiwi Fruit

Grapes: Any variety of table, wine, or raisin grape may be treated with any equipment listed in this section.

In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.

NOTE:

Applications should not be made when green shoots, canes, or foliage are in the spray zone.

Allow a minimum of 14 days between last application and harvest.

Product protected by

U.S. Pat. No. 4,405,531.

Other patents pending.

No license granted under any non-U.S. patent(s).

EPA Reg. No. 524-326-2935

GPL1100.04/CG

In case of a human health emergency involving this product, Call Collect, day or night, (314) 694-4000.

Distributed by
Wilbur-Ellis Company
P.O. Box 16458
Fresno, CA 93755

WEEDAR® 64

Brand Herbicide

RHÔNE-POULENC

The 2,4-D Amine Weed Killer

ACTIVE INGREDIENT:

2,4-Dichlorophenoxyacetic acid, dimethylamine salt*46.8%

INERT INGREDIENTS:53.2%

*2,4-Dichlorophenoxyacetic acid equivalent 38.9% by weight or 3.8 pounds per gallon

*Isomer specific by AOAC method No. 6.D01-5

E.P.A. Reg. No. 264-2AA

E.P.A. Est. No. 264-MO-01

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

PRECAUTION AL USARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicado ampliamente.

For PRODUCT USE Information Call 1-800-334-9745

For EMERGENCY Information ONLY Call 24 Hours A Day 1-800-334-7577.

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash skin with plenty of soap and water. Remove contaminated clothing. Get medical attention.

IF SWALLOWED: If patient is conscious and alert, give 2 to 3 glasses of water or milk to drink. If available, give one tablespoon of Syrup of Ipecac to induce vomiting. Alternatively, induce vomiting by touching back of throat with finger. Do not make an unconscious person vomit. Get medical attention.

IF IN EYES: Flush with water for at least 15 minutes. Get medical attention, **PREFERABLY AN OPHTHALMOLOGIST.**

IF INHALED: Move to an uncontaminated area. Get medical attention.

NOTE TO PHYSICIAN

This product contains a phenoxy herbicidal chemical. There is no specific antidote. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

PRECAUTIONARY STATEMENTS

DANGER

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. May be fatal if absorbed through the skin. Causes irreversible eye damage. Avoid breathing vapors or spray mist. Do not get in eyes, on skin, or on clothing. When handling this product, wear goggles or safety glasses, protective clothing and chemical resistant gloves. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water except as specified on this label. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas. Do not use the same spray equipment for other purposes unless thoroughly cleaned.

Do not contaminate water used for irrigation or domestic purposes (except as specifically recommended on this label) especially in areas where grapes, cotton, tomatoes or other susceptible plants are grown.

Do not treat irrigation ditches in areas where water will be used to overhead (sprinkler) irrigate susceptible crops especially grapes, tomatoes, tobacco, and cotton.

MIXING AND LOADING: Most cases of ground water contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of ground water supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent ground water contamination.

Do not apply WEEDAR® 64 Broadleaf Herbicide directly to, or permit to drift onto cotton, okra, grapes tomatoes, fruit trees, vegetables, flowers or other desirable crop or ornamental plants which are susceptible to 2,4-D herbicide. Do not apply near susceptible plants since very small quantities of the 2,4-D will cause severe injury during the growing or dormant periods. Crops contacted by WEEDAR® 64 Broadleaf Herbicide sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.

Do not apply when a temperature air inversion exists. Such a condition is characterized by little or no air movement and an increase in air temperature with an increase in height. In humid regions, a fog or mist may form. An inversion may be detected by producing a smoke column and checking for a layering effect. If questions exist pertaining to the existence of an inversion, consult with local weather services before making an application.

Use coarse sprays to minimize drift. Do not apply with hollow cone-type insecticide or other nozzles that produce fine spray droplets. Drift from aerial or ground application may be reduced by: (1) applying as near to the target as possible in order to obtain coverage; (2) by increasing the volume of spray mix per acre; (3) by decreasing the pounds of pressure at the nozzle tips; and (4) by using nozzles which produce a coarse spray pattern; (5) by not applying when wind is blowing toward susceptible crops or valuable plants.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read entire label before using this label.

GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application and until sprays have dried. Regular long-sleeved work clothing should be worn when working in treated fields. See PRECAUTIONARY STATEMENTS, STATEMENT OF PRACTICAL TREATMENT and NOTE TO PHYSICIAN for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: appropriate signal word (DANGER - PELIGRO), area treated with WEEDAR® 64 Broadleaf Herbicide, date of application, appropriate protective clothing, and reentry interval (i.e. until sprays have dried).

STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food or feed by storage or disposal. Store in original container in a dry secured storage area. Keep container tightly closed when not in use.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse or (equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL CAUTIONS AND RESTRICTIONS

Do not apply WEEDAR® 64 Broadleaf Herbicide through any type of irrigation system.

Do not use in or near a greenhouse.

RESTRICTIONS AND LIMITATIONS FOR USE ON CEREAL GRAINS

For aerial application on grain, apply WEEDAR® 64 Broadleaf Herbicide in 3 to 10 gallons of water per acre.

For ground application a minimum of 10 to 15 gallons of water per acre is recommended for proper spray coverage.

Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 2 weeks after treatment.

Do not feed treated straw to livestock if an emergency treatment as described below is applied.

RESTRICTIONS AND LIMITATIONS FOR USE ON CORN AND SORGHUM

Do not forage or feed fodder for 7 days following application.

RESTRICTIONS AND LIMITATIONS FOR USE IN FALLOWLAND AND CROP STUBBLE

Do not plant any crop for 3 months after treatment or until chemical has disappeared from the soil.

RESTRICTIONS AND LIMITATIONS FOR USE IN PASTURES AND RANGELANDS

Do not graze (dairy) cattle in treated areas for 7 days after application.

Do not cut forage for hay within 30 days of application.

Do not permit dairy animals or meat animals being finished for slaughter to forage treated fields within 3 days of slaughter.

RESTRICTIONS AND LIMITATIONS FOR USE ON CONSERVATION RESERVE PROGRAM AREAS:

Use at least 2 gallons of water per acre by air and 5 gallons of water per acre by ground.

Do not harvest or graze treated Conservation Reserve Program areas.

Do not apply to grasses in the boot to dough stage if grass seed production is desired.

RESTRICTIONS AND LIMITATIONS FOR USE ON GRASSES FOR SEED PRODUCTION

Do not graze dairy animals or cut forage for hay within 7 days of application.

RESTRICTIONS AND LIMITATIONS FOR USE ON NON-CROPLAND

Do not graze dairy animals for 7 days following application.

Use sufficient gallonage for thorough and uniform coverage.

RESTRICTIONS AND LIMITATIONS FOR USE IN STONE FRUIT AND NUT ORCHARDS

Do not apply to bare ground as injury may result.

Do not apply immediately before irrigation and withhold irrigation for 2 days before and for 3 days after treatment.

Do not allow spray to drift onto or contact foliage, fruit, stems, trunks of trees or exposed roots as injury may result.

Do not apply to newly established or young orchards. Trees must be at least 1 year old and in vigorous condition.

Do not apply during bloom.

Do not graze or feed cover crops from treated orchards.

Do not make more than 2 applications per year.

Do not harvest stone fruit within 40 days of application.

Do not harvest nuts within 60 days of application.

MIXING INSTRUCTIONS

Mix WEEDAR® 64 Broadleaf Herbicide only with water, unless otherwise directed on this label. Add about one-half the water to the mixing tank, then add WEEDAR® 64 with agitation and finally the rest of water with continuing agitation.

NOTE: Adding oil, wetting agent, or other surfactants to the spray may increase effectiveness on weeds but also may reduce selectivity to crops, resulting in crop damage.

COMPATIBILITY

If WEEDAR® 64 Broadleaf Herbicide is to be tank mixed with fertilizers or with other pesticides, compatibility should be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 qt) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing.

Read and follow all directions and precautions on this label and on the labels of any products for which a tank mixture is being considered.

APPLICATION PROCEDURES

Apply by air or ground equipment in sufficient gallonage to obtain adequate coverage, except as otherwise directed on this label.

Use 2 or more gallons of water per acre for aerial application and 10 or more gallons of water per acre for ground application.

GENERAL INFORMATION

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY DO NOT USE THIS PRODUCT.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply WEEDAR® 64 Broadleaf Herbicide only to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult your seed company, state Agricultural Extension Service or qualified crop consultant for advice.

Be sure that use of this product conforms to all applicable laws, rules and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator should become familiar with these laws, rules or regulations and follow them exactly.

GENERAL WEED LIST

Annual and Biennial Weeds

*beggarticks	*mallow (venice or little)	*Russian thistle
bullthistle	marshelder	Salsify (western or common)
coffeeweed	morningglory (common, ivy, wooly)	*smartweeds (annual species)
common cocklebur	*musk thistle(***)	sowthistles (annual or spiny)
common burdock	mustards (except blue mustard)	sunflower
common evening primrose	pepper weeds (except perennial)	*vervains
common lambsquarters	**pigweeds (Amaranthus spp.)	vetches
hairy galinsoga	prickly lettuce	wild carrot
jimsonweed	ragweed (common or giant)	wild lettuce
*knotweed	rough fleabane	wild parsnips

Perennial Weeds

*bindweed (hedge, field, European)	*goldenrod	**nettles (including stinging)
blue lettuce	*ground ivy	*orange hawkweed
*Canada thistle	healall	plantains
catnip	*hoary cress	sowthistle (perennial)
chicory	*ironweed	*vervains
dandelion	Jerusalem-artichoke	*wild garlic
*docks	many flowered aster	*wild onion
*dogbanes		

*These species may require repeated applications and/or use of the higher rate recommended on this product label even under ideal conditions for application.

**Control of pigweeds in the High Plains area of Texas and Oklahoma may not be satisfactory with this product.

***Not registered for control of musk thistle in California.

SPECIFIC USE DIRECTIONS

CEREAL GRAINS

CROP	AMOUNT OF WEEDAR® 64 PER ACRE	DIRECTIONS
Wheat, Barley, Oats and Rye (not underseeded with legumes) Postemergence Annual and biennial broadleaf weeds Perennial broadleaf weeds	1/2 to 2 pints* 1 to 2 pints*	Apply after grain is fully tillered (usually 4 to 8 inches high) but not forming joints in the stem. Do not spray grain in the boot to dough stage.
Wheat, Barley, Oats and Rye (underseeded with legumes)	1/4 to 1/2 pint*	Apply after grain is 8 inches tall. Do not spray grain in boot to dough stage. Do not spray alfalfa or sweet clover unless the infestation is severe and injury to these legumes can be tolerated.
Emergency weed control in Wheat Perennial broadleaf weeds	3 pints	Apply when weeds are approaching bud stage, after the grain dough stage. Do not spray during the boot to dough stage. The 3 pints per acre application can product injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury.

*Use the lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control problem justifies the grain damage risk. Do not apply WEEDAR® 64 to grain in the seedling stage.

CORN AND SORGHUM

CROP	AMOUNT OF WEEDAR® 64 PER ACRE	DIRECTIONS
CORN (Field and Sweet)		
Preplant	1 to 2 pints	To control emerged broadleaf weed seedlings or existing cover crops prior to planting corn, apply 7 to 14 days before planting. Do not use on light, sandy soil, or where soil moisture is inadequate for normal weed growth. Use high rate for less susceptible weeds or cover crops such as alfalfa.
Preemergence	2 to 3 pints	Apply 3 to 5 days after planting but before corn emerges. Do not use on light, sandy soils or where soil moisture is low.
Postemergence		Apply when weeds are small and corn is less than 8 inches tall (to top of canopy). When corn is over 8 inches tall, use drop nozzles and keep spray off foliage. Treat perennial weeds when they are in the bud to bloom stage. Do not spray corn in the tassel to dough stage. Corn treated with 2,4-D may become temporarily brittle. Winds or cultivation may cause stalk breakage during the period of time when the corn is brittle.
Annual broadleaf weeds	1/2 to 1 pint	
Perennial broadleaf weeds	1 to 1 1/2 pints	
Grain Sorghum (Milo)		
Postemergence	1 pint	Apply when sorghum is 6 to 15 inches tall. If sorghum is taller than 8 inches to top of the canopy, use drop nozzles and keep spray off the foliage. Do not treat during the boot, flowering or dough stage.

RICE, SUGARCANE, FALLOWLAND AND CROP STUBBLE

CROP	AMOUNT OF WEEDAR® 64 PER ACRE	DIRECTIONS
Rice	1 to 2 1/2 pints	Apply when rice is in the late tillering stage of development at the time of first joint development. Do not apply after panicle initiation, after rice internodes exceed one-half inch, at early seedling, early panicle, boot or heading stages. Consult local university or Agricultural Extension Service specialists for more specific information on rates and timing of application.
Sugarcane		
Preemergence	4 pints	Apply before canes appear for control of emerged broadleaf weeds. Apply after cane emerges and through lay-by. DO NOT USE IN CALIFORNIA.
Postemergence	1 1/2 to 2 pints	
Fallowland and Crop Stubble		
Annual broadleaf weeds	1 to 2 pints	Use the lower rate when weeds are small (2 to 3 inches tall) and actively growing. Use the higher rate on older and drought-stressed plants.
Biennial broadleaf weeds	2 to 4 pints	Spray while musk thistles or other biennial species are in the seedling to rosette stage and before flower stalks become apparent. The lower rate can be used in the spring during rosette stage. Use the highest rate in the fall or after flower stalks have developed.
Perennial broadleaf weeds	2 to 6 pints	Spray weed in the bud to bloom stage or while in good vegetative growth. Do not disturb treated areas for at least 2 weeks after treatment, or until tops are dead.
Wild garlic and onion in crop stubble	4 to 6 pints	Apply to new regrowth of wild garlic or onion which occurs in the fall following harvest of small grains, corn or grain sorghum.

ESTABLISHED GRASS PASTURES, RANGELAND, AND CONSERVATION RESERVE PROGRAM AREAS

WEEDS	AMOUNT OF WEEDAR® 64 PER ACRE	DIRECTIONS
Annual broadleaf weeds	2 pints	Apply when weeds are small and actively growing and prior to bud stage. Spray while musk thistles or other biennial species are in the seedling to rosette stage and before flower stalks become apparent. The lower rate can be used in the spring during rosette stage. Use the highest rate in the fall or after flower stalks have developed. Do not apply to newly seeded areas until grass is well established. Do not apply to grass in the early boot through milk stage if grass seed production is desired. Bentgrass and legumes may be injured by this treatment.
Biennial and perennial broadleaf weeds	2 to 4 pints	

CONSERVATION RESERVE PROGRAM AREAS

WEEDS	AMOUNT OF WEEDAR® 64 PER ACRE	DIRECTIONS
Annual broadleaf weeds	1/2 to 1 pint	Apply to actively growing annual broadleaf weeds. Use 1/2 to 1 pint when weeds are small; use higher rates on older weeds. Do not apply to young grasses with fewer than 6 leaves or prior to tillering, as excessive injury may result. Do not apply more than 1 pint until grasses are well established as excessive injury may result.
In young grasses		
In established grasses	1/2 to 2 pints	
Biennial and perennial broadleaf weeds	2 to 4 pints	Treat when biennial weeds are in the seedling to rosette stage and before flower stalks become apparent. Treat perennial weeds in the bud to bloom stage. Apply to actively growing weeds.
In established grasses		

GRASSES FOR SEED PRODUCTION

WEEDS	AMOUNT OF WEEDAR® 64 PER ACRE	DIRECTIONS
Annual and perennial broadleaf weeds	2 to 4 pints	Apply to established stands in spring from tiller to early boot stage. Do not spray in boot stage. New spring seedlings may be treated with the lower rate after grass seedlings have at least 5 leaves. Perennial weed regrowth may be treated in the fall. DO NOT USE IN CALIFORNIA.

NON-CROPLAND

fencerows, roadsides, drainage ditches, golf courses, cemeteries, parks, turfgrass, and other grass areas

WEEDS	AMOUNT OF WEEDAR® 64 PER ACRE	DIRECTIONS
Annual broadleaf weeds	2 to 4 pints	Treat when weeds are young and actively growing. Perennial weeds should be near the bud stage, but not flowering at application. Do not use on susceptible southern grasses such as St. Augustine. Do not apply to newly seeded areas until grass is well established. Bentgrass, clover, legumes and dichondria may be injured by this treatment.
Biennial and perennial broadleaf weeds	4 to 8 pints	

SPOT TREATMENT IN NON-CROP AREAS

Mix 2 to 3 fluid ounces of WEEDAR® 64 Broadleaf Herbicide in 3 gallons of water. Wet all weeds and stems thoroughly. For best results, treat when weeds are actively growing.

FORESTRY - TREE INJECTION

For controlling species such as alder, aspen, birch, blackgum, cherry, oak, sweetgum, and tulip poplar

Make injections as near to the root collar as possible, using one injection per inch of trunk dbh (4 1/2 feet). For resistant species such as hickory, injections should overlap. For best results, injections should be made during the growing season, May 15th through October 15th.

For Dilute Injection: Mix 1 gallon of WEEDAR® 64 Broadleaf Herbicide in 19 gallons of water for dilute injections.

For Concentrate Injections: Use 1 to 2 ml of concentrate WEEDAR® 64 Broadleaf Herbicide per injection. The injection bit must penetrate the inner bark.

STONE FRUIT AND NUT ORCHARDS

WEEDS IN CROP	AMOUNT OF WEEDAR® 64 PER ACRE	DIRECTIONS
Annual broadleaf weeds	3 pints	For control of weeds on the orchard floor, apply using coarse sprays and low pressure in sufficient volume of water to obtain thorough wetting of weeds. Treat when weeds are small and actively growing. Do not use on light, sandy soil. DO NOT USE IN CALIFORNIA.

WEEDS AND BRUSH IRRIGATION CANAL DITCHBANKS

(Seventeen Western States: Arizona, California, Colorado, Idaho, Kansas, Montana, Nebraska, New Mexico, Nevada, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington, and Wyoming).

For control of annual and perennial broadleaf weeds, apply 1 to 2 quarts of WEEDAR® 64 Broadleaf Herbicide per acre in approximately 20 to 100 gallons per acre. Treat when weeds are young and actively growing before the bud or early bloom stage. For harder-to-control weeds, a repeat spray after 3 to 4 weeks using the same rates may be needed for maximum results. Apply no more than two treatments per season.

For woody brush and patches of perennial broadleaf weeds, mix 1 gallon of WEEDAR® 64 in 150 gallons of water. Wet foliage thoroughly using about 1 gallon of solution per square rod.

SPRAYING INSTRUCTIONS

Apply with low pressure (10 to 40 psi) power spray equipment mounted on a truck, tractor, or boat. Apply while traveling upstream to avoid accidental concentration of chemical into water. Spray when the air is fairly calm, 5 mph or less. Do not use on small canals (less than 10 cfs) where water will be used for drinking purposes.

Boom spraying onto water surface must be held to a minimum and no cross-stream spraying to opposite banks should be permitted. When spraying shoreline weeds, allow no more than 2 foot overspray onto water with an average of less than 1 foot overspray to prevent introduction of greater than negligible amounts of chemical into the water.

Do not allow dairy animals to graze on treated areas for at least 7 days after spraying. Water within treated banks should not be fished.

AQUATIC WEED CONTROL

For use in ponds, lakes, reservoirs, marshes, bayous, drainage ditches, canals, rivers and streams that are quiescent or slow moving.

NOTICE TO APPLICATORS

State and Local Coordination: Before application, coordination and approval of local and state authorities may be required, either by letter of agreement or issuance of special permits for such use.

Fish Toxicity - Oxygen Ratio: Fish breathe oxygen in the water and a water - oxygen ratio must be maintained. Decaying weeds use up oxygen. To avoid fish kill from decaying plant material do not treat more than one half the lake or pond at one time. For large bodies of weed infested waters leave buffer strips of at least 100 feet wide and delay treatment of these strips for 4 to 5 weeks or until the dead vegetation has decomposed.

Wind Velocity - Ground or Surface Application: Do not apply when wind speeds are at or above 10 mph. Air Application: Do not apply when wind speeds are at or above 5 mph. The restrictions do not apply to subsurface applications used in weed control programs.

Irrigation: Only the use of treated waters for irrigation for three weeks after treatment unless an approved assay shows that the water does not contain more than 0.1 ppm 2,4-D acid. Do not treat irrigation ditches in areas where water will be used to overhead sprinkler irrigate susceptible crops especially grapes, tomatoes and cotton.

Potable Water: Delay the use of treated water for domestic purposes for a period of three weeks or until such time as an approved agency shows that the water contains no more than 0.1 ppm 2,4-D acid.

Water Hyacinth (*Eichornia crassipe*) - Directions For Use

WEEDAR® 64 will control water hyacinth with surface and air applications.

Amounts to Use: 2 to 4 quarts (4 lb. acid equivalent per gallon) per acre. **Spray the weed mass only.** Use 4 quarts when plants are matured or when the weed mass is dense.

When To Apply: Spray when water hyacinth plants are actively growing. Repeat as necessary to kill regrowth and hyacinth plants missed in the previous operation.

How To Use - Surface Application: Use power sprayers operated with a boom or spray gun mounted on a boat, tractor or truck. Thorough wetting of foliage is essential for maximum control. Use 100 to 400 gal./A of spray mixture. Special precautions such as the use of low pressure, large nozzles and thickening agents should be taken to avoid spray drift in areas of sensitive crops. For DIRECTA-SPRA™ operation use WEEDAR® 64 with 1 pint of drift control agent in 50 to 100 gallons of water. For other applications, follow the drift control agent label for mixing directions. **Air Application:** Use drift control spray equipment or thickening agents mixed into the spray solution. Apply 1.0 gallon per acre of WEEDAR® 64 through standard boom systems with a minimum of 5 gallons of spray mix per acre. For MICROFOIL® drift control spray systems, apply WEEDAR® 64 in 12 to 15 gallons spray mix per acre.

2,4-D Acid Equivalent	1/2 lb.	1 lb.	2 lbs.	3 lbs.	4 lbs.
WEEDAR® 64	1 pt.	2 pts.	2 qts.	3 qts.	4 qts.

Water Milfoil (*Myriophyllum spicatum*) - Directions For Use

For Eurasian Water Milfoil in programs conducted by the Tennessee Valley Authority in dams and reservoirs of the TVA system.

WEEDAR® 64 will control water milfoil with surface, subsurface and air applications.

How To Use: To control water milfoil when less than 5 gallons of concentrate per acre is recommended, dilute the concentrate with water to apply a minimum of 5 gallons of spray mix per acre. Do not treat within 1/2 mile of potable water intakes. Shoreline areas should be treated by sub-surface injection applied by boat to avoid aerial drift. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment washwaters.

Open Water Areas: To reduce contamination and prevent undue exposure to fish and other aquatic organism, do not treat water areas that are not infested with aquatic weeds.

Amounts To Use: Apply 2.5 to 10 gallons of WEEDAR® 64 per acre. The higher rate is used in areas of greater water exchange. These areas may require a repeat application.

When To Apply: For best results, apply in spring or early summer when milfoil starts to grow. This timing can be checked by sampling the lake bottom in areas heavily infested with weeds the year before.

Subsurface Application: Apply 2.5 to 10 gallons of WEEDAR® 64 per acre as a concentrate directly into the water through boat mounted distribution systems.

Surface Application: Apply 2.5 to 10 gallons of WEEDAR® 64 per acre in a minimum spray volume of 5 gallons mix per acre.

Air Application: Use drift control spray equipment or thickening agents mixed into the spray solution. Apply 2.5 to 10 gallons per acre of WEEDAR® 64 through standard boom systems with a minimum of 5 gallons of spray mix per acre. For MICROFOIL® drift control spray systems apply WEEDAR® 64 in 12 to 15 gallons spray mix per acre.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

THIS SPECIMEN LABEL IS INTENDED FOR USE ONLY AS A GUIDE IN PROVIDING GENERAL INFORMATION REGARDING THE DIRECTIONS, WARNINGS AND CAUTIONS ASSOCIATED WITH THE USE OF THIS PRODUCT. AS WITH ANY AGRICULTURAL CHEMICAL, ALWAYS FOLLOW THE LABEL INSTRUCTIONS ON THE PACKAGE BEFORE USING.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

Form No. 93482
Printed USA - 9/91

WEEDAR and MICROFOIL are registered trademarks of RHONE-POULENC.
DIRECTA-SPRA is a trademark of RHONE-POULENC.
©1991 Rhône-Poulenc Ag Company
Made in USA.

EPA Approval: 6/24/91

ENVIRONMENTAL ASSESSMENT

NOXIOUS WEED AND EXOTIC VEGETATION MANAGEMENT PLAN

**Montana Fish Wildlife, & Parks
Region One**

DECEMBER 1994

TABLE OF CONTENTS

	Page
LIST OF TABLES	iv
LIST OF FIGURES	vi
1.0 THE PURPOSE OF AND NEED FOR ACTION	1
1.1 Introduction	1
1.2 Noxious Weeds on Region One Properties	2
1.2.1 The Setting and Current Conditions	2
1.2.2 Noxious Weed Treatment History	3
1.3 Weed Management Goals	8
1.4 Programs and Legal Framework Affecting Region One's Weed Plan	8
1.4.1 Wildlife Management Area Management Plans	8
1.4.2 County Weed District 5-Year Plans	12
1.4.3 Other Noxious Weed Management Plans	12
1.4.4 Noxious Weed Trust Fund Environmental Impact Statement	16
1.4.5 Other Applicable Laws and Regulations	15
1.5 Decisions to be Made	15
1.6 Public Concerns and Issues	16
2.0 TREATMENT METHODS AND ALTERNATIVES	17
2.1 Treatment Methods	17
2.1.1 Chemical Methods	17
2.1.2 Non-chemical Methods	19
2.2 Description and Comparison of the Alternatives	19
2.2.1 Alternative 1: Status Quo	20
2.2.2 Alternative 2: Integrated Noxious Weed Management	20
2.2.3 Alternative 3: Chemical Methods	21
2.2.4 Alternative 4: Non-Chemical Methods	21
2.3 The Preferred Alternative	22
2.4 Cumulative Effects and Irreversible or Irretrievable Commitments	22
3.0 THE AFFECTED ENVIRONMENT AND ANALYSIS OF POTENTIAL IMPACTS	23
3.1 Air Quality	27
3.1.1 Current Conditions	27
3.1.2 Analysis of Potential Impacts	27
3.2 Geology and Soils	28
3.2.1 Current Conditions	28
3.2.2 Analysis of Potential Impacts	28

TABLE OF CONTENTS, continued

	Page
3.3 Groundwater	30
3.3.1 Current Conditions	30
3.3.2 Analysis of Potential Impacts	30
3.4 Surface Water	33
3.4.1 Current Conditions	33
3.4.2 Analysis of Potential Impacts	34
3.5 Fish and Wildlife	37
3.5.1 Current Conditions	37
3.5.2 Analysis of Potential Impacts	38
3.6 Vegetation	45
3.6.1 Current Conditions	45
3.6.2 Analysis of Potential Impacts	46
3.7 Recreation	49
3.7.1 Current Conditions	49
3.7.2 Analysis of Potential Impacts	49
3.8 Human Health	51
3.8.1 Current Conditions	51
3.8.2 Analysis of Potential Impacts	51
3.9 Cultural and Historic Resources	54
3.9.1 Current Conditions	54
3.9.2 Analysis of Potential Impacts	55
3.10 Environmental Review Criteria	57
 4.0 SUMMARY	 59
 5.0 LIST OF CONTRIBUTORS AND REVIEWERS	 62
 6.0 REFERENCES	 63
 7.0 APPENDICES	 68
APPENDIX A Herbicide Efficacy on State-Declared Noxious Weed Species Properties of Commonly Used Herbicides	
APPENDIX B Fish and Wildlife Species Recorded in Latilongs Corresponding with Fish, Wildlife & Parks Region One	

LIST OF TABLES

	Page
Table 1. Noxious weed species present and status on Fish, Wildlife & Parks Wildlife Management Areas in Region One	6
Table 2. Noxious weed species present and status on Fish, Wildlife & Parks Fishing Access in Region One, by drainage	9
Table 3. Noxious weed species present and status on Montana State Parks in Region One	13
Table 4. Total expenditures by Fish, Wildlife & Parks Region One for weed management on its Wildlife Management Areas, Fishing Access, and State Parks during fiscal years (July 1 - June 30) 1986 and 1992	3
Table 5. Herbicides proposed for use on Fish, Wildlife & Parks Region One properties, registered trade name and application rates on a per acre basis according to manufacturer label recommendations.	17
Table 6. Summary of environmental impacts resulting from the implementation of the Noxious Weed Management Plan emphasizing Integrated Noxious Weed Management	24
Table 7. Maximum concentrations not to be exceeded in drinking water for herbicides proposed for use by Fish, Wildlife & Parks in Region One. Information provided by the United States Environmental Protection Agency, Office of Drinking Water, 1990	37
Table 8. A partial list of fish and wildlife species in northwest Montana	38
Table 9. Median lethal dose (LD ₅₀) for domestic animals expressed in milligrams per kilogram for 3 herbicides proposed for use on Department of Fish, Wildlife and Park properties in Region One	41

LIST OF TABLES, continued

	Page
Table 10. Routine and extreme dose levels of 2,4-D and glyphosate compared to the median lethal dose (LD ₅₀) for various animal species. Table is adapted from the Bureau of Land Management Northwest Area Noxious Weed Control Program Final Environmental Impact Statement, 1985	43
Table 11. Plant species of concern on Fish, Wildlife & Parks properties in Region One, as identified by the Montana Natural Heritage Program, 1994	48
Table 12. Summary and comparison of environmental impacts resulting from the implementation of each Alternative	63

LIST OF FIGURES

	Page
Figure 1. Properties owned by or leased to Fish, Wildlife & Parks in Region One	4

1.0 THE PURPOSE OF AND NEED FOR ACTION

1.1 Introduction

As mandated by the County Noxious Weed Control Act of 1979 (MCA 7-22-2115), "it is unlawful for any person to permit any noxious weed to propagate or go to seed on his land..." Fish, Wildlife & Parks (FWP) owns or leases approximately 18,000 acres in northwest Montana. Administratively, this land area is known as Region One. FWP proposes to manage noxious weeds and exotic vegetation on its properties in Region One by implementing a Noxious Weed and Exotic Vegetation Management Plan in 1995. The Plan outlines an integrated approach to weed management and would be used to guide site-specific, weed treatments beginning in the summer of 1995. The Plan would also be used to guide weed treatments under cooperative weed management projects in which Region One may participate.

Overall direction for noxious weed management is provided by FWP's recently adopted Mission statement (FWP 1992):

"to provide for the stewardship of the fish, wildlife, parks and recreational resources of Montana, while contributing to the quality of life for the present and future generations."

The primary goal of the program continues to be that stated in a general FWP weed control policy adopted in 1983:

"to prevent to the extent feasible, the reproduction and distribution of agriculturally undesirable plant species throughout department lands or from department lands onto adjacent lands."

In accordance with the Montana Environmental Policy Act (MEPA), FWP must examine and identify potential impacts of its actions on the natural and human environments. This Draft Environmental Assessment (EA) describes the direct, indirect, and cumulative effects of the proposed implementation of the Noxious Weed and Exotic Vegetation Management Plan which is based on the concept of Integrated Noxious Weed Management. In addition, the EA presents alternatives to implementation of the Plan and describes the potential direct, indirect, and cumulative effects of implementation of these alternatives.

1.2 Noxious Weeds on Region One Properties

1.2.1 The Setting and Current Conditions

Exotic (non-native) weedy plants have been noted in Montana throughout this century. In the past few decades, some species have become significant components of local vegetation communities, actually dominating some sites. FWP draws a distinction between "weedy" species and "noxious" species. FWP recognizes the value of some weed species (e.g. common dandelion) to wildlife as food and/or cover. As declared in law, "noxious" weeds are those species which are very aggressive and successful competitors for nutrients, moisture, and space and thereby encroach into a vegetation community. Important factors related to weed encroachment include proximity to water, site disturbance caused by humans or some other agent, and seed dispersal. The Draft Programmatic Environmental Impact Statement for the State of Montana's Noxious Weed Trust Fund provides a discussion of the origins of noxious weeds as well as the environmental and economic effects once they become established (Montana Department of Agriculture [MDA] 1991).

FWP Region One boundaries overlap portions of Flathead, Lake, Lincoln, and Sanders counties. Region One includes 14 Wildlife Management and Wildlife Habitat Protection Areas, 41 Fishing Access and affiliated land and administrative sites, and 6 State Parks. Two FWP Parks affiliated properties are leased to other entities who are generally responsible for site maintenance. Likewise, FWP leases acreage from other landowners and we are responsible for maintenance. Figure 1 illustrates the locations of FWP properties within the Region.

Wildlife Management Areas (WMA's) are managed to meet the needs of wildlife, preserve habitat, and provide recreational opportunities. Fishing Access Sites (FAS's) provide public access to river corridors for recreation. State Parks (SP's) preserve unique natural, historic, and recreation resources. Noxious weeds are present on all but a few sites, but the number of noxious weed species and the patch size and degree of establishment vary considerably from site to site. Some patches consist of a few plants while others are larger and more dense. Tables 1, 2, and 3 summarize the status of noxious weeds on Region One properties.

Of the 15 State-declared noxious weed species, approximately 7 occur on Region One properties. These are Spotted knapweed, Russian knapweed, Leafy spurge, Whitetop, St. Johnswort, Sulphur cinquefoil and Canada thistle. An additional 12 species are declared noxious by the counties which overlap FWP Region One boundaries. These are identified in Appendix B of the Plan. Some of these species occur on FWP properties, with mullein, hound's tongue, and musk thistle being the most frequent.

Other noxious weed species which could potentially become established on Region One properties include purple loosestrife, Eurasian watermilfoil, and sulfur cinquefoil. Purple loosestrife is an aggressive aquatic weed that rapidly overtakes desirable wetland habitats. The State as well as the counties have taken an aggressive approach to the inventory and treatment of this species. Because many of Region One lands incorporate surface water bodies, whether

lakes or streams, purple loosestrife may become a problem in the future if detected and not managed.

Eurasian watermilfoil has not been identified east of the Continental Divide, but this species is an extremely aggressive water weed found in Oregon, Washington, and Alberta. It has made in-roads into Idaho along the Columbia River system and has been observed in the Missoula area (D. Workman 1992). It invades shallow shoreline areas, especially reservoirs; however it is not clear whether this species could survive the typically severe climate of Montana east of the Continental Divide.

Sulfur cinquefoil was recently identified in Lake County, and has been observed in recent inventories of one Region One property. This species rapidly invades rangelands. It may be more prevalent than previously thought because it resembles several native cinquefoils and is difficult to identify.

1.2.2 Noxious Weed Treatment History

Past weed management efforts on Region One properties have varied in method and intensity based on the noxious weed species present and its abundance. Typically, herbicide applications, mowing, and manual techniques (pulling or clipping) have been the primary treatment methods. Biological control experiments have also been initiated, with the first releases on Kokanee Bend Fishing Access in 1985. In 1989, biocontrol agents were released on several Fishing Access Sites. New releases and re-distributions have been conducted on additional Fishing Access Sites during the last few years. FWP has practiced an informal version of Integrated Noxious Weed Management since the passage of the County Noxious Weed Control Act in 1979. Table 4 illustrates FWP Region One weed management expenditures for materials, personnel, and contracted services during fiscal years 1986 and 1992.

Table 4. Total expenditures by Montana Fish, Wildlife & Parks Region One for weed management on its Wildlife Management Areas, Fishing Access Sites and State Parks combined during fiscal years (July 1 - June 30) 1986 and 1992.

Property Type	Fiscal Year 1992 ¹	Fiscal Year 1986 ¹
Wildlife Management Areas	\$2,901	\$unknown
Fishing Access	\$ 709	\$1,168
State Parks	\$4,752	\$1,944

¹ Total includes personnel, contracted services and materials.

Properties owned by or leased to Fish, Wildlife & Parks in Region One. Key to the Fishing Access Sites appears on the facing page.

- ② Fishing Access Site
△ State Park
▣ Wildlife Management Area
◇ Wildlife Habitat Protection Area

Area Enlarged

A horizontal number line with tick marks at 0, 10, and 20. The word "Miles" is written below the line. The segment between 0 and 10 is shaded with a stippled pattern.

Key to the Fishing Access Sites

CLARK FORK RIVER DRAINAGE

8. Flatiron Ridge	30. Thompson Chain of Lakes

FLATHEAD RIVER DRAINAGE

1. Ashley Creek	7. Ducharme	19. Old Steel Bridge	26. Spring Creek
2. Ashley Lake	11. Horseshoe Lake/Ferndale	20. Pressentine	27. Swan River
3. Beaver Lake	12. Kokanee Bend	21. Skyles Lake	28. Teakettle
4. Bigfork	13. Kuhns	22. Smith Lake	31. Walstad
5. Blanchard Lake	15. Loon Lake/Ferndale	23. Somers	32. Whitefish River
6. Bootjack Lake	17. Marlowe Springs	25. Sportsmans Bridge	33. Woods Bay

KOOTENAI RIVER DRAINAGE

9. Frank Lake	14. Loon Lake/Eureka	18. Moran Lake	29. Tetrault Lake
10. Glen Lake	16. Marl Lake	24. Sophie Lake	

Table 1. Noxious weed species present and status on Fish, Wildlife & Parks (FWP) Wildlife Management Areas in Region One.

FWP PROPERTY NAME and ACREAGE	WEED SPECIES PRESENT	WEED STATUS		
		SIZE ¹	DENSITY ²	LOCATION
Kootenai/Derozier- Woods Ranch (1,357)	spotted knapweed	25	MD	roadsides, meadows in section 2, fencelines
Kootenai/Falls (344.23)	spotted knapweed	15	LDA	roadsides, hay meadows, riverbank
Kootenai/West (920.12)	spotted knapweed	< 1	VLD	Has not been detected
Mount Silcox (1,552.3)	spotted knapweed	189	LBD	mostly in foothills
	St. Johnswort	3	VLD	very spotty
Ninepipe (3,142.36)	spotted knapweed	unk	LDB	roadsides,
	whitetop	unk	MD	roadsides,
	Russian knapweed	unk	LDB	old fields
	Canada thistle	unk	LDA	roadsides
Pablo (415.92)	spotted knapweed	unk	LDB	roadsides
	Canada thistle	unk	LDB	roadsides
Ray Kuhns (1,609.3)	spotted knapweed	unk	LDA	roadsides
RoundHorn (27)	spotted knapweed	5	MD	pastureland

¹ SIZE: total acres, including roadsides and scattered patches away from roads

² DENSITY: number of weed stems in the population

<i>VLD:</i>	<i>very low density</i>	--	<i>one or a few individual plants, accounting for a very small proportion of the vegetative ground cover; native species well represented</i>
<i>LDA:</i>	<i>low density - A</i>	--	<i>individual <u>plants clumped in small areas</u>, accounting for a small proportion of the vegetative ground cover; native species well represented</i>
<i>LDB:</i>	<i>low density - B</i>	--	<i>individual <u>plants scattered throughout site</u>, accounting for a small proportion of the vegetative ground cover; native species well represented</i>
<i>MD:</i>	<i>moderate density</i>	--	<i>individual <u>plants numerous</u>, accounting for a moderate proportion of the vegetative ground cover; native species present</i>
<i>HD:</i>	<i>high density</i>	--	<i>individual <u>plants abundant</u>, accounting for a high proportion of the ground cover; native species present but not well represented</i>
<i>VHD:</i>	<i>very high density</i>	--	<i>individual <u>plants very abundant</u>, accounting for almost all of the vegetative ground cover; native species almost absent</i>

1.3 Weed Management Goals

Region One's noxious weed and exotic vegetation management goals are derived from statutory requirements, FWP policies, as well as ecological considerations. They are to:

- Meet legal requirements (including the County Noxious Weed Control Act and the Montana Environmental Policy Act) to manage plants declared as noxious weeds by the State and by the Flathead, Lake, Lincoln, and Sanders county weed boards.
- Develop and implement an integrated approach to noxious weed management and a strategy to prioritize treatment areas.
- Attempt to contain, reduce or eradicate weed populations, depending on the location and status of the population.
- Prevent the introduction and establishment of new noxious weed species, and exotic plant species in Natural Zones of State Parks, to the extent feasible.
- Work cooperatively with adjacent landowners and the County Weed Boards to enable consistent and sound weed management efforts.
- Conduct treatments in an ecologically responsible manner to maintain fish, wildlife, and vegetation resources and protect human health and safety.
- Involve the public in the planning process.

1.4 Programs and Legal Framework Affecting Region One's Weed Plan

FWP Region One's implementation of its Weed Management Plan is subject to the policies and programs of this and other agencies, as well as to state and federal laws. These are briefly discussed below.

1.4.1 Wildlife Management Area Management Plans

Site specific integrated resource management plans for the WMA's are in various stages of development and/or revision. These plans detail management objectives, current conditions, and strategies to meet desired future conditions. Noxious weed management is an element of these plans. Region One's Weed Management Plan identifies goals and treatment strategies for each WMA which are consistent with each Area Management Plan.

Table 2. Noxious weed species present and status on Fish, Wildlife & Parks (FWP) Fishing Access Sites in Region One, by drainage.

FWP PROPERTY NAME and ACREAGE	WEED SPECIES PRESENT ¹	WEED STATUS SIZE ² DENSITY ³	
CLARK FORK RIVER			
Flatiron Ridge (24.9)	SK	20	LDB
Thompson Chain of Lakes (3,965.07)	SK, CT, SJ	1500, 900, 10	MD, MD, VLD
FLATHEAD RIVER			
Ashley Creek (18.25)	unknown		
Ashley Lake (31.56)	unknown		
Beaver Lake (2)	SK	1/4	VLD
Bigfork (0.69)	NONE		
Blanchard Lake (9.31)	SK,CT	1,4	LDA,LDB
Bootjack Lake (0)	unknown		
Ducharme (51.25)	SK	1/4	LDA
Horseshoe Lake/Ferndale (22.92)	unknown		
Kokanee Bend (172.34)	SK, LS, CT	15, 3, 1	LBD, LDA, VLD
Kuhns (82.59)	unknown		
Loon Lake/Ferndale (4.36)	unknown		
Marlowe Spgs (13.53)	unknown		
Old Steel Bridge/Kiwanis (122.3)	SK, CT	25, 2	MD, LDA
Pressentine (7.02)	SK, CT	4, 1/8	LDB, VLD
Skyles Lake (0.75)	SK	1/2	LBD
Smith Lake (3.16)	SK,CT	1/2, 1 1/2	LDA,LDB
Somers (2)	SK	1/64	LDA
Sportsmans Bridge (5.77)	SK	1/16	LDA
Spring Creek (240)	unknown		
Swan River (58.52)	SK	1/16	VLD
Teakettle (2.5)	SK	1/64	MD

FWP PROPERTY NAME and ACREAGE	WEED SPECIES PRESENT ¹	WEED STATUS	
		SIZE ²	DENSITY ³
Walstad (4.57)	CT	1/64	MD
Whitefish River (3.29)	NONE		
Woods Bay (11.7)	SK	1/2	VLD
KOOTENAI RIVER			
Frank Lake (88.88)	unknown		
Glen Lake (1.4)	NONE		
Loon Lake/Eureka (0.56)	unknown		
Marl Lake (3.02)	unknown		
Moran Lake (0)	unknown		
Sophie Lake (17.82)	SK	6	LDB
Tetrault Lake (7.88)	SK	5	HD
FISH HATCHERIES and ADMINISTRATIVE SITES			
Flathead Lake Salmon Hatchery (25.47)	SK	1/2	VLD
Jocko River Trout Hatchery (59.71)	SK	6	LDA
Lake Mary Ronan Spawn Station (40.27)	unknown		
Lore Lake Spawn Station (0)	unknown		
Libby Field Station (67.57)	SK,CT	11	LDA
Lone Lake Hatchery (4.33)	unknown		
Region One Headquarters Complex (3.27)	NONE		
Rogers Lake Spawn Station (0.63)	unknown		
Rose Creek Hatchery Site (20.2)	CT	10	LDA

¹ Weed species abbreviations:

SK = spotted knapweed	LS = leafy spurge	ST = sow thistle
CT = Canada thistle	MT = musk thistle	BT = bull thistle
MU = mullein	DT = dalmatian toadflax	SJ = St. Johnswort
OW = other weed-like vegetation		

² SIZE: total acres or partial acres for each species, respectively:

1 acre = 43,560 ft ² (200' x 200')	1/16 acre = 2,722 ft ² (50' x 50')
1/2 acre = 21,780 ft ²	1/32 acre = 1,361 ft ²
1/4 acre = 10,890 ft ² (100' x 100')	1/64 acre = 680 ft ² (25' x 25')
1/8 acre = 5,445 ft ²	< 1/64 acre estimation in square feet

³ DENSITY: number of weed stems in the population, respectively:

VLD: very low density	--	one or a few <u>individual plants</u> , accounting for a very small proportion of the vegetative ground cover; native species well represented
LDA: low density - A	--	individual <u>plants clumped in small areas</u> , accounting for a small proportion of the vegetative ground cover; native species well represented
LDB: low density - B	--	individual <u>plants scattered throughout site</u> , accounting for a small proportion of the vegetative ground cover; native species well represented
MD: moderate density	--	individual <u>plants numerous</u> , accounting for a moderate proportion of the vegetative ground cover; native species present
HD: high density	--	individual <u>plants abundant</u> , accounting for a high proportion of the ground cover; native species present but not well represented
VHD: very high density	--	individual <u>plants very abundant</u> , accounting for almost all of the vegetative ground cover; native species almost absent

1.4.2 County Weed District 5-Year Plans

The County Noxious Weed Control Act provides for the creation of County Weed Districts and a County Weed Board to implement noxious weed control programs within their respective counties. The Board is responsible for preparing a Noxious Weed Management Plan which identifies noxious weeds requiring treatment in that county (including those county-declared species) and outlines procedures for control actions. FWP is obliged to consult these plans.

Also, FWP is committed to coordinating with local County Weed Boards. The kinds of activities that typically require interagency coordination include release of biocontrol agents and "War on Weeds" community projects which are implemented with funds approved through the State's Noxious Weed Trust Fund Grant Program. Furthermore, FWP has contracted with County Weed Districts over the years to conduct herbicide applications on some of its properties. Contracting will likely continue in the future, subject to a particular District having the time and personnel resources to accept such contracts.

1.4.3 Other Noxious Weed Management Plans

The Montana Department of Transportation is developing a weed plan. This plan would have bearing on Region One's program if road reconstruction activities occurred or if weed management efforts were completed on state highways that border Region One lands.

Noxious weed management on FWP properties will be affected by the weed management activities of its neighbors. Some FWP properties in Region One are bordered by the Flathead and Kootenai National Forests and land administered by the National Park Service (NPS). The Flathead National Forest completed an Environmental Impact Statement on noxious weed control on the Spotted Bear and Hungry Horse Ranger Districts in 1991. The Kootenai National Forest operates an integrated weed control effort in conjunction with the Lincoln County Weed Control Plan. They are currently working on eradication of Rush skeletonweed on 9 sites associated with roadways. Dalmation toadflax and Leafy spurge control are priorities in the Eureka area. Glacier National Park completed its Exotic Vegetation Management Plan in February, 1991.

FWP Regions 2 and 8, bordering Region One to the south and southeast respectively, have already completed their Management Plan and EA process. Region 4, to the east, has also recently developed a weed management plan.

1.4.4 Noxious Weed Trust Fund Environmental Impact Statement

The Montana Department of Agriculture completed an EIS on the State's Noxious Weed Trust Fund Grants Program. If FWP were to participate in any grant program, either as a sponsor or as a cooperator, FWP's activities would be subject to the particulars of that document which include environmental review of proposed actions by MDA and fulfilling grant application requirements.

Table 3. Noxious weed species present and status on Montana State Parks in Region One.

FWP PROPERTY NAME and ACREAGE	WEED SPECIES PRESENT ¹	WEED SPECIES SIZE ² DENSITY ³	
Flathead Lake/Big Arm (217.49)	SK, CT,SC	150, 1, 1/64	LDB, HD, VLD
Flathead Lk/Finley Point (27.83)	SK, CT	2, 1/64	LDA, MD
Flathead Lake/Wayfarers (68.86)	SK, CT	10, 1/2	LDA, MD
Flathead Lk/West Shore (129.25)	SK	16	LDB
Flathead Lk/Wild Horse Is (2,163.92)	SK, MU	10, 10	LDA, LDA
	LS,OW	1/8, 1/64	MD, MD
	CT	1/64, 1/64, 1/64	MD, MD, MD
Flathead Lk/Yellow Bay (15)	SK	1/4	VLD
Lake Mary Ronan (76.08)	SK, CT	4, 1/16	LDB, LDA
Logan (17.58)	SK, CT	2, 10	LDA, HD
Lone Pine (182.35)	SK, CT, SJ	80, 10, 40	LDA, LDA, VLD
Thompson Falls (36.4)	SK, LS	12, 1/16	LDB, MD
Whitefish Lake (10.35)	NONE		

¹ Weed species abbreviations

SK = spotted knapweed	LS = leafy spurge	CT = Canada thistle
MU = mullein	DT = dalmatian toadflax	SJ = St. Johnswort
SC = sulphur cinquefoil	OW = other weed-like vegetation	

² SIZE: total acres or partial acres for each species, respectively:

1 acre = 43,560 ft ² (200' x 200')	1/16 acre = 2,722 ft ² (50' x 50')
1/2 acre = 21,780 ft ²	1/32 acre = 1,361 ft ²
1/4 acre = 10,890 ft ² (100' x 100')	1/64 acre = 680 ft ² (25' x 25')
1/8 acre = 5,445 ft ²	< 1/64 acre estimation in square feet

³ DENSITY: number of weed stems in the population

VLD: very low density	--	<i>one or a few <u>individual plants</u>, accounting for a very small proportion of the vegetative ground cover; native species well represented</i>
LDA: low density - A	--	<i>individual <u>plants clumped in small areas</u>, accounting for a small proportion of the vegetative ground cover; native species well represented</i>
LDB: low density - B	--	<i>individual <u>plants scattered throughout site</u>, accounting for a small proportion of the vegetative ground cover; native species well represented</i>
MD: moderate density	--	<i>individual <u>plants numerous</u>, accounting for a moderate proportion of the vegetative ground cover; native species present</i>
HD: high density	--	<i>individual <u>plants abundant</u>, accounting for a high proportion of the ground cover; native species present but not well represented</i>
VHD: very high density	--	<i>individual <u>plants very abundant</u>, accounting for almost all of the vegetative ground cover; native species almost absent</i>

1.4.5 Other Applicable Laws and Regulations

- Federal Insecticide, Fungicide, Rodenticide Act (P.L. 92-516, FIFRA) and Code of Federal Regulations 40 CFR (Part 171)
- Montana Environmental Policy Act (MCA 75-1-101 *et seq.*) and rules (ARM 12.2.401 *et seq.*)
- Montana Weed Control Act (MCA 80-7-801 *et seq.*)
- Montana Water Quality Act (MCA 75-5-101 *et seq.*)
- Montana Agricultural Chemical Groundwater Protection Act (MCA 80-15-100 *et seq.*)
- Montana Pesticides Act (MCA 80-8-801 *et seq.*)

1.5 Decisions to be Made

In accordance with MEPA, this Draft Environmental Assessment identifies potential impacts on the natural and human environment that may result from the proposed action, implementation of the Weed Management Plan (based on the concepts of Integrated Noxious Weed Management) or from the implementation of one of the reasonable Alternatives. It should provide the Region 1 Supervisor (decision maker) with adequate information to determine whether or not significant impacts would occur if the proposed action were to be implemented.

The Regional Supervisor must decide whether to continue to manage noxious weeds in the same manner and to the same extent as in the past (Status Quo Alternative), to accept one of the other three Alternatives presented in this document, or to accept one of the Alternatives with changes such as mitigation measures.

Once an Alternative is selected, the Regional Supervisor will decide whether or not the proposed action will have significant adverse impacts on the human and natural environments. If no significant adverse impacts would result, then the final EA will state the reasons for this finding and that the EA is the appropriate level of analysis. The Plan, revised to incorporate findings of the EA, would be implemented. If impacts are judged to be significant, an EIS may be completed to further investigate the proposed action.

1.6 Public Concerns and Issues

The following issues have been distilled from comment opportunities during various FWP public meetings, as well as scoping sessions related to FWP noxious weed management plans in other regions:

- What weed treatment methods does FWP propose to use? How will these methods affect the natural and human environment?
- Herbicide use can have negative impacts if handled improperly or used in sensitive areas. What are the potential impacts to resources such as air, groundwater, surface water, fisheries, wildlife, wildlife habitat, and native plant communities. What are the potential impacts to recreationists and FWP employees?
- Certain weed species have become established on Region One lands, while others have not. How will FWP Region One prioritize treatment activities?
- The noxious weed problem does not stop at FWP property boundaries. How will FWP address weed problems on adjacent lands? How will FWP work with other interested parties such as County Weed Boards, state and federal agencies, adjacent landowners, lessees, and volunteer groups?
- Poor land management practices such as those which extensively disturb the ground and soil surface facilitate noxious weed establishment. How will the FWP monitor and modify these activities to curtail the spread of noxious weeds on its properties?
- Public education and awareness about noxious weeds and their management are very important. What role will FWP have in public awareness? Also, what involvement will the public have in management decisions?

These issues are addressed herein, and the draft plan and environmental assessment will be publicized and made available for a 30-day public review and comment period. The documents will be available at FWP headquarters and regional offices, as well as at county libraries and weed district offices in the affected counties. If there is a significant amount of concern or opposition to the proposed alternative during the comment period, a public meeting will be held to hear and address any concerns. Following the review period, a Decision Notice will be issued incorporating comments and revisions.

2.0 TREATMENT METHODS AND ALTERNATIVES

2.1 Treatment Methods

The noxious weed management goals of FWP are to attempt to prevent the introductions of new noxious weed species and to discourage the spread of established species on FWP properties. In discussing how to meet these goals, FWP distinguishes between chemical and non-chemical treatment methods. These categories are outlined below. Various combinations of these methods are incorporated into the Alternatives which will be discussed subsequently.

2.1.1 Chemical Methods

Chemical treatment methods involve the use of herbicides to kill weeds directly or the application of soil sterilants which prevent the germination of noxious weed seeds. Over the past half-century, chemical herbicides have been developed to treat a variety of weed species. While the benefits of chemical methods include efficacy and low cost, there are valid concerns about any application of chemicals to the landscape.

The herbicides 2,4-D, picloram, glyphosate, and clopyralid are proposed for use on Region One lands. These herbicides may be used alone or in combination as recommended by label directions. Table 5 lists the chemicals and formulations proposed for use. Appendix A summarizes herbicide properties and lists the target noxious weed species and the efficacy of herbicide treatments. Other herbicides, which have been approved by the U.S. Environmental Protection Agency (EPA) and licensed for use in Montana, may be considered in the future as they become available.

Table 5. Herbicides proposed for use on Fish, Wildlife & Parks Region One properties, registered trade name, and application rates on a per acre basis according to manufacturer label recommendations.

Common Chemical Name	Registered Trade Name	Application Rates
2,4-D amine	Weedar 64, various	1 - 3 quarts
picloram	Tordon 22K	1 pint - 2 quarts
glyphosate	Ruler	16 ounces - 4 quarts
clopyralid	Stinger	1/4 - 1 1/3 pints
clopyralid	Transline	1/4 - 1 1/3 pints
clopyralid + 2,4-D	Curtail	2 - 2 2/3 pints
2,4-D ester	Hi-Dep	1 - 3 quarts

The use of herbicides would depend on the treatment objective, season of application, stage of plant growth, topography, species present, effectiveness of treatment, equipment limitations, and potential environmental impacts. Herbicide and application rate selection would depend on the target species, other vegetation present, presence of sensitive plant or animal species, soil type, presence of surface waters, wetlands, shallow groundwater or groundwater recharge areas, and proximity to human habitation or use sites. Appendix H in the Plan contains label instructions of each herbicide proposed for use.

Herbicides would always be applied according to manufacturer label directions, regardless of what Alternative is selected. Changes in application rates, recommended by research institutions, would be considered if these rates do not exceed label directions. Chemical applications would be supervised or conducted by licensed applicators and in compliance with Federal requirements.

Aerial applications of herbicide are not proposed under any circumstances. Vehicle-mounted sprayer (hand gun or boom) applications will be used primarily in open areas that are readily accessible by vehicle. Boom sprayers, which are fixed, provide less flexibility and would be used only where feasible to treat concentrated weed patches. The hand gun would be used for spot treatment of weeds. Spot treatments would be selected over broadcast methods. Regardless of spray method used, sprays would be applied in a manner that gives the best possible coverage with the least amount of drift. Hand applications would utilize backpack spraying and wiper application. Backpack sprayers would be used on small or scattered patches in rough terrain or environmentally sensitive areas. Contact systemic herbicides such as glyphosate would be used to treat individual plants or for seed bed preparation.

Precautions for use would include:

- During application periods, weather conditions (including forecasted precipitation events) and temperatures would be monitored by contractors and weed management personnel.
- Calibration checks would be conducted, when feasible, to ensure that equipment is functioning correctly.
- Label requirements would be followed for all herbicide applications; further precautions may be determined to be necessary during the pre-treatment review process.

2.1.2 Non-Chemical Methods

Manual methods involve grubbing or topping with hand tools and hand-pulling. These methods may effectively eradicate annual or biennial weed species in small or sparse populations, primarily by preventing seed production and distribution as long as treatment occurred prior to seed ripe. Annual re-treatment is required. Because these methods are labor intensive, their application would be limited primarily to environmentally or culturally sensitive areas and very small populations.

Cultural methods include mowing, tilling, reseeding, grazing, and other agricultural techniques. These methods may effectively reduce some noxious weed species, although implementation is often limited by topography and access. These methods can be costly and time consuming. Furthermore, disruption of the soil surface may create new microsites vulnerable to noxious weed encroachment. Burning, as a weed treatment method by itself, is generally of limited effectiveness.

Biological methods describe the introductions of host-specific natural enemies (insects or pathogens) on individual plants within a noxious weed population. Biological control agents can effectively stress noxious weeds and reduce seed production and viability. Use of biocontrol agents would not eliminate the species but can reduce the population to a tolerable level, particularly when used in combination with other treatment methods. Biological control may be the only method possible in some areas. These areas may include those that are inaccessible to mowers or where other treatment methods are not possible or cost effective to implement. Initial acquisition costs of biocontrol agents are high. However, as populations establish themselves on a site, insects can be collected and released to other sites.

2.2 Description and Comparison of the Alternatives

Four Alternatives for noxious weed management were considered. They are: Status Quo, Integrated Noxious Weed Management, Chemical Methods, and Non-Chemical Methods. The Montana Environmental Policy Act requires the consideration of a "No Action" alternative in which the proposed action (management of noxious weeds in this case) does not take place. However, under the County Noxious Weed Control Act, FWP is legally required to manage noxious weeds on its properties, as any other landowner in Montana. Thus the "No Action" alternative is not meaningful in the context of noxious weeds and will not be considered per se. Instead, FWP will consider the implementation of a "Status Quo" Alternative in which there is no change in the current management program. Four Alternatives for noxious weed management are described below. Each Alternative incorporates some or all of the treatment methods available: manual, cultural, biological, and chemical. However, each Alternative, in emphasizing a different method (s) would have different potential impacts to air quality, soils, water quality, wildlife/fisheries, vegetation, recreation, human health, and cultural/historic resources. Chapter 3.0 discloses these impacts.

2.2.1 Alternative 1 (Status Quo)

Under Alternative 1 (Status Quo), noxious weed management would continue along the course set over the past several years. Previous weed treatment activities in Region One included an emphasis on chemical methods, with some use of manual, cultural, and biological methods. Under this Alternative, no particular effort to integrate noxious weed management with other resource management programs would be made, nor would we plan a coordinated effort to manage noxious weeds. Field personnel would simply react to weed outbreaks as they occurred. Under this Alternative, chemicals methods would still be applied according to manufacturer label instructions, but the additional safeguards outlined in the Noxious Weed Management Plan would not be implemented. Biological control agents already introduced would run their courses, but insect populations would not be supplemented or redistributed to any great degree. Periodic consultation with County Weed Boards would occur but active participation in new, coordinated treatment initiatives or public awareness programs would not.

If this Alternative is selected, then noxious weeds would likely spread on and from Region One lands because this Alternative does not describe a proactive management strategy. Weed-free areas would likely be encroached upon as noxious weed seed reserves build up and species spread to ecological tolerance. This Alternative would not adequately protect state resources nor would FWP management goals and stewardship responsibilities be met. The potential loss of wildlife forage, the change in vegetation communities, and decreases of aesthetic quality would be significant.

2.2.2 Alternative 2 (Integrated Noxious Weed Management)

This Alternative calls for a program which integrates preventive measures with biological, manual, cultural, and chemical treatment methods in a proactive fashion. The Noxious Weed Management Plan details this approach, defined as Integrated Noxious Weed Management. Its goal is the long-term prevention and reduction of noxious weeds from FWP properties, whether for established populations or new ones. This approach permits consideration of and planning for other resource management objectives.

Alternative 2 provides the best opportunity for long-term weed management with minimal impact to the environment. The Plan promotes flexibility during the treatment design and planning processes. It allows a variety of treatment methods to be used as site-specific conditions warrant and incorporates many additional safeguards to ensure that other resources are not adversely affected by noxious weed treatment. Herbicide use would still present some risks, but they would be considerably less than those expected under other Alternatives. If this Alternative is selected, prospects for long-term prevention and reduction of noxious weeds on FWP properties are good -- while at the same time preserving other resources.

2.2.3 Alternative 3 (Chemical Methods)

Chemical herbicides would be the primary treatment method except where there is a high risk of environmental contamination or label directions preclude the use of herbicides. Biological control agents (insects and pathogens) already introduced would be permitted to run their course, but new introductions or re-distributions would not take place since herbicides may limit the viability and success of biocontrol agents. Manual or cultural treatment methods would not be considered. Measures to help prevent noxious weed encroachment could still be implemented.

Under this Alternative, the risk of environmental contamination through chemical over-application or accidental spill is highest, as is the chance of adversely affecting non-target plant or animal species. This would not be due to willful misuse, but rather because the total amount of chemicals used would be significantly higher under this Alternative than any other. Cumulative effects of chemical applications may occur if adjacent landowners also apply herbicides regularly. Ground- and surface water and the associated resources would be at greatest risk. Concerns for human health are the greatest under this Alternative.

If this Alternative is selected, prospects for long-term reduction are good because chemicals are generally effective. However chemical applications would be precluded in some settings because of manufacturer label restrictions (e.g. riparian zones). Thus noxious weeds would persist in such settings. In general, the integrity of other resources can not be guaranteed under this Alternative since the primary objective is the reduction of noxious weeds through chemical methods.

2.2.4 Alternative 4 (Non-Chemical Methods)

Weed management emphasis would be toward containment of current weed populations. This Alternative emphasizes manual, cultural, and biological control methods. Herbicides would not be used, nor would FWP Region One participate in cooperative programs which use or emphasize herbicides. Biological control methods would be accentuated and new pathogens would be introduced as they become available. Preventive methods that would not require major new funding by Region One could be implemented.

If this Alternative is selected, prospects for long-term success would be good only in areas where weeds patches are relatively small or well-confined and the rate of weed spread is relatively slow. Biological control treatments would have some impact on rate of spread but many researchers believe that more organisms (insects and pathogens) need to be developed and introduced to make biological control an effective tool (Story 1989). Also, several Region One properties are too small to sustain biocontrol organisms. In areas where populations are large or weed plants are well established with a seed source, prospects for long-term success under this Alternative are poor. The rate of spread will likely out-pace treatment activities. Coordination with adjacent landowners would be essential for this Alternative to be even marginally successful.

2.3 The Preferred Alternative

The preferred Alternative, (Integrated Noxious Weed Management), is briefly described below. The reader is referred to the Noxious Weed and Exotic Vegetation Management Plan for a more detailed discussion. Some elements of Alternative 2 are also incorporated under Alternatives 1, 3, and 4.

FWP's strategy is to employ the concept of Integrated Noxious Weed Management, as defined in the Plan. This concept is a systems approach to reduce pest damage to tolerable levels. This strategy seeks to tailor weed management activities to the treatment needs and site characteristics of a particular weed patch. First, weed management and treatment priorities are established. Then, treatment methods and techniques are selected, including natural predators and parasites, environmental modifications, and herbicides when necessary and appropriate. This approach allows the most flexibility for managing noxious weeds as site conditions change, treatment techniques improve, and funding opportunities arise. This approach is consistent with state law and the noxious weed plans developed by the counties.

2.4 Cumulative Effects and Irreversible or Irretrievable Commitments

In order to evaluate potential cumulative impacts to the environment, similar actions of adjacent landowners, federal, state, and county weed control boards will be considered. Of primary concern are the potential cumulative effects on ground- and surface waters resulting from herbicide application. Without coordinated efforts (communication, planning, and monitoring), contamination is possible.

The irreversible commitment of resources refers to the consumption or loss of resources that cannot be reversed, except possibly in the extreme long term. In the broadest sense, none of these Alternatives will result in the irreversible commitment of resources. However, some rare plant populations could be lost if proper mitigation measures are not followed, as would be the case under Alternatives 1, 3, and 4. Alternative 2 (Integrated Noxious Weed Management) is the only Alternative requiring such safeguards as sensitive plant inventory prior to chemical or cultural treatment activities.

The irretrievable commitment of resources refers to the consumption or loss of resources for a short period of time such as 50 to 100 years. In the absence of all noxious weed management activities, wildlife cover and forage, recreation values, and aesthetic values could be irretrievably committed. Continuation of the current management direction (Alternative 1, Status Quo) could result in the continued spread of noxious weeds and further alteration of resources that FWP is charged to maintain or enhance. Under Alternatives 1, 3, and 4, some rare plant populations could be impacted by treatment activities if proper mitigation measures are not followed. Only Alternative 2 (Integrated Noxious Weed Management) includes safeguards and guidelines to prevent the irretrievable commitment of resources or loss of rare plant populations.

3.0 THE AFFECTED ENVIRONMENT AND ANALYSIS OF POTENTIAL IMPACTS

This chapter provides the basis for comparing the Alternatives presented in Chapter 2. Based on the issues identified in Section 1.6, this section of the EA describes the natural resources that may be affected by implementation of the preferred Alternative or one of the other Alternatives. Potential impacts are addressed in two ways and analyzed in relation to the treatment methods proposed under each Alternative. First they are discussed generally for all Region One properties and for each Alternative because most impacts could occur on any site. Secondly, site specific impacts are discussed where specific resources may be negatively affected or where site conditions or characteristics may affect program implementation. The impacts of allowing noxious weeds to spread are also addressed. Table 6 summarizes the environmental impacts resulting from the selection of the preferred Alternative, or implementation of the Plan which emphasizes the concept of Integrated Noxious Weed Management.

Table 6. Summary of potential environmental impacts resulting from the implementation of the Weed Management Plan, emphasizing the concept of Integrated Noxious Weed Management.

	Unknown	None	Minor	Potentially Significant	Can Impact Be Mitigated
1. LAND RESOURCES Will the proposed action result in:					
a. Soil instability or changes in geologic substructures?			X		yes ¹
b. Disruption, displacement, erosion, compaction, moisture loss, or over-covering of soil which would reduce productivity or fertility?			X		yes ¹
c. Destruction, covering or modification of any unique geologic or physical features?		X			
d. Changes in siltation, deposition or erosion patterns that may modify the channel of a river or stream or the bed or shore of a lake?		X			
e. Exposure of people or property to earthquakes, landslides, ground failure, or other natural hazards?		X			
2. AIR Will the proposed action result in:					
a. Emission of air pollutants or deterioration of ambient air quality? (also see #13 (c)).			X		yes ¹
b. Creation of objectionable odors?			X		yes ¹
c. Alteration of air movement, moisture, or temperature patterns or any change in climate, either locally or regionally?		X			
d. Adverse effects on vegetation, including crops, due to increased emissions of pollutants?		X			

	Unknown	None	Minor	Potentially Significant	Can Impact Be Mitigated
3. WATER Will the proposed action result in:					
a. Discharge into surface water or any alteration of surface water quality, including but not limited to temperature, dissolved oxygen or turbidity?			X		yes ¹
b. Changes in drainage patterns or the rate and amount of surface runoff?			X		yes ¹
c. Alteration of the course or magnitude of flood water or other flows?		X			
d. Changes in the amount of surface water in any water body or creation of a new water body?		X			
e. Exposure of people or property to water related hazards such as flooding?		X			
f. Changes in the quality of ground water?			X		yes ¹
g. Changes in the quantity of ground water?		X			
h. Increase in risk of contamination of surface or ground water?			X		yes ¹
i. Effects on any existing water right or reservation?		X			
j. Effects on other water users as a result of any alteration in surface or ground water quality?			X		yes ¹
k. Effects on other water users as a result of any alteration in surface or ground water quantity?		X			
4. VEGETATION Will the proposed action result in:					
a. Changes in the diversity, productivity or abundance of plant species (including trees, shrubs, grass, crops, and aquatic plants)?			X		yes ¹
b. Alteration of a plant community?			X		yes ¹
c. Adverse effects on any unique, rare, threatened, or endangered species?			X		yes ¹
d. Reduction in acreage or productivity of any agricultural land?		X			
e. Establishment or spread of noxious weeds?		X			
5. FISH/WILDLIFE Will the proposed action result in:					
a. Deterioration of critical fish or wildlife habitat?		X			
b. Changes in the diversity or abundance of game animals or bird species?		X			
c. Changes in the diversity or abundance of nongame species?		X			
d. Introduction of new species into an area?		X			
e. Creation of a barrier to the migration or movement of animals?		X			
f. Adverse effects on any unique, rare, threatened, or endangered species?			X		yes ¹
g. Increase in conditions that stress wildlife populations or limit abundance (including harassment, legal or illegal harvest or other human activity)?		X			

	Unknown	None	Minor	Potentially Significant	Can Impact Be Mitigated
6. NOISE/ELECTRICAL EFFECT Will the proposed action result in:					
a. Increases in existing noise levels?		X			
b. Exposure of people to severe or nuisance noise levels?		X			
c. Creation of electrostatic or electromagnetic effects that could be detrimental to human health or property?		X			
d. Interference with radio or television reception and operation?		X			
7. LAND USE Will the proposed action result in:					
a. Alteration of or interference with the productivity or profitability of the existing land use of an area?		X			
b. Conflict with a designated natural area or area of unusual scientific or educational importance?		X			
c. Conflict with any existing land use whose presence would constrain or potentially prohibit the proposed action?		X			
d. Adverse effects on or relocation of residences?		X			
8. RISK/HEALTH HAZARDS Will the proposed action involve:					
a. Risk of an explosion or release of hazardous substances (including, but not limited to, oil, pesticides, chemicals, or radiation) in the event of an accident or other forms of disruption?			X		yes ¹
b. Affect an existing emergency response or emergency evacuation plan or create a need for a new plan?		X			
c. Creation of any human health hazard or potential hazard?			X		yes ¹
9. COMMUNITY IMPACTS Will the proposed action result in:					
a. Alteration of the location, distribution, density, or growth rate of the human population of an area?		X			
b. Alteration of the social structure of a community?		X			
c. Alteration of the level or distribution of employment or community or personal income?		X			
d. Changes in industrial or commercial activity?		X			
e. Changes in cultural diversity or uniqueness?		X			
f. Increased traffic hazards or effects on existing transportation facilities or patterns of movement of people and goods?		X			

	Unknown	None	Minor	Potentially Significant	Can Impact Be Mitigated
10. PUBLIC SERVICES/TAXES/UTILITIES Will the proposed action:					
a. Have an effect upon or result in a need for new or altered governmental services in any of the following areas: fire or police protection, schools, parks/recreational facilities, roads or other public facility maintenance, water supply, sewer or septic systems, solid waste disposal, health, or other governmental services? If any, specify:_____		X			
b. Have an effect upon the local or state tax base and revenues?		X			
c. Result in a need for new facilities or substantial alterations of any of the following utilities: Electric power, natural gas, other fuel supply or distribution systems, or communications?		X			
d. Result in increased use of any energy source?		X			
11. AESTHETICS/RECREATION Will the proposed action result in:					
a. Alteration of any scenic vista or creation of an aesthetically offensive site or effect that is open to public view?			X		yes ¹
b. Alteration of the aesthetic character of a community or neighborhood?		X			
c. Alteration of the quality or quantity of recreational opportunities and settings?			X		yes ¹
12. CULTURAL/HISTORIC RESOURCES Will the proposed action result in:					
a. Destruction or alteration of any site, structure or object of prehistoric, historic, or paleontological importance?			X		yes ¹
b. Physical change that would affect unique cultural values?			X		yes ¹
c. Effects on existing religious or sacred uses of a site or area?			X		yes ¹
13. SUMMARY EVALUATION OF SIGNIFICANCE					
Does the proposed action, considered as a whole:					
a. Have impacts that are individually limited, but cumulatively considerable? (A project may result in impacts on two or more separate resources which create a significant effect when considered together or in total.)		X			
b. Involve potential risks or adverse effects which are uncertain but extremely hazardous if they were to occur?		X			
c. Potentially conflict with the substantive requirements of any local, state, or federal law, regulation, standard or formal plan?		X			
d. Establish a precedent or likelihood that future actions with significant environmental impacts will be proposed?		X			
e. Generate substantial debate or controversy about the nature of the impacts that would be created?		X			

¹ Mitigative measures outlined in section 3.10 of the EA and section 2.0 of the Management Plan.

3.1 Air Quality

3.1.1 Current Conditions

Air quality in Montana is regulated through the Federal Clean Air Act and the Montana Clean Air Act. These laws establish ambient air quality and emission standards. Herbicides themselves are not regulated under the Federal Clean Air Act. Air quality within Region One is generally very good. Like other intermountain valleys in Montana, the Clark Fork, Flathead, Mission, Swan and Tobacco valleys experience frequent inversions. Likewise, these valleys can at times experience high particulate counts due to dust, vehicle emissions and/or wood smoke. Also, localized high winds, particularly in the Bad Rock Canyon area east of Columbia Falls, may create localized high particulate counts.

3.1.2 Analysis of Potential Impacts

General

Chemical treatment methods have the potential to affect air quality when chemical particles are suspended in the atmosphere. Impact severity depends on the type of chemical, the chemical concentration, and the duration of suspension. The drift of airborne spray particles may adversely affect air quality depending on the size of the particle and wind conditions. These effects are expected to be short-term.

Cultural treatment methods could temporarily increase dust levels if large areas were mowed, tilled, or reseeded. Biological and manual methods would not affect air quality. Overall, air quality would not be significantly affected by implementation of any of the Alternatives. Under Alternative 1 (Status Quo), Alternative 2 (Integrated Noxious Weed Management) and Alternative 3 (Chemical Methods), there would be short-term, minor effects associated with chemical odors in the immediate vicinity of a spray site. Chemical odors, resulting from a petroleum-based carrier medium in the herbicide, are released during the application process and could be perceived as unpleasant or offensive. The amount of chemical in the air would be elevated for only a short time. Additionally, some spray particles would drift away from the application location. Only Alternative 2 (Integrated Noxious Weed Management) minimizes these short-term impacts since guidelines for chemical applications (beyond those included on manufacturer labels) are included in the Plan. The Plan specifies that applications would be conducted when weather conditions (wind speed and turbulence) are suitable, nozzles on spray equipment would be adjusted to prevent aerosol dispersion, and herbicide formulations with relatively low volatility would be used. Alternatives 2 (Integrated Noxious Weed Management) and 4 (Non-Chemical) may increase dust levels as a result of cultivation or other agricultural practices which disrupt soils.

Site Specific Impacts

Region One properties that may be more significantly affected by changes in air quality would include those sites used frequently by the public or FWP employees. Examples include Flathead Lake (one of the most frequently visited State Parks in the system), Lone Pine State Park, and any of the Fishing Access Sites within a short drive of a major population center. Determining when low public use periods occur (for example, time of day or day of the week) at a particular site and restricting spray activities to those periods would reduce the potential for impact. Private property owners with residences near FWP property boundaries may also be affected by chemical treatments.

3.2 Geology and Soils

3.2.1 Current Conditions

Northwest Montana has a diverse geologic history. The landscape has been shaped by glacial, tectonic, erosional, and depositional forces. As such, the geology and soils of the region are highly variable. There are at least 16 different soil types in the area, derived from igneous, sedimentary, and metamorphic parent material (Montagne et al. 1982). Soil profile characteristics vary widely. Soils may be coarse or fine textured, falling into various classifications of sandy, silty, or clayey.

Some FWP sites have soils that are wet much of the year or have a water table within five feet of the surface. They are associated with streams, lakes, and depressions where groundwater is near the surface, where 'perched' water tables occur in clayey soils, or where irrigation practices maintain an elevated water table.

3.2.2 Analysis of Potential Impacts

General

Soils could be adversely affected under all Alternatives. The nature and extent of these impacts vary for each Alternative, depending on the method of noxious weed treatment emphasized. Soil properties themselves may dictate the feasibility of certain treatment methods.

Soil loss and reduced site productivity are the primary concerns with non-chemical treatment methods. Cultural treatments such as mowing and livestock grazing may involve a certain level of ground disturbance. This could reduce vegetative cover and expose the soil surface, making it susceptible to wind and water erosion. These activities could also cause soil compaction which would increase the potential for surface run-off as well as reduce the infiltration of water necessary for revegetation. Tilling and reseedling may leave soils temporarily exposed and more susceptible to erosion by wind and water or to subsequent noxious weed encroachment. Manual treatment methods could also reduce vegetative cover, with similar results. Biological methods are not expected to affect soils (BLM 1985).

Potential impacts to soils from chemical application include the accumulation of chemical residues (loading), the alteration of soil flora and fauna, and an increase in salt content. Soil contamination through loading is the primary concern with chemical treatment methods. The magnitude of impacts depends on several properties related to both the herbicide and the soil environment. Appendix A summarizes pertinent herbicide properties in soils.

Little is known of the consequences of chemical loading in Montana soils, although it's most likely to occur when persistent herbicides are used repeatedly (Sparks 1989). The breakdown and movement of herbicides in soil is influenced by chemical, physical, and microbial processes. By remaining in the soil environment, persistent herbicides can provide residual weed treatment for up to five years depending on the soil conditions and the herbicide. During that time there is the potential for herbicides to be transported within and between soil layers. Watson et al. (1989) documented movement of picloram to increased soil depth as application rate increased. Furthermore, 365 days after the application, the chemical was still detectable.

Herbicides which target broadleaf plants tend to give a competitive advantage to grass species. However, if grass populations are slow to recover, the soil surface may be exposed, thus increasing the erosion potential. The litter layer created by dead plant material may offset increased erosion risks by mitigating raindrop impact and promoting infiltration.

No evidence was found showing that herbicide use adversely affects soil organisms over the long-term (Audus 1976).

Under Alternative 1 (Status Quo), chemical applications would continue at their current intensity levels. Potential impacts would be those described above. The impacts may be more pronounced under this Alternative because chemical applications would proceed according to manufacturer label instructions without any additional safeguards, as would be the case under Alternative 2 (Integrated Noxious Weed Management). Noxious weeds would likely continue to proliferate and spread. Furthermore, if weeds become well established, a monoculture may result if weeds out-compete all other species. This, in turn, would increase the exposure of soil surfaces and cause an increase in soil erosion. (Rice et al. 1992).

Under Alternative 2 (Integrated Noxious Weed Management), the risk of soil contamination due to over-application of herbicides would be greatly reduced because other treatment methods would also be used. In addition, the concept of Integrated Noxious Weed Management tailors a treatment strategy to a particular site and its noxious weed status; thus, chemicals would not be applied in marginal settings where label instructions permit use but other resources values dictate otherwise. Under this Alternative, all chemical treatments would take place under the guidelines presented in the Plan. Impacts from manual and cultural methods would be related to alteration of vegetative cover and exposure of the soil surface, as described above.

Alternative 3 (Chemical Methods) entails the management of noxious weeds through the exclusive use of chemicals. The greater use of chemicals, especially those that are relatively persistent (e.g. picloram), could lead to chemical loading and thus soil contamination. In

addition, since chemicals would be the only method of treatment, there would be a greater risk of a chemical spill with the increased handling and transport under this Alternative. If non-target species are affected to a significant degree, the overall reduction in vegetative ground cover may increase exposure of the soil surface with the aforementioned potential outcomes.

Alternative 4 (Non-Chemical Methods) would eliminate the potential for herbicide impacts to soils. However, as discussed above, manual and cultural treatments may cause erosion and productivity losses.

Site Specific

Increased erosion could occur on any FWP site. Likewise chemical applications could cause the impacts described above on any site. Impacts to soil resources would be most pronounced on those sites where soils are fragile and/or not well-developed. All of the Alternatives could lead to erosion or soil contamination. However, each Region One site would be susceptible to erosion if noxious weeds are allowed to proliferate.

3.3 Groundwater

3.3.1 Current Conditions

Groundwater is an important resource. In addition to agriculture and commercial industries, it is used for municipal and rural residential purposes. Across Montana, 96% of the public water supplies are derived from groundwater (Montana Department of Health and Environmental Sciences [DHES] 1992). Groundwater is also important because of its interaction with surface waters through recharge areas. In fact, most perennial streams in Montana are recharged by groundwater. Thus, maintaining groundwater purity is important for many reasons. With localized exceptions, groundwater quality in Region One is good. Ten State Park wells and two Fishing Access wells in Region One provide visitors with drinking water where the source is groundwater.

3.3.2 Analysis of Potential Impacts

General

Herbicides can impact groundwater through point source discharges (e.g. accidental spills, uncontrolled releases during mixing, or releases at storage facilities). These impacts are often more severe than those of non-point source discharges. Deluca et al. (1989) documented contamination of domestic water supplies from point source discharges.

Non-point source releases of herbicides also have the potential to contaminate groundwater. Seven pesticides have been detected in groundwater studies in Montana. Of the seven, only 2,4-D and picloram are proposed for use on FWP Region One properties. Detection levels for all

seven pesticides were below established health protection levels (DHES 1992). Leaching, one type of non-point source release, may result from either the one-time application of a chemical unsuitable for the particular soil type or location or from the repeated application of chemicals to specific areas (loading). Several factors influence this occurrence. These include the herbicide's behavior in soil (i.e. persistence, mobility, adsorption, and microbial degradation), soil properties (texture, permeability, organic matter content, and depth), water solubility, depth to groundwater, and precipitation. Appendix A summarizes general herbicide properties for the proposed active ingredients. A brief summary of research findings to date for herbicides proposed for use on Region One properties follows.

The half-life of an herbicide is the time required for half the amount of the active chemical ingredients to become inert through degradation by natural processes. Half-life depends on application rate and the environment to which the chemical is introduced. According to the pesticides background statements put out by the U.S. Department of Agriculture [USDA] (1984), picloram can have a half-life of more than four years in arid regions and approximately one month under conditions of moist soils with a high organic matter content. This relatively long persistence in soil and a poor affinity to soil (Watson et al. 1989), could result in leaching if picloram is applied to shallow groundwater areas (within 10 feet of the surface) and/or highly permeable soils. Studies of various soil types indicate that picloram is usually confined to the upper 1 foot (30 cm) when application rates are low (less than 1 lb/acre) (USDA 1984, Rice et al. 1992). However, at rates of 3 to 9 lb/acre, picloram can readily move to depths greater than 3 feet (approximately 1 meter), even in relatively arid regions. The Plan recommends the Tordon 22K[®] formulation of picloram, which suggests rates of 1 lb/acre or less for broadleaf weeds on non-cropland, range, and pasture. Use would be precluded on lands with water tables that reach to within 5 feet of the soil surface.

Lateral movement of picloram has been reported in soils (National Research Council of Canada 1977). Also of concern are soils which are within 20 inches of bedrock. Soils "saturated" with herbicide could potentially leach through to bedrock, migrate along the bedrock surface, and arise downslope in off-target areas. Also, bedrock fissures could become conduits moving herbicides to groundwater. The extent to which this could occur is unknown.

Clopyralid persists for one to three months. Rice et al. (1992) found that clopyralid applied at rates of <1 lb/acre in bunchgrass communities did not persist for more than one year nor was the herbicide detectable below 25 cm. Because its adsorption to soil is low, leaching potential and mobility of clopyralid is high. The primary method of degradation is microbial activity in the soil; photo degradation is insignificant (Colby et al. 1989).

With a persistence of one to four weeks, 2,4-D is rapidly degraded by soil microbial activity. Photo degradation is minor. Since 2,4-D adsorption to soil is only moderate, it has a relatively high mobility and a moderate leaching potential (BLM 1985).

Glyphosate persists for up to one month. Glyphosate is readily adsorbed by soils, but microbial degradation of the chemical is slow. Leaching potential is low, but the potential for surface run-off is high (Colby et al. 1989).

Under Alternative 1 (Status Quo), herbicide use would continue at current levels of intensity. There would be a moderate risk for herbicide contamination of groundwater because herbicides with moderate to high leaching potential (2,4-D, clopyralid, and picloram) would still be used by FWP. The additional safeguards proposed in the Plan, however, would not be implemented. Biological control agents already introduced would not affect groundwater sources.

Under Alternative 2 (Integrated Noxious Weed Management), there is the potential to contaminate groundwater because herbicides with moderate to high leaching ability (picloram, 2,4-D, and clopyralid) would be used. However, the chemical applications would be conducted as specified by guidelines outlined in the Plan. The non-chemical treatment methods (cultural, biological, and manual) incorporated into this Alternative would not impact groundwater.

Alternative 3 (Chemical Methods) emphasizes herbicide use. Because this Alternative would increase the total volume of herbicides released into the environment, it presents the greatest risk to groundwater resources. Herbicide use would be outright prohibited in a number of settings because of the extent of sensitive groundwater areas, resulting in the proliferation of noxious weeds at these sites.

Alternative 4 (Non-Chemical Methods) would not affect groundwater resources associated with Region One properties.

Site Specific

To assess potential risks to groundwater on specific sites from chemicals, a procedure originally developed for the Lewis & Clark County Conservation District could be used in Region One (Lacey 1991). Risk of chemical contamination is determined for a site based on depth to seasonal or permanent water table, soil permeability, organic matter content, flooding hazard, and percent slope. A risk index based on these variables would be calculated on a scale between 1 and 100 and divided into four categories or classes of risk. Risk factors and classes would then be assigned to a soil map unit prepared by the Soil Conservation Service. With the exception of Flathead and Lake Counties, mapping for the counties in Region One is not yet complete. This procedure could be implemented in the future as soil mapping is completed.

Sites where fractured bedrock lies close to the soil surface are also of concern. Flathead Lake is an example of such a circumstance.

Due to the interplay between surface and groundwater and the gravelly nature of river floodplains in northwest Montana, Fishing Access Sites would be of concern as well. Furthermore, groundwater is used as a public drinking water supply on 12 FWP properties in Region One. They are:

Flathead Lake SP-Big Arm	Flathead Lake SP-Finley Point
Flathead Lake SP-Wayfarers	Flathead Lake SP-West Shore
Flathead Lake SP-Yellow Bay	Lake Mary Ronan SP
Logan SP	Lone Pine SP
Thompson Falls SP	Whitefish Lake SP
Somers FAS	Sophie Lake FAS

Broadscale applications with high herbicide rates are not recommended on these sites. Decreasing application frequencies and reducing the concentration of active ingredient in herbicide formulations would also reduce contamination risks.

3.4 Surface Water

3.4.1 Current Conditions

Many of the properties administered by FWP in Region One are associated with surface waters and have high recreation values associated with them. Surface water quality in Montana is generally good. Ninety percent of the total stream miles in Montana fully support all of their designated uses. However, those stream miles not fully supporting their designated uses are impacted in non-point source categories. In most cases, the impairment (s) is an alteration of flow, an increased level of suspended solids, and/or excessive siltation; however, in some circumstances, the impairment is related to mine tailings or mine waste water. Of those waterbodies in Region One with some level of impairment in these four categories, only one FAS is located on the Whitefish River which is impaired by excessive siltation.

3.4.2 Analysis of Potential Impacts

General

There is public and agency concern for chemical contamination of the environment, in general and water resources, in particular. Herbicides may reach surface waters through non-point sources such as surface runoff, groundwater discharge, spray drift, and erosion of soil containing herbicides. Point source releases may occur during cleaning, mixing or loading activities or from accidental spills. One other avenue of water contamination stems from the potential for cumulative impacts to surface water quality through repeated chemical treatment of a particular site.

The potential for non-point source discharge of herbicides to surface waters is a function of herbicide characteristics, application rate, soil type, slope, vegetation, the length of time between application and rainfall, and the presence or absence of an untreated buffer between the application site and surface water. Picloram and 2,4-D have both been reported to enter surface water through runoff (MDA 1991). However, a study conducted under field conditions by EPA measured runoff of picloram a short distance downstream from the application site. The authors concluded that picloram does not present a serious threat to surface water quality (Evans and Dvseja 1973). Because the characteristics of clopyralid are similar to picloram, the runoff potential of clopyralid should be similar. The chemical glyphosate strongly adsorbs to soil particles and would be unlikely to enter surface waters through leaching. However erosion of soil particles themselves to surface waters may cause the introduction of glyphosate after a precipitation event.

Little surface water contamination due to herbicide run-off from treated areas would be expected after major or minor storm events. After major storm events, instream dilution rates would be greatly enhanced whereas, during a minor storm event, runoff would be insufficient to mobilize any herbicides not bound to vegetation or soils. Sassman et al. (1984) found that the highest potential for herbicide contamination via run-off occurred when a moderate-sized storm event happened shortly after an area had been treated. In these situations, they found as much as three percent of the chemical applied to the watershed entered surface waters. However, in a highly porous, granitic watershed in Montana, Watson et al. (1989) were unable to detect any picloram in surface waters during 90 days of monitoring after application of 1 lb/acre of this herbicide.

Aerial drift has the lowest potential to cause measurable contamination of surface water. At the application rates proposed, incidental drift into water bodies would result in very low, usually undetectable, instream or in-lake concentrations of these substances. Furthermore, if label instructions are followed and treatment areas are carefully planned with use of buffer zones, even incidental drift into surface waters should not occur.

Point source discharge from accidental chemical spills pose the greatest risk to surface water quality. The introduction of large quantities could contaminate drinking water supplies and instream uses for several miles below the spill. But, even in this worst-case situation, surface

water contamination would be predominately short-term and acute rather than long-term. This is because the herbicides planned for use by Region One are low to moderate in toxicity and relatively short-lived in the environment.

All Federal and State laws and guidelines concerning the maintenance and enhancement of water quality are applicable to FWP. For example, as legal standards, the Federal Safe Drinking Water Act establishes maximum contaminant levels for potentially toxic substances in drinking water supplies (Table 7). The Environmental Protection Agency's [EPA] Gold Book provides additional protection guidelines for the health and welfare of aquatic life and wildlife (EPA 1986). Revisions to and/or inclusion of new substances in EPA's Gold Book occur periodically.

Other weed treatment methods may affect surface water quality by increasing sediment delivery to water courses. Intensive cultural treatments (e.g. tilling and reseeded) would produce the highest volume of sediment eligible for transport to a surface water course. The accumulation of sediment is negatively correlated with salmonid spawning success. Despite decreasing vegetative cover of a site, manual methods would not produce nearly as much overland runoff and erosion as tilling because manual methods are limited to very small populations whereas cultural methods are usually applied to a much larger land area. Biological methods would not affect surface water quality.

Under Alternative 1 (Status Quo), existing treatment regimes would continue. Surface water impacts would stem from the potential for chemical run-off from treated areas, accidental spills, or aerial drift during application. These risks are discussed above. Additional safeguards proposed under the Plan and embodied in Alternative 2 (Integrated Noxious Weed Management) would not be implemented under Alternative 1. Biological control agents already introduced on some properties would not impact surface waters.

Alternative 2 (Integrated Noxious Weed Management) attempts to minimize the environmental consequences of herbicide treatments by diversifying weed treatment methods to incorporate non-chemical methods. Where chemicals are applied, all precautions/restrictions proposed in the Plan which apply to shallow groundwater areas will be implemented during treatments near surface waters. For example, herbicide loading, mixing, and storage areas will be located at least 500 feet from open water or shallow groundwater. If mixing and loading of chemicals were to occur on site, open water would not be used for mixing to prevent contamination from back siphoning. Chemicals would not be applied less than 50 feet from any surface water unless a wick applicator is used. Picloram, the most persistent of the herbicides proposed for use, will not be applied on the immediate banks of a water body, below the high water level of a stream, or in identified sensitive groundwater areas. Biological, manual, and cultural methods would not impact surface waters, although cultural methods may temporarily increase sediment delivery to surface waters until vegetation has become established.

Alternative 3 (Chemical Methods) presents the highest probability of adversely impacting surface waters. The potential impacts would be as described above. Because chemical treatment is the only method proposed in this Alternative, the potential for accidental spills, run-off from treated areas, or aerial drift to contaminate surface waters is heightened.

Alternative 4 (Non-Chemical Methods) presents the least risk to surface waters because chemical treatment methods are not incorporated into this Alternative. There is the potential, however, to increase sediment delivery to surface waters in the short term from cultural treatment methods. Biological, manual, and cultural methods would not impact surface waters, although cultural methods may temporarily increase sediment delivery to surface waters until vegetation has become established.

Site Specific

Site specific impacts may occur on any property with surface water. Along with Fishing Access Sites, Wildlife Management Areas and State Parks also have perennial streams. In addition, those properties with surface waters used by the public for boating or swimming could be impacted. For example, Lake Mary Ronan is quite popular with the boating public in northwest Montana.

Table 7. Maximum concentrations not to be exceeded in drinking water supplies for herbicides proposed for use by FWP in Region One. Information provided by the United States Environmental Protection Agency, Office of Drinking Water, 1990.

Compound	Status of Listing	Maximum Contaminant Level ($\mu\text{g/l}$) ²	Drinking Water Formulation Equivalent Level ($\mu\text{g/l}$)
clopyralid	-	no data available	no data available
2,4-D/amine	P ¹	70	400
glyphosate	P	700	4000
picloram	P	500	2000

¹ P = proposed for listing

² $\mu\text{g/l}$ = parts per billion

3.5 Fish and Wildlife

3.5.1 Current Conditions

Roughly corresponding with FWP Region One boundaries, the northwest corner of Montana embodies some of the richest and most diverse flora and fauna across Montana. There are over 400 vertebrate species known to occur in latilongs 1, 2, 3, 13, 14, and 15, where a latilong is a unit of land encompassed by one degree of latitude and one degree of longitude. Appendix B, compiled from several sources as cited, lists the species that have been documented in the latilongs corresponding with Region One. Eight big game species, 68 other mammal species, over 300 bird species, 14 amphibian species, 9 reptile species, and 33 fish species are known to occur in this area (Bergeron et al. 1992, Holton 1990, Thompson 1982). With some exception, many of these species use Region One's properties for seasonal or year-round habitation. Waterfowl, upland bird and big game hunting, wildlife viewing, and fishing are primary recreational uses and management focuses of FWP Region One. Table 8 presents a partial list of fish and wildlife species which occur in Region One.

Table 8. A partial list of fish and wildlife species in northwest Montana.

Terrestrial	Raptors	Waterfowl	Upland Birds	Aquatic
white-tailed deer	bald eagle ¹	Canada goose	ring-necked pheasant	brown trout
mule deer	golden eagle ²	mallard	Ptarmigan	rainbow trout
mountain goat	peregrine falcon ¹	pintail	ruffed grouse	kokanee
bighorn sheep	osprey	gadwall	blue grouse	yellow perch
elk	Swainson's hawk	widgeon		westslope cutthroat ²
antelope	red-tailed hawk	green winged teal		mountain whitefish
moose	kestrel	blue winged teal		Yellowstone cutthroat ²
mink	northern harrier	great blue heron ²		brook trout
muskrat	ferruginous hawk ²	sandhill crane		artic grayling ²
beaver	rough-legged hawk	white pelican ²		lake trout
weasel		common loon ²		bull trout ^{1,2}
grey wolf		canvasback		largemouth bass
coyote		scaup		smallmouth bass
red fox		redhead		northern pike
mountain lion				lake whitefish
grizzly bear ¹				white sturgeon ^{1,2}
black bear				inland redband ^{1,2}
caribou				peamouth chub
				pygmy whitefish
				longnose sucker
				largescale sucker
				northern squawfish
				redside shiner
				burbot
				pumpkinseed
				bluegill
				sculpins

¹ Species listed under the Federal Endangered Species Act (ESA); the Kootenai River white sturgeon has been listed as endangered under ESA and by the U.S. Fish and Wildlife Service. The bull trout was petitioned for listing in 1993. The FWS concluded that listing was warranted for the species within the contiguous United States but listing was precluded until the process was complete on high priority species. The inland redband trout was recently petitioned and is undergoing a status review by the U.S. Fish and Wildlife Service. Westslope cutthroat have been reduced to less than 10 percent of their historic range.

² Species of Special Concern, as identified by the Montana Natural Heritage Program.

3.5.2 Analysis of Potential Impacts

General

Fish

Region One, as part of the Mission of FWP, is charged with protecting and enhancing fish and wildlife populations and their habitats. Fish populations suffer from the loss or degradation of habitat caused by resource development or other human activities. One threat to fish habitat is the impairment of water quality. Degradation of water quality may directly impact fish populations by preventing the successful completion of a life history stage. Indirect impacts on fishery resources result from negative effects on the aquatic community at lower trophic levels such as vegetation or insects. Furthermore, indirect impacts may result by obstructing the interplay between aquatic and terrestrial environments, as may occur, for example, with loss of the riparian vegetation stabilizing a streambank.

Under Alternative 1, (Status Quo), the existing treatment regimes would remain in place. Chemical treatments would continue at their current locations. Biocontrol agents would persist, as well. This Alternative may impact fishery resources if chemicals are introduced into surface waters. Introduction may result from leaching, accidental spill, aerial drift during application, or as run-off from treated areas after a precipitation event. The discussion of the potential impacts to surface waters summarizes these concerns. However, once herbicides are introduced into surface waters, the toxicity to fish and other aquatic life varies greatly depending upon the species in question, the chemical formulation, and percentage of active ingredient. According to criteria established by the U.S. Fish and Wildlife Service (1980), none of the chemicals proposed for use by FWP are highly toxic to fish and aquatic life (i.e. having an LC_{50} less than 1.0 parts per million [ppm], where LC_{50} is the concentration of toxicant that is fatal to 50% of the tested organisms over a specified time). Established populations of biocontrol agents are not expected to impact fishery resources.

Under Alternative 2 (Integrated Noxious Weed Management), all methods of noxious weed management may be utilized. The integrated approach will attempt to minimize impacts to fishery resources from chemical treatment activities through implementation of the guidelines contained in the Plan. The Plan further minimizes potential consequences to fish by requiring that only the amine formulation of 2,4-D and the Rodeo[®] formulation of glyphosate be applied near surface waters. Both of these herbicides have been shown to be only slightly toxic to trout (i.e. having LC_{50} values ranging from 100 to 1000 ppm). Also, these compounds readily adsorb to soils, which minimizes their mobilization into surface waters, and they do not persist in the environment for long periods.

Cultural methods which disturb the soil surface (e.g. plowing) may result in short-term increases in sediment delivery to a surface water until the site is revegetated. Increased sediment delivery may impact fish populations because excessive sedimentation is negatively correlated with

salmonid spawning success and the survival of fish embryos. Manual and biocontrol treatment methods likely would not increase sediment delivery or affect water quality and would not impact fishery resources.

Alternative 3 (Chemical Methods), in emphasizing chemical treatment methods, has the greatest potential to impact fishery resources. Direct impacts could occur if instream concentrations of herbicides exceeded levels that are toxic to resident fish. Indirect, food-chain impacts could occur if in-stream levels become elevated enough to become toxic to aquatic vegetation (particularly algae) or insects.

Because of its reliance on cultural, manual, and biological treatment methods, Alternative 4 (Non-Chemical Methods) has the greatest potential to increase sediment delivery to surface waters. Since manual and biological methods are not as successful at keeping pace with noxious weed seed production, cultural methods would likely comprise a large portion of the treatment activities. Some cultural methods (e.g. plowing) may increase surface runoff to surface waters, leading to increased sediment delivery which could affect spawning success. However, since chemicals would not be used, potential chemical contamination of surface waters would not be possible.

Wildlife

Wildlife populations also suffer from loss or degradation of habitat caused by resource development or other human activities. In the context of noxious weeds, human-influenced disturbance has affected wildlife habitat to the point that noxious weeds are encroaching upon some vegetation communities. Encroaching non-native weed species can alter the balance of structure and species composition in vegetation communities. Most native plant communities provide wildlife with diverse, high quality food and cover resources which could be reduced if vegetation communities become dominated by well-established noxious weeds. However, noxious weed treatment activities may also have some negative consequences to wildlife and wildlife habitat because vegetation is altered or removed. For example, a few wildlife species benefit directly from the presence of noxious weeds. Some birds eat weed seeds and others seek cover in dense weed patches. Small mammals have also been known to use weed seeds as a food source.

Wildlife populations and habitats can be positively and negatively affected by all Alternatives considered in this EA. Each Alternative includes some level of noxious weed treatment. Diminished grass and forb production due to noxious weed encroachment and proliferation may negatively affect habitats for many wildlife species during various seasons of the year. But under all Alternatives, weed treatment activities could disturb wildlife using an area. An indepth discussion of the positive and negative aspects of each Alternative follows.

Under Alternative 1, (Status Quo), the current noxious weed treatments would continue. Chemical applications would proceed and previously introduced biocontrol agents would be persist until they could no longer be sustained by the weed population or died of other causes.

Under this Alternative, weeds would likely flourish and expand their distributions. The detrimental effects of noxious weed encroachment into wildlife habitat have already been discussed. Since there would be some degree of chemical treatment activity, wildlife could be directly or indirectly affected by chemicals introduced into the environment as a result of weed management activities.

Chemical applications could affect wildlife directly through the ingestion of treated plants or by dermal absorption if an animal came into contact with treated vegetation. If properly applied, it is unlikely that any species would receive toxic doses of herbicides. However, in cases where animals are directly sprayed or feed exclusively on sprayed vegetation, some could receive toxic doses (BLM 1985). Toxicities of various chemicals considered for use cover a wide range. Tests on domestic animals have shown that the LD₅₀ (lethal dose to 50% of those tested) for 2,4-D is the lowest (most toxic) of pesticides considered in this assessment (Tables 9 and 10). For cattle (or for comparison, elk) grazing immediately after spraying 2,4-D at prescribed rates and only on sprayed vegetation, ingestion of the chemical would only be about 20% of the LD₅₀ (Lolo National Forest 1989). For dogs (or for comparison, coyotes), ingestion of a large quantity of meat from an animal that has eaten heavily on sprayed vegetation would result in less than 1/400 of the LD₅₀. It is difficult to extrapolate the level of effects on domestic animals to wildlife. Nonetheless, based on these data, it appears that neither elk nor coyotes could ingest enough of any pesticide through normal eating behavior to be fatal.

Table 9. Median lethal dose (LD₅₀) for domestic animals expressed in milligrams per kilogram for 3 herbicides proposed for use on Department of Fish, Wildlife and Park properties in Region One.

Species	Picloram	2,4-D	Glyphosate
mouse	2,000 - 4,000	368	²
(rat)	8,200	375	4,320
dog	-	100	-
cat	-	820	-
chicken	6,000	541	15,000 no effect
(mallard)	> 2,000	-	> 2,000
pigeon	-	668	-
rabbit	2,000	424	3,800
cow	540	100	-
(mule deer)	-	400 - 800	-

¹ The data in this table were taken from Lolo National Forest (1989); Sassman et al. (1984); USDA (1988). The reader should consult those documents for thorough discussions of herbicide toxicity.

² Data not available.

Monnig (1988) demonstrated that warm-blooded animals, when fed high concentrations of 2,4-D, picloram, and glyphosate, excrete the chemicals rapidly and do not retain pesticides in internal organs. Analysis performed for the final EIS on eradication of cannabis on federal lands showed that most large animals cannot receive lethal doses of herbicides (2,4-D or glyphosate) and that it is very unlikely in even the most extreme dose cases that small animals (such as mice or toads) would die (Table 10) (BLM 1985). This table demonstrates that in a worst-case situation, no listed animal would receive a dose greater than 11% of its median lethal dose for 2,4-D or 1% for glyphosate (BLM 1985). Thus, it seems that most wildlife species are not likely to be affected directly by ingestion of herbicides or herbicide-treated vegetation. However, none of these studies considered the possible impacts of chemical treatment in areas used for calving, fawning, nesting, or rearing of young. It is possible that young animals could be negatively affected through direct contact with or ingestion of herbicide. Additional research is required.

Bio-accumulation of contaminants in the food chain was of critical importance with the now-banned insecticides DDT and DDE, particularly for sensitive species such as raptors. It has been shown that herbicides proposed for use under the Plan are metabolized readily and excreted by mammals. Glyphosate, 2,4-D, and picloram do not bio-accumulate in fish and mammals (EPA 1983, 1986, 1988; USDA 1984). No evidence was found in the literature to support bio-accumulation problems from clopyralid.

Indirectly, chemical applications, may negatively affect wildlife over the short-term by killing native broadleaf plant species which are important sources of seasonal food and/or cover. Depending on the rate of application and the techniques used to apply herbicides, there can be losses of non-target vegetation important to wildlife species (BLM 1985). While woody species such as sagebrush and willows are highly valued for their uses by wildlife, they can be acutely affected by herbicides and the negative effects are long-term. In addition, wildlife may be temporarily displaced during treatment activities.

Currently established biocontrol agents are not expected to impact wildlife resources, either directly or indirectly.

Alternative 2 (Integrated Noxious Weed Management) combines manual, cultural, biological, and chemical treatments methods. By using an integrated approach to weed management, areas critical to wildlife populations during specific seasons could be treated at times or by techniques that would have the fewest negative impacts. Careful application of herbicides in sensitive areas, in conjunction with biological, manual, or cultural techniques, offers the best solutions for preserving the native vegetation component of wildlife habitat and the integrity of wildlife resources. In situations where woody species are important to wildlife management goals (e.g. Wildlife Management Areas), non-chemical methods will be emphasized.

Table 10. Routine and extreme dose levels of 2,4-D and glyphosate compared to the median lethal dose (LD₅₀) for various animal species. Table is adapted from the Bureau of Land Management Northwest Area Noxious Weed Control Program Final Environmental Impact Statement, 1985.

Species	Routine Case Dose	Extreme Case Dose	Median Lethal Dose LD ₅₀ ¹	Lab Animal
2,4-D				
Flicker	13.700	123.33	472 (T)	Pheasants
Dove	8.710	41.35	472 (T)	Pheasants
Jay	12.850	97.95	472 (T)	Pheasants
Kingfisher	0.017	0.42	472 (T)	Pheasants
Owl	7.930	23.63	472 (T)	Pheasants
Mouse	39.640	197.90	380 (BE)	Mouse
Rabbit	5.520	44.70	424 (BE)	Rabbit
Deer	1.140	13.76	400 (A)	Mule Deer
Fox	2.020	9.27	100 (A)	Dog
Toad	18.040	139.90	200 (M)	Tadpole
Snake	19.820	59.10	200 (M)	Toad
Eggs	9.420	47.10	No adverse effects at up to 35 mg/k	Hen eggs
Glyphosate				
Flicker	7.2100	64.90	4,640	Quail
Dove	4.5900	21.78	4,640	Quail
Jay	6.7600	51.51	4,640	Quail
Kingfisher	0.0087	0.22	4,640	Quail
Screech Owl	4.1700	12.44	4,640	Quail
Mouse	20.8600	104.20	4,000	Rat
Rabbit	2.9100	23.53	3,800	Rabbit
Deer	0.6000	7.22	3,800	Rabbit
Fox	1.0600	4.88	3,800	Rabbit
Toad	9.4900	73.65	- ²	
Snake	10.4300	31.10	-	
Bird Eggs	4.9600	24.78	-	

¹ Dose (mg/kg) lethal to 50% of those tested; BE = Butyl Ester, A = Acid, T = Technical, and M = Amine.

² No data available.

This approach integrates treatment techniques with natural resource values and assures that the proper technique will be used in sensitive areas to minimize impacts to other resources, including wildlife. The Plan's management objective for wildlife is to "protect existing, diverse wildlife habitats from encroachment by weeds while ensuring that threatened, endangered, rare, or unique species are not adversely affected by weed treatments." The Plan lists several management guidelines which describe the inherent mitigative aspects. These address specific and general approaches to treating sensitive areas that will be determined and delineated for each project, regardless of the treatment method.

Direct impacts to wildlife under Alternative 2 resulting from chemical applications are the same as those described for Alternative 1. Indirect impacts to wildlife resulting from chemical applications would also be similar to those described above. Other indirect impacts may result from temporary displacement because important seasonal vegetation may be disturbed during non-chemical treatments such as mowing.

The establishment of biocontrol agents is not expected to affect wildlife, either directly or indirectly.

Alternative 3 (Chemical Methods), emphasizes chemical methods in any situation requiring noxious weed treatment. Direct and indirect impacts resulting from this Alternative are similar to those discussed for Alternative 1. Furthermore, Alternative 3 poses the greatest risk to wildlife resources because the risks of over-application, chemical spills, and aerial drift during application are highest under this Alternative.

Under Alternative 4 (Non-Chemical Methods), non-chemical methods would be the only means of arresting the encroachment of noxious weeds into desirable vegetation communities. Vigorous application of manual and cultural methods under this Alternative, may temporarily delay the establishment of noxious weeds in weed-free areas, but would probably not effectively reduce weeds in vital habitats presently containing weeds. Direct impacts to wildlife of non-chemical methods include the short-term displacement of animals during treatment activities. Indirect impacts may result because non-target, native species may be impacted during treatment. Noxious weeds may encroach upon scarified, cultivated, or other disturbed soils if revegetation of native species is not rapid. The use of grazing livestock such as sheep to reduce and contain weeds, in addition to possibly displacing wildlife, may negatively affect non-target vegetation. If Alternative 4 (Non-Chemical Methods) is selected, weeds will likely expand and continue to encroach upon desirable plant communities.

Site Specific

Fish

Many properties in Region One (Fishing Access Sites in particular) incorporate surface waters which sustain aquatic communities, including sport fisheries. These sites could be adversely affected by inappropriately or improperly applied herbicides. If precautions and mitigation measures outlined in the Plan and incorporated into Alternative 2 are followed, the risk of negatively affecting the fishery resources is slight.

Species of Special Concern: As discussed previously, potential impacts to fishery resources exist, but can be mitigated under Alternative 2. Currently, only the Kootenai White Sturgeon occurring in Region One is listed under the Endangered Species Act, although the Bull Trout was deemed "warranted but precluded." Although not petitioned for listing, the westslope cutthroat trout and inland redband are considered Species of Special Concern. All three species have a limited distribution. Therefore, chemical treatment activities should be considered carefully on Fishing Access Sites in the Clark Fork, Flathead and Kootenai River drainages.

Wildlife

Direct negative effects to wildlife resources are likely to be minimal, as discussed previously. However, potential indirect effects of herbicide applications on wildlife vegetative food sources do exist, especially for broadleaf forbs and woody shrubs. If chemical treatments are not carefully conducted, important vegetative species could be affected. This would be particularly true on the Wildlife Management Areas which typically harbor ungulates during winter, the most stressful season for wildlife. Also of note are the shallow lake and wetland habitats associated with the Thompson Chain of Lakes and the riparian zones associated with Fishing Access Sites which provide high quality, diverse wildlife habitats. Emergent, floating, and submergent vegetation important to waterfowl and other species could also be negatively affected by improperly applied chemicals. If the additional safeguards outlined in the Plan (Alternative 2) are followed, the risk of negatively affecting wildlife resources is minimal.

Species of Special Concern: As discussed previously, potential toxic effects from herbicides is minimal for wildlife species in general. However, federally-designated threatened or endangered species require special consideration. Although not recognized under the Endangered Species Act, Species of Special Concern also require consideration. The primary concern for bald eagles and peregrine falcons is disturbance during nest building and incubation (March through June). The proximity of nests to Thompson Chain of Lakes FAS and Lake Mary Ronan and Flathead Lake State Parks makes implementation of weed management actions of some concern. Weed

treatments would usually occur late May through September. The level of disturbance should be no greater than that expected from typical recreation use at these sites. Mitigation measures include guidelines recommended by the Montana Bald Eagle Working Group (1991). The grizzly bear, another species with federal recognition, frequent more remote settings in the backcountry whereas the majority of noxious weed treatment activities occur near roads and trails. Nonetheless, the potential presence of grizzly bears should be appraised prior to weed treatment activities.

3.6 Vegetation

3.6.1 Current Conditions

The diverse geology and topography of northwestern Montana yields a diverse pattern of vegetation, including alpine meadows, forested slopes, shrub steppe, prairie grasslands, and wetland marshes. Vegetation resources on FWP Region One lands include rangelands, wetlands, riparian areas, and forested areas. Finer textured soils support short grass communities. Upland shrub/grassland communities include big sagebrush, fringed sagewort, and antelope bitterbrush with Rocky Mountain juniper and mountain mahogany occurring in rockier areas. Spruce - Fir forests with shrub understories are found at the higher elevations and on steeper slopes. Riparian communities include cottonwood, aspen, willow and numerous other mesic shrub species. Wetland marsh communities are found at Thompson Chain of Lakes FAS and Ducharme FAS to name just two.

During a survey of all Region One properties during the summer, 1993, noxious weeds were documented on all but a few properties, though with considerable variation in abundance. Spotted knapweed, Canada thistle, and leafy spurge were the most common noxious weed species encountered. Other "noxious" weed species declared by individual counties were also documented on FWP properties. The most common were hound's tongue, musk thistle, and mullein. Noxious weeds are strong competitors for moisture and soil nutrients. The possible effects of a decline in native vegetation resources on wildlife has already been discussed previously.

Plant Species of Special Concern

The Natural Heritage Program system, a partnership between The Nature Conservancy and state governments, was established in part as a clearinghouse to aid states in identifying and studying rare species. Rare plant species are protected under the Federal Endangered Species Act of 1973 (PL 93-205 as amended and rules 50 CFR Part 402). At this time, there are no threatened or endangered plant species on Region One properties. However, according to the Montana Natural Heritage Program, a number of plant species listed as Species of Special Concern (because of limited distribution or small populations) occur on Region One properties. Table 11 summarizes these.

Table 11. Plant Species of Special Concern on FWP properties in Region One, as identified by the Montana Natural Heritage Program, 1994.

Common Name	Scientific Name	Status ¹	Location
Twin Clover	<i>Trifolium latifolium</i>	G4, S1	Mount Silcox WMA
Rhombic-petaled Clarkia	<i>Clarkia rhomboidea</i>	G4, S1	Mount Silcox WMA
Columbia Water-meal	<i>Wolffia columbiana</i>	G4, S4	Ninepipe WMA
Spalding Campion	<i>Silene spaldingii</i>	G2, S1	Flathead Lake SP - Wild Horse Island
Slender Sedge	<i>Carex tinctoria</i>	G4, G5, SU	Flathead Lake SP - Yellow Bay
Giant Helleborine	<i>Epipactis gigantea</i>	G4, S2	Flathead Lake SP - Yellow Bay
Adder's Tongue	<i>Ophioglossum vulgatum</i>	G5, S1	Flathead Lake SP - Yellow Bay
Small Yellow Lady's Slipper	<i>Cypripedium calceolus var parviflorum</i>	G5, S2, S3	Whitefish Lake SP

¹ Global ranks:

- G2 imperiled; 6-20 occurrences
- G3 threatened throughout range
- G4 apparently secure
- G5 demonstrably secure
- GU possibly in peril, status uncertain

State ranks:

- S1 critically imperiled in Montana; ≤ 5 occurrences
- S2 critically imperiled in Montana; 6-20 occurrences
- S3 rare in Montana; ≥ 21 occurrences
- S4 apparently secure
- S5 demonstrably secure
- SU possibly in peril, status uncertain

3.6.2 Analysis of Potential Impacts

General

The nature and extent of impacts on vegetation resources as a result of noxious weed management would vary under each Alternative, as described below.

Under Alternative 1 (Status Quo), noxious weeds would be treated, but only at the current intensity levels. Thus, it is likely that noxious weeds would continue to spread and that new species would likely establish on FWP properties. Because treatment activities would be limited, it is also likely that noxious weeds would spread within Region One properties and beyond property boundaries. Under current direction, chemical treatments would continue. Chemical treatments may affect non-target vegetation. This could result from the use of non-selective herbicides, broad-scale application methods, or over application. Impact severity would depend

on the environmental conditions during application, application rate, method, carrier, and the herbicide. Herbicides which are selective for broadleaf weed species (such as 2,4-D and picloram) also kill native forbs, shrubs, and trees, thereby potentially decreasing natural diversity. Grasses are tolerant of most broadleaf herbicides and would benefit from treatment. Non-selective herbicides (glyphosate) can kill all vegetation, which creates a setting conducive for the re-establishment of weeds. Established biocontrol agents generally do not affect non-target vegetation resources because the agents are host-specific to a particular noxious weed species (Littlefield 1993). However, there was one reported incidence of an insect affecting a non-target species. Introduced to musk thistle, the seed weevil *Rhinocyllus conicus* also plagued a rare, native thistle called longstyled thistle (*Cirsium longistylum*) (Achuss and Schassberger 1991). The likelihood of a similar outcome in the future is reduced because now screening tests are more rigorous and include taxonomically related and morphologically similar native species (Littlefield 1993).

Alternative 2 (Integrated Noxious Weed Management), is expected to have the least impact on vegetation resources. This Alternative specifies the implementation of the Plan which allows for flexibility in approach and the use of optimum treatment methods for specific sites. Chemical treatments would still be used; however, the Plan outlines precautions and use standards that are designed to minimize or eliminate the potential adverse effects of herbicides on desirable vegetation. Special attention is given to riparian and wetland areas. If these guidelines are followed, then impacts would be low. Manual methods, used in localized situations where weed populations are small, would not adversely impact desirable vegetation. Cultural treatments such as mowing can be a short-term solution to prevent seed production. Short-term effects on desirable vegetation could occur. Tilling would be considered only for areas that had previously been cultivated because soil disturbance increases the likelihood of weed encroachment. Use of grazing animals to treat weeds would have little impact on desirable vegetation as long as herds are confined and use is carefully monitored since livestock will feed both on weeds and desirable vegetation. Indirectly, compaction of a site could slow the recovery of desirable plants. Because biological control organisms are host specific, no adverse impacts to desirable vegetation are expected. This Alternative is the best solution for the long-term management of noxious weeds and preservation of vegetation resources through implementation of the Plan.

Alternative 3 (Chemical Methods) is expected to have the greatest impact on vegetation resources. Since chemical treatments would be the sole method, non-target vegetation would be adversely impacted, particularly woody species. The type and severity of such impacts are discussed under Alternative 1. Although this Alternative would effectively reduce noxious weeds, the magnitude of impacts to non-target, desirable vegetation could be unacceptable.

Under Alternative 4 (Non-Chemical Methods), the emphasis would be on manual, cultural, and biological control methods. These methods would likely have some adverse impacts on non-target vegetation, as discussed for Alternative 2 (Integrated Noxious Weed Management). This Alternative would not lead to long-term success at managing noxious weeds. Rather, noxious weeds would likely proliferate.

Plant Species of Special Concern

Species of Special Concern are declared as such because their numbers are greatly reduced, their habitat is declining or threatened, or because populations occur in a limited or non-contiguous distribution. Studies addressing how noxious weeds affect species of concern are few. Lessica and Shelly (1991) showed reductions in reproductive output of sapphire rockcress (*Arabis fecunda*) where spotted knapweed had invaded. Currently, the rates of spread of spotted knapweed and leafy spurge appear to be the highest of the noxious weeds encountered on FWP Region One properties. It is known that noxious weeds are very competitive and could crowd out Species of Special Concern. For those rare species whose habitat coincides with areas susceptible to noxious weed encroachment, the threat may be very high. For example, Flathead Lake State Park - Wild Horse Island contains the Palouse Prairie habitat type association of Bluebunch Wheatgrass and Spalding Campion (*Agropyron spicatum* - *Silene spaldingii*). The Spalding Campion is rare throughout its range and therefore the Montana Natural Heritage Program indicates that the site should be evaluated for weed invasion.

Rare plants typically occur in very small populations and usually in non-contiguous distributions. Use of herbicides could cause a significant loss by eliminating a population (s) or reducing numbers within a population. Cultural methods could cause uprooting or physical damage, loss of seed production, or outright destruction. Identification of rare plant populations during treatment area planning would prevent losses. Identification of such populations is only incorporated into Alternative 2, through the Plan. None of the other Alternatives include any provisions for Species of Special Concern.

Site Specific

All Region One properties have desirable vegetation communities that could be impacted by improper herbicide treatments, whether it be in a natural field setting such as a Wildlife Management Area or a landscaped setting such as the Regional Headquarters Complex. Spot spraying and the use of appropriate herbicide formulations would reduce the potential for short-term injury or long-term plant community alteration. In the case of Species of Special Concern, impacts resulting from weed management actions could be acute in the absence of knowledge of these rare plants. Site specific impacts to populations of plant species of Special Concern could occur on any of the FWP properties supporting rare plants. Those plant species and the FWP properties on which they occur are listed in Table 11. Those sites are: Mount Silcox WMA, Ninepipe WMA, Flathead Lake SP - Wild Horse Island, Flathead Lake SP - Yellow Bay, and Whitefish Lake SP. Impacts to *Cypripedium* plants at Whitefish Lake State Park would not be necessary because there have been no noxious weeds identified within the park. Additionally, populations of *Silene spaldingii* on Wild Horse Island would not be impacted from chemical treatment methods on spotted knapweed and leafy spurge if they are carried out soon on the small patches elsewhere on the island before they are spread by wildlife. However, all rare plants listed in Table 11 could be affected by chemical treatments because these species are either broad-leaved or woody, characteristics which most herbicides are not selective against.

Additional safeguards outlined in the Plan would decrease the risk of negatively affecting these rare plants.

3.7 Recreation

3.7.1 Current Conditions

Northwest Montana draws much national and international attention for its natural resources and outdoor recreation opportunities. FWP properties in Region One are part of that attraction. These lands offer a wide variety of experiences such as waterfowl, upland game bird and big game hunting, year-round fishing, camping, picnicking, swimming, sailboarding, motorboating, canoeing, waterskiing, cross-country skiing, ice skating, and wildlife viewing. Other sites provide opportunities to experience historical settings (e.g. Wild Horse Island) or unique natural features (e.g. Flathead Lake State Park). In addition, FWP properties offer a wide variety of landscapes in which to recreate, ranging from river bottoms at Fishing Access Sites, bunchgrass slopes or forested hills on a Wildlife Management Area, or highly scenic forested lakeshores. Healthy, native plant communities are a significant part of the recreational experience, from both ecological and aesthetic perspectives.

The most heavily used roads, trails, campsites, and picnic areas have problems associated with soil compaction, braiding of trails, and noxious weeds resulting from disturbance. As indicated previously, noxious weeds occur on most FWP properties in Region One. Some recreationists find that noxious weeds detract from their enjoyment of an outdoor experience or particular site, while others don't seem to mind. Yet other recreationists are unaware of noxious weeds, how they are spread, and the repercussions of unchecked weed proliferation upon natural and recreational resources. Thus the impact of noxious weeds upon recreationists will depend on the perspective of the recreationist. However, most recreationists would likely be indirectly affected if weeds were allowed to spread unchecked. This is because of the concurrent changes in desirable vegetation communities.

3.7.2 Analysis of Potential Impacts

General

The impacts of noxious weed treatment upon recreation vary with the type of treatment and the type of recreation. Generally, recreation on FWP properties is oriented towards foot or horse travel as vehicles are restricted to established roads, trails, and parking areas. None of the Alternatives are expected to affect vehicular access or the driving of any motorized vehicle where they are permitted. However, while travelling on foot (or horse back) and pursuing any of the recreational activities listed above, recreationists may be impacted by the Alternatives, as described below.

Alternative 1 (Status Quo) would likely have a long-term negative impact on recreation resources. Under this Alternative, weed treatments would continue, but weeds would likely proliferate. Therefore, it has the potential to greatly affect recreation in the long run through changes to vegetation communities and, potentially, wildlife resources. Treatment of weeds as currently practiced would involve chemical applications and the established biocontrol agent releases. Of the two methods, chemical treatments would affect recreationists in the short-term, whereas biocontrol agents would not. In some situations, chemical applications may preclude public use for a period of time during and after the application. Similarly, some recreationists may find chemical odors unpleasant and offensive. Recreationists may be visually or aesthetically impacted since herbicides can affect non-target vegetation.

Alternative 2 (Integrated Noxious Weed Management) would have the fewest long-term, negative impacts to recreation. Under this Alternative, the Noxious Weed Management Plan would be implemented which combines a wide variety of weed treatments with site-specific planning with public awareness of the problems associated with noxious weeds. Although FWP's mission does not include weed education, the agency may cooperate with public awareness programs initiated by other organizations. Such initiatives would help recreationists identify problem areas, understand where the most sensitive areas exist, and give them the opportunity to be involved in weed management. FWP involvement may include a display at Regional Headquarters Offices or the distribution of a noxious weed information sheet or brochure through the vehicle, boat, and trailer registration system or through hunting and fishing license dealers. Other involvement may include range tours of the Wildlife Management Areas. This Alternative presents the best opportunity for the long-term management of noxious weeds.

Integrated Noxious Weed Management involves the use of chemical, biological, cultural, and manual methods of noxious weed treatment. The impacts of chemical treatments would be similar to those described above, but with the following exceptions. In contrast with Alternative 1, the Plan includes safeguards and guidelines for application which would protect other resource values such as recreation. These guidelines are intended to minimize short-term impacts on recreational resources and public use of FWP properties. Biological, manual, and cultural methods would not affect recreation.

Alternative 3 (Chemical Methods) would have the greatest negative impacts to recreation. By emphasizing herbicide treatments, this Alternative could cause the frequent closure of sites to public use during application, long-term adverse effects on non-target vegetation, and increased risk of exposure to chemicals by the public. Similar impacts as those described under Alternative 1 could be expected under this Alternative. Some degree of weed encroachment would still occur if treatments were not applied correctly or frequently enough.

Alternative 4 (Non-Chemical Methods) prescribes the use of biological, manual, and cultural methods to reduce noxious weed populations. These methods can be effective in certain situations. However, as indicated previously in the discussion of vegetation, cultural methods may actually lead to an increase in noxious weeds if not conducted properly. Manual methods are time-consuming. Biocontrol methods require longer time intervals for successful

management of weeds. For these reasons, this Alternative may not treat noxious weeds as effectively or efficiently as other Alternatives. Thus, over the long-term, recreation may be impacted through the alteration of desirable vegetation communities. In the short-term, however, this Alternative would have the least impact on recreation. Area closures required during chemical applications would not be necessary under this Alternative. Furthermore, biological, manual, and cultural methods would not affect recreational access or pursuits.

Site Specific

Site-specific impacts could be anticipated under those Alternatives which include some degree of chemical treatments and on those sites which sustain the greatest public use. Some of the most visited properties in Region One include Flathead Lake State Park, Lake Mary Ronan State Park, Fishing Access Sites near population centers, and Ray Kuhns Wildlife Management Area. Of the three Alternatives which include some provision for chemical treatments (1, 2, and 3), the site-specific impacts would be greatest under Alternative 1 (Status Quo) because no safeguards for non-target vegetation or other resources would be implemented. Alternative 3 (Chemical Methods) would also result in site-specific impacts at high visitation sites because herbicides would be the only method of treatment, leading to increased frequency of application. Likewise, safeguards would not be implemented under this Alternative either. Alternative 2 (Integrated Noxious Weed Management) would have the least impact on highly visited sites because safeguards would be implemented and because chemical treatments would be just one of the possible treatment methods.

3.8 Human Health

3.8.1 Current Conditions

Montana is recognized nationally for its clean, healthy environment. Implicit in a healthy environment is clean air and water, securing healthy living conditions for humans. As noted previously, Montana's air and water quality are generally good, with isolated exceptions.

3.8.2 Analysis of Potential Impacts

General

General impacts to human health could be anticipated if air and water quality are degraded for any reason. Of all the noxious weed treatment methods under consideration, chemical methods pose the highest risk to air and water quality. Earlier sections of this document specified potential impacts of noxious weed management activities on air and water resources in general and more specifically from chemical methods. The following discussion reviews these impacts from the perspective of human health. Those impacts would be most pronounced on sites

requiring the most intense noxious weed treatments and on those sites where use by the public is high.

Alternative 1 (Status Quo) describes the continuation of current noxious weed treatment methods and intensities: chemical applications without safeguards beyond manufacturer's labels and established biocontrol agents. Human health could be affected by herbicide use under this Alternative. Detailed discussions on this subject can be found in the following publications: Lolo National Forest Noxious Weed Management Draft EIS (1989); Human Health Risk Assessment of Herbicide Applications to Control Noxious Weeds and Poisonous Plants in the Northern Region of USDA Forest Service (Monnig 1988); Northwest Area Noxious Weed Control Program EIS, Final and Supplement (BLM 1985, 1987). We rely on these analyses and highlight several concepts: exposure routes, bio-accumulation, carcinogenic properties, and chemical additives.

The Northern Region assessment (Monnig 1988) is a "worst-case analysis" using several conservative assumptions that purposely over-estimate the likelihood and severity of exposure, accidents, and the resulting effects. Those assumptions include: higher application rates than label recommendations, continuous application over an entire project area, mixing and application errors that would not generally occur, human residences and food sources much closer than would occur in most projects, skin exposure estimates for workers failing to wear recommended protective clothing, and that there is no threshold or safe level for exposure to carcinogens (cancer-causing agents).

Human exposure routes are related to either direct contact with herbicides through spray drift, spills, and sprayed vegetation or to indirect contact through consumption of contaminated water, vegetables, fish, or grazing animals. The exposure risk is greatest for those workers who apply herbicides without the appropriate protective equipment such as clothing, respirators, and eye protection.

Unlike DDT, it appears that bio-accumulation does not occur with picloram, 2,4-D, and glyphosate. These herbicides are water soluble and are excreted rapidly. A small percentage of the population is hypersensitive to one or more of these chemicals and can suffer severe and long lasting symptoms after small doses. Workers are at the highest risk. Using proper application techniques and wearing recommended protective clothing can reduce this risk. Additional precautions would be implemented when herbicides are applied to high human use areas such as work places, campgrounds, and parks. Formulations of glyphosate and 2,4-D registered for use in human occupation areas would be used.

Of the chemicals proposed for use in Region One, 2,4-D presents the highest cancer risk based on animal feeding studies. However, this risk is considered to be very low. Again, chemical applicators would be at the highest risk. The Lolo National Forest EIS (1989) states that "although human carcinogenic risk levels for these herbicides appear to be quite low, scientific uncertainty still exists regarding the exact level of these risks." FWP employees are required to wear the appropriate protective equipment during applications to minimize their exposure.

Many chemical applications have been contracted out to private, commercial applicators in the past. The number of contracts would likely increase in the future under Alternatives 1,2, and 3. This would minimize exposure to FWP employees, but commercial applicators will then be increasing their exposure. However, commercial applicators usually have the ability to apply chemicals more safely because of their significant investment in safety equipment and clothing.

2,4-D is a chlorophenol-type herbicide in the dioxin family of chemicals. The isomers of dioxin vary in their toxicity but the "highly toxic dioxin" isomer found in 2,4,5-T is not found in 2,4-D (Linnainmaa 1983). The Canadian government appears to have avoided this issue by requiring that 2,4-D formulations have less than 10 parts per billion of any dioxin isomer. This has effectively left only the amine formulation of 2,4-D on the Canadian market (Tosine 1983).

Inert ingredients in herbicide formulations include surfactant, emulsifiers, diluents, and other additives. These have come under scrutiny for possible toxicological and carcinogenic effects. This concept has not been extensively researched. As of 1990, the EPA had listed about 115 inert substances which require further regulation or testing. Acute toxic effects of the POEA family of chemicals, a surfactant used in the Roundup® formulation of glyphosate, have been debated (O'Brien 1990). Furthermore, synergism, the possible interaction of herbicide ingredients with other chemicals in the environment (pollutants and other herbicides), has not been well studied.

This discussion points out the need for caution in the use of all herbicides, especially 2,4-D. However, this Alternative does not include any provisions for safeguarding other resources beyond that specified on chemical label instructions.

Established biocontrol agent populations would not impact human health. However, biocontrol agents are not expected to keep pace with the rate of spread of many noxious weed species and weed populations would likely expand. The spread and increase of noxious weeds populations may increase pollen levels and subsequently exacerbate human allergies.

Alternative 2 (Integrated Noxious Weed Management) also incorporates chemical methods, but such applications would be conducted with safeguards and consideration for other resource values. Such considerations would help diminish concern over exposure routes, bioaccumulation, carcinogenic properties, and chemical additives. Furthermore, chemical treatments would only be used in situations where other methods (biological, cultural, or manual) are not feasible. These non-chemical methods are not expected to have significant impacts on human health. Cultural methods may temporarily increase dust or pollen levels in the air, aggravating allergies. Manual and cultural methods pose some risk to workers. These risks include injuries typical of occupations requiring moderate to heavy physical labor such as lifting, pulling, walking over rough terrain, and operating equipment that may result in extremity injuries. These risks would be kept to a minimum if Occupational Safety and Health Administration work practices and standards are followed.

Under Alternative 3 (Chemical Methods), chemical applications would be the only means of treating noxious weeds. As such, this Alternative presents similar risks to human health as those discussed under Alternative 1. However, Alternative 3 poses the highest risk to human health because of the increased exposure to chemicals for both the public and chemical applicators as a result of the increased frequency of application and increased volume of chemicals introduced into the environment.

Alternative 4 (Non-Chemical) poses the lowest risk to human health because the most risky method of weed treatment (chemical) is not incorporated. The impacts of non-chemical treatment methods are highlighted in the discussion of Alternative 2. Human allergies may be exacerbated by dust from cultural methods or from elevated pollen levels as non-chemical methods may not keep noxious weeds in check as effectively as other Alternatives.

Site Specific

Site specific impacts to human health would be expected at those sites heavily visited by the public, where public use is concentrated such as in campgrounds, and where the intensity of noxious weed treatments is most pronounced. High public use is noted at the following sites: Flathead Lake State Park, Lone Pine State Park, and Fishing Access Sites close to population centers. Moreover, public use at these sites is often concentrated near campgrounds, picnic facilities, restrooms, and boat ramps in the case of Fishing Access Sites.

Those FWP properties with established noxious weed populations require the most intense treatments. Therefore, the potential impacts of exposure to chemicals for the public as well as for FWP employees is elevated. These properties are: Ninepipe Wildlife Management Area, Flathead Lake and Lone Pine State Parks, and Flatiron Ridge, Kokanee Bend, and Pressentine Fishing Access Sites.

3.9 Cultural and Historic Resources

3.9.1 Current Conditions

Cultural resources are important to this and future generations. Cultural resources under the care of FWP are typically associated with, but not exclusive to, the State Park System. In Region One, such cultural resources are preserved at Flathead Lake State Park. Homesteaded in the early 1900's, Wild Horse Island preserves a typical homesteader's quarters, barn, root cellar and orchard. Unfortunately, these cultural and historic sites are part of a greater landscape in which noxious weeds occur. All of these sites have noxious weeds growing within FWP property boundaries. The primary species are spotted knapweed and Canada thistle. Other "weedy" exotic plant species which are not designated as "noxious" do occur. The presence of noxious weeds may not directly threaten or harm a cultural or historic resource. However, their presence may affect the integrity of the flora and a visitor's perception or interpretation of the site.

3.9.2 Analysis of Potential Impacts

General

As described previously under recreation, the presence of noxious weeds may impair one's enjoyment of a site from an aesthetic point of view. In the case of cultural and/or historic sites, the impairment may be more serious. Where noxious weeds occur in a setting where cultural/historic resources are preserved, the treatment of noxious weeds may adversely affect the resource, depending on the method employed. In general, cultural methods, involving hard-edged tools or implements would have a greater potential for impact than the other methods. Tilling, reseeding, or mowing activities could damage or destroy surface and/or subsurface artifacts. As each cultural/historic resource within the State Park System is unique, the impacts of the various noxious weed management Alternatives are discussed on a site-specific basis.

Site Specific

Alternative 1 (Status Quo) is the continuation of previous noxious weed treatment efforts by chemical or biological means. This Alternative would not incorporate safeguards for other resource values. As such, cultural and historical resources may be impacted by a lack of consideration for their value during the planning stages of noxious weed treatment. Also, under this Alternative, newly-established noxious weeds would likely proliferate. Increases in noxious weeds would negatively affect the cultural and/or historic interpretation of a site. Chemical treatments could impact non-target vegetation. This would be especially detrimental at sites where the vegetation component contributes to the interpretation of the site, as is the case at Wild Horse Island. Cultural/historic sites in Region One would not be impacted by established biocontrol agents.

Alternative 2 (Integrated Noxious Weed Management) blends site treatment needs with resource values and appropriate methods to achieve noxious weed reduction. Possible methods include chemical, biological, manual, and cultural. This is the only Alternative which truly considers the importance of cultural/historic resources and attempts to achieve management objectives without impairing those values. Chemical treatments could impact non-target vegetation. This would be especially detrimental at sites where the vegetation component contributes to the interpretation of the site, such as Wild Horse Island. However, such impacts would be minimized by implementation of the guidelines outlined in the Plan. Cultural/historic sites would not be impacted by the establishment of biocontrol agents or by manual treatment methods. Cultural methods may have significant impacts if not properly planned or carried out. For example, tepee rings could be disrupted by mowing or tilling activities. But in requiring consideration of all site values, the Plan would minimize the possibility of disruption.

Alternative 3 (Chemical Methods) emphasizes the use of chemicals, without consideration of site values and resources. Impacts of chemical treatments under this Alternative would be similar to those described for Alternative 1, only the greater reliance on chemicals would put non-target

vegetation resources at even greater risk. Again, Wild Horse Island could suffer the most serious impacts.

Alternative 4 (Non-Chemical Methods) only permits the use of non-chemical treatments. Cultural/historic sites would not be impacted by the establishment of biocontrol agents or by manual treatment methods. Cultural methods may have significant impacts if not properly planned or carried out. For example, tepee rings could be disrupted by mowing or tilling activities.

3.10 Environmental Review Criteria

Future noxious weed management activities proposed under the Region One Weed Management Plan shall be reviewed in accordance with the following Environmental Review Criteria. In some cases, proposed activities may be determined by FWP to be categorically excluded from further environmental review. Those weed management activities for which the answer to every question below is "yes" are categorically excluded from subsequent review and can proceed under the current documentation. Activities triggering a "no" response to any of the questions below would prompt one of the following:

- 1) modification of the proposed activity to return it within the scope of the environmental analysis contained in this document
- 2) rejection of the proposed activity outright
- 3) initiation of a separate environmental review process for the proposed activity.

DESCRIPTION OF THE PROPOSED ACTIVITY:

ENVIRONMENTAL REVIEW CRITERIA	YES	NO
will the weed management activity be conducted using principles of good judgement?		
will the application of <u>non</u> -aquatic-labelled herbicides to running or standing water be avoided?		
will surfaced waters be buffered from the introduction of excessive amounts of sediment as a result of weed treatment activities?		
will herbicides be mixed and loaded at least 500 feet away from open water?		
will wick applicators be used when applying non-aquatic-labelled herbicides within 50 feet of surface waters?		
will aquatic-labelled herbicides be selected for areas with a sandy or gravelly soil consistency and for areas with water tables within 3 feet?		
will herbicide applications be avoided within 50 feet of high human use areas?		
will herbicide treatment sites be posted prior to and for 24 hours after herbicide application to notify the public that spraying activities will/have occurred?		
will the use of mechanized power equipment (e.g. mowers) be postponed when treatment sites are occupied by the public?		

ENVIRONMENTAL REVIEW CRITERIA	YES	NO
will herbicide manufacturer label instructions be followed at all times?		
will herbicides be applied by licensed applicators?		
will aerial herbicide applications be avoided?		
will herbicide applications be suspended when rainfall is expected within 24 hours, air temperatures exceed 85°F, or wind speed exceeds 8 m.p.h.?		
will herbicide spray equipment deliver a median drop diameter between 200 microns and 800 microns?		
is an Herbicide Emergency Response Plan in place?		
are biological control agents approved for release by the Animal and Plant Health Inspection Service?		
will riparian vegetation and surface water banks be protected from excessive disturbance resulting from weed management activity?		
will adverse impacts to rare, unique, threatened or endangered species resulting from weed treatment activities be avoided?		
will wildlife habitat be protected from excessive deterioration or disturbance during weed treatment activities?		
will Department personnel consider the impact of a weed treatment activity on wildlife populations, diversity, and distribution?		
will cultural/historic resources be protected from physical damage during weed treatment activities?		

ADDITIONAL COMMENTS:

4.0 SUMMARY

Exotic (non-native) weedy plants have been noted in Montana throughout this century. In the past few decades, some species have become significant components of local vegetation communities, actually dominating some sites. The most competitive and aggressive species have been termed "noxious." The Montana Legislature passed the County Noxious Weed Control Act in 1979, making it unlawful for a landowner to permit any noxious weed to propagate or go to seed on his/her land. This legislation also declared 15 non-native plant species as "noxious" and gave counties the authority to add species to the list as was locally necessary. Landowners are, by law, in compliance if a weed management plan has been prepared.

As a landowner of approximately 18,000 acres in northwest Montana, FWP, Region One, is required by local ordinances and state law to manage noxious weeds on its properties. Beyond the legal requirement, it is the mission of the FWP to "...provide for the stewardship of the fish, wildlife, parks, and recreational resources of Montana, while contributing to the quality of life for the present and future generations." Part of being a responsible steward of land resources involves the treatment of noxious weeds. FWP (Region One) prepared a Noxious Weed and Exotic Vegetation Management Plan which outlines an integrated approach to noxious weed management. The goal is the long-term reduction of weeds. It will be achieved using a variety of methods (including manual, cultural, biological, and chemical) in which a particular technique is matched with the treatment needs of a specific weed population, while considering and protecting the integrity of all resource values on a particular site.

In accordance with the Montana Environmental Policy Act, FWP must examine and identify potential impacts of its actions on the natural and human environment. This Draft Environmental Assessment describes the direct, indirect, and cumulative effects of the proposed implementation of the Weed Management Plan. The Environmental Assessment also presents Alternatives to the implementation of the Plan and describes potential impacts of these Alternatives. The Alternatives are:

Alternative 1 (Status Quo): noxious weed management activities would continue over the course set since passage of the County Noxious Weed Control Act, namely chemical treatments without any particular safeguards beyond those indicated in the manufacturer label instructions. Biological, manual, and cultural methods were also employed to a certain degree. However, no particular effort was made to plan or coordinate weed management efforts with other resource programs. This Alternative does not incorporate provisions for other resource values.

Alternative 2 (Integrated Noxious Weed Management): implementation of the Noxious Weed Management Plan which describes an approach integrating the management of noxious weeds with site resource values to achieve adequate weed management without negatively affecting other resources. Treatment methods would include manual, chemical, biological, and cultural.

Alternative 3 (Chemical Methods): emphasis on chemical treatment methods for all noxious weed management actions. No other methods would be used, with the exception of those biological control agents already introduced.

Alternative 4 (Non-Chemical Methods): emphasis on manual, cultural, and biological treatment methods for all noxious weed management actions. No chemical methods would be considered.

The preferred Alternative is 2, which would implement the Noxious Weed and Exotic Vegetation Management Plan. Potential impacts to the natural and human environments are evaluated. Resources considered during the evaluation were:

air quality	soils and geology	groundwater
surface water	fish and wildlife	vegetation
recreation	human health	cultural/historic resources

Table 12 summarizes and compares potential impacts resulting from each Alternative.

Table 12. Summary and comparison of environmental impacts resulting from the implementation of each Alternative.

	Alternative 1 No Action	Alternative 2 Integrated Approach	Alternative 3 Herbicide Approach	Alternative 4 No Chemical Approach
Air Quality	No significant effects.	Short-term, minor effects due to chemical vapor, noise, odors, dust or smoke in the project vicinity.	Short-term, minor effects due to chemical vapor, noise, or odors; potential greater than under Alternative 2.	Smoke or dust would increase particulate levels.
Soils	Potential to reduce total plant cover leading to increased erosion.	Reduced potential for contamination due to greater flexibility in choosing treatment methods.	Possible soil contamination due to lack of flexibility in choosing treatment methods.	Manual and mechanical treatments may cause erosion and losses in productivity.
Water Quality	Low potential for surface or ground water contamination.	Reduced potential for contamination due to greater flexibility in choosing treatment methods.	Greatest potential for chemical contamination to surface/ground waters.	Low potential for sedimentation if manual or mechanical treatments result in erosion.
Wildlife	Reduced wildlife habitat quality due to noxious weed invasion	Reduced potential for negative impacts with mitigative measures and management guidelines.	Greatest potential for chemical ingestion and dermal exposure to wildlife species.	Reduced wildlife habitat quality due to noxious weed invasion.
Fisheries	Low potential for sedimentation increases.	Minimal impacts expected due to treatment flexibility and implementation of mitigation measures.	Greatest potential to adversely affect fisheries.	Potential for increased sedimentation may adversely affect habitat quality.
Vegetation	Decline of desirable plant communities due to weed competition.	Lowest potential for impacts to native/domestic plant damage through mitigation.	Greatest potential for adverse impacts to non-target vegetation.	Some adverse effects expected to non-target vegetation due to plowing.
Recreation	Greatest potential to reduce recreation values.	Allows flexibility in treatment methods, while protecting ecological and aesthetic qualities.	Greatest potential to arrest encroachment or establishment of weeds.	Long-term physical removal of weeds required at each infested site.
Human Health	Increased potential for allergies, respiratory or skin ailments from weeds.	Possibility of exposure to chemicals or personal injury from equipment.	Increased possibility of exposure to chemicals.	Greatest potential for physical injury and respiratory ailments from smoke or dust.

5.0 LIST OF CONTRIBUTORS AND REVIEWERS

Dan Vincent	Region One, Regional Supervisor,
Harvey Nyberg	Region One, Regional Wildlife Manager
Carolyn Sime	Region One, Wildlife Biologist
Dan Casey	Region One, Wildlife Biologist
Jim Vashro	Region One, Regional Fisheries Manager
Dave Conklin	Region One, Regional Parks Manager
Jerry Sawyer	Region One, Parks Manager
Marty Watkins	Region One, Parks Manager
Merl Phillips	Region One, Park Maintenance Supervisor
Brian Marotz	Region One, Fisheries Program Officer
Mary Ellen McDonald	Program Officer, Parks Division
Jeff Erickson	Outdoor Recreation Planner, Parks Division
Steve Knapp	Habitat Bureau Chief, Wildlife Division
Thurston Dotson	Hatchery Bureau Chief, Fisheries Division
Mike Korn	Landowner-Sportsman Coordinator, Field Services Division
John Mundinger	Administrative Officer, Department Management
Bridgitt Erickson	Legal Counsel, Department Management

6.0 REFERENCES

- Achuff, P.L. and L. A. Schassberger. 1991. Weeds and rare native plants. Symposium, Montana Academy of Sciences. Billings, Montana.
- Audus, L.J. 1976. Herbicides, physiology, biochemistry, ecology. Editor. Second edition, vol. 2. Academic Press, Inc. London, England.
- Beaverhead National Forest. 1987. Environmental impact statement for noxious weeds and poisonous plant control. U.S. Department of Agriculture, Beaverhead National Forest. Dillon, Montana. 58pp. + appendices.
- Bergeron, D.C., C. Jones, D.L. Genter, and D. Sullivan. 1992. P.D. Skaar's Montana bird distribution, fourth edition. Special Publication, No. 2. Montana Natural Heritage Program, Helena. 116pp.
- Bureau of Land Management. 1985. Proposed Northwest Area noxious weed control program final environmental impact statement. U.S. Department of the Interior, Bureau of Land Management. Oregon State Office, Portland.
- Bureau of Land Management. 1987. Supplement to the Northwest Area noxious weed control program final environmental impact statement. U.S. Department of the Interior, Bureau of Land Management. Oregon State Office, Portland.
- Colby, S.R., E.R. Hill, N.E. Humburg, L.M. Kitchen, R.G. Lym, W.J. McAvoy, and R. Prasad. 1989. Herbicide handbook. 6th edition. Weed Science Society of America. Champaign, Illinois.
- Deluca, T., J. Larson, L. Torma, and G. Algard. 1989. A survey of pesticide residues in ground in Montana. Montana Department of Agriculture, Environmental Management Division. Helena, Montana.
- Environmental Protection Agency. 1983, 1986, and 1988. Pesticide fact sheets: 2,4-D, picloram, metsulfuron, and dicamba.
- Environmental Protection Agency. 1986. Quality criteria for water with updates (Gold Book). Publication 44015-86-001.
- Environmental Protection Agency. 1990. Office of Drinking Water. Drinking water regulations and health advisories.

- Evans, J.O. and D. Dvseja. 1973. Herbicide contamination of surface runoff water. Utah State University, Logan. Office of Research and Monitoring, U.S. Environmental Protection Agency, Washington, D.C. EPA-R2-73-266. 99pp.
- Gallatin National Forest. 1987. Noxious weed control final environmental impact statement. U.S. Department of Agriculture, Gallatin National Forest. Bozeman, Montana. 35pp. + appendices.
- Holton, G.D. 1990. A field guide to Montana fishes. Montana Department of Fish, Wildlife and Parks. 104pp.
- Lacey, C. 1991. Groundwater and agrichemical protection pilot project. Unpublished report and maps prepared for Lewis & Clark County Conservation District.
- Lavelle, D. 1988. Use and preference of spotted knapweed by elk and mule deer on two winter ranges in western Montana. Master's thesis, University of Montana, Missoula. 64pp.
- Lessica, P. 1993. Botanist. Personal communications with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Lessica, P. and J.S. Shelly. 1991. The effect of the introduced weed (*Centaurea maculosa*) on *Arabis fecunda*, a threatened Montana endemic. Unpubl. report. Montana Natural Heritage Program, Helena. 19pp. + appendices.
- Levine, C. 1993. Ecosystems Management Section, Water Quality Bureau, Montana Department of Health and Environmental Sciences. Personal communications with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Linnainmaa, K. 1983. Non-mutagenicity of phenoxy acid herbicides 2,4-dichlorophenoxyacetic acid [2,4-D] and 4-methyl-2-chlorophenoxyacetic acid [MCPA]. Pages 385-395 in Choudhary et al. Chlorinated dioxins and dibenzofurans in the total environment. Butterworth Publishers. Woburn, Massachusetts.
- Littlefield, J. 1993. Insect Quarantine Lab, campus of Montana State University. Personal communications with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Lolo National Forest. 1989. Noxious weed management draft environmental impact statement, amendment to the Lolo National Forest Plan. U.S. Department of Agriculture, Lolo National Forest. Missoula, Montana.
- Lolo National Forest. 1991. Noxious weed management final environmental impact statement, amendment to the Lolo National Forest Plan. U.S. Department of Agriculture, Lolo National Forest. Missoula, Montana.

- McKenna, D. 1993. Associate Research Hydrogeologist, Montana Bureau of Mines and Geology. Personal communications with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Monnig, E. 1988. Human health risk assessment of herbicide applications to control noxious weeds and poisonous plants in the Northern Region. U.S. Department of Agriculture Forest Service, Northern Region RPT 88-9. Missoula, Montana. 218pp.
- Montagne, C., L.S. Munn, G.A. Nielsen, J.W. Rodgers, and H.E. Hunter. 1982. Soils of Montana. Montana Agriculture Experiment Station Bulletin 744, Montana State University, Bozeman. 95pp.
- Montana Bald Eagle Working Group. 1991. Habitat management guide for bald eagles in northwestern Montana. 29pp.
- Montana Department of Agriculture. 1991. Draft noxious weed trust fund programmatic environmental impact statement. Helena, Montana.
- Montana Department of Fish, Wildlife and Parks. 1983. Weed control program for lands managed by the Montana Department of Fish, Wildlife and Parks. 4pp.
- Montana Department of Fish, Wildlife and Parks. 1992. Creating a Vision for the Future. Helena, Montana.
- Montana Department of Fish, Wildlife and Parks. 1993. Draft weed management plan for Region 2. Prepared by Land and Water Consulting Inc. Missoula, Montana. 34pp. + appendices.
- Montana Department of Fish, Wildlife and Parks. 1993. Draft environmental analysis for weed management for Region 2. Prepared by Land and Water Consulting Inc. Missoula, Montana. 29pp. + appendices.
- Montana Department of Fish, Wildlife and Parks. 1993. Draft integrated noxious weed management plan for Region 8. Prepared by OEA Research Inc. Helena, Montana. 78pp. + appendices.
- Montana Department of Fish Wildlife and Parks. 1993. Draft environmental assessment of the integrated noxious weed management plan for Region 8. Prepared by OEA Research Inc. Helena, Montana. 56pp. + appendices.
- Montana Department of Fish, Wildlife and Parks. 1994. Draft noxious weed management plan, Region 3. Montana Department of Fish, Wildlife and Parks, Bozeman. 78pp. + appendices.

- Montana Department of Health and Environmental Sciences. 1992. Montana water quality: the Montana 305(b) report. Ecosystems Management Section, Water Quality Bureau, Environmental Sciences Division. 42pp.
- Montana Natural Heritage Program. 1993. Database search for Species of Special Concern and threatened/endangered species in the vicinity of Montana Department of Fish, Wildlife and Parks properties in Region 3.
- National Research Council of Canada. 1977. Picloram: the effects of its use as a herbicide on environmental quality. Pages 34-36 in NRCC No. 13684. Associated Committee on Scientific Criteria for Environmental Quality. Quebec, Canada.
- Nelson, J., D. Wichman, P. Fay, and J. Connor. 1990. Montana weed control guide. Montana State University Extension Publication EB-23. Bozeman, Montana. 113pp.
- O'Brien, M. 1990. Roundup, Vision, POEA, and 1,4,dioxane: why full formulations are the problem. Journal of Pesticide Reform 9(4):14-15.
- Pence, D. 1993. Beaverhead National Forest. Personal communications with Carolyn Sime, Department of Fish, Wildlife and Parks.
- Phillips, G. 1993. Pollution Control Biologist and Water Quality Specialist for the Montana Department of Fish, Wildlife and Parks. Personal communications with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Rice, P.M., D.J. Bedunah, C.E. Carlson. 1992. Plant community diversity after herbicide control of spotted knapweed. U.S. Department of Agriculture Forest Service Intermountain Forest and Range Experiment Station. Research paper INT 460. Ogden, Utah. 6pp.
- Sassman, J., R. Pienta, M. Jacobs and J. Cioffi. 1984. Pesticide background statements. Volume 1, herbicides. Agriculture Handbook No. 633. U.S. Government Printing Office. Washington, D.C.
- Smith, Mark. 1993. Ecosystems Management Section, Water Quality Bureau, Montana Department of Health and Environmental Sciences. Personal communications with Carolyn Sime, Montana Department of Fish, Wildlife and Parks.
- Soucek, P. 1993. Gallatin National Forest. Personal communications with Carolyn Sime, Department of Fish, Wildlife and Parks.
- Sparks, D.L. 1989. Kinetics of soil chemical processes. Academic Press, Inc. San Diego, California.

- Story, J. 1989. Biological control: its potential for progress toward knapweed control in the next 10 years. Pages 237-238 in P.K. Fay and J.R. Lacey, editors. Proceedings of the 1989 knapweed symposium. Montana State University, Bozeman.
- Thompson, L. 1982. Distribution of Montana amphibians, reptiles, and mammals. Montana Audubon Council. Helena, Montana. 24pp.
- Tosine, H. 1983. Dioxins: a Canadian perspective. Pages 3-14 in Choudhary et al. Chlorinated dioxins and dibenzofurans in the total environment. Butterworth Publishers. Woburn, Massachusetts.
- U.S. Department of Agriculture. 1984. Pesticide background statements. Volume 1, herbicides. Agricultural Handbook 633. Washington, D.C.
- U.S. Department of Agriculture. 1988. Draft environmental impact statement for vegetation management in the Coastal Plain/Piedmont. Management Bulletin R8-MB 15. Southern Region, Atlanta, Georgia.
- U.S. Fish and Wildlife Service. 1980. Handbook of acute toxicity of chemicals to fish and aquatic invertebrates. Resource Publication No. 137. U.S. Department of the Interior, Fish and Wildlife Service. Washington, D.C.
- Watson, V.J., P.M. Rice, and E.C. Monnig. 1989. Environmental fate of picloram used for roadside weed control. Journal of Environmental Quality 18(2):198-205.
- Workman, D. 1992. Fisheries Manager, Montana Department of Fish, Wildlife and Parks. Personal communications with OEA Research Inc.

7.0 APPENDICES

APPENDIX A Herbicide Properties and Efficacy on Target Noxious Weed Species

APPENDIX B Fish and Wildlife Species Known to Occur in Region One of the Montana
Department of Fish, Wildlife and Parks

APPENDIX A

HERBICIDE EFFICACY ON STATE-DECLARED NOXIOUS WEED SPECIES

PROPERTIES OF COMMONLY USED HERBICIDES

Herbicide Efficacy on State-Declared Noxious Weed Species

	Efficacy Rating*				
	Excellent	Good	Fair	Poor	Insufficient data
Category 1					
Canada thistle	D,P,C	G	2,M,A	—	—
Dalmatian toadflax	—	D,P,G	2	—	—
Field bindweed	—	D,P,G,2	M	C	—
Leafy spurge	—	P	D,G	2	—
Diffuse knapweed	2,D,P	—	—	—	G
Russian knapweed	—	D,P,G,C	M	2	A
Spotted knapweed	D,P,G,C	—	2***	2	—
St. Johnswort (goatweed)	—	P	—	2	D,G
Whitetop (hoary cress)	—	D,G	2,A	P,M	—
Category 2					
Dyers woad	D**	2,P,G	—	—	—
Purple loosestrife	G	2,D	—	—	—
Sulfur cinquefoil	D,P,G	2	—	—	—
Category 3					
Rush skeletonweed	P	D	2	—	G
Yellow starthistle	—	—	—	—	—
Common crupina	—	—	—	—	—

- * Control ratings for the herbicides are as follows:

Excellent = >95% of weed population killed by single treatment

Good = one treatment per year maintains 85-94% suppression

Fair = 60-84% of weed population killed by single treatment

Poor = 10-59% of weed population killed by single treatment

Insufficient data = insufficient data exists to determine effectiveness

- ** C=clpyralid, D=dicamba, G=glyphosate, P=picloram, A=metsulfuron, M=MCPA, 2 = 2,4-D

- *** Data sources conflict on 2,4-D efficacy for spotted knapweed; Data ranges from poor to excellent.

References: USDI-Bureau of Land Management 1985; Nelson et al 1990; McCone 1991; Johnson 1991.

Properties of Commonly Used Herbicides

Active Ingredient (Trade Name®)	Selectivity	Surface Potential*	Soil Leaching Potential**	Persistence in Soil /Soil Half-life	Primary Degradation	Toxicity			
						Aquatic Organisms †			
						Trout	Bluegill	Rat	Animal ††
2,4-D (amine)	broadleaves	medium	medium/1	≤1 mo/10 days	microbial	100	168	no effect	2000
(ester)	broadleaves	medium	medium/1	≤1 mo/10 days	microbial	1.2	100	620	2025
picloram (Tordon 22K)	broadleaves	low	high/1	1-5 year/90 days	microbial	22.2	8.2	10,300	>2000
glyphosate (Rodeo)	non-selective	high	low/2	1 mo/30 days	microbial	1000	1000	>5000	...
(Roundup)	non-selective	high	low/2	1 mo/30 days	microbial	110	14	5400	...
dicamba (BANVEL)	broadleaves	low	high/1	3-12 mo/14 days	microbial	35	130	2629	>2510
clopyralid (Stinger)	broadleaves	low	high/1	1-4 mo/20 days	microbial	103.5	125.4	>5000	>4640
metsulfuron (Ally)	non-selective	medium	high/1	up to 34 mo/ 7-42 days	microbial	>150	>150	>5000	>2510
MCPA (amine)	broadleaves	low	high/1	1-6 mo/14 days	microbial	relatively	relatively	relatively	relatively
* Surface Potential - relative tendency to move with run-off.									
** Leaching Potential - relative tendency to dissolve in soil water so can move through soil profile/herbicide affinity to soil: 1-readily adsorbed;									
† not readily adsorbed									
‡ 96 hour LC-50 - the lethal pesticide concentration rate at which 50% of test animals will be killed: <1 mg/kg = highly toxic; 1-10 mg/kg = toxic;									
>10 mg/kg = slightly toxic. 48 hour LC-50 for dicamba. 1 mg/kg = 1 part per million (ppm).									
†† LD-50 - the dosage of toxicant, expressed in milligrams of toxicant/kg of animal body weight, required to kill 50% of the test organism in a specific									
test situation.									
References: USDA-Forest Service 1984; Weed Science Society of America 1989; US-EPA 1983, 1988; Lolo N.F. 1989; Whilson et al. 1989									

APPENDIX B

FISH AND WILDLIFE SPECIES RECORDED IN LATILONGS CORRESPONDING WITH Montana Fish, Wildlife & Parks Region One. OCCURRENCE RECORDS INDICATE AN OBSERVATION DURING THE BREEDING SEASON, WINTER SEASON, OR DURING MIGRATION.

Bird species recorded in latilongs 1, 2, 3, 13, 14, and 15. This list is based on the species and sequence from Montana Bird Distribution, 4th edition, published in 1992 by the Montana Audubon Council and the Montana Natural Heritage Program.

Region One BIRD LIST

Bird species recorded in administrative Region One of Montana Fish, Wildlife & Parks. This list is based on the species and sequence in Montana Bird Distribution, 4th Edition (Bergeron et al. 1992), with minor modifications based on more recent data (Casey, unpubl.). All species known to have occurred in latilong blocks 1,2,3,13,14, and 15 are listed except for those known to have occurred only outside the boundaries of Region One (e.g. east of the Continental Divide in latilong block 3). *Species in italics have been seen <20 times in Montana, or are quite rare in Region One.*

SPECIES	Status ^{a/}	SPECIES	Status
<i>Red-throated Loon</i>	<i>t</i>	Cinnamon Teal	B
<i>Pacific Loon</i>	<i>t</i>	Northern Shoveler	B
Common Loon	BW	Gadwall	BW
<i>Yellow-billed Loon</i>	<i>t</i>	Eurasian Wigeon	<i>t</i>
Pied-billed Grebe	BW	American Wigeon	BW
Horned Grebe	BW	Canvasback	BW
Red-necked Grebe	B	Redhead	BW
Eared Grebe	B	Ring-necked Duck	BW
Western Grebe	B	Greater Scaup	<i>t</i>
Clark's Grebe	<i>b</i>	Lesser Scaup	BW
American White Pelican	<i>t</i>	Harlequin Duck	B
Double-crested Cormorant	B	Oldsquaw	<i>t</i>
American Bittern	B	<i>Surf Scoter</i>	<i>t</i>
Great Blue Heron	BW	White-winged Scoter	<i>t</i>
<i>Great Egret</i>	<i>t</i>	Common Goldeneye	BW
<i>Snowy Egret</i>	<i>t</i>	Barrow's Goldeneye	BW
<i>Cattle Egret</i>	<i>t</i>	Bufflehead	BW
<i>Green-backed Heron</i>	<i>t</i>	Hooded Merganser	BW
<i>Black-crowned Night-heron</i>	<i>B</i>	Common Merganser	BW
<i>White-faced Ibis</i>	<i>t</i>	Red-breasted Merganser	<i>t</i>
Tundra Swan	tW	Ruddy Duck	BW
Trumpeter Swan	<i>b</i>	Turkey Vulture	B
<i>Mute Swan</i>	<i>t</i>	Osprey	B
Greater White-fronted Goose	<i>t</i>	Bald Eagle	BW
Snow Goose	<i>t</i>	Northern Harrier	BW
Ross' Goose	<i>t</i>	Sharp-shinned Hawk	BW
<i>Brant</i>	<i>t</i>	Cooper's Hawk	BW
Canada Goose	BW	Northern Goshawk	BW
Wood Duck	BW	<i>Red-shouldered Hawk</i>	<i>t</i>
Green-winged Teal	BW	<i>Broad-winged Hawk</i>	<i>t</i>
<i>American Black Duck</i>	<i>t</i>	Swainson's Hawk	B
Mallard	BW	Red-tailed Hawk	BW
Northern Pintail	BW	Ferruginous Hawk	B
<i>Garganey</i>	<i>t</i>	Rough-legged Hawk	tW
Blue-winged Teal	B	Golden Eagle	BW

SPECIES	Status	SPECIES	Status
American Kestrel	BW	Marbled Godwit	b
Merlin	BW	<i>Ruddy Turnstone</i>	<i>t</i>
Peregrine Falcon	BW	<i>Black Turnstone</i>	<i>t</i>
Gyr Falcon	tW	Sanderling	t
Prairie Falcon	BW	Semipalmated Sandpiper	t
Gray Partridge	BW	Western Sandpiper	t
<i>Chukar</i>	<i>BW</i>	Least Sandpiper	t
Ring-necked Pheasant	BW	<i>White-rumped Sandpiper</i>	<i>t</i>
Spruce Grouse	BW	Baird's Sandpiper	t
Blue Grouse	BW	Pectoral Sandpiper	t
White-tailed Ptarmigan	BW	Dunlin	t
Ruffed Grouse	BW	Stilt Sandpiper	t
Sharp-tailed Grouse	BW	<i>Buff-breasted Sandpiper</i>	<i>t</i>
Wild Turkey	BW	<i>Short-billed Dowitcher</i>	<i>t</i>
<i>Northern Bobwhite</i>	<i>b</i>	Long-billed Dowitcher	t
<i>Yellow Rail</i>	<i>t</i>	Common Snipe	BW
Virginia Rail	B	<i>American Woodcock</i>	<i>t</i>
Sora	B	Wilson's Phalarope	B
<i>Common Moorhen</i>	<i>t</i>	Red-necked Phalarope	t
American Coot	BW	<i>Parasitic Jaeger</i>	<i>t</i>
Sandhill Crane	B	<i>Long-tailed Jaeger</i>	<i>t</i>
Black-bellied Plover	t	Franklin's Gull	b
American Golden-plover	t	Bonaparte's Gull	t
Semipalmated Plover	t	<i>Mew Gull</i>	<i>t</i>
Killdeer	BW	Ring-billed Gull	BW
<i>Mountain Plover</i>	<i>t</i>	California Gull	BW
<i>Black-necked Stilt</i>	<i>b</i>	Herring Gull	tW
American Avocet	B	<i>Thayer's Gull</i>	<i>t</i>
Greater Yellowlegs	t	Glaucous Gull	t
Lesser Yellowlegs	t	<i>Glaucous-winged Gull</i>	<i>t</i>
Solitary Sandpiper	t	<i>Black-legged Kittiwake</i>	<i>t</i>
Willet	b	<i>Sabine's Gull</i>	<i>t</i>
Spotted Sandpiper	B	Caspian Tern	B
Upland Sandpiper	B	Common Tern	B
Whimbrel	t	Forster's Tern	B
Long-billed Curlew	B	Black Tern	B
<i>Hudsonian Godwit</i>	<i>t</i>	<i>Marbled Murrelet</i>	<i>t</i>

SPECIES	Status	SPECIES	Status
<i>Ancient Murrelet</i>	<i>t</i>	Downy Woodpecker	BW
Rock Dove	BW	Hairy Woodpecker	BW
<i>Band-tailed Pigeon</i>	<i>t</i>	<i>White-headed Woodpecker</i>	<i>t</i>
Mourning Dove	BW	Three-toed Woodpecker	BW
<i>Black-billed Cuckoo</i>	<i>b</i>	Black-backed Woodpecker	BW
<i>Yellow-billed Cuckoo</i>	<i>t</i>	Northern Flicker	BW
<i>Common Barn-owl</i>	<i>t</i>	Pileated Woodpecker	BW
Flammulated Owl	<i>b</i>	Olive-sided Flycatcher	B
Western Screech-owl	<i>b</i>	Western Wood-pewee	B
Great Horned Owl	BW	<i>Alder Flycatcher</i>	<i>t</i>
Snowy Owl	tW	Willow Flycatcher	B
Northern Hawk-owl	BW	Least Flycatcher	<i>b</i>
Northern Pygmy-owl	BW	Hammond's Flycatcher	B
Burrowing Owl	<i>b</i>	Dusky Flycatcher	B
Barred Owl	BW	Cordilleran Flycatcher	B
Great Gray Owl	BW	<i>Eastern Phoebe</i>	<i>t</i>
Long-eared Owl	BW	Say's Phoebe	B
Short-eared Owl	BW	<i>Ash-throated Flycatcher</i>	<i>t</i>
Boreal Owl	B	Western Kingbird	B
Northern Saw-whet Owl	BW	Eastern Kingbird	B
Common Nighthawk	B	<i>Scissor-tailed Flycatcher</i>	<i>t</i>
Common Poorwill	<i>b</i>	Horned Lark	BW
<i>Whip-poor-will</i>	<i>t</i>	Tree Swallow	B
Black Swift	B	Violet-green Swallow	B
Vaux's Swift	B	Northern Rough-winged Swallow	B
White-throated Swift	<i>b</i>	Bank Swallow	B
<i>Ruby-throated Hummingbird</i>	<i>t</i>	Cliff Swallow	B
Black-chinned Hummingbird	B	Barn Swallow	B
<i>Anna's Hummingbird</i>	<i>t</i>	Gray Jay	BW
Calliope Hummingbird	B	Steller's Jay	BW
<i>Broad-tailed Hummingbird</i>	<i>t</i>	Blue Jay	tW
Rufous Hummingbird	B	<i>Pinyon Jay</i>	<i>t</i>
Belted Kingfisher	BW	Clark's Nutcracker	BW
Lewis' Woodpecker	BW	Black-billed Magpie	BW
<i>Red-headed Woodpecker</i>	<i>b</i>	American Crow	BW
Red-naped Sapsucker	B	Common Raven	BW
Williamson's Sapsucker	B	Black-capped Chickadee	BW

SPECIES	Status	SPECIES	Status
Mountain Chickadee	BW	Red-eyed Vireo	B
Boreal Chickadee	BW	Tennessee Warbler	B
Chestnut-backed Chickadee	BW	Orange-crowned Warbler	B
Red-breasted Nuthatch	BW	Nashville Warbler	B
White-breasted Nuthatch	BW	Yellow Warbler	B
Pygmy Nuthatch	BW	<i>Chestnut-sided Warbler</i>	<i>t</i>
Brown Creeper	BW	<i>Magnolia Warbler</i>	<i>t</i>
Rock Wren	B	<i>Cape May Warbler</i>	<i>t</i>
Canyon Wren	BW	Yellow-rumped Warbler	BW
House Wren	B	Townsend's Warbler	B
Winter Wren	BW	<i>Blackburnian Warbler</i>	<i>t</i>
Marsh Wren	BW	<i>Palm Warbler</i>	<i>t</i>
American Dipper	BW	Blackpoll Warbler	<i>t</i>
Golden-crowned Kinglet	BW	Black-and-white Warbler	<i>t</i>
Ruby-crowned Kinglet	BW	American Redstart	B
<i>Eastern Bluebird</i>	<i>t</i>	Ovenbird	b
Western Bluebird	B	Northern Waterthrush	B
Mountain Bluebird	B	MacGillivray's Warbler	B
Townsend's Solitaire	BW	Common Yellowthroat	B
Veery	B	Wilson's Warbler	B
Swainson's Thrush	B	<i>Canada Warbler</i>	<i>t</i>
Hermit Thrush	B	Yellow-breasted Chat	B
American Robin	BW	<i>Scarlet Tanager</i>	<i>t</i>
Varied Thrush	BW	Western Tanager	B
Gray Catbird	B	<i>Rose-breasted Grosbeak</i>	<i>t</i>
<i>Northern Mockingbird</i>	<i>t</i>	Black-headed Grosbeak	B
<i>Sage Thrasher</i>	<i>t</i>	Lazuli Bunting	B
<i>Brown Thrasher</i>	<i>t</i>	<i>Indigo Bunting</i>	<i>t</i>
American Pipit	B	<i>Dickcissel</i>	<i>t</i>
<i>Sprague's Pipit</i>	<i>t</i>	<i>Green-tailed Towhee</i>	<i>t</i>
Bohemian Waxwing	BW	Rufous-sided Towhee	BW
Cedar Waxwing	BW	American Tree Sparrow	Tw
Northern Shrike	tW	Chipping Sparrow	B
Loggerhead Shrike	b	Clay-colored Sparrow	B
European Starling	BW	Brewer's Sparrow	B
Solitary Vireo	B	<i>Field Sparrow</i>	<i>t</i>
Warbling Vireo	B	Vesper Sparrow	B

SPECIES	Status	SPECIES	Status
Lark Sparrow	B	Western Meadowlark	BW
<i>Sage Sparrow</i>	<i>t</i>	Yellow-headed Blackbird	B
<i>Lark Bunting</i>	<i>b</i>	<i>Rusty Blackbird</i>	<i>tW</i>
Savannah Sparrow	B	Brewer's Blackbird	B
<i>Baird's Sparrow</i>	<i>b</i>	Common Grackle	B
Grasshopper Sparrow	B	Brown-headed Cowbird	B
Le Conte's Sparrow	B	Northern Oriole	B
Fox Sparrow	B	<i>Brambling</i>	<i>tW</i>
Song Sparrow	BW	Rosy Finch	BW
Lincoln's Sparrow	B	Pine Grosbeak	BW
<i>Swamp Sparrow</i>	<i>t</i>	<i>Purple Finch</i>	<i>t</i>
White-throated Sparrow	<i>tW</i>	Cassin's Finch	BW
<i>Golden-crowned Sparrow</i>	<i>t</i>	House Finch	BW
White-crowned Sparrow	B	Red Crossbill	BW
Harris' Sparrow	<i>tW</i>	White-winged Crossbill	BW
Dark-eyed Junco	BW	Common Redpoll	<i>tW</i>
Lapland Longspur	<i>tW</i>	<i>Hoary Redpoll</i>	<i>tW</i>
<i>Chestnut-collared Longspur</i>	<i>t</i>	Pine Siskin	BW
Snow Bunting	<i>tW</i>	American Goldfinch	BW
Bobolink	B	Evening Grosbeak	BW
Red-winged Blackbird	BW	House Sparrow	BW

^{a/} Status Codes:

- t Occurs as a transient, vagrant or winter visitor, with no evidence of breeding.
- b Circumstantial evidence of breeding (e.g. singing male in suitable habitat).
- B Confirmed evidence of breeding (e.g. nest with eggs, dependent young).
- W Confirmed overwintering - regular sightings at least one winter (12/15-2/15).

Fish species known to occur in northwest Montana. This list is based on A Field Guide to Montana Fishes by George Holton, published in 1990 by the Montana Department of Fish, Wildlife and Parks.

Black Bullhead
Longnose Sucker
Largescale Sucker
Peamouth
Northern Squawfish
Redside Shiner
Yellowstone Cutthroat Trout
Rainbow Trout
Brown Trout
Lake Trout
Burbot
Largemouth Bass
Smallmouth Bass
Mottled Sculpin
Pumpkinseed
Shorthead Sculpin
Spoonhead Sculpin

White Sturgeon
Lake Whitefish
Kokanee
Pygmy Whitefish
Bull Trout
Golden Trout
Westslope Cutthroat Trout
Mountain Whitefish
Brook Trout
Arctic Grayling
Bluegill
Yellow Perch
Northern Pike
Mosquitofish
Slimy Sculpin
Torrent Sculpin

Amphibian, reptile and mammal species recorded in latilongs 1, 2, 3, 13, 14, and 15. These lists are based on Distribution of Montana Amphibians, Reptiles, and Mammals by Larry S. Thompson, published in 1982 by The Montana Audubon Council.

AMPHIBIANS

Tiger Salamander
Long-Toed Salamander
Rough-skinned Newt
Coeur d'Alene (Van Dyke's) Salamander
Boreal (Western) Toad
Pacific Tree Frog
Leopard Frog

Boreal (western) Toad
Pacific Giant Salamander
Boreal Chorus Frog
Tailed Frog
Boreal Chorus Frog
Spotted Frog
Bullfrog

REPTILES

Western Skink
Painted Turtle
Common Garter Snake
Northern Alligator Lizard
Prairie Rattlesnake

Racer
Bull Snake
Rubber Boa
Western Garter Snake

MAMMALS

Masked Shrew
Vagrant Shrew
Water Shrew
Yuma Myotis
Fringed Myotis
Silver-Haired Bat
Townsend's Big-Eared Bat
Long-legged Myotis
Snowshoe Rabbit
Black-Tailed Jackrabbit
Red-Tailed Chipmunk
Hoary Marmot
Pika
Columbian Ground Squirrel
Red Squirrel
Northern Pocket Gopher
Pygmy Shrew
Deer Mouse
Gapper's Red-Backed Vole
Meadow Vole
Long-Tailed Vole
Water Vole
Muskrat
House Mouse
Porcupine
Gray Wolf
Black Bear
Raccoon
Fisher
Kit Fox
Mink
Badger
Striped Skunk
Mountain Lion
Bobcat
Mule Deer
Moose
Bison
Rocky Mountain Bighorn Sheep

Preble Shrew
Montane Shrew
California Myotis
Little Brown Myotis
Long-Eared Myotis
Big Brown Bat
Small-footed Myotis
Nuttall's Cottontail
White-Tailed Jackrabbit
Least Chipmunk
Yellow-Bellied Marmot
Richardson's Ground Squirrel
Yellow-pine Chipmunk
Golden-Mantled Ground Squirrel
Northern Flying Squirrel
Thirteen-lined Ground Squirrel
Beaver
Bushy-Tailed Woodrat
Heather Vole
Montane Vole
Northern Grasshopper Mouse
Northern Bag lemming
Norway Rat
Western Jumping Mouse
Coyote
Red Fox
Grizzly Bear
Marten
Ermin
Long-Tailed Weasel
Wolverine
Caribou
River Otter
Lynx
Elk
White-Tailed Deer
Pronghorn Antelope
Mountain Goat