AL.2.2000-279

C.2

Weeds *of the Prairies*

Weeds *of the Prairies*

Carol J. Bubar

Ph. D., P. Ag.
Olds College

Susan J. McColl

Ph. D., P. Ag.
University of
Saskatchewan

Linda M. Hall

Ph. D., P. Ag.
Alberta Agriculture, Food
and Rural Development

Published by:

Alberta Agriculture, Food and Rural Development
Publishing Branch
7000 - 113 Street
Edmonton, Alberta
Canada T6H 5T6

Editor: Chris Kaulbars
Graphic Designer: John Gillmore
Layout and Composition: Sherrill Strauss

Copyright © 2000. Her Majesty the Queen in Right of Alberta.
All rights reserved.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical photocopying, recording or otherwise without written permission from the Publishing Branch, Alberta Agriculture, Food and Rural Development.

ISBN 0-7732-6147-8

Contents

Acknowledgements v

Introduction 1

Identifying Weeds 2

Blue Flowered Species

Bluebur 10

Blueweed 12

Flax 14

Ground-ivy 16

Silky Lupine 18

Purple Flowered Species

Bull Thistle 22

Canada Thistle 24

Common Burdock 26

Creeping Bellflower 28

Hound's-tongue 30

Nodding Thistle 32

Purple Loosestrife 34

Russian Knapweed 36

Spotted Knapweed 38

Tall Larkspur 40

Two-grooved Milk-vetch 42

Pink Flowered Species

American Dragonhead 46

Cow Cockle 48

Field Scabious 50

Hemp-nettle 52

Henbit 54

Pale Smartweed 56

Prickly Rose 58

Prostrate Knotweed 60

Showy Milkweed 62

Stork's-bill 64

Water Smartweed 66

White Flowered Species

Baby's-breath 70

Bladder Campion 72

Canada Fleabane 74

Chickweed 76

White Flowered Species – continued

Cleavers	78
Common Pepper-grass	80
Corn Spurry	82
Death Camas	84
Diffuse Knapweed	86
Field Bindweed	88
Hairy Nightshade	90
Narrow-leaved Milk-vetch	92
Night-flowering Catchfly	94
Ox-eye Daisy	96
Pygmyflower	98
Round-leaved Mallow	100
Scentless Chamomile	102
Shepherd's Purse	104
Spotted water-hemlock	106
Stinkweed	108
White Clover	110
White Cockle	112
Wild Buckwheat	114
Wild Caraway	116
Wild Licorice	118
Wild Tomato	120

Yellow Flowered Species

Absinth	124
Argentine Canola	126
Ball Mustard	128
Black Henbane	130

Black Medick	132
Common Groundsel	134
Common Mullein	136
Dandelion	138
Dog Mustard	140
Early Yellow Locoweed	142
Flixweed	144
Goat's-beard	146
Golden-bean	148
Gumweed	150
Leafy Spurge	152
Narrow-leaved Hawk's-beard	154
Orange Hawkweed	156
Pasture Sage	158
Perennial Sow-thistle	160
Pineappleweed	162
Polish Canola	164
Prairie Sunflower	166
Prickly Lettuce	168
Purslane	170
Rough Cinquefoil	172
Spiny Annual Sow-thistle	174
Tall Buttercup	176
Tansy	178
Tumble Mustard	180
Wild Mustard	182
Wild Radish	184
Wood Whitlow-grass	186
Yellow Nut Sedge	188
Yellow Sweet Clover	190
Yellow Toadflax	192

Green Flowered Species

Barley	196
Barnyard Grass	198
Broad-leaved Plantain	200
Cocklebur	202
Downy Brome	204
False Ragweed	206
Field Dock	208
Field Horsetail	210
Foxtail Barley	212
Green Foxtail	214
Kochia	216
Lamb's-quarters	218
Persian Darnel	220
Proso Millet	222
Prostrate Pigweed	224
Quack Grass	226
Redroot Pigweed	228
Russian Thistle	230
Spear-leaved Goosefoot	232
Stinging Nettle	234
Tartary Buckwheat	236
Thyme-leaved Spurge	238
Wheat	240
Wild Oats	242

Glossary 245

Index 251

Organized by Common Name	251
Organized by Scientific Name	259
Organized by Family Name	263

Digitized by the Internet Archive
in 2016

<https://archive.org/details/weedsofprairies00buba>

*T*his book is the work of many photographers, weed scientists and agricultural experts. Written descriptions in the book have been compiled from the observations and research of many botanists and weed scientists. The authors would like to specifically acknowledge a number of individuals.

Acknowledgements

Photographers:

- Valerie Sowiak, AAFRD, Edmonton, Alberta
- Beth Hoar, AAFRD, Edmonton, Alberta
- Linda Hall, AAFRD, Edmonton, Alberta
- Sue McColl, University of Saskatchewan, Saskatoon, Saskatchewan
- Carol Bubar, Olds College, Olds, Alberta
- Susan Weaver, AAFC, Harrow, Ontario
- Doug Derksen, AAFC, Brandon, Manitoba
- Rick Holm, University of Saskatchewan, Saskatoon, Saskatchewan
- Bill Vanden Born, University of Alberta, Edmonton, Alberta
- Ian Morrison, University of Alberta, Edmonton, Alberta
- France Royer, Edmonton, Alberta
- Richard Dickinson, Toronto, Ontario

All have made contributions to the book.

Information on the typical location of weeds was contributed by:

- Dan Cole, AAFRD, Edmonton, Alberta
- Kelly Cooley, MD Pincher Creek, Pincher Creek, Alberta
- Doug Derksen, AAFC, Brandon, Manitoba
- Gary Martin, University of Manitoba, Winnipeg, Manitoba
- John Huffman, AAFRD, Grand Prairie, Alberta
- Vern Harms, University of Saskatchewan, Saskatoon, Saskatchewan
- Peter Regitinig, Roger's Sugar, Taber, Alberta

Weed distribution maps were produced with the able assistance of Stephen Bowkett, Land Resources Unit, Semiarid Prairie Agricultural Research Centre, Swift Current, Saskatchewan and Gordon Thomas, Saskatoon, Saskatchewan – both with Agriculture and Agri-Food Canada.

Introduction

Weeds of the Prairies was developed as a practical field guide for those who need a relatively non-technical aid to identify common weeds across the Canadian prairie provinces.

This book does not contain all the weeds, nor all the native plants that can be found on the prairies. However, it does include over 100 weeds that can be found in agricultural fields, pastures, rangeland, waste areas, gardens and roadsides.

The primary classification of weeds in this book is by flower color, which match the colors seen on the outside of the pages. Within each color group, species are alphabetized by the common name of the weed. Maps showing the per cent of fields where the weed is found are presented for many of the weed species in this book.

Each weed discussed has charts on life cycle and habitat. The life cycle chart describes whether the weed will grow as an annual, biennial and so on. The habitat charts show how likely the weed is to be found growing in certain areas. The relative frequency appears in the number of brush strokes beside the area description. Three strokes beside the term “conventional tillage” means the weed occurs fairly frequently in these areas. The range of brush strokes goes from zero – meaning not a habitat for this weed – to three – meaning a very common habitat.

At the end of the book, there is a complete species list that includes both the common and Latin names of the weed. In addition, we have provided an alphabetized plant family listing with the weeds of each family also alphabetized.

All attempts have been made to gather representative photographs of these weeds at several stages of growth. Where appropriate and available, photographs of infestations, key features and/or similar species are also presented. Descriptions and names were compiled carefully, but in any project of this size, errors can occur.

If you notice any errors or have information or photographs to add, please contact L.M. Hall, Alberta Agriculture, Food and Rural Development, 6903 - 116 Street, Edmonton, AB, T6H 4P2.

Identifying weeds

Identifying a weed can sometimes be a challenge. Knowing what the parts of the weed are called and being able to describe them can certainly help in identification. Here are some drawings of basic weed types: broadleaf weeds and grass weeds. In addition, a number of images here will show leaf types, leaf arrangements and so on to help with identification.

Broadleaf seedling

Broadleaf stem and leaf

Leaf attachment

leaf with petiole

sessile

Leaf arrangement

alternate

basal

opposite

whorled

Leaf type

simple palmate

compound palmate

simple pinnate

compound pinnate

trifoliate

Stipules at stem node

stipules

Leaf shapes

elliptical

lanceolate

linear

oval

ovate

spoon-shaped/
spatulate

Stem with ocrea

ocrea

Leaf margins

divided

finely divided

entire

lobed

scalloped

toothed

Growth habit

erect

twining

prostrate

Rhizome and stolon

rhizome

stolon

Broadleaf flower structure

Pea-type flower

standard

wing petal

keel

Inflorescence types

flower cluster in
leaf axil

involucral bracts

simple umbel

compound umbel

aster family – ray
flower

aster family – disk
flower

aster family – ray
and disk flower

raceme

spike

Grass collar

Grass spikelet with two florets

Blue

Flowered Species

Bluebur

Lappula squarrosa

Other names

Hitch-hiker, stickseed, stickweed

Family

Boraginaceae

Seedling

Cotyledons are oval with short petioles.

They are covered with short hair and have a small “bump” at the tip. The first true leaves emerge one at a time and appear folded in half.

They are narrowly ovate, densely hairy and have a distinct crease down the middle. Some people claim that the seedlings smell like a mouse-infested building. This seedling may be confused with night-flowering catchfly. Catchfly seedlings have a rounder cotyledon tip and true leaves that emerge in pairs. They also lack the crease and odor associated with bluebur seedlings.

Juvenile

Bluebur forms a low-growing rosette of leaves that are more spatulate than in the seedling, but that are also densely hairy and creased in appearance. Stems are erect, hairy and often much-branched. The leaves become reduced in size as they near the top of the plant.

Mature

Tiny, blue flowers are located in racemes with small bracts resembling leaves below each flower. The flowers are similar to those of the popular garden perennial “forget-me-not.” Each flower produces four bur-like seeds with two rows of sharp, hooked prickles. The prickly seeds readily stick to clothing and animal hair.

Key features

Densely hairy, creased leaves; tiny blue flowers; prickly, bur-like seeds

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling

Juvenile: rosette

Mature

Flower: tiny blue

Key feature: prickly seeds

Blueweed

Echium vulgare

Other names

Viper's bugloss, blue devil

Family

Borage *Boraginaceae*

Seedling

Cotyledons are oval and covered with fine hairs. The first true leaf is densely hairy and more elongated than the cotyledons. The leaf has a creased appearance.

Juvenile

Blueweed forms a large, black tap root and a basal rosette in the first year. The basal leaves are stalked, narrowly elongated and covered with stiff, spreading hairs. Stems are upright, and there may be one to several stems per tap root. The stem leaves are sessile and smaller than the rosette leaves. Both the stems and upper leaves are covered with a mixture of short and long stiff hairs. The longer hairs often have conspicuous swollen bases that make the stem appear speckled. Blueweed plants may cause rashes in sensitive individuals, so plants should be handled with care.

Mature

Flowers are typically reddish-purple in bud and turn bright blue after emergence. They are funnel-shaped with protruding stamens and are located in clusters in the axils of upper leaves. Each flower produces four angular, rough, grey to brown seeds.

Key features

Large tap root; dense, stiff hairs with swollen bases; bright blue flowers in clusters

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Juvenile: *rosette*

Infestation

Flower

Key feature: *flowers in clusters*

Flax

Linum usitatissimum

Family

Flax *Linaceae*

Seedling

Cotyledons are oval and slightly fleshy. They are larger than the true leaves, which are linear with pointy tips. The small, sessile leaves are alternate, but the first two may appear opposite.

Juvenile

Flax has a branched tap root and erect growth. There is usually one main stem and irregular, multi-branching at the top of the plant. Small, sessile leaves spiral alternately around the stem. Volunteer flax can be difficult to control in both cereals and broadleaf crops. Since it may remain green longer than the crop, it can interfere with harvesting and subsequent grain storage.

Mature

Flowers form at the ends of the upper branches and though usually blue, they can also be white or pale pink. Each flower opens for one day only. There are five petals, five sepals, five stamens and the developing fruit has five main segments, each divided by a partition. Full seed set is ten seeds per “boll” or capsule. The bolls change from green to brown as they ripen and hold mature seed for a long time without shattering. Mature seeds are oval, light brown (occasionally yellow) and very glossy due to a surface mucilage that causes the seed to feel sticky when wet.

Key features

Small, pointed, linear leaves alternately spiralled around the stem; blue flowers opening for one day only; mature “bolls” releasing seeds when machine-harvested

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides	
forage crops, hay fields	turf		waste areas	

Seedling

Flower

Key feature: *flax "boll"*

Juvenile

Mature

Ground-ivy

Glechoma hederacea

Other names

Creeping Charlie, gill-over-the-ground

Family

Mint *Lamiaceae*

Seedling

Seedlings are not commonly seen.

Juvenile

Stems are prostrate and square in cross-section. Propagation is largely by these creeping stems that root at the nodes. Ground-ivy can be very aggressive in turf, particularly in shady areas. Leaves are opposite with long petioles. The round to heart-shaped leaves have palmate venation and regularly-scalloped margins. Ground-ivy has an unpleasant, sour, minty odor. The prostrate growth and long petioles of ground-ivy distinguish it from henbit, a mint family weed with an erect growth habit and sessile upper leaves.

Mature

Short, flowering branches are produced from leaf axils of the prostrate stems. These branches are partially to fully erect. Small clusters of purple-blue flowers are in the leaf axils. Each flower is a small, two-lipped tube. Four tiny brown seeds are contained in a persistent calyx.

Key features

Square, prostrate stems that root at the nodes; opposite, petioled leaves with scalloped leaf margins; small, purple flowers produced in leaf axils of erect, flowering stems

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Juvenile: *prostrate, creeping growth habit*

Key feature: *opposite leaves with scalloped margins*

Mature

Flower: *clusters in leaf axils*

Silky Lupine

Lupinus sericeus

Family

Pea *Fabaceae*

Seedling

Seedlings are not commonly seen in their rangeland habitat.

Juvenile

Silky lupine is a tap-rooted legume with an erect stem covered in dense, silky hairs. The alternate leaves are compound palmate with dense, silky hairs covering the leaflets. The closely related species, silvery lupine, has leaflets that are sparsely hairy on the upper surface, though densely silvery-haired underneath.

Mature

Although a meadow of blooming lupines is a delight to hikers, both silky lupine and silvery lupine are poisonous to livestock. Silky lupine racemes have dense blue to bluish-purple flowers. The flower can also be blue and white bi-colored. Silvery lupine, on the other hand, has pale violet to almost white flowers arranged in less dense racemes. The large “standard” petal of a silky lupine flower is densely hairy on the back, while the standard petal of silvery lupine is hairless or nearly so. Pods are flattened and contain several seeds that are especially poisonous.

Key features

Compound palmate leaves covered on both sides with dense, silky hairs; deep blue, pea-type flowers in dense racemes; dense, silky hairs on the back of the “standard” petal

Similar species

Silvery lupine *Lupinus argenteus*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas		

Juvenile

Flower: *deep blue and pea-like*

Key feature: *compound palmate leaf*

Similar species: *Silvery lupine*

Mature

Purple

Flowered Species

Bull Thistle

Cirsium vulgare

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oval and lacking petioles. The first true leaf is oval to oblong with a fringe of spines. The second true leaf has dense, white hairs on the upper surface, which is an easy way to distinguish this weed from Canada thistle at the seedling stage.

Juvenile

Bull thistle forms a deep tap root and a large, flat, basal rosette in the first year. Rosette leaves are deeply lobed, and the lobes occur in clusters of two to three. Each lobe has long sharp spines on the margins and at the tip. Unlike other thistles, the upper leaf surface is prickly while the underside is covered with fine, wooly hair. Stems are erect, tall and often widely-branched. The stem surface is hairy and may have narrow, spiny, leaf-like “wings” on the middle and upper portions. The leaves on the upper stems become smaller toward the top of the plant.

Mature

Large, flask-shaped heads are found at the tops of branches. Each head has long, purple disk flowers enclosed by narrow, overlapping bracts tipped with a yellowish spine. The head size and long sharp spines on the involucre bracts help to separate this species from Canada thistle. The flowers produce a single, greyish-brown seed with dark lines on the surface. The seeds resemble those of Canada thistle except that they are larger and have a longer, more-branched pappus.

Key features

Large, flat, basal rosette in first year; leaves with lobes in clusters and a prickly surface; large, flask-shaped heads with long sharp spines on bract tips

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland	wetland	
reduced tillage		irrigation	roadsides
storage crops, hay fields		turf	waste areas

Juvenile

Flower: head, showing bracts with long sharp spines

Mature

Key feature: leaf-like “wings” on upper stem

Canada Thistle

Cirsium arvense

Other names

Creeping thistle

Family

Sunflower *Asteraceae*

Seedling

Seedlings have oblong to broadly oval, somewhat fleshy cotyledons. Shoots that emerge from underground root buds lack cotyledons. Both types have ovate first true leaves with sharp spines along the margins.

Juvenile

Canada thistle has deep, horizontal, creeping roots and often forms dense patches. It is largely the spreading underground roots that make control of this weed so difficult. The rosette leaves are usually smooth and irregularly-lobed with a spiny-toothed margin. However, plants vary in the extent of lobing, the length of spines on the margin and in the presence or absence of hair on the undersides of the leaves. Stems are erect, hollow and have many alternate, sessile leaves. The stem surface is usually smooth although spines may be present near the base of the plant.

Mature

Heads are numerous in clusters and are smaller than other thistle species. The disk flowers are pink-purple, occasionally white, and surrounded by narrow, flattened involucre bracts ending in short, weak prickles. Bull thistle is sometimes confused with Canada thistle. Its heads are much larger than Canada thistle, and the bracts are more slender and spine-tipped. In addition, the leaves of bull thistle have short prickles all over the surface. Canada thistle has male and female flowers in heads located on separate plants. The male heads are globe-shaped in contrast to the more flask-shaped female heads. Each female flower is capable of producing a single, tan-colored, slightly curved seed. The seeds have a feathery pappus and are readily dispersed by wind.

Key features

Deep, creeping roots; irregularly-lobed leaves with spines on margins only; small male and female heads on separate plants

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Juvenile: *irregularly-lobed leaves*

Seeds with feathery pappus

Mature

Flower: *bracts ending in head with weak prickles*

Common Burdock

Arctium minus

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oval and have noticeable veins. The first true leaves are petioled and ovate with entire to slightly wavy margins. The surface of the leaves is somewhat hairy and wrinkled.

Juvenile

Common burdock develops a thick, fleshy tap root and large basal rosette in the first year. The rosette leaves are ovate to triangular with a heart-shaped base and a wavy margin. Common burdock leaves are often very large and resemble rhubarb except that the undersides are woolly-hairy and the petioles are hollow. Stems are erect and often much-branched. They are rough-hairy, hollow and grooved lengthwise. Stem leaves are usually much smaller and less heart-shaped than the rosette leaves.

Mature

Heads are almost round and are found either singly or in clusters at the ends of branches. Each head has purple disk flowers surrounded by a receptacle with involucre bracts modified into narrow, hooked bristles. This structure forms the bur that aids in the dispersal of common burdock seeds by animals and man. The seeds found within the bur are oblong, smooth and mottled. They look somewhat like narrow, curved sunflower seeds.

Key features

Thick, fleshy tap root; large, heart-shaped basal leaves; receptacle with narrow, hooked bristles forming a bur

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides		
forage crops, hay fields	turf	waste areas		

Seedling

Juvenile: *large basal rosette*

Flower: *head with narrow, hooked bristles*

Mature

Key feature: *bur*

Creeping Bellflower

Campanula rapunculoides

Other names

Garden bluebell

Family

Bluebell *Campanulaceae*

Seedling

Cotyledons are oval. First true leaves are broadly ovate to heart-shaped. Both the leaf blades and petioles are hairy.

Juvenile

Creeping bellflower escapes from gardens and tends to choke out flower beds and creep into lawns. In addition to propagation by seed, it has persistent, fleshy rhizomes and storage tubers, making this an extremely difficult weed to eradicate. Young plants growing from the rhizomes are stemless. The leaves are heart-shaped with fine, irregular toothing and long petioles. Plants with stems are also produced. Stems are tall and erect with alternate leaves that become progressively smaller toward the top of the plant. Also, stem leaves become less heart-shaped and have shorter petioles toward the top of the plant.

Mature

The flowers are violet, five-pointed bells in long, leafy racemes at the tops of stems. The leaves of the raceme are smaller, narrower and sessile. This plant is quite attractive in bloom, and people tend to overlook its weedy nature at first. Each flower develops into a many-seeded, spherical capsule that opens with tiny pores. Seeds are light brown and shiny.

Key features

An urban weed with extensive, fleshy rhizomes and tubers; stemless young plants growing from rhizomes; attractive racemes of violet, bell-shaped flowers

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Juvenile: *heart-shaped leaves*

Mature

Flower: *bell-shaped flowers in raceme*

Mature leaf

Hound's-tongue

Cynoglossum officinale

Family

Borage *Boraginaceae*

Seedling

Cotyledons are oval with short stalks united at the base. The surface of each cotyledon is covered with short, stiff hairs directed toward the tip. The first true leaves are also hairy, especially on the undersurface. They have oval blades and elongated petioles.

Juvenile

Hound's-tongue produces a thick, deep tap root and basal rosette in the first year. The rosette leaves are petioled, broad and oblong, closely resembling the size and shape of a dog's tongue. They are also soft and velvety to the touch. In the second year, stems are formed that often branch near the top of the plant. The stem leaves are stalkless, narrower and more pointed than the basal leaves. The entire plant is covered with soft white hairs.

Mature

The reddish-purple or maroon flowers are located in racemes emerging from upper leaf axils. Each flower has five sepals joined together at the base to form a star. The petals are also fused into a funnel or cup shape. The flowers produce four triangular but rounded seeds with a flattened upper surface. The seeds are covered with short, barbed prickles that enable them to be easily attached to and spread by animals.

Key features

Broad basal leaves in the shape of a dog's tongue; soft white hairs over the entire plant; reddish-purple cup-shaped flowers in a raceme

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Juvenile

Flower: *reddish-purple flower*

Short, barbed prickles on seed

Mature

Key feature: *cup-shaped flowers in a raceme*

Nodding Thistle

Carduus nutans

Other names

Musk thistle

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oblong. The true leaves have irregularly lobed margins with spiny-tipped teeth. The leaves are light greyish-green and may be sparsely hairy.

Juvenile

Nodding thistle has a long, fleshy tap root and forms a large, flat rosette in the first year. The rosette leaves have deeply lobed, spiny margins. Stems elongate in the second year. They are usually sparsely branched and covered with winged prickles. Stem leaves are alternate and deeply lobed with spiny margins.

Mature

Large, showy flower heads are solitary at the ends of branches. Unlike the rest of the plant, the stems are naked just below the flower heads. The disk flowers of the head are bright purple. Bracts around each flower head are in loose whorls with strong sharp spines at the tip. The outer bracts are bent backwards, and most of the heads tend to nod. The seeds are shiny yellow-brown with a white pappus of long, unbranched hairs. The spread of nodding thistle has been somewhat limited by the introduction of an insect (*Rhinocyllus conicus* – a flower weevil) that feeds on the developing seeds.

Key features

Winged prickles along stem except below flower heads; large, nodding, purple flower heads; involucre bracts spine-tipped and bent backwards

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Juvenile: *rosette with deeply lobed leaves*

Key feature: *winged prickles on stems*

Mature

Flower: *head with spiny-tipped bracts*

Purple Loosestrife

Lythrum salicaria

Family

Loosestrife *Lythraceae*

Seedling

Seedlings are not commonly seen. Purple loosestrife forms large colonies in wetlands, marshy areas, ditches and roadsides. It is a major weed in eastern Canada, and infestations are becoming more frequent on the prairies.

Juvenile

In addition to a fibrous root system, purple loosestrife has thick, fleshy roots that can produce new, above-ground shoots. Established plants have a large crown that produces new buds each year. Although propagation is mainly from seed, stem pieces that have broken off or been mowed can also produce roots. Purple loosestrife has tall, branched, erect stems that are usually square in cross-section. Sessile leaves occur in twos or threes at the stem nodes. They are lanceolate in shape with entire margins. Larger leaves may have rounded lobes at the base. Fireweed is sometimes mistaken for purple loosestrife. The leaves of fireweed have short petioles and distinctive lateral veins that run parallel to the margins.

Mature

Bright magenta flowers are produced in long, dense, leafy spikes. The leaves in the inflorescence are alternate and smaller than lower leaves. Fireweed has less dense, non-leafy racemes. Individual flowers of purple loosestrife have five to seven petals and sepals that are united below the petals, while fireweed flowers have four petals. The seed capsule of purple loosestrife contains many tiny reddish-brown seeds. To say it is a prolific seed producer is an understatement. In one year, one plant may produce over two million seeds! The capsule is long and narrow containing seeds with tufts of silky hairs at the tips.

Key features

Leaves in twos or threes at stem nodes; square stems; showy, magenta flowers in long, dense, leafy spikes

Similar species

Fireweed *Epilobium angustifolium*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland	wetland
reduced tillage		irrigation	roadsides
forage crops, hay fields		turf	waste areas	

Juvenile

Flower: *magenta – with five to seven petals*

Infestation

Mature: *with dense, leafy spikes*

Similar species: *Fireweed*

Russian Knapweed

Centaurea repens

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oval. True leaves are hairy, greyish-green with shallow, irregularly-lobed margins. It is more commonly seen growing from the rootstock than as a seedling. Russian knapweed was introduced with alfalfa seed.

Juvenile

Russian knapweed has black to dark brown, scaly roots that penetrate the soil deeply and are widely spreading. Buds on the roots develop into leafy shoots and dense patches of this weed are a problem to control in rangeland and occasionally in field crops. The stems are much-branched and covered with matted, soft grey hairs to give a grey-green appearance. Lower leaves are alternate, sessile to clasping, shallowly and irregularly lobed or toothed. The upper leaves are smaller, usually entire and often end in a soft spine. Young leaves are grey-green and wooly, older leaves less so.

Mature

Spherical, flower heads are produced at the ends of upper, leafy branches. They appear silvery before opening. The receptacle holds only disk-type flowers that are purple to pinkish, becoming straw-colored with age. The whorled involucre bracts are shingle-like with entire margins. They are greenish at the base but papery and white at the tip. Seeds are hairless, white, oval and flattened. The pappus is composed of stiff white bristles, which soon fall off.

Key features

Black, scaly roots that extend deep and wide; papery bracts around heads; young leaves grey-green and wooly-hairy

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling: *grey-green, wooly leaves*

Juvenile: *growing from rootstock*

Infestation

Mature

Key feature: *upper leaves end in soft spine*

Spotted Knapweed

Centaurea maculosa

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oval with short stalks. The first true leaves are spoon-shaped with entire to wavy-edged margins.

Juvenile

Spotted knapweed forms a large tap root and rosette of leaves that are deeply divided into narrow segments. The leaves of its close relative, diffuse knapweed, are much more finely divided by comparison. In both species, the early leaves are covered with a thin layer of matted, wooly hair resembling a cobweb. Spotted knapweed produces erect, hairy stems that are typically less branched than diffuse knapweed. The upper stem leaves are almost linear. Both knapweeds contain a bitter-tasting chemical that makes them highly unpalatable, although not poisonous. They have taken over vast areas of rangeland in the interior of British Columbia and have greatly reduced the availability of desired forage.

Mature

Ovate heads are produced in clusters at the top of the plant. The tiny flowers are pink to purple, rarely white. From a distance, spotted knapweed looks a little like Canada thistle. The flowers are surrounded by oval bracts with a black-tipped fringe resembling a false eyelash. The involucral bracts of diffuse knapweed, on the other hand, are triangular and spine-tipped. Another knapweed species, yellow star-thistle, has bracts that are modified into narrow, sharp spines. Spotted knapweed seeds are narrowly oval, brown and have a "crown" of stiff bristles.

Key features

Deeply lobed, "cobwebby" basal leaves; generally purple flowers; oval bracts with a black-tipped fringe

Similar species

Yellow star-thistle *Centaurea solstitialis*

Life Cycle					
annual		winter annual		biennial
				creeping perennial
					simple perennial
Habitat					
conventional tillage		rangeland		wetland	
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Juvenile: *rosette with deeply divided leaves*

Mature

Flower: *heads with black fringed bracts*

Infestation

Tall Larkspur

Delphinium glaucum

Family

Buttercup *Ranunculaceae*

Seedling

Cotyledons are ovate and long-stalked. The true leaves have long petioles and a broad blade with three to five rounded lobes.

Juvenile

Tall larkspur produces tall, erect, usually hollow stems from a large, woody tap root. The stems are often covered with a bluish waxy coating or "bloom." The leaves are palmately divided into five primary lobes, which are in turn subdivided into narrow segments. They resemble a hand with the fingers stretched outwards. The leaves are usually smooth, but may be somewhat hairy on the margins and undersides. Although a highly attractive plant, tall larkspur is very poisonous, especially to cattle. Poisonings usually occur in early spring when the young plants are most toxic and before better forage is available.

Mature

The deep blue or purplish flowers are found on long racemes. Individual flowers are highly irregular with a pronounced downward-pointing spur. Each flower produces three pods that open to release irregularly-shaped brownish-purple seeds. The related species, low larkspur, is not only a much smaller plant than tall larkspur, but it has two-toned flowers that are white and bluish-purple.

Key features

Stems with bluish waxy bloom; palmately divided leaves; irregular, blue-purple flowers with a downward-pointing spur

Similar species

Low larkspur *Delphinium bicolor*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Mature

Flower: *flowering raceme*

Similar species: *Low larkspur – flowers*

Key feature: *palmately divided leaf*

Seedpods

Two-grooved Milk-vetch

Astragalus bisulcatus

Family

Pea *Fabaceae*

Seedling

Cotyledons are oval and the first true leaves are trifoliolate and long-stalked. Leaves produced later are pinnately compound.

Juvenile

Two-grooved milk-vetch forms deep, tap rooted plants that have several stems emerging from a woody, branched crown. Individual stems are coarse, tough and sparsely covered with bristly white hair. Leaves are pinnately compound with 17 to 27 narrowly oval leaflets. Each leaflet is white and hairy on the underside, but smooth or only sparingly hairy on the top. Normally a palatable plant, two-grooved milk-vetch can become poisonous when growing in dry, alkaline soils. In those situations, it may take up and concentrate high amounts of the mineral selenium in its tissues.

Mature

Dense, spike-like racemes are found on thick stalks at the top of the plant. The reddish-purple, pea-like flowers are often reflexed or downward-pointing. Each flower produces linear pods that are similarly reflexed. The common name for this weed comes from the characteristic two deep grooves found on the upper side of the pod. The pods open to release tiny, oval, reddish-brown to green seeds that have a notch on one side.

Key features

Pinnately compound leaves with 17 to 27 leaflets having hairy white undersides; dense, spike-like racemes of reddish-purple flowers; linear pods with two deep grooves on the upper side

Life Cycle					
annual	winter annual	biennial	creeping perennial	simple perennial	
Habitat					
conventional tillage	rangeland			wetland	
reduced tillage	irrigation			roadsides
forage crops, hay fields	turf			waste areas

Mature

Key feature: *dense, spike-like raceme*

Clump of plants

Pink

Flowered Species

American Dragonhead

Dracocephalum parviflorum

Family

Mint *Lamiaceae*

Seedling

The round cotyledons of American dragonhead have prominent, rounded basal lobes. The true leaves are opposite and oval shaped with scalloped margins. American dragonhead may be confused with henbit, which has similar seedling leaves. The rounded basal lobes of American dragonhead cotyledons are useful to distinguish it from henbit seedlings, which have pointed basal lobes on the cotyledons.

Juvenile

Stems are erect and square in cross-section with sparse hair. The leaves are opposite and petioled. Juvenile and mature leaves of American dragonhead are lanceolate and coarsely-toothed and are distinctly different from seedling leaves. The teeth often narrow into a spiny tip.

Mature

Small flowers are produced in dense spikes at the ends of branches. Flower color varies from almost white to pale pink or purple. The flower spikes of American dragonhead are soft to the touch, unlike hemp-nettle flowers that are very spiny. Each small flower produces four black seeds that are ridged and angled on one side and rounded on the other.

Key features

Scalloped margins of seedling leaves; square stem and opposite leaves with spiny-toothed margins on the adult; soft dense flower spikes

% of fields
where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

reduced tillage

forage crops, hay fields

rangeland

irrigation

turf

wetland

roadsides

waste areas

Seedling

Juvenile

Key feature: *dense flower spike*

Flower

Key feature: *opposite, coarsely-toothed leaves*

Cow Cockle

Vaccaria pyramidata

Other names

Soapwort

Family

Pink *Caryophyllaceae*

Seedling

Cow cockle has narrow, elongated, smooth cotyledons with short stalks. The first true leaves emerge in a pair. They look very similar to the cotyledons except the true leaves have a distinct crease down the center.

Juvenile

Stems are upright, much-branched and have prominent, swollen nodes. Leaves are opposite, sessile and joined at the base. They are oblong with pointed tips and entire margins. Both the stems and leaves are smooth and covered with a bluish-white waxy coating or "bloom." The leaves are thick and feel almost leathery to the touch.

Mature

Flowers are loosely grouped at the ends of the stems. Each flower has five notched, deep pink petals. The petals are surrounded by a green, flask-shaped calyx with five light green ribs. Fruits are brown, elongated capsules opening with four teeth. Each capsule contains many spherical black seeds with a bumpy surface. The seeds are a common contaminant of feed grain and are poisonous to livestock.

Key features

Opposite, smooth leaves; bluish-white waxy bloom; green, five-ribbed calyx

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Flower

Key feature: *five-ribbed calyx*

Juvenile: *with opposite leaves*

Mature

Field Scabious

Knautia arvensis

Other names

Blue buttons

Family

Teasel *Dipsacaceae*

Seedling

Cotyledons are oval and notched at the tip. The first true leaves are petioled and oval to oblong in shape.

Juvenile

Field scabious initially forms a deep tap root and a large, basal rosette. The rosette leaves are broadest in the middle and tapered at both ends. They frequently have a coarsely-toothed margin. Sturdy, erect stems are produced that branch sparingly at the top of the plant. The upper leaves are oppositely arranged and sessile. They are deeply and pinnately lobed into narrow, finger-like segments. The entire plant is covered with short, stiff hairs.

Mature

Small flowers are located in dense heads at the ends of long branches. The flowers may vary from pink to pale purple or even blue. Although it is tempting to place this weed in the *Asteraceae*, it does not belong to that family. Each tiny, funnel-shaped flower in the head produces a single, hairy, rectangular seed.

Key features

Upper leaves pinnately lobed and opposite; entire plant covered with short, stiff hairs; pink to blue flowers in dense heads

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Flower: *head of dense flowers*

Key feature: *short, stiff hairs*

Juvenile

Mature

Hemp-nettle

Galeopsis tetrahit

Family

Mint *Lamiaceae*

Seedling

The cotyledons of hemp-nettle have distinctly pointed basal lobes, long stalks and are slightly indented at the tip. True leaves are opposite, hairy and have coarsely-toothed margins. Seedlings of yellow-flowered hemp-nettle are indistinguishable from hemp-nettle.

Juvenile

The stem of hemp-nettle is square in cross section and covered with downward-pointing, bristly hairs. Although these hairs can penetrate the skin and cause irritation, they do not produce the extreme reaction caused by the specialized hairs of stinging nettle. Hemp-nettle also lacks rhizomes and stipules at the nodes, which are characteristic of stinging nettle. Hemp-nettle stems are conspicuously swollen below the nodes. The oppositely arranged leaves are elliptical with long petioles and are covered with stiff hairs.

Mature

Flowers are produced in clusters in the upper leaf axils. Each flower is a pink, two-lipped tube, although color can vary from white to purple to variegated. The five-pointed calyx surrounding each flower is extremely spiny, making hemp-nettle very unpleasant to touch at this stage. The four seeds produced by each flower remain inside the persistent calyx until they disperse at maturity. Seeds are shaped like a teardrop and have a grey, brown and black mottled surface. There is a round scar at the base of the seed. The mature plant of yellow-flowered hemp-nettle can be distinguished from hemp-nettle by its larger size and pale yellow flowers.

Key features

Stems swollen below the nodes; bristly, downward-pointing hairs covering the square stem; opposite leaves with long petioles

Similar species

Yellow-flowered hemp-nettle *Galeopsis speciosa*

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling

Juvenile

Key feature: *bristly hairs on square stem*

Mature

Flower

Henbit

Lamium amplexicaule

Family

Mint *Lamiaceae*

Seedling

Cotyledons of henbit are round and slightly indented at the top. Basal lobes of the cotyledons are pointed. The true leaves are opposite with scalloped margins and are similar to the first true leaves of American dragonhead. However, the cotyledons of American dragonhead have rounded basal lobes.

Juvenile

Stems of henbit are square in cross section and weakly erect. Branching occurs near the base of the plant. Leaves are oppositely arranged and have irregular, round-toothed margins. The lower leaves of henbit are long-stalked while upper leaves are sessile and clasp the stem. The erect stems and clasping upper leaves distinguish henbit from ground-ivy, which is mostly prostrate with long-stalked leaves.

Mature

Pink to purplish flowers are clustered in the axils of upper clasping leaves. Each small flower is a two-lipped tube. The upper lip is unlobed and has a tuft of hair on the top. The lower lip is two-lobed. Each flower produces four brown seeds with white dots.

Key features

Square, weakly erect stems; long-petioled lower leaves and clasping upper leaves; opposite leaves with irregular, round-toothed margins

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Key feature: *clasping upper leaves*

Mature

Flower: *clusters*

Pale Smartweed

Polygonum lapathifolium

Family

Buckwheat *Polygonaceae*

Seedling

Cotyledons are narrowly elliptical and joined at the base. True leaves emerge one at a time. They are lanceolate with short petioles and entire margins.

Juvenile

Stems are branched and vary from erect to partially prostrate in growth habit. They are slightly swollen above the nodes. The ocrea at each node is smooth and papery. The upper margin is either entire or has short hairs. The ocrea is key to differentiating both pale smartweed and green smartweed from lady's-thumb. Lady's-thumb has short hairs on the surface of the ocrea and a fringe of visible hair on the upper margin of the ocrea. Pale and green smartweed leaves may or may not have a purplish blotch on the upper surface. The blotch is usually present on lady's-thumb leaves. The first five to seven leaves of pale smartweed and green smartweed have white matted hair on the undersurface. Subsequent leaves have tiny, yellow glands dotted on the undersurface. Both these features are lacking on lady's-thumb leaves.

Mature

Flowers lack petals but sepals are often colored. Pale smartweed flowers are pinkish-white or greenish-white in dense, elongated spikes. Green smartweed has greenish-white flowers in erect spikes, which are shorter and plumper than those of pale smartweed. Lady's-thumb has pink-flowered spikes. Both pale smartweed and green smartweed have tiny, yellowish glands on the upper stems below the spikes. These glands are absent on upper stems of lady's-thumb. Seeds of pale and green smartweed are black, shiny and rounded with a pointy tip. The flattened sides are dented or hollow in the middle. Lady's-thumb seeds are three-sided or flattened, but not dented on the flattened sides.

Key features

Smooth, papery ocreas at swollen stem nodes; leaf undersurface with white matted hair or tiny, dotted glands; tiny glands on upper stems below spikes

Similar species

Green smartweed *Polygonum scabrum*; lady's-thumb *Polygonum persicaria*

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling

Juvenile

Key feature: *ocrea*

Mature

Similar species: *Green smartweed – spike*

Prickly Rose

Rosa acicularis

Other names

Wild rose

Family

Rose *Rosaceae*

Seedling

Seedlings are not commonly seen.

Juvenile

Three species of wild shrubby roses readily hybridize, making identification difficult. Prairie rose, *Rosa arkansana*, is a short, low shrub with few branches. Its stems mostly die back each year. It is easier to confuse prickly rose and Woods' rose, *Rosa woodsii*, which are much taller shrubs, multi-branched and have persistent stems from year to year. Prickly rose is a bushy shrub of pastures. It has erect, stout stems densely covered with thorns. Stems of Woods' rose have thorns that are scattered or only near the base. The compound pinnate leaves of both species are alternate with toothed leaflets. Leaves of prickly rose have three to seven leaflets, commonly five. Woods' rose normally has either five to seven or seven to nine leaflets. Prickles are absent below the stipules in prickly rose but are present in Woods' rose. Stipules of prickly rose are hairy and densely glandular while stipules of Woods' rose are sparsely so.

Mature

Flowers are pink, five petalled and usually solitary. The ovary and sepals of each flower develop into the fleshy fruit or "rosehip," which contains many seeds. The rosehip is round to pear-shaped with a distinct stalk or "neck" below. Woods' rose fruits are spherical but lack a neck.

Key features

Thorny stems; pinnately compound leaves, commonly with five leaflets; rosehips on distinct necks

Similar species

Prairie rose *Rosa arkansana*, Woods' rose *Rosa woodsii*

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

reduced tillage

forage crops, hay fields

rangeland

irrigation

turf

wetland

roadsides

waste areas

Juvenile

Key feature: *compound pinnate leaves with five leaflets*

Mature

Flower

Prostrate Knotweed

Polygonum aviculare

Other names

Knotweed, knotgrass, doorweed

Family

Buckwheat *Polygonaceae*

Seedling

Cotyledons are narrow and united at the base. The first true leaves are narrowly oval with short stalks.

Juvenile

As its name suggests, prostrate knotweed frequently spreads its stems flat over the ground in open areas.

However, when the weed grows in more densely populated sites, its stems may be partly or even fully erect. The thin, wiry stems have prominent, swollen nodes, giving them the appearance of a piece of string with knots tied in it. The silvery colored ocreas serve to further emphasize the swollen nodes. Stems are typically much-branched with small, alternate leaves that are broadest in the middle and narrow at both ends. The entire plant often has a bluish-green color. There are several closely related and very similar looking knotweed species. Striate knotweed, also found in Western Canada, may be distinguished from prostrate knotweed by its coarser, more erect stems and broader, more rounded leaves. All knotweeds have tough, deeply penetrating tap roots and thrive in poor, compacted soils; they will even do well in cracks in concrete.

Mature

The flowers are very small and inconspicuous. They lack petals but usually have five pink to purple, but occasionally green, sepals. The flowers are produced in leaf axils and are usually at least partially enclosed by the ocrea. Each flower produces a tiny, triangular dark brown seed that is surrounded by a papery hull.

Key features

Prostrate to semi-erect growth habit; long, wiry stems with swollen nodes covered with an ocrea; tiny, pink-purple axillary flowers

Similar species

Striate knotweed *Polygonum achoreum*

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

reduced tillage

forage crops, hay fields

rangeland

irrigation

turf

wetland

roadsides

waste areas

Seedling

Juvenile

Mature: *prostrate growth habit*

Flower: *small flowers in leaf axils*

Key feature: *swollen nodes with ocrea*

Showy Milkweed

Asclepias speciosa

Family

Milkweed *Asclepiadaceae*

Seedling

The cotyledons are oblong and have a shiny undersurface. True leaves are opposite with short hairs on the margin. Seedlings are rarely noticed in their roadside and pasture habitat.

Juvenile

Although showy milkweed is often listed as a poisonous species, livestock usually avoid it unless they are starving. It tends to spread slowly by creeping rootstocks to form clumps. Showy milkweed leaves appear greyish-green due to the soft hairs covering the surface. Leaves are oblong and leathery-thick. They are oppositely arranged on the stem and have very short petioles and stipules at the nodes. The whole plant is filled with a white, sticky latex.

Mature

The beautiful pink and white flowers are arranged in large, rounded umbels. Each flower has five outer petals and an inner five-pronged corona seated on the stamen tube. The flower stalks are covered with white wooly hairs. Each flower develops into either one or two large pods covered with wooly hairs and soft projections on the surface. The pod opens on one side at maturity to release many reddish-brown, flattened seeds. Each seed is equipped with a tuft of silky hair for wind distribution.

Key features

White, sticky latex in stem and leaves; opposite leaf arrangement; large, wooly-hairy seedpods that have soft projections on the surface and open on one side only at maturity

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides		
forage crops, hay fields	turf	waste areas		

Seedling

Juvenile

Flower: *round umbels*

Key feature: *large woolly-hairy seedpods*

Infestation

Stork's-bill

Erodium cicutarium

Other names

Redstem filaree

Family

Geranium *Geraniaceae*

Seedling

The three-lobed cotyledons of stork's-bill are unique. The true leaves are deeply and pinnately divided into segments, which are further lobed. Coarse, spreading hairs cover the leaf surface.

Juvenile

Stork's-bill has a tap root with stout lateral roots. The hairy stems are low and spreading, forming tangled clumps. Basal leaves are petioled and pinnately compound with deeply cleft leaflets. Stem leaves are fewer, smaller and sessile.

Mature

Flowers are in clusters, each with five pink-purple petals and five sepals with bristly tips. The styles of the pistil are united and elongate into a column or long "bill," hence the name stork's-bill. At maturity, the bill splits upward into five parts, each a one-seeded, long-tailed segment attached to a slender brown seed. When the tail dries, it resembles a coiled spring.

Key features

Three-lobed cotyledons on seedling; low, spreading growth habit; distinct bill of the fruit and coiled tail on the seed

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

reduced tillage

forage crops, hay fields

rangeland

irrigation

turf

wetland

roadsides

waste areas

Seedling

Juvenile: *pinnately compound leaves*

Key feature: "bill" of fruit

Mature

Flower: *pink-purple in cluster*

Water Smartweed

Polygonum amphibium

Family

Buckwheat *Polygonaceae*

Seedling

Seedlings are not commonly seen.

Juvenile

This species has both aquatic and terrestrial forms. The aquatic form is usually found growing in standing water while the terrestrial form is found in cultivated fields. Swamp smartweed, *Polygonum coccineum*, is a related and very similar species that also has both aquatic and terrestrial forms.

In cultivated fields, water smartweed is usually found in patches growing from extensive, creeping rootstocks. The stems have swollen nodes with prominent ocreas. Growth habit is erect or weakly erect. Leaves are alternate and petioled. They have entire margins and are approximately ovate, although shape varies. Leaves and stems are usually both hairy. The terrestrial form has a distinct “skirt” or spreading collar extending from the upper edge of the ocrea. The terrestrial form of swamp smartweed also has this feature. It is difficult to distinguish between the two species before flowering.

Mature

Although individual flowers lack petals, the calyx is bright pink to scarlet. The spike-like racemes are short and thick with hairless stalks below the inflorescence. The spike-like racemes of swamp smartweed are longer and more slender with hairy stalks below the inflorescence. Each flower produces one seed. Both species have brown to black, lens-shaped seeds.

Key features

In patches growing from creeping rootstocks; swollen nodes with prominent ocreas that may be “skirted”; bright pink to red flowers in spike-like racemes

Similar species

Swamp smartweed *Polygonum coccineum*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Juvenile

Key feature: *ocrea* at stem node

Mature

Flower: *spike-like raceme of bright pink flowers*

White Flowered Species

Baby's-breath

Gypsophila paniculata

Family

Pink Caryophyllaceae

Seedling

Cotyledons are linear. The true leaves are also linear, with entire margins and rounded tips. The leaves emerge in pairs and appear greyish-green. Seedlings are not often noticed.

Juvenile

Baby's-breath has a deep, penetrating root system. The plant is erect with freely-branching stems that are smooth and swollen at the nodes. Stems and leaves are covered with glandular hair to give a greyish-green, waxy "bloom." The leaves are opposite, entire-margined and linear to lanceolate with a single central vein.

Mature

Baby's-breath plants resemble large, white powder puffs along prairie roadsides. The many small, white-petalled flowers in an open-branched inflorescence give this delicate "puffy" appearance. Each flower develops into a multi-seeded capsule containing tiny black, snail-like seeds with minute bumps and ridges. The mature, brittle stem breaks at the base and tumbles with the wind to distribute seed.

Key features

Cotyledons and true leaves of seedling resemble each other; greyish-green "bloom" on stems and leaves due to glandular hair; delicate white "puffs" along prairie roadsides

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Mature

Flower: *small white flowers*

Key feature: *"white powder puffs"*

Infestation

Bladder Champion

Silene cucubalus

Family

Pink Caryophyllaceae

Seedling

The cotyledons and first true leaves resemble each other. They are oblong with entire margins, hairless and emerge two at a time.

Juvenile

Bladder champion has a deep, branching tap root. Like dandelion, new bladder champion plants can be produced if the crown of the plant is severed. Its much-branched stems are erect with swollen nodes. Leaves are opposite, sessile and joined at the base. They are long and narrow with a tapered tip. Both the stem and leaves are smooth and light green-to-whitish due to a waxy covering or “bloom” on the surface.

Mature

Flowers are found in branching clusters at the top of the plant. Each flower has five deeply-lobed white petals surrounded by a bladder-like calyx. The calyx may be light green or pink with either darker green or purplish net veins. The round seed capsules have six teeth at the top and are enclosed by the mature, loose, papery calyx. Seeds are kidney-shaped, grey and covered with tiny bumps. They are difficult to distinguish from the seeds of both white cockle and night-flowering catchfly.

Key features

Deep, branching tap root; opposite leaves and stems with a waxy bloom; bladder-like green or pink calyx

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland	wetland	
reduced tillage		irrigation	roadsides
forage crops, hay fields		turf	waste areas

Seedling

Juvenile: showing opposite leaves

Capsule with six teeth

Mature

Flower: with bladder-like calyx

Canada Fleabane

Erigeron canadensis

Other names

Horseweed

Family

Sunflower *Asteraceae*

Seedling

Canada fleabane has oval cotyledons and bright green, hairy leaves.

Juvenile

The rosette leaves are bright green, softly hairy and petioled. They are elliptical in shape and coarsely toothed.

Stems are erect and bristly-hairy. Stem leaves are alternate and often closely crowded. Middle and upper leaves are linear, smaller than basal leaves and are more harshly hairy than the young leaves. They taper to the stem and usually have no teeth on the margins. Crushed leaves have a carrot-like odor. Canada fleabane is becoming more and more frequent as a winter annual weed in fields with reduced tillage.

Mature

The numerous, fluffy-looking flower heads are on short branches at the top of the plant. The central, yellow disk flowers are surrounded by outer ray flowers. The ray flowers are white and may be partially or completely hidden by the several whorls of narrow, pointed bracts around each flower head. The yellowish seeds are oblong and flattened with a white pappus for wind dispersal.

Key features

Bristly-hairy stems and leaves; basal leaves toothed, upper leaves usually entire; numerous fluffy flower heads

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling

Juvenile: *rosette*

Flower: *numerous fluffy flower heads*

Mature

Key feature: *bristly-hairy stems*

Chickweed

Stellaria media

Family

Pink *Caryophyllaceae*

Seedling

Cotyledons are narrowly oval with a pinched tip. The first true leaves are opposite and broader than the cotyledons, although they too have a pinched and pointed tip.

Juvenile

Chickweed is low-growing with bright green, much-branched stems. Plants often form dense mats that are highly competitive with crop and garden plants. The stems may root at the nodes, making this weed very difficult to hand-pull. They are smooth except for a single row of fine, white hair. The opposite leaves also have a row of hair on the petioles. Leaves on the upper stems are usually stalkless. Chickweed can be distinguished from a related species, mouse-eared chickweed, by the leaves. In the latter species, the leaves are sessile and densely-hairy all over, resembling the ear of a mouse.

Mature

The small, star-shaped flowers are located at the tips of the stems or in branch axils. The flowers have five white, deeply-notched petals that are shorter than the surrounding hairy sepals. Each flower opens for only one day before a fruit develops. The seed capsules are egg-shaped and open by six teeth. Nearly round, reddish-brown seeds with a bumpy surface are produced throughout the growing season.

Key features

Low-spreading growth habit; single row of hairs on the stems and leaf petioles; tiny, star-shaped flowers

Similar species

Mouse-eared chickweed *Cerastium vulgatum*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Juvenile: *low-growing*

Mature

Flower: *deeply-notched petals*

Key feature: *hairy petioles, opposite leaves*

Cleavers

Galium aparine

Other names

Bedstraw

Family

Madder *Rubiaceae*

Seedling

Cotyledons are petioled and oblong with a notched tip. The first true leaves are narrow with a pointed tip, sessile and arranged in a whorl of four.

Juvenile

Cleavers has numerous, freely-branched stems that are weak and straggly. The plant has difficulty standing upright and is more often found lying on nearby vegetation. Stems are four-sided and covered with stiff, backward-pointing hairs that allow them to adhere to plants, animals and people alike. The leaves produced at this stage are usually in whorls of six to eight and are also covered with prickly-like hair. A close relative of cleavers, called false cleavers, has been identified in some areas of Alberta and Saskatchewan. While very similar in appearance, false cleavers can be distinguished by the leaves, which are notched at the tip. Both species are equally troublesome and difficult to control in crops.

Mature

Tiny, greenish-white flowers are located in clusters of two to five in leaf axils. While the flowers are very inconspicuous, they produce unique-looking fruits that appear as two seeds joined together. The weed's common name is derived from the fact that these two seeds break apart or "cleave" at maturity. Each greyish-brown seed is nearly round and covered with hooked bristles. They are similar in size to canola and are very difficult to screen out of that crop.

Key features

Narrow leaves with pointed tips arranged in whorls; weak, four-sided stems with prickly hair; fruits formed of two seeds joined together

Similar species

False cleavers *Galium spurium*

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

reduced tillage

forage crops, hay fields

rangeland

irrigation

turf

wetland

roadsides

waste areas

Seedling

Juvenile: *showing pointy-tipped leaves in whorls*

Mature: *straggly plants using crop for support*

Flower: *tiny, greenish-white*

Key feature: *two-seeded fruits*

Common Pepper-grass

Lepidium densiflorum

Other names

Pepperweed, green-flowered pepper-grass

Family

Mustard *Brassicaceae*

Seedling

Common pepper-grass has linear cotyledons with rounded tips. The first true leaves are petioled with deeply and irregularly lobed margins.

Juvenile

The leaves of the basal rosette are deeply and irregularly lobed with some variation from plant to plant. The stem is erect, covered with short hairs and very branched, especially near the top of the plant. Stem leaves are alternate, sessile and linear to lanceolate in shape. The margins are entire or occasionally toothed.

Mature

Common pepper-grass is not usually noticed until the pods are formed. The flowers are in a raceme, but are very inconspicuous as they often lack petals. Even if tiny white petals are present, they are shorter than the calyx, hence the other name “green-flowered pepper-grass.” The tiny seedpods are densely packed on the raceme. They are nearly round, flattened and notched at the top with a tiny bump in the middle of the notch. There are two seeds per pod, one on each side of the membranous wall inside the pod. The flattened, oblong seeds are bright reddish-yellow. At maturity, the brittle stem breaks off, and the plant rolls along the ground to scatter seed.

Key features

Basal rosette of petioled leaves with irregularly and deeply lobed margins; dense racemes of tiny, green seedpods; pods with two seeds

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Juvenile: *rosette with irregularly-lobed leaves*

Mature

Key feature: *dense, tiny seedpods*

Corn Spurry

Spergula arvensis

Family

Pink Caryophyllaceae

Seedling

Both cotyledons and true leaves are needle-like, erect and bright green. The true leaves emerge in whorls of at least three leaves per node.

Juvenile

Corn spurry may resemble young plants of field horsetail, a perennial weed that grows in a similar habitat. Unlike corn spurry, the stems of field horsetail are jointed with black-tipped bracts surrounding the nodes. It also has whorled branches, unlike the individual whorled leaves of corn spurry. The stems on corn spurry are bright green and have surfaces that range from smooth to finely hairy or even sticky in texture. The linear, somewhat fleshy leaves typically occur in whorls of six to ten, with small stipules below. The leaves appear either flat or slightly rounded in cross-section. Young corn spurry plants often exhibit a low, sprawling habit whereas fully grown plants may be either erect or spreading with single or much-branched stems.

Mature

Flowers are very small, but often numerous, and in clusters at the end of branches. The flowers are short-stalked and have five green sepals joined at the base and five white petals. The fruits are tiny capsules that tend to point downward at maturity. They split into five divisions to release small black, nearly round, flattened seeds. The seeds typically have a narrow white wing around the margin as well as whitish projections on the surface that resemble tiny pieces of chaff.

Key features

Bright green color; needle-like leaves; whorled leaf arrangement

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling

Juvenile

Flower: small, white

Mature

Key feature: needle-like leaves in whorls

Death Camas

Zygadenus gramineus

Family

Lily *Liliaceae*

Seedling

Seedlings are seldom noticed.

Juvenile

Death camas is extremely poisonous to livestock and humans. The related, more robust white camas is also poisonous, but less so. The early spring growth of both species is grass-like from the underground bulb that is particularly toxic. Both species are more likely to cause problems in the spring because the bulb may pull out more easily when the ground is wet. The long, narrow leaves of both species are keeled making them V-shaped in cross-section. Before flowering, death camas or white camas could be confused with prairie onion, which has U-chanelled leaves and a distinct onion smell when bruised. Death camas and white camas leaves do not have the strong onion odor.

Mature

The flowers of death camas are small with six creamy-white to yellow petals. They are arranged in a raceme that may be branched at the base. Flowers of the more showy white camas are larger, creamy-white to green in a less dense raceme that is unbranched at the base. White camas has a green, heart-shaped gland on each segment of the flower that can be used to distinguish it from death camas, which has a semicircular gland. Flowering plants should be easily distinguished from wild onion species, which have white or pink flowers arranged in simple umbels. The three-lobed capsule of death camas is elliptical and contains many seeds.

Key features

Long, narrow leaves V-shaped in cross-section; underground bulb; raceme of creamy-white to yellow flowers

Similar species

White camas *Zygadenus elegans*; prairie onion *Allium textile*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Mature

Flower: *raceme of creamy-white flowers*

Key feature: *bulb*

Capsules

Similar species: *White camas – flowers*

Diffuse Knapweed

Centaurea diffusa

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oval with short stalks. The first few true leaves are oblong with an entire margin while later leaves have shallow lobes.

Juvenile

Diffuse knapweed produces an elongated tap root and a rosette of much-divided, feathery-looking leaves. In contrast, the leaves of spotted knapweed are deeply lobed into narrow segments. The leaves of both species are covered with a thin layer of matted, wooly hair resembling a cobweb. Stems are erect, much-branched and hairy. Upper stem leaves are bract-like and entire. Although diffuse knapweed is not considered poisonous, it is extremely bitter-tasting and generally avoided by grazing animals. Both species of knapweed have almost formed monocultures in rangeland throughout parts of the British Columbia interior. Consequently, a major campaign is underway to minimize their spread into the prairies.

Mature

Numerous narrow heads are produced singly at the ends of branches. The flowers are generally white but occasionally pink-purple. They are surrounded by pale yellowish-green, triangular bracts ending in a stiff spine. The involuclral bracts are a key way to distinguish diffuse knapweed from its relatives, spotted knapweed and yellow star-thistle. Spotted knapweed heads have bracts with a black-tipped fringe while the bracts on yellow star-thistle heads are modified into slender, sharp spines. The seeds of diffuse knapweed are narrowly oval, light brown-to-black and lack a pappus.

Key features

Much-divided, "cobwebby" basal leaves; generally white flowers; triangular, spine-tipped bracts

Similar species

Yellow star-thistle *Centaurea solstitialis*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Flower: head, showing spine-tipped bracts

Infestation

Juvenile: *rosette*

Mature

Field Bindweed

Convolvulus arvensis

Family

Morning-glory *Convolvulaceae*

Seedling

Cotyledons are broad with a notched tip and visible venation. The true leaves are entire, petioled and arrow-shaped with blunt tips. Field bindweed is sometimes confused with wild buckwheat, which also has arrow-shaped leaves. However, wild buckwheat has linear cotyledons and more pointed leaf tips.

Juvenile

Field bindweed has persistent, spreading, white cord-like roots with buds that produce new shoots. Branched stems twine around the crop or form prostrate mats if there is no crop for support. Leaves are alternate, petioled and generally arrow-shaped with blunt tips. There is considerable variation in the overall leaf shape and the shape of the basal lobes. Wild buckwheat is also a twining plant and is often confused with field bindweed. However, wild buckwheat has ocreas at the stem nodes, and its arrow-shaped leaves tend to have more pointed tips.

Mature

Flowers are showy, white to pinkish and funnel-shaped with long stalks. Each flower lasts only one day. Flowers are usually solitary on stalks from leaf axils or in small groups. There is a pair of small narrow bracts on the flower stalks. Each flower produces a seed capsule with one to four seeds inside. The seeds are pear-shaped, appear dull grey-brown and have tiny bumps on the surface.

Key features

Twining stems; arrow-shaped leaves with blunt tips; funnel-shaped, white to pinkish flowers with long stalks

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Flower: funnel-shaped

Juvenile

Mature

Key feature: trailing stems

Hairy Nightshade

Solanum sarrachoides

Other names

Cupped nightshade

Family

Nightshade *Solanaceae*

Seedling

Cotyledons are stalked and narrowly oval with distinctly pointed tips. The first true leaves are ovate with wavy margins.

Juvenile

Hairy nightshade has stems that are usually erect but much-branched and spreading. Leaves are ovate to triangular with margins that range from entire to wavy and even toothed. As the name suggests, the entire plant is covered with dense, short hair as well as long, glandular hair making it sticky to the touch. Hairy nightshade is easily confused with black nightshade at the juvenile stage. While both species are hairy, black nightshade lacks the more prominent, sticky hairs.

Mature

The star-shaped flowers are smaller but closely resemble those of potato and tomato. They occur in axillary clusters of three to ten. The petals are either white or white tinged with purple. Each flower produces a berry that is initially green but turns yellowish-brown when fully ripe. The berries are partially surrounded or cupped by the sepals. Inside each berry are numerous light brown, tomato-like seeds. Once the fruits are formed, it is very easy to tell this weed from black nightshade. The berries of black nightshade are purple to black at maturity and are not cupped by the sepals. In both cases, people should avoid eating the immature green fruits since they are known to be highly poisonous.

Key features

Ovate to triangular leaves; long, glandular hairs that make plants sticky to the touch; yellowish-brown, cupped berries

Similar species

Black nightshade *Solanum nigrum*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides	
forage crops, hay fields	turf		waste areas	

Juvenile

Flower

Similar species: *Black nightshade* – berries

Mature

Key feature: *berries cupped by sepals*

Narrow-leaved Milk-vetch

Astragalus pectinatus

Family

Pea *Fabaceae*

Seedling

Seedlings of narrow-leaved milk-vetch are frequently overlooked in its rangeland habitat.

Juvenile

Narrow-leaved milk-vetch has a deep, stout tap root and an underground woody stem. Aerial stems are weakly erect to spreading, reddish at the base and found in clumps. The compound pinnate leaves usually have from nine to seventeen leaflets. The leaflets are linear and very narrow with flattened hairs. Pointed stipules are found at stem nodes. The whole plant has a distinctly unpleasant odor. It is unpalatable and therefore unlikely to poison livestock unless there is a severe shortage of forage. This species is a good indicator of high selenium soils.

Mature

Bloom occurs in late May to early June. Racemes are located in upper leaf axils. The pea-type flowers are cream-colored to pale yellow. Each flower forms a fleshy pod that is circular in cross-section. The pod becomes very dry and hard at maturity. The irregular, kidney-shaped seeds are tan to reddish-brown.

Key features

Clumps of weakly erect stems; unpleasant odor; compound pinnate leaves with narrow, linear leaflets

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Juvenile

Mature

Key feature: *compound pinnate, hairy leaf*

Flower: *flowering raceme*

Night-flowering Catchfly

Silene noctiflora

Other names

Sticky cockle

Family

Pink *Caryophyllaceae*

Seedling

Cotyledons are smooth and oblong. The first true leaves emerge two at a time and have dense, short hair on the surface and margins. This seedling is easily confused with bluebur. The true leaves of bluebur emerge one at a time and have a distinct crease down the center. In addition, bluebur cotyledons have a small "bump" at the tip.

Juvenile

Night-flowering catchfly often forms a rosette of opposite, densely-hairy leaves. They are wide at the tip with entire margins and broad but short petioles. These lower leaves are usually not sticky to the touch. Stems are erect with swollen nodes. They may branch sparingly near the top of the plant. Leaves on the upper plant are sessile and narrower at the tip than the basal leaves. The upper stems and leaves are covered with sticky, glandular hairs.

Mature

The flowers have five white-to-pink, notched petals and are usually open in the evening. The petals are enclosed in a sticky-hairy, tubular calyx with ten branched veins. Each flower produces a light brown capsule with six teeth at the top. Inside are many grey, kidney-shaped seeds covered with rows of minute bumps. The seeds are a frequent contaminant of timothy and alsike or red clover seed. Night-flowering catchfly is often mistaken for its close relative, white cockle, which is a biennial and not sticky to the touch. The calyx of white cockle has veins that are less-branched than night-flowering catchfly, and it forms a capsule opening by ten teeth.

Key features

Opposite, hairy leaves; sticky upper stems and flowers; a branched-veined calyx

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

reduced tillage

forage crops, hay fields

rangeland

irrigation

turf

wetland

roadsides

waste areas

Seedling

Juvenile: *rosette*

Flower: *sticky, branched-veined calyx (right)*

Mature

Seed capsule with six teeth

Ox-eye Daisy

Chrysanthemum leucanthemum

Other names

White daisy

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oval and stalkless. The first true leaves are broadly oval in shape with wavy, lobed margins.

Juvenile

Ox-eye daisy has shallow, short, creeping roots. A basal rosette is produced in the first year. The rosette leaves are petioled, spoon-shaped and coarsely-toothed. They are dark-green and have a smooth, almost shiny surface. Tall, sparsely-branched stems emerge in the second year from a thick crown. They are typically smooth, but may be sparingly hairy at the base. The upper stem leaves are narrower and more deeply-toothed than the basal leaves. They are sessile and often clasp the stem. All parts of the plant have a strong, unpleasant odor.

Mature

Numerous yellow disk and white ray flowers occur in dense, showy, daisy-like heads. Ox-eye daisy heads resemble those of scentless chamomile. The two weeds are easily distinguished by the spoon-shaped leaves and strong odor associated with ox-eye daisy. Each of the tiny disk flowers in the center of the head produces a single seed. The seeds are narrow and black with ten prominent white ribs. They have a small projection on one end, but lack the feathery pappus typical of many other *Asteraceae* weeds.

Key features

Short, creeping roots; spoon-shaped, coarsely-toothed basal leaves; strong, unpleasant odor

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Juvenile: *rosette*

Flower: *head*

Mature

Infestation

Pygmyflower

Androsace septentrionalis

Other names

Fairy candelabra

Family

Primrose *Primulaceae*

Seedling

The young seedling is extremely tiny and could easily be overlooked. It has tiny, oval cotyledons and sessile leaves.

Juvenile

The rosette is very tiny. It has elliptical leaves that are either hairless or covered with minute hairs. The leaf margins are usually entire but may be partly toothed. This plant is seen in early spring in stubble fields and gardens. Wood whitlow-grass has a tiny rosette and is found in a similar habitat. It can be distinguished from pygmyflower because it has branching hairs on the leaves.

Mature

Although several flowering stems arise from the rosette, usually one flowering stem is more well-developed than the others. The tiny white flowers are arranged in a simple umbel with bracts at the base. Each flower is tubular with five lobes that are slightly notched at the tip. When the petals fall, the five pointy lobes of the calyx resemble a tiny green star. Each flower develops into a many-seeded capsule.

Key features

Tiny leaves, flowers and overall short stature of the plant; early spring bloom; bracts at the base of the white-flowered umbel

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides		
forage crops, hay fields	turf	waste areas		

Seedling

Flower

Juvenile: *rosette*

Mature

Round-leaved Mallow

Malva rotundifolia

Other names

Cheeseweed

Family

Mallow *Malvaceae*

Seedling

Cotyledons are distinctly heart-shaped with a rounded tip and prominent veins. The first true leaves are long-stalked and round to kidney-shaped.

Juvenile

Round-leaved mallow is a low, spreading plant with long, much-branched, hairy stems. In densely populated areas, however, stems may be much more upright, even erect. The leaves at this stage usually appear five-lobed with a toothed margin. The teeth on the margin are frequently rounded at the tip, making the leaves look almost scalloped. They may be either smooth or sparingly hairy. The upper leaf surface has pronounced palmate venation and a wrinkled texture. There are small green stipules at the stem nodes.

Mature

Tiny, white or pale blue flowers are found in axillary clusters of one to three. Each flower produces a circular, disk-like fruit. Inside the fruit, seeds are arranged in a ring somewhat resembling the sections in an orange. Individual seeds are brown and wedge-shaped with a rough, pitted surface.

Key features

Round to kidney-shaped leaves with long petioles; long, spreading or erect much-branched stems; circular fruit with wedge-shaped seeds

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Key feature: *kidney-shaped leaves*

Infestation

Mature

Flower: *in axillary clusters*

Scentless Chamomile

Matricaria perforata

Other names

Scentless mayweed

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oval and sessile. The first true leaves are deeply-divided, often into five distinct segments. Both cotyledons and true leaves are smooth. Seedlings may be easily confused with flixweed. Flixweed cotyledons are stalked and narrower than scentless chamomile. In addition, the true leaves of flixweed are usually tri-lobed and hairy.

Juvenile

Scentless chamomile forms a rosette of finely-divided leaves. The leaves are almost fleshy in texture and have no odor when crushed. The sometimes confusing wild chamomile and pineappleweed have a pleasant aroma while ox-eye daisy and stinking mayweed both have a distinctly unappealing scent. In addition, ox-eye daisy can be distinguished from scentless chamomile by its spoon-shaped, lobed leaves. The stems produced by scentless chamomile are erect, smooth and often much-branched.

Mature

Small, daisy-like heads are found singly at the ends of the branches. Each head has tiny yellow disk flowers in the center surrounded by flat, white ray flowers. When the heads are ripe, the white flowers tend to point backwards, giving the entire head the appearance of a badminton shuttlecock. At first glance, the heads of wild chamomile look very much like scentless chamomile. With closer inspection, it is possible to see that the receptacle of scentless chamomile is more rounded than the conical-shaped receptacle of wild chamomile. Each yellow flower produces a seed that is dark brown with three white ribs on one side. The seeds do not have a hair-like pappus.

Key features

Finely-divided leaves; noticeable lack of odor; daisy-like heads at full maturity resemble a badminton shuttlecock

Similar species

Wild chamomile *Matricaria chamomilla*; stinking mayweed *Anthemis cotula*

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

reduced tillage

forage crops, hay fields

rangeland

irrigation

turf

wetland

roadsides

waste areas

Seedling

Juvenile

Flower: head

Mature

Similar species: *Wild chamomile*

Shepherd's-purse

Capsella bursa-pastoris

Family

Mustard *Brassicaceae*

Seedling

Cotyledons are oval to ovate with rounded tips. The first true leaves are entire to slightly lobed. Subsequent leaves have variable margins. Leaves are petioled and covered with star-shaped hairs intermingled with unbranched hairs.

Juvenile

The basal rosette leaf margins vary from slightly to deeply lobed. They can even be divided to the midrib. The extreme variability of the leaf margins can cause confusion when identifying this weed. Stems of shepherd's-purse are sparsely branched and usually covered with star-shaped hairs. Stem leaves are alternate, sessile and linear to lanceolate with entire margins or a few scattered teeth. The basal lobes of the leaves are pointed and clasp the stem.

Mature

The raceme is made up of small white flowers, each with four petals. The pod developing from each flower is triangular and flat. There is a notch at the tip with a small beak in the centre of the notch. Seeds are attached to both sides of the membranous wall inside the pod. The seeds are oblong, appear orange to reddish and have a pitted surface.

Key features

Star-shaped hairs on leaves and stems; upper leaves that clasp the stem with pointed basal lobes; triangular, flat seedpods

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling: *star-shaped hairs on leaf surface*

Juvenile: *rosette*

Key feature: *triangular seedpod*

Mature: *showing racemes*

Flower: *small, white, four petals*

Spotted Water-hemlock

Cicuta maculata

Family

Parsley *Apiaceae*

Seedling

Seedlings are seldom noticed.

Juvenile

Spotted water-hemlock is extremely poisonous to humans and livestock. Stock poisonings often occur in the spring when young plants are more easily pulled out of the ground. The poison is an oily liquid called cicutoxin that is found mainly in the fleshy roots and in horizontal cavities at the base of the stem. The tall, erect stem has purple spots or streaks. Spotted water-hemlock is often confused with the related but non-poisonous water-parsnip, *Sium suave*. Both species thrive in wet, marshy habitats. The leaves of water-parsnip are compound pinnate with finely toothed leaflets. In contrast, spotted water-hemlock leaves are twice-compound pinnate with coarsely toothed leaflets. Leaves are alternately arranged on the stem and have flared petioles at the base.

Mature

The tiny white flowers of both species are grouped into compound umbels. There are usually no bracts at the base of the main umbel of spotted water-hemlock, but there could be one or two. On the other hand, water-parsnip has several to many pointed bracts at the base of each main umbel. An individual flower can produce two brown seeds. Each seed has five ribs on one surface and is flat on the other side.

Key features

Purple-streaked stem; twice-compound pinnate leaves with coarsely toothed leaflets; bracts usually absent under main umbel

Similar species

Water-parsnip *Sium suave*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas		

Flower: *compound umbel of tiny, white flowers*

Toxin-filled cavities

Key feature: *twice compound pinnate leaf*

Similar species: *Water-parsnip – leaf*

Stinkweed

Thlaspi arvense

Other names

Penny cress, Frenchweed

Family

Mustard *Brassicaceae*

Seedling

Cotyledons are oblong with short stalks. The first true leaves are also oblong and have a smooth surface. Stinkweed may be distinguished from shepherd's-purse at the seedling stage by the absence of hair on the true leaves.

Juvenile

Stinkweed typically forms a basal rosette. The rosette leaves are petioled and have an entire to wavy margin. They give off a distinctive, strong, turnip-garlic odor when crushed. Stems are smooth, erect and often branched above. The upper stem leaves are smaller than the rosette leaves and have shallow, irregular teeth. They are sessile and often clasp the stem. Plants turn yellow soon after flowering.

Mature

Tiny, four-petalled white flowers are located in racemes at the tops of branches. Each flower produces a circular, flattened seedpod with "wings" and a narrow, deep notch at the top. At maturity, the seedpods turn brown and split open to reveal oval, reddish-brown to black seeds. The seeds are covered with rows of concentric ridges and resemble a human fingerprint.

Key features

Smooth surface of leaves and stems; turnip-garlic odor; flattened, winged, notched seedpod

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland	roadsides	waste areas
reduced tillage	irrigation			
forage crops, hay fields	turf			

Seedling

Juvenile: rosette

Mature

Flower: flowers and seedpods on raceme

Key feature: notched seedpod

White Clover

Trifolium repens

Other names

Dutch clover

Family

Pea *Fabaceae*

Seedling

Cotyledons are oval, and the first true leaf is long-stalked with a single blade indented at the tip. All subsequent leaves produced are trifoliate.

Juvenile

White clover is a shallow-rooted plant with extensive, creeping stems. When the stem nodes encounter the ground, they frequently root to form new shoots. The trifoliate leaves are characterized by very long petioles. The three leaflets are somewhat variable in shape, but typically broadest at the tip, which is often shallowly-notched. There is usually a pale, inverted V-shaped mark on the upper surface of each leaflet, but no hair. Each leaf has a pair of small, pointed stipules at the base that form a tube around the stem. While white clover is an important legume species in certain parts of Canada, in the prairies, it grows mostly as a weed of turfgrass.

Mature

Dense, round, head-like racemes are produced on long stalks. Each raceme contains between 20 and 40 white to pinkish pea-like flowers. The flowers form small pods with two to five squarish, yellow seeds.

Key features

Extensive creeping stems; trifoliate leaves with leaflets having an inverted V-shaped mark and shallow notched tip; dense, head-like racemes of white to pinkish pea-like flowers

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Mature

Flower: *pea-like flowers in dense racemes*

Key feature: *creeping stems*

Infestation

Key feature: *trifoliate leaf*

White Cockle

Silene alba

Family

Pink *Caryophyllaceae*

Seedling

Cotyledons are stalked, ovate in shape and somewhat constricted at the tips. The true leaves are softly hairy on both surfaces and margins. They are oppositely arranged, elliptical and have short, hairy petioles.

Juvenile

White cockle has a fleshy tap root and forms a rosette in the first year. Stems are erect, covered with hair and swollen at the nodes. The middle and upper leaves are sessile, opposite, elliptical and softly hairy. The stems and leaves of white cockle are not sticky to the touch like those of night-flowering catchfly.

Mature

The flowers of white cockle are large and showy, each with five notched white petals. Flowers on one plant are either all male or all female. The calyx surrounding male flowers is hairy with ten veins. The hairy calyx of female flowers has twenty veins, five of which are very distinct, the others less so. The female flower forms a light brown capsule that opens with ten teeth. The seeds are greyish, bumpy and kidney-shaped. Night-flowering catchfly has calyx veins that are more distinctly branched than white cockle. Also, the capsule of night-flowering catchfly opens with only six teeth. The overall stickiness of mature night-flowering catchfly is the easiest way to distinguish it from white cockle.

Key features

Opposite, softly hairy leaves; swollen nodes; large, white, showy flowers; capsule opening by ten teeth

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Juvenile

Mature

Flower

Infestation

Wild Buckwheat

Polygonum convolvulus

Family

Buckwheat *Polygonaceae*

Seedling

Cotyledons are linear and positioned at 120 degrees from each other. Tartary buckwheat can be distinguished at the seedling stage by its round cotyledons. The true leaves of wild buckwheat are arrow-shaped with pointy tips and pointed basal lobes. Leaves are alternately arranged and petioled with entire margins.

Juvenile

Wild buckwheat is tap-rooted and has a twining growth habit. The slender stem branches near the base and has ocreas at the stem nodes. The vines become entangled with the crop, causing lodging and making harvest difficult. Tartary buckwheat can be distinguished by its erect stems. Field bindweed, also a twining weed, has perennial, creeping rootstocks and lacks ocreas at the stem nodes.

Mature

Small flowers are produced in axillary and terminal racemes. In contrast, flowers of field bindweed are large and showy with white to pink funnel-shaped corollas. Although wild buckwheat flowers lack petals, there are five whitish-green sepals that tightly enclose a single seed. Each seed is dull, dark brown to black and triangular in cross-section. Tartary buckwheat seeds protrude from the sepals and are larger than seeds of wild buckwheat.

Key features

Twining stems; ocreas at stem nodes; arrow-shaped leaves with pointy tips

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Juvenile

Key feature: *ocrea*

Flower: *flowers in raceme*

Twining growth habit

Wild Caraway

Carum carvi

Family

Parsley *Apiaceae*

Seedling

Seedlings are not commonly seen in the field.

Juvenile

Wild caraway produces a large brown tap root and lacy-looking rosette in the first year. Tall, smooth, slender stems emerge from the crown the following spring. Each stem has pinnately compound leaves with leaflets that are once or twice divided into narrow, threadlike segments. This weed is a relatively new problem on the prairies, but has been found moving into ditches and roadsides adjacent to cultivated land.

Mature

Tiny white flowers are located in loose, compound umbels found at the top of the plant and at the ends of side branches. Wild caraway is sometimes mistaken for its relative, wild carrot. The two species can be distinguished easily by the bracts found beneath the inflorescence. Wild caraway is either bractless or has only one to three bracts while wild carrot has a distinct whorl of bracts, many of which are further divided. Wild caraway also tends to flower in spring or early summer, which is much earlier than wild carrot. The seeds of wild caraway are oblong, brown with five tan ribs and conspicuously scented.

Key features

Large, brown tap root; smooth pinnately compound leaves with leaflets of threadlike segments; white-flowered compound umbels with few or no bracts beneath

Similar species

Wild carrot *Daucus carota*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland	wetland
reduced tillage		irrigation	roadsides
forage crops, hay fields		turf	waste areas

Seedling

Key feature: *compound leaf*

Mature

Flower: *compound umbel of tiny, white flowers*

Infestation

Wild Licorice

Glycyrrhiza lepidota

Family

Pea *Fabaceae*

Seedling

Cotyledons are oblong and thickened. The first two true leaves have a single, oval blade on a short stalk. The surface of the leaves is covered with minute depressions while the margins have short, scattered hairs. All the remaining leaves produced are trifoliate.

Juvenile

Tall, erect, somewhat hairy stems arise from extensive, deep, creeping roots. The roots are highly aromatic when dug up, and the distinctive licorice scent gives this weed its common name. Leaves are pinnately compound with 11 to 19 elongated leaflets. The leaflets are broadest in the middle and have a prominently pointed tip. They are smooth on the upper surface with tiny, glandular dots on the underside.

Mature

Wild licorice forms erect, dense racemes on short stalks in the axils of upper leaves. The flowers are yellowish-white and pea-like in structure. Each flower forms a reddish-brown pod covered with hooked bristles and resembles a bur. The pods contain three to five large seeds.

Key features

Aromatic creeping roots with licorice scent; pinnately compound leaves with 11 to 19 leaflets, glandular-dotted below; reddish-brown bur-like pods

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland	wetland	
reduced tillage		irrigation	roadsides
forage crops, hay fields		turf	waste areas

Mature

Key feature: *bur-like pods*

Pinnately compound leaves

Infestation

Wild Tomato

Solanum triflorum

Other names

Cutleaf nightshade

Family

Nightshade *Solanaceae*

Seedling

Cotyledons are linear with a deep, noticeable midrib. The first true leaves have petioles and are ovate with wavy margins.

Juvenile

Wild tomato is a low spreading, mat-forming weed. Its stems are usually branched at the base and may have a smooth or slightly hairy surface. Older leaves are deeply lobed and typically sparsely hairy.

Mature

The flowers on wild tomato look very much like the flowers of its domestic relatives, tomato and potato. They are usually found in clusters of three in leaf and branch axils. Each flower has five white petals joined together in the shape of a star. The flowers produce smooth, round, green berries resembling miniature green tomatoes. As tempting as they may be, the berries should not be eaten since there have been some reports of people being poisoned by them. Inside each berry are light brown, ovate seeds that look exactly like tiny tomato seeds.

Key features

Low spreading, mat-forming growth habit; deeply lobed leaves, tomato-like flowers and berries

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling

Juvenile

Mature: *low, spreading mat*

Flower: *tomato-like*

Key feature: *berries*

Yellow

Flowered Species

Absinth

Artemisia absinthium

Other names

Wormwood

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are ovate and short-stalked.

The true leaves are covered with fine silvery hairs. The margins have rounded-to-blunt lobes. Leaves have a strong sage-like odor. Young seedlings of biennial wormwood may appear similar but have sharply-toothed lobing on the true leaves.

Juvenile

Absinth has a fibrous root system and develops a large rosette with long-petioled basal leaves. Leaf blades are deeply divided into many oblong, blunt-tipped segments. The covering of long silky hairs gives the leaves a light greenish-grey appearance. Stems are tall, erect and hairy. Leaves on the stem are alternate and decrease in hairiness towards the top of the plant. Upper leaves are almost sessile. They are much smaller than the basal leaves and have fewer lobes or none.

Mature

Flower heads are small, numerous and droop from short stalks in the leaf axils of upper branches. Each head holds many yellowish flowers, each capable of producing one seed. The receptacle and several whorls of bracts are densely hairy. Seeds are shiny brown with fine, white streaks and lack a pappus. Older plants become woody at the base. Absinth can be controlled by cultivation but can be weedy in overgrazed pastures and hay fields. It can produce tainted milk, and the pollen causes discomfort to some people.

Key features

Light greenish-grey leaves due to long silky hairs; leaf margins with rounded or blunt lobes; strong, sage-like odor

Similar species

Biennial wormwood *Artemisia biennis*

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling

Juvenile

Mature

Similar species: *Biennial wormwood* – seedling

Argentine Canola

Brassica napus

Family

Mustard *Brassicaceae*

Seedling

Cotyledons are broad and kidney-shaped. The undersides are smooth, in contrast to the wrinkled undersides of Polish canola cotyledons. The petioled leaves emerge one at a time. They have irregularly lobed margins and are usually hairless. The leaves are bluish-green and have a waxy surface.

Juvenile

Leaves of the basal rosette are large and irregularly lobed with the terminal lobe much larger than lobes near the base of the leaf. As in the seedling, they appear bluish-green and waxy. The stem is usually branched but less so than Polish canola. Typically, Argentine canola has about six branches. The stem leaves of Argentine canola are alternate, entire or sparsely toothed. The upper leaves are sessile and partially clasp the stem. Upper leaves of Polish canola fully clasp the stem.

Mature

The flowers are produced in terminal and axillary racemes. Each flower has four bright yellow petals. The petals are larger and lighter in color than Polish canola petals. The unopened, green flower buds on the raceme are visible above the opened flowers. Unopened flower buds of Polish canola remain below the opened flowers. Each flower produces a long, narrow pod with a membrane in the middle and seeds on either side. There is a short beak at the top of the pod. At maturity, the pods shatter easily to release seed. Seeds are spherical, brown to black and have a minutely rough surface. They are larger than Polish canola seeds.

Key features

Waxy, bluish-green leaves; upper stem leaves sessile but not fully clasping the stem; unopened flower buds visible above opened flowers

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Juvenile

Flower: bright yellow

Mature

Key feature: flower buds above opened flowers

Ball Mustard

Neslia paniculata

Family

Mustard *Brassicaceae*

Seedling

Cotyledons are petioled and almost round with shallow indentations at the apex. The first true leaves are oblong with round to slightly pointed tips.

Juvenile

Ball mustard forms a rosette of long, narrow leaves with entire or nearly entire margins. Stems are erect and often much-branched. The upper leaves are more arrow-shaped than the basal leaves and clasp the stem. The entire plant is usually covered with small, star-shaped hairs.

Mature

Bright yellow flowers are located in dense racemes at the ends of stems. Each flower is four-petalled and produces the characteristic ball-like seedpod that gives this weed its common name. The seedpods are long-stalked and yellowish-brown with a dimpled surface. Each pod contains a single seed although they rarely break open to expose the seed. Ball mustard seeds are difficult to separate from the seeds of Argentine canola.

Key features

Entire or nearly entire leaves; star-shaped hairs over the entire plant; ball-like seedpods

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling

Juvenile

Key feature: *ball-like seedpod*

Clasping upper leaves

Flower: *flowering raceme*

Black Henbane

Hyoscyamus niger

Family

Nightshade *Solanaceae*

Seedling

Cotyledons are long-stalked and oval with rounded tips. The first true leaves are ovate with wavy margins.

Juvenile

Black henbane typically develops a large tap root and basal rosette in the first year. The leaves are sessile and generally ovate to oblong with an irregularly lobed or toothed margin. The rosettes bolt to form very coarse, often tall stems. The whole plant is covered with long, sticky hairs that give off a foul smelling substance. Black henbane is a highly poisonous weed, but fortunately, the strong, unpleasant scent deters grazing animals.

Mature

Highly unusual, bell-shaped flowers are formed in crowded, one-sided spikes near the top of the plant. The petals have a conspicuous network of purple veins on a yellowish background and once seen, they are never forgotten. Each flower forms a light brown capsule full of small, kidney-shaped seeds. At full maturity, the capsules open at the top by a lid. They greatly resemble the shell of a peanut both in shape and texture.

Key features

Large irregularly lobed leaves; long, foul-smelling, sticky hairs; purple-veined yellowish flowers in one-sided spikes

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Juvenile: *rosette*

Flower: *purple-veined yellowish*

Mature

Seed capsules on spike

Black Medick

Medicago lupulina

Family

Pea *Fabaceae*

Seedling

Cotyledons are oval and stalkless. The first true leaf is simple, long-stalked and oval. The second leaf and all remaining leaves are compound with three small leaflets.

Juvenile

Black medick is prostrate with several to many wiry stems branching out from the base to form a wide, spreading mat. Both the leaves and stems are usually hairy. The trifoliate leaves are alternate with two sharply-pointed stipules at the base. Each leaflet has shallow teeth on the upper margin and a pointed tip. The middle leaflet has a definite stalk while the other two are sessile.

Mature

Flowers are very small, yellow and pea-like. They occur in dense, head-like racemes near the ends of the branches. The fruits are small, kidney-shaped seedpods that are black at maturity. Each seedpod contains a single yellow to olive-green seed.

Key features

Prostrate growth habit; trifoliate leaf with stalked, central leaflet; small, yellow pea-like flowers

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Juvenile: *showing prostrate growth*

Seedpods

Mature

Flower: *dense flower head*

Common Groundsel

Senecio vulgaris

Family

Sunflower Asteraceae

Seedling

Cotyledons are oblong with a grooved stalk. The first true leaves are oval with upward-pointing teeth along the margin. The petioles are also grooved, and the underside of the blades is sometimes pinkish.

Juvenile

Common groundsel forms a basal rosette of leaves that are typically lobed with a toothed margin. They have a slightly fleshy texture and are either smooth or sparingly hairy. The leaves have a prominent midrib underneath. Short, erect stems are produced that are branched and somewhat fleshy. The upper leaves are sessile and have irregularly lobed and toothed margins.

Mature

Small, cylindrical heads are in clusters at the ends of stems and branches. Each head has yellow disk flowers barely visible above the surrounding long, black-tipped, involucral bracts. A narrow tan-colored seed covered with short white hairs is produced by each flower. The seeds have a conspicuous white pappus.

Key features

Somewhat fleshy, lobed leaves; short, somewhat fleshy stems; black-tipped involucral bracts on the heads

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Juvenile

Mature

Flower: *flower heads and seeds*

Common Mullein

Verbascum thapsus

Family

Snapdragon *Scrophulariaceae*

Seedling

The cotyledons are ovate with a short stalk and tiny hairs. The first true leaves are oval, petioled and wooly. Common mullein is found on roadsides and pastures. It usually grows in sandy or gravelly soils. Seedlings are not commonly seen.

Juvenile

Common mullein is a tap-rooted biennial with a large rosette in the first year. The rosette is made up of large, whitish-green, wooly leaves that are narrower near the base.

Mature

The second-year flowering stem is wooly, tall and usually unbranched. The white-wooly leaves hug the stem, giving the stem a winged appearance. The yellow, saucer-shaped flowers are five-lobed and arranged in a dense, wooly, spike-like raceme at the top of the stem. Each flower develops into a spherical capsule with two compartments filled with many seeds. Each seed is six-sided, light to dark brown and has a deeply grooved surface. The dead flower stalks may remain standing for one or two years.

Key features

Large, white-wooly leaves; second year stem has winged appearance; yellow flowers densely produced in a thick, wooly, spike-like raceme

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Flower: *yellow flowers in raceme*

Juvenile: *rosette*

Key feature: *white-wooly leaves*

Infestation

Dandelion

Taraxacum officinale

Family

Sunflower *Asteraceae*

Seedling

A dandelion seedling has smooth, oval cotyledons. The first true leaves are oval with sparse, irregular toothing on the margins.

Juvenile

Dandelion has a deep, penetrating tap root that may be branched. The leaves are in a basal rosette throughout the life cycle, and there is no elongated, leafy stem. Margins of dandelion leaves are irregularly lobed and toothed, a feature that varies considerably from plant to plant. Young dandelions can be distinguished from perennial and annual sow-thistles by the lack of weak prickles on the margins. The leaf surface may be smooth or sparsely hairy. The entire plant is filled with a white, sticky juice.

Mature

Solitary flower heads are produced at the top of hollow, leafless stalks. The ray florets are bright yellow and showy. Involucral bracts around each head are green, linear and in two whorls. The outer whorl has shorter bracts that are bent backwards. The seed is yellowish-brown, long and slender with a long beak and a pappus for wind dispersal. Dandelions are traditionally a weed of lawns and turf but are becoming a problem in reduced tillage systems.

Key features

Hollow, leafless flower stalks with bright yellow flower heads; sticky, white latex throughout plant; leaves all basal

% of fields
where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

reduced tillage

forage crops, hay fields

rangeland

irrigation

turf

wetland

roadsides

waste areas

Seedling

Juvenile

Infestation

Mature

Key feature: *seed heads on leafless stalks*

Dog Mustard

Erucastrum gallicum

Family

Mustard *Brassicaceae*

Seedling

Cotyledons are oval with a small indentation at the tip. The first true leaves have simple, white hairs and wavy to slightly-toothed margins.

Juvenile

Dog mustard forms a rosette with somewhat hairy, oblong leaves. The leaves are deeply divided into coarse lobes with margins that may be wavy or irregularly-toothed. The basal leaves resemble the lower leaves of tumble mustard. Stems are erect and may be much-branched. The lower stems are covered with short, stiff, downward-pointing hairs.

Mature

Flowers are produced in elongated racemes at the top of the plant. Each flower has four pale yellow petals. The seedpods are long and narrow, four-sided and have a short beak. Dog mustard is the only weedy mustard that has flowers and seedpods originating from the axils of upper stem leaves. Oval, reddish-brown seeds are produced in a single row within each half of the seedpod.

Key features

Pinnately-divided leaves with coarse lobes; downward-pointing hairs on the lower stems; flowers and seedpods in the axils of upper leaves

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Juvenile: *rosette*

Flower: *pale yellow*

Mature

Key feature: *seedpods in upper leaf axils*

Early Yellow Locoweed

Oxytropis sericea

Family

Pea *Fabaceae*

Seedling

Early yellow locoweed has oval cotyledons and compound, silky-haired true leaves. The first true leaf has one leaflet only, but the third true leaf will usually have three leaflets. The leaves have long petioles with hairy stipules at the base. Growing on sandy, eroded pastures and roadsides, the seedlings are seldom noticed.

Juvenile

Early yellow locoweed has a thick tap root. There may be one or more basal rosettes in a clump, all coming from the branched, woody stem below ground. The leaflets are covered with soft, silky hairs giving the leaves a whitish-green appearance. There are 11 to 17 leaflets per leaf, arranged oppositely or nearly so. The stipules at the base of the leaves are also hairy. The plant is poisonous, causing erratic behaviour and even death in livestock.

Mature

Blooming early in May, the lemon-yellow or whitish-yellow flowers are in racemes at the top of leafless flower stalks. The “keel” of the pea-type flower has a prominent protrusion or “beak.” The calyx has both white and dark-colored hairs. Pods are leathery, covered with hair and contain kidney-shaped seeds.

Key features

Basal, whitish-green, hairy leaves with 11 to 17 leaflets; early May bloomer; pale to lemon-yellow flowers with a prominent beak on the “keel”

Similar species

It has not been clearly established which of the numerous related locoweed species are poisonous.

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Flower: *flowers in racemes*

Mature

Infestation

Flixweed

Descurainia sophia

Other names

Tansy mustard

Family

Mustard *Brassicaceae*

Seedling

Unlike many mustard weeds, flixweed has narrow, stalked cotyledons. The first true leaves are usually three-lobed, hairy and opposite. It is very easy to confuse flixweed with scentless chamomile at the seedling stage. Scentless chamomile has rounder, sessile cotyledons and smooth, usually five-lobed first true leaves.

Juvenile

Flixweed produces a basal rosette with greyish-green leaves that are finely divided. Stems are erect and often much-branched. The upper leaves are similar to, but much smaller than the rosette leaves. Both the stems and leaves are covered with star-shaped hairs.

Mature

Tiny, four-petalled, yellow flowers are produced in racemes at the tops of the stems. Long, narrow seedpods are formed on short stalks. Each seedpod contains two rows of minute, orange-colored seeds. Flixweed can be distinguished from the two related tansy mustard species by the seedpods. They are shorter, wider and longer-stalked in the tansy mustards than in flixweed.

Key features

Finely divided leaves; greyish-green color; narrow, short-stalked seedpods

Similar species

Green tansy mustard *Descurainia pinnata*; grey tansy mustard *Descurainia richardsonii*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Juvenile: rosette

Mature: plant with raceme

Key feature: finely divided leaves

Infestation

Goat's-beard

Tragopogon dubius

Other names

Western salsify, Johnny-go-to-bed-at-noon

Family

Sunflower *Asteraceae*

Seedling

Both cotyledons and true leaves are long and linear in shape. Seedling leaves are sessile with an entire margin.

Juvenile

Goat's-beard has a deep, thick tap root and leaves that are long, linear and clasping at the base. Very young leaves are covered with soft, downy hair. Older leaves almost appear grass-like, but with a more smooth and fleshy texture. Stems are smooth, hollow and contain a milky juice. They are usually tall and frequently have dead leaves around the base in the second year.

Mature

The heads of goat's-beard are dandelion-like, but much larger and more flat-topped. They are produced singly at the ends of stems and branches and usually face toward the morning sun. The heads will turn to follow the sun's path, but by afternoon, they are completely closed up. The stems directly below the heads are thickened, which is a key way to distinguish this weed from other species of goat's-beard. The ray flowers are a pale lemon yellow and are enclosed by long, narrow, involucre bracts. The outer flowers tend to open first so that those in the center may, at first glance, appear to be disks, although they are not. When seeds are produced, the heads are transformed into large, white, fluffy spheres. Each seed is narrow and ribbed with a long beak attached to an umbrella-like, feathery pappus.

Key features

Narrow, grass-like leaves; milky sap; large, white, fluffy seed heads

Life Cycle			
annual	winter annual	biennial	creeping perennial
simple perennial			
Habitat			
conventional tillage	rangeland	wetland	
reduced tillage	irrigation	roadsides	
forage crops, hay fields	turf	waste areas	

Seedling

Flower: *head*

Key feature: *large fluffy seed head*

Juvenile

Mature

Golden-bean

Thermopsis rhombifolia

Other names

Buffalo-bean

Family

Pea *Fabaceae*

Seedling

Seedlings are not commonly seen.

Juvenile

Golden-bean is usually found in large patches formed by stout, creeping rootstocks. Stems are erect, angled and smooth with upper branching. The leaves are alternate and trifoliate with two large, ovate stipules at the stem node. The leaflets may have flattened, silky hairs or be almost hairless.

Mature

Golden-bean is very common and one of the earliest bloomers on the prairies. It is seen in early May in rangelands and along roadsides. The pea-type flowers are bright, golden-yellow in short racemes. Each flower develops into a flattened, curved pod with several seeds. The pods, seeds and flowers of golden-bean have been reported to cause sickness in children, and the species is considered toxic to livestock.

Key features

Found in patches; bright, golden-yellow flowers blooming in early May; curved seedpods

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Juvenile

Flower: *golden-yellow, pea-like*

Key feature: *curved seedpod*

Mature

Gumweed

Grindelia squarrosa

Other names

Curlycup gumweed

Family

Sunflower *Asteraceae*

Seedling

Seedlings are not commonly seen.

Juvenile

Gumweed is a biennial or short-lived perennial with erect, freely-branching stems. They are hairless and frequently purple-tinged. The leaves are narrow with entire to toothed margins. The lower leaves are stalked while the upper are sessile and clasping. The leaves are hairless but covered with small glands that make the surface appear spotted. The entire plant is bitter-tasting and avoided by livestock.

Mature

The heads are small, daisy-like and occur in flattened clusters at the ends of stems and branches. Each head has bright yellow ray and disk flowers surrounded by narrow bracts with the tips curled backwards. The bracts are covered with a resinous “gummy” substance that makes them extremely sticky to the touch. The seeds of gumweed are cream-colored, ribbed and lack a noticeable pappus.

Key features

Purple-tinged stems; glandular, spotted leaves; sticky-gummy bracts with a curled tip

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Mature

Key feature: *gummy, curled bracts*

Flower: *head*

Toothed leaf margin

Leafy Spurge

Euphorbia esula

Family

Spurge *Euphorbiaceae*

Seedling

Cotyledons are linear in shape, pale green and veiny underneath. The seedling stem often has a few buds just above the soil surface. The first two leaves that develop appear to be opposite. Subsequent leaves are alternate, spiraling around the stem.

Juvenile

Stems are erect and hairless with sessile leaves that can be quite densely crowded. The leaves have entire margins and are linear to lanceolate. All parts of the plant contain a white, sticky latex. Although leafy spurge does produce viable seed, propagation and spread of the plant is mainly by extensive vertical and horizontal roots with buds that can produce new shoots. The vertical part of the root system can extend very deeply into the soil profile. Leafy spurge is poisonous to most livestock, but not to sheep.

Mature

Leafy bracts are at the base of the flowering clusters located at the top of the stem and upper branches. Flowers lack petals and are found in groups of several male flowers located near one female. The pair of greenish-yellow bracts under each grouping of flowers are almost round with pointy tips. The seed capsule produced from each female flower has three compartments, each with one seed. Capsules explode at maturity, ejecting the seed a few metres from the plant. Seed color varies from yellow to grey to brown. There is also a distinct line on one side of the seed and a yellowish bump near the base.

Key features

White latex in all parts of the plant; deep and spreading roots with buds; greenish-yellow bracts underneath the flower groupings

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Juvenile

Flower: clusters

Mature

Key feature: white latex in stem

Seed capsule and leafy bracts

Narrow-leaved Hawk's-beard

Crepis tectorum

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oval in shape with short stalks. The first true leaves are more elongated than the cotyledons and have prominent petioles. They often have a few distinct, downward-pointing teeth on the margins. Narrow-leaved hawk's-beard may be difficult to distinguish from dandelion at the seedling stage.

Juvenile

Narrow-leaved hawk's-beard forms a basal rosette, often in late summer or fall. The rosette leaves are narrow, but have variable margins ranging from sparsely-toothed to deeply lobed. Identification of this weed at the rosette stage is critical since the winter annual form is rapidly increasing across the prairies. Stems are erect and usually branched. They may be smooth or hairy and exude a milky juice when broken. The leaves on the upper stems are narrow and entire. They are sessile and sometimes also have pointed lobes that clasp the stem.

Mature

The bright yellow ray flowers are in small dandelion-like heads in clusters at the top of branches. The involucral bracts surrounding the receptacle are in several rows. The outer bracts are much shorter than the inner bracts. Each flower produces a dark purple seed with ten prominent ribs. The seeds are wind-dispersed with the aid of a hair-like pappus.

Key features

Winter annual habit; narrow stem leaves; milky juice in stem

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Juvenile: *rosette*

Infestation

Mature

Flower: *dandelion-like flower head*

Orange Hawkweed

Hieracium aurantiacum

Other names

Devil's paint-brush

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are round to oval in shape. The first true leaves are hairy with entire to finely-toothed margins.

Juvenile

Like strawberry, orange hawkweed has leafy runners that can root and produce new plants. Its leaves are mostly in a basal rosette. They are rounded at the tip and then narrow at the base into winged petioles. Both surfaces of the leaves are usually covered with stiff hairs. Stems are hairy, nearly leafless and contain a bitter, milky sap.

Mature

Dandelion-like heads are located in flat-topped clusters at the tops of stems. The ray flowers are orange-red, making this weed very easy to identify when mature. There are several other related hawkweed species in Canada, but all have yellow flowers. The involucral bracts surrounding the flowers appear blackish due to the gland-tipped hairs. Each flower produces one purplish-black, narrow, ribbed seed with a hair-like pappus.

Key features

Leafy runners; leafless, hairy stems; clusters of orange-red dandelion-like heads

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Juvenile: *basal rosette*

Infestation

Mature: *leafless, wiry stems*

Flower: *flower heads*

Pasture Sage

Artemisia frigida

Other names

Fringed sagebrush

Family

Sunflower *Asteraceae*

Seedling

The seedling has tiny oval cotyledons and hairy, silvery-grey leaves with slightly lobed margins.

Juvenile

Plants are low to the ground, with a woody base. Stems branch from the base of the plant, producing a mat-like appearance. The leaves are hairy, silvery-grey, finely divided and have a sage aroma. The lower leaves are petioled while the upper leaves are sessile, alternate and less finely divided. Pasture sage is an indicator of overgrazing, and the weed is quite drought tolerant.

Mature

Many small, yellowish flower heads are found on short flower stalks from upper leaf axils. The numerous, nodding heads have fertile florets in the centre and hairy bracts around the outside. Tiny, smooth, grey-brown seeds are produced in large amounts. There is no pappus.

Key features

Short statured plant with sage odor; silvery-grey, hairy leaves with finely divided margins; numerous, small, nodding heads

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Finely divided, silvery-grey leaves

Juvenile

Infestation

Mature

Flower: *small axillary flower heads*

Perennial Sow-thistle

Sonchus arvensis

Family

Sunflower Asteraceae

Seedling

Cotyledons are oval with slight indents at the tip. The first true leaves are spatulate with soft prickles along the margins. Seedlings are almost impossible to distinguish from other species of sow-thistle.

Juvenile

Perennial sow-thistle has extensive, horizontal, deep, creeping roots. These roots are light-colored, fleshy and easily broken. Each piece of broken root containing a bud can produce a new plant and potentially start a new infestation. Juvenile plants form rosettes of basal leaves that vary in shape from shallow to deeply lobed. The basal leaves may resemble dandelion but instead have winged stalks and prickles on the margin. Often, discrete patches of rosettes are found in an area, suggesting the perennial nature of this weed. Plants have erect stems that are hollow and contain a milky sap. They are smooth at the base but often have orange hairs near the top. Upper leaves are typically more deeply lobed than basal leaves. Although they also clasp the stem, the lobes at the base of the leaf are smaller than in the annual sow-thistle species.

Mature

Heads are found in clusters at the ends of branches. Each head is dandelion-like with bright yellow, ray flowers. The involucre bracts surrounding the heads are covered with dense, orange-colored hair. A related perennial sow-thistle species, *Sonchus uliginosus*, is similar in appearance but completely smooth on the upper stems and bracts. The heads of perennial sow-thistle are larger and more flask-shaped than those of the annual species. Seeds are brown with both lengthwise and crosswise ribs giving them a wrinkled look. They have a pappus and are easily moved by wind.

Key features

Fleshy, white, underground creeping roots; lobed leaves with a prickly margin; upper stems and bracts with orange hair

Similar species

Perennial sow-thistle *Sonchus uliginosus*

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

reduced tillage

forage crops, hay fields

rangeland

irrigation

turf

wetland

roadsides

waste areas

Seedling

Juvenile: *group of rosettes*

Infestation

Mature

Flower: *dandelion-like flower head*

Pineappleweed

Matricaria matricarioides

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oblong and stalkless. The first true leaves are long and narrow with scattered lobes along the margin.

Juvenile

Juvenile plants have leaves that are finely-divided, smooth and slightly fleshy to the touch. Pineappleweed greatly resembles scentless chamomile at this stage, but can be easily distinguished by the distinct pineapple odor when the leaves are crushed. Its stems are short, smooth and much-branched.

Mature

Ovate heads comprised of tiny, yellowish-green disk flowers are produced at the tops of branches. They lack the white, ray flowers associated with scentless chamomile and resemble miniature pineapples instead of daisies. Each flower produces a single, tiny, olive-to-brown seed with no obvious pappus.

Key features

Finely-divided leaves with pineapple odor; short, smooth stems; pineapple-shaped heads

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling

Juvenile

Flower: *pineapple-shaped flower heads*

Mature

Polish Canola

Brassica rapa

Other names

Turnip rape

Family

Mustard *Brassicaceae*

Seedling

Polish canola cotyledons are broad, kidney-shaped and wrinkled underneath. The leaves are green and hairy, lacking the waxiness and bluish-green tinge of Argentine canola. The leaves are alternate and petioled with irregularly and shallowly lobed margins. Although the base of the stem is usually hairy, this feature varies with cultivar. Polish canola is difficult to distinguish from wild mustard at the seedling and early juvenile stages. The base of the stem of wild mustard is usually more bristly-hairy than Polish canola. In addition, wild mustard stem nodes may develop purple coloration, even at a young age.

Juvenile

The basal rosette of Polish canola has green, hairy, petioled leaves. Typically, the rosette is smaller than the rosette of Argentine canola. Once the plant bolts, Polish canola is more branched than Argentine canola and may have up to 20 branches. The stem leaves are alternate, entire, sessile and totally clasp the stem with rounded basal lobes. In contrast, wild mustard stem leaves are coarsely toothed and sessile, but not clasping. Also, wild mustard stem nodes develop purple blotches while the nodes of Polish canola do not have this coloration.

Mature

Flowers are produced in terminal and axillary racemes. Each flower has four deep yellow petals, which are smaller than those of Argentine canola. Unopened flower buds are located below the opened flowers on the raceme. Each flower develops into a seedpod that is shorter than an Argentine canola pod. The beak at the top of the pod is longer than the beak on Argentine canola pods, and Polish canola seedpods have less tendency to shatter. Mature seeds of Polish canola are mixtures of brown and yellow seeds, which are smaller than Argentine canola seeds.

Key features

Green, hairy leaves; upper stem leaves fully clasping the stem; unopened flower buds below the opened flowers

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Juvenile: *rosette*

Key feature: *upper leaves fully clasping stem*

Flower: *flower buds below opened flowers*

Raceme of mature plant

Prairie Sunflower

Helianthus petiolaris

Family

Sunflower *Asteraceae*

Seedling

Seedlings have oblong cotyledons and oppositely arranged first true leaves. Leaves are petioled and lanceolate to ovate in shape. This plant is often not noticed until it blooms in its usual habitat along roadsides and edges of fields. Prairie sunflower is more frequently found on sandy soils while the related species, annual sunflower, is more typical of heavier soils.

Juvenile

Prairie sunflower is fibrous tap rooted with erect stems covered with short hairs, flattened to the surface. In contrast, stems of annual sunflower are rough and bristly-hairy to the touch. Leaf margins of prairie sunflower are entire or slightly wavy while those of annual sunflower are distinctly toothed. Prairie sunflower leaves are lanceolate to ovate and narrower at the base. Annual sunflower leaves are broadly ovate and often cordate at the base. Leaves of both species are mostly alternate, but lower ones may be opposite.

Mature

Flower heads of prairie sunflower are smaller in diameter than those of annual sunflower. Both species have bright yellow ray flowers around the perimeter. The central disk flowers are purplish to brown and flattened. The involucre bracts around the receptacle of prairie sunflower are lanceolate, gradually tapering to the tip and covered with short hairs. The disk bracts on the receptacle are white-bearded at the tip. Involucre bracts of annual sunflower are long-hairy on the surface with long marginal hairs. Disk bracts are hairy but lack the white-bearded tip.

Key features

Grows in sandy or gravelly locations; narrowly ovate blade tapering to the base with entire to wavy margins; disk bracts white-bearded and long marginal hairs lacking on involucre bracts

Similar species

Annual sunflower *Helianthus annuus*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Flower: *head*

Mature

Key feature: *wavy leaf margins*

Infestation: *roadside infestation*

Prickly Lettuce

Lactuca scariola

Other names

Compass plant

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oval and short-stalked. The first true leaves are shallowly lobed with soft prickles along the margin.

Juvenile

Prickly lettuce forms erect, mostly smooth, whitish-green stems that contain a milky juice. The lower leaves are deeply lobed, and the lobes tend to point backwards. The upper margin on each lobe is spiny-toothed while the lower margin is nearly entire. Upper leaves are usually less lobed and more pointed at the base than lower leaves. Unfortunately, the lobing on the leaves varies greatly from plant to plant. However, there are two very unusual features about the leaves of prickly lettuce to aid in identification. Firstly, most leaves have a single row of sharp prickles on the midrib on the underside. Secondly, the leaves are usually twisted at the base so that the blades are oriented vertically. Often, the leaf tips are aligned to point north-south, giving this weed the reputation of being a “compass plant.”

Mature

The heads are extremely small and occur in much-branched clusters at the top of the plant. Each head has five to twelve yellow ray flowers that turn blue when the plant is dried. A related species, *Lactuca pulchella*, is very similar to prickly lettuce except that its flowers are always blue, and it is a creeping perennial. Each flower of prickly lettuce produces a narrowly oval, ribbed seed with a long beak. The seeds have a white pappus and are readily wind-dispersed.

Key features

Sharp prickles on leaf midribs; vertically-aligned leaves; much-branched cluster of yellow heads

Similar species

Blue lettuce *Lactuca pulchella*

Habitat			Life Cycle
conventional tillage	rangeland	wetland	annual
reduced tillage	irrigation	roadsides	winter annual
forage crops, hay fields	turf	waste areas	biennial
			creeping perennial
			simple perennial

Seedling

Juvenile

Flower: *clusters of flower heads*

Mature

Key feature: *row of prickles on leaf midrib*

Purslane

Portulaca oleracea

Other names

Wild portulaca

Family

Purslane *Portulacaceae*

Seedling

The cotyledons are linear on short stalks. True leaves are succulent, teardrop-shaped and often reddish-green.

Juvenile

Purslane has a thick tap root and a prostrate growth habit. The smooth, fleshy stems form large circular mats on the soil surface. The leaves are sessile, succulent and alternately arranged, sometimes closely alternate. The fleshy, veinless leaves of purslane distinguish it clearly from other prostrate plants. Uprooted or broken-stemmed plants can re-establish roots, making purslane troublesome in gardens and row crops.

Mature

Flowers are yellow and usually have five petals. They are solitary or in small, axillary or terminal clusters and open only on bright, sunny mornings. Each flower develops into a spherical capsule that opens by a lid on top. The numerous seeds are black, shiny and bumpy. They are almost kidney-shaped with a white scar at one end.

Key features

Prostrate mats on soil surface; succulent leaves without veins; stems fleshy and often reddish

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Juvenile: *showing prostrate growth*

Flower: *yellow in leaf axil*

Mature

Rough Cinquefoil

Potentilla norvegica

Family

Rose *Rosaceae*

Seedling

Cotyledons are round. The first true leaves are unifoliate with only a few teeth on the margin.

Juvenile

Rough cinquefoil has erect to spreading stems that are usually branched at the top. They are covered with stiff, often reddish hairs. Leaves are trifoliate with three oblong leaflets and a pair of leaf-like stipules at the base. The lower leaves are petioled while the upper leaves are sessile. This weed is sometimes mistaken for strawberry at the juvenile stage. The leaves on rough cinquefoil have coarse, forward-pointing teeth on the margin surrounding each leaflet while strawberry leaflets are usually only toothed around the tip.

Mature

Flowers are found in clusters on short stalks emerging from leaf and branch axils. Each flower has five, pale yellow petals that are about the same size or even shorter than the green sepals. At first glance, it appears that there are ten sepals, but five of them are actually bracts. The flowers each produce a group of tiny, roughly oval-shaped seeds. Each seed is yellowish-brown with prominent branching ribs.

Key features

Hairy, often reddish stems; trifoliate leaves with toothed margins and stipules; yellow flowers with petals and sepals about the same length

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Key feature: *trifoliolate leaf*

Flower

Mature

Spiny Annual Sow-thistle

Sonchus asper

Other names

Prickly annual sow-thistle

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are fleshy and round to oval. The true leaves are oval to spoon-shaped and petioled. Margins of the true leaves have soft prickles. The seedlings of the two annual and two perennial sow-thistle species commonly found on the prairies are almost impossible to distinguish from each other.

Juvenile

Spiny annual sow-thistle is a tap rooted species with a white, sticky juice throughout the plant. The tap root distinguishes it from perennial sow-thistle, which has extensive, creeping roots. The rosette of spiny annual sow-thistle has glossy leaves with undulating, spiny-toothed margins. The stem is erect and branched at the top. Mid and upper alternate leaves clasp the stem with prominent, rounded, basal lobes. Spines on the leaf margins are unpleasant to touch. Leaves of annual sow-thistle, *Sonchus oleraceus*, are not glossy and have deeply lobed to divided margins. They have a broad, triangular, terminal lobe and are narrow near the base of the leaf, much like a winged petiole. They clasp the stem with basal, pointed lobes. Prickles on the leaf margins are weak, sparse and not unpleasant to the touch. Perennial sow-thistle leaves on the mid-stem have small, rounded, basal lobes but lack these on uppermost leaves.

Mature

The flower heads of both spiny annual sow-thistle and annual sow-thistle are almost triangular with a broad base and a narrow top. The heads of both species have pale yellow ray flowers. Heads of perennial sow-thistle can be distinguished by their much larger size and much deeper yellow flower color. Each flower of spiny annual sow-thistle produces a single, flattened brown seed with several lengthwise ribs. Annual sow-thistle seeds are similar, but have small cross-wrinkles in addition to the lengthwise ribs.

Key features

Tap root; leaves shiny with undulating, spiny-toothed margins and rounded basal lobes; triangular-shaped heads with pale yellow flowers

Similar species

Annual sow-thistle *Sonchus oleraceus*

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Seedling

Juvenile: rosette

Mature

Flower: triangular-shaped heads

Key feature: clasping leaves with spiny-toothed margins

Tall Buttercup

Ranunculus acris

Family

Buttercup *Ranunculaceae*

Seedling

Cotyledons are round to oval and long-stalked. The true leaves have long petioles and blades with three to five rounded lobes.

Juvenile

Tall buttercup has long, erect stems emerging from a short, but thick, rootstock. They have a hairy surface and are often branched near the top of the plant. The basal and lower leaves have long petioles and are softly hairy on both surfaces. The blade is palmately divided into five elongated lobes, each further divided into narrow, toothed segments. The upper leaves are small, nearly stalkless and usually three-lobed. Tall buttercup is highly unpalatable due to the presence of a bitter juice throughout the plant. If grazed, the juice can cause blistering and inflammation of the mouth and digestive system.

Mature

Bright yellow flowers with five rounded petals are located on long stalks at the top of the plant. Each flower produces a cluster of beige-yellow, egg-shaped, flattened seeds with a short, hooked tip.

Key features

Long, erect, hairy stems; hairy, palmately-divided leaves; bright yellow flowers on long stalks

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland	wetland
reduced tillage		irrigation	roadsides
forage crops, hay fields		turf	waste areas

Seedling

Flower: *bright yellow*

Infestation

Mature

Key feature: *palmately divided leaf*

Tansy

Tanacetum vulgare

Other names

Common tansy

Family

Sunflower *Asteraceae*

Seedling

The cotyledons and first true leaves are oval. The early leaves are petioled, hairy and have a slightly lobed margin. As leaves are produced, each one has progressively more lobes than the previous ones. Later, leaves tend to have a smooth surface.

Juvenile

Tansy has short rhizomes and tends to form large clumps of smooth, erect stems. The lower leaves are long-stalked and deeply cut into segments, which are, in turn, divided. They appear almost fern-like. Stem leaves are numerous, sessile and smaller than the basal leaves. Both types of leaves are covered with small glands and the entire plant is aromatic. Tansy is poisonous to livestock, but is usually avoided because of the strong odor.

Mature

Heads are in dense, round to flat-topped clusters. Each head is button-like and composed of many tightly packed, yellow disk flowers. The seeds are grey-tan, ribbed and lack an obvious pappus.

Key features

Short rhizomes; fern-like, aromatic leaves; yellow, button-like heads in clusters

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Juvenile

Clump with previous years' stalks

Infestation

Mature

Flower: *button-like flower heads*

Tumble Mustard

Sisymbrium altissimum

Family

Mustard *Brassicaceae*

Seedling

The cotyledons are stalked and oval with a slight notch at the tip. The first few true leaves are hairy and petioled with slightly lobed margins. Subsequent leaves have deeper lobing.

Juvenile

The basal rosette has large, petioled leaves covered with white, unbranched hairs. Margins are deeply lobed or divided to the midrib. The stem has white hairs at the base or can be hairless. The upper stem leaves are finely divided into narrow, threadlike segments that may have a few scattered teeth. The lower basal leaves of the rosette usually die before flowering occurs.

Mature

Pale yellow flowers with four petals are produced in short racemes at the ends of branches. The pods are very long, thin and smooth with no obvious beak at the tip. The small seeds are oblong, varying in color from yellow to reddish-yellow to olive-green. The mature stem becomes brittle and breaks at the base. The plant tumbles along the ground to distribute seed.

Key features

Lower leaves divided and upper leaves finely divided; pale yellow flowers; narrow pod that is much longer than other members of the *Brassicaceae*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Mature

Juvenile: rosette

Flower: pale yellow

Wild Mustard

Brassica kaber

Family

Mustard *Brassicaceae*

Seedling

The broad, kidney-shaped cotyledons are similar to both Polish and Argentine canola. The first true leaves are petioled, slightly lobed and hairy.

Juvenile

The stem is branched with bristly downward-pointing hair near the base and becomes less hairy or hairless near the top. Usually, the bases of Polish canola stems are less hairy. In the rosette stage, the stalked lower leaves of the plant have several deep lobes near the base of the blade and a large terminal lobe. There are sparse hairs, especially on the veins on the undersurface of the leaf. Upper leaves are sessile and coarsely toothed, but not clasping. Stem nodes and the bases of branches develop purple blotches, a feature that is useful to distinguish wild mustard from Polish canola, which does not develop this coloration.

Mature

The flowering racemes have bright yellow flowers with four petals. Each flower develops into a cylindrical, ribbed pod with a flattened beak that is about one third the length of the pod itself. There is often one seed at the base of the beak. The stalk at the base of the pod is very short and almost as thick as the pod. Seeds are spherical, brown to black and minutely netted under magnification.

Key features

Downward-pointing bristly hairs on lower stem; upper stem leaves coarsely toothed and sessile, but not clasping; purple blotches at stem nodes

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Juvenile: *rosette*

Flower: *flowering raceme*

Mature

Pods with flattened beaks

Wild Radish

Raphanus raphanistrum

Family

Mustard *Brassicaceae*

Seedling

Cotyledons are broad and kidney-shaped. The first true leaves usually have two small lobes at the base and slightly wavy margins. Wild radish is very difficult to distinguish from wild mustard at the seedling stage.

Juvenile

The basal rosette leaves are deeply-divided with a prominent, terminal lobe. They are frequently covered with a few, short, stiff hairs and are rough to the touch. Stems are erect and often freely-branched. They are covered with coarse, short hair, especially at the base. The upper stem leaves are narrower than the rosette leaves and usually have entire margins.

Mature

Flowers are produced in racemes at the ends of stems and branches. Most flowers have four pale yellow petals, although they may occasionally be white or even purple. Typically, the petals are conspicuously veined. Seedpods are long and narrow with a slender, pointed beak at the tip. They have distinctive constrictions between the seeds and break apart into segments at maturity. Each segment contains one oval, reddish-brown seed covered with fine net veins. Mature wild radish plants are easily confused with wild mustard. However, wild mustard flowers are smaller, a deeper yellow color and they lack the conspicuous veins on the petals. The seedpods of wild mustard also do not exhibit constrictions between the seeds.

Key features

Deeply-divided basal leaves; conspicuously-veined petals; constricted seedpods

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas	

Juvenile: *rosette*

Flower: *pale yellow in a raceme*

Mature

Infestation

Wood Whitlow-grass

Draba nemorosa

Other names

Yellow whitlow-grass

Family

Mustard *Brassicaceae*

Seedling

The seedling is very small and can easily be overlooked. It has ovate, sessile leaves.

Juvenile

The basal rosette has ovate leaves with two, three and four-pronged hairs. The leaf margins are finely toothed. Its tiny rosettes are similar in size to pygmyflower, and both species are often found in the same habitat. Wood whitlow-grass can be distinguished from pygmyflower by the branching hairs on the leaves. Pygmyflower leaves have minute, simple hairs or are hairless. Stems of wood whitlow-grass have simple and forked hairs.

Mature

This plant is seen very early in stubble fields, usually after it has bolted and flowered. Flower racemes have widely spaced flowers on long stalks. Each tiny yellow flower has four petals that are slightly notched at the tip. Each flower develops into a flattened, elliptical pod that points upward. The stalk on each pod is as long or longer than the pod itself. Flowering and seed set occur very early in spring.

Key features

Tiny rosette and short stature of mature plant; early bloom and seed set; stalk of each upward-pointing pod as long as or longer than the pod itself

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides		
forage crops, hay fields	turf	waste areas		

Seedling

Juvenile: rosette

Infestation

Mature

Flower: flowering raceme

Yellow Nut Sedge

Cyperus esculentus

Family

Sedge *Cyperaceae*

Seedling

Seedlings are seldom found in the field.

Juvenile

Yellow nut sedge is a creeping perennial weed with fibrous roots and long, light brown to white wiry rhizomes. The rhizomes may or may not end in dark brown, nut-like tubers. Both the tubers, when present, and rhizomes allow this weed to spread vigorously underground. Tall, unbranched, smooth stems are produced that are distinctly triangular in cross-section. The long, narrow leaves appear grass-like and are flat or somewhat folded. The leaves emerge from all three sides of the stem and occur mostly towards the base of the plant. Yellow nut sedge is considered one of the worst weeds in the world and has recently started to appear in moist areas in the eastern prairies.

Mature

Small, yellowish-brown flowers are grouped into flattened spikelets located in umbrella-like clusters at the tops of stems. There are usually three to four long, thin leaves located below each inflorescence. The seeds of yellow nut sedge are long, elliptical in shape and conspicuously three-sided.

Key features

Long, wiry rhizomes, often with brown tubers attached; triangular stems with narrow, grass-like leaves at the base; yellowish-brown flowers in umbrella-like clusters

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Juvenile

Flower: *yellowish-brown in clusters*

Mature

Infestation

Yellow Sweet Clover

Melilotus officinalis

Family

Pea *Fabaceae*

Seedling

Cotyledons are oblong with a round tip. The first true leaf has a single, broadly oval blade and a long, grooved stalk. Subsequent leaves are all trifoliate.

Juvenile

This biennial weed produces a large, deep tap root and basal rosette in the first year. The compound leaves are divided into three, narrowly oval leaflets. Only the terminal leaflet is stalked. The margins of all leaflets are toothed, almost to the base. Erect, often very tall, free-branching stems are formed in the second year. They typically have a smooth but strongly ridged surface. It is difficult to distinguish yellow sweet clover from its close relative, white sweet clover, until flowers and fruits are produced. Although both species of sweet clover are weedy in certain areas, they are also commonly grown as forage crops.

Mature

Yellow sweet clover has irregular, pea-like flowers in narrow, elongated racemes. Each yellow flower produces a small, egg-shaped pod with a wrinkled surface. The pods contain one or two olive green seeds that are also frequently purple-spotted. As the name suggests, white sweet clover forms white flowers that are smaller but similar in appearance to yellow sweet clover. Its pods have a net-veined surface and contain yellow colored seeds.

Key features

Trifoliate leaves with leaflet margins toothed almost to the base; yellow pea-like flowers in elongated racemes; small, egg-shaped pods with a wrinkled surface

Similar species

White sweet clover *Melilotus alba*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Mature

Similar species: *Yellow and white sweet clover together*

Flower: *elongated flower racemes*

Infestation

Yellow Toadflax

Linaria vulgaris

Other names

Butter-and-eggs, wild snapdragon

Family

Snapdragon *Scrophulariaceae*

Seedling

Yellow toadflax cotyledons are round with a very prominent rounded bump at the top. The first few true leaves are oval. Subsequent leaves are narrower and linear. This species is becoming more prevalent in reduced tillage systems.

Juvenile

Yellow toadflax spreads by extensive creeping roots that form dense patches. The stems are erect, hairless and branched above. The leaves are alternate, although often crowded on the stem. The linear leaves are narrowed at both ends. This feature helps to distinguish it from Dalmatian toadflax, which has lanceolate to broadly ovate leaves that clasp the stem. Before flowering, yellow toadflax can be distinguished from leafy spurge by breaking the stem.

Leafy spurge has a white milky latex inside while yellow toadflax does not. Yellow toadflax contains some toxic compounds but is usually avoided by grazing livestock.

Mature

Bright yellow flowers with orange centres are grouped into racemes at the ends of stems and branches. Each flower is “snapdragon-like” with an upper lip and a lower lip that has a conspicuous protrusion or “spur” extending downwards. Each flower develops into a seed capsule that releases many dark brown to black seeds. Each seed is flattened with a circular wing on the margin for wind distribution.

Key features

Found in patches due to creeping roots; crowded, linear leaves that are narrower at each end; bright yellow and orange flowers that are “snapdragon-like”

Similar species

Dalmatian toadflax *Linaria dalmatica*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Flower

Infestation

Mature

Similar species: *Dalmatian roadflax*

Green

Flowered Species

Barley

Hordeum vulgare

Family

Grass *Poaceae*

Seedling

Leaf blades and sheaths are hairless, although occasional hairs occur on the upper blade of some varieties. Barley has large white, conspicuous auricles that make it readily identifiable in the field. The ligule is long, white and membranous. Leaf blades are wide in comparison to wheat.

Juvenile

Barley occurs as a volunteer weed and can be a vigorous competitor. It has a fibrous root system, erect stems and often produces many tillers.

Mature

The inflorescence is a spike with either two or six rows of kernels. Two-row spikes are much thinner and flatter in appearance than six-row spikes. In both two-row and six-row varieties, there are three spikelets at each rachis node. Only the central spikelet in two-row types has a floret that develops into a grain or seed. The lateral spikelets are sterile. All three spikelets in six-row varieties have one floret that develops into one grain. The two tiny white glumes on the outside of each spikelet are awned, and there is a long awn or beard extending from the tip of each grain. The beard is either barbed or smooth and is broken off during threshing. Barley grains are straw-colored and can be hulled or hullless.

Key features

Large, white, claw-like auricles; awned glumes; beard extending from tip of grain either barbed or smooth

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas		

Seedling

Key feature: *large, claw-like auricles*

Mature: *panicle*

Two-row barley spike

Barnyard Grass

Echinochloa crusgalli

Family

Grass Poaceae

Seedling

The leaves of barnyard grass are completely hairless although the margins of the blade may be slightly rough. Auricles are absent. It is the only grass weed on the prairies that has no ligule, making this an excellent identification feature.

Juvenile

Barnyard grass has fibrous roots and a spreading, horizontal growth habit. Stem nodes in contact with the soil often form roots. The sheath of each leaf is rather flattened and may be purplish near the base.

Mature

The inflorescence is a tufted, compact panicle that is often purplish. The branches are covered with densely-crowded spikelets that are covered with stiff hairs. Only one of the two florets will develop into a seed. The sterile second floret often has a bristly awn extending from the tip. The seed is shiny, pale yellow to brown, flattened on one side and rounded on the other.

Key features

Auricles and ligule absent on hairless leaves; horizontal growth habit; bristly-looking panicles densely crowded with spikelets

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Mature

Infestation

Panicle

Key feature: *ligule absent*

Broad-leaved Plantain

Plantago major

Family

Plantain *Plantaginaceae*

Seedling

Cotyledons are oval. The true leaves are oval and petioled. They emerge one at a time to form a basal rosette.

Juvenile

The root is short and thick with many tough, lateral roots.

Broad-leaved plantain lacks an above-ground stem, and the leaves remain in a basal rosette throughout the life cycle. The leaf margins are entire or occasionally toothed. The oval to ovate leaves have thick petioles that are as long as the leaf blades. The leaves are prominently ribbed with three to seven main veins and may be smooth or slightly hairy. This weed is commonly found in lawns.

Mature

Flowers are arranged in narrow, cylindrical spikes at the top of leafless flower stalks. Depending on plant density, there may be several to several dozen flower stalks per plant. Broad-leaved plantain flowers and sets seed from spring to late autumn, and its pollen can cause hay fever. A small individual green flower develops into an egg-shaped capsule that splits to release numerous seeds. The seeds are dark brown to black and have various, angular shapes. There are threadlike ridges on the seed surface and a pale scar on one side.

Key features

Remains a basal rosette throughout life cycle; oval, prominently ribbed leaves with thick petioles as long as the blades; leafless flower stalks with green-flowered spikes

Life Cycle					
annual	winter annual	biennial	creeping perennial	simple perennial	
Habitat					
conventional tillage		rangeland		wetland	
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Mature

Flowering spike

Cocklebur

Xanthium strumarium

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are long and narrow, persisting at the base of the stem, sometimes until maturity. First true leaves are opposite, petioled and ovate with toothed margins. The remains of the bur below ground can be used to confirm seedling identification. Seeds and young seedlings are poisonous to livestock.

Juvenile

Cocklebur is tap rooted with erect, rough-hairy stems that are ridged and have purplish spots. Leaves have three main veins and are ovate to heart-shaped or triangular. Lower leaves are opposite; upper leaves are alternately arranged. Leaves and stems are rough to the touch due to glandular hairs. These hairs can cause a rash in allergic individuals.

Mature

Male and female flower heads are separate, but are on the same plant. They are located at the ends of the main stem and branches as well as in leaf axils. Male flower heads are spherical, non-spiny and produce pollen that causes an allergic reaction in some people. The female flower head is a green, spiny bur containing two flowers. The bur is covered with spines and has two beak-like spines at the tip. It becomes brown, hard and woody at maturity. The two seeds inside each bur are oblong, flattened and grey to dark brown, with one usually larger than the other.

Key features

Rough-hairy leaves and stems; ovate to heart-shaped or triangular leaves with three main veins; large, spiny burs

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides		
forage crops, hay fields	turf	waste areas		

Seedling

Juvenile

Mature

Key feature: *spiny burs*

Key feature: *rough-hairy stems*

Downy Brome

Bromus tectorum

Other names

Downy chess, cheat grass

Family

Grass *Poaceae*

Seedling

Leaf blades and sheaths of downy brome are pale green and covered with soft, dense hairs on both surfaces and margins. Venation is indistinct except for the midrib on the lower surface. Auricles are absent. The ligule is membranous, approximately 3 mm long and torn along the top.

Juvenile

The plant tillers to produce a tufted clump. The erect stems are slender and usually hairy, and the leaf sheaths are closed at the bottom and split at the top. The root system is fibrous, growing to a depth of about 30 cm. In some areas, downy brome is considered to be forage. Smooth brome, the related forage grass, is a perennial with branching rhizomes and smooth leaves with rough margins. The leaf texture and presence of rhizomes can be used to distinguish smooth brome from downy brome.

Mature

The panicle has long, drooping branches. It is pale green and becomes tinged with purple as it matures. Spikelets have hairy, uneven-sized outer glumes and are multi-floreted, each floret developing into one seed. The seeds are also softly hairy with a short, slender, rough awn emerging from just below the tip. Immature heads feel soft to the touch. Smooth brome panicles are not awned and hence, are easily distinguished from downy brome.

Key features

Soft, densely-hairy leaves; drooping, purple panicles; short, slender awns on seeds

Similar species

Smooth brome *Bromus inermis*

Life Cycle					
annual	winter annual	biennial	creeping perennial	simple perennial	
Habitat					
conventional tillage	rangeland		wetland		
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Juvenile

Key feature: *soft, dense hairs on leaves*

Mature: *panicle*

Similar species: *Smooth brome – panicle*

False Ragweed

Iva xanthifolia

Other names

Burweed, marsh elder

Family

Sunflower *Asteraceae*

Seedling

Cotyledons are oval and stalked. The first true leaves are opposite with variably lobed margins and a soft, hairy texture.

Juvenile

Stems are erect and much-branched. They are usually smooth, but occasionally hairy at the top. Leaves are mostly opposite with long, hairy petioles. The lower leaves are often three-lobed while the upper are broad, almost heart-shaped, with coarse, irregularly-toothed margins. Both surfaces of the leaves are covered with flattened hairs, giving them a velvety feel. The long petioles, toothed margins and ovate upper leaves may cause confusion with cocklebur at this stage. However, false ragweed has opposite, velvety leaves while the leaves on cocklebur are alternate and rough-textured.

Mature

The heads of false ragweed are small, greenish and located in loose, branching clusters at the top of the plant. Each head has male and female flowers surrounded by five pointed involucre bracts. The central flowers on the head are male and produce pollen that may cause hay fever in some people. The marginal flowers are female and each can produce one seed. Usually five seeds are found in a ring around the male flowers. The seeds are brown to black, very finely grooved and lack a pappus.

Key features

Velvety, opposite leaves; irregularly-toothed leaf margins; loose clusters of small green flower heads

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Mature

Juvenile: *showing velvety, opposite leaves*

Flower: *clusters of flower heads*

Field Dock

Rumex pseudonatronatus

Family

Buckwheat *Polygonaceae*

Seedling

Cotyledons are elliptical in shape. The first true leaf is petioled and broadly oval with entire margins.

Juvenile

Field dock is more abundant on the prairies than curled dock (*Rumex crispus*), which is more abundant in Eastern Canada. Western dock (*Rumex occidentalis*) is a native dock that is non-weedy but very similar to field dock. Western dock and field dock are extremely difficult to distinguish.

Field dock has a deep tap root. In the spring, new buds grow from the previous year's rosette. Stems are erect with ocreas at the nodes. Leaf arrangement is alternate. The narrow, lanceolate leaves have wavy-curved margins and taper to the base. Leaves of Western dock have round basal lobes and taper to the tip. Stems of both dock species are commonly red-tinged as maturity approaches.

Mature

Both field dock and Western dock have long, narrow, densely-flowered panicles. The stalk of each flower of field dock is distinctly jointed. Flower stalks of Western dock are unjointed. Each flower lacks petals but has six sepals in two whorls of three. The inner sepals are enlarged and form the winged valves of the fruit. These valves are heart-shaped in both species. The flower stalks of field dock are two to three times longer than the valves. Western dock flower stalks are about the same length as the valves. Projections called "tubercles" are absent on the valves of both species, although there may be a slight swelling on one valve of field dock. Seeds are brown and three-angled.

Key features

Ocreas at stem nodes; stalk of each flower distinctly jointed; heart-shaped valves of fruit that may have a swelling on one valve

Similar species

Western dock *Rumex occidentalis*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Mature

Juvenile

Key feature: ocrea at stem node

Dense panicle

Field Horsetail

Equisetum arvense

Other names

Scouring rush, pinegrass, mare's tail

Family

Horsetail *Equisetaceae*

Seedling

Field horsetail does not form seedlings because it is not a flowering, seed-producing plant. Instead of seedlings, stems are produced from underground rhizomes. These rhizomes often have attached tubers that may also break off to start a new plant.

Juvenile

A unique feature of field horsetail is that it has two distinctly different types of stems. Both types pull apart easily at the nodes. In early spring, the first stems to emerge are reproductive and are described more fully below. The secondary, vegetative stems are what most people recognize as field horsetail. They are green, hollow and more slender than the reproductive stems. Each node on the secondary stems has whorled branches and is surrounded by a small, toothed sheath. At first glance, field horsetail can be confused with corn spurry, another weed with a whorled arrangement. However, with closer inspection, it is easy to see that corn spurry has leaves, not branches, and that its stems are not easily separable at the nodes. It is also an annual that lacks rhizomes. Field horsetail is poisonous, particularly to young horses. They usually do not choose to eat it in the field, but may not be able to avoid it in hay.

Mature

The reproductive stems that emerge early in spring are white-brown and unbranched. They are hollow with each node surrounded by a sheath with black teeth. The stems are topped with a cone-like structure containing spores. Once the spores are released, these stems wither away, soon to be replaced by the secondary stems.

Key features

Green vegetative stems with whorled branches; brown reproductive stems topped by a spore-producing cone; both stems hollow with toothed sheaths and easily separable at the nodes

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Juvenile: *vegetative stem*

Key feature: *whorled branches*

Key feature: *spore-producing cones*

Infestation

Foxtail Barley

Hordeum jubatum

Other names

Wild barley

Family

Grass *Poaceae*

Seedling

The first few greyish-green leaves appear as thin, vertical wisps. They are covered with short, dense hairs. Venation is prominent and margins are rough. Auricles are absent or rudimentary. The membranous ligule is very short, about 1 mm, with fine hairs.

Juvenile

Foxtail barley is a fibrous-rooted, densely tufted perennial that spreads only by seed. The stems are erect and smooth while the leaf sheaths are split and hairy. Foxtail barley tolerates wet and moderately saline habitats.

Mature

The inflorescence is a dense, long-awned spike that may be greenish or purplish and nod to one side. It has a jointed rachis that breaks into sharply pointed segments at maturity. Each segment is made up of three spikelets, but only the central spikelet has one creamy-colored seed. In addition, each segment has seven awns with upward-pointing barbs that easily become attached to animals. If grazed, these barbed segments may become embedded in the animal's mouth and face, causing severe irritation, abscesses and even blindness.

Key features

Greyish-green, slender leaves; tufted growth; barley-like, densely-awned heads

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Juvenile

Mature

Spike

Key feature: *awned seeds*

Green Foxtail

Setaria viridis

Other names

Wild millet, millet

Family

Grass *Poaceae*

Seedling

The first tiny leaf of green foxtail emerges parallel to the ground. The leaf shape is linear to lanceolate with a pointy tip. The light green leaves are smooth and finely veined, with a distinct lower midrib. Margins and upper surface of leaf blades may feel slightly rough to the touch. The ligule is a distinct fringe of hair which is an excellent identification feature. Auricles are lacking and the margins of the split sheath are trimmed with hair.

Juvenile

Green foxtail has a fibrous root system and often occurs in dense populations. The leaf characteristics are especially useful to differentiate green from yellow foxtail. Yellow foxtail lacks hairy sheath margins. Also, yellow foxtail has long, kinky hairs on the upper surface of the leaves near the base of the blade. These hairs are completely lacking in green foxtail.

Mature

The soft, bristly spike-like panicle is green or purple-tinged and resembles a bottle brush. It has densely packed spikelets with groups of bristles attached below the spikelets. Each spikelet will produce only one seed, which is flattened on one side. The shiny seed is cream-colored to brown to black, mottled and has a hard, wrinkly surface. The seeds of green foxtail are smaller than those of yellow foxtail.

Key features

Ligule a fringe of hair; sheath margins trimmed with hair; bristly spike-like panicle that resembles a bottle brush

Similar species

Yellow foxtail *Setaria glauca*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Juvenile

Mature

Key feature: *bristly, spike-like panicle*

Key feature: *ligule is a fringe of hair*

Kochia

Kochia scoparia

Family

Goosefoot *Chenopodiaceae*

Seedling

Cotyledons are short and narrow. They are often, but not always, bright pink to magenta underneath. True leaves are sessile and linear with a pointed tip. They have entire margins and soft, long hairs on the surface. Many young leaves emerge almost simultaneously to form a rosette. When the stem elongates soon after, the leaves are alternately arranged on the stem.

Juvenile

Kochia has an erect, much-branched stem that is often purple-striped. The leaves of kochia are densely hairy and have a greyish-green appearance. Main stem leaves are larger than those on the lateral branches, which in turn have leaves that become smaller towards the branch tips.

Mature

Flowers of kochia lack petals, but have five winged sepals. They are green and inconspicuous in the leaf axils, often surrounded by clusters of long hairs. Each flower develops into a flattened, dull brown, egg-shaped seed with a broad groove on each side. Occasionally, the whole plant may develop bright red to magenta coloration in the fall to give a "burning bush" appearance.

Key features

Rosette appearance of the young plant; greyish-green leaves covered with dense, soft, long hairs; purple-striped stem

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides		
forage crops, hay fields	turf	waste areas		

Seedling

Juvenile

Key feature: *purple-striped stem*

Mature

Flower: *clusters in leaf axils*

Lamb's-quarters

Chenopodium album

Family

Goosefoot *Chenopodiaceae*

Seedling

Cotyledons are narrow, linear and often pink or purplish underneath. The first two true leaves appear opposite, but subsequent leaves are alternate. The greyish-green leaves are petioled and have mealy, white particles mainly on the lower leaf surfaces. Leaves are ovate to triangular with irregularly lobed or toothed margins.

Juvenile

The erect stem is branched, usually red or purple striped and ridged lengthwise. The leaves are variable in shape, and margins vary from toothed to lobed to entire. Net-seeded lamb's-quarters, *Chenopodium berlandieri*, is also very common in the prairie provinces. Although its leaves tend to be thicker, it is difficult to distinguish from lamb's-quarters until maturity.

Mature

Flowers are green and inconspicuous with no petals and five sepals. They are found in dense clusters at the tips of branches and crowded into leaf axils. Each flower develops into a shiny, black seed that is circular, flattened and covered with a plain, papery envelope. The envelope is etched with reticulate or netted markings in net-seeded lamb's-quarters. Also, the sepals of net-seeded lamb's-quarters are prominently keeled and do not completely cover the fruit at maturity. The sepals of lamb's-quarters are unkeeled and completely cover the mature fruit.

Key features

White, mealy particles on leaf undersurfaces; red or purple stripes on stems; dense clusters of tiny, greyish-green flowers

Similar species

Net-seeded lamb's-quarters *Chenopodium berlandieri*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Juvenile

Key feature: *striped stem*

Mature

Flower: *dense flower clusters*

Persian Darnel

Lolium persicum

Family

Grass *Poaceae*

Seedling

The young plant has very narrow leaf blades that are shiny or glossy. Auricles are absent on the first few seedling leaves. The ligule is membranous, short and ragged. Leaf blade margins and surfaces may be slightly rough to the touch.

Juvenile

Persian darnel is an erect, tufted annual with fibrous roots. The leaf sheath is split, smooth-textured and has prominent veins. The leaves usually have small, claw-like auricles on the middle and upper leaves of the plant.

Mature

The spike is composed of spikelets that are arranged edgewise to the rachis, making the inflorescence appear flat. There is only one outer glume on each spikelet except for the terminal spikelet, which has two glumes. Each spikelet is multi-floreted, and each floret produces a thin, light brown seed with a slender, wispy awn at the tip.

Key features

Shiny leaf blade; spikelets edgewise to the rachis; one glume per spikelet; terminal spikelet two-glumed

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Key feature: *spikelets edgewise to rachis*

Juvenile: *leaf with small auricles and short ligule*

Mature: *showing spikes*

Proso Millet

Panicum miliaceum

Other names

Broomcorn millet

Family

Grass *Poaceae*

Seedling

Proso millet seedlings resemble volunteer corn except that the leaves are narrower and covered with dense, stiff hairs. No auricles are present. The ligule is membranous at the base and fringed with hair along the top.

Juvenile

Proso millet is a plant that has escaped from cultivation. The plant is tufted with a fibrous root system. Stems are erect or partially reclining with erect tips. Witch grass is a related species, also densely hairy with the hairs bristling out at 90 degrees from the leaf blades and sheaths. The sheaths of witch grass leaves are often purplish, and the plant is much shorter in height than proso millet.

Mature

Proso millet is more easily distinguished from witch grass once it has headed out. The panicle of proso millet is compact and usually erect to drooping. Witch grass is an airy, delicate, loose panicle that may be up to half the height of the adult plant. Stems of witch grass break off below the head, and the panicle tumbles along the ground dropping seed. Proso millet panicles do not break off and are much coarser in appearance. The spikelets of proso millet are about twice the size of witch grass ones, although both produce only a single seed. Proso millet seeds are smooth and shiny, and they vary in color from yellow to olive to black.

Key features

Dense hairiness of leaves; erect to partially reclining growth habit; erect to drooping panicle with single-seeded spikelets

Similar species

Witch grass *Panicum capillare*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides		
forage crops, hay fields	turf	waste areas		

Seedling

Juvenile

Mature

Panicle

Key feature: *hairy leaves*

Prostrate Pigweed

Amaranthus blitoides

Family

Amaranthaceae

Seedling

Prostrate pigweed has narrow, linear cotyledons. The true leaves are alternate, petioled and are wider at the tip than at the base. There is a small indent at the leaf tip with a soft spine in the centre. The leaves have prominent white veins underneath. Prostrate pigweed cotyledons are longer in relation to the length of the first true leaves than are cotyledons of redroot pigweed at the same growth stage.

Juvenile

Prostrate pigweed forms prostrate, branching mats on the soil surface. Stems are fleshy and often reddish. Tumble pigweed, a related species, has similar leaves and prominent white venation but is erect with whitish-green stems. Other flat-growing species can be quite easily distinguished from prostrate pigweed. Purslane has fleshy leaves, lacking prominent venation. Prostrate knotweed has papery ocreas at the stem nodes, and thyme-leaved spurge has white latex in the stems.

Mature

Green, inconspicuous flowers are clustered in leaf axils. The flowers lack petals but have sepals and extra bracts around each flower. Although tumble pigweed also has green flowers in the leaf axils, the plant is obviously erect, so it is unlikely to be confused with prostrate pigweed. Tumble pigweed also has spiny-tipped bracts under each flower that are twice as long as the sepals. The sepals and extra bracts under each flower of prostrate pigweed are the same length.

Key features

Prostrate, branching mats on soil surface; leaves with soft spine at the tip and prominent, white veins underneath; inconspicuous, green flowers in leaf axils

Similar species

Tumble pigweed *Amaranthus albus*

% of fields where found

Life Cycle

annual

winter annual

biennial

creeping perennial

simple perennial

Habitat

conventional tillage

rangeland

wetland

reduced tillage

irrigation

roadsides

forage crops, hay fields

turf

waste areas

Seedling

Juvenile

Mature: *prostrate plant*

Flower: *clusters in leaf axils*

Similar species: *Tumble pigweed – mature*

Quack Grass

Elytrigia repens

Other names

Couch grass, twitch grass

Family

Grass *Poaceae*

Seedling

The first true leaves have hairy sheaths. Auricles are small and claw-like. The ligule is membranous but very short. The leaf margins and upper surface of the leaf blade are rough, and there may be sparse hair on the upper surface of the blade.

Juvenile

Quack grass is more frequently seen growing from a rhizome than as a young seedling. The rhizomes allow swift, vegetative propagation in the field, provide carbohydrate storage during the winter and make quack grass a difficult weed to control. The tough, whitish rhizomes are sharply pointed at the growing tip. Buds occur at the nodes, and a brown bract covers part of each internode. Buds grow into rhizome branches or above-ground shoots. There is often a noticeable M-shaped constriction near the leaf tips. Later leaves tend to be less hairy than earlier ones.

Mature

The broad surface of quack grass spikelets is against the rachis, making this spike easily distinguishable from Persian darnel, which has spikelets edgewise to the rachis. Each spikelet is multi-floreted with two glumes that may have short awns. The florets may also have a short awn on the tip. The seed is narrow and straw-colored at maturity.

Key features

Small, claw-like auricles; hairy sheaths on lower leaves; white rhizomes with brown bracts

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial

Habitat		
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides
forage crops, hay fields	turf	waste areas

Seedling

Juvenile

Key feature: *rhizome with bract and roots*

Spike

Key feature: *small, claw-like auricles*

Redroot Pigweed

Amaranthus retroflexus

Family

Amaranth *Amaranthaceae*

Seedling

Cotyledons are long and narrow, with short stalks. The first true leaves are ovate and shallowly notched at the tip. Close examination will reveal that the leaf mid-vein extends as a small bristle into the middle of the notch. Both the cotyledons and early true leaves are frequently reddish on the undersides. The young developing stem and tap root are also usually red.

Juvenile

Redroot pigweed has erect, simple-to-branched stems. The stems are usually hairy and green near the top of the plant and more smooth and reddish at the base. The tap root is often still pinkish-red at this stage. Leaves are alternate, long-petioled and somewhat diamond-shaped. The leaf margins vary from entire to slightly wavy, and it may sometimes be difficult to recognize the notch at the tip. The undersides of the leaves are sparsely hairy with prominent, white veins. The protruding veins and notched tip on the leaves help to distinguish redroot pigweed from lamb's-quarters, a weed often mistaken for it.

Mature

Tiny, green flowers are located in coarse, bristly spikes clustered at the top of the plant. A short, central spike is usually surrounded by several smaller, finger-like spikes pointing outwards. Small spikes are also frequently found in the axils of leaves. Each flower lacks petals but has three, spiny-tipped bracts below. The bracts are what give the inflorescence such a bristly, spiny appearance and feel. Each flower produces a tiny black, very shiny seed that is almost round with a narrow, flattened margin.

Key features

Red tap root and stem base; ovate to diamond-shaped leaves with a notched tip and prominent veins below; numerous, bristly spikes

Life Cycle					
annual	winter annual	biennial	creeping perennial	simple perennial	
Habitat					
conventional tillage	rangeland		wetland		
reduced tillage	irrigation		roadsides		
forage crops, hay fields	turf		waste areas		

Seedling

Juvenile

Infestation

Flower: *bristly spikes*

Key feature: *red stem and root on seedling*

Russian Thistle

Salsola pestifer

Family

Goosefoot *Chenopodiaceae*

Seedling

Cotyledons are slender, needle-like and succulent. The first two true leaves appear opposite, but subsequent leaves are alternate.

Russian thistle leaves are succulent with soft spines on the tips. This feature distinguishes the plant from corn spurry, which has needle-like leaves lacking the spiny tip.

Corn spurry leaves are also whorled. Seedlings of knawel have non-succulent, whorled or opposite leaves with hairs on the stems and leaves. Russian thistle leaves and stems lack hairs.

Juvenile

Russian thistle is erect and much-branched. It develops a bushy, spherical appearance as it matures. The stems are often red or purple striped. Young leaves are longer and more succulent than later leaves, which are wedge-shaped and tipped with a strong, sharp spine.

Mature

The small, inconspicuous green flowers are located in leaf axils. The flowers have no petals, but there are five papery sepals and two spiny-tipped bracts under each flower. The mature, round, bushy plant breaks off at ground level and rolls with the wind to drop its seed. The sepals remain attached around each seed as it falls off the plant. The dull grey seeds have a coiled embryo, which can be seen through an almost transparent seed coat.

Key features

Young leaves succulent, needle-like with soft spiny tips; mature leaves short, wedge-shaped with strong, sharp spines; red or purple striped stems

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Juvenile

Trampling plants

Mature

Flower: *flowers in leaf axils*

Spear-leaved Goosefoot

Monolepis nuttalliana

Family

Goosefoot *Chenopodiaceae*

Seedling

Cotyledons are linear with short stalks. The first true leaves usually have long petioles and are narrowly oval with two small “bumps” near the base. These bumps soon develop into triangular lobes, giving the leaves the appearance of a spearhead.

Juvenile

Spear-leaved goosefoot typically forms a basal rosette early in the spring. The spear-shaped leaves are thick and almost fleshy. Frequently, there is a light, powdery coating on the leaf undersides. The rosettes form low, spreading stems and branches that are smooth, succulent and often reddish. Spear-leaved goosefoot is similar to oak-leaved goosefoot, another low, spreading weed in this family. These two species can be easily distinguished by the leaves which are regularly and shallowly toothed around the margins in oak-leaved goosefoot. Although originally most common on saline soils, spear-leaved goosefoot is now widely adapted to gardens and cultivated fields across the prairies.

Mature

Small, inconspicuous flowers are located in axillary clusters. The flowers lack petals and instead have only one green, bract-like sepal. Each flower produces one tiny, flattened seed.

Key features

Spear-shaped leaves; fleshy texture of stems and leaves; low, spreading growth habit

Similar species

Oak-leaved goosefoot *Chenopodium glaucum*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland
reduced tillage	irrigation	roadsides		
forage crops, hay fields	turf	waste areas

Seedling

Juvenile

Similar species: *Oak-leaved goosefoot*

Mature

Flower: clusters in leaf axils

Stinging Nettle

Urtica dioica

Other names

American stinging nettle

Family

Nettle *Urticaceae*

Seedling

The cotyledons are oval with a notched tip. First true leaves are opposite and hairy with round-toothed margins. On the prairies, stinging nettle is usually found in moist, shady woodlands where seedlings are unlikely to be noticed. It can also be found in a variety of habitats including horticultural crops.

Juvenile

Stinging nettle has vigorous, spreading rhizomes and tall, erect, square-angled stems. Stems are equipped with specialized projections called “stinging hairs” that produce localized swelling, itchiness and numbness when touched. The leaves are opposite, ovate to lanceolate and toothed on the margins. The leaf petioles are long on the lower part of the plant, becoming shorter towards the top. Stinging hairs are also found on the lower leaf surfaces. There are stipules at the stem nodes. Hemp-nettle is sometimes confused with stinging nettle, but it lacks both stinging hairs and stipules.

Mature

The small, green flowers are in separate male and female clusters produced in the leaf axils. The spike-like chains of male flowers tend to be longer than the female ones. Individual flowers of hemp-nettle are much larger with a pink to purple corolla. The pollen of stinging nettle is one cause of summer hay fever. Seeds are tan to brown and egg-shaped.

Key features

Rhizomes; tall, square-angled stems with opposite leaves; flowers in spike-like chains in leaf axils; stinging hairs

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage		rangeland		wetland
reduced tillage		irrigation		roadsides
forage crops, hay fields		turf		waste areas

Seedling

Juvenile

Flower: *clusters in leaf axils*

Infestation: *mature*

Key feature: *stinging hairs*

Tartary Buckwheat

Fagopyrum tataricum

Family

Buckwheat *Polygonaceae*

Seedling

Cotyledons are almost round with rounded basal lobes and long stalks. Seedlings with attached cotyledons are easy to distinguish from wild buckwheat seedlings, which have linear cotyledons. The true leaves of tartary buckwheat are broadly triangular with wide-spreading basal lobes.

Juvenile

Tartary buckwheat is tap-rooted with erect green stems and ocreas at the stem nodes. Wild buckwheat is readily distinguished from tartary buckwheat by its weak, twining stems. Leaves of tartary buckwheat are alternate with long petioles on lower leaves and short petioles on upper leaves. The triangular leaves are often wider than they are long.

Mature

Flowers are green and inconspicuous in leaf axils and at the tips of stems. There are no petals, and the five sepals are unequal in length. The one seed that develops from each flower protrudes from the sepals. The seed is brown or dark grey, three-sided and rough on the edges. Each seed of wild buckwheat is securely held by the sepals and is smaller than that of tartary buckwheat.

Key features

Broad, triangular leaves; erect stems with ocreas at the stem nodes; seed protruding from the sepals

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides	
forage crops, hay fields	turf		waste areas	

Seedling

Juvenile

Flower

Mature

Thyme-leaved Spurge

Euphorbia serpyllifolia

Family

Spurge *Euphorbiaceae*

Seedling

Cotyledons of thyme-leaved spurge are ovate. The true leaves are also ovate, oppositely arranged and have finely toothed margins.

Juvenile

Thyme-leaved spurge forms prostrate mats on the soil surface. All parts of the plant are filled with a white, milky juice. This feature can be used to distinguish it from other mat-forming species such as prostrate knotweed, prostrate pigweed and purslane, which do not contain white latex. In addition, the leaves are opposite and finely toothed in contrast to the others, which have alternate leaves and entire leaf margins.

Mature

The flowers are small and resemble those of leafy spurge. The separate female flower forms a three-celled capsule with one seed in each compartment. The seeds are pitted and faintly ridged.

Key features

Prostrate mats; milky juice in stems and leaves; opposite, finely toothed leaves

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides	
forage crops, hay fields	turf		waste areas

Seedling

Juvenile

Mature: *prostrate plant*

Wheat

Triticum aestivum

Other names

Hard red spring wheat, bread wheat

Family

Grass *Poaceae*

Seedling

The leaf blade and sheath are hairless to the naked eye. The leaves are narrower than those of barley. The ligule is white, membranous and shorter than barley. Most characteristic are the small, claw-like auricles trimmed with hair. Durum wheat auricles are hairless.

Juvenile

Wheat is a tufted plant with fibrous roots. Volunteer wheat can also be a weed and will compete with the new crop.

Mature

The inflorescence is a spike with multi-floreted spikelets at the rachis nodes. The glumes are quite large and have characteristic shapes depending on the variety. The florets have beards or short awns extending from the tip of the floret. In contrast, durum wheat is typically bearded. The hull is removed during combining, and the grain is a reddish brown, oval to ovate to elliptical in shape. Durum grains are amber-colored, elliptical and larger than bread wheat kernels.

Key features

Hair on the margins of the auricles; leaf blades narrower than barley; spike with a single, multi-floreted spikelet at each rachis node

Similar species

Durum wheat *Triticum durum*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland	wetland		
reduced tillage	irrigation	roadsides		
forage crops, hay fields	turf	waste areas		

Seedling

Juvenile

Mature

Flower: *spike*

Key feature: *hairy auricles*

Wild Oats

Avena fatua

Family

Grass *Poaceae*

Seedling

Unlike most grasses, the seedling leaves of wild oats twist counter-clockwise. A tall, membranous ligule is easily visible at the collar region by the three to four leaf stage. Young plants may also be uprooted and checked for the easy-to-recognize seed, which is usually retained by the roots until later stages of growth.

Juvenile

Wild oats produces stems that are erect, smooth and hollow. Leaf blades are flat, broad and taper to a long thin point. They typically have a pronounced, light-colored midrib. Leaf sheaths are split and overlapping. Sparse hair is often present on both the blade and sheath in the collar region. Wild oats is very difficult to distinguish from cultivated oats at the juvenile stage.

Mature

The inflorescence of wild oats is a large, widely-branched panicle, often seen waving above the crop canopy. Each spikelet is usually comprised of two to three seeds surrounded by two large, papery glumes. The glumes remain attached to the panicle after the seeds have been dispersed. The seeds may vary greatly in color, from beige to brown to grey or black. They are usually hairy, especially around the base, but can be smooth. Despite these variations, all wild oat seeds are characterized by a large basal scar or "sucker mouth" and a prominent, bent, twisted awn. Seeds of tame oats usually are beige, smooth and lack the sucker mouth and long bent awn.

Key features

Leaves with counter-clockwise twist and tall, membranous ligule; large, widely-branched panicle inflorescence; seeds with sucker mouth and bent, twisted awn

Similar species

Oats *Avena sativa*

Life Cycle				
annual	winter annual	biennial	creeping perennial	simple perennial
Habitat				
conventional tillage	rangeland		wetland	
reduced tillage	irrigation		roadsides
forage crops, hay fields	turf		waste areas

Seedling

Mature: *panicles*

Key feature: *tall, membranous ligule*

Flower: *three spikelets*

Infestation

Glossary

Alternate – A leaf arrangement with one leaf per stem node.

Annual – A species that germinates in the spring, sets seed in the same year and then dies.

Aquatic – Growing in water.

Auricles – Claw-like, paired structures found at the collar in some grasses.

Awn – An extension at the tip of a floret or a glume in some grasses.

Axillary – Arising in a leaf axil.

Axil – see “Leaf Axil”

Axis – The central line of a plant or plant part.

Barb – A short, rigid bristle, often bent backwards.

Basal – At the base of a plant or plant structure. Often refers to the leaf arrangement of a rosette.

Beak – A prolonged tip.

Beard – A long awn.

Biennial – A species that germinates in the spring of the first year, producing a rosette that survives the winter in a dormant state. It resumes growth in the second year, flowers, sets seed and then dies.

Blade – The expanded, flat portion of a leaf above the sheath in grasses or above the petiole in broadleaf plants.

Bract – A modified leaf, reduced in size.

Bud – A dormant or unopened leaf, branch or flower, usually enclosed by protective bracts.

Bulb – An underground propagative structure consisting of fleshy bracts attached to a compressed stem.

Calyx – The collective term for the sepals of a flower.

Capsule – A type of fruit with inner compartments containing several to many seeds.

Clasping – Referring to the base of leaves, partially or entirely surrounding the stem.

Collar – Leaf tissue at the junction of the blade and sheath in grasses.

Compound Leaf – A leaf with a blade composed of two or more leaflets.

Compound Palmate Leaf – A compound leaf with leaflets radiating from a common point.

Compound Pinnate Leaf – A compound leaf with leaflets arranged on both sides of a central axis (rachis). Same as pinnately compound.

Compound Umbel – An inflorescence made up of many small umbels at the top of the main umbel.

Corolla – The collective term for the petals of a flower.

Corona – A structure between the stamens and the corolla in a Milkweed family flower.

Cotyledon – An embryonic leaf of a seed or seedling. The shape, texture and attachment of the two above-ground cotyledons of broadleaf weed seedlings are useful identification features.

Creeping Perennial – A species that survives for three or more seasons and, in that way, is similar to a simple perennial. However, a creeping perennial has a specialized method of vegetative propagation (rhizomes, stolons, budding rootstocks) in addition to seed production.

Crown – Shortened base of a stem below ground level.

Dicotyledon – A broad classification of flowering plants that are characterized by having two cotyledons in the seed, broad leaves with netted venation, flower parts in four's and five's and usually a taproot.

Disk Flower – Small, tube-like flower or floret found in the Sunflower family.

Divided – Refers to a leaf margin that is cut almost to the midrib.

Dormancy – A suspended or inhibited state of growth.

Elliptical – Elongated oval in shape.

Embryo – The rudimentary plant contained within the seed.

Endosperm – A food storage area in some seeds.

Entire – Refers to a leaf margin with no teeth or lobes.

Erect Stem – An upright or vertical stem.

Family – A grouping used in plant classification consisting of a number of similar genera.

Fibrous Root – A threadlike root. A plant with a fibrous root system does not have a main root.

Finely Divided – Refers to a leaf margin that is deeply cut into many slender segments, giving a lacy appearance.

Floral – Referring to a flower.

Floret – A small flower in the Grass or Sunflower family.

Flower Head – A type of inflorescence with many sessile flowers on a common, enlarged receptacle found in the Sunflower family. Also called a capitulum.

Flower Stalk – The stalk of a single flower, also called a pedicel.

Fruit – The mature, seed-containing structure that develops from a plant ovary.

Genus – A sub-division of a plant family. It consists of a number of similar plant species. (Plural of genus = genera)

Glabrous – Not hairy.

Glumes – A pair of bracts found at the base of a spikelet in grasses.

Heading – Stage of growth when the inflorescence is visibly emerging.

Hull – The collective term for the outer coverings of a grass family floret.

Inflorescence – Floral arrangement of a plant.

Internode – The part of a plant stem between two successive nodes.

Involucral Bract – A bract below the flower head in the Sunflower family.

Involucre – One or several whorls of bracts below a flower head in the Sunflower family.

Keel – Two united petals that form a pocket in a Pea family flower.

Keeled – Ridged, similar to the bottom of a boat.

Lanceolate – Shaped like the tip of a lance, broadest near the base and tapering to the tip.

Lateral – At or from the side.

Latex – White, milky juice in some plants.

Leaf Axil – The upper angle that a leaf makes with the stem to which it is attached.

Leaflet – One section of a compound leaf.

Ligule – A membranous or hairy projection at the junction of the leaf blade and leaf sheath in many grasses.

Linear – Long and narrow with almost parallel margins.

Lobed – Refers to an undulating leaf margin. Lobing varies from shallow to deep.

Margin – The edge of a leaf or other plant part.

Mealy – Granular texture.

Membranous – Similar to tissue paper or thin skin.

Midrib – The central vein of a leaf or other plant structure.

Monocotyledon – A broad classification of flowering plants characterized by having one cotyledon in the seed, narrow leaves with parallel venation, flower parts in three's and six's and a fibrous root system.

Node – Place on the stem from where leaves grow. Nodes are separated by internodes.

Ocrea – A sheath surrounding a stem node in the Buckwheat family.

Opposite – A leaf arrangement with two leaves per stem node, a paired leaf arrangement.

Oval – Widest at the centre, narrowing to both ends.

Ovary – A part of the pistil that contains the ovule(s), which ripen into seed(s).

Ovate – Egg-shaped.

Ovule – A part of the ovary that develops into a seed.

Palmate – Radiating pattern. A pattern of veins from a common point (as in a simple palmate leaf) or a pattern of leaflets radiating from a common point (as in a compound palmate leaf).

Panicle – A branched inflorescence in which the flowers or spikelets are attached by flower stalks to branches of the rachis.

Pappus – A calyx modified to form a tuft of bristles or hairs on a seed. The pappus is an adaptation to facilitate wind distribution of seed.

Petal – One of the parts of the corolla, often coloured.

Petiole – A leaf stalk.

Pinnate – Feather-like pattern. A pattern of veins emerging from both sides of a central vein (as in a simple pinnate leaf) or a pattern of leaflets arranged on both sides of a central axis (as in a compound pinnate leaf).

Pinnatifid – Divided in a pinnate pattern, but not to the central vein.

Pistil – The female part of a flower consisting of ovary, style and stigma.

Pod (Seedpod) – A type of fruit with many seeds.

Prickle – A weak, spiny outgrowth.

Prostrate Stem – A horizontal stem which grows flat or low to the ground.

Pubescent – Covered with hairs.

Raceme – A type of inflorescence in which the flowers or spikelets have stalks and are attached to a central rachis.

Rachis – The central axis of an inflorescence or a compound leaf.

Ray Flower – A flower or floret with a flat, strap-shaped petal, found in the Sunflower family.

Receptacle – The enlarged end of a flower stalk that holds the floral parts. In the Sunflower family, many flowers are attached to a greatly enlarged receptacle.

Reticulate – Referring to netted or non-parallel venation on a leaf or other plant part.

Rhizome – A horizontal, underground stem with buds and roots at the nodes.

Rib – A prominent vein or nerve.

Rosette – A cluster of radiating leaves at ground level.

Scale – A leaf or bract greatly reduced in size.

Scalloped – Refers to shallow, rounded lobes on a leaf margin.

Seed – A seed is the basic unit of plant propagation. It is a ripened ovule containing the embryo and surrounded by an outer coat. Endosperm may or may not be present.

Seedling – A young plant that has grown from a seed.

Sepal – One of the individual parts of the calyx.

Septum – A partition or wall, usually inside a fruit.

Sessile – Without a stalk.

Sheath – The lower, tubular portion of a grass leaf enclosing the stem.

Simple Leaf – A leaf with a single, unsegmented blade.

Simple Palmate Leaf – A simple leaf with veins radiating from a common point.

Simple Perennial – A plant species that survives for three or more seasons. Each spring the plant regrows from stored root and crown reserves. Seed production may occur in the first season and in each subsequent year. Spread of a simple perennial weed species is primarily by seed.

Simple Pinnate Leaf – A simple leaf with a central vein and smaller offshoots, similar to a feather.

Species – The basic unit of plant classification. Each weed in this book is a separate plant species.

Spike – An inflorescence in which sessile spikelets or flowers are attached to a central rachis.

Spikelet – A unit of the inflorescence in grasses, usually consisting of two outer glumes and one or more florets.

Spine – A sharp, rigid projection.

Stamen – The male, pollen-producing part of a flower.

Standard – The large, upper petal of a Pea family flower.

Star-shaped Hairs – Branched hairs that resemble a star.

Stem – The main supporting axis of a plant, divided into nodes and internodes.

Stigma – The top of a pistil, which is adapted (sticky, feathery) to receive pollen.

Stipules – Paired, leaf-like structures at a stem node.

Stolon – A creeping, above-ground, horizontal stem that roots at the nodes.

Style – Portion of a pistil between the ovary and the stigma.

Succulent – Fleshy.

Taproot – A thickened, vertical root.

Terminal – At the end of a stem or branch.

Terrestrial – Growing on land.

Toothed – A general term to describe a saw-like margin of a leaf or other structure.

Trifoliolate Leaf – A compound leaf with three leaflets.

Tuber – Enlarged portion of a stem or root.

Tubercle – A small, rounded projection, usually on a fruit or seed.

Tufted – In a clump.

Twining Stem – A weak stem that wraps around other plants or objects for support.

Umbel – A flat-topped inflorescence with the flower stalks radiating from a common point.

Valve – One section of the wall of a seedpod or seed capsule.

Veins – The visible transporting tissue of a leaf or other plant structure.

Venation – The pattern of veins on a leaf or other plant structure.

Whorl – A generic term for three or more plant structures arising from a common point.

Whorled – A type of leaf arrangement with three or more leaves per stem node.

Wing – A thin, flattened extension of a seed, fruit or other plant structure.

Wing Petal – One of the two lateral petals of a Pea family flower.

Winter Annual – A species that germinates in the fall and survives the winter as a dormant rosette. It resumes growth in the spring, sets seed in early summer and then dies.

Index

Organized by Common Name

Common name	Scientific name	Family name		Page
absinth	<i>Artemisia absinthium</i>	Sunflower	<i>Asteraceae</i>	124
American dragonhead	<i>Dracocephalum parviflorum</i>	Mint	<i>Lamiaceae</i>	46
American stinging nettle	<i>see stinging nettle</i>			234
annual sow-thistle	<i>similar to spiny annual sow-thistle</i>			174
annual sunflower	<i>similar to prairie sunflower</i>			166
Argentine canola	<i>Brassica napus</i>	Mustard	<i>Brassicaceae</i>	126
baby's-breath	<i>Gypsophila paniculata</i>	Pink	<i>Caryophyllaceae</i>	70
ball mustard	<i>Neslia paniculata</i>	Mustard	<i>Brassicaceae</i>	128
barley	<i>Hordeum vulgare</i>	Grass	<i>Poaceae</i>	196
barnyard grass	<i>Echinochloa crusgalli</i>	Grass	<i>Poaceae</i>	198
bedstraw	<i>see cleavers</i>			78
biennial wormwood	<i>similar to absinth</i>			124
black henbane	<i>Hyoscyamus niger</i>	Nightshade	<i>Solanaceae</i>	130
black medick	<i>Medicago lupulina</i>	Pea	<i>Fabaceae</i>	132
black nightshade	<i>similar to hairy nightshade</i>			90
bladder campion	<i>Silene cucubalus</i>	Pink	<i>Caryophyllaceae</i>	72
blue buttons	<i>see field scabious</i>			50
blue devil	<i>see blueweed</i>			12
blue lettuce	<i>similar to prickly lettuce</i>			168
bluebur	<i>Lappula squarrosa</i>	Borage	<i>Boraginaceae</i>	10
blueweed	<i>Echium vulgare</i>	Borage	<i>Boraginaceae</i>	12
bread wheat	<i>see wheat</i>			240
broad-leaved plantain	<i>Plantago major</i>	Plantain	<i>Plantaginaceae</i>	200
broomcorn millet	<i>see proso millet</i>			222
buffalo-bean	<i>see golden-bean</i>			148
bull thistle	<i>Cirsium vulgare</i>	Sunflower	<i>Asteraceae</i>	22
burweed	<i>see false ragweed</i>			206
butter-and-eggs	<i>see yellow toadflax</i>			192

Organized by Common Name

Common name	Scientific name	Family name		Page
Canada fleabane	<i>Erigeron canadensis</i>	Sunflower	<i>Asteraceae</i>	74
Canada thistle	<i>Cirsium arvense</i>	Sunflower	<i>Asteraceae</i>	24
cheat grass	<i>see downy brome</i>			204
cheeseweed	<i>see round-leaved mallow</i>			100
chickweed	<i>Stellaria media</i>	Pink	<i>Caryophyllaceae</i>	76
cleavers	<i>Galium aparine</i>	Madder	<i>Rubiaceae</i>	78
cocklebur	<i>Xanthium strumarium</i>	Sunflower	<i>Asteraceae</i>	202
common burdock	<i>Arctium minus</i>	Sunflower	<i>Asteraceae</i>	26
common groundsel	<i>Senecio vulgaris</i>	Sunflower	<i>Asteraceae</i>	134
common mullein	<i>Verbascum thapsus</i>	Snapdragon	<i>Scrophulariaceae</i>	136
common pepper-grass	<i>Lepidium densiflorum</i>	Mustard	<i>Brassicaceae</i>	80
common tansy	<i>see tansy</i>			178
compass plant	<i>see prickly lettuce</i>			168
corn spurry	<i>Spergula arvensis</i>	Pink	<i>Caryophyllaceae</i>	82
couch grass	<i>see quack grass</i>			226
cow cockle	<i>Vaccaria pyramidata</i>	Pink	<i>Caryophyllaceae</i>	48
creeping bellflower	<i>Campanula rapunculoides</i>	Bluebell	<i>Campanulaceae</i>	28
creeping Charlie	<i>see ground-ivy</i>			16
creeping thistle	<i>see Canada thistle</i>			24
cupped nightshade	<i>see hairy nightshade</i>			90
curlycup gumweed	<i>see gumweed</i>			150
cutleaf nightshade	<i>see wild tomato</i>			120
Dalmatian toadflax	<i>similar to yellow toadflax</i>			192
dandelion	<i>Taraxacum officinale</i>	Sunflower	<i>Asteraceae</i>	138
death camas	<i>Zygadenus gramineus</i>	Lily	<i>Liliaceae</i>	84
devil's paint-brush	<i>see orange hawkweed</i>			156
diffuse knapweed	<i>Centaurea diffusa</i>	Sunflower	<i>Asteraceae</i>	86
dog mustard	<i>Erucastrum gallicum</i>	Mustard	<i>Brassicaceae</i>	140

Organized by Common Name

Common name	Scientific name	Family name		Page
doorweed	<i>see prostrate knotweed</i>			60
downy brome	<i>Bromus tectorum</i>	Grass	<i>Poaceae</i>	204
downy chess	<i>see downy brome</i>			204
durum wheat	<i>similar to wheat</i>			240
Dutch clover	<i>see white clover</i>			110
early yellow locoweed	<i>Oxytropis sericea</i>	Pea	<i>Fabaceae</i>	142
fairy candelabra	<i>see pygmyflower</i>			98
false cleavers	<i>similar to cleavers</i>			78
false ragweed	<i>Iva xanthifolia</i>	Sunflower	<i>Asteraceae</i>	206
field bindweed	<i>Convolvulus arvensis</i>	Morning-glory	<i>Convolvulaceae</i>	88
field dock	<i>Rumex pseudonatronatus</i>	Buckwheat	<i>Polygonaceae</i>	208
field horsetail	<i>Equisetum arvense</i>	Horsetail	<i>Equisetaceae</i>	210
field scabious	<i>Knautia arvensis</i>	Teasel	<i>Dipsacaceae</i>	50
fireweed	<i>similar to purple loosestrife</i>			34
flax	<i>Linum usitatissimum</i>	Flax	<i>Linaceae</i>	14
flixweed	<i>Descurainia sophia</i>	Mustard	<i>Brassicaceae</i>	144
foxtail barley	<i>Hordeum jubatum</i>	Grass	<i>Poaceae</i>	212
Frenchweed	<i>see stinkweed</i>			108
fringed sagebrush	<i>see pasture sage</i>			158
garden bluebell	<i>see creeping bellflower</i>			28
gill-over-the-ground	<i>see ground-ivy</i>			16
goat's-beard	<i>Tragopogon dubius</i>	Sunflower	<i>Asteraceae</i>	146
golden-bean	<i>Thermopsis rhombifolia</i>	Pea	<i>Fabaceae</i>	148
green-flowered pepper-grass	<i>see common pepper-grass</i>			80
green foxtail	<i>Setaria viridis</i>	Grass	<i>Poaceae</i>	214
green smartweed	<i>similar to pale smartweed</i>			56
green tansy mustard	<i>similar to flixweed</i>			144
grey tansy mustard	<i>similar to flixweed</i>			144

Organized by Common Name

Common name	Scientific name	Family name		Page
ground-ivy	<i>Glechoma hederacea</i>	Mint	<i>Lamiaceae</i>	16
gumweed	<i>Grindelia squarrosa</i>	Sunflower	<i>Asteraceae</i>	150
hairy nightshade	<i>Solanum sarrachoides</i>	Nightshade	<i>Solanaceae</i>	90
hard red spring wheat	<i>see wheat</i>			240
hemp-nettle	<i>Galeopsis tetrahit</i>	Mint	<i>Lamiaceae</i>	52
henbit	<i>Lamium amplexicaule</i>	Mint	<i>Lamiaceae</i>	54
hitch-hiker	<i>see bluebur</i>			10
horseweed	<i>see Canada fleabane</i>			74
hound's-tongue	<i>Cynoglossum officinale</i>	Borage	<i>Boraginaceae</i>	30
Johnny-go-to-bed-at-noon	<i>see goat's beard</i>			146
knotgrass	<i>see prostrate knotweed</i>			60
knotweed	<i>see prostrate knotweed</i>			60
kochia	<i>Kochia scoparia</i>	Goosefoot	<i>Chenopodiaceae</i>	216
lady's-thumb	<i>similar to pale smartweed</i>			56
lamb's-quarters	<i>Chenopodium album</i>	Goosefoot	<i>Chenopodiaceae</i>	218
leafy spurge	<i>Euphorbia esula</i>	Spurge	<i>Euphorbiaceae</i>	152
low larkspur	<i>similar to tall larkspur</i>			40
mare's tail	<i>see field horsetail</i>			210
marsh elder	<i>see false ragweed</i>			206
millet	<i>see green foxtail</i>			214
mouse-eared chickweed	<i>similar to chickweed</i>			76
musk thistle	<i>see nodding thistle</i>			32
narrow-leaved hawk's-beard	<i>Crepis tectorum</i>	Sunflower	<i>Asteraceae</i>	154
narrow-leaved milk-vetch	<i>Astragalus pectinatus</i>	Pea	<i>Fabaceae</i>	92
net-seeded lamb's quarters	<i>similar to lamb's-quarters</i>			218
night-flowering catchfly	<i>Silene noctiflora</i>	Pink	<i>Caryophyllaceae</i>	94
nodding thistle	<i>Carduus nutans</i>	Sunflower	<i>Asteraceae</i>	32
oak-leaved goosefoot	<i>similar to spear-leaved goosefoot</i>			232

Organized by Common Name

Common name	Scientific name	Family name		Page
oats	<i>similar to wild oats</i>			242
orange hawkweed	<i>Hieracium aurantiacum</i>	Sunflower	<i>Asteraceae</i>	156
ox-eye daisy	<i>Chrysanthemum leucanthemum</i>	Sunflower	<i>Asteraceae</i>	96
pale smartweed	<i>Polygonum lapathifolium</i>	Buckwheat	<i>Polygonaceae</i>	56
pasture sage	<i>Artemisia frigida</i>	Sunflower	<i>Asteraceae</i>	158
penny cress	<i>see stinkweed</i>			108
pepperweed	<i>see common pepper-grass</i>			80
perennial sow-thistle	<i>Sonchus arvensis</i>	Sunflower	<i>Asteraceae</i>	160
perennial sow-thistle	<i>similar to arvensis</i>			160
Persian darnel	<i>Lolium persicum</i>	Grass	<i>Poaceae</i>	220
pineappleweed	<i>Matricaria matricarioides</i>	Sunflower	<i>Asteraceae</i>	162
pinegrass	<i>see field horsetail</i>			210
Polish canola	<i>Brassica rapa</i>	Mustard	<i>Brassicaceae</i>	164
prairie onion	<i>similar to death camas</i>			84
prairie rose	<i>similar to prickly rose</i>			58
prairie sunflower	<i>Helianthus petiolaris</i>	Sunflower	<i>Asteraceae</i>	166
prickly annual sow-thistle	<i>see spiny annual sow-thistle</i>			174
prickly lettuce	<i>Lactuca scariola</i>	Sunflower	<i>Asteraceae</i>	168
prickly rose	<i>Rosa acicularis</i>	Rose	<i>Rosaceae</i>	58
proso millet	<i>Panicum miliaceum</i>	Grass	<i>Poaceae</i>	222
prostrate knotweed	<i>Polygonum aviculare</i>	Buckwheat	<i>Polygonaceae</i>	60
prostrate pigweed	<i>Amaranthus blitoides</i>	Amaranth	<i>Amaranthaceae</i>	224
purple loosestrife	<i>Lythrum salicaria</i>	Loosestrife	<i>Lythraceae</i>	34
purslane	<i>Portulaca oleracea</i>	Purslane	<i>Portulacaceae</i>	170
pygmyflower	<i>Androsace septentrionalis</i>	Primrose	<i>Primulaceae</i>	98
quack grass	<i>Elytrigia repens</i>	Grass	<i>Poaceae</i>	226
redroot pigweed	<i>Amaranthus retroflexus</i>	Amaranth	<i>Amaranthaceae</i>	228
redstem filaree	<i>see stork's-bill</i>			64

Organized by Common Name

Common name	Scientific name	Family name		Page
rough cinquefoil	<i>Potentilla norvegica</i>	Rose	<i>Rosaceae</i>	172
round-leaved mallow	<i>Malva rotundifolia</i>	Mallow	<i>Malvaceae</i>	100
Russian knapweed	<i>Centaurea repens</i>	Sunflower	<i>Asteraceae</i>	36
Russian thistle	<i>Salsola pestifer</i>	Goosefoot	<i>Chenopodiaceae</i>	230
scentless chamomile	<i>Matricaria perforata</i>	Sunflower	<i>Asteraceae</i>	102
scentless mayweed	<i>see scentless chamomile</i>			102
scouring rush	<i>see field horsetail</i>			210
shepherd's purse	<i>Capsella bursa-pastoris</i>	Mustard	<i>Brassicaceae</i>	104
showy milkweed	<i>Asclepias speciosa</i>	Milkweed	<i>Asclepiadaceae</i>	62
silky lupine	<i>Lupinus sericeus</i>	Pea	<i>Fabaceae</i>	18
silvery lupine	<i>similar to silky lupine</i>			18
smooth brome	<i>similar to downy brome</i>			204
soapwort	<i>see cow cockle</i>			48
spear-leaved goosefoot	<i>Monolepis nutalliana</i>	Goosefoot	<i>Chenopodiaceae</i>	232
spiny annual sow-thistle	<i>Sonchus asper</i>	Sunflower	<i>Asteraceae</i>	174
spotted knapweed	<i>Centaurea maculosa</i>	Sunflower	<i>Asteraceae</i>	38
spotted water-hemlock	<i>Cicuta maculata</i>	Parsley	<i>Apiaceae</i>	106
stickseed	<i>see bluebur</i>			10
stickweed	<i>see bluebur</i>			10
sticky cockle	<i>see night-flowering catchfly</i>			94
stinging nettle	<i>Urtica dioica</i>	Nettle	<i>Urticaceae</i>	234
stinking mayweed	<i>similar to scentless chamomile</i>			102
stinkweed	<i>Thlaspi arvense</i>	Mustard	<i>Brassicaceae</i>	108
stork's-bill	<i>Erodium cicutarium</i>	Geranium	<i>Geraniaceae</i>	64
striate knotweed	<i>similar to prostrate knotweed</i>			60
swamp smartweed	<i>similar to water smartweed</i>	Rose	<i>Rosaceae</i>	66
tall buttercup	<i>Ranunculus acris</i>	Buttercup	<i>Ranunculaceae</i>	176
tall larkspur	<i>Delphinium glaucum</i>	Buttercup	<i>Ranunculaceae</i>	40

Organized by Common Name

Common name	Scientific name	Family name		Page
tansy	<i>Tanacetum vulgare</i>	Sunflower	<i>Asteraceae</i>	178
tansy mustard	see flixweed			144
tartary buckwheat	<i>Fagopyrum tataricum</i>	Buckwheat	<i>Polygonaceae</i>	236
thyme-leaved spurge	<i>Euphorbia serpyllifolia</i>	Spurge	<i>Euphorbiaceae</i>	238
tumble mustard	<i>Sisymbrium altissimum</i>	Mustard	<i>Brassicaceae</i>	180
tumble pigweed	similar to prostrate pigweed			224
turnip rape	see Polish canola			164
twitch grass	see quack grass			226
two-grooved milk-vetch	<i>Astragalus bisulcatus</i>	Pea	<i>Fabaceae</i>	42
Viper's bugloss	see blueweed			12
water smartweed	<i>Polygonum amphibium</i>	Buckwheat	<i>Polygonaceae</i>	66
water-parsnip	similar to spotted water-hemlock			106
western dock	similar to field dock			208
western salsify	see goat's beard			146
wheat	<i>Triticum aestivum</i>	Grass	<i>Poaceae</i>	240
white camas	similar to death camas			84
white clover	<i>Trifolium repens</i>	Pea	<i>Fabaceae</i>	110
white cockle	<i>Silene alba</i>	Pink	<i>Caryophyllaceae</i>	112
white daisy	see ox-eye daisy			96
white sweet clover	similar to yellow sweet clover			190
wild barley	see foxtail barley			212
wild buckwheat	<i>Polygonum convolvulus</i>	Buckwheat	<i>Polygonaceae</i>	114
wild caraway	<i>Carum carvi</i>	Parsley	<i>Apiaceae</i>	116
wild carrot	similar to wild caraway			116
wild chamomile	similar to scentless chamomile			102
wild licorice	<i>Glycyrrhiza lepidota</i>	Pea	<i>Fabaceae</i>	118
wild millet	see green foxtail			214
wild mustard	<i>Brassica kaber</i>	Mustard	<i>Brassicaceae</i>	182

Organized by Common Name

Common name	Scientific name	Family name		Page
wild oats	<i>Avena fatua</i>	Grass	<i>Poaceae</i>	242
wild portulaca	see purslane			170
wild radish	<i>Raphanus raphanistrum</i>	Mustard	<i>Brassicaceae</i>	184
wild rose	see prickly rose			58
wild snapdragon	see yellow toadflax			192
wild tomato	<i>Solanum triflorum</i>	Nightshade	<i>Solanaceae</i>	120
witch grass	similar to proso millet			222
wood whitlow-grass	<i>Draba nemorosa</i>	Mustard	<i>Brassicaceae</i>	186
Woods' rose	similar to prickly rose			58
wormwood	see absinth			124
yellow foxtail	similar to green foxtail			214
yellow nutsedge	<i>Cyperus esculentus</i>	Sedge	<i>Cyperaceae</i>	188
yellow star-thistle	similar to spotted knapweed			38
yellow star-thistle	similar to diffuse knapweed			86
yellow sweet clover	<i>Melilotus officinalis</i>	Pea	<i>Fabaceae</i>	190
yellow toadflax	<i>Linaria vulgaris</i>	Snapdragon	<i>Scrophulariaceae</i>	192
yellow whitlow-grass	see wood whitlow-grass			186
yellow-flowered hemp-nettle	similar to hemp-nettle			52

Organized by Scientific Name

Scientific name	Common name	Family name	Page
<i>Amaranthus blitoides</i>	prostrate pigweed	Amaranth	224
<i>Amaranthus retroflexus</i>	redroot pigweed	Amaranth	228
<i>Androsace septentrionalis</i>	pygmyflower	Primrose	98
<i>Arctium minus</i>	common burdock	Sunflower	26
<i>Artemisia absinthium</i>	absinth	Sunflower	124
<i>Artemisia frigida</i>	pasture sage	Sunflower	158
<i>Asclepias speciosa</i>	showy milkweed	Milkweed	62
<i>Astragalus bisulcatus</i>	two-grooved milk-vetch	Pea	42
<i>Astragalus pectinatus</i>	narrow-leaved milk-vetch	Pea	92
<i>Avena fatua</i>	wild oats	Grass	242
<i>Brassica kaber</i>	wild mustard	Mustard	182
<i>Brassica napus</i>	Argentine canola	Mustard	126
<i>Brassica rapa</i>	Polish canola	Mustard	164
<i>Bromus tectorum</i>	downy brome	Grass	204
<i>Campanula rapunculoides</i>	creeping bellflower	Bluebell	28
<i>Capsella bursa-pastoris</i>	shepherd's purse	Mustard	104
<i>Carduus nutans</i>	nodding thistle	Sunflower	32
<i>Carum carvi</i>	wild caraway	Parsley	116
<i>Centaurea diffusa</i>	diffuse knapweed	Sunflower	86
<i>Centaurea maculosa</i>	spotted knapweed	Sunflower	38
<i>Centaurea repens</i>	Russian knapweed	Sunflower	36
<i>Chenopodium album</i>	lamb's-quarters	Goosefoot	218
<i>Chrysanthemum leucanthemum</i>	ox-eye daisy	Sunflower	96
<i>Cicuta maculata</i>	spotted water-hemlock	Parsley	106
<i>Cirsium arvense</i>	Canada thistle	Sunflower	24
<i>Cirsium vulgare</i>	bull thistle	Sunflower	22
<i>Convolvulus arvensis</i>	field bindweed	Morning-glory	88
<i>Crepis tectorum</i>	narrow-leaved hawk's-beard	Sunflower	154

Organized by Scientific Name

Scientific name	Common name	Family name	Page
<i>Cynoglossum officinale</i>	hound's-tongue	Borage	30
<i>Cyperus esculentus</i>	yellow nut sedge	Sedge	188
<i>Delphinium glaucum</i>	tall larkspur	Buttercup	40
<i>Descurainia sophia</i>	flixweed	Mustard	144
<i>Draba nemorosa</i>	wood whitlow-grass	Mustard	186
<i>Dracocephalum parviflorum</i>	American dragonhead	Mint	46
<i>Echinochloa crusgalli</i>	barnyard grass	Grass	198
<i>Echium vulgare</i>	blueweed	Borage	12
<i>Elytrigia repens</i>	quack grass	Grass	226
<i>Equisetum arvense</i>	field horsetail	Horsetail	210
<i>Erigeron canadensis</i>	Canada fleabane	Sunflower	74
<i>Erodium cicutarium</i>	stork's-bill	Geranium	64
<i>Erucastrum gallicum</i>	dog mustard	Mustard	140
<i>Euphorbia esula</i>	leafy spurge	Spurge	152
<i>Euphorbia serpyllifolia</i>	thyme-leaved spurge	Spurge	238
<i>Fagopyrum tataricum</i>	tartary buckwheat	Buckwheat	236
<i>Galeopsis tetrahit</i>	hemp-nettle	Mint	52
<i>Galium aparine</i>	cleavers	Madder	78
<i>Glechoma hederacea</i>	ground-ivy	Mint	16
<i>Glycyrrhiza lepidota</i>	wild licorice	Pea	118
<i>Grindelia squarrosa</i>	gumweed	Sunflower	150
<i>Gypsophila paniculata</i>	baby's-breath	Pink	70
<i>Helianthus petiolaris</i>	prairie sunflower	Sunflower	166
<i>Hieracium aurantiacum</i>	orange hawkweed	Sunflower	156
<i>Hordeum jubatum</i>	foxtail barley	Grass	212
<i>Hordeum vulgare</i>	barley	Grass	196
<i>Hyoscyamus niger</i>	black henbane	Nightshade	130
<i>Iva xanthifolia</i>	false ragweed	Sunflower	206

Organized by Scientific Name

Scientific name	Common name	Family name	Page
<i>Knautia arvensis</i>	field scabious	Teasel	50
<i>Kochia scoparia</i>	kochia	Goosefoot	216
<i>Lactuca scariola</i>	prickly lettuce	Sunflower	168
<i>Lamium amplexicaule</i>	henbit	Mint	54
<i>Lappula squarrosa</i>	bluebur	Borage	10
<i>Lepidium densiflorum</i>	common pepper-grass	Mustard	80
<i>Linaria vulgaris</i>	yellow toadflax	Snapdragon	192
<i>Linum usitatissimum</i>	flax	Flax	14
<i>Lolium persicum</i>	Persian darnel	Grass	220
<i>Lupinus sericeus</i>	silky lupine	Pea	18
<i>Lythrum salicaria</i>	purple loosestrife	Loosestrife	34
<i>Malva rotundifolia</i>	round-leaved mallow	Mallow	100
<i>Matricaria matricarioides</i>	pineappleweed	Sunflower	162
<i>Matricaria perforata</i>	scentless chamomile	Sunflower	102
<i>Medicago lupulina</i>	black medick	Pea	132
<i>Melilotus officinalis</i>	yellow sweet clover	Pea	190
<i>Monolepis nutalliana</i>	spear-leaved goosefoot	Goosefoot	232
<i>Neslia paniculata</i>	ball mustard	Mustard	128
<i>Oxytropis sericea</i>	early yellow locoweed	Pea	142
<i>Panicum miliaceum</i>	proso millet	Grass	222
<i>Plantago major</i>	broad-leaved plantain	Plantain	200
<i>Polygonum amphibium</i>	water smartweed	Buckwheat	66
<i>Polygonum aviculare</i>	prostrate knotweed	Buckwheat	60
<i>Polygonum convolvulus</i>	wild buckwheat	Buckwheat	114
<i>Polygonum lapathifolium</i>	pale smartweed	Buckwheat	56
<i>Portulaca oleracea</i>	purslane	Purslane	170
<i>Potentilla norvegica</i>	rough cinquefoil	Rose	172
<i>Ranunculus acris</i>	tall buttercup	Buttercup	176

Organized by Scientific Name

Scientific name	Common name	Family name	Page
<i>Raphanus raphanistrum</i>	wild radish	Mustard	184
<i>Rosa acicularis</i>	prickly rose	Rose	58
<i>Rumex pseudonatronatus</i>	field dock	Buckwheat	208
<i>Salsola pestifer</i>	Russian thistle	Goosefoot	230
<i>Senecio vulgaris</i>	common groundsel	Sunflower	134
<i>Setaria viridis</i>	green foxtail	Grass	214
<i>Silene alba</i>	white cockle	Pink	112
<i>Silene cucubalus</i>	bladder campion	Pink	72
<i>Silene noctiflora</i>	night-flowering catchfly	Pink	94
<i>Sisymbrium altissimum</i>	tumble mustard	Mustard	180
<i>Solanum sarrachoides</i>	hairy nightshade	Nightshade	90
<i>Solanum triflorum</i>	wild tomato	Nightshade	120
<i>Sonchus arvensis</i>	perennial sow-thistle	Sunflower	160
<i>Sonchus asper</i>	spiny annual sow-thistle	Sunflower	174
<i>Spergula arvensis</i>	corn spurry	Pink	82
<i>Stellaria media</i>	chickweed	Pink	76
<i>Tanacetum vulgare</i>	tansy	Sunflower	178
<i>Taraxacum officinale</i>	dandelion	Sunflower	138
<i>Thermopsis rhombifolia</i>	golden-bean	Pea	148
<i>Thlaspi arvense</i>	stinkweed	Mustard	108
<i>Tragopogon dubius</i>	goat's-beard	Sunflower	146
<i>Trifolium repens</i>	white clover	Pea	110
<i>Triticum aestivum</i>	wheat	Grass	240
<i>Urtica dioica</i>	stinging nettle	Nettle	234
<i>Vaccaria pyramidata</i>	cow cockle	Pink	48
<i>Verbascum thapsus</i>	common mullein	Snapdragon	136
<i>Xanthium strumarium</i>	cocklebur	Sunflower	202
<i>Zygadenus gramineus</i>	death camas	Lily	84

Organized by Family Name

Family name		Common name	Scientific name	Page
Amaranth	<i>Amaranthaceae</i>	prostrate pigweed	<i>Amaranthus blitoides</i>	224
		redroot pigweed	<i>Amaranthus retroflexus</i>	228
Bluebell	<i>Campanulaceae</i>	creeping bellflower	<i>Campanula rapunculoides</i>	28
Borage	<i>Boraginaceae</i>	bluebur	<i>Lappula squarrosa</i>	10
		blueweed	<i>Echium vulgare</i>	12
		hound's-tongue	<i>Cynoglossum officinale</i>	30
Buckwheat	<i>Polygonaceae</i>	field dock	<i>Rumex pseudonatronatus</i>	208
		tartary buckwheat	<i>Fagopyrum tataricum</i>	236
		pale smartweed	<i>Polygonum lapathifolium</i>	56
		prostrate knotweed	<i>Polygonum aviculare</i>	60
		water smartweed	<i>Polygonum amphibium</i>	66
		wild buckwheat	<i>Polygonum convolvulus</i>	114
Buttercup	<i>Ranunculaceae</i>	tall larkspur	<i>Delphinium glaucum</i>	40
		tall buttercup	<i>Ranunculus acris</i>	176
Flax	<i>Linaceae</i>	flax	<i>Linum usitatissimum</i>	14
Geranium	<i>Geraniaceae</i>	stork's-bill	<i>Erodium cicutarium</i>	64
Goosefoot	<i>Chenopodiaceae</i>	kochia	<i>Kochia scoparia</i>	216
		lamb's-quarters	<i>Chenopodium album</i>	218
		Russian thistle	<i>Salsola pestifer</i>	230
		spear-leaved goosefoot	<i>Monolepis nutalliana</i>	232
Grass	<i>Poaceae</i>	barley	<i>Hordeum vulgare</i>	196
		barnyard grass	<i>Echinochloa crusgalli</i>	198
		downy brome	<i>Bromus tectorum</i>	204
		foxtail barley	<i>Hordeum jubatum</i>	212
		green foxtail	<i>Setaria viridis</i>	214
		Persian darnel	<i>Lolium persicum</i>	220
		proso millet	<i>Panicum miliaceum</i>	222
		quack grass	<i>Elytrigia repens</i>	226

Organized by Family Name

Family name		Common name	Scientific name	Page
Grass	<i>Poaceae</i>	wheat	<i>Triticum aestivum</i>	240
		wild oats	<i>Avena fatua</i>	242
Horsetail	<i>Equisetaceae</i>	field horsetail	<i>Equisetum arvense</i>	210
Lily	<i>Liliaceae</i>	death camas	<i>Zygadenus gramineus</i>	84
Loosestrife	<i>Lythraceae</i>	purple loosestrife	<i>Lythrum salicaria</i>	34
Madder	<i>Rubiaceae</i>	cleavers	<i>Galium aparine</i>	78
Mallow	<i>Malvaceae</i>	round-leaved mallow	<i>Malva rotundifolia</i>	100
Milkweed	<i>Asclepiadaceae</i>	showy milkweed	<i>Asclepias speciosa</i>	62
Mint	<i>Lamiaceae</i>	ground-ivy	<i>Glechoma hederacea</i>	16
		American dragonhead	<i>Dracocephalum parviflorum</i>	46
		hemp-nettle	<i>Galeopsis tetrahit</i>	52
		henbit	<i>Lamium amplexicaule</i>	54
Morning-glory	<i>Convolvulaceae</i>	field bindweed	<i>Convolvulus arvensis</i>	88
Mustard	<i>Brassicaceae</i>	common pepper-grass	<i>Lepidium densiflorum</i>	80
		shepherd's purse	<i>Capsella bursa-pastoris</i>	104
		stinkweed	<i>Thlaspi arvense</i>	108
		Argentine canola	<i>Brassica napus</i>	126
		ball mustard	<i>Neslia paniculata</i>	128
		dog mustard	<i>Erucastrum gallicum</i>	140
		flixweed	<i>Descurainia sophia</i>	144
		Polish canola	<i>Brassica rapa</i>	164
		tumble mustard	<i>Sisymbrium altissimum</i>	180
		wild mustard	<i>Brassica kaber</i>	182
		wild radish	<i>Raphanus raphanistrum</i>	184
		wood whitlow-grass	<i>Draba nemorosa</i>	186
Nettle	<i>Urticaceae</i>	stinging nettle	<i>Urtica dioica</i>	234
Nightshade	<i>Solanaceae</i>	hairy nightshade	<i>Solanum sarrachoides</i>	90
		wild tomato	<i>Solanum triflorum</i>	120

Organized by Family Name

Family name		Common name	Scientific name	Page
Nightshade	<i>Solanaceae</i>	black henbane	<i>Hyoscyamus niger</i>	130
Parsley	<i>Apiaceae</i>	spotted water-hemlock	<i>Cicuta maculata</i>	106
		wild caraway	<i>Carum carvi</i>	116
Pea	<i>Fabaceae</i>	silky lupine	<i>Lupinus sericeus</i>	18
		two-grooved milk-vetch	<i>Astragalus bisulcatus</i>	42
		narrow-leaved milk-vetch	<i>Astragalus pectinatus</i>	92
		white clover	<i>Trifolium repens</i>	110
		wild licorice	<i>Glycyrrhiza lepidota</i>	118
		black medick	<i>Medicago lupulina</i>	132
		early yellow locoweed	<i>Oxytropis sericea</i>	142
		golden-bean	<i>Thermopsis rhombifolia</i>	148
		yellow sweet clover	<i>Melilotus officinalis</i>	190
Pink	<i>Caryophyllaceae</i>	cow cockle	<i>Vaccaria pyramidata</i>	48
		baby's-breath	<i>Gypsophila paniculata</i>	70
		bladder campion	<i>Silene cucubalus</i>	72
		chickweed	<i>Stellaria media</i>	76
		corn spurry	<i>Spergula arvensis</i>	82
		night-flowering catchfly	<i>Silene noctiflora</i>	94
		white cockle	<i>Silene alba</i>	112
Plantain	<i>Plantaginaceae</i>	broad-leaved plantain	<i>Plantago major</i>	200
Primrose	<i>Primulaceae</i>	pygmyflower	<i>Androsace septentrionalis</i>	98
Purslane	<i>Portulacaceae</i>	purslane	<i>Portulaca oleracea</i>	170
Rose	<i>Rosaceae</i>	prickly rose	<i>Rosa acicularis</i>	58
		rough cinquefoil	<i>Potentilla norvegica</i>	172
Sedge	<i>Cyperaceae</i>	yellow nut sedge	<i>Cyperus esculentus</i>	188
Snapdragon	<i>Scrophulariaceae</i>	common mullein	<i>Verbascum thapsus</i>	136
		yellow toadflax	<i>Linaria vulgaris</i>	192
Spurge	<i>Euphorbiaceae</i>	thyme-leaved spurge	<i>Euphorbia serpyllifolia</i>	238

Organized by Family Name

Family name		Common name	Scientific name	Page
Spurge	<i>Euphorbiaceae</i>	leafy spurge	<i>Euphorbia esula</i>	152
Sunflower	<i>Asteraceae</i>	cocklebur	<i>Xanthium strumarium</i>	202
		false ragweed	<i>Iva xanthifolia</i>	206
		bull thistle	<i>Cirsium vulgare</i>	22
		Canada thistle	<i>Cirsium arvense</i>	24
		common burdock	<i>Arctium minus</i>	26
		nodding thistle	<i>Carduus nutans</i>	32
		Russian knapweed	<i>Centaurea repens</i>	36
		spotted knapweed	<i>Centaurea maculosa</i>	38
		Canada fleabane	<i>Erigeron canadensis</i>	74
		diffuse knapweed	<i>Centaurea diffusa</i>	86
		ox-eye daisy	<i>Chrysanthemum leucanthemum</i>	96
		scentless chamomile	<i>Matricaria perforata</i>	102
		absinth	<i>Artemisia absinthium</i>	124
		common groundsel	<i>Senecio vulgaris</i>	134
		dandelion	<i>Taraxacum officinale</i>	138
		goat's-beard	<i>Tragopogon dubius</i>	146
		gumweed	<i>Grindelia squarrosa</i>	150
		narrow-leaved hawk's-beard	<i>Crepis tectorum</i>	154
		orange hawkweed	<i>Hieracium aurantiacum</i>	156
		pasture sage	<i>Artemisia frigida</i>	158
		perennial sow-thistle	<i>Sonchus arvensis</i>	160
		pineappleweed	<i>Matricaria matricarioides</i>	162
		prairie sunflower	<i>Helianthus petiolaris</i>	166
		prickly lettuce	<i>Lactuca scariola</i>	168
		spiny annual sow-thistle	<i>Sonchus asper</i>	174
		tansy	<i>Tanacetum vulgare</i>	178
Teasel	<i>Dipsacaceae</i>	field scabious	<i>Knautia arvensis</i>	50

National Library of Canada
Bibliothèque nationale du Canada

3 3286 52099988 5

7 78753 06147 9

Printed in Canada

Alberta
AGRICULTURE, FOOD AND
RURAL DEVELOPMENT
AGDEX 640-4

**OLDS
COLLEGE**