**Ecological Land
Classification of
Big Hill Springs Provincial
Park**

NATURAL RESOURCES SERVICE
RESOURCE INFORMATION UNIT
BOW REGION

Publication No. 7403

ISBN: 0-7785-0418-2

**ECOLOGICAL LAND CLASSIFICATION
OF
BIG HILL SPRINGS PROVINCIAL PARK, ALBERTA**

Prepared For:

Natural Resources Service - Parks
Alberta Environmental Protection
Bow Region

By:

Ian Sutherland

Environmental Service
Alberta Environmental Protection
Bow Region

June, 1998

- ii

TABLE OF CONTENTS

1.0 INTRODUCTION.....	1
2.0 STUDY AREA DESCRIPTION.....	2
2.1 Location.....	2
2.2 Land Use History	2
2.3 Climate	3
2.4 Hydrology.....	3
2.5 Physiography and Bedrock Geology.....	4
2.6 Surficial Geology and Soils.....	4
2.7 Vegetation	6
2.8 Wildlife	7
3.0 METHODS.....	8
3.1 Field Survey	8
3.2 Data Analysis	8
3.3 Ecological Land Classification and Mapping	9
3.4 Ecosite Legend	9
4.0 RESULTS.....	10
4.1 Ecological Land Classification.....	10
4.2 Vegetation Community Types	10
Aspen Woodland.....	11
Balsam Poplar or Balsam Poplar – White Spruce Woodland	13
Shrubland – Willow dominated	16
Mixed Shrublands	19
Grasslands	23
Modified Grasslands	26
Grasslands -Wet meadows	29
4.3 Significant Plant Species.....	31
4.4 Significant Vegetation Cover Changes	31
5.0 ISSUES AND RECOMMENDATIONS	32
6.0 REFERENCES.....	35

Appendix 1: List of Vegetation Species

Appendix 2: List of Wildlife Observations

Appendix 3: Ecological Land Classification

LIST OF MAPS

1. Study Area Location
2. Map of Big Hill Springs Provincial Park
3. Changes in Vegetation Communities 1972-1997
4. Ecological Land Classification (in rear pouch)

ACKNOWLEDGEMENTS

Keith Ainsley (Resource Data Division, Edmonton) – project planning, field data collection;

Beth Cornish (Resource Data Division, Edmonton) – vegetation community typing;

Fine Zweier (Environmental Service, Southern East Slopes Region) – field data collection;

Margaret Bradley (Environmental Service, SES Region) – cartography;

Joyce Gould (Recreation & Protected Areas Division, Natural Resource Service) – vegetation inventory planning;

Rod Gow (District Ranger, NRS - SES Region) – project data; chief liaison;

Wanda Nadasde-Hogg (Park Ranger, NRS - SES Region) – project liaison;

Lyle Lambert (Seasonal Ranger, NRS - SES Region) – field data collection;

Jock Forster (Environmental Service, SES Region) – groundwater consulting;

Ebe Lorberg (Water Sciences Br., NRS, Edmonton) - groundwater consulting;

Steve Kassai and Olga Droppo (Calgary Field Naturalist Society) - wildlife data.

1. The first of the year was a very dry one, with little rain.

2. The second of the year was a very dry one, with little rain.

3. The third of the year was a very dry one, with little rain.

4. The fourth of the year was a very dry one, with little rain.

5. The fifth of the year was a very dry one, with little rain.

6. The sixth of the year was a very dry one, with little rain.

7. The seventh of the year was a very dry one, with little rain.

8. The eighth of the year was a very dry one, with little rain.

9. The ninth of the year was a very dry one, with little rain.

10. The tenth of the year was a very dry one, with little rain.

11. The eleventh of the year was a very dry one, with little rain.

1.0 INTRODUCTION

Big Hill Springs Provincial Park was established in 1957 to conserve a unique coulee environment and spring-fed creek system (AEP - Park File). In years since, a number of specialists and organizations have documented various aspects of the Park's resources including archaeological – cultural (Glenbow Foundation; University of Calgary: Park File), ecological (Wallis and Wershler 1972), hydrogeological (Curcio 1967, 1970; Borneuf 1983; Houseknecht 1984; Komex 1998), botanical (AEP 1994), and wildlife (Calgary Field Naturalist's Society).

The Park's landscape features contribute significantly to under-represented Natural History Themes in the Foothills Parkland Natural Subregion (Report 3: Alberta Protected Areas System Analysis [1994]; Alberta Environmental Protection, Edmonton).

A Park Management Plan (AEP 1996), currently at draft stage, raises a number of conservation issues and objectives, many of which are addressed in this update and synthesis of biophysical information, including:

- preservation of natural environment, features, species, and habitat;
- natural history interpretation;
- water quality and quantity;
- creekbank management;
- improvement of facilities and infrastructure (e.g. picnic sites, roads, trails);
- accommodate recreational activities;
- acquisition of select adjacent parcels of land;
- boundaries and fencing.

In this context, the overall objectives of this study were to:

- Complete a review of existing biophysical information;
- Conduct a biophysical inventory to current standards and scope;
- Conduct an analysis of any significant changes.

Secondary objectives included:

- Obtain new aerial photography of the Park and surrounding lands;
- Conduct an intensive baseline study of the springs and associated features;
- Make management recommendations.

2.0 STUDY AREA DESCRIPTION

2.1 Location

Big Hill Springs Provincial Park is located in the Big Hill Creek valley 15 km north of Cochrane in the north half of Section 29, Township 26, Range 3, west of the Fifth Meridian. It is accessible from Secondary Highway #567, east of the Junction with Highway #22 (Map 1).

It is situated, for the most part, in a side-coulee which leads from the Big Hill Creek valley and covers 32 hectares (AEP 1996) or 79 acres. Presently, the Park is distributed over two disjunct parcels of land – the main part established in 1957, and a 6-hectare parcel in the northwest corner of Section 29 (identified as the “northwest parcel” in this report) at the head of the coulee, acquired recently to preserve the primary springs. The two parcels are separated by about 150 meters of private land (Map 2). A full legal land survey has not been conducted to this point, and boundaries, fencelines, and area measurements are therefore approximate.

Biogeographically, the Park is in the Foothills Parkland Natural Subregion, at the northern end.

2.2 Land Use History

A constant source of water has long attracted people and wildlife to the Big Hill Springs area. Plains Indians hunted bison (evidently using the “jump” method) and camped near the springs (the AEP Park File documents findings of bone fragments and lithic tools). Lamoureux (1983) lists a “major” kill site and evidence of pictographs at Big Hill Springs, and “maybe” a major jump on the east face of Big Hill Springs coulee.

Ranchers arrived in the 1880’s to graze and water their cattle. [An anecdotal, but detailed history from this point to about 1970 is included in the Park File (AEP).] Industrial interest in the springs and creek began before the turn of the century: a creamery was in operation from 1891 to 1910, and a trout hatchery was tested in the 1951-1956 period.

Recreation, mainly picnicking became popular in the 1920’s and today is now the prime use of the area around the springs. Big Hill Springs Provincial Park was established in 1957. Increased visitor volume, heavy use, and environmental degradation into the 1970’s raised concerns among park officials, and was the impetus for early biophysical data collection and management planning (Alberta Recreation and Parks 1988; AEP Park File). In 1976 the Park was proposed for designation as a Preservation Park, and many conservation initiatives were tabled (Balogorodow 1976). A reconfiguration to “day use only” began in 1978-79 with the removal of an array of visitor services and facilities (park ranger, campground, playgrounds, roads, picnic sites, and electricity) and

ALBERTA

Edmonton
□

MAP 1

BIG HILL SPRINGS
PROV. PARK

★
□ Calgary

BIG HILL
SPRINGS
PROV. PARK

COCHRANE

TP 27

TP 26

TP 25 W5MER.

R.4

R.3

COCHRANE
LAKE

SCALE

Contour Interval 10 Metres

N

1A
To Calgary

RIVER

82

22

1A

766

114°30'00"

51°15'00"

Big Hill Springs Provincial Park

N 1/2 of Sec. 29 Tp. 26 Rg. 03 W5Mer.

LEGEND

- Park boundary
- ⊕ Spring or seep
- - - Trail
- * * * Fences (not all are shown)

reclamation of the sites. Day use exploration by visitors has affected the Park in terms of compacted, eroded, and denuded trails and paths (Alberta Recreation and Parks 1988). Today the facilities maintained include a parking lot, small picnic site, toilets, garbage bin, water well, and improved trails and footbridges.

Cattle ranching is the main land use in the surrounding area and at some sites within the Park under arrangements with a local rancher.

2.3 Climate

The Parkland Natural Region forms a biogeoclimatic transition zone between the drier grasslands of the plains and the moister coniferous forests of the Rocky Mountains. Mean annual precipitation is approximately 500mm in the northern part. Mean May to September precipitation is 290mm. The mean January temperature is -10°C and the mean July temperature is 14°C. The mean May to September temperature is 12-13°C and the frost-free period averages 90 days (Achuff 1994).

Conditions specific to the town of Cochrane, 8km to the southwest, are:

Temperature - Seasonal Averages	Jan. -11.8°C; April 3.3°C; July 16.4°C; Oct. 5.5°C
Frost Free Days	112
Annual Precipitation	50cm ; Rain: 28cm
Hours of Sunshine/year	2314

(from Town of Cochrane - Website Page 1997)

2.4 Hydrology

The hydrology and associated landforms are very distinctive and comprise a primary natural history theme of the Park (Lamoureux 1983). The Park is situated at the confluence of the spring-fed, first-order Big Hill Springs Creek and the somewhat larger Big Hill Creek, which flows down a glacial meltwater outlet from the township to the north, southwards to the Bow River at Cochrane (Map 1). Surficial drainage of the surrounding benchlands is either towards the ravine or into Big Hill Creek valley itself.¹

The Park is located in a groundwater discharge area, creating many springs and seeps, as well as unique hydrological conditions and features. The springs provide lower Big Hill Creek with its year-round source of water (Lamoureux 1983). The hydrology of the springs and Big Hill Springs Creek is treated more thoroughly in Komex (1998).

¹ Use of names and spelling of these creeks and valleys has been inconsistent. National (i.e. NTS) and provincial records and maps indicate **Bighill** Creek, but the spring-fed creek is officially unnamed (Alberta Environmental Protection, Resource Data Division; NAD83 Listing of Named Geographic Features 1997/02/12). The Park Management Plan (1996; in Draft) itself variously uses Big Spring Creek/coulee and Big Hill Spring Creek/valley; and Big Hill valley. Names used in this report (as per Alberta Recreation and Parks 1988) reflect widely-used colloquial versions.

2.5 Physiography and Bedrock Geology

Big Hill Springs Provincial Park is located at the western edge of the Interior Plains physiographic region where it meets the foothills of the Cordillera. The Interior Plains are composed of relatively undisturbed Phanerozoic sedimentary rocks, and in the Park the uppermost rock units are continental deposits of the Tertiary Porcupine Hills Formation overlying the Tertiary-Cretaceous Paskapoo Formation. These sediments were deposited as a result of erosion of the uplifting Rocky Mountains and foothills (during the late Cretaceous to early Eocene), and eastward transport of the sediments to form alluvial fans, stream, and lake sediments of the Alberta Syncline. As the sediments were deposited, the marine conditions that had dominated North America up to the end of the early Cretaceous were gradually replaced by continental conditions.

The (sedimentary) lithology of the Porcupine Hills Formation is sandstone (greywackes and sub-greywackes) and siltstone with minor shale. Significant outcrops are visible along ridges and valley walls throughout the area. The formation is composed of detrital quartz, chert, non-volcanic rock fragments and clastic carbonates (Yoon 1977; Park File).

2.6 Surficial Geology and Soils

During the Pleistocene Epoch, the Laurentide continental ice sheet advanced southward from the Hudson Bay and coalesced with the eastward-advancing Cordilleran ice sheet west of Calgary in the Big Hill Springs area. Deposition of glacial drift (till) and subsequent glacio-fluvial erosion modified the pre-existing land surface (Yoon 1977; Park File).

During glacial retreat about 10-12, 000 years ago, meltwater from the decaying ice margin to the north and east flowed into local topographic lows and deeply eroded the till blanket and soft underlying sandstone bedrock to create the Big Hill and Big Hill Springs coulees (Moran 1986; Alberta Recreation and Parks 1988).² The massive erosional process exposed underground aquifers, creating springs at a number of sites (Alberta Recreation and Parks 1988).

Today, the Park's complex and dramatic landscape can be considered composed of four primary elements:

- Till-blanketed Uplands. Moran (1986) describes this area as part of a larger unit of Spy Hill Drift (pebble-loam till) over the Porcupine Hills Formation sandstone. Found along upper south bank of the Big Hill Springs coulee, and extending to the south, the

² "Coulee" is used synonymously and interchangeably with "ravine" and "valley" in most instances here. According to Caboue (1996) it's a western Canadian term for a steep-sided prairie valley, but also refers to a broad-bottomed one of *glacial meltwater origin*, or to v-shaped gullies caused by more recent erosion.

moderately sloping, undulating benchland is about 1265m AMSL. Soils found at benchland sites were fine to medium-textured (clayey), mainly weak-profiled Calcareous Black Chernozems, and often gravelly-cobbly.³

- **Glaciofluvial Terrace.** The north bench above Big Hill Springs coulee is a glaciofluvial terrace about 30 meters below the morainal benchlands of Section 32 to the north.⁴ Of till parent material like the surrounding benchlands, soils found were fine to medium textured (clayey), mainly weak-profiled Calcareous Black Chernozems, and often gravelly-cobbly.⁵ Orthic Black Chernozems may be found in cultivated sites (tame pastures).
- **Coulee Walls.** The banks range from about 65m high in Big Hill valley to 15m high near the upper end of Big Hill Springs coulee, and range from 35-80% slope.⁶ Moran (1986) describes them as eroded bedrock outcrops of the non-marine Porcupine Hills formation, although at Big Hill Springs they are largely mantled with residuum, colluvium, or thin unconsolidated till veneers (Turchenek and Fawcett 1994). The dramatic, near-vertical bedrock face on the east bank of Big Hill Creek coulee towers just above – and outside - the Park's east boundary. Other smaller bedrock exposures are found in the Big Hill Springs ravine, mostly too small to map or indiscernible on airphotos. Slope failures (slumps and creep) are evident at many steeper sites, attesting to the generally weak bank stability afforded by the bedrock and till mantles in the area.

Significant ecological differences occur between the north and south-facing slopes in the highly incised Big Hill Springs Creek ravine. Slope, aspect, angle, and slope position are key determinants. Soils found on the heavily vegetated north-facing slope included primarily medium-textured (clay loams) and weakly developed or eroded Calcareous Black Chernozems with some Orthic Eutric Brunisols. On the south-facing bank, soil development appears minimal due to gradient, soil drainage, and insolation. Medium to coarse-textured Orthic Regosols were found on steep middle and upper slopes, and medium-textured Calcareous Dark Brown Chernozems on shallower, more vegetated lower slopes. In the Big Hill coulee, with moderately vegetated banks oriented east and west, soils on lower and mid-slopes tend towards Chernozemic (e.g. weakly-developed or eroded Orthic Blacks) in mainly fine to medium-textured colluvium.

³ Soils nomenclature and terminology is according to the Canadian System of Soil Classification – 2nd Edition; by Agriculture Canada, Pub. No. 1646 (1987).

⁴ An identical - and intersecting - feature is found above the east boundary of the Park in Big Hill Coulee. Bedrock control (i.e. fault-lines) likely accounts for the remarkable linearity of their outer risers or scarps.

⁵ Turchenek and Fawcett (1994) describe a landscape unit with *glaciolacustrine* veneer and some poorly-drained soils (Dunvargan-Maycroft) along the back of the terrace, extending slightly into the Park at points near the escarpment along the stepped northern boundary.

⁶ Lamoureux (1983) states that Big Hill coulee is the most deeply-incised glacial meltwater channel in the region, and is recorded as an Environmentally Significant Area for this and other biophysical reasons.

- **Valley Bottoms.** The floor of the Big Hill coulee is about 1200m AMSL, with relief nearly level at 2.5% towards the east (Big Hill Creek) and south. Moran (1986) and Turchenek and Fawcett (1994) classify this area as a silty-clayey fluvial floodplain. Although MacMillan (1987) calls the parent material here “recent lacustrine” in a depressional area, the specific physical characteristics remain undisputed.⁷

Coarse fluvial deposits from Big Hill Springs Creek overlie most of the Big Hill coulee within the Park boundary. A classic raised alluvial fan has formed, deflecting the normally meandering Big Hill Creek with comparatively erosion-resistant materials, over time, against the east coulee wall.⁸ Fluvial gravels in this feature are concentrated upslope (underlying the vicinity of the parking lot), and finer sediments downslope, towards the creek. Fans have formed at the bases of other nearby ravines and draws to the south, down the valley, and influenced the course of Big Hill Creek at those sites.

The forested Big Hill Springs ravine bottom, downcut in the old spillway, rises more steeply, about 5%, to the northwest and has numerous relict channels, terraces, and tufa deposits (Komex 1998). Soil types found in depressional sites like ravine floors reflect the effects of relatively fine-textured fluvial parent materials, low gradient, moist water regime, and typically lush vegetation. Medium-textured colluvium (at the edges) and alluvium (in the floodplain) are found in both valleys. Primary alluvial soils are Humic and Cumulic Regosols, sometimes gleyed, and Calcareous or Orthic Black Chernozems at raised and grassier sites.

Elevation within the Park boundary ranges from 1198 meters (AMSL) at the banks of Big Hill Creek, to 1272 meters on the highest benchlands above Big Hill Springs ravine.

2.7 Vegetation

The Foothills Parkland Natural Subregion generally forms a narrow transitional band between the grasslands of the Foothills Fescue Subregion and the forests of the Montane Subregion (within the Chernozemic soil regime). Desiccation by wind and low precipitation appear to be the main factors determining the extent of this Subregion.

Affected by topography and climate, there is a rapid transition from grassland with groves, to forest with grassy parks, to closed deciduous forest (Achuff 1994). This northern and largest segment of the Foothills Parkland is called the Black Diamond Upland ecodistrict, and characterized vegetatively as cleared land and shrub (Strong 1995). Strong (1992) and others contend that European settlement of the Parkland Natural Region reduced the normal frequency of fire, allowing former grasslands to be

⁷ Cauboue *et al* (1996) defines the sequential events of lacustrine deposition followed by partial reworking by fluvial processes as “fluviolacustrine”, appropriate where Big Hill Creek has meandered through.

⁸ Airphoto interpretation suggests that, under historical flow and deposition regimes, Big Hill Springs Creek should gradually migrate westwards, along the roadway, and that the confluence with Big Hill Creek will move northwards around the expanding fan. No changes in streamcourse were apparent between the 1985 and 1997 airphotos.

encroached by aspen and deciduous shrubs from the foothills. Today, little native grassland exists, having been encroached or replaced by crops and tame forage species.

Despite the extensive agriculture in the area, Big Hill coulee remains integral to a linear system of diverse green space environments extending from the Bow Valley to Highway 2 (which includes other coulees, valleys, and an abandoned railway line). Lamoureux (1983) recommends it as an “Environmental Priority Area”, and AEP (1997) identifies it as an extensive Provincial Environmentally Significant Area (ESA).

This vicinity near Calgary has recorded some of the most northerly occurrences of *Lupinus* spp. (lupines), *Danthonia* spp. (oat grass), and *Festuca idahoensis* (bluebunch fescue) (Achuff 1994).

The Park, at some sites used heavily by visitors prior to 1979, has undergone significant artificial revegetation with native and introduced species. Lin (1980) describes landscaping, sodding, seeding, tree planting (aspen and white spruce), shrub planting (“golden” willow and dogwood), maintenance, and other reclamation / rehabilitation activities during 1979-80. A large grove of *Caragana arborescens* (common caragana) and some *Syringa villosa* (late lilac) remains at the reclaimed site of a former park warden’s cabin. AEP (1994) notes the occasional presence of honeysuckle cultivars, and two widespread tame forage species, *Medicago sativa* (alfalfa) and *Phleum pratense* (timothy), probably spread from nearby pastures by winds and cattle.

2.8 Wildlife

An extensive list of animal species, primarily avifauna, observed in the Park can be found in Wallis and Wershler (1972).

Spalding (1980) states that the Richardson’s ground squirrel (also inventoried by Wallis and Wershler) is the most frequently seen mammal in the area, and that mink have also been observed. According to both references, common birds include the Savannah sparrow, Cedar Waxwing, Western Meadowlark, and Common Yellowthroat. Other noteworthy birds in the vicinity of the Park are the Great Blue Heron, Cooper’s Hawk, Prairie Falcon and Ruffed Grouse.⁹ Of these, the Cooper’s Hawk is considered rarest (Semenchuk 1992). Recent sightings of 132 bird species by members of the Calgary Field Naturalist’s Society in and near the Park are listed in Appendix 2. At least 21 of the species were apparently breeding in the area (evidenced by nests, fledglings, breeding pairs, and courtship and distraction displays). Two species listed are endangered: Piping Plover and Peregrine Falcon. Some mammal, reptile, and insect observations are also listed in Appendix 2. Beaver activity (new dam and lodge), not mentioned in reference materials, was observed just beyond the southeast corner during field checks in May, 1998. The dam backed-up Big Hill Creek in the Park.

⁹ Draw sites (small, shallow ravines) on the southwest side of the Park, were reported in 1974 to have up to three grouse lecks (AEP-Park File).

Lamoureux (1983) suggests Rainbow trout or Rainbow-Cutthroat hybrids may be present in Big Hill Creek as far upstream as the Park, since populations have been noted less than a mile to the south in wider reaches.

The Parkland has experienced significant habitat “fragmentation” from agricultural expansion. This has especially impacted forest-dwelling birds (AEP 1997).

3.0 METHODS

3.1 Field Survey

Initial stratification of the study area and selection of field sample sites was performed on 1:5000 panchromatic aerial photos taken on May 1, 1985 (i.e. leaf-off). Airphoto interpretation delineated ecological units that were distinctive by their geomorphic and vegetation patterns. Representative field sites were selected to identify and document, on standard field plot forms, the range of landform, soil, and vegetation conditions across the study area.

Field survey was carried out in early July 1997 (with some follow-up fieldwork in late September) according to methods and guidelines in the Ecological Land Survey Site Description Manual (CAPAF 1994). A total of 29 diverse sites throughout Big Hill Springs Provincial Park were visited on foot and documented.

A study of the Park’s hydrogeological characteristics and features, including the springs and creek was conducted at the same time under contract by Komex International Inc. of Calgary as a special component of the overall biophysical inventory. The study report is referenced as Komex, 1998.

Plant taxonomy follows Moss (1983) with common and scientific names referencing the Alberta Plants and Fungi Master Species List and Species Group Checklists (AEP 1993). Soil and landform terminology and classification used in this study are from CAPAF (1994), Cauboue *et al.* (1996), and the Canadian System of Soil Classification (CSCC) by the Agriculture Canada Expert Committee on Soil Survey (1987).

3.2 Data Analysis

A list of 126 species of vascular plants (excludes mosses and lichens) was compiled from field sampling, and is included as Appendix 1. The list is generally consistent with those in AEP (1994) and Wallis and Wershler (1972). Differences could be attributed to the number and variety of sites visited as well as to plant phenology during each survey.

These vegetation data were analysed and grouped into community types according to the “tabular method” described in Chapter 9 of Mueller-Dombois and Ellenberg (1974). This

method of floristic classification emphasizes dominance, constancy, and diagnostic or differentiating species in determining plant associations.

3.3 Ecological Land Classification and Mapping

The ELC is a hierarchical landscape classification and mapping system that subdivides the land surface into units with similar environmental / biophysical components. The map units that result from this process are characterized by distinctive recurring patterns of surficial material, landform, soil and vegetation. It serves, by means of data *interpretation* and analysis, to describe *typical* conditions and to generalize or simplify the landscape into units that may receive the same management.

New aerial photography of Big Hill Springs Provincial Park was acquired in June and July 1997 at scales of 1:5000 and 1:10 000. Subsequently, the ELC polygons were transferred from the 1985 photos to the new ones, with minor changes according to differences in vegetative cover evident on the 1997 coverage. Terrain, soils and vegetation data from the field sample sites were synthesized and used to develop the ELC scheme and to refine the polygon boundaries.

Map unit descriptions appearing on the ecosite (ELC) map (Map 4) represent the Ecosite level of the hierarchical classification system. This structure embraces the over-arching biophysical characteristics documented for the Natural Regions and Subregions (Achuff 1994), for their more local Ecodistricts (Strong 1995) and from airphoto and field data.

3.4 Ecosite Legend

Individual ecosite classes are described in spreadsheet format in Appendix 3. The study area lies within the Foothills Parkland Natural Subregion (#14), and more specifically the Black Diamond Upland (**BLA**) ecodistrict. Parent materials, the prime parameter used in delineating ecosites, are coded as follows:

- A Anthropogenic: Human-made or human-modified materials or features.
- C Colluvium: Unconsolidated materials moved by gravity; often at the base of a slope. Slopewash deposits or veneers are included.
- X (in association with C: "CX") Residuum: The product of *in situ* decomposition (e.g. weathering) of rock.
- F Fluvial: Related to streamflow and its associated erosional / depositional processes and features.
- FL Fluvio-Lacustrine: Lacustrine (or glaciolacustrine) deposits that have been partially re-worked by fluvial processes.
- GF Glacio-Fluvial: Pertaining to meltwater streams from wasting glacier ice and especially to the deposits and landforms produced by the streams.
- M Morainal (till): Loosely-consolidated rock debris formerly on or in a glacier, that accumulated in place as the surface ice was removed by ablation.

Numbers following the parent material identifier, such as 1,2,3, etc. represent specific combinations of landscape characteristics (slope steepness, drainage, soil subgroups, biota) occurring on the parent materials that more precisely describe the ecosite. Decimal numbers used in the Colluvial ecosite 2 show sub-categories primarily related to the composition and slope position of extensive and highly-varied Mixed Shrub communities in the Park.¹⁰

The slash ("/") marks used in the Vegetation Community descriptions separate vertical strata (e.g. canopy / shrubs / ground cover), and dashes ("-") are between dominant and sub-dominant (or in some cases co-dominant) species.

Slope classes, and soil subgroups and textures follow the Canadian System of Soil Classification. The seven soil drainage classes follow the CanSIS Manual for Describing Soils in the Field (Land Resource Research Institute 1983), and are reproduced in CAPAF (1994).

4.0 RESULTS

4.1 Ecological Land Classification

Map 4 shows a delineation of ecological units in Big Hill Springs Provincial Park distinguished primarily by geomorphic parameters (site, drainage, soils, etc.) and secondly by vegetation communities.

4.2 Vegetation Community Types

In most respects, the vegetative species and communities of the study area conform to those documented for the Foothills Parkland: *Agropyron* spp. (wheat grass) – *Poa* spp. (bluegrass) – *Danthonia parryi* (Parry's oat grass) communities dominate grasslands, with a large diversity of forb and grass species. *Populus tremuloides* (aspen) is generally dominant in the upland forests, with *Populus balsamifera* (balsam poplar) and *Picea glauca* (white spruce) occurring on shadier and/or moister sites. Distinctive in the area, willow groveland dominated by *Salix bebbiana* (Bebb's willow) (Achuff 1994) occurs in a variety of environments at Big Hill Springs ranging from medium-textured fluvial and fluvio-lacustrine material on imperfectly to poorly-drained sites as found in the coulee bottoms, to drier till-blanketed upland sites.

¹⁰ At this level of mapping, a *full* ecosite code would incorporate the Natural Subregion and ecodistrict identifiers (e.g. 14BLA.C2.1). An abbreviated version (e.g. C2.1) is shown on the ELC legend and map as all the ecosites here are in the same ecodistrict.

Field data from the 29 sites were classified into 25 different vegetation community types (4 sites were essentially duplicates), which can be grouped into generic categories of two forest types, two shrubland types, and three grassland types.

Vegetation community types are integral to the character of an ecosite. Although community types and landform units have typical associations, they, as with individual species, range from those occupying landscape-specific habitat to those common across a variety of environments. This is reflected in the Community Type descriptions that follow, and in a species-site list, Appendix 1.

The slash ("/") mark shown in the community type nomenclature separates vertical strata (e.g. canopy / shrubs / ground cover), and dashes ("-") are between dominant and sub-dominant (or in some cases co-dominant) species, by stratum. More typical conditions (e.g. in a range of slopes, moisture regimes, or set of soil subgroups) are named first in the community description tables. Proportions of a community's cover types can vary significantly between sites depending on local conditions.

Aspen Woodland

Three aspen community types were sampled in this study, representing the range that may be found mainly on the morainal uplands or colluvial coulee walls in the south part of the Park. Smaller and isolated stands were planted at the picnic sites, and can also be found on levees and banks along the lower reach of Big Hill Springs Creek. Shrubby understories, reflecting varying moisture and nutrient regimes, consist primarily of cow parsnip, prickly rose, with occasional snowberry.

4.2.1 Aspen / Cow parsnip

Below the Big Hill Springs coulee crest, in a mesic transition zone between the morainal aspen groves and the white spruce and balsam poplar-dominated units lower down, an aspen / cow parsnip community has established. About 5% white spruce are present in the overstory and understory, signifying the onset of a spruce-dominated climax community. Soil moisture may be retained (i.e. infiltration impeded) due to a shallow depth to bedrock.

Big Hill Springs Provincial Park Community Type: Aspen / Cow parsnip	
Sites: #23	Ecosites: C3

Moisture regime: mesic		Nutrient regime: permesotrophic	
Soil drainage: moderate		Soil subgroups: O.BL	
Slopes: 31-45%		Slope position, Aspect: Crest-Upper, NE	
Predominant / key species:			
Scientific name		Common Name	% Cover
Trees:	<i>Populus tremuloides</i> <i>Picea glauca</i>	aspen white spruce	30 5
Shrubs:	<i>Ribes oxycanthoides</i>	northern gooseberry	5
Forbs:	<i>Heracleum lanatum</i> <i>Viola canadensis</i>	cow parsnip western Canada violet	70 15
Gram- inoids:	<i>Elymus innovatus</i> <i>Poa pratensis</i>	hairy wild rye Kentucky bluegrass	15 15

4.2.2 Aspen / Prickly rose

Mature to old seral stands of aspen occupy areas adjacent to the aspen / cow parsnip communities. They occur on better-drained soils along the south side of the Park in morainal and colluvial parent materials. No significant understory has developed, and a variety of shrubs, mainly prickly rose and scattered white spruce, dominate this stratum.

Big Hill Springs Provincial Park Community Type: Aspen / Prickly rose			
Sites: #6, 22		Ecosites: C3, M2	
Moisture regime: mesic		Nutrient regime: mesotrophic	
Soil drainage: moderate - well		Soil subgroups: CA.BL	
Slopes: 10-45%		Slope position, Aspect: Mid, NE	
Predominant / key species:			
Scientific name		Common Name	% Cover
Trees:	<i>Populus tremuloides</i>	aspen	45-60
Shrubs:	<i>Rosa acicularis</i> <i>Ribes oxycanthoides</i> <i>Rubus idaeus</i> <i>Symphoricarpos albus</i>	prickly rose northern gooseberry wild red raspberry snowberry	20-35 5-10 5-10 3-8
Forbs:	<i>Viola canadensis</i> <i>Pyrola asarifolia</i>	western Canada violet common pink wintergreen	10-25 1-3

Gram- inoids:	<i>Elymus innovatus</i> <i>Poa pratensis</i>	hairy wild rye Kentucky bluegrass	1-15 5-10
------------------	---	--------------------------------------	--------------

4.2.3 Aspen / Snowberry-Prickly rose

Evidence of brush encroachment is found on grassy sites near the southwest edge of the Park above Big Hill Springs coulee. Here the aspen is open, short, and somewhat gnarly suggesting environmental stresses caused by insolation, moisture deficits, and competition with brush species. Other nearby aspen communities on till exhibit similar soil characteristics (horizons, thicknesses, textures) but here we have a typical grassland soil. Snowberry, an increaser, is most common in dry habitats, and its berries are toxic.

Big Hill Springs Provincial Park Community Type: Aspen / Snowberry – Prickly rose			
Sites: #20		Ecosites: M2	
Moisture regime: submesic		Nutrient regime: mesotrophic	
Soil drainage: moderate		Soil subgroups: CA.BL	
Slopes: 2-5%		Slope position, Aspect: Upper, SE	
Predominant / key species:			
Scientific name		Common Name	% Cover
Trees:	<i>Populus tremuloides</i>	aspen	20
Shrubs:	<i>Symphoricarpos albus</i> <i>Rosa acicularis</i> <i>Ribes oxycanthoides</i>	snowberry prickly rose northern gooseberry	30 25 5
Forbs:	<i>Heracleum lanatum</i> <i>Thalictrum venulosum</i>	cow parsnip veiny meadow rue	2 2
Gram- inoids:	<i>Poa pratensis</i> <i>Elymus innovatus</i>	Kentucky bluegrass hairy wild rye	20 3

Balsam Poplar or Balsam Poplar – White Spruce Woodland

Balsam poplar is associated with typically moister habitats than is aspen, such as depressions and fluvial terraces and floodplains. In the study area, ideal environments are found along Big Hill Springs Creek and the north-facing coulee wall above it. White spruce often shares the habitat, tending in many sites to succeed balsam poplar with a shade-tolerant climax community. The 3 communities described below are at differing successional stages, evidenced by the proportions of balsam and spruce.

4.2.4 Balsam poplar / Cow parsnip

This community is found along the Big Hill Springs Creek floodplain and coulee floor, from the picnic site west to mid-ravine, where white spruce become mixed in progressively shadier sites. Much of the area has a hummocky or terraced microrelief (created by fluvial erosion and deposition over time) creating a myriad of moist depressions and drier knolls. The balsam poplar favour the comparatively dry sites here, growing to over 20m, and the cow parsnip tend to the depressions. The Chernozem soil near the mouth of the coulee suggests this area was once grassed, perhaps used as pasture, and encroached by balsam poplar with a change in land use. No significant spruce understory was found at the sample site, however a few young aspen were noted. The *Viola*, *Geranium*, and *Delphinium* are common to moist woods.

Big Hill Springs Provincial Park Community Type: Balsam poplar / Cow parsnip			
Sites: #24		Ecosites: F5	
Moisture regime: mesic		Nutrient regime: permesotrophic	
Soil drainage: well		Soil subgroups: O.BL	
Slopes: 2-5%		Slope position, Aspect: Depression, SE	
Predominant / key species:			
Scientific name		Common Name	% Cover
Trees:	<i>Populus balsamifera</i>	balsam poplar	35
Shrubs:	<i>Populus balsamifera</i> <i>Ribes oxycanthoides</i>	balsam poplar northern gooseberry	5 5
Forbs:	<i>Heracleum lanatum</i> <i>Viola canadensis</i> <i>Delphinium glaucum</i> <i>Geranium richardsonii</i>	cow parsnip western Canada violet tall larkspur wild white geranium	15 8 5 5
Gram- inoids:	<i>Poa pratensis</i> <i>Bromus inermis</i>	Kentucky bluegrass awnless brome	10 8

4.2.5 Balsam poplar – White spruce / Willow / Horsetail

This mixedwood community is found associated with a variety of relatively moist sites: (1) in the shady riparian zone on the coulee floor where overbank flooding of Big Hill Springs Creek and saturated soil is common, (2) sites further upstream and closer to the creek than the Balsam / Cow parsnip community described previously, and (3) north-

facing coulee banks in the northwest parcel. Spruce may be expected to replace the seral Balsam poplar in drier sites. Species data below are from Site #13; #19 is similar.

Big Hill Springs Provincial Park Community Type: Balsam poplar - White spruce / Willow / Horsetail			
Sites: #13,19		Ecosites: F5, C4	
Moisture regime: subhydryc - mesic		Nutrient regime: permesotrophic	
Soil drainage: very poor - well		Soil subgroups: R.HG, CA.BL	
Slopes: 0-70%		Slope position, Aspect: Depression, lower-mid; N	
Predominant / key species:			
Scientific name		Common Name	% Cover
Trees:	<i>Populus balsamifera</i>	balsam poplar	10
	<i>Picea glauca</i>	white spruce	10
Shrubs:	<i>Salix bebbiana</i>	beaked willow	25
	<i>Populus balsamifera</i>	balsam poplar	5
	<i>Ribes oxycanthoides</i>	northern gooseberry	3
Forbs:	<i>Equisetum arvense</i>	common horsetail	20
	<i>Aster conspicuus</i>	showy aster	5
	<i>Epilobium angustifolium</i>	common fireweed	3
	<i>Veronica americana</i>	American brooklime	2
Gram-inoids:	<i>Carex</i> spp.	sedge	15
	<i>Poa pratensis</i>	Kentucky bluegrass	5
	<i>Catabrosa aquatica</i>	brook grass	3

4.2.6 White Spruce – Balsam poplar / Hairy wild rye

This community type is found at mid- to lower-slopes of the north-facing wall of Big Hill Springs ravine. The proportion of white spruce in the community is greater to the west up the ravine, and up-slope. The balsam poplar component is typical of lower slopes, while aspen would be found instead on drier uplands. It exemplifies a mature climax forest community for the study area.

Big Hill Springs Provincial Park Community Type: White spruce – Balsam poplar / Hairy wild rye	
Sites: #7	Ecosites: C5

Moisture regime: submesic		Nutrient regime: mesotrophic	
Soil drainage: well		Soil subgroups: CA.BL, O.EB	
Slopes: 46-70%		Slope position, Aspect: Mid, NE	
Predominant / key species:			
Scientific name		Common Name	% Cover
Trees:	<i>Picea glauca</i> <i>Populus balsamifera</i> <i>Populus tremuloides</i>	white spruce balsam poplar aspen	35 10-15 * 10
Shrubs:	<i>Picea glauca</i> <i>Sheperdia canadensis</i> <i>Ribes oxycanthoides</i> <i>Rosa acicularis</i> <i>Symphoricarpos albus</i> <i>Rubus idaeus</i>	white spruce buffalo berry northern gooseberry prickly rose snowberry wild red raspberry	3 7 7 6 5 5
Forbs:	<i>Aster conspicuus</i> <i>Viola canadensis</i>	showy aster western Canada violet	7 5
Gram-inoids:	<i>Elymus innovatus</i> <i>Poa pratensis</i>	hairy wild rye Kentucky bluegrass	25 10

* 10% denotes canopy; 15% denotes understory

Shrubland – Willow dominated

Willow (*Salix* spp.) occurs in a wide range of landscapes or ecological units such as woods, thickets, riverbanks, wetlands, and disturbed sites. They provide browse and bedding for herbivores. At Big Hill Springs, a recurring species is *Salix bebbiana* (beaked willow) encountered in wetland as well as upland sites, and in poor to rich nutrient regimes.¹¹ Willow is usually found here with secondary shrub species in areas once grassland and under a grazing regime, establishing first in drainages before spreading.

4.2.7 Willow – Cinquefoil / Bluegrass

This wetland community of 1-2m high willows straddles the lower reach of Big Hill Springs Creek in Big Hill coulee, between the road at the Park's north boundary and Big Hill Creek. It has formed on the outer fringe of the alluvial fan feature where creek water maintains a high soil moisture level. Aspens, some planted during landscaping, occupy

¹¹ At some sites, willow was recorded to the genus level only. Other willow species have been observed in the Park, mainly in wet areas: *S. pseudomonticola* (false mountain willow), *S. planifolia* (flat-leaved willow), *S. lucida* (shining willow), *S. maccalliana* (velvet-fruited willow), and *S. lutea* (yellow willow) (AEP 1994).

drier knolls and levees on the upstream edge of the unit. Gleyed medium-textured soil, with a high carbonate content indicates moist and nutrient-rich conditions. The cinquefoil, about .5m in height, is most dense in openings between the willows and around the edge of the community.

Big Hill Springs Provincial Park Community Type: Willow – Cinquefoil / Bluegrass			
Sites: #2		Ecosites: F2	
Moisture regime: subhydic		Nutrient regime: eutrophic	
Soil drainage: imperfect		Soil subgroups: GLCU.HR	
Slopes: 0-.5%		Slope position, Aspect: Toe-Depression, N/A	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Salix bebbiana</i>	beaked willow	30
	<i>Potentilla fruticosa</i>	shrubby cinquefoil	20
	<i>Ribes oxycanthoides</i>	northern gooseberry	15
Forbs:	<i>Achillea millefolium</i>	common yarrow	5
	<i>Mentha arvensis</i>	wild mint	5
	<i>Geum aleppicum</i>	yellow avens	3
	<i>Senecio</i> sp.	groundsel	3
Gram- inoids:	<i>Poa pratensis</i>	Kentucky bluegrass	40
	<i>Juncus balticus</i>	wire rush	20
	<i>Poa palustrus</i>	fowl bluegrass	10

4.2.8 Willow / Bluegrass – Rough fescue

This seral upland community is found on inclined to undulating till mantle above Big Hill Springs coulee to the south. The sample site, at the southwest corner in the northwest parcel of the Park, is affected by cattle grazing and brush encroachment. This vegetation association is a variant of other upland willow-dominated communities (like 4.2.9) along the southwest edge of the Park, and extending southward. Fescue is a decreaser under grazing and may be replaced by the more invasive bluegrass, especially on abused upland sites (Tannas 1997).

Big Hill Springs Provincial Park Community Type: Willow / Bluegrass – Rough fescue	
Sites: #18	Ecosites: M1
Moisture regime: mesic	Nutrient regime: permesotrophic
Soil drainage: moderate	Soil subgroups: O.BL
Slopes:10-30%	Slope position, Aspect: Mid, NE
Predominant / key species:	

Scientific name		Common Name	% Cover
Shrubs:	<i>Salix bebbiana</i>	beaked willow	30
	<i>Shepherdia canadensis</i>	Canada buffalo berry	10
	<i>Potentilla fruticosa</i>	shrubby cinquefoil	5
	<i>Betula glandulosa</i>	bog birch	5
	<i>Populus balsamifera</i>	balsam poplar	5
Forbs:	<i>Fragaria virginiana</i>	wild strawberry	8
Gram-inoids:	<i>Festuca scabrella</i>	rough fescue	15
	<i>Poa pratensis</i>	Kentucky bluegrass	15
	<i>Danthonia parryi</i>	Parry oat grass	5

4.2.9 Willow - Prickly rose / Hairy wild rye - Bluegrass

Beyond the aspen communities above Big Hill Springs coulee, willow-dominated groves spread up-slope to the south, just beyond the Park's irregular southwest boundary on undulating and gullied morainal uplands. Fingers of this community protrude into the Park through breaks between the aspen stands. Scattered white spruce are found on knolls and rises between gullies. The dominant shrub species found in the community favour relatively rich nutrient regimes as are found in calcareous Black Chernozems present here. Aspen could encroach in some sites of this community to form climatic climax stands, however succession would be slow in many parts due to dry conditions, and the willow would remain an edaphic climax.

Big Hill Springs Provincial Park Community Type: Willow - Prickly rose / Hairy wild rye - Bluegrass			
Sites: #21		Ecosites: M1	
Moisture regime: submesic		Nutrient regime: permesotrophic	
Soil drainage: moderate		Soil subgroups: CA.BL	
Slopes:10-30%		Slope position, Aspect: Mid, NE	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Salix bebbiana</i>	beaked willow	45 *
	<i>Rosa acicularis</i>	prickly rose	40
	<i>Symphoricarpos albus</i>	snowberry	20
	<i>Amelanchier alnifolia</i>	saskatoon	5
Forbs:	<i>Rubus idaeus</i>	wild red raspberry	5
	<i>Taraxacum officinale</i>	common dandelion	5
Gram-inoids:	<i>Elymus innovatus</i>	hairy wild rye	25
	<i>Poa pratensis</i>	Kentucky bluegrass	20

* Willow occupies the "tall shrub" stratum, while the others occupy the "low shrub".

Mixed Shrublands

Communities that fall into this category are as diverse as they are extensive at Big Hill Springs. Sites sampled were found to be dominated mainly by snowberry (buckbrush) or silver berry (wolf-willow) and to a lesser extent by prickly rose. Significantly, they occur mainly on hillsides within and near the Park perhaps due to lighter historical grazing regimes and/or sufficient moisture on the mid-to-lower slopes.

4.2.10 Snowberry – Prickly rose / Bluegrass

This community is found on a relatively moist lower hillslope of Big Hill coulee south the mouth of Big Hill Springs ravine (upslope from the parking lot). Shallow depth to bedrock and the lower slope position produce seeps that support lush growth at certain microsites.¹² While sparse, the moisture and nutrient-loving Cow parsnip was encountered on lower slopes. Scattered 1-2m willow sit over this knee-high shrub community giving it an open appearance on the hillside. Aspen are beginning to encroach this seral community from the fringes at many sites. The snowberry and rose are increasers; saskatoon is a decreaser, but a valuable component of native range (Tannas 1997).

Big Hill Springs Provincial Park Community Type: Snowberry - Prickly rose / Bluegrass			
Sites: #5		Ecosites: C2.1	
Moisture regime: subhygric		Nutrient regime: permesotrophic	
Soil drainage: moderate		Soil subgroups: O.BL	
Slopes: 31-45%		Slope position, Aspect: Lower, E	
Predominant / key species:			
Scientific name		Common Name	% Cover
Trees:			
Shrubs:	<i>Symphoricarpos albus</i>	snowberry	25
	<i>Rosa acicularis</i>	prickly rose	30
	<i>Amelanchier alnifolia</i>	saskatoon	15
	<i>Salix bebbiana</i>	beaked willow	5
	<i>Ribes oxycanthoides</i>	northern gooseberry	4
Forbs:	<i>Thalictrum venulosum</i>	veiny meadow rue	4
	<i>Anemone canadensis</i>	Canada anemone	3
Gram-inoids:	<i>Poa pratensis</i>	Kentucky bluegrass	35
	<i>Bromus inermis</i>	awnless brome	5

¹² This is approximately where ANHIC records indicate an observation of the relatively rare – and tracked – fern species *Pellaea glabella* (smooth cliff brake) in 1994. Its preference for habitat among limestone rocks (Moss 1992) would be met by the local tufa deposits.

4.2.11 Snowberry / Brome - Bluegrass

On the floor of Big Hill Springs ravine, the open and gently-sloping area supporting this community appears to have been an old clearing, perhaps pasture, now undergoing significant brush encroachment both from the creekbanks below and the aspen above. The sample site is on one of several fluvial terraces in a sequence extending the length of the coulee. Nutrient-rich, moist, with a deep alluvial topsoil, it provides ideal habitat for the invasive brome grass.

Big Hill Springs Provincial Park Community Type: Snowberry / Brome - Bluegrass			
Sites: #14		Ecosites: F6	
Moisture regime: subhygric		Nutrient regime: eutrophic	
Soil drainage: well		Soil subgroups: R.BL	
Slopes: 6-15%		Slope position, Aspect: Toe, E	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Symphoricarpos albus</i>	snowberry	15
	<i>Salix bebbiana</i>	beaked willow	5
	<i>Ribes oxycanthoides</i>	northern gooseberry	2
Forbs:	<i>Heracleum lanatum</i>	cow parsnip	5
	<i>Taraxacum officinale</i>	common dandelion	5
	<i>Urtica dioica</i>	common nettle	3
Gram-inoids:	<i>Bromus inermis</i>	awnless brome *	50
	<i>Poa pratensis</i>	Kentucky bluegrass	15
	<i>Carex sprengeii</i>	Sprengell's sedge	5

* Colloquially called "smooth" brome.

4.2.12 Snowberry / Bluegrass

This community is found on open lower slopes of both coulees in relatively arid environments (due to insolation, slope, and well-drained soil) on colluvial parent materials. The wide ratio range of snowberry to bluegrass in these sites (i.e. 15:80 to 75:10) suggests the effects of local site characteristics (e.g. seeps, aspect, microrelief, grazing regime) on competition between these two increaser species. The Dark Brown Chernozem was found on a predominantly shrubby site with a strong south aspect.

Big Hill Springs Provincial Park Community Type: Snowberry / Bluegrass	
Sites: #25,29	Ecosites: C2.1

Moisture regime: submesic		Nutrient regime: mesotrophic	
Soil drainage: well		Soil subgroups: CA.BL, CA.DB	
Slopes: 16-30%		Slope position, Aspect: Lower, S-W	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Symphoricarpos albus</i> <i>Ribes oxycanthoides</i> <i>Amelanchier alnifolia</i>	snowberry northern gooseberry saskatoon	15-75 0-3 0-3
Forbs:	<i>Geranium viscosissimum</i>	sticky purple geranium	1-5
Gram- inoids:	<i>Poa pratensis</i>	Kentucky bluegrass	10-80

4.2.13 Silverberry – Choke cherry / Bluegrass

Dense groves of silverberry are found on the lower colluvium-blanketed slopes of the east wall of Big Hill Creek coulee along the Park's east boundary, and in a few sites on the south-facing bank of Big Hill Springs ravine. These sites represent typical silverberry habitat. It's an increaser, resistant to moderate browsing and forms rather extensive colonies as climax grasses decline. Choke cherry is a decreaser, not as highly competitive (Tannas 1997). This indicates a silverberry / bluegrass edaphic climax community.

Big Hill Springs Provincial Park Community Type: Silverberry – Choke cherry / Bluegrass			
Sites: #9		Ecosites: C2.2	
Moisture regime: submesic		Nutrient regime: mesotrophic	
Soil drainage: moderate		Soil subgroups: O.R	
Slopes: 31-70%		Slope position, Aspect: Mid-Lower, W-S	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Eleagnus commutata</i>	silverberry	60 *
	<i>Prunus virginiana</i>	choke cherry	30
	<i>Symphoricarpos albus</i>	snowberry	25
	<i>Amelanchier alnifolia</i>	saskatoon	5

Forbs:	<i>Aster conspicuus</i> <i>Thalictrum venulosum</i>	showy aster veiny meadow rue	5 4
Gram-inoids:	<i>Poa pratensis</i> <i>Agropyron dasystachyum</i>	Kentucky bluegrass northern wheatgrass	20 5

* Silverberry occupies the “tall shrub” stratum, while the others occupy the “low shrub”.

4.2.14 Prickly rose - Silverberry / Bluegrass

This community occupies the middle hillside zone between the road and the colluvium-resuduum-outcrops unit (CX1) on the west crest of Big Hill Creek coulee. Buffalo berry is present at the upper fringes. There are signs of soil creep along this bank, perhaps exacerbated by cattle grazing-trampling over time, however it currently appears stable due to vegetative cover. The shrubs, all increasers, comprise a maturing edaphic climax community due to lack of moisture (caused by strong slopes, well-drained soil, and insolation) and poor soil development.

Big Hill Springs Provincial Park Community Type: Silverberry – Choke cherry / Bluegrass			
Sites: #28		Ecosites: C2.3	
Moisture regime: submesic		Nutrient regime: permesotrophic	
Soil drainage: well		Soil subgroups: CA.BL	
Slopes: 31-45%		Slope position, Aspect: Mid, E	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Rosa acicularis</i>	prickly rose	40
	<i>Eleagnus commutata</i>	silverberry	25
	<i>Symphoricarpos albus</i>	snowberry	5
Forbs:	<i>Apocynum androsaemifolium</i>	spreading dogbane	3
	<i>Geranium viscosissimum</i>	sticky purple geranium	3
Gram-inoids:	<i>Poa pratensis</i>	Kentucky bluegrass	10

4.2.15 Silverberry - Cinquefoil / Wheat grass - Sedge

This community is described using a composite of characteristics of two similar and nearby sites on the upper slopes on the east wall of Big Hill Springs coulee (above and just beyond the Park’s east boundary). These sites are just below the glaciofluvial terrace

(described in Section 2.6) on colluvial veneer over inclined weathered bedrock. The sedge, a drought-resistant increaser, sometimes found on such exposed areas and eroded slopes at high elevations, can be very effective in erosion control here due to its dense and tough root structure (Tannas 1997). Of the grasses - all increasers - June grass is native and a common grassland species. The sample sites are classed as young to mature edaphic climax communities.

Big Hill Springs Provincial Park Community Type: Silverberry – Cinquefoil / Wheat grass - Sedge			
Sites: #8,10		Ecosites: C2.2	
Moisture regime: submesic		Nutrient regime: mesotrophic	
Soil drainage: moderate		Soil subgroups: O.R, R.DB	
Slopes: 31-70%		Slope position, Aspect: Mid-Lower, W-S	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Eleagnus commutata</i>	silverberry	8-15
	<i>Amelanchier alnifolia</i>	saskatoon	5-15
	<i>Potentilla fruticosa</i>	shrubby cinquefoil	2-5
	<i>Prunus virginiana</i>	choke cherry	1
Forbs:	<i>Galium boreale</i>	northern bedstraw	3-30
	<i>Artemesia frigida</i>	pasture sagewort	3-10
Gram-inoids:	<i>Agropyron dasystachyum</i>	northern wheat grass	10-30
	<i>Poa pratensis</i>	Kentucky bluegrass	20
	<i>Carex filifolia</i>	thread-leaved sedge	15-20
	<i>Koeleria macrantha</i>	June grass	5-25

Grasslands

The four grassland communities described below represent a cross-section of essentially what remains of “native” or natural communities (opposed to tame or reclaimed sites) in an area long used for ranching. The presence and occasional dominance of introduced or non-native species like crested wheat grass and Kentucky bluegrass attest to the long-term effects of historical pasture seeding, dispersion by wind and animals, grazing regimes, and aggressive habitat competition by introduced / invader and increaser species such as brome. AEP (1997) describes that fescue grasslands, including those in the Parkland, are threatened by extensive cultivation and that any remaining areas should be considered endangered.

Grassland community types at Big Hill Springs are often quite variable by species composition, and spatially - as influenced by site-specific conditions including parent material, slope-aspect, microrelief, moisture, and land use.

4.2.16 Sand grass – Northern wheat grass – June grass

The sampled site is on an open mid-slope of the north bank of Big Hill Springs coulee, on medium to coarse soils. These factors combined with a strong south aspect contribute to the relative aridity of the site, hence its edaphic climax successional status. Although an increaser, sand grass is susceptible to trampling, with implications for human and cattle access and the site's slope stability. Northern wheatgrass, normally an increaser – and tolerant to grazing and trampling – may be a decreaser on such dry sites. June grass, often indicative of overgrazing, is an increaser (Tannas 1997).

Big Hill Springs Provincial Park Community Type: Sand grass – Northern wheat grass – June grass			
Sites: #26		Ecosites: C1	
Moisture regime: submesic-subxeric		Nutrient regime: mesotrophic	
Soil drainage: well-rapid		Soil subgroups: O.R	
Slopes: 46-70%		Slope position, Aspect: Mid, S	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Potentilla fruticosa</i>	shrubby cinquefoil	1
Forbs:	<i>Artemesia frigida</i>	pasture sagewort	2
	<i>Trifolium repens</i>	white clover	2
	<i>Phlox hoodii</i>	moss phlox	2
Gram-inoids:	<i>Calamovilfa longifolia</i>	sand grass	30
	<i>Agropyron dasystachyum</i>	northern wheatgrass	15
	<i>Koeleria macrantha</i>	June grass	15
	<i>Carex filifolia</i>	thread-leaved sedge	5

4.2.17 Northern wheat grass – Bluebunch fescue

This community shares the same ecological unit as the previous one, about 60m to the east along the north bank of Big Hill Springs coulee. Significant differences between sites are the shallower slope and finer-textured soil here facilitating Chernozemic soil development. Both the predominant grass species are increasers and can withstand grazing and trampling. The scattered shrubby cinquefoil on site may be seen as invasive and troublesome in range-lands, although they contribute to erosion control. This was the only sample site where the northern green bog-orchid (*Habenaria hyperborea*) was encountered – unusual in a dry grassland habitat, and uncommon in the Park (AEP 1994).

Big Hill Springs Provincial Park Community Type: Northern wheat grass – Bluebunch fescue	
Sites: #27	Ecosites: C1

Moisture regime: submesic-mesic		Nutrient regime: permesotrophic	
Soil drainage: well		Soil subgroups: CA,BL	
Slopes: 10-15%		Slope position, Aspect: Mid, S	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Potentilla fruticosa</i>	shrubby cinquefoil	1
Forbs:	<i>Thermopsis rhombofolia</i>	golden bean	5
	<i>Smiliacina stellata</i>	star-flowered Solomon's seal	5
	<i>Solidago missouriensis</i>	low goldenrod	3
Gram-inoids:	<i>Agropyron dasystachyum</i>	northern wheat grass	30
	<i>Festuca idahoensis</i>	bluebunch fescue	20
	<i>Koeleria macrantha</i>	June grass	5
	<i>Poa pratensis</i>	Kentucky bluegrass	5
	<i>Stipa comata</i>	needle-and-thread	5

4.2.18 Parry oat grass – Kentucky bluegrass

At the far northwest corner of the main Park parcel, the grassy landscape grades from a gently-undulating glaciofluvial terrace to upper colluvial slopes of Big Hill Springs coulee. The Chernozemic Black topsoil is deeper and fragment-free on the benchland portion. Light grazing use is evident. Both dominant grasses are increasers, although bluegrass is probably the more invasive and persistent, often replacing native species on abused upland sites in the region (Tannas 1997). Similarly, shrubby cinquefoil is invasive, adapted to a variety of soils, and can degrade good-quality grasslands. Successional status is a mature edaphic climax.

Big Hill Springs Provincial Park Community Type: Parry oat grass – Kentucky bluegrass			
Sites: #12		Ecosites: GF1, C1	
Moisture regime: mesic-submesic		Nutrient regime: mesotrophic	
Soil drainage: moderate		Soil subgroups: CA,BL	
Slopes: .5-30%		Slope position, Aspect: Crest-Upper, S	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Potentilla fruticosa</i>	shrubby cinquefoil	2
	<i>Rosa acicularis</i>	prickly rose	1
Forbs:	<i>Thalictrum venulosum</i>	veiny meadow rue	8
	<i>Sisyrinchium montanum</i>	common blue-eyed grass	3
	<i>Anemone patens</i>	prairie crocus	2

Gram-inoids:	<i>Danthonia parryi</i>	Parry oat grass	25
	<i>Poa pratensis</i>	Kentucky bluegrass	25
	<i>Koeleria macrantha</i>	June grass	5
	<i>Stipa viridula</i>	green needle grass	5

4.2.19 Kentucky bluegrass – Rough fescue

The sample site is located in the northwest parcel of the Park, low on the glaciofluvial terrace on the north side of the upper Big Hill Springs coulee. The soil is medium-textured Black Chernozemic. The site, long a pasture, apparently remains under a light grazing regime. Bluegrass is an increaser, often replacing natives on abused upland sites. Fescues are usually hardy, drought-resistant and suited to intensive grazing. Rough fescue is a valuable climax species, but slow to establish and is often a decreaser in many places (Tannas 1997). There is a rich variety of forbs represented in this mature edaphic climax community, although in small proportions.

Big Hill Springs Provincial Park Community Type: Kentucky bluegrass – Rough fescue			
Sites: #16		Ecosites: GF1	
Moisture regime: subxeric		Nutrient regime: mesotrophic	
Soil drainage: well		Soil subgroups: O.BL	
Slopes: 0-9%		Slope position, Aspect: Upper-Mid, S	
Predominant / key species:			
Scientific name		Common Name	% Cover
Forbs:	<i>Thermopsis rhombifolia</i>	golden bean	5
	<i>Artemesia frigida</i>	pasture sagewort	5
	<i>Achillea millefolium</i>	common yarrow	2
Gram-inoids:	<i>Poa pratensis</i>	Kentucky bluegrass	50
	<i>Festuca scabrella</i>	rough fescue	25
	<i>Agropyron dasystachyum</i>	northern wheatgrass	5
	<i>Koeleria macrantha</i>	June grass	2
	<i>Stipa viridula</i>	green needle grass	3

Modified Grasslands

Some sites in the Park, as discussed in previous sections, have experienced cultivation: seeding and / or reclamation and the introduction of non-native grass species. These correspond to former tame pastures recently acquired by the Park (i.e. the northwest parcel), and reclaimed facility sites in the lower part of the Park.

4.2.20 Northern wheat grass – June grass

This community was identified on the edge of the former Park headquarters facility site. Located on an elevated fluvial terrace in Big Hill Creek ravine, much of the central part was altered by the processes of site preparation, construction, landscaping, and finally, dismantling and reclamation. The peripheral vegetation appears essentially native and is synopsized in the table below.

The reclaimed portion has a heavy cover of brome grass (*Bromus* spp.) - invasive, but excellent for site stabilization (Tannas 1997), with a lesser amount of Kentucky bluegrass.¹³ There are remnant ornamental shrubs: mainly caragana, with some lilac and honeysuckle. Perhaps due to contouring and tillage at reclamation, as well its receiving slope position, this site is quite moist as evidenced by gleyed soil, lush brome growth, and balsam poplar shoots.

Big Hill Springs Provincial Park Community Type: Northern wheat grass – June grass			
Sites: #11		Ecosites: F6	
Moisture regime: subhygric		Nutrient regime: permesotrophic	
Soil drainage: imperfect		Soil subgroups: O.BL, O.HG	
Slopes: .5-2.5%		Slope position, Aspect: Toe-Depression, N/A	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Potentilla fruticosa</i>	shrubby cinquefoil	2
	<i>Juniperus horizontalis</i>	creeping juniper	2
Forbs:	<i>Thermopsis rhombifolia</i>	golden bean	3
	<i>Artemisia frigida</i>	pasture sagewort	5
	<i>Galium boreale</i>	northern bedstraw	20
Gram-inoids:	<i>Agropyron dasystachyum</i>	northern wheatgrass	25
	<i>Poa pratensis</i>	Kentucky bluegrass	15
	<i>Koeleria macrantha</i>	June grass	15
	<i>Danthonia parryi</i>	Parry oat grass	5

4.2.21 Awnless brome – Timothy

On an ecosite resembling the previous one, this grassland community is situated on the alluvial fan near the mouth of Big Hill Springs Creek where the old campground was. Lin (1980) describes the area as being covered with exotic grasses and weeds before reclaiming with a seed mix of crested wheat grass, Kentucky bluegrass, creeping red

¹³ Lin (1980) documents that the site was reclaimed with a seed mix of 30% creeping red fescue (*Festuca* sp.), 45% crested wheat grass (*Agropyron pectiniforme*), and 5% white clover (*Trifolium repens*). This composition is not evident today.

fescue and white clover (as well as with some transplanted aspen and spruce). Since then, brome and timothy have invaded and dominated due to the moist and rich soils. The site is criss-crossed with old vehicle trails and tracks.

Big Hill Springs Provincial Park Community Type: Awnless brome - Timothy			
Sites: #4		Ecosites: F4	
Moisture regime: subhygric		Nutrient regime: permesotrophic	
Soil drainage: well		Soil subgroups: CA.BL	
Slopes: 2-5%		Slope position, Aspect: Toe-Depression, E	
Predominant / key species:			
Scientific name		Common Name	% Cover
Forbs:	<i>Ranunculus acris</i>	tall buttercup	2
	<i>Trifolium pratense</i>	red clover	2
Gram-inoids:	<i>Bromus inermis</i>	awnless brome *	60
	<i>Phleum pratense</i>	timothy	30
	<i>Carex</i> spp.	sedge	5

* Colloquially called "smooth" brome.

4.2.22 Kentucky bluegrass - Awnless brome - Timothy

This community is found in a tame pasture near the centre of the northwest parcel, on the broad glaciofluvial terrace. Each grass species here is an increaser, with brome and timothy the more invasive, aggressive and persistent - and relatively more abundant in the moist depressions (shallow relict channels) at this site.

Big Hill Springs Provincial Park Community Type: Kentucky bluegrass – Awnless brome - Timothy			
Sites: #15		Ecosites: GF1	
Moisture regime: subhygric		Nutrient regime: mesotrophic	
Soil drainage: moderate		Soil subgroups: O.BL	
Slopes: 2-5%		Slope position, Aspect: Mid, SE	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Rosa acicularis</i>	prickly rose	1
Forbs:	<i>Thermopsis rhombifolia</i>	golden bean	5
	<i>Artemisia ludoviciana</i>	prairie sagewort	3
	<i>Geranium viscosissimum</i>	sticky purple geranium	3

Gram- inoids:	<i>Poa pratensis</i> <i>Bromus inermis</i> <i>Phleum pratense</i>	Kentucky bluegrass awnless brome timothy	80 5 5
------------------	---	--	--------------

Grasslands -Wet meadows

This category of vegetation community at Big Hill Springs is characterized by a site's proximity to water and high water table, and by a preponderance of hygrophytic graminoids such as sedges (*Carex* spp.) and rushes (*Juncus* spp.).

4.2.23 Tall buttercup / Wire rush

Above Big Hill Creek on the outer fringe of the alluvial fan, this meadow community has experienced cattle grazing (a tussocky surface) as well as effects from park visitors over time (since it surrounds the former campground). The soil, moist – as manifested by gleying in the profile, and rich in nutrients (calcareous) supports a lush vegetative cover. The introduced and invasive buttercup is common in such depressional and low meadows; and the rush, also an increaser is resistant to trampling and spreads when other species decline (Tannas 1997).

Big Hill Springs Provincial Park Community Type: Tall buttercup / Wire rush			
Sites: #3		Ecosites: F3	
Moisture regime: subhygric		Nutrient regime: permesotrophic	
Soil drainage: imperfect		Soil subgroups: GLCU.R	
Slopes: 2-5%		Slope position, Aspect: Toe-Depression, E	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Potentilla fruticosa</i>	shrubby cinquefoil	2
Forbs:	<i>Ranunculus acris</i>	tall buttercup	15
	<i>Potentilla anserina</i>	silverweed	5
	<i>Achillea millefolium</i>	common yarrow	4
Gram- inoids:	<i>Juncus balticus</i>	wire rush	25
	<i>Phleum pratense</i>	timothy	20
	<i>Poa pratensis</i>	Kentucky bluegrass	15

4.2.24 Water sedge

This riparian community is established on the floodplain of Big Hill Creek south of the confluence with Big Hill Springs Creek. Its thick rich, silty topsoil is subjected to periodic flooding and deposition. Trampling by cattle has degraded the soft creekbank with the formation of tussocks and small pits. Scattered willow in clumps and thickets surround the site upslope on the moister fringes of the alluvial fan. The sedge, normally a decreaser and somewhat susceptible to grazing pressure, can be good for soil-binding and erosion control in wet sites.

Big Hill Springs Provincial Park Community Type: Water sedge			
Sites: #1		Ecosites: F1	
Moisture regime: subhydric		Nutrient regime: eutrophic	
Soil drainage: imperfect - well		Soil subgroups: O.HR	
Slopes: 0-2.5 %		Slope position, Aspect: Depression-level, N/A	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Salix</i> spp.	willow	1
Forbs:	<i>Mentha arvensis</i>	wild mint	5
	<i>Potentilla anserina</i>	silverweed	3
Gram-inoids:	<i>Carex aquatilis</i>	water sedge	40
	<i>Poa pratensis</i>	Kentucky bluegrass	10

4.2.25 Kentucky bluegrass - Fowl bluegrass – Bebb's sedge

This edaphic climax community occurs adjacent to the uppermost spring discharge site on the floor of Big Hill Springs coulee, in the northwest parcel. Nominally tame pasture (as is the area just above it to the north), the site is significantly affected by year-round springwater, as well as by concentrated surface runoff in the spring and during rainy periods. Fowl bluegrass, unlike Kentucky bluegrass, is a decreaser and wetland species (Tannas 1997). This was the only site where the northern willow-herb (*Epilobium ciliatum*), generally common especially in wet and “springy places”, was encountered.

Big Hill Springs Provincial Park Community Type: Kentucky bluegrass – Fowl bluegrass – Bebb's sedge	
Sites: #17	Ecosites: F5

Moisture regime: subhygric		Nutrient regime: permesotrophic	
Soil drainage: well		Soil subgroups: O.HR	
Slopes: 2-5%		Slope position, Aspect: Depression, NW	
Predominant / key species:			
Scientific name		Common Name	% Cover
Shrubs:	<i>Salix</i> spp.	willow	5
Forbs:	<i>Epilobium ciliatum</i>	northern willow-herb	10
	<i>Veronica americana</i>	American brooklime	5
Gram-inoids:	<i>Poa pratensis</i>	Kentucky bluegrass	25
	<i>Poa palustrus</i>	fowl bluegrass	10
	<i>Phleum pratense</i>	timothy	5
	<i>Carex bebbii</i>	Bebb's sedge	5
	<i>Catabrosa aquatica</i>	brook grass	5

4.3 Significant Plant Species

Of the 126 vascular plant species encountered during this study, no rare or uncommon species (as tracked by ANHIC)¹⁴ were observed. Six species deemed Uncommon by AEP (1994) were observed, and are indicated in the Vegetation Species List, Appendix 1.¹⁵

The ANHIC database contains a number of observations of 2 rare species in or close to Big Hill Springs Provincial Park:

<i>Pellaea glabella</i>	smooth cliff brake (fern)
<i>Ranunculus aquatilis</i>	large-leaved white water crowfoot

4.4 Significant Vegetation Cover Changes

Changes in vegetative cover represent the most apparent, extensive, and rapid of biophysical changes in the Park.

A natural process of vegetation succession in a conservation area such as a park would normally be viewed as positive, and management practices would seek to foster this ecological evolution. Succession from the region's historical extensive grass cover to parkland in the absence of extensive grazing (and *with* fire suppression) is almost

¹⁴ Alberta Natural Heritage Information Centre database on rare plants. Administered by Alberta Environmental Protection, Natural Resources Service – Parks.

¹⁵ Wallis and Wershler recorded 5 of these 6 species too. It may suggest they aren't that "uncommon".

Big Hill Springs Provincial Park Changes in Vegetation Cover 1972-1997

Sec. 29 Tp. 26 Rg. 03 W5Mer.

Wallis & Werahler, 1972	Community Types, 1997
1 Grassland (Dry)	to Mixed Shrubs
2 Grassland (Dry)	to Aspen Woodland
3 Heavily Grazed Grassland	to Willow Shrubs
4 Heavily Grazed Grassland	to Mixed Shrubs
5 Heavily Grazed Grassland	to Balsam Poplar Woodland
6 Grassland (Wet)	to Balsam Poplar Woodland
7 Grassland (Wet)	to Aspen Woodland
8 Grassland (Wet)	to Mixed Shrubs
9 Immature Poplar Meadow	to Mature Willow Shrubs - Aspen
10 Silverberry	to Aspen Woodland
11 Willow - Riparian	to Balsam Poplar - White Spruce Woodland
12 Low Shrubs	to Aspen Woodland
13 Tall Willow	to Aspen Woodland
14 Tall Willow	to Mixed Shrubs
15 Aspen Woodland	to White Spruce - Balsam Poplar Woodland
16 Trampled and Mowed Grass	to Wet Meadow
17 Rock Outcrops	to Mixed Shrubs
18 Rock Outcrops	to Balsam Poplar - White Spruce Woodland

In the main part of the Park, areas and polygons with NO numbers have not experienced significant changes in vegetation cover.

inevitable. As succession proceeds, community structure becomes more stratified, biomass and organic matter accumulate, and nutrient cycling becomes more internal in the system. Food webs become more complex, species diversity increases to a point and net community production approaches equilibrium (Smith 1996).

The transition of vegetative cover from pioneer and seral communities such as grasses, shrubs and poplar to maturing climax communities like spruce stands is evident in many areas of the Park. Expansion of the spruce and poplar communities into old shrubland in the south of the Park and in Big Hill Springs ravine is the most significant in an areal sense. In Big Hill coulee, willow and mixed-shrub groves have matured and spread into former grassland pasture.

Apparent changes between current vegetation community distribution and that mapped by Wallis and Wershler 25 years ago (possibly using June, 1970 airphoto data) are indicated on Map 3. However, differences in approaches to ground cover characterization, classification, and mapping - rather than real change - may bear on the validity or extent of some of the "Changes" shown on Map 3 (e.g. 16, 17, 18). These differences would preclude a detailed comparison and analysis.

5.0 ISSUES AND RECOMMENDATIONS

The following are considered to be some issues that bear on the Park's ecological integrity, conservation of its natural history features, and enjoyment by the public. Most are self-evident and may merit further detailed study in order to develop suitable management approaches.

5.1 Land Use

As long as grazing is approved for certain Park lands, degradation, or at least suspension of some natural processes (e.g. vegetation succession, streamcourse development) may be expected. A *range*-specific inventory and assessment should be performed if cattle grazing is likely to continue within the Park, so ecologically-based guidelines can be established.

Impacts from visitor use, as well, were apparent at some sites and might be mitigated by facility / infrastructure enhancements, and signage appealing to a conservation ethic.

5.2 Fences and Boundaries

Wire fences were established in a pre- or early Park setting to control cattle grazing. Some fences in and around the Park appear to have become obsolete for that purpose, yet remain in-situ, in varying states of repair and decay. As well, most fencelines have little

correspondence with Park boundaries. Evidence of the Park's true boundaries on the ground is very limited and (with the fencing and unofficial warning signs) may unnecessarily deter visitors from experiencing its far corners.

5.3 Sensitive Sites

No ecosites or vegetation communities in the Park were analysed specifically as to their biological / ecological or physical vulnerability. Komex (1998) comments on the susceptibility of the tufa formations to erosion and wear, and on the effects of cattle and streambank trampling on water quality. If there are any significantly sensitive sites at Big Hill Springs, most will likely be in the relatively soft streamcourse, riparian, or wet areas.

5.4 Bank Stability

The steeper coulee walls are weak and physically sensitive to weathering, erosion, and trampling; therefore susceptible to mass movements including outright collapse. Numerous creep, slump, and rockfall deposits were observed. Access to the top and bottom of these sites should be restricted.

5.5 Rare Plants

It was suggested (pers. comm., Joyce Gould – Keith Ainsley) during the course of fieldwork, that the best prospects of finding any rare or uncommon plant species, probably **mosses**, would be in the ecologically unique spring or stream sites, and by a bryophyte specialist. As well, locations of the two rare plants already identified in the Park should be sought-out. A follow-up rare plant survey may be warranted.

5.5 Brush Encroachment

Brush encroachment involves *viable* rangelands being invaded by woody vegetation - brush and young trees, to a point where forage production is lost or unavailable for grazing. It's evident from comparing historical and contemporary airphotos and maps that this is occurring at many sites in the Park, particularly above the south bank of Big Hill Springs coulee, and in Big Hill coulee. Assessments should be made with a view to preserving grassland communities through pro-active measures, or to "let Nature take its course".

5.6 Adjacent Lands

It's apparent that with the two parcels of the Park not contiguous, and that because some of the Park's natural history features are minimally or not wholly enclosed by Park boundaries, that full and effective management of these features is constrained. Section 2.7 describes the link the Park plays in a regional green space network.

Big Hill Springs Provincial Park is a very small "island" - only 38 hectares, hardly ecologically viable on its own. It's vulnerable, and perhaps irreversibly affected by land use practices and industrial activities on nearby properties: farms, ranches, acreages, gravel operations, etc. Much of the local watershed drains through the Park, with implications for water quality and supply.

6.0 REFERENCES

- Alberta Environmental Protection (AEP). 1993. Alberta Plants and Fungi – Master Species List and Species Group Checklists. Pub. No. Ref. 75.
- Alberta Environmental Protection (AEP) – Parks. 1994. Big Hill Springs Provincial Park: Plants. Alberta's Watchable Wildlife Checklist Series. Government of Alberta.
- Alberta Environmental Protection (AEP). 1996. Big Hill Springs Provincial Park Management Plan. Natural Resources Service, Parks Management.
- Alberta Environmental Protection (AEP). 1997. The Parkland Natural Region of Alberta. Natural Resources Service, Recreation and Protected Areas Division, Natural Heritage Planning and Evaluation Branch.
- Alberta Environmental Protection (AEP). nd. Park File: Big Hill Springs Provincial Park. Natural Resource Service, Parks Management.
- Alberta Recreation and Parks. ca. 1988. Big Hill Springs Provincial Park. Curriculum guide. Copy held at University of Alberta, Edmonton.
- Blogorodow, P. *et al.* 1976. Big Hill Springs Provincial Park: Mini Master Plan. Regional Park Planning, Planning and Design Branch, Alberta Recreation, Parks and Wildlife.
- Borneuf, D. 1983. Springs of Alberta. Alberta Research Council.
- Canada-Alberta Partnership Agreement in Forestry (CAPAF). 1994. Ecological Land Survey Site Description Manual. Produced for Canadian Forest Service and Alberta Land and Forest Services; Alberta Environmental Protection
- Cauboue, M., Strong, W.L., Archambault, L. and Sims, R.A. 1996. Terminology of Ecological Land Classification in Canada. Natural Resources Canada, and Canadian Forest Service – Quebec. Information Report LAU-X-114E.
- Curcio, Michele P. 1967, 1970. Water Study and Report (for Big Hill Springs Provincial Park). Department of Lands and Forests, Parks Division. Government of Alberta.
- Houseknecht, Stephan J. 1984. Natural History Study of Mineral and Thermal Springs in Canada. TERA Environmental Consultants Ltd., Vancouver; Prepared for Parks Canada.
- Komex International Ltd. 1998. Hydrogeological Characterization of Big Hill Springs Provincial Park. Prepared for Alberta Environmental Protection, Southern East Slopes Region, Calgary.

Lamoureux, R.J. 1983. Environmentally Significant Areas of the Calgary Region. Prepared for the Calgary Regional Planning Commission

Lin, E., *et al.* 1980. Horticultural Management Plan for Big Hill Springs Provincial Park. Alberta Recreation and Parks – Parks Division.

MacMillan, R.A. 1987. Soil Survey of the Calgary Urban Perimeter. Alberta Soil Survey Report No. 45. Alberta Research Council, Edmonton.

Moss, E.H. 1983. Flora of Alberta. 2nd. Ed. Revised by J. Packer. University of Toronto Press.

Mueller-Dombois, D. and Ellenberg, H. 1974. Aims and Methods of Vegetation Ecology. John Wiley & Sons, Inc.

Semenchuk, G. (ed.). 1992. The Atlas of Breeding Birds of Alberta. Federation of Alberta Naturalists, Edmonton, Alberta.

Smith, R.L. 1996. Ecology and Field Biology - 5th Edition. Harper Collins College Publishers.

Spalding, D.A.E. (Ed.). 1980. A Nature Guide to Alberta. Provincial Museum of Alberta publication No.5. Hurtig Publishers, Edmonton.

Strong, W.L. 1992. Ecoregions and Ecodistricts of Alberta, Volume 1. Prepared by Ecological Land Surveys Ltd. for Alberta Forestry, Lands and Wildlife; Land Information Services Division; Edmonton

Strong, W.L. 1995. Ecodistricts of Alberta: Summary of Biophysical Attributes. Prepared for Alberta Environmental Protection; Resource Data Division; Edmonton

Tannas, K. ca. 1997. Common Plants of the Western Rangelands, Vols. I and II. Curriculum and Instructional Development Services, Lethbridge Community College.

Turchenek, L.W. and Fawcett, M.D. 1994. Soil Survey of the Municipal District of Rocky View No. 44, Alberta (excluding the Calgary Urban Perimeter). Alberta Soil Survey Report No.53. Alberta Research Council, Edmonton.

Wallis, C.A. and Wershler, C.R. 1972. An Ecological Survey of Big Hill Springs Provincial Park. Provincial Parks Planning Branch, Department of Lands and Forests, Alberta.

Appendix 1

BIG HILL SPRINGS PROVINCIAL PARK: VEGETATION SPECIES

The 126 species listed below were observed and recorded during field sampling in July 1997. This list does not constitute a complete or exhaustive list of species to be found in the Park, but would supplement other inventories.

Habitat Codes: A - Aspen Woodlands; G – Grasslands; Gm – Modified Grasslands; M – Moist Meadows; P – Balsam poplar or Balsam poplar–White spruce dominated Woodlands; R – Riparian areas; Sm – Mixed Shrublands; Sw – Willow-dominated Shrublands.
The habitat code *sequence* shown below for each species is not significant. U – Uncommon (AEP 1994).

<u>Species</u>	<u>Common Name</u>	<u>Habitat</u>	<u>Site #</u>
<i>Achillea millefolium</i>	common yarrow	M / Sm / Sw / G / Gm / A	2,3,5,9,11,15,16,17,18,20,21,25,28
<i>Actaea rubra</i>	red and white baneberry	A	6,22,23
<i>Agropyron dasystachyum</i>	northern wheatgrass	Sm / G	8,9,10,11,16,26,27,29
<i>Allium textile</i>	prairie onion	Sm / G	10,11,26,27,29
<i>Amelanchier alnifolia</i>	saskatoon	Sm / A / Sw	5,8,9,10,18,20,22,28,29
<i>Anemone canadensis</i>	Canada anemone	M / Sm / A	2,3,5,6,14
<i>Anemone multifida</i>	cut-leafed anemone	Sm / G	8,11,16
<i>Anemone patens</i>	prairie crocus	G / Gm	12,15,27
<i>Antennaria parvifolia</i>	small-leaved everlasting	G	11,27
<i>Apocynum androsaemifolium</i>	spreading dogbane	Sm	8,9,10,28
<i>Arctostaphylos uva-ursi</i>	common bearberry	Sw	18
<i>Artemisia frigida</i>	pasture sagewort	Sm / G	8,9,10,11,16,25,26,27,28
<i>Artemisia ludoviciana</i>	prairie sagewort	Sm / Gm	5,9,15,29
<i>Aster alpinus</i>	alpine aster	Sm	25
<i>Aster ciliolatus</i>	Lindley's aster	A / P / Sm / Sw / G	6,7,9,12,18,21,23,24
<i>Aster conspicuus</i>	showy aster	A / P / Sm / Sw	6,7,9,13,20,21,24
<i>Aster laevis</i>	smooth aster	Sm	28
<i>Aster spp.</i>	aster	Sm	5
<i>Astragalus dasyglottis</i>	purple milk vetch	G / Sw / A	11,12,18,20
<i>Astragalus tennellus</i>	loose-flowered milk vetch	Sm / G	8,10,26
<i>Betula glandulosa</i>	bog birch	Sw	18
<i>Bromus inermis</i>	awnless brome	A / G / Sm / Gm / P	4,5,8,9,10,14,15,16,24,25,28
<i>Calamagrostis canadensis</i>	bluejoint	A / G	6
<i>Calamovilfa longifolia</i>	sand grass	G	26
<i>Campanula rotundifolia</i>	harebell	Sm / G	8,27
<i>Caragana arborens</i>	common caragana	Gm	11
<i>Carex aquatilis</i>	water sedge	R	1
<i>Carex bebbii</i>	Bebb's sedge	M	17
<i>Carex filifolia</i>	thread-leaved sedge	Sm / G	8,10,26
<i>Carex spp.</i>	sedge	P / R	13
<i>Carex sprengelii</i>	Sprengel's sedge	A / P / Sm	7,14,23
<i>Catabrosa aquatica</i>	brook grass	P / R / M U	13,17
<i>Cerastium arvense</i>	mouse-ear chickweed	R / M / Sm / A / G / Sw	1,2,3,5,12,18,20
<i>Cirsium arvense</i>	Canada thistle	R / M / Sm / Sw / P	1,2,5,13,18,19,29
<i>Cirsium undulatum</i>	wavy-leaved thistle	Sm / G U	10,26,27,29
<i>Danthonia parryi</i>	Parry oat grass	G / Sw / Sm	11,12,18,25
<i>Delphinium glaucum</i>	tall larkspur	P	7,13,23,24
<i>Disporum trachycarpum</i>	fairy bells	A / P	6,7,22
<i>Dodecatheon pulchellum</i>	shooting star	M	3
<i>Eleagnus commutata</i>	silverberry	Sm	8,9,10,28
<i>Elymus innovatus</i>	hairy wild rye	A / P / Sm / Sw	6,7,9,20,21,22,23
<i>Epilobium angustifolium</i>	common fireweed	R / M / Sm / A / P	1,2,3,5,6,7,13,23,24
<i>Epilobium ciliatum</i>	northern willowherb	M	17
<i>Equisetum arvense</i>	common horsetail	R / M / P	1,2,13,17,24
<i>Erigeron flagellaris</i>	creeping fleabane	Sm / G	8,10,11,26
<i>Erigeron glabellus</i>	smooth fleabane	G	27
<i>Erigonum flavum</i>	yellow umbrella-plant	G	27
<i>Festuca idahoensis</i>	bluebunch fescue	G	27
<i>Festuca scabrella</i>	rough fescue	Sm / G / Gm / Sw / A	8,10,15,16,18
<i>Fragaria virginiana</i>	wild strawberry	M / A / Sm / Sw / P / G	3,6,7,9,12,18,20,21,22,24
<i>Gaillardia aristata</i>	gaillardia	Sm / G U	8,27
<i>Galium boreale</i>	northern bedstraw	M / Sm / Sw / A / P / G	3,5,6,7,8,9,10,11,16,18,20,21,22,24,25,26,27,28,29
<i>Gaura coccinea</i>	scarlet butterflyweed	Sm	10,29
<i>Geranium richardsonii</i>	wild white geranium	A / P	22,23,24
<i>Geranium viscosissimum</i>	sticky purple geranium	Sm / Gm	5,14,15,25,28,29
<i>Geum aleppicum</i>	yellow avens	R / M / P / A U	1,2,7,13,17,22,23

<i>Geum triflorum</i>	three-flowered avens	Sm / Sw / G / Gm / A / P	5,12,15,20,24
<i>Habenaria hyperborea</i>	nor. green bog orchid	G U	27
<i>Hackelia floribunda</i>	large-flowered stickseed	Sm	29
<i>Hackelia jessicae</i>	Jessica's stickseed	Sm / A	14,23
<i>Hedysarum alpinum</i>	alpine hedysarum	Sm / M / G / Gm / Sw / A	5,8,9,11,12,15,16,17,18,20,25,28
<i>Heracleum lanatum</i>	cow parsnip	Sm / A / P	5,6,7,13,14,20,23,24
<i>Hipperis vulgaris</i>	common mare's tail	R / stream	1
<i>Juncus balticus</i>	wire rush	M / Sm / P	2,3,5,13,17
<i>Juniperus horizontalis</i>	creeping juniper	Sm / G	8,10,11
<i>Koeleria macrantha</i>	June grass	Sm / G	8,10,11,16,26,27
<i>Lathyrus ochroleucus</i>	cream-colored vetchling	A / P / Sw	6,7,20,21,22,23
<i>Lilium philadelphicum</i>	western wood lily	Sw	21
<i>Lithospermum incisum</i>	narrow-leaved puccoon	Sm	8
<i>Lithospermum rudemale</i>	wooly gromwell	Sm / A / G	9,11,16,20,25,27,28
<i>Lonicera dioica</i>	twining honeysuckle	A / P	6,7
<i>Lonicera involucrata</i>	bracted honeysuckle	A / P	6,7
<i>Mentha arvensis</i>	wild mint	R / M / P	1,2,17,24
<i>Menyanthes trifoliata</i>	buck bean	R / M	1,2
<i>Mnium</i> sp.		R / M	1,2
<i>Monarda fistulosa</i>	wild bergamont	Sm	9,25,28
<i>Nasturtium officinale</i>	water cress	P / R	13
<i>Osmorhiza depauperata</i>	spreading sweet cicely	A	23
<i>Oxytropis splendens</i>	showy locoweed	Sm	8
<i>Phleum pratense</i>	timothy	M / Sm / P / Gm	3,4,9,13,15,17
<i>Phlox hoodii</i>	moss phlox	G	26
<i>Picea glauca</i>	white spruce	A / P / Sw	6,7,13,21,23,24
<i>Poa palustris</i>	fowl bluegrass	M / P	2,13,17
<i>Poa pratensis</i>	Kentucky bluegrass	M / Sw / Sm / A / P / G / Gm / R	1,2,3,5,6,7,8,9,11,12-18,20,21,22,23,24,25,27,28,29
<i>Populus balsamifera</i>	balsam poplar	Sm / A / P / Sw	5,6,7,13,24
<i>Populus tremuloides</i>	aspen	A / P / Sw	6,7,13,20,21,22,23
<i>Potentilla hippiana</i>	wooly cinquefoil	G	11
<i>Potentilla anserina</i>	silverweed	R / M	1,2,3,17
<i>Potentilla fruticosa</i>	shrubby cinquefoil	M / Sm / Sw / G	2,3,5,8,9,10,11,12,18,26,27,28
<i>Potentilla gracilis</i>	graceful cinquefoil	Gm / Sm	15,25,28
<i>Prunus virginiana</i>	choke cherry	Sm	8,9,10,28
<i>Pyrola asarifolia</i>	com. pink wintergreen	A	6,22,23
<i>Ranunculus acris</i>	tall buttercup	M / P / R	3,13,17
<i>Ribes oxycanthoides</i>	northern gooseberry	M / Sm / A / P / Sw	2,5,6,7,9,13,14,20,21,22,23,24,25,28
<i>Rosa acicularis</i>	prickly rose	Sm / A / P / G / Gm / Sw	5,6,7,9,10,12,15,18,20,21,22,23,24,25,28
<i>Rubus idaeus</i>	wild red raspberry	A / P / Sw	6,7,21,22,23,24
<i>Salix bebbiana</i>	beaked willow	M / Sw / Sm / P	2,5,13,14,18,21
<i>Salix</i> spp.	willow	R / P / G / M	1,6,12,17,24
<i>Scutellaria galericulata</i>	marsh skullcap	P / R	13
<i>Senecio canus</i>	prairie groundsel	M / Sm / G	3,8,10,27
<i>Senecio</i> sp.	ragwort / groundsel	M	2
<i>Senecio vulgaris</i>	common groundsel	P / R / M	13,17
<i>Shepherdia canadensis</i>	Canada buffalo berry	Sm / P / Sw / G	5,7,8,13,18,27,28
<i>Sisyrinchium montanum</i>	common blue-eyed grass	Sm / G / Gm	8,11,12,15,16,25
<i>Smilacina stellata</i>	Solomon's-seal	Sm / A / P / G / Sw	5,6,7,9,10,12,21,22,23,26,27,28
<i>Solidago missouriensis</i>	low goldenrod	G	27
<i>Solidago</i> spp.	goldenrod	G	12
<i>Spiraea betulifolia</i>	white meadowsweet	A / P	6,7
<i>Stipa columbiana</i>	Columbia needle grass	G / Sw	27,29
<i>Stipa comata</i>	needle-and-thread	G	27
<i>Stipa viridula</i>	green needle grass	Sm / G	10,12,16
<i>Symphoricarpos albus</i>	snowberry	Sm / Sw / A / P	5,6,7,9,10,14,18,20,21,22,23,25,28,29
<i>Taraxacum officinale</i>	common dandelion	R / M / A / Sm / P / Sw	1,3,6,9,13,14,17,20,21,22,23,24
<i>Thalictrum venulosum</i>	veiny meadow rue	M / Sm / A / G / Sw	3,5,6,9,12,14,18,20,28
<i>Thermopsis rhombofolia</i>	golden bean	Sm / G / Gm / M	8,10,11,12,15,16,17,27,28,29
<i>Thlaspi arvense</i>	stinkweed	Sm	14
<i>Trifolium pratense</i>	red clover	G	4,27
<i>Trifolium repens</i>	white clover	P / G / M	13,17,26
<i>Urtica dioica</i>	common nettle	A / P / M	6,7,13,14,17,22,23,24
<i>Veronica americana</i>	American brooklime	P / R-M U	13,17
<i>Vicia americana</i>	wild vetch	M / Sm / A / G / Sw	2,3,5,6,8,10,11,12,18,20,21,22,23
<i>Viola adunca</i>	early blue violet	Sm	5
<i>Viola canadensis</i>	western Canada violet	M / A / P / Sw	3,6,21,22,23,24
<i>Viola nephrophylla</i>	bog violet	R / M	1,2
<i>Zigadenus elegans</i>	white camas	M / Sm / G / Sw	2,3,8,11,12,16,18,21,27
<i>Zizia aptera</i>	heart-leaved Alexander	M / Sw / Sm / Gm	3,5,15,18,21

Appendix 2

Calgary Field Naturalist's Society Bird Record Project

Big Hill Springs Provincial Park - Bird sightings, 1987 to date

The following bird species were observed in or near the Park by members of the CFNS and recorded in the Bird Record Project database or in members' personal notes. Boldface entries indicate species confirmed or evidently breeding in the Park. Asterisks (*) indicate endangered species.

Alder Flycatcher	Eastern Kingbird	Orange-crowned Warbler
American Coot	Eastern Phoebe	Peregrine Falcon *
American Crow	European Starling	Pied-billed Grebe
American Dipper	Evening Grosbeak	Pileated Woodpecker
American Goldfinch	Gadwall	Pine Grosbeak
American Kestrel	Golden Eagle	Pine Siskin
American Redstart	Golden-crowned Kinglet	Piping Plover *
American Robin	Gray Catbird	Prairie Falcon
American Tree Sparrow	Great Blue Heron	Red Crossbill
American Widgeon	Great Horned Owl	Red-breasted Grosbeak
Bald Eagle	Greater Yellowlegs	Red-breasted Nuthatch
Bank Swallow	Green-winged Teal	Red-eyed Vireo
Barn Swallow	Hairy Woodpecker	Red-naped Sapsucker
Barrow's Goldeneye	Hermit Thrush	Red-necked Grebe
Black Tern	Hoary Redpoll	Red-Tailed Hawk
Black-billed Magpie	Hooded Merganser	Red-winged Blackbird
Black-capped Chickadee	House Sparrow	Ring-necked Duck
Blackpoll Warbler	House Wren	Rock Dove
Blue Jay	Killdeer	Rough-legged Hawk
Blue-winged Teal	Least Flycatcher	Ruby-crowned Kinglet
Bohemian Waxwing	LeConte's Sparrow	Ruffed Grouse
Boreal Chickadee	Lesser Scaup	Savannah Sparrow
Boreal Owl	Lesser Yellowlegs	Say's Phoebe
Brewer's Blackbird	Lincoln's Sparrow	Sharp-shinned Hawk
Broad-winged Hawk	Loggerhead Shrike	Sharp-tailed Grouse
Brown Creeper	Long-billed Curlew	Sharp-tailed Sparrow
Brown-headed Cowbird	MacGillivray's Warbler	Song Sparrow
Canada Goose	Mallard	Sora
Cape May Warbler	Merlin	Spotted Sandpiper
Cedar Waxwing	Mountain Bluebird	Swainson's Hawk
Chipping Sparrow	Mountain Chickadee	Swainson's Thrush
Cinnamon Teal	Mourning Warbler	Tennessee Warbler
Clay-coloured Sparrow	Northern Flicker	Tree Swallow
Cliff Swallow	Northern Goshawk	Vesper Sparrow
Common Raven	Northern Harrier	Warbling Vireo
Common Redpoll	Northern Oriole	Western Meadowlark
Common Snipe	Northern Pigmy Owl	Western Tanager
Common Yellowthroat	Northern Pintail	Western Wood-pewee
Cooper's Hawk	Northern Rough-winged Swallow	White-crowned Sparrow
Cordilleran Flycatcher	Northern Shoveler	White-throated Sparrow
Dark-eyed Junco	Northern Shrike	White-winged crossbill
Downy Woodpecker	Olive-sided Flycatcher	Willow Flycatcher

Wilson's Phalarope
Wilson's Warbler
Yellow Warbler
Yellow-bellied Sapsucker
Yellow-headed Blackbird
Yellow-rumped Warbler

Mammals sighted 1987 to date:

COMMON:

Coyote
Meadow Vole
Mule Deer
Pocket Gopher
Porcupine
Red Squirrel
Richardson's Ground Squirrel
Varying Hare (Snowshoe Hare)

LESS COMMON:

American Badger
Jumping Mouse
Long-Tailed Weasel
Moose

Reptiles:

Garter Snake

Observations provided by Steven Kassai and the
Calgary Field Naturalists Society, 1998.

Note: CFNS and members' wildlife sighting
records often have further data including exact
sites, dates, numbers, and special characteristics.

Insects sighted 1987 to date:

Butterflies:

Cabbage White
Clouded Sulphur
Common Alpine
Melissa Blue
Milbert's Tortoiseshell
Mourning Cloak
Painted Lady
Pearly Crescentspot
Prairie Ringlet
Red Admiral
Red-disked Alpine
Satyr Anglewing
Tiger Swallowtail
Western White
White Admiral
Wood Nymph

Others:

Cicada
Hawk Moth
Ichneumon Wasp
Showy Emerald Moth

Appendix 3: BIG HILL SPRINGS PROVINCIAL PARK

Ecosite	Vegetation Physiognomy	Vegetation Community Types	
A Anthropogenic	Unvegetated, OR sodded grounds with planted trees	Aspen and White spruce / Kentucky bluegrass	Upper alluvial fan
C1 Colluvial	Grasslands (including native remnants)	Northern wheat grass - Bluebunch fescue; Sand grass - Northern wheat grass - June grass	Upper south-facing and sandstone bedrock
C2.1	Mixed shrubs: Snowberry dominant; Some aspen encroachment at edges	Snowberry - Rose - Cinquefoil / Kentucky bluegrass - Brome - Northern wheat grass	Mid-to-lower coulee weathered sandstone
C2.2	Mixed shrubs: Silverberry dominant	Silverberry - Choke cherry - Rose - Cinquefoil / Sedge- Kentucky bluegrass	Mid-to-lower coulee weathered sandstone
C2.3	Mixed shrubs: Rose dominant	Rose - Silverberry - Buffaloberry / Kentucky bluegrass	Lower coulee wall weathered sandstone
C3	Aspen woodland	Aspen / Rose - Cow parsnip	North-facing, mid-slope blanket over incline
C4	Balsam poplar woodland	Balsam Poplar - White Spruce / Gooseberry	Lower north-facing sandstone bedrock
C5	White spruce-dominated woodland	White spruce - Balsam poplar - Aspen / Hairy wild rye	North-facing coulee bedrock and till. [C5]
CX1 Colluvial-Rock Residium	Unvegetated, to sparse grass and scattered shrubs	Snowberry - Rose / Kentucky bluegrass - Northern wheat grass	Upper coulee walls colluvium and residue
F1 Fluvial	Sedges with scattered willow-dominated shrub groves	Willow / Sedge	Creek channels and floodplains
F2	Willow-dominated shrubland	Willow - Cinquefoil - Aspen / Kentucky bluegrass - Rush	Lower alluvial fan and floodplains
F3	Open wet meadow	scattered Cinquefoil / Buttercup / Rush - Timothy	Alluvial fan periphery fragments. [F3]
F4	Modified (reclaimed) grassland with scattered transplanted aspen and spruce	Brome - Timothy	Upper alluvial fan.
F5	Moist riparian meadow and woodland	Balsam poplar - White spruce-Aspen / Cow parsnip - Willow - Horsetail / Fowl bluegrass - Sedge	Floodplain and bank undulating / terraced
F6	Open moist poplar forest with shrubs and grasses	scattered Aspen - Balsam poplar / Snowberry - Willow / Cow parsnip / Brome - Kentucky bluegrass	Fluvial terrace sequence tufa deposits in lower
FL1 Fluvio-Lacustrine	Wet meadow	scattered Cinquefoil / Buttercup / Rush - Timothy	Big Hill Creek coulee
GF1 Glacio-Fluvial	Grasslands and tame pastures	Oat grass - Kentucky bluegrass; Kentucky bluegrass - Rough fescue; Kentucky bluegrass - Brome - Timothy	Glacio-fluvial terrace coulee, mainly to the Fluvial (or GF) veneer
M1 Morainal	Willow-dominated shrubland	scattered White spruce / Willow - Rose - Snowberry / Hairy wild rye - Kentucky bluegrass - Rough fescue	Inclined, undulating, Big Hill Springs coulee
M2	Aspen woodland	Aspen / Rose - Snowberry	Upper north-facing rim blanket. [M2-u/V/R]

PARK ECOLOGICAL LAND CLASSIFICATION

Geomorphology [with unit codes]	Slope (%)	Soil Subgroups	Textures	Drainage Class	Comments
alluvial fan. Coarse gravels. [Ff-i]	0-2.5	N/A	N/A	N/A	Remaining visitor facilities include paved road, gravel parking lot, toilet-garbage and picnic sites
er south-facing coulee wall. Colluvial veneers over till sandstone bedrock. [Cv / M, Ri-s]	10-100	Calcareous Black, Orthic Regosol	gravelly- cobbly SiL,CL,SiC,L, LS	4	Sites 12b, 26, 27
o-lower coulee wall; colluvial blanket over inclined er sandstone. [Cv-M / Ri]	16-30	Calcareous and Orthic Black; and Calcareous Dark Brown	SiL,SiCL,L	3-4	Sites 5, 25, 29 Evidence of soil creep
o-lower coulee wall; colluvial blanket over inclined er sandstone. [Cv-M / Ri-s]	31-70	Orthic Regosol, Rego Dark Brown	rubbly - gravelly SC, SiCL, SiC,C	4	Sites 8,9,10; submesic. Evidence of soil creep
er coulee wall; colluvial veneers over inclined er sandstone. [Cv-M / Ri]	31-45	Calcareous Black	SiL,SiCL	3-4	Site 28; submesic Evidence of soil creep
er south-facing, midslope coulee wall and ledges; colluvial er over inclined bedrock. [Cv-b / Mi, Ri]	31-45	Orthic and Calcareous Black	SL,SCL,SiC,Si CL	3-4	Sites 6,23
er north-facing coulee wall; colluvial veneer over one bedrock and till. [Cv / M-Ri]	31-70	Calcareous Black	gravelly- rubbly SiCL	4-3	Site 19
er south-facing coulee wall; colluvial veneer over sandstone k and till. [Cv / Ri-s,V]	46-100	Calcareous Black, Orthic Eutric Brunisol	rubbly SiCL,SL	3-4	Site 7
er coulee walls; cliffs, sandstone outcrops; rubbly ium and residuum. [RXi-sMV]	46- >100	Orthic Regosols, non-soils	rubbly SL, SCL	3-4	Area of active weathering, erosion, creep, and rockfall
er channels and floodplains in Big Hill coulee. [Flo]	0-2.5	Orthic Humic Regosol	Si, SiCL	4-3	Site 1; trampled streambanks
er alluvial fan and streambank levees. [Ffl-t]	0-2.5	Gleyed Cumulic Humic Regosol	SiL, SCL, SiCL, CL	5	Site 2; tufa fragments in soil profile; aspen stands on creek levees
er al fan periphery; medium-textured over coarse nts. [Ffol-i]	2-5	Gleyed Cumulic Regosol	SCL,CSC	5	Site 3; tufa fragments in soil profile; extensive cattle grazing effects
er alluvial fan. [Ffl-i]	2-5	Calcareous Black	Gravelly SL,SCL,SC	3	Site 4; reclaimed campground; extensive cattle grazing effects
er plain and banks of Big Hill Springs Creek; dissected, ting / terraced. [Fou-IE]	.5-5	Calcareous Black, Rego Humic Gleysol, Orthic Humic Regosol	SCL,SC SiCL, SiC,	3-7	Sites 13, 17, 24; drier regimes on knolls or terraces, moister near creekbanks. Trampled banks at some sites
er terrace sequence above Big Hill Springs Creek; posits in lower ravine. [Ftl-i]	.5-15	Rego Black, Orthic Humic Gleysol	SiL, SiCL	3,5	Sites 11 (reclaimed), 14; includes relict channel, meander, and point bar in NW parcel used as pasture. F6 units mapped adjacently on ELC map reflect different terrace levels.
er ll Creek coulee (off the alluvial fan). [FLlo]	0-2.5	Rego Humic Gleysol, Terric Mesisol	CL, SiCL	6-7	Southeast corner of Park adjacent to Big Hill Creek
er fluvial terrace sequence above Big Hill Springs mainly to the north; gently undulating to inclined. (or GF) veneer over till / bedrock. [GFti]	0-9	Orthic and Calcareous Black	L, SiL, SiCL	3-4	Sites 12a,15,16
er d, undulating, and gullied morainial upland south of l Springs coulee; above M2. [Mbi-u-mV / R]	10-30	Calcareous and Orthic Black	gravelly- cobbly SiCL, SiC, C	4	Sites 18, 21
er north-facing rim of Big Hill Springs coulee; morainial . [Mbi-uV / R]	2-15	Calcareous Black	SiCL,SiC,C	4	Sites 20, 22. Some aspen-encroached grass & shrub sites.

Map 4

Ecological Land Classification of Big Hill Springs Provincial Park

N ½ of Sec.29 Tp.26 Rg.03 W5Mer.

ELC Units
Field Sample Sites
Park Boundary
Trails
Fences (not all are shown)

Digital Orthophoto Mosaic from June, 1997 1:5000 Airphotos.
Alberta Environmental Protection, Environmental Service, Calgary.

Ecosite	Vegetation Physiognomy	Vegetation Community Type	Geomorphology	Slope (%)	Soil Subgroups	Textures	Drainage Class	Comments
A Anthropogenic	Unvegetated, OR sodded grounds with planted trees.	Aspen and White spruce /Kentucky bluegrass	Upper alluvial fan. Coarse gravels. [Ff-i]	0-2.5	N/A	N/A	N/A	Remaining visitor facilities include paved road, gravel parking lot, toilet, garbage and picnic sites.
C1 Colluvial	Grasslands (including native remnants).	Northern wheat grass - Bluebunch fescue; Sand grass - Northern wheat grass - June grass	Upper south-facing coulee wall; colluvial veneers over till and sandstone bedrock. [Cv/M, Ri-s]	10-100	Calcareous Black, Orthic Regosol	gravelly-cobbly SiL,CL,SiC,L,LS	4	Sites 12b, 26, 27
C2.1	Mixed shrubs: Snowberry dominant. Some Aspen encroachment at edges.	Snowberry - Rose - Cinquefoil/Kentucky bluegrass - Brome - Northern wheat grass	Mid-to-lower coulee wall; colluvial blanket over inclined weathered sandstone. [Cv-M/Ri]	16-30	Calcareous and Orthic Black; and Calcareous Dark Brown	SiL,SiCL,L	3-4	Sites 5, 25, 29 Evidence of soil creep.
C2.2	Mixed shrubs: Silverberry dominant.	Silverberry - Choke cherry - Rose - Cinquefoil/Sedge - Kentucky bluegrass	Mid-to-lower coulee wall; colluvial blanket over inclined weathered sandstone. [Cv-M/Ri-s]	31-70	Orthic Regosol, Rego Dark Brown	rubby-gravelly SC,SiCL,SiC,C	4	Sites 8, 9, 10; submesic. Evidence of soil creep.
C2.3	Mixed shrubs: Rose dominant.	Rose - Silverberry - Buffaloberry /Kentucky bluegrass	Lower coulee wall; colluvial veneers over inclined weathered sandstone. [Cv-M/Ri]	31-45	Calcareous Black	SiL,SiCL	3-4	Site 28; submesic. Evidence of soil creep.
C3	Aspen woodland.	Aspen /Rose - Cow parsnip	North-facing, midlope coulee wall and ledges; colluvial blanket over inclined bedrock. [Cv-b/Mi,Ri]	31-45	Orthic and Calcareous Black	SL,SCL,SiC,SiCL	3-4	Sites 6, 23
C4	Balsam poplar woodland.	Balsam poplar - White spruce /Gooseberry	Lower north-facing coulee wall; colluvial veneer over sandstone bedrock and till. [Cv/M-Ri]	31-70	Calcareous Black	gravelly-rubby SiCL	4-3	Site 19
C5	White spruce - dominated woodland.	White spruce - Balsam poplar - Aspen /Hairy wild rye	North-facing coulee wall; colluvial veneer over sandstone bedrock and till. [Cv/Ri-s,V]	46-100	Calcareous Black, Orthic Eutric Brunisol	rubby SiCL,SL	3-4	Site 7
CX1 Colluvial-Rock Residuum	Unvegetated, to sparse grass and scattered shrubs.	Snowberry - Rose /Kentucky bluegrass - Northern wheat grass	Upper coulee walls; cliffs, sandstone outcrops; rubby colluvium and residuum. [RXi-sMV]	46->100	Orthic Regosols, non-soils	rubby SL,SCL	3-4	Area of active weathering, erosion, creep and rockfall.
F1 Fluvial	Sedges with scattered Willow - dominated shrub groves.	Willow /Sedge	Creek channels and floodplains in Big Hill coulee. [Flo]	0-2.5	Orthic Humic Regosol	Si,SiCL	4-3	Site 1; trampled streambanks.
F2	Willow - dominated shrubland.	Willow - Cinquefoil - Aspen /Kentucky bluegrass - Rush	Lower alluvial fan and streambank levees. [Ffi-t]	0-2.5	Gleyed Cumulic Humic Regosol	SiL,SCL,SiCL,CL	5	Site 2; tufa fragments in soil profile; aspen stands on creek levees.
F3	Open wet meadow.	Scattered Cinquefoil/Buttercup /Rush - Timothy	Alluvial fan periphery; medium-textured over coarse fragments. [Ffol-i]	2-5	Gleyed Cumulic Regosol	SCL,CSC	5	Site 3; tufa fragments in soil profile; extensive cattle grazing effects.
F4	Modified (reclaimed) grassland with scattered transplanted Aspen and Spruce.	Brome - Timothy	Upper alluvial fan. [Ffi-i]	2-5	Calcareous Black	gravelly SL,SCL,SC	3	Site 4; reclaimed campground; extensive cattle grazing effects.
F5	Moist riparian meadow and woodland.	Balsam Poplar - White spruce - Aspen /Cow parsnip - Willow - Horsetail/Fowl bluegrass - Sedge	Floodplain and banks of Big Hill Springs Creek; dissected, undulating /terraced. [Fou-tE]	.5-5	Calcareous Black, Rego Humic Gleysol, Orthic Humic Regosol	SCL,SC,SiCL,SiC	3-7	Sites 13, 17, 24; drier regimes on knolls or terraces, moister near creekbanks. Trampled banks at some sites.
F6	Open moist Poplar forest with shrubs and grasses.	Scattered Aspen - Balsam poplar /Snowberry - Willow /Cow parsnip /Brome - Kentucky bluegrass	Fluvial terrace sequence above Big Hill Springs Creek; tufa deposits in lower ravine. [Fti-i]	.5-15	Rego Black, Orthic Humic Gleysol	SiL,SiCL	3-5	Sites 11(reclaimed), 14; includes relict channel, meander and point bar in NW parcel used as pasture. F6 units mapped adjacently on ELC map reflect different terrace levels.
FL1 Fluvio-Lacustrine	Wet meadow.	Scattered Cinquefoil/Buttercup /Rush - Timothy	Big Hill Creek coulee (off the alluvial fan). [FLlo]	0-2.5	Rego Humic Gleysol, Terric Mesisol	CL,SiCL	6-7	Southeast corner of Park adjacent to Big Hill Creek.
GF1 Glacio-Fluvial	Grasslands and tame pastures.	Oat grass - Kentucky bluegrass; Kentucky bluegrass - Rough fescue; Kentucky bluegrass - Brome - Timothy	Glacio-fluvial terrace sequence above Big Hill Springs coulee, mainly to the north; gently undulating to inclined. Fluvial (or GF) veneer over till /bedrock. [GFti]	0-9	Orthic and Calcareous Black	L,SiL,SiCL	3-4	Sites 12a, 15, 16
M1 Morainal	Willow - dominated shrubland.	Scattered White spruce /Willow - Rose - Snowberry / Hairy wild rye - Kentucky bluegrass - Rough fescue	Inclined, undulating and gullied morainal upland south of Big Hill Springs coulee; above M2. [Mbi-u-mV/R]	10-30	Calcareous and Orthic Black	gravelly-cobbly SiCL,SiC,C	4	Sites 18, 21
M2	Aspen woodland.	Aspen /Rose - Snowberry	Upper north-facing rim of Big Hill Springs coulee; morainal blanket. [Mbi-uV/R]	2-15	Calcareous Black	SiCL,SiC,C	4	Sites 20, 22. Some aspen-encroached grass and shrub sites.

National Library of Canada
Bibliothèque nationale du Canada

3 3286 52098050 5