THE
VEGETATION OF MT. FUJI
(JAPAN)

BY
B. HAYATA, DR. SC.

F1
94
Jap
H

HARVARD UNIVERSITY

LIBRARY

OF THE

GRAY HERBARIUM

Received 26 March 1912

EX LIBRIS
THE PENNSYLVANIA
HORTICULTURAL SOCIETY

Pl. I. View of Mt. Fuji, seen from Ōmiya; Larix, evergreen conifer, deciduous broad leaved tree, and prairie, regions are distinctly visible. (Sketched from nature by the author in April 1903.)

THE
VEGETATION OF MT. FUJI
(JAPAN)

♦

理學博士 早田文藏著

富士植物帶論

富士山植物總目錄及ビ
富士山植物分布地圖

明治四十四年一月印行

東京 丸善株式會社

The Vegetation of Mt. Fuji

WITH A COMPLETE LIST OF PLANTS FOUND ON THE
MOUNTAIN AND A BOTANICAL MAP SHOWING
THEIR DISTRIBUTION

By

B. HAYATA, *Rigakuhakushi*

*Lecturer on Botany, in the Botanical Institute, College of Science,
Imperial University of Tōkyō.*

AUTHOR OF FLORA MONTANA FORMOSÆ, REVISIO EUPHORBIACEARUM
JAPONICARUM, COMPOSITÆ FORMOSANÆ ETC. AND ONE OF
THE AUTHORS OF ENÜMERATIO PLANTARUM
FORMOSANARUM.

WITH EIGHT PLATES, A COLOURED BOTANICAL MAP AND
THIRTY FIVE FIGURES IN THE TEXT.

THE
MARUZEN-KABUSHIKI-KAISHA

(Z. P. MARUYA & Co., LTD.)

TOKYO, OSAKA & KYOTO.

1911.

Printed at THE TOKYO TSUKIJI TYPE FOUNDRY.

T O K Y O

RECEIVED
GEORGE ENGELMANN
MAR 26 1912
Gray Herbarium
Harvard University

all rights reserved.

16197

CONTENTS.

INTRODUCTION.

	PAGE.
PHYSIOGRAPHY.	3
CLIMATE.	8
GENERAL ASPECT OF THE VEGETATION.	21
<i>a.</i> PLANT FORMATION IN PART I.... ..	21
<i>b.</i> PLANT FORMATION IN PART II.	29
<i>c.</i> PLANT FORMATION IN PART III.	34
<i>d.</i> PLANT FORMATION IN PART IV.	36
ZONATION OF THE PLANT-REGIONS.	38
<i>a.</i> PRAIRIE REGION.	39
<i>b.</i> DECIDUOUS BROAD LEAVED TREE REGION.	72
<i>c.</i> EVER-GREEN CONIFER REGION.... ..	88
<i>d.</i> LARIX REGION.	93
<i>e.</i> SALIX-ALNUS REGION.	95
<i>f.</i> HIGHER GRASS REGION.	97
GENERAL ASPECT OF THE FLORA OF THE MOUNTAIN. ...	101
INFLUENCE OF FIRE ON THE PRESENT VEGETATION. ...	107
ON THE ABSENCE OF PINUS PUMILA ON THE MOUNTAIN.	108
SUMMARY.	109
INDEX.... ..	111

Distant view of Mt. Fuji, seen from the Wada-pass ; the foreground shows a forest of *Pinus densiflora*, SIEB. et ZUCC.

(From the Atlas of Japanese Vegetation, by permission of Prof. M. MIYOSHI.)

In
Memory
of My Mother

Who

devoted Her Life to the Education of Her Three Sons and

by Whose Encouragement It was led to devote My

Own Life to the Study of Botany.

我等三人の兄弟を育て常に志を堅ふすべきを己に訓へ給ひし
亡き母君に報恩の紀念として此書を奉る

文

藏

明治戊申の歳亡き母君の第十三回の忌辰に當りて

INTRODUCTION.

ON the shore of the Pacific, towering to a height of more than twelve thousand feet, stands Mt. Fuji, symmetrical in shape and rich in vegetation, a fitting emblem of the Japanese nation. For several years past, I have taken a special interest in the botany of this beautiful mountain. The undulating prairie covering the gentle slope at the base, the dense forest stretching up to the middle, and higher up the dark growth of firs, are all so interesting and impressive that I have never tired of visiting it in every season of the year; observing how the vegetation changes as one makes the circuit of the mountain, or how the forest is differentiated according to height as one climbs to the peak. In summer, I have spent my vacations in the study of the flora, examining species after species, when all the prairie below is decorated with a great variety of flowers, and the broad leaved forest spreads out its foliage to the sunshine. But in winter, when the snow covers the peak and throws the dark forest of ever-green conifers into sharp relief, my attention has been turned to the investigation of the plant formations zone after zone, each conforming to a contour line which is in itself circular. No other mountain in the Empire shows so regular a vegetation.

In my first visit to this beautiful mountain some five years ago, when I was travelling around the gentle slope of the truncated cone, my attention was at once attracted to the dif-

ference between the forest zones on the different sides of the mountain. Then the question immediately arose what has caused this difference almost as symmetrical as Mt. Fuji itself?

The vegetation, as one may see from the base of the mountain, is roughly divided into six zones according to the altitude, viz. the prairie, the region of broad leaved trees, the ever-green conifer belt, the region of *Larix*, the stunted shrubbery, and the alpine stretches. But more broadly speaking, when only the sides of the mountain are considered, the regions of vegetation may be divided into two — northern and southern. On the southern side of the tree region, deciduous broad leaved trees predominate, but on the north the ever-green needle leaved trees are characteristic. The cause of this difference is an interesting subject for investigation.

In the course of my study, I have found that the different aspects of the tree regions are explainable by the differences of climate and soil according to altitude and exposure. In this paper, I have endeavoured also to enumerate all the plants found on the mountain and to arrange them in order according to regions, as explained in the map annexed.

In conclusion, I must express my hearty thanks to Prof. J. MATSUMURA, by whose special permission this work has been carried out. I am also indebted to Prof. M. MIYOSHI for his kindness to give me the privilege of reproducing some figures from his "Atlas of Japanese Vegetation"; to Dr. Y. SHIOZAWA by whose kindness I was able to secure all the data relating to the alternation of the forest regions; and to Mr. T. MAKINO who has kindly assisted me in determining the collections that I have made on the mountain. Nor should I forget to mention my obligations to Dr. T. OKADA who has liberally furnished me with all the meteorological data relating to the peak.

June 1907.

Fig. 1. Sketch-map of Mt. Fuji and its neighbouring peaks.
 ----- Provincial boundaries.

I. PHYSIOGRAPHY.

As vegetation has a close relation to the physiography of its locality, we must before going into details pause to consider the geology and climate of Fujiyama. Further, in order to study the relation of the flora of one mountain to that of another, we must necessarily know the geological relation between them. This chapter, therefore, will be devoted to the consideration of geology of this mountain and the neighbouring ranges, so far as is needful for investigating this subject.

Mt. Fuji, rising from the broad plain between the Provinces of Kai and Suruga, is a well defined elevation. It is

situated so that the meridian of $138^{\circ} 44'$ East of Greenwich, and the parallel of $35^{\circ} 21'$ North intersect near its summit. Attaining the height of 3778m. (approximately) above the sea level, this truncated cone adds a most graceful feature to the landscape of the Pacific coast. Its foot covers an extensive area, measuring 45 km. from N. to S. and 30 km. from E. to W. Combined with the elevations of the Myōkō volcanic group on the north, Mt. Yatsugatake in the middle, and far on the south Mts. Amagi and Izu-Shichitō,* it forms the Fuji volcanic chain, which divides the main-island into two parts. Among the volcanoes of this chain, Fujiyama, being the one formed most recently probably in the diluvian age, is thought to have the most recent vegetation.

It has Mt. Ataka at its very base, and Mt. Amagi some eighty miles to the south, while Mt. Kinbusen is on the north at a distance of about eighty miles. Further on, a hundred miles to the north-west, is Mt. Yatsugatake with Mt. Komagatake a little nearer, and Mt. Akaishi on the west at a distance of about eighty miles. All these mountains are supposed to have been already extinct, while Mt. Fuji was still active. And therefore, it is highly probable that all these elevations in the vicinity were crowned with rich vegetation while the young volcano was still bare of organic growth.

So much for the geology of the elevations in the vicinity. Turning now to the mountain itself, it has five lakes at its base, viz:—Yamanaka, Kawaguchi, Nishi, Shōji and Motosu. They are said with great probability to have been originally a semicircular lake skirting the base of the mountain. This lake was afterwards divided into five by the diminution of its water, and also by the intrusion of lava that flows from the volcano. These lakes give a great variety to the otherwise monotonous scenery of the basal slope, and especially on the north-western flank

* Seven islets near Prov. Izu.

where an extensive forest is found, owing to the underground water afforded by the lakes.

Although the mountain is entirely in a dormant stage, there is still observed some heat, 50°C in the rocks on the top,—an indication that the volcano has but recently ceased to be active. The inclination of the flank near the top is rather steep, measuring 34° – 32° , then gradually becoming a more and more gentle declivity of 25° – 17° , and at last most gently sloping league after league almost to a level. The shoulders of this volcanic cone are in all respects quite symmetrical, and its general outline is comparable to a logarithmic curve, so uniformly do the flanks slope in all directions.

The flanks are, as has been stated by Dr. T. HIRABAYASHI, all covered with lava broken into jaggy blocks, or invested with volcanic ashes. Generally speaking, the mountain is covered with lavas and cinders from the peak down to an elevation of 1500m., but the basal portion below is covered with tufas, sand and pebbles. It must be admitted however that there is a little local difference on the different sides,—for example, on the N. W. side the Aogigahara-lava occupies a very extensive area reaching down to an elevation of 1000m., while on the east the volcanic ashes and pebbles cover a broad area at the base and reach up as high as to elevation of 2500m.

Lavas are necessarily accompanied by scorïæ of various kinds. I may mention here that lava, when it consists of jaggy blocks, can absorb much water in its substance and is more likely to hold water upon its surface. When lava is thus broken into jaggy blocks, it is all the better for retaining water; because from a smooth plane surface water will run off rapidly; but with a rough surface, it has more chances of being retained. Not only has rough ground more power to hold water for the plants, but also it supplies the young seedlings with shelter sufficient to help their growth. But, if the lava consists of scorïa, the case is a little different. Scorïa is hard and glassy,

and it is rather difficult for plants to fasten their roots upon it even when there is a plentiful supply of water. Volcanic ashes or tufas can hold a considerable quantity of water; but an area covered with tufas is necessarily plane and smooth, and here in a heavy rainfall, the water is likely to run off from its surface. What makes the matter worse for plant-seedlings is that the area affords no place where plants can fasten their roots and is too flat to have any holes or shady places for the young plants. When strong and dry winds prevail, such ground at once loses all moisture and the plants find it difficult to grow. In this sense, in a surface covered with jagged blocks of lava, the plants can best find their rooting.

After stating so much, let us proceed to the consideration of the lavas of Mt. Fuji. According to Dr. T. HIRABAYASHI, some of the lavas belong to the eruption after the historical age, and therefore we are able to estimate their ages together with those of the vegetations upon them. There are seven streams of lava, each differing from the other, implying pre-historical and post-historical periods. They are as follows:— 1) the Ōsawa, 2) Mishima, 3) Ōmiya, 4) Karasunishi, 5) Yenkyō, 6) Aogigahara, and 7) Hōyē streams. The first four are pre-historical and the last three post-historical. All the lavas of this volcano are basic. The Yenkyō lava is said to have erupted in the year 880, the Aogigahara in the year 955, and the Hōyē lava in the year 1707.

Of all these lavas, the Aogigahara which stretches over the N.W. flank is the most porous and capable of supporting vegetation.* The other lavas are more or less dense and compact. The Hōyē lava consists of ashes or small particles of lava blocks, and is not fitted for plant growth. We, therefore, see the richest vegetation on the Aogigahara side and the poorest on the Hōyē side.

* The ground covered with this lava is especially fitted for the growth of conifers.

Pl. II. View of Mt. Fuji from Ukishima-muna. (Ukishima-swamp.)

The conifer region is seen quite dark below the treeless region clad with snow, and the region of deciduous broad leaved forests is seen less dark below the conifer region.

(After a photograph; a little brush is put in by the author, in order to show the plant regions clear.)

As we have stated above, the distribution of the lava has a close relation to the development of the forest. Independent of this matter, however, there must be certain differences of plant

Fig. 2. Sketch-map of Mt. Fuji and its vicinities; showing the positions of meteorological stations at the base of Mt. Fuji.

formation caused by climate. The next chapter is, therefore, devoted to the climate of the mountain.

II. CLIMATE.

As Mt. Fuji lies in the monsoon region, it has a heavy precipitation in the summer, with clear weather in the winter. Accordingly, its vegetation up to the height of 1500m. is mostly summer green forest formation. Above that line, conifers are found predominating over deciduous trees. In this chapter, we shall try to examine the climatic character of the different sides of the mountain. We have five meteorological stations, varying in altitudes from 100m. to 1000m., and located on different sides of Mt. Fuji. For the present investigation, it is necessary to take into account only the observations made at the altitude of about 1000m.; for the forest formation begins at that elevation. The five stations are not all in the tree regions, but it is possible for us to calculate the missing data from the known readings made at the base stations.

The localities of the five stations are shown in the accompanying sketch. They are as follows: 1) the Shiraito station, situated S.W. of the peak; 2) the Shōji station, N.W.; 3) the Ōmiya station, due S.; 4) the Goten station, S.E.; 5) the Nakano* station almost due E.

The climatic elements which must be taken into consideration in investigating plant formation are: 1) rainfall; 2) temperature; 3) humidity; 4) sunshine; 5) wind-direction. Let us consider them one by one.

1) **Rainfall.** It is a well known fact that other things being equal, the greater the rainfall the better for the vegetation. The following table shows the average rainfall for five years observed at the five stations.

* Nakano is located very near Yamanaka in the accompanying sketch.

Tab. I.

Stations	Latitude	Longitude	Altitude	Years	Jan.	Feb.	Mar.	Apr.
Shiraito	35° 17'	138° 35'	484 _m	1900-1904.	61.8	47.3	213.1	224.0
Ōmiya	35° 14'	138° 35'	150 _m	1899-1904.	60.7	55.6	216.4	205.5
Goten	35° 18'	138° 55'	456 _m	1900-1904.	81.8	34.4	144.9	244.0
Nakano near (Yamanaka)	35° 26'	138° 51'	986.4 _m	1901-1905.	75.4	90.1	248.1	225.5
Shōji	35° 31'	138° 37'	936.7 _m	1901-1905.	61.5	69.6	184.9	189.4

Stations	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Ann'l
Shiraito	307.1	275.2	409.9	270.2	278.5	238.5	121.6	117.0	2564.2
Ōmiya	237.6	231.4	386.3	178.3	251.5	249.8	98.6	105.3	2277.0
Goten	233.4	236.5	376.6	182.0	250.1	207.2	117.0	96.1	2204.0
Nakano near (Yamanaka)	259.3	345.7	489.9	266.7	331.9	318.5	125.0	159.1	2935.2
Shōji	246.0	320.2	415.9	247.5	301.5	251.8	92.0	109.5	2489.8

Tab. I. Containing the latitudes, longitudes and altitudes of the stations, the years of the observations, and the mean annual and monthly rainfalls (in millimeters), for the five stations.

It is very desirable to know the precipitation of the 1000m. altitude, as the forest regions begin at that point. We all

know that the rainfall, in the case of a mountain, increases up to the altitude of 1000m. and then decreases above that line. We can approximately calculate the rainfall of a flank of a mountain at any altitude, if we but know the rate of increase of rainfall for each 100m. The stations at Shiraito and Ōmiya are almost in a line from the top of the mountain. And therefore the difference of rainfall observed at these two stations will tell us the rate of increase of rainfall due to the altitude. The rate of increase in every 100m. thus calculated is 72mm. in annual mean of rainfall. Applying this rate to each station, we calculate the annual mean rainfall of the five directions at an altitude of 1000m., which amounts are shown in the following table.

Tab. II.

Annual mean rainfall at the altitude of 1000m. on the Shiraito side.		2936 mm.
„	„ Ōmiya side.	2889 „
„	„ Goten side.	2596 „
„	„ Nakano side.	2942 „
„	„ Shōji side.	2534 „

In this table we see the Nakano side leads in the amount of rainfall, next come the Shiraito and Ōmiya sides, while the Goten and Shōji sides have the least. This climatic character, however, seems to have but slight significance in explaining the present vegetation; for vegetation does not depend so much upon rainfall, as it does upon the soil, ground water, and humidity, which will be considered later.

However great the rainfall, if it comes all at once and runs away very rapidly, it will favour growth far less than frequent light rains. So the frequency of the rains should be taken into consideration.

Tab. III.

Months Stations	Jan.	Feb.	Mar.	Apr.	May	Jun.	July	Aug.	Sep.	Oct.	Nov.	Dec.	Ann'l
Shiraito	6	7	14	15	13	13	16	11	13	14	8	6	136
Ōmiya	5	6	13	12	13	13	13	5	10	11	6	4	111
Goten	5	5	14	12	12	10	14	6	11	12	8	5	114
Nakano	7	5	13	13	13	15	15	11	13	15	7	7	134
Shōji	6	6	12	14	14	17	16	13	14	14	8	7	141

Tab. III. Containing the average number of rainy days in each month, for the five years from 1901 to 1905 inclusive.

The figures of this table, being the result of observations made at different altitudes, do not give us any fair idea of the number of rainy days at the 1000m. altitude. We see, however, that these observations throw some light upon the investigation of the present plant-formation. The number of rainy days for Shiraito and Shōji are the greatest; hence they have the most abundant vegetation. Moreover, it may be urged that the Shiraito must have the most days of rain, for this station, notwithstanding its low elevation, has more rainy days than the Nakano station which lies almost at the altitude known to have the maximum precipitation. As will be shown afterwards, the most abundant growth of deciduous trees is found on the Shiraito side.

2) **Temperature.** The following table will give us the readings of the thermometers at the five stations. Here, however, we must consider the temperature at the altitude of 1000m., which temperature can be calculated from the known readings.

Tab. IV.

Stations	Years	Jan.	Feb.	Mar.	Apr.	May.	Jun.	Jul.	Aug.	Sept.	Oct.	Nov.	Dec.	Ann'l
Shiraito	1900-1904	6.3	6.3	9.3	13.9	17.9	21.0	23.4	26.5	23.4	18.3	12.9	8.4	15.6
Ōmiya	1899-1904	6.1	6.5	9.9	14.6	18.6	21.9	24.5	26.8	24.3	18.7	13.1	8.5	16.1
Goten.	1900-1904	3.2	3.9	7.4	13.2	16.5	20.7	22.6	26.0	23.3	16.9	11.1	6.0	14.2
Nakano	1900-1905	0.9	0.3	4.1	9.8	13.6	18.1	20.9	22.3	19.3	13.5	8.4	3.3	11.2
Shōji	1900-1905	0.3	0.6	4.4	9.8	13.8	18.3	21.2	22.3	20.0	13.6	7.6	3.0	11.3

Tab. IV. Containing the mean annual and monthly temperature (C.) for five stations; observations made daily at 10 A.M.)

From this table, we can calculate approximately the temperature at the height of 1000m. on the respective sides represented by the five stations. In making the calculation, it is necessary to know the gradient of temperature on Mt. Fuji; but owing to the scarcity of data, we are not able to obtain any reliable value of the gradient on the mountain. Fortunately, however, the gradient of temperature up to the 1000m. elevation is known with great approximity from observations made for long years on Mt. Tsukuba. As both mountains are situated on the Pacific coast, and stand near each other, the results obtained for Tsukuba will hold good for the basal portion of Mt. Fuji.

The following figures obtained by Dr. T. OKADA are given as the gradient of air temperature per 100m. on Mt. Tsukuba.

Tab. V.

Months	Jan.	Feb	Mar.	Apr.	May.	Jun.	Jul.	Aug.	Sept.	Oct.	Nov.	Dec.	Ann'l
Gradient	0.49	0.52	0.55	0.60	0.60	0.57	0.57	0.57	0.55	0.55	0.43	0.43	0.54

From Tables IV. and V., we obtain the following table showing the mean temperature for month and year at the 1000m. altitude on the respective sides of the mountain.

Tab. VI.

Months Sides	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Ann ^l
1000m. altitude on the Shiraito side.	3.8	3.6	6.5	10.8	14.4	18.1	20.5	23.6	20.6	15.5	10.7	6.2	12.8
„ „ Ōmiya side.	1.9	2.1	5.1	9.5	13.5	17.1	19.7	22.0	19.5	13.9	4.3	4.8	11.5
„ „ Goten side.	0.5	1.1	4.4	9.9	13.2	17.6	19.5	22.9	20.3	13.9	8.8	3.7	11.2
„ „ Nakano side.	0.9	0.3	4.1	9.8	13.6	18.1	20.9	22.3	19.3	13.5	8.4	3.3	11.2
„ „ Shōji side.	0.3	0.3	4.1	9.4	13.4	17.9	20.8	21.9	19.7	13.3	7.3	2.7	11.0

On looking at this table, we see that the temperature is the highest on the Shiraito side and next highest on the Ōmiya side. Other things being equal, the higher the temperature, the better plants grow. This is clearly shown in the plant formation of Mt. Fuji. On the Shiraito side, the forest is the richest, and the Ōmiya side comes next. On the Shōji side, the temperature is the lowest, but the forest is there favoured by the underground water due to the adjacent lake, and still more by the porous lava which covers all the ground*. On the Gotemba side, the temperature is comparatively high; but this advantage being offset by the poor soil, we have here the poorest vegetation.

3) **Humidity.** This most subtle thing plays an important rôle in plant formation. To my regret, we have no observations

* The case that the forest of conifers is favoured by porous lavas which cover the ground is everywhere met with in Japan. In Mt. Maccarinupri, we find conifer-forests flourish on lavas, while deciduous trees, on the ground covered by tufas and ashes.

Tab. VIII. (Temperature.)

Months Sides	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Ann'l
Altitude of 1000m. north.	-2.48	-1.3	2.1	8.1	12.8	17.2	20.4	21.6	17.3	10.4	6.0	0.4	9.3
Altitude of 1000m. south.	0.4	1.6	2.8	7.6	11.9	15.1	18.8	20.5	16.9	11.2	8.0	3.4	9.7

The maximum tensions E of the above temperatures are as follows :

Tab. IX. (Maximum tension = E .)

Months Sides	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Ann'l
E at the alti- tude of 1000m. on the north.	3.81	4.16	5.31	8.05	10.99	14.12	17.80	19.16	14.67	9.39	6.97	4.70	8.72
E at the alti- tude of 1000m. on the south.	4.70	5.12	5.58	7.78	10.36	12.76	16.12	17.91	14.30	9.9	7.99	5.82	8.96

The vapour tensions e_o at the two basal stations are shown in the following table. The observations cover the space of five years, 1896–1900 inclusive; and the figures are the mean of the readings made six times a day.

Tab. X. (Vapour tension e_o .)

Months Stations	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Ann'l
Kōfu	3.5	3.74	4.78	7.82	10.38	13.32	17.88	19.7	15.92	9.8	6.68	4.2	9.84
Numadzu	4.44	4.50	5.70	8.62	11.38	14.28	18.24	20.04	16.24	10.70	7.96	5.26	10.62

From Cale. I., Tab. X. and Formula I, we get the following table showing the vapour tensions e_h at the altitude of 1000m. on both northern and southern flanks.

Tab. XI. (Vapour tension e_h .)

Months Sides	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Ann'l
altitude of 1000m. on the north.	2.7	2.9	3.7	6.0	8.0	10.3	13.82	15.20	12.27	7.56	5.17	3.24	7.56
altitude of 1000m. on the south.	3.1	3.2	4.0	6.1	8.0	10.0	12.8	14.1	16.4	7.5	5.6	3.7	7.4

It is not vapour tension that plays an important part in the formation of forest; but it is saturation-deficit which is necessary and is the main cause in making forest.

From Tables IX and XI, we get the following table showing saturation-deficit $E - e_h$ on both flanks.

Tab. XII. (Saturation-deficit $E - e_h$.)

Months Sides	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Ann'l
1000m. altitude on the north.	1.01	1.26	1.61	2.05	2.99	3.82	3.39	3.96	2.4	1.83	1.8	1.46	1.16
1000m. altitude on the south.	1.6	1.9	1.58	1.68	2.36	2.76	3.32	3.81	2.9	2.4	2.39	2.12	1.56

Here we see in this table that, in January, February, September, October, November and December, i.e. in the so-called non-growing season, the southern flank is much drier than the northern; and contrariwise in the spring and summer, i.e. in the growing season. This difference of humidity clearly shows the difference of the vegetation of the two sides.

Fig. 3. View of Mt. Fuji seen from Shinshū. (N. E. side.) (after a phot.)

We have previously stated that the vegetation-regions of Mt. Fuji may be broadly divided into two according to side,— northern and southern. On the north, the ever-green conifers predominate, while on the south there is a most luxuriant growth of deciduous, broad-leaved trees. On the south, the air is dry in the winter, that does no harm to the deciduous trees. In the so-called growing season, the leaves of deciduous trees all come out, and it is rather dangerous for the trees when the air is dry, for they have such a broad surface of leaves for evaporation. In the winter, they have no leaves and, therefore, no surface for evaporation. In short, deciduous trees do not require moist air in the winter, but do in the spring and summer. In other words, deciduous trees have no fear of being dried up in the winter, for they have no leaves for transpiration; but in the summer when all their leaves are out, they are in great danger of losing too much water, if the air is dry. Therefore, deciduous trees can do well in a region where the air is drier in the

winter, and is moister in the summer. This, we see in Tab. XII., is the condition of the southern flank of Mt. Fuji.

In Tab. XII., we also see that, in the so-called non-growing season, i.e. in the winter, the air on the north side is rather humid, but in the spring and summer the air is comparatively dry. The ever-green conifers are as susceptible to a code of humidity in winter as in summer. But in the growing season, they have comparatively a smaller leaf-area for evaporation than the deciduous broad-leaved trees. They are, therefore, less in danger than the deciduous trees if the air is dry during the growing season. In short, the conifers can stand comparatively dry air in the growing season, but they can not stand it in winter. Therefore, the conifers will do well in a region where the air is comparatively moist in winter, and dry in spring and summer. This, as we see in Tab. XII., is the case with the region of the northern flank of Mt. Fuji.

As we have already stated, the temperature and rainfall can give but a faint explanation of the difference of the features of the forest-regions on the different sides. But a careful examination of the effects of humidity will prove it a most important reason for the differences found on the various flanks.

4) **Sunshine.** Lastly we shall consider the difference of the duration of sunshine on the different sides. The following table shows the possible duration of sunshine on both sides, granting that the observations at the Kofu station may answer for those of the northern side of Mt. Fuji, and the observations at the Numadzu station, those of the southern side of the mountain.

Tab. XIII.

Months Sides	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Ann'l
Kofu, northern side.	311	305	370	392	435	434	442	416	370	348	308	301	4432
Numadzu, southern side.	312	306	370	391	433	432	440	415	370	346	309	304	4431

The following table shows the actual duration of sunshine at the two stations.

Tab. XIV.

Months Sides	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug	Sep.	Oct.	Nov.	Dec.	Ann'l
Kōfu, northern side.	180	202	202	185	210	195	154	222	171	155	173	198	2246
Numadzu, southern side.	185	187	186	190	226	192	191	235	180	172	181	203	2306

Ratios of the possible duration to actual duration on both sides are shown in the following table.

Tab. XV.

Months Sides	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Ann'l
Kōfu, northern side.	58	66	55	47	49	45	35	53	46	45	56	66	52
Numadzu, southern side.	59	61	50	48	48	44	45	56	48	49	59	67	53

Upon considering the foregoing tables we find that through the year the weather is better on the southern side than on the northern. But it may be mentioned that in February and March the weather is better on the northern side than on the southern. This fact will at least favour conifers which have ever-green leaves even in the cold winter and have power to do starch-building even then; but will not benefit deciduous trees which have no leaves to enjoy the sunshine.

4) **Wind-direction.** It is true that the wind plays a somewhat important part in making forest. The fact is the gentler the wind the better for the forest. So, the velocity of the wind is the important point for us. We have, however, no information about the wind velocity at the base stations of the mountain.

Setting it aside, therefore, let us proceed to examine the direction of the wind.

As the wind is a means of conveying plant seeds to a distance, the discussion of the wind-direction will throw some light upon the questions whence and how plants were first brought to this volcano.

The following table shows the wind-direction at the different stations.

Tab. XVI.

Months Stations	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Ann'l
Shiraito	N	N	N	SE	SE	SE	SE	SE	N	N	N	N	SE
Ōmiya	S	S	S	S	S	S	S	S	S	S	S	S	S
Goten	S	S	S	S	S	S	S	S	S	S	S	S	S
Nakano	W	W	SW	S	S	S	S	S	S	S	S	SW	SW
Shōji	N	N	SE	SE	SE	SE	SE	SE	S	S	S	N	SE

Tab. XVI. Showing the wind-directions at the different stations.

As the mountain lies in the so-called monsoon region, the north wind prevails in the winter and the south wind in the summer. But the wind at the region under consideration is somewhat changed owing to the high mountain. Thus the prevailing wind throughout the year is the south wind.

As we have previously stated, the mountain must have been entirely bare, until it entered the extinct stage. After having become ready to give plants their stands, it must have received first some inhabitants from the neighbouring mountains. I have come on this occasion to consider that the wind is the only means of conveying plant-seeds to this mountain. The wind must,

therefore, play an important rôle in making up the flora of Mt. Fuji.

Here we must not forget to mention that in the middle layer of the air the constant wind is that from the south-west. In the case of Mt. Fuji, therefore, at the 2500m. elevation, the south-westerly wind prevails. It is only from southern mountains that the peak could have received alpine plants, and looking on the map we see that there is no high mountain on the south. In other words, there is no high elevation on the south to contribute alpine elements to Mt. Fuji's vegetation. This fact must go far towards explaining why the mountain is so poor in the alpine elements, as will be seen in the list. Such a plant as *pinus pumila* so common in the alpine regions in Japan is entirely wanting on this mountain.

III. GENERAL ASPECT OF THE VEGETATION.

In the preceding chapters, we have considered the physiography, geology and climate, of Mt. Fuji as far as is necessary for the discussion of the plant formation. In this chapter, we shall try to dwell upon the formation in general,—not to investigate the vegetation zone after zone, but to take a general glance over all the different parts of the mountain. In order to attain this end, it will be best to divide the mountain into four parts as shown in the accompanying sketch.

First, we shall consider these parts one by one; next we shall combine all parts into one in order to gain a general conception of the vegetation on the whole. Let us take Part I. for the first discussion.

a) Plant-formation in Part I.

On glancing over the volcano from the Ōmiya plain, at a distance of five miles from the base, we shall have a clear view of the southern flank with its abundant growth. The

arrangement of the vegetation according to the height is seen most clearly on this side.

Fig. 4. Sketch-map of Mt. Fuji; plant regions are divided to four parts.

As is shown in Plate I., the plant regions of the mountain are displayed very clearly by the different coloration of the formations; the truncated top, clad with pure white snow, or sometimes crowned with clouds; the hazy dark green region of the conifer-forest in the middle; the light red zone of the deciduous forest a little lower; then light brown to the base, and mile after mile of prairie-formation. The alpine stretches, shrubby growth, and *Larix*-formation, which lie above the conifer-forest, are but faintly seen. The deciduous forest enjoys the most luxuriant growth on this side.

As there is sufficient heat and considerable rainfall on this side, cultivation is carried on here and there even to the edge of the deciduous growth. The prairie-land on this side is said to have been formerly clad with a dense forest. It is known from the re-

cords that in former times clearings were made here, and more emphatically is this told by the roots or stumps buried in the soil.

Some portions in this part have been accurately surveyed by the Bureau of the Imperial Preserves for the first process of the afforestation now being carried out here. Through the kindness of Dr. Y. SHIOZAWA, expert of the Bureau, I was able to secure all the data with which I express numerically the degrees of the alternation of the forest-regions. The surveyed portions were divided into a number of parts, and the deciduous trees and conifers contained in some parts marked with asterisks were actually counted. The supply of timber contained in those parts is finally mentioned. This supply will approximately show the density of the forests.

The following tables show the relative frequency of both deciduous and coniferous trees in each part. The positions and the altitudes of the parts are given in the accompanying plate.

Tab. XVII.

Seet. I.

Tab. XVII. Showing the relative frequency of both deciduous and coniferous trees in Seet. I in the accompanying plate.

No.	Species	Larix	Picea and Abies	Deciduous broad leaved tree	Zelkova acuminata	No.	Species	Larix	Picea and Abies	Deciduous broad leaved tree	Zelkova acuminata
1			25%	75%		16			39%	57%	4%
2			23%	77%		17			28%	72%	
3			24%	74%		18			30%	70%	
4			35%	65%		19			18%	68%	14%
5			29%	71%		20			20%	71%	9%
6			29%	71%		21			20%	70%	10%
7			50%	50%		22			23%	72%	5%
8			33%	67%		23			29%	71%	
9			45%	55%		24			32%	68%	
10			34%	66%		25			21%	79%	
11			27%	73%		26			25%	75%	
12			29%	71%	4%	27			27%	63%	10%
13			21%	79%		28			24%	67%	9%
14			38%	58%	5%	29			20%	80%	
15			41%	54%							

Tab. XVIII.

Sect. II.

Tab. XVIII. Showing the relative frequency of both deciduous and coniferous trees in Sect. II. in the accompanying plate.

No.	Species	Larix	Abies, Tsuga	Abies, Tsuga and Picea	Deciduous broad leaved tree	Tsuga, Picea	No.	Species	Larix.	Abies, Tsuga	Abies Tsuga and Picea	Deciduous broad leaved tree	Tsuga, Picea
1		100%					32					100%	
2		100%					33					100%	
3		68%	31.6%		0.2%		34					100%	
4		45%	50%		5%		35		20%			80%	
5		2%		98%			36		8%			92%	
6				*96%	4%		37		8%			92%	
7		47%	41%		12%		38		43%			57%	
8		52%	44%		4%		39		27%			73%	
9		49%	49%		2%		40		15%			85%	
10			100%				41					100%	
11				100%			42		9%			91%	
12				*99%	1%		43					100%	
13		10%	80%		1%		44					100%	
14		9%	80%		11%		45					100%	
15			84%		16%		46		13%			87%	
16			65%		35%		47		16%			84%	
17					22%	78%	48		22%			78%	
18					20%	78%	49		19%			81%	
19			74%		26%		50		2%			98%	
20			78%		22%		51					100%	
21			81%		19%		52					100%	
22			62%		38%		53					100%	
23			27%		73%		54					100%	
24					100%		55		5%			95%	
25			52%		48%		56		10%			90%	
26			24%		76%		57					100%	
27			36%		64%		58					100%	
28			36%		64%		59					100%	
29			29%		71%		60					100%	
30			34%		66%		61					100%	
31			39%		61%		62					100%	

* including a small quantity of *Larix* in it.

Sect. II. (Continued.)

Nos. / Species	Larix.	Abies, Tsuga.	Abies, Tsuga and Picea.	Deciduous broad leaved trees.	Nos. / Species	Larix.	Abies, Tsuga.	Abies, Tsuga and Picea.	Deciduous broad leaved trees.
63				100%	75				100%
64				100%	76				100%
66				100%	77				100%
67				100%	78				100%
68				100%	79				100%
69				100%	80				100%
70				100%	81				100%
71				100%	82				100%
72				100%	83				100%
73				100%	84				100%
74				100%

Fig. 5. Pure stand of *Larix leptolepis* in No. 7 in Sect. III. (Photographed by the author).

Tab. XIX.

Sect. III.

Nos.	Species	Larix.	Abies, Tsuga.	Abies, Tsuga and Picea.	Abies.	Deciduous broad leaved trees.	Nos.	Species	Larix.	Abies, Tsuga.	Abies, Tsuga and Picea.	Abies.	Deciduous broad leaved trees.
1	100%						22						100%
2					100%		23						100%
3	6%				94%		24						100%
4					26%	74%	25						100%
5					100%		26					50%	50%
6	100%						27						100%
7	100%						28						100%
8						100%	29						100%
9			55%			45%	30						100%
10						100%	31						100%
11	100%						32						100%
12						100%	33						100%
13						100%	34						100%
14			52%			48%	35						100%
15						100%	36						100%
16						100%	37						100%
17						100%	38						100%
18						100%	39						100%
19						100%	40						100%
20					29%	71%	41						100%
21						100%	42						100%

Tab. XIX. Showing the relative frequency of both deciduous and coniferous trees in Sect. III. in the accompanying plate.

Fig. 6. Deciduous tree-forest in No. 22, in Sect. III. *Acer pictum* THUNE, *Stewartia monadelphica* S. et Z. and other trees are represented in the figure. (Photographed by the author.)

It is very interesting to arrange the parts marked with asterisks according to height, and to compare the density of the forest at the different altitudes. (cf. P. 24.)

As Sect. II. of this Part is the most typical on this side, I shall take this section for comparison.

Tab. XX.

Sect. II. (*Forest-growth.*)

Alt.					mean.
2250m.	(No. 6) 437				437
2000m.	(No. 7) 371	(No. 8) 406	(No. 16) 466	(No. 10) 830	516
1500m.	No. 42) 440	(No. 35) 443	(No. 38) 222	(No. 57) 266	343
1240m.	No. 69 167				167

The figures show timber-supply in 1 square km. measured in fm.

In this table, we see that the forest-zone, which extends from 2500m. downwards, attains its maximum growth at the altitude of 2000m., then decreasing gradually, comes to an end at the boundary of the forest, at an altitude of about 1000m.

Now considering the Conifers only, the following table shows the density of the conifer-forest according to height.

Tab. XXI.

Sect. II. (*Conifer-forest.*)

Alt.					Mean
2250m.	(No. 6) 425				425fm.
2000m.	(No. 7) 331	(No. 8) 406	(No. 16) 405	(No. 10) 830	495fm.
1500m.	(No. 42) 272	(No. 35) 68	(No. 38) 70	(No. 57) 0	102fm.
1250m.	(No. 69) 0				0
1000m.					0

In this table, we see that the Conifer-formation, which extends from 2500 m. downwards, attains its most luxuriant growth at an altitude of 2000m., then decreasing gradually, comes to an end at the 1250m. altitude.

The density of the deciduous forest according to height is shown in the following table.

PL. IV. View of Mt. Fuji seen from Lake Motosu, showing the plant regions found partly in Part IV. and partly in Part I. Conifer, deciduous tree, and prairie, regions are clearly seen.
(After a sketch by the author).

Tab. XXII.

Set. II. (*Deciduous forest*)

Alt.					Mean
2250m.	(No. 6) 12				12
2000m.	(No. 7) 39	(No. 8) 0	(No. 16) 61	(No. 10) 0	25
1500m.	(No. 42) 168	(No. 35) 374	(No. 38) 153	(No. 57) 266	240
1250m.	(No. 69) 167				167

In this table, we see that the deciduous forest which first comes in the shrubbery-formation at the high altitude of 2500m., increasing gradually, attains its maximum growth at the altitude of 1500m., then decreasing gradually, comes down to its boundary at the 1000m. altitude.

The following table shows the density of the forest-growth in general, and that of the coniferous and deciduous forest separately, according to the height.

Tab. XXIII.

b) *Plant-formation in Part II.*

Moving eastwards, let us discuss the formation of Part II. This is a rather limited area bordered by Mt. Ashitaka on the

south-west, and defined by the Kagosaka range on the north-east. From the Gotemba plain, we have a full view of this part. In the spring, when the snow still remains on the peak, the dark green belt of the pine-forest stands in clear relief against the back ground of the pure white snow. The belt is thick on the north, but becomes thinner and broader on the south, flanking downwards to the middle of the mountain, then giving way to the prairie which stretches far and wide to the fertile plain of Gotemba.

As is shown in the annexed map, the vegetation is the most scanty on this side. The treeless region, shown by the snow cap, stretches down as low as to about the 1200m. altitude, then giving way to *Picea*- or *Abies*-, but more especially to the red pine-, formation which is, however, not dense as in the case of other Parts, but very thin in every respect. This scarcity of vegetation is caused by the eruption of Hōyē which took place but recently, the ground being as yet not ready to produce a dense forest there.

The north-eastern portion of this Part (Sect. IV. in the accompanying sketch-map) was surveyed in connection with Part I. The following table will show the relative frequency of *Larix*, *Pinus densiflora*, and deciduous trees.

Tab. XXIV.

Sect. IV.

Species Nos.	Larix	Pinus densiflora,	Deciduous broad leaved trees	Species Nos.	Larix	Pinus densiflora,	Deciduous broad leaved trees
1	100%			6			100%
2	100%			7	27%		73%
3	100%			8	27%		73%
4	100%			9	100%		
5	55%		45%	10	100%		

Sect. IV. (Continued.)

Species Nos.	Larix.	Pinus densiflora.	Deciduous broad leaved trees.	Species Nos.	Larix.	Pinus densiflora.	Deciduous broad leaved trees.
11	100%			21		100%	
12	100%			22		100%	
13	67%	33%		23		100%	
14		100%		24		100%	
15	29%	71%		25		100%	
16		100%		26		100%	
17		100%		27		100%	
18		100%		28		100%	
19		100%		29		100%	
20		100%					

The following table will show the density of the forest-growth in general according to height.

Tab. XXV.

Sect. IV. (Forest-Growth)

Alt.			Mean
1800m.	(No. 2) 132		132 _f m.
1500m.	(No. 5) 131		131 _f m.
1300m.	(No. 7) 49		49 _f m.
1060m.	(No. 17) 68	(No. 18) 63	62 _f m.
940m.	(No. 24) 177		177 _f m.
840m.	0		0

As we see in the annexed map, the forest-growth comes into existence at an altitude of 2200m. and gradually increasing

as it descends, attains its most luxuriant growth at the 940m. altitude, then decreasing gradually, comes to an end at an altitude of 840m.

The density of the *Larix*-growth is shown in the following table.

Tab. XXVI.

Sect. IV. (*Larix leptolepis*)

Alt.			Mean
1800m.	(No. 2) 132		132
1500m.	(No. 5) 83		83
1300m.	(No. 7) 15		15
1060m.	(No. 17) 7	(No. 18) 4	5
940m.	(No. 24) 0.3		0.3

We see in this table that the *Larix*, which extends from an altitude of 2200m. downwards, attains its maximum growth at the height of 1800m., then gradually decreasing, comes almost to an end at an altitude of 940m.

The density of the growth of *Pinus densiflora* which stretches over this section is shown in the following table.

Tab. XXVII.

Sect. IV. (*Pinus densiflora*)

Altitudes			Mean
1800m.	(No. 21) 0		0
1500m.	(No. 5) 0		0
1300m.	(No. 7) 0		0
1060m.	(No. 17) 55	(No. 18) 31	53
940m.	(No. 24) 160		160

In this table, we see that *Pinus densiflora* which extends from 1300 m. downwards, attains its most luxuriant growth at an altitude of 970 m.

The density of the growth of the deciduous trees is as follows:—

Tab. XXVIII.

Sect. IV. (*Deciduous trees*)

Altitudes			Mean
1800m.	(No. 2) 0		0
1500m.	(No. 5) 46		46
1300m.	(No. 7) 37		37
1060m.	(No. 17) 6	(No. 18) 8	7
940m.	(No. 24) 17		17

The figures show timber-supply in 1 square hm. measured in 1 fm.

The maximum growth of the deciduous trees is seen at an altitude of 1500 m., but does not extend to any higher than an altitude of 1800 m.

The following table will show the density of the forest growth in general and that of the *Larix*, *Pinus* and deciduous trees, separately.

Tab. XXIX.

c) *Plant-formation of Part III.*

Now passing on to the north side, let us study Part III. The flank is generally like Part II., but has a far more developed forest-formation in the middle. The prairie is equally broad and extensive, not well suited to cultivation. This monotonous scenery of the prairie-formation is broken by lava-streams which show themselves here and there by the forests upon them.

Lakes are found on this side with water-plants such as *Potamogeton* and *Najas*. The general view of this grand flank is best obtained from the top of the Misaka pass. The colored plate annexed

Fig. 7. *Picea hondoensis* MAYR, with *Usnea longissima* ACH., about 1650 m. above the sea. (Yoshida side). (From the Atlas of Japanese Vegetation, by permission of Prof. M. MIYOSHI.).

showing the northern vegetation was made from a sketch by myself. It is a glorious sight to see the truncated cone, late in the spring when the snow covers the peak above, and the prairie below, but leaves the green of the pine-forest doubly dark against the pure white ground of snow. The deciduous tree-formation is on this

Fig. 8. Margin of the coniferous forest on the north side of the mountain. (Yoshida side).

side but faintly represented. The only vegetation distinguishable in the distance is the coniferous formation which surrounds the cone.

d.) Plant-formation of Part IV.

Turning our attention to the north-west side, let us consider the formation of Part IV. As is shown in the map annexed, the coniferous formation has here its most luxuriant growth. The Conifers stretch from the altitude of 2500 m. almost to the base. The deciduous forest is very poor and the prairie covers but a small area. A bird's eye view from the top of the Konno pass shows this part clad with ever-green conifers stretching like a level sea mile after mile of dark purple broken only by the mountain-range beyond. Broad leaved trees make patches here and there among the conifers, but are too scanty to produce any formation whatever. This coniferous region is called the "Aogi-

gahara"—the plateau of the ever-green trees. As we have previously explained, this beautiful formation is caused by the Aogigahara lava and the underground water supplied by the lakes.

Fig. 9. View of the north-western slope of the mountain, from Lake Shōji; the dark band at the base of the mountain is the Aogigahara Conifer-forest. (Photographed by the author).

Going further to the south-west through the Conifer-forest, we come down to the height of 1248 m. of Part I., where the formation of an immense deciduous forest opens before our eyes.

Thus, having completed our circuit of the mountain, let us consider the whole vegetation, combining the parts which we have discussed separately. Generally speaking, the formations are found to be more developed on the south side than on the north. Moreover, the deciduous forest predominates on the south, while on the north the Conifer-forest attains its most luxuriant growth. This difference of formation, as has been previously shown, is due to the physiography of the mountain on its different sides.

So much for the general aspect of the plant formation of

Mt. Fuji. The next chapter will be devoted to the consideration of the plant-regions zone by zone.

IV. ZONATION OF THE PLANT-REGIONS.

Climbing up through the vegetation of Mt. Fuji, we find that there are generally six regions according to altitude. They are 1) prairie*, 2) deciduous forest, 3) ever-green Conifers 4) *Larix*-region 5) *Salix-Alnus*-region and 6) alpine stretches. The deciduous forest is formed most abundantly on the south, but very scantily on the north and is likely to escape one's attention from a distance. The prairie-formation occupies a very spacious area on

Fig. 10. Prairie-formation on the Gotemba side. (Photographed by the author).

the south-east, while on the north-west, it is covered by the lower stretches of the Conifer-forest. Excepting these two small differences, the other formations are so equally developed on every side that they make nearly complete zones around the cone. The

* The term prairie is here used in the sense of grass-region.

local differences of these zones are shown in the map annexed. In this chapter we shall try to examine these regions one by one.

a) Prairie-Region (Basal region.)

This region mostly covers the gentle slope of the basal portion of the mountain. It is found on almost all sides, except the north-west, where the immense broad Conifer-forest stretches down almost to the base of the mountain.

The upper limitation of this low region is made very irregular by the intrusion of the forest-formation downwards. On the north-west side this boundary comes down as low as to about the 1000 m. altitude but on the south-east side, it goes up as high as to 1500 m. On the due south, it comes down to the 1000 m. line. On the north-east side, i.e. in Part III, there is seen the greatest extent of the prairie-formation. The upper limitation of this region on this flank lies at 1500 m. Two streams of lava crowned by the forest come down to the altitude of 1000 m. This undulating prairie is altogether monotonous broken only by the valleys which run into one another in their zigzag courses. Here and there tree-patches are seen along the streams. This side is especially suitable for the red pines, and they form thin forests which are in one part naturally made, but in another, formed by afforestation. Here on this side, I must not forget to mention, is the *Picea*-formation which is found quite isolated near lake Yamanaka. (cf. Pl. V.).

As we have already stated, the forest-formation regularly coincides with the lava-stream, that is clearly seen in the case of this *Picea*-formation.

Glancing at the geological map of the mountain, we see that on the north-east flank, a lava stream has flowed from the top down through the basal slope and has extended to the western extremity of lake Yamanaka. This stream is clearly traced by the forest-patches here and there upon it; for this

Fig. 11. Distant view of the pure stand of *Picea polita* near Lake Yamanaka; the foreground shows a cultivation of *Morus alba* LINN. (Photographed by the author).

Picea-formation is, after all, nothing but one of the forest-patches which flourish upon the porous lava of the stream.

Travelling along the base of the mountain on the north flank, one can not miss the noble forest of dark, green Conifers which extend over a distance of 2000 m., with a breadth of 1000 m. When I was journeying along this side, my attention was at once called to this great forest. On examining this secluded region, I found, to my astonishment, that it consists of a pure stand of *Picea polita*. This species is not rare on this mountain at rather high elevations, mixed with *Abies bicolor* and *Tsuga Sieboldi*. But to find its pure stand, so much broader than is seen elsewhere in the Empire, is something more than surprising.

The ground of this formation is all formed by large blocks of lava which are very porous in their structure, and very jagged in their surface. These characters of the lava have a very important relation to the forest-formation upon it. Moreover, underground water is plentifully supplied here by the adjacent lake. The Katsuragawa, an outlet of the lake, flows through this area of lava. Nothing is here wanting to make a luxuriant

Fig. 12. *Picea polita* in the pure stand near Lake Yamanaka.
(Photographed by the author).

growth. Towards the end of this beautiful formation, the forest becomes thinner and thinner until it gives way to the stretches of dwarf red pines. The accompanying photograph will give some idea of this pure stand.

We have one more forest-formation pushing down to this side of the prairie region. Looking at the northern area a little above, we see a stream of the Kenmarubi lava which is crowned, in the upper, with a Conifer forest, composed of *Picea* and *Abies*

and in the lower part, with young red pine. This forest is not so dense as that on the side of lake Yamanaka owing to the lack of the underground water. (cf. Map.)

On the south-west side, the Aogigahara lava comes down almost to the base of the mountain, and being nourished by the underground water, is covered by the dense coniferous forest of *Picea*, *Tsuga* and *Abies*.

Fig. 13. The Aogigahara forest at Shoji lake; the forest of the lake-side is seen in the back ground.
(Photographed by the author).

Generally speaking, all parts of the prairie-region have nearly the same plants. But when we consider the patches of forest which are found isolated in the prairie, some differences may be observed. On the southern flank, some deciduous broad trees come down into the prairie making patches here and there mostly along the valley and give some variety to the monotonous scenery on this side ; but on the northern flank, red pines play the part that the broad leaved trees take on the south. On the south, there is no group of pines on the prairie, nor is it likely that any

existed there in former times; for there is no evidence of one even in the relics buried in the soil. The red pine, therefore, does not seem to be suited to the south side, while the broad leaved trees grow better on this side. This fact plainly shows us that the southern prairie-side has sufficient temperature and humidity to maintain the broad leaved trees in the growing season, while the northern prairie-side has not. This is easily explainable by the climatic conditions of the different sides which we have already alluded to.

Fig. 14. *Cryptomeria*-avenue of the Asama-shrine (Yoshida).

The floral aspect of the prairie-region is almost the same on all sides. On the Jurigi side, there are some areas clad with *Arundinaria*, *Miscanthus* or sometimes with shrubbery. The plants found in the prairie-formation are as follows:—

Polypodiaceæ.

Latin names.	Japanese names.
<i>Adiantum monochlamys</i> EAT.	<i>Hakoneso</i> .
<i>Adiantum pedatum</i> LINN.	<i>Kujakushida</i> .
<i>Aspidium japonicum</i> MAKINO.	<i>Hariganewarabi</i> .
<i>Aspidium oligophlebium</i> CHRIST.	<i>Himewarabi</i> .
<i>Asplenium incisum</i> THUNB.	<i>Toranooshida</i> .
<i>Asplenium varians</i> HOOK et GREV.	<i>Iwatoranoo</i> or <i>Himotoranoo</i> .
<i>Athyrium nipponicum</i> HOOK.	<i>Inuwarabi</i> .
<i>Athyrium thelypteroides</i> DESV.	<i>Hakumōinode</i> or <i>Miyamashikeshida</i> .
<i>Athyrium yokoscense</i> MAKINO.	<i>Hebinonegoza</i> or <i>Koinuwarabi</i> .
<i>Blechnum Spicant</i> ROTH. var. sub- serrata LOWE.	<i>Shishigashira</i> .
<i>Coniogramme japonica</i> DIELS.	<i>Iwaganesō</i> .
<i>Cryptogramme japonica</i> THUNB.	<i>Tachishinobu</i> .
<i>Cystopteris japonica</i> LUERSS.	<i>Ushimewarabi</i> .
<i>Diplazium japonicum</i> BEDD.	<i>Shikeshida</i> .
<i>Diplazium lanceum</i> PRESL.	<i>Herashida</i> .
<i>Diplazium Wichuræ</i> METT.	<i>Nokogirishida</i> or <i>Yabukujaku</i> .
<i>Drymoglossum carnosum</i> HOOK.	<i>Mamezuta</i> .
<i>Microlepia hirsuta</i> PRESL.	<i>Inushida</i> .
<i>Microlepia marginalis</i> HANCE.	<i>Fumotoshida</i> .
<i>Microlepia Wilfordii</i> MOORE.	<i>Ōrenshida</i> .
<i>Nephrodium decursivo-pinnatum</i> BK.	<i>Gejigejishida</i> .
<i>Nephrodium dilatatum</i> DESV.	<i>Shiranewarabi</i> .
<i>Nephrodium Filix-mas</i> RICH.	<i>Memma</i> , <i>Oshida</i> or <i>Miyamainode</i> .
<i>Nephrodium Filix-mas</i> RICH. var. lacerum CHRIST.	<i>Kumawarabi</i> .
<i>Nephrodium Maximowiczii</i> BAKER.	<i>Nantaishida</i> .
<i>Nephrodium Phegopteris</i> BAUMG.	<i>Miyamawarabi</i> .
<i>Nephrodium prolixum</i> DIELS.	<i>Ibukishida</i> .
<i>Nephrodium sophoroides</i> DESV.	<i>Hoshida</i> .
<i>Nephrodium Thelypteris</i> DESV.	<i>Himeshida</i> or <i>Shorima</i> .

Fig. 15. Prairie-region on the north side of the mountain. (after a photograph).

Niphobolus linearifolius HOOK.
Odontosoria chinensis KUHN. var.
tenuifolia MAKINO.
Onoclea sensibilis LINN.
Polypodium hastatum THUNB.
Polypodium lineare THUNB.
Polypodium vulgare LINN. var. *japonicum* FR. et SAV.
Polystichum amabile (SM.)
Polystichum aristatum PRESL.
Polystichum craspedosorum DIELS.
Polystichum falcatum DIELS var.
Fortunei BAKER.
Polystichum tripterum SM.

Polystichum varium PRESL.
Pteridium aquilinum KUHN.
Pteris cretica LINN.
Pteris serrulata LINN.
Struthiopteris germanica WILLD.
Woodsia manchuriensis HOOK.

Birōdoshida.

Horashinobu.
Kōyawarabi.
Mitsudeuraboshi.
Nokishinobu.

Oshagojidenda. .
Ōbakanawarabi.
Kanawarabi or *Hosobakanawarabi.*
Tsurudenda.

Yabusotetsu.
Jumonjishida, Shūmokushida or *Mitsudekaguma.*
Itachishida.
Warabi.
Ōbanoinomotosō.
Inomotosō.
Kusasotetsu.
Fukuroshida.

<i>Woodsia polystichoides</i> EATON.	<i>Iwadenda.</i>
<i>Woodwardia radicans</i> SW.	<i>Komochishida.</i>

Schizæaceæ.

<i>Lygodium japonicum</i> SW.	<i>Kanikusa.</i>
-------------------------------	------------------

Osmundaceæ.

<i>Osmunda regalis</i> LINN. var. <i>japonica</i> MILDE.	<i>Zenmai.</i>
--	----------------

Ophioglossaceæ.

<i>Botrychium ternatum</i> SW.	<i>Fuyunohanawarabi.</i>
--------------------------------	--------------------------

Equisetaceæ.

<i>Equisetum arvense</i> L	<i>Sugina.</i>
----------------------------	----------------

Lycopodiaceæ.

<i>Lycopodium clavatum</i> LINN.	<i>Hikagenokazura.</i>
<i>Lycopodium serratum</i> THUNB.	<i>Tôgeshiba.</i>

Selaginellaceæ.

<i>Selaginella Kraussiana</i> A. Br.	<i>Kuramaçoke.</i>
<i>Selaginella involvens</i> SPRING.	<i>Iwahiba.</i>

Coniferæ.

<i>Abies firma</i> S. et Z.	<i>Momi.</i>
<i>Cephalotaxus drupacea</i> S. et Z.	<i>Inugaya.</i>
<i>Juniperus rigida</i> S. et Z.	<i>Muro or Nezu.</i>
<i>Pinus densiflora</i> S. et Z.	<i>Akamatsu.</i>
<i>Taxus baccata</i> LINN. subsp. <i>cuspidata</i> PILG.	<i>Ichii or Araragi.</i>
<i>Torreya nucifera</i> S. et Z.	<i>Kaya.</i>

Sparganiaceæ.

<i>Sparganium longifolium</i> TURCZ.	<i>Mikuri.</i>
--------------------------------------	----------------

Potamogetonaceæ.

Potamogeton crispus LINN.	<i>Ebimo.</i>
Potamogeton polygonifolius POUR.	<i>Irumushiro.</i>
Potamogeton Gaudichaudi CHAM.	<i>Sasabamo.</i>

Alismataceæ.

Sagittaria sagittifolia LINN.	<i>Omodaka.</i>
-------------------------------	-----------------

Hydrocharitaceæ.

Blixa caulescens MAX.	<i>Yanagisubuta.</i>
Hydrocharis Morsus-rauæ LINN.	<i>Dochikagami.</i>
Ottelia alismoides PERS.	<i>Mizuôbako.</i>

Gramineæ.

Agopyrum semicostatum NEES.	<i>Kamogigusa.</i>
Agrostis perennans TUCKERM.	<i>Yamanukabo.</i>
Alopecurus fulvus SM.	<i>Suzumenoteppô.</i>
Andropogon brevifolius SW.	<i>Ushikusa.</i>
Andropogon micranthus KUNTH. var. genuinus HACKEL.	<i>Himeaburasusuki.</i>
Andropogon Nardus LINN. var. Gær- ingii HACKEL.	<i>Ogarukaya.</i>
Arthraxon ciliare BEAUB.	<i>Kobunashiba.</i>
Arundinella anomala STEUD.	<i>Todashiba.</i>
Beckmannia erucæformis HOST.	<i>Minogome.</i>
Brachypodium silvaticum BEAUV.	<i>Miyamakamogigusa.</i>
Bromus japonicus THUNB.	<i>Suzumeno-chahiki.</i>
Bromus pauciflorus HACK.	<i>Kitsunegaya.</i>
Brylkinia caudata FR. SCHM.	<i>Hajigaeri.</i>
Calamagrostis arundinacea ROTH. var. sciuroides HACK.	<i>Saitôgaya.</i>
Calamagrostis Epigeios ROTH.	<i>Yamaawa.</i>
Diarrhena japonica FR. et SAV.	<i>Tatsunohige.</i>
Eleusine indica GÆRTN.	<i>Ohigiva.</i>
Eragrostis ferruginea BEAUV.	<i>Kazekusa.</i>
Eragrostis pilosa BEAUV.	<i>Nivahokori.</i>
Eriochloa villosa KUNTH.	<i>Narukobie.</i>
Festuca ovina LINN.	<i>Ushinokegusa.</i>

Glyceria remota FRIES var. *japonica*
HACK.

Glyceria tonglensis CLARK.

Imperata arundinacea CYR. var.
Königii HACK.

Isachne australis R. BR.

Leersia oryzoides SW. var. *japonica*
HACK.

Lophatherum gracile BRONGN. var.
elatum HACK.

Milium effusum LINN.

Miscanthus Matsumuræ HACK.

Miscanthus sinensis ANDR.

Muehlenbergia japonica STEUD.

Muehlenbergia japonica STEUD. var.
hakonensis HACK.

Oplismenus Burmanni BEAUB.

Panicum acroanthum STEUD.

Panicum Crus-Galli LINN.

Panicum indicum LINN.

Panicum sanguinale LINN.

Miyamaichigotsunagi.

Ōbanodojōtsunagi.

Fushigechigaya.

Chigozasa.

Sayanukagusa.

Sasakusa.

Ibukinukabo.

Kariyasumodoki.

Susuki.

Nezumigaya.

Tachinezumigaya.

Chizinizasa.

Nukakibi.

Nobie.

Hainumeri.

Mehijiwa.

Fig. 16. View of Mt. Fuji seen from Yoshida; forest-regions are clearly visible.

1. *Abies Veitchii*, 2. *Tsuga Sieboldi*, 3. *Larix leptolepis*.

(after a photograph).

Pl. V. View of Mt. Fuji, seen from the Fujigawa (Fuji-river); forest regions are faintly visible.

(After a photograph)

<i>Panicum sanguinare</i> LINN. var. <i>ciliare</i> DÖLL.	<i>Kemehijiwa.</i>
<i>Paspalum Thunbergii</i> KUNTH.	<i>Suzumenohie.</i>
<i>Pennisetum japonicum</i> TRIN.	<i>Chikarashiba.</i>
<i>Phalaris arundinacea</i> LINN.	<i>Kusayoshi.</i>
<i>Phragmites communis</i> TRIN.	<i>Yoshi, Ashi or Jishibari.</i>
<i>Poa annua</i> LINN.	<i>Suzumenokatabira.</i>
<i>Poa palustris</i> LINN. var. <i>strictula</i> HACK.	<i>Kawaraichigotsunagi.</i>
<i>Pollinia imberbis</i> NEES. <i>genuina</i> HACK.	<i>Ashiboso.</i>
<i>Pollinia nuda</i> (TRIN.) HACK.	<i>Sasagaya.</i>
<i>Polypogon Higegaweri</i> STEUD.	<i>Hiegairi.</i>
<i>Rottbøllia compressa</i> LIN. f. var. <i>japonica</i> HACK.	<i>Ushinoshippe.</i>
<i>Sasa albo-marginata</i> MAK. et SHIB.	<i>Kumazasa.</i>
<i>Sasa borealis</i> MAK. et SHIB.	<i>Suzutake.</i>
<i>Setaria glauca</i> BEAUB.	<i>Kin-enokoro.</i>
<i>Setaria pachystachys</i> F. et S.	<i>Hama-enokoro.</i>
<i>Setaria viridis</i> BEAUV.	<i>Enokorogusa.</i>
<i>Spodiopogon cotulifer</i> HACK.	<i>Aburasusuki.</i>
<i>Spodiopogon sibiricus</i> TRIN.	<i>Oburasusuki.</i>
<i>Sporobolus elongatus</i> R. BR.	<i>Nezumino-o.</i>
<i>Themeda Forskali</i> HACK. var. <i>japonica</i> HACK.	<i>Megarukaya.</i>
<i>Trisetum flavescens</i> BEAUB. var. <i>papillosum</i> HACK.	<i>Kanitsurigusa</i>
<i>Zoysia pungens</i> WILLD.	<i>Shiba</i>

Cyperaceæ.

<i>Bulbostylis barbata</i> KUNTH.	<i>Hatagaya</i>
<i>Carex arenicola</i> F. SCHMIDT.	<i>Kurokuzasuge</i>
<i>Carex Brownii</i> TUCK. var. <i>transversa</i> KUEK.	<i>Yuwasuge or Awabosuge.</i>
<i>Carex brunnea</i> THUNB.	<i>Nakirisuge</i>
<i>Carex caryophylla</i> LAUER subsp. <i>nervata</i> KUEK.	<i>Shibasuge.</i>
<i>Carex curvicolis</i> FR. et SAV.	<i>Yamanaruko.</i>
<i>Carex Gaudichaudiana</i> KUNTH var. <i>Thunbergii</i> KUEK.	<i>Azesuge.</i>

<i>Carex gibba</i> WAHL.	<i>Masukusa.</i>
<i>Carex incisa</i> BOOT.	<i>Tanisuge.</i>
<i>Carex ischnostachya</i> STEUD.	<i>Jūzusuge.</i>
<i>Carex japonica</i> THUNB.	<i>Higokusa.</i>
<i>Carex japonica</i> THUNB. var. <i>aphanolepis</i> KUEK.	<i>Sawasuge.</i>
<i>Carex japonica</i> THUNB. var. <i>chlorostachys</i> KUEK.	<i>Shirasuge.</i>
<i>Carex Morrowii</i> BOOT.	<i>Kansuge.</i>
<i>Carex Onoei</i> FR. et SAV.	<i>Harisuge.</i>
<i>Carex pseudo-conica</i> FR. et SAV.	<i>Ōetosuge.</i>
<i>Carex satsumensis</i> FR. et SAV.	<i>Aburashiba.</i>
<i>Carex stenantha</i> FR. et SAV.	<i>Iwasuge.</i>
<i>Cyperus difformis</i> LINN.	<i>Tamagayatsuri.</i>
<i>Cyperus Haspan</i> LINN.	<i>Mizuhanabi.</i>
<i>Cyperus Iria</i> LINN.	<i>Kogomegayatsuri.</i>
<i>Cyperus pilosus</i> VAHL.	<i>Ushigayatsuri</i> or <i>Onigaya'suri.</i>
<i>Cyperus rotundus</i> LINN.	<i>Hamasuge.</i>
<i>Eleocharis acicularis</i> R. BR.	<i>Matsubai.</i>
<i>Eleocharis japonica</i> MIQ.	<i>Iharii.</i>
<i>Fimbristylis autumnalis</i> RICH. et SCH.	<i>Himetentsuki.</i>
<i>Fimbristylis diphylla</i> VAHL.	<i>Tentsuki.</i>
<i>Fimbristylis miliacea</i> VAHL.	<i>Hideriko.</i>
<i>Fimbristylis squarrosa</i> VAHL.	<i>Azetentsuki.</i>
<i>Kyllingia brevifolia</i> ROTTE.	<i>Himekugu.</i>
<i>Lipocarpa microcephala</i> KTH.	<i>Hinjigayatsuri.</i>
<i>Mariscus Sieberianus</i> NEES.	<i>Kugu.</i>
<i>Pycreus globosus</i> REICH. var. <i>stricta</i> HOOK.	<i>Azegayatsuri.</i>
<i>Scirpus erectus</i> POIR.	<i>Hotarui.</i>
<i>Scirpus Cyperinus</i> KUNTH var. <i>concolor</i> MAK.	<i>Aburagaya.</i>

Araceæ.

<i>Acorus gramineus</i> AIT.	<i>Sekishō.</i>
<i>Arisæma serratum</i> SCHOTT. form.	
<i>Blumei</i> MAK.	<i>Tennanshō.</i>
<i>Pinellia tripartita</i> SCHOTT.	<i>Ōhange.</i>

Eriocaulaceæ.

Eriocaulon sexangulare LINN. *Mizutamiasō.*

Commelinaceæ.

Aneilema keisak HASSK. *Ibokusa.*
Commelina communis LINN. *Tsuyukusa.*

Pontederiaceæ.

Monochoria Korsakowii REGEL. et
MAACK. *Mizuaoi.*
Monochoria vaginalis PRESL. var.
plantaginea SOLMS.-LAUB. *Mizunagi.*

Juncaceæ.

Juncus effusus LINN. var. decipiens
BUCH. *Yi.*
Luzula campestris DC. var. capitata
MIQ. *Suzumenohie.*
Luzula plumosa E. MEY. *Nukaboshisō.*

Liliaceæ.

Allium japonicum REGEL. *Yamarakkio.*
Clintonia udensis TRAUTV. *Tsubameomoto.*
Disporum sessile DON. *Hochakusō.*
Disporum smilacinum A. GR. *Chigoyuri.*
Hemerocallis fulva LINN. *Yabukwanzō.*
Hosta cœrulea TRATT. form. lanci-
folia MATSUM. *Mizugibōshi.*
Lilium auratum LINDL. *Yamayuri.*
Lilium concolor SALISB var. parthe-
nion ELW. *Akahimeyuri.*
Lilium cordifolium THUNB. *Ubayuri.*
Lilium Maximowiczii REGEL. *Ko-oniyuri.*
Liriope graminifolia BAK. var. com-
munis MAX. *Yaburan.*
Metanarthecium luteo-viride MAXIM. *Nogiran*
Polygonatum latifolium DESF. var.
commutatum BAKER. *Narukoyuri.*

<i>Polygonatum officinale</i> ALL.	<i>Amadokoro.</i>
<i>Scilla japonica</i> BAKER	<i>Tsurubo.</i>
<i>Smilax china</i> LINN.	<i>Sarutoriibara.</i>
<i>Smilax herbacea</i> LINN.	<i>Shiode.</i>
<i>Smilax Sieboldi</i> MIQ.	<i>Yamakashiū.</i>
<i>Tricyrtis hirta</i> HOOK.	<i>Hototogisu.</i>
<i>Tricyrtis macropoda</i> MIQ.	<i>Yamahototogisu.</i>
<i>Veratrum Maackii</i> REGEL.	<i>Aoyagisō.</i>
<i>Veratrum nigrum</i> LINN.	<i>Shurosō.</i>

Amaryllidaceæ.

<i>Lycoris sanguinea</i> MAXIM.	<i>Kitsunenokamisori.</i>
---------------------------------	---------------------------

Dioscoreaceæ.

<i>Dioscorea gracillima</i> MIQ.	<i>Tachidokoro.</i>
<i>Dioscorea japonica</i> THUNB.	<i>Yamanoimo.</i>
<i>Dioscorea septemloba</i> THUNB.	<i>Momijidokoro.</i>

Iridaceæ.

<i>Iris japonica</i> THUNB.	<i>Shaga.</i>
<i>Iris sibirica</i> LINN. var. <i>orientalis</i> THUNB.	<i>Ayame.</i>

Orchidaceæ.

<i>Calanthe discolor</i> LINDL.	<i>Ebine.</i>
<i>Epipactis erecta</i> WETTST.	<i>Ginran.</i>
<i>Epipactis falcata</i> WETTST.	<i>Kinran.</i>
<i>Epipactis Thunbergii</i> A. GRAY.	<i>Suzuran.</i>
<i>Goodyera pendula</i> MAXIM.	<i>Tsurishūsuran.</i>
<i>Liparis auriculata</i> BLUME.	<i>Kumokirisō.</i>
<i>Neottia micrantha</i> LINDL.	<i>Himemuyōran.</i>
<i>Platanthera ussuriensis</i> MAXIM.	<i>Tonbosō.</i>
<i>Spiranthes australis</i> LINDL.	<i>Nejibana.</i>
<i>Oberonia japonica</i> MAK.	<i>Yōrakuran.</i>

Saururaceæ.

<i>Houttuynia cordata</i> THUNB.	<i>Dokudami.</i>
----------------------------------	------------------

Chloranthaceæ.

<i>Chloranthus serratus</i> R. et S.	<i>Futarishizuka.</i>
<i>Chloranthus japonicus</i> SIEB.	<i>Iitorishizuka.</i>

Salicaceæ.

<i>Populus tremula</i> LINN. var. <i>villosa</i> WESM.	<i>Yamanarashi.</i>
<i>Salix Caprea</i> LINN.	<i>Bakkoyanagi.</i>
<i>Salix japonica</i> THUNB.	<i>Shibayanagi.</i>

Betulaceæ.

<i>Alnus Yasha</i> MATSUM.	<i>Yashabushi.</i>
<i>Carpinus japonica</i> BLUME.	<i>Kumashide.</i>
<i>Carpinus yedoensis</i> MAXIM.	<i>Inushide.</i>
<i>Corylus heterophylla</i> FISH.	<i>Hashibami.</i>

Fagaceæ.

<i>Castanea vulgaris</i> LAM. var. <i>japo-</i> <i>nica</i> DC.	<i>Kuri.</i>
<i>Quercus acuta</i> THUNB.	<i>Akagashi.</i>
<i>Quercus cuspidata</i> THUNB.	<i>Shiinoki.</i>

Fig. 17. View of Mt. Fuji seen from Suzukawa, forest regions are faintly visible, (after a photograph.)

<i>Quercus dentata</i> THUNB.	<i>Kashiwa.</i>
<i>Quercus glandulifera</i> BLUME.	<i>Konara.</i>
<i>Quercus glauca</i> THUNB.	<i>Arakashi.</i>
<i>Quercus grosseserrata</i> BLUME.	<i>Mizunara.</i>
<i>Quercus serrata</i> THUNB.	<i>Kunnogi.</i>

Ulmaceæ.

<i>Celtis sinensis</i> PERS.	<i>Yenoki.</i>
------------------------------	----------------

Moraceæ.

<i>Fatoua pilosa</i> GAUD. var. <i>subcordata</i> RUPR.	<i>Kuwakusa.</i>
<i>Humulus japonicus</i> S. et Z.	<i>Kanamugura.</i>

Urticaceæ.

<i>Bœhmeria holosericea</i> BLUME.	<i>Oniyabumao.</i>
<i>Bœhmeria japonica</i> MIQ.	<i>Yabumao.</i>
<i>Bœhmeria japonica</i> MIQ. var. <i>tricuspis</i> HANCE.	<i>Akaso.</i>
<i>Bœhmeria nivea</i> BLUME.	<i>Mao.</i>
<i>Bœhmeria Sieboldiana</i> BLUME.	<i>Nagaboyabumao.</i>
<i>Bœhmeria spicata</i> THUNB.	<i>Koakaso.</i>
<i>Elatostema umbellatum</i> BLUME var. <i>majus</i> MAXIM.	<i>Uwabamisō.</i>
<i>Laportea bulbifera</i> WEDD.	<i>Mukagoirakusa.</i>
<i>Memoralis hirta</i> WEDD.	<i>Tsurumao.</i>
<i>Nanocnide japonica</i> BLUME.	<i>Katensō.</i>
<i>Pellionia radicans</i> WEDD.	<i>Sanshōsō.</i>
<i>Pilea petiolaris</i> BLUME.	<i>Miyamamuzu.</i>
<i>Urtica Thunbergiana</i> S. et Z.	<i>Irakusa.</i>

Loranthaceæ.

<i>Viscum album</i> LINN.	<i>Yadoriki.</i>
---------------------------	------------------

Santalaceæ.

<i>Thesium chinense</i> TURCZ.	<i>Kanabikisō.</i>
--------------------------------	--------------------

Aristolochiaceæ.

<i>Aristolochia debilis</i> S. et Z.	<i>Umanosuzukusa.</i>
<i>Aristolochia Kæmpferi</i> WILLD.	<i>Oba-umanosuzukusa.</i>

Polygonaceæ.

<i>Polygonum alatum</i> HAM.	<i>Tunisoba.</i>
<i>Polygonum cuspidatum</i> S. et Z.	<i>Itadori.</i>
<i>Polygonum esculentum</i>	<i>Soba.</i>
<i>Polygonum flaccidum</i> ROXB.	<i>Yanagitade.</i>
<i>Polygonum muricatum</i> LINN.	<i>Nagaba-unagizukami.</i>
<i>Polygonum Posumbu</i> HAM.	<i>Hanatade.</i>
<i>Polygonum Posumbu</i> HAM. var. <i>Blumei</i> (MEISN.)	<i>Inutade.</i>
<i>Polygonum senticosum</i> MEISN.	<i>Mamakonoshirinugui.</i>
<i>Polygonum Sieboldi</i> MEISN.	<i>Unagizuru.</i>
<i>Polygonum Thunbergii</i> S. et Z.	<i>Mizosoba.</i>
<i>Polygonum Thunbergii</i> S. et Z. var. <i>Maackianum</i> MAXIM.	<i>Sudekusa.</i>
<i>Polygonum viscosum</i> HAM. var. <i>ver-</i> <i>nicosum</i> MEISN.	<i>Nebaritade.</i>
<i>Rumex Acetosa</i> LINN.	<i>Suiba.</i>
<i>Rumex japonicus</i> MEISN.	<i>Gishigishi.</i>

Chenopodiaceæ.

<i>Chenopodium album</i> LINN.	<i>Akaza.</i>
--------------------------------	---------------

Amarantaceæ.

<i>Achyranthes bidentata</i> BL. var. <i>japonica</i> MIQ.	<i>Inokozuchi.</i>
<i>Amarantus Blitum</i> LINN.	<i>Inubiyu.</i>

Portulacaceæ.

<i>Portulaca oleracea</i> LINN.	<i>Suberihyu.</i>
---------------------------------	-------------------

Caryophyllaceæ.

<i>Arenaria serpyllifolia</i> LINN. var. <i>leptoclados</i> Guss.	<i>Nominotsuzuri.</i>
--	-----------------------

<i>Cerastium vulgatum</i> LINN. var.	
<i>glandulosum</i> KOCH.	<i>Miminagusa.</i>
<i>Cucubalus baccifer</i> LINN. var. <i>japonicus</i> MIQ.	<i>Nanbanhakobe.</i>
<i>Dianthus superbus</i> LINN.	<i>Kawaranadeshiko.</i>
<i>Lychnis Miqueliana</i> ROHR.	<i>Fushigurosenna.</i>
<i>Melandryum firmum</i> ROHR.	<i>Fushiguro.</i>
<i>Stellaria aquatica</i> SCOP.	<i>Ushihakobe.</i>
<i>Stellaria florida</i> FISCH. var. <i>angustifolia</i> MAXIM.	<i>Iwatsumekusa.</i>
<i>Stellaria media</i> LINN.	<i>Hakobe.</i>
<i>Stellaria uliginosa</i> LINN.	<i>Nominofusuma.</i>

Nymphæaceæ.

<i>Nuphar japonicum</i> DC.	<i>Kawahone.</i>
-----------------------------	------------------

Ceratophyllaceæ.

<i>Ceratophyllum demersum</i> LINN.	<i>Kingiyomo.</i>
-------------------------------------	-------------------

Magnoliaceæ.

<i>Illicium anisatum</i> LINN.	<i>Shikimi</i>
<i>Magnolia Kobus</i> DC.	<i>Kobushi.</i>
<i>Magnolia hypoleuca</i> S. et Z.	<i>Hô-noki.</i>
<i>Kadsura japonica</i> LINN.	<i>Sanekazura.</i>
<i>Schizandra nigra</i> MAXIM.	<i>Matsubusa.</i>

Ranunculaceæ.

<i>Aconitum Fischeri</i> REICH.	<i>Torikabuto.</i>
<i>Aconitum lycoctonum</i> LINN.	<i>Reiginsô.</i>
<i>Anemone cernua</i> THUNB.	<i>Okinagusa.</i>
<i>Anemone japonica</i> S. et Z.	<i>Shûmeigiku.</i>
<i>Aquilegia glandulosa</i> FISCH.	<i>Yamaodanaki.</i>
<i>Cimicifuga foetida</i> LINN. var. <i>simplex</i> HUTH.	<i>Sarashinashôma.</i>
<i>Clematis alpina</i> MILL.	<i>Miyamahanshōzuru.</i>
<i>Clematis apiifolia</i> DC.	<i>Botanzuru.</i>
<i>Clematis heracleifolia</i> DC. var. <i>stans</i> (S. et Z.)	<i>Kusabotan.</i>

<i>Clematis recta</i> LINN. var. <i>paniculata</i> (THUNB.)	<i>Senninsō.</i>
<i>Isopyrum stoloniferum</i> MAXIM.	<i>Shirokanesō.</i>
<i>Pæonia obovata</i> MAXIM.	<i>Yamashakuyaku.</i>
<i>Ranunculus acer</i> LINN.	<i>Kinpōge.</i>
<i>Ranunculus aquatilis</i> LINN. var. <i>flac-</i> <i>cidus</i> (PERS.) form. <i>Drouetii</i> HERM.	<i>Baikwamo.</i>
<i>Ranunculus japonicus</i> LUNGS.	<i>Kitsunenobotan.</i>
<i>Ranunculus sceleratus</i> LINN.	<i>Tagarashi.</i>
<i>Thalictrum minus</i> LINN. var. <i>elatum</i> LECOY.	<i>Akikaramatsu.</i>
<i>Thalictrum simplex</i> LINN. var. <i>affine</i> RGL.	<i>Nokaramatsu.</i>

Lardizabalaceæ.

<i>Akebia lobata</i> DCNE.	<i>Mutsuba-akebi.</i>
<i>Akebia quiuata</i> DCNE.	<i>Akebi.</i>

Berberidaceæ.

<i>Berberis Thunbergii</i> DC.	<i>Kotoritomarazu.</i>
<i>Epimedium macranthum</i> MORR. et DECNE.	<i>Ikarisō.</i>

Minispermaceæ.

<i>Cocculus Thunbergii</i> DC.	<i>Aotsuzurafuji.</i>
--------------------------------	-----------------------

Lauraceæ.

<i>Cinnamomum pedunculatum</i> NEES.	<i>Yabunikkei.</i>
<i>Lindera glauca</i> BLUME.	<i>Yamakōbashi.</i>
<i>Lindera obtusiloba</i> BLUME.	<i>Dankōbai.</i>
<i>Lindera præcox</i> BLUME.	<i>Aburachan.</i>
<i>Lindera umbellata</i> THUNB.	<i>Kuromogi.</i>
<i>Litsea glauca</i> SIEB.	<i>Shirodamo.</i>
<i>Machilus Thunbergii</i> S. et Z.	<i>Inugusu.</i>

Papaveraceæ.

<i>Chelidonium Majus</i> LINN.	<i>Kusanowō.</i>
<i>Corydalis decumbens</i> PERS.	<i>Yabu-engosaku.</i>

<i>Corydalis incisa</i> PERS.	<i>Murasakikeman.</i>
<i>Corydalis pallida</i> PERS.	<i>Miyamakikeman.</i>
<i>Macleya cordata</i> R. BR.	<i>Chanpagiku.</i>

Cruciferae.

<i>Arabis sagittata</i> DC.	<i>Yamahatazao.</i>
<i>Capsella bursa-pastoris</i> MENCH.	<i>Nazuna.</i>
<i>Cardamine hirsuta</i> LINN. var. <i>sylvatica</i> LINN.	<i>Tanetsukebana.</i>
<i>Dontstemon dentatus</i> BGE.	<i>Hanahatazao.</i>
<i>Draba nemorosa</i> LINN. var. <i>hebecarpa</i> LEDEB.	<i>Inunazuna.</i>
<i>Nasturtium montanum</i> WALL.	<i>Sukashitakabō.</i>
<i>Nasturtium palustre</i> DC.	<i>Inugarashi.</i>

Crassulaceae.

<i>Sedum Alfredi</i> HCE.	<i>Komochimannengusa.</i>
<i>Sedum Kantschiaticum</i> FISCH.	<i>Kirinsō.</i>
<i>Sedum lineare</i> THUNB.	<i>Onomannengusa.</i>
<i>Sedum Telephium</i> LINN. var. <i>purpureum.</i>	<i>Benkeisō.</i>
<i>Sedum verticillatum</i> LINN.	<i>Mitsubabenkeisō.</i>

Saxifragaceae.

<i>Astilbe chinensis</i> FR. et SAV. var. <i>japonica</i> MAXIM.	<i>Awamorishōma</i> or <i>Chidakesashi.</i>
<i>Astilbe Thunbergii</i> MIQ.	<i>Tori-ashishōma.</i>
<i>Deutzia gracilis</i> S. et Z.	<i>Hime-utsugi.</i>
<i>Deutzia scabra</i> THUNB.	<i>Utsugi.</i>
<i>Hydrangea hirta</i> S. et Z.	<i>Koajisai.</i>
<i>Hydrangea involucrata</i> SIEB.	<i>Tama-ajisai.</i>
<i>Hydrangea paniculata</i> SIEB.	<i>Norinoki.</i>
<i>Hydrangea scandens</i> MAXIM.	<i>Gotōzuru.</i>
<i>Parnassia palustris</i> LINN.	<i>Mumebachisō.</i>
<i>Ribes fasciculatum</i> S. et Z.	<i>Yabusanzashi.</i>
<i>Saxifraga cortusæfolia</i> S. et Z.	<i>Daimonjisō.</i>
<i>Saxifraga sarmentosa</i> LINN.	<i>Yukinoshita.</i>

Hamamelidaceæ.

Hamamelis japonica SIEB. et ZUCC. *Mansaku.*

Rosaceæ.

Agrimonia pilosa LEDEB.	<i>Kinmizuhiki.</i>
Aruncus silvester KOSTEL.	<i>Yamabukishōma.</i>
Cydonia japonica PERS. var. pyg- mæa MAXIM.	<i>Kusaboke.</i>
Fragaria indica ANDR.	<i>Hebiichigo.</i>
Geum japonicum THUNB.	<i>Daikonsō.</i>
Potentilla chinensis SER.	<i>Kawarasaiko.</i>
Potentilla cryptotænæa MAXIM.	<i>Mitsumoto.</i>
Potentilla Kleiniana W. et A.	<i>Ōebi-ichigo.</i>
Pourthiæa villosa DCNE.	<i>Kamatsuka.</i>
Pyrus Toringo SIEB.	<i>Zumi.</i>
Rosa multiflora THUNB.	<i>Noibara.</i>
Rosa Wichuraiana CREP.	<i>Terihanoibara.</i>
Rubus palmatus THUNB.	<i>Ki-ichigo.</i>
Rubus parvifolius LINN.	<i>Nawashiro-ichigo.</i>
Rubus phœnicolasius MAXIM.	<i>Urajiroichigo.</i>
Rubus rosæfolius SM.	<i>Baraichigo.</i>
Rubus Thunbergii S. et Z.	<i>Kusa-ichigo.</i>
Rubus trifidus THUNB.	<i>Kajiichigo.</i>
Sanguisorba officinalis LINN.	<i>Waremokō.</i>
Spiræa Blumei G. DON.	<i>Iwagasa.</i>
Spiræa bullata MAXIM.	<i>Koshimotsuke.</i>
Spiræa Thunbergii SIEB.	<i>Kogomebana.</i>
Stephanandra flexuosa S. et Z.	<i>Kogome-utsugi.</i>
Stephanandra Tanakæ FR. et SAV.	<i>Kana-utsugi.</i>
Ulmaria multijuga (MAXIM.)	<i>Shimotsukesō.</i>
Ulmaria purpurea (MAXIM.)	<i>Kioganoko.</i>

Leguminosæ.

Æschynomene indica LINN.	<i>Kusanemu.</i>
Albizzia Julibrissin BOIV.	<i>Nemunoki.</i>
Amphicarpæa Edgeworthii BENTH. var. japonica OLIV.	<i>Yabumame.</i>
Cæsalpinia sepiaria ROXB.	<i>Jakkei-ibara.</i>

<i>Cladrastis amurensis</i> B. et H. var.	
<i>floribunda</i> MAXIM.	<i>Inuenjū.</i>
<i>Crotalaria sessiliflora</i> LINN.	<i>Tinukimame.</i>
<i>Cassia mimosoides</i> LINN.	<i>Kawaraketsumei.</i>
<i>Desmodium Gardneri</i> BENTH.	<i>Obanusubito-hagi.</i>
<i>Desmodium laburnifolium</i> DC.	<i>Misonaoshi.</i>
<i>Desmodium Oldhami</i> Oliv.	<i>Fujikanzō.</i>
<i>Desmodium podocarpum</i> DC. var.	
<i>japonicum</i> MAXIM.	<i>Nusubito-hagi.</i>
<i>Desmodium polycarpum</i> DC.	<i>Kusakagi.</i>
<i>Indigofera pseudotinctoria</i> MATSUM.	<i>Komatsunagi.</i>
<i>Kraunhia floribunda</i> TAUB.	<i>Fuji.</i>
<i>Kraunhia japonica</i> TAUB.	<i>Natsufuji.</i>
<i>Lathyrus Davidii</i> HANCE.	<i>Itachisasage.</i>
<i>Lespedeza bicolor</i> TURCZ.	<i>Hiagi.</i>
<i>Lespedeza Buergeri</i> MIQ.	<i>Kihagi.</i>
<i>Lespedeza cyrtobotrya</i> MIQ.	<i>Marubahagi.</i>
<i>Lespedeza juncea</i> PERS. var. <i>sericea</i>	
HEMSL.	<i>Medohagi.</i>
<i>Lespedeza pilosa</i> S. et Z.	<i>Nekohagi.</i>
<i>Lespedeza stricta</i> H. et A.	<i>Yahazusō.</i>
<i>Lespedeza villosa</i> PERS.	<i>Inuhagi.</i>
<i>Lespedeza virgata</i> DC.	<i>Maki-ihagi.</i>
<i>Lotus corniculatus</i> LINN. var. <i>japonicus</i> RGL.	<i>Miyakogusa.</i>
<i>Pueraria Thunbergiana</i> BENTH.	<i>Kuzu.</i>
<i>Rhynchosia volubilis</i> LOUR.	<i>Tankirimame.</i>
<i>Sophora platycarpa</i> MAXIM.	<i>Yamayenjiū.</i>
<i>Vicia Cracca</i> LINN. var. <i>japonica</i> MIQ.	<i>Kusafuji.</i>
<i>Vicia pseudo-orobus</i> FISCH. et MEY.	<i>Obakusafuji.</i>
<i>Vicia sativa</i> LINN.	<i>Yahazu-endō.</i>
<i>Vicia unijuga</i> BR.	<i>Nanten-hagi.</i>

Geraniaceæ.

<i>Geranium nepalense</i> SWEET.	<i>Gennoshōko.</i>
<i>Geranium Sieboldi</i> MAXIM.	<i>Tachifūro.</i>
<i>Geranium Wilfordi</i> MAXIM.	<i>Kofūro.</i>

Oxalidaceæ.

<i>Oxalis corniculata</i> LINN.	<i>Katabami.</i>
---------------------------------	------------------

Linaceæ.

Linum possarioides PL. *Matsubanjin.*

Rutaceæ.

Bœninghausenia albiflora REICHE. *Arashigusa.*

Orixa japonica THUNB. *Kokusagi.*

Zanthoxylum pepitum DC. *Sanshū.*

Zanthoxylum schinnifolium S. et Z. *Inuzanshū.*

Simarubaceæ.

Picrasma quassioides BENX. *Nigaki.*

Polygalaceæ.

Polygala japonica HOUTT. *Himehagi.*

Euphorbiaceæ.

Acalypha australis LINN. *Inokigusa.*

Euphorbia adenochlora MORR. et
DCNE. *No-urushi.*

Euphorbia humifusa WILLD. *Nshikigusa.*

Euphorbia helioscopia LINN. *Tōdaigusa.*

Euphorbia pekinensis RUPR. *Tukatōdai.*

Euphorbia Sieboldiana MORR. et
DCNE. *Natsutōdai.*

Phyllanthus Matsumuræ HAYATA. *Himemikansō.*

Phyllanthus Urinaria LINN. *Komikansō.*

Securinega fluggeoides MUELL ARG. *Hitotsubahagi.*

Callitrichaceæ.

Callitriche japonica ENGEL. *Awagoke.*

Coriariaceæ.

Coriaria japonica A. GR. *Dokuutsugi.*

Buxaceæ.

Buxus sempervirens var. *japonica*
MAKINO. *Asamatsuge.*

Anacardiaceæ.

<i>Rhus semi-alata</i> MURR. var. <i>Osbeckii</i> DC.	<i>Fushinoki.</i>
<i>Rhus silvestris</i> S. et Z.	<i>Yamahaze.</i>
<i>Rhus Toxicodendron</i> LINN. var. <i>radicans</i> MIQ.	<i>Tsutaursushi.</i>
<i>Rhus trichocarpa</i> MIQ.	<i>Yamaursushi.</i>

Aquifoliaceæ.

<i>Ilex crenata</i> THUNB.	<i>Inutsuge.</i>
<i>Ilex integra</i> THUNB.	<i>Mochinoki.</i>
<i>Ilex rugosa</i> FR. SCHM.	<i>Tsurutsuge.</i>
<i>Ilex Sieboldi</i> MIQ.	<i>Unemodoki.</i>

Celastraceæ.

<i>Celastrus articulatus</i> THUNB.	<i>Tsuru-unemodoki.</i>
<i>Euonymus alata</i> K. KOCH.	<i>Nishikigi.</i>
<i>Euonymus europæa</i> L. var. <i>Hamil-toniana</i> MAXIM.	<i>Mayumi.</i>
<i>Euonymus japonica</i> THUNB.	<i>Masaki.</i>
<i>Euonymus oxyphylla</i> MIQ.	<i>Tsuribana.</i>
<i>Tripterigium Wilfordi</i> HOOK.	<i>Kurozumi.</i>

Staphyleaceæ.

<i>Euscaphis staphyleoides</i> S. et Z.	<i>Gonzui.</i>
---	----------------

Balsaminaceæ.

<i>Impatiens nolitangere</i> LINN.	<i>Kitsurifunesō.</i>
<i>Impatiens Textori</i> MIQ.	<i>Tsurifunesō.</i>

Sabiaceæ.

<i>Meliosma myriantha</i> S. et Z.	<i>Awabuki.</i>
<i>Meliosma tenuis</i> MAXIM.	<i>Miyamahōso.</i>

Rhamnaceæ.

<i>Berchemia racemosa</i> S. et Z.	<i>Kumayanagi.</i>
<i>Rhamnus japonica</i> MAXIM. var. <i>ge-nuina</i> MAXIM.	<i>Karoumemodoki.</i>

Vitaceæ.

<i>Vitis flexuosa</i> THUNB.	<i>Sankakuzuru.</i>
<i>Vitis heterophylla</i> THUNB.	<i>Nobudō.</i>
<i>Vitis inconstans</i> MIQ.	<i>Tsuta.</i>
<i>Vitis Thunbergii</i> S. et Z.	<i>Yebidsuru.</i>
<i>Vitis coignetiae</i> PULL.	<i>Kanebu.</i>

Theaceæ.

<i>Thea japonica</i> NOIS.	<i>Tsubaki.</i>
<i>Eurya japonica</i> THUNB.	<i>Hisakaki.</i>

Stachyuraceæ.

<i>Stachyurus præcox</i> S. et Z.	<i>Kifuji.</i>
-----------------------------------	----------------

Guttiferæ.

<i>Hypericum Ascyron</i> LINN.	<i>Tomoisō.</i>
<i>Hypericum chinense</i> LINN.	<i>Bijōyanagi.</i>
<i>Hypericum erectum</i> THUNB.	<i>Otogirisō.</i>
<i>Hypericum erectum</i> THUNB. var. <i>cæspitosum</i> MAKINO.	<i>Fuji-otogiri.</i>
<i>Hypericum japonicum</i> THUNB.	<i>Hime-otogiri.</i>
<i>Hypericum patulum</i> THUNB.	<i>Kinshibai.</i>

Elatinaceæ.

<i>Elatine orientalis</i> MAKINO.	<i>Mizuhakobe.</i>
-----------------------------------	--------------------

Violaceæ.

<i>Viola japonica</i> LANGSD.	<i>Kosumire.</i>
<i>Viola Patrinii</i> DC. var. <i>chinensis</i> GING.	<i>Sumire.</i>
<i>Viola silvestris</i> KIT. var. <i>grypoceras</i> A. GR.	<i>Tachitsubosumire.</i>

Thymelæaceæ.

<i>Daphne pseudo-mezereum</i> A. GR.	<i>Onishibari.</i>
<i>Wikstroemia Ganpi</i> MAXIM.	<i>Kozampi.</i>

Elæagnaceæ.

<i>Elæagnus glabra</i> THUNB.	<i>Tsurugumi.</i>
<i>Elæagnus macrophylla</i> THUNB.	<i>Marubagumi.</i>
<i>Elæagnus umbellata</i> THUNB.	<i>Akigumi.</i>

Lythraceæ.

<i>Lythrum Salicaria</i> LINN.	<i>Misohagi.</i>
<i>Rotala indica</i> KÖHNE var. <i>uliginosa</i>
Miq.	<i>Kikashigusa.</i>

Onagraceæ.

<i>Circæa quadrisulcata</i> MAXIM.	<i>Mizutamasō.</i>
<i>Epilobium angustifolium</i> LINN.	<i>Yanagisō.</i>
<i>Ludwigia prostrata</i> ROXB.	<i>Tagobō.</i>

Halorrhagidaceæ.

<i>Haloragis micrantha</i> R. BR.	<i>Arinotōgusa.</i>

Fig. 18. *Vitis Coignetiae* PULL. and *Angelica polyclada* FRANCH., Yoshida side, about 1200 m. above the sea. (From the Atlas of Japanese Vegetation, by permission of Prof. M. MIYOSHI.)

Araliaceæ.

<i>Aralia cordata</i> THUNB.	<i>Udo.</i>
<i>Aralia sinensis</i> LINN.	<i>Turanoki.</i>
<i>Acanthopanax divaricatum</i> S. et Z.	<i>Takanotsune.</i>
<i>Acanthopanax ricinifolium</i> S. et Z.	<i>Harigiri.</i>
<i>Acanthopanax spinosum</i> MIQ.	<i>Ukogi.</i>
<i>Hedera Helix</i> var. <i>colchica</i> C. KOCH.	<i>Kizuta.</i>
<i>Helwingia japonica</i> DIETR.	<i>Hanaikada.</i>

Umbelliferæ.

<i>Angelica Miqueliana</i> MAXIM.	<i>Yamazeri</i> or <i>Serimodoki.</i>
<i>Angelica polyclada</i> FRANCH.	<i>Shishiundo.</i>
<i>Bupleurum falcatum</i> LINN.	<i>Mishimasaiko.</i>
<i>Bupleurum sachalinense</i> FR. et SAV.	<i>Marubasaiko.</i>
<i>Chamæle tenera</i> MIQ.	<i>Sentôsô.</i>
<i>Cicuta virosa</i> LINN.	<i>Dokuzeri.</i>
<i>Cryptotænia japonica</i> HASSK.	<i>Mitsuba.</i>
<i>Hydrocotyle asiatica</i> LINN.	<i>Tsubokusa.</i>
<i>Hydrocotyle rotundifolia</i> ROXB.	<i>Chidomegusa.</i>
<i>Ligusticum acutilobum</i> S. et Z.	<i>Tôki.</i>
<i>Ænanthe stolonifera</i> DC.	<i>Seri.</i>
<i>Osmorhiza japonica</i> S. et Z.	<i>Yabuminjin.</i>
<i>Pimpinella calycina</i> MAXIM.	<i>Dakezeri</i> or <i>Tukanotsune.</i>
<i>Torilis japonica</i> DC.	<i>Yabujirami.</i>

Cornaceæ.

<i>Aucuba japonica</i> THUNB.	<i>Aoki.</i>
<i>Cornus ignorata</i> C. KOCH.	<i>Kumanomizuki.</i>
<i>Cornus macrophylla</i> WALL.	<i>Mizuki.</i>
<i>Marlea platanifolia</i> S. et Z.	<i>Urinoki.</i>

Clethraceæ.

<i>Clethra barbiuervis</i> S. et Z.	<i>Riöbu.</i>
-------------------------------------	---------------

Ericaceæ.

<i>Enkianthus nikœnsis</i> MAKINO.	<i>Aburatsutsuji.</i>
<i>Enkianthus japonicus</i> HOOK.	<i>Dôdantsutsuji.</i>
<i>Pieris japonica</i> D. DON.	<i>Asebi.</i>
<i>Pieris ovalifolia</i> D. DON.	<i>Kashioshimi.</i>
<i>Rhododendron indicum</i> SW. var. α	
<i>Kæmpferi</i> MAXIM.	<i>Yamatsutsuji.</i>
<i>Rhododendron sinense</i> SW.	<i>Rengetsutsuji.</i>
<i>Vaccinium bracteatum</i> THUNB.	<i>Shashampo.</i>
<i>Vaccinium Buergeri</i> MIQ.	<i>Ôbasunoki.</i>
<i>Vaccinium hirtum</i> THUNB.	<i>Sunoki</i> or <i>Usunoki.</i>

Myrsinaceæ.

<i>Ardisia japonica</i> BL.	<i>Yabukôji.</i>
-----------------------------	------------------

Primulaceæ.

<i>Lysimachia clethroides</i> DUBY.	<i>Torano-o</i> or <i>Okatorano-o.</i>
<i>Lysimachia decurrens</i> G. FORST.	<i>Miyamatagobô.</i>
<i>Lysimachia Fortunei</i> MAXIM.	<i>Numatorano-o.</i>
<i>Lysimachia japonica</i> THUNB.	<i>Konasubi.</i>
<i>Lysimachia vulgaris</i> LINN.	<i>Kusaredama.</i>

Styracaceæ.

<i>Styrax japonica</i> S. et Z.	<i>Egonoki.</i>
---------------------------------	-----------------

Symplocaceæ.

<i>Symplocos cratægoides</i> HAM.	<i>Sawafutagi.</i>
-----------------------------------	--------------------

Oleaceæ.

<i>Fraxinus Bungeana</i> DC. var. <i>pubi-</i> <i>nervis</i> WG.	<i>Toneriko.</i>
<i>Fraxinus longicuspis</i> S. et Z.	<i>Aotago.</i>
<i>Ligustrum japonicum</i> THUNB.	<i>Nezumimochi.</i>
<i>Osmanthus Aquifolium</i> B. et H.	<i>Hiragi.</i>

Loganiaceæ.

<i>Mitrasacme polymorpha</i> R. BR.	<i>Ainai.</i>
-------------------------------------	---------------

Gentianaceæ.

Crawfurdia fasciculata WALL.	<i>Tsururindō.</i>
Gentiana scabra BGE. var. Buergeri MAXIM.	<i>Rindō.</i>
Halenia sibirica BORK.	<i>Hana-ikari.</i>
Swertia bimaculata CLARK.	<i>Akebonosō.</i>
Swertia chinensis FRANCH.	<i>Senburi.</i>

Apocynaceæ.

Trachelospermum jasminoides LEM.	<i>Teikakazura.</i>
----------------------------------	---------------------

Asclepiadaceæ.

Cynanchum caudatum MAX.	<i>Ikema.</i>
Pycnostelma chinensis BGE.	<i>Suzusaiko.</i>
Tylophora nikiensis MATSUM.	<i>Ko-kamomezuru.</i>

Convolvulaceæ.

Calystegia sepium R. BR.	<i>Hirugao.</i>
Cuscuta japonica CHOIS. var. thyr- soidea ENGELM.	<i>Nenashikazura.</i>

Borraginaceæ.

Cynoglossum furcatum WALL.	<i>Orurisō.</i>
Trigonotis peduncularis BENTH.	<i>Tabirako.</i>

Verbenaceæ.

Callicarpa japonica THUNB.	<i>Murasakishikibu.</i>
Clerodendron trichotomum THUNB.	<i>Kusagi.</i>
Phryma leptostachya LINN.	<i>Hachokusō.</i>

Labiataæ.

Ajuga decumbens THUNB.	<i>Junihitoe.</i>
Calamintha chinensis BENTH.	<i>Kurumabana.</i>
Calamintha gracilis BENTH.	<i>Tōbana.</i>
Comanthosphace sublanceolata S. MOORE.	<i>Fujitenninsō.</i>
Elsholtzia cristata WILLD.	<i>Naginatakōjū.</i>

<i>Lamium album</i> LINN.	<i>Odorikosō.</i>
<i>Leonurus macranthus</i> MAXIM.	<i>Kisewata.</i>
<i>Mentha arvensis</i> LINN. var. <i>piperscens</i> HOLMES.	<i>Megusa.</i>
<i>Mosla grosseserrata</i> MAXIM.	<i>Mizokōjū</i> or <i>himijiso.</i>
<i>Mosla punctata</i> MAXIM.	<i>Inukōjū.</i>
<i>Plectranthus glaucocalyx</i> MAXIM. var. <i>japonicus</i> MAXIM.	<i>Hikiokoshi.</i>
<i>Plectranthus inflexus</i> VAHL.	<i>Yamahakka.</i>
<i>Plectranthus longitubus</i> MIQ.	<i>Akichōji.</i>
<i>Prunella vulgaris</i> LINN.	<i>Usubogusa.</i>
<i>Salvia japonica</i> THUNB. var. <i>bipinnata</i> FR. et SAV.	<i>Akinotamurasō.</i>
<i>Salvia nipponica</i> MIQ.	<i>Kotojisō.</i>
<i>Scutellaria indica</i> LINN. var. <i>japonica</i> MAXIM.	<i>Tatsunamisō.</i>
<i>Stachys aspera</i> MICHX. var. <i>japonica</i> MAXIM.	<i>Chorogidamashi.</i>
<i>Teucrium japonicum</i> WILLD.	<i>Nigakusa.</i>
<i>Teucrium stoloniferum</i> HAM. var. <i>Miquelianum</i> MAXIM.	<i>Tsurunigakusa.</i>

Solanaceæ.

<i>Lycium chinense</i> MILL.	<i>Kuko.</i>
------------------------------	--------------

Scrophulariaceæ.

<i>Euphrasia officinalis</i> LINN. var. <i>vulgaris</i> LEDEB.	<i>Kogomegusa.</i>
<i>Lindernia angustifolia</i> WETTS.	<i>Azetōgarashi.</i>
<i>Lindernia crustacea</i> (L.) F. v. M.	<i>Urikusa.</i>
<i>Lindernia pyxidaria</i> ALL.	<i>Azena.</i>
<i>Mazus rugosus</i> LOUR. var. <i>macranthus</i> F. et S.	<i>Sagigoke.</i>
<i>Mimulus nepalensis</i> BENTH.	<i>Mizohōzuki.</i>
<i>Pedicularis gloriosa</i> BISS. et MOOR.	<i>Hankwai-azami.</i>
<i>Pedicularis resupinata</i> LINN.	<i>Shiogamagiku.</i>
<i>Phteirospermum chinense</i> BGE.	<i>Koshiogama.</i>
<i>Scrophularia Patriniana</i> WYDL.	<i>Himeno-usutsubo.</i>
<i>Siphonostegia chinensis</i> BENTH.	<i>Hikiyomogi.</i>

Pl. VI. View of Mt. Fuji, seen from the Pacific coast; the coniferous, deciduous tree, and prairie, regions are visible. (After a photograph; a little brush is put in by the author in order to make the plant regions clear.)

<i>Veronica cana</i> WALL.	<i>Kuwagatasō.</i>
<i>Veronica longifolia</i> LINN.	<i>Ruritorano-o.</i>
<i>Veronica spuria</i> LINN.	<i>Yamatorano-o.</i>
<i>Veronica virginica</i> LINN.	<i>Kukaisō.</i>

Orobanchaceæ.

<i>Æginetia indica</i> ROXB.	<i>Nanbangiseru.</i>
------------------------------	----------------------

Gesneraceæ.

<i>Conandron ramondiioides</i> S. et Z.	<i>Iwatabako.</i>
---	-------------------

Acanthaceæ.

<i>Dicliptera crinita</i> NEES.	<i>Hagurosō.</i>
<i>Hygrophila lancea</i> MIQ.	<i>Oginotsune.</i>
<i>Justicia procumbens</i> LINN.	<i>Kitsunenomago.</i>

Plantaginaceæ.

<i>Plantago major</i> LINN. var. <i>asiatica</i> DCNE.	<i>Obako.</i>
---	---------------

Rubiaceæ.

<i>Galium aparine</i> LINN.	<i>Yayemugura.</i>
<i>Galium asprellum</i> MICHX.	<i>Hosoba-ya-emugura.</i>
<i>Galium brachypodium</i> MAXIM.	<i>Maruba-yotsubamugura</i>
<i>Galium gracile</i> BGE.	<i>Yotsubamugura.</i>
<i>Galium verum</i> LINN. var. <i>lacteum</i> . MAXIM.	<i>Kawaramatsuba.</i>
<i>Galium trifidum</i> LINN.	<i>Hosobayotsubamugura.</i>
<i>Gardenia florida</i> LINN.	<i>Kuchinashi.</i>
<i>Oldenlandia diffusa</i> ROXB.	<i>Futabamugura.</i>
<i>Oldenlandia hirsuta</i> LINN. f.	<i>Hashikagusa.</i>
<i>Pæderia tomentosa</i> BL.	<i>Hekusokazura.</i>
<i>Serissa foetida</i> COMM.	<i>Itakuchōge.</i>

Caprifoliaceæ.

<i>Abelia spathulata</i> S. et Z.	<i>Tsukubane-utsugi.</i>
<i>Diervilla floribunda</i> S. et Z.	<i>Nshikiutsugi.</i>
<i>Diervilla grandiflora</i> S. et Z.	<i>Itakone-utsugi.</i>

<i>Diervilla japonica</i> DC.	<i>Beni-utsugi</i> or <i>Tani-utsugi</i> .
<i>Diervilla Middendorffiana</i> CARR.	<i>Ukon-utsugi</i> .
<i>Lonicera gracilipes</i> MIQ.	<i>Uguisukagura</i> .
<i>Lonicera japonica</i> THUNB.	<i>Suikazura</i> .
<i>Lonicera Morrowii</i> A. GR.	<i>Kinginboku</i> or <i>Iyōtanboku</i> .
<i>Sambucus javanica</i> BL.	<i>Sokuzu</i> .
<i>Sambucus racemosa</i> LINN.	<i>Nivatoko</i> .
<i>Viburnum dilatatum</i> THUNB.	<i>Gamazumi</i> .
<i>Viburnum erosum</i> THUNB.	<i>Kobanogamazumi</i> .
<i>Viburnum Sieboldi</i> MIQ.	<i>Gomaki</i> .

Valerianaceæ.

<i>Patrinia scabiosæfolia</i> LINK.	<i>Omina-eshi</i> .
<i>Patrinia villosa</i> JUSS.	<i>Otoko-eshi</i> .

Dipsaceæ.

<i>Scabiosa japonica</i> MIQ.	<i>Matsumushisō</i> .
-------------------------------	-----------------------

Cucurbitaceæ.

<i>Actinostemma racemosum</i> MAXIM.	<i>Gokizuru</i> .
<i>Trichosanthes cucumeroides</i> MAXIM.	<i>Karasu-uri</i> .

Campanulaceæ.

<i>Adenophora verticillata</i> FISCH. var. <i>verticillata</i> FR. et SAV.	<i>Tsuriganeninjin</i> .
<i>Campanula punctata</i> LAM.	<i>Hotarubukuro</i> .
<i>Codonopsis lanceolata</i> B. et H.	<i>Tsuruninjin</i> .
<i>Platycodon grandiflorus</i> DC.	<i>Kikyō</i> .
<i>Phyteuma japonicum</i> MIQ.	<i>Shideshajin</i> .

Compositæ.

<i>Achillea ptarmicoides</i> MAXIM.	<i>Shirobananokogirisō</i> .
<i>Adenocaulon bicolor</i> HOOK.	<i>Nobuki</i> .
<i>Ainsliæ apiculata</i> SCH. BIP.	<i>Kikkōhaguma</i> .
<i>Ainsliæ acerifolia</i> SCH. BIP.	<i>Momijihaguma</i> .
<i>Anaphalis margaritacea</i> B. et H.	<i>Yamahahako</i> .
<i>Anaphalis yedensis</i> FR. et SAV.	<i>Kawarahahako</i> .
<i>Artemisia capillaris</i> THUNB.	<i>Kawarayomogi</i> .

<i>Artemisia japonica</i> THUNB.	<i>Otokoyomogi.</i>
<i>Artemisia Keiskeana</i> MIQ.	<i>Inuyomogi.</i>
<i>Artemisia vulgaris</i> LINN. var. <i>indica</i> MAXIM.	<i>Yomogi.</i>
<i>Aster fastigiatus</i> FISCH. et MEX.	<i>Himeshion.</i>
<i>Aster indicus</i> LINN.	<i>Yomena.</i>
<i>Aster indicus</i> LINN. var. <i>pinnatifidus</i> MAXIM.	<i>Yūgagiku.</i>
<i>Aster scaber</i> THUNB.	<i>Shirayamagiku.</i>
<i>Aster trinervius</i> ROXB.	<i>Yamashirogiku.</i>
<i>Atractylis ovata</i> THUNB.	<i>Okera.</i>
<i>Bidens pilosa</i> LINN.	<i>Sendangusa.</i>
<i>Bidens tripartita</i> LINN.	<i>Ta-ukogi.</i>
<i>Cacalia Krameri</i> (FR. et SAV.)	<i>Yaburegasu.</i>
<i>Carpesium abrotanoides</i> LINN.	<i>Yabutabako.</i>
<i>Carpesium triste</i> MAXIM.	<i>Gankubiyabutabako.</i>
<i>Chrysanthemum sinense</i> var. <i>japo-</i> <i>nicum</i> MAXIM.	<i>Rinnōgiku.</i>
<i>Centaurea atriplicifolia</i> (DC.)	<i>Yamabokuchi.</i>
<i>Cirsium japonicum</i> DC.	<i>No-azami.</i>
<i>Cirsium spicatum</i> (MAXIM.)	<i>Yama-azami.</i>
<i>Eclipta alba</i> HASSK.	<i>Takasaburō.</i>
<i>Erigeron annuus</i> PERS.	<i>Himejion.</i>
<i>Erigeron canadensis</i> LINN.	<i>Himemukashiyomogi.</i>
<i>Eupatorium japonicum</i> THUNB.	<i>Hiyodoribana.</i>
<i>Eupatorium Kirilowii</i> TURCZ.	<i>Sawahiyodori.</i>
<i>Gerbera Anandria</i> SCH. BIP.	<i>Senboniyari.</i>
<i>Gnaphalium japonicum</i> THUNB.	<i>Chichikogusa.</i>
<i>Gnaphalium multiceps</i> WALL.	<i>Hahakogusa.</i>
<i>Inula britannica</i> DC.	<i>Oguruma.</i>
<i>Inula salicina</i> LINN.	<i>Kasensō.</i>
<i>Lactuca debilis</i> MAXIM.	<i>Jishibari.</i>
<i>Lactuca denticulata</i> MAXIM.	<i>Yakushisō.</i>
<i>Lactuca stolonifera</i> BENTH.	<i>Iwanigana.</i>
<i>Lactuca Thunbergiana</i> MAXIM.	<i>Nigana.</i>
<i>Lactuca versicolor</i> SCH. BIP.	<i>Takasagosō.</i>
<i>Ligularia elivorum</i> MAXIM.	<i>Dakebuki.</i>
<i>Ligularia sibirica</i> CASS.	<i>Otakarakō.</i>
<i>Macroclidium robustum</i> MAXIM.	<i>Kashiwabahaguma.</i>
<i>Pertya scandens</i> SCH. BIP.	<i>Nagabano-kōyabōki.</i>

Picris hieracioides LINN. var. japonica RGL.	Kōzorina.
Saussurea Maximowiczii HERD.	Miyako-azami.
Senecio campestris DC.	Sawa-oguruma.
Senecio flammeus DC.	Kōrinkwa.
Senecio nemorensis LINN. var. Fuchsii KOCH.	Kion.
Serratula coronata LINN.	Tanurasō.
Solidago Virga-aurea LINN.	Akinokirinsō.

b) Deciduous broad leaved tree-Region.

Next to the prairie-region, comes the deciduous tree-region. As we have stated above, it is most abundantly formed on the south side, while it is the least so on the north, in consequence of the climatic characters which we have already alluded to. Under this heading, therefore, it is sufficient for us to consider its southern formation.

Fig. 19. *Fagus* and *Quercus* in the winter.

The lower boundary of this region lies, on the south-east, at an altitude of 870 m; but on the due south side, it climbs as high as 1400 m., and on some parts it comes up even to altitudes of 1600 m.—1700 m. On the due south where this formation attains its most luxuriant growth, we see the gigantic forest principally formed of *Fagus*, *Zelkova* and *Quercus*, with thick under-growth, the ground being sufficiently capable of holding water. The tall *Fagus* with climbing *Rhus* and *Euonymus* reaching to the top of the tree, the *Stewartia* bare of

Fig. 20. *Fagus* and *Quercus* in the summer.

bark, the thick foliage of *Quercus* and various shrubs forming a jungle over the under-growth,—all collectively form the most beautiful forest on this side.

A glance over this formation at the height of 1900 m. on the southern flank, will give us a general idea of it. The following description is drawn up from observations at the spot where I crossed in the summer of 1907. Over a little open space with the promontory of the Conifer-forest on the left side (east), the deciduous forest stretches away to the south upon a slope, so gentle that it seems at a distance perfectly flat, and the light green color of its foliage, in contrast with the dark green of the Conifers above, makes the forest still more remarkable. Far away, the prairie formation extends till it melts into the rice fields some miles distant. The several aspects of the vegetation of this region on the different sides are given in the accompanying photographs.

The forest-flora of this region is as follows:—

Juglandaceæ.

<i>Juglans Sieboldiana</i> MAXIM.	<i>Kurumi.</i>
<i>Pterocarya rhoifolia</i> S. et Z.	<i>Sawagurumi.</i>

Betulaceæ.

<i>Betula alba</i> LINN. var. <i>vulgaris</i> DC.	<i>Shirakanba.</i>
<i>Betula Ermanni</i> CHAM. var. <i>nipponica</i> MAXIM.	<i>Takekamba.</i>

Fagaceæ.

<i>Fagus sylvatica</i> LINN. var. <i>Sieboldi</i> MAXIM.	<i>Bunanoki.</i>
<i>Quercus aliena</i> BLUME.	<i>Naragashiva.</i>
<i>Quercus crispula</i> BLUME.	<i>Onara.</i>

Magnoliaceæ.

<i>Magnolia hypoleuca</i> S. et Z.	<i>Hōnoki.</i>
------------------------------------	----------------

Trochodendraceæ.

Cercidiphyllum japonicum S. et Z. *Katsura*.

Rosaceæ.

Pyrus aucuparia GÆRTN. var. *japonica* MAXIM. *Miyamananakamudo*.

Rutaceæ.

Phellodendron amurense RUPR. *Kikada*.

Aceraceæ.

Acer pictum THUNB. *Tsutanomiji*.
Acer palmatum THUNB. *Yamamomiji*.

Sabiaceæ.

Meliosma myriantha S. et Z. *Awabuki*.

Tiliaceæ.

Tilia cordata MIQ. *Shinanoki*.

Theaceæ.

Stewartia monadelphæ S. et Z. *Himeshara*.
Stewartia pseudocamellia MAXIM. *Natsutsubaki*.

Cornaceæ.

Cornus ignorata C. KOCH. *Kumanomizuki*.
Cornus Kousa BUERG. *Yamabōshi*.
Cornus macrophylla WALL. *Mizuki*.

Clethraceæ.

Clethra barbinervis S. et Z. *Ryōbu*.

Oleaceæ.

Fraxinus longicuspis S. et Z. *Ao-tago*.
Fraxinus Sieboldiana BLUME. *Shiōji*.

Caprifoliaceæ.

Viburnum furcatum BLUME.

Mushikari or *Obakamenoki*.

Fig. 21. Forest of broad leaved trees and Conifers, on the Yoshida side, about 1800 m. above the sea. *Viburnum furcatum* BLUME on the left, *Pyrus aucuparia* GÆRTN. var. *japonica* MAXIM. on the right, and backwards *Picea polita* CARR. *Abies brachyphylla* MAXIM. are more or less distinctly visible. (From the Atlas of Japanese Vegetation, by permission of Prof. M. MIYOSHI.)

The under-growth of the forests, shrubs and herbs found in the open places are as follows :—

Polypodiaceæ.

Adiantum pedatum LINN.

Kujakushida.

Aspidium muticum FR. et SAV.

Miyamakanavarabi.

<i>Athyrium nipponicum</i> BAKER.	<i>Inuwarabi.</i>
<i>Microlepia Wilfordi</i> MOORE.	<i>Ōrensishida.</i>
<i>Nephrodium dilatatum</i> DESV.	<i>Shiranewarabi.</i>
<i>Nephrodium Filix-mas</i> RICH.	<i>Memma, Oshida, Miyama-inode.</i>
<i>Nephrodium Filix-mass</i> RICH. var. <i>lacerum</i> CHRIST.	<i>Kumawarabi.</i>
<i>Nephrodium Filix-mass.</i> RICH. var. <i>polylepis</i> MAKINO.	<i>Miyamakumawarabi.</i>
<i>Nephrodium Filix-mas</i> Rich. var. <i>Sabæi</i> CHRIST.	<i>Miyama-itachishida.</i>
<i>Nephrodium glanduligerum</i> MAKINO.	<i>Hashigoshida.</i>
<i>Nephrodium hirtipes</i> HOOK.	<i>Iwahego.</i>
<i>Nephrodium Maximowiczii</i> BAKER.	<i>Nantaishida.</i>
<i>Nephrodium Phegopteris</i> BAUMG.	<i>Miyamawarabi.</i>
<i>Nephrodium thelypteris</i> DESV.	<i>Himeshida or Shorima.</i>
<i>Polypodium lineare</i> THUNB. var. <i>distans</i> MAKINO.	<i>Miyamanokishinobu.</i>
<i>Polypodium lineare</i> THUNB. var. <i>Onœi</i> MAKINO.	<i>Himenokishinobu.</i>
<i>Polypodium senanense</i> MAXIM.	<i>Miyama-uraboshi.</i>
<i>Polypodium trichomanoides</i> SW.	<i>Ōkuboshida.</i>
<i>Polystichum tripterum</i> SM.	<i>Shūmokushida.</i>

Hymenophyllaceæ.

<i>Hymenophyllum Wrightii</i> BOSCH.	<i>Kokeshinobu.</i>
<i>Trichomanes parvulum</i> POIR.	<i>Uchiwagoke.</i>

Osmundaceæ.

<i>Osmunda cinnamomea</i> LINN.	<i>Yamadorizenmai.</i>
---------------------------------	------------------------

Lycopodiaceæ.

<i>Lycopodium cryptomerinum</i> MAXIM.	<i>Sugiran or Iwasugi.</i>
<i>Lycopodium serratum</i> THUNB.	<i>Tōgeshiba.</i>

Gramineæ.

<i>Agrostis canina</i> LINN.	<i>Takanenukabo.</i>
<i>Agrostis flaccida</i> HACK.	<i>Miyamanukabo.</i>
<i>Agrostis perennans</i> TUCK.	<i>Yamanukabo.</i>
<i>Brylkinia caudata</i> FR. SCHM.	<i>Hajigaeri.</i>

<i>Calamagrostis sachalinensis</i> FR. SCHM.	<i>Himenogariyasu.</i>
<i>Calamagrostis villosa</i> MUT. var.	
<i>Langsdorffii</i> HACK.	<i>Iwagariyasu.</i>
<i>Festuca ovina</i> LINN.	<i>Ushinokegusa.</i>
<i>Milium effusum</i> LINN.	<i>Ibukinukabo.</i>
<i>Miscanthus Matsumure</i> HACK.	<i>Kariyasumodoki.</i>
<i>Muehlenbergia japonica</i> STEUD. var.	
<i>hakonensis</i> HACK.	<i>Tachinezumigaya.</i>
<i>Sasa albo-marginata</i> MAK. et SHIB.	<i>Kumazasa.</i>
<i>Sasa paniculata</i> MAK. et SHIB.	<i>Nemagaridake.</i>

Cyperaceæ.

<i>Carex Dœnitzii</i> BOECK.	<i>Kotanukiran.</i>
<i>Carex pseudo-conica</i> F. et S.	<i>O-itosuge.</i>
<i>Carex remota</i> LINN.	<i>Yabusuge.</i>

Juncaceæ.

<i>Juncus Maximowiczii</i> BUCH.	<i>Ito-inunohige.</i>
<i>Luzula campestris</i> DC. var. <i>sudetica</i>	
CLARK.	<i>Miyamasuzumenohie.</i>

Stemonaceæ.

<i>Croomia japonica</i> MIQ.	<i>Nabewari.</i>
------------------------------	------------------

Liliaceæ.

<i>Hosta cœrulea</i> TRATT. var. <i>lancifolia</i>	
MATSUM.	<i>Mizugibōshi.</i>
<i>Hosta Sieboldiana</i> ENGL. var. <i>longipes</i> MATSUM.	<i>Iwagibōshi.</i>
<i>Lilium avenaceum</i> FISCH.	<i>Takeshimayuri.</i>
<i>Lilium Hansonii</i> BAK.	<i>Kurumayuri.</i>
<i>Maianthemum Convallaria</i> WIGG. et	
ROTH.	<i>Maizurusō.</i>
<i>Metanarthecium foliatum</i> MAXIM.	<i>Nebarinogiran.</i>
<i>Paris quadrifolia</i> LINN. var. <i>obovata</i> RGL.	
<i>Paris tetraphylla</i> A. GR.	<i>Kurumabatsukubanesō.</i>
<i>Polygonatum latifolium</i> DESF. var.	<i>Tsukubanesō.</i>
<i>commutatum</i> BAK.	<i>Narukoyuri.</i>

Polygonatum officinale ALL.	<i>Amadokoro.</i>
Streptopus ajanensis TILING. var. japonica MAXIM.	<i>Takeshimaran.</i>
Trillium Smallii MAXIM.	<i>Eureisō.</i>
Veratrum album LINN. var. Lobelia- num BAK. f. japonica BAK.	<i>Baikeisō.</i>

Orchidaceæ.

Cypripedium debile REICH. f.	<i>Ko-atsumorisō.</i>
Ephiphianthus sachalinensis REICH. f.	<i>Ko-ichiyōran.</i>
Epipactis papillosa F. et S.	<i>Yezosuzuran.</i>
Gymnadenia cucullata REICH. f.	<i>Miyamamojizuri.</i>
Gymnadenia cyclochia KORS.	<i>Kamomoran</i> or <i>Ichiyōchidori.</i>
Liparis Krameri F. et S.	<i>Jigabachisō.</i>
Listera cordata R. BR.	<i>Kofutabaran.</i>
Listera Eschscholtziana CHAM.	<i>Futabaran.</i>
Platanthera angustifolia REICH. f.	<i>Mukagosō.</i>
Platanthera mandarinorum REICH. f.	<i>Yamasagisō.</i>
Platanthera ophryoides FR. SCHM.	<i>Kisochidori.</i>
Platanthera Yatabei MAXIM.	<i>Hosobanokisochidori.</i>
Oberonia japonica MAXIM.	<i>Yōrakuran.</i>
Orchis Chondradenia MAKINO.	<i>Onoiran.</i>
Yuania japonica MAXIM.	<i>Shōkiran.</i>

Salicaceæ.

Salix gracilistyla MIQ.	<i>Nekoyanagi.</i>
-------------------------	--------------------

Betulaceæ.

Corylus rostrata AIT. var. Sieboldiana MAXIM.	<i>Tsunohashibami.</i>
--	------------------------

Urticaceæ.

Laportea bulbifera WEDD.	<i>Mukago-irakusa.</i>
--------------------------	------------------------

Caryophyllaceæ.

Stellaria nemorum LINN. var. Bun- geana MAXIM.	<i>Miyamahakobe.</i>
---	----------------------

Ranunculaceæ.

<i>Aconitum Fischeri</i> REICH.	<i>Torikabuto.</i>
<i>Aconitum lycoctonum</i> LINN.	<i>Reizinsō.</i>
<i>Aquilegia glandulosa</i> FISCH.	<i>Yamaodamaki.</i>
<i>Cimicifuga biternata</i> MIQ.	<i>Miyamashōma.</i>
<i>Cimicifuga japonica</i> SPR.	<i>Mitsubashōma.</i>
<i>Clematis alpina</i> MILL.	<i>Miyamahanshōzuru.</i>
<i>Coptis brachypetala</i> S. et Z. var. <i>pygmæa</i> MIQ.	<i>Hosoba-ōren.</i>
<i>Isopyrum stoloniferum</i> MAXIM.	<i>Tsurushirokanesō.</i>
<i>Thalictrum aquilegifolium</i> LINN.	<i>Karamatsusō.</i>

Papaveraceæ.

<i>Corydalis decumbens</i> PERS.	<i>Yabu-engosaku.</i>
<i>Corydalis pallida</i> PERS.	<i>Miyamakikeman.</i>

Fig. 22. *Rodgersia podophylla* A. GRAY, and *Cimicifuga fetida* LINN. var. *simplex* HUTT., on the Yoshida side, about 17500 m. above the sea. The former plant is easily distinguishable in having its large whorled leaves, and the latter is very remarkable for its long spikes of white flowers. (From the Atlas of Japanese Vegetation, by permission of Prof. M. MIYOSHI.)

Saxifragaceæ.

Astilbe Thunbergii MIQ.	<i>Tōriashishōma.</i>
Chrysosplenium macrostemon MAXIM.	<i>Miyamanekonomesō.</i>
Hydrangea Hortensia DC. var. acuminata A. GR.	<i>Sawa-ajisai.</i>
Hydrangea Thunbergii SIEB.	<i>Amachanoki.</i>
Philadelphus coronarius LINN. var. Satsumi MAXIM.	<i>Satsuma-utsugi.</i>
Ribes alpinum LINN. var. japonicum MAXIM.	<i>Zarikomi.</i>
Ribes ambiguum MAXIM.	<i>Yashabishaku.</i>
Rodgersia podophylla A. GR.	<i>Yamaguruma.</i>
Schizophragma hydrangeoides S. et Z.	<i>Iwagarami.</i>
Tiarella polyphylla DON.	<i>Zutayakushū.</i>

Rosaceæ.

Amelanchier asiatica C. COCH.	<i>Zaifuriboku.</i>
Fragaria elatior EHRH.	<i>Miyamahebi-ichigo.</i>
Pourthiæa villosa DCNE.	<i>Kamatsuka.</i>
Prunus incisa THUNB.	<i>Mamezakura.</i>
Pyrus alnifolia LINN.	<i>Azuki-nashi.</i>
Pyrus Aria EHRH. var. Kamaonensis WALL.	<i>Urajironoki.</i>
Pyrus aucuparia GÆRTN. var. japonica MAXIM.	<i>Miyamananakamado.</i>
Pyrus Matsumuræ MAKINO.	<i>Nanakamado.</i>
Pyrus sambucifolia CH. et SCHL.	<i>Takanenanakamado.</i>
Rosa Lucie FR. et ROCH.	<i>Hai-ibara.</i>
Rubus morifolius SIEB.	<i>Kumaichigo.</i>
Rubus rosifolius SM.	<i>Bara-ichigo.</i>
Rubus Sieboldii BL.	<i>Hōroku-ichigo</i> or <i>Oni-ichigo.</i>
Spiræa bulbata MAXIM.	<i>Koshimotsuke.</i>
Spiræa japonica LINN. f.	<i>Shimotsuke.</i>

Geraniaceæ.

Geranium eriostemon FISCH.	<i>Gunnaifuuro.</i>
Geranium Wilfordi MAXIM.	<i>Kofuuro.</i>

Oxalidaceæ.

Oxalis acetosella LINN.	<i>Miyamakatabami.</i>
-------------------------	------------------------

Rutaceæ.

Zanthoxylum schiimifolium S. et Z. *Inuzanshū.*

Anacardiaceæ.

Rhus toxicodendron LINN. var. *radicaus* MIQ. *Tsuta-urushi.*

Aquifoliaceæ.

Ilex geniculata MAXIM. *Furinnunemodoki.*
Ilex macropoda MIQ. *Aohada.*

Celastraceæ.

Euonymus alata K. KOCH. var. *subtriflora* F. et S. *Komayumi.*
Euonymus melanantha F. et S. *Sawadatsu* or *Aogiku-mayumi.*
Euonymus sachalinensis MAXIM. *Murasaki-tsuribana.*

Staphyleaceæ.

Staphylea Bumalda S. et Z. *Mitsuba-utsugi.*

Aceraceæ.

Acer argutum MAXIM. *Asanohakaede.*
Acer carpiniifolium S. et Z. *Yamasheba.*
Acer cratægifolium S. et Z. *Urikaede.*
Acer japonicum THUNB. *Hauchiwakae.*
Acer micranthum S. et Z. *Kominekade.*
Acer nikøense MAXIM. *Chōjanoki.*
Acer purpurascens S. et Z. *Kajikade.*
Acer rufinerve S. et Z. *Urihadakade.*
Acer Sieboldianum MIQ. var. *microphyllum* MAXIM. *Ko-hauchiwakade.*
Acer spicatum LAM. var. *ukurunduense* MAXIM. *Ogarabana.*

Balsaminaceæ.

Impatiens nolitangere LINN. *Ki-tsurijune.*
Impatiens Textori MIQ. *Tsurifuncsō.*

Vitaceæ.

<i>Vitis Coignetiae</i> PULL.	<i>Gaebu</i> or <i>Yamabudō</i> .

Dilleniaceæ.

<i>Actinidia arguta</i> PLANCH.	<i>Sarunashi</i> .
<i>Actinidia Kolomikta</i> MAXIM.	<i>Miyamamatatabi</i> .
<i>Actinidia polygama</i> MIQ.	<i>Matatabi</i> .

Violaceæ.

<i>Viola Bissetii</i> MAXIM.	<i>Nagabasaishinsumire</i> .
<i>Viola japonica</i> LANGSD.	<i>Kosumire</i> .
<i>Viola Maximowicziana</i> MAKINO.	<i>Komiyamasumire</i> .
<i>Viola pinnata</i> LINN. var. <i>chærophyllodes</i> RGL.	<i>Ezo-sumire</i> .
<i>Viola variegata</i> FISCH.	<i>Miyamasumire</i> .
<i>Viola verecunda</i> A. GR. var. <i>semilunaris</i> MAXIM.	<i>Agisumire</i> .

Onagraceæ.

<i>Circea alpina</i> LINN.	<i>Miyamatanitade</i> .
<i>Circea erubescens</i> FR. et SAV.	<i>Tanitade</i> .
<i>Epilobium dahuricum</i> FISCH.	<i>Hime-akabana</i> .
<i>Epilobium japonicum</i> HASSK.	<i>Iwa-akabana</i> .

Araliaceæ.

<i>Aralia cordata</i> THUNB.	<i>Wudo</i> .
<i>Acanthopanax sciadophylloides</i> F. et S.	<i>Koshiabura</i> .
<i>Aralia repens</i> (MAXIM.) MAKINO.	<i>Tochibaninjin</i> .

Umbelliferæ.

<i>Angelica Florenti</i> F. et S.	<i>Shiraneinjin</i> .
<i>Angelica polymorpha</i> MAXIM.	<i>Shiranesenkiu</i> .
<i>Carum Tanakæ</i> FR. et SAV.	<i>Iwasentōsō</i> .
<i>Seseli Libanostis</i> KOCH.	<i>Ibukibōfū</i> .
<i>Sium Ninsi</i> LINN.	<i>Mukagoninjin</i> .

Fig. 23. Forest of deciduous trees, on the Yoshida side, about 1500 m. above the sea. *Aralia cordata* THUNB. on the right, *Fraxinus longicuspis* SIEB. et ZUCC. on the left, *Fagus*, *Quercus*, *Deutzia*, etc. are seen on the background. (From the Atlas of Japanese Vegetation by permission of Prof. M. MIYOSHI.)

Cornaceæ.

Cornus Kousa BUEG.

Yamabōshi.

Pirolaceæ.

Chimaphila japonica MIQ.

Mumegasasō.

Monotropia hypopitys LINN. var. *hirsuta* ROTH.

Shakujōbana.

Monotropa uniflora LINN.	<i>Ginryōsō.</i>
Pirola elliptica NUTT. var. minor MAXIM.	<i>Kobano-ichiyakusō.</i>
Pirola media SW.	<i>Maruba-ichiyakusō.</i>
Pirola renifolia MAXIM.	<i>Jinyō-ichiyakusō.</i>
Pirola rotundifolia L. var. albiflora MAXIM.	<i>Ichiyakusō.</i>
Pirola secunda LINN.	<i>Ko-ichiyakusō.</i>

Ericaceæ.

Andromeda campanulata MIQ.	<i>Furintsutsuji.</i>
Gaultheria adenothrix MAXIM.	<i>Akamono.</i>
Leucothoe Grayana MAXIM.	<i>Hanahirinoki.</i>
Menziesia pentandra MAXIM.	<i>Koyōrakutsutsuji.</i>
Pieris japonica D. DON.	<i>Asebi.</i>
Rhododendron dilatatum MIQ.	<i>Mitsubatsutsuji.</i>
Vaccinium bracteatum THUNB.	<i>Shushanpo.</i>

Diapensiaceæ.

Schizocodon soldanelloides S. et Z.	<i>Iwakagami.</i>
-------------------------------------	-------------------

Primulaceæ.

Trientalis europæa LINN.	<i>Tsumatorisō.</i>
Primula japonica A. GRAY.	<i>Kurinsō.</i>

Symplocaceæ.

Symplocos cratægoides HAM.	<i>Sawafutagi.</i>
----------------------------	--------------------

Styracaceæ.

Styrax japonica S. et Z.	<i>Egonoki.</i>
--------------------------	-----------------

Oleaceæ.

Ligustrum Ibota SIEB.	<i>Ibota.</i>
Ligustrum Ibota SIEB. var. ciliare MAXIM.	<i>Miyama-ibota.</i>

Gentianaceæ.

Halenia sibirica BORNK.	<i>Hana-ikari.</i>
-------------------------	--------------------

Labiatae.

<i>Calamintha multicaulis</i> MAXIM.	<i>Miyamatōbana.</i>
<i>Plectranthus inflexus</i> VAHL.	<i>Yamahakka.</i>

Scrophulariaceae.

<i>Lathræa japonica</i> MIQ.	<i>Yama-utsubo.</i>
<i>Pedicularis gloriosa</i> BISSET et MOOR.	<i>Hankwai-azami.</i>
<i>Pedicularis resupinata</i> LINN.	<i>Shōgamagiku.</i>

Gesneraceae.

<i>Conandron ramondioides</i> S. et Z.	<i>Iwatabako.</i>
--	-------------------

Rubiaceae.

<i>Galium brachypodium</i> MAXIM.	<i>Marubano-yotsubamugura.</i>
<i>Galium paradoxum</i> MAXIM.	<i>Yotsubamugura.</i>
<i>Pseudopyxis longituba</i> FR. et SAV.	<i>Shirobana-inamorisō.</i>

Caprifoliaceae.

<i>Diervilla japonica</i> DC.	<i>Beniutsugi.</i>
<i>Viburnum dilatatum</i> THUNB.	<i>Gamazumi.</i>
<i>Viburnum furecatum</i> BLUME.	<i>Mushikari.</i>
<i>Viburnum Wrightii</i> MIQ.	<i>Miyamagamazumi.</i>

Valerianaceae.

<i>Patrinia palmata</i> MAXIM.	<i>Kinreikwa.</i>
<i>Patrinia villosa</i> JUSS.	<i>Otoko-eshi.</i>

Campanulaceae.

<i>Adenophora remotiflora</i> MIQ.	<i>Sobana.</i>
<i>Adenophora verticillata</i> FR. et SAV.	<i>Tsuriganeninjin.</i>
<i>Adenophora polymorpha</i> LEDEB. var.	
<i>Lamarckii</i> TRAUTV.	<i>Himeshajin.</i>
<i>Peracarpa circæoides</i> H. FEE.	<i>Savagikyō.</i>
<i>Phyteuma japonicum</i> MIQ.	<i>Shideshajin.</i>

Compositae.

<i>Adenocaulon bicolor</i> HOOK.	<i>Nobuki.</i>
<i>Ainsliæ apiculata</i> SCH.-BIP.	<i>Kikkōhagama.</i>
<i>Aster dimorphophyllus</i> FR. et SAV.	<i>Tateyamagiku.</i>
<i>Aster scaber</i> THUNB.	<i>Shirayamagiku.</i>

<i>Aster trinervius</i> ROXB.	<i>Yamashirogiku.</i>
<i>Cacalia adenostylioides</i> FR. et SAV.	<i>Kanikōmori.</i>
<i>Cacalia delphiniifolia</i> SIEB. et ZUCC.	<i>Momijisō.</i>
<i>Cacalia farfaræfolia</i> SIEB. et ZUCC. var. farfaræfolia MAXIM.	<i>Miyamakōmorisō.</i>
<i>Cacalia hastata</i> LINN. var. glabra LEDEB.	<i>Yobusumaso.</i>
<i>Carpesium triste</i> MAXIM.	<i>Gankubiyabutabako.</i>
<i>Cirsium spicatum</i> (MAXIM.)	<i>Yama-azami.</i>
<i>Lactuca Raddeana</i> MAXIM.	<i>Yamanigana.</i>
<i>Macroclinidium robustum</i> MAXIM.	<i>Kashiwabahaguma.</i>
<i>Pertya scandens</i> SCH.-BIP.	<i>Nagabanokōyabōki.</i>
<i>Prenanthes acerifolia</i> (MAXIM.)	<i>Fuku-ōsō.</i>
<i>Saussurea Maximowiczii</i> HERD.	<i>Miyakoazami.</i>
<i>Saussurea Tanakæ</i> FR. et SAV. var. phyllolepis MAXIM.	<i>Tōhiren.</i>
<i>Saussurea triptera</i> MAXIM.	<i>Yahazuhigotai.</i>
<i>Senecio nemorensis</i> LINN. var. Fuchsii KOCH.	<i>Kion.</i>
<i>Solidago Virga-aurea</i> LINN.	<i>Akinokirinso.</i>

Fig. 24. Coniferous forest on the north side of the mountain; *Picea*, and *Abies*, with a few deciduous trees on the right.
(After a photograph.)

Fig. 25. Coniferous forest on the top of the Konnō pass, about 1500 m. above the sea.
(Photographed by the author.)

c) Ever-green Conifer-Region.

This is the most abundant vegetation on the mountain. As shown in the annexed map, the region forms a nearly complete belt around the middle of the mountain, the only intrusion of volcanic ashes being on the eastern side where this formation has almost disappeared. On the south, the formation begins at 1550 m. and stretches up to an elevation of 2300 m. On the north, it comes a little lower than on the south. Distant views of the region on the north and south are seen in Plates I., and VIII. As we have previously stated, there are some Conifer formations which come down to the prairie and are even isolated in it; such as the Aogigahara formation, the Kenmarubi formation, and the pure stand of *Picea polita* near lake Yamanaka.

On the south side of Part I., (cf. the foregoing sketch) on the lower boundary of this formation, there are found a few

Pl. VII. View of Mt. Fuji, seen from lake Saiko. (Nishilake.)

(After a sketch by the author.)

Abies firma. Ascending to an altitude of 1500 m., we see that the prevailing Conifer in this region is *Picea Veitchii* with a small amount of *Chamaecyparis*, *Tsuga* and *Picea hondōensis*. In the Kamiide district of the same part, the predominating tree is *Picea hondōensis* with the addition of some *Tsuga* and *Abies umbellata*. Going over to the due west side, *Picea Veitchii* begins to increase, occupying 50% of the whole, leaving 30% to *Tsuga*, and 20% to *Picea hondōensis*. Coming up to Part IV., we see here the famous forest of Aogigahara,—a grand growth stretching over the large area of 14 km. from

Fig. 26. Mt. Fuji from lake Shōji, the Aogigabara forest is seen at the base of the mountain, and on the left the promontory of the tertiary formation is covered with thin clumps of *Pinus densiflora*. The pine on the left in the foreground is *P. densiflora*.
(Photographed by the author.)

its upper to its lower end, and 5 km. from side to side. Although some local differences may be observed here, it is the *Tsuga* that controls the vegetation by so much as eighty percent, with much under-growth and with some additions of *Chamæcyparis*. This *Chamæcyparis* being tolerant of shade, can grow even in the dense forest of *Tsuga*. *Picea polita* and *Pinus parviflora* also intermingle here.

Fig. 27. Conifer forest of Aogigahara, near the top of the Konno pass. Conifers in the background are *Picea polita*, and *Tsuga Sieboldii*; the deciduous-tree in the foreground is *Alnus incana*.
(Photographed by the author.)

Turning to the due north side, the Conifer formation stretches from 2000 m. down to 1500 m.; and presents a grand distant view making a belt around the peak. The predominating tree on this side is *Tsuga* especially in the vicinity of Maruyama, i.e. 70% of *Tsuga* and 20% of *Larix*. A little eastwards from here, we come to Yakikomi (cf. map.) where *Larix* predominates to such an extent that it ranges from 1500 m. up

Fig. 28. The same; *Picea polita*.

(Photographed by the author.)

to 2200 m., thus intercepting this ever-green formation. This *Larix* formation is clearly distinguishable by its fresh green colour in the early spring.

Moving to the eastwards, we see that the forest gradually becomes thinner. Some deciduous trees appear below the height of 1500 m. They are *Fagus*, *Alnus*, and a very large quantity of *Sophora japonica* LINN. On the due east side i.e. the Subashiri side, *Larix* attains its most luxuriant growth, and extends through all tree regions.

On the Gotemba side, the Conifer formation is the least developed and in some parts there are even almost no Conifers at all; but in other parts there is a pure red pine formation which ranges from 800 m. up to 1200 m.

Fig. 29. Ever-green Conifer formation on the Subashiri side.

(Photographed by the author.)

The forest trees found in this region are as follows:—

Coniferæ.

Abies homolepis S. et Z.
Abies Veitchii LINDL.
Abies umbellata MAYR.

Dakemomi.
Shirabi.
Urajiromomi.

Fig. 33. Coniferous forest on the north side of the mountain, *Picea*, *Abies* and a few *Pyrus sambucifolia* on the right side. (After a photograph.)

Larix leptolepis GORD.

Picea Alcockiana CARR.

Picea ajanensis FISCH.

**Picea hondōensis* MAYR.

Picea polita CARR.

Pinus parviflora SIEB et ZUCC.

Tsuga diversifolia MAXIM.

Tsuga Sieboldiana CARR.

Karamatsu.

Iramomi.

Ezomatsu.

Tōhi.

Harinomi.

Himekomatsu.

Kometsuga.

Tsuga.

The undergrowth in this formation is almost the same as in the preceding formation.

d) *Larix* region.

Above a 2300 m. elevation of the Conifer region, the wind is too strong, cold, and dry for the growth of the ever

* I do not think that *Picea hondōensis* MAYR should be different from *Picea ajanensis* FISCH. The differences pointed out by MAYR in his "Abietinæ von Japan" is not very clear, and perhaps they should be regarded as individual characters.

green Conifers, and they give way to *Larix* which there forms the deciduous Conifer formation. *Larix*, being deciduous in the winter time, is tolerant of extreme dryness and severe cold. On the south, this *Larix* is found in as high as 2900 m. elevation. There it is seen clinging firmly to the ground, so as to prevent its being swept away by the wind as *Pinus pumila* does under the same conditions. This *Pinus* is very common in the alpine regions of Japan, but it is entirely absent on this mountain.

Fig. 31. *Larix leptolepis* on the western side (winter). (Photographed by the author.)

The plant-species found in this formation are almost the same as those in the following formation.

e) *Salix-Alnus Region.*

At about 2500 m. elevation, the *Larix* formation gives way to the shrubby-formation, such as *Salix*, *Alnus*, *Betula* and others. On the south this formation begins at 2600 m., mixed with *Larix*. On the north, it comes a little lower. The plants found in this region are as follows:—

Polypodiaceæ.

<i>Nephrodium gracilescens</i> BLUME. var.	
<i>glanduligerum</i> BAKER.	<i>Hashigoshida.</i>
<i>Nephrodium Phegopteris</i> BAUMG.	<i>Miyamawarabi.</i>

Coniferæ.

<i>Juniperus nipponica</i> MAXIM.	<i>Miyamanezu.</i>
<i>Larix leptolepis</i> GORD.	<i>Karamatsu.</i>

Cyperaceæ.

<i>Carex Dœnitzii</i> BÆCK.	<i>Kotanukiran.</i>
<i>Carex Wrightii</i> BÆCK. BULL.	<i>Himisuge.</i>

Gramineæ.

<i>Agrostis canina</i> LINN.	<i>Iwagariyasu.</i>
<i>Glyceria tonglensis</i> CLARK.	<i>Dojôtsunagi.</i>
<i>Miscanthus Matsumuræ</i> HACK.	<i>Kariyasumodoki.</i>

Salicaceæ.

<i>Salix Caprea</i> LINN.	<i>Bekkoyanagi.</i>
<i>Salix Sieboldiana</i> BL.	<i>Iwayanagi.</i>

Betulaceæ.

<i>Alnus incana</i> WILLD. var. <i>emarginata</i>	
MATSUM.	<i>Yakazuhanoki.</i>
<i>Alnus viridis</i> DC. var. <i>sibirica</i> RGL.	<i>Miyamahannoki.</i>
<i>Betula alba</i> LINN. var. <i>vulgaris</i> DC.	<i>Shirakanba.</i>
<i>Betula Ermanni</i> CHAM. var. <i>nipponica</i>	
MAXIM.	<i>Takekanba.</i>

Polygonaceæ.

<i>Polygonum cuspidatum</i> S. et Z.	<i>Itadori.</i>
<i>Polygonum polymorphum</i> LEDEB. var.	
<i>japonicum</i> MAXIM.	<i>Ontade.</i>

Caryophyllaceæ.

<i>Stellaria florida</i> FISCH. var. <i>angusti-</i>	
<i>folia</i> MAXIM.	<i>Iwatsumekusa.</i>

Ranunculaceæ.

<i>Aquilegia glandulosa</i> FISCH.	<i>Yamaodamaki.</i>
<i>Clematis alpina</i> MILL.	<i>Miyamahanshōzuru.</i>

Cruciferæ.

<i>Arabis serrata</i> F. et S.	<i>Fujihatazao.</i>

Rosaceæ.

<i>Aruncus silvester</i> KOSTEL.	<i>Yamabukishōma.</i>
<i>Fragaria elatior</i> EHRH.	<i>Shiobanahebiichigo.</i>
<i>Pyrus sambucifolia</i> CH. et SCHL.	<i>Takane-nanakamado.</i>
<i>Rosa acicularis</i> LINDL.	<i>Takanebara.</i>
<i>Rubus japonicus</i> FOCK.	<i>Goyō-ichigo.</i>
<i>Spiræa bracteata</i> MAXIM.	<i>Iwashimotsuke.</i>
<i>Spiræa bulbata</i> MAXIM.	<i>Koshimotsuke.</i>

Umbelliferæ.

<i>Angelica Florenti</i> F. et S.	<i>Shiraneninjin.</i>
<i>Angelica hakonensis</i> MAXIM.	<i>Iwaninjin.</i>
<i>Carum Tanakæ</i> FR. et SAV.	<i>Iwasentōsō.</i>

Ericaceæ.

<i>Leucothoe Grayana</i> MAXIM.	<i>Hanahirinoki.</i>
<i>Menziesia pentandra</i> MAXIM.	<i>Koyōrakutsutsuji.</i>
<i>Rhododendron Metternichii</i> S. et Z.	<i>Shakunange.</i>
<i>Vaccinium Vitis-idæa</i> LINN.	<i>Koke-momo.</i>
<i>Tripetaleia paniculata</i> S. et Z.	<i>Hotsutsuji.</i>

Pirolacæ.

Pirola media SW. *Maruba-ichiyakusō.*

Diapensiaceæ.

Schizocodon soldanelloides S. et Z. *Iwakagami.*

Oleaceæ.

Ligustrum Ibota SIEB. var. *ciliare*
MAXIM. *Miyama-ibota.*

Valerianaceæ.

Patrinia palmata MAXIM.

Orobanchaceæ.

Boschniakia glabra C.A. MEY. *Oniku.*

Campanulaceæ.

Adenophora remotiflora MIQ. *Sobana.*
Adenophora verticillata FR. et SAV. *Tsuriganeninjin.*

Compositæ.

<i>Aster trinervius</i> ROXB.	<i>Yamashirogiku.</i>
<i>Ainsliæa apiculata</i> SCH.-BIP.	<i>Kikkōhaguma.</i>
<i>Cacalia adenostylioides</i> FR. et SAV.	<i>Kanikōmori.</i>
<i>Cacalia delphiniifolia</i> SIEB. et ZUCC.	<i>Momijisō.</i>
<i>Cacalia farfaræfolia</i> SIEB. et ZUCC. var. farfaræfolia MAXIM.	<i>Miyamakōmorisō.</i>
<i>Saussurea Maximowiczii</i> HERD.	<i>Miyako-azami.</i>
<i>Saussurea triptera</i> MAXIM.	<i>Yahazuhigotai.</i>
<i>Solidago Virga-aurea</i> LINN.	<i>Akinotamurasō.</i>

f) Higher Grass Region.

Above the *Salix-Alnus* formation, there comes a small area of higher (alpine) grass formation. This formation is, however,

very poor on this mountain, owing to its recent habitation. The plants found here are as follows:—

Ophioglossaceæ.

Botrychium Lunaria Sw.

Himehanawarabi or *Hebinoshita*.

Gramineæ.

Deschampsia caryophyllea L.

Nukasusuki.

Cyperaceæ.

Carex Dœnitzii BECK.

Kotanukiran.

Carex stenantha FR. et SAV.

Iwasuge.

Carex Wrightii FR.

Himesuge.

Orchidaceæ.

Orchis Chondradenia MAKINO.

Onoe-ran.

Plantanthera conopsea SCHLECHT.

Chidorisô.

Fig. 32. *Potygonum cuspidatum* SIEB et ZUCC., on the Subashiri side, about 2000 m. above the sea. (From the Atlas of Japanese Vegetation, by permission of Prof. M. MIYOSHI.)

Salicaceæ.

Salix Sieboldiana BL. *Iwayanagi.*

Betulaceæ.

Alnus incana WILLD. var. *emarginata*
MATSUM. *Yahazuhannoki.*

Alnus viridis DC. var. *sibirica* RGL. *Miyamahannoki.*

Polygonaceæ.

Polygonum cuspidatum S. et Z. *Itadori.*

Polygonum debile MEISN. var. *triangularis* MEISN. *Miyamatanisoba.*

Polygonum polymorphum. LEDEB. var.
japonicum MAXIM. *Ontade.*

Caryophyllaceæ.

Stellaria florida FISCH. var. *angustifolia* MAXIM. *Iwatsumekusa.*

Ranunculaceæ.

Clematis alpina MILL. *Miyamahanshōzuru.*

Cruciferæ.

Arabis serrata F. et S. *Fujihatazao.*

Leguminosæ.

Astragalus adsurgens PALL. *Murasakimomenzuru.*

Astragalus reflexistipulus MIQ. *Momenzuru.*

Hedysarum esculentum LEDEB. *Tateyama-ōgi* or *Iwa-ōgi.*

Ericaceæ.

Cassiope lycopodioides DON. *Iwahige.*

Leucothœ Grayana MAXIM. *Hanahirinoki.*

Vaccinium Vitis-idæa LINN. *Kokemomo.*

Pirolaceæ.

Pirola media Sw.

Maruba-ichiyakusō.

Diapensiaceæ.

Schizocodon soldanelloides S. et Z.

Iwakagami.

Compositæ.

Artemisia pedunculosa MIQ.

Miyama-otokoyomogi.

Cirsium purpuratum (MAXIM.)

Fuji-azami.

Solidago Virga-aurea LINN.

Akinokirinsō.

Fig. 33. *Cirsium purpuratum* (MAXIM.), on the Subashiri side, about 2000 m. above the sea.
(From the Atlas of Japanese Vegetation by permission of Prof. M. MIYOSHI).

From 3220 m. upwards, there is no vegetation except a few mosses such as *Racomitrium canescens* BRID., *Stereodon brachycarpus* MITT. *S. plicatulus* LINDB., and a few lichens. On the

south, this grass region extends as high as 3220 m.; but on the north, it is limited under 3000 m. From this elevation upwards, the mountain is entirely barren and there is nothing to represent vegetation except a few lichens and mosses.

Fig. 34. Grass region on the north side of the mountain. (After a photograph).

V. GENERAL ASPECT OF THE FLORA OF THE MOUNTAIN.

Having arranged the plants according to their regions, let us consider the general aspect of the flora of the mountain. Summarizing all the flora, it is found to contain 900 species belonging to 125 families. The following table shows the number of species found in each region, and the sum of the species belonging to each family.

Families.	Regions.	Prairie.	Forest regions, including deciduous broad leaved trees, and evergreen conifers.	Shrubby regions including <i>Larix</i> and <i>Abies</i> <i>Salix</i> regions.	Higher grass region.	*Total number of species belonging to each family.
Hymenophyllaceæ			2			2
Polypodiaceæ		48	19	2		57
Schizæaceæ		1				1
Osmundaceæ		1	1			2
Ophioglossaceæ		1			1	2
Equisetaceæ		1				1
Lycopodiaceæ		2	2			3
Selaginellaceæ		2				2
Coniferae		6	11	2		18
Sparganiaceæ		1				1
Potamogetonaceæ		3				3
Alismataceæ		1				1
Hydrocharitaceæ		3				3
Gramineæ		59	12	3	1	66
Cyperaceæ		35	3	2	3	58
Araceæ		3				3
Eriocaulaceæ		1				1
Commelinaceæ		2				2
Pontederiaceæ		2				2
Juncaceæ		3	2			5
Stemonaceæ			1			1
Liliaceæ		22	13			32
Amaryllidaceæ		1				1
Dioscoreaceæ		3				3
Iridaceæ		2				2
Orchidaceæ		10	15		2	25

*It should be understood that there may exist the same species in different regions and therefore the total number of species belonging to each family is the sum of different species represented in different regions, but not the sum of figures given in the different columns.

Families. \ Regions.	Prairie.	Forest regions, including deciduous broad leaved trees, and evergreen and conifers.	Shrubby regions including <i>Larix</i> and <i>Abies</i> formations.	Higher grass region.	Total number of species belonging to each family.
Saururaceæ	1				1
Chloranthaceæ	2				2
Juglandaceæ		2			2
Salicaceæ	3	1	2	1	5
Betulaceæ	4	3	4	2	9
Fagaceæ	8	3			11
Ulmaceæ	1				1
Moraceæ	2				
Urticaceæ	13	1			13
Loranthaceæ	1				1
Santalaceæ	1				1
Aristolochiaceæ	2				2
Polygonaceæ	14		2	3	17
Chenopodiaceæ	1				1

Fig. 35. View of Mt. Fuji, seen from Ukishimanuma (Ukishima-swamp), forest regions are faintly visible.
(after a photograph).

Families.	Regions.		Forest regions, including deciduous broad leaved trees, and evergreen conifers.	Shrubbery regions including <i>Larix</i> and <i>Pinus</i> <i>Salix</i> formations.	Higher grass region.	Total number of species belonging to each family.
	Prairie.					
Amarantaceæ	2					2
Portulacaceæ	1					1
Caryophyllaceæ	10	1		1	1	12
Nymphæaceæ	1					1
Ceratophyllaceæ	1					1
Magnoliaceæ	5	1				5
Trochodendraceæ		1				1
Ranunculaceæ	18	9		2	1	22
Lardizabalaceæ	2					2
Berberidaceæ	2					2
Menispermaceæ	1					1
Lauraceæ	7					7
Papaveraceæ	5	2				6
Crucifereæ	7			1	1	8
Crassulaceæ	5					5
Saxifragaceæ	12	10				21
Hamamelidaceæ	1					1
Rosaceæ	26	15		7		43
Leguminosæ	32				3	35
Geraniaceæ	3	2				4
Oxalidaceæ	1	1				2
Linaceæ	1					1
Rutaceæ	4	2				5
Simarubaceæ	1					1
Polygalaceæ	1					1
Euphorbiaceæ	9					9
Callitrichaceæ	1					1
Coriariaceæ	1					1
Buxaceæ	1					2

Families. \ Regions.	Prairie.	Forest regions, including deciduous broad leaved trees, and evergreen conifers.	Shrubby regions including <i>Larix</i> and <i>Pinus</i> <i>Salix</i> formations.	Higher grass region.	Total number of species belonging to each family.
Anacardiaceæ	4	1			4
Aquifoliaceæ	4	2			6
Celastraceæ	6	3			9
Staphyleaceæ	1	1			2
Aceraceæ		12			12
Hippocastanaceæ		1			1
Balsaminaceæ	2	2			2
Sabiaceæ	2	1			2
Rhamnaceæ	2				2
Vitaceæ	5	1			5
Tiliaceæ		1			1
Dilleniaceæ		3			3
Theaceæ	2	2			4
Stachyuraceæ	1				1
Guttiferæ	6				6
Elatinaceæ	1				1
Violaceæ	3	6			8
Thymelæaceæ	2				2
Eleagnaceæ	3				3
Lythraceæ	2				2
Onagraceæ	3	4			7
Halorrhagidaceæ	1				1
Araliaceæ	7	3			9
Umbelliferæ	14	5	3		20
Cornaceæ	4	3			5
Clethraceæ	1	1			1
Pirolaceæ		8	1	1	8
Ericaceæ	9	7	5	3	18
Diapensiaceæ		1	1	1	1

Families.	Regions.		Forest regions, including deciduous broad leaved trees, and evergreen conifers.	Shrubby regions including <i>Larix</i> and <i>Pinus</i> <i>Salix</i> formations.	Higher grass region.	Total number of species belonging to each family.
	Prairie.					
Myrsinaceæ	1					1
Primulaceæ	5	2				7
Ebenaceæ				1		1
Symplocaceæ	1	1				1
Styracaceæ	1	1				1
Oleaceæ	4	4		1		7
Loganiaceæ	1					1
Gentianaceæ	5	1				5
Apocynaceæ	1					1
Asclepiadaceæ	3					3
Convolvulaceæ	2					2
Borraginaceæ	2					2
Verbenaceæ	3					3
Labiataæ	20	2				21
Solanaceæ	1					1
Scrophulariaceæ	15	3				16
Orobanchaceæ	1			1		2
Gesneriaceæ	1	1				1
Acanthaceæ	3					3
Plantaginaceæ	1					1
Rubiaceæ	11	3				13
Caprifoliaceæ	13	5				16
Valerianaceæ	2	2		1		3
Dipsaceæ	1					1
Cucurbitaceæ	2					2
Campanulaceæ	5	5		2		9
Compositæ	51	20		8	3	62

Among all the families, the largest is the Gramineæ having 65 species; next come the Polypodiaceæ and the Compositæ, then the Cyperaceæ, the Orchidaceæ, the Ranunculaceæ, and the Umbelliferæ, each having more than 20 species.

Since we have as yet no complete list of other mountain-flora in Japan, we are not yet in a position to make a comparison between the floras of the volcano and other mountains. Still we may be justified in saying that the flora of the present mountain is much more limited than that of other mountains.

VI. INFLUENCE OF FIRE ON THE PRESENT VEGETATION.

On the basal part of the mountain which is near inhabited districts, fires often occur, intentionally or accidentally, and

Fig. 36. View of Mt. Fuji, seen from Yamanaka; dark forests of ever-green conifers are in sharp relief.
(Phot. by the author).

sometimes destroy large portions of forest within a few hours. On this mountain, all the present prairie formations, except that of the Hōyē part, where the ground is not yet ready to accept plants, would have been a spacious forest formation but for being constantly burnt and cut over. Cutting is also an active factor in changing forests into prairie. In former times, there were, we are told, rich clumps of trees on the present prairie on all sides; but in places near inhabited districts the cutting took place little by little, and thus in course of time, formed a broad prairie. As far as my observation goes, fire can not extend far upwards from the base. The flanking is much steeper in the middle portion of the slope than in the base, and on the steep flank, a strong wind is always blowing down from the top, thus preventing fires from spreading towards the peak. So, the deciduous forests may be destroyed by fire, but the coniferous forests are generally free from such destruction, owing to their situation. After considering all the cases, I am forced to the conclusion that the present prairie is due to fire, or cutting over, except those parts on the Gotemba side where the volcanic ashes are too fresh to support vegetation.

VII. ABSENCE OF *PINUS PUMILA* ON THE MOUNTAIN.

Ascending as high as 2500 m., one is struck with the singular habit of the *Larix* in clinging to the ground, and stretching its arms over it, owing to the strong wind which here prevails all the year through. It is interesting to observe that this *Larix*, which in the lower elevations has the habit of a tall erect trunk, is here so stunted and crushed down to

the ground as to render it liable to be mistaken for *Pinus pumila*.*

In such high regions, *Pinus pumila* generally ranges over all the mountains of Japan. But on this mountain we have not found this pine thus far. The absence of the *Pinus* is worthy of special attention.

Considering the wind direction on the high mountains, where *Pinus pumila* grows, we find that the prevailing wind is south-westerly. But, of the neighbouring mountains which lie on the south-west of Mt. Fuji, there is none sufficiently high to support the *Pinus*. Therefore, the volcano has had no chance to get the seeds of *Pinus pumila* by the southwesterly wind, and this may be considered to be the cause of the absence of this tree on Mt. Fuji.

VIII. SUMMARY.

In the course of my several visits to the mountain, I have determined the difference of its vegetation according to altitude and exposure. So far as the exposure alone is concerned, the vegetation may be divided into two classes:—the northern and southern. On the northern slope, the conifer predominates, while on the south, the deciduous trees are in control. To ascertain the cause of this divergence, I have examined the physiography of the mountain and found that the divergence may be ascribed to the difference of the climatic factors, and principally to that of the saturation-deficits on the two sides. Next, I have considered the general aspect of the plant-formation, dividing the whole mountain into four parts, and describing

* The *Pinus* was erroneously recorded from the mountain by some authors.

the formations separately. On the southern side, I have found that the forest-formation attains its most luxuriant growth at the 2000 m. elevation. Next, I have dwelt upon each region more particularly and stated the maximum and minimum limitations of the growth. Then, the zonation of plant-regions has been discussed zone by zone. The species of each region are mentioned and the lists of plants given. I have also summarized all the plants in one table, and given a short note on the general aspect of the flora of the mountain. The plants contained in the table are in all 739 species belonging to 96 families. Next, I have considered the effect of fires, and come to the conclusion that the present prairie is mainly due to forest-fires. Lastly, I have considered the cause of the absence of *Pinus pumila* and stated that the wind direction is not suitable for conveying the seeds of the *Pinus* from the neighbouring mountains to the volcano.

[THE END.]

Pl VIII. View of Mt. Fuji, seen from the top of the Misaka-pass (Misaka-tôgô); forest regions are clearly seen between the snow, and prairie, regions.
(Sketched from nature by the author, in April 1902.)

Index.

- Abelia** *spathulata* S. et Z. 69
Abies. 41, 42
— *bicolor*. 40
— *brachyphylla* MAXIM. 76
— *firma* S. et Z. 46, 89
— *homolepis* S. et Z. 92
— *umbellata* MAYR. 92, 89
— *Veitchii* LINDL. 48, 92
ABSEN OF *Pinus Pumila* ON THE
MOUNTAIN. 108
Acalypha australis LINN. 61
Acanthaceæ. 69
Acanthopanax divaricatum S. et Z. 65
— *ricinifolium* S. et Z. 65
— *sciadophylloides* F. et S. . . . 65
— *spinosum* MIQ. 65
Aceraceæ. 75, 82
Acer argutum MAXIM. 82
— *carpinifolium* S. et Z. 82
— *cratægifolium* S. et Z. 82
— *japonicum* THUNB. 82
— *micranthum* S. et Z. 82
— *nikoëse* MAXIM. 82
— *palmatum* THUNB. 75
— *pictum* THUNB. 27, 75
— *purpurascens* S. et Z. 82
— *rufinerve* S. et Z. 82
— *Sieboldianum* MIQ. var. *micro-*
phyllum MAXIM. 82
— *spicatum* LAM. var. *ukuru-*
duense MAXIM. 32
Achillea ptarmicoides MAXIM. . . 70
Achyranthes bidentata BL. var.
japonica MIQ. 55
Aconitum Fischeri REICH. . . . 56, 80
— *lycoctonum* LINN. 66, 80
Acorus gramineus AIT. 50
Actinidia arguta PLANCH. 83
— *Kolomikta* MAXIM. 83
— *polygama* MIQ. 83
Actinostemma racemosum MAXIM. 70
Adenocaulon bicolor HOOK. . . . 70, 86
Adenophora polymorpha LEDEB.
var. *Lamarekii* TRAUTV. 86
— *remotiflora* MIQ. 86, 97
— *verticillata* FISCH. var. *verti-*
cillata FR. et SAV. 70, 86, 97
Adiantum monochlamys EAT. . . . 44
— *pedatum* LINN. 44, 76
Æginetia indica ROXB. 69
Æschynomene indica LINN. 59
Agopyrum semicostatum NEES. . . 47
Agrimonia pilosa LEDEB. 59
Agrostis canina LINN. 77, 95
— *flaccida* HACK. 77
— *perennans* TUCKERM. 47, 77
Ainsliea acerifolia SCH. BIP. . . . 70
— *apiculata* SCH. BIP. 70, 89, 97
Ajuga decumbens THUNB. 67
Akaishi Mt. 4
Akebia lobata DCNE. 57
— *quinata* DCNE. 57
Albizzia Julibrissin BOIV. 59
Alismataceæ. 47
Allium japonicum REGEL. 51
Alnus incana WILLD. var. *emar-*
ginata MATSUM. 95, 99
— *viridis* DC. var. *sibirica* REL. 95, 99

<i>Ahus Yasha</i> MATSUM.	53	<i>Arenaria serpyllifolia</i> LINN. var.	
<i>Alopecurus fulvus</i> SM.	47	<i>leptoclados</i> GUSS.	55
Alpine stretches.	38	<i>Arisæma serratum</i> SCHOTT. form	
Amagi Mts.	4	<i>Blumei</i> MAK.	55
Amarantaceæ.	55	Aristolochiaceæ.	55
<i>Amarantus Blitum</i> LINN.	55	<i>Aristolochia debilis</i> S. et Z.	55
Amaryllidaceæ.	52	— <i>Kæmpferi</i> WILLD.	55
<i>Amelanchier asiatica</i> C. COCH.	59	<i>Artemisia capillaris</i> THUNB.	70
<i>Amphicarpæa Edgeworthii</i> BENTH.		— <i>Keiskeana</i> MIQ.	71
var. <i>japonica</i> OLIV.	59	— <i>japonica</i> THUNB.	71
Anacardiaceæ.	62, 82	— <i>pedunculosa</i> MIQ.	100
<i>Anaphalis margaritacea</i> B. et H.	70	— <i>vulgaris</i> LINN. var. <i>indica</i>	
— <i>yedensis</i> FR. et SAV.	70	MAXIM.	71
<i>Andromeda campanulata</i> MIQ.	85	<i>Arthraxon ciliare</i> BEAUB.	47
<i>Andropogon brevifolius</i> Sw.	47	<i>Aruncus silvester</i> KOSTEL.	59, 96
— <i>micranthus</i> KUNTH. var. <i>genui-</i>		<i>Arundinella anomala</i> STEUD.	47
nus HACK.	47	<i>Arundinaria.</i>	44
— <i>Nardus</i> LINN. var. <i>Gœringii</i>		Asclepiadaceæ.	67
HACKEL.	47	<i>Aspidium japonicum</i> MAKINO.	44
<i>Aneilema keisak</i> HASSEK.	51	— <i>nuticum</i> FR. et SAV.	76
<i>Anemone cernua</i> THUNB.	56	— <i>oligophlebium</i> CHRIST.	44
— <i>japonica</i> S. et Z.	56	<i>Asplenium incisum</i> THUNB.	44
<i>Angelica Florenti</i> F. et S.	83, 96	— <i>varians</i> HOOK et GREV.	44
— <i>hakonensis</i> MAXIM.	96	<i>Aster dimorphophyllus</i> FR. et SAV.	86
— <i>Miqueliana</i> MAXIM.	65	— <i>fastigiatus</i> FISCH. et MEY.	71
— <i>polyclada</i> FRANCH.	64, 65	— <i>indicus</i> LINN.	71
— <i>polymorpha</i> MAXIM.	83	— <i>indicus</i> LINN. var. <i>pinnatifidus</i>	
<i>Aogigahara forest.</i>	37, 43	MAXIM.	71
<i>Aogigahara-lava.</i>	6, 5	— <i>scaber</i> THUNB.	71, 86
Apocynaceæ.	67	— <i>trinervius</i> ROXB.	71, 87, 97
Aquifoliaceæ.	62, 82	<i>Astilbe chinensis</i> FR. et SAV. var.	
<i>Aquilegia glandulosa</i> FISCH.	56, 80, 96	<i>japonica</i> MAXIM.	58
<i>Arabis sagittata</i> DC.	58	— <i>Thunbergii</i> MIQ.	58, 81
— <i>serrata</i> F. et S.	96, 99	<i>Astragalus adsurgens</i> PALL.	99
Araceæ.	50	— <i>reflexistipulus</i> MIQ.	101, 99
Araliaceæ.	65, 83	<i>Ataka Mt.</i>	4
<i>Aralia cordata</i> THUNB.	65, 83	<i>Athyrium nipponicum</i> HOOK.	44
— <i>cordata</i> THUNB.	84	— <i>nipponicum</i> BAKER.	77
— <i>repens</i> (MAXIM.) MAKINO.	83	— <i>thelypteroides</i> DESV.	44
— <i>sinensis</i> LINN.	65	— <i>yokoscense</i> MAKINO.	44
<i>Ardisia japonica</i> BL.	66	<i>Atractylis ovata</i> THUNB.	71
		<i>Aucuba japonica</i> THUNB.	65

Balsaminaceæ.	62, 82
Beckmannia cruciformis Host.	47
Berberidaceæ.	57
Berberis Thunbergii DC.	57
Berchemia racemosa S. et Z.	62
Betulaceæ.	53, 74, 79, 95, 99
Betula alba LINN. var. vulgaris	
DC.	74, 95
— Ermanni CHAM. var. nipponica	
MAXIM.	74, 95
Bidens pilosa LINN.	71
— tripartita LINN.	71
Blechnum Spicant ROTH. var. sub-	
serrata LOWE.	44
Blixa caulescens MAX.	47
Bœhmeria holosericea BLUME.	54
— japonica MIQ.	54
— japonica MIQ. var. tricuspidis	
HANCE.	54
— nivea BLUME.	54
— Sieboldiana BLUME.	54
— spicata THUNB.	54
Bœninghausenia albiflora REICHB.	61
Borraginaceæ.	67
Boschniakia glabra C.A. MEY.	97
Botrychium Lunaria Sw.	98
— ternatum Sw.	46
Brachypodium silvaticum BEAUV.	47
Bromus japonicus THUNB.	47
— pauciflorus HACK.	47
Brylkinia caudata FR. SCHM.	47, 77
Bulbostylis barbata KUNTH.	49
Bupleurum falcatum LINN.	65
— sachalinense FR. et SAV.	65
Buxaceæ.	61
Buxus sempervirens var. japonica	
MAKINO.	61

Cacalia	
adensotylloides FR. et	
SAV.	87, 97
— delphinifolia S. et Z.	87, 97

Cacalia farfaræfolia SIEB. et ZUCC.	
var. farfaræfolia MAXIM.	87, 97
— hastata LINN. var. glabra LEDEB.	87
— Krameri (FR. et SAV.	71
Campanulaceæ.	70, 86, 97
Calamagrostis arundinacea ROTH.	
var. sciuroides HACK.	47
— Epigeios ROTH.	47
— sachalinensis FR. SCHM.	78
— villosa MUT. var. Langsdorffii	
HACK.	78
Calamintha chinensis BENTH.	67
— gracilis BENTH.	67
— multicaulis MAXIM.	86
Calanthe discolor LINDL.	52
Callicarpa japonica THUNB.	67
Callitrichaceæ.	16
Callitriche japonica ENGEL.	61
Calystegia sepium R. BR.	67
Cæsalpinia sepiaria ROXB.	59
Campanula punctata LAM.	70
Caprifoliaceæ.	69, 76, 86
Capsella bursa-pastoris MÆNCH.	58
Cardamine hirsuta LINN. var. syl-	
vatica LINN.	58
Carex arenicola F. SCHMIDT.	49
— Brownii TUCK. var. transversa	
KUEK.	49
— brunnea THUNB.	49
— caryophylla LAUER subsp.	
nervata KUEK.	49
— curvicolis FR. et SAV.	49
— Deenitzii BOECK.	78, 75, 98
— Gaudichaudiana KUNTH var.	
Thunbergii KUEK.	49
— gibba WAHL.	50
— incisa BOOT.	50
— ischnostachya STEUD.	50
— japonica THUNB.	50
— — — var. aphanolepis KUEK.	50
— — — var. chlorostachys KUEK.	50
— Morrowii BOOT.	50

<i>Carex Onœi</i> FR. SAV.	50	<i>Cinnamomum pedunculatum</i> NEES.	57
— <i>pseudo-conica</i> FR. et SAV.	50, 78	<i>Circeæ alpina</i> LINN.	83
— <i>remota</i> LINN.	78	— <i>erubescens</i> FR. et SAV.	83
— <i>satsumensis</i> FR. et SAV.	50	— <i>quadrisulcata</i> MAXIM.	64
— <i>stenantha</i> FR. et SAV.	50, 98	<i>Cirsium japonicum</i> DC.	71
— <i>Wrightii</i> BÖCK. BULL.	95, 98	— <i>purpuratum</i> (MAXIM.)	100
<i>Carpesium abrotanoides</i> LINN.	71	— <i>spicatum</i> (MAXIM.)	71, 87
— <i>triste</i> MAXIM.	71, 87	<i>Chloranthus japonicus</i> SIEB.	53
<i>Carpinus japonica</i> BLUME	53	— <i>serratus</i> R. et S.	53
— <i>yedoensis</i> MAXIM.	53	<i>Cladrastis amurensis</i> B. et H. var.	
<i>Carum Tanakæ</i> FR. et SAV.	83, 96	<i>floribunda</i> MAXIM.	60
Caryophyllaceæ.	55, 79, 96, 99	<i>Clematis alpina</i> MILL.	56, 80, 96, 99
<i>Cassia mimosoides</i> LINN.	60	— <i>apiifolia</i> DC.	56
<i>Cassiope lycopodioides</i> DON.	101, 99	— <i>heracleifolia</i> DC. var. <i>stans</i> (S.	
<i>Castanea vulgaris</i> LAM, var. <i>japo-</i>		et Z.)	56
<i>nica</i> DC.	53	— <i>recta</i> LINN. var. <i>paniculata</i>	
Celastraceæ.	62, 82	(THUNB.)	57
<i>Celastrus articulatus</i> THUNB.	62	<i>Clerodendron trichotomum</i> THUNB.	67
<i>Celtis sinensis</i> PERS.	54	Clethraceæ.	65, 75
<i>Centūrea atriplicifolia</i> (DC.)	71	<i>Clethra barbinervis</i> S. et Z.	65, 75
<i>Cephalotaxus drupacea</i> S. et Z.	46	CLIMATE.	8
<i>Cerastium vulgatum</i> LINN. v.r.		<i>Clintonia ndensis</i> TRAUTV.	51
<i>glandulosum</i> KOCH.	46	<i>Cocculus Thunbergii</i> DC.	57
Ceratophyllaceæ.	56	<i>Codonopsis lanceolata</i> B. et H.	70
— <i>demersum</i> LINN.	56	<i>Comanthosphace sublanceolata</i> S.	
— <i>japonicum</i> S. et Z.	75	MOORE.	67
<i>Chamaecyparis.</i>	89, 90	Commelinaceæ.	51
<i>Chamaele tenera</i> MIQ.	56	<i>Commelina communis</i> LINN.	51
<i>Chelidonium Majus</i> LINN.	57	Compositæ.	70, 86, 97, 100
Chenopodiaceæ.	55	<i>Conandron ramondioides</i> S. et Z.	69, 86
<i>Chenopodium album</i> LINN.	55	Coniferæ.	46, 92, 93
<i>Chimaphila japonica</i> MIQ.	84	<i>Coniferous forest.</i>	36
Chloranthaceæ.	53	<i>Coniogromme japonica</i> DIELS.	44
<i>Chrysanthemum sinense</i> var. <i>japo-</i>		Convolvulaceæ.	67
<i>nicum</i> MAXIM.	71	<i>Coptis brachypetala</i> S. et Z. var.	
<i>Chrysosplenium macrostemon</i>		<i>pygmæa</i> MIQ.	80
MAXIM.	81	Coriariaceæ.	61
<i>Cicuta virosa</i> LINN.	65	<i>Coriaria japonica</i> A. GR.	61
<i>Cimicifuga biternata</i> MIQ.	80	Cornaceæ.	75, 84
— <i>foetida</i> LINN. var. <i>simplex</i>		<i>Cornus ignorata</i> C. KOCH.	65, 75
HUTH.	56, 80	— <i>Kousa</i> BUERG.	75, 84
— <i>japonica</i> SPR.	80	— <i>macrophylla</i> WALL.	65, 75

<i>Corydalis decumbens</i> PERS.	57, 80	<i>Desmodium podocarpum</i> DC. var.	
— <i>incisa</i> PERS.	58	<i>japonicum</i> MAXIM.	60
— <i>pallida</i> PERS.	58, 80	<i>Deutzia gracilis</i> S. et Z.	58
<i>Corylus heterophylla</i> FISH.	53	— <i>scabra</i> THUNB.	58
— <i>rostrata</i> AIT. var. <i>Seboldiana</i>		<i>Dianthus superbus</i> LINN.	56
MAXIM.	79	Diapensiaceæ.	85, 97, 100
Crassulaceæ.	58	<i>Diarrhena japonica</i> FR. et SAV.	47
<i>Crawfurdia fasciculata</i> WALL.	67	<i>Dicliptera crinita</i> NEES.	69
<i>Croomia japonica</i> MIQ.	78	<i>Diervilla floribunda</i> S. et Z.	69
<i>Crotalaria sessiliflora</i> LINN.	60	— <i>grandiflora</i> S. et Z.	69
Cruciferae.	58, 96, 99	— <i>japonica</i> DC.	70, 86
<i>Cryptogramme japonica</i> THUNB.	44	— <i>Middendorffiana</i> CARR.	70
<i>Cryptomeria-avenue.</i>	43	Dilleniaceæ.	83
<i>Cryptotenia japonica</i> HASSK.	65	Dioscoreaceæ.	52
<i>Cucubalus baccifer</i> LINN. var. <i>japo-</i>		<i>Dioscorea gracillima</i> MIQ.	52
<i>nicus</i> MIQ.	53	— <i>japonica</i> THUNB.	52
Cucurbitaceæ.	70	— <i>septemloba</i> THUNB.	52
<i>Cuscuta japonica</i> CHOIS. var. <i>thyr-</i>		<i>Diplazium japonicum</i> BEDD.	44
<i>soidea</i> ENGELM.	67	— <i>lanceum</i> PRESL.	44
<i>Cydonia japonica</i> PERS. var. <i>pyg-</i>		— <i>Wichuræ</i> METT.	44
<i>mæa</i> MAXIM.	59	Dipsaceæ.	70
<i>Cynanchum caudatum</i> MAX.	67	<i>Disporum sessile</i> DON.	51
<i>Cunoglossum furcatum</i> WALL.	67	— <i>smilacinum</i> A. GR.	51
Cyperaceæ.	49, 78, 95, 98	<i>Dontstemon dentatus</i> BGE.	58
<i>Cyperus difformis</i> LINN.	50	<i>Draba nomorosa</i> LINN. var. <i>hebe-</i>	
— <i>Haspan</i> LINN.	50	<i>carpa</i> LEDEB.	58
— <i>Iria</i> LINN.	50	<i>Drymoglossum carnosum</i> HOOK.	44
— <i>pilosus</i> VAHL.	50		
— <i>rotundus</i> LINN.	50		
<i>Cypripedium debile</i> REICHB f.	79	Eclipta alba HASSK.	71
<i>Cystopteris japonica</i> LUERSS.	44	Elæagnaceæ.	64
		<i>Elæagnus glabra</i> THUNB.	64
		— <i>macrophylla</i> THUNB.	64
		— <i>umbellata</i> THUNB.	64
Daphne pseudo-mezereum A. GR.	63	Elatinaceæ.	63
<i>Deciduous broad leaved tree-</i>		<i>Elatine orientalis</i> MELS.	45
<i>Region.</i>	72	<i>Elatostema umbellatum</i> BLUME var.	
<i>Deciduous forest.</i>	38	<i>majus</i> MAXIM.	54
<i>Deschampsia caryophylla</i> L.	98	<i>Eleocharis acicularis</i> R. BR.	50
<i>Desmodium Gardneri</i> BENTH.	60	— <i>japonica</i> MIQ.	50
— <i>laburnifolium</i> DC.	60	<i>Eleusine indica</i> GÆRTN.	47
— <i>Oldhami</i> Oliv.	60	<i>Elsholtzia cristata</i> WILLD.	67
— <i>polycarpum</i> DC.	60		

Enkianthus japonicus HOOK.	66
— nikensis MAKINO.	66
Ephiphianthus sachalinensis REICH. f.	79
Epilobium angustifolium LINN.	64
— dahuricum FISCH.	83
— japonicum HASSK.	83
— macranthum MORR. et DECNE.	57
Epipactis erceta WETTST.	52
— falcata WETTST.	52
— papillosa F. et S.	79
— Thunbergii A GRAY.	52
Equisetaceæ.	46
Equistum arvense L.	46
Eragrostis ferruginea BEAUV.	47
— pilosa BEAUV.	47
Ericaceæ.	66, 83, 96, 99
Erigeron annuus PERS.	71
— canadensis LINN.	71
Eriocaulaceæ.	51
Eriocaulon sexangulare LINN.	51
Eriochloa villosa KUNTH.	47
Euonymus alata K. KOCH.	62
— — — var. subtriflora. F. et S.	82
— europæa L. var. Hamiltoniana MAXIM.	62
— japonica THUNB.	62
— melanantha F. et S.	82
— oxphylla MIQ.	62
— sachalinensis MAXIM.	82
Eupatorium japonicum THUNB.	82
— Kirilowii TURCZ.	71
Euphorbiaceæ.	61
Euphorbia adenochlora MORR. et DCNE.	61
— helioscopia LINN.	61
— humifusa WILLD.	61
— pekinensis RUPE.	61
Euphrasia officinalis LINN. var. vul- garis LEDEB.	68
Eurya japonica THUNB.	63
Euscaphis staphyleoides S. et Z.	62

Ever-green Conifers.	33
— <i>Conifer-Region.</i>	88, 92
Fagaceæ.	53, 74
<i>Fagus</i> and <i>Quercus</i> in the winter.	72
— — in the summer.	73
— sylvatica LINN. var. Sieboldi MIXIM.	74
Fatoua pilosa GAUD. var. subcordata RUPE.	54
Festuca ovina LINN.	47, 78
Fimbristylis autumnalis RÆM. et SCH.	50
— diphylla VAHL.	50
— miliacea VAHL.	50
— squarrosa VAHL.	50
Fragaria elatior EHRL.	81, 96
— indica ANDR.	59
Fraxinus Bungeana DC. var. pubi- nervis WG.	66
— longicuspis S. et Z.	66, 75, 84
— Sieboldiana BLUME.	75
Fuji: its situation.	3
Galium aparine LINN.	69
— asprellum MICHX.	69
— brachypodium MAXI.	69, 86
— gracile BOE.	69
— paradoxum MAXIM.	86
— irifidum LINN.	69
— verum LINN. var. lacteum. MAXIM.	69
Gardenia florida LINN.	69
Gaultheria adenothrix MAXIM.	85
GENERAL ASPECT OF THE FLORA OF THE MOUNTAIN.	101
— — — VEGETATION.	21
Gentianaceæ.	67, 85
Gentiana scabra BGE. var. Buergeri MAXIM.	67
Geraniaceæ.	60, 81

Geranium eriostemon FISCH.	81	Hydrangea involucrata SIEB.	58
Geranium nepalense SWEET.	60	— paniculata SIEB.	58
— siebaldr MAXIM.	60	— scandens MAXIM.	58
— Wilfordi MAXIM.	60, 81	— Thunbergii SIEB.	81
Gerbera Anandria SCH. BIP.	71	Hydrocharitaceæ.	47
Gesneraceæ.	69, 68	Hydrocharis Morsus-ranæ LINN.	47
Geum japonicum THUNB.	59	Hydrocotyle asiatica LINN.	65
Glyceria remota FRIES. var. japo-		— rotundifolia ROXB.	65
nica HACK.	48	Hygrophila lancea MIQ.	69
— tonglensis CLARK.	48, 95	Hymenophyllaceæ.	77
Gnaphalium japonicum THUNB.	71	Hymenophyllum Wrightii BOSCH.	77
— muleiceps WALL.	71	Hypericum Ascyron LINN.	63
Goodyera pendula MAXIM.	52	— chinense LINN.	63
Goten station.	8	— erectum THUNB.	63
Gramineæ.	47, 77, 95, 98	— — — var. cæspitosum MAKINO.	63
Guttiferæ.	63	— japonicum THUNB.	63
Gymnadenia cucullata REICH. f.	79	— patulum THUNB.	63
— cyclochla KORS.	79		
Halenia sibirica BORK.	67, 85	Ilex crenata THUNB.	62
Halorrhagidaceæ.	64	— geniculata MAXIM.	82
Haloragis micrantha R. BR.	64	— integra THUNB.	62
Hamamelidaceæ.	69	— macropoda MIQ.	82
Hamamelis japonica SIEB. et ZUCC.	59	— rugosa FR. SCHM.	62
Hedera Helix var. colchica C.		— Sieboldi MIQ.	62
KOCH.	65	Illicium anisatum LINN.	56
Hedysarum esculentum LEDEB.	101, 99	Impatiens nolitangere LINN.	62, 82
Helwingia japonica DIETR.	65	— Textori MIQ.	62, 82
Hemerocallis fulva LINN.	51	Imperata arundinacea CYR. var.	
Higher Grass Region.	97	Kœnigii HACK.	48
Hosta corulea TRATT. form. lanci-		Indigofera pseudotinctoria MATSUM.	60
folia MATSUM.	51, 78	INFLUENCE OF FIRE ON THE PRESENT	
— Sieboldinna ENGL. var. longipes		VEGETATION.	107
MATSUM.	78	INTRODUCTION.	1
Houttuynia cordata THUNB.	52	Inula salicina LINN.	71
Hōyē lava.	6	Iridaceæ.	52
Humidity.	14	Iris japonica THUNB.	52
Humulus japonicus S. et Z.	54	— sibirica LINN. var. orientalis	
Hydrangea hirta S. et Z.	58	THUNB.	52
— Hortensia DC. var. acuminata		Isachne australis R. BR.	48
A. GR.	81	Isopyrum stoloniferum MAXIM.	57, 80
		Izu-Shichitō.	4

Juglandaceæ.	74	<i>Lespedeza bicolor</i> TUBCZ.	60
<i>Juglans Sieboldiana</i> MAXIM.	74	— <i>Buergeri</i> MIQ.	60
Juncaceæ.	51, 78	— <i>cyrtobotrya</i> MIQ.	60
<i>Juncus effusus</i> LINN. var. <i>decipiens</i>		— <i>junceæ</i> PERS. var. <i>sericea</i> HEMSLE.	60
BUCH.	51	— <i>pilosa</i> S. et Z.	60
<i>Juncus Maximowiczii</i> BUCH.	78	— <i>stricita</i> H. et A.	60
<i>Juniperus nipponica</i> MAXIM.	95	— <i>villosa</i> PERS.	60
— <i>rigida</i> S. et Z.	46	— <i>virgata</i> DC.	60
<i>Justicia procumbens</i> LINN.	69	<i>Leucothoe Grayana</i> MAXIM.	85, 96, 99
K		<i>Ligularia clivorum</i> MAXIM.	71
<i>Karasuishii.</i>	65	— <i>sibirica</i> CASS.	71
<i>Katsuragawa.</i>	40	<i>Ligusticum acutilobum</i> S. et Z.	65
<i>Kawaguchi.</i>	4	<i>Ligustrum Ibota</i> SIEB.	85
<i>Kadsura japonica</i> DINN.	56	— — — var. <i>ciliare</i> MAXIM.	85, 97
<i>Kenmarube lava.</i>	4	— <i>japonicum</i> THUNB.	66
<i>Kimbusen Mt.</i>	4	Liliaceæ.	51, 78
<i>Komagatake.</i>	4	<i>Lilium auratum</i> LINDL.	51
<i>Kraunhia floribunda</i> TAUB.	60	— <i>avenaceum</i> FISCH.	78
— <i>japonica</i> TAUB.	60	— <i>concolor</i> SALISB. var. <i>parthe-</i>	
<i>Kyellingia brevifolia</i> ROTTB.	50	nion ELW.	51
Labiataæ.	67, 86	— <i>cordifolium</i> THUNB.	51
<i>Lactuca debilis</i> MAXIM.	71	— <i>Hansonii</i> BAK.	78
— <i>denticulata</i> MAXIM.	71	— <i>Maximowiczii</i> REGEI.	51
— <i>Raddeana</i> MAXIM.	87	Linaceæ.	61
— <i>stolonifera</i> BENTH.	71	<i>Lindera glauca</i> BLUME.	57
— <i>Thunbergiana</i> MAXIM.	71	— <i>obtusiloba</i> BLUME.	57
— <i>versicolor</i> SCH. BIP.	71	— <i>præcox</i> BLUME.	57
<i>Lamium album</i> LINN.	68	— <i>umbellata</i> THUNB.	57
<i>Laportea bulbifera</i> WEDD.	54, 79	<i>Lindernia angustifolia</i> WETTS.	68
Lardizabalaceæ.	57	— <i>crustacea</i> (L) F. v. M.	68
<i>Larix leptolepis</i> GORD.	25, 48, 93, 94, 95	— <i>pyxidaria</i> ALL.	68
— <i>region.</i>	38, 93	<i>Linum possarioides</i> PL.	68
<i>Lathræa japonica</i> MIQ.	86	<i>Liparis auriculata</i> BLUME.	52
<i>Lathyrus Davidii</i> HANCE.	60	— <i>Krameri</i> F. et S.	79
Lauraceæ.	57	<i>Lipocarpa microcephala</i> KTH.	50
<i>Leersia oryzoides</i> SW. var. <i>japonica</i>		<i>Liriope graminifolia</i> BAK. var.	
HACK.	84	communis MAX.	51
Leguminosæ.	59, 99	<i>Listera cordata</i> R. BR.	79
<i>Leonurus macranthus</i> MAXIM.	68	— <i>Eschscholtziana</i> CHAM.	79
		<i>Litsea glauca</i> SIEB.	57
		Loganiaceæ.	66
		<i>Lonicera gracilipes</i> MIQ.	70

— japonica THUNB.	70	Mazus rugosus LOUR. var. macran-	
— Morrowii A. GR.	70	thus F. et S.	68
Lophatherum gracile BRONGN. var.		Melandryum firmum ROHR. . . .	56
elatum HACK.	48	Meliosma myriantha S. et Z. . 62,	75
Loranthaceæ.	54	— tenuis MAXIM.	62
Lotus corniculatus LINN. var. japo-		Memorialis hirta WEDD.	54
nicus RGL.	60	Mentha arvensis LINN. var. piper-	
Ludwigia prostrata BOXB. . . .	64	ascens HOLMES.	68
Luzula campestris DC. var. capitata		Menziezia pentandra MAXIM. . 85,	96
MIQ.	51	Metanarthecium foliatum MAXIM.	78
— campestris DC. var. sudetica		— luteo-viride MAXIM.	51
CLARK.	78	Microlepis hirsuta PRESL. . . .	44
— plumosa E. MEY.	51	— marginalis HANCE.	44
Lychnis Miqueliana ROHR. . . .	56	— Wilfordii MOORE.	77, 44
Lycium chinense MILL.	68	Milium effusum LINN.	48, 78
Lycopodiaceæ.	47, 77	Mimulus nepalensis BENTH. . . .	68
Lycopodium clavatum LINN. . . .	46	Minispermaceæ.	57
— cryptomerinum MAXIM.	77	Miscanthus Matsumure HACK. 48,	78,
— serratum THUNB.	46, 77		95
Lycoris sanguinea MAXIM.	52	— sinensis ANDR.	48
Lygodium japonicum SW.	46	Mishima.	6
Lysimachia clethroides DUBY. . .	66	Mitrasacme polymorpha R. BR. .	66
— decurrens G. FORST.	66	Monochoria Korsakowii REGEL. et	
— Fortunei MAXIM.	66	MAACK.	66
— japonica THUNB.	66	— vaginalis PRESL. var. plantaginea	
— vulgaris LINN.	66	SOLMS-LAUB.	51
Lytnaceæ.	64	Monotropa uniflora LINN.	85
Lythrum Salicaria LINN.	64	Moraceæ.	54
		Mosla grosseserrata MAXIM. . . .	68
Maccarinupri Mt.	13	— punctata MAXIM.	68
Machilus Thunbergii S. et Z. . .	57	Motosu.	4
Macleya cordata R. BR.	58	Muehlenbergia japonica STEUC. .	48
Macroclinidium robustum MAXIM.		— — — var. hakonensis HACK. 48,	78
.	71, 87	Myōkō Mt.	4
Magnoliaceæ.	56, 74	Myrsinaceæ.	69
Magnolia Kobus DC.	56		
— hypoleuca S. et Z.	56, 74	Nakano.	8
Maianthemum Convallaria WIGG.		Nanocnide japonica BLUME. . . .	54
et ROTH.	78	Nishi.	4
Mariscus Sieberianus NEES. . . .	50	Nasturtium montanum WALL. . .	58
Marlea platanifolia S. et Z. . . .	65	— palustre DC.	58

- Neottia microantha* LINDL. 52
Nephrodium decursivo-pinnatum
 BK. 44
 — *dilatatum* DESV. 44, 77
 — *Filix-mas* RICH. 44, 77
 — — — var. *lacerum* CHRIST. 44, 77
 — — — var. *polylepis* MAKINO. 77
 — — — var. *Sabæi* CHRIST. . . 77
 — *gracilescens* BLUME. var. *glandu-*
 ligerum BAKER. 77, 95
 — *hirtipes* HOOK. 77
 — *Maximowiczii* BAKER. . . . 44, 77
 — *Phegopteris* BAUMG. . . 44, 77, 95
 — *prolaxum* DIELS. 44
 — *sophoroides* DESV. 44
 — *Thelyteris* DESV. 44, 77
Niphobolus linearifolius HOOK. . 45
Nuphar japonicum DC. 56
Nymphæaceæ. 56

Oberonia japonica MAK. 52, 79
Odontosoria chinensis KUHN. var.
 tenuifolia MAKINO. 45
Oenanthe stolonifera DC. 65
Oldenlandia diffusa BOXB. . . . 69
 — *hirsuta* LINN. 69
Oleaceæ. 66, 75, 85, 97
Ōmiya. 6
 — *station.* 8
Onagraceæ. 64, 83
Onoclea sensibilis LINN. 45
Ophioglossaceæ. 46, 98
Oplisminus Barmanni BEAUR. . . 48
Orchidaceæ. 52, 79, 98
Orchis Chondradenia MAKINO. . 79, 98
Orobanchaceæ. 69, 97
Orixa japonica THUNB. 61
Osawa. 6
Osmunda regalis LINN. var. *japo-*
 nica MILDE. 46
Osmanthus Aquifolium B. et H. . 66

Osmorhiza japonica S. et Z. . . . 65
Osmundaceæ. 46, 77
Osmunda cinnmomea DINN. . . . 77
Ottelia alismoides PERS. 47
Oxalidaceæ. 60, 81
Oxalis acetosella LINN. 81
 — *corniculata* LINN. 60

Pæderia toniontosa BL. 69
Pæonia obovata MAXIM. 57
Papaveraceæ. 57, 80
Panicum acroanthum STEUD. . . . 48
 — *Crus-Galli* LINN. 48
 — *indicum* LINN. 48
 — *sanguinale* LINN. 48
 — — — var. *ciliare* DÖELL. . . 49
Parnassia palustris LINN. 58
Paris quadrifolia LINN. var. *obo-*
 vata RGL. 78
 — *tetraphylla* A. GR. 78
Paspalum Thunbergii KUNTH. . . 49
Patrinia palmata MAXIM. 86, 97
 — *scabiosæfolia* LINK. 70
 — *villosa* JUSS. 70, 86
Pedicularis gloriosa BISS. et Moor.
 68, 89
 — *resupinata* LINN. 65, 86
Pellionia radicans WEDD. 54
Pennisetum japonicum TRIN. . . . 49
Peracarpa circæoides H. FEE. . . 86
Pertya scandens SCH. BIP. 71, 87
Phalaris arundinacea LINN. 49
Phelledendron amurense RUPR. . 75
Philadelphus coronarius LINN. var.
 Satsumi MAXIM. 81
Phrangmites communis TRIN. . . . 49
Phyma leptostachya LINN. 67
Phyllanthus Matsumuræ HAYATA. 61
 — *Urinaria* LINN. 61
 PHYSIOGRAPHY. 3
Phteirospermum chinense BGE. . . 68

- Phyteuma japonium* MIQ. . . 70, 86
Picea 41, 42
Picea ajanensis FISCIL. 93
 — *Alcockiana* CARR. 93
Picea-formation. 39, 40
 — *hondōensis* MAYR. . . . 35, 89, 93
 — *polita* CARR. 10, 41, 91, 93
 — — CARR. pure stand of. . . . 40
 — *Veitchii*. 89
Picrasma quassioides BENN. . . . 61
Picris hieracioides LINN. var. *japo-*
nica RGL. 72
Pieris japonica D. DON. . . . 66, 85
 — *ovalifolia* D. DON. 66
Pilea petiolaris BLUME. 54
Pimpinella calycina MAXIM. . . . 65
Pinellia tripartita SCHOTT. . . . 50
Pinus densiflora S. et Z. . . . 46, 89
 — *parviflora* S. et Z. 93
 — *pumila*. 94
Pirolaceæ. 84, 97, 100
Pilola elliptica NUTT. var. *minor*
 MAXIM. 85
 — *media* Sw. 85, 97, 100
 — *renifolia* MAXIM. 85
 — *rotundifolia* L. var. *albiflora*
 MAXIM. 85
 — *secunda* LINN. 85
Plantaginaceæ. 69
Plantago major LINN. var. *asiatica*
 DCNE. 69
Plant-formation of Part I. . . 21
 — — — **II.** 29
 — — — **III.** 34
 — — — **IV.** 36
Platanthera angustifolia REICH. . 79
 — *conopsea* SCHLECHT. 98
 — *mandarinorum* REICH. f. . . . 79
 — *ophryoides* FR. SCHM. 79
Platanthera ussuriensis MAXIM. . 52
 — *Yatabei* MAXIM. 79
Platycodon gandiflorus DC. . . . 70
Plectranthus glaucocalyx MAXIM.
 var. *japonicus* MAXIM. 68
 — *inflexus* VAHL. 68, 86
 — *longitubus* MIQ. 68
Poa annua LINN. 49
 — *palustris* LINN. var. *strictula*
 HACK. 49
Pollinia imberhis NEES. *genuina*
 HACK. 49
 — *nuda* (TRIN.) KACK. 49
Polygalaceæ. 61
Polygala japonica HOUTT. . . . 61
Polygonatum latifolium DESF. var.
commutatum BAKER. 51, 78
 — *officinale* ALL. 52, 79
Polygonaceæ. 55, 96, 99
Polygonum alatum HAM. 55
 — *cuspidatum* S. et Z. . . . 55, 96, 98, 99
 — *debile* MEISN. var. *triangulare*
 MEISN. 99
 — *esculentum*. 55
 — *flaccidum* ROXB. 55
 — *muricatum* LINN. 55
 — *polymorphum* LEDEB. var. *japo-*
nicum MAXIM. 96, 99
 — *Posumbu* HAM. 55
 — var. *Blumei* (MEISN.) 55
 — *senticosum* MEISN. 55
 — *Sieboldi* MEISN. 55
 — *Thunbergii* S. et Z. 55
 — — — var. *Maackianum* MAXIM. 55
 — *viscosum* HAM. var. *vernicosum*
 MEISN. 55
Polypodiaceæ. 44, 76, 95
Polypodium hastatum THUNB. . . 45
 — *lineare* THUNB. 45
 — — — var. *distans* MAKINO. . . 77
 — — — var. *Onōi* MARINO. . . . 77
 — *senanense* MAXIM. 77
 — *trichomanoides* Sw. 77
 — *vulgare* LINN. var. *japonicum*
 FR. et SAV. 45

- Polypogon Higeaweri STEUD. 49
 Polystichum amabile (SM.) 45
 — aristatum PRESL. 45
 — crapedosorum DIELS. 45
 — falcatum var. Fortunei BAKEL. 63
 — tripterum SM. 45, 77
 — varium PRESL. 45
Pontederiaceæ. 51
 Populus tremula LINN. var. villosa
 WESM. 53
Portulaccaceæ. 55
 Portulacca LINN. 55
Potamogetonaceæ. 47
 Potamogeton crispus LINN. 47
 — Gamdichaudi CHAM. 47
 — polygonifolius POUR. 47
 Potentilla chinensis SER. 59
 — cryptotenæ MAXIM. 59
 — Kleiniana W. et A. 59
 Pourthiæa villosa DCNE. 59, 81
 Prairie. 38
 Prairie-Region (Basal region.) . 39, 45
 Prenanthes acerifolia (MAXIM.) . 87
Primulaceæ. 66, 85
 Primula japonica A. GRAY. 85
 Prunella vulgaris LINN. 68
 Prunus incisa THUNB. 81
 Pseudopyxis longituba FR. et SAV. 86
 Pteridium aquilinum KUHN. 45
 Pteris cretica LINN. 45
 — serrulata LINN. 45
 Pterocarya rhoifolia S. et Z. 74
 Pueraria Thunbergiana BENTH. . 60
 Pycnostemma chinensis BGE. 67
 Pyreus globosus REICHEB. var.
 stricta HOOK. 50
 Pyrus alnifolia LINN. 81
 — Aira EHRH. var. Kamaonensis
 WALL. 81
 — aucuparia GÆRTN. var. japonica
 MAXIM. 75, 76, 81
 — Matsumuræ MAKINO. 81
 Pyrus sambucifolia CH. et SCHL. 81, 96
 — Toringo SIEB. 59
Quercus acuta THUNB. 53
 — aliena BLUME. 74
 — cripula BLUME. 74
 — cuspidata THUNB. 53
 — dentata THUNB. 54
 — glandulifera BLUME. 54
 — glauca THUNB. 54
 — grosseserrata BLUME. 54
 — serrata THUNB. 54
Racomitrium canescens BRID. . . 100
 Rainfall. 8
Ranunculaceæ. 30, 56, 96, 99
 Ranunculus acer LINN. 57
 — aquatilis LINN. var. flacidus
 (PERS.) form. Drouetii HIERM. . 57
 — japonicus LUNGS. 57
 — scleratus LINN. 57
Rhamnaceæ. 62
 Rhamnus japonica MAXIM. var. ge-
 nuina MAXIM. 62
 Rhododendron indicum SW. var. α
 Kæmpferi MAXIM. 99
 — Metternichii S. et Z. 96
 — sinense SW. 66
 Rhus semi-alata MURR. var. Os-
 beckii DC. 62
 — Toxicodendron LINN. var. radi-
 cans MIQ. 62, 81
 — trichocarpa MIQ.
 Rhynchosia volubilis LOUR. 60
 Ribes alpinum LINN. var. japonicum
 MAXIM. 81
 Ribes ambiguum MAXIM. 81
 — fasciculatum S. et Z. 58
 Rodgersia podophylla A. GR. . 80, 81
Rosaceæ. 59, 75, 81, 96

<i>Rosa acicularis</i> LINDL.	96	<i>Saussurea Tanakæ</i> FR. et SAV. var.	
— <i>Luciæ</i> FR. et ROCH.	81	<i>phyllolepis</i> MAXIM.	87
— <i>multiflora</i> THUNB.	59	— <i>triptera</i> MAXIM.	87, 97
— <i>Wichuraiana</i> CREP.	59	<i>Sasa albo-marginata</i> MAK. et SHIB.	
<i>Rotala indica</i> KOEHNE var. <i>uliginosa</i>			49, 78
MIQ.	64	— <i>borealis</i> MAK. et SHIB.	49
<i>Rottboellia compressa</i> LIN. f. var.		— <i>paniculata</i> MAK. et SHIB.	78
<i>japonica</i> HACK.	49	Saxifragaceæ.	53, 81
<i>Rubus japonicus</i> FOCK.	96	<i>Saxifraga cortusæfolia</i> S. et Z.	58
— <i>morifolius</i> SIEB.	81	— <i>sarmentosa</i> LINN.	58
— <i>palmatus</i> THUNB.	59	<i>Scabiosa japonica</i> MIQ.	70
— <i>parvifolius</i> LINN.	59	Schizæaceæ.	46
— <i>phœnicolasius</i> MAXIM.	59	<i>Schizandra nigra</i> MAXIM.	56
— <i>rosæfolius</i> SM.	59, 81	<i>Schizocodon soldanelloides</i> S. et Z.	
— <i>Sieboldii</i> BL.	81		85, 97, 100
— <i>silvestris</i> S. et Z.	62	<i>Schizophragma hydrangeoides</i> S. et	
— <i>Thunbergii</i> S. et Z.	59	Z.	81
— <i>trifidus</i> THUNB.	59	<i>Scilla japonica</i> BAKER.	52
Rubiaceæ.	69, 86	<i>Scirpus Cyperinus</i> KUNTH. var. <i>con-</i>	
<i>Rumex Acetosa</i> LINN.	55	<i>color</i> MAK.	50
— <i>japonicus</i> MEISN.	55	— <i>erectus</i> POIR.	50
Rutaceæ.	61, 75, 82	Scrophulariaceæ.	68, 83
Sabiaceæ.	62, 75	<i>Scrophularia Patriniana</i> WYDL.	68
<i>Sagittaria sagittifolia</i> LINN.	47	— <i>indica</i> LINN. var. <i>japonica</i>	
Salicaceæ.	79, 53, 95, 99	MAXIM.	68
<i>Salix-Alnus</i> -region.	38, 95	<i>Securinega fluggeoides</i> MUELL ARG.	61
— <i>Caprea</i> LINN.	53, 95	<i>Sedum Alfreni</i> HCE.	58
— <i>gracilistyla</i> MIQ.	79	— <i>Kamtschaticum</i> FISCH.	58
— <i>japonica</i> THUNB.	53	— <i>lineare</i> KHUNB.	58
— <i>Sieboldiana</i> BL.	98, 99	— <i>Telephium</i> LINN. var. <i>pur-</i>	
— — — var. <i>bipinnata</i> FR. et SAV.	68	<i>pureum.</i>	58
<i>Salvia nipponica</i> MIQ.	68	— <i>verticillatum</i> LINN.	58
<i>Sambucus javanica</i> BL.	70	Selaginellaceæ.	46
— <i>racemosa</i> LINN.	70	<i>Selaginella involvens</i> SPRING.	46
<i>Sanguisorbia officinalis</i> LINN.	59	— <i>Krausiana</i> A. BR.	46
Santalaceæ.	45	<i>Senecio campestris</i> DC.	72
Saururaceæ.	52	— <i>flammeus</i> DC.	72
<i>Saussurea Maximowiczii</i> HERD.		— <i>namorensis</i> LINN. var. <i>Fuchsii</i>	
.	72, 87, 97	KOCH.	72, 87
		<i>Serissa foetida</i> COMM.	96
		<i>Serratula coronata</i> LINN.	72
		<i>Seseli Libanostis</i> KOCH.	83

<i>Setaria glauca</i> BEAUB.	49	<i>Stephanandra flexuosa</i> S. et Z.	59
— <i>pachystachys</i> F. et S.	49	— <i>Tanakæ</i> FR. et SAV.	59
— <i>viridis</i> BEAUV.	49	<i>Stereodon brachycarpus</i>	100
Shiraito station.	8	<i>Stewartia monadelpha</i> S. et Z.	27, 75
Shōji.	8	— <i>pseudocamellia</i> MAXIM.	75
— station.	8	<i>Streptopus ajanensis</i> TILING. var.	
Simarubaceæ.	61	— <i>japonica</i> MAXIM.	79
<i>Siphonostegia chinensis</i> BENTH.	68	<i>Struthiopteris germanica</i> WILLD.	45
<i>Sium Ninsi</i> LINN.	83	Styracaceæ.	65, 85
<i>Smilax china</i> LINN.	52	<i>Styrax japonica</i> S. et Z.	66, 85
— <i>herbacea</i> LINN.	52	SUMMARY.	109
— <i>Sieboldi</i> MIQ.	52	<i>Suzukawa</i>	53
Solanaceæ.	68	<i>Swertia chinensis</i> FRANCH.	67
<i>Solidago Virga-aurea</i> LINN.	72, 87, 67	— <i>bimaculata</i> CLARK.	67
.	100	Symplocaceæ.	66, 85
<i>Sophora japonica</i> . LINN.	91	<i>Symplocos cratægoides</i> HAM.	66, 85
<i>Sophora platycarpa</i> MAXIM.	60		
Sparganiaceæ.	46	Taxus baccata LINN. subsp. <i>suspi-</i>	
<i>Sparganium longifolium</i> TURCZ.	46	— <i>data</i> PILG.	46
<i>Spiranthes australis</i> LINDL.	52	Temperature.	11
<i>Spirea Blumei</i> G. DON.	59	<i>Teucrium japonicum</i> WILLD.	68
— <i>bracteata</i> MAXIM.	96	— <i>stoloniferum</i> HAM. var. <i>Mique-</i>	
— <i>bulbata</i> MAXIM.	59, 81, 96	— <i>lium</i> MAXIM.	68
— <i>japonica</i> LINN. f.	81	<i>Thalictrum aquilegifolium</i> LINN.	80
— <i>Thunbergii</i> SIEB.	59	— <i>minus</i> LINN. var. <i>elatum</i> LFCOY.	57
<i>Spodiopogon cotulifer</i> HACK.	49	— <i>simplex</i> LINN. var. <i>affine</i> RGL.	57
— <i>sibiricus</i> TRIN.	49	Theaceæ.	63, 75
<i>Sporobolus elongatus</i> R. BR.	49	<i>Thea japonica</i> NOIS.	63
<i>Stachys aspera</i> MICHX. var. <i>japonica</i>		<i>Themeda</i> Forskali HACK. var. <i>japo-</i>	
— MAXIM.	68	— <i>nica</i> HACK.	49
Stachyuraceæ.	63	<i>Thesium chinense</i> TURCZ.	54
<i>Stachyurus præcox</i> S. et Z.	63	Thymelæaceæ.	63
Stapyleaceæ.	62, 82	<i>Tiarella polyphylla</i> DON.	81
<i>Staphylea Bumalda</i> S. et Z.	82	Tiliaceæ.	75
<i>Stellaria aquatica</i> SCOP.	56	<i>Tilia cordata</i> MIQ.	75
— <i>florida</i> FISH. var. <i>angustifolia</i>		<i>Torilis japonica</i> DC.	65
— MAXIM.	56, 96, 99	<i>Torreya nucifera</i> S. et Z.	46
— <i>media</i> LINN.	56	<i>Trachelospermum jasminoides</i> LEM.	67
— <i>nemroum</i> LINN. var. <i>Bungeana</i>		<i>Trichosanthes cucumeroides</i> MAXIM.	70
— MAXIM.	79	<i>Trichomanes parvulum</i> POIR.	77
— <i>uliginosa</i> LINN.	56	<i>Tricyrtis macropoda</i> MIQ.	52
Stemonaceæ.	78		

- Trientalis europæa* LINN. 85
Tricyrtis hirta HOOK. 52
Trigonotis peduncularis BENTH. 67
Trillium Smallii MAXIM. 79
Tripetaleia paniculata S. et Z. 96
Tripterigyum Wilfordi HOOK. 62
Trisetum flavescens BEAUV. var.
papillosum HACK. 49
Trochodendraceæ. 75
Tsuga diversifolia MAXIM. 93
— *Sieboldiana* CARR. . 40, 48, 90, 93
Tylophora nikensis MATSUM. 67
- Ulmaceæ.** 54
Ulmaria multijuga (MAXIM.) 59
— *purpurea* (MAXIM.) 59
Umbelliferæ. 65, 83, 96
Urticaceæ. 54, 79
Urtica Thunbergiana S. et Z. 34
Usnea longissima ACH. 35
- Vaccinium bracteatum** THUNB. 66, 85
— *Buergeri* MIQ. 66
— *hirtum* THUNB. 66
— *Vitis-idaea* LINN. 96, 99
Valerianaceæ. 70, 86, 97
Veratrum album LINN. var. *Lobelia-*
num BAK. f. *japonica* BAK. 79
— *Maackii* REGEL. 52
— *nigrum* LINN. 52
Verbenaceæ. 67
Veronica can dina WALL. 69
— *longifolia* LINN. 69
— *spuria* LINN. 69
— *virginica* LINN. 69
Viburnum dilatatum THUNB. . 70, 86
— *erosum* THUNB. 70
— *furcatum* BLUME. 76, 86
— *Sieboldi* MIQ. 70
— *Wrightii* MIQ. 86
- Vicia Cracca* LINN. var. *japonica* MIQ. 60
— *pseudo-orobus* FISCH. et MEY. . 60
— *sativa* LINN. 60
— *unijuga* BR. 60
Violaceæ. 63, 83
Viola Bissetii MAXIM. 83
— *japonica* LANGSD. 63, 83
— *Maximowicziana* MAKINO. . . . 83
— *Patrinii* DC. var. *chinensis* GING. 63
— *pinnata* LINN. var. *Cherophyl-*
loides RGL. 83
— *silvestris* KIT. var. *grypoceras*
A. GR. 63
— *verecunda* A. GR. var. *semilu-*
naris MAXIM. 83
— *variegata* FOSCH. 83
Viscum album LINN. 54
Vitaceæ. 63, 83
Vitis Cignetiae PULL. 63, 64, 83
— *flexuosa* THUNB. 63
— *heterophylla* THUNB. 63
— *inconstans* MIQ. 63
— *Thunbergii* S. et Z. 63
- Wikstroemia Gampi** MAXIM. . . . 63
Wind-direction. 19
Woodsia manchuriensis HOOK. . . . 45
— *polystichoides* EATON. 46
Woodwardia radicans SW. 46
- Yamanaka.* 4
Yatsugatake, Mt. 4
Yenkyō-lava. 6
Yuania japonica MAXIM. 73
- Zanthoxylum pepiritum** DC. . . . 61
— *schinnifolium* S. et Z. 61, 82
Zelkova. 73
ZONATION OF THE PLANTREGIONS. 38
Zoysia pungens WILLD. 49

Supplements.

Chamæcyparis obtusa S. et Z.	<i>Hincki.</i>
Chamæcyparis pisifera S. et Z.	<i>Sawara.</i>
Zelkawa acuminata P.L.	<i>Keycki.</i>

著 作 權 登 錄

明治四十四年二月一日印刷
明治四十四年二月五日發行

著 作 者
早 田 文 藏
小石川區 原町百三十三番地

印刷兼發行者
丸 善 株 式 會 社
代表者 事務取締役 小柳津要人
東京市日本橋區通三丁目十四番地

印 刷 所
株會社 東京築地活版製造所
東京市京橋區築地二丁目十七番地

發 行 所
丸 善 株 式 會 社
東京市日本橋區通三丁目

丸善株式會社大阪支店 丸善株式會社京都支店
大阪市東區博勞町四丁目 京都市三條通麩屋町西入

ATLAS OF JAPANESE VEGETATION WITH EXPLANATORY TEXT

EDITED BY

Prof. M. MIYOSHI, D. Sc.

IN PORTFOLIO

Price, Each, Yen 1.00 - - Postage, Each 8 Sen.

CONTENTS

Set. I. (1-8). Cultivated and Semi-Cultivated Plants. I. „ II. (9-15). Vegetation of Nikko. I. „ III. (16-24). Vegetation of Riu-kiu. I. „ IV. (25-31). Cultivated and Semi-Cultivated Plants. II. „ V. (32-40). Vegetation of Nikko. II. „ VI. (41-46). Vegetation of Riu-kiu. II. „ VII. (47-53). Vegetation of Shikoku and its Vicinity. I.	Set. VIII. (54-62). Vegetation of Fuji. „ IX. 63-68). Vegetation of Sagahlin. I. „ X. (69-74). Vegetation of Sagahlin. II. „ XI. (75-79). Vegetation of Formosa. I. „ XII. (80-85). Vegetation of Formosa. II. „ XIII. (86-92). Coast Vegetation of Middle Japan. „ XIV. (93-101). Mountain Vegetation of Northern Japan.
--	---

INDEX PLANTUM JAPONICARUM SIVE ENUMERATIO PLANTARUM OMNIUM

EX INSULIS

KURILE, YEZO, NIPPON, SIKOKU, KIUSIU, LIUKIU, ET
FORMOSA HUCUSQUE CORNITARUM

Systematic et Alphabetice

DISPOSITA

ADJECTIS

SYNONYMIS SELECTIS, NOMINIBUS JAPONICIS,
LOCIS NATALIBUS.

AUCTORE

J. MATSUMURA.

Botanicus Professore in Universitate Imperiale Tokiense, Horti Directore.

IN 2 Vols.

DEMY 8 vo. CLOTH.

PRICE, ¥. 4.25

POSTAGE, 18 Sen.

VOL. 3 WILL APPEAR SHORTLY.

[illegible]

This book is due on the

QK369
.H413

16197

16197

LEGEND.

Al = Forest mainly represented by *Alnus*
 Al. Q. T. A. = Forest composed of *Alnus*, *Quercus* and small shrubs of *Taxus cuspidata*.
 A. P. T. = Forest mainly composed of *Alnus* and *Pinus densiflora*, with a small amount of *Fagus*.
 Co. B. = Coniferous forest with a considerable amount of *Betula*.
 F. = Forest mainly represented by *Fagus*.
 F. Q. = Forest mainly composed of *Fagus* and *Quercus*.
 F. Q. 1. 5. = Forest mainly composed of *Fagus* and *Quercus*, with a small amount of conifers.
 F. Q. Z. = Forest mainly composed of *Fagus* and *Quercus* with a small amount of *Zelkova acuminata*.
 F. C. = Forest mainly represented by *Fagus*, with a small amount of *Cladonia amurensis*.
 L. = Forest mainly represented by *Larix leptalepis*.
 L. 100% = Pure stand of *Larix leptalepis*.
 P. 70% = Forest mainly represented by *Pinus*.
 Pl. 1. L. = Afforested areas of *Pinus densiflora*.
 Pp. = Pure stand of *Pinus pitha*.
 Pl. Q. = Forest mainly represented by *Pinus densiflora* with a small amount of *Quercus serrata*.
 Pp. T. = Forest mainly composed of *Pinus pitha* and *Tsuga diversifolia* and *Sieboldi*.
 Q. = Forest mainly represented by *Quercus*.
 Q. P. = Forest mainly represented by *Quercus*, with a small amount of *Pinus densiflora*.
 T. = Forest mainly composed of *Tsuga Sieboldi* and *diversifolia*.
 T. 50% = Forest represented by *Tsuga*, with some *Abies* and *Pinus*.
 T. 90% = Forest mainly represented by *Tsuga*, with a small amount of *Abies*.
 T. A. = Forest composed of *Tsuga* and *Abies*.
 T. A. Q. C. = Forest composed of *Tsuga*, *Abies* and *Quercus*, with a small amount of *Chamaecyparis obtusa*.
 T. A. P. = Forest mainly composed of *Tsuga*, *Abies* and *Pinus*.
 T. F. Q. = Forest mainly represented by *Tsuga* with a small quantity of *Fagus* and *Quercus*.
 T. Q. P. = Forest mainly represented by *Tsuga* with some amount of *Quercus* and *Pinus densiflora*.
 Z. F. Q. = Forest with a considerable amount of *Zelkova acuminata* and some *Fagus* and *Quercus*.

● Base stations
 — Routes traversed by the author.
 --- Routes of public roads.

French Metres
 0 500 1000 1500 2000 2500 3000 3500 4000
 English Miles
 0 1 2 3 4 5 6 7 8 9 10

Botanical Map of Mt. Fuji

Compiled
by
B. HAYATA
Scale. 1:75,000