S
639.9
F2ub
1975
c. 2

FISH DIVISION
MONTANA FISH & GAME DEPT.
HELENA, MONTANA 59601

UPPER BLACKFOOT RIVER STUDY

A Premining Inventory of Aquatic and Wildlife Resources

STATE DOCUMENTS COLLECTION

APR 09 2003

MONTANA STATE LIBRARY
1515 E. 6th AVE.
HELENA, MONTANA 59620

Montana Department of Fish and Game
Environment and Information Division
in cooperation with
The Anaconda Company

By
Lester E. Spence
December 1975

MONTANA STATE LIBRARY

3 0864 1001 9019 1

ACKNOWLEDGMENTS

This project was made easier through assistance from several persons, including:

Mr. Ed Fickler, who provided interesting accounts of big game activities, past and present, in Alice Creek and surrounding areas; Dr. Robert Ream, University of Montana Forestry School, who was of great assistance in obtaining and constructing radio equipment used for elk telemetry; personnel in the Helena office of the U. S. Geological Survey who provided equipment and expertise used in stream-flow measurements; Dr. Don O. Loftsgaarden and Dr. Andrew Sheldon, University of Montana, who provided statistical services and advice; Dr. George Weisel, Zoology Department, University of Montana, who worked with the department in initiating the original water quality sampling program for the study; and personnel on the Lincoln Ranger District, Helena National Forest, who, in addition to providing alert and sure-footed horses, supplied pertinent data and hot coffee on cold winter sampling days. Their help is appreciated, as is that of Department of Fish and Game personnel who assisted on the project.

By agreement with the Montana Department of Fish and Game, the Anaconda Company funded approximately one-half the cost of this project for the first 2 years. This assistance, as well as that of Mr. Robert Dent, Anaconda Company biologist, is also appreciated.

TABLE OF CONTENTS

	Page
Acknowledgments	i
List of Tables.	iii
List of Figures	iv
Report Supplement	v
ABSTRACT.	1
RECOMMENDATIONS	2
ADDITIONAL STUDIES NEEDED	3
INTRODUCTION.	4
DESCRIPTION OF THE STUDY AREA	5
HISTORY AND METHODS	9
Water Quality.	9
Streamflow	13
Bottom Fauna	13
Fish	14
Wildlife	14
RESULTS	14
Water Quality.	14
Chemical Analyses	14
Biological Oxygen Demand.	26
Coliform Bacteria	26
Dissolved Oxygen.	27
Diurnal Variation of Dissolved Oxygen, Temperature and pH.	28
Miscellaneous Surface Water Temperatures.	28
Turbidity	28
Streamflow	31
Water Chemistry/Discharge Relationships.	35
U. S. Geological Survey Data	39
Acid Mine Seepage Surveys.	41
Metals in Stream Sediments	43
Bottom Fauna	44
Fish	49
Fish Population Surveys	49
Reproduction	54
Tag Returns	58
Metals in Fish Tissues.	61
Wildlife	61
General	61
Elk Distribution.	64
Elk Movement by Radio Telemetry	73
SUMMARY	81
LITERATURE CITED.	84
APPENDIX.	87

LIST OF TABLES

Table		Page
1	Number and percentage of heavy metal analyses with reported values less than the analytical detection level.	22
2	Values (mg/l) of total recoverable heavy metals which were greater than the detection level at selected sample stations	23
3	Maximum values and highest average values of metals reported at all 36 stations	24
4	Monthly DO levels from field analyses by the USGS at Pop's Place and Hogum stations	27
5	Variation in dissolved oxygen, percent saturation, water temperature and pH over a 24-hour period at six stations, 1968-1973	29
6	Summary of USGS analyses for metals (ug/l) at Pop's Place, water years 1969 and 1970.	40
7	Summary of USGS analyses for metals (ug/l) at Hogum, water years 1970-1973.	40
8	Partial summary of acid mine seepage water quality data obtained by the Anaconda Company in the Heddleston District, 1972 and 1973.	42
9	Levels of heavy metals in stream sediments and loss on ignition of organic matter, August 1973.	44
10	Diversity indices based on orders.	46
11	Diversity indices based on genera.	47
12	Water temperature and intragravel dissolved oxygen in Blackfoot River brown trout redds, 1972.	55
13	Water temperatures and dissolved oxygen in Alice Creek brook trout redds, 1972.	56
14	Characteristics of redds in Blackfoot River, 1973.	58
15	Tag returns from marked trout.	59
16	Change in size of trout recaptured a year after tagging. .	60
17	Highest spring counts of elk observed on Spencer Bar and vicinity, 1971-74.	71
18	Highest counts of elk observed on critical winter range, 1971-74.	72
19	Radio transmitter data	74
20	Elk banding data	74

LIST OF FIGURES

Figure		Page
1	Blackfoot River drainage. Numbers along main river are river miles above mouth.	6
2	Limits of glacial activity and direction of ice flow in Landers Fork and Alice Creek valleys	7
3	Location of proposed mining developments	10
4	Location of water quality stations	16
5	System for specifying water quality station locations. . .	17
6	Average concentrations (above detection level) of total recoverable iron, zinc, and copper at the seven main Blackfoot stations	25
7	Hydrographs of Blackfoot River at Pop's Place for water years 1969 and 1970.	32
8	Hydrographs of Blackfoot River at Hogum for water years 1971 and 1972.	33
9	Hydrograph of Blackfoot River at the Canyon station, 1972 and 1973, from periodic staff gage readings.	34
10	Relationship between discharge and dissolved solids, Blackfoot River at Pop's Place, water years 1969 and 1970 . . .	36
11	Relationship between discharge and dissolved solids, Blackfoot River at Hogum, water years 1971, 1972 and 1973 . . .	36
12	Relationship between discharge and bicarbonate, Blackfoot River at Pop's Place, water years 1969 and 1970.	37
13	Relationship between discharge and bicarbonate, Blackfoot River at Hogum, water years 1971, 1972 and 1973.	37
14	Relationship between discharge and sulfate, Blackfoot River at Pop's Place, water years 1969 and 1970.	38
15	Relationship between discharge and sulfate, Blackfoot River at Hogum, water years 1971, 1972 and 1973.	38
16	Location of bottom fauna collection stations	45
17	Fisheries stream survey and population estimate sample sites.	50
18	Principal elk winter ranges in the Blackfoot River valley above Lincoln.	63
19	Aerial sightings of one or more elk on winter and spring ranges, 1971-1974.	65
20	Location of proposed tailings pond with respect to the spring range and calving area locally known as Spencer Bar	66
21	Aerial sightings of newborn elk calves in June and July 1971-1974.	67
22	Landownership of spring range and calving areas.	68
23	Primary spring ranges and calving areas in the Alice Creek-lower Landers Fork vicinity.	69
24	Winter and spring ranges used by elk on private lands east of the continental divide.	70

REPORT SUPPLEMENT

As this report was being finalized, an event occurred in the Blackfoot River drainage which may alter some of the baseline information given in this report.

On or about June 19, 1975, a severe rainstorm added moisture to an already melting snowpack. The resulting heavy runoff, plus failure of the bypass facilities around the Mike Horse tailings dam filled the area behind the dam, overtopped and breached the structure. This sequence of events released approximately 100,000 tons of gray-colored, pyritic mine tailings deposits into the Blackfoot River.

The severity of the storm and high water conditions which washed out the access road greatly hampered repair efforts. Temporary repairs completed 9 days later allowed most of the water flowing through the tailings to be diverted around the tailings pond. However, seepage of water through the pond area continued to carry lesser amounts of tailings downstream 22 days after the dam failed. The Anaconda Company substantially rehabilitated the dam and installed new bypass structures in the fall of 1975 to prevent similar occurrences in the future. A new pond will be formed behind the repaired dam.

It is unknown how far downstream the tailings were carried, but for several days the gray-colored water was apparent at the mouth of Landers Fork, 15 miles downstream. High, muddy waters of Landers Fork obscured the dark gray color of the Blackfoot below their confluence to a point not noticeably different from the normal runoff conditions in the river. It is suspected the tailings were carried many more miles downstream in a more diluted form.

Cutthroat trout populations in the tailings pond and in the 300 feet of Beartrap Creek below the tailings dam were destroyed by the floodwaters. Also, the water quality, bottom fauna, fisheries and other data given in this report may have undergone changes due to mine tailings deposition in the stream channel. These changes will have occurred only in the main Blackfoot River and not in its tributaries. Biological and water quality sampling were conducted during the period by the Departments of Fish and Game and Health and Environmental Sciences and U. S. Forest Service. A summary report of the events and findings will be prepared by the Department of Fish and Game.

A possible result of the tailings pollution may be depression of aquatic biota further downstream than is presently occurring. This will not be known until further biological sampling is done on the river. If it is found that water quality and biological conditions are changed from that described in this report, further data will have to be collected to establish a new baseline of information to use in the event a mining operation begins in the Heddleston District.

Four fish ponds were constructed in the stretch of river between Anaconda Creek and Pass Creek during dam reconstruction. These ponds will improve the poor habitat conditions existing prior to dam failure, but only time will tell whether a fishery will develop in them.

ABSTRACT

This report is a compilation of chemical and biological data collected in anticipation of an open pit copper mining venture by the Anaconda Company in the headwaters of the Blackfoot River near Lincoln, Montana. Data are presented on stream water quality from 1968 through 1973, both laboratory and field analyses; streamflow; suspended sediment (turbidity); bottom fauna identified to lowest possible taxa; fish species present, their distribution and relative abundance, plus fish population estimates in selected stream sections; diurnal variations of dissolved oxygen; biological oxygen demand and coliform bacteria; continuous streamflow and temperature data and field and laboratory water quality data collected by the U. S. Geological Survey at two stations under contractual agreement with the Department of Fish and Game.

Information is presented on heavy metals concentrations in game fish and streambed sediments. Fish tagging and return data and data from spawning areas are also given.

General data on wildlife distribution, location and use of elk calving areas and winter ranges are presented. Elk migration patterns based on radio telemetry of three cow elk are given.

Water quality was generally good in the Blackfoot River and its tributaries. Degraded water quality existed in the extreme headwaters of the Blackfoot River due to acid mine drainage from the abandoned Mike Horse Mine and other tunnels. This lower water quality was reflected in bottom fauna populations which were poor in affected areas but good in areas unaffected by mine drainage. Diversity indices were calculated from bottom fauna data.

Principal fish species found in the study area were cutthroat trout, brook trout, brown trout and mountain whitefish. A total of 10 fish species was collected in the area. These included six salmonids, two suckers, one dace and one sculpin. Trout were found in all waters sampled except those with acid mine drainage in the Mike Horse Mine area. Intermittent fish populations were present in those streams which annually dry up due to natural lowering of the water table. Cutthroat and brook trout predominated in the Blackfoot River above Landers Fork, whereas brown trout predominated below. Mountain whitefish were present in most areas, but most were found in the Blackfoot River downstream from Poorman Creek.

Wildlife in the study area included elk, mule deer, white-tailed deer, moose, black and grizzly bears, mountain lions, bobcats, beavers and three species of mountain grouse. There have been unconfirmed reports of bighorn sheep in the area. Bald and golden eagles were also present, as were several species of waterfowl and furbearers.

Proposed mining facilities have the potential for causing water quality problems in the Blackfoot River and some of its tributaries.

A tailings pond is proposed to cover a portion of an elk calving area. Fisheries and wildlife habitat would be eliminated by the proposed open pit mine and other facilities. The increased human population in the small upper Blackfoot valley may have an additional adverse impact on fisheries and wildlife areas. Subdivisions along the Blackfoot River pose potential problems in maintaining the natural stream course due to channel and/or streambank modifications for flood protection.

Recommendations are given to prevent or mitigate damage to wildlife habitat with development.

An appendix to the report contains original basic data collected during the study.

RECOMMENDATIONS

1. The Anaconda Company or any other developer of the Heddleston properties should inform the Department of Fish and Game at the earliest stages of mining development so that environmental impacts can be determined before development plans proceed too far for reversal.
2. The mining developer should work closely with all federal, state and county agencies and local governing bodies to develop the mining complex with the least possible impact to the land, its wildlife and the social structure of the Lincoln valley.
3. Development of the mining complex will create the potential for serious sediment pollution problems as a result of road construction, removal of overburden, and ore excavation. Adequate measures to control stream sedimentation from any source should be undertaken by the mining developer. This is also true for any discharge of toxic metals which might occur.
4. A less damaging tailings pond site or, preferably, other waste disposal methods should be found. Hardscrabble Basin and Alice Creek should not be used for these purposes. Alternative and comparable tailings pond sites were not located in the area.
5. The proposed clarification pond is in a poor location from an environmental standpoint. Not only would the Blackfoot River have to be channelized for its construction, but the pond's location in the river floodplain has the potential for flood washout and spillage of metal deposits which would be intercepted from runoff from the stockpiles and mine area and transported to the pond. An alternative to pond use or a new location for the pond should be investigated.
6. Mining properties in the Alice Creek drainage not actually needed for mining purposes should be relinquished to the Department of Fish and Game for wildlife management in accordance with conditions agreed to in the March 16, 1970 letter from the Anaconda Company to the Director, Montana Fish and Game Department.

7. Follow-up water quality and biological monitoring should be initiated if extensive mining development should occur in the Mike Horse area. The cost of this program should be a responsibility of the mining developer.
8. Loss of elk calving range or other wildlife lands due to mining development, although irreplaceable, should be compensated for by acquisition of other wildlife lands, such as private land for winter big game range. This expense should be the responsibility of the mining developer.
9. Subdivision development in the Lincoln area poses a potential threat to the natural meandering course of the Blackfoot River by requiring streambank stabilization. Present and future planners should strongly consider the effect of river alteration on the stability of the river system and locate subdivisions away from riverbanks or other flood-prone areas.
10. The private lands shown as critical winter range in Figure 24 should be acquired if they become available from landowners willing to sell.
11. The feasibility of limiting motor vehicle access to the Spencer Bar calving area in May and June should be investigated.
12. Streamflow reservations in the project area should be submitted to the Department of Natural Resources under the 1973 Montana Water Use Act so that flows necessary to adequately sustain fish and wildlife can be obtained prior to mining development.

ADDITIONAL STUDIES NEEDED

1. Very little data were obtained on the Dolly Varden. Its status is practically unknown and more data should be gathered on its status, distribution and migratory habits.
2. Electrofishing indicated that mountain whitefish move within the river system on fall spawning migrations. More data should be gathered on the status and migratory habits of this species.
3. More data on spawning success, locations of spawning areas and the degree of spawning movement of resident cutthroat, brook and brown trout would increase the knowledge of these fisheries and should be obtained.
4. Additional data should be collected on fish populations in sections of the Blackfoot River and tributaries not sampled during this study to increase this knowledge for future planning purposes in the Blackfoot drainage.
5. The analysis of stream sediments for metal concentrations may shed new light on the origin of these metals and their effects on fish and wildlife. Additional sediment data should be obtained from the study area.

6. The three radioed elk studied during the period furnished good data on the migratory habits of elk which utilize the Scapegoat Wilderness. However, more elk should be banded with radio transmitters on both their winter and spring ranges to gain more complete data on these animals. The amount of information gained from radio telemetry is many times greater than that obtained from other marking methods and is well worth the additional expense of aerial tracking.
7. A study should be initiated to determine if there is competition for food between elk and domestic cattle and sheep utilizing the Spencer Bar area in the spring.
8. The effect of the high groundwater table and individual sewage disposal systems in the Lincoln area on Blackfoot River water quality is unknown. Data should be gathered on this potential problem.
9. A detailed chemical/biological study should be done on those portions of the Blackfoot River and tributaries affected by mine drainage to determine existing problem areas, what factors are depressing aquatic life, where the recovery zones are, and what causes the recovery. Such a study should include effects of mine drainage on water-oriented wildlife such as beavers, muskrats, mink, mice and shrews.

INTRODUCTION

This is the final report on the Upper Blackfoot River Study. A progress report was prepared by Dale (1971) under contractual agreement with the Department of Fish and Game. It included water quality data collected from February 1968 to March 1970. After issuance of the first report, a position of Environmental Planning Ecologist was established to continue gathering water quality data and expand the project to include biological data as well. This position was established in December 1970 under a project entitled "Hedderston Mining Development Project." The project title was later changed to "Upper Blackfoot River Study." Information contained in this second and final report includes all water quality and biological data obtained from February 1968 through August 1973.

At the time this study was begun in 1968, the Anaconda Company was actively considering development of an open pit copper and molybdenum mine in the Hedderston mining district. The district lies near the continental divide in the headwaters of the Blackfoot River near Lincoln, Montana. It was thus imperative that chemical and biological data be collected before development began to establish a "baseline" of existing ecological conditions. Mining on a commercial scale has not begun at this time, however, and it is unknown if or when development will occur.

The Blackfoot River drainage is an important hunting, fishing and recreational area. High water quality and other environmental standards must be maintained to perpetuate this use. State and federal regulations govern the maintenance of environmental quality, and adequate data are necessary to aid in development and enforcement of suitable standards.

DESCRIPTION OF THE STUDY AREA

The Blackfoot River flows 122 miles in a westerly direction from its source near the continental divide to its confluence with the Clark Fork River at Bonner, a small lumber industry town 7 miles east of Missoula, Montana (Figure 1). Principal tributary streams in downstream order are: Alice Creek, Landers Fork, Nevada Creek, North Fork, Monture Creek and Clearwater River. The source of the Blackfoot River is the confluence of Anaconda and Beartrap Creeks within the proposed mining area.

Mining in the upper Blackfoot valley began about 1865. Between 1865 and 1871, about \$7 million in high grade placer gold was taken from the gravels of Lincoln Gulch, 2 miles west of Lincoln. Mining activity at the source of the Blackfoot River dates from 1889 when William Heddleston and George Padbury discovered the Calliope lode. They built an arrastra on Pass Creek and recovered about \$11,000 in gold from the Calliope mine. Access to the Heddleston mining district was then limited to foot or horseback. Prospecting in later years produced very little gold, but some amounts of lead and other ores. Recovery of these metals required smelting, and the Mike Horse mill was eventually constructed on Mike Horse Creek in 1919. During the 1920's, as access and transportation improved, additional quantities of lead, silver and gold, and small but appreciable amounts of copper were mined and shipped to other smelting facilities. Mining continued intermittently through World War II, especially at the Mike Horse Mine. Operations at the Mike Horse ceased about 1953. In recent years the Anaconda Company has discovered a porphyry copper deposit in an area north of the Mike Horse Mine. There are several abandoned mineral prospects in the area, but there are no actively producing or developing mines (Pardee and Schrader 1933 and Mudge, et al. 1971). Core drilling and tunnel exploration by the Anaconda Company have recently occurred in the Mike Horse area. Several small prospectors are also active nearby.

Topography and geology of the Blackfoot valley near Lincoln were influenced by glacial activity. Glacial ice invaded the valley from the north at least twice. During the first advance ice moved down the broad valley north of Lincoln and was joined by ice from Landers Fork and Alice Creek which was moving down the Blackfoot valley. The main ice mass of the second advance moved down Copper Creek and Landers Fork, overflowed the east side of the Landers Fork valley in T 15N, R 7W, Section 18, continued southeastward and blocked the Alice Creek valley, forcing Alice Creek to cut a canyon in the rock on the east side of the valley. A unique area locally known as Spencer Bar in Hardscrabble Basin was formed by this glacial activity (Figure 2). (From Alden 1953 and Coffin and Wilke 1971. More detailed geological information is available in these references.)

Population of the Blackfoot valley is about 2,200 people. Agriculture and timber are the main industries. Most of the land adjacent to the river is private, mixed with some state and BLM land. There is very little national forest land adjacent to the river, but a good share of the mountainous terrain is on national forest land. Most surface water which makes up the Blackfoot River originates from tributaries which arise on and flow through national forest land.

Figure 1. Blackfoot River drainage. Numbers along main river are river miles above mouth.

Figure 2. Limits of glacial activity and direction of ice flow in Landers Fork and Alice Creek valleys. Shaded area is Spencer Bar (from Coffin and Wilke 1971).

Spencer Bar in Hardscrabble Basin was formed during the second glacial advance which spilled over from Landers Fork drainage (background). October 29, 1974.

The Blackfoot River is, for all practical purposes, "free-flowing" from source to mouth. There is a small diversion dam at the mouth of the stream which serves a wood products mill at Bonner. There is a small irrigation reservoir on Nevada Creek, a southern tributary. The Clearwater River flows through a series of natural lakes on its course down the valley. Waters of Landers Fork, North Fork and Monture Creek originate in wilderness or roadless areas. Logging activities are prevalent on both private and public lands in nearly every major subdrainage where this activity is allowed.

Hunting, fishing and other forms of recreation are important in the Blackfoot drainage. The Blackfoot River contains wild populations of rainbow, cutthroat, brook and brown trout, Dolly Varden and mountain whitefish. Elk, mule deer, white-tailed deer and black and grizzly bears inhabit the drainage, as do three species of mountain grouse (blue, Franklin's and ruffed).

A principal state highway (Route 200) runs the length of the Blackfoot drainage, providing access to recreation areas as well as a main transportation route for local and commercial vehicles between Missoula and Great Falls, two of the larger cities in the state.

There is currently a USGS (U. S. Geological Survey) gaging station near Bonner. Streamflow records at several discontinued stations are also available in USGS surface water records.

The basin has a drainage area of 2,290 square miles above the USGS gaging station 5.6 miles upstream from Bonner. The total drainage area is about 2,320 square miles.

Very little water is appropriated from the Blackfoot River at the present time. Water withdrawn from the stream is primarily used for private irrigation of pastures and haylands. An irrigation ditch about 1/8 mile below the mouth of Landers Fork diverts a significant amount of flow from the river during extreme low flow periods.

The proposed location of mining facilities determined the location of the sampling stations. The elevation of sampling stations ranged from 3,400 feet at Rainbow Bend to 5,050 feet at Pop's Place (Weizel and Newell 1970). Location of proposed mining facilities and a description of how the complex would operate are described by Coffin and Wilke (1971) from a report by the Montana Water Resources Board (1970). Figure 3 shows the location of these facilities as was known by those authors in 1971.

Previous studies related to impact of mining development in the Lincoln area have been done by Coffin and Wilke (1971) and Williams (1970).

HISTORY AND METHODS^{1/}

Water Quality

A water quality sampling program was begun by the Department of Fish and Game and University of Montana in February 1968, and was continued until March 1970. Five stations on the main Blackfoot River were sampled monthly, and tributary streams were sampled every 3 months for a total of 12-15 samples each month. Sampling at the Pop's Place station was discontinued in September 1968 when sampling there was assumed by the USGS under contract with the Department of Fish and Game. The USGS called this station "Blackfoot River near Lincoln, Montana."

Tributary sampling was discontinued in March 1970. Only the five main Blackfoot River stations continued to be sampled. These were: (1) Rainbow Bend, (2) Blackfoot Canyon camp (3.5 miles above Arrastra Creek), (3) Dalton, (4) Hogum and (5) Flesher. Field testing at all stations was also discontinued at this time. This schedule was maintained with few changes until February 1971 when sampling of Poorman Creek and Alice Creek again began. The Rainbow Bend station was discontinued at this time. Willow Creek (Flesher), and Hardscrabble Creek were added in October 1971 since they had never been sampled before.

^{1/} Specific methods for each sampling program described here can be found in the Appendices.

From: Coffin and Wilke 1971.
Base from U.S.G.S unedited advance prints

Figure 3. Location of proposed mining developments.

Analysis for "dissolved" as well as "total recoverable" heavy metals began in September 1971. This involved filtering and acidifying certain samples in the field. This practice was continued through the rest of the study. Analysis for the various forms of nitrogen and phosphorus began in March 1972 and continued until October 1972.

The USGS made periodic water quality and continuous temperature and discharge measurements at the gaging station at Pop's Place from October 1968 to September 1970 when the station was discontinued. A new station was established by USGS in October 1970 downstream at the Hogum Creek road bridge and was called "Blackfoot River below Alice Creek, near Lincoln, Montana." This station was operated until October 1975. The USGS collected continuous temperature and streamflow data, daily and monthly water quality data, and quarterly trace metals data at one or the other of these stations through September 1972. Beginning with the 1973 water year, quarterly analyses for trace metals were discontinued. Continuous streamflow and temperature measurements, and monthly water quality analyses were continued. During the 1974 water year, only continuous discharge and temperature data were collected.

USGS gaging station "Blackfoot River below Alice Creek, near Lincoln" (Hogum sample station) looking downstream. March 14, 1972.

Water quality samples were collected daily by a USGS observer at one or the other of the gaging stations through September 1972 when they were terminated. These daily samples were analyzed for specific conductance. On the basis of specific conductance, one to three samples representing extremes in total dissolved solids were selected and analyzed for common constituents.

Data from the two gaging stations are published in USGS Water Resources Data for Montana, Part 1 - Surface Water Records and Part 2 - Water Quality Records for the respective water years.

Water quality analyses not made by the USGS were performed by the State Department of Health and Environmental Sciences chemistry laboratory, Helena, Montana 59601.

BOD (Biochemical Oxygen Demand) sampling began at Canyon, Dalton and Hogum stations on the main river in February 1972. Samples were collected by the writer about every month and analyzed by the USGS in Helena. Beginning in October 1972 the USGS themselves collected and analyzed these samples monthly at the above three stations plus one additional station between Landers Fork and Lincoln (below 7-Up Pete Creek). Collections for coliform (total and fecal) determinations were also begun at these four stations by the USGS in October 1972.

In addition to irregular field determinations of DO (Dissolved Oxygen), diurnal DO measurements were made twice, once in September 1971 at six stations, and once in June 1972 at three stations. DO was determined and water temperatures were recorded every 4 hours for a 24-hour period at each station.

Intragravel DO was measured in brown and brook trout spawning redds in November 1972. Standpipes similar to those described by Terhune (1958) were placed in the spawning gravels at one location in the Blackfoot River below Poorman Creek and at the mouth of Alice Creek. Water velocities, stream discharge, surface water temperature and DO were measured, and DO and water temperatures were determined from water samples taken from within the standpipes. The approximate number of spawning redds was recorded in 1972 and 1973 in the Blackfoot River below Poorman Creek, and in 1972 in the Bartlett section on Alice Creek. Brown and brook trout were observed actively spawning over several of the redds in both places.

Measurements of turbidity were begun on a more regular basis in the spring of 1972 and continued in 1973. Sampling was intensified at the main monitoring stations during the spring runoff period, and reduced at other times of the year when the water was cleaner and fluctuated less.

Sediment samples from approximately the upper 2 inches of substrate in deposition zones of several streams and the main river were collected in 1973, stored in labeled cloth bags and analyzed for common heavy metals by the Montana Bureau of Mines and Geology laboratory in Butte.

The Anaconda Company conducted acid mine seepage surveys in the Heddleston District in 1972 and 1973. Some of their results are presented in this report.

Streamflow

Measurements of streamflow at locations other than USGS gaging stations began in March 1972. Porcelain-enameled iron staff gages were installed with assistance from USGS personnel at five locations: Canyon, Dalton and Flesher on the main river and on Poorman and Alice Creeks. In addition, a reference point was established on the Highway 200 bridge crossing Landers Fork and water height was measured with a steel tape and metal weight. The reference point was later discontinued due to channel instability at the measuring site. For the same reason there was no suitable location to install a staff gage at this site.

Bottom Fauna

Bottom fauna collections were begun in July 1968 at six Blackfoot River stations. Collections were also made in April and October 1969 at most of these same stations. These samples were collected in riffle habitat with a 2-foot-wide Needham kick screen which sampled a surface area of ± 4 square feet. One sample was collected at each station per sample period. A more intense sampling program began in March 1971. Quarterly collections were made in February, May (if water levels permitted), August and October at the following stations: Pop's Place, Flesher, Hogum, between Landers Fork and Lincoln, Dalton, Canyon and Rainbow Bend on the main river, plus Alice Creek and Poorman Creek. Periodic collections were also made on Landers Fork. A square-foot stream bottom sampler, modified after Waters and Knapp (1961) and having 38 meshes per inch (.018-inch mesh opening), was used for collection. This mesh size probably retained more insects than the U. S. Standard No. 30 sieve (.023-inch mesh size) used to define macroinvertebrates (APHA, ANWA, WPCF 1971). Three 1-square-foot samples were collected at each station in riffle habitat. The insects were preserved in glass jars containing 10 percent formalin. The samples were sorted to order. Volumes and numbers of insects in the samples were determined at the department's pollution control laboratory, Helena. The samples were then further sorted, recounted and identified to the lowest possible taxonomic group by a trained aquatic insect taxonomist working under contract with the Department of Fish and Game.

Insect collections made in 1968 and 1969 were identified to the lowest taxonomic group, but unfortunately the numbers of individuals in each group were not determined. Enumeration by lowest taxonomic group began with the March/April 1971 collections. Additional collections were made in August and October 1971, February, May, August and October 1972, and May 1973, but all of them have not been identified or enumerated below their orders.

Riffle areas were selected for sampling because they are the preferred type of habitat for bottom organisms and because life on riffles is more abundant than in pools. Riffles thus provide more information on which to base interpretation of conditions. Hard surfaces such as rocks and gravel are preferred attachment surfaces for organisms (Kittrell 1969). Also, it was felt that riffle data alone would be sufficient to show future changes in the quality or quantity of organisms due to pollution.

Needham and Usinger (1956) showed that a large number of samples was needed at a single station to give reliable (95 percent confidence) data on estimated total wet weight and estimated total numbers of bottom organisms. However, they found only two or three samples per station were needed to give 95 percent confidence that at least one member of each of the commonest genera of bottom insects would be present. It was therefore felt that three 1-square-foot samples would be adequate to establish insect diversity at the sample stations and would also be a sample size which could be managed.

Fish

Fish population estimates were made on eight sections of the Black-foot River: Pop's Place, Flesher, below Willow Creek (Flesher), Hogum, Stemple, Poorman-Dalton and Canyon. Population estimates in one section each of Alice Creek and Poorman Creek were also made. Trout over 9 inches in total length were marked with individually numbered white Floy-T tags to determine any significant movements. All fish were captured with a Fisher-Shocker (Model FS-101) or Smith-Root backpack electrofishing equipment. The Fisher-Shocker was floated downstream in a boat rigged for electrofishing.

Additional survey shocking to establish fish species composition, their relative numbers and size was also accomplished on the following streams: Beartrap Creek - 3 sections; Pass Creek - 2 sections; Anaconda Creek - 1 section; Shoue (Shave) Gulch - 1 section; Cadotte Creek - 1 section; Willow Creek (Flesher)- 3 sections; Alice Creek - 4 additional sections; Hardscrabble Creek - 3 sections; Hogum Creek - 1 section; Copper Creek - 2 sections; Landers Fork - 2 sections.

Twenty-five fish samples were collected in 1972 by electrofishing and analyzed for the more common heavy metals by EPA (U. S. Environmental Protection Agency).

Wildlife

Wildlife data collection centered on aerial and ground observations of elk using the spring calving areas on Spencer Bar, the Alice Creek drainage, portions of Landers Fork, and winter ranges in the upper Black-foot drainage and in the Middle Fork Dearborn River drainage east of the continental divide. Three cow elk were captured on their winter range east of the divide in April 1973 and fitted with radio transmitter collars. Their movements were followed on the ground and from aircraft during annual migrations to spring and summer range and return to winter range. Other wildlife observations were also made in the study area.

RESULTS

Water Quality

Chemical Analyses

Results of chemical analyses performed by the State Chemistry Laboratory on water samples from the 7 main river stations and 29 tributaries are given

in Appendix A (pg. A-1). The location of maximum and minimum values of these chemical properties and constituents are given in Appendix B. Laboratory detection levels for the constituents are given in Appendix C. Results of laboratory analyses excluding those which had values less than the analytical detection level are given in Appendix D.

Locations of the 36 stations sampled during the study are shown in Figure 4. Numbers correspond to the station locations given in Appendix E. The system used to describe station locations is based on the system of land subdivision used by the Bureau of Land Management and is shown in Figure 5.

Average total hardness, total alkalinity, bicarbonate and calcium values increased downstream between the Pop's Place and Canyon stations (river mile 119.4 to 84.7). However, values of these constituents were less at the Rainbow Bend station (river mile 9.7) than at the Canyon station.

Average total residue increased between Pop's Place and Canyon, but these values fluctuated among the six stations in the stream reach. Total residue was also less at Rainbow Bend than at Canyon.

The lower concentrations of hardness, alkalinity, residue and calcium at Rainbow Bend may be due to the entry of waters of East and West Twin Creeks upstream and on the same side of the river as the sample station. These two streams had the lowest average values of these constituents of any of the waters sampled.

The remaining chemical constituents analyzed for did not show any particular distribution pattern in the values obtained.

Average sulfates decreased in a downstream direction, but increased again at Rainbow Bend. Sulfates were higher at Pop's Place (average 41 mg/l) than at any other station sampled. This is due to acid mine drainage from the Mike Horse area. Sulfate is commonly formed from the oxidation of sulfide minerals which are common in the Mike Horse area. The sulfur in sulfide minerals such as pyrite (FeS_2) or chalcopyrite (CuFeS_2) is oxidized to sulfate (SO_4^{2-}) ions in water (Coffin and Wilke 1971). The second highest average sulfate level (22 mg/l) found was at Flesher station on the main river. This station is 4.8 miles downstream from Pop's Place. Average sulfate levels in tributaries ranged from zero to a maximum of 11 mg/l at Elk Creek and Union Creek. Both streams flow through agricultural lands. Other average values were: Willow Creek (Flesher) 7.3 mg/l; Hogum Creek, 8.1 mg/l; and Poorman Creek, 9.6 mg/l - all of which flow through old mining areas; and Spring Creek, 8.8 mg/l and Willow Creek, 9.9 mg/l which flow through agricultural land.

The major cation was calcium and the major anion was bicarbonate at all 36 stations sampled. However, when sulfate levels increased, such as at Pop's Place and Flesher, bicarbonate levels were reduced accordingly.

Bicarbonate alkalinity ranged from 6-230 mg/l at all stations. West Twin, East Twin, Sandbar Creek and Pop's Place had the lowest bicarbonate levels in that order. Average bicarbonate at main river stations ranged

Figure 4. Location of water quality stations. Shaded portion is principal study area. See Appendix E (p. A-34) for station descriptions.

Wells, springs, water-sampling locations and stream-gaging locations are assigned numbers based on the system of land subdivision used by the U. S. Bureau of Land Management. The number consists of 12 characters and describes the location by township, range, section and position within the section. The figure below illustrates the numbering method. The first three characters of the number give the township, the next three characters the range. The next two numbers give the section number within the township, and the next three letters describe the location within the quarter section (160-acre tract), the quarter-quarter section (40-acre tract), and the quarter-quarter-quarter section (10-acre tract).

These subdivisions of the 640 acre section are designated a, b, c, and d in a counterclockwise direction, beginning in the northeast quadrant. If there is more than one feature in a 10-acre tract, consecutive digits beginning with 2 are added to the number. For example, if a water-quality sample were collected in section 21, T 9N, R 20W, it would be numbered 09N20W21DAA2. The letters DAA indicate that the well is in the NE $\frac{1}{4}$ of the NE $\frac{1}{4}$ of the SE $\frac{1}{4}$, and the number 2 following the letters DAA indicates there is more than one water-quality sampling location in this 10-acre tract.

Figure 5. System for specifying water quality station locations.

from 61 at Pop's Place to 187 at Canyon. Average tributary levels ranged from 20 mg/l at West Twin Creek to 215 mg/l at Union Creek. West Twin Creek flows through boulder fill, whereas Union Creek flows through an alluvial valley and is used for irrigation of agricultural land (Weisel and Newell 1970). Union Creek does not have a good sport fish population, although other factors, such as excessive turbidity and water temperature, may be responsible.

Water quality station on Blackfoot River at Hogum Creek road, river mile 110.7. March 14, 1972.

Chloride and fluoride were found in lesser quantities than any of the other major ions.^{1/} Average fluoride levels were almost always lower than chloride levels. Out of 36 stations sampled, only Willow Creek (Fletcher), Hardscrabble Creek, Black Diamond Creek and Horsefly Creek had higher average fluoride than chloride levels (chloride was zero whereas fluoride was 0.10, 0.12, 0.20 and 0.10 mg/l, respectively). The maximum chloride levels found were 25 mg/l at Rainbow Bend and in tributaries 11 mg/l at Pass, Kleinschmidt and Union Creeks. The maximum fluoride level found on the main river was 0.50 mg/l at Pop's Place, and in tributaries 0.52 mg/l at Spring Creek.

^{1/} The major ions are calcium, magnesium, sodium, potassium, bicarbonate, carbonate, chloride, fluoride, sulfate.

The highest average chloride level on the main river was 4.8 mg/l at Rainbow Bend and in the tributaries 6.6 mg/l in Union Creek.

The highest average fluoride level found on the main river was 0.54 mg/l at Rainbow Bend, and in the tributaries 0.27 mg/l at East Twin Creek. Fluoride bearing rocks have a low solubility. The amount of fluorine (source of fluoride ions) in rocks is greater than the amount of chlorine; however, when all forms of both elements are considered, chlorine is much more abundant in the outer part of the earth (Hem 1970). Therefore, chloride would be expected to occur in higher concentrations than fluoride in natural water.

Total alkalinity values ranged from 35-189 mg/l on the main river and 5-226 mg/l in tributary streams. The lowest average value was found at Pop's Place (51 mg/l) on the main river. Lowest average tributary value was 17 mg/l at West Twin Creek. Average alkalinity values were over 100 mg/l at 5 of 7 main river stations (Pop's Place and Flesher were exceptions). In tributary streams, 14 of 29 stations had average alkalinity values above 100 mg/l. The remaining 15 had average values from 17-98 mg/l.

Sample station on Blackfoot River at Blackfoot Canyon Camp, river mile 84.7. August 24, 1971.

Total residue (dissolved solids) ranged from a minimum of 60 mg/l at Rainbow Bend to a maximum of 261 mg/l at Dalton on the main river and in tributary streams from 10 mg/l at West Twin Creek to 288 mg/l at

Hardscrabble Creek. Average main river values ranged from 114 mg/l at Flesher to 210 mg/l at 7-Up Pete. Average tributary values ranged from 21 mg/l at West Twin Creek to 208 mg/l at Union Creek.

Total hardness ranged from 60-340 mg/l at Pop's Place and Dalton, respectively, on the main river. Tributary values ranged from 7-310 mg/l in West Twin Creek and Union Creek, respectively. Average total hardness was lowest on the main river at Pop's Place (112 mg/l) and highest at Dalton (168 mg/l). Average tributary levels ranged from a low of 13 mg/l at East Twin Creek to a high of 194 mg/l at Union Creek. Hardness of water is due primarily to the presence of the cations calcium and magnesium. Water hardness is important to fish and aquatic life because toxic metals (copper, lead, zinc, etc.) are more toxic in soft water than in hard water. According to two references, the following values are used to classify water hardness (Brown, et al. 1970 and Dufor and Becker 1964).

<u>Total Hardness (As mg/l CaCO₃)</u>	<u>Classification</u>
0 - 60	Soft
61 -120	Moderately hard
121 -180	Hard
More than 180	Very hard

Water in the Blackfoot River drainage, therefore, ranges from soft to very hard.

Very little carbonate (CO₃) was found in any of the waters on a regular basis. Highest values were 12 mg/l at Rainbow Bend and 24 mg/l in Poorman Creek. Carbonates are less soluble in water than chlorides, sulfates and nitrates (McKee and Wolf 1963). They are not detectable when water has a pH of 8.3 or less.

Field pH levels determined with the Hach Model DR-EL field test kit showed a range of values between 6.6-8.7 for all waters sampled. The lowest values found were at Pop's Place on the main river (6.9), and in West Twin Creek (6.6), East Twin Creek (6.9) and Sandbar Creek (6.9).

Pop's Place and Sandbar Creek receive waters from mine shafts and tailings, which account for the low pH. The two low pH values in East and West Twin Creeks were not accounted for.

The next lowest pH was 7.1 at West Twin Creek. Pass Creek and Willow Creek both had pH 7.2 as low levels. Union Creek had a low level of pH 7.3. Highest pH levels were pH 8.7 at Dalton on the main river and the same in Poorman and Elk Creeks. Fourteen of 30 stations (6 main river stations and 24 tributaries) had maximum pH levels of 8.6. Most natural waters are buffered by CO₂ reactions, and their most effective buffering action occurs between pH 6.0-8.5. Unpolluted surface water generally has a pH between 6.5-8.5 (Hem 1970).

Main river levels of calcium ranged from a minimum of 10 mg/l at Pop's Place to a maximum at Canyon of 56 mg/l. Minimum tributary values

were 2 mg/l at both East and West Twin Creeks, and the maximum was 68 mg/l in Belmont Creek.

Magnesium ranged from zero at Canyon to 39 mg/l at Hogum on the main river, and in tributaries from zero at East and West Twin Creeks, Clearwater River and Pass Creek, to 51 mg/l in Union Creek. Magnesium was usually lower than calcium.

Sodium and potassium were reported together in laboratory data. Sodium + potassium values were highest on the main river at Canyon (33 mg/l) and lowest (zero) at all stations except 7-Up Pete (2.2 mg/l). In tributary streams, maximum values were 22 mg/l in Kleinschmidt and Elk Creeks and minimum values were zero in 23 of the 29 streams.

Alice Creek sample station as viewed upstream from Highway 200 bridge. August 1972.

Analyses of water samples for iron, arsenic, zinc, lead and copper revealed that in a significant number of samples these metals were not detected (Table 1).

Table 1. Number and percentage of heavy metal analyses with reported values less than the analytical detection level.

	Total Recoverable														
	Iron			Arsenic			Zinc			Lead			Copper		
	N	N<dl	%	N	N<dl	%	N	N<dl	%	N	N<dl	%	N	N<dl	%
Tributaries	198	108	54	94	80	85	204	152	74	62	54	87	202	165	82
Main River	209	91	44	81	67	83	209	102	49	53	47	89	209	162	78
Combined	407	199	49	175	147	84	413	254	62	115	101	88	411	327	80

N = total analyses

N<dl = number of analyses with reported values less than the detection level

% = percent of total analyses with reported values less than the detection level

Iron was detected more frequently than the other metals. Zinc, copper, arsenic and lead were the next most frequently detected in that order. Tributary streams, as a whole, had fewer detectable metals than did the main river. An exception was in the slightly more frequent occurrence of lead in tributary streams than in the main river. The occurrence of arsenic and lead were nearly the same in the main river and tributaries, probably because each of these metals was found infrequently and in very small amounts at all stations sampled.

By excluding the metals analyses having reported values less than the detection level, a new range and average value was calculated for each station (see Appendix D - p. A-18). The purpose is to show that heavy metals were not high at these stations, even when only those samples where the metal was detected are considered. These values are summarized in Table 2 for five selected stations in the upper 11 miles of the river. The location of these stations in the proposed mining complex (see Figures 3 and 4) makes them particularly suitable for showing any changes in water quality.

Table 2. Values (mg/l) of total recoverable heavy metals which were greater than the detection level at selected sample stations.

	B.F. @ Pop's Place (RM 119.4)	B.F. @ Fletcher (RM 114.6)	Alice Cr. (RM 113.2-0.9)	Hardscrabble Cr. (RM 112.5-0.5)	B.F. @ Hogum (RM 110.7)
Iron	(7)-87% ^{1/}	(32)-67%	(14)-50%	(8)-89%	(7)-27%
Range	.02-.27	.01-.50	.01-.90	.02-.28	.08-.60
Average	.12	.12	.18	.11	.26
Arsenic		(3)-15%	(3)-20%	(1)-12%	
Range	All <d1 ^{2/}	.001-.002	.001-.002	(.002) ^{3/}	All <d1
Average		.001	.002	-	
Zinc	(8)-100%	(40)-85%	(12)-40%	(4)-44%	(6)-24%
Range	.01-.44	.02-.94	.01-.72	.01-.04	.01-.03
Average	.22	.11	.11	.02	.02
Lead		(2)-13%	(4)-27%	(1)-12%	
Range	All <d1	.01-.01	.01-.01	(.01)	All <d1
Average		.01	.01	-	
Copper		(12)-25%	(9)-30%	(5)-56%	(4)-16%
Range	All <d1	.01-.18	.01-.06	.01-.02	.02-.06
Average		.03	.02	.01	.04

1/ Number and percent of total analyses showing detectable levels of the constituent.

2/ <d1 = less than detection level.

3/ Number in parentheses indicates only one sample value obtained.

The highest iron value detected was .90 mg/l in Alice Creek. However, this was an extreme value as shown by the average of .18 mg/l. Arsenic, when detected, was present in very small amounts. A maximum of .002 mg/l occurred at Flesher, Alice Creek and Hardscrabble Creek. The highest zinc value (.94 mg/l) was at the Flesher station on the main river. Again, the average of .11 mg/l shows this was an extreme value. The most lead detected was .01 mg/l at Flesher, Alice Creek and Hardscrabble Creek. The maximum copper value found was .18 mg/l at Flesher; however, the average of this station was .03 mg/l.

The maximum values and the highest average values of the above metals reported at all 36 stations in the study area are given in Table 3.

Table 3. Maximum values and highest average values of metals reported at all 36 stations.

	Total Recoverable Metal	Maximum Value (mg/l)	Reported At	Highest Average Value ^{1/} (mg/l)	Reported At
Blackfoot R.	Iron	1.5	Canyon	.24	Rainbow Bend
	Arsenic	.008	Rainbow Bend	(.008) ^{2/}	Rainbow Bend
	Zinc	.94	Flesher	.22	Pop's Place
	Lead	.01	Flesher, Dalton, Canyon	.01	Flesher, Dalton, Canyon
	Copper	.07	Dalton	.04	Hogum, Rainbow Bend
Tributaries	Iron	2.5	Sandbar Cr.	(.50) ^{2/}	North Fork
	Arsenic	.006	Beaver Cr.	(.006) ^{2/}	Beaver Cr.
	Zinc	.72	Alice Cr.	.24	Landers Fork
	Lead	.02	Poorman Cr.	.02	Poorman Cr.
	Copper	.12	E.Twin Cr.	(.12) ^{2/}	E.Twin Cr.

^{1/} Includes only those stations having more than one analysis performed.

^{2/} Numbers in parentheses indicate this was the only value greater than the detection level at this station.

Figure 6 illustrates the distribution of average concentrations of total recoverable iron, zinc and copper at the seven Blackfoot River stations based only on values greater than the detection level. Arsenic and lead

Figure 6. Average concentrations (above detection level) of total recoverable iron, zinc, and copper at the seven main Blackfoot River stations.

are not shown because they were found in very small quantities, when detected at all. If it were not for the 7-Up Pete station, average iron would be fairly level at all stations from Hogum downstream. The average of 0.4 mg/l at 7-Up Pete is based on only two samples of .03 and .06 mg/l. Further sampling would probably change this average, although Landers Fork may have an influence on metals levels at the station.

Zinc was highest at Pop's Place and Flesher, due to mine drainage from the Mike Horse area. Zinc was higher than iron at Pop's Place and almost the same at Flesher. Anaconda Company water quality data show that more zinc than iron came from the Mike Horse Mine. This tunnel also had the greatest flow into the Blackfoot River, and even though iron levels from other tunnels are higher than zinc levels, there is less water volume and these effluents do not always reach the river. Average copper fluctuated least of all and was highest at the Hogum station. No copper was detected at Pop's Place.

A bluish-gray color was always evident in the Blackfoot River near Pop's Place, giving the water a "hazy" appearance even though the water was clear. Willow Creek (Flesher) also had this appearance below (but not above) Sandbar Creek. Hogum Creek exhibited this appearance to a lesser degree. All these areas receive water from abandoned mines. This condition may be due to a colloidal hydroxide of a heavy metal. Hem (1970) states that "colloidal ferric hydroxide is commonly present in surface water, and small quantities may persist even in water that appears clean." This condition was not found in other streams sampled.

Biological Oxygen Demand (BOD)

Five-day BOD analyses were made at four main river stations: Hogum and 7-Up Pete above Lincoln and Dalton and Canyon below Lincoln. Results are given in Appendix F (p. A-35). Oxygen demand by decomposition of organic matter is very low at these stations. The highest BOD value recorded was 3.0 mg/l at Hogum on September 8, 1972. The next highest value was 2.5 mg/l at Hogum and 7-Up Pete. In no instance was zero BOD recorded, but in 25 percent of the analyses values were less than 1.0 mg/l. Sixty-one percent of the values were between 1.0-2.0 mg/l, and 14 percent were 2.0 mg/l or greater.

Coliform Bacteria

Total and fecal coliforms were determined at Hogum, 7-Up Pete, Dalton and Canyon stations during water year 1973 (Appendix G - p. A-38). Total coliform organisms reached 1700/100 ml once at Hogum and 1400/100 ml once at Canyon. Otherwise the greatest number found was 930/100 ml at Canyon. Out of 48 monthly analyses (12 at each station), 75 percent did not exceed 50/100 ml. The remaining analyses (excluding the three high values above) had coliform numbers between 0-440/100 ml. Nine percent were between 50-100/100 ml, 7 percent between 100-200/100 ml, and 4 percent were over 200/100 ml.

Fecal coliforms were much lower than total coliforms. Fecal coliforms reached a high of 15/100 ml at Dalton, followed by the next high of 14/100 ml at Canyon. Eighty-eight percent of the analyses had less than 10 fecal coliforms per 100 ml, and 73 percent were less than 5/100 ml.

Montana Water Quality Standards for B-D₁ waters, which include growth and propagation of salmonid fishes and associated aquatic life, require generally that the average number of total coliform organisms not exceed 1000/100 ml, and that fecal coliforms not exceed 200/100 ml.

It is unknown whether these coliform numbers are due to natural or domestic sources. There may be some seepage which reaches the river from septic tanks in the Lincoln area. However, from a coliform bacteria standpoint, water quality was very high at these four locations.

Dissolved Oxygen (DO)

Field analyses (Appendix H - pg. A-41) show minimum DO levels at 30 stations (DO not analyzed at 6 stations) were 7.0 mg/l. This minimum occurred at only 5 of the 30 stations (17 percent). Minimum DO was 8.0 mg/l or higher at 25 of 30 stations (83 percent), and 9 mg/l or higher at 14 stations (47 percent).

The highest DO level recorded was 14 mg/l at Kleinschmidt Creek, but this occurred only once. Lowest DO levels were found in Pass Creek, where the average was 8.0 mg/l (range 7-11 mg/l). Highest average level found was 12 mg/l in East Twin Creek (range 11-13). Three percent of the stations averaged 7 mg/l, 3 percent 8 mg/l, 7 percent 9 mg/l, 63 percent 10 mg/l, 20 percent 11 mg/l, and 3 percent 12 mg/l.

The USGS also made monthly field determinations of DO at the two gaging stations (Table 4).

Table 4. Monthly DO levels from field analyses by the USGS at Pop's Place and Hogum stations.

Water Year	No. of Analyses	Pop's Place			Hogum		
		Range	Ave.	Ave. Percent Satura- tion	Range	Ave.	Ave. Percent Satura- tion
1969	11	8.1 - 11.6	9.9	90			
1970	12	8.0 - 12.1	9.9	92			
1971	12				9.2 - 13.0	11.1	105
1972	11				8.2 - 13.4	10.8	101

The lower DO levels at Pop's Place may be due to: (1) the lack of much stream turbulence at the Pop's Place station; (2) passage of the river through a series of beaver ponds just above the Pop's Place station which may cause an increase in BOD from organic decomposition in the ponds; (3) the heavy metals from mine drainage creating a chemical oxygen demand (COD); or (4) steeper stream gradient above the Hogum station maintaining DO at higher levels by aeration.

Montana's Water Quality Standards classify the Blackfoot River and all tributaries as B-D1, which provides for the growth and propagation of salmonid fishes and associated aquatic life. Under this classification, dissolved oxygen "is not to be reduced below 7.0 milligrams per liter." The above data show this standard is presently being met at all sample stations on the river. However, it also appears that an increase in BOD could lower DO below the recommended level at those locations where DO readings of 7.0 mg/l were found.

Diurnal Variation of Dissolved Oxygen, Temperature and pH

Field analyses were made at 1 to 4-hour intervals for a 24-hour period at Pop's Place, Flesher, Alice Creek, Poorman Creek, Dalton and Canyon (Appendix I, p. A-48). Most of the data was obtained by the USGS at the Pop's Place gage beginning in 1968. Diurnal pH was obtained only at Pop's Place in 1968 and 1969.

The maximum diurnal variation in DO was 1.5 mg/l in August 1973 at Flesher (Table 5). Other variations ranged from a maximum of 1.4 mg/l at Dalton in June 1972 to a minimum of 0.4 mg/l at Pop's Place in August 1971. The lowest saturation value was 84 percent at Flesher in August 1973 and at Dalton in September 1971. The highest saturation value found was 108 percent at Poorman Creek, Dalton and Canyon in June 1972. The highest DO concentrations occurred when water temperatures were coolest in June 1972 and October 1968. Water temperatures associated with the DO variations reached a high of 67 F at Flesher in August 1973 and a low of 41 F at Pop's Place in October 1968. The greatest pH variation was 1.3 units, but this may be inaccurate (see footnote in Appendix I). Other pH values ranged from 7.3-8.6 with variations of 0.3-0.5 pH units.

Miscellaneous Surface Water Temperatures

Surface water temperatures taken periodically during the study are given in Appendix J, (p. A-53). The highest temperature recorded was 18.9 C (66 F) on the main Blackfoot River at the Flesher station on August 17, 1973. This was a very low water period in a dry year. Minimum temperature was 0 C (32 F) at several stations during the winter months.

Turbidity

Measurements of turbidity were made at 21 locations during the study (Appendix K, p. A-57). In nearly all cases, turbidity was related to streamflow, i.e., turbidity increased with increased streamflow. The highest turbidity recorded during the study was 220 JTU (Jackson Turbidity Units) in Landers Fork at Highway 200 during high water on June 3, 1972.

Table 5. Variation in dissolved oxygen, percent saturation, water temperature and pH over a 24-hour period at six stations, 1968-1973.

Station	Date	DO (mg/l)			Percent Saturation			Water Temp (F)			pH		
		Max	Min	Vari- ation	Max	Min	Vari- ation	Max	Min	Vari- ation	Max	Min	Vari- ation
B.F. River @ Pop's Place	9/9, 10/68	8.8	7.8	1.0	101	85	16	55	52	3	8.6	8.1	.5
	10/9, 10/68	10.4	9.8	.6	98	90	8	41	39	2	7.6	7.3	.3
	7/22, 23/69	8.4	7.9	.5	99	92	6	60	53	7	8.0	6.7*	1.3
	8/21, 22/69	8.0	7.3	.7	93	88	5	62	54.5	7.5	7.8	7.4	.4
	8/31, 9/1/71	8.1	7.7	.4	93	88	5	57	52	5			
	8/17, 18/73	8.0	7.2	.8	94	85	9	61	53	8			
B.F. River @ Flesher	8/31, 9/1/71	8.5	7.9	.6	98	88	10	58	50	8			
	8/17, 18/73	8.5	7.0	1.5	106	84	22	67	53	14			
Alice Creek	8/31, 9/1/71	8.4	7.8	.6	96	87	9	57	50	7			
	8/17, 18/73	8.8	7.6	1.2	105	86	19	64	50	14			
Poorman Cr.	9/1, 2/71	9.5	8.4	1.1	102	89	13	53	48	5			
	6/22, 23/72	10.6	9.5	1.1	108	102	6	54.5	45.5	9			
B.F. River @ Dalton	9/1, 2/71	8.9	8.0	.9	97	84	13	53	48	5			
	6/22, 23/72	10.3	8.9	1.4	108	97	11	56	47	9			
B.F. River @ Canyon	9/1, 2/71	8.9	8.4	.5	96	88	8	53	50	3			
	6/22, 23/72	9.8	8.8	1.0	108	98	10	55	49	6			

*May be in error due to improper calibration of DO meter (USGS field notation).

Nearly all the turbidity occurring in the main river below Landers Fork was due to suspended sediment from Landers Fork. The highest turbidity recorded on the main river above Landers Fork was 42 JTU at the Hogum station on May 17, 1972. In 1973, turbidity reached only 19 JTU at this station. The highest turbidity recorded at Flesher was 4.6 JTU in May 1972 and 5.2 JTU in May 1973. Highest of two values recorded at Pop's Place was 8.3 JTU.

Maximum turbidity recorded in Alice Creek was 31 JTU on May 17, 1972. A reading of 27 JTU on May 24, 1972 was due almost entirely to sediment from a mud slide washing into the stream from state land several miles upstream. The mud slide was caused by water saturation of the soil in a logged area with improperly placed roads.

Landers Fork contributes most of the suspended sediment to the upper Blackfoot River during spring runoff. Sediment originates from natural sources in the drainage. June 3, 1972.

Emphasis was placed on obtaining turbidity readings during spring runoff because this is when practically all the turbidity occurred. Sometimes the study area sustained a midwinter thaw which raised turbidity slightly. This occurred at Canyon on January 16, 1972 when a reading of 4.2 JTU was recorded. Turbidities were generally extremely low on the main river and most of its tributaries except during spring runoff. Turbidity was mainly due to suspended sediment. Higher than normal turbidity during low water periods was usually the result of man's activities in or adjacent to a stream. Rainstorms in summer and fall

did not increase streamflow greatly. Practically all winter precipitation went into storage as snow which did not run off in large amounts until spring.

Streamflow

Continuous discharge data were collected by the USGS at Pop's Place and Hogum gaging stations on the main river. Streamflow data at other locations were determined from staff gage readings and periodic flow measurements (Appendix L, p. A-63). Continuous discharge data were transformed into hydrographs and are shown in Figures 7 and 8 for both USGS stations. Low flow periods were generally from August-March, while higher flows occurred from April-July, although high flows began in mid-March in Water Year 1972 at the Hogum station.

Both of the gaging stations were upstream from the confluence of Landers Fork with the main Blackfoot River and they did not show the significance of Landers Fork in the timing and amount of runoff in the drainage. Landers Fork is a relatively long drainage (about 22 miles). Runoff waters originate in high mountain country along the continental divide. Consequently, the peak runoff in the Blackfoot River below Landers Fork came later in the season than did the peak at the gaging sites, which receive runoff from lower elevations and from a much smaller drainage area. Below the mouth of Landers Fork there were two peaks on the hydrograph - one for discharge of the Blackfoot above Landers Fork and another for discharge contributed by Landers Fork. This is illustrated by a hydrograph of discharge at the Canyon station which was based on periodic staff gage readings in 1972 and 1973 (Figure 9). There was a 2-week period between peaks in 1972. In 1973 the peak caused by Landers Fork was very low and somewhat indefinite, due to below-normal snowpack in Landers Fork drainage. In addition to snowpack, precipitation and temperature conditions during May and June determined the timing and peak of runoff.

Spring snowmelt raises the Blackfoot River to bank-full stage in Lincoln canyon. June 3, 1972.

Figure 7. Hydrographs of Blackfoot River at Pop's Place for water years 1969 and 1970.
From Coffin and Wilke 1971.

Figure 8. Hydrographs of Blackfoot River at Hogum for water years 1971 and 1972.

Figure 9. Hydrograph of Blackfoot River at the Canyon station, 1972 and 1973, from periodic staff gage readings.

Streamflow in the main Blackfoot River was continuous except for an intermittent stretch near Lincoln. Each fall and winter streamflow ceased in the vicinity of the Stemple road bridge in Lincoln to about 3 miles upstream. Aerial observations each year during the study revealed a dry streambed in this section. Flow recovery began below the Stemple bridge and was again normal at the mouth of Poorman Creek. Loss of streamflow was due to natural lowering of the groundwater level in the highly permeable alluvium. Coffin and Wilke (1971) discuss groundwater movement in the upper Blackfoot River.

Alice Creek also had some dry sections from late summer through winter due to fluctuating groundwater. The stream dried up beginning about 1/4 mile below the abandoned U. S. Forest Service guard station and ending about 2-1/2 miles downstream just above the mouth of Bear Creek. The exact upper and lower limits of the dry areas depended on seasonal climatic conditions. The lower end of Wildcat Gulch, a tributary in this stretch, was usually dry except during the spring runoff period. A small spring helped recharge Alice Creek in T 16N, R 7W, Sec 27 (SW $\frac{1}{4}$), near the mouth of Bear Creek.

Landers fork was dry in late summer above the mouth of Copper Creek in 1973, but not in 1974. Streamflow in 1973 below Copper Creek was primarily due to flow from Copper Creek. Groundwater again came to the surface before Landers Fork reached Highway 200, since flow at the highway was greater than either Copper Creek flow or Copper Creek + Landers Fork flow (see Appendix L - p. A-63). Landers Fork has reportedly been dry at Highway 200 in past years. This was never observed during the study, although the flow did get very low. Cadotte Creek, 7-Up Pete Creek, Bartlett Creek (Alice Creek tributary), and Blanchard Creek were observed to be dry by mid-summer each year. Elk Creek was dry during one winter sampling period. Other streams reached very low water levels, but were not observed to be dry during the study.

Water Chemistry/Discharge Relationships

Certain chemical ions were plotted against stream discharge on log/log paper to determine their relationships (Figures 10-15). Data were obtained from the USGS analyses at the Pop's Place gaging station for water years 1969 and 1970, and for the Hogum gaging station for the water years 1971 through 1973. Hydrographs indicated low flows were from August to March and high flows from April to July (see Figures 7 and 8). Data were plotted accordingly.

Dissolved solids showed a definite trend toward higher concentrations during low flow periods at both stations. The major ions, bicarbonate, calcium, magnesium and sodium as well as total hardness and specific conductance followed the same trend. Sulfate, fluoride and chloride did not show a definite trend. Potassium followed the same trend as sodium, except the points were more scattered, giving a less definite relationship.

Bicarbonate showed a good inverse relationship with flow at Hogum, but a poor relationship at Pop's Place. Conversely, sulfate at Pop's Place showed a good inverse relationship with flow, while a poor relationship was found at Hogum. The ratio of sulfate to bicarbonate at Pop's Place was higher (range .79-.96) than at Hogum (range .09-.11).

Figure 10. Relationship between discharge and dissolved solids, Blackfoot River at Pop's Place, water years 1969 and 1970.

Figure 11. Relationship between discharge and dissolved solids, Blackfoot River at Hogum, water years 1971, 1972 and 1973.

Figure 12. Relationship between discharge and bicarbonate, Blackfoot River at Pop's Place, water years 1969 and 1970.

Figure 13. Relationship between discharge and bicarbonate, Blackfoot River at Bonum, water years 1971, 1972 and 1973.

Figure 14. Relationship between discharge and sulfate, Blackfoot River at Pop's Place, water years 1969 and 1970.

Figure 15. Relationship between discharge and sulfate, Blackfoot River at Hogan, water years 1971, 1972 and 1973.

pH at Pop's Place ranged between 7.1-8.0 over the 2-year period, whereas at Hogum the range was 6.4-8.5. A plot of pH against discharge at Hogum showed slightly higher pH levels (more basic) during low flow periods than during high flow periods.

There was no relationship between dissolved iron and discharge at either station. The highest dissolved iron levels found were during the high water period. However, the next highest values were found during low water.

The highest levels of dissolved nitrogen ($\text{NO}_2 + \text{NO}_3$ as N) and ortho-phosphorous were found during the high water period, although there were also high values during low water. There was no apparent relationship between these constituents and streamflow.

U. S. Geological Survey Data

Data discussed in this section are from the two USGS gaging stations operated during the study: "Blackfoot River near Lincoln" (Pop's Place station) for water years 1969 and 1970, and "Blackfoot River below Alice Creek near Lincoln" (Hogum station) for water years 1971 through 1973. Data for these water years are published by the USGS in Water Resources Data for Montana, Part 2, Water Quality Records, for the respective water years.

Discussion of USGS data will include only continuous water temperatures, suspended sediment, and metals. Other chemical constituents have previously been discussed.

Continuous thermograph records at Pop's Place show daily water temperatures ranged from 0 C during winter months to 17 C (62.6 F) during July and August in water year 1969. The average daily maximum temperature in August was 15 C (59 F). The continuous record is incomplete for July. During water year 1970 the minimum temperature was again 0 C in winter, but reached a maximum of 18 C (64.4 F) one day each in July and August. The average daily maximum temperature in July was 15.7 C (60.3 F) and in August 15.1 C (59.2 F).

The minimum water temperature at Hogum in water year 1971 was 0 C during winter months and reached a maximum of 21 C (69.8 F) in July and 21.5 C (70.7 F) in August. The average maximum temperature in July was 17.8 C (63 F) and in August 19.2 C (66.6 F). Summer water temperatures in water year 1972 were slightly cooler than in 1971 - maximum temperature in July was 19.5 C (67.1 F) and in August 21 C (69.8 F). Average maximum temperatures in July and August were 16.6 C (61.9 F) and 17.6 C (63.7 F). A maximum temperature of 18 C (64.4 F) or above was reached each day for 14 consecutive days in July and 25 consecutive days in August 1971. In 1972 the longest period of 18 C or more was 9 days. However, during these hot spells, water temperatures cooled off 7-11 C (12.6-19.8 F) each day in both 1971 and 1972.

Instantaneous suspended sediment analyses made by the USGS show low concentrations prevail most of the year at both gaging stations. Concentrations at Pop's Place for water year 1969 ranged from 1-16 mg/l and for water year 1970 from 3-18 mg/l. The highest concentrations occurred during spring runoff. Concentrations at Hogum ranged from 1-18 mg/l in water year 1971 and from 1-249 in water year 1972. As with turbidity, the highest concentrations occurred during spring runoff. The day after the maximum concentration (249 mg/l) was recorded at Hogum, a turbidity reading of 42 JTU was recorded. Both readings were taken at a flow of 900 cfs.

Metals considered most significant are iron, zinc, lead, arsenic, copper and cadmium. From 5-28 analyses were made each year for iron, and 2-4 analyses were made for the other metals. Tables 6 and 7 show values obtained. Other metals analyses not tabulated here can be found in USGS Water Resources Data for Montana, Part 2, Water Quality Records.

Table 6. Summary of USGS analyses for metals (ug/l) at Pop's Place, water years 1969 and 1970.

Water Year		Fe	Zn	Pb	As	Cu	Cd
1969	Range	10-80	450-550	2-3	None	6-7	11-55
	Avg	28	500	2.7		6.5	39
1970	Range	29-262	None	None	None	None	None
	Avg	119					

Table 7. Summary of USGS analyses for metals (ug/l) at Hogum, water years 1970-1973.

Water Year		Fe	Zn	Pb	As	Cu	Cd
1971	Range	0-110	None	None	0-4	None	0
	Avg	26			1		0
1972	Range	0-50	10-30	0-4	0	0-14	0
	Avg	19	20	1.5	0	4.8	0
1973	Range	9-150	10-20	3-3	0-2	1-8	0
	Avg	36.5	15	3	1	4.5	0

Acid Mine Seepage Surveys

Chemical surveys were made in 1972 and 1973 by the Anaconda Company to determine the location and extent of acid mine seepage from abandoned mine tunnels in the Heddleston mining district. A partial summary of the company's survey reports is given in Table 8, with original laboratory data given in Appendix M (p. A-67). This summary includes only those laboratory results for total recoverable parameters from an unfiltered water sample. The company also analyzed filtered and acidified samples for total dissolved constituents, developed acidity titration curves and made general observations of the mine tunnels. The latter data are in Anaconda Company files.

It is apparent from data in Table 8 that high concentrations of metals enter the Blackfoot River from mine tunnels and tailings. Zinc and cadmium were highest from the Mike Horse tunnel in both years. Iron was highest from the Paymaster tunnel in 1972 (Paymaster had no flow in 1973). Iron, zinc, copper, aluminum and manganese occurred in the largest concentrations. Nickel, chromium, arsenic, lead and cadmium occurred in lesser amounts.

Mine drainage from the abandoned Mike Horse Mine contributes significant amounts of heavy metals to the upper Blackfoot River. June 15, 1971.

Table 8. Partial summary of acid mine seepage water quality data obtained by the Anaconda Company in the Heddleston District, 1972 and 1973.^{1/}

Source	Date	Lab pH	Tot. Alk	Tot. Hard.	Calc. Tot. Res.	HCO ₃	SO ₄	Ca	Mg	Cl	Fe	Zn	Cd	As	Pb	Cu	Cr	Ni	Mn	Al	Est. Flow (gpm)
Mike Horse Tunnel	8/ 4/72	6.33	50	1031	1352	61	928	150	158	2	20	52	.094	.006	.22	1.45	<.02	.10	26.6	.2	300
	7/12/73	6.8	85	581	964	61	652	96	83	0	9	26.5	0.55	.04	.05	.26	.005	.067	15.3	.24	300
Anaconda Tunnel	8/ 4/72	2.96	0	327	626	0	485	68	38	1.4	28	6.20	.028	.016	.08	4.30	<.02	.05	12.6	4.9	5
	7/12/73	2.77	0	251	582	2/	466	54	28	0	8.3	4.30	.043	<.001	.405	1.75	.003	.050	15.5	4.2	1
Tunnel No. 3	8/ 4/72	3.02	0	74	268	0	211	16.8	7.8	3.2	17.2	.18	<.010	.013	<.02	30.4	<.02	.06	.31	7.7	30
	7/12/73	2.84	0	28	296	2/	252	1.3	6.0	0	19	.26	.004	.04	.005	28.5	.009	.085	.43	10.0	2
Edith No. 2 Tunnel	8/ 4/72	3.12	0	70	253	0	189	25	2.2	2.2	13.4	.58	.012	.005	.12	5.08	<.02	.05	.78	4.7	7
	7/12/73	2.98	0	57	238	2/	188	10.4	7.6	0	10	.61	.013	<.001	.148	3.70	.002	.056	1.00	5.7	1
Paymaster Tunnel	8/ 4/72	2.62	0	174	919	0	696	48	13.3	11.2	120	3.40	.048	.021	.10	21.0	<.02	.08	4.56	15.8	5
Mine drainage from Swamp Gulch near Pop's Place	8/ 4/72	6.19	20	76	145	25	67	18.4	7.3	2.0	15.6	.16	.014	.001	.08	.23	<.02	<.02	1.6	1.4	10
	4/27/73	3.80	0	50	112	0	80	12.0	4.8	0	18	1.6	.04	<.01	.21	.60	-	-	-	-	-
Hearttrap Cr. below Tailings Pond	8/ 4/72	8.24	102	117	202	125	34	22	15.2	1.9	.1	.08	<.010	.008	<.02	<.02	<.02	<.02	.08	.2	
	7/12/73	8.08	85	88	158	104	26	11	14.7	0	.083	.065	.002	.03	.03	.013	.001	.007	-	-	
Drill Hole No. DZ74-135 (Flowing Hole)	8/ 4/72	-	0	58	46	2/	-	12	6.8	-	14.2	.42	<.010	.005	<.02	.25	<.02	.02	.79	2.0	5

^{1/} Results expressed as Total Recoverable in mg/l.

^{2/} Not given in data. However, when total alkalinity is zero, bicarbonate (HCO₃) is also zero, since alkalinity is composed almost entirely of HCO₃ and CO₃ (carbonate). CO₃ is not present below pH 8.3.

^{3/} Fish and Game sample analyzed by State Chemistry Laboratory, Helena. (Not shown in Appendix M).

Note: Viking tunnel had no flow in 1972 or 1973. Paymaster tunnel, Drill Hole No. DZ74-135, and the mine drainage from Swamp Gulch had no flow in July 1973.

A comparison between seepage from mine tunnels and water from Beartrap Creek below the tailings pond showed a great difference in water quality. Water quality in Beartrap Creek was quite high compared with that from the tunnels, and it compared favorably with water quality at stations further downstream. It is also of interest to note that cutthroat trout lived in this short stretch of Beartrap Creek, whereas no fish lived in any of the mine seepages. No fish were found in Beartrap Creek below its confluence with the Mike Horse tunnel seepage due to the poor quality of water from this tunnel.

It is apparent that a water quality problem exists in the headwaters of the Blackfoot River. This poor quality water causes a reduction in stream biota in the main river downstream from the mining areas (see Fish and Bottom Fauna sections of this report).

Metals in Stream Sediments

Stream sediment samples weighing approximately 5 pounds each were collected at 10 stations in August 1973 and analyzed for copper, nickel, lead, cadmium, zinc and iron (Table 9). A 25-pound sample was collected at the Hogum station in 1974 and analyzed more extensively than were the 1973 samples. Results of these analyses are given in Appendix N (p. A-76). Loss of organic matter on ignition at 750 C (LOI @ 750 C) was determined for the 1973 samples, but not for the 1974 sample. Only the sediment fraction passing through a Tyler standard sieve No. 100 was analyzed in 1973, whereas 10 fractions between sieve No. +16 and -325 were analyzed in the 1974 sample.

With the exception of a higher zinc value at the Hogum station (259 ug/g), the highest levels were found at stations associated with mine drainage. The Blackfoot River at Pop's Place had the highest levels of all metals except iron, followed by Flesher, the next station downstream. Poorman Creek had a slightly higher copper level than did Flesher, which may be related to past mining activities upstream from the station. The highest iron was found in Willow Creek (Flesher). Observations of Sandbar Creek, a Willow Creek tributary upstream from the sample point, showed this stream to be the probable source of iron.

It is interesting to note the high cadmium level at Pop's Place. Cadmium usually occurs in extremely low amounts in natural water, because it occurs as the sulfide CdS in a rare mineral called greenockite. Cadmium used in industry is produced entirely from zinc ores. The ratio of zinc to cadmium in these ores is about 200 to 1 (Briscoe 1949). The high zinc levels from mine drainage in the Mike Horse area indicate that the source of the high cadmium levels at Pop's Place are from zinc ores in the Mike Horse area.

The concentration of metals in sediments at stations which did not directly receive mine drainage of some kind were generally well below those levels from locations receiving wastes.

Willow Creek (Flesher) had the highest loss on ignition of organic matter. This reinforces observations made of suspended matter during water

Table 9. Levels of heavy metals in stream sediments and loss on ignition of organic matter, August 1973.

Station	Micrograms/Gram Dry Weight						LOI Percent @ 750 C
	Cu	Ni	Pb	Cd	Zn	Fe	
Pop's Place	700	50	400	282.5	4080	36,600	7.85
Flesher	80	30.5	83	47.5	1000	34,600	5.98
Willow Creek(Flesher)	73	18	39.5	1.2	152	40,600	10.70
Alice Creek	42	12	32.5	<1	35	31,700	5.10
Hogum ^{1/}	47	11.5	29.5	1.2	259	25,700	4.55
	47	21.5	29.5	1.0	259	Not reanalyzed	
Landers Fork	29	10.5	28	1	41.5	25,700	7.33
Below 7-Up Pete Cr.	25	12	24.5	1.0	77.5	29,700	5.90
Poorman Creek	88	20	70	1	73.5	24,700	9.40
Dalton	23	10	28	<1	53	26,700	4.55
Canyon	24	10	47.5	<1	52	25,700	4.80

^{1/} Same sample analyzed twice. See paragraph 5 of letter in Appendix N (pg. A-76) for explanation of variation in results.

quality sampling of this stream. Even during low water periods, a heavy yellow-brown residue was retained on the 0.45 micron filter used to filter the water samples. This residue was always more pronounced than that found at other stations sampled the same day. A portion of this residue is apparently from organic matter, which also becomes deposited in the stream sediments.

The second sediment sample collected at the Hogum station in May 1974 weighed 25 pounds, much heavier than previous samples. Analyses on the 10 different particle sizes were made to determine if metals were carried in different amounts on various sizes of sediment particles. The results showed a sharp increase in metals concentration occurred between the -200 and -230 fractions, indicating that analyzing this size sediment would give better results than analyzing only the -100 fraction in determining metals content.

Bottom Fauna

Bottom fauna collections were made at 11 stations during the study (Figure 16). The numbered station descriptions are given in Appendix O (p. A-82). Identifications were made to the lowest taxonomic group possible on samples collected in 1968 and 1969, March/April 1971, August 1971 and October 1971 (Appendix P - p. A-83). However, as was previously stated, the 1968 and 1969 samples were not enumerated by species.

Figure 16. Location of bottom fauna collection stations.

Diversity indices were computed for these collections according to the method of Wilhm and Dorris (1968) and these indices are also given in Appendix P (p. A-83). The formulae and computer program used in their computation are given in Appendix Q (p. A-114). The basic data given in Appendix P are summarized by station in Appendix R (p. A-116). Appendix S (p. A-126) is a summary of those bottom fauna collections which were not identified below their orders.

The number of genera, number of individuals, and their diversity indices, by station, are shown in Tables 10 and 11. For the 1968 and 1969 samples, the diversity indices were computed on number of individuals in an order rather than number of individuals in a genus. When numbers in a genus were used instead of numbers in an order, the diversity index had a higher value.

Table 10. Diversity indices based on orders.^{1/}

Station	Date	No. of Orders	No. of Indiv.	Dmin	D	Dmax	R
Pop's Place	7/16/68	4	75	0.30	1.37	2.0	0.37
	4/29/69	2	2	Dmin = D = Dmax			Undefined
	10/ 1/69	2	99	0.08	0.24	1.0	0.82
Flesher	7/16/68	5	179	0.20	1.93	2.32	0.18
Hogum	7/16/68	5	1653	0.03	1.79	2.32	0.23
	4/29/69	5	486	0.09	1.86	2.32	0.21
	10/ 1/69	5	574	0.07	1.49	2.32	0.37
Dalton	7/16/68	5	1502	0.03	1.26	2.32	0.46
	10/ 1/69	5	540	0.08	1.71	2.32	0.27
Canyon	7/16/68	5	589	0.07	0.90	2.32	0.63
	10/ 1/69	5	670	0.06	1.62	2.32	0.31
Rainbow Bend	7/18/68	5	209	0.17	1.80	2.32	0.24
Alice Cr. at Highway 200	8/24/71	5	427	0.09	1.98	2.32	0.15

^{1/} Approximate 4 sq. ft. sample collected with a Needham kick screen (mesh size = 16 meshes/inch).

Table 11. Diversity indices based on genera.^{1/}

Station	Date	No. of Genera	No. of Indiv.	Dmin	D	Dmax	R
Pop's Place	8/20/71	16	161	0.81	2.42	4.0	0.50
	10/19/71	12	195	0.51	2.51	3.58	0.35
Flesher	3/19/71	32	1062	0.33	3.76	5.00	0.27
	8/20/71	32	1522	0.24	3.01	5.00	0.42
Hogum	3/19/71	53	6024	0.12	4.01	5.73	0.31
	8/23/71	47	4522	0.14	3.56	5.55	0.37
Between Landers Fk. & Lincoln	8/23/71	40	1808	0.26	3.57	5.32	0.35
	10/26/71	44	2530	0.22	3.58	5.46	0.36
Dalton	3/19/71	43	1638	0.31	3.48	5.43	0.38
	8/20/71	35	1291	0.31	3.33	5.13	0.37
Canyon	3/18/71	44	2395	0.23	3.34	5.46	0.40
	8/20/71	28	922	0.33	2.24	4.81	0.57
Rainbow Bend	11/ 2/71	34	889	0.42	3.11	5.09	0.42
Alice Cr. above Nat'l. Forest boundary	8/24/71	25	1211	0.92	2.65	4.64	0.45
Alice Cr. at Highway 200	3/19/71	41	2538	0.20	4.12	5.36	0.24
	8/24/71	50	2620	0.24	4.42	5.64	0.23
Poorman Cr.	4/14/71	45	2780	0.20	4.28	5.49	0.23
	8/23/71	43	3650	0.15	3.41	5.43	0.38

^{1/} Three 1 sq. ft. samples collected with a modified Waters round square foot streambottom sampler (mesh size = 38/inch). Data from all three samples combined.

Theoretical maximum diversity (Dmax) exists if each individual belongs to a different species and theoretical minimum diversity (Dmin) exists if all individuals belong to the same species. The distribution of individuals among species lies between these extremes in most communities and diversity is intermediate. Redundancy (R) is an expression of the dominance of one or more species, and is inversely proportional to the wealth of species

(Wilhm and Dorris 1968). The numerical value for diversity (D) lies between D_{max} and D_{min} and shows the actual diversity of the aquatic community as compared to the maximum and minimum diversities which could have occurred in the given sample. When diversity is high, redundancy is low and vice versa.

Data indicate that bottom fauna are depressed at the Pop's Place station. This is due to past and present acid mine drainage from the Mike Horse mine and other mine tunnel seepages above the station. The depression is probably due to a combination of factors, including metals in stream sediments and in surface flow. Metals in sediments may cause direct mortalities of aquatic organisms living on the streambed, and/or may interfere with their reproductive cycles. The metals may also affect the eggs laid by trout in spawning gravels.

The bottom fauna population was better at the Flesher station than at Pop's Place as a result of improved water quality. However, cadmium and zinc levels in sediments were still fairly high, suggesting that bottom fauna may still be somewhat depressed. Except for Pop's Place, Alice Creek above the national forest boundary (which goes dry each year) and Canyon stations, fewer species were found at Flesher than at other stations in 1971 (see Table 11). The reason for lower numbers at Canyon on that date may be due to a slight change in the sample site between March and August, or sampling variability.

The highest number of aquatic insect species (53) for a single sampling period was found in the Blackfoot River at Hogum section of the river. September 14, 1972.

Rainbow Bend showed the greatest volume of insects per square foot, although numbers were usually lower than at other stations sampled during the period. This was because the largest form of insect, the giant stonefly *Pteronareys californica*, was regularly found at this station but was not regularly found at other stations. *P. californica* was found at Canyon and Dalton stations in 1968 and 1969 but not in 1971. One live adult *P. californica* was found at Dalton in June 1972. The tentative upper limit of distribution of this species is the Dalton station.

Statistical analysis of bottom fauna data can take several forms. However, because of variations between life cycles, diversity indices and other population data must be compared from samples taken at the same time of year. Since only 1 year's data have been analyzed taxonomically to date, it is not possible to compare samples on this basis, except to compare general longitudinal species distribution in the drainage. Should mining actually occur in the drainage, bottom fauna samples should be collected at the same times of year during, and for a period following, construction and operation of the mining complex. Statistical comparisons can then be made.

Fish

Fish Population Surveys

Stream surveys established species composition and relative abundance of game and nongame fish in the study area (Appendix T - p. A-129). Population estimates were made in nine stream sections (Appendix U - p. A-139). The locations of these sampling sites are shown in Figure 17. The Petersen mark-recapture method (as described by Vincent 1971 and 1974) was used for assessing fish populations.

Ten species of game and nongame fish were found in the study area (see Appendix T). Cutthroat and brook trout were the principal species found in the Blackfoot River and tributaries above Landers Fork, although whitefish were also present. Only one small brown trout was caught in the Blackfoot above Landers Fork (Hogum section). Brown trout were most abundant in the Blackfoot River below Lincoln. Mountain whitefish were probably next in abundance, although good number estimates were not obtained. Year-round populations of brown trout were found in the river only from about the mouth of Poorman Creek downstream.

For about 3 miles above Lincoln the main river went dry each year from late fall until early spring. Fish populations in this section were, therefore, intermittent. According to local residents, game fish occupied this section during the summer, but numbers or predominant species were unknown.

Poorman Creek contained cutthroat and brook trout, with cutthroat most numerous. One small brown trout was caught in Poorman Creek in the section near Circle City. A few brown trout were also observed in the mouth of Poorman Creek during the fall spawning season, probably entering from the main river. Poorman Creek flows through a series of beaver dams

Figure 17. Fisheries stream survey and population estimate sample sites.

Cutthroat trout are an important game fish in the upper Black-foot River. This 1-pound trout was caught in the Hogum section of the river. USGS gaging station is in the background. September 14, 1972.

just above its mouth, which may restrict movement of brown trout from the main river. Dolly Varden also inhabited Poorman Creek, but it is not known if they are migrants from the main river.

There were no fish in Beartrap Creek below its confluence with mine drainage from Mike Horse Creek, or in the main Blackfoot River below Anaconda Creek. However, there was a good cutthroat trout population in Beartrap Creek above the mine drainage (between the tailings dam and Mike Horse Creek) because of much better water quality (see Appendix M - p. A-67).

Both cutthroat and brook trout were observed in beaver ponds near Pass Creek, indicating improved water quality. The beaver ponds are in the proposed open pit area. If water quality were to be improved from the Mike Horse mine drainage, fish populations would be poor in the stretch of stream between Mike Horse Creek and Pass Creek because physical habitat is small and has been badly damaged by past mining and roadbuilding activities. Spawning would be the only potential use of this section unless habitat is improved.

Anaconda Creek contained cutthroat and brook trout. Cutthroat were most numerous. Only cutthroat were found in Shoue (Shave) Gulch. Both streams are within the Mike Horse mine area, but neither stream had obvious water quality problems.

It is apparent from water quality data collected by the Anaconda Company, and fish population data, that drainage from the Mike Horse mine is a large contributor to the degraded Blackfoot River headwaters. It also appears that high zinc levels (and possibly cadmium) are the principal cause of this water quality degradation.

The tailings pond on Beartrap Creek contained cutthroat trout which were larger than most fish found in nearby streams. The pond was not sampled, but visual observations were made of surfacing fish. Also, local fishermen verified the cutthroat species present from fish catches. Beartrap Creek above the pond also contained small cutthroat trout. Larger habitat size and better growing conditions than in small streams are factors which favor greater fish production in the pond. An attempt to reclaim polluted waters from the Mike Horse mine area should consider the possible tradeoffs involved in using the pond as part of the treatment system. Use of the pond to treat mine drainage would eliminate the trout fishery, but might improve the polluted river downstream for several miles. Mine drainage should be rehabilitated, but alternative methods which would avoid use of the tailings pond as part of the treatment system should be investigated.

Mine drainage into Willow Creek (Fletcher) from Sandbar Creek appears to have some detrimental effects on the fishery. Fish shocking in Willow Creek above and below Sandbar Creek produced cutthroat and brook trout in both areas. However, the proportion of cutthroat was higher above Sandbar Creek than below (2.2:1 vs 1.4:1). More trout per 1000 feet of stream were also found above Sandbar Creek than below (233 vs 156 per 1000 feet). This was not a population estimate, but nevertheless appeared significant. Physical habitats were comparable. The appearance of Willow Creek (Fletcher)

below Sandbar Creek was similar to that of the main river at Pop's Place which was affected by mine drainage. This is an indication that similar causal conditions existed in both places. Since trout were found in both affected streams, it appears that reduced water quality did not have a direct killing effect on the fish, even though their numbers were below that which would normally be expected. The population depression may be related to metals in the stream and streambed which reduce optimum conditions for insect and fish reproduction, both of which are necessary to sustain a wild fish population. Physical habitat was good in both streams and this was not believed to be a limiting factor. A more detailed study should be conducted on this problem.

Fish sampling and insect data showed water quality in the Blackfoot River to be greatly improved at the Flesher sampling station. However, Willow Creek (Flesher) enters less than 1/2 mile below this station. There may be further water quality reduction from Willow Creek, since the Blackfoot River fish population estimate was higher above than below Willow Creek. Number of trout per 1000 feet of stream was also a little higher above than below, and there were almost three times more cutthroat above than below Willow Creek.

Pass Creek supported a population of small cutthroat and brook trout. Cutthroat were most abundant. Physical habitat was too small to produce very many catchable-size fish.

Cadotte Creek dried up each year about a mile before it reached the Blackfoot River. However, beaver dams above this point supported a fishable population of brook trout. No other fish species were collected in this section.

Alice Creek supported a good, fishable cutthroat/brook trout population. Streamflow was intermittent for about 2-1/2 miles above Bear Creek and fish populations were lower when streamflow was present. Shocking in this stretch at several locations produced some cutthroat and brook trout during the summer. However, a return visit in the late fall showed this same stream section to be dry. A year-round cutthroat population existed above the intermittent area near the abandoned U. S. Forest Service guard station. A few brook trout also inhabited this stretch of stream. Cutthroat predominated in all sampled sections of Alice Creek except in the beaver ponds on the Fickler Ranch. Here brook trout were most numerous. Dolly Varden may migrate into Alice Creek for spawning, although only a few small fish were taken in shocking samples.

Brook trout were the most abundant game fish in Hardscrabble Creek. Only two cutthroat were found in three shocking sections. There was beaver activity in Hardscrabble Basin which created pond habitat for brook trout. Sculpins were very numerous in Hardscrabble Creek below the beaver ponds.

Copper Creek contained a population of small cutthroat, brook trout and Dolly Varden. A few catchable trout were caught during this study and a 5-pound Dolly Varden was caught while sampling in 1968 (Spence 1970). Copper Creek may be a spawning and rearing stream for Dolly Varden which migrate from the main river.

Landers Fork was a poor stream for resident trout in its lower reaches. The 1964 flood destroyed portions of the stream, leaving wide, unstable sections of channel. A few deep holes contained cutthroat, brook trout, and Dolly Varden which were not readily sampled by electrofishing. Landers Fork reportedly contains a migratory Dolly Varden population which comes from the main Blackfoot River. The extent of this run is unknown. However, a natural vertical falls about 15 feet high on the Silver King Ranch (T 15N, R 8W, Sec. 12) prevents migrating fish from reaching upper Landers Fork. As previously noted, Landers Fork went dry above Copper Creek in 1973. There was a small flow here in 1974. Thus a resident fishery in this section is precluded, and flow conditions would dictate whether Dolly Varden could migrate through the area every year.

The status of the Dolly Varden is unknown in the study area. It is believed adult fish move from the Blackfoot River into Landers Fork, Copper Creek, North Fork, Monture Creek and possibly Alice Creek to spawn in the fall. They may also spawn in the main river and other tributaries. Several adult Dolly Varden were caught in the Poorman-Dalton and Hogum sections of the main river during the study. As previously mentioned, one large Dolly was caught in Copper Creek. No large Dolly Varden were caught in Landers Fork, but local fishermen are known to fish for them in the fall below the barrier falls. A complete study should be undertaken to establish the status, distribution, migratory habits and significance of the Dolly Varden in the Blackfoot River drainage.

The migratory behavior of mountain whitefish in the upper Blackfoot River is also not precisely known. There was some indication from shocking data that whitefish were more likely to be present in certain stream sections in September than earlier in the season. Schools of whitefish exhibiting spawning behavior were observed in Alice Creek and in two locations on the main Blackfoot River in the fall of the year. Population estimates for whitefish were attempted in Alice Creek in 1971, and in the Blackfoot River, Hogum section, in 1972. However, insufficient numbers of whitefish were recaptured in these sections to make an estimate.

Reproduction

Young-of-the-year trout were found in all stream sections sampled, including the main Blackfoot River. The only artificial stocking of these waters in recent years has been with hatchery-raised catchable rainbow trout. Very few of these fish were captured from areas where they were planted, and it was concluded that all waters in the study area maintained themselves by natural reproduction. The following paragraphs describe the location and characteristics of some spawning areas in the Poorman-Dalton sections of the Blackfoot River and in Alice Creek in 1972 and 1973:

On November 13, 1972, 43 trout redds were identified in the 5600-foot Poorman-Dalton shocking section of the Blackfoot River downstream from the mouth of Poorman Creek. Both brown and brook trout were observed on some of the redds. Approximately one half the redds were in the upper one-third of the section.

Two redds near the mouth of Poorman Creek had up to eight large brown trout on them at various times. One trout was marked with a white T-tag. The tag number was not seen, so the tagging location is unknown. The fish could have been tagged in the Poorman-Dalton section, or at the Stemple road bridge 1-1/4 miles upstream, since some of the 21 brown trout tagged at the latter location are known to have moved downstream after tagging (see section on tag returns).

Standpipes (Terhune 1958) were placed in the above two redds to measure intragravel DO (dissolved oxygen). Two standpipes were placed in each redd, one near the upstream end and one near the downstream end. One redd measured 5 feet wide by 13 feet long and the other was about the same size. This measurement includes the entire area of disturbed gravel which may have contained more than one egg deposition area. Standpipes were driven about 6 inches into the gravel and allowed to stabilize for 48 hours before measurements were taken.

Surface water DO and DO in water taken from inside the standpipes were determined with a Hach Model A1-36-WR test kit. Velocity and depth of surface water were made with a Price Model AA current meter. A summary of data is given in Table 12.

Table 12. Water temperature and intragravel dissolved oxygen in Blackfoot River brown trout redds, 1972.

Standpipe No.	Water Depth (ft)	Velocity (fps) @ 0.6 depth	Surface			Intragravel		
			DO (ppm)	Temp (°F)	% Sat.	DO	Temp	% Sat.
1-1 ^{1/}	1.5	1.11	8.5	42	80	7.5	42	71
1-2	1.3	1.26	8.5	42	80	7.5	42	71
2-1	1.2	1.18	8.5	42	80	8.0	43	76
2-2	1.0	1.18	8.5	42	80	7.5	42	71

1/ 1-1 = Redd #1 upper standpipe; 1-2 = Redd #1 lower standpipe

In addition to the redds seen in the main river, three redds were seen in the mouth of Poorman Creek and were believed to be brown trout redds.

Four standpipes were placed in known brook trout redds in the mouth of Alice Creek on November 14, 1972. One standpipe was placed in each redd

and was allowed to stabilize for 25 hours before being sampled. Sampling techniques previously given were used here also. Sampling results are given in Table 13.

Table 13. Water temperatures and dissolved oxygen in Alice Creek brook trout redds, 1972.

Standpipe No.	Water Depth (ft)	Velocity (fps) @ 0.6 depth	Surface			Intragravel		
			DO (ppm)	Temp (°F)	% Sat.	DO	Temp	% Sat.
1	.55	.439	9.0	40.5	84	6.0	40 ^{1/}	56
2	.45	.818	9.0	40.5	84	6.5	39.5	61
3	.85	.375	9.0	40.5	84	6.5	39.5	61
4	1.6	.319	9.0	40.5	84	6.5	39.5	61

^{1/} A spring with slightly warmer water entered just above standpipe No. 1

Brown trout spawning sites in the Blackfoot River below Poorman Creek in which dissolved oxygen and water temperature measurements were made. Note standpipes in center of photo. November 13, 1972.

One pair of small Dolly Varden was observed in a group of brook trout about 150 yards upstream from the Alice Creek sample site. The fish showed indications of spawning activity.

The gravels in all spawning areas were loose and quite easily moved. They also contained fair amounts of silt which washed out easily when disturbed. The intragravel DO levels were, however, suitable to sustain eggs and fry during development. Water temperatures were also not limiting in this respect.

Brook trout spawning site near mouth of Alice Creek. Note standpipes at center of photo. November 14, 1972.

On November 15, 1973, 43 trout redds were again counted in the 5600-foot Poorman-Dalton section. Several pairs of brown trout were spooked off the redds. Two browns with white T-tags were observed. No brook trout were seen. In both 1972 and 1973 the distribution of redds was similar, although there were more or fewer redds in any specific area of stream.

On December 3, 1973, 36 of the above redds were measured for velocity and depth characteristics (Table 14). Measurements were taken at one point immediately upstream from any disturbed area, to obtain the approximate velocity and depth of the spawning area before digging began. There may have been more than one redd in the disturbed area, but the single measurement sufficed for all. Redds were recorded as being located in a

(1) pool, (2) pool tail, (3) riffle, (4) run, (5) riffle corner. These designations were based on depth, apparent velocity and streambed characteristics. A pool consisted of deep, slow water, usually with a smooth surface. A pool tail was where water left a pool over a riffle and usually had a "V" shape. A riffle was shallow, choppy, fast water over a rough, rocky substrate. A run was smoother than a riffle and somewhat deeper. A riffle corner comprised an area inside the main current of the riffle as it turned a corner, with some flow escaping to the inside of the curve which had less velocity than the main flow. Redds associated with pools were almost always near the pool tail, where velocity was greater and depth shallower. The streamflow at time of measurement on December 3, 1973 was 18 cfs, which was about half what it was on November 13, 1972 (37 cfs). Flow measurements were made with a Price Model AA current meter. Velocities were measured at 0.6 of total depth.

Table 14. Characteristics of redds in Blackfoot River, 1973.

Redd Location	No. of Redds	Depth (ft)		Velocity (fps)	
		Range	Mean	Range	Mean
Pool	6	.9 - 1.7	1.26	.241 - 1.11	.75
Pool Tail	15	.5 - .95	.67	.539 - 2.10	1.20
Riffle	11	.45 - .6	.53	.615 - 2.29	1.20
Run	1	1.5	-	2.00	-
Riffle corner	3	.6 - .9	.73	.875 - 1.29	1.14
All areas	36	.45 - 1.7	.75	.241 - 2.29	1.18

Nineteen of the 36 redds were confined to 3 distinct areas. The rest were single redds or in pairs at different locations. Streambed materials in spawning areas were loose and composed of silt, sand, pebbles and small rocks. Spawning sites were closely associated with escape cover. No redds were found where there was a lack of nearby cover, even though these sites looked as good for spawning as the others. One stretch of several hundred yards was unused where the stream flowed over gravels with no deep pools, undercut banks or log jams (i.e., no cover).

Tag Returns

Returns of trout marked with individually numbered tags gave some indication of the extent of movement in the system. Returns also indicated that the main river is not necessarily dependent on tributary streams for recruitment. A total of 463 trout over 9 inches long was marked in 11 stream sections (Table 15).

Table 15. Tag returns from marked trout.

Stream or Main River Section	Number Tagged	Number Returned	Percent Return
Cadotte Creek	4	0	0
Flesher Section	2	0	0
Willow Creek	3	0	0
Below Willow Cr. Section	7	0	0
Alice Creek	17	1	5.9
Hogum Section	15	2	13.3
Copper Creek	2	0	0
Stemple Section	23	2	8.6
Poorman Creek	16	3	18.8
Poorman-Dalton Section	269	13	4.8
Canyon Section	105	0	0
Total	463	21	4.5

The number of tags voluntarily returned by fishermen was 21 or 4.5 percent. There was some movement out of the sections where fish were tagged, but most fish were caught within the same section where they were tagged. Those which did leave the section moved both upstream and downstream. The farthest movement was by a brown trout which traveled about 4-1/2 river miles downstream from the Stemple bridge in Lincoln to the Dalton Mountain bridge west of Lincoln. The upper part of this section became dewatered in the fall from natural causes and could account for this movement.

Of the 21 tagged fish returned, 1 was a Dolly Varden, 13 were brown trout, 6 were cutthroat trout and 1 was a brook trout. All but two of the brown trout returned were marked in the Poorman-Dalton section of the main river. The other two were marked in the Stemple section. One cutthroat and the Dolly Varden were marked in the Hogum section of the main river. Two cutthroat were from the Poorman-Dalton section, two cutthroat and the brook trout were from Poorman Creek and one cutthroat was from Alice Creek.

Sixty-two percent of the tag returns were from brown trout tagged in the Poorman-Dalton section. The majority of the 463 trout tagged were marked in this section, because more trout over 9 inches long were captured here than in any of the other stream sections sampled.

Eleven trout were recaptured by shocking in the Poorman-Dalton section and one in Alice Creek the year after tagging. All of these fish were recaptured in the same section where they were tagged. Changes in growth over the period are shown in Table 16. All but two of the trout gained in length and all but two gained in weight. One fish lost in both length and weight, one gained nothing in length and only slightly in weight, and one gained in length while losing weight. The biggest increase in length was 2.6 inches, and in weight 0.87 pounds.

Table 16. Change in size of trout recaptured a year after tagging.

Stream or River Section	Tag No.	Sp.	Date Tagged	Approx. Age	Date Recap.	Total Length (in.) at		Total Weight (lbs.) at		Change in	
						Tagging	Recap.	Tagging	Recap.	Length	Weight
Poorman-											
Dalton	2-1029	LL	8/26/71	4	8/15/72	14.6	14.5	1.15	.98	-0.1	- .17
"	2-1076	LL	8/26/71	4	9/ 1/72	17.5	17.5	1.71	1.73	0	+ .02
"	2-1091	LL	9/ 2/71	4	8/15/72	12.9	14.2	.78	1.06	+1.3	+ .28
"	2-1096	LL	9/ 2/71	4	8/15/72	13.3	14.3	.86	1.13	+1.0	+ .27
"	2-1101	LL	9/ 2/71	3	8/15/72*	11.8	14.0	.56	.94	+2.2	+ .38
"	2-1105	Eb	9/ 2/71	2 or 3	8/22/72	11.2	11.5	.56	.51	+0.3	- .05
"	2-1119	LL	9/ 2/71	4	8/15/72	14.8	15.3	1.26	1.42	+0.5	+ .16
"	2-1120	LL	9/ 2/71	4	8/15/72	15.0	17.5	1.23	2.10	+2.5	+ .87
"	2-1121	LL	9/ 2/71	4	8/22/72	15.0	15.6	1.27	1.43	+0.6	+ .16
"	2-1129	LL	9/ 2/71	4	8/22/72	14.1	15.6	1.09	1.44	+1.5	+ .35
"	2-1253	LL	8/15/72	2 or 3	9/13/73	10.0	12.6	.38	.75	+2.6	+ .37
Alice Cr.	2-1298	Eb	8/21/72	3	9/ 7/73	9.6	11.0	.40	.55	+1.4	+ .15

*This fish was recaptured again on 9/1/72 in the same section.

Tagging data indicate that resident brown trout in the main river near Lincoln do not undertake long migrations to spawn in tributaries. There probably are short movements within a stream section to find suitable spawning sites. If tributaries are locally accessible, some movement into them may occur. For example if Poorman Creek had been more accessible, brown trout may have used it more extensively for spawning. Spawning may also occur in the Keep Cool and Beaver Creek system. Dolly Varden and mountain whitefish may undertake longer spawning migrations, but adequate supporting data are not currently available. Observations on spring spawning cutthroat trout were limited because of the difficulty of sampling during high water conditions.

General observations of fisherman use in the Lincoln area suggest that fishing pressure is generally low on the main river and tributaries. The low number of tag returns tends to support these observations. Pressure is concentrated at the better access sites, particularly on weekends, but is greatly reduced on less accessible portions of the river.

Metals in Fish Tissues

Twenty-five fish samples from five different stream locations were collected in 1972 and analyzed by EPA for cadmium, copper, lead, mercury, nickel, zinc and arsenic. Methods and results are given in Appendix V (p. A-143). Iron was not analyzed due to a laboratory error.

Concentrations of all metals except arsenic were uniformly lowest in fish from the Blackfoot River, Poorman-Dalton section. Arsenic was no different than in samples from the other locations. However, some of the lowest cadmium, lead, mercury, and zinc levels were found in fish from the Hogum section of the main river. The highest levels of cadmium, lead, mercury and nickel were found in fish from Poorman Creek. Copper and zinc levels were highest in both Poorman Creek and Alice Creek. The metal found in the greatest quantity in all samples was zinc. Van Meter (1973) found zinc in liver tissues to be higher than other metals analyzed and suggested the higher concentrations are related to a greater need in metabolic functions. There was no apparent trend in arsenic levels at any location. The single highest level of arsenic was found in a cutthroat from Poorman Creek.

Background data on levels of metals in fish tissues from Montana streams which are unaffected by man's activities are insufficient to make valid comparisons with the data in Appendix V (p. A-143). More such data will be needed from such state waters, and it would be valuable to make additional collections in the upper Blackfoot drainage to increase the sample size for later comparison if a mining development occurs in the Mike Horse area.

Wildlife

General

Wildlife in the study area included elk, mule deer, white-tailed deer, moose, black and grizzly bears, mountain lions, bobcats, beavers and three

species of mountain grouse (Franklin's, blue and ruffed). There were unconfirmed reports of bighorn sheep along the continental divide. Bald and golden eagles and ospreys were also observed. Several species of waterfowl were found on streams and beaver ponds, and in intermittent marshes in the spring.

Both white-tailed and mule deer wintered in the principal study area. The main wintering area for whitetails was along the Blackfoot River in the 12-mile long canyon below Lincoln. Mule deer wintered at higher elevations in the canyon. Some white-tailed deer also wintered along lower Poorman Creek, and in scattered lowland areas in the Lincoln valley.

Small bands of elk regularly wintered at several locations in the Lincoln area (Figure 18). The same elk may have moved from one range to another, depending on the time of year and severity of the winter.

There were few moose observed in the study area. Two moose were observed wintering in Indian Meadows in February 1974, and one was seen on Highway 200 in Lincoln Canyon in June 1974. Only occasional reports of moose were received from adjacent areas.

Three grizzly bears were observed by others in the Spencer Bar calving area during the summer of 1972.

Beaver were prevalent in the upper Blackfoot River. They occupied most swampy areas created by geologic dams (see Coffin and Wilke 1971). They also built their dams across flowing streams. One of the areas beaver inhabited would be covered by the proposed clarification pond, another by the proposed open pit mine, and a third area would be covered by the proposed tailings pond in Hardscrabble Basin.

Geologic dams aid beaver in creating ponds along the upper Blackfoot River. July 23, 1973.

Figure 18. Principal elk winter ranges in the Blackfoot River valley above Lincoln.

Breeding and nesting waterfowl, Wilson's snipe and nongame birds occupied the proposed tailings pond site in the spring. Snowmelt created a temporary wetland in part of this grassland flat in Hardscrabble Basin. The extent of the wetland depended on the amount of spring runoff in the basin. After the wetland dried up later in the season it was grazed by domestic sheep.

Elk Distribution

Seasonal observations of elk were made by aircraft from 1971 through 1974 (Appendix W - p. A-146). These observations on winter and spring ranges were plotted to the nearest quarter section and are shown in Figure 19. Most spring flying effort was spent observing locations of elk in the Alice Creek-lower Landers Fork area. A portion of the calving area near Spencer Bar may be used for a tailings pond in the mining operation (Figure 20). Areas adjacent to Spencer Bar itself would also be influenced by the tailings pond. Newborn calves were observed for 3 years throughout this area in June and July (Figure 21). Most calving occurred on private lands (Figure 22). However, adjacent state and national forest lands were also utilized to some extent during the calving period. Delineation of primary spring ranges and calving areas, based on aerial observations and radio telemetry information, is shown in Figure 23. Surrounding areas were probably also used for calving, but the largest concentrations of elk were found in the areas shown in Figure 23.

Aerial observations made during the winter as well as in the spring months showed that only a few elk spent the entire winter in the Spencer Bar area. These elk utilized the Alice Mountains as winter range (see Figure 18). Greater numbers of elk were found in this wintering area in the fall and early winter, but observations indicated this was just a temporary early wintering area which elk used prior to migrating to winter range east of the continental divide.

Spencer Bar and vicinity was well covered by snow all winter until about mid-April when the snow began to recede. However, elk did not begin to arrive on the calving area until the last of April or first week in May. Vegetation had just begun to green up at this time. Elk continued to arrive during May and the maximum numbers occurred from mid-May to the last week in May (Table 17). By this time vegetation had usually greened up well over the entire area. "Green up" time depended on snow conditions and seasonal weather patterns.

Elk utilized the calving areas primarily during May and June. By July 1 it was difficult to find many elk, even from the air.

Fewer elk were observed on the Spencer Bar area in the spring of 1973 as compared to the other years. This may be related to the mild winter of 1972-73. Snowpack was very light, particularly east of the continental divide on critical winter range (Figure 24). Fewer elk were observed on this winter range in 1972-73 than in the previous two winters (Table 18).

Feed on Spencer Bar may provide a needed "boost" to elk who winter on an area where sufficient feed is difficult to obtain during hard winters. Since more winter range was available in 1972-73 due to the mild winter, the elk may not have required this boost quite as much because winter feed was more available.

Figure 20. Location of proposed tailings pond with respect to the spring range and calving area locally known as Spencer Bar.

R8W

R7W

T 15 N

Figure 21. Aerial sightings of newborn elk calves in June and July 1971-1974. Location and number of calves observed.

R8W

R7W

T 15 N

===== Silver King Ranch	===== The Anaconda Company
===== State of Montana	===== Sieben Ranch Company
===== Champion International	===== Sieben Livestock Company
===== Elk Park Development Co.	===== Other private land
===== Fickler Ranch	

Figure 22. Land ownership of spring range and calving areas.

Figure 23. Primary spring ranges and calving areas in the Alice Creek-lower Landers Fork vicinity. Shaded area is Spencer Bar.

Table 17. Highest spring counts of elk observed on Spencer Bar and vicinity, 1971-74.^{1/}

Spring of	Date	Number of Flights ^{2/}	Highest Elk Count
1971	May 24	11	93
1972	May 24	5	92
1973	May 31	5	35
1974	May 14	3	104

^{1/} Includes only the area shown on maps in Figures 20-22.

^{2/} Total of both winter and spring flights.

Elk begin to utilize Spencer Bar about May 1, the time of "green up."

Table 18. Highest counts of elk observed on critical winter range, 1971-74.

Winter of	Number of Flights	Highest Elk Count	Date
1970-71	5	195	February 9, 1971
1971-72	2	200	April 19, 1972
1972-73	4	142	April 19, 1973
1973-74	11/	146	April 30, 1974

1/ Flight made with Cessna 180 instead of Supercub.

Elk and deer winter range east of continental divide.
October 29, 1974.

Elk Movement by Radio Telemetry

On April 23, 1973 two mature cow elk were trapped and banded with radio transmitter collars. These elk will be referred to as CH-6 and CH-9, which correspond to the receiver channels on which their signals were received. On April 26, 1973 a third elk, CH-2, a yearling cow was banded after being captured from a helicopter using a "Cap-Chur-Gun" and SUCOSTRIN (Succinyl choline chloride). All three elk were banded on critical winter range east of the continental divide (see Figure 24).

Radio transmitters and the receiver used in tracking were obtained from AVM Instrument Company, 810 Dennison Drive, Champaign, Illinois 61820. The receiver used was the Model LA-12 powered by eight 1.5-volt penlight batteries. The receiver also had a connection for an outside power source. Transmitters were Model SB-2 powered by four ZM12T2 mercury batteries. Activated transmitters were first sealed in beeswax, then further sealed in an acrylic compound poured into a rubber mold containing the transmitter and batteries. The completed transmitter package was attached by small brass bolts (previously sealed into the acrylic) to a 6-inch wide collar made of canvas webbing. Brightly colored nylon material made by SAFLAG (Safety Flag Company of America) was stitched to the webbing into individual color designs which could be recognized from aircraft. The designs consisted of a solid color background with a 1-1/2-inch stripe of contrasting color stitched around the center of the band. The radio antenna was placed beneath this center stripe and tied in place with nylon cord. Pertinent banding data and characteristics of the radio transmitters are shown in Tables 19 and 20.

Radio transmitter collars like these were placed on three cow elk during the study. April 1973.

Table 19. Radio transmitter data.

Elk No.	Channel	Frequency (KHz)	Beeps per Minute
CH-2	2	150.831	121
CH-6	6	150.942	74
CH-9	9	151.005	119

Table 20. Elk banding data.

Elk No.	Sex	Age	Banding Date	Ear Tag Nos.		Total weight of Collar (lbs)
				Right	Left	
CH-2	F	Yearling	4/26/73	5045A	5044A	2.10
CH-6	F	Mature	4/23/73	R0928	R0927	2.20
CH-9	F	Mature	4/23/73	R0931	R0929	2.08

Radio tracking was conducted both from the ground and from aircraft. The antenna used for ground tracking was a hand-held Model 23, 3-element yagi antenna modified to the specific transmitter frequency range. The antenna was obtained from Hy-gain Electronics Corporation, 8601 N.E. Highway 6, Lincoln, Nebraska 68505.

An aircraft antenna system adapted from Craighead et al. (1971), was mounted on a Cessna 180 aircraft having a metal skin. It is a two-element driven array consisting of two 1/4-wavelength grounded whip antennas spaced 1/4-wavelength apart and attached to the underside of the fuselage. A 1/4-wavelength delay line in series with one of the antennas causes a 90-degree phase shift in one signal before it is combined with the signal from the other antenna. The delay line is a short length of coaxial cable enclosed in a small switch box. A toggle switch connects the delay line to either the right or left antenna. The system provides a stronger signal through the antenna on the side of the aircraft on which the radioed animal is located. By switching from side to side the aircraft can "home in" on the animal by a series of turns.

Flight dates, whether radioed elk were observed or not, and the number of other elk present near the radioed elk are tabulated in Appendix X (p. A-149). Maps of radioed elk movements are shown in Appendix Y (p. A-151).

By May 3, CH-6 and CH-2 were moving off the winter range. Both animals crossed the continental divide near Lewis and Clark Pass and reached spring range between lower Alice Creek and Landers Fork within 24 hours. This was the time between the aerial sighting on May 3 and the ground location on May 4. CH-9 remained east of the divide a few days longer, but arrived on the Spencer Bar area by May 7.

CH-6 eventually crossed Landers Fork and spent about 67 days in and near a burned area between Landers Fork and Copper Creek. She had a calf which was observed twice. CH-6 was always with or close by several other cows, and later in the season, cows with calves. By July 14, CH-6 had left the spring range and calving area and traveled to summer range on Red Mountain and North Fork Copper Creek in the Scapegoat Wilderness area. By November 11 she had retraced her steps to former spring range between Copper Creek and Landers Fork and by January 3 had recrossed the continental divide onto winter range. A winter range observation indicated she still had her calf by her side.

The movements of CH-2 were quite erratic during the spring of 1973. After reaching lower Landers Fork, she wandered throughout the area with other elk and then recrossed the continental divide into the East Fork Falls Creek (Dearborn River) drainage. She spent about 65 days in this area traveling widely back and forth. Finally she crossed into the West Fork and Middle Fork Falls Creek drainages and spent the remainder of the summer near the continental divide below Caribou Peak. By November 11 CH-2 had crossed the continental divide again to the Silver King Mountain area. She was still west of the divide on December 5 and by January 3, 1974 she was again on winter range east of the divide. CH-2 and CH-6, although widely separated during late spring and summer, were observed close together on their winter range.

During much of her travels, CH-2 was quite elusive and evaded visual observation. She spent much of her time in heavy timber. When she was seen, she was usually by herself, or at least no other animals were seen nearby.

CH-9 spent 30 days on and near Spencer Bar in company with several other cow elk and did not move as widely as did CH-2 and CH-6. However, when she finally did move off the spring calving area, she went in a hurry. On June 5 she was still on the calving area, but on June 6 she was located at a point up Landers Fork near Lone Mountain, having traveled about 8 airline miles overnight. The flight on June 6 was the last observation made on CH-9 due to a radio malfunction. She was searched for several times, but not found. Finally on October 17 or 18, 1973 she was killed by a hunter near Meadow Lake in T 16N, R 9W, Sec. 18 (center of south half). A bull calf, believed by the hunter's guide to belong to the dead cow was also killed at the same time and place. However, on July 11, 1973 a backpacker observed several cow elk, one wearing a "red" neckband, on the southwest side of Bugle Mountain in the Scapegoat Wilderness, west of, and across the drainage from where CH-9 was killed. CH-9 did have a bright red stripe on her blue collar which showed up readily. Previous observations of this collar showed that the

red color showed up better than the blue color, and, depending on light conditions, frequently made the entire collar appear red. Since no other banded elk were known to be in this area, it is assumed that the hiker observed CH-9, and that CH-9 traveled at least as far into the Scapegoat Wilderness as Bugle Mountain.

CH-6 survived the winter of 1973-74, moving to several locations on the winter range during the period. She ranged west toward the continental divide for a short time, then returned to the main range. Between April 30 and May 4, 1974 she again crossed the continental divide on almost the same date as in 1973 and arrived on spring range. She was first found on May 4 in the same location where she spent most of her time in the spring of 1973 on the ridge between Landers Fork and Copper Creek.

On May 7 and 14 she was found about 2 miles east of there on the ridge between Landers Fork and Hardscrabble Creek in the McDonald Meadows area. She was never observed there in 1973, although she did move east of Landers Fork at least once, on June 6 of that year. She was in company with several other cows and yearlings on spring range in 1974, and observations indicated that one of the yearlings was hers. CH-6 did not have a new calf with her in 1974, as far as could be determined.

The movements of CH-6 in 1974 were quite different than in 1973. She did not move into the high country summer range as early in 1974 as she did in 1973. She stayed on the spring range until sometime between August 15 and September 3. While on the spring range she was seen with other cows and calves and also by herself. On September 3 and 14 she was in the Indian Meadows area. She was broadly located in Red Creek near Red Mountain on October 4, but the weather was too bad to determine a more exact location. Two subsequent flights on October 29 and November 14 failed to locate her signal. Finally on December 12 she was again located between Landers Fork and Copper Creek. She could not be located on January 15 and January 31, 1975, and not until February 26, 1975 was she again located, near Lewis and Clark Pass. She was still in this area on March 28, 1975, the last aerial sighting made on her prior to completion of this report. Her radio signal was practically inaudible on the last flight and it is believed the usefulness of her transmitter was about gone. However, a report of a cow elk wearing a "white" neckband was received by the game warden in Lincoln. The observation was made by a hiker in August 1975 on or near Bugle Mountain in the Scapegoat Wilderness. The neckband on CH-6 was blue with a white stripe which gave a "whitish" or "bluish-white" appearance from a distance. The hiker could have observed CH-6. No further observations of CH-6 have been made as of the date of this report.

CH-2 also made it through the 1973-74 winter. Although she did not travel as widely as CH-6, she did move to several different locations during the winter period. In the spring of 1974, in contrast to 1973, she did not cross to the west side of the continental divide, but went directly into East Fork Falls Creek, where she spent considerable time in 1973 prior to moving to higher summer range. She was observed in the East Fork on both May 7 and May 14, 1974. In 1973 she was west of the divide in the Landers Fork area between May 4 and May 11, but by May 14 had recrossed the divide and was on her way to the East Fork area.

Other movements of CH-2 in 1974 were also different than in 1973. After moving into the East Fork of Falls Creek she made her way into the West Fork and Middle Fork Falls Creek where she had spent considerable time in 1973. However, she suddenly left this area. On July 15 she was in this area but on July 19 she was observed all alone on the prairie about 4 miles east of Bean Lake, having traveled a distance of 13 air miles in 4 days. A week later she was back in the mountains with other elk and spent the rest of the summer in the Table Mountain-Falls Creek Ridge areas. Radio contact was lost between May 14 and June 27 for some unknown reason, so her whereabouts during this period are unknown. On October 4, 1974 she was still on the east slope of the continental divide, but by October 29 she had moved into the backcountry near Caribou Peak - the same area she occupied during the summer of 1973. She was still in this area on December 12. By January 15 she had moved into the Table Mountain area. She finally spent the rest of the winter of 1974-75 with other elk in the Cuniff Creek area near the Dearborn River. She did not return to the winter range where she was banded like she did in the winter of 1973-74. The last time she was observed prior to report completion was on August 24, 1975 (observations made since March 28, 1975 are not plotted on the maps in Appendix Y). Her radio was still operating, but at a significantly reduced level.

Both CH-2 and CH-6 eventually occupied the same summer range areas in 1974 as they used in 1973. The primary difference was in the time of occupancy. Both animals had similar movement behavior in 1974. They both stayed "outside" of their 1973 summer areas until late in the fall when both were observed in the areas they had occupied in 1973. Again, the exact location of CH-6 on October 4 was not determined, even though it was believed she was in the Red Mountain area. This behavioral difference may have been due to differences in climatic conditions in 1973 and 1974. The summer of 1973 was very dry, whereas 1974 was wetter. It may be that the elk did not have to move to higher summer range in 1974 because of the wetter summer conditions. However, the fall of 1974 was long and dry and may have forced the elk to higher range at a later time than was necessary in 1973.

Radio telemetry data and aerial observations of elk established that Spencer Bar, Hardscrabble Basin and adjacent lands in lower Alice Creek and Landers Fork are important spring range and elk calving areas. Since only 3 elk from a wintering herd of over 100 animals were fitted with radios, and since 2 of the 3 animals migrated to and spent time on or near the Spencer Bar area, it is evident that this range is an important link in the life cycle of this herd. Elk utilizing this range are the same animals hunted in the fall in, and adjacent to, the Scapegoat Wilderness. Any decline in available spring range due to mining activities may have an adverse impact on animals in this herd.

Loss of spring range might be offset by an improvement in feed on the winter range east of the continental divide (see Figure 24). Cattle presently graze these mostly private lands heavily all summer, reducing available feed for wintering elk. Acquisition and/or improvement of critical elk winter range could provide for a healthier herd and a subsequent increase in calf production.

Cessna 180 aircraft used in tracking radioed elk. May 14, 1973.

The potential increase in number of people in the Lincoln valley because of mining development could have an impact on elk utilizing the spring range. Local residents already drive into the Spencer Bar area to view the elk during the calving season. Local residents state, from their observations over many years, that the elk move farther away from roads as this public activity increases. Pickup-type campers were observed during the study to have stayed overnight in the middle of Spencer Bar during the spring period. The influx of several hundred people into the valley could increase public use of the calving area and further reduce the useful amount of area available for calving purposes. In fact, increased public use may be a significant factor as to whether the calving ground could continue to provide one of the basic needs of elk - security from harassment.

The calving areas shown in Figure 23 apparently supply basic needs for elk over and above those which can be supplied by other nearby areas. Otherwise concentrations of elk would have been found in these other areas. Thus there appears to be no alternative area(s) which can furnish these required spring needs. Any loss of the calving areas because of mining development means either greater concentrations of elk on remaining areas or a reduction in animals using the range.

Vegetated openings and adjacent timber cover provide food and security for elk on Spencer Bar. May 21 & 26, 1971.

Based on preliminary plans of the Anaconda Company, the tailings pond will cover a portion of the calving ground and eliminate elk use there. Should the proposed pond become a reality, elk use of the remaining calving area might be enhanced by controlling public or company access to it during the calving period. Should a tailings pond not be built, another consideration for improvement of the calving area is the tailings pond site itself. According to local residents and prior to the advent of increased vehicle use of the Alice Creek road, elk regularly utilized this grassland flat and adjacent hills east of the pond site. Very few elk presently use this area, possibly due to easy public access coupled with a lack of suitable escape cover. Thus there is potentially available a portion of spring range which could be used by elk if access to it were restricted during the spring. May and June are the critical months when elk occupy the area. If motorized vehicles could be controlled in Hard-scrabble Basin during this period, increased elk use might be realized. Suitable agreements with private landowners would have to be made for this to be accomplished.

Easy public access to Spencer Bar and vicinity may influence successful use of this elk calving area. June 3, 1971.

Domestic sheep are grazed by the Sieben Ranch Company and/or Sieben Livestock Company on their own and other private lands in the calving area (see Figure 22). Sheep usually arrive on the area about July 1, and are removed in September. By the time the sheep arrive, most elk have left the calving area and gone to higher summer range. However, some elk spent the summer in the vicinity of the calving area. Other private lands in this area are grazed by cattle during the summer.

Suitable feed and other necessary habitat conditions on Spencer Bar and vicinity are apparently the reason the elk move there from east of the divide in the spring. Since there is more moisture west of the divide than on the east side, and competition from cattle is less, grasses and forbs are able to provide a needed "boost" for elk which usually winter on deficient feed. It is, therefore, important to maintain the productivity of the area for this purpose.

No range analysis was made in the calving area. In the spring, elk have first use of the newly growing grasses and forbs prior to sheep and cattle use. However, the extent of food competition, if any, between the elk, cattle and sheep should be investigated. Observations of the area gave the impression that overgrazing was not a major problem except at sheep-holding areas and around salt licks.

SUMMARY

With a few exceptions, water quality is high in the Blackfoot River and its tributaries. The Blackfoot River at Pop's Place and Willow Creek (Fletcher) have lower water quality due to mine drainage, and Willow Creek (Fletcher) may affect water quality in the main river below their confluence. Elk Creek and Union Creek, in the lower river, have lowered water quality due to agricultural activities.

All of the waters sampled had calcium and bicarbonate as the major ions. Acid mine drainage in the Mike Horse area caused an increase in sulfates at Pop's Place which were greater than in any of the other waters sampled. However, bicarbonates were still greater than sulfates at this station. Anaconda Company water quality data showed calcium, magnesium and sulfate to be the major ions in seepage water from mine tunnels. Poor quality water from mine seepage is improved by the time it reaches Pass Creek, but there are still chemical problems below this point for several miles.

The heavy metals iron, zinc, copper, arsenic and lead occur in the waters sampled, but these metals could not be detected in a significant number of the samples chemically analyzed. When metals were detected, iron occurred most frequently, followed by zinc, copper, arsenic and lead. Sediment analyses showed high concentrations of iron even in streams which have not been affected by mining activities. Thus iron occurrence in surface waters is from both natural and manmade sources.

Biological oxygen demand is low in the waters sampled. The highest BOD recorded was 3.0 mg/l at the Hogum station on the main river.

The numbers of coliform bacteria present in surface water were generally low at the sample stations. In two cases total coliforms exceeded 1000/100 ml, but 75 percent of the samples did not exceed 50/100 ml - the maximum amount recommended for Montana A-closed drinking water. Eighty-eight (88) percent of the analyses showed less than 10 fecal coliforms per 100 ml, well below the recommended maximum limit of 200/100 ml for B-D₁ waters.

Dissolved oxygen concentrations at all stations sampled were 7.0 mg/l or higher, and were 8.0 mg/l or higher at 83 percent of the stations sampled. These levels meet present Montana Water Quality Standards for B-D₁ waters. Diurnal oxygen variations over a 24-hour period ranged from 0.4-1.5 mg/l at six stations, and 56 percent of the variations were less than 1.0 mg/l.

pH ranged from 7.3-8.6 units with variations of 0.3-0.5 pH units over the 24-hour period.

Turbidity values recorded were extremely low at most stations except during spring runoff. Natural runoff from Landers Fork contributes almost all of the sediment to the Blackfoot River above Lincoln and downstream as far as Nevada Creek. Nevada Creek and the North Fork of the Blackfoot River also add to the sediment load of the main river. Most tributaries contribute some suspended sediment during high water, but are extremely clear (often less than 1.0 JTU turbidity) at other times of the year. Poor logging practices and road building on state land in Section 16, T 15N, R 7W on Alice Creek caused massive slumping of water-saturated soil into the creek, resulting in turbidity increases at the Highway 200 sample station. Corrective measures by the state of Montana should be undertaken on this problem. The turbidity of the main river at Flesher reached a maximum of 5.2 JTU in 1973 and was only 4.6 JTU in 1972. An increase in turbidity at this station will be a good indicator of man-caused sediment problems above the station.

Continuous temperature records kept by the U. S. Geological Survey at two gaging stations showed that surface water temperatures reached a maximum of 64.4 F in July and August at Pop's Place, and a maximum of 70.7 F at Hogum in August. Maximum summer temperatures may prevail for 2-3 weeks at a time, but the daily maximum temperature is cooled by 13-20 F during the night.

Hydrographs of streamflow in the main river above Landers Fork showed maximum flows occurred generally between April and July and minimum flows between August and March.

Landers Fork contributes most of the total discharge to the upper river and flows usually peak in the first week of June. The Blackfoot above Landers Fork may peak 2-3 weeks earlier, depending on air temperature and precipitation.

Bottom fauna populations were sampled at 11 stations. Populations are reduced at Pop's Place on the main river due to mine drainage upstream. Although the number of species and number of individuals was substantially reduced at Pop's Place, they had both greatly increased at the Flesher station 5 miles downstream, suggesting substantial recovery. The highest number of species found during a single sampling period was 53 at the Hogum station on the Blackfoot River. The next highest was 50 species in Alice Creek at highway 200. Samples from other stations contained from 25-45 species each. The highest number of species found at Pop's Place and Flesher was 16 and 32, respectively. The effect of Willow Creek (Flesher) on bottom fauna in the Blackfoot River is not known.

Trout populations were found in all streams sampled in the study area, except the Blackfoot River between Beartrap Creek and Pass Creek in the Mike Horse area. Catchable-sized trout are limited by physical habitat in the smaller streams, but nevertheless provided a fishery. Larger trout were found in the main river, especially below Lincoln. Indications were that natural reproduction occurred in all streams containing fish, and that resident cutthroat, brook and brown trout did not migrate long distances to spawn. Dolly Varden and mountain whitefish may move longer distances to spawn, but sufficient data to determine this were not obtained. The presence of trout in nearly all waters further indicates high water quality.

It is evident from electrofishing and bottom fauna data that fish and bottom fauna populations are depressed in the Blackfoot River near Pop's Place. Anaconda Company data established that high levels of metals, particularly zinc from the Mike Horse tunnel, flowed into the Blackfoot River from abandoned mines in the Mike Horse area.

However, the quality of water in the Blackfoot improves by the time it reaches the mouth of Pass Creek. Both cutthroat and brook trout occupied these waters. This water quality improvement is probably due to dilution and precipitation of toxic metals by surface and ground water inflow at the mouth of Pass Creek. However, chemical conditions were still not good enough below Pass Creek near Pop's Place to sustain a fishery consistent with the good physical habitat present in the stream. The exact downstream extent of the depressed biota is unknown, but it was greatly improved at the Flesher station 5 miles downstream.

Stream sediment analyses suggest that high concentrations of metals, particularly cadmium and zinc, in the stream substrate may have an indirect toxic effect on the stream biota. Immature insect forms living on the substrate may accumulate these metals in their systems from direct contact or through ingestion of other food organisms which live on the metal-laden substrate. The result could be a direct mortality of the more sensitive forms, or egg mortality during reproduction. Since trout also lay eggs in stream substrates during spawning, certain metals may have the same effect on them - increased mortality by absorption of the metals into deposited eggs. The result would be reduced recruitment of young fish to the stream population. Young-of-the-year trout were captured from the stream, indicating that reproduction was at least partially successful. Another possibility is that the lack of insect life reduced the fish food supply so that only the most competitive fish survived to adulthood.

Fish populations in the Blackfoot River below the mouth of Willow Creek (Flesher), when compared with those above, indicated some adverse influence of Willow Creek on the fishery. This would be due to mine drainage from Sandbar Creek which enters Willow Creek 3 miles upstream from its mouth.

Fish and wildlife species will be affected by the proposed mining development by direct loss of habitat. Increased pressure from a larger human population may also affect these populations.

Aerial and ground observations of elk distribution showed that Spencer Bar and adjacent lands are important elk calving areas. These areas are spring ranges for elk which winter on private land east of the continental divide and summer in or near the Scapegoat Wilderness area. This fact was documented by the use of radio collars placed on three cow elk in April 1973. The movements of two of these elk were monitored by aircraft until March 1975. The radio on the third elk stopped transmitting in June 1973, and the animal was killed by a hunter in October 1973 in the Scapegoat Wilderness.

A portion of the Spencer Bar elk calving area would be encroached upon by a proposed tailings pond and increased public use of the area during the spring calving period could have an adverse impact if development should occur.

Beaver habitat will be lost in the areas used for the proposed open pit mine, tailings pond and clarification pond. Reservoirs created by the tailings pond and clarification pond will alter existing fisheries. Metallic wastes from mill tailings may make the active tailings pond unsuitable for trout species which presently occupy waters in that area.

Present and potential subdivisions along the Blackfoot River near Lincoln will likely be more in demand than at present if the population of the valley increases. Such demand will pose an added threat to the natural meandering course of the river. Streambank stabilization projects will likely be requested by riverfront property owners. The cumulative result of such protection measures will be increased channel erosion and loss of fish habitat. Lincoln sits in a flood-prone area and any tampering with the streams could increase the flood hazard.

Proper planning by the mining company and local planning groups would greatly reduce the impact of development on the land, its fish and wildlife and the social structure of the Lincoln area. Fish and wildlife habitats which are destroyed or altered by mining development should be compensated for by purchase of other lands at the developer's expense. Lands in the Alice Creek area not actually needed for mining purposes should be turned over to the Montana Department of Fish and Game for management in accordance with the March 16, 1970 letter of agreement from the Anaconda Company to the Director, Montana Fish and Game Department.

LITERATURE CITED

- Alden, W. C. 1953. Physiography and glacial geology of western Montana and adjacent areas. U. S. Geol. Surv. Prof. Paper 231. 200 p.
- APHA, AWWA, WPCF. 1971. Standard methods for the examination of water and wastewater. 13th ed. Am. Publ. Health Assn., Washington, D. C. 874 p.
- Brown, E., M. W. Skougstad and M. J. Fishman. 1970. Techniques of water-resources investigations of the U. S. Geological Survey. Book 5 - laboratory analysis. Chapter A1 - Methods for collection and analysis of water samples for dissolved minerals and gases. 160 p.

- Briscoe, H. T. 1949. General chemistry for colleges. 4th ed. Houghton Mifflin Co., Boston. 773 p.
- Coffin, D. L. and K. R. Wilke. 1971. Water resources of the upper Blackfoot River valley, westcentral Montana. Report No. 1 of technical report series of the Water Resources Division, Dept. Natl. Res. and Conser., Helena, Mont. 82 p.
- Craighead, J. J., F. J. Craighead, Jr. and J. R. Varney. 1971. Development of satellite related biotelemetry equipment. Prog. Rept., NASA Res. Grant NGR 27-002-006. Mont. Coop. Wildl. Res. Unit, Univ. Montana, Missoula. 38 p. mimeo.
- Dale, A. 1971. Progress report, Blackfoot River water quality study. Mont. Fish and Game Dept. 29 p. mimeo.
- Dufor, C. N. and E. Becker. 1964. Public water supplies of the 100 largest cities in the United States, 1962. U. S. Geol. Surv. Wat. Supply Paper 1812. 364 p.
- Hem, J. D. 1970. Study and interpretation of the chemical characteristics of natural water. U. S. Geol. Surv. Wat. Supply Paper 1473. 363 p.
- Kittrell, F. W. 1969. A practical guide to water quality studies of streams. Fed. Wat. Poll. Cont. Admin., Washington, D. C. 135 p.
- McKee, J. E. and H. W. Wolf. 1963. Water quality criteria. 2nd ed. Publ. No. 3-a, Calif. Wat. Res. Contr. Bd. 548 p.
- Montana Water Resources Board. 1970. The Anaconda Company's proposal for development of the Heddleston mining district, Lewis and Clark County, Montana. Statement by Douglas G. Smith, Director, MWRB, at public briefing, Helena, March 18, 1970. 9 p. mimeo.
- Mudge, M. R., R. L. Earhart, K. C. Watts, Jr., E. T. Tuchek, W. L. Rice and D. L. Peterson. 1971. Mineral resources of the Lincoln backcountry area, Powell and Lewis and Clark Counties, Montana. U. S. Geological Survey open file report, compiled by U. S. Geol. Surv. and U. S. Bur. of Mines. 165 p.
- Needham, P. R. and R. L. Usinger. 1956. Variability in the macrofauna of a single riffle in Prosser Creek, California, as indicated by the Surber sampler. Hilgardia 24(14):383-409.
- Pardee, J. T. and F. C. Schrader. 1933. Metalliferous deposits of the greater Helena mining region, Montana. U. S. Geol. Surv. Bull. 842. U. S. Govt. Print. Off., Washington, D. C. 318 p.
- Spence, L. E. 1970. Western Montana fishery study. Fed. Aid in Fish. Restor. Proj. F-12-R-15. April 1, 1968 to March 31, 1969. 22 p. mimeo.
- _____. 1974. An inexpensive surface water dissolved oxygen sampler. Prog. Fish. Cult. 36(1): p. 26.

- Terhune, L. B. D. 1958. The mark IV groundwater standpipe for measuring seepage through salmon spawning gravel. J. Fish. Res. Bd. Can. 15: 1027-1063.
- Van Meter, W. P. 1973. Heavy metal concentrations in fish tissues. OWRR Project No. A-053-Mont entitled "Cadmium, lead, and zinc concentrations in the fish of the upper Clark Fork River." *In* Annual Report of Montana University Joint Water Resources Center, Mont. St. Univ., Bozeman. p. 21-29.
- Vincent, R. 1971. River electrofishing and fish population estimates. Prog. Fish. Cult. 33(3):163-169.
- _____. 1974. Addendum to river electrofishing and fish population estimates. Prog. Fish. Cult. 36(3): p. 182.
- Waters, T. F. and R. J. Knapp. 1961. An improved streambottom fauna sampler. Trans. Amer. Fish. Soc. 90(2):225-226.
- Weisel, G. F. and R. L. Newell. 1970. Quality and seasonal fluctuation of headwater streams in western Montana. Bull. 38, Mont. Forest and Conservation Exper. Sta., School of Forestry, Univ. of Mont., Missoula. 16 p.
- Wilhm, J. L. and T. C. Dorris. 1968. Biological parameters for water quality criteria. Bioscience 18(6):477-481.
- Williams, A. S. (editor). 1970. Lincoln area study, analyses and proposals. Center for Planning and Economic Development, Mont. St. Univ., Bozeman. 126 p.

APPENDIX

APPENDIX A

Results of Laboratory Analyses

Laboratory analyses were conducted by the State Chemistry Laboratory, Helena. Only a partial record is given herein for those stations which had more than 12 samples analyzed. These stations are: Main River at Flesher, Hogum, Dalton, Canyon and Rainbow Bend; tributaries at Alice Creek and Poorman Creek. For these stations 12 sampling dates were selected which included the maximum, minimum and approximate average total dissolved solids for the station, with the remainder selected randomly over the sample period, but with the stipulation that each of the 12 months of the year are represented.

All values are given in mg/l except turbidity (JTU). Total hardness, alkalinity, CO_3 and HCO_3 are reported as mg/l CaCO_3 . Alkalinity values are total and are calculated from HCO_3 and CO_3 values as follows (Hem, 1970, Pg 84):

- (1) HCO_3 value $\times 0.8202$ = bicarbonate alkalinity as CaCO_3
- (2) CO_3 value (if any) $\div 0.4917 \times 0.8202$ = carbonate alkalinity as CaCO_3
- (3) (1) + (2) = total alkalinity as CaCO_3

Metals analyses are reported as total recoverable.

A zero entry means the value was less than the analytical detection level (except CO_3 , which is the true value if pH was below 8.3). A blank in a column means no analysis was performed.

Prior to February, 1971 sodium and potassium values were determined by calculating the difference between the sum of the determined anions and the sum of the determined cations in milliequivalents/liter. Direct determinations of these values by atomic absorption began in February, 1971 and resulted in increased accuracy in the cation-anion balance.

Water samples were collected in flowing water beneath the water surface in midstream whenever possible. During the period when only "total recoverable" metals were determined, a 1-liter Nalgene bottle was filled by submerging it beneath the water surface. "Dissolved" metals analyses required additional quantities of water, so a 2-quart plastic jug was filled at the same time that the 1-liter bottle was filled. The 1-liter sample was used for total recoverable metals and required no treatment. Water in the 2-quart jug was filtered through a .45 micron membrane filter into two 1-liter Nalgene bottles. One bottle received no further treatment. Three milliliters of 1:1 concentrated nitric acid (HNO_3) were added to the second bottle and the bottle shaken. Labels containing pertinent collection data were attached to each sample bottle.

Samples for nitrogen and phosphorus determination were collected in glass jars and immediately chilled in ice water to 4 C. The samples were transported to the laboratory the same day and analyses were immediately begun for those N and P compounds which are unstable.

APPENDIX A continued

BLACKFOOT RIVER AT POP'S PLACE

	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na+K	Fe	As	Zn	Pb	Cu				
										Alk													
2-17-68	0	186	90	0	73	79	3	0.0	0.3	60	28	5	26	0.08			0.40		0.00				
3-17-68	0	156	110	0	49	74	3	2.2	0.50	40	20	15	5	0.24	0.00		0.44		0.00				
4-21-68		76	70	0	61	10	2	0.00	0.10	50	10	11	0	0.04	0.00		0.07	0.00	0.00				
5-25-68	0	80	60	0	43	29	1	.1	0	35	12	7	5	0	0		.22		0				
6-20-68	0	100	65	0	61	13	1.5	0.07	0.20	50	22	2	6	0.02	0.00		0.30		0.00				
7-22-68	0	142	160	0	72	37	9	0	0.04	59	13	36		0.2	0.00		.01		0.00				
9-9-68	3	146	115	0	64	45	4	0		52	29	10	0	0.1			0.2		0				

BLACKFOOT RIVER AT FLESHER PASS ROAD

A-2

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na+K	Fe	As	Zn	Pb	Cu				
										Alk													
2-17-68	0	124	70	0	100	21	3	0	.10	82	20	5	18	.08			.05		0				
3-17-68	0	130	90	0	73	43	2	3.6	.28	60	20	10	9	0	0		.06		0				
7-16-69	0	130	110	0	92	2	9	0	.10	75	20	15	12	0			.10		.38				
10-7-69		124	190	0	104	39	8	0	.02	85	24	32	0	.10			0		0				
1-20-70	0	90	90	0	100	3	5	0	.15	82	25	5	5	.04			.02		0				
4-27-70		130	140	0	130	24	1.5	0	.20	107	30	20	0	0			.04		0				
5-22-70		100	80	0	80	21	.5	.10	.10	66	12	12	6	.10			.04		0				
6-23-71		110	87	0	130	0	0	.10	.30	107	20	8.6	1.5	.02			.21		0				
9-21-71		80	85	0	79	0	0	.40	.20	65	20	8.5	1.5	0			.02		.01				
11-22-71		110	83	0	85	37	0	.01	.20	70	20	8.2	2	.09			.12	0	.02				
12-21-71		100	93	0	98	11	0	.04	.01	80	21	9.6	1.8	.02			.10	0	.02				
8-17-73		152	92	0	106	11.9	1	.05	.10	87	21	9.6	2.4	.20	0	0	0	0	0				

APPENDIX A continued

BLACKFOOT RIVER AT DALTON MTN ROAD BRIDGE

Date	Turb	TSS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk.	Ca	Mg	Na+K	Fe	As	Zn	Pb	Cu
2-17-68	0	170	135	6	195	0	2	0	.10	170	28	16	17	.09		.02		0
4-21-68		150	105	0	200	0	4	0	.06	164	20	13	30	0	0	0	0	0
5-25-68	0	114	112	0	146	8	1	.10	0	120	29	10	3	0	0	0		0
6-20-68	9	82	102	0	140	6	4	.10	.10	115	40	1	10	.01	0	0		0
10-8-69		166	340	0	207	12	7	0	0	170	52	27	0	.10		.05		0
1-20-70	0	150	150	0	190	5	3	.10	.15	156	40	10	12	.04		0		0
7-21-70		150	190	0	210	2	0	0	.10	172	40	22	0	.10		.02		.03
8-20-70		160	240	0	200	8	0	0	.20	164	44	32	0	.10		0		.03
9-22-70		180	230	0	200	2	0	0	0	164	48	27	0	0		.06		0
11-22-71		150	110	0	200	7	0	0	.20	164	37	12	2.1	.01	0	0	0	.02
12-21-71		160	140	0	200	2.5	0	0	.10	164	39	12	2.1	0		.01	0	.02
3-5-73		261	165	0	193	7.5	0	.27	.10	158	45	13	2.5	.03	0	0	0	0

A-3

BLACKFOOT RIVER AT BLACKFOOT CANYON CAMP

Date	Turb	TSS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk.	Ca	Mg	Na+K	Fe	As	Zn	Pb	Cu
2-17-68	0	180	130	0	210	0	1	0	.10	172	40	8	16	.06		0		0
5-25-68	0	110	112	0	122	10	1	.30	0	100	27	10		.60	0	0		0
6-20-68	9	130	103	0	140	14	2	.30	.10	115	38	1	16	.05	0	0		0
10-27-68		168	210	0	201	16	9	0	0	165	48	12	0	0		0		0
11-29-68		174	210	0	207	4	9	0	.16	170	44	32	0	0		0		0
1-20-70	0	160	155	0	195	5	3	0	.15	160	45	10	8	.06		0		0
7-21-70		180	140	0	200	2	0	0	.20	164	40	10	12	.02		.03		.06
8-20-70		166	150	0	190	8	0	.10	.20	156	44	17	0	.30		0		.03
9-22-70		150	210	0	200	6	0	0	.10	164	48	29	0	0		.03		0
4-27-71		110	110	0	140	10	1	.05	.10	115	32	7	1.4	.40		.30		.06
12-21-71		170	150	0	200	3.2	0	0	.10	164	42	12	2.2	0		0		.03
3-5-73		260	165	0	192	7.5	0	.40	.10	157	45	13	2	.02	0	0	0	0

APPENDIX A continued

BLACKFOOT RIVER AT HOGUM CREEK ROAD BRIDGE

Date	Turb	TSS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu					
										Alk														
2-17-68	0	130	105	0	110	0	2	0	.10	115	28	9	6	0			0		0					
4-21-68		130	90	0	116	20	3	0	0	95	24	8	13	0	0	0	0	0						
6-20-68	2	106	77	0	98	14	2.5	.30	.14	80	27	2	10	.14	0		.03		0					
5-3-69	5	96	110	0	85	8	2	0		70	24	12	0	.20	0		0		0					
7-16-69	0	108	130	0	116	9	6	0	.08	95	28	15	0	0			.02		.04					
10-7-69		128	210	0	143	19	8	0	0	117	32	32	0	0			0		0					
1-20-70	0	125	125	0	150	8	3	0	.15	123	30	10	10	.10			0		0					
3-27-70	4	142	110	0	145	7	3	0	.30	119	26	11	10	0			0		0					
8-20-70		130	180	0	150	9	0	0	.20	123	28	27	0	0			0		.03					
9-22-70		130	180	0	150	10	0	0	.10	123	32	24		0			.02		0					
11-22-71		130	108	0	140	14	0	0	.20	115	26	10	2.3	.02	0		.03	0	.02					
12-21-71		120	110	0	140	8.6	0	.13	.10	115	28	11	2.4	0			.01		.01					

A-4

BLACKFOOT RIVER BELOW 7-UP PETE CREEK

Date	Turb	TSS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
										Alk									
3-7-73		210	131	0	151	11	0	.05	.10	124	33	12	2.2	.06	0		.01	0	.01
8-20-73		210	135	2	157	0	0	.17	.10	133	36	11	2.4	.03	0	0	0	0	

APPENDIX A continued

BLACKFOOT RIVER AT RAINBOW BEND

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk.	Ca	Mg	Na+K	Fe	As	Zn	Pb	Cu
2-17-68	0	116	100	0	143	6	3	0.00	0.1	117	32	5	13	.09		0		0
5-25-68	0	110	80	0	98	7	1	.10	0	80	19	8	3	.50	0	0		0
6-20-68	8	60	84	0	110	6	3	.60	.10	90	31	1	9	.05	0	0		0
10-19-68	6	114	140	12	140	9	7	0	.38	135	28	17	7	0		0		0
7-16-69	0	118	130	0	140	3	6	0	.08	115	32	12	0	0		.03		0
11-29-69		110	180	0	171	2	8	0	.10	110	36	22	0	.12		0		0
1-20-70	0	135	125	0	160	7	3	.10	.15	131	30	10	13	.12		0		0
3-27-70	20	158	130	0	180	8	3	.60	.40	118	34	11	15	.20		0		0
1-21-70		170	110	0	180	3	.5	.60	.10	118	32	13	20	.01		0		0
8-20-70		150	190	0	170	6	0	0	.10	139	36	24	0	.02		0		.03
9-22-70		160	200	0	170	9	0	0	0	139	36	27	0	0		.06		0
12-17-70		170	200	0	180	7	0	.04	.10	118	44	22	0	.10		.08		0

A-5

PASS CREEK

	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na+K	Fe	As	Zn	Pb	Cu																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
--	------	-----	------	-----	------	-----	----	-----	---	--------------	----	----	------	----	----	----	----	----	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

SANDBAR CREEK

																		Calc.									
		Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Alk	Ca	Mg	Na+K	Fe	As	Zn	Pb	Cu								
7-16-69		27	88	150	0	55	11	7	0	0.10	15	12	29	0	2.5		0.01		0.02								

APPENDIX A continued

WILLOW CREEK (FLESHER)

Date	Turb	TSS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
										Alk									
10-20-71		70	69	0	100	4.5	0	.10	.10	82	16	6.8	2	.16			0		.02
6-20-72		66	52	0	72	9.4	0	.26	.10	65	12	5.1	1.1	.27	.002	.01	.01	.01	
8-31-72		70	74	0	86	0	0		.12	70	19	6.1	1.8		0	.01	0	0	
9-26-72		80	75	0	83	.8	0	.13	.10	68	20	6.1	1.7		.002	0	0	0	
10-31-72		88	93	0	80	3.7	.5	.13	.10	66	28	5.2	2.2	.33	0	0	0	0	
3-7-73		130	78	0	81	21	0	.25	.10	66	18	8	2.3	.20	0	.01	0	0	
5-11-73		92	49	0	57	11	0	.28	.05	47	11	5.2	6.8	.20		0	0	0	
8-17-73		135	80	0	101	2.9	1.0	.10	.10	83	19	8	2.8	.29	0	0	0	0	

ALICE CREEK

Date	Turb	TSS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
										Alk									
2-17-68	0	120	75	0	153	0	2	0	.10	125	44	4	2	0			0		0
1-2-69	0	120	140	0	140	0	5	0	0	115	24	20	0	0			0		0
6-16-69	0	126	120	0	145	0	4	0	.38	119	20	17	2	0			0		0
3-26-71		120	130	0	150	7	0	.05	.30	123	28	15	1.5	0			.03		0
4-27-71		80	90	0	100	4	0	.10	.10	82	24	7		.20			.72		.05
9-21-71		110	110	0	150	2.1	0	.20	.30	123	28	10	1.7	0					0
11-22-71		110	110	0	150	7	0	.04	.20	123	27	9.9	2	.90	0		.10	0	.03
12-21-71		100	110	0	150	0	0	.02	.10	123	27	10	1.8	0			0	0	.01
5-22-72		68	69	0	92	4	0	.36	.10	75	17	6.2	.9	.15	0	0	0	0	
7-19-72		120	100	0	150	4.1	0	.20	.10	123	24	9.6	1.1	.06	.002	.02	.01	.01	
10-31-72		110	110	0	140	0	0	.09	.10	115	30	8.8	2	0	0	0	0	0	.01
8-17-73		214	128	0	165	3.7	0	.04	.10	135	30	13	2.1	0	0	0	0	0	0

HARDCRABBLE CREEK

Date	Turb	TSS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
										Alk									
10-20-71		170	150	0	210	0	0	.10	.10	172	38	14	4.2	.03			0		.01
3-23-72		130	110	0	150	0	0	.11	.10	123	28	9.7	2.8	.21	0		0	0	0
4-20-72		120	110	0	160	8.2	0	.14	.22	131	28	9.6	2.8	.09	0		0	0	.02
5-22-72		110	85	0	130	6	0	2.00	.10	107	22	7.2	2.3	.08	0		0	0	.01
7-19-72		170	140	0	210	3.6	0	.34	.10	172	34	12	3.2	.02	.002	.01	.01	.01	
11-29-72		288	178	0	214	6.5	.5		.10	176	47	15	4.2	0	0		.01	0	.02
3-5-73		258	161	4	188	7	0	.09	.10	161	40	15	3.8	.11	0		.01	0	0

APPENDIX A continued

HORSEFLY CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
10-20-71		66	50	0	73	0	0	.10	.10	60	16	2.6	3.4	.08		0		.01

BLACK DIAMOND CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
10-20-71		70	68	0	100	0	0	.10	.20	82	19	4.7	3.5	.18		0		.22

HOGUM CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
10-7-69		132	170	0	131	22	10	0	.18	107	28	24	0	.40		0		0
1-20-70	0	120	105	0	135	8	2	0	.20	111	30	5	13	.20		0		0
5-22-70		60	50	0	80	7	0	0	.30	66	12	5	10	.20		0		0
6-16-70		70	80	0	70	2	0	0	.30	57	20	7	0	0				
8-20-70		120	160	0	120	7	0	.20	.30	98	28	22	0	.50		0		.03
9-22-70		110	150	0	120	7	0	.20	.10	98	20	24	0	.10		.07		0
10-21-70		120	150	0	110	6	0	.10	.20	90	28	20	0	.16		.06		0
12-17-70		130	160	0	130	8	0	.06	.10	107	32	20	0	.30				
1-15-71		130	190	0	130	7	0	.04	.20	107	28	29	0	.20		.03		0
2-2-71		50	70	0	50	7	0	.10	.30	66	20	5	4.1	.30		.03		0

LANDER'S FORK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
2-17-68	0	146	120	0	180	0	3	0	.10	148	36	9	12	0		0		0
5-25-68	0	130	100	0	172	6	1	.15	0	100	22	10	6	0	0	0		0
9-29-68	4	162	155	0	174	3	3	0		143	42	12	0	0		0		0
1-2-69	0	128	170	0	172	3	5	0	.28	141	36	15	0	0		0		0
3-28-69	0	130	150	0	152	12	4	0		125	36	15	0	0		0		0
6-16-69	0	116	130	0	151	0	4	0	.36	124	28	15	0	0		.16		0
8-27-69	0	132	210	0	177	1	6	0	0	145	36	39	0	0	0	0		0
11-29-69	0	124	190	0	85	5	9	0	0	70	40	22	0	.22		0		0
2-24-71		130	150	0	160	0	0	.10	.20	131	36	15	1	0		.03		0
3-5-73		218	139	0	163	5	0	.12	.10	134	36	12	1.5	0	0	0	0	0
5-10-73		150	96	2	116	3	0	.15	.02	88	28	8.8	1.1	.31	0	0	0	0

APPENDIX A continued

POORMAN CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	P	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
2-17-68	0	140	95	0	119	11	2	0	.20	122	30	5	19	0		0		0
5-25-68	0	100	96	0	121	12	1	.10	0	99	28	6	9	0	0	0		0
1-2-69	1	134	150	0	154	4	4	0	.02	126	36	15	0	0		0		0
3-26-71		120	110	0	150	16	0	.05	.30	123	40	2	1.3	0		.02		0
1-27-71		110	110	0	130	13	0	.05	.10	107	36	5	1	.20		.22		.02
8-23-71		140	130	0	160	1	0	.10	.20	131	37	9	2	0		.01		0
9-21-71		130	126	0	160	8	0	.20	.30	131	36	9	2.1	0		0		.02
12-21-71		150	130	0	160	9	0	0	.10	131	35	9.2	1.8	0		0	0	.01
6-20-72		130	110	0	170	12	0	.30	.10	139	31	7.2	.9	.03	.001	.01	.02	.01
7-19-72		140	120	0	160	12	0	.22	.10	131	31	8.1	1.3	.06	.002	.02	.01	.01
10-31-72		130	120	0	140	6	0	.13	.10	115	37	7.2	2	0	.001	0	0	0
11-29-72		209	126	0	146	16	.5		.10	120	37	8.1	1.5	0	0	.01	0	0

SPRING CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	P	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
2-17-68	0	180	150	0	220	0	3	0.0	0.1	150	40	12	17	0.08		0.00		0.00
5-25-68	0	180	148	0	201	4	1	.2	0	165	38	12	13	0	0	0		0
9-9-68	3	170	170	0	214	0	4	0		176	46	14	5	0.0		0		0
1-2-69	2	166	180	0	214	0	5	0.3	0.52	176	44	17	2	0		0.0		0.0
3-28-69	10	180	160	0	195	15	4	0.3	0.1	160	48	10	6	0.1		0		0
6-16-69	0	162	210	0	195	0	5	0	0.36	160	32	32	0	0		0		0
8-27-69	0	190	260	0	207	5	7	0	0	170	48	34	0	0	0	0		0
11-29-69	0	168	220	0	220	11	8	0	0.16	150	44	27	0	0.08		0.00		0.02

KEEP COOL CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	P	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
2-17-68	0	170	130	0	201	0	2	0.0	0.1	165	34	11	18	0.00		0.00		0.00
5-25-68	0	180	120	0	177	7	1	.1	0	115	35	8	16	.2	0	0		0
9-9-68	4	142	167	0	177	5	3	0		115	50	11	0	0.0		0		0
1-2-69	3	160	190	0	189	0	6	0	0.2	155	44	20	0	0.3		0.0		0.0
3-28-69	17	136	150	0	152	0	5	0		125	44	10	0	0		0		0
6-16-69	0	160	145	0	182	0	6	0	0.32	140	40	11	7	0		.02		0
8-27-69	0	164	260	0	189	3	9	0	0.02	155	44	36	0	0	0	0		0
11-29-69		164	190	0	201	1	8	0	0.08	165	48	17	0	0.10		0.00		0.00

APPENDIX A continued

BEAVER CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na+K	Fe	As	Zn	Pb	Cu
2-17-68	0	150	120	0	186	0	2	0.0	0.1	152	40	5	16	0.00		0.00		0.00
5-25-68	0	110	76	0	104	8	1	.1	0	85	25	2	12	0	0	0		0
9-9-68	4	158	160	0	180	2	4	0		118	48	10	0	0.0		0		0
1-2-69	2	142	140	0	168	0	5	0	0	128	44	7	4	0		0.0		0.0
3-28-69	8	136	140	0	146	7	6	0.2		120	44	7	0	0.1		0		0
6-16-69	0	132	130	0	145	1	5	0	0.38	119	32	12	0	0		0		0
8-27-69	0	170	270	0	189	1	9	0	0	155	48	37	0	0	0.006	0		0
11-29-69		170	200	0	201	2	10	0	0.08	165	52	17	0	0.12		0.01		0.04

A9

WILLOW CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na+K	Fe	As	Zn	Pb	Cu
3-17-68	5	100	55	0	88	11	3	3.6	0.38	72	14	5	12	0.12	0.00	0.00	0.00	0.00
6-20-68	15	100	73	0	98	6	2	0.50	0.14	80	27	1	9	0.12	0.00	0.00		0.00
10-19-68	4	104	100	0	116	8	7	0		95	28	7	7	0.1		0		0
2-8-68	6	114	110	0	112	10	5	0		92	24	12	0	0.1		0.0		0.0
5-3-68	20	78	70	0	49	8	3	0		40	16	7	0	0.5	0	0		0
7-16-68	17	88	70	0	73	17	8	0	0.10	60	20	5	9	0.4		0.01		0.02
10-7-69		138	180	0	122	15	8	0	0	100	28	27	0	0		0.02		0.0
1-20-70	0	90	75	0	90	4	3	0.1	0.15	74	25	5	0	0.14		0		0

SAUERKRAUT CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na+K	Fe	As	Zn	Pb	Cu
10-20-71		50	39	0	61	0	0	.20	0	50	12	2.3	3	.02		.01		.02

APPENDIX A continued

ARRASTRA CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu				
3-17-68	0	136	90	0	122	3	1	2.5	0.38	100	20	10	7	0.00	0.00	0.00	0.00	0.00				
6-20-68	0	72	51	0	79	4	3.5	0.20	0.00	65	19	1	8	0.12	0.00	0.00		0.00				
10-19-68	5	128	110	0	140	6	5	0		115	28	10	9	0.0		0		0				
2-8-69	6	102	110	0	126	4	2	0		103	28	10	0	0.0		0.0		0.03				
5-3-69	33	90	120	0	104	4	2			85	28	12	0	0	0	0		0				
7-16-69	0	88	110	0	116	1	6	0	0.06	95	20	15	0	0		0.01		0.04				
10-7-69		116	170	0	134	14	9	0	0.1	110	28	24	0	0.1		0.0		0.0				
1-20-70	0	105	95	0	130	2	3	0	0.10	107	30	5	9	0.06		0		0				

NORTH FORK BLACKFOOT RIVER

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu				
4-21-68		150	130	0	183	0	2	0.34	0.06	150	28	15	10	0.00	0.00	0.00	0.00	0.00				
6-22-68	0	136	105	0	158	2	7	0	0.02	130	38	2	17	0.5	0.00	0		0.00				
11-30-68	6	142	150	0	175	2	4	0.7		144	44	10	0	0		0		0				
5-3-69	0	106	140	0	128	5	4	0		105	32	15	0	0	0	0		0				
7-16-69	0	110	130	0	140	1	6	0	0.06	115	32	12	0	0		0.01		0.02				
10-7-69		132	220	0	177	12	8	0	0	115	32	24	0	0		0.0		0.0				
1-20-70	0	135	130	0	170	5	2	0	0.10	139	30	10	14	0		0.05		0				

KLEINSCHMIDT CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu				
3-17-68	0	160	130	0	201	5	2	6.2	0.28	165	30	13	22	0.00	0.00	0.00	0.00	0.00				
6-20-68	0	140	130	0	183	6	3	0.50	0.10	150	51	0	15	0.00	0.00	0.00	0.00	0.00				
10-19-68	4	140	150	6	189	6	4	0.2		165	28	20	12	0.0		0		0				
2-8-69	1	156	150	0	182	5	4	0		146	36	14	6	0.0		0.0		0.02				
5-3-69	4	144	170	0	152	5	4	0		125	36	20	0	0	0.001	0		0				
7-16-69	0-2	156	190	0	189	2	7	0	0.06	155	36	24	0	0		0.02		0.00				
10-7-69		152	230	0	180	13	11	0	0	148	32	37	0	0		0.0		0.0				
1-20-70	0	120	130	10	150	4	3	0.5	0.10	140	30	15	2	0.04		0		0.				

APPENDIX A continued

WARREN CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	P	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
4-21-68		140	75	0	128	0	3	0.50	0.06	105	24	4	16	0.10	0.00	0.00	0.00	0.00
6-22-68	8	122	180	0	115	3	5	0	0.00	94	33	24		0.3	0.00	0.00		0.00
11-30-68	8	132	140	0	144	3	4	0.4	0.2	118	28	17	0	0.2		0		0

MONTURE CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	P	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
3-17-68	0	90	80	0	107	6	2	6.0	0.00	88	20	8	9	0.00	0.00	0.00	0.00	0.00
6-20-68	2	46	34	0	43	0	1	0.50	0.14	35	12	1	1	0.00	0.00	0.00		0.00
10-19-68	4	90	90	0	113	7	6	0	0	93	20	10	8	0.0		0.00		0.00
2-8-69	4	124	130	0	147	3	4	0		120	32	10	4	0.1		0.0		0.0
5-3-69	13	62	90	0	67	7	3	0		55	16	12	0	0.16	0	0		0
7-16-69	0	54	70	0	73	1	6	0	0.06	60	16	7	0	0		0.01		0.00
10-7-69		108	180	0	116	11	9	0	0	95	20	32	0	0		0.0		0.0
1-20-70	0	95	95	0	120	0	2	0.10	0.10	98	25	10	0	0.04		0		0

A-11

COTTONWOOD CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	P	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
4-21-68		150	125	0	127	20	3	0.50	0.00	104	28	14	20	0.00	0.00	0.00	0.00	0.00
7-22-68	0	176	180	0	187	2	6	0	0.00	153	46	16		0.0	0.00	0		0.00
11-30-68	7	154	160	0	182	2	4	0.5		119	34	18	0	0		0		0
5-3-69	7	132	140	0	128	7	7	0		105	36	12	0	0.1	0	0		0
7-16-69	0	164	170	0	177	1	6	0	0.10	115	40	15	0	0		0.02		0.00

CLEARWATER RIVER

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	P	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
3-17-68	5	86	70	0	97	3	3	8.9	0.00	80	14	9	10	0.10	0.00	0.00	0.00	0.00
6-20-68	0	60	48	0	73	0	3	0.50	0.04	60	19	0	8	0.00	0.00	0.02		0.00
10-19-68	4	90	80	0	110	4	3	0		90	20	7	6	0.0		0		0
2-8-69	6	84	80	0	98	2	4	0		80	16	10	3	0.0		0.0		0.03
5-3-69	10	80	80	0	73	4	3	0		60	20	7	0	0	0	0		0
7-16-69	0	64	90	0	85	0	6	0	0.08	70	20	10	0	0		0.01		0.00
10-7-69		90	160	0	122	7	8	0	0	100	28	22	0	0		0.0		0.0
1-20-70	0	105	95	0	125	0	1	0.10	0.15	102	30	5	5	0.06		0		0

APPENDIX A continued

BLANCHARD CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu				
										Alk													
4-21-68		60	35	0	55	0	2	0.00	0.00	15		8	4	5	0.07	0.00	0.03	0.00	0.00				
7-22-68	Dry											Dry											
11-30-68	8	86	100	0	98	0	4	0.1		80		18	13	0	0		0		0				

ELK CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu				
										Alk													
3-17-68	65	180	115	0	174	14	2	3.2	0.28	163		38	5	22	0.62	0.00	0.00	0.00	0.00				
6-20-68	18	186	120	0	165	12	3	0.60	0.28	135		44	2	17	0.25	0.00	0.00		0.00				
10-19-68	4	168	150	6	171	13	5	0		150		40	12	10	0.0		0		0				
5-3-69	285	128	110	0	92	7	4	0		75		28	10	0	0.9	0	0		0				
7-16-69	24	136	130	0	146	7	8	0	0.18	120		32	12	4	0		0.02		0.00				
10-7-69		188	220	0	189	13	9	0	0.1	155		44	27	0	0.1		0.2		0.0				

EELMONT CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu				
										Alk													
4-21-68		110	80	0	116	0	3	0.00	0.00	95		24	5	10	0.05	0.00	0.00		0.00				
7-22-68	0	160	210	0	173	0	5	0	0.00	112		68	10		0.4	0.00	0.00		0.00				
11-30-68	7	154	190	0	175	0	4	0		114		44	20	0	0		0		0				

UNION CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc.		Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu				
										Alk													
3-17-68	44	190	105	0	155	11	6	1.2	0.10	127		24	11	20	0.40	0.00	0.00		0.00				
6-20-68	25	242	165	0	232	8	2	0.60	0.24	190		62	2	19	0.45	0.00	0.00		0.00				
10-19-68	10	242	210	6	256	11	6	0		220		48	22	1	0.16		0		0				
2-8-69	9	206	210	0	238	8	5	0		195		44	24	2	0.1		0.0		0.02				
3-28-69	200	150	120	0	122	18	6	0		100		28	12	5	0.8		0		0				
6-16-69	15	168	175	0	207	0	7	0	0.28	170		28	26	3	0.36		.06		0				
8-27-69	20	242	310	0	232	11	10	0.9	0.1	190		40	51	0	0.4	.006	0		0				
11-29-69		220	260	0	275	12	11	0	0.12	226		48	34	0	0.22	-	0.00		0.02				

APPENDIX A continued

GOLD CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
4-21-68		100	95	0	104	0	2	0.00	0.00	85	20	11		0.00	0.00	0.00	0.00	0.00
7-22-68	0	110	125	0	122	4	5	0	0.00	100	32	11		0.0	0.00	0.00		0.00
11-30-68	7	100	120	0	119	0	4	0		98	28	12	0	0.4		0		0

EAST TWIN CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
3-17-68	11	44	10	0	27	8	2	2.7	0.44	22	4	0	12	0.18	0.00	0.00	0.00	0.12
6-20-68	0	24	8	0	24	0	2	0.20	0.10	20	2	1	5	0.06	0.00	0.00		0.00
10-19-68	7	50	20	0	36	5	3	0		30	8	0	9	0.0		0.01		0

WEST TWIN CREEK

Date	Turb	TDS	Hard	CO3	HCO3	SO4	Cl	NO3	F	Calc. Alk	Ca	Mg	Na&K	Fe	As	Zn	Pb	Cu
3-17-68	3	24	10	0	21	0	2	0.23	0.34	17	4	0	5	0.08	0.00	0.00	0.00	0.00
6-20-68	0	10	7	0	6	0	2	0.00	0.04	5	3	0	0	0.05	0.00	0.00		0.00
10-19-68	7	30	40	0	34	4	2	0	0.26	28	2	8	0	0.0		0		0

APPENDIX B

Minimum and Maximum Values of Chemical Constituents (mg/l) and Where the Values Occurred

Turbidity (JTU)

Main River - Minimum <dl* at all stations. Maximum 20 at Rainbow Bend.

Tributaries - Minimum <dl at Hogum Creek. Maximum 285 at Elk Creek.

(The above values are from infrequent (usually monthly) sampling and probably do not present a true turbidity picture (particularly higher values). See section on turbidity for more complete data.)

Total Residue

Main River - Minimum 60* at Rainbow Bend. Maximum 261 at Dalton.

Tributaries - Minimum 10 at West Twin Creek. Maximum 288 at Hard-scrabble Creek.

Total Hardness

Main River - Minimum 60 at Pop's Place, Flesher, Canyon. Maximum 340 at Dalton.

Tributaries - Minimum 7 at West Twin Creek. Maximum 310 at Union Creek.

Total Alkalinity (Calculated)

Main River - Minimum 35 at Pop's Place. Maximum 189 at Dalton.

Tributaries - Minimum 5 at West Twin Creek. Maximum 226 at Gold Creek.

*<dl = less than analytical detection level

Carbonate, CO_3^-

Main River - Minimum <dl at all stations. Maximum 12 at Rainbow Bend.

Tributaries - Minimum <dl at all stations. Maximum 24 at Poorman Creek.

(Unpolluted, natural waters normally have extremely low amounts of CO_3 due to a buffering capacity which changes the CO_3 to HCO_3 (bicarbonate) below pH 8.3.)

Bicarbonate HCO_3^-

Main River - Minimum 43 at Pop's Place. Maximum 230 at Dalton.

Tributaries - Minimum 6 at West Twin Creek. Maximum 275 at Union Creek.

(The principal anion in all waters sampled was HCO_3^- .)

Sulfate, SO_4^-

Main River - Minimum <dl at Flesher, Hogum, Dalton, Canyon. Maximum 79 at Pop's Place.

(The higher values found at Pop's Place are due to acid mine drainage from sulfide ore in several old mines in the Heddleston District.)

Tributaries - Minimum <dl at 21 of 29 stations. Maximum 22 at Hogum Creek.

APPENDIX B continued

Chloride, Cl^-

Main River - Minimum <dl at all but Pop's Place (1.0). Maximum 25 at Rainbow Bend.
Tributaries - Minimum <dl at 8 of 29 stations. Maximum 11 at Pass Creek, Kleinschmidt Creek, and Union Creek.

(High chloride levels are usually one sign of a stream polluted by domestic sewage.)

Nitrate, NO_3^-

Main River - Minimum <dl at all stations except 7-up Pete (.05). Maximum 5.3 at Hogum.

Tributaries - Minimum <dl at 24 of 29 stations. Maximum 8.90 at Clearwater River.

Fluoride, F^-

Main River - Minimum <dl at all stations except 7-up Pete (.10). Maximum .50 at Pop's Place.

Tributaries - Minimum <dl at 17 of 29 stations. Maximum 2.0 at Hardscrabble Creek.

Calcium, Ca^{+2}

Main River - Minimum 10 at Pop's Place. Maximum 56 at Canyon.
Tributaries - Minimum 2 at both East Twin and West Twin Creeks.
Maximum 68 at Belmont Creek.

(Calcium was the most abundant cation found in the waters sampled.)

Magnesium, Mg^{+2}

Main River - Minimum <dl at Canyon. Maximum 39 at Hogum.
Tributaries - Minimum <dl at Pass Creek, Clearwater River, East Twin Creek and West Twin Creek. Maximum 51 at Union Creek.

(Magnesium was the second most abundant cation found in the waters sampled.)

Sodium and Potassium, $\text{Na}^{+} + \text{K}^{+}$

Main River - Minimum <dl at all stations except 7-up Pete (.03). Maximum 33 at Canyon.

Tributaries - Minimum <dl at 23 of 29 stations. Maximum 22 at Kleinschmitt and Elk Creeks.

Total Recoverable Iron, Fe

Main River - Minimum <dl at all stations except 7-Up Pete (.03). Maximum 1.5 at Canyon.

Tributaries - Minimum <dl at 20 of 29 stations. Maximum 2.5 at Sandbar Creek.

Dissolved Iron, Fe

Main River (analyzed at 3 of 6 stations only) - Minimum <dl at all stations except 7-up Pete (.03). Maximum .19 at Dalton.

Tributaries (analyzed at 7 of 29 stations only) - Minimum <dl at Alice Creek, Hardscrabble Creek, Landers Fork, and Poorman Creek.
Maximum .20 at Willow Creek (Fletcher).

Total Recoverable Arsenic, As

Main River - Minimum <dl at all stations. Maximum .008 at Rainbow Bend.

Tributaries - Minimum <dl at all stations. Maximum .006 at Beaver Creek.

APPENDIX B continued

Total Recoverable Zinc, Zn^{+}

Main River - Minimum <dl at all stations except Pop's Place (.01).
Maximum 0.94 at Flesher.

Tributaries - Minimum <dl at 27 of 29 stations. Maximum 0.72 at
Alice Creek.

Dissolved Zinc, Zn^{+}

Main River (analyzed at 4 of 7 stations only) - Minimum <dl at
all four stations. Maximum 0.17 at Flesher.

Tributaries (analyzed at 8 of 29 stations only) - Minimum <dl at
all 8 stations. Maximum 0.02 at Alice Creek and Hardscrabble Creek.

Total Recoverable Lead, Pb^{+}

Main River - Minimum <dl at all stations. Maximum 0.01 at Flesher,
Dalton and Canyon.

Tributaries (analyzed at 20 of 29 stations only) - Minimum <dl at
all 20 stations. Maximum of 0.02 at Poorman Creek.

Dissolved Lead, Pb^{+}

Main River (analyzed at 4 of 7 stations only) - Minimum <dl at
all 4 stations. Maximum 0.01 at 3 of the 4 stations.

Tributaries (analyzed at 5 of 29 stations only) - Minimum <dl at
all stations. Maximum 0.01 at 4 of 5 stations.

Total Recoverable Copper, Cu^{+}

Main River - Minimum <dl at all stations. Maximum 0.18 at Flesher.

Tributaries - Minimum <dl at 24 of 29 stations. Maximum 0.12 at
East Twin Creek.

Dissolved Copper, Cu^{+}

Main River (analyzed at 4 of 7 stations only) - Minimum <dl at all
4 stations. Maximum 0.02 at all stations except 7-up Pete (<dl)

Tributaries (analyzed at 8 of 29 stations only) - Minimum <dl at
5 of 8 stations. Maximum 0.02 at 4 of 8 stations.

Nitrogen as Nitrite NO_2-N

Main River (analyzed at 3 of 7 stations only) - Minimum <dl at all
stations. Maximum 0.002 at Dalton.

Tributaries (analyzed at 4 of 29 stations only) - Minimum <dl at all
stations. Maximum 0.003 at Alice Creek.

Nitrogen as Ammonia, NH_3-N

Main River (analyzed at 3 of 7 stations only) - Minimum and maximum <dl
at all stations.

Tributaries (analyzed at 4 of 29 stations) - Minimum and maximum <dl
at all stations.

Nitrogen as Nitrate $\text{NO}_3\text{-N}$

Main River (analyzed at 3 of 7 stations only) - Minimum 0.01 at Flesher and Canyon. Maximum 0.60 at Canyon.
Tributaries - (analyzed at 4 of 29 stations only) - Minimum 0.01 at Alice Creek. Maximum 0.44 at Hardscrabble Creek.

Organic Nitrogen

Main River (analyzed at 3 of 7 stations only) - Minimum <dl at Flesher. Maximum 0.85 at Dalton.
Tributaries (analyzed at 4 of 29 stations only) - Minimum <dl at Willow Creek (Flesher) and Alice Creek. Maximum 1.5 at Poorman Creek.

Total Phosphate, PO_4^{--}

Main River (analyzed at 3 of 7 stations only) - Minimum <dl at all 3 stations. Maximum 0.19 at Canyon.
Tributaries (analyzed at 4 of 29 stations only) - Minimum 0.01 at Alice Creek and Poorman Creek. Maximum 0.10 at Poorman Creek.

Ortho Phosphate, PO_4^{--}

Main River (analyzed at 3 of 7 stations only) - Minimum <dl at all 3 stations. Maximum 0.04 at Canyon.
Tributaries (analyzed at 4 of 29 stations only) - Minimum <dl at Poorman Creek and Hardscrabble Creek. Maximum 0.08 at Hardscrabble Creek.

APPENDIX C

Analytical Detection levels of Chemical Constituents

<u>Constituents</u>	<u>Minimum Detection Level (mg/l)</u>	
Residue, total	10	
Hardness, total	.1	
CO ₃ (above pH 8.3)	6	Prior to July 1972
	1.2	since July 1972
HCO ₃	6	Prior to July 1972
	1.2	since July 1972
SO ₄	1.0	
Cl	1.0	Prior to June 1972
	.5	since June 1972
NO ₃	.01	
F	.05	
Ca	.1	
Mg	.1	
Na	.1	since Feb. 1971*
Fe	.01	
As	.001	
Zn	.01	
Pb	.01	
Cu	.01	
NO ₂ -N	.001	
NH ₃ -N	.01	
NO ₃ -N	.01	
Org. N	.01	
PO ₄ , total	.01	
PO ₄ , Ortho	.01	

*Prior to February 1971, Na was calculated. Since then it has been measured directly by atomic absorption.

Note: In all cases except NO₂-N and Arsenic (As), results are reported to two significant figures. NO₂-N and As are reported to three significant figures.

APPENDIX D

Summary of Laboratory Analyses with Chemical Results Less Than and Greater Than the Analytical Detection Level

Samples collected during the period February 1968 through August 1973. Main river stations are given in descending order downstream followed by tributaries in descending order downstream.

RM	=	River Mile
N	=	Sample size, or total number of laboratory analyses made during the period.
d1	=	Analytical detection level of chemical constituent (see Appendix C).
N<d1	=	Number of analyses with chemical results less than the detection level.
%<d1	=	Percent of analyses with chemical results less than the detection level.
Range>d1	=	Lowest (L) and highest (H) chemical results which were greater than the detection level.
$\bar{X}>d1$	=	Average value of chemical results which were greater than the detection level

All values in mg/liter except turbidity (JTU).

A blank space in a column means all values were less than the detection level. Numbers in parentheses indicate only one value was greater than the detection level. A dash (-) indicates no data were collected. When all spaces below a dash (-) are blank, this indicates no data were collected.

Analyses were done in accordance with Standard Methods for the Examination of Water and Wastewater, current edition.

APPENDIX D continued

Turbidity (NTU)	Total Residue	Total Hardness	Total Alkalinity (Calc.)											Total Recoverable					Dissolved			
				CO ₂	HCO ₃	SO ₄	Cl	NO ₃	F	Ca	Mg	Na		Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

Blackfoot River at Pop's Place (RM 119.4)

N	6	8	8	8	8	8	8	8	8	8	7	8	6	8	2	8	-	-	-	-
N <dl	5	0	0	0	0	0	0	4	1	0	0	2	1	6	0	2	8			
% <dl	83	0	0	0	0	0	0	50	13	0	0	29	13	100	0	100	100			
Range >dl	(3)	76	60	35	0	43	10	.07	.04	10	2	5	.02		.01					
L																				
H		186	160	62	0	76	79	9	2.2	.20	29	16	26	.27		.44				
I >dl	(3)	130	96	51	0	62	41	3.4	.60	.18	20	12	8.8	.12		.22				

Blackfoot River at Fleisher Road (RM 114.6)

N	-	41	41	41	41	41	41	41	36	41	41	39	48	20	48	15	48	13	15	14	15	
N <dl	-	0	0	0	0	0	2	19	20	3	0	0	13	16	17	8	13	36	3	3	13	9
% <dl	-	0	0	0	0	0	5	46	49	8	0	0	33	33	85	15	87	75	23	20	87	60
Range >dl	-	80	60	55	0	67	2	.5	.01	.01	12	2	1	.01	.001	.02	.01	.01	.01	.03	(.01)	.01
L																						
H	-	152	220	125	0	152	43	9	3.6	.34	32	37	18	.5	.002	.94	.01	.18	.11	.17		.02
I >dl	-	115	111	80	0	98	22	4.2	.29	.15	22	14	4.8	.12	.001	.11	.01	.03	.04	.09	.01	.01

Blackfoot River at Hogum Road (RM 110.7)

N	15	26	26	26	26	26	26	26	21	26	26	25	26	7	25	2	25		
N <dl	9	0	0	0	0	0	1	4	20	5	0	0	11	19	7	19	2	21	
% <dl	67	0	0	0	0	0	4	15	77	24	0	0	44	73	100	76	100	84	
Range >dl	2	96	76	65	0	79	2	.5	.1	.08	14	2	1	.08		.01		.02	
L	9	142	230	125	0	152	20	8	5.3	.36	32	39	21	.6		.01		.05	
H	4.7	124	128	107	0	131	9.3	3.7	1.0	.20	27	16	8.5	.26		.02		.04	

APPENDIX D continued

Turbidity (NTU)	Total Residue	Total Hardness	Total Alkalinity (Calc.)	CO ₂	HCO ₃	SO ₄	Cl	NO ₃	F	Ca	Mg	Na	Total Recoverable					Dissolved			
													Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

Blackfoot River Below 7-Up Pete Creek (RM 106.2)

N	-	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2		
	-	0	0	0	0	0	1	2	0	0	0	0	0	0	2	1	2	1	0	1	2	2
	-	0	0	0	0	0	0	50	100	0	0	0	0	0	0	100	50	100	50	0	50	100
Range >dl	L	-	210	131	124	0	151	(11)	.05	.1	33	11	2.2	.03	(.01)	(.01)	.03	(.01)				
	H	-	210	135	133	2	157		.17	.1	36	12	2.4	.06			.06					
I >dl	-	210	133	129	1	154	(11)		.11	.1	34	12	2.3	.04	(.01)	(.01)	.04	(.01)				

Blackfoot River at Dalton Mountain Road Bridge (RM 95.9)

N	17	42	41	42	41	42	42	41	41	36	41	41	41	48	20	50	15	50	12	14	13	14
N <dl	9	0	0	0	0	0	9	15	24	6	0	0	16	24	15	29	13	37	7	12	11	9
% <dl	53	0	0	0	0	0	21	37	59	17	0	0	39	50	75	58	87	74	58	86	85	64
Range >dl	L	1	88	102	110	0	134	1.6	.5	.03	.06	20	1	1.3	.01	.002	.01	.01	.01	.01	.01	.01
H	9	261	340	189	6	230	12	7	2.8	.3	52	37	30	1.3	.002	.59	.01	.07	.19	.01	.01	.02
I >dl	2.1	161	167	154	.24	187	5.8	3.7	.29	.16	39	16	6.8	.26	.002	.08	.01	.02	.06	.02	.01	.01

Blackfoot River at Blackfoot Canyon Camp (RM 84.7)

N	14	40	40	40	40	40	40	40	40	36	40	40	38	47	19	47	15	47	11	14	13	14
N <dl	7	0	0	0	0	0	8	17	20	4	0	1	17	21	14	26	13	34	7	10	11	7
% <dl	50	0	0	0	0	0	20	43	50	11	0	3	45	45	74	55	87	72	64	71	85	50
Range >dl	L	1.5	110	60	100	0	122	.4	.5	.01	.02	24	1	1.2	.02	.001	.01	.01	.01	.01	.01	.01
H	10	260	250	180	6	220	16	9	1.4	.4	56	32	33	1.5	.007	.30	.01	.06	.06	.01	.01	.02
I >dl	4.6	166	166	154	.2	187	5.5	3.0	.26	.17	41	16	8.5	.27	.003	.05	.01	.02	.04	.01	.01	.01

APPENDIX D continued

Turbidity (FTU)	Total Sediment	Total Hardness	Total Alkalinity (calc.)	CO ₃	HCO ₃	SO ₄	Cl	NO ₃	F	Ca	Mg	Na	Total Recoverable					Dissolved			
													Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

Blackfoot River at Rainbow Bend (RM 9,7)

N	17	30	30	30	30	30	30	30	30	26	30	30	30	30	7	29	2	29	-	-	-	-
N < d1	7	0	0	0	0	0	3	8	16	5	0	0	16	10	6	19	2	25				
% < d1	41	0	0	0	0	0	10	27	53	19	0	0	53	33	86	66	100	14				
L	2	60	80	85	0	98	4	5	.04	.02	19	1	1	.01	(.008)	.01	.02					
Range > d1	20	170	230	148	12	180	10	31	4.0	.44	46	24	20	1.0	.08	.06						
H	8	137	148	128	.4	155	5.8	4.8	.54	.17	33	16	7.5	.24	(.008)	.04	.04					

Pase Creek (RM 120,4-0,8)

N	7	8	8	8	8	8	8	8	8	6	8	8	7	8	2	8	-	8	-	-	-
N < d1	5	0	0	0	0	0	4	0	6	2	0	1	5	8	2	6	-	8	-	-	-
% < d1	71	0	0	0	0	0	50	0	75	33	0	13	71	100	100	75	-	100	-	-	-
L	1.5	68	68	50	0	61	2	1	.2	.10	16	5	1		.10	-					
Range > d1	15	100	160	90	0	110	5	11	.6	.60	32	29	6								
H	8.1	85	99	77	0	94	3	5.6	.60	.24	.23	12	3.5		.14	-					

Sandbar Creek (RM 114,3-2,8-0,1)

N	1	1	1	1	1	1	1	1	1	1	1	1	1	1	-	1	-	1	-	-
N < d1	0	0	0	0	0	0	0	0	1	0	0	0	1	0	-	0	-	0	-	0
% < d1	0	0	0	0	0	0	0	0	100	0	0	0	100	0	-	0	-	0	-	0
L	(27)	(88)	(150)	(45)	(0)	(55)	(11)	(7)	(.10)	(12)	(29)	(2.5)	-	(.01)	-	(.02)	-	-	-	-
Range > d1																				
H																				
% > d1	Same as Range > d1																			

APPENDIX D continued

Turbidity (JTU)	Total Hardness	Total Hardness	Total Alkalinity (CaCl ₂)	CO ₃	HCO ₃	SO ₄	Cl	NO ₃	F	Ca	Mg	Na	Total Recoverable					Dissolved			
													Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

Willow Creek (Floater) (RM 114.3-0.0)

		-	8	8	8	8	8	8	8	8	8	8	8	6	6	8	7	8	6	8	7	8	
N		-	0	0	0	0	0	1	6	0	0	0	0	0	4	5	6	6	0	7	6	6	
N < d1		-	0	0	0	0	0	13	75	0	0	0	0	0	57	63	86	75	0	88	86	75	
% < d1		-	66	49	47	0	57	.8	.5	.10	.05	11	5.2	1.1	.16	.002	.01	(.01)	.01	.07	(.01)	(.01)	.01
Range > d1	L	-	135	93	83	0	101	21	1.0	.28	.10	28	8.0	6.8	.33	.002	.01	.02	.20	.01	.01	.01	
I > d1	H	-	91	71	68	0	83	7.6	.8	.19	.09	18	6.4	2.6	.24	.002	.01	(.01)	.02	.17	(.01)	(.01)	.01

Aline Creek at Highway 200 (RM 113.2-0.9)

N	7	30	30	30	30	30	30	30	29	28	30	30	30	28	15	30	15	30	13	15	14	15
N < d1	5	0	0	0	0	0	13	21	8	4	0	0	5	14	12	18	11	21	8	13	12	11
% < d1	71	0	0	0	0	0	43	70	28	14	0	0	17	50	80	60	73	70	62	87	86	73
Range > d1	3	68	69	61	0	74	1	.5	.02	.03	16	4	.9	.01	.001	.01	.01	.01	.02	.01	.01	.01
I > d1	9	214	240	135	0	165	11	7	.16	.38	44	19	5.2	.90	.002	.72	.01	.06	.07	.02	.01	.01
I > d1	6	120	116	110	0	135	5.7	3.9	.15	.15	27	13	1.6	.18	.002	.11	.01	.02	.05	.02	.01	.01

Hardscrabble Creek (RM 112.5-0.5)

N	-	9	9	9	9	9	9	9	9	9	9	9	9	9	8	9	8	9	9	9	8	9
N < d1	-	0	0	0	0	0	3	7	0	0	0	0	0	1	7	5	7	4	2	8	7	6
% < d1	-	0	0	0	0	0	33	78	0	0	0	0	0	11	88	56	88	14	78	11	13	33
Range > d1	-	110	85	107	0	130	1.6	.5	.09	.07	22	7.2	2.3	.02	(.002)	.01	(.01)	.01	.02	(.02)	(.01)	.01
	-	288	178	181	6	214	8.2	1.0	2.0	.20	47	16	7.2	.28	.04	.02	.02	.16	.02	.16	.01	.01
	-	198	140	154	1.6	184	5.6	.75	.37	.11	35	12	3.9	.11	.002	.02	.01	.01	.06	.02	.01	.01

APPENDIX D continued

Turbidity (FTU)	Total Residue	Total Hardness	Total Alkalinity (Calc.)											Total Recoverable					Dissolved			
				CO ₂	HCO ₃	SO ₄	Cl	NO ₃	F	Ca	Mg	Na		Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

Borefly Creek (RM 112.2-0.5)

N	-	1	1	1	1	1	1	1	1	1	1	1	1	1	-	1	-	1	1	1	-	1
N < dl	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0
% < dl	0	0	0	0	0	0	100	100	0	0	0	0	0	0	0	100	0	0	100	0	0	0
Range > dl	(66)	(50)	(60)	(0)	(73)				(0.1)	(0.1)	(16)	(2.6)	(3.4)	(.08)				(.01)	(.04)			(.02)
H	Same as Range > dl																					
I > dl	Same as Range > dl																					

Black Diamond Creek (RM 110.3-1.2-0.0)

N	-	1	1	1	1	1	1	1	1	1	1	1	1	1	-	1	-	1	1	1	-	1
N < dl	0	0	0	0	0	1	1	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0
% < dl	0	0	0	0	0	100	100	0	0	0	0	0	0	0	100	0	0	100	0	0	100	0
Range > dl	(70)	(68)	(82)	(0)	(100)			(0.1)	(.20)	(39)	(4.7)	(3.5)	(.18)				(.02)	(.14)			(.01)	
H																						
I > dl	Same as Range > dl																					

Bogus Creek (RM 110.3-0.2)

N	-	10	10	10	10	10	10	10	10	10	10	10	10	10	10	-	8	-	8	-	-	-	-
N < d1	0	0	0	0	0	0	0	8	4	0	0	0	0	7	1	4	7						
% < d1	0	0	0	0	0	0	0	80	40	0	0	0	0	70	10	50	88						
Range > d1	L	60	70	57	0	70	2	2	.04	.08	12	5	4.1	.10	.03	(.03)							
	H	132	190	111	0	135	22	10	.20	.30	32	29	13	.50	.07								
I > d1		108	128	91	0	111	8.1	6.0	.12	.21	25	16	9.0	.26	.05	(.03)							

APPENDIX D continued

Turbidity (JTU)	Total Residue	Total Hardness	Total Alkalinity (Calc.)					NO ₃	F	Ca	Mg	Na	Total Recoverable					Dissolved			
				CO ₂	HCO ₃	SO ₄	Cl						Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

Landers Fork (RM 107.2-0.8)

W	8	12	12	12	12	12	12	12	12	12	12	12	12	12	5	12	1	12	3	3	3	3
N <dl	7	0	0	0	0	0	2	4	7	3	0	0	6	10	5	10	3	12	2	3	3	3
% <dl	88	0	0	0	0	0	17	11	58	30	0	0	50	81	100	81	100	100	67	100	100	100
Range >dl	(4)	116	96	70	0	85	1	1	.15	.02	22	8.8	1.0	.22		.03			(.10)			
L		225	210	148	2	180	12	9	.42	.36	42	39	12	.31		.46						
H	(4)	150	146	125	.17	152	4.3	4.4	.24	.16	34	16	4.4	.27		.24			(.10)			

Poorman Creek (RM 99.4-2.2)

N	7	28	28	28	28	28	28	28	24	26	27	27	27	26	14	28	13	26	12	14	13	14
N <dl	4	0	0	0	0	0	1	18	9	1	0	0	6	17	9	18	11	18	9	11	11	9
% <dl	57	0	0	0	0	0	4	64	38	4	0	0	22	66	64	64	85	69	75	79	85	64
Range >dl	1	100	95	70	0	86	1.0	.5	.01	.02	20	2	.9	.03	.001	.01	.01	.01	.01	.01	.01	.01
	10	209	230	172	24	210	18	8	.63	.36	60	20	19	.37	.002	.32	.02	.01	.06	.01	.01	.02
L																						
H	5	137	128	121	1.2	145	9.6	3.6	.20	.16	36	9.1	2.8	.12	.002	.08	.02	.02	.03	.01	.01	.01
>dl																						

Spring Creek (RM 96.6-0.5-1.3)

N	8	8	8	8	8	8	8	8	7	8	8	8	8	2	8	-	8	-	-	-	-
N <dl	5	0	0	0	0	0	4	0	5	2	0	0	1	5	2	8	-	7			
% <dl	33	0	0	0	0	0	50	0	63	29	0	0	38	63	100	100	-	86			
Range >dl	2	162	148	160	0	195	4	1	.20	.10	32	10	2	.08		-	(.02)				
L	10	190	260	180	0	230	15	8	.30	.52	48	34	17	.10		-					
H	5	175	187	171	0	208	8.8	4.6	.27	.25	42	20	8.6	.09			(.02)				
>dl																					

APPENDIX D continued

Turbidity (JTU)	Total Residue	Total Hardness	Total Alkalinity (CaCO ₃)	CO ₂	HCO ₃	SO ₄	Cl	NO ₃	F	Ca	Mg	Na	Total Recoverable					Dissolved			
													Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

Keep Cool Creek (RM 96.6-1.0)

N	7	8	8	8	8	8	8	8	6	8	8	8	8	2	8	-	8	-	-	-	-
N <dl	4	0	0	0	0	0	4	0	7	1	0	0	5	5	2	7	-	8			
% <dl	57	0	0	0	0	0	50	0	87	17	0	0	63	63	100	87	-	100			
Range >dl	3	136	120	125	0	152	1	1	(.10)	.02	34	8	7	.10	(.02)	-					
H	17	180	260	165	0	201	7	9	.34	50	26	18	.30			-					
X >dl	8	160	169	150	0	184	4	5	(.10)	.17	42	16	14	.20		(.02)	-				

Beaver Creek (RM 96.6-0.7-0.2)

N	7	8	8	8	8	8	8	8	6	8	8	8	8	2	8	-	8	-	-	-	-
N <dl	4	0	0	0	0	0	2	0	6	3	0	0	5	6	3	7	-	7			
% <dl	57	0	0	0	0	0	25	0	75	50	0	0	63	75	50	87	-	87			
Range >dl	2	110	76	85	0	104	1	1	.10	.08	25	2	4	.10	(.006)	(.01)	-	(.04)			
H	8	170	270	165	0	201	8	10	.20	.38	52	37	16	.12			-				
X >dl	5	146	155	135	0	165	3.5	5.2	.15	.19	42	12	11	.11	(.006)	(.01)	-	(.04)			

Willow Creek (RM 94.2-1.2)

N	7	8	8	8	8	8	8	8	5	8	8	8	8	1	8	1	8	-	-	-	-
N <dl	1	0	0	0	0	0	0	0	5	1	0	0	4	1	1	6	1	7			
% <dl	14	0	0	0	0	0	0	0	63	20	0	0	50	13	100	75	100	88			
Range >dl	4	78	55	40	0	49	4	2	.10	.10	14	1	7	.10		.01	(.02)				
H	20	138	180	100	0	122	17	8	3.6	.38	28	27	12	.50		.02					
X >dl	11	102	92	77	0	94	9.9	4.9	1.4	.19	23	8.6	9.2	.21		.02	(.02)				

APPENDIX D continued

Turbidity (NTU)	Total Residue	Total Hardness	Total Alkalinity (Calc.)											Total Recoverable					Dissolved			
				CO ₃	HCO ₃	SO ₄	Cl	NO ₃	F	Ce	Mg	Na		Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

Seaukraut Creek (RM 93.7-0.2)

N	-	1	1	1	1	1	1	1	1	1	1	1	1	1	-	1	-	1	1	1	-	1
N <dl	-	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	-	0	0	1	-	0
S <dl	-	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	-	0	0	100	-	0
Range >dl	-																					
H	-	One Sample Only																				
I >dl	-	(50)	(79)	(50)	(0)	(61)	(0)	(0)	(.2)	(0)	(12)	(2.1)	(1)	(.02)	-	(.01)	-	(.02)	(.02)	-	(.01)	

Arroyo Creek (RM 79.9-0.1)

N	7	8	8	8	8	8	8	8	8	8	8	8	8	8	3	8	1	8				
N <dl	4	0	0	0	0	0	0	0	0	4	5	1	0	0	4	5	1	7	1	6		
S <dl	57	0	0	0	0	0	0	0	0	50	71	20	0	0	50	63	100	88	100	75		
Range >dl	5	72	51	65	0	79	1	1	.20	.06	19	1	7	.06	(.01)				.01			
H	33	136	170	115	0	140	14	9	2.5	.38	30	24	9	.12						.04		
I >dl	2	105	107	98	0	119	4.8	3.9	1.4	.16	25	11	8.2	.09	(.01)					.04		

Kleinschmidt Creek (RM 51.8-5.6-0.1)

N	7	8	8	8	8	8	8	8	8	8	8	8	8	8	1	8	1	8				
N <dl	3	0	0	0	0	0	0	0	0	4	1	0	1	3	7	2	7	1	7			
S <dl	43	0	0	0	0	0	0	0	0	50	20	0	13	38	88	67	88	100	88			
Range >dl	1	120	130	125	6	150	2	2	.20	.06	28	13	2	(.04)	(.001)	(.02)			(.02)			
H	4	160	230	165	10	201	13	11	6.2	.28	51	37	22									
I >dl	3	146	160	150	2	178	5.8	4.8	1.8	.14	35	20	11	(.04)	(.001)	(.02)			(.02)			

APPENDIX D continued

Turbidity (JTU)	Total Residue	Total Hardness	Total Alkalinity (Calc.)	CO ₃	HCO ₃	SO ₄	Cl	NO ₃	F	Ca	Mg	Na	Total Recoverable					Dissolved			
													Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

North Fork Blackfoot River (RM 51.8-5.6)

N	5	7	7	7	7	7	7	7	5	7	7	7	7	3	7	3	7				
N <dl	4	0	0	0	0	0	1	0	5	1	0	0	4	6	3	5	3	6			
% <dl	67	0	0	0	0	0	14	0	71	20	0	0	57	86	100	71	100	86			
Range >dl	(6)	106	105	105	0	140	1	2	.34	.02	28	2	10	(.50)		.01	(.02)				
L		150	220	150	0	183	12	8	.70	.10	44	34	17			.05					
H																					
I >dl	(6)	130	144	132	0	162	4.5	4.7	.52	.06	34	14	14	(.50)		.03	(.02)				

Warren Creek (RM 48.2-4.4)

N	2	1	2	1	3	3	3	3	1	1	3	3	2	3	2	3	1	3			
N <dl	0	0	0	0	0	0	1	0	1	1	0	0	1	0	2	3	1	3			
% <dl	0	0	0	0	0	0	33	0	33	33	0	0	50	0	100	100	100	100			
Range >dl	8	122	75	94	0	115	3	3	.10	.06	24	4	(16)	.10							
L	8	140	180	118	0	144	3	5	.50	.20	33	24		.30							
H																					
I >dl	8	131	132	106	0	129	3	4	.45	.11	29	15	(16)	.20							

Monture Creek (RM 44.2-3.0)

N	7	8	8	8	8	8	8	8	8	6	8	8	8	8	3	8	1	5			
N <dl	3	0	0	0	0	0	2	0	5	2	0	0	4	5	1	7	1	5			
% <dl	43	0	0	0	0	0	25	0	63	33	0	0	50	61	100	88	100	100			
Range >dl	2	46	34	35	0	43	1	1	.10	.06	12	1	1	.04		(.01)					
L	13	126	180	120	0	147	11	9	6.0	.14	32	32	9	.16							
H																					
I >dl	6	84	96	80	0	98	5.8	4.1	2.2	.08	20	11	5.5	.07		(.01)					

APPENDIX D continued

Turbidity (NTU)	Total Residue	Total Hardness	Total Alkalinity (calc.)	CO ₂	HCO ₃	SO ₄	Cl	NO ₃	F	Ca	Mg	Na	Total Recoverable					Dissolved			
													Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

Cottonwood Creek (RM 41.6-0.6)

	1	5	5	5	5	5	5	5	3	5	5	4	5	3	5	1	5	
N																		
N < d1	2	0	0	0	0	0	0	0	3	2	0	0	3	4	3	4	1	5
% < d1	50	0	0	0	0	0	0	0	60	67	0	0	75	80	100	80	100	100
Range > d1	7	132	125	104	0	127	1	3	5	(.1)	28	12	(20)	(.10)		(.02)		
	7	176	180	153	0	187	20	7	5	40	18							
H	7	155	155	131	0	160	6.4	5.2	50	(.1)	37	15	(20)	(.1)		(.02)		

Clearwater River (RM 33.8-3.0)

N	7	8	8	8	8	8	8	8	8	8	8	8	8	8	3	8	1	8	
N < d1	3	0	0	0	0	0	3	0	5	2	0	1	3	6	3	6	1	7	
% < d1	43	0	0	0	0	0	38	0	61	40	0	13	38	75	100	75	100	88	
Range > d1	4	60	70	60	0	73	2	1	10	0.4	14	5	3	.06		.01	(.03)		
	10	105	160	102	0	125	7	8	8.9	.15	30	22	10	.10		.02			
E > d1	6	82	88	80	0	98	4	3.9	3.2	.09	21	10	6.4	.08		.02	(.01)		

Blanchard Creek (RM 33.8-2.6-0.2)

	1	2	2	2	2	2	2	2	2	1	2	2	2	2	1	2	2	2	
N																			
N < d1	0	0	0	0	0	0	2	0	1	1	0	0	1	1	1	1	2	2	
% < d1	0	0	0	0	0	0	100	0	50	100	0	0	50	50	100	50	100	100	
Range > d1	(8)	60	35	45	0	55	2		(.1)	8	4	(5)		(.07)		(.03)			
		86	100	80	0	98	4			18	13								
H																			
I > d1	(8)	73	68	62	0	77	3		(.1)	13	9	(5)		(.07)		(.03)			

APPENDIX D continued

Turbidity (JTU)	Total Residue	Total Hardness	Total Alkalinity (calc.)	CO ₂	HCO ₃	SO ₄	Cl	NO ₃	F	Ca	Mg	Na	Total Recoverable					Dissolved			
													Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu

Elk Creek (RM 27.9-1.0)

N	5	6	6	6	6	6	6	6	4	6	6	6	6	3	6	1	6				
N <dl	0	0	0	0	0	0	0	4	0	0	0	2	2	3	4	1	6				
S <dl	0	0	0	0	0	0	0	67	0	0	0	33	33	100	67	100	100				
Range >dl	4	128	110	75	0	92	7	2	.60	.10	28	2	4	.10		.02					
H	285	188	220	155	6	189	14	9	3.2	.28	44	27	22	.90		.20					
X >dl	79	164	141	130	1	156	11	5.2	1.9	.21	38	11	13	.67		.11					

Belmont Creek (RM 21.2-0.1)

N	2	3	3	3	3	3	3	3	2	3	3	2	3	2	3	1	3				
N <dl	1	0	0	0	0	0	3	0	3	2	0	0	1	1	2	3	1	3			
S <dl	50	0	0	0	0	0	100	0	100	100	0	0	50	33	100	100	100	100			
Range >dl	(7)	110	80	95	0	116	3		24	5	(10)		.05								
H		160	210	144	0	175	5		68	20			.40								
X >dl	4	141	160	127	0	155	4		45	12	10		.23								

Gold Creek (RM 13.0-0.0)

N	2	3	3	3	3	3	3	3	2	3	3	1	3	2	3	1	3				
N <dl	1	0	0	0	0	0	2	0	3	2	0	0	1	2	2	3	1	3			
S <dl	50	0	0	0	0	0	67	0	100	100	0	0	100	67	100	100	100	100			
Range >dl	(7)	100	95	85	0	104	(4)	2	20	11			(.40)								
H		110	125	100	0	122	5		32	12											
X >dl	(7)	103	113	94	0	115	(4)	3.7	27	11			(.40)								

APPENDIX D continued

Turbidity (JTU)	Total Residue	Total Hardness	Total Alkalinity (Calc.)												Total Recoverable					Dissolved			
				CO ₂	HCO ₃	SO ₄	Cl	NO ₃	F	Ca	Mg	Na	Fe	As	Zn	Pb	Cu	Fe	Zn	Pb	Cu		

Union Creek (RM 12.3-0.0)

N	7	8	8	8	8	8	8	8	8	8	8	8	8	3	8	1	8	
N < dl	0	0	0	0	0	0	1	0	5	0	0	0	2	0	2	7	1	6
% < dl	0	0	0	0	0	0	13	0	63	0	0	0	25	0	67	88	100	75
Range) dl	9	150	105	100	0	122	8	2	.60	.10	.24	2	1	.10	(.004)	(.06)		.02
H	200	242	310	226	0	275	18	11	1.2	.28	.62	51	20	.80				.02
I > dl	46	208	124	177	0	215	11	6.6	.90	.17	.40	23	8.3	.36	(.004)	(.06)		.02

West Twin Creek (RM 10.3-0.0)

N	3	3	3	3	3	3	3	3	3	3	3	3	3	2	3	1	3	
N < dl	1	0	0	0	0	0	1	0	1	0	0	2	0	1	2	2	1	2
% < dl	33	0	0	0	0	0	33	0	33	0	0	67	0	33	100	67	100	67
Range) dl	7	24	8	20	0	24	5	2	.20	.10	2	(1)	5	.06		(.01)		(.12)
H	11	50	20	30	0	36	8	3	2.7	.44	8	12		.18				
I > dl	9	39	13	24	0	29	6.5	2.3	1.4	.27	4.7	1	8.7	.12		(.01)		(.12)

West Twin Creek (RM 10.0-0.0)

N	3	3	3	3	3	3	3	3	3	3	3	3	3	2	3	1	3	
N < dl	1	0	0	0	0	0	2	0	2	0	0	2	2	1	2	3	1	1
% < dl	33	0	0	0	0	0	67	0	67	0	0	67	67	33	100	100	100	100
Range) dl	3	10	7	5	0	6	(4)	2	(.21)	.04	2	(8)	(5)	.05				
H	7	30	40	28	0	34	2		.34	4				.08				
I > dl	5	21	19	17	0	20	(4)	2	(.23)	.21	3	(8)	(5)	.06				

APPENDIX D continued

Summary of Laboratory Analyses for N + P Compounds with Chemical
Results Less Than and Greater Than the Analytical Detection Level

Blackfoot River at Fleisher Road (Rm 114.6)

	NO ₂ -N	NH ₃ -N	NO ₃ -N	Org. N	Total PO ₄	Ortho PO ₄
N	6	6	6	6	7	6
N < dl	5	6	0	1	3	4
% < dl	83	100	0	17	43	67
Range > dl L	(.001)		.01	.08	.01	.01
H			.08	.39	.03	.01
X > dl	(.001)		.03	.28	.02	.01

A-12

Blackfoot River at Dalton Mountain Road (Rm 95.9)

	NO ₂ -N	NH ₃ -N	NO ₃ -N	Org. N	Total PO ₄	Ortho PO ₄
N	7	7	8	7	7	7
N < dl	6	7	0	0	2	3
% < dl	86	100	0	0	29	43
Range > dl L	.002		.02	.18	.01	.01
H	.002		.16	.85	.17	.03
X > dl	.002		.06	.44	.05	.02

Blackfoot River at Blackfoot Canyon Camp (Rm 84.7)

	NO ₂ -N	NH ₃ -N	NO ₃ -N	Org. N	Total PO ₄	Ortho PO ₄
N	7	7	7	7	7	6
N < dl	7	7	0	0	2	3
% < dl	100	100	0	0	29	50
Range > dl L			.01	.20	.01	.01
H			.60	.75	.19	.04
X > dl			.17	.38	.06	.02

Willow Creek (Fleisher) (Rm 114.2 = 0.0)

	NO ₂ -N	NH ₃ -N	NO ₃ -N	Org. N	Total PO ₄	Ortho PO ₄
N	2	2	2	2	2	2
N < dl	2	2	0	1	0	0
% < dl	100	100	0	50	0	0
Range > dl L			.03	(.39)	.02	.01
H			.03		.02	.02
X > dl			.03	(.39)	.02	.02

APPENDIX D continued

Alice Creek at Highway 200 (Rm 113.2 - 0.9)

	NO ₂ -N	NH ₃ -N	NO ₃ -N	Org. N	Total PO ₄	Ortho PO ₄
N	7	7	7	7	7	7
N < dl	6	7	0	1	1	2
% < dl	86	100	0	14	14	29
Range > dl L	(.003)		.01	.03	.01	.01
H			.14	.50	.06	.02
X > dl	(.003)		.05	.30	.02	.02

Peorinen Creek (Rm 99.4 - 2.2)

	NO ₂ -N	NO ₃ -N	NO ₃ -N	Org. N	Total PO ₄	Ortho PO ₄
N	7	7	5	7	7	7
N < dl	4	7	0	0	0	1
% < dl	57	100	0	0	0	14
Range > dl L	.001		.02	.20	.01	.01
H	.002		.07	1.5	.10	.03
X > dl	.002		.51	.12	.02	.03

A-33

Hardacre Creek (Rm 112.5 - 0.5)

	NO ₂ -N	NH ₃ -N	NO ₃ -N	Org. N	Total PO ₄	Ortho PO ₄
N	5	5	4	5	5	5
N < dl	5	5	0	0	0	1
% < dl	100	100	0	0	0	20
Range > dl L			.03	.27	.03	.02
H			.45	.50	.08	.08
X > dl			.16	.42	.06	.05

APPENDIX E
Water Quality Station Locations and River Mileages

Station No.	Name	Location 1/	River Miles Above Mouth 2/	Station No.	Name	Location 1/	River Miles Above Mouth 2/
1	Blackfoot River at Rainbow Bend	13N 17W 2 con	9.7	22	Spring Creek	14N 9W 23 bob	96.6 - 0.5 - 1.3
2	West Twin Creek	13N 17W 2 cad	10.0 - 0.0	23	Pocraman Creek	14N 9W 36 ddd	99.4 - 2.2
3	East Twin Creek	13N 17W 2 dbe	10.3 - 0.0	24	Blackfoot River below 7-up Pete Creek	14N 8W 14 odd	106.2
4	Union Creek	13N 16W 6 dac	12.3 - 0.0	25	Landers Fork	14N 8W 12 eac	107.2 - 0.8
5	Gold Creek	13N 17W 6 bbd	13.0 - 0.0	26	Hogum Creek	14N 7W 8 acb	110.3 - 0.2
6	Belmont Creek	14N 16W 24 bod	21.2 - 0.1	27	Black Diamond Creek	14N 7W 8 dda	110.3 - 1.2 - 0.0
7	Kik Creek	14N 15W 36 aed	27.9 - 1.0	28	Blackfoot River at Hogum Creek Road	14N 7W 5 odd	110.7
8	Blanchard Creek	14N 14W 5 dsa	33.8 - 2.6 - 0.2	29	Horsefly Creek	14N 7W 4 ada	112.2 - 0.5
9	Clearwater River	14N 14W 4 bdb	33.8 - 3.0	30	Handscrabble Creek	15N 7W 33 aed	112.5 - 0.5
10	Ottomwood Creek	15N 13W 29 obb	41.6 - 0.6	31	Alice Creek	15N 7W 27 dbc	113.2 - 0.9
11	Menture Creek	15N 13W 24 bdb	44.2 - 3.0	32	Willow Creek (Flasher)	15N 7W 34 daa	114.3 - 0.0
12	Warren Creek	15N 12W 35 ada	48.2 - 4.4	33	Sandbar Creek	14N 6W 6 cdd	114.3 - 2.8 - 0.1
13	North Fork Blackfoot River	14N 11W 6 acb	51.8 - 5.6	34	Blackfoot River at Flasher Road	15N 7W 35 bca	114.6
14	Kleinschmidt Creek	14N 11W 6 aod	51.8 - 5.6 - 0.1	35	Blackfoot River at Pop's Place	15N 6W 19 aac	119.4
15	Arresta Creek	14N 10W 30 aab	79.9 - 0.1	36	Pace Creek	15N 6W 17 ddd	120.4 - 0.8
16	Blackfoot River at Blackfoot Canyon Camp	14N 10W 34 bdc	84.7		Source (Junction of Anaconda Creek and Beartrap Creek)	15N 6W 27 bbb	122.0
17	Sauerkraut Creek	14N 9W 29 cad	93.7 - 0.1				
18	Willow Creek	14N 9W 28 adc	94.2 - 1.2				
19	Blackfoot River at Dalton Mountain Road	14N 9W 28 aba	95.9				
20	Beaver Creek	14N 9W 22 bda	96.6 - 0.7 - 0.2				
21	Keep Cool Creek	14 N 9W 22 abc	96.6 - 1.0				

1/ Location method given in Figure 5.

2/ Mileages obtained from the following sources: Mouth to Willow Creek from USGS map "Plan and Profile of Blackfoot River, to Mile 95, and Tributaries, Montana, Miscellaneous Data Sites", dated 1937. Willow Creek to Source, measured from the following USGS 7 1/2 minute quadrangle maps: Moose Creek, 1948; Lincoln, 1948; Swede Gulch, 1968; Stemple Pass, 1968; Cadotte Creek, 1963; Rogers Pass, 1963.

APPENDIX F

Results of BOD Analyses

Samples were collected by the writer from February 1972 through September 1972 and analyses were made by the USGS, Helena. Collections and analyses were both made by the USGS during water year 1973 (October 1972 - September 1973).

A 1 liter water sample was collected and supersaturated with oxygen. Dissolved oxygen was then measured. Five days later dissolved oxygen content was again measured. The difference between the first and second dissolved oxygen levels was the five day BOD.

APPENDIX F continued

	<u>Date</u>	<u>Time</u>	<u>5-day BOD (mg/l)</u>
	Blackfoot River at Hogum Road Bridge (USGS gage "below Alice Creek near Lincoln")		
1972	2-2	1330	1.8
	3-23	1200	1.1
	4-14	1010	1.2
	5-19	1050	1.8
	8-2	1020	1.1
	9-8	1010	3.0
	10-25	1130	1.3
	11-28	1500	2.3
	12-27	1330	1.1
1973	1-23	1130	0.6
	2-27	1330	1.2
	3-28	1330	0.6
	4-25	1400	1.4
	5-21	1800	0.9
	6-18	1620	2.5
	7-16	1200	1.8
	8-13	1500	1.0
	9-11	1600	0.6

	<u>Date</u>	<u>Time</u>	<u>5-day BOD (mg/l)</u>
	Blackfoot River below 7-up Pete Creek		
1972	10-26	1010	1.0
	11-28	1115	2.5
	12-27	1140	1.4
1973	1-23	1615	1.1
	2-27	1010	0.8
	3-28	0935	1.1
	4-25	1100	1.5
	5-21	1300	0.8
	6-18	1200	2.0
	7-16	1115	1.8
	8-13	1145	0.7
	9-11	1230	0.5

APPENDIX F continued

	<u>Date</u>	<u>Time</u>	<u>5-day BOD (mg/l)</u>
	Blackfoot River at Dalton Mountain Road Bridge		
1972	2-2	0945	1.3
	3-23	0945	1.6
	4-14	0935	1.3
	5-19	0945	1.8
	8-2	0935	1.3
	9-8	0945	2.0
	10-26	0900	1.4
	11-28	1015	1.9
	12-27	1040	1.6
A-37 1973	1-23	Invalid - analytical error	
	2-27	0900	1.6
	3-28	0900	0.8
	4-25	1000	0.8
	5-21	1120	1.2
	6-18	1040	2.0
	7-16	1000	1.4
	8-13	1020	0.8
	9-11	1040	0.6

	<u>Date</u>	<u>Time</u>	<u>5-day BOD (mg/l)</u>
	Blackfoot River at Blackfoot Canyon Camp		
1972	2-2	no sample collected	
	3-23	0815	1.2
	4-14	0915	1.5
	5-19	0915	1.6
	8-2	0920	1.0
	9-8	0925	1.9
	10-26	0800	1.4
	11-28	0920	2.4
	12-27	0945	1.3
1973	1-23	1420	1.4
	2-27	0745	0.9
	3-28	0815	0.9
	4-25	0900	1.3
	5-21	0930	1.2
	6-16	0915	2.2
	7-16	0900	1.4
	8-13	0930	0.6
	9-11	0935	0.4

APPENDIX G

Results of Coliform Bacteria Analyses

Collection and analysis of water samples were performed by the USGS, Helena, Montana. Sample preparation and incubation was done in the field using a portable test kit. The membrane filter technique was used to prepare the water sample for incubation. Results are reported in total coliforms per 100 ml and fecal coliforms per 100 ml of water sample.

APPENDIX G continued

	Date	Time	Temp (°C)		Total Coliforms per 100 ml	Fecal Coliforms per 100 ml		Date	Time	Temp (°C)		Total Coliforms per 100 ml	Fecal Coliforms per 100 ml	
			Air	Water						Air	Water			
A-39	Blackfoot River at Hogan Road Bridge (USGS gage below Alice Creek near Lincoln)							Blackfoot River below 7-up Pete Creek						
	1972	10-26	1130	10.5	3.5	1	0	1972	10-26	1010	2.0	3.5	12	0
		11-28	1500	-1.5	0.0	6	0		11-28	1115	-4.5	0.0	10	1
		12-27	1550	5.5	1.5	7	0*		12-27	1140	4.5	3.0	0	0*
	1973	1-23	1130	-5.0	0.0	1	0	1973	1-23	1615	---	1.0	2	0
		2-27	1330	8.0	2.5	0	0		2-27	1010	5.5	3.0	6	4
		3-28	1130	4.0	3.5	0	0		3-28	0935	3.5	0.5	2	0
		4-25	1400	7.5	5.5	0	0		4-25	1100	7.5	5.0	12	0
		5-21	1800	14.0	11.0	28	2*		5-21	1300	11.0	7.0	13	0
		6-18	1620	11.0	9.0	12	0*		6-18	1200	7.5	5.5	10	2
		7-16	1200	25.5	15.0	136	0		7-16	1115	24.0	11.5	28	1
		8-13	1500	30.5	15.5	8	11		8-13	1145	25.5	12.0	75	6
	9-11	1600	25.0	15.5	1700	1		9-11	1230	24.0	11.0	170	2	

*Reported values based on non-ideal count of colonies.

*Reported value based on non-ideal count of colonies.

APPENDIX G continued

	Date	Time	Temp (°C)		Total Coliforms per 100 ml	Fecal Coliforms per 100 ml		Date	Time	Temp (°C)		Total Coliforms per 100 ml	Fecal Coliforms per 100 ml	
			Air	Water						Air	Water			
A-40	Blackfoot River at Dalton Mtn Road Bridge							Blackfoot River at Blackfoot Canyon Camp						
	1972	10-26	0900	0.5	5.5	32	4							
		11-28	1015	-12.0	1.5	49	3	1972	10-26	0800	1.5	5.5	20	0
		12-27	1040	5.5	4.5	23	6*		11-28	0920	-10.0	1.0	1400*	2
									12-27	0945	4.0	4.0	18	2
	1973	1-23	1500	---	5.0	2	0	1973	1-23	1420	---	2.0	2	0
		2-27	0900	8.0	4.0	120	15		2-27	0745	1.5	4.5	14	2
		3-28	0900	2.0	3.5	3	0		3-26	0815	1.0	3.0	0	0
		4-25	1000	8.0	6.0	11	6		4-25	0900	6.5	6.5	24	4
		5-21	1120	10.0	7.0	74	4		5-21	0930	8.5	6.5	260	9
		6-18	1040	6.0	6.5	24	6		6-18	0915	6.5	6.5	50	14
		7-16	1000	19.0	10.5	48	13		7-16	0900	18.0	11.5	43	9
		8-13	1020	20.5	10.5	84	10		8-13	0930	20.5	11.0	61	2
		9-11	1040	20.5	9.5	440	10		9-11	0935	15.0	10.0	930	9

*Reported value based on non-ideal count of colonies.

*Reported value based on non-ideal count of colonies.

APPENDIX H

Results of Field Analyses

Analyses were made with a HACH Model DR-EL water analysis kit. Air and water temperatures were determined with a Taylor pocket thermometer. All values except temperature and pH are given in mg/l. Alkalinity and hardness are reported as mg/l CaCO_3 .

APPENDIX II continued

BLACKFOOT RIVER AT POP'S PLACE

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COLL
2-17-68	1130	0	0	6.9	11	10	55	110	2	2	p.c.	B.D.
3-17-68	0900	1	1	7.3	12	15	65	105	0	2	snow	"
4-21-68	0930	1	-1	7.6	11	15	25	15	0	2	snow	"
5-26-68	1300	6	7	7.7	9	35	30	60	0	2	cloud	"
6-21-68	1030	8	12	8.1	9	40	10	75	0	2	p.c.	"
7-22-68	1210	11	10	8.2	8	60	55	95	0	2	p.c.	"
9-10-68	1015	10	20	8.0	9	60	50	95	0	2	p.c.	"

BLACKFOOT RIVER AT HOGIN CREEK BRIDGE

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COLL
2-17-68	1500	1	1	8.1	12	110	75	115	5	2	cloud	B.D.
3-17-68	1015	2	1	8.6	11	95	65	105	0	2	snow	"
4-21-68	0715	2	0	8.1	10	85	60	95	0	2	snow	"
5-26-68	0215	7	9	8.2	11	60	15	70	5	2	cloud	"
6-21-68	0815	10	11	8.3	11	80	60	80	2	2	p.c.	"
7-22-68	1745	17	18	8.4	9	105	65	105	0	2	p.c.	"
9-12-68	1325	13	25	8.3	9	110	70	105	0	2	clear	"
10-20-68	0830	1	0	7.9	11	110	70	110	0	2	p.c.	"
12-1-68	1130	0	-7	8.0	11	120	75	120	0	2	p.c.	"
1-3-69	1130	0	-8	7.8	10	115	70	110	0	2	p.c.	"
2-9-69	road closed by snow											
3-29-69	1005	0	-2	8.0	11	110	65	110	0	2	p.c.	"
5-1-69	1015	5	6	7.9	10	80	50	80	0	2	cloudy	"
6-17-69	1120	11	20	8.0	10	90	60	90	0	2	clear	"
7-17-69	1110	11	23	7.9	9	90	55	90	0	2	clear	"
8-27-69	1120	11	25	8.0	9	120	70	120	0	2	clear	"
10-8-69	1030	6	7	8.6	9	80	110				p.c.	L.S.
11-29-69	1630	3.5	-0.5	8.6	12		120				clear	L.S.
1-20-70	1700	-0.5	0	8.6	11.0						Snow	"

BLACKFOOT RIVER AT FLESHER PASS ROAD

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COLL
2-17-68	1300	0	3	7.8	11	80	50	95	2	2	p.c.	B.D.
3-13-68	0915	1	1	8.2	11	55	50	90	0	2	snow	"
4-21-68	0815	1	-1	8.1	11	55	50	85	0	2	snow	"
5-26-68	1145	7	8	8.2	10	55	15	70	0	2	cloud	"
6-21-68	0930	10	11	8.4	10	60	50	80	0	2	p.c.	"
7-22-68	1415	16	17	8.4	9	80	60	100	0	2	p.c.	"
9-10-68	1135	10	24	8.3	9	75	60	100	0	2	clear	"
10-20-68	1015	1	0	7.7	10	70	60	95	0	2	cloud	"
12-1-68	1030	0	-7	7.5	11	75	60	100	0	2	p.c.	"
1-3-69	not taken, snow too deep											
2-9-69	not taken, snow too deep											
3-29-69	0910	0	-2	7.4	11	80	50	85	0	2	p.c.	"
5-1-69	0920	5	-6	7.6	9	70	15	75	0	2	cloud	"
6-17-69	1020	10	21	7.9	10	70	50	90	0	2	clear	"
7-17-69	1010	12	22	7.8	8	70	55	85	0	2	clear	"
8-27-69	0925	11	16	7.2	8	85	70	95	0	2	clear	"
10-8-69	0915	1.5	1.5	8.5	10	50	110				p.c.	L.S.
11-29-69	1600	0	0	8.5	12		100				clear	L.S.
1-20-70	1650	-0.5	0.5	8.5	10.0						p.c.	"

APPENDIX 1 continued

BLACKFOOT RIVER AT GALTON MTN ROAD BRIDGE

BLACKFOOT RIVER AT BLACKFOOT CANYON CAMP

DATE	TIME	R ₂ O TEMP	AIR TEMP	PH	D.O.	AIR	Ga HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL	DATE	TIME	R ₂ O TEMP	AIR TEMP	PH	D.O.	AIR	Ga HARD	TOTAL HARD	TURB	CO ₂	WEATHER
2-18-68	1215	4	6	8.3	11	165	60	160	0	2	rain	B.D	2-18-68	1130	2	5	8.4	12	155	60	160	0	2	rain
3-17-68	1130	4	1	8.4	11	165	110	160	0	2	snow	"	3-17-68	1100	5	1	8.4	11	160	110	150	4	2	snow
4-21-68	1015	6	8	8.4	11	160	110	160	0	2	pc	"	4-20-68	1430	7	10	8.5	11	155	110	150	0	2	p.c.
5-25-68	1100	7	13	8.0	9	115	75	120	8	2	p.c.	"	5-25-68	1000	7	10	8.3	9	110	75	110	12	2	p.c.
6-20-68	1900	13	18	8.4	9	115	80	110	8	2	p.c.	"	6-20-68	1815	14	17	8.4	11	120	80	110	9	2	p.c.
7-22-68	2030	13	16	8.4	8	160	105	155	0	2	p.c.	"	7-23-68	0815	9	12	8.4	9	160	110	155	0	2	p.c.
9-9-68	1130	13	21	8.3	8	155	110	160	0	2	clear	"	9-10-68	1520	14	25	8.3	9	165	110	160	0	2	clear
10-19-68	1530	8	9	7.9	9	165	120	170	0	2	p.c.	"	10-19-68	1115	6	8	8.1	11	170	115	160	0	2	p.c.
12-1-68	1215	1	-3	8.0	9	180	115	170	0	2	p.c.	"	12-1-68	1315	2	3	8.1	11	170	115	160	0	2	p.c.
1-2-69	1300	1	0	7.9	10	175	120	160	0	2	cloud	"	1-2-69	1200	0	0	7.9	9	165	115	160	0	2	cloud
2-9-69	0905	2	-2	8.0	10	170	120	160	0	2	snow	"	2-8-69	1605	2	-2	8.1	12	165	120	165	0	2	p.c.
3-28-69	1500	8	9	8.0	10	170	115	155	0	2	p.c.	"	3-28-69	1215	5	8	8.1	12	165	120	155	0	2	p.c.
5-1-69	1655	10	12	8.0	10	130	90	120	0	2	p.c.	"	5-1-69	1520	8	12	8.1	10	125	75	70	2	0	clear
6-16-69	1410	12	23	8.0	9	135	95	130	0	2	clear	"	6-16-69	1355	11	21	8.0	10	110	100	115	2	2	clear
7-17-69	1310	13	26	7.9	9	140	95	130	0	2	p.c.	"	7-16-69	1550	15	26	8.0	9	145	95	135	0	2	p.c.
8-26-69	1410	13	31	8.0	7	170	120	160	0	2	clear	"	8-26-69	1355	11	23	8.1	8	170	115	160	0	2	clear
10-8-69	1130	8	11.5	8.6	8		130	180			cloudy	S	10-7-69	1510	9.5	16	8.6	10		120	160			p.c.
11-29-69	1130	4.5	4	8.6				160			p.c.	L.S.	11-29-69	1300	2.5	2	8.6	12			160			clear
1-20-70	0930	2.5	0	8.7	10.0						O'cast	"	1/20-70	1440	3.0	1.5	8.6	8.5						O'cast

APPENDIX 4 continued

BLACKFOOT RIVER AT RAINCOW BEND

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COLL
2-18-68	1530	0	6	8.5	10	135	80	130	0	2	rain	B.D.
3-16-68	0815	1	1	8.1	11	115	70	115	1	2	p.c.	"
4-20-68	0900	4	4	8.3	11	110	70	105	0	2	p.c.	"
5-25-68	0745	2	8	8.4	9	80	50	80	20	2	p.c.	"
6-20-68	1030	13	18	8.3	10	90	55	85	8	2	p.c.	"
7-22-68	1330	17	26	8.6	9	130	80	120	0	2	clear	"
9-9-68	1200	14	23	8.4	10	145	85	130	0	2	clear	"
10-19-68	0815	1	-1	8.1	10	135	90	130	0	2	clear	"
11-30-68	1100	0	1	8.1	12	135	85	130	0	2	p.c.	"
1-2-69	0930	0	-7	7.9	11	110	85	135	0	2	cloud	"
2-8-69	0930	1	0	7.8	12	130	90	120	0	2	cloud	"
3-28-69	1000	3	8	7.9	12	135	80	110	22	2	p.c.	"
5-3-69	0920	7	7	7.9	11	100	65	95	10	2	clear	"
6-16-69	1130	13	23	8.1	10	110	70	105	2	2	clear	"
7-16-69	0930	16	17	8.1	10	115	70	110	0	2	clear	"
8-26-69	1115	16	22	8.2	7	110	95	135	0	2	clear	"
10-7-69	0920	7	2.5	8.6	10		100	170			clear	4.5
11-29-69	1000	0	1.5	8.65	13			110			clear	4.5
1-20-70	1000	0	3.5	8.6	11.5						p.c.	"

PASS CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COLL
2-17-68	1015	2	0	7.6	9	65	10	65	8	2	clear	B.D.
5-26-68	1130	6	7	8.0	8	15	25	15	0	2	cloud	"
9-10-68	0815	9	15	8.1	7	75	10	75	0	2	p.c.	"
1-3-69	0810	2	-11	7.2	9	75	10	65	0	2	p.c.	"
1-29-69	0750	0	-5	7.3	11	60	10	60	0	2	p.c.	"
6-17-69	0900	8	16	7.2	9	70	10	60	0	2	clear	"
8-27-69	0840	11	7	7.2	7	75	50	75	0	2	clear	"
11-29-69	1530	3.5	3.5	8.4				80			clear	4.5

SANDBAR CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COLL
7-17-69	0955	14	20	6.9	7	10	25	15	27	2	clear	B.D.

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COLL
2-17-68	1100	1	3	8.0	11	115	70	115	7	2	p.c.	B.D.
5-26-68	1045	7	9	8.3	11	90	60	85	4	2	cloud	"
9-10-68	1230	10	24	8.3	9	120	75	115	0	2	clear	"
1-3-69	1015	0	-10	7.8	10	120	75	115	0	2	p.c.	"
3-28-69	1630	4	4	7.7	10	115	65	110	0	2	p.c.	"
6-17-69	1055	11	21	7.9	10	105	70	100	0	2	clear	"
8-27-69	1230	14	26	7.9	10	125	80	130	0	2	clear	"
11-30-69	1000	0.5	2	8.4				120			clear	4.5

HOGIM CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COLL
10-8-69	1000	5	3.5	8.5	9.0		60	100			p.c.	4.5
1-20-70	1630	-0.5	0	8.5	11.0						snow	"

LANDER'S FORK

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COLL
2-17-68	1215	1	0	8.4	10	110	85	110	6	2	snow	B.D.
5-26-68	0815	14	8	8.3	10	130	75	130	5	2	cloud	"
9-10-68	1110	10	25	8.2	10	145	90	135	0	2	clear	"
1-2-69	1150	4	-2	8.1	10	145	90	140	0	2	p.c.	"
3-28-69	1655	6	5	8.0	10	140	90	135	0	2	p.c.	"
6-17-69	1250	9	21	7.9	10	115	80	100	0	2	clear	"
8-27-69	1105	9	19	7.9	8	115	95	115	0	2	clear	"
11-30-69	0910	3	-3.5	8.6				110			clear	4.5

ATPND A ... Inland

ATPNDTX 1 continued

BEAVER CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
2-18-68	1100	0	2	8.1	11	110	130	110	0	2	rain	B.D.
5-25-68	1630	8	8	8.2	10	80	65	80	8	2	rain	"
9- 8-68	1500	13	20	8.5	10	155	120	115	0	2	clear	"
1- 2-69	1100	0	-2	7.8	10	130	100	120	0	2	cloud	"
3-28-69	1120	4	9	7.9	11	130	100	115	0	2	p.c.	"
6-16-69	1515	13	23	8.0	10	130	75	100	0	2	clear	"
8-26-69	1705	16	28	8.0	9	155	115	150	0	2	clear	"
11-30-69	1120	0	0.5	8.6				160			clear	< 5

FOORMAN CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
2-18-68	0715	2	1	8.1	11	120	85	125	0	2	snow	"
5-25-68	1215	6	12	8.1	10	95	75	105	18	2	p.c.	"
9- 8-68	1820	10	21	8.4	9	125	90	125	0	2	p.c.	"
1- 2-69	1700	3	0	8.1	10	130	90	130	0	2	p.c.	"
3-28-69	1715	7	7	8.1	10	120	90	120	0	2	p.c.	"
6-16-69	1715	13	20	8.0	10	120	90	110	0	2	clear	"
8-27-69	1320	13	26	8.1	9	130	100	125	0	2	clear	"
11-29-69	1100	5	7.5	8.7				130			clear	< 5

WILLOW CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
3-16-68	1630	6	9	8.0	12	55	10	55	10	2	p.c.	B.D.
6-20-68	1200	16	17	8.0	8	80	50	75	15	2	p.c.	"
10-12-68	1620	6	9	7.8	9	50	55	75	2	2	p.c.	"
2- 9-69	0815	0	-2	7.2	9	85	50	70	0	2	snow	"
5- 1-69	1605	11	12	7.4	10	45	25	10	15	2	clear	"
7-17-69	1215	10	25	7.8	8	55	35	50	17	2	clear	"
10- 7-69	1610	9	13	8.6	11		60	100			p.c.	< 5
1-20-70	0920	-0.5	0	8.4	10.0						D'cast	"

SPRING CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
2-18-68	0300	3	1	8.1	10	175	110	175	0	2	snow	B.D.
5-25-68	1615	9	11	8.1	11	160	105	170	0	2	cloud	"
9- 9-68	1650	13	20	8.1	9	175	110	170	0	2	p.c.	"
1- 2-69	1520	4	0	8.1	10	180	120	170	0	2	cloud	"
3-28-69	1315	7	9	8.0	10	180	120	175	0	2	p.c.	"
6-16-69	1620	10	20	8.0	10	175	115	165	0	2	clear	"
8-26-69	1510	10	29	8.1	9	180	120	170	0	2	clear	"
11-30-69	1035	3.5	1.5	8.6				180			clear	< 5

KEEP COOL CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
2-18-68	1000	0	1	8.4	11	160	105	150	0	2	snow	B.D.
5-25-68	1630	11	11	8.4	10	115	90	110	10	2	p.c.	"
9- 9-68	1600	13	20	8.4	9	150	100	110	0	2	clear	"
1- 2-69	1500	0	0	7.8	10	155	105	145	0	2	cloudy	"
3-28-69	1150	4	9	7.2	11	115	100	135	0	2	p.c.	"
6-16-69	1605	15	21	7.9	10	110	95	130	0	2	clear	"
8-26-69	1610	15	28	8.0	9	160	105	150	0	2	clear	"
11-30-69	1050	0	1.5	8.6				170			clear	< 5

ARRASTRA CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	W	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
3-17-68	1100	4	2	8.4	11	100	65	95	0	2	rain	B.D.
6-20-68	1710	11	21	8.1	11	60	30	60	0	2	p.c.	"
10- 20-68	1315	6	9	8.2	12	105	60	100	0	2	cloud	"
2-8- 69	1530	2	-2	8.0	11	100	60	100	0	2	p.c.	"
5- 3-69	1010	7	12	8.0	11	90	50	85	0	2	clear	"
7-16-69	1505	15	26	7.9	9	100	55	95	0	2	p.c.	"
10- 7-69	1500	9	19	8.5	10		60	110			p.c.	< 5
1-20-70	1400	3.5	2.0	8.6	11.0						D'cast	"

APPENDIX A (continued)

KLEINSCHMIDT CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
1-12-68	1200	6	2	8.6	10	165	95	150	2	2	rain	B.D.
6-20-68	1815	16	21	8.6	10	150	90	115	0	2	p.c.	"
10-20-68	1115	9	8	8.3	10	155	90	115	0	2	cloud	"
2-8-69	1410	6	-5	8.2	12	115	90	135	0	2	p.c.	"
5-1-69	1405	11	15	8.1	11	115	90	110	0	2	clear	"
7-16-69	1115	16	26	8.1	10	160	95	115	0-2	2	clear	"
10-7-69	1115	13	12	8.6	10		80	150			clear	4.5
1-20-70	1310	6.5	2.0	8.6	10.0						0'cast	"

MONTURE CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
1-16-68	1520	7	9	8.3	12	80	15	75	0	2	cloud	B.D.
6-20-68	1600	10	24	8.1	11	110	20	30	2	2	p.c.	"
10-19-68	1320	6	8	8.0	12	80	15	80	0	2	p.c.	"
2-8-69	1315	0	-5	7.7	11	110	75	115	0	2	p.c.	"
5-1-69	1215	6	18	7.5	11	55	30	15	0	2	p.c.	"
7-16-69	1320	12	22	7.8	9	65	30	55	0	2	clear	"
10-7-69	1300	8	15	8.1	11		50	100				4.5
1-20-70	1130	1.0	1.5	8.6	9.0*						WATN Cloudy	4.5

* Oxygen could not be fixed immediately. Could have been some O₂ loss.

NORTH FORK BLACKFOOT RIVER

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
1-21-68	1230	6	7	8.3	10	115	80	110	0	2	p.c.	B.D.
7-21-68	0915	8	15	8.5	10	110	80	130	0	2	p.c.	"
11-20-68	2030	6	0	7.9	10	110	80	135	0	2	p.c.	"
5-2-69	1330	6	15	8.0	11	120	70	115	0	2	p.c.	"
7-16-69	1226	12	26	7.9	10	120	70	115	0	2	clear	"
10-7-69	1310	10	22	8.5	9.0		90	150			clear	4.5
1-20-70	1315	6.0	1.5	8.6	9.0						0'cast	"

COTTONWOOD CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
1-21-68	1120	6	4	8.1	11	115	100	115	0	2	p.c.	B.D.
7-22-68	1600	11	26	8.5	11	105	100	150	0	2	p.c.	"
11-30-68	1500	0	2	8.1	11	160	100	150	0	2	p.c.	"
5-1-69	1205	6	12	8.0	11	135	80	120	2	2	p.c.	"
7-16-69	1215	12	22	8.0	9	155	95	150	0	2	clear	"

CLEARWATER RIVER

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
1-21-68	1310	6	8	8.2	10	90	65	85	30	2	p.c.	B.D.
7-23-68	1030	10	19	8.4	9	95	60	90	8	2	p.c.	"
11-10-68	1250	0	3	7.8	10	110	65	105	5	2	p.c.	"

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
1-16-68	1100	1	11	8.2	11	75	15	70	0	2	p.c.	B.D.
6-20-68	1200	16	25	8.2	11	60	15	50	0	2	p.c.	"
10-19-68	1300	7	8	7.9	12	80	50	70	0	2	p.c.	"
2-8-69	1215	0	-7	7.6	11	70	15	65	0	2	p.c.	"
5-1-69	1115	7	12	7.9	11	60	35	55	0-2	2	p.c.	"
7-16-69	1120	17	23	7.8	9	65	15	60	0	2	clear	"
10-7-69	1110	10.5	7	8.6	9		50	90			clear	4.5
1-20-70	1130	1	3.5	8.6	10.5						p.c.	4.5

APPENDIX 1, continued

BLANCHARD CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
4-21-68	1220	5	6	8.2	11	10	25	15	0	2	p.c.	B.D.
7-21-68	1210	stream dry										
11-30-68	1130	0	2	7.9	10	80	15	75	0	2	p.c.	"

GOLD CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
4-20-68	1030	3	6	8.1	11	80	50	75	0	2	p.c.	B.D.
7-22-68	1130	11	29	8.5	9	100	65	95	0	2	clear	"
11-30-68	1200	0	2	8.0	11	100	65	95	0	2	p.c.	"

UNION CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
3-16-68	1130	4	15	8.3	12	120	70	120	10	2	p.c.	B.D.
6-20-68	1200	14	22	8.5	9	195	110	180	25	2	p.c.	"
10-19-68	1130	2	3	8.1	9	210	120	200	15	2	clear	"
2-8-69	1030	0	14	7.6	10	195	110	185	5	2	cloud	"
3-28-69	1015	1	8	7.3	8	105	70	100	120	2	p.c.	"
6-17-69	1550	21	28	8.1	9	160	90	145	15	2	clear	"
8-26-69	1210	16	21	8.1	7	180	115	170	20	2	clear	"
11-29-69	1050	0	7.5	8.6				200			p.c.	45

* Flood Stage

EAST TWIN CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
3-16-68	1015	3	10	7.3	13	10	5	10	4	2	p.c.	B.D.
6-20-68	1200	8	19	8.1	11	15	5	10	0	2	p.c.	"
10-19-68	1045	2	3	6.9	11	15	5	15	0	2	clear	"

WEST TWIN CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
3-16-68	1000	3	5	7.1	12	10	5	10	0	2	p.c.	B.D.
6-20-68	1115	8	18	7.7	11	5	2	2	0	2	p.c.	"
10-19-68	1000	2	0	6.6	11	2	2	5	0	2	clear	"

ELK CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
3-16-68	1315	6	15	8.3	11	125	90	125	60	2	p.c.	B.D.
6-20-68	1110	11	22	8.3	10	135	105	130	18	2	p.c.	"
10-19-68	1215	1	7	8.1	10	150	110	140	10	2	clear	"
2-8-69	no sample due to ice											
5-1-69	1030	4	12	7.9	12	95	60	80	28	2	p.c.	"
7-16-69	1015	12	21	7.9	8	130	85	110	24	2	clear	"
10-1-69	1030	5	1	8.7	11	130	130	170			clear	45
1-20-70	1100	Creek frozen solid, no sample taken										

BELMONT CREEK

DATE	TIME	H ₂ O TEMP	AIR TEMP	pH	D.O.	ALK	Ca HARD	TOTAL HARD	TURB	CO ₂	WEATHER	COL
4-20-68	1230	3	11	8.4	11	75	70	90	0	2	p.c.	B.D.
7-21-68	1320	11	24	8.5	10	110	120	145	0	2	clear	"
11-30-68	1300	0	2	8.1	11	145	110	140	0	2	p.c.	"

APPENDIX I

Blackfoot River near Lincoln (Pop's Place), USIS
 Analyses (Elevation 5130 feet aal)

Results of Diurnal Dissolved Oxygen and
 Water Temperature Determinations

Analyses made at approximately four-hour intervals for 24 hours using the aside modification of the Winkler method according to "Standard Methods for the Examination of Water and Wastewater," 13th ed. 1971.

Water samples were collected beneath the water surface using a homemade Kemmerer type water sampler (Spencer, 1974).

Analyses at Pop's Place by the USGS were made with a YSI D.O. Meter checked at beginning and end of period by modified Winkler method.

Saturation percentages calculated from nomographs showing percent saturation as a function of dissolved oxygen (ppm) and water temperature (°F). Mean sea level (msl) elevation of stations rounded to nearest 200 feet for use with nomographs.

1968

Date	Time	Water Temp. (°F)	D.O. (mg/l)	Percent Saturation	pH
9-9	1200	52	8.8	96	8.1
	1400	54	8.8	99	8.3
	1600	55	8.8	101	8.5
	1800	55	8.6	99	8.4
9-10	2000	55	8.2	94	8.5
	2200	54	7.8	88	8.2
	2400	54	8.0	89	8.5
	0200	54	8.0	89	8.4
	0400	52	8.0	88	8.4
	0600	52	8.0	88	8.4
	0800	50	8.0	85	8.6
	1000	52	8.4	93	8.6
	1200	52	8.8	96	8.4
	1200	39	10.4	96	7.3
10-9	1400	39	10.4	96	7.3
	1600	41	10.2	98	7.4
	1800	41	9.8	94	7.3
	2000	41	9.8	94	7.3
	2200	41	10.0	95	7.3
	2400	39	9.8	90	7.3

APPENDIX I continued

(continued)

Date	Time	Water Temp (°F)	D.O. (mg/l)	Percent Saturation	pH	Date	Time	Water Temp (°F)	D.O. (mg/l)	Percent Saturation	pH	Q (cfs)
10-10	0200	39	9.8	90	7.4	7-23	1000	53	8.4	93	7.7	13.6
	0400	39	9.8	90	7.3		1100	54	-	-	-	13.6
	0600	39	9.8	90	7.4		1200	55	8.4	95	7.8	13.6
	0800	39	10.0	93	7.4		1400	57	8.2	96	7.8	13.3
	1000	39	10.0	93	7.6		1600	59	8.0	96	7.9	12.9
	1200	39	10.2	95	7.4		1800	60	8.0	98	7.9	12.9
						8-21	1200	55	7.9	92	7.8	5.3
							1400	58	7.7	92	7.8	5.3
							1600	61	7.5	92	7.8	
							1800	62	7.4	92	7.7	
							2000	61	7.3	89	7.7	
							2200	60	7.3	88	7.5	
							2400	59	7.4	88	7.5	
						8-22	0200	57	7.5	88	7.4	
							0400	56	7.6	88	7.4	
							0600	55	7.7	89	7.4	5.3
							0800	54.5	7.9	89	7.5	
							1000	54.5	7.9	90	7.6	
							1200	56	8.0	93	7.5	

		1969											
Date	Time	Water Temp (°F)	D.O. (mg/l)	Percent Saturation	pH	Q (cfs)							
7-22	1800	59	8.0	96	6.7*	13.6							
	1900	59	8.0	96	7.7	12.9							
	2000	59	8.0	96	7.6	12.9							
	2100	59	-	-	7.7	12.5							
	2200	59	7.9	94	8.0	12.5							
	2400	57	8.0	94	7.6	12.9							
7-23	0130	56	8.0	93	7.8	12.9							
	0230	55	8.0	92	7.7	13.3							
	0400	55	8.0	92	7.6	13.6							
	0600	54	8.2	92	7.6	13.6							
	0800	53	8.2	92	7.6	13.6							

*Probably due to failure to properly calibrate pH meter (USGS field notation).

APPENDIX I continued

Blackfoot River at Top's Place (Elevation 5130 feet msl)
(Fish and Game Analyses)

1971

Date	Time	Water Temp. (°F)	D.O. (mg/l)	Percent Saturation
8-31	2010	57	7.7	91
9-1	0045	54	7.7	88
	0400	54	7.9	91
	0815	52	8.0	89
	1245	53	8.1	90
	1625	56	8.0	93
	2025	54	7.8	89

1973

05-y

8-17	1240	58	7.8	94
	1635	61	7.5	93
	2035	60	7.3	90
8-18	2440	57	7.2	85
	0440	55	7.5	87
	0840	53	7.8	87
	1235	56.5	8.0	93

Blackfoot River at Fleisher Road (Elevation ± 5000 feet msl)
(Fish and Game Analyses)

1971

Date	Time	Water Temp. (°F)	D.O. (mg/l)	Percent Saturation
8-31	1945	57	7.9	92
	2400	54	7.9	89
9-1	0445	51	8.2	89
	0835	50	8.2	88
	1225	52	8.5	94
	1555	58	8.3	98
	1955	54	7.9	90

1973

8-17	1220	59	8.4	101
	1615	67	8.1	106
	2010	61	7.2	98
8-18	2420	58	7.0	84
	0420	55	7.4	85
	0820	53	8.0	89
	1220	57	8.5	99

APPENDIX I continued

Alice Creek (Elevation $\pm$ 5000 feet msl) Fish and Game Analyses1971

<u>Date</u>	<u>Time</u>	<u>Water Temp (°F)</u>	<u>D.O. (mg/l)</u>	<u>Percent Saturation</u>
8-31	2025	54	7.8	89
	0030	54	8.0	91
9-1	0425	50	8.1	87
	0800	50	8.3	89
	1200	55	8.4	96
	1610	57	8.1	94
	2010	53	7.9	88

1972

8-17	1200	58	8.8	105
	1600	64	7.7	98
	2000	57.5	7.6	89
	2400	53	7.7	86
8-18	0400	50.5	8.0	97
	0800	50	8.3	89
	1200	58	8.1	96
	1310	61	8.0	98

Poorman Creek (Elevation $\pm$ 4600 feet msl)
Fish and Game Analyses1971

<u>Date</u>	<u>Time</u>	<u>Water Temp (°F)</u>	<u>D.O. (mg/l)</u>	<u>Percent Saturation</u>
9-1	1140	50	9.3	99
	1525	53	9.3	102
	1900	52	8.6	94
	2320	50	8.4	89
9-2	0230	no sample collected		
	0725	48	8.8	92
	1100	50	9.5	101

1972

<u>Date</u>	<u>Time</u>	<u>Water Temp (°F)</u>	<u>D.O. (mg/l)</u>	<u>Percent Saturation</u>	<u>Discharge (cfs)</u>
6-22	1230	49	10.4	108	35
	1710	54.5	9.6	108	35
	2030	53	9.5	104	32
	2425	49	9.9	104	32
6-23	0425	46	10.1	102	35
	0835	45.5	10.3	102	35
	1220	46	10.6	107	36

A-51

APPENDIX I continued

Blackfoot River at Dalton Mountain Road
(Elevation $\pm$ 4400 feet msl)

Fish and Game analyses

1971

Date	Time	Water Temp (°F)	D.O. (mg/l)	Percent Saturation
9-1	1100	50	8.8	92
	1455	53	8.6	94
	1845	53	8.9	97
	2250	52	8.0	86
9-2	0230	no sample collected		
	0910	48	8.1	84
	1045	49	8.9	93

1972

					Discharge (cfs)
6-22	1210	50	10.3	108	1050
	1615	54.5	9.3	104	1050
	2000	56	8.9	101	960
	2400	52	9.0	97	940
6-23	0400	49	9.3	97	940
	0805	47	9.6	98	960
	1155	47	9.7	99	1025

Blackfoot River at Blackfoot Canyon Camp (Elevation $\pm$ 4400 feet msl)

Fish and Game Analyses

1971

Date	Time	Water Temp (°F)	D.O. (mg/l)	Percent Saturation
9-1	1030	50	8.7	91
	1430	53	8.8	96
	1825	52	8.8	95
	2225	52	8.4	91
9-2	0230	no sample collected		
	0625	50	8.4	88
	1025	50	8.9	93

1972

					Discharge (cfs)
6-22	1130	50	9.8	103	930
	1635	54.5	9.5	107	920
	1935	54.5	9.7	108	910
	2330	55	8.8	98	900
6-23	0330	53	9.1	98	880
	0735	50	9.3	98	880
	1130	49	9.6	100	900

APPENDIX J

Miscellaneous Surface Water Temperatures

Temperatures given herein were obtained periodically during the study. All temperatures are given in degrees centigrade and were obtained with a Taylor pocket thermometer. Conversion of °F to °C was done where necessary.

These temperatures are in addition to temperature data given in Appendix H, Field Analyses.

Temperatures from main Blackfoot River stations are given first in descending order downstream, followed similarly by tributary streams.

APPENDIX J continued

Year	Date	Time	H ₂ O Temp
------	------	------	-----------------------

Blackfoot River at Pop's Place

1972	11-1	--	0.6
1973	5-14	--	6.3
	8-17	1245	14.4

Blackfoot River at Flasher Road Bridge

1971	3-26	1310	0.0
	4-27	1125	1.0
	5-21	1450	4.5
	6-23	1420	15.0
	8-23	1545	17.0
	9-21	1410	5.0
	11-22	1345	0.0
	12-21	1000	0.0
1972	3-23	1400	2.0
	4-20	1200	4.5
	5-22	1210	6.0
	6-20	0915	8.5
	7-19	1230	7.5
	8-31	1300	12.8
	9-26	1145	2.2
	10-31	1200	0.0
	11-1	--	0.0
	11-29	1400	0.0
1973	3-7	1230	0.0
	5-14	1150	8.3
	8-17	1520	18.9

Year	Date	Time	H ₂ O Temp	Year	Date	Time	H ₂ O Temp
------	------	------	-----------------------	------	------	------	-----------------------

Blackfoot River at Boyan Road Bridge

1971	3-26	1230	3.5	1972 (cont'd)	11-3	--	4.4
	4-27	1040	1.5		11-28	1500	0.0
	5-21	1320	5.0		12-27	1550	1.5
	6-23	1445	17.5	1973	1-23	1130	0.0
	8-23	1355	14.0		2-27	1330	2.5
	9-21	1300	7.5		3-28	1130	3.5
	11-22	1235	4.0		4-25	1400	5.5
	12-21	1140	1.0		5-21	1800	11.0
1972	2-2	1330	0.0		6-18	1620	9.0
	3-23	1200	2.5		7-16	1200	15.0
	5-19	1050	5.0		8-13	1500	15.5
	8-2	1020	12.2		8-20	1045	11.7
	9-8	1010	7.2		9-11	1600	15.5
	10-25	1130	3.5				

Blackfoot River below 7-up Pete Creek

1972	10-26	1010	3.5	1973 (cont'd)	4-25	1100	5.0
	11-3	--	3.9		5-21	1300	7.0
	11-28	1115	0.0		6-18	1200	5.5
	12-27	1140	3.0		7-16	1115	11.5
1973	1-23	1615	1.0		8-13	1145	12.0
	2-27	1010	3.0		8-20	1120	9.4
	3-7	1020	0.3		9-11	1230	11.0
	3-28	0935	0.5				

A-55

[illegible]

APPENDIX J continued

<u>Year</u>	<u>Date</u>	<u>Time</u>	<u>H₂O Temp</u>	<u>Year</u>	<u>Date</u>	<u>Time</u>	<u>H₂O Temp</u>	<u>Year</u>	<u>Date</u>	<u>Time</u>	<u>H₂O Temp</u>	<u>Year</u>	<u>Date</u>	<u>Time</u>	<u>H₂O Temp</u>
<u>Alice Creek at Highway 200</u>								<u>Landers Fork at Highway 200</u>							
1971	3-26	1300	3.0	1972 (con't'd)	6-20	1030	9.5	1972	11-3	--	5.0				
	4-27	1105	1.5		7-19	1150	8.5	1973	3-5	1135	4.4				
	5-21	1400	5.0		8-31	1330	14.0		5-19	1135	5.0				
	6-23	1430	16.5		9-26	1040	4.4		8-20	0950	8.3				
	8-23	1620	16.0		10-31	1100	2.8								
	9-21	1335	8.5		11-2	--	3.3								
	11-22	1315	2.5		11-29	1330	1.1	1971	3-26	1200	5.0	1972 (con't'd)	5-22	1000	5.0
	12-21	1041	1.0	1973	3-5	1240	2.8		4-27	1330	4.0		6-20	1245	8.0
1972	3-23	1315	3.5		4-24	--	8.9		5-21	1650	5.5		7-19	0945	7.5
	4-20	1100	5.0		5-16	1400	10.7		6-23	1525	13.0		8-31	1440	12.8
	5-22	1130	5.0		8-17	1115	13.3		8-23	--	10.0		9-26	1000	6.1
									9-21	1515	10.0		10-26	--	6.3
									11-22	1105	4.0		10-31	1020	4.4
									12-21	1235	4.0		11-29	1200	3.9
<u>Hardscrabble Creek</u>								<u>Boorman Creek</u>							
1971	10-20	--	5.5	1973	3-5	1420	1.1	1972	3-23	1100	3.5	1973	3-5	1045	2.2
1972	3-23	1230	2.0		4-24	--	7.2		4-20	0940	3.5		8-20	1215	10.6
	4-20	1030	3.5		5-19	1000	10.6								
	5-22	1100	8.0		8-18	1115	12.2								
	7-19	1100	9.0												
	11-25	1300	0.0												

APPENDIX K

Results of Turbidity Measurements

Depth-integrated samples were collected with a Model DR-48 hand-suspended sediment sampler. Samples were thoroughly mixed and split into three 30 milliliter glass vials, with caps, and transported to the laboratory. Turbidity was determined with a Hach Model 2100-A laboratory turbidimeter within 24 hours of collection. Turbidity in each of the three vials was measured, and the average of the three readings became the final figure used. Turbidity values were rounded off in accordance with "Standard Methods for the Examination of Water and Wastewater," 13 edition, 1971, page 352.

Turbidity measurements made at the beginning of the study in 1968 were made in the field with a Hach Model DR-EL field test kit. These measurements are probably less accurate than those obtained with the Model 2100-A meter.

Streamflow was determined from staff gage height reading at time of sample collection, or from USGS streamflow records (daily mean, CFS). Where no gage was installed, a brief description of flow characteristics is given.

Turbidity of Union Creek near McNamara Landing

Date	Turb (JTU)	Flow
1972 3-13	50	High and muddy

Turbidity of Clearwater River at Highway 200

Date	Turb (JTU)	Flow
1972 4-18	2.3	High (not measured)

Turbidity of Cottonwood Creek on B-C Game Range

Date	Turb (JTU)	Flow
1972 4-10	2.1	High and clear
5-16	14	Flooding, turbid
6-5	4.8	High and flooding, partially clear

Turbidity of North Fork Blackfoot River at Highway 200

Date	Turb (JTU)	Flow
1972 6-3	150	High, fast, muddy
6-9	130	Flooding, muddy
6-20	9.4	High, milky-clear

Turbidity of Arroyo Creek at Highway 200

Date	Turb (JTU)	Flow
1972 5-18	3.5	Clear
6-7	2.7	High and clear

APPENDIX K continued

Turbidity of Blackfoot River at Blackfoot Canyon Camp

Date	Turb (JTU)	Flow (cfs)
1972 2-14	1.4	--
3-17	4.4	--
4-28	8.9	520
5-8	3.9	620
5-17	100	2830
5-18	100	3100
5-24	31	2150
5-31	90	2700
6-3	140	3500
6-5	120	3175
6-7	110	3050
6-9	110	3100
6-13	55	2360
6-16	37	1488
6-20	23	1045
6-27	7.3	790
7-10	3.5	574
1973 2-14	0.65	90 (est.)
5-19	55	930
5-22	19	840
5-31	4.8	565
6-4	2.5	540
6-8	4.9	610
1-16	4.2	± 100 (est.)
5-28	47	1500

Turbidity of Willow Creek at Dalton Min Road

Date	Turb (JTU)	Flow
1972 3-17	8.5	Slightly flooding
5-8	9.2	Not too high, murky
6-3	60	Flooding, culvert washed out
6-5	40	Flooding
6-27	7.1	Not too high, murky

Turbidity of Blackfoot River at Dalton Min Bridge

Date	Turb (JTU)	Flow (cfs)
1972 3-1	.70	--
3-17	1.8	--
4-28	7.1	460
5-8	2.6	580
5-17	100	2475
6-3	170	2950
6-5	150	2800
6-7	120	2625
6-9	130	2625
6-13	60	1950
6-16	35	1550
6-27	6.5	810
7-10	2.5	600
1973 2-14	1.0	80
5-19	40	875
5-22	15	750
5-31	4.0	475
6-4	2.4	455
1974 5-28	45	1400

APPENDIX K continued

Turbidity of Beaver Creek at Highway 200

Date	Turb. (JTU)	Flow
1972 3-1	1.0	Low and clear, ice covered

Turbidity of Spring Creek at Highway 200

Date	Turb. (JTU)	Flow
1972 3-1	.75	Low and clear

Turbidity of Foreman Creek at Stemple Road

Date	Turb. (JTU)	Flow (cfs)
1972 2-14	.85	--
3-1	.80	--
3-15	1.6	--
5-8	2.9	29
5-17	31	217
6-3	24	200
6-7	8.8	130
6-13	9.3	62
6-16	5.0	52
6-20	3.5	41
1973 2-14	.45	--
5-17	1.9	7.8
5-22	2.8	23
5-31	1.0	12.5
6-4	.90	9

Turbidity of Blackfoot River below 7-up Pete Creek

Date	Turb. (JTU)	Flow
1973 2-22	.50	Very low, clear
5-17	34	High, fast, murky

APPENDIX K continued

Turbidity of Lander's Fork of Blackfoot

	<u>Date</u>	<u>Turb (JTU)</u>	<u>Flow</u>
A-60	1972 3-1	.30	Low and clear
	3-17	.55	Low and clear
	5-17	50	$\pm$ 1,100 cfs
	5-24	7.1	Rising, milky gray
	5-31	120	High, muddy brown
	6-3	220	$\pm$ 2,000 cfs
	6-5	150	High, muddy brown
	6-7	130	High, muddy brown
	6-9	120	High, muddy brown
	6-13	50	Falling, light brown
	6-16	26	Falling, clearing up, brownish gray
	6-20	12	Falling, clearing up, milky green
	6-27	4.5	Falling, clear
	7-10	2.0	Low and clear
	11-22	.40	Low and clear
1973	2-22	.65	19 cfs
	5-17	45	Rising, murky
	5-19	40	High, fast, milky brown
	5-22	15	$\pm$ 665 cfs, milky green
	5-24	13	Fairly high, milky green
	5-31	4.4	Not too high, greenish clear
	6-4	2.2	Falling, clear
	6-8	5.5	$\pm$ 2 inches rise, clearish green
	6-11	2.9	Fell $\pm$ 2 inches, colorless green
	1974 5-28	45	Rising, light brown

Turbidity of Copper Creek

<u>Date</u>	<u>Turb (JTU)</u>	<u>Flow</u>
1972 5-24	1.7	High and fast, clear
6-5	6.0	High and fast, gray-green
1973 6-8	2.0	Mod. high, fast, greenish clear

Turbidity of Blackfoot River at Hogan Bridge
(USGS gage below Aline Creek near Lincoln)

<u>Date</u>	<u>Turb (JTU)</u>	<u>Flow (cfs)</u>
1972 3-17	3.7	55
4-28	7.4	
5-17	42	900
5-31	11	509*
6-3	6.4	463*
6-7	3.8	309*
6-9	3.1	279*
6-13	2.4	179*
7-10	1.2	52*
11-22	.80	25*
1973 2-22	.80	11*
5-14	3.3	165*
5-17	19	268*
5-19	4.7	268*
5-22	2.7	193*
5-31	1.2	85*
6-4	1.1	69*
1974 5-28	5.2	382

*Mean flow for the day from U.S.G.S. records.

APPENDIX K continued

Turbidity of Hogum Creek at Hogum Road Bridge

<u>Date</u>	<u>Turb (JTU)</u>	<u>Flow</u>
1972 6-9	5.5	Clear, slightly milky green

Turbidity of Hardscrabble Creek at Highway 200

<u>Date</u>	<u>Turb (JTU)</u>	<u>Flow</u>
1972 4-28	4.3	Flooding
5-17	2.5	Slight flooding, clear
11-22	.7	Low and clear, ice-covered
1973 3-5	.7	Low and clear, ice-covered
4-24	1.6	Rising, clear
5-17	1.0	Low and clear
5-19	1.0	Low and clear

P
5Turbidity of Alise Creek at Highway 200

<u>Date</u>	<u>Turb (JTU)</u>	<u>Flow (cfs)</u>
1972 3-1	.55	--
3-17	1.6	--
4-10	1.9	48
4-28	7.7	102
5-17	31	450
5-19	13	240
5-24	27	190
6-3	4.9	215
6-9	3.6	95
6-13	2.6	71
11-22	1.1	8
1973 2-22	.70	5.7
4-24	1.4	11
5-14	3.1	50
5-16	14	78
5-17	15	94
5-19	4.5	105
5-22	2.6	35
5-31	1.3	28
6-4	.90	27

APPENDIX K continued

Turbidity of Willow Creek (Fletcher) at confluence
with Blackfoot River

<u>Date</u>	<u>Turb (JTU)</u>	<u>Flow</u>
1972 6-3	2.8	High, colorless
11-22	4.2	Low, clear, oily appearance
1973 4-24	7.5	Rising, grayish green
5-14	2.6	High, fast, oily blue, clear

Turbidity of Blackfoot River at Pop's Place

<u>Date</u>	<u>Turb (JTU)</u>	<u>Flow</u>
1972 3-15	8.3	Rising, orange color, detritus
1973 5-14	1.9	High, clear with oily color

Turbidity of Blackfoot River at Fletcher Road

<u>Date</u>	<u>Turb (JTU)</u>	<u>Flow (cfs)</u>
1972 3-1	1.3	--
4-10	1.1	51
5-5	1.5	112
5-17	4.6	337 +
5-31	.25	149
6-3	1.3	125
6-9	.90	80
6-13	1.1	46
11-22	1.2	-- (gage frozen in ice)
1973 2-22	1.0	-- (gage frozen in ice)
4-24	5.2	42
5-14	1.1	66
5-16	2.6	95
5-19	1.6	100
5-22	1.2	66
5-31	1.1	34
6-4	.95	29

APPENDIX L

Streamflow Record

Records are based on periodic staff gage readings and flow measurements. Measurements were made with a Price model AA or a pygmy current meter according to USGS recommended techniques. Gage height was plotted against flow to form a stage-discharge curve from which the flow from other gage height readings was obtained. Annual high flows were measured by USGS personnel.

The staff gages were 4 inches wide, in sections 3.33 feet long, and were marked at 1.0 foot, 0.10 foot, and 0.02 foot intervals.

Staff gage height was recorded at each station whenever water quality, turbidity or insect samples were collected, and at other times the station was visited. Staff gages were read as frequently as possible during spring runoff periods.

An asterick (*) indicates a flow measurement was made that day.

APPENDIX L continued

Streamflow based on periodic staff gage readings and flow measurements

Blackfoot River at Plesher							
	Date	Gage Ht. (ft.)	Flow (CFS)		Date	Gage Ht. (ft.)	Flow (CFS)
1972	3-24	1.47	88*	1973	4-24	.97	42*
	4-10	1.17	51*		5-1	.91	38
	4-13	1.10	50		5-14	1.26	66
	4-17	1.00	43		5-16	±1.54	95
	4-20	.96	40		5-17	±1.62	104
	4-26	1.36	75		5-19	±1.58	100
	5-5	±1.70	112		5-22	1.26	66
	5-17	2.40	337*		5-23	1.16	56
	5-19	±2.25	200		5-31	.86	34
	5-22	±2.10	170		6-4	.79	29*
	5-31	±1.96	149		7-23	.29	5
	6-3	±1.78	125		7-26	.27	4*
	6-9	1.40	80		8-7	.24	4
	6-13	1.12	46		8-17	.16	3
	6-16	.95	44*		9-11	.31	6
	6-19	.90	37		9-25	.28	5
	6-20	.87	35				
	7-10	.50	13*				
	7-19	.74	26				
	8-9	.34	7*				
	8-25	.36	8				
	8-31	.28	5				
	9-13	.34	7				
	9-26	.30	6				
	10-18	.28	4*				

Streamflow based on periodic staff gage readings and flow measurements

Alice Creek at Highway 200								
	Date	Gage Ht. (ft.)	Flow (CFS)		Date	Gage Ht. (ft.)	Flow (CFS)	
1972	3-24	1.12	23*		2-22	.54	5.7*	
	3-30	1.11	23		4-24	.78	11	
	4-10	1.47	48*		4-27	.84	13	
	4-13	1.40	40		5-1	.79	12	
	4-14	1.35	37		5-11	1.50	50	
	4-17	1.29	33		5-14	1.50	50	
	4-20	1.25	31		5-16	± 1.80	78	
	4-26	1.63	60		5-17	± 1.90	94	
	4-28	1.93	102*		5-19	± 1.95	105	
	5-4	1.94	102		5-21	± 1.72	68	
	5-16	± 2.64	430*		5-22	± 1.26	35	
	5-17	2.70	450		5-23	± 1.55	54	
	5-19	± 2.55	310		5-31	1.17	29	
	5-22	± 2.50	280		6-4	1.08	27*	
	5-24	± 2.30	190		7-23	.66	8	
	5-31	± 2.36	215		7-26	.65	8*	
	6-3	± 2.26	177		8-7	.60	7	
	6-9	1.90	95		8-14	.61	7	
	6-13	1.68	71		9-25	.58	6.5	
	6-16	1.50	50		9-26	.57	6	
	7-10	1.04	21					
	8-9	.85	15*					
	8-25	.83	13					
	9-13	.80	12					
	10-18	.73	9.7*					
	11-22	.65	8					
	1973	2-1	.61	7				

APPENDIX L continued

Streamflow in Landers Fork and Copper Creek based only on flow measurements^{1/}

Streamflow based on periodic staff gage readings and flow measurements

Poorman Creek			(1) Landers Fork above Copper Creek			(2) Copper Creek		(3) Landers Fork at Highway 200	
Date	Gage Ht. (ft.)	Flow (CFS)	Date	Gage Ht. (ft.)	Flow (CFS)	Date	Flow (CFS)	Date	Flow (CFS)
1972 3-22	.92	21*	9-26	.78	5			4-19-72	40
3-23	.98	23	10-19	.76	5*			5-8-72	160
4-5	.82	11*	10-31	.74	5			5-16-72	1,100
4-10	.94	19	1973 2-27	.52	1 gage reset-new rating begins			6-2-72	2,000
4-13	.90	16	3-5	.54	1			6-10-72	1,990
4-17	.83	10	4-26	.62	2.8			8-9-72	136
4-20	.83	10	5-17	.78	7.8			10-18-72	26
4-25	.87	14	5-19	.89	13.5			2-22-73	19
5-5	±1.10	32	5-22	.99	23	5-21-73	388	5-21-73	565*
5-8	1.06	29	5-23	.97	19	6-18-73	131	6-18-73	238*
5-16	1.88	171*	5-31	.87	12.5	7-25-73	15	7-25-73	73**
5-17	±2.05	217	6-4	.81	9*	9-27-73	4	9-27-73	36**
5-22	±1.70	117	7-23	.54	2				
6-3	1.92	169*	7-24	.54	2*				
6-3	±2.00	200	8-7	.50	1-2 (poor curve)				
6-7	±1.75	130	8-14	.49	1-2 (poor curve)				
6-13	±1.42	62	8-20	.47	1				
6-16	1.35	52*	9-25	.46	1				
6-20	1.22	41	9-27	.46	1				
7-10	1.00	18							
7-19	1.00	18							
8-7	.90	12.5							
8-10	.88	12*							
8-31	.82	8.6							
9-13	.81	8.4							

*Flow not measured at Highway 200
This is probably the approximate flow at Highway 200 (Copper + Landers) during this high water period because water table is saturated.

**Copper (2) plus Landers (1) does not equal flow of Landers at Highway 200 (3) due to ground water secession. Landers (1) is dry below measuring site above Copper Creek and Copper Creek (2) contributes most of flow. Ground water secession at Highway 200 (3) was 32% on 7-25-73 and 45% on 9-27-73.

^{1/} Extreme channel instability in Landers Fork precluded normal use of staff gage or other reference points.

APPENDIX I continued

Streamflow based on periodic staff gage readings and flow measurements

Blackfoot River at Dalton Mtn Bridge

	Date	Gage Ht.(ft)	Flow (CFS)		Date	Gage Ht.(ft)	Flow (CFS)
1972	3-22	1.02	152*	1973	8-10	±1.25	221*
	3-23	1.18	190		8-25	1.18	190
	3-29	1.16	185		8-31	±1.10	170
	3-30	1.13	178		9-8	.96	133
	4-5	1.17	185		9-13	±.98	140
	4-10	±1.45	285		10-19	.95	125*
	4-13	1.42	275		10-31	.89	118
	4-14	1.40	265		11-29	.81	100
	4-17	1.39	260		2-1	±.69	77
	4-19	1.36	252*		2-14	.70	80
	4-20	±1.35	245		2-15	.68	72*
	4-26	±1.56	330		3-5	.67	70
	4-28	±1.78	460		4-26	±.67	70
	5-4	±1.88	520		5-3	±.66	70
	5-8	±1.98	580		5-11	±1.00	142
	5-16	3.90	2350*		5-17	±1.86	505
	5-17	±4.20	2475		5-19	±2.42	875
	5-19	±4.15	2425		5-21	±2.35	850
	5-22	±4.10	2385		5-21	±2.46	889*
6-2	4.80	3330*	5-22	2.26	750		
6-3	±4.65	2950	5-23	±2.10	670		
6-5	±4.50	2800	5-31	±1.80	475		
6-7	±4.35	2625	6-4	1.76	455		
6-9	±4.35	2625	6-8	±1.90	540		
6-13	±3.66	1950	6-18	1.68	432		
6-16	±3.20	1550	7-23	±.94	126		
6-20	±2.86	1240	7-24	.94	126*		
6-27	2.38	810	8-7	±.89	118		
7-10	±2.00	600	8-14	.86	112		
7-19	±1.86	505	8-20	.82	103		
8-2	±1.46	310	9-25	.78	95		
			9-28	.79	81*		

Streamflow based on periodic staff gage readings and flow measurements

Blackfoot River at Blackfoot Canyon Camp

	Date	Gage Ht.(ft.)	Flow (CFS)	Date	Gage Ht.(ft.)	Flow (CFS)	Date	Gage Ht.(ft.)	Flow (CFS)	
1972	3-22	1.53	244*	7-19	2.48	529	5-23	2.66	760	
	3-29	1.59	265	7-27	2.06	400	5-24	2.60	740	
	3-30	1.55	250	7-31	1.92	357	5-31	2.15	565	
	4-5	1.63	267*	8-2	1.92	357	6-4	2.10	540	
	4-10	2.01	385	8-7	1.77	315	6-8	2.28	610	
	4-13	1.96	370	8-10	1.70	287*	6-18	1.99	471*	
	4-14	1.90	350	8-25	1.60	265	7-23	1.03	156	
	4-17	1.93	362	8-31	1.48	234	7-24	1.03	156*	
	4-20	1.81	327	9-8	1.35	197	8-7	.96	143	
	4-26	2.12	415	9-13	1.37	203	8-14	.90	133	
	4-28	2.44	520	9-26	1.33	193	8-20	.85	132	
	5-4	2.59	561	10-19	1.30	172	9-14	.87	127	
	5-8	2.78	620	10-31	1.25	171	9-25	.83	118	
	5-17	±5.80	2830*	11-29	1.14	142	9-28	.82	120*	
	5-18	±6.10	3100	1973	2-1	.94	102			
	5-19	±5.90	2850	2-14	.88	90				
	5-22	±5.70	2700	2-15	.90	99*				
	5-24	±5.30	2150	3-5	.94	135				
	5-31	±5.70	2700	4-20	.94	102				
	6-2	6.50	3670*	4-26	.96	106				
	6-3	±6.40	3500	5-3	.93	101				
	6-5	±6.15	3175	5-7	.96	106				
	6-7	±6.05	3050	5-11	1.28	180				
	6-9	±6.10	3100	5-14	1.22	220				
	6-13	±5.46	2360	5-16	1.50	310				
	6-16	4.58	1488	5-17	2.15	560				
	6-20	4.08	1045	5-19	±3.10	930				
	6-27	3.30	790	5-21	3.00	895				
	7-10	2.63	574	5-21	3.16	1000*				
	7-18	2.30	462	5-22	2.87	840				

APPENDIX M

Laboratory analyses made by the Anaconda Company on acid mine seepage in the Heddleston Mining District in 1972 and 1973.

Cation - anion balances do not agree in these analyses. The computer program which analyzed the laboratory data did not take into account milliequivalents for H^+ from extremely low pH or high concentrations of heavy metals in the cation summation. However, individual values of the parameters given are correct, according to the Anaconda Company.

Appendix H (continued)
THE ANACONDA COMPANY
 HEADQUARTERS - GREAT SALT LAKE

DATE: 6/24/72 PROJECT: HEADQUARTERS MINE - SEEPAGE PLANT
 REMARKS: BY: DENT FILE

FIG. #1

SCHEMATIC OF TUNNEL LOCATIONS

ENVIRONMENTAL ENGINEERING DEPARTMENT
 THE ANACONDA COMPANY BUTTE, MONTANA 59701
 WATER QUALITY ANALYSIS

LOCATION AND STATE: MONTANA
 PROJECT: LEWIS & CLARK
 LOCATION: HEADQUARTERS MINE
 DATE SAMPLED: 06-24-72
 TIME SAMPLED: 1145
 ANALYST: GPK (E)
 FIELD MEASUREMENTS: 5, GPK (E)
 PROPERTY OR OPERATION: HEADQUARTERS
 LABORATORY NUMBER: 72-1149
 SAMPLE OR BOTTLE NUMBER: 1695
 STATISTICAL NUMBER: 1695
 SAMPLE TYPE: DRAINAGE
 SAMPLE HANDLING: 4100
 PRINCIPAL USE OF WATER: RECREATION

PROJECT: HEADQUARTERS MINE
 REMARKS: PAYMASTER TUNNEL DRAINAGE IRON PPT & DENSE ALGAE MAT

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED					
	MG/L	MEQ/L		MG/L	MEQ/L
CALCIUM (CA)	49.	2.409	BICARBONATE (HCO3)	3.	0.1
MAGNESIUM (MG)	13.3	1.097	CARBONATE (CO3)	0.	0.0
SODIUM (NA)	8.4	0.365	HYDROXIDE (OH)	0.	0.000
POTASSIUM (K)	1.20	0.031	CHLORIDE (CL)	11.2	0.310
TOT. IRON (FE)	120.	6.446	SULFATE (SO4)	676.	14.494
MANGANESE (MN)	4.56	0.166	NITRATE (NO3)	.00	0.0
ALUMINUM (AL)	19.8	1.797	FLUORIDE (F)	.87	0.046
SILICA (SiO2)			PHOSPHATE (PO4)		
TOTAL CATIONS		12.262	TOTAL ANIONS		14.353
STANDARD DEVIATION OF CATION-ANION BALANCE		0.12 SIGMA			
LABORATORY PH		2.62	CARBONATE HARDNESS AS CaCO3		0
FIELD TEMPERATURE DEGREES F		44.	NON-CARB. HARDNESS AS CaCO3		174
CALCULATED DISSOLVED SOLIDS		919.1	TOTAL HARDNESS AS CaCO3		174
EVAPORATED SOLIDS AT 105 C		1104.	TOTAL ALKALINITY AS CaCO3		0
SPECIFIC CONDUCTANCE IN MICROMHOS/CM AT 25 C		1690.0	FIELD TURBIDITY (JTU)		
ACIDITY AS CaCO3			TOTAL FILTERABLE SOLIDS		
			DISSOLVED OXYGEN AS O2		
SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)					
ARSENIC (AS) .021		COPPER (CU) 21.0		NICKEL (NI) .04	
CADMIUM (CD) .048		LEAD (PB) .10		ZINC (ZN) 3.40	
CHROMIUM (CR) <.02		MERCURY (HG)			

ADDITIONAL PARAMETERS
 AA CALCIUM (CA) 40.
 AA MAGNESIUM (MG) 16.50

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS LABELED TO-TOTAL RECOVERABLE
 PERCENT REACTANCE VALUES
 ANALYSIS REQUESTED BY GPK
 DATE ANALYSIS COMPLETED 8-14-72
 ANALYST MAC
 NOTE: IF CATION-ANION BALANCE RELATED TO THIS ANALYSIS REFER TO NUMBER 72-1149

Appendix M (continued)

ENVIRONMENTAL ENGINEERING DEPARTMENT
THE ANACONDA COMPANY BUTTE, MONTANA 59701

WATER QUALITY ANALYSIS
NATION OR STATE MONTANA
PROVINCE OR COUNTY LEWIS & CLARK
LOCATION
DATE SAMPLED 09-04-72
TIME SAMPLED 1405
WATER FLOW RATE 5. GPM (F)
FIELD MEASUREMENTS
PROPERTY OR OPERATION HEDDLESTON
LABORATORY NUMBER 72-1149
SAMPLE OR BOTTLE NUMBER 597
STATION NUMBER
SAMPLE TYPE DRAINAGE
SAMPLE HANDLING 4103
PRINCIPAL USE OF WATER

PROJECT HEDDLESTON NO. 2 TUNNEL
REMARKS ALGAE SPARSE NO IRON PPT

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED
CALCIUM (CA) 394 3.972 BICARBONATE (HCO3) 0.0 MEQ/L
MAGNESIUM (MG) 39.0 3.167 CARBONATE (CO3) 0.0 MEQ/L
SODIUM (NA) 3.6 0.244 HYDROXIDE (OH) 0.0 MEQ/L
POTASSIUM (K) 1.14 0.029 CHLORIDE (CL) 1.4 0.237
TOTAL IRON (FE) 20 1.504 SULFATE (SO4) 460.0 9.696
MANGANESE (MN) 12.6 0.459 NITRATE (NO3) 0.0 0.0
ALUMINUM (AL) 4.9 0.545 FLUORIDE (F) 0.01 0.032
SILICA (SI02) PHOSPHATE (PO4)

TOTAL CATIONS 9.315 TOTAL ANIONS 9.768
STANDARD DEVIATION OF CATION-ANION BALANCE 1.476 SIGMA
LABORATORY PH 2.96 CARBONATE HARDNESS AS CaCO3 0
FIELD TEMPERATURE DEGREES F 52.0 NON-CARB. HARDNESS AS CaCO3 327
CALCULATED DISSOLVED SOLIDS 626.0 TOTAL HARDNESS AS CaCO3 327
EVAPORATED SOLIDS AT 105 C 740.0 TOTAL ALKALINITY AS CaCO3 0
SPECIFIC CONDUCTANCE IN MICROHMS/CM AT 25 C 1140.0 FIELD TURBIDITY (JTU)
ACIDITY AS CaCO3 DISSOLVED OXYGEN AS (O2)
SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
ARSENIC (AS) 0.005 COPPER (CU) 0.09 NICKEL (NI) 0.09
CADMIUM (CD) 0.020 LEAD (PB) 0.08 ZINC (ZN) 6.20
CHROMIUM (CR) <0.02 MERCURY (HG)

ADDITIONAL PARAMETERS
AA CALCIUM (CA) 58.0 AA MAGNESIUM (MG) 26.20

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS LABELED TR-TOTAL RECOVERABLE
SAMPLED BY RLD PERCENT REACTANCE VALUES
ANALYSIS REQUESTED BY GMM CA MG NA K CL SC4 HCO3 CO3 NO3
DATE ANALYSIS COMPLETED 8-14-72 50 44 5 0 0 100 0 0 0
ANALYST HMC OSC MEAS. OSC CALC.
NOTE: IN CORRESPONDENCE RELATED TO THIS ANALYSIS REFER TO NUMBER 72-1149

ENVIRONMENTAL ENGINEERING DEPARTMENT
THE ANACONDA COMPANY BUTTE, MONTANA 59701

WATER QUALITY ANALYSIS
NATION OR STATE MONTANA
PROVINCE OR COUNTY LEWIS & CLARK
LOCATION
DATE SAMPLED 09-04-72
TIME SAMPLED 1225
WATER FLOW RATE 7. GPM (F)
FIELD MEASUREMENTS
PROPERTY OR OPERATION HEDDLESTON
LABORATORY NUMBER 72-1145
SAMPLE OR BOTTLE NUMBER 227
STATION NUMBER
SAMPLE TYPE DRAINAGE
SAMPLE HANDLING 4103
PRINCIPAL USE OF WATER

PROJECT HEDDLESTON NO. 2 TUNNEL
REMARKS ALGAE SPARSE SOME IRON PPT

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED
CALCIUM (CA) 25.0 1.240 BICARBONATE (HCO3) 0.0 MEQ/L
MAGNESIUM (MG) 2.2 0.179 CARBONATE (CO3) 0.0 MEQ/L
SODIUM (NA) 14.2 0.618 HYDROXIDE (OH) 0.0 MEQ/L
POTASSIUM (K) 1.28 0.033 CHLORIDE (CL) 2.2 0.061
TOTAL IRON (FE) 13.9 6.720 SULFATE (SO4) 189.0 3.932
MANGANESE (MN) 0.78 0.028 NITRATE (NO3) 0.0 0.0
ALUMINUM (AL) 4.7 0.523 FLUORIDE (F) .51 0.027
SILICA (SI02) PHOSPHATE (PO4)

TOTAL CATIONS 3.340 TOTAL ANIONS 4.019
STANDARD DEVIATION OF CATION-ANION BALANCE 4.15 SIGMA
LABORATORY PH 3.12 CARBONATE HARDNESS AS CaCO3 0
FIELD TEMPERATURE DEGREES F 53.0 NON-CARB. HARDNESS AS CaCO3 70
CALCULATED DISSOLVED SOLIDS 292.9 TOTAL HARDNESS AS CaCO3 70
EVAPORATED SOLIDS AT 105 C 394.0 TOTAL ALKALINITY AS CaCO3 0
SPECIFIC CONDUCTANCE IN MICROHMS/CM AT 25 C 610.0 FIELD TURBIDITY (JTU)
ACIDITY AS CaCO3 DISSOLVED OXYGEN AS (O2)
SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
ARSENIC (AS) 0.005 COPPER (CU) 5.08 NICKEL (NI) 0.09
CADMIUM (CD) 0.012 LEAD (PB) .12 ZINC (ZN) .58
CHROMIUM (CR) <0.02 MERCURY (HG)

ADDITIONAL PARAMETERS
AA CALCIUM (CA) 14.2 AA MAGNESIUM (MG) 11.22

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS LABELED TR-TOTAL RECOVERABLE
SAMPLED BY RLD PERCENT REACTANCE VALUES
ANALYSIS REQUESTED BY GMM CA MG NA K CL SC4 HCO3 CO3 NO3
DATE ANALYSIS COMPLETED 8-14-72 60 9 30 2 2 98 0 0 0
ANALYST HMC OSC MEAS. OSC CALC.
NOTE: IN CORRESPONDENCE RELATED TO THIS ANALYSIS REFER TO NUMBER 72-1145

Appendix M (continued)

THE ANACODIA COMPANY
 ENVIRONMENTAL ENGINEERING DEPARTMENT
 BUTTE, MONTANA 59701
 WATER QUALITY ANALYSIS

NATION OR STATE MONTANA
 PROPERTY OR OPERATION HEADLESTON
 PROV. OR COUNTY LEAVELAND
 LOCATION
 DATE SAMPLED 08-04-72
 TIME SAMPLED 1300
 DATE ANALYSIS 08-14-72
 ANALYST MAC
 FIELD MEAS METHOD 30. GPR (F)
 SAMPLE NUMBER 1300
 STATION NUMBER
 SAMPLE TYPE
 SAMPLE HANDLING
 DRAINAGE
 4100
 PRINCIPAL USE OF WATER RECREATION

PROJECT HEADLESTON RD
 REMARKS TUNNEL NO 3 SPACE ALGAE LIGHT IRON PRECIPITATE

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED		MG/L		MG/L		MG/L	
CALCIUM (CA)		16.4	0.240	BICARBONATE (HCO3)		0.	0.0
MAGNESIUM (MG)		7.4	0.135	CARBONATE (CO3)		0.	0.0
SODIUM (NA)		3.2	0.138	HYDROXIDE (OH)		0.	0.009
POTASSIUM (K)		1.30	0.033	CHLORIDE (CL)		3.2	0.089
TOT. IRON (FF)		17.2	0.424	SULFATE (SO4)		211.	4.885
MANGANESE (MN)		0.1	0.011	NITRATE (NO3)		0.00	0.0
ALUMINUM (AL)		7.7	0.155	FLUORIDE (F)		0.33	0.017
SILICA (SiO2)				PHOSPHATE (PO4)			
TOTAL CATIONS		3.445		TOTAL ANIONS		4.452	
STANDARD DEVIATION OF CATION-ANION BALANCE		8.24 SIGMA					
LABORATORY PH		3.02		CARBONATE HARDNESS AS CaCO3		0	
FIELD TEMPERATURE DEGREES F		40.1		NON-CAR. HARDNESS AS CaCO3		74	
CALCULATED DISSOLVED SOLIDS		200.4		TOTAL HARDNESS AS CaCO3		74	
EVAPORATED SOLIDS AT 105 C		344.		TOTAL ALKALINITY AS CaCO3		10	
SPECIFIC CONDUCTANCE IN MICROHMS/CM AT 25 C		400.0		FIELD TURBIDITY (JTU)			
ACIDITY AS CaCO3				TOTAL FILTERABLE SOLIDS			
				DISSOLVED OXYGEN AS 100%			

SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
 ARSENIC (AS) .013 COPPER (CU) 30.4 NICKEL (NI) .06
 CADMIUM (CD) <.010 LEAD (PB) <.02 ZINC (ZN) .18
 CHROMIUM (CR) <.02 MERCURY (HG)

ADDITIONAL PARAMETERS
 AA CALCIUM (CA) 2.3
 PH/FIELD 4.4
 AA MAGNESIUM (MG) 4.75

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS LABELED TR-TOTAL RECOVERABLE
 SAMPLED BY RLD
 ANALYSIS REQUESTED BY GWH
 DATE ANALYSIS COMPLETED 8-14-72
 ANALYST MAC
 DSC MEAS.
 DSC CALC.
 NOTE: IN CORRESPONDENCE RELATED TO THIS ANALYSIS REFER TO NUMBER 72-1148

ENVIRONMENTAL ENGINEERING DEPARTMENT
 THE ANACODIA COMPANY
 BUTTE, MONTANA 59701
 WATER QUALITY ANALYSIS

NATION OR STATE MONTANA
 PROPERTY OR OPERATION HEADLESTON
 PROV. OR COUNTY LEAVELAND
 LOCATION
 DATE SAMPLED 08-04-72
 TIME SAMPLED 1300
 DATE ANALYSIS 08-14-72
 ANALYST MAC
 FIELD MEAS METHOD 300. GPR (F)
 SAMPLE NUMBER 1300
 STATION NUMBER
 SAMPLE TYPE
 SAMPLE HANDLING
 DRAINAGE
 4100
 PRINCIPAL USE OF WATER

PROJECT HEADLESTON RD
 REMARKS MIKE HARVE MINE TUNNEL NO 3 ALGAE-IRON PPT - TURBID

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED		MG/L		MG/L		MG/L	
CALCIUM (CA)		150.	7.480	BICARBONATE (HCO3)		61.	1.500
MAGNESIUM (MG)		159.	13.072	CARBONATE (CO3)		0.	0.0
SODIUM (NA)		3.8	0.165	HYDROXIDE (OH)		0.	0.020
POTASSIUM (K)		1.20	0.031	CHLORIDE (CL)		1.6	0.045
TOT. IRON (FF)		20.	1.074	SULFATE (SO4)		928.	19.315
MANGANESE (MN)		26.6	0.963	NITRATE (NO3)		1.6	0.003
ALUMINUM (AL)		2	0.022	FLUORIDE (F)		.99	0.052
SILICA (SiO2)				PHOSPHATE (PO4)			
TOTAL CATIONS		22.743		TOTAL ANIONS		20.420	
STANDARD DEVIATION OF CATION-ANION BALANCE		-5.27 SIGMA					
LABORATORY PH		6.33		CARBONATE HARDNESS AS CaCO3		50	
FIELD TEMPERATURE DEGREES F		42.		NON-CAR. HARDNESS AS CaCO3		991	
CALCULATED DISSOLVED SOLIDS		1351.6		TOTAL HARDNESS AS CaCO3		1031	
EVAPORATED SOLIDS AT 105 C		1454.		TOTAL ALKALINITY AS CaCO3		50	
SPECIFIC CONDUCTANCE IN MICROHMS/CM AT 25 C		1460.0		FIELD TURBIDITY (JTU)			
ACIDITY AS CaCO3				TOTAL FILTERABLE SOLIDS			
				DISSOLVED OXYGEN AS 100%			

SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
 ARSENIC (AS) .004 COPPER (CU) 1.45 NICKEL (NI) .10
 CADMIUM (CD) .094 LEAD (PB) 1.22 ZINC (ZN) 52.
 CHROMIUM (CR) <.02 MERCURY (HG)

ADDITIONAL PARAMETERS
 AA CALCIUM (CA) 150.
 PH/FIELD 5.
 AA MAGNESIUM (MG) 134.00

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS LABELED TR-TOTAL RECOVERABLE
 SAMPLED BY RLD
 ANALYSIS REQUESTED BY GWH
 DATE ANALYSIS COMPLETED 8-14-72
 ANALYST MAC
 DSC MEAS.
 DSC CALC.
 NOTE: IN CORRESPONDENCE RELATED TO THIS ANALYSIS REFER TO NUMBER 72-1148

Appendix N (continued)

ENVIRONMENTAL ENGINEERING DEPARTMENT
THE ANACONDA COMPANY BUTTE, MONTANA 59711
WATER QUALITY ANALYSIS

NATION OR STATE MONTANA
PROV. OR COUNTY LEWIS & CLARK
LOCATION
DATE SAMPLED 08-04-72
TIME SAMPLED 1700
WATER FLOW RATE 10.0 GPM (G)
FLOW METER METHOD
PROPERTY OR OPERATION HEADLUSTON
LABORATORY NUMBER 72-1146
SAMPLE OR BOTTLE NUMBER 204
STATION NUMBER
SAMPLE TYPE DRAINAGE
SAMPLE NUMBER 4109
PRINCIPAL USE CH WATERS

PROJECT HEADLUSTON
REMARKS MINE DRAINAGE NEAR POPS PLACE

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED

	MG/L	MG/L	MG/L	MG/L
CALCIUM (CA)	18.4	0.720	DICARBONATE (HCO ₃)	25.0
MAGNESIUM (MG)	7.3	0.599	CARBONATE (CO ₃)	0.0
SODIUM (NA)	6.4	0.270	HYDROXIDE (OH)	0.0
POTASSIUM (K)	0.60	0.015	CHLORIDE (CL)	2.0
TOT. IRON (FE)	15.6	0.838	SULFATE (SO ₄)	57.0
MANGANESE (MN)	1.6	0.058	NITRATE (NO ₃)	12.0
ALUMINUM (AL)	1.4	0.159	FLUORIDE (F)	0.015
SILICA (SIO ₂)			PHOSPHATE (PO ₄)	

TOTAL CATIONS 2.864
STANDARD DEVIATION OF CATION-ANION BALANCE -6.93 SIGMA

LABORATORY PH 6.19
FIELD TEMPERATURE DEGREES F 58.0
CALCULATED DISSOLVED SOLIDS 145.4
EVAPORATED SOLIDS AT 105 C 90.0
SPECIFIC CONDUCTANCE IN MICROHMS/CM AT 25 C 170.0
ACIDITY AS CaCO₃

SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
ARSENIC (AS) .001 COPPER (CU) .23 NICKEL (NI) <.02
CADMIUM (CD) .014 LEAD (PB) .08 ZINC (ZN) .16
CHROMIUM (CR) <.02 MERCURY (HG)

ADDITIONAL PARAMETERS
AA CALCIUM (CA) 18.4 AA MAGNESIUM (MG) 7.00

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS Labeled TR-TOTAL RECOVERABLE
SAMPLED BY RLO
ANALYSIS REQUESTED BY GMA PERCENT REACTANCE VALUES
DATE ANALYSIS COMPLETED 08-14-72 51 38 15 1 3 75 22 0 0
ANALYST MAC OSC PEAS. OSC CALC.
NOTE: IN CORRESPONDENCE RELATED TO THIS ANALYSIS REFER TO NUMBER 72-1146

ENVIRONMENTAL ENGINEERING DEPARTMENT
THE ANACONDA COMPANY BUTTE, MONTANA 59711
WATER QUALITY ANALYSIS

NATION OR STATE MONTANA
PROV. OR COUNTY LEWIS & CLARK
LOCATION
DATE SAMPLED 08-04-72
TIME SAMPLED 1620
WATER FLOW RATE
FLOW METER METHOD
PROPERTY OR OPERATION HEADLUSTON
LABORATORY NUMBER 72-1147
SAMPLE OR BOTTLE NUMBER 565
STATION NUMBER
SAMPLE TYPE DRAINAGE
SAMPLE NUMBER 41
PRINCIPAL USE CH WATERS

PROJECT HEADLUSTON M O
REMARKS NEAR TPAP CREEK BELOW TAILINGS POND

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED

	MG/L	MG/L	MG/L	MG/L
CALCIUM (CA)	22.0	1.008	DICARBONATE (HCO ₃)	12.5
MAGNESIUM (MG)	15.2	1.250	CARBONATE (CO ₃)	0.0
SODIUM (NA)	3.5	0.152	HYDROXIDE (OH)	0.0
POTASSIUM (K)	0.94	0.024	CHLORIDE (CL)	1.9
TOT. IRON (FE)	1.1	0.005	SULFATE (SO ₄)	34.0
MANGANESE (MN)	0.08	0.003	NITRATE (NO ₃)	0.02
ALUMINUM (AL)	0.2	0.022	FLUORIDE (F)	0.010
SILICA (SIO ₂)			PHOSPHATE (PO ₄)	

TOTAL CATIONS 2.545
STANDARD DEVIATION OF CATION-ANION BALANCE 1.80 SIGMA

LABORATORY PH 8.24
FIELD TEMPERATURE DEGREES F 59.0
CALCULATED DISSOLVED SOLIDS 202.5
EVAPORATED SOLIDS AT 105 C 120.0
SPECIFIC CONDUCTANCE IN MICROHMS/CM AT 25 C 230.0
ACIDITY AS CaCO₃

SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
ARSENIC (AS) <.008 COPPER (CU) <.02 NICKEL (NI) <.02
CADMIUM (CD) <.010 LEAD (PB) <.02 ZINC (ZN) .08
CHROMIUM (CR) <.02 MERCURY (HG)

ADDITIONAL PARAMETERS
AA CALCIUM (CA) 21.8 AA MAGNESIUM (MG) 15.60
PH, FIFLO 5.0

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS Labeled TR-TOTAL RECOVERABLE
SAMPLED BY RLO
ANALYSIS REQUESTED BY GMA PERCENT REACTANCE VALUES
DATE ANALYSIS COMPLETED 08-14-72 43 90 6 1 2 25 73 0 0
ANALYST MAC OSC PEAS. OSC CALC.
NOTE: IN CORRESPONDENCE RELATED TO THIS ANALYSIS REFER TO NUMBER 72-1147

Appendix H (continued)

ENVIRONMENTAL ENGINEERING DEPARTMENT
THE BRACONCA COMPANY BUTTE, MONTANA 59701
WATER QUALITY ANALYSIS

NATION OR STATE MONTANA PROPERTY OR OPERATION HEDDLESTON
PROV. OR COUNTY LEWIS & CLARK LABORATORY NUMBER 73-1690
LOCATION 07-12-73 SAMPLE OR BOTTLE NUMBER 205
DATE SAMPLED 07-12-73 ANALYST RLD
TIME SAMPLED 1200 SAMPLE TYPE DRAINAGE
WATER FLOW RATE 300. GPM (E) SAMPLE HANDLING 41
WATER FLOW RATE 300. GPM (E) PRINCIPAL USE OF WATER
FLOWS METHOD (MN)

PROJECT HEDDLESTON
REMARKS ONTLE HOLE 0274-135 FLOWING HOLE

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED	MG/L	NEG/L
CALCIUM (CA)	14.2	0.000
MAGNESIUM (MG)	0.4	0.000
SODIUM (NA)	0.4	0.000
POTASSIUM (K)	1.40	0.000
TOT. IRON (FE)	14.2	0.000
MANGANESE (MN)	0.70	0.000
ALUMINUM (AL)	2.0	0.000
SILICA (SI)	0.222	0.000

TOTAL CATIONS	243.1	TOTAL ANIONS	0.200
STANDARD DEVIATION OF CATION-ANION BALANCE	2.4%	SIGMA	
LABORATORY PH	42.0	CARBONATE HARDNESS AS CaCO ₃	0
FIELD TEMPERATURE DEGREES F	59.9	NON-CARB. HARDNESS AS CaCO ₃	59
CALCULATED DISSOLVED SOLIDS	0	TOTAL HARDNESS AS CaCO ₃	59
EVAPORATED SOLIDS AT 105 C	0	TOTAL ALKALINITY AS CaCO ₃	0
SPECIFIC CONDUCTANCE IN MICROHMS/CM AT 25 C	0	FIELD TURBIDITY (JTU)	0
ACIDITY AS CaCO ₃	0	TOTAL FILTERABLE SOLIDS	0
		DISSOLVED SOLIDS (TDS)	0
SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)			
ARSENIC (AS)	0.05	NICKEL (NI)	0.02
COPPER (CU)	0.05	ZINC (ZN)	0.02
CHROMIUM (CR)	0.05		
MERCURY (HG)	0.05		

ADDITIONAL PARAMETERS
AA CALCIUM (CA) 12.0 AA MAGNESIUM (MG) 0.42

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS Labeled TR-TOTAL RECOVERABLE
ANALYSIS REQUESTED BY RLD PERCENT REACTANCE VALUES
DATE ANALYSIS COMPLETED 8-14-72
ANALYST RLD
THIS ANALYSIS PERTAIN TO FORM 72-1139

ENVIRONMENTAL ENGINEERING DEPARTMENT
THE BRACONCA COMPANY BUTTE, MONTANA 59701
WATER QUALITY ANALYSIS

NATION OR STATE MONTANA PROPERTY OR OPERATION HEDDLESTON
PROV. OR COUNTY LEWIS & CLARK LABORATORY NUMBER 73-1690
LOCATION 07-12-73 SAMPLE OR BOTTLE NUMBER 205
DATE SAMPLED 07-12-73 ANALYST RLD
TIME SAMPLED 1200 SAMPLE TYPE DRAINAGE
WATER FLOW RATE 300. GPM (E) SAMPLE HANDLING 41
WATER FLOW RATE 300. GPM (E) PRINCIPAL USE OF WATER
FLOWS METHOD (MN)

PROJECT HEDDLESTON WATER QUAL ENV ENG DEPT
REMARKS NIKE HORSE TUNNEL

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED	MG/L	NEG/L	MG/L	NEG/L
CALCIUM (CA)	96.0	0.000	BICARBONATE (HCO ₃)	1000.0
MAGNESIUM (MG)	83.0	0.000	CARBONATE (CO ₃)	0.0
SODIUM (NA)	1.40	0.000	HYDROXIDE (OH)	0.0
POTASSIUM (K)	1.75	0.000	CHLORIDE (CL)	0.0
TOT. IRON (FE)	9.0	0.000	SULFATE (SO ₄)	652.0
MANGANESE (MN)	0.857	0.000	NITRATE (NO ₃)	0.52
ALUMINUM (AL)	0.027	0.000	FLUORIDE (F)	0.680
SILICA (SI)	0.0	0.000	PHOSPHATE (PO ₄)	0.0

TOTAL CATIONS 17.655 TOTAL ANIONS 1000.0
STANDARD DEVIATION OF CATION-ANION BALANCE 2.4%

LABORATORY PH	6.80	CARBONATE HARDNESS AS CaCO ₃	85
FIELD TEMPERATURE DEGREES F	52.0	NON-CARB. HARDNESS AS CaCO ₃	496
CALCULATED DISSOLVED SOLIDS	964.5	TOTAL HARDNESS AS CaCO ₃	581
EVAPORATED SOLIDS AT 105 C	1156.0	TOTAL ALKALINITY AS CaCO ₃	85
SPECIFIC CONDUCTANCE IN MICROHMS/CM AT 25 C	1140.0	FIELD TURBIDITY (JTU)	0
ACIDITY AS CaCO ₃	0	TOTAL FILTERABLE SOLIDS	0
		DISSOLVED SOLIDS (TDS)	0

SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
ARSENIC (AS) 0.04 COPPER (CU) 0.05 NICKEL (NI) 0.01
CHROMIUM (CR) 0.05 MERCURY (HG) 0.05 ZINC (ZN) 0.01

ADDITIONAL PARAMETERS
AAANBENEDSON (CR) 0.0
TR SILVER (AG) 0.01

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS Labeled TR-TOTAL RECOVERABLE
ANALYSIS REQUESTED BY RLD PERCENT REACTANCE VALUES
DATE ANALYSIS COMPLETED 07-20-73
ANALYST RLD
THIS ANALYSIS PERTAIN TO FORM 72-1139

Appendix M (continued)

ENVIRONMENTAL ENGINEERING DEPARTMENT
THE ANACONDA COMPANY BUTTE, MONTANA 59701
WATER QUALITY ANALYSIS

NATION OR STATE MONTANA PROPERTY OR OPERATION HEDDLESTON
PRCV. OR COUNTY LEMISCLARK LABORATORY NUMBER 73-1687
LOCATION SAMPLE OR BOTTLE NUMBER 150G
DATE SAMPLED 07-12-73 STATION NUMBER
TIME SAMPLED 1145 SAMPLE TYPE DRAINAGE
WATER FLOW RATE GPM IE SAMPLE HANDLING 41
FLOW MEAS METHOD INH PRINCIPAL USE OF WATER

PROJECT HEDDLESTON WATER QUAL ENV ENG DEPT
REMARKS ANACONDA TUNNEL

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED

PARAMETERS	MG/L	MEQ/L	MG/L	MEQ/L
CALCIUM (CA)	54.	2.700	BICARBONATE (HCO3)	
MAGNESIUM (MG)	28.	2.316	CARBONATE (CO3)	
SODIUM (NA)	3.70	0.161	HYDROXIDE (OH)	
POTASSIUM (K)	1.40	0.036	CHLORIDE (CL)	0.0
TOT. IRON (FE)	8.3	0.446	SULFATE (SO4)	466.
MANGANESE (MN)	15.5	0.364	NITRATE (NO3)	0.54
ALUMINUM (AL)	4.2	0.467	FLUORIDE (F)	0.438
SILICA (SIO2)			PHOSPHATE (PO4)	0.0
TOTAL CATIONS	6.490		TOTAL ANIONS	9.228
STANDARD DEVIATION OF CATION-ANION BALANCE	9.99	SIGMA		
LABORATORY PH	2.77	CARBONATE HARDNESS AS CaCO3	0	
FIELD TEMPERATURE DEGREES F	64.0	NON-CARB. HARDNESS AS CaCO3	251	
CALCULATED DISSOLVED SOLIDS	582.6	TOTAL HARDNESS AS CaCO3	251	
EVAPORATED SOLIDS AT 105 C	712.	TOTAL ALKALINITY AS CaCO3	0	
SPECIFIC CONDUCTANCE IN MICROHMS/CM AT 25 C	1110.0	FIELD TURBIDITY (JTU)		
ACIDITY AS CaCO3		TOTAL FILTERABLE SOLIDS		
		DISSOLVED OXYGEN AS (O2)		

SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
ARSENIC (AS) <.001 COPPER (CU) 1.75 NICKEL (NI) .050
CADMIUM (CD) <.043 LEAD (PB) .405 ZINC (ZN) 4.30
CHROMIUM (CR) <.003 MERCURY (HG)

ADDITIONAL PARAMETERS
ARABENNESUM (CRS) 84.0
TR SILVER (AG) .010

NOTE. PARAMETERS ARE TOTAL DISSOLVED UNLESS LABELED TR-TOTAL RECOVERABLE
SAMPLED BY RLO PERCENT REACTANCE VALUES
ANALYSIS REQUESTED BY RLO CA MG NA K CL SC4 HCO3 CO3 NO3
DATE ANALYSIS COMPLETED 07-20-73 52 44 3 1 0 100 0 0 0
ANALYST LAB OSC MEAS. OSC CALC.

NOTE. IN CORRESPONDENCE RELATED TO THIS ANALYSIS REFER TO NUMBER T3-1687

ENVIRONMENTAL ENGINEERING DEPARTMENT
THE ANACONDA COMPANY BUTTE, MONTANA 59701
WATER QUALITY ANALYSIS

NATION OR STATE MONTANA PROPERTY OR OPERATION HEDDLESTON
PRCV. OR COUNTY LEMISCLARK LABORATORY NUMBER 73-1686
LOCATION SAMPLE OR BOTTLE NUMBER 1780
DATE SAMPLED 07-12-73 STATION NUMBER
TIME SAMPLED 1100 SAMPLE TYPE DRAINAGE
WATER FLOW RATE 1. GPM IE SAMPLE HANDLING 41
FLOW MEAS METHOD INH PRINCIPAL USE OF WATER

PROJECT HEDDLESTON WATER QUAL ENV ENG DEPT
REMARKS E01H NO 2 TUNNEL

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED

PARAMETERS	MG/L	MEQ/L	MG/L	MEQ/L
CALCIUM (CA)	10.4	0.520	BICARBONATE (HCO3)	
MAGNESIUM (MG)	7.6	0.623	CARBONATE (CO3)	
SODIUM (NA)	8.50	0.370	HYDROXIDE (OH)	
POTASSIUM (K)	6.2	0.155	CHLORIDE (CL)	0.0
TOT. IRON (FE)	10.0	0.537	SULFATE (SO4)	180.
MANGANESE (MN)	1.00	0.036	NITRATE (NO3)	0.22
ALUM. (NAI AL)	5.7	0.634	FLUORIDE (F)	0.42
SILICA (SIO2)			PHOSPHATE (PO4)	0.0
TOTAL CATIONS	2.879		TOTAL ANIONS	3.931
STANDARD DEVIATION OF CATION-ANION BALANCE	6.61	SIGMA		
LABORATORY PH	2.98	CARBONATE HARDNESS AS CaCO3	0	
FIELD TEMPERATURE DEGREES F	60.	NON-CARB. HARDNESS AS CaCO3	57	
CALCULATED DISSOLVED SOLIDS	237.6	TOTAL HARDNESS AS CaCO3	57	
EVAPORATED SOLIDS AT 105 C	414.	TOTAL ALKALINITY AS CaCO3	0	
SPECIFIC CONDUCTANCE IN MICROHMS/CM AT 25 C	666.0	FIELD TURBIDITY (JTU)		
ACIDITY AS CaCO3		TOTAL FILTERABLE SOLIDS		
		DISSOLVED OXYGEN AS (O2)		

SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
ARSENIC (AS) <.001 COPPER (CU) 3.70 NICKEL (NI) .056
CADMIUM (CD) .013 LEAD (PB) .148 ZINC (ZN) .61
CHROMIUM (CR) .602 MERCURY (HG)

ADDITIONAL PARAMETERS
ARABENNESUM (CRS) 10.4
TR SILVER (AG) .001

NOTE. PARAMETERS ARE TOTAL DISSOLVED UNLESS LABELED TR-TOTAL RECOVERABLE
SAMPLED BY RLO PERCENT REACTANCE VALUES
ANALYSIS REQUESTED BY RLO CA MG NA K CL SC4 HCO3 CO3 NO3
DATE ANALYSIS COMPLETED 07-20-73 31 37 22 9 0 100 0 0 0
ANALYST LAB OSC MEAS. OSC CALC.

NOTE. IN CORRESPONDENCE RELATED TO THIS ANALYSIS REFER TO NUMBER T3-1686

Appendix M (continued)

ENVIRONMENTAL ENGINEERING DEPARTMENT
THE ANACONDA COMPANY BUTTE, MONTANA 59701
WATER QUALITY ANALYSIS

NATION OR STATE MONTANA PROPERTY OR OPERATION HEGGLESTON
PRCV. OR COUNTY LEWIS & CLARK LABORATORY NUMBER 73-1684
LOCATION LOCATION SAMPLE OR BOTTLE NUMBER 1556
DATE SAMPLED 07-12-73 SAMPLE TYPE GRAINAGE
TIME SAMPLED 1130 SAMPLE HANDLING 41
WATER FLOW RATE 2. GPM (E) PRINCIPAL USE OF WATER
FLOW MEAS METHOD (MN)

PROJECT HEGGLESTON WATER QUAL ENV ENG
REMARKS TUNNEL NO 3

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED

MG/L	MEQ/L	MG/L	MEQ/L
CALCIUM (CA)	1.3	0.064	BICARBONATE (HCO3)
MAGNESIUM (MG)	0.0	0.495	CARBONATE (CO3)
SODIUM (NA)	1.60	0.070	HYDROXIDE (OH)
POTASSIUM (K)	4.70	0.120	CHLORIDE (CL)
TOT. IRON (FE)	19.0	1.021	SULFATE (SO4)
MANGANESE (MN)	.43	0.016	NITRATE (NO3)
ALUMINUM (AL)	10.0	1.112	FLUORIDE (F)
SILICA (SIO2)			PHOSPHATE (PO4)

TOTAL CATIONS 2.898 TOTAL ANIONS 5.276
STANDARD DEVIATION OF CATION-ANION BALANCE 9.99 SIGMA

LABORATORY PH 2.84 CARBONATE HARDNESS AS CaCO3 0
FIELD TEMPERATURE DEGREES F 42. NON-CARB. HARDNESS AS CaCO3 28
CALCULATED DISSOLVED SOLIDS 296.1 TOTAL HARDNESS AS CaCO3 28
EVAPORATED SOLIDS AT 105 C 486. TOTAL ALKALINITY AS CaCO3 0
SPECIFIC CONDUCTANCE IN FIELD TURBIDITY (JTU)
MICROHMS/CM AT 25 C 855.0 DISSOLVED OXYGEN AS (O2)

SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
ARSENIC (AS) .04 COPPER (CU) 28.9 NICKEL (NI) .085
CADMIUM (CD) .004 LEAD (PB) .005 ZINC (ZN) .26
CHROMIUM (CR) .009 MERCURY (HG)

ADDITIONAL PARAMETERS
AAAMGUESUM (CDB) 8.0
TR SILVER (AG) .001

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS LABELED TR-TOTAL RECOVERABLE
SAMPLED BY RLO PERCENT REACTANCE VALUES
ANALYSIS REQUESTED BY RLO CA PG NA K CL SCA HCO3 CO3 MG3
DATE ANALYSIS COMPLETED 07-20-73 4 46 116 0 100 0 0 0
ANALYST LAB OSC MEAS. OSC CALC.
NOTE: IN CORRESPONDENCE RELATED TO THIS ANALYSIS REFER TO NUMBER 73-1684

ENVIRONMENTAL ENGINEERING DEPARTMENT
THE ANACONDA COMPANY BUTTE, MONTANA 59701
WATER QUALITY ANALYSIS

NATION OR STATE MONTANA PROPERTY OR OPERATION HEGGLESTON
PRCV. OR COUNTY LEWIS & CLARK LABORATORY NUMBER 73-1682
LOCATION LOCATION SAMPLE OR BOTTLE NUMBER 264
DATE SAMPLED 07-12-73 STATION NUMBER
TIME SAMPLED 1230 SAMPLE TYPE STREAM
WATER FLOW RATE 2. GPM (E) SAMPLE HANDLING 41
FLOW MEAS METHOD (MN) PRINCIPAL USE OF WATER

PROJECT HEGGLESTON WATER QUALITY ENV ENG DEPT
REMARKS BEAR TRAP CREEK BELOW TAILINGS POND

PARAMETERS REPORTED IN MILLIGRAMS PER LITER EXCEPT AS INDICATED

MG/L	MEQ/L	MG/L	MEQ/L
CALCIUM (CA)	11.0	0.548	BICARBONATE (HCO3)
MAGNESIUM (MG)	14.7	1.210	CARBONATE (CO3)
SODIUM (NA)	1.25	0.054	HYDROXIDE (OH)
POTASSIUM (K)	.75	0.019	CHLORIDE (CL)
TCT. IRON (FE)	.083	0.004	SULFATE (SO4)
MANGANESE (MN)	.069	0.003	NITRATE (NO3)
ALUMINUM (AL)	.02	0.002	FLUORIDE (F)
SILICA (SIO2)			PHOSPHATE (PO4)

TOTAL CATIONS 1.841 TOTAL ANIONS 2.258
STANDARD DEVIATION OF CATION-ANION BALANCE 3.01 SIGMA

LABORATORY PH 8.08 CARBONATE HARDNESS AS CaCO3 85
FIELD TEMPERATURE DEGREES F 54.0 NON-CARB. HARDNESS AS CaCO3 3
CALCULATED DISSOLVED SOLIDS 158.4 TOTAL HARDNESS AS CaCO3 88
EVAPORATED SOLIDS AT 105 C 210. TOTAL ALKALINITY AS CaCO3 85
SPECIFIC CONDUCTANCE IN FIELD TURBIDITY (JTU)
MICROHMS/CM AT 25 C 240.0 TOTAL FILTERABLE SOLIDS
ACIDITY AS CaCO3 DISSOLVED OXYGEN AS (O2)

SELECTED TRACE METALS (RECOVERABLE BY ACID DIGESTION OF UNFILTERED SAMPLE)
ARSENIC (AS) .03 COPPER (CU) .013 NICKEL (NI) .007
CADMIUM (CD) .002 LEAD (PB) .03 ZINC (ZN) .665
CHROMIUM (CR) .001 MERCURY (HG)

ADDITIONAL PARAMETERS
AAAMGUESUM (CDB) 16.0
TR SILVER (AG) .001

NOTE: PARAMETERS ARE TOTAL DISSOLVED UNLESS LABELED TR-TOTAL RECOVERABLE
SAMPLED BY RLO PERCENT REACTANCE VALUES
ANALYSIS REQUESTED BY RLO CA PG NA K CL SCA HCO3 CO3 NO3
DATE ANALYSIS COMPLETED 07-20-73 30 46 3 1 0 24 75 0 0
ANALYST LAB OSC MEAS. OSC CALC.
NOTE: IN CORRESPONDENCE RELATED TO THIS ANALYSIS REFER TO NUMBER 73-1682

APPENDIX N

Results of Metals Analyses of
Stream Sediments, 1973

November 12, 1973

MONTANA BUREAU OF MINES AND GEOLOGY

BUTTE, MONTANA 59701

November 15, 1973

Mr. Lter Spence
Box 27
Ovando, MT 59854MF60 Stream Sediment Samples
Received 10-11-73

Field No.	Cu	Ni	Pb	Cd	Zn	Fe	LOI @ 750°C
1	24	10	47.5	<1	52	25,700	4.80
2	23	10	28	<1	53	26,700	4.55
3	88	20	70	1	73.5	24,700	9.40
4	29	10.5	28	1	41.5	25,700	7.33
5	47	11.5	29.5	1.2	259	25,700	4.55
Repeat	47	21.5	29.5	1.0	259	not done - not enough sample.	
6	42	12	32.5	<1	35	31,700	5.10
7	80	30.5	83	47.5	1,000	34,600	5.98
8	700	50	400	282.5	4,080	36,600	7.85
9	73	18	39.5	1.2	152	40,600	10.70
10	25	12	24.5	1.0	77.5	29,700	5.90

Respectfully submitted,

L. A. Megelin
Chief Chemist

LAN/np

Note: Results expressed in micrograms per gram, dry weight. LOI = Loss
of organic matter on ignition in percent.

ITEMIZED INVOICE -- Stream Sediment Samples

Lab No.

Sample No.

1	0920	Blackfoot R @ Canyon Camp	8-14-73
2	1100	Blackfoot R @ Dalton Mtn. Bridge	8-14-73
3	1150	Footman Ck	8-14-73
4	1340	Landers Fork of Blackfoot R	8-14-73
5	1520	Blackfoot R @ Hogeim Rd Bridge	8-14-73
6	1545	Alice Ck @ Highway 200, Bridge	8-14-73
7	1615	Blackfoot R @ Fleisher	8-14-73
8	1650	Blackfoot R @ Pop's Place	8-14-73
9	1440	Willow Creek (Fleisher)	8-14-73
10	1140	Blackfoot R. below 7-up	8-20-73

10 Stream Sediment Samples @ \$6 \$60.00

A-75

APPENDIX H (continued)

MONTANA BUREAU OF MINES AND GEOLOGY
BUTTE, MONTANA 59701

April 25, 1974

Mr. Lister Spence
Montana Fish and Game Commission
Box 27
Ovando, Montana 59854

Dear Lister:

Please find under separate cover the canvas sample sack as requested. Please return it with approximately 25 pounds of sediment.

We will complete the following investigations that may assist you in understanding the relationship of metal ion concentration versus particle size. We will separate the particles with a series of standard sieves and analyze each series or fraction for Cu, Pb, Zn, Cd, and Ni, and will also provide you with analyses showing what percent each fraction or group of particles represents in the sample. The sieve analysis and trace metal analyses will be done in duplicate to reveal homogeneity and replication of values.

I have included a copy of the report, dated November 15, 1973, to clarify how these values were reported. The metals are reported in micrograms per gram, meaning that 1×10^{-6} grams is one microgram (1 ppm) or $\frac{.000001 \text{ grams}}{1,000,000 \text{ grams}}$. For example - Field No. 1 shows 24 for Cu.

This means that each gram of sediment of -100 mesh size contains .000024 grams of elemental copper, commonly referred to as parts per million or formally expressed as 24 micrograms/gram ($\mu\text{g/g}$). The L.O.I. (loss on ignition) is expressed in percent (Weight/Weight). For example - Field No. 1 shows 4.80 L.O.I. @ 750°. This indicates that each gram of -100 mesh sediment loses .048 grams upon heating the sample to 750°C. I apologize for not clarifying this on the report.

On Field sample No. 5, as discussed on the telephone, the only significant change was that of Ni from 11.5 $\mu\text{g/g}$ to 21.5 $\mu\text{g/g}$. I shall try to explain in more detail why replicate analyses do not always agree closely. First, homogeneity of the sample is difficult to obtain without more elaborate preparation, which may be justified in your investigations.

To approach homogeneity, the bulk of the sample must consist of particles of almost uniform size. Normally this requires pulverization of the sample to a very fine particle,

Mr. Lister Spence
Page two
April 25, 1974

-325 mesh. The Bureau and other investigators have found that by removing the -100 mesh fraction from soils and sediments they begin to approach homogeneity to the degree that metal concentration profiles are well enough defined to allow interpretation of their mobilities. After giving this some thought, do you feel that you must have a more precisely defined profile, approaching absolute trace metal values? If so, we can pulverize the -100 material or the entire sediment sample to -325 to assure homogeneity.

Respectfully yours,

L. A. Wegelin
Chief Chemist

LAW:wh
Enclosure

MONTANA BUREAU OF MINES AND GEOLOGY

BUTTE, MONTANA 59701

July 17, 1974

Mr. Liter Spence
Montana Fish & Game Commission
Helena, Montana 59601

Dear Liter:

I am reporting the results of our investigations on the large sediment sample. To prepare the sample for analysis it was dried at 110°C to constant weight, blended to assure homogeneity, and split into four representative portions.

The particle size or sieve analyses were performed on two of the four split portions, and the elemental determinations were performed in duplicate on each fraction of the two sieve analyses. I did this to reveal repetition and homogeneity and to provide enough statistical information for standard deviation. The data are shown in four tables. Table I converts mesh (Tyler, U.S. Number) to actual opening dimension of each sieve. Table II shows trace-metal relationship to particle size.

Table III shows each fraction or group of particles in terms of weight percent of the sample. Table IV shows the standard deviation of the analyses versus particle size.

I have enclosed a graph relating particle size to trace-metal concentrations. In the graph, the metals show a sharp increase in concentration from the -200 to -230 fractions. Looking at this, your investigations may justify using the -200 mesh particle rather than the -100 mesh particle. In looking at Table IV you will notice how the standard deviation of the analyses improves as the particle becomes smaller. I believe I mentioned this to you previously. Again, the standard deviation, no matter what fraction or particle size, would be improved tremendously if the samples were pulverized to -325. This again may be another consideration for future work.

I hope you find this report in order and informative, and should you have any questions please feel free to inquire.

Yours truly,

Laurence A. Wegelin
Laurence A. Wegelin
Chief Chemist

LAW/jp

A-17

APPENDIX N continued

TABLE I

Tyler Sieve (mesh)
U.S. Standard No.

Opening in millimeters

-16	1.190
-20	.841
-40	.420
-60	.250
-100	.149
-140	.105
-200	.074
-230	.063
-325	.044

A-9

Opening in inches

Tyler Sieve (mesh)
Standard U.S. No.

	Cu	Pb	Zn	Cd	Ni	Fe
.0469 Sieve Test +16	25	11.5	120	.05	11.5	15,470
A +16	36.5	13	130	.05	14	13,970
.0331 Sieve Test +16	26.5	29	115	.05	11.5	13,470
B +16	<u>45.5</u>	<u>20</u>	<u>1725</u>	<u>.05</u>	<u>11</u>	<u>13,470</u>
.0098 Mean	33.38	18.38	134.38	.05	12.00	14,095.00
.0059 Std. Deviation	9.56	7.99	26.17	.00	1.35	946.46
.0041 Sieve Test -16	38	16	137.5	.05	13.5	18,960
A -16	37	18.5	140	.10	13.5	18,460
.0029 Sieve Test -16	38	22	145	.05	12	14,970
B -16	<u>32.5</u>	<u>16.5</u>	<u>127.5</u>	<u>.05</u>	<u>12</u>	<u>16,970</u>
.0025 Mean	36.38	18.25	137.50	.06	12.75	17,340.00
.0017 Std. Deviation	2.63	2.72	7.36	.03	.87	1,791.89
Sieve Test -20	38	22	157.5	.05	13.5	18,960
A -20	38	20	142.5	.05	13	18,960
.0025 Sieve Test -20	37	21	172.5	.08	14.5	20,460
B -20	<u>36</u>	<u>18</u>	<u>160</u>	<u>.08</u>	<u>13</u>	<u>20,460</u>
.0025 Mean	37.25	20.25	158.13	.07	13.50	19,710.00
.0017 Std. Deviation	.96	1.71	12.31	.02	.71	866.03
Sieve Test -40	36	24.5	175	.05	14	21,460
A -40	36	32.5	180	.05	14	21,460
.0017 Sieve Test -40	38	21	175	.08	13.5	19,960
B -40	<u>45</u>	<u>21.5</u>	<u>175</u>	<u>.08</u>	<u>13.5</u>	<u>19,960</u>
.0017 Mean	38.75	24.88	176.25	.07	13.75	20,710
.0017 Std. Deviation	4.27	5.31	2.50	.02	.29	866.03

TABLE II

Metal Concentration/Particle Size Relationship

APPENDIX N continued

Sieve Test A	-60 -60	36.5 35.5	19.5 17	140 425	.05 .05	11.5 19	16,970 15,970	Sieve Test A	-230 -230	95 105.5	32.5 36	250 262.5	.05 .05	14.5 15	19,960 19,960
Sieve Test B	-60 -60	39 33	22.5 24	152.5 142	.05 .05	11.5 11	17,960 18,460	Sieve Test B	-230 -230	102 111	26.5 23	192.5 187.5	.08 .08	11.5 11.5	20,960 20,460
Mean		36.00	20.75	214.88	.05	13.25	17,340.00	Mean		103.38	29.50	223.13	.07	13.13	20,335.00
Std. Deviation		2.48	3.12	140.19	.00	3.84	1,103.41	Std. Deviation		6.70	5.85	38.64	.02	1.88	478.71
Sieve Test A	-100 -100	30.5 39.5	20 22.5	157.5 155	.05 .05	9 11	17,470 16,470	Sieve Test A	-325 -325	122.5 125.5	34 60	270 270	.13 .13	15 15	24,450 24,450
Sieve Test B	-100 -100	39 34	24 22	147.5 145	.08 .08	11 11	18,960 18,960	Sieve Test B	-325 -325	127.5 124	35 36	262.5 262.5	.13 .13	14.5 15	24,950 24,450
Mean		35.75	22.12	151.25	.07	10.50	17,965.00	Mean		124.88	41.25	266.25	.13	14.88	24,575.00
Std. Deviation		4.29	1.65	5.95	.02	1.00	1,219.30	Std. Deviation		2.14	12.53	4.33	.00	.25	250.00
Sieve Test A	-140 -140	40 35.5	20 19	612.5 140	.05 .05	42 9	18,460 18,460	Note: All concentrations are reported as micrograms per gram (ppm).							
Sieve Test B	-140 -140	46 40	24 24	150 152.5	.08 .08	10.5 10.5	19,960 20,460								
Mean		40.38	21.75	263.75	.07	18.00	19,335.00								
Std. Deviation		4.30	2.63	232.56	.02	16.02	1,030.00								
Sieve Test A	-200 -200	53.5 60	25 22.5	157.5 182.5	.05 .05	9.5 9	19,460 18,960								
Sieve Test B	-200 -200	55 54.5	23 24	165 167.5	.08 .08	11 11	19,460 18,460								
Mean		55.75	23.63	168.13	.07	10.13	19,563.71								
Std. Deviation		2.90	1.11	10.48	.02	1.03	478.71								

APPENDIX B continued

TABLE III
Sieve Analysis

	Sieve Test	Tyler Sieve (mesh)		Grams Recovered	% (W/W)				
		U.S. Standard No.							
A-20	A	+16		271.0	14.8	A	-230	29.7	1.6
	B	+16		237.8	13.3	B	-230	28.5	1.6
	A	-16		149.9	8.2	A	-325	34.2	1.9
	B	-16		147.4	8.3	B	-325	31.7	1.8
	A	-20		226.3	12.3	Sieve Test A			
	B	-20		220.7	12.4				
	A	-40		403.9	22.0	Total Recovered Wt. Total Beginning Wt. Percent Recovered			
	B	-40		427.6	24.0				
	A	-60		466.7	25.4	Sieve Test B			
	B	-60		440.7	24.7				
	A	-100		145.0	7.9	Total Recovered Wt. Total Beginning Wt. Percent Recovered			
	B	-100		141.2	7.9				
A	-140		90.2	4.9					
B	-140		88.8	5.0					
A	-200		17.1	.9					
B	-200		18.3	1.0					
							Total Recovered Wt.		
							1,834.0		
							1,836.3		
							99.87		
							Total Recovered Wt.		
							1,782.7		
							1,784.0		
							99.92%		

TABLE IV
Standard Deviation/Particle Size

	<u>Cu</u> Mean & Std. Dev.	<u>Pb</u> Mean & Std. Dev.	<u>Zn</u> Mean & Std. Dev.	<u>Cd</u> Mean & Std. Dev.	<u>Ni</u> Mean & Std. Dev.	<u>Fe</u> Mean & Std. Dev.
+16	33.38 9.56	18.38 7.99	134.38 26.17	.05 .00	12.00 1.35	14,095.00 946.48
-16	36.38 2.63	18.25 2.72	137.50 7.36	.06 .03	12.75 .87	17,340.00 1,791.89
-20	37.25 .96	20.25 1.71	158.13 12.31	.07 .02	13.50 .71	19,710.00 866.00
-40	38.75 4.27	24.88 5.31	176.25 2.50	.07 .02	13.75 .29	20,710 866.03
-60	36.00 2.48	20.75 3.12	214.88 140.19	.05 .00	13.25 3.84	17,340.00 1,103.41
-100	35.75 4.29	22.12 1.65	151.25 5.95	.07 .02	10.50 1.00	17,965.00 1,219.30
-140	40.38 4.30	21.75 2.63	263.75 232.56	.07 .02	18.00 16.02	19,335.00 1,030.00
-200	55.75 2.90	23.63 1.11	168.13 10.48	.07 .02	10.13 1.03	19,563.71 478.71
-230	103.38 6.70	29.50 5.85	223.13 38.64	.07 .02	13.13 1.88	20,335.00 478.71
-325	124.88 2.14	41.25 12.53	266.25 4.33	.13 .00	14.88 .25	24,575.00 250.00

APPENDIX O
Bottom Fauna Station Locations

Station Number	Station Name	Location	ELEVATION (ft. msl)
1	Blackfoot River at Pop's Place	Across from Pop's Place Bar, 1 mile downstream from mouth of Pass Creek	5130
2	Blackfoot River at Flesher	Upstream from culvert at crossing of the river by state highway 279	4960
3	Alice Creek above National Forest Boundary	At Bear Gulch road crossing of creek 1.1 mile above national forest boundary	5400
4	Alice Creek at Highway 200	50 ft. upstream from highway bridge	4950
5	Blackfoot River at Hogum	50 ft. upstream from USFS bridge crossing river on Hogum Creek road	4815
6	Landers Fork at Highway 200	300 ft. upstream from highway bridge	4750
7	Blackfoot River between Landers Fork and Lincoln (below Landers)	At end of county road 1.4 miles downstream from mouth of Landers Fork	4680
8	Poorman Creek	0.1 mile south of where Poorman Creek first crosses Stemple road about 3 miles south of Lincoln	4640
9	Blackfoot River at Dalton	300 ft. upstream from Dalton Mountain road bridge west of Lincoln	4460
10	Blackfoot River at Canyon	50 ft. upstream from USFS cement bridge over river at Blackfoot Canyon campground	4400
11	Blackfoot River at Rainbow Bend	At small island in river which is reached by taking pulloff to the left 1/8 mile downstream from Rainbow Bend and going 100 yards to river	3400

APPENDIX P

Kinds and abundance of bottom fauna collected from the Blackfoot River and certain tributaries, 1968 - 1971, including diversity indices.

Organisms were sorted to order, counted and volumes determined at the Pollution Control Laboratory, State Department of Health and Environmental Sciences, Helena. Sorted organisms were further identified to lowest taxa and recounted by Ms. Patricia Ann Mitchell, Department of Zoology, University of Alberta, Edmonton, Alberta, Canada. (1968 and 1969 samples were identified but not recounted.)

Identifications have been made only on the following samples to date: All of 1968 and 1969; March/April 1971; August 1971; and October/November 1971. The remaining samples collected in 1972 and 1973 have only been sorted and counted to order.

The 1968 and 1969 samples were ± 4 square feet in area and were obtained with a two-foot wide Needham kick screen. All other samples were obtained with a modified Waters round square foot stream bottom sampler (Waters 1961). Three one-square foot samples were collected at each station. Each square foot sample which has been identified is shown in this appendix. Numbers (i.e., #1, #2, #3) following station name are the individual square foot sample numbers.

Diversity indices were calculated using the method of Wilhm and Dorris (1968). These indices are given for each ± 4 square foot sample in 1968 and 1969, and each one square foot sample in 1971. The 1971 samples have indices given on only one sheet. However, indices are for the three one-square foot samples combined. Only one index (for the entire station) is given for the 1968 and 1969 data because the numbers in each species were not determined. This index is based on the number of orders and the number of individuals in each order. The 1971 samples are based on the number of species and number of individuals in each species. An index is given for each order as well as for the entire station. Diversity indices were not computed for all the October/November 1971 samples because the identification data were received too late to be processed prior to report completion.

Only the five major insect orders (Ephemeroptera, Plecoptera, Tricoptera, Diptera, and Coleoptera) were used in diversity index calculations. Items with a check mark (✓) preceding them were omitted from the diversity index computations due to their indefinite or non-specific identification. The diversity index computed for the entire station (three square feet combined) is shown as "Station D. I."

In the cases where an organism could be identified only to family, each family was assumed to possess at least one species and the numbers in the family were used as if they comprised only one species. The same holds true for genera if the species were unknown but shown as "spp." indicating more than one unidentifiable species was present.

Figures shown as "Original Total" are numbers of organisms of all kinds (including noninsects and fish) in a sample as counted by Pollution Control Laboratory personnel. "Final Total" was determined by Ms. Mitchell after identifications were made. The variations in numbers may be due to inaccurate original count or deterioration of insects between counts. Except for 1968 and 1969, "Final Total" was used for computing diversity indices, since it was believed to be more accurate than "Original Total".

Diversity indices were computed by Dr. Don O. Loftsgaarden, Professor of Mathematics, University of Montana, Missoula. His explanation of the procedure and computer program used is given in Appendix Q.

D = diversity index. DMAX is the maximum diversity index which could occur in the given sample, DMIN is the minimum diversity index which could occur, and R is redundancy, or a measure of how evenly or unevenly the numbers of organisms are distributed in the sample. When redundancy is high at a station, diversity is low and vice versa. DMAX and DMIN are limits which allow one to position D and aid in its interpretation. Redundancy is always between 0 and 1. Diversity can range from less than 1 in heavily polluted areas to more than 3 in clean water areas. Values from 1 to 3 are found in areas of moderate pollution (Wilhm and Dorris 1968).

A comparison of total numbers of organisms collected per square foot with the Needham kick screen and the modified Waters round-square-foot sampler (at those stations where both were used simultaneously) shows the Waters sampler to be more efficient in collecting total numbers of insects. The mesh size used in the Needham screen was 16 meshes per inch compared to 38 meshes per inch for the Waters sampler, and more insects were retained by the latter. Also the design and use of the Waters sampler precludes much loss of insects around the screen and enables one to retain more insects in a sample.

APPENDIX P CONTINUED

Blackfoot River at Pop's Place 7/16/68

#1 Plecoptera 2 T

Paraperla, Alloperla or Kathroperla spp. (too small to identify)

Trichoptera 7 .2 cc

Rhyacophila angelita Banks
Brachycentrus sp.

Coleoptera 1 T

Zaitzevia sp.

Diptera 34 .1 cc

Tendipedidae:
Hydrobaenus sp.

#2 Plecoptera 2 T

Nemoura besanetsa Ricker
Paraperla, Alloperla or Kathroperla spp. (too small to identify)

Trichoptera 5 .2 cc

Rhyacophila angelita Banks
Brachycentrus sp.
Hydropsyche sp.
Arctopsyche sp.

Station D.I.

D = 1.37131 DMIN = 0.385691
DMAX = 2. R = 0.37186

Coleoptera 7 .05 cc

Zaitzevia sp.

#1 And #2 combined

Diptera 17 T

Tendipedidae
Hydrobaenus sp.

Blackfoot at Pop's Place 4/29/69 pH 7.9

Plecoptera 1

Nemoura cinctipes Banks

Ephemeroptera 1

Cinygmula sp.

D = DMIN = DMAX

R is undefined

Blackfoot River at Pop's Place 10/1/69

Plecoptera 95, 0.3 cc

Hastriella brevis ?
Nemoura cinctipes Banks
Alloperla sp.
Isogenus sp.

Station D.I.

D = 0.244142 DMIN = 0.14620E-2 = .0814620
DMAX = 1. R = 0.022893

Trichoptera 4, 0.1 cc

Parapsyche sp.
Rhyacophila acropedes or vao
Arctopsyche sp.

APPENDIX P CONTINUED

BLACK-LOT RIVER AT POP'S PLACE #1
1 sq. ft. 8-20-71 L. Spence

See No diversity computed

EPHEMEROPTERA 1 T

Baetis sp.

1 Total 1
D = 1.36479 DMH = 0.26683
DIAK = 2.32193 DM = 0.475337
2
3
4 Total 34

PLANEPTERA 35 T

Memoria clintones
Agonostoma sp.
Allopietia sp.

TRICHOPTERA 2 T 5+2 D. DIAK = DIAK, R is undefined

✓ Pupae

1 of each species

2 Total 2
D = 2.32492 DMH = 0.927075
DIAK = 2.32193 DM = 0.212932

DIPTERA 20 T

Simuliidae
Campoclastus sp.
Helicidius sp.
Orthocentrus sp. 1

3
4
6
7 Total 20

Original Total 58 Final Total 57

Identified and Enumerated by
Patricia Ann Mitchell
MS, University of Montana, 1968
January 28, 1973

Patricia Ann Mitchell

Station D.I.

D = 2.4173 DMH = 0.812958
DIAK = 4.00000 DM = 0.496293

BLACK-LOT RIVER AT POP'S PLACE #2
1 sq. ft. 8-20-71 L. Spence

PLANEPTERA 45 T

Memoria clintones
Agonostoma sp.
Agonostoma sp.
Allopietia sp.
Allopietia sp.

29
3
1
9
2 Total 44

TRICHOPTERA 2 T

Hydroptila areolaris? (measure of species)
Hydroptila areolaris

1
1 Total 2

DIPTERA 4 T

Orthocentrus sp. 1
Orthocentrus sp. 5

3
1 Total 4

COLEOPTERA 4 T

Optioeremus sp. adult
Heterilinus convolutus adult
Salixia sp. adult

1
2
1 Total 4

Original Total 55 Final Total 54

Identified and Enumerated by
Patricia Ann Mitchell
MS, University of Montana, 1968
Dept. of Entomology
University of Alberta, Edmonton
January 28, 1973

Patricia Ann Mitchell

BLACK-LOT RIVER AT POP'S PLACE #3
1 sq. ft. 8-20-71 L. Spence

PLANEPTERA 48 T

Memoria clintones
Agonostoma sp.
Agonostoma sp.
Allopietia sp.

29
2
1
8 Total 49

DIPTERA 2 T

Simuliidae
Campoclastus sp. 1

1
1 Total 2

COLEOPTERA 1 T

Heterilinus convolutus adult

1 Total 1

Original Total 51 Final Total 52

Identified and Enumerated by
Patricia Ann Mitchell
MS, University of Montana, 1968

January 28, 1973

Patricia Ann Mitchell

APPENDIX P CONTINUED

BLACKFOOT RIVER AT POP'S PLACE #2
1 sq. ft. 10-19-71 L. Spence

BLACKFOOT RIVER AT POP'S PLACE #3
1 sq. ft. 10-19-71 L. Spence

BLACKFOOT RIVER AT POP'S PLACE #1
1 sq. ft. 10-19-71 L. Spence

PLECOPTERA 43 0.1	
<u>Hemania sinclairi</u>	23
<u>Desmopis brevicornis</u>	1
<u>Leptania sp.</u>	7
<u>Alloperla sp.</u>	10
<u>Paraperla frontalis</u>	2
	Total 43
TRICHOPTERA 2 T	
<u>Arctopsyche grandis</u>	1
<u>Parapsyche alberta</u>	1
	Total 2
DIPTERA 2 T	
<u>Metriocnemus sp.</u>	1
Unknown Chironomidae	1
	Total 2

Original Total 47 Final Total 47

PLECOPTERA 34 0.1	
<u>Hemania sinclairi</u>	12
<u>Desmopis brevicornis</u>	5
<u>Leptania sp.</u>	4
<u>Alloperla sp.</u>	12
<u>Paraperla frontalis</u>	1
	Total 34
COLEOPTERA 1 T	
<u>Zaiteria sp. adult</u>	1
	Total 1
DIPTERA 1 T	
Could not find a dipteran	

S=1 No diversity computed

Original Total 36 Final total 36

PLECOPTERA 87 0.2	
<u>Hemania sinclairi</u>	27
<u>Desmopis brevicornis</u>	12
<u>Leptania sp.</u>	13
<u>Alloperla sp.</u>	32
<u>Paraperla frontalis</u>	2
	Total 86
TRICHOPTERA 2 T	
<u>Arctopsyche grandis</u>	1
<u>Hydropsychidae (head missing)</u>	1
	Total 2
DIPTERA 26 T	
<u>Anisobla sp.</u>	1
Unknown Chironomidae	24
	Total 25

Original Total 115 Final Total 113

Identified and Enumerated by
Patricia Ann Mitchell
MS, University of Montana, 1968
July 31, 1973

Patricia Ann Mitchell

Identified and Enumerated by
Patricia Ann Mitchell
MS, University of Montana, 1968
July 31, 1973

Patricia Ann Mitchell

Identified and Enumerated by
Patricia Ann Mitchell
MS, University of Montana, 1968
July 31, 1973

Patricia Ann Mitchell

A-86

Station D.I.

b = 8.50616 DIVIN = 7.503176
BIAX = 3.55496 DE = 2.55596

Blackfoot at Fletcher Pass Road bridge 7/16/68

Hemiptera 22 .4 cc

Classocoris subulosa
Allograpta sp.
Acronycta pacifica
Neocera clausii
Acronycta sp.

Trichoptera 24 .4 cc

Hydropsyche van der aarwaden
St. acutipila sp.
St. acutipila sp.

Ephemeroptera 10 .1 cc

Ephemerella longimanus (Zanon)
Heptagenia sp.
Heptagenia sp.

Coleoptera 12 .05 cc

Tritrebia sp.
Baris concolor Le Conte
Le Conte sp.
Valtomerus prob. quadrinotatus

Diptera 91 .4 cc

Simuliidae
Simulium sp.
Simulium
Simulium

APPENDIX P CONTINUED

Blackfoot at Fletcher Pass Road bridge 7/16/68

Hemiptera 42, (no del. given)

Classocoris clausii Banks
Allograpta sp.
Acronycta pacifica Banks
Neocera clausii Banks
Acronycta sp.

Trichoptera 8,

Hydropsyche sp.
St. acutipila sp.

Ephemeroptera

Ephemerella sp.

Diptera 34,

Simulium sp.
Simulium
Simulium
Simulium

Blackfoot at Fletcher Pass Road bridge 8/2/68

Hemiptera 8, 0.3 cc

Classocoris subulosa Banks
Allograpta sp.
Acronycta pacifica Banks
Neocera sp.
Acronycta sp.

Trichoptera 7, 0.1 cc

Hydropsyche sp.
St. acutipila sp.

Ephemeroptera 1, 1

Ephemerella longimanus (Zanon)

Diptera

Simulium sp. 1

Diptera 10, 1

Simulium
Simulium sp.
Simulium
Simulium sp.

APPENDIX P CONTINUED

BACKWATER RIVER AT FIFTEEN FEET ABOVE CULVERT 1

3/19/71

1 sq. ft. sample from middle of riffle above culvert. 1. Spence, 1430

hemiptera 99 .1	D = 1.38395 D1/D = 0.10277
beetle sp. -43	DIAC = 1.53496 DI = 0.175214
<u>Scaphisoma longimanus</u> -18	Total 86
<u>Scaphisoma</u> sp. -25	
ecoptera 166 .4	

<u>Memura</u> sp. prob. <u>hesaneta</u> -66	
<u>Cynipis</u> sp. -11	
<u>Triclistus</u> <u>caerulea</u> -1	
<u>Chrysomelid</u> <u>viridis</u> -2	
<u>Chrysomelid</u> sp. -48	D = 2.83047 D1/D = 0.355177
<u>Memura</u> <u>cinclipes</u> -17	DIAC = 3.45943 DI = 0.21163
<u>Memura</u> sp. -4	
<u>Chrysomelid</u> <u>frontalis</u> -2	
<u>Chrysomelid</u> <u>brunneicollis</u> -5	
<u>Chrysomelid</u> sp. -3	
<u>Chrysomelid</u> <u>brunneicollis</u> -2	Total 161
Unknown <u>Pisicoptera</u> -2	

ichneumonids 36 .1	
<u>Myrmica</u> <u>scabroscus</u> -1	D = 1.66666 D1/D = 1.569689
<u>Hydrophilus</u> sp. -21	DIAC = 2.58496 DI = 0.451104
<u>Hydrophilus</u> sp. -2	
<u>Hydrophilus</u> sp. -30	Total 37

ptera 203 .2	
<u>Cardinalis</u> sp. -3	
<u>Hydrophilus</u> sp. -1	
<u>Hydrophilus</u> sp. -97	
<u>Hydrophilus</u> sp. -9	
<u>Hydrophilus</u> sp. -1	D = 2.15534 D1/D = 0.234755
<u>Hydrophilus</u> sp. -1	DIAC = 3.52496 DI = 0.432241
<u>Hydrophilus</u> sp. -14	
<u>Hydrophilus</u> sp. -3	
<u>Hydrophilus</u> sp. -2	
<u>Hydrophilus</u> sp. -11	
<u>Hydrophilus</u> sp. -1	Total 182

Original Total 210
Final Total 202

Station D.I.

D = 2.75111 D1/D = 1.334923
DIAC = 3.45943 DI = 0.21163

BACKWATER RIVER AT FIFTEEN FEET ABOVE CULVERT 1

3/19/71

1 sq. ft. sample from center of riffle 50' above culvert. 1. Spence.

<u>Sphenoptera</u> 29 .1	
<u>Scaphisoma</u> sp. -17	
<u>Scaphisoma</u> <u>longimanus</u> -2	
<u>Scaphisoma</u> sp. -2	Total 21

<u>Pisicoptera</u> 62 0.2	
<u>Memura</u> sp. prob. <u>hesaneta</u> -30	
<u>Memura</u> <u>cinclipes</u> -12	
<u>Chrysomelid</u> sp. -11	
<u>Cynipis</u> sp. -7	
<u>Chrysomelid</u> sp. -1	
<u>Chrysomelid</u> <u>subulosa</u> -1	
<u>Chrysomelid</u> <u>brunneicollis</u> -1	
<u>Chrysomelid</u> sp. -1	Total 64

<u>Trichoptera</u> 8 .1	
<u>Hydrophilus</u> sp. -5	
<u>Hydrophilus</u> sp. -2	
<u>Hydrophilus</u> sp. -1	Total 8

<u>Diptera</u> 111 0.1	
<u>Hydrophilus</u> sp. 3	11
<u>Hydrophilus</u> sp. -3	3
<u>Hydrophilus</u> sp. -1	1
<u>Hydrophilus</u> sp. -1	1
<u>Hydrophilus</u> sp. -11	11
<u>Hydrophilus</u> sp. -79	Total 109

Original total 210
Final Total 202

BACKWATER RIVER AT FIFTEEN FEET ABOVE CULVERT 1

3-19-71

1 sq. ft. sample from center of riffle, 75 ft. above culvert. 1. Spence.

<u>Sphenoptera</u> 98 0.1	
<u>Cynipis</u> sp. -16	
<u>Scaphisoma</u> sp. -17	
<u>Scaphisoma</u> <u>longimanus</u> -4	Total 67

<u>Pisicoptera</u> 149 0.5	
<u>Chrysomelid</u> <u>caerulea</u> or <u>curvata</u> -2	
<u>Memura</u> <u>cinclipes</u> -20	
<u>Memura</u> sp. prob. <u>hesaneta</u> -103	
<u>Chrysomelid</u> sp. -17	
<u>Chrysomelid</u> sp. -10	
<u>Chrysomelid</u> sp. -20	
<u>Chrysomelid</u> sp. -2	
<u>Chrysomelid</u> <u>subulosa</u> -1	
<u>Chrysomelid</u> <u>brunneicollis</u> -1	
<u>Chrysomelid</u> <u>brunneicollis</u> -1	Total 211
Unknown small <u>Pisicoptera</u> -1	

<u>Trichoptera</u> 25 0.2	
<u>Chrysomelid</u> sp. -6	
<u>Chrysomelid</u> <u>caerulea</u> -3	
<u>Chrysomelid</u> sp. -9	
<u>Chrysomelid</u> <u>brunneicollis</u> -1	
<u>Chrysomelid</u> sp. -1	
<u>Chrysomelid</u> <u>brunneicollis</u> -1	Total 23

<u>Diptera</u> 105 0.1	
<u>Chrysomelid</u> sp. -12	
<u>Chrysomelid</u> sp. -10	
<u>Chrysomelid</u> sp. -16	
<u>Chrysomelid</u> sp. -1	
<u>Chrysomelid</u> sp. -9	
<u>Chrysomelid</u> sp. -1	
<u>Chrysomelid</u> sp. -19	Total 99

Original Total 210
Final Total 202
199

APPENDIX P CONTINUED

BLACKFOOT RIVER AT FLESHER PASS ROAD BRIDGE #1

1 sq. ft. 10-19-71
1245 L. Spence

EPHEMEROPTERA 22 T			
<u>Baetis</u> sp.	16		
<u>Glyptotendipes</u> sp.	5	Total 21	
PLECOPTERA 130 0.2			
<u>Nemoura glaucoptera</u>	21		
<u>Capnia</u> sp.	2		
<u>Leuctra</u> sp.	20		
<u>Agnetis</u> sp.	5		
<u>Acroneuria</u> sp.	79		
<u>Alloperla</u> sp.	2	Total 131	
<u>Paraperla frontalis</u>			
TRICHOPTERA 49 0.1			
<u>Hydropsyche</u> sp.	13		
<u>Brachycentrus</u> sp.	34		
<u>Stenonema</u> acropodae or var	1		
<u>Limnephilia</u> #2	1	Total 49	
DIPTERA 161 0.1			
<u>Antocha</u> sp.	88		
<u>Paratendipes</u> sp.	64		
<u>Tanypterus</u> sp.	4		
<u>Orthocentrus</u>	2	Total 158	
Original Total 362	Final Total 359		

BLACKFOOT RIVER AT FLESHER PASS ROAD BRIDGE #2

1 sq. ft. 10-19-71
L. Spence

EPHEMEROPTERA 60 T			
<u>Baetis</u> sp.	40		
<u>Glyptotendipes</u> sp.	15	Total 55	
PLECOPTERA 145 0.6			
<u>Nemoura glaucoptera</u>	34		
<u>Capnia</u> sp.	6		
<u>Agnetis</u> sp.	11		
<u>Acroneuria</u> sp.	2		
<u>Alloperla</u> sp.	2		
<u>Paraperla frontalis</u>	2		
<u>Alloperla</u> sp.	81	Total 144	
TRICHOPTERA 54 0.1			
<u>Hydropsyche</u> sp.	14		
<u>Stenonema</u> acropodae or var	1		
<u>Limnephilia</u> #2	1		
<u>Stenonema</u> acropodae or var	2		
<u>Brachycentrus</u> sp.	33		
<u>Limnephilia</u> #2	2		
<u>Hydropsyche</u> sp.	1	Total 54	
DIPTERA 137 0.2			
<u>Antocha</u> sp.	72		
<u>Paratendipes</u> sp.	59		
<u>Orthocentrus</u>	3		
<u>Tanypterus</u> sp.	1	Total 135	
UNKNOWN 2 T			
<u>Nematode</u>	2	Total 2	
Original Total 398	Final Total 390		

BLACKFOOT RIVER AT FLESHER PASS ROAD BRIDGE #3

1 sq. ft. 10-19-71
1300 L. Spence

EPHEMEROPTERA 169 T			
<u>Baetis</u> sp.	101		
<u>Glyptotendipes</u> sp.	37		
<u>Stenonema</u> acropodae or var	3	Total 141	
PLECOPTERA			
<u>Nemoura glaucoptera</u>	26		
<u>Capnia</u> sp.	20		
<u>Leuctra</u> sp.	15		
<u>Agnetis</u> sp.	1		
<u>Acroneuria</u> sp.	4		
<u>Alloperla</u> sp.	2		
<u>Paraperla frontalis</u>	2		
<u>Alloperla</u> sp.	4		
<u>Alloperla</u> sp.	83	Total 160	
TRICHOPTERA 45 0.1			
<u>Hydropsyche</u> sp.	3		
<u>Stenonema</u> acropodae or var	1		
<u>Limnephilia</u> #2	1		
<u>Stenonema</u> acropodae or var	2		
<u>Brachycentrus</u> sp.	1		
<u>Limnephilia</u> #2	35	Total 45	
DIPTERA 313 0.3			
<u>Antocha</u> sp.	5		
<u>Paratendipes</u> sp.	4		
<u>Orthocentrus</u>	101		
<u>Tanypterus</u> sp.	1		
<u>Tanypterus</u> sp.	85		
<u>Orthocentrus</u>	4		
<u>Orthocentrus</u>	27	Total 317	
ANNELEDA 2 T			
<u>Oligochaeta</u>	2	Total 2	
Original Total 694	Final Total 665		

Enumerated and Identified by
/s/ Patricia Ann Mitchell
Department Zoology
University of Alberta

/s/ Patricia Ann Mitchell

APPENDIX P CONTINUED

ALICE CASEY & DAVE NATIONAL FOREST MOUNTAIN 41
8-24-71

1 sq. ft. sample L. Spence

ALICE CASEY & DAVE NATIONAL FOREST MOUNTAIN 42
8-24-71

1 sq. ft. L. Spence

EPHEDRUMPTERA 156 0.5 cc	
Smorua decurtatus	16
Steth. sp.	17
Chrysomelid sp.	110
Chrysomelid doddsi	3
Chrysomelid coloradensis	7
Meloidae sp.	1
Total	154

PLECOPTERA 44 T	
Neuroptera breviscula	45
Alloptera sp. or Arthropoda sp.	1
Total	46

TRICHOPTERA 6 T	
No trichoptera	

DIPTERA	
Orthocentrus sp. 5 (all same species)	6
Total	6

ANNELEIDA 8 T	
Annelida	8
Total	8

Original Total 214 Final Total 214

EPHEDRUMPTERA 378 1.0	
Steth. sp.	73
Chrysomelid sp.	259
Chrysomelid decurtatus	13
Chrysomelid doddsi	13
Chrysomelid coloradensis	13
Chrysomelid breviscula	1
Meloidae sp.	2
Total	368

PLECOPTERA 76 0.1	
Neuroptera breviscula	77
Smorua cinctipes	1
Arthropoda sp. (too small to identify further)	2
Total	82

TRICHOPTERA 0	
Hydroptila sp.	1
Total	1

MEPTERA 15 T	
Orthocentrus sp. 1	1
Orthocentrus sp. 5	11
Neuroptera sp.	1
Hydroptila sp. (probably)	2
Total	15

SEMITATA 10 T	
Annelida - Oligochaeta	10
Total	10

Original Total 479 Final Total 479

Identified and Enumerated by
Patricia Ann Mitchell
MS Univ. of Montana, Missoula, 1968

Patricia Ann Mitchell
January 22, 1973

ALICE CASEY & DAVE NATIONAL FOREST MOUNTAIN 41
8-24-71

D = 1.171 DIAM = 1.171
DIAM = 2.1593 DIAM = 1.171

EPHEDRUMPTERA 399 1.6	
Steth. sp.	174
Smorua cinctipes	81
Chrysomelid sp.	61
Chrysomelid doddsi	1
Chrysomelid coloradensis	13
Chrysomelid breviscula	17
Meloidae sp.	1
Total	364

PLECOPTERA 114 0.1	
Neuroptera cinctipes	23
Neuroptera breviscula	45
Arthropoda sp.	2
Arthropoda sp.	10
Unknown Plecoptera (too small to identify)	1
Total	114

TRICHOPTERA 4 T	
Classonema sp.	2
Psychomyidae	2
Total	4

DIPTERA 49 T	
Neuroptera sp.	10
Orthocentrus sp. 1	7
Orthocentrus sp. 5	5
Orthocentrus sp.	3
Hydroptila sp.	24
Total	49

COLEOPTERA 3 T	
Zelotes sp. adult 1, larva 1	2
Chrysomelid sp. adult	1
Total	3

ANNELEIDA 20 T	
Annelida - Oligochaeta	20
Total	20

Original Total 589 Final Total 581

Identified and Enumerated by
Patricia Ann Mitchell
MS, 1968, University of Montana

Dept. of Entomology
University of Alberta, Edmonton

January 28, 1973

Patricia Ann Mitchell

Station D.I.

D = 2.04755 DIAM = 2.04755
DIAM = 4.04755 DIAM = 2.04755

Identified and Enumerated by
Patricia Ann Mitchell
MS Univ. of Montana, Missoula, 1968
Dept. of Entomology
University of Alberta
Edmonton, Alberta, Canada
January 18, 1973

A-91

Patricia Ann Mitchell

Sample #2 lost in Helena lab.

- ALICE 2456 at HIGH, Y 200, BRIDGE #3

3/19/11

1 sq. ft. sample from center of riffle 75' above bridge. L.Spence.

Ephemeroptera 565 1.1

Total 26b

Ephezerella infrequens 97
Ephezerella doddsi 9
Ephezerella sp. 2
✓ Ephezerella sp. too small to identify 29
Rhithrogena sp. 13
Speoron longicarinus 14
Cinygula spp. 172
Baetis sp. 173
Paraleptophlebia sp. 14

Total 525

Total 2005

Plecoptera	525	1.0
------------	-----	-----

Total 13%

Total 540

Trichoptera	320	0.6
-------------	-----	-----

Total 270

Total 312

Diptera	396	1.5
---------	-----	-----

Original Total 902
Final Total ~~336~~
117

Pericoma sp. 45
Antocha sp. 29
Hexatoma sp. 2
Simuliidae 2

Total 347

637

$$D = 4.1217 \quad D(11) = 0.2023$$

$$D(12) = 5.35755 \quad D(13) = 0.239657$$

<i>Cardiocladius</i> sp.	62
<i>Hydrobaenus</i> sp. 1	63
<i>Hydrobaenus</i> sp. 2	20
<i>Hydrobaenus</i> sp. 3	49
<i>Metriocnemus</i> sp.	36
<i>Dianesa</i> sp. larvae	67
<i>Dianesa</i> sp. pupae	51
<i>Brillia</i> sp. 1	
<i>Pentaneura</i> sp. 1	
Other Tenthredinidae	26

Original total 1812
Final total ~~1803~~
1744

APPENDIX P CONTINUED

ALICE CRICK AT HIGHTWAY 200 Bridge #1
1 so. ft. 8-24-71 L. Spence

ALICE CRICK AT HIGHTWAY 200 Bridge #1
1 so. ft. 8-24-71 L. Spence

ALICE CRICK AT HIGHTWAY 200 Bridge #3
1 so. ft. 8-24-71 L. Spence

EPHEDROPTERA	285	0.2		
Baetis sp.	13			
Chironomus sp.	73			
Chironomus sp.	25	D = 2.24656 DMIN = 0.132387		
Paratrichoptera sp.	7	DIAX = 3.45943 R = 0.36469		
Chironomus sp.	10			
Chironomus sp.	2			
Chironomus sp.	9			
Chironomus sp.	11			
Chironomus sp.	4			
Chironomus sp.	1			
Chironomus sp.	1	Total 275		

PLECOPTERA	120	0.5		
Amphipoda sp.	13			
Amphipoda sp.	73	D = 1.55081 DMIN = 0.192418		
Amphipoda sp.	5	DIAX = 3.16993 R = 0.843112		
Amphipoda sp.	4			
Amphipoda sp.	7	Total 126		

TRICHOPTERA	102	0.5		
Hydropsyche sp.	39			
Hydropsyche sp.	32	D = 2.39339 DMIN = 0.239339		
Hydropsyche sp.	10	DIAX = 3.78844 R = 0.397843		
Hydropsyche sp.	7			
Hydropsyche sp.	1			
Hydropsyche sp.	1			
Hydropsyche sp.	2			
Hydropsyche sp.	1	Total 99		

DIPTERA	133	0.2		
Beetles sp.	2			
Beetles sp.	15			
Beetles sp.	41			
Beetles sp.	31	D = 2.89884 DMIN = 0.194938		
Beetles sp.	9	DIAX = 3.78844 R = 0.397843		
Beetles sp.	1			
Beetles sp.	1			
Beetles sp.	1			
Beetles sp.	1	Total 154		

COLEOPTERA	39	0.1		
Chironomus sp.	25	D = 1.39139 DMIN = 0.08405		
Chironomus sp.	3	DIAX = 2.78844 R = 0.397843		
Chironomus sp.	9			
Chironomus sp.	1			
Chironomus sp.	1	Total 50		

HYMENOPTERA	2	0.2		
Beetles sp.	2	Total 2		
Cottus 1	1	Original Total 672 Final Total 704		

EPHEDROPTERA	248	0.1		
Baetis sp.	80			
Chironomus sp.	112			
Chironomus sp.	1			
Chironomus sp.	12			
Chironomus sp.	9			
Chironomus sp.	34			
Chironomus sp.	1			
Chironomus sp.	1	Total 260		

PLECOPTERA	187	0.2		
Amphipoda sp.	13			
Amphipoda sp.	5			
Amphipoda sp.	100			
Amphipoda sp.	17			
Amphipoda sp.	2			
Amphipoda sp.	1	Total 145		

TRICHOPTERA	248	0.1		
Hydropsyche sp.	111			
Hydropsyche sp.	73			
Hydropsyche sp.	24			
Hydropsyche sp.	5			
Hydropsyche sp.	5			
Hydropsyche sp.	33			
Hydropsyche sp.	1			
Hydropsyche sp.	1			
Hydropsyche sp.	20	Total 251		

DIPTERA	287	0.5		
Beetles sp.	106			
Beetles sp.	4			
Beetles sp.	6			
Beetles sp.	2			
Beetles sp.	58			
Beetles sp.	1			
Beetles sp.	58			
Beetles sp.	4			
Beetles sp.	2			
Beetles sp.	45			
Beetles sp.	17			
Beetles sp.	4			
Beetles sp.	27	Total 338		

COLEOPTERA	94	0.1		
Chironomus sp.	19			
Chironomus sp.	13			
Chironomus sp.	4	Total 50		

1 species - pale head, mentum

EPHEDROPTERA	293	0.1		
Baetis sp.	55			
Chironomus sp.	159			
Chironomus sp.	7			
Chironomus sp.	15			
Chironomus sp.	33			
Chironomus sp.	4			
Chironomus sp.	9			
Chironomus sp.	1	Total 308		

PLECOPTERA	163	0.1		
Amphipoda sp.	41			
Amphipoda sp.	83			
Amphipoda sp.	2			
Amphipoda sp.	1			
Amphipoda sp.	15			
Amphipoda sp.	1	Total 143		

TRICHOPTERA	215	0.2		
Hydropsyche sp.	133			
Hydropsyche sp.	17			
Hydropsyche sp.	2			
Hydropsyche sp.	1			
Hydropsyche sp.	5			
Hydropsyche sp.	24			
Hydropsyche sp.	2			
Hydropsyche sp.	16	Total 213		

DIPTERA	156	0.4		
Beetles sp.	3			
Beetles sp.	41			
Beetles sp.	3			
Beetles sp.	7			
Beetles sp.	5			
Beetles sp.	6			
Beetles sp.	1			
Beetles sp.	2			
Beetles sp.	22			
Beetles sp.	1	Total 213		

COLEOPTERA	86	0.1		
Chironomus sp.	9			
Chironomus sp.	3			
Chironomus sp.	4			
Chironomus sp.	1	Total 23		

Original Total: 914 Final Total: 964

*Appears to be 1 species - pale head, light mentum

APPENDIX P CONTINUED

ALICE CREEK AT HIGHWAY 200 #1

10-19-71

1 sq. ft.

1315

L. Spence

EPHEMEROPTERA 290 0.5

Heatis spp. 69
Clavicornia sp. 111
Elithronema sp. 21
Stenonema sp. 10
E. infrequens 97
E. libialis 12
Paraleptocolebia sp. 19

Total 299

PLECOPTERA 145 0.5

Nemoura cinclipes 15
Gamla sp. 12
Leuctra sp. 8
Alloperla sp. 91
Aronia pacifica 4
Aronia californica 1

Total 131

TRICHOPTERA 269 0.9

Hydropsyche sp. 146
Arctopsyche grandis 7
Glossopsyche sp. 64
Brachycentrus sp. 19
Hydropsyche sp. 5
M. yemalis 2
 pupae (pupae usually are same as above larvae)

Total 244

DIPTERA 465 T + 42

Hemiteles sp. 5
Antocha sp. 131
Pericoma sp. 152
 Orthocid #2 (one easily recognizable species that lives in the blue-green like *Hemiteles*)
 Other Orthocidinae 40
 Tanytarsini 1
 Brullia sp. 1

Total 492

COLEOPTERA 73 0.1

Oligoneurus sp. 38
Stenelmis sp. 18
Stenelmis sp. 17

Total 73

ANNELIDA 2 T

Oligochaeta - family Naididae 2
 Original Total 1286 Final Total 1241

ALICE CREEK AT HIGHWAY 200 #2

10-19-71

1 sq. ft.

1330

L. Spence

EPHEMEROPTERA 210 0.4

Heatis spp. 69
Clavicornia sp. 51
Elithronema sp. 2
Stenonema sp. 10
E. infrequens 61
E. libialis 8
Paraleptocolebia sp. 5

Total 207

PLECOPTERA 110 0.1

Nemoura cinclipes 56
Gamla sp. 5
Leuctra sp. 15
Aronia sp. (small) 1
Alloperla sp. 26
Aronia sp. 1

Total 104

TRICHOPTERA 318 1.9

Hydropsyche sp. 158
Arctopsyche grandis 18
Glossopsyche sp. 37
Brachycentrus sp. 56
Hydropsyche sp. 33
M. yemalis 1
Chamaea sp. 1
Limnephilidae sp. 2 9
 pupae 3

Total 316

DIPTERA 411 0.2 + 60

Hemiteles sp. 3
Antocha sp. 129
Pericoma sp. 104
 Orthocid #2 137
 other orthocidinae 17
 Brullia sp. 1
 Simuliidae 2
 Naididae 1

Total 458

COLEOPTERA 38 T

Oligoneurus sp. 30
Stenelmis sp. 5
Stenelmis sp. 3

Total 38

UNKNOWN 1 T

Nematode 1

Total 1

Original Total 1148

Final Total 1124

ALICE CREEK AT HIGHWAY 200 #3

10-19-71

1 sq. ft.

275 0.5

L. Spence

EPHEMEROPTERA 275 0.5

Heatis spp. (2 species present) 35
Clavicornia sp. 106
Elithronema sp. 7
Stenonema sp. 11
E. infrequens 26
E. libialis 1
Paraleptocolebia sp. 12

Total 269

PLECOPTERA 119 0.3

Nemoura cinclipes 6
Gamla sp. 8
Leuctra sp. 20
Aronia sp. (small) 4
Alloperla sp. (prob. pacifica) 3
Aronia sp. 69
 Small Chloroperlidae (maybe *paraperla*) 5
Arctopsyche - too small to be sure

Total 115

TRICHOPTERA 344 0.8

Hydropsyche sp. 116
Arctopsyche grandis 9
Glossopsyche sp. 136
Brachycentrus sp. 5
Hydropsyche sp. 12
M. yemalis 4
Limnephilidae #1 2
 pupae 2

Total 335

DIPTERA 700 0.2 + 107

Hemiteles sp. 1
Antocha sp. 158
Pericoma sp. 4
 Orthocid #2 221
 other orthocid chironomidae 360 (including those in *Orthocid*)
 Brullia sp. 3
Stenelmis sp. 2
 Tanytarsini 1

Total 810

COLEOPTERA 60 T

Oligoneurus sp. 39
Stenelmis sp. 6
Stenelmis sp. 15

Total 60

OTHER

1 T

Original Total 1606

Final Total 1589

/s/ Patricia Ann Mitchell

/s/ Patricia Ann Mitchell
 May 9, 1974

APPENDIX P CONTINUED

Blackfoot River at Howe's Creek Road Bridge 7/15/63

Plecoptera 53 .7 cc

Glaucenia sabinea Banks
Acronema pacifica Banks
Kathoperla sp.
Acronema theodora Kuehnen and Claassen
Enocla cinctipes Banks
Alloperla sp.

Trichoptera 155 1.3 cc

Brachycentrus sp.
Hyacophila acropodes or ves
Hyacophila angelita Janks
Hydrophoca sp.
Limnephila (keys to Astenophylax in key, but isn't.)
Psychovia flavida (hagen) or lurida (Ross)
Anagrus sp.
Lepidostoma sp.
Gehrotrichia sp.

Ephemeroptera 503 1.4 cc

Paraletochlebia vaciva ?
Ephemerella fl-villina McDunnough
Ephemerella tisi-lis McDunnough
Stocrus lili-viridis (Luton)
Gastis spp.
Cinygula sp.

Coleoptera 124 .05 cc

Talitrea sp.
Lara sp.
Parpus sp. prob. concolor
Optiocervus sp.
Pterilampus sp. prob. corpulentus
Chrysomelinae adult, prob. not aquatic

Diptera 813 2.0 cc

Hexatoma sp.
Antocha sp.
Simulidae
Phabiodontia sp. ?
Tendipedidae
Pentecura sp.
Tanytarsus sp.

Station D.I.

Blackfoot River at Hogan Creek Road Bridge 4/29/65

Plecoptera 65. 1.10 cc

Acronurus pacifica Banks
Classenia sabulosa Banks
Nemoura bicarata Hicker
Nemoura distinctus Banks
Illoperla sp.
Cynia distincta
Cynia sp. prob. elongata or confusa
Isoperla sp.

Trichoptera 42, 0.5 cc

Rhyacophila acropedee or vao
Glossosoma sp.
Lepidostoma sp.
Hydropsyche sp.
Brachycentrus sp.
Limnephilidae

Ephemeroptera 254, 1.4 cc

Isotia spp.
Ephemerella infrequens McDunnough
Glyptotendipes sp.
Ephemerus longicauda (Sutcliffe)
Ephemerella grandis angens McDunnough
Paraleptochleba vacillans Sutcliffe ?
Rhyacophila sp. (too small for proper identification)
Ephemerella sp. (too small for proper identification)

Coleoptera 14. 106 cc

Harpus concolor LeJoute
Opiasorvus prob. quadrinaculatus
Heterolophus prob. corpulentus

Diptera 91, 0.1 cc

Pericoma sp.
Simalilloae
Tendipedidae
Metriocnemus sp.
Carolecladius sp.

✓ Shells of *Stagnicola palustris* spicile 10. 0.1 cc

Station D.I.

Blackfoot River at near lower end of the ...

Plecoptera 49

Hasteria sp.
Arcynopteryx ovellata Frison (or curvata)
Aronauria pacifica Banks
Ceomys dist. not a or confusa

Trichoptera 399

Rhyacophila acronodes Banks or var. alina
Brachycentrus sp.
Lepidostoma sp.
Hydroscapha sp.
Manaphilidae

Ephemeroptera 32

Echinorella grandis ingens 'cDannough
Paraleptoclella sp., prob. vaciva gator
Clypeura sp.
Echinorella sp.

Coleoptera 50

Heterlirmius compulentes (Le Conte)
Optioservus sp. prob. quadriraculatus
Zaitzevia no.

Diptera 44

Antocha sp.
Rhabdonastix sp.
Pericoma sp.
Gnathochorax sp.
Simuliidae

Station D.I.

D = 1.49871 DelIN = 7.33555E-2
D1FX = 2.32193 ? = 0.36975

0738855

FILE TOTAL 3091

APPENDIX P CONTINUED

BLACKFOOT RIVER BETWEEN LANDER'S FORK AND LINCOLN #2

1 sq. ft. L. Spence

BLACKFOOT RIVER BETWEEN LANDER'S FORK AND LINCOLN #2

1 sq. ft. 12 Spores

Blackfoot River between Lander's Ford and Little Falls, ID.

1. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 840. 841. 842. 843. 844. 845. 8

^a L. 2000.

Species	Count	Percentage
<i>Bastie</i> sp.	88	
<i>Clypeola</i> sp.	41	
<i>Stomus depressus</i>	50	
<i>Stomarella socialis</i>	96	
<i>Stomarella coloradensis</i>	8	
<i>Stomarella tidalia</i>	26	
<i>Stomarella infrenans</i>	2	
<i>Stomarella gracilis lymnae</i>	1	
<i>Stomarella bairdi</i>	1	
<i>Stomarella</i> sp.	151	
Total	464	

PLECOPTERA	54	0.1	
<i>Neoura cinctipes</i>	3		
<i>Pteronarcys badia</i>	1		
<i>Capnia</i> sp. ?	4	D = 1.29781	DNH = 0.3365995
<i>Acronemia pacifica</i>	1	D = 3.16993	H = 0.672841
<i>Oligoneuria subulosa</i>	1		
<i>Alopiura</i> sp.	33		
<i>Perlodina</i> (too small)	5		
<i>Isogenus</i> sp.	3		
		Total	54

TRICHOPTERA	118	0.8	
<u>Rhyacophila acropetala</u>	2		
<u>Glossosoma</u> sp.	40	D = 1.75332	SDIN = 0.23529
<u>Brachycentrus</u> sp.	7	DIAT = 3.0	0.451129
<u>Arctopsycha grandis</u>	5		
<u>Tinodes</u> sp.	14		
✓ <u>Pupae</u>	4	Total	119

DIPTERA 65 0.1		
A-99	<u>Hexatoma</u> sp.	2
	<u>Napididae</u> #2	1
	<u>Paricosia</u> sp.	1
	<u>Simulium</u>	2
	<u>Caricoidia</u> sp.	24
	<u>Metricnemus</u> sp	3
	<u>Tanylarus</u> sp.	2
	<u>Brillia</u> sp.	3
	Unknown Chironomidae	15
		Total 63

COLEOPTERA 9 T 5=1 No diversity Computed
Optionservus sp. lar-1, ad-8 9 Total 9
 Original Total 715, Final Total 709

Enumerated and Identified by
Patricia Ann Mitchell
April 30, 1973

Station D.I.

$D = 3.57293$ $\text{Gilt} = 1.0511$ C
 $D167 = 5.26135$ $Dm = 1.76501$

SPERMATOPHYTES	300	0.4
<u>Beilsia</u> sp.		50
<u>Cloezella</u> sp.		4
<u>Umbrocarina</u> sp.	116	
<u>Scorpus longimanus</u>		2
<u>Scorpus decentius</u>		31
<u>Scorpus Godardii</u>		74
<u>Scorpus trilineatus</u>		7
<u>Scorpus colerudensis</u>		3
<u>Scorpus infrenus</u>		1
<u>Scorpus grande</u>		1
		Total 291

PLECOPTERA 62 0.1		
<u>Nemoura cinctipes</u>	3	
<u>Pteronarcys badia</u>	4	
<u>Agronema pacifica</u>	2	
<u>Arsenoperla</u> sp.	1	
<u>Alloperla</u> sp.	51	Total 61

TRICHOPTERA	68	0.2	
<u>Glossosoma</u> sp.	27		
<u>Hyacophila acropodes</u>	3		
<u>Brachycentrus</u> sp.	1		
<u>Tinodes</u> sp.	5		
<u>Arctopsyche grandis</u>	24		
✓ Pupae	4		Total 64

DIPTERA 23 T.	
Simuliidae	12
<u>Hexatoma</u> sp.	1
<u>Cardiocladius</u> sp.	6
<u>Brillia</u> sp.	1
<u>Mesocneme</u> sp.	2
Unknown Chironomidae	3
	Total 25

COLEOPTERA	3	T	
<u>Optioservus</u> sp. adulte	3		Total 3

Original Total 456 Final Total 444

Identified and Enumerated by
Patricia Ann Mitchell
MS, Univ. of Montana, 1968
April 30, 1973

Patricia Ann Mitchell

EPHAKRON PERA	456	0.3
<u>Bastis</u> sp.	60	
<u>Clypeola</u> sp.	5	
<u>Anthrenus</u> sp.	246	
<u>Ephemerella</u> <u>addol</u>	93	
<u>Ephemerella</u> <u>coloradensis</u>	3	
<u>Ephemerella</u> <u>randis incens</u>	5	
<u>Ephemerella</u> <u>tibialis</u>	5	
<u>Ephemerella</u> <u>infrenans</u>	1	
<u>Eucross</u> <u>decentius</u>	25	Total 443

PLICOPTERA	68	0.1	
<u>Alloperla</u> sp.		61	
<u>Pteronarcetta</u> badia		3	
<u>Acroneuria</u> pacifica		1	
<u>Acroneuria</u> sp. ? (too small)		2	
<u>Hexura</u> clintopes	5		
			Total 72

TRICHOPTERA 116 0.3		
<u>Areopsyche grandis</u>	26	
<u>Hydropsyche</u> sp.	5	
<u>Glossosoma</u> sp.	73	
<u>Hyatrichia acronotus</u> or <u>Var</u>	1	
<u>Brachycentrus</u> sp.	3	
<u>Limnephilidae</u>	1	
Unknown - Psychomyiidae?	1	
✓ Pupae	4	Total 116

DIPTERA	39	0.3
<i>Nematoma</i> sp.		5
<i>Simuliidae</i>		1
<i>Muscidae</i> larva?		1
<i>Cardiocladius</i> sp.		17
<i>Anatrus</i> sp.		5
<i>Brillia</i> sp.		1
<i>Kellicottia</i> sp.		1
<i>Orthocentrus</i> sp. 1 (* <i>Hydrobaenus</i>)		3
		Total 34

COLEOPTERA	6	†		
<i>Ootiosenus</i> sp.			6	Total 6
NEMATODA	3	?		
Nematoda			3	Total 3

Original Total 688 Final Total 672

BLACKFOOT RIVER BETWEEN LANDER'S FORK AND LINCOLN #2
10-26-71
1 sq. ft. _____ L. Spence

620	0.9	
<i>Glyptotendipes</i> sp.	76	D = 1.58868 EFIN = 0.034997
<i>Chironomus</i> spp.(prob.more than 1)	416	DIFF = 2.58496 N = 0.2079
<i>Beetia</i> sp.	32	
<i>Epheuerella doddsi</i>	57	
<i>Epheuerella infrequens</i>	34	Total 615

OTRA	111	0.6	
<u>Nemora sinicae</u>	4		
<u>Cepia</u> sp.	14		
<u>Brachyura</u> sp.	21		
<u>Isopoda</u> sp.	3	D = 2.2064	BIH = 0.52612
<u>Parlodica</u>	1	IAH = 3.9039	D = 2.8457
<u>Arenaria californica</u>	2		
<u>Arenaria shondra</u> (probably)	2		
<u>Alloperla</u> sp.	63	Total	109

D = 1.23745 DMN = 9.23440E-2 = .092443
 DMAX = 3.16993 R = 8.611774
 KOTTERA 162 0.5
Arctopernha grandis 10
Glossosoma sp. 109
Hydroporche sp. 50
 Limnephilidae Genus 1 3 Total 172

27	0.3	
Simuliidae		3
Hexatoma spp.		6
Dicranota sp.	1 D = 2.1021	DOIN = 3.672346
Antocha sp.	1 DIAC = 3.45943	T = 2.453383
Empididae Genus 1		1
Perisoma sp.		1
Chironomidae		12
Chironomus sp.		1
Chironomidae unknowns		2
	Total 27	

	D = 1.37695	DNIN = 1.37295
OPTERA 2 T	DNAX = 1.58496	D = 1.
Ontosarvus sp. adulta	2	Total 2

LIDA 3 T		
Oligochaetes	3	Total 8

Station D.I.

SPRINGCROFTERA	565	1.2	
<i>Bastia</i> sp.		42	
<i>Cinyxula</i> sp.		67	
<i>Rhithrotrama</i> sp.		342	
<i>Schemerella</i> <i>doddsi</i>		55	
<i>Schemerella</i> <i>grandis</i> <i>insens</i>		2	
<i>Schemerella</i> <i>infrequens</i>		67	
			Total 575

PLECOPTERA	65	0.7	
<u>Pteronarcys badia</u>			2
<u>Speobytena</u> sp.			17
<u>Capnia</u> sp.			1
<u>Agromyza curvata</u> or <u>parallel</u>			1
<u>Agromyza</u> <u>sinuata</u>			1
<u>Agromyza</u> sp.			1
✓ <u>Parolides</u> (too small)			1
<u>Agrionema californica</u>			1
<u>Agrionema pacifica</u>			4
<u>Classenia rubra</u>			1
<u>Alloperla</u> sp.			21
<u>Centella</u>			2
			61

TRICHOPTERA	292	0.5	
<u>Glossosoma</u> sp.			186
<u>Arctopsyche grandis</u>			21
<u>Hydropsyche</u> sp.			84
<u>Chracophila acropodae</u> or <u>var</u>			1
			Total 292

Hexatoma sp.	10	
Antocha sp.	2	
Eupidiidae #2	5	
Simuliidae	1	
Pericoma sp.	25	Total 43

COLEOPTERA	1	7		
<u>Zaitsevia</u> sp. lar.			1	Total 1
✓ NEMATODA	1	7		
Nematoda			1	Total 1

Original Total ⁹⁶⁶~~967~~ Final Total 973

Identified and Enumerated by
Patricia Ann Mitchell, M.S.
Department of Zoology
University of Alberta
Edmonton, Alberta, Canada

Patricia Ann McShell

EPHEMEROPTERA	582	1.0	
<i>Pactis</i> sp.		119	
<i>Cinygmula</i> sp.		78	
<i>Rithrogena</i> sp.		257	
<i>Sphenerehella</i> <u>doddsi</u>		42	
<i>Sphenerehella</i> <u>grandidi ingens</u>		2	
<i>Sphenerehella</i> <u>infraguana</u>		85	Total 583

PLECOPTERA		84	0.2
	<u>Brachyptera</u> sp.	4	
	<u>Gania</u> sp.	12	
	<u>Nesoura clinctipes</u>	10	
	<u>Arcynopteryx curvata</u> or <u>parallela</u>	1	
	<u>Isonurus festalis</u>	2	
	<u>Isonurus</u> sp.	3	
	<u>Acronura pacifica</u>	1	
	<u>Classania subulosa</u>	1	
	<u>Alloperla</u> sp.	45	
✓	Plecoptera too small to identify	4	Total 83

XENICTOPTERA		515	1.1
<i>Glossosoma</i> sp.		331	
<i>Archipsyche grandis</i>		37	
<i>Hydroxysyche</i> sp.		142	
<i>Hydropsyche</i> <i>acropedes</i> of <i>var</i>		1	
<i>Hydropsyche</i> sp.		1	
<i>Rhyacops</i> sp.		1	
<i>Limnephilidae</i> Genus 2		2	
			Total 516

DIPTERA		54	0.2	
<u>Hexatoma</u> spp.			6	
<u>Empididae</u> #4			1	
<u>Empididae</u> #5			1	
<u>Pericoma</u> sp.			25	
<u>Cardiophilus</u> sp.			1	
Unknown Orthoclad Chironomidae		20		Total 5

COLEOPTERA	2	T	
<u>Naryus concolor</u> adult	1		
<u>Oxiosernus</u> sp. lar.	1		Total 2

✓ NEMATODA	3	T	
Nematoda		3	Total 3
✓ COOTUS SP.	3		
		3	Total 3
Original Total	1243	Final Total	1244

Enumerated and Identified by
Patricia Ann Mitchell
July 27, 1973
- Patricia Ann Mitchell

APPENDIX D CONTINUED

POORMAN CREEK #1
8-23-71
Spence 1 sq. ft.

[illegible]

EDOPTERA 159 0.9 ee
Nemoura cinctipes 135 D = 1.88555 DMW = 0.104372
Acroneuria pacifica 14 DMW = 2.55496 W = 0.61654
Alloperla sp. 3
Eucanopsis bravipecta (probably) 9 Total 161

41	<u>Arctophila grandis</u>	
36	<u>Diodes</u> sp. (Psychomyiidae)	D = 1.33205 EW11 = 1.25027
4	<u>Hyacophila acropetor</u> of YMO	DAK = 2.8735 T = 1.33229
1	<u>Hyacophila</u> sp.	
1	<u>Glossosoma</u> sp.	
1	✓Pupa	Total 84

PERA	290	0.1 cc	D ₁ = 7.01119	D ₁ M ₁ = 7.05331E-2 = 0.07331
			DMAX = 3.787044	R = 0.576836
Stenulidae			110	
Rapididae Genus 1			2	
Rapidus sp.			14	
Orthocladus sp. 2			18	
Myrtarus sp.			7	
Cardiocladius sp.			2	
Brillia sp.			69	
Metriocnemus sp.			21	
Unknown Chironomidae			62	
				Total 305

DIPTERA 38 T	
<u>Heterelmis</u> sp. larvae 14, adults 2	16
<u>Otitus</u> sp. larvae 2, adults 3	5
<u>Zaitzeva</u> sp. larvae 1	1
<u>Erychius</u> sp. (Halipidas) larvae	3
	Total 25

CELIDA 6 T		
Oligochaeta	14	Total 14
	Original Total 932	Final Total 914

Identified and Enumerated by
Patricia Ann Mitchell
University of Alberta
March 19, 1973

Station D.I.

EPHEDROPTICIA	281	0.2 cc

<i>Rattia</i> spp.	145
<i>Cinnyula</i> sp.	55
<i>Rhithrona</i> sp.	36
<i>Speorus</i> <i>deceptivus</i>	7
<i>Speorus</i> <i>longinamus</i>	7
<i>Ephemerella</i> <i>doddsi</i>	3
<i>Ephemerella</i> <i>hyatrix</i>	6
<i>Ephemerella</i> <i>grandis</i>	1
<i>Ephemerella</i> <i>tibialis</i>	1
<i>Amelopteryx</i> sp.	4
	Total 266

<u>Nemoura cinetipes</u>	53	
<u>Alloperla</u> sp.	12	
<u>Acronuria pacifica</u>	10	
<u>Bucconossia brevicauda</u> (probably)	1	Total 76

TRICHOPTERA	60	0.4	ee	
<u>Arctopsyche</u>	<u>grandis</u>			38
<u>Procladius</u>	sp.			10
<u>Hydropsyche</u>	<u>seropedes</u> or <u>vao</u>			2
<u>Hydropsyche</u>	sp.			8
<u>Glossosoma</u>	sp.			2
				Total 60

DIPTERA 1065 0.8 cc	
<u>Anopheles</u> sp.	1
<u>Siaulidae</u>	989
<u>Empididae Genus 1</u>	8
<u>Orthocentrus</u> sp. 2	1
<u>Cardinalis</u> sp.	1
<u>Salix</u> sp.	21
<u>Hydrophilus</u> sp.	8
<u>Unknown Chironomidae</u>	29
<u>Pericoma</u> sp.	1
	Total 10

COLLEPTERA	31	7	
<u>Heterischnus</u> sp.		29	
<u>Narceus concolor</u>		2	Total 31

ANNEJIDA 13 T		
Oligochaeta	13	Total 13

Enumerated and Identified by
Patricia Ann Mitchell
University of Alberta, Edmonton
March 20, 1973

POORMAN CREEK #3
8-23-71
1 sq. ft. L. Spence

SPERMATOPHYTES	342	0.2 cc
<u>Betula</u> spp.	187	
<u>Corymbia</u> sp.	61	
<u>Specorus deceptivus</u>	4	
<u>Specorus longimanus</u>	3	
<u>Rhithroena</u> sp.	18	
<u>Paraleptophlebia</u> sp.	6	
<u>Ephemerella</u> spp.	3	
<u>Ephemerella</u> <u>gossii</u>	7	
<u>Ephemerella</u> <u>gracilis</u>	5	
<u>Ephemerella</u> <u>tibialis</u>	7	
<u>Ephemerella</u> <u>brachy</u>	10	

PLECOPTERA	#34	0.5 cc	
<u>Pteronarcissa</u> sp.			2
<u>Acronesia</u> <u>pacifica</u>			34
<u>Nemoura</u> <u>cinctipes</u>			208
<u>Alloperla</u> sp.			13
<u>Isogenus</u> sp.			3
<u>Stenonema</u> <u>brevicauda</u> ?			1
			261

TRICHOPTERA 78 0.3 ec		
<u>Elysiophila</u> sp.	6	
<u>Elysiophila</u> <u>acronodes</u> or <u>yno</u>	8	
<u>Arctopsyche</u> <u>granile</u>	23	
<u>Pinodes</u> sp.	27	
<u>Limnephilidae</u> Genus 2	6	
<u>Hydroptilidae</u>	1	
✓ pupae	2	Total 73

STATION	S16	0.2 ce	
Stimuliidae			200
Nepitidae Genus #1			30
Nepitidae Genus #2			3
Nepitidae Genus #3			1
Anelasma sp.			1
Peridroma sp.			20
Orthocladus sp. 2			20
Grilla sp.			67
Meloidonemus sp.			26
Sisyrtus sp.			8
Caridiodon sp.			2
Pentaneura sp.			1
Unknown Diptera			160
			Total 500

COLEOPTERA 74 ♀	
<u>Marpus</u> sp. larvae	9
<u>Optioservus</u> sp. lar-3, ad-3	6
<u>Zeisleria</u> sp. adult	1
<u>Heterelmis</u> sp. ad-2, lar-50	52
<u>Erychius</u> sp.	3
	Total 71

ANNELEIDA	6	7	
Oligochaeta		16	Total 16

Original Total	1250	Final Total	1254

APPENDIX P CONTINUED

BLACKFOOT RIVER AT DALTON MOUNTAIN ROAD BRIDGE #2
1 sq. ft. left bank 8-20-71 L. Spence

EPHEMEROPTERA 180 0.2			
<i>Baetis</i> spp.	154	D = 1.34794 DMN = 0.102447	
<i>Ephemerella</i> sp.	17	DMAX = 2.68735 R = 0.529147	
<i>Stenonema</i> sp.	1		
<i>Ephemerella</i> sp.	4		
<i>Ephemerella grandis</i>	5	Total 181	
PLECOPTERA 4 T			
<i>Leuctra</i> sp.	1	D = 2.02229 DMN = 0.351178	
<i>Alloperla</i> sp.	1	DMAX = 2.53496 R = 0.324365	
<i>Pteronarcys frontalis</i>	2	Total 4	
TRICHOPTERA 59 T			
<i>Glossosoma</i> sp.	33		
<i>Hydropsyche</i> sp.	2	D = 1.92379 DMN = 0.215031	
<i>Arctopsycha grandis</i>	4	DMAX = 2.56496 R = 0.271955	
<i>Brachycentrus</i> sp.	19		
<i>Limnephilid</i> Genus 2	1	Total 59	
DIPTERA 111 0.2			
<i>Atherix variegata</i>	8	D = 1.74933 DMN = 0.092977	
<i>Simuliidae</i>	99	DMAX = 3.05943 R = 0.537723	
Unknown Chironomidae	4		
<i>Tipulidae</i>	2	Total 113	
COLEOPTERA 2 T			
<i>Zaitsevius</i> sp. adult	1	D = 1.34886 DMN = 0.444193	
<i>Cholesterus</i> sp. larva	1	DMAX = 2.32193 R = 0.572939	
Total 2			
NEMATODES 2 T			
None			
Total 1			
Original Total 359		Final Total 360	

Identified and Enumerated by
Patricia Ann Mitchell
University of Alberta, Edmonton
May 23, 1973

A-106
Patricia Ann Mitchell
station D.T.
D = 0.32391 DMN = 0.37355
DMAX = 0.102447 R = 0.37257

BLACKFOOT RIVER AT DALTON MOUNTAIN ROAD BRIDGE #2
1 sq. ft. right bank 8-20-71 L. Spence

EPHEMEROPTERA 220 0.1			
<i>Baetis</i> spp.	147		
<i>Cinygmula</i> sp.	5		
<i>Stenonema</i> sp.	47		
<i>Ephemerella tibialis</i>	6		
<i>Ephemerella grandis</i>	12		
<i>Ephemerella</i> sp.	5	Total 222	
PLECOPTERA 21 T			
<i>Alloperla</i> sp.	9		
<i>Pteronarcys</i> sp.	6		
<i>Stenonema</i> sp.	1		
<i>Pteronarcys pacifica</i>	1	Total 17	
TRICHOPTERA 103 0.1			
<i>Brachycentrus</i> sp.	60		
<i>Glossosoma</i> sp.	12		
<i>Arctopsycha grandis</i>	17		
<i>Limnephilid</i> Genus 1	1		
<i>Limnephilid</i> Genus 2	4		
<i>Hydropsyche</i> sp.	7	Total 101	
DIPTERA 68 0.1			
<i>Atherix variegata</i>	3		
<i>Hexatoma</i> sp.	1		
<i>Simuliidae</i>	5		
<i>Epididae</i> Genus #1	4		
<i>Epididae</i> Genus #2	1		
<i>Helicoverpa</i> sp.	14		
<i>Cholesterus</i> sp.	1		
<i>Cholesterus</i> sp. (= <i>Cholesterus</i>)	1		
<i>Brachycentrus</i> sp.	1		
Unknown Chironomidae	32	Total 64	
COLEOPTERA 22 T			
<i>Nephus</i> sp. larva	1		
<i>Stenonema</i> sp. larva	1		
<i>Cholesterus</i> sp. lar-12, ad-3	15		
<i>Cholesterus</i> sp. lar-4, ad-1	5	Total 22	
NEMATODES 38 T			
Oligochaeta	38	Total 38	

Original Total 472, Final Total 469

Enumerated and Identified by
Patricia Ann Mitchell
MS, Univ. Montana, 1968
University of Alberta, Edmonton
May 26, 1973
Patricia Ann Mitchell

BLACKFOOT RIVER AT DALTON MOUNTAIN ROAD BRIDGE #3
1 sq. ft. right bank 8-20-71 L. Spence

EPHEMEROPTERA 242 0.2			
<i>Baetis</i> spp.	146		
<i>Ephemerella</i> sp.	68		
<i>Cinygmula</i> sp.	1		
<i>Ephemerella</i> sp.	7		
<i>Ephemerella tibialis</i>	4		
<i>Ephemerella grandis</i>	14	Total 240	
PLECOPTERA 16 T			
<i>Pteronarcys</i> sp.	6		
<i>Alloperla</i> sp.	7		
<i>Stenonema</i> sp.	2		
<i>Pteronarcys pacifica</i>	3	Total 18	
TRICHOPTERA 53 T			
<i>Brachycentrus</i> sp.	27		
<i>Glossosoma</i> sp.	8		
<i>Arctopsycha grandis</i>	8		
<i>Limnephilid</i> Genus 1	3		
<i>Limnephilid</i> Genus 2	3		
<i>Hydropsyche</i> sp.	4	Total 53	
DIPTERA 198 0.2			
<i>Atherix variegata</i>	9		
<i>Simuliidae</i>	118		
<i>Epididae</i> #1	2		
<i>Epididae</i> #2	3		
<i>Cholesterus</i> sp.	16		
<i>Cholesterus</i> sp. lar-9, ad-3	1		
Unknown Chironomidae	28	Total 177	
COLEOPTERA 21 T			
<i>Cholesterus</i> sp. lar-4, ad-1	1		
<i>Cholesterus</i> sp. ad-1	1		
<i>Cholesterus</i> sp. lar-9, ad-3	12	Total 18	
and 3 empty exuviae			
OLIGOCHAETA			
Oligochaeta	19	Total 19	
Original Total 530		Final Total 525	

Identified and Enumerated by
Patricia Ann Mitchell
University of Alberta, Edmonton
May 26, 1973
Patricia Ann Mitchell

APPENDIX P CONTINUED

BLACKFOOT RIVER AT DALTON MOUNTAIN ROAD BRIDGE #1

1 sq. ft. 10-26-71 L. Spence

EPHEMEROPTERA	540	1.2	
<i>Heptagenia</i> spp. (2 species)	76		
<i>Stenonema</i> sp.	39		
<i>Hydropsyche</i> sp.	413		
<i>Hydropsyche</i> grandis	8		
<i>Hydropsyche</i> grandis incana	8		
<i>Hydropsyche</i> incana	8		
Total	552		

PLECOPTERA	125	0.9	
<i>Heurogaster</i> sinuatus	8		
<i>Pteronarcissa</i> badia	22		
<i>Pteronarcissa</i> sp.	2		
<i>Pteronarcissa</i> sp.	2		
<i>Pteronarcissa</i> sp.	4		
<i>Pteronarcissa</i> frontalis	63		
<i>Pteronarcissa</i> sp.	3		
<i>Pteronarcissa</i> sp.	8		
<i>Pteronarcissa</i> pacifica	3		
<i>Pteronarcissa</i> californica	1		
Total	115		

TRICHOPTERA	198	0.7	
<i>Hydropsyche</i> sp.	22		
<i>Hydropsyche</i> grandis	18		
<i>Hydropsyche</i> sp.	6		
<i>Glossosoma</i> sp.	148		
<i>Glossosoma</i> sp.	2		
<i>Glossosoma</i> sp.	1		
Total	197		

DIPTERA	35	0.1	
<i>Atopsyche</i> sp.	5		
<i>Ephydra</i> #1	3		
<i>Atopsyche</i> sp.	23		
<i>Orthoclad</i> Chironomidae	4		
Total	36		

COLEOPTERA	4	T	
<i>Oligoneurus</i> sp.	4		
Total	4		

HEMPTODA	6	T	
<i>Oligoneurus</i> sp.	1		
<i>Hemiptera</i>	4		
Total	5		

ODONATA	1		
Total	1		

Original Total 908

Final Total 909

BLACKFOOT RIVER AT DALTON MOUNTAIN ROAD BRIDGE #2

1 sq. ft. 10-26-71 L. Spence

EPHEMEROPTERA	512	0.5	
<i>Heptagenia</i> spp.	39		
<i>Stenonema</i> sp.	32		
<i>Hydropsyche</i> sp.	421		
<i>Hydropsyche</i> grandis	2		
<i>Hydropsyche</i> grandis incana	4		
<i>Hydropsyche</i> incana	2		
<i>Hydropsyche</i> sp.	12		
Total	514		

PLECOPTERA	122	0.7	
<i>Heurogaster</i> sinuatus	12		
<i>Pteronarcissa</i> badia	13		
<i>Pteronarcissa</i> sp.	2		
<i>Pteronarcissa</i> sp.	1		
<i>Pteronarcissa</i> sp.	6		
<i>Pteronarcissa</i> sp.	2		
<i>Pteronarcissa</i> sp.	19		
<i>Pteronarcissa</i> pacifica	63		
<i>Pteronarcissa</i> californica	1		
Total	118		

TRICHOPTERA	150	1.1	
<i>Hydropsyche</i> sp.	28		
<i>Hydropsyche</i> grandis	39		
<i>Hydropsyche</i> sp.	55		
<i>Glossosoma</i> sp.	18		
<i>Glossosoma</i> sp.	5		
<i>Glossosoma</i> sp.	3		
Total	148		

DIPTERA	17	T	
<i>Atopsyche</i> sp.	7		
<i>Ephydra</i> #1	1		
<i>Atopsyche</i> sp.	3		
<i>Orthoclad</i> Chironomidae	1		
Total	17		

AMPHIBIA	17	T	
<i>Oligoneurus</i> sp.	17		
Total	17		

Original Total 818

Final Total 814

Patricia Ann Mitchell

BLACKFOOT RIVER AT DALTON MOUNTAIN BRIDGE #3

1 sq. ft. 10-26-71 L. Spence

EPHEMEROPTERA	680	0.9	
<i>Heptagenia</i> spp. (2)	35		
<i>Stenonema</i> sp.	39		
<i>Hydropsyche</i> sp.	552		
<i>Hydropsyche</i> grandis	7		
<i>Hydropsyche</i> grandis incana	3		
<i>Hydropsyche</i> incana	3		
<i>Hydropsyche</i> sp.	10		
<i>Hydropsyche</i> sp.	1		
Total	690		

PLECOPTERA	97	0.8	
<i>Heurogaster</i> sinuatus	9		
<i>Pteronarcissa</i> badia	1		
<i>Pteronarcissa</i> sp.	5		
<i>Pteronarcissa</i> sp.	9		
<i>Pteronarcissa</i> pacifica	22		
<i>Pteronarcissa</i> frontalis	1		
<i>Pteronarcissa</i> sp.	48		
Total	95		

TRICHOPTERA	163	1.0	
<i>Hydropsyche</i> sp.	20		
<i>Hydropsyche</i> grandis	35		
<i>Glossosoma</i> sp.	91		
<i>Glossosoma</i> sp.	2		
<i>Glossosoma</i> sp.	6		
Total	150		

DIPTERA	43	0.1	
<i>Atopsyche</i> sp.	19		
<i>Ephydra</i> #1	3		
<i>Atopsyche</i> sp.	10		
<i>Orthoclad</i> Chironomidae	3		
<i>Orthoclad</i> Chironomidae	3		
Total	41		

COLEOPTERA	2	T	
<i>Oligoneurus</i> sp.	1		
<i>Hemiptera</i> sp.	1		
Total	2		

HEMPTODA	5	T	
<i>Oligoneurus</i> sp.	5		
Total	5		

ODONATA	2		
Total	2		

Original Total 992

Final Total 972

Patricia Ann Mitchell

Plecoptera 7 .05 cc

Agreonaria pacifica Banks
Agreonaria californica Fernald
Agreonaria sp. (probably); or *Agreonaria* sp.

APPENDIX P CONTINUED

Trichoptera 6 .1 cc

Station D.I.

Hydropsyche sp. D = 5.96237 D:LN = 7.82575D = 2 = .0772.75
Hydropsyche sp. D:LN = 2.72193 D = 7.53177
Hydropsyche sp.

Ephemeroptera 475 1.9 cc

Ephemerella longicauda (Ston)
Ephemerella sp.
Ephemerella sp. (2 types)
Ephemerella sp. prob. *undulata* (Banks)
Ephemerella (Ston) *undulata* (Banks)
Ephemerella sp.
Ephemerella sp.

Coleoptera 9 7

Orthocentrus sp. prob. *quadriculatus*

Diptera 92 .6 cc

Phorcia sp. (3 spp.) ✓ Aquatic insect 4 spp. .1 cc
Phorcia sp. (3 spp.) ✓ Hemiptera 1 7
Phorcia sp. ✓ Gastropoda 71 7
Phorcia sp.
Hydropsyche sp.

Plecoptera 71

Agreonaria pacifica Banks
Agreonaria californica Banks
Agreonaria sp. (4 spp.)
Agreonaria sp. (4 spp.)
Agreonaria californica Fernald
Agreonaria sp. (4 spp.)
Agreonaria sp. (4 spp.)
Agreonaria sp. (4 spp.)

Trichoptera 139

Hydropsyche sp.
Hydropsyche sp.
Hydropsyche sp.
Hydropsyche sp.
Hydropsyche sp.

Ephemeroptera 401

Ephemerella sp.
Ephemerella sp.
Ephemerella sp.
Ephemerella sp.
Ephemerella sp.

Coleoptera 13

Orthocentrus sp. (4 spp.)
Orthocentrus sp. (4 spp.)

Diptera 59

Phorcia sp.
Phorcia sp.
Phorcia sp.
Phorcia sp.
Phorcia sp.
Phorcia sp.

Hydropsyche sp.

Station D.I.

✓ Oligochaeta 2 spp.

✓ Planaria

D = 1.58033 D:LN = 6.46337D = 2 = .0444.7
 D:LN = 2.35192 D = 6.30536

APPENDIX P CONTINUED

BLACKFOOT RIVER AT BLACKFOOT CANYON CAMP #1

1 sq. ft. with round sampler L. Spence

I = 1.15179 D.I.H. = 7.02363E-27 0.72834
D.I.H. = 0.53491 T = 0.570593

EPHEMEROPTERA	186	0.1
<i>Heptagenia</i> spp.	60	
<i>Heptagenia</i> sp.	112	
<i>Heptagenia tibialis</i>	11	
<i>Heptagenia tibialis</i> sp.	1	Total 184

PLECOPTERA	28	0.1
<i>Acroneuria pacifica</i>	2	
<i>Acroneuria pacifica</i> or <i>maculata</i>	10	D = 1.97292 D.I.H. = 0.314212
<i>Nemoura glauca</i>	2	D.I.H. = 0.30735 T = 0.413152
<i>Acroneuria pacifica</i>	1	
<i>Alloperla</i> sp.	13	Total 28

TRICHOPTERA	9	7
<i>Hydropsyche</i> sp.	2	
<i>Hydropsyche</i> sp.	1	D = 0.03089 D.I.H. = 1.44551
<i>Arctopsyche grandis</i>	4	D.I.H. = 0.53495 T = 0.432527
<i>Trichoptera</i> sp.	1	
<i>Macropodella</i> sp.	1	Total 9

DIPTERA	42	0.1
<i>Hexatoma</i> sp.	33	
<i>Hexatoma</i> sp.	1	
<i>Hexatoma</i> sp.	1	D = 0.04914 D.I.H. = 0.591439
<i>Hexatoma</i> sp.	2	D.I.H. = 0.314212 T = 0.413152
<i>Hexatoma</i> sp.	4	
<i>Hexatoma</i> sp.	4	
<i>Hexatoma</i> sp.	1	Total 46
Unknown Chironomidae	3	

COLEOPTERA	1	7
<i>Oligoneurus</i> sp. larva	1	Total 1
HEMPTERA	28	7
<i>Oligoneurus</i>	23	Total 23

Original Total 294 Final Total 291

Identified and Enumerated by
Patricia Ann Mitchell
MS, University of Montana 1968
May 27, 1973

Station D.I.

D = 0.04914 D.I.H. = 0.591439
D.I.H. = 0.413152 T = 0.413152

BLACKFOOT RIVER AT BLACKFOOT CANYON CAMP #2

1 sq. ft. from center of riffle L. Spence

EPHEMEROPTERA	370	0.2
<i>Heptagenia</i> spp.	127	
<i>Heptagenia</i> sp.	236	
<i>Heptagenia tibialis</i>	3	
<i>Heptagenia tibialis</i>	1	Total 367

PLECOPTERA	16	7
<i>Nemoura glauca</i>	1	
<i>Alloperla</i> sp.	11	
<i>Arctopsyche curvata</i> or <i>parvula</i>	1	Total 14
<i>Acroneuria pacifica</i>	1	

TRICHOPTERA	6	7
<i>Glossosoma</i> sp.	4	
<i>Arctopsyche grandis</i>	2	Total 6

DIPTERA	32	0.1
<i>Hexatoma</i> spp.	13	
<i>Alloperla</i> sp.	5	
<i>Hexatoma</i> sp.	1	
<i>Hexatoma</i> sp.	4	
<i>Hexatoma</i> sp.	7	Total 30

COLEOPTERA	1	7
<i>Oligoneurus</i> sp. adult	1	Total 1
Original Total 425		Final Total 416

Identified and Enumerated by
Patricia Ann Mitchell
University of Alberta, Edmonton
May 28, 1973
Patricia Ann Mitchell

BLACKFOOT RIVER AT BLACKFOOT CANYON CAMP #3

1 sq. ft. from left side of riffle L. Spence

EPHEMEROPTERA	202	0.1
<i>Heptagenia</i> spp.	211	
<i>Heptagenia</i> sp.	87	
<i>Heptagenia tibialis</i>	2	
<i>Heptagenia tibialis</i>	1	
<i>Heptagenia tibialis</i> sp.	1	Total 204

PLECOPTERA	11	0.2
<i>Acroneuria pacifica</i>	3	
<i>Alloperla</i> sp.	4	
<i>Acroneuria pacifica</i>	1	
<i>Acroneuria pacifica</i>	2	Total 10

TRICHOPTERA	4	7
<i>Arctopsyche grandis</i>	3	
<i>Glossosoma</i> sp.	1	Total 4

DIPTERA	18	0.3
<i>Atherix varians</i>	8	
<i>Chironomidae</i> unknown	3	
<i>Hexatoma</i> spp.	5	
<i>Simuliidae</i>	2	Total 18

HEMATODA	5	7
<i>Hematoda</i>	1	
<i>Oligoneurus</i>	2	Total 3

plus 2 chironomid and tubes

Original Total 240 Final Total 232

Identified and Enumerated by
Patricia Ann Mitchell
University of Alberta, Edmonton
May 28, 1973
Patricia Ann Mitchell

APPENDIX P CONTINUED

BLACKFOOT RIVER AT BLACKFOOT CANYON CAMP #1

1 sq. ft. 10-26-71 L. Spence

EPHEMEROPTERA	520	0.8		
<i>Baetis</i> sp. (one species)	36			
<i>Chironomus</i> sp.	125			
<i>Stygomyia</i> sp.	339			
<i>Stygomyia</i> sp.	2			
<i>Stygomyia</i> sp.	2			
<i>Stygomyia</i> sp.	42			
<i>Stygomyia</i> sp.	2			
			Total	526
PLECOPTERA	72	0.4		
<i>Nemoura cinctipes</i>	2			
<i>Nemoura</i> sp.	2			
<i>Leuctra</i> sp.	18			
<i>Leuctra</i> sp.	4			
<i>Pteronarcys</i> sp.	6			
<i>Pteronarcys</i> sp.	4			
<i>Pteronarcys</i> sp.	4			
<i>Pteronarcys</i> sp.	32			
			Total	72
TRICHOPTERA	122	0.2		
<i>Hydropsyche</i> sp.	67			
<i>Hydropsyche</i> sp.	3			
<i>Hydropsyche</i> sp.	43			
<i>Hydropsyche</i> sp.	2			
<i>Hydropsyche</i> sp.	7			
			Total	122
DIPTERA	24	0.1		
<i>Hexatoma</i> sp.	9			
<i>Asota</i> sp.	1			
<i>Asota</i> sp.	1			
<i>Asota</i> sp.	3			
<i>Asota</i> sp.	2			
<i>Asota</i> sp.	2			
			Total	24
COLEOPTERA	1	T		
<i>Optioeruvus</i> sp.	1			
			Total	1
NEMATODA	3	T		
<i>Nematoda</i>	3			
			Total	3
ANNELIDA	26	T		
<i>Oligochaeta</i>	26			
			Total	26
COTYLS	1	T		
			Total	1
			Original Total	769
			Final Total	774

BLACKFOOT RIVER AT BLACKFOOT CANYON CAMP #2

1 sq. ft. 10-26-71 L. Spence

EPHEMEROPTERA	575	0.9		
<i>Baetis</i> sp.	53			
<i>Chironomus</i> sp.	151			
<i>Stygomyia</i> sp.	278			
<i>Stygomyia</i> sp.	1			
<i>Stygomyia</i> sp.	1			
<i>Stygomyia</i> sp.	76			
<i>Stygomyia</i> sp.	4			
<i>Stygomyia</i> sp.	1			
			Total	565
PLECOPTERA	104	0.7		
<i>Nemoura cinctipes</i>	19			
<i>Nemoura</i> sp.	4			
<i>Leuctra</i> sp.	8			
<i>Leuctra</i> sp.	9			
<i>Leuctra</i> sp.	29			
<i>Leuctra</i> sp.	8			
<i>Leuctra</i> sp.	30			
			Total	107
TRICHOPTERA	324	1.0		
<i>Hydropsyche</i> sp.	234			
<i>Hydropsyche</i> sp.	25			
<i>Hydropsyche</i> sp.	44			
<i>Hydropsyche</i> sp.	14			
			Total	317
DIPTERA	30	0.2		
<i>Hexatoma</i> sp.	7			
<i>Asota</i> sp.	10			
<i>Asota</i> sp.	8			
<i>Asota</i> sp.	1			
<i>Asota</i> sp.	3			
			Total	29
COLEOPTERA	2	T		
<i>Optioeruvus</i> sp.	1			
<i>Optioeruvus</i> sp.	1			
			Total	2
ANNELIDA	23	T		
<i>Oligochaeta</i>	23			
			Total	23
NEMATODA	3	T		
<i>Nematoda</i>	3			
			Total	3
			Original Total	1061
			Final Total	1046

Patricia Ann Mitchell

BLACKFOOT RIVER AT BLACKFOOT CANYON CAMP #3

1 sq. ft. 10-26-71 L. Spence

EPHEMEROPTERA	344	0.5		
<i>Baetis</i> sp.	37			
<i>Chironomus</i> sp.	71			
<i>Stygomyia</i> sp.	170			
<i>Stygomyia</i> sp.	52			
<i>Stygomyia</i> sp.	1			
			Total	331
PLECOPTERA	71	0.4		
<i>Nemoura cinctipes</i>	3			
<i>Nemoura</i> sp.	8			
<i>Leuctra</i> sp.	9			
<i>Leuctra</i> sp.	18			
<i>Leuctra</i> sp.	2			
<i>Leuctra</i> sp.	26			
<i>Leuctra</i> sp.	4			
			Total	70
TRICHOPTERA	223	0.4		
<i>Hydropsyche</i> sp.	154			
<i>Hydropsyche</i> sp.	20			
<i>Hydropsyche</i> sp.	4			
<i>Hydropsyche</i> sp.	43			
<i>Hydropsyche</i> sp.	1			
			Total	224
DIPTERA	31	0.2		
<i>Hexatoma</i> sp.	9			
<i>Asota</i> sp.	3			
<i>Asota</i> sp.	11			
<i>Asota</i> sp.	1			
<i>Asota</i> sp.	5			
<i>Asota</i> sp.	1			
			Total	30
NEMATODA	9	T		
<i>Nematoda</i>	2			
			Total	2
OLIGOCHAETA	7	T		
<i>Oligochaeta</i>	7			
			Total	7
COTYLS	1	T		
			Total	1
			Original Total	679
			Final Total	664

Patricia Ann Mitchell

/ Patricia Ann Mitchell

Blackfoot River at Rainbow Bend 7/18/68Plecontera 23 5.5 ccPteronarcys californica NewportClaassenia sabulosa BanksTrichoptera 76 .7 ccGlossosoma sp.Hydropsyche sp.Arctopsyche sp.

Station D.I.

Ephemeroptera 85 .25 cc

D = 1.33433 DMIN = 0.174355

DMAK = 2.32193 R = 0.241046

Epeorus longimanus (Eaton)Baetis spp. (2 types)Ephemerella flavilinea McD.Ephemerella tibialis McD.Ephemerella hecuba (Eaton)Rhithrogena undulata (Banks) (prob.)Coleoptera 1 TOptioservus sp.✓ Gastropoda 3 .15 ccDiptera 24 .30 ccPhysa sp.Antocha sp.✓ Oligochaeta 5 TProtonotaria fitchii Osten SackenAtherix variegata WalkerHexatoma sp.Tendipeidae

APPENDIX P CONTINUED

BLACKFOOT RIVER AT RAINBOW BRID #1

1 sq. ft. 11-2-71 L. Spence

HEMiptera 181 0.2	
<i>Baetis</i> sp. (probably <i>bicaudatus</i>)	6
<i>Acanthias</i> sp.	1
<i>Paraleptophlebia</i> sp.	1
<i>Stygomyia</i> sp.	3
<i>Stygomyia</i> sp.	28
<i>Stygomyia</i> sp. (too small to ident. further)	19
<i>Stygomyia</i> sp. (similar to <i>infrequens</i>)	112
Total	177

HEMiptera 75 0.3	
<i>Rhyacophila brevicauda</i> (probably)	28
<i>Rhyacophila californica</i>	1
<i>Rhyacophila parvula</i> or <i>parvella</i>	1
<i>Rhyacophila</i> sp.	28
<i>Rhyacophila californica</i>	8
<i>Rhyacophila</i> sp.	6
Total	74

HEMiptera 80 0.3	
<i>Rhyacophila</i> sp.	74
<i>Rhyacophila</i> sp.	1
<i>Rhyacophila</i> sp.	2
Total	77

HEMiptera 47 0.4	
<i>Rhyacophila</i> sp.	1
<i>Rhyacophila</i> sp.	2
<i>Rhyacophila</i> sp.	2
<i>Rhyacophila</i> sp.	1
<i>Rhyacophila</i> sp.	15
<i>Rhyacophila</i> sp.	28
Total	46

HEMiptera 4 T	
<i>Rhyacophila</i> sp. lar.	5
Total	4

HEMiptera 10 T	
<i>Rhyacophila</i> sp.	9
<i>Rhyacophila</i> sp.	1
Total	10

Original Total 397 Final Total 557

Identified and Enumerated by
Patricia Ann Mitchell
Department of Zoology
University of Alberta
Edmonton, Alberta, Canada
July 30, 1973

Station D.I.

D = 3.11882 WMIN = 0.015179
DIAM = 0.02746 DM = 0.056355

BLACKFOOT RIVER AT RAINBOW BRID #2

1 sq. ft. 11-2-71 L. Spence

HEMiptera 103 0.1	
<i>Baetis</i> sp.	4
<i>Paraleptophlebia</i> sp.	2
<i>Stygomyia</i> sp.	2
<i>Stygomyia</i> sp.	12
<i>Stygomyia</i> sp.	18
<i>Stygomyia</i> sp. (similar to <i>infrequens</i>)	42
<i>Stygomyia</i> sp.	3
Total	103

HEMiptera 22 0.5	
<i>Rhyacophila californica</i>	3
<i>Rhyacophila brevicauda</i>	2
<i>Rhyacophila acuticollis</i>	1
<i>Rhyacophila</i> sp.	6
<i>Rhyacophila californica</i>	1
<i>Rhyacophila</i> sp.	2
<i>Rhyacophila</i> sp. (small)	2
Total	19

HEMiptera 150 0.4	
<i>Rhyacophila</i> sp.	144
<i>Rhyacophila</i> sp.	1
<i>Rhyacophila</i> sp.	1
Total	146

HEMiptera 11 T	
<i>Rhyacophila</i> sp.	2
<i>Rhyacophila</i> sp.	1
<i>Rhyacophila</i> sp.	8
Total	11

HEMiptera 3 T	
<i>Rhyacophila</i> sp.	3
Total	3

Original Total 289 Final Total 282

Identified and Enumerated by
Patricia Ann Mitchell
MS, Univ. of Montana, 1968
July 30, 1973

Patricia Ann Mitchell

BLACKFOOT RIVER AT RAINBOW BRID #3

1 sq. ft. 11-2-71 L. Spence

HEMiptera 104 T	
<i>Baetis</i> sp.	14
<i>Stygomyia</i> sp.	30
<i>Stygomyia</i> sp.	13
<i>Stygomyia</i> sp.	1
<i>Stygomyia</i> sp. (similar to <i>infrequens</i>)	41
Total	101

HEMiptera 21 0.5	
<i>Rhyacophila brevicauda</i> ?	1
<i>Rhyacophila californica</i>	1
<i>Rhyacophila</i> sp.	1
<i>Rhyacophila</i> sp.	2
<i>Rhyacophila</i> sp.	8
<i>Rhyacophila</i> sp.	1
<i>Rhyacophila californica</i>	7
<i>Rhyacophila</i> sp.	1
Total	22

HEMiptera 104 0.8	
<i>Rhyacophila</i> sp.	96
<i>Rhyacophila</i> sp.	4
<i>Rhyacophila</i> sp.	1
<i>Rhyacophila</i> sp.	1
Total	102

HEMiptera 8 T	
<i>Rhyacophila</i> sp. larva (probably)	1
<i>Rhyacophila</i> sp.	1
<i>Rhyacophila</i> sp.	6
Total	8

HEMiptera 1 T	
<i>Rhyacophila</i> sp. larva	1
Total	1

HEMiptera 1	
<i>Rhyacophila</i> sp.	1
Total	1

Original Total 239 Final Total 235

Identified and Enumerated by
Patricia Ann Mitchell
MS, University of Montana, 1968
July 31, 1973

Patricia Ann Mitchell

Appendix Q

DIVERSITY INDEX COMPUTATION PROCEDURE

By
Don O. Loftsgaarden

These comments refer to a paper by Wilhm and Dorris, Bioscience 18(6), 477-481, June 1968 entitled "Biological Parameters for Water Quality Criteria." In particular, comments refer to columns 2 and 3 on page 478.

The value of $\bar{d}$ (referred to as D in the program) used for the calculations is given at the top of column 3.

$$\bar{d} = - \sum_{i=1}^s \left(\frac{n_i}{n} \right) \log_2 \left(\frac{n_i}{n} \right) \quad (1)$$

This $\bar{d}$ value follows from the last $\bar{H}$ value at the bottom of column 2 by substituting sample values for population values (i.e., n_i for N_i , n for N , etc.).

This last $\bar{H}$ value follows from the first $\bar{H}$ value in column 2 by approximating the factorials.

$\bar{d}_{\min}$ and $\bar{d}_{\max}$ are given at the top of column 3 but still have factorials in them, making them impossible to use when n_i and n are large. However, the maximum $\bar{d}$ value occurs when all s species occur equally often, i.e., when

$$n_i = \frac{n}{s} \quad i = 1, 2, \dots, s.$$

But this implies that

$$\frac{n_i}{n} = \frac{\frac{n}{s}}{n} = \frac{1}{s} \quad \text{for } i = 1, 2, \dots, s.$$

If we make this substitution for n_i in formula (1) above, we find that $\bar{d}_{\max}$ is as follows:

$$\bar{d}_{\max} = - \log_2 \left(\frac{1}{s} \right) \quad (2)$$

This is the formula used in the program.

The minimum $\bar{d}$ value occurs when $s-1$ of the species have $n_i = 1$ and the other species has all of the rest of the insects or a total of $n-(s-1)$. Substituting these values in (1) above we arrive at

$$\bar{d}_{\min} = - \left(\frac{s-1}{n} \right) \log_2 \left(\frac{1}{n} \right) - \frac{n-s+1}{n} \log_2 \frac{n-s+1}{n} \quad (3)$$

This is the formula used in the program.

Reviewing, in the program $\bar{d} = D$, $d_{\max} = D \text{ MAX}$, and $d_{\min} = D \text{ MIN}$. Formulas (1), (2), and (3) above are used to do the computations. The redundancy is computed using the formula given in the paper, i.e.,

$$R = \frac{D_{\max} - D}{D_{\max} - D_{\min}}$$

Note: A copy of the program used in the computations is attached.

DINDEL

03:03

17-APR-74

```

5 REM STATION 9
10 REM PUT H,G, AND N(1) IN DATA LINES 340,341,ETC
100 READ A,M,S
110 IF M=999 GO TO 400
110 MAT READ H(S)
120 DEF FNL(A) = LN(A)/LN(2)
130 LET D=0
150 FOR I=1 TO S
160 LET N=N(1)/M
170 LET D = D+N*FNL(X)
180 NEXT I
200 LET N=(S-1)/M
210 LET Y=1/M
220 LET Z=(M-S+1)/M
230 LET W = N*FNL(Y) + Z*FNL(Z)
240 LET V = (-1)*W
260 LET Q = (-1)*FNL(1/5)
280 LET R=(G-D)/(G-V)
290 PRINT "PROBLEM 1.D. = "A;"D="D;"DMIN="W
291 PRINT "DMAX="G;"E="R
295 PRINT
310 GO TO 100
340 DATA 9,540,5
341 DATA 69,211,229,5,26
350 DATA 10,999,10
400 END

```

*A = 9 = Problem I.D., N = 540 = TOTAL No. BUGS, S = 5 = No. species
 N(1) = 69 = No. bugs species 1, N(2) = 211 = No. bugs species 2, etc
 (Actually for the example shown species should be read order).*

READ?

→ This is an A, M, S used to tell program to stop.
 If M=999 program stops, see line 110 in program.

Data from more than one problem can be run at the same time. Data for the additional problems is put in program lines 342-359 using the format prescribed in lines 340 and 341 above.

Summary of kinds and abundance of bottom fauna identified from the Blackfoot River and certain tributaries from 1968-1971.

Collection methods are explained in Appendix P.

Numbers in each species were not determined in 1968 or 1969, but were determined in 1971.

Volumes given were determined by order in the Pollution Control Laboratory, Helena.

Final totals were determined by Ms. Patricia Ann Mitchell during species identification and are the same as those given in Appendix P.

Numbers in parentheses following the number of individuals in a genus show the number of species which were identified within that genus at the station. (In 1968 and 1969 data, these numbers are in parentheses after an (x), which simply indicates the presence of a genus.) Numbers given in 1968 and 1969 data are total number of individuals in the order.

Volumes less than 0.01 cc are shown as T (trace). Each T represents the volume less than 0.01 cc in a one square foot sample.

APPENDIX R continued

Number and Kinds of Major Aquatic Insects Collected with Kick Screen ($\pm 4 \text{ ft}^2$ Sample) from Blackfoot River
in 1968 and 1969

| Location
Month/Yr
No. of Samples	Pop's			Fleisher			Hogum			Dalton			Canyon			Rainbow		
	7/68	4/69	10/69	7/68	4/69	10/69	7/68	4/69	10/69	7/68	4/69	10/69	7/68	4/69	10/69	7/68	4/69	10/69
<u>Plecoptera</u> (Stoneflies)	4	UN	95	22	8	42	53	65	49	17		69	7		78	23		
Nemoura	x	x	x	x	x(2)	x	x	x(2)	x	x					x			
Capnia								x(2)	x									
Pteronarcys										x		x		x		x		
Pteronarcocella										x		x			x			
Arcynopteryx									x			x			x			
Isogenus			x		x					x		x						
Isoperla			x					x										
Alloperla	x?			x	x	x	x	x		x?		x			x			
Hastaperla			x	x					x				x					
Paraperla	x					x									x			
Acroneuria				x			x	x	x	x		x	x		x			
Glaassenia				x	x	x	x	x							x	x		
Kathroperla							x			x?								
<u>Ephemeroptera</u> (Mayflies)	0	2		10	1	UN	508	254	32	432		229	475		401	85		
Heptagenia																		
Cinygmula			x		x	x	x	x	x	x			x					
Epeorus				x	x		x	x		x			x			x		
Rithrogena								x		x		x	x		x	x		
Baetis				x			x	x		x		x	x		x	x		
Paraleptophlebia							x	x	x									
Ameletus																		
Ephemerella							x(2)	x(3)	x(2)	x(4)		x(3)	x(3)		x(3)	x(3)		
<u>Tricoptera</u> (Caddis flies)	12		4	24	7	8	155	62	399	24		211	6		139	76		
Rhyacophila	x		x	x			x	x	x	x		x						
Glossosoma								x				x			x	x		
Anagapetus							x											
Sortosa																		
Tinodes																		

APPENDIX R continued
Number and Kinds of Major Aquatic Insects Collected from Blackfoot River in 1968 and 1969

Location	Pop's			Fletcher			Hogum			Dalton			Canyon			Rainbow		
Month/Yr	7/68	4/69	10/69	7/68	4/69	10/69	7/68	4/69	10/69	7/68	4/69	10/69	7/68	4/69	10/69	7/68	4/69	10/69
No. of Samples	2	1	1	1	1	1	1	1	1	1	None	1	1	None	1	1	None	None
Hydropsyche	x				x			x	x			x	x			x	x	
Arctopsyche	x		x			x					x				x	x		
Psychomyia							x											
Parapsyche			x															
Ochrotrichia							x											
Lepidostoma					x		x	x	x			x						
Brachycentrus	x			x		x	x	x	x	x		x	x			x		
Limnephilidae							x	x	x	x		x						
<u>Coleoptera (Beetles)</u>	8			32	UN		124	14	50	12		5	9			13	1	
Zaitzevia	x			x	x		x		x									
Narvus				x			x	x				x						
Lara				x			x											
Optioservus				x			x	x	x	x		x	x			x	x	
Heterolimnius							x	x	x	x		x				x		
Cleptelmis										x								
<u>Diptera (True flies)</u>	51			91	10	24	813	91	44	443		26	92			39	24	
Hexatoma				x		x	x		x	x			x			x	x	
Antocha					x		x										x	
Rhabdomastix							x											
Chelifera																		
Pericoma								x	x									
Protoplasa										x		x				x	x	
Atherix										x		x				x	x	
Tanytarsus							x											
Pentaneura							x											
Metriocnemus								x										
Hydrobaenus	x												x			x		
Brillia					x													
Cardiocladius								x										
Diaenosa								x										
Simuliidae					x	x	x		x	x			x			x		
Tendipedidae						x				x						x	x	
Empididae													x					

A-118

UN = Unknown
? = Identity not positive
() = Indicates more than one species

Number and Volume of Major Aquatic Insects Collected from Blackfoot River with
Kick Screen in 1968 and 1969 by Order.

Location	Fon's		Flasher		Bogum		Dalton		Canyon		Rainbow	
Year	1968	1969	1968	1969	1968	1969	1968	1969	1968	1969	1968	1969
No. of Samples	2	2	1	2	1	2	1	1	1	1	1	None
Plecoptera												
No.	4	95	22	50	53	114	17	69	7	78	23	
Vol. ^{2/}	T	0.3+	0.4	-	0.7	-	0.9	-	0.05	-	5.5	
Ephemeroptera												
No.	0	2	10	1	508	32	432	229	475	401	85	
Vol.	-	-	0.1	T	1.4	-	1.6	-	1.9	-	0.25	
Tricoptera												
No.	12	4	24	15	155	399	24	211	6	139	76	
Vol.	0.4	0.1	0.2	UN-0.1	1.3	-	0.2	-	0.1	-	0.7	
Coleoptera												
No.	8		32		124	50	12	5	9	13	1	
Vol.	T-0.05		0.05		0.05	-	0.05	-	T	-	T	
Diptera												
No.	51	2	91	34	813	44	443	26	92	39	24	
Vol.	T-0.1	T	0.4	UN-T	2.0	-	0.9	-	0.6	-	0.3	

^{1/}Each sample is ± 4 square feet

^{2/}Volume in cc

T = Trace ($< .01$)

- = Vol. not known

UN = Vol. not given for one sample

APPENDIX B continued

March/April 1971

| Location
No. sq. ft.
sampled | Fletcher
3 | Alice Creek
2 | Hogum
3 | Portman Creek
3 | Dalton
3 | Canyon
3 |
|------------------------------------|---------------|------------------|------------|--------------------|-------------|-------------|
| PLECOPTERA | | | | | | |
| Number of Individuals | | | | | | |
| Pteronarcissa | - | - | - | 2 | 5 | 15 |
| Nemoura | 248(2) | 446(2) | 501(2) | 69 | 1 | 9 |
| Capnia | 38 | 78 | 161 | - | 4 | 8 |
| Isoperla | - | - | 40 | 38 | 13 | 44 |
| Argyropteryx | 3 | 1 | 1 | - | - | 5 |
| Acrocnemia | 3 | 1 | 7 | 97 | 11 | 19(2) |
| Classania | 2 | - | 6 | - | 1 | 10 |
| Alloperla | 105 | 135 | 221 | 25 | 68 | 82 |
| Brachyptera | 4 | 1 | - | - | - | 1 |
| Isogenus | 19 | 9 | 5 | 16 | 3 | - |
| Barlaerla | - | - | - | 18 | 6 | 19 |
| Paraperla | 2 | - | - | - | - | - |
| Chloroperlidae | 10 | - | 3 | - | - | - |
| Unknown | 3 | 3 | - | - | 2 | - |

021-y

Pinal Total	436	674	954	265	114	212
Total Volume (cc)	1.1	1.2	2.4	3.9	0.7	2.2
Ave. no./sq. ft	146	337	318	88	38	71
Ave. vol./sq. ft
(cc)	0.4	0.6	0.8	1.3	0.2	0.7

1/
1 sample (1 sq. ft.) lost prior to identification

March/April 1971

| Location
No. sq. ft.
sampled | Fletcher
3 | Alice Creek
2 | Hogum
3 | Portman Creek
3 | Dalton
3 | Canyon
3 |
|------------------------------------|---------------|------------------|------------|--------------------|-------------|-------------|
| EPHEMEROPTERA | | | | | | |
| Baetis | 107 | 320 | 461(2) | 856(2) | 132(2) | 133(3) |
| Paraleptophlebia | - | 17 | 64 | 22 | 2 | 9 |
| Rhyacophila | - | 199(3) | 538(3) | 238(4) | 81(4) | 146(4) |
| Rhyacophila | - | 24 | 3 | 70 | 34(2) | 915 |
| Cinygmula | 43 | 210 | 1461 | 317 | 25(2) | 21 |
| Rhyacophila | 24 | 27 | 219 | 160 | - | 3 |
| Ameletus | - | - | 4 | - | - | 1 |
| Total | 174 | 787 | 2790 | 1663 | 583 | 1222 |
| Total Volume (cc) | 0.2+T | 1.3 | 1.3 | 2.7 | 1.4 | 2.1 |
| Ave. no./sq. ft | 58 | 394 | 917 | 554 | 194 | 407 |
| Ave. vol./sq. ft
(cc) | 0.07+T | 0.65 | 0.43 | 0.9 | 0.47 | 0.7 |

TRICOPTERA

Glossosoma	1	81	235	35	419	295
Brachycentrus	20	20	34	1	100	100
Hydropsyche	38	307	849	1	128	141
Arctopsyche	3	9	3	140	55	8
Rhyacophila	4	31(2)	88(3)	78(2)	1	-
Psychrocnema	-	1	-	-	-	-
Lepidostoma	-	-	188	-	-	2
Hydroptila	-	-	-	1	-	-
Limnephilidae	2	12	15	7	5	9

Total	68	461	1412	263	708	555
Total Volume (cc)	0.3+T	1.1	2.1	4.3	2.5	1.9
Ave. no./sq. ft	23	231	471	88	236	185
Ave. vol./sq. ft
(cc)	0.1+T	0.55	0.7	1.4	0.83	0.6

APPENDIX R continued

March/April 1971

March/April 1971

| Location
No. sq. ft.
sampled | Flesher
3 | Alice Creek
2 | Hogum
3 | Poorman Creek
3 | Dalton
3 | Canyon
3 |
|------------------------------------|--------------|------------------|------------|--------------------|-------------|-------------|
| DIPYTERA | | | | | | |
| Simuliidae | 131 | 5 | 45 | 8 | 1 | 324 |
| Atherix | - | - | - | - | 17 | 28 |
| Hexatoma | 1 | 2 | 8 | 1 | 7 | 29(3) |
| Epididae | 2 | - | 9(3) | 88(3) | 2 | 4(3) |
| Trillia | 22 | 1 | 66(2) | 19 | 3 | 8 |
| Rhabdomastix | - | - | - | - | - | 2 |
| Hydrobatus | | | | | | |
| (Orthocladus)* | 165(2) | 284(3) | 469(3) | 60(3) | 6 | 4(2) |
| Dicranota | 1 | - | 3 | 4 | 2 | 3 |
| Metriocnemus | 1 | 53 | 10 | 30 | 15 | 3 |
| Antocha | 37 | 58 | 42 | 34 | 5 | 1 |
| Cardiocladius | 3 | 86 | 30 | 1 | 9 | - |
| Pericoma | 24 | 63 | 118 | 147 | 3 | - |
| Calopsoceta | | | | | | |
| (Tanytarsus)* | 1 | - | 2 | 26 | - | - |
| Pentaneura | - | 1 | 3 | - | 3 | - |
| Diamesa | - | 11 | - | - | - | - |
| Oricoptopus | | | | | | |
| (Orthocladus)* | - | - | 1 | 24 | - | - |
| Helicidae | - | 3 | - | - | - | - |
| Chironomidae | - | 50 | 57 | 25 | 8 | 2 |

Total	388	617	863	467	81	408
Total Volume (cc)	0.4	2.0	0.8	0.4	0.3	1.2
Ave. no./sq. ft.	129	309	288	156	27	136
Ave. vol./sq. ft.
(cc)	0.13	1.0	0.27	0.13	0.1	0.4

*Names in parentheses are the most recent Genus names, and are used in August 1971 data.

COLLEPTERA

Optioservus	-	25	59	5	23	7
Heterlimnius	-	8	2	111	3	-
Salixia	-	2	10	-	-	-
Claytonia	-	-	-	-	5	-
Brychius	-	-	-	1	-	-
Total	0	35	71 1/2	117	31	7
Total Volume (cc)	-	7	0.2	0.2+T	T	T
Ave. no./sq. ft.	-	17	35	39	10	2
Ave. vol./sq. ft.
(cc)	-	7	0.1	0.07+T	T	T

1/ one vial missing from sample

| Location
No. sq. ft.
sampled | Flesher
3 | Alice Creek
2 | Hogum
3 | Poorman Creek
3 | Dalton
3 | Canyon
3 |
|------------------------------------|--------------|------------------|------------|--------------------|-------------|-------------|
| OLEDOCHARTA and
NEMATODA | 0 | 1 | 0 | 44 | 71 | 12 |
| Total Volume (cc) | | T | | 1.9+T | 0.1+T | T |
| Grand Total | 1066 | 2576 | 6050 | 2863 | 1600 | 2652 |
| Volume (cc) | 2.0 | 5.6+T | 6.8 | 11.5+T | 5.0+T | 7.4+T |
| Ave. no./sq. ft. | 355 | 1288 | 2017 | 954 | 533 | 854 |
| Ave. vol./sq. ft.
(cc) | 0.67 | 2.80+T | 2.24 | 3.84 | 1.66 | 2.47+T |

APPENDIX R continued

August 1971

| Station
No. sq. ft.
sampled | Pop'a | Pleather | Alice Creek
0 100 | Alice Creek
0 N.P. Rudy | Bogum | Below
Lauders Fork | Peckman Creek | Salton | Canyon |
|-----------------------------------|-------|----------|----------------------|----------------------------|--------|-----------------------|---------------|--------|--------|
| 2 | 3 | | | | 3 | 3 | 3 | 3 | 3 |
| Number of Individuals | | | | | | | | | |
| PLECOPTERA | | | | | | | | | |
| Pteronarcissa | - | - | - | - | 11 | 11 | 2 | 12 | 3 |
| Nemoura | 88 | 643 | 113 | 24 | 174 | 11 | 398 | 3 | 3 |
| Captia | - | 5 | 6 | - | 1 | - | - | - | - |
| Imania | - | 2 | 39(2) | 17(2) | 3 | 1 | - | - | 11 |
| Acronarcissa | 1 | 1 | 15(2) | 1 | 10(1) | 9 | 38 | 4 | 3 |
| Classania | - | 2 | - | - | 25 | 1 | - | - | 1 |
| Alloparia | 21 | 192 | 256 | - | 366 | 145 | 28 | 17 | 28 |
| Brachyptera | - | - | - | - | - | - | - | - | - |
| Isogonus | 13 | 2 | 3(2) | - | 2 | 3 | 3 | 1 | - |
| Nectoparia | - | - | - | - | - | - | - | - | - |
| Paraparia | 4 | - | - | - | - | - | - | 2 | 1 |
| Europtania | - | - | - | 204 | - | - | 11 | - | - |
| Chloroparidae | - | - | - | - | - | - | - | - | - |
| Peristodidae | - | 1 | - | - | - | 5 | - | - | - |
| Unknown | - | 3 | - | 2 | - | - | - | - | - |
| Final Total | 127 | 872 | 434 | 248 | 580 | 187 | 488 | 39 | 52 |
| Total Volume (cc) | 3T | 1.1 | 0.8 | 0.24T | 2.0 | 0.3 | 1.44T | 3T | 0.34T |
| Ave no./sq. ft. | 62 | 291 | 145 | 83 | 193 | 62 | 166 | 13 | 17 |
| Ave vol./sq. ft. (cc) | T | 0.37 | 0.27 | .074T | 0.67 | 0.1 | .474T | T | 0.14T |
| EMBIPTERA | | | | | | | | | |
| Swella | 1 | 70 | 267 | 171 | 204 | 198 | 307 | 447 | 298 |
| Paralutophlebia | - | - | 34 | - | - | - | 14 | - | 1 |
| Epimeria | - | 1 | 145(4) | 86(4) | 137(3) | 37(4) | 34(4) | 37(3) | 18(3) |
| Rhithrogena | - | - | 33 | - | 3 | 513 | 60 | 132 | 637 |
| Climacia | - | - | 344 | 337 | 390 | 52 | 256 | 6 | - |
| Epurota | - | - | 2 | 91(2) | - | 108(2) | 33(2) | 1 | - |
| Amelura | - | - | - | 3 | - | - | - | - | - |
| Final Total | 1 | 71 | 843 | 890 | 736 | 1198 | 926 | 643 | 733 |
| Total Volume (cc) | T | 0.1+2T | 0.4 | 2.2 | 0.3 | 2.1 | 0.4 | 0.3 | 0.4 |
| Ave no./sq. ft. | 0.3 | 24 | 281 | 297 | 252 | 399 | 309 | 214 | 252 |
| Ave vol./sq. ft. (cc) | T | 0.034T | 0.13 | 0.73 | 0.1 | 0.7 | 0.2 | 0.17 | 0.13 |
| TRICOPTERA | | | | | | | | | |
| Glossosoma | - | - | 41 | 2 | 33 | 140 | 3 | 53 | 5 |
| Brachycentrus | - | 34 | 17 | - | 189 | 11 | - | 106 | 2 |
| Hydropsyche | - | 49 | 283 | - | 1790 | 5 | - | - | 1 |
| Amphipsyche | - | 30 | 72 | - | 86 | 102 | 102 | 29 | 9 |
| Rhyacophila | 1 | 7 | 20(3) | 1 | 44(2) | 4 | 24(2) | - | - |
| Psychotria | - | - | - | - | - | - | - | - | - |
| Lepidostoma | - | 1 | 29 | - | 17 | - | - | - | - |
| Hydropsylla | - | - | - | - | - | - | 1 | - | - |
| Limoniphilidae | - | 17(2) | 62(2) | - | 120 | 1 | 6 | 121(2) | 1 |
| Parapsyche | 1 | - | - | - | 3 | - | - | - | - |
| Tonina | - | - | 1 | - | - | 19 | - | - | - |
| Chimarra | - | - | 1 | - | 1 | - | 73 | 13 | 1 |
| Chimarrillo | - | - | 1 | - | - | - | - | - | - |
| Psychomyiidae | - | - | - | 2 | 4 | - | - | - | - |
| Attagapetus | - | - | - | - | 1 | - | - | - | - |
| Unknown | 2 | 2 | 37 | - | 16 | 13 | 3 | - | - |
| Final Total | 4 | 161 | 563 | 3 | 2286 | 297 | 217 | 213 | 19 |
| Total Volume (cc) | 3T | 0.3 | 1.1 | 3T | 3.2 | 1.3 | 0.74T | 0.142T | 3T |
| Ave no./sq. ft. | 1.3 | 34 | 188 | 1.7 | 743 | 99 | 72 | 71 | 4.3 |
| Ave vol./sq. ft. (cc) | T | 0.17 | 0.37 | T | 1.07 | 0.43 | 0.234T | 0.034T | T |

APPENDIX R continued

August 1971

| Station
No. sq. ft.
sampled | Pop'a
3 | Pleaser
3 | Alice Creek
@ 100
3 | Alice Creek
@ N.P. Body
3 | Hogus
3 | Below
Landers Fork
3 | Peorman Creek
3 | Detton
3 | Canyon
3 |
|-----------------------------------|------------|--------------|---------------------------|---------------------------------|------------|----------------------------|--------------------|-------------|-------------|
| Number of Individuals | | | | | | | | | |
| DIPTERA | | | | | | | | | |
| Simuliidae | 4 | 14 | 30 | - | 96 | 15 | 1299 | 222 | 4 |
| Atheria | - | - | - | - | - | - | - | 10 | 15 |
| Macalima | - | 1 | 9 | - | 4 | 8 | - | 1 | 31 |
| Empididae | - | - | 3 | - | - | - | 34(3) | 7(2) | 3(2) |
| Brillia | 6 | 25 | 15 | 24 | 22 | 5 | 157 | 1 | 2 |
| Rhachidomyia | - | - | - | - | - | - | - | - | - |
| Orthocladus | 12 | 27 | 156(3) | 30(2) | 95(2) | 3 | 39 | - | - |
| Dicranota | - | - | - | - | - | - | - | - | - |
| Metriocnemus | - | 11 | 88 | 11 | 79 | 6 | 55 | 30 | - |
| Antocha | - | 47 | 162 | - | 11 | 2 | 2 | - | - |
| Cardinalia | 6 | - | 31 | 3 | 18 | 67 | 3 | 4 | - |
| Pericoma | - | 55 | 158 | - | 49 | 1 | 24 | - | - |
| Tanytarsus | - | 25 | 6 | - | 150 | 17 | 15 | - | 4 |
| Pantophora | - | 1 | - | - | - | - | 1 | 3 | - |
| Clunias | - | - | - | - | - | - | - | - | - |
| Helinae | - | - | - | - | - | - | - | - | - |
| Chironomidae | - | 189 | 49 | - | 36 | 18 | 251 | 64 | 13 |
| Helidae | - | 2 | - | - | - | - | - | - | - |
| Polomya | - | - | - | 2 | - | - | - | - | - |
| Muscid | - | - | - | - | - | 1 | - | - | - |
| Tipulidae | - | - | - | - | - | - | - | 2 | - |
| Final Total | 26 | 397 | 707 | 70 | 342 | 122 | 1882 | 354 | 94 |
| Total Volume (cc) | 37 | 37 | 1.1 | 37 | 0.24T | 0.44T | 1.1 | 0.5 | 0.5 |
| Ave no./sq. ft. | 8.7 | 132 | 236 | 23 | 181 | 41 | 427 | 118 | 31 |
| Ave vol./sq. ft. (cc) | T | T | 0.37 | T | 0.074T | 0.134T | 0.37 | 0.17 | 0.17 |
| 1/ Formerly called Hydrobaenus | | | | | | | | | |
| 2/ Formerly called Calogasteria | | | | | | | | | |
| COLEOPTERA | | | | | | | | | |
| Optocerosus | 1 | 1 | 75 | 1 | 184 | 18 | 11 | 28 | 2 |
| Metasternus | 3 | - | 28 | - | 7 | - | 97 | 2 | - |
| Zaitzevia | 1 | 26 | 19 | 2 | 181 | - | 2 | 1 | - |
| Clapallia | - | - | - | - | 7 | - | - | 10 | - |
| Strophus | - | - | 1 | - | - | - | 6 | - | - |
| Lare | - | 1 | - | - | - | - | - | - | - |
| Harpus | - | - | - | - | 11 | - | 11 | 1 | - |
| Final Total | 5 | 28 | 123 | 3 | 390 | 18 | 127 | 42 | 2 |
| Total Volume (cc) | 27 | 37 | 0.14T | T | 0.14T | 37 | 37 | 37 | 27 |
| Ave no./sq. ft. | 1.7 | 9.3 | 41 | 1 | 130 | 6 | 42 | 14 | 0.7 |
| Ave vol./sq. ft. (cc) | T | T | 0.034T | T | 0.034T | T | T | T | T |
| OLIGONEURA and MINUTIDAE | | | | | | | | | |
| Final Total | - | 4 | 2 | 38 | 4 | 3 | 45 | 57 | 26 |
| Total Volume (cc) | - | T | 27 | 27 | T | T | 37 | 37 | T |
| Ave no./sq. ft. | - | 1.3 | 0.67 | 13 | 1.3 | 1 | 14 | 19 | 9 |
| Ave vol./sq. ft. (cc) | - | T | T | T | T | T | T | T | T |
| Grand Total | | | | | | | | | |
| No. | 163 | 1333 | 2672 | 1234 | 4560 | 1825 | 3693 | 1348 | 948 |
| Vol. | 117 | 1.74T | 3.544T | 2.44T | 5.844T | 4.143T | 3.844T | 1.1410T | 1.247T |
| Ave no./sq. ft. | 54 | 351 | 891 | 418 | 1520 | 808 | 1231 | 449 | 316 |
| Ave vol./sq. ft. (cc) | 1.3 | 0.374T | 1.174T | 0.844T | 1.934T | 1.374T | 1.274T | 0.374T | 0.447T |

APPENDIX R continued

October/November 1971

Station No., sq. ft. sampled	Pop's	Flesher	Allen Creek & 200	Hogus	Below Lander's Fork	Fourteen Creek	Dalton	Canyon	Rainbow Bend
3	3	3	3	3	3	3	3	3	3
Number of Individuals									
PLECOPTERA									
Pteronarcys	-	-	-	-	-	-	-	-	5
Pteronarcys	-	-	-	-	-	-	-	-	23
Heurore	62	81	77	38	14	150	29	24	1
Opesia	-	28	25	8	35	6	4	6	-
Leptocryptus	-	2	-	21	16	17	17	65	44
Leptocryptus	-	3	5	-	3(2)	-	-	4	1
Leptocryptus	-	7(2)	9(2)	27	9(3)	62	50(2)	16	17(2)
Leptocryptus	-	-	-	15	-	-	-	-	-
Leptocryptus	54	243	186	237	129	79	174	88	10
Leptocryptus	-	-	-	-	42	4	6	21	1
Leptocryptus	24	15	43	13	9(2)	15	3	-	-
Leptocryptus	-	-	-	-	-	-	-	-	-
Leptocryptus	5	6	-	-	2	-	5	2	-
Leptocryptus	18	-	-	-	-	-	-	-	31
Leptocryptus	-	-	5	-	-	-	-	-	-
Leptocryptus	-	-	-	-	2	-	-	-	2
Leptocryptus	-	46	-	-	-	-	-	-	-
Leptocryptus	-	-	-	-	4	-	-	-	-
Final Total	163	435	350	339	253	261	338	249	115
Total Volume (cc)	0.4	1.2	0.9	1.2	1.3	7.1	2.4	1.5	1.3
Ave. no./sq. ft.	54	145	137	113	84	823	109	83	38
Ave. vol./sq. ft. (cc)	0.13	0.60	0.30	0.40	0.43	2.37	0.8	0.5	0.43

1/ Total includes 780 Plecoptera which dried up in one square foot sample and were not identified.

EPHEMEROPTERA									
Heurore	-	157	173(2)	258(2)	193	1037(2)	150(2)	106	84
Heurore	-	-	36	10	-	99	7	7	-
Heurore	-	3	268(5)	53(13)	344(3)	180(4)	87(4)	174(3)	198(2)
Heurore	-	-	12	12	1015	164	1366	707	90
Heurore	-	97	268	528	221	1301	130	347	5
Heurore	-	-	-	-	-	-	1	-	52
Heurore	-	-	-	-	-	-	1	-	8
Heurore	-	-	-	-	-	-	-	-	1
Heurore	-	-	-	-	-	-	-	1	-
Final Total	-	217	775	1342	1773	2809	1736	1422	381
Total Volume (cc)	-	37	1.4	0.9	3.1	2.8	2.6	2.2	0.347
Ave. no./sq. ft.	-	72	258	447	591	934	579	474	127
Ave. vol./sq. ft. (cc)	-	7	0.47	0.3	1.0	0.93	0.87	0.73	0.147
TRICHOPTERA									
Glossosoma	-	2	237	329	626	164	294	132	1
Heurore	-	102	133	86	1	9	29	30	-
Heurore	1	30	420	1398	275	5	70	455	314
Heurore	2	-	30	-	68	355	92	48	-
Heurore	-	6(2)	34(2)	47(2)	7(2)	137(2)	6	-	2
Heurore	-	-	-	-	-	-	-	-	-
Heurore	-	-	-	83	-	4	-	1	-
Heurore	-	1	-	-	-	-	-	-	-
Heurore	-	3	5(2)	45(3)	5(2)	1	13	7	-
Heurore	1	-	-	-	-	-	-	-	5
Heurore	-	-	-	-	1	-	-	-	-
Heurore	-	-	9	-	-	-	-	-	-
Heurore	-	-	-	-	-	-	-	-	-
Heurore	-	-	-	-	-	-	-	-	1
Heurore	-	-	-	-	-	-	-	-	-
Heurore	-	-	7	7	-	-	-	-	2
Final Total	4	148	895	2004	980	661	504	663	325
Total Volume (cc)	27	0.3	3.6	4.4	2.1	6.3	2.8	1.6	1.5
Ave. no./sq. ft.	1.3	49	298	668	329	227	168	221	108
Ave. vol./sq. ft. (cc)	27	0.1	1.2	1.47	0.7	2.1	0.93	0.53	0.5

APPENDIX R continued

October/November 1972

Station No. of Pl. sampled	Pop's	Fisher	Alice Creek # 250	Hogum	Below Louisa Park	Peckman Creek	Dalton	Canyon	Rainbow Bend
	3	3	3	3	3	3	3	3	3
Number of Individuals									
DEPTERA									
Phallidae	-	-	62	2	4	9	38	16	-
Alaria	-	-	-	-	-	-	6	28	-
Coenocorys	-	5	9	5	22	-	1	25	1
Agallidae	-	1	7	8	8(1)	26(2)	7	-	-
Trilobella	-	-	5	-	-	7	-	-	-
Rhabdosomatina	-	32(several)	774(several)	3518(several)	22	256(several)	8	6	15
Oribolaelus	-	4	-	1	1	7	-	1	-
Diprasiidae	-	-	2	-	-	12	-	-	-
Metriocampa	1	-	418	151	3	340	31	4	1
Actinidae	-	351	-	-	2	-	-	-	-
Cardinalius	-	-	481	326	62	183	3	-	-
Parionina	-	208	-	-	-	-	-	-	-
Tanytarsus	-	-	-	-	-	-	-	-	4
Falsimura	-	-	-	-	-	-	-	-	-
Cixiidae	-	-	-	-	-	-	-	-	-
Belinidae	-	-	-	-	-	-	-	-	-
Chironomidae	25	-	-	-	-	-	-	3	40
Dolichopodidae	-	-	-	-	-	-	-	-	-
Tricorythidae	-	8	-	1	-	-	-	-	-
Tanytarsus	-	1	2	3	-	2	-	-	-
Falsimura	-	-	-	-	-	-	-	-	-
Final Total	27	610	1760	4045	124	876	94	83	65
Total Volume (cc)	37	0.6	0.4 + T	2.7	1.1	1.0	0.2 + T	0.5	0.4 + 2T
Ave. no./sq. ft.	9	2.03	587	1348	41	292	31	28	22
Ave. Vol/sq. ft. (cc)	T	0.2	0.13 + $\frac{T}{3}$	0.9	0.37	0.33	0.06 + $\frac{T}{3}$	0.17	0.13 + $\frac{2T}{3}$
COLEOPTERA									
Optioerinae	-	-	107	161	3	7	5	2	-
Delicliminae	-	-	38	7	-	181	1	-	-
Salicaria	1	-	26	78	1	2	-	1	5
Cryptorhinae	-	-	-	1	-	-	-	-	-
Hydrobiae	-	-	-	-	-	21	-	-	-
Larae	-	-	-	4	1	7	-	-	-
Barypa	-	-	-	-	-	-	-	-	-
Final Total	1	0	171	251	5	218	6	3	5
Total Volume (cc)	7	0	0.1+2T	0.3	3T	0.2+T	2T	3T	2T
Ave. no./sq. ft.	0.3	0	57	84	1.7	73	2	1	1.7
Ave. vol./sq. ft. (cc)	$\frac{7}{3}$	0	0.03+ $\frac{2T}{3}$	0.10	T	0.06+ $\frac{T}{3}$	$\frac{2T}{3}$	T	T
OLEDOGASTRA and BRACHYPTERA									
Final Total	0	4	3	11	7	101	27	64	14
Total Volume (cc)	0	2T	3T	3T	3T	0.9	3T	3T	3T
Ave. no./sq. ft.	0	1.3	1	3.7	2.3	34	9	21	4.7
Ave. vol./sq. ft.	0	$\frac{2T}{3}$	T	T	T	0.3	T	T	T
Grand Total									
No.	195	1414	3954	7992	3142	7147 ^{2/3}	2695	2264	905
Vol.	0.4+8T	2.1+5T	6.4+8T	9.5+3T	7.6+5T	18.3+T	8.0+6T	5.8+8T	3.5+8T
Ave. no./sq. ft.	65	471	1318	2664	1547	2182	898	828	302
Ave. vol./sq. ft. (cc)	0.33+2T	0.7+ $\frac{5T}{3}$	2.1+2T	3.17+T	2.53+ $\frac{5T}{3}$	6.1+ $\frac{T}{3}$	2.67+2T	1.93+2T	1.17+ $\frac{8T}{3}$

2/ Includes 780 Plecoptera which were not identified. See footnote 1/.

Appendix S

Summary of numbers and volumes (by order) of bottom fauna collected from the Blackfoot River and tributaries in 1971, 1972 and 1973 which were not identified to species. Volume in cc. T = trace

| Date | | <u>Pop's</u> | <u>Flesher</u> | <u>Alice Cr.
@ N. F.</u> | <u>Alice Cr.
@ 200</u> | <u>Hogum</u> | <u>Landers</u> | <u>Below
Landers</u> | <u>Poorman
Creek</u> | <u>Dalton</u> | <u>Canyon</u> | <u>Rainbow
Bend</u> |
|-------------|------|--------------|----------------|------------------------------|----------------------------|--------------|----------------|--------------------------|--------------------------|---------------|---------------|-------------------------|
| PLECOPTERA | | | | | | | | | | | | |
| Feb/
Mar | No. | | 912 | | 2282 | 942 | | 309 | 482 | 225 | 368 | 241 |
| 1972 | Vol. | | 2.4 | | 4.4 | 1.9 | | 1.0+T | 2.5 | 2.7 | 1.7 | 7.4 |
| May | No. | 50 | 133 | | 168 | 234 | 84 | 143 | 164 | 154 | 51 | |
| 1972 | Vol. | 3T | 0.4 | | 0.2+T | 0.9 | 0.3+T | 0.7 | 5.2 | 1.2 | 0.1+2T | |
| Aug | No. | 75 | 964 | 93 | 551 | 484 | | 294 | 231 | 212 | 195 | 173 |
| 1972 | Vol. | 0.1+2T | 1.7 | 0.1+2T | 1.4 | 1.8 | | 1.4 | 1.5 | 0.5 | 2.2 | 9.7 |
| Oct/
Nov | No. | 233 | 881 | | 477 | 463 | 255 | 362 | 708 | 158 | 391 | 278 |
| 1972 | Vol. | 0.6 | 3.3 | | 1.1 | 4.0 | 0.1+2T | 0.8 | 2.6 | 2.2 | 3.8 | 6.2 |
| May | No. | 63 | 75 | | 292 | | 89 | | | | 141 | |
| 1973 | Vol. | 3T | 3T | | 1.4 | | 0.2+T | | | | 1.1 | |

EPHEMEROPTERA

Feb/
Mar	No.		571		1541	1789		179	1893	1000	1214	1359
1972	Vol.		0.3+T		2.2	1.4		0.8+T	2.7	2.6	2.2	2.7
May	No.	0	30		347	94	391	671	1439	546	301	
1972	Vol.	0	3T		0.9	0.2+T	3.9	2.2	6.0	1.3	2.3	
Aug	No.	0	166	788	783	621		1894	1645	595	1052	851
1972	Vol.	0	0.1+2T	2.6	1.0	0.9		4.2	1.0	0.7	0.8	0.8
Oct/
Nov	No.	0	89		716	574	796	1621	1265	758	1510	1129
1972	Vol.	0	3T		1.2	0.9	1.2	2.8	2.3	2.2	2.6	1.5
May	No.	3	5		889		1204				1738	
1973	Vol.	3T	2T		2.6		11.0				3.3	

| <u>Date</u> | <u>Pop's</u> | <u>Flesher</u> | <u>Alice Creek
@ N. F.</u> | <u>Alice Creek
@ 200</u> | <u>Hogum</u> | <u>Landers</u> | <u>Below
Landers</u> | <u>Poorman
Creek</u> | <u>Dalton</u> | <u>Canyon</u> | <u>Rainbow
Bend</u> |
|-------------|--------------|----------------|--------------------------------|------------------------------|--------------|----------------|--------------------------|--------------------------|---------------|---------------|-------------------------|
|-------------|--------------|----------------|--------------------------------|------------------------------|--------------|----------------|--------------------------|--------------------------|---------------|---------------|-------------------------|

TRICOPTERA

Feb/
Mar	No.			786	1164		90	506	779	628	740
1972	Vol.			3.8	2.5		0.6	3.5	1.0	1.6	2.7
		296									
		1.4									
May	No.	0	9	377	136	688	16	316	422	134	
1972	Vol.	0	3T	1.9	0.7	6.7	0.1+2T	6.7	1.3	0.2+T	
Aug	No.	1	173	11	406	681	281	218	82	93	808
1972	Vol.	T	0.7	3T	2.7	1.4	2.4	0.3	0.1+2T	0.2+T	3.6
Oct/
Nov	No.	4	180		961	354	192	253	431	379	766
1972	Vol.	2T	0.4		3.0	1.9	0.8	2.9	5.5	1.5	4.4
May	No.	0	34	665		36				505	
1973	Vol.	0	0.2+T	3.0		1.7				2.0	

DIPTERA

Feb/
Mar	No.			2642	5483		41	607	169	94	448
1972	Vol.			3.2	2.1		1.5	1.1	1.1+T	0.5	0.6
		402									
		1.0									
May	No.	0	384	778	700	62	476	1991	472	174	
1972	Vol.	0	0.5	1.5	1.5	0.2+T	1.5	4.3	2.4	0.1+2T	
Aug	No.	11	1000	68	571	716	5715	800	496	234	129
1972	Vol.	3T	1.8	0.2+T	1.4	0.2+T	8.7	0.6	1.3+T	0.8	0.2+T
Oct/
Nov	No.	17	144	640	508	7	104	284	81	320	131
1972	Vol.	3T	0.1+2T	2.0	0.5	3T	0.9	0.9	0.5	1.1	0.4+T
May	No.	0	104	329		473				116	
1973	Vol.	0	0.1+2T	0.9		1.0				0.7	

APPENDIX S continued

| <u>Date</u> | <u>Pop's</u> | <u>Flesher</u> | <u>Alice Creek</u>
<u>@ N. F.</u> | <u>Alice Creek</u>
<u>@ 200</u> | <u>Hogum</u> | <u>Landers</u> | <u>Below</u>
<u>Landers</u> | <u>Poorman</u>
<u>Creek</u> | <u>Dalton</u> | <u>Canyon</u> | <u>Rainbow</u>
<u>Bend</u> |
|-------------|--------------|----------------|--------------------------------------|------------------------------------|--------------|----------------|--------------------------------|--------------------------------|---------------|---------------|-------------------------------|
|-------------|--------------|----------------|--------------------------------------|------------------------------------|--------------|----------------|--------------------------------|--------------------------------|---------------|---------------|-------------------------------|

COLEOPTERA

Feb/
Mar
1972	No.
Vol.		2
2T	102
0.2+T	159
0.3		5
2T	141
0.3	38
0.1	2
2T	2
2T
May
1972	No.
Vol.	0
0	0
0	71
3T	49
3T	0
0	2
T	314
0.4	4
3T	1
T	
Aug
1972	No.
Vol.	0
0	30
2T	1
T	79
0.2+T	181
0.1+2T	11
2T	240
0.1+2T	120
0.1+2T	13
3T	11
3T
Oct/
Nov
1972	No.
Vol.	0
0	4
2T		83
3T	119
0.1+2T	0
0	8
3T	86
0.2+T	3
2T	5
3T
May
1973	No.
Vol.	0
0	31
0.1+2T		21
3T		0
0			14
2T	

OLIGOCHAETA, NEMATODA & OTHERS

Feb/
Mar
1972	No.
Vol.		7
2T	15
2T	0
0		1
T	44
0.6	25
3T	32
3T	58
3T
May
1972	No.
Vol.	0
0	0
0	9
0.1+3T	4
2T	40
4T	1
T	91
2.4+2T	2
2T	0
0	
Aug
1972	No.
Vol.	0
0	16
3T	123
1.0+T	19
3T	5
2T	20
4T	61
3T	12
4T	7
2T	0
0
Oct/
Nov
1972	No.
Vol.	1
T	3
2T		6
3T	6
2T	0
0	11
4T	11
0.3+2T	3
2T	48
5T
May
1973	No.
Vol.	0
0	36
0.1+3T		7
0.2+3T		5
2T			0
0	

APPENDIX T

Species composition, size, and relative abundance of game and nongame fish from stream surveys, 1971-1973.

Fish were collected from larger waters with a Fisher-Shocker electrofishing unit mounted in a boat. Small streams were sampled with a Smith-Root Type IV backpack electrofisher. Total length of fish was measured to the nearest .10 inch, and weight to the nearest .01 pound.

Common and scientific names for the fish found in the study area and the abbreviations used are given below. Sculpin refers to the slimy sculpin, and Dace to the longnose dace.

<u>Abr.</u>	<u>Common Name</u>	<u>Scientific Name</u> ^{1/}
Ct	cutthroat trout	<u>Salmo clarki</u> Richardson
Rb	rainbow trout	<u>Salmo gairdneri</u> Richardson
LL	brown trout	<u>Salmo trutta</u> Linnaeus
Eb	brook trout	<u>Salvelinus fontinalis</u> (Mitchell)
DV	Dolly Varden	<u>Salvelinus malma</u> (Walbaum)
Wf	mountain whitefish	<u>Prosopium williamsoni</u> (Girard)
FSu	longnose sucker	<u>Catostomus catostomus</u> (Forster)
CSu	large scale sucker	<u>Catostomus macrocheilus</u> Girard
Sculpin	slimy sculpin	<u>Cottus cognatus</u> Richardson
Dace	longnose dace	<u>Rhinichthys cataractae</u> (Valenciennes)

^{1/} In accordance with "Common and Scientific Names of Fishes from the United States and Canada", 3rd ed., American Fisheries Society Special Publication No. 6, 1970.

APPENDIX T continued

<u>Stream</u>	<u>Location</u>			<u>Section</u>	<u>Species</u>	<u>No. in Sample</u>	<u>Length (in)</u>		<u>Weight (lbs)</u>		<u>Age Classes</u>
	<u>T</u>	<u>R</u>	<u>S</u>	<u>Length (ft)</u>			<u>Range</u>	<u>Average</u>	<u>Total</u>	<u>Average</u>	<u>Present</u>
Blackfoot River
at Pop's Place
August, 1973	15N	6W	19	1200	Ct	9	1.6-6.7	5.2	.49	.05	
					Eb	57	1.9-7.9	3.9	1.96	.03	
					Sculpin	4	3.1-4.7	3.7	.10	.03	
Blackfoot River -
below Pop's Place
section <u>1</u> /
Sept-Oct, 1971	15N	6W	19,	3486	Ct	219	1.8-9.8	4.4	9.81	.04	0 - III
					Eb	134	2.2-10.4	5.0	9.44	.07	0 - III
	15N	7W	24		FSu	2	5.7-6.7	-	.21	.10	
					Sculpin	6	not recorded		not recorded		
	Blackfoot River
Flesher section ^{1/}
Aug-Sept, 1973	15N	7W		35	2455	Ct	185	1.5-10.9	3.7	6.55
Eb				108			2.0-9.1	3.8	3.81	.04	
Wf				10			2.7-3.3	3.0	.10	.01	
FSu				8			4.5-6.5	5.4	.53	.07	
Sculpin				43			1.2-4.3	3.1	.66	.02	
Blackfoot River -
below Willow Creek
Section <u>1</u> /
Aug-Sept, 1973	15N	7W	34,
35	1500	Ct	69	1.6-11.7	4.8	4.53	.06	
					Eb	102	2.0-11.0	4.4	5.30	.05	
					Wf	1		2.6		.01	
					FSu	14	4.3-7.1	5.7	1.06	.08	
					Sculpin	64	1.7-5.3	2.5	.59	.01	
					Dace	1	4.3			.04	

1/ See population estimate results in Appendix U .

APPENDIX T

continued

| <u>Stream</u> | <u>Location</u> | | | <u>Section
Length (ft)</u> | <u>Species</u> | <u>No. in
Sample</u> | <u>Length (in)</u> | | <u>Weight (lbs)</u> | | <u>Age Classes
Present</u> |
|-------------------------------------|-----------------|----------|----------|--------------------------------|----------------|--------------------------|--------------------|--------------------------------------|---------------------|----------------|--------------------------------|
| | <u>T</u> | <u>R</u> | <u>S</u> | | | | <u>Range</u> | <u>Average</u> | <u>Total</u> | <u>Average</u> | |
| Blackfoot River
Hogum section 1/ | 14N | 7W | 5 | 4100 | Ct | 131 | 1.7-13.5 | 3.7 | 8.05 | .06 | 0 - IV |
| | | | | | Eb | 74 | 2.7-9.9 | 5.2 | 5.18 | .07 | 0 - II |
| | | | | | Rb(H)* | 3 | 9.4-10.6 | 10.1 | 1.25 | .42 | |
| | | | | | LL | 1 | | 6.5 | | .08 | II |
| | | | | | DV | 7 | 6.5-16.8 | 9.6 | 2.99 | .43 | II - IV |
| Aug-Sept, 1972 | | | | | Wf | 106 | 2.6-15.9 | 10.0 | 52.24 | .49 | 0 - V |
| | | | | | FSu | 9 | 4.7-6.7 | 5.8 | .66 | .07 | |
| | | | | | CSu | 5 | 5.3-11.6 | 8.1 | 1.30 | .26 | |
| | | | | | Sculpin | 422 | 1.1-4.8 | (only smallest and largest measured) | | | |
| | | | | | Dace | 19 | 2.5-5.6 | 4.3 | .61 | .03 | |
| * * * | | | | | | | | | | | |
| Sept, 1973 1/ | 14N | 7W | 5 | 4100 | Ct | 87 | 1.7-13.6 | 4.4 | 5.84 | .07 | 0 - III |
| | | | | | Eb | 84 | 2.6-8.1 | 3.8 | 2.67 | .03 | 0 - II |
| | | | | | Rb(H)* | 18 | 10.8-12.3 | 11.4 | 10.35 | .58 | |
| | | | | | DV | 3 | 6.8-13.6 | 11.2 | 1.98 | .66 | I, II, IV |
| | | | | | Wf | 31 | 2.7-13.5 | 3.8 | 2.05 | .07 | 0, III, IV |
| | | | | | FSu | 6 | 2.5-6.4 | 4.7 | .32 | .05 | |
| | | | | | Dace | 6 | 3.0-5.0 | 4.1 | .17 | .03 | |
| | | | | | Sculpin | numerous | not measured | | not measured | | |

*Hatchery raised to catchable size before stocking

APPENDIX T

continued

[illegible]

continued

<u>Stream</u>	<u>Location</u>			<u>Section</u>	<u>Species</u>	<u>No. in Sample</u>	<u>Length (in)</u>		<u>Weight (lbs)</u>		<u>Age Classes Present</u>
	<u>T</u>	<u>R</u>	<u>S</u>	<u>Length (ft)</u>			<u>Range</u>	<u>Average</u>	<u>Total</u>	<u>Average</u>	
<u>Beartrap Creek</u>											
(1) above White Hope Mine
August 16, 1972	15N	6W	27	400	Ct	13	2.9-4.4	3.5	.25	.02	
	two sections combined										
(2) between tailings dam and Mike Horse Creek
July 26, 1972	15N	6W	27	300	Ct	36	4.0-9.4	6.9	4.74	.13	
(3) below Mike Horse Creek
July 26, 1972	15N	6W	27	400	No fish caught or observed. Mine drainage enters from Mike Horse Creek.						
Blackfoot River below confluence of Beartrap and Anaconda Creeks
August 16, 1972	15N	6W	28	350	No fish caught or observed. Mine drainage still prevails in this section.						

APPENDIX T continued

| <u>Stream</u> | <u>Location</u> | | | <u>Section
Length (ft)</u> | <u>Species</u> | <u>No. in
Sample</u> | <u>Length (in)</u> | | <u>Weight (lbs)</u> | | <u>Age Classes
Present</u> |
|---|-----------------|----------|-----------------------|--|---------------------|--------------------------|-------------------------------|-------------------|----------------------|-------------------|--------------------------------|
| | <u>T</u> | <u>R</u> | <u>S</u> | | | | <u>Range</u> | <u>Average</u> | <u>Total</u> | <u>Average</u> | |
| Anaconda Creek
July 26, 1972 | 15N | 6W | 27 | 250 | Ct
Eb | 37
2 | 1.7-7.0
5.5-6.9 | 4.8 | 1.95
.21 | .05
.11 | |
| Shoue (Shave)
Gulch
July 26, 1972 | 15N | 6W | 21 | 800 | Ct | 116 | 2.2-6.7 | 4.3 | 4.17 | .04 | 0 - II |
| Pass Creek
July 27, 1972 | 15N | 6W | 16 | 920 | Ct | 83 | 1.4-6.4 | 3.1 | 1.30 | .02 | 0 - II |
| | | | two sections combined | | Eb | 30 | 3.5-6.6 | 3.5 | 1.58 | .05 | 0 - II |
| | | | | | Sculpin | 3 | 1.5-3.0 | 2.3 | .02 | .01 | |
| Cadotte Creek
August 29, 1973 | 15N | 7W | 13 | shocked in
scattered pools
and beaver dams
only | Eb | 13 | 7.1-10.7 | 8.5 | 3.73 | .29 | |
| Willow Creek
(Fletcher) | | | | | | | | | | | |
| (1) above Sandbar
Creek
August 21, 1973 | 14N | 6W | 8 | 1400 | Ct
Eb
Sculpin | 224
102
167 | 2.6-7.7
1.9-8.6
1.5-4.5 | 4.2
3.9
2.8 | 7.72
3.06
1.03 | .03
.03
.01 | |
| (2) below
Sandbar Creek
August 22, 1973 | 14N | 7W | 1 | 900 | Ct
Eb
Sculpin | 82
58
166 | 1.6-9.0
2.3-7.9
1.8-4.6 | 4.9
4.9
2.6 | 4.30
3.00
1.09 | .05
.05
.01 | |

* * *

* * *

APPENDIX T

continued

| <u>Stream</u> | <u>Location</u> | | | <u>Section
Length (ft)</u> | <u>Species</u> | <u>No. in
Sample</u> | <u>Length (in)</u> | | <u>Weight (lbs)</u> | | <u>Age Classes
Present</u> |
|---|-----------------|----------|----------|---------------------------------|----------------|--------------------------|--------------------|----------------|---------------------|----------------|--------------------------------|
| | <u>T</u> | <u>R</u> | <u>S</u> | | | | <u>Range</u> | <u>Average</u> | <u>Total</u> | <u>Average</u> | |
| (3) at mouth
August 24, 1973 | 15N | 7W | 35 | 1028 | Ct | 48 | 1.8-11.3 | 5.1 | 2.93 | .06 | |
| | | | | | Eb | 36 | 2.3-11.3 | 5.5 | 3.15 | .09 | |
| | | | | | FSu | 2 | 4.6-7.0 | | .17 | .08 | |
| | | | | | Sculpin | 38 | 1.8-4.3 | 2.7 | .21 | .00+ | |
| <u>Alice Creek</u> | | | | | | | | | | | |
| (1) at USFS Guard
Station
July 25, 1972 | 16N | 7W | 14 | 790
(.21 acre) | Ct | 94 | 2.0-8.6 | 5.7 | 6.85 | .07 | 0 - II |
					Eb	1	9.0		.39		
					Sculpin	16	3.0-4.3	3.8	.43	.03	
* * *											
(2) below Wildcat
Gulch $\frac{1}{2}$ mile
August 1, 1972	16N	7W	22	500	Ct	1	4.5		.05		
					Eb	1	8.2		.22		
					Sculpin	23	2.5-4.2	3.3	.40	.02	
* * *											
(3) above National
Forest Boundary
$\frac{1}{2}$ mile
August 1, 1972	16N	7W	27	490
two sections
combined	Ct	54	2.1-9.0	3.9	1.93	.03	0 - II
					Eb	15	5.1-10.0	6.5	2.02	.13	I - III
					Sculpin	5	2.4-3.8	3.0	.08	.02	
* * *											
(4) on Fickler
Ranch
August 17, 1972	15N	7W	4	700	Ct	12	3.0-8.3	4.8	.58	.05	0 - II
					Eb	73	1.9-11.3	5.6	7.64	.10	0 - III
					Wf	4	7.8-12.9	10.7	1.97	.49	I, II, V
					FSu	5	3.2-10.0	5.4	.40	.08	
					Sculpin	25	1.7-4.1	3.0	.40	.02	

APPENDIX T

continued

| <u>Stream</u> | <u>Location</u> | | | <u>Section
Length (ft)</u> | <u>Species</u> | <u>No. in
Sample</u> | <u>Length (in)</u> | | <u>Weight (lbs)</u> | | <u>Age Classes
Present</u> |
|---|-----------------|----------|----------|--------------------------------|----------------|---|--------------------|-----------------------|---------------------|----------------|--------------------------------|
| | <u>T</u> | <u>R</u> | <u>S</u> | | | | <u>Range</u> | <u>Average</u> | <u>Total</u> | <u>Average</u> | |
| (5) Bartlett
Section <u>1</u> /
September, 1971 | 15N | 7W | 27 | 1880 | Ct | 353 | 1.7-9.7 | 4.8 | 18.74 | .05 | 0 - III |
					Eb	163	2.5-12.5	4.8	9.66	.06	0 - II
					DV	1		6.8		.10	
					Wf	77	2.5-10.2	4.6	4.19	.05	0 - III
					FSu	1		4.3		.04	
					Sculpin	214	?-5.0	only largest measured			
* * *											
(6) Bartlett
Section <u>1</u> /
August, 1972	15N	7W	27	2779	Ct	271	1.6-10.5	4.9	16.69	.06	0 - III
					Eb	166	2.1-10.3	4.8	10.61	.06	0 - III
					DV	2	5.4-7.1		.15	.08	
					Wf	7	2.3-7.1	3.9		.03	
					FSu	10	5.0-7.3	5.8		.07	
					CSu	1		8.5		.23	
Sculpin						numerous-not counted or measured					
* * *											
(7) Bartlett
Section <u>1</u> /
September, 1973	15N	7W	27	2295	Ct	384	1.6-10.5	4.4	16.85	.04	0 - III
					Eb	273	2.1-12.0	4.1	12.16	.04	0 - III
					DV	2	6.8-9.9		.47	.24	II, II
					Wf	10	2.5-3.5	3.1		.04	all age 0
					FSu	19	3.8-6.1	4.7		.04	
					CSu	26	3.7-6.9	5.1		.06	
Sculpin						numerous
4.4-5.8 4.7 (only six were measured)					
<u>Hardscrabble Creek</u>											
(1) at Spencer Bar
July 31, 1972	15N	7W	21	450	Eb	1		9.6		.36	
					FSu	15	1.4-5.5	3.6	.33	.02	

* * *

APPENDIX T continued

| <u>Stream</u> | <u>Location</u> | | | <u>Section
Length (ft)</u> | <u>Species</u> | <u>No. in
Sample</u> | <u>Length (in)</u> | | <u>Weight (lbs)</u> | | <u>Age Classes
Present</u> | |
|--|-----------------|----------|-----------|--------------------------------|----------------|--------------------------|--|----------------|---------------------|----------------|--------------------------------|--|
| | <u>T</u> | <u>R</u> | <u>S</u> | | | | <u>Range</u> | <u>Average</u> | <u>Total</u> | <u>Average</u> | | |
| (2) up Alice
Creek road $\frac{1}{2}$ mile
(below beaver dams)
August 1, 1972 | 15N | 7W | 28,
33 | 400 | Ct | 2 | 3.5-3.7 | | .03 | .02 | | |
					Eb	93	1.5-6.7	5.1	2.90	.03		
					Wf	3	2.4-5.1	4.0	.06	.02		
					FSu	20	3.5-5.8	4.6	.74	.04		
					Sculpin	203	1.5-4.6 (only largest and smallest measured)					
Many additional small sculpins not collected.												
* * *												
(3) near mouth
(north of highway)
July 31, 1972	15N	7W	33	430	Eb	64	1.8-7.6	3.9	1.95	.03	0 - II	
					Wf	6	2.4-5.4	3.7	.10	.02	0, I	
					FSu	8	3.7-5.3	4.5	.24	.03		
					Sculpin	154	1.6-4.0	2.6	1.44	.01		
Hogum Creek
September 10, 1973	14N	7W	8	1051	Ct	52	1.9-8.4	4.8	2.67	.05		
					Eb	34	2.3-8.3	5.6	2.81	.08		
					DV	2	5.5-7.7		.21	.10		
					FSu	1		5.0		.05		
					Sculpin	59	1.6-4.2	2.7	.43	.01		
<u>Copper Creek</u>												
(1) at Sucker
Creek Road
August 23, 1973	15N	8W	25,
26	1225	Ct	5	4.7-11.5	8.1	1.44	.29		
					DV	9	4.0-8.3	5.3	.47	.05		
					Sculpin	25	2.3-4.7	3.2	.37	.01		
* * *												
(2) near mouth
August 22, 1973	15N	8W	36	2100	Ct	2	3.2-3.4		.02	.01		
					Eb	3	2.4-3.0	2.6	.01	.00+		
					DV	10	2.4-6.0	3.6	.19	.02		
					Wf	1		2.9		.00+		
					Sculpin	23	2.3-3.7	3.0	.26	.01		

APPENDIX T continued

| Stream | Location | | | Section
Length (ft) | Species | No. in
Sample | Length (in) | | Weight (lbs) | | Age Classes
Present |
|--|----------|-----|----|------------------------|---------|---|--|--------------------------------------|--------------|---------|------------------------|
| | T | R | S | | | | Range | Average | Total | Average | |
| <u>Landers Fork</u> | | | | | | | | | | | |
| (1) below Copper Creek road bridge
September 10, 1973 | 15N | 8W | 36 | 1500 | | | No fish caught or observed. Habitat water present only in pools, with very little flow in between. Stream dry at end of section. | | | | |
| * * * | | | | | | | | | | | |
| (2) at Highway 200
September 10, 1973 | 14N | 8W | 12 | 2500 | Ct | 15 | 2.9-7.3 | 4.9 | .85 | .06 | |
| | | | | | Eb | 7 | 3.4-8.6 | 6.9 | 1.19 | .17 | |
| | | | | | DV | 2 | 2.9-6.5 | | .08 | .04 | |
| | | | | | Sculpin | 12 | 2.2-4.4 | 3.0 | .17 | .01 | |
| <u>Poorman Creek</u> | | | | | | | | | | | |
| (1) Circle City
Section 1/
September, 1972 | 13N | 8W | 6 | 1760 | Ct | 570 | 1.6-10.3 | 4.8 | 30.31 | .05 | 0 - III |
					Eb	77	2.6-11.2	4.5	4.08	.05	0 - II
					DV	2	7.1-11.3		.68	.34	both age II
					Wf	36	3.7-13.9	10.6	20.65	.57	0 - III
					LL	1		4.8		.04	and older
					Sculpin	112	1.3-5.4	(only largest and smallest measured)			
* * *											
(2) at Circle City
September 13, 1973	13N	8W	6	500	Ct	80	1.6-11.3	4.8	4.67	.06	
					Eb	42	2.0-8.0	3.3	.96	.02	
					DV	5	12.3-16.4	14.6	5.58	1.12	
					Wf	1		9.1		.25	
					Sculpin	numerous - not collected					
Arastra Creek
July 24, 1972	15N	10W	24	±200	Ct	15	5.0-8.7	6.7	2.02	.13	
					DV	2	8.8-12.0		.76	.38	
						(5 Ct and 1 DV caught by hook and line)					

APPENDIX U

Population estimates were derived by the Petersen mark-recapture technique using the method described by Vincent (1971 and 1974). Estimates made prior to September 28, 1973 were derived from the following formula using a computer program:

$$N = \frac{(M+1)(C+1)}{R+1}$$

Where

N = Population estimate
M = Number of fish marked
C = Number of fish in recapture sample
R = Number of marked fish in recapture sample (C)

Estimates made subsequent to September 28, 1973 were derived by using Chapman's (1951) modification of the Petersen formula in the computer program as follows:

$$N = \frac{(M+1)(C+1)}{R+1} - 1$$

In this study, however, scales were collected at 1.0 inch length intervals instead of 1/10 inch intervals as described by Vincent (1974).

When possible, the population was divided into more than one length group for purposes of making the estimate. Whether this was possible depended on the number of marked recoveries (R) obtained for the various sizes of fish.

Where recoveries were low, length groups often had to be combined to obtain a reliable estimate of the total fish population. The purpose of this procedure was to obtain the most reliable estimate of the total population. Consequently length group breakdown is not necessarily consistent for the same station for different years because of differences in efficiency of sampling the various sizes of fish each year. Twenty scales were collected for each inch-class whenever possible.

Confidence limits are given at the 80 percent probability level. Figures in parentheses are the confidence limits in percent. The level of accuracy sought was 25 percent.

Confidence limits for the total are not the sum of the confidence limits for each component; rather they are approximately the square root of the sum of their squares.

Total-number figures are only for those fish species and age classes which could be estimated. In some cases other game fish were present in the population but were too few in number to be reliably estimated, or they were not thought to be permanent residents of the stream section.

Species	Age Class	Average Length (in)	Average Weight (lbs)	Number Estimate	Weight Estimate (lbs)
Blackfoot River below Pop's Place, 1971
(Section Length 3486 feet)					
Ct	0	Present, but recoveries insufficient for estimate			
	I	4.4	.03	131	3.87
	II	6.2	.09	89	8.32
	III and older	7.5	.18	13	2.27
Total				233	14.46
Total per 1000 feet				67	4.15
Eb	0	Present, but recoveries insufficient for estimate			
	I	4.7	.04	50	1.96
	II	6.4	.11	112	12.29
	III and older	7.9	.20	22	4.31
Total				184±45 (24%)	18,56±5 (27%)
Total per 1000 feet				57	5.32
Grand Total - yearlings and older				417	33.02
Grand Total per 1000 feet - yearlings and older				120	9.47

APPENDIX II continued

Species	Age Class	Length Group	Number Estimate	Weight Estimate (lbs)
Blackfoot River, Fleisher Section, 1973
(Section Length 2455 feet)				
Ct	not aged	1.5 - 4.9	380	6
	aged	5.0 - 10.9	58	6
Total			438±99 (23%)	12±2 (17%)
Total per 1000 feet			178	4.89
Eb	not aged	2.0 - 4.9	231	3
	aged	5.0 - 9.1	35	3
Total			266±84 (32%)	6±2 (33%)
Total per 1000 feet			108	2.44
Grand Total			704±130 (18%)	18±3 (17%)
Grand Total per 1000 feet			287	7.33

Blackfoot River, below Willow Creek Section, 1973
(Section Length 1500 feet)				
Ct	not aged	1.6 - 4.9	60	1
	aged	5.0 - 11.7	39	5
Total			99±20 (20%)	6±1 (17%)
Total per 1000 feet			66	4
Eb	not aged	2.0 - 4.9	193	4
	aged	5.0 - 11.0	42	5
Total			235±74 (31%)	9±2 (22%)
Total per 1000 feet			157	6
Grand Total			334±77 (23%)	15±2 (13%)
Grand Total per 1000 feet			223	10

Species	Age Class	Average Length (in)	Average Weight (lbs)	Number Estimate	Weight Estimate (lbs)
Alice Creek - Bartlett Section, 1971
(Section Length 1880 feet)					
Ct	0	3.0	.01	295	3.52
	I	4.7	.04	360	14.07
	II	6.8	.12	102	12.27
	III and older	9.0	.28	6	1.71
Total				763±131 (17%)	31.57±4 (13%)
Total per 1000 feet				406	16.79
Eb	0	3.2	.01	105	1.55
	I	5.1	.06	204	11.22
	II and older	7.4	.19	41	7.97
Total				350±74 (21%)	20.74±4 (19%)
Total per 1000 feet				186	11.03
Grand Total				1113±150 (13%)	52.31±6 (11%)
Grand Total per 1000 feet				592	27.82
Grand Total - yearlings and older				713	47.24
Grand Total per 1000 feet - yearlings and older				379	25.12

Alice Creek - Bartlett Section, 1972
(Section Length 2779 feet)					
Ct	0	3.5	.02	338	6.52
	I	5.4	.06	190	11.41
	II	7.7	.18	54	9.54
	III and older	9.4	.32	10	3.18
Total				592±114 (19%)	30.65±5 (16%)
Total per 1000 feet				213	11.03
Eb	0	2.7	.01	119	1.27
	I	5.0	.05	141	6.85
	II	7.3	.17	37	6.39
	III and older	8.6	.29	9	2.60
Total				306±65 (21%)	17.11±3 (18%)
Total per 1000 feet				110	6.16
Grand Total				898±131 (15%)	47.76±6 (13%)
Grand Total per 1000 feet				511	17.16
Grand Total - yearlings and older				441	30.97
Grand Total - yearlings and older per 1000 feet				159	14.36

APPENDIX II continued

Species	Age Class	Average Length (in)	Average Weight (lbs)	Number Estimate	Weight Estimate (lbs)
Alice Creek - Bartlett Section, 1973
(Section Length 2295 feet)					
Ct	0	Present, but recoveries insufficient for estimate			
	I	5.5	.06	155	9.11
	II and older	7.5	.17	45	7.71
Total				204±20 (10%)	16.82±2 (12%)
Total per 1000 feet				91	7.33
Eb	0	2.9	.01	427	3.73
	I	5.4	.06	107	6.37
	II	8.3	.26	19	4.89
	III and older	12.0	.81	1	0.88
Total				554±109 (20%)	15.87±2 (13%)
Total per 1000 feet				241	6.92
Grand Total				763±111 (14%)	32.69±3 (9%)
Grand Total per 1000 feet				332	14.24
Grand Total - yearlings and older				336	28.96
Grand Total - yearlings and older per 1000 feet				146	12.62

191-V

Blackfoot River - Hugum Section, 1972
(Section Length 4100 feet)					
Ct	0	2.6	.01	203	2.99
	I	6.8	.13	19	2.65
	II and older	10.6	.55	9	4.79
Total				231±52 (22%)	10.38±1 (10%)
Total per 1000 feet				56	2.53
Eb	0	3.5	.02	46	0.89
	I	6.4	.10	57	5.76
	II and older	9.2	.33	2	0.71
Total				105±20 (19%)	7.36±1 (14%)
Total per 1000 feet				26	1.80
Grand Total				336±55 (16%)	17.74±1 (7%)
Grand Total per 1000 feet				82	4.33
Grand Total - yearlings and older				28	13.86
Grand Total - yearlings and older per 1000 feet				7	3.38

Species	Age Class	Average Length (in)	Average Weight (lbs)	Number Estimate	Weight Estimate (lbs)
Blackfoot River - Hugum Section, 1973
(Section Length 4100 feet)					
Ct	0	2.8	.01	90	1.18
	I	5.3	.07	59	4.08
	II	9.2	.34	3	1.15
	III and older	13.6	1.12	1	1.28
Total				153±42 (27%)	7.69±1 (13%)
Total per 1000 feet				37	1.88
Eb	0	3.3	.01	178	2.55
	I	6.8	.13	22	2.95
	II and older	8.1	.21	5	1.00
Total				205±68 (34%)	6.50±2 (31%)
Total per 1000 feet				50	1.38
Grand Total				358±81 (23%)	14.19±2 (14%)
Grand Total per 1000 feet				87	3.46
Grand Total - yearlings and older				90	10.46
Grand Total - yearlings and older per 1000 feet				22	2.55

Pooman Creek - Circle City Section, 1972
(Section Length 1760 feet)					
Ct	0	3.8	.02	743	16.32
	I	6.0	.08	314	24.39
	II	7.7	.18	54	9.75
	III and older	9.6	.35	13	4.52
Total				1124±123 (11%)	54.98±6 (11%)
Total per 1000 feet				639	31.24
Total - yearlings and older				361	38.66
Total - yearlings and older per 1000 feet				216	21.96

APPENDIX II continued

Species	Age Class	Average Length (in)	Average Weight (lbs)	Number Estimate	Weight Estimate (lbs)
Blackfoot River, Poorman-Dalton Section, 1971
(Section Length 5350 feet)					
Eb	0	3.4	.02	219	3.89
	I	6.5	.11	58	6.42
	II and older	9.6	.36	10	3.73
	Total			287±122 (42%)	14,046 (43%)
Total per 1000 feet				54	2.62
LL	0	3.1	.02	530	8.01
	I	6.3	.10	576	58.96
	II	8.0	.21	106	21.91
	III	11.5	.65	116	75.13
	IV and older	14.2	1.07	147	197.40
Total				1475±408 (28%)	321,41±70 (22%)
Total per 1000 feet				275	60.08
Grand Total				1762±426 (24%)	335,45±70 (21%)
Grand Total per 1000 feet				329	62.70
Grand Total - yearlings and older				1013	323.55
Grand Total - yearlings and older per 1000 feet				189	60.47

Blackfoot River, Poorman-Dalton Section, 1972
(Section Length 5700 feet)					
Ct	0	4.3	.03	74	2.32
	I	8.8	.25	9	2.18
	II and older	11.3	.56	5	2.89
Total				88±24 (27%)	7.39±2 (27%)
Total per 1000 feet				15	1.30
Eb	0	Present, but recoveries insufficient for estimate			
	I	6.5	.10	93	9.69
	II and older	9.7	.36	15	5.50
Total				108±27 (25%)	15.19±4 (26%)
Total per 1000 feet				19	2.66
LL	0	Present, but recoveries insufficient for estimate			
	I	5.7	.07	834	32.34
	II	10.0	.38	84	32.05
	III and older	14.1	1.14	198	224.93
Total				716±111 (16%)	269,52±45 (16%)
Total per 1000 feet				126	50.79
Grand Total				912±117 (13%)	312.10±55 (14%)
Grand Total per 1000 feet				160	54.75
Grand Total - yearlings and older				836	309.78
Grand Total - yearlings and older per 1000 feet				147	54.35

Species	Age Class	Average Length (in)	Average Weight (lbs)	Number Estimate	Weight Estimate (lbs)
Blackfoot River, Canyon Section, 1971
(Section Length 10725 feet)					
LL	0	Present, but recoveries insufficient for estimate			
	I	4.2	.04	193	6.88
	II	7.3	.16	116	18.08
	III and older	14.6	1.19	277	329.49
Total - yearlings and older				586±139 (24%)	354,45±84 (24%)
Total - yearlings and older per 1000 feet				55	33.05

REPLY TO
ATTN OF

APPENDIX V
U.S. ENVIRONMENTAL PROTECTION AGENCY

REGION X
1200 SIXTH AVENUE
SEATTLE, WASHINGTON 98101

Concentration of Metals in
Samples of Fish from Blackfoot
River drainage

U. S. Environmental Protection Agency
Region X Laboratory
15345 N. E. 36th Street
Redmond, Washington 98052

November 2, 1973

Mr. Litter E. Spence, Jr.,
Environmental Planning Ecologist
Department of Fish and Game
State of Montana
Box 27
Ovanda, Montana 59854

Dear Mr. Spence:

In the attached table are results for metal determinations on fish tissue sent to us about the middle of August.

Please note that the last three digits in the laboratory number correspond to your sample number. Since the size of the fish was relatively small and upon thawing the tissue tended to be in a disintegrated state, it was not possible to determine metals in the various organs.

It is hoped that this information will assist you in your study.

Very truly yours,

Arnold R. Gahler

Arnold R. Gahler
Supervisory Chemist

arg/jp

Attachment

cc: F. Nelson
G. G'Neal

Concentrations expressed in micrograms per gram, wet weight.

(composite of 7 fish)

| St. No. | Section and Date | Species | Length
(in) | Weight
(lbs) | Cd | Cu | Pb | Hg | Ni | Zn | As |
|---------|------------------------|---------|----------------|-----------------|-----|------|----|-----|-----|----|-----|
| 33001 | Indian Creek | WF | 11.7 | .53 | .24 | 3.61 | 4 | .23 | 4.8 | 17 | .13 |
| 33002 | Indian Creek | WF | 11.7 | .10 | .30 | 1.88 | 2 | .06 | 3.4 | 12 | .16 |
| 33003 | L-25-72 | CF | 11.7 | .16 | .15 | 1.67 | 1 | .03 | 1.3 | 6 | .23 |
| 33004 | L-25-72 | CF | 10.6 | .16 | .24 | 1.38 | 1 | .02 | 1.6 | 20 | .12 |
| 33005 | " | CF | 8.3 | .18 | .18 | 6.59 | 3 | .01 | 3.6 | 25 | .42 |
| 33006 | " | CF | 7.8 | .17 | .25 | 1.58 | 2 | .03 | 1.5 | 26 | .15 |
| 33007 | Indian Creek | WF | 11.9 | .58 | .10 | 1.36 | 2 | .02 | 1.6 | 7 | .15 |
| 33008 | Indian Creek | WF | 10.6 | .40 | .10 | .78 | 2 | .04 | 1.2 | 21 | .11 |
| 33009 | (below Bartlett Creek) | CF | 8.9 | .29 | .15 | 1.27 | 2 | .03 | 1.7 | 24 | .15 |
| 33010 | L-25-72 | CF | 7.8 | .20 | .20 | 1.66 | 2 | .04 | 1.6 | 21 | .11 |
| 33011 | " | CF | 9.0 | .31 | .15 | 1.98 | 2 | .04 | 1.7 | 24 | .11 |
| 33012 | " | CF | 6.7 | .11 | .30 | 2.77 | 2 | .02 | 3.8 | 24 | .25 |
| 33013 | Blackfoot River | CF | 14.3 | .11 | .15 | .78 | 1 | .05 | 1.5 | 7 | .16 |
| 33014 | (Indian Creek) | CF | 11.3 | .58 | .10 | .68 | 1 | .02 | 1.3 | 6 | .40 |
| 33015 | 9-1-72 | CF | 12.8 | .77 | .10 | .78 | 1 | .03 | 1.3 | 7 | .25 |
| 33016 | " | CF | 13.9 | .12 | .10 | .78 | 1 | .02 | 1.3 | 7 | .14 |
| 33017 | " | CF | 11.7 | .17 | .10 | .69 | 1 | .02 | 1.3 | 6 | .14 |
| 33018 | " | CF | 11.4 | .61 | .10 | .69 | 1 | .02 | 1.3 | 6 | .14 |
| 33019 | " | CF | 11.7 | .61 | .10 | .69 | 1 | .02 | 1.3 | 6 | .14 |
| 33020 | " | CF | 11.4 | .61 | .10 | .69 | 1 | .02 | 1.3 | 6 | .14 |
| 33021 | Blackfoot River | CF | 11.4 | .120 | .10 | 1.28 | 1 | .03 | 2.5 | 7 | .11 |
| 33022 | (Indian Creek) | CF | 12.6 | .17 | .10 | .79 | 1 | .02 | 3.0 | 7 | .02 |
| 33023 | 9-1-72 | WF | 11.5 | 1.16 | .10 | 1.09 | 1 | .02 | .9 | 6 | .16 |
| 33024 | " | WF | 12.1 | .75 | .10 | 1.58 | 1 | .02 | 2.6 | 6 | .15 |
| 33025 | Indian Creek | CF | 11.5 | .15 | .15 | .69 | 1 | .02 | .6 | 13 | .09 |

APPENDIX V continued
ENVIRONMENTAL PROTECTION AGENCY

REGION X
1200 SIXTH AVENUE
SEATTLE, WASHINGTON 98101

U.S. Environmental Protection Agency
Region X Laboratory
15345 N.E. 36th Street
Redmond, Washington 98052

January 28, 1974

Mr. Liter E. Spence, Jr.
Planning Ecologist
Department of Fish and Game
State of Montana
Ovando, Montana 59854

Dear Mr. Spence,

In reply to your letter concerning the iron values on fish tissue samples, no iron values were obtained by the laboratory. When the samples were being digested prior to analysis by atomic absorption, the chemist unfortunately placed some acid not specially purified for iron in the digestion mixture so that the iron values would not have been reliable.

We apologize that this occurred, but the other values should be reliable.

Very truly yours,

Arnold R. Gahler

Arnold R. Gahler
Supervisory Chemist

AG:jn

U.S. ENVIRONMENTAL PROTECTION AGENCY
REGION X

1200 SIXTH AVENUE
SEATTLE, WASHINGTON 98101

Region X Laboratory
1555 Alaskan Way South
Seattle, Washington 98134
May 15, 1975

REPLY TO
ATTN OF:

Liter E. Spence
Habitat Evaluation Leader
State of Montana
Department of Fish & Game
Helena, Montana 59601

Dear Mr. Spence:

In response to your letter of May 6, 1975 we reviewed the bench data sheets. Each fish sample was homogenized thoroughly in a Waring blender and then dried in an oven at 60°C. After the samples were dried, they were mixed thoroughly again. A 5.00 gram portion of dried fish was digested with concentrated nitric and hydrochloric acids until all organic matter was digested. This solution was then diluted to a known volume with deionized water and analyzed directly with an Atomic Absorption Spectrophotometer for cadmium, copper, lead, nickel, and zinc (EPA). Half of this solution, which represented 2.50 grams of dried fish, was redigested with concentrated sulfuric acid to remove all nitric acid and was analyzed for arsenic by the Gutzeit method (Standard Methods.) A 1.00 gram portion of the dried fish was digested and analyzed for mercury by the cold-flame technique (EPA).

The following table summarizes both wet-weight and dry-weight results for all parameters requested:

The following references were used:
(EPA) Methods for Chemical Analysis of Water and Wastes, 1971

(Standard Methods) Standard Methods for the Examination of Water and Wastewater, 13th Edition, 1971.

If we can be of further assistance, please let us know.

Sincerely,

Robert H. Rieck

Robert H. Rieck
Chemist

cc: Mr. Arnold Gahler
Mr. Francis Nelson

APPENDIX V continued

| Lab
Number	Lead		Mercury		Nickel		Zinc		Arsenic		Cadmium		Copper	
	Wt Basis	Ory Wt Basis	Wt Basis	Ory Wt Basis	Wt Basis	Ory Wt Basis	Wt Basis	Ory Wt Basis	Wt Basis*	Ory Wt Basis*	Wt Basis	Ory Wt Basis	Wt Basis	Ory Wt Basis
33001	4	16	.23	.89	4.8	19.4	17	68	.13	.51	.24	.99	3.61	14.62
33002	2	9	.06	.23	3.5	14.0	12	49	.16	.64	.30	1.20	1.88	7.60
33003	1	3	.03	.11	1.3	5.2	6	26	.23	.95	.15	.60	1.67	6.75
33004	1	3	.02	.09	1.6	6.4	15	62	.14	.56	.10	.40	1.38	5.60
33005	3	11	.04	.16	3.8	15.3	28	112	.62	2.51	.24	.98	6.59	26.67
33006	2	9	.03	.12	5.4	22.0	26	104	.15	.60	.25	1.00	1.58	6.40
33007	< 1	2	.02	.09	1.8	7.1	7	29	.08	.35	.10	.39	1.36	5.49
33008	1	3	.02	.07	.9	3.5	9	36	.18	.71	.10	.39	.78	3.14
33009	2	6	.04	.14	1.6	6.3	21	83	.14	.56	.15	.60	1.27	5.16
33010	2	8	.03	.12	3.6	14.4	22	90	.15	.60	.20	.80	1.68	6.80
33011	2	6	.04	.14	1.7	6.8	14	57	.14	.56	.15	.60	1.98	8.00
33012	2	6	.04	.15	1.0	4.0	22	90	.11	.44	.15	.60	1.48	5.99
33013	2	9	.02	.07	3.8	15.2	26	104	.19	.76	.30	1.20	2.77	11.20
33014	1	3	.02	.09	.8	3.2	7	30	.25	.99	.15	.60	.78	3.16
33015	1	3	.03	.12	1.5	6.0	7	28	.16	.64	.10	.40	.79	3.20
33016	1	3	.02	.08	1.3	5.2	6	26	.40	1.60	.10	.40	.59	2.40
33017	< 1	2	.03	.12	.8	3.2	7	28	.25	1.00	.10	.40	.79	3.19
33018	1	3	.02	.08	.6	2.4	6	23	.14	.56	.10	.40	.49	2.00
33019	< 1	2	.02	.08	.7	2.8	6	26	.13	.51	.10	.40	.69	2.80
33020	1	3	.01	.05	1.3	5.2	8	33	.58	2.31	.10	.40	.89	3.59
33021	1	3	.02	.09	2.5	10.0	7	30	.11	.44	.10	.40	1.28	5.20
33022	1	3	.03	.12	3.0	12.0	5	22	.02	.08	.10	.40	.79	3.20
33023	< 1	2	.02	.09	.9	3.6	8	32	.16	.64	.10	.40	1.09	4.40
33024	< 1	3	.03	.10	2.6	10.4	6	24	.15	.60	.10	.40	1.98	8.00
33025	1	3	.02	.10	.6	2.4	13	54	.09	.35	.15	.60	.89	3.59

* Results are expressed as micrograms per gram.

APPENDIX W

Location of Elk Observed From Aircraft, 1971-1974

Year	Flight Date	Number Observed and Classification ^{1/}	Drainage	Location				Year	Flight Date	Number Observed and Classification ^{1/}	Drainage	Location			
				TN	RW	S	± Sex.					TN	RW	S	± Sex.
1971	1-13	4 Co	Pass Creek	15	6	8		1971	4-13 cont.	6	Bedrock Creek	16	6	5	SW
		30 HCo	Alice Creek	16	7	13				11	Bedrock Creek	17	6	31	NW
		14	Green Creek	16	7	13				8	Bedrock Creek	17	6	32	NW
		20 HCo	Green Creek	16	6	7,18				3	Bedrock Creek	17	6	32	center
										1	Little Skunk Creek	17	6	30	NW
	2-9	136 HCo	Bedrock Creek	16	6	3				11	Little Skunk Creek	17	6	19	NW
			Hardgrove Creek	17	6	34,35				8	Little Skunk Creek	17	6	20	SE
		2 B	Tom's Gulch	16	7	6				36	Green Creek	16	6	16	SW
		1 C	M. Pk Dearborn	16	6	30				14	M. Pk. Dearborn	16	6	15	NW
		2 B, 19 C, 8 c	Telephone Gulch	16	7	24				95	M. Pk. Dearborn	16	6	15	NE
	3-2	1 B	Bedrock Creek	16	6	5			5-12	1	Tom's Gulch	15	7	6	SW
		34 B, 134 C, 27 c	Bedrock Creek	16	6	3				4	Blackfoot	15	7	31	SE
		2 C, 2 c	Tom's Gulch	15	7	6				13	Hardscabble Creek	15	7	8	SE
		24	Telephone Gulch	16	7	23	NE			1	Hardscabble Creek	15	7	17	NE
		9	Green Creek	16	7	13	NW			2	Bartlett Creek	15	7	2	NW
	4-2	117	Bedrock Creek	16	6	3	NW			3	Tom's Gulch	16	7	32	SW
		25	Green Creek	16	6	4	SW			4	Alice Creek	16	7	27	NW
		12 B	Bedrock Creek	16	6	5	SE			1	Alice Creek	16	7	10	SW
		1 B	Landers Fork	15	7	6	SW			2	Alice Creek	16	7	13	NW
			Landers Fork	15	7	6	SE			14	M. Pk. Dearborn	16	6	29	SE
	4-8	33	Alice Creek	16	7	13	NW			2	M. Pk. Dearborn	16	6	15	SW
		8	M. Pk Green Creek	16	6	7	NE			17	M. Pk. Dearborn	16	6	15	SE
		5	Tom's Gulch	15	7	5	NE			9	M. Pk. Dearborn	16	6	10	SW
		12	Green Creek	16	7	13	NE			10	M. Pk. Dearborn	16	6	10	SE
		4	Alice Creek	16	7	14	NE			10	Green Creek	16	6	17	SW
1972	4-13	9	Green Creek	16	6	16	NE			2	Green Creek	16	6	17	NE
		35	Green Creek	16	6	9	NE			1	Green Creek	16	6	8	SE
		6	Bedrock Creek	17	6	31	SE			2	Green Creek	16	6	6	SW
		8	Bedrock Creek	17	6	30	SW			8	Green Creek	16	7	12	SW
		2	Bedrock Creek	17	6	30	SE			2	Green Creek	16	7	1	NW
	5-24	9	Telephone Gulch	16	7	23	NE			17	E. Pk. Falls Creek	17	7	36	NE
		4	Tom's Gulch	15	7	5	SE			3	Big Skunk Creek	17	6	18	NW
		6	M. Pk Green Creek	16	6	12	SE			3	Big Skunk Creek	17	6	18	SE
		13	M. Pk Green Creek	16	6	7	NE			59	Hardscabble Creek	15	7	8, 17, 20	SE
		2	M. Pk Green Creek	16	6	6	SE			1	Landers Fork	15	7	18	SW
										3	Landers Fork	15	7	7	SW
										2	Landers Fork	15	8	12	SE
										16	Landers Fork	15	8	12	SE
										1	Landers Fork	15	8	1	SW
										6	Landers Fork	15	7	31	NW

^{1/} B = Bull C = Cow c = calf @ = newborn calf

no symbol by number = unclassified

APPENDIX W continued

Year	Flight Date	Number Observed and Classification ^{1/}	Drainage	Location			Year	Flight Date	Number Observed and Classification ^{1/}	Drainage	Location			
				TN	NS	± Sec.					TN	NS	± Sec.	
1971	5-24 cont.	2	Tom's Gulch	15	7	6 NW	1972	5-24 cont.	21	Landers Fork	15	7	18 NE	
		3	Tom's Gulch	15	7	6 center			12	Landers Fork	15	7	6 SW	
		8	Hogum Creek	14	7	16 or 22			5	Tom's Gulch	15	7	6 NE	
	5-25	2	Black Diamond Creek	14	7	10 or 11			1	Landers Fork	15	8	1 NE	
		5	Landers Fork	16	9	14 NE			6	Blackfoot	15	7	30 SE	
		1	Hardscrabble Creek	15	7	20 SW			1	Landers Fork	15	7	30 SW	
	6-6	9	Hardscrabble Creek	15	7	30 center			6	Blackfoot	15	7	31 NE	
		2 C, 1 @	1	Alice Creek	15	7			21 NW	1	Blackfoot	15	7	31 SE
		1	Alice Creek	15	7	22 NW			10	Alice Creek	16	7	28 center	
	1	5	Hardscrabble Creek	15	7	17 NW			4	Telephone Gulch	16	7	23 SW	
		4	Hardscrabble Creek	15	7	8 SW			4	Alice Creek	16	7	14 NE	
		1	Hardscrabble Creek	15	7	8 SW			3	Green Creek	16	7	12 SE	
	12, 1 @	1	Hardscrabble Creek	15	7	17 NW			8	N. Pk. Green Creek	16	6	7 NE	
		6	Landers Fork	15	7	18 NE			5	Big Skunk Creek	17	6	7 SW	
		1	Landers Fork	15	7	18 NW		6-3	7	Landers Fork	15	7	30 SE	
	7 C, 3 @	1	Landers Fork	15	7	7 center SE			5	Landers Fork	15	7	30 NE	
		4	Landers Fork	15	7	7 NE			7	Landers Fork	15	7	30 NE	
		1	Tom's Gulch	15	7	6 SW			1	Hardscrabble Creek	15	7	19 NW	
	6-15	1	Landers Fork	15	7	18 NW			4	Landers Fork	15	7	18 NE	
		1	Hardscrabble Creek	15	7	16 SW			2	Hardscrabble Creek	15	7	17 NE	
		1	Landers Fork	15	7	31 NE			3	Landers Fork	15	7	18 SW	
	7-1	4	Hardscrabble Creek	15	7	17 NW			7	Landers Fork	15	7	18 NW	
		2	Landers Fork	15	8	36 NE			2	Landers Fork	15	7	7 SW	
		1	Landers Fork	15	7	7 SW			2	Landers Fork	15	8	1 SE	
1972	4-19	46	Bedrock Creek	16	6	3 NE	1		Hardscrabble Creek	15	7	16 NW		
		10	N. Pk. Dearborn	16	6	2 NW	1		Landers Fork	16	9	11 SE		
		53	Green Creek	16	6	10 NE	8		Alice Creek	16	7	14 NE		
	5-4	54	Green Creek	16	6	11 NW	6-7	7	Hardscrabble Creek	15	7	8 SE		
		37	Hardgrove Creek	17	6	35 SW		1	Tom's Gulch	15	7	5 SW		
		15	Big Skunk Creek	17	6	7 SW		4	Tom's Gulch	15	7	7 NE		
	5-24	2	Tom's Gulch	17	6	17 SW		9	Landers Fork	15	7	7 SE		
		7	Tom's Gulch	15	7	6		4	Landers Fork	15	7	18 NW		
		2	Landers Fork	15	9	14 NE		1	Landers Fork	15	8	1 SE		
	7	3	Hardscrabble Creek	15	7	8 NW		5	Landers Fork	15	8	1 SW		
		2	Hardscrabble Creek	15	7	17 NW		4	Alice Creek	16	7	12 SW		
		23	Hardscrabble Creek	15	7	17 SW		6	Green Creek	16	7	13, 18 center section line		
	1	1	Hardscrabble Creek	15	7	17 SE		6-20		1	Alice Creek	15	7	16 SE
		3	Hardscrabble Creek	15	7	20 NW				1	Hardscrabble Creek	15	7	17 SW
		4	Alice Creek	15	7	16 center				10	Landers Fork	15	7	18 NE
	1	1	Alice Creek	15	7	16 SW				1	Landers Fork	15	7	7 center
		1	Alice Creek	15	7	22 NW								
		1	Hardscrabble Creek	15	7	19 NE								
1	Hardscrabble Creek	15	7	19 SW										

APPENDIX W continued

Year	Flight Date	Number Observed and Classification ¹	Drainage	TN	RM	S	1/4 Sec.
1972	6-20	cont. 1
2 C, 1 G.	Landers Fork	15	7	7	NW
		1	Alice Creek	16	7	15	SW
		1	Alice Creek	16	7	10	SW
1973	3-7	±100	Alice Creek	16	7	15	SE
		6	Alice Creek	16	7	22	NE
		3	Alice Creek	16	7	14	NW
		15	Alice Creek	16	7	22	SW
	3-19	440	Willow Creek	13	9	2	NW
			Alice Creek	16	7	15	SE
			Alice Creek	16	7	22	NE
		3	Alice Creek	16	7	22	SW
		37	Big Skunk Creek	17	6	7	SW
	4-3	58	Blackfoot	14	8	16,17	
		1	Alice Creek	16	7	11	SW
		13	Telephone Gulch	16	7	23	NE
		7	Bedrock Creek	16	6	3	NE
		15	Green Creek	16	6	4	NE
		116	Green Creek	16	6	9,10	
	4-19	20	Alice Creek	16	7	22	center
		7	Green Creek	16	6	4	SW
		136	Bedrock Creek	17	6	33	SW
	5-31	1	Bedrock Creek	16	6	3	NE
		5	Hardscrabble Creek	15	7	30	SE
		2	Hardscrabble Creek	15	7	30	SE
		3	Hardscrabble Creek	15	7	30	SW
		5	Landers Fork	15	7	30	NE
		1	Landers Fork	15	7	30	SE
		1	Alice Creek	15	7	22	NW
		1	Alice Creek	16	7	12	SW
		2	Hardscrabble Creek	15	7	20	SW
		1	Hardscrabble Creek	15	7	20	SE
		4	Landers Fork	15	7	18	SW
		3	Hardscrabble Creek	15	7	17	NE
		1	Landers Fork	15	7	18	NE
		4	Hardscrabble Creek	15	7	8	SW
		1	Landers Fork	15	7	7	NW
		2	Landers Fork	15	7	1	SE

A-148

Year	Flight Date	Number Observed and Classification	Drainage	TN	RM	S	1/4 Sec.
1974 ²	4-30	2	Landers Fork	15	7	7	SW
		8	Hardscrabble Cr.	15	7	17	NW
		1	Hardscrabble Cr.	15	7	20	NE
	5-14	37	Landers Fork	15	8	1	SE
		3	Hardscrabble Cr.	15	7	17	SW
		34	Hardscrabble Cr.	15	7	19	NE
		5	Landers Fork	15	7	29	NW
		11	Blackfoot	15	7	32	NW
		14	Blackfoot	15	7	32	NE
	6- 1	1	Hardscrabble Cr.	15	7	7	NW
		9	Hardscrabble Cr.	15	7	18	NW
		3	Hardscrabble Cr.	15	7	18	NW
		10	Hardscrabble Cr.	15	7	18	SE
		16	Hardscrabble Cr.	15	7	17	SW
		5C, 1G	Hardscrabble Cr.	15	7	17	NE
		5	Hardscrabble Cr.	15	7	16	SW
		3	Hardscrabble Cr.	15	7	28	NW
		6	Hardscrabble Cr.	15	7	29	NW
		1	Hardscrabble Cr.	15	7	29	SW
		1	Hardscrabble Cr.	15	7	31	SW
		1	Hardscrabble Cr.	15	7	31	SW
		11	Landers Fork	15	8	25	SW
		3	Landers Fork	15	8	36	NE

²Data for 1974 is not considered as complete as 1971-1973 data, since observations in 1974 were made from a Cessna 180 instead of a Supercub. However, the observations actually shown in 1974 are accurate.

APPENDIX K

Additional elk observed with radioed elk CH-2, CH-6 and CH-9 on telemetry flights. Flights which were not completed due to bad weather are not shown.

<u>Elk CH-2</u>				<u>Elk CH-2 continued</u>			
<u>Year</u>	<u>Flight Date</u>	<u>Radioed Elk Seen?</u>	<u>Number of Additional Elk Seen with Radioed Elk1/</u>	<u>Year</u>	<u>Flight Date</u>	<u>Radioed Elk Seen?</u>	<u>Number of Additional Elk Seen with Radioed Elk1/</u>
1973	Banded on 4-26			1974	1- 3	No	30-40
	5- 3	Yes	Not Recorded		2- 7	Yes	Not Recorded
	5-11	Yes	13+		3- 9	Yes	20
	5-14	Yes	10		4- 4	No	15-20
	5-17	No			4-17	Yes	51
	5-19	Yes	6-10		4-30	Yes	43
	5-21	No	Several		5- 4	No	6-8
	5-24	No			5- 7	No	
	5-29	Yes	6		5-14	Yes	14
	6- 2	Yes	6		5-28	Signal Not Located	
	6- 6	Yes	None		6- 1	Signal Not Located	
	6- 7	Location Not Attempted			6-12	Location Not Attempted	20 Q5
	6-11	Yes	1		6-27	Yes	4-6+
	6-20	No			7- 6	Yes	
	6-22	Yes	None		7-15	No	
	6-27	No			7-19	Yes	None
	7- 3	No	4 C, 2 Q		7-26	Yes	3 C, 3 Q
	7-10	No	1		8- 5	Yes	15-20 Q5
	7-14	No			8-15	Yes	4 C, 1 Q
	7-19	No			9- 3	No	
	7-25	No			9-14	No	
	8- 1	No			10- 4	No	
	8- 9	No			10-29	No	
	8-20	No	1 B, 2 C		11-14	No	
	8-27	No	8 Q5, 1B	1975	12-12	Yes	20
	9- 4	No	4		1-15	Yes	85
	9- 7	Yes	8-10		1-31	Yes	14
	9-28	Yes	1 C, 1 Q		2-26	Yes	63
	10- 8	No	4		3-28	Yes	
	10-17	No					
	11-11	No					
A-189	11-14	No					
	12- 5	No					

1/Includes animals in immediate proximity even though radioed elk may not have been seen.

APPENDIX I continued

Elk CH-6

Year	Flight Date	Radioed Elk Seen?	Number of Additional Elk Seen with Radioed Elk
1973	Banded on 4-23		
	5- 3	Yes	Not Recorded
	5-11	Yes	4-5
	5-14	Yes	1
	5-17	Yes	2
	5-19	No	1
	5-21	No	3
	5-24	Yes	2
	5-29	No	2
	6- 2	Yes	8
	6- 6	Yes	None
	6- 7	Location Not Attempted	
	6-11	Yes	8
	6-20	No	5 C, 5 C
	6-22	Yes	18 C, 8 C
	6-27	Yes	6 C, 7 C
	7- 3	Yes	7 C, 2 C
	7-10	Yes	6 C, 4 C
	7-14	Yes	6 C, 3 C
	7-19	Yes	None
	7-25	Yes	9-10
	8- 1	Yes	8 C, 6 C
	8- 9	No	
	8-20	No	
	8-27	Yes	1 C, 1 B
	9- 4	No	
	9- 7	Yes	5 C
	9-28	No	2 B, 1 C
	10- 8	Yes	1 B, 3-4 C
	10-17	Yes	1 C
	11-11	No	
	11-14	No	
	12- 5	Yes	8-10
1974	1- 3	No	30-40
	2- 7	Yes	Not Recorded
	3- 9	Yes	20
	4- 4	No	6
	4-17	Yes	34
	4-30	Yes	30

Elk CH-6 continued

Year	Flight Date	Radioed Elk Seen?	Number of Additional Elk Seen with Radioed Elk
1974	5- 4	No	
	5- 7	No	2
	5-14	Yes	10
	5-28	No	
	6- 1	Yes	1
	6-12	Yes	4
	6-27	No	Several-Numbers Not Recorded
	7- 6	Yes	1
	7-15	Yes	3 C, 1 C
	7-19	Yes	1 C
	7-26	Yes	None
	8- 5	No	
	8-15	Yes	None
	9- 3	No	
	9-14	No	1
	10- 4	Too Windy	
	10-29	Signal not located	
	11-14	Signal not located	
	12-12	Yes	2 C, 1 C
	1-15	Yes	3 C, 2 B
1975	1-31	Signal not located	
	2-26	Yes	53
	3-28	Yes	52

Elk CH-9

Year	Flight Date	Radioed Elk Seen?	Number of Additional Elk Seen with Radioed Elk
1973	Banded on 4-23		
	5- 3	No	
	5-11	Yes	1
	5-14	Yes	2
	5-17	Yes	4
	5-19	Yes	2
	5-21	No	1
	5-24	No	1
	5-29	Yes	1
	6- 2	Yes	3
	6- 6	No	
	6- 7	Lost Radio Signal - Animal Killed by Hunter, October 17 or 18, 1973.	

Appendix Y (continued)

Movements of radioed elk Q-6 from April 23, 1973 to January 3, 1974.

Appendix 1 (continued)

Movements of redcedar elk G-6 from January 8, 1974 to March 20, 1975.