REPTILE LIFE

WALFORD B. JOHNSON
& STANLEY C. JOHNSON B.A.

d
G

L

1.

d

G.

Presented by the Executors of
Norman Douglas Simpson

1890 - 1974

Gowans's Nature Books

THE object of these little books is to stimulate a love for nature and a desire to study it. The first volumes of the series have been extremely successful, and the publishers hope to make the others equally interesting and beautiful.

- No. 1.—WILD BIRDS AT HOME. Sixty Photographs from life, by Chas. Kirk, of British Birds and their Nests.
- No. 2.—WILD FLOWERS AT HOME. First Series. Sixty Photographs from Nature, by Cameron Todd.
- No. 3.—WILD FLOWERS AT HOME. Second Series. By Cameron Todd.
- No. 4.—BUTTERFLIES AND MOTHS AT HOME. Sixty Photographs from life, by A. Forrester.
- No. 5.—WILD BIRDS AT HOME. Second Series, by Charles Kirk.
- No. 6.—FRESHWATER FISHES. Sixty Photographs from life, by Walford B. Johnson and Stanley C. Johnson, B.A.,
- No. 7.—TOADSTOOLS AT HOME. Sixty Photographs of Fungi, by Somerville Hastings, F.R.C.S.
- No. 8.—OUR TREES & HOW TO KNOW THEM. Sixty Photographs by Charles Kirk.
- No. 9.—WILD FLOWERS AT HOME. Third Series. By Somerville Hastings, F.R.C.S.
- No. 10.—LIFE IN THE ANTARCTIC. Sixty Photographs by Members of the Scottish Antarctic Expedition.
- No. 11.—REPTILE LIFE. Sixty Photographs by Walford B. Johnson and Stanley C. Johnson, B.A.
- No. 12.—SEA-SHORE LIFE. By the same.

Others in Preparation

PRICE:

PARCHMENT COVER, 6D. NET. EACH VOLUME.

POSTAGE, 1D. EACH.

GOWANS & GRAY, Ltd., London & Glasgow

The Children's Bookshelf

PRETTY little Books with about a dozen
coloured illustrations by well-known artists.

No. 1.—THE UGLY DUCKLING, by Hans Andersen.
Coloured illustrations by Stewart Orr.

No. 2.—LITTLE KLAUS AND BIG KLAUS, by Hans
Andersen. Coloured illustrations by Charles Pears.

No. 3.—THE PIED PIPER OF HAMELIN, by
Robert Browning. Coloured illustrations by Charles
Robinson.

No. 4.—THE STORY OF CHANTICLEER. Re-told
from Chaucer, by R. Brimley Johnson. Coloured
illustrations by Stewart Orr.

PRICE, 6D. EACH IN PARCHMENT COVERS. POST FREE 7D.

GOWANS & GRAY, Ltd., London & Glasgow

Natural History Museum Library

000097233

Pocket Anthologies

Compiled by ADAM L. GOWANS, M.A.

33rd Thousand.

No. 1.—The Hundred Best Poems (Lyrical) in the English Language.

15th Thousand.

No. 2.—The Hundred Best Poems (Lyrical), in the English Language. Second Series.

10th Thousand.

No. 3.—The Book of Love: One hundred o. the best love-poems in the English language.

6th Thousand.

No. 4.—The Hundred Best Blank Verse Passages in the English Language.

In Preparation.

No. 5.—Poetry for Children. One hundred of the best poems for the young in the English language.

No. 6.—The Ways of God. An unconventional anthology of poems on the great problems of existence.

Others in Contemplation.

PRICES:

PAPER, 6D. NET.

CLOTH, 1S. NET.

LEATHER, 2S. NET.

POSTAGE, 1D.

GOWANS & GRAY, Ltd., London & Glasgow

The Haunted and the Haunters;

OR,

THE HOUSE AND THE BRAIN

BY

LORD LYTTON

This masterpiece in little has been called, and probably is, the best ghost story in the world. It would certainly be difficult to find any to surpass it in its bold and effective introduction of the supernatural, its equally daring, yet consistent and logical, explanation of uncanny appearances.

The present is an exact reprint of the tale as it originally appeared in *Blackwood's Magazine* for 1859. The incomplete version which is printed in Lytton's *Collected Works* leaves many of the incidents very imperfectly explained.

Price, in Pretty Parchment Covers, 6d. Net. Post Free, 7d.

THE
HEAVENLY FOOTMAN;

OR,

A DESCRIPTION OF THE MAN

THAT GETS TO HEAVEN

BY

JOHN BUNYAN

“The Heavenly Footman” is the best known of Bunyan’s shorter works. Vigorous, racy, and directly personal in its appeal, it is a sermon which, delivered orally, could never fail to be deeply impressive. And it will command the attention of him who merely reads it better than most sermons he is likely to hear.

The text followed is that printed by Mr. George Offor in the Works of Bunyan, published by Messrs. Blackie & Son in 1860.

THE MARRIAGE RING;

OR,

THE MYSTERIOUSNESS AND

DUTIES OF MARRIAGE

BY

JEREMY TAYLOR

The Treatise on "The Marriage Ring" occupies Nos. XVII. and XVIII. of the famous "Twenty-five Sermons preached at the Golden Grove; being for the Winter Half-year." These two Sermons have always been the most popular of the great prose-writer's. The text followed is the best, *i.e.*, that of Eden's revised and corrected issue (1862) of Heber's edition of the Whole Works, Taylor's own notes only being given.

Price, in Pretty Parchment Covers, 8d. Net. Post Free, 7d.

THE
LADY OF LYONS;

OR,
LOVE AND PRIDE

A PLAY
BY
LORD LYTTON

"The Lady of Lyons" is a thoroughly enjoyable, if unsophisticated, play, and has held the stage since it was written. Though not, as a whole, of great literary merit, it rises in a few blank verse passages to a high poetic level.

The present is an exact reprint of the first edition, a few obvious misprints having been corrected.

THE TOWER OF NESLE

A PLAY

BY

ALEX. DUMAS, PÈRE

One of the most striking examples of the Romantic drama. A most exciting play in which every speech, every word tells with tremendous force. Thrilling to read, one can imagine the sensation it produced when first acted, at a time when the antagonism between the supporters of the Classical and Romantic drama was at its height.

Price, In Pretty Parchment Covers, 6d. Net. Post Free, 7d.

GOWANS'S NATURE BOOKS, No. 11

REPTILE LIFE

CARSON & NICOL, PRINTERS, GLASGOW.

BLOCKS BY HISLOP & DAY, EDINBURGH.

SHARP-NOSED CROCODILE
(*Crocodylus americanus*)

W. B. Johnson, Photo.

REPTILE LIFE

*Sixty photographs from life, by Walford B. Johnson and
Stanley C. Johnson, B.A.*

GOWANS & GRAY, LTD.

5 ROBERT STREET, ADELPHI, LONDON, W.C.

58 CADOGAN STREET, GLASGOW

1907

AGENTS FOR THE SALE OF THIS SERIES

France: A. Perche, 45 Rue Jacob, Paris.

Germany: W. Weicher, Windmühlenweg, I. III, Leipzig.

Switzerland: E. Frankfurter, 12 Grand-Chêne, Lausanne.

Belgium: Spineux & Cie, 62 Montagne de la Cour, Brussels.

Scandinavia: A. & O. Schedin, Lund, Sweden.

Spain and Portugal: Ferreira & Oliveira, Lda, Rua do Ouro 132, Lisbon.

Holland: Kirberger & Kesper and Jacs. G. Kobbers, Amsterdam.

„ *A. Abrahams, The Hague.*

ALL the photographs in the present volume, like those in "Freshwater Fishes," by the same authors, are from life. Particular attention is drawn to the fact that Messrs. Johnson have been successful in obtaining photographs of examples of ALL the British reptiles.

SHARP-NOSED CROCODILE

W. B. Johnson, Photo.

W. B. Johnson, Photo.

INDIAN CROCODILE
(*Crocodylus palustris*)

CROCODILE (SWIMMING ON THE SURFACE)

W. B. Johnson, Photo.

BRAZILIAN TORTOISE
(*Testudo tabulata*)

W. B. Johnson, Photo.

GRECIAN TORTOISE
(*Testudo graeca*)

W. B. Johnson, Photo.

DENTATED HINGED TORTOISE
(*Cinixys erosa*)

W. B. & S. C. Johnson, Photo.

CAROLINA BOX-TURTLE
(*Cistudo carolina*)

W. B. Johnson, Photo.

EUROPEAN POND-TORTOISE
(*Emys orbicularis*)

W. B. Johnson, Photo.

DERBIAN STEMOTHERE
(*Sternothoerus derbianus*)

W. B. & S. C. Johnson, Photo.

SNAPPING TURTLE
(*Chelydra serpentina*)

W. B. & S. C. Johnson, Photo.

BUNGOMA RIVER TURTLE
(*Emyda granosa*)

W. B. & S. C. Johnson, Photo.

BUNGOMA RIVER TURTLE (SWIMMING)

W. B. Johnson, Photo.

WALL-GECKO
(*Tarentola mauritanica*)

W. B. & S. C. Johnson, Photo.

WALL-GECKO ON TRUNK OF A TREE

W. B. & S. C. Johnson, Photo.

RED-THROATED ANOLIS
(*Anolis carolinensis*)

W. B. Johnson, Photo.

CHAMAELÉON ANOLIS
(*Anolis equestris*)

W. B. Johnson, Photo.

W. B. Johnson, Photo.

HORNED LIZARD
(*Phrynosoma cornutum*)

W. B. Johnson, Photo.

HORNED LIZARD

COMMON SCHELTOPUSIK
(*Ophisaurus apus*)

W. B. Johnson, Photo.

COMMON SCIAENOPS

W. B. & S. C. Johnson, Photo.

BLINDWORM (BRITISH)
(*Amphisbaena*)

W. B. Johnson, Photo.

W. B. & S. C. Johnson, Photo.

PEARL LIZARD
(*Lacerta ocellata*)

SPANISH EYED-LIZARD
(*Lacerta ocellata*)

W. B. Johnson, Photo.

SPANISH EYED-LIZARD

W. B. & S. C. Johnson, Photo.

GREEN LIZARD
(*Lacerta viridis*)

W. B. & S. C. Johnson, Photo.

GREEN LIZARD

W. B. & S. C. Johnson, Photo.

GREEN LIZARD

W. B. & S. C. Johnson, Photo.

GREEN LIZARD

W. B. & S. C. Johnson, Photo.

GREEN LIZARD

W. B. & S. C. Johnson, Photo.

SAND LIZARD
(*Lacerta agilis*)

W. B. Johnson, Photo.

CONTINENTAL SAND LIZARD
(*Lacerta agilis*)

W. B. & S. C. Johnson, Photo.

VIVIPAROUS LIZARD (BRITISH)
(*Lacerta vivipara*)

W. B. & S. C. Johnson, Photo.

WALL LIZARD
(*Lacerta muralis*)

W. B. & S. C. Johnson, Photo.

WALL LIZARD

W. B. & S. C. Johnson, Photo.

WALL LIZARD

W. B. & S. C. Johnson, Photo.

WALL LIZARD

W. B. & S. C. Johnson, Photo.

ACANTHUS-FOOYED LIZARD
(*Acanthodactylus velox*)

W. B. Johnson, Photo.

ACANTHUS-FOOTED LIZARD

W. B. Johnson, Photo.

BLUE LIZARD
(*Gerrhonotus cernuus*)

W. B. & S. C. Johnson, Photo.

COMMON SKINK
(*Scincus officinalis*)

W. B. Johnson, Photo.

CUNNINGHAM SKINK
(*Eyenia cunninghami*)

W. B. Johnson, Photo.

CUNNINGHAM SKINK

W. B. & S. C. Johnson, Photo.

W. B. Johnson, Photo.

COMMON CHAMELEON
(*Chamaeleon vulgaris*)

COMMON CHAMELEON

W. B. Johnson, Photo.

COMMON BOA
(*Boa constrictor*)

W. B. Johnson, Photo.

RINGED SNAKE (BRITISH)
(*Tropidonotus natrix*)

W. B. Johnson, Photo.

DICE SNAKE
(*Tropidonotus tessellatus*)

W. B. Johnson, Photo.

DARK GREEN SNAKE
(*Zamenis gemonensis*)

W. B. Johnson, Photo.

ÆSCULAPIAN SNAKE
(*Coluber longissimus*)

W. B. Johnson, Photo.

ÆSCULAPIAN SNAKE

W. B. Johnson, Photo.

W. B. Johnson, Photo.

ÆSCULAPIAN SNAKE

LEOPARD SNAKE
(*Coluber leopardinus*)

W. B. Johnson, Photo.

LEOPARD SNAKE

W. B. Johnson, Photo.

BULL OR PINE SNAKE
(*Coluber melanoleucus*)

W. B. Johnson, Photo.

SMOOTH SNAKE (BRITISH)
(*Coronella austriaca*)

W. B. Johnson, Photo.

SMOOTH SNAKE (BRITISH)

W. B. Johnson, Photo.

KING SNAKE
(*Coronella getula*)

W. B. Johnson, Photo.

HOOP SNAKE

(Erythrolamprus asculapii)

W. B. Johnson, Photo.

COMMON VIPER OR ADDER (BRITISH)
(*Vipera berus*)

W. B. Johnson, Photo.

Some Notes on the Reptiles

OF WHICH

PHOTOGRAPHS APPEAR IN

THE FOREGOING PAGES . .

BY

WALFORD B. JOHNSON.

To the average person the word "Reptile" means anything between snakes and slugs or crocodiles and worms, and still to a large number induces thoughts of something bideous and repulsive, or even dangerous. Properly, however, this term should only be applied to cold-blooded vertebrate creatures, which are born into the world with all the main features of their parents' form and shape already developed. The Common Frog, which is regarded by many as one of the most typical of this order, is really not a reptile at all. It first sees the light as a tadpole, and is then as much unlike its parents as it is possible to imagine. On account of these transformations which it undergoes in the early part of its existence, the frog, in common with the toad, newt, salamander, and many others, is distinguished from true reptiles. Now a lizard is born usually from an egg, and escapes from the shell as an exact representation, in miniature, of its parents in everything but colour. This, coupled with the fact that it is cold-blooded and vertebrate, at once brings the lizard under the order of Reptilia. In short an animal is only a reptile when it possesses a backbone, is cold-blooded, and is born structurally similar to its parents, providing also it breathes at all periods of its existence by means of lungs.

Every member of this order is covered with some kind of scaly armour, the plates varying between the tiny granules of the chamæleon and the large shields which protect the tortoise. We may divide all reptiles roughly into four great families:—Crocodiles, Tortoises, Lizards, and Snakes.

Crocodiles, as every one knows, when of any size are extremely dangerous, becoming morose and wantonly destructive to animal life; but if taken young they may be easily tamed, becoming then quite gentle and good tempered. They inhabit all the tropical parts of the Old World, spending an almost aquatic life, and preying on any terrestrial creatures, which they are strong enough to pull under water and drown. Their throats for the size of the animals are surprisingly small, which necessitates the tearing of their food into small pieces. This is accomplished very quaintly by firmly holding the victim in the jaws, while pieces are torn from it by the claws of the hind foot. Crocodiles are chiefly confined to the Old World, but here is a species found in Jamaica, this being the one figured in our

illustrations. Alligators, Caimans, etc., are almost entirely confined to tropical America. All the members of the Crocodilia are born from eggs, which are usually deposited in masses of decaying vegetation on the margins of the rivers and lakes they inhabit. The heat produced in the fermentation helps to bring about the incubation of the eggs. Alligators are in most respects similar to crocodiles, but the typical member of the latter group is a much more scaly and rougher skinned animal than the average alligator. Some crocodiles are so much like their congeners and *vice-versa* that it is a very difficult matter to discriminate between them. If, however, the following formula be remembered no further difficulty will be experienced in their separation. Close to the front of the lower jaw of a crocodile there is a tusk on either side which, instead of being covered by the upper jaw, still shows when the mouth is shut. In the upper jaw there is a depression or notch, and it is into this that the tusk fits. Alligators invariably lack this dental arrangement.

Tortoises.—The regularity with which these creatures make their appearance in bird-shops during the early summer has made them neither feared nor respected. These Grecian Land Tortoises are usually purchased to clear the house of "black-beetles," or to rid the garden of slugs. Curiously enough though this is a popular notion, no tortoise of this species has ever been known to prey on either of the above pests, not even when dying from starvation, which they are too frequently left to do. Instead of clearing the garden of slugs they are usually discovered clearing it of lettuce, strawberries, etc., as they are confirmed vegetarians. These reptiles have no teeth but a very bird-like beak, the jaws closing on one another like a pair of scissors. As they are toothless, they do not masticate their food but bite off portions of a convenient size to swallow. The carnivorous tortoises tear their victims to pieces with the help of their fore-legs. These creatures are born from almost spherical hard-shelled eggs which are laid in trenches in the sand. The holes are dug out by the hind feet of the tortoise which, after depositing the eggs, refills the trench with the dug-out sand and leaves their incubation to the sun. With the approach of the cold season these reptiles cease to feed, and, burrowing down deeply into the ground, they pass the winter in a state of torpor, again waking to activity with the return of the warmth of spring.

Lizards.—This family is by far the most widespread of all the reptile groups. Its members are found in almost every country in the world which is not embraced by the Polar Regions. They inhabit alike dry arid deserts and tropical jungle marshes, and several races exclusively frequent the higher elevations in mountainous districts.

It is this section of the Reptilia which displays the greatest diversity in colouration. Most lizards are hatched in the ordinary way, but several species are born alive. Our common English Lizard is an example of this latter group. Though these reptiles are so abundant, only one division comprising but a few species contains members capable of giving a poisonous injection. These are the American Heloderms which are fortunately never met with in the Old World. The outer skin of these reptiles is periodically cast, not, as in the case of snakes in one piece, but is shredded off as the animals dart in and out the coarse herbage, amongst which they live. The illustration on page 32 shows a green lizard beginning to slough on the right leg. There is, however, one lizard which sheds its coat entire, and that one is the Blue Lizard of America figured on page 44.

Snakes.—Of all reptiles, and indeed animals generally, the snake is the least understood by the average person, and is consequently the most distrusted. Many people still regard all these creatures as vindictive and poisonous, and perversely refuse to be persuaded otherwise. The ancients were quite at a loss to account for the method of a snake's progression, for it seemed to them to travel by means of some invisible power, for they saw no limbs. Because it was too much for them, therefore, witchcraft was considered to be at the bottom of it. This coupled with the fact that some snakes were known to inflict poisonous wounds gave rise to the silly prejudice which still finds many disciples to this day. A number of snakes are deadly poisonous, but the vast majority are known to be quite harmless. It is a common, though erroneous, belief that every snake is anxious at all times to attack human beings. The most dangerous of these would be far more anxious to escape than to bite. If, however, retreat is cut off, the cornered reptile would fiercely strike in its own defence.

The scaly covering of most snakes is so beautifully glossy that many people are given the impression that they are wet and slimy. All ophidians cast their skins several times during the year, and the process is a very curious one. The colours of the body gradually begin to darken, and the creature is then comparatively lethargic. After about a week of this the skin over the eyes becomes quite opaque and bluish, and the ventral scales assume the same shade. Several days are passed in this state, the snake being all the time quite out of condition and restless, refusing to take food. At the end of this period the eyes once more glisten with their natural colours, and the whole of the animal appears to have ridded itself of the old cuticle. However this is not so. The reptile now begins to rub its lips on all the rough objects in its path until at last the skin of the jaws is rasped back. When once started, the loose covering is further drawn off by the snake creeping under rough sticks and bark. These catch in the old covering and hold it while the animal draws itself out. In this way a healthy snake will make a perfect cast, although the slough will be turned inside out.

A serpent's method of taking food is also very interesting. If a live mouse be placed in a cage containing a hungry leopard snake, the former will be immediately seized by the head in such a way that the victim cannot bite its assailant. The coils of the ophidian quickly twist round the unfortunate mouse and in a few seconds crush all signs of life from it. When satisfied that it is quite dead, the reptile proceeds to work its mouth over the head of the carcase by a curious alternate action of the sides of the jaws. Now, instead of being jointed together by a bony process, as in other animals, the jaws of a snake are united by an elastic ligament. This allows the creature to swallow prey many times larger than its own head. As the jaws work over the mouse, the ligament stretches sufficiently to let the victim (still quite whole) pass down into the stomach of the reptile. A swelling afterwards usually indicates the exact position in the snake of the swallowed animal.

Note.—The figures at the end of each description indicate the average lengths usually attained by the respective animals.

The Sharp-nosed Crocodile comes from Jamaica and the West Indies generally. It is one of the few crocodiles inhabiting the New World. This species is characterised by the narrow and elongated snout, which has a distinct median ridge running throughout its length (10 to 15 feet).

The Indian Crocodile is the typical member of this family, and at the same time one of the most fierce. It would be quite possible for a man to go amongst a number of full-grown alligators and receive no hurt; but to go unarmed too near a single specimen of a crocodile would result in disaster, especially if it were of the above species. This monster inhabits the whole of India and the adjacent islands, living exclusively in fresh water. Its food consists of any animal it is strong enough to master, terrestrial victims being dragged under water and drowned before they are eaten. When the dry season comes round these crocodiles *æstivate*, burying themselves in the mud during the drought, where in a state of torpor they await the refilling of the water-holes (12 to 16 feet).

Brazilian Tortoise.—This species is one of the most handsome of all the tortoise tribe. The shields of the carapace are of a golden yellow in the centre, bordered by a broad margin of a brown or blackish tint. The head and legs are also of a dark colour, but relieved by a number of bright red spots. As its name implies it is a native of Brazil and the warmer parts of South America. There it dwells in the forest jungles, living on herbs and fruits ($1\frac{1}{2}$ to 2 feet).

Grecian Tortoise.—In the late spring and early summer months it is a common sight in London, and other large towns, to see a number of these creatures displayed in bird-shops and on barrows in market streets. There they spend their time in slowly and laboriously crawling over one another, or surveying the situation with calm indifference. Under such conditions they look anything but attractive. If, however, a specimen be required it may be made to become a really interesting and active creature. In choosing a tortoise of any kind take it up and carefully examine its eyes, refuse it if they are not glossy black and wide open. Then look to its neck and legs, and be satisfied that they are in no way lacerated. The shell, both carapace and plastron, should also receive attention, as specimens are not infrequently badly cracked in transit. When satisfied as to its external condition its head or one of its legs, which are sure to be hanging out, should be touched. If that member be sharply withdrawn it is a sign of good vitality, if the reptile remains indifferent to such liberties it is usually in weak health. After a satisfactory purchase has been made it should be taken home and bathed in tepid water, when if a healthy one it will become far more lively. After scrubbing its shell it should be dried on a cloth, and put in such a position that it can freely drink. Sometimes Grecian tortoises will drink for an hour or more. When it has been satisfied on this point a little fresh lettuce or dandelion leaf should be offered it. It may refuse to eat at first, but if the weather be warm healthy herbivorous tortoises seldom refuse lettuce. Under such conditions land tortoises will soon become anything but inactive and uninteresting (12 to 15 inches).

Dentated Hinged Tortoise.—This reptile is a native of New Guinea where it serves as a "table delicacy" to the natives. A very interesting point concerning this tortoise is the natural hinge with which the hinder part of its carapace is furnished. Our photograph clearly shows this flexible ligament. It allows the creature to securely close at will the cavity through which the hind legs usually protrude. The animal is also quite safe from a frontal attack when the head is withdrawn and covered by the hard scaly fore legs. This species, like the Brazilian Tortoise, will do well in English gardens during the summer months if regularly fed with such food as lettuce, gooseberries, and cherries (14 to 16 inches).

Carolina Box-tortoise.—This one is also a hinged tortoise, but here it is the plastron, not the carapace, that is flexible. When the animal is alarmed or sleeping, the head and limbs are withdrawn and the movable portions of the shell, both before and behind, are tightly held against the carapace. In this position only the shell is visible, the softer parts of the tortoise being completely hidden and protected. It is this ability to close that has earned for these creatures their popular title of Box-tortoises. The Carolina species is chiefly nocturnal and inhabits the United States and Mexico. There it lives, for the most part in marshy districts where it can find abundance of water insects, worms, and such creatures which form its staple diet (10 to 12 inches).

European Pond Tortoise.—This is the reptile commonly sold by dealers as the Water Tortoise or the Freshwater Turtle. Its colour is a bluish black above, profusely spotted or splashed with yellow. The under shell is, as a rule, a pale chrome with darker edgings to the shields. It can be bought very cheaply in London and elsewhere during the summer months, and will do very well if supplied with plenty of water which it can enter or leave at will. It feeds beneath the surface on all kinds of fish, water insects, and worms, and in captivity will readily partake of small pieces of raw meat (10 to 12 inches).

Derbian Sternothera.—The Sternotheres are terrapins very similar to the last species in appearance, feeding and living in the same way. They are, however, usually stouter built and have shorter tails. The Derbian variety is almost black with light spots on the head and fore legs. As it spends by far the greater part of its time in water, its shell is often a veritable forest of vegetation on account of the green plants, or algæ as they are called, which grow in profusion on the carapace (1½ to 2 feet).

The Snapping Turtle is a native of North America, where it grows to quite a large size for a terrapin. Someone has fancied a resemblance between it and an alligator, for it is often called an Alligator-terrapin. It is almost exclusively aquatic and carnivorous (4 to 4½ feet).

The Bungoma River Turtle is one of the soft tortoises. They have earned this name "soft" on account of the leathery and spongy shell which contains them. The one figured in our illustrations, when closed, looked like the top of a huge toadstool in colour and shape, for it is almost circular, and a dark brown above and pinkish below. Like most of the aquatic tortoises, it is entirely carnivorous, living on the soft-bodied creatures which share its haunts (10 to 12 inches).

Wall Gecko.—This little lizard for many centuries has been credited with almost every kind of evil. People did not understand it, and consequently, as is usual in such cases, they invented anecdotes to make up for their lack of knowledge. If anything went wrong in the countries where they lived it was put down to the geckos. The disease of leprosy was gravely attributed to a gecko having crawled over the affected person. Needless to say these little creatures are perfectly harmless in every way. Their habits, however, are indeed very curious. The one figured was able to run along a ceiling, in an inverted position of course, as fast as a mouse could travel along the ground. It could also run up and down a window-pane like a fly. Some of these lizards are able to give vent to a kind of chirp-like noise, which is said to sound like "Gec—ko," hence the

popular name of the family. The colour of the wall gecko is very variable, being sometimes straw-coloured with darker blotches, and often dark brown with blackish marks. When alarmed these creatures are able to break off their tails, but whether they are shot off intentionally, or merely knocked off against some object in their hurried flight, is not properly known. The tail, however, grows again in about a season, but this one is smooth, not rough and belted like the old one. If our illustrations be consulted it will be observed that from the tip of the tail to almost the root it is smooth, but just above this part it is clearly marked in sharp pointed belted sections. The smooth tail is the reproduced one. The original tail was to the tip like the portion seen just above the smooth part. Wall geckos live very well in captivity, where they may be fed on almost any kind of insect life. They bask most of the day time, sleeping in the hottest sunshine, but towards evening come out to feed (6 to 8 inches).

Red-throated Anolis.—This is one of the commonest lizards of the West Indies. Its chief interest lies in the fact that it has an even greater power of changing colour than the *Chamæleon*. It is, according to some writers, common everywhere it exists at all. This creature lives among the branches of trees, where it is tolerably safe from its enemies, on account of its colour harmonising so well with the foliage. It may be kept in captivity with other hardy lizards, and treated in the same way (10 to 12 inches).

The Chamæleon Anolis comes from Cuba, and is very similar to the preceding species, but larger. Its colour is anything between black and pea-green. The lower jaw is furnished with an extendable pouch, which the animal widely distends when alarmed. With pouch displayed this creature looks truly dangerous, and it is thought that this dodge is supposed to strike terror into the heart of its would be tormentors, and cause their hurried departure. In captivity "black beetles" seem to be its favourite "dish" ($1\frac{1}{2}$ to 2 feet).

The Horned Lizard, sometimes called the Californian Toad on account of its toad like appearance, is an inhabitant of the dry sandy portions of Mexico and Central America. There it spends its time lying on bare patches basking in the scorching sun. Its food, which is usually taken at dusk, consists of slow moving sand beetles. The creature is of a dull brown colour, blotched with a darker tint of the same hue, and is therefore not easily seen as it lies motionless on stony ground. If however it be disturbed, from an orifice below the eye it squirts a jet of its own blood at the intruder. This method of self-protection is very curious, and possibly the enemy, on seeing blood flow on its own body imagines it has been wounded by the formidable spines, and immediately considers discretion the better part of valour (4 to 6 inches).

The Common Scheltopusik, though very snake-like, is unmistakably a lizard. Scheltopusiks, Blindworms, and a few others are so much like serpents that a few notes on the separation of legless lizards from snakes would possibly be interesting. All but a very few snakes have immovable eyelids. Those of the limbless lizards are perfectly movable. The tongue of a snake is quite different from that of a lizard. In the former it is like two equal pieces of string sewn together for half their lengths with the free portions separated into a fork. In the latter it is like a tape with a deep V-shaped piece cut out of one end. Besides the snake is furnished with a notch in the lower jaw which enables it to put out its tongue without separating the jaws. A lizard has to open its mouth to protrude

that member. Again, all snakes have, on the under surface, a row of very broad scales. These shields extend from side to side of the snake, and there is never more than one row. Here and there it may be that a scale is divided into two, but there are only a very few so divided. Now, if a blindworm be examined in the same way, it will be noticed that its under surface is protected by some eight or ten rows of scales, moreover they will feel quite smooth if stroked the wrong way. If, however, these plates be rubbed from tail to head in a snake, the finger will keep on catching on their loose rough edges. Then again legless lizards move quite clumsily compared with snakes.

The Scheltopusik is quite harmless and will live well in captivity. It is a native of the woods which cover the western part of the Continent, where it preys upon snails, mice, and worms. Its colour is a shining reddish brown, with a pale yellowish head. Like all the lizards, its tail becomes very brittle when the animal is alarmed. On account of its possessing these two properties of glass—brittleness and lustre—it has been called popularly the Glass Snake (30 to 40 inches).

The Blindworm, also called the slow-worm, is not blind or slow, neither is it a worm. Rustics have called it a worm because of its short cylindrical body; blind and slow on account of its remaining quite motionless when disturbed. They considered that it must be either unable to see at all or be very lethargic to remain quiet when in danger, for most creatures rush off under such circumstances. The little animal tries no doubt to pass off as a dead stick. Whenever found it is ruthlessly destroyed, for most countrymen believe this harmless reptile to be even more venomous than the adder itself. Almost all its time, especially towards evening, it is looking out for and devouring slugs, and is therefore one of the animals to be fostered on crop lands, yet most farmers kill as many as they can find. In captivity the blindworm will live very well soon becoming quite tame. It should be kept in a case well supplied with damp earth and moss into which the little creature will burrow. Small slugs and worms should constitute its food. It must have access to warm (not baking) sunshine, and when basking a careful examination will show it has eyes that are anything but blind. These reptiles may be caught in plenty in almost any rural district in south of England, as well as all over the Continent. They usually lie under old stones, boards, or fallen trunks of trees (12 to 14 inches).

The Pearl Lizard and Spanish Eyed Lizard.—These two reptiles are really the same species, but in different stages of development. The Eyed variety is covered with a number of blue spots edged with brown on a dull green background. As the lizard grows older and bigger the spots on the back gradually die away, while those on the sides become larger. Their blue colouration becomes lighter and lack the brownish margins. Later the spots die away altogether and the lizard, which by that time has reached about two feet, becomes a uniform dull green, with occasional splashes of a faint yellow. In these last two stages it is often known as the Pearl Lizard (1 to 2 feet).

The Green Lizard is frequently reputed to be British, but no satisfactory proof of this has been so far forthcoming. However, it is very common all over the warmer parts of the Continent. It forms several well defined varieties, photographs of which will be seen in the foregoing pages. This lizard is of a delicate emerald tint, sometimes devoid of markings, sometimes closely studded with black

spots. Often there is a double row of dark brown blotches edged with white running down the back. Frequently these markings merge into one another forming a continuous line. In fact it would be a difficult task to find two specimens with the same pattern. As this reptile inhabits grassy slopes it will be seen how admirably its colouration is adapted to its protection. Like all the members of its class this lizard is particularly fleet. The speed, however, cannot be maintained for long, but as the animal is seldom far from its hole it can cover just that distance with wonderful agility. When caught it bites furiously, and very often breaks its tail as well, but the nip though painful for the moment does not even cause the teeth to break the skin ($1\frac{1}{2}$ to 2 feet).

The Sand Lizard is a truly British reptile. Its appearance is, however, uncertain, and its distribution is very local. On the Continent it is abundant, forming several varieties there. This lizard is much blunter headed than its green relative just described, the tail too is proportionally shorter. The darker markings on the back are chocolate-brown relieved with white spots, while the under surface is greenish closely spotted with black. This will at once separate emerald varieties of the sand lizard from the green lizard, in which the ventral surface is a uniform yellowish colour (8 to 10 inches).

The Viviparous Lizard is the common brown one, frequently met with on sunny days all over the British Isles, not excepting even Ireland. It is fairly common on dry heaths where there is plenty of heather cover. The black spotted brown colour is very difficult to see against the bare ground where it delights to bask (6 to 8 inches).

Wall Lizards are very abundant all over the Continent. They may be seen scampering up and down the loose stone walls of gardens in the hottest sunshine, and hiding in nooks and crannies such rockeries afford. They seem to form endless varieties, the colours ranging from light brown with yellow spots to dark green relieved with patches of black (7 to 9 inches).

The last six lizards described are all very suitable for the vivarium and may be kept together quite safely if they are all of about one size. There might be some cannibal tragedies if this were not the case, especially where the sand lizard is caged. They should be liberally fed on flies, grasshoppers, beetles, worms, and spiders.

Acanthus-footed Lizard.—This little animal comes from North Africa and is usually a light brownish-yellow colour, with bright piercing eyes and a tail much longer than its body. It lives chiefly among stones where it subsists on the sand-beetles to be obtained in such places. It derives both its popular and scientific names from the fact that its hind feet are fringed something like the leaves of the conventional acanthus flower which adorns certain kinds of architecture. It will live very well in a vivarium where it can get access to plenty of sunshine (8 to 10 inches).

The Blue Lizard is an inhabitant of Central America. If the illustration be compared with that of the *schelopusik*, it will be noticed that both animals have the lateral fold running from the neck to the root of the tail. In fact, we might almost be justified in calling the blue lizard a *schelopusik* with limbs as the two are closely related. The animal in question thrives very well in an ordinary vivarium where it will eat almost any small moving prey (1 to $1\frac{1}{2}$ feet).

Common Skink.—Unlike the gecko the common skink has been prized for many centuries as a form of medicine. If swallowed alive it cured consumption! If dried and powdered it was supposed to cure some other disease!! The idea is still maintained in some parts of Arabia where portions of its body are administered as a drug. It frequents sandy deserts in North Africa and Central Asia where it subsists almost entirely upon beetles and flies. The colour is a shining rich brown broken with splashes of white and yellow. This is a good vivarium inmate, thriving well in a sunny situation if it has plenty of sand in which it delights to burrow (4 to 6 inches).

The Cunningham Skink is a native of the wooded parts of Australia, living upon insects, and to a certain extent fruits. Though quite harmless the appearance suggests anything but this, for it is thickly covered by a number of prickly scales, which will be noticed on referring to our illustration. In captivity this creature is exceptionally hardy. It seems to prefer a dry case, and it is particularly partial to ripe bananas ($1\frac{1}{2}$ to 2 feet).

Common Chamæleon.—Of all the lizards this species is the most peculiar. Though very broad when viewed laterally, its body is as thin as a paper-knife when seen from above. As these creatures are tree-dwellers their feet are specially adapted for firmly grasping the boughs, and often the prehensile tail assists in making sure of its foothold. The movements are usually sedate and deliberate, but there are times when the creature will indulge in a little violent exercise. When disturbed a healthy wild specimen will open its mouth almost at right angles, hissing loudly the while and woe betide the hand that too closely approaches, for it will sharply receive a most painful nip like that of a pair of pincers. The tongue is by far the most wonderful part of the animal. If a hungry chamæleon be supplied with a butterfly, it will first survey the situation, then slowly and deliberately move towards the unfortunate insect till within about seven or eight inches of it. Then with a sudden leaning forward of the shoulders it slightly opens the mouth and presses the tongue against the upper jaw. This is followed by the protrusion of the tongue for about an inch, then an instant after it is flung like lightning against the victim and as quickly withdrawn, but with the insect sticking to the tip. The whole process occupies but a few seconds. The pressing of the tongue against the upper jaw, previous to striking, is considered to be for the purpose of supplying the tip of that member with an adhesive substance which causes any insect so touched to stick to it. But what has given rise to so much fable and story about this creature is the power of changing its colour, and even its pattern. However there are several, if not many, other animals which have this ability to a greater degree than the chamæleon itself. There are a certain number of shades of black, brown, yellow, and green which it can assume, but it has not the power often accredited to it of changing to any hue it desires. At night when asleep the common species is always yellow; in the full glare of the sun it is usually dark brown. If, however, it is frightened a large number of emerald green spots often make their appearance. In a state of nature the colour usually harmonises with the foliage. Yet another peculiar faculty is the chamæleon's. To make up for the stiffness of the neck the eyes are wonderfully active in their sockets, and furthermore while one is fixed upon an object before the animal the other can be studiously watching some insect behind. The two eyes are capable of independent vision. At first this was considered to be peculiar to chamæleons alone, but other animals

have since been observed to possess this strange power. The smooth blenny, abundant in our rock pools at low tide, may be cited as an example. The common *chamæleon* is a native of Africa and India, and feeds exclusively upon soft-bodied moving creatures, as insects and spiders (6 to 12 inches).

The Common Boa is the snake usually depicted in the old natural history books swallowing an ox or some animal of such dimensions. Though it not infrequently reaches a length of a dozen feet, such prey is quite beyond its mastery. However, this snake has been known to swallow a small antelope or a fair-sized dog, but rats and birds are the common prey. In the South American forests which constitute its home the day time is passed in a lethargic condition among the boughs of trees to which its body bears a great resemblance. But when night comes on, it becomes very active and ever ready to seize any unfortunate rat which indiscreetly ventures too near the supposed branch. Before swallowing, the prey is crushed to death in the powerful coils of the reptile. In spite of their great strength these snakes are singularly good-tempered and make admirable pets. The one photographed was nearly eight feet long, and though it strongly resented "sitting" for the camera, it never once tried to use its formidable teeth when it was being made to assume a suitable attitude (8 to 12 feet).

The Ringed Snake is one of the three British ophidians, only one of which, the viper or adder, is poisonous. As the staple diet is frogs, it is commonly met with in the neighbourhood of water between the months of April and September. Toads seem to be invariably refused, but newts and fish are greedily swallowed when opportunity affords them. This snake may be easily distinguished from the viper, to which it really bears very little resemblance, by the yellow coloured mark just behind the head. Besides the olive colour of the snake is quite different from the reddish brown of the adder. On account of this yellow blotch just referred to the creature has been called the Collared Snake, while Grass Snake and British Water Snake are two others of its appellations. The Ringed Snake makes a very good vivarium specimen, as it will, of course, stand our vacillating climate quite well and being exceedingly handsome is a real acquisition there. The ground colour, though very variable, is usually an olive green, marked with several rows of dark patches. The ventral surface is bluish in light specimens and almost black in darker varieties. This snake altogether lacks the dark zigzag chain of markings invariably found down the back of the viper. As this creature is neither poisonous nor a constrictor, it is unable to kill its prey and therefore always swallows the victims alive (2 to 4 feet).

The Dice Snake is closely allied to the grass snake which it resembles in habits in almost every particular. Though it does not occur in Britain, like the last species, it is found abundantly all over the Continent. Though quite innocuous, brown varieties of it sometimes resemble the viper, but a closer examination will quickly dispel all doubt on the subject. It always lacks the yellow collar of the grass snake, and the ground colour ranges from dull green above with a bluish tint below to reddish brown on the back with a pinkish ventral surface. It is quite as hardy as the grass snake and will live as well in captivity where it should be fed on live newts and fish (2 to 3 feet).

The Dark Green Snake is a native of Southern Europe. It is often called the Angry Snake on account of its habit, when wild, of

furiously biting when alarmed. However, as the nip seldom draws blood, and the reptile rapidly becomes tame in captivity, it is very suitable for the vivarium, where it should be fed on live young mice and lizards. This species is very variable, some specimens being quite yellow closely spotted with black, while others are very dark with but a few light marks near and about the head. But in all the varieties the tail seems to be marked with seven or eight dark parallel longitudinal lines (2 to 3 feet).

Æsculapian Snake.—This reptile has great historical interest, for in the time of the early Romans it was believed to be the messenger of Æsculapius, the God of Healing. Its appearance was always considered the omen of some gracious action on the part of the supposed deity. This belief seems to have taken such a firm root in the imagination of the superstitious Romans that many ancient writers record the fact that the Legions always carried a number of these sacred reptiles when on their empire making expeditions. Of course the snakes have absolutely no power over disease, and neither had any of their ancestors. This species will do very well in a vivarium if fed regularly on young mice, birds, or lizards. Its colour is usually dark brown above and greyish-yellow below with a very few yellow markings on the head. When wild, it bites furiously, but this vice soon leaves it if the reptile be treated reasonably in captivity (2 to 4 feet).

The Leopard Snake is one of the most beautiful of the European reptiles. The colour in the typical specimen (which somewhat suggests that of a leopard) is a pinkish-yellow, spotted with large bright red blotches. The ventral surface is of a greyish-blue tint and so glossy that it is beautifully iridescent when viewed obliquely. This snake, too, is very spiteful when first caught, but, like the others already mentioned, it soon ceases to resent being handled if properly treated. It is tolerably hardy in captivity, but its colouration is the chief factor in its popularity as a vivarium specimen. Young mice and lizards are its favourite food (2 to 3 feet).

Bull Snake.—This reptile comes from North America where it feeds for the most part on rats and birds. It derives its strange name from its supposed power of bellowing like a bull. This, like most popular notions concerning snakes, is probably a myth. In colour it is usually creamy blotched with large irregular patches of dark brown. It is very hardy and, in spite of its large size and great strength, is quite good-tempered. Sometimes this reptile is known as the Pine Snake (3 to 5 feet).

Smooth Snake.—This little reptile, which seldom exceeds two feet, is the rarest of Britain's three ophidians. [Of course, like all other snakes, it is absent from Ireland]. Its discovery in these Islands was of comparatively recent date and since then it has been captured in several of the British counties. The New Forest seems to be its headquarters this side of the Channel, and it is exceedingly rare even there. However, it is fairly common, though local, on the Continent. At first sight it much resembles the viper, but instead of the single zigzag chain of markings on the back of the adder it has a double row of black blotches. The ground colour is dark grey, and the ventral surface a bluish black. In a state of nature it feeds on lizards and blindworms, usually constricting them before swallowing. In captivity, where it is very hardy, it seems to refuse any other food (1 to 2 feet).

King Snake.—This is another of the North American ophidians which is hardy enough to withstand the rigours of the amateur's vivarium. It is allied to the British smooth snake, but grows much larger than its European relative. Small specimens are the most suitable for reptile cases, for the snake is somewhat prone to cannibalism. The colours are yellow and black, delicately mottled in bands across the creature's body. Birds, rats, and mice seem to be its natural food (4 to 5 feet).

Hoop Snake.—The first impression that one has when viewing a motionless specimen of the above is that it is some fantastic design in colour worked upon a china snake, and furthermore that the artist who modelled it has never seen a real specimen of any reptile. However, such ideas are short-lived for the restless creature is sure to move before long. It never seems to be happy unless it has twisted itself into a sort of complicated letter S, with its curious unsnake-like head buried somewhere beneath the coils. A minute or two later it will have found this position not to its liking and will have proceeded to try another of the same variety. The scales are tightly packed on it, somewhat like those of a blindworm, and are so glossy that the creature looks like a piece of highly polished porcelain. The colours are bright red and shining black scattered almost at random all over the dorsal surface. On account of this its popular name is often the Red and Black Snake. Some old writers have affirmed that when this ophidian (which is absolutely harmless) wished to chase other animals for the purpose of poisoning them, it would place its tail in its mouth and, with the circle or hoop so formed, bowl along after its fleeing victims with lightning like rapidity! Possibly this legend has been the means of naming this creature the Hoop Snake. It will do well in the summer time in captivity if fed on young rats and mice (3 to 6 feet).

The Viper.—Fortunately Britain only nurtures this one poisonous reptile, though of the three indigenous species it is by far the commonest. It is usually met with in dry situations, such as pine woods or heather moors, where it spends a good deal of its time out in the open basking in the sunshine. The grass snake, on the other hand, is usually found near ditches or in damp places generally. It is quite easy to distinguish between the two species. In the first place the grass snake has a conspicuous yellow bar across the neck, while the adder is always without any light marking there. Instead it has a clumsy V-shaped dark blotch on the head, the back too of the latter is ornamented with a dark chain of zig-zag markings, which is never seen in the former. It has instead usually four rows of small deep green spots. The adder is more often than not of a reddish-brown, while the British grass snake is olive green. If a viper be cornered without chance of escape it coils itself up in such a position that any moment it can throw its head violently against its assailant. Hissing strongly it awaits results. When its enemy is close enough the head is darted forward with open mouth so that the fangs are thrust into the intruder's flesh, and are as rapidly withdrawn. If the enemy be a human being, for the first few minutes nothing will be observed save two tiny punctures in the skin. But in a very short time a sharp pain will be felt, and the flesh in the neighbourhood of the bite will begin to swell rapidly. This will continue until, perhaps, the whole of the bitten limb will be affected. If the person be healthy, before a week is out a complete recovery will have been made, but even then the experience is not particularly pleasant. Of course where such things as mice, lizards, or birds are struck death results in a few minutes, and this is often the method in

which a viper secures its prey. The fangs which inflict the wound are very interesting. In the upper jaw there are two teeth, which are so jointed that they can be closed into the gums in the roof of the mouth, or turned so that their points come forward and project a little beyond the top lip. These teeth, which are the fangs, are moreover tubular even to their points, and at the other ends are connected, by means of ducts, to glands which secrete the venom. When the viper wishes to inflict a poisonous wound it opens its jaws almost at right angles, throws forward the fangs, and plunges them into the flesh of its victim. As the teeth enter the skin the poison glands contract by means of powerful muscles, forcing the venom contained there down the ducts and out of the hollow teeth deep into the wound. The fangs are then instantly withdrawn, the whole operation occupying but a few seconds. This species cannot be recommended as a vivarium specimen, though it is exceedingly handsome (12 to 15 inches).

If any who read these notes would care to keep a vivarium, there are several preliminary precautions to be observed if success is to attend the effort. An empty glass aquarium will make a tolerably good case, but a little ingenuity and a Tate sugar-box will make a better one. In the former all the ventilation will have to come down through the top, in the latter suitable holes can be made in the sides near the bottom as well, which will keep the air circulating perfectly throughout. This second method is very much more healthful than the first. The bottom of the case should be covered with moss, and it is a good plan to keep one portion of it quite dry and the other part very moist. In this way the creatures can choose which they prefer to lie under, and their choice will not always be the same. Virgin cork bark should be liberally supplied for shade and hiding places, and a few small branches will add to the natural look of the case. A vessel full of clean water should always be there in such a place that the inmates can drink without the slightest difficulty. The vivarium should never be allowed to remain in strong sunshine unless the creatures within are known to naturally live in such a tropical atmosphere. Tortoises and alligators will live together comfortably, but snakes should never be confined with lizards or many of the latter will disappear.

The Author is indebted to the Proprietor of the Bedford Conservatories, Covent Garden Market, London, for the loan of many of the rarer specimens figured in the illustrations.

Gowans's Nature Books

THE object of these little books is to stimulate a love for nature and a desire to study it.

The first volumes of the series have been extremely successful, and the publishers hope to make the others equally interesting and beautiful.

- No. 1.—WILD BIRDS AT HOME. Sixty Photographs from life, by Chas. Kirk, of British Birds and their Nests.
- No. 2.—WILD FLOWERS AT HOME. First Series. Sixty Photographs from Nature, by Cameron Todd.
- No. 3.—WILD FLOWERS AT HOME. Second Series. By Cameron Todd.
- No. 4.—BUTTERFLIES AND MOTHS AT HOME. Sixty Photographs from life, by A. Forrester.
- No. 5.—WILD BIRDS AT HOME. Second Series, by Charles Kirk.
- No. 6.—FRESHWATER FISHES. Sixty Photographs from life, by Walford B. Johnson and Stanley C. Johnson, B.A.,
- No. 7.—TOADSTOOLS AT HOME. Sixty Photographs of Fungi, by Somerville Hastings, F.R.C.S.
- No. 8.—OUR TREES & HOW TO KNOW THEM. Sixty Photographs by Charles Kirk.
- No. 9.—WILD FLOWERS AT HOME. Third Series. By Somerville Hastings, F.R.C.S.
- No. 10.—LIFE IN THE ANTARCTIC. Sixty Photographs by Members of the Scottish Antarctic Expedition.
- No. 11.—REPTILE LIFE. Sixty Photographs by Walford B. Johnson and Stanley C. Johnson, B.A.
- No. 12.—SEA-SHORE LIFE. By the same.

Others in Preparation

PRICE:

PARCHMENT COVER, 6D. NET. EACH VOLUME.

POSTAGE, 1D. EACH.

GOWANS & GRAY, Ltd., London & Glasgow

Gowans's Art Books

THIS Series, published at a very low price and containing **sixty** beautifully-printed reproductions of the best pictures of the great masters, is intended to give the lover of art a general idea of the style and characteristics of the most famous painters of the world.

No. 1.—THE MASTERPIECES OF RUBENS.

No. 2.—THE MASTERPIECES OF VAN DYCK.

No. 3.—THE MASTERPIECES OF REMBRANDT.

No. 4.—THE MASTERPIECES OF RAPHAEL.

No. 5.—THE MASTERPIECES OF REYNOLDS,

No. 6.—THE MASTERPIECES OF TENIERS.

No. 7.—THE MASTERPIECES OF THE EARLY
FLEMISH PAINTERS.

No. 8.—THE MASTERPIECES OF TITIAN.

No. 9.—THE MASTERPIECES OF FRANZ HALS.

No. 10.—THE MASTERPIECES OF MURILLO.

No. 11.—THE MASTERPIECES OF WOUVERMAN.

No. 12.—THE MASTERPIECES OF VELAZQUEZ.

Others in Preparation.

PRICES:

PARCHMENT COVER, 6D. NET.

CLOTH, 1s. NET.

LEATHER, 2s. NET.

POSTAGE 1D.

GOWANS & GRAY, Ltd., London & Glasgow

Gowans's Books for the Young

A SERIES of prettily got up little volumes containing old favourites and other books sure to become favourites.

Ready.

No. 1.—THE SEVEN CHAMPIONS OF CHRISTENDOM. By W. H. G. Kingston.

No. 2.—THE HEROES. Greek Fairy Tales for my Children, by Charles Kingsley.

No. 3.—THE STORY OF PRINCE AHMED AND THE FAIRY PERIE BANOU (from the Arabian Nights).

Others in Contemplation.

PRICE: PICTURE BOARDS, CLOTH BACK, 6d. NET.
POSTAGE, 1d.

GOWANS & GRAY, Ltd., London and Glasgow