582.35 (410)

FRA

AN
ANALYSIS
of the
BRITISH FERNS
AND
THEIR ALLIES.

by

G. W. FRANCIS, F.L.S.

THIRD EDITION.

TO
SIR WILLIAM JACKSON HOOKER, LL.D., F.R., A. & L.S.

&c. &c. &c.

AND
REGIUS PROFESSOR OF BOTANY IN THE UNIVERSITY OF GLASGOW.

SIR,

To you, who stand so far before all other Writers in a practical knowledge of British Botany, and who have made the Ferns so particularly the subject of your attention, I beg respectfully to dedicate this little Work. I am aware that it is too small to be worthy of your attention; but I am anxious to take as early an opportunity as possible of offering my homage to those brilliant talents which have contributed so essentially to diffuse a love of Botany; that energy without which even talents are unavailing, and that urbanity of manners and liberality of feeling for which Botanists have always been celebrated.

That you may long be spared the full enjoyment of all your mental and physical faculties, to cheer your Friends and to instruct the World, is the ardent wish of

Sir,

Your most obedient Servant,

THE AUTHOR.

PREFACE.

“I acknowledge no authority but that of observation.”—LINN.

This motto was my governing principle in writing the following work on the “British Ferns and other Allies;” and in adopting it I hope that I shall neither be accused of arrogance, neglect of the opinion of others, nor yet of unnecessarily varying the details of science. Should the reader ask, Why I write at all? I answer, because the only book ever published upon this subject, (Bolton’s “*Filices Britannicæ*,”) has long been out of print; and so much difference of opinion exists as to the identity of some species, and the arrangement of others, that I thought a plain and practical synopsis like the present would be useful to the tyro, if not to the practical botanist.

The materials from which it has been compiled are these:—I inspected all the herbaria to which I had access; gathered wild and cultivated fronds wherever I could procure them; and wrote to most of our first-rate botanists for specimens, remarks, and habitats. All these being collected, arranged, and studied, they were described and engraved without reference to any series of plates or descriptions whatever. I then collated these with the works of Linnæus, Willdenow, Sprengel, Swartz, Pursh, Withering, Smith, Hooker, Lightfoot, Hudson, &c. &c., and wherever there was a difference between myself and others I searched again for the truth; and, if still in doubt, have been careful to record the disparity.

The long introductory matter explains all that is known of the internal structure, not only of the indigenons species, but of foreign also; and as it tends to induce in the mind a philosophical knowledge of the plants afterwards detailed, I flatter myself that the part devoted to this will not be the least valuable to the student of nature.

The manner in which the object has been accomplished it is necessary to explain more fully: and, first, as to the illustrative plates—they are small, for the sake of economy, and are intended chiefly to indicate the habit of the plants, while the magnified parts show their detail. They might have been finer as works of art; but, had they been executed by an engraver, minute as they are, they would perhaps have been less botanically accurate, as the smallest variation in many of them would have materially altered their character; and, therefore, although a first and an untutored attempt at etching, I have preferred executing them myself, especially as by so doing I should save a large expense, perhaps not be refunded by the sale of the work. The engravings of genera is a new feature in illustration, and it is hoped a useful one.

In the record and detail of species the following order is observed:—First, the Latin and English name, and reference to figure. Secondly, those essential characters which alone are necessary for discriminating the species, and which alone the true botanist will find it convenient to consult. The synonymes and references to figures in other works which follow, give a history of the plant, and enable the student to refer elsewhere, if in doubt. The description may be considered collateral evidence, while the remaining parts will show him the varieties to which his plant is subject, the cause of them, its particular and general distribution, and the peculiarities attached to it.

In the part of the work which treats of the genera, the reader will find, first, the derivation of the genus, and a concise account of its general characteristics, and under it the arrangement of the species, according to their obvious distinctions.

In the essential characters of the different species, as few words as possible have been used, and those few pure and scientific. In the synonymes, which go back to the time of Linnæus, (or in some few instances a little before,) the names of authors only are given, unless they have called a plant by different names in different of their works, when the works themselves are also specified. In the descriptive

part, and discriminating remarks which follow it, pure scientific detail, has not been so much aimed at as obvious differences and popular observations. The habitats have been collected from every authentic source which was attainable by me; a vast number will be found which have not been recorded before, and those few which are contained in previous publications have most of them been lately authenticated.

Information of this varied and local kind cannot, of course, be expected from any one's unassisted labours. I have therefore had recourse in the latter part to the assistance of friends, and I cannot speak too highly of the kindness and warmth with which my advances have been received, and without which indeed very much interesting matter must of necessity have been omitted. I hope that I have acknowledged in every instance the remarks thus received. My obligations are particularly due to H. C. Watson, Esq., who lent me the numerous lists, and manuscript localities, the results of his own observations, or communicated by botanical correspondents whilst he was preparing the "New Botanist's Guide;" also for the valuable remarks of Mr. W. Wilson, Mr. W. Leighton, Dr. Murray, Rev. W. Bree, Mr. W. Pamplin, Mr. Bevis, and Mr. Castle.

And now, kind reader, I leave the work in your hands, concluding with the words of Linnæus to Haller:—"If you have remarked errors in me your superior wisdom must pardon them. Who errs not while perambulating the domains of nature? Who can observe every thing with accuracy? Correct me as a friend, and I as a friend will requite the kindness."
(*March 1, 1837.*)

The above remarks formed the preface to the first edition. The alterations introduced into the present are such as my own further study of this tribe, or the remarks of various kind friends have suggested. The plates have been all re-engraved, and of a larger size, consequently they are more numerous; and the plants, which it was almost impossible to express in the former very minute sketches, are more clearly defined. The general character of the plates are, however, similar, those of both editions having been drawn from almost in every instance the very same specimens. The wood-cuts, now for the first time introduced, will, it is believed, be found useful.

The introduction contains much new and valuable matter on the organic structure of certain species, which the laborious researches, and the acumen of Messrs. Henderson, Valentine, &c., have enabled me to avail myself of. Mr. H. C. Watson also has kindly contributed remarks of real interest on the geographical distribution of the various tribes; and Mr. Bevis, of the Royal Botanic Garden, supplied me with the ground-work of the appendix,—my own cultivation of the class having been very limited. The habitats have been arranged with more regard to latitude, but they have not been materially increased. That portion of the work devoted to scientific description has, no less than the other parts, been carefully corrected, yet the alterations will be found very few and unimportant.

My opinion of the identity and distinctness of certain species remains unaltered; yet I am aware that some writers differ from me, and would unite two of the Polypodiums; *Asplenium alternifolium*, and *Ruta muraria*, &c., founding their opinion upon specimens preserved in herbaria. I venture to protest against a too general reliance being placed upon this source of information. Not but that reference to celebrated herbaria is most valuable, in addition to research; but this is all—a knowledge of plants can never be acquired by such means, because herbaria do not testify the circumstances of aspect, season, altitude, moisture, soil, and other influences, to which plants are subject. I make this remark chiefly in reference to the *Cysteas*, *Woodsias*, some of the *Aspidiums*, &c., about which botanists differ more than about any other genera. It is true that numerous fronds may be found of an intermediate character between two species; still, unless circumstances of growth be corresponding to all, this does not prove identity between the remote examples. This observation particularly applies to Ferns, because the diagnoses of the species rest mainly upon the shape of the frond, and this is the part which is alone subject to variation. The herbaria of even the most celebrated botanists are not always to be depended upon for accuracy. Plants are often received named—the name without examination is almost as frequently considered correct, and that specimen forms henceforth a criterion whereby to name others; thus error is continued and multiplied. To these causes I am inclined to attribute the contrariety of opinion which exists relative to our Ferns; and I may add, perhaps, to our Willows, Roses, and Brambles.

It may, perhaps, be expected that I should have said something relative to the new system of classifying Ferns according to their venation. I have not done so in the body of the work, because I do not find the system either correct, convenient, or practicable. For example, the veins of the British *Trichomanes* and *Hymenophyllum* are precisely similar, yet few would assign all these to the same genus. The veins of *Polypodium vulgare* and *phegopteris* are very different from each other, yet the plants are conveniently placed together. The veins of the latter plant resemble those of *Aspidium oreopteris*, yet I cannot consent to unite the two into one species. As to the veins of *Grammitis ceterach*, Mr. Newman shows them as anastomosing; Mr. Presl as distinct and unattached at their extremities. Mr. Smith, Curator of Kew Gardens, and who has paid much attention to the subject, says, "that neither of my figures, (and which are copied, the one from Newman, the other from Presl,) is correct." And as to my own opinion, I confess I cannot make them out at all to my satisfaction, and that is the case generally with the coriaceous Ferns, particularly after having been dried for the herbarium.

June 1st, 1842.

A third edition of this work being now called for, I am sorry that I have it not in my power to offer any additional remarks upon the plants described. I would not have it inferred, however, that I consider my book either without error or complete, but merely that having been, since the issue of the second edition, resident on the Continent, and at all times much occupied, I have made no additions to my own stock of Fern knowledge; and moreover I never expected that a third edition would be called for, and therefore made no preparation for it.

I have to remark, however, that I have been blamed by one or two botanists for not discarding some of the present names, and adopting the more ancient, *Lastræa*, and other. I desire to pay the greatest deference to the opinions of others, and am most happy to learn from all, as to matters of fact at all times, yet a change of names is always perplexing, and in this case unnecessary. I cannot admit either that the amount of evil is very greatly less in re-establishing an obsolete nomenclature than in coining a new one. As to the mere intention assigned by some of restoring their due honor to ancient botanists, it is a principle good in theory, but which cannot be carried out; and would be of much injury to botany if it could: as we should be soon overwhelmed with terms now forgotten, and which are generally less appropriate and significant than the more modern appellations.

GEORGE FRANCIS, FLS.

July 1st, 1847.

INTRODUCTION.

A FERN is a flowerless plant which has a fibrous root, vascular stem, nerved leaves, reticulated cuticle furnished with stomata; and which bears spores as fruit in capsular receptacles.

The Ferns and their Allies form the first order of the Linnæan class Cryptogamia, and the structure of them shows so exactly an intermediate character between the Vasculares and Cellulares, that all systems of classification have assigned them this station among vegetables. They are without flowers, have but imperfectly-formed vessels, and no deposition of real woody fibre, they therefore cannot with propriety be arranged with Phænogamous plants; while their semi-vascular texture and fully-developed leaves show their organization to be greatly above that of any other order of Cryptogamia.

Although the True Ferns have a direct analogy with the Palmæ and Cycadææ, the connexion between them and other orders is more apparent in the Pteroides or Fern Allies, particularly the Equiseta and Lycopodia. The Equiseta are nearly connected with several orders of Flowering plants. In their hollow, jointed, silicious stems, they resemble the Grasses; in other respects the Coniferæ and Amentacææ, approaching the one by means of the genus Casuarina, and the other by that of Ephedra, nor are they far removed in structure from the Charas; thus connecting also the Ferns and the Algæ. The other of the Fern Allies, the Lycopodia, were considered by the earlier botanists as Mosses, so slightly do they differ from that tribe, not only in habit, but in many important characteristics.

Thus the tribes under consideration, which are divided according to the modern system into FILICALES, LYCOPODALES, and EQUISETALES; the first the True Ferns, the others the Pteroids or Fern Allies, altogether form valuable, because well-connecting links in the great chain of nature.

The scientific division of the Ferns into tribes and orders was long a desideratum in botany. The earlier schemes are too vague for the present state of science. That of Willdenow, in which he depends upon external characters alone, is still however used ; it divides the whole Ferns as follows :—

1. GONOPTERIDES, which includes only the genus *Equisetum*.
2. STACHYOPTERIDES, including *Lycopodium*, *Botrychium*, and *Ophioglossum*.
3. SCHISMATOPTERIDES, containing only *Osmunda*.
4. FILICES, which comprises all the Dorsal and Marginal Ferns.
5. HYDROPTERIDES, (Water Ferns,) containing *Isoetes* and *Pilularia*.

Valuable as the system of Willdenow was, compared to all previously adopted, it is very far inferior to that of Sir J. E. Smith, improved as this has been by Mr. R. Brown, Mr. Kaulfuss, and others. Here, not only the external character of the fructification, but the structure of the fruit itself, and of its envelopes, is considered of importance as a guide to essential characters ; while the shape, division, and habit of the plant are used only in the discrimination of species : thus classifying the Ferns as much as possible by the same laws as those which govern higher orders of vegetation.

FILICALES.

ANNULATÆ, which have their capsules or thecæ ringed.

Thecæ in clusters at the back of the frond, bursting irregularly and transversely. Ring vertical or a continuation of the footstalk of the theca. Stems solid, their vernation circinate.	}	POLYPODIACEÆ.
---	---	---------------

Thecæ in two-valved receptacles on the margin of a frond, bursting irregularly and vertically. Ring oblique. Vernation circinate. Stems solid.	}	HYMENOPHYLLACEÆ.
--	---	------------------

EXANNULATÆ, which are without a ring to their capsules.

Thecæ in clusters, terminating a leafy frond, bursting at a longitudinal suture, two valved, transparent, petioled, striated. Vernation circinate. Stems solid ..	}	OSMUNDACEÆ.
---	---	-------------

Thecæ in spikes or racemes attached to a leafy frond, bursting at a transverse suture, two valved, opaque, sessile, smooth. Vernation straight. Stems hollow ..	}	OPHIOGLOSSACEÆ.
---	---	-----------------

LYCOPODALES.

Thecæ of two kinds, indehiscent, inclosed within the base of radical leaves. Leaves hollow and filiform. Stem none, Vernation straight. (<i>Water plants</i>)	}	ISOETACEÆ.
--	---	------------

Thecæ of one kind, coriaceous, scaly, seated near the base of radical leaves. Leaves hollow and filiform. Stem long and creeping. Vernation circinate	}	MARSILEACEÆ.
---	---	--------------

Thecæ of two kinds, axillary in a leafy spike or stem, two to four valved, sessile, free, dehiscent at a regular fissure. Stem solid, leafy, creeping or upright	}	LYCOPODIACEÆ.
--	---	---------------

EQUISETALES.

Thecæ in terminal conical catkins, bursting at a longitudinal fissure. Spores attached to four filaments. Stems leafless, striated, hollow, jointed	}	EQUISETACEÆ.
---	---	--------------

POLYPODIACEÆ.

(Including *Grammitis*, *Polypodium*, *Woodsia*, *Cistopteris*, *Aspidium*, *Asplenium*, *Scolopendrium*, *Blechnum*, *Pteris*, *Cryptogramma*, *Adiantum*.)

POLYPODIACEÆ, Br., D.C., Kaulf., Bory, Hook., Grev., Mack., &c.;—
 FILICES VERÆ, Willd., Linn, Schreb, Juss.;—FILICES ANNULATÆ, Hoffm.;
 FILICES GYRATÆ, Web., Mohr., Swtz.;—FILICES DORSIFERÆ, Smith;—
 FILICALES, Lindl.;—PTERIDALES, FILICALES PHYLLOPTERIDES, EPIPHYL-
 LOSPERMEÆ, &c. &c.

STRUCTURE.*—A FERN consists of root, frond, and fructification. The rhizomas or subterranean stems, as well as the fibrils or true roots, are included under the first term; while the frond comprises every part above the ground, except the fruit and its appendages; and is subdivided into rachis or stem,† and pinnae or leaves, which latter are generally more or less compound, lobed, or indented.

THE ROOT of all our native Ferns is perennial and fibrous. The fibres are stout, generally hairy or scaly, and in many instances furnished at the extremity with hoods or sheaths, the use of which is not very obvious. Modern botanists agree with Sprengel in believing them the organs of absorption, as the hoods of the *Lemna* and some other of our water plants. Röth maintained that they were mere defensive organs, intended to prevent the introduction of the grosser fluids, and to shield the extremity of the fibre from injury. In many cases the fibres issue from a crown, and form a tufted root; in others from thick stems, which in the British species creep under ground, sometimes to a considerable distance. These rhizomas or creeping stems are furnished with buds, irregularly disposed upon their surface; the uppermost ones yielding fronds, while those below produce as invariably radical fibres.

THE RACHIS is sometimes smooth, at others scaly or hairy, sometimes wholly clothed with leaf-like expansions; at others void of them at the lower part. When cut transversely, it is seen to consist first of a cuticle; then we find a hard, woody, green, brown, or black bark, the space within being filled with cellular tissue. Longitudinally through the tissue run bundles of sap vessels, most of which are true spirals; not, however, formed of a cylindrical thread, coiled up as in more

* In illustration of the structure of all the tribes, the reader is referred to the illustrations of genera and their explanation. The stomata afterwards spoken of may very easily be seen in any under part of the cuticle, merely by tearing it off and submitting it to the microscope. The arrangement of the vessels in the stem is apparent to the unassisted eye in any transverse section of it; and to view the spiral vessels it is only necessary to take two pins, and having thrust them through one of the bundles of vessels, separate them a little from each other, and in the cleft thus made the spirals will appear distinct when considerably magnified.

† In the progress of the work it has not been thought necessary to make a difference between the rachis and stipes, nor to divide the part under ground into root and rhizoma, the first term of each being sufficient.

perfect plants, but rather of a flat band, like a riband rolled spirally on a cylinder. Mixed with these vessels, which are real tracheæ, are tubular perforated ducts. The whole, thus compounded of the two sorts, is sometimes collected into a close bundle, but more generally into a cylindrical sinuous ring, either hollow or filled with cellular tissue, and surrounded with a dark membrane. The number of these fascicles bears considerable relation to the size of the frond; thus in *Pteris aquilina*, there are eight or ten; in *Aspidium aculeatum*, five; in *Polypodium vulgare*, three; while in the minuter species there is but one, which then occupies the centre of the rachis. When several bundles are present, no general rule can be given for their position, (though constant in the same species,) so varied are they in shape, size, and distance from each other. It is thought that the depression so often visible on one side of the rachis is occasioned by the absence of vessels on that part.

The cellular substance appears to have no tendency to arrange itself in strata, nor do the vessels increase in number as the plant increases in age. The stems, therefore, contain no real wood; the nearest approach to it being the hardened cuticle and the ducts themselves. They increase very little in diameter, but grow longitudinally throughout their whole length.

THE FROND is in its leafy part thin, veiny, and green. The veins do not extend longitudinally through the leaf in any species, as in the Monocotyledones, but diverge in a forked form, (dichotomously divided,) from the base of the leaf, or from the midrib; differing, however, from those in dicotyledonous plants in not containing woody fibre, and in being uniform in size throughout all their ramifications, and therefore more properly called nerves than veins. The divisions of the frond are for the most part constant in the same species, but varied in their size and number by external circumstances; the primary causes of which are superabundance or deficiency of nutriment, while temporary heat or moisture, exposure, shelter, or season of the year, occasion other but less striking irregularities. Even these causes have but little effect over numerous kinds, and very seldom in any case do they occasion so great an alteration of ordinary characters, as to throw doubt upon the species. (See *Cistopteris fragilis*.) The Ferns are several years before they come to maturity, until which their essential characters are not always obvious. Thus young plants of *Aspidium Filix-mas* very much resemble *Woodsia ilvensis*; they are first pinnatifid, then pinnate, afterwards when perfect nearly doubly pinnate. Also when a Fern has its barren fronds different from those which are fertile, the latter are more contracted, as if the sap which expanded the leaves of the one was employed in nourishing the fruit of the other.

THE VERNATION.—The circinate vernation, or curling up of the unexpanded frond, which prevails in all the dorsal Ferns, is almost peculiar to this tribe and one of their allies, being found in only two other orders; namely, the Palmæ and Cycadææ. If the frond be simple, so is the vernation, resembling a flat spiral spring; but when the frond is subdivided, the vernation becomes equally compound, the larger divisions first opening, and by degrees the branches, pinnæ, and lobes.

THE SCALES so visible upon some species, have been thought merely an excrescent growth caused by superabundant sap exuding from the surcharged pores. Sprengel supposed that they were part of the epidermis itself, lacerated by the pressure of the juices beneath. Perhaps both of these opinions are correct, a part of the epidermis of the frond being first detached, and afterwards nourished in the same manner as animal hair, and although void of life yet increasing from the base.

THE CUTICLE of the leafy portion of the frond presents a reticulated appearance, (the meshes having wavy sides,) and is furnished on the under surface with respiratory stomata, similar in form and function to those of Flowering Plants. The number of these on a given space is in a great degree accordant with the rapidity of the frond's withering when gathered. They are very abundant in *Aspidium filix-fœmina*, *Aspidium dilatatum*, and *Polypodium Dryopteris*. Thus is explained the cause of the drooping habit of this last and some other species. Be it observed, however, that in *Grammitis ceterach* and *Aspidium lobatum* they are still more numerous, yet these latter plants do not wither so soon, a circumstance that may easily be accounted for from the frond of both being thicker, the deprivation of an equal quantity of water not producing so great an effect.

THE REPRODUCTION of Ferns is a subject involved in much obscurity. Hedwig, Bernhardt, and others, have proposed theories to explain this intricate matter, but without success. That the Ferns have no visible flowers is evident, but that they have some apparatus analogous to stamens, is maintained by most of our first botanists. So keen has been the search for these in the present tribe, that every part of the plant has been subjected to the minutest investigation: not only the thecæ, their ring, and their cover, but the spiral vessels of the rachis, the stomata upon their cuticle, and the glands which are sometimes found attending upon them.

Sprengel long ago stated that the young sori, or rather that the swelled extremity of the veins of *Polypodium vulgare*, which sometimes remain abortive, and at others produce thecæ, were filled with oblong-shaped bodies of a greyish color, which he considered to be stamens, and as yielding pollen in the same manner as the external stamens of flowering plants. These being attended upon by young ovules, the latter became impregnated and grew to perfection, while the pollen masses were decayed or absorbed. This opinion was in a great degree and for a long time disregarded, (perhaps because of his imperfect figures,) and the grey bodies considered to be abortive capsules. The following remark, however, from "The Annals of Natural History," given in the synopsis of a paper read before the Academy of Sciences at Berlin, March, 1840, by Professor Link, throws a stronger light upon the subject. "The part which Sprengel years ago indistinctly figured, and which Blume and Presl at present consider to be male organs of fructification, have been more accurately examined by Professor Link, and illustrated by drawings. They are long hollow filaments, separated by septa into articulations generally simple, rarely ramified; the last articulation is thicker,

and filled with a delicate granular mass. It may also at times be observed that this mass is exuded at the last articulation, and surrounds this as a crust. These parts are frequently longer than the capsules, and are easily distinguished from the young capsules." The late Professor Don, who wrote the above, adds—"It is certainly probable that they are the stamina of Ferns, and indeed Link found them, after frequent search, in most of the Ferns which he submitted to microscopical examination." Mr. Henderson, in an interesting paper on "The Germination of Ferns,"* denies that there is any impregnation in Ferns, Mosses, or Equiseta; yet, in an after paper read before the Linnæan Society on "The Reproductive Organs of Equisetum," completely confirms the above view. He states that he has found two kinds of granules in the unripe thecæ of Ferns, Lycopodium, and Ophioglossum; that the one kind is mostly absorbed during the maturing of the other; and by submitting each to the test of iodine, he proved the one to be amylaceous or starchy—the other more of the nature of pollen. Thus the matter seems set at rest; and as Mr. Henderson has found similar pollen granules in the thecæ of Mosses and Jungermanniæ; in the apothecia of Lichens; in the lamellæ of Agarics, and in the perithecia of some other Fungi, his experiments assist much in elucidating other tribes, as well as that under especial consideration.

SEEDS OR SPORES, AND THEIR GERMINATION.—The small, round, rough grains contained in the thecæ, considered formerly as gemmæ or buds, are now known as seeds, yet differing from common seeds in many respects. They have no cotyledons, but are a mass of cellular substance. Instead of sending upwards a plumule, and downwards a radical, from fixed points, they grow indifferently from any part of their surface; that most exposed to light shooting into the future frond, while the deeper point propels the root. Owing to these differences the seeds have been called, not only here but in all the tribes of Cryptogamic vegetables, spores (or sporules) rather than seeds. They retain their vitality for many years, and those brushed from the dried plants of an herbarium will grow long after the specimens have been gathered, coming up first with a small crown or bud, from which soon issues a peculiar shaped frond not unlike that of a Lichen, or rather like that of Marchantia, and differing much from the fronds of future growth. When this has expanded to a certain size, according to the species, the centre of it, both below and above, becomes thicker; the lower part elongates into a root, while the upper part assumes a gyrate or circinate form, and gradually unfolds itself into an upright frond, of the same texture though much less divided than those afterwards produced.

THECA AND SORUS.—The spores in all the species are contained in capsules or thecæ, each of which opens at a transverse irregular fissure, and is furnished with a jointed spring, nearly surrounding it, and by the elasticity of which the capsule is torn open and the spores dispersed. The thecæ are collected into linear, oblong, or circular clusters, called sori, of which Professor Link thus writes:†—

* "Mag. of Zoology and Botany," v. 1., p. 340.

† "Ann. of Nat. History," August 1840.

“The sorus is in general situated on a receptacle which, when roundish, consists entirely of short spiral vessels, so called, vermicoid bodies, similar to the thickened extremities of the leaf nerves, which might therefore be regarded as abortive receptacles, (*query* thecæ.) In the elongated receptacles, spiral vessels are also met with.” The sori are in some tribes of the Ferns naked, but in the generality covered with a scale or indusium, of shape similar to themselves. I remarked in the first edition, that the origin of this integument was undoubtedly similar to that of the scales, namely, disrupted epidermis. Microscopic observations, however, induce me to doubt this assigned origin for the indusia, and to consider them as distinct organs, as much so indeed as the calyx of a flowering plant, or the calyptra of a Moss. Whether they arise from the vermicoid bodies of Link, just spoken of, or not, I have been prevented from observing. These certainly attend the genus *Woodsia*, and are intermingled with the thecæ, the genus having no real indusium. In the genera *Cystopteris*, of *Aspidium* *Adiantum*, *Pteris*, &c., the reticulation of the indusium and epidermis is very different, and the former is not furnished with stomata. In many genera this organ may be supposed a part of the frond itself turned over upon the thecæ, as in *Adiantum* and *Pteris*, but I believe the reticulation is very distinct.

NUMBER AND DISTRIBUTION.—For the following remarks upon this subject I am indebted to H. C. Watson, Esq., than whom no botanist has more studied the subject. He says,—“Dorsiferous Ferns are found in every part of Britain, except on the summits of the loftier mountains, and in small spaces of the lower grounds, whence they are banished by local peculiarities of the soil or surface. But overlooking these merely local exceptions, of trifling extent, *Polypodiaceæ* may be stated to range over the whole of Britain, from south to north; from east to west; and from the shores of the sea almost to the summits of the highest hills; in which latter situation their absence is to be attributed rather to the bleak exposure than to the absolute height. The number of our dorsiferous Ferns will be estimated variously, according to the views entertained with respect to union or division of reputed species, but 35 is the number most generally received. These bear a proportion to flowering plants, (reckoning the latter at 1400,) of 1 to 40. The order has a great numerical predominance over the other orders of *Filicales*, all taken together; the proportion of its species being to those of the other three orders, as 6 to 1. And since the most abundant and widely-ranging species of Ferns are also to be found amongst the *Polypodiaceæ*, the effect of this order in the general vegetation of our island much exceeds that of the allied orders of Ferns. Of the six species referred to other orders, one is exclusively an inhabitant of Yorkshire, (if it really be there still;) namely, *Trichomanes brevisetum*; a second, *Hymenophyllum Tunbrigense*, is local; a third, *H. Wilsoni*, though much more plentiful, is limited to the northern and western counties; whilst the other three, the *Osmunda*, *Botrychium*, and *Ophioglossum*, though widely scattered through Britain, are by no means so generally present as many of the *Polypodiaceæ*. Several of the British dorsiferous Ferns are so widely and plentifully scattered throughout the island, that

there can be no doubt about their existence in every county of Britain ; although published records will not enable any one to make the assertion on evidence. Others, on the contrary, are much more limited in their range, being absent from the southern, or northern, or lowland counties of England. Only a few species can be called decidedly rare. The mountain valleys below 1500 or 2000 feet are the head quarters of Polypodiaceæ ; very few species wholly shunning the mountain tracts, and a still less number being exclusively confined to the higher parts of the hills. The specimens are numerous amongst the mountain valleys, in the western counties, and in the vicinity of the coast. On the contrary, they are few in marshes, on low plains, dry moors, exposed downs, and places destitute of woods or other shelter from sun and wind. Apparent exceptions to the favorable or unfavorable effects of any of these conditions, may usually be explained by excess in some other and counteracting one. Thus, the salt spray and violent winds of the western shores are highly unfavorable to the growth of most Ferns, though otherwise their humidity of atmosphere would be favorable ; and accordingly whilst the exposed shores and cliffs may be almost without Ferns, caves and sheltered ravines in their immediate vicinity may be numerously tenanted. For the most part, however, even situations that are unsuitable to the majority of Ferns, have their own appropriate species. Thus, *Asplenium marinum* flourishes on cliffs exposed to the sea ; *Pteris aquilina* and *Blechnum* often grow on the unsheltered heaths, in places open to sun and wind ; *Asplenium rutamuraria* and *As. Adiantum nigrum* live in the crevices of dry walls and rocks. There are, however, no aquatic Ferns, and scarcely any of the Polypodiaceæ, that can be designated marsh plants ; unless that occupant of swampy bogs, *Aspidium thelypteris*, be called a tenant of the marshes. A light friable soil, and more especially that formed by the decay of tree leaves, mosses, or other vegetables, is suitable to the roots of most Ferns ; but some delight in limestone soils, as *Grammitis ceterach*, *Polypodium calcareum*, and *Cistopteris fragilis* ; whilst the *Asplenium septentrionale* and *Woodsia ilvensis* seem to affect the basaltic trap and the harder primary rocks. The operations of human industry have greatly interfered with the natural distribution of Ferns in this country. They have been banished from our roads, corn fields, meadows, and artificial pastures ; and the cutting of peat, and burning of heath and furze, often check the growth of species fitted to thrive in places where these operations are performed. On the other hand, our hedge-banks, loose stone fences, old buildings, and neglected quarries, frequently become artificial fernetums, by affording suitable habitats for several species. These remarks on the distribution of Polypodiaceæ will be understood to apply to the island of Great Britain only. Ireland has twenty-six species of dorsiferous Ferns ; but of their range and distribution within that island little is known. All those of Ireland are natives also of England.*

* A very interesting and extended paper upon the distribution of our Ferns, by Mr. Watson, will be found in " The Transactions of the Edinburgh Botanical Society." Part II. The above remarks, together with others of a corresponding character, inserted at other places, were kindly contributed to the Author, in writing, some time since. They may not agree, therefore, in precise words with the Edinburgh paper.

VIRTUES.—The uses of the Ferns are not very conspicuous. Their bitter principle renders them unpalatable to all creatures. Neither men nor brutes employ any species as an article of food, unless driven by the necessity of hunger;* and even the little insects that infest herbaria refuse to prey upon them. They are not, however, wholly useless, either in medicine or the arts. Their nauseous taste renders them efficacious in expelling intestinal worms; some of them have been used as a substitute for hops in brewing, and with better success than most other plants, on account of the tannin and gallic acid they contain, precipitating the feculent matter in the wort. The same constituent principles renders them also serviceable in preparing kid and other light leathers, and when burnt they yield much comparatively pure potass. The dried fronds of the common brakes are valuable to pack fruit in, and as they retain moisture less, are much better than straw to shield garden plants from frost. Except for these uses, the Ferns have been but little employed, unless, indeed, for those purposes to which most plants when dry are available, namely, for thatch, for fodder, and for fuel.†

HYMENOPHYLLACEÆ.

(Containing the Genera *Hymenophyllum* and *Trichomanes*.)

TRICHOMANOIDEÆ, *Kaulf.*;—FILICES DESCISCENTES, *Spreng.*;—PART OF GYRATEÆ, ANNULATEÆ, POLYPODIACEÆ, GLEICHENIACEÆ, FILICES VERÆ, HYMENOPHYLLEÆ, &c. of *Authors*.

STRUCTURE.—The plants contained in this order long maintained a situation among the dorsal Ferns, though improperly, because their fruit is not dorsal but marginal, growing in a distinct and differently organised receptacle. The annulus corresponds in its functions, jointed appearance, and elasticity, to those of the last order, except that instead of its being a continuation of the foot-stalk of the theca, it is placed obliquely or transversely, and of consequence the theca bursts vertically. For this reason, *Hymenophyllum* and *Trichomanes* form an order separate from the Polypodiaceæ. Besides the difference in the fruit, the texture of the leaves is much more cellular than in the last order. The stem of each native genus is quite smooth and round, and contains but one bundle of spiral vessels; this is solid and forms an axis. The thecæ arise from the veins

* Professor Henslow was kind enough to point out to me some time since that I had forgotten the circumstance of the New Zealanders living mainly upon Fern roots. It is true that they do so; still Ferns are a sorry food, and now that the colonists have taught the natives the art of cultivation, Fern roots are becoming less and less an article of consumption. That hunger alone induced the islanders to use these roots as food, may be inferred from the circumstance, that they were ready enough to work for the first settlers merely to be supplied with the commonest European grain or pulse, though the Fern grew abundantly on every side, and might have been procured and prepared comparatively without labor or expense.

† For the proper and modern culture of the Ferns, see the Appendix.

still more evidently than even in the preceding order, as the receptacle is in the place of a lobe of the leaf. The laminæ of the lobe contracted form the valves of the receptacle, and its vein or nerve exists as a central column, covered with fructification; in *Hymenophyllum* terminated thereby; in *Trichomanes* the vein is prolonged much beyond the thecæ. The root of *Trichomanes* is thick, black, and very hairy; that of *Hymenophyllum* very long, creeping, and matted together.

DISTRIBUTION.—“*Hymenophyllaceæ*, more impatient of drought than many of the *Polypodiaceæ*, delight in shaded situations, where they are bedewed by frequent mists and the spray of waterfalls; growing on damp rocks and stones, half buried amongst the Mosses that accompany them; and like the Mosses rapidly shrinking or expanding with variations of moisture. Two of the three native species are rare in Britain, especially the *Trichomanes*, which is peculiar to Yorkshire, if it be not now extinct in England. Another, *Hymenophyllum Tunbridgense*, is found in various counties of England and Scotland, varying from Devon and Sussex to the river Clyde in Scotland. The much more abundant *H. Wilsoni*, was so long confounded with the former species, that it has been supposed a rarer plant than it really is, for the greater number of localities assigned for *H. Tunbridgense* belong to *H. Wilsoni*. This latter species ranges from Cornwall, northward, to Sutherland, and rises on the mountains of Wales nearly to 2000 feet, and not improbably more.”—*Mr. Watson's MS.*

OSMUNDACEÆ.

(Contains only the genus *Osmunda*.)

OSMUNDACEÆ, *Br., Kaulf., Lind., Hook., Agard.*;—OSMUNDEÆ, *Spreng.*;—SCHISMATOPERIDES, *Willd.*;—SPURIE GYRATEÆ, *Swz.*;—RIMATEÆ, *Mohr.*;—ACROGYRATEÆ, *Bernh.*;—EXANNULATEÆ of *Modern Authors.*

STRUCTURE.—This order varies very little from the two former, except in the fructification. The rachis is similar in veneration and structure—the mixed vessels are arranged in the indigenous species in part of a circle like the letter ©. The cuticle of the stem is void of stomata, but on the under surface of the foliaceous part of the frond they are round, abundant, and very large. In the fruit, a great difference exists between this and the Annulate Ferns. The thecæ are transparent and wrinkled, but not ringed; they are not torn asunder irregularly, but open at a distinct longitudinal fissure, and are thus shown to be regularly two-valved. They are not fixed at the back of a leafy frond, but densely clustered at the top into a large erect raceme. The leaflets of the upper part of the frond are also seen partly changed into thecæ of similar character, thus proving the origin of the whole. (See *Osmunda*.)

DISTRIBUTION.—"Here the distribution of an order is that of a single species only, the *Osmunda regalis*, which prevails chiefly in the south-west of England, and perhaps the south-west of the Highland tract; being very frequent on the coasts of Cornwall, and occurring in plenty about some of the lochs of Argyleshire, and on adjacent islands. The range of the *Osmunda* extends the whole length and breadth of Britain; but whether it be found on the islets north of Sutherland, or west of Islay, is yet unknown. Still, there are considerable tracts along the eastern side of England and Scotland, where it is either extremely scarce or wholly wanting. The *Osmunda* differs from most of the dorsiferous Ferns in being adapted to thrive in marshy places, and to flourish on the sea shores, sometimes only just above high water line. Notwithstanding that it endures the boreal climate of Sutherland, (a county in which the specimens are said to be small,) the *Osmunda* is perhaps never seen much above the sea level in England, although the general vegetation of the coast in Sutherland approximates closely to what is observed at 1000 or 1200 feet of elevation in England. In respect to frequency of occurrence, the *Osmunda* may rank nearly on the level of its allies, the *Botrychium* and *Ophioglossum*, though rather less frequent than either of these."—*Mr. Watson's MS.*

OPHIOGLOSSACEÆ.

(Contains *Botrychium* and *Ophioglossum*.)

OPHIOGLOSSACEÆ, *Br., Lind., &c.*;—OPHIOGLOSSEÆ, *Spreng.*;—FILICES, *Linn., Smith, Hook., &c.*;—STACHIOPTERIDES, *Willd.*;—BIVALVA, *Hoffm.*; VALVATÆ, *Web., Mohr.*;—AGYRATEÆ, *Swz.*

STRUCTURE.—The plants contained in this order are very near in general structure to that of the last, yet in some important particulars they differ very materially. Their root is smooth, fibrous and yellow, not creeping nor hairy; and gives rise to one or at most two fronds only, which issue from the ground with a straight and not circinate vernation. The frond half way up divides into a leafy expansion. The thecæ are sessile, opaque, ringless, smooth, collected into a simple or compound spike, and are supposed to arise, as in the last-described order, from the leaf itself. The thecæ open by a regular transverse fissure, emitting smooth, yellow, very minute seeds; those of *Botrychium* in twos or threes. The roots of both genera are perennial, the stems herbaceous and hollow. The stem of *Botrychium* containing its ducts in two bundles near the centre; that of *Ophioglossum* in from five to seven bundles, seated between two cylindrical cuticles, and by their pressure forcing the inner one into a tortuous form.

DISTRIBUTION.—"The two plants comprehended in this small order differ from most of the Polypodiaceæ, by growing chiefly in more open situations; their upright habit perhaps rendering them less adapted to banks and rocks.

Both range widely through Britain, and are about equally frequent, holding an intermediate place between the rare and the common plants. The *Ophioglossum* prevails chiefly in England, decreasing in frequency northward. The *Botrychium*, on the contrary, is abundant on the hills and moors of the north, and becomes a rare plant in the south, and especially in the south-east of England. *Ophioglossum* ranges from the south of England, as Devon and Sussex, northward at least to Moray; and, if we may rely upon Barry's 'History of Orkney,' to those islands also. The *Botrychium* is scarce on the south side of the Thames and Bristol Channel, but is stated to grow in North Devon and in Hampshire, though not introduced into the Flora of the former county. The stations in the south of England of course indicate that both species will grow at a low elevation, and remote from the mountain tract, even in the warmest part of Britain; and both also thrive amidst the mountains in the north of England and Scotland; but the *Botrychium* probably rises to a much greater height on the hills, as it occurs on the Breadalbane mountains, near Killin, at the estimated height of 1000 yards, whilst no very high elevation for the *Ophioglossum* appears on record. Of the two, the *Botrychium* is the least frequent, or seems to be so on account of its more boreal and Alpine tastes."—*Mr. Watson's MS.*

ISOETACEÆ.

(Containing *Isoetes* only.)

LYCOPODIACEÆ, *Lind., Decan., Brongn.*;—MARSILEACEÆ, *Hook.*;—MISCELLANEÆ, PART OF RHIZOSPERMÆ, RHIZOPTERIDES, HYDROPTERIDES, &c.

The genus *Isoetes* has in all arrangements of British plants been associated with *Pilularia*, on account of their both being water plants, both having round and filiform leaves, and bearing two kinds of grains or capsules; but, except in these particulars, they are totally different from each other. The roots of *Isoetes* are tufted, composed of round, smooth, branched fibres; its leaves grow from a crown, and consist of four hollow tubes, united together; but so brittle are they, that the cells are often broken into each other by the pressure used in drying the plants, and therefore the leaf generally appears a single tube, divided into cells by transverse dissepiments. It is so swelled at the base that the joint or cell next the root becomes a receptacle for the fruit, which being of two kinds, as in *Pilularia*, are considered analogous to them; viz. pollen and spores respectively, the former in fine powdery grains in the inner leaves—the real spores or seeds being confined to those on the outside of the plant. These larger globules are not single round spores, but each is composed of three or four spores, joined together. Their junction shows at the apex three radiating lines, which were for a long time considered as the hilum of the seed. The attachment and arrangement of the globules within the receptacle are very beautiful and remarkable. The

theca when cut across exhibits a number of transverse bars, to which the spores are attached by little foot-stalks, there being four on each bar, set crosswise with each other. The leaves are said to have stomata, and to be circinate in veneration, but neither of these is the case. Being a submersed water plant, of course it is without stomata, and Martius expressly says, " veneration not circinate, but only a little bent." An observation confirmed to me by four or five botanists of eminence.

DISTRIBUTION.—" The single species of this order, a submerged aquatic, can of course grow only where there is a suitable home in the waters. Probably it may require also that the water be of low temperature in summer, since most of its habitats are the Highland lakes ; although, according to Mr. Griffith, it extends southward into Shropshire. There are very few other English and Welsh counties that produce it, and these few are all mountainous ; namely, Caernarvonshire, Denbighshire, Cumberland, and Northumberland. In most counties north of the Firths of Forth and Clyde, we may find it in the lakes, some of them being situate 600 yards or upwards above the sea."—*Mr. Watson's MS.*

MARSILEACEÆ.

(Including only *Pilularia*.)

PART OF THE MARSILEACEÆ, *Br.*, *Brongn.*, *Decan.*, *Hook.*, *Grev.* ;—HYDROPTERIDES, *Willd.* ;—RHIZOSPERMÆ, *Roth* ;—RHIZOPTERIDES, *Mart.* ;—RADICALIA, *Hoffm.* ;—RHIZOCARPÆ, *Batsch*.

The stem of *Pilularia*, which is the only English genus of this order, is creeping, and set at intervals with leaves, roots, and fruit. The leaves or petioles, as some call them, are curled up in veneration, as in the Polypodiaceæ, have stomata upon their cuticle, and a cross section of them shows that they are divided longitudinally into various cells, separated from each other by septa radiating from the centre, and forming by their union a kind of axis, composed of dotted ducts or spiral vessels. Thus the structure of the leaf although without a central cavity, is in a great degree analogous to that of the stem of the Equisetaceæ, and also to the leaf of *Isoetes*. The roots and stems are similarly constructed.

The thecæ are round, coriaceous, brown, hairy, and divided into four cells. They contain globules of two kinds ; the first small round grains, said to be pollen, but which others consider as abortive capsules. These occupy principally the upper part of each theca ; in the lower are found much larger grains, which are oval, rather pointed, contracted in the middle, and at their apex have a conical projection. These are true spores, and as well as the former are contained in membranous bags. Since writing the above, a very valuable paper by Mr. Valentine, upon " The Germination and General Structure of *Pilularia*," was read before the Linnæan Society, and is to be found in their " Transactions,"

for 1841, page 483, from which I extract the following remarks:—"The first external sign of germination is the appearance of four cells projecting through the apex. The enlarged cellular mass then distends the conical projection, and at length appears with four of its cells projecting beyond the general mass, and compressed into a quadrangular form. These projecting cells soon harden, and acquire a reddish brown hue—soon after which little fibrillæ or rootlets begin to shoot from one side. They are simply articulated tubes, or elongated cells applied end to end, each produced from one of the cells of the germ. The germ now gradually points in two places, which are by no means fixed, but occur in various situations, according to the position of the sporule in respect to light. These two points gradually lengthen, and if dissected, each will be found to consist of a closed sheath, containing in one instance the leaf, in the other the root, in the form of a conical process like a finger in a glove. Besides this sheath which embraces the upper part of the root, there is an exceedingly delicate expansion which closely embraces the extremity of the root like a cap. After the leaf has grown to be many times the length of the sporule, or about two lines long, another leaf grows from the germ close to the first, to which it is in all respects similar; and then a bud begins to be developed from some indefinite part of the germ, and like the leaves and root form within a sheath. This bud is covered by a peculiar kind of jointed hairs, whose attachments are lateral at a short distance from their bases."

Mr. Valentine then shows the origin and progress of the sporules within the theca, making afterwards the following pointed remark:—"This account of *Pilularia* shows that it is incorrect to say of *Aerogens* that germination takes place at no fixed point, but upon any part of the surface of the spores; for it is quite certain in this instance that germination invariably takes place at a fixed spot, which may be pointed out before germination has commenced." This view exactly accords with some experiments I have instituted, and with the remarks of Dr. Lloyd, who read a most interesting paper upon this subject, before the British Association, in 1836. The statement in "*Lindley's Nat. Sys.*" ed. 2, page 416, that the theca never produces but one plant, is not correct, for if the theca be broken and its contents scattered, numerous of the spores will germinate.

DISTRIBUTION.—"The solitary species of this order belonging to Britain is widely distributed, being found in Devon and Sussex, and extending at intervals from those counties northward to Sutherland. It is either frequently overlooked, or is otherwise a scarce plant in the south of England, increasing in plenty in the northern counties; probably the drier climate of the south-east of England, causing the small pools to diminish much or entirely to dry up in summer, is adverse to the existence of a plant that is adapted to grow within the shallow margins of ponds and lakes. Of its range in altitude, little seems to have been yet ascertained."—*Mr. Watson's MS.*

LYCOPODIACEÆ.

(Comprises only Lycopodium.)

LYCOPODIACEÆ, *Br., Decan., Hook., Lindl., Burn.*;—LYCOPODINEÆ, *Swz.*;
 LYCOPODEÆ, *Spreng.*;—BIVALVIA, *Hoffm.*;—VALVATÆ, *Web., Mohr.*;—
 STACHIOPTERIDES, *Willd.*

STRUCTURE.—The Lycopodia resemble the Mosses in habit, the Ferns in vascular structure and foliaceous texture, and the Marsileaceæ in fruit. The stems are rigid, leafy throughout their whole extent, not subterranean, but upright or trailing along the ground, frequently to the distance of many feet, and throwing out short, stiff, smooth radicles wherever they touch the soil. A transverse section of the root shows the longitudinal ducts to be compressed into an axis. In the stem they are arranged, as in the Ferns, into various cylindrical bundles, the centre of which is filled with a cellular tissue, looser than the remaining part; the ducts in this order are flat spiral bands, minutely dotted. The stomatæ on the cuticle of the leaves are very abundant, the cuticle itself being reticulated, not as in the Ferns, but into regular four-sided meshes.

The thecæ are sessile, in the axils of the leaves, of two kinds, one two-celled, opening at a longitudinal fissure, containing very fine smooth resinous grains, which are supposed by most botanists to be pollen, but by Professor Lindley to be abortive thecæ. The other kind is three or four-valved, opening at a transverse line, and contains from three to five round, slightly tuberculated grains, many times larger than the preceding. That these are the true spores of the plant is evident from their germination, and Willdenow says that he has seen the smaller grains grow also; if so, they likewise must be spores, but as they are so very different in size and appearance, it is supposed that some mistake has arisen. Mr. Salisbury, in vol. 12 of the “*Linnæan Transactions*,” describes the germination of one species, which presents strange anomalies, throwing out a radicle and plumelet, in a manner similar to the monocotyledonous plants, and yet appearing immediately afterwards with two leaves, which he represents as cotyledons.

DISTRIBUTION.—“The Lycopodiaceæ for the most part affect exposed situations on open heaths, or the summits and sloping acclivities of mountains; although without altogether shunning more shady homes. *L. selaginoides* inclines to the most humid situations, growing frequently in the crevices of dripping rocks, about waterfalls, and in swampy ground, where water oozes from the sides of hills. *L. inundatum* occurs in analogous situations, on lower or more southern heaths and commons. The other four species choose drier dwelling places; *L. selago* and *alpinum* bearing the rude exposure of the mountain summits; and *L. annotinum* and *clavatum* being more frequent on the sloping acclivities than on the summits; but none of these four are exclusively restricted to such situations. The order is pre-eminently boreal and alpine, only one species,

L. inundatum, being at all frequent in the low counties of the south-east of England, and decreasing in abundance towards the northern and hilly counties; whilst all the rest prevail in the Scottish Highlands, and decrease in frequency in a southward course."—*Mr. Watson's MS.*

VIRTUES.—Seldom used in medicine, where safer drugs are attainable. The Orkney Islanders use *L. selago* and *clavatum* as a cattle remedy; it is said to cure sheep of vermin and of different entaneous disorders; in the human subject it is an emetic and purgative. The pollen is highly inflammable, and was once imported in some abundance from Germany and Sweden to imitate lightning at the theatres, but latterly powdered rosin has been substituted. *Lycopodium clavatum* is said to be valuable in dyeing woollen cloths, and for making mats it is admirable; and the Poles make a decoction of its leaves as a remedy for the disorder called *Plica polonica*. The pollen is wetted with so much difficulty that when spread on the top of the water in a basin, a finger may be plunged to the bottom without becoming wet.

EQUISETACEÆ.

(Comprises only *Equisetum*.)

EQUISETACEÆ, *Decan., Ag., Kaulf., Lind., Hook., Grev., Brong.*;—GONOPTERIDES, *Willd.*;—PART OF FILICES, MISCELLANÆÆ, &c., of *Authors*.

CLASS.—These plants differ widely from all those hitherto described, and certainly approach very much nearer to flowering plants than Ferns themselves. In fact their relation to the Coniferæ is so strong, both in external and internal structure, and their analogy with some other orders so apparent, that I continue them among the Fern allies more in accordance with the opinion of others than my own. Yet I cannot consider them Dicotyledonous plants, as Professor Lindley has done, because their germination is essentially like that of the Cellulares, and their reproductive organs have no analogy with those of flowering plants. In fact, *Equisetum* forms a perfectly distinct order, and cannot be allied with any other.

STRUCTURE.—The stems, which are partly beneath and partly above the surface of the ground, are when young filled with very loose cellular tissue: the moisture of this soon drying up, they become hollow. They are set at intervals with joints, attended by toothed sheaths, are regularly channelled or striated, rigid, and covered with fine particles of silex, particularly at the ridges of the striæ. The depressed part of each channel has two longitudinal rows of minute holes or open pores, very different from the usual stomatæ, and much resembling the pores which Dr. Mohl, of Munich, represents as occurring in the woody tissue of the Coniferæ, now a well-known characteristic of that

order, and which it will be recollected there, as here, are arranged in longitudinal lines. A transverse section of the stem shows that between the outer and inner cuticle is a circle (and in *Equisetum fluviatile* two circles, alternating with each other), of tubes, distinct from each other, but terminating at every joint of the stem; fresh tubes of a similar character being found at every other joint. Around these tubes, and especially towards the outside of the whole stem, and seated immediately under the cuticle, are the sap vessels of the plant, which appear flat, spiral, perforated ducts, as in the other orders. The inner surface of the stem is frequently more silicious than the outer, forming a very beautiful object even to the naked eye. So abundant is this deposit in *Equisetum hyemale*, that, after the vegetable matter has been removed by maceration, the silix has been sufficiently abundant to retain the form of the plant. M. John, of Berlin, states that the stems contain full 13 per cent. of silica. The following interesting particulars of the silicious cuticle of *Equisetum* is given in Dr. Greville's excellent "*Flora Edinensis*," p. 214. "On subjecting a portion of the cuticle to the analysis of polarized light under a high magnifying power, Dr. Brewster detected a beautiful arrangement of the siliceous particles, which are distributed in two lines parallel to the axis of the stem, and extending over the whole surface. The greater number of the particles form simple straight lines, but the rest are grouped into oval forms, connected together like the jewels of a necklace, by a chain of particles forming a sort of curvi-linear quadrangle; these rows of oval combinations being arranged in pairs. Many of those particles which form the straight lines do not exceed the 500th part of an inch in diameter. Dr. Brewster also observed the remarkable fact, that each particle has a regular axis of double refraction. In the straw and chaff of wheat, barley, oats, and rye, he noticed analogous phenomena, but the particles were arranged in a different manner, and "displayed figures of singular beauty." From these data, the learned Doctor concludes, "that the crystalline portions of silix, and other earths which are found in vegetable films, are not foreign substances of accidental occurrence, but are integral parts of the plant itself, and probably perform some important function in the processes of vegetable life."—*Brewster, MS.*

REPRODUCTION.—The reproductive organs are borne in a terminal spike or catkin, composed of hexagonal or octagonal shields; from each side of which depend hollow, scale-like follicles, opening inwardly, and emitting green, ovate spores, to each of which is attached four club-shaped filaments. When the spores are immature, the filaments are twisted tightly round them, but when ripe they become exceedingly elastic and hygrometrical, so much so, that the irritability occasioned by a change of temperature or moisture, causes the spores first to burst the thecæ which bear them, and afterwards to scatter themselves to a considerable distance.

A valuable paper upon the formation of the spores, by Mr. Henderson, was read before the Linnæan Society, in June 1840; and will be found in their "*Transactions*," vol. xviii. p. 567. The experiments of this gentleman exactly

confirm the above view, and give also the origin of the minute grains found upon the elaters or elastic filaments; from this paper, it appears, that when the integument of the spore shows the spiral lines, which it will afterwards break into the elaters, it contains a greenish colored fluid, mixed with some minute granules. The spores then becomes darker, the granules increase, the liquid is absorbed, leaving the granules which it contained sticking in masses to the spores, and to the separated portions of the integument. It is these masses of granules, when found adhering to the filaments in the ripened state of the spore, that have been taken for pollen grains, and such the observations of the author prove them to be. He says, (p. 571) "On comparing these granules with those contained in the unopened anthers of flowering plants, they appear to me to be in every respect identical; in the thecæ they seem to occupy a similar place with those in the cells of the anthers, and they decrease in like manner during the progress to maturity of the pollen grain and of the spore."

DISTRIBUTION.—"Nearly equal in number with the Lycopodiaceæ, the plants included under this order contrast against them in their distribution. Whilst the former are described in general terms as plants of the moors and mountains, the Equisetaceæ belong more especially to marshes, fields, and woods. Both orders, or genera, are alike widely distributed through Britain; but whilst the Lycopodiums prevail high on the mountains, the Equisetums are more abundantly bestowed upon the lower grounds, or the plains and valleys; the latter also evince more of a maritime, or even littoral tendency than the Lycopodiums; being often seen in abundance on and near the sea shore."—*Mr. Watson's MS.*

USES.—They are harmless to cattle, but refused on account of their husky, rigid texture. They are useless as medicines, but employed in the arts; their siliceous cuticle rendering them valuable as polishing substances for marble, wood, ivory, and even metals.

GENERA.

THE first order of the Ferns is conveniently divided into the sub-orders *Nudæ* and *Indusiataæ*, according as the sori are naked or covered with an indusium. The arrangement of the species into Genera depends upon the shape and position of the sori, together with the nature, the adherence, and the manner of opening of the indusium when there is one. In the other orders the same principles are adopted, as far as their structure will admit.

As the orders of the Fern Allies contain each but one British genus, the characters of that genus are but a recapitulation of those of the order itself.

NUDÆ.

Sori linear or oblong, scattered	GRAMMITIS.
Sori round, scattered	POLYPODIUM.

INDUSIATÆ.

Indusium distinct from the Frond.

Sorus round. Indusium cleft into capillary segments ..	WOODSIA.
Sorus round. Indusium cucullate or bladder-shaped ..	CISTOPTERIS.
Sorus round. Indusium peltate, either round or reniform	ASPIDIUM.
Sorus linear or oblong, transverse, solitary. Indusium from lateral veins, opening towards the midrib	} ASPLENIUM.
Sorus linear, transverse, in twin masses. Indusia folding over each other, and opening outwardly.....	
Sorus linear, longitudinal, close to the midrib. Indusia opening outwardly	} BLECHNUM.

Cover part of the Frond itself reflexed.

Sorus linear, continued around the margin of the frond	PTERIS.
Sorus on transverse veins, near to the margin	CRYPTOGRAMMA.
Sorus in distinct spots, attached to the cover itself.....	ADIANTUM.

Thecæ in pitcher-shaped, one-valved receptacles	TRICHOMANES.
Thecæ in compressed, two-valved receptacles	HYMENOPHYLLUM.

Thecæ petiolcd, reticulated, terminating a leafy frond ..	OSMUNDA.
Thecæ sessile, smooth, in a separate compound spike ..	BOTRYCHIUM.
Thecæ sessile, smooth, in a separate simple spike	OPHIOGLOSSUM.

Thecæ attached to the root, free, and indehiscent	PILULARIA.
Thecæ imbedded in the base of the leaves	ISOETES.
Thecæ of two kinds, axillary in a leafy spike, or stem..	LYCOPodium.
Thecæ of one kind, in catkins, terminating a leafless stem	EQUISETUM.

GRAMMITIS. Swz. GRAMMITIS.

(γρᾰμμῖτις, a line; alluding to the linear fructification.)

A shows the under surface of the frond of *Grammitis Ceterach*. B, the veins according to Newman. C, the veins according to Presle. D, position of the fruit. E, an unopened theca. F, a theca scattering its spores. G, the spores. H, a scale. I, cuticle and stomata. J, transverse section of the rachis.

GRAMMITIS CETERACH.

SCALY GRAMMITIS. SCALY HART'S-TONGUE. MILTWAST.

(Plate 1, fig. 1.)

CHA.—Frond pinnate or pinnatifid, scaly beneath. Lobes alternate, confluent, blunt, entire.

SYN.—*Grammitis ceterach*, Swz., Hook., Mack.—*Asplenium* or *ceterach*, Ger., Ptm., Ray.—*Asplenium ceterach*, Linn., Huds., Sibth., Lightf., Bolt., With., Spreng.—*Gymnopteris ceterach*, Bernh.—*Scolopendrium ceterach*, Roth. Galp., Smith.—*Ceterach officinarum*, Willd. Decan. Neum.

FIG.—E.B. 1244.—Park. 1046, f. 1—Ger. 978.—Lobel, 807.—Bolt. 12 (bad).

DES.—Root perennial, fibrous, black, tufted. Fronds many from the same root, herbaceous, 3 to 6 inches high, blunt, of a thick texture, dark green above, covered with brown scales beneath. Lobes confluent at their base, round, entire, alternate at the lower part of the frond, flat only when young, afterwards curved inwards towards the main rib, thereby exposing more the fructification. Thecae all the summer, at first concealed by the scales, afterwards bursting through them in oblong, transverse masses, without covers, but surrounded by very delicate, white, membranous scales.

SIT.—On rocks, old walls, &c., chiefly in the South of England.

HAB.—ENG.: Near Lancaster, Mr. W. Wilson. Common about Settle, Yorkshire, Mr. J. Tatham. On limestone rocks in Lath-kill-dale, Derby, Mr. J. E. Bowman. On a wall at Newton, near Melbourne, Derbys., Rev. A. Bloom. Dovedale, Derby, Mr. T. S. Scholes. Walls at Ludlow, about the quarries, Salop, Mr. J. S. Bayly. Old wall near Cowley, Oxon, Mr. Baxter. Wall at Tocknells, near Painswick, Glouc., Mr. Merrick. Martock, Somers., Mr. T. H. Cooper. Stapleton Quarries, near Bristol, Mr. Anderson. Cheddar, Mr. W. C. Trevelyan. Malvern Abbey, Mr. W. Christy. Bath, Mr. C. C. Babington. On the tower of Old Alresford Church, Hants, Mr. Forder. Walls at Winchester, chiefly to the E. and N.E. of the City, Mr. W. Pamplin. Topsham and elsewhere in Devon, Mr. Kingston. On the bridge over the Tamar, in the road

from Callington to Tavistock, *Jones's Tour*.—WAL.: Denbighshire (rare), *Mr. J. E. Bowman*. Walls of a ruin at Treborth, near Bangor, *Mr. W. Wilson*.—IRE.: Ruins of Saggard Church, *Mr. Ketly*. Walls near Cork, also near Kilkenny, and in county Clare, *Mr. Mackay*. Cave-hill, *Mr. Templeton*. Headford, Galway, *Mr. Shuttleworth*. *Abundant at Boringlough, near Dublin.*

GEO.—Holland, Spain, France, Switzerland, Nassau, Jena, Leipsic, and other parts of Germany, the Tyrol, Sicily, and the Canary Islands.

Walls at Whitechurch, — Ellen cone —

POLYPODIUM. Linn. POLYPODY.

(πολύς many, and ποὺς, ποδός, a foot; from its numerous roots.)

A, pinnule of natural size of *Polypodium vulgare*. B, magnified section of a sorus. C, front view of ditto. D, longitudinal section of rachis. G, transverse ditto. E, spiral perforated duct. F, vernal, rhizoma and rootlets. H, I, theca and spore.

*Sprengel enumerates no less than 250 species of this genus; all of them are herbaceous, some a few inches only, and others several feet in height. Inhabitants of most parts of the world, particularly of the islands within the Tropics; several are found on the continent of America, and a few are confined to China. Only four species are British.**

1.—POLYPODIUM VULGARE.

COMMON POLYPODY. POLYPODY OF THE OAK. WALL FERN.

(Plate 1, fig. 2.)

CHA.—Fronde pinnatifid, lanceolate. Lobes oblong, obtuse, somewhat serrated. Rachis smooth. Root hairy.

SYN.—*Polypodium vulgare*, *Tourn.*, *Ger.*, *Park.*, *Ray*, *Linn.*, *Huds.*, *Lightf.*, *Plum.*, *Swz.*, *Spreng.*, *With.*, *Smith*, *Hook.*, *Mack.*, *Gray*, &c.

FIG.—*E.B.* 1149.—*Flo. Dan.* 1060.—*Woodv. Med. Bot. supp.* 271.—*Ger.* 467.—*Bolt.* 18.—*Plu. Fil.* t. A f. 2.

DES.—Root, or rather rhizoma, creeping horizontally, covered with scales, and numerous stout, branched, hairy fibres. Rachis quite smooth, yellow, void of lobes half way up. Frond from 6 to 12 inches high, lanceolate, scarcely contracting below. Lobes oblong, obtuse, and slightly serrated, sometimes wanting the serratures, at others acuminate, while occasionally they are found very much cut and divided. Sori naked, yellow, large,

* The number of species in a genus is always subject to variation, particularly in one so extensive as Polypody, as newly-discovered plants are always adding to the number, while different classification often divides one genus into many.

prominent, and arranged in straight lines equally distant from the margin and the midrib of the lobe; each sorus terminating one of the branches of a transverse vein. The plant is perennial and the fruit found throughout the summer.

- β . (*P. cambricum*, Linn.) Frond ovate; pinnule ovate, and deeply cleft.
- γ . (*sinuatum*.) Frond ovate, or triangular; pinnules proliferous.
- δ . (*serratum*.) Pinnules distinctly and often doubly serrated.
- ϵ . (*acutum*.) Pinnules pointed; fronds long; both narrow.
- θ . (*bifidum*.) Pinnules cleft at the point.

Mr. Mackay remarks, in his "*Flora Hibernica*," that the Irish plant is somewhat different from the *Polyp. Cambricum* of Linnæus. It is in fact our variety γ , which is the same as the *Pol. Virginianum* of Pursh, and intermediate between the usual state of the plant and the *Cambricum*; it bears fruit copiously, whereas the real *Cambricum* is usually without fruit, both in its wild and cultivated state. We might expect this, indeed, from the feather-like appearance of the plant, and the dilation of its lobes, a too great expansion of leaf being here as elsewhere detrimental to the production of fruit. The foregoing observation was made in distinct reference to a frond, of which C in the annexed cut is an exact representation; but a plant still more nearly approaching Linnæus's *Pol. Cambricum* is in Sir J. Smith's herbarium, marked as from Ireland. A pinnule is represented in the Fig. D, copied from the original specimen, an admirable figure of the whole frond, as well as of the *Cambricum* is in "*Newman's Ferns*," p. 22. One pinnule of the latter is represented at B, and a whole frond of it, from my herbarium, at A. The other varieties are shown at E F and G.

VIR.—Although still retained in the pharmacopœias, it is scarcely, if at all, used in medicine at the present day. It is feebly astringent, of a bitter and nauseous taste, and has been considered efficacious in catarrhal disorders, and against worms, in doses of from one to two drams of the dried root.

HAB.—The common states of the plant (α and δ .) are generally distributed over the United Kingdom, on trees, walls, banks, and rocks.— β . On the rocks in some parts of North Wales, but without fruit.—Braid Hall, near Edinburgh, *Mr. Brown*. At Chepstow, Monm., *Sir J. E. Smith*.— γ . Woods at Dulwich (1835), *Mr. Saunders and Mr. W. Pamplin*. South Isles of Arran (1806), *Mr. Mackay*. In the Dargle, county of Wicklow, *Miss Fitton*. Innisfallen Island, Killarney, *Mr. Kelly*.—South side of King's Park, Edinburgh, *Mr. Brown*. ϵ . Rocks in North Wales, *With*. Meadows near Maldon, and other meadows near Ewell, Surrey, *Mr. J. Bevis*. Cobham Park, Kent.—*G. P.*

GEO.—Found in most of the middle parts of Europe and North America.

2.—POLYPODIUM PHEGOPTERIS.

BEECH FERN. WOOD POLYPODY. SUN FERN.

(Plate 1, fig. 3.)

CHA.—Frond bipinnatifid. Lower pinnæ deflexed. Lobes obtuse, entire, hairy.

SYN.—Polypodium phegopteris of *Linn.*, *Willd.*, *Swz.*, *Spreng.*, *Huds.*, *Lightf.*, *Bolt.*, *With.*, *Smith*, *Hook.*, *Mack.*, *Newm.*—Polystichum phegopteris, *Roth.*—Polypodium latebrosum, *Gray*, *Salisb.*

FIG.—*E.B.* 2224.—*Bolt.*, 20 (*not good.*)—*Flo. Dan.* 1241.

DES.—Root perennial, hairy, slender, creeping horizontally. Frond triangular, herbaceous, erect, hairy, 6 to 12 inches high. Pinnæ opposite, very acute, adnate, the lower pair bent forwards, pendulous, and distant from the pair next above them. The lobes of all are obtuse, entire, and directed towards the point of the pinna, particularly the two lowest, which with those on the opposite pinna form a cross. The rachis is smooth, and without pinnæ on the lower half. Sori round, distinct, very small, brown, and seated around the margin of the lobes.

The pendulous character of the lower pinnæ, and the cruciform direction of their bases are most apparent in vigorous plants, and serve as characters which immediately distinguish this plant from its congeners.

SIT.—In moist woods and rocky dells, chiefly in mountainous countries.

HAB.—ENG.: Rocks at the foot of Cheviot, above Langley Ford, *Mr. Winch.* Cawsey Dean, Durham, *Mr. R. B. Bowman.* Around Keswick, Cumb., *Mr. H. C. Watson.* Wensley-dale, Yorks., *Mr. J. Ward.* Common about Settle, Yorks., *Mr. J. Tatham.* Prestwich Clough and Boghart Clough, Lancashire, *Mr. Merrick.* Egerton Moss, near Bolton, *Mr. W. Christy.* Rocks at the Belle Hag, one mile from Sheffield, *G. F.* Norwood, Surrey, and near Brentford, Middx., *Mr. J. Bevis.* Lidford Fall, Beckey Fall, Dartmoor, Devon, *Jones's tour.* Isle of Man, *Mr. E. Forbes.*—WAL.: Llanberris, first and second field towards Snowdon, *Mr. C. C. Babington.* Capel Curig, North Wales, *Mr. T. H. Cooper.* Frequent in Caern., not at any considerable elevation, *Mr. W. Wilson.*—SCO.: Grampians, Aberdeensh., Red Caird Hill, W. of Invernesshire up to 1150 yds. Forfarshire, Sutherland, Dumbarton, and other parts of the Highlands, *Mr. H. C. Watson.* Moray, and Rosshire, *Rev. G. Gordon.* Ben Lomond, *Professor Henslow.* Ruberslaw, Jedburgh, &c.—Campsie, near Glasgow, *Mr. T. H. Cooper.*—IRE.: Powerscourt Waterfall, (right-hand side,) *Mr. O. Kelly.* Waterfall above Lough Eske. Co. of Donegal, and at other places in the northern counties, *Mr. Mackay.*

GEO.—Throughout Germany, and indeed most European countries as far North as Lapland, but not in the South countries. Linnæus received specimens from Canada.

about the Greenside mine. Helvellyn, Cumberland 1849
G. F. & J. B. Corn

3.—POLYPODIUM DRYOPTERIS.

TENDER THREE-BRANCHED POLYPODY.

(Plate 1, fig. 4.)

CHA.—Frond tri-pinnate, tender. Branches drooping. Lobes obtuse, erenate. Sori distinct.

SYN.—*Polypodium dryopteris* of Linn., Willd., Swz., Ehrh., Huds., Bolt., Lightf., Hull., Galp., With., Newn., Smith, Hook., Grev., Mack.—*Polypodium Pulchellum*, Gray. *Polystichum Dryopteris*, Roth.

FIG.—*E.B.* 616 (*excellent*).—*Bolton*, 28 (*bad.*)—*Gerard*, 974.—*Park*, 1044.

DES.—Root perennial, creeping, black, slender, slightly hairy. Fronds herbaecous, scattered, tender, drooping, smooth, and of a light green color. The three branches (of which the middle may be considered a continuation of the stem) are bent backwards, and doubly pinnate. Lobes crenate, oblong, obtuse. Main stem 6 to 8 inches long below the branches, quite smooth, except at the very base. Sori nearly marginal, scattered, remaining perfectly distinct.

SIT.—Dry stony places, chiefly in mountainous countries of the north.

HAB.—ENG.: Rocks at the foot of the Cheviot, above Langley Ford, *Mr. Winch*. Wooded banks of the White Adder, between the Retreat and Elm Cottage, Berwickshire, *Dr. Johnston*. Durham, *Mr. R. B. Bowman*. Near Yoxhall Lodge, in Needwood Forest, Staffordsh., *Mr. C. C. Babington*. Cumberland, up to 500 yards of elevation, *Mr. H. C. Watson*. Higher part of the Tees, *Mr. J. Hogg*. Egerton Moor, near Bolton, *Mr. W. Christy*. Dean Church, Clough, near Bolton, *Mr. J. Martin*. Dry places near Lancashire, (sparingly), at Hill Cliff, Cheshire, and at Warrington, *Mr. Rylands*. Boghart Hole Clough, and Prestwich Clough, Lanc., *Mr. Merrick*. Rocks at the Belle Hag, Sheffield, *G. F. Froddesley Hill*, Salop, *Rev. W. Corbett*. N. side of Titterstone Clee Hill, Salop, *Mr. E. Lees*. Near Richmond, Yorks., *Mr. J. Ward*. Near Bristol, *Miss Worsley*.—WAL.: Craig Breidden, Montgomerysh., *Mr. J. E. Bowman*. Rhaiadr-y-Wenol-Twll Du, Caernarvonsh., *Mr. C. C. Babington*. Frequent in N. Wales, and observed near Twll Du at an elevation of 1000 feet and upwards, *Mr. W. Wilson*. Just leaving Llangollen, on a slate rock, *Mr. W. Wilson* and *Mr. Bowman*.—SCO.: Moray, Rossh., *Rev. G. Gordon*. Perthshire, Forfarshire, Aberdeenshire, *Mr. H. C. Watson*. Hawthorn Dean, near Edinburgh, *Mr. T. H. Cooper*.—IRE.: On the mountains of Mourne, Turk Mountain, Killarney, Mam-turk, Cunnamara, Tullamore Park, &c., *Mr. Mackay*.

GEO.—Throughout great part of Europe and North Asia.

4.—POLYPODIUM CALCAREUM.

RIGID THREE-BRANCHED POLYPODY. LIME POLYPODY.

(Plate 1, fig. 5.)

CHA.—Frond tri-pinnate, rigid. Branches upright. Lobes obtuse, deeply crenate.

SYN.—Polypodium calcareum, Swz., Willd., Smith, Hook., Pursh., Galp.—
Polypodium dryopteris, Bolt., Dicks.—Polypodium dryopteris β ,
With., 2nd. edit.—Polypodium Robertianum, Hoffm.—Nephrodium
dryopteris, Michx.

FIG.—E.B., 1525.—Bolt. 1.—Ger. 1135.

DES.—This is so similar to the last, that when dried they are scarcely to be distinguished, hence, the doubt of their claim as distinct species; but when growing, the eye will instantly see the difference between the two. The *P. calcareum* is known from its root being thicker and less creeping, its frond rather larger in size, much more rigid, quite upright, and of a dark green color; its lobes more deeply cut, and stem more sealy towards the base, and on the upper part sprinkled over with fine white minute hairs; its sori are browner and more numerous. The minute pubescence seen on this species is most observable on luxuriant and fresh plants. It is, I believe, a constant and decided character, in which opinion I am supported by Mr. Wilson and Mr. Babington, than whom few are better able to form a correct judgment. The latter gentleman observes, that the microscope shows every particle of this pubescence or meanness to be a minute, stalked gland.

Mr. Newman, in his beautifully-illustrated book on Ferns, blends this with the former species; yet his figures show strongly-marked differences. Both of us no doubt argue according to our respective means of observation; and after a very careful re-examination of both the species in my own and Sir J. Smith's herbarium, and also as growing in Kew Gardens, I see no reason to alter a single word in the above descriptive characters. Whether the size of the two be of moment is the only thing which appears of little certainty; but the rigid erect habit, and dark color are very characteristic, even without noticing the pubescence. Indeed, whatever doubt I may have of the distinctness of certain others of the Ferns, I cannot for an instant consider the present and former species identical; let it be observed, however, that in the case of these, as well as some other species, the same herbarium often contains but the commoner plant, which is the *P. dryopteris*, yet some of the specimens may be under the name of *P. calcareum*, and too often does it happen, that the receiver of a specimen, taking for granted that such is correctly named, makes it a guide for his own future judgment, and thus an error becomes perpetuated.

HAB.—Arncliffe and Gordale, Yorks., Mr. R. B. Bowman. Near Lancaster, Mr. Gibson. Sheddin Clough, three miles from Burnley, Lanc., Mr. Leyland. Common about Settle, Yorks., Mr. J. Tatham. Matlock Bath, Derbys., Dr. Howitt. Road-side under the Lover's Leap, near Buxton, Derbys., Mr. H. C. Watson. Cheddar Cliffs, Somers., Mr. W. Christy. Box Quarries, near Bath. Mr. Flower. Not found in either Scotland or Ireland. *Burnington Comb. Kennet. C. 26.*
Westminster near Cheddar 1850

GEO.—Recorded by Pursh and Michaux as occurring throughout North America from Canada to Pennsylvania, and no doubt this is correct, as the description of Pursh so exactly accords with our plant; though Swartz says that it is found in England only.

WOODSIA, Br. HAIR-FERN.

(Named in honor of Mr. J. Woods, an English Botanist.)

A, portion of a frond of *Woodsia ilvensis*, natural size. B, ditto enlarged. C, sorus longitudinally divided. D, indusium. E, one portion of ditto. F, scale. G, theca. H, spores.

Mr. Brown first separated from the *Polypodiums*, &c., this very distinct genus, which contains only two British and four foreign species, all very small plants, and natives of mountainous regions. The indusium, if such it can be called, is very singular and beautiful: it is attached under the mass of thecae—inclusing them at first in a bag, it then becomes split into numerous segments, which look like hairs interspersed with the thecae, and were so considered until Mr. Brown showed their true nature in “*Trans. Linn. Soc.*” vol. xi.

WOODSIA ILVENSIS.

OBLONG WOODSIA. HAIRY WOODSIA. DOWNY HAIR-FERN.

(Platc 1, fig. 6, A.)

CHA.—Frond pinnate, oblong, scaly. Pinnæ oblong, blunt, deeply cut, crenate.

SYN.—*Woodsia ilvensis*, *Brown, Smith, Hook., Spreng.*—*Polypodium ilvense*, *Swz., Willd., Schk.*—*Acrostichum ilvense*, *Linn., Huds., Ehrh.*—*Polypodium Arvonium* of *With.*, in description but not in references.*

FIG.—*E. B. Supp.* 2616.—*Flo. Dan.* 391.—*Pluk. Phyt.* 281, fig. 4, (good.)

DES.—Root perennial, tufted, black, smooth. Fronds numerous, 1 to 4 inches high, covered with capillary, brownish-white scales. Rachis scaly; the lower third of it without pinnæ, the upper two-thirds containing six to eight pairs, placed nearly opposite to each other. Larger pinnæ cut into from four to six blunt segments on each side. Sori scattered, convex, consisting of five or six roundish thecae. Cover torn into a few capillary divisions.

Mr. Sowerby observes, that the capillary segments of the indusium are not so numerous as in the next species, and the thecae more spherical. The plant cultivated and formerly sold at the London nurseries, under the name of *Woodsia ilvensis*, is *Notholaena distans*, a plant in every respect different from ours, which is much smaller, and less white and downy than that New Holland species.

SIT.—On rocks in mountainous countries.

HAB.—Higher parts of the Tees, *Mr. J. Hogg*. Rocks, (near where *Oxytropis campestris* grows,) between Glen Dole and Glen Phee, in the Clova Mountains,

* I cannot refer to *Withering's* *Polypodium Arvonium* and *ilvense* with certainty, as his description of these two plants is very obscure and far from characteristic.

Forfarshire, at 550 yards of elevation, *Mr. H. C. Watson*, (from which station it is larger than the Welsh plant.) On the Basaltic Rocks, called Falcon Clints, near Caldron Spout, Teesdale, *Mr. R. B. Bowman*. Glydes-vawr, near Lyn-y-cwm, *Mr. Winch*. Last seen in July, 1836, by *Mr. W. Wilson*.

GEO.—Found in different parts of Germany, as on the Alps of Salzburg and Carinthia, the Giant and Hartz Mountains, &c. In Sweden, Norway, and the Isle of Elba or Ilva, (whence the name *Ilvensis*;) also in Italy, Siberia, and on the Pyrenees. Pursh says from Canada to Virginia, but it may be much doubted if our plant be here indicated.

WOODSIA HYPERBOREA.

ROUND-LEAVED WOODSIA.

(Plate 1, fig. 6, B.)

CHA.—Frond pinnate, oblong, nearly smooth. Pinnæ triangular, blunt, deeply crenate.

SYN.—*Woodsia hyperborea*, *Br., Hook., Smith, E. Fl., Galp.*—*Acrostichum Alpinum*, *Bolt.*—*Ceterach Alpinum*, *Lam., Decan.*—*Polypodium hyperboreum*, *Swz., Willd., Spreng., Smith in E. B.*

FIG.—*E. B.* 2023.—*Bolt.* 42.—“*Linn. Trans.*” vol. xi.—*Pluk. Phyt.* 89, f. 5.

DES.—Root perennial, fibrous, black, tufted, and very long, giving rise to many oblong fronds, from 2 to 4 inches high. Lower part of the stem covered with light brown capillary scales. Eight or ten pairs of pinnæ, only the two or three lower pairs opposite, and these not constantly so, all nearly smooth, bluntly triangular, deeply crenate, or cut into two or three segments on each side. The upper half of each pinna larger than the other, and in luxuriant specimens cut into lobes near the stem. Sori from six to ten on each pinna, placed near the edge, light brown, very large, and often confluent.

From the very numerous segments of the indusium, a sorus appears like a bunch of hairs. The discriminating character is, however, chiefly the less cut, shorter, and more alternate pinnæ. The plant known as *Woodsia hyperborea* by gardeners is in reality a large variety of *Woodsia Ilvensis*, known as such before the separation of the present from that species.

SIT.—Found only on the highest rocks and mountains of Wales and Scotland.

HAB.—Ben Lawers, *Dr. Murray* and *Mr. W. Wilson*. Clova Mountains, *Mr. G. Don*. Craig Chailleach, Perthsh., *Mr. Maughan*. Mael Ghyrddy, Perthsh., and on Snowdon, below Bwlch-y-Saeth (Clowwyn-y-Garnedd), at an elevation of 2500 feet and upwards, very sparingly, *Mr. W. Wilson*. Mr. C. C. Babington, says, “I was not able to find this plant on Glydr Fawr, Caernarvonshire, July 1835, although in company with J. Roberts, Esq., of Bangor, who knew its station well. It is, I fear, exterminated in that place.” I searched for it in the same spot in 1837, and a botanical friend in 1840, but both without success.

GEO.—Lapland, Germany? France? (*Swz.*) Lulea, in Lapland (*Spreng.*) Canada, and high mountains of Pennsylvania and Virginia.

CISTOPTERIS, Bern. BLADDER-FERN.

(κυστος, a bladder, ὤψις, a fern; the indusiums being like bladders.)

A, one of the pinnæ of the frond of *Cistopteris fragilis*. B, a lobe magnified. C, young sori and indusia. D, sori cut transversely. E, theca. F, seed. G, indusium magnified from Bauc's "*Genera Filicum*." H, ditto from Schott's "*Genera Filicum*."

The genus is distinguished by its indusiums being inflated like bags, not being attached by a central column, but only by the edge nearest this rachis, and finally, either quite bent back or thrown off altogether. They first open on the top, or on the side nearest the apex of the frond or pinna.

1.—CISTOPTERIS DENTATA.

TOOTHED BLADDER FERN.

(Plate 2, fig. 1.)

CHA.—Frond bipinnate, oblong, lanceolate. Pinnæ ovate, lanceolate. Pinnules ovate, obtuse, crenate. Sori distinct.

SYN.—*Cystea dentata*, Eng. Flo.—*Cyathea dentata*, Eng. Bot., Dav. W. Bot. Galp.—*Polypodium dentatum*, Dicks., With., Hull.—*Aspidium dent.*, Swz., Willd., Hook. in Fl. Sco., Decan.—*Athyrium dentatum*, Gray.

FIG.—E. B., 1588.—*Pluk. Phyt.* 179, f. 5 (a cultivated specimen).—Bolt 27.

DES.—Root tufted, black, fibrous. Fronds numerous, oblong, lanceolate, 6 to 9 inches high, herbaceous. Stem slender, smooth, green except at the lower part, winged near the apex, without pinnæ for one-third of its height, above this bearing about fourteen pairs, opposite to each other. Pinnæ ovate, blunt, length twice their width, their main rib winged. Pinnules about ten pairs in the larger pinnæ, decurrent, ovate, obtuse, crenate or toothed, very rarely cut into distinct lobes, unless in luxuriant specimens, when the frond becomes wider, the pinnules very deeply cut, and sometimes petioled, but never losing their ovate, roundish, blunt form. Sori scattered, and always remaining distinct; Sir J. E. Smith says confluent, but this does not agree with any of my specimens, though probably in hot weather they may be found so.

Our plant *Cistopteris dentata* is to be known from every state of *Cistopteris fragilis*, in the shape of its frond and pinnules, which in this are very much blunter, rounder, and less divided; its rachis also is shorter and less brittle, and the whole smaller than the next species.

1

2

3

4

5

6

SIT.—On rocks in the north of England and Wales; also in Scotland.

HAB.—Ben Lawers, Perthshire, *Mr. R. Maugham*. Cadair Idris, and on rocks near Wrexham, *Mr. J. E. Bowman*. Rocks near Barmouth, *Mr. Purton*. Snowdon, *Mr. C. C. Babington*. Craig Breidden, Montgomerysh., *Rev. A. Bloxam*. Castle Dinas, *Mr. W. Leighton*. Common about Settle, *Mr. J. Tatham*, and *Mr. Chorley*. β . On lofty hills in the North, *Sir J. E. Smith*. Near Llanberris, Caern., *Mr. Lloyd*. Cordale, in Craven, *Mr. Curtis*. Downton, in Herefords.,—

GEO.—Common in Germany, Switzerland, Dauphiny, Prussia, Holland, Verona, &c.

2.—CISTOPTERIS FRAGILIS.

BRITTLE BLADDER FERN.

(Plate 2, fig. 2.)

CHA.—Frond twice-pinnate, lanceolate. Pinnæ lanceolate. Pinnules ovate, pointed, deeply cut, toothed, decurrent.

SYN.—*Cystea fragilis*, *E. Fl.*—*Cistopteris fragilis*, *Hook. in Br. Fl.*, *Mack.*, *Bernh.*—*Aspidium fragile*, *Swz.*, *Hook. in Fl. Sco.*, *Willd.*, *Grev.*, *Lightf.*—*Polypodium fragile*, *With.*, *Linn.*, *Huds.*, *Bolt.*, *Hoffm.*, *Ehrh.*, *Dick.*—*Cyathea fragilis*, *Roth*, *Smith in E. B.*, &c. *Galp.*—*Cyclopteris fragilis*, *Schrad.*, *Gray*.

FIG.—*E. B.* 1587—*Bolt.* 45-46.—*Flo. Dan.* 401.

DES.—Root black, fibrous, and tufted. Fronds numerous, deciduous, bright green, from 6 to 12 inches high, twice-pinnate, lanceolate, pointed, and finely tapering towards the apex. Rachis very brittle and shining, of a dark brown or black color on the lower part, and quite smooth, except a tuft of scales at the very base. Pinnæ opposite, pointed, about twenty pairs, confined to the upper half of the rachis, and growing nearly at right angles to it. Their length more than twice their width, except the lower pair, which are also distant from the next above them. Pinnules alternate, acute, deeply lobed, crenate or bluntly acute, decurrent and tapering more or less at the base. Sori numerous, confluent, black when young, afterwards a shining brown, and found throughout the summer. Indusium white, with an irregular margin, and soon obliterated or thrown off by the growing thecæ.

In general habit resembling the last species, but instantly to be distinguished by the shape of the frond, which is sharper and longer pointed, as is also the case with the pinnæ and pinnules; the whole is also much more divided, all the larger pinnules being cleft, and not merely toothed, as in every state of *Cistopteris dentata*. The stem is also darker, longer, and more brittle, and the sori so numerous as soon to become confluent.

No Ferns are more altered by circumstances than this genus, hence the difficulty of distinguishing the species. The varieties, however, are not distinct in themselves, as they may all sometimes be found upon the same plant, and different seasons produce differently-shaped and more finely-divided fronds. For example, those which arise in ordinary seasons alone answer the above description; a cold

spring occasions barren fronds, the pinnules of which are rounded, delicate, wide, crenate, and running much into each other, while long-continued drought or warm weather occasions those fronds which arise in summer to be much smaller, much yellower, more entire, and the sori more crowded. In the extreme state it may be described as follows:—Fronde linear, oblong. Pinnæ blunt, pinnate, ovate or round, toothed, quite covered with sori. If the summer continue very wet and cold, the fronds do not take the above character, but have broader and darker colored pinnules; in this case exactly resembling the cultivated *Cistopteris dentata*, except in the shape of the frond itself.

β. (*angustata*.) Frond oblong, ovate. Pinnæ ovate, pointed.

Cyathea angustata, *E. B. and E. F.*—*Polypodium rhæticum*, *Dick., Bolt.*—*Aspidium rhæticum*, *Willd.*—By no means the *Polypodium rhæti.* of *Linnaeus*, nor the *Polypodium tenue* of *Hoffm.*, which is the *Aspidium intermedium* of modern authors.

Very distinct as a variety, not a species. It differs from the usual state of the plant only in a rather larger and broader frond, with pinnules doubly toothed and slightly pointed.

SIT.—On alpine rocks and other lofty situations.

HAB.—ENG.: Near Richmond, Yorks., *Mr. J. Ward.* About Settle, Yorks., *Mr. J. Tatham.* Cumberland, Ruins of Peveril Castle, Castleton, and the Lover's Leap, near Buxton, Derbys., *Mr. H. C. Watson.* Matlock, Derbys., *Dr. Howitt.* Cheddar, Somers.,* *Mr. W. C. Trevelyan.* Nottinghamsh., *Mr. T. H. Cooper.* Near Bristol, *Miss Worsley.* At Exwick, near Exeter, *Mr. Jacob.*—WAL.: Cave at Clogwyn Coch, Snowdon, and rocks above Cwn Idwel, near Tŷll Du, *Mr. W. Wilson.* Near Wrexham, Denbighsh., *Mr. J. E. Bowman.*—SCO.: Aberdeenshire, *Mr. H. C. Watson.* Moray and Rosshire, *Rev. G. Gordon.* Near Maens, Berwicksh., *Rev. A. Baird.* Sutherland and the Kincardineshire Coast, *Dr. Murray.* Near Killin, *Mr. W. Wilson.*—IRE.: Rocks and mountains of Kerry, *Mr. Mackay.* Lough Inn, and Lough Derryclare, Cunnemara, *Mr. Shuttleworth.*

GEO.—Common in Germany, Saxony, Switzerland, Holland, &c.

Dundry. Winifred. Burnington. Cheddar alps. super. near. West town bridge. 1855. Apr. 23. From the walls of Stoke Churchyard. Landover, Glentworth. Ellen and Clidore. 1856. (1856) C.

3.—CISTOPTERIS ALPINA.

ALPINE BLADDER-FERN. LACINIATED BLADDER-FERN.

(Plate 2, fig. 3.)

CHA.—Fronde tri-pinnate, ovate, lanceolate. Pinnules ovate, blunt. Segments linear, obtuse, toothed.

SYN.—*Cistopteris alpina*, *Hook. in Br. Fl., Desv.*—*Cistopteris regia*, *Bernh.*—*Cyathea incisa*, *Smith in E. Bot., Galp.*—*Cyathea alpina*, *Roth.*—*Cystea regia*, *Smith in E. Fl. & Fl. Br.*—*Polypodium regium*, *Lin., Hüll.*—*Polypodium trifidum*, *With.*—*Polypodium alpinum*, *Jacq., Schk.*—*Athyrium alpinum*, *Spreng.*—*Athyrium regium*, *Gray.*—*Aspidium alpinum*, *Swz. Willd., Hook. in Fl. Sco.*

FIG.—*E. B.* 163.—*Jacq. Icon. vol. 3 t. 742.*—*Sequier Pl. Veron. supp. 1, 3.*

DES.—Root black, fibrous, tufted. Frond tri-pinnate, ovate, or ovato-lanceolate, herbaceous, 2 to 6 inches high. Pinnæ about ten or twelve pairs, set rather alternately, except the lower pair, their length not above twice their width. Larger pinnules broadly

ovate, or wedge-shaped, repeatedly cut into broad linear segments. Sori small, scattered, seated nearly at the apex of the segments. Margin of the indusium entire.

These marks clearly indicate this to be a distinct species, far removed from both the others, and in cultivation instead of approaching the *fragilis* or *dentata*, it becomes yet more different, as the pinnules increase in length, but scarcely in width, as in the former cases. In general habit our present species is by far the tenderest and most numerously cleft, with a shorter and less brittle rachis than *Cistopteris dentata* or *fragilis*.

The late Professor Don thought the *Cistopteris regia* and *Cistopteris alpina* to be essentially different, but Sir W. J. Hooker speaks confidently of the Layton plant being precisely the same as that represented by Jacquin and Schkuhr, which are the same as the *alpina* of Don; and as our plant at the present time has the wedge-shaped pinnules, said by Mr. Don, to be peculiar to the *Cistopteris regia*, we are bound to conclude that formerly, when the plant was vigorous, it took one character, and now that it is but struggling for existence it assumes the other. Indeed luxuriant plants lately received from Low Layton, though the kindness of Mr. E. H. Bulton, who gathered it as lately as 1840, confirm to me the accuracy of Sir W. J. Hooker's view upon the subject. Mr. W. Pamplin, of Queen Street, Soho, an indefatigable botanist, is the re-discoverer of this plant, and kindly furnished me with specimens gathered in 1835. The first account we have of the plant as British is by Mr. Forster, in Symon's "Synopsis," published in 1793.

HAB.—Wall at Low Layton, Essex, 1836, *Mr. W. Pamplin*. Caernarvonsh, *Mr. J. E. Bowman*. Cwm Idwel, *Mr. Griffiths*. On Snowdon, near the Copper Mine, *Mr. Winch*. Ben Lawers, *Mr. Maughan*. Rocks at the Dropping Well, Knaresborough, *Mr. W. Christy*.

GEO.—Jena, Oldenburgh, and other parts of Germany, Italy, &c.

ASPIDIUM, Swz. SHIELD FERN.

(ασπίς, a shield; the indusium being of this form.)

A, pinnules of *Aspidium lonchitis*. B, portion of ditto, showing the fruit magnified. C, transverse section of a sorus. D, ditto of the stem. E, scale magnified. F, theca and spore.

A widely-distributed and extensive genus, of not less than from 160 to 170 species, all of them herbaceous, some evergreen, others deciduous. The indusium is either reniform and fixed at the sinus, when they belong to the genus *Nephrodium* of Brown, or else orbicular and peltate, which is the true character of *Aspidium*. The greater number of the British *Aspidia* somewhat differ from the

* Mr. W. Wilson writes me, that the Welch stations refer to *Cistopteris fragilis*. I have also received *Cistopteris dentata* from Craig Breidden, under the name of *Alpina*.

true character of the genus, as their indusiums, though orbicular, have a deep lateral notch, which occasions them to appear somewhat reniform, and hence also they in some degree cease to be peltate; but the variation is not so great as to render it advisable to separate them into two genera.

1.—ASPIDIUM LONCHITIS.

ROUGH ALPINE SHIELD-FERN.

(Plate 2, fig. 4.)

CHA.—Frond pinnate. Pinnæ lunate, bristly-serrate. Rachis scaly.

SYN.—*Aspidium lonchitis*, Swz., Willd., Hook., Smith, Mack., Galp., Spreng., Sehk.—*Aspidium asperum*, Gray.—*Polypodium lonchitis*, Linn., Bott., With., Huds., Lightf.—*Polystichum lonchitis*, Roth., Decan., Hoffm., Newm.

FIG.—*E. B.* 797.—*Bott.* 19.—*Flo. Dan.* 497.—*Park.* 1042.—*Ger.* 979.

DES.—Root tufted, black, fibrous. Fronds 6 to 12 inches high, numerous, dark green, arranged in a circle around the crown of the root, very rigid, not growing upright, but generally half decumbent, forming a flat, cup-shaped plant. Rachis scaly, clothed with pinnæ nearly to its base. The pinnæ are numerous, crowded, stalked, alternate, smooth above, slightly scaly beneath, crescent-shaped, with an auricle on the upper side of the base of each, serrated, with the serratures ending in a bristle, that part of the pinna above its midrib much larger than the lower portion, in position rather declining and bent forwards, so that they very often approach those on the opposite side of the rachis, the back of the frond being outwards. Sori confined to the upper third of the frond, arranged in single rows, black or brown, and very large. Cover orbicular, notched, attached at the centre, and soon becoming shrivelled.

Sir J. E. Smith says, that "this plant dwindles rather than becomes luxuriant when cultivated," as it often is on rock-work, &c., forming a curious, rigid, and pretty plant, not in any way altered from its original characteristics, except becoming less spinous. The American is more spinous than our plant.

HAB.—In situations above 1000 yards, probably 1100 yards above the sea level, on the Breadalbane mountains, Perthshire, and plentiful almost every where in the Highland valleys, and on the declivities of the mountains. Scarce in England, nor have I ever seen it here. Craig Chailleach, Perth., and Clova mountains, Forfarsh., *Mr. H. C. Watson*. Falcon Clints, near Cauldron Spout, Teesdale, *Mr. R. B. Bowman*. Glen Isla, Forfarsh., *Mr. W. Brand*. Aberdeenshire, *Dr. Murray*. Moray and Rosshire, *Rev. G. Gordon*. Base of Benmore, Sutherland, *Dr. Johnston*. Very large in Glen Fee, *Mr. W. Wilson*. Clogwyn-y-Garnedd, Snowdon, *Mr. C. C. Babington*. Higher part of the Tees, *Mr. J. Hogg*. Common about Settle, Yorks., *Mr. J. Tatham*.—IRE.: In a glen E. of Lough Eske, Donegal; and on Glenade Mountain, Leitrim, *Mr. E. Mackay*. Brandon Mountain, *Mr. W. Wilson*.

GEO.—Silesia, Bavaria, the Tyrol, Switzerland, Sweden, Norway, and the Aleutian Islands.

2.—ASPIDIUM LOBATUM.

CLOSE-LEAVED, PRICKLY SHIELD-FERN.

(Plate 2, fig. 5.)

CHA.—Frond bipinnate. Lobes decurrent, spinulose, elliptical, that next the rachis very large.

SYN.—*Aspidium lobatum*, Swz., Gray, Willd., Schk., Smith, Hook. in *Bri. Fl.*, not in *Flo. Scot.*, Forst., Galp., Mack.—*Polypodium lobatum*, Huds.—*Polypodium aculeatum*, Bolt., With.

FIG.—*E. B.* 1563.—*Bolt.*, 26, f. 1, (a full-grown,) f. 2 (a young plant.)

DES.—Root tufted. Fronds growing from a circle, rigid, glaucous green, from 15 inches to 2 feet high, evergreen, perfectly ovate. Lower pinnæ crowded, so as to overlap each other; sometimes, however, the frond is elongated at the lower part, when the pinnæ are proportionably distant. Rachis stout, scaly, and with pinnæ to the very base. Pinnæ short, alternate, lanceolate, pointed, and curved upward, therefore somewhat lunate. Smaller pinnules running much into each other, the larger slightly auricled, decurrent, and that next the rachis so much larger than the rest as to project over its next neighbour, and also partly to conceal the base of the pinna next above it; the inner edge of all the larger lobes running parallel to the rachis, and at a little distance from it, so that if held up, a line of light will appear on each side of the rachis, except near the base, where the first lobes are set very close to the main stem, whence perhaps its name of close-leaved. Sori large, in single rows, confined to the top of the frond. Cover orbicular, fixed by the centre, persistent, but easily knocked off.

β (*lonchitidoides*.) Pinnules combined, forming nearly a pinnate frond. *Filix lonchitidi affinis*, Ray. *A. aculeatum* β, Smith in *E. Fl.* *A. lobatum*, Hook. in *Br. Fl.* FIG.—*Pluk*, *Phyt*, t. 180, f. 3. (good.)

This species is distinguished from the following, for which alone it can be taken, by the decurrent lobes; and as Sir J. E. Smith very rightly observes, “by the much shorter, more crowded, and less scaly pinnæ.” Added to which, the lobes are more entire, being but slightly auricled, very convex, thick, and of a glaucous color, furnished with a less number of and smaller bristly serratures, sometimes wanting them entirely at the sides. The sori also are more confined to the top of the frond, and larger than in *A. aculeatum*. The variety *lonchitidoides* is not very scaly, and in form and size exactly intermediate between this species and *A. lonchitis*.

SIT.—On shady banks and damp hedge rows, chiefly in the north.

HAB.—Extremely common in Scotland and in the north of England, gradually losing itself towards the south, and becoming more and more intermingled with *A. aculeatum*, which in its turn is superseded still more southerly by *A. angulare*. In the middle and south of England, its recorded habitats are

F

See Notes at the end of this Volume —

Leicestershire, *Rev. A. Bloxam*. Common about Settle, Yorksh., *Mr. J. Tatham*. Pottery Car, near Doncaster, *Mr. S. Appleby*. Matlock, Derbysh., *Dr. Howitt*. At Studley, Sambourne, Overley, and Weatherly, Warwicksh., *Rev. W. Bree*. Lane leading to the Vachè from Chalfont, Bucks, *Mr. A. Halley*. Near Bristol, *Miss Worsley*. Near Dorking, Surrey; in Hants, &c. *Mr. W. Pamplin*. Near Yarmouth, *Mr. Paget*. Sussex and S. Kent, *Rev. G. E. Smith*. WAL.: Near Wrexham, Denbigh, *Mr. J. E. Bowman*.—IRE.: Collinglen, near Belfast, *Mr. J. Templeton*. Hermitage, County Wicklow, *Dr. Osborne*. County of Derry, *Mr. D. Moore*. β Glen Fee, Clova Mountains, *Mr. W. Wilson*. Braid Woods, near Edinburgh, *Mr. H. Cooper*. (*Quarry, & Stakewood near Bursleigh*)
East of the place & cont.
Cheddar Cliffs July 1854

GEO.—Germany, Switzerland, &c.

3.—*ASPIDIUM ACULEATUM*.*

COMMON PRICKLY SHIELD-FERN.

(Plate 2, fig. 6.)

CHA.—Frond bipinnate, broadly lanceolate. Lobes petioled, ovate, distinctly auricled, aristate. Rachis scaly.

* SYN.—*Aspidium aculeatum*, Swz., Willd., Hook. in *B. Fl.* ed. 4, Smith, Galp., Mack., Gray.—*Aspidium lobatum*, Hook. in *Fl. Scot.*, Schk.—*Polypodium aculeatum*, Linn., Huds., Lightf., Ehrh.—*Polystichum aculeatum*, Roth., Decan.

FIG.—*E.B.* 1562.—*Pluk. Phyt.*, 180 f. 1. (*not good*.)

DES.—Root tufted. Fronds numerous, perfectly lanceolate, evergreen, bipinnate. Pinnæ alternate, gradually tapering, close together, their midribs covered with hair-like scales. Lobes ovate, distinctly petioled, serrate, spinulose, and with an auricle on the upper side at the base of each; that next the rachis larger than the rest, but not so much so as in the last species; all remaining distinct from each other nearly to the point of the pinna, although sometimes so crowded as to overlap. Rachis clothed with pinnæ to its base, and very sealy. Sori distinct, brown, small. Cover orbicular, fixed by its centre, soon withering.

* As many gentlemen, distinguished for their botanical knowledge, consider the *Aspid. aculeatum* and *A. angulare* as distinct species, it is necessary that I should state the reasons why I have blended these two plants together. I have not been guided by any desire of innovation, believing unsteadiness of nomenclature and of classification to be the bane of science, but because after the most careful examination of specimens from all parts of Great Britain where they grow, and after consulting all the most celebrated practical botanists that I have the honor to be acquainted with, I have found it absolutely impossible to draw the line of demarcation between the plants. To delineate extreme states of any variable plant is easy enough, but where there is so regular a gradation from the robust pointed pinnales to the blunt and delicate ones, the difficulties of discrimination are insurmountable. Also, upon writing to various gentlemen for specimens and habitats, I have received the same plant repeatedly under the two names, and it is very remarkable that the habitats received invariably refer to both varieties, though they have not always been received from the same person. Thus Dr. Johnston says, that both grow at Pease Bridge, Berwickshire; Mr. Bowman says of *Aculeatum*, near Richmond, Yorkshire; while Mr. J. Tatham notes the same place as a station for the *Angulare*. Thus doubts arise if the same or different plants are indicated. The name *Angulare* appears, however, by far the more commonly given to it, and I should for this reason have preferred it to *Aculeatum*, in deference to the opinion of my countrymen; but foreigners give the name *Angulare* to an Hungarian Fern very different from ours, and as *Aculeatum* is the specific name of all authors who have

* species combined the two Ferns, and is by far more expressive. I have adopted it. This Fern is *A. angulare* of Smith & of Hooker, and therefore it is not synonymous with the *A. aculeatum* of those Authors—it is best distinguished by having the pinnales very distinctly petioled nearly to the extremity of the pinna, whilst in *A. aculeatum*, or *A. lobatum*, the pinnales are all decurrent. See *Natural History of Botany*.

1

2

3

4

5

6

7

This plant varies much in the sharper or blunter shape of the lobes of the leaves, for which reason it is sometimes extremely difficult to decide if a frond be of this species or the former. Luxuriant plants assume much the appearance of *Lobatum*, as the large pinnules become slightly decurrent: but in this state they become somewhat deeply cut, or even compound, while in the last species they are truly entire, losing their serratures instead of becoming more cleft by culture.

β (*angulare.*) Pinnules short, blunt, distinctly auricled. Rachis very chaffy. *Aspidium angulare*, *Smith* in *E. Fl.*, *Hook.*, *Mack.*, *Willd.* *A. aculeatum* β . *Smith* in *Fl. Br.* FIG.—Plate 2, f. 6 β . *E. B. Supp.* 2776.

γ (*linearis*) Pinnules linear and very sharp pointed. FIG. Pl. 2, f. 6 γ .

These are well marked varieties, yet not sufficiently distinct either in habit or character to constitute separate species. The var. β has, when luxuriant, its lower and larger pinnæ compound; when it becomes of course subtripinnate, and larger, (but not comparatively more robust,) thereby differing from the first or normal state of the plant, which alone approaches the last species in occasionally decurrent and convex pinnules.

SIT.—Common in hedge rows, damp banks, &c., chiefly in the south.

HAB.—SCO.: Peasebridge, *Dr. Johnston*. ENG.: Near Richmond, Yorks., *Mr. J. Tatham*. Burton Wood, near Warrington, Lanc.; and in Cheshire, *Mr. Rylands*. Ulverscroft Priory, Charnwood Forest, *Rev. A. Bloxam*. Isle of Man, *Mr. Forbes*. Derbyshire, *Dr. Howitt*. Warwickshire, *Rev. W. T. Bree*. Somerset, *Mr. A. Southby*. Little Worley Common, Essex, *Mr. R. Castles*. About Tonbridge Wells and elsewhere, Kent, (abundant,) and near Bramshot, Hants, *Mr. W. Pamplin*. Osterley Park, Lampton Lane, and Sion Lane, near Brentford, Midd., *Mr. J. Bevis*. Near Hastings, *Mr. W. C. Trevelyan*. Sussex, *Rev. G. E. Smith*. Kingsteignton, *Mr. Anderson*. Near Gurnet Bay, Isle of Wight, *Prof. Henslow*. —WAL.: Near Wrexham, Denbighs., *Mr. J. E. Bowman*. Cickle, near Beaumaris, Anglesea, *Mr. W. Leighton*. Near Bangor and Caernarvon, *Mr. W. Wilson*. IRE.: Colin Glen, Belfast, *Mr. Mackay*. Hedgebanks, near Carrickfergus, *Mr. F. Whittle*. β Intermixed with and even more common in the extreme south of the kingdom than the first state of the plant.— γ Near Clonmell, *Mr. G. S. Gough*.

GEO.—Europe generally, Arabia, Cape of Good Hope, North Africa, on the Green Mountains, Vermont, and other places in North America.

4.—ASPIDIUM THELYPTERIS.

MARSH SHIELD-FERN.

(Plate 3, fig. 1.)

CHA.—Frond pinnate, erect. Pinnæ linear, lanceolate, smooth. Segments mucronated. Sori small. Root creeping.

SYN.—*Aspidium thelypteris*, *Swz.*, *Willd.*, *Smith*, *Hook.*, *Galp.*, *Mack.*, *Pursh.*—*Polypodium thelypteris*, *Linn.*, *E. B.*, *Dicks.*, *Ehrh.*, *With.*, *Lightf.*, (not of *Huds.*)—*Acrostichum thelypteris*, *Linn.*, *Bolt.*—*Athyrium thelypteris*, *Spreng.*—*Polystichum thelypteris*, *Rotk.*—*Lastrea thelypteris*, *Presl.*, *Newm.*

FIG.—*E. B.* 1018.—*Flo Dan. t.* 760.—*Bolt.* 43, 44.—*Newm.*, page 46.

DES.—Root creeping, furnished with long, black, slender, rather smooth runners, giving rise at various points along their surface to black radical fibres, and erect, light green, smooth ovate, or (when fertile) oblong fronds, each from 6 to 12 inches long, having a

slender, and generally smooth rachis. Pinnæ linear-lanceolate, pointed, deeply pinnatifid, petioled, opposite. Segments oblong, obtuse, occasionally with a very small point; the first upper segment on each pinna much longer than the others. Sori in continued longitudinal lines near the margin of each segment, small, brown or black, at first distant, afterwards confluent. Cover thin, white, round, kidney-shaped, fastened near the centre, and soon lost among the growing thecæ. The barren fronds differ much from those which are fertile; they are altogether wider, shorter and flatter, with the pinnæ horizontal, and rachis void of pinnæ half way up. The fertile fronds have two-thirds of the rachis covered with pinnæ: which are more numerous, deflexed, and curled, particularly at the point. The edges of the pinna, folding over the lines of sori, give it an acute appearance.

The only British Fern with which it is possible to confound this is *Aspidium oreopteris*, from which it differs in its smaller size, lighter color, more ovate frond not contracting so much below, the folded segments of the pinnæ, and its creeping root. This last character will distinguish it from all our other species of this genus, it being the only one of which the root is not tufted. It is by no means easily cultivated, nor frequent in fruit when wild, as the fertile fronds do not rise till late in the season. While undergoing the process of desiccation for the herbarium, the elasticity of the annulus of the theca is very apparent, bursting with violence, and scattering the spores in all directions and to a considerable distance.

HAB.—Common in Scotland, *Sir W. J. Hooker*. Learmouth Bogs, Northum., *Mr. Winch*. Near Settle, Yorks., *Mr. J. Tatham*. Allesley, Warw., *Rev. W. Bree*. Knutsford Moor and New Church Bog, near Over, Cheshire, *Mr. W. Wilson*. Oxtou Bogs, Notts., *Dr. Howitt*. Windsor Park and Sunning Hill Wells, Berks., *Mr. J. Bevis*. Valley below Cæsar's Camp on Wimbledon Common, planted there some years ago by *Mr. Tyton*. Bog on Waterdown Forest, near Tunbridge Wells (1835), *Mr. Pamplin*. Somersct., *Mr. Southby*. Belton, Suffolk, *Mr. Paget*. Sussex, *Mr. Borrer*. Border of Lake near Red Wharf, Anglesea, *Mr. W. Wilson*. Beaumaris, Anglesea, *Mr. J. E. Bowman*. Marshes at Glencree, County of Wicklow; and Mucruss, Killarney, *Mr. Mackay*.

GEO.—Pomerania, Mecklenburgh, Prussia, Denmark, Sweden, N. and S. Africa, and in all the United States, but seldom with fruit.

Abundant on Woodmoor, Somersct. (S.)

5.—ASPIDIUM OREOPTERIS.

HEATH SHIELD-FERN.

(Plate 3, fig. 2.)

CHA.—Frond pinnate, lanceolate. Pinnæ glandulous, deeply cleft. Segments blunt, entire. Root tufted.

SYN.—*Aspidium oreopteris*, *Suz.*, *Willd.*, *Smith*, *Hook.*, *Galp.*, *Spreng.*, *Mack.*, *Schk.*—*Aspidium odoriferum*, *Gray.*—*Polypodium oreopteris*, *Ehrh.*, *Dicks.*, *With.*, *Hull*, *Sibth.*, *Hoffm.*, *Linn.*—*Polypodium thelypteris*, *Huds.*, *Bolt.*, *Lichtf.*, *Hedw.*—*Polystichum montanum*, *Deccau.*—*Lastræa oreopteris*, *Presl*, *Newm.*

FIG.—*E. B.* 1019.—*Flo. Dan.* 1121—*Bolt.* 22, f. 1 and 2.

DES.—Root tufted, large, black, scaly, fibrous. Fronds several, growing in a circle from a crown, finely lanceolate, tapering at both ends. Rachis covered with fine hair on the upper part, and with a few scattered scales on the lower, delicate green, with a deep channel on the upper side. Pinnæ extending nearly all along the rachis, more or less alternate, sessile, deeply pinnatifid, tapering to a fine point, on the upper side smooth, on the under side hairy particularly about the main rib, and covered with yellowish, shining glands, smelling of turpentine. Segments very numerous, flat, blunt and entire. Sori marginal, at length confluent, covering all the pinnæ. Cover thin, white, kidney-shaped, soon shrivelling up.

The fresh plant may instantly be known from all its congeners by the smell emitted when drawn through the hand, or by holding it up to the light, in which situation it shows very plainly translucent, minute points, very similar to those seen in *Hypericum perforatum*; though, be it observed, that unfavorable situation and cold weather will often prevent the formation of, if not obliterate these odorous pores. They are most abundant when the plants grow in sunny, but not too dry localities. This Fern can only be mistaken for *As. thel.* or *Asp. Fil.-mas*; it has already been distinguished from the former in describing that plant, from the latter it may easily be known by its more elegant shape, its smaller size and more delicate structure, no less than by its greater smoothness in every part, particularly its rachis. The segments of the pinnæ also are not crenate, as in *Filix-mas*, and the sori, which in that are large, distinct, and confined to the lower half of the segment, are in this plant small, closer together, more numerous, and continued throughout the whole length of the segment, very near the margin.

SIT.—On heaths and in shady lanes, not uncommon in the north.

HAB.—SCO.: Glen Isla, Forfarsh., *Mr. W. Brand.* Common in Sutherland, *Dr. Johnston.* Banks of Loch Tay, *Mr. T. H. Cooper.* Aberdeenshire, but not common, *Dr. Murray.* Foot of Craig Challeach, &c., *Mr. W. Wilson.*—ENG.: Near Chapel Weardale, Durham; and Cawsey Dean, near Newcastle, *Mr. R. B. Bowman.* Keswick, and near Lodore Waterfall, Cumbl., *Mr. H. C. Watson.* By the Tees, *Mr. J. Hogg.* Near Richmond, Yorks., *Mr. J. Ward.* Coleshill Heath and Corley, Warw., *Rev. W. Bree.* Near Warrington, *Mr. W. Wilson.* Dethick Moor, and near Riley, Derbys., *Dr. Howitt.* Isle of Man, *Mr. Forbes.* Dallington Heath, near Northampton, *Mr. Anderson.* N. side of Shotover Hill, Oxfordsh., *Mr. Baxter.* Oxtou and Eddingley Bogs, Notts; and Hartswell, near Farnsfield, *Mr. T. H. Cooper.* Somerset, *Mr. A. Southby.* Bradwell, Suffolk, *Mr. Turner.* Sussex and Kent, *Rev. G. E. Smith.* Bailey's Hill, between Brasted and Tunbridge, (1835,) *Mr. Pamplin.*—WAL.: Near Wrexham, Denbighshire, *Mr. J. E. Bowman.* Llanberris and Nant Gwynedd, Caernarvonsh., *Mr. C. C. Babington.* Frequent in Caernarvonsh., *Mr. W. Wilson.*—IRE.: Powerscourt Deer Park and Waterfall, Mangerton Mountain, *Dr. Osborne.* Lough Corril, Galway, *Mr. Shuttleworth.* Plentiful in Ireland, *Mr. Mackay.*

GEO.—Germany, Italy, Switzerland, Prussia, &c.

abundant about Linton. A Devon - also on the Peak Hill, Sidmouth
as at Conkane Ferry, Gloucestershire. C. J. 1850 - 112 1050 -

6.—*ASPIDIUM FILIX-MAS*.

MALE FERN.

(Plate 3, fig. 3.)

CHA.—Frond pinnate, broadly lanceolate. Pinnæ alternate, deeply pinnatifid. Segments obtuse, crenate. Rachis scaly.

SYN.—*Aspidium Filix-mas*, Swz., Willd., Smith, Hook., Galp., Mack.—*Polypodium Filix-mas*, Linn., Huds., Bolt., Woodv., Dicks., Ehrh., Ger., With., Lightf.—*Polystichum Filix-mas*, Roth, Decan.—*Polystichum callipteris*, Bernh.—*Lastræa Filix-mas*, Presl, Newm.

FIG.—*E. B.* 1458.—*Bolt.* 24.—*Woodv.* 49.—*Flo. Lon.* 40.—*Newm.*, page 51.

DES.—Root large, tufted, black, and scaly. Fronds growing centrally from a crown, broadly lanceolate, pinnate. Pinnæ lanceolate, pointed, alternate, smooth, except on the under side of the midrib, of a bright green, regularly tapering, curved upwards, and very deeply cleft. Segments oblong, obtuse, slightly crenate at the sides, copiously at the end, very close together, but not overlapping each other. Sori confined to the upper half of the frond, and to the lower half of each segment of the pinnæ, round, large, and very prominent. Cover large, orbicular, with a notch on one side, at first white and transparent, afterwards opaque, and of a fine reddish brown, covering the thecæ even till they are fully ripe.

The large size, robust appearance, and decided character of this plant, obtained for it very early and very aptly the name of Male Fern. Medicinal properties of some importance have been ascribed to it, and apparently with justice. It is retained in most of the pharmacopœias of Europe as a specific for the larger kinds of intestinal worms, and used very extensively for that purpose by the faculty on many parts of the Continent, and if the employment of it has been discontinued here, it is not because of its inutility, but from the discovery of other remedies equally potent and better understood. The stem and roots are bitter and astringent, and have been used instead of hops.

β (*variegatum*.) White, tipped and edged with green, (same habit.)

γ (*recurvum*.) Pinnæ crisped, turned down. Frond small. Rachis smooth.

δ (*spinsum*.) Pinnules serrate, smaller blended together, larger auricled.

The above states of the plant appear constant, besides which it is sometimes found with a cormus, some inches above the ground; Mr. W. Wilson has seen it thus in Caernarvonshire, and Mr. Mackay in Wicklow. A singular variety with the upper pinnæ remarkably compound or branched has been observed in Bore-hill Lane, below Dorking, Surrey, by Mr. W. Pamplin. Also Mr. T. Clarke, Jun., of Bridgewater, has been so kind as to send me from King's Cliff Valley, four miles from that town, several fronds of a very large variety, which is found there in considerable abundance. It is of a very dark color, has sori along the whole pinnule, and the pinnules themselves are all deeply serrated along their margin. Mr. Clarke also writes me, that Sir W. J. Hooker confirms his, and I

may add, my opinion also, that this plant is a variety of *Filix-mas*, though Sir William observes that Schkuhr, who found it near Dresden, looked upon it as a new species, and figured and described it as *A. erosum*. The continental *A. Filix-mas* is usually more crenate or serrate than ours.

SIT.—Hedge-banks, &c., and in shady lanes throughout the kingdom.

HAB.—I have received numerous habitats from most of the English and Scottish counties, from the extreme south to the Orkney Islands, and yet in some places this plant is rare. Inchnadamff, in Sutherland, is one of these.— β : Near Keswick, Cumberland, *Mr. H. C. Watson*.— γ : Not very uncommon in dry situations in the south.— δ : Bomcre Pool and Sutton Spa, both near Shrewsbury, *Mr. W. Leighton*. Nettlecomb, Somerset, *Mr. W. C. Trevelyan*.

GEO.—North America, throughout Europe, and in Africa.

7.—ASPIDIUM CRISTATUM.

CRESTED SHIELD-FERN.

(Plate 3, fig. 4.)

CHA.—Frond pinnate. Pinnæ opposite, pinnatifid, oblong, obtuse. Segments ovate, decurrent, crenate, bristled.

SYN.—*Aspidium cristatum*, Swz., Willd., Smith, Hook., Spreng., Galp., Mack., Schk., Pursh.—*Polypodium cristatum*, Linn., Afzel in Stockh. Trans. for 1787.—(Not of Bolt., With., or Huds.)—*Polystichum cristatum*, Roth, Decan, Hoffm.—*Polypodium callipteris*, Ehrh., Hoffm.—*Lastræa cristata*, Presl, Newm.

FIG.—Hook. in *Flo. Lon.*, new ser. 113.—E. B. 2125, (not 1949.)—Newm. page 54.

DES.—Root tufted. Fronds erect, rigid, yellowish green, bipinnate, oblong, blunt. Pinnæ opposite, eight to fourteen pairs, very distant from each other, short, ovate, oblong, obtuse, very deeply pinnatifid or rather pinnate at their lower part. Segments ovate, crenate, each crenature furnished with two or three small sharp points or bristles, the principal vein in each segment slightly crooked, but the midrib of the whole pinna straight. Rachis slightly scaly only towards the lower part, where for about one-third of its height it is otherwise naked. Sori large, very distinct, black at first, afterwards brown. Cover white when young, very thick, circular, with a lateral notch, and fixed by the centre.

Few plants have occasioned more discussion than this. The difficulty has arisen chiefly because sufficient stress has not been laid upon the simply pinnate character of the frond; had this been regarded more, *Aspidium spinulosum* would not so often have been confounded with it. The *cristatum*, besides being less divided, has a more obtuse, more linear frond, and contracts very much below. The sori of *cristatum* are comparatively much larger and less numerous, and their covers persistent, not hidden by the capsules. It very nearly resembles the American *Aspidium goldianum*.

HAB.—This is one of the rarest Ferns, not only here but on the Continent. The only recorded habitats of it in this country are the Lows in Holt-heath, Norfolk, *Rev. R. B. Francis*. On bogs among alder bushes, at Westleton,

Suffolk, *Mr. Dary*. Oxtou Bogs Notts, *Dr. Howitt* and *Mr. T. Cooper*; and lately discovered on Edgefield Heath, and at Fritton, Norfolk, by *Mr. Wigham*, of Norwich. Even one of these habitats may, perhaps, be now expunged, as *Mr. Dennes* informs me it is thirty years since it was last found at the Lows in Holt-heath. It was stated on page 70, of the first edition, that I had reason to believe that this plant grew on Wimbleton Common; this was an error of judgment or of memory in my informant. It does not grow there, but the *A. spinulosum* does. *Mr. Mackay* admits it into the Irish Flora, as growing in the grounds of Sir H. Gough, at Rathronan, near Clonmel, found there by *Mr. G. S. Gough*, in 1835; he says that the Irish plant is acutely serrate.

GEO.—Oldenburgh, Bremen, Mecklenburgh, Hanover, and other parts of Germany. New York to Virginia.

8.—ASPIDIUM RIGIDUM.

RIGID SHIELD-FERN.

(Plate 3, fig. 5.)

CHA.—Frond bipinnate. Pinnæ alternate. Lobes oblong, decurrent, tridentate. Rachis scaly.

SYN.—*Aspidium rigidum*, *Hook.* in *Bri. Flo.*, ed. 3 and 4, *Swezt.*, *Schk.*—*Aspidium spinulosum*, *Hook.* in *Bri. Flo.*, ed. 1.—*Polypodium rigidum*, *Hoffm.*—*Polystichum rigidum*, *Decan.*—*Polystichum strigosum*, *Roth.*—*Lastrea rigida*, *Presl*, *Newm.*

FIG.—*E. B. supp.* 2724.—*Schk. fil. t.* 38.—*Newm. page* 56.

DES.—Root tufted. Rachis thick, rigid, very scaly all the way up. Frond lanceolate, not contracted below, erect, from one to two feet high. Pinnæ tapering, alternate, very close together, from thirty to forty pairs, their stipes very much thickened at their union with the rachis. Lobes distinct, decurrent, oblong, blunt, tridentate, but *not spinulose*, their midrib waved. Sori large and abundant, chiefly on the upper part of the frond. Indusium round reniform, persistent, with a glandular margin, white at first, lead-colored afterwards, covering the whole mass of thecae, &c.

Much diversity of opinion has existed respecting the identity of this very distinct plant, a small state of the *spinulosum* being very often sent for it. Its generally *alternate* pinnæ would be perhaps sufficient to distinguish the two, but in other respects it differs essentially from that more common species. The rachis of the *rigidum* is very scaly and very much thicker than in the *spinulosum*, its pinnæ much more numerous and nearer together, the lower pair not broader than the rest, the lobes of all quite decurrent, and not by any means *spinulose*, besides which the indusia are very large, and so different, as at once to distinguish the two plants; in addition to which it may be remarked, that *Aspidium rigidum* is much darker in color than the *spinulosum*, as it is also than the *cristatum*. It is intermediate between the last and next species in the number of its divisions, but does not resemble either of them in habit or appearance. *Mr. Newman*, and the late Professor *Don*, both support me in the identity of this species with the *Aspidium rigidum* of *Schkuhr*, specimens from whom I have seen. *Mr. Newman* justly remarks that, "when cultivated, it assumes a more diffuse and lax appearance, and is not so like *Schkuhr's* figure as the plant from *Settle*."

HAB.—Found by *Rev. W. Bree*, in 1815, on Ingleborough, on a natural platform, near the foot of the mountain, and towards the neighbouring village. This was, I believe, the only situation recorded for this fern, at the publication of my first edition in 1837. Since then it has been sought after and found in three or four places, considerably distant from each other; and there is reason to suppose that it is pretty generally distributed all over the Ingleborough range, towards the foot of the hills. Thus *Mr. W. Wilson* finds it at Wharnside. *Mr. Chorley* has kindly communicated to me specimens from near Settle, where he and *Mr. J. Tatham* find it abundantly. Also other fronds of the true plant have reached me from *Miss Beever*, a young and enthusiastic botanist, who finds it at Arnside Knot, not far from Silverdale.

GEO.—Switzerland, Prussia, Germany, &c.

8.—ASPIDIUM SPINULOSUM.

PRICKLY SHIELD-FERN.

(Plate 3, fig. 6.)

CHA.—Frond bipinnate. Pinnæ opposite. Lobes finely cut, spinulose. Rachis nearly smooth, white.

SYN.—*Aspidium spinulosum*, Willd.—*Polypodium spinulosum*, Swz., Retz.—*Polypodium cristatum*, Hoffm., Schreb.—*Polypodium spinosum*, Schr.—*Polypodium dentatum*, Moench.

FIG.—*E. B.* 1460.—*Flo. Dan.* 707.—*Pluk. Phyt.* 181, f. 2, (*a young plant*,) *Schk. fil.* 48.

DES.—Frond ovate or oblong, always erect and flat. Pinnæ very nearly opposite, smooth, and distinct, as are also the lobes, which are rarely convex. Segments oblong, pointed, doubly serrate, and spinulose. Rachis nearly smooth, swelled at its ramifications, of a whitish color, and generally covered with black dots. Sori scattered, small. Indusium small, brown, soon shrivelling up.

This plant goes by various names among British botanists. It is repeatedly considered and sent as *Aspidium cristatum*, (which see, page 39,) and is such of some authors, but not of Smith, Hooker, or Mackay. It is also confounded with the much rarer *Aspidium rigidum*, the diagnostics of which are very distinct; and with the next species, *Aspidium dilatatum*, it is often considered identical, though sufficiently different, both wild and cultivated, in habit, texture, and color. Our present plant is narrower than the *dilatatum*, of a less number of pinnæ, flat, erect, rigid in habit, of a very light green color, the midrib of the lobes more zigzag and prominent, the lower pinnæ rarely twice pinnate, the indusium glandulous, and the whole plant much more delicate.

It should be observed, that the above remarks are not intended to apply to that plant which Sir J. E. Smith's herbarium contains, and which Sir W. J. Hooker describes as a variety of *dilatatum*, under the above name. The spinulosum of northern botanists, of Sir J. E. Smith, and of the Liverpool Botanic Garden, is, in reality, but a variety of the next, and closely approaches to the recurvum of Bree, and dumetorum of Smith, if not identical with them. The plant here intended to be described is altogether different, and in cultivation retains precisely the character of the wild plant, never approaching in the most remote degree the

Aspidium dilatatum, though the mountain form of this latter plant has the lower pinnae much abbreviated.

In a variety of *spinulosum* given me by Mr. J. Merrick, of Manchester, the lobes on the *upper* side of each pinna are much larger than those on the lower; also, it may be remarked, that in dry situations the lobes will become convex, but this is by no means common.

SIT.—On wet moors, sides of pools and ponds, wet hedge-rows, &c.

HAB.—SCO.: Moray and Rosshire, *Rev. G. Gordon*. Aberdeenshire, *Dr. Murray*. Dumbartonshire, *Mr. J. Hooker*. Auchindenny Woods, Edinburgh, *Mr. Watson*. Isle of Man, *Mr. Forbes*. Near Richmond, Yorks., *Mr. J. Ward*. Ingleborough, Yorkshire, *Rev. W. Bree*. In a small state at Woolston Moss, Lanc., and Newchurch Bog, near Over, Cheshire, *Mr. W. Wilson*. Titterstone Clee Hills, Shrops., *Mr. J. S. Bayly*. Bomere Pool, Salop, *Mr. C. Babington*. Warwicksh., *Rev. W. S. Bree*. Derbys., *Dr. Howitt*. Pottery Car, near Doncaster, *Mr. S. Appleby*. Dallington Heath, near Northampton, *Mr. Anderson*. Norfolk, *Miss Bell*. Near the Windmill, and near the Spring-well, on Wimbledon Common, *Mr. W. Pamplin*. Barnes Common, Surrey (near the Water-house), *Mr. Castles*. Abundant in Essex, *Mr. J. Bevis*. Common in Kent, *Mr. W. Pamplin*. Tonbridge, Kent, *Mr. W. C. Trevelyan*. Sussex and S. Kent, *Rev. G. E. Smith*. Wood near Dunsford Bridge, Devon, *Mr. Jacob*. Near Torquay, *Dr. Greville*.—WAL.: Aber, Caern., *Mr. Leighton*. Near Wrexham, Denb., *Mr. J. E. Bowman*. *Note*.—I cannot say whether the Scotch and Welsh habitats refer to the above plant, or to the *Spinulosum* of the Br. Fl.

GEO.—Switzerland, Dauphiny, Salzburg, Darmstad, and North America.

9.—ASPIDIUM DILATATUM.

GREAT SHIELD FERN. DILATED SHIELD FERN.

(Plate 3, fig. 7.)

CHA.—Frond tripinnate, triangular. Pinnæ opposite, lobes deeply dentate, spinulose, petioled. Rachis scaly.

SYN.—*Aspidium dilatatum*, *Willd.* *Spreng.*, *Forst.*, *Galp.*, *Gray*.—*Aspidium spinulosum*, *Swz.*, *Sibth.*, *Hook.* (*not a*), *Mack.*, *Schk.*—*Polypodium cristatum*, *With.*, *Bolt.*, *Huds.*, *Ehrh.*, *Moench.*, *Lightf.*—*Polypodium dilatatum*, *Hoffm.*, *Mull.*—*Polystichum multiflorum*, *Roth.* *Lastrea dilatata*, *Presl.* *Newm.*

FRG.—*E. B.* 1461.—*Bolt.* 23.—*Schk. fil.* 47. *Newm. p.* 59. 61.

DES.—Root black, tufted. Frond tripinnate, triangular, from a few inches to 2 feet high, dark green, and drooping. Pinnæ opposite, smooth, oblong, obtuse, pinnate, except the lower pair which are doubly pinnate. Lobes ovate, pointed, convex, deeply but irregularly serrated and spinulose, petioled, their midribs straight. Rachis covered with broad, brown scales. Sori all the summer, distinct. Indusiums soon becoming obliterated, round, with a lateral notch.

A very variable plant, altered much by cultivation and circumstances; thus if it grow in a situation which is wet in the spring and dried up in the summer, as on the margin of a pond, it will become var. β , very dark, large, and quite

drooping. Continued wet will elongate the frond and separate the pinnæ and lobes as in var. γ . A young plant is only twice pinnate and flat. A dry and rocky, or a confined situation will render the frond small and less divided, the lobes blunt, deflexed, and drooping: thus starved it becomes the *Aspidium dumetorum* of Smith (var. δ). I know not the nature of the habitats in which the recurved var. (ϵ) of Bree grows, and can only regret that botanists do not record the circumstances, as well as the places, in which plants are found. The varieties *recurvum* and *dumetorum* are, I believe, not altered by cultivation, and Sir J. E. Smith implies, in his description of the latter, that its spores produce the same variety.

α (*dilatatum*) Frond sub-tripinnate, triangular, ovate. Pinnules petioled.

β (————) Frond tripinnate, deflexed, triangular. Pinnules convex.

γ (————) Frond tripinnate, triangular, elongated. Pinnules somewhat decurrent, and distant from each other.

δ (*dumetorum*) Frond small, triangular, drooping. Pinnules blunt.

ϵ (*recurvum*, Bree.) Frond small. Pinnules concave, and dark green.

Newm., p. 61.

SIT. and HAB.— $\alpha \beta \gamma$. Very common in damp hedge-rows and swampy woods, ascending to an elevation of 1000 yards in many parts of the Highlands, and probably even to 1200 yards on the Cairngorum range, Mr. H. C. Watson.— δ . : Derbyshire (rare), Mr. J. E. Bowman and Dr. Howitt. Common about Settle, Yorks., Mr. J. Tatham. Black Rock, Cromford, Derb., G. F. Ben-na-Baird, Aberdeensh., Mr. H. C. Watson. Powercourt Waterfall, and side of Djouce Mountain, Ireland (abundant), Mr. Mackay. ϵ . Plentiful about Penzance, Cornwall, Rev. W. Bree.

GEO.—Common throughout Europe. and from Pennsylvania to Virginia.

The glory of the British Highlands - very fine in cultivation -

ASPLENIUM. Linn. SPLEENWORT.

(ασπληγον, a medicine to cure disorders of the spleen, from α and σπλην.)

A, part of the frond of *Asplenium marinum*. One pinnule, showing the veins and origin of the fruit, the others the sori in different states. B, part of a pinnule magnified. C, the same cut transversely. D, under eutete. E, transverse section of the stem. F, indusium. G, theca and spore. H, young plant.

Sori linear at first, afterwards oblong; indusium linear, attached to a transverse vein, and opening on the opposite part of the sorus towards the central nerve of the pinna. This is a well marked and extensive genus, of which Sprengel enumerates no less than 151 species; of these ten only are British, which are very little or not at all altered by culture, they are therefore less liable to run into varieties than some other genera. It is only when the sori are in a young state that many species can be known to belong to this genus, as the indusia are so delicate that they are soon lost among the sori, which in many of the smaller species at last appear like round or oblong spots.

1.—ASPLENIUM SEPTENTRIONALE.

FORKED SPLEENWORT.

(Plate 4, fig. 1.)

CHA.—Frond simply partite. Segments linear, sharply toothed at their extremity.

SYN.—*Asplenium septentrionale*, Swz., Willd., Hull, Hoffm., Hook., Smith, Galp., Gray.—*Aerostichum septentrionale*, Linn., Bolt., Dicks., Ehrh., With., Huds., Lightf.—*Scelopendrium septentrionale*, Roth.

FIG.—*E. B.* 1017.—*Flo. Dan.* 60.—*Bolt.* 8.—*Flo. Lon.* 162.—*Ger.* 1561.—*Newm.*, p. 73.

DES.—Fronds very numerous, upright when young, drooping afterwards, rigid, 1 to 3 inches high, cleft near the top into two or three linear sharp-pointed alternate segments, which in proportion to their size are furnished at or near their extremity with from one to three acute, but not spinous teeth. Sori one on each side of the segment, nearly longitudinal, concealed at first by a white indusium, attached at the outer edge; afterwards the swelling sori throw back the indusia, covering the whole of the segment, and finally curving and contorting it in a curious manner.

Mr. H. C. Watson writes thus: "Although quite a northern fern I observed the young fronds destroyed by a frost of 25 degrees Fahr. in April, 1835. The plant had been under a glass in a cold frame during winter, where the temperature inside had risen a few degrees higher than outside by day, and had never been allowed to sink to the freezing point at night, in consequence of a thick covering of mats."

SIT.—On the rocky clefts of mountains, chiefly in the north. Not in Ireland.

HAB.—ENG.: In situations probably exceeding 1000 feet in height in Cumberland, where it occurs sparingly on rocks, between the vale of Newlands and Borrodale.—WAL.: Craig Ddw. (a mile above Llanberris Church), Caern., Mr. C. C. Babington. Snowdon, (rare), Mr. J. E. Bowman. Llyn-y-ewm, N. Wales, Mr. W. Wilson.—SEO.: Arthur's Seat, Edinburgh, (above the rail-road), Jedburg, &c., Mr. H. C. Watson. Blackford Hill, Edin., Mr. W. Brand.

GEO.—Holland, Switzerland. Not unfrequent throughout Europe.

1856. - Abundant on Wetten Rocks in Valley of the Aar near Basle, Germany. C. E. & Co.

2.—ASPLENIUM ALTERNIFOLIUM.

ALTERNATE-LEAVED SPLEENWORT.

(Plate 4, fig. 2.)

CHA.—Frond pinnate. Pinnæ alternate, wedge-shaped, notched.

SYN.—*Asplenium alternifolium*, Smith, Hook., Dicks., With., Galp., Jacq.
—*Asplenium germanicum*, Willd., Hoffm., Ehrh., Lam., Weis., Gray.
—*Phyllitis heterophylla*, Moench.—*Scelopendrium alternifolium*, Roth.
—*Asplenium Breyonii*, Retz.

FIG.—*E. B.* 2258.—*Jacq. Misc. t. 5. f. 2.*—*Brey. Cent.* 11. 91.—*Newm.* p. 71. β β .

1

2

3

4

5

6

DES.—Fronde from 1 to 3 inches high, very light green, upright, delicate, about half covered with pinnæ, which are alternate and wedge-shaped; the larger partly three-cleft, the smaller bluntly notched at the end only. Rachis dark at the base only. Sori two to four on each pinna, small, light brown, becoming confluent, but not occupying the whole under surface. Indusium entire on the margin.

This species is intermediate between the last and *Asplenium Ruta-muraria*, although of a more delicate and erect habit than either; its color also is much lighter and its sori smaller and less confluent. When once seen it cannot possibly be mistaken for any of the numerous casual forms of *Asplenium Ruta-muraria*.

The plants sold under this name in the nurseries around London are the true species derived from some plants brought wild from Scotland, about 14 years ago, and given to the Countess De Vande, at Bayswater, and from her garden distributed around. It quite retains its character in cultivation.

HAB.—Found originally by Mr. Dickson on rocks in the south of Scotland, two miles from Kelso on the Tweed. Now existing at Dunkeld, in Perthshire, Mr. Bishop and Dr. Macnab. Very sparingly near Dunfermline, Fifeshire, Dr. Dewar.

GEO.—Germany, Sweden, and Switzerland, where it is quite an alpine plant.

3.—ASPLENIUM RUTA-MURARIA.

WALL RUE. RUE-LEAVED SPLEENWORT. TENTWORT.

(Plate 4, fig. 3.)

CHA.—Fronde bipinnate. Pinnæ alternate. Pinnules ovate or wedge-shaped, with rounded notched extremities.

SYN.—*Asplenium Ruta-muraria*, Linn., Willd., Hook., Smith, Bolt., Bull., Ehrh., Huds., With., Galp., Lightf.—*Asplenium murale*, Bernh., Gray. *Scolopendrium Ruta-muraria*, Roth.—*Phyllitis Ruta-muraria*, Moench. *Ruta-muraria*, Bau., Ray., Ger., Plum., Newm.

FIG.—E. B. 150.—Bolt. 16.—Fl. Dan. 190.—Bull., Fr. 195.—Plum. Fil. t. A. f. 3.—Newm. Brit. Ferns., p. 71. not β. β.

DES.—Root tufted, black, very long. Frond from 1 to 4 inches high, dull green. Rachis green, except at the very base. Pinnæ confined to the upper half, from three to five or six in number, placed alternately, for the most part distinctly three cleft. Pinnules ovate in small fronds, wedge-shaped in the larger; their tips rounded and erenate, or unequally notched. Barren fronds broader and shorter. Sori dark brown, finally confluent, and covering the whole under surface. Indusium uneven at the margin.

SIT.—On walls, ruins, rocks, and other similar situations.

HAB.—Very generally distributed over the United Kingdom, though there are a few districts where it is scarcely found. I believe Berwickshire is one of these, nor is it by any means plentiful in Norfolk or Suffolk.

GEO.—Most parts of Europe, and from New York to Carolina in America.

4.—ASPLENIUM MARINUM.

SEA SPLEENWORT.

(Plate 4, fig. 4.)

CHA.—Frond oblong, pinnate. Pinnæ obtuse, serrate, slightly auricled above. Rachis winged.

SYN.—*Asplenium marinum*, Linn., Willd., Huds., Bolt., Dicks., Lightf., With., Galp., Smith, Hook, Maek., Gray, Newm., &c.

FIG.—*E. B.* 392.—*Lob. Ic.* 814.—*Fl. Lon.* 60.—*Bolt.* 15.—*Ger.* 1143—*Newm.* 75.

DES.—Root very thickly tufted, black, with stout fibres. Frond 6 to 9 inches high, pinnate, irregularly oblong, obtuse. Rachis winged all the way down, black, shining, smooth, without pinnæ at the lower part, above bearing about twenty on each side, mostly alternate, obtuse, about an inch long in the middle of the frond, running at the base into the wing of the rachis, therefore slightly decurrent; the upper side of each generally auricled, the lower side proportionably truncated. Sori large, transverse, at first linear, then oblong, but never confluent. Indusium white or of a pale brown.

SIT.—Upon maritime rocks, or in caves by the sea side, and in one or two inland situations.

HAB.—ENG: Marsden Rocks, Durham, *Mr. R. B. Bowman*. Isle of Man, *Mr. Forbes*. Above the Black Rocks at the entrance of the Mersey, (Cheshire side,) *Mr. H. C. Watson*. Liverpool, (near the Dingle,) *Mr. Merrick*. Still at Hulme Stone Quarry, (otherwise called Winwick Stone Delph,) near Warrington, where Bolton gathered it, (*v. Bolt. Fil. loc. cit.*) *Mr. W. Wilson*. In this place Mr. Shaw, of Bollington, many years ago found a curious variety, with a much more divided frond than is usual, and which remains with him distinct in culture. Sussex, *Mr. Borrer*. West of Cornwall, *Professor Henslow*.—WAL.: Anglesea, *Mr. J. E. Bowman*. Near the South Stack Light-house, Holyhead, *Mr. C. C. Babington*. Ormeshead, and near Bangor, *Mr. W. Wilson*.—SCO.: Parish of Nigg, Rosshire, *Mr. Briehan*. Near Port Patrick, Wigtownshire, *Dr. Balfour*. Moray, *Rev. G. Gordon*. Isle of Staffa, *Mr. J. Dovaston*. Near Eyemouth, Berwickshire, *Rev. A. Baird*. Frequent on the whole line of the Berwickshire coast, *Dr. G. Johnston*. Fife and Aberdeenshire, (common,) *Dr. Murray*. Isle of Arran, *Mr. T. H. Cooper*.—IRE.: Sutton side of Howth Mountain, Underwood, Killiney Hill, &c., *Dr. Osborne*. Derrinane, county Kerry, *Mr. Kelly*. Abundant on the southern and western coasts, *Mr. Maekay*.

GEO.—Barbary, Canary Islands, Spain, St. Helena, West Indies, Islands of the Archipelago, &c. There is but little difference between our plant and Dr. Hooker's species *Asplenium obtusatum*.

None at Western Cape-More - abundant at Cape Town -

5.—ASPLENIUM TRICHOMANES.

COMMON MAIDEN-HAIR SPLEENWORT. WALL SPLEENWORT.

(Plate 4, fig. 5.)

CHA.—Frond pinnate, linear. Pinnæ subrotund, crenate. Rachis black.

SYN.—*Asplenium trichomanes*, Linn., Willd., Michx., Woodv., Bolt., Dicks., Ehrh., Lightf., Smith, Hook., With., Spreng., Huds.—*Asplenium saxatile*, Salisb., Gray.—*Asplenium trichomanoides*, Schkr., (not Michx.)—*Asplenium melanocaulon*, Willd., Pursh.—*Trichomanes*, Ray, Fuchs., Tillands., Bauh., Plum., Park.—*Phyllitis rotundifolia*, Moench., Newm.

FIG.—*E. B.* 576.—*Flo. Lon.* 156.—*Bolt.* 13.—*Flo. Dan.* 119.—*Woodv.* 201.—*Ger.* 1146.—*Plum. t. B. f.* 1.—*Newm.*, p. 80.

DES.—Fronds tufted, linear, pinnate, 2 to 4 inches high, dark green, very rigid, quite smooth, with a purplish-black shining rachis, channelled in front. Pinnæ from twenty to thirty pairs, opposite or alternate, (generally the former,) obtuse, crenate, of a round or oval form, very distinct from each other all the way up, and sessile, or very nearly so. Sori two to six on each pinna, placed transversely, very dark colored, finally confluent, often covering the whole under surface. *Abundant at Brigs Crag, & around Inverness.*

HAB.—Common on rocks, old walls, &c., in most parts of the United Kingdom; not only on the main land, but the Isles of Anglesca, Man, Wight, Sheppy, and the Channel Isles, yet by no means frequent in the N and NE. of Scotland.

GEO.—Throughout Europe. In Jamaica. In Japan and other parts of Asia. Canada, Pennsylvania, and high mountains of Carolina.

6.—ASPLENIUM VIRIDE.

GREEN MAIDEN-HAIR SPLEENWORT. GREEN RIBBED SPLEENWORT.

(Plate 4, fig. 6.)

CHA.—Frond pinnate, linear. Pinnæ roundish-deltoid, crenate. Rachis green.

SYN.—*Asplenium viride*, Huds., Willd., Roth., Dicks., Ehrh., Bolt., Smith, Hook., With., Spreng., Galp., Lightf., Gray, Newm.

FIG.—*E. B.* 2257.—*Bolt.* 14.—*Flo. Dan.* 1289.—*Pluk. Phy.* 89, f. 6.—*Newm.* 78.

DES.—Fronds numerous, pinnate, linear, from 3 to 6 inches high, of a very light green color. Pinnæ petioled, alternate, the upper ones ovate, the lower roundly triangular, attached to the rachis by the centre of one of the sides, which is somewhat truncate, the other two sides being regularly and deeply crenate, sometimes doubly so. Rachis quite green, except at the lower part. Sori reddish brown, two to six on each pinna, confined to the middle of it, finally becoming confluent, but even then not extending to the margin.

This is immediately distinguished from the last by the lighter color of all its parts, its less spreading sori, and differently-shaped and alternate pinnæ; added to which, the pinnæ on the lower part of the frond are generally distant, and those near the top of the frond crowded, while the whole is much more delicate and elegant. Sometimes the frond is divided into two, as represented in Bolton, t. 2, f. 3, when it becomes the *Trichomanes ramosum* of authors; but this branching is an accidental circumstance, and by no means constant; it therefore

does not constitute a variety, more especially as not more than two or three branched fronds are found upon a plant, all the rest being of the common character and appearance.

SIT.—On rocks, not farther south than Yorkshire, or perhaps Derbyshire.—ENG.: On rocks in Northumberland, *Mr. Winch*. Mazebeck Scars, Westmorl., and Gordale, Yorkshire, *Mr. R. Bowman*. Near Halifax, Yorkshire, *Mr. R. Leylands*. Near Ais-la-Beek, and Richmond, Yorks., *Mr. J. Ward*. Settle, *Mr. Chorley*.—WAL.: Cader Idris, *Mr. J. E. Bowman*. Snowdon, *Mr. C. C. Babington*. Twll. Du, Caern., *Mr. T. H. Cooper*. Not uncommon on the Welch mountains, *Mr. W. Wilson*.—SCO.: Rosshire, *Rev. G. Gordon*. Cawder Woods, Nairns, *Mr. W. Staples*. Base of Benmore, Sutherlandsh., *Dr. Johnston*. Far too common in the Highlands to need the specifying of stations, *Mr. H. C. Watson*.—IRE.: Turk Mountain, Killarney, Ben Bulbin, county of Sligo; and on the Donegal Mountains, near Lough Eske, *Mr. Mackay*.

GEO.—Germany, Holland, Switzerland, France; very rare, except on the mountains of Tyrol and Carinthia.

1867—very luxuriant on the Rocky sides of a torrent between Glion & les Aumonts, Vaud.

7.—ASPLENIUM FONTANUM.

SMOOTH ROCK SPLEENWORT.

(Plate 5, fig. 1.)

x CHA.—Frond bipinnate. Pinnæ oblong, blunt, alternate. Pinnules wedge-shaped, cleft, and toothed. Rachis winged.

SYN.—*Asplenium fontanum*, *Hook.*, *Smith*, *Bernh.*—*Aspidium fontanum*, *Willd.*, *Suez.*, (not of *Schkr.*)—*Polypodium fontanum*, *Linn.*, *Huds.*, *Bolt.*, *With.*—*Athyrium fontanum*, *Gray*.

FIG.—*E. B.* 2024.—*Lob. Ic.* 810, 1.—*Bolt.* 21, (*bad.*)—*Newm.*, page 4.

DES.—Root tufted, long, black. Frond lanceolate, bipinnate, evergreen, 2 to 6 inches high. Rachis winged throughout. Pinnæ alternate, ovate, oblong, those in the middle of the frond from a quarter to half an inch long, formed of six or eight pinnules placed alternately. Pinnules short, broadly wedge-shaped, petioled, so very deeply cleft at the sides and toothed at the apex as to become nearly pinnate. Seldom more than two sori upon each pinnule, which soon extend over the whole surface of it.

Our present species most resembles *Asplenium lanceolatum*, the shape of the frond being nearly the same. The fontanum, however, is much more delicate, and smaller in all its parts, of a very dark green color, its pinnules not half the size, and of a very different shape to those of the lanceolatum, besides which its winged rachis is of itself a sufficient diagnostic. It is very much more difficult to distinguish it from *Asplenium Halleri*, a species that is very rare on the Continent, and for which our fontanum is very generally sold.

HAB.—Supposed to be now extinct in England; it was once found on Amer-sham Church, in Buckinghamshire, and at Wybourn, in Westmoreland. I have been informed that living plants were found at a waterfall in either Northumberland or Westmoreland, 14 or 16 years ago, and also that it once grew on Ahwick Castle; but if so, it is no longer found there.

GEO.—Saxony, Switzerland, South Europe, and Sicily.

x This *Asplenium Halleri* are unquestionably the same, & the latter name should be adopted—1867 Feb 25 & 26. Found by me abundantly on the Rocks of Mt D'Arrel, above Roche Canton de Vaud. (S. F.)
Also in the Gorge of Montreux, & on Mt Sonchaud, above Chillon.

1

2

3

4

5

6

8.—ASPLENIUM LANCEOLATUM.

LANCEOLATE SPLEENWORT.

(Plate 5, fig. 2.)

CHA.—Frond lanceolate, bipinnate. Pinnæ and pinnules obovate, sharply toothed at the apex.

SYN.—Asplenium lanceolatum, *Huds., Swz., Hoffm., Willd., Smith, Hook., Forst., With., Galp., Gray.*—*Phyllitis lancifolia, Moench.*

FIG.—*E. B.* 240.—*Ger. Herb.* 1135.—*Newm., page 66.*

DES.—Frond lanceolate, bipinnate, from 3 to 6 inches high, upright in habit, and of a light green color. Rachis green, minutely hairy, not winged, void of pinnæ below. ⁺Pinnæ opposite, from twelve to twenty pairs, the lower pair short, distant from the next, and often slightly drooping. *Pinnules ovate, sharply serrated and pointed, the smaller confluent, the larger petioled and tapering at the base, particularly that on the upper side next the rachis. Sori light brown, one or two near the middle of each lobe, at first linear, afterwards round, but very rarely or never covering the whole under surface. ⁺*Consider this ~~opposite~~ character of no value; the pinnæ being as frequently alternate as opposite.*

† This Fern has been repeatedly confounded with *Asplenium Adiantum nigrum*, though there is a very great dissimilarity between them; our present species is of a different shape, color, size, and habit, its divisions less numerous, the naked part of its stem shorter, and its sori less extended: in fact they vary in almost every particular.

SIT.—On rocks, &c. in the south of England, and in Wales.

HAB.—On the walls of the Church of St. Sanceret, near the Land's End, Cornwall, *Jones's Tour.* Abundant around Penzance and St. Ives, *Mr. H. C. Watson.* Scilly Islands, *Mr. W. C. Trevelyan.* Sussex, *Mr. Borrer.* High rocks, near Tunbridge Wells, (1835,) *Mr. W. Pamplin.* Near Barmouth, (plentiful,) *Mr. J. E. Bowman* and *Mr. W. Wilson.*

GEO.—Azores, Bohemia, Hungary, France.

July - about Penzance, G. E. B. - E. B. - in the Woods at Stapleton - Gloucestershire
also on the permanent rocks

9.—ASPLENIUM ADIANTUM NIGRUM.

BLACK MAIDEN-HAIR. SHINING SPLEENWORT.

(Plate 5, fig. 3)

CHA.—Frond tripinnate, subdeltoid. Pinnæ alternate. Pinnules inciso-serrate, blunt. Rachis winged, black.

SYN.—*Asplenium Adiantum nigrum, Linn., Willd., Smith, Hook., Mack., Bolt., Roth, Huds., With., Galp., Bernh., Lightf.*—*Asplenium lucidum, Gray, Salisb.*—Black spleenwort, *Newm.*

FIG.—*E. B.* 1950.—*Flo. Dan.* 250.—*Bolt.* 17.—*Ger.* 1137.—*Newm., page 68.*

sp. lanceolatum, - the pinnæ are attached to the Rachis at right Angles - but in A. Adiantum nigrum the pinnæ form an acute Angle with the Rachis -

DES.—Fronde tripinnate, ovate or deltoid, 4 to 8 inches high, dark green, rigid, and erect. Rachis black, smooth, slightly winged, clothed with pinnae only on the upper half. Pinnae alternate, those only on the lower part twice pinnate, the lowermost the largest. Pinnules deeply cleft, tapering at their base, sharply serrated at and near the top. Sori linear at first, round at last, covering the whole under surface of the frond.

α Fronds rigid, tripinnate only at the lower part. (*The common plant.*)

β Fronds delicate, tripinnate throughout. (*Not β of Smith.*)

Sir J. E. Smith, in conformity with the old authors, makes another variety, differing only from the common plant in having long fronds and distant pinnae; but I leave any one to say if it be anything more than a drawn up plant of the common species, found as it was, solitary, in a dark cave.

HAB.—α.: Common through the United Kingdom, on walls, rocks, &c. I have habitats from the Orkney Islands, and from those in the English Channel, from the eastern as well as from the western counties, from Wales, Scotland, and Ireland. Dr. Murray writes me, "Not common in the north of Scotland." —β.: Limestone rocks at Mueruss, Killarney, Mr. Mackay, Miss Hutchins, and Dr. Taylor. Mount Cahir-Courcee, six miles from Tralee, Mr. W. Andrews.

GEO.—Italy, France, Germany, Madeira, and high mountains of Carolina.

Bristington ~~near~~ about the World's Rock, Weston super Mare, a beautiful

10.—ASPLENIUM FILIX-FŒMINA.

LADY FERN.

(Plate 5, fig. 4.)

CHA.—Fronde broadly lanceolate, bipinnate. Pinnae tapering, pointed. Pinnules oblong, inciso-serrate. Rachis smooth.

SYN.—*Asplenium Filix-fœmina*, Hook., Mack., Spreng., Bernh.—*Aspidium Filix-fœmina*, Swz., Willd., Smith, Hook. in *Fl. Sco.*, Galp.—*Polypodium Filix-fœmina*, Linn., Lightf., Hud., Bolt., Dicks., With.—*Polypod. ovato-crenatum*, Hoffm.—*Athyrium Fil-fœm.*, Roth, Decan., Presl., Neum.

FIG.—*E. B.* 1459.—*Flo. Dan.* 1346.—*Bolt.* 25.—*Phuk. Phyt.* 180, f. 4. *Newm.*, p. 63.

DES.—Root large, tufted. Rachis without scales, green (rarely purple), the naked part very short. Frond bipinnate, broadly lanceolate, long-pointed, and tapering at the base, 12 to 20 inches high, dark green, very delicate in habit, often recurved. Pinnae alternate, from twenty to forty pairs, oblong, tapering gradually to a point, the lower ones sometimes drooping. Pinnules very numerous, oblong, rather blunt, pinnatifid, or inciso-serrate, the serratures minutely toothed, but not aristate, the lower pair close to and parallel with the rachis. Sori solitary, near the base of the lobes, at first linear-reniform, at length round, but not confluent. Indusium jagged, white, oblong or reniform.

β Rachis red and somewhat scaly. (This is the character the plant bears in Switzerland.)

γ (*Aspid. irriguum*, Sm.) Frond narrow, pinnæ distant, deeply cleft.

δ Frond broad and small, pinnæ and pinnules short and few, nearly white.

All the varieties of this Fern are so very tender (particularly the var. γ), that they shrivel up and become withered almost immediately upon being gathered. Under the name of *Aspidium irriguum*, I have received fronds (without fruit) of very different habit, marked γ and δ, neither of them by any means a distinct species, perhaps not even a constant variety, as the former appears to me rather a plant drawn up either by a confined situation or excess of moisture, while the other is perhaps a young plant only, and its very light color an adventitious circumstance. The beauty of this common plant occasioned its name of Lady Fern, contrasting as it does with the robust habit of *Filix-mas* or Male Fern.

SIR.—Its natural habitation is swampy woods and damp hedge-rows; or, as Sir Walter Scott incidentally remarks in his novel of “Waverley,”—

“Where the copse-wood is the greenest,
Where the fountain glistens sheenest,
Where the morning dew lies longest,
There the Lady Fern grows strongest.”

HAB.—Pretty freely distributed over the southern and midland counties of England and Ireland, though it is by no means abundant in North Wales or North Scotland, except in particular neighbourhoods.—β: Frequent in moist woods in Kent, *Mr. W. Pamplin*.—γ: Ruberslaw, Jedburgh; Aber, Caern.; and near the English Bridge, Shrewsbury, *Mr. Leighton*. Marsh at Mucruss, Killarney, *Mr. Mackay*. In some boggy woods belonging to Eridge Park, Tunbridge Wells, (1835), *Mr. W. Pamplin*.—δ: Prestwich Carr, near Manchester, *Mr. Merrick*, who gave me a specimen, (5 inches high.)

GEO.—Throughout Europe, and from Canada to Virginia, in North America.

Abundant & beautiful in Kew's Ram. Ferns, & in The Ceylon Woods.

SCOLOPENDRIUM, Swz. HART'S-TONGUE.

A, portion of a frond of *Scolopendrium vulgare*, showing the origin of the fruit from lateral veins, and with its ordinary appearance. B, transverse section of the twin masses of fruit, with their folding indusiums while in a young state. C, ripened fruit, in which the sori have become confluent, and thrown back the covers. D, theca and spores. E, theca opened. F, transverse section of the rachis.

The sorus of this small genus appears to have two indusiums, at first folded over each other, and afterwards thrown back in contrary directions; but in fact the sorus itself is no less double, two of them growing together so closely as to form in appearance but one mass; this is transverse, and seated between those lateral veins to which the two covers are attached.

SCOLOPENDRIUM VULGARE.

COMMON HART'S-TONGUE.

(Plate 5, fig. 5.)

CHA.—Frond ligulate, acute, entire, cordate at the base. Rachis sealy.

SYN.—*Scolopendrium vulgare*, *Smith*, *Hook.*, *Spreng.*, *Mack.*, *Gray.*—*Asplenium scolopendrium*, *Linn.*, *Huds.*, *Bolt.*, *Woodw.*, *Ehrh.*—*Asplenium elongatum*, *Salisb.*—*Scolopendrium officinarum*, *Swz.*, *Willd.*, *Pursh.*

FIG.—*E. B.* 1150.—*Bolt.* 11.—*Flo. Lon.* 67.—*Ger.* 1138.—*Schk. fl.* 83.

DES.—Root tufted. Fronds numerous, a foot high, strap-shaped, pointed, the base of them heart-shaped, smooth, except the lower part of the rachis and sometimes the midrib, which are very sealy. Sori attached to oblique transverse veins, always in twin united masses, each having its cover attached; the one at the upper side, the other at the lower, and when young folding over each other in the middle. The sori are oblong, distant from each other, and chiefly at the upper part of the frond.

This plant is very apt to become differently cleft and crisped, remaining so under cultivation, and bearing fruit copiously in that state; hence the following varieties are noticed.

α (*vulgare*). Frond ligulate, flat and single pointed.

β (*crispum*). Frond crisped and curled along the margins.

γ (*multifidum*). Frond much cleft at the top.

ϵ (*linearis*). Frond very long and narrow.

The above, except β , can scarcely be considered distinct varieties, but should rather be accounted monstrosities, particularly γ , which is produced by over abundance of food and warmth; thus if the common state of the plant be transplanted to rich soil in a green-house, it will rarely ever fail to produce fronds cleft more or less towards the apex. These same plants, if again thrown out and neglected, will return to their original state. The spores of all the varieties will produce the common plant.

Sir J. E. Smith remarks, that "the whole plant has a nauseous scent when bruised, and is of a mucilaginous and acid taste." It is now discarded from the regular practice of medicine, but frequently still sold in our herb shops, being used as an ointment for burns, &c., and taken internally as an astringent.

SIT.—In damp ruins, rocks, wells, &c.

HAB.—Scarcely a common Fern, though abundant in some places, particularly in the south and west of both England and Ireland, but decreasing in quantity northwards. Isle of Man, *Mr. E. Forbes*. Near Braunston, Leicestershire (rare), *Rev. A. Bloxam*. In Wagg Lane, Congleton, Cheshire; also at Buxton, Matlock, and Dove Dale, Derbyshire, *Mr. H. C. Watson*. Near Leeds, *Mr. H. Denny*. Near Richmond and Settle, Yorkshire, *Mr. J. Tatham*. Three varieties on Pottery Car, near Doncaster, *Mr. Appleby*. Hawkstone, Salop, *Mr. J. S. Bayly*. Abundant about Twickenham, Whitton, Hounslow, Brentford, &c., Middlesex; also at Barnes Common and Wimbledon Common, Surrey, *G. F.*

Arniston Woods, Edinburgh, *Mr. W. Brand.* Cawdor Woods, *Mr. W. Stables.* Moray, *Rev. G. Gordon.* Sutherland, Aberdeenshire, and Kincardineshire, but by no means common, *Dr. Murray.* Orkney, *Dr. Gillies.* Near Wrexham, Denb., *Mr. J. E. Bowman.* Castell Aber, Lleiniog, Anglesea, *Mr. W. Leighton.* —*y:* Caernarvon Castle, *Mr. J. F. M. Dovaston.* Carreg Onan, Anglesea, *Mr. W. Leighton.* *Luxuriant & abundant in our woods at Bristlington*

GEO.—Not found in the northern countries of Europe. In Germany as far north as Grimmia. Very rare in North America, being, according to Pursh, found only in one place, viz. New York.

BLECHNUM, Linn. HARD FERN.

(From βλεχνον, a Greek name for a Fern.)

A, portion of a fertile frond of *Blechnum boreale*. B, pinnule magnified, showing the covers or indusiums. C, transverse section of sorus, pinnule, and indusia. D, theca and spores. E, transverse section of rachis.

A genus of thirty-one species, known by bearing its fruit in closely united masses, not on transverse veins, as in *Scolopendrium*, but one on each side, and close to the midrib of the pinnule. Covers attached on the outer side of each mass, opening on the inner side, but not folding over each other, as in the last genus.

BLECHNUM BOREALE.

HARD FERN. ROUGH SPLEENWORT.

(Plate 5, fig. 6.)

CHA.—Frond pinnate, erect. Pinnæ linear, entire. Rachis smooth.

SYN.—*Blechnum boreale*, Swz., Willd., Spreng., Smith, Hook., Mack., Galp., Gray.—*Blechnum spicant*, Roth., With.—*Osmunda spicant*, Linn., Bolt., Hedw., Ehrh., Lightf.—*Osmunda borealis*, Salisb.—*Lonchitis aspera*, Ray, Ger.—*Acrostichum nemorale*, Lam. Fl. Fr.—*Acrostichum spicant*, Sibth., Vill.—*Asplenium spicant*, Bernh.—*Onoclea spicant*, Hoffm.—*Lomaria spicant*, Desv., Newm., Presl.

FIG.—E. B. 1159.—Bolt. 6.—Flo. Dan. 99.—Ger. 1140.—Schk. fl. 110.

DES.—Root black, tufted, scaly, with stout fibres. Rachis smooth and polished. Fertile fronds numerous, erect, strap-shaped, tapering at each end, about a foot high. Pinnæ linear, dilated somewhat at the base, in some degree falcate, distant from each other, and alternate, wholly covered on the under side with fruit. Barren fronds lanceolate, shorter than those which are fertile, and growing more on the outside of the plant, their pinnæ oblong, curved upward, and placed close together at their bases, but scarcely dilated at that part. Sori continued in an uninterrupted line from the base

to the point of each pinna, one on each side of the midrib. Indusium attached to very near the edge of the pinna, opening on the side nearest the midrib.

While young the back of the lobe shows only the midrib and two irregularly-edged white covers; afterwards these bend back and turn brown, and as in our species no leafy expansion appears outside the lines of thecae, but the cover seems to be the edge of the frond reversed, it might be taken at first sight for a *Pteris*, yet upon examination a narrow extension of the frond will be seen beyond the insertion of the indusiums. A curious variety of *Blechnum boreale* is found by Miss Beever, near Ambleside. Its lobes are much distorted, serrated, toothed, or deeply crenate. I have ventured to name and figure a portion of one of the fronds kindly sent me by Miss Beever.

β (*stricta*.) Frond linear, pinnules abbreviated, and with irregular margins.

SIT.—On sandy heaths, hedge-rows, stony places, &c.

HAB.—Spread throughout England, Scotland, and Ireland, in the last country especially in the counties of Wicklow and Clare. It ascends to 700 yards in Cumberland, 800 in Forfarshire, and much higher on the Cairngorum Mountains, in Aberdeenshire, where it probably attains to situations of the height of 1200 or 1300 feet, Mr. H. C. Watson.

GEO.—Common in Germany, Denmark, Norway, Sweden, and N. W. coast of America. *Conchium woods, Glostershire*—

PTERIS, Linn. BRAKES.

(πτερίς, a Fern; from πτερίξ, a feather.)

A, part of one of the divisions of the frond. B, the same magnified, showing the continued indusium. C, transverse and perspective view of part of a pinnule. D, theca and spore. E, outer indusium magnified, showing its ciliated margin. F, transverse section of the rachis near the root. G, ditto of the creeping rhizoma.

A very extensive genus, comprising no less than 120 species, most of them from warm climates. One species only is British. The fructification is borne in a continued line along the margin of the frond, which appears to be turned over so as to form a continued indusium, but which upon microscopic examination is seen to be of different and more delicate structure; an inner indusium is also present in ours and some other species, which many botanists consider a necessary character of a *Pteris*, and that its absence or presence might serve to divide the genus into two.

1

2

3

4

5

6

British Museum (Natural History)
Cromwell Road, London SW7 5BD
Telephone 01-589 6323

British Museum (Natural History)
**Lectures, films and
gallery tours**

May June July 1978

General information

Education Officer **Frank H Brightman**
Lecturers **Joyce Pope, John Stidworthy, Anne McCord**

Most of the tours and lectures are intended for a general audience,
but they are not suitable for very young children.

Lectures and film shows are held in the Lecture Hall and begin at 3pm.
All lectures are illustrated with films or colour slides.

Gallery tours begin at the Main Entrance at 3pm.

Programmes may be subject to alteration.

The Museum is open weekdays 10am–6pm, Sundays 2.30pm–6pm.
It is near South Kensington Underground Station or may be reached by
Buses 74, 14, 30, 45, 49 and 39A (Saturdays only).

RH Hedley
Director

The picture of *Linnaea borealis* is part of our commemoration of the bicentenary of the death of Linnaeus, the great Swedish botanist on whose work our present system of naming plants and animals is founded. Although it is unusual for a scientist to call a species after himself, Linnaeus, who gave the first scientific description of this little plant, which grows in Scandinavia and is sometimes found in northern Scotland, did so. He explained that he named the plant as he did because it is small, obscure and short lived, like himself. Such modesty appears to have been unusual in Linnaeus, who is reputed to have been inordinately vain and to have had his portrait painted more than any of his contemporaries, including the king of Sweden.

British Museum (Natural History)

Lectures, films and gallery tours

May

Wednesday 3	Introducing Birds	Lecture/Joyce Pope
Saturday 6	From Zermatt to the Glaciers	Special Lecture/Desmond & Marjorie Parrish
Wednesday 10	Polymorphism	Tour/Joyce Pope
Saturday 13	The Natural History of the Burren	Special Lecture/Maryangela Keane
Wednesday 17	Giants of the Past	Tour/John Stidworthy
Saturday 20	The Colonial Naturalist	Film Show
Wednesday 24	Introducing Insects	Tour/Joyce Pope
Saturday 27	Dinosaurs	Lecture at 11am/Joyce Pope
	Migration	Lecture/Joyce Pope
Monday 29	Conservation in Britain	Lecture/Joyce Pope
Wednesday 31	Sea Monsters of the Past	Lecture/Joyce Pope

June

Saturday 3	Life in Mountains	Lecture/Anne McCord
Monday 5	Rhinos and their Relatives	Lecture/John Stidworthy
Wednesday 7	Animals and Man	Lecture/John Stidworthy
Saturday 10	The Cat Family	Lecture/John Stidworthy
Wednesday 14	Cuckoos	Tour/Joyce Pope
Saturday 17	Dinosaurs	Lecture at 11am/Joyce Pope
	Jellyfishes and their Relatives	Lecture/Joyce Pope
Wednesday 21	The Ice Age	Tour/Anne McCord
Saturday 24	Birds and Man	Lecture/Anne McCord
Wednesday 28	Mammals of the Past	Tour/John Stidworthy

July

Saturday 1	Linnaeus	Lecture/Frank Brightman
Wednesday 5	Famous Fossils	Tour/Anne McCord
Saturday 8	The Evolution of Man	Lecture/Anne McCord
Wednesday 12	Insects and Flowers	Tour/Joyce Pope
Saturday 15	Dinosaurs	Lecture at 11am/Joyce Pope
	Ladybirds	Lecture/Joyce Pope
Wednesday 19	Animals Colours	Tour/John Stidworthy
Saturday 22	Life Between the Tides	Tour/John Stidworthy
Tuesday 25	Life in Hedgerows	Lecture/Anne McCord
Wednesday 26	Pollution	Lecture/Anne McCord
Saturday 29	The Garden Naturalist	Lecture/Anne McCord

The Burren, in the north of County Clare, is a hundred square miles of natural rock garden. The clefts of the bare limestone are the home of many rare plants, such as the maidenhair fern, and over 800 other species of plant are found there. Mrs Maryangela Keane, a noted naturalist from Co. Clare, will be describing the extraordinary geological formation, and the flora and fauna of the Burren, along with the occupation of the area by ancient man.

Jellyfishes are primitive creatures, surviving from the distant geological past, yet they are among the most successful of water-living creatures, found throughout the oceans of the world. They are totally soft bodied, yet they are flesh eaters, capable of subduing quite large prey with their paralysing stings and sometimes even human being are killed by them. This lecture will show some of the variety and beauty to be found among jellyfishes and their relatives. It will be illustrated by film 'Moon Jelly', a new film from Japan, which is about our commonest and least harmful jellyfish.

The brightly coloured ladybirds are lucky symbols to many people. Why they should have their curious name, for none is a bird and only about half are female, why they should be among the very few beetles to be more than tolerated by most people, why they are red and black and why we have in the last few years suddenly found huge numbers of ladybirds in our countryside, towns and houses, are all questions to be discussed in this lecture, which will be illustrated by slides and the French film 'Aphid-eating Ladybirds'.

PTERIS AQUILINA.

BRAKES. BRACHEN. FEMALE FERN.

(Plate 6, fig. 1.)

CHA.—Frond thrice pinnate. Larger pinnules pinnatifid, smaller entire. Rachis smooth.

SYN.—*Pteris aquilina*, Linn, and all modern Authors.—*Filix fœmina*, Ray, Ger.—*Asplenium aquilina*, Bernh.—*Pteris caudata* β, Schk.

FIG.—E. B. 1679.—Ger. 1128.—Bolt. 10, (all bad).

DES.—Root long and creeping, black and smooth when old, tomentose and brown when young. Rachis smooth, shining, without pinnæ on the lower part, tapering and black near its junction with the root. Fronds annual, erect, rigid, repeatedly divided, 2 to 5 feet high. Pinnæ opposite, more and more divided downwards, the smallest entire, the next pinnatifid, still lower ones pinnate, pinnato-pinnatifid, and twice pinnate. Pinnules opposite below, alternate above, oblong, blunt, connected to the midrib by their whole base, that terminating the pinna much larger than the others near it. Sori in a continued line around every sinuosity of the pinna. Indusium adhering to the margin of the frond, within which is another cover, contrary or opposite to the outer one, and in like manner fringed. When the young fronds first uncoil themselves they are densely downy.

If the stem be cut across near the root, it exhibits the bundles of vessels very plainly, in the form of an oak tree, or, as Linnæus thought, a spread eagle; hence its name *Aquilina*. This is seen in the generic wood-cut above, where also is a transverse section of the rhizoma, showing a totally different arrangement of vessels. The circumstance of the curious arrangement of vessels of the stem was a matter of notoriety at a very early period. Thus we find in a most rare little book, entitled, "A Dialogue or Communication of two persons devysed or set forth, in the Latine Tongue, by the noble and famous elarke Desiderius Erasmus, intituled, The Pilgrimage of pure Devotyon, newly translatyd into Englishe." (no date, supposed to be 1551), is the following curious passage: "Peraventure they ymagyne the symylytude of a tode to be there; evyn as we suppose when we cutte the fearne stalke there to be an egle."

This Fern is useful for many purposes, independently of the anthelmintic and astringent properties the herbalists attach to it. It is the favorite haunt of the deer tribe. As it is very long before it rots, and does not harbour insects, it is excellent as thatch; it does not hold moisture so much as straw, and is therefore better as litter for cattle, and as a cover to preserve plants from frost. It is also very excellent to lay fruit upon, or to pack it in, as it does not communicate any mustiness. Containing tannin, it is useful in the preparation of the lighter kinds of leather, and affords excellent potash when burnt. Its harsh texture and astringent taste render it unpalatable to cattle, though the roots are sought for

by pigs, and have even been dried and ground for bread, but only in times of the greatest scarcity. Upon being boiled, they yield a strong mucilage. The peasants of most parts of the kingdom assert their right to it as fuel, and use it chiefly to heat their ovens, a purpose for which it is well adapted, as it burns furiously. It is so valuable to the farmer of Germany for cattle fodder, that it is an article of ready sale there, and the cutting of it subject to very severe forest laws.

It remains dormant during more than half the year, the fronds not appearing till the middle of May, and being cut off with the first slight frost of autumn. It is also very impatient under culture: to remove a root otherwise than with a considerable quantity of earth attached to it, or in any season but that of its torpidity, would assuredly destroy it, as would also cutting down the fronds three or four seasons in succession. The remarkable paucity of young fern plants, of almost every species, must have struck the attention of most botanists. A single frond of *Pteris aquilina* produces more seed than any number the mind can conceive; millions of fronds do often extend over a waste, or park, yet how rarely is a young plant to be discovered any where. Indeed, had young plants been frequent, our ancestors could scarcely have imbibed the notion that they yielded no seed, or that it was a rarity, and only to be procured at the exact hour of the night on which John the Baptist was born. Pliny says, "of fern be two kinds, and they bear neither flower nor seed." Culpepper writing upon this Fern, which was in his time called Female Fern, "the seed of which," he observes, "some authors hold to be so rare," says, "such a thing there is, I know, and may be easily had upon Midsummer eve, and for aught I know, two or three days after it, if not more." The supposed circumstance of its seeding upon a single night, occasioned it to be called in Brown's pastoral ballads (1613).—

"The wondrous, one-night-seeding fern,"

Butler alludes to this superstitious notion. Hudibras, Part III, cant. iii. 3, 4.

"That spring like fern, that insect weed,
Equivocally without seed."

Absurd as these notions are, they were not wholly exploded in the time of Addison. He laughs at a doctor "who was arrived at the knowledge of the green and red dragon, and had discovered the female fern seed." Then again, in the dawn of botany and medicine, when affinities and antipathies, or as it was called the doctrine of signatures, was supposed to rule all things, we find that this Fern must be good for reed wounds, (punctured wounds) because, Dioscorides saith, "the fern dieth if the reed be planted about it; and, contrarywise, that the reed dieth if it be compassed with fern," which, as Gerard justly tells us, "is vaine to thinke that it hapneth by any antipathic or naturall hatred, and not by reason that this fern prospereth not in moist places, nor the reed in dry." Another result of the admirable and scientific reasoning of Dioscorides was once prevalent in this country, that, because Fern seed was invisible, therefore forsooth, those who carried it about them were rendered invisible also. This circumstance relative to Fern seed is alluded to in Beaumont and Fletcher's "Fair Maid of the Inn:"

"——— Had you Gyges' ring?
Or the herb that gives invisibility?"

Again, in Ben Jonson's play of the "New Inn:"

"——— I had
No medicine, Sir, to go invisible,
No Fern seed in my pocket."

Also, in Shakspeare's Henry IV., Part I., though here spoken ironically, Gads-hill says, "We have the receipt for Fern seed, we walk invisible."

Several other country adages attach themselves to the Fern, as the following :—

"When the Fern is as high as a spoon,
You may sleep an hour at noon;
When the Fern is as high as a table,
You may sleep as long as you're able."

Passing however these absurdities, of which many others might have been adduced, we may remark that very few of our poetical writers have thought the fern tribe worth their attention. Miss Twamley, however, is an exception; she has many passages in the "Romance of Nature," and other works, which relate to them. She speaks of "the Fan-like Ferns, which seem poised still and sleepily until the morn returns." In another place,

The Ferns too, are waving all stately here,
With seed-stored fronds thickly laid;
And shedding, when hastily brushed by the deer,
Their light, fertile dust o'er the glade.

Sir.—Upon barren heaths, in parks and woods, contenting itself occasionally with any soil or situation; it delights, however, in sand and strong loam, while it shuns the limestone and chalk districts; thus, if I recollect rightly, it is scarcely found on Salisbury Plain, nor do I remember meeting it any where in Kent, except in sandy spots. Be it observed, however, that it is not wholly excluded from chalk and limestone, as I have seen it occasionally on both. It is not fond of a lofty situation, as, according to Mr. Watson, it is not found in places more than 500 or 600 yards above sea level.

Geo.—Generally distributed over Europe, and in North America. The American species varies a little from ours, being rather more finely divided, somewhat ciliated, and earlier in growth.

CRYPTOGRAMMA, Br. ROCK-BRAKE.

(From κρυπτος, covered, γραμμα, a line; from the concealed lines of theææ.)

A, portion of a fertile frond of *Cryptogramma crispa*. B, a pinnule somewhat enlarged. C, ditto with the lateral margins thrown back to show the position of the fruit. D, transverse section of the pinnule. E, pinnule of a barren frond. F, a variety of ditto. G, theca thrown open and spores.

Our only plant was long considered a *Pteris*, because, although very different in habit, it has, like that genus, its fruit situated near the edge of the frond in an apparent continued line, the reflexed edge forming its cover. It differs, however, in having its sori not continued along all the undulations of the whole frond, but confined to the sides of the separate lobes.

CRYPTOGRAMMA CRISPA.

ROCK-BRAKE. STONE-FERN. CRISPED-FERN. PARSLEY-FERN.

(Plate 6, fig. 2.)

CHA.—Frond thrice pinnate. Fertile pinnules oblong, blunt : barren ones wedge-shaped, cleft, crenate.

SYN.—*Cryptogramma crispa*, Hook. in *Br. Fl.*, Mack.—*Pteris crispa*, Linn. MSS., Willd., Swz., Hull, With., Smith, Hook. in *Fl. Sco.*,—*Osmunda rupestris*, Salisb.—*Osmunda crispa*, Linn. in *Sp. Pl.*, Huds., Lightf., Bolt.—*Stegania onocleoides*, Gray.—*Onoclea crispa*, Roth., Hoffm.—*Allosorus erispus*, Bernh., Kaulf., Spreng.

FIG.—*E. B.* 1160.—*Bolt.* 7.—*Flo. Dan.* 496.—*Pluk. Phyt. t.* 5. *f.* 2.—*Newm.* 18.

DES.—Root slightly creeping, long and fibrous. Frond thrice pinnate, deciduous, of a very lively green color, 3 to 12 inches high. Rachis slender, smooth, and shining. Barren pinnules wedge-shaped or roundish, deeply cut and crenate, pinnæ nearly opposite, but not always so, four or five pairs. Fertile fronds taller and more robust, but less expanded than the barren ones ; their pinnæ more inclined to be alternate. Pinnules oblong, elliptic, blunt, their crenate sides turned over upon the sori, which are in lines along each side of the lobe, distinct only for a very short time at first, then very confluent and crowded.

SIT.—Southey calls this plant the “ Mountain Parsley ; ” an appellation which well expresses its tender habit, its delicate, lively color, and its numerous, finely cut, and crisped leaves. Covering large patches as it sometimes does on the tops of rocky mountains, it adds a bright gleam of verdure and of beauty to its romantic but barren dwelling place, and becomes an oasis of rich fertility upon the precipitous face of the otherwise sterile rock.

HAB.—From 200 yards upwards to a considerable elevation in Caernarvonshire (top of Snowdon). In Cumberland from 200 or 300 yards to 1040 yards. In the Highlands, from the low valleys to 1100 yards on Ben-na Baird. More common in the lake district of England than in Scotland, but frequent in several parts of the latter, *Mr. H. C. Watson*. Breiddon Hill (12 miles west of Shrewsbury), *Mr. J. E. Bowman*. Greenfield, Saddleworth, *Mr. J. Merrick*. Higher parts of the Tees, *Mr. Hogg*. Common about Settle, Yorkshire, *Mr. J. Tatham*. Skiddaw, Helvellyn, Saddleback, Grassmoor, Vale of Newlands, &c., Cumberland, *Mr. H. C. Watson*. On rocks at the foot of Cheviot, above Langley Ford, *Mr. Winch*. Near Lancaster, *Mr. W. Wilson*.—WAL. : Mount Glyder, Mount Snowdon, and Mynydd Mawr, Caernarvonshire, *Mr. C. C. Babington*. Cader Idris, *Mr. Purton*. North Wales, (abundantly), *Mr. W. Christy*.—SCO. : Rosshire, *Rev. G. Gordon*. Glen Tilt and Blair Athol, Perthshire, *Mr. W. Brand*. Not rare in Sutherland, *Dr. Murray*.—IRE. : Abundant on the Mourne Mountains, *Mr. Mackay*.

GEO.—Lapland, Germany, Switzerland, Pyrenees, Silesia, Sweden, Jutland, Norway, Dauphiny, Holland.

1849. Aug. 3 abundant about Windermere.—GEORGE J. F. BENNETT com.
1856. — abundant on Snowdon. At M. Weymouth all of May Childers com.

ADIANTUM, Linn. MAIDEN-HAIR.

(From, α against, and $\delta\alpha\upsilon\tau\alpha$, moisture; the plants never being wet.)

A. pinnules of *Adiantum capillus-veneris*, showing the position of the sori and indusiums. B, an indusium removed, showing the attachment of the sori, one indusium covering several. C, theca and ring. D, spore.

A very beautiful, delicate, and interesting genus of sixty-three species, indigenous to the southern countries of Europe and the tropical regions, this country being the northern limit of them all. The sori are arranged in spots along the margin of the pinnules, and covered by part of the frond reflexed.

ADIANTUM CAPILLUS-VENERIS.

TRUE MAIDEN-HAIR.

(Plate 6, fig. 3.)

CHA.—Frond twice pinnate. Pinnules alternate, wedge-shaped, lobed, on capillary petioles. Indusium oblong.

SYN.—*Adiantum capillus-veneris*, Linn., Willd., Smith, Bolt., Dieks., Hook., Mack.—*Adiantum capillus*, Swz.—*Adiantum fontanum*, Salisb., Gray.—*Adiantum coriandrifolium*, Lam.—*Capillus-veneris verus*, Dill. in Ray's Syn., Ger.

FIG.—E. B. 1564.—Bolt. 29.—Jaeq. Misc. t. 7.—Ger. 1143 (bad).—Newm. 9.

DES.—Root slightly creeping and very hairy. Rachis slender, shining, rigid, purplish-black, without pinnæ on the lower part. Pinnæ alternate, in young fronds lobed only, afterwards pinnate. Pinnules wedge-shaped, crenate or cleft at the top, alternate. Sori marginal, in spots, one near the end of each lobe of the pinnule; the apex of which is turned over, forming a white, oblong cover, to which the fruit itself is attached.

The manner of the expansion of this plant is very singular and interesting. The young frond is but slightly circinate in vernalion, appearing at first with only one or two small, wedge-shaped pinnules; after a time these split into lobes, which lobes become wider, long-stalked, and detached from each other, forming separate wedge-shaped pinnules, exactly similar to those from which they were detached, and if the plant be luxuriant, these again divide in a similar manner; thus some fronds are found pinnate, others twice, and sometimes thrice pinnate. The whole plant forms an interesting object for the microscope, particularly the membranous indusium, which is beautifully veined. The ring of the theca also is very different from that of any other British Fern. (See end of the genus.)

VIR.—The properties of *Adiantum* are very uncertain. Its use is said to give name to the syrup *Capillaire*. It has neither fragrance nor flavor, and when boiled yields only a little mucilage.

HAB.—Port Kerig, Glamorganshire (verified 1834). Banks of the Carron, a rivulet in Kincardineshire, *Professor Beallie*. In a small cave on the east side of Carrach Gladden, a cove on the north coast of Cornwall, between Hayle and St. Ives, *Professor Henslow*. Isles of Arran, county of Galway, *Dr. Osborne*. At Wrisbeg, on a rock facing south-west on the shore of Loch Bulard, *Mr. C. C. Babington*.

GEO.—South Europe, Isles of Bourbon, Teneriffe, Jamaica, and Hispaniola.

1840. Mar: 29—Malta. abundant. *Ellen combe in the legends in other damp places*
 1843. White Pebble Bay. *Ellen combe. Ellen & our Children combe*
 1853. July Damp Rocks on Coast at Lelant. Cornwall. *Ellen & our Children combe*

HYMENOPHYLLUM, Swz. FILMY PERN.

(*υμνη*, a membrane, *φυλλον*, a leaf; or the membranous-leaved Fern.)

A, part of a frond of *Hymenophyllum Tunbridgense*. B, the same slightly increased to show the veins of the frond, and the origin and character of the fruil. C, sorus magnified, and one of its covers removed. D, theca with transverse ring. E, the same opened. F, spores.

In this small and delicate genus, a lobe of the pinna is contracted into the fruil and its receptacles, the lamina of the lobe forming two valves, inclosing between them the midrib, to near the end of which are attached several ringed and petioled thecæ, the annulus of which does not coincide with the petiole, but is placed transversely. (See Introduction).

1.—HYMENOPHYLLUM TUNBRIDGENSE.

TUNBRIDGE FILMY-FERN.

(Plate 6, fig. 4.)

CHA.—Frond pinnate. Pinnæ pinnatifid, erect. Lobes serrated. Rachis winged. Involucre orbicular, serrated at the top.

SYN.—*Hymenophyllum Tunbridgense*, *Smith, Willd., Hook., Mack., Swz., Gray.*—*Trichomanes Tunbridgense*, *Linn., Huds., With., Bolt., Lightf.*
 —*Trichomanes pulchellum*, *Satish.*

FIG.—*E. B.*—162.—*Hook.* in *Flo. Lon.* 71.—*Bolt.* 31.—*Flo. Dan.* 954.—*Hedw.* 3.—*Forst* in *Flo. Tonb.* (excellent.) *Newm.* p. 92.

DES.—Root black, fibrous, hairy, extensively creeping, rather upon than under the surface of the ground. Rachis naked on the lower part, capillary, black, broadly winged all the way down. Fronds solitary, at intervals along the creeping stem or root, 1 to 2 inches high, of a light green color. Pinnæ alternate, growing quite upright, their veins dichotomously branched. Lobes sharply

serrated or toothed, linear and blunt pointed, running into each other, and seated chiefly on the upper side of what may be called the midrib of the pinna, but not wholly confined to that side, as in the next species. Receptacles formed from and in the place of the last lobe, on the upper side of each pinna; thus they appear in two rows, one on each side of the rachis. The receptacle is composed of two flat or slightly convex, roundish valves, folding over each other, and sharply serrated at the points; between which is a free column covered with thecæ.

SIT.—On damp, shady rocks, generally among moss.

HAB.—On the moist and shady sides and fissures of the various rocks near Tunbridge Wells, viz., the High Rocks, and the rocks in Eridge Park (abundant, 1835), *Mr. W. Pamplin*. Clefts of the rocks at Wistman's Wood, Dartmoor; rocks by Dunsford Bridge, Becky Fall, &c., Devon, *Flo. Dev.* Greenfield, near Saddleworth (very rare), *Mr. W. Wilson*. Near Halifax, *Mr. Leyland*. Near Cader Idris and Dolgelle, *Mr. Bowman*. Very abundant and fine near the Upper Lake, Killarney, *Mr. W. Wilson*. Powerscourt Waterfall, Glencree, and other places in the county of Wicklow, *Mr. Mackay*.

GEO.—This and probably the next species are scattered over Europe from Italy to Norway.
Abundant in the Granite Cairns about Pizgance—Ellen and Children came—

2.—HYMENOPHYLLUM WILSONI.

NORTHERN FILMY-FERN. SCOTTISH FILMY-FERN.

(Plate 6, fig. 5.)

CHA.—Frond pinnate. Pinnæ semi-pinnatifid, recurved. Lobes serrate. Rachis not winged. Receptacle ovate, entire.

SYN.—Hymenophyllum Wilsoni, *Hook in Br. Flo., Mack., Newm.*

FIG.—*E. B., suppl. 2686. Newm., p. 94.*

DES.—Rachis rigid, capillary, winged at the top. Frond 1 or 2 inches high, dark green. Pinnæ alternate, bent backwards, growing horizontally rather than vertically as in the last species, besides which the lobes curve downwards, so that when the edge of them is looked at they have a falcate appearance, although they are oblong and blunt, and it may be added, very sharply serrated. When in fruit, all the leafy expansions turn in one direction, and the fruit in the opposite. The receptacles are situated as in the last species, but are larger, very convex, perfectly ovate, and entire.

The absence of wings to the rachis, the different habit, the semi-pinnatifid character of the pinnæ, and the entire, convex receptacles, serve to distinguish this from Hymenophyllum Tunbridgense, with which it was confounded, until shown to be distinct by that accurate botanist, Mr. W. Wilson.—Miss Beaver has sent me fronds from Coniston, which are much branched.

SIT.—On moist alpine rocks, near waterfalls, &c.

HAB.—Waterfall above Ambleside, Westmoreland, *Mr. J. Bowerbank*. Black Rocks of Great End, in the Scawfell range, and at Scale Force, near Buttermere, Cumberland, (1833,) *Mr. H. C. Watson*. Greenfield, near Saddleworth, and near Silverdale, Lancashire, *Miss Beever*.—WAL.: On Snowdon, near Llanberis Pass, and on the adjacent mountains, especially near Tŷll Du, *Mr. W. Wilson*. Rocks about Nant Phrancon, in situations from 200 to 650 yards of elevation, *Mr. H. C. Watson*. On rocks, near the Rhydol, Montgomeryshire, at a plank over a dangerous gulph of the river Pont Bren, *Mr. E. Lees*.—SCO.: Finlarig Burn, near Killin, Perthshire, *Mr. Wilson*. Argyleshire, *Mr. J. Hooker*.—IRE.: At Killarney, (very plentiful,) *Mr. Wilson*. Shanafolia Mountain, *Mr. C. C. Babington*. Kerry Mountains, Cunnemara, &c., *Mr. Mackay*.

1853 July. Abundant on the mossy granite Boulders between Pinguine Lake &c.

TRICHOMANES, Linn. BRISTLE-FERN.

(Τριξ, τριχος, a hair, and μακρος, loose or long; from the free hairs which terminate the receptacles.)

A, pinnule of *Trichomanes brevisetum*. B, portion of ditto with fruit, enlarged. C, ditto, still more greatly enlarged, to show the loose cellular structure of the frond. D, longitudinal section of the sorus magnified. E, theca, with transverse ring. F, spores.

All the species of this beautiful genus, amounting to forty-six in number, are very cellular and tender, their fruit attached to the midrib of a lobe, as in the last genus, but here the receptacle is one-valved, and the midrib not terminated by the thecæ, and confined within the receptacle, but projecting much beyond it, and like a hair in appearance. We have but one species, and that very rare.

TRICHOMANES BREVISETUM.

SHORT-STILED BRISTLE-FERN. CUP GOLDBLOCKS.

(Plate 6, fig. 6.)

CHA.—Frond thrice pinnatifid. Lobes linear, entire. Rachis winged. Receptacles urceolate.

SYN.—*Trichomanes brevisetum*, *Hort. Kew.*, *Hook.* in *B. Fl.*, *Smith* in *E. Fl.*, *Mack.*—*Trichomanes alatum*, *Hook.* in *Flo. Lon. N. S.*, *Swz.*, (Not of *Willd.*)—*Trichomanes pyxidiferum*, *Huds.*, *Bolt.*, *With.*, *Hull.*—*Hymenophyllum alatum*, *Smith* in *E. B.*, *Willd.*—*Hymenophyllum Tunbridgensæ* β , *Smith* in *Fl. Br.*

FIG.—*E. B.* 1417.—*Ray. Syn. t.* 3, *f.* 3.—*Bolt.* 30.—*Flo. Lon.* 53.—*Newm.*, page 88.

DES.—Root very thick, black, and densely hairy. Rachis smooth and winged all the way down. Frond pellucid, membranous, dichotomously branched in all its parts, 6 to 12 inches high, dark green. Pinnæ alternate, twelve or fourteen pair, vertical, much cleft, lobes ultimately linear, but every where running much into each

1

2

3

5

4

other, their veins conspicuous, prominent, and beautifully branched. Receptacles pitcher-shaped, taking the place of lobes, but not confined to those nearest the main stem, as in the last genus.

HAB.—Near Killarney, in several situations, *Mr. W. Wilson*. Hermitage, in the county of Wicklow, *Mr. Nuttall*. Powerscourt Waterfall, *Mr. Mackay*. Once found in Ballinhasy Glen, near Cork, by *Mr. J. Drummond*. Its situation at Turk Waterfall, Killarney, is thus described by *Mr. Newman*:—"I here found it to the left of the seat whence tourists take the first view of the fall. About 15 yards higher up the stream, the rocky bank on the left projects into the river; this projection is only to be approached by leaping from stone to stone, along the bed of the torrent, which in time of flood is rather an exciting and ticklish operation: you are so close to the fall as to be covered by the spray, and the roar is almost deafening. Having reached the projection, the botanist must ascend it by means of the roots and branches, a feat very readily performed; and there is a little platform at the top, where he can stand very comfortably; and while so standing, he will find the rocky bank just on a level with his eyes, completely covered with Trichomanes, the dark green fronds hanging heavily down, dripping with wet, and if the sun happen to shine, begemmed with sparkling drops." Glendine, near Youghal, county of Cork, is another habitat for this plant, as discovered by *Mr. Ball*, of Dublin. It grows here in great luxuriance.

GEO.—St. Domingo, Jamaica, the Caribbees, Madeira, &c.

Waterfall, Killarney with Orchideaceae. Aug. 1839 -

ly. Valencia - Ireland. (a doubtful habitat) -

Valpurga Island. Iberian &c. & Tristram's Cornish (H. & J.) & 8, 10, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935, 936, 937, 938, 939, 940, 941, 942, 943, 944, 945, 946, 947, 948, 949, 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, 984, 985, 986, 987, 988, 989, 990, 991, 992, 993, 994, 995, 996, 997, 998, 999, 1000.

OSMUNDA, Linn. ROYAL-FERN.

(*Osmund*, *Sax.*, strength; this being the largest and strongest of our Ferns (?))

A, portion of a frond of *Osmunda regalis*, natural size. B, mass of fruit magnified. C, theca separated. D, ditto splitting open. E, spores. F, transverse section of the stem. G, cuticle of a pinnule. H, cuticle of the stem.

In this genus the upper part of the leafy frond becomes changed into a compound spike of fructification, without any indusium, receptacle, or annulus. *Osmunda* is a small genus, the species of which very much resemble each other in size and character. They are natives of Europe and North America.

OSMUNDA REGALIS.

ROYAL-FERN. WATER-FERN. FLOWERING-FERN.

(Plate 7, fig. 1.)

CHA.—Frond bipinnate. Pinnules oblong, nearly entire, slightly auricled. Sori terminal.

SYN.—*Osmunda regalis*, Linn., Willd., Bolt., and all modern botanists.

FIG.—*E. B.* 209.—*Bolt.* 5.—*Flo. Dan.* 217.—*Flo. Lon.* 150.—*Ger.* 1131.—*Newm.*, page 97.

DES.—Root a thick, short, scaly, and fibrous tuber. Rachis smooth, rigid, upright. Fronds several, 3 to 6 feet high, bright green, twice pinnate. Pinnæ distant, nearly opposite. Pinnules almost sessile, oblong, blunt, with waved or slightly crenate edges, frequently auricled. Those pinnæ on the top of the frond are either wholly or partially changed into fructification, when they appear like a compound spike, each bunch of which seems composed of a number of circular bundles of capsules. The thecæ are petioled and beautifully reticulated. Spores nearly globular.

VIR.—The internal parts of the root, as well as the young fronds, were once used in pharmacy as a cure for bruises, and as conferring strength.

SIT.—In wet woods, swampy moors, &c. (See *Introduction*, page 11.)

HAB.—SCO.: Head of Loch Fine, to the N. E. of Inverary, Argyshire, and near Loch Lomond, (Dumbarton side,) *Mr. H. C. Watson*. At the side of the loch at Inchnadamph, Sutherlandshire, *Dr. Johnston*. Aberdeenshire and coast of Kincardineshire, *Dr. Murray*.—ENG.: Warwickshire, *Rev. W. Bree*. Ellesmere Lakes and West Felton, Salop, *Mr. W. Leighton*. Plentiful at Speke, near Liverpool, *Mr. T. B. Hall*. Chat Moss, *Mr. W. Christy*. Woolston Moss and other places near Warrington, Lancashire, *Mr. W. Wilson*. Isle of Man, *Mr. Forbes*. Pottery Car, near Doncaster, *Mr. S. Appleby*. Near Leeds, *Mr. Denny*. Bulwell, Notts, (near the upper mill,) *Mr. T. H. Cooper*. Norfolk, *Miss Bell*. Kavanah's Wood, Great Warley Common; also near the barracks, on Little Warley Common, *Mr. R. Castle*. Near Leith Hill, Surrey; and in several places from 5 to 8 miles S. W. of Dorking, *Mr. W. Pamplin*. On Bagshot Heath, *Mr. J. Lloyd*. In a wet shady spot, by the river side, between Frimley Village and Frimley Green, Surrey; also sparingly on Esher Common, by the entrance to the lane leading thence towards Epsom, *Mr. H. C. Watson*. Tonbridge, *Mr. Trevelyan*. Corner of the lake at Uckfield, Sussex, near Chudleigh, on the banks of the Teign; also near Ivy Bridge on the Erme; and on the Goonhilly Downs, about St. Ives, *Jones's Tour*. Isle of Wight, *Rev. G. E. Smith*.—IRE.: Mucruss Abbey, *Mr. Kelly*. Castlebar, Mayo, *Dr. Osborne*. Kelly's Glen, Co. Dublin, &c., *Mr. Mackay*.

GEO.—Europe, chiefly the northern parts, and all the United States.

*3 July - About Poughkeepsie with Ellen & our Children -
Abundant in Virginia - Rare in Wood at Cleveland - Jones*

BOTRYCHIUM, Linn. MOON-WORT.

(From *βολος*, a bunch; as its fruit is borne in clusters.)

A, fertile branched spike of fruit of *Botrychium lunaria*. B, part of ditto enlarged. C, ditto with the thecæ opened. D, spores. E, transverse section of the stem.

The fruit in this somewhat extensive genus is produced upon a compound spike distinct from the leafy expansion, though attached to it at the stem. The thecæ are opaque and sessile. There is only one British species. (See *Introduction*.)

BOTRYCHIUM LUNARIA.

COMMON MOON-WORT.

(Plate 7, fig. 2.)

CHA.—Frond pinnate, solitary. Lobes flabelliform, erenate.

SYN.—*Botrychium lunaria*, Swz., Willd., Hook., Mack., Smith in *E. Fl.*, Gray.—*Osmunda lunaria*, Linn., Smith in *Fl. Br.* and *E. B.*, Bolt., Lam., Dicks., Ehrh.—*Osmunda lunata*, Salisb.—*Lunaria minor*, Ger., Ray, Matth., Camer., Fuchs., Gesner, &c.FIG.—*E. B.* 318.—*Bolt.* 4.—*Flo. Dan.* 18, f. 1.—*Flo. Lon.* 66.—*Newm.*, p. 100.

DES.—Root of thick, smooth, yellow fibres. Frond of a dull, yellowish green, 2 to 6 inches high, rarely more than one from a root, quite smooth in every part. Stem hollow, rather succulent, half way up it divides into two branches, one being a pinnatifid or pinnate frond, the other the fruit. Pinnules of the leafy part five or six pair, opposite, decurrent, fan-shaped, regularly erenate. Fruit covering the upper part of the other branch of the stem in a compound spike, not in aggregate clusters, as in *Osmunda*, but separate, though nearly touching each other, and arranged in single lines along the branches of the spike. The thecae are opaque, sessile, round, smooth, yellow at first, afterwards brown. Spores oval, smooth, generally attached to each other in pairs.

VIR.—Its virtues are more imaginary than real, more magical than physical. Its name *Lunaria*, or Moon-wort, is taken from the shape of the leaves, and gathered by the light of the moon, was said to “doo wonders.” Gerard mentions a remarkable instance of the properties attributed to it by the alchemists and witches, “that it will loose locks, and make them fall from the feet of horses that do grase where it doth grow;” “too drowsie a dream” for even the credulous Gerard to believe; but he adds, that it is “singular for wounds.”

SIT.—In pastures chiefly in the northern and mountainous countries.

HAB.—SCO.: Bernerside Hill, W. of Berwickshire, *Mr. W. Baird*. South side of Loch Tay, and ascending to 3000 feet on adjacent mountains; Clova and Pentland Hills, &c. *Mr. H. C. Watson*. Blair Athol, Perthshire, *Mr. W. Brand*. Moray, *Rev. G. Gordon*. Orkney, *Rev. C. Clouston*. Aberdeenshire, *Dr. Murray*.—ENG.: Sea banks near Tynemouth, Northumberland, *Miss Hancock*. Higher Tees, *Mr. J. Hogg*. Newcastle Town Moor, *Mr. R. Bowman*. Common about Settle, Yorkshire, *Mr. J. Tatham*. Warwickshire, *Rev. W. Bree*. Southport, Lancashire, *Mr. Rylands*. Greenfield, near Manchester, *Mr. J. Merrick*. Cheshire and Derbyshire, *Mr. W. Wilson*. Pottery Car, *Mr. S. Appleby*. Clifton, Norton, Fiskerton, Newstead, and Sherwood Forest, Notts, *Mr. T. H. Cooper*. Shotover Hill, Oxen, *Mr. Baxter*. Linton, Cambridgeshire, *Mr. C. C. Babington*. Near Titchborne, Hants (1836), *Mr. Forder*. Leith Hill, Surrey; Shirley Common, near Croydon, Surrey; between Dartford and Foot's Cray, Kent; and S. W. of Petersfield, Hants, *Mr. W. Pamplin*. Deep Dean, near Dorking, *Mr. J. Nash*. South Kent, *Rev. G. E. Smith*. Near Barnstaple, Devon (1836).—WAL.: Near Wrexham, *Mr. J. E. Bowman*. Near Rodney's Pillar, Montgomeryshire, *Rev. A. Bloxam*. Craig Breidden, *Mr. Dovaston*.

GEO.—Throughout North Europe and North Asia.

77. July 1.—Collected in tolerable abundance by me on Skatsheloc Hill, part of the Mendips, above Sidcot School, about 15 min from 15 miles & 2 from Exeter.
C. E. Fox, H. O. Stephens & L. Butler com. S—

OPHIOGLOSSUM, Linn. ADDER'S-TONGUE.

(From *οφις*, a serpent, and *γλῶσσα*, a tongue.)

A, fertile and barren frond of *Ophioglossum vulgatum*. B, fertile frond after it has shed its spores. C, cuticle. D, transverse section of the stem. E, root and veneration. F, spores.

This genus bears its fruit in a simple spike attached to a leafy frond. The thecae are connected not only to each other, but attached by their whole base to the stem which bears them; when ripe they open transversely. There are twelve foreign species of this genus, inhabitants of Europe and North America.

OPHIOGLOSSUM VULGATUM.

COMMON ADDER'S-TONGUE.

(Plate 7, fig. 3.)

CHA.—Frond entire, solitary, ovate, obtuse.

SYN.—*Ophioglossum vulgatum* of most Botanists.—*Ophiog. ovatum*, Salisb.

FIG.—*E. B.* 108.—*Bolt.* 3.—*Flo. Lon.* 78.—*Flo. Dan.* 147.—*Ger.* 404.—*Sckh.* 153.—*Newm.* p. 103.

DES.—Root composed of a few stout, yellow, smooth fibres, running horizontally. Frond of one entire, upright leaf, ovate, blunt, 2 to 6 inches high, of a lurid green color. Stem tapering downwards, and hollow. Fruit in a single, unbranched, stalked, and pointed spike, connected with the leafy expansion. Thecae yellow, opaque, sessile, in two single rows, connected with each other, so that after the round, smooth, yellow seeds are dispersed, a number of transverse clefts are seen along each side of the spike. Sometimes found with more than one spike, at other times the leaf-like frond is deeply cleft at the top.

VIR.—It is considered by the country people as valuable to form an ointment for wounds, and for this purpose is gathered by baskets-full; for be it observed that in some parts of the country it is almost as abundant as the herbage among which it grows.

SRR.—In meadows and moist pastures in most parts of the kingdom. (See *Introduction*.)

HAB.—SCO.: Dalmeny Woods, near Edinburgh, Mr. W. Brand. Orkney, Rev. C. Clouston. Balmuto, Miss Boswell. Carlowrie, Mr. Falconer. ENG.:—Middleton-one-row, Durham, Mr. R. Borman. Round Howe, near Richmond, Yorkshire, Mr. J. Ward. West Felton, Salop, Mr. W. Leighton. Warwickshire, Rev. W. Bree. Field behind Heawood Hall, Alderley, Cheshire, Mr. H.

Watson. Near Warrington, Lancashire, *Mr. W. Wilson*. Near Braunston, Leicestershire, *Rev. A. Bloxam*. Heanor, Derbyshire, and Colwiek, Notts, *Dr. Howitt*. Pottery Car, *Mr. S. Appleby*. Near Bristol, *Miss Worsley*. Somerset, *Mr. A. Southby*. Norfolk, *Miss Bell*. At the side of the pond, in Wike Farm, Sion Lane, Isleworth; near the ladder stile, Osterley Park, near Brentford, Middlesex; and 4 miles south of Dorking (abundant), *Mr. J. Bevis*. Various parts of Surrey, Kent, Herts, and Hants, *Mr. W. Pamplin*. Meadows of Long Leet, Wiltshire, *Mr. Rowden*. Sussex, *Mr. H. C. Watson*. Near Slateford, Near Barnstaple, Devon, *Mr. J. Nash*—*WAL.*: Near Wrexham, *Mr. J. E. Bowman*.—*IRE.*: Lawn of the Observatory, Dunsink, *Mr. Kelly*. Not unfrequent in Ireland, *Mr. Mackay*.

GEO.—Throughout Europe, and from New York to Pennsylvania in North America.

America.
Copies in Stockwood have reflected at base of Hower Hill - Wilmington
Hence our children come

PILULARIA.—PILL-WORT.

(From *Pilula*, a little pil; from the shape of its seed-vessels.)

A, frond of *Pilularia globulifera* magnified, showing the vernalion, rhizoma, and roots in different states. B, cross section of the root. C, of the seed vessel. D, one quarter of ditto still further magnified, showing the spores and bags of granules. E, spore and abortive granules, the latter burst open. F, spore. G, point of ditto in germination. H, front view of the end of ditto in germination. I and J, germination still further advanced. K, cuticle of the stem. L, membrane which divides the root into various cells. M, hair from an involucre; a, its point of attachment. The germination taking place from a determinate point, shows that the word spore is as relative to the *Pilularia* a misnomer, and that the reproductive grains are real seeds. These figures are taken from Mr. Valentine's paper mentioned in the Introduction.

PILULARIA GLOBULIFERA.

CREeping PILL-WORT. PEPPER-GRASS.

(Plate ~~47~~ fig~~5~~ ~~10~~) 4.

СНА.—Leaves filiform. Stem creeping. Receptacles coriaceous, hairy, nearly radical.

SYN.—*Pilularia globulifera* of all botanists.

FIG.—*E. B.* 521.—*Bolt.* 40.—*Flo. Dan.* 223.—*Hook.* in *Flo. Lon.* 83.

DES.—Stem very long, cylindrical, and creeping close to the ground, throwing off at intervals of half an inch or more several

simple, very small, smooth, radical fibres, and from the same part upwards from two to six filiform, hollow, green leaves, about 2 inches long, among which, at their base, grow one or two receptacles, not radical, as generally said, but attached to the *upper* part of the stem, and therefore, although near the roots, not attached to them. The receptacles are round like a pepper-corn, (hence the name of the plant,) brown and hairy. The spores are oblong, contracted in the middle, and slightly pointed at one end.

SIT.—Pools of water, edges of lakes, &c. not uncommon.

HAB.—SCO.: Near Inverskin, Sutherland, *Mr. Campbell*. Marshy ground between the village of Currie and the Pentland Hills, Edinburgh, *Mr. H. Watson*. Near Slateford, Forfarshire, *Mr. W. Brand*.—ENG.: Prestwich, near Northumberland, *Mr. R. Bowman*. Coleshill Pool, Warwickshire, *Rev. W. Bree*. Bomere Pool, Salop, *Rev. E. Williams*. Near Richmond, Yorkshire, *Mr. J. Ward*. Beam Heath, near Nantwich, Cheshire, *Mr. J. E. Bowman*. Once plentiful at Bartington Heath, Cheshire, and still found at Baguley Moor, in the same county, *Mr. W. Wilson*. Filby and Hopton Commons, near Yarmouth, *Mr. Paget*. Once and perhaps still in the ponds at Roehampton, Surrey; and on Iver Heath, Middlesex, *G. F.* In a small pool between Okeshot Hill and Claremont Park, Surrey, *Mr. H. C. Watson*. Sussex, *Rev. G. E. Smith*. Grosvenor and Roche, Cornwall, *Jones's Bot. Tour*.—WAL.: Near Llanfaeloy, Anglesea; and border of Llyn Idwel, Caernarvonshire, *Mr. W. Wilson*.

GEO.—Most parts of Europe.

ISOETES, Linn. QUILLWORT.

(From 1705, equal, and 1705, the year; the plant being evergreen.)

PLATE OF GENERA, FIG. XVIII.

A, lower part of a plant of *Isoetes lacustris*, natural size. B, portion of the filiform leaf, much magnified. C, receptacle of the larger kind of fruit. D, receptacle of the smaller granules. E, spore magnified. F, arrangement of four spores upon one of the transverse bars of the receptacles, as described in the Introduction, p. 12. G, section of the receptacle which bears fertile spores. H, section of the abortive sporules.

ISOETES LACUSTRIS.

EUROPEAN QUILLWORT. MERLIN'S GRASS. *Plate 7. Fig. 5.*

CHA.—Leaves subulate, bluntly quadrangular, formed of four transversely-jointed longitudinal cells.

SYN.—*Isoetes lacustris* of all modern botanists.

FIG.—*E. B.*—1084.—*Flo. Lon. N. S.* 131.—*Bolt.* 41.—*Flo. Dan.* 191.—*Sehk. fil.* 173.

DES.—Root tufted, composed of long, branched, smooth fibres. Leaves radical, tufted, filiform or subulate, 2 to 4 inches high, light green, and very brittle. Receptacles formed of the base of the leaves: the outer, which are also the larger and older leaves, bearing perfect seeds; the inner and younger leaves produce finer granules, as explained in the *Introduction*.

MR. W. Wilson finds two varieties in Wales; the one densely tufted, with slender, erect leaves, the other with broader and widely-spreading leaves. The former of these, Dr. Hooker thinks may be the *Isoetes setacea* of Bosc. Sprengel says, "that the plant grows at the bottom of carp ponds, where it would not be of very easy access, did not the fish assist the botanist by disengaging it from the mud, when it is found floating at the edges of the pond."

SIT.—Found only in the extreme north of Wales, north of England, and in Scotland, which is a curious circumstance, because submersed water plants are not in general so strictly confined to particular latitudes or altitudes.

HAB.—SCO.: Loch Callader, Aberdeenshire, and Loch Brandy, Forfarshire, *Mr. W. Brand.* Loch Whirral, Forfarshire, *Dr. Graham.* Loch Tay, Perthshire, *Dr. Greville.* Most of the Scottish Lakes, *Mr. H. C. Watson.*—IRE.: Lakes in the Rosses, Donegal, *Rev. Mr. Murphy.*—ENG. and WAL.: Prestwick Carr, Northumberland, *B. G. Ulswater,* Cumberland, *Mr. Williams.* Coniston lake, *Miss Beever.* In Llyn-y-cwm, Plynnon Frich (Snowdon), Eake Ogwan, and Llanberris lakes; also in Floutern Tarn, between Scale Face and Whitehaven, *Mr. W. Wilson.* Lakes of Denbighshire, *Mr. J. E. Bowman.*

GEO.—More copious in Sweden and Denmark than elsewhere. New York and northwards in America.

LYCOPODIUM, Linn. CLUB-MOSS.

(From *λυκος*, a wolf, and *πους, ποδος*, a foot; the ends of the stems appearing like the hairy feet of some animals.)

A, spike of fruit of *Lycopodium selaginoides*, natural size. B, two leaves or bracts of ditto; one showing the larger grains, the other the receptacle for the smaller. C, receptacle opening and scattering the granules. D, cuticle of a leaf. E, section of the stem. F, spore. G, abortive granule.

A very extensive genus of no less than 140 species, found in all parts of the world, some in the hotter, and others in the colder countries. Six only are natives of Britain, and these are far inferior in beauty of appearance to many of foreign growth. The *Lycopodiums* were always taken for and called Mosses

by the old botanists; and they do indeed resemble that tribe in many of their external characters, having sessile, smooth, entire, or at most serrated leaves. Their fruit, however, is greatly different; most species of the *Lycopodiums* bear it in terminal scaly spikes, in a few others, among which is our *Lycopodium selago*, the fruit is not confined to the apex of the branches, but is found in the axils of the leaves throughout the whole plant. The root grows from every part of the stem which touches the ground.

LYCOPodium: CLAVATUM.

COMMON CLUB-MOSS. FOX GLOVE. STAG'S-HORN. WOLF'S-CLAW.

(Plate 8, fig. 1.)

CHA.—Stem trailing. Leaves linear, incurved, hair-pointed. Theca in naked, stalked, double spikes. Scales ovate, serrate.

SYN.—*Lycopodium clavatum* of most botanists.—*Muscus clavatus* sen *Lycopodium*, *Ger. Park.*, &c.—*Lycopodium officinale*, *Neek.*—*Lepidotis clavata*, *Beaur.*

FIG.—*E. B.* 224.—*Ger.* 1562.—*Phytologist.* No. 1. p. 1.

DES.—Root fibrous, scattered. Stem branched, several feet long, lying on the ground, bright green. Leaves crowded, linear, curved, with a long, diaphanous, hair-like point. Spike of fruit cylindrical, usually in pairs, yellow, about an inch long, supported upon a rigid, upright, long stem, which is void of leaves, but set at intervals with whorls of very fine, short setæ. Scales of the spikes broadly ovate, pointed, and dentate or serrate. Theca large, round, one to three, attached to the base of each scale, and filled with a very fine yellow powder.

VIR.—For the virtues of this plant, see page 16; in addition to which it is said to be used to ameliorate wines, but its emetic properties render this doubtful.

SIT. and HAB.—On hill-sides, particularly in the northern part of the kingdom, but not ascending to so lofty a situation as some other species. Hoy Hill, Orkney, *Rev. C. Clouston.* Plentiful in the Highlands, in Cumberland, and in North Wales, *Mr. H. C. Watson.* Charlewood Forest, Leicestershire, *Rev. A. Bloom.* Todmorden, Lancashire, *Mr. W. Wilson.* Settle, Yorks., *Mr. J. Tatham.* Derbyshire, *Dr. Howitt.* Coleshill, Warwickshire, *Rev W. Bree.* Notts, *Mr. T. H. Cooper.* Oxfordshire, *Mr. Baxter.* Somerset, *Mr. A. Southby.* Woking Heath, Surrey, between the Canal and Railway, 2 or 300 yards S.W. of the Station-house, *Mr. H. C. Watson.* Lane between Dorking and Leith Hill, and on Addington Hills, beyond Croydon, Surrey; also on the high heathy ground above Tring, Herts, *Mr. W. Pamplin.* Sussex, *Rev. G. E. Smith.* —IRE.: Kelly's Glen, Ballypascorney, and other places on the Dublin Mountains, *Mr. Mackay.* *Dunfermline, Berwick, &c.*

GEO.—In most of the northern parts of Europe and Asia, and from Canada to Pennsylvania in America.

1

2

3

4

5

6

2.—LYCOPODIUM ANNOTINUM.

INTERRUPTED CLUB-MOSS.

(Plate 8, fig. 2.)

CHA.—Stem procumbent. Leaves in five rows, lanceolate, acute, spreading. Spikes simple, scales broadly ovate, imbricated.

SYN.—*Lycopodium annotinum*, Linn., Willd., Spreng., Smith, Hooker, Ehrh., Huds., Lightf., With., Pursh, Gray.—*Lepidotis annotina*, Beauv.

FIG.—*E. B.* 1727.—*Fl. Dan.* 127.—*Dill. Musc.* 63, f. 9.—*Schk. fil.* 162.

DES.—Root of stout and scattered fibres. Stem very long and trailing, dichotomously branched, of a dullish green color, and extending in length from year to year. Branches simple or nearly so when fruitful, upright at first, afterwards becoming decumbent. Leaves in five rows, lanceolate, acute, spreading, entire or very slightly serrated. Fertile spike solitary, sessile, terminal, an inch long, scales very short, very broad, pointed, and imbricated.

Sir J. E. Smith says, that “the scales of the spike of one season falling off, the stem thus left naked gives rise the following season to leaves, but these not being so numerous as in the other parts of the plant, the stem acquires an interrupted habit.” I cannot reconcile this to the appearance of my specimens, but rather suppose that, as in the former species the spike wholly falls off, and the next year’s shoot puts forth more vigorous leaves than those which terminated the whole stem, thus giving the jointed appearance which the plant presents; but I have never seen it in a fresh state, and therefore write this with hesitation.

SIR.—On the highest Welch and Scottish mountains.

HAB.—Pretty frequent between 500 and 850 yards on the mountains of Clova, and the W. of Aberdeenshire; I have never seen it above 900 or below 400 yards; Glen Dole, Forfar, and mountains adjacent; Ben-na-Baird, Loch-na-garr, &c. Aberdeen, *Mr. H. C. Watson*. Freewater, Rosshire, *Mr. Staples*. Hoy Hill, Orkney, *Rev. C. Clouston*. Summit of Cairngorum, *Sir W. J. Hooker*. Still found on Glyder Vawr, Snowdon, but reduced to a solitary root, and when last seen, (1836,) without fructification, *Mr. W. Wilson*. Charnwood Forest, Leicestershire, *Rev. A. Bloxam*. This is remarkable as being the only English habitat recorded. (See “*Naturalist*,” vol. ii. page 135.) Not in Mr. Mackay’s “*Flora Hibernica*.”

GEO.—Europe in mountainous countries; in America, from Canada to Pennsylvania; also in N. Asia.

3.—LYCOPODIUM INUNDATUM.

MARSH CLUB-MOSS.

(Plate 8, fig. 3.)

CHA.—Stem creeping. Branches simple, erect. Leaves and scales linear, acute, curved upwards. Spikes solitary.

SYN.—*Lycopodium inundatum* of botanists.—*Plananthus inundatus*, Beauv.

FIG.—*E. B.* 239.—*Flo. Dan.* 336.—*Dillen Musc* 62, f. 7, (good.)

DES.—Stem very closely appressed to the ground, 1 to 2 inches long, and but slightly branched, bearing fibrous roots all along its lower surface. Branches simple, barren ones decumbent, fertile ones upright. Leaves irregularly placed, crowded, linear lanceolate, acute, all turned upwards. Spikes solitary, terminal, green, 1 inch long, quite erect. Scales linear, dilated at the base, curved upwards, entire or with one or two teeth only in luxuriant specimens.

This plant, which, like the rest of its tribe, is perennial, shows very strikingly the manner of growth of all the creeping species, though there are few of them so rapid in their decay as this. It creeps along the ground, and grows at one end as it decays at the other; thus if its habitat be a level piece of mud, as it generally is on commons, &c., the effect is easily seen in a black mark or line of the decayed plant, sometimes for many inches beyond where it is then vegetating. It ceases to grow in the winter, but continues to decay; thus very many plants are exterminated, and only the vigorous ones have strength to put forth new foliage, of these a very small portion generally remains, and thus it is that the plants are always small in the early part of the season.

SIT.—On wet moors and commons, particularly where turf has been pared.

HAB.—Near Loch Lee, Nairnshire, *Mr. W. Stables*. Near Craig Darrock, Rosshire, *Rev. G. Gordon*. Delamere Forest, Bartington Heath; and Bagueley Moor, Cheshire, *Mr. W. Wilson*. Coleshill, Warwickshire, *Rev. W. Bree*. Valley near Cæsar's Camp, Wimbledon Common, Surrey; bogs near Titchborne Church, Hants, (1836;) Putney Heath; Bagshot Heath; Shirley Common, Surrey; and Keston Heath, Kent, *Mr. W. Pamplin*. Esher Common and Cobham Common, Surrey, *Mr. R. Castle*. Filby, Belton, Yarmouth, Norfolk, *Mr. Paget*. Sussex and South Kent, *Rev. G. E. Smith*. Bovey Heathfield, Devon, *Mr. Babington*. Not in Ireland.

GEO.—Europe, Isles of Bourbon (?) Canada to New York.

4.—LYCOPODIUM SELAGINOIDES.

PRICKLY CLUB-MOSS. MOUNTAIN-MOSS.

(Plate 8, fig. 4.)

CHA.—Stems procumbent. Leaves lanceolate, acute. Spikes large, solitary. Scales ovate, deeply toothed.

SYN.—*Selaginella spinosa*, Beauv.—*Lycopodium selaginoides* of modern authors.—*Bernhardia spinosa*, Gray.

FIG.—*E. B.* 1148.—*Flo. Dan.* 70.—*Dill. Mus.* 68, f. 1.—*Schk. fil.* 165.

DES.—Stems creeping, slightly branched, 2 or 3 inches long. Barren branches delicate, recumbent, simple. Fertile branches upright, rigid, bearing a solitary spike. Leaves lanceolate, acute, toothed, imbricated, bright green. Spike large, oblong, cylindrical,

yellowish, and terminal. Scales much larger and wider than the leaves, and deeply toothed, spreading widely on account of the very large capsules.

This plant shows very well the two sorts of capsules; those in the lower part being what in the *Introduction* are called spores, while the upper capsules contain only a fine powdery mass, considered pollen by some authors, and abortive seeds by others. Mr. T. G. Rylands, alluding to some specimens gathered on Seaforth Common, near Liverpool, says truly, "that plants of this species are of a brighter color than the rest, and that when growing they appear to form small thick tufts about an inch in height and diameter."

SRT.—On mountain sides, and in moist alpine situations.

HAB.—Hoy Hill, Orkney, *Rev. C. Clouston*. Ben Lawers, *Mr. H. C. Cooper*. Ben Lomond, *Mr. W. Leighton*. North coast of Sutherland, at the sea level, *Dr. Johnston*. Rare in Aberdeenshire, *Dr. Murray*. Moray and Rosshire, *Rev. G. Gordon*. Abundant in the Highlands, rising to situations of 1000 yards or more in height, on the Breadalbane mountains, Perthshire. Pretty frequent on the hills of Cumberland, as around Borrowdale, Keswick, Derwentwater, &c., *Mr. H. C. Watson*. Near Richmond, Yorkshire, *Mr. J. Ward*. In wet places among sand-hills on the coast of Anglesea, near Aberffraw, and on the coast of Lancashire, near Southport, *Mr. W. Wilson*. Higher parts of the Tees, *Mr. J. Hogg*. Caernarvonshire, *Mr. J. E. Bowman*. Wyn-ddur, Arddu, Snowdon, *Mr. C. Babington*. Capel Curig, *Dr. Howitt*. Llauberris Pass and Nant Phraeon, *Mr. Watson*. Various parts of Ireland, *Mr. H. C. Mackay*.

Geo.—Sprengel says, only found in Europe at Bremen, Oldenburgh, Silesia, Bavaria, and Switzerland. In Canada and New Hampshire, in America, according to Pursh, who says the American is smaller than the English plant.

5.—LYCOPODIUM ALPINUM.

SAVINE-LEAVED CLUB-MOSS.

(Plate 8, fig. 5.)

CHA.—Stem procumbent, branches fascieled, flat at top. Leaves and scales in four rows.

SYN.—*Lycopodium alpinum of almost all botanists*.—*Lycopodium sabinæ-folium, Pursh*.—*Lepidotis alpina, Beauv.*

FIG.—*E. B.* 234.—*Flo. Dan.* 79.—*Dill. Mus.* 58, f. 2.

DES.—Roots scattered, long, with stout, branched, downy fibres. Stem 2 to 4 feet in length, creeping quite close to the ground, very rigid, irregularly leafy. Branches alternate, set along the stem at uncertain intervals, in an upright, rigid, close or fan-shaped fascicle, level at the top. Leaves blunt, oblong, imbricated in four rows, rather convex. Spikes terminating all the older branches, erect, an inch or less in length, and compact. Scales pointed, broad at the base, tapering upwards, with waved edges, sometimes with two or three teeth, flatter and less rigid than the leaves.

VIR.—According to Sir W. J. Hooker it is used to dye woollen cloths of a yellow color.

SIT.—On the grassy sides of mountains.

HAB.—At 1000 yards of elevation on Carnedd David, Caernarvonshire, probably 1200 yards in Aberdeenshire; also to the summit of Ben Hope, in Sutherland, at 1000 yards or thereabouts, where the climate is probably less genial than that at 1200 yards in Aberdeenshire; to 1150 yards on Ben Nevis, and descending to the base of the mountains. Too plentiful on all the mountain tracts of Scotland to call for particular localities. On most of the Cumberland and Yorkshire mountains, *Mr. H. C. Watson*. Somerset, *Mr. A. Southby*. Near Todmorden, Lancashire, at a very low elevation (a single root only), *Mr. W. Wilson*.—IRE.: Agila and Barnesmore mountains, Donegal, *Mr. E. Murphy*. Barnesmoor Mountain, and Mourne Mountain, *Mr. Mackay*. Brandon Mountain, *Mr. W. Wilson*. *on the Foreland Hill, Devon.*

GEO.—All the northern and mountainous part of Europe, as Lapland, Germany, Switzerland, Pyrenees, the Tyrol, Sweden, Norway, Russia, &c. Also in Canada and Siberia.

6.—LYCOPODIUM SELAGO.

FIR CLUB-MOSS. UPRIGHT FIR-MOSS.

(Plate 8, fig. 6.)

CHA.—Stem erect, dichotomously branched, flat at top. Leaves in eight rows. Thecae axillary.

SYN.—*Plananthus selago*, *Beauv.*—*Selago vulgaris*, *Dillw.*—*Lycopodium abietiforme*, *Gray.*—*Lycopodium selago* of other botanists.

FIG.—*E. B.* 233.—*Flo. Dan.* 104.—*Dill. Mus.* t. 56, f. 1.

DES.—Root tufted, fibrous. Stems 2 to 6 inches high, growing quite erect, one issuing only from the root, and this becoming divided dichotomously until they form a cluster of from six to ten ultimate divisions; the upper fruitful branches are, however, scarcely more than forked. Leaves in eight rows, of a dark shining green color, crowded, lanceolate, entire, acute, convex on the outer side, a little spreading, and curved upwards. The fruit is not borne in a terminal spike, as in the other species, but in the axils of the common leaves, all down the upper part of the stem. Capsules large, kidney-shaped, regularly two-valved, opening by a transverse fissure, and scattering minute, yellow, globular, smooth spores.

This plant is likewise viviparous, producing not only capsules of seeds, but occasionally also curious petioled buds, which consist of three or four differently-sized ovate leaves; they are irregularly placed in the axils of the common leaves, that is, in the place of the capsules.

SIT.—On mountain sides, &c.

HAB.—It attains the summit of Ben-na-Muich-dhn, the loftiest of the northern Grampians or Cairngorum range, and the second summit of Britain (4320 feet). Common everywhere on the hilly tracts of Britain, especially the Scottish Highlands.—ENG.: Helvellyn, Skiddaw, &c.; on the loftiest rocks of Dartmoor, and above Edale Chapel, Derbyshire, *Mr. H. C. Watson*. Common about Settle, *Mr. J. Tatham*. Coleshill, Warwickshire (rare), *Rev. W. Bree*. Wensley dale, Yorkshire, *Mr. J. Ward*. Once seen on Woolston Moss, near Warrington, *Mr. Wilson*. Waldron Down, Sussex, and near Bristol, *Miss Worsley*. *Peak Hill, Sidmouth, Devon 1856.*

1

2

3

4

5

6

8

7

Shotover Hill, Oxon, Mr. Baxter. Mansfield Forest, near the Blidworth Gate, Mr. T. H. Cooper.—WAL.: Frequent on the Welch mountains, where a variety is found with the leaves widely spreading, Mr. W. Wilson.—IRE.: Lough Bray and mountains, in the south of Ireland (frequent), *Irish Flora*. Known in Kerry as Virgin Mary's Furze.

GEO.—Over Europe and North America. (Not in Pursh.)

EQUISETUM, Linn. HORSE-TAIL.

A, spike of fruit of *Equisetum palustre*. B, portion of the stem and branches of ditto. C, receptacle magnified. D, under side of a scale of ditto, showing the scattering of the spores. E, spore in its young state. F, ditto more advanced, and unfolding its filiform appendages. G, ditto spread out and with abortive granules attached to the filaments. H, abortive granules. I, transverse section of the stem of *Equisetum variegatum*. J, ditto of *Equisetum fluviatile*. K, longitudinal section of ditto. L, sheath of *Equisetum Drummondii*. M, longitudinal section of stem of *Equisetum limosum*. N, particles of silex on *Equisetum Drummondii*.

A widely distributed but not very extensive genus, which inhabits for the most part temperate and cold countries. The species now living are all small plants, but the fossil remains of the *Equisetaceæ* show that at some former period gigantic specimens must have been frequent. Our plants prefer watery situations and strong soil. They may be divided into sections as follows:—

- * Fertile stems naked, succeeded by branched barren ones.
- ** Fertile stems branched from their first growth.
- *** Fertile stems not branched at first, but finally becoming so.
- **** Fertile stems always remaining simple, barren stems the same.

Equisetum fluviatile, *Drummondii*, and *arvense*, belong to the first section; *E. sylvaticum* to the second; *E. palustre* and *limosum* to the third; *E. variegatum* and *hyemale* to the fourth.

1.—EQUISETUM FLUVIATILE.

GREAT HORSE-TAIL. WATER HORSE-TAIL.

(Plate 9, fig. 1.)

CHA.—Barren stems erect, with thirty to forty branches in each whorl. Fertile stems with loose sheaths, having numerous teeth.

SYN.—*Equisetum fluviatile*, Linn., Willd., Smith, Hook., Bolt., Huds., Lightf., With., Gray.—*E. telmateia*, Ehrh., Flo. Dan.—*E. cburncum*, Roth, Schr.—*E. majus*, Ray, Ger.—*E. maximum*, Lam.

FIG.—*E. B.* 2022.—*Bolt.* 36, 37.—*Ger. Her.* 1113.—*Flo. Dan.* 1469.

DES.—Barren stem 2 to 4 feet high, quite erect, white, succulent, surrounded by whorls of from thirty to forty branches. Branches rapidly growing upon the stem as soon as it issues from the ground, giving it soon a broad-topped appearance. In its future growth this blunt character is lost, the main stem becoming elongated, and the branches are then long, slender, simple, jointed, ascending, with channels along their surface, and at the angles of these other very minute ones. Fertile stems 4 to 6 inches high, arising in March or April, and decaying as the barren stems arise, reddish white, extremely succulent, and wholly without branches at any time. Their sheaths four to six in number, are nearly an inch long, and generally so close together as to overlap each other, very deeply, sharply, and numerous toothed. Catkin large and conical.

Withering says, “fertile stems sometimes leafy.” He ought rather to have said, barren stems sometimes fruitful; as a catkin is often found in the middle or latter part of summer terminating it, particularly if the weather has been dry for some time previously; in fact it may be produced at any time with such cultivated plants as grow in pots, merely by removing the pots from the watery situation in which they are usually placed into a drier spot of ground. Mr. W. Wilson attributes this state of the plant to drought as here stated, and adds that he has seen a specimen gathered near Bangor where this catkin was topped by a prolongation of the branched frond, (July, 1836.)

The name Fluvatile is not so applicable to this species as it would have been to some others; it is rarely found on the banks of rivers or ponds, nor do I remember ever having seen it growing in the water. It rather affects strong loamy damp ground, clayey banks, and swampy bogs.

HAB.—Very abundant in some parts of England, as about London, in Hants, Bucks, &c.; but Mr. Watson thinks scarcely a common plant generally.

GEO.—Europe, Siberia, North America.

2.—EQUISETUM DRUMMONDII.

BLUNT-TOPPED HORSE-TAIL.

(Plate 9, fig. 2.)

CHA.—Barren stem blunt, erect, with about twelve branches. Fertile stems with prickly-toothed sheaths.

SYN.—*Equisetum Drummondii*, Hook. in *E. B. suppl.*; Mack., *Fl. Hib.*

FIG.—*E. B. suppl.*, t. 2777.

DES.—Barren stem exceedingly delicate, finely tapering upwards, very rough on the angles, with white and shining particles of silice, 12 inches high, of a pale, lightish green, particularly the scales, which widen upwards, six or eight in number, rather close together, with long, black, terminal teeth.

This plant differs from *Equisetum arvense* in its more glaucous green color very much more delicate habit both of stem and branches, and blunter outline. The fertile frond is much more rigid in texture, with harder, whiter, and more numerous-toothed sheaths, and the points of the teeth are more diaphanous than in the next species.

HAB.—First found by Mr. T. Drummond at Wolf Hill, the seat of W. Thompson, Esq., near Belfast.

3.—EQUISETUM ARVENSE.

CORN HORSE-TAIL.

(Plate 9, fig. 3.)

CHA.—Barren stem taper-pointed, decumbent. Sheaths of the fertile stem three or four, distant, loose.

SYN.—*Equisetum arvense*, Linn., Willd., Smith, Hook., Bolt., Ehrh., Huds., Lightf., With., Pursh., Mack., Gray.—*Equisetum segetale*, Ger.

FIG.—E. B. 2020.—Bolt. 34.—Flo. Lon. 64.—Ger. Her. 1114.—Park. 1202.

DES.—Root branched, creeping. Main stem of the barren frond procumbent, tapering to the end, sometimes very long, pointed, rough, with whorls of branches all the way down, and forked at the base; that is to say, two or more fronds springing from the same part of the root. Branches simple, varying much in number, fourteen or sixteen, if all are present, but generally not more than eight or ten, or even six, at the upper part of the frond. They are dark green, rough, four channelled, with simple angles. Fertile fronds appearing before the others, light brown, with four or five distant, deeply-toothed sheaths. The teeth are sharp, ribbed, and rather dark colored, particularly round their edges. As the fertile frond comes to maturity, the sheaths decay from the point downwards; thus their black tooth is often tipped with white, and surrounded with a diaphanous membrane, particularly the upper sheaths, which are larger and longer than those below.

The plant puts on very different characters in different circumstances; sometimes it appears as a cylindrical pointed stem, without any branches—this is its early state, for it does not throw out branches immediately, as in *Equisetum fluviatile*, therefore it never appears with a densely leafy, obtuse frond as that does. On strong soil, and in shady situations, as when hid among growing corn, the branches become exceedingly long, scattered in habit, and often geniculated, in which state it is represented by Gerard as *Equisetum segetale*. The closeness and number of the sheaths of the fertile stem are by no means a sure diagnostic of any of our species; the remarks respecting them, therefore, are more general than specific information.

SIT. and HAB.—In corn fields, sandy banks, waste ground, &c. This species is so common, and so difficult to eradicate, as to be a very troublesome weed.

GEO.—Found equally in Europe, Asia, and North America.

4.—EQUISETUM SYLVATICUM.

WOOD HORSE-TAIL.

(Plate 9, fig. 4.)

CHA.—Stem erect. Branches compound, deflexed. Sheaths loose.

SYN.—*Equisetum sylvaticum* of all modern botanists.FIG.—*E. B.* 1874.—*Bolt.* 32, 33.—*Flo. Dan.* 1182.—*Schk. fl.* 166.

DES.—Stem erect, from 6 inches to 2 feet high, branched, bright green. Branches compound, slender, smooth, drooping at the ends, and whorled, the lower part of the stem without branches. Catkins ovate, erect, stalked, and terminating the stem, borne early in the season, and dying away long before the remaining part, as is the case with all the following species; it is very rarely, however, found in fruit. Sheaths deeper colored than the stem.

SIR.—In woods and shady places, chiefly in the North,

HAB.—Rosshire and Moray, *Rev. G. Gordon*. Orkney, *Rev. C. Clouston*. Frequent in the Highlands of Scotland, *Mr. W. Witson*. Near Richmond, Yorkshire, *Mr. Ward*. About Settle, Yorkshire, (scarcely), *Mr. J. Tatham*. Near Leeds, *Mr. Denny*. Forge Valley, near Scarborough, Yorkshire; near the ruins of Dale Abbey, and Southwood, near Calke Abbey, Derbyshire, *Rev. A. Bloxam*. Egerton, near Bolton, *Mr. W. Christy*. Cumberland, Cheshire, Lancashire, *Mr. Watson*. Benthall Edge, Salop, *Mr. W. Leighton*. Cromford Moor, Derbyshire, *Dr. Howitt*. In Bagley Wood, between Oxford and Abingdon, *Mr. W. Baxter*. Hampstead Heath and fields towards Hendon, Middlesex, *Mr. W. Pamplin*. Sussex, *Rev. G. E. Smith*. Somerset, *Mr. Southby*. Moist woods, Kelly's Glen, Ballynascorney, *Mr. O. Kelly*. Abundant in the North of Ireland, *Mr. Mackay*.

GEO.—All Germany, Prussia, Holland, and Switzerland. From New York to Virginia, &c., and in North Asia.

5.—EQUISETUM LIMOSUM.

SMOOTH NAKED HORSE-TAIL.

(Plate 9, fig. 5.)

CHA.—Stem erect, naked or branched, smooth. Sheaths short, appressed. Teeth numerous.

SYN.—*Equisetum limosum*, *Linn.*, *Willd.*, *Smith*, *Hook.*, *Bolt.*, *Huds.*, *Lightf.*, *With.*, *Mack.*, *Gray*.—*Equisetum polymorphum*, *Schr.*—*Equisetum heleocharis*, *Ehrh.*

FIG.—*E. B.* 929.—*Flo. Dan.* 1184.—*Bolt.* 38.

DES.—Root much creeping, with scattered fibres. Stem erect, quite smooth, striated, but not channelled, generally naked, but sometimes putting out a few branches late in the season, which are smooth, simple, and ascending. Catkin terminal, broad and short, for the most part sessile in the upper sheath. Sheaths short, close pressed to the stem, with very numerous short brown teeth.

Often confounded with *Equisetum palustre*, of which by some of the older botanists it was considered only a variety. It is, however, very distinct, and may easily be distinguished by not bearing branches till late in the season, after the catkin has decayed; its branches also are less numerous, shorter, and either scattered over the plant or in irregular whorls; it has shorter and more numerous toothed sheaths, which are pressed close to the stem. The whole plant is smoother, and has shorter, thicker, and nearly sessile catkins.

SIT.—In low swampy ground, sides of streams, &c.

HAB.—Not so common as *Equisetum palustre*, but pretty generally distributed. Moray and Rosshire, *Rev. G. Gordon*. Tees, *Mr. J. Hogg*. Cheshire, Lancashire, and Cumberland, *Mr. H. C. Watson*. Wensley Dale, Yorkshire, *Mr. J. Ward*. Needwood Forest, Staffordshire, and Gamlingay Bogs, Cambridgeshire. River Severn, near Shrewsbury, Hancott Pool, ditto, *Mr. W. Leighton*. Near Wrexham, *Mr. J. E. Bowman*. Warwickshire, *Rev. W. Bree*. Leicestershire, *Rev. A. Bloxam*. Derbyshire, *Dr. Howitt*. Norfolk, *Miss Bett*. Somerset, *Mr. Southby*. Sussex and South Kent, *Rev. G. E. Smith*. Near Bristol, *Miss Worsley*. Frequent in Ireland, *Mr. Mackay*.

GEO.—Holland, Switzerland, and other parts of Europe.

6.—EQUISETUM PALUSTRE.

MARSH HORSE-TAIL.

(Plate 9, fig. 6.)

CHA.—Stem erect, naked or branched, rough. Sheaths long, loose. Teeth few and long.

SYN.—*Equisetum palustre* of all English botanists.—*Equisetum nodosum*, Schr.—*Equisetum ramosum*, Schl.

FIG.—*E. B.* 2021.—*Bolt.* 35.—*Flo. Dan.* 1183.—*Lob. Icon.* 795.—*Ger. Her.* 1114.—*Schk.* 168, 169.

DES.—Root creeping. Stem upright, branched throughout, 6 to 12 inches high, dark green, deeply channelled. Branches five-sided, simple, ascending, six to ten in a whorl, a less number of and shorter branches upwards. Catkins terminal, cylindrical, tapering, on a long stalk, erect, found in May and June, sometimes before the branches, at other times appearing long after the stem becomes branched. Sheaths large, loose, with a few long tapering black teeth.

β (*alpinum*.) smaller, upper branches abortive.

γ (*polystachion*.) upper branches elongated and fruitful.

The second variety is always found in such situations as convince us that its peculiar conformation arises from its being nipped by frost or cropped by cattle, especially as when thus proliferous, the main stem is almost always injured at the top; a proof that here, as often is the case with Flowering Plants, the early flowers being by any cause destroyed, the plant makes an effort to repair the loss at a later season of the year by producing others.

SIT and HAB.—Very common in ponds, wet valleys, water-courses, &c.
 β :—Breadalbane Mountain, Perthshire, at 3000 feet high, *Mr. H. C. Watson*.
 γ : No certain habitat of this can be given, because it is an accidental state of the plant, and not a permanent variety. I have often found it in Richmond Park, Surrey, and by the side of the Lea River at Stratford, Essex.

GEO.—Common throughout Europe, and in North America.

7.—EQUISETUM VARIEGATUM.

VARIEGATED ROUGH HORSE-TAIL.

(Plate 9, fig. 7.)

CHA.—Stems procumbent, rough. Sheaths black at top. Teeth few, white, and persistent.

SYN.—*Equisetum variegatum*, *Willd.*, *Schk.*, *Smith*, *Hook.*, *Mack.*—*Equisetum arenarium*, *of authors.*—*Equisetum tenue*, *Hopp.*

FIG.—*E. B.* 1987.

DES.—Root very woolly. Stem branched at the base only, rather procumbent in habit, 4 to 6 inches long, of a green color, rough and channelled. Catkins terminal, ovate, at first black and sessile, afterwards long stalked, yellow and brown. Sheaths of the stem widening at top, black only at their upper part, which is sharply, but not numerously toothed, the upper sheath of the stem being much larger and more spreading than the rest.

Its smaller size, recumbent habit, differently-colored sheaths, with their prominent and permanent teeth, serve to distinguish this from *E. hyemale*.

SIT.—On the sandy sea-shore in the north of the kingdom.

HAB.—ENG.: Sand hills on the Cheshire coast, between Hoylake and the Rock Fort, *Mr. H. C. Watson*. Wardrew, Northumberland, (abundant,) *Mr. Winch*. Southport, Lanc., *Mr. W. Wilson*. Near the Powder Magazine, in Wallasey, opposite Liverpool, *Mr. J. E. Bowman*. Bootle Sands, near Liverpool, *Mr. Rylands*. Near Winch Bridge, Teesdale, *Mr. Bowman*.—SCO.: Sands of Barry, Forfarshire, *Dr. Greville*. Near Avoch, Rosshire, *Rev. G. Gordon*.—IRE.: Portmarnock, opposite Baldoyle, *Dr. Taylor*. Mueruss, Killarney (a tall var.), *Mr. W. Wilson*. Moist banks near a waterfall at the upper end of Colin Glen, Belfast, *Mr. Mackay*. Ballyharrigan Glen, near Dimgiven, *Mr. D. Moore*.

GEO.—Switzerland, Italy, France, Alsatia, &c.

8.—EQUISETUM HYEMALE.

ROUGH HORSE-TAIL. SHAVE GRASS. DUTCH RUSH.

(Plate 9, fig. 8.)

CHA.—Stem erect, rough, deeply striated. Sheaths short, appressed, black at each end. Teeth deciduous.

SYN.—*Equisetum hyemale*, *Linn.*, *Willd.*, *Smith*, *Hook.*, *Lightf.*, *Ehrh.*, *Huds.*, *With.*, *Pursh*, *Mack.*, *Gray*. (Not of *Bory.*)—*Equisetum nudum*, *Ray*, *Gerard*.

FIG.—*E. B.* 915.—*Hook.* in *Flo. Lon.* 161.—*Ger. Her.* 1113.—*Bolt.* 39.—*Schk. fil.* 172.

DES.—Root black, branched. Stems erect, of a very dark green, without whorls of branches, but forked and divided at the base, 2 to 3 feet high, regularly and numerous furrowed. Sheaths 2 to 3 inches distant from each other, very closely pressed to the stem, short, with a black rim at the top and bottom of each. Teeth of the scales black and deciduous.

It is surprising that this plant, so valuable in a general as well as a commercial point of view, is not cultivated along our sandy coasts, where it would grow luxuriantly and rapidly, forming a strong embankment, and yielding a considerable profit. The Dutch are well acquainted with the value of its long and matted roots in restraining the wasting effects of the ocean, which would soon undermine their dykes were it not for the *Equisetum hyemale* which is planted upon them. At the proper season it is cut down and exported to other countries, where its naked and flinty stems are used for polishing domestic utensils, furniture, marble, &c. It is here sold as Dutch rush, (not Dutch rushes, which are *Scirpus glaucus*, or sometimes *Scirpus palustris*; the former being used for the bottoms of chairs, the latter by coopers to stop leakages.)

So abundant is the silex upon both the inner and outer cuticle of the stem, that it is said the whole of its vegetable matter may be removed without destroying the shape of the plant. Every part of it is a very beautiful object under the microscope.

SIT.—In woods and boggy places; rather rare, particularly in the South.

HAB.—ENG.: Hawthorn Dean, Durham, *Mr. T. H. Cooper*. Scotswood Dean, near Newcastle, *Mr. Bowman*. Near Over, Cheshire, *Mr. W. Wilson*. Common near Halifax, *Mr. R. Leyland*. In a dell at Bitterley, below the Clce Hills, Salop. Forge valley, near Scarboro', Yorkshire. In a small stream at the bottom of Grace Dieu Wood, Charnwood Forest, Leicestershire, *Rev. A. Bloxam*. South Kent, *Rev. G. E. Smith*.—WAL.: Near Wrexham, *Mr. J. E. Bowman*.—SCO.: Edinburgh, in the stream just below Roslyn Castle, *Mr. H. C. Watson*. Moray and Rosshire, *Rev. G. Gordon*. Wood at Corra Linn, Lanarks, *Mr. C. C. Babington*.—IRE.: Tyrone, *Mr. Shuttleworth*. Wood at Leislip Castle, near Dublin. Powerscourt, and around Dublin, *Mr. Mackay*.

GEO.—All Germany, Holland, and Switzerland. From Canada to Virginia, and in Asia.

APPENDIX.

CULTIVATION OF FERNS GENERALLY.

THIS tribe of plants was, but a few years ago, scarcely known in cultivation. Lately, however, it has been considerably sought after; and as little has hitherto been written on the culture of Ferns, or the selection of species, the following general observations will, I trust, be acceptable. It is right to observe, that for a great part of the list, and some of the remarks which follow, I am indebted to Mr. Bevis, of the Botanic Garden, Regent's Park, a well-known and ardent cultivator of the Ferns.

Sowing.—For sowing Fern seed the spring of the year is to be preferred. The pots in which it is to be sown should be of a small size, (say 4s.,) both for the sake of convenience, and because they hold less moisture, and allow a better circulation. Fill the pots half full of fine broken pot-shreds; that is, garden pots broken into small pieces. Over these lay a portion of Sphagnum, or other porous moss; and then fill the pot to within half an inch of the top, with soil prepared in the following manner:—Take three parts of loose or unadhesive peat earth, (that is to be chosen which lies about an inch beneath the surface, where it is not too much decayed;) put it into a pan of boiling water to kill any vegetable matter that may be alive, or small worms, which become very troublesome as the Fern seeds vegetate; then let it get dry enough to rub through the hand, but not very finely; to this powdered peat add one-third of white sand, mixing both articles well together. When the pots are filled to within half an inch of the top, give them a little water, on which sow the seeds, taking care not to water them afterwards over the top. The seeds should be sown thinly—if thickly sown, or too much water be given, they are apt to *fog*, or kill each other, before they are large enough for potting or pricking off. After sowing lay a piece of common glass over the top of the pot, and set it in a saucer, taking care not to let the saucer be without water, and place it in a light but shaded place. When the plants have shown the first leaf, a little air may be admitted—after which, should they be too thick, they may be thinned by taking them up in small patches with the point of a knife, and transplanting them into another pot, prepared as the former, but made fine by sifting the soil; when large enough they may be divided a second time. Many persons mix a great portion of broken brick in the seed pot, upon which the seeds vegetate well, but the difficulty of transplanting therefrom is very great. It is of course necessary to know whether the species belong to the green-house or stove, that the vegetating seed may be placed in a congenial climate, observing only that a dry air and direct sunshine is to be, as much as possible, avoided.

After-cultivation.—Ferns of different habit require a different treatment. This is in some degree accordant with their natural soil and places of growth, yet not wholly so. Numerous of the British Ferns, although they naturally live in bleak and exposed situations, yet when under culture require some degree of protection. So also notwithstanding some of them seem naturally to prefer the

interstices of brick walls, such for example, as *Grammitis ceterach*, they will not flourish in brick rubbish. To treat this little understood part of the subject intelligibly, and to show the extent to which the foregoing observation applies, it is advisable to divide the Ferns into various distinct sections, as follows:—

STOVE AND GREEN-HOUSE FERNS.

Ferns with rhizomas.—These in their native habitats are, in many instances, epiphytes, deriving their nourishment chiefly from the air, or from other scanty sources. If potted they require much less water than others, the soil should be porous, and the pots should be filled half full of turfy peat. Most of this division thrive very well suspended on blocks of wood, in a warm moist atmosphere, but should be watered very sparingly in the winter season. The small, creeping, entire-leaved species thrive as epiphytes, as they run a long way in the season. These are readily increased by cuttings. If potted they seldom show any fructification, owing to their stunted habits.

Ferns with crowns.—Those Ferns with crowns from which the fronds issue, require a soil made finer, with a greater depth. Care must be taken in not covering their crowns, which is certain death to the greater number of the species; they likewise require more frequent watering as they are sooner affected by drought. Many of this section produce bulbs upon their leaves, from which they readily increase. Others root from the tip of their fronds, without forming bulbs; such is the case with many of the *Aspleniums*. This section requires care in dividing, as by this mode they seldom make good plants; it is preferable to increase them from seed, as they grow more freely, and make finer plants. The *Gymnogrammas* should be potted in loam, as they are very apt to damp if potted in peat soil; in fact, most Ferns which have various-colored leaves prefer a loamy soil. No kind of manure should be at any time given to Ferns; even weak lime water should be avoided, as great havoc is sometimes made with it.

Ferns with thick fleshy roots.—Of these there are but a few species. All those of *Marattia* have a curious sealy cormus, resembling that of the genus *Zamia*, from which are produced strong thick fleshy roots. They prefer loam and peat, and are easily injured by shifting, as the soil is very apt to drop from them when they are turned out of the pot. If kept too hot they are apt to drop their leaves; they do best in a temperature not exceeding 60°. *Danæas* resemble the last genus, and require the same treatment. Care must be taken to give them free draining.

Arborescent Ferns.—In general these do not thrive well; they require a very moist atmosphere. Their stems should always be bound with *Sphagnum*, or they soon look very sickly, as they are covered all up their stem with spongeoles, by which they, in a great degree, receive their nourishment. They should be well drained, and freely syringed on their stems to keep the moss moist.

Numerous of the Ferns under stove culture are infested with a species of thrips; others have a rusty appearance, often laid to that insect, but which appears to arise from being kept too warm—the green-house species, mixed with those properly belonging to the stove, being always first attacked. Other insects seldom attack them, with the exception of brown scale and slugs, which are soon destroyed. The best remedy for the thrips is the vapor arising from sulphur sprinkled over the flues or pipes.

HARDY FERNS.

Hardy Ferns should always have a prepared soil on a sloping north bank, where they are seen to the greatest advantage. Some of the strong *Aspidiums* will grow in any common soil, where the small species would soon perish. They prefer a shady place, but do not like the drip of trees, nor stagnant water about them. The border or soil should be made of one-fourth coarse grit or river sand, with three parts peat or bog earth, chopped well in pieces, but not sifted—this should be a foot deep. If a very damp place, the border should have a layer of broken brick below to drain it, if intended for the more choice species; for although *Osmundas*, *Aspidium thelipteris*, and *Blechnum boreale*, will grow in the water, they thrive better a small distance off, and you gain the advantage of growing all the species in the same border. Some of the smaller species should be raised on mounds above the border to keep them drier; the best way is to place four or five stones edgewise, thereby forming a hollow in the centre, putting a little drainage in the bottom. Small species prefer shallow soil; they are likewise benefited by placing a bell glass over them, to retain the humidity of the atmosphere. Many of them it is difficult to find situations suitable for, without covering both in summer and winter; such for instance as *Adiantum capillus-Veneris*; *Asplenium alternifolium*, *viride*, *trichomanes*, *septentrionale*, and *marinum*; *Hymenophyllum Wilsoni* and *Tunbridgense*; *Trichomanes brevisetum*; *Ceterach*, and many small foreign species; likewise the *Lycopodiums*, the hardy species of which may all be grown. *Hymenophyllums* and *Trichomanes* require but little soil; they thrive best fastened on a piece of porous stone, over which has been shaken a little sand. They should always be covered with a glass, and kept very moist and shaded, being inhabitants of wet dripping rocks. Many of the other small species grow in drier situations, even on sunny walls, but they are always finer in the shade; from such situations they are difficult to remove, owing to the roots penetrating the crevices of the wall or rock, and take a long time to get thoroughly established in a new situation. The *Botrychiums* are also removed with difficulty; they require a good drainage.

Fern Houses.—The plan adopted by Mr. Ward, (of Wellclose Square, London,) is deserving of particular attention from many causes, independent of the cultivation of the Ferns. The principle established by this excellent and well-known botanist is, that a constant renewal of air is not necessary for the well-being of plants. Thus if a plant be inclosed in a glass case, watered, and then the case closed up air-tight, the moisture which evaporates having no means of flying off will be condensed on the sides of the case, and trickling down will moisten the plants a second time, only to be evaporated and condensed again and again each succeeding day. Also, the air which is necessarily included in the cases does not become unfit for the use of the plant. That these are established facts may be easily proved by planting a Fern or a Moss in a phial, well corking and sealing the phial, and suffering it to remain in this state for a length of time. This may appear curious rather than useful, and on so small a scale as that of a phial it really is so; but the same principle holds good to any convenient extent, and a glass jar of many gallons, or a box with a glass top of any moderate size, may be thus stored with numerous plants, and made to form a highly interesting parlor ornament, and that without the plants requiring the least care or atten-

tion, except to remove decayed parts, or train up a too-exuberant growth. Even were the discovery capable of no further extent, it would be most invaluable in the transport of plants from one country to another, preserving them from the vicissitude of season, from the effect of salt spray, (so detrimental to most plants,) and from the inattention of their temporary guardians. Indeed Mr. Ward's air-tight cases are now universally employed for the transit of living plants. We would advert, moreover, to the injurious effect of a contaminated atmosphere upon plants, and remind our city friends of their repeated disappointments in window culture, or of their abortive attempts to ruralize their back court-yard. Mr. Ward's plan ensures them success; it is only necessary to cover it with glass—to have but one door of ingress, and that seldom to be used—and to stock the covered space with any plants that can endure a shady situation, (among which the Ferns stand pre-eminent,) and they have at once a beautiful green-house. It may, perhaps, be but a glass closet attached to the outside of an ordinary window, or it may extend the width of the house—in either case success is certain. Those who are desirous of learning more upon the subject may consult a little work by Mr. Ward, entitled, "Growing of Plants in Closed Cases;" and which is just published by Mr. Voorst, Paternoster Row.

SELECT LIST OF FERNS.

STOVE FERNS.

ACROSTICHUM

crinitum
flagellifolium
lingua
nicotianifolium
simplex
scolopendrium
stemaria
villosum
latifolium
dimorphum
aureum
longifolium

ANTROPHIUM

lanceolatum

ADIANTUM

eocinnum
cordatum
eristatum
cuneatum
falcatum
fragile
macrophyllum
obliquum
obtusatum
pubescens
radiatum
serrulatum
tenerum
trapeziforme
varium

ALSOPHILA

Bisphamii

ASPLENIUM

alatum
auriculatum
biauratum
bisectum
brasiliense

contiguum
cuneatum
cultrifolium
compressum
cicutarium
dentatum
fragrans
laciniatum
monanthum
pulchrum
pumilum
pubescens
praemorsum
radicans
rhizophorum
salicifolium
Sheppardii
striatum
serratum
zamiæfolium

ALLANTODIA

Scandiua

ANEMLA

phyllitides
laciniata
eollina
radicans

ASPIDIUM

appendiculatum
albo-punctatum
aristatum
coriaceum
crinitum
exaltatum
falcatum
fraxinifolium
hispidum
hypocrepis
lucens
maerum
macrophyllum

mollis
mucronatum
parasiticum
patens
pectinalum
pennigerum
pubescens
pungens
parasiticum
rhyzophyllum
semicordatum
serra
Sprengelii
tuberosum
trifoliatum
unitum

BLECHNUM

angustifolium
brasiliense
eocorvadense
gracile
hiatum
intergerrimum
longifolium
lanecolum
occidentale
pectinatum
polypodiifolium

CERATOPTERIS

thalictroides
CHEILANTHES
dicksonoides
lendigera
microphylla
tenuifolia
farinosa

CÆNOPTERIS

rhyzophylla
trilobata
funiculacea

DIDYMOCHLÆNA

simosa
DANÆA
alata
DIPLAZIUM
arboreum
decussatum
grandifolium
plantagineum

DICKSONIA

auricoma
arborea

DAVALLIA

corcovadense

GYMNOGRAMMA

chrysophylla
calomclanos
ochracea
peruviana
polypodioides
rufa
sulphurea
trifoliata
tartarica
asplenoides

HEMIONITIS

palmata

LYCIDIUM

circinatum
microphyllum
volubile

LOMARIA

falcata
longifolia
nuda
Plumieri

LYCOPODIUM

apothecium
circinale
ciliare
stoloniferum
Willdenowii

cordifolium
MENISCIUM
 sorbifolium
 palustre
MARATTIA
 laevis
 alata
NOTHOCHLÆNA
 nivea
 rufa
 sinuosa
 trichomanoides
NEPHOBOLUS
 nervata
 pertusa
 sinensis
OPIHOGLOSSUM
 petiolatum
 reticulatum
PSILOTUM
 triquetrum
POLYBOTRYA
 cervina
 cylindrica
PLEOPELTIS
 maerocarpa
 latifolia
 angusta
PTERIS
 arguta
 biaurita
 cruciata
 flexuosa
 discolor
 denticulata
 gigantea
 geraniifolia
 grandifolia
 hastata
 intrimarginalis
 leptophyllus
 brasiliensis
 nemoralis
 palmata
 pedata
 Plumieri
 læta
 sagittifolia
POLYPODIUM
 aureum
 angustifolium
 ariolatum
 Barometz
 curvatum
 crenatum
 scandens
 dissimile
 diversifolium
 decumanum
 deflexum
 effusum
 hastatum
 difforme
 irioides
 incanum
 lævigatum
 lycopodioides
 lygodioides
 loraceum
 myrsifolium
 otides
 olivaceum
 fraxinifolium
 piloselloides
 exiguum
 pectinatum
 phymatodes
 glaucum
 quercifolium
 repens

repandum
 rhyzophyllum
 reptans
 ramosum
 polyanthos
 serpens
 Schkuhrii
 trichomanoides
 tetragonum
 phyllitides
 sphorodoearpon
 cjeutarium
 pendulum
 enoodes
 latipes

GREEN-HOUSE FERNs.

ACROSTICHUM
 alaicorne
ADIANTUM
 assimile
 capillus-veneris
 formosum
 hispidulum
 reniforme
 moritzianum
 athiopieum
 venosum
ALSOPHILA
 australis
ASPLENIUM
 acutum
 diversifolium
 flabellifolium
 lucidum
 nidus
 palmatum
 bipartitum
 Petrarchæ
ALLANTODIA
 australis
 umbrosa
 strigosa
 axillaris
 tenera
ASPIDIUM
 aciculatum
 elongatum
 auriculatum
 lætevirens
BALANTIMUM
 eulcium
BLECHNUM
 australe
 serrulatum
 striatum
 cartilagineum
CIBOTIUM
 Bellardieri
CHEILANTHES
 fragrans
 odora
 caudata
DOODIA
 aspera
 media
 maxima
 eandata
 kunthiana
DAVALLEA
 canariensis
 gibberosa
 pyxydata
 dubia
LOMARIA

nuda
 lanceolata
 procera
 minor
 Pattersonii
LINDSÆA
 media
 triehomancides
NOTHOCILÆNA
 lanuginosa
 marantæa
 distans
 pumila
 tenera
 hirsuta
 profusa
NEPHOBOLUS
 rupestris
 confluent
PHYSEMATIUM
 molle
PTERIS
 cretica
 erenata
 esculenta
 falcata
 hastata
 longifolia
 serrulata
 vespertilionis
 umbrosa
 tremula
POLYPODIUM
 dripanum
 Bellardieri
 proliferum
 tenellum
TODIA
 africana
 australis
TRICHOMANES
 brevisetum
WOODWARDIA
 radicans
HYMENOPHYLLUM
 Wilsoni
 Tunbridgense

HARDY FERNS.

ALLOSURUS
 crispus
ADIANTUM
 pedatum
ASPLENIUM
 adiantum nigrum
 ebumum
 filix-femina
 fontanum
 alternifolium
 lanceolatum
 Michauxii
 marinum
 ruta-muraria
 septentrionale
 trichomanes
 viride
 athyrium
ASPIDIUM
 aculeatum
 acrostichoides
 atomarium
 bulbiferum
 cristatum
 dilatatum
 dimetorum ?
 fuscatum
 fragrans
 filix-mas
 Goldianum
 intermedium
 irriguum ?
 lobatum
 laneastriense
 lonchitis
 montanum
 marginale
 novæboracense
 obtusum
 oreopteris
 rigidum
 rhyzophyllum
 spinulosum
 thelypteris
BLECHNUM
 boreale
BOTRYCHIUM
 dissectum
 fumaroides
 lunaria
 virgineum
CHEILANTHES
 vestita
CYSTOPTERIS
 dentata
 fragilis
 regia
DICKSONIA
 pilosiuscula
GRAMMITIS
 ceterach
LYCOPODIUM
 alpinum
 annotinum
 complanatum
 elavatum
 denticulatum
 dendroideum
 helveticum
 inundatum
 selago
 sclaginoides
OPIHOGLOSSUM
 vulgatum
OSMUNDA
 cinnamomea
 interrupta
 regalis
 spectabile
ONOCLEA
 sensibilis
 obtusiloba
PTERIS
 aquilina
 atropurpurea
 caudata
POLYPODIUM
 alpestre
 calcareum
 dryopteris
 hexagonopterum
 phegopteris
 virginicum ?
 vulgare
SCOLOPENDRIUM
 officinarum
STRUTHIOPTERIS
 germanica
 pennsylvanica
WOODSIA
 Perminiana
 Ilvensis
 hyperborea
 rubra
WOODWARDIA
 blechnoides
 virginica

INDEX

OF ORDERS, GENERA, AND SPECIES.

	Page		Page		Page
<i>Aerostichum alpinum</i>	27	<i>filix-fœmina</i>	50	<i>regia</i>	30
<i>ilvense</i>	26	<i>fontanum</i>	48	<i>EQUISETALES</i>	2
<i>nemorale</i>	53	<i>germanicum</i>	44	<i>EQUISETACEÆ</i>	2, 16
<i>septentrionale</i>	44	<i>Halleri</i>	48	<i>EQUISETUM</i>	75
<i>spicant</i>	53	<i>irriguum</i>	50	<i>arcnarium</i>	80
<i>thelypteris</i>	35	<i>lanceolatum</i>	49	<i>arvense</i>	77
ADIANTUM	59	<i>lucidum</i>	49	<i>Drummondii</i>	76
<i>capillus</i>	59	<i>marinum</i>	46	<i>eburneum</i>	75
<i>capillus-veneris</i>	59	<i>melanocaulon</i>	47	<i>fluviatile</i>	75
<i>coriandrifolium</i>	59	<i>murale</i>	45	<i>helcocharis</i>	78
<i>fontanum</i>	59	<i>obtusatum</i>	46	<i>hyemale</i>	80
<i>Allosorus crispus</i>	58	<i>ruta-muraria</i>	45	<i>limosum</i>	78
ANNULATÆ	2	<i>saxatile</i>	47	<i>majus</i>	75
ASPIDIUM	31	<i>scolopendrium</i>	52	<i>maximum</i>	75
<i>aculeatum</i>	34	<i>septentrionale</i>	44	<i>nodosum</i>	79
<i>alpinum</i>	30	<i>spicant</i>	53	<i>nudum</i>	80
<i>angulare</i>	35	<i>trichomanes</i>	46	<i>palustre</i>	79
<i>asperum</i>	32	<i>trichomanoides</i>	47	<i>polymorphum</i>	78
<i>cristatum</i>	39	<i>viride</i>	47	<i>ramosum</i>	79
<i>dentatum</i>	28	<i>Athyrium alpinum</i>	30	<i>scgetale</i>	77
<i>dilatatum</i>	42	<i>dentatum</i>	28	<i>sylvaticum</i>	78
<i>dumetorum</i>	43	<i>filix-fœmina</i>	50	<i>telmateia</i>	75
<i>erosum</i>	39	<i>fontanum</i>	48	<i>tenue</i>	80
<i>filix-fœmina</i>	50	<i>regium</i>	30	<i>variegatum</i>	80
<i>filix-mas</i>	38	<i>thelypteris</i>	35	EXANNULATÆ	2
<i>fontanum</i>	48	<i>Bernhardia spinosa</i>	72	<i>Filix-fœmina</i>	55
<i>fragile</i>	29	BLECHNUM	53	<i>Filix lonchitidi affinis</i>	33
<i>fragrans</i>	36	<i>boreale</i>	53	GENERA	19
<i>goldianum</i>	39	<i>spicant</i>	53	GRAMMITIS	20
<i>irriguum</i>	50	BOTRYCHIUM	64	<i>ceterach</i>	20
<i>lincaris</i>	35	<i>lunaria</i>	65	<i>Gymnopteris ceterach</i>	20
<i>lobatum</i>	33, 34	<i>Capillus venciis verus</i>	59	HYMENOPHYLLACEÆ	2
<i>lonchitidoides</i>	33	<i>Ceterach alpinum</i>	27	HYMENOPHYLLUM	60
<i>lonchitis</i>	32	<i>officinarum</i>	20	<i>alatum</i>	62
<i>odoriferum</i>	36	CISTOPTERIS	28	<i>Tunbridgense</i>	60
<i>oreopteris</i>	36	<i>alpina</i>	30	<i>Tunbridgense</i> β	62
<i>recurvum</i>	38, 43	<i>dentata</i>	28	<i>Wilsoni</i>	61
<i>rheticum</i>	30	<i>fragilis</i>	29	ISOETACEÆ	2
<i>rigidum</i>	40	<i>regia</i>	30	ISOETES	68
<i>spinulosum</i>	41	CRYPTOGRAMMA	57	<i>lacustris</i>	68
<i>spinulosum</i>	40, 42	<i>crispa</i>	58	<i>setacea</i>	69
<i>thelypteris</i>	35	<i>Cyclopteris fragilis</i>	29	<i>eristata</i>	39
ASPLENIUM	43	<i>Cyathea alpina</i>	30	<i>Lastræa dilatata</i>	42
<i>adiantum nigrum</i>	49	<i>angustata</i>	30	<i>filix-mas</i>	38
<i>alternifolium</i>	44	<i>dentata</i>	28	<i>oreopteris</i>	36
<i>aquilina</i>	55	<i>fragilis</i>	29	<i>rigida</i>	40
<i>Brcynii</i>	44	<i>incisa</i>	30	<i>thelypteris</i>	35
<i>ceterach</i>	20	<i>Cystea dentata</i>	28	<i>Lepidotis alpina</i>	73
<i>elongatum</i>	52	<i>fragilis</i>	29	<i>annotina</i>	71

	Page		Page		Page
<i>clarata</i>	70	PILULARIA	67	<i>virginianum</i>	21
<i>Lomaria spicant</i>	53	<i>globulifera</i>	67	<i>vulgare</i>	21
<i>Lonehitis aspera</i>	53	<i>Plananthus inundatus</i> ..	72	<i>Polystichum aculeat</i> ...	34
<i>Lunaria minor</i>	65	<i>selago</i>	74	<i>callipteris</i>	38
LYCOPODALES ..	2	POLYPODIACEÆ 2, 3		<i>cristatum</i>	39
LYCOPODIACEÆ 2, 15		POLYPODIUM	21	<i>dryopteris</i>	24
LYCOPODIUM	69	<i>aculeatum</i>	33, 34	<i>filix-mas</i>	38
<i>abietiforme</i>	74	<i>alpinum</i>	30	<i>lonehitis</i>	32
<i>alpinum</i>	73	<i>arronieum</i>	26	<i>montanum</i>	36
<i>annotinum</i>	71	<i>calcareum</i>	24	<i>multiflorum</i>	42
<i>clavatum</i>	70	<i>callipteris</i>	39	<i>phegopteris</i>	23
<i>inundatum</i>	71	<i>cambrieum</i>	22	<i>rigidum</i>	40
<i>officinale</i>	70	<i>cristatum</i> ..	41, 42, 39	<i>strigosum</i>	40
<i>sabinaefolium</i>	73	<i>dentatum</i>	28	<i>thelypteris</i>	56
<i>selaginoides</i>	72	<i>dentatum (Moench)</i> ..	41	PTERIS	54
<i>selago</i>	74	<i>dilatatum</i>	42	<i>aquilina</i>	55
MARSILEACEÆ 2, 13		<i>dryopteris</i>	24	<i>caudata</i> β	55
<i>Muscus elavatus</i>	70	<i>dryopteris</i>	25	<i>erispa</i>	58
<i>Nephrod. dryopteris</i> ..	25	<i>filix-femina</i>	50	SCOLOPENDRIUM ..	51
<i>Onoclea spicant</i>	53	<i>filix-mas</i>	38	<i>alternifolium</i>	44
<i>crispa</i>	58	<i>fontanum</i>	48	<i>eeterach</i>	20
OPHIOGLOSSACEÆ 2		<i>fragile</i>	29	<i>offeinatum</i>	52
OPHIOGLOSSUM ..	66	<i>hyperboreum</i>	27	<i>phyllitis</i>	52
<i>vulgatum</i>	66	<i>Ilvense</i>	26	<i>ruta muraria</i>	45
<i>oratum</i>	86	<i>latebrosum</i>	23	<i>septentrionale</i>	44
ORDERS	2	<i>lobatum</i>	33	<i>vulgare</i>	52
OSMUNDACEÆ 2, 10		<i>lonehitis</i>	32	<i>Selaginella spinosa</i> ..	72
OSMUNDA	63	<i>oreopteris</i>	36	<i>Selago vulgaris</i>	74
<i>borcalis</i>	53	<i>ovato-erenatum</i> ..	50	<i>Stegania onocleoides</i> ..	58
<i>crispa</i>	58	<i>pulehellum</i>	24	<i>Trichomanes</i>	47
<i>lunaria</i>	65	<i>phegopteris</i>	23	TRICHOMANES ..	62
<i>lunata</i>	65	<i>regium</i>	30	<i>alatum</i>	62
<i>regalis</i>	63	<i>rhæticum</i>	30	<i>brevisetum</i>	62
<i>rupestris</i>	58	<i>rigidum</i>	40	<i>pulehellum</i>	60
<i>spicant</i>	53	<i>robertianum</i>	25	<i>pyxidiferum</i>	62
<i>Phyllitis heterophylla</i> ..	44	<i>spinosum</i>	41	<i>Tunbridgense</i>	60
<i>lanceifolia</i>	49	<i>spinulosum</i>	41	WOODSIA	26
<i>rotundifolia</i>	47	<i>thelypteris</i>	35, 36	<i>hyperborea</i>	27
<i>ruta-muraria</i>	45	<i>trifidum</i>	30	<i>Ilvensis</i>	26

APPENDIX

TO

THE PHYTOLOGIST

FOR 1851.

ART. I.—*Synoptical Table of the British Ferns.*

By EDWARD NEWMAN.

Obs. — This is confessedly but a crude attempt to arrange our ferns in accordance with their natural affinities. After maturely considering the associations upon which heretofore so much stress has been laid, I have arrived at the conclusion that they are unsatisfactory; and yet, in abandoning such divisions as Polypodium of the Linnean school; as Aspidium, so earnestly advocated by Smith and Hooker in this country, and equally great names on the Continent; and as the more recent combination called Lastrea by Presl and John Smith, and formerly adopted by myself; I feel as one groping his way in the dark, and shall eagerly avail myself of any friendly hand that may be held out to support and direct my tottering and uncertain footsteps.

VEGETABLES are divided by botanists into four primary groups, one of which, called Exogens, is single, and the others, severally called Endogens, Acrogens, and Thallogens, are double; that is, each of them contains two groups, which, in intimate structure, are alike, but in certain less important characters differ.

Acrogens are either Filicoid, or ferns and their allies; or Muscoid, or mosses and their allies.

Filicoid Acrogens are divided into seven secondary groups, called Orders; which however consist of a single one and three pairs: so that the divisions of the secondary, are in fact numerically identical with those of primary groups.

Primary groups are called *classes*, secondary groups *orders*. The seven orders of Filicoid Acrogens are these:—Polypodiaceæ; Osmundaceæ and Ophioglossaceæ; Marsiliaceæ and Lycopodiaceæ; Equisetaceæ and Characeæ.

The present paper treats only of the first, second and third of these orders, the British species of which are so few in number, that it is extremely difficult so to arrange them as to give any idea of a connected series.

Order. — POLYPODIACEÆ, *R. Brown.*

Plants composed of fibrous roots, solid simple rhizoma, and flat leafy fronds which rise with a circinate veneration and bear capsules in clusters on their back or edges. The capsules are provided with an elastic jointed ring. The divisions of this order are still obscure, and require further investigation; the subjoined arrangement is confessedly imperfect, but will probably be found convenient to those whose attention is chiefly confined to the European ferns.

Family. — ADIANTEÆ.

The ultimate divisions of the frond generally stipitate and leaf-like but without a mid-vein: clusters of capsules small, nearly circular, seated on the reflexed bleached margin: no apparent involucre.

Genus. — ADIANTUM, *Linneus.*

No mid-vein: veins of divisions of the frond variously branched, free at extremity: involucre not apparent: clusters of capsules nearly orbicular and situate on a bleached reflexed margin.

ADIANTUM CAPILLUS-VENERIS.

Adiantum Capillus-Veneris, *Linn. Sp. Pl.* 1559; *Lightf. Fl. Scot.* 679; *Huds. Fl. Ang.* 460; *Bolt. Fil. Brit.* 24, t. 29; *With. Arr.* 781; *Sm. E. F.* iv. 320, *E. B.* 1564; *Mack. Fl. Hib.* 344; *Newm. N. A.* 9, *F.* 83; *Hook. and Arn.* 576; *Bab.* 416.

Stipes black, shining, about the same length as the frond; frond deltoid, pinnate; pinnæ alternate, pinnate, pinnules stalked, leaf-like.

Sea-coast of Devonshire, Cornwall, Glamorganshire, Isle of Man, and South Isles of Arran.

Family. — PTERIDÆ.

Ultimate divisions of frond with a distinct mid-vein, lateral veins branched and united at their extremities by a distinct marginal vein, on which the capsules are placed in a continuous line, and are covered by the bleached reflexed epidermis.

Genus. — EUPTERIS, *Newman*.

Mid-vein distinct, lateral veins anastomosing at the margin, forming a marginal vein: involucre attached to inner side of marginal vein, linear, its margin split into capillary segments: capsules attached in a linear series to the marginal vein, exterior to the involucre: epidermis prolonged, bleached, reflexed, split into capillary segments and covering the capsules in the manner of an involucre.

EUPTERIS AQUILINA.

Pteris aquilina, *Linn. Sp. Pl.* 1533; *Lightf. Fl. Scot.* 657; *Huds. Fl. Ang.* 451; *Bolt. Fil. Brit.* 16, t. 10; *With. Arr.* 765; *Sm. E. F.* iv. 318, *E B.* 1679; *Mack. Fl. Hib.* 343; *Newm. N. A.* 11, *F.* 93; *Hook. and Arn.* 575; *Bab.* 415.

Eupteris aquilina, *Newm. Phytol.* ii. 278.

Rhizoma creeping; stipes long; frond deltoid, tripinnate, erect.

Abundant everywhere except on chalk.

Obs. — There are several other natural divisions of the Linnean genus *Pteris*, but neither of them contains British species. Messrs. Houlston and Moore, in their "Descriptive List of Cultivated Ferns," now in course of publication in the 'Gardeners' Magazine of Botany,' treat all the species of *Eupteris* as identical, thus making one species

cosmopolitan : in this conclusion, however, I am hardly prepared to agree.

Family. — BLECHNEÆ.

The ultimate divisions of the frond sessile, having a distinct mid-vein, the lateral veins anastomose in a linear series on each side of the mid-vein : capsules seated in a continuous line on that side of these anastomosing veins which is nearest the mid-vein, covered by a continuous linear lateral involucre, which opens towards the mid-vein.

Genus. — LOMARIA, *Willdenow*.

Mid-vein distinct, lateral veins anastomosing in a linear series on each side, parallel to the mid-vein, and emitting free branches to the margin : involucre linear, opening towards the mid-vein ; capsules in a linear series on the inner side of each anastomosing vein.

LOMARIA SPICANT.

Osmunda spicant, *Linn. Sp. Pl.* 1522 ; *Lightf. Fl. Scot.* 654 ; *Huds. Fl. Ang.* 450 ; *Bolt. Fil. Brit.* 8, t. 6.

Osmunda spicanthus, *With. Arr.* 763.

Blechnum spicant, *With. Arr.* 765.

Lomaria spicant (*Desv.*), *Newm. N. A.* 9, *F.* 89.

Blechnum boreale (*Swartz*), *Sm. E. F.* iv. 316, *E. B.* 1159 ; *Mack. Fl. Hib.* 343 ; *Hook. and Arn.* 575 ; *Bab.* 415.

Fronds of two kinds : fertile fronds erect, linear, pinnate ; pinnæ distant, reflexed, narrow, linear ; the lower portion of the stipes naked : barren fronds prostrate, lanceolate, pinnatifid ; pinnæ close, flat, broad, blunt.

Common on damp soils.

Family. — ASPLENIEÆ.

A large and varied group, approaching very nearly to the last, but constantly differing : the capsules are seated in linear clusters on one side of lateral veins, whose direction is always at an angle with the median line of the pinnule, so that in no instance can two or more of these lines of

capsules form a continuous line : each cluster is covered more or less completely by a linear lateral involucre.

Genus. — NOTOLEPEUM, *Newman*.

Mid-vein of pinnules present, lateral veins alternate, branched, branches anastomosing among themselves and with the branches of the next lateral vein : clusters of capsules on the first anterior branch of each lateral vein, and all of them directed towards the apex of the pinnule, except in the first lateral veins, both branches of which bear clusters, the anterior cluster directed as usual, but the posterior towards the midrib of the frond, these are therefore placed back to back : each cluster is accompanied by a narrow and nearly erect involucre, occupying the same position with regard to the capsules as that of a true *Asplenium* : the back of the frond is densely clothed with pointed overlapping scales.

Obs.—This genus has long been noticed as distinct, but I have seen no description in which the characters are correctly given. A new name seemed needful ; the names of *Scolopendrium*, *Grammitis*, and *Gymnogramma*, successively employed, have been severally restricted to very different genera, and that of *Ceterach* seems objectionable, as belonging of right to the species. A mode of overcoming the difficulty certainly exists in a repetition of the name, as *Ceterach Ceterach*, and this plan has been largely followed in the nomenclature of fishes, but I think it has not hitherto found its way into the science of Botany. Whenever it shall be introduced, there can be no doubt that the name of *Notolepeum* must give way.

NOTOLEPEUM CETERACH.

Asplenium Ceterach, *Linn. Sp. Pl.* 1538 ; *Lightf. Fl. Scot.* 661 ; *Huds. Fl. Ang.* 452 ; *Bolt. Fil. Brit.* 20, t. 12 ; *With. Arr.* 767.

Scolopendrium Ceterach, *Sm. E. F.* iv. 315, *E. B.* 1244.

Grammitis Ceterach (*Swartz*), *Mack.* 337.

Ceterach officinarum (*Willd.*), *Newm. N. A.* 28, *F.* 293 ; *Hook. and Arn.* 566 ; *Bab.* 415.

Notolepeum Ceterach, *Newm. F.* 9.

Stipes shorter than the frond; frond linear-lanceolate, pinnatifid, divisions waved or lobed; back of frond ferruginous with the dense covering of scales.

On limestone rocks, very local: on mortared walls, common in the West of England and South of Ireland.

Genus. — *PHYLLITIS*, Newman.

Lateral veins twice or thrice bifurcate, free at the extremity: capsules in linear series upon the anterior and posterior branches, on the anterior directed towards the apex of the frond, on the posterior towards its base, always in pairs, *i. e.*, when the anterior branch of a lateral vein bears a line of capsules, the posterior branch of the lateral vein next before it also bears a line of capsules corresponding in length, and the two lines or series form a confluent mass of capsules, covered by two involucres, which face each other, and even in an early stage of growth overlap and appear united.

Obs.—The name *Phyllitis* was employed by Ray and other eminent botanists of the pre-Linnean era, but I believe has not been used as generic since the introduction of the binominal nomenclature. I consider that the name of *Scolopendrium* should be confined, as intended by its author, to the species. As in the preceding instance, the repetition of the name, thus, *Scolopendrium Scolopendrium*, would be the strict application of the law of priority. The genus is generally acknowledged as distinct.

PHYLLITIS SCOLOPENDRIUM.

Asplenium Scolopendrium, Linn. *Sp. Pl.* 1537; *Lightf. Fl. Scot.* 660; *Huds. Fl. Ang.* 452; *Bolt. Fil. Brit.* 18, t. 11; *With. Arr.* 766.

Scolopendrium vulgare (*Sym. Syn.*), *Sm. E. F.* iv. 314, *E. B.* 1150; *Mack. Fl. Hib.* 342; *Newm. N. A.* 28, *F.* 289; *Hook. and Arn.* 574; *Bab.* 415.

Phyllitis Scolopendrium, *Newm. F.* 10.

Stipes shorter than frond; frond pendulous, linear, strap-shaped, entire.

Common on stone walls, hedge-banks, &c., in damp places.

Genus. — AMESIUM, *Newm.*

Ultimate divisions without a distinct mid-vein : veins of ultimate divisions very few, sparingly branched, free at the extremities : involucres narrow, linear, frequently facing each other as in the preceding genus, but rarely overlapping.

Obs.—Roth unites the species of this genus with those of *Scolopendrium*, all other authors with *Asplenium*, from which, however, they appear to me abundantly distinct.

* *Fronde simple.*

AMESIUM SEPTENTRIONALE.

Acrostichum septentrionale, *Linn. Sp. Pl.* 1524 ; *Lightf. Fl. Scot.* 656 ; *Huds. Fl. Ang.* 450 ; *Bolt. Fil. Brit.* 12, t. 8 ; *With. Arr.* 764.

Asplenium septentrionale, *Sm. E. F.* iv. 308, *E. B.* 1017 ; *Newm. N. A.* 27, *F.* 269 ; *Hook. and Arn.* 572 ; *Bab.* 41.

Amesium septentrionale, *Newm. F.* 10.

Stipes and frond of equal length, their separation indistinct ; frond linear, narrow, gradually diminishing into the stipes, sometimes forked, apices of divisions bifid ; clusters of capsules in two, three, or four long linear series.

A small and rare fern, in rocky and mountainous places. Somerset, Caernarvon, Cumberland, Northumberland, Edinburgh.

** *Fronde linear, pinnate.*

AMESIUM GERMANICUM.

1770. *Asplenium germanicum*, *Weiss, Pl. Cr.* 299 ; *Willd. Sp. Pl.* v. 330 ; *Hoffm. Deutschl. Fl.* ii. 13 ; *Ehrh. Crypt.* 43 ; *Presl. Tent. Pteridog.* 108 ; *Newm. F.* 265 ; *Bab.* 414.

1779. *Asplenium Breynii*, *Retz, Obs. Bot. fasc. i.* 32 ; *Sw. Syn. Fil.* 85.

1781. *Asplenium alternifolium*, *Wulfen. Jacq. Misc.* ii. 51 ;
With. Arr. 768 ; *Sm. E. F.* iv. 309, *E. B.* 2258 ; *Newm.*
N. A. 27 ; *Hook. and Arn.* 573.

Amesium germanicum, *Newm. F.* 10.

Stipes shorter than frond ; frond linear, pinnate ; pinnæ alternate, distant, linear, ascending, bifid or trifid at the apex ; clusters of capsules linear.

A small and extremely rare plant. Caernarvonshire, Perthshire.

*** *Frond deltoid.*

AMESIUM RUTA-MURARIA.

- Asplenium Ruta-muraria*, *Linn. Sp. Pl.* 1541 ; *Lightf. Fl. Scot.* 665 ; *Huds. Fl. Ang.* 453 ; *Bolt. Fil. Brit.* 28, t. 16 ; *With. Arr.* 769 ; *Sm. E. F.* iv. 309, *E. B.* 150 ; *Newm. N. A.* 27, *F.* 261 ; *Hook. and Arn.* 573 ; *Bab.* 414.

Amesium Ruta-muraria, *Newm. F.* 10.

Stipes longer than frond ; frond deltoid, composed of a few diamond-shaped, stalked, leaf-like divisions ; clusters of capsules linear, becoming confluent and entirely covering the divisions of the frond.

A small fern, common on rocks and mortared walls.

Genus. — ASPLENIUM.

Mid-vein distinct ; lateral veins simple or branched : involucre linear, attached to the side of the vein, its free margin sometimes jagged, but not split into capillary segments.

* *Frond linear, pinnate.*

ASPLENIUM TRICHOMANES.

1753. *Asplenium Trichomanes*, *Linn. Sp. Pl.* 1540 ; *Huds. Fl. Ang.* 452 ; *Bolt. Fil. Brit.* 22, t. 15 ; *With. Arr.* 768 ; *Sm. E. F.* iv. 305, *E. B.* 576 ; *Mack. Fl. Hib.* 341 ; *Newm. N. A.* 28, *F.* 285 ; 109 ; *Hook. and Arn.* 573 ; *Bab.* 414.

1776. *Asplenium Trichomanoides*, *With. Bot. Arr. Veg.* 653; *Lightf. Fl. Scot.* 662.

Stipes generally shorter than the frond, purple throughout; frond pinnate; pinnæ distant, stalked, ovate; clusters of capsules linear, nearly black.

Common on rocks, walls and hedge-banks.

ASPLENIUM VIRIDE.

1753. *Asplenium Trichomanes ramosum*, *Linn. Sp. Pl.* 1541; *With. Bot. Arr. Veg.* 654.

1762. *Asplenium viridi*, *Huds. Fl. Ang.* 385.

1777. *Asplenium viride*, *Lightf. Fl. Sc.* 663; *Huds. Fl. Ang.* 453; *Bolt. Fil. Brit.* 24, t. 14; *With. Arr.* 768; *Sm. E. F.* iv. 306, *E. B.* 2257; *Mack. Fl. Hib.* 341; *Newm. N. A.* 28, *F.* 281; *Hook. and Arn.* 573; *Bab.* 414.

Stipes somewhat shorter than the frond, purple at the base, otherwise green; frond very narrow, linear, pinnate; pinnæ stalked, distant, diamond-shaped, toothed; clusters of capsules linear, at last confluent, rust-coloured.

On rocks in mountain districts only.

Obs.—In this instance the earliest name, *Asplenium Trichomanes ramosum*, is abandoned, because evidently given as that of a variety, not of a species. The second in date is also abandoned, because of its ungrammatical termination, subsequently corrected by the author himself.

ASPLENIUM MARINUM.

Asplenium marinum, *Linn. Sp. Pl.* 1540; *Lightf. Fl. Scot.* 664; *Huds. Fl. Ang.* 453; *Bolt. Fil. Brit.* 26, t. 15; *With. Arr.* 769; *Sm. E. F.* iv. 307, *E. B.* 392; *Mack. Fl. Hib.* 341; *Newm. N. A.* 27, *F.* 275; *Hook. and Arn.* 573; *Bab.* 414.

Adiantum Trapeziforme, *Huds. Fl. Ang.* 460; *With. Bot. Arr. Veg.* 655; but certainly not of *Linn. Sp. Pl.* 1559, as cited by early English authors.

Adiantum Trapeziforme, *Berk. Syn.* 309.

Stipes shorter than frond; frond pinnate; pinnæ stalked, ovate, serrated; clusters of capsules linear, rust-coloured, always separate.

Common on rocks by the sea.

*** *Fronde deltoid.*

ASPLENIUM ADIANTUM-NIGRUM.

Asplenium Adiantum-nigrum, *Linn. Sp. Pl.* 1541; *Lightf. Fl. Scot.* 666; *Huds. Fl. Ang.* 454; *Bolt. Fil. Brit.* 30, t. 17, 3; *With. Arr.* 770; *Sm. E. F.* iv. 310, *E. B.* 1950; *Mack. Fl. Hib.* 342; *Newm. N. A.* 27, *F.* 255; *Hook. and Arn.* 573; *Bab.* 414.

Stipes longer than frond; frond elongate, deltoid, pinnate; lowest pair of pinnæ always longest, all the pinnæ pinnate; clusters of capsules linear, approximate to mid-rib.

*** *Fronde lanceolate.*

ASPLENIUM LANCEOLATUM.

Asplenium lanceolatum, *Huds. Fl. Ang.* 454; *With. Arr.* 770; *Sm. E. F.* iv. 311, *E. B.* 240; *Newm. N. A.* 27, *F.* 249; *Hook. and Arn.* 573; *Bab.* 414.

"? *Asplenium obtusatum*, *Guss.*" *Bab. MSS.*

Stipes shorter than frond; frond semi-erect, lanceolate, pinnate; lowest pair of pinnæ shorter than the second pair, all pinnate or pinnatifid; clusters of capsules at first linear, afterwards circular, distant from midrib.

Local, and mostly maritime; chiefly upon stone walls. Caernarvonshire, Cornwall, Devon, Gloucestershire, Kent, Merionethshire, Sussex.

Genus. — *ATHYRIUM*, *Roth.*

Ultimate divisions generally distinct and leaf-like, each with a distinct mid-vein: lateral veins always branched: involucre crescent-shaped, its free margin split into capillary segments. The rhizoma or cormus is long-enduring, suberect, and often of very large size; it sometimes increases laterally, but generally from the centre, and occasionally, in very old individuals, it becomes erect and trunk-like, as in the tree-ferns. There is probably some difference between the species in this respect, but the subject has not obtained the attention of botanists.

Obs. — This genus has a very great similarity, in general habit, to that which immediately follows; a similarity also extending more or less to several genera of the following family: it is comprised in the genus *Aspidium* of Smith: it is now almost universally regarded as comprising but a single species, the *Polypodium Filix-femina* of Linneus, and the following synonymes are those of the genus.

Polypodium Filix-femina, *Linn. Sp. Pl.* 1551; *Lightf. Fl.*

Scot. 673; *Huds. Fl. Ang.* 458; *With. Arr.* 778.

Polipodium Filix-femina, *Bolt. Fil. Brit.* 46, t. 25.

Aspidium Filix-fœmina (*Swartz*), and *Aspidium irriguum*,
Sm. E. F. iv. 295, 6, *E. B.* 1459 and *E. B. S.* 2199.

Asplenium Filix-fœmina (*Bernh.*), *Mack. Fl. Hib.* 342;
Hook. and Arn. 574.

Athyrium Filix-fœmina (*Roth*), *Newm. F.* 420; *Bab.* 413.

* *Frond deltoid.* (No ascertained British species).

(*ATHYRIUM DELTOIDEUM.*

Aspidium crenatum, *Sommerf. in Vet. Ac. Handlung.* 104.

Cystopteris crenata, *Fries, Nov. Fl. Suec.* 165; *Hook. Sp. Fil.* i. 200.

Stipes as long as the frond, pretty thickly clothed towards the base with dark brown lanceolate scales; frond deltoid, resembling that of *Eupteris aquilina*, but of smaller size, pinnate; pinnæ stipitate, lanceolate, pinnate; in the first pair of pinnæ the pinnules on the lower are longer than those on the upper side; pinnules pinnate, their lobes distant, but united at the base by the winged midrib of the pinnule, blunt, slightly lobed on the upper margin; clusters of capsules confined to the upper portion of the frond, rather scattered in an oblique series on each side of the midrib, and rather near it; involucre nearly linear, attached to the side of the capsuliferous vein, and opening towards the midrib, its free margin ragged.

North of Europe, in thick woods. Found by Mr. R. B. Bowman, in the Pass of Kringelen, in Norway.

Obs.—There exist a *Polypodium crenatum*, an *Aspidium crenatum*, and an *Athyrium Filix-femina*, var. *crenatum*; under these circumstances, a change of name will, I hope, be allowed. The plant is here

introduced as being likely to occur in Scotland, and also to show that the genus *Athyrium* is not restricted to the lanceolate form of frond).

**** *Frond lanceolate.***

ATHYRIUM OVATUM.

1795. *Polypodium dentatum*, Hoffm. *Deutsch. Flor.* ii. 6.

1800. *Athyrium ovatum*, Roth, *Fl. Germ.* iii. 64; Newm. *F.* 420, *Phytol.* iv. 368.

Athyrium Filix-femina, var. *dentatum*, Newm. *F.* 203, ad partem.

Asplenium Filix-fœmina, var. *latifolium*, Hook. and Arn. 574; Bab. 413; Houlston and Moore, *Gard. Mag. of Bot.* iii. 262.

Frond lax, flaccid, dark green, lanceolate, pinnate; pinnae scarcely ascending, approximate, flattened, pinnate; pinnules ovate, distinctly stalked, crowded, overlapping, lobed at the base, toothed at the apex; clusters of capsules elongate as in *Asplenium*, in a series on each side of the pinnule but *distant from it*. A large plant, two to three feet high.

Rare; found in the neighbourhood of Keswick by Miss Beever and Miss Wright.

Obs. — The prior name of *dentatum* is discarded on account of its identity with another *Polypodium dentatum*.

ATHYRIUM MOLLE.

Polypodium molle, Schreber, *Spic. Flor. Lips.* 70; Ehrh. *Crypt.* 9; Hoffm. *Deutsch. Flor.* ii. 6.

Athyrium molle, Roth, *Flor. Germ.* iii. 61; Newm. *N. A.* 26.

Athyrium Filix-femina, var. *molle*, Newm. *F.* 242.

Athyrium Filix-femina, γ. Bab. 413.

Frond lax, flaccid, bright green, broadly lanceolate, pinnate; pinnae scarcely ascending, approximate, flattened, pinnate, subpinnate, or pinnatifid; pinnules blunt, approximate, connected by the wing of the midrib; clusters of capsules in a series on each side of the midrib of the pinnule, and *very near it*. Generally a small plant, twelve to

eighteen inches high, when larger somewhat more divided, and then it is the *Polypodium trifidum* of Hoffman.

Common in woods.

ATHYRIUM INCISUM.

Polypodium incisum, Hoffm. *Deutsch. Flor.* ii. 6.

Athyrium Filix-femina, Roth, *Flor. Germ.* iii. 65.

Athyrium Filix-femina, var. *incisum*, Newm. *F.* 243.

Athyrium Filix-femina β. Bab. 413.

Frond suberect, subrigid, dull green, lanceolate, pinnate; pinnæ subdistant, pinnate; pinnules deeply incised or lobed, divisions dentate; clusters of capsules close, owing to the greater subdivision of frond not ranged in series, but crowded and finally confluent. A large plant, two to four feet high, and proportionably broad.

Common in wet woods. A much more beautiful fern than either of the preceding, and the type of the genus.

ATHYRIUM CONVEXUM.

Athyrium rhœticum, Roth. *Flor. Germ.* iii. 67; Newm. *N.*

A. 26.

Athyrium Filix-femina, var. *convexum*, Newm. *F.* 245.

Athyrium Filix-femina, α. Bab. 413.

Frond pale green, erect, rigid, linear lanceolate, pinnate; the stipes and rachis semipellucid, and often beautifully coloured with purple or red; pinnæ distant, at first ascending, then spreading, and finally deflexed, extremely acute; pinnules distant, very narrow, linear, entirely unconnected, their margin convolute; clusters of capsules subrotund, close to the midrib of the pinnule, and finally covering their under surface, and themselves partially covered by the convolute margin of the pinnules.

Not uncommon in exposed localities.

Obs.—The seedlings of these plants are particularly abundant near the parent; they constitute the *A. rhœticum*, var. *minus* of Roth, and the *Aspidium irriguum* of Smith.

Family. — POLYPODIÆ.

A large and varied group. Capsules seated in circular clusters directly on the back of the lateral veins.

Genus. — PSEUDATHYRIUM, *Newm.*

Involucre wanting: clusters of capsules small, at first distinct, but often crowded when mature: first and second pinnules on each pinna, both above and below, of nearly corresponding size: ultimate divisions pointed but without spines: precisely the habit of *Athyrium*.

PSEUDATHYRIUM ALPESTRE.

Aspidium alpestre, *Hoppe, Taschenb.* (1805), 216; *Schkuhr*, 58.

Polypodium alpestre, *Koch.*

Aspidium rhæticum, *Swartz, Syn. Fil.* 59.

Polypodium rhæticum, *Woods, Tour. Fl.* 423.

Pseudathyrium alpestre, *Newm. Phytol.* iv. 370.

Rhizoma tufted; stipes shorter than frond; frond elongate, lanceolate, pinnate; pinnae pinnate; pinnules approximate, lobed, lobes notched.

Scotland. Three specimens are in the herbarium of Mr. Watson, gathered by himself, one in Canlochen Glen, Forfarshire, a second on Ben Aulder, and a third on "mountains near Dalwhinnie." Mr. Watson thinks it possible that, as he went from Dalwhinnie to Ben Aulder, both the specimens last mentioned may be from the latter locality. Common in alpine districts of Europe, and will in all probability prove so in Scotland. Lightfoot, in describing the clusters of capsules in the Scotch specimens of *Filix-femina*, says they first appear as "*distinct round dots*;" he had probably seen the present species.

Obs. — In *Schkuhr's 'Handbook'* there is a reference to plate 60, which plate is named *Aspidium umbrosum*, but certainly represents this species.

Genus. — POLYSTICHUM, Schott.

Involucre circular, scale-like, its margins free, attached by its centre : first upper pinnule of each pair greatly larger than the second, and larger than the first and second lower pinnules ; all the ultimate divisions ending in an acute spine.

Obs.—The genus *Polystichum*, as proposed by Roth, is almost identical with *Aspidium* of Willdenow, but was subsequently restricted by Schott to the typical species, and those having a similar circular involucre. Although as regards British species the genus thus restricted is remarkably distinct, yet among exotic ferns we find species almost precisely intermediate between this and the following genus.

POLYSTICHUM ANGULARE.

Polypodium aculeatum, *Lightf. Fl. Scot.* 675 ; *Huds. Fl. Ang.* 459.

Aspidium angulare, *Willd. Sp. Pl.* v. 257 ; *Sm. E. F.* iv. 291, *E. B. S.* 2776 ; *Mack. Fl. Hib.* 339 ; *Hook. and Arn.* 568.

Polystichum angulare, *Newm. N. A.* 25, *F.* 173 ; *Bab.* 412.

Rhizoma tufted ; stipes one-third as long as the frond, densely clothed with large, red, chaffy scales ; frond drooping, graceful, broad lanceolate, lax, feathery, pinnate ; pinnæ very numerous, linear, distant, pinnate ; pinnules distinct, stalked, often distant, auricled at the base, rounded at the apex, serrated, spined.

Common in woods and shady lanes.

POLYSTICHUM ACULEATUM.

Polypodium aculeatum, *Linn. Sp. Pl.* 1552.

Polypodium lobatum, “affinis præcedenti (*P. angulare*, *l. c.*) an distincta sit species ?” *Huds. Fl. Ang.* 459 ; *With. Bot. Arr. Veg.* 651.

Polipodium aculeatum, *Bolt. Fil. Brit.* 48, t. 26. “*Polipodium lobatum*, *Hall. Hist.* 1712, and *Fl. Ang.* 459, is doubtless a young plant of *Polipodium aculeatum* ; of this I am certain from observation.”—*Bolt. l. c.*

Aspidium lobatum (Swartz), *Sm. E. F.* iv. 291; *Mack. Fl. Hib.* 338; *Hook. and Arn.* 568.

Polystichum aculeatum (Roth), *Newm. N. A.* 25, *F.* 169; *Bab.* 411.

Rhizoma tufted; stipes very short; frond rigid, horizontal, leathery, linear lanceolate, pinnate; pinnæ numerous, pinnatifid, divisions decurrent.

Common everywhere.

POLYSTICHUM LONCHITIS.

Polypodium Lonchitis, *Linn. Sp. Pl.* 1548; *Lightf. Fl. Scot.* 668; *Huds. Fl. Ang.* 455; *With. Arr.* 773; *Sm. E. B.* 797.

Polipodium Lonchitis, *Bolt. Fil. Brit.* 34, t. 19.

Aspidium Lonchitis (Swartz), *Sm. E. F.* iv. 284; *Mack. Fl. Hib.* 338; *Hook. and Arn.* 568.

Polystichum Lonchitis (Roth), *Newm. N. A.* 25, *F.* 163; *Bab.* 411.

Rhizoma tufted; stipes very short; frond linear, rigid, leathery, pinnate; pinnæ entire, auricled at the base, serrated, very spiny; clusters of capsules circular, crowded, often confluent, confined to the upper half of the frond.

On mountains, rare. Caernarvonshire, Durham, Yorkshire, Forfarshire, Perthshire, Kerry, Sligo.

Genus. — LOPHODIUM, *Newm.*

Involucre nearly circular, scale-like, its direction oblique to the plane of the frond, its margin with a conspicuous notch, and its attachment at this notch: first upper pinnule generally larger than the second, and always greatly less than the first lower pinnule: all the ultimate divisions ending in a point: cormus or rhizoma large, massive, and long enduring.

* *Frond deltoid; points of divisions spine-like.*

LOPHODIUM (RECURVUM) FENESECH.

1790. *Polipodium cristatum*, *Bolt. Fil. Brit.* 42, ad partem, *i. e.*, t. 23.

1832. *Nephrodium fœnesecii*, *Lowe, Camb. Phil. Trans.* iv. 7, ad partem, fortè omninò.

Aspidium dilatatum, var. *recurvum*, *Bree, Mag. Nat. Hist.* iv. 162.

1843. *Aspidium recurvum*, *Bree, Phytol.* i. 773.

Lastrea recurva, *Newm. N. A.* 23, *F.* 225.

Lastrea fœnesecii, *Watson, Phytol.* ii. 568; *Bab.* 411.

Aspidium spinulosum, γ. *Hook. and Arn.* 57.

Lophodium recurvum, *Newm. Phytol.* iv. 371.

Rhizoma tufted, large, crown unusually broad; stipes as long as frond, woody, clothed with long, narrow, lacinated scales; frond elongate, triangular (being exactly that of *Asplenium Adiantum-nigrum*), drooping, elegant, pale delicate green when young, its under surface sprinkled over with sessile, pellucid glands (which probably cause the powerful scent for which this species is remarkable), pinnate; lowest pair of pinnæ longest stalked, all pinnate; all the divisions of the frond concave; involucre jagged, without stalked glands; clusters of capsules round, crowded, covering every part of the frond.

Common in Ireland and Cornwall; occurring in Devonshire, N. Wales, Cumberland, Sussex, &c. sparingly.

*** Frond linear-lanceolate; points of divisions spine-like.*

LOPHODIUM MULTIFLORUM.

Polypodium cristatum, *Linn. Sp. Pl.* 1551, ad partem; *Huds. Fl. Ang.* 390 (1762), (I place this synonyme here without hesitation, not simply from the accordance of Hudson's specific character, but because that author specially cites Hampstead Heath as the locality, and the present species has existed there from Hudson's time to the present, and no other form or supposed species has ever been found there); *Id.* 457, (1778); *Lightf. Fl. Scot.* 670; *Bolt. Fil. Brit.* 42, ad partem, (the second variety refers to this species); *With. Arr.* 778.

Polystichum multiflorum, *Roth, Fl. Germ.* iii. 87.

Aspidium dilatatum, *spinulosum*, and *dumetorum*, *Sm. E.*

F. iv. 292, 3, 4, and also of *Smith's Herbarium*, now in the possession of the Linnean Society.

Aspidium spinulosum, *Mack. Fl. Hib.* 340 ; var. β . *Hook. and Arn.* 571.

Lastrea dilatata, *Newm. N. A.* 23 ; *Bab.* 411.

Lastrea multiflora, *Newm. F.* 215.

Lophodium multiflorum, *Newm. Phytol.* iv. 371.

Rhizoma tufted ; stipes very stout, nearly as long as the frond, densely clothed with long pointed scales, which are dark brown along the middle but pale at the edges ; frond glandular when young, very large, deep green, drooping, ovate-lanceolate, pinnate ; lowest pair of pinnæ shorter than the second, third, fourth or fifth, pinnæ pinnate ; pinnales pinnate or pinnatifid ; ultimate divisions serrated, spined ; all the divisions of the frond convex ; involucre nearly circular, fringed with stalked glands ; clusters of capsules circular, covering every part of the frond.

Common everywhere.

Obs. — Two or more doubtful species of *Lophodium* occupy this place ; viz., *L. glandulosum*, probably identical, as suggested to me by Mr. Moore, with *Lastrea maculata* of Deakin, *L. collinum*, &c. The characters of these are still insufficiently ascertained.

LOPHODIUM SPINOSUM.

Polypodium cristatum, *Linn. Sp. Pl.* 1551, ad partem.

1770. *Polypodium Filix-femina* γ . *spinosa*, *Weiss, Crypt.* 316.

1800. *Polystichum spinosum*, *Roth, Fl. Germ.* iii. 91.

Lastrea spinosa, *Newm. N. A.* 21, *F.* 209.

Lastrea spinulosa, *Bab.* 410.

Lophodium spinosum, *Newm. Phytol.* iv. 371.

Rhizoma stout, slowly but extensively creeping ; stipes as long as frond, clothed sparingly, except at the base, with broad, rounded, pale brown, diaphanous scales ; frond slightly drooping, elongate, linear, pinnate ; pinnæ rather distant, winged, pinnate ; pinnales at the base of the pinnæ separated from the midrib by a deep notch, towards the apex of the pinnæ decurrent, all lobed, the lobes serrated

and spined; divisions at the apex of the frond narrow, their terminations acute; all the divisions of the frond flat; involucre nearly circular, its margins waved, not torn, nor furnished with teeth or stalked glands; clusters of capsules circular, crowded, sometimes confluent, confined to the upper part of the frond.

Common in damp woods in England. I have not seen it from Scotland or Ireland.

Obs.—This very common fern has totally escaped the notice of Smith, Mackay, and Hooker and Arnott. Mr. Moore, in copying my figures of the paleæ and involucres of multiflora and spinosa, has unhappily transposed them. Babington is the only British author to whom I can refer.

LOPHODIUM ULIGINOSUM.

Aspidium spinulosum, var. *uliginosum* (*A. Braun*), *Döll, Rhein. Flor.* 17, 18. Vide *Phytol.* iii. 101.

Lastrea uliginosa, *Newm. Phytol.* iii. 679.

Lastrea cristata, var. *uliginosa*, *Moore, Phytol.* iv. 149, in *Rep. Bot. Soc. Ed.*

Lastrea cristata, *Bab.* 410, ad partem.

Aspidium spinulosum, *Hook. and Arn.* 571, ad partem.—

“The plant under the name of *L. uliginosa* in the Royal Gardens, corresponds with our *A. spinulosum*, α .”—*Hook. and Arn. l. c.* The plant here referred to as cultivated at Kew, is correctly named as my *L. uliginosa*; I mention this to show, from the evidence of all parties concerned, that *Aspidium spinulosum*, α ., *Hook. and Arn.*, and my *Lophodium uliginosum* are identical.

Lophodium uliginosum, *Newm. Phytol.* iv. 371.

Rhizoma tufted; veneration simply circinate, fronds erect, rigid, linear lanceolate, of two kinds, the fertile resembling those of the preceding, the barren those of the following species, pinnate; pinnæ also pinnate.

Not uncommon; bogs and boggy heaths, in company with the following species.

LOPHODIUM CALLIPTERIS.

Polypodium cristatum, *Linn. Sp. Pl.* 1551, ad partem.

1788. *Polypodium Callipteris*, *Ehrhart, Beitrage*, iii. 77, *Crypt.* 53; *Hoffm. Deutsch. Fl.* ii. 6.

1800. *Polystichum cristatum*, *Roth, Fl. Germ.* iii. 84.

Aspidium cristatum, *Sm. E. F.* iv. 289, *E. B.* 2125; *Hook. and Arn.* 569.

Lastrea Callipteris, *Newm. F.* 12.

Lastrea cristata, *Newm. N. A.* 21, *F.* 203; *Moore, Phytol.* iv. 149, ad partem; *Bab.* 410, ad partem.

Lophodium Callipteris, *Newm. Phytol.* iv. 371.

Rhizoma very stout, slowly creeping, often extending two or three feet; stipes branched, as long as the frond, sparingly clothed with short, broad, pale, semi-hyaline scales; frond very erect, narrow, linear, pinnate; pinnae rather distant, short, somewhat triangular, pinnatifid, from five to eight pairs usually of the same length, but the fifth, sixth, seventh or eighth pair somewhat longest, and the others gradually approaching them in length, the distance between each pair gradually decreasing from the base towards the apex; pinnules generally decurrent, oblong, lobed, serrated, blunt or rounded at the apex, the lower pinnules generally larger and longer than the upper; involucre flat, its margin irregular; clusters of capsules crowded, often confluent, confined to the upper part of the frond.

On boggy ground, in Cheshire, Norfolk, Nottingham, and Suffolk; very local.

*** *Involucre more convex and more completely reniform; points of divisions obtuse.*—*Dryopteris*, *Schott.*

LOPHODIUM FILIX-MAS.

Polypodium Filix-mas, *Linn. Sp. Pl.* 1551; *Lightf. Fl. Sc.* 671; *Huds. Fl. Ang.* 458; *With. Arr.* 775.

Polipodium Filix-mas, *Bolt. Fil. Brit.* 44, t. 24.

Aspidium Filix-mas (*Swartz*), *Sm. E. F.* iv. 288, *E. B.* 1458; *Mack.* 340; *Hook. and Arn.* 569.

Aspidium cristatum, *Sm. E. B.* 1949; *Mack.* 340.

Dryopteris Filix-mas, *Schott, Fil.*

Lastrea Filix-mas, *Newm. N. A.* 19, *F.* 197; *Bab.* 410.

Rhizoma tufted ; stipes short, densely clothed with reddish scales ; fronds semi-erect, lanceolate, pinnate ; pinnæ numerous, pinnate ; pinnules blunt, serrated ; involucre very perfect, without stalked glands ; clusters of capsules less crowded nearer the midrib, absent from the lower pinnæ.

Common everywhere.

Obs.—Two apparent species, when better understood, may be introduced here, *Lophodium erosum*, the *Aspidium erosum* of Schkuhr, and *L. abbreviatum*, the *Polystichum abbreviatum* of Decandolle.

LOPHODIUM (RIGIDUM) FRAGRANS.

1753. *Polypodium fragrans*, *Linn. Sp. Pl.* 1089, (1st edition) ; *Huds. Fl. Ang.* 388, (1st edition) ; *With. Arr.* 650 ; *Villars, Hist. Pl. Dauph.* iii. 843.

1795. *Polypodium rigidum*, *Hoffm. Deutsch. Flor.* ii. 16.

1810. *Polystichum strigosum*, *Roth, Fl. Germ.* iii. 86.

Aspidium fragrans, *Gray, Nat. Arr.* ii. 9.

Aspidium rigidum (*Swartz*), *Hook. E. B. S.* 2724 ; *Hook. and Arn.* 569.

Lastrea rigida (*Presl*), *Newm. N. A.* 19, *F.* 191 ; *Bab.* 411.

Rhizoma tufted ; stipes much shorter than the frond, densely clothed with reddish scales ; frond semi-erect, glandulose, sweet-scented, lanceolate, pinnate ; pinnæ very numerous ; pinnules oblong, obtuse, serrated ; involucre very perfect, fringed with stalked glands ; clusters of capsules very crowded, covering the pinnules, absent from the lower pinnæ.

Not uncommon upon limestone rocks in the North of England.

Genus. — HEMESTHEUM.

Clusters of capsules on both branches of the lateral veins, and equidistant from the mid-vein, so as to form a continuous submarginal series, which, in the first division, is completely covered by the revolute margin of the pinnule ; in the second division, nearly so : involucre instable, sometimes totally absent, at other times small, indistinct, subreniform, evanescent : first upper pinnule longer

than the first lower, both slightly longer than the second and following pinnules: ultimate divisions without a point.

Obs. — This genus has been carefully defined by Schott under the name of *Thelypteris*; but I object, for the reason before stated, to the transfer of a name from a species to a genus.

HEMESTHEUM THELYPTERIS.

Acrostichum Thelypteris, *Linn. Sp. Pl.* 1528; *Bolt. Fil.*

Brit. part ii. 78, t. 43, 44; *With. Bot. Arr. Veg.* 649.

Polypodium Thelypteris, *Linn. Mant.* 505; *Huds. Fl. Ang.* 457; *With. Arr.* 776; ? *Lightf. Fl. Scot.* 674.

Polystichum Thelypteris, *Roth. Fl. Germ.* iii. 77.

Aspidium Thelypteris (*Swartz*), *Sm. E. F.* iv. 285; *Mack. Fl. Hib.* 340; *Hook. and Arn.* 569, (excl. syn. "E. B. 1018," which represents *G. Phegopteris*).

Lastrea Thelypteris (*Bory*), *Newm. N. A.* 19, *F.* 183; *Bab.* 409.

Thelypteris palustris, *Schott, Fil.*

Rhizoma creeping; fronds of two kinds, both erect, on long smooth stipes, lanceolate, pinnate; pinnæ pinnatifid, lower pinnæ equalling the rest in length; pinnules blunt, entire, in fertile fronds with convolute margins covering the capsules; lateral veins branched; involucre small, and present on scarcely more than half the clusters; capsules in circular clusters, scarcely marginal.

In marshes, very local.

HEMESTHEUM (OREOPTERIS) MONTANUM.

Polypodium fragrans, *Linn. Mant.* ii. 307; *Huds. Fl. Ang.* 457, (2nd edition). The reader will see that the altered description no longer agrees with *L. rigida*.

1781. *Polypodium montanum*, *Vogler, Diss. de Pol. mont.*

1788. *Polypodium Oreopteris*, *Ehrh. Beitr.*; *Dicks. Tr. Linn. Soc.* i. 181, (1791); *With. Arr.* 775.

Polipodium Thelypteris, *Bolt. Fil. Brit.* 40, t. 22.

Polystichum montanum, *Roth. Flor. Germ.* iii. 74.

Aspidium Oreopteris (*Swartz*), *Sm. E. F.* iv. 286, *E. B.* 1019; *Mack. Fl. Hib.* 339; *Hook. and Arn.* 569.

Lastrea Oreopteris, *Newm. N. A.* 17, *F.* 187; *Bab.* 410.

Rhizoma tufted ; stipes very short, chaffy ; frond semi-erect, lanceolate, much attenuated at the base, pinnate ; pinnæ pinnatifid, divisions rounded, lower pinnæ very short, deltoid, obtuse ; veins and capsules as in the preceding ; involucre sometimes quite obvious.

Mountains and moist woods, frequent.

Obs.—Each species is the type of a group of species, and might be regarded as a genus, but the intervention of *H. Novaboracense* tends very much to unite them.

Genus. — GYMNOCARPIUM, *Newman*.

Ultimate divisions of the frond with branched lateral veins free at the extremity, bearing clusters of capsules on all the branches : involucre none. In the British species the plant extends by means of a rapidly extending, slender, subterranean, stolon-like rhizoma.

GYMNOCARPIUM PHEGopteris.

Polypodium Phegopteris, *Linn. Sp. Pl.* 1550 ; *Lightf. Fl. Scot.* 669 ; *Huds. Fl. Ang.* 456 ; *With. Arr.* 775 ; *Sm. E. F.* iv. 282, *E. B.* 2224 ; *Mack. Fl. Hib.* 337 ; *Newm. F.* 115 ; *Hook. and Arn.* 566.

Polipodium Phegopteris, *Bolt. Fil. Brit.* 36, t. 20.

Aspidium Thelypteris, *Sm. E. B.* 1018.

Lastrea Phegopteris, *Newm. N. A.* 17, *F.* 13.

Polypodium ? Phegopteris, *Bab.* 408.

Gymnocarpium Phegopteris, *Newm. Phytol.* iv. 371.

Polystichum Phegopteris, *Roth, Flor. Germ.* iii. 72.

Rhizoma creeping ; stipes long ; frond ovate-deltoid, pinnate, drooping ; first pair of pinnæ distinct, turned back, the rest united at the base, pointing forwards, all pinnatifid ; veins, capsules and involucres as in the preceding ; colour dull green ; stem concolorous, rather scaly.

By mountain rills and waterfalls, and in wet woods : common in Scotland, North of England, and Wales ; rare in Ireland.

GYMNOCARPIUM ROBERTIANUM.

1785. *Polypodium Dryopteris*, *Bolt. Fil. Brit.* 53, t. 1.

1795. *Polypodium Robertianum*, *Hoffm. Deutsch. Fl.* ii. 10. — “Fronde triangulari, foliolis ternis bipinnatis: pinnis pinnulisque inferne pinnatifidis. Stipes glaucus, uno latere sulcatus. Frons tenera. Uterque nudo oculo subtili tomento ad lentem brevissimis glandulis obsitus. Odor debilis Geran. Robert. Fructif. minuta.”—*Hoffm. l. c.*

1804. *Polypodium calcareum*, *Sm. Fl. Brit.* 1117, *E. F.* iv. 283, *E. B.* 1525; *Newm. F.* 131; *Hook. and Arn.* 567.

Lastrea calcarea, *Newm. N. A.* 17.

Lastrea Robertiana, *Newm. F.* 13.

Polypodium ? *calcareum*, *Bab.* 409.

Gymnocarpium Robertianum, *Newm. Phytol.* iv. 371.

Rhizoma creeping; stipes erect; frond elongate-deltoid, glandular-mealy; lower pinnae bipinnate, upper pinnae pinnate only; colour dull green, stipes concolorous.

Among loose stones in limestone districts, not common.

GYMNOCARPIUM DRYOPTERIS.

Polypodium Dryopteris. *Linn. Sp. Pl.* 1555; *Lightf. Fl. Scot.* 678; *Huds. Fl. Ang.* 460; *With. Arr.* 780; *Sm. E. F.* iv. 283, *E. B.* 616; *Mack. Fl. Hib.* 338; *Newm. F.* 123; *Hook. and Arn.* 567.

Polypodium Dryopteris, *Bolt. Fil. Brit.* 52, t. 28.

Polystichum Dryopteris, *Roth. Fl. Germ.* iii. 80.

Lastrea Dryopteris, *Newm. N. A.* 15, *F.* 13.

Polypodium ? *Dryopteris*, *Bab.* 409.

Gymnocarpium Dryopteris, *Newm. Phytol.* iv. 371.

Rhizoma creeping; stipes erect, glabrous; frond triple, deltoid, smooth, the three branches pinnate; pinnae pinnatifid; lateral veins usually simple; involucre generally wanting; clusters of capsules near the extremity of each lateral vein, forming a marginal series; colour a bright green, stipes purplish.

An exquisitely beautiful little fern. Common in mountain districts, otherwise rare.

Genus. — CYSTOPTERIS, *Bernhardi*.

Mid-vein of ultimate divisions distinct but sinuous, lateral veins branched, free: involucre attached almost beneath the mass of capsules, half-way between the mid-vein and the extremity, directed at first backwards, then upwards, then forwards, and almost covering the circular mass of young capsules like a hood, its anterior margin split into unequal and often capillary segments.

CYSTOPTERIS (MONTANA) ALLIONI.

1789. *Polypodium montanum*, *Allion. Pedem.* n. 2410.

Cyathea montana, *Roth, Flor. Germ.* iii. 100.

Cystopteris montana (*Link*), *Newm. Phytol.* i. 671, *N. A.* 15, *F.* 13 & 159; *Hook. and Arn.* 572; *Bab.* 413.

Rhizoma creeping; stipes erect, longer than the frond; frond deltoid.

Apparently a rare, certainly a local fern; hitherto observed only in Scotland. Found in 1836 on Ben Lawers, by Mr. W. Wilson: in a ravine called Corrach Dh'Ouflach, in 1841, by Messrs. W. Gourlie and W. Adamson: in the same place by Mr. Borrer and Dr. Walker-Arnott in 1850: and again by the Rev. W. Little in 1851.

Obs.—I think the name of *Polypodium montanum* was given to this plant against all the rules of botanical nomenclature, Vogler having given that name to another species eight years previously.

CYSTOPTERIS FRAGILIS.

Polypodium fragile, *Linn. Sp. Pl.* 1553; *Lightf. Fl. Scot.* 677; *Huds. Fl. Ang.* 459; *Bolt. Fil. Brit.* 50, t. 27 & 46; *With. Arr.* 779.

1762. *Polypodium rhæticum*, *Huds. Fl. Ang.* 458; *With. Arr.* 780; *Bolt. Fil. Brit.* part ii. 80, t. 45, but certainly not of *Linn. Sp. Pl.* 1552, as cited by early English authors.

1779. *Polypodium polymorphum*, *Villars, Dauph.* iii. 847.

1793. *Polypodium dentatum*, *Dicks. Crypt. fasc.* iii. 1, t. 7, f. 1, *Id. H. Sicc. fasc.* 16; *With. Arr.* 776.

1796. *Polypodium trifidum*, *With. Arr.* 779.

Cyathea fragilis, *Roth, Flor. Germ.* iii. 94.

Cystea fragilis, *Sm. E. F.* iv. 298, *E. B.* 1587.

Cystea dentata, *Sm. E. F.* iv. 300, *E. B.* 1588.

Cystea angustata, *Sm. E. F.* iv. 301.

Cystea regia, *Sm. E. F.* iv. 302, ad partem, *i. e.*, excl. the plant found on the garden-wall at Low Layton, which has not been gathered wild in Britain.

Cistopteris fragilis, *Mack. Fl. Hib.* 341.

Cystopteris fragilis, *Newm. N. A.* 15, *F.* 13, 149; *Hook. and Arn.* 572; *Bab.* 412.

? *Cystopteris dentata*, *Bab.* 412, ad partem, *i. e.*, excl. *Dickieanum*, *Sim.*

Rhizoma quasi-tufted, but increasing laterally; stipes shorter than the frond; frond erect, lanceolate, bipinnate; pinnæ ascending; pinnules distinct.

A small, elegant, and fragile fern, common in Wales, the North of England, Scotland, and parts of Ireland.

Obs.—*Cystopteris alpina* of Desveux, Hooker and Arnott, and Babington, being also the *C. regia* of Smith in part, and the *C. incisa* of 'English Botany,' is not a true native.

CYSTOPTERIS DICKIEANA.

Cystopteris Dickieana, *Sim, in Gard. Journ.* 308, 1848.

Cystopteris fragilis, α . *Dickieana*, *Moore, Bot. Gaz.* i. 310.

Cystopteris dentata, *Bab.* 412, ad partem, *i. e.*, excl. *dentata*, *Sm.*

Rhizoma tufted; stipes much shorter than the frond; frond ovate-lanceolate; pinnæ crowded, overlapping, twisted as in *Polystichum Lonchitis*, scarcely pinnatifid, never pinnate, very broad and obtuse, their divisions slightly notched; clusters of capsules small, round, remaining distinct, submarginal; involucre generally wanting.

Rare; found by Dr. Dickie in a cave by the sea near Aberdeen.

Obs.—This little fern is better known to cultivators than to field-botanists. It has exactly the habit of a *Woodsia*: I only know it as cultivated, and then it appears perfectly distinct. It is reproduced readily from seed, and loses none of its distinguishing characters.

Genus. — *WOODSIA, R. Br.*

Mid-vein of ultimate divisions indistinct, lateral veins branched, free: involucre seated near the extremity of each branch, its base inclosing the base of the capsules, its margin split into capillary segments, which mingle with the capsules.

WOODSIA ILVENSIS.

? *Acrostichum Ilvense*, *Linn. Sp. Pl.* 1528.

Acrostichum Ilvense, *Huds. Fl. Ang.* 451; *Bolt. Fil. Brit.* 14, t. 9.

Polypodium arvonicum, *With. Arr.* 774.

Woodsia Ilvensis, *R. Br. Tr. Linn. Soc.* xi. 173; *Sm. E. F.* iv. 322, *E. B. S.* 2616; *Newm. N. A.* 13, *F.* 137; *Hook. and Arn.* 567.

Woodsia Ilvensis, *Bab.* 409, ad partem.

Rhizoma tufted; stipes sometimes as long as the frond, but generally shorter, distinctly articulated towards the base; frond erect, lanceolate, pinnate; pinnæ oblong, subopposite, lobed or pinnatifid, scaly.

A very small and rare fern. Found on rocks in Caernarvonshire by Mr. W. Wilson; in Durham by Mr. Backhouse; in Forfarshire by Mr. Watson and Mr. Wilson.

WOODSIA ALPINA.

Polypodium fontanum, *Herb. Linn. certè.*

1776. *Acrostichum Ilvense*, *With. Arr.* 649.

1790. *Acrostichum alpinum*, *Bolt. Fil. Brit.* 76, t. 42.

1793. *Acrostichum hyperboreum*, *Liljeblad, St. Tr.* 201, t. 8.

Woodsia hyperborea, *R. Br. Tr. Linn. Soc.* xi. 173; *Sm. E. F.* iv. 323, ? *E. B.* 2023; *Hook. and Arn.* 567.

Woodsia Ilvensis, *Bab.* 409, ad partem.

Woodsia alpina, *Newm. N. A.* 13, *F.* 143.

Rhizoma tufted; stipes shorter than the frond, articulated; frond narrow, linear, pinnate; pinnæ distant, deltoid, blunt, lobed.

A very small and rare fern. Found on rocks in Caernarvonshire and Perthshire by Mr. W. Wilson.

Genus. — ALLOSORUS, *Bernhardi*.

Mid-vein distinct, lateral veins free : involucre not apparent : capsules in circular clusters near the extremity of the lateral veins, which are often divided : epidermis prolonged, bleached, reflexed, entire, and covering the capsules in the manner of an involucre.

ALLOSORUS CRISPUS.

Osmunda crispa, *Linn. Sp. Pl.* 1522 ; *Lightf. Fl. Scot.* 655 ;
Huds. Fl. Ang. 450 ; *Bolt. Fil. Brit.* 10, t. 7.

Pteris crispa, *With. Arr.* 764 ; *Sm. E. F.* iv. 319, *E. B.* 1160.

Cryptogramma crispa, *Mack. Fl. Hib.* 343 ; *Hook. and Arn.* 575.

Allosorus crispus (*Bernh.*), *Newm. N. A.* 13, *F.* 103 ; *Bab.* 408.

Rhizoma prostrate ; stipes as long as the frond ; fronds of two kinds, both deltoid, and divided into numerous leaf-like stipitate divisions.

A small plant. Stony mountain regions in Scotland and the North of England.

Genus. — CTENOPTERIS.

Mid-vein distinct : lateral veins of the pinnæ or pinnules branched, free, swollen or capitate at their extremities ; the anterior branch simple, generally terminating midway between the mid-vein and the margin, bearing a cluster of capsules at its extremity ; the posterior branch is twice or thrice dichotomously divided, the capitate extremities usually forming a line parallel to the margin : involucre none : rhizoma usually attached by means of its roots to the surface of a rock, the bark of a tree, &c., thus always having a pseudo-parasitic or climbing appearance, cylindrical, branched, extending itself at the extremities, at first densely clothed with paleæ, but as these fall off becoming smooth and naked ; of slow growth, tough and very enduring, here and there marked with nearly circular scars, the site of fallen fronds, which, though persistent through the winter, are deciduous in early summer, falling off at a basal articulation. (See *Phytol.* ii. 274).

Obs.—This genus is indicated by Presl under the title of *Polypodium* § *Ctenopteris*, but I am not sufficiently acquainted with the host of exotic species which that author has placed in his sectional divisions to express any opinion as to their affinity. The genus, as restricted above, is sufficiently extensive, and has not, as far as my information extends, been previously isolated.

CTENOPTERIS VULGARIS.

Polypodium vulgare, Linn. *Sp. Pl.* 1544; *Lightf. Fl. Scot.* 667; *Huds. Fl. Ang.* 455; *With. Arr.* 773; *Sm. E. F.* iv. 280, *E. B.* 1149; *Mack. Fl. Hib.* 337; *Newm. N. A.* 13, *F.* 111; *Hook. and Arn.* 566; *Bab.* 408.

Polipodium vulgare, Bolt. *Fil. Brit.* 32, t. 18.

Polypodium Ctenopteris vulgare, Presl, *Tent. Pteridog.* 179.

Stipes articulated at the base, of nearly equal length with the frond; frond pinnatifid.

Walls, trees and hedge-banks, common.

Obs.—The *Polypodium cambricum* of all authors is referrible to this species.

Obs.—The beautiful *Davallia canariensis*, which, together with the extensive family to which it belongs, is intermediate between *Polypodieæ* and *Hymenophylleæ*, may possibly occur in the South-west of Ireland.

Family. — HYMENOPHYLLEÆ.

Frond appearing to consist of branched veins, each accompanied throughout by a membranous wing or margin: cluster of capsules nearly spherical, seated on one of these veins which projects beyond the edge of the leaf, the cluster being inclosed in a kind of cup-like involucre.

Genus. — TRICHOMANES, Linneus.

Involucre elongate, somewhat urn-shaped: capsuliferous vein projecting beyond it in the form of a long stiff bristle.

TRICHOMANES SPECIOSUM.

Trichomanes speciosum, Willd. *Sp. Pl.* v. 514; *Newm. N. A.* 29, *F.* 305.

Trichomanes brevisetum, R. Br. in Ait. *Hort. Kew.* v. 529; *Sm. E. F.* iv. 324; *Mack. Fl. Hib.* 344.

Trichomanes Europæum, *Sm. in Rces' Encyc.* xxxvi.

Trichomanes alatum, *Hook. Flor. Lond.* t. 53.

Hymenophyllum alatum, *Sm. E. B.* 1417.

Trichomanes radicans, (*Newm. in litt.* 1838, non *Swartz*);
Hook. and Arn. 576; *Bab.* 416.

Rhizoma hairy, creeping extensively; stipes about as long as the frond; frond pendulous, triangular, twice or thrice pinnate.

In Ireland only. Shaded wet glens and the vicinity of waterfalls in Cork and Kerry.

Genus. — HYMENOPHYLLUM, *Smith.*

Involucre shorter and rounder than in the preceding genus, and bivalved: the receptacle or capsuliferous vein not longer than the involucre.

HYMENOPHYLLUM TUNBRIDGENSE.

Trichomanes Tunbridgense, *Linn. Sp. Pl.* 1561; *With. Arr.* 781.

? *Trichomanes pyxidiferum*, *Bolt. Fil. Brit.* 56, t. 30.

Trichomanes Tunbridgense, *Huds. Fl. Ang.* 461, ad partem.

Hymenophyllum Tunbridgense, *Sm. E. F.* iv. 326, *E. B.* 162; *Mack. Fl. Hib.* 345; *Newm. N. A.* 29, *F.* 321; *Hook. and Arn.* 577; *Bab.* 416.

Rhizoma filiform, creeping extensively; frond drooping, pinnate; pinnæ alternate, composed of four or five alternate, dichotomously forked divisions, consisting, as in *Trichomanes*, of a membranous wing and a median stiff wiry vein, margins of the wing serrated; involucre in the axil of the pinnæ, flattened, serrated at top.

A small moss-like plant, growing on rocks, local.

HYMENOPHYLLUM UNILATERALE.

Trichomanes Tunbridgense, *Lightf. Fl. Scot.* 681; *Huds. Fl. Ang.* 461, ad partem; *Bolt. Fil. Brit.* 58, t. 31.

1810. *Hymenophyllum unilaterale*, *Willd. Sp. Pl.* v. 521; *Newm. F.* 14.

1830. *Hymenophyllum Wilsoni*, *Hook. Brit. Flor.* 446, *E. B. S.* 2686; *Mack. Fl. Hib.* 345; *Newm. N. A.* 29, *F.* 325; *Hook. and Arn.* 577; *Bab.* 416.

Pinnæ secund, unilateral ; involucre pear-shaped, entire at the top until mature, then dehiscent, the valves widely separating, otherwise as in *H. Tunbridgensis*.

A small moss-like plant. On rocks in mountain regions.

Order. — OSMUNDACEÆ, *R. Brown*.

Plants composed of fibrous roots, solid simple rhizoma, and flat leafy fronds which rise with a circinate vernation. Fructification upon a portion of the frond in which the veins alone remain, the parenchyma being apparently represented by clustered, globose, reticulated capsules, which are not provided with an elastic ring : involucre none.

Genus. — OSMUNDA, *Linneus*.

Characters those of the order.

OSMUNDA REGALIS.

Osmunda regalis, *Linn. Syst. Nat.* 1521 ; *Lightf. Fl. Scot.* 653 ; *Huds. Fl. Ang.* 449 ; *Bolt. Fil. Brit.* 6, t. 5 ; *With. Arr.* 763 ; *Sm. E. F.* iv. 327, *E. B.* 209 ; *Mack. Fl. Hib.* 345 ; *Newm. N. A.* 29, *F.* 331 ; *Hook. and Arn.* 578 ; *Bab.* 417.

Rhizoma very large, tufted ; stipes woody, as long as the frond ; frond nearly erect, and, including the stipes, four to ten feet high, pinnate ; pinnæ opposite, spreading, pinnate ; pinnules alternate, ovate, stalked, very entire ; terminal panicle of capsules golden coloured, large, very conspicuous.

Wet places, very local.

Order. — OPHIOGLOSSACEÆ.

Plants composed of succulent and comparatively stout roots, some of which travel horizontally in the manner of stolons, succulent stipes, and branched fronds with straight vernation. Frond composed of two branches, the outer leafy, the inner, which it seems to inclose at its base in the manner of a spathe, entirely capsuliferous : capsules large, without reticulations, ring, or involucre, opening by a transverse fissure.

Genus. — BOTRYCHIUM, Swartz.

Frond produced annually *within* the base of the fertile frond of the preceding year: exterior or barren branch variously divided; interior or capsuliferous branch also much divided: capsules spherical, sessile, crowded.

BOTRYCHIUM LUNARIA.

Osmundla lunaria, *Linn. Sp. Pl.* 1519; *Lightf. Fl. Scot.* 652; *Huds. Fl. Ang.* 449; *Bolt. Fil. Brit.* 4, t. 4; *With. Arr.* 762; *Sm. E. B.* 318.

Botrychium lunaria, *Sm. E. F.* iv. 328; *Mack. Fl. Hib.* 346; *Newm. N. A.* 30, *F.* 337; *Hook. and Arn.* 578; *Bab.* 417.

Root stout, spreading, sparingly branched; stipes erect; fertile branch of the frond a racemose panicle, the barren branch pinnate; pinnæ three to seven pairs, flabelliform, with crenate margins.

A dwarf plant. Heaths, rather local.

Genus. — OPHIOGLOSSUM, Linneus.

Frond produced annually *exterior* to the base of the last year's frond: exterior or barren branch a simple undivided spathe; interior or fertile branch a simple, erect, stalked, pointed spike, in the substance of which two parallel series of large spherical capsules are embedded; these open by a transverse fissure, giving the spike a serrated appearance.

OPHIOGLOSSUM VULGATUM.

Ophioglossum vulgatum, *Linn. Sp. Pl.* 1518; *Lightf. Flor. Scot.* 651; *Huds. Flor. Ang.* 449; *Bolt. Fil. Brit.* 2, t. 3; *With. Arr.* 761; *Sm. E. F.* iv. 330, *E. B.* 108; *Mack. Fl. Hib.* 346; *Newm. N. A.* 30, *F.* 349; *Hook. and Arn.* 578; *Bab.* 417.

Stipes erect; fertile branch of the frond an erect, club-shaped, pointed spike; the barren branch an entire ovato-lanceolate leaf.

A dwarf plant, growing on heaths and in meadows.

Aspidium lobatum - A. aculeatum, and A. angulare.

These 3 Species of Smith & Hooker & others, are in my opinion properly regarded by Mr. Francis as only constituting two Species.

A. lobatum of Hooker being synonymous with A. lobatum of Francis. A. aculeatum of Hooker, synonymous with A. aculeatum of Francis, -

A. angulare of Hooker, synonymous with A. aculeatum, variety angulare of Francis.

A. lobatum is a very distinct species, characterized by recurrent pinnales - the pinnae are more crowded than in the other, & the pair of pinnales on each pinna next to the rachis are much larger than the rest. The Stipes is shorter, and the frond is of a dark green, narrower & more rigid than A. aculeatum.

A. aculeatum, the pinnales are distinctly petioled. - The frond is larger & broader than that of lobatum, less rigid, stipes longer, the pinnae less crowded - the lowest pair of pinnales on each pinna not much exceeding in size those above it. The varieties

of this species appear to result much from exposure to the Sun & light, when it is found more delicate in structure & more lax, than when growing in a moist & shady place - there are several grades of variety, but nothing in my opinion sufficiently distinct

to constitute two species. I consider A. aculeatum, var. angulare,

of Francis to be a rather smaller, more lax & more delicate yellowish-green frond, having its pinnales more crowded,

more rounded & compact, & more at right angles to the

midribs of the Pinnae, - the pinnales also are more distinctly

petiole to the extremity of the pinna. I prefer the specific
name *A. aculeatum* for the reason assigned by Frezzeris, viz,
because *A. angulare* is the name given to a different fern on
the Continent, which Sprengel describes as having a tripinnate
frond.

Aspidium dilatatum, varietas recurvum, Bree's Fern. - The
pinnales of this Fern are always incurved, instead of
being recurved, giving to the upper surface of each frond a of
each separate pinna a concave appearance, & I have therefore
always considered that this Fern should be named
A. dilatatum, varietas incurvum: By Babington it is
regarded as a distinct species, the *Lastrea Fenisecii*. The name
dilatatum, varietas recurvum, would be more properly applied
to varietas B of A. dilatatum in Francis' British Ferns, in which the
frond and the pinnales are recurved, giving to the upper
surface of the frond a convex appearance -

Botrychium Lunaria - I collected this Fern in July 1851
at Shutehoe Hill, a Spur of the Mendips, about 2 miles from
Bristol & 15 miles from Bristol. - it was growing there not sparingly -
Asplenium septentrionale grows abundantly on the loose
stone walls about Culbone, between Porlock & Lynton, Somersetshire, & also
on the wall at Oare, near Lynton -

Adiantum Capillus Veneris - Collected for me by Mrs Barlow on the
wall of Llanrhaeadr Castle, which is below Cardiff on the Coast of
Wales - also on the rocks about it - Also collected by me
abundantly in July 1853, growing in wet shaded crevices & small
pools on the Coast below Lelant Church, between Hayle & St Ives,
Cornwall - & again on wet rocks at Carrageh Gladden Cove, nearer to St Ives -

Hymenophyllum Wilsoni & H. Tunbridgeense. Both of these Ferns collected (in July 1853) in tolerable abundance on Barn Gallowa, a fine pile of Granite Rocks, between purslane & Gurnards' head - 6 miles from the former.

Asplenium Fontanum. This Fern was collected in 1842, by Mr W. Hutcheson (a Gardener, now residing at Zeeppark, near Margate) on Shaded Rocks by the Sea, 2 miles NE of Stonehaven, near Aberdeen, & it was in considerable abundance (this I don't believe, 77)

Gymnogramma leptophylla. - The localities for this Fern are given in the phytologist, Volume 4, pages 600, 716, 816, 914, 973. at the last mentioned page it is stated to be widely distributed in Guernsey, Jersey, - growing on the banks of exposed lanes, having a southern aspect - more especially in those places where the moistened soil induces the growth of *Marchantia*, - in company with which it particularly thrives, & in Moss. The principal localities for it, are near St. Helier, near St. Aubin's, and in several places near St. Lawrence. I collected this *Gymnogramma* on the Rocks of Gibraltar in March 1834, & in March 1841, in the woods about Lake Avernus, n^r. Naples. This Fern is well figured by Hooker in the "Scenes Filicum" Vol: 1. Pl: 25.

March 24/55 Received from Mr Hammond of 27, Don Terrace, Jersey, young plants of *Gymnogramma*, collected in that Island by his sister. They grew & must have matured seed in my Fern-house, - as I had fine plants the next year, - altho' this an annual plant.

Aspidium spinulosum, var^s A of Hooker, is A. spinulosum
of Francis - and A. spinulosum, var^s B of the (i.e. dilatatum)
the A. dilatatum of Francis. I agree with the latter that
these are distinct species, - for several years I have cultivated
from both, - viz A. spinulosum of Francis, collected by the Rev. Wilson,
mistake for A. cristatum, on Bausey Heath, nr. Lygon, & the same
species from Mylor near Falmouth - also A. dilatatum of Francis
which abounds in the Bristlington House Woods, whilst I have
never met with the former in this neighbourhood or district.
In A. spinulosum, the fronds are more erect, narrower,
of a lighter colour, of more delicate structure, & less subdivided
than in A. dilatatum, - whilst its ^{Rachis} is less scaly & slenderer.
The different habit of these species, when growing side by side, is
very marked. The fronds of A. dilatatum spread out like
the foliage of a Palm tree.

Aug 23/69. - Aspl² Septentrionale. - Bull, Gardener of Lord,
march at Cauncelegr near Chudleigh, informs me that this
occurs on Limestone Rocks on that property, -

Asplenium Fontanum. Hooker & other English Botanists
consider this to be the same as polytrichum fontanum of Linnaeus
& Hudson, Bolton & Withering; The linear Sorus, with an Inclusion,
proves it to be an Asplenium. In Lowe's British Ferns
Asplenium Halleri is given as the Synonym of A. fontanum. on
the Authority of Brown, Sprengel & DeCandolle. Francis, at
page 48, treats them as distinct - but he is clearly in error -
In the Spring of 1866 I found this plant abundantly in the Digne
above Roche, in the woods above Montbrun March, & above the Castle of
Chillon, & also in other places in the Valley of the Rhone. In the
Botanical Works of the district it is termed A. Halleri - and
A. fontanum is not mentioned. It entirely agrees with the description
of the latter by Francis, - for altho he describes it as from 2 to 6
inches high, whilst the Ferns of the plants in the Valley of the Rhone
are generally from 5 to 7 inches high, & occasionally, 9 or 10 inches,
yet I have found in these same localities many specimens
in full fructification & development not more than 3 inches long.
I have for several years previously cultivated in England the
A. Halleri & A. fontanum of Hieronymus & DeCandolle.
and having considered them to be the same -

The Celebrated N. Haller, the Contemporary of Linnaeus, resided
at Roche. It is probable that, in this district at least, he discovered
the plant & that it was named after him. I prefer this

specific Name would have found it near
a spring, or in humid places. but generally in the
June 1859. Mr Wilks of Torquay, states that *A. frutescens* has
been found by the Rev^d Smith - growing on a wall at
Guspreston - near Leeds - but until I have seen & verified the
plant I shall utterly disbelieve the fact.

Since I made this last entry I have received from the Rev^d H. C. Chandler
a letter in which is the following statement (dated July 21. 1859).

The old person, Mr Smith of Guspreston, informs me that the plant is not
indigenous, but grows freely in their little grounds - It was brought originally,
he says from the Pyrenees.

28
Orange