3411.730 624 T. 3

Lepidopterorum Catalogus

editus a

F. Bryk

Pars 84:

A. Klima

Pyralididae:

Subfam.: Scopariinae, Nymphulinae.

Dr. W. Junk

Verlag für Naturwissenschaften

's-Gravenhage

20. XII. 1937.

Lepidopterorum Catalogus

- Pars 1: Ch. Anrivillius, Chrysopolomidae. 1911. 4 p.
2: A. Pagenstecher, Callidulidae. 1911. 14 p.
3: A. Pagenstecher, Libytheidae. 1911. 12 p.
4: H. Wagner et R. Pfizner, Hepialidae. 1911. 26 p.
5: E. Strand, Noctuidae: Agaristinae. 1912. 82 p.
6: E. Meyrick, Adelidae, Micropterygidae, Gracilariidae. 1912. 68 p.
7: H. Zerny, Syntomidae. 1912. 179 p.
8: L. B. Prout, Geometridae: Brephinae, Oenochrominae. 1912. 94 p.
9: P. Mabille, Hesperidae: Subf. Pyrrhopyginae. — McDunnough, Megathymidae. 1912. 22 p.
10: E. Meyrick, Tortricidae. 1912. 86 p.
11: H. Eltringham et K. Jordan, Nymphalidae: Subfam. Acraeinae. 1913. 65 p.
12, 18, 21, 23: H. Wagner, Sphingidae. 1913—1919. 420 p.
13: E. Meyrick, Carposinidae, Heliodinidae, Glyphipterygidae. 1913. 53 p.
14: L. B. Prout, Geometridae: Subfam. Hemitheinae. 1913. 192 p.
15: K. W. von Dalla Torre, Castniidae: Subfam. Castniinae, Neocastniinae, Pemphigostolinae. 1913. 28 p.
16: E. Strand, Brahmaeidae. — H. G. Dyar et E. Strand: Megalopygidae, Daleceridae, Epipyropidae. 1913. 35 p.
17: E. Meyrick, Pterophoridae, Ormeodidae. 1913. 44 p.
19: E. Meyrick, Hyponomeutidae, Plutellidae, Amphitheridae. 1914. 64 p.
20: K. W. von Dalla Torre, Thyrididae. 1914. 55 p.
22, 24, 26: E. Strand, Arctiidae: Arctiinae, Nolinae, Lithosiinae. 1919—1922. 900 p.
25: K. W. von Dalla Torre, Cymatophoridae. 1921. 38 p.
27: F. Bryk, Baroniidae, Temnopalpidae, Parnassiidae. 1923. 247 p.
28: K. W. von Dalla Torre et E. Strand, Lepidarbelidae. — E. Strand, Heterogyniidae. 1923. 14 p.
29: K. W. von Dalla Torre, Cossidae. 1923. 63 p.
30: K. W. von Dalla Torre, Epipleminidae, Uraniidae. 1924. 57 p.
31: K. W. von Dalla Torre et E. Strand, Aegeriidae. 1925. 282 p.
32: R. van Eecke, Cochlidiionidae (Limacodidae). 1925. 81 p.
33, 71: H. Burgeff et F. Bryk, Zygaenidae. 1926—1936. 332 p.
34: K. W. von Dalla Torre et E. Strand, Psychidae. 1929. 215 p.
35, 37, 39: F. Bryk, Papilionidae. 1929—1930. 675 p.
36: H. Neustetter, Heliconiidae. 1929. 136 p.
38, 40, 41, 44: H. Stichel, Riodinidae 1930—1931. 795 p.
42: F. Bryk, Dioptidae. 1930. 65 p.
43, 46, 48: M. Gaede, Satyridae. 1931. 759 p.
45: F. Bryk, Pericopinae. 1931. 57 p.
47, 64, 69, 74: H. H. Shepard, Hesperidae: Subfam. Pyrginae. 1931—1936. 679 p.

Fortsetzung siehe 3. Umschlagseite.

Ochrab-
ZMLA 622

Lepidopterorum Catalogus

editus a

F. Bryk

Pars 84:

A. Klima

Pyralididae:

Subfam.: Scopariinae et Nymphulinae.

Dr. W. Junk

Verlag für Naturwissenschaften

's-Gravenhage

1937.

INHALTS-VERZEICHNIS

	Seite
Fam. Pyralididae	3
Subf. Scopariinae	7
Subf. Nymphulinae	57

Fam. **Pyralididae.**

Auct. A. Klima.

- Pyrales* Linné, Syst. Nat. (ed. 10), p. 533 (1758). —
 — Scopoli, Ent. Carn., p. 239 (1763).
Pyralides [Schiffermüller et Denis], Syst. Verz. Schm.
 Wien, p. 118 (1776).
Pyralides [*Pyralides Pseudogeometrae (ambiguæ, genui-*
nae, consobrinae, Pseudobombyces (conformæ, dissimiles),
Pseudonoctuae (aequivocæ, veræ), Legitimæ (nitidæ, aqua-
ticæ, personatæ, communes, sociatæ)] Hübner, Samml.
 Eur. Schm., 6. Horde, p. 1 (1796).
Lepidoptera Pyralidiformia Schrank, Fauna Boica,
 2 (2), p. 161 (1802).
Nocturni (pars), *Phalaenites* (pars) Latreille, Hist. Nat.
 Crust. Ins., 14, p. 142, 208 (1805).
Nocturna (pars), *Pyralites* Latreille, Gen. Crust. Ins., 4,
 p. 189, 192, 216 (1809).
Nocturna (pars), *Pyralites* + *Crambites* Latreille, Consid.
 Gen. Ordre Nat. Anim., p. 345, 349, 367, 369 (1810).
 Trib. *Pyralides* (pars), Fam. *Pyralida* + *Tineida* (pars)
 Leach, Edinb. Encycl., 9, p. 135 (1815).
Pyralidae Leach in Samouelle, Entom. Useful Comp.,
 p. 254 (1819).
Diplochilota (pars), *Pyralides* Billberg, Enum. Ins., p.
 — (1820).
Pyralides Trib. *Geometriformes* [Stirpes *Salice* (Fam.
Dubiae, Probabiles, Egregiae, Mirabiles, Venustulae), *Her-*
pyzontes (Fam. *Curvipalpes, Rectipalpes*)], Trib. *Difformes*
 [Stirpes *Idiae* (Fam. *Monstrosæ, Fallaces, Nitidæ*), *Helia-*
cæ (Fam. *Purpureæ, Fuscæ, Nigræ, Virides*)], Trib.
Vulgares [Stirpes *Palpitæ* (Fam. *Ambiguæ, Teneræ, Re-*
pandatæ, Unicolores, Nobiles, Notatæ), *Elophilæ* (Fam.
Strigatæ, Ornatae), *Tetrachilæ* (Fam. *Margaritiferae, Di-*
stinctæ, Aeratæ, Dubiæ, Obsoletæ, Politæ)] Hübner, Verz.
 bek. Schm., p. 339—372 (1826). — Stephens, Ill. Brit. Ent.
 Haustellata, 4, p. 416, 417 (1834) (1835).
Pyralidae Stephens, Syst. Cat. Brit. Ins., p. 157 (1829).
 Fam. *Notturni* (pars), Sezione *Pyralidi* O. G. Costa,
 Fauna Regno Napoli (1832—36).
Pyralina (pars), *Pyralites* + *Crambites* Newman, Entom.
 Mag., 2, p. 384 (1834).
Semidiurna (pars), *Pyralidae* Stephens, Ill. Brit. Ent.
 Haustellata, 4, p. 9, 397 (1834).
 Trib. *Pyralæ* (Subtrib. *Hercynites, Ennychites, Pyralites,*
Nymphulites, Scopulites, Botytes, Cledeobytes, Aglossites,
Herminites) + Trib. *Schoenobidæ* + Trib. *Crambidae* Du-
 ponchel, Cat. Méth. Lep. Eur., p. 194, 313, 314 (1844).
Crambides + *Pyralidides* Herrich-Schaeffer, Syst. Be-
 arb. Schm. Eur., 4, p. 3—6 (1848), p. 118 (1850).

Pyralidae (Subfam. *Herminidi*, *Cledeobidi*, *Aglossidi*, *Nymphulidi*, *Botydi*, *Scopulidi*, *Ennychidi*) Stephens, List Spec. Spec. Brit. Anim., 5, p. 230 (1850).

Pyralidoidae Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 5, p. 41 (1852).

Pyralites, Trib. *Squamosae* (Fam. *Odontidae*), + *Pulverulentae* (Fam. *Pyralidae*, *Cleodobidae*) + *Luridae* (Fam. *Hercynidae*, *Ennychidae*, *Asopidae*, *Steniidae*, *Hydrocampidae*, *Spilomelidae*, *Margarodidae*, *Botyidae*) + *Plicatae* (Fam. *Scoparidae*) Guenée, Delt. et Pyral., p. 97 (1854).

Pyralidina (pars), *Pyralites* (Fam. *Odontidae*, *Pyralidae*, *Cledeobidae*, *Ennychidae*, *Asopidae*, *Steniidae*, *Hydrocampidae*, *Botyidae*, *Nolidae*, *Choreutidae*) + *Crambites* (Fam. *Eudoreidae*, *Galleridae*, *Phycitidae*, *Crambidae*, *Cholephoridae*) Stainton, Man. Brit. Butterfl. Moths, 2, p. 123, 132, 159 (1859).

Pyralides, Trib. *Pyralites* [Subtrib. *Squamosae* (Fam. *Odontidae*) + *Pulverulentae* (Fam. *Pyralidae*, *Cledeobidae*) + *Luridae* (Fam. *Hercynidae*, *Ennychidae*, *Asopidae*, *Steniidae*, *Hydrocampidae*, *Spilomelidae*, *Margarodidae*, *Botyidae*) + *Plicatae*] + Trib. *Crambites* [Fam. *Galleridae*, *Phycitidae*, *Crambidae*] Walker, List Spec. Ins. Brit. Mus., 17, 18, 19, p. 255—1022 (1859); 27, p. 1—202 (1863); 34, p. 1224—1533 (1865); 35, p. 1701—1763 (1866).

Pyralidina + *Crambina* Wocke, Cat. Lep. Eur. Microlep., p. 85 (1861).

Pyralidinen (Fam. *Chrysaugidae*, *Hamalochroidae*, *Semniidae*, *Pyralididae*, *Homophysidae*) + *Crambinen* Lederer, Wien. Ent. Monatschr., 7, p. 256 (1863).

Pyralidina (Fam. *Pyralididae*, *Botidae*, *Chilonidae*, *Crambidae*, *Phycideae*, *Galleriae*) Heinemann, Kleinschm., 1, H. 2, p. 1 (1865).

Fam. *Pyralides* (Gruppen *Pyralididae*, *Botyidae*, *Chilonidae*, *Crambidae*, *Phycideae*, *Galleridae*) Wallengren, Oefvers. Kgl. Vet. Akad. Förh., 8, p. 965 (1871).

Pyralidina (*Pyralididae*, *Acentropodidae*, *Chilonidae*, *Crambidae*, *Phycideae*, *Galleriae*) Wocke, Cat. Lep. Eur. Faun. Microlep., p. 201 (1871).

Fam. *Pyralidina* (*Pyralidae*, *Botyidae*, *Crambidae*, *Galleridae*, *Phycidae*) Snellen, Vlinders Nederl., 2 (1), Microlep., p. 6 (1882).

Trib. *Pyrales* [Fam. *Botydidae*, *Margaronidae*, *Asopidae*, *Spilomelidae*, *Ennychidae*, *Hercynidae*, *Steniidae*, *Siculidae*, *Hydrocampidae*, *Pyralidae*, *Scoparidae*] + Trib. *Crambites* [Fam. *Phycitidae* (Subfam. *Phycitinae*, *Anerastinae*), *Schoenobidae*, *Crambidae*, *Ancylolomidae*, *Galleridae*] Swinhoe et Cotes, Cat. Moths India, 5, p. 591 (1889).

Pyralidina, Fam. *Pyralidae* (Subfam. *Pyraustinae*, *Scopariinae*, *Schoenobiinae*, *Hydrocampinae*, *Acentropodinae*, *Homophysinae*, *Musotiminae*, *Chrysauginae*, *Endotrichinae*, *Pyralidinae*, *Epipaschiinae*) + Fam. *Crambidae* (Subfam. *Oxychirotnae*, *Crambinae*, *Ancylolominae*, *Phycitinae*, *Galleriinae*) Ragonot, Ann. Soc. Ent. Fr., (s. 6) 10, p. 445 (1890) (1891).

Pyralidina, Fam. *Phycitidae* (p. 361) + *Galleriidae* (p. 383) + *Crambidae* (p. 386) + *Pyraustidae* (p. 392) + *Pyralididae* (p. 425) Meyrick, Handb. Brit. Lep., p. 360 (1895).

- Pyralidae* (Subfam. *Chrysauginae*, *Pyralidinae*, *Phycitinae*, *Schoenobiinae*, *Crambinae*) Druce, Biol. Centr.-Amer. Het., 2, p. 188 (1895, 1896), p. 545 (1899).
- Pyralidae* (Subfam. *Galleriinae*, *Crambinae*, *Schoenobiinae*, *Anerastiinae*, *Phycitinae*, *Epipaschiinae*, *Chrysauginae*, *Endotrichinae*, *Pyralinae*, *Hydrocampinae*, *Scopariinae*, *Pyraustinae*) Hampson, Fauna Brit. India Moths, 4, p. 1 (1896).
- Pyrales* (*Odontidi*, *Pyralididi*, *Cledeobiidi*, *Hercynidi*, *Ennychidi*, *Agroteridi*, *Steniidi*, *Hydrocampidi*, *Margarodesidi*, *Botysidi*, *Scopariidi*, *Chilonidi*, *Crambidi*, *Nephopterycidi*, *Galleriidi*) Acloque, Faune France, 3, p. 249, 343—349 (1897).
- Pyralidae* (Subfam. *Galleriinae*, *Crambinae*, *Schoenobiinae*, *Anerastiinae*, *Phycitinae*, *Epipaschiinae*, *Chrysauginae*, *Endotrichinae*, *Pyralinae*, *Hydrocampinae*, *Scopariinae*, *Pyraustinae*) Hampson, Proc. Zool. Soc. Lond., p. 591 (1898).
- Stemmatoneopoda* (pars) Karsch, Ent. Nachr., 24, p. 302, 303 (1898).
- Galleriidae* (p. 415) + *Crambidae* (p. 416) + *Schoenobiidae* (p. 418) + *Epipaschiidae* (p. 421) + *Chrysaugidae* (p. 426) + *Endotrichidae* (p. 426) + *Pyralidae* (p. 429) + *Nymphulidae* (p. 437) + *Pyraustidae* (p. 452) (Subfam. *Hymeniinae*, *Pinaciinae*, *Dichocrociinae*, *Syleptinae*, *Margaroniinae*, *Pyraustinae*) Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2 (1900).
- Pyralidae* (Subfam. *Galleriinae*, *Crambinae*, *Schoenobiinae*, *Anerastiinae*, *Phycitinae*, *Epipaschiinae*, *Chrysauginae*, *Endotrichinae*, *Pyralinae*, *Hydrocampinae*, *Scopariinae*, *Pyraustinae*) Staudinger et Rebel, Cat. Lep. Pal. Faun., 2, p. 1, (1901).
- Pyralidae* (Subfam. *Pyraustinae*, *Nymphulinae*, *Scopariinae*, *Pyralinae*, *Chrysauginae*, *Schoenobiinae*, *Crambinae*, *Galleriinae*, *Epipaschiinae*, *Phycitinae*, *Anerastiinae*) Fernald et Hulst in Dyar, List North Amer. Lep. (Bull. U. S. Nat. Mus., nr. 52), p. 371 (1902).
- Pyralidina* (pars), *Pyralites* (Fam. *Pyraustidae*, *Pyralidae*, *Hydrocampidae*, *Endotrichidae*, *Scopariidae*) + *Phycitidae* + *Anerastiidae* + *Crambidae* + *Galleriidae* Barrett, Lep. Brit. Isl., 9, p. 160, 161 (1904); 10, p. 1 (1905).
- Pyralidae* (Subfam. *Galleriinae*, *Crambinae*, *Schoenobiinae*, *Anerastiinae*, *Phycitinae*, *Epipaschiinae*, *Endotrichinae*, *Pyralinae*, *Hydrocampinae*, *Scopariinae*, *Pyraustinae*) Rebel in Spuler, Schm. Eur., 2, p. 189 (1906).
- Pyralididae* (Subfam. *Galleriinae*, *Crambinae*, *Schoenobiinae*, *Anerastiinae*, *Phycitinae*, *Pyralidinae*, *Hydrocampinae*, *Scopariinae*, *Pyraustinae*) Wahlgren, Svensk Insekt-fauna, 10 (2), p. 6 (1915).
- Microfrenatae* (pars) Heymons in Brehm, Tierleben, Vielfüßler, Insekten u. Spinnenkerfe, p. 217 (1915).
- Pyralidoidea* (pars), *Pyralididae* (Subfam. *Galleriinae*, *Crambinae*, *Pyralinae*, *Phycitinae*, *Pyraustinae*, *Epipaschiinae*) Mosher, Bull. Illinois State Lab. Nat. Hist., 12, art. 2, p. 32, 69, 72 (1916).
- Pyralidae* (Subfam. *Pyraustinae*, *Nymphulinae*, *Scopariinae*, *Pyralinae*, *Chrysauginae*, *Schoenobiinae*, *Crambinae*, *Galleriinae*, *Macrothecinae*, *Epipaschiinae*, *Endotrichinae*, *Phycitinae*, *Anerastiinae*) Barnes et Mc Dunnough, Check List Lep. Bor. Amer., p. 128 (1917).
- Pyralidina* (pars) (*Galleriinae*, *Crambinae*, *Schoenobiinae*, *Anerastiinae*, *Phycitinae*, *Epipaschiinae*, *Chrysauginae*, *En-*

- dotrichinae*, *Pyralinae*, *Hypotiinae*, *Hydrocampinae*, *Scopariinae*, *Pyraustinae*) Janse, Checklist South Afr. Lep. Het., p. 141 (1917).
- Pyralidae* Subfam. *Tineinae* (= *Galleriinae*), *Crambinae*, *Siginae* (= *Schoenobiinae*), *Hyposotropinae* (= *Anerastiinae*), *Anerastiinae* (= *Phycitinae*), *Pococerinae* (= *Epipaschianae*), *Semniinae* (= *Chrysauginae*), *Endotrichinae*, *Pyralinae*, *Nymphulinae* (= *Hydrocampinae*), *Scopariinae*, *Agroterinae* (= *Pyraustinae*) Hampson, Novit. Zool., 25, p. 386 (1918).
- Pyralidae* (Subfam. *Pyralinae*, *Endotrichinae*, *Schoenobiinae*, *Hydrocampinae*, *Scopariinae*, *Pyraustinae*, *Anerastiinae*, *Phycitinae*, *Crambinae*, *Galleriinae*) Börner in Brohmer, Fauna v. Deutschland, 2. Aufl., p. 331, 338, 339, 348 (1920).
- Pyralidoidea* (pars) Fam. *Pyralididae* (Subfam. *Schoenobiinae*, *Chrysauginae*, *Galleriinae*, *Macrothecinae*, *Glaphyriinae*, *Pyraustinae*, *Nymphulinae*, *Scopariinae*, *Pyralidinae*, *Ancylolominae*, *Crambinae*, *Epipaschiinae*, *Phycitinae*, *Anerastiinae*) Forbes, Lep. New York p. 39, 521 (1923).
- Pyraloidea* (pars), Fam. *Pyralidae* [Subfam. *Pyralinae* (Trib. *Galleriini*, *Pyralini*) + *Pyraustinae* (Trib. *Pyraustini*, *Crambini*)] Börner in Brohmer, Fauna v. Deutschland, 3. Aufl., p. 366, 376 (1925).
- Pyralidina* (pars), Fam. *Pyralididae* (Trib. *Pyraustini*, *Scopariini*, *Nymphulini*, ? *Hypotiini*, *Pyralidini*, *Endotrichini*, *Chrysaugini*, *Pococerini*, *Phycidini*, *Anerastiini*, *Schoenobiini*, *Crambini*, *Galleriini*, ? *Oxychirotini*, ? *Atterini*, *Brachypodicolini*) Handlirsch, Handb. Entom., 3, p. 900—904 (1925).
- Pyralidina* (pars), Fam. *Phycitidae* (p. 376) + *Galleriidae* (p. 399) + *Crambidae* (p. 402) + *Pyraustidae* (p. 414) + *Pyralididae* (p. 443) Meyrick, Revis. Handb. Brit. Lep., p. 29, 375 (1928).
- Crambidae* (= *Pyralidae*) (Subfam. *Crambinae*, *Phycitinae*, *Anerastiinae*, *Schoenobiinae*, *Hypotiinae*, *Endotrichinae*, *Aglossinae*, *Nymphulinae*, *Scopariinae*, *Pyraustinae*) De Joannis, L'Amat. Papill., 4, p. 264 (1929).
- Crambidae* (= *Pyralidae*) (Subfam. *Crambinae*, *Anerastiinae*, *Phycitinae*, *Schoenobiinae*, *Hypotiinae*, *Endotrichinae*, *Aglossinae*, *Nymphulinae*, *Scopariinae*, *Pyraustinae*) + *Galleriidae* De Joannis, L'Amat. Papill., 6, p. 6, 7, 139 (1932).
- Pyralididae* (Subfam. *Galleriinae*, *Crambinae*, *Schoenobiinae*, *Anerastiinae*, *Phycitinae*, *Epipaschiinae*, *Endotrichinae*, *Pyralinae*, *Hydrocampinae*, *Scopariinae*, *Pyraustinae*) Hering, Tierwelt Mitteleur., Ergbd. 1, p. 253 (1932).
- Pyralidae* (Subfam. *Galleriinae*, *Crambinae*, *Schoenobiinae*, *Anerastiinae*, *Phycitinae*, *Endotrichinae*, *Pyralinae*, *Hydrocampinae*, *Scopariinae*, *Pyraustinae*) Eckstein, Schmetterl. Deutschl., 5, p. 45, 49 (1933).
- Crambidae* [Subfam. *Anerastiinae* (= *Hyposotropinae*), *Phycitinae*, (= *Anerastiinae*), *Crambinae*, *Schoenobiinae*, *Hypotiinae*, *Endotrichinae*, *Aglossinae*, *Nymphulinae*, *Scopariinae*, *Pyraustinae* (= *Agroticinae*)] + *Galleriidae* Lhomme, Cat. Lep. France Belge, 2, fasc. 1, p. 5, 161 (1935).
- Pyralidina* (pars), Fam. *Pyralididae* (Subfam. *Pyraustinae*, *Scopariinae*, *Nymphulinae*, *Pyralidinae*, *Endotrichinae*, *Chrysauginae*, *Pococerinae*, *Phycitinae*, *Anerastiinae*, *Macrospiratinae*, *Schoenobiinae*, *Crambinae*, *Galleriinae*, *Macrothecinae*) Zerny et Beier in Kükenthal et Krumbach, Handb. Zool., 4 (2), Ins. 2, Lief. 8, p. 1678, 1700—1702 (1936).

1. Subfam. **Scopariinae** Ragonot

Ragonot, Ann. Soc. Ent. Fr., (s. 6) 10, p. 445, 446, 453 (1890) (1891). — Hampson, Fauna Brit. India Moths, 4, p. 2, 241 (1896); Trans. Ent. Soc. Lond., p. 223, (1897). — Pagenstecher, Zoologica (Chun) H. 29, p. 184 (1900). — Staudinger et Rebel, Cat. Lep. Pal. Faun., 2, p. 50 (1901). — Fernald in Dyar, List North Amer. Lep. (Bull. U. S. Nat. Mus., nr. 52), p. 397 (1902). — Rebel in Spuler, Schm. Eur., 2, p. 189, 223 (1906). — Dyar, Journ. N. York Ent. Soc. 14, p. 99 (1906). — Wahlgren, Svensk Insektfauna, 10 (2), p. 11, 49 (1915). — Barnes & Mc. Dunnough, Check List Lep. Bor. Amer., p. 136 (1917). — Börner in Brohmer, Fauna v. Deutschl., 2. Aufl., p. 348 (1920). — Forbes, Lep. N. York, p. 524, 581, f. 352, 353, 354 (1923). — De Joannis, L'Amat. Papill., 4, p. 285 (1929); 6, p. 8, 20 (1932). — Hering, Tierwelt Mitteleur. Ergbd. 1, p. 284 (1932). — Eckstein, Schm. Deutschl., 5, p. 45, 63 (1933). — Lhomme, Cat. Lep. France Belge, 2, fasc. 1, p. 105 (1935). — Zerny & Beier in Kükenenthal & Krumbach, Handb. Zool., 4 (2), Ins. 2, Lief. 8, p. 1701 (1936).

Scoparidae (pars) Guenée, Delt. et Pyral., p. 412, 416 (1854). — Walker, List Spec. Lep. Ins. Brit. Mus., 18, p. 811 (1859); 34, p. 1496 (1865). — Swinhoe & Cotes, Cat. Moths India, 5, p. 666 (1889).

Scopariadae Meyrick, Trans. Ent. Soc. Lond., p. 342 (1884); Trans. Proc. N. Zeal. Inst., 17, p. 68, 69 (1865).

Scopariidi Acloque, Faune France, 3, p. 346 (1897).

Scopariidae Barrett, Lep. Brit. Isl., 9, p. 306 (1904).

Scoparianae Hampson, Proc. Zool. Soc. Lond., p. 591 (1898). — Janse, Check-list South Afr. Lep. Het., p. 157 (1917). — Hampson, Novit. Zool., 25, p. 386 (1918).

Scopariini Handlirsch in Schröder, Handb. Entom., 3, p. 902 (1925).

Botidae (part.) Heinemann, Kleinschm., 1, H. 2, p. 8, 18 (1865).

Botydae (part.) Wallengren, Oefvers. Kgl.-Vet. Akad. Förh., 8, p. 965, 971 (1871). — Snellen, Vlinders Nederland, 2 (1), Microlep., p. 19 (1882).

Crambidae (part.) Duponchel, Cat. Méth. Lep. Eur., p. 313 (1844). — Wollaston, Ann. & Mag. Nat. Hist., (S. 3), 1, p. 119 (1858) (part.).

Crambides (part.) Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 3—6 (1848).

Crambina (part.) Wocke, Cat. Lep. Eur. Microlep., p. 85 (1861).

Eudoridi Stephens, List Spec. Brit. Anim., 12, p. 2 (1852).

Eudoreidae Stainton, Man. Brit. Butterfl. Moths, 2, p. 159 (1859).

Pyralididae (part.) Lederer, Wien. Ent. Monatschr., 7, p. 334 (1863). — Wocke, Lat. Lep. Eur. Faun., p. 201 (1871).

Pyralidinae (part.) Druce, Biol. Centr.-Amer. Het., 2, p. 200 (1895, 1896); p. 545 (1899).

Pyraustinae (part.) Trib. *Pyraustini* (part.) Börner in Brohmer, Fauna v. Deutschl., 3. Aufl., p. 377 (1925).

Pyraustidae (part.) Meyrick, Trans. Ent. Soc. Lond., p. 436 (1890); Handb. Brit. Lep., p. 397 (1895); Revis. Handb. Brit. Lep., p. 375, 414 (1928). — Tillyard, Ins. Austral. & N. Zeal., p. 430, 431 (1926).

1. Genus **Micraglossa** Warren

Warren, Ann. Mag. Nat. Hist., (s. 6), 8, p. 65 (1891) (Typus generis: *M. scoparialis* Warr.). — Hampson, Fauna Brit. India Moths, 4, p. 241 (1890); Trans. Ent. Soc. Lond., p. 224 (1897).

Microglossa Hampson, Ann. Mag. Nat. Hist., (S. 8) 20, p. 279, 280 (1917) (*M. philippinensis* Hampson, *M. cupritincta* Hampson.).

scoparialis Warren, Ann. Mag. Nat. Hist., (s. 6) 8, Assam, p. 65, 66 (♀) (1891). — Hampson, Fauna Sikkim, Brit. India Moths, 4, p. 242, f. 146 (♂) Südchina (1896); Trans. Ent. Soc. Lond., p. 224, fig. (Typus: (♂) (1897). Darjeeling)

convatalis Klunder van Giejn, Tijdschr. Ent., 55, Key-Ins. p. 352 (♂) (1912).

aenealis Hampson, Trans. Ent. Soc. Lond., p. 224 Assam (1897); Journ. Bombay Nat. Hist. Soc., 12, (Khasis), p. 705 (1899).

flavidalis Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, Westchina p. 20 (♂) (1907).

distictalis Hampson, Journ. Bombay Nat. Hist. Soc., Ceylon 18, p. 575 (♀) (1908).

philippinensis Hampson, Ann. Mag. Nat. Hist., (s. Philippinen 8) 20, p. 279 (♂) (1917) (*Microglossa*).

cupritincta Hampson, Ann. Mag. Nat. Hist. (S. 8) Holl. 20, p. 280 (♂) (1917) (*Microglossa* err. typ.). Neu-Guinea

7

2. Genus **Eclipsiodes** Meyrick

Meyrick, Trans. Ent. Soc. Lond., p. 343 (1884). — Hampson, Trans. Ent. Soc. Lond., p. 224, 225 (1897).

(Typus generis: *E. crypsixantha* Meyr.)

crypsixantha Meyr., Trans. Ent. Soc. Lond., p. 343 S. Australien, (♂ ♀) (1884). — Hampson, Trans. Ent. Soc. Lond., p. 225, fig. (♂) (1897). — Queensland, Lower, Proc. Linn. Soc. N. S. Wales, 14 (24), p. 91 (1899). N. S. Wales

homora Turner, Trans. Proc. Roy. Soc. S. Austral., N. W. 32, p. 104 (♂) (1908). Australien

drosera Meyrick, Trans. Ent. Soc. Lond., p. 245 Victoria (♂ ♀) (1887). — Hampson, Trans. Ent. Soc. London, p. 225 (1897).

icelomorpha Turner, Trans. Proc. Roy. S. Austral., W. Australien 32, p. 105 (♂) (1908).

- cuprealis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, Victoria
p. 20 (♂) (1907).
striatalis Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, W. Australien
p. 21 (♂) (1907).

Species incertae sedis.

- acrocapna** Turner, Proc. Roy. Soc. Queensland, Queensland
27, p. 51 (♂) (1915).
argolina Lower, Proc. Linn. Soc. N. S. Wales (1901), N. S. Wales
26, p. 665 (♀) (1902). (Typus:
Broken Hill)
dapsilis Turner, Trans. Proc. Roy. Soc. S. Austral., Queensland
32, p. 105 (♂ ♀) (1908).
epigypsa Lower, Proc. Linn. Soc. N. S. Wales, 26 N. S. Wales
(1901), p. 665 (♀) (1902). (Typus:
Broken Hill)
leuconota Lower, Proc. Linn. Soc. N. S. Wales, 26 S. Australien
(1901), p. 666 (♀) (1902). (Typus:
Semaphore)
meliphyrta Turner, Trans. Proc. Roy. Soc. S. Queensland
Austral., 32, p. 106 (♂) (1908).
paumleta Turner, Trans. Proc. Roy. Soc. S. Austral., Queensland
32, p. 106 (♂) (1908).
semigilva Turner, Trans. Proc. Roy. Soc. Victoria, Victoria
35, p. 51 (♂) (1922). (Typus:
Daytrop)

14

3. Genus **Cholius** Guenée

Guenée, Eur. Microlep. Index meth., p. 94 (1845). — Chapman,
Trans. Ent. Soc. Lond. 1911, p. 507 (1912).

(Typus generis: *Ch. ochrealis* [Schiff. & Denis]).

- ochrealis** [Schiffermüller & Denis], Syst. Verzeichn. Südwestl.
Schm. Wien, p. 123 (1776) (*Pyralis*). — Deutschland,
Hübner, Samml. Eur. Schm., 1 (*Pyrallides*), t. Österreich,
8, f. 47 (1796) (*Pyralis*). — Zeller, Linn. Ungarn,
Ent., 1, p. 271 (1846) (*Eudorea*). — Her- Frankreich,
rich-Schaeffer, Syst. Bearb. Schm. Eur., 4, Belgien,
p. 49 (*Tin.*), t. 14, f. 99 (1848) (*Eudorea*). N. u. M.
— Guenée, Delt. et Pyral., p. 431 (1854) Italien,
(*Scoparia*). — Heinemann, Kleinschm., 1, N. Balkan,
H. 2, p. 24 (1865) (*Scoparia*). — Rebel in Mazedonien,
Spuler, Schm. Eur., 2, p. 222 (1906) (*Scoparia*). — S.O.-
— Chapman, Trans. Ent. Soc. Lond., Rußland,
1911, p. 507 (1912) (*Cholius*). — M. He- Taurus,
ring, Tierwelt Mitteleur. (Ergbd. 1), p. Kleinasien,
284 (1932) (*Scoparia*). — Eckstein, Schm. Marokko
Deutschl., 5, p. 63 (1933) (*Scoparia*). (Gr. Atlas).
silacella Hübner, Samml. Eur. Schm., 8.
Horde, p. 37, f. 232 (1796) (*Tinea*). —
Zincken, Germar, Mag. Ent., 2, p. 107
(1817). — Treitschke, Schm. Eur., 9 (1),
p. 130 (1832) (*Chilo*).

simplicella De la Harpe, Bull. Soc. Vaud. Sci. Nat., 6, p. 402 (1861) (*Eudorea*); Mitt. Schweiz. Ent. Ges., 1 (4), f. 89 (1863) (*Eudorea*).

subsp. *cremealis* Zerny, Deutsche Ent. Zeitschr. Iris, 48, p. 9 (1934). Nörtl. Libanon

leucopeplalis Hampson, Trans. Ent. Soc. Lond., p. 396, t. 3, f. 7 (1900) (*Pionea*). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 63 (1901) (*Pionea*). — Müller-Rutz, Mitteil. Schweiz. Ent. Ges., 14 (5), p. 186, 190 (1928) (*Scoparia*). Sarawschan, Fergana, W. Thian, Aksu

2

4. Genus **Scoparia** Haworth

Haworth, Lep. Brit., 3, p. 498 (1812). — Guenée, Delt. & Pyral., p. 416 (1854). — Walker, Cat. Lep. Het. Brit. Mus., 19, p. 814 (1859). — Ledereer, Wien. Ent. Monatschr., 7, p. 347 (1863). — Heinemann, Kleinschm., 1, H. 2, p. 20 (1865). — Wallengren, Öfvers. K. Vet.-Akad. Förh., 28 (8), p. 973 (1871). — Meyrick, Trans. Ent. Soc. Lond., p. 346 (1884); Trans. Proc. N. Zeal. Inst., 17, p. 70 (1884). — Moore, Lep. Ceyl., 3, p. 354 (1886). — Meyrick, Handb. Brit. Lep., p. 399, 319 (1895). — Hampson, Fauna Brit. India Moths, 4, p. 242 (1896); Trans. Ent. Soc. Lond., p. 224, 229 (1897). — Acloque, Faune France, 3, p. 347 (1897). — Barrett, Lep. Brit. Isl., 9, p. 306 (1904). — Dyar, Journ. N. York Ent. Soc., 14, p. 99 (1906). — Rebel in Spuler, Schmett. Eur., 2, p. 223 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 501, t. 35—44 (genit.). — Wahlgren, Svensk Insektfauna, 10 (2), p. 49 (1915). — Tullgren & Wahlgren, Svenska Ins., p. 388, 395 (1922). — Forbes, Lep. N. York, p. 581, f. 353—355 (1923). — Müller-Rutz, Mitt. Schweiz. Ent. Ges., 13, p. 431, f. 1—3 (genit.) (1925). — Tillyard, Ins. Austr. & N. Zealand, p. 431 (1926). — Meyrick, Revis. Handb. Brit. Lep., p. 416 (1928). — Hudson, Butt. Moths N. Zeal., p. 182 (1928). — Börner in Brohmer, Fauna Deutschl., (ed. 3), p. 377; p. 376, f. 594 (Gehörblase) (1925); id., ibid., (ed. 4), p. 389, f. 731 (Gehörblase) (1932). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 284 (1932). — Eckstein, Schm. Deutschl., 5, p. 63 (1933).

(Typus generis: *cembrae* Haw.).

Eudorea Curtis, Brit. Ent., 4 (fasc. 42), p. 170 (1827) (typ. *E. dubitata* Hübn.). — Duponchel, Hist. Nat. Léop., 8 (2), p. 282 (1831). — Stephens, Ill. Brit. Ent. Haust., 4, p. 298 (1834). — Zeller, Linn. Entom., 1, p. 265 (1846). — Stainton, Man. Brit. Moths, 2, p. 160 (1859). — De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 77—89 (1863).

Lissophanes Warren, Ann. Mag. Nat. Hist., (s. 6) 8, p. 67 (1891) (typus: *ceramica* Warr.).

Scopaea Haworth, Lep. Brit., 4, p. 498 (1812) (*S. cembrae* Haw.).

Xeroscopa Meyrick, Trans. Ent. Soc. Lond., p. 349 (1884) (typus: *ejuncida* Knaggs); Trans. Proc. N. Zeal. Inst., 17, p. 190 (1885). — Hampson, Trans. Ent. Soc. Lond., p. 224, 225 (1897).

Anarpiia Chapman, Trans. Ent. Soc. Lond. 1911, p. 507 (1912) (typus: *pyrenaealis* Dup.).

- Dipleurina* Chapman, Trans. Ent. Soc. Lond. 1911, p. 507 (1912) (typus: *crataegella* Hübner.).
Eudoria Chapman, Trans. Ent. Soc. Lond. 1911, p. 507 (1912) (typus: *murana* Curt., nom. emend.).
Scoparona Chapman, Trans. Ent. Soc. Lond., p. 507 (1912) (typus: *centuriella* Schiff. & Den.).
Witlesia Chapman, Trans. Ent. Soc. Lond. 1911, p. 507 (1912) (typus: *pallida* Steph.).

a) Palaearktische Region.

- centuriella* [Schiffermüller & Denis], Syst. Verzeichn. Alpen, Schm. Wien, p. 319 (1776) (*Tinea*). — schles.
 Fabricius, Ent. Syst., 3 (2), p. 289 (1794). Gebirge,
 — Treitschke, Schm. Eur., 9 (1), p. 49 (1832). Schweiz,
 (*Galleria*); 10 (3), p. 268 (1835) (*Galleria*). Frankreich,
 — Hübner, Samml. Eur. Schm., 8. Horde Finnland,
 t. 35, f. 239 (1796) (*Tin.*). — Zincken, Ger- Schweden,
 mar, Mag. Ent., 4, p. 245 (1821). — Zeller, Lappland,
 Linn. Entom., 1, p. 275, f. 3 (1846) (*Eudo- Polarregion*,
rea). — Herrich-Schaeffer, Syst. Bearb. Sibirien,
 Schm. Eur., 4, p. 44 (*Tin.*), t. 14, f. 93, 94 Kaukasus;
 (1848) (*Eudorea*). — De la Harpe, Mitth. Labrador,
 Schweiz. Ent. Ges., 1 (4), p. 78 (1863) Alaska,
 (*Eudorea*). — Heinemann, Kleinschm., 1, H. Canada,
 2, p. 24 (1865). — Wallengren, Öfvers. K. Vereinigte
 Vet.-Akad. Förh., 28 (8), p. 973 (1871). Staaten
 Aurivillius, Grönlands Insektfauna, 1, p. 23, (Arizona,
 t. 2, f. 6 (1890). — Hampson, Trans. Ent. Massachu-
 Soc. Lond., p. 234 (1897). — Dyar, Proc. setts,
 Wash. Acad. Sci., 2, p. 498 (1900); Proc. Montana,
 U. S. Nat. Mus., 27, p. 918 (1904). — Rebel N. York,
 in Spuler, Schm. Eur., 2, p. 223, 1906. — Utah,
 Dyar, Journ. N. York Ent. Soc., 14, p. 101 Colorado)
 (♂ ♀) (1916). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 35, f. 2—5 (1912)
 (*Scoparona*). — Wahlgren, Svensk Insekt-
 fauna, 10 (2), p. 50, 51, t. 3, f. 58 (1915).
 — Forbes, Lep. N. York p. 582, f. 352, 356
 (1923). — Müller-Rutz, Mitt. Schweiz. Ent.
 Ges., 13, p. 432, f. II (genit.) (1925). —
 Löfqvist & Karvonen, Not. Ent., 7, p. 40 n.
 79 (1927). — De Joannis, L'Amat. Pap., 6, p.
 33 (1932). — M. Hering, Tierwelt Mitteleur.
 (Ergbd. 1), p. 284 (1932). — Eckstein,
 Schm. Deutschl., 5, p. 63 (1933).
borealis Lefèvre, Ann. Soc. Ent. Fr., 5,
 p. 400, t. 10, f. 9 (1835) (*Eudorea*). — Du-
 ponchel, Hist. Nat. Léop. Suppl., 4, p. 114,
 t. 60, f. 1 (1842) (*Eudorea*).
numeralis Zetterstedt, Ins. Lapp., p. 971
 (1839) (*Scopula*).
muneralis Hampson, Trans. Ent. Soc. Lon-
 don, p. 234 (1897) (*Eudorea*; ex synonym.).
quadratella Zetterstedt, Ins. Lapp., p. 997
 (1839) (*Phycis*).
caecalis Walker, Cat. Lep. Het. Brit. Mus.,
 16, p. 36 n. 24 (♂) (1858) (*Hyppena*) (Neu-
 schottland).

- centurialis* Guenée, Delt. & Pyral., p. 418 (1854).
- caliginosalis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1460 (1865) (*Scopula*).
- ninguidalis* Hulst, Trans. Amer. Ent. Soc., 13, p. 147 (1886).
- albisinuatella* Fernald, Smith's List Lep. Bor. Amer. nr. 4129 (1891), nec Packard, Proc. Boston Soc. Nat. Hist., 11, p. 53 (1865). — Hampson, Trans. Ent. Soc. Lond., p. 234 (1897), nec Packard, l. c.
- frigidella* Hampson, Trans. Ent. Soc. Lond., p. 234 (1897), nec Packard, Proc. Boston Soc. Nat. Hist., 11, p. 53 (1865).
- forma *confluella* Krulikowsky, Soc. Ent., 23, p. 18 (1909). — Caradja, Deutsch. Ent. Zeitschr. Iris, 30, p. 22 (1916). Ostrubland (Kasan), Amurgebiet, Ussuri
- cembrae** Haworth, Lep. Brit., 4, p. 498 (1812) (*Scopea*). — Stainton, Man. Brit. Butt. Moths, 2, p. 161 (1859). — Heinemann, Kleinschm., 1, H. 2, p. 27 (1865). — Knaggs, Ent. Mon. Mag., 5, t. 1, f. 4 (1869). — Berce, Faune Ent. Fr. Lep., 6, p. 216 (1878). — Snellen, Vlinders Nederl., 2, p. 29 (1882). — Leech, Brit. Pyral., p. 13, t. 14, f. 1 (1886). — Snellen, Tijdschr. v. Ent., 32, p. 31 (1889); 37, p. 2 (1894). — Meyrick, Handb. Brit. Lep., p. 423 (1895). — Hampson, Trans. Ent. Soc. Lond., p. 235 (1897). — Barrett, Lep. Brit. Isl., 9, p. 307, t. 409, f. 4, 4a (1904). — Rebel in Spuler, Schm. Eur., 2, p. 223 (1910). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 503, 507, t. 39, f. 24—27 (1912). — Wahlgren, Svensk Insekt-fauna, 10 (2), p. 50, 51 (1915). — Löfqvist & Karvonen, Not. Ent., 7, p. 40 sub n. 80 (1927) (= ? *zelleri*). — Meyrick, Revis. Handb. Brit. Lep., p. 440 (1928). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 286 (1932).
- cembrella* Stephens, Ill. Brit. Ent. Haust., 4, p. 299 (1834) (*Eudorea*). — Wood, Index Entom., p. 298, f. 1440 (1838). — Zeller, Linn. Ent., 1, p. 312 (1846) (*Eudorea*).
- subfusca* Stephens, Ill. Brit. Ent. Haust., 4, p. 299 (1834) (*Eudorea*). — Wood, Index Entom., p. 208, t. 46, f. 1441 (1838). — Zeller, Linn. Entom., 1, p. 313 (1846) (*Eudorea*).
- cembralis* Guenée, Delt. & Pyral., p. 421 (1854).
- erralis* Hampson, Trans. Ent. Soc. London, p. 235 (1897) (ex synon.).
- scotica* White, Ent. Mon. Mag., 8, p. 169 (1872). — Leech, Brit. Pyral., p. 14, t. 4, f. 3 (1886). — South, Entom., 23, p. 273 (1890).
- England, Schottland, Finnland, Estland, Schweden, Belgien, Holland, Frankreich, Schweiz, Deutschland, Österreich, Böhmen, Schlesien, Ungarn, Slawonien, Bosnien, Herzegowina, Dalmatien, Rumänien, Mazedonien, Albanien, N. Italien, N. Spanien, Marokko (Gr. Atlas), N. Persien

- var. *zelleri* Wocke, Jahrb. Schles. Ges. Vaterl. Cultur, p. 96 (1854). — Knaggs, Entom. Ann., p. 126 (1869). — Heinemann, Kleinschmetz., 1, H. 2, p. 26 (1865). — Berce, Faune Ent. Fr. Lep., 6, p. 199 (1878). — Leech, Brit. Pyral., p. 13, t. 14, f. 2 (1886). — Hampson, Trans. Ent. Soc. London, p. 235 n. 13 (1897). — South, Entom., 23, p. 272 (1890). — Barrett, Lep. Brit. Isl., 9, p. 308, t. 409, f. 40 (1904) (*cembrae* var.). — Zerny, Ann. Naturh. Hofmus. Wien, 28, p. 324, t. 26, f. 28 (1914). — Löfqvist & Karwonen, Notul. Ent., 7, p. 40 (1927) (spec.). — M. Hering, Tierwelt Mitteleur. (Erg.-bd. 1), p. 286 (1932). — Eckstein, Schmetterl. Deutschl., 5, p. 63 (1933).
- Biologie:** Wood, Ent. Mon. Mag., 25, p. 126 (1888). — Buckler, Larvae Brit. Butt., 9, p. 183 (1901).
- anatolica** Caradja, Deutsche Ent. Zeitschr. Iris, 30, p. 22 (1917).
- coprista** Meyrick, Exot. Micr., 5, p. 18 (♂) (1936).
- asphodeliella** De la Harpe, Neue Denkschr. allg. Schweiz. Ges. Naturw., 14 (7), p. 35 (1855) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 6, p. 143 (1855) (*Eudorea*). — De la Harpe, Mitt. Schweiz. Ent. Ges., 1 (4), p. 81 (1863) (*Eudorea*). — Zerny, Ann. Naturh. Hofmus. Wien, 28, p. 324 (1914).
- cembrae* (part.) Hampson, Trans. Ent. Soc. London, p. 235 (1897).
- ambigualis** Treitschke, Schm. Eur., 7, p. 184 (1829) (*Hercyna*). — Duponchel, Hist. Nat. Léop., 8 (2), p. 288, t. 229, f. 5a-c (1831) (*Eudorea*). — Zeller, Linn. Entom., 1, p. 289 (1846) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 45, (*Tin.*), t. 16, f. 108 (1848) (*Eudorea*). — Guenée, Del. & Pyral., p. 420 (1854). — Stainton, Man. Brit. Butt. Moths, 2, p. 161 (1859). — Heinemann, Kleinschm., 1, H. 2, p. 28 (1865). — Wallengren, Öfvers. K. Vet.-Akad. Förh., 28 (8), p. 974 (1871). — Berce, Faune Ent. Fr. Lep., 6, p. 198 (1878). — Snellen, Vlinders Nederl., 2, p. 28 (1882). — Leech, Brit. Pyral., p. 14, t. 14, f. 5 (1886). — Meyrick, Handb. Brit. Lep., p. 423 (1895). — Hampson, Trans. Ent. Soc. London, p. 234 (1897). — Knaggs, Entom., 33, p. 109, fig. (1900). — Barrett, Lep. Brit. Isl., 9, p. 312, t. 410, f. 2, 2a, b (1904). — Rebel in Spuler, Schm. Eur., 2, p. 223, t. 82, f. 43 (1906). — Chapman,
- England,
Deutschland,
Österreich,
Ungarn,
Slavonien,
Slavonien,
Bosnien,
Herzegowina
N. Persien
- Kleinasien
(Ak Chebir),
Syrien
- Libanon
(Typus:
Arayah)
- Dalmatien,
Schweiz
- England,
Shetland Ins.,
Schweden,
Lappland,
Finnland,
Estland,
Rußland
St. Petersburg,
Mitteluropa,
Schweiz,
Frankreich,
Belgien,
Andalusien,
Taurus,
Kleinasien,
Syrien,
N. Afrika,
Japan,
Labrador

- Trans. Ent. Soc. Lond. 1911, p. 507, f. 36, f. 8, 10, 11, t. 37, f. 12, 13 (1912). — Zerny, Ann. Hofmus. Naturh. Wien, 28, p. 325, t. 26, f. 29 (1914). — Wahlgren, Svensk Insektfauna, 10 (2), p. 50, 51 (1915). — Filipjev, Notul. Ent., 4, p. 9, f. 1a—c, 3 (genit.), (1925). — Müller-Rutz, Mitteil. Schweiz. Ent. Ges., 13, p. 432, f. 1a (genit.) (1925). — Löfqvist & Karvonen, Not. Ent., 7, p. 40 n. 81 (1927). — Meyrick, Revis. Handb. Brit. Lep., p. 441 (1928). — Gerasimov, Ann. Mus. Zool. Acad. Leningrad, 31, p. 26, t. 8, f. 5, 7 (1930). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 286 (1932). — Eckstein, Schm. Deutschl., 5, p. 63 (1933). — Zerny, Zeitschr. Öst. Ent. Ver., 20, p. 34, t. 3, f. A (genit.) (1935).
- dubitalis* Stephens (nec Hübner), Ill. Brit. Ent. Haust., 4, p. 299 (1834). — Wood, Index Entom., p. 208, f. 1442 (1838).
- ambiguella* Treitschke, Schm. Eur., 9 (1), p. 138 (1832) (*Chilo*); 10 (3), p. 272 (1835) (*Chilo*). — De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 82 (1863) (*Eudorea*).
- incertalis* Guenée, Delt. & Pyral., p. 420 (1854).
- conicella* De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 80 (1863) (*Eudorea*); Neue Denkschr. allg. Schweiz. Ges. Naturw., 20 (2), p. 41 (1864) (*Eudorea*).
- ancipitella* De la Harpe, Neue Denkschr. allg. Schweiz. Ges. Naturw., 14 (7), p. 41, f. 5 (1855); Mitth. Schweiz. Ent. Ges., 1 (4), p. 81 (1863) (*Eudorea*); Neue Denkschr. allg. Schweiz. Ges. Naturw., 20 (2), p. 45 (1864) (*Eudorea*).
- ? var. *atomalis* (Doubleday List) Stainton, Entom. Ann., p. 42 (1855); Man. Brit. Butt. Moths, 2, p. 161 (1859). — Heinemann, Kleinschm., 1, H. 2, p. 24 (1865). — Knaggs, Ent. Mon. Mag., 5, t. 1, f. 8 (1869). — Leech, Brit. Pyral., p. 15, t. 14, f. 6 (1886). — Banks, Ent. Mon. Mag., 26, p. 7, 98 (1890). — Knaggs, Entom., 33, p. 110, fig. (1900). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 51, nr. 949 b (1901). — Barrett, Lep. Brit. Isl., 9, p. 314, t. 410, f. 3, 3a (1904). — Rebel in Spuler, Schm. Eur., 2, p. 223 (1906). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 286 (1932) (*ambiguella* forma).
- ? *taeniatella* Teich, Arbeit. Naturf. Ver. Riga N. F., 6, H., p. 73 (1889); Korr.-Blatt Naturf. Ver. Riga, 45, p. 44 (1902).
- ? *erralis* Guenée, Delt. & Pyral., p. 421 (1854). — De Joannis, Ann. Soc. Ent. Fr., 88, p. 32 (1919).
- England,
Holland,
Livland,
Faeroer Ins.,
Belgien,
Elsaß,
Ural

- var. *octavianella* Mann, Wien. Ent. Monatschr., 3, p. 164 (1859) (*Eudorea*). — De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 86 (1863) (*Eudorea*). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 51, nr. 949a (1901). — Rebel in Spuler, Schm. Eur., 2, p. 223 (1910). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 286 (1932) (*ambigualis* forma).
- ambigualis* (part.) Hampson, Trans. Ent. Soc. Lond., p. 235 (1897).
- ? *signella* Teich, Arbeit. Naturf. Ver. Riga N. F., 6. H., p. 72 (1889); Korr.-Blatt Naturf. Ver. Riga, 45, p. 44 (1902).
- forma *crossi* Bankes, Ent. Mon. Mag., 45, p. 58 (1909). England
- var. *nigra* Hamfelt, Ark. Zool., 10, nr. 25, p. 2 (1917). Faeroer Ins.
- forma *bifascialis* Turati, Atti Soc. Ital. Sci. Nat., 62, p. 63, t. B, f. 23 (1923). Apenninen
- subsp. *kwangtungialis* Caradja, Mem. Acad. Rom., (s. 3) 3, mem. 7, p. 336 (1925). Südchina
- Biologie: Buckler, Larvae Brit. Butt. Moths, 9, p. 189 (1901) (*Eudorea*).
- ulmella** Knaggs (Dale i. l.), Ent. Mon. Mag., 3, p. 217, fig. (1867); v. 5, p. 293, t. 1, f. 12 (1869). — Mason, Ent. Mon. Mag., 23, p. 163, fig. (1886) (*Eudorea*). — Leech, Brit. Pyral., p. 15, t. 14, f. 8 (1886). — Knaggs, Entom., 33, p. 114, fig. (1900). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 51, nr. 949c (1901) (*ambigualis* var.). — Barrett, Brit. Lep. Isl., 9, p. 315, t. 410, f. 4, 4a (1904). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 37, f. 14, 15, 16 (1912). — Rebel in Spuler, Schm. Eur., 2, p. 223 (1910) (*ambigualis* var.). — Zerny, Ann. Naturh. Hofmus. Wien, 28, p. 325, t. 26, f. 30 (1914). — Meyrick, Entom., 54, p. 51 (1921); id., Revis. Handb. Brit. Lep., p. 441 (1928). — Löfqvist & Karvonen, Not. Ent., 27, p. 40 n. 82 (1927).
- ambigualis* (part.) Hampson, Trans. Ent. Soc. London, p. 235 (1897).
- conspicualis* Hodgkinson, Ent. Mon. Mag., 18, p. 134 (1881) (*Eudorea*); Entom., 14, p. 223, t. 14, f. 8, 9 (1881). — Leech, Brit. Pyral., p. 15, t. 14, f. 7 (1886). — South, Entom., 23, p. 273 (1890).
- Morphologie: Filipjev, Notul. Ent., 5, p. 9—13, f. 1, 2, 4 (genit.) (1925).
- Biologie: Buckler, Larvae Brit. Butt., 9, p. 188 (1901) (*Eudorea*).
- klinckowstroemi** Hamfelt, Ark. Zool., 10, nr. 25, p. 25, p. 3, f. 1—3 (1917). Faeroer Ins.
- crucigera** Gerasimow, Annu. Mus. Zool. Acad. Leningrad, 31, p. 26, t. 8, f. 3, 4 (genit.), t. 15, f. 4 (♀) (1930). Buchara

Sicilien,
Sierra
Nevada,
Belgien,
Elsaß,
? Holland,
? Livland

England
Faeroer Ins.
Apenninen
Südchina
England,
Österreich,
N. Ungarn,
Schweiz,
Finnland,
Rußland,
Faeroer Ins.,
Amur, Ost-
turkestan,
Alai,
Ussuri,
Sajowan

- microdentalis** Hampson, Ann. Mag. Nat. Hist., (s. Japan 7) 19, p. 22 (1907).
- basistrigalis** Knaggs, Ent. Mon. Mag., 3, p. 1, fig. (1866); 5, p. 293, t. 1, f. 6 (1869). — England, Sweden, Faeroer Ins.,
Leech, Brit. Pyral., p. 14, t. 14, f. 4 (1886). — Belgien, Frankreich,
Tutt & Briggs, Ent. Mon. Mag., 26, p. 51 (1890). — Porritt, Ent. Mon. Schweiz,
Mag., 26, p. 88 (1890). — South, Entom., Österreich,
23, p. 271 (1890). — Meyrick, Handb. Brit. Ungarn,
Lep., p. 423 (1895) (= *ambigualis*). — Schlesien,
Bower, Ent. Mon. Mag., 31, p. 273 (1895). N. Italien,
— Knaggs, Entom., 33, p. 112, fig. (♂ ♀) Istrien,
(1900). — Barrett, Lep. Brit. Isl., 9, p. Dalmatien,
309, t. 410, f. 1, 1a, b (1904). — Rebel Bosnien,
in Spuler, Schm. Eur., 2, p. 223 (1906). Herzegowina,
Chapman, Trans. Ent. Soc. Lond. 1911, p. Sieben-
502, 503, 507, t. 38, f. 19—23 (1912). — bürgen,
Zerny, Ann. Naturh. Hofmus. Wien, 28, Rumänien,
p. 325, t. 26, f. 31 (1914). — Meyrick, Re- Bulgarien,
vis. Handb. Brit. Lep., p. 441 (1928). — Mazedonien,
M. Hering, Tierwelt Mitteleur. (Ergbd. 1), Kleinasien,
p. 286 (1932) (*ambigualis* forma). Seealpen,
ambigualis (part.) Hampson, Trans. Ent. West-
Soc. London, p. 235 (1897). pyrenäen
- var. *syriaca* Caradja, Deutsche Ent. Zeit- Syrien
schr. Iris, 30, p. 22 (1917).
- Biologie: Buckler, Larvae Brit. Butt., 9, p. 188 (1901) (*Eudorea*).
- incertalis** Duponchel, Hist. Nat. Lép., 8 (2), p. 286, Südeuropa,
t. 229, f. 4 (1831) (*Eudorea*). — Spanien
Zeller, Linn. Entom., 1, p. 279, t. 2, f. 4 (1836) (Sierra
(*Eudorea*). — Herrieh-Schaeffer, Syst. Be- Nevada),
arb. Schm. Eur., 4, p. 45, (*Tin.*), t. 17, f. Frankreich,
118 (1848); (*Pyr.*) t. 22, f. 156 (1849). — Tunis,
Guenée, Delt. & Pyral., p. 419 (1854). — Algier,
De la Harpe, Mitth. Schweiz. Ent. Ges., Marokko,
1 (4), p. 82 (1863) (*Eudorea*). — Berce, (Gr. Atlas),
Faune Ent. Fr. Lep., 6, p. 193 (1878). — Kleinasien,
De Joannis, Ann. Soc. Ent. Fr., 84, p. 78 Syrien,
(1915). — Müller-Rutz, Mitt. Schweiz. Ent. Palästina,
Ges., 13, p. 432, f. I b (genit.) (1925). Kaukasus,
incertella Duponchel, Cat. Méth. Lép. Eur., Armenien
p. 319 (1844).
- ab. *pyrenaealis* Duponchel, Hist. Nat. Lép. Spanien,
Suppl., 4, p. 115, t. 60, f. 2a, b (1842) Süd-
(*Eudorea*). — Rebel in Spuler, Schm. Eur., 2, frankreich,
p. 223 (1906). — Chapman, Trans. Ent. Soc. Italien,
Lond. 1911, p. 507, t. 40, f. 33—35 (1912) Dalmatien,
(*Anarpiä*). Balkan,
pyrenaeella Duponchel, Cat. Méth. Lép. Cypern,
Eur., p. 320 (1844). Kleinasien,
pyrenaeicalis Guenée, Delt. & Pyral., p. Amasien,
419 (1854). — Chrétien, Le Natural., 26, Libanon,
p. 45 (1904). Syrien,
Palästina,
Mauretanien

- prinzi* Schawerda, Verh. z. b. Ges. Wien, Mostar
63, p. 169 (1913) (*incertalis* ab.).
- mandschurica* Christoph, Bull. Soc. Nat. Moscou, Ussurigebiet
56 (1), p. 8 (1881).
- ingratella* Zeller, Linn. Entom., 1, p. 283, f. 5 (1846) Österreich,
(*Eudorea*). — Herrich-Schaeffer, Syst. Be- Ungarn,
arb. Schm. Eur., 4, p. 45, (*Tin.*), t. 15, f. Elsaß,
105, 106 (1848). — De la Harpe, Mitth. Schweiz,
Schweiz. Ent. Ges., 1 (4), p. 82 (1863) Dalmatien,
(*Eudorea*). — Heinemann, Kleinschm., 1, Rumänien,
H. 2, p. 27 (1865). — Zeller, Verh. z. b. Griechenland,
Ges. Wien, 18, p. 595 (1868). — Hampson, land,
Trans. Ent. Soc. London, p. 235 (1897); Balkan,
id., Journ. Bombay Nat. Hist. Soc., 15, p. Südrußland,
212 (1903). — Rebel in Spuler, Schm. Eur., Estland,
2, p. 223 (1910). — Chapman, Trans. Ent. Kleinasien,
Soc. Lond. 1911, p. 507 (1912) (*dubitalis* Anatolien,
var.). — M. Hering, Tierwelt Mitteleur. Armenien,
(Ergbd. 1), p. 286 (1932). Syrien,
ingratalis Guenée, Delt. & Pyral., p. 422 Kaschmir,
(1864). Marokko
(Gr. Atlas)
- var. *nevadalis* Caradja, Deutsche Ent. Zeit- Sierra
schr. Iris, 30, p. 22 (1917). Nevada
- dubitalis* Hübner, Samml. Eur. Schm., 6 (*Pyralides*), Mittel- u.
t. 8, f. 49 (1796) (*Pyralis*). — Treitschke, Südeuropa,
Schm. Eur., 7, p. 183 (1829) (*Hercyna*). — England,
Duponchel, Hist. Nat. Lép. 8 (2), p. 290, Schweden,
t. 229, f. 6 (1831) (*Eudorea*). — Zeller, Livland,
Linn. Entom., 1, p. 285, f. 6 (1846) (*Eudo- Estland,*
rea). — Herrich-Schaeffer, Syst. Bearb. Finnland,
Schm. Eur., 4, p. 45, (*Tin.*), t. 15, f. 107 Belgien,
(1848) (*Eudorea*). — De la Harpe, Mitth. Süd-
Schweiz. Ent. Ges., 1 (4), p. 86 (1863) frankreich,
(*Eudorea*). — Heinemann, Kleinschm., 1, H. Spanien,
2, p. 31 (1865). — Wallengren, Öfvers. K. Italien,
Vet.-Akad. Förh., 28 (2), p. 974 (1871). — Dalmatien,
Berce, Faune Ent. Fr. Lep., 6, p. 200 Balkan,
(1878). — Snellen, Vlinders Nederl., 2, p. Kleinasien,
27 (1882). — Leech, Brit. Pyral., p. 16, Syrien,
t. 14, f. 9 (1886). — Meyrick, Handb. Brit. Palästina
Lep., p. 423 (1895). — Hampson, Trans. Ent.
Soc. London, p. 235 (1897). — Fryer, Ent.
Mon. Mag., 35, t. 1, f. 3 (1899). — Barrett,
Lep. Brit. Isl., 9, p. 318, t. 410, f. 5, 5a
(1904). — Rebel in Spuler, Schm. Eur., 2,
p. 223, t. 82, f. 43, Nachtr.-Taf. 9, f. 58,
(1906). — Adkin, Proc. S. Lond. Ent. Nat.
Hist. Soc., p. 82 (1907). — Chapman, Trans.
Ent. Soc. Lond. 1911, p. 503, 504, f. A
(*nervatura* al. ant.), p. 507, t. 36, f. 6, 7a
9, t. 41, f. 40 (1911). — Wahlgren, Svensk
Insektfauna, 10 (2), p. 50, 51 (1915). —
Löfqvist & Karvonen, Not. Ent., 7, p. 40
n. 83 (1927). — Meyrick, Revis. Handb.
Brit. Lep., p. 440 (1928). — M. Hering,
Tierwelt Mitteleur. (Ergbd. 1), p. 286, f.

- 504 (1932). — Eckstein, Schm. Deutschl., 5, p. 63, t. 1, f. 25 (1933).
- pyralella* Hübner, Samml. Eur. Schm., 8. Horde, p. 32, t. 24, f. 167 (1796) (*Tinea*). — Wood, Index Entom., p. 203, t. 46, f. 1443 (1838). — Stainton, Man. Brit. Butt. Moths, 2, p. 161 (1859).
- ? *confusella* De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 87 (1863) (*Eudorea*).
- ingratella* Knaggs (nec Zeller, 1846), Ent. Mon. Mag., 5, t. 1, f. 2 (1869). — Leech, Brit. Pyral., p. 16, t. 14, f. 10 (1886) (*dubitalis* var.). — South, Entom., 23, p. 273 (1890). — Barrett, Lep. Brit. Isl., 9, p. 319, t. 410, f. 5 b (1904) (*dubitalis* var.). — Adkin, Proc. S. Lond. Ent. Nat. Hist. Soc., p. 82 (1907).
- ab. *tristrigella* Stephens, Ill. Brit. Ent. England, Haust., 4, p. 300 (1834) (*Eudorea*). — Elsaß Wood, Index Entom., p. 208, t. 46, f. 1444 (1838) (*Eudorea*). — Zeller, Linn. Entom., 1, p. 314 (1846) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 5, (*Tim.*) t. 14, f. 98 (1848) (*Eudorea*). — Guenée, Delt. & Pyral., p. 422 (1854). — Lhomme, Cat. Lep. Fr. Belg., 2 (1), p. 107 (1935).
- ab. *purbeckensis* Banks, Proc. Dors. F. Engl. Küste, Club, 10, p. 202, t. 1, f. 8 (1889). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, Elsaß, p. 51, nr. 954 b (1901). — Lhomme, Cat. Belgien Lep. Fr. Belg., 2 (1), p. 107 (1935).
- alba* Tutt, Ent. Rec., 11, p. 142 (1901).
- dubitalis* var. Fryer, Ent. Mon. Mag., 35, t. 1, f. 1, 2, 4 (1899).
- var. *ivanalis* Krulikovsky, Rev. Russe d' Ostrubland Ent., 9, p. 113 (1909).
- dubitalis* Eversmann, Fauna Lep. Volgo-Ural., p. 473 (1844) (*Hercyna*).
- ab. *inspersalis* Galvagni, Verh. z. b. Ges. Österreich Wien, 70, p. 64 (1920).
- forma *australis* Müller-Rutz, Mitt. Ent. Schweiz Ges. Zürich, 5, p. 336 (1920).
- perplexella** Zeller, Isis, p. 165 (1839) (*Eudorea*); Südost-Linn. Entom., 1, p. 282 (1846) (*Eudorea*). Europa — Herrich-Schaeffer, Syst. Bearb. Schm., (Krain, Eur., 4, p. 45, (*Tim.*) t. 16, f. 110—112 Fiume, (1848). — Heinemann, Kleinschm., 1, H. N. Italien, 2, p. 25 (1865). — De la Harpe, Mitth. Dalmatien, Schweiz. Ent. Ges., 1 (4), p. 79 (1863) Griechenland, (*Eudorea*). — Hampson, Trans. Ent. Soc. (Frankreich Lond., p. 234 (1897). — Rebel in Spuler, (Hautes Schm. Eur., 2, p. 224 (1906). — Chapman, Pyr.), Trans. Ent. Soc. Lond. 1911, p. 507, t. 40, f. 31, 32, t. 44, f. 69, 71 (1911). Kleinasien
- perplexalis* Guenée, Delt. & Pyral., p. (Brussa) 423 (1854).
- absconditalis** Christoph in Romanoff, Mém. Lép., 3, Armenien p. 17, t. 1, f. 8 (1887).

- staudingeralis** Mabille, Ann. Soc. Ent. Fr. (s. 4) 9, p. 58, t. 2, f. 6 (1869) (*Eudorea*). — Spanien, Korsika, Hampson, Trans. Ent. Soc. Lond., p. 234 Sardinien (1897).
 ab. *bytinskiella* Schawerda, Intern. Ent. Zeitschr., 28, p. 183 (1934) (*perplexella* var.); Zeitschr. Öst. Ent. Ver., 21, p. 64 (1936) (*staudingeralis* ab.).
- poliophacalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 21 (1907). Syrien (Libanon), ? Taurus
- manifestella** Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 46, (*Tin.*) t. 15, f. 104 (1848) (*Eudorea*). — De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 80 (1863) (*Eudorea*). — Heinemann, Kleinschm., 1, H. 2, p. 32 (1865). — Zeller, Verh. z. b. Ges. Wien, 18, p. 596 (1868). — Hampson, Trans. Ent. Soc. London, p. 235 (1897). — Rebel in Spuler, Schm. Eur., 2, p. 224 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 37, f. 17, 18, t. 44, f. 72 (1912). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 286 (1932).
manifestalis Guenée, Delt. & Pyral., p. 423 (1854).
zelleri Rebel, Ann. Naturh. Hofmus. Wien, 25, p. 392 (1911) (nec Wocke, 1854).
 forma *fasciata* Turati, Natur. Sicil., 23, p. 325, t. 4, f. 33, 34 (1921). S. Schweiz
 forma *italica* Turati, Natur. Sicil., 23, p. 326, t. 4, f. 35—38 (1921). Mittel-Italien
 forma *ticinensis* Müller-Rutz, Mitt. Ent. Ver. Zürich, 5, p. 336 (1920). S. Schweiz
- dicteella** Rebel, Ann. Naturh. Hofm. Wien, 30, p. 149, t. 4, f. 4 (1916). Kreta
- sibirica** Lederer, Verh. z. b. Ver. Wien, v. 3, p. 384, t. 7, f. 2 (1853). Altai, Ural
- phaeoleuca** Zeller, Linn. Entom., 1, p. 306, f. 13 (1846) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 49 (1848) (*Tin.*) t. 29, f. 204 (1850). — Stainton, Man. Brit. Butt. Moths, 2, p. 162 (1859). — De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 85 (1863) (*Eudorea*). — Heinemann, Kleinschm., 1, H. 2, p. 37 (1865). — Berce, Faune Ent. Fr. Lep., 6, p. 204 (1878). — Snellen, Tijds. v. Ent., 40, p. 327, t. 12, f. 7 (1898). — Rebel in Spuler, Schm. Eur., 2, p. 224 (1916). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 39, f. 28—30 (1912). — Zerny, Ann. Naturh. Wien, 28, p. 326, t. 26, f. 32 (1914). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 286 (1932).
phaeoleucalis Guenée, Delt. & Pyral., p. 423 (1854). Alpen, Österreich, Schweiz, Holland, Mehadia, Krain, Bosnien, Herzegowina, Montenegro, Bulgarien, Albanien, N. O. Persien

- vesuntialis* Guenée, Delt. & Pyral., p. 424 (1854).
- sciaphilella* De la Harpe, Neue Denkschr. Schweiz. Ges. Naturw., 14 (1), p. 35 (1855).
- ?*parella* Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 5, (*Tin.*) t. 15, f. 102 (1848).
- murana* Rebel, Ann. Naturh. Hofmus. Wien, 18, p. 302 (1903); 25, p. 395 (1911), nec Curtis (1828).
- ambigualis* Rebel, Ann. Naturh. Hofmus. Wien, 27, p. 323 (1913), nec Treitschke (1829).
- forma *gratiatella* Turati, Atti Soc. Ital. Sci. Nat., 53, p. 583 (1914). N. Italien, Schweiz
- fuscella** Turati, Atti Soc. Ital. Sci. Nat., 53, p. 584, f. 5 a, b (1914). — Müller-Rutz, Mitt. Ent. Zürich, 5, p. 336 (1920); Jahresb. Naturf. Ges. Graubündens, t. 2, f. 1, 2 (1925—26). N. Italien, Schweiz
- vallesialis** Duponchel, Hist. Nat. Lép., 8 (2), p. 284, t. 229, f. 3 (1831) (*Eudorea*). — Zeller, Linn. Entom., 1, p. 300, f. 9, 10 (1846) (*Eudorea*). — Herr.-Schaeff., Syst. Bearb. Schmett. Eur., 4, p. 47, (*Tin.*) t. 15, f. 103 (1848). — Heinemann, Kleinschm., 1, H. 2, p. 33 (1865). — Berce, Faune Ent. Fr. Lép., 6, p. 202, t. 5, f. 6 (1878). — De Joannis, Ann. Soc. Ent. Fr., 84, p. 76 (1915). Westalpen, Schweiz, Frankreich, Karpathen, Ostpyrenäen
- valesialis* Hampson, Trans. Ent. Soc. London, p. 235 (1897) (part.). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 52, nr. 961 a (1901). — Rebel in Spuler, Schm. Eur., 2, p. 224 (1901). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 43, f. 58 (1912) (*Eudoria*). — Müller-Rutz, Mitt. Schweiz. Ent. Ges., 13, p. 432, f. III a (genit.) (1925). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 287 (1932).
- forma *octonella* Zeller, Isis, p. 175 (1839) (*Eudorea*); Linn. Entom., 1, p. 300 (1846) (*Eudorea*). — De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 82 (1863) (*Eudorea*); Neue Denkschr. Schweiz. Ges. Naturw., 20 (2), p. 42 (1864) (*Eudorea*). — Caradja, Deutsche Ent. Zeitschr. Iris, 30, p. 23 (1917). — Lhomme, Cat. Lep. Fr. Belg., 2 (1), p. 109, (1935). Ostalpen, Karpathen
- forma *stauderi* Mitterberger, 27. Jahresber. Wien. Ent. Ver., p. 159 (1917). Österreich
- parella** Zeller, Isis, p. 175 (1839) (*Eudorea*); Linn. Entom., 1, p. 303, f. 11 (1846) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 47, (*Tin.*) t. 15, f. 150, t. 17, f. 119, 120 (1848). — Müller-Rutz, Mitteil. Schweiz. Ent. Ges., 13, p. 433 (1925); 13, p. 504 (1927). Hochalpen, Schweiz

- valesialis* (part.) Hampson, Trans. Ent. Soc. London, p. 235 (1897).
- forma *imparella* De la Harpe, Neue Denkschr. Schweiz. Ges. Naturw., 20 (2), p. 42 (1864). — Millière, Icon. Chen. Lep., 1, p. 403, t. 50, f. 2 (1864). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 51, nr. 961 b (1901). — Müller-Rutz, Mitt. Schweiz. Ent. Ges., 12, p. 505 (1917); 13, p. 433 (1927). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 287 (1932). — Lhomme, Cat. Lep. Fr. Belg., 2 (1), p. 109 (1935).
- gallica** Peyerimhoff, Pet. Nouv., p. 332 (1873). — Millière, Icon. Chen. Lep., 3, p. 338, t. 142, f. 3, 4 (1874). — Berce, Faune Ent. Lep. Fr., 6, p. 215 (1878). — Fettig, Cat. Lep. Alsac., p. 11 (1882). — Hampson, Trans. Ent. Soc. London, p. 235 (1897). — Rebel in Spuler, Schm. Eur., 2, p. 224 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 41, f. 39, t. 44, f. 73, 74 (1912). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 286 (1932).
- alpina** Curtis, Ann. Mag. Nat. Hist. (s. 2), v. 5, p. 117 (1850). — Stainton, Man. Brit. Butt. Moths, 2, p. 163 (1859). — Knaggs, Ent. Mon. Mag., 5, t. 1, f. 18 (1869). — Vaughan, Entom., 13, t. 4, f. 1, 2 (1880). — Leech, Brit. Pyral., p. 15, t. 15, f. 8 (1886). — Hampson, Trans. Ent. Soc. London, p. 236 (1897). — Meyrick, Handb. Brit. Lep. p. 421 (1895). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 52, nr. 963 (1901). — Barrett, Lep. Brit. Isl., 9, p. 336, t. 412, f. 3, 3a, b (1904). — Rebel in Spuler, Schm. Eur., 2, p. 224 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 43, f. 57 (1912) (*Eudoria*). — Longstaff, Butt.-Hunt. Many Lands, p. 21 (1912). — Wahlgren, Svensk Insektfauna, 10 (2), p. 50, 52 (1915). — Meyrick, Revis. Handb. Brit. Lep., p. 438 (1928).
- borealis* Tengström (nec Lefébre (1835)), Bidr. Finl. Fauna, p. 102 (1842). — Teich, Korr.-Blatt Naturf. Ver. Riga, 45, p. 45 (1902). — Hampson, Trans. Ent. Soc. London, p. 236 (1897). — Löfqvist & Karvonen, Notul. Ent., 7, p. 40 (1927).
- gracilalis* Stainton, Man. Brit. Butt. 2, p. 163 (1859). — Heinemann, Kleinschm., 1, H. 2, p. 34 (1865). — Knaggs, Ent. Mon. Mag., 5, t. 1, f. 16 (1869). — Mason, Ent. Mon. Mag., 25, p. 162 (1888). — Briggs, Ent. Mon. Mag., 25, p. 185 (1888). — Barrett, Lep. Brit. Isl., 9, p. 337, t. 412, f. 30 (1904) (*alpina* var.).
- imparella* Wocke (nec De la Harpe), Ent. Zeit. Stettin, 25, p. 201 (1864) (1863).
- Hochalpen,
Schweiz,
Karpauthen
- Frankreich,
Elsaß,
Schweiz
- Nordeuropa
(Gebirge
Schottlands,
Shetland
Ins.,
Norwegen,
Schweden,
Lappland,
Nordkap,
Finnland,
Estland)

- var. *lapponica* Caradja, Deutsche Ent. Zeitschr. Iris, 30, p. 23 (1917). Schweden, Lappland
 Biologie: Buckler, Larvae Brit. Lep., 9, p. 199 (1901) (*Eudorea*).
- vafra** Meyrick, Ent. Mon. Mag., 49, p. 158 (♂) (1913); Revis. Handb. Brit. Lep., p. 439 (1928). England (Chester)
- petrophila** Standfuss sen., Ent. Zeit. Stettin, 9, p. 312 (1848) (*Eudorea*); Bresl. Zeitschr. Ent., 3, p. 25, t. 2, f. 5 (1849). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 6, p. 144 (1855) (*Eudorea*). — Heinemann, Kleinschmetz., 1, H. 2, p. 35 (1865). — Berce, Faune Ent. Fr. Lep., 6, p. 203 (1878). — Hampson, Trans. Ent. Soc. Lond., p. 236 (1897). — Rebel in Spuler, Schm. Eur., 2, p. 22 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507 (1912) (= *sudetica* Zell.). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 287 (1928). — Eckstein, Schm. Deutschl., 5, p. 63 (1933). Alpen, Schlesien (Riesengebirge), Elsaß, Schweiz, Frankreich, Bosnien, Herzegowina
- subsp. *balkanica* Rebel, Sitzber. Akad. Gebirg. Wien, 1, 126, p. 798 (1917). Albanien
- isochroalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) Japan 19, p. 22 (♀) (1907).
- sudetica** Zeller, Isis, p. 175 (1839) (*Eudorea*); Ent. Zeit. Stettin, 4, p. 148 (1843) (*Eudorea*); Linn. Entom., 1, p. 304, f. 12 (1846) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 48, (*Tin.*) t. 16, f. 116, 117, (1848). — De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 85 (1863) (*Eudorea*). — Heinemann, Kleinschm., 1, H. 2, p. 36 (1865). — Wallengren, Öfvers. K. Vet.-Akad. Förh., 28 (8), p. 978 (1871). — Berce, Faune Ent. Fr. Lep., 6, p. 204 (1878). — Hampson, Trans. Ent. Soc. London, p. 236 (1897) (part.). — Rebel in Spuler, Schm. Eur., 2, p. 224 (1906). — Chapman, Trans. Ent. Soc. Lond., 1911, p. 504, f. B (nervat. al. anter.), p. 507, t. 42, f. 50, 51, 52, 53 (1912) (*Eudoria*). — Wahlgren, Svensk Insektfauna, 10 (27), p. 50, 52 (1915). — Löfqvist & Karvonen, Not. Ent., 7, p. 40 n. 85 (1927). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 287 (1932). — Eckstein, Schm. Deutschl., 5, p. 63 (1933). Gebirge Mittel- u. Nord-europas (excl. England), Polarregion, Schweden, Lappland, Estland, Finnland, Frankreich, Geb. Bulgariens
- sudeticalis*** Duponchel, Hist. Nat. Léop. Suppl., 4, p. 427, t. 84, f. 7 (1842). — Guenée, Delt. & Pyral., p. 427 (1854).
- var. *luzialis* Guenée, Delt. & Pyral., p. 427 (1854). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 52, nr. 965 (1901). — Lhomme, Cat. Lep. Fr. Belg., 2 (1), p. 110 (1935). Pyrenäen
- sudetica*** (part.) Hampson, Trans. Ent. Soc. London, p. 236 (1897).

- var. *livonica* Zeller, Linn. Entom., 1, p. 304 (1846). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 52, nr. 965 b (1901). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 287 (1932) (spec. dist.). — Lhomme, Cat. Lep. Fr. Belg., 2 (1), p. 110 (1935). Livland, Lappland, Kanin, Estland, N.-Deutschland)
- talboti** Schmidt, Ann. Mag. Nat. Hist., (s., 10), 14, p. 544, t. 17, f. 6 (♂) (1934). Marokko (Typus: Gr. Atlas)
- ancipitalis** Zerny, Zeitschr. Öst. Ent. Ver., 20, p. 33, t. 3, f. 13, 14 B (genit.) (1935). Castilien
- stenota** Wollaston (Zeller i. l.), Ann. Mag. Nat. Hist., (s. 3) 1, p. 119 (1858) (*Eudorea*). — ? Rebel, Ann. Naturh. Hofmus. Wien, 7, p. 248 (1892). — Hampson, Trans. Ent. Soc. Lond., p. 236 (1897). — Bethune-Baker, Trans. Ent. Soc. Lond., p. 582 (1894). — Rebel, Ann. Naturh. Hofmus. Wien, 11, p. 113 (1896). Madeira (Typus), ? Canaren
- decorella* Stainton, Ann. Mag. Nat. Hist., (s. 3) 3, p. 210 (1859) (*Eudorea*). — Bethune-Baker, Trans. Ent. Soc. Lond., p. 581 (1894).
- melanographa** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 24 (♂) (1907). Azoren
- stenota* Warren (nec Wollaston, 1858), Nov. Zool. Tring, 12, p. 447 (1905).
- semiamplalis** Warren, Nov. Zool. Tring, 12, p. 446 (♂) (1905). Azoren
- scoriella** Wollaston (Zeller i. l.), Ann. Mag. Nat. Hist., (s. 3), 1, p. 119 (1858) (*Eudorea*). — Hampson, Trans. Ent. Soc. Lond., p. 236 (1897). — Warren, Nov. Zool., 12, p. 446 (1905). — Bethune-Baker, Trans. Ent. Soc. London, p. 582 (1894). Madeira
- wollastoni** Bethune-Baker, Trans. Ent. Soc. Lond., p. 583 (♂) (1894). Madeira
- murana** Curtis, Brit. Ent., 4, p. 170 (1828). — Stephens, Ill. Brit. Ent. Haust., 4, p. 301, (1834). — Wood, Index Entom., p. 209, t. 46, f. 1447 (1838). — Zeller, Linn. Entom., 1, p. 316 (1846) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 6, p. 143 (1855) (*Eudorea*). — Stainton, Man. Brit. Butt. Moths, 2, p. 162 (1859). — De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 83 (1865) (*Eudorea*). — Heinemann, Kleinschm., 1, H. 2, p. 38 (1865). — Wallengren, Öfvers. K. Vet.-Akad. Förh., 28 (8), p. 975 (1871). — Zeller, Ent. Zeit. Stettin, p. 81 (1878). — Berce, Faune Ent. Fr. Lep., 6, p. 207 (1878). — Leech, Brit. Pyral., p. 17, t. 15, f. 1 (1886). — Meyrick, Handb. Brit. Lep., p. 421 (1895). — Hampson, Trans. Ent. Soc. London, p. 236 (1897) (part.). — Barrett, Lep. Brit. Isl., 9, p. 320, t. 411, f. 1, 1a (1904). — Rebel in Spuler, Schm. Eur., 2, p. 224 (1906). — England, Schweden, Polarregion, arkt. Norwegen, Gebirg. Mittel-Europa, Alpen, Schweiz, Frankreich, Balkan

- Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 41, f. 44—46 (1912) (*Eudoria*). — Zerny, Ann. Naturh. Hofmus. Wien, 28, p. 326, t. 26, f. 33 (1914). — Wahlgren, Svensk Insektfauna, 10 (2), p. 50, 52, t. 3, f. 59 (1915). — Löfqvist & Karvonen, Not. Ent., 7, p. 40 n. 86 (1927) („Hauptform“). — Meyrick, Revis. Handb. Brit. Lep., p. 439 (1928). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 287 (1932). — Eckstein, Schm. Deutschl., 5, p. 63 (1933).
- muralis* Guenée, Delt. & Pyral., p. 425, t. 10, f. 9 (1854).
- parella* Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 5, (*Tim*), t. 15, f. 101 (1848) (*Eudorea*).
- ?*delphinatalis* Guenée, Delt. & Pyral., p. 425, t. 10, f. 9 (1854).
- var. *tuoniana* Hoffmann, Ent. Zeitg. Stett., 54, p. 130, 141 (1893). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 52, nr. 969b (1901). — Löfqvist & Karvonen, Not. Ent., 7, p. 40 n. 86 (1927) (*S. murana* var.).
- murana* (part.) Hampson, Trans. Ent. Soc. London, p. 236 (1897).
- var. *ranica* Strand, Arch. Naturg., 85 A, 4, p. 38 (1920).
- forma *pseudomurana* Müller-Rutz, Mitteil. Alpen Schweiz. Ent. Ges., 13, p. 434 (1925).
- subsp. *nevadensis* Zerny in Rebel & Zerny, Zeitschr. Ost. Ent. Ver., 12, p. 112 (1927). Sierra Nevada
- Biologie: Buckler, Larvae Brit. Lep., 9, 193 (1901) (*Eudorea*). — Sich, Entom., 43, p. 343 (1910).
- liebmanni** Petry, Ent. Zeit. Stett., 65, p. 245 (1904). Korsika
- virescens** Rebel, Verh. z. b. Ges., 46, p. 174 (♀) (1896). — Caradja, Deutsche Ent. Zeitschr. Iris, 30, p. 23 (♂) (1917). Marokko
- lineola** Curtis, Brit. Ent., 4, p. 170 (1828). — Stephens, Ill. Brit. Ent. Haust., 4, p. 300 (1834). — Wood, Index Entom., p. 208, t. 46, f. 1446 (1838). — Zeller, Linn. Entom., 1, p. 315 (1846) (*Eudorea*). — Curtis, Ann. Mag. Nat. Hist., (s. 2) 5, p. 115 (1851). — Stainton, Man. Brit. Butt. Moths, 2, p. 162 (1859). — Knaggs, Ent. Mon. Mag., 5, t. 1, f. 17 (1869). — Berce, Faune Ent. Fr. Lep., 6, p. 208 (1878). — Leech, Brit. Pyral., p. 17, t. 15, f. 6 (1886). — Meyrick, Handb. Brit. Lep., p. 421 (1895). — Hampson, Trans. Ent. Soc. London, p. 236 (1897). — Barrett, Lep. Brit. Isl., 9, p. 332, t. 412, f. 1, 1a, b (larva) (1904). — Rebel in Spuler, Schm. Eur., 2, p. 224 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 43, f. 61—63 (1912) (*Eudoria*). — Meyrick, Revis. Handb. Brit. Lep., p. 438 (1928). England, Mittel- u. West-Frankreich, Sardinien, Gravosa, Amasien, N. Spanien, Algier, Marokko

lineolalis Guenée, Delt. & Pyral., p. 426 (1854).

Biologie: Porritt, Ent. Mon. Mag., 21, p. 101, 236 (1884). — Buckler, Larvae Brit. Butt., 9, p. 195, t. 155, f. 1 (1901) (*Eudorea*).

resinea Haworth, Lep. Brit., 4, p. 499 (1812) (*Scopaea*). — Curtis, Brit. Ent., 4, f. 170 (1828) (*Eudorea*). — Stephens, Ill. Brit. Ent. Haust., 4, p. 301 (1834) (*Eudorea*). — Wood, Index Entom., p. 209, t. 46, f. 1448 (1838) (*Eudorea*). — Zeller, Linn. Entom., 1, p. 316 (1846) (*Eudorea*). — Stainton, Man. Brit. Butt. Moths, 2, p. 162 (1859). — Heinemann, Kleinschm., 1, H. 2, p. 42 (1865). — Berce, Faune Ent. Fr. Lep., 6, p. 209 (1878). — Snellen, Vlinders Nederl., 2, p. 32 (1882). — Leech, Brit. Pyral., 17, t. 15, f. 2 (1886). — Meyrick, Handb. Brit. Lep., p. 420 (1895). — Hampson, Trans. Ent. Soc. Lond., p. 236 (1897). — Barrett, Lep. Brit. Isl., 9, p. 325, t. 441, f. 3 (1904). — Rebel in Spuler, Schm. Eur., 2, p. 224 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 43, f. 65, t. 44, f. 68, 75 (1912) (*Eudoria*). — Meyrick, Revis. Handb. Brit. Lep., p. 437 (1928). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 284, f. 503 (1932). — Eckstein, Schmett. Deutschl., 5, p. 63 (1933).

resinalis Guenée, Delt. & Pyral., p. 429, t. 10, f. 11 (1854).

vandalicella Herrich-Schaeffer (Zeller MS.), Syst. Bearb. Schm. Eur., (Pyr.), t. 22, f. 157, p. 288 (1849); 6, p. 143 (1855). — De la Harpe, Neue Denkschr. Schweiz. Ges. Naturw., 14 (7), p. 39 (1855); Mitt. Schweiz. Ent. Ges., 1 (4), p. 87 (1863) (*Eudorea*).

vaudalicella Hampson, Trans. Ent. Soc. London, p. 236 (1897) (ex syn.).

orientalis Alphéraky, Trudy Ent. Ross., 10, p. 26 (*resinae* var.).

forma *pernigrata* Turati, Atti Soc. Ital. Sci. Nat., 62, p. 64, t. 3, f. 25 (1923) (*Dipleurina*).

Biologie: Porritt, Ent. Mon. Mag., 23, p. 248 (1886). — Buckler, Larvae Brit. Butt., 9, p. 192 (1901) (*Eudorea*).

laetella Zeller, Linn. Entom., 1, p. 298, f. 8 (1846) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 48, (Tin.), t. 16, f. 115 (1848). — De la Harpe, Mitt. Schweiz. Ent. Ges., 1 (4), p. 87 (1863) (*Eudorea*). — Heinemann, Kleinschm., 1, H. 2, p. 42 (1865). — Andersen, Ent. Tidskr., 18, p. 111 (1897). — Hampson, Trans. Ent. Soc. London, p. 236 (1897). — Rebel in Spuler, Schm. Eur., 2, p. 224 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 43, f. 64, 66 (*laetalis*), t. 44, f. 70 (1912). — Wahlgren, Svensk Insektfauna, 10 (2), p.

England,
Irland,
Dänemark,
Mittel- u.
Südeuropa
(Belgien,
Frankreich,
Schweiz,
Nord- u.
Mittel-
italien,
Korsika,
Sardinien,
N.-Griechenland,
Herzegowina,
Spanien,
Algier,
Portugal),
Marokko
(Gr.
Atlas),
Syrien,
Sibirien

Apenninen

Deutschland,
Österreich,
Schweiz,
Frankreich,
Korsika,
Dalmatien,
Bosnien,
Herzegowina,
Rumänien,
Polen,
Livland,
Estland,

- 50, 52, t. 3, f. 60 (1915). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 287 (1932). — Eckstein, Schm. Deutschl., 5, p. 63 (1933).
laetalis Guenée, Delt. & Pyral., p. 427 (1854).
- tafiarella** Chrétien, Bull. Mus. Nat. Hist. Nat. Paris, 14, p. 358 (♂) (1908). Canaren
- coeruleotincta** Hampson, Ann. Mag. Nat. Hist. (s. 8) 20, p. 281 (♀) (1917). Canaren
- truncicolella** Stainton, Brit. Tin. et Pteroph., p. 2 (1849) (*Eudorea*); Man. Brit. Butt. Moths, 2, p. 161 (1859). — Heinemann, Kleinschm., 1, H. 2, p. 39 (1865). — Berce, Faune Ent. Fr. Lep., 6, p. 210, t. 5, f. 7 (1878). — Snellen, Vlinders Nederl., 2, p. 30 (1882). — Leech, Brit. Pyral., p. 17, t. 16, f. 11 (1886). — Meyrick, Handb. Brit. Lep., p. 422 (1895). — Hampson, Trans. Ent. Soc. Lond., p. 236 (1897). — Barrett, Lep. Brit. Isl., 9, p. 323, t. 411, f. 2, 2a (1904). — Rebel in Spuler, Schm. Eur., 2, p. 225 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 42, f. 47—49 (1912) (*Eudoria*). — Wahlgren, Svensk Insektafauna, 10 (2), p. 50, 52 (1915). — Löfqvist & Karvonen, Not. Ent., 7, p. 40 n. 87 (1927). — Meyrick, Revis. Handb. Brit. Lep., p. 440 (1928). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 287, f. 505 (1932). — Eckstein, Schmett. Deutschl., 5, p. 63 (1933).
- mercurella* Treitschke, Schm. Eur., 9 (1), p. 133 (1832) (*Chilo*); v. 10 (3), p. 272 (1835) (*Chilo*). — Zeller, Linn. Entom., 1, p. 293, f. 1 (1846) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 48 (1848). — De la Harpe, Mitt. Schweiz. Ent. Ges., 1 (4), p. 80, (1863) (*Eudorea*). — Wallengren, Öfvers. K. Vet.-Akad. Förh., 28 (8), p. 975 (1871).
- murcurella* Hampson, Trans. Ent. Soc. Lond., p. 236 (1897) (ex syn.).
- mercurialis* Guenée, Delt. & Pyral., p. 428 (1854).
- ?anissella* Millière, Icon. Chen. Lep., 1, p. 401, t. 59, f. 1 (1864). — Berce, Faune Ent. Fr. Lep., 6, p. 212 (1878).
- ?pallida* Banks, Ent. Mon. Mag., 31, p. 49 (1895).
- Biologie: Porritt, Ent. Mon. Mag., 18, p. 106 (1881). — Buckler, Larvae Brit. Butt., 9, p. 189 (1901) (*Eudorea*).
- ichinosawana** Matsumura, Journ. Coll. Agric. Hokkaido Imp. Univ., 15, p. 185 (1925). Sachalin
- crataegella** Hübner, Samml. Eur. Schm., 8. Horde, p. 32, f. 231 (1796) (*Tinea*). — Zeller, Linn. Entom., 1, p. 296 (1846) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 48, (*Tim.*), t. 16, f. 113, 114 (1848) (*Eudorea*). — Stainton, Man. Brit. Butt. Moths, 2, p. 161 (1859). — De la
- Schweden,
Finnland
- England,
Schweden,
Lappland,
Finnland,
Estland,
Belgien,
Schweiz,
Frankreich,
N. u.
Mittel-
Italien,
Bosnien,
Herzegowina,
Rumänien,
N.W. u. S.O.
Rußland,
N. Kaukasus,
Ural,
Ussuri-
gebiet,
West-
China,
Japan
- Europa,
Kleinasien,
Ostsibirien,
Syrien,
Ussuri,
Westchina

Harpe, Mitt. Schweiz. Ent. Ges., 1 (4), p. 88 (1863) (*Eudorea*). — Heinemann, Kleinschm., 1, H. 2, p. 41 (1865). — Wallengren, Öfvers. K. Vet.-Akad. Förh., 28, (8), p. 977 (1871). — Berce, Faune Ent. Fr. Lep., 6, p. 211 (1878). — Snellen, Vlinders Nederl., 2, p. 30 (1882). — Leech, Brit. Pyral., p. 18, t. 15, f. 4 (1886). — Meyrick, Handb. Brit. Lep., p. 422 (1895). — Hampson, Trans. Ent. Soc. London, p. 236 (1897). — Barrett, Lep. Brit. Isl., 9, p. 327, t. 411, f. 4a, 4b (larva) (1904). — Rebel in Spuler, Schm. Eur., 2, p. 225, t. 82, f. 44 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 41, f. 41—43 (1912) (*Dipleurina*). — Wahlgren, Svensk Insektafauna, 10 (2), p. 511, 53, t. 3, f. 61 (1915). — Müller-Rutz, Mitt. Schweiz. Ent. Ges., 13, p. 432, f. III b, (genit.), (1925). — Löfqvist & Karvonen, Not. Ent., 7, p. 40 n. 88 (1927). — Meyrick, Revis. Handb. Brit. Lep., p. 439 (1928). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 288, f. 506 (1932). — Eckstein, Schm. Deutschl., 5, p. 64 (1933). *crataegalis* Guenée, Delt. & Pyral., p. 428 (1854).

Biologie: Heyden, Ent. Zeit. Stettin, 26, p. 376 (1865). — Buckler, Larvae Brit. Butt., 9, t. 155, f. 5 (1901) (*Eudorea*). — Sorhagen, Allg. Zeitschr. Ent., 6, p. 243 (1901).

x-signata Filipjev, Jahrb. Martj. Staatsmus., 5 (1), Minussinsk p. 28 (1927).

sachalinensis Matsumura, Journ. Coll. Agric. Hokkaido Imp. Univ., 15, p. 185 (1925).

frequentella Stainton, Man. Brit. Butt. Moths, 2, England, p. 162 (1859). — Herrich-Schaeffer, Syst. Schweden, Bearb. Schm. Eur., 6, p. 144 (1855) (*Eudorea*). — Heinemann, Kleinschm., 1, H. 2, p. 40 (1865). — Snellen, Vlinders Nederl. 2, p. 31 (1882). — Wallengren, Öfvers. K. Vet.-Akad. Förh., 28 (8), p. 976 (1871). — Meyrick, Handb. Brit. Lep., p. 422 (1895). — Hampson, Trans. Ent. Soc. London, p. 236 (1897). — Rebel in Spuler, Schm. Eur., 2, p. 225 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 42, f. 54—56 (1912) (*Eudoria*). — Wahlgren, Svensk Insektafauna, 10 (2), p. 52, 53 (1915). — Müller-Rutz, Mitt. Schweiz. Ent. Ges., 13, p. 432, f. III a (genit.), (1925). — Löfqvist & Karvonen, Not. Ent., 7, p. 40 n. 89 (1927). — Meyrick, Revis. Handb. Brit. Lep., p. 439 (1928). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 288 (1932). — Eckstein, Schm. Deutschl., 5, p. 64 (1933). *crataegella* Herrich-Schaeffer, Syst. Bearb. Schmett. Eur., 4, p. 48, (*Tin.*), t. 16, f. 114 (1848). (*Eudorea*). — Caradja, Deutsche Ent. Zeitschr. Iris, 30, p. 23 (1917). Sachalin

- mercurea* Haworth, Lep. Brit., p. 499 (1812) (*Scopaea*). — Curtis, Brit. Ent., 4, t. 170, (1828) (*Eudorea*).
- mercurella* Stephens, Ill. Brit. Ent. Haust., 4, p. 302 (1834). — Wood, Index Entom., p. 209, t. 46, f. 1499 (1838). — Leech, Brit. Pyral., p. 18, t. 15, f. 3 (1886). — Barrett, Lep. Brit. Isl., p. 329, t. 411, f. 5, 5a, b (1904).
- crataegalis* Duponchel, Hist. Nat. Lép., 8 (2), p. 292, t. 229, f. 7 (1831).
- var. *portlandica* Stainton (Dale i. l.), Syst. Englische Küsten
Cat. Brit. Tin. et Pteroph., p. 3 (1849). — Curtis, Ann. Mag. Nat. Hist. Nat. (s. 2), 5, p. 115 (1851). — Stainton, Ent. Mon. Mag., 21, p. 275 (1885). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 53, nr. 976 b (1901). — Barrett, Lep. Brit. Isl., 9, p. 330, t. 411, f. 5c—e (1904). — Lhomme, Cat. Lep. Fr. Belg., 2 (1), p. 112 (1935).
- phaecoleuca* Stainton (nec Zeller), Man. Brit. Butt. Moths, 2, p. 162 (1859). — Knaggs, Ent. Mon. Mag., 5, p. 293, t. 1, f. 10 (1869). — Leech, Brit. Pyral., p. 19, t. 15, f. 5 (1886).
- ab. *concinella* Curtis, Ann. Mag. Nat. Hist., (s. 2) 5, p. 115 (1851). — Staudinger & Rebel, Cat. Lep. Pal. Faun., 2, p. 53, nr. 976 b (1911). — Lhomme, Cat. Lep. Fr. Belg., 2 (1), p. 113 (1935). England, Belgien
- Biologie: Porritt, Ent. Mon. Mag., 22, p. 260 (1886). — Buckler, Larvae Brit. Butt., 9, p. 190, t. 155, f. 6 (1901) (*Eudorea*). — Rebel in Spuler, Schm. Eur., 2, Nachtr.-Taf. 9, f. 59 (1906).
- berytella** Rebel, Verh. z. b. Ges. Wien, 61, p. 148 (1911). Syrien
- acuminatella** Stainton, Ann. Mag. Nat. Hist., (s. 3) 3, p. 210 (1859). Madeira
- luteusalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 23 (♀) (1907). Azoren
- gilvescens** Rebel, Ann. Naturh. Hofmus. Wien, 31, p. 49 (♂ ♀) (1917). Canaren
- angustea** Stephens, Ill. Brit. Ent. Haust., 4, p. 302 (1834). — Wood, Index Entom., p. 209, t. 46, f. 1450 (1839). — Stainton, Man. Brit. Butt. Moths, 2, p. 163 (1859) (*Eudorea*). — Knaggs, Ent. Mon. Mag., 5, t. 1, f. 19 (1869). — Borce, Faune Ent. Fr. Lep., 6, p. 212 (1878). — Leech, Brit. Pyral., p. 30, t. 15, f. 7 (1886). — Hampson, Trans. Ent. Soc. Lond., p. 236 (1897). — Meyrick, Handb. Brit. Lep., p. 421 (1895). — Rebel, Ann. Naturh. Hofmus. Wien, 13, p. 374 (1898). — Barrett, Lep. Brit. Isl., 9, p. 334, t. 412, f. 2, 2a, b, c (larva) (1904). — Rebel in Spuler, Schm. Eur., 2, p. 225 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 43, f. 59, 60 (1912) (*Eudoria*). — Longstaff, Butt.-Hunt. Many Lands, Südeuropa (excl. Südrußland), England, Shetland Ins., Kleinasien, Madeira, Canaren, Azoren, Mauretanien

- p. 168, 172 (1912). — Meyrick, Revis. Handb. Brit. Lep., p. 438 (1928). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 288 (1932).
- coarctata* Zeller, Linn. Entom., 1, p. 308, f. 14 (1846) (*Eudorea*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 46, t. 17, f. 121 (1848) (*Eudorea*). — De la Harpe, Mitt. Schweiz. Ent. Ges., 1 (4), p. 88 (1863) (*Eudorea*).
- coarctalis* Guenée, Delt. & Pyral., p. 430 (1854). — Millière, Icon. Chen. Lép., 3, p. 88, t. 109, f. 12—16 (1871).
- amissella* Millière, Icon. Chen. Lép., 1, p. 401, t. 50, f. 1 (1864).
- var. *napolitana* Millière, Icon. Chen. Lép., Súditalien 3, p. 90, t. 109, f. 17 (1871).
- Biologie: Porritt, Ent. Mon. Mag., 22, p. 209 (1886). — Buckler, Larvae Brit. Butt., 9, p. 196, t. 155, f. 8 (1901) (*Eudorea*).
- pallida** Stephens, Ill. Brit. Ent. Haust., 4, p. 300 (1834). — Zeller, Linn. Entom., 1, p. 314 (1846) (*Eudorea*). — Wood, Index, Entom., p. 208, t. 46, f. 1445 (1838). — Stainton, Man. Brit. Butt. Moths, 2, p. 163 (1859). — De la Harpe, Mitth. Schweiz. Ent. Ges., 1 (4), p. 88 (1863) (*Eudorea*). — Heinemann, Kleinschm., 1, H. 2, p. 43 (1865). — Knaggs, Ent. Mon. Mag., 5, t. 1, f. 1 (1869). — Berce, Faune Ent. Fr. Lép., 6, p. 214 (1878). — Snellen, Vlinders Nederl., 2 (1), p. 33 (1882). — Leech, Brit. Pyral., p. 21, t. 15, f. 9 (1886). — Meyrick, Handb. Brit. Lep., p. 423 (1895). — Hampson, Trans. Ent. Soc. Lond., p. 235 (1897). — Barrett, Lep. Brit. Isl., 9, p. 339, t. 412, f. 4, 4a, b (1904). — Rebel in Spuler, Schm. Eur., 2, p. 225 (1906). — Chapman, Trans. Ent. Soc. Lond. 1911, p. 507, t. 40, f. 36—38, t. 44, f. 67 (1912) (*Witlesia*). — Wahlgren, Svensk Insektauna, 10 (2), p. 51, 53 (1915). — Meyrick, Revis. Handb. Brit. Lep., p. 440 (1928). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 284 (1932). — Eckstein, Schm. Deutschl., 5, p. 64 (1933).
- oertzeniella* Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 49, (*Tin.*), t. 14, f. 97 (1848) (*Eudorea*).
- forma *denigrata* Costantini, Neue Beitr. Syst. Insektenk., 2, nr. 12, p. 105 (1923) (*Whitesia pallida* fa.). — Turati, Atti Soc. Ital. Sci. Nat., 62, p. 65, t. B, f. 27 (1923) (*Whitesia*).
- obsoleta** Staudinger, Hor. Soc. Ent. Ross., 15, p. 163 (♂, ♀) (1880). England, Orkney Ins., Schweden, Lövland, Estland, Mittel-Europa (Belgien, Frankreich, Schweiz, N. u. S. Italien), N. u. O. Spanien, Süd-Rußland, Lenkoran
- Syrien (Typus: Kerasdere)

Species Incertae sedis.

- aequipennalis** Warren, Nov. Zool. Tring, 12, p. 444 (1905). Azoren (Typus: Terceira)

- coecimaculalis** Warren, Nov. Zool. Tring, 12, p. 445 (♂, ♀) (1905). Azoren
(Typus: Calheta)
- interlinealis** Warren, Nov. Zool. Tring, 12, p. 446 (♀) (1905). Azoren
(Typus: Caldeira)
- ab. *pallidimarginalis* Warren, Nov. Zool. Tring, 12, p. 446 (1905). Azoren

b) Orientalische Region.

- metaleucalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 23 (♂) (1907). Westchina
- promiscua** Wileman & South, Entom., 52, p. 268 (♀) (1919). Formosa
- pulveralis** Snellen, Trans. Ent. Soc. Lond., (1890). — Hampson, Fauna Brit. India Moths, 4, p. 242 (1896). Sikkim, Darjeeling
- medinella** Snellen, Trans. Ent. Soc. Lond., (1890). — Hampson, Fauna Brit. India Moths, 4, p. 242 (1896). Sikkim
- submedinella** Caradja, Acad. Rom. Mem. Sect. Sti-int., (s. 3) 4, mem. 8, p. 407 (1927). Westchina
- albifusalis** Hampson, Ann. Mag. Nat. Hist. (s. 7) 19, p. 24 (♀) (1907); Journ. Bombay Nat. Hist. Soc., 18, p. 575 (♀) (1908). Ceylon
- olivaris** Hampson, Ill. Typ. Het. Brit. Mus., 8, p. 131, t. 154, f. 11 (1891); Fauna Brit. India Moths, 4, p. 243 (♀) (1896). Nilgiris
- crocealis** Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 212 (♀) (1903). Nilgiris
- delicatalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1500 (1865). — Hampson, Fauna Brit. India Moths, 4, p. 243 (1896); id., Trans. Ent. Soc. London, p. 237 n. 126 (1897). Sikkim, Nilgiris
- rufostigma** Hampson, Ill. Typ. Het. Brit. Mus., 8, p. 131, t. 154, f. 4 (1891); Fauna Brit. India Moths, 4, p. 243 (1896); id., Trans. Ent. Soc. Lond., p. 237 (1897). Nilgiris
- medlorufalis** Hampson, Fauna Brit. India Moths, 4, p. 243 (1896); Trans. Ent. Soc. Lond., p. 237 (1897). Dharmasala, Sikkim
- phaealis** Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 213 (1903). Bombay, Belgaum
- polialis** Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 213 (♀) (1903). Assam
- murifficalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 826 (1859) (Ceylon). — Hampson, Ill. Typ. Het. Brit. Mus., 9, p. 182, t. 174, f. 10 (1890); Fauna Brit. India Moths, 4, p. 244, f. 147 (♂) (1890); Trans. Ent. Soc. Lond., p. 229, fig. (♂) 237 (1897); in Forbes, Nat. Hist. Sokotra, p. 335, t. 20, f. 25 (1903). N.W.-Himalaya, Ceylon, Westchina, Sokotra
- truncicolella* Butler (nec Stainton, 1849), Ill. Typ. Het. Brit. Mus., 7, p. 20 (1899).

- congestalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 826 n. 34 (♀) (1859). — Hampson, Ill. Typ. Lep. Het. Brit. Mus., 9, p. 182, t. 174, f. 2 (1893). Ceylon, W.-Himalaya, West- u. Zentralchina
- murificalis** (part.) Hampson, Trans. Ent. Soc. London, p. 237 (1897).
- straminealis** Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 213 (♂) (1903). Simla
- vinotinctalis** Hampson, Fauna Brit. India Moths, 4, p. 244 (♂) (1896); id., Trans. Ent. Soc. Lond., p. 237 n. 130 (1897). Nilgiris, West- u. Zentral-China
- canicostalis** Hampson, Fauna Brit. India Moths, 4, p. 244 (1896); id., Trans. Ent. Soc. Lond., p. 237 n. 131 (1897). Nilgiris
- ochrotalis** Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 213 (♂) (1903). Nilgiris
- barbipennis** Hampson, Trans. Ent. Soc. Lond., p. 238 (1897) (Gunong Ijau). Malakka

Species incertae sedis.

- fulvosignalis** Snellen, Tijds. v. Ent., 23, p. 204 (♀) (1880); v. 26, t. 6, f. 10 (1884). Celebes (Typus: Makassar)
- nugalis** Snellen, Tijds. v. Ent., 23, p. 205 (1880); v. 26, t. 6, f. 11 (1884). Celebes (Typus: Maros)
- stötzneri** Caradja, Acad. Rom. Mem. Sect. Stiint., (s. 3), 4, mem. 8, p. 407 (1927). Westchina (Szetschwan)
- xanthomelas** De Joannis, Ann. Soc. Ent. Fr., 98, p. 655, t. 3, f. 13 (1929). Tonkin, Kwangtung

c) Aethiopische und madagassische Region.

- nigritalis** Walker in Mellis's St. Helena, p. 190 (1875). — Wollaston, Ann. Mag. Nat. Hist., (s. 5) 3, p. 333 (1879). — Hampson, Trans. Ent. Soc. London, p. 237 n. 115 (1897). St. Helena
- lucidalis** Walker in Mellis's St. Helena, p. 190 (1875). — Wollaston, Ann. Mag. Nat. Hist., (s. 5) 3, p. 334 (1879). St. Helena
- lusiadalis** Hampson, Trans. Ent. Soc. London, p. 237 n. 116 (1897).
- similis** Wollaston, Ann. Mag. Nat. Hist., (s. 5), 3, p. 333 (1879). — Hampson, Trans. Ent. Soc. Lond., p. 237 n. 117 (1897). St. Helena
- helenensis** Wollaston, Ann. Mag. Nat. Hist., (s. 5) 3, p. 334 (1879). — Hampson, Trans. Ent. Soc. Lond., p. 237 (1897). St. Helena
- scintillulalis** Wollaston, Ann. Mag. Nat. Hist., (s. 5) 3, p. 335 (1879). — Hampson, Trans. Ent. Soc. Lond., p. 237 (1897). St. Helena
- transversalis** Wollaston, Ann. Mag. Nat. Hist., (s. 5) 3, p. 336 (1879). — Hampson, Trans. Ent. Soc. Lond., p. 237 (1897). St. Helena

- contemptalis** Walker, Cat. Lep. Het. Brit. Mus, 34, Südafrika
p. 1499 (♂) (1865). — Hampson, Trans. Ent.
Soc. Lond., p. 237 (1897).
- dulcis** De Joannis, Soc. Ent. Fr. Livre Centen., p. Mauritius
45, t. 23, f. 21 (♀) (1932).
- resinodes** De Joannis, Soc. Ent. Fr. Livre Centen., Réunion
p. 451, t. 23, f. 20 (♂, ♀) (1932).
- nolalis** Felder & Rogenhofer, Reise Novara, (Het. 2), Südafrika
p. 17, t. 134, f. 41 (1874). — Hampson,
Trans. Ent. Soc. London, p. 237 (1897).
- musificalis** Walker, vide sub b) Orientalische Re-
gion, p. 30.
- phaeopalpia** Hampson, Ann. Mag. Nat. Hist., (s. Natal
8) 20, p. 282 (♂) (1917).

Species Incertae sedis.

- allicola** Meyrick, Exot. Microlep., 4, p. 546 (♂) Uganda
(1935). (Typus:
Ruwenzori)
- benigna** Meyrick, Trans. Ent. Soc. Lond., p. 366 Mauritius
(1910).
- idiogama** Meyrick, Exot. Microlep., v. 4, p. 545 Uganda
(♂, ♀) (1935). (Typus:
Ruwenzori)
- thyrophanta** Meyrick, Exot. Microlep. 4, p. 245 Uganda
(♂) (1932). (Typus:
Ruwenzori)

d) Nearktische Region.

- centuriella** [Schiff. & Den.] vide Paläarkt. Region,
p. —.
- rectilinea** Zeller, Verh. z. b. Ges. Wien, 24, p. 427, Vancouver
t. 12, f. 3 (1874). — Hampson, Trans. Ins.,
Ent. Soc. Lond., p. 234 (1897). — Dyar, Kalifornien
Proc. U. S. Nat. Mus., 27, p. 918 (1904);
Journ. N. York Ent. Soc., 14, p. 100 (1906).
refugalis Hult, Trans. Amer. Ent. Soc.,
13, p. 148 (1886).
- commortalis** Dyar, Insec. Insc. Menst., 9, p. 67 Brit.
(♂) (1921); Proc. U. S. Nat. Mus., 74, art. Columbien,
24, p. 1 (1929). Victoria,
Canada
- expallidalis** Dyar, Journ. N. York Ent. Soc., 14, Nevada
p. 101 (1906).
- rigidalis** Barnes & Mac Dunnough, Contr. Hist. Arizona
Lep. N. Amer., 1 (5), p. 35, t. 3, f. 7
(1912).
- denigata** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, Arizona,
p. 1 (♂) (1929). Kalifornien
- normalis** Dyar, Ent. News Philad., v. 15, p. 71 (♀) Neu-
(1904); Journ. N. York Ent. Soc., 14, p. Mexiko
102 (1906). (Typus:
Blulat)

- delphusa** Druce, Biol. Centr.-Amer. Het., 2, p. 279, t. 64, f. 1 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 234 (1897). — Dyar, Journ. N. York Ent. Soc., 14, p. 102 (1906); Proc. U. S. Nat. Mus., 74, art. 24, p. 2 (1929).
 Mexiko, Guatemala, Arizona, Colorado
- smithi** Druce, Biol. Centr.-Amer. Het., 2, p. 279, t. 64, f. 3 (1896) (Mexiko). — Hampson, Trans. Ent. Soc. Lond., p. 234 (1897).
- sabura** Druce, Biol. Centr.-Amer. Het., 2, p. 279, t. 64, f. 4 (1896) (Mexiko).
- flexuosa** Dyar, Proc. U. S. Nat. Mus., 54, p. 370 (♀) (1918) (Mexiko).
- penumbralis** Dyar, Journ. N. York Ent. Soc., 14, p. 103 (1906). — Forbes, Lep. N. York, p. 533, f. 357 (1923).
 Ontario, N. Hampshire, N. Jersey, N. York, Pennsylvanien
- cinereomedia** Dyar, Ent. News Philad., 15, p. 72 (♂, ♀) (1904); Journ. N. York Ent. Soc., 14, p. 103 (1906). — Forbes, Lep. N. York, p. 533, f. 359 (1923).
 Pennsylvanien, N. York (Typus: New Brighton)
- schwarzalis** Dyar, Journ. N. York Ent. Soc., 14, p. 103 (♂) (1906).
 Arizona (Typus: Santa Rita Mts.)
- strigalis** Dyar, Journ. N. York Ent. Soc., 14, p. 104 (1906). — Forbes, Lep. N. York, p. 533 (1923).
 Ontario (Typus: Grinsby), N. Hampshire bis Georgia, N. York, Pennsylvanien, Maryland
- nominatalis** Hulst, Trans. Amer. Ent. Soc., 13, p. 148 (1886) (Vancouver Ins.). — Hampson, Trans. Ent. Soc. Lond., p. 233 (1897). — Dyar, Proc. U. S. Nat. Mus., 27, p. 918 (1904); 74, art. 24, p. 2 (1929).
 Brit. Colombien, Utah, Colorado, Arizona, N. Hampshire, Mexiko, Guatemala
- spaldingalis** Barnes & Mc Dunnough, Contr. Nat. Hist. Lep. N. Amer., 1 (5), p. 34 (♂, ♀), t. 3, f. 1, 4 (1912).
 (Typus: Eureka (Utah))
- albertalis** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 2 (♂) (1929).
 Brit. Colombien

- tricoloralis** Dyar, Ent. News Philad., 15, p. 72 Brit.
(♂, ♀) (1904); Proc. U. S. Nat. Mus., 27, Colombien,
p. 818 (1904); Journ. N. York Ent. Soc., Washington,
14, p. 105 (1906); Proc. U. S. Nat. Mus., Colorado,
74, art. 24, p. 2 (1929). Kalifornien
rufitinctalis Hampson, Ann. Mag. Nat. (Typus:
Hist., (s. 7) 19, p. 21 (1907). Wellington)
- echo** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, Brit.
p. 3 (♂) (1929). Colombien
- leucophthalma** Dyar, Proc. U. S. Nat. Mus., 74, Brit.
art. 24, p. 3 (♂) (1929). Colombien
- basalis** Walker, Cat. Lep. Het. Brit. Mus., 34, Vereinigte
p. 1497 (♂) (1859) (Nord-Amerika). — Staaten
Hampson, Trans. Ent. Soc. Lond., p. 234 (Maine,
(1897). — Dyar, Journ. N. York Ent. Soc., N. Hamp-
14, p. 105 (1906). — Forbes, Lep. N. York, shire,
p. 583, f. 353, 358 (1923). — Dyar, Proc. N. York,
U. S. Nat. Mus., 74, art. 24, p. 3 (1929). Penn-
biplagiatis Walker, Cat. Lep., Het. Brit. sylvanien,
Mus., 34, p. 1499 (♂) (1865). — Hampson, Iowa,
Trans. Ent. Soc. Lond., p. 234 (1897). Maryland,
libella Grote, Bull. U. S. Geol. Surv., 4, Virginia,
p. 675 (1878). N. Carolina,
bronzalis Barnes et McDunnough, Contr. Florida,
Nat. Hist. Lep. N. Amer., 5 (1), p. 48 Montana,
(1922) (Kalifornien). Arizona,
cervalis McDunnough, Canad. Ent., 59, Kalifornien),
p. 267 (1927). (Brit. Columbien). Kanada,
Mexiko
- ab. *palloralis* Dyar, Journ. N. York Ent. Arizona,
Soc., 14, p. 106 (1906). Neu-
Mexiko,
Colorado
- ab. *fernaldalis* Dyar, Ent. News Philad., Brit.
15, p. 72 (♂, ♀) (1904); Proc. U. S. Nat. Colombien
Mus., 27, p. 818 (1904). — Dyar, Journ. N. (Typus:
York Ent. Soc., 14, p. 106 (1906). Kaslo,
Brit. Col.)
- ab. *obispalis* Dyar, Journ. N. York Ent. Kalifornien
Soc., 14, p. 106 (1906). (Typus:
San Louis
Obispo)
- jonesalis** Dyar, Insec. Insc. Menst., 3, p. 88 (♀) Bermuda
(1915). Ins.
- truncatalis** McDunnough, Canad. Ent., 54, p. 36 Kanada
(1922). — Forbes, Lep. N. York, p. 584 (Ontario)
(1923).
- pacificalis** Dyar, Insec. Insc. Menst., 9, p. 66 (♂, Brit.
♀) (1921) (*S. basalis* subsp.); Proc. U. S. Colombien,
Nat. Mus., 74, art. 24, p. 4 (1929). Alaska

- alaskalis* Barnes et Benjamin, Contr. Nat. Hist. Lep. N. Amer., 5 (1), p. 49 (♂) (1922).
- torniplagalis* Dyar, Journ. N. York Ent. Soc., 12, p. 105 (1904); 14, p. 106 (1906). Washington, Colorado
- alialis* Barnes et Mc Dunnough, Contr. Hist. Nat. Lep. N. Amer., 1 (5), p. 35, t. 3, f. 6 (1912). Neu-Mexiko (Typus: St. Wingate)
- lugubralis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1498 (♂) (1865). (Hudson - Bai). — Hampson, Trans. Ent. Soc. Lond., p. 233 (1897). — Dyar, Journ. N. York Ent. Soc., 14, p. 104 (1906). — Forbes, Lep. N. York, p. 583, f. 360 (1923). — Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 5 (1929). Kanada, Alaska, Vereinigte Staaten (N. Hampshire, Minnesota, Washington, Arizona, Kalifornien)
- phycitinalis* Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 4 (♂) (1929). Brit. Colombien

Species incertae sedis.

- atropicta* Hampson (Zeller MS.), Trans. Ent. Soc. Lond., p. 233 (♂) (1897). — Dyar, Journ. N. York Ent. Soc., 14, p. 107 (1906). Vereinigte Staaten

e) Neotropische Region.

- chordactis* Meyrick, Trans. Ent. Soc. Lond., p. 272 (♂) (1887). — Hampson, Trans. Ent. Soc. Lond., p. 233 (1897). Patagonien (Typus: Magellanstraße)
- strigigramma* Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 281 (♂) (1917). Peru
- ragonoti* Butler, Trans. Ent. Soc. Lond., p. 59 (1883) (*Mella*; ad *Phycidas* adfert.). Peru, Chile, Juan Fernandez (Typus: Valparaiso)
- glauculalis* Hampson, Trans. Ent. Soc. Lond., p. 233 (♂) (1897). — Enderlein, Svenska Akad. Handl., 48 (3), p. 40, f. 3 (1912). Falklands Ins.
- tersella* Zeller, Ent. Zeit. Stettin, 33, p. 476, t. 2, f. 9 (1872). — Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 5 (1929). Colombien (Bogotá)
- atricuprea* Hampson, Ann. Mag. Nat. Hist. (s. 8) 20, p. 281 (♂) (1917). Colombien, Jamaica
- ceramica* Warren, Ann. Mag. Nat. Hist. (s. 6) 8, p. 67 (1891) (*Lissophanes*). Peru

- dispersa** Butler, Trans. Ent. Soc. Lond., p. 58 (1883). — Hampson, Trans. Ent. Soc. London, p. 233 n. 77 (1897). Chile
(Typus: sine patria)
- pusilla** Dyar, Proc. U. S. Nat. Mus., 47, p. 320 (1914); 74, art. 24, p. 5 (1929). Panama
- subtersa** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 5 (♂) (1929). Guatemala, Costa Rica, Mexiko, West-Indien, Brasilien
- extincta** Dyar, Insec. Insc. Menst., 9, p. 67 (♂) (1921); Proc. U. S. Nat. Mus., 74, art. 24, p. 5 (1929). Mexiko
(Typus: Mexico City)
- albipunctata** Druce, Biol. Centr.-Amer. Het., 2, p. 563, t. 101, f. 2 (1899). — Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 8, (1929). Costa Rica, Guatemala
- tricolor** Zeller, Ent. Zeitg. Stettin, 33, p. 473, t. 2, f. 10 (1872). — Hampson, Trans. Ent. Soc. London, p. 233 n. 78 (1897). — Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 6 (1929). Colombien (Bogotá)
- molliculella** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 6 (♂) (1929). Mexiko
- stereostigma** Dyar, Proc. U. S. Nat. Mus., 54, p. 369, (♀) (1918); 74, art. 24, p. 6 (1929). Mexiko, Guatemala
- multifacies** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 6 (♀) (1929). Guatemala, Mexiko
- inexoptata** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 7 (♂) (1929). Mexiko
- ycarda** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 7 (♀) (1929). Guatemala, Mexiko
- biscutella** Zeller, Ent. Zeitg. Stettin, 33, p. 474, t. 2, f. 8 (1872). — Hampson, Trans. Ent. Soc. Lond., p. 233 (1897). — Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 7 (1929). Colombien, Costa Rica, Guatemala, Mexiko
- albifrons* Druce, Biol. Centr.-Amer. Het., 2, p. 278, t. 64, f. 2 (1896), nec Zeller, Stett. Ent. Zeit., 33, p. 476, nota 2 (1872) (*Scoparia* spec.).
- anagantis** Dyar, Proc. U. S. Nat. Mus., 54, p. 376 (1918); 74, art. 24, p. 7 (1929). Mexiko
- crassuliscula** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 8 (♂) (1929). Brasilien, Franz. Guayana
- imparilis** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 8 (♂) (1929). Jamaica
- vinasalis** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 8 (♂) (1929). Costa Rica

- excursalis** Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 8 (♂) (1929). Costa Rica
ulmaya Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 9 (♂) (1929). Guadeloupe
longipennis Zeller, Ent. Zeitg. Stettin, 33, p. 479, t. 2, f. 11 (1872). — Hampson, Trans. Ent. Soc. Lond., p. 233 (1897). — Dyar, Proc. U. S. Nat. Mus., 74, art. 24, p. 9 (1929). Colombien (Bogotá), Guatemala, Costa Rica, Mexiko

delphusa Druce (vide Nearktische Region, p. 33).

nominatais Hulst (vide Nearktische Region, p. 33).

Species incertae sedis.

- anadonta** Dyar, Proc. U. S. Nat. Mus., 54, p. 369 (♂) (1918). Mexiko
antarcticalis Staudinger, Erg. Hamb. Magalh. Sammelr., 2, Lep., p. 106 (♂) (1898). Feuerland
cyclophora Dyar, Proc. U. S. Nat. Mus., 54, p. 370 (1918). Mexiko

f) Hawaiische Region.

- tetranesa** Meyrick, Fauna Hawai., 1, p. 247, 249, t. 7, f. 3 (♀) (1899). Hawai (Typus: Haleakale)
rhombias Meyrick, Fauna Hawai., 1, p. 249 (♂, ♀) (1899). Hawai (Typus: Kaholua-mano)
amphicypellia Meyrick, Fauna Hawai., 1, p. 250, t. 7, f. 4 (♂) (1899). Hawai (Typus: Mth. Waimea)
oxythyma Meyrick, Fauna Hawai., 1, p. 250, (♂, ♀) (1899). Hawai (Typus: Mts. Waimea)
balanopsis Meyrick, Fauna Hawai., 1, p. 250, t. 7, f. 5 (♂, ♀) (1899). Hawai (Typus: Kona)
passalota Meyrick, Fauna Hawai., 1, p. 251 (♂, ♀) (1899). Hawai (Typus: Haleakale)
orthoria Meyrick, Fauna Hawai., 1, p. 251, t. 7, f. 6 (♂, ♀) (1899). Hawai (Typus: Olaa)
gonodecta Meyrick, Fauna Hawai., 3, p. 362 (1904) (n. nov.) Hawai (Typus: Mts. Waimea)
mesoleuca Meyrick, Fauna Hawai., 1, p. 252 (♂, ♀) (1899) (nec Meyrick, Trans. Ent. Soc. Lond., p. 237 (1888)).

- mesoleuca** Meyrick, Trans. Ent. Soc. Lond., p. 237 (1888); Fauna Hawai., 3, p. 363 (♂, ♀) (1904). Hawai (Typus: Oahu)
- miantis** Meyrick, Fauna Hawai., 1, p. 252 (♂) (1899). Hawai (Typus: Molokai)
- antimacha** Meyrick, Fauna Hawai., 1, p. 252 (♀) (1899). Hawai (Typus: Lahaina)
- tyraula** Meyrick, Fauna Hawai., 1, p. 253, t. 7, f. 7 (♀) (1899). Hawai (Typus: Kaholua-mano)
- pentaspila** Meyrick, Fauna Hawai., 1, p. 253 (♂, ♀) (1899). Hawai (Typus: Kaala)
- pachysema** Meyrick, Trans. Ent. Soc. Lond., p. 236 (♀) (1888) (*Xeroscopa*); Fauna Hawai., 1, p. 254 (1899). Hawai (Typus: Haleakale)
- thalamias** Meyrick, Fauna Hawai., 1, p. 254 (♂) (1899). Hawai (Typus: Waimea)
- loxocentra** Meyrick, Fauna Hawai., 1, p. 255, t. 7, f. 9 (♀) (1899). Hawai (Typus: Kona)
- halirrhoa** Meyrick, Fauna Hawai., 1, p. 255 (♂, ♀) (1899). Hawai (Typus: Olaa)
- aetias** Meyrick, Fauna Hawai., 1, p. 256, t. 7, f. 10 (♂, ♀) (1899). Hawai (Typus: Olaa)
- melanocephala** Meyrick, Fauna Hawai., 1, p. 256 (♂) (1899). Hawai (Typus: Kona)
- formosa** Butler, Ann. Mag. Nat. Hist., (s. 5) 7, p. 331 (♂) (1881) (*jucunda* var.). — Meyrick, Trans. Ent. Soc. Lond., p. 237 (1888) (*Xeroscopa*); Fauna Hawai., 1, p. 257 (♂, ♀) (1899). Hawai
- struthias** Meyrick, Fauna Hawai., 1, p. 257, f. 7, f. 11 (♂) (1899). Hawai (Typus: Mts. Waimea)
- triacma** Meyrick, Fauna Hawai., 1, p. 258, t. 7, f. 12 (♂, ♀) (1899). Hawai (Typus: Mts. Waimea)

- jucunda** Butler, Ann. Mag. Nat. Hist., (s. 5) 7, p. 331 (♀) (1881). — Meyrick, Trans. Ent. Soc. Lond., p. 238 (♀) (1888) (*Xeroscopa*); Fauna Hawai., 1, p. 258 (♂, ♀) (1899). Hawai
- crycerodes** Meyrick, Fauna Hawai., 1, p. 259, t. 7, f. 13 (♂, ♀) (1899). Hawai
(Typus:
Mts.
Waimea)
- crataca** Meyrick, Fauna Hawai., 1, p. 259 (♂, ♀) (1899). Hawai
(Typus:
Olaa)
- catactis** Meyrick, Fauna Hawai., 3, p. 363 (♂, ♀) (1904). Hawai
(Typus:
Oahu)
- ombrodes** Meyrick, Trans. Ent. Soc. Lond., p. 234 (♀) (1888) (*Xeroscopa*); Fauna Hawai., 1, p. 260 (pars), f. 7, f. 14 (♂, ♀) (1899); 3, p. 364 (1904). Hawai
- fragosa** Meyrick, Trans. Proc. N. Zeal. Ins., 42, p. 91 (♀) (1909). Sonntags-
Ins.
(Denham-
Bai)
- siderina** Meyrick, Fauna Hawai., 1, p. 260, t. 7, f. 15 (♂, ♀) (1899). Hawai
(Typus:
Kauai)
- parachlora** Meyrick, Fauna Hawai., 1, p. 260, t. 7, f. 16 (♀) (1899). Hawai
(Typus:
Olaa)
- ianthes** Meyrick, Fauna Hawai., 1, p. 261, t. 7, f. 17 (1899); 3, p. 364 (1904). Hawai
(Typus:
Kauai)
- gymnopis** Meyrick, Ent. Mon. Mag., (s. 2) 15 (40), p. 131 (1904). Hawai
- marmarias** Meyrick, Fauna Hawai., 1, p. 261 (♂, ♀) (1899). Hawai
(Typus:
Olaa & Hilo)
- aeolias** Meyrick, Fauna Hawai., 1, p. 262 (♂) (1899). Hawai
(Typus:
Oahu)
- nectarias** Meyrick, Fauna Hawai., 1, p. 262, t. 7, f. 18 (♀) (1899). Hawai
(Typus:
Hilo & Lanai)
- nectarioides** Swezey, Proc. Hawai Ent. Soc., 2, p. 273 (♂, ♀) (1912). Hawai
(Typus:
Molokai)
- bucolica** Meyrick, Fauna Hawai., 1, p. 263 (♂) (1899). Hawai
- isopiaca** Meyrick, Ent. Mon. Mag., (s. 2) 15 (40), p. 132 (1904). Hawai
(Typus:
Kona)

- pyrseutis** Meyrick, Fauna Hawai., 1, p. 263 (1899); Hawai
3, p. 364 (1904). (Typus:
Hilo &
Kilauea)
- erebochalca** Meyrick, Fauna Hawai., 1, p. 264 Hawai
(♂, ♀) (1899). (Typus:
Maui, Hilo
& Molokai)
- oenopsis** Meyrick, Fauna Hawai., 1, p. 264, t. 7, Hawai
f. 19 (♂, ♀) (1899). (Typus:
Molokai)
- nyctombra** Meyrick, Fauna Hawai., 1, p. 264 (♀) Hawai
(1899). (Typus:
Halakala)
- clonodes** Meyrick, Fauna Hawai., 1, p. 265, t. 7, Hawai
f. 20 (♂, ♀) (1899). (Typus:
Kauai &
Oahu)
- thyellopis** Meyrick, Fauna Hawai., 1, p. 265 (♂, ♀) Hawai
(1899). (Typus:
Hilo, Olan)
- melichlora** Meyrick, Fauna Hawai., 1, p. 266, t. 7, Hawai
f. 21 (♂, ♀) (1899). (Typus:
Kona)
- zophochlora** Meyrick, Fauna Hawai., 1, p. 266 Hawai
(♂, ♀) (1899). (Typus:
Molokai)
- hawaiensis** Butler, Ann. Mag. Nat. Hist., (s. 5) 7, Hawai
p. 330 (1881). — Meyrick, Trans. Ent. Soc.
Lond., p. 235 (1888) (*Xeroscopia*); id., Fauna
Hawai., 1, p. 267 (1899); 3, p. 364 (1904).
- empeda** Meyrick, Fauna Hawai., 1, p. 267 (♂, ♀) Hawai
(1899). (Typus:
Halakala
& Lanai)
- demodes** Meyrick, Trans. Ent. Soc. Lond., p. 234 Hawai
(♀) (1888) (*Xeroscopia*); id., Fauna Hawai., 1,
p. 268 (♀) (1899); 3, p. 264 (1904).
- dactylopa** Meyrick, Fauna Hawai., 1, p. 268, t. 7, Hawai
f. 22 (♂, ♀) (1899). (Typus:
Molokai)
- geraea** Meyrick, Fauna Hawai., 1, p. 269, t. 7, Hawai
f. 22 (♂, ♀) (1899). (Typus:
Oahu &
Kauai)
- epimystis** Meyrick, Fauna Hawai., 1, p. 269 (♂) Hawai
(1899).

- lycopodiae** Swezey, Proc. Hawai. Ent. Soc., 2 (3), Hawai
p. 104, t. 3, f. 2 (♂, ♀) (1910).
Biologie: Swezey, l. c., p. 104, 105 (me-
tam.) (1910).
- frigida** Butler, Ann. Mag. Nat. Hist., (s. 5) 7, Hawai
p. 331 (1881). — Hampson, Trans. Ent.
Soc. London, p. 233 (1897). — Meyrick,
Fauna Hawai., 1, p. 270 (♂, ♀) (1899).
- peronettis** Meyrick, Fauna Hawai., 1, p. 270 (♀) Hawai
(1899).
- ischulias** Meyrick, Trans. Ent. Soc. Lond., p. 235 Hawai
(1888) (*Xeroscopa*); id., Fauna Hawai., 1, p.
271, t. 7, f. 24 (♂, ♀) (1899).
- melanopsis** Meyrick, Trans. Soc. Ent. Lond., p. 233 Hawai
(1888) (*Xeroscopa*); id., Fauna Hawai., 1, p.
271 (1899).
- montana* Butler, Trans. Ent. Soc. Lond.,
p. 41 (188) (*S. coarctata* var.?).
- macrophanes** Meyrick, Trans. Ent. Soc. Lond., p. Hawai
231 (♀) (1888); id., Fauna Hawai., 1, p. 271
(♀) (1899).
- meristis** Meyrick, Fauna Hawai., 1, p. 272 (♂, ♀) Hawai
(1899); 3, p. 365 (1904). (Typus:
frigida Meyrick (nec Butler, 1881), Trans. Kona &
Ent. Soc. Lond., p. 231 (1888). Molokai)
- halmaea** Meyrick, Fauna Hawai., 1, p. 272, t. 7, Hawai
f. 25 (1899); 3, p. 365 (1904). (Typus:
Kauai)
- platyscia** Meyrick, Fauna Hawai., 1, p. 273, t. 7, Hawai
f. 26 (♂, ♀) (1899). (Typus:
Kauai,
Molokai)
- religiosa** Meyrick, Fauna Hawai., 3, p. 365 (♂, ♀) Hawai
(1904). (Typus:
Oahu)
- probolaea** Meyrick, Fauna Hawai., 1, p. 273, t. 7, Hawai
f. 27 (♂, ♀) (1899). (Typus:
Kauai)
- omilehlopls** Meyrick, Fauna Hawai., 1, p. 274, t. Hawai
7, f. 28 (♂, ♀) (1899). (Typus:
Kauai, Olaa
& Hilo)
- venosa** Butler, Ann. Mag. Nat. Hist., (s. 5) 7, Hawai
p. 332 (1881). — Meyrick, Trans. Ent. Soc.
Lond., p. 233 (188?) (*Xeroscopa*); id., Fauna
Hawai., 1, p. 274 (♂, ♀) (1899) (*Sco-
paria*).

g) Austrofasmanische, melanesische und polynesische Subregion.

- anthracas** Meyrick, Trans. Proc. N. Zeal. Inst., Tasmanien
17, p. 74 (♂, ♀) (1884). — Hampson,
Trans. Ent. Soc. Lond., p. 233 (1897).

- chiasta** Meyrick, Trans. Proc. N. Zeal. Ins., 17, N.S.Wales,
p. 74 (♂) (1885). — Hampson, Trans. Ent. Victoria
Soc. Lond., p. 233 n. 65 (1897).
- eumeles** Meyrick, Trans. Proc. N. Zeal. Inst., 17, N.S.Wales
p. 75 (♂) (1885). — Hampson, Trans.
Ent. Soc. Lond., p. 232 (1897).
- platymera** Lower, Trans. Roy. Soc. S. Austral., Tasmanien
29, p. 107 (♂) (1905).
- protorthra** Meyrick, Trans. Ent. Soc. Lond., p. 450 N.S.Wales
(♂, ♀) (1885). — Hampson, Trans. Ent.
Soc. Lond., p. 233 (1897).
- aphrodes** Meyrick, Trans. Proc. N. Zeal. Inst., 17, N.S.Wales,
p. 76 (♀) (1885). — Hampson, Trans. Ent. Queensland
Soc. Lond., p. 232 n. 62 (1897).
- epicryma** Meyrick, Trans. Proc. N. Zeal. Inst., 17, S.Australien
p. 76 (♂) (1885). — Hampson, Trans. Ent.
Soc. Lond., p. 232 (1897).
- exhibitilis** Walker, Cat. Lep. Het. Brit. Mus., 34, N.S.Wales
p. 1500 (♀) (1865). — Meyrick, Trans. Proc.
N. Zeal. Ins., 17, p. 77 (♂, ♀) (1885).
— Hampson, Trans. Ent. Soc. Lond., p. 232
(1897). — Tillyard, Ins. Austral. & N. Zeal.,
p. 431, t. 28, f. 56 (1925).
- charopoea** Turner, Trans. Proc. Roy. Soc. S. Austral., Queensland
32, p. 106 (♂) (1908).
- syntaraeta** Meyrick, Trans. Proc. N. Zeal. Inst., 17, N.S.Wales,
p. 77 (♂, ♀) (1885). — Hampson, Trans. Tasmanien
Ent. Soc. Lond., p. 232 (1897). — Longstaff,
Butt.-Hunt. Many Lands, p. 440 (1912).
- synapta** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Tasmanien
p. 78 (♀) (1885). — Hampson, Trans. Ent.
Soc. Lond., p. 232 (1897).
- homala** Meyrick, Trans. Proc. N. Zeal. Inst., 17, S.Australien
p. 79 (♂, ♀) (1885). — Hampson, Trans.
Ent. Soc. Lond., p. 232 (1897).
- eremitis** Meyrick, Trans. Proc. N. Zeal. Inst., 17, S.Australien
p. 79 (♂, ♀) (1885). — Hampson, Trans.
Ent. Soc. Lond., p. 232 (1897).
- perierga** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Tasmanien
p. 80 (♂, ♀) (1885). — Hampson, Trans.
Ent. Soc. Lond., p. 232 (1897).
- gomphota** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Tasmanien
p. 80 (♂) (1885). — Hampson, Trans.
Ent. Soc. London, p. 232 (1897). — Longstaff,
Butt.-Hunt. Many Lands, p. 440 (1912).
- anaplecta** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Tasmanien
p. 89 (♂, ♀) (1885). — Hampson, Trans.
Ent. Soc. London, p. 231 (1897).
- spelaea** Meyrick, Trans. Proc. N. Zeal. Inst., 17, N.S.Wales,
p. 89 (♂, ♀) (1885). — Hampson, Trans. Victoria,
Ent. Soc. London, p. 231 (1897). Tasmanien,
- pusilla* Rosenstock, Ann. Mag. Nat. Hist., S.Australien
(s. 5) 16, p. 427 (1885).
- acropola** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Tasmanien
p. 101 (♂) (1885). — Hampson, Trans.
Ent. Soc. Lond., p. 230 (1897).
- eleodoralis** Walker, Cat. Lep. Het. Brit. Mus., 18, N.S.Wales,
p. 793 (♀) (1859) (*Scopula*) (Sydney). — Victoria,
Hampson, Trans. Ent. Soc. Lond., p. 230 Tasmanien

- (1897). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 1011 (♂, ♀) (1885).
oxygona Meyrick, Trans. Ent. Soc. Lond., p. 381 Victoria,
 (♂, ♀) (1897). S.Australien,
 Tasmanien,
plagiolis Meyrick, Trans. Ent. Soc. Lond., p. 247 Tasmanien,
 (♀) (1887). — Hampson, Trans. Ent. Soc. Victoria,
 Lond., p. 230 (1897). — Turner, Proc. Linn. N.S.Wales
 N. S. Wales, 56, p. 344 (1931).
ochrophora Turner, Proc. Roy. Soc. Queens-
 land, 27, p. 52 (♀) (1915).
philonephes Meyrick, Trans. Proc. N. Zeal. Inst., Victoria,
 17, p. 110 (♀) (1885). S.Australien

Species incertae sedis.

- anisophragma** Lower, Trans. Roy. Soc. S. Austral., N.S.Wales
 25, p. 67 (♂) (1901).
anthomera Lower, Trans. Roy. Soc. S. Austral., S.Australien,
 20, p. 157 (♂) (1896). N.S.Wales
arcta Lucas, Proc. Roy. Soc. Queensland, 13, p. 84 Queensland
 (♂) (1898). Typus:
 Brisbane)
axiolecta Turner, Proc. Roy. Soc. Victoria, 35, p. Tasmanien
 53 (♀) (1922).
chrysomieta Meyrick, Trans. Ent. Soc. Lond., 77, Marquesas
 p. 166 (♂, ♀) (1929). Ins. (Typ.:
 Hiva Oa)
chrysopetra Meyrick, Trans. Ent. Soc. Lond., 77, Gesellschafts-
 p. 169 (♂) (1929). Ins. (Typ.:
 Tahiti)
citrocosma Meyrick, Trans. Ent. Soc. Lond., 77, Marquesas
 p. 166 (♂, ♀) (1929). Ins. (Typ.:
 Hiva)
clerica Meyrick, Trans. Ent. Soc. Lond., 77, p. Marquesas
 167 (♂, ♀) (1929). Ins. (Typ.:
 Hiva Oa)
commercialis Meyrick, Trans. Ent. Soc. Lond., 77, Marquesas
 p. 167 (♂, ♀) (1929). Ins. (Typ.:
 Hiva Oa)
crocospila Turner, Proc. Roy. Soc. Victoria, 35, Victoria
 p. 53 (♀) (1922). (Typus:
 Gisborne)
crypterythra Lower, Trans. Roy. Soc. S. Austral., N.S.Wales
 25, p. 67 (1901) (*Eclipsiodes*).
deliniens Lucas, Proc. Roy. Soc. Queensland, 13, Queensland
 p. 84 (♂, ♀) (1898). Typus:
 Brisbane)

- emmetropis** Turner, Proc. Roy. Soc. Queensland, 27, N.S.Wales
p. 52 (♂) (1915). (Typus:
Kościuszko-
berg)
- encausta** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Tasmanien
p. 111 (♂) (1885).
- eutacta** Turner, Proc. Linn. Soc. N. S. Wales, 56, Queensland
p. 343 (♂) (1931). (Typus:
Nation.Park)
- exterminata** Meyrick, Trans. Ent. Soc. Lond., 77, Austral. Ins.
p. 169 (♂, ♀) (1929). (Typ.: Rapa)
- favilliferella** Walker, Cat. Lep. Het. Brit. Mus., Australien,
35, p. 179 (1866) (*Nephopteryx*). — Mey- Tasmanien
rick, Trans. Proc. N. Zeal. Inst., 17, p.
119 (1885).
- gethosyna** Turner, Proc. Roy. Soc. Victoria, 35, Queensland
p. 53 (♀) (1922). (Typus:
Nat. Park)
- hypoxantha** Lower, Trans. Roy. Soc. S. Austral., S.Australien,
20, p. 158 (♀) (1896). Victoria
- ischnoptera** Turner, Proc. Roy. Soc. Victoria, 35, Victoria
p. 51 (♂) (1922). (Typus:
Ringwood)
- ithyntis** Turner, Proc. Roy. Soc. Victoria, 35, p. N.S.Wales,
52 (♂, ♀) (1922). Victoria
- leucomela** Lower, Trans. Roy. Soc. S. Austral., S.u. W.-
17, p. 165 (♂, ♀) (1885). Australien
- lichenopa** Lower, Proc. Linn. Soc. N. S. Wales, N.S.Wales
22, p. 18 (♂, ♀) (1897).
- melanoxantha** Turner, Proc. Roy. Soc. Victoria, Queensland
35, p. 52 (♀) (1922). (Typus:
Nat. Park)
- mesogramma** Lower, Proc. Linn. Soc. N. S. Wales, N.S.Wales
25, p. 408 (♀) (1900). (Typus:
Broken Hill)
- metacrossa** Hampson, Ann. Mag. Nat. Hist., (s. 8) Holl.
20, p. 280 (♂) (1917). Neu-Guinea
- nephelitis** Meyrick, Trans. Ent. Soc. Lond., p. 247 Australien
(♂, ♀) (1887) (*Xeroscopa*).
- niphетodes** Turner, Proc. Linn. Soc. N. S. Wales, Queensland
56, p. 343 (♂, ♀) (1931). (Typus:
Bunya Mts.)
- officialis** Meyrick, Trans. Ent. Soc. Lond., 77, p. Marquesas
168 (♂, ♀) (1929). Ins. (Typ.:
Hiva Oa)
- opostactis** Meyrick, Trans. Ent. Soc. Lond., 77, Marquesas
p. 168 (♂, ♀) (1929). Ins. (Typ.:
Fatu Hiva)

- philorhpha** Meyrick, Trans. Ent. Soc. Lond., 77, p. 168 (♂, ♀) (1929). Gesellschafts-
Ins. (Typ.:
Tahiti)
- psednopa** Meyrick, Trans. Ent. Soc. Lond., 77, p. 169 (♀) (1929). Austral. Ins.
(Typ.: Rapa)
- schizodesma** Lower, Proc. Linn. Soc. S. Austral., 24, p. 90 (♂, ♀) (1899). N.S.Wales,
S.Australien
- spectacularis** Meyrick, Trans. Ent. Soc. Lond., 77, p. 167 (♂, ♀) (1929). Marquesas
Ins. (Typ.:
Hiva Oa)
- stenopa** Lower, Proc. Linn. Soc. N. S. Wales, 26 (1901), p. 66 (♂) (1902). S.Australien
(Typus:
Blackerood)
- susanae** Lower, Proc. Linn. Soc. N. S. Wales, 25, p. 38 (♂, ♀) (1900). N.S.Wales
- threnodes** Meyrick, Trans. Ent. Soc. Lond., p. 246 (♂, ♀) (1887). W.Australien
- tricitra** Meyrick, Exot. Microlep., 3, p. 583 (1930). Holl.
Neu-Guinea
- tristicta** Turner, Proc. Roy. Soc. Victoria, 35, p. 54 (♂) (1922). N.S.Wales
(Typus:
Ebor)
- tritocirrha** Turner, Trans. Proc. Roy. Soc. S. Austral., 42, p. 286 (1918). Norfolk Ins.

h) Neuseeländische Subregion.

- thyridias** Meyrick, Trans. Ent. Soc. Lond., p. 229 (♂) (1905). — Hudson, Butt. Moths N. Neuseeland
Zeal., p. 183, t. 22, f. 36 (♀) (1928). (Typus:
Wellington?)
- oreas** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 81 (♂) (1885). — Hampson, Trans. Ent. Neuseeland
Soc. Lond., p. 232 (1897). — Hudson, Butt. Moths N. Zeal., p. 183 (1928).
- philerga** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 81 (♂, ♀) (1885). — Hampson, Trans. Neuseeland
Ent. Soc. Lond., p. 232 (1897). — Long-
staff, Butt.-Hunt. Many Lands, p. 452, 469, 479 (1912). — Hudson, Butt. Moths N. Zeal., p. 183, t. 21, f. 12 (♀) (1928).
- meliturga** Meyrick, Trans. Ent. Soc. Lond., p. 228 (♂, ♀) (1905). — Hudson, Butt. Moths N. Neuseeland
Zeal., p. 183, t. 22, f. 12 (♂) (1928). (Typus:
Auckland &
Wellington)
- chlamydota** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 82 (♂, ♀) (1885). — Hampson, Trans. Neuseeland
Ent. Soc. Lond., p. 232 (1897). — Hudson,
Butt. Moths N. Zeal., p. 184, t. 19, f. 28 (♂) (1928).

- triclera** Meyrick, Trans. Ent. Soc. Lond., p. 230 (♀) (1905). — Hudson, Butt. Moths N. Zeal., p. 184, t. 19, f. 27 (♀) (1928). Neuseeland (Typus: Wellington)
- hemiplaca** Meyrick, Trans. Proc. N. Zeal. Inst., 21, p. 155 (♂) (1889). — Hudson, Butt. Moths N. Zeal., p. 184, t. 22, f. 11 (♀) (1928). Neuseeland
- dochmia** Meyrick, Trans. Ent. Soc. Lond., p. 229 (♂) (1905). — Hudson, Butt. Moths N. Zeal., p. 184, t. 22, f. 2 (♂) (1928). Neuseeland (Typus: Lake Wakatipu)
- minusculalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1503 (♂) (1865). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 82 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 232 (1897). — Hudson, Butt. Moths N. Zeal., p. 184, t. 22, f. 38 (♀) (1928). Neuseeland
- minnalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1504 (♀) (1865). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 83 (1885). — Hampson, Trans. Ent. Soc. Lond., p. 232 (1897). — Longstaff, Butt.-Hunt. Many Lands, p. 461, 469 (1912). — Hudson, Butt. Moths N. Zeal., p. 185, t. 22, f. 37 (♀) (1928). Neuseeland
- chimcria** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 84 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 232 (1897). — Hudson, Butt. Moths N. Zeal., p. 185, t. 2, f. 36 (larva), t. 22, f. 43 (1928). Neuseeland
- dinodes** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 85 (♂) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 232 (1897). — Hudson, Butt. Moths N. Zeal., p. 185, t. 22, f. 3 (♀) (1928). Neuseeland
- parmifera** Meyrick in Chilton, Subantarctic Isl. N. Zeal., 1, p. 72, t. 2, f. 2 (1909). — Tillyard, Ins. Austral. & N. Zeal., p. 432, t. 31, f. 8 (1926). — Hudson, Butt. Moths N. Zeal., p. 186, t. 22, f. 13 (♂) (1928). Auckland Ins.
- acharis** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 85 (♂) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 232 (1897). — Hudson, Butt. Moths N. Zeal., p. 186, t. 21, f. 14 (♀) (1928). Neuseeland
- zophochlaena** Meyrick, Trans. Proc. N. Zeal. Inst., 54, p. 162 (♂) (1923). — Hudson, Butt. Moths N. Zeal., p. 186, t. 49, f. 3 (♂) (1928). Neuseeland
- animosa** Meyrick, Trans. Proc. N. Zeal. Inst., 46, p. 103 (♀) (1904). — Hudson, Butt. Moths N. Zeal., p. 186, t. 45, f. 5 (♂) (1928). Neuseeland
- molifera** Meyrick, Trans. Proc. N. Zeal. Inst., 56, p. 415 (♀) (1926). — Hudson, Butt. Moths N. Zeal., p. 187, t. 51, f. 18 (♀) (1928). Neuseeland
- cymatias** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 86 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. London, p. 232 (1897). — Long-

- staff, Butt.-Hunt. Many Lands, p. 469 (1912).
 — Hudson, Butt. Moths N. Zeal., p. 187, t. 22, f. 40 (♀) (1928).
- microphthalma** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 87 (♂) (1885). — Hampson, Trans. Ent. Soc. London, p. 232 (1897). — Hudson, Butt. Moths N. Zeal., p. 187, t. 5, f. 8 (♀) (1928). Neuseeland
- hemicycla** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 87 (♀) (1885). — Hampson, Trans. Ent. Soc. London, p. 232 (1897). — Tillyard, Ins. Austr. & N. Zeal., p. 432, t. 31, f. 7 (1926). — Hudson, Butt. Moths N. Zeal., p. 187, t. 22, f. 7 (♀) (1928). Neuseeland
- xysmatias** Meyrick, Trans. Proc. N. Zeal. Inst., 39, p. 110 (♂) (1907). — Hudson, Butt. Moths N. Zeal., p. 187, t. 21, f. 5 (♂) (1928). Neuseeland
- ergatis** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 88 (♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 231 (1897). — Hudson, Butt. Moths N. Zeal., p. 188, t. 22, f. 28 (♂) (1928). Neuseeland
- autochroa** Meyrick, Trans. Proc. N. Zeal. Inst., 39, p. 110 (♂) (1907). — Hudson, Butt. Moths N. Zeal., p. 188, t. 21, f. 16 (♀) (1928). Neuseeland
- cinefacta** Philpott, Trans. Proc. N. Zeal. Inst., 56, p. 391 (♂, ♀) (1926). — Hudson, Butt. Moths N. Zeal., p. 188, t. 48, f. 7 (♂) (1928). Neuseeland
- eucapna** Meyrick, Trans. Proc. N. Zeal. Inst., 20, p. 65 (1888). — Hudson, Butt. Moths N. Zeal., p. 188, t. 22, f. 6 (♀) (1928). Neuseeland
- lychnophanes** Meyrick, Trans. Proc. N. Zeal. Inst., 57, p. 697 (♀) (1927). — Hudson, Butt. Moths N. Zeal., p. 188, t. 52, f. 10 (♀) (1928). Neuseeland
- critica** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 88 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 231 (1897). — Hudson, Butt. Moths N. Zeal., p. 189, t. 22, f. 42 (♂) (1928). Neuseeland
- gracilis** Philpott, Trans. Proc. N. Zeal. Inst., 55, p. 209 (♂, ♀) (1924). — Hudson, Butt. Moths N. Zeal., p. 189, t. 50, f. 27 (♂) (1928). Neuseeland
- characta** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 90 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 231 (1897). — Longstaff, Butt.-Hunt. Many Lands, p. 479 (1912). — Hudson, Butt. Moths N. Zeal., p. 189, t. 22, f. 4 (♀) (1928). Neuseeland
- ustimacula** Felder et Rogenhofer, Reise Novara (Het.) 2, p. 17, t. 135, f. 17 (1877). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 91 (1885). — Hampson, Trans. Ent. Soc. Lond., p. 231 (1897). — Hudson, Butt. Moths N. Zeal., p. 189, t. 22, f. 39 (♂) (1928). Neuseeland

- conifera* Butler, Cist. Ent., 2, p. 493 (1879).
pongalis Felder et Regenhöfer, Reise Novara, (Het.), Neuseeland
 p. 17, t. 137, f. 33 (1877). — Meyrick,
 Trans. Proc. N. Zeal. Inst., 17, p. 91 (♂,
 ♀) (1885). — Hampson, Trans. Ent. Soc.
 Lond., p. 231 (1897). — Hudson, Butt.
 Moths N. Zeal., p. 189, t. 22, f. 1 (♀)
 (1928) (*psammitis*).
melanaegis Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
 17, p. 92 (♂, ♀) (1885). — Hampson,
 Trans. Ent. Soc. Lond., p. 231 (1897). —
 Hudson, Butt. Moths N. Zeal., p. 190, t.
 21, f. 51 (♀) (1928).
trapezophora Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
 17, p. 93 (♀) (1885). — Hampson, Trans.
 Ent. Soc. Lond., p. 231 (1897). — Tillyard,
 Ins. Austral. & N. Zeal., p. 432, t. 31, f. 9
 (1926). — Hudson, Butt. Moths N. Zeal.,
 p. 190, t. 21, f. 27 (♂) (1928).
philetaera Meyrick, Trans. Proc. N. Zeal. Inst., 17, Neuseeland
 p. 93 (♂) (1885). — Hampson, Trans.
 Ent. Soc. Lond., p. 231 (1897). — Hudson,
 Butt. Moths N. Zeal., p. 190 (1928).
galactalis Hudson, Ent. Mag. Mon., 49, p. 250 Neuseeland
 (1913); Butt. Moths N. Zeal., p. 190, t. 21,
 f. 46 (♂) (1928).
locularis Meyrick, Trans. Proc. N. Zeal. Inst., 44, Neuseeland
 p. 188 (♂) (1912). — Hudson, Bull. Moths
 N. Zeal., p. 190, t. 26, f. 15 (♂) (1928).
torodes Meyrick, Trans. Ent. Soc. Lond., p. 568 Neuseeland
 (1901). — Hudson, Butt. Moths N. Zeal.,
 p. 191 (1928).
triscellis Meyrick in Chilton, Subantarctic Isl. N. Auckland Ins.
 Zeal., 1, p. 71, t. 2, f. 11 (1909). — Hudson,
 Butt. Moths N. Zeal., p. 191, t. 21, f. 4
 (♀) (1928).
periphanes Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
 17, p. 94 (♂) (1885). — Hampson, Trans.
 Ent. Soc. Lond., p. 231 (1897). — Long-
 staff, Butt.-Hunt. Many Lands, p. 461 (1912).
 — Hudson, Butt. Moths N. Zeal., p. 191,
 t. 21, f. 3 (♀) (1928).
colpota Meyrick, Trans. Proc. N. Zeal. Inst., 20, Neuseeland
 p. 65 (1888). — Hampson, Trans. Ent.
 Soc. Lond., p. 231 (1897). — Hudson, Butt.
 Moths N. Zeal., p. 191, t. 21, f. 1 (♂)
 (1928).
choristis Meyrick, Trans. Proc. N. Zeal. Inst., 39, Neuseeland
 p. 111 (♂) (1907). — Hudson, Butt. Moths
 N. Zeal., p. 191, t. 21, f. 2 (♀) (1928).
phalerias Meyrick, Trans. Ent. Soc. Lond., p. 230 Neuseeland
 (♀) (1905). — Hudson, Butt. Moths N.
 Zeal., p. 192, t. 21, f. 6 (♀) (1928).
quaestoria Meyrick, Trans. Proc. N. Zeal. Inst., 60, Neuseeland
 p. 487 (♂) (1929).
diphtheralis Walker, Cat. Lep. Het. Brit. Mus., Neuseeland
 34, p. 1501 (♀) (1865). — Meyrick, Trans.
 Proc. N. Zeal. Inst., 17, p. 94 (1885). —
 Hampson, Trans. Ent. Soc. London, p. 231

- (1897). — Longstaff, Butt.-Hunt. Many Lands, p. 462 (1912). — Hudson, Butt. Moths N. Zeal., p. 184, t. 21, f. 20 (♂) (1928).
- submarginalis** Walker, Cat. Lep. Het. Brit. Mus., 27, p. 48 (1863) (*Hypochalcia*). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 95 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 231 (1897). — Longstaff, Butt.-Hunt. Many Lands, p. 452, 470, 474, 479 (1912). — Hudson, Butt. Moths N. Zeal., p. 192, t. 21, f. 7—9 (1928).
- maoriella** Walker, Cat. Lep. Het. Brit. Mus., 35, p. 1720 (1866) (*Nephopteryx*).
- ? **linealis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1503 (♀) (1865) (*Nephopteryx*). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 119 (1885). — Hampson, Trans. Ent. Soc. Lond., p. 232 (1897).
- cataxesta** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 96 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 231 (1897). — Hudson, Butt. Moths N. Zeal., p. 193, t. 21, f. 19 (♂) (1928).
- asaleuta** Meyrick, Trans. Proc. N. Zeal. Inst., 39, p. 111 (♀) (1907). — Hudson, Butt. Moths N. Zeal., p. 193, t. 21, f. 17 (♂) (1928).
- tetracycla** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 97 (♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 231 (1897). — Hudson, Butt. Moths N. Zeal., p. 193, t. 21, f. 21 (♀) (1928).
- gyrotoma** Meyrick, Trans. Proc. N. Zeal. Inst., 41, p. 7 (♂) (1909). — Hudson, Butt. Moths N. Zeal., p. 193, t. 21, f. 18 (♂) (1928). — Philpott, Trans. Proc. N. Zeal. Inst., 62, p. 29 (1931).
- repercussa** Philpott, Trans. Proc. N. Zeal. Inst., 60, p. 301 (♂, ♀) (1929).
- indistinctalis** Walker, Cat. Lep. Het. Brit. Mus., 27, p. 48 (1863) (*Hypochalcia*). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 97 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 226 (1897) (*Xeroscopa*). — Longstaff, Butt. Hunt. Many Lands, p. 452 (1912). — Hudson, Butt. Moths N. Zeal., p. 193, t. 21, f. 10 (♂) (1928).
- rakaiensis** Knaggs, Ent. Mon. Mag., 4, p. 80 (1867). — Hampson, Trans. Ent. Soc. Lond., p. 231 (1897).
- bisnualls** Hudson, Butt. Moths N. Zeal., p. 194, t. 21, f. 45 (♀) (1928).
- chalicodes** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 98 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 231 (1897). — Longstaff, Butt.-Hunt. Many Lands, p. 461 (1912). — Hudson, Butt. Moths N. Zeal., p. 194, t. 22, f. 13 (♀) (1928).

- ciserodes* Meyrick, Trans. Proc. N. Zeal. Inst., 52, p. 30 (♂) (1920).
- psammitis** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 99 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 230 (1897). — Tillyard, Ins. Austral. & N. Zeal., p. 432, f. 33 (caput) (1926). — Hudson, Butt. Moths N. Zeal., p. 194, t. 24, f. 51 (1928). Neuseeland, Chatham Ins.
- leptalea** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 98 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 230 (1897). — Longstaff, Butt.-Hunt. Many Lands, p. 452 (1912). — Hudson, Butt. Moths N. Zeal., p. 194 (1928). Neuseeland, Chatham Ins.
- leptophaea* Meyrick, Trans. Ent. Soc. Lond., p. 277 (1902). (Chatham Ins.).
- flmbriata** Philpott, Trans. Proc. N. Zeal. Inst., 49, p. 243 (1917). — Hudson, Butt. Moths N. Zeal., p. 194, t. 45, f. 3 (♂) (1928). Neuseeland
- epicomia** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 99 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 230 (1897). — Hudson, Butt. Moths N. Zeal., p. 195, t. 21, f. 33 (♀) (1928). Neuseeland, Kermadec u. Auckland Ins.
- teredayi** Knaggs, Ent. Mon. Mag., 4, p. 80 (1867). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 100 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 230 (1897). — Longstaff, Butt.-Hunt., p. 455, 475 (1912). — Philpott, Trans. Proc. N. Zeal. Inst., 49, p. 221 (1917). — Tillyard, Ins. Austral. & N. Zeal., p. 432, t. 31, f. 11 (1926). — Hudson, Butt. Moths N. Zeal., p. 195, t. 21, f. 34 (♂) (1928). Neuseeland
- moanalis* Felder et Rogenhofer, Reise Novara (Het.), p. 17, t. 137, f. 34 (1877).
- vulpecula** Meyrick, Trans. Proc. N. Zeal. Inst., 57, p. 697 (♀) (1896). — Hudson, Butt. Moths N. Zeal., p. 195, t. 21, f. 32 (♀) (1928). Neuseeland
- acompa** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 100 (♂) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 230 (1897). — Hudson, Butt. Moths N. Zeal., p. 195, f. 45, f. 6 (♂) (1928). Neuseeland
- cyptastis** Meyrick, Trans. Proc. N. Zeal. Inst., 41, p. 7 (♂, ♀) (1909). — Hudson, Butt. Moths N. Zeal., p. 195, t. 22, f. 30 (♂) (1928). Neuseeland
- illota** Philpott, Trans. Proc. N. Zeal. Inst., 51, p. 224 (♂, ♀) (1919). — Hudson, Butt. Moths N. Zeal., p. 196, t. 20, f. 12 (♂) (1928). Neuseeland
- Biologie: Chappell, Trans. Proc. N. Zeal. Inst., 60, p. 262 (1929).
- manganeutis** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 102 (♂) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 230 (1897). — Hudson, Butt. Moths N. Zeal., p. 196 (1928). Neuseeland
- cryptsinoa** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 102 (♂) (1885). — Hampson, Trans.

- Ent. Soc. Lond., p. 230 (1897). — Meyrick, Trans. Proc. N. Zeal. Inst., 53, p. 334 (1921). — Hudson, Butt. Moths N. Zeal., p. 196, t. 22, f. 9 (♂) (1928).
- agana* Meyrick, Trans. Proc. N. Zeal. Inst., 44, p. 119 (1912).
- alopecias** Meyrick, Trans. Ent. Soc. Lond., p. 570 (♂, ♀) (1901). — Hudson, Butt. Moths N. Zeal., p. 196 (1928). Neuseeland (Typu: Mount Cook)
- axena** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 103 (♂) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 230 (1897). — Longstaff, Butt.-Hunt. Many Lands, p. 472, 482 (1912). — Hudson, Butt. Moths N. Zeal., p. 196, t. 44, f. 22 (♂) (1928). Neuseeland
- pachyerga** Meyrick, Trans. Proc. N. Zeal. Inst., 57, p. 697 (♂) (1927). — Hudson, Butt. Moths N. Zeal., p. 196, t. 52, f. 9 (♂) (1928). Neuseeland
- exilis** Knaggs, Ent. Mon. Mag., 4, p. 81 (1867). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 104 (♂) (1885). — Hudson, Butt. Moths N. Zeal., p. 197, t. 21, f. 44 (♀) (1928). Neuseeland
- turneri** Philpott, Trans. Proc. N. Zeal. Inst., 58, p. 362 (♂) (1928). Neuseeland
- steropaea** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 103 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. London, p. 230 (1897). — Hudson, Butt. Moths N. Zeal., p. 197, t. 22, f. 8 (♂) (1928). Neuseeland
- elaphra** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 105 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 230 (1897). — Hudson, Butt. Moths N. Zeal., p. 197, t. 22, f. 5 (♀) (1928). Neuseeland
- atmogramma** Meyrick, Trans. Proc. N. Zeal. Inst., 47, p. 202 (♂) (1915). — Hudson, Butt. Moths N. Zeal., p. 197, t. 45, f. 25 (♂) (1928). Neuseeland
- paltomacha** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 105 (♂) (1885). — Hampson, Trans. Ent. Soc. London, p. 230 (1897). — Longstaff, Butt.-Hunt. Many Lands, p. 471, 474 (1912). — Hudson, Butt. Moths N. Zeal., p. 197, t. 21, f. 36 (♂), 37 (♀) (1928). Neuseeland
- deltophora** Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 106 (♂) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 230 (1897). — Hudson, Butt. Moths N. Zeal., p. 197, t. 21, f. 39 (♀) (1928). Neuseeland
- pallidula** Philpott, Trans. Proc. N. Zeal., 58, p. 363 (♂, ♀) (1928). Neuseeland
- sabulosella** Walker, Cat. Lep. Het. Brit. Mus., 27, p. 178 (1863) (*Crambus*). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 106 (♂, ♀) (1885). — Hampson, Trans. Ent. Soc. Lond., p. 230 (1897). — Hudson, Butt. Moths N. Zeal., p. 198, t. 21, f. 38 (♂) (1928). Neuseeland, Chatham Ins., Auckland Ins

- panopla** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Neuseeland
p. 107 (♂, ♀) (1885). — Hampson, Trans.
Ent. Soc. Lond., p. 230 (1897). — Hudson,
Butt. Moths, p. 198, t. 21, f. 43 (♂)
(1928).
- clavata** Philpott, Trans. Proc. N. Zeal. Inst., 44, Neuseeland
p. 116 (♂) (1912). — Hudson, Butt. Moths,
p. 198, t. 21, f. 40 (♀) (1928).
- trivirgata** Felder et Rogenhofer, Reise Novara Neuseeland
(Het.), p. 12, t. 137, f. 29 (1877) (*Cram-*
bus). — Meyrick, Trans. Proc. N. Zeal.
Inst., 17, p. 107 (♂, ♀) (1885). — Hamp-
son, Trans. Ent. Soc. Lond., p. 230 (1897).
— Tillyard, Ins. Austr. & N. Zeal., p. 432,
t. 31, f. 10 (1926). — Hudson, Butt. Moths
N. Zeal., p. 198, t. 21, f. 42 (♂) (1928).
- augastis** Meyrick, Trans. Proc. N. Zeal. Inst., 39, Neuseeland
p. 112 (♂) (1907). — A. Longstaff, Butt.
Hunt. Many Lands, p. 469 (1912). — Hud-
son, Butt. Moths N. Zeal., p. 188, t. 24,
f. 8 (♀) (1928).
- petrina** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Neuseeland
p. 111 (♂, ♀) (1885) (*Xeroscopa*). — Hamp-
son, Trans. Ent. Soc. Lond., p. 226 (1897)
(*Xeroscopa*). — Hudson, Butt. Moths N.
Zeal., p. 199, t. 21, f. 11, 12 (1928).
- contexta** Philpott, Trans. Proc. N. Zeal. Inst., 62, Neuseeland
p. 28 (♂, ♀) (1931).
- legionaria** Philpott, Trans. Proc. N. Zeal. Inst., Neuseeland
58, p. 362 (♂, ♀) (1928).
- limatula** Philpott, Trans. Proc. N. Zeal. Inst., 61, Neuseeland
p. 436 (♂, ♀) (1930).
- halopis** Meyrick in Chilton, Subantarctic Isl. N. Auckland Ins.
Zeal., 1, p. 72, t. 2, f. 3 (1909). — Hudson,
Butt. Moths N. Zeal., p. 199, t. 24, f. 50
(♂) (1928).
- cyameuta** Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
17, p. 112 (♂, ♀) (1885). — Hampson,
Trans. Ent. Soc. Lond., p. 226 (1897)
(*Xeroscopa*). — Longstaff, Butt.-Hunt. Many
Lands, p. 477 (1912) (*Xeroscopa*). — Hud-
son, Butt. Moths N. Zeal., p. 199, t. 21,
f. 54 (♀) (1928).
- falsa** Philpott, Trans. Proc. N. Zeal. Inst., 55, Neuseeland
p. 208 (♂, ♀) (1924). — Hudson, Butt.
Moths N. Zeal., p. 200, t. 50, f. 23 (♂)
(1928).
- autumna** Philpott, Trans. Proc. N. Zeal. Inst., 58, Neuseeland
p. 83 (♂, ♀) (1927).
- declivis** Philpott, Trans. Proc. N. Zeal. Inst., 50, Neuseeland
p. 126 (♂) (1918). — Hudson, Butt. Moths
N. Zeal., p. 200, t. 44, f. 23 (♀) (1928).
- dryphactis** Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
43, p. 61 (1911). — Hudson, Butt. Moths
N. Zeal., p. 200, t. 21, f. 53 (♀) (1928).
- astragalota** Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
17, p. 113 (♂, ♀) (1885) (*Xeroscopa*). —
Hampson, Trans. Ent. Soc. Lond., p. 226
(1897) (*Xeroscopa*). — Longstaff, Butt.-

Hunt. Many Lands, p. 479 (1912). — Hudson, Butt. Moths N. Zeal., p. 200, t. 24, f. 49 (♂) (1928).

- rotuella** Felder et Rogenhofer, Reise Novara, p. 7, Neuseeland
t. 137, f. 30 (1874) (*Crambus*). — Meyrick,
Trans. Proc. N. Zeal. Inst., 17, p. 113 (♀)
(1885) (*Xeroscopa*). — Hampson, Trans. Ent.
Soc. Lond., p. 226 (1897) (*Xeroscopa*). —
Longstaff, Butt. - Hunt. Many Lands, p.
469 (1912) (*Xeroscopa*). — Hudson, Butt.
Moths N. Zeal., p. 200, t. 21, f. 41 (♂)
(1928).
- scripta** Philpott, Trans. Proc. N. Zeal. Inst., 50, Neuseeland
p. 126 (♂, ♀) (1918). — Hudson, Butt.
Moths N. Zeal., p. 200, t. 44, f. 24 (♂)
(1928).
- harpalea** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Neuseeland
p. 114 (1885) (*Xeroscopa*). — Hampson,
Trans. Ent. Soc. London, p. 226 (1897)
(*Xeroscopa*). — Hudson, Butt. Moths N.
Zeal., p. 201 (1928).
- ejuncida** Knaggs, Ent. Mon. Mag., 4, p. 81 (1867) Neuseeland
(*Xeroscopa*). — Meyrick, Trans. Proc. N.
Zeal. Inst., 17, p. 115 (♂, ♀) (1885). —
Hampson, Trans. Ent. Soc. London, p. 226
fig. (♂) (1897), (*Xeroscopa*). — Hudson,
Butt. Moths N. Zeal., p. 201, t. 21, f. 52
(♀) (1928).
- niphosphora** Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
17, p. 115 (♂, ♀) (1885) (*Xeroscopa*). —
Hudson, Butt. Moths N. Zeal., p. 201, t.
24, f. 9 (♂) (1928).
- niphospora* Hampson, Trans. Ent. Soc.
Lond., p. 226 (1897) (*Xeroscopa*).
- apheles** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Neuseeland
p. 115 (♂, ♀) (1885). — Hampson, Trans.
Ent. Soc. Lond., p. 226 (1897) (*Xeroscopa*).
— Hudson, Butt. Moths N. Zeal., p. 201
(1928).
- aspidota** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Neuseeland
p. 115 (1885). (*Xeroscopa*). — Hampson,
Trans. Ent. Soc. Lond., p. 226 (1897) (*Xero-*
scopa). — Tillyard, Ins. Austral. & N. Zeal.,
p. 431, f. 6 (1926). — Hudson, Butt. Moths
N. Zeal., p. 201, t. 24, f. 10 (♀) (1928).
- sideraspis** Meyrick, Trans. Ent. Soc. Lond., p. 231 Neuseeland
(♂, ♀) (1905). — Hudson, Butt. Moths
N. Zeal., p. 202, t. 24, f. 24 (♀) (1928).
(Typus:
Mount
Earnslaw &
Humboldt
Range)
- nomeutis** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Neuseeland
p. 116 (♂, ♀) (1885). — Hampson, Trans.
Ent. Soc. Lond., p. 227 (1897) (*Xeroscopa*).
— Hudson, Butt. Moths N. Zeal., p. 202,
t. 50, f. 28 (♂) (1928).
- famularis** Philpott, Rec. Auckland Inst. Mus., 1, Neuseeland
p. 3 (♀) (1930).

- pura** Philpott, Trans. Proc. N. Zeal. Inst., 55, Neuseeland
p. 208 (♂, ♀) (1924). — Hudson, Butt.
Moths N. Zeal., p. 202, t. 24, f. 39 (♂),
40 (♀) (1928).
- caliginosa** Philpott, Trans. Proc. N. Zeal. Inst., Neuseeland
50, p. 127 (♂) (1918). — Hudson, Butt.
Moths N. Zeal., p. 202, t. 45, f. 2 (♀)
(1928).
- parachalca** Meyrick, Trans. Ent. Soc. Lond., p. 569 Neuseeland
(♂) (1901). — Hudson, Butt. Moths N.
Zeal., p. 202, t. 22, f. 10 (♀) (1928).
- organea** Meyrick, Trans. Ent. Soc. Lond., p. 569 Neuseeland
(♂) (1901). — Hudson, Butt. Moths N.
Zeal., p. 202, t. 22, f. 27 (♂) (1928).
- pascoella** Philpott, Trans. Proc. N. Zeal. Inst., Neuseeland
p. 43 (♂, ♀) (1920). — Hudson, Butt.
Moths N. Zeal., p. 203, t. 48, f. 8 (♀)
(1928).
- epicremna** Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
17, p. 117 (♂) (1885). — Hampson, Trans.
Ent. Soc. Lond., p. 226 (1897) (*Xeroscopa*).
— Hudson, Butt. Moths N. Zeal., p. 203,
t. 46, f. 4 (♀) (1928).
- luminatrix** Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
41, p. 8 (♂, ♀) (1919). — Hudson, Butt.
Moths N. Zeal., p. 203, t. 21, f. 48 (♂)
(1928).
- legnota** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Neuseeland
p. 117 (♂, ♀) (1885) (*Xeroscopa*). —
Hampson, Trans. Ent. Soc. Lond., p. 227
(1897) (*Xeroscopa*). — Hudson, Butt. Moths
N. Zeal., p. 203, t. 21, f. 49 (♂) (1928).
- octophora** Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
17, p. 118 (♂, ♀) (1885). — Hampson,
Trans. Ent. Soc. Lond., p. 227 (1897) (*Xero-*
scopa). — Hudson, Butt. Moths N. Zeal.,
p. 203, t. 24, f. 21 (♀) (1928).
- chalara** Meyrick, Trans. Ent. Soc. Lond., p. 570 Neuseeland
(♂, ♀) (1901). — Hudson, Butt. Moths
N. Zeal., p. 204, t. 24, f. 41 (♀) (1928).
- sinuata** Philpott, Trans. Proc. N. Zeal. Inst., 61, Neuseeland
p. 436 (♂, ♀) (1930).
- fumata** Philpott, Trans. Proc. N. Zeal. Inst., 47, Neuseeland
p. 198 (♂) (1915). — Hudson, Butt. Moths
N. Zeal., p. 204, t. 44, f. 11 (♂) (1928).
- asterisca** Meyrick, Trans. Proc. N. Zeal. Inst., 17, Neuseeland
p. 118 (♂, ♀) (1885) (*Xeroscopa*). — Hud-
son, Butt. Moths N. Zeal., p. 204, t. 21,
f. 47 (♀) (1928).
- asterica* Hampson, Trans. Ent. Soc. Lond.,
p. 227 (1897) (*Xeroscopa*).
- leucogramma** Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
17, p. 119 (1885) (♂, ♀) (*Xeroscopa*). —
Hampson, Trans. Ent. Soc. Lond., p. 227
(1897) (*Xeroscopa*). — Hudson, Trans. Proc.
N. Zeal., p. 204, t. 22, f. 41 (♀) (1928).

Species incertae sedis.

- oculata** Philpott, Trans. Proc. N. Zeal. Inst., 58, Neuseeland
p. 82 (♂, ♀) (1927).
parca Philpott, Trans. Proc. N. Zeal. Inst., 58, Neuseeland
p. 362 (♂, ♀) (1928).
subditella Walker, Cat. Lep. Het. Brit. Mus., 35, Neuseeland
p. 1720 (♂) (1866) (*Nephopteryx*).
subita Philpott, Trans. Proc. N. Zeal. Inst., 44, Neuseeland
p. 116 (1911) (*Orocrambus*); 62, p. 28
(1931).
sylvestris Clarke, Trans. Proc. N. Zeal. Inst., 56, Neuseeland
p. 418 (♀) (1926).
tuicana Clarke, Trans. Proc. N. Zeal. Inst., 56, Neuseeland
p. 418 (♂, ♀) (1926).
ustiramis Meyrick, Trans. Proc. N. Zeal. Inst., Neuseeland
62, p. 95 (♂) (1931).
 392

5. Genus Tetraprosopus Butler

Butler, Ann. Mag. Nat. Hist., (s. 5) 9, p. 97 (1882). — Meyrick,
Trans. Proc. N. Zeal. Inst., 17, p. 108 (1885). — Hampson, Trans.
Ent. Soc. Lond., p. 224, 227 (1897).

(Typus generis: *meyrickii* Butl.)

- meyrickii** Butler, Ann. Mag. Nat. Hist., (s. 5) 9, N.S.Wales,
p. 97 (1882). — Meyrick, Trans. Proc. N. Victoria,
Zeal. Inst., 17, p. 108 (♂, ♀) (1885). — S.Australien
Hampson, Trans. Ent. Soc. Lond., p. 228,
fig. (♂) (1897).
 Biologie: Meyrick, Trans. Proc. N. Zeal.
Inst., 17, p. 109 (188).
paracycla Lower, Proc. Linn. Soc. N. S. Wales, N.S.Wales
26 (1901), p. 667 (♀) (1902). (Typus:
2 Broken
Hill)

6. Genus Promylaea Meyrick

Meyrick, Fauna Hawai., 1, p. 227 (1899) (Typus generis: *P. pyropae*
Meyr.).

- pyropa** Meyrick, Fauna Hawai., 1, p. 227, t. 6, Hawai
f. 9 (♀) (1899).
quadrifascia Swezey, Proc. Hawai Ent. Soc., 8, Hawai
p. 523 (1924).
 Biologie: Swezey, Proc. Hawai Ent. Soc.,
2, p. 221 (1912).
 2

7. Genus Mestolobes Butler

Butler, Trans. Ent. Soc. Lond., p. 37 (1882). — Hampson, Trans.
Ent. Soc. Lond., p. 224, 228 (1897).

(Typus generis: *M. oenone* Butl. = *M. abnormis* Butl.)

a) Hawaiische Region.

- amethystias** Meyrick, Fauna Hawai., 1, p. 229, Hawai
t. 6, f. 10 (♀) (1899).
chrysomolybda Meyrick, Fauna Hawai., 1, p. 230, Hawai
f. 11 (♂, ♀) (1899).

- chrysomolybdoides** Swezey, Proc. Hawai Ent. Soc., Hawai 4, p. 380, fig. (♀) (1920).
- xanthoscia** Meyrick, Fauna Hawai., 1, p. 230 (♀) Hawai (1899).
- crypsichrysa** Meyrick, Fauna Hawai., 1, p. 230 (♂, Hawai ♀) (1899).
- erynnis** Meyrick, Fauna Hawai., 1, p. 231, f. 12 Hawai (♂) (1899).
- arctura** Meyrick, Fauna Hawai., 1, p. 231 (♀) Hawai (1899).
- sirina** Meyrick, Fauna Hawai., 1, p. 231 (♂) (1899). Hawai
- epidelta** Meyrick, Fauna Hawai., 1, p. 231, f. 13 Hawai (♀) (1899).
- iochrysa** Meyrick, Fauna Hawai., 1, p. 232, f. Hawai 14 (♀) (1899).
- droseropa** Meyrick, Fauna Hawai., 1, p. 232 (♀) Hawai (1899).
- ombrias** Meyrick, Fauna Hawai., 1, p. 233, f. 15 Hawai (♂, ♀) (1899).
- banausa** Meyrick, Fauna Hawai., 1, p. 233 (♂, Hawai ♀) (1899).
- minusecula** Butler, Ann. Mag. Nat. Hist., (s. 5) 7, Hawai p. 325 (1881) (*Boreophila*). — Meyrick, Trans. Ent. Soc. Lond., p. 230 (1888). — Hampson, Trans. Ent. Soc. Lond., p. 229 (1897). — Meyrick, Fauna Hawai., 1, p. 233 (1899).
- simæthina* Butler, Trans. Ent. Soc. Lond., p. 38 n. 9 (1882).
- ochrias** Meyrick, Fauna Hawai., 1, p. 234 (♂) Hawai (1899).
- scleropis** Meyrick, Fauna Hawai., 1, p. 234, f. 16 Hawai (♂, ♀) (1899).
- eurylyca** Meyrick, Fauna Hawai., 1, p. 235 (♂) Hawai (1899).
- orthrias** Meyrick, Fauna Hawai., 1, p. 235, t. 6, Hawai f. 17 (♂) (1899).
- pessias** Meyrick, Fauna Hawai., 1, p. 236, t. 6, Hawai f. 18 (♂, ♀) (1899).
- chimoniás** Meyrick, Fauna Hawai., 1, p. 236 (♂) Hawai (1899).
- aphrias** Meyrick, Fauna Hawai., 1, p. 237 (♂, ♀) Hawai (1899).
- mesacma** Meyrick, Fauna Hawai., 1, p. 237, t. 6, Hawai f. 19 (♂) (1899).
- chorolychna** Meyrick, Fauna Hawai., 1, p. 237 Hawai (♂) (1899).
- perixantha** Meyrick, Fauna Hawai., 1, p. 238, f. Hawai 20 (♂, ♀) (1899).
- semichroa** Butler, Trans. Ent. Soc. Lond., p. 39 Hawai (1882). — Meyrick, Trans. Ent. Soc. Lond., p. 229 (1888); Hampson, *ibid.*, p. 229 (1897). — Meyrick, Fauna Hawai., 1, p. 238 (♂) (1899).
- pragmatica** Meyrick, Fauna Hawai., 1, p. 239 (♂) Hawai (1899).
- abnormis** Butler, Trans. Ent. Soc. Lond., p. 35 Hawai (♀) (1882) (*Metasia?*). — Meyrick, Trans. Ent. Soc. Lond., p. 228 (1888). — Hampson,

- Trans. Ent. Soc. Lond., p. 228 fig. (♂),
p. 229 (1897). — Meyrick, Fauna Hawai,
1, p. 239 (1899).
oenone Butler, Trans. Ent. Soc. Lond., p.
37 (♂) (1882).
autodoxa Meyrick, Fauna Hawai, 1, p. 239, t. 6, Hawai
f. 21 (♂, ♀) (1899); 3, p. 361 (1904).
antichora Meyrick, Fauna Hawai, 1, 3, p. 361 (♂) Hawai
(1904).
homalopa Meyrick, Fauna Hawai, 1, p. 240 (♂, ♀) Hawai
(1899).

Species incertae sedis.

- quadrifasciata* Swezey, Proc. Hawai Ent. Soc., 4, Hawai
p. 381 (♀) (1920).
sicaria Meyrick, Ent. Mon. Mag., (s. 2) 15 (40), Hawai
p. 131 (1904).

b) Polynesische Subregion.

- acleropa* Meyrick, Bull. Bishop Mus., 114, p. 340 Marquesas
(1935). Ins.
isoterma Meyrick, Bull. Bishop Mus., 114, p. 340 Marquesas
(1935). Ins.
procelecaris Meyrick, Bull. Bishop Mus., 114, p. 341 Marquesas
(1935). Ins.
35

8. Genus **Dasyscopa** Meyrick

Meyrick, Trans. Ent. Soc. Lond., p. 464 (1894). — Hampson,
Fauna Brit. India Moths, 4, p. 245 (1896); Trans. Ent. Soc.
Lond., p. 224, 239 (1897).

(Typus generis: *homogenes* Meyr.)

- homogenes* Meyrick, Trans. Ent. Soc. Lond., p. 464 (♂) (1894). — Hampson, Fauna Brit. Assam,
India Moths, 4, p. 245, f. 148 (♂) (1896); Malakka,
Trans. Ent. Soc. Lond., p. 239, fig. (♂) Sumbawa
(1897). (Typus: Sumbawa)
planilinealis Warren, Ann. Mag. Nat. Typus:
Hist., (s. 6) 17, p. 207 (♂) (1896) (Sco- Khasias
paria).
polysemalis Hampson, Trans. Ent. Soc. Lond., p. Lifu,
239 (♂) (1897). Loyalty
2. Ins.
456

2. Subfam. **Nymphulinae**.

Fernald in Dyar, List North Amer. Lep. (Bull. U. S. Nat. Mus.,
nr. 52), p. 395 (1902). — Dyar, Journ. N. York Ent. Soc., 14,
p. 72 (1906). — Barnes et Mc. Dunnough, Chock List Lep. Bor.
Amer., p. 135 (1917). — Hampson, Novit. Zool., 25, p. 386 (1918).
— Forbes, Lep. N. York, p. 524, 574 (1928). — De Joannis, L'Amat,

- Papill., 4, p. 281, 285 (1929); 6, p. 8, 19 (1932). — Karny, Biol. Wasserins., p. 165 (1934). — Lhomme, Cat. Lep. France Belge, 2, fasc. 1, p. 99 (1935). — Zerny et Beier in Kükenthal et Krumbach, Handb. Zool., 4 (2), Ins. 2, Lief. 8, Lp. 1701 (1936).
- Nymphulites* Duponchel, Cat. Lep. d'Eur., p. 201 (1844).
- Nymphulidi* (pars) + *Botydi* (pars) Stephens, List Spec. Brit. Anim., 5, p. 236, 238 (1850).
- Nymphulidae* Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 437 (1900).
- Nymphulini* Handlirsch in Handb. Entom., 3, p. 902 (1925).
- Botidae* (pars) Heinemann, Kleinschm., 1, H. 2, p. 8, 18 (1865).
- Botydae* (pars) Wallengren, Oefvers. Kgl.-Vet. Akad. Förh., 8, p. 965, 971 (1871). — Snellen, Vlinders Nederland, 2 (1), Microlep., p. 19 (1882).
- Botydidae* (pars) (p. 591) + *Asopidae* (pars) (p. 624) + *Spilomelidae* (pars) (p. 633) + *Hydrocampidae* (pars) (p. 647) Swinhoe et Cotes, Cat. Moths India, 5 (1889).
- Crambides* (pars) Herrich-Schäffer, Syst. Bearb. Schm. Eur., p. 3—6 (1848).
- Crambina* (pars) Wocke, Cat. Lep. Eur. Microlep., p. 85 (1861).
- Hydrocampinae* Hampson, Fauna Brit. India Moths, 4, p. 187 (1896); Trans. Ent. Soc. Lond., p. 130 (1897); Proc. Zool. Soc. Lond., p. 591 (1898). — Pagenstecher, Zoologica (Chun), H. 29, p. 170 (1900). — Standinger et Rebel, Cat. Lep. Pal. Faun., 2, p. 48 (1901). — Rebel in Spuler, Schm. Eur., 2, p. 189, 223 (1906). — Wahlgren, Svensk Insektfauna, 10 (2), p. 11, 45 (1915). — Janse, Check-list South Afr. Lep. Hot., p. 156 (1917). — Börner in Brohmer, Fauna v. Deutschl., 2. Aufl., p. 348 (1920). — Tullgren & Wahlgren, Svenska Ins., p. 394 (1922). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ. Sapporo, 22, p. 138 (1928); Ins. Matsum., 3, p. 128 (1929). — M. Hering, Tierwelt Mitteleur. (Ergbd., 1), p. 281 (1932). — Eckstein, Schm. Deutschl., 5, p. 45, 61 (1933).
- Hydrocampinae* (p. 445, 446, 456) + *Musotiminae* (p. 445, 446, 457) Ragonot, Ann. Soc. Ent. Fr., (s. 6) 10 [1890] (1891).
- Musotimidae* Meyrick, Trans. Ent. Soc. Lond., p. 217 (1886); p. 436, 468 (1890).
- Pyralidae* (pars) Stephens, Syst. Cat. Brit. Ins., p. 157 (1829).
- Pyralites* (pars) Newman, Ent. Mag., 2, p. 384 (1834).
- Pyralites* (pars) + *Nymphulites* Duponchel, Cat. Méth. Lep. Eur., p. 199, 201 (1844).
- Piralidi* (pars) O. G. Costa, Fauna Regno Napoli, 1832—36).
- Pyralididae* (pars) Lederer, Wien. Ent. Monatschr., 7, p. 334 (1863). — Wocke, Cat. Lep. Eur. Faun., p. 201 (1871).
- Pyralidinae* (pars) Druce, Biol. Centr.-Amer. Het., 2, p. 200 (1895) (1896), p. 545 (1899).
- Pyraustidae* (pars) Meyrick, Handb. Brit. Lep., p. 397 (1895); Revis. Handb. Brit. Lep., p. 375, 414 (1928).
- Pyraustidae* (pars) + *Hydrocampidae* (pars) Barrett, Lep. Brit. Isl., 9, p. 161, 284 (1904).
- Pyraustinae* (pars) Trib. *Pyraustini* (pars) Börner

- in Brohmer, Fauna v. Deutschl., (3. Aufl.), p. 377 (1925);
id. (4. Aufl.) p. 389 (1932).
Steniadae (pars) (p. 146, 232) + *Hydrocampidae*
(pars) (p. 146, 254) + *Botyidae* (pars) (p. 146, 315)
Guenée, Delt. et Pyral. (1854).
Steniadae (pars) + *Hydrocampidae* (pars) Stainton,
Man. Brit. Butterfl. Moths, 2, p. 412, 413 (1859).
Steniadae (pars) (p. 406, 1325) + *Hydrocampidae*
(p. 426, 1331) + *Spilomelidae* (pars) (p. 463, 1343)
+ *Margarodidae* (pars) (p. 489, 1353) + *Botyidae*
(pars) (p. 545, 1380) Walker, List Spec. Ins. Brit. Mus.,
17, 18 (1859); 34 (1865).
Steniidi (pars) + *Hydrocampidi* + *Botysidi*
(pars) Acloquo, Faune France, 3, p. 345 (1897).

1. Genus **Gargela** Walker

Walker, Cat. Lep. Het. Brit. Mus., 29, p. 815 (1864). — Hamp-
son, Ann. Mag. Nat. Hist., (s. 7) 18, p. 374 (1906).

(Typus generis: *G. renatusalis* Walk.)

- Mixophyla* Meyrick, Trans. Ent. Soc. Lond., p. 269 (1887)
(Typus generis: *M. erminea* Meyr.).
Mixophila Hampson, Fauna Brit. India Moths, 4, p. 190
(1896); Trans. Ent. Soc. Lond., p. 132, 134 (1897). — Pagen-
stecher, Zoologica, H. 29, p. 171 (1900). — C. Water-
house, Index Zool., 1, (1880—1900), p. 228 (1902).
Angonia Snellen, Tijds. v. Ent., 36, p. 54 (1893) (*Crambi-
dae*; typus generis: *A. crambidalis* Snell.).
Zebronia (part.) Walker, Cat. Lep. Het. Brit. Mus., 17, p.
484 n. 27 (1859) (*Z. renatusalis* Walk.).
Crambus (part.) Moore, Lep. Ceylon, 3, p. [379], 380
(1886) (*C. ermineus* Moore).

renatusalis Walker, Cat. Lep. Het. Brit. Mus., 17, p. 484 n. 27 (♂) (1859) (*Zebronia*). (Bor-
neo). — Hampson, Fauna Brit. India Moths,
4, p. 190, f. 113 (1896) (*Mixophila*); Trans.
Ent. Soc. Lond., p. 135, fig. (♂) (1897).
— Caradja, Bull. Acad. Roum., 15, p. 178
(1932).

O.Himalaya,
Birma,
Süd- u.
Ostchina,
Ceylon,
Borneo,
Java,
Sumatra,
Sumbawa,
Neu-Guinea

subpurella Walker, Cat. Lep. Het. Brit.
Mus., 29, p. 815 (1864) (*Gargela*) (Cey-
lon).

ermineus Moore, Lep. Ceyl., 3, p. 380,
t. 184, f. 7 (1886) (*Crambus*) (Ceylon).

erminea Meyrick, Trans. Ent. Soc. Lond.,
p. 269 (1887); p. 11 (1894) (*Mixophila*).

crambidalis Snellen, Tijds. v. Ent., 36, p.
56, t. 3, f. 1, (1893) (*Angonia*). (Java,
Ceylon).

renatalis Meyrick, Trans. Ent. Soc. Lond.,
p. 470 (1894) (*Mixophyla*).

niphostola Hampson, Ann. Mag. Nat. Hist., (s. 8) Fergusson
19, p. 362 (♀) (1917). Ins.

trilinealis Hampson, Trans. Ent. Soc. Lond., p. 135 Fergusson
(♂) (1897) (*Mixophila*). Ins.

- apicalis** Pagenstecher, Zoologica, H. 29, p. 172 (1900) Bismarck-Archipel (Neu-Pommern)
- cuprealis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 374 (♂) (1906). Borneo & Pulo Laut
- xanthocasis** Meyrick, Trans. Ent. Soc. Lond., p. 81 (♀) (1897) (*Mixophyla*) (Sangir). — Hampson, Journ. Bombay Nat. Hist. Soc., 12, p. 700, t. 1, f. 8 (1899); id., Trans. Ent. Soc. Lond., p. 135 (1897) (*Mixophyla*). NW.-Himalaya, Pulo Laut, Sangir, Fergusson Ins., Neuguinea
- chrysias** Meyrick, Trans. Ent. Soc. Lond., p. 82 (♂, ♀) (1897) (*Mixophyla*). — Hampson, Trans. Ent. Soc. London, p. 135 (1897) (*Mixophyla*). Amboina (Typus), Banda Ins., Neuguinea
- arcualis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 374 (♂) (1906). Brit. Neuguinea (Typus: Milne Bay)
- obliquivitta** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 362 (1917). 9 Molukken (Typus: Batchian)

2. Genus **Argyractis** Hampson

Hampson, Trans. Ent. Soc. Lond., p. 132, 135 (1897); Ann. Mag. Nat. Hist., (s. 7) 18, p. 375 (1906).

Oxyelophila Forbes, Ent. News, 33, p. 102, t. 5, f. 3 (1922) (subgen.) (typ. sgen.: *A. (O.) callista* Forb.).

Cataclysta (part.) Hampson, Trans. Ent. Soc. Lond., p. 150, 151, 152 n. 45 (1897).

(Typus generis: *A. argentilincalis* Hamps.)

a) **Orientalische Region.**

- malayalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 375 (♀) (1906). Selangor (Malakka) (Typus: Kuala Lumpur)

b) **Aethiopische und Madagassische Region.**

- pervenustalis** Hampson, Trans. Ent. Soc. Lond., p. 152 (♂) (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 377 (1906) (*Argyractis*). Westafrika (Goldküste)

- leucostola** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 363 (♂, ♀) (1917). Brit. Zentr.-Afrika (Typ.: Mt. Mlanje)
- triopalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 377 (♂, ♀) (1906). Sierra Leone (Typus), Nigeria, Old Calabar, Warri
- pentopalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 378 (♀) (1906). Sierra Leone
- albipunctalis** Hampson, Trans. Ent. Soc., p. 152 (♂) (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 381 (1906) (*Argyractis*). Madagaskar
- supercilialis** Hampson, Trans. Ent. Soc. Lond., p. 152 (♂, ♀) (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 381 (1906) (*Argyractis*). Madagaskar
- coloralis** Guenée, Delt. et Pyral., p. 265 (1854) (*Cataclysta*). — Hampson, Trans. Ent. Soc. Lond., p. 153 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 381 (1906) (*Argyractis*). Mauritius
- capensis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 383 (♂, ♀) (1906). Mashonaland, Natal, (Typus: ♀ Kapkolonie)
- lithocharis** Meyrick, Exot. Microlep., 5, p. 8 (♂, ♀) (1936). Belg. Congo
- mimicalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 366 (♂, ♀) (1917). Sierra Leone, N. Nigeria (Typus: ♀ Borgu)
- nyasalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 367 (♂) (1917). Brit. O. Afrika, Tanganyika-Territ. (Typus: Blantyre)
- nandinalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 383 (♀) (1906). Brit. O. Afrika (Typus: Eb Urru)
- tetropalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 384 (♂, ♀) (1906). Nigeria (Typus: Ogbomoso, Loruba)
- nigerialis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 388 (♂) (1906). Nigeria (Typus: Warri)

- leucostrialis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 388 (♂) (1906). Sierra Leone
nymphulalis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 389 (♂) (1906). Natal
 (Typus:
 Kimbolton)

Species incertae sedis.

- acera** Strand, Arch. Naturg., 78 A H. 12, p. 77 Goldküste
 (1912) (*Cataclysta*). — Gaede, Mitt. Zool.
 Mus. Berlin, 8, p. 390 (1917) (*Argyractis*).
dualalis Gaede, Ent. Zeit. Stettin, 77, p. 13 (♂) Kamerun
 (1916).
enryxantha Meyrick, Exot. Microlep., 5, p. 7 (♂) Belg. Congo:
 (1936). (Typus: Fala)
lithorma Meyrick, Exot. Microlep., 5, p. 7 (♂) Belg. Congo
 (1936). (Typus:
 Elisabeth-
 ville)
periopis Hampson, Proc. Zool. Soc. Lond., p. 494, Rhodesia
 t. 40, f. 22 (♂) (1910).
sambesica Strand, Arch. Naturg., 75 A H. 1, p. 384 Rhodesia
 (1909) (*Cataclysta*). — Gaede, Mitt. Zool.
 Mus. Berlin, 8, p. 390 (1917) (*Argyractis*).

c) Nearktische Region.

- callista** W. T. M. Forbes, Ent. News, 33, p. 102, t. 5, Texas
 f. 3 (nervat.) (1922) (*A. Oxyelophila*). (Typus: N.
 Braunfels)

d) Neotropische Region.

- lophosomalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 376 (♂) (1906). Brasilien
 (Rio
 Janeiro)
 (Typus:
 Organ Mts.,
 Tijuca)
danaealis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 376 (♂) (1906). Mexiko
 (Typus:
opulentialis Druce (nec Lederer, 1863), Tabasco
 Biol. Centr.-Amer. Het., 2, p. 272 (1896) (Teapa))
 (part.) (*Cataclysta*).
chrysopalis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 378 (♂) (1906). Mexiko
 (Typus:
opulentialis Druce (nec Lederer, 1863), Teapa
 Biol. Centr.-Amer. Het., 2, p. 272 (1896) (Tabasco))
 (part.) (*Cataclysta*).
aeglesalis Walker, Cat. Lep. Het. Brit. Mus., 19, Brasilien
 p. 952 (♂) (1859). — Hampson, Trans. (Typus: Rio
 Ent. Soc. Lond., p. 151 (1897) (*Cataclysta*); Janeiro)
 Ann. Mag. Nat. Hist., (s. 7) 18, p. 379
 (1906) (*Argyractis*).

- hamiferalis** Hampson, (nec Walker, 1859), Ann. Mag. Nat. Hist., (s. 7) 18, p. 379 (♂, ♀ (1906)). Mexiko, Panama (Typus: Vera Cruz, Atoyac)
- inaurata** Cramer, Pap. Exot., 4, p. 135, t. 359, f. G (1782) (*Cataclysta*). — Hampson, Trans. Ent. Soc. Lond., p. 151 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 380 (1906). Surinam, Brit. Guyana
- aialis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 953 (♂) (1859) (*Cataclysta*). (Rio Janeiro). — Hampson, Trans. Ent. Soc. Lond., p. 150 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 380 (1906) (*Argyractis*). — Druce, Biol. Centr.-Amer. Het., 2, p. 272, t. 63, f. 12 (1896) (*Cataclysta*). *gemmiferalis* Lederer, Wien. Ent. Monatschr., 7, p. 454, 487, t. 18, f. 8 (1863) (*Cataclysta*). — Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1333 (1865) (*Cataclysta*). Zentral-Amerika, Panama, Westindien, Venezuela, Brasilien (Rio Janeiro), Mexiko
- flavivittalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 363 (♂) (1917). Peru (Typus: S. Domingo)
- maronialis** Schaus, Proc. Ent. Soc. Wash., 26, p. 93 (♂) (1924). Franz. Guayana (Typus: St. Jean, Maroni-Fluß)
- cathanal** Schaus, Proc. Ent. Soc. Wash., 26, p. 94 (♂) (1924). Costa Rica (Typus: Guapiles)
- maguilalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 94 (♀) (1924). Franz. Guayana (Typus: St. Jean)
- bedealis** Schaus, Proc. Ent. Soc. Wash., 26, p. 95 (♀) (1924). Brasilien (Typus: Pernambuco)
- aclistalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 288 (♀) (1914) (*Cataclysta*). Panama
- cabimalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 289 (♀) (1914) (*Catalysta*). Panama
- aroalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 95 (♀) (1924). Venezuela (Typus: Aroa)
- amethystina** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, p. 672 (♀) (1912). Costa Rica
- mignonalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 290 (1914) (*Cataclysta*). Panama

- aztecalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 96 Mexiko
(♂) (1924). (Typus: Jalapa)
- parvissimalis** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, p. 673 (1912). Costa Rica
- puralis** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, Costa Rica
p. 674 (♀) (1912).
- tristalis** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, Costa Rica
p. 674 (♀) (1912).
- cineralis** Schaus, Proc. U. S. Nat. Mus., 30, p. 133 Mexiko
(1906).
- triumphalis** Schaus, Ann. Mag. Nat. Hist., (s. 8) Cota Rica,
9, p. 674 (♀) (1912). — Dyar, Proc. U. S. Panama
Nat. Mus., 47, p. 288 (1914) (*Cataclysta*).
- esperanzalis** Schaus, Proc. Ent. Soc. Wash., 26, Costa Rica
p. 96 (♀) (1924). (Typus: Esparanzo)
- guadarensis** Schaus, Proc. U. S. Nat. Mus., 30, Mexiko
p. 133 (1906).
- brunneodora** Dyar, Proc. U. S. Nat. Mus., 47, Panama
p. 289 (1914) (*Cataclysta*).
- herminalis** Schaus, Proc. U. S. Nat. Mus., 30, p. Mexiko
134 (1906).
- jalapalis** Schaus, Proc. U. S. Nat. Mus., 30, p. Mexiko
134 (♀) (1906). — Dyar, Proc. U. S. Nat.
Mus., 47, p. 290 (1914) (*Cataclysta*).
- orizabalalis** Schaus, Proc. U. S. Nat. Mus.,
30, p. 136 (♀) (1906) (*Cataclysta*). — Dyar,
Proc. U. S. Nat. Mus., 47, p. 290 (1914).
- cylolalis** Schaus, Proc. U. S. Nat. Mus., 30, p. 134 Mexiko
(1906).
- sinitalis** Schaus, Proc. U. S. Nat. Mus., 30, p. Mexiko
135 (1906).
- volcanalis** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, Costa Rica
p. 675 (♀) (1912).
- albilbasalis** Schaus, Ann. Mag. Nat. Hist., (s. 8) Costa Rica,
9, p. 670 (♂) (1912). Panama
- auspicatalis** Schaus, Ann. Mag. Nat. Hist., (s. 8) Costa Rica,
9, p. 671 (♀) (1912). — Dyar, Proc. U. S. Panama
Nat. Mus., 47, p. 288 (1914) (*Cataclysta*).
- valstanalis** Schaus, Proc. Ent. Soc. Wash., 26, p. Venezuela
97 (♀) (1924). (Typus: Merida)
- schwarzalis** Schaus, Proc. Ent. Wash., 26, p. 97 Guatemala
(♀) (1924). (Typus: Trece Aguas, Ata Vera Paz)
- chejelalis** Schaus, Proc. Ent. Soc. Wash., 26, p. Guatemala
98 (♀) (1924). (Typus: Chejel)

- tamanalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 99 (♀) (1924). W.Kolumbien
(Typus:
Juntas
del Rio
Tamana &
Rio
San Juan)
- ulfridalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 99 (♂) (1924). Westindien
(Typus:
Dominica)
- pavonialis** Hampson, Trans. Ent. Soc. Lond., p. 150 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 380 (1906) (*Argyractis*). Guadeloupe,
Brasilien,
Peru
(Callao)
- fulvicinctalis** Hampson, Trans. Ent. Soc. Lond., p. 150 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 380 (1906) (*Argyractis*). Jamaika
- axis** Hampson, Ann. Mag. Nat. Hist., (s. 6) 16, p. 343 (1895) (*Cataclysta*); Trans. Ent. Soc. Lond., p. 149 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 380 (1906) (*Argyractis*). Westindien
(N. Grenada)
Peru
- constellalis** Hampson, Trans. Ent. Soc. Lond., p. 151 (♂) (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 380 (1906) (*Argyractis*). Kolumbien
(Typus:
Brasilien,
St. Martha)
- niphoplagalalis** Hampson, Trans. Ent. Soc. Lond., p. 136 (1897). Panama,
Dominica
- catenalis** Guenée, Delt. et Pyral., p. 267 (1854) (*Cataclysta*) (Brasilien). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 440 (♀) (1859). — Lederer, Wien. Ent. Monatschr., 7, p. 454, t. 18, f. 9 (1863) (*Cataclysta*). — Hampson, Trans. Ent. Soc. Lond., p. 149 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 380 (1906) (*Argyractis*). Mexiko,
Brasilien,
Panama
- rinconadalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 100 (♀) (1924). Mexiko
(Typus:
Rinconada)
- lucianalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 100 (♀) (1924). Costa Rica
(Typus:
Juan Vinas)
- chalcistis** Dognin, Ann. Soc. Ent. Belg., 54, p. 118 (1910). Peru
- nitens** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, p. 672 (♀) (1912). Costa Rica
- gontranalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 101 (♀) (1924). Mexiko
(Typus:
Orizaba)
- aengusalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 101 (♂) (1924). Brasilien
(Typus: Rio
de Janeiro)

- alemundalis** Schaus, Proc. Ent. Soc. Wash., 26, Costa Rica
p. 102 (♀) (1924). (Typus:
Peralta)
- serapionalis** Schaus, Proc. Ent. Soc. Wash., 26, Guatemala
p. 102 (♀) (1924). (Typus:
Cayuga)
- plusialis** Herrich-Schaeffer, Ver. Regensb., 25, p. Cuba
26 (1871) [Sep. Lief. 3, p. 66 (1871)]
(*Cataclysta*). — Hampson, Trans. Ent. Soc.
Lond., p. 153 (1897) (*Cataclysta*).
- psalmoidalis** Schaus, Proc. Ent. Soc. Wash., 26, Cuba
p. 103 (♀) (1924).
- leucogonialis** Hampson, Ann. Mag. Nat. Hist., (s. Panama
7) 18, p. 330 (♀) (1906). (Typus:
cerrussalis Druce (nec Felder et Rogen- Chiriqui)
hofer, 1874), Biol. Centr.-Amer. Het., 2,
p. 273 (1896) (part.) (*Cataclysta*).
- cuprescens** Hamps., Ann. Mag. Nat. Hist., (s. 8) Ecuador
17, p. 364 (♀) (1917). (Typus:
Rio Verde)
- productalis** Hampson, Ann. Mag. Nat. Hist., (s. Peru
8) 19, p. 364 (♀) (1917). (Typus:
Carabaya)
- argyrophora** Hampson, Ann. Mag. Nat. Hist., (s. Kolumbien
8) 19, p. 365 (♀) (1917). (Typus:
Condoto Prov.
Choko)
- longipennis** Hampson, Ann. Mag. Nat. Hist. (s. 7) Mexiko
18, p. 381 (♀) (1906) (*Argyractis*). — Barnes (Typus:
et Benjamin, Contr. Nat. Hist. Lep. N. Orizaba,
Amer., 5, p. 190 (1922) (*Cataclysta*). Jalapa),
cronialis Druce, Biol. Centr.-Amer. Het., Arizona
2, p. 274 (1896) (part.) (*Cataclysta*). —
Hampson, Trans. Ent. Soc. Lond., p. 149
(1897) (*Cataclysta*).
- brunneosulfusa** Hampson, Ann. Mag. Nat. Hist., Ecuador
(s. 8) 19, p. 365 (♀) (1917). (Typus:
Banos,
Rio Pastaza)
- annulalis** Guene, Delt. et Pyral., p. 266 (1854) Brasilien,
(*Cataclysta*). — Hampson, Trans. Ent. Soc. Mexiko,
Lond., p. 149 (1897) (*Cataclysta*); Ann. Mag. Guatemala,
Nat. Hist., (s. 7) 18, p. 382 (1906) (*Ar- Honduras,*
gyractis). Costa Rica,
Panama,
Venezuela
- nigrifusalis** Dognin, Hét. Nouv. Amer. du Sud, Kolumbien
fasc. 3, p. 65 (♂) (1911).
- symphonalis** Dyar, Proc. U. S. Nat. Mus., 27, p. Panama
290 (1914) (*Cataclysta*).
- multipecta** Dyar, Proc. U. S. Nat. Mus., 47, p. 291 Panama
(1914) (*Cataclysta*).

- samealis** Felder et Rogenhofer, Reise Novara (Het.), p. 5, t. 136, f. 14 (1874) (*Cataclysta*). Kolumbien
- gratalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1335 (♀) (1865) (*Cataclysta*) (St. Domingo). — Hampson, Trans. Ent. Soc. Lond., p. 149 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 382 (1906) (*Argyractis*). Mexiko, Westindien (Haiti), Guatemala, Costa Rica, Panama, Kolumbien, Brit. Guayana, Brasilien
- cerrussalis** Felder et Rogenhofer, Reise Novara (Het.), p. 5, t. 136, f. 8 (1874) (*Cataclysta*) (Bogotá). — Druce, Biol. Centr.-Amer. Het., 2, p. 273 (1896) (*Cataclysta*).
- schistopalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 382 (1906). Mexiko (Typus: Tabasa, Teapa)
- aialis** Druce (nec Walker, 1859), Biol. Centr.-Amer. Het., 2, p. 272 (1896) (part.) (*Cataclysta*).
- malcusalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 103 (♀) (1924). Westindien (Typus: St. Domingo)
- climacusalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 104 (♀) (1924). Peru (Typus: Tiguri, Carabaya)
- zamoralis** Schaus, Proc. Ent. Soc. Wash., 26, p. 104 (♀) (1924). Ecuador (Typus: Zamora)
- lulesalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 105 (♂) (1924). Argentina (Typus: Luleo, Tucuman)
- alvealis** Felder et Rogenhofer, Reise Novara (Het.), p. 5, t. 136, f. 3 (1874) (*Cataclysta*) (Amazonas). — Hampson, Trans. Ent. Soc. Lond., p. 150 (1897) (*Cataclysta*). — Schaus, Proc. Ent. Soc. Wash., 26, p. 106 (1924) (*Argyractis?*). Brasilien
- laurentialis** Schaus, Proc. Ent. Soc. Wash., 26, p. 105 (♀) (1924). Franz. Guayana (Typus: St. Laurent, Maroni-Fluß)
- castusalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 106 (♀) (1924). Mexiko (Typus: Jalapa)
- tapajosalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 107 (♀) (1924). Brasilien (Typus: Rio Tapajos)
- dodalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 107 (♀) (1924). Brit. Guayana (Typus: Tumatumari-Fluss)

- gordianalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 107 (♀) (1924). Peru, Brasilien (Typus: Rio Puramaya (Peru); Rio Solimoes, Bras.)
- odoalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 108 (♀) (1924). Brasilien (Typus: Rio Teffé, Rio Solimoes)
- benezetalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 109 (♀) (1924). Brasilien (Typus: Teffé, Rio Solimoes)
- conallalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 109 (♀) (1924). Brit. Guayana
- onyxalis** Hampson, Trans. Ent. Soc. Lond., p. 149 (1897) (*Cataclysta*) (nom. nov.); id., Ann. Mag. Nat. Hist., (s. 7) 18, p. 385 (1906) (*Argyraetis*). Mexiko, Guatemala, Westindien, Venezuela
- divulsalis* Druce (nec Walker, 1859), Biol. Centr.-Amer. Het., 2, p. 272, t. 63, f. 13 (1896) (*Cataclysta*).
- pyropalis** Guenée, Delt. et Pyral., p. 265 (1854) (*Cataclysta*). — Hampson, Trans. Ent. Soc. Lond., p. 153 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 385 (1906) (*Argyraetis*). Brasilien
- caesoalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 952 (♂) (1859) (*Cataclysta*). Brasilien
- coesalis* Hampson, Trans. Ent. Soc. Lond., p. 149 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 385 (1906) (*Argyraetis*).
- insulalis** Walker, Trans. Ent. Soc. Lond., (s. 3) 1, p. 123 (1862) (*Cataclysta*). — Hampson, Trans. Ent. Soc., Lond., p. 150 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 385 (1906) (*Argyraetis*). Haiti, Brasilien
- pantheralis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 442 (♀) (1859) (*Cataclysta*). — Hampson, Trans. Ent. Soc. Lond., p. 151 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 385 (1906) (*Argyraetis*). Brasilien
- premalis** Druce, Biol. Centr.-Amer. Het., 2, p. 272, t. 63, f. 11 (1896) (*Cataclysta*). — Hampson, Trans. Ent. Soc. Lond., p. 151 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist. (s. 7) 18, p. 385 (1906) (*Argyraetis*). Mexiko, (Typus: Teapa in Tabasco), Brasilien, Argentina
- albulalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 385 (♂) (1906). Jamaika (Typus: Runaway Bay)

- subornata** Hampson, Trans. Ent. Soc. Lond., p. 151 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 386 (1906) (*Argyractis*). Brasilien (Typus: São Paulo), Argentinien
- harpalis** Snellen, Tijds. v. Ent., 43, p. 296, t. 17, f. 1 (1901) (*Cataclysta*). — Hampson, Ann. & Mag. Nat. Hist., (s. 7) 18, p. 386 (1906). — Dyar, Proc. U. S. Nat. Mus., 47, p. 287 (1914) (*Argyractis*). — Forbes, Ent. News, 33, p. 102 (1922) (subg. *Oxyelophila*). Mexiko, Guatemala, Costa Rica, Panama
- endoralis* Druce (nec Walker, 1859), Biol. Centr.-Amer. Het., 2, p. 275 (1896) (part.) (*Hydrocampa*).
- necomalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 287 (♂) (1914). — Forbes, Ent. News, 33, p. 102 (1922) (subg. *Oxyelophila*). Panama
- ticonalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 287 (♀) (1914). — Forbes, Ent. News, 33, p. 102 (1922) (subg. *Oxyelophila*). Panama
- lanceolalis** Hampson, Trans. Ent. Soc. Lond., p. 140 (1897) (*Nymphula*). — Forbes, Ent. News, 33, p. 102 (1922) (subg. *Oxyelophila*). Brasilien (Typus: Espiritu Santo)
- melanograpta** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 367 (♂) (1917). Brit. Guayana (Typus: Demerara)
- micropalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 386 (♂) (1906). Mexiko (Typus: Teapa)
- endoralis* Druce (nec Walker, 1859), Biol. Centr.-Amer. Het., 2, p. 275 (1896) (part.) (*Cataclysta*). (Tabasco)
- argenteolinealis** Hampson, Trans. Ent. Soc. Lond., p. 136; p. 135 (fig. ♂) (1897). Brasilien (Typus: Espiritu Santo)
- phaeopastalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 368 (♀) (1917). Kolumbien (Typus: Choko)
- parthenodalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 387 (♂) (1906). Argentinien (Typus: Tucuman (Los Vasquez))
- iasusalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 951 (♀) (1859) (*Oligostigma*?). — Hampson, Trans. Ent. Soc., p. 136 (1897); Ann. Mag. Nat. Hist., (s. 7) 18, p. 387 (1906) (*Argyractis*). Brasilien
- phoxopteralis* Snellen, Tijds. v. Ent., 43, p. 295, t. 16, f. 11, 12 (1901) (*Cataclysta*).
- metazonalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 388 (♂) (1906). Brasilien (S. Paulo)

Species incertae sedis.

- decoralis** Dognin, Ann. Soc. Ent. Belg., 49, p. 65 Ecuador (1905).
leucistis Dognin, Ann. Soc. Ent. Belg., 50, p. 184 Ecuador (♂) (1906).
tessimalis Dyar, Insec. Insc. Menstr., 14, p. 148 Mexiko (♂) (1926). (Typus: Chiapas, Tabasco)
trespasalis Dyar, Insec. Insc. Menstr., 14, p. 147 Mexiko (♀) (1926). (Typus: Chiapas, Tabasco)
 127

3. Genus Eristena Warren

Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 149 (1896). — Hampson, Fauna Brit. India Moths, 4, p. 190 (1896); Trans. Ent. Soc. Lond., p. 132, 136 (1897).

(Typus generis: *E. postalbalis* Hampson).

- postalbalis** Hampson, Ill. Typ Sp. Lep. Het. Brit. Ceylon Mus., 9, p. 175, t. 174, f. 3 (1891); Fauna Brit. India Moths, 4, p. 191, f. 114 (♀) (1891); Trans. Ent. Soc. Lond., p. 137, fig. (♀) (1897).
tenebrifera Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 368 (♂) (1917). Holl. Neu-Guinea (Typus: Mikika-River)
tetralitha Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 368 (♀) (1917). Holl. u. Brit. Neu-Guinea (Typus: Mikika-River)
murinalis Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 150 (1896). — Hampson, Fauna Brit. India Moths, 4, p. 191 (1896); Trans. Ent. Soc. Lond., p. 137 (1897). Assam (Khasis)
oligostigmatalis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 389 (♂, ♀) (1906); Journ. Bombay Nat. Hist. Soc., 18, p. 572 (1908). Andamanen
straminealis Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 209 (♂) (1903). Sikkim
trigonalis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 390 (♂, ♀) (1906). Neu-Guinea (Typus: Kapaur)

Species incertae sedis.

- grisealis** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 148 (♂) (1915). Holl. Neu-Guinea
melanolitha Turner, Trans. Roy. Soc. S. Austral., 32, p. 81 (♂) (1908). Queensland

4. Genus **Arxama** Walker

Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1183 (♀) (1865). — Hampson, Trans. Ent. Soc. Lond., p. 132, 137 (1897).

(Typus generis: *subcervinalis* Walker).

monetalis Snellen, Tijds. v. Ent., 23, p. 244 (♀) Pulo Laut,
(1880) (*Cymoriza*); 27, p. 49, t. 5, f. 3 Celebes
(1884) (*Cymoriza*). — Hampson, Trans. Ent.
Soc. Lond., p. 138 (1897).

atralis Hampson, Trans. Ent. Soc. Lond., p. 138 Malakka
(1897). (Typus: *Gunony Ijou*)

subcervinalis Walker, Cat. Lep., Het. Brit. Mus., Ceram
34, p. 1183 (♀) (1865). — Hampson, Trans.
Ent. Soc. Lond., p. 138, p. 137, fig. (♂)
(1897). — Swinhoe, Cat. East. Austral. Lep.
Het., 2, p. 437, t. 6, f. 11 (1900).

ochrocealis Hampson, Ann. Mag. Nat. Hist., (s. 7) N. Borneo
18, p. 390 (♀) (1906).

cretacealis Hampson, Ann. Mag. Nat. Hist., (s. 7) Salomon Ins.
18, p. 390 (♂) (1906).

5

5. Genus **Nymphula** Schrank

v. Schrank, Fauna Boica, 2 (2), p. 162 (1802). — Hübner, Verz.
bek. Schmett., p. 362 n. 3465 (1816) [1827] (typus generis:
N. potamogalis). — Treitschke, Schm. Eur., 7, p. 107 (1829). —
Duponchel, Hist. Nat. Léop. Fr., 8 (2), p. 158 (1831). — Herrich-
Schaeffer, Schm. Eur., 4, p. 10 (1848). — Moore, Lep. Ceylon,
3, p. 304 (1885). — Meyrick, Trans. Ent. Soc. Lond., p. 465
(1890); Handb. Brit. Lep., p. 402 (1895). — Hampson, Fauna Brit.
India Moths, 4, p. 191 (1896); Trans. Ent. Soc. Lond., p. 132, 138
(1897). — Pagenstecher, Zoologica, H. 29, p. 173 (1900). — Hering,
Stettin. Ent. Zeit., 64, p. 61 (1903). — Dyar, Journ. N. York Ent.
Soc., 14, p. 78, (1906). — Rebel in Spuler, Schm. Eur., 2, p. 221
(1906). — Ulmer, Uns. Wass. ins., p. 79, f. 60 (larva), p. 80, f. 62
(larva), f. 63 (pes larvae) (1911). — Meyrick, Trans. Proc. N.
Zeal. Inst., 45, p. 38 (1913). — Wahlgren, Svensk Insektafauna,
10 (2), p. 46 (19115). — Tullgren & Wahlgren, Svenska Ins.,
p. 388, 394 (1922). — Forbes, Lep. N. York, p. 578 (1928). —
Hudson, Butt. Moths N. Zeal., p. 177 (1928). — Meyrick, Revis.
Handb. Brit. Lep., p. 418 (1928). — Shibuya, Journ. Fac. Agr.
Hokkaido Imp. Univ., 22, p. 139, 140 (1928); Ins. Matsum., 3,
p. 123 (1929). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 281
(1932). — Eckstein, Schm. Deutschl., 5, p. 62 (1933).

Hydrocampa Latreille, Fam. Nat. Règne Anim., p. 478
(1825) (typus: *nymphacuta* L.). — Guenée, Delt. et Pyral.,
p. 273 (1854). — Lederer, Wien. Ent. Monatschr., 7, p. 451
(1863). — Heinemann, Kloinschm., 1, H. 2, p. 102 (1865).
— Wallengren, Oefvers. K. Vet.-Akad. Förh., 28 (8),
p. 1005 (1871). — Snellen in Veth, Midden Sumatra, 4,
p. 75 (1880). — Meyrick, Trans. Ent. Soc., p. 465 (1890);
Handb. Brit. Lep., p. 403 (1895). — Cannaviello, Bol.
Soc. Esp. Nat., 2, p. 164 (1902). — Barrett, Lep. Brit.
Isl., 9, p. 290 (1904). — Tullgren & Wahlgren, Svenska

- Ins., p. 388, 394 (1922). — Meyrick, Revis. Handb. Brit. Lep., p. 419 (1928).
- Hydrocampus* Berthold, Letr. Nat. Fam. Tierr., p. 485 (1827).
- Paraponyx* Hübner, Verz. bek. Schmett., 5, p. 362 (1826) (part.; sine fixatione typi generis). — Stephens, Ill. Brit. Ent. Haust., 4, p. 41 (1834). — Lederer, Wien. Ent. Monatschr., 7, p. 452 (1863). — Wallengren, Oefvers. K. Vet.-Akad. Förh., 28 (8), p. 1007 (1871). — Snellen, in Veth, Midden-Sumatra, 4, p. 76 (1880); Vlinders Nederl., 2 (1), p. 76 (1882). — Moore, Lep. Ceylon, 3, p. 301 (1885).
- Paraponyx* Guenée, Delt. et Pyral., p. 268 (1854). — Walker, List Lep. Het. Brit. Mus., 17, p. 452 (1859). — Stainton, Man. Brit. Butterfl. Moths, 2, p. 144 (1859). — Heinemann, Kleinschm., 1, H. 2, p. 105 (1865). — Barrett, Lep. Brit. Isl., 9, p. 287 (1904). — Wahlgren, Svensk Insektfauna, 10 (2), p. 47 (1915). — Tullgren & Wahlgren, Svenska Ins., p. 388, 394 (1922). — M. Hering, Tierwelt Mitteleur., (Ergbd. 1), p. 282 (1932). — Eckstein, Schm. Deutschl., 5, p. 63 (1933).
- Synclita* Lederer, Wien. Ent. Monatschr., 7, p. 448 (1863) (typus: *gurgitalis* Led.).
- Nymphacella* Grote, Nov. Amer. Ent., 1, p. 97 (1880) (typus: *maculalis* Clem.).
- Hygraula* Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 129 (1885) (typus: *nitens* Butl.).
- Hydreuretis* Meyrick, Trans. Ent. Soc. Lond., p. 435 (1885) (typus: *tullialis* Walk.).
- Metoecca* Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 145 (1896) (typus: *foedalis* Guenée).
- Nausinoë* (part.) Hübner, Verz. bek. Schmett., p. 363 n. 3466 (1816) [1826] (*N. nymphaealis* Schiff.).
- Kasania* Krulikovsky, Rev. Russe Ent., 9, p. 316 (1909) (typus: *arundinalis* Eversm.).
- Oligostigma* Walker, Cat. Lep. Het. Brit. Mus., 17, p. 429 gen. 4 (1859). — Hampson, Trans. Ent. Soc. London, p. 167 (1897) (Typus: *O. junceale*). — Dyar, Journ. N. York Ent. Soc., 14, p. 88 (1906).
- Biologie: Portier, Arch. Zool. Expér., (s. 5) 8, p. 89—379 (1912). — Weleh, Ann. Ent. Soc. Amer., 9, p. 159—190, t. 7—9 (1916). — Efflatoun, Bull. Soc. Ent. Egypte, 6, p. 29 (1928). — Welsh, Trans. Amer. Microsc. Soc., 41, p. 29, figs. (1922) (Respir. system d. Larven). — Forbes, Lep. N. York, p. 576 (1923) (tab. anal. larv.).
- (Typus generis: *nymphacata* L.)

a) Paläarktische Region.

- rivularis** Duponchel, Hist. Nat. Lép., 8 (2), p. 341, Süd-
t. 233, f. 5 (1831). — Zeller, Stettin. Ent. Frankreich,
Zeit., 10, p. 233 (1849). — Herrich-Schäffer, N. O.
Syst. Bearb. Schm. Eur., 4, p. 11 (1848). Deutschland,
— Guenée, Delt. et Pyral., p. 275 (1854) Nieder-
(*Hydrocampus*). — Heinemann, Kleinschm., Österreich,
1, H. 2, p. 104 (1865) (*Hydrocampus*). — N. Italien,
— Berce, Faune Ent. Fr., 6, p. 121 (1878). Korsika,
— Hampson, Trans. Ent. Soc. Lond., p. 139 Griechenland,
(1897). — Cannaviello, Bol. Soc. Esp. Hist. land,
Nat., 2, p. 171 (1902). — Rebel in Spuler, Belgien,
Schm. Eur., 2, p. 221 (1906). — Wahlgren, Schweden,
Svensk Insektfauna, 10 (2), p. 46, t. 2, Norwegen

f. 49 (1915). — M. Hering, Tierwelt Mitteleur. (Ergbd., 1), p. 282 (1932). — Eckstein, Schm. Deutschl., 5, p. 62 (1933).

- stagnata** Donovan, Nat. Hist. Brit. Ins., 11, p. 10, t. 363, f. 3 (1906) (*Phaena*). — Guenée, Delt. et Pyral., p. 276 (1854) (*Hydrocampa*). — Stainton, Man. Brit. Butt. Moths, 2, p. 145 (1859) (*Hydrocampa*). — Lederer, Wien. Ent. Monatschr., 7, p. 451 (1863) (*Hydrocampa*). — Heinemann, Kleinschm., 1, H. 2, p. 103 (1865) (*Hydrocampa*). — Berce, Faune Ent. Fr. Lép., 6, p. 120 (1878). — Snellen, Vlinders Nederl., 2, (1), p. 75 (1882) (*Parapynx*). — Reuter, Acta Soc. Faun. Flor. Fennica, 15 (5), p. 17 (1899) (*Hydrocampa*). — Meyrick, Trans. Ent. Soc. Lond., p. 466 (1890); Handb. Brit. Lep., p. 402 (1895). — Hampson, Trans. Ent. Soc. Lond., p. 139 (1897). — Rebel in Spuler, Schm. Eur., 2, p. 221, t. 81, f. 17 (1906). — Wahlgren, Svensk Insektfauna, 10 (2), p. 46, t. 2, f. 50 (1915). — Tullgren & Wahlgren, Svenska Insekt., p. 394 (1922). — Meyrick, Revis. Handb. Brit. Lep., p. 419 (1928). — Shibuya, Ins. Matsum., 3, p. 121, 124, t. 5, f. 1 (1929). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 282, f. 501 (1932). — Eckstein, Schm. Deutschl., 5, p. 62, t. 1, f. 19 (1933). — Karny, Biol. Wasserins., p. 167 (1934).
- stagnalis* Guenée, Delt. et Pyral., p. 276 (1854) (*Hydrocampa*). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 458 (1858) (*Hydrocampa*). — Barrett, Lep. Brit. Isl., 9, p. 294, t. 408, f. 3 a—e (1904) (*Hydrocampa*).
- potamogalis* Hübner, Samml. Auserl. Vög. Schm., p. 28 (1793); Samml. Eur. Schm., 6. Horde, p. 19 (*Pyral.*), t. 13, f. 82 (1796) (*Pyralis*); id., Verz. bek. Schmett., p. 362 n. 3465 [1816] (1826) (*Nymphula*). — Schrank, Fauna Boica, 2 (2), p. 62 (1802).
- nymphaealis* Treitschke, Schm. Eur., 7, p. 141 (1829). — Duponchel, Hist. Nat. Lép., 8 (2), p. 177, t. 222, f. 3 (1831) (*nymphalis*). — Zetterstedt, Ins. Lapp., p. 973 (1839). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 11 (1848). — Wallengren, Oefvers. K. Vet.-Akad. Förh., 28 (8), p. 1006 (1871).
- Biologie: Buckler, Ent. Mon. Mag., 16, p. 7 (1877); Larvae Brit. Lep., 9, p. 101, t. 152, f. 1 (1901).
- nota: South, Malayan Agr. Journ., 14 (6), p. 146—159 (1926).
- nymphaeata** Linné, Syst. Nat. (ed. 10), p. 529 (♂) (1758) (*Phalaena Geometra*); Fauna Suec. ed. 2, p. 340 (1761) (*Geometra*). — Fabricius, Ent. Syst., 3 (2), p. 316 (1794) (*Phalaena*). — Stephens, Ill. Brit. Ent. Mitteleuropa, Nordeuropa, (N. Spanien, Süd-Frankreich, Belgien, Italien, Schwyz, Dalmatien, Bosnien, Herzegowina, Rumänien, England, Finnland, Schweden, Lappland), Sibirien, Uralsk, S. Ural, Amur, Kleinasien, Armenien, Japan
- Mitteleuropa, Livland, Estland, Finnland, Südskan-

- Haust., 4, p. 39 (1834). — Wood, Index Entom., p. 123, t. 28, f. 800 (1839) (*Hydrocampa*). — Westwood et Humphrey, Brit. Moths, 2, p. 97, t. 76, f. 21—23 (1843) (*Hydrocampa*). — Stainton, Man. Brit. Butt. Moths, 2, p. 145 (1859) (*Hydrocampa*). — Lederer, Wien. Ent. Monatschr., 7, p. 451 (1863). — Heinemann, Kleinschm., 1, H. 2, p. 104 (1865) (*Hydrocampa*). — Wallengren, Oefvers. K. Vet.-Akad. Förh., 28 (8), p. 1006 (1871). — Berce, Faune Ent. Fr. Lép., 6, p. 118, t. 3, f. 13 (1878). — Snellen, Vlinders Nederl., 2 (1), p. 74 (1882) (*Hydrocampa*). — Meyrick, Handb. Brit. Lep., p. 403 (1895). — Hampson, Trans. Ent. Soc. Lond., p. 139 (1897). — Sharp in Harmer & Shipley, Cambridge Nat. Hist., 6 (Ins. 2), p. 421, 423 (1899) (*Hydrocampa*). — Cannaviello, Bol. Esp. Soc. Nat. Hist., 2, p. 169 (1902). — Rebel in Spuler, Schm. Eur., 2, p. 221, t. 81, f. 18 (1906). — Wesenberg-Lund, Ins.-liv. ferske Vand., p. 202, fig. 161 (imago), p. 205, f. 163 A, B, C (larva), p. 206, 207, 439 (1915) (*Hydrocampa*). — Tullgren & Wahlgren, Svenska Ins., p. 394, f. 315 g (1922) (*Hydrocampa*). — Börner in Brohmer, Fauna Deutschl., (ed. 3), p. 377 (1925). — Wahlgren, Svensk Insektfauna, 10 (2), p. 47, t. 2, f. 51 (1915). — M. Hering, Okol.blattmin. Ins.-larven, p. 179 (1926) (*Hydrocampa*). — Meyrick, Revis. Handb. Brit. Lep., p. 420 (1928). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 282 (1932). — Eckstein, Schm. Deutschl., 5, p. 62, t. 1, f. 20 (1933). — Karny, Biol. Wasserins., p. 160, f. 103 A (larva), p. 167, 174, f. 108 B (larva) (1934). — Börner in Brohmer, Fauna Deutschl., (ed. 4), p. 389 (1932). — Amsel, Mitt. Zool. Mus. Berlin, 20 (2), p. 288, t. 9, f. 23 (1935).
- nymphacalis* (Schiffermüller et Denis), Verz. Schm. Wien, p. 121 (1776) (*Pyralis*). — Hübner, Samml. Eur. Schm., 6. Horde (*Pyral.*), t. 13, f. 83 (1796) (*Pyralis*). — Schrank, Fauna Boica, 2 (2), p. 63 (1802). — Hübner, Verz. bek. Schmett., p. 363 n. 3467 [1816] (1826). (*Nausinoë*). — Treitschke, Schm. Eur., 7, p. 141, (1829). — Guenée, Delt. et Pyral., p. 275 (1854) (*Hydrocampa*). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 457 (1858) (*Hydrocampa*). — Barrett, Lep. Brit. Isl., 9, p. 291, t. 408, f. 2 a—e (1904) (*Hydrocampa*).
- potamogata* Linné, Syst. Nat. ed. 10, p. 529 (♀) (1758) (*Geometra*). — Donovan, Nat. Hist. Brit. Ins., 9, t. 363, f. 1 (1806) (*Phalaena*). — Stephens, Ill. Brit. Ent. Haust., 4, p. 38 (1834) (*Hydrocampa*). —
- dinavien,
England,
Irland,
N. Spanien,
Italien,
Südrussland,
Ciskaukasien,
Kleinasien,
Palästina,
N.O. Sibirien.
Bucharä,
Mongolei,
Sarawtschan,
Amur,
Ussuri,
Ostchina,
Mauretanien,
Balearn

- Wood, Index Entom., p. 122, t. 28, f. 799 (1839) (*Hydrocampa*).
- potamogalis* Treitschke, Schm. Eur., 7, p. 143 (1829). — Duponchel, Hist. Nat. Léop., 8 (2), p. 172, t. 222, f. 1, 2 (1831) (*Hydrocampa*). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 11 (1848).
- rivulata* Scopoli, Ent. Carn., p. 573 (1763) (*Hydrocampa*).
- tripunctalis* O. G. Costa, Fauna Reg. Napoli Lep. Nott., p. 3, t. 4, f. 1 (1832). — Cannaviello, Bol. Soc. Esp. Nat. Hist., 2, p. 170 (1902).
- lunulalis* O. G. Costa, Fauna Reg. Napoli Lep. Nott., p. 3, t. 4, f. 3 (1832). — Cannaviello, Bol. Soc. Esp. Nat. Hist., 2, p. 171 (1902).
- transversalis* O. G. Costa, Fauna Reg. Napoli Lep. Nott., p. 4, t. 7, f. 5 (non 2) (1832). — Cannaviello, Bol. Soc. Esp. Nat. Hist., 2, p. 171 (1902).
- fa. *umbrata* Meder, Schrift. Naturw. Ver. Schleswig-Holstein, 20, p. 359 (1934). Ins. Amrum
- subsp. *latifasciata* W. Rothschild, Journ. Bombay Nat. Hist. Soc., 28, p. 178 (♀) (1921). Persien
- subsp. *benesignata* Caradja, Mem. Acad. Rom., (s. 3) 3, mem. 7, p. 328 (1925). Ost- u. Mittelchina
- Anatomie: Stitz, Zool. Jahrb. Anat., 14, p. 143—147 (1900).
- Biologie: Buckler, Ent. Mon. Mag., 12, p. 210 (1876). — Sorhagen, Berl. Ent. Zeitschr., 26, p. 150 (1882). — Haye, Proc. Ent. Soc. Lond., p. XVIII, (1900). — Buckler, Larvae Brit. Lep., 9, p. 85, t. 151, f. 3 (1901). — Sorhagen, Allg. Zeitschr. Ent., 6, p. 277 (1901). — Rebel in Gmelin, Schm. Eur. Nachtr. Taf. 9, f. 52 (1906). — Ege, Vidensk. Meddel., 76, p. 29—42 (1923) (Respir.-syst. larv. et pup.); Physiol. paprs. dedic. A. Krogh, p. 25—39, fig. (1926).
- noxia: Supino, Rendic. R. Inst. Lombardo Scienze e Lettere (s. 2) 49, nr. 2, 3, p. 108—114 (1916).
- hederalis** Amsel, Mitt. Zool. Mus. Berlin, 20 (2), p. 288, t. 9, f. 24 (1935). Palästina
- auralis* Osthelder, Mitt. Münch. Ent. Ges., 24, p. 94 (1935) (*nymphacata* subsp.).
- interruptalis** Pryer, Cist. Ent., 2, p. 233, t. 4, (1877) (*Hydrocampa*); id., Trans. Asiat. Soc. Jap., 13, p. 62 (1885) (*Hydrocampa*). — Hampson, Trans. Ent. Soc. Lond., p. 139 (1897). — Leech, Trans. Ent. Soc. Lond., p. 432 (1901). — Wileman, Trans. Ent. Soc. Lond., p. 370 (1911). — Shibuya, Ins. Matsum., 3, p. 124, t. 5, f. 2 (1929). — Matsumura, 6000 Ill. Ins. Japan, p. 1043, fig. (♀) (1931). West-, Mittel-, Ost- u. Süd-China, Ussuri, Korea, Japan

- subsp. *separatalis* Leech, Entom., 22, p. 71, t. 4, f. 2, 13 (1889) (*Hydrocampa interruptalis* var.). — Shibuya, Ins. Matsum., 3, p. 125 (1929).
 Biologie: Kobayashi, Insect World, 31, p. 334—338, fig. (1927).
- corculina** Butler, Ill. Typ. Sp. Het. Brit. Mus., 3, p. 75, t. 59, f. 7 (1879) (*Oligostigma*). — Pryer, Trans. Asiat. Soc. Jap., 13, p. 61 (1885) (*Oligostigma*). — Hampson, Trans. Ent. Soc. Lond., p. 167 (1897) (*Oligostigma*). — Leech, Trans. Ent. Soc. Lond., p. 436 (1901) (*Oligostigma*). — Suzuki, List Spec. Hanazono Ent. Labr., p. 25 (1915). — Matsumura, Journ. Coll. Agr. Hokkaido Imp. Univ., 15, p. 187 (1925) (*Oligostigma*). — Shibuya, Ins. Matsum., 3, p. 125, t. 5, f. 3 (1929). — Matsumura, 6000 Ill. Ins. Japan, p. 1043, fig. (♂) (1931).
- distinctalis* Ragonot, Ann. Soc. Ent. Fr., 63, p. 174 (1894) (*Oligostigma stagnata* var.). — Staudinger, Deutsche Ent. Zeitschr. Iris, 10, p. 343 nota (1897). — Snellen, Tijdschr. v. Ent., 43, p. 286 (1901).
- nawalis* Wileman, Trans. Ent. Soc. Lond., p. 373 (1911) (*Aulacodes*). — Shibuya, Ins. Matsum., 3, p. 125 (1929) (= *N. curculina*).
- bifurcalis** Wileman, Trans. Ent. Soc. Lond., p. 373, t. 31, f. 23 (♂) (1911) (*Parthenodes*). — Shibuya, Ins. Matsum., 3, p. 124, 125, t. 5, f. 4 (1929). — Matsumura, 6000 Ill. Ins. Japan, p. 1043, fig. (♂) (1931).
- fengwhanal** Pryer, Cist. Ent., 2, p. 235, t. 4, f. 11 (1877) (*Lepyrodes*). — Hampson, Trans. Ent. Soc. Lond., p. 140 (1897). — Leech, Trans. Ent. Soc. Lond., p. 432 (1901). — Shibuya, Ins. Matsum., 3, p. 124, 126, t. 5, f. 5 (1929). — Caradja, Bull. Acad. Rom., 15, p. 147 (1932). — Matsumura, 6000 Ill. Ins. Japan, p. 1043, fig. (♂) (1931).
- foedalis** Guenée, Delt. et Pyral., p. 228, t. 4, f. 7 (1854) (*Isopteryx*). (Indien). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 402 (1859) (*Isopteryx*) (Hindustan). — Lederer, Wien. Ent. Monatschr., 7, p. 449 (1863) (*Isopteryx*). — Snellen, Tijds. v. Ent., 15, p. 96 (1872) (*Isopteryx*); id., ibid., 27, p. 47 (1884) (*Isopteryx*). — Moore, Lep. Ceylon, 3, p. 306 (1886) (*Isopteryx*). — Ribbe, Deutsche Ent. Zeitschr. Iris, 1, p. 89 (1886) (*Isopteryx*). — Pagenstecher, Jahrb. Nassau. Ver. Naturk., p. 175 (1886). — Meyrick, Trans. Ent. Soc. Lond., p. 229 (1887) (*Pterygismus*); p. 470 (1894) (*Pterygismus*). — Snellen, Trans. Ent. Soc. Lond., p. 638 (1890) (*Pterygismus*). — Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 145 (1896) (*Metoeca*). — Hampson, Fauna Brit. India Moths, 4,
- Korea,
Japan
- Amur,
Mongolei,
Japan,
Sachalin
- Korea,
Japan
- Nord-,
Mittel- u.
Ost-China,
Japan,
Korea
- Japan,
Orientalische
Region
(Indien,
Ostchina,
Tonkin,
Ceylon,
Celebes,
Java,
Borneo,
Sumbawa,
Pulolaut,
Malakka,
Philippinen),
Australische
Region
(Australien,
Neu-Guinea,

- p. 192 (1896); id., Trans. Ent. Soc. Lond., p. 140 (1897). — Pagenstecher, Zoologica, H. 29, p. 172 (1900). — E. Hering, Stett. Ent. Zeit., 64, p. 64 (1903) (*Pterygisus*). — Leech, Trans. Ent. Soc. Lond., p. 433 (1911). — Strand, Ent. Mitt., 8, p. 60 (1919). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 141, 142 (1928); Ins. Matsum., 3, p. 124, 125 (1929).
- tencellalis* Guenée, Delt. et Pyral., p. 228 (1854) (*Isopteryx*) (Cayenne). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 401 (1859) (*Isopteryx*). — Lederer, Wien. Ent. Monatschr., 7, p. 449 (1863) (*Isopteryx*).
- leucothoalis* Walker, Cat. Lep. Het. Brit. Mus., 17, p. 400 (♀) (1859) (*Isopteryx*?) (Vereinigte Staaten).
- oethonalis* Walker, Cat. Lep. Het. Brit. Mus., 17, p. 484 (♀) (1859) (*Zebronia*) (Sarawak).
- scitalis* Lederer, Wien. Ent. Monatschr., 7, p. 451, 483 (1863) (*Hydrocampa*) (Amboina, Java).
- solitalis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1340 (1865) (*Hydrocampa*) (Java, Amboina).
- epispila* Meyrick, Trans. Ent. Soc. Lond., p. 257 (1886) (*Physematia*?) (Fiji).
- forma *cantonialis* Caradja, Acad. Rom. Mem. Sect. Stiint., (s. 3) 3, mem. 7, p. 328, t. 1, f. 31 (1925).
- affinialis** Guenée, Delt. et Pyral., p. 270 (1854) (*Paraponyx*). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 454 (1859) (*Paraponyx*). — Lederer, Wien. Ent. Monatschr., 7, p. 452 (1863) (*Paraponyx*). — Butler, Proc. Zool. Soc. Lond., p. 167 (1885) (*Paraponyx*). — Swinhoe, Proc. Zool. Soc. Lond., p. 524 (1884); p. 868 (1885); p. 458 (1886) (*Paraponyx*). — Hampson, Fauna Brit. India Moths, 4, p. 193 (1896); Trans. Ent. Soc. Lond., p. 141 (1897). — Rebel, Bull. Soc. Ent. Egypte, 19, p. 186 (1926).
- incommoda* Butler, Proc. Zool. Soc. Lond., p. 180 (1881) (*Oligostigma*) (Sokotra).
- fusco-marginata* Bethune-Baker, Trans. Ent. Soc. Lond., p. 48, t. 1, f. 17 (1894) (Aegypten). — Rebel, Deutsche Ent. Zeitschr. Iris, 26, p. 81 (1912). — Zerny, Ann. Naturh. Hofmus., 18, p. 323 (1914). — Caradja, Deutsche Ent. Zeitschr. Iris, 30, p. 21 (1917). — Rebel, Bull. Soc. Roy. Ent. Egypte, 10, p. 186 (1926).
- var. *alaicalis* Caradja, Deutsche Ent. Zeitschr. Iris, 30, p. 21 (1917).
- turbata** Butler, Trans. Ent. Soc. Lond., p. 586 (1881) (*Paraponyx*). — Pryer, Trans. Asiatic Soc. Jap., 13, p. 61 (1885) (*Paraponyx*). — Meyrick, Trans. Ent. Soc. Lond., p. 10, 470 (1894). — Hampson, Fauna Brit.

Samoa,
Fiji Ins.,
Aru Ins.,
Bismarck-
Archipel),
Aethiopische
Region
(Afrika,
Angola,
Madagaskar),
Nearktische
Region
(Vereinigte
Staaten),
Neotropische
Region
(Panama,
Br. u. Fr.
Guayana,
Brasilien)

Canton

Syrien,
Palästina,
Aegypten,
Indien,
Malakka,
Sokotra,
Aru Ins.

Alai

Amur,
Ussuri,
West-,
Zentral-u.
Süd-China,

- India Moths, 4, p. 192 (1896) (part.); Trans. Ent. Soc. Lond., p. 141 (1897). — Leech, Trans. Ent. Soc. Lond., p. 433 (1901). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 141, 143 (1928); Ins. Matsum., 3, p. 124, 126, t. 5, f. 6 (1929). — Matsu-mura, 6000 Ill. Ins. Jap., p. 1044, fig. (♂) (1931).
- floralis* Leech, Entom., 22, p. 71, (♂), t. 4, f. 1 (1889) (*Leparodes*); Trans. Ent. Soc. Lond., p. 433, (1901). (Typus: ♀ Tsuruga & Fushiki)
- sultschana* Ragonot, Ann. Soc. Ent. Fr., 63, p. 174 (1894) (*Hydrocampa* (*Nymphula*)). — Staudinger et Rebel, Cat. Lep. Pal. Faun., 2, p. 49, nr. 915 (1901) (N.). — Zerny, Ann. Naturh. Hofmus. Wien, 28, p. 324 (1914) (*Parthenodes*). — Chrétien, Ann. Soc. Ent. Fr., 85, p. 448 (1917). — Caradja, Deutsche Ent. Zeitschr. Iris, 30, p. 20 (1917) (*Nymphula*); id., Mem. Acad. Rom. Sect. Stiint., (s. 3) 3, mem. 7, p. 333 (1925) (*Parthenodes*).
- sultschana* Hampson, Trans. Ent. Soc. Lond., p. 384 (♀) (1900) (*Parthenodes*). — Leech, Trans. Ent. Soc. Lond., p. 433 (1901) (= *Nymphula turbata* Butl.). — Wileman, Trans. Ent. Soc. Lond., p. 374 (1911) (= *Nymphula turbata* Butl.). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 143 (1928) (= *Nymphula turbata* Butl.). (Typus: ♀ Amurland, Sutschan)
- Biologie: Fletcher, Sci. Repts. Agric. Res. Inst. Pusa 1921—22, p. 51—67 (1922) (Schädling).
- enixalis* Swinhoe, Proc. Zool. Soc. Lond., p. 869 (♀) (1885) (*Isopteryx*) (Bombay). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 141, 143, t. 6, f. 3 (1928); Ins. Matsum., 3, p. 124, 126, t. 5, f. 7 (1929). (Typus: ♀ Ostchina, Indien, Japan, Formosa)
- linealis* Moore, Deser. Ind. Lep. Atkinson, p. 210 (♂) (1888) (*Cymoriza*) (Calcutta).
- nigralbalis* Caradja, Mem. Acad. Rom. Sect. Stiint., (s. 3) 3, mem. 7, p. 329 (1925) (*diffflualis* subsp.); Bull. Acad. Roum., 15, p. 147 (1932); Deutsche Ent. Zeitschr. Iris, 47, p. 153 (1934) (= *enixalis* Swinh.).
- stratiotata* Linné, Syst. Nat. ed. 10, p. 529 (1758) (*Phalaena Geometra*); Fauna Suecica, 2. ed., 4, p. 341 (1761) (*Geometra*); Syst. Nat., ed. 12, 1 (2), p. 873 (1767) (*Hydrocampa*). — Curtis, Brit. Ent., 11, t. 495 (1834) (*Hydrocampa*). — Stephens, Ill. Brit. Ent. Haust., 4, p. 41 (1834) (*Paraponyx*). — Wood, Index Entom., p. 123, f. 802 (1839) (*Hydrocampa*). — Stainton, Man. Brit. Butt. Moths, 2, p. 144 (1859) (*Paraponyx*). — Heinemann, Kleinschm., 1, H. 2, p. 106 (1865) (*Paraponyx*). — Lederer, Wien. Ent. Monatschr., 7, p. 452 (1863) (*Paraponyx*). (Typus: ♀ Mitteleuropa, Livland, Finnland, Irland, Estland, Schweden, Süd-Frankreich, N. Spanien, N. Italien, Istrien, Kroatische Küste,

- Wallengren, Oefvers. K. Vet.-Akad. Förh., 28 (8), p. 1008 (1871) (*Paraponyx*). — Berce, Fauna Ent. Fr. Lep., 6, p. 116 (1878). — Snellen, Vlinders Nederl., 2 (1), p. 77 (1882). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897). — Meyrick, Handb. Brit. Lep., p. 403 (1895). — Sharp, in Harmer & Shipley, Cambridge Nat. Hist., 6, Ins. (2), p. 423 (1899) (*N. (Paraponyx)*). — Cannaviello, Bol. Soc. Esp. Hist. Nat., 2, p. 172 (1902). — Rebel in Spuler, Schm. Eur., 2, p. 221, t. 81, f. 19 (1906). — Ulmer, Uns. Wasserins., p. 83 (1911) (*Paraponyx*). — Wahlgren, Svensk Insektafauna, 10 (2), p. 47, t. 2, f. 52 (1915) (*Paraponyx*). — Wesenberg-Lund, Ins. Liv. Ferske Vand., p. 202, f. 160 (imago), p. 210, 211, f. 165 (larva) (1915) (*Paraponyx*). — Börner in Brohmer, Fauna Deutschl., (ed. 3), p. 377 (1925). — Meyrick, Revis. Handb. Brit. Lep., p. 419 (1928) (*Paraponyx*). — Hering, Tierwelt, Mitteleur. (Ergbd., 1), p. 282 (1932) (*Paraponyx*). — Börner in Brohmer, Fauna Deutschl., (ed. 4), p. 389 (1932). — Eckstein, Schm. Deutschl., 5, p. 62 (1933) (*Paraponyx*). — Karny, Biol. Wasser-Ins., p. 160, f. 103 B (larva), p. 166, 173 (1934) (*Paraponyx*).
stratiotalis [Schiffermüller et Denis], Verz. Schm. Wien, p. 123 (1776) (*Pyrallis*). — Hübner, Samml. Eur. Schm., 6. Horde (*Pyr.*), t. 13, f. 87 (1796) (*Pyrallis*). — Hübner, Verz. bek. Schm., p. 362 n. 3461 ([1816]—1826). (*Paraponyx*). — Treitschke, Schm. Eur., 7, p. 137 (1829) (*Nymphula*). — Duponchel, Hist. Nat. Lep., 8 (2), p. 183, t. 222, f. 6 (1831). — Stephens, Ill. Brit. Ent. Haust., 4, p. 41 (1834). — Eversmann, Fauna Lep. Volgo-Ural., p. 464 (1844). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 10 (1848). — Guenée, Delt. et Pyral., p. 269 (1854) (*Paraponyx*). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 453 (1859) (*Paraponyx*). — Barrett, Lep. Brit. Isl., 9, p. 287, t. 408, f. 1, 1 a, b (larva) (1904).
paludata Fabricius, Ent. Syst., 3 (2), p. 213 n. 312 (1794) (*Phalaena*) (Anglia).
paludalis Schrank, Fauna Boica, 2 (2), p. 62 (1802).
ab. fasciata Teich, Korr. Blatt. Naturf. Livland Ver. Riga, 51, p. 37 (1908).
subsp. *amanica* Osthelder, Mitt. Münchn. Syrien Ent. Ges., 24, p. 95 (1935).
Biologie: Ochsenheimer & Treitschke, Schm. Eur., 7, p. 137 (1829). — Buckler, Ent. Mon. Mag., 12, p. 160 (1875); Larvae Brit. Lep., 9, p. 79, t. 151, f. 2 (1901). — Sorhagen, Allg. Zeitschr. Ent., 6, p. 277 (1901). — Herzegowina, Bosnien, Rumänien, Bulgarien, Südrussland, Kleinasien, Issyk Kul, Iligebiet, Russ. Turkestan, Uralsk

- Rebel in Spuler, Schm. Eur. Nachtr.-Taf. 9, f. 58 (1906).
- vittalis** Bremer, Lep. Ost-Sib., p. 66, t. 6, f. 3 (1864) (*Oligostigma*). — Pryer, Trans. Asiat. Soc. Jap., 13, p. 61 (1885) (*Oligostigma*). — Meyrick, Trans. Ent. Soc. Lond., p. 466 (1890). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897). — Leech, Trans. Ent. Soc. Lond., p. 434 (1901). — Wileman, Trans. Ent. Soc. Lond., p. 371 (1911). — Shibuya, Ins. Matsum., 3, p. 124, 126, t. 5, f. 8 (1929). — Matsumura, 6000 Ill. Ins. Japan, p. 1044, fig. (♂) (1931).
- regularis* Pryer, Cist. Ent., 2, p. 234, t. 4, f. 8 (1877) (*Oligostigma*).
- Biologie: Marumo, Oyo Dobuts. Zasshi Tokyo, 1 (1), p. 45—46, fig. (1929) (larva).
- nivalis** [Schiffermüller et Denis], Verz. Schm. Wien, p. 316 (1776) (*Phalaena*). — Hampson, Trans. Ent. Soc. Lond., p. 143 (1897). — Rebel in Spuler, Schm. Eur., 2, p. 221 (1906). — M. Hering, Tierwelt Mitteleuropa, (Ergbd. 1), p. 282 (1932) (*Paraponyx*). — Eckstein, Schm. Deutschl., 5, p. 62 (1933).
- candidata* Fabricius, Mant. Ins., 2, p. 215 (1787) (*Phalaena*); Ent. Syst., 3 (2), p. 321 (1794) (*Phalaena*).
- nivealis* Hübner, Samml. Eur. Schm., 6. Horde (*Pyral.*), t. 21, f. 141 (1896) (*Pyralis*); id., Verz. bek. Schmett., p. 362 n. 3460 (1816) [1826] (*Paraponyx*). — Ochsenheimer & Treitschke, Schm. Eur., 7, p. 133 (1829). — Eversmann, Fauna Lep. Volgo-Ural., p. 463 (1844). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 111 (1848). — Heinemann, Kleinschm. 1, H. 2, p. 105 (1865). — Wallengren, Oefvers. K. Vet.-Akad. Förh., 28 (8), p. 1008 (1871).
- algalis** Lucas P. H., Explor. Alg., 3 (6), p. 404 (1849) (*Hydrocampu*). — Guenée, Delt. et Pyral., p. 271 (1854) (*Paraponyx*).
- arundinalis** Eversmann, Bull. Soc. Imp. Nat. Moscou, 15, p. 561, t. 6, f. 12 (1842); Fauna Volgo-Ural., p. 469 (1844). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 11, f. 95 (1848). — Hampson, Trans. Ent. Soc. Lond., p. 144 (1897). — Krulikovsky, Rev. Russe Ent., 9, p. 316 (1909) (*Kasania*).
- ussuriensis** Rebel, Deutsche Ent. Zeitschr. Iris, 24, p. 6, t. 1, f. 7 (♀) (1910). — Caradja, Deutsche Ent. Zeitschr. Iris, 30, p. 21 (♂) (1917) (*ussurialis*). — Shibuya, Ins. Matsum., 3, p. 124, 127, t. 5, f. 9 (1929). — Caradja, Bull. Acad. Roum., 15, p. 147 (1932). — Matsumura, 6000 Ill. Ins. Japan., p. 1044, fig. (♂) (1931).
- Amur,
Mandschurei,
Ost- u.
Mittelchina,
Korea,
Japan
- Bayern,
Österreich,
Ungarn,
Rumänien,
Bosnien,
Südrussland,
S. Ural,
Uralsk,
Kirghisen-
steppe
- Mauretanien
(Algier)
- S. O.-
Rußland,
N.W.-
Sibirien,
S. Ural,
Uralsk,
Kirghisen-
steppe
- Ussuri,
Amur,
O.- u. S.-
China
(Typus: ♀
Chaborowka)

- munakatae* Matsumura, Oyo Konohugaku, p. 548 (1920). — Shibuya, Ins. Matsum., 2 (2), p. 97 (1927) (= *N. ussuriensis*).
- fluctuosalis** Zeller, Lep. Micropt. Caffr., p. 27 (1852) (*Nymphula*) (Natal). — Hampson, Ill. Typ. Sp. Het. Brit. Mus., 8, p. 40 (1891) (*Paraponyx*); 9, p. 51 (1893). — Meyrick, Trans. Ent. Soc. Lond., p. 11 (1894). — Hampson, Fauna Brit. India Moths, 4, p. 193, f. 115 (♂) (1896); Trans. Ent. Soc. Lond., p. 143; p. 138, fig. (♂) (1897). — Matsumura, Ill. Ent. Zeit. Berlin, 24, p. 381 (1900). — Hering, Stettin. Ent. Zeit., 64, p. 61 (1903). — Wileman, Trans. Ent. Soc. Lond., p. 371 (1911). — Strand, Ent. Mitt., 8, p. 102 (1919). — Matsumura, Oyo Konchugaku, p. 548, t. 22, f. 7 (1920). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 141, 144 (1928); Ins. Matsum., 3, p. 124, 127, t. 5, f. 11 (1929). — Matsumura, 6000 Ill. Ins. Japan, p. 1043, fig. (♂) (1931).
- linealis* Guenée, Delt. et Pyral., p. 271 (1854) (*Paraponyx*) (Ostindien). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 454 (1859) (*Paraponyx*). — Lederer, Wien. Ent. Monatschr., 7, p. 452 (1863) (*Paraponyx*). — Snellen in Veth, Midden-Sumatra, 4, p. 76 (1880); Tijds. v. Ent., 27, p. 48 (1884) (*Paraponyx*). — Moore, Lep. Ceylon, 3, p. 302 (1885) (*Paraponyx*). — Meyrick, Trans. Ent. Soc. Lond., p. 212 (1888) (*Paraponyx*).
- chrysippusalis* Walker, Cat. Lep. Het. Brit. Mus., 17, p. 432 (1859) (*Oligostigma*) (China).
- obitalis* Walker, Cat. Lep. Het. Brit. Mus., 17, p. 432 (1859) (*Oligostigma*). (Ceylon). — Wallace et Moore, Proc. Zool. Soc. Lond., p. 364 (1866) (*Oligostigma*).
- curta* Butler, Ent. Mon. Mag., 15, p. 270 (1879) (*Oligostigma*) (Hawaii).
- oryzalis* Wood-Mason, Rico Pest of Burma, Calcutta (1885) (*Paraponyx*).
- Biologie: De Joannis, Bull. Soc. Ent. Fr., p. 137—139 (1913) (Mimicry).
- noxia*: Shiraki, Injur. Ins. Formos, 1, p. 128, t. 32, f. 2 (1910). — Duport, Bull. Econ. Indochine (Hanoi-Haiphong) n. s., nr. 99 (1912), nr. 102, 105 (1913).
- depunctalis** Guenée, Delt. et Pyral., p. 274 (1884) (*Hydrocampa*). (Ostindien). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 461 (1859) (*Hydrocampa*). — Lederer, Wien. Ent. Monatschr., 7, p. 451 (1863) (*Hydrocampa*). — Swinhoe, Proc. Zool. Soc. Lond., p. 869 (1885) (*Hydrocampa*). — Moore, Lep. Ceylon, 3, p. 305, (1886) (*Nymphula*). — Swinhoe, Trans. Ent. Soc., p. 288 (1890) (*Hydrocampa*). — Meyrick, Trans. Ent. Soc. Lond., p. 11 (1894) (*Nymphula*). — Hamp-

Tropische
u. subtrop.
Gebiete,
Indien,
Ceylon,
Westchina,
Tonkin,
Formosa,
Japan,
Sumatra,
Hawai,
Australien,
Sandwich
Ins.,
Afrika
(Port.
Ostafrika,
Natal)

Gibraltar,
Marokko,
Indien,
Burma,
Südchina,
Tonkin,
Formosa,
Borneo,
Java,
Sumatra,
Ceylon,

- son, Fauna Brit. India Moths, 4, p. 195 (1896); Trans. Ent. Soc. Lond., p. 144 (1897). — E. Hering, Stettin. Ent. Zeit., 64, p. 64 (1903). — Strand, Ent. Mitt., 8, p. 102 (1919). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 142, 145, t. 6, f. 11 (1928). — Matsumura, 6000 Ill. Ins. Japan, p. 1043, fig. (♂) (1931).
- stagnalis* Zeller, Lep. Micr. Gaffr., p. 26 (1852) (nom. praeocc.). — Lederer, Wien. Ent. Monatschr., 7, p. 452 (1863) (*Parapoynx*). — Hampson, Ill. Typ. Sp. Lep. Het. Brit. Mus., 8, p. 40 (1891) (*Parapoynx*); 9, p. 51 (1893) (*Parapoynx*). — Kaye, Proc. Ent. Soc. Lond., p. XVIII (1900) (*Hydrocampa*).
- decussalis* Walker, Cat. Lep. Het. Brit. Mus., 17, p. 481 (♂, ♀) (1859) (*Zebronia?*). (Ceylon). — Meyrick, Trans. Ent. Soc. Lond., p. 433 (1885) (*Parapoynx*) (Queensland, Ceylon).
- dianalis* Schaus, Proc. U. S. Nat. Mus., 30, p. 1377 (1905) (*Parapoynx*). (S. Paulo).
- vestigialis* Snellen, in Veth, Middeu-Sumatra, 4, p. 78 (1880); Tijds. v. Ent., 27, p. 50 (1884). — Möschler, Abh. Senckenb. Ges., 16 (1), p. 318 (1890). (Portoriko).
- Biologie*: De Joannis, Bull. Soc. Ent. Fr., p. 137—139 (1913) (Mimicry).
- noxia*: Green, Rep. Gov. Ent. Dep. Agric. Ceylon pt. 4, p. 2—5 (1911/12). — Fletcher, Madras Dep. Agr. Bull., 3, nr. 67, 10 pp. (1913); Rep. Gov. Ent. Dep. Agr. Madras Pres., p. 36—41 (1913). — Mackenna, Rep. Progr. Agr., India f. 1915/16, p. 50—56 (1916). — Mac Swiney, Rep. Agr. Dep. Assam f. 1918/19, p. 6—8 (1919). — Hutson, Rep. Ent. Ceylon Dep. Agr. Admin, f. 1919, p. C8—C10 (1920). — Van Hall, Meded. Inst. Plantenziekten, Buitenzorg, nr. 39 (1920), nr. 46 (1921), nr. 61 (1923). — Pillai, Short Notes Ins. Pests Crops Travancore Dep. Agr. Trivandrum 53, pp. (1921). — Ramakrishna Ayyar, Rep. Proc. 4, Ent. Meeting Pusa, p. 49—55 (1921). — Subramania Jyer, Journ. Mysore Agr. Exp. Union, Bangalore, 3, n. 3, 4, p. 189—194 (1921). — South, Agr. Bull. FMS Kuala Kumpur, 9, nr. 4, p. 284—289 (1921). — Scott, Rep. Agr. Dep. Assam, p. 6 (1923). — South, Malayan Agr. Journ., 10, nr. 10—12, p. 264—248, 1922; 11, nr. 4, p. 89—96 (1923). — Jack, Malayan Agr. Journ., 11, nr. 5, 6, p. 103—119, 139—161 (1923). — Hector, Rep. Dep. Agr. Bengal (1919/20); append. 5, p. 3 (1920/21); p. 37—41 (1922/23); p. 1—5 (Calcutta) (1921—24). — Gosh, Rep. Proc. 5, Ent. Meeting Pusa, p. 89 (1924). — South et Birkinshaw, Malayan Agr. Journ., 12, nr. 2, p. 32—50 (1921). — Susainathan, Rep. 5, Ent. Mee-

Celebes,
Philippinen,
Australien,
Portug.
Ostafrika,
Südafrika,
Malediven,
Südamerika
(S. Paulo),
Portorico

Ceylon

ting Pusa, p. 25—33 (1924). — South, Malayan Agr. Journ., 13, nr. 7, p. 189—202 (1925). — Jardine, Reeve et Jebaratnam, Ceylon Adm. Rep. Agr., p. D17—19 (1924); p. D19—D20 (1926) (Colombo). — Gosh, Dep. Agr. Burma, nr. 16, 19—21, 25—29, 33, 35—42, 48 (1925). — Mac Kerral, Rep. Oper. Dep. Agr. Burma, 34 pp. (1925). — Gupta, Ann. Rep. Dep. Agr. Assam, p. 26—27 (1926). — Gosh, Rep. Ent. Mandalay, Sericult. Work f. (1925), 10+8 p. (1925) (Rangoon). — Kunhi Kannan, Journ. Mysore Agr. Exp. Union, 8, nr. 1, p. 11—17 (1926) (Bangalore). — Gupta, Rep. Dep. Agr. Assam 1928/29, p. 44—45 (1929). — Leefmans, Meded. Inst. Plantenziekten, nr. 75 (Buitenzorg), 96 pp. (1929). — Miller et Pagden, Malayan Agr. Journ., 18, nr. 6, p. 289—292 (1930). — Corbett, Proc. 4. Pac. Sci. Congr. Java, 4, p. 133—135 (1930) (Batavia). — Ramakrishna Ayyar, Agric. Livestock India, 3, pt. 4, p. 341—351 (Delhi) (1933).

takamukai Shibuya, Ins. Matsum., 3, p. 124, 127, t. 5, f. 10 (♂) (1929).

Japan
(Typus:
Kiushu)

orientalis Filipjev, Trav. Inst. Zool. Acad. Sci. URSS, 2, p. 15, fig. (1934).

Ussuri

b) Orientalische Region.

nymphacata Linné vide sub a) Paläarktische Region p. 73.

interruptalis Pryer vide sub a) Paläarktische Region p. 75.

nigrolinealis Pryer, Cist. Ent., 2, p. 233, t. 4, f. 6 (1877). — Hampson, Trans. Ent. Soc. Lond., p. 139 (1897). — Caradja, Mem. Acad. Rom. Sect. Stiint. s. 3, 3, mem. 7, p. 328 (1925). subsp. *sordidior* W. Rothschild, Journ. Bombay Nat. Hist. Soc., 28, p. 178 (♂) (1921).

Zentral- u.
Westchina

Persien

bifurcalis Wileman vide sub a) Paläarktische Region p. 76.

nigra Warren, Ann. Mag. Nat. Hist., s. 6, 8, p. 221 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 140 (1897); Journ. Bombay Nat. Hist. Soc., 12, p. 701 (1899). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 141, 142 (1928).

Indien
(Khasis),
Formosa

hampsoni South, Trans. Ent. Soc. Lond., p. 420 (♀), t. 14, f. 28 (♀) (1901) (*Trichophyesis*). — Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 209 (1903) (*Nymphula*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 23, 142, 1928.

Assam,
Ceylon,
Westchina,
Formosa
(Typus:
Ichang;
[Khasis])

- fengwhanal**is Pryer, vide sub a) Paläarktische Region p. 76.
- foedalis** Guenée, vide sub a) Paläarktische Region p. 77.
- spilomelalis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 403 (♂) (1859) (*Isopteryx*). — Hampson, Trans. Ent. Soc. Lond., p. 140 (1897) (= *foedalis* Guen.). — Swinhoe, Cat. East. Austral. Lep. Het., 2, p. 438 (1900) (= *foedalis* Guen.). — W. Rothschild, Rep. B. O. U. Exped., 2, nr. 15, p. 119 (1915) (spec. dist.). Ceylon (Typus), Holl. Neu-Guinea
- foedalis* (part.) auctorum.
- affinialis** Guenée, vide sub a) Paläarktische Region p. 77.
- sinicalis** Hampson, Trans. Ent. Soc. Lond., p. 141 (1897). — Lecch, Trans. Ent. Soc. Lond., p. 432 (1901). Ostchina, Tonkin (Typus: Chekiang)
- manilensis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 369 (♂) (1917). Philippinen
- responsalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1326 (♀) (1865) (*Diasemia*). (Moreton-Bai). — Meyrick, Trans. Ent. Soc. Lond., p. 207 (♂) (1887) (*Parapoynx*). — Swinhoe, Trans. Ent. Soc. Lond., p. 288 (1890) (*Hydrocampa*). — Hampson, Ill. Typ. Sp. Lep. Het. Brit. Mus., 9, p. 51 (1893) (*Nymphula*). — Meyrick, Trans. Ent. Soc. Lond., p. 470 (1894). — Hampson, Fauna Brit. India Moths, 4, p. 192 (1893); Trans. Ent. Soc. Lond., p. 141 (1897). — Leeche, Trans. Ent. Soc. Lond., p. 432 (1901). — Hering, Stettin. Ent. Zeit., 64, p. 63 (1903). Indien, Burma, Tonkin, Ceylon, Sumatra, Java, Lombok, Celebes, Australien, Queensland
- marmorata* Meyrick, Trans. Ent. Soc. London, p. 434 (♀) (1885) (*Parapoynx*). (Queensland, N. S. Wales). (Typus: ♀ Duaringa)
- crisonalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 961 (1859) (*Hydrocampa*). (Ceylon). — Hampson, Ill. Sp. Typ. Sp. Lep. Het. Brit. Mus., 9, p. 176, t. 174, f. 8 (1893) (*Parapoynx*); Fauna Brit. India Moths, 4, p. 192 (1897); Trans. Ent. Soc., p. 141 (1897). Ost- u. Südchina, Ceylon, Java, Sumatra, Celebes, Australien, Melville Ins.
- hebraicalis** Snellen, Tijds. v. Ent., p. 23, p. 240 (1880) (*Parapoynx*); 27, p. 48, t. 4, f. 11 a (caput) (1884) (*Parapoynx*). — E. Hering, Stettin. Ent. Zeit., 64, p. 62 (1903) (spec. dist.). (Typus: Makassar)
- myina* Meyrick, Trans. Ent. Soc. Lond., p. 432 (1885) (*Parapoynx*). (Queensland). — E. Hering, Stett. Ent. Zeit., 64, p. 62 (1903) (bon. spec.). (Typus: ♂, Duaringa)
- diffualis** Snellen, in Veth, Midden-Sumatra, 4 (1), 8, p. 76 (1880); Tijds. v. Ent., 27, p. 47 (1884). — Hampson, Trans. Ent. Soc. Lond., p. 146 (1897). — E. Hering, Ent. Zeit. Stettin, 64, p. 63 (1903) (= *responsalis* Walk.). Sumatra, Java, Philippinen

- definitalis** Strand, Ent. Mitt., 8, p. 61, (♂, ♀) (1919). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 141, 143 (1928). Formosa
- turbata** Butler, vide a) Paläarktische Region p. 77.
- enixalis** Swinhoe, vide a) Paläarktische Region p. 78.
- fregonalis** Snellen, Tijds. v. Ent., 23, p. 241 (♂, ♀) (1880); 27, p. 49, t. 4, f. 12 (1884) (*Parapoynx*). — Hampson, Trans. Ent. Soc. Lond., p. 145 (1897). — E. Hering, Ent. Zeit. Stettin, 64, p. 62 (1903). Celebes, Sumatra, Java (Typus: Bonthain, Makassar)
- diminutalis** Snellen, Tijds. v. Ent., 23, p. 242 (♂, ♀) (1880) (*Parapoynx*); 27, p. 48, t. 5, f. 1 (1884) (*Parapoynx*). — Meyrick, Trans. Ent. Soc. Lond., p. 470 (1894). — Hampson, Fauna Brit. India Moths, 4, p. 193 (1896); Trans. Ent. Soc. Lond., p. 142 (1897). — Pagenstecher in Kükenthal, Forschungsreise, p. 465 (1897) (*Parapoynx*). — E. Hering, Stett. Ent. Zeit., 64, p. 62 (1903). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 142, 144 (1928). Indien, China, Tonkin, Formosa, Ceylon, Sumatra, Java, Celebes, Lombok, Borneo, Philippinen, Port. Ostafrika (Typus: Makassar)
- vitalis** Bremer, vide sub a) Paläarktische Region p. 80.
- ussuriensis** Rebel, vide sub a) Paläarktische Region p. 80.
- fluctuosalis** Zeller, vide sub a) Paläarktische Region p. 81.
- votalis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 433 (♂) (1859) (*Oligostigma*). (Ceylon). — Moore, Lep. Ceylon, 3, p. 302 (1885) (*Parapoynx*). — Swinhoe, Trans. Ent. Soc. Lond., p. 285 (1890) (*Parapoynx*). — Hampson, Ill. Typ. Sp. Lep. Het. Brit. Mus., 8, p. 40 (1891) (*Parapoynx*); id., ibid., 9, p. 51, t. 174, f. 4 (1893) (*Parapoynx*); id., Fauna Brit. India Moths, 4, p. 194 (1896) (*Nymphula*); id., Trans. Ent. Soc. Lond., p. 142 (1897). Indien, Burma, Tonkin, Ceylon
- leucostola** Hampson, Fauna Brit. India Moths, 4, p. 194 (♂) (1896); Trans. Ent. Soc. Lond., p. 143 (1897). Burma (Typus: Rangoon)
- fusalis** Hampson, Fauna Brit. India Moths, 4, p. 195 (1896); Trans. Ent. Soc. Lond., p. 143 (1897). Assam (Typus: Nāgas)
- incurvalis** South, Trans. Ent. Soc. Lond., p. 434 (♂) (1901). Zentralchina
- osculatrix** Meyrick, Exot. Microlep., 4, p. 394 (♂, ♀) (1933). Siam, S.O. Borneo, Queensland (Brisbane)
- fuscicostalis** Hampson, Fauna Brit. India Moths, 4, p. 195 (♂) (1896); Trans. Ent. Soc. Lond., p. 143 (1897). Bengalen (Calcutta)

- depunctalis** Guenée, vide sub a) Paläarktische Region p. 81.
- slmplalis** Snellen, Trans. Ent. Soc. Lond., p. 638 (1896). — Hampson, Fauna Brit. India Moths, 4, p. 195 (♂) (1896); Trans. Ent. Soc. Lond., p. 144 (1897). Sikkim
- grisealis** (Hampson, Journ. Bombay Nat. Hist. Soc., 21, p. 1261 (♀) (1912). Ceylon
- uxorialis** Strand, Ent. Mitt., 8, p. 102 (♂) (1919). Formosa
— Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 141, 145 (1928).
- litanalis** Walker, Cat. Lep. Hct. Brit. Mus., 18, p. 706 (♂) (1859) (*Botys?*), (Sarawak). — Borneo, Assam
Hampson, Journ. Bombay Nat. Hist. Soc., 18, p. 572 (1908).
titanalis Hampson, Trans. Ent. Soc. Lond., p. 144 (1897).
stenialis Warren, Ann. Mag. Nat. Hist., (s. 6) 8, p. 68 (1891) (*Endotricha*).
- seriopunctalis** Hampson, Trans. Ent. Soc., p. 144 (1897). Amboina, Buru, Fergusson Ins.
- unguicalis** Snellen in Veth, Midden-Sumatra, 4 (1), 8, p. 77 (1880). — E. Hering, Ent. Zeit. Sumatra, Java, Celebes
Stettin, 64, p. 61 (1903).
- cuneoalis** Snellen, Tijds. v. Ent., 23, p. 243 (♂) (1880) (*Parapoynx*); 27, p. 48, t. 5, f. 2 (1884) (*Parapoynx*). — Pagenstecher, Jahrb. Ver. Nassau, 37, p. 283 (1886) (*Parapoynx*). Aru, Amboina, Celebes (Typus: Bonthain)
- coenosalis** Snellen, Tijds. v. Ent., 38, p. 157 (♂) (1895) (*Hydrocampa*). — Hampson, Trans. Ent. Soc. Lond., p. 145 (1897). Celebes (Typus: S. Celebes, Bantimorong)
- fulguralis** Canadja, Deutsche Ent. Zeitschr. Iris, 47, p. 153 (1934). China (Kwangtung)

c) Aethiopische und Madagassische Region.

- foedalis** Guenée, vide sub a) Paläarktische Region p. 76.
- affinialis** Guenée, vide sub a) Paläarktische Region p. 77.
- diminutalis** Snellen, vide sub b) Orientalische Region p. 35.
- fluctuosalis** Zeller, vide sub a) Paläarktische Region p. 81.
- depunctalis** Guenée, vide sub b) Orientalische Region p. 81.
- plumbefusalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 370 (♀) (1917). Sudan
- clrcealis** Walker, Cat. Lep. Hct. Brit. Mus., 17, p. 460 (♂) (1859) (*Hydrocampa*). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897). Südafrika (Typus) Madagaskar

Species incertae sedis.

- lutelvittalis** Mabille, C. R. Ent. Belg., 23, p. XXVI (1880). Madagaskar
minimalis Saalmüller, Ber. Senckenb. Ges., p. 298 (1879) (*Hydrocampa*); Lep. Madag., 2, p. 505 (1891). Madagaskar
minoralis Mabille, C. R. Ent. Belg., 25, p. LXIII (1881) (*Paraponyx*). Madagaskar
ophiaula Meyrick, Exot. Microlep., 5, p. 8 (♂, ♀) (1936). Belg. Congo
pseudofoedalis Strand, Arch. Naturg., 73, A H. 12, p. 76 (1912). Kamerun, Span. Guinea

d) Australische Region.

- foedalis** Guenée, vide sub a) Paläarktische Region p. 76.
spilomelalis Walker, vide sub b) Orientalische Region p. 84.
affinialis Guenée, vide sub a) Paläarktische Region p. 77.
responsalis Walker, vide sub b) Orientalische Region p. 84.
crisonalis Walker, vide sub b) Orientalische Region p. 84.
dicentra Meyrick, Trans. Ent. Soc. Lond., p. 431 (1885) (*Paraponyx*). — Hampson, Trans. Ent. Soc. London, p. 142 (1897). — E. Hering, Stettin. Ent. Zeit., 64, p. 63 (1903). Australien, Queensland
pallida Butler, Trans. Ent. Soc. Lond., p. 423 (1886) (*Oligostigma*).
osculatrix Meyrick, vide sub b) Orientalische Region p. 85.
nitens Butler, Cist. Ent., 2, p. 556 (1880) (*Paraponyx*). — Meyrick, Trans. Proc. N. Zeal. Inst., 17, p. 130 (1885) (*Hygraula*); 20, p. 63, 1 (1888) (*Paraponyx*). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897). — Longstaff, Butt. Hunt. Many Lands, p. 452, 474 (1912). — Hudson, Butt. Moths N. Zeal., p. 177, t. 19, f. 22 (♂) (1928). Australien, N.S. Wales, Neuseeland
tullialis Walker, Cat. Lep. Het. Brit. Mus., 17, p. 462 (♀) (1859) (*Hydrocampa*). — Meyrick, Trans. Ent. Soc. Lond., p. 436 (1885) (*Hydreuretis*). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897). Australien, N. S. Wales
expatrialis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 391 (♀) (1906). Neu-Guinea, Fergusson Ins.
fluctuosalis Zeller, vide sub a) Paläarktische Region p. 81.
dentizonalis Hampson, Trans. Ent. Soc. Lond., p. 143 (♂) (1897). Queensland (Typus: Cedar Bay)
medusalis Walker, Cat. Lep. Het. Brit. Mus., 17, p. 486 (♀) (1859) (*Zebronia?*) (Moreton-Bai). — Meyrick, Trans. Ent. Soc. Lond., Australien

- p. 438 (1885) (= *decussalis* Walk.). — Hampson, Trans. Ent. Soc. Lond., p. 144 (1897).
- depunctalis** Guenée, vide sub b) Orientalische Region p. 81.
- seriopunctalis** Hampson, vide sub b) Orientalische Region p. 86.
- polystictalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 391 (♀) (1906). Brit. u. Holl. Neu-Guinea, Ron Ins.
- metastictalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 370 (♀) (1917). D'Entrecasteaux Ins. (Goodenough Ins.)
- pleonaxalis** Hampson, Trans. Ent. Soc. Lond., p. 144 (1897). Humboldt-Bai Fergusson Ins., Neu-Guinea

Species incertae sedis.

- aristodera** Turner, Trans. Roy. Soc. S. Austral., 32, p. 83 (1908). Queensland
- cuneoalis** Snellen, vide sub b) Orientalische Region p. 86.
- diplopa** Lower, Trans. Roy. Soc. S. Austral., 27, p. 62 (1903). N. S. Wales
- epimochla** Turner, Trans. Roy. Soc. S. Austral., 32, p. 83 (♂) (1918). Queensland
- euryscia** Meyrick, Trans. Ent. Soc. Lond., p. 435 (♂) (1885) (*Hydreuretis*). Australien
- pudica** Lucas, Proc. Linn. Soc. N. S. Wales (s. 2), 8, p. 157 (♂, ♀) (1893) (*Paraponyx*). W. Australien
- sinuosa** Lucas, Proc. Linn. Soc. N. S. Wales (s. 2) 7, p. 261 (1892). Queensland
- stenoides** Butler, Ann. Mag. Nat. Hist., (s. 5) 10, p. 235 (1882) (*Hydrocampa*). — Pagensteher, Zoologica H. 29, p. 172 (1900). Herzog von York Ins.
- tenebralis** Lower, Proc. Linn. Soc. N. S. Wales, 26, p. 664 (♂) (1902). Queensland
- terranea** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 147 (♀) (1915). Holl. Neu Guinea

e) Nearktische Region.

- foedalis** Guenée, vide sub a) Paläarktische Region p. 76.
- maculalis** Clemens, Proc. Acad. Nat. Sci. Phil., 12, p. 218 (♂) (1860) (*Sironia*). — Grote, New Ch. List N. Amer. Moths, p. 54 (1882) (*Paraponyx*). — Hart, Bull. Ill. Lab. Nat. Hist., 4, p. 167 (1895) (*Nymphaeella*). — Dyar, Journ. N. York Ent. Soc., 14, p. 78, 79 (♂, ♀) 1906). — Strand, Arch. Naturg., 85 A H. 12, p. 131 (1920) — Forbes, Lep. N. York, p. 577 (1923). Vereinigte Staaten, (N. York, Massachusetts, Florida), Ontario
- ab. *foeminalis* Dyar, Journ. N. York Ent. Soc., 14, p. 80 (♂) (1906). — Forbes, Lep. N. York, p. 578 (1923). Florida (Typus: Palm Beach)

- ab. *masculinalis* Dyar, Journ. N. York Ent. Soc., 14, p. 80 (♀) (1906). — Forbes, Lep. N. York, p. 578 (1923). Florida
(Typus:
Coccanut
Grove)
- seminivella* Walker, Cat. Lep. Het. Brit. Mus., 35, 1717 (♀) (1866) (*Nephopteryx*).
— Hampson, Trans. Ent. Soc. Lond., p. 142 (1897).
- dispar* Grote, Nov. Amer. Ent., 1, p. 97 (1880) (*Nymphacella*).
- Biologie: Dyar, Journ. N. York Ent. Soc., 14, p. 80 (1906). — Forbes, Psyche, 17, p. 219—226, t. 12 (1910). — Welch, Ann. Ent. Soc. Amer., 9, p. 160—181, f. 1—10, 12—19 (1916); 17, p. 395—402 (1924). — Welch et Sehon, Ann. Ent. Soc. Amer., 21, p. 243—258 (1928).
- allionealis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 453 (♂, ♀) (1859) (*Paraponyx*). — Hart, Ill. Stat. Lab. Nat. Hist., 4, p. 173 (1895) (*Paraponyx*). — Hampson, Trans. Ent. Soc., p. 142 (1897). — Fernald, Bull. 52, U. S. Nat. Mus., p. 4491 (1902). (*Nymphula*). — Dyar, Journ. N. York Ent. Soc., 14, p. 80 (♂, ♀) (1906). — Forbes, Lep. N. York, p. 578 (1923). Vereinigte Staaten
(Georgia,
Wiskonsin,
Illinois,
Massachusetts,
Ohio,
Florida)
- itealis* Walker, Cat. Lep. Het. Brit. Mus., 17, p. 458, n. 5 (♀) (1859) (*Hydrocampa*). (Georgia & U.S.A.). — Hampson, Trans. Ent. Soc., p. 142 (1897). — Forbes, Lep. N. York, p. 578 (1903) (*allionealis* race).
- cretacealis* Lederer, Wien. Ent. Monatschr., 7, p. 452, 484 (1863) (*Paraponyx*).
- plenilinealis* Grote, Papilio, 1, p. 18 (1881). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897).
- aptalis* Lederer, Wien. Ent. Monatschr., 7, p. 452, 485 (1863) (*Paraponyx*).
- fluctuosalis* (part.) Hampson, Trans. Ent. Soc. Lond., p. 143 (1897).
- obscuralis** Grote, Papilio, 1, p. 18 (1881) (*Oligostigma*). — Hart, Bull. Ill. Stat. Lab. Nat. Hist., 4, p. 167, t. 1, f. 1—7 (1895) (*Paraponyx*). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897). — Holland, Moth Book, p. 399, t. 48, f. 10 (1903). — Dyar, Journ. N. York Ent. Soc., 14, p. 81 (♂, ♀) (1906). — Forbes, Lep. N. York, p. 578 (1923). Vereinigte Staaten
(Maine,
Massachusetts,
N. York,
Pennsylvanien,
Ohio,
Illinois,
Wiskonsin,
Florida)
- Biologie: Dyar, Journ. N. York Ent. Soc., 14, p. 81, 82 (1906). — Forbes, Psyche, 17, p. 219—226, t. 12 (1910).
- seminealis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 430 (♂) (1859) (*Oligostigma*). — Dyar, Journ. N. York Ent. Soc., 14, p. 89 (♂, ♀) (1906) (*Oligostigma*). — Forbes, Lep. N. York, p. 578 (1923). Vereinigte Staaten
(Massachusetts,
Pennsylvanien bis
Florida)

- semineale* Hampson, Trans. Ent. Soc. Lond., p. 167 (1897) (*Oligostigma*).
- tedyuscongalis* Clemens, Proc. Acad. Nat. Sci. Phil., 12, p. 216 (1860) (*Eustales*). — Grote, New ch. list. N. Amer. moths, p. 64 (1882) (*Paraponyx*).
- juncealis* Fernald (nec Guenée) Bull. 52 U. S. Nat. Mus., p. 396, n. 4502 (1902) (*Oligostigma*).
- Biologie: Forbes, Psyche, 17, p. 219—226, t. 16 (1910).
- badiusalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 955 (1859) (*Cymoriza*) (Kanada). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897). — Dyar, Journ. N. York Ent. Soc., 14, p. 82 (♂, ♀) (1906). — Strand, Arch. Naturg., 85, A H. 12, p. 131 (1920). — Forbes, Lep. N. York, p. 578 (1923).
- curviferalis* Walker, Cat. Lep. Het. Brit. Mus., 34, 331 (1865) (*Oligostigma*). —
- albalis* Robinson, Ann. N. York Lyc. Nat. Hist., 9, p. 153, t. 1, f. 3 (1869) (*Oligostigma*). — Hart, Bull. Ill. Stat. Lab. Nat. Hist., 4, p. 173 (1895).
- vittatalis** Dyar, Journ. N. York Ent. Soc., 14, p. 89 (♂) (1906) (*Oligostigma*). Florida (Biscayne Bay)
- obliteralis** Walker, Cat. Lep. Het. Brit. Mus., 17, 399 (♀) (1859) (*Isopteryx*?). — Hart, Bull. Ill. Stat. Lab. Nat. Hist., 4, p. 176, t. 2, f. 7—12 (1895) (*Hydrocampa*). — Hampson, Trans. Ent. Soc., p. 141 (1897) (*Nymphula*). — Fernald, Bull. 52, U. S. Nat. Mus., n. 4496 (1902). — Dyar, Journ. N. York Ent. Soc., 14, p. 83 (♂, ♀) (1906). — Forbes, Lep. N. York, p. 577 (1923).
- obscuralis* Möschler, Verh. z. b. Ges. Wien, 31, p. 432, t. 18, f. 39 (1881) (*Paraponyx*).
- proprialis* Fernald, Ent. Amer., 4, p. 37 (1888) (*Hydrocampa*).
- Biologie: Dyar, Journ. N. York Ent. Soc., 14, p. 84 (1906).
- nomophilalis** Dyar, Journ. N. York Ent. Soc., 14, p. 84 (♂, ♀) (1906). Florida, Texas
- gyralis** Hulst, Trans. Amer. Ent. Soc., 13, p. 159 (1886) (*Hydrocampa*). — Hart, Bull. Ill. Stat. Lab. Nat. Hist., 4, p. 175 (1895) (*Hydrocampa*). — Fernald, Bull. 52 U. S. Nat. Mus., n. 4489 (1902) (*Nymphula*). — Dyar, Journ. N. York Ent. Soc., 14, p. 85 (♂, ♀) (1906). — Forbes, Lep. N. York, p. 577 (1923). Georgia, Florida, Illinois, N. York, Massachusetts
- dentilinea* Hampson, Trans. Ent. Soc. Lond., p. 139 (♂) (1897).
- Biologie: Dyar, Journ. N. York Ent. Soc., 14, p. 86 (1906). — Forbes, Psyche, 17, p. 219—226, t. 12 (1910).
- serrilinealis** Barnes et Benjamin, Contr. Nat. Hist. Ontario Lep. N. Amer., 5, p. 190 (1924).

- nebulosalis** Fernald, Ent. Amer., 3, p. 127 (1887) Florida
(*Hydrocampa*). — Hampson, Trans. Ent. Soc. Lond., p. 145 (1897). — Dyar, Journ. N. York Ent. Soc., 14, p. 86, 87 (1906).
- icciusalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 971 (♂) (1859) (*Leucochroma*). — Hart, Bull. Ill. Stat. Lab. Nat. Hist., 4, p. 176 (1895) (*Hydrocampa*). — Hampson, Trans. Ent. Soc. Lond., p. 139 (1897). — Dyar, Journ. N. York Ent. Soc., 14, p. 87 (1906). — Strand, Arch. Naturg., 85, A H. 12, p. 131 (1920). — Forbes, Lep. N. York, p. 576 (1923).
- faulalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 973 (♀) (1859) (*Leucochroma*). (Patria?).
- formosalis** Clemens, Proc. Acad. Nat. Sci. Phil., 12, p. 217 (1860) (*Hydrocampa*?). — Packard, Amer. Nat., 18, p. 824 (1884) (*Hydrocampa*).
- genuialis** Lederer, Wien. Ent. Monatschr., 7, p. 451, 483 (1863) (*Hydrocampa*). (N. Amerika, Venezuela).
- genuinalis** Lederer, Wien. Ent. Monatschr., 7, t. 18, f. 2 (1863) (*Hydrocampa*).
- pacalis** Grote, Papilio, 1, p. 17 (1881) (*Hydrocampa*).
- Biologie: Dyar, Journ. N. York Ent. Soc., 14, p. 87 (1906). — Forbes, Psyche, 17, p. 219—226, t. 12 (1910). — Welch, Ann. Ent. Soc. Amer., 9, p. 181—184, f. 11 (1916); id., New ch. list, N. Amer. moths, 54 (1882). — Fernald, Smith List Lep. Bor. Amer., n. 4175 (1891).
- ekthlipsis** Grote, Canad. Ent., 8, p. 111 (1876) Canada
(*Hydrocampa*). — Hart, Ill. Stat. Lab. Nat. Hist., 4, p. 176 (1895) (*Hydrocampa*). — Vereinigte
Hampson, Trans. Ent. Soc. Lond., p. 139 Staaten
(1897). — Fernald, Bull. 52 U. S. Nat. (Maine, N.
Mus., n. 4490 (1902). — Dyar, Journ. N. Hampshire,
York Ent. Soc., 14, p. 88 (1906). — Massa-
Forbes, Lep. N. York, p. 576 (1923). chusetts,
N. York,
Illinois)

Species incertae sedis.

- australis** Hulst, Trans. Amer. Ent. Soc., 13, p. 168 Florida
(1885) (*Hydrocampa*). — Hampson, Trans.
Ent. Soc. Lond., p. 145 (1897).
- camalis** Quaintance, Bull. Florida Exp. Stat., nr. Florida
45, p. 68 (1898).

f) Neotropische Region.

- aetalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. Brasilien
954 (♂) (1859) (*Cymoriza*). — Hampson,
Trans. Ent. Soc. Lond., p. 139 (1897).
- lotialis** Felder et Rogenhofer, Reise Novara,
p. 14, t. 135, f. 4 (1874).

- adiantealis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 949 (♂) (1859) (*Hydrophysa*). (Rio Janeiro). — Hampson, Trans. Ent. Soc., p. 140 (1897). Brasilien
- abrotalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 956 (♂) (1859) (*Cymoriza?*). — Hampson, Trans. Ent. Soc., p. 140 (1897). Brasilien
- foedalis** Guenée, vide sub a) Paläarktische Region p. 76.
- obliteralis** Walker, vide sub e) Nearktische Region p. 90.
- gurgitalis** Lederer, Wien. Ent. Monatschr., 7, p. 449, t. 17, f. 17 (1863) (*Synclita*). (Venezuela). — Hampson, Trans. Ent. Soc. Lond., p. 141 (1897). Brasilien, Venezuela, Surinam
- modestalis* Lederer, Wien. Ent. Monatschr., 7, p. 449, t. 18, f. 1 (1863) (*Synclita*). (Surinam).
- leucoplagalalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 369 (♂) (1917). Mexiko
- zoilusalis* Druce (nec Walker), Biol. Centr.-Amer. Het., 2, p. 558 (1895), part. (Bocchoris).
- filigeralis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1633 (♂) (1865) (*Hydrocampa?*). — Hampson, Trans. Ent. Soc. Lond., p. 141 (1897). Brasilien (Typus: Ega)
- endoralis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 459 (1859) (*Hydrocampa*). (Honduras). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897). Mexiko, Guatemala, Honduras, Costa Rica, Panama, Kolumbien, Uruguay, Bahama Ins.
- distinctalis* Snellen, Tijds. v. Ent., 18, p. 261, t. 14, f. 14 (1875) (*Parapoynx*). (Kolumbien).
- hermeasalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 945 (1859) (*Isopteryx?*) (Venezuela). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897). Südamerika, Panama
- hydrothlonalis** Snellen, Tijds. v. Ent., 18, p. 262, t. 14, f. 15 (1875) (*Parapoynx*). — Hampson, Trans. Ent. Soc. Lond., p. 142 (1897). Südamerika
- allionealis** Walker, vide sub e) Nearktische Region p. 89.
- perlalis** Hampson, Trans. Ent. Soc. Lond., p. 143 (♂) (1897). Guadeloupe, Peru
- depunctalis** Guenée, vide sub b) Orientalische Region p. 81.
- flavicostralis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 371 (♂) (1917). Poru
- graphicalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 371 (♀) (1917). Peru

Species incertae sedis.

- avertinalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 109 (♂) (1924). Cuba (Typus: Baracoa)
- effrenatalis** C. Berg, Ent. Zeit. Stettin, 37, p. 352, (1876). Uruguay

- flavialis** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, p. 677 (♀) (1902). Costa Rica
- franciscalis** Schaus, Proc. U. S. Nat. Mus., 30, p. 139 (1906). Brasilien (Paraná)
- impuralis** Feller et Rogenhofer, Reise Novara (Het.), p. 10, t. 135, f. 2 (1874). (*Glyphodes?*) Haiti (S. Domingo).
- indomitalis** Berg, Ent. Zeit. Stettin, 37, p. 352 (1876). Uruguay
- infirmalis** Möschler, Abh. Senckenb. Ges., 14 (3), p. 82 (1886) (*Parapoynx*). Jamaika
- pampenealis** Dyar, Ins. Insc. Menstr., 12, p. 20 (1924). Mexiko
- paulalis** Schaus, Proc. U. S. Nat. Mus., 30, p. 138 (1906). Brasilien (S. Paulo)
- rugosalis** Möschler, Abh. Senckenb. Ges., 16, p. 318 (1890) (*Parapoynx*). Portoriko
- vitrealis** Felder et Rogenhofer, Reise Novara (Het.), p. 14, t. 135, f. 11 (1874). Guayana
- 113

6. Genus **Araeomorpha** Turner

Turner, Trans. Roy. Soc. S. Austral., 32, p. 82 (1908) (generotypus: *A. atmota* Turn.).

- atmota** Turner, Trans. Roy. Soc. S. Austral., 32, p. 82 (♂, ♀) (1908). Queensland

1

7. Genus **Margarosticha** Lederer

Lederer, Wien. Ent. Monatschr., 7, p. 454 (1863). — Hampson, Trans. Ent. Soc. Lond., p. 132, 146 (1897).

(Typus generis: *pulcherrimalis* Led.)

- pulcherrimalis** Lederer, Wien. Ent. Monatschr., 7, p. 454, t. 18, f. 11 (1863). — Hampson, Trans. Ent. Soc. Lond., p. 146, fig. (♂) (1897). Amboina, Ceram
- gaudialis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 372 (♂) (1917). Admiralitäts-Ins.
- bimaculalis** Snellen, Tijds. v. Ent., 23, p. 245 (1880); 27, p. 50, t. 5, f. 5 (1884). — Meyrick, Trans. Ent. Soc. Lond., p. 474 (1894). — Hampson, Trans. Ent. Soc. Lond., p. 146 (1897). Celebes, Mindanao
- australis** Felder et Rogenhofer, Reise Novara (Het.), p. 5, t. 136, f. 10 (1874). — Hampson, Trans. Ent. Soc. Lond., p. 147 (1897). Australien, Queensland
- sphenotis* Meyrick, Trans. Ent. Soc. Lond., p. 204 (1897). — Lucas, Proc. Roy. Soc. Queensland, 13, p. 81 (1898).
- euprepialis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 372 (♂) (1917). Queensland

leucozonalis Hampson, Trans. Ent. Soc. Lond., p. 147 (♂) (1892). Fergusson Ins. Neu-Guinea

argyrograpta Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 373 (♀) (1917). Bismarck-Archipel (Rook Ins.)

Species incertae sedis.

distributa Lucas, Proc. Roy. Soc. Queensland, 13, p. 81 (♀) (1898). Queensland

plumbealis Kenrick, Proc. Zool. Soc. Lond., p. 544, t. 68, f. 9 (1912). Holl. Neu-Guinea

9

8. Genus **Cataclysta** Hübner.

Hübner, Verz. Schm., 5, p. 363 (1826). — Stephens, III. Brit. Ent. Haust., 4, p. 39 (1834). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 7 (1848). — Guenée, Delt. et Pyral., p. 263 (1854). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 435 (1859). — Stainton, Man. Brit. Butt. Moths, 2, p. 143 (1859). — Lederer, Wien. Ent. Monatschr., 7, p. 453 (1863). — Heinemann, Kleinschm., 1, H. 2, p. 106 (1865). — Moore, Lep. Ceylon, 3, p. 299 (1885). — Meyrick, Trans. Ent. Soc. Lond., p. 219 (1886); p. 466 (1890); Handb. Brit. Lep., p. 402 (1895). — Hampson, Fauna Brit. India Moths, 4, p. 196 (1896); Trans. Ent. Soc. Lond., p. 132, 147 (1897). — Pagenstecher, Zoologica H. 29, p. 173 (1900). — E. Hering, Ent. Zeit. Stettin, 64, p. 65 (1903). — Barrett, Lep. Brit. Isl., 9, p. 234 (1904). — Rebel in Spuler, Schm. Eur., 2, p. 221 (1906). — Wahlgren, Svensk Insekthäfa, 10 (2), p. 67 (1915). — Tullgren & Wahlgren, Svenska Ins., p. 388 (1922). — Meyrick, Revis. Handb. Brit. Lep., p. 418 (1923). — M. Hering, Tierwelt Mitteleur. Ergbd. 1, p. 282 (1932). — Eckstein, Schm. Deutschl., 5, p. 62 (1933).

Strepsinoma (part.) Meyrick, Trans. Ent. Soc. London, p. 85—86 (1897) (typus: *amaura* Meyrick).

Chrysendeton Grote, Papilio, 1, p. 16 (1881) (typus: *medicinalis* Grote).

Anydraula Meyrick, Trans. Ent. Soc. Lond., p. 427 (1885) (typus: *glycerialis* Walk.).

Elophila Dyar, Journ. N. York, Ent. Soc. 14, p. 90 (1906) (nec Hübner, Tentamen, 2, 1806). — Forbes, Lep. N. York, p. 579, f. 347, 349 (1923).

(Typus generis: *C. lemnalis* [Schiff. & Den.] = *lemnata* Linné.)

a) Paläarktische Arten.

blandialis Walker, Cat. Lep. Het. Brit. Mus., 17, p. 448 (♀) (1859) (Ceylon). — Moore, Lep. Ceylon, 3, p. 299, t. 179, f. 15 (1885). — Hampson, Fauna Brit. India Moths, 4, p. 197, f. 116 (♂) (1896); Trans. Ent. Soc. Lond., p. 148; p. 147, fig. (♂) (1897). — Leech, Trans. Ent. Soc. Lond., p. 435 (1901). Japan, Südchina, Formosa, Indien (Dharmasala), Calcutta, Nilgiris,

- E. Hering, Stettin, Ent. Zeit., 64, p. 65 (1903). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 146 (1928); Ins. Matsum., 3, p. 128, t. 5, f. 12 (1929).
- patnalis* Felder et Rogenhofer, Reise Novara (Het.), p. 5, t. 136, f. 7 (♂) (1877). — Pagenstecher, Jahrb. Nassau. Ver. Naturk., p. 118 (1888) (Amboina).
- stipalis* Snellen in Veth, Midden-Sumatra, 4, p. 78, t. 5, f. 11, 11a (1880) (*Nymphicula*). — Pagenstecher, Zoologica, H. 29, p. 174 (1900) (*Nymphicula*).
- cuneifera* Moore, Lep. Ceylon, 3, p. 300, t. 179, f. 14 (1885).
- junctalis* Hampson, Ill. Typ. Sp. Het. Brit. Mus., 8, p. 140, t. 155, f. 24 (1891) (Nilgiris). — Makumo, Journ. Col. Agr. Tokyo Imp. Univ., 8, p. 187 (1913).
- mesorophna* Meyrick, Trans. Ent. Soc. Lond., p. 10 (1894). — Hampson, Fauna Brit. India Moths, 4, p. 197 (♂), 566 (1896) (Shan Staaten).
- xanthobathra* Meyrick, Trans. Ent. Soc. Lond., p. 474 (1894) (*Anydraula*). (Pulo Trans.). — W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 118 (1915).
- perirrorata* Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 374 (1917) (Goldküste, S. Nigeria, Br. Zentr. Afrika, Mashonaland, Kapkolonie).
- forma *nigritalis* Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 178, t. 174, f. 5 (1893).
- blandialis* (part.) Hampson, Trans. Ent. Soc., p. 148, sub. n. 1 (1897).
- forma *bombayensis* Swinhoe et Cotes, Cat. Moths India, 5, p. 651 (1889) (n. nov.).
- dilucidalis* Walker (nee Guenée, 1854), Cat. Lep. Het. Brit. Mus., 17, p. 446, n. 18 (1859). (Ceylon).
- lemnata** Linné, Syst. Nat., ed. 10, p. 529 (1758) (*Phalaena Geometra*); ed. 12, 1 (2), p. 874 (1767) (*Geometra*). — Fabricius, Ent. Syst., 3 (2), p. 215, n. 319 (1794) (*Phalaena*). — Donovan, Nat. Hist. Brit. Ins., 8, t. 266, f. 1, 2 (1806) (*Phalaena*). — Stephens, Ill. Brit. Ent. Haust., 4, p. 40 (1834) (*Hydrocampa*). — Wood, Index Entom., p. 123, t. 28, f. 801 (1839) (*Hydrocampa*). — Stainton, Man. Brit. Butt. Moths, 2, p. 144 (1859). — Lederer, Wien. Ent. Monatschr., 7, p. 454 (1863). — Heine-mann, Kleinschm., 1, H. 2, p. 107 (1865). — Wallengren, Oefvers. K. Vet.-Akad. Förh., 28 (8), p. 1009 (1871). — Berce, Faune Ent. Fr. Lep., 6, p. 114, t. 3, f. 11 (1878). — Snellen, Vlinders Nederl., 2 (1), p. 78 (1882). — Leech, Brit. Pyral. et Pteroph., p. 47, t. 6, f. 6 (1886). — Meyrick, Handb.
- Bombay),
Ceylon,
Borneo,
Celebes,
Sumatra,
Java,
Amboina;
Neu-Guinea,
Fidji Ins.,
Bismarck-
Arch.;
Portug.
O.-Afrika,
Goldküste,
Br. Zentr.-
Afrika,
Mashona-
land,
S. Nigeria,
Kapkolonie
- Süd- u.
Ostchina
(Lienping,
Canton,
Kwangtung)
- Sangir,
Tonkin
- Mittel- u.
Nordeuropa
(England,
Irland,
Schweden,
Finnland,
Frankreich,
Belgien,
Italien,
Rumänien,
Bulgarien,
Thrazien,
Mazedonien,
Süd-
russland),
Kleinasien,
Lenkoran,
Armenien,
Balearen

- Brit. Lep., p. 402 (1895). — Hampson, Trans. Ent. Soc. Lond., p. 148 (1897). — Sharp in Harmer & Shipley, The Cambridge Nat. Hist., 6 (Ins. 2), p. 423 (1899). — Rebel in Spuler, Schm. Eur., 2, p. 221, t. 81, f. 20 (1906). — Wahlgren, Svensk Insektafauna, 10 (2), p. 47, t. 2, f. 53 (1915). — Wesenberg-Lund, Ins. Liv. Ferske Vand., p. 203, 204, f. 162 a—b (larva, chrysalis) (1915). — Tullgren & Wahlgren, Svenska Ins., p. 394, f. 315 (1922). — Börner in Brohmer, Fauna Deutschl., (ed. 3), p. 377 (1925); id., ibid., (ed. 4), p. 389 (1932). — Meyrick, Revis. Handb. Brit. Lep., p. 418 (1928). — M. Hering, Tierwelt Mitteleur. (Ergbd. 1), p. 282, f. 502 (1932). — Eckstein, Schm. Deutschl., 5, p. 62, t. 1, f. 21 (1933). — Karny, Biol. Wasserins., p. 166, 174, f. 108 B, C (larva, chrysal.) (1934).
- lemnalis* [Schiffermüller et Denis], Verz. Schm. Wien, p. 123 (1775) (*Pyrallis*). — Hübner, Samm. Eur. Schm., p. 48, t. 13, f. 83, 84 (1796) (*Pyrallis*); id., Verz. bekannt. Schmett., p. 363, n. 3468 ([1816]—1826) (*Cataclysta*). — Ochsenheimer & Treitschke, Schm. Eur., 7, p. 134 (1829) (*Nymphula*); 10 (3), p. 30 (1835) (*N.*). — Dup., Hist. Nat. Léop., 8 (2), p. 179, t. 222, f. 4, 5 (1831). — Eversmann, Fauna Lep. Volgo-Ural., p. 464 (1844). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 7 (1848). — Guenée, Delt. et Pyral., p. 267 (1854). — Freyer, Neuere Beitr. Schm. Kunde, 7, p. 140, t. 680, f. 1 (1858) (*Pyrallis*). — Cannaviello, Bol. Soc. Esp. Hist. Nat., 2, p. 172 (1902) (*Hydrocampa*). — Barrett, Lep. Brit. Isl., 9, p. 284, t. 107, f. 4, 4a, b (larva) (1904).
- uliginata* Fabricius, Ent. Syst., 3 (2), p. 214, n. 317 (1794) (*Phalaena*) (Dania).
- var. *ochracea* Hauder, Verh. z. h. Ges. Wien, 60, p. (8) (1910). — Rebel et Zerny, Denkschr. Akad. Wien, 103, p. 130 (1931). Ober-
Österreich,
Albanien
- var. *confirmata* Krulikovsky, Mater. Faun. Flor. Ross. Moskau, 9, p. ? (1909). — Peters, Lep. Fauna Estland, 2. Aufl., p. 399 (1924). Wjatka,
Ural, N.O.-
Persien
- subsp. *brunneospersa* Osthelder, Mitt. Münchner Ent. Ges., 24, p. 95 (1935). N. Syrien
- Anatomie: Griffith, Trans. Ent. Soc. Lond., p. 129, t. 4, f. 16 (1898) (frenulum).
- Biologie: Treitschke, Schm. Eur., 7, p. 134 (1829). — Buckler, Ent. Mon. Mag., 12, p. 102 (1875); Larvae Brit. Butt., 9, p. 74, t. 151, f. 1 (1901). — Sorhagen, Allg. Zeitschr. Ent., 6, p. 277 (1901). — Rebel in Spuler, Schm. Eur. Nachtr.-Taf. 9, f. 54 (1906). — Chapman, Entom., 38, p. 1—5, 38—43, t. 1, f. 1—9 (1905). — Solowiow, Zeitschr. wiss. Insektenbiol., 19, p. 174—

176 (1924). — Natanson-Grodzinska, C. R. Acad. Cracovie, nr. 7, p. 9 (1932); Bull. Int. Acad. Cracovie B 2, p. 132—140, fig. (1932).

- midas** Butler, Trans. Ent. Soc. Lond., p. 585 (1881). Amur,
— Preyer, Trans. Asiat. Soc. Jap., 13, p. 62 Ussuri,
(1885). — Hampson, Trans. Ent. Soc., p. Zentral-
152 (1897). — Leech, Trans. Ent. Soc. Lond., china,
p. 435 (1901). — Shibuya, Ins. Matsum., 3, Japan,
p. 128, t. 5, f. 13 (1929). Korea

b) Orientalische Region.

blandialis Walker, vide sub a) Paläarktische Region, p. 94.

amboinalis Hampson, Ann. Mag. Nat. Hist., (s. 8) Amboina
19, p. 374 (1917).

fuscalis Hampson, Ill. Typ. Sp. Lep. Het. Brit. Mus., 9, p. 178, t. 174, f. 13 (♀) (1893); Bhutan,
Fauna Brit. India Moths, 4, p. 197 (♀) Südchina,
(1896); Trans. Ent. Soc. Lond., p. 148 Tonkin,
(1897). Ceylon,
Borneo

nyctopis Meyrick, Trans. Ent. Soc. Lond.,
p. 475 (1894).

atriterminalis Hampson, Ann. Mag. Nat. Hist., (s. 8), 19, p. 458 (♀) (1917). Celebes,
Sangir

drusiualis Walker, Cat. Lep. Het. Brit. Mus., 17, p. Borneo,
450 (♂) (1859). (Borneo). — Hampson, Australien,
Trans. Ent. Soc. Lond., p. 148 (1897) (C. Queensland,
Anydraula). Fidji

drusialis Lederer, Wien. Ent. Monatschr.,
7, p. 454 (1863). — Meyrick, Trans. Ent.
Soc. Lond., p. 429 (1885) (*Anydraula*).

meropalis Walker, Cat. Lep. Het. Brit. Mus., 17, Borneo
p. 462 (♀) (1859) (*Hydrocampa?*). — Hamps., (Sarawak)
Trans. Ent. Soc. Lond., p. 223 (1897) (*Hydrocampa?*). — Swinhoe, Cat. East. Austral.
Lep. Het. Oxford Mus., 2, p. 440 (1900).

actoralis Walker, Cat. Lep. Het. Brit. Mus., 19, p. 962 (♀) (1859) (*Hydrocampa?*).
— Hampson, Trans. Ent. Soc., p. 223 (1897) (*Hydrocampa?*).

trimacula Hampson, Ill. Typ. Sp. Lep. Het. Brit. Mus., 8, p. 139, t. 155, f. 14 (1891); Fauna S. Indien,
Brit. India Moths, 4, p. 197 (♀) (1896); id., Nilgiris
Trans. Ent. Soc. Lond., p. 151 (1897).

angulata Moore, Lep. Ceylon, 3, p. 300, t. 181, f. Ceylon,
12 (1885). — Hampson, Fauna Brit. India Formosa
Moths, 4, p. 196 (♂) (1896); id., Trans.
Ent. Soc. Lond., p. 152 (1897). — Shibuya,
Journ. Fac. Agr. Hokkaido Imp. Univ., 22,
p. 146, 147 (1928).

midas Butler, vide sub a) Paläarktische Region p. 97.

- croesusalis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 449 (♂) (1859). (Sarawak). — Hampson, Fauna Brit. India Moths, 4, p. 196 (♂) (1896); Trans. Ent. Soc. Lond., p. 191 (1897) (*Strepsinoma*). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 448, t. 7, f. 14 (1900) (*Strepsinoma*). — *trigonalis* Swinhoe, Ann. Mag. Nat. Hist., (s. 6) 16, p. 304 (1895). *croesalis* Meyrick, Trans. Ent. Soc. Lond., p. 86 (1897) (*Strepsinoma*). subsp. *angustalis* Caradja, Mem. Acad. Röm. Sect. Stiint., (s. 3), 3, mem. 7, p. 329 (1925). (Canton)

Species incertae sedis.

- manilensis** Sauber in Semper, Schm. Philipp, 2, p. 654 (1902) (*Nymphicula*). Luzon

c) Aethiopische und Madagassische Region.

- blandialis** Walker, vide sub a) Paläarktische Region, p. 94.
euclidialis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 392 (♂) (1906). Nigeria
nyasalis Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 457 (1917). Nyassaland

Species incertae sedis.

- argyrochrysalis** Mabilie, Ann. Soc. Ent. Fr., 68, p. 743 (1899) (1900). Madagaskar
callichromalis Mabilie, Bull. Soc. Zool. Fr., 3, p. 94 (1878). — Hampson, Trans. Ent. Soc. Lond., p. 153 (1897). Madagaskar
leroi Strand, Arch. Naturg., 8, A H. 10, p. 107 (1914). Sudan (Bahr el Abied)
pusillalis Saalmüller, Ber. Senckenb. Ges., p. 295 (1879); Lep. Madag., 2, p. 504 (1891). — Hampson, Trans. Ent. Soc. London, p. 153 (1897). Madagaskar
tripunctalis Snellen, Tijds. v. Ent., 15, p. 98, t. 8, f. 1 (1872). — Hampson, Trans. Ent. Soc., p. 153 (1897). — Strand, Ent. Mitt., 4, p. 185 (1915). Guinea, Kamerun

d) Australische Region.

- blandialis** Walker, vide sub a) Paläarktische Region, p. 94.
nigristriata Hampson, Ann. Mag. Nat. Hist., (s. 8), 19, p. 374 (1917). Holl. Neu-Guinea, Ross-Ins., Admiraltäts Ins., Louisiaden, St. Aignan)
queenslandica Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 376 (♂, ♀) (1917). Queensland

- drusiualis** Walker, vide sub b) Orientalische Region, p. 97.
- glyceerialis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 450 (♀) (1859). — Meyrick, Trans. Ent. Soc. Lond., p. 423 (1885) (*Anydraula*). — Hampson, Trans. Ent. Soc. Lond., p. 148 (1897). Australien, N. S. Wales
- lampetialis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 451 (♂) (1859). Australien (Moreton Bai)
- repetitalis** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 205 (1896). Queensland, Melville Ins.

Species incertae sedis.

- cyanolitha** Meyrick, Trans. Ent. Soc. Lond., p. 220 (♀) (1886) (*Anydraula*). — Hampson, Trans. Ent. Soc. Lond., p. 153 (1897). Fidji Ins.
- hexalitha** Meyrick, Trans. Ent. Soc. Lond., p. 219 (1886). — Hampson, Trans. Ent. Soc. Lond., p. 153 (1897). Fidji Ins.
- pericempa** Turner, Proc. Roy. Soc. Queensland, 27, p. 36 (♂) (1916). N. Australien (Port Darwin)
- psathyrodes** Turner, Trans. Roy. Soc. S. Austral., 32, p. 83 (♂) (1908). Queensland
- tamsi** nom. nov. Samoa
dialitha Tams (nec Dyar, 1914), Ins. Samoa, part. 3, fasc. 4, p. 259, t. 17, f. 6 (1935).

e) Nearktische Region.

- fulcalis** Clemens, Proc. Acad. Nat. Sci. Phil., 12, p. 217 (1860). — Hampson, Trans. Ent. Soc. Lond., p. 149 (1897) (*Cataclysta*). — Dyar, Journ. N. York Ent. Soc., 14, p. 94 (1906) (*Elophila*). — Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 382 (1906) (*Argyrectis*). — Dyar, Insec. Insc. Menstr., 5, p. 75 (1917) (*Elophila*). — Barnes et Mc Dunning, Contr. Nat. Hist. Lep. N. Amer., 4, p. 169 (1918). — Forbes, Lep. N. York, p. 580 (1923). — Comstock, Introd. Ent., (ed. 2), p. 649 (1925) (*Elophila*). Vereinigte Staaten (Massachusetts, N. York, New Jersey, Pennsylvania, Ohio, Maryland, Texas, Neumexiko, Arizona, Washington, Oregon, Texas), Jamaica
- angulatalis* Lederer, Wien. Ent. Monatsschr., 7, p. 453, 486 (1863).
 Biologie: Lloyd, Journ. N. York Ent. Soc., 27, p. 263, t. 25 (1929) (*Elophila*).
- saternalis** Dyar, Insec. Insc. Menstr., 5, p. 75 (♀) (1917) (*Elophila*). Texas, Arizona
- truekecalis** Dyar, Insec. Insc. Menstr., 5, p. 76 (♀) (1917) (*Elophila*). Nevada, Oregon

- opulentalis** Lederer, Wien. Ent. Monatschr., 7, p. 453, 486, t. 18, f. 7 (1863). — Hampson, Trans. Ent. Soc. Lond., p. 151 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 379 (1906) (*Argyractis*). — Dyar, Insec. Insc. Menstr., 5, p. 76 (1917).
- confusalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1334 (1865) (*Cataclysta*). (Nordamerika). — Barnes et Mc Dunnough, Contr. Nat. Hist. Lep. N. Amer., 2 (3), p. 133, t. 8, f. 11 (1913) (*Argyractis?*) (Panama); 2 (5), p. 215 (1914) (*Elophila*).
- divisalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1336 (1865) (*Cataclysta*) (Pará).
- avernalis** Grote, Trans. Kansas Acad. Sci., 8, p. 53 (1878) (*Chrysendeton*). — Fernald, Smith's List n. 4157 (1891) (*Cataclysta*). — Hampson, Trans. Ent. Soc. Lond., p. 153 (1897). — Mc Dunnough, Canad. Ent., 48, p. 28 (1916) (Arizona, N. Mexiko).
- fulicalis** (part.) Hampson, Trans. Ent. Soc. London, p. 149 (1897).
- moniligeralis** Lederer, Wien. Ent. Monatschr., 7, p. 454, 487, t. 18, f. 10 (1863) (*Cataclysta*). (N. Amerika). — Hampson, Trans. Ent. Soc. Lond., p. 149 (1897) (*Cataclysta*) (W. Indien, Honduras, Kolumbien); Ann. Mag. Nat. Hist., (s. 7) 18, p. 385 (1906) (*Argyractis*). — Dyar, Insec. Insc. Menstr., 14, p. 77 (1917) (*Elophila*).
- lossonalis** Dyar, Journ. Ent. Soc. N. York, 14, p. 93 (1906) (*Elophila*).
- claudialis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 437 (♂) (1869) (*Cataclysta*). (Vereinigte Staaten). — Fernald, Smith's List, nr. 4153 (1891) (*Chrysendeton*). — Hart, Bull. Ill. Stat. Lab. Nat. Hist., 4, p. 174 (1895) (*Chrysendeton*). — Hampson, Trans. Ent. Soc. Lond., p. 151 (1897) (*Cataclysta*). — Dyar, Journ. N. York Ent. Soc., 14, p. 92 (1906) (*Elophila*). — Barnes et Mc Dunnough, Contr. Nat. Hist. Lep. N. Amer., 2, p. 215 (1914) (*Elophila*).
- brunnildalis** Dyar, Journ. N. York Ent. Soc., 14, p. 91 (♀) (1906) (*Elophila*).
- bifascialis** Robinson, Ann. Lyc. Nat. Hist. N. York, 9, p. 153, t. 4, f. 3 (1869). — Hampson, Trans. Ent. Soc. Lond., p. 149 (1897) (*Cataclysta*); Ann. Mag. Nat. Hist., (s. 7) 18, p. 376 (1906) (*Argyractis*). — Dyar, Journ. N. York Ent. Soc., 14, p. 91 (1906) (*Elophila*). — Forbes, Lep. N. York, p. 580 (1923).
- ab. kearfottalis** Dyar, Journ. N. York Ent. Soc., 14, p. 92 (1906) (*Elophila*).
- drumalis** Dyar, Journ. N. York Ent. Soc., 14, p. 92 (♀) (1906) (*Elophila*).
- medicinalis** Grote, Papilio, 1, p. 15 (1881) (*Cataclysta*) (Illinois). — Hampson, Trans. Ent.
- Vereinigte Staaten (Pennsylvania, Arizona, Neu-Mexiko), Mexiko, Brit. Colombien, Guayana, Brasilien
- Florida, Westindien, Honduras, Kolumbien
- Illinois, Florida, N. Carolina, Distr. Columbia, Pennsylvania, St. Domingo
- Kalifornien
- N. York bis Wisconsin, Maryland, Virginia, Texas, N. Mexiko
- Arizona (Phoenix)
- Florida (Fort Drum)
- Illinois, Missouri,

- Soc. Lond., p. 149 (1897). (*Cataclysta*). — Distr.
Barnes et Mc Dunnough, Contr. Nat. Columbia
Hist. Lep. N. Amer., 2 (5), p. 215 (1914). —
Forbes Lep. N. York, p. 579 (1923).
claudialis Dyar, Journ. N. York Ent. Soc.,
14, p. 92 (1906).
- magnificalis** Hübner, Samml. Eur. Schm., 6. Horde Kanada
Zünsler, p. 18, t. 16, f. 104 (1796) (*Pyralis*). (Quebeck),
(England err.); id., Verz. bek. Schmett., Vereinigte
p. 363, n. 3469 ([1816]—1826) (*Cataclysta*). — Staaten
Treitschke, Schm. Eur., 7, p. 140 (1829) (Massa-
(*Nymphula*). — Stephens, Ill. Brit. Ent. chussets,
Haust., 4, p. 395 (1835) (*Cataclysta?*). — Pennsylv-
Hampson, Trans. Ent. Soc. Lond., p. 148 vanien,
(1897) (*Cataclysta*). — Dyar, Journ. N. York N. York,
Ent. Soc., 14, p. 93 (1906) (*Elophila*). — Maine)
Forbes, Lep. N. York, p. 579 (1906) (*Elo-
phila*).
- lamialis* Walker, Cat. Lep. Het. Brit. Mus.,
17, p. 436 (♂) (1859) (Vereinigte Staaten).
— Fernald, Smith's List., nr. 4154 (1891)
(*Chrysendeton*).
- heliopalisis* Clemens, Proc. Acad. Nat. Sci.
Phil., 12, p. 218 (1860) (*Cataclysta?*). —
Grote, New Ch. List., p. 54 (1882) (*Chry-
sendeton*).
- helopalisis* Hampson, Trans. Ent. Soc. Lon-
don, p. 149 (1897) (ex-synon.).
- cronialis** Druce, Biol. Centr.-Amer. Het., 2, 274, t. Mexiko
63, f. 18 (♂) (1896). — Hampson, Trans. (Typus:
Ent. Soc. Lond., p. 149 (1897) (*Cataclysta*); Jalapa);
Ann. Mag. Nat. Hist., (s. 7) 18, p. 381 Arizona
(1906) (*Argyractis*). — Dyar, Journ. N.
York Ent. Soc., 14, p. 95 (1906) (*Elophila*).
- schaefferalis** Dyar, Journ. N. York Ent. Soc., 14, p. Arizona
95 (♀) (1906) (*Elophila*). (Palmerlee)
- daemonialis** Dyar, Proc. Ent. Soc. Wash., 10, p. Pennsylv-
112 (1908). vanien

Species incertae sedis.

- cancellalis** Dyar, Insec. Insc. Menstr., 5, p. 77 Texas
(♀) (1917) (*Elophila*).
- imitabilis** Dyar, Insec. Insc. Menstr., 5, p. 78 (♂) Florida
(1917) (*Elophila*).
- irroratalis** Dyar, Insec. Insc. Menstr., 5, p. 79 (♀) Florida
(1917) (*Elophila*).
- plevie** Dyar, Insec. Insc. Menstr., 5, p. 78 (♀) Mexiko,
(1917) (*Elophila*). N. Carolina

f) Neotropische Region.

- fulicalis** Clemens, vide sub e) Nearktische Region p.
99.
- opulentalis** Lederer, vido sub o) Nearktische Re-
gion p. 100.
- moniligeralis** Lederer, vide sub e) Nearktische Re-
gion p. 100.

- claudialis** Walker, vide sub e) Nearktische Region p. 100.
- romanalis** Druce, Biol. Centr.-Amer. Het., 2, p. 273, t. 63, f. 14 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 149 (1897). Mexiko (Typus: Amula in Guerrero)
- cronialis** Druce, vide sub e) Nearktische Region p. 101.
- tessellalis** Hampson, Trans. Ent. Soc. Lond., p. 149 (1897). Brasilien (S. Paulo)
- cumalis** Druce, Biol. Centr.-Amer. Het., 2, p. 273, t. 63, f. 15 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 150 (1897). Guatemala, Panama (Typus: El Reposo (Guatem.), Chiriqui Vulkan (Costa Rica))
- divulsalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1336 (♂) (1865). — Hampson, Trans. Ent. Soc. Lond., p. 150 (1897). Brasilien, Amazonas (Typus: Ega)
- chalcitis** Felder et Rogenhofer, Reise Novara (Het.), p. 5, t. 136, f. 9 (1874). — Hampson, Trans. Ent. Soc. Lond., p. 150 (1897). Brasilien
- guenealis** Snellen, Tijds. v. Ent., 18, p. 260, t. 14, f. 13 (1875) (*Paraponyx*).
guenealis Hampson, Trans. Ent. Soc. Lond., p. 150 (1897). Brasilien, Argentinien
trilinealis Warren, Trans. Ent. Soc. Lond., p. 295 (1889).
- melaternalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 393 (♀) (1906). Brasilien
- glycysalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 288 (1914) (*Cataclysta*). Panama Costa Rica, Fr. Guayana
- pulcherialis** Druce, Biol. Centr.-Amer. Het., 2, p. 274, t. 63, f. 16 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 151 (1897). Costa Rica (Typus: Vulkan de Irazu)
- ilialis** Walker, Cat. Lep. Het. Brit. Mus., 17, 441 (♀) (1859) (Pará). — Hampson, Trans. Ent. Soc. Lond., p. 151 (1897). Brasilien (Typus: Vera Paz)
- pomperialis** Druce, Biol. Centr.-Amer. Het., 2, p. 276, t. 63, f. 24 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 152 (1897). Guatemala, Costa Rica (Typus: Vulkan Irazu)
- obliquifascia** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 458 (♀) (1917). Brasilien (Pernambuco)
- chionostola** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 458 (♀) (1917). Peru

- sumptuosalis** Möschler, Abh. Senckenb. Ges., 16, p. 319 (1890). — Hampson, Trans. Ent. Soc. Lond., p. 153 (1897). Portorico
- azadasalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 110 (♂) (1924). Brit. Guyana
Typus: Essex
nebo-Fluss,
Rockstone
- mangholdalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 110 (♂) (1924). Ecuador
(Typus:
Zaruma)
- phocosalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 111 (♂) (1924). Peru
- miralis** Möschler, Abh. Senckenb. Ges., 16 (1), p. 319 (1896). Portorico
- anicctalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 111 (♂) (1924). Cuba
(Typus:
Santiago)
- bronachalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 112 (♂) (1924). Cuba oder
Porto Rico
- peraltalis** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, p. 673 (♂) (1912). — Dyar, Proc. U. S. Nat. Mus., 47, p. 289 (1914). Costa Rica
- argyrolepta** Dyar, Proc. U. S. Nat. Mus., 47, p. 289 (1914). Panama
- iolepta** Dyar, Proc. U. S. Nat. Mus., 47, p. 290 (♀) (1914). Panama
- unilinealis** Dyar, Proc. U. S. Nat. Mus., 47, p. 290 (♂) (1914). Panama
- autobella** Dyar, Proc. U. S. Nat. Mus., 47, p. 291 (♀) (1914). Panama
- vacuolata** Dyar, Proc. U. S. Nat. Mus., 47, p. 291 (♂) (1914). Panama
- dialitha** Dyar, Proc. U. S. Nat. Mus., 47, p. 291 (♂) (1914). Panama
- zelota** Dyar, Proc. U. S. Nat. Mus., 47, p. 292 (♀) (1914). Panama

Species incertae sedis.

- bijonalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 287 (♀) (1914). Panama
- complicatalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 289 (♀) (1914). Panama
- dasymalis** Dyar, Insec. Insc. Menstr., 14, p. 148 (1926). Mexiko
- jaliscalis** Schaus, Proc. U. S. Nat. Mus., 40, p. 135 (1906). Mexiko
- minimalis** Herrich-Schaeffer, Ver. Regensb., 25, p. 26 (1871) (Sep. p. 65). Cuba
- scalaris** Schaus, Proc. U. S. Nat. Mus., 30, p. 136 (♀) (1906). Brasilien
(Paraná)

9. Genus **Metaclysta** Hampson

Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 455 (1906).

tetrommata Hampson, Ann. Mag. Nat. Hist., (s. 7) Louisiaden
18, p. 455 (♂) (1906). (St. Aignan)

1

10. Genus **Symphonia** Hampson

Hampson, Fauna Brit. India Moths, 4, p. 198 (1896); Trans.
Ent. Soc. Lond., p. 132, 153 (1897).

(Typus generis: *multipictalis* Hamps.)

multipictalis Hampson, Fauna Brit. India Moths, Ceylon
4, p. 195, f. 117 (♂) (1896); Trans. Ent.
Soc. Lond., p. 154, fig. (♂) (1897).

synophanoses Klunder van Gijen, Tijd. v. Ent., Java
55, p. 350 (1912) (1913).

albiocularis Hampson, Ann. Mag. Nat. Hist., (s. 7) S.-Nigeria
18, p. 391 (♂) (1906).

leucostictalis Hampson, Ann. Mag. Nat. Hist., (s. 7) Brit.
18, p. 456 (♂) (1906). Neu-Guinea

trivitalis Warren, Ann. Mag. Nat. Hist., (s. 6) Indien,
16, p. 302 (1895) (*Bocchoris*) (Assam). — Assam
Hampson, Fauna Brit. India Moths, 4, p. (Khasis),
282 (♂) (1896) (*Bocchoris*). — Shibuya, Formosa,
Journ. Fac. Agr. Hokkaido Imp. Univ., 22, Ceylon
p. 24, 147 (1928) (*Symphonia*).

fenestralis Warren, Ann. Mag. Nat. Hist.,
(s. 6) 17, p. 105 (1896) (*Synclera*).

5

11. Genus **Musofima** Meyrick

Meyrick, Trans. Ent. Soc. Lond., p. 288 (1884). — Ragonot, Ann.
Soc. Ent. Fr., (s. 6) 10, p. 458 (1890) (1891). — Hampson, Fauna
Brit. India Moths, 4, p. 194 (1896); Trans. Ent. Soc. Lond.,
132, 154 (1897). — Meyrick, Trans. Proc. N. Zeal. Inst., 45, p. 38
(1912) (1913). — Hudson, Butt. Moths N. Zeal., p. 177 (1928).

? *Diathrausta* Felder & Rogenhofer, Reise Novara, (Het.),
p. 8 (1874) (sine fixat. typi generis).

(Typus generis: *aduncalis* Feld. et Rogenh.)

suffusalis Hampson, Ill. Het. Typ. Sp. Brit. Mus., Süd-China,
9, p. 178, t. 174, f. 20 (1893); Fauna Brit. Süd-Indien
India Moths, 4, p. 199 (1897); Trans. Ent. (Bangalore),
Soc. Lond., p. 155 (1897). Ceylon,
Luzon

fuscalis Snellen, Tijd. v. Ent., 43, p. 298, t. 17, f. 2, Java,
3 (♂) 1901). Assam

- francki** Caradja, Mom. Acad. Rom. Sect. Stiint., (s. 3) 4, mem. 8, p. 404 (1927) (*franckeii* err. typ.). Westchina
- aduncalis** Felder et Rogenhofer, Reise Novara (Het.), p. 8, t. 135, f. 11 (1874) (*Dia-thrausta*?). — Meyrick, Trans. Ent. Soc. Lond., p. 289 (1884) (*Musotima*). — Hampson, Trans. Ent. Soc. Lond., p. 155 (1897). — Hudson, Butt. Moths, N. Zeal., p. 177, t. 19, f. 19 (♂) 20 (♀) (1928). Neuseeland
- decoralis** Snellen, Tijds. v. Ent., 43, p. 299, t. 17, f. 4, 5 (♂) (1901). Java
- accleralis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 403 (♂) (1859) (*Isopteryx*) (Ceylon). — Moore, Lep. Ceyl., 3, p. 303 (1886) (*Cy-moriza*). — Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 180, t. 174, f. 24 (1893) (*accleralis*); Fauna Brit. India Moths, 4, p. 200, f. 118 (♂) (1896); Trans. Ent. Soc. Lond., 155, fig. (♂) (1897). — Marumo, Journ. Col. Agr. Tokyo Imp. Univ., 8, p. 187 (1923). — Shibuya, Ins. Matsum., 3, p. 135 (1929). Japan, Indien (Nilgiris), Ceylon
- nubilalis** R. South, Trans. Ent. Soc. Lond., p. 435, t. 14, f. 27 (1901). — Caradja, Mem. Acad. Rom. Sect. Stiint., (s. 3) 3, mem. 7, p. 329 (1925) (? = *accleralis*); Bull. Acad. Roum., 15, p. 147 (1932). Zentral-China
- ochropteralis** Guenée, Delt. et Pyral., p. 230 (1854) (*Isopteryx*) (Patria?). — Hampson, Trans. Ent. Soc. Lond., p. 155 (1897). Australien, N.S.Wales, Tasmanien
- acrias** Meyrick, Trans. Ent. Soc. Lond., p. 289 (1884). — Hampson, Trans. Ent. Soc. Lond., p. 155 (1897). Australien, Tasmanien
- persinualis** Hampson, Trans. Ent. Soc. Lond., p. 155 (1897). Amboina, Tenimber, Fergusson Ins., Queensland
- fuscidiscalis** Hampson, Trans. Ent. Soc. Lond., p. 156 (1897). Fergusson Ins.
- nitidalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1317 (♀) (1865) (*Isopteryx*). — Meyrick, Trans. Ent. Soc. Lond., p. 290 (1884). — Hampson, Trans. Ent. Soc. Lond., p. 155 (1897). — Hudson, Butt. Moths N. Zeal., p. 177, t. 19, f. 18 (♂), tab. frontisp., f. 22 (1928) (ovum). Queensland, N.S.Wales, Victoria, West-Australien, Auckland-Ins., Neuseeland
- timaralis** Felder et Rogenhofer, Reise Novara, (Het.), p. 8, t. 135, f. 23 (1874) (*Dia-thrausta*). (Neuseeland) (Typus: „Australia“)
- Biologie: Chapell, Trans. Proc. N. Zeal. Inst., 60, p. 560—562, f. 1—4 (1930).

Species incertae sedis.

- callidryas** Turner, Proc. Roy. Soc. Victoria, 35, p. 49 (♂, ♀) (1922). Queensland

- inerustalis** Snellen, Tijds. v. Ent., 38, p. 159 (1894). Java,
— E. Hering, Ent. Zeit. Stettin, 64, p. 54 Fergusson
(1903). Ins.
- instrumentalis** Swinhoe, Ann. Mag. Nat. Hist., (s. Assam
6) 14, p. 209 (1894) (*Ambia*). — Hampson, (Khasis)
Fauna Brit. India Moths, 4, p. 204 (♂)
(1896) (*Ambia*); Trans. Ent. Soc. Lond.,
p. 160 (1897) (*Ambia*). — Snellen, Tijds. v.
43, p. 300, t. 17, f. 6 (1900) (*Musotima*).
- leptorrhoda** Turner, Trans. Proc. Roy. Soc. S. Queensland
Austral., 32, p. 85 (♂, ♀) (1908).
- stictochroa** Turner, Proc. Roy. Soc. Queensland, 27, N.S.Wales
p. 37 (♀) (1915).

17

12. Genus **Baeophila** Turner

Turner, Trans. Roy. Soc. S. Austral., 32, p. 81 (1908).

Ambia (part.) Hampson, Trans. Ent. Soc. Lond., p. 159
(1897) (*A. albipunctalis* Warr.).

(Typus generis: *selenias* Turn.)

- selenias** Turner, Trans. Roy. Soc. S. Austral., 32, Queensland
p. 81 (♂) (1908).
- albipunctalis** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, Assam
p. 202 (1896). — Hampson, Fauna Brit. (Khasis),
India Moths, 4, p. 203 (♀) (1896) (*Ambia*); Ceylon
Trans. Ent. Soc. Lond., p. 159 (1897) (*Ambia*).
bia).
- ellipes** Tams, Ins. Samoa, pars 3 (fasc. 4), p. 259, Samoa
t. 17, f. 5 (1935).

3

13. Genus **Panofima** Meyrick

Meyrick, Exot. Microlep., 4, p. 540 (♂) (1934).

- copidosema** Meyrick, Exot. Microlep., 4, p. 540 Belg. Congo
(♂) (1934). (Gebirge b.
1 Nyamu-
kubi)

14. Genus **Cosmophylla** Turner

Turner, Trans. Roy. Soc. S. Austral., 32, p. 85 (1908).

- oxygramma** Turner, Trans. Roy. Soc. S. Austral., Queensland
32, p. 86 (♂) (1908).

1

15. Genus **Eugauria** Snellen

Snellen, Tijds. v. Ent., 43, p. 290 (1901). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 24 (1928).

Cataclysta (part.) Hampson, Trans. Ent. Soc. London, p. 153 (1897) (*C. albidentata* Hamps.).

albidentata Hampson, Trans. Ent. Soc. Lond., p. 153 (♀) (1897) (*Cataclysta*) (Java). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 24, 147 (1928). Java, Formosa

compactalis Snellen, Tijds. v. Ent., 43, p. 291, t. 16, f. 10 (1901) (Java).

1

16. Genus **Susumia** Marumo

Marumo, Oyo Dobuts. Zasshi, 2, p. 41 (1930) [nov. gen.].

Epimima (part.) Meyrick, Trans. Ent. Soc. Lond., p. 235 (1886) [= *Marasmia* Led., Pyraustinae]. — Shibuya, Ins. Matsum., 3, p. 135 (1929).

Bradynomorpha Matsumura, Inj. Ins. Jap. Dainippon Gaichuzensho, p. 514 (1920) (sine descr.).

(Typus generis: *exigua* Butl.)

daisensis Shibuya, Ins. Matsum., 3, p. 135, t. 5, f. 28 (♂) (1929). Japan
(Typus:
Honshu
(Mt.
Daisen)

nawae Matsumura, Inj. Ins. Dainippon Gaichuzensho, p. 514, t. f. 188 (1920) (*Bradynomorpha*). — Shibuya, Ins. Matsum., 3, p. 135, t. 5, f. 21 (1929) (*Epimima*). Japan

admixtalis Matsumura, Oyo Konchugaku, p. 549, t. 21, f. 10 (1920) (*Bradina*).

exigua Butler, Ann. Mag. Nat. Hist., (s. 5) 4, p. 453 (1879) (*Samoa*). — Hampson, Proc. Zool. Soc. Lond., p. 639 (1898) (*Marasmia*). — Leech, Trans. Ent. Soc., p. 453 (1901) (*Marasmia*). — Matsumura, Cat. Ins. Jap., p. 208 (1905). — Shibuya, Ins. Matsum., 3, p. 135 (1929) (*Epimima*). — Marumo, Oyo Dobuts. Zasshi, 2, p. 41, fig. (1930) (*Susumia*). Japan

3

17. Genus **Cymoriza** Guenée

Guenée, Delt. et Pyral., p. 271 (1854). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 455 (1859). — Lederer, Wien. Ent. Monatschr., 7, p. 453 (1863). — Moore, Lep. Ceylon, 3, p. 302 (1885). — Hampson, Fauna Brit. India Moths, 4, p. 200 (1896); Trans. Ent. Soc. Lond., p. 133, 156 (1897).

Cymorrhiza Hering, Ent. Zeitg. Stettin, 64, p. 54 (1903).

(Typus generis: *irrectalis* Guen.)

- irrectalis** Guenée, Delt. et Pyral., p. 272, part. (1854) N.O.Indien,
(Ostindien). — Hampson, Fauna Brit. India Bhotan,
Moths, 4, p. 201 (♀) (1896); Trans. Ent. Sikkim,
Soc. Lond., p. 157 (1897). — E. Hering, Ent. Südechina,
Zeit. Stettin, 64, p. 54 (1903) (*Cymorrhiza*). Formosa,
— Shibuya, Journ. Fac. Agr. Hokkaido Imp. Sumatra
Univ., 22, p. 148, t. 5, f. 1 (1928).
- hoenel** Caradja, Deutsche Ent. Zeit. Iris, 47, p. 154 Ostchina
(1933). (Kwangtung)
- interruptalis** Wileman et R. South, Entom., 50, Formosa
p. 175 (1917) (*Ambia*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 24,
148 (1928).
- taiwanalis** Shibuya, Journ. Fac. Agr. Hokkaido Imp. Formosa
Univ., 22, p. 148, t. 5, f. 9 (♂) (1928):
- ustalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. Himalaya,
1529 (♀) (1865) (*Oligostigma*) (Java). — Dharmasala,
Hampson, Fauna Brit. India Moths, 4, p. Kulu,
201, f. 119 (♂) (1896); Trans. Ent. Soc., Sikkim,
p. 157; p. 156 (fig. ♂) (1897). Java,
irrectalis (part.), Guenée, Delt. et Pyral., Brit.
p. 272, t. 9, f. 4 (1854), (Ostindien). Neu-Guinea
fascialis Warren, Ann. Mag. Nat. Hist.,
(s. 6) 18, p. 223 (1896) (Khasis).

Species incertae sedis.

- gigantalis** Janse, Bull. Hill Mus., 1, p. 500 (♂) Ceram
(1924).
- isorrhoa** Meyrick, Exot. Microlep., 5, p. 7 (♂) Witu
(1936). (French Isl.)
- paraolis** Schaus, Proc. U. S. Nat. Mus., 30, p. 137 Brasilien
(1906). (S. Paulo)

8

18. Genus **Pseudolifhosia** Hampson

Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 1 (1907).

- sehausi** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, Mexiko
p. 1 (♂, ♀) (1907).

1

19. Genus **Omphaloptera** Hampson

Hampson, Trans. Ent. Soc. Lond., p. 133, 157 (1897).

(Typus generis: *randalis* Druce.)

- completalis** Dognin, Ann. Soc. Ent. Belg., 48, p. Peru
126 (♀) (1904).
- randalis** Druce, Biol. Centr.-Amer. Het., 2, p. 275, Mexiko
t. 63, f. 21 (1896) (*Hydrocampa*). — Hamp- (Typus:
son, Trans. Ent. Soc. Lond., p. 157, fig. Teapa in
(♂) (1897). Tabasco)
- stigmatalis** Dognin, Ann. Soc. Ent. Belg., 48, p. 127 Peru
(♂) (1904).

virginalis Dognin, Ann. Soc. Ent. Belg., 48, p. 126 Ecuador
(♀) (1904).

4

20. Genus **Thysanoidma** Hampson

Hampson, Ill. Typ. Sp. Het. Brit. Mus., 8, p. 142 (1891); Fauna
Brit. Ind. Moths, 4, p. 202 (1896); Trans. Ent. Soc. Lond., p.
132, 157 (1897).

octalis Hampson, Ill. Typ. Sp. Het. Br. Mus., 8, Assam,
p. 142, t. 156, f. 19 (1891); Fauna Brit. Indien,
India Moths, 4, p. 202, f. 120 (♀) (1896); (Nilgiris),
Trans. Ent. Soc. Lond., p. 158, fig. (♀) Südchina,
(1897). — Shibuya, Journ. Fac. Agr. Hok- Formosa
kaido Imp. Univ., 22, p. 149 (1928).

1

21. Genus **Teternia** Meyrick

Meyrick, Proc. Linn. Soc. N. S. Wales, s. 2, 4, p. 109 (1890). —
Hampson, Trans. Ent. Soc. Lond., p. 132, 158 (1897).

terminitis Meyrick, Proc. Linn. Soc. N. S. Wales, Australien
s. 2, 4, p. 1109 (1890). — Hampson, Trans.
Ent. Soc. Lond., p. 158, fig. (♂) (1897).

1

22. Genus **Ephormotris** Meyrick

Meyrick, Exot. Microlep., 4, p. 393 (♂) (1933).

octopis Meyrick, Exot. Microlep., 4, p. 393 (♂) Malakka
(1933). (Typus:

1

Kuala
Lumpur)

23. Genus **Parambia** Dyar

Dyar, Proc. U. S. Nat. Mus., 47, p. 292 (1914).

(Typus generis: *gnomosynalis* Dyar.)

gnomosynalis Dyar, Proc. U. S. Nat. Mus., 47, p. Panama,
292 (1914). Fr. Guyana

glenealis Dyar, Proc. U. S. Nat. Mus., 47, p. 292 Panama,
(1914). Fr. Guyana

cedroalis Schaus, Proc. Ent. Soc. Wash., 26, p. Cuba
116 (♀) (1924). (Typus:

3

Alto del
Cedro (ad
Santiago))

24. Genus **Ambia** Walker

Walker, Cat. Lep. Het. Brit. Mus., 19, p. 957 (♂) (1859). — Hampson, Fauna Brit. India Moths, 4, p. 202 (1896) (part.). — Trans. Ent. Soc. Lond., p. 132, 159 (1897). — Dyar, Journ. N. York Ent. Soc., 14, p. 89 (1906).

Opistheidicta Warren, Ann. Mag. Nat. Hist., (s. 6), 5, p. 478 (1890). — E. Hering, Ent. Zeit. Stettin, 64, p. 50 (1903).

Oligernis Meyrick, Trans. Ent. Soc. Lond., p. 470 (1894). — E. Hering, Ent. Zeitg. Stettin, 64, p. 50 (1903).

Leucogephyra Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 219 (1896).

(Typus generis: *ptolycusalis* Walk.).

a) **Paläarktische Region.**

- colonalis** Bremer, Lep. Ost-Sib., p. 67, t. 6, f. 4 (1864) (*Hydrocampa*). — Meyrick, Trans. Ent. Soc. Lond., p. 469 (1890) (*Musotima*). — Hampson, Fauna Brit. India Moths, 4, p. 204 (1896); Hampson, Trans. Ent. Soc. Lond., p. 159 (1897). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 150, t. 6, f. 27 (1928). Amur, Ostchina (Kwangtung), N.W.-Himalaya (Dharmasala), Formosa
- locuples** Butler, Ill. Typ. Sp. Het. Brit. Mus., 7, p. 97, t. 135, f. 7 (1889) (*Oligostigma*).
- dives** Butler, Ill. Typ. Spec. Het. Brit. Mus., 7, p. 97, t. 135, f. 8 (1889) (*Oligostigma*).
- thyridialis** Lederer, Verh. z. b. Ges. Wien, 5, p. 219, t. 4, f. 2 (1855) (*Nymphula*). — Hampson, Trans. Ent. Soc. Lond., p. 166 (1897). Syrien, Palästina, Ägypten

b) **Orientalische Region.**

- tenebrosalis** Hampson, Fauna Brit. India Moths, 4, p. 203 (♂) (1896); Trans. Ent. Soc. Lond., p. 159 (1897). Ceylon
- caeruleata** Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 179, t. 174, f. 19 (1893); Journal Bombay Nat. Hist. Soc., 18, p. 573 (1908). Ceylon
- cyanealis** Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 210 (♂) (1903).
- xantholeuca** Hampson, Fauna Brit. India Moths, 4, p. 203 (♀) (1896); Trans. Ent. Soc. Lond., p. 159 (1897). Ceylon, Borneo
- colonalis** Bremer, vide sub a) Paläarktische Region p. 110.
- leucochrysa** Meyrick, Trans. Ent. Soc. Lond., p. 471 (1894). — Hampson, Trans. Ent. Soc. Lond., p. 160 (1897). Borneo
- ptolycusalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 958 (♂) (1859) (*Ambia*) (Sarawak). — Hampson, Trans. Ent. Soc. Lond., p. 160 (1897). — E. Hering, Ent. Zeit. Stettin, 64, p. 51 (1903). Borneo, Celebes, Philippinen, Anstralien, Queensland

- eromenalis* Snellen, Tijds. v. Ent., 23, p. 276 (1880) (*Caenostola*); 27, p. 6, t. 3, f. 4 (1884) (*Caenostola*). — Meyrick, Trans. Ent. Soc. Lond., p. 206 (1887) (*Paraponyx*). — Snellen, Tijds. v. Ent., 43, p. 297 (1901) (*Thysanoidma*). — Swinhoe, Ann. Mag. Nat. Hist., (s. 7) 7, p. 135 (1901) (*Oligostigma*).
- endophthalma** Meyrick, Trans. Ent. Soc. Lond., p. 470 (1894). — Hampson, Trans. Ent. Soc. Lond., p. 160 (1897). Borneo.
- mesoscotalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 456 (1906). Banda Ins., Luisiaden (St. Aignan)
- palliolatalis** Swinhoe, Trans. Ent. Soc. Lond., p. 287 (1890). — Hampson, Fauna Brit. India Moths, 4, p. 204 (1896). Assam, Burma
- palliolatatis* Hampson, Trans. Ent. Soc. Lond., p. 160 (1897).
- complicata** Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 222 (1896). — Hampson, Journ. Bombay Nat. Hist. Soc., 12, p. 700 (♀) (1896); Trans. Ent. Soc. Lond., p. 160 (1897). Assam
- fulvobasalis** Snellen, Tijds. v. Ent., 23, p. 244 (1880) (*Cymoriza*). — Hampson, Trans. Ent. Soc. Lond., p. 160 (1897). Borneo, Celebes
- semifascialis** Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 219 (1896) (*Leucogephyra*). — Hampson, Journ. Bombay Nat. Hist. Soc., 12, p. 701 (1899); Trans. Ent. Soc., p. 161 (♂) (1897). Assam (Khasis)
- rhabdotalls** Hampson, Trans. Ent. Soc. Lond., p. 161 (♂) (1897). Amboina
- iambealis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 950 (♂) (1859) (*Oligostigma*). — Hampson, Fauna Brit. India Moths, 4, p. 205, f. 121 (♂, ♀) (1896). Assam, Ceylon
- iambesalis* Hampson, Trans. Ent. Soc. Lond., p. 162, p. 159, fig. (♂) (1897).
- fulvalis* Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 180, t. 174, f. 26 (♀) (1893) (*Musotima*).
- albiflavicalis* Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 223 (1896) (*Cymoriza*).
- interstrigalis** Hampson, Trans. Ent. Soc. Lond., p. 162 (1897). Amboina, Fergusson Ins., Neu-Guinea
- atristrigails** Hampson, Trans. Ent. Soc. Lond., p. 162 (♂) (1897). Amboina, Fergusson Ins., Neu-Guinea
- leucoplaca** Meyr., Trans. Cat. Ent. Soc. Lond., p. 85 (1897) (*Nymphula*). — Hampson, Trans. Ent. Soc. Lond., p. 163 (1897). Sangir, Fergusson Ins., Neu-Guinea

- marginalis** Moore, Desce. Lep. Ins. Atkinson, p. 211 (1887). — Hampson, Fauna Brit. India Moths, 4, p. 205 (1896); Trans. Ent. Soc. Lond., p. 164 (1897). — Caradja, Mem. Acad. Rom. Sect. Stiint, s. 3, 3, mem. 7, p. 330 (1925). N.O.-Indien, Sikkim, Assam, Ost- und Südchina, Borneo
- albicomma* Swinhoe, Ann. Mag. Nat. Hist., (s. 6) 14, p. 210 (1894) (*Cymoriza*).
- fulvalis* Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 224 (1896) (*Cymoriza*).
- albobasalis** Caradja, Deutsche Ent. Zeitschr. Iris, 47, p. 154 (1933). Ostchina (Kwangtung)
- magnificalis** Swinhoe, Ann. Mag. Nat. Hist., (s. 6) 16, p. 299 (1895). — Hampson, Fauna Brit. India Moths, 4, p. 206 (♂) (1896); Trans. Ent. Soc. Lond., p. 164 (1897). N.O.-Indien
- interruptalis* Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 224 (1896) (*Cymoriza*) (Khasis).
- amoyalis** Caradja, Mem. Acad. Rom. Sect. Stiint, s. 3, 3, mem. 7, p. 330, t. 1, f. 32 (1925). Südchina
- melanistis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 464 (♀) (1917). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 150 (1928). Formosa
- trichostylalis** Hampson, Trans. Ent. Soc. Lond., p. 164 (1897). Celebes
- poritialis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 951 (♀) (1859) (*Oligostigma*) (Patria?). — Moore, Lep. Ceylon, 3, p. t. 179, f. 11 (1885) (*Cymoriza*). — Hampson, Fauna Brit. India Moths, 4, p. 206 (1896); Trans. Ent. Soc. Lond., p. 164 (1897) (*A. Opisthedeicta*). — E. Hering, Ent. Zeit. Stettin, 64, p. 50, (1903). S.O.-Indien, Ceylon, Sumatra, Java, Celebes
- pedionoma** West, Nov. Zool., 36, p. 213 (♀) (1931). Philippinen
- lobophoralis** Hampson, Fauna Brit. India Moths, 4, p. 207 (♂) (1896). Assam, Sikkim, Pulo Laut
- iriusalis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 434 (1859) (*Oligostigma?*) (Sarawak). — Hampson, Trans. Ent. Soc. Lond., p. 165 (1897) (*A. (Opisthedeicta)*). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 441, t. 16, f. 19 (1900). — E. Hering, Ent. Zeit. Stettin, 64, p. 50 (1903). Borneo, Pulo Laut, Sumatra
- irialis* Meyrick, Trans. Ent. Soc. Lond., p. 470 (1894) (*Nymphula*).
- tigridalis** Hampson, Trans. Ent. Soc. Lond., p. 165 (♀) (1897). Pulo Laut
- cataclystalis** Hampson, Trans. Ent. Soc. Lond., p. 166 (♀) (1897). — E. Hering, Ent. Zeit. Stettin, 64, p. 51 (1903). Pulo Laut, Sumatra
- dohrni** E. Hering, Ent. Zeit. Stettin, 64, p. 51, t. 1, f. 30 (♀) (1903). Sumatra
- lipocosmalis** Snellen, Tijds. v. Ent., 43, p. 286, t. 16, f. 4, 5 (1900) (*Hydrocampa*). — E. Hering, Ent. Zeit. Stettin, 64, p. 50 (1903). Sumatra, W.-Java

c) Aethiopische Region.

- chalcichroalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 457 (♀) (1906). Kapkolonie
melanalis Hampson, Ann. Mag. Nat. Hist. (s. 7) 18, p. 458 (1906). Kapkolonie
albiflavalis Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 462 (♂) (1917). S.-Nigeria (Lagos)
albomaculalis Hampson, Trans. Ent. Soc. Lond., p. 165 (♀) (1897) (*A. (Opisthedeicta)*). Goldküste (Aburi)

d) Australische Region.

- tendicularis** Rebel, Mitt. Zool. Mus. Hamburg, 32, p. 135, t. 1, f. 1 (1915). — Trans., Ins. Samoa pars 3, fasc. 4, p. 260 (1935).
chrysogramma Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 459 (1917).
mesoseotalis Hampson, vide sub b) Orientalische Region p. 111.
cymophoralis Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 459 (♂) (1917). Louisiaden (St. Aignan);
oligalis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 457 (1906). Louisiaden (St. Aignan), Goodenough Ins.
oculalis Hampson, Trans. Ent. Soc. Lond., p. 160 (♂) (897). Australien (Typus: Mt. Macedon)
interstrigalis Hampson, vide sub b) Orientalische Region p. 111.
atristrigalis Hampson, vide sub b) Orientalische Region p. 111.
leucoplaca Meyrick, vide sub b) Orientalische Region p. 111.
fulvicolor Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 462 (♂) (1917). Brit. Neu-Guinea
fusalis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 466 (♂) (1906). Bali
vagilincalis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 461 (1906). Holl. Neu-Guinea, Louisiaden (St. Aignan)
aulacophora Hampson, Trans. Ent. Soc. Lond., p. 165 (1897). Fergusson Ins.

Species incertae sedis.

- klossi** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 119 (♂) (1915). Holl. Neu-Guinea
novaguincensis Kenrick, Proc. Zool. Soc. Lond., p. 519, t. 68, f. 6 (1912). Holl. Neu-Guinea
obliquistriga W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 119 (♂) (1915). Holl. Neu-Guinea

- parachrysis** Meyrick, Exot. Microlep., 4, p. 608 (♂) Fidji Ins.
(1935). (Typus: Wunidawa)
- punetimarginata** W. Rothschild, Rep. B. O. U. Holl.
Exp., 2, nr. 15, p. 118 (♂) (1915). Neu-Guinea
- rothschildi** nom. nov.
marginalis W. Rothschild, (nec Moore, Holl.
1887), Rep. B. O. B. Exp., 2, nr. 15, p. Neu-Guinea
119 (♂) (1915).
- schlostochaeta** Tams, Ins. Samos, pars 3 (fasc. 4), p. Samoa
261, t. 15, f. 8 (♂) (1935).

e) Nearktische Region.

- striatalis** Dyar, Journ. N. York Ent. Soc., 14, p. Florida
90 (♂, ♀) (1906). (Charlotte Harbor)

f) Neotropische Region.

- debalis** Druce, Biol. Centr.-Amer. Het., 2, p. 276, Guatemala
t. 63, f. 23 (♂) 1896) (*Hydrocampa*). — (Vera Paz),
Hampson, Trans. Ent. Soc. Lond., p. 160 Costa Rica
(1897). (Irazu),
Panama
(Chiriqui)
- dendalis** Druce, Biol. Centr.-Amer. Het., 2, p. 277, Guatemala
t. 63, f. 27 (1896) (*Hydrocampa*?). — (Typus: San
Hampson, Trans. Ent. Soc. Lond., p. 160 Geronimo)
(1897).
- cantiusalis** Schaus, Proc. Ent. Soc. Wash., 26, p. Peru
115 (♀) (1924). Col.
Perene)
- picalis** Hampson, Trans. Ent. Soc. Lond., p. 160 Brasilien
(1897). (S. Paulo)
- tenella** Hampson, Trans. Ent. Soc. Lond., p. 161 Peru
(♀) (1897). (Typus: Callao)
- rufitincta** Hampson, Ann. Mag. Nat. Hist., (s. 8) Peru
19, p. 460 (♀) 1917).
- hemigrammalis** Hampson, Ann. Mag. Nat. Hist., Bolivien
(s. 8) 19, p. 461 (♀) (1917).
- sufetulodes** Hampson, Ann. Mag. Nat. Hist., (s. 8) Peru
19, p. 461 (♀) (1917).
- perornatalis** Schaus, Ann. Mag. Nat. Hist., (s. 9) 8, Costa Rica
p. 677 (♀) (1912).
- semilunalis** Hampson, Trans. Ent. Soc. Lond., p. Brasilien
163 (♂) (1897). (Typus: Petropolis)
- niveiplagalis** Hampson, Ann. Mag. Nat. Hist., (s. Peru
8) 19, p. 463 (♂) (1917).
- fulvinctalis** Hampson, Trans. Ent. Soc. Lond., p. Peru,
163 (1897) (Callao, Peru). Costa Rica,
Panama

- bolusalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 957 (♂) (1859) (*Cymoriza*). — Hampson, Trans. Ent. Soc., p. 163 (1897). Brasilien
- flavalis** Warren, Trans. Ent. Soc. Lond., p. 293 (1889). — Hampson, Trans. Ent. Soc. Lond., p. 163 (1897). Brasilien, Fr.Guyana, Panama
- elphegalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 112 (♂) (1924). Br. Guyana (Typus: Essequibo-Fluß)
- leucostictalis** Hampson, Ann. Mag. Nat. Hist., (s. 6) 16, p. 344 (♀) (1895); Trans. Ent. Soc. Lond., p. 163 (1897). Grenada
- phaeochoalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 458 (1906). Jamaica
- leucocymalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 459 (♂) (1906). Jamaica
- oedizonalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 459 (1906). Jamaica
- damescalis** Guenée, Delt. et Pyral., p. 272 (1854) (*Cymoriza*). — Hampson, Ent. Soc. Lond., p. 163 (1897). Brasilien
- bocusalis* Walker, Cat. Lep. Het. Brit. Mus., 19, p. 955 (♀) (1859) (*Cymoriza*).
- albitessellalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 460 (♂) (1906). Jamaica
- argentistriata** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 464 (1917). W.-Colombien
- albibasalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 461 (♂) (1906). Surinam, Brasilien
- phaeozonalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 462 (1906). Mexiko
- metalophota** Hampson, Trans. Ent. Soc. Lond., p. 166 (♀) (1897) (*A. (Opisthedeicta)*). Jamaica, Bahama Ins.
- argyrectalis** Schaus, Ann. Mag. Nat. Hist., (s. 9) 8, p. 677 (♀) (1912). Costa Rica
- heptopalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 1, p. 490 (♂) (1908). Brasilien (Itaparica)
- jonesalis** Schaus, Proc. U. S. Nat. Mus., 30, p. 137 (1906) (*Cymoriza*); Proc. Ent. Soc. Wash., 26, p. 114 (1924) (*Ambia*). Brasilien (S. Paulo)
- mareonalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 113 (♂) (1924). Peru (Typus: Yahuar-mayo)
- asaphalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 114 (♀) (1924). Peru (Typus: Tignri, Carabaya)

- arnoulalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 114 (♂) (1924). Mexiko (Typus: Jalapa)
- symphorasalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 115 (♀) (1924). Guatemala (Typus: Chejel)
- clianalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 115 (♂) (1924). Panama (Typus: Geldersland)

Species incertae sedis.

- enallassalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 294 (♀) (1914). Panama
- enantialis** Dyar, Proc. U. S. Nat. Mus., 47, p. 294 (♀) (1914). Panama
- enarexalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 294 (♀) (1914). Panama
- foveocosta** Dyar, Proc. U. S. Nat. Mus., 47, p. 293 (♂) (1914). Panama
- fulvalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 293 (♂) (1914). Panama
- fuscalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 293 (♀) (1914). Panama
- intortalis** Schaus, Ann. Mag. Nat. Hist., (s. 9) 8, p. 677 (♀) (1912). Costa Rica
- paigniodesalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 293 (♀) (1914). Panama
- phaecomeralis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 465 (♀) (1917). Bolivien

94

25. Genus **Oligostigma** Guenée

Guenée, Delt. et. Pyral., p. 260 (1854). — Snellen, Tijds. v. Ent., 19, p. 186 (1876). — Hampson, Fauna Brit. India Moths, 4, p. 207 (1896); Trans. Ent. Soc., p. 132, 167 (1897). — E. Hering, Ent. Zeit. Stettin, 64, p. 56 (1903). — Dyar, Journ. N. York Ent. c., 14, p. 88 (1906).

Hemilba Swinhoe, Ann. Mag. Nat. Hist., (s. 7) 8, p. 24 (1901). — Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 210 (1903).

(Typus generis: *juncealis* Guen.)

a) Orientalische Region.

- angulipennis** Hampson, Ill. Typ. Sp. Het. Brit. Südindien Mus. 8, p. 139, t. 155, f. 6 (1891); Trans. Ent. Soc. Lond., p. 167 (1897).
- angulipenne* Hampson, Fauna Brit. India Moths, 4, p. 209 (♀) (1896).
- alicialis** Hampson, Ann. Mag. Nat., (s. 7) 18, p. 463 (♀) (1906); Journ. Bombay Nat. Hist. Soc., 8, p. 573, t. E, f. 15 (♀) (1908). Ceylon

- niveinotatum** Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 176, t. 174, f. 32 (1893); Fauna Brit. India Moths, 4, p. 210 (1896); Trans. Ent. Soc. Lond., p. 167 (1897). Ceylon
- excisa** Swinhoe, Ann. Mag. Nat. Hist., (s. 7) 8, p. 24 (1901) (*Hemiloba*). — Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 210 (♂) (1903) (*Oligostigma*). Assam
- hapilista** Swinhoe, Trans. Ent. Soc. Lond., p. 20, t. 1, f. 11 (1892) (*Cataclysta*). — Hampson, Fauna Brit. India Moths, 4, p. 207 (1896); Trans. Ent. Soc. Lond., p. 168 (1897). — E. Hering, Stettin. Ent. Zeit., 64, p. 60 (1903). — Shibuya, Journ. Fac. Agr. Hokkaido, 22, p. 150, 151 (1928). Assam (Khasis), Indien, Formosa, Sumatra
- albidivisa** Warren, Ann. Mag. Nat. Hist., (s. 6), 18, p. 221 (1896) (*Oligostigma*) (Khasis).
- fuscifusale** Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 177, t. 174, f. 28 (1893); Fauna Brit. India Moths, 4, p. 209 (♂) (1896); Trans. Ent. Soc. Lond., p. 168 (1897). Assam, Ceylon
- syagrusalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 953 (♀) 1859 (*Cataclysta*). — *syagrusale* Hampson, Trans. Ent. Soc. Lond., p. 168 (1897). Borneo
- melanodes** Meyrick, Trans. Ent. Soc. Lond., p. 473 (1894).
- conspurcatalis** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 202 (1896) (*Ambia*). Assam (Khasis), Sangir, Holl. Neu-Guinea
- conspurcatale** Hampson, Fauna Brit. India Moths, 4, p. 209 (♂) (1896); Trans. Ent. Soc. Lond., p. 168 (1897).
- parvalis** Moore, Proc. Zool. Soc. Lond., p. 616 (1877); Trans. Ent. Soc. Lond., p. 168 (1897). — Hering, Ent. Zeit. Stettin, 64, p. 60 (1903). Andamanen, Borneo, Sumatra
- parvale** Hampson, Fauna Brit. India Moths, 4, p. 209 (♀) (1896).
- fumibasale** Hampson, Fauna Brit. India Moths, 4, p. 208 (♀) (1896); Trans. Ent. Soc. Lond., p. 168 (1897). Ceylon
- andreustalis** Hampson, Journ. Bombay Nat. Hist. Soc., 21, p. 1261, t. G, f. 43 (♀) (1912). Madras
- insectalis** Pryer, Cist. Ent., 2, p. 234, t. 4, f. 7 (1877). China (Shanghai)
- insectale** Hampson, Trans. Ent. Soc. Lond., p. 168 (1897).
- bilinealis** Snellen, Tijds. v. Ent., 19, p. 196, t. 8, f. 1, a—c (1876). Indien (Punjab, Bengal, Dharmasala, Calcutta)
- bilineale** Hampson, Fauna Brit. India Moths, 4, p. 208 (1896); Trans. Ent. Soc. Lond., p. 168 (1897).

- villidalis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 435 (♀) (1859) (*Oligostigma*?) (Sarawak). — Meyrick, Trans. Ent. Soc. Lond., p. 470 (1894) (*Nymphula*); p. 85 (1897) (*Nymphula*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 151 (1928). — Tams, Ins. Samoa, pars 3, fasc. 4, p. 261 (1935). Assam, Formosa, Ceylon, Java, Borneo, Sangir, Fidji, Samoa, Australien, Queensland
- villidale** Hampson, Fauna Brit. India Moths, 4, p. 208 (♂) (1896); Trans. Ent. Soc. Lond., p. 168 (1897).
- sacadasalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 963 (♀) (1859) (*Hydrocampa*) (Moreton-Bai).
- unilinealis** Snellen, Tijds. v. Ent., 19, p. 197, t. 8, f. 2, a, b (1876). — Meyrick, Trans. Ent. Soc. Lond., p. 221 (1886) (*Anydracula*).
- melanotalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 463 (1906); Journ. Bombay Nat. Hist. Soc., 18, p. 573 (1908). Ceylon
- discalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 464 (♂) (1906). Celebes
- orthoteles** Meyrick, Trans. Ent. Soc. Lond., p. 472 (1894). Sumbawa
- camptoteles** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 464 (1906). Tambora
- picalis** Guemée, Delt. et Pyral., p. 274 (1854) (*Hydrocampa*). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 461 (1859) (*Hydrocampa*). — Lederer, Wien. Ent. Monatschr., 7, p. 451 (1863) (*Hydrocampa*). Indien, Ceylon
- picala** Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 177, t. 174, f. 23 (1893); Fauna Brit. India Moths, 4, p. 210, f. 122 (♂, ♀) (1896); Trans. Ent. Soc. Lond., p. 167, fig. (♀) (1897).
- dianale** Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 177, t. 174, f. 27 (1893); Fauna Brit. India Moths, 4, p. 211 (♂) (1896); Trans. Ent. Soc. Lond., p. 169 (1897). Ceylon
- hemicryptis** Meyrick, Trans. Ent. Soc. Lond., p. 84 (♂, ♀) (1897). — Hampson, Trans. Ent. Soc. Lond., p. 169 (1897). Sangir
- camptozonale** Hampson, vide sub c) Australische Region p. 120.
- palleuca** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 465 (♂) (1906). Borneo
- flavipictalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 469 (♀) (1917). Singapore
- pulchellale** Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 178, t. 174, f. 30 (1893); Fauna Brit. India Moths, 4, p. 211 (♂) (1896); Trans. Ent. Soc. Lond., p. 170 (1897). Assam, Ceylon
- ornata** Moore, Lep. Ceylon, 3, p. 299, t. 181, f. 11 (1885) (*Cataclysta*). Ceylon
- ornatum** Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 51 (1893); Fauna Brit. India

Moths, 4, p. 212 (1896); Trans. Ent. Soc. Lond., p. 170 (1897).

auropunctalis Hampsen, Journ. Bombay Nat. Hist. Soc., 15, p. 210 (♀) (1903); Ann. Mag. Nat. Hist. (s. 7) 18, p. 466 (1906); Journ. Bombay Nat. Hist. Soc., 18, p. 573 (1908). Indien (Bhotan, Madras), Java

var. *javanica* Strand, Ent. Mitt., 3, p. 276 (1914). Java

chrysozonalis Hampsen, Journ. Bombay Nat. Hist. Soc., 21, p. 1261 (♀) (1912). Madras

bifurealis Pryer, Cist. Ent., 2, p. 232, t. 4, f. 14 (1877). Südechina, — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 151 (1928). Formosa, Assam,

bifurcale Hampsen, Fauna Brit. India Meths, 4, p. 212 (♂) (1896); Trans. Ent. Soc. Lond., p. 170 (1897). — Caradja, Mem. Acad. Rem. Sect. Stiint., (s. 3) 3, mem. 7, p. 331 (1925). Burma, var. *szechuanalis* Caradja, Ark. Zool., 27, A, nr. 8, p. 77 (1935). China (N.O.-Szechuan)

albifuscalis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 466 (♂) (1906); Journ. Bombay Nat. Hist. Soc., 18, p. 571 (♂) (1908). Assam (Khasis)

aracale Hampson, Trans. Ent. Soc. Lond., p. 170 (1897); Journ. Bombay Nat. Hist. Soc., 12, p. 701 (1899). Ceylon

Biologie: Hampson, Proc. Ent. Soc. Lond., p. X (1900); Ent. Men. Mag., s. 2, 11 (30), p. 214 (1900).

Species incertae sedis.

adjunctalis Snellen, Tijds. v. Ent., 38, p. 158, t. 6, f. 17, 18 (1895). — E. Hering, Stettin. Ent. Zeit., 64, p. 59 (1903). Java, Lombok, Flores, Celebes

angustalis Sauber in Semper, Schm. Philipp., 2, p. 655 (♀) (1902). Philippinen

excisalis Snellen, Tijds. v. Ent., 43, p. 289, t. 16, f. 9 (1901). — E. Hering, Stettin. Ent. Zeit., 64, p. 59 (1903). Java, Sumatra

falcatalis Snellen, Tijds. v. Ent., 43, p. 287, t. 16, f. 7, 8 (1901). — Hering, Stettin. Ent. Zeit., 64, p. 59 (1903). Java, Sumatra

kwangtungialis Caradja, Mem. Acad. Roum. Sect. Stiint., (s. 3) 3, mem. 7, p. 331, t. 1, f. 33 (1925). Südechina

minutalis Caradja, Bull. Acad. Roum., 15, p. 148 (♂, ♀) (1932). Südechina

quinquinalis Snellen, Tijds. v. Ent., 35, p. 176 (1892). Java, Celebes, Aru Ins.

sanguinisepta Swinhoe, Ann. Mag. Nat. Hist., (s. 7) 8, p. 25 (♂) 1901. Bernéo

b) Aethiopische und Madagassische Region.

- rufiterminalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) Madagaskar
19, p. 467 (♀) (1917).
irialis Walker, Cat. Lep. Het. Brit. Mus., 17, p. Westafrika
444 (♂) (1859) (*Cataclysta*) (Sierra Leone).
irialis Hampson, Trans. Ent. Soc. Lond.,
p. 168 (1897).
piperitalis Hampson, Ann. Mag. Nat. Hist., (s. 8) N.-Nigeria
19, p. 467 (♀) (1917).
blpunctalis Hampson, Ann. Mag. Nat. Hist., (s. 7) Nigeria
18, p. 463 (♀) (1906).
flavialbalis Hampson, Ann. Mag. Nat. Hist., (s. 8) Madagaskar
19, p. 468 (♀) (1917).

c) Australische Region.

- centrimacula** Hampson, Ann. Mag. Nat. Hist., (s. 8) Queensland
19, p. 465 (1917).
conspurcatalis Warren, vide sub a) Orientalische
Region p. 117.
dominulalis Walker, Cat. Lep. Het. Brit. Mus., 34, Neu-Guinea
p. 1338 (1865) (*Cataclysta*).
dominulalis Hampson, Trans. Ent. Soc.
Lond., p. 168 (1897) (*Oligostigma*).
villidalis Walker, vide sub a) Orientalische Re-
gion p. 118.
argylineale Hampson, Trans. Ent. Soc. Lond., p. Queensland
168 (1897).
polydectalis Walker, Cat. Lep. Het. Brit. Mus., 17, Australien,
p. 451 (♂) (1859) (*Cataclysta*). — Meyrick, N.S.Wales,
Trans. Ent. Soc. Lond., p. 430 (♂) (1885) Fidji Ins.
(*Paraponyx*); id., ibid., p. 221 n. 49 (1886)
(*Paraponyx*).
polydectale Hampson, Trans. Ent. Soc.
Lond., p. 169 (1897).
chrysota Meyrick, Trans. Ent. Soc. Lond., p. 221 Fidji Ins.
(♂) (1886) (*Paraponyx*).
leucomma Hampson, Ann. Mag. Nat. Hist., (s. 8) Queensland
19, p. 468 (♂, ♀) (1917).
camptozonale Hampson, Trans. Ent. Soc. Lond., Amboina,
p. 169 (♂, ♀) (1897). Fergusson
Ins.,
Queensland,
West-
Australien

ectogonalis Hampson, Ann. Mag. Nat. Hist., (s. 7) Louisiaden
18, p. 466 (♀) (1906). (St.Aignan)
tripletale Hampson, Trans. Ent. Soc. Lond., p. 170 Fergusson Ins.
(1897).

Species incertae sedis.

- argyrotoxis** Pagenstecher, Jahrb. Nassau. Ver. Na- Aru Ins.
turk., 39, p. 176 (1886).
barbararcha Meyrick, Exot. Microlep., 4, p. 240 Fidji Ins.
(♂) (1932). (Typus:
Lantoka)

- candidalis** Pagenstecher, Jahrb. Nassau. Ver. Na- Aru Ins.
turk., 39, p. 176 (♀) (1886).
orphanalis Pagenstecher, Jahrb. Nassau. Ver. Na- Aru Ins.
turk., 39, p. 177 (1886).
praestabilis Pagenstecher, Jahrb. Nassau. Ver. Na- Aru Ins.
turk., 39, p. 177 (♂, ♀) (1886).

d) Neotropische Region.

- cryptalis** Druce, Biol. Centr.-Amer. Het., 2, p. 274. Mexiko,
t. 63, f. 19 (1896) (*Cataclysta*). Costa Rica,
Panama
cryptale Hampson, Trans. Ent. Soc., p. 167
(1897).
profusalis Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, Costa Rica
p. 978 (1912).
peruviansis Hampson, Ann. Mag. Nat. Hist., (s. 8) Peru
19, p. 466 (1917).
juncealis Guenée, Delt. et Pyral., p. 261, t. 4, f. 6 Brasilien,
(1854) (Cayenne). — Schaus, Proc. Ent. Soc. Fr. Guyana;
Wash., 26, p. 117 (1924). — Forbes, Lep. N. Florida
York, p. 578 (1923) (*Nymphula*).
junceale Hampson, Trans. Ent. Soc. Lond.,
p. 167 (1897).
odrianalis Schaus, Proc. Ent. Soc. Wash., 26, p. Brasilien
116 (♂) (1924). (Typus:
S. Paulo)
flavimarginalis Warren, Trans. Ent. Soc. Lond., Brasilien
p. 294 (1889).
flavimarginale Hampson, Trans. Ent.
Soc. Lond., p. 167 (1897).
exhibitilis Walker, Trans. Ent. Soc. Lond., (s. 3) Brasilien
1, p. 123 (1862).
exhibitale Hampson, Trans. Ent. Soc.
Lond., p. 168 (1897).
phaedralis Walker, Cat. Lep. Het. Brit. Mus., 17, Brasilien
p. 443 (1859) (*Cataclysta*).
phaedrale Hampson, Trans. Ent. Soc. Lond.,
p. 168 (1897).
fulvicolor Hampson, Ann. Mag. Nat. Hist., (s. 8) Peru
19, p. 469 (♂) (1917).
grisealis Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, Jamaika
p. 465 (♂) (1906).

Species incertae sedis.

- andeotalis** Schaus, Proc. Ent. Soc. Wash., 26, p. Brasilien
117 (♀) (1924). (Typus:
Castro
Parana)
duealis Schaus, Proc. U. S. Nat. Mus., 30, p. 136 Brasilien
(1906). (S. Paulo)
electralis Dyar, Proc. U. S. Nat. Mus., 47, p. 295 Panama
(♂) (1914).
mollitalis Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, Panama
p. 678 (1912). Costa Rica,
Panama

- purifacialis** Dyar, Proc. U. S. Nat. Mus., 47, p. 296 (♀) (1914).
semimarginalis Dyar, Proc. U. S. Nat. Mus., 47, p. 295 (♂) (1914).
tigrinalis Dyar, Proc. U. S. Nat. Mus., 47, p. 295 (1914).

74

26. Genus **Ridleyana** Hampson

Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 469 (1906).

Teratauxta E. Hering (Hampson i. l.), Ent. Zeitg. Stettin, 62, p. 345 (1901).

- paradoxa** E. Hering, Ent. Zeit. Stettin., 62, p. 345 (♂) (1901) (*Teratauxta*); 64, t. 1, f. 28 (1903) (*Teratauxta*). — Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 470 (1906) (*Ridleyana*).

1

27. Genus **Parthenodes** Guenée

Guenée, Delt. et Pyral., p. 252 (1854). — Hampson, Fauna Brit. India Moths, 4, p. 216 (1896); Trans. Ent. Soc. Lond., p. 134, 181 (1897).

- Paracymoriza* Warren, Ann. Mag. Nat. Hist., (s. 6) 6, p. 479 (1890) (Type: *vagalis* Walk.).
Gethosyne Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 221 (1896) (Type: *acquivocalis* Warr.).
Stenicula Snellen, Tijds. v. Ent., 43, p. 284 (1901) (Type: *flavicaput* Snellen). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 153 (1928).

(Typus generis: *hydrocampalis* Guenée).

a) Paläarktische Region.

- prodigalis** Leech, Entom., 22, p. 70, t. 4, f. 16 (1889) (*Cataclysta*). — Hampson, Trans. Ent. Soc. Lond., p. 183 (1897). — Leech, Trans. Ent. Soc. Lond., p. 438 (1901). — Caradja, Mem. Acad. Rom. Sect. Stiint., s. 3, 4, mem. 8, p. 405 (1927). — Shibuya, Journ. Fac. Agr. Imp. Univ., 22, p. 153, t. 4, f. 32 (1928); Ins. Matsum., 3, p. 129, t. 5, f. 15 (1929). Zentral- u. Westchina, Japan, Formosa, Korea
 var. *cataclystalis* Strand, Ent. Mitt., 8, p. 105 (1919) (*Oligostigma*) (spec.). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 25, 153, 154 (1928) (*prodigalis* var.). Formosa

b) Orientalische Region.

- inextricata** Moore, Deser. Lep. Ins. Atkinson, p. 210, t. 7, f. 7 (1887). — Hampson, Fauna Brit. India Moths, 4, p. 218 (1896); Trans. Ent. Soc. Lond., p. 181 (1897). (*P. (Paracy-* Sikkim, Assam, Ostchina (Kwangtung)

- moriza*)). — Hering, Stettin. Ent. Zeit., 64, p. 55 (1903).
- exsolvalis* (part.) Snellen, Tijds. Ent., 38, p. 156 (1895) (*Hydrocampa*).
- vagalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1530 (1865) (*Oligostigma*) (Java). — Hampson, Fauna Brit. India Moths, 4, p. 219, f. 124 (♂) (1896); Trans. Ent. Soc. Lond., p. 182, fig. (♂) (1897) (*P. (Paracymoriza)*).
Indien
(Punjab,
Dharmasala,
Nilgiris),
Java,
Ostchina
(Kwangtung)
- aurantialis** Swinhoe, Ann. Mag. Nat. Hist., (s. 6) 16, p. 304 (1895). — Hampson, Fauna Brit. India Moths, 4, p. 219 (1896); Trans. Ent. Soc. Lond., p. 182 (1897) (*P. (Paracymoriza)*).
Assam
(Khasis),
Süd- und
Ostchina
- gangeticalis** Lederer, Wien. Ent. Monatschr., 7, p. 453, 485, t. 18, f. 5 (non 4) (1863) (*Parapolyx*). — Hampson, Fauna Brit. India Moths, 4, p. 219 (1896) (*P. ?*); id., Trans. Ent. Soc., p. 184 (1897).
Indien
- oxygona** Meyrick, Trans. Ent. Soc. Lond., p. 470 (1894). — Hampson, Trans. Ent. Soc. Lond., p. 182 (1897) (*P. (Paracymoriza)*).
Sumbawa
- loricatalis** Lederer, Wien. Ent. Monatschr., 7, p. 453, 486 (♀) (1863) (*Cymoriza*). — Hampson, Trans. Ent. Soc. Lond., p. 182 (1897).
Amboina
- olivalis** Hampson, Ill. Typ. Sp. Het. Brit. Mus., 8, p. 140, t. 155, f. 8 (♂) (1891); Fauna Brit. India Moths, 4, p. 216 (♂, ♀) (1897); Trans. Ent. Soc. Lond., p. 182 (1897) (*P. (P.)*).
- dentifascialis* Hampson, Ill. Typ. Sp. Brit. Mus., 8, p. 141, t. 155, f. 21 (♀) (1891) (*Paracymoriza*).
- exsolvalis** Snellen, Midden-Sumatra, 4, Nat. Hist., 2 (8), Lep., p. 76 (1880) (*Hydrocampa*); Tijds. v. Ent., 38, p. 156, t. 6, f. 13, 14 (1895). — Hampson, Fauna Brit. India Moths, 4, p. 217 (1896); Trans. Ent. Soc. Lond., p. 182 (1897) (*P. (P.)*).
Sikkim,
Assam,
Burma,
Tonkin,
Sumatra,
Java
- rivularis** Moore, Deser. Lep. Ins. Atkinson, p. 210, t. 7, f. 8 (1887) (*Cymoriza*). — E. Hering, Ent. Zeit. Stettin, 64, p. 55 (1903) (spec. dist.).
Bengalen,
Nepaul,
Himalaya,
Kulu
- exsolvalis* Snellen, Trans. Ent. Soc. Lond., p. 639 (1890); Tijds. v. Ent., 38, p. 156 (1895); 39, p. 69 (1896) (spec. dist.). — Hampson, Fauna Brit. India Moths, 4, p. 217 (1896); id., Trans. Ent. Soc. Lond., p. 182 (1897) (*P. (P.)*).
- immanis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 470 (♀) (1906).
N.-Borneo
- stellata** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 203 (1896). — Hampson, Fauna Brit. India Moths, 4, p. 217 (♂) (1896); Trans. Ent. Soc. Lond., p. 183 (1897) (*P. (P.)*).
Assam
(Khasis),
Süd- und
Zentralchina

- latifascialis** Warren, Ann. Mag. Nat. Hist., (s. 6) Assam
18, p. 220 (1896) (*Nymphula*). — Hampson,
Trans. Ent. Soc. Lond., p. 183 (1897) (*P.*
(*P.*)); id., Journ. Bombay Nat. Hist. Soc.,
12, p. 702 (1899). (Khasis)
- albifascialis** Hampson, Ill. Typ. Sp. Het. Brit. Mus., Assam,
8, p. 140, t. 156, f. 1, 9, 1893; Fauna Brit. Nilgiris
India Moths, 4, p. 217 (♂) (1896); Trans.
Ent. Soc. Lond., p. 183 (1897) (*P.* (*P.*))
- phlegetonalis** Snellen, Tijds. v. Ent., 38, p. 155, Java
t. 6, f. 9, 10 (1895) (*Hydrocampa*). — E.
Hering, Ent. Zeit. Stettin, 64, p. 55 (1903)
(spec. dist.).
- albifascialis** (part.) Hampson, Trans. Ent.
Soc. Lond., p. 183 (1897).
- semialbida** Warren, Ann. Mag. Nat. Hist.,
(s. 6) 17, p. 203 (1896) (*Paracymoriza*).
- nigriplaga** Swinhoe, Ann. Mag. Nat. Hist., (s. 6) Assam
14, p. 209 (1894) (*Nymphula*). — Hampson,
Fauna Brit. India Moths, 4, p. 218 (♂)
(1896); Trans. Ent. Soc. Lond., p. 183 (1897).
- ectopalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) N.-Borneo
18, p. 472 (♂) (1906).
- aequivocalis** Warren, Ann. Mag. Nat. Hist., (s. 6) Assam
18, p. 221 (1896) (*Gethosyne*). — Hampson,
Trans. Ent. Soc. Lond., p. 183 (1897); Journ.
Bombay Nat. Hist., 12, p. 702 (♂) (1899)
(*P.* (*P.*)). (Khasis)
- prodigalis** Leech, vide sub a) Paläarktische Re-
gion p. 122.
- distinctalis** Leech, Entom., 22, p. 67, t. 4, f. 5 West- u.
(1889) (*Botys*). — Hampson, Trans. Ent. Zentral-
Soc. Lond., p. 183 (1897) (*P.* (*P.*)). China
- var. *taiwanalis* Wileman et R. South, Formosa
Entom., 50, p. 176 (1917) (*Parthenodes tai-
wanalis*). — Shibuya, Journ. Fac. Agr. Hok-
kaido Univ. Imp., 22, p. 25, 153, 154 (1928).

Species incertae sedis.

- flavicaput** Snellen, Tijds. v. Ent., 43, p. 285, t. 16, Java
f. 2, 3 (1900) (*Stenricula*).
- parallelalis** Sauber in Semper, Schm. Philipp, 2, p. Philippinen
657, t. 66, f. 7 (♂, ♀) (1902) (*Paracymo- (Luzon)*
riza).
- pallidalis** R. South, Trans. Ent. Soc. Lond., p. 439, Zentral-
t. 14, f. 29 (1901). — Caradja, Bull. Acad. China
Roum., 15, p. 148 (1932).
- triangularis** R. South, Trans. Ent. Soc. Lond., p. West- u.
438, t. 14, f. 26 (1901). Ostchina
(Kwangtung)

c) Aethiopische und Madagassische Region.

- seotalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, Sambesi
p. 470 (♀) (1906).
- africalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) Nigeria,
18, p. 470 (♂) (1906). Nyassaland

- albiceps** Hampson, Ann. Mag. Nat. Hist., (s. 8) Transvaal
20, p. 211 (♀) (1917).
- angularis** Hampson, Trans. Ent. Soc. Lond., p. 184 Madagaskar,
(♂) (1897). D.O.Afrika
(Tanganyika-
Terr.)
(Typus:
Madagascar)
- upupalis** Guenée in Maillard, Notes sur l'île de la Réunion
Réunion, p. 63 (1863) (*Cymoriza*). — Hamp-
son, Trans. Ent. Soc. London, p. 184 (1893).

d) Australische Region.

- eetargyralis** Hampson, Trans. Ent. Soc. Lond., p. Fergusson
182 (♀) (1897) (*P. (Paracymoriza)*). Ins.
- eugethes** Tams, Ins. Samoa, pars 3 (fasc. 4), p. 261, Samoa
t. 15, f. 7, (♀) (1935).
- rectangulalis** Kenrick, Proc. Zool. Soc. Lond., p. 77, Brit.
t. 4, f. 58 (1907). Neu-Guinea

e) Neotropische Region.

- melanicalis** Hampson, Ann. Mag. Nat. Hist., s. 8, Kolumbien
20, p. 209 (♀) (1917).
- mesoleucalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) Peru
20, p. 209 (♀) (1917).
- paralleloidalis** nom. nov. Peru
parallelalis Hampson, (nec Sauber, 1902),
Ann. Mag. Nat. Hist., (s. 8) 20, p. 210 (♀)
(1917).
- rufalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, Kolumbien
p. 210 (♀) (1917).
- mediocinctalis** Hampson, Trans. Ent. Soc. Lond., p. Rio de
183 (♀) (1897) (*P. (P.)*). Janeiro
- dabinsalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 121 Mexiko
(♀) (1924).
- gualbertalis** Schaus, Proc. Ent. Soc. Wash., 26, p. Surinam
121 (♀) (1924).
- caepiosalis** Walker, Cat. Lep. Het. Brit. Mus., 19, Brasilien
p. 949 (♀) (1859) (*Hydrophysa*) (Rio Ja-
neiro). — Hampson, Trans. Ent. Soc. Lond.,
p. 183 (1897) (*P. (P.)*).
- chaleialis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, Brasilien
p. 471 (♀) (1906). (S. Paulo)
- nymphulalis** Hampson, Ann. Mag. Nat. Hist., (s. Mexiko,
7) 18, p. 472 (1906). Panama
- hydroecampalis** Guenée, Delt. et Pyral, p. 253, t. 9, Fr.Guyana,
f. 5 (1854) (Cayenne). — Hampson, Trans. Brasilien
Ent. Soc. Lond., p. 183 (1897) (*P. (P.)*).

Species incertae sedis.

- berthalis** Schaus, Proc. U. S. Nat. Mus., 30, p. 140 Brasilien
(1906). (Paraná)

- briocusalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 120 (♀) (1924).
incultalis Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, p. 679 (♂) (1912).
 46

28. Genus **Daulia** Walker

Walker, Cat. Lep. Het. Brit. Mus., 19, p. 975 (♀) (1859). — Hampson, Fauna Brit. India Moths, 4, p. 220 (1896); Trans. Ent. Soc. Lond., p. 134, 184 (1897).

Girtexta Swinhoe, Trans. Ent. Soc. Lond., p. 285 (1890).

(Typus generis: *afralis* Walker).

- afralis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 975 (♀) (1859) (Sarawak). — Hampson, Fauna Brit. India Moths, 4, p. 220, f. 125 (♂) (1896); Trans. Ent. Soc. Lond., p. 185; p. 184, fig. (♂) (1897). — Meyrick, Trans. Ent. Soc. Lond., p. 89 (1897). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 154 (1928); Ins. Matsum., 3, p. 129, t. 5, f. 16 (1929).
argentuosalis Swinhoe, Trans. Ent. Soc. Lond., p. 286 (1890) (*Girtexta*) (Rangoon). — Hampson, Ill. Typ. Sp. Het. Brit. Mus., 8, p. 41 (1891).
argyrostotalis Hampson, Journ. Bombay Nat. Hist. Soc., 21, p. 1262 (♂) (1912).
aurantialis Hampson, Fauna Brit. India Moths, 4, p. 221 (♂) (1896); Trans. Ent. Soc. Lond., p. 185 (1897).
subaurealis Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1343 (♀) (1865) (*Lepyrodes*) (Kapland). — Hampson, Trans. Ent. Soc. Lond., p. 185 (1897).
argyrophoralis Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 2 (1907).
 5

29. Genus **Margarochroma** Warren

Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 164 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 134, 185 (1897); Journ. Bombay Nat. Hist. Soc., 12, p. 703 (1899).

(Typus generis: *pictalis* Warren).

- fuscalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 2 (♂) (1907).
pictalis Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 165 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 185, fig. (♂) (1897); Journ. Bombay Nat. Hist. Soc., 12, p. 703 (♂) (1899).
 2

30. Genus **Brevicella** Kenrick

Kenrick, Proc. Zool. Soc. Lond., p. 549 (1912).

emarginata Kenrick, Proc. Zool. Soc. Lond., p. 549, Holl.
t. 68, f. 8 (1912). Neu-Guinea
1

31. Genus **Auxolophosis** Meyrick

Meyrick, Exot. Microlep., 4, p. 397 (♀) (1933).

ioxanthias Meyrick, Exot. Microlep., 4, p. 397 (♀) Fidji Ins.
(1933). (Typus:
1 Wunidawa)

32. Genus **Massepha** Walker

Walker, Cat. Lep. Het. Brit. Mus., 17, p. 488 (1859). — Hampson, Fauna Brit. India Moths, 4, p. 253 (1896); Proc. Zool. Lond. Soc., p. 594, 615 (1898) (ad *Pyraustinas*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 25 (1928) (ad *Hydrocampinas*).

Blepharomastix (part.) Druce, Biol. Centr.-Americ., Het. 2, p. 270 (1896) (*B. romalis* Druce).

Physematia Snellen, Tijds. v. Ent., 23, p. 240 (1880) (*Ph. pollutalis* Snell.).

Ceratoclasia Druce, Biol. Centr.-Amer., Het. 2, p. 554 (1899).

(Typus generis: *absolutalis* Walker).

a) **Orientalische Region.**

absolutalis Walker, Cat. Lep. Het. Brit. Mus., 17, Indien,
p. 489 (1859) (Ceylon). — Hampson, Fauna Tonkin,
Brit. India Moths, 4, p. 253, f. 151 (♂) Ostchina
(1896); Proc. Zool. Soc. Lond., p. 615, f. (Kwangtung)
14 (♂) (1898). — Shibuya, Journ. Fac. Formosa,
Agr. Hokkaido Imp. Univ., 22, p. 156 (1928). Ceylon,
— Meyrick, Exot. Microlep., 4, p. 394 Java,
(1933). Celebes

pollutalis Snellen, Tijds. v. Ent., 23, p. 240 (♂) [Typus: Maros (Celebes); Ambarawa (Java)] (1880) (*Physematia*); 27, p. 47, t. 4, f. 10 (♂) (caput) (1884) (*Physo-*
somatia); id.; ibid., 28, p. 48 (1885) (*Physo-*
matia).

marginalis Swinhoe, Ann. Mag. Nat. Hist., (s. 7) 17, p. 287 (1906).

nowa: Fletcher, Scient. Rep. Agric. Res. In. Pusa, 1916/17 (Calcutta), p. 71—90 (1917).

bengalensis Moore, Deser. Lep. Ins. Atkinson, p. N. Ost-
211 (1887). — Hampson, Fauna Brit. India Indien
Moths, 4, p. 254 (1896). (Calcutta),
(Khasis)
Assam

- ambialis** Hampson, Journ. Bombay Nat. Hist. Soc. N.-Indien, 15, p. 214 (1903). Assam, Südchina
- talboti** Caradja, Mem. Acad. Rom. Sect. Stiint., (s. 3) Westchina 4, p. 407 (1927).
- carbonalis** Warren, Ann. Mag. Nat. Hist., (s. 6) Assam 17, p. 148 (1896) (*Stenia*). — Hampson, (Khasis) Fauna Brit. India Moths, 4, p. 254 (♂) (1896).
- rufescens** Hampson, Journ. Bombay Nat. Hist. Soc., Madras 21, p. 1263, t. G, f. 44 (♀) (1912).
- rectangularis** Caradja, Deutsche Ent. Zeit. Iris, 47, p. China 157 (1934).

b) Aethiopische Region.

- entephriadia** Hampson, Proc. Zool. Soc. Lond., p. Warri, 616 (♀) (1898). Niger
- griscalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) Goldküste 20, p. 373 (♂) (1912).
- longipennis** Hampson, Ann. Mag. Nat. Hist., (s. 8) Kamerun 9, p. 167 (♂) (1912).
- tessmanni** Gaede, Mitt. Zool. Mus. Berlin, 8, p. Kamerun, 391 (♂) (1917). Span. Guinea
- flavimaculata** Gaede, Mitt. Zool. Mus. Berlin, 8, Span. p. 391 (♂) (1917). Guinea

c) Australische Region.

- phoenicobapta** Hampson, Proc. Zool. Soc. Lond., Fergusson p. 615, t. 49, f. 3 (♂) (1898). Ins., Neu-Guinea

d) Neotropische Region.

- gracilis** Hampson, Proc. Zool. Soc. Lond., p. 616 Espiritu (♂) (1898). Santo
- grammalis** Guenée, Delt et Pyral., p. 249 (1854) Fr. u. Br. (*Stenia*) (Cayenne). — Hampson, Proc. Zool. Guyana, Soc. Lond., p. 616 (1898). Panama, Kolumbien, Brasilien
- asiusalis** Walker, Cat. Lep. Het. Brit. (Typus: Mus., 18, p. 626 (♂) (1859) (*Botys*) (Rio Pará) Janeiro).
- romalis** Druce, Biol. Centr.-Amer. Het., 2, (Typus: p. 270, t. 63, f. 4 (♂) (1896) (*Blepharomastix*). Panama: Chiriqui)
- syngamoides** Hampson, Ann. Mag. Nat. Hist., (s. 8) 9, p. 106 (♂) (1912). Argentinien
- cuprescens** Hampson, Ann. Mag. Nat. Hist., (s. 8) 9, p. 166 (♂) (1912). Brasilien (S. Paulo)

- fulvalis** Hampson, Proc. Zool. Soc. Lond., p. 616, f. 49, f. 2 (♂) (1898). Brasilien (Castra Parana)
- lupa** Druce, Biol. Centr.-Amer. Het., 2, p. 554, t. C, f. 20 (1899) (*Ceratoclasia*). — Dyar Proc. U. S. Nat. Mus., 47, p. 257 (1914) (*Massepha*). Guatemala, Panama

20

33. Genus **Pelena** Moore

Moore, Lep. Ceylon, 3, p. 386 (1886).

- Luma* Hampson, Fauna Brit. India Moths, 4, p. 229 (1896), part; Trans. Ent. Soc. Lond., p. 186 (1897).
Loxocoryx Meyrick, Trans. Ent. Soc. Lond., p. 6 (1894) (Type: *sericea* Butl.).
Pelina Hampson, Trans. Ent. Soc. London, p. 186 (1897) (ex synon.).
 (Typus generis: *unicolor* Moore).
- unicolor** Moore, Lep. Ceylon, 3, p. 386, t. 184, f. 16 (1886) (*Pelena*). — Hampson, Fauna Brit. India Moths, 4, p. 229 (♀) (1896) (*Luma*); Trans. Ent. Soc. Lond., p. 187 (1897) (*Luma*).
- uniformis* Hampson, Ill. Typ. Sp. Het., 9, p. 171, t. 173, f. 7 (1893) (*Lygropia*).
- sericea** Butler, Ann. Mag. Nat. Hist., (s. 5) 4, p. 451 (1869) (*Deana*). — Pryer, Trans. Asiat. Soc. Japan, 13, p. 59 (1885) (*Deana*). — Meyrick, Trans. Ent. Soc. Lond., p. 6 (1894) (*Loxocoryx*). — Hampson, Fauna Brit. India Moths, 4, p. 229, f. 134 (♂) (1896) (*Luma*); Trans. Ent. Soc. Lond., p. 186, fig. (♂), (1897) (*Luma*). — Leech, Trans. Ent. Soc. Lond., p. 441 (1901) (*Luma*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 157 (1928) (*Pelena*); Ins. Matsum., 3, p. 129, t. 5, f. 17 (1929). Süd- u. Zentral-China, Tonkin, Assam (Khasis), Burma, Japan, Formosa
- obscuralis** Swinhoe, Ann. Mag. Nat. Hist., (s. 6) 16, p. 300 (1895) (*Loxocoryx*). — Hampson, Fauna Brit. India Moths, 4, p. 230 (1896) (*Luma*); Trans. Ent. Soc. Lond., p. 187 (1897) (*Luma*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 157 (1928) (*Pelena*). Indien, Assam, Formosa
- photias** Meyrick, Trans. Ent. Soc. Lond., p. 458 (1994) (*Tylosteges*). — Hampson, Proc. Zool. Soc. Lond., p. 496 (1898) (*Nacoleia*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 26, 158 (1928) (*Pelena*). Indien, Pulo Laut, Borneo, Formosa
- bimaculalis** Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 157, t. 6, f. 28 (♂) (1928). Formosa

5

34. Genus **Pelinopsis** Dognin

Dognin, Ann. Soc. Ent. Belg., 49, p. 67 (1905).

pachyzancloides Dognin, Ann. Soc. Ent. Belg., 49, Ecuador
p. 67 (♂) (1905).

1

35. Genus **Luma** Walker

Walker, Cat. Lep. Het. Brit. Mus., 27, p. 121 (1863). — Hampson, Fauna Brit. India Moths, 4, p. 229 (1896); Trans. Ent. Soc. Lond., p. 132, 186 (1897). (part.).

Zebronia Leech, Entomologist, 22, p. 71 (1899) (*Z. ornatalis* Leech).

(Typus generis: *anticalis* Walker).

- anticalis** Walker, Cat. Lep. Het. Brit. Mus., 27, p. 121 (1863). — Hampson, Trans. Ent. Soc. Lond., p. 186 (1897). Borneo
- trilinealis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1351 (1865) (*Zebronia*). — Hampson, Trans. Ent. Soc. Lond., p. 186 (1897) (*Luma*). — Swinhoe, Cat. East. Austral. Lep. Het. Mus. Oxford, 2, p. 448, t. 7, f. 10 (1900) (*Pelina*). Ceram
- macropsalis** Hampson, Trans. Ent. Soc. Lond., p. 186 (1897). Fergusson Ins.
- flavalis** Hampson, Ill. Typ. Sp. Het. Brit. Mus., 9, p. 171, t. 173, f. 14 (1893); Fauna Brit. India Moths, 4, p. 229 (♀) (1896); Trans. Ent. Soc. Lond., p. 187 (1897). Ceylon
- flavimarginalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 10, p. 2 (1907); Journ. Bombay Nat. Hist. Soc., 18, p. 575 (1908). Ceylon
- holoxantha** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 2 (♀) (1907). Mashonaland
- ornatalis** Leech, Entom., 22, p. 71, t. 4, f. 12 (1899) (*Zebronia*). — Hampson, Journ. Bombay Nat. Hist. Soc., 12, p. 703 (1899); Trans. Ent. Soc. Lond., p. 187 (1897). West-, Zentral- u. Südechina, Assam
- longidentalis** Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 211 (1903). Bhotan, Key Ins.
- albifascialis** Hampson, Trans. Ent. Soc. Lond., p. 187 (♂) (1897). Jamaica
- trimaculata** Hampson, Trans. Ent. Soc. Lond., p. 187 (♂) (1897). Pula Laut
- tyranthes** Meyrick, Trans. Ent. Soc. Lond., p. 89 (1897). — Hampson, Trans. Ent. Soc. Lond., p. 187 (1897). Sangir

36. Genus **Orphnophanes** Lederer

Lederer, Wien. Ent. Monatschr., 7, p. 428 (1863). — Hampson, Fauna Brit. India Moths, 4, p. 230 (1896); Trans. Ent. Soc. Lond., p. 132, 187 (1897). — Pagenstecher, Zoologica H. 29, p. 17 (1900).

Syntomodora Meyrick, Trans. Ent. Soc. Lond., p. 467 (1894) (Typus: *S. thoasalis* Walk.). — E. Hering, Ent. Zeit. Stettin, 64, p. 39 (1903).

(Typus generis: *eucerusalis* Walker).

- eucerusalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 1002 (♂) (1859) (*Botys*) (Sarawak). Borneo, Sikkim,
eucerasalis Hampson, Fauna Brit. India Moths, 4, p. 231, f. 135 (♂) (1896); Trans. Ent. Soc. Lond., p. 188, fig. (♂) (1897). — Amboina, Sumatra, Woodlarch
 E. Hering, Ent. Zeit. Stettin, 64, p. 38 (1903). Ins., St. Aignan, Fergusson
productalis Lederer, Wien. Ent. Monatschr., 7, p. 428, t. 16, f. 7 (1863). — Ins.,
 Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1487 (1865). — Snellen, Trans. Ent. Soc. Lond., p. 625 (1890). Brit.
euceralis Meyrick, Trans. Ent. Soc. Lond., p. 469 (1894) (nom. emend.). Neu-Guinea
 S.O.-Borneo
- laevis** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 206 (1896) (*Parapoynx*?). — Assam
 Hampson, Fauna Brit. India Moths, 4, p. 230 (♂) (1896); Trans. Ent. Soc. Lond., p. 188 (1897). (Khasis)
- thoasalis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 692 (1859) (*Botys*) (Sarawak). — Sikkim,
 Meyrick, Trans. Ent. Soc. Lond., p. 467 (1894) (*Syntomodora*). — Assam,
 Hampson, Fauna Brit. India Moths, 4, p. 230 (♂) (1896); Trans. Andamanen,
 Ent. Soc. Lond., p. 188 (1897). — E. Hering, Borneo,
 Ent. Zeit. Stettin, 64, p. 39 (1903) (*Syntomodora*). Java, Buru,
 Nias, Holl.
 Neu-Guinea
- albissignalis** Hampson, Fauna Brit. India Moths, 4, p. 231 (♂) (1896); Trans. Ent. Soc. Lond., p. 188 (1897). Sikkim,
 Assam,
 Java,
 Philippinen
- inconsequens** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 204 (1896) (*Syntomodora*). — N.O.-Indien,
 Hampson, Fauna Brit. India Moths, 4, p. 231 (♂) (1896); Trans. Ent. Soc. Lond., p. 188 (1897). — Assam,
 Swinhoe, Ann. Mag. Nat. Hist., (s. 7) 8, p. 136 (1901) (*Tabidia*). — Pulo Laut,
 E. Hering, Ent. Zeit. Stettin, 64, p. 40 (1903) (*Syntomodora*). Sumatra,
 Queensland
- strigatalis** Pagenstecher, Zoologica H. 29, p. 178, t. 1, f. 16 (1900). Bismarck-
 Archipel
 (Neu-Pommern),
 Shortlands
 Ins.

37. Genus **Niphostola** Hampson

Hampson, Fauna Brit. India Moths, 4, p. 222 (1896); Trans. Ent. Soc. Lond., p. 133, 189 (1897).

(Typus generis: *micans* Hamps.).

- micans** Hampson, Fauna Brit. India Moths, 4, p. 222, f. 127 (♂) (1896); Trans. Ent. Soc. Lond., p. 189, fig. (♂) (1897). Assam
- punctata** Swinhoe, Trans. Ent. Soc. Lond., p. 156 (♂) (1904). Borneo
(Typus: Santabung)

38. Genus **Perisyntrocha** Meyrick

Meyrick, Trans. Ent. Soc. Lond., p. 469 (1894). — Hampson, Trans. Ent. Soc. Lond., p. 133, 189 (1897). — Pagenstecher, Zoologica H. 29, p. 175 (1900).

(Typus generis: *P. alienialis* Walk. = *anialis* Walk.).

- anialis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 710, n. 287 (♂) (1859) (*Botys*) (Sarawak). Celebes,
— Hampson, Trans. Ent. Soc., Lond., p. 190 Sula, Buru,
(1897) (*Perisyntrocha*). — Pagenstecher, Neu-
Zoologica H. 29, p. 175 (1900). Britannia,
Bismarek-
Archipel,
Neu-Guinea,
Australien,
Melville Ins.
- alienalis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1351 (♀) (1865) (*Zebronia*). — (Makian,
Meyrick, Trans. Ent. Soc. Lond., p. 469 Celebes,
(1894) (*Perisyntrocha*). N.Guinea)
- circumdatalis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1435 (♀) (1865) (*Botys*) (Sula).
- cuneolalis* Snellen, Tijds. v. Ent., 23, p. 243 (♂) (1880); 27, p. 48, t. 5, f. 2 (1884) (Typus: Makassar)
(*Paraponyx*).
- picata* Butler, Ann. Mag. Nat. Hist., (s. 5) 10, p. 235 (1882) (*Stegothyris*) (Neu-Britannia).
- ossealis** Hampson, Fauna Brit. India Moths, 4, p. 222, f. 128 (♂) (1896); Trans. Ent. Soc. Lond., p. 190; p. 189 fig. (♂) (1897). N.O.-
Himalaya,
Assam,
Sikkim
- flavalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 211 (1917). Kamerun
- suffusa** Kenrick, Proc. Zool. Soc. Lond., p. 548, t. 68, f. 4 (1912). Holl.
Neu-Guinea

39. Genus **Eranistis** Meyrick

Meyrick, Trans. Proc. N. Zeal. Inst., 42, p. 70 (1909).

pandora Meyrick, Trans. Proc. N. Zeal. Inst., 42, p. 70 (♀) (1909). Kermadec Ins.

1

40. Genus **Strepsinoma** Meyrick

Meyrick, Trans. Ent. Soc. Lond., p. 85 (1897). — Hampson, Trans. Ent. Soc. Lond., p. 133, 190 (1897).

(Typus generis: *amaura* Meyrick.)

a) **Orientalische Region.**

amaura Meyrick, Trans. Ent. Soc. Lond., p. 85 (♂, ♀) (1897). — Hampson, Trans. Ent. Soc. Lond., p. 190, fig. (♂) (1897). Sumbawa, S.O.-Borneo

sphenactis Meyrick, Trans. Ent. Soc. Lond., p. 86 (♂, ♀) (1897). — Hampson, Trans. Ent. Soc. Lond., p. 190 (1897). Sangir

hapilistalis Strand, Ent. Mitt., 8, p. 105 (♀) (1919) Formosa
(*Oligostigma*?). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 24, 152 (1928) (*Strepsinoma*).

b) **Australische Region.**

albimaculalis W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 124 (♀) (1915). Holl. Neu-Guinea

albiplagialis W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 124 (♂) (1915). Holl. Neu-Guinea

aulacodoidalis W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 123 (♀) (1915). Holl. Neu-Guinea

ectopalis Hampson, Trans. Ent. Soc. Lond., p. 190 (1897). Fergusson Ins.

fuliginosa W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 124 (♂) (1915). Neu-Guinea

grisealis W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 124 (♂) (1915). Holl. Neu-Guinea

9

41. Genus **Duponchelia** Zeller

Zeller, Isis, p. 588 (1847). — Hampson, Trans. Soc. Ent. Lond., p. 133, 191 (1897).

(Typus generis: *fovealis* Zell.)

fovealis Zeller, Isis, p. 588 (1847). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 8, t. 1, f. 3 (Pyr.) (1848). — Guenée, Delt. et Pyral., p. 239 (1854). — Hampson, Trans. Ent. Soc. Lond., p. 191, fig. (♂) (1897). — Süd- und West-europa, (Süd-Frankreich,

- Rebel in Spuler, Schm. Eur., 2, p. 222 (1906). Italien,
Dalmatien,
canuialis Millière, Icon. Chen. Lep., 2, Balkan,
p. 426, t. 95, f. 5—7 (1868) (*Stenia*). Korsika,
griseata Butler, Ann. Mag. Nat. Hist., (s. Kreta,
4) 16, p. 415 (1875) (*Hymenia*). Cypem,
Spanien),
Canaren,
W.-Mau-
retamen
(Gr. Atlas),
Algier,
Ägypten,
Kleinasien,
Syrien,
Südarabien
(Aden),
Port.
O.-Afrika,
Congo,
Natal
- caldalis** Oberthür, Etud. Ent., 12, p. 36, t. 9, f. 39 Mauretanien,
(1888). Algier
- fumidalis** Turati, Atti Soc. Ital. Sci. Nat., 73, p. Cyrenaica
189, fig. (1934).
3

42. Genus **Stegothyris** Lederer

Lederer, Wien. Ent. Monatschr., 6, p. 427 (1863). — Hampson,
Fauna Brit. India Moths, 4, p. 223 (1896); Trans. Ent. Soc. Lond.,
p. 133, 192 (1897).

(Typus generis: *fasciculalis* Zell.)

- fasciculalis** Zeller, Lep. Caffr., p. 63 (1852). — Südafrika,
Hampson, Trans. Ent. Soc. Lond., p. 192 Madagaskar
(1897).
- diagonalis** Guenée, Delt. et Pyral., p. 201 (1854) Indien,
(*Salbia*) (Java?). — Snellen in Veth, Mid- Himalaya,
den-Sumatra, 4, p. 74 (1880); Trans. Ent. Burma,
Soc. Lond., p. 625 (1890). — Hampson, Südchina,
Fauna Brit. India Moths, 4, p. 223, f. 129 Formosa,
(♂) (1896); Trans. Ent. Soc. Lond., p. Sumatra,
192, fig. (♂) (1897). — E. Hering, Ent. Java,
Zeit. Stettin, 64, p. 40 (1903). — Shibuya, Lombok,
Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. Sumbawa
158, t. 6, f. 5 (1928).
- transversalis** Lederer, Wien. Ent. Mo-
natschr., 7, p. 427, 482, t. 16, f. 5 (1863)
(Venezuela?).
- plagalalis** Moore, Proc. Zool. Soc. Lond., p.
96 (1867) (*Botys*).
- atralis** Pagenstecher in Voeltzkow, Reise Ostafrika, Komoren
2, p. 131 (1907).
3

43. Genus **Dracaenura** Meyrick

Meyrick, Trans. Ent. Soc. Lond., p. 227 (1886). — Hampson, Trans. Ent. Soc. Lond., p. 133, 192 (1897). — Pagenstecher, Zoologica H. 29, p. 178 (1900).

(Typus generis: *stenosoma* Feld. et Rogenh.)

- stenosoma** Felder [et Rogenhofer], Reise Novara Fidji Ins.
(Het.) 2 (2), p. 16, t. 134, f. 16 (1874)
(*Platamonia*?). — Meyrick, Trans. Ent. Soc.
Lond., p. 227 (1886). — Hampson, Trans.
Ent. Soc. Lond., p. 192 (fig. (♂) 193 (1897).
- versicolor** Pagenstecher, Zoologica H. 29, p. 177, t. 2, Bismarck-
f. 36 (♂) (1900). Archipel
(Neu-
Pommern)
- prosthenialis** Hampson, Trans. Ent. Soc. Lond., p. Neu-Guinea
193 (♂) (1897). (Humboldt-
Bai)
- metaleuca** Hampson, Ann. Mag. Nat. Hist., (s. 8) Brit.
20, p. 211 (♂) (1917). Neu-Guinea
- leucoprocta** Hampson, Trans. Ent. Soc. Lond., p. Neu-Hebriden
193 (♂) (1897).
- myota** Meyrick, Trans. Ent. Soc. Lond., p. 230 Neu-
(♂) (1886). — Hampson, Trans. Ent. Soc. Hebriden,
Lond., p. 194 (1897). Fidji Ins.
- agramma** Meyrick, Trans. Ent. Soc. Lond., p. 229 Samoa Ins.
(♂, ♀) (1886). — Hampson, Trans. Ent.
Soc. Lond., p. 194 (1897). — Rebel, Denk-
schr. Akad. Wiss. Wien, 85, p. 430 (1910; 2.
Beih. Jahrb. Hamb. Wiss. Anstalt, 32, p.
146 (1915). — Tams, Ins. Samoa, pars 3,
fasc. 4, p. 262 (1935).
- ab. *dolia* Tams, Ins. Samoa, pars 3, fasc. 4, p. Samoa Ins.
262 (1935).
- adela** Tams, Ins. Samoa, pars 3, fasc. 4, p. 263, t. Samoa Ins.
15, f. 3 (♂) (1935).
- horochroa** Meyrick, Trans. Ent. Soc. Lond., p. 229 Neu-Guinea,
(♂) (1886). — Hampson, Trans. Ent. Soc. Buru
Lond., p. 194 (1897). (Typus:
Port
Moresby)
- chrysochroa** Hampson, Ann. Mag. Nat. Hist., (s. 7) Holl.
19, p. 3 (♂) (1907). Neu-Guineaa
- pelochra** Meyrick, Trans. Ent. Soc. Lond., p. 228 Fidji Ins.
(♂, ♀) (1886). — Hampson, Trans. Ent.
Soc. Lond., p. 194 (1897).
- argyrogaster** Butler, Trans. Ent. Soc. Viti
Lond., p. 431 (1886) (*Botys*).

- aegialitis** Meyrick, Trans. Proc. N. Zeal. Inst., 42, Kermadec
p. 71 (♂) (1910). Ins.
- asthenota** Meyrick, Trans. Ent. Soc. Lond., p. 229 Fidji Ins.
(♀) (1886). — Hampson, Trans. Ent. Soc.
Lond., p. 194 (1897).
- cineticorpus** Hampson, Trans. Ent. Soc. Lond., p. Loyalty Ins.
194 (♂) (1897).
- tagiadialis** Hampson, Trans. Ent. Soc. Lond., p. 194 Fergusson
(1897). Ins.
- albonigralis** Hampson, Trans. Ent. Soc. Lond., p. Brit.
194 (♂) (1897). Neu-Guinea
- semialbalis** W. Rothschild, Rep. B. O. U. Exp., 2, Holl.
nr. 15, p. 123, t. 2, f. 39 (♀) (1915). Neu-Guinea

Species incertae sedis.

- arfakalis** Swinhoe, Ann. Mag. Nat. Hist., (s. 8) 20, Holl.
p. 158 (♂, ♀) (1917); s. 8, 2, p. 94 (1918). Neu-Guinea
- griseotinctalis** W. Rothschild, Rep. B. O. U. Exp., Holl.
2, nr. 15, p. 123 (♀) (1915). Neu-Guinea
- plebejifascialis** W. Rothschild, Rep. B. O. U. Exp., 2, Holl.
nr. 15, p. 122 (♂) (1915). Neu-Guinea
- pseudopelochra** W. Rothschild, Rep. B. O. U. Exp. Holl.
2, nr. 15, p. 123 (♂) (1915). Neu-Guinea
- torridalis** Kenrick, Proc. Zool. Soc. Lond., p. 77, Brit.
t. 3, f. 67 (1907). Neu-Guinea
- unicoloralis** Kenrick, Proc. Zool. Soc. Lond., p. Brit.
77, t. 4, f. 68 (1907). Neu-Guinea
23

44. Genus **Osphranfis** Meyrick

Meyrick, Trans. Ent. Soc. Lond., p. 87 (1897). — Hampson, Trans.
Ent. Soc. Lond., p. 240 (1897).

- paraphaea** Meyrick, Trans. Ent. Soc. Lond., p. 88 Talaut
(♂) (1897).

1

45. Genus **Pythagorea** Meyrick

Meyrick, Trans. Ent. Soc. Lond., 77, p. 162 (1929).

- categorica** Meyrick, Trans. Ent. Soc. Lond., 77, Gesellsch.-
p. 162 (♂, ♀) (1929). Ins.
1 (Typus:
Tahiti)

46. Genus **Hymenoptychis** Zeller

Zeller, Vet. Akad. Handl. (Lep. Caffr.), p. 63 (1852). — Snellen, Tijds. v. Ent., 27, p. 40 (1884). — Meyrick, Trans. Ent. Soc. Lond., p. 225 (1886). — Hampson, Fauna Brit. India Moths, 4, p. 223 (1896); Trans. Ent. Soc. Lond., p. 133, 195 (1897). — Pagenstecher, Zoologica H. 29, p. 175 (1900).

Syrbatis Walker, Cat. Lep. Het. Brit. Mus., 27, p. 119 (1863).

(Typus generis: *sordida* Zeller.)

- sordida** Zeller, K. Vet. Akad. Handl. Lep. Caffr., Indien
p. 65 (1852) (Natal). — Lederer, Wien. (Penang,
Ent. Monatschr., 7, p. 426, t. 16, f. 2 Bengalen),
(1863). — Walker, Cat. Lep. Het. Brit. Birma,
Mus., 34, p. 1329 (1865). — Snellen, Tijds. Malakka,
v. Ent., 27, p. 40 (1884). — Meyrick, Celebes,
Trans. Ent. Soc. Lond., p. 226 (1886); p. Ceylon,
514 (1889). — Hampson, Ill. Typ. Sp. Het. Borneo,
Brit. Mus., 8, p. 40 (1891); 9, p. 45 (1893); Neu-Guinea,
Fauna Brit. India Moths, 4, p. 224, f. Fidji,
130 (♂) (1896); Trans. Ent. Soc. Lond., Philippinen,
p. 195, fig. (♂) (1897). — Pagenstecher, Bismarck-
Zoologica H. 29, p. 175 (1900). Archipel,
tipuliformis Walker, Cat. Lep. Het. Buru,
Brit. Mus., 27, p. 120 (1863) (*Syrbatis*) Shortland,
(Sarawak). Marshall,
pterophoralis Walker, Cat. Lep. Het. Ellice
Brit. Mus., 34, p. 1413 (♀) (1865) (*Botys*) Ins.,
(Penang). — Swinhoe, Trans. Ent. Soc. Australien,
Lond., p. 272 (1890) (*Tatobotys*). Tasmanien,
Queensland,
D. O.- und
Port. O.-
Afrika,
Südafrika,
Madagaskar
- dentilinealis** Snellen, in Veth, Midden-Sumatra, 4, Java,
(1) 8, Lep., p. 74 (1880); Tijds. v. Ent., Sumatra,
27, p. 40 (1884). — Hampson, Trans. Ent. Celebes,
Soc. Lond., p. 195 (1897). — Pagenstecher, Philippinen,
Zoologica H. 29, p. 176 (1900). — Hering, Bismarck-
Ent. Zeit. Stettin, 64, p. 44 (1903). Archipel
(Neu-
Pommern)
- phryganidalis** Pagenstecher, Jahrb. Nassau. Ver. Na- Aru Ins.
turk., 38, p. 174 (1885).
- scalpellalis** Pagenstecher, Jahrb. Nassau. Ver. Na- Aru Ins.
turk., 38, p. 173 (1885).

4

47. Genus **Eoöphila** Swinhoe

Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 442 (1900). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 152 (1928).

Theila Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 443 (1900) (Typus: *gibbosalis* Guen.).

Aulacodes (part. Hampson, Trans. Ent. Soc. Lond., p. 172, 176 (1817) (*Au. peribocalis* Walk.; *A. gibbosalis* Guenée).

(Typus generis: *peribocalis* Walk.)

- peribocalis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 446 (1859) (*Cataclysta*) (Hindostan). — Hampson, Ill. Typ. Sp. Lep. Het. Brit. Mus., 8, p. 40 (1891) (*Oligostigma*); Fauna Brit. India Moths, 4, p. 215, f. 123 (♂) (1896) (*Aulacodes*); Trans. Ent. Soc. Lond., p. 172, fig. (♂) (1897) (*Aulacodes*). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 442 (1900) (*Eoöphila*).
- halialis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 447 (♀) (1859) (*Cataclysta*) (China). — Swinhoe, Proc. Zool. Soc. Lond., p. 868 (1885) (*Cataclysta*).
- sobrina** Pryer, Cist. Ent., 2, p. 232, t. 4, f. 3 (1875) (*Cataclysta*) (Chekiang).
- papulalis** Snellen, Trans. Ent. Soc. Lond., p. 640 (1890) (*Oligostigma*) (Sikkim).
Biologie: Pruth, Rec. Indian Mus., 30, p. 353—356, t. 11, f. 1—8 (1928).
- conjunctalis** Wileman et R. South, Entom., 50, p. 176 (♂, ♀) (1917) (*Aulacodes*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 25, 152 (1928) (*Eoöphila*).
- aulacodealis** Strand, Ent. Mitt., 8, p. 106 (1919) (*Oligostigma*).
- gibbosalis** Guenée, Delt. et yPral., p. 262 (1854) (*Oligostigma*) (Ostindien). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 431 (1859) (*Oligostigma*). — Snellen, Tijds. v. Ent., 19, p. 189, 207, t. 9, f. 12a—e (1876) (*Oligostigma*); 27, p. 49 (1884) (*Oligostigma*). — Hampson, Trans. Ent. Soc. Lond., p. 176 (1897) (*Aulacodes*). — Swinhoe, Cat. East. Austral. Lep. Het. Mus. Oxford, 2, p. 443 (1900) (*Thecla*). — Sauber in Semper, Schm. Philipp, 2, p. 657 (1902). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 153, t. 6, f. 12 (1928).
- plicatalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1332 (1865) (*Oligostigma*) (Gelebes).
- tripunctalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1531 (♂, ♀) (1865) (*Oligostigma*) (Java). — Meyrick, Trans. Ent. Soc. Lond., p. 473 (1894) (*Oligostigma*).
- hirsuta** Semper, Schm. Philipp, 2, p. 657, t. 60, f. 6 (♂) (1902) (*Thecla*).
- snelleni** Semper, Schmett. Philipp., 2, p. 656, t. 66, f. 4, 5 (♂, ♀) (1902).
- waigaoalis** Swinhoe, Cat. East. Austral. Lep. Het. Mus. Oxford, 2, p. 442, t. 6, f. 29 (♂) (1900) (*Eoöphila*). — Hampson, Trans. Ent. Soc. Lond., p. 177 (1897) (sine descr.) (*Aulacodes*).
- Indien,
Himalaya,
Birma,
West-,
Zentr-, Ost-
und
Südchina,
Tonkin,
Aden,
Ceylon
- Formosa
- Indien,
Himalaya,
Sikkim,
Formosa,
Celebes,
Sumatra,
Java,
Sumbawa,
Philippinen,
Brit.
Neu-Guinea
- Philippinen
- Philippinen
- Waigau

48. Genus **Aulacodes** Guenée

Guenée, Delt. et Pyral., p. 258 (1854). — Hampson, Fauna Brit. India Moths, 4, p. 212 (1896); Trans. Ent. Soc. Lond., p. 133, 171 (1897).

Hydrophysa Guenée, Delt. et Pyral., p. 259 (1854) (Typus: *psyllalis* Guenée).

Chnaura Lederer, Wien. Ent. Monatschr., 7, p. 435 (1863) (Typus: *octavialis* Led. = *secutalis* Walk.).

(Typus generis: *aechmialis* Guen.).

a) **Orientalische Region.**

- polydora** Meyrick, Trans. Ent. Soc. Lond., p. 82 Celebes,
(♂) (1897) (*Oligostigma*). — Hampson, Sangir
Trans. Ent. Soc. Lond., p. 172 (1897). (Typus:
ochreipicta Meyrick (nec Moore 1887), Sangir)
Trans. Ent. Soc. Lond., p. 472 (1894) (*Oligostigma*).
- crassicornalis** Guenée, Delt. et Pyral., p. 261 (1854) Java,
(*Oligostigma*) (Java). — Snellen, Tijds. v. Sumatra,
Ent., 19, p. t. 9, f. 8a—e (1876) (*Oligostigma*). — Hampson, Trans. Ent. Soc. Lombok,
Lond., p. 172 (1897). — E. Hering, Ent. Celebes
Zeit. Stettin, 64, p. 57 (1903) (*Oligostigma*).
- javanalis** Guenée, Delt. et Pyral., p. 262
(1854) (*Oligostigma*) (Java).
- bipunctalis** Walker, Cat. Lep. Het. Brit. (Java)
Mus., 34, p. 1531 (♂, ♀) (1865) (*Oligostigma*) (Java).
- Biologie: Liftinck, Treubia, 14, p. 132—
134, f. 1 (1932).
- aureolalis** Snellen, Tijds. v. Ent., 19, p. 200 (♂, ♀), Java,
t. 8, f. 5, a—e (1876) (*Oligostigma*). — Sumbawa
Hampson, Trans. Ent. Soc. Lond., p. 173 (Typus:
(1897) (*Aulacodes*). — E. Hering, Ent. Zeit. Java)
Stettin, 64, p. 59 (1903) (*Oligostigma*).
- colonialis** Guenée, Delt. et Pyral., p. 262 (1854) Bombay,
(*Oligostigma*) (Bombay). — Snellen, Tijds. Sikkim
v. Ent., 19, p. 193, t. 8, f. 8, a—d (1886)
(*Oligostigma*). — Hampson, Fauna Brit.
India Moths, 4, p. 213 (♂) (1896); Trans.
Ent. Soc. Lond., p. 173 (1897).
- idiotis** Meyrick, Trans. Ent. Soc. Lond., p. 472 (♂) Celebes
(1894). — Hampson, Trans. Ent. Soc. Lond.,
p. 173 (1897).
- metriodora** Meyrick, Trans. Ent. Soc. Lond., p. 83 Talaut
(♂) (1897) (*Oligostigma*). — Hampson,
Trans. Ent. Soc. Lond., p. 173 (1897).
- endosaris** Meyrick, Trans. Ent. Soc. Lond., p. 472 Pulo Laut
(♂) (1894) (*Oligostigma*). — Hampson,
Trans. Ent. Soc. Lond., p. 173 (1897).
- ceratucha** Meyrick, Trans. Ent. Soc. Lond., p. Pulo Laut
471 (♂, ♀) (1894) (*Oligostigma*). — Hamp-
son, Trans. Ent. Soc. Lond., p. 173 (1897).

- sejunctalis** Snellen, Tijds. v. Ent., 19, p. 207 (♂, ♀) t. 9, f. 11, a—c (1876) (*Oligostigma*); in Veth, Midden-Sumatra, 4 (1) 8, p. 77 (1880) (*Oligostigma*). — Hampson, Fauna Brit. India Moths, 4, p. 213 (1896) (*Aulacodes*); Trans. Ent. Soc. Lond., p. 174 (1897) (*Aulacodes*). — E. Hering, Ent. Zeit. Stettin, 64, p. 58 (1903) (*Aulacodes*).
delicata Moore, Lep. Ceylon, 3, p. 556, t. 215, f. 5 (♂) (1887) (*Cataclysta*).
- simplicialis** Snellen, Tijds. v. Ent., 19, p. 201 (♂), t. 8, f. 6 (1876) (*Oligostigma*).
plicatilis Walk., Hampson, Trans. Ent. Soc. Lond., p. 174 (1897).
 Biologie: Muir et Kershaw, Proc. Ent. Soc. Lond., p. XL, F. 1—4 (1909).
- parapomasalis** Hampson, Trans. Ent. Soc. Lond., p. 175 (♂) (1897).
- hemimelaena** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 470 (♀) (1917).
- hamalis** Snellen, Tijds. v. Ent., 19, p. 199 (♂), t. 8, f. 4, a—c (1876) (*Oligostigma*); in Veth, Midden-Sumatra, 4 (1) 8, p. 77 (1888) (*Oligostigma*). — Hampson, Fauna Brit. India Moths, 4, p. 214 (1896); Trans. Ent. Soc. Lond., p. 175 (1897). — E. Hering, Ent. Zeit. Stettin, 64, p. 58 (1903) (*Oligostigma*).
- melanops** Hampson, Fauna Brit. India Moths, 4, p. 214 (♂) (1896); Trans. Ent. Soc. Lond., p. 175 (1897).
- ochripieta** Moore, Descr. Lep. Ins. Atkinson, p. 209 (1887) (*Cataclysta*). — Caradja, Mem. Acad. Rom. Sect. Stiint., s. 3, 3, mem. 7, p. 332 (1925) (*Aulacodes*).
hamalis (part.) Hampson, Trans. Ent. Soc. Lond., p. 175 (1897).
- sinensis** Hampson, Trans. Ent. Soc. Lond., p. 176 (♂, ♀) (1897). — Leech, Trans. Ent. Soc. Lond., p. 437 (1901).
- saturatalis** Snellen, Trans. Ent. Soc. Lond., p. 639 (1890) (*Oligostoma*). — Hampson, Fauna Brit. India Moths, 4, p. 215 (1896).
saturalis Hampson, Trans. Ent. Soc. Lond., p. 176 (1897).
- simplex** West, Nov. Zool., 36, p. 214 (♀) (1931).
- dominalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1337 (1865) (*Cataclysta*). — Hampson, Fauna Brit. India Moths, 4, p. 216 (1896); Trans. Ent. Soc. Lond., p. 176 (1897).
peribocalis Hampson, Ill. Typ. Sp. Lep. Het. Brit. Mus., 8, p. 40, t. 156, f. 4 (1891) (*Oligostigma*). (nec Walker).

N.-Indien,
Himalaya,
Sikkim,
Ceylon,
Sumatra
(Typus:
Pundjab)

Java

Flores, Alor

Philippinen

N. u. W.-
Himalaya,
Nilgiris,
Sikkim,
Assam,
Tonkin,
Sumatra
(Typus:
Pundjab)

Sikkim

N.W.-
Himalaya,
Nilgiris,
West- und
Südechina

West- und
Zentralchina
(Typus:
Omci-Shan)

Sikkim

Philippinen

Südindien,
Nilgiris

- klilmai** Bryk (n. novum).
tripunctalis Snellen (nec Walker, 1865), Java,
 Tijds. v. Ent., 19, p. 205, t. 9, f. 9 (1876) Sumbawa
 (*Oligostigma*). — Hampson, Trans. Ent. Soc. (Typus:
 Lond., p. 176 (1897). Malang &
 Ardjoeno)
- mormodes** Meyrick, Trans. Ent. Soc. Lond., p. 83 Borneo,
 (1897) (*Oligostigma*). — Hampson, Trans. Sangir
 Ent. Soc. Lond., p. 176 (1897).
- splendens** West, Novit. Zool., 36, p. 214 (♂) (1931). Philippinen
- latifascialis** Snellen, Tijds. v. Ent., 19, p. 202, t. 8, Celebes
 f. 7, a, b (1876) (*Oligostigma*). — Hamp-
 son, Trans. Ent. Soc. Lond., p. 177 (1897).
- argyropis** Meyrick, Trans. Ent. Soc. Lond., p. 473 Pulo Laut,
 (♂, ♀) (1894) (*Oligostigma*). — Hampson, Celebes
 Trans. Ent. Soc. Lond., p. 179 (1897). — (Typus)
 Hering, Ent. Zeit. Stettin, 64, p. 59 (1903).
- basilissa** Meyrick, Trans. Ent. Soc. Lond., p. 474 Sumbawa,
 (♂) (1894); Hampson, ibid., p. 179 (1897). Timor,
 Buru
 (Typus:
 Sumbawa)
- gephyrotis** Meyrick, Trans. Ent. Soc. Lond., p. 84 Talaut
 (♀) (*Oligostigma*). — Hampson, Trans. Ent.
 Soc. Lond., p. 179 (1897).

Species incertae sedis.

- dilucidalis** (Boisduval) Guérin, Règne Anim. Ic. Java
 Ins., t. 90, f. 17, p. 524 (1844).
nectalis Snellen, Tijds. v. Ent., 19, p. 206,
 t. 9, f. 10 (1876) (*Oligostigma*).
- fuscicostalis** W. Rothschild, Novit. Zool., 22, p. 224 (♀) (1915). Ceram
- laminialis** Hampson, Trans. Ent. Soc. Lond., p. 437 Zentral- u.
 (1901). Westchina
- mimeticalis** Caradja, Mem. Acad. Rom. Sect. Stiint., Südchina
 s. 3, 3, mem. 7, p. 332, t. 1, f. 34 (1925).
- reductalis** Caradja, Mem. Acad. Rom. Sect. Stiint., Südchina
 s. 3, 3, mem. 7, p. 332, t. 1, f. 35 (1925).
- stresemanni** W. Rothschild, Novit. Zool., 22, p. 223 Ceram
 (♂, ♀) (1915).

b) Australische Region.

- trichoceralis** Hampson, Trans. Ent. Soc. Lond., p. 174 (♂) (1897). Neu-Guinea
 (Typus:
 Humboldt-
 Bay)
- cyclozonalis** Hampson, Ann. Mag. Nat. Hist., (s. Brit.
 7) 18, p. 467 (♀) (1906). Neu-Guinea
- heptopis** Hampson, Trans. Ent. Soc. Lond., p. 174 Fergusson
 (♀) (1897). Ins.

- quadriplagiata** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 471 (♂) (1917). Goodenough Ins.
- siennata** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 206 (1896) (*Oligostigma*). — Hampson, Trans. Ent. Soc. Lond., p. 177 (1897).
subsp. *intensa* W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 120 (1915). Holl. Neu-Guinea
- costifascialis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 171 (1917). Goodenough Ins., Rook Ins.
- nigriplagialis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 472 (1917). Holl. Neu-Guinea
trigonalis ab. 1, Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 390 (1906) (*Eristena*).
- dolichoplagia** Hampson, Ann. Mag. Nat. Hist., (s. 8) 19, p. 472 (♂) (1917). Holl. Neu-Guinea
- acroperalis** Hampson, Trans. Ent. Soc. Lond., p. 177 (1897). Fergusson Ins., Brit. Neu-Guinea, Zentral-Ceram (Typus: Fergusson)
- diopsalis** Hampson, Trans. Ent. Soc. Lond., p. 177 (1897). Amboina, Fergusson Ins., Brit. Neu-Guinea
- subsp. *major* W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 120 (♀) (1915). Holl. Neu-Guinea
- mesoscialis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 201 (1917). Holl. Neu-Guinea
- hemithermalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 202 (♂, ♀) (1917). Holl. Neu-Guinea
- chrysoxantha** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 202 (♀) (1917). Holl. Neu-Guinea
- metazonalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 203 (♀) (1917). Bismarck-Archipel (Rook Ins.)
- aurantipennis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 204 (♂) (1917). Holl. Neu-Guinea
- argenteopieta** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 204 (1917). Holl. Neu-Guinea
- brunnealis** Hampson, Trans. Ent. Soc. Lond., p. 178 (♀) (1897). Holl. Neu-Guinea (Typus: Humboldt-Bay)

- similis** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 147 (♀) (1915). Holl.
Neu-Guinea
- mediofascialis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 205 (♀) (1917). Holl.
Neu-Guinea
- cervinalis** Hampson, Trans. Ent. Soc. Lond., p. 178 (1897). Neu-Guinea
(Typus:
Humboldt-
Bay)
- subsp. *distincta* W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 120 (♀) (1915). Holl.
Neu-Guinea
- junctiscriptalis** Hampson, Trans. Ent. Soc. Lond., p. 178 (♂, ♀) (1897). Fergusson
Ins.,
Neu-Guinea
- metataxalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 467 (♂) (1906). Neu-Guinea
- thermichrysis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 205 (♂) (1917). Bismarck-
Archipel
(Ins. Rook)
- coniferalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 206 (1917). Holl.
Neu-Guinea
- flavifascialis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 207 (♀) (1907). Holl.
Neu-Guinea
- argentimaculalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 207 (♂) (1917). Holl.
Neu-Guinea
- pentopalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 468 (♀) (1906). Salomon Ins.
- longiplagialis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 208 (♀) (1917). Holl.
Neu-Guinea
- obliquivitta** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 208 (♀) (1917). Holl.
Neu-Guinea
- secutalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1291 (♂) (1865) (*Syngamia*) (Mysol). — Mysol,
Salomon
Hampson, Trans. Ent. Soc. Lond., p. 100 Ins., Neu-
(1897). Hebriden
- octavialis* Lederer, Wien. Ent. Monat-
schr., 7, p. 435 (1863) (*Chnaura*).
- goniophoralis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 18, p. 463 (♂) (1906). Holl.
Neu-Guinea
- bipunctalis** Kenrick, Proc. Zool. Soc. Lond., p. 76, t. 3, f. 57 (1907). Brit. u.
Holl.
Neu-Guinea

Species incertae sedis.

- eupselias** Meyrick, Trans. Ent. Soc. Lond., 77, p. 161 (♂) (1929). Marquesas
Ins.
(Typus:
Tahuate)

- lunalis** Kenrick, Proc. Zool. Soc. Lond., p. 76, t. 3, f. 55 (1907). Brit. Neu-Guinea
- nephelanthopa** Meyrick, Exot. Microlep, 4, p. 499 (♀) (1934). Fidji Ins. (Typus: Wunidawa)
- ovomaculalis** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 121 (♀) (1915). Holl. Neu-Guinea
- postbasalis** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 122 (♀) (1915). Holl. Neu-Guinea
- pulchralis** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 121 (♂) (1915). Holl. Neu-Guinea
- purpurcalis** Kenrick, Proc. Zool. Soc. Lond., p. 76, t. 3, f. 56 (1907). Br. u. Holl. Neu-Guinea
- rufocastanea** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 148 (♂) (1915). Holl. Neu-Guinea
- triplaga** Lower, Trans. Roy. Soc. S. Austral., 27, p. 62 (1903). Queensland
- uniplagialis** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 121 (♂) (1915). Holl. Neu-Guinea
- wollastoni** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 121, t. 2, f. 58 (♀) (1915). Holl. Neu-Guinea

c) Neotropische Region.

- psyllalis** Guenée, Delt. et Pyral., p. 259, t. 6, f. 11 (1854) (*Hydrophysa*) (Brasilien). — Hampson, Trans. Ent. Soc. Lond., p. 179 (1897). Westindien, Grenada, Trinidad,
- scutthesalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 950 (1859) (*Oligostigma?*) (Brasilien). Venezuela, Brasilien, Guatemala
- tortalis** Lederer, Wien. Ent. Monatschr., 7, p. 452, 484, t. 18, f. 3 (1863) (*Hydrocampa*) (Venezuela). — Druce, Biol. Centr. Amer. Het., 2, p. 277 (1896).
- obtusalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 296 (♂) (1914). Panama
- congallalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 117 (♀) (1924). Guatemala (Typus: Cayuga)
- julittalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 118 (♀) (1924). Pauama (Typus: Alhajudo)
- aechmialis** Guenée, Delt. et Pyral., p. 258 (1854) (*Aulacodes*) (Cayenne). Westindien, Fr. Guyana,
- oechmialis** Hampson, Trans. Ent. Soc. Lond., p. 179 (1897). Venezuela, Brasilien,
- fragmentalis** Lederer, Wien. Ent. Monatschr., 7, p. 453, t. 18, f. 6 (1863) (*Parapognax*). Guatemala, Panama

- gothicalis* Felder et Rogenhofer, Reise Novara (Het.), p. 15, t. 136, f. 13 (1874) (*Parapognæ?*) (Amazonas).
- bosoralis* Druce, Biol. Centr.-Amer. Het., 2, p. 278, t. 63 (♂) f. 31 (1896) (*Hydro-campa*). (Typus: Guatemala (Teleman in Vera Paz))
- plumipcdalis* Hampson, Ann. Mag. Nat. Hist., (s. 6) 16, p. 343 (1895) (*Hydro-campa*).
- halitalis** Felder et Rogenhofer, Reise Novara (Het.), p. 15, t. 136, f. 12 (1874) (*Parapognæ*) (Amazonas). — Hampson, Trans. Ent. Soc. Lond., p. 180 (1897). Brasilien (Amazonas)
- convoluta** Hampson, Trans. Ent. Soc. Lond., p. 180 (♂) (1897). Amazonas (Typus: Javary-Fluß)
- hodevalis** Druce, Biol. Centr.-Amer. Het., 2, p. 278, t. 63, f. 30 (♂) (1896). — Hampson, Trans. Ent. Soc. Lond., p. 180 (1897). Guatemala (Typus: San Geronimo)
- citronalis** Druce, Biol. Centr.-Amer. Het., 2, p. 277, t. 63, f. 28 (♂) (1896). — Hampson, Trans. Ent. Soc. Lond., p. 100 (1897). Mexiko (Omiteme), Costa Rica (Volcan de Irazu)
- traversalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 296 (♂) (1914). Panama
- nissenalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 118 (♀) (1924). Peru (Typus: Callao)
- fiachmalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 118 (♀) (1924). Panama (Typus: Alhajuelo)
- grimbaldalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 119 (♀) (1924). Guatemala (Typus: Cayuga)
- semicircularis** Hampson, Trans. Ent. Soc. Lond., p. 180 (♀) (1897). Amazonas, Espiritu Santo
- calistoalis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 442 (♀) (1859) (*Cataclysta*) (Santarem). — Hampson, Trans. Ent. Soc. Lond., p. 181 (1897). Südamerika
- odulphalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 119 (♀) (1924). Guatemala (Typus: Cayuga)
- thetysalis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 440 (♀) (1859) (*Cataclysta*) (Villa Nova). — Hampson, Trans. Ent. Soc. Lond., p. 181 (1897). Brasilien

- adjutrealis** Schaus, Proc. Ent. Soc. Wash., 26, p. 120 (♂) (1924). Peru
(Typus:
Col. Perene)

Species incertae sedis.

- confusalis** Schaus, Proc. U. S. Mus., 30, p. 139 (1906). Brasilien
(Paraná)
- delicata** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, p. 679 (♂) (1912). Costa Rica
- moralis** Schaus, Proc. U. S. Mus., 30, p. 138 (1906). Brasilien
(S. Paulo)
- pampalis** Schaus, Proc. U. S. Nat. Mus., 30, p. 139 (1906). Brasilien
(Paraná)
- reversalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 295 (♂) (1914). Panama
- templalis** Schaus, Proc. U. S. Nat. Mus., 30, p. 138 (1906). Brasilien
(Paraná)

99

49. Genus **Decticogaster** Snellen

- Snellen, Tijds. v. Ent., 23, p. 230 (Sept.) (1880); 42, p. 84 (1899).
— Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 159 (1928).
(Typus generis: *D. zonulalis* Snell. = *Botys varanensis* Walk.)

Tatobotys Butler, Proc. Zool. Soc. Lond., p. 686 (Dez.) (1880) (Typus: *argillacea* Butl. = *janapalis* Walk.). — Hampson, Fauna Brit. India Moths, 4, p. 224 (1896); Trans. Ent. Soc. Lond., p. 196 (1897). — Pagenstecher, Zoologica II. 29, p. 176 (1900).

Cometura Meyrick, Trans. Ent. Soc. Lond., p. 226 (1886) (Typus: *C. picogramma* Meyr. = *Botys biannulalis* Walk.). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 445 (1900).

- janapalis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 676 (♀) (1859) (*Botys*) (Ceylon). — Indien,
Moore, Lep. Ceylon, 3, p. 342 (1886) (*Ha- Birma,*
palia?). — Hampson, Fauna Brit. India Ostchina
Moths, 4, p. 224, f. 131 (♂) (1896) (*Tato- (Kwangtung)*
botys); Trans. Ent. Soc. Lond., p. 196, fig. Tonkin,
(♂) (1897) (*Tatobotys*). — Pagenstecher, Formosa,
Zoologica H. 29, p. 176 (1900). — Shibuya, Ceylon,
Journ. Fac. Agr. Hokkaido Imp. Univ., 22, Borneo,
p. 159 (1928) (*Decticogaster*). Ceram,
japanalis Snellen, Tijds. v. Ent., 42, p. 84 Philippinen,
(1899) (*Tatobotys*). Bismarek-
cunealis Walker, Cat. Lep. Het. Brit. Mus., Archipel,
34, p. 1420 (♀) (1865) (*Botys*) (Ceram). — Salomon
Swinhoe, Trans. Ent. Soc. Lond., p. 272 Ins.,
(1890) (*Tatobotys*). Australien,
argillacea Butler, Proc. Zool. Soc. Lond., Queensland
p. 686 (1881) (*Tatobotys*). — Meyrick, Trans.
Ent. Soc. Lond., p. 214 (1887) (*Lonchodes*).

- ceramochra* Meyrick, Trans. Ent. Soc. Lond., p. 443 (♂) (1885) (*Lonchodes*); p. 214 (1887) (*Lonchodes*) (= *argillacea* Butl.). Typus: Duaringa (Queensland)
- varanesalis* Walker, Cat. Lep. Het. Brit. Mus., 18, p. 719 (1859) (*Botys*) (Sarawak). — Hampson, Fauna Brit. India Moths, 4, p. 225 (♂) (1896) (*Tatobotys*); Trans. Ent. Soc., p. 196 (1897) (*Tatobotys*). Birma, Borneo
- zonulalis* Snellen, Tijds. v. Ent., 23, p. 231 (1880) (*Decticogaster*); 27, p. 40, t. 3, f. 11 (1884) (*Decticogaster*). — Swinhoe, Trans. Ent. Soc. Lond., p. 272 (1890) (*Decticogaster*). (Typus: Balangnipa, Makassar)
- biannulalis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1439 (♂) (1865) (*Botys*) (Sula). — Hampson, Trans., Ent. Soc. Lond., p. 197 (1897) (*Tatobotys*). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 445 (1900) (*Cometura*). — Hampson, Journ. Bombay Nat. Hist. Soc., 18, p. 574 (1908) (*Tatobotys*). — Swinhoe, Ann. Mag. Nat. Hist., (s. 7) 17, p. 287 (1906). Birma, Tonkin, Sula Ins., Neu-Hebriden, Philippinen, Fidji Ins., Samoa
- picrogramma* Meyrick, Trans. Ent. Soc. Lond., p. 226 (♂, ♀) (1886) (*Cometura*) (Fidji). — Swinhoe, Trans. Ent. Soc. Lond., p. 276 (1890) (*Cometura*). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 445 (1900) (= *lirisalis* Walk.). Typus: Fidji
- gemella* Moore, Descr. Lep. Ins. Atkinson, p. 208 (1888) (*Hedylepta*) (Calcutta).
- janapalis* (part.) Hampson, Trans. Ent. Soc. Lond., p. 196 (1897) (= *gemella*) (*Tatobotys*).
- albivenalis* Hampson, Trans. Ent. Soc. Lond., p. 197 (♀) (1897) (*Tatobotys*). Bonin Ins., (Typus), ? Buru, Batchian, Salomon Ins., Philippinen
- aurantialis* Hampson, Trans. Ent. Soc. Lond., p. 197 (♂) (1897) (*Tatobotys*).
- tanyscia* West, Nov. Zool., 36, p. 215 (♂) (1931) (*Tatobotys*).

Species incertae sedis.

- amoyalis* Caradja, Bull. Acad. Roum., 15, p. 148 (♂) (1932) (*Cometura*). Amoy
- angustalis* Caradja, Deutsche Ent. Zeitschr. Iris, 47, p. 155 (1934). Südchina (Kwangtung)
- depalpalis* Strand, Ent. Mitt., 8, p. 130 (♂) (1919) (*Tatobotys*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 159, 160 (1928) (*Decticogaster*). Formosa

- leuculalis** Snellen, Tijds. v. Ent., 42, p. 83 (♂) t. 4, f. 8 (♂) (1899) (*Cometura*). Loyalty Ins.
- lirialis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 1003 (♀) (1859) (*Botys*) (Sarawak). — Meyrick, Trans. Ent. Soc. Lond., p. 468 (1894) (*Cometura*). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 445 (1900) (*Cometura*).
 Hinter-Indien, Borneo, Philippinen (Luzon), Fidji Ins., Viti Ins.
- janapalis* (part.) Hampson, Trans. Ent. Soc. London, p. 196 (1887) (*Tatobotys*).
- bistrigulalis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1415 (1865) (*Botys*) (Sarawak). — Hampson, Trans. Ent. Soc. Lond., p. 196 (1894) (= *janapalis* Walk.). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 445 (1900).
- vibrata** Meyrick, Trans. Ent. Soc. Lond., 77, p. 161 (♂) (1929) (*Tatobotys*).
 12 Marquesas Ins.
 (Typus: Hiva Oh)

50. Genus **Almonia** Walker

Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1495 (♂) (1865). — Hampson, Fauna Brit. India Moths, 4, p. 225 (1896); Trans. Ent. Soc. Lond., p. 133, 197 (1897).

Dodanga Moore, Lep. Ceylon, 3, p. 331 (1886). (Typus generis: *D. lobipennis* Moore = *Botys truncatalis* Walk.).

(Typus generis: *A. truncatalis* Walk.)

- truncatalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1440 (♀) (1865) (*Botys*) (Java). — Meyrick, Trans. Ent. Soc. Lond., p. 86 (1897) (*Dodanga*). — Hampson, Fauna Brit. India Moths, 4, p. 226, f. 132 (♂) (1896) (*Almonia*); Trans. Ent. Soc. Lond., p. 198 (1897). — Swinhoe, East. Austral. Lep. Het. Oxford Mus., 2, p. 446 (1900).
 Indien (Nilgiris), Ceylon, Java, Sula Ins., N.-Celebes, Sangir
- onustalis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1495 (♂) (1865) (*Almonia*) (Sula).
- lobipennis* Moore, Lep. Ceylon, 3, p. 332, t. 182, f. 3 (1885) (*Dodanga*) (Ceylon). — Hampson, Fauna Brit. India Moths, 4, p. 226 (♂) (1896) (*Almonia*); Trans. Ent. Soc. Lond., p. 198 (1897). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 446, (1900) (= *truncatalis* Walk.).
- cristata** Hampson, Ill. Typ. Sp. Lep. Het. Brit. Mus., 8, p. 136, t. 156, f. 7 (1891); Fauna Brit. India Moths, 4, p. 226 (♂) (1896); Trans. Ent. Soc. Lond., p. 198 (1897).
 Ganjam, Nilgiris
- atratalis** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 125, t. 2, f. 40 (1915).
 3 Holl. Neu-Guinea

51. Genus **Bradina** Lederer

Lederer, Wien. Ent. Monatschr., 7, p. 424 (1863). — Hampson, Fauna Brit. India Moths, 4, p. 226 (1896); Trans. Ent. Soc. Lond., p. 113, 198 (1897). — Pagenstecher, Zoologica H. 29, p. 177 (1900).

Erilita Lederer, Wien. Ent. Monatschr., 7, p. 426, (1863) (Typus: *modestalis* Led.). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 446 (1900). — E. Hering, Ent. Zeit. Stettin, 64, p. 46 (1903).

Pleonectusa Lederer, Wien. Ent. Monatschr., 7, p. 426 (1863) (Typus: *admixtalis* Walk.). — E. Hering, Ent. Zeit. Stettin, 64, p. 46 (1903). — Moore, Lep. Ceylon, 3, p. 286 (1885).

Physematia Lederer, Wien. Ent. Monatschr., 7, p. 447 (1863) (Typus: *concordalis* Lederer).

Trematarcha Meyrick, Trans. Ent. Soc. Lond., p. 233 (1886) (Typus: *erilitalis* Feld. et Rogenh. = *rectiferalis* Walker).

Epichronistis Meyrick, Trans. Ent. Soc. Lond., p. 260 (1886) (Typus: *acrospila* Meyrick).

Epichromistis Hampson, Trans. Ent. Soc. London, p. 198 (1897) (ex synon.).

(Typus generis: *impressalis* Leder.)

a) Paläarktische Region.

atopalis Walker, Cat. Lep. Het. Brit. Mus., 18, p. 664 (♂) (1859) (*Botys*) (Shanghai). — Nord-, Pryer, Trans. Asiat. Soc. Jap., 13, p. 66 West-, (1885) (*Botys*). — Hampson, Trans. Ent. Zentral- u. Soc. Lond., p. 200 (1897) (*Bradina*). — Südchina, Leech, Trans. Ent. Soc. Lond., p. 439 (1901). Korea, — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Japan, Univ., 22, p. 160, 161, t. 6, f. 13 (1928); Formosa Ins. Matsum., 3, p. 130, t. 5, f. 18 (1929). — Caradja, Bull. Acad. Roum., 15, p. 148 (1932).

damasalis Walker, Cat. Lep. Het. Brit. Mus., 18, p. 668 (♂) (1859) (*Botys*) (Shanghai).

aulacodialis Strand, Ent. Mitt., 8, p. 109 (♂) (1919) (*Bradina*) (Formosa). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 161 (1928).

andresi Rebel, Deutsche Ent. Zeitschr. Iris, 26, p. 81; p. 82, f. 10 (♂) (1912). Ägypten (Typus: ♂ Choubrah)

megesalis Walker, Cat. Lep. Het. Brit. Mus., 18, p. 663 (♂) (1859) (*Botys*) (N. China). — Nord-, Hampson, Trans. Ent. Soc. Lond., p. 200 Zentral- u. (1897). — Leech, Trans. Ent. Soc. Lond., Westchina, Japan p. 439 (1901). — Shibuya, Ins. Matsum.,

3, p. 130 (1929). — Caradja, Bull. Acad. Roum., 15, p. 148 (1932).

- admixtalis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 665 (♂, ♀) (1859) (*Botys*) (Ceylon). — Lederer, Wien. Ent. Monatschr., 7, p. 426, 481 (1863) (*Pleonectusa*). — Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1480 (1865) (*Pleonectusa*). — Moore, Lep. Ceylon, 3, p. 286, t. 180, f. 13 (1885) (*Pleonectusa*). — Snellen, Trans. Ent. Soc. Lond., p. 624 (1890) (*Pleonectusa*). — Hampson, Ill. Typ. Sp. Lep. Het. Brit. Mus., 8, p. 40 (1891) (*Pleonectusa*); 9, p. 50 (1893) (*Pleonectusa*); Fauna Brit. India Moths, 4, p. 227 (1896); Trans. Ent. Soc. Lond., p. 201 (1897). — Pagenstecher, Zoologica, H. 29, p. 177 (1900). — Swinhoe, Cat. East. Austral., Lep. Het. Oxford Mus., 2, p. 446 (1900) (*Erilita*). — Leech, Trans. Ent. Soc. Lond., p. 440 (1901). — E. Hering, Ent. Zeit. Stettin, 64, p. 46 (1903) (*Pleonectusa*). — Marumo, Journ. Col. Agr. Tokyo, 8, p. 187 (1923). — Caradja, Mem. Acad. Rom. Sect. Stiint., (s. 3) 4, mem. 8, p. 405 (1927). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 16 (1928); Ins. Matsum., 3, p. 130, t. 5, f. 19 (1929).
- panaeusalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 998 (♂) (1859) (*Botys*) (Ceylon).
- tabidalis** Lederer, Wien. Ent. Monatschr., 7, p. 426, 481 (1863) (*Pleonectusa*) (Amboina). — Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1480 (1865) (*Pleonectusa*). — Snellen in Veth, Midden-Sumatra, 4, p. 77 (1880); Tijds. v. Ent., 27, p. 40 (1884); Trans. Ent. Soc. Lond., p. 624 (1890). — Pagenstecher, Jahrb. Nassau Ver. Naturk., p. 112 (1888). — E. Hering, Ent. Zeit. Stettin, 64, p. 47 (1903) (*Pleonectusa*).
- sodalis** Lederer, Wien. Ent. Monatschr., p. 426, 481 (1863) (*Pleonectusa*) (Amboina). — Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1480 (1865) (*Pleonectusa*). — Pagenstecher, Jhb. Nassau. Ver. Naturk., 41, p. 277 (1888); in Kückenthal, Forschungsreise Lep., p. 464 (1897); Zoologica H. 29, p. 177 (1900).
- leptogastralis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1432 (♂) (1865) (*Botys*) (Ceram).
- pallidalis** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 147 (1896) (*Pleonectusa*) (Assam).
- lophophoralis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 9, p. 435 (♂) (1912) (*Nacoleia*). — Wilman, Trans. Ent. Soc. Lond., p. 379 (1911) (*Nacoleia*) (non descr.) (Hondo). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 121, 130, 131 (1928) (*Bradina*).

Japan,
Malediven,
West- u.
Südchina,
Formosa,
Indien,
Tonkin,
Perak,
Malakka,
Singapore,
Hainan,
Sumatra,
Ceylon,
Java, Ceram,
Celebes,
Amboina,
Nikobaren,
Chagos Ins.,
Philippinen,
Queensland,
Bismarck-
Archipel,
Bali,
Louisaden
(St. Aignan),
Shortland
Ins.,
Fergusson
Ins.,
Port. O.-
Afrika,
Liberia,
Goldküste,
Nyassaland,
Natal,
Madagaskar,
Fernando Po,
Komoren

Singapore,
Japan

b) Orientalische Region.

- remipes** Hampson, Trans. Ent. Soc. Lond., p. 199 (♂) (1897). Pulo Laut
- impressalis** Lederer, Wien. Ent. Monatschr., 7, p. 425, t. 15, f. 16 (1863) (Amboina). — Mysol, Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1372 (1865). — Pagenstecher, Jahrb. Nassau, Sula, Buru, Ver. Naturk., p. 112, (1886); p. 158 (1888). Aru Ins., Bismarck-Archipel, — Hampson, Trans. Ent. Soc. Lond., p. 200 (1897). — Pagenstecher, Zoologica, H. Holl. u. Brit. 29, p. 117 (1900). Neu-Guinea, Ellice Ins.
- selectalis** Lederer, Wien. Ent. Monatschr., 7, p. 425, 481, t. 16, f. 1 (1863). — Hampson, Trans. Ent. Soc. Lond., p. 200 (1897). Amboina, Buru
- concordalis** Lederer, Wien. Ent. Monatschr., 7, p. 447, t. 2, f. 27, t. 17, f. 14 (1863) (*Physematia*) (Nikobaren). — Hampson, Fauna Brit. India Moths, 4, p. 228 (1896); Trans. Ent. Soc. Lond., p. 200 (1897). — Strand, Ent. Mitt., 8, p. 132 (1919). Südchina, Nikobaren
- var. *defloralis* Strand, Ent. Mitt., 8, p. 133 (1919). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 160, 162 (1928). Formosa
- atopalis** Walker, vide sub a) Paläarktische Region, p. 149.
- adhaesalis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 664 (♂) (1859) (*Botys*) (Ceylon, Shanghai). — Moore, Lep. Ceylon, 3, p. 287, t. 180, f. 14 (1885) (*Plconectusa*). — Hampson, Fauna Brit. India Moths, 4, p. 227 (♀) (1896); Trans. Ent. Soc. Lond., p. 200 (1897). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 447 (1900) (*Erilita*). — E. Hering, Ent. Zeit. Stettin, 64, p. 46 (1903) (*Erilita*). Südiudieu, Ceylon, Borneo, Sula, Sumatra, Java, ? Japan
- aeclusalis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 687 (♂) (1859) (*Botys*) (Sarawak).
- teutalis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 690 (♀) (1859) (*Botys*) (Sarawak).
- canalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1429 (♀) (1865) (*Botys*). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 447 (1900) (*Erilita*). Mysol
- translinealis** Hampson, Fauna Brit. India Moths, 4, p. 228, f. 133 (1896); Trans. Ent. Soc. Lond., p. 199, fig. (♂); p. 200 (1897). — R. South in Leech, Trans. Ent. Soc. London, p. 440 (1901). N.W.-Himalaya
- rectilinealis** R. South in Leech, Trans. Ent. Soc. Lond., p. 440, t. 15, f. 1 (1901). Zentral- u. Westchina
- megesalis** Walker, vide sub a) Paläarktische Region, p. 149.

- bicoloralis** Hampson, Fauna Brit. India Moths, 4, Birma
p. 228 (♂) (1896); Trans. Ent. Soc. Lond.,
p. 200 (1897).
- consortialis** Caradja, Mem. Acad. Rom. Sect. Stiint., Westchina
(s. 3), 4, mem. 8, p. 405 (1927). (Szetschwan)
- melanoperas** Hampson, Fauna Brit. India Moths, 4, Birma,
p. 227 (♂) (1896); Trans. Ent. Soc. Lond., West-
p. 200 (1897). — Caradja, Mem. Acad. Rom. Mittel- u.
Sect. Stiint., (s. 3) 3, mem. 7, p. 334 (1925). Südchina,
— Shibuya, Journ. Fac. Agr. Hokkaido Imp. Formosa,
Univ., 22, p. 161 (1928). Brit.
Neu-Guinea
- subsp. *sinensis* Caradja, Mem. Acad. Rom. West-
Sect. Stiint., (s. 3) 3, mem. 7, p. 334 (1925). Mittel- u.
Südchina
- subpurpurescens** Warren, Ann. Mag. Nat. Hist., N.O.-Indien,
(s. 6) 17, p. 147 (1896) (*Pleonectusa*). — Bhotan,
Hampson, Fauna Brit. India Moths, 4, p. Assam
227 (♂) (1896); Trans. Ent. Soc. Lond., (Khasis)
p. 201 (1897).
- macaralis** Walkor, Cat. Lep. Het. Brit. Mus., 18, Perak,
709 (♂) (1859) (*Botys*) (Sarawak). — Mey- Borneo,
rick, Trans. Ent. Soc. Lond., p. 468 (1894) Java,
(*Pleonectusa*). — Hampson, Trans. Ent. Soc. Amboina,
Lond., p. 201 (1897). Sula, Salo-
mon Ins.
- admixtalis** Walker, vide sub a) Paläarktische Region,
p. 150.
- geminalis** Caradja, Mem. Acad. Rom. Sect. Stiint., West- u.
(s. 3) 4, mem. 8, p. 405 (1927); Deutsche Südchina
Ent. Zeitschr. Iris, 47, p. 155 (1934). (Szetschwan,
Kwangtung)
- dentalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) Sumba,
19, p. 4 (1907). Selaru
- extenuatalis** Walker, Cat. Lep. Het. Brit. Mus., 34, Sula,
p. 1424 (♀) (1865) (*Botys*) (Sula). — Mey- Amboina,
rick, Trans. Ent. Soc. Lond., p. 86 (1897) Mysol,
(*Pleonectusa*). — Hampson, Trans. Ent. Soc. Ceram
Lond., p. 201 (1897) (*Bradina*).
- leptomeralis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1429 (♂) (1865) (*Botys*) (Mysol,
Ceram).
- melesalis** Walker, Cat. Lep. Het. Brit. Mus., 18, Indien
p. 705 (♂) (1859) (*Botys*) (Sarawak). — (Sind,
Meyrick, Trans. Ent. Soc. Lond., p. 465 Haiderabad),
(1894) (Metasia). — Hampson, Trans. Ent. Borneo,
Soc. Lond., p. 201 (1897) (*Bradina*). Sumbawa,
sagittalis Swinhoe, Ann. Mag. Nat. Hist., Ceylon,
(s. 6) 16, p. 301 (1895) (*Pleonectusa*). — Westafrika
Hampson, Fauna Brit. India Moths, 4, p. 228
(1896).
- pumilialis** Hampson, Ann. Mag. Nat. Hist., (s. 7) Sumbawa,
19, p. 4 (1907). Pura

- modestalis** Lederer, Wien. Ent. Monatschr., 7, p. 426, t. 16, f. 3 (1863) (*Erilita*) (Amboina).
 — Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1373 (1865) (*Erilita*). — Meyrick, Trans. Ent. Soc. Lond., p. 216 (1887) (*Pleonectusa*).
 — Pagenstecher, Jahrb. Nassau. Ver. Naturk., 41, p. 112 (1888) (*Pleonectusa*). — Hampson, Trans. Ent. Soc. Lond., p. 202 (1897) (*Bradina*). — Pagenstecher, Zoologica, H. 29, p. 177 (1900) (*Erilita*).
 Java,
 Celebos,
 Sula,
 Amboina,
 Buru,
 Ceram,
 Bismarck-Arch.,
 Salomon
 Ins.,
 Ellice
 Ins.,
 Shortland
 Ins.,
 Neu-Guinea,
 Aru,
 Samoa Ins.
- chalinota** Meyrick, Trans. Ent. Soc. Lond., p. 233 (♂) (1886) (*Pleonectusa*). (Salomon-, Ellice Ins.)
- erilitoides** Strand, Entt. Mitt., 8, p. 108 (♀) (1919). Formosa
 — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 160, 162 (1928).

Species incertae sedis.

- aureolalis** De Joannis, Bull. Soc. Ent. Fr., p. 198 (1899). — Fryer, Trans. Linn. Soc. Lond., 15, p. 25 (1912). Seychellen
- ceramica** W. Rothschild, Nov. Zool., p. 224 (♂) (1915). Ceram
- intermedialis** Caradja, Bull. Acad. Roum., 15, p. 149 (♂ ♀) (1932). Südchina
- nigripunctalis** R. South in Leech, Trans. Ent. Soc. Lond., p. 440 (♀), t. 15, f. 25 (♀) (1901). Westchina
 — Caradja, Mem. Acad. Rom. Sect. Stiint., (s. 3) 3, mem. 7, p. 334 (1925). (Typus: Chia-ting-fu)
- opacusalis** Swinhoe, Trans. Ent. Soc. Lond., p. 156 (♂) (1904). Borneo

c) Aethiopische Region.

- admixtalis** Walker, vide sub a) Paläarktische Region, p. 150.
- acrospila** Meyrick, vide sub d) Australische Region, p. 154.
- melesalis** Walker, vide sub b) Orientalische Region, p. 152.
- hemiphaealis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 5 (♂) (1907). Brit. O.-Afrika
- sordidalis** Dewitz, Verh. Akad. Leop.-Car., 42, p. 88, t. 3, f. 11 (1880) (*Botys*). Lagos,
 Togo,
 Kamerun,
 Span. Guinea

d) Australische Region.

- sempnopa** Meyrick, Trans. Ent. Soc. Lond., p. 234 (♂) (1886) (*Trematarcha*). — Hampson, Trans. Ent. Soc. Lond., p. 199 (1897). Fidji Ins., Samoa Ins. (Typus: Rotumah Island)
- rectiferalis** Walker, Trans. Ent. Soc. Lond., (s. 3) 1, p. 126 (♂) (1862). (*Botys*). — Hampson, Trans. Ent. Soc. Lond., p. 199 (1897). Neu-Caledonien, Neu-Hebriden, Fidji Ins. (Typus: Neu-Caledonien)
- erilitalis** Felder et Rogenhofer, Reise Novara, (Het.), p. 13, t. 135, f. 26 (1874) (*Marasmia*). — Meyrick, Trans. Ent. Soc. Lond. p. 234 (1886) (*Trematarcha*).
- impressalis** Lederer, vide sub b) Orientalische Region, p. 151.
- affinis** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 125 (♂) (1915). Holl. Neu-Guinea
- costalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 3 (♀) (1907). Salomon Ins.
- leucura** Hampson, Trans. Ent. Soc. Lond., p. 200 (1897). — Rebel, 2. Beih. Jahrb. Hamb. Wiss. Anstalt, 32, p. 151 (1915). Fidji Ins. (Typus), Samoa
- chlorionalis** Tams, Ins. Samoa, pars (3 fasc.) 4, p. 264, t. 15, f. 4 (♂) (1935). Samoa
- pycnolopha** Tams, Ins. Samoa, pars 3, (fasc. 4), p. 264, t. 15, f. 1 (♂) (1935). Samoa
- mannasalis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 726 (♂) (1859) (*Botys*) (Moreton-Bai). — Hampson, Trans. Ent. Soc. Lond., p. 201 (1897). Australien
- leptolopha** Tams, Ins. Samoa, pars 3 (fasc. 4), p. 265, t. 15, f. 2 (♂) (1935). Samoa
- admixtalis** Walker, vide sub a) Paläarktische Region, p. 150.
- macaralis** Walker, vide sub b) Orientalische Region, p. 152.
- glaucinalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 3 (1907). Brit. u. Holl. Neu-Guinea, Salomon Ins.
- chloroscia** Meyrick, Trans. Ent. Soc. Lond., p. 232 (♂) (1886) (*Pleonectusa*). — Hampson, Trans. Ent. Soc. Lond., p. 201 (1897). Tonga Ins.
- nigropunctata** Pagenstecher, Zoologica, H. 29, p. 177 (19300). Shortland Ins.
- acrospila** Meyrick, Trans. Ent. Soc. Lond., p. 261 (♂, ♀) (1886) (*Epichronistis*). — Hampson, Trans. Ent. Soc. Lond., p. 201 (1897). Mauritius, Fidji (Typus), Tonga, Mirikoi, Malediven

- xanthalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 212 (♂, ♀) (1917). Louisiaden (St. Aignan), Admiraltäts-Ins.
- angusta** Butler, Ann. Mag. Nat. Hist., (s. 5) 10, p. 234 (1882) (*Asopia*). — Hampson, Trans. Ent. Soc. Lond., p. 201 (1897). Neu-Britannien, Herzog von York Ins., Freundschafts-Ins.
- parallela** Meyrick, Trans. Ent. Soc. Lond., p. 231 (1886) (*Pleonectusa*). — Hampson, Trans. Ent. Soc. Lond., p. 201 (1897). Fidji Ins.
- horatius** Butler, Trans. Ent. Soc. Lond., p. 430 (1886) (*Botys*). (Viti)
- aurata** Butler, Ann. Mag. Nat. Hist., (s. 5) 20, p. 124 (1887) (*Pleonectusa*). — Hampson, Trans. Ent. Soc. Lond., p. 201 (1897). Salomon Ins.
- metaleucalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1372 (♀) (1835). — Meyrick, Trans. Ent. Soc. Lond., p. 231 (1886) (*Pleonectusa*). — Hampson, Trans. Ent. Soc. Lond., p. 201 (1897). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 447, t. 6, f. 27 (1900) (*Erilita*). Fidji Ins.
- albigenitalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 212 (♂) (1917). Salomon Ins.
- argentata** Butler, Ann. Mag. Nat. Hist., (s. 5) 20, p. 124 (1887). — Hampson, Trans. Ent. Soc. Lond., p. 201 (1897). Salomon Ins.
- puuctilinealis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 4 (♂) (1907). Fidji Ins.
- antisema** Meyrick, Trans. Ent. Soc. Lond., p. 263 (♂) (1886) (*Isopteryx*). — Hampson, Trans. Ent. Soc. Lond., p. 201 (1897). Neu-Hebriden
- neuralis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 4 (♂) (1907). — Rebel, 2. Beih. Hamb. Wiss. Anstalt, 32, p. 135 (1915). Samoa
- perlucidalis** Hampson, Trans. Ent. Soc. Lond., p. 201 (♀) (1897). Marquesas Ins.
- modestalis** Lederer, vide sub b) Orientalische Region, p. 153.

Species incertae sedis.

- chaleophaea** Meyrick, Exot. Microlep., 4, p. 243 (♂) (1932). Fidji Ins. (Typus: Lolotu)
- cirrhophanes** Meyrick, Exot. Microlep., 4, p. 242 (♂, ♀) (1932). Fidji Ins. (Typus: Lautoka)
- craterotoxa** Meyrick, Exot. Microlep., 4, p. 241 (♀) (1932). Fidji Ins. (Typus: Lautoka)
- fuliginosa** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 124 (♀) (1915). Holl. Neu-Guinea

- grisealis** W. Rothschild, Rep. B. O. U. Exped., 2, Holl.
nr. 15, p. 124 (♂) (1915). Neu-Guinea
- haplomorpha** Meyrick, Exot. Microlep., 4, p. 242 Fidji Ins.
(♂) (1932). (Typus:
Lautoka)
- leptographa** Meyrick, Exot. Microlep., 4, p. 243 (♂) Fidji Ins.
(1932). (Typus:
Lautoka)
- miantodes** Meyrick, Exot. Microlep., 4, p. 242 (♀) Fidji Ins.
(1932). (Typus:
Lautoka)
- parbattoides** Tams, Ins. Samoa, pars 3 (fasc. 4), Samoa
p. 266, t. 6, f. 26 (♂, ♀) (1935).
- porphyroclista** Meyrick, Exot. Microlep., 4, p. 499 Fidji Ins.
(♂) (1934). (Typus:
Vunidawa)
- postbicoloralis** W. Rothschild, Rep. B. O. U. Exp., 2, Holl.
nr. 15, p. 125 (♀) (1915). Neu-Guinea
- stigmophanes** Meyrick, Exot. Microlep., 4, p. 242 Fidji Ins.
(♂, ♀) (1932). (Typus:
Lautoka)
- tormentifera** Meyrick, Trans. Ent. Soc. Lond., 77, Marquesas
p. 161 (♂, ♀) (1929). Ins.
- trispila** Meyrick, Trans. Ent. Soc. Lond., p. 232 Fidji Ins.
(♂) (1886) (*Pleonectusa*). — Hampson,
Trans. Ent. Soc. London, p. 202 (1897).

e) Neotropische Region.

- agraphalis** Guenée, Delt. et Pyral., p. 244 (1854) Brasilien
(*Stenia*). — Hampson, Trans. Ent. Soc.
Lond., p. 200 (1897).
- hodesalis** Walker, Cat. Lep. Het. Brit. Mus., 19, Brasilien
p. 960 (♂) (1859) (*Hydrocampa?*). — Hamp- (Typus:
son, Trans. Ent. Soc. Lond., p. 202 (1897) Pará)
(*B. (Erilita)*).
- desumptalis** Walker, Cat. Lep. Het. Brit. Mus., 34, Brasilien
p. 1532 (1865) (*Hydrocampa?*) (Ega). — (Typus:
Hampson, Trans. Ent. Soc. Lond., p. 202 Ega)
(1897).
- purpurascens** Hampson, Ann. Mag. Nat. Hist., (s. Jamaica
7) 19, p. 5 (♀) (1907).

Species incertae sedis.

- antenoralis** Walker, Cat. Lep. Het. Brit. Mus., 19, Brasilien
p. 893 (♂) (1859) (*Pyralis*). — Hampson,
Trans. Ent. Soc. London, p. 202 (1897).
- externalis** Walker, Trans. Ent. Soc. Lond.,
(s. 3) 1, p. 119 (1862) (*Pyralis*).

- finbaralis** Schaus, Proc. Ent. Soc. Wash., 26, p. 122 (♀) (1924). Mexiko
(Typus:
Cordoba)
- aaronalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 122 (♂) (1924). Guatemala
(Typus:
Cayuga)
- hemmingalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 123 (♀) (1924). Jamaica
- melltnialis** Druce, Biol. Centr.-Amer. Het., 2, p. 559, t. 101, f. 12 (♂) (1899) (*Epichronistis*). Guatemala
(Typus:
Dueñas)
79

52. Genus **Synfrifa** Dognin

Dognin, Ann. Soc. Ent. Belg., 49, p. 66 (1905).

- umbralis** Dognin, Ann. Soc. Ent. Belg., 49, p. 66 (♂) (1905). Ecuador
(Typus: ♂
Zamora)
1

53. Genus **Nicaria** Snellen

Snellen, Tijds. v. Ent., 23, p. 229 (1880).

- latisquamalis** Snellen, Tijds. v. Ent., 23, p. 230 (♂) (1880); 27, p. 39, t. 3, f. 10 a, b (1884). Celebes
(Typus:
Bonthain)
1

54. Genus **Coptobasis** Lederer

Lederer, Wien. Ent. Monatschr., 7, p. 429 (1863). — Hampson, Fauna Brit. India Moths, 4, p. 232 (1896); Trans. Ent. Soc. Lond., p. 133, 202 (1897). — Pagenstecher, Zoologica, H. 29, p. 179 (1900). — E. Hering, Ent. Zeit. Stettin, 62, p. 295 (1901).

(Typus generis: *sulcialis* Walker)

- mesopsectralis** Hampson, Trans. Ent. Soc. Lond., p. 203 (♂) (1897) (Amboina). — E. Hering, Ent. Zeit. Stettin, 62, p. 296 (1901). Pulo Laut,
Sumatra,
S.-Borneo,
S.-Celebes,
Natuna Ins.,
Amboina,
Fergusson
Ins., Holl.
u. Brit.
Neu-Guinea,
St. Aignan
- sulcialis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 684 (♂) (1859) (*Botys*) (Sarawak). — Lederer, Wien. Ent. Monatschr., 7, p. 429 (1863). — Snellen, Tijds. v. Ent., 27, p. 42 (1884). — Pagenstecher, Jahrb. Nassau. Ver. Natk., 41, p. 113 (1888). — Snellen, Trans. Ent. Soc. Lond., p. 625 (1890). — Hampson, Fauna Brit. India Moths, 4, p. 232, f. 136 (♂) (1896); Trans. Ent. Soc. Lond., p. 203 (1897). — Pagenstecher, Zoologica, H. 29, Assam,
Sikkim,
Borneo,
Celebes,
Sumatra,
Neu-Guinea,
Amboina,
Bismarck-
Archipel

- p. 179 (1900). — E. Hering, Ent. Zeit. Stettin, 62, p. 296 (1901).
megapteralis Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1407 (♀) (1865) (part.) (*Botys*) (Ceram & Darjeeling).
monocromalis Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1419 (1865) (Sula). — Hampson, Celebes, Sula Ins. Trans. Ent. Soc. Lond., p. 203 (1897) (*monochromalis*). — E. Hering, Ent. Zeit. Stettin, 62, p. 296 (1901).
lophocera Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 5 (♂) (1907). S.-Celebes
dentalis A. Pagenstecher, Zoologica, H. 29, p. 179 Bismarck-Archipel (1900). Neu-lauenburg)
5

55. Genus **Temnobasis** Gaede

Gaede, Ent. Zeit. Stettin, 77, p. 131 (1916).

- simialis* Gaede, Ent. Zeit. Stettin, 77, p. 131 (♂, ♀) Kamerun (1916). (Typus: ♂ Buea)
1

56. Genus **Dialthraustodes** Hampson

Hampson, Fauna Brit. India Moths, 4, p. 233 (1896); Trans. Ent. Soc. Lond., p. 133, 203 (1897).
(Typus generis: *leucotrigona* Hampson)

- leucotrigona* Hampson, Fauna Brit. India Moths, 4, Nilgiris, p. 233, f. 138 (♂) (1896); Trans. Ent. Soc. Seychellen Lond., p. 203, fig. (♂) (1897).
hemilencalis Hampson, Trans. Ent. Soc. Lond., p. Queensland 204 (♂) (1897).
fulvofusa Hampson in Leech, Trans. Ent. Soc. Lond., Assam p. 442 (♂) (1901). — Leech, Trans. Ent. (Khasis), Soc. Lond., p. 442 (1901). — Hampson, Zentralchina Journ. Bombay Nat. Hist. Soc., 15, p. 211 (Typus: (♂) (1903). Ichang)
similis Hampson, Journ. Bombay Nat. Hist. Soc., Assam 15, p. 211 (♀) (1903). (Khasis)
4

57. Genus **Coelorhyncidia** Hampson

Hampson, Fauna Brit. India Moths, 4, p. 232 (1896); Trans. Ent. Soc. Lond., p. 133, 204 (1897).
(Typus generis: *ovulalis* Hampson)

- ovulalis* Hampson, Fauna Brit. India Moths, 4, p. N.W.- 232, f. 317 (♂) (1896); Trans. Ent. Soc. Himalaya, Lond., p. 204, fig. (♂) (1897). Simla, Dharmasala

- elathealis** Walker, Cat. Lep. Het. Brit. Mus., 18, Brasilien
p. 615 (♂) (1859) (*Botys*) (Amazonas). —
Hampson, Trans. Ent. Soc. Lond., p. 204
(1897).
- flammealis** Hampson, Ann. Mag. Nat. Hist., (s. 8) Ecuador
20, p. 212 (♂) (1917).
- pandaralis** Walker, Cat. Lep. Het. Brit. Mus., 18, Brasilien
p. 618 (♀) (1859) (*Botys*) (Amazonas). —
Hampson, Trans. Ent. Soc. Lond., p. 204
(1897).
- trifidalis** Hampson, Trans. Ent. Soc. Lond., p. 205 Amboina
(1897).
- purpurea** Hampson, Ann. Mag. Nat. Hist., (s. 7) Amboina
19, p. 6 (♂) (1907).
- cuprescens** Hampson, Ann. Mag. Nat. Hist., (s. 8) Holl.
20, p. 213 (♂) (1917). Neu-Guinea
- nitidalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, Holl. u.
p. 6 (♂, ♀) (1917). Brit.
8 : Neu-Guinea

58. Genus **Geshna** Dyar

Dyar, Journ. N. York Ent. Soc., 14, p. 95 (1906). — Forbes, Lep.
N. York, p. 581, f. 350 (1923).

(Typus generis: *G. cannalis* Quaint.)

- cannalis** Quaintance, Bull. Florida Exp. Stat., nr. 45, Florida
p. 68 (1898) (*Hydrocampa*). — Dyar, Proc.
Ent. Soc. Wash., 4, p. 463 (1901) (*Nym-*
phula); Journ. N. York Ent. Soc., 14, p. 96
(1906).
Biologie: Dyar, Journ. N. York Ent. Soc.,
14, p. 96 (1906).
- primordialis** Dyar, Journ. N. York Ent. Soc., 14, Vereinigte
p. 97 (1906). — Forbes, Lep. N. York, Staaten
p. 581 (1923). (Illinois,
Penn-
sylvanien,
N. Jersey,
Maine,
N. York,
Texas,
N. Carolina),
Kanada
(Quebec,
Ontario)

2

59. Genus **Diathrausta** Lederer

Lederer, Wien. Ent. Monatschr., 7, p. 438 (1863). — Meyrick, Trans.
Ent. Soc. Lond., p. 218 (1887). — Hampson, Fauna Brit. India
Moths, 4, p. 233 (1896); Trans. Ent. Soc. Lond., p. 133, 205 (1897).
— Dyar, Journ. N. York Ent. Soc., 14, p. 97 (1906). — Forbes, Lep.
N. York, p. 580, f. 351 (1923).

(Typus generis: *profundalis* Led.)

Syngamia Wileman, Trans. Ent. Soc. Lond., p. 377 (1911).

a) Paläarktische Region.

- brevifascialis** Wileman, Trans. Ent. Soc. Lond., p. 377 (1911) (*Syngamia*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 163, t. 5, f. 12 (1928); Ins. Matsum., 3, p. 131, t. 5, f. 20 (1929). Japan, Formosa
- picata** Leech (nec Butler, 1889), Trans. Ent. Soc. Lond., p. 442 (1901).

b) Orientalische Region.

- profundalis** Lederer, Wien. Ent. Monatschr., 7, p. 438, t. 17, f. 7 (1863) (Amboina). — Hampson, Fauna Brit. India Moths, 4, p. 233, f. 139 (♂) (1896); Trans. Ent. Soc. Lond., p. 205, fig. (♂) (1897). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 23, p. 163, t. 6, f. 21 (1928). Indien, Sikkim, Südchina, Tonkin, Formosa, Ceylon, Sumbawa, Amboina, Buru
- var. **conicalis** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 143 (1896). — Hampson, Trans. Ent. Soc. London, p. 205 (1897). Assam (Khasis)
- var. **oblitura** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 143 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 205 (1897). Assam (Khasis)
- picata** Butler, Ill. Typ. Sp. Lep. Het. Brit. Mus., 7, p. 94, t. 134 (1889). — Hampson, Fauna Brit. India Moths, 4, p. 234 (1896); Trans. Ent. Soc. Lond., p. 205 (1897). N.W.-Himalaya (Dharmasala), West-, Süd- u. Ostchina
- plumbealis** Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 174 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 206 (1897); Journ. Bombay Nat. Hist. Soc., 12, p. 704 (1899). Assam, West-, Süd- u. Ostchina
- var. **talboti** Caradja, Mém. Acad. Roum., 10, p. 47 (1927) (*Massepha*); ibid., 14, p. 208 (1931) (*Diathrausta*).

c) Aethiopische Region.

- fulviceps** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 214 (1917). Nyassaland

d) Nearktische und Neotropische Region.

- griseifusa** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 213 (♂) (1917). Brasilien (S. Catharina)
- nerinalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 928 (1859) (♀) (*Desmia*) (Brasil.). — Hampson, Trans. Ent. Soc. Lond., p. 205 (1897). Kanada (Ontario), Vereinigte Staaten, Mexiko, Panama, Brasilien, Ecuador, Trinidad

- fa. *nerinalis* Walker, l. c. — Dyar, Insec. Insc. Menstr., 1, p. 102 (1913) (*D. nerinalis* „Form“). Brasilien (Typus: Castro, Paraná)
- fa. *reconditalis* Walker, Cat. Lep. Het. Brit. Mus., 19, p. 943 (♂) (1859) (*Hymenia* spec.) (N. York). — Hampson, Trans. Ent. Soc. Lond., p. 205 (1897) (*Diathrausta*). — Dyar, Journ. N. York Ent. Soc., 14, p. 97 (1906). (spec. dist.); Insec. Insc. Menstr., 1, p. 100 (1913) (*nerinalis* forma). — Forbes, Lep. N. York, p. 580 (1923) (spec. dist.). Vereinigte Staaten (Massachusetts, N. York, Colorado, Neu-Mexiko, Connecticut, N. Jersey, Pennsylvanien, Arizona)
- minutalis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1297 (♂) (1865) (*Aediodes*) (Nordamerika).
- octomaculalis* Fernald, Ent. Amer., 3, p. 127 (1887) (*Diathrausta*).
- pisusalis* Fernald (nec Walker, 1859), Smith's List Lep. Bor. Amer., nr. 3985 (1891).
- fa. *harlequinialis* Dyar, Insec. Insc. Menstr., 1, p. 100 (1913) (*D. nerinalis* „form“). Colorado, Neu-Mexiko, Arizona, Mexiko (Typus: Tehuacan (Mexiko))
- montana* Haimbach, Ent. News, 26, p. 323, t. 12, f. 6 (1915) (Colorado). — Mc Dounough, Ent. News, 26, p. 471 (1915).
- fa. *cubanalisis* Dyar, Insec. Insc. Menstr., 1, p. 101 (1913). Cuba (Typus: Baracoa)
- fa. *minutalis* Druce, Biol. Centr.-Amer. Het., 2, p. 560, t. 101, f. 18 (1899) (*Diasemia?*). — Dyar, Insec. Insc. Menstr., 1, p. 101 (1913) (*D. nerinalis* „Form“). Mexiko (Typus: Orizaoa)
- fa. *cacalis* Dyar, Insec. Insc. Menstr., 1, p. 101 (1913) (*D. nerinalis* „Form“). Mexiko, Colombia oder Ecuador
- fa. *angustella* Dyar, Insec. Insc. Menstr., 1, p. 101 (1913) (*D. nerinalis* „Form“). Panama (Typus: Trinidad River)
- delicata* Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 218 (1896). — Hampson, Trans. Ent. Soc. Lond., p. 206 (1897). Cayenne
- cymialis* Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 6 (♂) (1907). Brasilien (Paraná)
- daeckcalis* Haimbach, Ent. News, 26, p. 322, t. 12, f. 5 (1915). — Forbes, Lep. N. York, p. 581 (1923). N. Jersey

e) Australische Region.

leucographa Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 213 (♀) (1917). D'Entrecasteaux Ins. (Goodenough Ins.)

lypera Tams, Ins. Samoa pars 3 (fasc. 4), p. 267, Samoa t. 17, f. 4 (♀) (1935).
12

60. Genus **Deuterophysa** Warren

Warren, Trans. Ent. Soc. Lond., p. 272 (1889). — Hampson, Trans. Ent. Soc. Lond., p. 133, 206 (1897).
(Typus generis: *costimaculalis* Warren)

costimaculalis Warren, Trans. Ent. Soc. Lond., p. 273 (1889). — Hampson, Trans. Ent. Soc. Lond., p. 206, fig. (♂) (1897).

micralis Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 7 (♂) (1907). Jamaica

grisealis Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 214 (♂) (1917). Br. Zentralafrika

furvitermen Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 215 (1917). Br. Zentralafrika

obregonalis Schaus, Proc. Ent. Soc. Wash., 26, p. 123 (♂) (1924). Mexiko
5

61. Genus **Lasiogyia** Hampson

Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 7 (1907).

xanthozonata Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 7 (♂) (1907). Fergusson Ins.
1

62. Genus **Microphysefica** Hampson

Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 215 (1917).

peperita Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 215 (♂) (1917). Mexiko
1

63. Genus **Epistenia** Chrétien

Chrétien, Ann. Soc. Ent. Fr., 79, p. 524 (1910).

Stenia (part.) auctorum.

bruguieralis Duponchel, Hist. Nat. Lep. Fr., 8 (2), p. 320, t. 232, f. 3 (1831) (*Stenia*). — Südeuropa (Spanien, Herrich-Schaeffer, Syst. Bearb. Schm. Eur., Süd-

4, p. 14, f. 84 (1848) (*Stenia*). — Zeller, Isis, p. 22, 569 (1847) (*Stenia*). — Guenée, Delt. et Pyral., p. 245 (1854) (*Stenia*). — Berce, Faune Ent. Fr. Lep., 6, p. 105, t. 3, f. 8 (1878) (*Stenia*). — Hampson, Trans. Ent. Soc. Lond., p. 207 (1897) (*Stenia*). — Rebel in Spuler, Schm. Eur., 2, p. 222 (1906) (*Stenia*). — Chrétien, Ann. Soc. Ent. Fr., 79, p. 524 (1910) (*Epistenia*).

frankreich,
Italien,
Korsika,
Kreta,
Cypern,
Balkan,
Dalmatien),
Kleinasien,
Amasien,
Libanon,
Syrien,
Palästina,
Mauretanien,
Westasien,
Holl.
Neu-Guinea

ab. *infernalis* Zerny, Deutsche Ent. Zeitschr. Iris, 41, p. 137, t. 2, f. 13 (1927).

tenebrosa W. Rothschild, Nov. Zool., 35, p. 233 (1929) (*bruguieralis* subsp.). — Zerny, Mem. Soc. Sci. Nat. Maroc, 42, p. 122 (1935).

subsp. *bahrutalis* Amsel, Veröff. Deutsch. Kolon. & Übersee-Mus. Bremen, 1 (2), p. 209, t. 11, f. 4 (♂) (1933).

Süd-
Andalusien,
Marokko
(Gr. Atlas)

Totes Meer

daralis Chrétien, Ann. Soc. Ent. Fr., 79, p. 523 (*Stenia*), p. 524 (*Epistenia*) (1911).

Tunis,
S.-Algier
(Biskra),
Marokko
(Gr. Atlas),
Jerusalem

2

64. Genus **Stenia** Duponchel

Duponchel, Cat. Méth. Lep. Eur., p. 201 (1845). — Guenée, Delt. et Pyral., p. 241 (1854). — Stainton, Man. Brit. Butt. Moths, 2, p. 143 (1859). — Walker, Cat. Lep. Het. Brit. Mus., 17, p. 413 (1859). — Lederer, Wien. Ent. Monatschr., 7, p. 421 (1863). — Heinemann, Kleinschm., 1, H. 2, p. 100 (1865). — Wallengren, Oefvers. K.-Vet.-Akad. Förh., 28 (8), p. 1004 (1871). — Meyrick, Trans. Ent. Soc. Lond., p. 462 (1890); Handb. Brit. Lep., p. 404 (1895). — Hampson, Fauna Brit. India Moths, 4, p. 234 (1896); Trans. Ent. Soc. Lond., p. 206 (1897). — Barrett, Lep. Brit. Isl., 9, p. 297 (1904). — Rebel in Spuler, Schm. Eur., 2, p. 222 (1906). — Dyar, Journ. N. York Ent. Soc., 14, p. 99 (1906). — Wahlgren, Svensk Insektfauna, 10 (2), p. 48 (1915). — Meyrick, Revis. Handb. Brit. Lep., p. 421 (1928). — M. Hering in Brohmer & Ehrmann & Ulmer, Tierwelt Mitteleur. (Ergbd. 1), p. 282 (1932). — Eckstein, Schm. Deutschl., 5, p. 62 (1933).

Arnica Guenée, Delt. et Pyral., p. 240 (1854) (*Typus: nervosalis* Guenée).

amaurophanes Lederer, Wien. Ent. Monatschr., 7, p. 422 (1863) (*Typus: stigmosalis* Herr.-Schaeff.).

Sozoa Walker, Cat. Lep. Het. Brit. Lep., 34, p. 1373 (1865) (*Typus: costalis* Walk. = *ianthealis* Walk.).

Steniodes Snellen, Tijds. v. Ent., 18, p. 244 (1875) (Typus: *gelliasalis* Walk.); 42, p. 77 (1899). — E. Hering, Ent. Zeit. Stettin, 62, p. 317 (1901). — Dyar, Journ. N. York Ent. Soc., 14, p. 99 (1906).

Symmoracma Meyrick, Trans. Ent. Soc. Lond., p. 469 (1894) (Typus: *spodinopa* Meyrick).

(Typus generis: *punctalis* (Schiff. et Den.)).

a) Paläarktische Region.

- intervacatalis** Christoph, Hor. Soc. Ent. Ross., 12, p. 221, 276 (♂, ♀), t. 7, f. 47 (1877); id. in Romanoff, Mém., 3, p. 42, t. 2, f. 11 (1887).
Armenien,
Palästina,
Persien
(Typus:
Schabrud)
- punctalis** [Schiffermüller & Denis], Syst. Verz. Schm. Wien, p. 123 (1776) (*Pyralis*). — Hübner, Samml. Eur. Schm. (*Pyr.*), t. 21, f. 140 (1796) (*Pyralis*). — Fabricius, Ent. Syst., 3 (2), p. 222, f. 794 (*Phalacna*). — Treitschke, Schm. Eur., 7, p. 130 (1829); 10 (3), p. 28 (1835) (*Nymphula*). — Herrich-Schäffer, Syst. Bearb. Schm. Eur., 4, p. 14, t. 4, f. 24 (1848). — Guenée, Delt. et Pyral., p. 248 (1854). — Stainton, Man. Brit. Butt. Moths, 2, p. 143 (1859). — Lederer, Wien. Ent. Monatschr., 7, p. 422 (1863). — Heinemann, Kleinschm., 1, H. 2, p. 101 (1865). — Wallengren, Oefvers. K. Vet.-Akad. Förh., 28 (8), p. 1005 (1871). — Berce, Faune Ent. Fr. Lep., 6, p. 104 (1878). — Meyrick, Handb. Brit. Lep., p. 405 (1895). — Hampson, Fauna Brit. India Moths, 4, p. 234 (1896); Trans. Ent. Soc. Lond., p. 207 (1897). — Barrett, Lep. Brit. Isl., 9, p. 297, t. 409, f. 1, 1, a, b (1904). — Rebel in Spuler, Schm. Eur., 2, p. 222, t. 81, f. 13 (1906). — Wahlgren, Svensk Insektafauna, 10 (2), p. 48, t. 2, f. 54 (1915). — Meyrick, Revis. Handb. Brit. Lep., p. 421 (1928). — M. Hering in Tierwelt Mitteleuropas (Ergbd. 1), p. 282 (1932). — Eckstein, Schm. Deutschl., 5, p. 62 (1933).
- aetnealis* Duponchel, Hist. Nat. Lep., 8 (2), p. 199, t. 223, f. 4 (1831) (Asopia).
- longipetalis* Curtis, Brit. Ent., 7, p. 312 (1830) (*Margaritia*). — Stephens, Ill. Brit. Ent. Haust., 4, p. 55 (1834) (*Margaritia*). — Wood, Index Entom., p. 126, t. 28, f. 825 (1839) (*Margaritia*).
- lorquinalis* Guenée, Delt. et Pyral., p. 138 (1854) (*Cledeobia*).
- fa. mutantalis* Caradja, Ent. Rundschau, 45, p. 26 (1928); Peters, ibid., 45, p. 30, nota (1928).
Sicilien
- var. *flavipunctalis* Hampson, Trans. Ent. Soc. Lond., p. 385, t. 3, f. 8 (1900) (*Stenia* spec.). — Żerny, Deutsche Ent. Zeitschr. Iris, 41, p. 137, t. 2, f. 12 (1927) (*punctalis* var.).
Aragonien,
Andalusien,
Ost-Pyrenäen,
Portugal

- subsp. *zernyi* Bryk (n. nov.).
gigantalis Zerny (nec Druce, 1896), Mem. Soc. (Sci. Nat. Maroc, 42, p. 123 (1935) (*S. punctalis* subsp.).
 Biologie: Hellins, Ent. Men. Mag., 21, p. 121 (1884). — Buckler, Larvae Brit. Lep., 9, p. 70, t. 150, f. 7 (1901).
concoloralis Oberthür, Etud. Ent., 1, p. 68, t. 2, f. 6 (1876); 3, p. 46 (1878) (*Scopula*). — Standinger, Deutsche Ent. Zeitschr. Iris, 5, p. 294 (1892). Südsanien, Mauretanien
afrella Turati, Atti Soc. Ital. Sci. Nat., 63, p. 145, t. 5, f. 37 (♂) (1924). Cyrenaica
ferrealis Hampson, Trans. Ent. Soc. Lond., p. 385 (1900). S.-Mesopotamien, Syrien, Armenien
fuscocillialis Ragonot, Ent. Mon. Mag., 17, p. 230 (1881). — Mitterberger, Zeitschr. Wiss. Insektenbiol., 11, p. 175 (1915). S.O.-Frankreich, Aragonien, Portugal, Süditalien (Typus: Silves)
stigmosalis Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 14, (Pyr.), t. 4, f. 20 (1848). — Guenée, Delt. et Pyral., p. 259 (1854). — Hampson, Trans. Ent. Soc. Lond., p. 207 (1897). — Rebel in Spuler, Schm. Eur., 2, p. 222 (1906). Südost-Europa (Kroatien, Bulgarien, Rumänien, Rumelien, Thrazien, Balkan, Süd-Rußland), Frankreich, Kleinasien, Amasien, Libanon, Armenien, Westasien
amoenialis Christoph, Bull. Soc. Imp. Nat. Moscou, 56 (1), p. 30 (1881) (*Amaurophanes*). — Hampson, Trans. Ent. Soc. Lond., p. 208 (1897). Ussuri, Süd- und Ostchina
nervosalis Lucas, Explor. Alg., 3, p. 399, t. 4, f. 6 (1849). — Guenée, Delt. et Pyral., p. 241 (1854). — Hampson, Trans. Ent. Soc. Lond., p. 208 (1897). Algier, Süd-Frankreich, Sicilien, Cyrenaica
 fa. *pygmaealis* Amsel, Veröff. Deutsch. Kolon. u. Uebersee-Mus., 1 (2), p. 255 (1935) (gen. aest.). Tel-Aviv
charonialis Walker, Cat. Lep. Het. Brit. Mus., 17, p. 372 (♂) (1859) (*Asopia*) (Shanghai). — Hampson, Trans. Ent. Soc. Lond., p. 221 (1897) (*Mabra*). — Leech, Trans. Ent. Soc. Lond., p. 445 (1901) (*Mabra*). — Caradja, Mem. Acad. Rom. Sect. Stiint., (s. 3) 4, Amur, West- u. Mittelchina, Japan, Korea, Formosa

- mem., 8, p. 407 (1927) (*Mabra*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 164 (1928); Ins. Matsum., 3, p. 131, t. 5, f. 22 (1929).
dissipatalis Christoph, Bull. Soc. Imp. Nat. Moscou, 56 (1), p. 28 (1881) (*Stenia*?).
metasialis Rebel, Ann. Naturh. Hofmus. Wien, 30, p. 148, t. 4, f. 6 (1916). Kreta
alluminialis Turati, Atti Soc. Ital. Sci. Nat., 73, p. 190 (1934). Cyrenaica

b) Orientalische Region.

- costipunctalis* Snellen, Tijds. v. Ent., 42, p. 77 (♂, ♀), t. 4, f. 4 (♂), 5 (antenna ♂) (1899) (Typus: *Steniodes*). — E. Hering, Ent. Zeit. Stettin, 62, p. 317, 318 (1901). Java (Tegal)
- punctalis* (Schiff. & Denis), vide sub a) Paläarktische Region p. 164.
- nigriscripta* Swinhoe, Ann. Mag. Nat. Hist., (s. 6) 16, p. 302 (1895) (*Mabra*). — Hampson, Trans. Ent. Soc. London, p. 221 n. 4 (1898). Indien, Assam, Formosa
 — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 164 (1928).
- charonialis* Walker, vide sub a) Paläarktische Region p. 165.
- rubriceps* Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 212 (♂) (1903). Dharmasala, Tenimber
- grisealis* Hampson, Bull. Liverpool Mus., 2, p. 39 (1899); Journ. Bombay Nat. Hist. Soc., 15, p. 212 (1903); in Forbes, Nat. Hist. Sokotra, p. 335, t. 20, f. 4, 23 (1903). Indien, Sokotra
- minoralis* Snellen, Tijds. v. Ent., 23, p. 222 (♀) (1880) (*Auxomitia*); 26, p. 137, t. 8 (♀), 8a (caput) (1883) (*Auxomitia*); 43 (1900) p. 270, 271 (1901) (*Auxomitia*). — Pagenstecher, Zoologica H. 29, p. 191 (1901). — Malakka, Singapore, Südchina, Formosa, Ceylon,
profanalis Hampson (nec Walk.), Ill. Typ. Sp. Lep. Het. Brit. Mus., 9, p. 175, t. 174, f. 22 (1893). Borneo, Java, Sumbawa,
spodinopa Meyrick, Trans. Ent. Soc. Lond., p. 469 (1894) (*Symmoracma*). — Hampson, Fauna Brit. India Moths, 4, p. 234, f. 140 (♂) (1896); Trans. Ent. Soc. Lond., p. 207, fig. (♂) (1897) (*S. (Steniodes)*). — Buru, Celebes, Bismarck-Archipel, Philippinen (Typus: Takalar, Bonthain)
 — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 164 (1928).
spodinopella Strand, Ent. Mitt., 8, p. 130 (1919) (*spodinopa* forma). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 26 (1928).
- hieralis* Swinhoe, Trans. Ent. Soc. Lond., p. 156 (♂) (1904). Borneo (Typus: Sarawak)
- paediusalis* Walker, vide sub c) Aethiopische und Madagassische Region p. 167.

c) Aethiopische und Madagassische Region.

- baltealis** Mabille, C. R. Ent. Soc. Belg., 25, p. LXII (1881). Madagaskar
- desertalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 8 (♀) (1907). Kapkolonie
- grisealis** Hampson, vide sub b) Orientalische Region p. 166.
- modestalis** Saalmüller, Ber. Senckenb. Ges., p. 299 (1879); Lep. Madag. (2), p. 503 (1891). — Hampson, Trans. Ent. Soc. Lond., p. 208 (1897). Madagaskar
- paediusalis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 788 (1859) (*Scopula*) (Congo). — Hampson, Fauna Brit. India Moths, 4, p. 235 (1896). Westafrika, Port. O.-Afrika, Aden, N.O.-Indien
- poediusalis** Hampson, Trans. Ent. Soc. Lond., p. 298 (1897).
- phaeospilalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 12 (♂) (1907). Brit. O.-Afrika
- pulchellalis** Mabille, Ann. Soc. Ent. Fr., s. 5, 9, p. 337 (1879). — Hampson, Trans. Ent. Soc., p. 208 (1897). Madagaskar
- triflexalis** Gaede, Ent. Zeit. Stettin, 77, p. 131 (♀) (1916). Kamerun
- uniflexalis** Mabille, Ann. Soc. Ent. Fr., (s. 5) 9, p. 336 (1879). — Gaede, Mitt. Zool. Mus. Berlin, 8, p. 396 (1917). — Hampson, Trans. Ent. Soc. Lond., p. 208 (1897). Madagaskar

d) Australische Region.

- punctalis** [Schiff. & Den.], vide sub a) Paläarktische Region p. 164.
- tenebrosalis** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 126 (♂) (1915). Holl. Neu-Guinea

e) Neotropische Region.

- gelliasalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 988 (♀) (1859) (*Botys*) (Rio Janeiro). — Hampson, Trans. Ent. Soc. Lond., p. 207 (1897) (*St. (Steniodes)*). — Dyar, Journ. N. York Ent. Soc., 14, p. 99 (1906) (*Steniodes*). — Westindien, Jamaica, Gronada, Brasilien (Rio de Janeiro), Florida, Texas
- lutealis** Snellen, Tijds. v. Ent., 18, p. 245, t. 13 (♂, ♀), f. 17, 18 (1875) (*Steniodes*). (Typus: Ubaque)
- nennuisalis** Sehaus, Proc. Ent. Soc. Wash., 26, p. 124 (♀) (1924). Peru (Typus: Trujillo)
- retractalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 216 (♀) (1917). Br. Guyana

- declivalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 285 (1914). Costa Rica,
Panama,
Venezuela
- var. *indianalis* Dyar, Proc. U. S. Nat. Mus., 47, p. 286 (1914). Westindien,
Cuba,
Jamaica,
Portorico,
Grenada
- dominicalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 124 (♀) (1924). Westindien
(Typus:
Dominica)
- ceddalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 125 (♂) (1924). Cuba
(Typus:
Cayamas)
- midalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 125 (♀) (1924). Ecuador
(Typus:
Queveds)
- marialis** Schaus, Proc. Ent. Soc. Wash., 26, p. 125 (♀) (1924). Guatemala
(Typus:
Volean de
S. Maria)
- styxalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 126 (♂) (1924). Fr. Guyana
(Typus:
St. Jean)
- glaucinalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 266 (♀) (1917). Bolivien
- fusalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 265 (1917). Kolombien
- acuminalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 286 (1914). Pauama,
Costa Rica
- obscura** Warren, Trans. Ent. Soc. Lond., p. 291 (♂) (1889) (*Sozoa*). — Hampson, Trans. Ent. Soc. Lond., p. 207 (1897) (*St. (Steniodes)*). Brasilien
(Typus:
Pupunha)
- ianthealis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 523 (♂) (1859) (*Margaronia*) (Bogotá). — Hampson, Trans. Ent. Soc. Lond., p. 207 (1897). Venezuela,
Kolombien
- costalis* Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1373 (♂) (1865) (*Sozoa*). (Bogotá,
Venezuela)
- adustalis* Felder et Rogenhofer, Reise Novara (Het.), p. 9, t. 135, f. 32 (1874). (Ercta?) (Bogotá)
- costalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 8 (1907). Guatemala,
Costa Rica,
Kolombien,
Panama
- colubralis* Druce, Biol. Centr.-Amer. Het., 2, p. 268 (1895), part. (*Blepharomastix*), nec Guen. (1854).
- cronanalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 126 (♂) (1924). Peru
(Typus:
El Cam-
pamiento)

- bademusalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 126 (♂) (1924). Peru
(Typus:
Col Perene)
- carthaghalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 127 (♂) (1924). Peru
(Typus:
Col Perene)
- primolalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 127 (♀) (1924). Surinam
(Typus:
Geldersland)
- guianalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 128 (♀) (1924). Fr. Guyana
(Typus:
St. Jean)
- turiafalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 128 (♂) (1924). Peru?
- caclamalalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 128 (♂) (1924). Kolombien
- mononalis** Dyar, Proc. U. S. Nat. Mus., 54, p. 369 (♀) (1918). Mexiko
- cayugalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 128 (♂) (1924). Guatemala
(Typus:
Cayuga)
- herrereralis** Schaus, Proc. Ent. Soc. Wash., 26, p. 129 (♀) (1924). Guatemala
(Typus:
Volcan de
S. Maria)
- caulealis** Schaus, Proc. Ent. Soc. Wash., 26, p. 129 (♂) (1924). Guatemala
(Typus:
Volcan de
S. Maria)
- beuvealis** Schaus, Proc. Ent. Soc. Wash., 26, p. 129 (♀) (1924). Guyana
(Typus:
St. Laurent)
- branealis** Schaus, Proc. Ent. Soc. Wash., 26, p. 130 (♀) (1924). Jamaica
(Typus:
Kingston)
- semifuscalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 9 (1907). Ecuador
- interruptalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 9 (♀) (1907). Ecuador
- fusculunalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 9 (♂) (1907). Guatemala,
Costa Rica,
Panama
colubralis Druce (nec Guen. (1854)), Biol.
Centr.-Amer. Het., 2, p. 268, part. (*Blepharomastix*).
- irroratalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 10 (♂) (1907). Guatemala,
Brasilien
(Rio de
Janeiro)

- sauralis** Guenée, Delt. et Pyral., p. 242 (1854). — Cayenne
Hampson, Trans. Ent. Soc. Lond., p. 207
(1897).
- biannulalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) Brasilien
19, p. 10 (♂) (1907). (Rio de Janeiro)
- aphenice** Hampson, Ann. Mag. Nat. Hist., (s. 7) Mexiko
19, p. 11 (♂) (1907).
phenice Druce (nec Cram.), 1780, Biol.
Centr.-Amer. Het., 2, p. 249 (1895).
- mallaleuca** Hampson, Ann. Mag. Nat. Hist., (s. 7) Brasilien
19, p. 11 (♂) (1907). (Paraná, S. Paulo)
- glgantal** Druce, Biol. Centr.-Amer. Het., 2, p. Guatemala
269, t. 63, f. 5 (♂) (1896) (*Blepharomastix*). — Hampson, Trans. Ent. Soc. Lond.,
p. 208 (1897). (Sinanja in Vera Paz)

Species incertae sedis.

- ambialis** Schaus, Proc. Ent. Soc. Wash., 26, p. 123 Mexiko
(♂) (1924). (Typus: Jalapa, Guatemala)
- annulalis** Dognin, Ann. Soc. Ent. Belg., 50, p. S.O.-Peru
184 (♀) (1906). (Typus: Carabaya)
- astenialis** Dyar, Insec. Insc. Menstr., 10, p. 15 Mexiko
(♂, ♀) (1922). (Typus: Mexico City)
- batrachalis** Guenée, Delt. & Pyral., p. 243, t. 9, Brasilien
f. 8 (1854) (*Nacoleia*).
- benetinctalis** Dyar, Proc. U. S. Nat. Mus., 47, p. Mexiko
397 (1914).
- carillalis** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, Costa Rica
p. 681 (♂) (1912).
- colubralis** Guenée, Delt. et Pyral., p. 242 (1854). Fr. Guayana,
(Cayenne). Panama, Trinidad,
ineffectalis Walker, Trans. Ent. Soc. Brasilien
Lond., (s. 3) 1, p. 126 (1862) (*Botys*);
Cat. Lep. Het. Brit. Mus., 34, p. 1390 (1865)
(Amazonas). (Espirito Santo)
- electralis* Walker, Cat. Lep. Het. Brit.
Mus., 18, p. 600 (♂) (1859) (*Botys*) (Ega).
- melitealis* Walker, Cat. Lep. Het. Brit.
Mus., 18, p. 602 (♂) (1854) (*Botys*) (Brasilien).
- hyperochalis** Dyar, Proc. U. S. Nat. Mus., 47, p. Panama
285 (1914).
- laccrtalis** Guenée, Delt. & Pyral., p. 244 (1854) Brasilien
(*Nacoleia*).

- lubricalis** Dognin, Ann. Soc. Ent. Belg., 49, p. 66 Ecuador
(♂) (1905).
- pacificalis** Schaus, Ann. Mag. Nat. Hist., (s. 8) Costa Rica
9, p. 681 (♂) (1912).
- poasalis** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, Costa Rica
p. 681 (♂) (1912).
- prophaealis** Dognin, Ann. Soc. Ent. Belg., 57, p. Peru
415 (1913).
- samealis** Dyar, Proc. U. S. Nat. Mus., 47, p. 286 Panama,
(♀) (1914). Fr. Guyana,
- saponalis** Guenée, Delt. et Pyral., p. 243 (1854) Fr. Guyana,
(Cayenne). Trinidad,
Panama

79

65. Genus **Eurytorna** Meyrick

Meyrick, Trans. Ent. Soc. Lond., p. 251 (1886). — Hampson, Trans. Ent. Soc. Lond., p. 133, 208 (1897).

- heterodoxa** Meyrick, Trans. Ent. Soc. Lond., p. 252 Fidji Ins.
(♂) (1886). — Hampson, Trans. Ent. Soc.
Lond., p. 208 (1897).

1

66. Genus **Pilelocera** Lederer

Lederer, Wien. Ent. Monatschr., 7, p. 431 (1863). — Hampson, Fauna Brit. India Moths, 4, p. 236 (1896); Trans. Ent. Soc. Lond., p. 133, 209 (1897). — Pagenstecher, Zoologica H. 29, p. 181 (1900). — Dyar, Journ. N. York Ent. Soc., 14, p. 98 (1906).

- Rinecera** Butler, Mem. Nat. Acad. Sci., 2, p. 95 (1884)
(Typus: *mirabilis* Butl. = *signiferalis* Wallgr.).
- Danaga** Moore, Lep. Ceylon, 3, p. 272 (1885) (Typus:
concisalis Moore).
- Ptilaeola** Meyrick, Trans. Ent. Soc. Lond., p. 244 (1886)
(Typus: *ulophanes* Meyrick).
- Erebangela** Meyrick, Trans. Ent. Soc. Lond., p. 245
(1886) (Typus: *melanauges* Meyrick).
- Diploctyla** Meyrick, Trans. Ent. Soc. Lond., p. 246 (1886)
(Typus: *ochrosema* Meyrick). — Pagenstecher, Zoologica
H. 29, p. 182 (1900).
- Strepsimela** Meyrick, Trans. Ent. Soc. Lond., p. 249
(1886) (Typus: *xanthosoma* Meyr.).
- Graphicopoda** Butler, Trans. Ent. Soc. Lond., p. 421
(1886) (Typus: *melanauges* Meyr.).
- Normatholepis** Butler, Trans. Ent. Soc. Lond., p. 424
(1886) (Typus: *erebina* Butl.).
- (Typus generis: *violalis* Lederer.)

a) Paläarktische Region.

- aegimiusalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 929 (1859) (*Desmia*) (Sarawak). — Sarawak). — Hampson, Fauna Brit. India Moths, 4, p. 236, f. 142 (♂) (1896); Trans. Ent. Soc. Lond., p. 212; p. 209, fig. (♂) (1897). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 449 (1900). — Pagenstecher, Zoologica H. 29, p. 182 (1900). — Leech, Trans. Ent. Soc. Lond., p. 443 (1901). — Strand, Ent. Mitt., 8, p. 131 (1919). — Marumo, Journ. Col. Agr. Tokyo Imp. Univ., 8, p. 188 (1923). — W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 185 (1915) (*Almonia*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 166 (1928); Ins. Matsum., 3, p. 132, t. 5, f. 23 (1929).
- collaris** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1293 (♂) (1865) (*Desmia*) (Morty). — Meyrick, Trans. Ent. Soc. Lond., p. 467 (1894) (*Ptilaeola*).
- cincta** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1293 (1865) (*Desmia*) (Neu-Guinea, Mysol).
- mysolalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1294 (♂) (1865) (*Desmia*) (Mysol); 34, p. 1299 (♂) (1865) (*Aediodes*) (Mysol).
- flavomaculalis** Pagenstecher, Jahrb. Nassau. Ver. Naturk., 37, p. 279, t. 7, f. 8 (1884) (*Piletocera*?) (Sep. p. 132).
- flavomaculata** Hampson, Trans. Ent. Soc. Lond., p. 212 (1897) (ex syn.).
- pullatalis** Swinhoe, Proc. Zool. Soc. Lond., p. 420 (♀) (1889) (*Danaga*) (Andamanen). — Caradja, Mem. Acad. Rom. Sect. Stiint., (s. 3) 3, mem. 7, p. 335 (1925) (*aegimiusalis* forma) (Amoy).
- formosalis** Strand, Arch. Naturg., 85 A H. 12, p. 134 (1920) (*Piletocera*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 26 (1928).
- sodalis** Leech, Entom., 22, p. 71, t. 4, f. 6 (1889) (*Desmia*). — Hampson, Trans. Ent. Soc. Lond., p. 213 (1897). — Leech, Trans. Ent. Soc. Lond., p. 442 1(901). — E. Hering, Ent. Zeitschr. Stettin, 64, p. 47 (1903). — Marumo, Journ. Coll. Agr. Tokyo Imp. Univ., 8, p. 188 (1923). — Shibuya, Ins. Matsum., 3, p. 132, t. 5, f. 24 (1929).
- opacalis** Rebel, Bull. Soc. Roy. Ent. Egypte, 19, p. 186 (1926).
- Syrien,
Japan,
Südchina,
Sikkim,
Assam
(Khasis),
Formosa,
Borneo,
Sumatra,
Neu-Guinea,
Mysol,
Buru,
Bismarck-
Archipel,
Andamanen,
Shortland
Ins.,
Queensland
- Japan,
Ostchina,
Sumatra,
Java
- Ägypten
(Typus:
Bacos)

b) Orientalische Region.

- megaspilalis** Hampson, Trans. Ent. Soc. Lond., p. 211 (♂) (1897).
- torsicostalis** Hampson, Trans. Ent. Soc. Lond., p. 212 (1897).
- Batchian
Amboina,
Holl.
Neu-Guinea

- violalis** Lederer, Wiener Ent. Monatschr., 7, p. 431, t. 16, f. 15, t. 5, f. 14, 35 (1863) (Amboina). — Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1485 (1865). — Hampson, Trans. Ent. Soc. Lond., p. 212 (1897). Amboina, Ceram, Neu-Guinea
- aegimlusalis** Walker, vide sub a) Paläarktische Region, p. 172.
- reducta** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1296 (♂) (1865) (*Desmia*). — Hampson, Trans. Ent. Soc. Lond., p. 212 (1897). Ceram, Mysol, Buru (Typus: Mysol)
- hadesialis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 14 (♂) (1907). Penang
- albicinctata** Hampson, Trans. Ent. Soc. Lond., p. 212 (♂, ♀) (1897). Batchian, Shortland Ins.
- batjanalis** Strand, Arch. Naturg., 85, A H. 12, p. 134 (1920). Batchian
- octosemalis** Hampson, Fauna Brit. India Moths, 4, p. 236 (♂) (1896); Trans. Ent. Soc. Lond., p. 213 (1897). Birma
- orientalis** Snellen, Tijds. v. Ent., 23, p. 233 (1880); 27, p. 43, t. 4, f. 3, 3 a (1884) (*Aediodes*)! — Hampson, Trans. Ent. Soc. Lond., p. 213 (1897). Celebes, Java (Typus: Celebes)
- javanalis** Snellen, Tijds. v. Ent., 43 (1900), p. 278 (♂, ♀), t. 15, f. 17 (♂) (1901) (*Diploptyla*). Java
- albifunata** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 143 (♀) (1896) (*Diploptyla?*). — Hampson, Fauna Brit. India Moths, 4, p. 237 (♀) (1896); Trans. Ent. Soc. Lond., p. 213 (1897). Assam (Khasis)
- discalis** Hampson, Journ. Bombay Nat. Hist. Soc., 15, p. 212 (♀) (1903). Ceylon
- flexiguttalis** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 144 (1896). — Hampson, Fauna Brit. India Moths, 4, p. 238 (♂, ♀) (1896); Trans. Ent. Soc. Lond., p. 213 (1897). Assam, Tonkin
- maculifrons** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 269 (♂) (1917). Assam (Khasis)
- sodalis** Leech, vide sub a) Paläarktische Region, p. 172.
- confinalis** Caradja, Mem. Acad. Rom. Sect. Stiint., (s. 3) 4, mem. 8, p. 406 (1927). Westchina
- costipunctata** Warren, Ann. Mag. Nat. Hist., (s. 6) 18, p. 217 (1896) (*Danaga*). — Hampson, Fauna Brit. India Moths, 4, p. 237 (1896); Trans. Ent. Soc. Lond., p. 213 (1897). Birma, Borneo
- illectalis** Walker (nec Walk. 1859), Cat. Lep. Het. Brit. Mus., 19, p. 931 (♂) (1859) (*Desmia?*) (Sarawak) (nomen praeocc.).
- conceisalis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 345 (♀) (1859) (*Desmia*) (Ceylon). — Moore, Lep. Ceylon, 3, p. 272, t. 179, f. 3 (1885) (*Danaga*). — Hampson, Ill. Typ. Sp. Lep. Het. Brit. Mus., 8, p. 39 (1891); 9, p. 49 (1893) (*Danaga*); Fauna Brit. India Südindien, (Nilgiris), Ceylon, Borneo

- Moths, 4, p. 237 (1896); Trans. Ent. Soc. Lond., p. 213 (1897).
- discisignalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 270 (♀) (1917). Assam (Khasis)
- illectalis** Walker, Cat. Lep. Het. Brit. Mus., 17, p. 362 (♂) (1859) (*Salbia*?) (Sarawak). — Hampson, Trans. Ent. Soc. Lond., p. 213 (1897). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 450, t. 8, f. 21 (1900).
- a ceo alis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 929 (♂) (1859) (*Desmia*?) (Sarawak).
- zebinalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 932 (♀) (1859) (*Desmia*) (Sarawak). — Meyrick, Trans. Ent. Soc. Lond., p. 467 (1894) (*Ptilacola*). — Hampson, Trans. Ent. Soc. Lond., p. 213 (1897). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 451, t. 7, f. 13 (1900). Borneo
- latalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1294 (♀) (1865) (*Botys*). — Hampson, Trans. Ent. Soc. Lond., p. 213 (1897). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 450, t. 7, f. 15 (1900). Sula
- basalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1295 (♂) (1865) (*Botys*). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 450 (1900). Sula
- latalis** (part.) Hampson, Trans. Ent. Soc. Lond., p. 213 (1897).
- tellesalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 933 (♀) (1859) (*Desmia*) (Sarawak). — Meyrick, Trans. Ent. Soc. Lond., p. 86 (1897) (*Diploctyla*). — Hampson, Trans. Ent. Soc. Lond., p. 213 (1897) (part.). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 450, t. 8, f. 14 (1900). Borneo, Sangir
- parvalis** Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1300 (♀) (1865) (*Aedides*). Sula
- purvalis** Hampson, Trans. Ent. Soc. Lond., p. 213 (1897) (ex synon.).
- tellesalis** Hampson, Trans. Ent. Soc. Lond., p. 213 (1897) (part.).
- lycopusalis** Walker, Cat. Lep. Het. Brit. Mus., 19, p. 930 (♀) (1859) (*Desmia*?) Sarawak). — Hampson, Trans. Ent. Soc. Lond., p. 213 (1897). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 450, t. 7, f. 23 (1900). Borneo
- viola** Butler, Ann. Mag. Nat. Hist., (s. 5) 10, p. 234 (♂) (1882) (*Pyrausta*) (Herz. v. Yrk Ins.). — Hampson, Trans. Ent. Soc. Lond., p. 213 (1897) (*Piletocera*). — Pagenstecher, Zoologica, H. 29, p. 187 (1900). Amboina, Banda Ins., Bismarck-Archipel, Herzog v. York Ins.
- aurantialis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 271 (1917). Batchian

- elongalis** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 145 (1896) (*Thysanodesma*). — Hampson, Fauna Brit. India Moths, 4, p. 238 (♂) (1896); Trans. Ent. Soc. Lond., p. 213 (1897). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 166 (1928). Indien, Assam (Khasis), Formosa
- flavidiscalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 272 (♂) (1917). Sikkim
- metochrealis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 272 (1917). Ceylon
- conchylia** Meyrick, Trans. Ent. Soc. Lond., p. 468 (♂) (1894) (*Nausinoe*). — Hampson, Trans. Ent. Soc. Lond., p. 214 (1897). Pulo Laut
- cumulalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 16 (♂) (1907). Borneo
- micralis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 16 (♂) (1907). Borneo
- costifascialis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 273 (♂) (1917). Malakka (Selangor)
- nudicornis** Hampson, Trans. Ent. Soc. Lond., p. 214 (♂) (1897); Journ. Bombay Nat. Hist. Soc., 12, p. 704 (1899). Assam (Typus: Khasis)

Species incertae sedis.

- immundalis** R. South in Leech, Trans. Ent. Lond., p. 460, t. 15, f. 4 (1901) (*Nacoleia*). — Caradja, Bull. Acad. Poum., 15, p. 149 (1932) (*Piletocera*). Zentralchina (Typus: Chang-yang- & Ichang)
- litanalis** Walker, Cat. Lep. Het. Brit. Mus., 18, p. 706 (♀) (1859) (*Botys?*) (Sarawak): — Hampson, Trans. Ent. Soc. Lond., p. 144 (1897) (*Nymphula litanalis*). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 438 (1900) (*Nymphula*). — De Joannis, Ann. Soc. Ent. Fr., 98, p. 654 (1929) (*Piletocera*). Borneo, Tonkin
- stenialis** Warren, Ann. Mag. Nat. Hist., (s. 6) 8, p. 68 (1891) (*Endotricha*).
- quadralis** Snellen, Tijdschr. v. Ent., 43 (1900), p. 280 (♂) (1901) (*Diplotyla*). Java (Typus: W. Java, Preanger)

c) Aethiopische Region.

- signiferalis** Wallengren, Wien. Ent. Monatschr., 4, p. 175 (1860) (*Isopteryx*). — Hampson, Trans. Ent. Soc. Lond., p. 210 (1897) (*P. Rinecera*). — Rebel, Denkschr. Akad. Wiss. Wien, 85, p. 430 (1910); 2. Beih. Jahrb. Hamb. Wiss. Anstalt, 32, p. 146 (1915). — Tams, Ins. Samoa, pars 3, fasc. 4, p. 269 (1935). Natal, Buru, Fidji, Ellice Ins., Carolinen, Marquesas Ins., Austral. Ins. (Rapa, Rurutu),
- barbicornis** Felder et Rogenhofer, Reise Novara (Het.), p. 6, t. 136, f. 1, (1874) (*Ceratoclasia*).

- mirabilis* Butler, Mem. Nat. Acad. Sci., p. 95 (1884) (*Rinecera*); id., Trans. Ent. Soc. Lond., p. 424 sub n. 69 (1886) (*Rinecera*). Gesellschaft.-Ins. (Tahiti), Samoa,
- nigrescens* Butler, Trans. Ent. Soc. Lond., p. 424 (1886) (*Rinecera*). (Typus: Viti)
- lanceolalis* Guenée, Delt. et Pyral., p. 246, t. 10, f. 8 (1854) (*Stenia*) (Kap d. gut. Hoffg.). Südafrika
— Hampson, Trans. Ent. Soc. Lond., p. 213 (1897) (*P. (Danaga)*).
- anesalis* Walker, Cat. Lep. Het. Brit. Mus., 18, p. 640 (1859) (Cape).
- rufulalis* Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 18 (1907). Sierra Leone
- flavalis* Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 274 (1917). Nyassaland

d) Australische Region.

- ochrosema* Meyrick, Trans. Ent. Soc. Lond., p. 209 (♂) (1897) (*P. (Diptolyta)*). — Hampson, Trans. Ent. Soc. Lond., p. 209 (1897) (*P. (Diptolyta)*). Neu-Hebriden (Typus), Fidji Ins.
- chloronata* Meyrick, Trans. Ent. Soc. Lond., p. 514 (♂, ♀) (1889) (*Diptolyta*). — Hampson, Trans. Ent. Soc. Lond., p. 209 (1897) (*P. (Diptolyta)*). — Pagenstecher, Zoologica, H. 29, p. 183 (1900). Neu-Guinea, Bismarck-Archipel (Typus: Mt. Obra)
- cyclopila* Meyrick, Trans. Ent. Soc. Lond., p. 248 (♂) (1886) (*Diptolyta*). — Hampson, Trans. Ent. Soc. Lond., p. 209 (1897) (*P. (Diptolyta)*). — Rebel, Denkschr. Akad. Wiss. Wien, 85, p. 430 (1910); 2. Beih. Jahrb. Hamb. Wiss. Anstalt, 32, p. 145 (1915). Samoa
- signiferalis* Wallengren, vide sub c) Aethiopische Region, p. 175.
- steffanyi* Tams, Ins. Samoa, part. 3 (fasc. 4), p. 267, t. 17, f. 1, 2 (1935). Samoa
- rechingeri* Tams, Ins. Samoa, part. 3 (fasc. 4), p. 267, t. 17, f. 3 (1935). Samoa
- albipictalis* Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 12 (♂) (1907). Salomon Ins.
- stygialis* Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 13 (♂) (1907). Salomon Ins.
- scotochroa* Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 266 (♂) (1917). Salomon Ins.
- albiventralis* Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 266 (♂) (1917). Salomon Ins.
- purpureofusa* Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 266 (♂) (1917). Salomon Ins.
- denticostalis* Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 13 (♂) (1907). Salomon Ins.

- infernalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 14 (♂) (1907). Salomon Ins.
- xanthosoma** Meyrick, Trans. Ent. Soc. Lond., p. 249 (♂, ♀) (1886) (*Strepsimela*). — Hampson, Trans. Ent. Soc. Lond., p. 210 (1897) (*P. Strepsimela*). — Rebel, Denkschr. Akad. Wiss. Wien, 85, p. 430 (1910); 2. Beih. Jahrb. Hamb. Wiss. Anstalt, 32, p. 146 (1915). Samoa
- pseudadelpha** Meyrick, Trans. Ent. Soc. Lond., p. 233 (♂, ♀) (1887) (*Strepsimela*). — Hampson, Trans. Ent. Soc. Lond., p. 210 (1897). Fidji Ins.
- vestigialis** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 144 (1896) (*Diploptyla*) (Queensland). — Hampson, Trans. Ent. Soc. Lond., p. 210 (1897). — Rebel, Denkschr. Akad. Wiss. Wien, 85, p. 430 (1910); 2. Beih. Jahrb. Hamb. Wiss. Anstalt, 32, p. 146 (1915). — Tams, Ins. Samoa, part. 3 (fasc. 4), p. 269 (1935). Queensland, Fergusson Ins., Holl. Neu-Guinea, Salomon Ins., Samoa
- epipercialis** Hampson, Trans. Ent. Soc. Lond., p. 210 (♂) (1897). Fergusson Ins., Holl. Neu-Guinea
- atrata** Pagenstecher, Zoologica H. 29, p. 182, t. 1, f. 5 (1900) (*Diploptyla*). Bismarck-Archipel (Neu-Pommern)
- erebina** Butler, Trans. Ent. Soc. Lond., p. 425, t. 10, f. 7 (1886) (*Hormatholepis*). — Hampson, Trans. Ent. Soc. Lond., p. 211 (1897). Fidji Ins. (Typus: Viti)
- melanauges** Meyrick, Trans. Ent. Soc. Lond., p. 246 (♂, ♀) (1886) (*Erebangela*). — Hampson, Trans. Ent. Soc. Lond., p. 211 (1897) (*P. Erebangela*). Fidji Ins.
- hecate* Butler, Trans. Ent. Soc. Lond., p. 421 (1886) (*Graphicopoda*). Tonga
- torsicostalis** Hampson, vide sub b) Orientalische Region, p. 172.
- plumbicostalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 266 (♂) (1917). Holl. Neu-Guinea
- violalis** Lederer, vide sub b) Orientalische Region, p. 173.
- aeginiusalis** Walker, vide sub a) Paläarktische Region, p. 172.
- ruficeps** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 267 (♂) (1917). Brit. Neu-Guinea
- violascens** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 267 (♂) (1917). Holl. Neu-Guinea, Fergusson Ins.
- albimixtalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 268 (♂) (1917). Queensland
- microdentalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 14 (♀) (1907). Woodlark Ins.
- reducta** Walker, vide sub b) Orientalische Region, p. 173.

- albicinctata** Hampson, vide sub b) Orientalische Region, p. 173.
- stenipteralis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 268 (♂) (1917). Brit. Neu-Guinea
- leucocephalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 269 (♂, ♀) (1917). Bismarck-Archipel (Rook Ins.)
- leucogastralis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 269 (♂) (1917). Holl. Neu-Guinea
- bisignalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 270 (♀) (1917). Bali
- bistrigalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 271 (♀) (1917). Holl. Neu-Guinea
- viola** Butler, vide sub b) Orientalische Region, p. 174.
- dactyloptila** Meyrick, Trans. Ent. Soc. Lond., p. 243 (♂, ♀) (1886) (*Semioceros*). — Hampson, Trans. Ent. Soc. Lond., p. 213 (1897) (*P. (Danaga)*). Fidji Ins.
- phaeocraspedalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 15 (♀) (1907). Salomon Ins.
- rotundalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 15 (♂) (1907). Bali, Talaut, Woodlark Ins., Louisiaden (St. Aignan)
- ulophanes** Meyrick, Trans. Ent. Soc. Lond., p. 245 (♂, ♀) (1886) (*Ptiacina*). — Hampson, Trans. Ent. Soc. Lond., p. 213 (1897) (*P. (Ptiacola)*). Fidji Ins.
- mesophaealis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 273 (♀) (1917). Holl. Neu-Guinea
- holophacalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 274 (1917). Holl. Neu-Guinea
- analytodes** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 274 (♂) (1917). Bismarck-Archipel (Rook Ins.)

Species incertae sedis.

- albescens** Rebel, Mitt. Zool. Mus. Hamburg, 32, p. 135, t. 1, f. 11 (1915). — Tams, Ins. Samoa pars 3, fasc. 4, p. 270, t. 16, f. 10, 11, text-fig. 7 (1935). Fidji Ins.
- argopis** Meyrick, Trans. Ent. Soc. Lond., p. 248 (♀) (1886) (*Diploptyla*). — Hampson, ibid., p. 214 (1897). Samoa
- cnneaspila** Meyrick, Exot. Microlep., 4, p. 398 (♀) (1933). Fidji Ins. (Typus: Vunidawa)
- inconspicualis** Kenrick, Proc. Zool. Soc. Lond., p. 78, t. 4, f. 69 (1907). Brit. Neu-Guinea

- meeckii** Lucas, Proc. Linn. Soc. N. S. Wales, 8, p. 159 (♂) (1893) (*Diploptyla*). — Hampson, Trans. Ent. Soc. Lond., p. 215 (1897). West-Australien
- microcentra** Meyrick, Trans. Ent. Soc. Lond., p. 249 (♀) (1886) (*Strepsimela* (?)). — Hampson, Trans. Ent. Soc. Lond., p. 215 (1897). Fidji Ins.
- nasonia** Meyrick, Exot. Microlep., 4, p. 398 (♂) (1933). Fidji Ins.
(Typus: Vunidawa)
- ocelligera** Meyrick, Exot. Microlep., 4, p. 241 (♂, ♀) (1932). Fidji Ins.
(Typus: Vunidawa & Suva)
- rhopalophora** Meyrick, Exot. Microlep., 4, p. 501 (♂) (1934). Fidji Ins.
(Typus: Vunidawa)

e) Neotropische Region.

- bufalls** Guenée, Delt. et Pyral., p. 245 (1854) (*Stenia*) (Cayenne). — Hampson, Trans. Ent. Soc. Lond., p. 214 (1897) (*Piletocera*). — Druce, Biol. Centr.-Amer. Lep. Het., 2, p. 561, t. 101, f. 21 (1899). — Dyar, Journ. N. York Ent. Soc., 14, p. 98 (♂) (1906). Fr. u. Holl. Guyana, Westindien, Panama, Florida
- plebeialis** Walker (nec Lederer, 1863), Cat. Lep. Het. Brit. Mus., 34, p. 1385 (1865) (*Botys*). — Dyar, Proc. U. S. Nat. Mus., 47, p. 285 (1914).
- stercoralls** Möschler, Verh. z. b. Ges. Wien, 31, p. 419, f. 29 (1881) (*Botys*). — Dyar, Proc. U. S. Nat. Mus., 47, p. 285 (1914). Surinam, Bermuda Ins.
- bufalis** (part.) Hampson, Trans. Ent. Soc. Lond., p. 214 (1897). — Dyar, Journ. N. York Ent. Soc., 14, p. 98 (1906).
- ranalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 17 (♂) (1907). Brasilien (Unt. Amazonas)
- simplicialis** Barnes et Mc Dunnough, Contr. Lep. Nat. Hist. Lep. N. Amer., 2 (4), p. 175 (♂, ♀), t. 4, f. 1 (♀) (1913). — Dyar, Proc. U. S. Nat. Mus., 47, p. 285 (1914). Florida, Panama, Costa Rica (Typus: ♀ Everglade, Florida)
- albicollialis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 17 (♀) (1907). Brasilien (Unt. Amazonas)
- fulvalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 18 (♀) (1907). Brit. Guyana

Species incertae sedis.

- agathanalis** Schaus, Proc. Ent. Soc. Wash., 26, p. 130 (♂) (1924). Panama (Typus: Tabernilla)

- grammalis** Guenée, Delt. et Pyral., p. 249 (1854) Cayenne
(*Stenia*). — Hampson, Trans. Ent. Soc. London, p. 215 (1897).
inconspicua Schaus., Ann. Mag. Nat. Hist., (s. 8) Costa Rica
9, p. 680 (♀) (1912).
99

67. Genus **Glaucoda** Karsch

Karsch, Ent. Nachr., 26, p. 247 (1900).

- transparitalis** Karsch, Ent. Nachr., 26, p. 248 (♂, ♀), Togo,
(1900). Kamerun,
Spanisch
Guinea,
Congo

1

68. Genus **Aediodina** Strand

Strand, Ent. Mitt., 8, p. 133, 134 (1919).

Aediodes Lederer (nec Guenée, 1854), Wien. Ent. Monatschr., 7, p. 433 (1863).

- quaternalis** Lederer, Wien. Ent. Monatschr., 7, p. 434, 483, t. 17, f. 1 (1863) (*Aediodes*) (Amboina). — Hampson, Proc. Zool. Soc. Lond., p. 654 (1898) (*Bocchoris*). — Strand, Ent. Mitt., 8, p. 133 (1919) (*Aediodina*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 167 (1928). Indien,
Java,
Amboina,
Formosa,
Queensland

1

69. Genus **Alufefa** Swinhoe

Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 457 (1900).

- aequalis** Walker, Cat. Lep. Het. Brit. Mus., 34, Neu-Guinea
p. 1299 (♂) (1865) (*Aediodes*). — Hampson,
Trans. Ent. Soc. Lond., p. 214 (1897) (*Piletocera*). — Swinhoe, Cat. East. Austral. Lep.
Het. Oxford Mus., 2, p. 457, t. 7, f. 9 (1900).

1

70. Genus **Neogenesis** Hampson

Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 19 (1907).

- flaviplagialis** Hampson, Ann. Mag. Nat. Hist., (s. 7) Brit. u.
19, p. 19 (1907). — Kenrick, Proc. Zool. Soc. Holl.
Lond., p. 548 (1912). Neu-Guinea

1

71. Genus **Heringiella** Berg

C. Berg, Com. Mus. B. Aires, 1, p. 17 (1898) (n. nov.).

Heringia v. Hedemann, Ent. Zeit. Stettin, 55, p. 293
(1894) (nec Rondani, 1856).

- mendica** von Hedemann, Ent. Zeit. Stettin, 55, p. 294 Dänisch-
(1894) (*Heringia*). Westindien

1

72. Genus **Ellogima** Turner

Turner, Proc. Roy. Soc. Queensland, 24, p. 149 (1913).

macroperalis Hampson, Trans. Ent. Soc. Lond., p. Queensland
214, (1897) (*Piletocera*). — Turner, Proc. (Typus:
Roy. Soc. Queensland, 24, p. 149 (♂) (1913) Cedar Bay,
(*Ellogima*). Cooktown)

1

73. Genus **Camptomastix** Warren

Warren, Ann. Mag. Nat. Hist., (s. 6) 9, p. 439 (1892).

Camptomastix Hampson, Fauna Brit. India Moths, 4,
p. 238 (1896); Trans. Ent. Soc. Lond., p. 134, 215 (1897);
Journ. Bombay Nat. Hist. Soc., 18, p. 575 (1908).

(Typus generis: *hisbonalis* Walker)

turbatalis Christoph, Bull. Soc. Imp. Nat. Moscou, Ussuri,
56 (1), p. 31 (1881) (*Orphnophanes*). — Amur
Hampson, Trans. Ent. Soc. London, p. 189
(1897). — Zerny, Ann. Naturh. Hofmus.
Wien, 28, p. 324 (1914) (*Stenia*). — Caradja,
Deutsche Ent. Zeitschr. Iris, 30, p. 21 (1917)
(*Camptomastix*).

hisbonalis Walker, Cat. Lep. Het. Brit. Mus., 18, Indien
p. 707 (♀) (1859) (*Botys*) (Sarawak). — (Himalaya,
Warren, Ann. Mag. Nat. Hist., (s. 6) 9, p. 439 Simla),
(1892) — Hampson, Fauna Brit. India Moths, Birma,
4, p. 239, f. 143 (♂) (1896) (*Camptomastix*); Assam,
Trans. Ent. Soc. Lond., p. 215, fig. (♂) West- u.
(1897). — Leech, Trans. Ent. Soc. Lond., p. Mittelchina,
443 (1901). — Marumo, Journ. Coll. Agr. Japan,
Tokyo Imp. Univ., 8, p. 183 (1923). — Formosa,
Shibuya, Journ. Fac. Agr. Hokkaido Imp. Tonkin
Univ., 22, p. 167, t. 6, f. 29 (1928); Ins.
Matsum., 3, p. 133, t. 5, f. 25 (1929).

pacalis Leech, Entom., 22, p. 69 (♀), t. 4, (Typus:
f. 15 (♀) (1889) (*Botys*) (China). — Warren, Ningpo)
Ann. Mag. Nat. Hist., (s. 6) 9, p. 439
(1892) (*Camptomastix*).

longipalpis Butler, Ill. Typ. Sp. Lep.
Het. Brit. Mus., 7, p. 95, t. 135, f. 4 (1889)
(*Diploptyla*) (Dharmasala).

areolifera Strand, Deutsche Ent. Zeit-
schr. Iris, 32, p. 64 (1918) (*Thliptoceras*).
— Shibuya, Journ. Fac. Agr. Hokkaido Imp.
Univ., 22, p. 26 (1928).

exuvialis Snellen, Trans. Ent. Soc. Lond., p. 627 Sikkim,
(1890) (*Diploptyla*). — Hampson, Fauna Brit. Assam
India Moths, 4, p. 239 (♂) (1896).

baracoalis Schaus, Proc. Ent. Soc. Wash., 26, p. Cuba
123 (♀) (1924). (Typus:
Baracoa)

4

74. Genus **Hydropionea** Hampson

Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 275 (1917).

Clupeosoma (part.) Hampson, Trans. Ent. Soc. London, p. 216 (1897).

Sufetula (part.) Druce, Biol. Centr.-Amer., Lep. Het., 2, p. 204 (1895) (*S. melliculalis*).

(Typus generis: *melliculalis* Led.)

melliculalis Lederer, Wien. Ent. Monatschr., 7, p. 414, t. 15, f. 6 (*Sparagmia*?) (Brasilien). — Zentr.-u. Druce, Biol. Centr.-Amer. Lep. Het., 2, p. 204 (1895) (*Sufetula*). Südamerika (Guatemala, Costa Rica, Venezuela),
melliculale Hampson, Trans. Ent. Soc. London, p. 217 (1897) (*Clupeosoma*); Ann. Mag. Nat. Hist., (s. 8) 20, p. 275 (1917) ?Philippinen (*Hydropionea*).

fenestralis Barnes et Mc Dunnough, Contr. Nat. Hist. Lep. N. Amer., 2 (6), p. 235 (♂, ♀), t. 1, f. 11 (1914) (*Diasemia*?); 4, p. 169 (1917). — Dyar, Insec. Insc. Menstr., 5, p. 79 (1917). Arizona (Typus: St. Catalina)

oblectalis Hulst, Trans. Amer. Ent. Soc., 13, p. 154 (1886) (*Loxostege*). — Barnes et Mc Dunnough, Contr. Nat. Hist. Lep. N. Amer., 4, p. 170 (1918) (*Hydropionea*). Arizona
eumoros Dyar, Insec. Insc. Menstr., 5, p. 79 (1917) (*Clupeosoma*). — Barnes et Mc Dunnough, Contr. Nat. Hist. Lep. N. Amer., 4, p. 170 (1918) (*Hydropionea*).

3

75. Genus **Zagiridia** Hampson

Hampson, Trans. Ent. Soc. Lond., p. 134, 216 (1897).

(Typus generis: *noctualis* Hamps.)

noctualis Hampson, Trans. Ent. Soc. Lond., p. 216, fig. (♂) (1897). S.O.-Borneo, Pulo Laut (Typus)

rufalis Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 275 (♀) (1917). Woodlark Ins.

2

76. Genus **Clupeosoma** Snellen

Snellen, Tijds. v. Ent., 23, p. 203 (1880). — Hampson, Trans. Ent. Soc. Lond., p. 134, 216 (1897) (part.) — Pagenstecher, Zoologica H. 29, p. 183 (1900).

Hydrorybina Hampson, Fauna Brit. India Moths, 4, p. 239 (1896) (Typus: *polusalis* Walk.). — Swinhoe, Ann. Mag. Nat. Hist., (s. 7) 8, p. 139 (1901).

(Typus generis: *pellucidalis* Snell.)

a) Paläarktische Region.

- pryeri** Butler, Trans. Ent. Soc. Lond., p. 588 (1881) Japan
(Anemosa). — Pryer, Trans. Asiat. Soc. (Typus:
 Japan, 13, p. 67 (1885) *(Anemosa)*. Yokohama)
 Hampson, Trans. Ent. Soc. Lond., p. 217
 (1897); Leech, *ibid.*, p. 444 (1901). — Mat-
 sumura, Cat. Ins. Jap., p. 205 (1905). —
 Shibuya, Ins. Matsum., 3, p. 133, t. 5, f. 26
 (1929).

b) Orientalische Region.

- polusalis** Walker, Cat. Lep. Het. Brit. Mus., 18, Sikkim,
 p. 703 (♀) (1859) *(Botys)* (Sarawak). — Birma,
 Hampson, Fauna Brit. India Moths, 4, p. Ostchina,
 239, f. 144 (♂) (1896) *(Hydrorybina)*. — Tonkin,
 Swinhoe, Ann. Mag. Nat. Hist., (s. 7) 8, p. Borneo,
 139 (1901) *(Hemiscopsis)*. — E. Hering, Ent. Sumatra,
 Zeit. Stettin, 64, p. 48 (1903). Holl.
polusale Hampson, Trans. Ent. Soc. Lond., Neu-Guinea,
 p. 217, fig. (♂) (1897) *(Clupeosoma)*. — Queensland
 Swinhoe, Cat. East. Austral. Lep. Het. Ox-
 ford Mus., 2, p. 451, t. 7, f. 20 (1900)
(Clupeosoma).
 Biologie: Caradja, Deutsche Ent. Zeit-
 schr. Iris, 47, p. 156 (1933).
- violascens** Hampson, Ann. Mag. Nat. Hist., (s. 8) Borneo,
 20, p. 275 (♀) (1917). Pulo Laut
- bicolor** Moore, Descr. Lep. Ins. Atkinson, p. 207 Indien
 (1887). — Hampson, Fauna Brit. India (Bengalen),
 Moths, 4, p. 240 (♂, ♀) (1896) *(Hydrorybina)*; Trans. Ent. Soc. Lond., p. 217 (1897). Assam,
rubralis Warren, Ann. Mag. Nat. Hist., Ceylon
 (s. 6) 17, p. 97 (1896) *(Eurycreon?)*.
- pellucidalis** Snellen, Tijds. v. Ent., 23, p. 204 (♂, Celebes
 ♀) (1880); 26, p. 124, t. 6, f. 9, 9a (caput) (Typus:
 (1883). — Pagenstecher, Zoologica H. 29, Makassar),
 p. 183 (1900). Sumbawa,
pellucidale Hampson, Trans. Ent. Soc. Tenimber,
 Lond., v. 217 (1897). Dili,
 Fergusson
 Ins.,
 Bismarck-
 Archipel
 (Neu-
 Pommern)
- glaucinalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) Singapore,
 20, p. 278 (♀) (1917). Sumbawa

c) Aethiopische Region.

- noordimimalis** Hampson, Ann. Mag. Nat. Hist., Goldküste
 (s. 8) 20, p. 276 (♂) (1917).

d) Australische Region.

- polusalis** Walker, vide sub b) Orientalische Region, p. 183.
- brunneofusalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 276 (♀) (1917). Holl. Neu-Guinea
- pellucidalis** Snellen, vide sub b) Orientalische Region, p. 183.
- atristriata** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 277 (♂, ♀) (1917). Timur, Bismarck-Archipel (Rook Ins.)
- laniferalis** Hampson, Ann. Mag. Nat. Hist., (s. 7) 19, p. 19 (♂) (1907). Louisiaden (St. Aignan)
- rufistriata** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 277 (♂) (1917). Holl. Neu-Guinea
- astrigalis** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 278 (♂) (1917). Queensland
- metachryson** Hampson, Trans. Ent. Soc. Lond., p. 218 (♂) (1897). Fergusson Ins.

Species incertae sedis.

- lampra** Tams, Ins. Samoa pars 3 (fasc. 4), p. 271, t. 15, f. 9 (♀) (1935). Samoa
- mimetica** Lower, Trans. Roy. Soc. S. Austral., 27, p. 65 (♀) (1903). West-australien
- photina** Tams, Ins. Samoa, pars 3 (fasc. 4), p. 272, t. 15, f. 10 (♀) (1935). Samoa
- rhodea** Lower, Trans. Roy. Soc. S. Austral., 29, p. 107 (♀) (1905). Queensland
- strigatalis** W. Rothschild, Rep. B. O. U. Exp., 2, nr. 15, p. 126 (♂) (1915). Holl. Neu-Guinea

e) Neotropische Region.

- dentata** Druce, Biol. Centr.-Amer. Lep. Het., 2, p. 203 (♂), t. 60 f. 13 (1895) (*Sufetula*). Zentral-Amerika
- dentatum* Hampson, Trans. Ent. Soc. Lond., p. 217 (1897). (Guatemala) (Typus: Vera Paz)
- rusina** Druce, Biol. Centr.-Amer. Lep. Het., 2, p. 203, t. 60, f. 14 (1895) (*Sufetula*). Mexiko
- rusinum* Hampson, Trans. Ent. Soc. Lond., p. 217 (1897). (Typus: Las Vigas)
- basale** Hampson, Trans. Ent. Soc. Lond., p. 217 (♂) (1897). Barbados

Species incertae sedis.

- barnesalis** Dyar, Insec. Insc. Menstr., 11, p. 27 (♂, ♀) (1923). Guatemala (Typus: Santa Maria Volcano)

- brevicans** Dyar, Insec. Insc. Menstr., 11, p. 26 Guatemala
(♂, ♀) (1923). (Typus:
Santa Maria
Volcano)
- lavinia** Schaus, Ann. Mag. Nat. Hist., (s. 8) 9, p. 680 Costa Rica
(♀) (1912).
- protopennis** Dyar, Insec. Insc. Menstr., 11, p. 27 Mexiko
(♂) (1923). (Typus:
Zacualpan)
- pseudopis** Dyar, Proc. U. S. Nat. Mus., 47, p. 396 Mexiko
(♀) (1914).
- schausalis** Dyar, Insec. Insc. Menstr., 11, p. 26 Guatemala
(♂, ♀) (1923). (Typus:
Santa Maria
Volcano)
- sufflexalis** Dyar, Proc. U. S. Nat. Mus., 47, p. 396 Mexiko
(1914).

28

77. Genus **Perinephela** Hübner

Hübner, Verz. bek. Schm., p. 357, n. 3418 (1816) [1826] (part.). —
Meyrick, Revis. Handb. Brit. Lep., p. 424 (1928).

- Perinephela* Heinemann, Kleinschm., 1, H. 2, p. 97 (1865).
— Rebel in Spuler, Schm. Eur., 2, p. 222 (1906). — Wahl-
gren, Svensk Insektfauna, 10 (2), p. 48 (1915). — M.
Hering, Tierwelt Mitteleur. (Ergbd., 1), p. 282 (1932). —
Eckstein, Schm. Deutschl., 5, p. 62 (1933).
Perinephila Hampson, Trans. Ent. Soc. Lond., p. 219,
220 (1897) (sectio I = subgenus). — Staudinger et Rebel,
Cat. Lep. Pal. Faun., (ed. 3) 2, p. 50 (1901).
Perinephile Barrett, Lep. Brit. Isl., 9, p. 234 (1904).
Psammotis (part.) Hampson, Trans. Ent. Soc. London,
p. 219 (1897).

(Typus generis: *lancaulis* (Schiff. et Den.))

- lancaulis** [Schiffermüller et Denis], Syst. Verz. Schm. England,
Wien, p. 121 (1775) (*Pyralis*). — Hübner, Schweden,
Verz. bek. Schm., p. 357 (1826) (*Perinephela*). Dänemark,
— Treitschke, Schm. Eur., 7, p. 79 (1829) Livland,
(*Botys*). — Duponchel, Hist. Nat. Lep. Fr., Mittel- u.
8 (2), p. 111, t. 216, f. 4, 5 (1831) (*Botys*). Südeuropa
— Wood, Index Entom., p. 124, t. 28, f. 809 (Frankreich,
(1839) (*Margaritia*). — Herrich-Schaeffer, Belgien,
Syst. Bearb. Schm. Eur., 4, p. 29 (1848) N.-Spanien,
(*Botys*). — Guenée, Delt. et Pyral., p. 338 Ostpyrenäen,
(1854) (*Botys*). — Stainton, Man. Brit. Butt. Italien,
Moths, 2, p. 149 (1859). — Heinemann, Bosnien,
Kleinschm., 1, H. 2, p. 97 (1865). — Berce, Herzego-
Fauna Ent. Fr. Lep., 6, p. 135 (1878). — wina,
— Snellen, Vlinders Nederl., 2 (1), p. 70 (1882). Dalmatien,
— Meyrick, Trans. Ent. Soc. Lond., p. 445 Rumänien),
(1890); Handb. Brit. Lep., p. 407 (1895). — Cis-
Hampson, Trans. Ent. Soc. Lond., p. 219, kaukasien,

- fig. (♂), 220 (1897) (*Psammotis*) (*Perinephela*). — Leech, Trans. Ent. Soc. Lond., p. 444 (1901) (*Psammotis*). — Barrett, Lep. Brit. Isl., 9, p. 234, t. 402, f. 4, 4a, b (larva) (1904). — Rebel in Spuler, Schm. Eur., 2, p. 222, t. 82, f. 41 (1906). — Wahlgren, Svensk Insektafauna, 10 (2), p. 48, t. 2, f. 55 (1915). — Meyrick, Revis. Handb. Brit. Lep., p. 424 (1928). — Shibuya, Ins. Matsum., 3, p. 133, t. 5, f. 27 (1929). — M. Hering, Tierwelt Mitteleur. (Ergbd., 1), p. 282 (1932). — Eckstein, Schm. Deutschl., 5, p. 62, t. 1, f. 22 (1933).
- glabralis* Hübner, Samml. Eur. Schmett. Pyr., p. 22, t. 10, f. 65 (♀), t. 18, f. 117 (♂) (1796) (*Pyralis*); Verz. bek. Schm., p. 357 (1816) [1826] (*Perinephela*).
- longalis* Haworth, Lep. Brit., p. 379 (1821) (*Pyralis*).
- Biologie: Heyden, Ent. Zeit. Stettin, 23, p. 172 (1862) (*Botys*). — Buckler, Ent. Mon. Mag., 12, p. 277 (1876) (*Botys*). — Larvae Brit. Lep., 9, p. 125, t. 153, f. 4 (1904). — Rebel in Spuler, Schm. Eur., 2, Nachtr.-Taf., 9, f. 55 (1906).
- doerriesi** Staudinger, Deutsche Ent. Zeitschr. Iris, 5, t. 3, f. 23 (1892); 6, p. 81 (1893) (*Perinephela*?). — Hampson, Trans. Ent. Soc. London, p. 220 (1897) (*Psammotis*).
- Amur,
Askold Ins.,
Ussuri,
Westchina

2

78. Genus **Zebronia** Hübner

Hübner, Verz. bek. Schm., p. p. 361 (1816) [1826] (part.). — Hampson, Trans. Ent. Soc. Lond., p. 123, 218 (1897).

(Typus generis: *foederalis* Hbn. = *phenice* Cram.).

- phenice** Cramer, Pap. Exot., 4, t. 382, f. G (1780). — Hampson, Trans. Ent. Soc. Lond., p. 219 (1897). — Butler, Proc. Zool. Soc. Lond., p. 443 (1898). — Kheil, Mem. Soc. Esp. Hist. Nat., 1, p. 181 (1905). — Strand, Arch. Naturg., 78 A, II. 1, p. 80 (1912).
- foederalis* Hübner, Verz. bek. Schm., p. 361, n. 3456 (1826) (*Zebronia*).
- podalirialis* Guenée, Delt. et Pyral., p. 281 (♂) (1854) (*Spilomela*) (Senegal, Gabun).
- podalivalis* Hampson, Trans. Ent. Soc. Lond., p. 219 (1897) (ex synonym.).
- phenicealis* Guenée, Delt. et Pyral., p. 281 (1854) (*Spilomela*) (Surinam).
- lathurusalis* Walker, Cat. Lep. Het. Brit. Mus., 19, p. 969 (1859) (*Zebronia*) (Sierra Leone).
- var. *mayottensis* Pagenstecher in Voeltzkow, Reise Ostafrika, 2, p. 132 (1907).
- West- u.
Zentralafrika
Congo,
Uganda,
Kamerun,
Brit.-
Deutsch- u.
Port.
O.-Afrika,
Span.
Guinea,
Abessinien,
Rhodesia,
Mauritius,
Komoren,
?Surinam
- Komoren

botydis Strand, Ent. Mitt., 4, p. 187 (1915).

Kamerun

semizebralis Walker, Cat. Lep. Het. Brit. Mus., 34, p. 1345 (♀) (1865) (*Zebronia*) (Bogotá).
Costa Rica,
Panama

3

79. Genus **Glyphandra** Karsch

Karsch, Ent. Nachr., 26, p. 249 (1900).

biincisalis Karsch, Ent. Nachr., 26, p. 249 (1900). Togo,
N.-Kamerun,
Span.
Guinea

1

80. Genus **Galadra** Walker

Walker, Cat. Lep. Het. Brit. Mus., 32, p. 640 (1865).

rhomboidalis Walker, Cat. Lep. Het. Brit. Mus., Neu-Guinea
32, p. 641 (1865).
rhomboidata Swinhoe, Cat. East.
Austral. Lep. Het. Oxford Mus., 2, p. 449,
t. 7, f. 6 (1900).

1

81. Genus **Hyaloplaga** Warren

Warren, Ann. Mag. Nat. Hist., (s. 6) 9, p. 432 (1892). — Hamp-
son, Fauna Brit. India Moths, 4, p. 235 (1896); Trans. Ent. Soc.
Lond., p. 133, 219 (1897).

pulchralis Moore, Proc. Zool. Soc. Lond., p. 869 Indien,
(1867) (*Hydrocampi*) (Sikkim). — Warren, Sikkim
Ann. Mag. Nat. Hist., (s. 6) 9, p. 432
(1892) (*Hyaloplaga*). — Hampson, Fauna
Brit. India Moths, 4, p. 235, f. 141 (♂)
(1896); Trans. Ent. Soc. Lond., p. 219, fig.
(♂) (1897).
var. *reducta* Caradja, Bull. Acad. Roum., China
14, p. 208 (1931). (Kwanhshien)

82. Genus **Psammofis** Hübner

Hübner, Verz. bek. Schm., p. 350, n. 3353 (1816) [1826]. —
Lederer, Wien. Ent. Monatschr., 7, p. 382 (1893). — Heinemann,
Kleinschm., 1, H. 2, p. 92 (1865) (*Psamotis*). — Wallengren,
Oefvers. K. Vet.-Akad. Förh., 28 (8), p. 1000 (1871). — Meyrick,
Trans. Ent. Soc. Lond., p. 448 (1890); Handb. Brit. Lep., p. 411
(1895). — Hampson, Trans. Ent. Soc. Lond., p. 134, 219 (1897)
(part.). — Rebel in Spuler, Schm. Eur., 2, p. 222 (1906). —
Wahlgren, Svensk Insektafauna, 10 (2), p. 48 (1915). — Caradja,
Deutsche Ent. Zeitschr. Iris, 30, p. 21 (1917). — Meyrick, Revis.
Handb. Brit. Lep., p. 427 (1928). — M. Hering, Tierwelt Mitteleur.
(Ergbd., 1), p. 284 (1932). — Eckstein, Schm. Deutschl., 5, p. 63
(1933).

- Lemia* Duponchel, Cat. Méth. Lep. Eur., p. 205 (1846)
(Typus: *pulveralis* Hübn.).
Lemiodes Guenée, Delt. et Pyral., p. 401 (1854) (Typus:
pulveralis Hübn.).
Hapalia (part.) Hübner, Verz. bek. Schmett., p. 355, n.
3397 (1816) [1826] (*H. hyalinialis* Hbn.).
(Typus generis: *pulveralis* Hübn.)

a) Paläarktische Region.

- pulveralis** Hübner, Samml. Eur. Schm. Pyral., p. Mittel- u.
29, t. 17, f. 109 (1796) (*Pyralis*); Verz. Nordeuropa,
Schm., p. 350, n. 3353 (1816) [1826]. — Frankreich,
Stephens, Ill. Brit. Ent. Haust., 4, p. 65 Spanien,
(1834) (*Margaritia*). — Treitschke, Schm. Ural,
Eur., 7, p. 63 (1829) (*Scopula*); 10 (3), p. Bucchara,
11 (1835) (*Scopula*). — Duponchel, Hist. Armenien,
Nat. Lep., 8 (2), p. 94, t. 215, f. 4 (1831) N.-Persien,
(*Scopula*). — Stephens, Ill. Brit. Ent. Haust., Amurgebiet,
4, p. 55 (1834). — Wood, Index Entom., Japan
p. 126, t. 28, f. 826 (1839) (*Margaritia*). —
Duponchel, Cat. Méth. Lep. Eur., p. 205
(1845) (*Lamia*). — Herrich-Schaeffer, Syst.
Bearb. Schm. Eur., 4, p. 27, f. 17, 18 (1848)
(*Botys*). — Guenée, Delt. et Pyral., p. 401
(1854) (*Lemiodes*). — Lederer, Wien. Ent.
Monatschr., 7, p. 382 (1863). — Heine-
mann, Kleinschm., 1, H. 2, p. 92 (1865). —
Wallengren, Oefvers. K. Vet.-Akad. Förh.,
28 (8), p. 1000 (1871). — Berce, Fauna
Ent. Fr. Lep., 6, p. 187, t. 5, f. 1 (1878).
— Snellen, Vlinders Nederl., 2 (1), p. 63
(1882). — Leech, Brit. Pyral., p. 41, t.
5, f. 7 (1886). — Meyrick, Trans. Ent.
Soc. Lond., p. 449 (1890); Handb. Brit.
Lep., p. 411 (1895). — Hampson, Trans.
Ent. Soc. Lond., p. 220 (1897). — Rebel
in Spuler, Schm. Eur., 2, p. 222 (1906).
— Wahlgren, Svensk Insektauna, 10 (2),
p. 49, t. 2, f. 56 (1915). — Meyrick, Revis.
Handb. Brit. Lep., p. 427 (1928). — Shi-
buya, Ins. Matsum., 3, p. 134, t. 5, f. 14
(1929). — M. Hering in Brohmer & Ehr-
mann & Ulmer, Tierwelt Mitteleur. (Ergbd.
1), p. 284 (1932). — Eckstein, Schm.
Deutschl., 5, p. 63 (1933).
fa. rubetra (D. Lucas) Lhomme, Cat. Lep. Frankreich
France Belg., 2 (1), p. 104 (1935).
var. grisealis Staudinger, Hor. Ent. Ross.,
7, p. 193, t. 2, f. 10 (1870) (*Botys pulveralis*
var.). — Hampson, Trans. Ent. Soc. Lond.,
p. 220 (1897). — Rebel in Spuler, Schm.
Eur., 2, p. 222 (1906).
Biologie: Disqué, Ent. Zeit. Stettin, 51, p.
57 (1890). — Rebel in Spuler, Schm. Eur.
Nachtr.-Taf. 9, f. 56 (1906).
Andalusien,
Griechen-
land,
Trans-
sylvanien,
Bulgarien,
Albanien,
Mazedonien,
Südrußland,
Armenien,
Kleinasien,
Sajangebiet,
Alai

- hyalinalis** Hübner, Samml. Eur. Schm. 6. Horde, p. 23, t. 11, f. 74 (1796). — Schrank, Fauna Boica, 2 (1), p. 62 (1802). — Hübner, Verz. bek. Schmett., p. 355, n. 3397 (1816) [1826] (*Hapalia*). — Treitschke, Schm. Eur., 7, p. 100 (1829) (*Botys*); 10 (3), p. 23 (1835) (*Botys*). — Duponchel, Hist. Nat. Léop., 8 (2), p. 119, t. 217, f. 3 (1831) (*Botys*). — Stephens, Ill. Brit. Ent. Haust., 4, p. 50 (1834). — Wood, Index Entom., p. 125, t. 28, f. 814 (1839) (*Margaritia*). — Eversmann, Fauna Lep. Volgo-Ural., p. 458 (1844). — Herrich-Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 28 (1848). — Guenée, Delt. et Pyral., p. 336 (1854) (*Botys*). — Stainton, Man. Brit. Butt. Moths, 2, p. 149 (1859) (*Botys*). — Heinemann, Kleinschm., 1, H. 2, p. 71 (1865) (*Botys*). — Berce, Fauna Ent. Fr. Lep., 6, p. 132, t. 4, f. (1878). — Snellen, Vlinders Nederl., 2 (1), p. 49 (1882). — Meyrick, Handb. Brit. Lep., p. 411 (1895). — Hampson, Trans. Ent. Soc. Lond., p. 220 (1897). — Rebel in Spuler, Schm. Eur., 2, p. 222, t. 82, f. 42 (1906). — Wahlgren, Svensk Insektafauna, 10 (2), p. 49 (1915). — Mitterberger, Zeitschr. Wiss. Insektenbiol., 11, p. 175 (1915). — Turati, Atti Ital. Sci. Nat., 63, p. 146, t. 5, f. 38, (1924). — Meyrick, Revis. Handb. Brit. Lep., p. 428 (1928). — M. Hering in Brohmer & Ehrmann & Ulmer, Tierwelt Mitteleur. (Ergbd. 1), p. 284 (1932). — Eckstein, Schm. Deutschl., 5, p. 63, t. 1, f. 23 (1933).
- Biologie: Hellins, Ent. Mon. Mag., 21, p. 99 (1884) (*Botys*). — Buckler, Larvae Brit. Lep., 9, p. 120, t. 153, f. 2 (1901). — Voukassovich, C. R. Soc. Biol., 96, p. 235—257 (1927) (Parasiten u. Hyperparasiten).
- decoloralis** Turati, Atti Soc. Ital. Sci. Nat., 63, p. 146, t. 5, f. 39 (♀) (1924). Cyrenaica

b) Orientalische Region.

- spissalis** Guenée, Delt. et Pyral., p. 326 (1854) (*Botys*) (Java). — Lederer, Wien. Ent. Monatschr., 7, p. 392 (1863) (*Pachynoa*). — Hampson, Proc. Zool. Soc. Lond., p. 198 (1899) (*Polygrammodes*). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 26, 168 (1928) (*Psammotis*). Indien, Java, Formosa

c) Australische Region.

- trygoda** Meyrick, Trans. Ent. Soc. Lond., p. 381 (♂) (1897). Queensland (Typus: Brisbane)

83. Genus **Eurrhyncha** Hübner

Hübner, Verz. bek. Schm., p. 360, n. 3451 (1816) [1826]. — Heinemann, Kleinschm., 1, H. 2, p. 57 (1865). — Wallengren, Oefvers. K. Vet.-Akad. Förh., 28 (8), p. 981, (1871). — Hampson, Trans. Ent. Soc. Lond., p. 134, 220 (1897). — Rebel in Spuler, Schm. Eur., 2, p. 222 (1906). — Wahlgren, Svensk Insektfauna, 10 (2), p. 49 (1915). — Forbes, Lep. N. York, p. 581 (1923). — Meyrick, Revis. Handb. Brit. Lep., p. 423 (1928). — M. Hering in Brohmer & Ehrmann & Ulmer, Tierwelt Mitteleur., (Ergbd. 1), p. 284 (1932). — Eckstein, Schm. Deutschl., 5, p. 63 (1933).

(Generotypus: *Eu. urticalis* Hbn. = *urticata* L.).

urticata Linné, Fauna Suecica, 2. ed., p. 340 (1761) Europa, Klein-
(*Geometra*). — Fabricius, Ent. Syst., 3 (2), Kleinasien,
p. 312 (1794). — Donovan, Nat. Hist. Brit. Armenien,
Ins., 10, p. 64, t. 349, f. 2 (1801) (*Phalaena*). Westsibirien,
— Stephens, Ill. Brit. Ent. Haust., 4, p. 44 Sajan- u.
(1834) (*Botys*). — Wood, Index Entom., p. Iligebiet,
123, t. 28, 805 (1839). — Stainton, Man. Brit. Altai,
Butt. Moths, 2, p. 150 (1859) (*Botys*). — Ussuri,
— Lederer, Wien. Ent. Monatschr., 7, p. 364 Amur,
(1863). — Heinemann, Kleinschm., 1, H. 2, West-
p. 58 (1865). — Wallengren, Oefvers. K. Zentral- u.
Vet.-Akad. Förh. 28 (8), p. 982 (1871). — Südchina;
— Berce, Fauna Ent. Fr. Lep., 6, p. 140, Nordamerika
t. 4, f. 4 (1878). — Leech, Brit. Pyral., 28, (Mac Nab's
t. 3, f. 8 (1886). — Meyrick, Handb. Brit. Ins., Nova
Lep., 407 (1895). — Hampson, Trans. Ent. Scotia)
Soc. Lond., p. 221, fig. (♂) (1897). —
— Leech, Trans. Ent. Soc. Lond., p. 444 (1901).
— Rebel in Spuler, Schm. Eur., 2, p. 222,
t. 81, f. 60 (1906). — Wahlgren, Svensk
Insektenfauna, 10 (2), p. 49, t. 3, f. 57
(1915). — Peters, Lep. Fauna Estland, 2.
Aufl., p. 400 (1924) (ab. melan.). — Forbes,
Lep. N. York, p. 581 (1923). — Parlin,
Maine Nat., 4, p. 56 (1924). — Comstock,
Introd. Ent., (ed. 2), p. 649 (1927). —
— Johnson, Psyche (Boston), 34, p. 176 (1927).
— Meyrick, Revis. Handb. Brit. Lep., p.
423 (1928). — M. Hering in Tierwelt Mittel-
eur. (Ergbd. 1), p. 284 (1932). — Eckstein,
Schm. Deutschl., 5, p. 63, t. 1, f. 24 (1933).

urticalis [Schiffermüller et Denis], Verz.
Schm. Wien, p. 121 (1776) (*Pyralis*). —
Hübner, Samml. Eur. Schm. 6. Horde, p. 21
(Pyr.), t. 12, f. 78 (1896). — Schrank, Fauna
Boica, 2 (1), p. 60 (1802). — Treitschke,
Eur. Schm., 7, p. 106 (1829) (*Botys*). —
Duponchel, Hist. Nat. Lep., 8 (2), p. 107,
t. 216, f. 3 (1831) (*Botys*). — Guenée, Delt.
et Pyral., p. 342 (1854) (*Botys*). — Herrich-
Schaeffer, Syst. Bearb. Schm. Eur., 4, p. 27
(1848) (*Botys*).

hortulata Linné, Syst. Nat., ed. 10, p. 529
(1758) (*Phalaena*). — Snellen, Vlinders Ne-
derl., 2 (1), p. 37 (1882).

ab. *dissoluta* Skala, Ent. Zeit., 42, p. 105 (1928).

Biologie: Treitschke, Schm. Eur., 7, p. 106 (1829). — Buckler, Larvae Brit. Lep., 9, p. 136, t. 153, f. 8 (1901). — Sorhagen, Allg. Zeitschr. Ent., 6, p. 243 (1901). — Rebel in Spuler, Schm. Eur., 2, Nachtr.-Taf., 9, f. 56 (1906). — Speyer, Arb. Biol. Reichsanst. Land- u. Forstwirt., 20, p. 190 (1932).

84. Genus **Mabra** Moore

Moore, Lep. Ceylon, 3, p. 280 (1885). — Hampson, Fauna Brit. India Moths, 4, p. 240 (1896); Trans. Ent. Soc. Lond., p. 134, 221 (1897).

Neurophruda Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 463 (1896) (Typus: *N. daulialis* Warr.).

Neophruda Hampson, Trans. Ent. Soc. Lond., p. 221 (1897) (ex synon.).

(Typus generis: *eryxalis* Walker)

a) Paläarktische Region.

eryxalis Walker, Cat. Lep. Het. Brit. Mus., 17, p. 371 (♂) (1859) (*Asopia*) (Ceylon). — Japan, Südcina, Tonkin, Indien, Birma, Assama, Formosa, Korea, Ceylon, Borneo, Singapur, Sumatra, Philippinen, Chagos Ins.
Moore, Lep. Ceylon, 3, p. 280, t. 179, f. 4 (1885) (*Mabra*). — Swinhoe, Trans. Ent. Soc. Lond., p. 277 (1890) (*Asopia*). — Hampson, Fauna Brit. India Moths, 4, p. 240, f. 145 (♂) (1896); Trans. Ent. Soc. Lond., p. 221, fig. (♂) (1897). — Swinhoe, Cat. East. Austral. Lep. Het. Oxford Mus., 2, p. 451 (1900). — E. Hering, Ent. Zeit. Stettin, 64, p. 49 (1903). — Shibuya, Journ. Fac. Agr. Hokkaido Imp. Univ., 22, p. 163 (1928); Ins. Matsum., 3, p. 134 (1929).
velatalis Snellen in Veth, Midden Sumatra, 4, Lep., p. 63, t. 5, f. 4, 4a (1880) (*Botys*).

b) Orientalische Region.

eryxalis Walker, vide sub a) Paläarktische Region, p. 191.
nigriscripta Swinhoe, Ann. Mag. Nat. Hist., (s. 6) 16, p. 302 (1895). — Hampson, Fauna Brit. India Moths, 4, p. 241 (1896); Trans. Ent. Soc. Lond., p. 221 (1897). Assam (Khasis), Süd-China
fauculalis Walker, Cat. Lep. Het. Brit. Mus., 19, p. 962 (♂) (1859) (*Hydrocampa*) (Sarawak). — Indien, Sikkim, Assam (Khasis), Borneo
Meyrick, Trans. Ent. Soc. Lond., p. 465 (1894) (*Stenia*). — Hampson, Trans. Ent. Soc. Lond., p. 221 (1897) (*Mabra*); Journ. Bombay Nat. Hist. Soc., 12, p. 704 (1899).
— Swinhoe, Ann. Mag. Nat. Hist., (s. 7) 17, p. 286 (1906).
fuscipennalis Hampson, Trans. Ent. Soc. Lond., p. 221 (♀) (1897); Journ. Bombay Nat. Hist. Soc., 12, p. 704 (♀) (1899). Assam (Typus: Khasis)

- daulialis** Warren, Ann. Mag. Nat. Hist., (s. 6) 17, p. 464 (1896) (*Neurophruda*); Journ. Bombay Nat. Hist. Soc., 12, p. 705 (♂) (1899); Trans. Ent. Soc. Lond., p. 222 (1897).
obealis Snellen, Tijds. v. Ent., 41 (1898), p. 181 (♂, ♀), t. 8, f. 11 (♀) (1899) (*Botys*). — E. Hering, Ent. Zeit. Stettin, 64, p. 49 (1903).
 Assam
 Sumatra,
 Java
 (Typus:
 Java,
 Tegal)

c) Neotropische Region.

- garzettalis** Felder et Rogenhofer, Reise Novara (Het.), 2 (2), p. 4, t. 135, f. 1 (1874) (*Blepharomastix*?).
garzettalis Hampson, Trans. Ent. Soc. Lond., p. 221 (1897).
metallescens Felder et Rogenhofer, Reise Novara (Het.), 2 (2), p. 4, t. 137, f. 58 (1874) (Botys?). — Hampson, Trans. Ent. Soc. Lond., p. 222 (1897).
 Brasilien
 Kolombien
 (Bogotá)

9

85. Genus **Neomabra** Dognin

Dognin, Ann. Soc. Ent. Belg., 49, p. 67 (1905).

- nitidalis** Dognin, Ann. Soc. Ent. Belg., 49, p. 67 (♂) (1905).
 Ecuador

1

86. Genus **Genoloba** Walsingham

Walsingham, Ent. Mon. Mag., 21, p. 176 (1885). — Meyrick, Ent. Mon. Mag., 25, p. 372 (1889). — Hampson, Trans. Ent. Soc. Lond., p. 134, 222 (1897); Journ. Bombay Nat. Hist. Soc., 21, p. 1262 (1912).

Agrammia Guenée, Delt. et Pyral., p. 405 (1854).
 (Typus generis: *obliteralis* Walk.)

a) Australische Region.

- obliteralis** Walker, Cat. Lep. Het. Brit. Mus., 30, p. 945 (1864) (*Pterophorus*). — Walsingham, Ent. Mon. Mag., 21, p. 176 (1885) (ad *Pterophorides* adfert.). — Hampson, Trans. Ent. Soc. Lond., p. 222, fig. (♂) (1897).
 Australien

b) Orientalische Region.

- cuprescens** Hampson, Ann. Mag. Nat. Hist., (s. 8) 20, p. 279 (♀) (1917).
taprobana Hampson, Journ. Bombay Nat. Hist. Soc., 21, p. 1762 (1912).
 Sumbawa
 Ceylon

c) **Neotropische Region.**

- iridalis** Guenée, Delt. et Pyral., p. 406 (1854) Cayenne
(*Agrammia*).
matronalis Guenée, Delt. et Pyral., p. 405 (1854) Brasilien
(*Agrammia*).
5

87. Genus **Homophysodes** Dyar

- Dyar, Proc. U. S. Nat. Mus., 47, p. 296 (1914).
morbida Dyar, Proc. U. S. Nat. Mus., 47, p. 296 Panama
(1914).
1

88. Genus **Escandia** Dyar

- Dyar, Proc. U. S. Nat. Mus., 47, p. 297 (1914).
fimbrialis Dyar, Proc. U. S. Nat. Mus., 47, p. 297 Panama
(♀) (1914).
1

Genera incertae sedis.

89. Genus **Cilaus** De Joannis

- De Joannis, Livre Centenn. Soc. Ent. Fr., p. 449 (1932).
longinasus De Joannis, Livre Centenn. Soc. Ent. Fr., Reunion
p. 450, t. 23, f. 19 (♀) (1932).
1

90. Genus **Clepsicosma** Meyrick

- Meyrick, Trans. Proc. N. Zeal. Inst., 20, p. 63 (1888); 45, p. 45
(1912) (1913). — Hudson, Butt. Moths N. Zeal., p. 204 (1928).
iridia Meyrick, Trans. Proc. N. Zeal. Inst., 20, p. 64 Neuseeland
(1888). — Hudson, Butt. Moths N. Zeal.,
p. 204, t. 24, f. 22 (♂) (1928).
1

91. Genus **Eustenia** Snellen

- Snellen, Tijds. v. Ent., 42, p. 79 (1899).
acuminatalis Snellen, Tijds. v. Ent., 42, p. 80, t. 4, W.-Java
f. 6, 7 (♂, ♀) (1899). (Buitenzorg);
1 Saleijer
bei Celebes

92. Genus **Hylebafis** Turner

- Turner, Trans. Roy. Soc. S. Austral., 32, p. 84 (1908).
scintillifera Turner, Trans. Roy. Soc. S. Austral., Queensland
32, p. 84 (♂, ♀) (1908).
1

93. Genus **Malaciofis** Meyrick

Meyrick, Exot. Microlep., 4, p. 541 (1934).

thiogramma Meyrick, Exot. Microlep., 4, p. 541 Belg. Congo
(♀) (1934). (Typus:
1 Elisabeth-
ville)

94. Genus **Nannomorpha** Turner

Turner, Trans. Proc. Roy. Soc. S. Austral., 32, p. 80 (1908).

alychnopa Turner, Trans. Proc. Roy. Soc. S. Queensland
Austral., 32, p. 80 (♀) (1908).
1

95. Genus **Pseudoligosigma** Strand

Strand, Arch. Naturg., 85 A, H. 12, p. 132 (1920).

incisa Strand, Arch. Naturg., 85 A, H. 12, p. 132 Trinidad
(♂) (1920).
1

96. Genus **Tholerasfis** Turner

Turner, Proc. Roy. Soc. Queensland, 27, p. 36 (1915).

elaphra Turner, Proc. Roy. Soc. Queensland, 27, Queensland
p. 36 (♀) (1915).
1
1229

Addenda**Scoparia** Haworth (p. 10)

phycophanes Meyrick, Exot. Microlep., 5, p. 110 (♂) Irak
(1937). (Rawaduz
Gorge)

rupestris Meyrick, Exot. Microlep., 5, p. 109 (♂, ♀) Irak
(1937). (Rayat)

Argyracfis Hampson (p. 60)

dorophanes Meyrick, Exot. Microlep., 5, p. 101 Belg. Congo
(♀) (1937). (Eala)

scioxantha Meyrick, Exot. Microlep., 5, p. 101 (♂) Belg. Congo
(1937). (Elisabeth-
ville)

sphragidaema Meyrick, Exot. Microlep., 5, p. 102 Belg. Congo
(♀) (1937). (Elisabeth-
ville)

Index.

Genera, Subgenera.

- Aediodes* Lederer 180
Aediodina Strand 180
Agrammia Guenée 192
Almonia Walker 148
Alutefa Swinhoe 180
Amaurophanes Lederer 163
Ambia Hampson 110
Ambia Walker 110
Anarpio Chapman 11
Angonia Snellen 59
Anydraula Meyrick 94
Argyractis Hampson 60, 194
Arnia Guenée 163
Arxama Walker 71
Aracomorpha Turner 93
Aulacodes Guenée 139
Aulacodes Hampson 138
Auxolophotis Meyrick 127

Badinomorpha Matsumura 107
Baeoptila Turner 106
Blepharomastix Druce 127
Bradina Lederer 149
Brevicella Kenrick 127

Camptomastix Warren 181
Camptomastix Hampson 181
Cataclysta Hübner 94
Cataclysta Hampson 60, 107
Ceratoclasia Druce 127
Chnaura Lederer 139
Cholius Guenée 9
Chrysendeton Grote 94
Cilaus De Joannis 193
Clepsicosma Meyrick 193
Clupeosoma Hampson 182
Clupeosoma Snellen 182
Coelorrhynchidia Hampson 158
Cometura Meyrick 146
Coptobasis Lederer 157
Cosmophylla Turner 106
Crambus Moore 59
Cymoriza Guenée 107
Cymorrhiza Hering 107

Danaga Moore 171
Dasyseopa Meyrick 57

Daulia Walker 126
Decticogaster Snellen 146
Deuterophysa Warren 162
Diathrausta Fekler [& Rogenhofers] 104
Diathrausta Lederer 159
Diathraustodes Hampson 158
Dipleurina Chapman 11
Diplotyla Meyrick 171
Dodanga Moore 148
Dracaenura Meyrick 135
Duponchelia Zeller 133

Eclipsiodes Meyrick 8
Ellogima Turner 181
Elophila Dyar 94
Ecöphila Swinhoe 137
Ephormotris Meyrick 109
Epichromistis Hampson 149
Epichronistis Meyrick 149
Epinima Meyrick 107
Eranistis Meyrick 133
Erebangela Meyrick 171
Erlita Lederer 149
Eristena Warren 70
Epistenia Chrétien
Escandia Dyar 193
Eudorea Curtis 11
Eudoria Chapman 11
Eugauria Snellen 107
Eurrhynpara Hübner 190
Eurytorna Meyrick 171
Eustenia Snellen 193

Galadra Walker 187
Gargela Walker 59
Genoloba Walsingham 192
Gethosyne Warren 122
Girtexia Snellen 126
Geshna Dyar 159
Glanecoda Karsch 180
Glyphandra Karsch 187
Graphycopoda Butler 171

Hapalia Hübner 188
Hemiloba Swinhoe 116
Heringia v. Hedemann 180

- Heringiella* Berg 180
Homophysodes Dyar 193
Hormatholepis Butler 171
Hyaloplaga Warren 187
Hydreuretis Meyrick 72
Hydrocampa Latreille 71
Hydrocampus Berthold 72
Hydrophysa Guenée 139
Hydropionea Hampson 182
Hydrorybina Hampson 182
Hygraula Meyrick 72
Hylebatus Turner 193
Hymenoptychis Zeller 137

Kasania Krulikovsky 72

Lasiogyia Hampson 162
Lemia Duponchel 188
Lemiodes Guenée 188
Leucogephyra Warren 110
Lissophanes Warren 11
Loxocoryx Meyrick 129
Luma Hampson 129
Luma Walker 130

Mabra Moore 191
Malaciotis Meyrick 194
Margarochroma Warren 126
Margarosticha Lederer 93
Massepha Walker 127
Mestolobes Butler 55
Metaclysta Hampson 104
Metoea Warren 72
Micraglossa Warren 8
Microglossa Hampson 8
Microphysetica Hampson 162
Mixophila Hampson 59
Mixophyla Meyrick 59
Musotima Meyrick 104

Nannomorpha Turner 194
Nausinoe Hübner 72
Neogenesis Hampson 180
Neomabra Dognin 192
Neophruda Hampson 191
Neurophruda Warren 191
Nicaria Snellen 157
Niphostola Hampson 132
Nymphaecella Grote 72
Nymphula Schrank 71

Oligernis Meyrick 110
Oligostigma Guenée 116
Oligostigma Walker 72
Omphaloptera Hampson 108
Opistheidicta Warren 110
Orphnophanes Lederer 131

Osphrantis Meyrick 136
Oxyelophila Forbes 60

Panotima Meyrick 106
Paracymoriza Warren 122
Parambia Dyar 109
Paraponyx Guenée 72
Paraponyx Hübner 72
Parthenodes Guenée 122
Pelena Moore 129
Pelina Hampson 129
Pelinopsis Dognin 130
Perinephela Hübner 185
Perinephele Hübner 185
Perinephila Hampson 185
Perinephile Barrett 185
Perisyntrocha Meyrick 132
Physematia Lederer 149
Physematia Snellen 127
Piletocera Lederer 171
Pleonectusa Lederer 149
Promylaca Meyrick 55
Psammotis Hampson 185
Psammotis Hübner 187
Pseudoligostigma Strand 194
Pseudolithosia Hampson 108
Ptilaeola Meyrick 171
Pythagorea Meyrick 136

Ridleyana Hampson 122
Rinecera Butler 171

Scoparia Haworth 10, 194
Scoparona Chapman 11
Scopea Haworth 10
Sozoa Walker 163
Stegothyris Lederer 134
Stenia Duponchel 163
Stenia auct. 162
Stenacula Snellen 122
Steniodes Snellen 164
Strepsimela Meyrick 171
Strepsinoma Meyrick 94
Strepsinoma Meyrick 133
Sufetula Druce 182
Susumia Marumo 107
Symmoracma Meyrick 164
Symphonia Hampson 104
Synclita Lederer 72
Syngamia Wileman 159
Syntomodora Meyrick 131
Syntrita Dognin 157
Syrbatis Walker 137

Tatobotys Butler 146
Temnobasis Gaede 158
Teratauxta Hering 122
Termatarcha Meyrick 149

Tetraprosopos Butler 55
 Tetrernia Meyrick 109
 Theila Swinhoe 137
 Tholerastis Turner 194
 Thysanoidma Hampson 109

Witlesia Chapman 11
 Zagiridia Hübner 182
 Zebronia Leech 186
 Zebronia Walker 59

Species, Subspecies, Varietates, Aberrationes, Formae.

- aaronalis Schaus (*Bradina*) 157
 abispalis (ab.) Dyar (*Scoparia*) 34
 abnormis Butler (*Scoparia*) 56
 abrotalis Walker (*Nymphula*) 93
 absconditalis Christoph (*Scoparia*) 18
 absolutalis Walker (*Massepha*) 127
 acclalaris Walker (*Musotima*) 105
 accra Strand (*Argyrectis*) 62
 acealis Walker (*Pileocera*) 174
 acharis Meyrick (*Scoparia*) 46
 acleropa Meyrick (*Scoparia*) 57
 aclistalis Dyar (*Argyrectis*) 63
 acolias Meyrick (*Scoparia*) 39
 acompa Meyrick (*Scoparia*) 50
 acrias Meyrick (*Musotima*) 105
 acrocampa Turner (*Eclipsiodes*) 9
 acroperalis Hampson (*Aulacodes*) 142
 acropola Meyrick (*Scoparia*) 42
 acrospila Meyrick (*Bradina*) 153, 154
 actias Meyrick (*Scoparia*) 38
 actoralis Walker (*Cataclysta*) 97
 acuminalis Dyar (*Stenia*) 168
 acuminatalis Snellon (*Eustenia*) 193
 acuminatella Stainton (*Scoparia*) 28
 adela Tams (*Dracaenura*) 135
 adhaesalis Walker (*Bradina*) 151
 adiantalis Walker (*Nymphula*) 92
 adjunctalis Snellen (*Oligostigma*) 119
 adjutrealis Schaus (*Oligostigma*) 146
 admixtalis Matsunura (*Susumia*) 107
 admixtalis Walker (*Bradina*) 150, 152, 153, 154
 aduncalis Felder & Rogenhofer (*Musotima*) 105
 adustalis Felder & Rogenhofer (*Stenia*) 168
 aelis Druce (*Argyrectis*) 67
 aelis Walker (*Argyrectis*) 63
 aechmialis Guenée (*Aulacodes*) 144
 aeclusalis Walker (*Bradina*) 151
 aegialitis Meyrick (*Dracaenura*) 136
 aegimiusalis Walker (*Pileocera*) 172, 173, 177
 aeglesalis Walker (*Argyrectis*) 62
 aenealis Hampson (*Micraglossa*) 8
 aengusalis Schaus (*Argyrectis*) 65
 aequalis Walker (*Alutefa*) 180
 aequipennalis Warren (*Scoparia*) 29
 aequivocalis Warren (*Parthenodes*) 124
 aetalis Walker (*Nymphula*) 91
 aetnealis Duponchel (*Stenia*) 164
 affinalis Guenée (*Nymphula*) 77, 84, 86, 87
 affinis W. Rothschild (*Bradina*) 154
 afralis Walker (*Daulia*) 126
 afrella Turati (*Stenia*) 165
 africalis Hampson (*Parthenodes*) 124
 agana Meyrick (*Scoparia*) 51
 agathanalis Schaus (*Pileocera*) 179
 agramma Meyrick (*Dracaenura*) 135
 agraphalis Guenée (*Bradina*) 156
 alaicalis (var.) Caradja (*Nymphula*) 77
 alaskalis Barnes & Benjamin (*Scoparia*) 35
 alba Tutt (*Scoparia*) 18
 albalis Robinson (*Nymphula*) 90
 albertalis Dyar (*Scoparia*) 33
 albescens Rebel (*Pileocera*) 178
 albibasalis Hampson (*Ambia*) 115
 albibasalis Schaus (*Argyrectis*) 64
 albiceps Hampson (*Parthenodes*) 125
 albicillialis Hampson (*Pileocera*) 179

- albicinctata Hampson (*Pileocera*) 173, 178
 albicomma Swinhoe (*Ambia*) 112
 albidentata Hampson (*Eugauria*) 107
 albidivisa Warren (*Oligostigma*) 117
 albifascialis Hampson (*Luma*) 130
 albifascialis Hampson (*Parthenodes*) 124
 albifascialis Hampson (*Parthenodes*) 124
 albiflavialis Hampson (*Ambia*) 113
 albiflavicalis Warren (*Ambia*) 111
 albifrons Druce (*Scoparia*) 36
 albifusalis Hampson (*Scoparia*) 30
 albifuscalis Hampson (*Oligostigma*) 119
 albigenitalis Hampson (*Bradina*) 155
 albilunata Warren (*Pileocera*) 173
 albiaculalis W. Rothschild (*Strepsinoma*) 133
 albimixtalis Hampson (*Pileocera*) 177
 albioculalis Hampson (*Symphonia*) 104
 albipictalis Hampson (*Pileocera*) 176
 albiplagiialis W. Rothschild (*Strepsinoma*) 133
 albipunctalis Hampson (*Argyrectis*) 61
 albipunctalis Warren (*Baeoptila*) 106
 albipunctata Druce (*Scoparia*) 36
 albisignalis Hampson (*Orphnophanes*) 131
 albisinuatella Fernald (*Scoparia*) 12
 albitessellalis Hampson (*Ambia*) 115
 albivenalis Hampson (*Decticogaster*) 147
 albiventralis Hampson (*Pileocera*) 176
 albobasalis Caradja (*Ambia*) 112
 albomaculalis Hampson (*Ambia*) 113
 albonigralis Hampson (*Druacynura*) 136
 albulalis Hampson (*Argyrectis*) 68
 alemundalis Schaus (*Argyrectis*) 66
 algiralis Lucas (*Nymphula*) 80
 alialis Barnes & Mc Dunnough (*Scoparia*) 35
 alicialis Hampson (*Oligostigma*) 116
 alienalis Walker (*Perisyntrocha*) 132
 allicola Meyrick (*Scoparia*) 32
 allionealis Walker (*Nymphula*) 89, 92
 alluminialis Turati (*Stenia*) 166
 alopecias Meyrick (*Scoparia*) 51
 alpina Curtis (*Scoparia*) 21
 alvealis Felder & Rogenhofer (*Argyrectis*) 67
 alychnopa Turner (*Nannomorphia*) 194
 amana (subsp.) Osthelder (*Nymphula*) 79
 amaura Meyrick (*Strepsinoma*) 133
 ambialis Hampson (*Massepha*) 128
 ambialis Schaus (*Stenia*) 170
 ambigualis Hampson (*Scoparia*) 13
 ambigualis Hampson (*Scoparia*) 15, 16
 ambigualis Rebel (*Scoparia*) 20
 ambiguelia Treitschke (*Scoparia*) 14
 amboinalis Hampson (*Cataclysta*) 97
 amethystias Meyrick (*Mestolobes*) 55
 amethystina Schaus (*Argyrectis*) 63
 amissella Millièrre (*Scoparia*) 29
 amoenialis Christoph (*Stenia*) 165
 amoyalis Caradja (*Decticogaster*) 147
 amoyalis Caradja (*Ambia*) 112
 amphilicella Meyrick (*Scoparia*) 37
 anadonta Dyar (*Scoparia*) 37
 anagantis Dyar (*Scoparia*) 36
 analtodes Hampson (*Pileocera*) 178
 anaplecta Meyrick (*Scoparia*) 42
 anatolica Caradja (*Scoparia*) 13
 ancipitalis Zeryy (*Scoparia*) 23
 ancipitella De la Harpe (*Scoparia*) 14
 andeolalis Schaus (*Oligostigma*) 121
 andresi Rebel (*Bradina*) 149
 andreusialis Hampson (*Oligostigma*) 117
 angularis Hampson (*Parthenodes*) 125

angulata Moore (*Cataclysta*) 97
angulatalis Lederer (*Cataclysta*) 99
angulipenne Hampson (*Oligostigma*) 116
angulipennis Hampson (*Oligostigma*) 116
angusta Butler (*Bradina*) 155
angustalis (subsp.) Caradja (*Cataclysta*) 98
angustalis Caradja (*Dectiogaster*) 147
angustalis Sauber (*Oligostigma*) 119
angustea Stephens (*Scoparia*) 28
angustella (fa.) Dyar (*Dithrausta*) 161
anialis Walker (*Perisynthrocha*) 132
anicitalis Schaus (*Cataclysta*) 103
animosa Meyrick (*Scoparia*) 46
anisophragma Lower (*Scoparia*) 43
anissella Millière (*Scoparia*) 26
annulalis Dognin (*Stenia*) 170
annulalis Guenée (*Argyrectis*) 66
antarcticalis Staudinger (*Scoparia*) 37
antenoralis Walker (*Bradina*) 156
anthomera Lower (*Scoparia*) 43
anthracias Meyrick (*Scoparia*) 41
anticalis Walker (*Luma*) 130
antichora Meyrick (*Mestolobes*) 57
antimacha Meyrick (*Scoparia*) 38
antisema Meyrick (*Bradina*) 155
apheles Meyrick (*Scoparia*) 53
aphenice Hampson (*Stenia*) 170
aphrias Meyrick (*Mestolobes*) 56
aphrodes Meyrick (*Scoparia*) 42
apicalis Pagenstecher (*Gargela*) 60
aptalis Lederer (*Nymphula*) 89
araeale Hampson (*Oligostigma*) 119
arcta Lucas (*Scoparia*) 43
arctura Meyrick (*Mestolobes*) 56
arcualis Hampson (*Gargela*) 60
areolifera Strand (*Campomastix*) 181
arfakalis Swinhoe (*Dracaenura*) 136
argentata Butler (*Bradina*) 155
argenteopicta Hampson (*Aulacodes*) 142
argentilinealis Hampson (*Argyrectis*) 69

argentimaculalis Hampson (*Aulacodes*) 143
argentistriata Hampson (*Ambia*) 115
argentuosalis Swinhoe (*Daulia*) 126
argillacea Butler (*Dectiogaster*) 146
argolina Lower (*Eclipsiodes*) 9
argopis Meyrick (*Pileocera*) 178
argylineale Hampson (*Oligostigma*) 120
argyractalis Schaus (*Ambia*) 115
argyrochrysalis Mabilie (*Cataclysta*) 98
argyrogaster Butler (*Dracaenura*) 135
argyrograpta Hampson (*Margarosticha*) 94
argyrolepta Dyar (*Cataclysta*) 103
argyrophora Hampson (*Argyrectis*) 66
argyrophoralis Hampson (*Daulia*) 126
argyropis Meyrick (*Aulacodes*) 141
argyrostrotalis Hampson (*Daulia*) 126
argyrotoxalis Pagenstecher (*Oligostigma*) 120
aristodera Turner (*Nymphula*) 88
arnoulalis Schaus (*Ambia*) 116
aroalis Schaus (*Argyrectis*) 63
arundinalis Eversmann (*Nymphula*) 80
asaleuta Meyrick (*Scoparia*) 49
asaphalis Schaus (*Ambia*) 115
asiusalis Walker (*Massepha*) 128
asphodeliella De la Harpe (*Scoparia*) 13
aspidota Meyrick (*Scoparia*) 53
astentialis Dyar (*Stenia*) 170
asterica Hampson (*Scoparia*) 54
asterisca Meyrick (*Scoparia*) 54
asthenota Meyrick (*Dracaenura*) 136
astragalota Meyrick (*Scoparia*) 52
astrigalis Hampson (*Clupeosoma*) 184
atmogramma Meyrick (*Scoparia*) 51
atmota Turner (*Aracomorpha*) 93
atomalis (var.) Stainton (*Scoparia*) 14
atopalis Walker (*Bradina*) 149, 151
atralis Hampson (*Arxama*) 71

- atralis Pagenstecher (*Stegothyris*) 134
 atrata Pagenstecher (*Piletocera*) 177
 atratalis W. Rothschild (*Almonia*) 148
 atricuprea Hampson (*Scoparia*) 35
 atristriata Hampson (*Clupeosoma*) 184
 atristrigalis Hampson (*Ambia*) 111
 atriterminalis Hampson (*Cataclysta*) 97
 atropicta Hampson (*Scoparia*) 35
 augastis Meyrick (*Scoparia*) 52
 aulacodealis Strand (*Eoöphila*) 138
 aulacodialis Strand (*Bradina*) 149
 aulacodoidalis W. Rothschild (*Strepsinoma*) 133
 aulacophora Hampson (*Ambia*) 113
 auralis Osthelder (*Nymphula*) 75
 aurantialis Hampson (*Daulia*) 126
 aurantialis Hampson (*Dectiogaster*) 147
 aurantialis Hampson (*Pileto-cera*) 174
 aurantialis Swinhoe (*Parthenodes*) 123
 aurantipennis Hampson (*Aulacodes*) 142
 aurata Butler (*Bradina*) 155
 aureolalis De Joannis (*Bradina*) 153
 aureolalis Snellen (*Aulacodes*) 139
 auropunctalis Hampson (*Oligostigma*) 119
 auspicatalis Schaus (*Argyrectis*) 64
 australis Felder & Rogenhofer (*Margarosticha*) 93
 australis Hulst (*Nymphula*) 91
 australis (fa.) Müller-Rutz (*Scoparia*) 18
 autobella Dyar (*Cataclysta*) 103
 autochroa Meyrick (*Scoparia*) 47
 autodoxa Meyrick (*Mestolobes*) 57
 autumnna Philpott (*Scoparia*) 52
 avernalis Grote (*Cataclysta*) 100
 avertinalis Schaus (*Nymphula*) 92
 axena Meyrick (*Scoparia*) 51
 axiolecta Turner (*Scoparia*) 43
 axis Hampson (*Argyrectis*) 65
 azadalis Schaus (*Cataclysta*) 103
 aztecalis Schaus (*Argyrectis*) 64
 bademusalis Schaus (*Stenia*) 169
 badiusalis Walker (*Nymphula*) 90
 bahrlutalis (subsp.) Amsel (*Epi-stenia*) 163
 balanopis Meyrick (*Scoparia*) 37
 balcanica (subsp.) Rebel (*Scoparia*) 22
 baltealis Mabille (*Stenia*) 167
 banausa Meyrick (*Mestolobes*) 56
 baracocalis Schaus (*Campomastix*) 181
 barbararcha Meyrick (*Oligostigma*) 120
 barbicornis Felder & Rogenhofer (*Pileto-cera*) 175
 barbipennis Hampson (*Scoparia*) 31
 barnesalis Dyar (*Clupeosoma*) 184
 basale Hampson (*Clupeosoma*) 184
 basalis Walker (*Pileto-cera*) 174
 basalis Walker (*Scoparia*) 34
 basilissa Meyrick (*Aulacodes*) 141
 basistrigalis Knaggs (*Scoparia*) 16
 batjanalis Strand (*Pileto-cera*) 173
 batrachalis Guenée (*Stenia*) 170
 bedealis Schaus (*Argyrectis*) 63
 benesignata (subsp.) Caradja (*Nymphula*) 75
 benetinctalis Dyar (*Stenia*) 170
 benezetalis Schaus (*Argyrectis*) 68
 bengalensis Meyrick (*Massepha*) 127
 benigna Meyrick (*Scoparia*) 32
 berthalis Schaus (*Parthenodes*) 125
 berytella Rebel (*Scoparia*) 28
 beuvealis Schaus (*Stenia*) 169
 biannulalis Walker (*Dectiogaster*) 147
 biannulalis Walker (*Stenia*) 170
 bicolor Moore (*Clupeosoma*) 183
 bicoloralis Hampson (*Bradina*) 152
 bifascialis Robinson (*Cataclysta*) 100
 bifascialis (fa.) Turati (*Scoparia*) 15
 bifurcale Hampson (*Oligostigma*) 119
 bifurcalis Pryer (*Oligostigma*) 119

- bifurcalis Wileman (*Nymphula*) 76, 83
 biincisalis Karsch (*Glyphandra*) 187
 bijonalis Dyar (*Cataclysta*) 103
 bilineale Hampson (*Oligostigma*) 117
 bilinealis Snellen (*Oligostigma*) 117
 bimaculalis Shibuya (*Pelena*) 129
 bimaculalis Snellen (*Margarosticha*) 93
 biplagialis Walker (*Scoparia*) 34
 bipunctalis Hampson (*Oligostigma*) 120
 bipunctalis Kenrick (*Aulacodes*) 143
 bipunctalis Walker (*Aulacodes*) 139
 biscutella Zeller (*Scoparia*) 36
 bisignalis Hampson (*Piletocera*) 178
 bisinualis Hudson (*Scoparia*) 49
 bistrigalis Hampson (*Piletocera*) 178
 bistrigulalis Walker (*Dectiogaster*) 148
 blandialis Hampson (*Cataclysta*) 95
 blandialis Walker (*Cataclysta*) 94, 97, 98
 bocusalis Walker (*Ambia*) 115
 bolusalis Walker (*Ambia*) 115
 bombayensis (fa.) Swinhoe & Cotes (*Cataclysta*) 95
 borealis Lefebvre (*Scoparia*) 11
 borealis Tengström (*Scoparia*) 21
 bosoralis Druce (*Aulacodes*) 145
 botydis Strand (*Zebronia*) 187
 branealis Schaus (*Stenia*) 169
 brevicans Dyar (*Clupeosoma*) 185
 brevifascialis Wileman (*Diatrausta*) 160
 briocusalis Schaus (*Parthenodes*) 126
 bronacbalis Schaus (*Cataclysta*) 103
 bronzalis Barnes & Mc Dunnough (*Scoparia*) 34
 bruguieralis Duponchel (*Epi-stenia*) 162
 brunnealis Hampson (*Aulacodes*) 142
 brunneodora Dyar (*Argyrectis*) 64
 brunneofusalis Hampson (*Clupeosoma*) 184
 brunneospersa (subsp.) Osthelder (*Cataclysta*) 96
 brunneosuffusa Hampson (*Argyrectis*) 66
 brunnilidalis Dyar (*Cataclysta*) 100
 bucolica Meyrick (*Scoparia*) 39
 bufalis Guenée (*Piletocera*) 179
 bufalis Hampson (*Piletocera*) 179
 bytinskiella (ab.) Schawerda (*Scoparia*) 19
 cabimalis Dyar (*Cataclysta*) 63
 cacalis (fa.) Dyar (*Diatrausta*) 161
 caclamalis Schaus (*Stenia*) 169
 caecalis Walker (*Scoparia*) 11
 caepiosalis Walker (*Parthenodes*) 125
 caeruleata Hampson (*Ambia*) 110
 caesoalis Walker (*Argyrectis*) 68
 caidalis Oberthür (*Duponchelia*) 134
 caliginosa Philpott (*Scoparia*) 54
 caliginosalis Walker (*Scoparia*) 12
 callichromalis Mabille (*Cataclysta*) 98
 callidryas Turner (*Musotima*) 105
 callista Forbes (*Argyrectis*) 62
 callistoalis Walker (*Aulacodes*) 145
 camalis Quaintance (*Nymphula*) 91
 camptoteles Hampson (*Oligostigma*) 118
 camptozonale Hampson (*Oligostigma*) 118, 120
 canalis Walker (*Bradina*) 151
 cancellalis Dyar (*Cataclysta*) 101
 candidalis Pagenstecher (*Oligostigma*) 121
 candidata Fabricius (*Nymphula*) 80
 canicostalis Hampson (*Scoparia*) 31
 cannalis Quaintance (*Geshna*) 159
 cantiusalis Schaus (*Ambia*) 114
 cantonalis (fa.) Caradja (*Nymphula*) 77
 canuialis Millièrè (*Duponchelia*) 134
 capensis Hampson (*Argyrectis*) 61
 carbonalis Warren (*Massepha*) 128
 carillalis Schaus (*Stenia*) 170
 carthagalis (var.) Schaus (*Stenia*) 169
 castusalis Schaus (*Argyrectis*) 67

- cataclystalis Hampson (*Ambia*) 112
cataclystalis (var.) Strand (*Parthenodes*) 122
 catactis Meyrick (*Scoparia*) 39
 cataxesta Meyrick (*Scoparia*) 39
 categorica Meyrick (*Pythagorea*) 136
 catenalis Guenée (*Argyrectis*) 65
 cathanalis Schaus (*Argyrectis*) 63
 caulealis Schaus (*Stenia*) 169
 cayugalis Schaus (*Stenia*) 169
 ceddalís Schaus (*Stenia*) 168
 cedroalis Schaus (*Parambia*) 109
 cembrae Hampson (*Scoparia*) 13
 cembrae Haworth (*Scoparia*) 12
 cembralis Guenée (*Scoparia*) 12
 cembrela Stephens (*Scoparia*) 12
 centrimacula Hampson (*Oligostigma*) 120
centurialis Guenée (*Scoparia*) 12
 centuriella Schiffermüller & Denis (*Scoparia*) 11, 32
 ceramica W. Rothschild (*Bradina*) 153
 ceramica Warren (*Scoparia*) 35
 ceramochra Meyrick (*Dectiogaster*) 147
 ceratucha Meyrick (*Aulacodes*) 139
cerrussalis Druce (*Argyrectis*) 66
cerrussalis Felder & Rogenhofer (*Argyrectis*) 67
cervalis Mc Dunnough (*Scoparia*) 34
 cervicalis Hampson (*Aulacodes*) 143
 chalara Meyrick (*Scoparia*) 54
 chalcialis Hampson (*Parthenodes*) 125
 chalcichroalis Hampson (*Ambia*) 113
 chalcistis Dognin (*Argyrectis*) 65
 chalcitis Felder & Rogenhofer (*Cataclysta*) 102
 chalcophora Meyrick (*Bradina*) 155
 chalicodes Meyrick (*Scoparia*) 49
 chalinota Meyrick (*Bradina*) 153
 characta Meyrick (*Scoparia*) 47
 charonialis Walker (*Stenia*) 165, 166
 charopaea Turner (*Scoparia*) 42
 chejelalis Schaus (*Argyrectis*) 64
 chiasta Meyrick (*Scoparia*) 42
 chimeria Meyrick (*Scoparia*) 46
 chimonias Meyrick (*Mestolobes*) 56
 chionostola Hampson (*Cataclysta*) 102
 chlamydota Meyrick (*Scoparia*) 45
 chlorionalis Tams (*Bradina*) 154
 chloronota Meyrick (*Pileocera*) 176
 chordactis Meyrick (*Scoparia*) 35
 choristis Meyrick (*Scoparia*) 48
 chorolychna Meyrick (*Mestolobes*) 56
 choroscia Meyrick (*Bradina*) 154
 chryseis Meyrick (*Nymphula*) 37
 chrysias Meyrick (*Gargela*) 60
 chrysippusalis Walker (*Nymphula*) 81
 chrysochroa Hampson (*Dracae-nura*) 135
 chrysogramma Hampson (*Ambia*) 113
 chrysomicta Meyrick (*Scoparia*) 43
 chrysomolybda Meyrick (*Mestolobes*) 55
 chrysomolybdoides Swezey (*Mestolobes*) 56
 chrysopalis Hampson (*Argyrectis*) 62
 chrysopetra Meyrick (*Scoparia*) 43
 chrysota Meyrick (*Oligostigma*) 120
 chrysoxantha Hampson (*Aulacodes*) 142
 chrysozonalis Hampson (*Oligostigma*) 119
 cilianalis Schaus (*Ambia*) 116
 cincta Walker (*Pileocera*) 172
 cincticorpus Hampson (*Dracae-nura*) 136
 cinefacta Philpott (*Scoparia*) 47
 cineralis Schaus (*Argyrectis*) 64
 cinereomedia Dyar (*Scoparia*) 33
 circealis Walker (*Nymphula*) 86
 circumdatalis Walker (*Perisintrocha*) 132
 cirrhophanes Meyrick (*Bradina*) 155
 ciserodes Meyrick (*Scoparia*) 50
 citrocossa Meyrick (*Scoparia*) 43
 citronalis Druce (*Aulacodes*) 145
 claudialis Dyar (*Cataclysta*) 101
 claudialis Walker (*Cataclysta*) 100, 101
 clavata Philpott (*Scoparia*) 52
 cleodoralis Walker (*Scoparia*) 42
 clerica Meyrick (*Scoparia*) 43
 climacusalis Schaus (*Argyrectis*) 67
 clonodes Meyrick (*Scoparia*) 40

- coarctalis* Guenée (*Scoparia*) 29
coarctata Zeller (*Scoparia*) 29
coecimaculalis Warren (*Scoparia*) 30
coenosalis Snellen (*Nymphula*) 86
coeruleotincta Hampson (*Scoparia*) 26
coesalis Hampson (*Argyrectis*) 68
collaris Walker (*Pileocera*) 172
colonalis Bremer (*Ambia*) 110
colonialis Guenée (*Aulacodes*) 139
coloralis Guenée (*Argyrectis*) 61
colpota Meyrick (*Scoparia*) 48
colubralis Druce (*Stenia*) 168, 169
colubralis Guenée (*Stenia*) 170
commercialis Meyrick (*Scoparia*) 43
commortalis Dyar (*Scoparia*) 32
compactalis Snellen (*Eugauria*) 107
completalis Dognin (*Omphaloptera*) 108
complicata Warren (*Ambia*) 111
complicatalis Dyar (*Cataclysta*) 103
conallalis Schaus (*Argyrectis*) 68
conchylia Meyrick (*Pileocera*) 175
concinella (ab.) Curtis (*Scoparia*) 28
concisalis Walker (*Pileocera*) 173
concoloralis Oberthür (*Stenia*) 165
concordalis Lederer (*Bradina*) 151
confinalis Caradja (*Pileocera*) 173
confirmata (var.) Krulikovsky (*Cataclysta*) 96
conflucella (fa.) Krulikovsky (*Scoparia*) 12
confusalis Schaus (*Aulacodes*) 146
confusalis Walker (*Cataclysta*) 100
confusella De la Harpe (*Scoparia*) 18
congallalis Schaus (*Aulacodes*) 144
congestalis Walker (*Scoparia*) 31
conicalis (var.) Warren (*Dia-thrausta*) 160
conicella De la Harpe (*Scoparia*) 14
conifera Butler (*Scoparia*) 48
coniferalis Hampson (*Aulacodes*) 143
conjunctalis Wileman & South (*Eoöphila*) 138
consortialis Caradja (*Bradina*) 152
conspicualis Hodgkinson (*Scoparia*) 15
conspurcatale Hampson (*Oligostigma*) 117
conspurcatalis Warren (*Oligostigma*) 117, 120
constellalis Hampson (*Argyrectis*) 65
contemplalis Walker (*Scoparia*) 32
contexta Philpott (*Scoparia*) 52
convatalalis Klunder (*Micraglossa*) 8
convoluta Hampson (*Aulacodes*) 145
copidosema Meyrick (*Panotima*) 106
coprista Meyrick (*Scoparia*) 13
coreulina Butler (*Nymphula*) 76
costalis Hampson (*Bradina*) 154
costalis Hampson (*Stenia*) 168
costalis Walker (*Stenia*) 168
costifascialis Hampson (*Aulacodes*) 142
costifascialis Hampson (*Pileocera*) 175
costimaculalis Warren (*Deuterophysa*) 162
costipunctalis Snellen (*Stenia*) 166
costipunctata Warren (*Pileocera*) 173
crambidalis Snellen (*Gargela*) 59
crassicornalis Guenée (*Aulacodes*) 139
crassiuscula Dyar (*Scoparia*) 36
crataca Meyrick (*Scoparia*) 39
crataegalis Duponchel (*Scoparia*) 28
crataegalis Guenée (*Scoparia*) 27
crataegella Herrich-Schaeffer (*Scoparia*) 27
crataegella Hübner (*Scoparia*) 26
craterotoxa Meyrick (*Bradina*) 155
cremealis Zerny (subsp.) (*Cholius*) 10
cretacealis Hampson (*Arxama*) 71
cretacealis Lederer (*Nymphula*) 89
cristalis Walker (*Nymphula*) 87
cristata Hampson (*Almonia*) 148
critica Meyrick (*Scoparia*) 47
crocalis Hampson (*Scoparia*) 30

- erocospila* Turner (*Scoparia*) 43
croesalis Meyrick (*Cataclysta*) 98
croesusalis Walker (*Cataclysta*) 98
cronanalis Schaus (*Stenia*) 168
cronialis Druce (*Argyrectis*) 66
cronialis Druce (*Cataclysta*) 101, 102
crossi (fa.) Banks (*Scoparia*) 15
crucigera Gerasimow (*Scoparia*) 15
cryerodes Meyrick (*Scoparia*) 39
crypterythra Lower (*Scoparia*) 43
cryptichrysa Meyrick (*Mestobes*) 56
cryptinoa Meyrick (*Scoparia*) 50
cryptixantha Meyrick (*Eclipsides*) 8
cryptale Hampson (*Oligostigma*) 121
cryptalis Druce (*Oligostigma*) 121
cubanalis (fa.) Dyar (*Diathrausta*) 161
cumalis Druce (*Cataclysta*) 102
cumulalis Hampson (*Piletocera*) 175
cunealis Walker (*Decticogaster*) 136
cuneifera Moore (*Cataclysta*) 95
cuneolalis Snellen (*Nymphula*) 86, 88
cuneolalis Snellen (*Perisyntrocha*) 132
cuprealis Hampson (*Gargela*) 60
cuprealis Hampson (*Eclipsiodes*) 9
cuprescens Hampson (*Argyrectis*) 66
cuprescens Hampson (*Coelorhynchidia*) 159
cuprescens Hampson (*Genoloba*) 192
cuprescens Hampson (*Massepha*) 128
cupritincta Hampson (*Micraglossa*) 8
curta Butler (*Nymphula*) 81
curviferalis Walker (*Nymphula*) 90
cyameuta Meyrick (*Scoparia*) 52
cyanealis Hampson (*Ambia*) 110
cyanolitha Meyrick (*Cataclysta*) 99
cyclophora Dyar (*Scoparia*) 37
cyclopila Meyrick (*Piletocera*) 176
cyclozonalis Hampson (*Aulacodes*) 141
cyloialis Schaus (*Argyrectis*) 64
cymatias Meyrick (*Scoparia*) 46
cymialis Hampson (*Diathrausta*) 161
cymophoralis Hampson (*Ambia*) 113
cyptastis Meyrick (*Scoparia*) 50
dabiusalis Schaus (*Parthenodes*) 125
dactyliopa Meyrick (*Scoparia*) 40
dactyloptila Meyrick (*Piletocera*) 178
daeckkealis Haimbach (*Diathrausta*) 161
daemonalis Dyar (*Cataclysta*) 101
daisensis Shibuya (*Susumia*) 107
damasalis Walker (*Bradina*) 149
damescalis Guenée (*Ambia*) 115
danaealis Hampson (*Argyrectis*) 62
dapsilis Turner (*Eclipsiodes*) 9
daralis Christoph (*Epistenia*) 163
dasymalis Dyar (*Cataclysta*) 103
daulialis Warren (*Mabra*) 192
debalis Druce (*Ambia*) 114
declivalis Dyar (*Stenia*) 168
declivis Philpott (*Scoparia*) 52
decoloralis Turati (*Psammotis*) 189
decoloris Dognin (*Argyrectis*) 70
decoloris Snellen (*Musotima*) 105
decorella Stainton (*Scoparia*) 23
decussalis Walker (*Nymphula*) 82
definitalis Strand (*Nymphula*) 85
defloralis (var.) Strand (*Bradina*) 151
delicata Moore (*Aulacodes*) 140
delicata Schaus (*Aulacodes*) 146
delicata Warren (*Diathrausta*) 161
delicatalis Walker (*Scoparia*) 30
deliniens Lucas (*Scoparia*) 43
delphinatalis Guenée (*Scoparia*) 22
delphusa Druce (*Scoparia*) 33, 37
deltophora Meyrick (*Scoparia*) 51
demodes Meyrick (*Scoparia*) 40
dendalis Druce (*Ambia*) 114
denigata Dyar (*Scoparia*) 32
denigrata (fa.) Costantini (*Scoparia*) 29
dentalis Hampson (*Bradina*) 152
dentalis Pagenstecher (*Coptobasis*) 158
dentata Druce (*Clupeosoma*) 184
dentatum Hampson (*Clupeosoma*) 184

- denticostalis Hampson (*Pileto-
cera*) 176
 dentifascialis Hampson (*Parthe-
nodes*) 123
 dentilinea Hampson (*Nymphula*)
 90
 dentilinealis Snellen (*Hymeno-
ptychis*) 137
 dentizonalis Hampson (*Nymphu-
la*) 87
 depalpalis Strand (*Decticogaster*)
 147
 depunctalis Guenée (*Nymphula*)
 81, 86, 88, 92
 desertalis Hampson (*Stenia*) 167
 desumptalis Guenée (*Bradina*)
 156
 diagonalis Guenée (*Stegothyris*)
 134
 dialitha Dyar (*Cataclysta*) 103
 dialitha Tams (*Cataclysta*) 99
 dianale Hampson (*Oligostigma*)
 118
 dianalis Schaus (*Nymphula*) 82
 dicentra Meyrick (*Nymphula*) 87
 dicteella Rebel (*Scoparia*) 19
 diffualis Snellen (*Nymphula*) 84
 dilucidalis Guenée (*Aulacodes*)
 141
 dilucidalis Walker (*Cataclysta*) 95
 diminutalis Snellen (*Nymphula*)
 85, 86
 dinodes Meyrick (*Scoparia*) 46
 diopsalis Hampson (*Aulacodes*)
 142
 diphtheralis Walker (*Scoparia*) 48
 diplopa Lower (*Nymphula*) 88
 discalis Hampson (*Oligostigma*)
 118
 discalis Hampson (*Pileto-cera*) 173
 discisignalis Hampson (*Piletoce-
ra*) 174
 dispar Gaede (*Nymphula*) 89
 dispersa Butler (*Scoparia*) 36
 dissipatalis Christoph (*Stenia*)
 166
 dissoluta (ab.) Skala (*Eurrhy-
para*) 191
 distictalis Hampson (*Micra-
glossa*) 8
 distincta (subsp.) W. Rothschild
 (*Aulacodes*) 143
 distinctalis Leech (*Parthenodes*)
 124
 distinctalis Ragonot (*Nymphula*)
 76
 distinctalis Snellen (*Nymphula*)
 92
 distributa Lucas (*Margarosticha*)
 94
 dives Butler (*Ambia*) 110
 divisalis Walker (*Cataclysta*) 100
 divulsalis Druce (*Argyrectis*) 68
 divulsalis Walker (*Cataclysta*)
 102
 dochmia Meyrick (*Scoparia*) 46
 dodalis Schaus (*Argyrectis*) 67
 doerriesi Strand (*Perinephela*)
 186
 dohrni E. Hering (*Ambia*) 112
 dolia (ab.) Tams (*Dracaenura*;
 135
 dolichoplagia Hampson (*Aulaco-
des*) 142
 dominalis Walker (*Aulacodes*) 140
 dominicalis Schaus (*Stenia*) 168
 dominulale Hampson (*Oligo-
stigma*) 120
 dominulalis Walker (*Oligostig-
ma*) 120
 dorophanes Meyrick (*Argyrectis*)
 194
 drosera Meyrick (*Eclipsiodes*) 8
 droseropa Meyrick (*Mestolobes*)
 56
 drumalis Dyar (*Cataclysta*) 100
 drusialis Lederer (*Cataclysta*) 97
 drusiusalis Walker (*Cataclysta*)
 97, 99
 dryphactis Meyrick (*Scoparia*) 52
 dualalis Gaede (*Argyrectis*) 62
 dubitalis Eversmann (*Scoparia*)
 18
 dubitalis Fryer (*Scoparia*) 18
 dubitalis Hübner (*Scoparia*) 17
 dubitalis Stephens (*Scoparia*) 14
 ducalis Schaus (*Oligostigma*) 121
 dulcis De Joannis (*Scoparia*) 32
 eanesalis Walker (*Pileto-cera*) 176
 echo Dyar (*Scoparia*) 34
 ectargyralis Hampson (*Parthe-
nodes*) 125
 ectogonalis Hampson (*Oligostig-
ma*) 120
 ectopalis Hampson (*Parthenodes*)
 124
 ectopalis Hampson (*Strepsinoma*)
 133
 effrenatalis Berg (*Nymphula*) 92
 ejuncida Knaggs (*Scoparia*) 53
 ekthlipsis Grote (*Nymphula*) 91
 elaphra Meyrick (*Scoparia*) 51
 elaphra Turner (*Tholerastis*) 194
 elathealis Walker (*Coelorrhynci-
dia*) 159
 electralis Dyar (*Oligostigma*) 121

- electralis* Walker (*Stenia*) 170
ellipes Tams (*Baeoptila*) 106
elongalis Warren (*Piletocera*) 175
elphегalis Schaus (*Ambia*) 115
emarginata Kenrick (*Brevicella*) 127
enmetropis Turner (*Scoparia*) 44
empeda Meyrick (*Scoparia*) 40
enallassalis Dyar (*Ambia*) 116
enantialis Dyar (*Ambia*) 116
enarexalis Dyar (*Ambia*) 116
encapna Meyrick (*Scoparia*) 47
encausta Meyrick (*Scoparia*) 44
endopthalma Meyrick (*Ambia*) 111
endoralis Druce (*Argyrectis*) 69
endoralis Walker (*Nymphula*) 92
endosaris Meyrick (*Aulacodes*) 139
enixalis Swinhoe (*Nymphula*) 78, 85
enneaspila Meyrick (*Piletocera*) 178
entephiadia Hampson (*Massepha*) 128
epicomia Meyrick (*Scoparia*) 50
epicremna Meyrick (*Scoparia*) 54
epieryma Meyrick (*Scoparia*) 42
epidelta Meyrick (*Mestolobes*) 56
epigypsa Lower (*Eclipsiodes*) 9
epinochla Turner (*Nymphula*) 83
epimystis Meyrick (*Scoparia*) 40
epipercialis Hampson (*Piletocera*) 177
episbila Meyrick (*Nymphula*) 77
erebina Butler (*Piletocera*) 177
erebochalea Meyrick (*Scoparia*) 40
eremitis Meyrick (*Scoparia*) 42
ergatis Meyrick (*Scoparia*) 47
erilitalis Felder & Rogenhofer (*Bradina*) 154
erilitoides Strand (*Bradina*) 153
erminea Meyrick (*Gargela*) 59
ermineus Moore (*Gargela*) 59
eromenalis Snellen (*Ambia*) 111
erralis Guenée (*Scoparia*) 14
erralis Hampson (*Scoparia*) 12
erynuis Meyrick (*Mestolobes*) 56
eryxalis Walker (*Mabura*) 191
esperanzalis Schaus (*Argyrectis*) 64
euceralis Meyrick (*Orphnophanes*) 131
eucerasalis Hampson (*Orphnophanes*) 131
eucerusalis Walker (*Orphnophanes*) 131
euclidialis Hampson (*Cataclysta*) 98
eugethes Tams (*Parthenodes*) 125
eumeles Meyrick (*Scoparia*) 42
eumoros Dyar (*Hydropionea*) 182
euprepialis Hampson (*Margarosticha*) 93
eupselias Meyrick (*Aulacodes*) 143
eurylea Meyrick (*Mestolobes*) 56
euryscia Meyrick (*Nymphula*) 88
euryxantha Meyrick (*Argyrectis*) 62
eutaeta Turner (*Scoparia*) 44
excisa Swinhoe (*Oligostigma*) 117
excisalis Snellen (*Oligostigma*) 119
excursalis Dyar (*Scoparia*) 37
exhibitale Hampson (*Oligostigma*) 121
exhibitilis Walker (*Oligostigma*) 121
exhibitilis Walker (*Scoparia*) 42
exigua Butler (*Susumia*) 107
exilis Knaggs (*Scoparia*) 51
expallidalis Dyar (*Scoparia*) 32
expatrialis Hampson (*Nymphula*) 87
exsolvalis Snellen (*Parthenodes*) 123
exsolvalis Snellen (*Parthenodes*) 123
extenuatalis Walker (*Bradina*) 152
externinata Meyrick (*Scoparia*) 44
externalis Walker (*Bradina*) 156
extincta Dyar (*Scoparia*) 36
exuvialis Snellen (*Campomastix*) 181
falcatalis Snellen (*Oligostigma*) 119
falsa Philpott (*Scoparia*) 52
famularis Philpott (*Scoparia*) 53
fascialis Warren (*Cymoriza*) 108
fasciata (ab.) Teich (*Nymphula*) 79
fasciata (fa.) Turati (*Scoparia*) 19
fasciculalis Zeller (*Stegothyris*) 134
fauculalis Walker (*Mabura*) 191
faulalis Walker (*Nymphula*) 91
favilliferella Walker (*Scoparia*) 44
fenestralis Barnes & Mo Dunning (*Hydropionea*) 182
fenestralis Warren (*Symphonia*) 104

- fenghwanalis Pryer (*Nymphula*) 76, 84
 feredayi Knaggs (*Scopula*) 50
 fernaldalis (ab.) Dyar (*Scoparia*) 34
 ferrealis Hampson (*Stenia*) 165
 fiachmalis Schaus (*Aulacodes*) 145
 filigeralis Walker (*Nymphula*) 92
 fimbrialis Dyar (*Escandia*) 193
 fimbriata Philpott (*Scoparia*) 50
 finbaralis Schaus (*Bradina*) 157
 flammealis Hampson (*Coelorrhynchidia*) 159
 flavalis Hampson (*Luma*) 130
 flavalis Hampson (*Perisyntrocha*) 132
 flavalis Hampson (*Piletocera*) 176
 flavalis Warren (*Ambia*) 115
 flavialbalis Hampson (*Oligostigma*) 120
 flavialis Schaus (*Nymphula*) 93
 flavicaput Snellen (*Parthenodes*) 124
 flavicostalis Hampson (*Nymphula*) 92
 flavidalis Hampson (*Micraglossa*) 8
 flavidiscalis Hampson (*Piletocera*) 175
 flavifascialis Hampson (*Aulacodes*) 143
 flavimaculata Gacde (*Massepha*) 128
 flavimarginale Hampson (*Oligostigma*) 121
 flavimarginalis Hampson (*Luma*) 130
 flavimarginalis Warren (*Oligostigma*) 121
 flavipictalis Hampson (*Oligostigma*) 118
 flaviplagiialis Hampson (*Neogenesis*) 180
 flavipunctalis (var.) Hampson (*Stenia*) 164
 flavivittalis Hampson (*Argyrestis*) 63
 flavomaculalis Pagenstecher (*Piletocera*) 172
 flavomaculata Hampson (*Piletocera*) 172
 flexiguttalis Warren (*Piletocera*) 173
 flexuosa Dyar (*Scoparia*) 33
 floralis Leech (*Nymphula*) 78
 fluctuosalis Zeller (*Nymphula*) 81, 85, 86, 87
 fluctuosalis Hampson (*Nymphula*) 89
 foedalis Guenée (*Nymphula*) 76, 84, 86, 87, 88
 foedalis auct. (*Nymphula*) 84
 foederalis Hübner (*Zebronia*) 186
 foeminalis (ab.) Dyar (*Nymphula*) 88
 formosa Butler (*Scoparia*) 38
 formosalis Clemens (*Nymphula*) 88
 formosalis Strand (*Piletocera*) 172
 fovealis Zeller (*Duponchelia*) 133
 foveocosta Dyar (*Ambia*) 116
 fragmentalis Lederer (*Aulacodes*) 144
 fragosa Meyrick (*Scoparia*) 39
 franciscalis Schaus (*Nymphula*) 93
 francki Caradja (*Musotima*) 105
 fregonalis Snellen (*Nymphula*) 85
 frequentella Stainton (*Scoparia*) 85
 frigida Butler (*Scoparia*) 41
 frigida Meyrick (*Scoparia*) 41
 frigidella Hampson (*Scoparia*) 12
 fulguralis Caradja (*Nymphula*) 86
 fulcalis Clemens (*Cataclysta*) 99, 101
 fulcalis Hampson (*Cataclysta*) 100
 fuliginosa W. Rothschild (*Bradina*) 155
 fuliginosa W. Rothschild (*Strepsinoma*) 133
 fulvalis Dyar (*Ambia*) 116
 fulvalis Hampson (*Ambia*) 111
 fulvalis Hampson (*Massepha*) 129
 fulvalis Hampson (*Piletocera*) 179
 fulvalis Warren (*Ambia*) 112
 fulviceps Hampson (*Diathrausta*) 169
 fulvicinctalis Hampson (*Argyrestis*) 65
 fulvicolor Hampson (*Ambia*) 113
 fulvicolor Hampson (*Oligostigma*) 121
 fulvitinctalis Hampson (*Ambia*) 114
 fulvobasalis Snellen (*Ambia*) 111
 fulvofusa Hampson (*Diathraustodes*) 158
 fulvosignalis Snellen (*Scoparia*) 31

- fumata Philpott (*Scoparia*) 54
 fumibasale Hampson (*Oligostigma*) 117
 fumidalis Turati (*Duponchelia*) 134
 furvitermen Hampson (*Deuterophysa*) 162
 fusalis Hampson (*Ambia*) 113
 fusalis Hampson (*Nymphula*) 85
 fusalis Hampson (*Stenia*) 168
 fusalis Dyar (*Ambia*) 116
 fusalis Hampson (*Cataclysta*) 97
 fusalis Hampson (*Margorochroma*) 126
 fusalis Snellen (*Musotima*) 104
 fuscella Turati (*Scoparia*) 20
 fuscicostalis Hampson (*Nymphula*) 85
 fuscicostalis W. Rothschild (*Aulacodes*) 141
 fuscidiscalis Hampson (*Musotima*) 105
 fuscifusale Hampson (*Oligostigma*) 117
 fuscilunalis Hampson (*Stenia*) 169
 fuscipennalis Hampson (*Mabra*) 191
 fuscocillialis Ragonot (*Stenia*) 165
 fuscocomarginata Bethune-Baker (*Nymphula*) 77

 galactalis Hudson (*Scoparia*) 46
 gallica Peyerimhoff (*Scoparia*) 21
 gangeticalis Lederer (*Parthenodes*) 123
 garzellalis Hampson (*Mabra*) 192
 garzettalis Felder & Rogenhofer (*Mabra*) 192
 gaudialis Hampson (*Margarosticha*) 93
 gelliasalis Walker (*Stenia*) 167
 gemella Moore (*Decticogaster*) 147
 geminalis Caradja (*Bradina*) 152
 gemmiferalis Lederer (*Argyrectis*) 63
 genuialis Lederer (*Nymphula*) 91
 genuinalis Lederer (*Nymphula*) 91
 gephyrotis Meyrick (*Aulacodes*) 141
 geraea Meyrick (*Scoparia*) 40
 gethosyna Turner (*Scoparia*) 44
 gibbosalis Guenée (*Eoöphila*) 138
 gigantalis Druce (*Stenia*) 170
 gigantalis Janse (*Cymoriza*) 108
 gigantalis Zerny (*Stenia*) 165
 gilvescens Rebel (*Scoparia*) 28
 glabralis Hübner (*Perinephela*) 186
 glaucinalis Hampson (*Bradina*) 154
 glaucinalis Hampson (*Clupeosoma*) 183
 glaucinalis Hampson (*Stenia*) 168
 glauculalis Hampson (*Scoparia*) 35
 glenecalis Dyar (*Parambia*) 109
 glycerialis Walker (*Cataclysta*) 99
 glycisalis Dyar (*Cataclysta*) 102
 gnomosynalis Dyar (*Parambia*) 109
 gomphota Meyrick (*Scoparia*) 42
 goniophoralis Hampson (*Aulacodes*) 143
 gonodecta Meyrick (*Scoparia*) 37
 gontranalis Schaus (*Argyrectis*) 65
 gordianalis Schaus (*Argyrectis*) 68
 gothicalis Felder & Rogenhofer (*Aulacodes*) 145
 gratiatella (fa.) Turati (*Scoparia*) 20
 gracilalis Stainton (*Scoparia*) 21
 gracilis Hampson (*Massepha*) 128
 gracilis Philpott (*Scoparia*) 47
 grammalis Guenée (*Massepha*) 128
 grammalis Guenée (*Pileocera*) 180
 graphicals Hampson (*Nymphula*) 92
 gratalis Walker (*Argyrectis*) 67
 grimbaldalis Schaus (*Aulacodes*) 143
 grisealis Hampson (*Deuterophysa*) 162
 grisealis Hampson (*Massepha*) 128
 grisealis Hampson (*Nymphula*) 86
 grisealis Hampson (*Oligostigma*) 121
 grisealis Hampson (*Stenia*) 166, 167
 grisealis W. Rothschild (*Bradina*) 156
 grisealis W. Rothschild (*Eristena*) 70
 grisealis W. Rothschild (*Strepsinoma*) 133

- grisealis* (var.) Staudinger
(Psammotis) 188
griseata Butler (*Duponchelia*)
 134
griseifusa Hampson (*Diathrausta*) 160
griseotinctalis W. Rothschild
(Dracaenura) 136
guadarensis Schaus (*Argyractis*)
 64
gualbertalis Schaus (*Parthenodes*) 125
guenealis Snellen (*Cataclysta*)
 102
guencealis Hampson (*Cataclysta*)
 102
guianalis Schaus (*Stenia*) 169
gurgitalis Lederer (*Nymphula*)
 92
gymnopis Meyrick (*Scoparia*) 39
gyralis Hulst (*Nymphula*) 90
gyrotona Meyrick (*Scoparia*) 49

hadesialis Hampson (*Pileocera*)
 173
halialis Walker (*Eoöphila*) 138
halirrhoa Meyrick (*Scoparia*) 38
halitalis Felder & Rogenhofer
(Aulacodes) 145
halmaea Meyrick (*Scoparia*) 41
halopis Meyrick (*Scoparia*) 52
hamalis Hampson (*Aulacodes*)
 140
hamalis Snellen (*Aulacodes*) 140
hamiferalis Hampson (*Argyractis*)
 63
hampsoni South (*Nymphula*) 83
hapilista Swinhoe (*Oligostigma*)
 117
hapilistalis Strand (*Strepsinoma*)
 133
haplomorpha Meyrick (*Bradina*)
 156
harlequinialis (fa.) Dyar (*Diathrausta*) 161
harpalea Meyrick (*Scoparia*) 53
harpalis Snellen (*Argyractis*) 69
hawaiensis Butler (*Scoparia*) 40
hebraicalis Snellen (*Nymphula*)
 84
hecate Butler (*Pileocera*) 177
hederalis Amsel (*Nymphula*) 75
helenensis Wollaston (*Scoparia*)
 31
heliopalisis Clemens (*Cataclysta*)
 101
helopalisis Hampson (*Cataclysta*)
 101

hemicyptis Meyrick (*Oligostigma*) 118
hemicycla Meyrick (*Scoparia*) 47
hemigrammalis Hampson (*Ambia*) 114
hemileucalis Hampson (*Diathraustodes*) 158
hemimelaena Hampson (*Aulacodes*) 140
hemiphaealis Hampson (*Bradina*)
 153
hemiplaca Meyrick (*Scoparia*) 46
hemithermalis Hampson (*Aulacodes*) 142
hemmingalis Schaus (*Bradina*)
 157
heptopalis Hampson (*Ambia*) 115
heptopalis Hampson (*Aulacodes*)
 141
hermeasalis Walker (*Nymphula*)
 92
herminalis Schaus (*Argyractis*)
 64
herreralis Schaus (*Stenia*) 169
heterodoxa Meyrick (*Eurytorna*)
 171
hexalitha Meyrick (*Cataclysta*) 99
hieralis Snellen (*Stenia*) 166
hirsuta Semper (*Eoöphila*) 138
hisbonalis Walker (*Campomastix*)
 181
hodevalis Druce (*Aulacodes*) 145
hoenei Caradja (*Cymoriza*) 108
holophaealis Hampson (*Pileocera*) 178
holophaealis Hampson (*Pileocera*) 178
holoxantha Hampson (*Luma*) 130
homala Meyrick (*Scoparia*) 42
homalopa Meyrick (*Mestolobes*)
 57
homogenes Meyrick (*Dasyscopa*)
 57
homora Turner (*Eclipsiodes*) 8
horatius Butler (*Bradina*) 155
horochroa Meyrick (*Dracaenura*)
 135
hortulata Linné (*Eurrhypara*)
 190
hyalinalis Hübner (*Psammotis*)
 189
hydrocampalis Guenée (*Parthenodes*) 125
hydrothionalis Snellen (*Nymphula*) 92
hyperochalis Dyar (*Stenia*) 170
hypoxantha Lower (*Scoparia*) 44

- iambealis* Walker (*Ambia*) 111
iambesalis Hampson (*Ambia*) 111
ianthealis Walker (*Stenia*) 168
ianthes Meyrick (*Scoparia*) 39
iasusalis Walker (*Argyractis*) 69
icciusalis Walker (*Nymphula*) 91
icelomorpha Turner (*Eclipsiodes*) 8
ichinosawana Matsumura (*Scoparia*) 26
idiogama Meyrick (*Scoparia*) 32
idiotis Meyrick (*Aulacodes*) 139
ilialis Walker (*Cataclysta*) 102
illectalis Walker (*Piletocera*) 173
illectalis Walker (*Piletocera*) 174
illota Philpott (*Scoparia*) 50
imitabilis Dyar (*Cataclysta*) 101
immanis Hampson (*Parthenodes*) 123
immundalis South (*Piletocera*) 175
imparcilla (fa.) De la Harpe (*Scoparia*) 21
imparella Wocke (*Scoparia*) 21
imparilis Dyar (*Scoparia*) 36
impressalis Lederer (*Bradina*) 151, 154
impuralis Felder & Rogenhofer (*Nymphula*) 93
inaurata Cramer (*Argyractis*) 63
incertalis Duponchel (*Scoparia*) 16
incertalis Guenée (*Scoparia*) 14
incertella Duponchel (*Scoparia*) 16
incisa Strand (*Pseudoligostigma*) 194
incommoda Butler (*Nymphula*) 77
inconsequens Warren (*Orphnophanes*) 131
inconspicua Schaus (*Piletocera*) 180
inconspicualis Kenrick (*Piletocera*) 178
incrustalis Snellen (*Musotima*) 106
incultalis Schaus (*Parthenodes*) 126
incurvalis R. South (*Nymphula*) 85
indianalis (var.) Dyar (*Stenia*) 168
indistinctalis Walker (*Scoparia*) 49
indomitalis Berg (*Nymphula*) 93
ineffectalis Walker (*Stenia*) 170
inexoptata Dyar (*Scoparia*) 36
inextricata Meyrick (*Parthenodes*) 122
infernalis Hampson (*Piletocera*) 177
infernalis (ab.) Zerny (*Epistenia*) 163
infirmalis Möschler (*Nymphula*) 93
ingratalis Guenée (*Scoparia*) 17
ingratella Knaggs (*Scoparia*) 18
ingratella Zeller (*Scoparia*) 17
insectale Hampson (*Oligostigma*) 117
insectalis Pryer (*Oligostigma*) 117
inspersalis (ab.) Galvagni (*Scoparia*) 18
instrumentalis Swinhoe (*Musotima*) 106
insulalis Walker (*Argyractis*) 68
intensa (subsp.) W. Rothschild (*Aulacodes*) 142
interlinealis Warren (*Scoparia*) 30
intermedialis Caradja (*Bradina*) 153
interruptalis Hampson (*Stenia*) 169
interruptalis Pryer (*Nymphula*) 75, 83
interruptalis Warren (*Ambia*) 112
interruptalis Wileman & South (*Cymoriza*) 108
interstrigalis Hampson (*Ambia*) 111, 113
intervacatalis Christoph (*Stenia*) 164
intortalis Schaus (*Ambia*) 116
iochrysa Meyrick (*Mesolobes*) 56
iolepta Dyar (*Cataclysta*) 103
ioxanthias Meyrick (*Auxolophotis*) 127
irialis Meyrick (*Ambia*) 112
iridalis Guenée (*Genoloba*) 193
iridia Meyrick (*Clepsicosma*) 193
irisale Hampson (*Oligostigma*) 120
irialis Walker (*Oligostigma*) 120
iriusalis Walker (*Ambia*) 112
irrectalis Guenée (*Cymoriza*) 108
irrectalis Guenée (*Cymoriza*) 108
irroratalis Dyar (*Cataclysta*) 101
irroratalis Hampson (*Stenia*) 169
ischnias Meyrick (*Scoparia*) 41
ischnoptera Turner (*Scoparia*) 44
isochroalis Hampson (*Scoparia*) 22
isoplaca Meyrick (*Scoparia*) 39

- isorrhoea Meyrick (*Cymoriza*) 108
 isoterma Meyrick (*Mestolabes*) 57
 italica (fa.) Turati (*Scoparia*) 19
 itealis Walker (*Nymphula*) 89
 ithyntis Turner (*Scoparia*) 44
 ivanalis (var.) Krulikovsky (*Scoparia*) 18

 jalapalis Schaus (*Argyrectis*) 64
 jaliscalis Schaus (*Cataclysta*) 103
 janapalis Hampson (*Decticogaster*) 147, 148
 janapalis Walker (*Decticogaster*) 146
 japanalis Snellen (*Decticogaster*) 146
 javanalis Guenée (*Aulacodes*) 139
 javanalis Snellen (*Pileocera*) 173
 javanica (var.) Strand (*Oligostigma*) 119
 jonesalis Dyar (*Scoparia*) 35
 jonesalis Schaus (*Ambia*) 115
 jucunda Butler (*Scoparia*) 39
 julitalis Schaus (*Aulacodes*) 144
 junceale Hampson (*Oligostigma*) 121
 juncealis Fernald (*Nymphula*) 90
 juncealis Guenée (*Oligostigma*) 121
 junctalis Hampson (*Cataclysta*) 95
 junctiscriptalis Hampson (*Aulacodes*) 143

 kearfottalis (ab.) Dyar (*Cataclysta*) 100
 klimai Bryk (*Aulacodes*) 141
 klinckowstroemi Hamfeldt (*Scoparia*) 15
 klossi W. Rothschild (*Ambia*) 113
 kwangtungialis Caradja (*Oligostigma*) 119
 kwangtungialis (subsp.) Caradja (*Scoparia*) 15

 lacertalis Guenée (*Stenia*) 170
 laetalis Guenée (*Scoparia*) 26
 laetella Zeller (*Scoparia*) 25
 levalis Warren (*Orphnophanes*) 131
 lamialis Walker (*Cataclysta*) 101
 laminalis Hampson (*Aulacodes*) 141
 lampetialis Walker (*Cataclysta*) 99

 lampra Tams (*Clupeosoma*) 184
 lancealis Schiffermüller & Denis (*Perisyntrocha*) 185
 lanceolalis Guenée (*Pileocera*) 176
 lanceolalis Hampson (*Argyrectis*) 69
 laniferalis Hampson (*Clupeosoma*) 184
 lapponica (var.) Caradja (*Scoparia*) 22
 latalis Hampson (*Pileocera*) 174
 latalis Walker (*Pileocera*) 174
 lathurusalis Walker (*Zebronia*) 186
 latifascialis Snellen (*Aulacodes*) 141
 latifascialis Warren (*Parthenodes*) 124
 latifasciata (subsp.) W. Rothschild (*Nymphula*) 75
 latisquamalis Snellen (*Nicaria*) 157
 laurentialis Schaus (*Argyrectis*) 67
 lavinia Schaus (*Clupeosoma*) 185
 legionaria Philpott (*Scoparia*) 52
 legnota Meyrick (*Scoparia*) 54
 lemnalis Schiffermüller & Denis (*Cataclysta*) 96
 lemnata Linné (*Cataclysta*) 95
 leptalea Meyrick (*Scoparia*) 50
 leptogastralis Walker (*Bradina*) 150
 leptographa Meyrick (*Bradina*) 156
 leptolopha Tams (*Bradina*) 154
 leptomerallis Walker (*Bradina*) 152
 leptophaea Meyrick (*Scoparia*) 50
 leptorrhoda Turner (*Musotima*) 106
 leroii Strand (*Cataclysta*) 98
 leucistis Dognin (*Argyrectis*) 70
 leucocephalis Hampson (*Pileocera*) 178
 leucochrysa Meyrick (*Ambia*) 110
 leucoecymalis Hampson (*Ambia*) 115
 leucogastralis Hampson (*Pileocera*) 178
 leucogonialis Hampson (*Argyrectis*) 66
 leucogramma Meyrick (*Scoparia*) 54
 leucographa Hampson (*Dithrausta*) 162
 leucomela Lower (*Scoparia*) 44

- leucomma Hampson (*Oligostigma*) 120
 leuconota Lower (*Eclipsiodes*) 9
 leucopeplalis Hampson (*Cholius*) 10
 leucophthalma Dyar (*Scoparia*) 34
 leucoplaca Meyrick (*Ambia*) 111, 113
 leucoplagalis Hampson (*Nymphula*) 92
 leucoprocta Hampson (*Dracaenura*) 135
 leucostictalis Hampson (*Ambia*) 115
 leucostictalis Hampson (*Symphonia*) 104
 leucostola Hampson (*Argyrectis*) 61
 leucostola Hampson (*Nymphula*) 85
 leucostriatalis Hampson (*Argyrectis*) 62
 leucothoalis Walker (*Nymphula*) 77
 leucotrigona Hampson (*Diathraustodes*) 158
 leucozonalis Hampson (*Margarosticha*) 94
 leuculalis Snellen (*Decticogaster*) 148
 leucura Hampson (*Bradina*) 154
 libella Grote (*Scoparia*) 34
 lichenopa Lower (*Scoparia*) 44
 liebmanni Petry (*Scoparia*) 24
 limatula Philpott (*Scoparia*) 52
 linealis Guenée (*Nymphula*) 81
 linealis Moore (*Nymphula*) 78
 linealis Walker (*Scoparia*) 49
 lineola Curtis (*Scoparia*) 24
 lineolalis Guenée (*Scoparia*) 25
 liodesalis Walker (*Bradina*) 156
 lipocosmalis Snellen (*Ambia*) 112
 lirisalis Walker (*Decticogaster*) 148
 litana's Walker (*Nymphula*) 86
 litana's Walker (*Piletocera*) 175
 lithocharis Meyrick (*Argyrectis*) 61
 lithorma Meyrick (*Argyrectis*) 62
 livonica (var.) Zeller (*Scoparia*) 23
 lobipennis Moore (*Almonia*) 148
 lobophoralis Hampson (*Ambia*) 112
 locularis Meyrick (*Scoparia*) 48
 locuples Butler (*Ambia*) 110
 longalis Haworth (*Perisyntrocha*) 186
 longidentalis Hampson (*Luma*) 130
 longinasus De Joannis (*Cilaus*) 193
 longipalpis Butler (*Campomastix*) 181
 longipedalis Curtis (*Stenia*) 164
 longipennis Hampson (*Argyrectis*) 66
 longipennis Hampson (*Massepha*) 128
 longipennis Zeller (*Scoparia*) 37
 longiplagialis Hampson (*Aulacodes*) 143
 lophocera Hampson (*Coptobasis*) 158
 lophophoralis Hampson (*Bradina*) 150
 lophosomal's Hampson (*Argyrectis*) 62
 loricalalis Lederer (*Parthenodes*) 123
 lorquinialis Guenée (*Stenia*) 164
 lotialis Felder & Rogenhofer (*Nymphula*) 91
 loxocentra Meyrick (*Scoparia*) 38
 lubricalis Dognin (*Stenia*) 171
 luciana's Schaus (*Argyrectis*) 65
 lucida's Walker (*Scoparia*) 31
 lugubralis Walker (*Scoparia*) 35
 lulesalis Schaus (*Argyrectis*) 67
 luminatrix Meyrick (*Scoparia*) 54
 lunalis Kenrick (*Aulacodes*) 144
 lunulalis Costa (*Nymphula*) 75
 lupa Druce (*Massepha*) 129
 lusedalis Hampson (*Scoparia*) 31
 lutealis Snellen (*Stenia*) 167
 luteivittalis Mabilie (*Nymphula*) 87
 luteusalis Hampson (*Scoparia*) 28
 luzialis (var.) Guenée (*Scoparia*) 23
 lychnophanes Meyrick (*Scoparia*) 47
 lycopodiae Swezey (*Scoparia*) 41
 lycopusalis Walker (*Piletocera*) 174
 lypera Tams (*Diathrausta*) 162
 macaralis Walker (*Bradina*) 152, 154
 macroperalis Hampson (*Ellogima*) 181
 macrophanes Meyrick (*Scoparia*) 41
 macropsalis Hampson (*Luma*) 130
 maculalis Clemens (*Nymphula*) 88

- maculifrons Hampson (*Pileocera*) 173
 magnificalis Hübner (*Cataclysta*) 101
 magnificentalis Swinhoe (*Ambia*) 112
 maguillalis Schaus (*Argyrectis*) 63
 major (subsp.) W. Rothschild (*Aulacodes*) 142
 malayalis Hampson (*Argyrectis*) 60
 malousalis Schaus (*Argyrectis*) 67
 mallaleuca Hampson (*Stenia*) 179
 mandschurica Christoph (*Scoparia*) 17
 manganeutis Meyrick (*Scoparia*) 50
 mangholdalis Schaus (*Cataclysta*) 103
 manifestalis Guenée (*Scoparia*) 19
 manifestella Herrieh-Schaeffer (*Scoparia*) 19
 manilensis Hampson (*Nymphula*) 84
 manilensis Sauber (*Cataclysta*) 98
 mannusalis Walker (*Bradina*) 154
 maoriella Walker (*Scoparia*) 39
 marconalis Schaus (*Ambia*) 115
 marginalis Moore (*Ambia*) 112
 marginalis W. Rothschild (*Ambia*) 114
 marginalis Swinhoe (*Massepha*) 127
 marialis Schaus (*Stenia*) 168
 marmarias Meyrick (*Scoparia*) 39
 marmorea Meyrick (*Nymphula*) 84
 maronialis Schaus (*Argyrectis*) 63
 masculinalis (ab.) Dyar (*Nymphula*) 89
 matronalis Guenée (*Genoloba*) 193
 mayottensis (var.) Pagenstecher (*Zebronia*) 186
 medicinalis Grote (*Cataclysta*) 100
 medinella Snellen (*Scoparia*) 30
 mediocinctalis Hampson (*Parthenodes*) 125
 mediofascialis Hampson (*Aulacodes*) 143
 mediorufalis Hampson (*Scoparia*) 30
 medusalis Walker (*Nymphula*) 87
 meeckii Lucas (*Pileocera*) 179
 megapteralis Walker (*Coptobasis*) 158
 megaspilalis Walker (*Pileocera*) 172
 megesalis Walker (*Bradina*) 149, 151
 melanaegis Meyrick (*Scoparia*) 48
 melana'is Hampson (*Ambia*) 113
 melanauges Meyrick (*Pileocera*) 177
 melanicalis Hampson (*Pileocera*) 125
 melanistis Hampson (*Ambia*) 112
 melanocephala Meyrick (*Scoparia*) 38
 melanodes Meyrick (*Oligostigma*) 117
 melanographa Hampson (*Scoparia*) 23
 melanograptia Hampson (*Argyrectis*) 69
 melanolitha Turner (*Eristena*) 70
 melanoperas Hampson (*Bradina*) 152
 melanopis Meyrick (*Scoparia*) 41
 melanops Hampson (*Aulacodes*) 140
 melanota'is Hampson (*Oligostigma*) 118
 melanoxantha Turner (*Scoparia*) 44
 melatorna'is Hampson (*Cataclysta*) 102
 melesa'is Walker (*Bradina*) 152, 153
 melichlora Meyrick (*Scoparia*) 40
 melinhyrta Turner (*Eclinsiodora*) 9
 melitealis Walker (*Stenia*) 170
 meliturga Meyrick (*Scoparia*) 45
 mell'iculale Hampson (*Hydropi-nea*) 182
 mellicula'is Lederer (*Hydropi-nea*) 182
 mell'inalis Druce (*Bradina*) 157
 mendica v. Hedemann (*Heringiella*) 180
 mercurialis Guenée (*Scoparia*) 26
 mercurea Haworth (*Scoparia*) 28
 mercurella Stephens (*Scoparia*) 28
 mercurella Treitschke (*Scoparia*) 26
 meristis Meyrick (*Scoparia*) 41
 meropa'is Walker (*Cataclysta*) 97
 mesacma Meyrick (*Mestolobes*) 56

- mesogramma Lower (*Scoparia*) 44
 mesoleuca Meyrick (*Scoparia*) 37
 mesoleuca Meyrick (*Scoparia*) 38
 mesoleucalis Hampson (*Pileto-
cera*) 125
 mesophaealis Hampson (*Pileto-
cera*) 178
 mesopsectralis Hampson (*Copto-
basis*) 157
 mesorhpha Meyrick (*Cataclysta*)
 95
 mesosialis Hampson (*Aulacodes*)
 142
 mesoscotalis Hampson (*Ambia*)
 111, 113
 metachryson Hampson (*Clupeoso-
ma*) 184
 metacrossa Hampson (*Scoparia*)
 44
 metaleuca Hampson (*Dracae-
nura*) 136
 metaleucalis Hampson (*Scoparia*)
 30
 metaleucalis Walker (*Bradina*)
 155
 metallescens Felder & Rogenho-
 fer (*Mabra*) 192
 metalophota Hampson (*Ambia*)
 115
 metasialis Rebel (*Stenia*) 165
 metastictalis Hampson (*Nym-
phula*) 88
 metataxalis Hampson (*Aulacodes*)
 143
 metazonalis Hampson (*Argyrac-
tis*) 69
 metazonalis Hampson (*Aulaco-
des*) 142
 metochrealis Hampson (*Pileto-
cera*) 175
 metriodora Meyrick (*Aulacodes*)
 139
 meyrickii Butler (*Tetraproso-
pus*) 55
 miantis Meyrick (*Scoparia*) 38
 miantodes Meyrick (*Bradina*) 156
 micans Hampson (*Niphostola*) 132
 micralis Hampson (*Deuterophy-
sa*) 162
 micralis Hampson (*Pileto-
cera*) 175
 microcentra Meyrick (*Pileto-
cera*) 179
 microdentalis Hampson (*Pileto-
cera*) 177
 microdentalis Hampson (*Scopa-
ria*) 16
 micropalis Hampson (*Argyractis*)
 69
 microphthalma Meyrick (*Scopa-
ria*) 47
 midalis Schaus (*Stenia*) 168
 midas Butler (*Cataclysta*) 97
 mignonalis Dyar (*Argyractis*) 63
 mimetica Lower (*Clupeosoma*)
 184
 mimeticalis Caradja (*Aulacodes*)
 141
 mimicalis Hampson (*Argyractis*)
 61
 minimalis Herrich-Schaeffer (*Ca-
taclysta*) 103
 minimalis Saalmüller (*Nymphu-
la*) 87
 minoralis Mabilie (*Nymphula*) 87
 minoralis Snellen (*Stenia*) 166
 minualis Walker (*Diathrausta*)
 161
 minualis Walker (*Scoparia*) 46
 minuscula Butler (*Mestolobes*) 56
 minusculealis Walker (*Scoparia*)
 46
 mifutalis Caradja (*Oligostigma*)
 119
 minutalis (fa.) Druce (*Dia-
thrausta*) 161
 mirabilis Butler (*Pileto-
cera*) 176
 miralis Möschler (*Cataclysta*) 103
 moanalis Felder & Rogenhofer
 (*Scoparia*) 50
 modestalis Lederer (*Bradina*) 153,
 155
 modestalis Lederer (*Nymphula*)
 92
 modestalis Saalmüller (*Stenia*)
 167
 molifera Meyrick (*Scoparia*) 46
 molliculella Dyar (*Scoparia*) 36
 mollitalis Schaus (*Oligostigma*)
 121
 monetalis Snellen (*Arxama*) 71
 moniligeralis Lederer (*Cata-
clysta*) 100, 101
 monocromalis Walker (*Coptoba-
sis*) 158
 monomma Warren (*Luma*) 163
 mononalis Dyar (*Stenia*) 169
 montana Butler (*Scoparia*) 41
 montana Haimbach (*Diathrausta*)
 161
 moralis Schaus (*Aulacodes*) 146
 morbidalis Dyar (*Homophysodes*)
 193
 mormodes Meyrick (*Aulacodes*)
 141
 multifacies Dyar (*Scoparia*) 36
 multipicta Dyar (*Argyractis*) 66

- multipictalis Hampson (*Symphonia*) 104
 munakatae Matsumura (*Nymphula*) 81
 muneralis Hampson (*Scoparia*) 11
 muralis Guenée (*Scoparia*) 24
 murana Curtis (*Scoparia*) 23
 murana Hampson (*Scoparia*) 24
 murana Rebel (*Scoparia*) 20
 murcurella Hampson (*Scoparia*) 26
 murificalis Hampson (*Scoparia*) 31
 murificalis Walker (*Scoparia*) 30, 32
 murinalis Warren (*Eristena*) 70
 mutanalis (fa.) Caradja (*Stenia*) 164
 myina Meyrick (*Nymphula*) 84
 myota Meyrick (*Dracaenura*) 135
 mysolalis Walker (*Pileocera*) 172
 nandinalis Hampson (*Argyrectis*) 61
 napolitalis (var.) Millière (*Scoparia*) 29
 narcissusalis Walker (*Nymphula*) 72
 nasonia Meyrick (*Pileocera*) 179
 nawae Matsumura (*Susumia*) 107
 navalis Wileman (*Nymphula*) 76
 nebulosalis Fernald (*Nymphula*) 91
 necomalis Dyar (*Argyrectis*) 69
 nectalis Snellen (*Aulacodes*) 141
 nectarias Mevrick (*Scoparia*) 39
 nectarioides Swezey (*Scoparia*) 39
 nennuisalis Schaus (*Stenia*) 167
 nephelanthopa Meyrick (*Aulacodes*) 144
 nephelitis Mevrick (*Scoparia*) 44
 nerinalis Walker (*Diathrausta*) 160
 nerinalis (fa.) Walker (*Diathrausta*) 161
 nervosalis Lucas (*Stenia*) 165
 neuralis Hampson (*Bradina*) 155
 nevadalis (var.) Caradja (*Scoparia*) 17
 nevadensis (subsp.) Zerny (*Scoparia*) 31
 nigerialis Hampson (*Argyrectis*) 61
 nigra (var.) Hamfelt (*Scoparia*) 15
 nigra Warren (*Nymphula*) 83
 nigrabalis Caradja (*Nymphula*) 78
 nigrescens Butler (*Pileocera*) 176
 nigrifusalis Dognin (*Argyrectis*) 66
 nigriplaga Swinhoe (*Parthenodes*) 124
 nigriplagialis Hampson (*Aulacodes*) 142
 nigripunctalis R. South (*Bradina*) 153
 nigriscripta Swinhoe (*Mabra*) 191
 nigriscripta Swinhoe (*Stenia*) 166
 nigristriata Hampson (*Cataclysta*) 98
 nigritalis (fa.) Hampson (*Cataclysta*) 95
 nigritalis Walker (*Scoparia*) 31
 nigrolinealis Pryer (*Nymphula*) 83
 nigropunctata Pagenstecher (*Bradina*) 154
 ninguidalis Hulst (*Scoparia*) 12
 niphetodes Turner (*Scoparia*) 44
 niphoplagalis Hampson (*Argyrectis*) 65
 niphosphora Meyrick (*Scoparia*) 53
 niphospora Hampson (*Scoparia*) 53
 niphostola Hampson (*Gargela*) 59
 nissonalis Schaus (*Aulacodes*) 145
 nitens Buttler (*Nymphula*) 87
 nitens Schaus (*Argyrectis*) 65
 nitidalis Dognin (*Neomabra*) 192
 nitidalis Hampson (*Coelorrhynchidia*) 159
 nitidalis Walker (*Musotima*) 105
 nivalis Schiffermüller & Denis (*Nymphula*) 80
 nivealis Hübner (*Nymphula*) 80
 niveinotatum Hampson (*Oligostigma*) 117
 niveiplagalis Hampson (*Ambia*) 114
 noctualis Hampson (*Zagiridia*) 182
 nolalis Felder & Rogenhofer (*Scoparia*) 32
 nomeutis Mevrick (*Scoparia*) 53
 nominatalis Hulst (*Scoparia*) 33, 37
 nomophilalis Dyar (*Nymphula*) 90
 noordimimalis Hampson (*Clupeosoma*) 183
 normalis Dyar (*Scoparia*) 32
 novaguineensis Kenrick (*Ambia*) 113
 nubilalis R. South (*Musotima*) 105

- nudicornis Hampson (*Pileoce-*
ra) 175
 nugal's Snellen (*Scoparia*) 31
 numeral's Zeller (*Scoparia*) 11
 nyasalis Hampson (*Argyractis*)
 61
 nyasalis Hampson (*Cataclysta*)
 98
 nyctombra Meyrick (*Scoparia*)
 40
 nyctopis Meyrick (*Cataclysta*) 97
 nymphaealis Schiffermüller &
 Denis (*Nymphula*) 74
 nymphaealis Treitschke (*Nym-*
phula) 73
 nymphaeata Linné (*Nymphula*)
 73, 83
 nymphula's Hampson (*Argyrac-*
tis) 62
 nymphulalis Hampson (*Parthe-*
nodes) 125

 obealis Snellen (*Mabra*) 192
 obisalis (ab.) Dyar (*Scoparia*)
 34
 obital's Walker (*Nymphula*) 81
 oblectalis Hulst (*Hydropionea*)
 182
 obliquifascia Hampson (*Cata-*
clysta) 102
 obliquistriga W. Rothschild (*Ambia*)
 113
 obliquivitta Hampson (*Aulacodes*)
 143
 obliquivitta Hampson (*Gargela*)
 60
 oblitera's Walker (*Genoloba*) 192
 oblitera's Walker (*Nymphula*)
 90, 92
 obliturata (var.) Warren (*Dia-*
thrausta) 160
 obregonalis Schaus (*Deuterophy-*
sa) 162
 obscura Warren (*Stenia*) 168
 obscuralis Grote (*Nymphula*) 89
 obscuralis Möschler (*Nymphula*)
 90
 obscuralis Swinhoe (*Pelena*) 129
 obsoleta Staudinger (*Scoparia*) 29
 obtusalis Dyar (*Aulacodes*) 144
 ochracea (var.) Hauser (*Cata-*
clysta) 96
 ochreal's Schiffermüller & Denis
 (*Cholius*) 9
 ochreipicta Meyrick (*Aulacodes*)
 139
 ochrias Meyrick (*Mestolobes*) 56
 ochripicta Moore (*Aulacodes*) 140

 ochrocealis Hampson (*Arxama*)
 71
 ochrophora Turner (*Scoparia*) 43
 ochropteralis Guenée (*Musotima*)
 105
 ochrosema Meyrick (*Pileocera*)
 176
 ochrotalis Hampson (*Scoparia*) 31
 ocelligera Meyrick (*Pileocera*)
 179
 octalis Hampson (*Thysanoidma*)
 109
 octavialis Lederer (*Aulacodes*)
 143
 octavianella (var.) Mann (*Sco-*
paria) 15
 octomaculalis Fernald (*Dia-*
thrausta) 161
 octonella (fa.) Zeller (*Scoparia*)
 20
 octophora Meyrick (*Scoparia*) 54
 octopis Meyrick (*Ephormotris*)
 109
 octosemalis Hampson (*Pileocera*)
 173
 oculalis Hampson (*Ambia*) 113
 oculata Philpott (*Scoparia*) 55
 odoalis Schaus (*Argyractis*) 68
 odrianalis Schaus (*Oligostigma*)
 121
 odulphalis Schaus (*Aulacodes*) 145
 oechmialis Hampson (*Aulacodes*)
 144
 oedizonalis Hampson (*Ambia*) 115
 oenone Butler (*Mestolobes*) 57
 oenopis Meyrick (*Scoparia*) 40
 oertzeniella Herrich-Schaeffer
 (*Scoparia*) 29
 oethonalis Walker (*Nymphula*) 27
 officialis Meyrick (*Scoparia*) 44
 oligalis Hampson (*Ambia*) 113
 oligostigmalis Hampson (*Eriste-*
na) 70
 olivalis Hampson (*Parthenodes*)
 123
 olivaris Hampson (*Scoparia*) 30
 ombrias Meyrick (*Mestolobes*) 56
 omichlopi's Meyrick (*Scoparia*) 41
 onustalis Walker (*Almonia*) 148
 onyxalis Hampson (*Argyractis*)
 68
 opacalis Rebel (*Pileocera*) 172
 opacusalis Swinhoe (*Bradina*) 153
 ophiaula Meyrick (*Nymphula*) 87
 opostactis Meyrick (*Scoparia*) 44
 opulentalis Druce (*Argyractis*) 62
 opulenta's Lederer (*Cataclysta*)
 100, 101
 oreas Meyrick (*Scoparia*) 45

- organea* Meyrick (*Scoparia*) 54
orientalis Alphéraky (*Scoparia*) 25
orientalis Filipjev (*Nymphula*) 83
orientalis Snellen (*Piletocera*) 173
orizabalis Schaus (*Argyrectis*) 64
ornata Moore (*Oligostigma*) 118
ornatalis Leech (*Luma*) 130
ornatum Hampson (*Oligostigma*) 118
orphanalis Pagenstecher (*Oligostigma*) 121
orthoria Meyrick (*Scoparia*) 37
orthoteles Meyrick (*Oligostigma*) 118
orthrias Meyrick (*Mestolobes*) 56
oryzalis Wood-Mason (*Nymphula*) 81
osculatrix Meyrick (*Nymphula*) 85, 87
ossealis Hampson (*Perisyntrocha*) 132
ovomaculalis W. Rothschild (*Aulacodes*) 144
ovulalis Hampson (*Coelorhynchidia*) 158
oxygona Meyrick (*Parthenodes*) 123
oxygona Meyrick (*Scoparia*) 43
oxygramma Turner (*Cosmophylla*) 106
oxythyma Meyrick (*Scoparia*) 52

pacalis Grote (*Nymphula*) 91
pacalis Leech (*Campomastix*) 181
pachyerga Meyrick (*Scoparia*) 51
pachysema Meyrick (*Scoparia*) 38
pachyzancloides Dognin (*Pelina*) 130
pacificalis Dyar (*Scoparia*) 34
pacificalis Schaus (*Stenia*) 171
paediualis Walker (*Stenia*) 166, 167
paigniodesalis Dyar (*Ambia*) 116
palleuca Hampson (*Oligostigma*) 118
pallida Banks (*Scoparia*) 26
pallida Butler (*Nymphula*) 87
pallida Stephens (*Scoparia*) 29
pallidalis R. South (*Parthenodes*) 124
pallidalis Warren (*Bradina*) 150
pallidimarginalis (ab.) Warren (*Scoparia*) 30
pallidula Philpott (*Scoparia*) 51
palliolatalis Swinhoe (*Ambia*) 111

palliolatalis Hampson (*Ambia*) 111
palloralis (ab.) Dyar (*Scoparia*) 34
paltomacha Meyrick (*Scoparia*) 51
paludalis Schrank (*Nymphula*) 79
paludata Fabricius (*Nymphula*) 79

pampalis Schaus (*Aulacodes*) 146
pampenealis Dyar (*Nymphula*) 93
panaeusalis Walker (*Bradina*) 150
pandaralis Walker (*Coelorhynchidia*) 159

pandora Meyrick (*Eranistis*) 133
panmicta Turner (*Eclipsiodes*) 9
panopla Meyrick (*Scoparia*) 52
pantheralis Walker (*Argyrectis*) 68

papulalis Snellen (*Eoöphila*) 138
parachalca Meyrick (*Scoparia*) 54
parachlora Meyrick (*Scoparia*) 39
parachrysis Meyrick (*Ambia*) 114
paracycla Lower (*Tetraprosopus*) 55

paradoxa E. Hering (*Ridleyana*) 122
parallela Meyrick (*Bradina*) 155
parallelalis Hampson (*Parthenodes*) 125
parallelalis Sauber (*Parthenodes*) 124

paralleloidalis Klima (*Parthenodes*) 125
paraphaca Meyrick (*Osphrantis*) 136

parapomasalis Hampson (*Aulacodes*) 140

parbattoides Tams (*Bradina*) 156
parca Philpott (*Scoparia*) 55

parella Herrich-Schaeffer (*Scoparia*) 20, 24
parella Zeller (*Scoparia*) 20
parmifera Meyrick (*Scoparia*) 46
paroalis Schaus (*Cymoriza*) 108
parthenodalis Hampson (*Argyrectis*) 69

parvale Hampson (*Oligostigma*) 117

parvalis Moore (*Oligostigma*) 117
parvalis Walker (*Piletocera*) 174
parvissimalis Schaus (*Argyrectis*) 64

pascoella Philpott (*Scoparia*) 54
passalota Meyrick (*Scoparia*) 37
patnalis Felder & Rogenhofer (*Cataclysta*) 93

paulalis Schaus (*Nymphula*) 93

- pavonialis Hampson (*Argyrectis*) 65
 pedionoma West (*Ambia*) 112
 pellucidale Hampson (*Clupeosoma*) 183
 pellucidalis Snellen (*Clupeosoma*) 183, 184
 pelochra Meyrick (*Dracaenura*) 135
 pentaspila Meyrick (*Scoparia*) 38
 pentopalis Hampson (*Argyrectis*) 61
 pentopalis Hampson (*Aulacodes*) 143
 peperita Hampson (*Microphysetica*) 162
 peraltalis Schaus (*Cataclysta*) 103
 peribocalis Hampson (*Aulacodes*) 140
 peribocalis Walker (*Eoöphila*) 138
 pericompa Turner (*Cataclysta*) 99
 perierga Meyrick (*Scoparia*) 42
 periopis Hampson (*Argyrectis*) 62
 periphanes Meyrick (*Scoparia*) 48
 perirrorata Hampson (*Cataclysta*) 95
 perixantha Meyrick (*Mestolobes*) 56
 perlalis Hampson (*Nymphula*) 92
 perlucidalis Hampson (*Bradina*) 155
 pernigralis (fa.) Turati (*Scoparia*) 25
 pernumbralis Dyar (*Scoparia*) 33
 peronetus Meyrick (*Scoparia*) 41
 perornatalis Schaus (*Ambia*) 114
 perplexalis Guenée (*Scoparia*) 18
 perplexella Zeller (*Scoparia*) 18
 persinualis Hampson (*Musotima*) 105
 peruviensis Hampson (*Oligostigma*) 121
 pervenustalis Hampson (*Argyrectis*) 60
 pessias Meyrick (*Mestolobes*) 56
 petrina Meyrick (*Scoparia*) 62
 petrophila Standfuss sen. (*Scoparia*) 22
 phaealis Hampson (*Scoparia*) 30
 phaedrale Hampson (*Oligostigma*) 121
 phaedralis Walker (*Oligostigma*) 121
 phaeochroalis Hampson (*Ambia*) 115
 phaeocraspedalis Hampson (*Pileotocera*) 178
 phaeoleuca Stainton (*Scoparia*) 28
 phaeoleuca Zeller (*Scoparia*) 19
 phaeoleucalis Guenée (*Scoparia*) 19
 phacomeralis Hampson (*Ambia*) 116
 phaeopalpia Hampson (*Scoparia*) 32
 phaeopastalis Hampson (*Argyrectis*) 69
 phaeospilalis Hampson (*Stenia*) 167
 phaeozonalis Hampson (*Ambia*) 115
 phalerias Meyrick (*Scoparia*) 48
 phenice Cramer (*Zebonia*) 186
 phenice Druce (*Stenia*) 170
 phenicealis Guenée (*Zebonia*) 186
 philerga Meyrick (*Scoparia*) 45
 philetaera Meyrick (*Scoparia*) 48
 philippinensis Hampson (*Micraglossa*) 8
 philonophes Meyrick (*Scoparia*) 43
 philorhpha Meyrick (*Scoparia*) 45
 phlegetonalis Snellen (*Parthenodes*) 124
 phocosalis Schaus (*Ambia*) 103
 phoenicobapta Hampson (*Massepha*) 128
 photias Meyrick (*Pelena*) 129
 photina Tams (*Clupeosoma*) 184
 phoxopteralis Snellen (*Argyrectis*) 69
 phryganidalis Pagenstecher (*Hymenoptychis*) 137
 phycitinalis Dyar (*Scoparia*) 35
 phycophanes Meyrick (*Scoparia*) 194
 picale Hampson (*Oligostigma*) 118
 picalis Guenée (*Oligostigma*) 118
 picalis Hampson (*Ambia*) 114
 picata Butler (*Diathrausta*) 160
 picata Butler (*Perisyntrocha*) 132
 picata Leech (*Diathrausta*) 160
 picrogramma Meyrick (*Decticogaster*) 147
 pictalis Warren (*Margarochroma*) 126
 piperitalis Hampson (*Oligostigma*) 120
 pisusalis Fernald (*Diathrausta*) 161
 plagalis Moore (*Stegothyris*) 134
 plagiotis Meyrick (*Scoparia*) 43
 planilinealis Warren (*Dasyscopa*) 57
 platymera Lower (*Scoparia*) 42

- platyscia Meyrick (*Scoparia*) 41
 plebeialis Walker (*Pileocera*) 179
 plebeijfascialis W. Rothschild
 (*Dracaenura*) 136
 plenilinealis Grote (*Nymphula*)
 89
 pleonaxalis Hampson (*Nymphu-*
la) 88
 plevie Dyar (*Cataclysta*) 101
 plicatalis Walker (*Eoöphila*) 138,
 140
 plumbealis Kenrick (*Margaro-*
sticha) 94
 plumbealis Warren (*Dia-*
thrausta) 160
 plumbefusalis Hampson (*Nym-*
phula) 86
 plumbicostalis Hampson (*Pileto-*
cera) 177
 plumipedalis Hampson (*Aulaco-*
des) 145
 plusialis Herrich-Schaeffer (*Ar-*
gyractis) 66
 poasalis Schaus (*Stenia*) 171
 podalirialis Guenée (*Zebronia*)
 186
 podalivalis Hampson (*Zebronia*)
 186
 poediusalis Hampson (*Stenia*)
 167
 polialis Hampson (*Scoparia*) 30
 poliophaealis Hampson (*Scoparia*)
 19
 pollutalis Snellen (*Massepha*)
 127
 polusale Hampson (*Clupeosoma*)
 183
 polusalis Walker (*Clupeosoma*)
 183, 184
 polydectale Hampson (*Oligostig-*
ma) 120
 polydectalis Walker (*Oligostig-*
ma) 120
 polydora Meyrick (*Aulacodes*) 139
 polysemalis Hampson (*Dasyco-*
pa) 57
 polystictalis Hampson (*Nymphu-*
la) 88
 pomperialis Druce (*Cataclysta*)
 102
 pongalis Felder & Rogenhofer
 (*Scoparia*) 48
 poritialis Walker (*Ambia*) 112
 porphyroclista Meyrick (*Bradi-*
na) 156
 portlandica (var.) Stainton (*Sco-*
paria) 28
 postalbalis Hampson (*Eristena*)
 70
 postbasalis W. Rothschild (*Aula-*
codes) 144
 postbicoloralis W. Rothschild
 (*Bradina*) 156
 potamogalis Hübner (*Nymphula*)
 73
 potamogalis Treitschke (*Nym-*
phula) 75
 potamogata Linné (*Nymphula*)
 74
 praestabilis Pagenstecher (*Oligo-*
stigma) 121
 pragmatica Meyrick (*Mestolo-*
bes) 56
 premalis Druce (*Argyractis*) 68
 primolalis Schaus (*Stenia*) 169
 primordialis Dyar (*Geshna*) 159
 prinzi Schawerda (*Scoparia*) 17
 probolaea Meyrick (*Scoparia*) 41
 procelearis Meyrick (*Mestolobes*)
 57
 prodigalis Leech (*Parthenodes*)
 122, 124
 productalis Hampson (*Argyrac-*
tis) 66
 productalis Lederer (*Orphno-*
phanes) 131
 profanalis Hampson (*Stenia*) 166
 profundalis Lederer (*Diathra-*
usta) 160
 profusalis Schaus (*Oligostigma*)
 121
 promiscua Wileman & South
 (*Scoparia*) 30
 prophaelis Dognin (*Stenia*) 171
 proprialis Fernald (*Nymphula*)
 90
 prosthenialis Hampson (*Dracae-*
nura) 135
 protopennis Dyar (*Clupeosoma*)
 185
 protorthra Meyrick (*Scoparia*) 42
 pryeri Butler (*Clupeosoma*) 183
 psalmoidalis Schaus (*Argyractis*)
 66
 psammitis Meyrick (*Scoparia*) 50
 psathyrodes Turner (*Cataclysta*)
 99
 psednopa Meyrick (*Scoparia*) 45
 pseudadelpha Meyrick (*Pileoce-*
ra) 177
 pseudofodalis Strand (*Nymphu-*
la) 87
 pseudomurana (fa.) Müller-Rutz
 (*Scoparia*) 24
 pseudopelochra W. Rothschild
 (*Dracaenura*) 136
 pseudopis Dyar (*Clupeosoma*) 185
 psyllalis Guenée (*Aulacodes*) 144

- pterophoralis* Walker (*Hymenoptychis*) 137
ptolycusalis Walker (*Ambia*) 110
pudica Lucas (*Nymphula*) 88
pulchellale Hampson (*Oligostigma*) 118
pulchellalis Mabilie (*Stenia*) 167
pulcherialis Druce (*Cataclysta*) 102
pulcherrimalis Lederer (*Margarosticha*) 93
pulchralis Moore (*Hyaloplaga*) 187
pulchralis W. Rothschild (*Aulacodes*) 144
pullatalis Swinhoe (*Pileocera*) 172
pulveralis Hübner (*Psammotis*) 188
pulveralis Snellen (*Scoparia*) 30
pumiliialis Hampson (*Bradina*) 152
punctalis [Schifferm. & Denis] (*Stenia*) 164, 166, 167
punctata Swinhoe (*Niphostola*) 132
punctilinealis Hampson (*Bradina*) 132, 155
punctimarginata W. Rothschild (*Ambia*) 114
pura Philpott (*Scoparia*) 54
puralis Schaus (*Argyractis*) 64
purbeckensis (ab.) Banks (*Scoparia*) 18
purifactualis Dyar (*Oligostigma*) 122
purpurascens Hampson (*Bradina*) 156
purpurea Hampson (*Coelorhynchidia*) 159
purpurealis Kenrick (*Aulacodes*) 144
purpureofusa Hampson (*Pileocera*) 176
purvalis Hampson (*Pileocera*) 174
pusilla Dyar (*Scoparia*) 36
pusilla Rosenstock (*Scoparia*) 42
pusillalis Saalmüller (*Cataclysta*) 98
pycnolopha Tams (*Bradina*) 154
pygmaealis (fa.) Amsel (*Stenia*) 165
pyrallela Hübner (*Scoparia*) 18
pyrenaecalis (ab.) Duponchel (*Scoparia*) 16
pyrenaecella Duponchel (*Scoparia*) 16
pyrenaicalis Guenée (*Scoparia*) 16
pyropa Meyrick (*Promylaea*) 55
pyropalis Guenée (*Argyractis*) 68
pyrseutis Meyrick (*Scoparia*) 40
quadralis Snellen (*Pileocera*) 175
quadratella Zetterstedt (*Scoparia*) 11
quadrifascia Swezey (*Promylaea*) 55
quadrifasciata Swezey (*Mestolobes*) 57
quadriplagiata Hampson (*Aulacodes*) 142
quaestoria Meyrick (*Scoparia*) 48
quaternalis Lederer (*Aediodina*) 180
queenslandica Hampson (*Cataclysta*) 98
quinquinalis Snellen (*Oligostigma*) 119
ragonoti Butler (*Scoparia*) 35
rakaiensis Knaggs (*Scoparia*) 49
ranalis Hampson (*Pileocera*) 179
randalis Druce (*Omphaloptera*) 108
ranica (var.) Strand (*Scoparia*) 24
rechingeri Tams (*Pileocera*) 176
reconditalis (fa.) Walker (*Dithrausta*) 161
rectangulalis Caradja (*Massepha*) 128
rectangulalis Kenrick (*Parthenodes*) 125
rectiferalis Walker (*Bradina*) 154
rectilinea Zeller (*Scoparia*) 32
rectilinealis R. South (*Bradina*) 151
reducta (var.) Caradja (*Hyaloplaga*) 187
reducta Walker (*Pileocera*) 173, 177
reductalis Caradja (*Aulacodes*) 141
refugalis Hulst (*Scoparia*) 32
regularis Fryer (*Nymphula*) 80
religiosa Meyrick (*Scoparia*) 41
remipes Hampson (*Bradina*) 151
renatalis Meyrick (*Gargela*) 59
renatusalis Walker (*Gargela*) 59
repercussa Philpott (*Scoparia*) 49
repetitalis Warren (*Cataclysta*) 99
resinalis Guenée (*Scoparia*) 25
resinea Haworth (*Scoparia*) 25
resinodes De Joannis (*Scoparia*) 32

- responsalis Walker (*Nymphula*)
 84, 87
 retractalis Hampson (*Stenia*) 167
 reversalis Dyar (*Aulacodes*) 146
 rhabdotalis Hampson (*Ambia*)
 111
 rhodea Lower (*Clupeosoma*) 184
 rhombias Meyrick (*Scoparia*) 37
 rhomboidalis Walker (*Galadra*)
 187
rhomboidata Swinhoe (*Galadra*)
 187
 rhopalophora Meyrick (*Pileto-*
cera) 175
 ridopalis Swinhoe (*Coptobasis*)
 222
 rigidalis Barnes & Mc Dunnough
 (*Scoparia*) 32
 rinconadalis Schaus (*Argyrectis*)
 65
 rivularis Duponchel (*Nymphula*)
 72
 rivularis Meyrick (*Pileocera*)
 123
rivulata Scopoli (*Nymphula*) 75
 romalis Druce (*Massepha*) 128
 romanalis Druce (*Cataclysta*) 102
 rothschildi Klima (*Ambia*) 114
 rotuella Felder & Rogenhofer
 (*Scoparia*) 52
 rotundalis Hampson (*Pileocera*)
 178
rubetra (fa.) D. Lucas (*Psam-*
motis) 188
rubralis Warren (*Clupeosoma*)
 183
 rubriceps Hampson (*Stenia*) 166
 rufalis Hampson (*Parthenodes*)
 125
 rufalis Hampson (*Zagiridia*) 182
 rufescens Hampson (*Massepha*)
 128
 ruficeps Hampson (*Pileocera*)
 177
 rufistriata Hampson (*Clupeoso-*
ma) 184
 rufiterminalis Hampson (*Oligo-*
stigma) 120
 rufitincta Hampson (*Ambia*) 114
 rufitinctalis Hampson (*Scoparia*)
 34
 rufocastanea W. Rothschild (*Au-*
lacodes) 144
 rufostigma Hampson (*Scoparia*)
 30
 rufulalis Hampson (*Pileocera*)
 176
 rugosalis Möschler (*Nymphula*)
 93
 rusina Druce (*Clupeosoma*) 184
rusinum Hampson (*Clupeosoma*)
 184
 rupestris Meyrick (*Scoparia*) 194
 sabulosella Walker (*Scoparia*) 51
sabura Druce (*Scoparia*) 33
sacadasalis Walker (*Oligostigma*)
 118
sachalinensis Matsumura (*Scopa-*
ria) 27
sagittalis Swinhoe (*Bradina*) 152
sambesica Strand (*Argyrectis*)
 62
samealis Dyar (*Stenia*) 171
samealis Felder & Rogenhofer
 (*Argyrectis*) 67
sanguinsecta Swinhoe (*Oligostig-*
ma) 119
saponalis Guenée (*Stenia*) 171
saternalis Dyar (*Cataclysta*) 99
saturalis Hampson (*Aulacodes*)
 140
saturatalis Snellen (*Aulacodes*)
 140
saurialis Guenée (*Stenia*) 170
scalaris Schaus (*Cataclysta*) 103
scalpellalis Pagenstecher (*Hy-*
menoptychis) 137
schaefferalis Dyar (*Cataclysta*)
 101
schausalis Dyar (*Clupeosoma*) 185
schausi Hampson (*Pseudolithosia*)
 108
schistochaeta Tams (*Ambia*) 114
schistopalis Hampson (*Argyrec-*
tis) 67
schizodesma Lower (*Scoparia*) 45
schwarzalis Dyar (*Scoparia*) 33
schwarzalis Schaus (*Argyrectis*)
 64
sciaphilella De la Harpe (*Scop-*
aria) 20
scintillifera Turner (*Hylebatis*)
 193
scintillulalis Wollaston (*Scoparia*)
 31
scioxantha Meyrick (*Argyrectis*)
 194
scitalis Lederer (*Nymphula*) 77
scleropis Meyrick (*Mesolobes*) 56
scoparialis Warren (*Micraglossa*)
 8
scoriella Wollaston (*Scoparia*) 23
scotalis Hampson (*Parthenodes*)
 124
scotica White (*Scoparia*) 12
scotochroa Hampson (*Pileocera*)
 176

- scripta* Philpott (*Scoparia*) 53
scutthesalis Walker (*Aulacodes*) 144
secutalis Walker (*Aulacodes*) 143
sejunctalis Snellen (*Aulacodes*) 140
selectalis Lederer (*Bradina*) 151
selenias Turner (*Baeoptila*) 106
semialbalis W. Rothschild (*Dra-caenura*) 136
semialbida Warren (*Parthenodes*) 124
semiamplalis Warren (*Scoparia*) 23
semichroa Butler (*Mestolobes*) 56
semicircularis Hampson (*Aulacodes*) 145
semifascialis Warren (*Ambia*) 111
semifuscalis Hampson (*Stenia*) 169
semigilva Turner (*Eclipsiodes*) 9
semilunalis Hampson (*Ambia*) 144
semimarginalis Dyar (*Oligostigma*) 122
semineale Hampson (*Nymphula*) 90
seminealis Walker (*Nymphula*) 89
seminivella Walker (*Nymphula*) 89
semizebralis Walker (*Zebronia*) 187
semnopa Meyrick (*Bradina*) 154
separatalis (subsp.) Leech (*Nymphula*) 76
serapionalis Schaus (*Argyractis*) 66
sericea Butler (*Pelena*) 129
seriopunctalis Hampson (*Nymphula*) 86, 88
serrilinealis Barnes & Benjamin (*Nymphula*) 90
sibirica Lederer (*Scoparia*) 19
sicaria Meyrick (*Mestolobes*) 57
sideraspis Meyrick (*Scoparia*) 53
siderina Meyrick (*Scoparia*) 39
siennata Warren (*Aulacodes*) 142
signella Teich (*Scoparia*) 15
signiferalis Wallengren (*Piletocera*) 175, 176
silacella Hübner (*Cholius*) 9
simacethina Butler (*Mestolobes*) 56
simialis Gaede (*Temnobasis*) 158
similis Hampson (*Diathraustodes*) 158
similis W. Rothschild (*Aulacodes*) 143
similis Wollaston (*Scoparia*) 31
simplalis Snellen (*Nymphula*) 86
simplex West (*Aulacodes*) 140
simplicella De la Harpe (*Cholius*) 10
simplicialis Barnes & Mc Dounough (*Piletocera*) 179
simplicialis Snellen (*Aulacodes*) 140
sinensis (subsp.) Caradja (*Bradina*) 152
sinensis Hampson (*Aulacodes*) 140
sinicalis Hampson (*Nymphula*) 84
sinitalis Schaus (*Argyractis*) 64
sinuata Philpott (*Scoparia*) 54
sirina Meyrick (*Mestolobes*) 56
slossonalis Dyar (*Cataclysta*) 100
smithi Druce (*Scoparia*) 33
snelleni Semper (*Eoöphila*) 138
sobrina Pryer (*Eoöphila*) 138
sodalis Lederer (*Bradina*) 150
sodalis Leech (*Piletocera*) 172, 173
solitalis Walker (*Nymphula*) 77
sordida Zeller (*Hymenoptychis*) 137
sordidalis Dewitz (*Bradina*) 153
sordidior (subsp.) W. Rothschild (*Nymphula*) 83
spaldingalis Barnes & Mc Dounough (*Scoparia*) 33
spectacularis Meyrick (*Scoparia*) 45
spelaea Meyrick (*Scoparia*) 42
sphenactis Meyrick (*Strepsinoma*) 133
sphenotis Meyrick (*Margarosticha*) 93
sphragidacma Meyrick (*Argyractis*) 194
spilomelalis Walker (*Nymphula*) 84, 87
spissalis Guenée (*Psammotis*) 189
splendens West (*Aulacodes*) 141
spodinopa Meyrick (*Stenia*) 166
spodinopella Strand (*Stenia*) 166
stagnalis Guenée (*Nymphula*) 73
stagnalis Zeller (*Nymphula*) 82
stagnata Donovan (*Nymphula*) 72
stauderi (fa.) Mitterberger (*Scoparia*) 20
staudingeralis Mabilie (*Scoparia*) 19
steffanyi Tams (*Piletocera*) 176

stellata Warren (*Parthenodes*)
123
stenialis Warren (*Nymphula*) 86
stenialis Warren (*Pileotocera*) 175
steniodes Butler (*Nymphula*) 88
stenoiperalis Hampson (*Pileotocera*) 178
stenomma Felder & Rogenhofer
(*Dracaenura*) 135
stenopa Lower (*Scoparia*) 45
stenota Warren (*Scoparia*) 23
stenota Wollaston (*Scoparia*) 23
stercoralis Möschler (*Pileotocera*)
179
stereostigma Dyar (*Scoparia*) 36
steropaea Meyrick (*Scoparia*) 51
stictochroa Turner (*Musotima*)
106
stigmatalis Dognin (*Omphaloptera*) 108
stigmophanes Meyrick (*Bradina*)
156
stigmatosalis Herrich-Schaeffer
(*Stenia*) 165
stipalis Snellen (*Cataclysta*) 95
stötzneri Caradja (*Scoparia*) 31
straminealis Hampson (*Eristena*)
70
straminealis Hampson (*Scoparia*)
31
stratitotalis Schiffermüller & Denis
(*Nymphula*) 79
stratitotata Linné (*Nymphula*) 78
stresemanni W. Rothschild (*Aulacodes*) 141
striatilis Dyar (*Ambia*) 114
striatalis Hampson (*Eclipsiodes*)
9
strigalis Dyar (*Scoparia*) 33
strigatalis Pagenstecher (*Orphnophanes*) 131
strigatalis W. Rothschild (*Clupeosoma*) 184
strigigramma Hampson (*Scoparia*)
35
struthias Meyrick (*Scoparia*) 38
stygalis Hampson (*Pileotocera*)
176
styxalis Schaus (*Stenia*) 168
subaurealis Walker (*Daulia*) 126
subcervinalis Walker (*Arxama*)
71
subditella Walker (*Scoparia*) 55
subfusca Stephens (*Scoparia*) 12
subita Philpott (*Scoparia*) 55
submarginalis Walker (*Scoparia*)
49
submedinella Caradja (*Scoparia*)
30

subornata Hampson (*Argyrectis*)
69
subpurella Walker (*Gargela*) 59
subpurpurascens Warren (*Bradina*) 152
subtersa Dyar (*Scoparia*) 36
sudetica Hampson (*Scoparia*) 22
sudetica Zeller (*Scoparia*) 22
sudeticalis Duponchel (*Scoparia*)
22
sufetulodes Hampson (*Ambia*)
114
sufflexalis Dyar (*Clupeosoma*)
185
suffusa Kenrick (*Perisyntrocha*)
132
suffusalis Hampson (*Musotima*)
104
sulcialis Walker (*Coptobasis*) 157
sultschana Ragonot (*Nymphula*)
78
sumptuosalis Möschler (*Cataclysta*) 103
supercilialis Hampson (*Argyrectis*) 61
susannae Lower (*Scoparia*) 45
sutschana Hampson (*Nymphula*)
78
syagrusale Hampson (*Oligostigma*)
117
syagrusalis Walker (*Oligostigma*)
117
sylvestris Clarke (*Scoparia*) 55
symphonalis Dyar (*Argyrectis*)
66
symphorasalis Schaus (*Ambia*)
116
synapta Meyrick (*Scoparia*) 42
syngamoides Hampson (*Massepha*) 128
synophanoses Klunder (*Symphonia*) 104
syntaracta Meyrick (*Scoparia*)
42
syriaca (var.) Caradja (*Scoparia*) 16
szechuanalis (var.) Caradja (*Oligostigma*) 119

tabidalis Lederer (*Bradina*) 150
taeniatella Teich (*Scoparia*) 14
tafirella Chrétien (*Scoparia*) 26
tigiadalis Hampson (*Dracaenura*) 136
taiwanalis Shibuya (*Cymoriza*)
108
taiwanalis (var.) Wileman & South
(*Pileotocera*) 124

- takamukai Shibuya (*Nymphula*) 83
 talboti (var.) Caradja (*Diatkrausta*) 160
 talboti Caradja (*Massepha*) 128
 talboti Schmidt (*Scoparia*) 23
 tamanalis Schaus (*Argyrectis*) 65
 tamsi Klima (*Cataclysta*) 99
 tanyscia West (*Decticogaster*) 147
 tapaiosalis Schaus (*Argyrectis*) 67
 taprobana Hampson (*Genoloba*) 192
 tedyscongalis Clemens (*Nymphula*) 90
 tellesalis Hampson (*Piletocera*) 174
 tellesalis Walker (*Piletocera*) 174
 templalis Schaus (*Aulacodes*) 146
 tendicularis Rebel (*Ambia*) 113
 tenebralis Lower (*Nymphula*) 88
 tenebrifera Hampson (*Eristena*) 70
 tenebrosa W. Rothschild (*Epistenia*) 163
 tenebrosalis Hampson (*Ambia*) 110
 tenebrosalis W. Rothschild (*Stenia*) 167
 tenella Hampson (*Ambia*) 114
 tenellalis Guenée (*Nymphula*) 77
 terranea W. Rothschild (*Nymphula*) 88
 terminitis Meyrick (*Tetrernia*) 109
 tersella Zeller (*Scoparia*) 33
 tessellalis Hampson (*Cataclysta*) 102
 tessimalis Dyar (*Argyrectis*) 70
 tessmanni Gaede (*Massepha*) 128
 tetracycla Meyrick (*Scoparia*) 49
 tetralitha Hampson (*Eristena*) 70
 tetranesa Meyrick (*Scoparia*) 37
 tetrommata Hampson (*Metaclysta*) 104
 tetropalis Hampson (*Argyrectis*) 61
 teutalis Walker (*Bradina*) 151
 thalamias Meyrick (*Scoparia*) 38
 thermichrysis Hampson (*Aulacodes*) 143
 thetysalis Walker (*Aulacodes*) 145
 thiogramma Meyrick (*Malaciotis*) 194
 thoasalis Walker (*Orphnophanes*) 131
 threnodes Meyrick (*Scoparia*) 45
 thyellopis Meyrick (*Scoparia*) 40
 thyridialis Lederer (*Ambia*) 110
 thyridias Meyrick (*Scoparia*) 45
 ticinensis (fa.) Müller-Rutz (*Scoparia*) 19
 ticonalis Dyar (*Argyrectis*) 69
 tigrimalis Hampson (*Ambia*) 112
 tigrinalis Dyar (*Oligostigma*) 122
 timaralis Felder & Rogenhofer (*Musotima*) 105
 tipuliformis Walker (*Hymenoptychis*) 137
 titanalis Hampson (*Nymphula*) 86
 tormentifera Meyrick (*Bradina*) 156
 torniplagalis Dyar (*Scoparia*) 35
 torodes Meyrick (*Scoparia*) 48
 torridalis Kenrick (*Dracaenura*) 136
 torsicostalis Hampson (*Piletocera*) 172, 177
 tortalis Lederer (*Aulacodes*) 144
 translinealis Hampson (*Bradina*) 151
 transparitalis Karsch (*Glaucoda*) 180
 transversalis Costa (*Nymphula*) 75
 transversalis Lederer (*Stegothyris*) 134
 transversalis Wollaston (*Scoparia*) 31
 trapezophora Meyrick (*Scoparia*) 48
 traversalis Dyar (*Aulacodes*) 145
 trespassalis Dyar (*Argyrectis*) 70
 triacma Meyrick (*Scoparia*) 38
 triangularis R. South (*Parthenodes*) 124
 trichoceralis Hampson (*Ambia*) 141
 trichostylalis Hampson (*Ambia*) 112
 tricitra Meyrick (*Scoparia*) 45
 triclera Meyrick (*Scoparia*) 46
 tricolor Zeller (*Scoparia*) 36
 tricoloralis Dyar (*Scoparia*) 34
 trifidialis Hampson (*Coelorrhynchidia*) 159
 triflexalis Gaede (*Stenia*) 167
 trigonalis Hampson (*Aulacodes*) 142
 trigonalis Hampson (*Eristena*) 70
 trigonalis Swinhoe (*Cataclysta*) 98

- trilinealis Hampson (*Gargela*) 59
 trilinealis Walker (*Luma*) 130
 trilinealis Warren (*Cataclysta*) 102
 trimacula Hampson (*Cataclysta*) 97
 trimaculata Hampson (*Luma*) 130
 triopalis Hampson (*Argyrectis*) 61
 triplaga Lower (*Aulacodes*) 144
 tripletale Hampson (*Oligostigma*) 120
 tripunctalis Costa (*Nymphula*) 75
 tripunctalis Snellen (*Aulacodes*) 141
 tripunctalis Snellen (*Cataclysta*) 98
 tripunctalis Walker (*Eoöphila*) 138
 triscelis Meyrick (*Scoparia*) 48
 trispila Meyrick (*Bradina*) 156
 tristalis Schaus (*Argyrectis*) 64
 tristicta Turner (*Scoparia*) 45
 tristrigella (ab.) Stephens (*Scoparia*) 18
 tritocirra Turner (*Scoparia*) 45
 triumphalis Schaus (*Argyrectis*) 64
 trivirgata Felder & Rogenhofer (*Scoparia*) 52
 trivitalis Warren (*Symphonia*) 104
 truckeealis Dyar (*Cataclysta*) 99
 truncatalis Mc Dunnough (*Scoparia*) 34
 truncatalis Walker (*Almonia*) 148
 truncicolella Butler (*Scoparia*) 30
 truncicolella Stainton (*Scoparia*) 26
 trygoda Meyrick (*Psammotis*) 189
 tuicana Clarke (*Scoparia*) 55
 tullialis Walker (*Nymphula*) 87
 tuoniana (var.) Hoffmann (*Scoparia*) 24
 turbata Butler (*Nymphula*) 77, 85
 turbatalis Christoph (*Camptomastix*) 181
 turiatalis Schaus (*Stenia*) 169
 turneri Philpott (*Scoparia*) 51
 tyrantes Meyrick (*Luma*) 130
 tyraula Meyrick (*Scoparia*) 38
 tyrophanta (*Scoparia*) 32
 ulfridalis Schaus (*Argyrectis*) 65
 uliginata Fabricius (*Cataclysta*) 96
 ulmaya Dyar (*Scoparia*) 37
 ulmella Knaggs (*Scoparia*) 15
 ulophanes Meyrick (*Piletocera*) 178
 umbralis Dognin (*Syntrita*) 157
 umbrata (fa.) Meder (*Nymphula*) 75
 unguiculis Snellen (*Nymphula*) 86
 unicolor Moore (*Pelema*) 129
 unicoloralis Kenrick (*Dracaenura*) 136
 uniflexalis Mabille (*Stenia*) 167
 uniformis Hampson (*Pelema*) 129
 unilinealis Dyar (*Cataclysta*) 103
 unilinealis Snellen (*Oligostigma*) 118
 uniplagiatis W. Rothschild (*Aulacodes*) 144
 upupalis Guenée (*Piletocera*) 125
 urticalis Schiffermüller & Denis (*Eurrhypara*) 190
 urticata Linné (*Eurrhypara*) 190
 ussuriensis Rebel (*Nymphula*) 80, 85
 ustalis Walker (*Cymoriza*) 108
 ustimacula Felder & Rogenhofer (*Scoparia*) 47
 ustiramis Meyrick (*Scoparia*) 55
 uxorialis Strand (*Nymphula*) 86
 vacuolata Dyar (*Cataclysta*) 103
 vafra Meyrick (*Scoparia*) 22
 vagalis Walker (*Parthenodes*) 123
 vagilinealis Hampson (*Ambia*) 113
 valesialis Hampson (*Scoparia*) 20, 21
 vallesialis Duponchel (*Scoparia*) 20
 valstanalis Schaus (*Argyrectis*) 64
 vandaliella Herrich-Schaeffer (*Scoparia*) 25
 varanesalis Walker (*Dectiogaster*) 147
 vandaliella Hampson (*Scoparia*) 25
 velatalis Snellen (*Mabra*) 191
 venosa Butler (*Scoparia*) 41
 versicolor Pagenstecher (*Dracaenura*) 135
 vestigialis Snellen (*Nymphula*) 82

- vestigialis Warren (*Piletocera*) 177
 vesuntialis Guenée (*Scoparia*) 20
 vibrata Meyrick (*Decticogaster*) 148
 villidale Hampson (*Oligostigma*) 118
 villidalis Walker (*Oligostigma*) 118, 120
 vinasalis Dyar (*Scoparia*) 36
 vinotinctalis Hampson (*Scoparia*) 31
 viola Butler (*Piletocera*) 174, 178
 violalis Lederer (*Piletocera*) 173, 177
 violascens Hampson (*Clupeosoma*) 183
 violascens Hampson (*Piletocera*) 177
 virescens Rebel (*Scoparia*) 24
 virginalis Dognin (*Omphaloptera*) 109
 vitrinalis Felder & Rogenhofer (*Nymphula*) 93
 vittalis Bremer (*Nymphula*) 80, 85
 vittatalis Dyar (*Nymphula*) 90
 volcanalis Schaus (*Argyractis*) 64
 votalis Walker (*Nymphula*) 85
 vulpecula Meyrick (*Scoparia*) 50
 waigaoalis Swinhoe (*Eoöphila*) 138
 wollastoni Bethune-Baker (*Scoparia*) 23
 wollastoni W. Rothschild (*Aulacodes*) 144
 xanthalis Hampson (*Bradina*) 155
 xanthobathra Meyrick (*Cataclysta*) 95
 xanthocasis Meyrick (*Gargela*) 60
 xantholeuca Hampson (*Ambia*) 110
 xanthomela De Joannis (*Scoparia*) 31
 xanthoscia Meyrick (*Mestolobes*) 56
 xanthosoma Meyrick (*Piletocera*) 177
 xanthozonata Hampson (*Lasio-ggia*) 162
 x-signata Filipjev (*Scoparia*) 27
 xysmatias Meyrick (*Scoparia*) 47
 ycarda Dyar (*Scoparia*) 36
 zamoralis Schaus (*Argyractis*) 67
 zebinalis Walker (*Piletocera*) 174
 zelleri Rebel (*Scoparia*) 19
 zelleri (var.) Wocke (*Scoparia*) 13
 zelota Dyar (*Cataclysta*) 103
 zernyi (subsp.) Bryk (*Stenia*) 165
 zoilusalis Druce (*Nymphula*) 92
 zonulalis Snellen (*Decticogaster*) 147
 zophochlaena Meyrick (*Scoparia*) 46
 zophochlora Meyrick (*Scoparia*) 40

- In Druck:
- 85: H. Schüssler, Eupterotidae, Epicopeiidae.
86: H. Stichel, Nymphalidae I.
87: H. H. Shepard, Hesperidae: Subfam. Hesperinae II.
88: M. Gaede, Oecophoridae.

In Vorbereitung:

F. Bryk,	Noctuidae,	Arctiidae:	Microarctiinae.
W. A. Collier,	Geometridae (partes ultimae.)		
J. Draeseke,	Lycaenidae	[Continuatio].	
M. Gaede,	Incurvariidae	Opostegidae	Elachistidae
	Tischeriidae	Tineidae	Cosmopterygidae
	Heliozelidae	Colcophoridae	Lyonetiidae
	Nepticulidae	Momphidae	Ceniosomidae.
A. Klima,	Pyrilidae:	Subfam. Pyraustinae,	Phycitinae, Aner-
	stiinae, Schoenobiinae,	Macropiratinae,	Crambinae, Galle-
	riinae, Macrothecinae.		
H. Schüssler,	Thaumetopoeidae,	Bombycidae,	Endromididae,
	Morphidae.		
H. H. Shepard,	Hesperiidae	[Continuatio].	
† H. Stichel,	Nymphalidae	[Continuatio].	
H. Zerny,	Pyrilidae:	Subfam. Pyralidinae,	Endrotrichinae,
	Chrysauginae,	Pococerinae.	

Nunmehr liegt vollständig vor:

Die Blatt-Minen Mittel- und Nord-Europas

einschließlich Englands

von

Professor **Dr. Martin Hering**
(Universität Berlin)

**Bestimmungs-Tabellen der von Insekten-
Larven aller Ordnungen erzeugten Minen in
den wildwachsenden und in den im Haushalt
des Menschen kultivierten Pflanzen.**

XII und 631 Seiten mit 7 Tafeln und
500 Text-Abbildungen

Minen von
Apfelblatt-Motten

II Subscriptions-Preis des gesamten Werkes
brosch. RM. 66.—, gebunden RM. 71.—

Die Urteile der Fachgelehrten darüber sind:

Prof. Dr. M. Wolff-Eberswalde (*Der Deutsche Forstwirt* 1935, Bd. 17, pag. 1173 und 1936, Bd. 18, pag. 618): „... Das erste brauchbare Bestimmungswerk... Unfehlbarkeit der Führung... — Wir Zoologen schätzen das Werk als wichtigste und willkommenste Neuerscheinung der letzten Jahrzehnte.“

F. Heikertinger-Wien (*Koleopterologische Rundschau* 1936, Bd. 21, Nr. 6, pag. 238): „... eines der wertvollsten und willkommensten Werke, die auf entomologischem Gebiete in den letzten Jahren erschienen sind, eine Grundlage für alle Weiterforschung... ungeheuerere Bedeutung...“

Adolf Hoffmann-Wien (*Entomologischer Anzeiger* 1935, Nr. 45, pag. 292): „... Meisterhand in der Behandlung dieses schwierigen Stoffes... ein Werk deutscher Gründlichkeit, deutschen Fleißes und Könnens...“

Prof. Dr. K. Escherich-München (*Zeitschrift f. angewandte Entomologie* 1936, pag. 655): „Für uns angewandte Entomologen wird es einen wichtigen Platz im täglich benützten Handapparat einnehmen.“

Lieferung 1 wird auf Wunsch gern zur Ansicht zugesandt.

Bitte Prospekt zu verlangen.