59
801
FL
42
56

17 F

THE
- 1705 -
PUBLISHED BY

PLANTÆ FAVERSHAMIENSES.

A
C A T A L O G U E
OF THE
MORE PERFECT
P L A N T S
Growing spontaneously about
FAVERSHAM,
IN THE
COUNTY OF KENT.

WITH AN
A P P E N D I X
Exhibiting a SHORT VIEW of the
FOSSIL BODIES
OF THE ADJACENT
ISLAND OF SHEPEY.

By EDWARD JACOB, Esq; F.S.A.

Consider the Lillies of the Field how they grow — Matt. vii. 28.

L O N D O N :

Printed for the AUTHOR; by J. MARCH, on Tower-Hill.

Sold by B. WHITE, at HORACE'S-HEAD, FLEET-STREET;
and T. EVANS, PATER-NOSTER-ROW.

M,DCC,LXXVII.

AROLD
AROLD YOUNG
HARVARD
UNIVERSITY

Nov. 1912

27759

РОДИТЕ
МУЗЕУМ
ОБРАЗОВАН
УЧЕНИЦИ

T O
Mr. JOHN WHITE,
MERCHANT OF LONDON,
IN A GRATEFUL SENSE OF HIS
LONG AND SINGULAR
FRIENDSHIP.

THIS LITTLE WORK
IS MOST RESPECTFULLY INSCRIBED
BY HIS EVER OBLIGED,
AND VERY HUMBLE SERVANT,

EDWARD JACOB.

Faversham,
Aug. 10, 1777.

*Nescio quid fascinauerit nostros ut tota mens
trahatur ad peregrinas plantas, illæ vero quas
pedibus conculcamus, omnino relinquuntur.*

C. LINNÆI Bibl. Botan. p. 24.

P R E F A C E.

THE ensuing Collection of PLANTS was begun many Years since upon the Basis of the late Rev. Mr. Bateman's Catalogue, with whom, in the early part of Life, the Author made several Botanical Excursions hereabouts; and it hath gradually kept increasing ever since, as the constant Visits, in the way of his Profession, to different Parts, gave him opportunities of searching after them. It was ever intended for his own private Use and instructive Entertainment, till lately, at the recommendation of several Gentlemen well versed in Botany, it hath been revised and digested, and is now offered to the Lovers of that Science, as a means, however small, of promoting the topical Knowledge of English Plants; if it shall prove agreeable to their Taste, he will think himself amply rewarded for his Pains.

The great Promoter of its publication was the late worthy Richard Warner, Esq; of Woodford-Row, in Essex, who not only permitted the
free

P R E F A C E.

free Use of his Plantæ Woodfordienses, (without the Aid of which, and Mr. Hudson's excellent Flora Anglica, it would never have been undertaken) but also assisted the Author in the Prosecution of it, and above all, introduced him to the acquaintance of that accurate Botanist, Mr. Alchorne, Assay-Master of the Mint, whose kind Informations and Corrections have been of the greatest Advantage.

The principal Inducement for the Publication was to rectify the Account given in Mr. Blackston's Specimen Botanicum, taken from Mr. Bateman's M.S. Catalogue of near Two Hundred Plants growing about this Town, because many of the rare ones at this time, are not to be found in the Places therein referred to, which have of late been often diligently searched in hope of retrieving them, as they would have added much to this Work, but in vain.

In the Course of about Seventy Years since that Catalogue was made, many Causes and Accidents may be assigned for the loss of some of the rare Plants there mentioned, yet none greater than the industrious hand of improving Husbandry. Mr. Bateman might possibly have been mistaken in others,

P R E F A C E.

others, and possibly too some may be overlooked by the Author, who is not so conceited as to think he hath discovered all the Plants growing round him. Those Gentlemen who are fond of the Study must be sensible, that a single Person, whose necessary attention in the different Branches of his Profession was fully engaged, could not afford sufficient Leisure, however fervent his Inclination might be, to search at proper Seasons every corner of a District Five or Six Miles Diameter, when even the great Mr. Ray could make considerable Additions to the first Catalogue of his Cambridgeshire Plants, and yet leave many unobserved, to be added thereto by Dr. Martyn, and others.

In order, however, to supply his Deficiencies, the Author hath inserted some few rare Plants discovered at a greater distance in the Eastern parts of the County, either by himself, or communicated to him by his obliging Friends, Mr. Alchorne, the Rev. Mr. Jones, of Pluckley, and Mr. Boys of Sandwich, for which Favours he begs they will be pleased to accept his most grateful Acknowledgements.

As

P R E F A C E.

As he hath composed his Catalogue upon Mr. Warner's alphabetical Plan, he hopes the Preference given to Mr. Ray's Names will be pardoned by the more scientific Botanists, especially as he hath printed an Index of those of Linnæus.

The Appendix, giving a short Account of the Shepey Fossils, it is presumed will not be deemed an improper Addition, by the Lovers of Natural History, as hitherto no particular one hath been offered to their Notice, although so well deserving of it.

The Plate of the Author is the unexpected Donation of an obliging Friend.

The ABBREVIATIONS explained.

R. S. J. Raii Synopsis Stirpium Britannicarum.
Ed. III. 1724.

H. F. Gul. Hudsoni Flora Anglica. 1762.

A. Annual. B. Biennial. P. Perennial. S. Shrub, or Tree.

The Month (as March, &c.) denotes the Time of the Plants flowering.

* This Mark placed before a Plant, although found wild, shews it most probably to have originated from Seed dispersed by Birds, or otherwise from Gardens.

 PLANTÆ FAVERSHAMIENSES.

A

ABSINTHIUM marinum album. R. S. 188.

ARTEMISIA *maritima*. H.F. 311.

White Sea-Wormwood.

In the Salt Marshes—very common. August. P.

ABSINTHIUM vulgare. R. S. 188.

ARTEMISIA *Absinthium*. H.F. 311.

Common Wormwood.

In Farm-yards and on dry banks—common. August. P.

* ACER majus. R. S. 407.

ACER *Pseudo-Platanus*. H.F. 378.

The greater Maple or Sycamore Tree.

In a Hedge near Brenly and Hanslets Forstal — very uncommon. May. S.

ACER minus. R. S. 470.

ACER *campestre*. H. F. 379.

The Common Maple.

In Hedges—very common. May S.

B

ABRACINOS.

ABRACINUM

ABRACINUM

ABRACINUM

ACINOS. R. S. 238.

THYMUS *Acinos*. H. F. 230.

Wild Basil.

On Badgen Downs — plentifully. July. A.

ADIANTUM nigrum Officinarum. R.S. 126.

ASPLENIUM *Adiantum-nigrum*. H.F. 386.

Common black Maiden-hair.

On the hollow sandy Banks in Byfing Wood and
Hernhill—not common. P.

AGRIFOLIUM. R. S. 466.

ILEX *Aquifolium*. H. F. 505.

The Holly Tree.

In Woods and Hedges—common. May. S.

AGRIMONIA. R. S. 202.

AGRIMONIA *Eupatoria*. H. F. 180.

Agrimony.

On the Sides of Fields and under Hedges — very
common. June. S.

ALCEA vulgaris. R. S. 252.

MALVA *Alcea*. H. F. 268.

Vervain Mallow.

In Pastures at Graveney and Hernhill — not com-
mon. August. P.

ALLIUM

ALLIUM sylvestre. R. S. 369.

ALLIUM vineale. H. F. 121.

Crow-Garlick.

On the Wall that leads to Thorn Creek — plentifully.
June. P.

ALLIUM sylvestre latifolium. R. S. 370.

ALLIUM ursinum. H. F. 122.

Ramsons.

Amongst the Alders near Hernhill-Church — plentifully. May. P.

ALNUS. R. S. 442.

BETULA *Alnus*. H. F. 506.

The Common Alder.

In moist Places—very common. July. S.

ALSINE hirsuta Myosotis. R. S. 349.

CERASTIUM *vulgatum*. H. F. 175.

Narrow-leaved Mouse-ear Chickweed.

In Fields and Meadows—common. May. A.

ALSINE hirsuta Myosotis latifolia præcocior.

R. S. 348.

CERASTIUM *viscosum*. H. F. 175.

The broader leaved Mouse-ear Chickweed.

On the Walls of the Abbey—common. April. A.

ALSINE major repens perennis. R. S. 347.

CERASTIUM aquaticum. H. F. 177.

Great Marsh Chickweed.

In Davington Brooks, and at Goodnestone — not common. July. P.

ALSINE marina foliis portulacæ. R.S. 351.

ARENARIA peploides. H. F. 167.

Sea Chickweed.

On Shellnefs in Shepey—not common. July. P.

ALSINE minor multicaulis. R. S. 349.

ARENARIA serpyllifolia. H. F. 167.

The least Chickweed.

On Roofs of old Houses and on old Walls — common. May. A.

ALSINE spergula dicta major. R. S. 351.

SPERGULA arvensis. H. F. 177.

Corn Spurrey.

On sandy Grounds in Hern-Hill—common. July. A.

ALSINE spergulæ facie media. R. S. 351.

ARENARIA marina. H. F. 169.

Sea Spurrey.

On the Paths to Holly Shore—common. June. A.

ALSINE

ALSINE foliis caryophylleis. R. S. 344.

SAGINA erecta. H. F. 64.

The least Stichwort.

In Nagden Marshes—not common. May. A.

ALSINELLA muscosa flore repens. R. S. 345.

SAGINA procumbens. H. F. 63.

Pin Pearl-wort, Chickweed breakstone.

On old Walls about the Town—common. June. A.

ALTHÆA vulgaris. R. S. 252.

ALTHÆA officinalis. H. F. 267.

Marsh-Mallow.

In the Marshes at Clapgate and Goodnestone—common. August. P.

ANACAMPSEROS, vulgo Faba crassa. R. S. 269.

SEDUM Telephium. H. F. 170.

Orpine, or Live-long.

In Byfing and Cockset Woods—not common. August. P.

ANAGALLIS flore phoeniceo. R. S. 282.

ANAGALLIS arvensis. H. F. 73.

Male Pimpernel.

In Fields and Gardens—common. May—August. A.

ANAGALLIS lutea. R. S. 282.

LYSIMACHIA nemorum. H. F. 73.

Yellow Pimpernel.

In Jud's Wood, Ospringe—uncommon. June. P.

ANBLATUM Cordi five Aphyllon. R. S.*288.

LATHRÆA Squamaria. H.F. 233.

Toothwort.

Under Hedges near Mr. Jennings's House in Pluckley
—very uncommon. April. P.

ANEMONE nemorum. R. S. 259.

ANEMONE nemorosa. H. F. 208.

Wood Anemony.

In Woods and Hedges—very common. April. P.

ANGELICA sylvestris. R. S. 208.

ANGELICA sylvestris. H. F. 103.

Wild Angelica.

Near Stone-Bridge, and in Byfing Wood—not
common. July. P.

ANGELICA sylvestris minor feu erratica. R.S. 208.

ÆGOPODIUM Podagraria. H. F. 111.

Herb Gerard, Gout-weed, or Ash-weed.

In Gardens and under Hedges, near Boughton Church
—not common. May. P.

ANONIS maritima procumbens. R. S. 332.

ONONIS repens. H. F. 273.

Creeping Rest-harrow.

On the Sand Downs near Deal—uncommon. July. P.

ANONIS

ANONIS spinosa flore purpureo. R. S. 332.

ONONIS *spinosa*. H. F. 273.

Rest-harrow, Cammock, Petty Whin.

In the Marshes at Clapgate—common. July. P.

ANONIS non spinosa. R. S. 332.

ONONIS *arvensis*. H. F. 273.

Corn Rest-harrow, or Cammock.

In Corn Fields at Preston and Ospringe—common.
July. P.

ANTIRRHINUM majus. R. S. omittitur.

ANTIRRHINUM *majus*. H. F. 239.

The greater Snap-Dragon.

On old Walls about the Town—common. June to
September. P.

APARINE. R. S. 225.

GALIUM *Aparine*. H. F. 57.

Cleavers, or Goose-grass.

Under Hedges—very common. August. A.

APARINE palustris minor Parisiensis. R. S. 225.

GALIUM *uliginosum*. H. F. 56.

Marsh Goose-grass.

At the Bottom of Jud's Wood in Ospringe — un-
common. August. P.

APIUM

APIUM palustre et officinarum. R. S. 214.

APIUM graveolens. H. F. 111.

Smallage.

About the Sluice Bridge and the Keys—very common.
August. B.

AQUILEGIA flore simplici. R. S. 273.

AQUILEGIA vulgaris. H. F. 207.

Common Columbines.

In Badgen Wood, and beyond Whitehill, Ospringe
—not common. June. P.

ARTEMISIA vulgaris. R. S. 190.

ARTEMISIA vulgaris. H. F. 312.

Mugwort.

Under Hedges and on the Sides of Fields—very common.
August. P.

ARUM. R. S. 266.

ARUM maculatum. H. F. 342.

Wake-Robin, Cuckow-Pint, Lords and Ladies.

In shady Places and under Hedges—very common.
May. P.

ASPERULA. R. S. 224.

ASPERULA odorata. H. F. 55.

Woodroof.

In Ospringe Woods—plentifully. May. P.

ASPLE-

ASPLENIUM. R. S. 118.

ASPLENIUM *Ceterach*. H. F. 385.

Spleen-wort, Milt-waft.

*On Lenham Church Wall — lately lost on the Walls
of our Abbey. P.*

ASTER *maritimus cæruleus*. R. S. 175.

ASTER *Tripolium*. H. F. 319.

Sea Star-wort.

In the Salt Marshes—plentifully. August. P.

ASTER *arvensis cæruleus acris*. R. S. 175.

ERIGERON *acre*. H. F. 314.

Blue flowered Flea-bane.

On old Walls and on dry Banks—common. August. P.

ASTER *maritimus flavus*, *Crithmum chrysanthemum dictus*. R. S. 174.

INULA *chrythmoides*. H. F. 320.

Golden Sampire.

In the Marshes of Shepey—uncommon. August. P.

ASTRAGALUS *luteus*. R. S. 326.

ASTRAGALUS *glycyphyllos*. H. F. 281.

Wild Liquorice, or Liquorice Vetch.

*Under Hedges in Peasedown, near Whitehill, Ospringe
—uncommon. July. P.*

ATRIPLEX

ATRIPLEX angustifolia maritima dentata.
R. S. 152.

ATRIPLEX ferrata. H. F. 377.

Narrow leaved Orache.

Near Sheerness in Shepey—*uncommon*. August. A.

ATRIPLEX angustissimo et longissimo folio.
R. S. 153.

ATRIPLEX litoralis. H. F. 378.

Grass-leaved Orache.

Near Ramsgate Pier in Thanet — *uncommon*.
August. A.

ATRIPLEX maritima. R. S. 152.

ATRIPLEX laciniata. H. F. 376.

Jagged Sea Orache.

Near Sheerness in Shepey—*uncommon*. August. A.

ATRIPLEX fylvestris angustifolia. R. S. 151.

ATRIPLEX patula. H. F. 377.

Narrow leaved wild Orache.

In Gardens and on Dunchills—*common*. August. A.

ATRIPLEX fylvestris folio hastato seu deltoide.
R. S. 151.

ATRIPLEX hastata. H. F. 377.

Wild Orache with a spear pointed Leaf.

In Gardens and waste Places — *common*. September. A.

ATRIPLEX

ATRIPLEX *maritima scopariæ folio.* R. S. 153.

ATRIPLEX *litoralis.* H. F. 378. β

Grass-leaved Orache.

Near Margate Pier in Thanet — uncommon.

August. A.

ATRIPLEX *maritima fruticosa.* R. S. 153.

ATRIPLEX *portulacoides.* H. F. 376.

Sea Purslane.

Near Harty Ferry—plentifully. August. P.

ATRIPLEX *maritima perennis folio deltoide seu triangulari.* R. S. 152.

ATRIPLEX *bastata.* H. F. 377. β

Wild Sea Orache.

In the Brent Marshes—common. August. A.

B.

BACCHARIS, *monspeliensium.* R. S. 179.

CONYZA *squarrosa.* H. F. 314.

Plowman's Spikenard.

On Beacon Hill and dry Banks—not uncommon.

August. B.

BALLOTE. R. S. 244.

BALLOTA *nigra.* H. F. 227.

Stinking Horehound.

Under Hedges—very common. August. A.

BELLA.

BELLADONNA. R. S. 265.

ATROPA *Belladonna*. H. F. 79.

Deadly Night-shade, or Dwale.

*Amongst the Bushes under Preston-House Wall —
very uncommon. June and July. P.*

BELLIS *sylvestris minor*. R. S. 184.

BELLIS *perennis*. H. F. 320.

Common Daisy.

*In Meadows — very common. From March to Oc-
tober. P.*

BERBERIS *dumetorum*. R. S. 465.

BERBERIS *vulgaris*. H. F.

The Berberry-bush, or Pipperidge-bush.

*In a Hedge beyond Queen Court, Ospringe — un-
common. May. S.*

BETONICA. R. S. 238.

BETONICA *Officinalis*. H. F. 220.

Wood Betony.

In Woods and under Hedges — common. July. P.

BETA *sylvestris maritima*. R. S. 38.

BETA *vulgaris*. H. F. 93.

Sea Beet.

*On the Sea Walls at Ham and Graveney — uncommon.
August. B.*

BETULA.

BETULA. R. S. 443.

BETULA *alba*. H. F. 354.

The Birch Tree.

In Woods—common. April. S.

BIFOLIUM majus, feu Ophrys major quibusdam. R. S. 385.

OPHRYS *ovata*. H. F. 338.

Common Twayblade.

In Cockfet and Jud's Woods—common. June. P.

BLITUM Atriplex sylvestris dictum. R. S. 154.

CHENOPODIUM album. H. F. 91.

Common wild Orache.

In uncultivated Places, and on Rubbish—common.
August. A.

BLITUM foetidum Vulvaria dictum. R. S. 156.

CHENOPODIUM Vulvaria. H. F. 92.

Stinking Orache.

Under Walls about the Town—uncommon. August. A.

BLITUM Ficus folio. R. S. 155.

CHENOPODIUM *serotinum*. H. F. 91.

Late flowering or Fig-leaved Blite.

In Gardens and on Rubbish—common. A.

BLITUM Kali minus album dictum. R. S. 156.

CHENOPODIUM *maritimum*. H. F. 92.

Sea Blite, or white Glass-wort.

In the Salt Marshes—common. August. A.

BLITUM perenne Bonus Henricus dictum.

R. S. 156.

CHENOPODIUM *Bonus Henricus*. H. F. 89.

Common English Mercury, or All-good.

By Way-sides in Ospringe—not uncommon. July. P.

BLITUM Pes anserinus dictum. R. S. 154.

CHENOPODIUM *murale*. H. F. 90.

Common Goose-foot, or Sow-bane.

In Gardens and on Rubbish—not uncommon.

August. A.

BLITUM Pes anserinus dictum acutiore folio.

R. S. 154.

CHENOPODIUM *rubrum*. H. F. 90.

Sharp-leaved Goose-foot, or Sow-bane.

On Dunghills and Rubbish—common. August. A.

BLITUM *rubrum minus*. R. S. 157.

AMARANTUS *Blitum*. H. F. 356.

The least Blite.

On Rubbish and in Gardens—common. August. A.

BORRAGO

BORRAGO hortensis. R. S. 228.

BORRAGO officinalis. H. F. 68.

Bororage.

On old Walls, and by Road-sides near the Town—
not common. June. P.

BORRAGO sempervirens. R. S. 227.

ANCHUSA sempervirens. H. F. 66.

Evergreen Alkanet.

On Rubbish near Preston Church Yard — very un-
common. June. P.

BRASSICA maritima arborea sive procerior ra-
mosa. R. S. 293.

BRASSICA oleracea. H. F. 253.

Sea Cabbage.

On the Cliffs between Deal and Dover — plentifully.
April. B.

BRYONIA alba. R. S. 261.

BRYONIA alba. H. F. 373.

White Briony.

In Hedges — common. May. P.

BUGLOSSA sylvestris minor. R. S. 227.

LYCOPSIS arvensis. H. F. 69.

Small wild Bugloss.

In Corn-fields near Preston Church — uncommon.

May. A.

BUGLOSS.

BUGLOSSUM *arvense* annuum. Lithospermi
folio. R. S. 227.

LITHOSPERMUM *arvense*. H. F. 66.
Bastard Alkanet.

In Corn-fields in Preston and Hernhill — common
July. A.

BUGULA. R. S. 245.

AJUGA *reptans*. H. F. 219.
Bugle.

In Byfing Wood — common. May. P.

BULBOCASTANUM. R. S. 209.

BUNIUM *Bulbocastanum*. H. F. 105.
Earth-nut, or Hog-nut.

In Byfing Wood and Hernhill — common. June. P.

BUPLEURUM *minimum*. R. S. 221.

BUPLEURUM *tenuissimum*. H. F. 97.
The least Hare's-ear.

Near Sheerness in Shepey — uncommon. August. A.

BUPLEURUM *perfoliatum rotundifolium* an-
nuum. R. S. 221.

BUPLEURUM *rotundifolium*. H. F. 97.
Thorow-wax.

In Corn-fields near Brogdale in Ospringe — uncommon.
July. A.

BUTO-

BURSA Pastoris. R. S. 306.

THLASPI *Bursa Pastoris*. H. F. 247.

Shepherd's Purse.

In Hop Grounds, and by Road-sides — very common.

May to September. A.

BUTOMUS. R. S. 273.

BUTOMUS *umbellatus*. H. F. 152.

The flowering Rush, or Water Gladiole.

In Dykes near Pluckley — uncommon. June. P.

* BUXUS. R. S. 445.

BUXUS *sempervirens*. H. F. 355.

The Box Tree.

A few large Trees near Houses in Sheldwich — very uncommon. S.

C.

CAKILE quibusdam, aliis Eruca marina et Rha-
phanus marinus. R. S. 307.

BUNIAS *Cakile*.

Sea Rocket.

*Near Sheerness in Shepey, and Cliff's End in
Thanet — uncommon.* June. A.

CALAMINTHA humilior folio rotundiore.
R. S. 243.

GLECHOMA *hederacea*. H. F. 244.

Ground Ivy. Alehoof.

In shady Places, and under Hedges — very common.
May. P. CALA-

CALAMINTHA vulgaris. R. S. 243.

MELISSA *Calamintha*. H. F. 230.

Common Calamint.

By the Sides of Roads, and under Hedges — common.
August. P.

CAMPANULA rotundifolia. R. S. 277.

CAMPANULA *rotundifolia*. H. F. 80.

The lesser round leaved Bell Flower.

By Way-sides in Ospringe and Boughton — not common.
August. P.

CAMPANULA vulgarior foliis urticæ. R. S. 276.

CAMPANULA *Trachelium*. H. F. 81.

Canterbury Bells, or Great Throat-wort.

In Woods and Hedges — very common. — With a white
Flower in Hernhill — very uncommon. July. P.

CAPRIFOLIUM germanicum. R. S. 458.

LONICERA *Periclymenum*. H. F. 80.

Common Honey-suckle, or Woodbine.

In Woods and Hedges — very common. April to
August. S.

CARDAMINE. R. S. 299.

CARDAMINE *pratensis*. H. F. 256.

Common Ladies Smock, or Cuckow-flower.

In moist Meadows — very common. April. P.

CARDI-

CARDAMINE parviflora. H. F. 257.

Small flowered Ladies Smock.

In Meadows near Pluckley—uncommon. April. A.

CARDIACA. R. S. 239.

LEONURUS Cardiacæ. H. F. 228.

Mother-wort.

Near Cockfet, in Ospringe — very uncommon.

July. B.

CARDUUS acaulos minore purpureo flore.

R. S. 195.

CARDUUS acaulos. H. F. 308.

Dwarf Carline Thistle.

Upon Beacon Hill—not uncommon. July. P.

CARDUUS caule crispo. R. S. 194.

CARDUUS crispus. H. F. 306.

Thistle upon Thistle.

By the Road-sides, and on dry Banks in Ospringe — common. June. A.

CARDUUS lanceolatus. R. S. 195.

CARDUUS lanceolatus. H. F. 305.

Spear-Thistle.

In uncultivated Ground, and under Hedges — common.

July. B.

CAR.

CARDUUS Mariæ. R. S. 195.

CARDUUS *marianus*. H. F. 307.

Milk Thistle, or Lady's Thistle.

*By the Way-side at Davington, and by the Sea Walls
beyond Nagden—not uncommon. July. A.*

CARDUUS Mariæ hirsutus non maculatus.

R. S. 194.

CARDUUS *marianus*. H. F. 307. β.

Ladies Thistle without Spots.

In Graveney Marshes — uncommon. July. A.

CARDUUS nutans. R. S. 193.

CARDUUS *nutans*. H. F. 306.

Musk Thistle.

On Badgen Downs — plentifully. July. B.

CARDUUS palustris. R. S. 194.

CARDUUS *palustris*. H. F. 306.

Marsh Thistle.

In the Marshes — common. July. B.

CARDUUS spinosissimus capitulis minoribus.

R. S. 194.

CARDUUS *acanthoides*. H. F. 306.

Wetted Thistle with small Flowers.

Upon Graveney Marsh Sea Wall—common. July. A.

CAR-

CARDUUS stellatus. R. S. 176.

CENTAUREA *Calcitrapa*. H. F. 326.

Star Thistle.

In Graveney Marshes — common. July. A.

CARDUUS tomentosus *Acanthium dictus vul-*
garis. R. S. 196.

ONOPORDUM *Acanthium*. H. F. 308.

Cotton Thistle.

In uncultivated Places, and by the Road-side — common.
July. B.

CARDUUS vulgarissimus *viarum*. R. S. 194.

SERRATULA *arvensis*. H. F. 305.

Common Way-Thistle, or Creeping Thistle.

In Fields and Byways — very common. July. P.

CARLINA *sylvestris quibusdam, aliis Atractilis*.
R. S. 175.

CARLINA vulgaris. H. F. 309.

The common wild Carline Thistle.

In dry Fields and Pastures — common. June. B.

CARYOPHYLLATA. R. S. 253.

GEUM *urbanum*. H. F. 198.

Herb Bennet, or Common Avens.

In Woods, and under Hedges — common. July. P.

CARY-

ABRIDGED
ABRIDGED
HARVARD
UNIVERSITY

CARYOPHYLLATA montana purpurea. R. S.

253.

GEUM rivale. H. F. 198.

Water Avens.

In a Wood near Barber's Mill in Hothfield — very uncommon. July. P.

CARYOPHYLLUS holosteus arvensis glaber flore majore. R. S. 346.

STELLARIA *Holostea*. H. F. 166.

The greater Stich-wort.

Under Hedges, and on the Sides of Fields — very common. April. P.

CARYOPHYLLUS holosteus arvensis glaber flore minore. R. S. 346.

STELLARIA *graminea*. H. F. 166.

The lesser Stich-wort.

Under Hedges, chiefly on a dry Soil — common. July. P.

CARYOPHYLLUS holosteus arvensis medius. R. S. 347.

STELLARIA *graminea*. H. F. 166. β .

The middle Stich-wort.

Upon Beacon Hill — not uncommon. July. P.

CARYOPHYLLUS latifolius barbatus minor annuus flore minore. R. S. 337.

DIANTHUS *Armeria*. H. F. 161.

Deptford Pink.

On Beacon Hill, and dry Pastures in Ospringe — not uncommon. July. A. CAS-

CASTANEA. R. S. 440.

FAGUS *Castanea*. H. F. 359.

The Chesnut Tree.

In Byling and other Woods—not uncommon. May. S.

CATANANCE leguminosa quorundam. R. S.
325.

LATHYRUS *Nissolia*. H. F. 275.

Crimson Grass-Vetch.

On Beacon Hill — not uncommon. June. A.

CAUCALIS minor flosculis rubentibus. R. S. 219.

CAUCALIS *Anthriscus*. H. F. 99.

Hedge Parsley.

In bushy Grounds, and on Ham Wall—not uncommon.
July. B.

CAUCALIS fegetum minor, Anthriscus hispido
similis. R. S. 220.

CAUCALIS *arvensis*. H. F. 98.

Small Corn Parsley.

In Fields, and amongst Corn—common. August. A.

CENTAURIUM luteum perfoliatum. R. S. 287.

BLACKSTONIA *perfoliata*. H. F. 146.

Yellow Centory.

In dry chalky Fields — common. July. A.

CENTAURIUM minus. R. S. 286.

GENTIANA *Centaurium*. H. F. 88.

Lesser Centory.

In dry Pastures—very common. July to October. A.

*With a white Flower, near Minster in Shepey —
but uncommon.*

CERASUS *sylvestris fructu rubro*. R. S. 463.

PRUNUS *Avium*. H. F. 187.

Common wild Cherry.

In Woods — common. April. S.

CERASTIUM *hirsutum minus parvo flore*.

R. S. 348.

CERASTIUM *semi-decandrium*. H. F. 175.

The least Mouse-ear Chickweed.

On Walls and dry Soils — common. April. A.

CEREFOLIUM *sylvestre*. R. S. 207.

CHÆROPHYLLUM *temulum*. H. F. 108.

Wild Chervil.

Under Hedges — very common. July. A.

CHAMÆMELUM *foetidum*. R. S. 185.

ANTHEMIS *Cotula*. H. F. 223.

Stinking Mayweed.

In Corn Fields — too common. June. A.

CHA-

CHAMÆMELUM inodorum. R. S. 185.

MATRICARIA inodora. H. F. 322.

Field Feverfew.

On the sides of paths in Corn-fields — common.

August. A.

CHAMÆMELUM odoratissimum repens flore
simplici. R. S. 185.

ANTHEMIS nobilis. H. F. 323.

Sweet-scented Camomile.

On Charing Heath — uncommon. August. P.

CHAMÆMELUM vulgare. R. S. 184.

MATRICARIA Chamomilla. H. F. 322.

Corn Feverfew.

In cultivated Fields amongst Corn — common. June. A.

CHELIDONIUM minus. R. S. 246.

FICARIA verna. H. F. 214.

Pilewort, or lesser Celandine.

In Meadows — very common. April. P.

CHENOPODIUM Betæ folio. R. S. 157.

CHENOPODIUM polyspermum. H. F. 92.

Upright round leaved Blite, or Allseed.

In the Conduit Row — not uncommon. August. A.

D

CHENO-

CHENOPODIUM erectum foliis triangularibus
dentatis, spicis e foliorum alis plurimis longis
erectis tenuibus. R. S. 155.

CHENOPODIUM urbicum. H. F. 89.

Upright Blite.

In the Conduit Row, and in Ditches—common. Sep-
tember. A.

CHENOPODIUM Stramonii folio. R. S. 154.

CHENOPODIUM hybridum. H. F. 90.

Maple-leaved Blite.

On Dungbills—not common. August. A.

CHRYSANTHEMUM segetum. R. S. 182.

CHRYSANTHEMUM segetum. H. F. 321.

Corn Marigold.

In Corn-fields — common. July. A.

CICHOREUM sylvestre. R. S. 172.

CICHOREUM Intybus. H. F. 303.

Wild Succory.

*In uncultivated Places, and by Sides of Fields—very
common.* August. P.

CICUTA. R. S. 215.

CONIUM maculatum. H. F. 100.

Hemlock.

In shady Places near the Town—common. July. A.

CICU-

CICUTARIA tenuifolia. R. S. 215.

ÆTHUSA *Cynapium*. H. F. 107.

The lesser Hemlock, or Fools Parsley.

In Gatefield, and in Gardens — not uncommon. September. A.

CICUTARIA vulgaris. R. S. 207.

CHÆROPHYLLUM *sylvestre*. H. F. 108.

Wild Cicely, or Cow-weed.

Under Hedges, and in Fields — common. June. A.

CIRCÆA lutetiana. R. S. 289.

CIRCÆA *lutetiana*. H. F. 9.

Enchanters Night-shade.

In Byfing Wood — common. July. P.

CLEMATIS daphnoides major. R. S. 268.

VINCA *major*. H. F. 77.

The greater Periwinkle.

In Hedges, near Houses at Ore and Sindal Farm — uncommon. April. P.

CLEMATIS latifolia seu Atragene quibusdam.
R. S. 258.

CLEMATIS *Vitalba*. H. F. 217.

Great Wild Climber, or Traveller's Joy.

In Hedges — very common. July. P.

CLINOPODIUM Origano simile. R. S. 239.

CLINOPODIUM *vulgare*. H. F. 228.

Great wild Basil.

In Hedges — very common. July. P.

COCHLEARIA folio sinuato. R. S. 303.

COCHLEARIA *anglica*. H. F. 248.

Sea Scurvy-grass.

On the Sea Walls near Thorn—common. May. B.

CONVOLVULUS major. R. S. 275.

CONVOLVULUS *sepium*. H. F. 74.

Great Bindweed.

Under Hedges—very common. With a reddish purple Flower, in the Brent Hedge; but very uncommon. August. P.

CONVOLVULUS maritimus Soldanella dictus.

R. S. 276.

CONVOLVULUS *Soldanella*. H. F. 75.

Sea Bindweed.

On the Sand Downs near Sandwich — not common.

July. P.

CONVOLVULUS minor vulgaris. R. S. 275.

CONVOLVULUS *arvensis*. H. F. 74.

Small Bindweed.

In Fields, and by Road-sides—very common. July. P.

CONIZA

CONYZA media. R. S. 174.

INULA dysenterica. H. F. 320.

Middle Fleabane.

About the Abbey Pond — common. August. P.

CONYZA minor. R. S. 174.

INULA Pulicaria. H. F. 320.

Small Fleabane.

About the Conduit-row, and at Downe's-forstal in
Hernhill — not uncommon. September. A.

CORNUS fæmina. R. S. 460.

CORNUS sanguinea. H. F. 53.

The female Cornel, Dog-berry Tree, or
Prick-wood.

In Hedges — common. June. S.

CORYLUS fylvestris. R. S. 439.

CORYLUS Avellana. H. F. 360.

The Hazle-nut Tree.

In Woods and Hedges — very common. March. S.

COTYLEDON vera radice tuberosa. R. S. 271.

COTYLEDON Umbilicus Veneris. H. F. 169.

Navel-wort, Kidney-wort, or Wall Penny-wort.

On Tenterden Church, and in a Stone Pit at
Boughton Monchelsea — very uncommon. June. P.

CRACCA. R. S. 322.

VICIA *Cracca*. H. F. 277.

Tufted Vetch.

In the Hedges of the Beacon Farm — uncommon.

July. A.

CRACCA minor. R. S. 322.

ERVUM *hirsutum*. H. F. 280.

Small wild Tare, Tine-Tare, or hairy Tare.

In Rickefy-lane Hedge, Selling — not uncommon.

July. A.

CRACCA minor filiquis singularibus flosculis
cærulefcentibus. R. S. 322.

ERVUM *tetraspermum*. H. F.

Tine-Tare with smooth Pods, or smooth Tare.

In Fields and under Hedges — common. June. A.

CRITHMUM *marinum*. R. S. 217.

CRITHMUM *maritimum*. H. F. 101.

Sampire.

*On the Cliffs between St. Margaret's and Dover —
plentifully. August. P.*

CRUCIATA. R. S. 223.

VALANTIA *Cruciata*. H. F. 375.

Cross-wort, or Mug-weed.

In Hedges in Preston and Ospringe — not uncommon.

June. F.

CUSCUTA

CUSCUTA major. R. S. 281.

CUSCUTA *europæa*. H. F. 89.

Dodder, Hell-weed, Devil's-guts.

Upon leguminous Plants — common. July. A.

CYANUS. R. S. 198.

CENTAUREA *Cyanus*. H. F. 325.

Blue Bottles, or Corn Bottles.

In Corn-fields — very common. July. A.

CYNOGLOSSUM. R. S. 226.

CYNOGLOSSUM *officinale*. H. F. 67. β.

Great Hound's Tongue.

By the Road Sides near Boughton Chalk Pits — not common. June. P.

CYNOGLOSSUM *folio virenti*. R. S. 226.

CYNOGLOSSUM *officinale*. H. F. 67. β.

The lesser green leaved Hound's Tongue.

Near Sandwich — not common. June. P.

D.

DAMASONIUM *stellatum* Dalechampii.

R. S. 272.

ALISMA *Damasonium*. H. F. 138.

Star-headed Water Plantain.

In a Field-pond of Mr. Jacob's Farm, at East Church in Shepey — very uncommon. June to August. P.

DAU.

DAUCUS vulgaris. R. S. 218.

DAUCUS *Carota*. H. F. 99.

Wild Carrot, or Bird's Nest.

In Pastures, and by Way-sides—very common. July. B.

* DELPHINIUM *segetum* flore caruleo.

R. S. 273.

DELPHINIUM *Consolida*. H. F. 207.

Wild Larkspur.

In King's Field, amongst Corn — very uncommon.

June. A.

DENS-LEONIS. R. S. 170.

LEONTODON *Taraxacum*. H. F. 297.

Dandelion.

In Meadows and Pastures—very common. May. P.

DENS-LEONIS *hirsutus*, Hieracium dictus.

R. S. 127.

LEONTODON *hispidum*. H. F. 297.

Rough Dandelion, or Dandelion Hawk-weed.

In Meadows and Pastures — common. June. P.

DIGITALIS *purpurea*. R. S. * 283.

DIGITALIS *purpurea*. H. F. 240.

Purple Fox Glove.

In Woods and Hedges — common. July. B.

DIPSACUS

DIPSACUS minor seu Virga pastoris. R. S. 192.

DIPSACUS pilosus. H. F. 49.

Small wild Teasel, or Shepherd's Rod.

In the Road Hedge leading from Plomford to Badgen

Downs — uncommon. July or August. B.

DIPSACUS sylvestris seu Labrum Veneris.

R. S. 192.

DIPSACUS sylvestris. H. F. 49.

Wild Teasel.

In uncultivated Places, and under Hedges — very

common. July. B.

E.

ECHIUM alterum sive Lycopsis anglica. R. S. 228.

ECHIUM anglicum. H. F. 70.

English Viper's Bugloss.

By Waysides near Whitehill in Ospringe — not un-

common. August. P.

ECHIUM vulgare. R. S. 218.

ECHIUM vulgare. H. F. 69.

Viper's Bugloss.

On barren Ground, and upon old Walls — very common.

July. P.

EQUISETUM arvense longioribus setis. R. S.

130.

EQUISETUM arvense. H. F. 380.

Corn Horse-tail.

In moist Fields amongst Corn — common. March. P.

EQUISETUM majus. R. S. 130.

EQUISETUM fluviatile. H. F. 381.

River Horse-tail.

In the River about the Powder Mills — common.

May. P.

EQUISETUM palustre. R. S. 131.

EQUISETUM palustre. H. F. 380.

The lesser Marsh Horse-tail.

In the Corn Fields near Nagden — not uncommon.

June. P.

EQUISETUM sylvaticum. R. S. 130.

EQUISETUM sylvaticum. H. F. 380.

Wood Horse-tail.

In the swampy Ground near Ore Mill — not common.

May. P.

ERICA Brabantica folio Coridis hirsuto quaterno.

R. S. 471.

ERICA tetralix. H. F. 144.

Cross-leaved Heath.

At the Bottom of Jud's Wood — not common. Au-

gust. S.

ERICA tenuifolia. R. S. 471.

ERICA cinerea. H. F. 144.

Fine leaved Heath.

Upon Charing Heath — not common. August. S.

ERICA

ERICA vulgaris. R. S. 470.

ERICA vulgaris. H. F. 144.

Common Heath, or Ling.

In Woods—very common. June to September. P.

ERUCA aquatica. R. S. 297.

SISYMBRIUM *sylvestre*. H. F. 258.

Water Rocket.

In Ham Ponds near Sandwich—uncommon. June. P.

ERUCA hirsuta filiqua cauli appressa. Erysimum
vulgare. R. S. 298.

ERYSIMUM *officinale*. H. F. 250.

Hedge Mustard.

Under Hedges, and by Road Sides — very common.
May. A.

ERUCA lutea feu Barbarea. R. S. 297.

ERYSIMUM *Barbarea*. H. F. 251.

Winter Cress or Rocket.

Near the Powder Mills—not uncommon. May. P.

ERYNGIUM marinum. R. S. 222.

ERYNGIUM *maritimum*. H. F. 95.

Eryngo, or Sea Holly.

On the Shore between Graveney and Sea Salter —
uncommon. July. P.

ERYSIMUM

ERYSIMUM *Sophia dictum.* R. S. 298.

SYSIMBRIUM *Sophia.* H. F. 259.

Flix-weed.

By the Road Sides near Sittingbourn — uncommon.

June. A.

EUONYMUS *vulgaris.* R. S. 468.

EUONYMUS *europæus.* H. F. 84.

The Spindle Tree, or Prick-wood.

In Woods and Hedges — common. May. S.

EUPATORIUM *Cannabinum.* R. S. 179.

EUPATORIUM *cannabinum.* H. F. 310.

Common Hemp Agrimony, or Dutch Agrimony.

About the Abbey Pond — common. August. P.

EUPHRASIA. R. S. * 284.

EUPHRASIA *officinalis.* H. F. 234.

Eye-bright.

In most upland Pastures — very common. August. A.

EUPHRASIA *pratensis rubra.* R. S. * 284.

EUPHRASIA *Odontites.* H. F. 234.

Red Eye-bright.

In the Ways and Corn Fields about Preston — very common. August. A.

FAGUS.

F.

FAGUS. R. S. 439.

FAGUS *sylvaticus*. H. F. 360.

The Beech Tree.

In Woods — common. May. S.

* FEGOPYRUM. R. S. 144.

POLYGONUM *Fagopyrum*. H. F. 150.

Buck-wheat, or Brank.

In Corn fields near Beacon Hill — uncommon. July. A.

FEGOPYRUM *scandens sylvestre*. R. S. 144.

POLYGONUM *Convolutus*. H. F. 149.

Black Bind-weed.

In Corn fields and Hedges — not uncommon. June to September. A.

FERRUM *equinum germanicum* filiquis in summitate. R. S. 326.

HIPPOCREPIS *comosa*. H. F. 281.

Tufted Horse-shoe Vetch.

On chalky Ground in Ospringe — common. July. A.

FILIPENDULA. R. S. 259.

SPIRÆA *Filipendula*. H. F. 190.

Common Dropwort.

Upon Beacon Hill — not uncommon. July. P.

E

FILIX

FILIX Fœmina. R. S. 124.

PTERIS *aquilina*. H. F. 384.

Female Fern, or Common Brakes.

In Woods — very common. August. P.

FILIX mas vulgaris. R. S. 120.

POLYPODIUM *Filix mas*. H. F. 389.

Common Male Fern.

On shady Benks under Hedges — common. P.

FILICULA saxatilis ramosa maritima nostras.

R. S. 125.

PTERIS *aquilina*. H. F. 384. β.

Small-branched Stone-Fern.

On Walls about the Town — uncommon. August. P.

FCENICULUM vulgare. R. S. 217.

ANETHUM *Fœniculum*. H. F. 110.

Common Fennel, or Finkle.

By the Road-side as you enter the Town — not uncommon. August. P.

FRAGARIA. R. S. 254.

FRAGARIA *vesca*. H. F. 194.

Strawberry.

In Woods — very common. April. P.

FRAGRA-

FRAGARIA sterilis. R. S. 254.

FRAGARIA sterilis. H. F. 195.

Barren Strawberry.

On barren Lands — very common. April. P.

FRANGULA five Alnus nigra baccifera. R.

465.

RHAMNUS Frangula. H. F. 83.

The black Berry-bearing Alder.

In Badgen Wood — not common. May. S.

FRAXINUS. R. S. 469.

FRAXINUS excelsior. H. F. 379.

The common Ash Tree.

In Woods and Hedges — common. April. S.

FUMARIA vulgaris. R. S. 204.

FUMARIA officinalis. H. F. 270.

Fumitory.

In Fields and Gardens — common. May. A.

FUMARIA alba latifolia. R. S. 204.

FUMARIA claviculata. H. F. 270.

Climbing Fumitory.

In moist Hedges in Davington and Ospringe — not common. August. A.

G.

GALE frutex odoratus septentrionalium. R.S. 443.

MYRICA *Gale*. H. F. 368.

Sweet Willow, Goule, Dutch Myrtle.

On Willsborough Lees, near Ashford — plentifully.

May. S.

GALEOPSIS legitima Dioscoridis. R. S. 237.

STACHYS *sylvatica*. H. F. 227.

Hedge Nettle.

In Woods and Hedges—very common. August. A.

GALLIUM luteum. R. S. 224.

GALLIUM *verum*. H. F. 55.

Yellow Ladies Bedstraw, or Cheese-renning.

On the Sides of Fields—very common. July. P.

GENISTA angulosa trifolia. R. S. 474.

SPARTIUM *scoparium*. H. F. 271.

Common Broom.

In Woods and dry Fields — very common. June. P.

GENISTA minor aspalathoides. R. S. 475.

GENISTA *anglica*. H. F. 272.

Needle Furze, or Petty Whin.

Amongst Furze which grows on wet Land — not common. May. P.

GENISTA

GENISTA spinosa vulgaris. R. S. 475.

ULEX europæus. H. F. 272.

Furze, Whins, or Gorfe.

On Beacon Hill—*very common*. March to June. P.

GENISTELLA tinctoria. R. S. 474.

GENISTA tinctoria. H. F. 272.

Dyers-weed, Green-wood.

In the Marshes near Holly Shore — *not common*.
July. P.

GENTIANELLA fugax autumnalis. R.S. 275.

GENTIANA Amarella. H. F. 87.

Autumnal Gentian.

In dry Pastures and Ospringe Chalk Pits — *not uncommon*. August. A.

GERANIUM Cicutæ folio inodorum. R. S. 357.

GERANIUM cicutarium. H. F. 262.

Field Crane's-bill without Scent, or Hemlock-leaved Crane's-bill.

By Road sides, and on old Walls — *not uncommon*.
April to June. A.

GERANIUM columbinum. R. S. 359.

GERANIUM molle. H. F. 265.

Common Dove's-foot Crane's-bill.

Under Hedges — *very common*. June. A.

GERANIUM columbinum humile flore cæruleo.

R. S. 359.

GERANIUM *pusillum*. H. F. 266.

Small flowered Dove's-foot Crane's-bill.

By Waysides leading to Thorn — common. June. A.

GERANIUM columbinum majus foliis dissectis.

R. S. 359.

GERANIUM *dissectum*. H. F. 266.

Jagged leaved Dove's-foot Crane's-bill.

In dry Fields in Ospringe — not uncommon. May to July. A.

GERANIUM columbinum dissectis foliis pediculis
florum longissimis. R. S. 359.

GERANIUM *columbinum*. H. F. 266.

Long-stalked Dove's-foot Crane's-bill.

Upon Oldwives Lees — not common. July. A.

GERANIUM robertianum. R. S. 358.

GERANIUM *robertianum*. H. F. 264.

Herb Robert.

In Hedges, and on Rubbish — very common. June. B.

GERANIUM *pusillum* *supinum* *maritimum*.

R. S. 357.

GERANIUM *maritimum*. H. F. 263.

Sea Crane's-bill.

On the Sand Downs near Deal — not common.

August. P.

GLAUX

GLAUX *maritima*. R. S. 285.

GLAUX *maritima*, H. F. 86.

Sea Milkwort, or black Saltwort.

In the Brent Marshes — common. July. P.

GNAPHALIUM *longifolium humile ramosum*
capitulis nigris. R. S. 181.

GNAPHALIUM *uliginosum.* H. F. 313.

Black-headed, long leaved, low branched cud-
weed.

In moist Places in Hernhill — not uncommon.
August. A.

GNAPHALIUM *anglicum.* R. S. 180.

GNAPHALIUM *sylvaticum.* H. F. 313.

Upright Cudweed.

In Judd's Wood — common. August. B.

GNAPHALIUM *minimum.* R. S. 181.

FILAGO *montana.* H. F. 328.

The least Cudweed.

In dry sandy Meadows, and among Corn — common.
July. A.

GNAPHALIUM *minus seu Herba impia.*
R. S. 180.

FILAGO *germanica.* H. F. 328.

Common Cudweed.

In dry Meadows and by Road sides — common. July. A.

GROS-

* GROSSULARIA. Miller Botan Officinal. 220.

RIBES *Uva crispa*. Lin. System. Nat. Tom. II.
p. 184.

The Gooseberry Bush.

In several Hedges beyond White Hill, Ospringe.
April. S.

H.

HERA communis major et minor. R. S. 459.

HERA *Helix*. H. F. 85.

Climbing, or berried Ivy, also barren and creeping
Ivy.

In Woods and Hedges — very common. October. S.

HELIANTHEMUM vulgare. R. S. 341.

CISTUS *Helianthemum*. H. F. 205.

Dwarf Cistus, or little Sunflower.

On dry chalky Banks — very common. June. P.

HELENIUM. R. S. 176.

INULA *Helenium*. H. F. 319.

Elicampane.

On the moist Ground near the Half-way House to Can-
terbury — very uncommon. August. P.

HELLEBORUS niger hortensis. R. S. 271.

HELLEBORUS *viridis*. H. F. 215.

Wild black Hellebore.

On the Cliff beyond Westfield in Pluckley — un-
common. April. P. HEL-

HELLEBORASTER maximus. R. S. 271.

HELLEBORUS *fætidus*. H. F. 215.

Great bastard black Hellebore, Bear's-foot, or
Setter-wort.

By the Road side, up the Chalk-hill, about a Mile
North-west from Charing—*uncommon*. March. P.

HELLEBORINE *altera atro-rubente flore.*
R. S. 383.

SERAPIAS *latifolia*. H. F. 341. β .

Broad-leaved bastard Hellebore, a Variety.

In Ospringe Woods—*uncommon*. August. P.

HELLEBORINE *latifolia montana*. R. S. 383.

SERAPIAS *latifolia*. H. F. 341.

Broad leaved bastard Hellebore.

In King's Wood — *uncommon*. July. P.

HELLEBORINE *latifolia flore albo clauso.*

R. S. 384.

SERAPIAS *longifolia*. H. F. 341.

White flowered bastard Hellebore.

In Badgen and Jud's Woods—*uncommon*. June. P.

HERBA Paris. R. S. 264.

PARIS *quadrifolia*. H. F. 150.

Herb Paris, True-love, or One-berry.

In Woods near Doddington—*not common*. May. P.

HES.

HESPERIS allium redolens. R. S. 293.

ERYSIMUM *Alliaria*. H. F. 251.

Jack by the Hedge, or Sauce alone.

Under Hedges — very common. May. P.

HIERACIUM *Echioides capitulis Cardui benedicti*. R. S. 166.

PICRIS *Echioides*. H. F. 294.

Ox's Tongue, or Lang de Boeuf.

In Hedges and Pastures going to Thorn Creek — common. July. A.

HIERACIUM fruticosum angustifolium majus. R. S. 168.

HIERACIUM *umbellatum*. H. F. 300.

Narrow leaved bushy Hawk-weed.

In Ospringe Chalk Pits, and by Way-sides at Dunkirk — uncommon. August. P.

HIERACIUM fruticosum latifolium hirsutum. R. S. 187.

HIERACIUM *sabaudum*. H. F. 300.

Bushy Hawk-weed with broad rough Leaves.

In Hedges about Sheldwich — not common. July. P.

HIERACIUM longius radiculatum. R. S. 165.

HYPOCHÆERIS *radicata*. H. F. 302.

Long-rooted Hawk-weed.

On Beacon Hill — common. June. P.

HIER-

HIERACIUM luteum glabrum five minus hirsutum. R. S. 165.

CREPIS testorum. H. F. 301.

Smooth Succory Hawk-weed.

In Fields and on old Walls — common. June to September. A.

HIERACIUM maximum Chondrillæ folio asperum. R. S. 166.

CREPIS biennis. H. F. 301.

Rough Succory Hawk-weed.

In Hedges near Sittingbourn — uncommon. August. B.

HIERACIUM minus præmorsa radice. R. S. 164.

LEONTODON autumnale. H. F. 297.

Hawk-weed with bitten Roots, or yellow Devil's Bit.

In Corn fields in Hernhill — common. August. P.

HORMINUM sylvestre Lavendulæ flore. R. S. 237.

SALVIA verbenaca. H. F. 9.

Common English wild Clary, or Oculus Christi.

In Davington and Ore Church-Yards — not common. May. P.

HOTTONIA. R. S. 285.

HOTTONIA palustris. H. F. 72.

Water Violet, or Gilliflower.

About the Decoy Ponds — uncommon. In Dykes near Deal — common. June. P. HYA

HYACINTHUS *anglicus*. R. S. 373.

HYACINTHUS *non scriptus*. H. F. 123.

English Hyacinth, or Hare-bells.

In Woods and under Hedges—very common. April. P.

HYDROCOTYLE *vulgaris*. R. S. 222.

HYDROCOTYLE *vulgaris*. H. F. 96.

Marsh Penny-wort, or White-rot.

In the Marshes about Luddenham and Ore — not uncommon. May. P.

HYOSCYAMUS *vulgaris*. R. S. 274.

HYOSCYAMUS *niger*. H. F. 77.

Common Henbane.

By Road sides near the Town—common. June. P.

HYPERICUM. R. S. 342.

HYPERICUM *perforatum*. H. F. 290.

Saint John's-wort.

Under Hedges, and by Road sides — very common. July. P.

HYPERICUM *Androsæmum dictum*. R. S. 343.

HYPERICUM *hirsutum*. H. F. 291.

Tutsan, or hairy St. John's-wort.

In hollow Lanes in Boughton—uncommon. July. P.

HYPE-

HYPERICUM Ascyron dictum caule quadrangulo. R. S. 344.

HYPERICUM *quadrangulum*. H. F. 292.
Saint Peter's-wort.

In Davington Brooks—not uncommon. July. P.

HYPERICUM pulchrum Tragi. R. S. 342.

HYPERICUM *pulchrum*. H. F. 290.

Upright Saint John's-wort.

In Byfing Wood — common. July. P.

HYPERICUM maximum Androsæmum vulgare dictum. R. S. 343.

HYPERICUM *Androsæmum*. H. F. 291.

Tutfan, or Park-leaves.

In a Hedge near Provender Wood — uncommon.
July. P.

HYPERICUM minus supinum. R. S. 343.

HYPERICUM *humifusum*. H. F. 290.

The least trailing Saint John's-wort.

On sandy Lands in Ore—common. July. P.

HYPERICUM *eleganissimum non ramosum folio lato*. R. S. 343.

HYPERICUM *montenum*. H. F. 291.

Imperforate Saint John's-wort.

In Byfing Wood — uncommon. July. P.

J.

JACEA major. R. S. 198.

CENTAUREA *Scabiosa*. H. F. 325.

Great Knap-weed, or Matfellow.

On the sides of Paths — common. July. P.

JACEA nigra. R. S. 198.

CENTAUREA *Jacea*. H. F. 326.

Common Knap-weed, or Matfellow.

On the sides of Fields — very common. August. P.

JACOBÆA latifolia palustris sive aquatica.

R. S. 178.

SENECIO *aquaticus*. H. F. 317.

Broad-leaved Marsh or Water Ragwort.

In the Brents — common. August. P.

JACOBÆA senecionis folio incano perennis.

R. S. 177.

SENECIO *erucifolius*. H. F. 317.

Hoary Perennial Ragwort with Groundsel Leaves.

In Byfing Wood — not common. July. P.

JACOBÆA vulgaris. R. S. 177.

SENECIO *Jacobæa*. H. F. 316.

Common Ragwort.

In Meadows and Pastures — very common. July. P.

IRIS palustris lutea. R. S. 374.

IRIS *Pseudacorus*. H. F. 13.

Yellow Water Flower de luce.

In moist boggy Places — very common. June. P.

IRIS sylvestris quam Xyrim vocant. R. S. 375.

IRIS *fœtidissima*. H. F. 13.

Stinking Gladdon.

By Sandgate Castle near Folkestone — very uncommon.

July. P.

JUNIPERUS vulgaris baccis parvis purpureis.

R. S. 444.

JUNIPERUS *communis*. H. F. 372.

The common Juniper.

On dry chalky Commons — very common. May. S.

K.

KALI geniculatum perenne fruticosius procumbens. R. S. 136.

SALICORNIA *europæa*. H. F. 1. β.

Perennial Marsh Sampire, or jointed Glaſs-wort.

In the Isle of Shepey — uncommon. August. P.

KALI spinosum cochleatum. R. S. 159.

SALSOLA *Kali*. H. F. 33.

Prickly Glaſs-wort.

On the Shore between Graveney and Seafalter — not common. July. A.

KNAWEL. R. S. 159.

SCLERANTHUS *annuus*. H. F. 160.

German Knot-grass, or annual Knavel.

On barren dry sandy Soils—very common. August. A.

L.

LACTUCA *agnina* feu *Valerianella foliis ferratis*.

R. S. 201.

VALERIANA *Locusta*. H. F. 12.

Small Corn Sallet, or Valerian with jagged leaves.

In Fields among Corn, and on old Walls—common.

May. A.

LACTUCA *fylvestris murorum*. R. S. 162.

PRENANTHES *muralis*. H. F. 296.

Ivy-leaved wild Lettuce.

In Hedges near Wilderton—uncommon. July. P.

LACTUCA *fylvestris minima*. R. S. 162.

LACTUCA *saligna*. H. F. 296.

The least wild Lettuce.

On Banks of Ditches near Sheerness—uncommon.

August. A.

LAMIUM *album*. R. S. 240.

LAMIUM *album*. H. F. 225.

White Archangel or Dead Nettle.

Under Hedges—very common. May. P.

LAMIUM

LAMIUM rubrum. R. S. 240.

LAMIUM rubrum. H. F. 225.

Red Archangel or Dead Nettle.

On Rubbish, and sides of Fields — very common.

May. A.

LAMIUM cannabinum floribus albis verticillis
purpurascentibus. R. S. 241.

GALEOPSIS *Tetrait*. β . H. F. 226.

White flowered Hemp-leaved Dead Nettle.

*In Hedges between Plomford and Badgen Downs —
not common. August. A.*

LAMIUM cannabinum folio vulgare. R. S. 240.

GALEOPSIS *Tetrait*. H. F. 226.

Nettle Hemp, or Hemp-leaved Dead Nettle.

In Byfing Wood — not common. August. A.

LAMIUM cannabinum folio, flore amplo luteo, labio
purpureo. R. S. 241.

GALEOPSIS *Tetrait* δ . H. F. 226.

Hemp-leaved Dead Nettle with a parti-coloured
Flower.

*In Mr. Jacob's new Wood-hedge at Nackington —
very uncommon. August. A.*

LAMIUM folio caulem ambiente majus et minus.
R. S. 240.

LAMIUM amplexicaule. H. F. 225.

Great Henbit.

In Hop-grounds — very common. June. A.

LAMIUM luteum. R. S. 242.

GALEOPSIS *Galeobdolon*. H. F. 226.

Yellow Archangel, Dead Nettle, or yellow Nettle
Hemp.

In Woods and under shady Hedges — very common.

May. P.

LAMPSANA. R. S. 173.

LAPSANA *communis*. H. F. 303.

Nipple-wort.

In moist Cornfields, and by Way-sides — very common.

July. A.

LAPATHUM *acetosum repens lanceolatum*.

R. S. 143.

RUMEX *Acetofella*. H. F. 136.

Sheep's Sorrel.

On dry sandy or gravelly Ground — very common.

July. P.

LAPATHUM *acetosum vulgare*. R. S. 143.

RUMEX *Acetosa*. H. F. 136.

Common Sorrel.

In Meadows and Pastures — very common. June. B.

LAPATHUM *acutum*. R. S. 142.

RUMEX *acutus*. H. F. 134.

Sharp-pointed Dock.

In uncultivated Places — very common. June. P.

LAPA-

LAPATHUM acutum minimum. R. S. 141.

An RUMICIS *acuti* varietas. H. F. 134.

The lesser sharp-pointed Dock.

In Woods — not uncommon. June. P.

LAPATHUM folio acuto crispo. R. S. 141.

RUMEX *crispus*. H. F. 134.

Sharp-pointed Dock, with curled Leaves.

On the Mud Walls beyond the Sluice — common.

July. P.

LAPATHUM folio acuto rubente. R. S. 142.

RUMEX *sanguineus*. H. F. 133.

Blood-wort.

In Davington Brooks — uncommon. July. B.

LAPATHUM maximum aquaticum sive Hydro-lapathum. R. S. 140.

RUMEX. *Britannica*. H. F. 135.

Great Water Dock.

In Ditches and boggy Places — very common. August. P.

LAPATHUM pulchrum Bononiense sinuatum.

R. S. 142.

RUMEX *puleher*. H. F. 134.

Fiddle-Dock.

By the sides of Foot-ways about the Town — not uncommon. June. P.

LAPA-

LAPATHUM viride. R. S. 141.

An RUMICIS *acuti* varietas. H. F. 134.

The greener leaved Dock.

In shady Woods and moist Ground — not uncommon.

June. P.

LAPATHUM vulgare folio obtuso. R. S. 141.

RUMEX *obtusifolius*. H. F. 134.

Common broad-leaved Dock.

In moist Places, and by Road sides — very common.

July. P.

LAPPA major Arctium Dioscoridis. R. S. 197.

ARCTIUM *Lappa*. H. F. 304.

Burdock, or Clot burr.

Under Hedges, and by Road-sides — very common.

August. B.

LAPPA major capitulo glabro maximo. R. S. 196.

ARCTIUM *Lappa*. β . H. F. 304.

Greater Burdock, with a larger Flower.

By Road-sides, and under Hedges — common.

August. B.

LAPPA major capitulis parvis glabris. R. S. 197.

ARCTIUM *Lappa*. H. F. 304.

Burdock with small Heads.

By Road sides in Hern-hill — not uncommon. Au-

gust. B.

LAPPA

LAPPA major montana capitulis tomentosis.

ARCTIUM *Lappa*. H. F. 304.

Greater Burdock with woolly Heads.

In the Way-sides near Luddenham Court—uncommon.

August. B.

LATHYRUS luteus sylvestris dumetorum. R. S.

320.

LATHYRUS *pratensis*. H. F. 277.

Tare everlasting, or common yellow bastard Vetchling.

In Woods and Hedges — common. August. A.

LATHYRUS major latifolius. R. S. 319.

LATHYRUS *latifolius*. H. F. 276.

Broad-leaved everlasting Pea.

In a Field Hedge at Copton, and by the Road-side near Boughton Street—uncommon. July. P.

LATHYRI majoris species flore rubente et albido.
R. S. 319.

LATHYRUS *sylvestris*. H. F. 276.

Narrow-leaved everlasting Pea.

In a Hedge of the Beacon Farm—uncommon. August. P.

LAUREOLA. R. S. 465.

DAPHNE *Laureola*.

Dwarf Laurel, or Spurge-Laurel.

In Woods and under Hedges — common. March. P.

LENS

LENS palustris. R. S. 129.

LEMNA minor. H. F. 345.

Duck's Meat.

In Ponds and standing Waters — very common.

June. P.

LENTICULA aquatica trifulca. R. S. 129.

LEMNA trifulca. H. F. 344.

Ivy-leaved Duck's Meat.

In the Ditches of Ham Marsh—common. June. P.

LENTICULA palustris major. R. S. 129.

LEMNA polyrrhiza. H. F. 345.

Greater Duck's Meat.

In a Ditch near Waterham—uncommon. July. A.

LEPIDIDIUM latifolium. R. S. 304.

LEPIDIDIUM latifolium. H. F. 244.

Dittander, or Pepperwort.

Near the King's Head Key in the Town—very uncommon. July. P.

LEUCANTHEMUM vulgare. R. S. 184.

CHRYSANTHEMUM *Leucanthemum*. H. F.

321.

The greater Daisie, or Ox-eye.

In Pastures, and amongst Corn—common. May. P.

LEUCO-

LEUCOJUM luteum vulgo Cheiri flore simplici.

R. S. 291.

CHEIRANTHUS *Cheiri*. H. F. 250.

Wall-flower, or wild Chier.

On old Walls — very uncommon. May. P.

* LEUCOJUM bulbosum trifolium minus.

Bauhini Pinacis. 56.

GALANTHUS *nivalis*. Lin. Syst. Natur.

II. p. 234.

The Snow Drop.

In Pastures at South-street, and at Davington —

very common. February. P.

LIGUSTRUM. R. S. 465.

LIGUSTRUM *vulgare*. H. F. 3.

Privet, or Prim.

In Woods and Hedges — very common. May. S.

LILLIUM convallium. R. S. 264.

CONVALLARIA *majalis*. H. F. 126.

Lilly Convally, or May Lilly.

In the Blean Woods — plentifully. May. P.

LIMNOPEUCE. R. S. 136.

HIPPURIS *vulgaris*. H. F. 1.

Mare's Tail.

In the Powder-Mill Waters — uncommon. May. P.

LIMO-

LIMONIUM. R. S. 200.

STATICE *Limonium*. H. F. 114.

Sea Lavender.

In the Salt Marshes—very common. August. P.

LINARIA *Elatine dicta folio acuminato.*

R. S. * 282.

ANTIRRHINUM *Elatine*. H. F. 237.

Sharp-pointed Fluellin.

In Corn Fields — common. September. A.

LINARIA *Elatine dicta folio subrotundo.*

R. S. * 282.

ANTIRRHINUM *spurium*. H. F. 237.

Round-leaved Fluellin.

In the Corn fields of Preston—not uncommon. August. A.

LINARIA *lutea vulgaris*. R. S. * 281.

ANTIRRHINUM *Linaria*. H. F. 238.

Common yellow Toad-flax.

On the sides of Fields and Roads — very common. July. P.

LINUM *sylvestre angustifolium floribus dilute purpurascentibus vel carneis.* R. S. 362.

LINUM *tenuifolium*. H. F. 116.

Narrow-leaved Wild Flax.

Upon Beacon Hill — not common. June. P.

LINUM

LINUM sylvestre cæruleum perenne erectius flore
et capitulo majore. R. S. 362.

LINUM *perenne*. H. F. 115.

Perennial Blue Flax.

In the Isle of Shepey — uncommon. June. P.

LINUM sylvestre catharticum. R. S. 362.

LINUM *catharticum*. H. F. 116.

Purging Flax, Wild Dwarf Flax, Mill-mountain
On sandy and chalky Soils — very common. June. A.

LITHOSPERMUM, seu Miliun Solis. R. S. 228.

LITHOSPERMUM *officinale*. H. F. 65.

Gromwell, Gromil, or Graymill.

In Badgen Wood, and by Road sides — not uncommon.
June. P.

LONCHITIS aspera. R. S. 118.

OSMUNDA *Spicant*. H. F. 382.

Rough Spleen-wort.

In the Woods near Dunkirk — uncommon. July. P.

LOTI corniculatæ major species. R. S. 334.

LOTUS *corniculata*. H. F. 288. γ.

The greater Bird's-foot Trefoil.

Under Hedges near Byling Wood — not common.
July. P.

LOTUS corniculata glabra minor. R. S. 334.

LOTUS *corniculata*. H. F. 288.

Bird's-foot Trefoil.

Under Hedges, and in Fields in Ospringe——common.

July. P.

LUPULUS mas et foemina. R. S. 137.

HUMULUS *Lupulus*. H. F. 369.

Hops, the male and female.

In Hedges in Preston — not uncommon. July. P.

LUTEOLA. R. S. 366.

RESEDA *Luteola*. H. F. 181.

Wild Woad, Yellow Weed, or Dyers Weed.

In Davington Church-Yard, and on barren Lands——

not uncommon — June. A.

LYCHNIS major noctiflora Dubrensis perennis.

R. S. 340.

CUCUBALUS *viscosus*. H. F. 163.

Dover Campion.

On the Cliffs between St. Margaret's and Dover——

plentifully. July. P.

LYCHNIS maritima repens. R. S. 337.

SILENE *amœna*. ? H. F. 164.

Sea Campion.

In the Isle of Shepey. — uncommon. August. P.

LYCH-

LYCHNIS plumaria sylvestris simplex. R.S. 338.

LYCHNIS Flos cuculi. H. F. 174.

Meadow Pink, Wild Williams, Cuckow Flower.

In moist Pastures — very common. June. P.

LYCHNIS Saponaria dicta. R. S. 339.

SAPONARIA officinalis. H. F. 160.

Common Soap-wort.

In Hedges near Sindal and Keatings, and beyond
Sheldwich Church — uncommon. July. P.

LYCHNIS fegetum major. R. S. 338.

AGROSTEMMA Githaco. H. F. 173.

Cockle.

In Corn Fields — common. June. A.

LYCHNIS supina maritima Ericæ facie. R.S. 338.

FRANKENIA levis. H. F. 119.

Smooth Sea-Heath.

Near Minster in Shepey — uncommon. August. P.

LYCHNIS sylvestris angustifolia caliculis turgidis
striatis. R. S. 341.

SILENE concidea. H. F. 165.

Narrow-leaved Campion.

On the Sandhills near Deal — uncommon. July.

LYCHNIS sylvestris flore albo. R. S. 339.

LYCHNIS dioica. H. F. 174.

Wild white Campion.

In Woods and Hedges — not uncommon. June. P.

LYCHNIS sylvestris flore rubello. R. S. 339.

LYCHNIS dioica. H. F. 174. β.

Red flowered wild Campion.

In Woods and Hedges — common. June. P.

LYCHNIS sylvestris, quæ Behen album vulgo.

R. S. 337.

CUCUBALUS Behen. H. F. 163.

Spatling Poppy, Bladder Campion, or white
Corn Campion.

In dry Pastures, and in Corn fields—common. July. P.

LYCOPUS palustris glaber. R. S. 236.

LYCOPUS europæus. H. F. 8.

Water Horehound.

*On the Banks of the River, and in boggy Places —
common. July. P.*

LYSIMACHIA campestris. R. S. 311.

EPILOBIUM montanum. H. F. 141.

The greater smooth-leaved Willow-herb, or
Loose-strife.

*By the sides of Ditches, and under moist Hedges —
common. July. B.*

LYSI-

LYSIMACHIA lutea. R. S. 282.

LYSIMACHIA vulgaris. H. F. 72.

Yellow Willow-herb, or Loose-strife.

On the swampy Ground near Stone Bridge — very uncommon. August. P.

LYSIMACHIA filiquosa glabra media five minor.
R. S. 311.

EPILOBIUM tetragonum. H. F. 141.

Middle smooth-leaved Willow-herb, or Loose-strife.

In moist Ditches, and near the River — common.
July. P.

LYSIMACHIA filiquosa glabra minor angustifolia. R. S. 311.

EPILOBIUM palustre. H. F. 141.

The least smooth coddled Willow-herb, or Loose-strife.

In moist Places — common. July. P.

LYSIMACHIA filiquosa hirsuta magno flore.
R. S. 311.

EPILOBIUM ramosum. H. F. 141.

Great flowered Willow-herb, called Codlings and Cream.

On the sides of the shooting Meadow Stream — common.
July. P.

LYSIMACHIA filiquosa hirsuta parvo flore.
R. S. 311.

EPILOBIUM hirsutum. H. F. 140.

The lesser hairy coddled Willow-herb, or Loose-strife.

On the sides of the shooting Meadow Stream — common.
July. P.

LYSI-

LYSIMACHIA speciosa, quibusdam Onagra
dicta. R. S. 310.

EPILOBIUM angustifolium. H. F. 140.
Rosebay Willow-herb.

In a Wood Hedge beyond the five Mile Stone to Charing
— *very uncommon.* July. P.

M.

MALUS sylvestris. R. S. 452.

PYRUS Malus. H. F. 189.

The Crab Tree.

In Hedges — not uncommon. May. S.

MALVA sylvestris minor. R. S. 251.

MALVA rotundifolia. H. F. 268.

Small wild or Dwarf Mallow.

On dry Banks by Road sides — common. July to
October. A.

MALVA vulgaris. R. S. 251.

MALVA sylvestris. H. F. 268.

Common Mallow.

Under Hedges, by Way sides, and on Rubbish — very
common. July. B.

MARRUBIUM album. R. S. 239.

MARRUBIUM vulgare. H. F. 228.

White Horehound.

On Sheldwich Lees — not common. July. P.

MATRI-

MATRICARIA. R. S. 187.

MATRICARIA *Parthenium*. H. F. 321.

Feverfew.

Under Hedges, and on old Walls—common. June. P.

MELAMPYRUM *sylvaticum* flore luteo. R. S.
* 286.

MELAMPYRUM *sylvaticum*. H. F. 236.

Common Cow-wheat with a yellow Flower.

In Byfing Wood—very common. June to August. A.

MELILOTUS *vulgaris*. R. S. 331.

TRIFOLIUM *Melilotus officinalis*. H. F. 282.

Common Melilot.

Amongst the Corn at Halfto—plentifully. July. B.

* MELISSA *officinalis*. Miller Botan. Off. 290.

MELISSA *officinalis*. Lin. System. Nat. II.

p. 400.

Common Garden Balm.

Under Hedges in Boughton and Davington — un-
common. July. P.

MENIANTHES *palustre triphyllum latifolium*
et angustifolium. R. S. 285.

MENYANTHES *trifolia*. H. F. 71.

Marsh Trefoil, or Buckbean.

In the moist Meadows of the Abbey—common. July. P.

MENTHA

MENTHA aquatica feu Syfimbrium. R. S. 233.

MENTHA *aquatica*. H. F. 223.

Water-mint.

In moist Places, and by the River side — very common.

August. P.

MENTHA feu Calamintha aquatica. R. S. 232.

MENTHA *arvensis*. H. F. 223.

Water-mint with whirled coronets.

On the sides of Ditches, and in Corn fields — common.

August. P.

MENTHA spicata angustifolia glabra spica latiore.

R. S. 234.

MENTHA *spicata*. H. F. 221.

Spear-mint.

On the Side of the River opposite the Powder Stove —

very uncommon. August. P.

MENTHASTRUM folio rugoso rotundiore spontaneum flore spicato odore gravi. R. S. 234.

MENTHA *rotundifolia*. H. F. 221.

Round leaved Horse-mint.

In a Meadow behind King's Mill — uncommon. Au-

gust. P.

MENTHASTRUM spicatum folio longiore candicante. R. S. 234.

MENTHA *longifolia*. H. F. 221.

Long leaved Horse-mint.

On the Banks of the River above Ospringe Church —

not uncommon. August. P.

MER-

MERCURIALIS annua glabra vulgaris, Mercurialis mas et Foemina. R. S. 139.

MERCURIALIS annua. H. F. 371.

French Mercury.

On Rubbish and in Gardens — not common. September. A.

MERCURIALIS perennis repens. Cynocrambe dicta. R. S. 138.

MERCURIALIS perennis. H. F. 371.

Dog's Mercury.

In Woods and under Hedges — very common. April. P.

MESPILUS Apii folio sylvestris spinosa, five Oxyacantha. R. S. 453.

CRATÆGUS Oxyacantha. H. F. 188.

The white Thorn, or Hawthorn, called Quick.

In Woods and Hedges — very common. May. S.

MESPILUS Apii folio sylvestris non spinosa. R. S. 453.

CRATÆGUS torminalis. H. F. 188.

The common or wild Service-tree, or Sorb.

In the Woods about Chart — uncommon. April or May. S.

MILLEFOLIUM vulgare. R. S. 183.

ACHILLEA Millefolium. H. F. 324.

Common Yarrow, Millfoil, or Nose-bleed.

In Meadows and Pastures — very common. June. P.

MOL-

MOLLUGO montana minor Gallio albo similis.
R. S. 224.

GALIUM montanum. H. F. 56.

Small mountain bastard Madder, or mountain
Ladies Bedstraw.

In the Corn fields about Sindal—not uncommon. July. P.

MOLLUGO vulgarior. R. S. 223.

GALIUM Mollugo. H. F. 56.

Wild Madder, or great bastard Madder.

In Hedges and bushy Places—very common. July. P.

MOLLUGINIS vulgarioris varietas minor.
R. S. 224.

GALIUM palustre. H. F. 57.

White Ladies Bedstraw.

On the sides of the shooting Meadow Stream—common.
July. P.

MOSCHATELLINA foliis fumarizæ bulbosæ.
R. S. 267.

ADOXA Moschatellina. H. F. 150.

Tuberous Moschatell.

In Woods and Hedges—very common. April. P.

MYOSOTIS scorpioides hirsuta. R. S. 229.

MYOSOTIS scorpioides. H. F. 65.

Moufe-ear Scorpion grass.

In Woods, on old Walls and dry Pastures—common.
April to August. P. MYOSO-

MYOSOTIS scorpioides palustris. R. S. 229.

MYOSOTIS *scorpioides*. H. F. 65. δ .

Water Scorpion-grass.

On the sides of Ore Mill-pond—not common. April
to August. P.

MYRRHIS sylvestris feminibus asperis. R. S. 220.

SCANDIX *Anthriscus*. H. F. 108.

Small Hemlock - Chervil with rough Seeds.

Under Hedges, and by the sides of Ditches — common.
May. A.

N.

NAPUS sylvestris. R. S. 295.

BRASSICA *Napus*. H. F. 253.

Wild Navew.

Amongst Corn, and by the sides of Ditches — common.
May. B.

NARCISSUS sylvestris pallidus calyce luteo.
R. S. 371.

NARCISSUS *Pseudo Narcissus*. H. F. 123.

Wild English Daffodil.

In the Orchards near Selling-Street—very uncommon.
April. P.

NASTURTIUM supinum capsulis verrucosis, five
Coronopus Ruellii. R. S. 304.

COCHLEARIA *Coronopus*. H. F. 248.

Swine's Cress.

On the sides of Roads and Foot-paths — very common.

July. A.

NEPETA

NEPETA major vulgaris. R. S. 237.

NEPETA *Cataria*. H. F. 220.

Nep, or Cat-mint.

Under the West Hedge of King's-Field — not common.

July. P.

NUMMULARIA. R. S. 283.

LYSIMACHIA *Nummularia*. H. F. 73.

Money-wort, or Herb Two-pence.

In moist Hedges at Luddenham, and near Provender

— not common. June. P.

NUMMULARIA minor flore purpurascens.

R. S. 283.

LYSIMACHIA *tenella*. H. F. 73.

Purple flowered Money-wort.

In moist Meadows by the Abbey — not uncommon.

July. P.

NYMPHÆA alba. R. S. 368.

NYMPHÆA *alba*. H. F. 206.

White Water Lilly.

In Ham Ponds near Sandwich — uncommon. July. P.

NYMPHÆA lutea. H. F. 368.

NYMPHÆA *lutea*. H. F. 206.

Yellow Water Lilly.

In Ore Mill Pond — uncommon. July. P.

OENAN-

O.

OENANTHE *aquatica*. R. S. 210.

OENANTHE *fistulosa*. H. F. 104.

Water Dropwort.

In the Ditches near Fairbrook—common. July. P.

OENANTHE *Cicutæ facie*. Lobellii. R.S. 210.

OENANTHE *crocata*. H. F. 105.

Hemlock Dropwort.

*In the Water-lane between Sitingbourn and Milton—
very uncommon. June. P.*

ONOBRYCHIS feu Caput gallinaceum. R.S. 327.

HEDYSARUM *Onobrychis*. H. F. 281.

Saintfoin or Cock's-head.

On the chalky Hills — common. June. P.

OPHIOGLOSSUM. R. S. 128.

OPHIOGLOSSUM *vulgatum*. H. F. 382.

Adders Tongue.

*In Meadows at Ospringe and Luddenham — not un-
common. May. P.*

OPULUS *Ruellii*. R. S. 460.

VIBURNUM *Opulus*. H. F. 112.

Water Elder, or Cherrywood.

*In Badgen Wood, and in Hedges — not uncommon.
June. S.*

H

CICHIS

ORCHIS abortiva rufa sive Nidus avis. R. S. 382.

OPHRYS *Nidus avis*. H. F. 338.

Birds-nest Orchis.

In Byfing and Jud's Woods — very uncommon.

June. P.

ORCHIS alba bifolia minor calcari oblongo.

R. S. 380.

ORCHIS *bifolia*. H. F. 333.

Butterfly Orchis.

In Badgen and Cockset Woods — not common.

May. P.

ORCHIS anthropophora oreades. R. S. 397.

OPHRYS *anthropophora*. H. F. 340.

The Green Man Orchis.

On chalky Banks — common. June. P.

ORCHIS fuciflora galea et alis purpurascens.

R. S. 379.

OPHRYS *apifera*. H. F. 340.

The Bee Orchis.

In Meadows in Ospringe — not uncommon. June. P.

ORCHIS galea et alis fere cinereis. R. S. 378.

ORCHIS *militaris*. H. F. 335.

The Man Orchis.

On chalky bushy Banks beyond Whitehill, Ospringe — not common. June. P.

ORCHIS

C. Hall sc.

Orchis & Morio femina flore pleno

ORCHIS magna latifolia galea fusca vel nigricante.

R. S. 378.

ORCHIS *purpurea*. H. F. 334.

The purple Man Orchis.

In Cockset and Jud's Woods—uncommon. May. P.

ORCHIS morio foemina. R. S. 377.

ORCHIS *morio*. H. F. 333.

Female Orchis, or Female Fools Stones.

In Meadows — very common. In a Meadow near Cades in Ospringe, some of a Rose Colour, some white, and a very few with double Flowers.
May and June. P.

ORCHIS morio mas foliis maculatis. R. S. 376.

ORCHIS *mascula*. H. F. 333.

Male Orchis, or Male Fools Stones.

In Hedges and Meadows—very common. May. P.

ORCHIS myodes galea et alis herbidis. R.S. 379.

OPHRYS *muscifera*. H. F. 340.

The Fly Orchis.

Among the Bushes in Ospringe Parsonage Meadows—
not uncommon. June. P.

ORCHIS odorata moschata. R. S. 378.

OPHRYS *Monorchis*. H. F. 339.

Yellow or Musk Orchis.

On the chalky Banks of Ospringe Parsonage Mea-
dows — uncommon. June. P.

ORCHIS palmata minor flore luteo-viridi.

R. S. 381.

SATYRIUM viride. H. F. 337.

Frog Satyrion or Orchis.

On dry grassy Banks near Whitehill ——— uncommon.

June. P.

ORCHIS palmata pratensis latifolia longis calcaribus. R. S. 380.

ORCHIS latifolia. H. F. 335.

Male handed Orchis, or Satyrion royal.

On moist boggy Grounds about Ore Mill ——— not uncommon. June. P.

ORCHIS palmata speciosiore thyrsifolia maculato. R. S. 381.

ORCHIS maculata. H. F. 335.

Female handed Orchis, or Satyrion royal.

In Chalk Pits — common. June. P.

ORCHIS pannonica. R. S. 377.

ORCHIS ustulata. H. F. 334.

Little purple flowered Orchis.

On the South side of Chatham Hill ——— plentifully.
May. P.

ORCHIS purpurea spica congesta pyramidalis. R. S. 377.

ORCHIS pyramidalis. H. F. 334.

Purple late flowering Orchis.

On chalky Banks in Ospringe — common. July. P.

ORCHIS *spiralis alba odorata*, five *Triorchis*.

R. S. 378.

OPHRYS *spiralis*.

Triple Ladies Traces.

In Ospringe Parsonage Meadows ——— *uncommon*.

September. P.

ORCHIS five *Testiculus sphegodes hirsuto flore*.

R. S. 380.

OPHRIS *apifera*. H. F. 340. β .

Humble Bee Orchis with green Wings.

On the Banks of Whitehill in Selling ——— *uncommon*.

April. P.

ORIGANUM *Onites*. R. S. 236.

ORIGANUM *Onites*. H. F. 229.

Pot Marjoram.

In Waldershare Park ——— *very uncommon*. August. P.

ORIGANUM *vulgare spontaneum*. R. S. 236.

ORIGANUM *vulgare*. H. F. 229.

Wild Marjoram.

Under Hedges, and in bushy Places ——— *very common*.

July. P.

ORNITHOPODIUM *radice nodosa*. R. S. 326.

ORNITHOPUS *perpusillus*. H. F. 280.

Birds Foot.

In Broom fields at Hernhill ——— *common*. May to Au-

gust. P.

OROBANCHE major Garyophyllum olens.

R. S. * 288.

OROBANCHE major. H. F. 232.

Broom-rape.

In Broom fields — not uncommon. June. B.

OROBUS sylvaticus foliis oblongis glabris.

R. S. 324.

OROBUS tuberosus. H. F. 274.

Wood Pea, or Heath Pea.

In Jud's Wood — very common. May. P.

OSTRYA Ulmo similis, fructu in umbilicis foliaceis.

R. S. 451.

CARPINUS. Betulus. H. F. 360.

The Horn-beam Tree, or Horse-Beech.

In Woods — common. May. S.

OXYS alba. R. S. * 281.

OXALIS Acetofella. H. F. 173.

Wood Sorrel.

In Byfing and Jud's Woods — common. June. A.

P.

PAPAVER corniculatum luteum. R. S. 309.

CHELIDONIUM Glaucium. H. F. 202.

Yellow horned Poppy.

On the Beach at Sea-Salter — common. July. A.

PAPAVER

PAPAYER corniculatum luteum. Chelidonia
dictum. R. S. 309.

CHELIDONIUM *majus*. H. F. 201.

The greater Celandine.

Among Rubbish, and on old Walls—common. June. P.

PAPAYER laciniato folio capitulo brevior glabro
annuum, Rhœas dictum. R. S. 308.

PAPAYER *Rhœas*. H. F. 203.

Red Poppy, or Corn Poppy.

In Corn fields — very common. July. A.

PARIETARIA. R. S. 158.

PARIETARIA *officinalis*. H. F. 376.

Pellitory of the Wall.

On old Walls — very common. July. P.

PARONYCHIA vulgaris. R. S. 292.

DRABA *verna*. H. F. 243.

Common Whitlow Grass.

On old Walls and dry Pastures — very common.

April. A.

PASTINACA sylvestris latifolia. R. S. 206.

PASTINACA *fativa*. H. F. 109.

Wild Parsnep.

*On the chalky Bank by the Roadside near Wilderton—
uncommon.* July. B.

PEDI-

PEDICULARIS feu Crista galli lutea. R. S.
* 284.

RHINANTHUS *Crista galli*. H. F. 234.

Yellow Rattle, or Cocks-comb.

In Byfing Wood, and by Road fides—common. July. A.

PEDICULARIS palustris rubra elatior. R. S.
* 284.

PEDICULARIS *palustris*. H. F. 236.

Great Marsh red Rattle, Cocks-comb, Loufe-wort.

About the Powder Mills at Ore—not common. June. A.

PEDICULARIS pratensis rubra vulgaris. R. S.
* 284.

PEDICULARIS *sylvatica*. H. F. 236.

Red Rattle, Cocks-comb, Loufe-wort.

In Jud's Wood — common. June. P.

PENTAPHYLLOIDES *Argentina dicta*. R. S.
256.

POTENTILLA *Argentina*. H. F. 195.

Wild Tansey, or Silver-weed.

By the fides of Roads and Paths—common. July. P.

PENTAPHYLLUM erectum foliis profunde
— fectis subtus argenteis flore luteo. R. S. 255.

POTENTILLA *argentea*. H. F. 196.

Tormentil Cinquefoil.

In a Road hedge near Newnham.—uncommon. May. P

PENTA-

PENTAPHYLLUM vulgatissimum. R. S. 225.

POTENTILLA reptans. H. F. 197.

Common Cinquefoil, or Five-fingered Grass.

By the sides of Roads and Paths—common. June. P.

PENTAPTEROPHYLLON aquaticum, flosculis
ad foliorum nodos. R. S. 316.

MYRIOPHYLLUM verticillatum. H. F. 357.

Verticillated Water-millfoil.

In the Ditches of Graveney Marsh — not uncommon.
July. P.

PERCEPIER Anglorum. R. S. 159.

APHANES arvensis. H. F. 60.

Pursley-piert.

Upon barren Lands and old Walls—common. May. A.

PERSICARIA latifolia geniculata caulibus macu-
latis. R. S. 145.

POLYGONUM Persicaria. H. F. 147. β.

Dead or spotted Arsmart.

In Fields, and on Rubbish — common. August. A.

PERSICARIA maculosa. R. S. 145.

POLYGONUM Persicaria. H. F. 147,

Dead or spotted Arsmart.

In Meadows and Ditches—very common. Sept. A.

PERSI-

PERSICARIA mitis major foliis pallidioribus.
R. S. 145.

POLYGONUM *pensylvanicum*. H. F. 148.

Greater dead Arsmart with pale Leaves.

On a Dungbill at Staple Street — uncommon.
August. A.

PERSICARIA falicis folio perennis, Potamogiton
angustifolium dicta. R. S. 145.

POLYGONUM *amphibium*. H. F. 147.

Perennial Willow-leaved Arsmart.

On the sides of the Abby Ditches — not common.
July. P.

PERSICARIA vulgaris acris, feu Hydropiper.
R. S. 144.

POLYGONUM *Hydropiper*. H. F. 148.

Lakeweed, Arsmart, Water-pepper.

In moist Places and wet Ditches — very common.
August. A.

PETASITES. R. S. 179.

TUSSILAGO *Petasites*. H. F. 315.

Common Butter Bur, Pestilent-wort.

Near Waterham in Hernhill — uncommon. March. P.

* PETROSELINUM vulgare. Miller. Botan.
Officinal. 324.

APIUM *Petroselinum*. Linnæi System. Natur.
II. p. 217.

Common Parsley.

On the West Church-yard Wall — very uncommon.
June. B. PEU-

PEUCEDANUM. R. S. 206.

PEUCEDANUM *officinale*. H. F. 101.

Hog-fennel, or Sulphur-wort.

On the Wall leading to Thorn Creek — plentifully.

June. P.

PELLANDRIUM vel Cicutaria aquatica quorundam. R. S. 215.

PELLANDRIUM *aquaticum*. H. F. 107.

Water Hemlock.

On boggy Ground between Staple-Street and Hernhill Church — uncommon. July. B.

PHYLLITIS feu Lingua cervina vulgo. R. S. 116.

ASPLENIUM *Scolopendrium*. H. F. 384.

Hart's Tongue.

In shady Lanes, hollow Ways, and on Walls — common. F.

PILOSELLA repens. R. S. 170.

HIERACIUM *Pilosella*. H. F. 298.

Common creeping Mouse-ear.

On dry sandy Banks — common. May. A.

PIMPINELLA Saxifraga. R. S. 213.

PIMPINELLA *major*. H. F. 110.

Great Burnet-saxifrage.

In a thick Hedge beyond Whitehill — uncommon.

August. P.

PIMP:-

PIMPINELLA *Saxifraga minor*. R. S. 213.

PIMPINELLA *Saxifraga*. H. F. 111. β .

Small Burnet-saxifrage with divided Leaves.

In Meadows in Hernhill — not uncommon. August. P.

PIMPINELLA *Saxifraga minor foliis sanguiforbæ*. R. S. 213.

PIMPINELLA *Saxifraga*. H. F. 111.

The lesser round leaved Burnet-saxifrage.

In dry Pastures — common. August and September. P.

PLANTAGO *aquatica*. R. S. 257.

ALISMA *Plantago*. H. F. 137.

Great Water Plantain.

In moist Places, and on the sides of Ponds — common. August. P.

PLANTAGO *foliis laciniatis, Coronopus dicta*.

PLANTAGO *Coronopus*. H. F. 52.

Buck's-horn Plantain, Star of the Earth.

In the Abby Meadows — very common. August. A.

PLANTAGO *latifolia vulgaris*. R. S. 314.

PLANTAGO *major*. H. F. 51.

Great Plantain, Way-bread.

By Way sides — very common. July. P. The *Plantago panicula sparsa*, found in *Perry field*, is an elegant variety of this Plant, but very uncommon. PLAN-

PLANTAGO major incana. R. S. 314.

PLANTAGO media. H. F. 51.

Hoary Plantain.

By Road sides — common. July. P.

PLANTAGO marina. R. S. 315.

PLANTAGO Loeftingii. H. F. 52.

Sea Plantain.

On the sides of the Ditches and Sea Walls — common.
July. P.

PLANTAGO maritima. H. F. 52.

Narrow leaved Plantain.

In the Brent Marshes — not uncommon. July. P.

PLANTAGO quinquenervia. R. S. 314.

PLANTAGO lanceolata. H. F. 52.

Rib-wort, or Rib wort Plantain.

In Fields and Meadows — very common. June to
August. P.

POLYGALA. R. S. * 287.

POLYGALA vulgaris. H. F. 271.

Milk-wort.

In dry Fields, and under Hedges — very common.
June. P.

POLYGONUM *brevi angustoque folio*. R. S. 146.

POLYGONUM *aviculare*. H. F. 149. β .

Knot-grass with a small narrow Leaf.

By Road sides—common. June to September. A.

POLYGONUM *mas vulgare*. R. S. 146.

POLYGONUM *aviculare*. H. F. 149.

Common Knot-grass.

In uncultivated Ground, and by Way sides — very common. June to October. A.

POLYGONUM *oblongo angustoque folio*.
R. S. 146.

POLYGONUM *aviculare*. H. F. 149. γ .

Knot-grass with a more oblong narrow Leaf.

In Corn fields at Hernhill — common. June to September. A.

POLYPODIUM. R. S. 117.

POLYPODIUM *vulgare*. H. F. 387.

Common Polypody.

In Byfing Wood, and on Banks of hollow Ways — common. P.

POPULAGO; five *Caltha palustris*. R. S. 272.

CALTHA *palustris*. H. F. 214.

Marsh Marigold.

In the Brent Marshes—very common. April. P.

POPUL-

POPULUS alba. R. S. 446.

POPULUS alba. H. F. 370.

The White Poplar, or Abele Tree.

In moist Woods, and boggy Places — common.

March. S.

POPULUS Lybica Plinii. R. S. 446.

POPULUS tremula. H. F. 370.

The Asp, or Trembling Poplar.

In moist Places in Hernhill — not uncommon.

March. S.

POPULUS major. R. S. 446.

POPULUS nigra. H. F. 370.

The Black Poplar.

In moist boggy Places in Hernhill — not uncommon.

March. S.

PORTULA. R. S. 368.

PEPLIS Portulaca. H. F. 128.

Water Purslane.

In the Brent Meadows — not uncommon. September. P.

POTAMOGETON foliis pennatis. R. S. 150.

MYRIOPHYLLUM spicatum. H. F. 357.

Feathered or spiked Water Millfoil.

In the Powder Mill Waters, and Graveney Marsh Dykes — common. July. P.

POTAMOGITON seu Fontalis crispa. Tribulus
aquaticus. R. S. 149.

POTAMOGETON *crispum*. H. F. 61.

The greater Water Caltrops.

In the Powder Mill Waters—common. June. P.

POTAMOGETON seu Fontalis media lucens.
R. S. 149.

POTAMOGETON *ferratum*. H. F. 61.

The lesser Water Caltrops, or Frog's Lettuce.

In the River near the Powder Stove—not uncommon.
June. P.

POTAMOGITON gramineum latiusculum foliis
et ramificationibus dense stipatis. R. S. 149.

POTAMOGETON *gramineum*. H. F. 62.

Grass leaved Pond weed.

*In the shallow Waters about the Powder Mills — not
common.* P.

POTAMOGITON maritimum ramosissimum gran-
diusculis capitulis, capillaceo folio nostras.
R. S. 150.

POTAMOGETON *marinum*. H. F. 63.

Sea Pond-weed.

In Dykes near Sheerness — uncommon. August. P.

POTAMOGITON millefolium, seu foliis gramineis
ramosum. R. S. 150.

POTAMOGETON *peñtinatum*. H. F. 62.

Fine or Fennel-leaved Pond-weed.

In Ospringe River — not common. June. P.

POTAMOGITON perfoliatum. R. S. 149.

POTAMOGETON *perfoliatum*. H. F. 60.

Perfoliated Pond-weed.

In Ospringe River — *common*. July. P.

POTAMOGITON rotundifolium. R. S. 148.

POTAMOGETON *natans*. H. F. 60.

Broad leaved Pond-weed.

In a Pond near Uplees — *uncommon*. August. P.

PRIMULA pratensis inodora lutea. R. S. 284.

PRIMULA *veris*. H. F. 70. β .

Great Cowslips, or Oxslips.

In thick Hedges beyond Whitehill — *not common*.

April. P.

PRIMULA *veris* major. R. S. 284.

PRIMULA *veris*. H. F. 71. α .

Common Pagils, or Cowslips.

In Meadows and Hedges — *very common*. April. P.

PRIMULA *veris* vulgaris. R. S. 284.

PRIMULA *vulgaris*. H. F. 70.

Common Primrose.

In Woods and Hedges — *very common*. February to

May. P.

PRUNELLA. R. S. 238.

PRUNELLA *vulgaris*. H. F. 231.

Common Self-heal.

In Meadows — common. August. P.

PRUNUS *sylvestris*. R. S. 48.

PRUNUS *spinosa*. H. F. 186.

The Black Thorn, or Sloe Tree.

In Hedges — very common. April. S.

PRUNUS *sylvestris major*. R. S. 462.

PRUNUS *infinitia*. H. F. 186.

The black Bullace Tree.

In some Hedges near Rickefey-Lane in Selling —
uncommon. April. S.

PRUNUS *sylvestris fructu majore albo*. R.S. 462.

PRUNUS *infinitia*. H. F. 186. γ.

The white Bullace Tree.

In Hedges at Cades in Ospringe, and at Selling —
uncommon. April. S.

PRUNUS *sylvestris fructu rubro acerbo et ingrato*.

R. S. 463.

PRUNUS *infinitia*. H. F. 186. β.

The Scad Tree.

In Hedges — not uncommon. April. S.

PTARMICA. R. S. 183.

ACHILLEA *Ptarmica*. H. F. 325.

Sneeze-wort, Bastard Pellitory, or Goose-tongue.

In Davington Brooks — *uncommon*. August. P.

PULEGIUM. R. S. 235.

MENTHA *Pulegium*. H. F. 224.

Penny-royal, or Pudding-grafs.

In a small Pond at Elvyland in Ospringe — *uncommon*.

PYRASTER seu Pyrus sylvestris. R. S. 452.

PYRUS *communis*. H. F. 189.

The Wild Pear Tree.

In the Wood leading from Keneways to Wilderton —
very uncommon. April. S.

Q.

QUERCUS latifolia. R. S. 440.

QUERCUS *Robur*. H. F. 359.

The common Oak Tree.

In Woods — *very common*. April. P.

QUERCUS latifolia mas quæ brevi pediculo est.

R. S. 440.

QUERCUS *Robur*. H. F. 359. β.

The Oak Tree.

In Woods and Hedges — *common*. April. S.

RADI-

R.

RADICULA sylvestris feu palustris. R. S. 301.

SISYMBRIUM *amphibium*. H. F. 258. α.

Water Radish.

About the Powder-Mill Waters — not common.

June. P.

RADIOLA vulgaris serpyllifolia Millegrana!
minima. R. S. 345.

LINUM *Radiola*. H. F. 117.

The least Rupture-wort, or All-seed.

*On sandy Banks between Ore and Luddenham —
common. A.*

RANUNCULO five Polyanthemo aquatili albo
affine. R. S. 250.

RANUNCULUS *aquatilis*. H. F. 213. γ.

Fennel-leaved Crow-foot.

In Running Waters — common. May. P.

RANUNCULUS aquaticus albus, circinatis
tenuissime divisis foliis, floribus ex alis longis
pediculis innixis. R. S. 249.

RANUNCULUS *aquatilis*. H. F. 213. δ.

Finest leaved Water Crow-foot.

In the Brent Dykes — common. April to June. P.

RANUN-

RANUNCULUS aquatilis. R. S. 249.

RANUNCULUS *aquatilis*. H. F. 213.

Various leaved Water Crow-foot.

In the shooting Meadow Stream — common.

May. P.

RANUNCULUS aquatilis hederaceus albus.

R. S. 249.

RANUNCULUS *hederaceus*. H. F. 212.

Ivy leaved Water Crow-foot.

By the Wall that leads to Thorn — common. May. P.

RANUNCULUS aquatilis omnino tenuifolius.

R. S. 249.

RANUNCULUS *aquatilis*. H. F. 213. β .

Fine leaved Water Crow-foot.

In Cspringe River near Whitehill — common. April
to June. P.

RANUNCULUS arvorum. R. S. 248.

RANUNCULUS *arvensis*. H. F. 212.

Corn Crow-foot.

In Corn fields — very common. June. A.

RANUNCULUS bulbosus. R. S. 247.

RANUNCULUS *bulbosus*. H. F. 211.

Round rooted or bulbous Crow-foot, or Butter-
cups

In Meadows — very common. May. P.

RANUN-

RANUNCULUS flammeus major. R. S. 250.

RANUNCULUS *Lingua*. H. F. 210.

Great Spear-wort.

In the old Haven near Sandwich — very uncommon.

May. P.

RANUNCULUS flammeus minor. R. S. 250.

RANUNCULUS *Flammula*. H. F. 210.

The lesser Spear-wort.

In the shooting Meadow Stream — common. June to
September. P.

RANUNCULUS nemorosus dulcis secundus
Tragi. R. S. 248.

RANUNCULUS *Auricomus*. H. F. 211?

Sweet wood Crow-foot, or Goldilocks.

In Woods and under Hedges — common. April. P.

RANUNCULUS palustris. R. S. 249.

RANUNCULUS *sceleratus*. H. F. 212.

Round leaved Water Crow-foot.

On the Banks of the Brent Marshes — common.
June. A.

RANUNCULUS pratensis erectus acris. R. S.
248.

RANUNCULUS *acris*. H. F. 211.

Upright Meadow Crow foot.

In Meadows — very common. June. P.

RANUNCULUS pratensis repens. R. S. 247.

RANUNCULUS *repens*. H. F. 210.

Common creeping Crow-foot.

On Beacon Hill — common. May. P.

RAPA

RAPA fativa rotunda. R. S. 294.

BRASSICA *Rapa*. H. F. 253.

Turneps.

In Corn fields, and by Waysides — not uncommon.

April. B.

RAPISTRUM arvorum. R. S. 295.

SINAPIS *arvensis*. H. F. 260.

Charlock, or Wild Mustard.

In Fields among Corn—common. May. A.

RAPISTRUM flore luteo, filiqua glabra articulata.

R. S. 296.

RHAPHANUS *Rbaphanistrum*. H. F. 252. β .

Yellow flowered Charlock with a jointed Cod.

In Fields among Corn — common. July. A.

RAPUNCULUS *Scabiotæ capitulo cæruleo*.

R. S. 278.

JASIONE *montana*. H. F. 329.

Hairy Sheep's Scabious.

By the Waysides near Dunkirk — uncommon.

August. A.

RESEDA vulgaris. R. S. 366.

RESEDA *lutea*. H. F. 181.

Base Rocket.

In Boughton Chalk Pits — not common. July. A.

RHAM-

RHAMNOCIDES fructifera, foliis Salicis, baccis
leviter flavescentibus. R. S. 445.

HIPPOPHAE *Rhamnoides*. H. F. 368.

Sea Buck Thorn, or Sallow Thorn.

*In Shepey, uncommon; near Sandown Castle —
plentifully. April. S.*

RHAMNUS catharticus. R. S. 466.

RHAMNUS *catharticus*. H. F. 83.

Buck Thorn, or Purging Thorn.

*In Perry Wood and Badgen Down Hedges — not
common. May. S.*

RHAPHANISTRUM siliqua articulata glabra
majore et minore. R. S. 296.

RHAPHANUS *Rhaphanistrum*. H. F. 252.

White flowered Charlock with a jointed Cod.

In Fields among Corn — common. June. A.

RHAPHANUS ruficanus. R. S. 301.

COCHLEARIA *Armoracia*. H. F. 248.

Horfe Radish.

*On the Banks of the River, above the Sluice, and of
Ore Stream—uncommon. June. P.*

RIBES nigrum vulgo dictum folio olente. R. S. 456.

RIBES *nigrum*. H. F. 85.

Black Currants, or Squinancy Berries.

*In the boggy Oster Ground near Stone Bridge — un-
common. May. S.*

RIBES

* *RIBES vulgare fructu rubro.* R. S. 465.

RIBES rubrum. H. F. 84.

Currants.

In some thick Hedges in Ospringe — not common.

May. S.

ROSA sylvestris altera minor, flore albo nostras.

R. S. 455.

ROSA arvensis. H. F. 192.

White flowered Dog's Rose.

In Hedges — very common. June. S.

ROSA sylvestris inodora seu canina. R. S. 454.

ROSA canina. H. F. 192.

Red flowered wild Dog's Rose, or Hep Tree.

In Hedges — very common. June. S.

* *ROSA sylvestris odora.* R. S. 454.

ROSA Eglanteria. H. F. 191.

Sweet Briar.

In a Hedge near Ore-Stray — very uncommon.

June. S.

ROSA pumila spinosissima, foliis Pimpinellæ

glabris flore albo. R. S. 455.

ROSA spinosissima. H. F. 191.

The Burnet Rose.

On the North side of Chatham Hill — uncommon.

June. S.

K

RUBE-

RUBEOLA arvensis repens cærulea. R. S. 225.

SHERARDIA arvensis. H. F. 54.

Little Field Madder.

In Cades new Wood, and in Broom fields in Hernhill — not uncommon. May. A.

RUBEOLA vulgaris quadrifolia lævis floribus purpurascens. R. S. 225.

ASPERULA Cynanchica. H. F. 55.

Squinancy Wort.

In Ospringe Chalk Pits, and on Badgen Down — July. P.

RUBUS Idæus spinosus fructu rubro. R. S. 467.

RUBUS Idæus. H. F. 193.

The Raspberry Bush.

In Stocking Wood, near Lees Court, and in Perry Wood — not common. May. S.

RUBUS major fructu nigro. R. S. 467.

RUBUS fruticosus. H. F. 193.

The Bramble, or Blackberry Bush.

In Woods and Hedges — very common. June. S.

RUBUS minor fructu cæruleo. R. S. 467.

RUBUS cæsius. H. F. 193.

The small Bramble, or Dew-berry Bush.

In Woods and Hedges — common. July. S.

RUSCUS.

RUSCUS. R. S. 262.

RUSCUS *aculeatus*. H. F. 373.

Knee Holly, or Butcher's-broom.

In Ofspringe Woods and Hedges—common. April. P.

RUTA *muraria*. R. S. 122.

ASPLENIUM *Ruta muraria*. H. F. 386.

Wall Rue, White Maiden-hair, Tentwort.

*On the Buttrices near the South Door of the Church —
very uncommon.* P.

S.

SALICARIA *vulgaris purpurea foliis oblongis*.

R. S. 367.

LYTHRUM *Salicaria*. H. F. 179.

Purple-spiked Loosestrife, or Willow-herb.

*In the swampy Ground near Stone-bridge — not un-
common.* August. P.

SALICORNIA *geniculata annua*. R. S. 136.

SALICORNIA *europæa*. H. F. 1.

Marsh Sampire, Jointed Glass-wort, or Salt-wort.

In the Marshes as you go to Thorn — common. August. A.

SALIX. R. S. 447.

SALIX *alba*. H. F. 366.

The common white Willow.

In moist Places about the Powder Mills—not common.

April. S.

SALIX

SALIX folio articulato splendente flexilis. R. S. 448.

SALIX amygdalina. H. F. 363.

The round leaved shining Willow, or Almond-leaved Willow.

In a Hedge as you enter Sheldwich Road—uncommon.
May. S.

SALIX folio longissimo. R. S. 450.

SALIX viminalis. H. F. 366.

The Osier.

In moist Places—common. April. S.

SALIX latifolia rotunda. R. S. 449.

SALIX caprea. H. F. 366.

The common round-leaved Sallow.

In Woods and Hedges—common. April. S.

SALIX pumila foliis utrinque candicantibus et lanuginosis. R. S. 447.

SALIX arenaria. H. F. 364.

Sand Willow.

On the Sand Downs near Deal—plentifully. June. S.

SAMBUCUS. R. S. 461.

SAMBUCUS nigra. H. F. 112.

Common Elder.

In Hedges—very common. May. S.

SAMBU-

SAMBUCUS humilis, feu Ebulus. R. S. 461.

SAMBUCUS *Ebulus*. H. F. 113.

Dwarf Elder, Wall-wort, or Dane-wort.

By the Road sides near Boughton-Street — not uncommon. July. P.

SAMOLUS Valerandi. R. S. 283.

SAMOLUS *Valerandi*. H. F. 79.

Round-leaved Water Pimpernel.

In the Brent Marshes — common. June. P.

SANGUISORBA minor R. S. 203.

POTERIUM *Sanguisorba*. H. F. 358.

Burnet.

On the chalky Hills in Ospringe — common. July. P.

SANICULA five Diapensia. R. S. 221.

SANICULA *europæa*. H. F. 96.

Sanicle.

In Woods — very common. May. P.

SAXIFRAGA aurea. R. S. 158.

CHRYSOSPLENIUM *oppositifolium*. H. F.

156.

Golden Saxifrage.

In Jud's Wood — not common. April. P.

SAXIFRAGA rotundifolia alba. R. S. 354.

SAXIFRAGA granulata. H. F. 159.

White Saxifrage, or Sen-green.

Upon Beacon Hill — uncommon. May. P.

SAXIFRAGA verna annua. R. S. 354.

SAXIFRAGA tridactylites. H. F. 159.

Rue-leaved Saxifrage, or Whitlow-grass.

On Houses and Walls — very common. April. A.

SCABIOSA major communior folio laciniato.

R. S. 191.

SCABIOSA arvensis. H. F. 50.

Common Field Scabious.

In moist Fields among Corn, and on Bank sides — common. August. P. With a proliferous Flower in Badgen Wood, uncommon.

SCABIOSA minor vulgaris. R. S. 191.

SCABIOSA Columbaria. H. F. 50.

The lesser Field Scabious.

In dry hilly Places — not uncommon. July. P.

SCABIOSA radice succisa, flore globofo. R. S. 191.

SCABIOSA Succisa. H. F. 50.

Devil's Bit.

In Eysing Wood and the Abby Meadows — common.

August. P.

SCAN-

SCANDIX femine rostrato vulgaris. R. S. 207.

SCANDIX *Pecten Veneris*. H. F. 107.

Shepherd's Needle, or Venus Comb.

In Fields among Corn—very common. July. A.

SCORODONIA feu Salvia agrestis. R. S. 245.

TEUCRIUM. *Scorodonia*. H. F. 219.

Wood Sage.

In Byfing Wood — very common. July. P.

SCROPHULARIA aquatica major. R. S. * 283.

SCROPHULARIA *aquatica*. H. F. 239.

Water Betony, Water Figwort.

On the Banks of the River, and in moist Ditches—
common. July. P.

SCROPHULARIA major. R. S. * 283.

SCROPHULARIA *nodosa*. H. F. 239.

Common knobby-rooted Figwort.

In moist Hedges — common. July. P.

SEDUM minimum non acre flore albo. R. S.
270.

SEDUM *annuum*. H. F. 172.

Mountain Stone-crop.

On the Beach near Sandown Castle — plentifully.
August. P.

SEDUM

SEDUM minus hæmatoides. R. S. 269.

SEDUM *reflexum*. H. F. 170. β.

Yellow Stone-crop, or Prick-madam.

On the Abby Walls — common. July. P.

SEDUM minus luteum ramulis reflexis. R. S.

270.

SEDUM *reflexum*. H. F. 170.

Yellow reflexed Stone-crop.

On the Abby Walls — common. July. P.

SEDUM parvum acre flore luteo. R. S. 270.

SEDUM *acre*. H. F. 171.

Stone-crop, or Wall-pepper.

On Walls and Houses — not uncommon. July. P.

SEDUM *sexangulare*. H. F. 171.

Insipid Stone-crop.

Near Sheernefs — plentifully. July. B.

SEMPERVIVUM majus. R. S. 269.

SEMPERVIVUM *teſtorum*. H. F. 185.

House-leek, or Sengreen.

On Walls and tiling of Houses — not uncommon.

July. P.

SENECIO

SENECIO vulgaris. R. S. 178.

SENECIO vulgaris. H. F. 315.

Common Groundfel, or Simson.

In Fields and under Hedges — very common.

May. A.

SERPILLUM citratum. R. S. 230.

THYMUS Serpyllum. H. F. 229. ε.

Lemon Thyme.

On Badgen Down — not common. July. P.

SERPILLUM vulgare. R. S. 230.

THYMUS Serpyllum. H. F. 229.

Mother of Thyme.

On dry billy Ground — very common. July. P.

SERRATULA. R. S. 196.

SERRATULA tinctoria. H. F. 304.

Saw-wort.

Near the late Decoy Ponds at Graveney — uncommon.

July. P.

SESELI pratense nostras. R. S. 216.

SESELI Caruifolia. H. F. 106.

Meadow Saxifrage.

*In the low Meadows in Hernhill, and in Shepey —
not uncommon. August. P.*

SIDERI-

SIDERITIS anglica strumosa radice. R. S. 242.

STACHYS palustris. H. F. 227.

Clowns Allheal.

In moist Places, and near the River — common.

August. P.

SIDERITIS arvensis rubra. R. S. 242.

GALEOPSIS Ladanum. H. F. 225.

Narrow leaved Allheal, or Iron-wort.

In Ospringe Corn fields — common. August. A.

SIDERITIS humilis lato obtuso folio. R. S. 242.

GLECHOMA arvensis. H. F. 224.

Upright Ground Ivy.

In Ore Corn fields — common. August. A.

* SINAPI album filiqua hirsuta, semine albo vel
ruffo. R. S. 295.

SINAPIS alba. H. F. 260.

White Mustard.

By the Wayside in the Brent Marsh — very uncommon.

August. A.

SINAPI fativum secundum. R. S. 295.

SINAPIS nigra. H. F. 260.

Common Mustard.

*In Shepey amongst Corn, and by Waysides — plenti-
fully.* July. A.

SISYM-

SISYMBRIUM Cardamine seu Nasturtium aquaticum. R. S. 300.

SISYMBRIUM *Nasturtium*.

Water Cress.

By the River sides — very common. July. B. or P.

SISYMBRIUM hirsutum. R. S. 233.

MENTHA *aquatica*. H. F. 223. γ .

Water Mint.

On the sides of the River — very common. July. P.

SIUM aromaticum. Sison officinarum. R. S. 211.

SISON *Anomum*. H. F. 104.

Bastard Stone Parsley.

In a Hedge near Thorn — not uncommon. August. P.

SIUM arvense five segetum. R. S. 211.

SISON *segetum*. H. F. 104.

Corn Parsley, or Honewort.

In moist Hedges near Ewell — not uncommon. July. B.

SIUM latifolium foliis variis. R. S. 211.

SIUM *latifolium*. H. F. 103.

Great Water Parsnep.

In marshy Ground, and by the River side — common. August. P.

SIUM

SIUM pufillum foliis variis. R. S. 212.

SISON *inundatum*. H. F. 104.

The least Water Parsnep.

In the Dykes of Nagden Marsh — not common.
June. A.

SIUM umbellatum repens. R. S. 211.

SIUM *nodiflorum*. H. F. 103.

Creeping Water Parsnep.

In the Abby Ditches — common. August. P.

SMYRNIUM. R. S. 208.

SMYRNIUM *Olusatrum*. H. F. 109.

Alexanders.

In a Hedge as you enter Graveney Marsh — uncommon.
June. B.

SOLANUM lignosum seu Dulcamara. R.S. 265.

SOLANUM *Dulcamara*. H. F. 78.

Woody Night-shade, or Bitter-sweet.

In moist Woods and Hedges — common. July. P.

* SOLANUM pomo spinoso oblongo flore calathoides, Stramonium vulgo dictum. R.S. 266.

DATURA *Stramonium*.. H. F. 78.

Thorn Apple.

On Dungbills — very uncommon. August. A.

SOLA-

SOLANUM vulgare. R. S. 265.

SOLANUM nigrum. H. F. 78.

Common Night-shade.

On Rubbish and Dungbills — very common. July. A.

SONCHUS asper laciniatus. R. S. 163.

SONCHUS oleraceus. H. F. 294. γ .

Prickly Sow-thistle with jagged Leaves.

On Rubbish and in cultivated Grounds — common.

August: A.

SONCHUS asper non laciniatus. R. S. 163.

SONCHUS oleraceus. H. F. 294. δ .

Prickly Sow-thistle with less jagged Leaves.

In cultivated Grounds, and on Rubbish — common.

June to August. A.

SONCHUS lævis. R. S. 162.

SONCHUS oleraceus. H. F. 294. α .

Common Sow-thistle.

On Rubbish, in Fields and cultivated Grounds — very common. June to September. A.

SONCHUS lævis minor paucioribus laciniis.
R. S. 163.

SONCHUS oleraceus. H. F. 294. β .

Smooth Sow-thistle with less jagged Leaves.

On Rubbish, in Fields and cultivated Grounds —

June to September. A.

L

SONCHUS

SONCHUS repens, multis Hieracium majus.
R. S. 163.

SONCHUS arvensis. H. F. 295.

Tree Sow-thistle.

In Fields and Hedges — common. August. P.

SORBUS sylvestris foliis domesticæ similis.
R. S. 452.

SORBUS aucuparia. H. F. 189.

Mountain Ash, Quicken Tree, Roan Tree.

In the Blean Woods — not uncommon. May. S.

SPARGANIUM non ramosum. R. S. 437.

SPARGANIUM erectum. H. F. 346. β .

Bur-reed not branched.

*In the Ditches of Ham and Graveney Marshes —
common. July. P.*

SPARGANIUM ramosum. R. S. 437.

SPARGANIUM erectum. H. F. 346.

Great branched Bur-reed.

In Ponds near Ewell — not uncommon. July. P.

SPERGULA maritima flore parvo cæruleo, semine
vario. R. S. 351.

ARENARIA marina. H. F. 169. γ .

Small flowered Sea-spurrey.

In Shepey Island — not uncommon. July. P.

SPER-

SPERGULA purpurea. R. S. 163.

ARENARIA rubra. H. F. 169.

Purple Spurrey, Purple flowered Chickweed.

On sandy Grounds in Hernhill — not uncommon.

July. A.

SPHONDYLUM. R. S. 205.

HERACLEUM *Sphondylium*. H. F. 102.

Cow Parsnep.

On the sides of Fields, and in Hedges — common.

July. B.

STATICE montana minor. R. S. 203.

STATICE *Armeria*. H. F. 114.

Thrift, or Sea Gilly-Flower.

In all the Salt Marshes—very common. July. P.

STELLARIA. R. S. 289.

CALLITRICHE *verna*. H. F. 2.

Vernal Star wort, Water wort, Star-headed Chickweed.

In the Powder Mill Streams—common. April. A.

STELLARIA aquatica foliis longis tenuissimis.

R. S. 290.

CALLITRICHE *autumnalis*. H. F. 2.

Autumnal Star-wort.

In the River opposite the Gunpowder Stove — not uncommon. September. A.

STRATIOTES foliis Afari, femine rotundo.
R. S. 290.

HYDROCHARIS *Morsus ranæ*. H. F. 372.
Frog-bit.

In the Dykes of Ham and Graveney Marshes —
common. June. P.

SYMPHYTUM magnum. R. S. 230.

SYMPHYTUM *officinale*. H. F. 68.
Comfrey.

On the Banks of the Powder Mill Waters — uncommon.
July. P.

T.

TAMNUS *racemosa*, flore minore luteo pallescente. R. S. 262.

TAMUS *communis*. H. F. 369.
Black Briony.

Under Hedges, and among Buskes — common. June. P.

TANACETUM. R. S. 188.

TANACETUM *vulgare*. H. F. 310.
Common Tanfy.

By Way sides, near Queen Court and Uplees — very
uncommon. August. P.

TAXUS. R. S. 445.

TAXUS *baccata*. H. F. 372.
The Yew Tree.

In Woods and Hedges on a chalky Soil — common.
April. S. THA-

THALICTRUM seu Thaliectrum majus. R. S.
203.

THALICTRUM *flavum*. H. F. 216.

Meadow Rue.

In the shooting Meadow Stream — uncommon.
June. P.

TITHYMALUS characias Amygdaloides. R. S.
312.

EUPHORBIA *Amygdaloides*. H. F. 184.

Wood Spurge.

*In Woods, and under Hedges in Ospringe — very
common.* April. P.

TITHYMALUS helioscopius. R. S. 313.

EUPHORBIA *Helioscopia*. H. F. 183.

Sun Spunge, or Wart-wort.

In Meadows, and on Dungbills — common. July. A.

TITHYMALUS leptophyllos. R. S. 313.

EUPHORBIA *exigua*. H. F. 182.

Dwarf Spurge, Small annual Spurge.

In Corn fields and Gardens — common. July. A.

TITHYMALUS parvus annuus, foliis subrotundis
non crenatis, Peplus dictus. R. S. 313.

EUPHORBIA *Peplus*. H. F. 182.

Petty Spurge.

In cultivated Grounds — common. July. A.

TITHYMALUS platyphyllos Fuchsii. R. S.
312.

EUPHORBIA platyphyllos. H. F. 184.

Broad leaved Spurge.

In Corn fields near Thorn Creek — common.
July. P.

TORMENTILLA. R. S. 257.

TORMENTILLA erecta. H. F. 197.

Tormentil Septfoil.

In Woods and Hedges — common. June. P.

TORMENTILLA reptans. R. S. 257.

TORMENTILLA reptans. H. F. 198.

Creeping Tormentil.

In Broom fields at Hernhill — common. July. P.

TRAGOPOGON luteum. R. S. 171.

TRAGOPOGON pratense. H. F. 293.

Yellow Goat's Beard, Go to Bed at Noon.

In Fields and Pastures — common. June. B.

TRAGOPOGON purpureum. R. S. 172.

TRAGOPOGON porrifolium. H. F. 293.

Purple Goat's Beard.

In a Marsh near Sheernefs — uncommon.
June. B.

TRICHO.

TRICHOMANES. R. S. 119.

ASPLENIUM *Trichomanes*. H. F. 385.

English black Maiden-Hair.

On the Walls of the Church and Abby——not common. P.

TRIFOLIUM *arvense humile spicatum* feu

Lagopus. R. S. 330.

TRIFOLIUM *arvense*. H. F. 285.

Hare's Foot, or Hare's Foot Trefoil.

Upon Beacon Hill — not uncommon. July. A.

TRIFOLIUM *cochleatum folio cordato maculato*.

R. S. 333.

MEDICAGO *arabica*. H. F. 288.

Heart Trefoil, or Clover.

In the Abby Meadows——not uncommon. June. A.

TRIFOLIUM *cochleatum modiolis spinosis*.

R. S. 333.

MEDICAGO *arabica*. H. F. 288. γ.

Hedge-hog Trefoil, or Clover.

In Pastures near Sheernefs——not common. May. A.

TRIFOLIUM *fragiferum*. R. S. 329.

TRIFOLIUM *fragiferum*. H. F. 286.

Strawberry Trefoil.

In the Abby Meadows — common. August. P.

TRIFO-

TRIFOLIUM flosculis albis, in glomerulis oblongis asperis, cauliculis proxime adnatis.

R. S. 329.

TRIFOLIUM *scabrum*. H. F. 285.

Oval headed Trefoil.

In Pastures in Shepey — common. June. A.

TRIFOLIUM lupulinum alterum minus. R. S. 330.

TRIFOLIUM *procumbens*. H. F. 287.

The lesser Hop Trefoil, Decumbent Trefoil.

In Meadows — common. May to August. P.

TRIFOLIUM lupulinum minimum. R. S. 331.

TRIFOLIUM *filiforme*. H. F. 287.

Small Trefoil.

On gravelly Banks beyond South Street — uncommon.

May. A.

TRIFOLIUM pratense album. R. S. 327.

TRIFOLIUM *repens*. H. F. 283.

Dutch Clover, White flowered Meadow Trefoil.

In Fields and Meadows — very common. May to September. P.

TRIFOLIUM pratense luteum capitulo Lupuli, vel agrarium. R. S. 330.

TRIFOLIUM *agrarium*. H. F. 286.

Hop Trefoil.

On Beacon Hill — not uncommon. June. A.

TRIFO-

TRIFOLIUM pratense purpureum. R. S. 328.

TRIFOLIUM *pratense*. H. F. 284.

Common Purple or Honey-suckle Trefoil.

In Fields and Meadows — very common. May to
September. P.

TRIFOLIUM purpureum majus, folis longioribus
et angustioribus, floribus saturatioribus.
R. S. 328.

TRIFOLIUM *medium*. H. F. 284.

Long leaved Purple Trefoil with deeper coloured
Flowers.

In Byfing Wood, and by Field sides in Ospringe —
not uncommon. July. P.

TRIFOLIUM stellatum glabrum. R. S. 329.

TRIFOLIUM *maritimum*. H. F. 284.

Teasel-headed Trefoil.

On Shellnefs in Shepy — uncommon. June. P.

TURRITIS muralis minor. R. S. 294.

TURRITIS *hirsuta*. H. F. 254.

Hairy Tower Mustard.

On old Walls at Colkins in Boughton — very uncommon. May. B.

TUSSILAGO. R. S. 173.

TUSSILAGO *Farfara*. H. F. 315.

Common Colt's Foot.

In moist Fields — very common. April. P.

TYPHA.

TYPHA. RS. 436.

TYPHA *latifolia*. H. F. 345.

Great Cat's Tail, or Reed Mace.

On the sides of Ponds, and in Dykes — very common.
July. P.

TYPHA *palustris media*. R. S. 436.

TYPHA *angustifolia*. H. F. 345.

Narrow leaved Cat's Tail.

In the Ditches at Ewell and Nagden — not uncommon.
July. P.

V.

VALERIANA *sylvestris major*. R. S. 200.

VALERIANA *officinalis*. H. F. 12.

Great wild Valerian.

On the Banks of Ofpringe River — not uncommon.
June. P.

VALERIANA *sylvestris minor*. R. S. 200.

VALERIANA *dioica*. H. F. 12.

Small wild or Marsh Valerian.

On the Banks of the shooting Meadow Stream —
common. June. P.

VALERIANELLA *arvensis præcox humilis*
femine compresso, sive Lactuca agnina.
R. S. 201.

VALERIANA *Locusta*. H. F. 12.

Lamb's Lettuce, or Corn Sallad.

In Corn fields, and rich cultivated Ground — common.
May. A. VALE-

VALERIANELLÆ vulgaris species major, ferotina. R. S. 201.

VALERIANA *Locusta*. H. F. 13. β.

Late flowering Lambs Lettuce, or Corn Sallad.

In the Corn fields between Ore and Harty-ferry — not common. July. A.

VERBASCUM flore albo parvo. R. S. 287.

VERBASCUM *Lychnitis*. H. F. 76.

Hoary Mullein, or white Mullein.

On old Walls at Davington — not common. July. B.

VERBASCUM mas latifolium luteum. R. S. 287.

VERBASCUM *Thapsus*. H. F. 75.

Great white Mullein, High Taper, Cows Lungwort.

On dry Banks by Road sides — common. July. B.

VERBASCUM nigrum flore parvo, apicibus purpureis. R. S. 288.

VERBASCUM *nigrum*. H. F. 76.

Sage-leaved black Mullein.

On the sides of Chalk fields in Ospringe — very uncommon. As you ascend St. Martin's-Hill near Canterbury — plentifully. July. P.

VERBENA vulgaris. R. S. 236.

VERBENA *officinalis*. H. F. 505.

Vervain.

By Road sides and Foot paths — very common. July. P.

VERBE-

VERBESINA feu Cannabina aquatica flore minus
pulchro elatior. R. S. 187.

BIDENS tripartita. H. F. 309.

Trifid Water-hemp Agrimony.

In marshy watery Places — common. July. A.

VERBESINA pulchriore flore luteo. R. S. 187.

BIDENS cernua. H. F. 309.

Whole leaved Water-hemp Agrimony.

In Dykes near the Town Key—common. August. A.

VERONICA aquatica angustifolia minor. R. S.
280.

VERONICA scutellata. H. F. 5.

Narrow leaved Water Speedwell, or Brooklime.

In wet Ditches and marshy Ground—common. June. P.

VERONICA aquatica longifolia media. R. S. 280.

VERONICA Anagallis. H. F. 5.

The middle long leaved Water Speedwell, or
Brooklime.

In the Abby Ditches — common. July. P.

VERONICA aquatica rotundifolia Becabunga
dicta minor. R. S. 280.

VERONICA Becabunga. H. F. 4.

Common Brooklime.

*In wet Ditches and small running Waters — very
common. May. P.*

VERONICA

VERONICA Chamædrys sylvestris dicta. R. S. 281.

VERONICA Chamædrys. H. F. 5.

Wild Germander.

In Fields and Meadows — very common. May. P.

VERONICA floribus singularibus, in oblongis
pediculis, Chamædryfolia. R. S. 279.

VERONICA agrestis. H. F. 6.

Germander Speedwell, or Chickweed.

In Fields, on Rubbish, and in cultivated Ground —
common. May. A.

VERONICA flosculis singularibus Hederulæ folio,
Morsus gallinæ minor dicta. R. S. 280.

VERONICA hederifolia. H. F. 6.

Ivy leaved Speedwell, or small Henbit.

In dry Fields, and on old Walls — common. April. A.

VERONICA mas supina et vulgarissima. R. S.
281.

VERONICA officinalis. H. F. 4.

The Male Speedwell, or Fluellin.

In barren dry Fields — common. May. P.

VERONICA pretenfis minor ; sive Betonica Pauli.
R. S. 279.

VERONICA serpyllifolia. H. F. 4.

Little, or smooth Speedwell, or Paul's Betony.

In Fields and Meadows — common. May. P.

VIBURNUM. R. S. 460.

VIBURNUM *Lantana*. H. F. 112.

The pliant Mealy, or Way-faring Tree.

In Hedges — common. May. S.

VICIA. R. S. 320.

VICIA *sativa*. H. F. 278.

Common Vetch, or Tare.

In Fields among Corn — not uncommon. June. A.

VICIA *sepium perennis*. R. S. 320.

VICIA *sepium*. H. F. 278.

Bush Vetch.

In bushy Places and Hedges — common. May. P.

VICIA *minima præcox Parisiense*. R. S. 321.

ERVUM *solanense*. H. F. 279.

Spring Tare.

In Ospringe Chalk Pits — common. April. P.

VICIA *sylvestris* sive *Cracca major*. R. S. 321.

VICIA *lathyroides*. H. F. 279.

Strangle Tare, or Wild Vetch.

In sandy Fields in Hernhill — not uncommon.

May. A.

VINCA

VINCA *Pervinca minor*. R. S. 268.

VINCA *minor*. H. F. 77.

Periwinkle.

In Cades Wood, and Hedges in Preston and Ospringe
— *common* April. P. With a white Flower,
near the 39th Mile Stone — *very uncommon*.

VIOLA *bicolor arvensis*. R. S. 366.

VIOLA *tricolor*. H. F. 331. β .

Pansies, Heart's-ease, Two Faces under a Hood.
In sandy Corn fields at Hernhill — *common*. Au-
gust. A.

VIOLA *martia inodora sylvestris*. R. S. 364.

VIOLA *canina*. H. F. 331.

Wild or Dogs Violet.

In Woods and under Hedges — *very common*. April. P.

VIOLA *martia hirsuta inodora*. R. S. 365.

VIOLA *hirta*. H. F. 330.

Hairy Violet.

In the new Wood at Cades in Ospringe — *plentifully*.
March. P.

VIOLA *martia alba*. R. S. 364.

VIOLA *odorata*. H. F. 330. β .

White Violet.

Under dry Hedges and Banks, by Way sides — *not un-*
common. March. P.

VIOLA *maritia purpurea*. R. S. 364.

VIOLA *odorata*. H. F. 330.

Purple sweet Violet.

Under Hedges, and in Fields — very common.

March. P.

VIOLA *tricolor*. R. S. 361.

VIOLA *tricolor*. H. F. 331.

Pansies, Three Faces under a Hood.

In Corn fields at Ore, and in Gardens — common.

June. A.

VIRGA *aurea*. R. S. 176.

SOLIDAGO *Virga aurea*. H. F. 318.

Common Golden Rod.

In Weeds and Hedges — common. August. P.

VISCUM. R. S. 464.

VISCUM *album*. H. F. 367.

Mistle, or Mistletoe.

On Trees, especially in Orchards — common.

May. S.

VULNERARIA *rustica*. R. S. 325.

ANTHYLLIS *Vulneraria*. H. F. 273.

Kidney Vetch, or Ladies Finger.

About Byfing Wood — common. July. P.

ULMA

U.

ULMARIA. R. S. 259.

SPIRÆA *Ulmaria*. H. F. 191.

Meadow sweet.

In moist Meadows in Davington—common. July. P.

ULMUS folio glabro.

ULMUS *glabra*. H. F. 95. β.

The smooth leaved or Wych Elm.

In Hedges—not uncommon. April. S.

ULMUS folio latissimo scabro. R. S. 469.

ULMUS *glabra*. H. F. 95.

The Wych Hazel, or broad-leaved Elm.

In Hedges at Throwleigh—uncommon. April. S.

ULMUS *vulgatissima*, folio lato scabro. R. S.
468.

ULMUS *campestris*. H. F. 94.

The common Elm.

In Hedges—very common. April. S.

URTICA minor. R. S. 140.

URTICA *urens*. H. F. 355.

The lesser Stinging Nettle.

In cultivated and uncultivated Ground—very common.

August. A.

URTICA racemifera major perennis. R. S. 139.

URTICA dioica. H. F. 355.

The common Stinging Nettle.

Under Hedges, and on Rubbish — very common
July. P.

PLANTS OMITTED.

* CALENDULA officinalis. Lin. Syst. Nat. II.

P. 577.

Garden Marigold.

On the Beach at St. Margaret's at Cliff; plentifully.

And in a Field at Woodnesborow — very
uncommon. June. A.

CRAMBE maritima brassicæ folio. R. S. 307.

CRAMBE maritima. H. F. 260.

Sea Colewort.

At St. Margaret's at Cliff — uncommon. June. P.

HYDROCERATOPHYLLON folio aspero quatuor cornibus armato R. S. 135.

CERATOPHYLLUM demersum. H. F. 357.

Horned Pond-weed.

In Marsh Dykes — Common. June. P.

* MYAGRUM. R. S. 302.

MYAGRUM sativum. H. F. 242.

Gold of Pleasure.

Near Sandwich, among the Flax — plentifully.

June. A.

TRI-

TRIFOLIUM luteum lupulinum. R. S. 331.

MEDICAGO lupulina. H. F. 288.

Melilot Trefoil.

In Paths near Westwood — not common. May to August. A.

CAUCULIS nodosa echinato femine. R. S. 220.

TORDILIUM nodosum. H. F. 98.

Knotted Parsley.

By the Road side at Minster in Shepey — not common May. A.

LINARIA antirrhinum dicta. R. S. 283.

ANTIRRHINUM minus. H. F. 238.

The least Toad Flax.

In a Corn field near Whitehill, Selling — uncommon. A.

SENECIO minor latiore folio, five montana.

R. S. 178.

SENECIO sylvaticus. H. F. 316.

Mountain Groundsel.

In Badgen Wood — not common. August. A.

OENANTHE aquatica triflora. R. S. 220:

OENANTHE fistulosa. H. F. 104. β .

The lesser Water Dropwort.

In Ditches at Graveney — common. July. P.

1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900

1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930

1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960

1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990

1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020

FOSSILIA SHEPEIANA.

A
SHORT VIEW
OF THE
FOSSIL BODIES
Native and Extraneous,
OF THE
ISLAND
OF
SHEPEY,
IN THE
COUNTY OF KENT.

Quicquid sub terra est in apricum proferet ætas.

INTRODUCTION.

THE Island of Shepey, with the annexed Peninsulas of Harty and Elmly, surrounded by the Waters of the Thames, the Medway, and the Sea, is upwards of Thirty Miles in Circumference, about Thirteen in Length from Shellness to Sheerness, and in its greatest Breadth about Six. It hath long been noted for producing large Quantities of Sheep (from whence probably it derived its Name) as well as Corn; and exhibits to the inquisitive Naturalist a most desirable Spot, by affording on its Surface many rare Plants, and more especially in the Bowels of its Northern Cliffs, so great a Quantity and Variety of Fossils, both native and extraneous, as are scarcely to be paralleled.

These Cliffs are in Length about Six Miles from East to West, gradually declining at each End; the more elevated Parts whereof reach about two-thirds of their extension, and are, at the very highest of them about Minster, not less than Thirty Yards in perpendicular Height above the Beach or Shore, and being composed of Clay, and constantly washed at their Basis by the
Tides

Tides which beat against them more especially when driven by strong Easterly Winds, they are continually wasting more or less, and falling down upon the Shore, where the Clay being by degrees washed away, the Fossils are left in great Abundance. So great is the Loss of the Land at the highest Parts, that sometimes near an Acre of it hath sunk down at once, though these Falls are indeed extraordinary.

Minster, Shurland, and Warden, are the three Manors to which these Cliffs appertain, the Proprietors whereof hire them out to different Owners of Coperas Works, who employ the neighbouring Poor to collect the Pyrites or Coperas Stone upon the Beach, at the Rate of about a Shilling per Buskel, which they deposit in Heaps, when collected, until a Quantity sufficient to load a Hoy is procured, when it is measured and paid for.

These Coperas Gatherers, from the Author's first Visits were induced to preserve and carry home every Thing that seemed to their Apprehension somewhat particular, and having since met with Encouragement, and acquired some Information relative to what was most desirable by the Curious, they have generally from that Time reserved such, until Inquiries have been made for them. Many rare Specimens, both native and extraneous, far beyond any equal Space of Ground, perhaps

perhaps the Universe can produce, are by these means preserved. Indeed, more may thus be obtained, at a very easy Expence, than by the most diligent Fossilists alone, as they cannot allow due Time to examine the Shore sufficiently. This is evidently confirmed by comparing the Specimens from this Coast, recited in the Works of our most celebrated * Naturalists, with those in the ensuing Catalogue. The Author doubts not that many curious Specimens have escaped his Inquiries, and as the Cliffs are daily dropping their Contents more or less, he is fully assured that there will always remain a Variety sufficient to gratify every Lover of the Study, whenever they may choose to take a Trip thither. But alas! One disagreeable Circumstance attending a considerable Part of the Fossils here collected, is, that they are so much impregnated with Pyritical Matter, that after being for some Time placed in a Cabinet, the Salts thereof shoot and entirely destroy them. Happy would it be, could some certain Remedy be discovered whereby this Accident might be prevented. The Loss of many valuable Specimens by this Cause, together with his Distance from any inquisitive and able Naturalists, at last induced the Author to dispose of his whole Collection to INGHAM FOSTER, Esq;

* Dr. Grew, Mr. Lihuyd, Mr. Petiver, Dr. Woodward,
Mr. Dale, and Dr. Hill.

Merchant of London, where it now forms no inconsiderable Part of that Gentleman's very valuable Cabinet.

He, many Years since, entertained Thoughts of publishing a descriptive Catalogue of all the Fossils there, by him collected, but was prevented, as well by more necessary Attachments as the Difficulty of the Undertaking: However, it having been intimated to him that a concise Account of them would not be deemed an unacceptable Appendix to his PLANTÆ FAVERSHAMIENSES, in which several of the Shepey Plants are recited, as no special Account of these Fossils hath been offered to the Public, and as those that have been already noted by the before cited Authors are dispersed in their Works, and as many others have not as yet been published, he complied with the Proposal, and set about revising his Notes and Observations, and now offers this short Account of the Discoveries made there in his annual Visits for above these Thirty Years past; at first solely for the Purpose of collecting these elegant Bodies, and afterwards to superintend his private Affairs, having acquired some Property in the Island, not far distant from the Cliffs.

What the candid Reader may expect in the following Catalogue is, a general Account of the Variety of the Fossil Bodies, both native and extraneous, reduced into

some Order under particular Heads, with Notice of their degrees of Rarity. If what is offered shall afford any Information worthy the attention of the Lovers of Natural History, the Author will think himself sufficiently rewarded for this Attempt of exhibiting the Out-lines of a Work which merits the Abilities of a Person of more Judgment and greater Leisure, to enlarge and perfect, than he pretends to be possessed of. He cannot conclude however, without observing, that from the contemplation of so great a Variety of extraneous Fossils discovered in these Cliffs, which were evidently the produce of very different Climates, he thinks himself rationally induced to believe, that nothing short of an universal Deluge could be a Cause adequate to such an Effect.

E. J.

FOSSILIA SHEPEIANA.

NATIVE FOSSILS.

THE CLIFFS.

THESE are composed of three different Strata of brown Clay; the uppermost is of the lightest Colour, and most friable; the lowest darkest, and of a closer Texture.

PYRITÆ.

Copras Stones, are the most frequent of all the Fossils of the Cliffs: They are of various Forms and Sizes; round, botryoide, flat, varying in every Degree, as well in Figure as Colour, and impregnating most of the other Fossils found here.

LUDUS HELMONTII:

The Waxed Vein; Nodules of which are lodged in the Cliffs in great Numbers; they are generally flat and oval, and of various Sizes, sometimes exceeding two Feet in length, all are covered with Coats

in the Cliffs, but are frequently seen denudated on the Beach; the *Tali* are of different Sizes and Forms, generally they pass strait from one flat Surface of the Nodule to the other; in some there is a double Set of them, running only half through the Body, with intermediate sparry Partitions; which being generally of a waxen Colour, at first gave this Nodule its Name. The Number of the *Tali* in different Nodules are indeterminate; both the *Tali* and *Septa* vary in Colour in different Specimens — they are very common.

BEZOAR MINERALE.

Bodies, which are crustated, ferrugineous, and of a rounded Form, covered with various Crusts of the Colour of the Clay of the Cliffs, and of different Sizes — These are common.

C A L C U L I.

Pebbles. Amongst the numbers on the Beach, there have been some Specimens collected, which by polishing have proved very elegant.

S E L E N I T E S.

Selenite, is found here in great Plenty, chiefly of the columnar sort, some Specimens exceed six Inches in length, but most are of shorter dimensions,

fions, which are common. The rhomboidal Sort is not near so common.

LEPASTRUM. D. HILL.

The Starry Waxed Vein, is a Fossil peculiar to these Cliffs, not having as yet been discovered elsewhere to the Author's Knowledge. Dr. Hill hath made from their Forms three Sorts of it, to which others might be added if nicely attended to. It is ever affixed to the Septa of the Ludus Helmontii, where the Fissures are so far divided as to permit the Stars to shoot. The largest Specimens do not exceed three Inches in Length, and these are very rare; those are generally of a much less Size that are common.

SUCCINUM.

Amber, is very uncommon; a few Specimens only, and those small and transparent, are to be found here.

EXTRANEOUS FOSSILS.

VEGETABLES.

LIGNUM FOSSILE.

Fossil Wood, in large Masses, seemingly of Oak from its Grain, even to above three Feet in Length,

is plentifully dispersed on the Beach, and generally impregnated with pyritical Matter.

EQUISETUM.

Horse-tail, Joints of the naked Species, are sometimes found even to the Length of three Inches, but they are uncommon.

FRUCTUS VARII.

Fossil Fruits, of very different Kinds, are here collected, in Variety and Number exceeding all that have been mentioned by Writers on this Branch of Natural History: Many of them have been drawn and compared with recent ones, in a particular Account given of them by the Author's late learned Friend, Dr. *James Parsons*, in the *Philosophical Transactions* — Vol. L. Part I. p. 396. These have by some been imagined mere Sports of Nature amongst the Coperas; but as all the Specimens have not the misfortune to be impregnated by it, some few, and those very elegant ones, being entirely free from it, and petrified, that Conceit without further Discussion, must fall to the Ground.

ARISTÆ.

Ears of Corn, some small Species, are sometimes to be met with, imperfect and impregnated with the Pyrites, but they are not common.

MYCE-

M Y C E T I T Æ.

The Mushroom Stone. Here are two Kinds, the Conoide and Discoide, both small, and the first is by far the most frequently to be found.

A N I M A L I U M P A R T E S.

E L E P H A S.

The Elephant. A considerable Number of the Bones of this very large Animal have been discovered on the Eastern Shore, a Mile from the Cliffs, on the Author's Manor of *Nutts*, in *Laysdown*; an Account of which may be seen in the Philosophical Transactions — Vol. xlviii. part ii. page 626.

T E S T U D O.

The Tortoise. Two varieties have been found under the Cliffs, one of a more convexed Form, the other more depressed; both are extremely rare.

P I S C I U M P A R T E S.

The Heads of Fish, from a very small Size to that of a middling Cod, are not unfrequent.

Palates of Fish, some oval, others of a squarish form, when perfect, or nearly so, are very uncommon;

common ; but small Portions of them, especially of the square Sort are common : To these last, Mr. *Llbuyl* hath given the whimsical Name of *Scopula littoralis*, and to those of the first kind, *Bufo nitæ*. In the Gentleman's Magazine for May, 1755, p. 409, is a Plate exhibiting both Kinds, in fine Preservation.

Teeth of Fish, usually called *Sharks Teeth*, are plentifully met with, except those of a large Size, which are very uncommon.

Back Bones of Fish, are very common in separate Joints, but with three or more united, are uncommon.

Parts of the Bodies of Fish, with the Scales on them, are not very uncommon, but none entire have come to the Author's Knowledge.

Tails of Fish, are frequently collected, and some few with a Vertebra united to them ; there are very few of a large Size.

CANCRI MACROURI.

Lobsters, are not uncommon, but none of the Specimens are perfect ; the Bodies and Tails united are not so frequent as when separated ; the Claws are also found separately of various Sizes, but generally much injured.

Prawns

Prawns, are very rare; they are nearly perfect, and considerably larger than those taken recent on the opposite Coast of the Isle of *Thanet*.

CANCRI BRACHYURI.

Fossil Crabs. Before the Publication of Dr. *Hill's* History of *Fossils*, these were extremely rare in *England*; but since that Time, thousands of Specimens here collected, have passed through the Author's Hands. There are two or three Varieties, but all common except very fair Specimens, which are still very rare; their Size varies from that of a small Bean, to that of the common *Crab* of the Coast; some of them, especially the smaller ones, are impregnated by the *Pyrites*.

ECHINITÆ.

Echinites, here are the *Ovarius*, *Galeatus*, *Pileatus*, *Cordatus*, and *Placentiformis*; but all of them are very rare, especially the last.

The Spines, are of the foliated Kind, and that Sort named by Mr. *Llbuys*, *Volvola*—both are rare.

ASTERIAS.

The Star Fish of five Rays, has not yet been found entire, and its single Rays are uncommon.

ASTROITÆ

ASTROITÆ ET ASTEROPODIUM.

Star Stones, with depressed Rays, and the Roots of them are very uncommon; the Specimens of the latter Kind were so imperfect, that without better Information the Author was at a Loss how to place them in his Collection.

BELEMNITES.

Belemnite, only one Specimen, and that imperfect, hath hitherto been discovered.

CONCHYLIA.

TUBULI MARINI.

Worm Shells, of various contorted Forms, generally of a small Size, some with their Shells, others denudated, frequently impregnated with the Coperas, are common.

LAPIS SYRINGOIDES.

The piped Waxen Vein, externally appears like the *Ludus Helmontii*, being found in incrusted Nudoles; they are evidently Oak that had first been eroded by Sea Worms, and afterwards their Perforations filled with Pyritical Matter and Spar,
similar

similar to the *Waxen Vein*, surrounding each Perforation ; these are very common.

CONCHYLIA UNIVALVIA.

DENTALIA.

The Tooth Shells, generally small, about an Inch long, curving towards the Apex in the more perfect Specimens, some covered with their Shells, and others denudated ; both Sorts are common.

PATELLA.

The Limpet, is a very rare Fossil, having only once been collected ; the Specimen was small, of the striated Sort, and somewhat imperfect.

NAUTILUS.

The chambered Nautilus, is a common Fossil of these Cliffs. These vary in Size from that of Hazel-Nut to that of a Man's Head, most commonly they have their pearly Shell, and are often immersed in Nodules ; there are also some small ones found of a ferrugineous Colour ; very few of the large ones are perfect, having received Injury by compression.

ORTHO-

ORTHOCERATITES.

Small Fragments seemingly of this *Genus* are found here, but very uncommon; the Chambers or Partitions appear to be united by a Dove-tail Suture.

TUBULI CONCAMERATI.

Cylindrical Bodies, with several Concamerations passing transversely through them, are generally of the Size of a Swan's Quill, and about two or three Inches in Length, some petrified, others wholly pyritical, and are not uncommon.

COCHLEÆ.

Snail Shells, with a short as well as with a produced Clavicle, some with their Shells, others denuded, are generally of a small Size, and impregnated by the *Pyrites*, and are uncommon.

TROCHI.

The Topshells, are of different Forms, some flatter, others more elevated, of various Sizes, some with Shells on, others denuded, most are impregnated with *Pyrites*, and are common.

CONCHA

CONCHA VENERIS.

The Cowry. This is small, denudated, pyritical, and extremely rare.

BUCCINA.

Whilkes, are found in considerable Quantities, varying in Form and Size, but generally small.

The Opercula are sometimes, though rarely seen.

NERITÆ.

The Nerites, are common; and generally impregnated with Pyrites.

TURBINES.

The Screw Shells, and casts of them without the Shell, of various Sizes, and mostly impregnated with Pyrites, are very common.

CONCHYLIA BIVALVIA.

OSTREA.

Fossil Oysters; generally of a small Size, are found here, but very uncommon; although the whole Coast is covered with recent ones.

O

CONCHÆ

CONCHÆ ANOMIÆ.

These Shells in great variety, tho' generally of a small Size, and impregnated with the Pyrites, are very common.

PECTUNCULI.

Cockles, smooth, striated, and fasciated, of a small Size, are found in tolerable Plenty; but fair and large Specimens are uncommon.

TELLINÆ.

The Wedge Shell. Small Specimens of which, generally denudated and pyritical, are very common.

MUSCULI.

Muscles. These are extremely rare, a few large Specimens, and one only of the common Muscle have been hitherto met with; although the Coast abounds with the latter.

PINNÆ.

The Sea Wing. Only small Fragments of this Shell have as yet been discovered, and these are very rare.

F I N I S.

I N D E X

OF THE

ENGLISH NAMES.

A.

	Page
Abele Tree	87
Adder's Tongue	73
Agrimony	2
Dutch, or common Hemp	36
Alder, common	3
black Berry-bearing	39
Alexanders	108
Alkanet, evergreen	15
bastard	16
Ale-hoof	17
All-good	14
All-heal, Clowns	106
narrow leaved	106
All-feed, upright Blite	25
least Rupture-wort	92
Anemony Wood	6
Angelica, wild	6
Apple-Thorn	108
Archangel, red	53
yellow	54
white	52
Arsmart,	

	Page
Arfinart, common	82
dead or spotted	81
greater, with pale leaves	82
with spotted stalks	81
willow-leaved, perennial	82
Ash Tree	39
mountain	110
Ash-weed	6
Asp, or Aspen	87
Avens, common	21
water	22
B.	
Ba'm, garden	67
Basil, wild	2
great wild	28
Base Rocket	95
Batchelor's-buttons	91
Bear's-foot	45
Bedstraw, Ladies-yellow	40
Beech Tree	37
Beet, sea	12
Bell-flower, lesser round leaved	18
Berberry Bush	12
Betony, water	103
wood	12
Bindweed, black	37
great	28
small	28
sea	28
Birch Tree	13
Bird's Foot	77
trefoil	62
the greater	61

	Page
Bird's Nest, or Wild Carrot	32
Bitter-sweet	108
Blackberry	98
Blite, late flowering, or Fig-leaved	13
the least	14
maple leaved	26
upright	26
round-leaved, or All-feed	25
sea	14
Blood-wort	55
Borage	15
Bottle, blue	31
white	64
Box Tree	17
Brakes, common	38
Bramble, or Briar, common	98
small	98
Briar, sweet	97
Briony, black	112
white	15
Broom, common	40
rape	78
Brooklime, common	120
middle, long leaved	120
narrow leaved	120
Buckbean	67
Buckthorn	96
sea	96
Buck's-horn plantain	84
Buck-wheat, or Brank	37
Bugle	16
Bugloss, small wild	15
vipers	33
English vipers	33
Bullace	

	Page
Bullae Tree, black	90
white	90
Burdock	56
with smaller heads	56
taller, with a larger flower	56
with wooly heads	57
Burnet	101
Burnet-saxifrage, great	83
lesser round leaved	84
small	84
Bur-reed, great branched	110
not branched	110
Bush Vetch	122
Butcher's Broom	99
Butter-bur, common	82
Butter-cups	93

C.

Cabbage, sea	15
Calamint, common	18
Calf's-foot	7
Caltrops water, greater	88
lesser	88
Cammock	7
Camomile, sweet-scented creeping	25
Campion-bladder	64
corn, white	64
greater corn	64
red flowered	64
wild white	64
Dover	62
sea	62
narrow leaved	63
	Can-

I N D E X.

v

	Page
Canterbury Bells	18
Carrot, wild	32
Cat-mint	
Cat's-tail, great	118
lesser	118
Celandine, greater	79
lesser	25
Centory, lesser	24
lesser, with a white Flower	24
yellow	23
Charlock, white	96
or white Mustard	95
yellow	95
Cheese-rening	40
Chier, wild	59
Cherry wild, common	24
Cherry-wood	73
Chervil, small Hemlock	71
wild	24
Chesnut Tree	23
Chick-weed, breakstone	5
least	4
marsh, great	4
mouse-ear, broader leaved	3
narrow leaved	3
least	24
sea	4
purple flowered	111
star-headed	111
Chickweed or Speedwell	121
Cicely, wild	27
Cinque-foil, common	81
Tormentil	80
Cistus,	

	Page
Cistus, dwarf, or little Sun-flower	44
Clary, common English wild	47
Climber, great wild	27
Cleavers	7
Clot-bur	56
with smaller heads	56
taller, with a purple flower	56
with wooly heads	57
Clover, Dutch	116
Clown's All-heal	106
Cockle	63
Cock's comb, great marsh, red rattled	80
red rattle	80
yellow	80
Cock's-head	73
Codlings and Cream	65
Colewort, sea	126
Colt's-foot, common	117
Columbines	8
Comfrey	112
Corn, bottles	31
Feverfew	25
Marygold	26
Mint	68
Parsley, small	23
Sallad	118
late flowering	119
small	52
Cornel, female	29
Cotton Thistle	21
Cow's Lung-wort	119
Cow Parsnep	111
Cowslip, great	89
common	89
Cow-	

I N D E X.

vii

	Page
Cow-weed	27
Cow-wheat, common, wild	67
Crab Tree	66
Crane's-bill, dove's-foot	41
field	41
hemlock leaved	41
jagged leaved	42
long stalked	42
small flowered	42
sea	42
Cresses, water	107
winter, or Rocket	35
Crimson Grass Vetch	23
Cross-wort	30
Crowfoot, common creeping	94
corn	93
finest leaved	92
ivy leaved	93
round, rooted, or bulbous	93
sweet-wood	94
water fennel leaved	92
fine leaved	93
round leaved	94
various leaved	93
upright meadow	94
Crow-garlick	3
Cuckow flower	63
pint	8
Cudweed, black headed	43
common	43
least	43
long leaved, low branched	43
upright	43
Cur-	

	Page
Currants, black	96
red	97
D.	
Daffodil, wild English	71
Daifie, common	12
greater	58
Dandelion	32
rough, or Dandelion Hawkweed	32
Deptford Pink	22
Devil's-bit	102
yellow	47
guts	31
Dewberry	98
Dittander	58
Dock, common broad leaved	56
great water	55
greener leaved	56
red leaved	55
fiddle	55
sharp pointed	54
leffer	55
with curled leaves	55
Dodder	31
Dogberry	29
Dog's Mercury	69
Rose, red flowered	144
white flowered	143
Violet	123
Dove's-foot, common	41
jagged leaved	42
small flowered	42
Dropwort	37
	Drop-

I N D E X.

ix

	Page
Dropwort Hemlock	73
water	73
Duck's-meat	58
greater	58
ivy leaved	58
Dwale	12
Dwarf Elder	101
Dyer's Weed	41

E.

Earth-nut	16
Eglantine	97
Elder, common	100
Dwarf	101
water	73
Elecampane	44
Elm, broad leaved	125
common	125
smooth-leaved, or Wych	125
Enchanter's Nightshade	27
Eryngo, Sea Holly	35
Eyebright	36
red	36

F.

Fennel, common	38
Hog, or Sulphur-wort	38
Fern, common male	38
female, or common brakes	38
small branched stone	38
Feverfew	67
corn	25
field	25

Figwort,

	Page
Figwort, common knobby rooted	103
water	103
Finkle	38
Five leaved grafs	81
Flax, purging or wild dwarf	61
narrow leaved wild	60
perennial blue	61
Fleabane, middle	29
small	29
blue flowered	9
Flix-weed	36
Flower-de-luce, yellow water	51
Fluellin, round leaved	60
sharp pointed	60
Fool's Parsley	27
stones, female	75
male	75
Foxglove, purple	32
French Mercury, male and female	69
Frog-bit	112
lettuce	88
Fumitory	39
great white	39
Furze	41
needle	40

G.

Gentian, dwarf autumnal	41
Germander-speedwell, or Chickweed	121
wild	121
Gilli-flower, sea	111
Gladdon, stinking	51
	Glaſs.

I N D E X.

xi

	Page
Glass-wort, white	14
jointed	51
prickly	51
Goat's-beard, purple	114
yellow	114
Gooseberry	44
Gold of Pleasure	126
Goldilocks	94
Golden rod, common	124
Goose-foot, common	14
sharp leaved	14
Goose-grass	7
marsh	7
Goose Tongue	91
Gorse	41
Goule	40
Grass-vetch, crimson	23
Green-wood	41
Go to bed at noon	114
Gout-weed	6
Gromwell, or Gromile	61
Ground Ivy	17
upright	106
Groundsel, common	105
mountain	127

H.

Hare bells	48
Hare's ear, the least	16
Hare's foot	115
Hart's tongue	83
Hawk nut, or Hog-nut	16
Hawk-	

	Page
Hawkweed, with bitten roots	47
bushy	46
narrow leaved bushy	46
long rooted	46
smooth succory	47
rough succory	47
Hawthorn	69
Hazel-nut Tree	29
Hazel-wych	125
Heart's-ease	123
Three faces under a hood	124
Heath common, or Ling	35
cross leaved	34
fine leaved	34
Pea, or wood pea	78
Hedge mustard	35
nettle	40
parsley	23
Hellebore, great bastard black	45
wild black	44
broad leaved bastard	45
white flowered bastard	45
Hell-weed	31
Hemlock	26
Hemlock Chervil, small with rough leaves	71
dropwort	73
lesser, or Fool's parley	27
water	83
Hemp-agrimony, common or Dutch	36
Water trifid	120
whole leaved	120
Henbane, common	48
Henbit, great	53
small	121
Hep-	

Page

Hep-tree	97
Herb, Bennet, or common Avens	21
Gerard	6
Paris	45
Robert	42
Two-pence	72
High-taper	119
Hog-fennel	83
Hog-nut	16
Holly-tree	2
Hone-wort	107
Honeysuckle, common	18
trefoil	117
Hop, male and female	62
trefoil	116
Horehound, stinking	11
water	64
white	66
Horn-beam Tree, or Horse-beech	78
Horse-radish	96
Horse-tail, corn	33
lesser, marsh	34
river	34
wood	34
Horse-shoe Vetch	37
Hound's-tongue, great	31
lesser green leaved	31
House-leek	104
Hyacinth, English	48

I.

Jack-by-the-hedge	46
Iron-wort	106
b	Ivy,

	Page
Ivy, climbing or berried	44
Juniper, common	51

K.

Kidney-vetch	124
wort	29
Knapweed, common	50
great	50
Knawel, annual	52
Knee-holly	99
Knot-grafs, common	86
German	52
oblong narrow leaved	86
small narrow leaved	86

L.

Ladies Bedstraw, mountain	70
white	70
yellow	40
Smock	18
small flowered	19
Finger	124
Thistle	20
unspotted	20
Traces, triple	77
Lamb's lettuce	118
late flowering	119
Lang-de-beuf	46
Larkspur, wild	32
Lavender, sea	60
Laurel, dwarf	57
Lettuce,	

	Page
Lettuce, Ivy leaved wild	52
least wild	52
Lilly-convally	59
water white	72
yellow	72
Ling	35
Live-long	5
Lords and Ladies	8
Loose-strife, greater smooth-leaved	64
least smooth-leaved coddled	65
lesser hairy coddled with small flowers	65
middle smooth-leaved	65
purple spiked	99
yellow	65
Loufewort, common	80
marsh, red rattle, Cock's-comb	80
Liquorice, wild or Vetch	9

M.

Madder, little field	98
great bastard	70
small mountain	70
wild, or great bastard	70
Madwort, German	19
Maiden hair, white	99
common black	2
English	115
Mallow, common	60
small wild, or dwarf	66
marsh	5
Maple, common	1
greater, or Sycamore	1
b 2	Mares's-

	Page
Mare's-tail	59
Marigold, corn	26
marsh	86
garden	126
Marjoram, pot	77
wild	77
Matfellow, common	50
great	50
May Lilly	59
weed, stinking	24
Meadow Pink	63
Rue	113
Sweet	125
Medick-Vetchling	73
Melilot	67
trefoil	127
Mercury, common English	14
dog's	69
French, male and female	69
Milfoil	69
water, feathered or spiked	87
verticillated	81
Milk-wort	85
sea	43
Mill-mountain	61
Mint, Nep or Nap	72
long leaved horse	68
round leaved horse	68
spear	68
water	68
with whorled coronets	68
Missel or Misseltoe	124
Money-wort	72
purple flowered	72
Moscatel,	

	Page
Moscatel, tuberous	70
Mother-wort	19
Mother of thyme, common	105
Mountain Ash	110
Mouse-ear scorpion grass	70
common creeping	83
water scorpion grass	71
Mug-weed	30
Mug-wort	8
Mullein, great white	119
hoary or white	119
sage leaved black	119
Musk Thistle	20
Mustard, common	106
hedge	35
tower, lesser	117
white	106
Myrtle, Dutch	40

N.

Navel-wort	29
Navew, wild	71
Needle Furze	40
Nep or Nap, or Cat-Mint	72
Nettle, dead	52
hedge	40
hemp-leaved, with a parti-coloured flower	53
white flowered	53
red	53
yellow	54
stinging, common	126
lesser	125
b 3	Nettle-

	Page
Nettle-hemp, white flowered	53
fair flowered	53
yellow	54
Nightshade, common	109
deadly	12
Enchanter's	27
woody	108
Nipple-wort	54
Nose-bleed	69
Nut-earth, Kipper, Pig, Hawk or Hog	16

O.

Oak Tree, common	91
with acorns on short-foot stalks	91
Oculus Christi, or Christ's eye	47
One Berry	45
Orach, common wild	13
grass leaved	10
narrow leaved wild	10, 11
with a spear pointed leaf	10
jagged sea	10
wild sea narrow leaved	10
wild sea	11
stinking	13
Orchis, female	75
female handed	76
male, or male fool's stones	75
male handed	76
butterfly	74
bee	74
bee hairy, or humble bee	77
fly	75
frog	76

Orchis,

Orchis, man green	74
purple	75
Neapolitan	74
little purple flowered	76
purple late flowering	76
yellow or musk	75
bird's nest	74
ladies traces	77
Orpine	5
Ofier	100
Ox-eye	58
lips	89
tongue	46

P.

Pagils, common	89
Panflies, two faces under a hood	123
three faces under a hood	124
Park-leaves	49
hairy	48
Parfley, garden	82
corn or Honewort	107
corn small	23
hedge	23
knotted	127
stone bastard	107
Parfnep, Cow	111
great water	107
least water	108
water creeping	108
wild	79
Paul's Betony	12
Pea,	

	Page
Pea, everlasting broad leaved	57
narrow leaved	57
wood or heath	78
Pear, wild	91
Pellitory of the wall	79
bastard	91
Penny royal	91
Penny-wort, marsh	48
wall	29
Pepper-wort	58
Pepper, water	82
Periwinkle, the greater	27
the lesser	123
Pestilent Wort	82
Petty Whin	40
Pig-nut or Hog-nut	16
Pilewort	25
Pimpernel, male	5
yellow	5
round leaved water	101
Pin Pearl-wort	5
Pink, Deptford	22
meadow	63
Pipperidge Bush	12
Plantain, Buck's-horn	84
great	84
hoary	85
ribwort	85
sea	85
narrow leaved sea	85
water great	84
star-headed	31
Polypody	86
	Pond-

Pond-weed, broad leaved	89
fine or fennel leaved	88
grafs leaved	88
horned	126
perfoliated	89
fea	88
Poplar Tree, black	87
white	87
trembling	87
Poppy, Corn	79
red	79
spatling	64
yellow horned	78
Plowman's spikenard	11
Prick-madam	104
Prick-wood, Cornel-tree	29
Spindle tree	36
Primrose, common	89
Privet or Prim	59
Pudding-grafs	91
Purflane, water	87
fea	11
Pursley Piert	81

Q.

Quick	69
Quicken Tree	110

R.

Radish, Horfe	96
Water	92
Ragwort,	

	Page
Ragwort, broad leaved marsh	50
common	50
hoary perennial with Groundfel leaves	50
Ramsons	3
Raspberry	98
Rattle, yellow	80
red marsh	80
Reed Mace	118
lesser	118
Rest-harrow, Corn	7
spinous	7
creeping	6
Ribwort Plantain	85
Roan Tree	110
Rocket, or Winter Cresses	35
base	95
water	35
sea	17
Rose-bay, Willow herb	66
Rose, Burnet	97
dog's red flowered	97
white flowered	97
Rue, meadow	113
wall	99
whitlow grafs	102
Rupture-wort, the least	92
Rush, flowering	17

S.

Sage wood	103
Saint-foin	73
	Saint

Saint John's-wort	48
hairy	48
imperforate	49
least trailing	49
upright	49
Saint Peter's-wort	49
Sallow, common round leaved	100
Salt-wort, annual	99
black	43
Sampire	30
golden	9
marsh	99
perennial	51
Sanicle	101
Satyrion Royal, male handed	76
female handed	76
Sauce alone	46
Saw-wort	105
Saxifrage, common golden	101
meadow	105
white	102
Rue-leaved	102
Scabious, common field	102
lesser	102
Sheep's hairy	95
Scad Tree, or Scad Plumb	90
Scorpion-grafs	70
water	71
Sea Bindweed	28
Buckthorn or fallow	96
Cabbage	15
Campion	62
Colewort	126
Pond-weed	88
	Sea

	Page
Sea Holly	35
Heath smooth	63
Gilly-flower	111
Rocket	17
Lavender	60
Plantain	85
narrow leaved	85
Spurrey, small flowered	110
Starwort	9
Scurvy-grass	28
Self-heal	90
Sengreen, white	103
common	104
Septfoil	114
Service Tree, common	69
Setter-wort	45
Shepherd's Needle	103
Purfe	17
Rod	33
Silver Weed	80
Simson	105
Sloe Tree	90
Smallage	8
Snap-dragon, greater	7
Sneeze-wort	91
Snow-drop	59
Soap-wort, common	63
Sorb Tree	69
Sorrel, common	54
Sheep's	54
Wood	78
Sow-bane	14
	Sow-

I N D E X.

xxv

	Page
Sow-thistle, common	109
prickly with jagged leaves	109
with less jagged leaves	109
smooth with less jagged leaves	109
Tree	110
Spear-thistle	19
Spear-wort, great	94
lesser	94
Speedwell, or Chickweed	121
Germander	121
ivy-leaved	121
little smooth	121
male	121
water, narrow leaved	120
with long leaves	120
Spikenard, Plowman's	11
Spatling Poppy	64
Spindle Tree	36
Spleenwort, or Miltwaste	9
rough	61
Spurge, broad leaved	114
dwarf, small annual	113
Laurel	57
petty	113
Sun, or Wart-wort	113
Wood	113
Spurrey, Corn	4
purple	111
sea	4
small flowered	110
Squinancy wort	98
Star-headed Water Chickweed	111
Star of the Earth	84
Thistle	21
2	Star-

	Page
Star-wort, autumnal	111
vernal	111
sea	9
Stich-wort, greater	22
least	5
lesser	22
middle, a variety	22
Stonecrop, yellow	104
Wall Pepper	104
yellow reflexed	104
mountain	103
insipid	104
Stone Parsley, bastard	107
Strangle Tare	122
Strawberry	38
barren	39
trefoil	115
Succory, wild	26
Sulphur-wort	83
Sun flower, little	44
Sweet Briar	97
Swine's Cress	71
Sycamore	1

T.

Tansey, common	112
wild	80
Tare, common	122
everlasting	57
hairy	30
small wild	30
smooth	30
spring	122
Tare,	

Tare, strangle	122
tine	30
with smooth pods	30
Teasel, wild	33
small wild	33
Tent-wort	99
Thistle, common way, or creeping	21
Carline, dwarf	19
wild	21
Cotton	21
Marsh	20
Milk, or Lady's	20
without spots	20
Musk	20
Spear	19
Star	21
up n Thistle	19
Weked, with small flowers	20
Thorn Apple	108
Thorn, black	90
white	69
common purging	90
sea buck	96
Thorough wax	16
Three faces under a hood	124
Thrift	111
Throat-wort, great	18
Thyme, mother of	105
Lemon	105
Tine Tare	30
with smooth pods	30
Toad Flax, common yellow	60
the least	127
Tooth-wort	6

	Page
Tormentil	114
Cinquefoil	80
creeping	114
Tower Mustard, hairy	117
Traveller's Joy	27
Tree, pliant mealy	122
Trefoil, Bird's foot	62
greater	61
common purple	117
Hare's-foot	115
Heart	115
Hedge-hog	115
Hop	116
lesser	116
long leaved purple	117
Marsh	67
Melilot	127
oval headed	116
Teasel headed	117
small	116
Strawberry	115
white flowered meadow	116
Triple Ladies Traces	77
Tutsan	49
hairy	48
Turnep	95
Tway-blade, common	13
Two faces under a hood	123

V.

Valerian, great wild	118
jagged leaved	52
small wild or marsh	118
Venus-	

I N D E X.

xxix

	Page
Venus-comb	103
Vervain	119
mallow	2
Vetch Bush	122
Crimson grass	23
common	122
tufted	30
wild	122
spring	122
tufted Horse shoe	37
Kidney, or Lady's finger	124
Vetchling, common yellow bastard	57
Medick	73
Violet, Dog's or wild	123
purple sweet	124
Throat wort leaved	123
white	123
water	47
hairy	123
Viper's Bugloss	33
English	33

W.

Wake Robin	8
Wall-flower	59
Penny-wort	29
Pepper	104
Rue	99
wort	101
Wart-wort	113
Water Betony	103
Caltrops, greater	88
lesser	88

Water

	Page
Water Crow-foot, fennel leaved	92
fine leaved	93
finest leaved	92
Ivy leaved	93
round leaved	94
various leaved	93
Cresses	107
Dock, great	79
Dropwort	73
Hemlock	73
Elder	73
Figwort	103
Gladiole	17
Hemlock	83
Hemp agrimony, trifid	120
with an undivided leaf	120
Horehound	64
Lilly, white	72
yellow	73
Mill-foil, feathered or spiked	87
verticillated	81
Mint	107
hairy	107
with whorled coronets	68
Parfnep, creeping	108
great	107
least	108
Pepper	82
Plantain, great	84
star-headed	31
Purslane	87
Ragwort	50
Water	

I N D E X.

	xxx
	Page
Water Speedwell, middle long leaved	120
narrow leaved	120
Wort	111
Violet	47
Way-bread	84
Way-faring Tree	122
Weed, yellow	62
Whin	41
petty	7
White-glaswort	14
White-rot	48
White-thorn	69
Whitlow-grafs, common	79
rue leaved	102
Williams, wild	63
Willow, common white	99
round eared shining	100
sand	100
sweet	40
Willow-herb, purple spiked	99
great flowered	65
greater smooth leaved	64
least smooth leaved coddled	65
lesser hairy coddled with small	
flowers	65
middle smooth leaved	65
Rose bay	66
yellow	65
Winter Cresses	35
Woad, wild	62
Wood Anemone	6
Betony	12
bine	18
roof	8

Wood

	Page
Wood Pea	78
Sorrel	78
waxen	59
Sage	103
Spurge	113
Wormwood, common	I
white sea	I
Wort, white glaſs	14
Wych Elm	125
Hazel	125

Y.

Yarrow, common	69
Yellow-weed, common	62
Yew Tree	112

I N D E X

OF THE

L I N N Æ A N N A M E S.

A.	Page
Acer campestre	1
Pseudo platenus	1
Achillea Millefolium	69
Ptarmica	91
Adoxa Moschatellina	70
Aegopodium Podagraria	6
Aethusa Cynapium	27
Agrimonia Eupatoria	2
Agrostemma Githago	63
Ajuga reptans	16
Alisma Damasonium	31
Plantago	84
Allium vineale	3
Urfinum	3
Althæa officinalis	5
Amarantus Blitum	14
Anagallis arvensis	5
Anchusa sempervirens	15
Anemone nemorosa	6
Anethum	

	Page
Anethum Fœniculum	38
Angelica sylvestris	6
Anthemis Cotula	24
nobilis	25
Anthyllis Vulneraria	124
Antirrhinum minus	127
Elatine	60
Linaria	60
majus	7
spurium	60
Aphanes arvensis	81
Apium graveolens	8
Petroselinum	82
Aquilegia vulgaris	8
Arctium Lappa β . γ . δ .	56, 57
Arenaria marina β .	4
marina γ .	110
rubra	111
peplodes	4
serpyllifolia	4
Artemisia Absinthium	1
maritima	1
vulgaris	8
Arum maculatum	8
Asperula odorata	8
Cynanchica	98
Asplenium Adiantum nigrum	2
Ceterach	9
Ruta muraria	99
Scolopendrium	83
Trichomanes	115
Asttragalus glycyphyllos	9
Aster Tripolium	9
	Atriplex

Atriplex hastata	10
hastata β .	11
laciniata	10
litoralis β .	11
litoralis	10
patula	10
ferrata	10
portula coides	11
Atropa Belladonna	12

B.

Ballota nigra	11
Bellis perennis	12
Berberis vulgaris	12
Beta vulgaris	12
Betonica officinalis	12
Betula alba	13
Alnus	3
Bidens tripartita	120
cernua	120
Borago officinalis	15
Brassica Napus	71
Rapa	95
oleracea	15
Blackstonia perfoliata	23
Bryonia alba	15
Bunium Bulbocastanum	16
Bunias Cakile	17
Bupleurum rotundifolium	16
tenuissimum	16
Butomus umbellatus	17
Buxus sempervirens	17

C.

	Page
Calendula officinalis	126
Callitriche autumnalis	111
verna	111
Caltha palustris	86
Campanula rotundifolia	18
Trachelium	18
Cardamine parviflora	19
pratensis	18
Carduus acanthoides	20
acaulos	19
crispus	19
lanceolatis	19
Marianus	20
Marianus β .	20
nutans	20
palustris	20
Carlina vulgaris	21
Carpinus Betulus	78
Caucalis Anthriscus	23
arvensis	23
Centaurea Calcitrapa	21
Cyanus	31
Jacea	50
Scabiosa	50
Cerastium aquaticum	4
femidecandrium	24
viscosum	3
vulgatum	3
Ceratophyllum demersum	126
Chærophyllum sylvestre	27
temulum	24

Cheiran-

	Page
Cheiranthus Cheiri	59
Chelidonium majus	79
Glaucium	78
Chenopodium album	13
Bonus Henricus	14
hybridum	26
maritimum	14
murale	14
polyspermum	25
rubrum	14
ferotinum	13
Vulvaria	13
urbicum	26
Chrysanthemum Leucanthemum	58
fegetum	26
Chrysofplenium oppositifolium	101
Cichoreum Intybus	26
Circœa Lutetiana	27
Cistus Helianthemum	44
Clematis Vitalba	27
Clinopodium vulgare	28
Cochlearia anglica	28
Armoracia	96
Coronopus	71
Conium maculatum	26
Convallaria majalis	59
Convolvulus arvensis	28
sepium	28
Soldanella	28
Conyza squarrosa	11
Cornus sanguinea	29
Corylus Avellana	29
Cotyledon Umbilicus Veneris	29
Crambe maritima	126

	Page
Cratægus Oxyacantha	69
torminalis	69
Crepis tectorum	47
biennis	47
Crithmum maritimum	30
Cucubalus Behen	64
viscosus	62
Cuscuta europæa	31
Cynoglossum officinale	31
officinale β .	31

D.

Daphne Laureola	57
Datura Stramonium	108
Daucus Carota	32
Delphinium Consolida	32
Dianthus Armeria	22
Digitalis purpurea	32
Dipsacus pilosus	33
sylvestris	33
Draba verna	79

E.

Echium anglicum	33
vulgare	33
Epilobium angustifolium	66
hirsutum	65
montanum	64
palustre	65
ramosum	65
tetragonum	65

Equi-

I N D E X.

xxxix

Page

Equisetum arvense	33
fluviatile	34
palustre	34
sylvaticum	34
Erica cinerea	34
Tetralix	34
vulgaris	35
Erigeron acre	9
Ervum hirsutum	30
solonienſe	122
tetraspermum	30
Eryngium maritimum	35
Eryſinum Alliaria	46
Barbarea	35
officinalis	35
Euonymus europæus	36
Eupatorium cannabinum	36
Euphorbia Amygdaloides	113
exigua	113
Helioscopia	113
Peplus	113
platyphyllos	114
Euphraſia officinalis	36
Odontites	36

F.

Fagus Caſtanea	23
ſylvaticus	37
Ficaria verna	25
Filago germanica	43
montana	43
Fragraria ſterilis	39
veſca	38

d 2

Frankenia

	Page
Frankenia lævis	63
Fraxinus excelsior	39
Fumaria officinalis	39
claviculata	39

G.

Galanthus nivalis	59
Galeopsis Ladanum	105
Tetrahit	53
β.	53
δ.	53
Galeobdolon	54
Galium Aparine	7
uliginosum	7
verum	40
Moliugo	70
montanum	70
palustre	70
Genista anglica	40
tinctoria	41
Gentiana Amarella	41
Centaurium	24
Geranium cicutarium	41
columbinum	42
dissectum	42
molle	41
pusillum	42
maritimum	42
robertianum	42
Geum urbanum	21
rivale	22
Glaux maritima	43
Glechoma	

	Page
Glechoma arvensis	106
hederacea	17
Gnaphalium sylvaticum	43
uliginosum	43

H.

Hedera Helix	44
Hedysarum Onobrychis	73
Helleborus foetidus	45
viridis	44
Heracleum Spondylium	111
Hieracium Pilosella	83
fabaudum	46
umbellatum	46
Hippocrepis comosa	37
Hippophae Rhamnoides	96
Hippuris vulgaris	59
Hottonia palustris	47
Humulus Lupulus	62
Hyacinthus non scriptus	48
Hydrocharis Morfus ranæ	112
Hydrocotyle vulgaris	48
Hyoscyamus niger	48
Hypericum Androsæmum	49
hirsutum	48
humifusum	49
montanum	49
perforatum	48
pulchrum	49
quadrangulum	49
Hypochæris radicata	46

I.

	Page
Jasione montana	95
Ilex Aquifolium	2
Inula crithmoides	9
dysenterica	29
Helenium	44
Pulicaria	29
Juniperus communis	51
Iris foetidissima	51
Pseudacorus	51

L.

Lactuca saligna	52
Lamium album	52
amplexicaule	53
rubrum	53
Lapsana communis	54
Lathræa squamaria	6
Lathyrus latifolius	57
Nissolia	23
pratensis	57
sylvestris	57
Lemna minor	58
polyrrhiza	58
trifolca	58
Leontodon autumnale	47
hispidum	32
Taraxacum	32
Leonurus Cardiaca	19
Lepidium latifolium	59
Ligustrum vulgare	59
Linum	

	Page
Linum catharticum	61
perenne	61
Radiola	92
tenuifolium	60
Lithospermum officinale	61
arvense	16
Lithrum Salicaria	99
Lonicera Periclymenum	18
Lotus corniculata	62
γ.	61
Lychnis dioica	64
β.	64
Flos cuculi	63
Lycopsis arvensis	15
Lycopus europæus	64
Lyfimachia nemorum	5
tenella	72
vulgaris	65
Nummularia	72

M.

Malva Alcea	2
rotundifolia	66
fylvestris	66
Marrubium vulgare	66
Matricaria Chamomilla	25
inodora	25
Parthenium	67
Medicago arabica	115
arabica γ.	115
lupulina	127
Melampyrum sylvaticum	67

Melissa

	Page
Melissa Calamintha	18
officinalis	67
Mentha aquatica	68
γ.	107
arvensis	68
longifolia	68
rotundifolia	68
spicata	68
Pulegium	91
Menyanthes trifolia	67
Mercurialis annua	69
perennis	69
Myrica Gale	40
Myosotis scorpioides	70
δ.	71
Myriophyllum spicatum	87
verticillatum	81
Myagrū sativum	126

N.

Narcissus Pseudo-Narcissus	71
Nepeta Cataria	72
Nymphæa alba	72
lutea	72

O.

Oenanthe crocata	73
fistulosa	73
β.	127
Ononis arvensis	7
repens	6
spinosa	7

Onopor-

	Page
Onopordum Acanthium	21
Ophioglossum vulgatum	73
Ophrys Anthropophora	74
apifera	74
apifera β .	77
Monorchis	75
Nidus avis	74
muscifera	75
ovata	13
spiralis	77
Orchis bifolia	74
latifolia	76
mascula	75
maculata	76
militaris	74
Morio	75
purpurea	75
pyramidalis	76
ustulata	76
Origanum onites	17
vulgare	77
Ornithopus perpusillus	77
Orobanche major	78
Orobis tuberosus	78
Osmunda spicant	61
Oxalis Acetosella	78

P.

Papaver Rhœas	79
Parietaria officinalis	79
Paris quadrifolia	45
Pastinaca sativa	79
	Pedicu-

	Page
Pedicularis palustris	80
sylvatica	80
Peplis Portulaca	87
Pteris aquilina	38
aquilina β .	38
Peucedanum officinale	83
Phellandrium aquaticum	83
Picris Echioides	46
Pimpinella major	83
saxifraga	84
β .	84
Plantago Coronopus	84
lanceolata	85
Leoflingii	85
major	84
maritima	85
media	85
Polygala vulgaris	85
Polygonum amphibium	82
aviculare	86
β . et γ .	86
Convolvulus	37
Fagopyrum	37
Hydropiper	82
pensylvanicum	82
Persicaria	81
β .	81
Polypodium Filix mas	83
vulgare	86
Populus alba	87
nigra	87
tremula	87

	Page
Potentilla Argentina	80
argentea	80
reptans	81
Potamogeton crispum	88
natans	89
marinum	88
gramineum	88
pectinatum	88
perfoliatum	89
ferratum	88
Poterium Sanguisorba	101
Prenanthes muralis	52
Primula veris α .	89
β .	89
vulgaris	89
Prunella vulgaris	90
Prunus Avium	24
infititia	90
infititia β . et γ .	90
spinosa	90
Pyrus Malus	66
communis	91

Q.

Quercus Robur	91
β .	91

R.

Ranunculus acris	94
aquatilis	93
aquatilis γ . δ .	92
aquatilis β .	93
arvensis	93

Ranun-

	Page
Ranunculus Auricomus	94
bulbosus	93
Flammula	94
hederaceus	93
Lingua	94
repens	94
sceleratus	94
Raphanus Raphanistrum	96
	β.
	95
Reseda lutea	95
Luteola	62
Rhamnus catharticus	96
Frangula	39
Rhinanthus Crista galli	80
Rosa arvensis	97
canina	97
Eglanteria	97
spinosissima	97
Ribes rubrum	97
nigrum	96
Uva crispa	44
Rubus cæsius	98
fructicosus	98
Idæus	98
Rumex Acetosa	54
Acetosella	54
acutus	54
Britannicus	55
crispus	55
obtusifolius	56
pulcher	55
sanguineus	55
Rumicis acuti varietas	55, 56
Ruscus	

Ruscus aculeatus

Page

99

S.

Sagina erecta

5

procumbens

5

Salix alba

99

arenaria

100

amygdalina

100

capræa

100

vimenalis

100

Salicornia europæa β .

51

europæa

99

Salsola Kali

51

Salvia Verbenaca

47

Sambucus Ebulus

101

nigra

100

Samolus Valerandi

101

Sanicula europæa

101

Saponaria officinalis

63

Satyrium viride

76

Saxifraga trydaetylites

102

granulata

102

Scabiosa arvensis

102

Columbaria

102

Succisa

102

Scandix Anthriscus

71

Pecten Veneris

103

Sedum acre

104

annuum

103

reflexum

104

reflexum β .

104

Telephium

5

sexangulare

104

c

Scleran-

	Page
Scleranthus annuus	52
Sempervivum tectorum	104
Senecio aquaticus	50
erucifolius	50
Jacobæa	50
fylvaticus	127
vulgaris	105
Serapias latifolia β .	45
latifolia	45
longifolia	45
Scrophularia aquatica	103
nodosa	103
Serratula arvensis	21
tinctoria	105
Seseli Caruifolia	105
Sherardia arvensis	98
Silene amœna	62
conoidea	63
Sinapis alba	106
arvensis	95
nigra	106
Sison Amomum	107
inundatum	108
segetum	107
Sisymbrium amphibium α .	92
Nasturtium	107
Sophia	36
fylvestre	35
Sium latifolium	107
nodiflorum	103
Smyrnium Olusatrum	108
Solanum Dulcamara	108
nigrum	109
Solidago Virga aurea	124

I N D E X.

	li
	Page
Sonchus arvensis	110
oleraceus α . β . γ . δ .	109
Sorbus aucuparia	110
Sparganium erectum β .	110
Spartium scoparium	40
Spergula arvensis	4
Spiræa Filipendula	37
Ulmaria	125
Stachys palustris	106
sylvatica	40
Stellaria graminea	22
graminea β .	22
Holoosteia	22
Statice Armeria	111
Limonium	60
Symphytum officinale	112

T.

Tamus communis	112
Tanacetum vulgare	112
Taxus baccata	112
Teucrium Scorodonia	103
Thalictrum flavum	113
Thlaspi Bursa Pastoris	17
Thymus Acinos	2
Serpyllum	105
Serpyllum ϵ .	105
Tordylium nodosum	127
Tormentilla erecta	114
reptans	114
Tragopogon pratense	114
porrifolium	114
c 2	Trifolium

	Page
Trifolium agrarium	116
arvense	115
filiforme	116
fragiferum	115
medium	117
Melilotus officinalis	67
maritimum	117
pratense	117
procumbens	116
repens	116
scabrum	116
Tussilago Farfara	117
Petasites	82
Turritis hirsuta	117
Typha latifolia	118
angustifolia	118

V.

Valantia Cruciata	30
Valeriana dioica	118
officinalis	118
Locusta	118
Locusta α .	52
Verbascum Lychnitis	119
nigrum	119
Thapsus	119
Verbena officinalis	119
Veronica agrestis	121
Anagallis	120
arvensis	121
Becabunga	120
Chamædrys	121
hederifolia	121

Veronica

I N D E X.

liii

Page

Veronica officinalis	121
serpyllifolia	121
scutellata	120
Viburnum Lantana	122
Opulus	73
Vicia Cracca	30
Lathyroides	122
fativa	122
sepium	122
Vinca major	27
minor	123
Viola canina	123
hirta	123
odorata	124
odorata β .	123
tricolor; et β .	123
Viscum album	124

U.

Ulex europæus	41
Ulmus campestris	125
glabra; et β .	125
Urtica dioica	126
urens	126

1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9
10	10	10
11	11	11
12	12	12
13	13	13
14	14	14
15	15	15
16	16	16
17	17	17
18	18	18
19	19	19
20	20	20
21	21	21
22	22	22
23	23	23
24	24	24
25	25	25
26	26	26
27	27	27
28	28	28
29	29	29
30	30	30
31	31	31
32	32	32
33	33	33
34	34	34
35	35	35
36	36	36
37	37	37
38	38	38
39	39	39
40	40	40
41	41	41
42	42	42
43	43	43
44	44	44
45	45	45
46	46	46
47	47	47
48	48	48
49	49	49
50	50	50
51	51	51
52	52	52
53	53	53
54	54	54
55	55	55
56	56	56
57	57	57
58	58	58
59	59	59
60	60	60
61	61	61
62	62	62
63	63	63
64	64	64
65	65	65
66	66	66
67	67	67
68	68	68
69	69	69
70	70	70
71	71	71
72	72	72
73	73	73
74	74	74
75	75	75
76	76	76
77	77	77
78	78	78
79	79	79
80	80	80
81	81	81
82	82	82
83	83	83
84	84	84
85	85	85
86	86	86
87	87	87
88	88	88
89	89	89
90	90	90
91	91	91
92	92	92
93	93	93
94	94	94
95	95	95
96	96	96
97	97	97
98	98	98
99	99	99
100	100	100

1	1	1
2	2	2
3	3	3
4	4	4
5	5	5

E R R A T A.

The candid Reader is desired to correct these
Errors and Omissions.

Page	line
4	20 <i>after marina add</i> β.
17	9 <i>for Pluckley read</i> Hedcorne.
28	19 <i>after Sandwich add</i> and on Shellness.
36	4 <i>after Sittingbourn add</i> and on the Standard Key.
42	25 <i>after Deal add</i> and on Shellness.
49	20 <i>for elegantissimum read</i> elegantissimum.
—	22 <i>for montenum read</i> montanum.
56	22 <i>after 304 add</i> γ.
57	2 <i>after 304 add</i> δ.
59	5 <i>for uncommon read</i> common.
—	12 <i>for common read</i> uncommon.
73	4 <i>for Fairbrook read</i> Luddenham.
76	21 <i>after plentifully add</i> on Badgen Downs, sparingly.
88	14 <i>for granineum read</i> gramineum.
144	16 <i>for uncommon read</i> common.

Omitted to be inserted in their proper Places.

ANTIRRHINUM angustifolium sylvestre.

R. S. * 283.

ANTIRRHINUM Orontium. H. F. 239.

The least Snapdragon, or Calf's-snout.

In a Corn field at Luddenham — uncommon.

July. A.

CASSIDA palustris vulgatiores flore cæruleo.

R. S. 244.

SCUTELLARIA galericulata. H. F. 231.

Hooded Willow-herb.

In a moist Hedge at Chartham Hatch — uncommon.

August. P.