QGE

782

S68X

1926

INVERT.

ZOOL.

9QE
782
56DX
1926
March 1, 1926

Am. Mus. Nat. Hist.

SMITHSONIAN INSTITUTION

AMERICAN SILURIAN CRINOIDS

BY

FRANK SPRINGER

LAS VEGAS, NEW MEXICO

ASSOCIATE IN PALEONTOLOGY, U. S. NATIONAL MUSEUM

(Publication 2871)

CITY OF WASHINGTON
PUBLISHED BY THE SMITHSONIAN INSTITUTION

1926

The Lord Baltimore Press
BALTIMORE, MD., U. S. A.

CONTENTS

	PAGE
INTRODUCTION	I
Early Geologists	I
The Tennessee Area	I
The Indiana Area	6
The New York Area	7
The Foreign Silurian	8
Origin of the work	9
Collections used	10
Acknowledgments	11
Order CAMERATA	12
Family DIMEROCRINIDAE	12
<i>Dimerocrinus</i>	12
<i>Eudimerocrinus</i> , new genus	14
<i>Cyphocrinus</i>	14
<i>Gazacrinus</i>	15
<i>Lamptrocrinus</i>	19
<i>Genus and species indet.</i>	21
Family RHODOCRINIDAE	22
<i>Paragazacrinus</i> , new genus	22
<i>Paulocrinus</i> , new genus	22
<i>Wilsonicrinus</i> , new genus	23
<i>Eudimicrocrinus</i> , new genus	24
<i>Genus and species indet.</i>	24
<i>Lyriocrinus</i>	24
Family MELOCRINIDAE	25
<i>Macrostylocrinus</i>	25
<i>Melocrinus</i>	27
<i>Cytocrinus</i>	28
<i>Mariacrinus</i>	28
<i>Alloocrinus</i>	30
<i>Patellioocrinus</i>	31
<i>Genus and species indet.</i>	32
<i>Laurelocrinus</i> , new genus	32
<i>Clonocrinus</i>	34
Family CALYPTOCRINIDAE	35
<i>Eucalyptocrinus</i>	35
<i>Callicrinus</i>	39
Family BATOCRINIDAE	41
<i>Carpocrinus</i>	42
<i>Desmidocrinus</i>	43
<i>Cylicoocrinus</i>	43
<i>Periechocrinus</i>	44
<i>Saccocrinus</i>	45
<i>Aorocrinus</i>	47
Family PLATYCRINIDAE	48
<i>Lyonicrinus</i> , new genus	48
<i>Myrtillocrinus</i>	49
<i>Coccocrinus</i>	51
<i>Culicocrinus</i>	51
<i>Genus and species indet.</i> , No. 1.	51
<i>Genus and species indet.</i> , No. 2.	51
<i>Genus and species indet.</i> , No. 3.	52
<i>Hapalocrinus</i>	52
<i>Brahmacrinus</i>	53
<i>Nyctocrinus</i>	54
<i>Marsipocrinus</i>	55

	PAGE
Order FLEXIBILIA	64
Family LECANOCRINIDAE	64
<i>Lecanocrinus</i>	65
<i>Anisocrinus</i>	65
<i>Hormocrinus</i>	66
<i>Asaphocrinus</i>	66
<i>Pycnosaccus</i>	67
Family SAGENOCRINIDAE	68
<i>Sagenocrinus</i>	68
Family ICHTHYOCRINIDAE	68
<i>Ichthyocrinus</i>	68
<i>Clidochirus</i>	69
Family TAXOCRINIDAE	69
<i>Protaxocrinus</i>	69
<i>Gnorimocrinus</i>	70
Order INADUNATA	71
LARVIFORMIA	71
<i>Pisocrinus</i>	72
<i>Triacrinus</i>	74
<i>Zophocrinus</i>	81
<i>Tiaracrinus</i>	81
<i>Mysticocrinus</i>	83
Family HETEROCRINIDAE	85
<i>Myelodactylus</i>	85
Family CALCEOCRINIDAE	88
Morphology and evolutionary sequence	88-103
Analysis of Genera	104
<i>Cremacrinus</i>	105
<i>Eucheiocrinus</i>	111
<i>Calceocrinus</i>	115
<i>Halysiacrinus</i>	121
Family PETALOCRINIDAE	126
<i>Petalocrinus</i>	126
<i>Crotalocrinus</i>	128
Family CYATHOCRINIDAE	131
<i>Thalamocrinus</i>	131
<i>Ampheristocrinus</i>	132
<i>Lecythiocrinus</i>	133
<i>Cyathocrinus</i>	133
<i>Parisocrinus</i>	135
<i>Gissocrinus</i>	135
<i>Botryocrinus</i>	135
INSERTAE SEDIS	137
<i>Parastephanocrinus</i>	139
<i>Stephanocrinus</i>	139
BLASTOIDEA AND CYSTIDEA	141
<i>Troostocrinus</i>	141
<i>Tetracystis</i>	141
<i>Stribalocystites</i>	142
<i>Lysocystites</i>	142
<i>Caryocrinus</i>	143
PERSONAL RECORD	144
BIBLIOGRAPHY OF AUTHOR'S WORKS	146
A. Scientific	146
1. In Collaboration	146
2. In Sole Authorship	146
B. Miscellaneous	149
INDEX	167
EXPLANATION OF PLATES	173

AMERICAN SILURIAN CRINOIDS

By FRANK SPRINGER

ASSOCIATE IN PALEONTOLOGY, U. S. NATIONAL MUSEUM

INTRODUCTION

Western Tennessee has been classic ground for American paleontologists ever since Gerard Troost, State Geologist of that state, brought to the 1849 meeting of the American Association for the Advancement of Science a series of fossil crinoids collected by him in Decatur and neighboring counties, and submitted through Professor Louis Agassiz¹ an imposing list of genera and species new to science, of which he had prepared a monograph with full descriptions and numerous illustrations. For want of funds to defray the expense, the monograph was not published during the author's lifetime, but remained buried until 1909, when it was issued by the United States National Museum as Bulletin 65, the MS., together with the collection, having been left by Troost to the Smithsonian Institution with a view to publication. By that time most of the new forms had been described by other authors, and the credit for the original discoveries announced in his list was lost to the one by whom it was so well deserved.

In 1847 came the eminent German paleontologist, C. Ferdinand Roemer, who became deeply interested in the prolific fauna of the Tennessee Silurian, and during a sojourn of several weeks made extensive collections in Decatur County. The results of his researches were published in 1860 in his well known monograph entitled "Die Silurische Fauna des Westlichen Tennessee," 4to, Breslau, Germany. Many new species were brought out, and the work has been regarded as a standard for the area of which it treats.

The Tennessee Area

The region under consideration lies along the Tennessee river and within its drainage. It abounds in natural exposures in the bluffs of the river and its tributaries, and in numerous glades where the fossiliferous limestone was exposed on the surface and disintegrated by erosion of the softer overlying clays and marls. Fossils in great numbers were weathered out upon those glades, which produced the greater part of the specimens obtained by the early collectors.

Following the two pioneers above mentioned, Prof. J. M. Safford as State Geologist made extensive studies of the Silurian formations, and his accounts

¹ Proc. Am. Assn. Adv. Sci. (2), VIII, read Aug. 1849, vol. 2, 1850, pp. 59-64.

of the stratigraphy were published in the *American Journal of Science* for 1861, and in his Report of 1869. He gave to the Niagaran rocks the collective name *Meniscus*, which he subdivided into two portions, a lower or variegated bed, and an upper, or sponge-bearing bed. He also made important collections, which are now in the Museum of Vanderbilt University at Nashville.

In consequence of the publication of the works of these early geologists, the Silurian area became an attractive field for collectors, and was visited from time to time by many ardent fossil hunters, among them Col. Sidney S. Lyon, Prof. A. H. Worthen, of Illinois, Dr. Carl Rominger, of Michigan, Mrs. J. M. Milligan, who resided in Decatur County for a considerable time, and my old associate, Charles Wachsmuth, who made two journeys along the Tennessee river during his sojourns in the South in the '80's. He was greatly impressed with the richness of the fossil beds, and the opportunities which they offered for intensive collecting, which the state of his health did not permit him to utilize in that then rather wild country—although even with his limited facilities he made important acquisitions.

In more recent times the geology of the western Tennessee region has been studied in detail by Dr. Aug. F. Foerste, whose valuable paper on "Silurian and Devonian Limestones of Western Tennessee" was published in 1903.¹ He recognized in Safford's variegated beds the Clinton, Osgood, Laurel and Waldron formations of the northern areas, and at the top of this division two new formations, the Lego and Dixon beds; while to the upper or sponge-bearing bed of the *Meniscus* he applied the new name *Brownsport*. Still later investigations were made by Prof. W. F. Pate, of Lebanon, Kentucky, and Dr. R. S. Bassler, of the United States National Museum. Their joint work published in 1908, entitled "The Late Niagaran Strata of West Tennessee,"² contains the latest and fullest information on the subject. It was based upon several seasons of study and collecting by Mr. Pate, the last two of them in my service, supplemented by a careful review of the stratigraphy by the two authors together in a field excursion made expressly for the purpose in 1907. Many sections of typical localities are given, and the paper should be consulted for a more detailed presentation of the facts than is attempted here.

Pate and Bassler found that the *Brownsport* and succeeding beds, which have furnished by far the greater part of the fossils from the glade region, are separable into four principal divisions, well marked faunally and lithologically for which they proposed the formation names *Beech River*, *Bob*, *Lobelville* and *Decatur*. The tabulation opposite of the several classifications as made by them will be useful in tracing the horizons as mentioned in the works of different authors.

¹ *Journal of Geology*, 11, pp. 554-715.

² *Proc. U. S. Nat. Mus.*, vol. 34, pp. 407-432.

The collections made by Wachsmuth in the glades of Decatur County, and his observations on the mode of occurrence of the fossils, gave me a very strong impression of the possibilities of that region for the paleontologist. In 1905 I purchased the collection of Professor Pate, containing among other material considerable obtained by him from the glades of Decatur and Perry counties. From him I secured additional authentic information regarding the local conditions and prospects for further collecting, on the basis of which I arranged with

Niagaran Strata of West Tennessee

	SAFFORD, 1869.	FOERSTE, 1903.	PATE AND BASSLER, 1908.
Niagaran	Meniscus		Decatur
			Lobelville { Coral zone Bryozoan zone
		Gant	Brownsport { Conchidium zone Dyctyonella zone Uncinulus zone
		Brownsport	
			Beech River { Eucalyptocr. zone Troostocr. zone Coccoocr. zone
		Dixon	Dixon
		Lego	Lego
		Waldron	Waldron
		Laurel	Laurel
	Variegated bed	Osgood	Osgood
		Clinton	Clinton

him to make a systematic campaign in that area the following season. All the crinoids that had been found up to that time had been in the form of specimens weathered out upon the glades, and consisted mainly of the loose calices with which the arms or finer structures were rarely preserved. If the beds from which they came could be located in place, there would be a possibility, in accordance with our long experience in the Burlington limestone, of finding some colonies with the crinoids imbedded in the matrix as originally deposited. Accordingly, Mr. Pate, at the outset of his excursion in the summer of 1906, was instructed to trace the crinoids by the fragments to their layer, and then dig.

The first result of this method was to locate along the bluffs of Beech River in Decatur County an exposure of shales and shaly limestone some 60 feet in thickness, containing the beds from the erosion of which the most productive of the glades were derived. Preliminary tests showed that some layers of this contained crinoids in place in good preservation, and thereupon a force was organized for extensive quarrying, with proper blasting equipment, and the work of uncovering the fossiliferous layers was carried on for the remainder of the summer under Mr. Pate's energetic direction. It was resumed the following summer, during which by persistent quarrying and blasting the favorable exposures were thoroughly excavated, and the accessible crinoid-bearing strata exhausted. These were chiefly in the upper shales, now known as the *Eucalyptocrinus* zone. Some smaller exposures in the vicinity produced the name-giving genera of the *Troostocrinus* and *Coccoocrinus* zones. Several years later Frederick Braun was sent for a final inspection of the Tennessee river area, especially the weathered quarry dumps, which yielded some excellent additional specimens.

The occurrence of the crinoids as disclosed by these quarrying operations was rather irregular, and much material was moved that proved to be barren. But in the aggregate the work of the two seasons was eminently successful, yielding a total of ninety-five species of crinoids, of which fifty-nine were new, distributed among thirty-three genera, besides a blastoid and a few cystids. The crinoids from the shales were usually silicified, and for the most part in excellent preservation. The matrix was readily removed by preparation, so that many of them when ready for study had the crown freely exposed on one or all sides. In some cases considerable colonies were found; for example, the hitherto rare "*Coccoocrinus*" *bacca*, of which about a thousand calices were recovered from a bed of shaly limestone of limited area. Taking the collection as a whole, it may be said without exaggeration that more finely preserved crinoids were secured by Mr. Pate's two summers' campaign than had been found by all the collectors in the Tennessee Silurian during the sixty years preceding.

Many rare forms, hitherto known only from isolated or fragmentary specimens, occurred in numbers sufficient for a thorough elucidation of their characters, and an even greater number were entirely new. This was conspicuously true for the *Flexibilia*, the *Calceocrinidae* and several genera of *Camerata*, of which there were forms and specimens such as had never been seen before. Details of these, however, will be better shown by the figures upon the plates.

Upon the broader question of the parallelism of the crinoidal fauna with that of the north European Silurian, the collection furnishes a wealth of information confirmatory of the existence of a migrational connection between the two areas. In several rare and specialized genera, as well as in some of abundant occurrence, species are found which except for difference in matrix can scarcely be distinguished from the English or Swedish forms. The general facies of the fauna proclaims a close relationship, and this impression is reinforced by the similarity of many of the other fossils.

The prominent exposures thus worked belong to the Beech River formation of the Brownsport group as designated by Pate and Bassler in their paper already mentioned, which they further subdivided into three zones called in ascending order *Coccocrinus* zone, *Troostocrinus* zone and *Eucalyptocrinus* zone. Of these the last, or upper, zone was by far the most important, the greater part of the crinoids having come from it. The other two are chiefly characterized

General Section Tennessee, etc. Niagaran

TENNESSEE					OTHER AREAS		EUROPE	
Silurian	Niagaran	Devonian Helderbergian		Linden				
				Decatur limest.				
		Brownsport Beech River	Lobelville, coral zone		Louisville ls. Kentucky, Ind.			
			Bob, limest. and shale					
			Eucalyptocrinus zone, shale			Bainbridge ls., Mo.		
				Troostocrinus zone, shale and limest.				
				Coccocrinus zone, shaly limest.			Cedarville dol. Ohio	
		Dixon, shale and red limest.		Racine dol. Illinois				
		Lego, limest.		Hopkinton dol. Iowa				
		Waldron, shale		Indiana				
		Laurel, massive limest.		Indiana				
		Osgood, shale and limest.		Rochester shale New York				
				Wenlockian				

by the forms from which their names are given. Above the shales is a considerable thickness of the Decatur limestone, which yielded only a few specimens here, but at a locality along the bluffs of the Tennessee river some remarkably distinct forms were obtained. At other places in the immediate vicinity the lower Niagaran formations occur, from Osgood to Dixon, so that the principal members are represented in their proper sequence within a short distance.

To facilitate reference to the horizons mentioned in the descriptions, I give above the general section of the Silurian formations of Tennessee as determined by Pate and Bassler, together with the equivalents which may enter into the discussions.

The Indiana Area

Next to Tennessee the most important western Silurian area is in Indiana, which is the typical region for the three lower Niagaran formations. The first important development there was made by James Hall and under his direction at Waldron, in the shale beds which bear its name. Among a wealth of other fossils these produced a remarkable crinoidal fauna, containing numerous new forms which were described by Hall in several publications from 1869 to 1882.

Underlying the Waldron, at the town of St. Paul in the same vicinity, is an immense exposure of the Laurel formation, a massive limestone of considerable thickness, which has yielded a crinoid fauna of a facies markedly different from those of the other Silurian members. An excellent collection from this area was made by the Doctors Howard, which was acquired by the Geological Survey of Indiana and formed the basis of descriptions by S. A. Miller in the 17th and 18th Annual Reports. Dr. C. C. Washburn also made an extensive collection from which a choice selection of crinoids passed into my hands, the main collection being afterwards acquired by the University of Chicago. Good crinoid material was obtained at St. Paul, as well as in the Wabash area to the north and the Osgood beds to the south, by A. C. Benedict, and in the Waldron shale at Harts-ville by Dr. Moses Elrod, both of which were subsequently acquired by me, as was also the large collection of Mr. John F. Hammell, of Madison, Indiana, which contained important material from all these formations.

The northern area of Wabash, Grant and Madison counties presents differences in the crinoid content from the Laurel which suggest a connection, in part at least, with the Racine dolomite of the Chicago area. The most characteristic species of the St. Paul rocks are not found there, although some of the northern species are thought to occur at St. Paul. The geology of the Indiana Niagaran has been carefully studied by Dr. Foerste, and his papers on the subject will be found fully instructive.¹

Being greatly impressed with the possibilities of the St. Paul exposures of the Laurel formation, and the striking character of its fauna as indicated by the earlier collections, I made some special efforts from time to time to secure a more complete representation of its contents. The crinoids were chiefly contained in the softer interbedded layers between the beds of massive limestone which were extensively quarried for commercial purposes. These layers constituted the débris of the quarries left in scattered dumps, in which the specimens became exposed by weathering from year to year, affording the main resource of the collectors. No prolific colonies were uncovered, and no great quantity of specimens was observed, but the preservation was excellent, and new or rare forms were always to be expected. Careful overhauling of this material was made under my direction during several years by Charles S. Beachler in 1887,

¹ 21st Ann. Rep. Dep. Geol. Indiana, 1896, pp. 213-288; 22d *ibid.*, 1897, pp. 195-255.

Frederick Braun in 1911, and, most important of all, during the three seasons of 1915, 1916, and 1917 by my former assistant, the late Dr. Herrick E. Wilson, who had before that made some collections there for the University of Chicago. Local conditions were favorable during his campaigns, which were fairly successful, and among his acquisitions were some of the important types described in this work.

The New York Area

In the eastern Silurian area the classic locality is Lockport, New York. Extensive exposures of the Rochester shale yielded a rich and varied fauna, among which were many new forms of crinoids which formed the types of genera and species described by Hall in the earlier volumes of the *Paleontology of New York*. The pioneer collections made by Professor Hall and Colonel Jewett ultimately passed into the possession of the American Museum of Natural History at New York, and the State Museum at Albany, where I have been afforded abundant opportunity for their study, as well as the types of Hall's numerous species from Waldron. Another fine collection from the same formation at Grimsby, Ontario, was assembled by the eminent financier, the late Sir Edmund Walker, which was also placed at my disposal, and is now in the University of Toronto.

The most intimate and protracted study of the Lockport beds, shales and underlying limestones, was made by the resident physician, Dr. E. N. S. Ringueberg. During a residence of more than thirty years he accumulated a large collection which formed the basis of several important publications by him from 1882 to 1890, in which a number of new genera and species were described.¹ The crinoids and cystids, including the types of Ringueberg's species, were acquired by me, and form an important part of the material studied. In addition to this, I have the fruits of three campaigns in the years 1910, 1912 and 1914 at Lockport by the veteran collector, Frederick Braun, in my service. In the course of an intensive study of the shales he carried on large quarrying operations where the conditions were found favorable. Some of his acquisitions were of the greatest importance, notably among the *Flexibilia* and the aberrant genus *Myelodactylus*, as fully described in my monograph of the former group of 1920, and in the paper on *Unusual Forms of Fossil Crinoids*, of 1926.

While the Lockport fauna does not fall strictly within the limits of this work, some consideration of it is essential for purposes of comparison, as the formation is to be correlated with the Osgood of the Indiana area.

¹ Two New Species of Crinoids from Shale of the Niagara Group. *Jour. Cin. Soc. Nat. Hist.*, Oct., 1882, p. 119; New Fossils from the Niagara Period. *Proc. Acad. Nat. Sci. Phil.*, 1884, p. 144; New Genera and Species of Fossils from the Niagara Shales. *Bull. Buff. Soc. Nat. Sci.*, 1886, p. 5; Some New Species of Fossils from the Niagara Shales. *Proc. Acad. Nat. Sci. Phil.*, 1888, p. 131; The Niagara Shales of Western New York. *Am. Geol.*, May, 1888, p. 263; The Calceocrinidae: A Revision of the Family, and Description of New Species. *Ann. New York Acad. Sci.*, 1889, p. 388; The Crinoidea of the Lower Niagara Limestone at Lockport. *Ann. N. Y. Acad. Sci.*, July, 1890, p. 301.

The Chicago Area

Another important Silurian area to be considered by way of comparison is that of the Racine dolomite at Chicago, which has produced a considerable crinoidal fauna, heretofore fully worked up and described by Stuart Weller¹ and A. W. Slocum.² In this formation the calcareous test of the crinoids has been largely dissolved by chemical action, so that for the most part only the internal casts are preserved. For this reason close comparison of characters is usually difficult. But the general facies of the fauna is very important, differing in many respects from those of southern Indiana and Tennessee, and paralleling that of the European Silurian in the presence of the highly specialized genus *Crotalocrinus*. In this it is reinforced by the occurrence in the equivalent Hopkinton dolomite of Iowa of the equally specialized *Petalocrinus*, also of Gotland.

The Foreign Silurian

In view of the close parallelism between the American Silurian and that of northern Europe before alluded to, it became desirable to secure an adequate representation of the crinoids from the two principal localities, Dudley in England, and the island of Gotland, Sweden. I had the advantage during a sojourn in Europe in 1887 of personal examination and study of the great Silurian collections in the British Museum, and in the Dudley Museum, and also of several private collections at Dudley during a visit to that field, especially those of Mr. Charles Holcroft, now in the Birmingham Museum, of Mr. John Gray, since acquired by the British Museum, and of Mr. William Madely. Through the good offices of Messrs. Holcroft and Madely, and from other local collectors, I was able to secure considerable useful material from that famous locality, which has been increased by purchases and exchanges since. From Gotland the more common species were obtained from time to time from dealers, and rarer ones from the Riks Museum at Stockholm by exchange, thanks to the liberality of my good friend Prof. G. Lindström. In more recent years I secured the services of local collectors on the island, first of Prof. G. Klinteberg, and afterwards of Mr. A. Florin, an experienced collector, who knew the geology of the island thoroughly, whom I engaged for a season's operations with blasting equipment. This resulted in some valuable acquisitions in the way of rare, and even new, forms, which have aided materially in the comparative study of the faunas.

All of the material thus accumulated forms part of my personal collection in the United States National Museum, where it will remain in perpetuity.

The total number of species herein illustrated and discussed is 198 belonging to 63 genera; of these 74 species are new and 12 undetermined. They are distributed as follows: Camerata, total 103, new 43, undetermined 12; Flexibilia, total 18, new 3; Inadunata, total 77, new 28.

¹ Bull. Chicago Acad. Sci., 4, pt. 1, 1900.

² Field Columbian Museum, 2, Geol. Ser., No. 10, 1908

Origin of the Work

The idea of a systematic work upon the American Silurian crinoids was first suggested by the account of the wonderful crinoidal fauna of Gotland by Angelin in his *Iconographia Crinoideorum*, and afterward by Bather in the *Crinoides* of Gotland, and by my observations upon the English collections from the Dudley area above mentioned. The collections made by Wachsmuth in Tennessee and Beachler in Indiana indicated a certain correspondence between the Silurian crinoidal faunas of the two continents which offered an inviting subject for investigation, and it was probably this consideration which induced the beginning of efforts toward the assembling of material with reference to it. It was not until the discovery of the rich collections made by Pate in 1906-7 that the plans took definite shape with the starting of drawings by Mr. Chapman, concurrent with the work upon the *Flexibilia*. These were prosecuted from time to time in a desultory way, as the demands of business and of other work permitted, and were continued as additional material was acquired. Preliminary studies and notes were made in the same irregular way, but it was only in 1923 that I found myself in a position to assemble the drawings, and begin the final preparation of the text.

At this point the work was suspended by a protracted physical breakdown incident to an impaired heart, from which a slow recovery has enabled me to resume it at intervals. That I have been able to bring the work to a conclusion is a result that was not expected; and for which I am profoundly thankful.

In conformity with the preparatory field-work, this treatise is chiefly limited to the Silurian faunas of Tennessee and Indiana. The crinoids of the Lockport field have been thoroughly treated by Hall and Ringueberg, and a few of them since by myself. Weller's and Slocum's works upon the Chicago crinoids cover that field exhaustively. Therefore, except for incidental reference to a few forms for special reasons, it has not been deemed necessary to include those of either field. For a similar reason I am omitting most of the Waldron species, which were so fully described by Hall.

The effort is made to reduce the treatment of species to a simple presentation of the essential facts, avoiding so far as possible the formal recital of characters which often becomes irksome. In some instances, where a form or group has been the subject of extended discussion hitherto, or where the acquisition of new material has thrown important light upon forms or questions before obscure, full latitude is employed. This will account for some differences in method that may be noticeable.

In the descriptions it has not been thought necessary to include tables of dimensions. They are usually only a pedantic incumbrance, inasmuch as the same information is more conveniently furnished by the drawings, which are

always made to scale upon a photographic basis, and are of natural size except as indicated in the legend.

While the chief purpose of this treatise is to bring to light the new genera and species which have been discovered within the areas hereinbefore indicated, it is also hoped to increase our knowledge of forms previously known where new information has been gained by later discoveries. But where such species have already been adequately illustrated and described, they will not be included here.

As to synonymy, Dr. Bassler's Bibliographic Index of Ordovician and Silurian Fossils, published by the United States National Museum as Bulletin 94, in 1915, brings references to all essential literature relative to American genera and species down to that date in the greatest detail and most convenient form. The publication has been widely distributed, and is available in every scientific library and in the hands of all working paleontologists. It is therefore unnecessary to encumber these pages with the usual full citations of authors previous to that date, inasmuch as a simple reference to that work will afford all the needful information. I am thus at liberty to limit my references covering dates previous to 1915 to such of the treatises as may for the sake of clarity or convenience be deemed of special importance in the history of the form under consideration.

In the notation of Horizon and Locality in connection with species, I am avoiding the needless repetition of the term "Silurian," inasmuch as all species herein described, unless otherwise stated, are from that horizon, and reference to the respective formations is usually quite sufficient to identify the geological position.

The material upon which this work is based is derived from the following sources, consisting of collections either made for, or acquired by, myself, and now in the United States National Museum, and those belonging to other owners which I have been privileged to study:

Author's Collections Used

ROCHESTER SHALE; Lockport, New York

By Frederick Braun, 1910, 1912, 1914. From Dr. E. N. S. Ringueberg, collection, Lockport.

OSGOOD, LAUREL AND WALDRON; Indiana and Tennessee

By Charles S. Beachler, 1887; St. Paul and Waldron. Braun, 1911; at St. Paul. Bassler and Braun, at Newsom, Tennessee. Herrick E. Wilson, 1915, 1916, 1917; St. Paul, Indiana.

From Dr. Moses Elrod, collection; Hartsville, Indiana. Dr. C. C. Washburn; Waldron, Indiana. Mr. John F. Hammell, collection; Madison, Indiana, including that of A. C. Benedict; St. Paul and other southern localities and northern area.

LATE NIAGARAN (BROWNSPORT) ; Tennessee

By Charles Wachsmuth, 1885 to 1888. W. F. Pate, 1906, 1907. Braun, 1911.

From Col. S. S. Lyon, collection, 1859 to 1860. Mrs. J. M. Milligan, collection; Jacksonville, Illinois. Pate, collection; before 1906.

OTHER COLLECTIONS STUDIED

American Museum Natural History, New York, Hall Collection; Lockport. State Museum, New York, Albany, Hall and Jewett collections; Lockport and Waldron. Sir Edmund Walker, Toronto; Grimsby, Ontario. Walker Museum, University of Chicago; Chicago area and St. Paul. University of Illinois, Champaign; Worthen collection. Vanderbilt University, Nashville; Safford collection. United States National Museum: Containing Troost collection, Tennessee; Rominger collection, Tennessee and Michigan; Mr. Ulrich's collection of Ordovician from Minnesota.

British Museum; London. Dudley Museum; Dudley, England.

All of these last mentioned collections were most generously placed at my disposal for study, for which I am deeply indebted to the owners and authorities in charge of them.

I am also under renewed obligation to Hon. Charles D. Walcott, Secretary of the Smithsonian Institution; to Dr. Geo. P. Merrill, head of the Division of Geology in the National Museum, for their continued interest in my researches; to Dr. R. S. Bassler, Curator of Paleontology, for his unflagging assistance during their prosecution; and to Miss Margaret W. Moodey, Registrar in the Division of Geology, who has direct charge of my collection and library, and has administered my office in the National Museum during my protracted disability, when I was endeavoring to complete the later stages of this work through the medium of correspondence. This involved numerous commissions in the way of selecting and sending specimens, books and notes, looking up references, and attending to the endless details incident to such investigations. All of these were executed by her, under advice of Dr. Bassler when needed, with a zeal and intelligent interest that met, and often anticipated, every requirement.

The drawings for the plates have been in the course of execution during a period of more than fifteen years, by Mr. Kenneth M. Chapman, of Santa Fé, New Mexico, the artist of many of my former works. They are based upon photographs made by himself at first, and in the later stages by Dr. Bassler and Dr. C. E. Resser. To his keen observation I owe their accuracy of details, and as works of art the drawings speak for themselves.

Except as otherwise stated, the originals of all the drawings, including the types of the new species herein described, are in the author's collection in the United States National Museum.

Order CAMERATA Wachsmuth and Springer

CRINOIDEA IN WHICH THE LOWER BRACHIALS TAKE PART IN THE DORSAL CUP. ALL PLATES OF CALYX UNITED BY CLOSE SUTURE AND IMMOVABLE. MOUTH AND FOOD-GROOVES COMPLETELY COVERED, THE COVERING PIECES OF THE LATTER FREQUENTLY INCORPORATED IN THE TEGMEN. ANAL OPENING EXCENTRIC OF SUBCENTRAL, FREQUENTLY AT THE END OF A TUBE. ARMS UNISERIAL OR BISERIAL, AND PINNULATE.

The crinoids of this order constitute more than half of those herein treated, embracing 31 genera of which 7 are new, and 43 new species out of a total of 103. Important new light is shed upon some forms heretofore not well understood such as *Gazacrinus*, *Lampterocrinus* and *Lyonicrinus*. The most prolific of the genera is *Marsipocrinus*, hitherto accounted rare, which now appears from the Tennessee area to the number of 10 species, some of them in a preservation and with a wealth of individuals before unknown.

Family DIMEROCRINIDAE Bather

This family, embracing substantially all dicyclic Camerata with truncate post. B in which the RR are in contact except at the anal side, while it has an extreme range from Ordovician to Devonian, is essentially a Silurian form, of which we have four well marked genera in the present collection.

DIMEROCRINUS Phillips

Plate 1

Dimerocrinites Phillips in Murchison's Sil. Syst., 1839, p. 674.

Dimerocrinus D'Orbigny, Prodr. Pal., 1, 1850, p. 46.—Wachsmuth and Springer, Rev. Pal., 2, 1881, pp. 184, 197.—Bather, Treatise on Zool. (Lankester), 3, 1900, p. 198.—Zittel-Eastman, Textb. Pal., 2d Ed., 1913, p. 187.—Jaekel, Phylogenie und System, 1918, p. 41.—Bassler, Bibliogr. Index, Bull. 92, U. S. Nat. Mus., 1915, p. 438.

Thysanocrinus Hall, Pal. New York, 2, 1852, pp. 188, 355.—Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 190.

Glyptaster Hall, Pal. New York, 2, 1852, p. 187; 28th Rep. New York St. Mus., 1879, p. 131.—Wachsmuth and Springer, Rev. Pal., 2, 1881, p. 193.

Eucrinus Angelin, in part, Icon. Crin. Suecc., 1878, p. 198.

Calyx rather elongate; IBB 5; post. B truncate; first anal plate in line with RR, followed by 3 in second range; iBr in several ranges; anus without a tube; arms biserial, simple, 2 or 4 to the ray, directed upwards; column round.

Genotype. *Dimerocrinus decadactylus* Phillips.

Distribution. Silurian; England, Gotland, America.

This genus is well represented in the English Silurian under Phillips' type from Dudley, in Gotland by several species under the name *Eucrinus*, and in America by a series of species under *Thysanocrinus* and *Glyptaster*, to which must now be added important new forms from the Tennessee collections. In the monograph of Crinoidea Camerata of 1897 priority was given to the name *Thysanocrinus*, a course which has since been modified in favor of the perfectly recognizable type described by Phillips.

Dimerocrinus planus new species

Plate 1, figs. 1-7

A finely preserved species, with elongate conical calyx, smooth surface, and very stout arms 2 to the ray; the interbrachial system is well developed in several ranges, by which the brachials are incorporated to the third or fourth IIBr. IBB small, rarely visible from a side view. Anus directly through the tegmen. Column long, composed of uniformly short columnals, averaging about .2 mm.

This is a conspicuous fossil in the Tennessee Silurian, of which besides the seven specimens figured there are a number of others showing similar characters. In having only ten arms it is closer to the Dudley species than to those from Gotland.

Horizon and locality. Beech River formation, Brownsport group, Niagaran; Decatur County, Tennessee.

As collections are made, it is not always possible to give the subdivision from which the species is derived. For the Beech River formation, by far the greater part of the specimens are from the upper, *Eucalyptocrinus*, zone.

Dimerocrinus milliganae (Miller and Gurley)

Plate 1, fig. 8

Glyptaster milliganae Miller and Gurley, Bull. 10, Illinois St. Mus., 1896, p. 87, pl. 5, figs. 7-9.

I am figuring the type of this Tennessee species, formerly in the collection of Mrs. J. M. Milligan. Only the calyx is preserved, but it is well differentiated by the prominent ridges along the basal and radial series, more angular contour generally, and larger IBB.

Horizon and locality, same as last.

Dimerocrinus nodobasis new species

Plate 1, figs. 9, 10

Similar to *D. planus* but differing in the presence of sharp angular cusps upon the basals; the IBB are extremely small, entirely concealed by the top columnal.

Horizon and locality, same as last.

Dimerocrinus inornatus (Hall)

Plate 1, figs. 11, 12

Glyptaster inornatus Hall, Trans. Alb. Inst., 4, 1863, p. 205; 28th Rep. New York St. Mus., 1879, p. 134, pl. 14, figs. 1-6.—Bassler, Bibliogr. Index, 1915, p. 439.

A massive species with strongly nodose basals, producing a stellate base. From the Waldron formation, Hartsville, Indiana. *D. (Glyptaster) occidentalis* (Hall), 28th Rep. New York St. Mus., 1879, p. 133, pl. 13, figs. 7-11, a much more delicate species with sharp surface sculpturing, is from the same formation at Waldron.

To the foregoing American species may be added *Rhodocrinus halli* Lyon, from the Louisville limestone, a doubtful species from the Waldron, three species from the Racine dolomite, five from the Rochester shale, and one from the Helderbergian—all as listed in Bassler's bibliography.

The following five species of Angelin from Gotland, *Eucrinus interradians*, *ornatus*, *laevis*, *quingangularis* and *speciosus*, with 4 arms to the ray, are referred to this genus, and probably two or more of them are synonyms.

Eudimerocrinus multibrachiatus new genus and species

Plate I, figs. 13, 13a, b, 14

The difference in arm structure between this and all species referred to *Dimerocrinus* is so great that to be consistent with the practice followed in other cases, it seems imperative to propose a new genus for its reception. While in those forms the arms are always simple, not branching beyond the calyx, here they undergo several bifurcations at wide angles up to as many as three, giving 6 or 8 final branches. These are all stout, and biserial both above and below the bifurcations—a type of arm highly distinctive of several genera, such as *Abacocrinus*, *Clonocrinus*, *Megistocrinus*, etc. In the composition of the calyx there is no difference, and with specimens consisting of the calyx alone it would be impossible to point out any character for distinction more than specific. But we have two unusually fine specimens, with the arms almost completely preserved to the numerous bifurcations, and both entirely freed from the matrix, so that the characters of the calyx are fully disclosed on all sides. The surface structure is remarkably clear and distinct, with strong sculpturing, many of the plates sharply projecting and almost spinose.

If in this species the interbranchials extended down to a connection with the basals, it might fairly go with *Eucrinus venustus* and *E. minor* of Angelin, for which Wachsmuth and Springer established the genus *Anthemocrinus*¹ under the Rhodocrinidae.

Horizon and locality. Beech River formation, Brownsport group, Niagaran; Decatur County, Tennessee.

CYPHOCRINUS S. A. Miller

Plate I

Cyphocrinus Miller, 18th Ann. Rep. Dep. Geol. Indiana, 1894, p. 304 (adv. sheets 1892).—Bather, Treatise on Zool., 3, 1900, p. 199.—Weller, Bull. Chicago Acad. Sci., 4, pt. 1, p. 75 (synonymy discussed).—Zittel-Eastman, Textb. Pal., 2d Ed., 1913, p. 187.—Bassler, Bibliogr. Index, 1915, p. 340.
Hyptiocrinus Wachsmuth and Springer, Am. Geol., 10, 1892, p. 138; N. A. Crin. Cam., 1897, p. 200.

Calyx constructed as in *Dimerocrinus*, but low, wide, and recurving from a narrow obconical base to a spreading, inverted bowl, with arms directed downwards; axillary and central tegmen plates spiniferous.

Genotype. *Cyphocrinus gorbyi* S. A. Miller.

Distribution. Silurian; not known outside of America.

¹ Rev. Pal., 2, 1881, p. 208. See also Jaekel, Phylogenie und System, 1918, p. 40, where he claims that *E. minor* may have only 4 BB.

Cyphocrinus gorbyi S. A. Miller*Plate 1, figs. 15-18*

Cyphocrinus gorbyi Miller, 18th Ann. Rep. Indiana Dep. Geol., 1894 (adv. sheets, 1892, p. 51), p. 305, pl. 7, figs. 14-16.—*Hyptiocrinus typus* Wachsmuth and Springer, Am. Geol., 10, 1892, p. 138; N. A. Crin. Cam., 1897, p. 201, pl. 19, figs. 6 a-c.

This genus, while contrasting in the hemispheric and drooping contour widely with the preceding, clearly exhibits the essential characters of the family in the construction of the calyx. The arms have not been found beyond slight remnants, from which it is evident that they were biserial, stout and unbranched; and from the appearance of the extremely elongate spinous ambulacra upon the tegmen surmounting the rays, and the direction of the arm-openings, it might be inferred that they were of the recumbent type, with the pinnules directed outward.

Four specimens of the type species, which is quite abundant at the type locality, are figured in order to show some details more fully than appeared in the original description. This applies especially to the interbrachial openings at the margin of the tegmen for fixed pinnules to the number of 4 or 6 between the rays and their divisions. These openings, formerly interpreted as "respiratory pores," are now traced in several Camerate genera with unbranched arms to a connection with the arm-groove.¹

Weller has described a species, *C. chicagoensis*, from the Racine dolomite of the Chicago area, but otherwise the form has only been found in the Laurel limestone at St. Paul, Indiana.

This and the next following form were independently described by Wachsmuth and Springer and S. A. Miller in 1892. The Wachsmuth and Springer names were published in the American Geologist for September, 1892, but owing to some delay in printing, that number of the magazine was not actually issued until some weeks after its purported date. Meantime Mr. Miller issued advance sheets of the 18th Rep. Geol. Indiana containing his new names, which were specially distributed to his correspondents. The claim was made that this distribution antedated the delivery of the magazine to its subscribers. Dr. Weller in connection with his Chicago species considered the question in the light of such evidence as could be obtained, and decided in favor of the priority of Miller's names, which I have accepted.

Horizon and locality. Laurel limestone, St. Paul, Indiana.

GAZACRINUS S. A. Miller*Plate 2*

Gazacrinus Miller, 18th Rep. Geol. Dep. Indiana, 1894, p. 303 (adv. sheets, 1892, p. 49), N. A. Geol. Pal., 1st App., 1892, p. 679; *ibid.*, 2d App., 1894, p. 746.—Weller, Bull. Chicago Acad. Sci., 4, pt. 1, 1900, p. 78, fig. 38 (synonymy discussed).—Bather, Treatise on Zool., 3, 1900, p. 188, fig. 119.—Zittel-Eastman, Textb. Pal., 2d Ed. 1913, p. 187.—Jaekel, Phylogenie und System, 1918, pp. 41, 80.—Bassler, Bibliogr. Index, 1915, p. 545.

Idiocrinus Wachsmuth and Springer, Am. Geol., 10, 1892, p. 135; N. A. Crin. Cam., 1897, p. 202.

Base concave, IBB confined to the cavity; iBr in cup limited to a single large plate in each interradius connecting with tegmen, that of posterior side following the truncate B; arms biserial, simple, directed upwards. Anus without a tube. Tegmen composed of a few plates, crowned with a central pyramid.

¹ See Springer, *Scyphocrinus*, Smithsonian Inst., 1917, pp. 33-37.

Genotype. *Gazacrinus inornatus* S. A. Miller.

Distribution. Silurian to Devonian; America. Not recognized in the foreign Silurian.

The position and relations of this genus have been misunderstood. Bather placed it among the Flexibilia. Jaekel in his latest attempt at the systematic arrangement of the crinoids, *Philogenie und System*, 1918, does not fully agree with this reference, but in order to be on the safe side places it in each of his two most widely distinct grand divisions; first under his subclass Cladocrinoidea, equivalent to the Camerata, in a family next to the Dimerocrinidae (p. 41), and then in his other subclass, Pentacrinoidea (p. 80), in a family Gazacrinidae under the suborder Calpocrinites—an association of types about as incongruous as could possibly be imagined.

As originally described, partly through errors of definition and partly through limitation of material, the essential characters of the genus were not fully disclosed, it being known only by two species from a single locality, specimens of which were rare and imperfect. Later acquisitions have furnished much new information, giving us now a series of seven or more species ranging through several formations of the Silurian in different localities and up into the Devonian. These have added greatly to our knowledge of structures unsuspected before.

From them it now appears that the genus has ten heavy, biserial arms, directed upward and closely abutting. The tegmen is unlike that of others of the family, and is a distinct variation from that of the Camerata generally, being composed of a few large interambulacral plates, surmounted by a set of triangular interlocking plates thought to be orals which vary in different species, being suturally closed or even anchylosed in some to form a solid pyramid with angular vertical ridges forming grooves into which the arms may fit. The various forms of this structure are well shown on plate 2, and from these it may be seen that there is a strong analogy between this tegmen and that of the Calyptocrinidae, as will appear by comparison with the figures of *Eucalyptocrinus crassus* on plate 7, figs. 19, 20. This will furnish the desired line of connection with the Camerate type. See also Wachsmuth and Springer, N. A. Crin. Cam., 1897, pl. 81, figs. 12-15; and Hall, 28th Rep. New York St. Mus., 1879, pl. 19, figs. 1, 3, 4, 5.

The interpretation of the pyramid of plates that we call orals is involved in considerable doubt. They are buried under closely fitting arms, which would affect the regularity of their development, and there is manifest in some species a strong tendency to fusion, by which the relation of the component plates may be greatly modified.

The definition of the genus by Mr. Miller, and description of the type species, are erroneous as to the composition of the base in the definite statement that it has "no subradials"—that is, that it is a monocyclic crinoid, which is a serious and fundamental error. This was due to loose observation, and the lack of careful preparation of the specimens to bring out the infrabasals at the bottom of the cavity, where they are plainly to be seen in more than a dozen of my specimens. Wachsmuth and Springer's definition of *Idiocrinus*, based on the same species as Miller's under a different name, specified the 5 small IBB at the bottom of the cavity, and the fact is shown by numerous examples of this and other species. It is also stated by Miller that the arms are composed of a single series of flattened plates, which was pure guess-work, and contrary to the fact as now known.

Gazacrinus inornatus S. A. Miller

Plate 2, figs. 1-9

Gazacrinus inornatus Miller, 18th Rep. Indiana Dep. Geol., 1894 (adv. sheets, 1892, p. 49), p. 303, pl. 5, figs. 9, 10, 15-17.—*Idiocrinus elongatus* Wachsmuth and Springer, Am. Geol., 10, 1892, p. 136; N. A. Crin. Cam., 1897, p. 203, pl. 18, figs. 8 a-c.

Specimens small, average height to width at top of iBr 12 x 11 mm. Calyx elongate conical. Base narrow with BB higher than wide, curving abruptly into

a cavity forming an inverted cone with the small IBB at the bottom. RR very large. iBr and IIBr projecting over into the tegmen, but exteriorly the brachials pass insensibly into the large biserial arms, two to the ray, which abut closely both between the rays and their divisions. Ventral pyramid somewhat variable in shape, asymmetric, consisting of five similar plates with a smaller central one interposed; or the central plate may represent the posterior oral pushed in between the others by anal structures, with the posterior triangular plate flanked by two others connecting forming the border of the anal opening. These pyramid plates abut by their proximal edges upon the bent over processes formed by extensions of the IIBr and iBr plates. Surface smooth.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Gazacrinus magnus new species

Plate 2, figs. 10, 11, 12

This species, from the same horizon and locality as the preceding, is known only from the tegminal pyramid, which is of great size, and exhibits in a remarkable degree the sharp ridges—5 or 10 in number—forming the border of vertical grooves or compartments in which the arms may rest as in *Eucalyptocrinus*. There are 3 specimens. Fig. 12 has only 5 broad grooves, while fig. 10 has 10, with extremely sharp dividing ridges. This pyramid is formed by the sutural union of 5 elongate plates, as is clearly shown by the interior view of the same specimen in fig. 10a. This is further confirmed by fig. 11, which shows a single detached plate of a similar and larger pyramid. Here there is no question of a central plate as in the preceding species.

Horizon and locality, same as last.

Gazacrinus depressus new species

Plate 2, figs. 13, 14

A crushed specimen from St. Paul of different type from the preceding, rather resembling the Tennessee form.

Horizon and locality, same as last.

Gazacrinus ventricosus (Wachsmuth and Springer)

Plate 2, figs. 15-16

Idiocrinus ventricosus Wachsmuth and Springer, Am. Geol., 10, p. 137; N. A. Crin. Cam., 1897, p. 205, pl. 18, figs. 9 a-b.

Specimens very small, height to width at top of IIBr about 2 by 2.5 mm. Calyx low, subglobose, but with base excavate and IBB sunken in the cavity. iBr large, elongate, projecting inward where their extensions support the tegminal pyramid, which here consists usually of anchylosed plates by the fusion

of which the identity of the component plates is lost; in this condition the solidified pyramid is frequently found isolated, while the calyx of the species is extremely rare and the arms unknown. These detached pyramids are usually strongly marked with radiating ridges and grooves, as shown in figs. 16, 16a, which converge exteriorly into a fused globular mass at the center. From such examples the inference would be that in the course of the modification of the tegmen incident to compression by the closely packed arms the original function of the structure had been lost. Surface smooth with calyx plates moderately convex.

These isolated bodies when first observed were supposed by Miller and by Wachsmuth and Springer to represent the ventral structure of *Pisocrinus*, specimens of which were numerous in the same bed.

Horizon and locality, same as last.

Gazacrinus ramifer (Roemer)

Plate 2, figs. 17-21

Eucalyptocrinus ramifer Roemer, Sil. Fauna Westl. Tennessee, 1860, p. 51, pl. 4 a, b.—*Callicrinus ramifer* Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 358.

A large species; calyx wider than high, average about 10 by 12 mm. Base broad, excavate, the plates strongly marked by a stellate rim bordering the cavity, which is here wide and shallow. Other ridges passing from BB to RR divide the surface into prominent triangles. IBB fairly large. Arms long and heavy, directed upward, lying parallel but not so closely abutting as in the type species. The ventral pyramid is relatively very large, with only a few sharp ridges, as shown in fig. 19a.

It is an interesting fact that Roemer when describing this species, one of the rarest known to the early Tennessee collectors, should have been led by an indefinable resemblance in the calyx alone to refer it to *Eucalyptocrinus*, and that Wachsmuth and Springer, upon the evidence of a solitary specimen which they had obtained, while in much doubt as to its generic relations, recognized a possible affinity with the other Calyptocrinid, *Callicrinus*.

The shales of the Beech River formation have yielded some remarkably fine specimens of this species, throwing new light on the relations of the genus, free from the matrix and exposing arms and surface characters at all sides. The clue to the actual structure first afforded by this material was later confirmed by the finding of the equally instructive specimens of the type species at St. Paul.

Horizon and locality. Beech River formation, Niagaran; Decatur and Wayne counties, Tennessee.

Gazacrinus milliganæ (Miller and Gurley)

Plate 2, figs. 22-24

Thysanocrinus milliganæ Miller and Gurley, Bull. 8, Illinois St. Mus., 1896, p. 51, pl. 3, figs. 23-25.

Similar to the preceding, and perhaps only a variety of it, represented by imperfect specimens. I am figuring Miller and Gurley's type specimen and a

couple of others that may perhaps go with it. The chief difference observable is that here the base is less deeply excavate, and the BB, instead of being raised into a stellate ridge, are sharply nodose, forming a well defined pentagon.

Horizon and locality. Beech River formation, Niagaran; from the glades in Decatur County, Tennessee.

Gazacrinus stellatus new species

Plate 2, figs. 25 a-c

This species is the only instance in which the genus has been recognized from beyond the limits of the Silurian. The unique specimen upon which it is based comes from the Linden formation of the Helderbergian, the lower beds of which in some Tennessee localities lie in very close relation to the Decatur limestone at the top of the Niagaran, where it almost seems as if there is an intermingling of typical fossils. The species, while in general similar in size and proportions to *G. ramifer*, is clearly distinct in detail. The basal cavity is sharply defined, the edge forming an acutely pointed pentagon with straight sides. The BB and other calyx plates are covered with coarse pustules, which tend to coalesce below into lines forming raised triangles on the radials, but with more isolated rugose markings on the other plates.

Horizon and locality. Linden formation, Helderbergian, Lower Devonian; Hardin County, Tennessee.

Two species have been described by Weller from the Racine dolomite of the Chicago area, and one by Hall from the Rochester shale at Lockport as *Thysanocrinus* has been referred to the genus. For list and synonymy, see Bassler, Bibliogr. Index, 1915, p. 545.

LAMPTEROCRINUS Roemer

Plate 3

Lampteroocrinus Roemer, Sil. Fauna Westl. Tennessee, 1860, p. 40.—Wachsmuth and Springer, Rev. Pal. 2, 1881, pp. 184, 199; 3, 1885, p. 101; N. A. Crin. Cam., 1897, p. 207.—Bather, Treatise on Zool., 3, 1900, p. 199, fig. 124.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 187.—Jaekel, Phylogenie und System, 1918, p. 41. For further references see Bassler, Bibliogr. Index, 1915, p. 689.

Calyx subturbinate, similar to that of *Dimerocrinus* but asymmetrical owing to bulging at anal side; iBr few, anal interradius wider than the others; rays produced into 5 tubular extensions, bearing uniserial, pinnuliferous arms at each side, with more than one pinnule to a brachial; tegmen composed of numerous convex plates passing into a strong, subcentral tube; column sharply pentagonal.

Genotype. *Lampteroocrinus Tennesseensis* Roemer.

Distribution. Silurian; America.

In the Revision of the Palaeocrinoidea, pt. 2, p. 200, and in N. A. Crinoidea Camerata, p. 207, Wachsmuth and Springer described the brachial arrangement of this genus as consisting of lateral appendages, in which the brachials, from primibrachs up, roofed by covering pieces were extended into single rigid tubes, or free rays, giving off the arms laterally. This

was an inference by Wachsmuth of a structure which had not actually been seen, based upon analogy of the Batocrinid genus *Stegancrinus*. It was criticised by Bather in his review of the *Camerata* monograph, 6th notice, *Geol. Mag.*, March, 1899, p. 125, calling attention to the fact that our interpretation had not yet received confirmation.

This objection can now be met by the proved fact. *Lampterocrinus* is one of the best known and abundant crinoids in the Tennessee rocks, and from Roemer's time on the calices were to be found in collections generally. They are firm, strong fossils, with thick plates solidly cemented together, well adapted to resist weathering in the glades. From these rotund bodies the other parts are readily detached, so that until now no one has ever seen any vestige of the arm structure. By the fortunate occurrence of some specimens intact in the Beech River shales, protected by the matrix which could be removed by preparation, we are now enabled to exhibit the complete crown with all the appendages in their natural position. In the remarkable series of specimens here studied, there are two with the tubular rays and side arms and the anal tube in place, and four others with parts of the radial extensions favorably located for inspection. There are five heavy tubes composed of the primibrachs and higher brachials curved like arm plates, with their covering plates, fringed with very long, slender arms given off alternately from each side, and bearing delicate pinnules—precisely the structure of *Stegancrinus sculptus* of the Burlington limestone. The side-arms are uniserial, and each ossicle bears two pinnules, one from each side.

While the type species remains as the chief representative of the genus, the new material has produced a well defined new one with a variant for a possible third.

Hall described a small species *L. parvus*, from the Waldron shale, and also one from the Racine dolomite of the Chicago area, from which three species are also described by Weller. Rowley described another from the Missouri Silurian. None of these throw any light on the structure. See Bassler's Index for list with references.

***Lampterocrinus tennesseensis* Roemer**

Plate 3, figs. 1-6

Lampterocrinus tennesseensis Roemer. *Sil. Fauna Westl. Tennessee*, 1860, p. 37, pl. 4, figs. 1 a, b.—Wachsmuth and Springer, *N. A. Crin. Cam.*, 1897, p. 209.—Bassler, *Bibliogr. Index*, 1915, p. 690.

This original species, so well known in all collections by weathered specimens from the glades, is strongly characterized by its elongate form, sharp stellate sculpture, and prominence of the anal interradius, which is larger than the others, and curves over into the tegmen where it passes into the strong sub-central tube, while the lower part is more or less swollen so as to produce a decided asymmetry. This anal "hump" is a most conspicuous character, by which the weathered calices are usually recognized at a glance. The sculpture takes the form of highly raised ridges passing from center to center of the plates, outlining a variety of triangles and hexagonal figures. At the anal side these often connect in such a way that they form a strong median ridge running the full length of the interradius from the posterior basal to the anal tube.

The complete crown, free on both sides, was imbedded in a shaly limestone, where several instructive fragments of arms also occurred showing their mode of connection and relation to the tubular rays, and especially their bipinnulation, as in fig. 6. This peculiar structure, at one time considered to be an impossibility in crinoid morphology (*Crin.* Gotland, 1893, p. 36), is now known to exist in several genera. I figure on plate 7 a specimen of *Carpocrinus* from Gotland in which the two pinnules to a brachial are well developed; and the structure

was fully described in a Devonian genus in my paper on *Dolatocrinus*,¹ with 2, 3 and 4 pinnules to a brachial. Since then it has been further observed by Miss Goldring, Devonian Crinoids of New York, 1924, in the genera *Clarkeocrinus*, p. 180, and *Liparocrinus*, p. 397, with like numbers of pinnules to the brachial.

This leading species for the Tennessee Silurian was recognized by Troost long before Roemer's time, and fully described under the name *Balanocrinites* in his unpublished monograph.

Horizon and locality. Shaly limestone of Beech River formation, Niagaran; Decatur and adjoining counties, Tennessee.

***Lampterocrinus sculptus* new species**

Plate 3, figs. 7, 8

A variant of the preceding, from the same horizon and locality, with a lower and more spreading calyx, more delicate sculpture and more numerous ridges.

***Lampterocrinus roemeri* new species**

Plate 3, figs. 9-14

Differs from the type species in the much plainer character of the surface markings. The ridges are fewer and more prominent, and the rhomboidal areas enclosed by them are almost uniformly smooth, and not crossed by other ridges; thus the stellate appearance is largely wanting. The specimens all come from a soft shale at a little different level from the former species.

In addition to the small crown with arms and anal tube, fig. 10, there are 4 calices with parts of the radial appendages attached, as in fig. 9, and 20 other good specimens, in all of which the characters above mentioned are constant. Here also may be seen the sharply pentagonal stem characteristic of the genus, with concave sides, as shown in figs. 9 and 10. I have called this species *L. roemeri*, in honor of the founder of the genus.

Horizon and locality. Soft shale of the Beech River formation, Niagaran; Tuck's Mill, Decatur County, Tennessee.

————— genus and species indet.

Plate 3, figs. 15, 15a

This fragment, of which there is not enough of the calyx left to indicate its generic relations, is figured on account of the very unusual feature of a hexagonal stem, almost unknown among crinoids. Aside from its six-sided section, the stem in detail strongly resembles that of *Lampterocrinus*, and the specimen may be only a sport from that genus.

Horizon and locality, same as last.

————— genus and species indet.

Plate 1, figs. 19-22

Stem fragments of unknown crinoid from St. Paul, Indiana.

¹ Bull. 115, U. S. Nat. Mus., 1921, pp. 5, 8.

Family RHODOCRINIDAE Roemer

This dicyclic family, similar to the Dimerocrinidae except that the radials are separated all around, is introduced to receive a number of peculiar forms falling within its limits for which it has been necessary to propose new genera.

Paragazacrinus rotundus new genus and species

Plate 4, figs. 1-4

On account of the strong resemblance of the tegmental parts this small species was at first supposed to be a variety of *Gazacrinus*; but when it was found that in three good specimens the RR were separated all around by iBr truncating all 5 BB that interpretation became untenable, and there seemed no recourse except to place it as a new genus under the Rhodocrinidae. There are 5 IBB sunken in an indented pit, and 4 or 5 large iBr in the cup separating the rays up to the tegmen, which branch and become free on the first IIBr, giving 10 arms in all, which are not further known. They rest in grooves bounded by ridges forming a ventral pyramid which is surmounted by an elongate spine. This is supported by 5 tegmen plates, the exact relation of which is difficult to determine, but they are thought to be interambulacral in position. The three rotund calices are figured, together with a detached central spine, as to which the question remains open whether it may not belong to *Gazacrinus*, although it is believed that the spine is a separate element covering the pyramid. The two forms are from the same locality and formation, but whether from the same layer is not known.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Paulocrinus biturbinatus new genus and species

Plate 4, figs. 5, 5a-c

This genus and species are proposed for the reception of a solitary calyx, which both in general form and in cup structure is totally unlike any other known from the Silurian. It is large, robust, with truncate conical cup spreading to the arm-bases, and from there contracting to an opposite cone consisting of a many-plated tegmen passing into a large, subcentral anal tube. The calyx plates are thick and strongly elevated, those of the tegmen rather nodose, and those of the cup more or less acuminate, and ornamented by radiating ridges.

The most peculiar thing about this form is the base, forming the truncate end of the dorsal cone. There are 5 IBB surrounding an enormous lumen, and no two of them alike, 3 smaller and pentagonal on the posterior and left posterior sides; the other 2 hexagonal and truncated by BB are larger than the former 3 combined. A similar inequality pervades the BB but reversed in position as if compensatory, the two larger being the posterior and left posterior (see diagram, fig. 5c). Also the RR are more or less unequal, and asymmetric in

form, two of them connecting with the IBB. Arm openings 4 to the ray, directed horizontally from a projecting rim; the r. post. ray bifurcates on the first IBr, but by way of compensation this is followed by more than two IIBr; the arms not preserved.

All this points to an abnormal specimen, or sport—but of what? The form and general structure of the calyx are positive and well defined, and in all the collections from St. Paul and the Tennessee localities nothing has been found with which it can be compared. Therefore it will have to stand upon the above description until the normal form is shown by further discovery.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Wilsonicrinus discoideus new genus and species

Plate 4, figs. 6, 6a

This is another Rhodocrinid straggler from the St. Paul beds, known only by the single type specimen, and apparently without close relatives. The calyx is a broad, wheel-shaped disk, with smooth plates, slightly convex below and contracting above to a large subcentral anal tube. It has 4 IBB sunken into a small indented cavity, BB fairly equal in size, truncate above and connecting with single large iBr. Arm-openings 10, small and widely separated, issuing from a series of cuneate IIBr.

The outstanding feature of this genus is the fixed pinnules originating upon the longer face of the IIBr to the number of 4, 5 or 6 on each side in the form of more or less irregular pinnulars, which by contact with the IIBr and with each other are incorporated in the cup; they lead directly to interbrachial openings for free pinnules independent of the arms, emerging at the margin of the tegmen between the rays and their subdivisions. These are very conspicuous, corresponding in number to the incorporated pinnules. Thus the interbrachial areas, above and beyond the single large iBr plates which abut upon the truncated BB, are occupied by these fixed pinnules, which have every appearance of higher iBr. For a full discussion of the fixed pinnules in various genera and the interpretation of their openings at the margin of the tegmen, see my papers on *Scyphocrinus*¹ and *Dolatocrinus*.²

The type specimen was found by my former assistant, the late Dr. Herrick E. Wilson, who recognized its highly distinctive characters, and the generic name is proposed in his memory.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Emperocrinus indianensis Miller and Gurley

Plate 4, figs. 7, 7a

Emperocrinus indianensis Miller and Gurley. Bull. 6, Illinois St. Mus., 1895, p. 43, pl. 4, figs. 16, 17.—Bather, Treatise on Zool., 3, 1900, p. 202.—Bassler, Bibliogr. Index, 1915, p. 476.

Another rare form, of simpler construction than the preceding, which I am illustrating from a new specimen. Instead of being discoid, the calyx is pentagonal, and the iBr areas, instead of being filled to the margin with the incorporated pinnules, are depressed, contracted and limited to the single large

¹ Smithsonian Publication No. 2440, 1917, pp. 33-37, 40-46; pl. 9, figs. 5-6.

² Bull. 115, U. S. Nat. Mus., 1921, pp. 24, 25, pl. 7, figs. 2-5.

plate. There are a few openings for free interbrachial pinnules, apparently only one to each arm on the outer side of the ray, lying close under the arm-base. These structures were not brought out in the original description, and whereas the number of IBB is there said to be three, it is distinctly five in my specimen. Miller and Gurley treated the genus as belonging to the Flexibilia.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Rhodocrinidae genus and species indet.

Plate 4, figs. 8, 8a

A fragment with only part of the base, attached in a slanting position to a very large stem. There are 4 IBB at the bottom of an indented cavity as in several of the associated forms, and entirely covered by the stem. None of the others of this family here described have any part of the stem preserved, and it may be that, large as it seems, this is the normal type for the group. But the sloping attachment to the calyx is unusual. The BB are truncate for a connection with the succeeding iBr, separating the RR, so it probably belongs here.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Lyriocrinus melissa Hall

Lyriocrinus melissa, Hall, 28th Rep. New York St. Mus. Nat. Hist., 1879, p. 139, pl. 15, figs. 18-27; 11th Ann. Rep. Indiana Dep. Geol., 1882, p. 269, pl. 14, figs. 18-27, pl. 15, fig. 11.—Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 263, pl. 11, figs. 4a-f.—For complete synonymy, Bassler, Bibliogr. Index, Bull. 92, U. S. Nat. Mus., 1915, p. 774; also for the genus, and other species, p. 773.

This very typical Rhodocrinid species is one of the leading fossils in the Waldron formation, occurring abundantly at the type locality, Waldron, Indiana, and in the rich exposure of Waldron shale at Newsom, near Nashville, Tennessee; it has also been identified in the Racine dolomite of the Chicago area. Reference to the thorough discussion and illustration of the species by Hall and by Wachsmuth and Springer will afford all needful information, and it is accordingly omitted from consideration here.

Another fine species, *L. dactylus*, was described by Hall from the Rochester shale at Lockport, of which unusually perfect specimens from Grimsby, Ontario, are in the Sir Edmund Walker collection in the University of Toronto. I have an undescribed specimen from Dudley, indicating the presence of the genus in the English Silurian.

Family MELOCRINIDAE Zittel

Of this extensive monocyclic family, with the radials in contact all around, the collections here in hand contain eight genera, two of them foreigners now first recognized in this country, one which though known before has developed an unexpected wealth of species, and one new.

MACROSTYLOCRINUS

Plate 4

Macrostylocrinus Hall, Pal. New York, 2, 1852, p. 203.—Wachsmuth and Springer, Rev. Pal., 2, 1881, pp. 96, 102; N. A. Crin. Cam., 1897, p. 285.—Bather, Treatise on Zool., 3, 1900, p. 162.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 191.—Jaekel, Phylogenie und System, 1918, p. 36.—Bassler, Bibliogr. Index, Bull. 92, U. S. Nat. Mus., 1915, p. 782.

BB 3 unequal; anal interradius wider than the others above level of RR; arms 10, simple, biserial.

Genotype. *Macrostylocrinus ornatus* Hall.

Distribution. Silurian and Devonian; America, England.

This typically Silurian genus, but now known to occur in the Lower Devonian,¹ which has hitherto been accounted as one of the rare fossils, is well represented in the present collections, especially for illustrating the structure of the tegmen, heretofore unknown. Several species are figured, all but one previously described, and of these it is not deemed necessary to repeat the descriptions.

***Macrostylocrinus fasciatus* Hall**

Plate 4, figs. 9, 9a, b

Macrostylocrinus fasciatus Hall, 28th Rep. New York St. Mus., 1879, p. 130, pl. 13, figs. 5, 6.—Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 288, pl. 22, fig. 13.—Bassler, Bibliogr. Index, 1915, p. 782.

The specimen here figured has the fasciculated striae, rarely well preserved, perfectly distinct, and shows correctly the full dimensions of the calyx and the form and proportions of all the plates. The important feature, however, is that it has the tegmen intact, which has not been found before. Usually the upper part is broken away, so that the great elongation of the cup does not appear. The tegmen is composed of a great number of small smooth plates, traversed from the center to arm-bases by narrow, almost linear, ambulacra. The small anal opening lies at the extreme margin, where it projects between the two posterior rays, connecting with the median row of anal plates on the dorsal side. Radial series sharply defined by a median ridge. Total height of calyx 13 mm.

Horizon and locality. Waldron shale, Niagaran; Hartsville, Indiana.

¹ Springer, Unusual Forms of Fossil Crinoids, Proc. U. S. Nat. Mus., Vol. 67, 1926, p. 35.

Macrostylocrinus granulosus Hall*Plate 4, figs. 10-13*

Macrostylocrinus striatus var. *granulosus* Hall, 28th Rep. New York St. Mus., 1879, p. 129.—Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 289, pl. 22, figs. 15a, b.—Bassler, Bibliogr. Index, 1915, p. 782, 783.

There is here illustrated a form with turbinate calyx, expanding upward as most frequent in the genus, in which the tegmen is also beautifully preserved in two specimens. This is of a somewhat different type, in accord with the more spreading form of the calyx, in which the ambulacra tend to become wider, and the anal opening very much larger; it is situated in the depressed posterior interambulacrum between the margin and the center, and in one specimen is closed by a pyramid of small plates. The granulose surface, completely devoid of striae, together with the general form and proportions, seems to indicate the reference to this species. It is very small, as is best seen by comparing the drawings, which are mostly double size.

Horizon and locality. Waldron shale, Niagaran; Madison, Indiana, Newsom and perhaps Decatur County, Tennessee.

Macrostylocrinus laevis new species*Plate 4, figs. 14-19*

This name is suggested to designate a smooth form occurring rather plentifully in the Beech River formation, having relatively small RR and wide IBr, and in which we have the stout biserial arms partly preserved.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

Macrostylocrinus meeki (Lyon)*Plate 4, figs. 20-22*

Actinocrinus meeki Lyon, Proc. Acad. Nat. Sci. Phil., 1861, p. 411, pl. 4, figs. 4a, b.—*Macrostylocrinus meeki*, Wachsmuth and Springer, Rev. Pal., 2, p. 103; N. A. Crin. Cam., 1897, p. 290, pl. 22, figs. 16a, b.

A broader and more rotund calyx than usual in the genus, and notable for the sharp median ridges which follow the rays from radials into arm-bases; it may also be more or less striate.

Horizon and locality. Louisville limestone; Jefferson County, Kentucky, and Black River formation, Niagaran; Decatur County, Tennessee.

(?) Macrostylocrinus pustulosus new species*Plate 4, figs. 25, 25a*

It is not clear that this unique specimen belongs to the genus. The general contour and pustulose ornament would seem to be exceptional, while the arrangement of the anal side is consistent with the reference. In the large side of the RR it bears considerable resemblance to the early Platycrinidae, and if we had the tegmen preserved it might be found to go with *Culicocrinus*.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

Of other species there is *M. striatus* Hall from the Waldron, three from the Racine dolomite, one from the Rochester shale (see Bassler's Index, p. 782); and a thoroughly representative species from England, *M. anglicus*, by Jackel. *M. indianensis* Miller and Gurley, from St. Paul, does not belong to the genus, and may be a *Patelliocrinus*. I am figuring (pl. 4, fig. 24) a complete specimen of the type species, *M. ornatus*, from Lockport, New York.

MELOCRINUS Goldfuss

Plate 5

Melocrinites Goldfuss, Petref. Germ., 1826, p. 197.

Melocrinus, Agassiz, Ann. Nat. Hist., 1838, p. 447.—Wachsmuth and Springer, Rev. Pal., 2, 1881, p. 118; N. A. Crin. Cam., 1897, p. 292.—Bather, Treatise on Zool., 3, 1900, p. 161.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 190.—Jackel, Phylogenie und System, 1918, p. 32.—Goldring, Devon. Crin. New York, 1923.—For complete list of references and table of synonyms, Bassler, Bibliogr. Index, Bull. 92, U. S. Nat. Mus., 1915, p. 794.

BB 4, unequal; iBr numerous, anal interradius but slightly distinct; rays produced into two main rami, giving off unilateral, pinnulate arms, uniserial or biserial, to the outside of the dichotomy: the rami may be separate (Silurian species), or more or less fused by their inner margins (Devonian species); column round.

Genotype. *Melocrinus hieroglyphicus* Goldfuss.

Distribution. Devonian and Silurian; Germany, Gotland, England, America.

Few genera have had so much written about them as this, or have been treated under so many different names, as will be seen by the list of references in the Bibliographic Index. The most recent discussion is that by Miss Goldring in the monograph of the Devonian Crinoids of New York, of which this genus and its allies form a most important part of the fauna. Hitherto it has been almost negligible among American Silurian fossils, the few species described being neither characteristic nor well represented. That which gives a renewed interest for this work is the discovery of a single specimen in the Beech River formation showing a wide distribution of the typical form, and the parallelism of its occurrence in this country and the north European Silurian. Very conspicuous species of what I regard as the Silurian type of the genus have been found in England and Gotland, some of them described by Angelin. Now for the first time we have a parallel to them from the American Silurian.

Melocrinus tennesseensis new species

Plate 5, fig. 1

Although the calyx plates of this unique specimen are somewhat disturbed, the decisive features of the rays are well preserved, and the general aspect of the fossil points unmistakably to a close relationship with the typical forms of England and Gotland. For comparison I am figuring two specimens of an abundant species from Dudley, England, which if not elsewhere described may be taken as *M. spectabilis* of Angelin (pl. 5, figs. 2, 3). The remarkable similarity of type in the two forms is apparent at a glance,—even the stem of one being almost a duplicate of the other. Evidently the calyx of the American species is much the shorter, the base of the cup less turbinate than in the English form, and the iBr in the latter more numerous and developed.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

Melocrinus oblongus Wachsmuth and Springer*Plate 5, figs. 5, 6*

Melocrinus oblongus Wachsmuth and Springer, N. A. Crin. Cam., p. 200, pl. 22, figs. 9, 12.

Confined to the higher Niagaran beds. Two specimens are figured especially to show the division of the rays into two rows of brachials, for comparison with the similar elongate form referred to Roemer's *Cytocrinus* with only one such row.

Horizon and locality. Louisville limestone, Louisville, Kentucky, and Bob formation, Niagaran; Decatur County, Tennessee.

Other Niagaran species are *M. obconicus* Hall, of the Waldron beds, *M. aequalis* S. A. Miller, of the Laurel; one from Missouri by Rowley, one from the Racine dolomite of the Chicago area and one from the Rochester shale at Lockport.

Melocrinus onondaga new species*Plate 5, fig. 4*

I figure here what appears to be a very young stage of the genus from the middle Devonian, which is interesting for the remarkable shortness of the calyx compared with the well developed arms.

Horizon and locality. Onondaga formation, Middle Devonian; Sylvania, Ohio.

Cytocrinus laevis Roemer*Plate 5, figs. 7, 7a, 8*

Cytocrinus laevis Roemer, Sil. Faun. Westl. Tennessee, 1860, p. 46, pl. 4, figs. 2a-c.—*Melocrinus roemeri* Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 301, pl. 22, figs. 11a, b.—Bassler, Bibliogr. Index, 1915, pp. 795, 796.

The brachials in this form are extended into 5 trunks composed of a single series of plates, which is a good generic distinction from similar lobate species such as *Melocrinus oblongus*, according to the practice followed in other groups; while the arms themselves have not been seen, the direct succession of primi-brachs into the arm opening shows how they must be.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

MARIACRINUS Hall*Plate 5*

Mariacrinus Hall, Am. Jour. Sci. (2), 25, 1858, p. 278; Pal. New York, 3, 1861, pp. 104, 139.—Wachsmuth and Springer, Rev. Pal., 2, 1881, pp. 114, 118, 233; *ibid.*, 3, 1885, p. 326; N. A. Crin. Cam., 1897, p. 281.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 190.—Goldring, Devon. Crin. New York, 1923, p. 109.—Bassler, Bibliogr. Index, 1915, p. 785.

BB 4; similar to *Melocrinus*, but arms free, branching once or twice, with alternating pinnules or ramules.

Genotype. *Mariacrinus plumosus* Hall.

Distribution. Silurian, Devonian; America, England, Gotland.

Mariacrinus carleyi (Hall)*Plate 5, figs. 9, 10*

Glyptocrinus carleyi Hall, Trans. Alb. Inst., 4, 1863, p. 203; 28th Rep. New York St. Mus., 1879, p. 132, pl. 14, figs. 7-10.—*Mariacrinus carleyi*, Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 282, pl. 22, figs. 2a-c.—Bassler, Bibliogr. Index, 1915, pp. 785, 786.

A very elongate, turbinate species, widely distributed and found at all the principal Waldron localities, and also in the Laurel. The specimen from the latter here figured lacks the sharp radiating ridges upon the lower plates usual in the Waldron specimens, the granulose surface predominating, but in the other characters the two agree.

Waldron shale, Niagaran; Waldron and Hartsville, Indiana, and Newsom, Tennessee; also Laurel limestone; St. Paul, Indiana.

Mariacrinus aureatus S. A. Miller*Plate 5, figs. 11, 11a*

Mariacrinus aureatus Miller, 17th Rep. Indiana Dep. Geol., 1892, p. 644 (adv. sheets, 1891), pl. 6, fig. 36.

A broadly rotund species, in almost every way the reverse of the preceding, especially in the highly sculptured ornament which covers the entire cup with delicate, stellate figures.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Mariacrinus sp.*Plate 5, fig. 13*

This specimen, associated with *Melocrinus* in the beds at Dudley, illustrates the presence of the two closely allied genera there as here; it shows the arms branching in the usual way, which is the distinguishing character.

Horizon and locality. Wenlockian, Silurian; Dudley, England.

(?) Mariacrinus rotundus new species*Plate 5, figs. 12, 12a*

This is one of some strange forms from the topmost Silurian, occurring in close relation to the Lower Devonian. It is so utterly different in general aspect from any other species of this or allied genera that it is placed here with much doubt. If seen in a well weathered condition in a miscellaneous collection, the specimen would without hesitation be taken for *Dizygocrinus rotundus* of the Burlington limestone. Yet according to the structure of the cup as plainly delineated, with no anal plates, and radials in contact all around, it must belong to the Melocrinidae, having abnormally small basals which cannot be defined in the narrow indented cavity, compensated by relatively large radials. The highly rounded tegmen is filled with numerous but very definite plates pierced by a small subcentral anal opening, while the iBr in the cup are but few.

Horizon and locality. Decatur limestone, Niagaran; below Grandview in the bluffs of the Tennessee river, Perry County, Tennessee. Associated with *Aorocrinus nodosus*, infra.

ALLOCRINUS Wachsmuth and Springer*Plate 6*

Allocrinus Wachsmuth and Springer, Geol. Surv. Illinois, 8, 1890, p. 206; N. A. Crin. Cam., 1897, p. 306.—Bather, Treatise on Zool., 3, 1900, p. 162.—Zittel-Eastman, Textb. Pal., 2d ed. 1913, p. 191.—Jackel, Phylogenie und System, 1918, p. 36.—Bassler, Bibliogr. Index, 1915, p. 25.

BB 3, unequal. Dorsal cup low and broad. No anal plates; iBr 1, large. Arms 10, uniserial.

Genotype. *Allocrinus typus* Wachsmuth and Springer.

Distribution. Silurian; America.

This genus was proposed upon the basis of very scant material, consisting of a calyx and a detached set of arms. The extremely low and broad contour of the former with its concave base, and the massive character of the latter, marked it as thoroughly distinct, confirmed later by a second species from a different horizon, also described from a poor specimen. While hoping for further information, I was agreeably surprised at the wealth of new material disclosed by the Tennessee collections, containing not only numerous excellent specimens of the type species, but also of at least two others, among them some with almost the complete crinoid beautifully preserved. Only a few of these are shown by the drawings.

Allocrinus typus Wachsmuth and Springer*Plate 6, figs. 1-4*

Wachsmuth and Springer, Geol. Surv. Illinois, 8, 1890, p. 207, pl. 14, figs. 7-8; N. A. Crin. Cam., 1897, p. 307, pl. 24, figs. 7a, b.

The long, broad and somewhat convex uniserial brachials give to the arms an aspect of massiveness and simplicity. Having a shallow, concave base, with radials forming part of the concavity and projecting into a prominent rim at the margin, the calyx is fairly broad and expands upward to the arm-bases. Cup plates all strongly nodose. With one of the well preserved crowns we are able to show the round stem, with its rather long columnals, to its full length, with terminal roots attached indicating attachment to a soft bottom.

Horizon and locality. Beech River formation, Niagaran; Decatur and Wayne counties, Tennessee.

Allocrinus ponderosus new species*Plate 6, figs. 5, 6*

Similar to the type, but differing in the extremely ponderous arms, widely swollen in the middle, combined with a very short and narrow calyx.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

Allocrinus longidactylus new species*Plate 6, figs. 7-10*

Distinguished by extreme length and relative slenderness of the arms, which impart a different aspect to the complete crown. Thus the arms lie much

farther apart and afford a view of the close fringe of pinnules, which are scarcely visible in the preceding species. The calyx is relatively broader, and the nodes upon the cup plates are very conspicuous, especially the single large iBr which by its central node stands out distinct from the others. One specimen is characterized by excessively large radials.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

***Allocrinus benedicti* S. A. Miller**

Plate 6, figs. 11, 12

Allocrinus benedicti Miller, 17th Ann. Rep. Indiana Dep. Geol., 1892, p. 647, pl. 7, fig. 1.—Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 308, pl. 24, figs. 8a, b.

Miller's figure gave no idea of generic or specific characters, being made from a fragment. I am figuring two good crowns to show the form as it occurs in the Laurel, in which the calyx plates are more broadly rounded and the arms rather less robust than in the type.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

PATELLIOCRINUS Angelin

Plate 6

Patelliocrinus Angelin, Icon. Crin. Suec., 1878, p. 1.—Zittel, Handb. Pal., 1879, p. 368.—Wachsmuth and Springer, Rev. Pal., 2, 1881, p. 100.—Bather, Treatise on Zool., 3, 1900, p. 162.—Zittel-Eastman, Textb. Pal., 3d ed. 1913, p. 191.—Jaekel, Phylogenie und System, 1918, p. 36.

BB 3, unequal, Dorsal cup elongate. No anal plates; iBr few. Arms biserial or cuneate uniserial.

Genotype. *Patelliocrinus pachydactylus* Angelin.

Distribution. Silurian; Gotland, America.

This Gotlandian genus, now for the first time doubtfully recognized in America, while having a simple calyx with 3 BB and a large iBr with anal side not differentiated, is widely separated from the preceding by its elongate, conical cup and biserial arms. Angelin described 9 species in which there is considerable confusion of generic characters. The figures cannot be depended on (Bather, Crinoidea of Gotland, p. 5) for details, and it is hard to tell exactly what should be considered as the type. But I have taken *P. pinnulatus*, pl. 24, fig. 5, as a fair representative, and there is among several of the species a general similarity which seems to warrant the reference of our species, in default of anything better.

***Patelliocrinus ornatus* new species**

Plate 6, figs. 13-15a

The surface, as shown by several specimens, is almost invariably ornate, with variations in intensity due to erosion or growth. It is best shown by figs. 15, 15a, from a free specimen unusually well preserved, in which the stout biserial arms and crowded pinnules are in place. The relatively small radials

are common to all the specimens. It is probable that the specimen named by Miller and Gurley *Macrostylocrinus indianensis*, Bull. 6, Illinois St. Mus., pl. 3, fig. 33, belongs to this genus, but the figure is too indistinct for comparison.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Patelliocrinus rugosus new species

Plate 6, figs. 16-18

With its strong rugose sculpturing, this species may be distinguished from the preceding by the great preponderance in size of the radials.

Horizon and locality, same as last.

Patelliocrinus laevis new species

Plate 6, figs. 19, 20

Differs from the two preceding by its perfectly smooth surface and greater elongation of calyx.

Horizon and locality, same as last.

————— genus and species indet.

Plate 6, figs. 21-21b

An apparently complete crown, having a low calyx of smooth, rounded plates, three small unequal BB, large RR followed by a single IBr which is axillary, filling most of the distal face of the R, and supporting two stout, biserial arms. There seems to be a single iBr, mostly not very well defined, but with the posterior one the largest, and no other indication of anal structures. It would seem that with so many details available it should not be difficult to determine the systematic position of this fossil, but after repeated efforts, aided by consultation with colleagues, I am still obliged to confess my inability to treat it intelligently. It is doubtless a new genus, but where to place it, I do not know. So I am giving the facts as I and the artist see them, and leaving it to others to continue the guessing.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

LAURELOCRINUS new genus

Plate 7

This genus is proposed for an assemblage of peculiar forms from the St. Paul beds, which will not fit into any already known. But for the tripartite base it might go under *Clonocrinus*, but that would mean rather wide variation from the type. Our specimens, while in considerable number, are all deficient in the upper part of the calyx and crown, so that we are without information

as to the arms, or even with certainty as to the higher brachials. The characters are an elongate calyx, cylindrical or slightly contracting upwards; base composed of 3 small BB, unequal, forming with the RR a broad, flattened, or shallow concave disk; RR very large, curving below into the base and upward into the sides of the cup; IBr and iBr have peculiar variations—the latter usually one elongate plate with or without a small one in succession. Four species are indicated, based upon rather minor differences visible in the specimens as preserved.

Genotype. *Laurelocrinus paulensis* new species.

Distribution. Silurian; America.

***Laurelocrinus paulensis* new species**

Plate 7, figs. 1-8

A large, robust species, with broad base and sides rising nearly straight; first IBr quadrangular, usually about square, and iBr elongate, 6- or 8-sided, followed by a second one much narrower; or exceptionally IBr may be elongate and iBr acuminate, not followed by a second; axillary IBr short, followed by 3 IIBr, the third axillary, indicating probably four arms to the ray. BB limited to a shallow cavity, occupied largely by the column-facet with very large lumen. Surface smooth.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

***Laurelocrinus wilsoni* new species**

Plate 7, figs. 9-13

A much smaller form than the preceding, even more so than the drawings indicate at first glance, those of the latter being natural size and of this enlarged by one half; and it is of similar contour; first IBr hexagonal, iBr 5-sided, followed by one or two others. Arms apparently 4 to the ray. Surface in lower part tending to be striate.

Horizon and locality, same as last.

***Laurelocrinus gibbosus* new species**

Plate 7, figs. 14-16

A similar calyx but much more rounded, with extremely large RR and iBr, and small quadrangular IBr, all strongly gibbous; base somewhat convex, with small column facet and pentagonal lumen.

Horizon and locality, same as last.

Laurelocrinus spinoradialis new species*Plate 7, figs. 17-17b*

A medium sized species, with RR armed with sharp spines projecting downward; first IBr quadrangular, short; iBr 2 in succession, very elongate, 8- or 9-sided; IIBr 3 in series, the first very large, the next two narrow and divided by an iIBr, the third axillary, indicating 4 arms to the ray. Based on the unique specimen figured.

Horizon and locality, same as last.

CLONOCRINUS Quenstedt (not Oehlert)*Plate 7*

Clonocrinus Quenstedt, Petref. Deutschl., 4, 1876, p. 638 (not Oehlert, 1879).—Bather, Treatise on Zool., 3, 1900, p. 162.—Zittel-Eastman, Textb. Pal., 2d ed. 1913, p. 190.—Jackel, Phylogenie und System, 1918, p. 32.

Corymbocrinus Angelin, Icon. Crin. Suecc., 1878, p. 18.—Zittel, Handb. Pal., 1879, p. 373.—Wachsmuth and Springer, Rev. Pal., 3, 1885, p. 106.

BB 4, hidden by column; iBr few, one large followed by small ones in vertical row; no anals. Arms dichotomous, biserial below and above the bifurcations.

Genotype. *Eucalyptocrinus polydactylus* McCoy.

Distribution. Silurian; Gotland, England, America.

This genus is regarded as leading from the Melocrinidae towards the Calyptocrinidae, having a similar distribution of the cup plates, but with iBr beyond the first not paired, and the arms not resting in compartments between partitions. It is essentially a Gotlandian form, of which six species have been described under the name *Corymbocrinus*. The genus is doubtfully recognized here from some imperfect specimens derived from a formation which has produced a number of new and strange forms, one of which is figured.

Clonocrinus occidentalis new species*Plate 7, figs. 18, 18a*

Has deeply sunken basal cavity, with the small BB at the bottom; and the iBr following the first in single vertical succession.

This may be compared with *Corymbocrinus laevis* Ang., pl. 23, fig. 20.

Horizon and locality. Decatur limestone; near Perryville, Tennessee.

Family CALYPTOCRINIDAE Angelin

Monocyclic. Dorsal cup usually with 4 BB and concave base; lower brachials and interbrachials incorporated in cup, which above the base is perfectly pentamerous; plates of calyx usually limited to a definite number; radials in contact all around; no anals; arms, usually 20, resting in pairs in compartments formed by vertical partitions borne by processes rising from the fixed iBr and iIIBr to varying heights; tegmen elevated in a central anal tube, and composed of 4 circlets of large plates variously shaped bearing the partitions. Silurian and Devonian.

For instructive diagrams illustrating the structure, see N. A. Crin. Cam., 1897, p. 331.

EUCALYPTOCRINUS Goldfuss*Plates 7, 8, 9*

Eucalyptocrinites Goldfuss, Petref. Germ., 1826, p. 214.

Eucalyptocrinus, Agassiz, Ann. Nat. Hist., 1838, p. 447.—Wachsmuth and Springer, Rev. Pal., 3, 1885, p. 127; N. A. Crin. Cam., 1897, p. 332.—Bather, Treatise on Zool., 3, 1900, p. 164.—Weller, Bull. Chicago Acad. Sci., 4, pt. 1, 1900, p. 102.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 192.—Jaekel, Phylogenie und System, 1918, p. 32.—Complete reference with list of synonyms and species, Bassler, Bibliogr. Index, 1915, pp. 501-507.

Calyx elongate to wide spreading; BB 4, at bottom of an inverted funnel formed by RR at the sides, followed by IBr, IIBr and IIIBr, 2 each; iBr one large 8- or 10-sided, angular above, supporting a pair of elongate plates of second range extending upward between the arm-bases to a connection with the partitions; a single iIIBr of similar form rests upon the truncate second IBr and connects with the partition in line with the ray. No anals. Arms biserial, paired, resting in 10 vertical compartments bounded by partitions supported by processes arising from the iBr and iIIBr plates extending the full height of the tegmen and meeting above in a circlet of peripheral plates. Tegmen composed usually of 4 ranges of plates, tubular, expanding upward into a canopy having an inner set of plates variable in number, often 4, surrounding a central anal opening, either direct, or produced into a short tube.

Genotype. *Eucalyptocrinus rosaceus* Goldfuss.

Distribution. Silurian; Gotland, England, America. One species in Middle Devonian, Germany.

This prolific Silurian genus, first described from a Devonian species, which still remains the only one known from that epoch, has by the growth of research become the leading known crinoid of its typical horizon, both in variety and distribution. Some 40 species have now been described from the European and American areas, including 7 from Gotland, 3 from England, and 30 from the Niagaran of America, some of which are probably synonyms. The American species are distributed among the subdivisions of the Niagaran as follows: Rochester 4, Waldron 7, Racine 9, Beech River 8, Decatur 2. The last 2 are now first described, one of them marking a new departure from the type. The Tennessee collections

from the Beech River formation and the Waldron shale at Newsom have yielded exceptionally fine material of already known species, some of which I am illustrating with new figures, where the characters are better shown than in the originals. Several of the leading forms are so amply treated in the published accounts that further attention is not needed. A full list of the species with their references is given in Bassler's Index. This genus is remarkable for the absolute uniformity of its cup plates. In all the numerous species there is scarcely a variation from the number, form and distribution of the plates as above stated.

***Eucalyptocrinus elrodi* S. A. Miller**

Plate 8, figs. 1, 2

Eucalyptocrinus elrodi Miller, 17th Rep. Indiana Dep. Geol., 1892, p. 650 (adv. sheets, p. 40), pl. 7, figs. 9, 10.—Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 339, pl. 81, figs. 7a, 8-13.—Bassler, Bibliogr. Index, 1915, p. 503, 504.

The colony of the Waldron shale at Newsom seems to have been particularly favorable to the growth of this and the succeeding species, they having attained a size unprecedented in the genus. I figure here two calices far exceeding those from the type locality, and in which the pustulose ornament in two styles is wonderfully sharp, and the calyx plates are delineated with the clarity of a diagram.

Horizon and locality. Waldron shale, Niagaran; Hartsville and Waldron, Indiana; Newsom, Tennessee.

***Eucalyptocrinus magnus* Worthen**

Plate 8, fig. 3

Eucalyptocrinus magnus Worthen, Geol. Surv. Illinois, 6, 1875, p. 501, pl. 25, fig. 3.—Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 348, pl. 82, figs. 7, 8.—Bassler, Bibliogr. Index, 1915, p. 505.

This figure of a cup 75 mm. in diameter is given to show the maximum size that may be attained by the genus. It is not an isolated example, as the species is typically large, with cup low, broad and smooth, and specimens are not uncommon of approximately the dimensions of this. It is abundant in the Waldron beds at Newsom, and a characteristic fossil of them, not having been found at the Indiana localities, although it is reported from the Racine dolomite of the Chicago area. For the opposite extreme in size reference should be had to *E. ovalis*, of which some specimens are less than 5 mm. in diameter, and which rarely exceed 10 or 12 mm. (see N. A. Crin. Cam., pl. 82, fig. 4).

Horizon and locality. Waldron shale, Niagaran; Newsom, Tennessee.

***Eucalyptocrinus lindahli* Wachsmuth and Springer**

Plate 8, figs. 4-5

Eucalyptocrinus lindahli Wachsmuth and Springer, Am. Geol., 1892, p. 139; N. A. Crin. Cam., 1897, p. 347, pl. 82, fig. 9.—Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 47, pl. 12, figs. 5, 6.—Bassler, Bibliogr. Index, 1915, p. 504.

A rare species, which was unknown to collectors from the time of Troost, who exhibited in 1849 what he rightly called a magnificent specimen under the

name *Eucalyptocrinus splendidus* which remained buried in his unpublished monograph until 1909, down to the date of Wachsmuth and Springer's description. Only two or three specimens have been found, all remarkably true to type, from which I am giving figures of a very mature and what may be a young individual. The depressed interbrachial partitions wholly lack the thickening and wide extension of those of *E. milliganae*; instead they are narrow and knife-like, with a peculiar splitting or doubling at the extreme upper end—seen also in *E. ventricosus*.

Horison and locality. Beech River formation, Niagara; Wayne and Decatur counties, Tennessee.

***Eucalyptocrinus milliganae* Miller and Gurley**

Plate 8, figs. 6-8

Eucalyptocrinus milliganae, Miller and Gurley, Bull. 10, Illinois St. Mus., 1896, p. 88, pl. 5, figs. 4-6.—Bassler, Bibliogr. Index, 1915, p. 505.

This is one of the best defined species of the genus, but Miller and Gurley's type specimens were imperfect and failed to show its most distinctive character, the wing-like projections of the interbrachial partitions, as is seen in the three specimens I have figured, especially fig. 6. These are selected out of a dozen or more crowns almost equally perfect, from a colony in the uppermost shale of the Beech River formation at Tuck's Mill, which by their prominence gave to that bed the name *Eucalyptocrinus* zone. These specimens vary greatly in size, from 20 to 50 mm. in height of crown, but without exception they have the projecting partitions more or less indicated by which the species is readily distinguished. This is modified by growth, being more pronounced in the older individuals. Along with this character the turbinate calyx, broadly truncate at the base, and the smooth surface, are substantially as described, and the median swelling of the arms, narrowing to fine points above, is to be noted; they do not fold over the tegmen or encroach upon it in any way, but the widening partitions curve over to a junction with the central parts, forming a broad, flat circular roof, with all its plates well exposed. This is in marked contrast with *E. ventricosus* in which the exposed part of the tegmen is reduced to a small disk.

Horizon and locality. Beech River formation. *Eucalyptocrinus* zone, Niagara; Tuck's Mill, Decatur County, Tennessee.

***Eucalyptocrinus ventricosus* Wachsmuth and Springer**

Plate 8, figs. 9-12

Eucalyptocrinus ventricosus Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 341, pl. 83, figs. 11, 12.—Bassler, Bibliogr. Index, 1915, p. 507.

The most abundant species in the Beech River formation, being found at almost all the localities throughout the Tennessee river area, but only at the Tuck's mill beds have such finely preserved specimens been recovered as those

I am figuring. It is notable always for the full rotundity of contour, the arms infolding closely over the margin of the tegmen, with slightly visible partitions, leaving exposed only a small disk in the center, in strong contrast to the wide-spreading roof of *E. milliganae*.

Horizon and locality. Beech River formation, Niagaran; Decatur, Wayne and Perry counties, Tennessee.

***Eucalyptocrinus* sp.**

Plate 8, fig. 13

A form from the Beech River formation, Decatur County, Tennessee, with cup once and a half wider than high, and unusually tumid plates, probably different from the other species.

***Eucalyptocrinus pernodosus* new species**

Plate 9, figs. 1-4

This species, from the uppermost Niagaran, stands out distinct from the others by its extremely low and broad cup, twice as wide as high, relatively enormous RR, and the high convexity of its cup-plates. In mature specimens, as in figs. 1 and 2, this goes to the extreme of nodosity, the plates being medially produced into spinous projections resembling pegs. The crown is unusually elongate, exceeding the longest of *E. milliganae* by 20 per cent, but this excessive length is strictly in the arms, which are also stout, with little taper, but folding over at the tips between the narrow partitions, which are thus left well exposed surrounding a relatively small median disk. This and the forms following are the only examples of the genus as yet found in this formation. In contour of cup the species may be compared with *E. depressus* of S. A. Miller, from the Racine dolomite, figured in N. A. Crin. Cam., pl. 82, fig. 3.

Horizon and locality. Decatur limestone, Niagaran; quarries above Perryville, Tennessee.

***Eucalyptocrinus sculptilis* new species**

Plate 9, figs. 5-6a

This companion species, from the same horizon as the last, is the reverse of it in most of the characters. Instead of the broad, low cup, it is here of the subturbinate type frequent in the genus, and in place of the striking convex or nodose elevation of the plates, there is an equally striking surface marked by a delicate sculpturing extending from the RR to the uppermost cup-plates. The singular appearance of the stem shown in fig. 5 is due to a peculiar spreading of the long nodal ossicles, exposing between them a greater number of short internodals. The structure is evidently similar to that seen in the Devonian *Dolatocrinus*, as is fully explained and illustrated in my paper upon that genus,

Bull. 115, U. S. Nat. Mus., 1921, pp. 16-18, pls. 9, 10. Presumably the long rounded columnals in the stem of fig. 5 are nodals similarly overgrown, but without separation to expose the intervening plates.

Horizon and locality. Decatur limestone, Niagaran; Bluffs near Perryville, Tennessee.

Eucalyptocrinus sp.

Plate 9, figs. 7, 7a

A form from the same horizon as the preceding having a perfectly smooth surface and turbinate calyx, which may be a new species, but the characters are not sufficiently pronounced for description, and there is a little uncertainty as to the number of lbr.

Horizon and locality, same as last.

Eucalyptocrinus crassus Hall

Plate 7, figs. 19, 20

Eucalyptocrinus crassus Hall, Trans. Albany Inst., 4, 1863, p. 197.—Bassler, Bibliogr. Index, 1915, p. 503.

The most abundant American species, amply illustrated in various publications noted in the Index. The two figures are given here merely for comparison with *Gasacrinus* in regard to the ventral pyramid. The structure is further shown in the Camerata monograph, pl. 81, figs. 12, 14, 15.

Horizon and locality. Waldron shale; the specimens figured are from Hartsville, Indiana, but the species also occurs abundantly at Waldron, and at Newsom, Tennessee, and is reported from the Racine dolomite of the Chicago area.

CALLICRINUS D'Orbigny

Plate 9

Callicrinites D'Orbigny, Prodr. Pal., 1, 1849, p. 45.

Callicrinus Pictet, Traité Pal., 2d ed., 4, 1857, p. 301.—Wachsmuth and Springer, Rev. Pal., 3, 1885, p. 135; N. A. Crin. Cam., 1897, p. 353.—Weller, Jour. Geol., 5, 1897, p. 774; Bull. Chicago Acad. Sci., 4, pt. 1, 1900, p. 117, fig. 47.—Bather, Treatise on Zool., 3, 1900, p. 164, fig. 77.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 192.—Complete reference and list of species, Bassler, Bibliogr. Index, 1915, p. 158.

Similar to *Eucalyptocrinus* in the cup, but basal concavity usually broader; tegmen contracting to a decanter shape, and partitions only extending part way to ends of arms; large spinous processes frequently given off from plates of cup and tegmen, the upper circlet of the latter often extended in four quadrant-shaped horizontal projections, sometimes to a considerable length.

Genotype. *Eugeniocrinites costatus* Hisinger.

Distribution. Silurian; Gotland, England, America.

This closely allied form, also typically Gotlandian, has added eight more species to the Calyptocrinid type for that area, besides one or two perhaps undescribed, from England. It also became well established in the American Silurian, where if judged by the number of reported species it would appear to be one of the leading crinoids, limited, with one exception, to a single formation or its equivalent, the Racine dolomite of the Chicago area, from which nine species have been described. As these species are nearly all based upon internal casts of the calyx, in which some of the characters are obscure or not shown at all, there is probably

in this genus, as well as in *Eucalyptocrinus*, some duplication. The generic relations of the species were at first not understood, some being described under *Eucalyptocrinus* and some as a strange organism called by Hall *Cryptodiscus*, now known to be appendages of the *Callicrinus* tegmen. Weller was the first to point out this fact, and his studies upon the abundant material of the Chicago area did much toward clarifying our understanding of the American occurrences.

Most instructive figures for the explanation of the genus are given by Bather in the Treatise on Zoology (Lankester), 3, 1900, p. 163.

The exceptional species from the Laurel formation described by Wachsmuth and Springer from a single specimen can now be further illustrated, and some of its characters better defined. From fragments we know that there were other species in the same beds.

Callicrinus beachleri Wachsmuth and Springer

Plate 9, figs. 8-13

Callicrinus beachleri Wachsmuth and Springer, Am. Geol., 10, 1892, p. 140; N. A. Crin. Cam., 1897, p. 355, pl. 83, figs. 14a, b.

The series of figures from six specimens here given are especially directed to the surface characters of the dorsal cup, which is notable for the extremely high and sharp wavy ridges which traverse the plates from radials to arm-bases. In the type specimen this was much obscured by hard matrix, but these specimens occurred in a soft granular limestone readily removed by preparation. The smooth, decanter-shaped pyramid enclosed by the arms, with the very short partitions at the lower margin, are also well shown.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Callicrinus sp.

Plate 9, figs. 14-15

These are the terminal quadrant-shaped plates of the tubular tegmen which abut by fours, with a small anal opening in the middle. First described by Hall as *Cryptodiscus*, they have been thoroughly explained by Weller in his paper for the Chicago Academy, and Bather's figures 2, 3 and 4, on page 163 of the Lankester Zoology, show how they go together.

Horizon and locality. Laurel limestone, St. Paul, Indiana.

Callicrinus sp.

Plate 9, fig. 16

This is another of the quadrant plates which were only connected with their fellows by short sutures, and beyond that were produced into greatly elongated, rounded appendages. Doubtless these types represented different species, perhaps some of those described by Weller from the Chicago area, but we have only the fragments of this and the preceding form, which are too meagre for comparison of species.

Horizon and locality, same as last.

Family BATOCRINIDAE Wachsmuth and Springer

Monocyclic. Base hexagonal. BB 3, equal. Lower brachials and inter-brachials incorporated in dorsal cup; radials in contact except at anal side; proximal anal plate in line with RR, heptagonal, followed by three in second range; rays usually branching within the cup, and the free arms simple, pinnulate; first primibrach quadrangular except in *Periechocrininae*. Ordovician to Carboniferous.

Jaekel, *Phylogenie und System*, 1918, p. 34, goes back to the plan of Angelin, and while rejecting Monocyclica and Dicyclica as grand divisions of the crinoids, proposes monocyclic suborders based upon the number of basals; but after listing those of 4 and 5 BB as Tetramera and Pentamera, he throws the remainder, whether with 3, 2, or undivided basals, into a single suborder, *Miomera*. Under this he puts together all Batocrinidae and Actinocrinidae into the younger family, Actinocrinidae, a short-lived derivative of the other, on the ground that the distinguishing character between them, viz.: that the first has three and the last two anals in the second range "would scarcely be sufficient even for separating genera." In this he completely ignores the almost unrivaled constancy of the characters by which these two families are separated according to the accepted classification.

It is now well known that among the crinoids modifications of the anal structures yield characters of the highest taxonomic value. A character in zoological classification derives its value, as has been repeatedly pointed out by the most eminent authorities, not from its physiological importance, but from the uniformity with which, however unimportant functionally, it prevails throughout many different species, and is common to a great number of forms and not common to others. Here the family Batocrinidae, extending from the Ordovician to the later Lower Carboniferous, with nearly 300 species, distributed among 24 genera which differ among themselves by many striking and conspicuous features, is held together by a definite structure of the anal area, without the slightest deviation save an occasional sporadic specimen. A perfectly evident transitional genus leads to the Actinocrinidae, the character of which is equally constant and invariable in over 100 species, and 7 thoroughly well defined genera.

No one looking at a complete collection exhibiting the entire assemblage of genera and species under these two forms, or at the ample illustrations of them as given in the *Camerata* monograph, will doubt for a moment that the far-reaching characters by which they are defined mark the lines of descent. For an author of a systematic work to say that they would hardly serve for separating genera, is pure dogmatic assertion, made without the slightest consideration of the facts.

Subfamily CARPOCRININAE Bather

BB 3; RR rather large; IBr quadrangular; IIBr usually passing into the free arms, mostly 2, occasionally 3, 4, or 5 to the ray; iBr definitely arranged; anal plates in vertical row; tegmen composed of numerous small plates, with orals in the center compressed; column round.

CARPOCRINUS Müller

Plate 7

- Carpocrinus* Müller, Monatsb. Berlin Acad. Wiss., 1, 1848, p. 208; Verh. Naturh. Verein, 12, 1855, p. 19.—Wachsmuth and Springer, Rev. Pal., 2, 1881, p. 105 (full synonymy, pp. 107, 108).—Bather, Treatise on Zool., 3, 1900, p. 166.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 194.—Jaekel, Phylogenie und System, 1918, p. 37.
- Phoenicocrurus* Austin, Ann. Mag. Nat. Hist., 11, 1843, p. 205.—*Abracrinus* D'Orbigny, Prodr. Pal., 1, 1850, p. 47.—*Habrocrinus* Angelin, Icon. Crin. Suecc., 1878, p. 3; *Pionocrinus* Ang., *ibid.*, p. 4; *Leptocrinus* Ang., *ibid.*, p. 3.

Calyx rotund with protuberant base; iBr usually in several ranges; anals in prominent ridge; arms stout, not over two to the ray, with brachials uniserial, quadrangular or slightly cuneate, bearing a pinnule on each side, sometimes two at the longer face.

Genotype. *Actinocrinites simplex* Phillips.

Distribution. Silurian; Gotland, England, America.

This genus, treated under several names, is one of the most prolific of the north European Silurian, from which no less than nineteen species have been described, all occurring in Gotland, and at least one also in England. A full list is given by Wachsmuth and Springer, Rev. Pal., 2, pp. 107-8. Representing as it does an early and simple Batocrinid type, some migrational extension into the American Silurian might well be expected, and in fact was looked for with confidence as the Niagaran fauna of Indiana and Tennessee began to show its remarkable development. So far this expectation has been realized by only a single species, which is believed to be fully characteristic as judged by the calyx alone. For comparison, and to illustrate the accompanying type of arms with their doubled pinnules, I am figuring on plate 7 a complete crown of *C. simplex*, the species which is common to England and Gotland.

Carpocrinus sculptus new species

Plate 7, figs. 21-21b

Except for the absence of arms, the specimen is very perfect, the calyx being entirely free including the tegmen; even of the arms there are short stumps in one ray, enough to show that they have the number, stoutness and uniserial brachials characteristic of the genus. The calyx is well rounded, with protuberant base, proportionally more elongate and with more ranges of iBr than usual. Median row of anal plates very conspicuous, passing up into the tegmen to an opening midway from the center and next to the posterior oral, bordered by a circlet of small convex plates. Tegmen strongly marked by 5 very distinct orals, 4 small partly enclosing the much larger posterior one, and extremely conspicuous ambulacra with a double series of covering plates, passing outward and branching to the 10 arm-openings; interambulacra small and numerous. Surface of cup plates very ornately sculptured; this is sharp and distinct on the part which was imbedded in the soft matrix, fig. 21, but in fig. 21a, owing to erosion, the surface appears perfectly smooth.

Two of Angelin's species, *Habrocrinus ornatissimus* and *H. ornatus*, have a similar ornamentation, but are otherwise different.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

DESMIDOCRINUS Angelin*Plate 7*

Desmidocrinus Angelin, Icon. Crin. Suecc., 1878, p. 5.—Wachsmuth and Springer, Rev. Pal., 2, 1881, p. 108.—Bather, Treatise on Zool., 3, 1900, p. 166.—Zittel-Eastman, 2d ed., 1913, p. 194.—Jaekel, Phylogenie und System, 1918, p. 37.

Calyx as in *Carpocrinus*, but with iBr few, anals not in prominent row, arms irregular, 3, 4, or perhaps 5, to the ray, uniserial; brachials quadrangular.

Genotype. *Desmidocrinus pentadactylus* Angelin.

Distribution. Silurian; Gotland, England, America.

Desmidocrinus laurelianus new species*Plate 7, figs. 24-24b*

This and the succeeding species are referred to the above Gotlandian genus because of the increase of arms above 10, and the smaller iBr areas. Here the latter consist chiefly of a single large plate. In both the anal interradius is fully developed, but not in the form of a raised vertical row as in *Carpocrinus*. In this the ray branches on the second IBr. Cup plates are convex and smooth.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Desmidocrinus dubius new species*Plate 7, fig. 25*

Distinguished from the preceding by the mode of arm-branching, which here occurs first on about the sixth IBr, and perhaps again on one ramus, thus giving 4 or 5 arms to the ray. The calyx is more elongate and cup more turbinate, and the reference to the genus is made with some doubt.

Horizon and locality, same as last.

Cylicocrinus canaliculatus S. A. Miller*Plate 7, figs. 26-29*

Cylicocrinus canaliculatus Miller, 18th Ann. Rep. Indiana Dep. Geol., 1894, p. 285 (adv. sheets, 1892, p. 31), pl. 5, figs. 13, 14.—As *Barrandocrinus*, Wachsmuth and Springer, N. A. Crin. Cam., 1897, pp. 484, 485.—Bather, Treatise on Zool., 3, 1900, p. 166.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 195.—Bassler, Bibliogr. Index, 1915, p. 101.

The mystery surrounding the relations of the genus founded upon this species as type has not been solved by any of the recently acquired material, by which our information is still limited to the dorsal cup below the arm-bases; it remains insufficiently known for a generic definition. It has the calyx of the Batocrininae, similar in all visible respects to that of the Swedish *Barrandocrinus* with the recumbent arms—a similarity which led Wachsmuth and Springer to think it might possibly belong to that genus. The specimens I have since obtained show a thin, flange-like projection at the base which seem to me inconsistent with the close enveloping of the cup by the pendent arms such as is indicated by the various figures of *Barrandocrinus*, for example in my recent paper on Unusual Forms of Fossil Crinoids, Proc. U. S. Nat. Mus., 67, 1926, pl. 9, figs. 6-7.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Cylicocrinus spinosus new species*Plate 7, figs. 30, 30a*

Founded upon an isolated hexagonal base with prominent spinose projections, differing thus from all the other specimens. May be only a variant of the type.

Horizon and locality, same as last.

Subfamily PERIECHOCRININAE Bather

BB 3. IBr hexagonal; iBr numerous, passing into iAmb; tegmen composed of numerous small undifferentiated plates.

PERIECHOCRINUS Austin*Plate 10*

Periechocrinites Austin, Ann. Mag. Nat. Hist., 10, 1842, p. 109.

Periechocrinus McCoy, Syn. British Foss., 1855, p. 56.—Wachsmuth and Springer, Rev. Pal., 2, p. 127 (list of species); N. A. Crin. Cam., 1897, p. 519.—Bather, Treatise on Zool., 3, 1900, pp. 166, 168.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 194.—Jaekel, Phylogenie und System, 1918, p. 35.—Bassler, Bibliogr. Index, 1915, pp. 953, 954 (references, synonyms and list of American species).

Calyx elongate, slender, expanding to the arm-bases; cup-plates long and thin, with prominent axial folds along radial series; rays bifurcate two or three times within the cup, leading to from 20 to 40 free arms, biserial, and usually not branching after becoming free.

Genotype. *Actinocrinites moniliformis* Phillips.

Distribution. Silurian; Gotland, England, America.

Of the large number of Silurian species which have been referred to this genus, at least one is from England, 9 are from Gotland, and 11 from North America. The latter are distributed as follows: Rochester shale, 1; Laurel limestone, 2; Waldron shale, 1; Brownsport, 3; Racine dolomite, 4. Besides these are two from the Lower Carboniferous, and one in Europe from the Devonian which may go better under *Saccocrinus*.

The typical form, as represented by the abundant and well known English species, *P. moniliformis*, is well marked and readily distinguished by the extremely elongate, slender cup, thin plates, unusually high RR and IBr with their conspicuous axial folds running to the arm-bases. Several species of that type have also been described from America, but along with them are some others which while having the same general calyx arrangement are wanting in the above characters, having the cup-plates lower, thicker, more convex and without folds. Taking as an example of this type such a form as the Carboniferous *P. whitei*, pictured in N. A. Crin. Cam., pl. 6, fig. 1, one is impressed with the force of the doubt with which it is listed. In the Zittel-Eastman Textbook, edition of 1913, I undertook to revive Hall's discarded genus *Saccocrinus* as a receptacle for some of these non-characteristic species, and am making the distinction accordingly here for two species. An entirely new form for the Tennessee Silurian turned up in the Beech River formation, clearly belonging to the English and Swedish type, but unfortunately too much injured for a specific definition.

Periechocrinus tennesseensis (Hall)*Plate 10, figs. 1-4*

Saccocrinus tennesseensis Hall, Geol. Surv. Ohio, Pal., 2, 1875, p. 125, pl. 6, fig. 10.—*Periechocrinus tennesseensis*, Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 528, pl. 50, fig. 4 (figured as Roemer's type of *Saccocrinus speciosus*).—Wood, Bull. 64, U. S. Nat. Mus., 1909, pp. 76, 78, pl. 6, fig. 10.—*Saccocrinus speciosus* Roemer (not Hall), Sil. Fauna Westl. Tennessee, 1860, p. 42, pl. 3, figs. 3a-c.—Bassler, Bibliogr. Index, 1915, p. 956.

The species is well known to collectors in the Tennessee area from the earliest times. Both Troost and Roemer recognized it, and in the modern collections it is not uncommon. It is rather on the border line between the two allied genera as they are here defined. I am figuring four specimens in order to show some of the minor variations which must be considered in any critical comparison of the described species. Some show the axial folds by obscure traces, and some scarcely at all; but all have the slender calyx and narrow turbinate base usually seen in the genus. The arms vary in different individuals not otherwise distinguishable, from 4 to 5 to the ray. The species has been found at the principal localities of the Brownsport group.

Horizon and locality. Beech River and perhaps other formations of the Brownsport, Niagaran; Decatur, Perry and Wayne counties, Tennessee.

Periechocrinus sp.*Plate 10, figs. 5, 5a*

The crushed and damaged specimen here figured is of a distinctly different type from the preceding, and has much the character of the typical English form; it is the only one thus far found that seems to approach the European type so closely; if in better condition for observation it would doubtless bear close comparison with some of those species. The calyx walls are very thin, and the axial folds leading to the arms sharp and prominent. In the number of the branching, biserial arms, which seem to run to 7 or 8 to the ray, the specimen shows a significant resemblance to Angelin's figures of *P. moniliformis*, Icon. Crin., pl. 19, figs. 14, 14b.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

SACCOCRINUS Hall*Plate 10*

Saccocrinus Hall, Pal. New York, 2, 1852, p. 205; 28th Rep. New York St. Mus., 1879, p. 127.—Roemer, Sil. Fauna Westl. Tennessee, 1860, p. 44.—Meek and Worthen, Geol. Surv. Illinois, 3, 1868, p. 347; *ibid.*, 5, 1873, p. 394.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 194.—Bassler, Bibliogr. Index, 1915, p. 954.

Calyx arranged like *Periechocrinus*, but the arms from about 20 (exceptionally 10) openings branch after becoming free, and are biserial both below and above the bifurcations. Plates usually thicker and more convex, and without axial folds.

Genotype. *Saccocrinus speciosus* Hall.

Distribution. Silurian. Devonian and Lower Carboniferous. America, Europe.

One of the Gotlandian species, *P. laevis* (Ang. not Portlock) = *P. minor* Wachsmuth and Springer, Icon. Crin., pl. 18, fig. 11, differs from the dominant type in a way that would lead toward this genus. The Carboniferous species before alluded to would be much more at home in this restricted company, and the larger genus will be better balanced and more consistent in its characters with such forms as the first one here recorded eliminated.

***Saccocrinus benedicti* S. A. Miller**

Plate 10, figs. 6-14

Saccocrinus benedicti, Miller, 17th Ann. Rep. Indiana Dep. Geol., 1892, p. 283 (adv. sheets, p. 29), pl. 5, fig. 1.—*Periechocrinus ornatus*, Wachsmuth and Springer, N. A. Crin. Cam., 1897, pl. 51, fig. 7 (not pl. 50, figs. 3a, b).

A strongly marked species, abundant at St. Paul and obtained in excellent preservation minus the arms. The contrast between it and *P. tennesseensis* figured on the same plate is most striking. There is a stout, ovoid calyx, with rather broad base, thick plates of which the surface varies from simple convex to rugose, nodose, and even perhaps stellate ornament. The arm-openings stand out boldly from the calyx at about the third IIBr, thus limiting the free arms to two to the ray; how they branch, if they do, beyond that is unknown. In this small number of arms the species differs widely from all others of the genus thus far known. The one specimen with stellate ornamentation, fig. 14, is probably of a different species, but lacks too much in the arm region for definition.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

***Saccocrinus cuspidatus* new species**

Plate 10, fig. 15

A form partaking more of the general shape of *Periechocrinus* than the preceding, but with a less slender, far larger and more robust calyx; while the shorter cup-plates with their sharp cusps instead of folds differentiate it completely, and the latter character from all other species otherwise associated. The arms apparently become free with the IIIBr, which would give an average of four to the ray, probably subject to variation. The specimen is from a higher formation than the others mentioned.

Horizon and locality. Louisville limestone, Niagaran; Jefferson County, Kentucky.

In the Laurel limestone at St. Paul, there is another large species, ill defined, described by Miller and Gurley as *Saccocrinus umbrosus*, Bull. 6, Illinois St. Mus., 1895, p. 24, pl. 2, figs. 13, 14.

Subfamily BATOCRININAE

Tegmen broad, well differentiated, plates large and heavy, forming a rigid roof; arms not branching beyond the cup; biserial; first IBr usually quadrangular. Silurian to Lower Carboniferous.

This extensive subfamily, typically Carboniferous, and hitherto considered as having originated in the Devonian, must now be moved a step farther back in the geological scale and credited to the terminal Silurian, as is proved by the occurrence of a well defined species in the Tennessee area.

Aorocrinus nodosus new species

Plate 10, figs. 16-16b

Calyx rotund, with slightly protuberant base; plates elevated into sharp nodes; iBr few, not connecting with tegmen except at the anal side. Anus sub-central without a tube. Tegmen plates smooth, and none of them spiniferous.

The single specimen upon which this species is based was found in the uppermost formation of the Tennessee Niagaran, the Decatur limestone, in a bed which is closely associated with the overlying Linden with its characteristic Helderbergian species of the Devonian *Scyphocrinus*. Nevertheless the presence in the same layer of such an unquestionable Niagaran form as *Caryocrinus ornatus* leaves no room for doubting the stratigraphic position of our species, which is thus in accord with the great difference in its facies from that of the hitherto known Devonian forms. The extremely rotund and robust calyx and sharply nodose cup-plates produce a distinctly new type, for which, however, we may have a connecting link in a species from the Hamilton recently acquired, and herewith figured.

It has been contended that the genus *Aorocrinus* as founded by Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 470, should give way to *Coclocrinus* of Meek and Worthen. While this may be true for some Lower Carboniferous species, that genus was founded upon a type with very broad, concave base, from which most of the forms referred to *Aorocrinus* are sufficiently differentiated, but which may also have a forerunner in the new Devonian species here figured.

Horizon and locality. Decatur limestone, Niagaran; below Grandview, 10 miles above Clifton on the Tennessee River, Perry County, Tennessee. Associated with *Mariocrinus rotundus*.

Aorocrinus clarkensis new species

Plate 10, figs. 17, 17a

This isolated specimen from the Devonian is figured for comparison with the preceding species. Whether it belongs to this genus is not clear, in the absence of the parts in the arm-zone. It is derived from the Hamilton beds of Clark County, Indiana.

Family PLATYCRINIDAE Roemer

Monocyclic. RR in contact all around. No anal. BB 3, unequal, forming a pentagon.

This monocyclic family is intermediate between the typical Camerata and the Inadunata. As in the latter, the dorsal cup consists chiefly of basal and radial structures, those of inter-radial origin being in the later stages largely confined to the tegmen. It had a long geological range, from Silurian to Lower Carboniferous, and in the latter its typical forms were fully developed, in which the plates above the radials enter but slightly into the cup, and the stem becomes remarkably specialized with an elliptical section and transverse ridge. It culminated with a great number of characteristic species, in which the dominant characters may be seen and studied to best advantage.

The Silurian and Devonian stage exhibits some marked variations, but in the main embraces forms of the simpler and more primitive types, showing the least departure from the Camerate plan. While they are for the most part rare, the present collections have yielded abundant representatives of two of the subdivisions. In each the material is of exceptional importance, because the specimens belonging to some forms occur in such numbers and preservation as to afford the means of comparative studies not hitherto possessed on either continent, requiring in one case the recognition of a new genus.

Section I. Subfamily COCCOCRININAE Bather

Rays with two or more primibrachs.

The division here first discussed, embracing forms with 2 or more IBr, represents the simplest form of the Camerata, the calyx consisting only of 3 BB forming a pentagon, 5 RR, 2 x 5 IBr, 5 iBr and 5 orals. The tegmen is almost completely occupied by 5 large triangular orals forming a closed pyramid analogous to the larval stage of the crinoids. Anal opening is lateral, in the interradio-oral suture, and the arms usually two to the ray. Column round, no sign of the elliptic columnals having yet appeared.

LYONICRINUS new genus

Plate 11

Pro *Coccocrinus bacca* Roemer, Sil. Fauna Westl. Tennessee, 1860, p. 51.—Cf. *Coccocrinus* Joh. Müller, in Wirtgen und Zeiler, Verh. Naturh. Verein Rheinh., 12, 1855, p. 20.—Zittel, Handb. Pal., 1, 1879, p. 347.—Wachsmuth and Springer, Rev. Pal., pt. 2, 1881, p. 58; pt. 3, 1885, p. 114; N. A. Crin. Cam., 1897, p. 738.—Bather, Treatise on Zool., pt. 3, 1900, p. 156.—Jaekel, Krinoid. Deutschl., 1895, p. 95; Phylogenie und System, 1918, p. 91.—Zittel-Eastman Textb. Pal., 1913, p. 199.—Bassler, Bibliogr. Index, 1915, p. 248.

Dorsal cup as in the Platycrinidae. Small basal in left posterior position. IBr 2. IIBr more than 2. Tegmen composed of 4 large, triangular interambulacral plates (? orals) surrounding a central space, perhaps for posterior oral, with open clefts between them for ambulacra; plus one large posterior plate flanked by 2 smaller ones supporting a protuberant anal opening. Arms ten, unbranched, uniserial, with quadrangular brachials.

Genotype. *Coccocrinus bacca* Roemer.

Distribution. Silurian; America.

This genus is founded upon the species *Coccocrinus bacca* Roemer, of the Tennessee Silurian. Much has been written about the genus *Coccocrinus*, and it still remains illy under-

stood, partly owing to scarcity of material, and partly to the inherent obscurity of all the early Platyocrinidae because the dorsal cup is so simple and alike in so many genera. It has been shifted about in classification; but most of the later authorities agree in placing it among the Platyocrinidae along with *Culicocrinus*. From this Jaekel in his work on Phylogenie und System, 1918, dissents, and while placing *Culicocrinus* among the Platyocrinidae greatly restricted, transfers *Coccoocrinus* to an entirely different order, and ranks it in a suborder Hyocrinites along with Mesozoic and Recent forms such as *Eudesicrinus*, *Hyocrinus*, *Ptilocrinus*, etc.

The typical species, *C. rosaceus*, described by Roemer in 1844 as *Platyocrinus*, was made by Johannes Müller the type of *Coccoocrinus* in 1855. It has a tegmen constructed mainly of five large triangular plates, interlocking and forming a pyramid somewhat resembling that of *Haplocrinus*, having grooves along their edges; but with this difference, that whereas in *Haplocrinus* the ventral pyramid rests directly upon the shoulders of the radials, in *Coccoocrinus* there is a set of plates interposed called "suborals," interbrachial in position, on which the triangular plates rest in direct succession. The anus is at the line of junction between the posterior interbrachial and the triangular plate, each being notched for the opening. The structure is well shown by figures 1 and 2 on plate 11, the first being from what is probably the original of Schultze's figure in his monograph of the Eifel Echinoderms, and the last from the specimen figured by Wachsmuth and Springer in the *Camerata* monograph, pl. 3, fig. 14, obtained by me at Gerolstein in 1887. There is a slight difference in the contour of the tegmen of the two specimens, and in the appearance of the grooves, partly due to difference in preservation—the last being somewhat crushed—but the essentials are the same. It is a Middle Devonian species.

The nearest allied form to *Coccoocrinus* is *Culicocrinus*, described by Johannes Müller from the Lower Devonian, which has likewise a tegmen composed of 5 large orals, but they are closely united by suture, without any groove along their edges; they are also preceded by plates interbrachial in position (pl. 11, figs. 3, 3a, 4, 4a, b, c). The chief difference between the two genera, so far as can be ascertained from the material available, is said to be that *Culicocrinus* has heavy, biserial arms, while in *Coccoocrinus*, the arms are supposed to be uniserial and rather delicate, and that the former has the lower brachials more deeply incorporated in the cup than the latter.

The tegmen of *Culicocrinus* is paralleled by that of *Myrtillocrinus* Hall, from the Onondaga of New York, in being composed of closely interlocking oral plates, but without any underlying interbrachial structures (pl. 11, figs. 5, 5a).

In 1860 F. Roemer described the species *Coccoocrinus bacca*, from the Silurian of Tennessee. It was a rare form, and he had but few specimens, which disclosed nothing of the tegmen; but he saw the beginning of the interambulacral plates, which he and subsequent authors took to be the "suborals" as in *C. rosaceus*. Little addition to his knowledge was obtained by later collectors, Wachsmuth in his Tennessee excursions finding but four specimens, one of which preserved some of those plates, but not intact. This was figured in the *Camerata* monograph, pl. 75, figs. 15a-c. It has been assumed that the interbrachially situated plates here, as in *Coccoocrinus*, were followed by a pyramid of triangular plates substantially covering the ventral side. Much of the discussions of the generic position of *Coccoocrinus* hitherto has been based on this species, now here separated.

Among the collections made for me in Tennessee in 1906-7 is an extraordinary lot of material of this species from which considerable new information has been derived. There are in all about 1,000 specimens, from two colonies not far apart. They occurred mostly in a shaly and limestone deposit disintegrating into clay, forming part of the lower member of the Beech River formation designated by Pate and Bassler as the *Coccoocrinus* zone. The specimens chiefly consisted of the calyx only, but a few were found with more or less of the arms attached, uniserial, and ten in number. Many of the specimens have the interambulacral

tegmen plates in good condition, together with some additional structures connected with the anus, and traces of ambulacra.

In numerous examples the base is well exposed, so that the position of the small basal can be observed. Jaekel and Bather have stated that this is not a reliable character for the genus, and Jaekel declares it is not constant in *C. bacca*. These specimens show that it is thoroughly constant for the species, but that its position is not in accordance with the general rule in crinoids with an unequally tripartite, monocyclic base, namely, left anterior (or 7 o'clock). On the contrary it is here uniformly at the left posterior (10 o'clock, pl. 11, figs. 8, 21, 22), whereas in *C. rosaceus*, as shown by my specimens, it is normal. The former appears to be the case in the specimen figured in the Camerata monograph, pl. 75, figs. 15a-c; but through misunderstanding a note was inserted in the explanation of fig. 15a stating that it is drawn with the anal side at the left lower corner (which would make it according to rule), whereas it is actually at the right lower corner. Fig. c, the posterior view, shows this by the position of the interbasal suture, which would be to the right of the interradian suture if the explanation was correct.

Now the tegmen of *C. bacca* appears to be considerably different from that of the typical *C. rosaceus*. Instead of being suboral plates succeeded by the large triangular plates, as in that species, and as heretofore supposed in this, the interambulacra are themselves almost triangular, the lower face obtusely angular to fit directly into the notch formed by the sloping shoulders of the radials, while above they are extended to an acute angle, so that when in normal position they would form a low pyramid. But instead of interlocking at the apex they leave a considerable open space in the middle, and are also not in contact laterally, but are parted by open clefts running between their edges to the arm bases, along which traces of side pieces of ambulacra can occasionally be seen (pl. 11, figs. 6 to 18).

The large tegmen plates are analogous in position to orals, and that is probably the proper term for them. Four of them are of like shape and size, but the fifth remains to be accounted for. The central vacant space seen in all the specimens is of the proper size and position to lodge the posterior oral, pushed in between the other four by the anal structures, as in the tegmen of many Camerata illustrated by Wachsmuth and Springer on plate 3 of their monograph, and precisely as seen in that of *Culicocrinus* figured herein on plate 11, figure 4c. It is evident that this tegmen was of rather fragile construction, as the central plate is invariably wanting, and the ambulacra only represented by traces. Thus the tegmental structure differs also from that of *Haplocrinus* and *Culicocrinus*, in the fact that the plates of the ventral pyramid in those genera are closely joined, those of *Culicocrinus* without any grooves, and those of *Haplocrinus* with superficial grooves at their edges, not for ambulacra, but as mere receptacles for the arms, as herein now shown for the first time in *H. mespiliformis* (pl. 25, figs. 28, 29, 29a. See also Devonian Crinoids of New York, pl. 40, figs. 10-14).

There is also a very material difference in the anal structure of the two forms. In *Coccocrinus* there is a simple opening at the base of the posterior oral. In *C. bacca* the anus is at the end of a short protuberance, of which we often find one plate, and sometimes one or two more succeeding it, in place above a large posterior plate flanked by a smaller plate at either side. In one specimen four plates in this series are visible, leading to the opening. The two flanking plates served as a strong support, for we frequently see them, and the intervening posterior plate standing firmly upright, while all other plates of the tegmental arch have fallen down (pl. 11, figs. 10-20).

The evidence is complete to prove that the structure of *C. bacca* is different from that of *C. rosaceus* in three major characters, viz.: the composition of the tegmen, the anal structure, and the position of the small basal. From this it follows that they cannot be retained in the same genus. Accordingly I am proposing for the Tennessee species the new genus *Lyonicrinus*, in memory of Col. Sidney S. Lyon, who was an ardent student of the crinoids, and author of many notable genera and species from the Kentucky-Indiana area.

Lyonicrinus bacca (Roemer)*Plate II, figs. 6-23*

Coccocrinus bacca F. Roemer, Sil. Fauna Westl. Tennessee, 1860, p. 51, pl. 4, figs. 5a-c.—Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 739, pl. 75, fig. 15.—Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 29, pl. 4, figs. 9.—Bassler, Bibliogr. Index, 1915, p. 248.

The species is sufficiently discussed in the remarks upon the genus.

Horizon and locality. *Coccocrinus* zone of Beech River formation, Brownsport group, Niagaran; Decatur and Perry counties, Tennessee.

Coccocrinus rosaceus (F. Roemer)*Plate II, figs. 1, 1a, b; 2*

Platycrinus rosaceus Roemer, Rhein. Ueberg., 1844, p. 63, pl. 3, fig. 3.—*Coccocrinus rosaceus* Joh. Müller, Verh. Naturh. Verein Rheinf. 12, 1855, p. 21, pl. 7, figs. 5a-c.—Schultze, Echin. Eifl. Kalk., 1867, p. 89, pl. 12, fig. 13.—Wachsmuth and Springer, N. A. Crin. Cam., 1897, pl. 3, fig. 14.

For comparison with *Lyonicrinus bacca*. Middle Devonian; Eifel, Germany.

Culicocrinus nodosus (Wirtgen and Zeiler)*Plate II, figs. 3, 3a*

Platycrinus nodosus Wirtgen and Zeiler, Verh. Naturh. Verein Rheinf. 12, 1855, p. 15, pl. 6, figs. 23.—*Culicocrinus nodosus* Joh. Müller, *ibid.*, p. 24, pl. 8, figs. 1-4.

See discussion under *Lyonicrinus*.

Lower Devonian; Coblenz, Germany.

(?) Culicocrinus spinosus new species*Plate II, figs. 4, 4a-c*

This name is given to a solitary specimen from Tennessee, with much doubt as to its generic affinities, but chiefly on account of the five large spiniferous interlocking orals, which differ from those of the typical species in having the posterior plate flanked on the outer side by anal structures and smaller lateral plates, but nevertheless present considerable similarity in proportion and general arrangement. The primibrachs are also less deeply incorporated in the cup.

Beech River formation, Brownsport group, Niagaran; Decatur County, Tennessee.

Platycrinidae indet., No. 1*Plate II, figs. 24, 24a, b*

A dorsal cup with massive plates, and large BB. Beech River formation; Decatur County, Tennessee.

Platycrinidae indet., No. 2*Plate II, fig. 25*

Calyx with large interambulacral plates in position. Laurel limestone; St. Paul, Indiana.

Platycrinidae indet., No. 3*Plate 11, figs. 26, 27, 28, 29*

A small species rather common in the Laurel limestone at St. Paul, Indiana.

The three foregoing species are figured in order to preserve a record of their occurrence, hoping that future discoveries may throw more light upon them.

Hapalocrinus Jaekel emend. Bather*Plates 11, 12*

Hapalocrinus Jaekel, Krin. Deutschl., 1895, p. 37.—Bather, Geol. Mag., 4, 1897, p. 342; Treatise on Zool., 3, 1900, p. 156.

Platycrinidae with 2 or more IBr; orals small; arms uniserial with cuneate brachials tending to biserial, branching once or twice; column large, bearing strong cirri.

Genotype. *Hapalocrinus elegans* Jaekel.

Distribution. Silurian to Lower Devonian; England, Gotland, Germany, America.

The specimens which I have referred to this genus are notable for their profuse development of arms and pinnules, and of strong cirri occurring in definite whorls, at intervals extending apparently the full length of the column. The generic characters of these forms are not very certain. Some of them might perhaps better be placed under *Cordylocrinus* of Angelin, but the distinctions are hard to point out with confidence, and I have chosen the easier course by throwing them all under the one form, which is strongly typified by the "*Platycrinus*" *retiaris* Phillips of the English Silurian. This arrangement would seem to include both uniserial and biserial arms—a variation that may occur among the later Platycrinidae within the same species, or even the same individual, incident to growth, as seen for example in *P. huntsvillae*. A very strongly marked species from the Lower Devonian is figured for comparison.

Hapalocrinus gracilis new species*Plate 11, figs. 30, 31, 32*

With heavy cuneate arms and strong pinnules, and rather high tegmen plates.

Hapalocrinus cirrifer new species*Plate 12, figs. 1, 2, 3, 4*

Remarkable for the long, slender arms, in some specimens almost thread-like; the calyx is large, but details of its structure are obscure.

Hapalocrinus pinnulatus new species*Plate 12, fig. 5*

Calyx narrow and relatively smaller, but bearing much stronger arms and pinnules.

Hapalocrinus tuberculatus new species*Plate 12, fig. 6*

Arms and pinnules similar to the preceding, but calyx rotund, and with tubercular surface.

Hapalocrinus tennesseensis new species*Plate 12, fig. 7*

Crown much displaced, with sharply uniserial arms and quadrangular brachials.

All the foregoing five species from the Beech River formation, Brownsport group, Niagara; Decatur County, Tennessee.

Hapalocrinus devonicus new species*Plate 12, fig. 8*

Unusually strong stem and cirri, large calyx, and tubercular surface.

Keyser formation, Helderbergian, Lower Devonian; Keyser, West Virginia.

Brahmacrinus elongatus new species*Plate 12, figs. 9-15*

The genus to which this species is referred was proposed by Sollas,¹ to receive a Platycrinid form from the British Lower Carboniferous of true Camerate type, having the lower brachials completely incorporated in the cup by means of solid interbrachials. Some of the earlier American species of similar type, such as *P. eriensis*,² might well be placed under it, and it affords a practicable resting place for the present singularly distinct form.

It may be defined as a Platycrinid with large IBr and IIBr incorporated in the cup by iBr in two ranges, huge BB, and relatively smaller RR than in *Platycrinus*. Along with this it is remarkable for having the base divided into 3 subequal plates, which are sharply excavate into a saucer-shaped depression surrounding a rather small stem. This has a minute axial opening, which is a Platycrinid characteristic, but there is no sign of elliptic columnals. The peculiar excavate base is seen in several species of *Platycrinus*—such as *P. huntsvillae* and others shown in N. A. Crin. Cam., pl. 70, fig. 12; pl. 73, figs. 11, 12.

In the present species the elongate, rather cylindrical, form of the calyx is different from that of the Platycrinidae generally, as is also the great size of the basals. Apparently there are two arms to the ray, which probably branch again. There are two good specimens of the calyx, and several bases.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

¹ Quart. Jour. Geol. Soc. London, 56, 1900, p. 264, pl. 16, figs. 1, 2.

² N. A. Crin. Cam., 1897, p. 676. See also as to the British form, p. 644.

INSERTAE SEDIS

Nyctocrinus magnitubus new genus and species*Plate 12, figs. 16-21*

Among the Tennessee collections there is a form abundantly represented by well preserved specimens, of which after frequent reviews I have been unable to fix the systematic position. It is a dicyclic crinoid of Camerate aspect, with 5 IBB, suggesting the Dimeroocrinidae; but it has the RR in contact all around, with no anal in the cup, which excludes that family as well as the Rhodocrinidae. The dicyclic base excludes the Melocrinidae, and other Camerate families that might be suggested. Except for this, there would be a somewhat near approach to the Platycrinidae. There are large RR, with one or two IBr, filling only part of the distal face, leaving well defined shoulders supporting large iBr chiefly above the line of RR (perhaps more properly tegminal iAmb.), of which the posterior one seems to be rather the largest. Basals are large, curving into a shallow concavity, in which the 5 IBB form a broad, inverted cone. Arms are stout, biserial, 2 or 4 to the ray with occasionally another bifurcation. The most remarkable character is a long, spinose anal tube, which projects beyond the limits of the arms.

Except for the well defined interbrachial plates resting on the radials, this form might fall under some of the dicyclic Inadunata; but with our present knowledge it may as well be left in an intermediate position. In addition to the five specimens figured in which more or less of the arms and anal tube are shown, and the two giving perfect internal and external views of the base, there are half a dozen calices in which the same type of dicyclic base clearly appears—so that the definition of characters here given may be regarded as beyond question. It is a singular fact that all the specimens in which the tube is shown differ from all others of the formation, even those in the same matrix, in being very dark in color—which suggests the generic name.

Beech River formation, Brownsport group, Niagaran; Decatur County, Tennessee.

Section 2. Subfamily MARSIPOCRININAE Bather

Rays with only one primibrach.

The next subdivision, with rays limited to a single primibrach, is abundantly represented by its earliest genus, remarkable for the manner in which its brachials to the axillary secundibrach are buried in the cup, so as to produce a thoroughly Camerate structure, and for the constancy with which its characters are maintained throughout numerous species, now to be described.

MARSIPOCRINUS Bather*Plates 13-19*

- Marsupiocrinites* Phillips (not de Blainville, 1830), in Murchison's Sil. Syst., 1839, p. 672.—Austin, Ann. Mag. Nat. Hist., 10, 1842, p. 109.
- Marsupiocrinus* McCoy, British Pal. Rocks and Foss., 1854, p. 54.—Wachsmuth and Springer, Rev. Pal., 2, 1881, pp. 63, 230; N. A. Crin. Cam., 1897, p. 730.—Weller, Bull. Chicago Acad. Sci., 4, pt. 1, 1900, p. 137.
- Marsupiocrinus* Bather, Quart. Jour. Geol. Soc. London, 45, 1889, p. 173; Treatise on Zool., 3, 1900, p. 156, fig. 70.—Bassler, Bibliogr. Index, Am. Ord. Sil. Foss., 1915, p. 786.
- Cupellaecrinites* Troost, Am. Jour. Sci., 1849, p. 419; Proc. Am. Assn. Adv. Sci., 1850, p. 61 (not defined).—*Cupellaecrinus* Meek and Worthen (not Steininger), Proc. Acad. Nat. Sci. Phil., 1865, p. 161; Geol. Surv. Illinois, 2, 1866, p. 172.—Shumard, Trans. Acad. Sci. St. Louis, 2, 1866, p. 361; p. 387, footnote.—Meek, Am. Jour. Sci., 1866, p. 118.

Platycrinidae with one IBr, small, axillary, and IIBr on either side of it resting on RR and wholly incorporated in dorsal cup, followed by another axillary in case of further division; one large iBr supported by shoulders of RR, flanked by smaller ones at upper corners which may be restricted to the tegmen; arms stout, biserial, two or four to the ray, with large pinnules; tegmen composed of numerous Amb. and iAmb. plates; anus subcentral, directly through the tegmen, without a tube; orals small, unsymmetric, pushed anteriorly, the posterior plate between the other four; stem circular in section, lumen large, usually pentagonal or pentapetalous; small basal at l. ant. side.

Genotype. *Marsupiocrinus coelatus* Phillips.

Distribution. Silurian to Lower Devonian; England, Sweden, America.

The name *Marsupiocrinus*, proposed by Phillips for the English species, was preoccupied by de Blainville. Although Troost's Ms. name, *Cupellaecrinites*, given for the form afterwards described by Roemer as *Platycrinus tennesseensis*, was adopted and published by Meek and Worthen and by Shumard, it is held invalid by most authors for conflict with *Cypellaecrinus* of Steininger. Bather's modification of Phillips's name, proposed to remedy the existing confusion, has been generally accepted.

While the European Silurian has furnished some excellent examples for illustrating the structure of the genus, especially those figured by Angelin from Gotland, the little that has been published in regard to it in America has been far from commensurate with its importance as now known. This has been due to the extreme rarity of specimens heretofore available. Wachsmuth and Springer for the Camerate monograph had very meagre material, either of our own or in other collections. We had only two specimens with the tegmen preserved,

which were considered quite wonderful, and the Troost collection which has since become accessible was not much better supplied. The few calices that had been picked up on the glades consisted mostly of the dorsal cup, or of the basal plates only, with the surface more or less smoothed by erosion, and such as were found were usually assumed to belong to Roemer's species, *M. tennesseensis*. The publication later of Troost's monograph (Bull. 64, U. S. Nat. Mus., 1909) indicated the existence of a wider range of forms than had been previously known, and brought out a number of new specific names, but the preservation of the specimens was for the most part poor, and not much aid was given toward fixing the diagnostic characters of the species. The elaborate original descriptions were of specimens rather than of species.

The collection now assembled furnishes for the first time adequate material for the study of the American forms of the genus, which seems to have flourished profusely in the Tennessee area. Most of it was derived from the shales of the *Eucalyptocrinus* zone of the Beech River formation. These yielded numerous specimens from the soft matrix, in exquisite preservation, admitting the most delicate preparation, by which the surface ornament, details of tegmen, and in many cases the arms and pinnules, were brought to light with a perfection never before seen in this country. The study of this collection has resulted in the recognition of ten species belonging to this field alone, of which three are new, while five of those proposed by Troost are confirmed by ample evidence among the new material. Thus we are able to place the leading species upon a firm footing, and to bring out in detail the great diversity by which the genus is characterized upon a well defined structural plan. In other areas a species has been described by Weller from the Racine dolomite at Chicago, and one by Hall from the Ohio Silurian, as well as from the Helderbergian of New York, by which the range of the genus is extended into the Devonian. Of the European Silurian there is a fine species from England, and four from Gotland described by Angelin, some of which may be synonyms. These forms for the most part differ consistently from the American species. A notable fact is the extreme rarity of this genus in the Laurel formation of St. Paul, where its presence is barely indicated by a few fragments.

No genus of fossil crinoids is more clearly defined than *Marsipocrinus*. The general form, with broadly arched tegmen, and the peculiarity by which the secundibrachs rest directly upon the radials and are thus, as well as by contact with the large interbranchials, incorporated in the cup, impart a certain physiognomy by which it is readily recognized. Typically the calyx is low and broad in the proportion of 1 to 3, or 1 to 2, with somewhat less difference in the English species. From others of the family it is also decisively set off by its many-plated tegmen and large stem-lumen.

Within the genus there are several prominent specific characters by which the limits of the species are fairly well defined:

1. Form of calyx, which may be either—*a*, concavo-convex or hemispheric, with radial facets directed below the horizontal, and tegmen exceeding dorsal cup in size; or *b*, bi-convex, with radial facets intermediate and tegmen about equaling dorsal cup; or *c*, convex conical; dorsal cup deep convex, tegmen low conical, constricted, narrower than cup.

2. Basal pentagon flat or convex with raised margin, to concave with basals at bottom of cavity.

3. Interbranchial plate projecting with corrugated arch, or smooth without projection.

4. Tegmen either—*a*, sharply tubercular, or with raised, angular, diamond-shaped plates, and ambulacra not conspicuous; or, *b*, smooth, or with rounded plates, and ambulacra well defined.

5. Surface markings either—*a*, coarse granular; *b*, corrugated, with coarse wrinkles or tubercles; *c*, tubercles confluent forming more or less interrupted lines or coarse striae part way on plates; *d*, fine striae traversing the plates; *e*, entirely smooth.

6. Number of arms, 2 to the ray as in most American species, or 4 to the ray as usual in the European species.

Marsipocrinus rosaeformis (Troost)*Plate 13, figs. 1-8; 15, figs. 9, 9a*

Cupellaecrinites rosaeformis Troost, Am. Jour. Sci., 1849, p. 419; Proc. Am. Assn. Adv. Sci., 1850, p. 61 (nom. nud.).—*Cupellaecrinus rosaeformis* Shumard, Trans. Acad. Sci. St. Louis, 2, 1866, p. 361.—*Marsipocrinus rosaeformis* (Troost) Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 37, pl. 13, figs. 5, 6.

A large species, typical specimens 12 mm. height of calyx by 30 mm. width, to a maximum of 15 by 40 mm. Calyx concavo-convex or hemispheric; if resting on a plane surface will touch by radials or arm-bases. Basal pentagon flat, relatively small, about one third the total diameter, usually with a small rim around the column-facet. RR horizontal, with facets directed out or downward; axillary IBr very small, triangular, not over one third the width of the radial, flanked by large transversely elongate IIBr which rest upon the radial and meet by their short inner faces above the axillary; they curve around on the outer side, abutting upon the large iBr and the smaller ones above, and are followed by higher brachials leading to the biserial arms. iBr large, arched, projecting in a curved lip outward and upward, often strongly corrugated or wrinkled, giving to the calyx a decagonal outline as seen from above; a much smaller plate abreast of the iBr at either side rests upon the IIBr, sometimes limited to the tegmen. Arms stout, biserial, two to the ray. Tegmen broadly arched, larger than dorsal cup; plates strongly nodose, shading to angular in some specimens, somewhat obscuring the ambulacra, which are not conspicuous. Surface ornament of cup very strong, either in form of a simple corrugation with wrinkles or tubercles not forming lines, or more or less confluent in radiating lines forming triangles, or parallel striae.

This is the most prominent species in the Tennessee collections, represented by many fine specimens mostly obtained from the excavations. Comparison of Troost's type specimen in the National Museum, as figured in 1909, although in poor condition, established that the hemispheric calyx is the typical form, and enables us to assign to the species the other characters so fully illustrated by the new material. The specimen figured herein, pl. 15, figs. 9, 9a, reproduced from my work on *Scyphocrinus*, 1917, pl. 9, figs. 5, 5a, as *M. tennesseensis*, is probably referable to this species.

Horizon and locality. Chiefly *Eucalyptocrinus* zone of the Beech River formation, Brownsport group, Niagaran; Tuck's Mill, near Decaturville, Decatur County, Tennessee.

Marsipocrinus tennesseensis (Roemer)*Plate 14, figs. 1-6*

Platycrinus tennesseensis Roemer, Sil. Fauna Westl. Tennessee, 1860, p. 35, pl. 3, figs. 4a-f.—*Platycrinus* (*Cupellaecrinus*) *tennesseensis* Shumard, Trans. Acad. Sci. St. Louis, 1866, p. 362.—*Marsupocrinus tennesseensis*, Wachsmuth and Springer, Rev. Pal., 2, 1881, p. 65; Proc. Acad. Nat. Sci. Phil., 1888, p. 373, pl. 20, fig. 7; N. A. Crin. Cam., 1897, p. 731, pl. 75, figs. 16a, b.—Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 31, pl. 9, figs. 5, 6.—Bassler, Bibliogr. Index, 1915, p. 788.

Similar to *M. rosaeformis* in size, general proportions and form of iBr plate, but with calyx bi-convex, and dorsal cup and arched tegmen about equal.

Basal pentagon proportionally larger; surface ornament similar but usually less pronounced. Radial facets directed above the horizontal. Tegmen plates raised, somewhat nodose, angular, or smooth; ambulacra fairly defined.

Roemer's *P. tennesseensis* as described by him contains two species, of which his figures 4a, b, c, must be taken for the type. His specimen was weathered, and surface of both cup and tegmen probably eroded. But the characters above specified are sufficient to identify the species, of which Wachsmuth and Springer figured a well preserved example (N. A. Crin. Cam., pl. 75, fig. 16, but not fig. 17).

Horizon and locality. Beech River formation, Brownsport group, Niagaran; *Eucalyptocrinus* zone and glades, Decatur County, Tennessee.

Marsipocrinus striatus (Wachsmuth and Springer)

Plate 14, figs. 7-13

Marsupiocrinus striatus Wachsmuth and Springer, N. A. Crin. Cam., 1897, p. 732, pl. 75, fig. 18 only. Not *Cupellaecrinus striatus* Troost, nor *Marsipocrinus striatus*, Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 33, pl. 9, figs. 13, 14 = *M. inflatus*.

A somewhat smaller species than the preceding, varying from 12 to 14 mm. height of calyx to 15 to 26 mm. diameter. Calyx bi-convex, dorsal cup usually higher than tegmen. Basal pentagon shallowly depressed, about one half the total width. Radial facets directed upwards. iBr usually not projecting, smooth, curving over edge of tegmen. Tegmen low, plates rounded, smooth and but little raised, leaving ambulacra well defined. Surface of cup usually striate, continuous from plate to plate, sometimes rather corrugate with striae more obscure—much dependent upon preservation.

A considerable number of specimens answer to this description, and agree fairly well with the type of Wachsmuth and Springer, pl. 75, fig. 18. It is a variable form, tending to shade into other species, but the strong striate ornament and prominent exposure of ambulacra are the features chiefly relied on. In specimens from the glades the striae are usually much obscured by weathering, for which allowance must be made.

Horizon and locality. Beech River formation, Brownsport group, Niagaran; Tuck's Mill and on glades, Decatur County, Tennessee.

Marsipocrinus inflatus (Troost)

Plate 15, figs. 1-8

Cupellaecrinus inflatus Troost, Proc. Am. Assn. Adv. Sci., 1850, p. 61.—*Marsupiocrinus striatus*, Wachsmuth and Springer, N. A. Crin. Cam., 1897, pl. 75, fig. 17.—Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 34.

A fairly large species, with calyx relatively high, the average being about 20 mm., by 30 mm. wide. Dorsal cup convex and rather deep; basal pentagon large, about three fifths the total diameter, margins usually notched at the inter-basal sutures, and sometimes with slight protrusions at the middle of the sides. iBr not projecting, but flush with calyx wall and curving over into the tegmen.

Arms stout, two to the ray, with pinnule openings between their bases sometimes seen. Tegmen low conical, abruptly constricted above the arm-bases, and distinctly narrower than the cup; plates flat or low convex, smooth or studded with small tubercles; ambulacra not conspicuous. Surface of cup ornate, usually corrugate, sometimes with ridges forming striae, or lines parallel to margin of plates; sutures often sharply beveled.

A well marked species, usually readily recognized by the high calyx, large basal pentagon, and a certain massiveness not apparent in most other species. The constricted tegmen is very decisive, but this is seldom preserved, and with only the dorsal cup the large pentagon and vertical sides are usually sufficient. In well preserved specimens the sculpturing is highly ornate. This is clearly the form described by Troost under the names *C. striatus* and *C. inflatus*, as shown by his type of the latter, and the figures of the former in the monograph of 1909, pl. 9, figs. 13, 14. The first name being preoccupied by Wachsmuth and Springer, the last should stand for the species. A specimen of it was figured by us as *M. tennesseensis* (N. A. Crin. Cam., 1897, pl. 75, fig. 17 only).

The enlarged details of a specimen of this species shown by figure 8 of plate 15, along with another which may be of *M. rosaeformis*, are reproduced from my works on *Scyphocrinus*, 1917, pp. 33-38, pl. 9, figs. 5a-c, and *Dolatocrinus*, Bull. 115, U. S. Nat. Mus., 1921, p. 24, pl. 2, figs. 3, 4. They were given for demonstrating the true nature and function of the interbrachial apertures observed in various genera of the Camerata, and frequently designated as "respiratory pores," which are now seen to be the openings for pinnules, the ambulacra leading to which are shown in some specimens of other species. For a full discussion of this subject, and the significance of these specimens, reference should be had to those papers, at the pages indicated.

Horizon and locality. Beech River formation, chiefly *Eucalyptocrinus* zone, Brownsport group, Niagaran; Tuck's Mill and glades, Decatur County, Tennessee.

Marsipocrinus verneuili (Troost)

Plate 16, figs. 1-6

Cupellacrinites verneuili Troost, Proc. Am. Assn. Adv. Sci., 1850, p. 61 (nom. nud.).—*Marsipocrinus verneuili* (Troost), Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 32, pl. 9, figs. 10, 11, 12.

A small species, average diameter of calyx about 15 mm. Calyx bi-convex, with shallow concave base, height to width about as 1 to 2. Basal pentagon about half the total diameter, broadly excavate, bounded by a conspicuous, obtusely triangular rim, much larger than the column-facet, flaring outward to a sharp edge. iBr not projecting, bent over into the tegmen. Arms tapering, two to the ray. Tegmen lower than cup, with low, smoothly rounded plates; ambulacra narrow, well defined, with some outer divisions running to pinnule openings. Surface of cup coarse striato-corrugate or rugose.

This species as represented by Troost's type is well defined, and is confirmed by seven specimens in the present collection. The flaring rim around the column is a striking character, seen to some extent in other species, but in none so conspicuous as this.

Horizon and locality. Beech River formation of Brownsport group, *Eucalyptocrinus* zone, Niagaran; Tuck's Mill, Decatur County, Tennessee.

Marsipocrinus excavatus new species*Plate 16, figs. 7-7b*

Calyx plano-convex, with basal pentagon abruptly sunken around a small, round column-facet. iBr projecting in small points. Tegmen conical, higher than cup; plates low, covered with small tubercles. Surface markings on cup coarse, with irregular strong wrinkles and obscure striae. Founded upon a single specimen 13 mm. high by 28 mm. wide, with basal pentagon about one third the total diameter.

Horizon and locality. *Eucalyptocrinus* zone of Beech River formation, Brownsport group, Niagaran; Tuck's Mill, Decatur County, Tennessee.

Marsipocrinus concavus new species*Plate 16, fig. 8*

Calyx with deep funnel-shaped basal cavity, involving part of the radials. Basal pentagon very small, almost covered by column at bottom of cavity, bordered by a thin rim. Radials long, extending far down into the basal cavity. Surface ornament strong, composed of some irregular striae, and concentric ridges roughly parallel to margins of pentagon. Tegmen and other parts unknown.

A thoroughly distinct species, differing from all others as *Pisocrinus quinquelobus* differs from species with large basals. Unfortunately represented by only a single imperfect specimen consisting of basals and radials.

Horizon and locality. *Eucalyptocrinus* zone of Beech River formation, Brownsport group, Niagaran; Tuck's Mill, Decatur County, Tennessee.

Marsipocrinus stellatus (Troost)*Plate 17, figs. 1-10*

Cupellaecrinites stellatus Troost, Am. Jour. Sci., 1849, p. 419; Proc. Am. Assn. Adv. Sci., 1850, p. 61 (nom. nud.).—*Cupellaecrinus stellatus* Shumard, Trans. Acad. Sci. St. Louis, 1866, p. 361.—*Marsipocrinus stellatus* (Troost) Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 36, pl. 10, figs. 1, 2.

A rather small species, with dorsal cup perfectly smooth, ranging from 8 and 10 mm. height of calyx to 16 and 24 mm. width. Calyx bi-convex, broadly spreading, with base flat or slightly convex. Basal pentagon about two fifths the total diameter; interbasal sutures often obscure. iBr with rather sharp projecting lip. Arms directed horizontal, long and slender toward the extremities, two to the ray. Tegmen broadly rounded, smooth in interambulacral regions, or with scattering sharp nodes; ambulacra conspicuous from middle to arm-bases, with some outer branches leading to pinnule openings. Surface of cup entirely devoid of ornament. Stem stout, tapering distally dividing into terminal branches for attachment.

The smooth surface in this species is not a matter of erosion, but is a thoroughly definite character, as is shown by at least a dozen specimens in the collection, mostly well preserved

in the matrix and several with arms and stem attached, in none of which is there any trace of surface markings. The tendency is toward ankylosis of the basals. Along with these characters the low and broadly spreading calyx makes the species readily recognized.

Horizon and locality. *Eucalyptocrinus* zone of Beech River formation, Brownsport group, Tuck's Mill, Decatur County, Tennessee.

Marsipocrinus striatissimus new species

Plate 18, figs. 1-7

A very large species, attaining a diameter of 50 or 60 mm., with extremely thin plates, so that specimens are rarely found undistorted, or with anything but the basal pentagon preserved. Calyx low and broadly spreading, with arm facets directed about horizontal. Base convex, with narrow rounded rim about column facet. iBr projecting but little, mostly smooth. Tegmen low, usually crushed, with plates level, smooth, or covered with small nodes or tubercles; ambulacra not conspicuous. Surface of cup marked by an exceedingly fine striation, with straight lines crossing the sutures from plate to plate.

Roemer's figure 4d may be of this species, and it was also this which Wachsmuth and Springer had in mind when in describing their much smaller and more robust *M. striatus* they stated that it sometimes reached 6 cm. in width. It is not uncommon on the glades as fragments only, characterized by large size and the extreme thinness of the plates. The crushed specimen figured on plate 18, figures 1, 1a, is the nearest to a complete calyx that is known. Two fairly robust dorsal cups have the characteristic delicate striation, while in the specimen with arms it is almost obliterated by pressure.

Horizon and locality. *Eucalyptocrinus* zone of Beech River formation, Brownsport group, Niagaran; Tuck's Mill and glades, Decatur County, Tennessee.

Marsipocrinus magnificus (Troost)

Plate 19, figs. 1-1b

Cupellaeocrinites magnificus Troost, Proc. Am. Assn. Adv. Sci., 1850, p. 61 (nom. nud.).—*Cupellaeocrinus magnificus* Shumard, Trans. Acad. Sci. St. Louis, 2, 1866, p. 361.—*Marsipocrinus magnificus* (Troost) Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 34, pl. 10, figs. 5-7.

A very large and robust species, with four arms to the ray. Calyx bi-convex; height 27 mm. width 40 mm. Base about flat, with a thickened rounded rim surrounding the column-facet; basal pentagon about three fifths the diameter of calyx. Two secundibrachs in succession meet above the triangular primibrach, the second one axillary, followed by a second division of the ray, giving four arm openings instead of two as in all the preceding species. iBr very large, broadly shield-shape, truncate above, and but little projecting. Tegmen composed of strong convex or nodose plates, with ambulacra not conspicuous. Surface of cup deeply sculptured, with beveled sutures, and coarse tubercles coalescing into prominent ridges to the corners of the basal pentagon, and from there across the radials, forming conspicuous triangles with apex at the primibrach.

Known only from the fine and perfect calyx of the type specimen in the Troost collection, and distinguished from all other known American Silurian species by having four arms to the ray, as in the English and Swedish species. But it is remarkable that whereas in the latter the interbrachial plate is compressed by the increased number of arms so that it is elongate and pointed above, in the present species it is wide and broadly truncate above as in the 2-armed forms, the expansion at the arm bases being accommodated by the great convexity of the tegmen. In fact, aside from the number of arms this species bears not the slightest resemblance to the European forms. That other species with 4 arms existed in the Tennessee area is indicated by two fragments from Wayne County, insufficient for description.

Horizon and locality. Brownsport group, Niagaran; Decatur County, Tennessee.

Marsipocrinus turbinatus new species

Plate 17, figs. 11-13

Among all the collections from the Laurel formation at St. Paul the four fragmentary bases here figured are the only specimens found belonging to this genus. They are of a wholly different type from all the other known species, with their turbinate base indicating a more or less elongate calyx; but the pentagonal form, with the rim surrounding the column-facet, and the large lumen, would seem to settle the generic affinities of the species.

Horizon and locality. Laurel formation, Niagaran; St. Paul, Indiana.

Marsipocrinus coelatus Phillips

Plate 19, figs. 2-7

Marsupiocrinites coelatus Phillips, Murchison's Sil. Syst., 1839, p. 672, pl. 18, fig. 1.—Austin, Ann. Mag. Nat. Hist., 1842, p. 109.—Bather, Lankester Zool., pt. 3, p. 157, fig. 70.

A medium sized species. Calyx convex-conical, ranging from 7 to 15 mm. height to 12 to 30 mm. width. Dorsal cup rather deep, broadly rounded below; basal pentagon nearly flat, sometimes excavate, about one third the total width. RR mostly horizontal in mature specimens; IBr rather large, triangular; IIBr 2 in series, the second one axillary for a second division, followed by 2 or 3 brachials passing into the biserial arms. Arms 4 to the ray, directed vertically, fairly stout. iBr elongate, narrowing to an apex between the rays, almost completely enveloped by the lower brachials and incorporated in the dorsal cup. Tegmen low conical or rounded, composed of numerous small plates, often covered by a gastropod commensally attached over the anus; ambulacra not prominent, but distinctly marked by small covering plates, with outer branches running to interbrachial pinnule openings. Surface of cup striato-corrugate, marked in well preserved specimens by strong ridges crossing the sutures, but more often obsolete. Stem stout and apparently fixed.

An abundant and well known species of the English Silurian, here figured for comparison; it has been best illustrated heretofore by Bather in the Lankester Zoology, part 3, page 157, figure 70. Specifically it bears little resemblance to any American form, while it is a fine representative of the 20-armed type which seems to have predominated in the European

area, in contrast to that of 10 arms chiefly characteristic of the American rocks, from which it is also sharply distinguished by its elongate, pointed interbrachial plate.

A feature of this species not heretofore mentioned to my knowledge is the tendency to grow in clusters, several individuals in a group being attached by their stout stems to corals or other fixed objects. I am illustrating this by a slab from Dudley, shown by a reduced photograph on plate 19, with nine stems springing from a common base, four of them with the crowns attached and visible, and the others with the crowns imbedded in the matrix. Such a multiple fixation is unusual among crinoids, but here it seems to be a character more than merely incidental, as other specimens are known, one very fine one, for example, in the Dudley Museum, and another in the collection of the late Mr. Charles Holcroft, now in the Museum of Birmingham.

Horizon and locality. Wenlockian group, Silurian; Dudley, England.

Of other described species there are one by Weller from the Racine dolomite, one by Hall from the dolomite at Cedarville, Ohio, one by Hall from the Helderbergian of New York; and four by Angelin from Gotland.

Order FLEXIBILIA Zittel

CRINOIDS WITH A FLEXIBLE CALYX IN WHICH THE LOWER BRACHIALS ARE LOOSELY INCORPORATED INTO THE DORSAL CUP EITHER BY LATERAL UNION WITH EACH OTHER, BY MEANS OF INTERBRACHIALS, OR OF A LOOSE SKIN STUDED WITH CALCAREOUS PARTICLES. ALL PLATES BEYOND THE RADIALS UNITED BY LOOSE SUTURE, AND MORE OR LESS MOVABLE. MOUTH AND TEGMINAL FOOD GROOVES EXPOSED. BASE DICYCLIC, WITH THREE INFRA-BASALS, UNEQUAL. ARMS NON-PINNULATE. STEM ROUND.

In view of the fact that this order of the Crinoidea has recently been treated monographically in my work on the Crinoidea Flexibilia, 2 volumes 4to, Smithsonian Publication 2501, 1920, in which nearly all the species occurring in the areas here under consideration have been fully described and their generic relations discussed, it is not deemed necessary to repeat the descriptions, or to do more than give a list of the genera and species with references to that work where the complete bibliography will be found, together with a selection of the principal figures—for the greater convenience of students of the Silurian faunas.

Of the Flexibilia group there have been recognized from this area 10 genera and 18 species, all but two of the genera from Tennessee. Three of the species are new; 11 are from Tennessee, 6 from Indiana, and one from Ohio. They are distributed among the four families of the order as follows: Lecanocrinidae 12, Sagenocrinidae 2, Ichthyocrinidae 2, Taxocrinidae 2. Thus the great preponderance of this type as found in the present collections belongs to the small, robust and better preserved Lecanocrinidae.

Of the ten genera, nine are common to both Europe and America. Of these, six were formerly considered to be exclusively Swedish and English. One newly described genus is represented by species from both continents. One highly typical Gotlandian and English genus, *Pycnosaccus*, hitherto barely recognized in this country, appeared in the Beech River formation in considerable abundance, one species of which can scarcely be distinguished from its foreign prototype. *Sagenocrinus*, another specialized form of the European Silurian, is represented here by a thoroughly typical species. And a third, *Gnorimocrinus*, now appears for the first time in America. Besides these important contributions from the Tennessee Silurian there are two well defined new genera, *Hormocrinus* and *Asaphocrinus*, in the form of finely preserved specimens, the first of which also occurs in Europe as new species. Thus the Flexibilia, usually the rarest of the crinoids, take their place among this fauna as one of its most important constituents.

Suborder SAGENOCRINOIDEA

Posterior interradius either not differentiated or containing anal plates, not usually formed into a tube, the first of which is incorporated in the calyx by sutural union with adjacent brachials for at least part of its height on both sides; posterior basal, if differentiated, truncate or angular, and sutureally connected with succeeding anal plate.

Family LECANOCRINIDAE Springer

Infrabasals more or less erect, forming an essential part of the calyx wall; crown usually short, rotund.

LECANOCRINUS Hall*Plate 20*

Lecanocrinus Hall, Pal. New York, 2, 1852, p. 199.—Springer, Crinoidea Flexibilia, 1920, p. 125.

Rays in contact except at the anal side. Radial rhombic, obliquely to lower left of r. post. R. Anal x alone. Arms dichotomous, interlocking.

Genotype. *Lecanocrinus macropetalus* Hall.

Distribution. Silurian to Devonian; Gotland, England, America.

Lecanocrinus pusillus Hall*Plate 20, figs. 1-5*

Lecanocrinus pusillus Hall, Trans. Albany Inst., 4, 1863, p. 200.—Springer, Crinoidea Flexibilia, 1920, p. 131, pl. 1, figs. 1-13.

Waldron shale; Waldron, Indiana; Newsom, Tennessee.

Lecanocrinus pisiformis (Roemer)*Plate 20, figs. 6-12*

Poteriocrinus pisiformis Roemer, Sil. Fauna Westl. Tennessee, 1860, p. 54, pl. 4, figs. 7a-d.—*Lecanocrinus pisiformis* Springer, Crinoidea Flexibilia, 1920, p. 135, pl. 1, figs. 14-36.

Beech River formation; Decatur County, Tennessee.

Lecanocrinus elongatus new species*Plate 20, figs. 13, 14*

Similar to *L. pisiformis*, but crown more elongate, and basals relatively higher.

Laurel limestone; St. Paul, Indiana.

Lecanocrinus meniscus Springer*Plate 20, figs. 15-15b*

Springer, Crinoidea Flexibilia, 1920, p. 140, pl. 1, figs. 37a-c.

Beech River formation; Decatur County, Tennessee.

Three species have been described from the Rochester shale, one from the Racine, and one from the Bainbridge of Missouri; and 10 species described as *Lecanocrinus* have been referred to other genera. See Bassler's Bibliogr. Index, page 692.

ANISOCRINUS Angelin*Plate 20*

Anisocrinus Angelin, Icon. Crin. Suecc., 1878, p. 13.—Springer, Crinoidea Flexibilia, 1920, p. 160.

Rays above radials separated by solid plates arched over by brachials. RA irregularly under r. post. R, above line of BB. Anal x and one large iBr alone or followed by others. Arms dichotomous.

Genotype. *Anisocrinus interradiatus* Angelin.

Distribution. Gotland, America.

Anisocrinus greenei (Miller and Gurley)*Plate 20, figs. 16, 17*

Lecanocrinus greenei Miller and Gurley, Bull. 8, Illinois St. Mus., 1896, p. 52, pl. 3, fig. 28.—*Anisocrinus greenei*, Springer, Crinoidea Flexibilia, 1920, p. 163, pl. 10, figs. 8, 9.

Louisville limestone; Jefferson County, Kentucky; Beech River formation, Decatur County, Tennessee.

There are also figured for comparison:

Anisocrinus oswegoensis (Miller and Gurley), plate 20, figure 18, from Crinoidea Flexibilia, plate 10, figures 7a-c. Oswego, Illinois.

Anisocrinus angelini Wachsmuth and Springer, plate 20, figure 19, from Crinoidea Flexibilia, plate 10, figures 4a, b. Gotland, Sweden.

HORMOCRINUS Springer*Plate 22*

Hormocrinus Springer, Crinoidea Flexibilia, 1920, p. 160.

Rays above radials separated by solid plates followed by perisome. No RA. Anal α and large iBr followed by others abreast and perisome, or directly by perisome. Arms dichotomous.

Genotype. *Centrocrinus tennesseensis* Worthen.

Distribution. Silurian; America, England, Gotland.

Hormocrinus tennesseensis (Worthen)*Plate 22, figs. 1-3b*

Centrocrinus tennesseensis Worthen, Geol. Surv. Illinois, 8, 1890, p. 96, pl. 14, fig. 1.—*Hormocrinus tennesseensis*, Springer, Crinoidea Flexibilia, 1920, p. 167, pl. 14, figs. 1-5.

Beech River formation; Decatur County, Tennessee.

ASAPHOCRINUS Springer*Plate 20*

Asaphocrinus Springer, Crinoidea Flexibilia, 1920, p. 174.

Rays above radials separated by perisome only. RA rhombic, obliquely to left of r. post. R. Anal α followed by others passing into a tube. No iBr, areas filled with perisome, sometimes not exposed. Arms dichotomous.

Genotype. *Asaphocrinus bassleri* Springer.

Distribution. Silurian; America.

Asaphocrinus bassleri Springer*Plate 20, figs. 20, 21*

Springer, Crinoidea Flexibilia, 1920, p. 178, pl. 10, figs. 10-14.

Beech River formation; Decatur County, Tennessee.

Asaphocrinus minor new species*Plate 20, fig. 22*

An imperfect specimen lacking the basal parts, referred with doubt to this genus. The general proportions, distribution of arms, and extent of anal structures, are in favor of it.

Laurel limestone; St. Paul, Indiana.

PYCNOSACCUS Angelin*Plate 21*

Pycnosaccus Angelin, Icon. Crin. Suecc., 1878, p. 13.—Springer, Crinoidea Flexibilia, 1920, p. 180.

Rays above radials separated by perisome only. RA rhombic, obliquely to left of r. post. R. Anal α alone, followed by perisome. IBr 1 to 4. Arms dichotomous.

Genotype. *Cyathocrinites scrobiculatus* Hisinger.

Distribution. Silurian to Devonian; Gotland, England, America.

The first indication of the presence of this genus in America was given by the imperfect specimen described by Weller in 1900 as *P. americanus* from the Racine dolomite. It has since appeared probable that the species described by Hall as *Lecanocrinus caliculus* from the Clinton of New York belongs here. Recent discoveries have extended the range of the genus into the Lower Devonian, and perhaps into the middle Devonian of Canada. The material from the Beech River formation of Tennessee not only furnished fine specimens for illustrating the general habitus of the genus, but has thrown a flood of light upon its detailed structure, by means of which its actual characters became for the first time understood.

Pycnosaccus patei Springer*Plate 21, figs. 1-4*

Springer, Crinoidea Flexibilia, 1920, p. 186, pl. 12, figs. 1-9.

Beech River formation; Decatur County, Tennessee.

Pycnosaccus welleri Springer*Plate 21, figs. 5-7*

Springer, Crinoidea Flexibilia, 1920, p. 186, pl. 13, figs. 3-5.

Beech River formation; Decatur County, Tennessee.

Pycnosaccus laurelianus new species*Plate 21, figs. 8, 8a*

A small species (the figures are double size), with more robust arms and smaller radial than the Tennessee forms, and a peculiar upward widening of the anal plate not observed in others.

Laurel limestone; St. Paul, Indiana.

Pycnosaccus dubius Springer*Plate 21, figs. 9, 9a*

Springer, Crinoidea Flexibilia, 1920, p. 189, pl. 13, figs. 11a, b.

Beech River formation; Decatur County, Tennessee.

Family SAGENOCRINIDAE Roemer emend. Springer

Infrabasals more or less recumbent, taking little part in calyx wall. Crown usually elongate. Rays above radials partly or wholly separated.

SAGENOCRINUS Austin*Plate 22*

Sagenocrinites Austin, Ann. Mag. Nat. Hist., 1842, p. 110.—*Sagenocrinus* Springer, Crinoidea Flexibilia, 1920, p. 215.

RA obliquely under r. post. ray, more or less between BB. Post. B truncate. Anals and iBr numerous. Arms dichotomous, not abutting.

Genotype. *Actinocrinites? expansus* Phillips.*Distribution.* Silurian; England, Gotland, America.**Sagenocrinus clarki** Springer*Plate 22, figs. 4, 4a*

Springer, Crinoidea Flexibilia, 1920, p. 220, pl. 19, figs. 4a-d.

Beech River formation; Decatur County, Tennessee.

Sagenocrinus americanus Springer*Plate 22, fig. 5*

Springer, Am. Geol., 30, 1902, p. 86; Crinoidea Flexibilia, 1920, p. 221, pl. 19, fig. 5.

Waldron shale; Waldron, Indiana.

Family ICHTHYOCRINIDAE Angelin emend. Wachsmuth and Springer

Infrabasals horizontal, not appearing externally and taking no part in the calyx wall. Crown elongate or rotund. Rays widening upward to accommodate expansion of calyx, but in close contact, usually interlocking or abutting above interbrachials.

ICHTHYOCRINUS Conrad*Plate 22*

Ichthyocrinus Conrad, Jour. Acad. Nat. Sci. Phil., 8, 1842, p. 279.—Springer, Crinoidea Flexibilia, 1920, p. 264.

Rays in contact all around. RA in form of R under r. post. R. No anal or iBr. Post. B not differentiated. Arms dichotomous, interlocking.

Genotype. *Ichthyocrinus laevis* Conrad.*Distribution.* Silurian to Devonian; America, England, Gotland, Bohemia.

Ichthyocrinus subangularis Hall*Plate 22, figs. 12, 13**Ichthyocrinus subangularis* Hall, Trans. Albany Inst., 4, 1860, p. 201.—Springer, Crinoidea Flexibilia, 1920, p. 281, pl. 34, figs. 1-10.

Waldron shale; Waldron, Indiana.

CLIDOCHIRUS Angelin*Plate 21**Clidochirus* Angelin, Icon. Crin. Suecc., 1878, p. 12.—Springer, Crinoidea Flexibilia, 1920, p. 294.

Rays in contact except at anal side. RA in form of R under r. post. R. Anal α alone or followed by others. Post B truncate. Arms dichotomous, interlocking.

Genotype. *Clidochirus pyrum* Angelin.*Distribution.* Silurian to Devonian: Gotland, America.**Clidochirus americanus** Springer*Plate 21, figs. 10, 11*

Springer, Crinoidea Flexibilia, 1920, p. 298, pl. 37, figs. 7, 8.

Clinton; Dayton, Ohio.

Suborder TAXOCRINOIDEA

Family TAXOCRINIDAE Wachsmuth emend. Springer

Posterior interradius always differentiated and occupied by anal plates in a tube-like series, none of which are incorporated in the calyx. All anal plates from post. B up separated by perisome from adjacent brachials at one or both sides. Distal face of post. B not suturally connected with anal plate. Rays above radials partly or wholly separated all around. Crown usually elongate.

PROTAXOCRINUS Springer*Plate 22**Protaxocrinus* Springer, Crinoidea Flexibilia, 1920, p. 345.

RA in form of R under r. post. R. iBr few. Rays wholly separated by iBr or perisome. Arms dichotomous.

Genotype. *Taxocrinus ovalis* Angelin.*Distribution.* Ordovician, Silurian, Devonian; Gotland, America.

Protaxocrinus robustus Springer*Plate 22, figs. 6, 7*

Springer, Crinoidea Flexibilia, 1920, p. 349, pl. 45, figs. 9-11.

Beech River formation; Decatur County, Tennessee.

GNORIMOCRINUS Wachsmuth and Springer*Plate 22**Gnorimocrinus* Wachsmuth and Springer, Rev. Pal., 1, 1879, p. 50.—Springer, Crinoidea Flexibilia, 1920, p. 353.

RA rhombic, obliquely to left of r. post. R. iBr few, rays wholly separated by iBr or perisome. Arms dichotomous.

Genotype. *Taxocrinus expansus* Angelin.*Distribution.* Silurian; Gotland, America.**Gnorimocrinus cirrifer** Springer*Plate 22, figs. 8-10*

Springer, Crinoidea Flexibilia, 1920, p. 355, pl. 47, figs. 7-12.

Beech River formation; Decatur County, Tennessee.

Gnorimocrinus varians Springer*Plate 22, figs. 11, 11a*

Springer, Crinoidea Flexibilia, 1920, p. 356, pl. 47, figs. 13-17.

Beech River formation; Decatur County, Tennessee.

Order INADUNATA Wachsmuth and Springer

CRINOIDEA WITH A RIGID CALYX IN WHICH THE BRACHIALS ARE FREE (OR SOMETIMES LOOSELY CONNECTED) ABOVE THE RADIALS. NO INTERRADIALS OR INTERBRACHIALS EXCEPT AT THE POSTERIOR SIDE, AND BRACHIALS NEVER NORMALLY INCORPORATED IN THE CALYX. PLATES OF THE CUP UNITED BY CLOSE SUTURE. MOUTH SUBTEGMINAL; FOOD-GROOVES SUPRATEGMINAL, BUT MAY BE CLOSED BY FIXED AMBULACRAL PLATES. ARMS WITH OR WITHOUT PINNULES.

Of this order by far the most important development resulting from the present collections is in the Calceocrinidae, which in their peculiar anatomy stand out apart from all other crinoids. This has not only furnished the material for the first satisfactory comparison of European and American forms, but also has produced an example of evolutionary modification, with the process clearly visible, that is without precedent among the crinoids. This succession involves a long course from the Ordovician to the Lower Carboniferous, and in the somewhat elaborate treatment of the family which its history has dictated, it has been found necessary to include rather full discussion and illustration based upon those epochs in addition to the Silurian.

Another strange form, *Myelodactylus*, with its coiled bilateral stem, has proved to be prolific in instructive material, revealing close relationship with some of the European species. The genus *Gissocrinus*, hitherto known only from Gotland and England, is represented here by at least two species, one of them of a type decidedly new and distinct.

Among the Larviformia, the puzzling genus *Zophocrinus* is now illustrated more completely than ever before, and doubtful points of its structure cleared up; and the mysterious *Mysticocrinus* re-figured and described. The inconspicuous and widely distributed *Pisocrinus* is represented by such a wealth of new material as to require a complete review of the numerous species already known.

Suborder LARVIFORMIA Wachsmuth and Springer

Monocyclic (with exceptions). Calyx above the base consisting only of radials and orals, without anal plates, and usually without visible ambulacra; plates immovably united by close suture. Arms non-pinnulate, simple and uniserial. Tegmen when known consisting chiefly of orals.

Family PISOCRINIDAE Angelin

Basals three to five; radials five, unequal, r. post. and r. ant. compound, l. post. and ant. much the largest, l. ant. reduced.

PISOCRINUS De Koninck

Plates 23, 24, 25

Pisocrinus De Koninck, Bull. Acad. Roy. Belgique, 2me. ser., 4, 1858, p. 93.—Angelin, Icon. Crin. Suecc., 1878, p. 20.—Wachsmuth and Springer, Rev. Pal., 3, 1886, p. 172.—S. A. Miller, Jour. Cin. Soc. Nat. Hist., 1879, p. 10; Am. Geol., 1890, p. 356; 17th Ann. Rep. Dep. Geol., Indiana, 1892, p. 636-642 (adv. sheets, pp. 26-32).—Bather, Crin. Gotl., 1893, p. 21; Treatise on Zool., 3, 1900, p. 149, fig. 62.—Jaekel, Zeitschr. d. D. Geol. Gesell., 1900, p. 482; Phylogenie und System, 1918, p. 89.—Etheridge, Rec. Australian Mus., 1904, p. 289.—Zittel-Eastman, Textb. Pal., 1913, p. 208.—Thomas, Proc. Iowa Acad. Sci., 22, 1916, p. 291.—Bassler, Bibliogr. Index, 1915, p. 980.

Triacrinus Ringueberg (not Munster), Proc. Acad. Nat. Sci. Phil., 1884, p. 144.

Crinoids of very small size. BB 5, unequal; r. p. and r. a. truncate distally and smaller than the other three, which are angular above. RR unequal and unsymmetric; l. post. and ant. large, widest below, and meeting BB by truncate faces, the other three RR small, angular below and not touching BB; r. post. R. transversely bisected, the large lower segment serving as inferradial for it and the r. ant. R. thus forming the RA, which separates them both from BB; this plate and the two large RR constitute the greater part of the calyx. Arms 5, unbranched non-pinnulate, composed of very long brachials beyond the first, which is short; they spring from indented facets at the middle of the radials bounded by processes or partitions at either side projecting upward and inward; the processes are various in form and size, from low, narrow, rectangular, to high, wide and of spear-head shape, that on the posterior side, formed by the right and left posterior radials, being lower and wider than the others, and curved inward for support of the anal tube. Tube crescentic or triangular in section, resembling an arm. Tegmen arched by 5 solid orals which interlock at the center, meet closely at their lateral faces, and connect with the radial processes. Stem round, with short columnals, thickened in the middle. Surface usually smooth, exceptionally tubercular. (See generic diagram, pl. 23, fig. 46.)

Genotype. *Pisocrinus pilula* De Koninck.

Distribution. Silurian; Gotland, England, America, Australia.

Pisocrinus is a form of cosmopolitan distribution, being found almost wherever Silurian strata are exposed, not only in Europe and America, but also in the antipodes, two species having been described from Australia. A characteristic fossil of the Wenlockian beds of England and Sweden, from the former of which it was first described, the genus has been recognized from American rocks in New York, Ohio, Indiana, Illinois, Iowa, Missouri and Tennessee, in almost all the subdivisions of the Silurian from the Clinton and Osgood to the Racine dolomite and the Louisville. In some of these areas it is an important guide fossil. Species are numerous, and in some of them individuals are abundant. Comparison of European and American species shows a striking correspondence in some of them, which contributes important evidence of a migrational connection by way of a northern route between the faunas of northern Europe and the United States during Silurian epochs.

For a full discussion of the relations of *Pisocrinus* and the literature concerning it, reference should be had to Dr. Bather's Crinoidea of Gotland, 1893. The parallelism of European and American species is recognized, being emphasized by the statement on page 27

that except for the difference in matrix and mode of fossilization he could not undertake to discriminate between some of the species from the two continents. This agrees with my own observations made while assembling a representative series of foreign species for comparison, as is shown by forms herein illustrated on plate 25.

Bather's studies upon the morphology of *Pisocrinus* were very important, especially in placing upon a logical basis the orientation of the calyx, which had not theretofore been properly understood. Thus the relation of the peculiar large plate lying underneath the right posterior and right anterior radials as the radial became clear. He recognized three described species in the European Silurian, and noted an undescribed species from Dudley.

In America eleven species have been described, of which at least one is a synonym. Certain of these species are beautifully distinct, so that there is not the least difficulty in recognizing them, even from inferior specimens; such, for example, as *P. gcmiformis* and *P. baccula*, occurring at the same locality in the Laurel formation, and *P. quinculobus* and *P. tennesseensis* of the Brownsport; whereas *P. campana* is so variable and widely distributed that some varieties are easily confused with other species, and one is inclined to see among them an admixture of European forms.

For one of so small and inconspicuous a habitus, this genus possesses some remarkably distinctive characters. In the unequal distribution of radials, it partakes of the singular type of irregularity which characterises its family, as well as the Calceocrinidae and several other monocyclic genera, such as *Heterocrinus*, *Haplocrinus*, *Myocrinus*, *Catilocrinus*, etc., in which two of them—and those always the same, l. post. and ant.—are simple, larger than the others and of a different shape, while the other three are transversely divided or much reduced in size. The projecting processes between the radial facets are peculiar to this and a few other genera having narrow, unbranched arms; their varying shape offers useful specific characters; the projections are often built up about equally from the apposed radials, in which case the interrational suture bisects the process, while in others the division may be unequal, or the projection entirely confined to one of the radials.

The inequality and difference in shape of the basals is a character in which this genus differs from most other crinoids, even from those of its own family.

The arms are characterized by a great length of brachials—an unusual feature which is repeated in some of the contemporaneous Calceocrinidae. This fact led S. A. Miller into some curious notions about the brachial appendages of *Pisocrinus*, he declaring with characteristic assertiveness that they were not arms, but something else of unknown function for which he coined a new term, "arm blades." The arms vary considerably in length in different species; in some it seems as if they were limited almost to a single brachial, while in others they extend to a dozen or more. The ventral furrow in some species is broad and deep, fringed with side pieces so large as to produce a considerable resemblance to pinnules (pl. 24, figs. 8, 9).

The structure of the tegmen, which was formerly misunderstood—certain isolated, radiately folded, conical plates being supposed to belong to it which we now know to be the fused orals of *Gazaerinus*—has been made clear by the discovery of two specimens in which it is perfectly preserved, consisting of five closely apposed orals similar to those of *Symbathocrinus*.

The characters available for the discrimination of species are for the most part well defined. They are:

1. General form of calyx, whether *a*, conical, elongate, or short and wide; *b*, ovoid, expanding upward or contracting; *c*, globose; *d*, lobed or round. In some species these are constant and reliable, while in others with a wide range of variation all of the first three may be included.
2. Base, whether *a*, convex with small indented column-facet; *b*, concave, broad and shallow, or with deep, funnel-shaped pit; *c*, truncate, with column-facet almost to full width.

3. Basals, whether *a*, large, visible completely in side view, or only by points of the larger ones; *b*, small, hidden at bottom of a deep cavity.
4. Radial facets, whether wide or narrow, plain or dove-tailed.
5. Processes, whether *a*, low and narrow, straight; *b*, angular, widening upward or inward; or *c*, high, spear-head shape.

No discussion of the genus *Pisocrinus* would be complete without consideration of the Devonian forms from the Eifel described by Münster¹ and Schultze² as *Triacrinus*, in which the arrangement of calyx plates is the same, except that it is said to have only three basals instead of five. This probably holds good for *T. altus*, but in *T. depressus* the basals vary from 3 to 5; and in my specimens those with 3 are rather more rare than those with 5. In other ways this species shows a considerable similarity to *P. tennesseensis*. The tri-partite base of the Devonian genus is an evident derivative by fusion from the 5 basals of its Silurian ancestor, and the occasional occurrence of 4 and 5 basals a reversion to the older type. I am giving a series of figures of both species on plate 25, showing these facts.

The other member of the family, *Calycanthocrinus*, also from the Devonian, represents a development of the same plan of radial arrangement from *Triacrinus* in the direction of the Catillocrinidae, by means of an increased number of arm-bearing segments.

In the description of species of *Pisocrinus* the terms "large" and "small" are only relative, all forms of the genus being very small as compared with crinoids generally. The figures on the plates are mostly double size.

***Pisocrinus gemmiformis* S. A. Miller**

Plate 23, figs. 1-8a

Pisocrinus gemmiformis Miller, Jour. Cin. Soc. Nat. Hist., 1879, p. 113, pl. 9, figs. 6a-c; 17th Ann. Rep. Indiana Dep. Geol., 1892, p. 636, pl. 6, figs. 10-12, 24-25 (adv. sheets, 1891, p. 26).—Foerste, Jour. Geol., 1903, p. 562.—Slocum, Field Columb. Mus., 2, Geol. Ser., 1908, p. 278, pl. 84, figs. 1-4.

A small species: diameter of maximum specimen at lower third of cup 6 mm., usually much smaller, ranging down to 2.5 mm. Calyx ovoid, truncate above, contracting at arm-facets; smaller specimens globose. Base broadly rounded, with cavity for column small, abruptly sunken. BB large, well exposed in side view, where the larger ones reach one third the height of calyx. Radial facets wide; processes low, narrow, rectangular, not widening inward. Arms short and heavy. Tegmen arched by 5 interlocking orals, closely bounded by the short first brachials.

One of the two most characteristic species of the Laurel at St. Paul, where it occurs in abundance, mostly less than 4 mm. in diameter; the habitus is small, larger specimens being the exception. The very low, narrow processes between the facets, coupled with the prominent basals, are the distinctive characters. The type of the original description in 1879 was said to be from the lower part of the Niagara group, in Ripley County, Indiana, and in that of 1891 in the 17th Indiana, the type locality was given as Osgood, and the species said to be "now known from Madison and other places in the Niagara group of Indiana." Thus it might have been derived from the Osgood or Laurel formations. According to Foerste (Jour. Geol., 1903, p. 562) it extends from the Osgood to Laurel in Tennessee. Slocum identifies it in the Racine of the Chicago area, but the altered condition of the dolomite specimens may prevent a close comparison. It is also reported from the Bainbridge limestone of

¹ Beiträge z. Petrefaktenk., 1839, p. 3.

² Echin. Eifel. Kalk., 1866, p. 106.

Ste. Genevieve County, Missouri. As the first American species of *Pisocrinus* to be described, there is a tendency to identify as *P. gemmiformis* any small, globose form, which may in some cases be the younger stage of other species. I have found this to be the case in the Tennessee collections. The one thing certain is that the species is characteristic of the typical Laurel formation. The oral plates are shown in figs. 8, 8a.

Horizon and locality. Laurel formation and perhaps Osgood, Niagaran; St. Paul, Osgood, Madison, Indiana, and possibly other localities as noted. I have figured a specimen from Gotland, plate 25, figure 13, which has all the characters of this species.

***Pisocrinus baccula* Miller and Gurley**

Plate 23, figs. 9-15

Pisocrinus baccula Miller and Gurley, Bull. 7, Illinois St. Mus., 1895, p. 79, pl. 5, figs. 23-26.

A relatively large species, maximum specimens 5 mm. high by 8 mm. wide. Calyx globose to ovoid, widest in lower third, contracting upwards. RR and RA curving sharply inward, forming a broad triangular base by which the calyx rests. BB almost wholly concealed in shallow concavity, only points of larger plates sometimes visible from side; outline of base seen from below obtusely triangular. Radial facets narrow, contracting upwards; processes large, spear-head shape. Arms longer than in preceding species.

The most characteristic species of the Laurel limestone at St. Paul, and not recognized elsewhere. It is associated with *P. gemmiformis*, but whether in the same layers cannot be ascertained—presumably not, as the formation is thick, and the material of the quarry dumps, from which most of the specimens are collected, is intermingled from different levels. It differs from that species in almost every essential character, especially in the radial facets and processes, and is recognizable at a glance in adult specimens from all other species by its broad, triangular base. It is abundant at the type locality, and is found in various sizes from 3 to 8 mm. in diameter.

Horizon and locality. Laurel formation, Niagaran; St. Paul, Indiana.

***Pisocrinus granulosus* Rowley**

Plate 23, figs. 37-39

Pisocrinus granulosus Rowley, Am. Geol., 1904, p. 260, pl. 16, figs. 1-3.

A medium sized species. Calyx globose, contracting at the arm-bases, about as high as wide, a maximum specimen being 5.5 by 6.5 mm., smaller ones about equal. Base convex, with very small column-facet, sharply indented; basals of good size, angles of largest ones visible from side view. Radial facets small, dove-tailed; processes large, broadly rounded, angular to spear-head shape. Surface smooth to granulose, or tubercular.

Professor Rowley's types were from brownish or reddish clays in the Bainbridge formation of Ste. Genevieve County, Missouri, associated with his other species, *P. sphericus*, and some other forms, to which there is to be added *P. tennesseensis* as recently discovered by Dr. Weller. Among my Tennessee collections is a series of specimens from red clays of the Dixon formation at the top of Safford's variegated bed, just below the Beech River. They partake of the color of the matrix, in which they differ from all the other Tennessee occur-

rences; and they undoubtedly belong to the present species, having the same more or less granulose surface, depending upon preservation, some being exceptionally tubercular. My figures on plate 23 are made from Tennessee specimens.

Horizon and locality. Dixon and Bainbridge formations, Niagaran; Decatur County, Tennessee, and Ste. Genevieve County, Missouri.

***Pisocrinus campana* S. A. Miller**

Plate 24, figs. 6-27

Pisocrinus campana Miller, 17th Ann. Rep. Indiana Dep. Geol., 1892, p. 642, pl. 11, figs. 4, 5 (adv. sheets, 1891, p. 32).—*Pisocrinus* sp. undet., Wachsmuth and Springer, N. A. Crin. Cam., pl. 8, fig. 10.

A rather large species. Calyx bell-shaped, conical to ovoid, expanding upward to the arm-facets; height of typical specimen 5 mm., width at base 3.5 mm., at facets 6 mm. Base truncate, narrow, with small indented column facet. Basals large, extending well up the sides, all five plainly exposed. Radial facets wide; processes low, rather narrow, square, usually widening inward, rarely obtusely angular. Arms long, composed of 5, 6 or more long brachials, 7 mm. or more in length, with sutures well defined.

A widely distributed species, variable in general form, but with two outstanding characters; the small rectangular processes and high basals, the combination of which, especially when reinforced by the expanding calyx, offers good criteria for identification. The long arms, as shown in 5 specimens, is also a good character, in contrast to the short appendages of several other species. No other species, excepting *P. pocillum* of Gotland, has such uniformly prominent basals, which in some cases rise to one third the height of the calyx at the arm bases. In fact there is a tendency of the angular basals to rise still higher to a connection with the shorter radials, which normally they do not touch. This has been observed in many specimens, especially from the type locality. The connection may be established by the downward extension of one or more of the short radials. This abnormality, which I have illustrated by three figures on plate 24, has not been observed in any other species.

As stated, with the expanding and bell-shaped forms the identification is easy, but those with a lower calyx, ovoid to globose, are confusing; if they have fairly high basals, we may call them *campana*, while those with basals but little visible will have to go into *benedicti*. There will be an intermediate zone in which the distinction is shadowy.

The typical area for this species is in Wabash, Grant and Madison counties in northern Indiana, where it is associated with the other two species described at the same time, *P. gorbyi* and *P. benedicti*. The formation is a dolomitic limestone with interbedded clays or shale, in which the few species of crinoids chiefly occur, thought to be equivalent to either the Racine or the Waukesha formations of the Chicago area.

The material studied by Mr. Miller when describing these species was collected by A. C. Benedict and later acquired by me in the Hammell collection, including the types of this and *P. gorbyi*. Specimens more or less typical, in considerable variety, but all having the characteristic high basals, have been recognized in various localities of the Tennessee area, ranging from the Laurel to the Lobelville formations.

Pisocrinus campana is to the American rocks what *P. pilula* is to the European—a pro-tem species which includes in its variations the principal forms of calyx from conical through ovoid to globose; to which our species adds another by expanding at the distal margin until it is bell-shaped. If casts of specimens from the two regions were intermingled, some of them would be difficult to separate. Bather's observations in the Crinoidea of Gotland, page 30, upon the various forms of *P. pilula* would apply with equal force to the present species, and

his opinion as to the probable identity of some of them with *P. benedicti* and some other American species will be shared by every one who has looked for reliable characters for distinguishing them in our own collections.

Horizon and locality. Dolomites of Wabash, Marion, Anderson, Northern Indiana; Osgood and Laurel; St. Paul and other localities in southern Indiana. Laurel and Brownsport formations; Rise Mill and Flatwoods, Perry County; Martin's Mill, Sinking Creek, Wayne County; Tuck's Mill and various glades in Decatur County, Tennessee.

***Pisocrinus benedicti* S. A. Miller**

Plate 24, figs. 28-36

Pisocrinus benedicti Miller, 17th Ann. Rep. Indiana Dep. Geol., 1892, p. 639, pl. 6, figs. 13-16 (adv. sheets, 1891, p. 29).—Slocum, Field Columb. Mus., 2, Geol. Ser., 1908, p. 279, pl. 84, figs. 8-11.

Any rather small, globose or ovoid form, expanding upward, with fairly wide radial facets and low, square or slightly widening processes, in which the basals are only slightly visible—in other words, which is not *campana*—may be taken as this species. To judge by the material from the type localities in northern Indiana used by Mr. Miller and bearing his labels, it is a rather generalized form, without any very distinctive characters. I have referred to it with doubt specimens from the Brownsport group in Tennessee, with a considerable range of variation in general form from ovoid to oblate. Slocum has recognized it in the dolomites of the Chicago area, and more recently in MS. from the Bainbridge formation in Missouri. None of these identifications are made with much confidence because the species as described is unsatisfactory.

Horizon and locality. Dolomites of Wabash and Madison counties, northern Indiana; Brownsport group of Decatur, Perry counties, Tennessee; Bainbridge limestone, Ste. Genevieve County, Missouri.

***Pisocrinus quinquelobus* Bather**

Plate 23, figs. 16-29

Haplocrinites hemisphericus Troost, Am. Jour. Sci., 1849, p. 420; Proc. Am. Assn. Adv. Sci., 1850, p. 61 (nom. nud.).—*Pisocrinus gorbyi* (in part) Miller, 17th Rep. Ind. Dep. Geol., 1892, p. 640, pl. 6, figs. 21-23.—*Pisocrinus quinquelobus* Bather, Crin. Gotl., 1893, pl. 27; Am. Geol., 1896, p. 184.—Slocum, Field Columb. Mus., 2, Geol. Ser., 1908, p. 280, pl. 84, figs. 5-7.—*Pisocrinus milliganae* Miller and Gurley, Bull. 7, Ill. St. Mus., 1896, p. 80, pl. 5, figs. 27, 28.—Miller, N. A. Geol. Pal., 2d App., 1897, p. 749, figs. 1369, 1370.—Foerste, Jour. Geol., 1903, p. 712.—Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 28.

A medium sized species, exceptionally attaining a width of 8 mm. Calyx much wider than high, measured from the level of the arm-facets, height to width of an average specimen being 3 by 5.5 mm.; cup conical, 5-lobed with interradial depressions, expanding upward, and the large radials recurving into an inverted basal funnel. Basals entirely hidden at bottom of funnel, covered by a narrow column. Radial facets narrow, dove-tailed to receive wedge-shaped processes; processes very large spear-head shape, sometimes petaloid. Arms

short, with not over two or three long brachials beyond the first short one, tapering to a pointed apex.

This is the most common species of the Tennessee area, occurring by the hundreds at various localities in the Brownsport group. It is what Miller and Gurley undertook to describe as *P. milliganae*, but they entirely missed the prime character of the hidden basals, by which alone it is distinguished from *P. gorbyi*, and which it shares with *P. sphericus* of Rowley. Although the authors declare emphatically that their species has "large basals, seen in side view even plainer than in *P. gorbyi*," this does not appear in their figures; and the type specimens, formerly in the collection of Mrs. Milligan and now in my possession, show the large radials recurving deep into the basal pit. It is a singular fact that it should have remained for a foreign author, purely in an incidental way, to give the valid name to the most abundant of all the *Pisocrinus* species; but Bather's description is perfectly correct, and will hold.

The characters are distinctive, as appears by several figures on plate 23 showing the form in all stages, especially the exact location and minute size of the basals as seen in the fractured cross-section at figure 29. The great size and deep undercutting of the spear-head processes is a striking feature of this species. In some specimens, such as figure 18, they are almost as high as the remaining portion of the cup, and in general they occupy over a third the total height of the cup measured to their angular points.

The wealth of material in hand affords full information regarding the arms, of which I figure 4 specimens. All exhibit the same elongated brachials, with a tendency to a decided taper. Aside from its typical region in Tennessee, the species has been recognized sparsely in Missouri and Illinois.

Horizon and locality. Brownsport group of the Niagaran. Collected in all glades in Decatur County, and in excavations at 4 different localities along Beech River, and at 6 other localities in Perry and Wayne counties, Tennessee. In collections from the glades there may be some intermingling with *P. sphericus*, but wherever *quinelobus* was found in place it did not intergrade from lobate to round. Also in the Bainbridge limestone, Ste. Genevieve County, Missouri, and Racine dolomite of the Chicago area. In the sunken basals this species finds a parallel in *P. ollula* of Gotland, as shown by figures 8, 9, 10 of plate 25.

***Pisocrinus gorbyi* S. A. Miller**

Plate 23, figs. 40-45

Pisocrinus gorbyi Miller, 17th Ann. Rep. Indiana Dep. Geol., 1892, p. 640, pl. 6, figs. 17-20 (adv. sheets, 1891, p. 30); N. A. Geol. Pal., 1st App., 1892, p. 681, fig. 1243.—Rowley, Am. Geol., 1904, p. 269, pl. 16, figs. 4-7.

A small species. Similar to *P. quinelobus* except that the basals, instead of being hidden in a cavity, form a triangle, partly occupying a shallow depression, and are more or less visible in a side view. Radial facets rather narrow; processes large, lance-head shape. Surface usually smooth, perhaps sometimes tubercular.

Miller described under this species specimens from two remote areas, northern Indiana and western Tennessee, which agreed in having a lobate calyx. The Tennessee specimens were afterwards separated by Miller and Gurley under the name *P. milliganae*, which they proposed for the prolific species already named by Bather *P. quinelobus*. The original description holds good for the Indiana form, which is fairly common at its type locality. It is readily distinguished from the other two species of that area by its lobed calyx, and it is important

to note that the basals are visible from a side view, which is the only character that separates it from the Tennessee form. This appears in Mr. Miller's figure 18, which is confirmed by the type specimen, now in my possession. With it are about 20 others from the Benedict collection, in several of which this character is shown, so that it may be accepted as reliable, although in the majority of specimens as found the basal plates cannot be made out.

The species has been recognized at two localities in Tennessee in the upper part of the Brownsport group, and in the Bainbridge limestone of Missouri.

Horizon and locality. In dolomite of Wabash and Madison counties, Northern Indiana; Lobelville formation; Flatwoods and Rise Mill, Perry County, Tennessee; Bainbridge limestone; Ste. Genevieve County, Missouri.

***Pisocrinus sphericus* Rowley**

Plate 23, figs. 30-36

Pisocrinus sphericus Rowley (*P. globosus*? Ringuenberg), Am. Geol., No. 34, 1904, p. 270, pl. 16, figs. 8, 9.

Similar to *P. quinquelobus* in having the same minute basals, hidden at the bottom of a deep cavity, and very large lanceolate processes; but with calyx globose in contour instead of lobate, not expanding upward, and perfectly round as seen from above or below. The distinction between the two forms is supported by ample evidence, and in about 60 specimens of this species from a single locality, ranging from small to maximum size, there is no tendency to a lobate form. In Tennessee it occurs chiefly in the uppermost beds of the Beech River formation, probably passing up into the Lobelville; its horizon is usually different from that of *quinquelobus*. It also occurs in the Bainbridge formation of Ste. Genevieve County, Missouri, from which Rowley figured it with doubt as *P. globosus* of Ringuenberg, but added tentatively a name of his own, which will stand.

Horizon and locality. Brownsport formation, Niagaran; clays at top of the Beech River, and base of Lobelville; near Rise Mill, Perry County, and in Decatur and Wayne counties, Tennessee. Bainbridge formation; Ste. Genevieve County, Missouri.

***Pisocrinus tennesseensis* (Roemer)**

Plate 24, figs. 1-5

Symbathocrinus tennesseensis Roemer, Sil. Fauna Westl. Tennessee, 1860, pl. 4, figs. 6a, b.—Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 26.—Bassler, Bibliogr. Index, 1915, p. 1249.—*Pisocrinus tennesseensis* Wachsmuth and Springer, Rev. Pal., 3, 1886, pp. 166, 174.

Attaining a large size—the largest known species. Calyx conical, usually wider than high. Column facet large, shallow, sharply indented, leaving a thin rim at the margin. Base broadly truncate; basals almost entirely confined to the column facet, only a small point of one occasionally seen. Radial facets very wide, narrowing inwards; processes flat above, remarkably narrow at the connection with the radials, and widening both inward and upward, giving the effect of a mortise or dove-tailed union with the facets. They are so much undercut that it is difficult to remove the matrix without prying them off. Specimens

are usually large, small ones rarely seen; maximum calyx 6 mm. high at arm facets, 9 mm. wide at that level and 5 mm. wide at the base. Arms unknown.

This is a very characteristic species of the Brownsport group of Tennessee; is widely distributed, occurring at numerous localities and ranging through several formations. It has lately been discovered by Dr. Weller, with thoroughly representative specimens, in the Bainbridge limestone of Missouri. While nowhere abundant, it is well represented in the collections, and is remarkably constant in its characters. The only species at all resembling it is *Triacrinus depressus* of the Eifel Devonian, through which the evident succession of the two genera can be readily traced, specimens of it showing variations of 3 to 5 basals.

Pisocrinus tennesseensis is not listed in Bassler's Bibliographic Index, nor was it recognized by Miss Wood in the Troost Monograph, being reported in both under Roemer's name *Symbathocrinus*, notwithstanding Wachsmuth and Springer in part 3 of their Revision, had in two places definitely referred the species to *Pisocrinus*.

Horizon and locality. Dixon beds and through the Brownsport, Niagaran. At four localities along Beech River in Decatur County, at Martin's Mill and another locality in Wayne County, and one in Perry County, Tennessee. Also Bainbridge limestone, Ste. Genevieve County, Missouri.

***Pisocrinus pyriformis* (Ringueberg)**

Plate 24, fig. 37

Triacrinus pyriformis Ringueberg, Proc. Acad. Nat. Sci. Phil., 1884, p. 145, pl. 3, fig. 1.

***Pisocrinus globosus* (Ringueberg)**

Plate 24, fig. 38

Triacrinus globosus Ringueberg, Proc. Acad. Nat. Sci. Phil., 1884, p. 146, pl. 3, fig. 2.—*Pisocrinus globosus* S. A. Miller, N. A. Geol. Pal., 1889, p. 269.—Rowley, Am. Geol., 1894, p. 270, pl. 16, figs. 8, 9.

These two species, from the Clinton of Lockport, New York, bear a close relation to some of the western forms and are occasionally identified as occurring in that area; I am therefore figuring the type specimens for comparison.

***Pisocrinus pilula* De Koninck**

Plate 25, figs. 1-6

Bather, Crinoidea of Gotland, 1893, p. 27, for full synonymy.

A series of figures of this protean species from the Wenlockian of Gotland and England is given for comparison with our *P. campana*. Also for comparison with other American species, *P. ollula* Angelin, figures 7-10; *P. pocillum* Angelin, figures 11, 11a; both from Gotland and fully described in Bather's work; and two specimens of undescribed species from Dudley, England, figures 12, 12a and 13-14.

ZOPHOCRINUS S. A. Miller

Plate 25

Zophocrinus Miller, 17th Ann. Rep. Indiana Dep. Geol., 1892, p. 642 (adv. sheets, 1891, p. 32).—Weller, Bull. Chicago Acad. Sci., 4, 1900, p. 152, fig. 57.—Bather, Treatise on Zool., pt. 3, 1900, p. 151, fig. 63.—Slocum, Field Columb. Mus., 2, Geol. Ser., 1908, p. 284, fig. 5.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 158.—Bassler, Bibliogr. Index, 1915, p. 1339.

Monocyclic. Calyx elongate pyriform; BB 3, large, 2 equal and one larger; RR 4, the one on the posterior side larger than the others, probably representing r. and l. post. RR fused. Tegmen solid, composed of 5 spade-shaped orals, interlocking at the center and extending outward narrower to the periphery, where they connect with the radials; post. oral the largest, pushed in between the points of the other four; five groups of narrow plates resembling brachials, usually three in each group, closely united, two meeting behind the third, lie between the peripheral extensions of the orals; they spring from the radials and their distal ends are marked by fossae, indicating articular facets for slender arms. Surface smooth. Column-facet small. Specimens small, ranging from 5 mm. high to a maximum of 10 mm.

Genotype. *Zophocrinus howardi* S. A. Miller.

Distribution. Niagaran; America; not observed in Europe.

Chiefly known from the type species, from which the characters of the genus are taken; it occurs typically in the Laurel limestone, at St. Paul, Indiana, but has been found in Tennessee, and two species have been described from the Chicago area.

This singular form is remarkable for the fact that whereas the dorsal cup is composed above the base of four radial elements, evidently formed by fusion of two from the primitive five, the tegmen is constructed strictly on the quinquerradial plan, with 5 large orals meeting toward the center like those of *Haplocrinus*; but unlike the latter connecting with the dorsal cup by extensions of a different shape, between which are groups of narrow plates supposed to be brachials. These groups, while connecting with the 4 large radials, do not seem to stand in any definite relation to them, as they do not, except occasionally, spring from their middle. Hence it would seem that the disturbance in radiation internally was greater than would be caused by a simple fusion of two radials, as in that case there should be a group of brachials from the middle of each of the three equal original radials, and two from the larger one. The anal opening is not always distinctly shown, but seems to lie closely alongside the peripheral extension of the posterior oral; apparently this does not always coincide with the larger radial.

Now it happens that the essential structures by which *Zophocrinus* differs so markedly from other crinoids are found in the Devonian genus *Tiaracrinus* from the Eifel. This was described by Schultze in 1867 (Mon. Echin. Eifl., p. 226, pl. 13, fig. 8) from a single species, *T. quadrifrons*, but with a complete misconception of its characters. The tegmen with its border of narrow marginal plates he mistook for a stem segment, and figured his specimen upside down, so that the small cavity for the column-facet was conceived to be the tegmental opening where the calyx plates almost came together. This fact was noted by Zittel in 1879¹ and has been further mentioned by Oehlert in connection with the description of a new species, *T. soyei*.²

¹ Handb. Pal., 1, p. 425.

² Bull. Geol. Soc. France, Ser. 3, 1882, p. 359.

I have given on plate 25, figures 27-27c, careful figures of the specimen used by Schultze as the type, to show the remarkable parallelism with *Zophocrinus*. The two forms differ widely in certain details. *Tiaracrinus* has a very small, concave base, apparently of 3 plates, and the surface of the cup is covered by 4 series of sharp, transverse folds passing from one radial to another, ending in deep pits forming vertical rows separated by a wide groove along the middle of each radial—in contrast to the perfectly smooth surface of *Zophocrinus*. The cup has 4 unequal radials, but instead of the posterior one being largest, it is sometimes the smallest.

But the solid tegmen is constructed on precisely the same principle as that of *Zophocrinus*, being composed of 5 orals meeting at the center and interlocking, with extensions or connections reaching to the periphery between which are 5 groups of extremely narrow plates, presumably brachials, springing from the radials and forming with the interposed oral extensions a conspicuous border around the margin. Instead of 3 to a group, the number of brachials here is 11 or 12, and they do not coincide in position with the radials. Thus the 4 radials, quinquepartite tegmen, and the peripheral border of narrow brachials, are the essential characters common to the two forms. The enlarged figure of this tegmen on plate 25 shows some indefinite small plates at the outer angles of the orals which are not understood. As the structures are preserved in this unique specimen, some of the sutures are obscure, and these small plates are not to be depended on.

The interpretation of the peripheral border of narrow plates in *Zophocrinus* has been somewhat of a puzzle. Those of *Tiaracrinus* represent the same element without doubt, the only difference being their greater number, amounting to 55 or 60 instead of about 15. That they are brachials, in view of their form, and position relative to the radials, seems highly probable, and Bather's suggestion that they may be compared with the arms of *Catillocrinus* accords with my own view. In both cases there would be the unusual feature of several arms originating upon a single radial, and in the *Catillocrinidae* we have a range in number of arms—from 12 to 57—similar to that in the two genera under consideration, viz.: 15 to 55 or 60.¹ The distinct presence of fossae upon the distal face of the brachials in *Zophocrinus* indicates that they must have borne articulated arms.

As to the systematic position of these genera Bather and Jaekel are in substantial agreement, placing *Zophocrinus* among the Monocyclic Inadunata, while referring *Tiaracrinus* to the Cystidea (Lankester Zool., 1900, pp. 151, 20; Phylogenie und System, 1918, pp. 89, 99), the latter based upon the belief that the rows of pits up and down the middle of the radials represent pore-rhombs. There is no suggestion of pores on the smooth surface of *Zophocrinus* to call for a reference to the cystids, and the remarkable identity in essential calyx structure would seem to preclude placing the two genera in different divisions of echinoderms. It may be that the extreme specialization of the tegmen and arms common to the two is an independent development, but it seems more reasonable, and in better accord with the facts as now disclosed, to regard them both as aberrant crinoids, in some way related, and modified from the Silurian to the Devonian.

***Zophocrinus howardi* S. A. Miller**

Plate 25, figs. 19-26

Zophocrinus howardi S. A. Miller, 17th Ann. Rep. Indiana Dep. Geol., 1892, p. 642 (adv. sheets, 1891, p. 33), pl. 6, figs. 26-28; N. A. Geol. Pal., 1st App., 1892, p. 683, fig. 1254.—Weller, Bull. Chicago Acad. Sci., 4, pt. 1, 1900, p. 152, pl. 15, fig. 13.—Bather, Treatise on Zool., pt. 3, 1900, p. 151, fig. 63.—Grabau and Shimer, N. A. Index Foss., 2, 1910, p. 474, fig. 1785.

Type of the genus. The characters are fully stated in the general remarks. The species is rather abundant at the type locality, St. Paul, Indiana, but speci-

¹ Springer, On the family Catillocrinidae, Smithsonian Misc. Coll., vol. 76, 1923, p. 20.

mens showing the structure of the tegmen and marginal brachials clearly are rare. The foregoing description is made from twelve specimens in which the groups of three brachials and the interposed orals are distinctly shown. In these small specimens the sutures in the tegmen are usually rather obscure. Mr. Chapman's enlarged figure 22 on plate 25 was made many years ago after a careful study of two well preserved specimens. A specimen subsequently acquired, figures 21, 21a, having the oral plates slightly parted, confirms the original figure in every particular. The species ranges in size from 5 to 10 mm. height to the top of the radials; the latter figure is rare, the usual height being from 6 to 8 mm.

Horizon and locality. Laurel limestone; St. Paul and Greensboro, Indiana; and Beech River formation; Flatwoods, Perry County, Tennessee; perhaps Racine dolomite; Lemont, Illinois.

Zophocrinus globosus Slocom, Field Columb. Mus., Geol. Ser. 2, 1908, p. 285, pl. 85, figs. 15-19, from near Lemont, Illinois, and *Zophocrinus pyriformis* Slocom, *ibid.*, page 285, plate 85, figures 12-14, from Romeo, Illinois, are both referred to the Racine division of the Niagaran.

MYSTICOCRINUS Springer

Plate 26

Springer, Amer. Jour. Sci., 46, 1918, p. 666.

Calyx globose, rigid, with no indication of loose suture or flexibility; lower brachials do not take part in calyx wall. Dicyclic; IBB 3, the small plate in r. post. position. RA in primitive position below r. post. R. being the lower segment of that radial, which (and no other) is compound. RR unequal, with lateral processes projecting between arm-bases; facets curved and excavated, not filling distal face of radials. Anal plate 1, large, angular above, projecting above level of radials; opening unknown. Arms uniserial, branching once, composed of only a few brachials. Tegmen narrow, probably covered by a pyramid of orals as in *Pisocrinus* and *Symbathocrinus*.

Genotype. *Mysticoocrinus wilsoni* Springer.

Distribution. Silurian; not known outside of America.

This diminutive crinoid, described by me in 1918 from a very perfect specimen, and of which a fragment of another slightly larger is known, is of a wholly novel type, with such a complex assemblage of characters that its systematic position is uncertain. It is an Inadunate, superficially resembling the Larviformia, and upon a preponderance of characters coming nearer to that group than to any other. The primitive radianal is against it, and the dicyclic base also, although the latter may fall within an admitted exception. It would seem to be intermediate between the Larviformia and the Fistulata; in the primary composition of the cup, especially in the radianal under the right posterior radial, it is similar to the Dendrocrinidae, although wholly unlike them in the dwarfed arms and interbrachial processes, both of which are Larviformia characteristics. The 3 IBB, with small plate at right posterior, is a character of the Flexibilia, from which it is excluded on other grounds.

Aside from this combination of general characters, the most striking feature is the inequality of the radials, and the distribution of the processes on radial and anal plates in such a way that there is an arrow-head projection between the arm-bases in every interbrachial area. Processes analogous to these, of various shapes, are characters of *Pisocrinus*.

***Mysticocrinus wilsoni* Springer**

Plate 26, figs. 1a-g

Springer, Amer. Jour. Sci., 46, 1918, p. 666, pl. II, figs. 1-8.

Calyx very small, the size of small *Pisocrinus*, about 2.7 mm. height and width of type, with but little height added by the infolding arms; flattened below, strongly constricted at the arm-bases. IBB forming a shallow saucer, externally concave, the smaller r. post. plate being more than one fifth the area of the pentagon. BB strongly convex, bent outward and upward, forming about one third the height of the cup; post. B heptagonal, wider than high, subangular below, truncate above; r. post. and r. ant. BB hexagonal, the other two pentagonal. RA pentagonal, truncate above to meet r. post. R, and beveling the lower corner of anal. Anal α very large, sagittaeform, higher than wide, beveled laterally by the RA; apex acuminate, rising almost to the height of first secundibrachs; offsets of arrow-head resting upon and interlocking with shoulders of post. RR.

RR unequal and differing in shape; angular processes project nearly as high as the anal between the arm-bases from three of the RR, viz., at one (outer) side of each posterior and at both sides of the anterior R, while the laterals have none; each process has an offset like those of the anal interlocking with adjacent RR. Radial facets deeply curved, occupying a median position except on the two post RR, in which the single outer processes occupy about one-half the distal width of the plate and the facets toward the posterior side the remaining half. Ant. R. the largest plate in the cup; it and the two post. RR widen upwards, while the two lateral RR correspondingly diminish in width, and by reason of the absence of processes are shorter than the former; the R. post. R is the smallest, although with the RA added it is about equal to and slightly longer than l. post. R. Arms unequal, uniserial, branching once, infolding and apparently very short; the two post. and probably the ant. arms have 2 IBr, while the laterals seem to have but one; the axillaries are followed by series of 3 or more subquadrangular brachials which taper gradually. Anus and tegmen unknown. Stem circular.

Horizon and locality. Laurel limestone, at top of chert; St. Paul, Indiana.

Suborder FISTULATA Wachsmuth and Springer

Monocyclic or dicyclic. Posterior interambulacrum usually more or less extended into a strongly plated anal tube or sac. Arms pinnulate or non-pinnulate, usually uniserial, but biserial in some of the later genera. Tegmen composed of numerous plates, either orals with supra tegmental ambulacra passing over their edges, and interambulacra, or more or less undifferentiated plates.

Family HETEROCRINIDAE Zittel

Monocyclic. Basals five. One or more of radials compound. Anal α usually resting on left shoulder of r. post. R, supporting a tube. Arms five, non-pinnulate, uniserial, dichotomous or heterotomous, or may bifurcate with ramules; proximal IBr usually full width of RR. Tegmen little known.

MYELODACTYLUS Hall*Plate 27*

Myelodactylus Hall, Pal. New York, 2, 1852, p. 191.—Angelin, Icon. Crin. Suecc., 1878, p. 11.—Wachsmuth and Springer, Rev. Pal., 1, 1879, p. 146.—Weller, Bull. Chicago Acad. Sci., 4, 1900, pp. 45, 68.—Springer, Unusual forms of Fossil Crinoids, Proc. U. S. Nat. Mus., vol. 67, 1926, pp. 3-32, pls. 1-6. *Herpetocrinus* Salter, Catal. Camb. Sil. Foss., 1873, p. 118.—Bather, Crin. Gotl., 1893, p. 36, text-fig. 12, pl. 1, figs. 24-77; Amer. Geol., 1895, p. 214; Nat. Sci., 12, 1898, p. 339; Treatise on Zool., 3, 1900, p. 146, fig. 49.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 212.—Bassler, Bibliogr. Index, 1915, p. 844.—Thomas, Proc. Iowa Acad. Sci., 26, 1919, p. 481.

An Inadunate crinoid with coiled stem, more or less completely enveloping the crown, circular in proximal region; bilaterally symmetric, elliptic or sub-crescentic in middle and distal regions; with two rows of cirri along the margins or back of the bilateral part. Crown similar to that of *Iocrinus*.

Genotype. *Myelodactylus convolutus* Hall.

Distribution. Silurian to Lower Devonian; America, England, Gotland.

Having recently in the paper on "Unusual Forms of Fossil Crinoids," above cited, discussed this remarkable form and its congeners at considerable length, with descriptions and illustrations of the known American species, it is unnecessary to go into details concerning it here. Inasmuch, however, as some of the most important of the new material belongs to the Tennessee fauna under consideration, and for completeness must be given a place in this work, it has been thought best to reproduce a selection from the principal figures, assembled upon a single plate the better to show at a glance the variety and range of this greatly specialized genus. To this end a list of the species as figured will be given with a brief résumé of the salient characters, while for more complete information the reader is referred to the paper alluded to.

Myelodactylus ammonis (Bather)*Plate 27, figs. 1-5a*

Herpetocrinus ammonis Bather, Crin. Gotl., 1893, p. 49, pl. 2, figs. 54-63.—*Myelodactylus ammonis* Springer, Unusual Forms, 1926, p. 10, pl. 2, figs. 1-9.

Coil very close; stem short, tapering to a point; cirri short, alternating from broad end of successive columnals.

Beech River formation; Decatur County; and Waldron shale; Newsom, Tennessee. *M. gorbyi* Miller and Gurley, 17th Rep. Indiana Dep. Geol., 1891, p. 72, from near Nashville, Tennessee, is too indefinite for comparison.

Myelodactylus convolutus Hall*Plate 27, figs. 6-8*

Myelodactylus convolutus Hall, Pal. New York, 2, 1852, p. 192, pl. 45, figs. 5, 6.—Springer, Unusual Forms, 1926, pp. 8, 16, pl. 1, figs. 1-8.—*Herpetocrinus convolutus* Bather, Crin. Gotl., 1893, p. 48, pl. 2, figs. 50-53.

Coil open and broadly convolute distalwards; stem long; columnals short, quadrangular, uniform; cirri flat, regularly paired on successive columnals.

Rochester shale; Lockport, New York; Laurel limestone, St. Paul, Indiana; and other Silurian horizons and localities. *M. bridgeportensis* S. A. Miller, Jour. Cin. Soc. Nat. Hist., 1880, p. 141, figs. 2a-e, from the Racine dolomite, and *Eomyelodactylus rotundatus* Foerste, Bull. 19, Sci. Lab. Dennison Univ., 1919, p. 19, pl. 1, fig. 8, pl. 2, fig. 3 are thought to belong to this species.

Myelodactylus brevis Springer*Plate 27, figs. 9, 9a*

Springer, Unusual Forms, 1926, p. 10, figs. 9, 9a.

Like *convolutus*, but with coil close and stem short, tapering to a point.

Beech River formation; Decatur County, Tennessee.

Myelodactylus fletcheri (Salter)*Plate 27, figs. 10, 10a, b*

Herpetocrinus fletcheri Salter, Catal. Camb. Sil. Foss., 1873, p. 118.—Bather, Crin. Gotl., 1893, p. 46, pl. 1, figs. 34-49, and p. 182, under fig. 38.—*Myelodactylus fletcheri* Springer, Unusual Forms, 1926, p. 10, pl. 1, figs. 10-12c.

A species from the Wenlockian of Dudley, England, and occurring also in Gotland. Similar to *M. convolutus*, but with cirri round and bead-like. The specimen figured is notable for being the only one of the genus from the Silurian which shows the exact structure of the crown, of Iocrinid type, with only 4 rays. It was noticed in Bather's work of 1893, on p. 182.

Myelodactylus extensus Springer*Plate 27, figs. 11-18*

Springer, Unusual Forms, 1926, p. 14, pl. 3, figs. 1-13.

Like *ammonis*, except that the coil is open beyond the proximal region, and stem extended for a considerable distance in a broad curve toward the distal end.

Beech River formation; Decatur County, Tennessee.

Myelodactylus brachiatus Hall*Plate 27, fig. 19*

Myelodactylus brachiatus Hall, Pal. New York, 2, 1852, p. 232, pl. 45, figs. 7a-e.—Springer, Unusual Forms, 1926, p. 14, pl. 4, figs. 1-10.

Coil open, circular part of stem very long and slender, with crown tending to be free of the coil; cirri limited to the distal end, few, long, branching, and alternating at intervals of several columnals.

Rochester shale; Lockport, New York.

Myelodactylus keyserensis Springer*Plate 27, figs. 20, 20a*

Springer, Unusual Forms, 1926, p. 19, pl. 6, figs. 1-3.

Coil open; cirri numerous, long, slender, paired on successive columnals; crown large, with long arms branching repeatedly, swelling between the two rows of closely packed cirri; rays 5, of *Iocrinus* type.

Keyser formation, Helderbergian; Lower Devonian; Keyser, West Virginia.

Myelodactylus schucherti Springer*Plate 27, fig. 21*

Springer, Unusual Forms, 1926, p. 21, pl. 5, figs. 9-9c.

Coil close, circular part of stem thick and long; cirri round, short, paired on successive columnals.

Linden formation, Helderbergian, Lower Devonian; Benton County, Tennessee.

Myelodactylus nodosarius (Hall)*Plate 27, figs. 22-24*

Brachiocrinus nodosarius Hall, Pal. New York, 3, 1859, p. 118, pl. 5, figs. 5-7; pl. 6, figs. 1-3.—Goldring, Devon. Crin. New York, 1923, p. 332, pl. 41, figs. 1-4.—*Herpetocrinus nodosarius* Bather, Amer. Geol., 16, 1895, p. 213.—*Myelodactylus nodosarius* Springer, Unusual Forms, 1926, p. 20, pl. 5, figs. 1-8.

Coil open; stem elongate, terminating in a bulbous enlargement; cirri few, short, thicker than the stem and thickest in the middle, alternating at intervals of 1 to 5 columnals, composed of a few rounded, bead-like cirrals.

New Scotland formation, Helderbergian, Lower Devonian; Schoharie County, New York.

CALCEOCRINIDAE Meek and Worthen

Calceocrinidae Meek and Worthen, Geol. Surv. Illinois, 5, 1873, p. 502.—Wachsmuth and Springer, Rev. Pal., 3, 1886, p. 273.—Ringueberg, Ann. New York Acad. Sci., 4, 1889, p. 388.—Bather, Crin. Gotl. 1893, p. 54; Treatise on Zool., 3, 1900, p. 147.—*Cremacrinidae* Ulrich, 14th Ann. Rep. Geol. Surv. Minnesota, 1886, p. 105.

Monocyclic Inadunata, with crown pendent or recumbent on the stem, in which the radials are connected with the base by muscular articulation instead of by suture, and in which bilateral symmetry, at first disturbed, has been progressively established by the location of the stem first to the right of the anal tube, and afterwards in the same plane; anterior and left posterior radials large and simple; left anterior, right posterior and right anterior radials originally compound, the last two finally obliterated or non-arm-bearing; anal α shifted over the right posterior radius, and supporting a massive anal tube; arms of two types, one (1. ant.) median, marking the plane of bilateral symmetry, and three lateral in early stage, afterwards, and finally, becoming definitely fixed at two, which are peculiarly modified by a doubly heterotomous mode of branching, wholly distinct from that of the median arm; basals first 4, later reduced by fusion to 3; calyx antero-posteriorly compressed, especially at the junction of radials and basals, where it is almost linear; transverse section about middle of radials triangular.

Ordovician to Lower Carboniferous; America and Europe.

This singularly modified crinoid differs from all others by having the base united to the radials by a hinged muscular articulation, allowing motion of the crown above the base up and down in the plane of its bilateral symmetry. By this character, without precedent among the echinoderms, the family is set apart from the remainder of the class as a distinct zoological unit. Its morphology has been studied by Wachsmuth and Springer, Ringueberg, and Bather. The most extensive study is that of Bather, contained in his monograph on the Crinoidea of Gotland, 1893, which embodied an elaborate treatise on the family Calceocrinidae, with the advantage of the fine material from the Swedish and English Silurian, and is enriched with a wealth of beautiful figures drawn by the master hand of Georg Liljevall. No Silurian material at all comparable to that had been available to the American authors, nor has there been since until the collection herein described was assembled. In the treatment of the family, I have availed myself of Bather's work, and of his later treatise in the Lankester Zoology, part 3, 1900, page 147.

The literature on the subject is difficult to follow, aside from the obscurity inherent in the anomalous structure, because the authors have employed different terminology for the parts which are peculiar to it. It is therefore necessary, before entering upon the discussion of a type of which the superficial appearance in its later stage is so widely different from that of other crinoids, to settle the orientation of the calyx. This depends chiefly upon the relation of the radials, which are so greatly affected by the modifications that have occurred. Wachsmuth and Springer and Ringueberg treated the radius which is located between the two large radials, and thus occupies a conspicuous median position, and which had been called "dorsal" by Hall, as the anterior. Bather, on the other hand, regarded the median radius

as the left anterior, the anterior being the large plate adjoining it to the left. I have adopted this plan, as being the logical deduction from the facts about which there is no dispute.

The type under consideration is beyond question closely related to the Heterocrinidae, with which it was contemporaneous in the early stage, and of which it was evidently an offshoot. E. Billings described the first two species of it as *Heterocrinus*. In most of the early monocyclic genera of that type having unequal radials, such as *Heterocrinus*, *Ectenocrinus*, *Ohioocrinus*, and also those of other groups, as *Pisocrinus*, *Triacrinus*, *Mycocrinus* and *Catilloocrinus*, it is always the anterior and left posterior radials that are large and simple, while the others are compound or much reduced in size. (See pl. 23, fig. 46.)

Having thus the essential characters of the Heterocrinidae, and the two large radials with the arms they bear being clearly identified as the anterior and left posterior, it necessarily follows that in this family the arm or radius which lies between them, and which on the side away from the anal tube occupies the median position, is the *left anterior*. With this fact established to begin with, we are prepared to trace the modifications which followed the reversal of the crown upon the stem.

The first departure from the primitive type was the disappearance of one basal and the r. post. arm, and the location of the stem at one side of the anal tube, and to the right of the r. post. radial; the plane marked by the stem and l. ant. radial is one of imperfect bilateral symmetry. R. post. arm being now absent, its place is occupied by the anal tube. Next the r. ant. arm also disappears in the later genera, in which the stem shifts so as to coincide with the anal tube in the median posterior position, thus producing complete bilateral symmetry; the r. post. and r. ant. superradials fuse to form a transversely extended subanal¹ plate, sometimes T-shaped, underlying the anal σ , which is the direct support, or beginning of the anal tube; later the subanal piece atrophies, and the anal σ or tube rests upon the corresponding inferradials; finally also the left basals fuse into a single plate. The simple radials (l. post. and ant.) increase in size, forming the sides of the cup, and bending around toward the posterior, where the tube is, as well as in the opposite direction where they eventually meet between the two segments of l. ant. compound radial.

These segments are quite variable in shape and size, and the variation is decidedly progressive. In the Ordovician forms the lower one, inferradial, is elongate quadrilateral, rather broad, and connects by its distal face with the roughly triangular superradial. A similar connection, with a gradual decrease in width of the inferradial, is maintained in some Silurian forms; in others the two segments are separated, both becoming triangular, and this structure remains fixed and constant throughout the Devonian and Lower Carboniferous. The proximal margin of this and the adjoining radials next to the hinge is often strongly denticulate.

The arm system becomes peculiarly modified, both in general plan and minor details. It consists of two distinct types, one of which is limited to a single arm, the left anterior, while the other contains two or more arms, first unsymmetric and afterwards equally balanced on either side of it. The former is an element *sui generis*, standing alone, distinguished from all the others by its median location and exclusive characters. By reason of its size, position, and relation to the other arms, it dominates the anterior region of the crown, and as the diametrical opposite of the reversed stem fixes the plane of bilateral symmetry somewhat after the manner of a spinal column. In view of its strictly median position between those borne by the two large, simple radials, I have found it convenient in discussion to designate this as the "median arm," rather than as the "dorsal" as is done by Hall and some other authors, that term being objectionable because currently used among the echinoderms in an entirely different sense. Of course the term "left anterior" is retained as the correct technical expression.

¹ I prefer the term "subanal," which is self-explanatory and correct in fact, to that of "T-piece" as employed by some authors, which is misleading in that the plate is scarcely ever found of that shape.

The median arm is borne by the l. ant. R, and it varies greatly in size; it is sometimes much the largest, furnishing the main controlling element in the movement of the crown; frequently it remains simple, extending distally beyond the ends of the other arms; it may branch, usually only once in the earlier forms, but sometimes twice or even more in some of the later; also it varies in the number and length of brachials, from 2 or 3 of extreme length to 20 or more short ones; occasionally this arm is relatively slender, and even shorter than the lateral arms. But always it is a unit by itself, without any corresponding element.

The two side arms borne by the large, simple radials, each at first symmetrically divided into two equal rami, are later extraordinarily developed into numerous branches springing from successive diminishing axillaries having unequal articulating faces, each giving off from the shorter, inner face a branch called an "axil-arm," and from the longer, outer face the next axillary. The latter constitute the main ramus of each of these arms, composed of a series of 3 to 8 axillaries, termed "main-axils," which lie side by side and curve around transversely toward the anal tube in form of an arch; the branch (axil-arm) springing from each main-axil on the shorter side toward the median arm gives off ramules in alternate succession, of which those on the inner side are often invisible from the exterior of the folded arm, while those on the outer side, especially the first one, may reach the height of the crown and form the chief visible part of the arm. This plan of arm-branching may be called the "axil-arm system."

The base also undergoes notable modifications. In the first remove from the ancestral form, concurrent with the asymmetric location of the stem and the disappearance of the r. post. arm, the r. post. basal atrophies, and is only rarely, perhaps doubtfully, represented by a minute triangular relic; thus there are left 4 BB, which form a semicircular base, hinged by muscular articulation to the radials, with the hinge on the straight side and the larger r. ant. and post. BB on the curve. All the basals are in contact with the stem as usual in crinoids; the median two next to the hinge, l. post. and l. ant. BB, called the left basals, enter or touch the stem-facet by narrow points, and their outer sides make a reverse curve, giving a peculiar and characteristic outline. In the next, 3-armed, stage, with the stem shifted back to the median position, the 4 basals remain as before, the two left BB in contact with the stem by their points, and preserving the same general shape. But in the next, axil-arm, stage these basals are completely detached from the stem, and fuse into a single plate which is usually triangular with straight sides in the Silurian forms, and finally becomes curved in the Carboniferous. Thus while in the Ordovician stage and its immediate Silurian successor the four basals are connected with the stem as in their ancestor when erect upon the stem, a progressive change occurs with the establishment of the axil-arm system, by which two of them are entirely withdrawn from contact with the stem or its facet. These differences are further important because they can be recognized from the isolated base alone, which is often the only part recovered in the fossils.

These various stages form an evolutionary series, the members of which may for the present be designated as A, B, C and D. It begins in the Ordovician with Form A, having 3 lateral arm-bearing rays, hereinafter for convenience called simply arms, the stem alongside the anal tube, the lateral axillary primibrachs equal-faced (in which three characters it differs from all the rest of the family), also 4 basals unfused, the r. post. and r. ant. inferradials separated from one another and from the anal σ by the corresponding superradials, and the two segments of l. ant. R broadly connected; is modified in the Silurian with Forms B and C, having the lateral arms reduced to two, balanced, the stem shifted to the plane of the anal tube, the primibrachs unequal-faced; the left basals in the former divided, and in the latter fused into a single triangular plate; and it culminates in the Devonian and Lower Carboniferous with Form D, in which, along with the foregoing characters of C, the fused l. post. and l. ant. basals finally form a curved plate as wide as the hinge, the r. post. and r. ant. inferradials have come together underneath the anal σ , the superradials which as the subanal piece

Lateral arms as in C; inferradials of vanished arms meet under *x*; segments of l. ant. R always separated. BB 3, the fused left basal more or less curved, not touching stem. Devonian to Lower Carboniferous. *Halysiocrinus*.

Lateral arms 2, balanced, with axil-arms borne on short faces of successive axillaries, arching toward anal tube; inferradials of vanished arms separated by subanal piece; segments of l. ant. R usually separated. 3 BB, l. ant. and l. post. fused to a triangular plate, not touching stem. Silurian. *Calceocrinus*.

R. ant. arm vanished, leaving median and 2 lateral arms balanced; stem shifted to plane of anal tube; segments of l. ant. R narrowly connected; 4 BB, all touching stem. Silurian. *Eucheirocrinus*.

Crown bent upon stem; r. post. arm and basal vanished, leaving median and 3 lateral arms, unsymmetric, anal tube, and 4 BB all touching stem; stem at side of tube; segments of l. ant. R broadly connected. Ordovician to Silurian. *Cremacrinus*.

Probable ancestral form like *Heterocrinus*, with 5 arm-bearing RR, 2 simple, 3 compound; crown erect upon stem; anal tube springing from r. post. R. Ordovician. Dotted line indicates arm and basal first to be eliminated.

FIG. 1. The Series of Calceocrinidae
Read from below up

before separated them having disappeared, and the two segments of l. ant. radial, having as a rule become disconnected in C, are now widely and permanently separated by the two large radials meeting between them. In B the lateral arm structure is of an intermediate stage, while C, which is abundantly represented in the fauna herein described, has in common with D the axil-arm structure, and differs from it in the fact that in most species the two inferradials are still separated by the subanal piece formed by the fusion of their respective superradial segments, and also in the size and shape of the fused basals.

This may be better understood by reference to the accompanying series of diagrams on page 91 constructed as if the cup were opened between the anterior and left anterior radials and spread out flat; this puts the left anterior ray always at the left end, with the anal tube and stem at the front. The relative position of the respective parts is maintained throughout.

In the Ordovician type, which passes over into the Silurian, the side arms branch so that the ray divides equally upon the axillary primibrach into two main divisions, which are symmetric as between themselves. From these ramules are given off from successive brachial series alternately on opposite sides after the usual manner of the heterotomous arm.

In the other leading Silurian form, C, as before stated, the axil-arm system, a wholly different and unprecedented plan of arm structure, was developed, in which, instead of the rays dividing symmetrically into two equal rami, they divided upon the primibrach from unequal articulating faces, the shorter one giving off an axil-arm and the longer one another axillary, likewise unsymmetrical, from which the same process was repeated several times. Thus the branching was by successive axillaries from one side of the ray outward from the median arm toward the anal tube, producing a somewhat fan-shaped arrangement of the ray on either side. This modification changed the general aspect of the crinoid, and the relation of its calyx and arm plates, to such an extent that in order to understand the descriptions it is desirable to have a little clearer account of what happened during the course of the evolution from Ordovician to Carboniferous times.

Starting, for example, with the cup of *Heterocrinus bellewillensis* of W. R. Billings, as given opposite O at the bottom of the series, we have a circlet of five arm-bearing radial plates, three of them compound and two simple, together with an anal tube springing from the r. post. radial, all underlain by 5 basals. The plane of bilateral symmetry in this circlet bisects the anal tube and the simple anterior radial, with two radials on either side, one simple and one compound at the left, and two compound ones at the right. By reason of the stemward bending of the crown, and consequent pushing of the anal tube to the right, the two arms to the right of it were one after the other crowded out; the superradials and inferradials were modified until the arm-bearing segments disappeared or ceased to function, and there were left only three arm-bearing radials, the l. post., l. ant. and ant. RR in that order. The l. ant. R became the base of the median arm; the other two, having greatly increased in size, supported the side arms, which by their numerous branches spread around toward each other, and toward the anal tube, usually overlapping and concealing it from view. This resulted in a flattened or elliptic ring of calyx plates, consisting of a median and two lateral arm-bearing radials and the anal series, in which the radial antipodal to the anal tube was now the left anterior instead of the anterior as in the ancestral stage, leaving one of the large simple radials at either side; so that the plane of bilateral symmetry would lie from the stem and anal tube to the median arm.

Thus the goal toward which this family group was tending was a tri-radial form with a perfect bilateral symmetry. It was attained by (1) complete elimination of the ancestral r. post. and r. ant. rays through atrophy of their superradials and incorporation of their inferradials with the anal structures; (2) restoration of the stem to the plane of the anal tube; along with these occurred (3) the equal development of the axil-arm system on either side of the median arm; and (4) the permanent detachment of l. ant. inferradial from the ray by the meeting of the two large radials above it.

Such is the general type of the prevailing post-Ordovician forms, of which there is herein illustrated a series of typical Silurian and Carboniferous species. Orienting the crinoid in the usual way by viewing it from the anal side, the large anterior radial would be said to lie at the right side, and the left posterior at the left; each supports a many-branched arm, and between them is the left anterior, median radial supporting its arm of different type which is frequently unbranched. But for descriptive purposes, on account of the very peculiar mode of branching of these side arms in Silurian and later forms, it has been found more convenient to view the crinoid from the conspicuous median arm, and while retaining the designation of the arms themselves as l. post. and ant., according to the conventional plan, to reckon direction of their component parts from the median arm outward and around toward the anal side. Therefore from this view the anterior arm lies toward the left, and its branches as seen from the exterior diminish in size from the median arm stemward, such branches as are given off in that direction being termed in the literature "adanal," "outer," or "left," while those that lie toward the median arm, and therefore away from the stem and anal side, are termed "abanal," "inner," or "right." In describing the corresponding details of the arm at the opposite side (left posterior) these "right" and "left" directions are reversed.

The Axil-arm System

With this distinction borne in mind, some of the confusion in the descriptive literature is clarified, and we are now in a position to give a more detailed account of the mode of branching which was finally attained in the family, and which forms one of its most striking characters. In this later and final stage of arm structure the axil-arms, borne upon the outer and shorter faces of the main-axils, are composed of a succession of brachials in series of two or more each, called for convenience Alphabrachs, Betabrachs, Gammabrachs, etc. The distal, or upper, Alphabrach is an axillary, of which the inner articulating face bears a ramule, rarely seen; the outer face carries the main arm with the second series of brachials, Betabrachs, the upper or distal one of which gives off from the outer side an unbranched ramule, and from the inner the third series of brachials, Gammabrachs; and so on alternately—a ramule and then a continuation of the main arm, until the latter terminates with an equal bifurcation.

The outer ramules of each axil-arm always tend to be larger than the inner, as explained by Bather, so that the first ramule on the extreme outer, or adanal, side, with the Betabrachs that support it, becomes the most conspicuous branch of the arm, while the rest of the arm is correspondingly diminished and sometimes completely hidden from view. Thus it results that the Alphabrachs, the outer Betabrachs, and the outer ramule appear to form a continuous arm, often all that is seen from the exterior, while the inner Betabrachs, Gammabrachs and the series which follow them, if any, although morphologically the main element of the axil-arm, are often inconspicuous or even invisible. In the closed condition of the arms as usually seen in the fossils, they are frequently folded under the adjoining outer ramules, while the latter constitute the visible axil-arms, lying parallel and gradually diminishing in length and thickness toward the anal side and stem. Especially the first or lowest ramule, given off by the axillary Alphabrach toward the median arm, is usually obscured in form C, but is more freely exposed in the later species of Form D.

The axil-arm is a direct development from the equally divided, ramule-bearing arm of Form A, as may be seen by comparing it with the arms shown in figures 9, 11, and 13 of plate 28, in which the alternate ramules from about each second or third brachial reach the full height of the crown—the essential difference being that in the earlier form both rami of each arm branched that way, and the main branch maintained its preponderance over the ramules. From the symmetrical ray of Form A, with its two equal and similar rami composed of successive ramule-bearing brachial series, and dividing upon an equal-faced primibrach, to the fully developed axil-arm system of Form C, borne upon an unequal-faced primibrach, with nothing of the original arm-divisions left visible but a set of closely packed parallel

ramules, seems a long jump. But the intermediate stage is beautifully shown in the Gotland species, *Calceocrinus gotlandicus*, where a stemward diminishing series of rami similar to those of Form A is combined with a succession of unequal-faced axillaries following an unsymmetric primibrach. The three stages are shown in the accompanying text-figures.

If in *b* the left ramule of each ramus enlarges, and all parts to the right are dwarfed and hidden behind it, as in certain species of Form C, the arm system of *c* will result.

The prime morphological factor at the bottom of these modifications is the change in shape of the lateral axillary primibrach (IAX) from equal- to unequal-faced, from which the unequal division of the ray necessarily follows. This actually begins with Form B, in the typical species of which the articulating faces of the primibrach differ in length about as 4 to 3, giving rise to an unsymmetric division within the ray. Therefore we have in this a third broad character by which Form A is differentiated from all the rest of the family.

FIG. 2. Three Stages in Development of Axil-Arm

- a. *Cremacrinus tubuliferus*. Equal rami from symmetric IBr.
 b. *Calceocrinus gotlandicus*. Unequal rami from unsymmetric IBr.
 c. *Calceocrinus foerstei*. Axil-arm system, with only parallel ramules visible from short faces of IBr and successive axillaries.

Other Progressive Characters

Thus the axil-arm as a post-Ordovician development is a most conspicuous character marking the evolutionary succession of the family in geological time. It is now a further striking fact that there is correlated with it another character more often seen because it appears in the calyx plates, which are the most frequently preserved in the fossils.

That is the greater or less separation of the two large simple radials by the connection, or want of connection, of the upper and lower segments of the l. ant. radial. The change was incident to the transverse extension of those radials to form the support of the axil-arms, so that the two structures underwent a parallel progression. In all the Ordovician species the segments were broadly connected, the inferradial being an elongate, quadrangular plate; this was constant in Form A, of which some species passed over into the Silurian. In Form B of the Silurian the connection was maintained, with a tendency to modification by the narrowing of the inferradial at the top, leading toward an acute-angled apex. In the Silurian Form C the complete change was effected by the transformation of the inferradial from a quadrangular to a triangular plate, and the consequent severing of connection between the two segments. Exceptionally as a survival the Ordovician type persists in this form, as in *Calceocrinus foerstei*, *C. interpres* and *C. pinnulatus*; and rarely in other species the two segments may touch by their apices. Finally with Form D in the Devonian and Carboniferous the separation of the segments by the interposition of the simple radials was completely established, and became a constant character of the genus *Halysiocrinus*.

The evidence on this point is remarkably decisive. Among upwards of 400 specimens of Form D from the American Devonian and Lower Carboniferous in which these plates are visible, there is not a single case in which the left anterior radial is not completely and widely separated into two triangular segments, with the anterior and left posterior radials meeting between them. These are distributed as follows: Helderbergian 1; Oriskany 1; Onondaga 4; Hamilton 10; typical Lower Burlington 6; Upper Burlington 37; Keokuk 58; the remaining 300 are from the Knobstone, or New Providence shale, equivalent to the Lower Burlington in part, in which at several localities the flattened cups of united radial plates and detached anchylosed basals occur in great numbers.

Thus it is seen that the two structures, the axil-arm and the divided segments, arose together, and continued in conjunction to the end of the family. The presence of the two segments unconnected proclaims the axil-arm almost without exception, which is the important fact because nearly always in evidence.

The axil-arm system of branching as above described is modified in the Gotland species, *Calceocrinus pinnulatus* of Bather; here the arms are composed of two main trunks with cuneate brachials, from each of which a pinnule is given off at the longer side, thus forming two similar rows of alternating pinnules, or ramules, after the manner of *Decadocrinus*. This is so different from all other species in the family that it will undoubtedly, as Bather has suggested, form the type of a new genus.

The branching of the median arm is also a feature of considerable variability. Dr. Bather in the Crinoidea of Gotland, page 72, stated that "the l. ant. arm gradually bifurcates at a greater distance from the cup, and eventually ceases to branch altogether." Examination of the American material, with the important additions now available, shows that the modification in this character is not attended by any such regularity.

In the Ordovician Form A the arm is either simple or bifurcates once at varying heights. In Form C, while the arm in most species as herein described is unbranched, there is one from the same locality in which it bifurcates three or four times. In the Carboniferous species the tendency in regard to the branching of this arm is the reverse of that above supposed; *Halysicrinus dactylus* from the Burlington, for example, branches once and then again in one ramus, producing 3, or perhaps 4, finials; while in *H. nodosus*, the most prominent species from the later Keokuk, represented in the collection by numerous specimens, the median arm divides twice and three times, and sometimes even once more, to a maximum of 8, or 10, finials. Thus there is a progression in this character from a median arm simple or with a single bifurcation in the Ordovician, to one of three or more bifurcations in the Carboniferous.

In a similar way the primary branches of the lateral arms increase in number from the 3 of Form B to the 5 or more axil-arms of Forms C and D.

So also in regard to the form and proportions of the flattened cup as seen from the left anterior side. In Forms A and B it is usually subquadrangular, about as high as wide (pl. 28, figs. 5a, 14); it changes in many species of Form C to a trapezoidal form wider than high, and widening toward the proximal margin (pl. 29, figs. 7a, 12a); in mature specimens of Form D the cup has spread out transversely until the margin at the hinge may exceed the height as 3.5 to 2 (pl. 30, figs. 8, 9, 10).

Thus there were at least six definite phylogenetic modifications from the Ordovician to the Carboniferous, which were more or less concurrent, viz:

1. Stem from unsymmetric lateral to median position.
2. Number of basals from 4 to 3.
3. Number of lateral arms from 3 to 2.
4. Segments of l. ant. R from connected to disconnected.
5. Median arm from simple to many-branched.
6. Lateral arms from equal bifurcation to axil-arm system.

Now these progressive modifications that have occurred in the phylogeny of the family are paralleled and recapitulated to some extent in the life history of the individual, as will be seen in the description of some of the later species. While not absolutely uniform, but marked by some exceptions and recessions, yet in general the facts indicate a well defined progression within this family from the early to the later geological stages comparable to that which occurred from the young to the mature stages of individual growth in the later forms.

The Anal Tube

The anal tube, although often concealed from view by the posterior arm branches, is usually large, and is composed on the posterior side of massive plates, somewhat resembling brachials; they do not entirely enclose the tube, but are crescentic in section, outwardly curved and leaving a space at the anterior side which is filled by an integument of small plates, similar to what has been described in *Catillocrinus* and *Synbathocrinus*.¹ The anal α is always a large, convex plate, constructed like the tube plates. The statement of Wachsmuth and Springer² that the tube extends to the tips of the arms and perhaps beyond their limits, was doubted by Bather (Crin. Gotl., p. 71), but is confirmed for the Silurian species by the new material (pl. 29, figs. 7b, 8b, 15b). Of this the most remarkable example of all is the new species *Cremacrinus tubuliferus*, in which the tube is not only of great size, but may bend outward like a bow, so that for a part of its length it is free from the arms.³

Shifting of the Stem

Aside from the reversal in direction, the shifting of the stem in this family is a remarkable phenomenon, which I think has not been noticed heretofore by authors. Correlated as it is with other changes of first rank, it becomes a character of much significance in the evolution of the group. It has to do directly with the equilibrium of the organism, which is so notable a feature among paleozoic echinoderms.

In the 5-rayed ancestral form, the plane of bilateral symmetry is from the anal tube to the anterior ray, leaving two arm-bearing radials on each side of it. When in connection with the bending of the crown upon the base the right posterior arm is eliminated, and the reversed stem falls into its place, the left anterior arm becomes the dominant factor on the anterior side, and the plane fixed by these two bisects the calyx so as to leave an arm and the anal tube on one side to balance the two arms remaining on the other. With the elimination of one of these two arms in the next stage, this balance would be disturbed, but for the fact that it is at the same time restored by the stemward torsion of the parts from the right to fill the vacancy, and the consequent shifting of the stem to a median position directly in front of the anal tube, thus establishing a plane of perfect bilateral symmetry with one large arm-bearing radial on either side. This being the condition of stable equilibrium, it is maintained to the end of the group. Thus it is seen that the location of the stem in its two positions has a definite connection with the changes accompanying the progressive modifications within the family, and marking the broad separation of Form A from all other forms.

The Hinge

The outstanding character of this family, by which it is broadly set off from all other known crinoids, is the mobility of the crown upon the base incident to the connection of the radials with the consolidated basals by muscular articulation instead of the usual suture, by

¹ Springer, On the Family Catillocrinidae, Smithsonian Misc. Coll., Vol. 76, No. 3, 1923, pp. 17-20.

² Revision of the Palaecrinoidea, pt. 3, 1886, p. 281.

³ See also Miss Goldring's description of the tube in *Halysicrinus secundus*, Devonian Crinoids of New York, 1924, p. 339, pl. 41, fig. 6.

means of which the crown could swing back and forth like a door upon its hinge. The anatomical feature most frequently discussed in relation to such movements of the crown toward or from the stem, as has been very fully described in Dr. Ringueberg's excellent paper on the Calceocrinidae, is the externally conspicuous hinge-like joint formed by deep grooves along the straight transverse edges of the consolidated basal and radial plates, which "has externally the appearance of a gaping fissure, but is closed internally by the exact approximation of the inner edges of the grooves." The fissure is formed by the articular margins of the l. ant. inferradial and adjoining radials meeting the apposed basals, bordered on either side by a broad and deep fossa, and divided by a straight line of contact, the approximated edges of which may form a fulcral ridge. Descriptions of this hitherto have dwelt chiefly upon the articular surface as usually seen from the anterior side, which occupies an equal space upon the faces of the respective plates. These spaces are often filled with numerous small, irregular plates, which have been described by Ulrich and by Bather, the latter under the term "supplementary plates," with several good figures, and which are here shown by figure 20a, on plate 28. They are of a somewhat scaly appearance, not meeting by suture, and have a darker color and denser composition than the other calyx plates. In some specimens similar plates are also observed in the gaping articulations of the axillaries or brachials. Their office is not clear. That the hinge represents a muscular articulation, with the usual combination of muscle and ligament, there can be no doubt.

This structure would provide the power for one movement of the crown. But to produce a complete up and down motion from a recumbent to a partially erect position and vice versa, there must be a reciprocal mobility and muscular control between basals and radials at the opposite side. Abundant evidence of this is seen in the fossils, sometimes in well-preserved crowns where the base stands at a considerable angle from the radials, but especially in the numerous weathered specimens of consolidated bases and united radials, separated from one another along the line of the basi-radial contact. Instead of a sutural union between the basals and the two large inferradials at the posterior side, the edges of these plates are rounded or beveled, and there are marks of muscular attachment on the inner surfaces similar to those along the external fissure. Thus the opposing musculature is furnished for motion by way of flexure or extension either way.

The mechanism of the hinge can be better understood by reference to the figures on plate 30, part of them based upon specimens composed from the detached elements above mentioned, and some upon crowns with the parts in position as found in the fossils. This structural material is chiefly of the abundant species *Halysiocrinus perplexus* Shumard, from the basal Lower Carboniferous. Figures 4 and 5 show the anterior aspect of the calyx when the articulated parts are opened as far as they will go, almost to the same plane, with the transverse grooves forming the two sides of the gaping fissure which lodged the exterior muscles actuating the movement in that direction. The floor of the grooves next to the dividing line is marked by bands of striae vertical to the margin. In figure 6 the exact opposite is seen, showing the disposition of the parts at the interior, and their appearance when fully extended. The joint-faces of the median and two lateral arms are shown, with their muscular markings, and by contrast the rounded edges of the two large radials which would be in apposition to the corresponding edges of the basals when closed, but without any sutural union. Here the relation of all the plates composing the calyx may be seen, as well as the cavity which lodges the viscera. From the margins of the apposed basal and radial plates wide bands of vertical striae extend inward, marking the attachment of powerful muscles by the contraction of which the crown was closed down upon the base and brought into the position in which it is usually found, as in figures 17, 18, 19. (Best seen with a low magnifier.)

Starting with these two examples, we can readily trace the relation of the parts in several positions incident to different degrees of opening or closing. In figure 16a, from a complete crown, in the most frequent position of rest, the basals are firmly closed against the radials

and the inferradials of the lost arms, here meeting underneath the anal series to form a sub-anal piece. In figure 7 they are opened to about a right angle; while in figure 15 the very unusual condition is seen of the crown widely opened out almost to an erect position upon the stem. Some other figures, 8, 9, 10, are given showing in part the same structures, but more especially to illustrate the change in shape of the calyx due to growth, from elongate with nearly parallel sides in the younger individuals, to relatively short and widely spreading downward toward the base in the older.

Mobility of the Crown

With the establishment of the tri-radial symmetry in the group, the organism after an early beginning seems to have finally adapted itself to the recumbent habit as the position of rest, from which it could readily change as occasion required. The widely gaping articulations between radials and arms, and between many of the successive brachials, secured in addition to the larger movement upon the base the maximum of flexibility above the hinge line, whereby the arms could be flexed or extended by the contraction of their own muscles. Change of position of the crown was further facilitated by the curvature of the stem at its proximal end, as shown by the presence of wedge-formed columnals, by the articulation of which it could be bent or straightened to some extent. The closed condition is by far the most frequent as found in the fossils. Evidence of habitual pressure of the crown upon the stem is not uncommon in the form of deep indentations upon the anal plates, or in the Ordovician form upon the interray to the right of them, as shown in figures 17 of plate 28; 24 of plate 29, and 14, 16a of plate 30. In others equally firmly closed, the indentations are not present, the pressure being relieved by the slight proximal bending of the stem.

Excellent illustrations of the adaptability of the crown to these movements are afforded by figures 5 of plate 28 and 17, 18, 19 of plate 30, with the crown lying closely recumbent upon the stem, and figure 21 of plate 28 in which it stands almost at a right angle, and figure 15 of plate 30 before mentioned in which it is extended almost in the same line.

The expression "bending of the crown upon the stem" recalls the structure of the preceding Heterocrinid derivative, *Myelodactylus*, to which the same term might be applied. There, also, is some diminution of radials due to pressure or a cramped position, accompanied by a specialization of stem in its way equally remarkable with this. But the change in direction of the crown is accomplished by simple curvature of the stem; there is nothing in the structure analogous to the hinge and the consequent mobility of the crown upon the base.

Mode of Life

The question arises, what is the relation between the extreme departure of the crinoids of this family from the usual habitus of crinoids and their mode of life? Primarily, all such modifications have to do with the food supply and the mode of obtaining it. In some genera, as *Torynocrinus*, the arms are bent from the calyx to one side, from which presumably the food-bearing currents came. In others, such as *Monobrachiocrinus* and *Abrachiocrinus*, the arms have progressively been reduced in number and finally eliminated entirely, because, as is supposed, the crinoids lived in an environment where the food was brought to them by the currents in such quantities that arms were no longer needed to gather it.

Here, in the form now before us, not only the arms were affected by the adaptation to unusual conditions, but the entire crown above the base was forced out of its natural upright position on the stem. It has hitherto been generally assumed that the crown had become habitually pendent upon the stem, so that while the stem itself remained upright, the normal position of the crown was reversed, and the arms pointed downward, with, however, some power in the crinoid, by means of the hinge between basals and radials and the muscular articulation, to raise itself to a horizontal or partially erect position. This supposition was

expressed by statements such as "the crinoid hangs downward along the column," and "the upper part of the crinoid could be bent upward and brought into an erect position." Such a lifting of the entire crown upon so small a fulcrum would seem to call for a muscular equipment rather greater than is indicated by the evidence in the fossil state.

A more recent interpretation has been given by Jaekel (*Phylogenie und System*, 1918, pp. 86-88), which seems to accord better with the observed facts. He considers these crinoids, which he groups as a suborder Calceocrinites, to be reef-dwellers, living exposed to tidal currents, the stem lying recumbent upon the bottom, where when at rest the crown was closed down at one side like a barnacle. When feeding, the arms opened out toward the current, for which but little movement of the crown as a whole upon the base was required. When the tide turned, the arms would readily close down, the larger ones serving as a protection for the more delicate parts against injury from the ebb-current. Thus the crown habitually was in a horizontal position to start with, from which only a relatively small movement was needed beyond the ordinary opening of the arms to enable the organism to obtain food from above like other crinoids. The theory is graphically presented by Jaekel's figures, 81, 83, representing the closed and open conditions.

This interpretation presupposes a sessile condition by means of a distal attachment of the stem to the reefs. Some evidence of this is seen in specimens from Dudley, England, in which the stem is sometimes found rather short with a foreign object attached. In the Tennessee specimens it is not so evident, the few stems preserved being fairly long, and apparently tapering towards a point.

With the recumbent position as that of rest, in which the crinoid habitually reclined except when feeding, the diminution in size of the arms toward the posterior side, which became such a marked character of the post-Ordovician forms, is accounted for as the result of a cramped position, pressure, and loss of motion.

Evolution in the Act

Returning to the evolutionary modifications exhibited in this group, the study of the material now in hand has brought to light some remarkable facts. The most striking movement within the family, next to the development of the axil-arm, was the elimination of the fourth arm from Form A, concurrent with the shifting of the stem back to its normal position in the plane of the anal tube, and the restoration of bilateral symmetry in the resulting 3-armed crinoid, thus establishing a condition in which all the other forms differed in a large way from the typical Ordovician form. Such a change in characters of major importance, involving some of the larger factors in the anatomy of the group, would be expected to occur in a gradual way, scarcely discernible in individual cases. Hence we were prepared to observe results rather than the process by which they were obtained.

When preparing the finely preserved Tennessee material belonging to this form, I was surprised to find in some instances a marked difference in the size of the lateral arms. Upon following this up in seven of the nine specimens herein described under three species, it was disclosed that in every one of them the right anterior arm was smaller than the anterior arm adjoining it. The diminution was in various degrees, in some cases amounting to well-defined atrophy by which the arm was reduced to less than one fourth the size of its adjacent fellow. The reduction is decisively shown by the figures on plate 28, especially figs. 3*b* and 4*b*, in which the smaller arm occupies but a small part of the space on that side; in the former specimen its branches are like very short pinnules, in contrast to the long and robust ramules of the other arm alongside; and in the latter it lies against the large anal tube with scarcely more than a remnant of its primitive dimensions. In another species, figure 7 of plate 28, having both arms spread like a fan in excellent position for observation, the area of the anterior arm is about five times that of the dwarf. In every case, with the two lateral arms of the anterior side lying side by side for comparison, the difference in general dimensions,

in width of brachials, length and stoutness of ramules, and where observable in number of branchings, is perceptible at a glance without the need of measurements.

Now the arm which is thus dwarfed is always the right anterior, the one which is destined to disappear in the course of the modifications which produced the 3-armed form of subsequent and some contemporaneous types, and the manner of the change by which that result was attained is apparent before our eyes. Upon first discovering this fact, I assumed that the change took place wholly in the Silurian; but examination of the Ordovician species, especially some exceptionally well preserved specimens from the Trenton of Kirkfield in Ontario, showed that the tendency to atrophy of the fourth arm began much farther back. There are eight specimens of *C. articulatus* Billings or a closely allied species, in which the two adjacent arms are seen, and in all of them the diminution of the right anterior arm is almost as great as in the Tennessee species, as will be seen in figures 9 to 13 of plate 28. It is observable in varying degree in some of the other Ordovician species in which the arms can be compared, and in some the fourth arm seems to be as large as the others. Bather (Crin. Gotl., p. 61) observed that this arm was smallest in *Castocrinus*.

Thus it is evident that we have in this instance a concrete example of the process by which the change from one genus to another was accomplished in the course of geological time. The atrophy of the two arms was clearly the direct result of pressure incident to the reversal and recumbent position of the crown against the stem, and when the axil-arm system was established, the same influence was reflected in the reduced size of the branches in the stemward direction.

Distribution

So far as known, Form A does not occur in the European areas. Its presence in the Silurian of Tennessee is a new and unexpected discovery. It is in the Silurian development of the family, as evidenced by the forms included under Form C, that the most interesting facts relating to distribution are found, bearing on the relation between the European and American faunas. About seven species of *Calceocrinus* have been recognized from Gotland, and there are at least two or three from England. Of the American species only those from the Rochester and Waldron formations of New York and Indiana have been heretofore figured, and these are mostly not in good condition for comparative study.

The Beech River formation of Tennessee has yielded remarkable material mostly in fine preservation, embracing three well-marked species suitable for comparison with European forms. Of these two, *C. foerstei* and *C. bassleri*, are of similar types to certain Swedish forms, but still are differentiated by good specific characters. Of the Swedish species, *C. gotlandicus*, *C. pinnulatus*, and *C. nitidus* are thoroughly distinct. Thus there are left *C. pugil*, *C. tucanus* and *C. interpres* on the other side, and *C. foerstei* and *C. bassleri* from here, through which a migrational connection may confidently be traced.

No member of the family has yet been found in the European Devonian or Carboniferous, but in America it occurs in most of the principal formations from the Ordovician to the middle of the Lower Carboniferous, viz.: Black River, Trenton, of the Ordovician; Rochester, Waldron, Laurel, Brownport of the Silurian; Helderbergian, Oriskany, Onondaga, Hamilton of the Devonian; New Providence, Burlington and Keokuk of the Lower Carboniferous.

It is recognized that among the known species belonging to this family, from all three of the countries where it occurs, there are some which might form the types of new genera. One of these has been suggested by Dr. Bather, but without proposing a name. Mr. Ulrich has defined several others, for which he proposed names orally at the New Haven meeting of the Paleontological Society in 1912. It is not my purpose to enter upon these questions, which it is hoped will be taken up again by those authors with a view to publication.

Nomenclature

The broader taxonomic questions relative to this family, as presented by the hitherto published literature, are somewhat complex, and have given rise to considerable difference of opinion. They have been discussed by Wachsmuth and Springer,¹ Ulrich,² Ringueberg³ and Bather.⁴

The first name to be employed for a member of the family was *Calceocrinus* by Hall in 1852,⁵ based on the common semicircular basal piece, to which no specific name was given. In 1860⁶ he proposed the genus *Cheirocrinus*, with *C. chrysalis*, a Silurian species, as genotype. This name being preoccupied, Meek and Worthen in 1869 and 1873⁷ proposed *Eucheirocrinus* to replace it, with the same species for genotype. At the same time they described and figured a species as *Calceocrinus bradleyi*,⁸ and in connection with it proposed the family name Calceocrinidae, saying: "It is evident that this remarkable genus differs so widely from all the other known types that it must be regarded as belonging to an entirely distinct and unnamed family, which might be called Calceocrinidae." In 1879⁹ Hall published a Silurian species as *Calceocrinus stigmatus*, with a description, figures, and diagram which he said "illustrate the generic structure."

In 1886 (op. cit., p. 107) Ulrich proposed the genera *Cremacrinus*, with the Ordovician *C. punctatus* as genotype; *Deltacrinus*, based upon *Cheirocrinus clarus* Hall, of the Middle Devonian, as genotype; and *Halysiocrinus* with *C. dactylus* Hall, of the Lower Carboniferous, as genotype. He regarded *Calceocrinus* as not established, on the ground that it was described without a type species, and therefore proposed the name Cremacrinidae for the family instead of Calceocrinidae, although at the same time he did not entirely discard *Calceocrinus*, but listed under it four species which all have the family characters.

In 1889 Ringueberg (op. cit., p. 392) proposed the genus *Castocrinus* for the reception of the Ordovician species described by Billings and W. R. Billings, and some new species of his own; also *Proclivocrinus* for a Silurian form clearly identical with *Eucheirocrinus*. He also redescribed *Calceocrinus* from specimens identical with Hall's original, together with a new species called *C. typus* showing the correlated characters, to serve as the type species. These forms were illustrated with good figures and diagrams, by which the generic structures were clearly shown.

Wachsmuth and Springer in part 3 of their Revision of the Palaeocrinoidea, 1886, pp. 273-281, recognized *Calceocrinus* as valid. Bather in Crinoidea of Gotland, 1893, pp. 54-99, accepted *Castocrinus*, *Eucheirocrinus*, *Halysiocrinus*, and upheld *Calceocrinus*, under which he described seven species from the Swedish Silurian. Both of these authors, as well as Ringueberg, adopted Meek and Worthen's family name Calceocrinidae.

When preparing the revision of the chapter on the Crinoidea for the Zittel-Eastman Textbook of Palaeontology, edition of 1913, I recorded *Eucheirocrinus* and *Halysiocrinus*, but included *Cremacrinus* and *Deltacrinus* in place of *Castocrinus* and *Calceocrinus*. Re-examination of the facts, with more detailed study of the material in hand than I was able to give it at that time on account of the pressure of other work, has led me to the opinion that the arrangement then adopted should be modified in some respects.

Cremacrinus as proposed in 1886 was described as having a dorsal or centrodorsal (median) and two lateral arms, symmetric, one on each side. No mention was made of the

¹ Revision of the Palaeocrinoidea, pt. 3, 1886, pp. 273-281.

² 14th Ann. Rep. Geol. Surv. Minnesota, 1886, pp. 105-112.

³ Ann. New York Acad. Sci., 4, 1889, pp. 388-406 (separate 1-19).

⁴ Crinoidea of Gotland, 1893, pp. 54-99.

⁵ Pal. New York, 2, 1852, p. 85, figs. 5, 6.

⁶ 13th Rep. New York St. Cab. Nat. Hist., pp. 121-122.

⁷ Proc. Acad. Nat. Sci. Phil., 1869, p. 173; Geol. Surv. Ill., 5, 1873, pp. 443, 502.

⁸ *Ibid.*, Geol. Surv. Ill., 5, 1873, p. 502, pl. 14, fig. 9.

⁹ 28th Rep. N. Y. St. Mus. Nat. Hist., 1879, p. 147, pl. 19, figs. 9-11.

fourth arm, which is clearly specified in the description and figure of *Castocrinus*. But it now appears upon further examination in the light of subsequently acquired material that the fourth arm was actually present in the type specimen of *Cremacrinus punctatus*, now in the U. S. National Museum. Therefore the species of *Cremacrinus* and *Castocrinus* are generically identical, and *Cremacrinus*, being the earlier in date, will be valid for the species falling under Form A. This conclusion is reinforced by the further fact, not mentioned in the description of either, that all species referred to the two genera in which it can be seen have the stem in the position at the right of the anal tube, which is such a fundamental character of Form A.

That the fourth arm was not mentioned in the original description of *Cremacrinus* was owing to the imperfection of the type specimen upon which the genus and species were founded. This was apparently well preserved, and showed the median and two lateral arms, one on each side, as described. At the right of the anal tube, however, is a vacant space about the size of a lateral arm where the specimen had been attached to the rock, and from the information now afforded by several other specimens of the same species since obtained, it is evident that the missing arm was imbedded in the matrix and pulled off in collecting (pl. 28, figs. 16, 16a). With our present understanding of the structure, traces of the fourth arm may now be recognized in the type, but with no other guide than the imperfect original it was natural that the author should describe only what he could see, and frame his diagnosis accordingly.

For the forms which had formerly been referred to *Calceocrinus*, and which would now fall under Form C, preference was given to *Deltacrinus*, which as before stated was proposed to take the place of *Calceocrinus*, rejected because lacking a type species. It was based upon *Cheirocrinus clarus* Hall for genotype, a Middle Devonian species represented by the single type specimen from the Hamilton of New York. The species is so completely non-characteristic of the Silurian forms belonging to Form C, that I cannot reconcile the genus founded upon it with the highly distinctive characters for which they are so conspicuous. It entirely lacks the arched axil-arm system by which Forms C and D are distinguished from all other crinoids (text figs. 1C, D, and pl. 29, figs. 7, 8, 9, 14, 20). This cannot be shown better than by the remarks of the author himself in his paper of 1886, from which I take the following extracts:

On page 110, under *Halysiocrinus*, it is said:

"In the formation of the dorsal side of the calyx this genus is precisely *Deltacrinus*. The only difference so far detected is found in the number of arms. In *Cremacrinus* we have three primary radials, and in *Deltacrinus* five; while the species for which the above generic term is proposed have eleven, one large central arm on the dorsal side and ten smaller ones, the first pieces of which project abruptly outward and extend in a curved series transversely around the ventral side. Their inner ends articulate with the ventral arch."

By comparing my diagrams of Forms C and D in text-figure 1, and various figures of *Calceocrinus* on plate 29, it will be seen that they have exactly the characters which are above ascribed to *Halysiocrinus* as distinguishing it from *Deltacrinus*. That the two forms cannot go together is further emphasized by the author in the continuation of his discussion on page 112:

"There are at least three American species that have the characters ascribed to *Halysiocrinus*. These are the *C. dactylus* and *C. nodosus* Hall, and the *C. wachsmuthi* M. & W., the first and last from the Burlington limestone and the second from the Keokuk Gr. Besides these the *Cheirocrinus Gotlandicus* of Angelin is an unquestionable member of the genus In the construction of the body and in the possession of a strong dorsal arm, these species do not differ from *Deltacrinus*. The lateral arms, however, differ conspicuously from all the species of both *Cremacrinus* and *Deltacrinus* in being much more numerous, subequal, and extending completely around the ventral side The ventral arms give to these crinoids a different appearance from that presented by the more simple species of *Cremacrinus* and *Deltacrinus*. In my opinion they constitute an important deviation from the type of those genera, and fully warrant generic separation."

That is to say, in the opinion of the author himself, species having the axil-arm system, borne on short faces of successive axillaries, arching around to the anal side—Forms C and D of my diagrams—do not belong under *Deltacrinus*, and should be separated from it generically.¹

Therefore, whatever else *Deltacrinus* may be, it cannot include the Silurian species which come under Form C. They are, however, clearly covered by the genus *Calceocrinus* Hall, if not as described and figured by him in 1879 under the name *Calceocrinus stigmatus*, then certainly as amended and redescribed by Ringueberg in 1889 under *C. halli* as the original unnamed species of Hall, and a new species, *C. typus*, showing the correlated characters, for a genotype. I have no doubt that Ringueberg's identification of Hall's original *Calceocrinus* was correct, and that it is congeneric with *C. halli* and *C. typus* (pl. 29, figs. 17, 18, 19). There are two known generic forms of the hinged crinoid in the New York Silurian, and they can be distinguished by the base alone, without the aid of other characters:

1. With 4 basals; in which the two left basals, those in the middle next to the hinge, retain the characters of the Ordovician form, being in contact with the stem by their narrow points, their outer sides making a sinuous curve, and being still divided but with a tendency to coalesce and their dividing suture to become obsolete. This is Hall's *Cheirocrinus* (*Eucheirocrinus* M. and W.). His figure in the 13th Report, 1860, page 123, is incorrect in that it does not show the two small basals touching the stem-facet or divided, though it does indicate the curving of the sides; but both the missing characters appear in the type specimen, as here figured, plate 29, figure 1c, confirmed by numerous other identical specimens from the same locality, some of them figured by Ringueberg (his pl. 10, figs. 6, 7b, 13), and herein (pl. 29, fig. 2). Errors in the original description or figure due to incorrect observation or imperfect material cannot control the definition where the actual fact is shown by the type and identical specimens. The habitus of this base is thoroughly characteristic, recognizable at a glance; any New York Silurian species with that type of base must be *Cheirocrinus*, and any one with a different type of base cannot be *Cheirocrinus*.

2. With 3 basals; in which the two left basals are detached from the stem and fused into a single triangular plate. Any species with such a base cannot be *Cheirocrinus* as determined by the type, regardless of how the name may have been applied by authors, including Hall himself, who listed species having both forms of base under *Cheirocrinus* of the 13th Report, and shifted them to *Calceocrinus* in the 28th Report. But such a base is precisely that of *Calceocrinus halli*, *C. typus*, or any of the numerous species falling under Form C (perhaps excepting one in Gotland). Hall's original *Calceocrinus* must be one or the other of these two forms. His figure (Pal. New York, 2, pl. 85, fig. 5) shows a base with a triangular plate fitting into a much larger somewhat lunate-shaped piece, having the column-facet widely separate from the triangular plate (pl. 29, fig. 18). The description speaks of only two plates visible, but adds that the plates are so closely ankylosed as to obscure or obliterate the lines of suture.

The fact that he mentions only two plates is not material, for we know there must have been a suture from the apex of the triangle through the stem-facet, as is the case in every species of the entire family; so that the lunate-shaped piece into which the triangular plate fitted necessarily consisted of the two large basals, making the three which were correlated with the axil-arm system of Forms C and D. It is immaterial, because the undivided triangular plate is the decisive character, which excludes *Cheirocrinus*, defines what the original *Calceocrinus* is, and determines the fact that it must be congeneric with *C. halli* and *C. typus*. It cannot be anything else.

Therefore Ringueberg's identification of *Calceocrinus* as his *C. halli* was founded upon an unmistakable diagnostic character of generic value, clearly expressed in the original de-

¹ *Deltacrinus dactylus* and *D. nodosus*, Smithsonian Misc. Coll., vol. 76, No. 3, 1923, p. 19, pl. 5, figs. 20, 21, should be written *Halysiacrinus*.

scription and figure; and he was entitled to amend the genus by setting out the other correlated characters which go with it. Thus there can be no reasonable doubt of the validity of the genus as of one or the other of the above dates.

With the genus thus validated and recognized, it would seem that the family name Calceocrinidae proposed by Meek and Worthen in 1873, and adopted by De Loriol, Wachsmuth and Springer, Ringueberg, S. A. Miller, P. H. Carpenter and Bather, will have to stand.

From these considerations it follows that the four stages of the family herein discussed are logically expressed by the following genera, the relations of which, in their broader aspects, may be thus summarized:

Analysis of the Genera

- Radials 2 simple, 3 compound, connected with basals by muscular articulation instead of suture, admitting mobility of crown upon base as by a hinge, and consequent bending upon the stem to a pendent or recumbent position CALCEOCRINIDAE.
- Form A. Arm-bearing rays 4 (median and 3 lateral); stem at right of anal tube, and bilateral symmetry disturbed. IAx equal-faced; arms heterotomous; BB 4. Ordovician to Silurian..... *Cremacrinus*.
- All others. Arm-bearing rays 3 (median and 2 lateral); stem in plane of anal tube, and bilateral symmetry restored; IAx unequal-faced.
- Form B. Lateral arms unequally dichotomous, with few branches; BB 4; Silurian *Eucheirocrinus*.
- C and D. Lateral arms heterotomous, branching numerously with axil-arm system from unequal-faced axillaries; BB 3.
- Form C. R. post. and r. ant. inferradials separated by subanal piece; main branch of axil-arms may be hidden by outer ramule; segments of l. ant. R usually separated; fused basal usually triangular and narrower than hinge; median arm usually simple. Silurian..... *Calceocrinus*.
- Form D. Inferradials meet under anal π ; main branch of axil-arm always more or less visible; segments of l. ant. R. always separated; fused basal usually curved and about as wide as hinge; median arm frequently branching. Devonian to Carboniferous..... *Halysiocrinus*.

CREMACRINUS Ulrich*Plate 28*

Cremacrinus Ulrich, 14th Ann. Rep. Geol. Surv. Minnesota, 1886, p. 107.—*Castocrinus* Ringueberg, Ann. New York Acad. Sci., 4, 1889, p. 392.—Bather, Crin. Gotl., 1893, pp. 56, 61; Treatise on Zool., Echinoderma, 3, 1900, p. 148.—Zittel, Grundzüge Pal., 1, 1910, p. 152.—*Cremacrinus* Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 213.—Bassler, Bibliogr. Index, Bull. 92, U. S. Nat. Mus., 1915, p. 288.

Arm-bearing rays 4. Calyx unsymmetric. Stem not in plane of anal tube, but lying to the right of it. Median (l. ant.) arm simple, or branching once; the inferradial of its compound radial quadrilateral, connecting with superradial; primibrach always short. Lateral arms 3, two on ant. side and one on l. post., each dividing into two equal and symmetrical rami, heterotomous, bearing alternate ramules at intervals of 2, 3, or sometimes 4 brachials; primibrachs 2, the axillary with articulating faces equal. R. post. superradial supporting anal tube, and r. ant. superradial the r. ant. arm, which is usually smaller than ant.; the two superradials meeting above the stem-facet; r. post. and r. ant. inferradials separated from one another and from anal α by their respective superradials. Basals 4, all in contact with the stem; the two left BB entering into, or touching by their points the stem-facet; also much narrower than the others, and their outer sides making a reverse curve. Proximal margin of radials bordering hinge often strongly denticulate.

Genotype. *Cremacrinus punctatus* Ulrich.

Distribution. Ordovician and Silurian; America.

A widely distributed form of the Trenton or underlying horizons, occurring at various localities in Canada, New York, Minnesota and Kentucky, from which seven species have been described, and passing up into the Silurian in Tennessee, with four hitherto undescribed species herein illustrated. It is not known outside of America. The genus forms a section by itself, broadly differentiated from all others of the family by its 3 heterotomous lateral arms and asymmetric stem.

Cremacrinus ulrichi new species*Plate 28, figs. 1, 2*

Crown relatively large for the genus, about 40 mm. high in two specimens. Median (l. ant.) arm fairly strong at the base, about as long as the adjacent lateral arms, and tapering to a narrow point; consisting of about 7 brachials, long and slender beyond the short IBr; inferradial elongate rectangular, broadly connecting with superradial. Main rami of ant. and l. post. arms giving off from successive series of 2 or 3 brachials—Alphabrachs, Betabrachs, etc.—alternately at each side a large ramule the full length of the arm, composed of one short and several long ossicles, ending with an equal bifurcation, to a total of 6 or 7 ramules to each ramus. R. ant. arm much smaller than the others, and

its two slender rami having fewer branches. Lateral IBr 2. Anal tube not exposed. Stem in largest specimen 14.5 cm. long, diminishing distalwards; columnals about 2 mm. long except in the proximal part, where they become much shorter and cuneate in the curve.

This species is founded upon two crowns, one of which has the stem nearly complete. Both are exposed from the anterior side only, so that we are without information as to the character of the anal series and tube. The relative length and slenderness of the brachials impart a certain lightness and delicacy of construction which is in contrast to that of the accompanying species. In superficial appearance of this and the following species there is a considerable resemblance to the *Calceocrinus gotlandicus* of Angelin, but the fine illustrations of that species in Dr. Bather's work give no indication of any of the generic characters which are here so pronounced; its mode of arm-branching, however, represents an intermediate stage between the two equal rami of *Cremacrinus* and the complete axil-arm system of the typical *Calceocrinus* and *Halysicrinus*.

The specific name is in honor of the eminent geologist and paleontologist, Mr. E. O. Ulrich, of the United States Geological Survey and the National Museum.

Horizon and locality. *Eucalyptocrinus* zone of the Beech River formation, Brownsport group, Niagaran; Tuck's Mill, Decatur County, Tennessee.

***Cremacrinus tubuliferus* new species**

Plate 28, figs. 3, 3a, b, 4, 4a, b, 5, 5a, 6

Crown uniformly small, ranging in height from 20 to 23 mm. in five specimens. Cup elongate, quadrangular, but little constricted; average height 6 mm. to width at hinge of 4.5 mm. Median (l. ant.) arm short and slender, tapering to a narrow point and not reaching the height of the adjacent lateral arms; with few very long brachials, sometimes only 2 or 3 in all beyond the short primibrach; inferradial truncate, broadly connecting with triangular superradial. Main rami of adjacent (ant. and l. post.) lateral arms have from 6 to 9 axillary brachials each, nodose and prominent, giving off alternate ramules with very long ossicles beyond the first, reaching the full height of the arms, from successive series of two or three brachials—Alphabrachs, Betabrachs, etc.—ending in an equal bifurcation; thus there are 12 to 18 finials to each of these arms; r. ant. arm similarly divided into two rami, but greatly dwarfed in size, much narrower at the primibrach, shorter than the other arms and having fewer and smaller ramules. In one specimen where the anterior arm is 18 mm. long, the r. ant. arm bifurcates finally at about 9 mm.; and in two specimens having both arms perfectly exposed the area covered by the anterior arm is about four times that of the r. ant. Lateral IBr. 2. Anal series extended into a massive tube, exceeding in diameter any of the arms, and apparently equaling them in length, sometimes curving outward like a bow. Stem prominent at right of the tube, stout, with columnals beyond the proximal curve 1.5 to 2 mm. long; in one speci-

men it is about complete, attaining a length of 9.5 cm., and tapering to a narrow distal end.

The outstanding character of this remarkable species is the relatively enormous, curved anal tube. It is represented by a wonderful series of six specimens, four of them having the tube intact, which are all figured, in order to show that the dwarfing of the fourth arm is no mere sporadic occurrence but a definite process; a fifth, with the stem complete, has the tube plates displaced, and the sixth is fragmentary. Another striking character of this and some of the associated species is the great length of brachials in the median arm, and of the ossicles in the ramules—except the first which is always short—reaching sometimes five times their width; this would seem to make for lessened flexibility in the arms. In two of the specimens, figures 3 and 4, I have been able to free the crowns completely from the matrix, so that in both all the lateral arms may be inspected and compared, and the abnormal position of the stem clearly seen—thus furnishing a full exposition of the essential characters of the genus; in one the tube is visible from both sides.

Horizon and locality. *Eucalyptocrinus* zone of Beech River formation, Brownsport group, Niagaran; Tuck's Mill, Decatur County, Tennessee.

***Cremacrinus decatur* new species**

Plate 28, fig. 7

Similar to *C. tubuliferus*, but larger, the crown being 35 mm. high, with distal end incomplete. Median arm relatively stout, quite in contrast to that of the two preceding species; 2 long brachials of about 3.5 mm. are preserved beyond the short primibrach, and judging by the taper there were probably one or two more. Anterior arm has 6 axillary brachials in each ramus, giving off ramules with very long ossicles after the first alternately to a total of 12, and perhaps one or two more at the broken distal end; interposed brachials mostly in series of two, the upper one often long enough for a third; r. ant. arm less than one fourth the size of ant., and apparently branching only two or three times; IBr 2. Anal plates and tube not visible, only the anterior side being exposed. Stem stout, with long columnals, those beyond the proximal curve being 2.5 mm. long.

The species is based upon a single very perfect crown, slightly broken at the distal end, and imbedded at one side; if it had a curved anal tube it would stand rather close to the last species, but the strong median arm is characteristic. The difference in size of the anterior and right anterior arms is most pronounced, the area covered by the former being about five times that of the latter. It is from a much higher horizon than any of the other species, in a formation that has yielded a number of crinoids with markedly distinct characters.

Horizon and locality. Decatur limestone, Niagaran; near Rise Mill, Perry County, Tennessee.

***Cremacrinus simplex* new species**

Plate 28, figs. 8, 8a

Of a very different aspect from any of the preceding, in the small number of arm-branches, and their dichotomous bifurcation, which might well warrant a reference to a new genus. Crown small, 28 mm. in height. Median arm fairly

stout, reaching the full height of the crown; it consists of 4 very long brachials beyond the short primibrach, tapering to a narrow end; l. ant. inferradial quadrilateral, fairly wide. IBr of lateral arms 2. L. post. arm dividing on second primibrach into two rami, the inner one bearing a long branch on the second Alphabrach, and bifurcating on the second Betabrach into two equal branches; the outer ramus divides in a similar way, but begins on the first Alphabrach. Thus there are only 6 branches on that side, in contrast to the 12 to 18 ramules of other species, and these are rather of the dichotomous type than the heterotomous. The arm arrangement on the anterior side is different; the anterior arm divides into two large apparently unbranched rami, and the r. ant. arm, decidedly smaller than the former, has an unbranched inner ramus, while the outer branch, next to the anal series, divides again into slender finials scarcely more than half the length of the other arms. Brachials on both sides long and slender, similar to those of the median arm. Anal series strongly rounded, to the left of stem. Stem stout, asymmetric in position, with fairly long columnals, about 1.2 mm. beyond the curve.

The species is founded upon a single specimen, associated with the first two preceding; it is a complete crown, exposed on both sides, and having a short section of stem attached. It may be regarded as an intermediate stage between Forms A and B, the latter being distinctly foreshadowed in the arm arrangement, while it wholly lacks the ramule-bearing feature of the former. Yet the presence of the fourth arm, and the asymmetric position of the stem, determine its close relation to the Ordovician type, beyond question, thus furnishing another unmistakable link in the chain of evolutionary modifications so conspicuous in the history of this family.

Horizon and locality. *Eucalyptocrinus* zone of Beech River formation, Brownsport group, Niagaran; near Tuck's Mill, Decatur County, Tennessee.

***Cremacrinus articulatus* (Billings)**

Plate 28, figs. 9-13

Heterocrinus articulatus Billings, Geol. Surv. Canada, Dec. 4, 1859, p. 51, pl. 4, fig. 8.—*Castocrinus articulatus* Ringueberg, Ann. New York Acad. Sci., 4, 1889, p. 395, pl. 10, fig. 4.—Springer, A Trenton Echinoderm Fauna, Mem. 15-P, Geol. Surv. Canada, 1911, p. 25.

I am illustrating this species rather fully chiefly to show the incipient dwarfing of the r. ant. arm in the Ordovician stage of the genus. It has been known heretofore only from very scant material, consisting solely of the imperfect type specimen from Ottawa, Canada, which was refigured by Ringueberg. The single arm preserved shows very clearly the strongly nodose axillaries which constitute its leading character. Our knowledge of this rare form has been enlarged in recent years by the acquisition of a fine series of specimens from Kirkfield, in Ontario, in exquisite preservation, by which all the characters are brought out in detail. These are supplemented by an excellent specimen from an equivalent horizon in Kentucky. By a selection from this material the heterotomous structure of the two equal rami of the lateral arms, with their

5 to 7 ramules given off at intervals of 2, 3 or 4 brachials, is thoroughly shown, and the fourth arm, beginning the process of atrophy by which it is eventually eliminated, can be seen and studied. The area which it occupies is about one half that of the permanent lateral arms. The median arm is long and slender, with 17 to 20 short brachials, either ending beyond these in a narrow point or in an equal bifurcation of very slender finials. The cup is but little constricted, with height and width about the same. Surface in well preserved specimens somewhat punctate.

Horizon and locality. Trenton (Curdsville), Ordovician; Ottawa and Kirkfield, Canada, and Woodford County, Kentucky. This horizon is somewhat higher than that of *C. punctatus*.

***Cremacrinus kentuckiensis* (Miller and Gurley)**

Plate 28, figs. 14, 14a, 15

Calceocrinus kentuckiensis Miller and Gurley, Bull. 5, Illinois State Museum, 1894, p. 29, pl. 2, figs. 24, 25.

This species, of somewhat coarser construction than *C. articulatus*, is introduced with two specimens from the Trenton of Kentucky on account of the striking manner in which they illustrate the peculiar relation in this genus of the stem and anal tube. In figure 15 the tube occupies the middle space between the two arms at the posterior side, which in crinoids generally would be in the plane of bilateral symmetry. But that is not the case here, because the arm that is seen lying to the right of it is the fourth, or odd, arm, thus giving two arms on that side as against only one on the other. This would result in a complete loss of balance between the respective sides—a condition that is contrary to the whole tendency of the echinoderm organization—were it not that the equilibrium is preserved by the location of the stem in the plane of imperfect bilateral symmetry passing through it and the median arm, leaving the anal tube in contact with the left posterior arm as it was before; thus these two form a counterpoise to the two arms at the right, and the stem, instead of the anal tube, occupies the true middle position with regard to the calyx as a whole, as shown in figure 14a. When the odd arm atrophies in the next stage of the family, there must be a different arrangement in order to maintain a balance between the two sides, which have now become equal, with one large simple radial at each side. This is effected by the shifting of the stem into the same plane with the anal tube, thus establishing the perfect bilateral symmetry characteristic of the succeeding genera.

The r. ant. arm in both specimens is markedly smaller than the other lateral arms, as may be seen in figure 15, where one ramus of each arm adjoining the anal tube is visible on either side and their relative sizes may be compared. The median arm in this species is relatively shorter and stouter than in *C. articulatus*, and bifurcates on about the 14th brachial into fairly strong branches.

Horizon and locality. Trenton (Curdsville), Ordovician; near High Bridge, Mercer County, Kentucky.

Cremacrinus punctatus Ulrich*Plate 28, figs. 16-20**Cremacrinus punctatus* Ulrich, 14th Ann. Rep. Geol. Surv. Minnesota, 1886, p. 107, fig. 1.

This species is figured here in order to correct the error in the original description due to imperfect material, by which it appeared to have only two lateral arms, one on each side. It has the four arms characteristic of the genus, one median and three lateral, unsymmetric. The fourth arm, right anterior, which owing to injury is missing in the type (fig. 16a), is present in several specimens subsequently acquired from the same locality and horizon, as shown by figures 17, 18, and 19, but in none of them are all the side arms preserved to their full length. So far as can be observed, the fourth arm does not appear to be materially smaller than its fellows, which would lead to the inference that in this species, derived from an earlier formation than the Canadian species, the process of elimination by dwarfing of the fourth arm had not yet begun.

The median arm is preserved to the end in one specimen, where it consists of 21 brachials, the last one axillary, and becomes very slender. One ramus of the left posterior arm is also intact, showing 7 ramules, with two more added by a terminal bifurcation. The position of the stem to the right of the anal tube in the hollow of the right posterior interray is admirably shown by figure 17, with the deep indentation resulting from contact when the crown was recumbent. Another specimen, figure 20a, gives an excellent picture of the hinge, with its denticulate margin and numerous "supplementary plates." The cup in this species is slightly constricted, widening downwards, height to greatest width as 6 to 6.5; primibrachs 2; brachials in series of 2 and 3.

Horizon and locality. Black River (Decorah), Ordovician, below the horizon of *C. articulatus*; Minneapolis, Minnesota.

Additional species referable to *Cremacrinus* are *Calceocrinus furcillatus* and *C. rugosus* of W. R. Billings, and *Castocrinus billingsianus* Ringueberg, all from the Trenton of Canada.

EUCHEIROCRINUS Meek and Worthen*Plate 29*

Cheirocrinus Hall (not Eichwald), 13th Rep. New York St. Cab. Nat. Hist., 1860, p. 122.—*Eucheirocrinus* Meek and Worthen, Proc. Acad. Nat. Sci. Phil., 1869, p. 73; Geol. Surv. Illinois, 5, 1873, pp. 443, 502.—Bather, Crinoidea of Godland, 1893, pp. 56, 61, fig. 13b.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 213.—Bassler, Bibliographic Index, Bull. 95, U. S. Nat. Mus., 1915, p. 508.—*Proclivocrinus* Ringueberg, Ann. New York Acad. Sci., 4, 1889, p. 396 (sep., p. 9).

Arm-bearing rays 3, the r. ant. arm having entirely disappeared. Stem shifted back to median position in plane of anal tube, and perfect bilateral symmetry restored. Median arm simple or branching; segments of its l. ant. compound radial usually narrowly connected, inferradial widening downward. Lateral arms symmetric, one on a side; unequally dichotomous, dividing into two branches, which bifurcate once or twice again; r. post. and r. ant. super-radials fused to form subanal piece supporting anal tube, and separating the respective inferradials from one another and from anal α . Lateral primibrachs usually 2, the axillary with unequal faces. Basals 4, as in *Cremacrinus*, with suture between left BB sometimes obscure.

Genotype. *Cheirocrinus chrysalis* Hall.

Distribution. Silurian; America, England.

This Silurian form has thus lost the fourth arm, and recovered the symmetric position of the stem, but has not yet acquired the doubly heterotomous mode of branching of the two lateral rays, which marks the axil-arm system—although the unequal-faced primibrach probably forms the first step towards it. The arm structure, although in the main rather simple, is subject to considerable variation. In the typical form the lateral arm divides on the axillary second primibrach, and branches again. The median arm typically is simple, or only divides high up, while in *E. ontario* and the English species *E. anglicus* it branches near the calyx and at least twice above. In *E. chrysalis* this arm is relatively massive, and in the other two species its robustness is compensated by the lower and more numerous bifurcations. Thus the arms in this genus may be considered to be in a state of flux, leading toward the axil-arm plan. In the connection of the segments of l. ant. radial it represents a rather intermediate stage between the Ordovician *Cremacrinus* and the Devonian and Carboniferous *Halysiocrinus*, the tendency being for the inferradial to become narrower upward. The base remains substantially as in *Cremacrinus*, the unfused left BB together being small, touching the stem, and having the same characteristic curved outline, with a tendency to coalesce, which became a fixed character in the succeeding genera.

The genus as now understood has a considerable distribution, from New York, Canada, Indiana and Tennessee in America to Dudley in England. It is essentially Silurian, but it may be that *Calceocrinus barrandeii* Walcott, from the Trenton, should be referred to it.

Eucheirocrinus chrysalis (Hall)

Plate 29, figs. 1, 1a-c, 2, 3

Cheirocrinus chrysalis Hall, 13th Rep. New York St. Cab. Nat. Hist., 1860, p. 122, diagrams 1-5; 28th Rep., 1879, p. 147, fig. 2.—*Eucheirocrinus chrysalis*, Meek and Worthen, Proc. Acad. Nat. Sic. Phil., 1869, p. 73; Geol. Surv. Illinois, 5, 1873, p. 443, 502.—*Proclivocrinus chrysalis* Ringueberg, Ann. New York Acad. Sci., 4, 1880, p. 399 (12), pl. 10, figs. 2, 13.—*Proclivocrinus radculus* Ringueberg, *ibid.*, p. 397 (10), pl. 7, fig. 6.

The genotype. To facilitate comparison of the generic characters I am figuring the type specimen from the Rochester shale of the Silurian at Lockport, New York, together with some others from the same locality. In the type the suture dividing the left BB is indistinct, but can be seen in strong sunlight. In these specimens the outer articulating face of the axillary primibrach exceeds the inner in width about as 4 to 3. I also give herewith some diagrams to illustrate the variations in arm structure of this and the two species showing the greatest departure from type, *E. ontario* and *E. anglicus*.

FIG. 3. Varieties of Arm-Branching in *Eucheirocrinus*. a, *E. chrysalis*; b, *E. ontario*; c, *E. anglicus*

Eucheirocrinus minor new species

Plate 29, figs. 5, 5a, b

A small species. Crown 39 mm. high, of which the calyx occupies 6.5 mm. Has 3 arms, and stem in plane of anal series. Median arm fairly stout, tapering, as long as the lateral arms, consisting chiefly of two long brachials, four or five times as long as wide, preceded by the short primibrach and followed by a third brachial shorter and narrowing to a point; segments of 1. ant. radial connected, with inferradial elongate quadrilateral. Lateral arms 2, one on each side, dividing on second primibrach into 2 rami, dichotomous, branching somewhat unequally at intervals of 3 or 4 brachials to about 6 finials, the ramus next to the anal tube being the smallest. Stem long, slender, about three times the length of the crown, tapering gradually to a narrow distal end; columnals in proximal part about 1 mm. long and wide, becoming longer with half the width distally.

In form and proportions of the median arm this species differs strongly from the type, and resembles those of the associated species of *Cremacrinus* occurring in the same formation; but the absence of a fourth arm and the median position of the stem clearly exclude it from

that genus, while the structure of the lateral arms is substantially that of *Eucheirocrinus*, with a difference, however, in the proportionate size of the branches. The specimen figured is unique, and only the left posterior side is fully exposed; but I was able to free the opposite side sufficiently to show that it has but one lateral arm. It is an intermediate form of uncertain position.

Horizon and locality. *Eucalyptocrinus* zone of Beech River formation, Brownsport group, Niagaran; Tuck's Mill, Decatur County, Tennessee.

***Eucheirocrinus ontario* Springer**

Text-fig. 3b

Eucheirocrinus ontario Springer, Geol. Surv. Canada, Mem. 3, No. 91, 1919, p. 127, fig. 6.

This species, described by me some years ago, is of somewhat earlier age than the type; it is inserted for comparison of arm structure, especially of the median arm, which bifurcates on the third brachial, and the branches at least once more at like intervals beyond. This arrangement is shown by text-figure 3*b* under *E. chrysalis*. From the Cataract formation, Medinan; Stony Creek, Ontario, Canada.

***Eucheirocrinus anglicus* new species**

Plate 29, figs. 6, 6a; text-fig. 3c

This small species from the English Silurian is described here on account of the extraordinary variation from the type which it exhibits in the branching of the median arm. This bifurcates on the first and only primibrach, and then once or twice more to the number of at least 6 ultimate branches; the lateral arms are irregular, both their rami and their branches being of an intermediate character (text-fig. 3*c*). The branching of the median arm on the primibrach is paralleled only by the species *Calceocrinus pinnulatus* Bather (Crin. Gotl., pl. 4, fig. 133) of Gotland; but whereas in that species the lateral arms have the usual two primibrachs, here the single axillary primibrach extends to the lateral arms also, thus duplicating the otherwise unique character of another Gotland species, *C. nitidus*, which occurs also in England in the same formation and locality as the present species. In the profuse branching of the median arm this species recalls a similar departure from type in *Calceocrinus bifurcatus*.

The species is based upon a single specimen, unique so far as I know, from Dudley, England, which I obtained many years ago from the well known dealer, Robert Damon of Weymouth. It is notable for the rather ponderous character of the arm branches, the great predominance of the median arm, and the relatively small size of the calyx. It has a short stem, apparently about half the length of the crown, which seemed to end amid the debris of some other fossils so that it could not be determined whether the stem was attached to a foreign object or not. The columnals increase in length rapidly toward the distal end, where they become twice as long as wide. In this respect the form has a rather juvenile aspect. The crown, which is considerably flattened by pressure, has a height of 20 mm.

Horizon and locality. Wenlockian, Silurian; Dudley, England.

Eucheirocrinus indianensis (S. A. Miller)

Plate 29, figs. 4, 4a, b

Calceocrinus indianensis S. A. Miller, 17th Ann. Rep. Indiana Dep. Geol., 1892, p. 645, pl. 6, fig. 37 (adv. sheets, p. 35).

Founded upon a very imperfect specimen with only part of the calyx preserved, but enough to show the narrow and tapering inferradial of the left anterior ray, and the 4 basals, indicating the probable presence of the genus among the many rare forms of its locality. No others have been found.

Horizon and locality. Laurel formation; St. Paul, Indiana.

CALCEOCRINUS Hall em. Ringueberg*Plate 29*

Calceocrinus Hall, Pal. New York, 2, 1852, p. 352, pl. 85, figs. 5, 6; 28th Rep. New York St. Cab. Nat. Hist., 1879, p. 146, figs. 1, 2; 12th Ann. Rep. Indiana Dep. Geol., 1882, p. 281.—Shumard, Trans. Acad. Sci. St. Louis, 2, 1866, p. 358.—Meek and Worthen, Geol. Surv. Illinois, 5, 1873, pp. 443, 502, pl. 14, fig. 9.—Ulrich, 14th Ann. Rep. Geol. Surv. Minnesota, 1886, pp. 104, 113.—Wachsmuth and Springer, Rev. Pal., 3, sec. 2, 1886, p. 274.—Ringueberg, Ann. New York Acad. Sci., 4, 1889, p. 401.—Bather, Crinoidea of Gotland, 1893, p. 67; Treatise on Zool., 3, 1900, p. 147.—Jaekel, Phylogenie und System, 1918, p. 86.—*Deltacrinus*, Bassler, Bibliogr. Index, Bull. 92, U. S. Nat. Mus., 1915, pp. 156, 393.

Arm-bearing rays 3. Calyx bilaterally symmetric, with stem in plane of anal tube. Median arm simple or branching, supported by compound l. ant. radial, the segments of which are usually separated. Lateral arms one on each side, supported by the large radials, branching from unequal faces of the axillary primibrach and of successive diminishing axillaries curving around transversely toward the anal side, each axillary supporting at the shorter, inner face an axil-arm composed of successive series of 2 or 3 brachials, and bearing alternate ramules, and at the longer, outer face the next axillary; primibrachs of lateral arms usually 2. R. post. and r. ant. superradials fused to form a subanal piece underlying anal α , usually separated from the large radials by their respective inferradials, or they may be eliminated. Basals reduced to 3 by fusion of l. post. and l. ant. BB, the fused plate never or rarely entering the stem-facet. The foregoing characters apply also to the genus *Halysiocrinus*.

R. post. and r. ant. inferradials usually not in contact, interposed between anal α or subanal piece and the large radials. The two segments of l. ant. compound radial exceptionally connected by short suture, but usually entirely separated as triangles by the large radials meeting between them. Fused left basal usually triangular with straight sides, and narrower than the hinge in the American species. Median arm usually simple.

Genotype. *Calceocrinus typus* Ringueberg.

Distribution. Silurian; America, England, Gotland.

The characters stated in the first paragraph are common to this and the next following genus. The axil-arm system characteristic of the two is a feature unknown in other crinoids; in this by atrophy and fusion the two non-arm-bearing radials have become to a large extent eliminated, leaving the suprabasal portion of the calyx to consist, in addition to plates of the anal area, almost entirely of the two large simple radials, and the more or less interposed l. ant. compound radial, upon the former of which the one-sided, thoroughly unique type of arm-structure has been established. This begins in the Silurian with numerous species in the Wenlockian of Sweden and England, and in the Niagaran of New York, Indiana, Missouri and Tennessee, extending through the Devonian into the Lower Carboniferous, where it culminates with the end of the family in the Keokuk and Warsaw beds of the Mississippi Valley. The extreme stage, in which the brachial series of the main arm are subordinated to the over developed outer ramules and thus hidden from view at the exterior, is confined to certain species of *Calceocrinus*, viz.: *C. foerstei*, *C. bifurcatus* and *C. typus* of the American rocks, and *C. tenax*, *C. nitidus* and *C. interpretis* of the European; but never occurs, so far as known, in *Halysiocrinus*, in which some portion of the main arm is always more or less ex-

posed. Thus while a species with the main arms partly exposed might belong to either genus, we can be fairly certain that one in which these parts are hidden by the outer ramules does not belong to *Halysicrinus*.

Coincident with the development of the axil-arm system and the enlargement of the simple radials came the separation of the two segments of the 1. ant. compound radial. This, as before shown, was a progressive feature, appearing first in the Silurian among species of this genus *pari passu* with the axil-arms, and becoming established as an invariable structure in the later stages of the family. It is a thoroughly reliable character for distinguishing the two genera here defined from those preceding them, as is also the fusion of the two left basals into one triangular or convex plate not connected with the stem, this being especially valuable because determinable from the base alone. The minor differences by which the two are separated are not conspicuous, but the division, confirmed as it is by stratigraphic considerations, is useful, and the characters if carefully observed are usually sufficient for identifying them. In some of the Gotland species the fused left basal is almost, and rarely quite, as wide as the hinge.

Calceocrinus foerstei new species

Plate 29, figs. 7a, b, 8a, b, 9, 10, 11

Crown of medium size, ranging from 37 to 50 mm. in height, and fairly stout. Cup as seen from 1. ant. side relatively short, obtusely trapezoidal, but little constricted; height to the width at hinge as 1 to 1.4. Median (1. ant.) arm very stout, simple, and longer than the axil-arms; brachials averaging 3.4 mm. high by 4 mm. wide in a large specimen; segments of 1. ant. R connected, infer-radial elongate quadrilateral. Lateral IBr 2. Main-axils of lateral arms 5 to 8. Axil-arms stout, with brachials of exposed ramules as wide as high, about 2.5 mm. in large specimen; outer (adanal) ramules following Betabrachs closely apposed, decreasing regularly in size stemwards, and forming the chief visible arm-branches; when closed they hide from view the main arms, as well as the inner (abanal) ramules. Anal tube stout, composed of large segments resembling brachials, projecting beyond the axil-arms; anal x large, strongly rounded, may be slightly indented by pressure of column; subanal piece fairly large, wider than high, usually more or less indented. Stem stout, apparently rather short, the longest one preserved beginning to taper at 5.5 cm.; columnals about 1.5 mm. high and 3 mm. wide in middle portion; with but little curvature, the stem lying closely parallel to the anal series.

This species is described upon the evidence of eight crowns, mostly well preserved, and one having the short stem probably to near the distal end; three of them are figured. In the exceptional presence of a quadrangular inferradial and the resulting connection of the segments of 1. ant. R., it shares with the Silurian species of *Cremacrinus* in the survival of this primitive Ordovician structure, and in this character differs from all other species of this genus except *C. pinnulatus* and *interpres* of Gotland. In the approximation of the outer ramules of the axil-arms it is of the type of the Gotlandian *C. tenax*. One specimen, figure 10, somewhat distorted, differs from the others in the proportions of the brachials and axil-arms, but upon the more essential characters appears to belong with the species. It was figured under the impression that it might have to be separated. Figure 11 is from a set of arms of unusual size which exhibits especially well the characteristic disposition of the axil-arm in the extreme stage.

The specific name is given in recognition of the extensive and valuable researches of Dr. August F. Foerste in the Silurian and Devonian formations of Tennessee.

Horizon and locality. *Eucalyptocrinus* zone of Beech River formation, Brownsport group, Niagaran; Tuck's Mill, Decatur County, Tennessee.

***Calceocrinus bassleri* new species**

Plate 29, figs. 12a, b, 13, 14; pl. 28, fig. 21

Crown of large size, from 50 to 56 mm. in height. Cup seen from left anterior side obtusely subquadrangular, relatively short and wide; height to width at hinge line in maximum specimen 1 to 1.5; but little constricted. Median (l. ant.) arm simple, stout, much thicker and longer than the axil-arms; brachials about 3 mm. high; segments of l. ant. R triangular, widely separated by the large RR abutting between them, exceptionally touching by their apices. Lateral IBr 2. Main-axils of lateral arms 5 to 8, brachials in series of 2 to 4; inner Betabrachs (abanal) and following series more or less exposed alongside the outer ramules, their brachials moniliform. Anal tube projecting beyond arms, not so stout as in preceding species; anal α large, strongly rounded, but little indented by column; subanal piece rather short, usually widening upward. Stem long, stout, but little curved next to calyx; columnals very short, at 25 mm. from base averaging .7 mm. long and 3 mm. diameter.

This species is founded upon six crowns, four of which are figured, one with almost the complete stem attached. It differs from the preceding species in having the segments of l. ant. R disconnected as usual in the genus, in the exposure of the main branch of the axil-arms, and in the short columnals of the stem. It is to be compared with *C. pugil* and *C. tucanus* of Gotland, with which it is evidently closely related, the chief differences being its larger size and greater number of axil-arms. In the relative exposure of the main axil-arm it is closer to those species than the published figures would indicate, as is shown by a specimen from Gotland in my collection herein figured, plate 29, figure 21.

This fine species is named in honor of Dr. R. S. Bassler, of the United States National Museum, whose field-work during several seasons has greatly enlarged our knowledge of the geological structure of Tennessee.

Horizon and locality. *Eucalyptocrinus* zone of Beech River formation, Brownsport group, Niagaran; Tuck's Mill, Decatur County, Tennessee.

***Calceocrinus bifurcatus* new species**

Plate 29, figs. 15, 15a, b

Crown of medium size, about 45 mm. high, of slender form as compared with preceding species. Cup widening basalward, height 7 mm., width 9 mm., and but little constricted. Median (l. ant.) arm slender and very deep, bifurcating on the third brachial above the primibrach, and again on the second, the inner branch dividing again, thus giving 6 or 8 finials; all the brachials are long and narrow, the lower three being 4.5 mm. long by 1.5 wide; segments of l. ant. R widely separated by large RR, the superradial triangular and the inferradial

narrowly acuminate. Lateral IBr 2. Main-axils of lateral arms 6. Outer ramules of axil-arms closely apposed, concealing the main branches and inner ramules. Anal plates strongly convex, partly obscured by the closely adhering stem. Tube not visible. Stem fairly strong, with columnals 1.5 mm. long at about midway of the crown.

Only the single type specimen is known, and that is from a slightly lower horizon than the preceding species. The crown is in the recumbent position, closely covering the stem which is almost enveloped by the smaller axil-arms. The striking character by which the species differs from almost all others of the genus is the repeated bifurcation of the median arm, in which it anticipates the later Carboniferous forms.

Horizon and locality. *Coccoecrinus* zone of the Beech River formation, Brownsport group, Niagaran; West of Tuck's Mill, Decatur County, Tennessee.

Calceocrinus stigmatus Hall

Plate 29, fig. 16

Cheirocrinus stigmatus Hall, Trans. Alb. Inst., 4, 1863, p. 225; *Calceocrinus stigmatus* Hall, 28th Rep. New York St. Mus., 1879, p. 147, pl. 19, figs. 9-11; 11th Ann. Rep. Indiana Dep. Geol., 1882, p. 281, pl. 19, figs. 9-11.

Copy of Hall's figure of the type, to show the occurrence of the form in the Waldron formation of the Silurian. Only the radial part of the calyx is preserved, showing a wide separation of the segments of l. ant. compound radial. This will distinguish it from *Eucheirocrinus*, but not from *Halysiocrinus*, and the other characters are not in sufficient detail for comparison. The geological position, however, raises a fair presumption against the latter. The species is extremely rare, no other specimens having been found.

Waldron shale, Niagaran; Waldron, Indiana.

Calceocrinus typus Ringueberg

Plate 29, figs. 17, 17a

Calceocrinus typus Ringueberg, Ann. New York Acad. Sci., 4, 1889, p. 402, pl. 10, fig. 8.

Ringueberg's type specimen, now in my collection, is figured to illustrate the typical form as recognized among the New York specimens. It has the axil-arms of the type of *C. foerstei* and *C. tenax*, and the disconnected segments of l. ant. R.

From the Rochester shale of the Niagaran; Lockport, New York, where it is one of the rarest fossils.

In addition to the two foregoing species Ringueberg described *Calceocrinus bidentatus* and *C. contractus*, both from the Niagaran at Lockport, New York.

Calceocrinus Hall

Plate 29, fig. 18

Hall, Pal. New York, 2, 1852, p. 352, pl. 85, fig. 5.

The original figure upon which the genus was proposed. After Hall. It is figured here for comparison with *C. halli* and *C. typus*, identified as congeneric with this.

Horizon and locality. Rochester shale; Lockport, New York.

Calceocrinus halli Ringueberg*Plate 29, fig. 19*

Ringueberg, Ann. New York Acad. Sci., 4, 1889, p. 403, pl. 10, fig. 9.

Type of the species which Ringueberg considered identical with Hall's original. The consolidated left basal, fused into a single triangular plate not touching the stem, is the dominant character in both, fully establishing their generic identity, and by reason of its exceptional width their specific identity also.

Horizon and locality, same as last.

Calceocrinus alleni Rowley

Amer. Geol., 34, 1904, p. 275, pl. 16, figs. 30-33.

Described from fragments of calyx only, which show only the probable generic characters. Bainbridge formation. Niagaran; Ste. Genevieve County, Missouri.

Calceocrinus nitidus Bather*Plate 29, fig. 20*

Calceocrinus nitidus Bather, Crin. Gotl., 1893, p. 91, pl. 3, figs. 117-122.—*Synchiocrinus anglicus* Jaekel, Phylogenie und System, 1918, p. 86, figs. 81-83.

I am illustrating this species from an unusually fine specimen in my collection from England, in which the crown is more completely preserved than in the types. In the principal one of these the median arm has only 8 brachials remaining, whereas here they reach a total of 15. The arm is decidedly stouter and longer than those adjoining it, and has a graceful curvature parallel to the 6 visible axil-arms, of which only the outer or adanal ramules are seen, the main branches being entirely concealed underneath them. The species is remarkable for having only a single lateral primibrach, in which it differs from all others in the family known to me, with the exception of the singular English species herein described as *Eucheirocrinus anglicus*.

Although described from Gotland, *C. nitidus* is apparently the most common species of this genus in the English Silurian, as among my own material are no less than four good specimens, besides two casts from specimens in British collections, in all of which the characters are remarkably constant. One of these has the stem in place, but incomplete, for a distance of once and a half the length of the crown, which is closely recumbent upon it; the columnals in the more distal portion have increased to 2 mm. in length for 3 mm. diameter.

This species is undoubtedly the same form for which Jaekel in his *Phylogenie und System*, 1918, has encumbered the literature with two useless synonyms as *Synchiocrinus anglicus*.

Horizon and locality. Wenlockian, Silurian; Gotland, Sweden, and Dudley, England.

Calceocrinus pugil Bather*Plate 29, fig. 21**Calceocrinus pugil* Bather, Crinoidea of Gotland, 1893, p. 82, pl. 3, figs. 98-104.

Figured from a specimen in my collection from Gotland for comparison with *C. bassleri* from Tennessee, which resembles it in the relative visibility of the main branches of the axil-arms. The exposure in that respect is somewhat greater in this specimen than in those figured by Dr. Bather, and the ossicles in the main branches are more distinctly nodose. In decided contrast to the preceding species, the median arm in this is slender and short, shorter than the adjoining axil-arms; in the principal type specimen this arm has 8 brachials visible, with room for only 1 more, and here the number is apparently limited to 8.

Horizon and locality. Wenlockian, Silurian; Gotland, Sweden.

HALYSIOCRINUS Ulrich*Plates 29, 30*

Cheirocrinus Hall (not Eichwald), 13th Rep. New York St. Cab., 1860, p. 123.—*Halysiocrinus* Ulrich, 14th Ann. Rep. Geol. Surv. Minnesota, 1886, p. 110.—Bather, Crin. Gotl., 1893, p. 61, fig. 13b.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 214.

Arms and stem as in *Calceocrinus*, except that the axil-arms do not reach the extreme stage, and the main branch is always more or less exposed. The two segments of l. ant. R triangular, and always completely separated by the simple radials. Subanal piece atrophied or hidden, its place being taken by the posterior shifting of r. post. and r. ant. inferradials, which have met, supporting anal x directly. Fused left basal usually convex toward the stem, and nearly or quite the full width of hinge. Median arm frequently branching.

Genotype. *Cheirocrinus dactylus* Hall.

Distribution. Devonian to Carboniferous; America.

As stated under *Calceocrinus*, the first paragraph of the diagnosis of that genus applies to this, in which the axil-arm system is equally characteristic, but developed in a somewhat different way, the exposure of the main branch being here a constant character, as is also the separation of the segments of l. ant. R. There may also be noted an increasing tendency to branching of the median arm. The difference in the structure of the posterior side, marking for this genus the final stage in the elimination of the radials of the vanished arms, in which their inferradial remnants are pushed to a position directly underneath the anal tube, is a reliable criterion; and the greater width and convex form of the fused basal is a useful guide in practice, being fairly established in the Devonian and apparently constant in the Carboniferous.

The genus is essentially an American form, being represented by species in the Helderbergian, Onondaga and Hamilton of the Devonian, and the New Providence, Burlington, Keokuk and Warsaw of the Lower Carboniferous. It has not thus far been recognized in Europe.

Halysiocrinus keyserensis new species*Plate 29, fig. 22*

This earliest known representative of its genus, from the Lower Devonian, while it has the predominant outer ramules and the dwarfed main branches somewhat exposed, as usual in the genus, differs from the axil-arm type generally by having the axillary primibrach and the succeeding main-axils about equal-faced. The median arm is long and simple, with brachials moniliform, and the axil-arms four in number. The anal plates at base of tube are large and broadly rounded, but the tube itself is not exposed. In the nearly equal bifurcation upon the main-axils there is some resemblance to *Deltacrinus clarus* and *Halysiocrinus secundus* of the Middle Devonian, as figured and redescribed in the recent volume on the Devonian Crinoids of New York, plate 41.

Based upon a unique specimen from the Keyser formation of the Helderbergian, at Keyser, West Virginia.

Halysiocrinus marylandensis (Ohern)

Plate 29, figs. 23, 23a, b

Calceocrinus marylandensis Ohern, Maryland Geological Survey, Lower Devonian, 1913, p. 253, pl. 40, figs. 1-3.

This species, described for the Maryland Geological Survey, is figured here to show the occurrence of the genus in the Oriskany division of the Lower Devonian. It is remarkable for its robust proportions in contrast to the other Devonian forms, but the generic characters are thoroughly well established. There are at least six axil-arms springing from extremely unequal-faced main-axils, and the main arm-branches are fairly well exposed alongside the large outer ramules. The median arm is simple to the eleventh brachial, and of unusual size, as are also the plates of the anal series. The whole aspect of the crinoid is one of massiveness not before observed in the family.

Only the type specimen is known, from the Oriskany sandstone at Cumberland, Maryland.

Halysiocrinus carinatus new species

Plate 29, figs. 24, 24a, 25, 26

This Hamilton species is introduced in order further to show the course of development of the family during the Devonian, and especially as a fine example of the indentation of the large anal plates by pressure on the stem. It is remarkable for the extraordinary number of axil-arms, 10 on each side, the narrowness of the 1. ant. superradial, and slenderness of the median arm following it, of which only a few brachials are preserved; this is very small and keeled, in contrast to the broad, rounded arm of most other species. The space gained by the reduction in width of this arm affords room for the increased number of main-axils, as a result of which the lateral arm-facets are directed more nearly upward than in other species; this character is constant in specimens from the three localities. The axil-arms are also slender, with elongate brachials in two or more well exposed branches. The anal plates are very large, and the indentation made by the stem is sharp and deep, and could only have been produced by strong and persistent pressure, such as would result from a habitual recumbent position of the crown upon the stem. No mere temporary suspension of the crown alongside the stem would account for it. The anal tube is sharply angular, tapering to a point below the distal end of the arms; this form of tube is a striking character of the species which has not been seen in any other, and reminds one of the pointed tube in some species of *Zeacrinus*, leaving the position of the anal opening in doubt. Surface of radials and basals corrugated or pustulose.

The species is founded on two crowns from the Traverse formation of northern Michigan; the type here figured, 38 mm. in height, is sharply and accurately preserved, except for

the distal part of the median arm. Two calices are also figured of the same or a closely allied species from equivalent horizon in other areas.

Horizon and locality. Hamilton group, Middle Devonian; Alpena, Michigan, New Buffalo, Iowa, and Clark County, Indiana.

Halysiocrinus perplexus (Shumard)

Plate 30, figs. 4-13

Cheirocrinus (*Calceocrinus*?) *perplexus* Shumard, Trans. St. Louis Acad. Sci., 1866, 2, p. 358.

Some specimens of this species are figured in connection with the explanation of the hinge already given, for which it offers unusual facilities. It occurs in the New Providence shale of the basal Lower Carboniferous,¹ and is found chiefly on weathered exposures at Button Mould and other knobs south of Louisville in Kentucky. Complete crowns are unknown, but the two characteristic elements of the united radial plates and separate consolidated basals have been collected by hundreds, among which are many that furnish a clear exposition of the structure of these parts. They vary in size from cups 10 mm. high by 6 mm. wide at the top and 8 mm. at the hinge line, to those 20 mm. high, which increase in width from 14 mm. at the top to 33 mm. at the hinge—thus showing a tendency to become wider basalward with age. Some differences may be noted among the specimens which might indicate the presence of more than one species, such as the greater or less projection of the superradial, and variation in surface from pustulose to smooth; but the latter is largely a matter of erosion, and in default of more complete material I have referred them all to the original species.

Horizon and locality. New Providence shale, Mississippian; Button Mould Knob, and other localities widely distributed in Kentucky and Tennessee, and Clark County, Indiana.

Halysiocrinus dactylus (Hall)

Plate 30, figs. 1, 2, 2a, b, 3, 3a

Cheirocrinus dactylus Hall, 13th Rep. New York St. Cab. Nat. Hist., 1860, p. 123, figs. 1, 2.—*Halysiocrinus dactylus* Ulrich, 14th Ann. Rep. Geol. Surv. Minnesota, 1886, p. 110.—*Cheirocrinus ventricosus* Hall, 13th Rep., supra, p. 123.—*Halysiocrinus ventricosus* Bather, Crin. Gotl., pp. 65, 66, pl. 4, figs. 141, 142.

I am illustrating this Lower Carboniferous species because it has been designated as the type of the genus, and for the further reason that doubt has been expressed as to its standing, some authors thinking it may be a synonym of Hall's *C. ventricosus* from the same formation—both having been described in the same publication. I am therefore figuring the type specimen in the Museum of Comparative Zoology, together with some others from my own collection showing further details. The type, also well figured by Hall, gives a complete picture of the generic characters, whereas *C. ventricosus* was de-

¹ Knobstone of authors, and by some erroneously referred to the Keokuk. See Springer, The Crinoid Fauna of the Knobstone formation. Proc. U. S. Nat. Mus., Vol. 41, 1911, p. 178.

scribed without figure from an isolated calyx only, without any distinctive characters. As there is nothing in the description to establish priority, it is deemed proper to hold it for the species which is recognizable.

This is the dominant species of the genus in the Burlington limestone, where it is found in excellent preservation. It is represented in the collection by upwards of 20 specimens, which exhibit some of the variations to be expected in a prolific species. Normally it has 4 axil-arms, but there may be 3 or 5; the median arm branches on the 6th to the 8th or 9th brachial, usually with a single bifurcation, but sometimes with two; the ramules of the lateral arms usually occur at intervals of 4 brachials, but occasionally 3 or 5, mostly in two series but may be three, always exposed. The form of the cup changes with maturity of the individual, that of the younger ones being higher and narrower than in the older; as a rule the cup is more elongate than in other species, the average of four specimens being 9 mm. height by 8.5 mm. width at the hinge. A striking example of the difference in this and other respects due to age is furnished by an extremely young individual associated with this species and presumably belonging to it, which I have illustrated on plate 30. The complete crown is only 12 mm. high, and the stem, also about complete, 22 mm. long; the cup is 4.5 mm. high, 2.5 mm. wide across the l. ant. side at the top, diminishing to 1.7 mm. at the hinge; on the l. post. side, as originally exposed, there is but one arm, unbranched, and on the anterior side, as seen by removing the crown from the matrix, there are two branches, besides a curving anal tube, also visible from the first position. Enlarged figures were made from both these views (pl. 30, figs. 3, 3a).

Horizon and locality. Upper Burlington limestone, Mississippian; Burlington, Iowa.

Halysiocrinus nodosus (Hall)

Plate 30, figs. 14-21

Cheirocrinus nodosus Hall, 13th Rep. New York St. Cab. Nat. Hist., 1860, p. 124; Phot. Plates of Crinoids, 1872, pl. 6, fig. 15.—*Halysiocrinus nodosus* Ulrich, 14th Ann. Rep. Geol. Surv. Minnesota, 1886, p. 112.—*Deltacrinus nodosus*, Spr., Smithsonian Misc. Coll., 76, 1923, pl. 5, fig. 21.

This species may be taken to represent the culmination of the family in the Lower Carboniferous. At least one species from a later horizon is known, which is rare and shows a reversionary tendency. The present form is selected for illustration to complete the picture of the type we have been discussing, because it is the most striking species of all the Carboniferous in its characters, is found in remarkably fine preservation, and in such abundance as to offer excellent facilities for the study of variations within the limits of a well defined species. The large size, robust habitus, sharp nodosity, curvature and frequent branching of the median arm, and free exposure of main axil-arms, are conspicuous and decisive characters.

I am illustrating it on plate 30 rather fully, first, because the recumbent habitus, with the crown pressing closely down over the stem, is so admirably shown, for example by figure 19 in which the arms seem actually to envelope the stem in a presumably horizontal position. In figure 17, the suggestion that the crown might close down like a barnacle is brought forcibly to mind; it is here strongly curved in the plane of its bilateral symmetry, with the large median arm at the perimeter; it is a peculiar posture, which might be described by the word "humped," and the resulting general form, when seen from the side, is semi-elliptical. This is the typical form for the species, and while there are some specimens not otherwise distinguishable in which the "hump" is not so pronounced, it is a fact that out of about forty well preserved

crowns in the material now in hand, considerably more than half of them have the decided appearance of a tightly strung bow, and practically all of these have the stem lying close underneath on the side opposite the curve. No crinoid with such a curvature of the crown could conceivably occupy a habitual position either erect, or hanging down along an erect stem; whereas the form here exhibited lends much plausibility to the idea of a recumbent habitus. A single specimen, figure 15, already alluded to in the discussion, shows the crown almost fully opened upon the hinged articulation with the base, but this is believed to be incidental, due to some spasmodic movement at death.

It also illustrates the great and final extension of the two large radials toward the tube, they being now separated only by the inferradials of the vanished arms, which are here reduced to a relatively small size, and with which, indeed, they may be occasionally fused. Along with this, the visible anal plates are much reduced, being but little exposed at the base of the tube, in strong contrast to their condition as usually seen in the Ordovician, Silurian and Devonian species; nevertheless the tube extends to the full height of the arms (fig. 16a).

Secondly, the species exemplifies in a striking manner the variations that must be expected and allowed for in a prolific form, especially one that is marked by such strong characters that it is readily recognized, even in many cases from fragments. As above stated, there are about forty good specimens in the collection with the crown intact, all from a single restricted colony in the Keokuk group. They range in size from 70 mm. height of crown down to 20 mm., so that we have the opportunity to compare the condition of young and adult. Upon tabulating the characters as exhibited by this series of specimens, it is found that as to several of them there is a definite progressive variation according to the maturity of the individual:

The proportions of the cup. Typically as seen from the left anterior radius this becomes wider toward the hinge-line, where its width exceeds the height from there to the top of the superradial. A composite of five mature specimens gives 15 mm. height by 16 mm. width, with a maximum of 17 mm. by 20; whereas in the young of about 20 mm. height of crown these proportions are reversed to 8 mm. height by 5 mm. width (pl. 30, fig. 20).

The bifurcation of the median arm. Here this reaches its culmination for the family to a maximum of 8 and 10 branches. It bifurcates throughout the species on from the fifth to the tenth brachial—in 70 per cent on the seventh or eighth, which is the normal—but the number of branches differs with growth. Specimens of from 35 to 70 mm. height of crown have from 6 to 8 (rarely 10) finials, while those of 20 to 35 mm. have from 2 to 4; in the youngest individuals only about 20 mm. high, where the median arm narrows almost to a point, there is evidence of a single bifurcation upon the sixth or seventh brachial.

In like manner the number of axil-arms increases with age, specimens above about 40 mm. in height having 5 and 6 on a side, the former number predominating; those from 30 to 40 mm. have usually 4, and below 30 mm. 3. The juvenile condition in this and the preceding species, with their small number of lateral arm-branches, recalls the arm-structure of *Eucheirocrinus*, from which the axil-arm system may have been derived. In the present species the Alphabrachs and Betabrachs range in their series from 3 to 5, the prevailing number being 4.

From this résumé of the facts resulting from a careful tabulation, it is evident that most of these variations are progressive with individual growth, and that, in this form at least, such differences would not form any sure basis for the separation of species.

The progressive variations thus exhibited due to growth in the individual are analogous to those which occurred in the same characters during the life of the family from the earlier to the later geological stages.

Horizon and locality. Keokuk group, Mississippian; Indian Creek, Montgomery County, Indiana; and rare in the vicinity of Keokuk, Iowa.

Family PETALOCRINIDAE Weller and Davidson

Fistulata in which IBB are minute and probably fused, and in which the branches of each arm, from IAx to the finials inclusive, are fused into rigid arm-fans, articulating with the cup by means of a free IBr. No anal.

Petalocrinus Weller and Davidson*Plate 26*

Petalocrinus Weller and Davidson, Jour. Geol., 4, 1896, pp. 166-170.—Bather, Quart. Jour. Geol. Soc. London, 54, 1898, pp. 401-441, pls. 25, 26; Treatise on Zool., pt. 3, 1900, p. 175, fig. xci.—Thomas, Proc. Iowa Acad. Sci., 22, 1916, p. 289.

A dicyclic Inadunate crinoid, with rays multibrachiate, fused into rigid, fan-shaped appendages, smooth and amorphous at the dorsal side, and grooved for the ambulacra at the ventral, articulated to the radials through a single short primibrach; with pentamerous symmetry; IBB minute, hidden by column; 5 BB similar in form; RR 5, equal, with no anals. Stem subcircular.

Genotype. *Petalocrinus mirabilis* Weller and Davidson.

Distribution. Silurian; America and Gotland.

In the fusion of its many-branched arms into solid appendages this genus differs from all other crinoids. Described by the discoverers, Weller and Davidson, in 1896 from the Silurian of Iowa, it later formed the subject of an elaborate memoir by Bather in 1898, based upon the type and other material from America loaned to him for study, together with specimens from the corresponding horizon in Gotland. From the latter he described three new species and from the former two, making a total, including the type, of six species. To his complete and exhaustive discussion of the morphology and systematic relations of this unique form I would refer the student for all needful information.

The genus is introduced here because my collections from St. Paul, Indiana, have yielded several excellent specimens confirmatory of one of Bather's species. For the better understanding of this, I am giving some instructive illustrations of the type species, *P. mirabilis*, especially one of an unusually fine specimen from the original locality in the rare condition of having the calyx and appendages, which are called arm-fans, in place. For the use of this I am indebted to the thoughtfulness of Prof. A. O. Thomas, of the University of Iowa.

Of the six described species, all but the type are represented only by the isolated arm-fans; and of the type, while these fragments are numerous, only three or four specimens have been found with the fans in position and the calyx preserved. From this lack of other elements for comparison, specific characters must be sought in the relative proportions of the arm-fans, their shape as determined by the angle formed by their sides with one another, and the number and distribution of the ventral grooves. The fans are somewhat paddle-shaped, and the facet for articulation at the narrow proximal end is almost the width of the radial.

Petalocrinus mirabilis Weller and Davidson*Plate 26, figs. 12-14*

Petalocrinus mirabilis Weller and Davidson, Jour. Geol., 4, 1896, p. 166.—Bather, Quart. Jour. Geol. Soc. London, 54, 1898, p. 427, pl. 26, figs. 37-56.

Dorsal cup bowl-shaped, flattened dorsally, height about half its width which varies from 3.25 to 6 mm.; diameter of crown with fans outstretched

about 20 mm. Arm-fans bilaterally symmetrical in number and distribution of grooves; average of 30 isolated fans gives length 10.6, width 13 mm., with range of variation from 7.75 to 14.5 mm. length and 7 to 16 mm. width. Ventral grooves, finial, mostly 16, rarely 14. Angle of arm-fans 51° to 83° , normal variation 71° to 78° ; ridges usually narrower than the grooves.

The above figures are taken from Bather's tabulation of a large number of specimens. They are confirmed by a series of about 80 fans in my own hands which indicate a range of 18 to 30 mm. in diameter of crown, with slight difference of average in some other respects. Combining the data there is thus established a limit for the species not exceeding 14 mm. length and 16 mm. width of fan, and not less than 50° nor more than 83° angle of the sides. A fair representation of its form and proportions is given by the figures on plate 26. Figure 13 shows the smooth, amorphous dorsal surface of a set of isolated fans, artificially arranged, and figure 12 the sharply grooved ventral furrows of another set. The ten pieces have a range of 10-13 mm. length and 10-14 mm. width. It is therefore a relatively small species. In figure 14 we have a double size view of the complete calyx with all the arm-fans in position, but the dorsal side is stripped off, leaving only the inner surface of the ventral grooves visible.

There is a very close resemblance between this species and the Gotland form, *P. visbyensis* of Bather, which is of similar small size and number of grooves, but has a greater range of variation.

Horizon and locality. Hopkinton stage of Niagaran; near Monticello, Jones County, Iowa. The original of figure 14 was collected by the late Professor Samuel Calvin, in 1899.

***Petalocrinus inferior* Bather**

Plate 26, figs. 15-16

Petalocrinus inferior Bather, Quart. Jour. Geol. Soc. London, 54, 1898, pl. 26, fig. 57, text-fig. 10.

From the American material loaned him Bather described two new species, *P. inferior* from the type locality in Jones County, Iowa, and *P. longus* from St. Paul, Indiana, each based upon a single arm-fan in the Walker Museum, University of Chicago, from approximately equivalent Niagaran horizons, the former perhaps a little the higher.

Among my collections from the Laurel formation at St. Paul are four specimens of arm-fans of dimensions far exceeding those of the type species, but which at once invite comparison with the two above mentioned. Three of them exhibit remarkable uniformity of characters, as shown by the following data: Length of arm-fan 23, 28, 28 mm.; greatest width 22, 25, 26 mm.; number of finial grooves 30, 30, 28; angle of fan 70° , 65° , 68° . The grooves are divided into two closely symmetrical halves.

Two of these specimens are figured; the first, plate 26, figures 15, 15a, enlarged, is a very well preserved fan with the grooved ventral and smooth dorsal surfaces exposed free of the matrix. The second, figure 16, natural size, shows only the ventral surface; and in both the division of the ventral grooves into two symmetrical halves, and the paddle-shaped handle of the fan at the proximal end, are well marked, as they are also in the third specimen of about the same size and form, not figured.

Of Bather's two American species we would naturally expect to find the prototype of our specimens in the one from St. Paul, *P. longus*; but the disparity in dimensions, and its extreme attenuation of form as indicated by its angle of only 38° , must clearly differentiate it; while his Iowa species, closely approximate in size, with 28 grooves in symmetrical halves, and angle of 70° , seems to fit the case exactly. Therefore I am referring the three specimens confidently to *P. inferior*.

The fourth specimen, not figured as it only shows the amorphous dorsal side, is much larger and more elongate than the others, being 45 mm. in length, with angle of 55° , probably belongs to *P. longus*, but the absence of other characters makes comparison unsatisfactory.

Horizon and locality. Hopkinton and Laurel formations; Monticello, Iowa, and St. Paul, Indiana; Dr. Foerste has also obtained a well marked specimen in the equivalent dolomite at Cedarville, Ohio.

Family CROTALOCRINIDAE Angelin

Genus **CROTALOCRINUS** Austin

Plate 26

Crotalocrinus Austin, Ann. Mag. Nat. Hist., vol. 10, 1842, p. 109.—Wachsmuth and Springer, Rev. Pal., 3, 1886, p. 145; Proc. Acad. Nat. Sci. Phil., 1888, pp. 364-390, pls. 19, 20.—Bather, Treatise on Zool., pt. 3, 1900, p. 176.—Weller, Jour. Geol., 10, 1902, p. 532.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 216.—Bassler, Bibliogr. Index, vol. 1, 1915, p. 292.

While there is a certain analogy between this and the preceding genus owing to the fact that in both the arm-branches do not move independently as usual in crinoids, but are more or less connected laterally within the rays, upon which ground the two were thought by Weller when describing *Petalocrinus* to be closely related, there are also broad differences between them. Here the five rays, instead of being rigidly fused, have their many branches connected at short intervals so as to form a flexible network, in which the outlines of the brachials are plainly delineated on both dorsal and ventral surfaces; these are connected laterally by points of attachment from near the middle of each ossicle, with open spaces between them, forming innumerable elongate meshes; owing to these lateral projections the brachials have the form of a cross with short arms; they are long and flat on the dorsal surface, and deeply grooved for the ambulacra on the ventral. This pliant network is capable of considerable motion by way of folding and stretching, and in one species, *C. pulcher*, forms five broad, reticulate leaves which when closed over the calyx overlap one another; and in another, *C. rugosus*, it is continuous for some distance around the calyx, and when spread out may be inrolled from the distal margin.

In the formation of the calyx the two are also strongly different. Instead of having pentamerous symmetry as in *Petalocrinus*, the dorsal cup of *Crotalocrinus* is divided by an anal plate in line with the radials, after the manner of the Cyathocrinidae, and the infrabasals are large and prominent. In *Petalocrinus* the ray is articulated to the calyx by means of a single primibrach filling the entire radial facet, whereas in *Crotalocrinus* the lower brachials as far as the third or fourth order rest within the radial facet, and are more or less rigidly incorporated into the dorsal cup by sutural connection among themselves, with the radials and with tegmen plates. It is only higher up in the rays, where the plates are no longer in contact

with the radials, that the muscular articulation admitting motion begins. The systematic position of the genus has been differently interpreted by authors, and is evidently intermediate, pointing on the one hand to a connection with the Camerata, and on the other with the Inadunata, with the preponderance in favor of the latter.

A more comprehensive discussion of this remarkable genus, with ample illustrations, may be found in Wachsmuth and Springer's paper of 1888, entitled "*Crotalocrinus*, its Structure and Zoological Position."¹ It contains an especially fine picture of the form having the reticulate rays continuous part way and inrolled from the ends, for which I have not room enough on my plate. The genus, although rare, is one of the most characteristic of the specialized forms in the Silurian of England and Gotland.

Crotalocrinus has not been found thus far within the area embraced in this work, but it is liable to appear at any time, as it has been recognized in the Racine dolomite of the Chicago area by Weller, and recently in the Silurian of Gaspe by Schuchert, thus adding another significant link to the chain of connection of the American Silurian with that of northern Europe. For this reason, as well as for comparison with the other highly specialized *Petalocrinus*, I have thought it would be of service to give some figures showing the characteristics of the type.

***Crotalocrinus cora* (Hall)**

Plate 26, fig. 17

Cyathocrinus cora Hall, 20th Rep. New York St. Mus., 1868, p. 324, pl. 11, figs. 13, 14; rev. ed., 1870, p. 366.—*Crotalocrinus americanus* Weller, Bull. 4, Chicago Acad. Sci. Nat. Hist. Surv., 1900, p. 143, pl. 14, fig. 1; *Crotalocrinus cora*, Jour. Geol., 10, p. 532, pl. 3, figs. 1-5.—Slocom, Field Columb. Mus., 2, Geol. Ser., 10, 1908, p. 292, pl. 86, figs. 3, 4.—Bassler, Bibliogr. Index, vol. 1, 1915, p. 292.

This species occurs in the Racine dolomite at Racine, Wisconsin, and at Chicago, described at an early day but its generic affinities not recognized until long after. Its history and characters are fully set forth in Dr. Weller's paper above cited, reinforced by further illustrations by Slocom, leaving no doubt of the correctness of his reference to the genus. By way of confirmatory evidence I give a figure of another specimen from Chicago showing an intensity of surface sculpture similar to that of the English species following.

***Crotalocrinus rugosus* (J. S. Miller)**

Plate 26, figs. 18-20

Cyathocrinus rugosus Miller, Nat. Hist. Crin., 1821, p. 89, with plate.—*Crotalocrinus rugosus* Austin, Ann. and Mag. Nat. Hist., vol. 11, 1843, p. 198.—Angelin, Icon. Crin. Suec., 1878, p. 26, pls. 7, fig. 4; 17, figs. 8, 8a (not figs. 3, 3a, b).—Wachsmuth and Springer, Proc. Acad. Sci. Phil., 1888, pp. 364, 386, pl. 19, figs. 1-5.

I am figuring a calyx of this species from the type locality at Dudley, England, for comparison with our Chicago specimens in the matter of surface markings and arrangement of plates, plate 26, figure 18. There are also two instructive structural figures, 19 and 20, from a Gotland specimen, the first showing the arrangement of the lower brachials in contact with the radial, and the second the mode of connection between the brachials of abutting arm-branches with the projecting processes midway and the open spaces between them forming the flexible net work; and also the covering plates of the ambulacra.

¹ Proc. Acad. Nat. Sci. Phil., Nov., 1888, pp. 364-390; pls. 19, 20.

Crotalocrinus pulcher (Hisinger)*Plate 26, fig. 21*

Cyathocrinus pulcher Hisinger, Leth. Suecc., Supp. 2, 1840, p. 6, pl. 29, figs. 5a, b.—*Crotalocrinus pulcher* Angelin, Icon. Crin. Suecc., 1878, p. 26, pls. 7, figs. 5-7; 8, figs. 1-9; 17, figs. 1a-d; 25, figs. 8-20.—Wachsmuth and Springer, Proc. Acad. Nat. Sci. Phil., 1888, p. 386, pl. 20, figs. 1a, b.

The figure of a very complete specimen of this species is given to afford a general idea of the formation and arrangement of the reticulate flexible appendages into which the rays are formed by the lateral connection of the arm-branches as illustrated in figure 21. In this form the network is divided into five leaf-like fronds which when folded overlap one another. The one in which the reticulation is continuous around the calyx and inrolled from the distal margin is shown in the Wachsmuth and Springer paper, plate 19, figure 1a, which should be consulted for comparison. The species occurs both in Gotland and England.

Family CYATHOCRINIDAE Roemer

Dicyclic; IBB usually 5. Radial facets usually semicircular, less than width of radial; union with brachials by incomplete articulation, without transverse ridge. Arms uniserial, non-pinnulate, mostly dichotomous. Tegmen composed of rather large orals, rigid ambulacrals, which often encroach upon and obscure the other plates.

Thalamocrinus Miller and Gurley*Plate 26*

Miller and Gurley, Bull. 7, Illinois St. Mus., 1895, p. 82.—Bather, Treatise on Zool., 3, 1900, p. 204.

Dicyclic. IBB 5, large, about one third the height of cup; BB still larger, nearly half the height of cup; RR much shorter, not over one sixth the height of cup; radial facets curved, shallow, narrower than RR, directed upward. RA small, rhombic, abutting on *x*, supported by post. and r. post. BB. Anal *x* small, in line with RR. Arms slender, with elongate brachials, uniserial, number unknown.

Genotype. *Thalamocrinus ovatus* Miller and Gurley.

Distribution. Silurian to Devonian; America.

This genus falls under the Carabocrininae subdivision of the Cyathocrinidae, with RA similar to that of *Palaeocrinus*. It is rather near to *Homocrinus*, as interpreted by Bather in connection with his Gotland species, Crin. Gotl., p. 101, pl. 4, figs. 143-147, but has narrower facets and more simple arms. Miller and Gurley mention the "radials or third series of plates" as six, thus including what we know as the anal *x*. They missed the important element, the radianal, both in their descriptions and figures; but it appears plainly in all four of the species, including some of the specimens examined by the authors from the collection of Mrs. Milligan. It is a rare form, and most of the principal specimens are figured here.

Thalamocrinus ovatus Miller and Gurley*Plate 26, figs. 2-5*

Miller and Gurley, Bull. 7, Illinois St. Mus., 1895, p. 82, pl. 5, figs. 29-31.—Miller, N. A. Geol. Pal., 2d App., 1897, p. 754, fig. 1402.

Characterized by its elongate ovoid form, which is well maintained among several specimens, the great height of BB, and constriction at the distal margin, whereby the space is reduced to very narrow facets for slender arms. In the only specimen with any part of the arms preserved they are shattered so that little can be seen except the displaced brachials, which are long, narrow and rounded; it is evident that the arms were long and slender, probably simple or only branching once.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

Thalamocrinus cylindricus Miller and Gurley*Plate 26, figs. 6-9*

Miller and Gurley, Bull. 7, Illinois St. Mus., 1895, p. 82, pl. 5, figs. 32, 33.—Miller, N. A. Geol. Pal., 2d App., 1897, p. 754, fig. 1401.

Differs from the type species in its narrow, cylindrical form, greater relative height of IBB, and less constriction at the distal margin.

Horizon and locality, same as last.

Thalamocrinus globosus new species*Plate 26, figs. 10-10b*

Calyx short and almost perfectly spherical, but otherwise similar to *T. ova-tus*, of which it may be only a variety. This and the two preceding species have a narrow column-facet, followed by an expanding base.

Horizon and locality, same as last.

Thalamocrinus elongatus new species*Plate 26, fig. 11*

In the unique specimen of this Devonian form of the genus the RR are missing, but otherwise the cup above the column shows the essential elements, including the RA, similar to the foregoing species. But the basal parts are widely different; instead of the BB curving inward to a narrow facet, they are enveloped to about their full width by the proximal ossicle of a large column which expands broadly towards the calyx. In general contour it resembles *T. cylindricus*, but is considerably larger than any of the other species.

Horizon and locality. Linden formation, Helderbergian, Lower Devonian; Benton County, Tennessee.

(?) Ampheristocrinus typus Hall*Plate 31, fig. 1*

Hall, 11th Ann. Rep. Indiana Dep. Geol., 1882, p. 278, pl. 15, figs. 17, 18.—Wachsmuth and Springer, Rev. Pal., 3, 1886, p. 219.—Bassler, Bibliogr. Index, 1915, p. 32.

A complete crown with elongate, slender and sharply angular cup; long, delicate arms bifurcating at a rather wide angle three or four times; radial facets curved and very narrow; arms uniserial with rather long, rounded brachials. IBB elongate, their number uncertain. Anal tube long, projecting beyond the arms. Stem stout and long, composed of short, alternating columnals.

This genus and species were founded upon a dorsal cup only, without either arms or stem. It is characterized by 3 IBB and a prominent anal area with an oblique RA, and large anal α . Neither of these is shown by the unique specimen figured, which is imbedded upon the posterior side, thus leaving the generic characters in doubt. But the calyx with its sharp

longitudinal angularity and its extremely narrow radial facets is so remarkably like the type, that we can readily conceive that the delicate arms would go with it, and that the other characters probably are present. The genus has been but little known, and this specimen certainly shows how it ought to look, so I am venturing to place it accordingly, despite the difference in horizon.

Horizon and locality. Beech River formation; Decatur County, Tennessee; type is from Waldron shale, Waldron, Indiana.

(?) ***Lecythiocrinus problematicus*** new species

Plate 31, figs. 11, 11a, b

Lecythiocrinus White, Proc. U. S. Nat. Mus., 2, 1880, p. 256, pl. 1, figs. 4, 5.—Worthen, Geol. Surv. Illinois, 7, p. 317, pl. 31, fig. 8.—Wachsmuth and Springer, Rev. Pal., 3, 1886, p. 228.

Calyx small, low and broad. IBB 3, the small plate in r. post. position. BB unequal, all rather small except post. B, which is much the largest, rising high up between RR. RR large and massive, with facets deeply horse-shoe-shaped; dorsal canal about the middle, and food groove at upper margin. Anal opening directly through the calyx wall at the junction of post. B and the two post. RR, carved about equally out of each of them. There is no plate above it analogous to an anal α , the two RR abutting directly. Tegmen apparently composed of interlocking orals, the sutures of which cannot be made out; ambulacra with alternating covering pieces pass from the radials under the orals.

While this species has a strong superficial resemblance to some of the Devonian forms embraced in the subfamily Gasterocominae, it is definitely excluded from them by its lack of undivided infrabasal disk and peripheral axial canals, which are so characteristic of them. A later genus with which it would seem to agree in part is the little known *Lecythiocrinus* from the American Upper Carboniferous, having 3 IBB and a simple axial canal, but the radial facets are directed differently. Although not stated in the diagnosis, this genus is now known to have the anal opening through the dorsal cup as in the Gasterocominae, as was indicated by Worthen when describing *L. adamsi*, and confirmed by specimens in my possession since discovered. Comparison should also be made with *Hypocrinus* of Beyrich from the Permian of Timor, fully described by Wanner in his treatise on the Timor crinoids. The reference here made is admittedly doubtful.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

CYATHOCRINUS Miller

Plate 31

Cyathocrinus J. S. Miller, Nat. Hist. Crin., 1821, p. 85.—Wachsmuth and Springer, Rev. Pal., 1, 1879, p. 79; 3, 1886, p. 21.—Bather, Crin. Gotl., 1893, p. 126; Treatise on Zool., 3, 1900, p. 173.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 220.—Bassler, Bibliogr. Index, 1915, p. 316 (complete synonymy).

IBB 5. No RA. Anal α in line with RR; anus at end of a tube short and rounded, or long with valvular pyramid at distal end. Radial facets horse-shoe-shaped, directed outward. Arms dichotomous, branching numerously at rather wide angles.

Genotype. *Cyathocrinus planus* Miller.

Distribution. Silurian to Lower Carboniferous; Europe and America.

Cyathocrinus decatur new species*Plate 31, fig. 2*

A complete crown, with elongate cup perfectly smooth and rounded. IBB elongate, probably 5, anal side not exposed. BB large; RR small, facets distinctly facing outward, making arm-bases horizontal. Arms stout, with three or four wide-angled bifurcations; brachials longer than wide. Stem wide, the full width of base, with columnals lengthening distally until as long as wide.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

Cyathocrinus cf. striolatus Angelin*Plate 31, figs. 4-7*

Cyathocrinus striolatus Angelin, Icon. Crin. Suecc., 1878, p. 23.—Bather, Crin. Gotl., 1893, p. 136, pl. 7, figs. 218-221.

Dorsal cup forming a truncate cone with nearly straight sides; plates folded into sharp ridges radiating from plate to plate, leaving distinct pits at the corners; surface smooth to tubercular. IBB fairly high, with lower margin slightly raised into a rim, forming a broad, truncate base. BB large. RR large with fairly wide, curved facets. Anal α small, scarcely half the size of RR. Axial canal large.

I have referred the four specimens here figured to the Gotland species with considerable confidence, but the arms if known might disclose a specific difference. In three of them the surface markings are sharp, but in the fourth they are smoothed by erosion. The IBB are not uniform in size, one being usually smaller than the others, and in the specimen of figure 6 there are six IBB, thus indicating a tendency to instability in this element.

Horizon and locality. Laurel limestone; St. Paul, Indiana.

Cyathocrinus wilsoni new species*Plate 31, figs. 3, 3a*

Distinguished by its conical calyx, expanding from a very small column-facet, and the great length of its deeply rounded brachials; these are not entirely uniform, and some are more than twice as long as others. RR facets deeply excavate; arms bifurcating widely at very short intervals in lower part, some with only one brachial. IBB 5, small, and RR much larger than BB. Anal α of true Cyathocrinid type, but smaller than usual.

Horizon and locality. Laurel limestone; St. Paul, Indiana. Named for the late Dr. Herrick E. Wilson, who collected the type.

Parisocrinus siluricus new species*Plate 31, figs. 8, 8a*

Parisocrinus Wachsmuth and Springer, Rev. Pal., 1, 1879, p. 115; *ibid.*, 3, 1886, p. 221.—Springer, New Amer. Foss. Crin., Mem. M. C. Z. Harvard, 1911, p. 157.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 218.

This primarily Lower Carboniferous type, occurring sparsely in the Devonian, must now be moved a step further back in the geological scale as the result of the discovery of the fine species here illustrated. The genus has been described as a Cyathocrinid with Poteriocrinid anal side—that is, it has curved radial facets only part the width of the radials, and non-pinnulate arms, combined with a large radianal; as a further Cyathocrinid character it has the anal opening at the distal end of a large tube. These characters are strongly emphasized in the present species. It has a smooth, expanding turbinate calyx with 5 large IBB, large RR with facets directed upward; rather stout arms bifurcating twice or oftener, with fairly long brachials. The tube is large in proportion to the calyx, as usual in the genus, and is heavily plated to the end. RA is only partly shown in figure 15a, but it is nearly as large as *x*. Column large, composed of alternating columnals.

Horizon and locality. Laurel limestone; St. Paul, Indiana.

Gissocrinus Angelin*Plate 32*

Gissocrinus Angelin, Icon. Crin. Suecc., 1878, p. 10.—Wachsmuth and Springer, Rev. Pal., 1, p. 89.—Bather, Crin. Gotl., 1893, p. 152; Treatise on Zool., 3, 1900, p. 175.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 220.

Similar to *Cyathocrinus*, but with less than 5 IBB, and a tendency to greater surface sculpture and variation in arms.

Genotype. *Gissocrinus typus* Bather.

Distribution. Silurian; Gotland, England, America.

This genus, one of the most prominent Inadunate crinoids of Gotland and England, has not hitherto been known in America. It is now represented in our Silurian fauna by several species, one of which presents characters wholly novel. Nine species are recognized from Gotland, two of which are also well known in England, and one is common at Dudley only. The Gotland species are all fully described in Bather's work of 1893, above cited.

Gissocrinus lyoni new species*Plate 32, figs. 1, 1a*

A rather massive species, of the type of the English species *G. goniadactylus* Phillips (*Cyathocrinites goniadactylus*, in Murchison's Silurian System, 1839, p. 671, pl. 17, fig. 1). IBB 3. Cup rather low, broadly spreading. Arms stout,

angular at the back, branching but few times. It is based on a single specimen from a higher horizon than those following, and I have been able to expose the crown fully for both lateral and dorsal views.

Horizon and locality. Louisville limestone; Jefferson County, Kentucky.

Gissocrinus delicatus new species

Plate 32, fig. 2

A small species, of the type of *G. typus* Bather, Crin. Gotl., p. 155, but with cup more slender and turbinate, and arms more delicate with numerous bifurcations. IBB apparently 3, followed by very elongate BB. Anal tube extremely long, projecting beyond the tips of the arms. There is only the one specimen, very well preserved.

Horizon and locality. Beech River formation; Decatur County, Tennessee.

Gissocrinus magnibrachiatus new species

Plate 32, figs. 3-6

Of very large size, far larger than any other known species; crown so far as preserved about 75 mm. high. Calyx small, crushed in the only specimen preserving it and its composition unknown. Arms of relatively enormous size compared with the calyx, composed of long, broad, low convex and very thin brachials with smooth surface, ranging from 10 to 14 mm. high to 7 to 10 mm. wide, with frequent bifurcations in the lower part, even from successive brachials; articulating faces of brachials connected by 1, 2, or occasionally 3 small patelloid plates, most frequently 2 in succession, arising from deeply curved facets occupying less than half the width of the brachial; the distal faces of the axillaries have 2 pairs of patelloid plates abreast and near the lateral margins, while on the other faces the patelloid is in the middle; they are as thick as the brachials, and pass through to the inside. The positions of these connecting plates coincide exactly with the course of the ventral furrows or food grooves on the inner side of the brachials, which branch upon the axillaries and are doubled at the distal end; the grooves are extremely narrow, not over one fifth the total width of the brachial, and are closed by a double row of small, alternating covering plates not over half a millimeter long and wide.

Having assembled an excellent series of specimens of the principal English species, the study of these in connection with Bather's exhaustive illustrations of the Swedish forms gave me a mental picture of this as a genus of wide variations, especially in the arms, under which the unexpected was always liable to occur. With this in mind, when the first specimens of this extraordinary form were disclosed among the Tennessee collections, I labeled them *Gissocrinus* without any very definite reason for it. As other specimens appeared showing complete constancy of characters hitherto without precedent, I began to wish for assistance toward the understanding of it, and accordingly sent a photograph of one of the most instruc-

tive specimens to Dr. Bather at the British Museum, requesting his opinion on the tentative identification I had made. His answer was that upon first looking at the picture, before reading anything I had written about it, he had said "*Gissocrinus*," but without knowing why. So in view of our scant knowledge concerning the structure of the calyx, these two concurrent guesses will have to be taken as the chief authority for the generic reference.

The most remarkable feature of the species is the small oval connecting plates at the brachial articulations, sometimes two, or even three, in succession, and located precisely where the food grooves, whether one or two, cross the faces of the brachials; they are not confined to the dorsal side, but extend entirely through to the interior, where they are equally prominent, as is beautifully shown by figure 6. I know of nothing else like them in crinoid morphology. The term "patelloid plates" which I have applied to them will doubtless recall the term as used by Hall for some supposed intervening plates connecting successive brachials in *Forbesiocrinus* and other genera; but those, as I have shown at length in the *Flexibilia* monograph of 1920, pages 44, 112, 239, etc., are not plates, but only superficial processes from the proximal face of brachials at the dorsal side, which sometimes get broken off and thus may resemble plates. This crinoid must have been of fragile construction, as indicated by the extreme thinness of the wide-spreading brachials; this is well shown by figures 3 and 6, where several of them are seen superimposed in two or more layers. They recall the paper-like calyx plates of *Uintacrinus*.

Horizon and locality. Beech River formation; Tuck's Mill, Decatur County, Tennessee.

***Gissocrinus approximatus* new species**

Plate 32, figs. 7-9

Fragmentary brachials found at St. Paul indicate that forms similar to the last exist in the Laurel limestone, with an important difference in the apparent absence of the patelloid plates. We have isolated axillary brachials with food-grooves both open and covered, as figured, one of them of even greater length than in the Tennessee specimens.

Horizon and locality. Laurel limestone; St. Paul, Indiana.

***Gissocrinus quadratus* new species**

Plate 32, figs. 10, 11

I figure under this name an axillary plate and the two following it which are nearly rectangular, and clearly devoid of any patelloid plates, and also a series of nearly square plates with the food-grooves covered. No other parts of this or the preceding species were found, but their occurrence is important as evidence of the wide distribution and variable nature of this peculiar type.

Horizon and locality, same as last.

***Botryocrinus tenuidactylus* new species**

Plate 31, fig. 9

Botryocrinus Angelin, Icon. Crin. Suecc., 1878, p. 24.—Wachsmuth and Springer, Rev. Pal., 1, 1879, p. 97; 3, 1886, p. 191.—Bather, Ann. Mag. Nat. Hist., 7, 1891, p. 392; *ibid.*, 9, 1892, p. 189; Treatise on Zool., 3, 1900, p. 113, figs. 21, 179.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 221.—Bassler, Bibliogr. Index, 1915, p. 130.

This genus is chiefly distinguished from the Cyathocrinidae by its mode of arm-branching, which is mostly of the heterotomous type, the rays dividing

into two main rami which give off ramules, sometimes approaching pinnulation. It also has the RA small, rhomboidal, in the oblique position. RR facets curved, less than width of plate. Our specimen unfortunately lacks the calyx, but the set of arms figured has a facies not dissimilar to that of the English species *B. ramosus*, but the arms are more slender and brachials longer. The ramules are long, and in turn bear pinnules which are apparently confined to the side next to the main arm. No other specimen was found.

Two species from the Waldron shale described by Hall from the calyx only as *Dendrocrinus nucleus* and *Cyathocrinus polyxo*, have been referred to *Botryocrinus*, and I am figuring a calyx perhaps of the latter, from Hartsville, Indiana, seen from the anal side, to show the type which might be expected to go with these arms, plate 31, figure 10.

Horizon and locality. Beech River formation; Decatur County, Tennessee.

INSERTAE SEDIS

Parastephanocrinus typus new genus and species*Plate 31, figs. 12-12c*

Dicyclic. Calyx small, low and broad, height to width 5 to 7 mm., with a concave base. IBB 5, at the bottom of the cavity. BB large, elongate, forming more than half the bulk of the calyx, curving upwards, and elevated into high median ridges which branch to the next plates; post. B rises above the others to about half the height of RR, curving inward to the anal opening. RR much smaller exteriorly, but massive, extending deeply in toward the oral center, and forking into strong projecting processes at either side which meet and abut in pairs, except at the posterior interradius where they are separated by the anus; they are deeply indented medially by grooves for ambulacra toward the central cavity, with narrow, round facets facing outward, and occupying about one third the width of the plate. Anal opening at the margin between RR and post. B. Plates of tegmen unknown, but the central space was probably filled by solid orals.

The first thought on looking at the ventral side of this crinoid is of the ever-puzzling *Stephanocrinus*. The deep ambulacral grooves, and the paired processes of the forked radials, are substantially on the same plan; the open space at the center is just right to accommodate a set of interlocking orals, with covered ambulacra passing under them. But the dicyclic base and concave cup must exclude it from that genus, and the only recourse is to propose a new genus for its reception; in doing so, I recognize how limited our understanding of it is. From the form and position of the facets it is improbable that the arms could have been like those of *Stephanocrinus*; but at all events they were small. As an aid to the discussion, I give some figures showing the leading characters of *Stephanocrinus*.

Horizon and locality. Louisville limestone, Niagaran; 1¾ miles east of Bob, Decatur County, Tennessee.

Stephanocrinus angulatus Conrad*Plate 31, figs. 13-16*

Stephanocrinus Conrad, Jour. Acad. Nat. Sci., Phil., 8, 1842, p. 278; p. 279, pl. 15, fig. 18.—Wachsmuth and Springer, Rev. Pal., 3, 1886, p. 291.—Bather, Treatise on Zool., 3, 1900, p. 97, fig. 2, p. 145.—Zittel-Eastman, Textb. Pal., 2d ed., 1913, p. 207.—Jaekel, Phylogenie und System, 1918, p. 109.—Wanner, Permischen Blastoiden von Timor, 1924, p. 186.—Bassler, Bibliogr. Index, 1915, p. 1187 (complete synonymy and list of species).

This genus is of debatable systematic position, and has been variously considered by different authors as a crinoid, a cystid or a blastoid. In favor of the first view are the branching, biserial arms; while the forked radials, resemblance of the orals to the deltoids, and the orientation of the small basal in the r. ant. interrarial position instead of the l. ant., as in crinoids with 3 basals, are all characters which tend toward the blastoids. I am figuring some specimens of the type species from Lockport, New York, and another from Tennessee, for

comparison with the preceding species, to show the extremes in general form of calyx, variation in the forked radials, the orals covering the central space of the tegmen, and a detail of the arm clusters.

S. angulatus Conrad, the type, is from the Rochester shale, and *S. gemmiformis* Hall, plate 31, figures 17-19, a species of wide distribution and in many ways the opposite of the former, was described from the same horizon and locality, as well as from the Waldron formation of Indiana, and is also found at Newsom, and other localities in Tennessee.

The BLASTOIDEA and CYSTIDEA

It has not been my intention to make a critical study of these two classes; yet in view of their intimate association with the crinoids in the formations here discussed, it seems advisable to illustrate the more prominent forms which have appeared in the collections. These include two species of blastoids and seven of cystids.

Troostocrinus reinwardti (Troost)*Plate 33, figs. 1-8*

Pentremites reinwardti Troost, Trans. Geol. Soc. Pennsylvania, 1835, p. 224, pl. 10, figs. 9-12; 5th Geol. Rep. Tennessee, 1840, p. 55.—*Pentatremites reinwardti* Roemer, Sil. Fauna Westl. Tennessee, 1860, p. 60, pl. 3, figs. 2a-c.—*Pentremites (Troostocrinus) reinwardti* Shumard, Trans. Acad. Sci. St. Louis, 2, 1866, pp. 384, 385.—Bassler, Bibliogr. Index, 1915, p. 1305; complete synonymy of genus and species.

This species has been well known by weathered specimens in the Tennessee glades from the earliest times, but it is only with the present collections that its prolific occurrence in a definite horizon became manifest. Here in the middle zone of the Beech River shales it was obtained in place in large numbers, being so characteristic that the generic name was attached to that subdivision. It is probable that most of those from the glades were derived from erosion of the same bed. The species has a wide distribution throughout the Tennessee area, and has also been reported from the Louisville limestone of Kentucky.

While specimens from the glades usually have the extremely narrow base broken off, the preservation of those found in the shales is excellent, presenting in a well defined type a considerable variation in size and proportions. I am figuring a representative selection ranging from 10 to 45 mm. in height; and along with these a unique specimen with the brachioles intact—a condition not before seen.

Horizon and locality. *Troostocrinus* zone of the Beech River formation, Niagaran; Tuck's Mill and on glades in Decatur, Wayne and Perry counties, Tennessee.

Troostocrinus sanctipaulensis Foerste*Plate 33, fig. 9*

Ohio Jour. Sci., 21, 1920, p. 64, pl. 1, fig. 16.

An imperfect specimen from the Laurel limestone at St. Paul; differing but little from the preceding.

Tetracystis fenestratus Schuchert*Plate 33, figs. 10-14*

Echinoencrinites fenestratus Troost, Am. Jour. Sci., (2) 8, 1849, p. 419 (not defined); Bull. 64, U. S. Nat. Mus., 1909, p. 8.—*Tetracystis fenestratus* Schuchert, Smithsonian Misc. Coll., 47, 1904, p. 219, pl. 34, figs. 6-8.—Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 8.—Bassler, Bibliogr. Index, 1915, p. 1263.

This form was recognized by Troost in 1849, and formed the subject of an elaborate description in his then unpublished monograph, as now appears in the U. S. National Museum Bulletin 64. In the meantime Schuchert's description was published, based upon Troost's solitary specimen, with proper credit to him. The species is quite rare, not having been found by any of the collectors since Troost and Roemer until the material now in hand was obtained in the *Eucalyptocrinus* zone at the Tuck's Mill excavations. This consists of eight specimens, of which I am figuring five, giving lateral views from four positions and one ventral. The circular cross-section of the calyx, the three abutting pairs of pectinirhombs, the four open ambulacral grooves traversing the surface longitudinally, are all thoroughly shown. The grooves were bordered by about 11 pairs of brachioles marked by their slightly raised bases,

and in one instance traces of the brachioles themselves are seen in place. The hydropore exhibits a tendency to doubling, some specimens having two pores. The surface of the plates is deeply sculptured into rugose ridges or folds, mostly vertical to the sutures; the preservation of this ornament is variable, but in several specimens it is very sharp and distinct. In the last three characters there are some differences from the type as described, which may be due to individual variation, or upon closer study may warrant the proposal of a new species.

It is interesting to note, as bearing upon the relation of the American and European Silurian, that Angelin described and figured a remarkably similar form from the Wenlockian of Gotland,¹ and referred it to the American species *Lepadocrinus gebhardi* Conrad. Another of similar type has been described from the English Silurian under the name *Pseudocrinus quadrifasciatus*, illustrated by Bather in Lankester Zoology, page 62. On the other hand *Caryocrinus*, the most prolific American type, is not represented in the European Silurian.

Horizon and locality. Beech River formation, *Eucalyptocrinus* zone, Niagaran; Decatur County, Tennessee.

Stribalocystites gorbyi S. A. Miller

Plate 33, figs. 15-18

Stribalocystites gorbyi Miller, 18th Ann. Rep. Indiana Dep. Geol., 1894, p. 265, pl. 2, figs. 3-8.—Bassler, Bibliogr. Index, 1915, p. 1209, for genus and species.

This peculiar genus, which is said by the author to have no arms, is represented by a series of good specimens of this small species from the type locality. It is notable for the extremely small size of the tegmen, which is composed of only 5 or 6 plates without any fixed plan of arrangement. Arm-facets are certainly present, though obscure, 3 to 5 in number. Otherwise the structure is that of *Caryocrinus*, the folds and raised pores being in a variety of conditions. In general form and appearance of the calyx in a lateral view the species seems quite similar to *Caryocrinus bulbulus*.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Lysocystites sculptus (S. A. Miller)

Plate 33, figs. 19-22

Lysocystites Miller, N. A. Geol. Pal., 1889, p. 259.—Bather, Treatise on Zool., 3, 1900, p. 70.—*Aethocystites sculptus* Miller, 18th Ann. Rep. Indiana Dep. Geol., 1894, p. 264, pl. 2, fig. 2 (adv. sheets, 1892, p. 10).—*Lysocystites sculptus* Foerste, Ohio Jour. Sci., vol. 21, 1920, p. 44, fig. 3.

Originally described from imperfect material which left most of the characters in doubt. In 1920 Dr. Foerste, from a study of the specimens here figured together with the types, worked out the plate system of this species, showing that it belongs to the genus *Lysocystites* proposed by Miller for the preoccupied *Echinocystites* and referred by Bather to the family Cryptocrinidae of the rare order Aporita, without pores or rhombs. The cup consists of three circlets of plates, the first containing 3 unequal plates, the equivalent of 5 with two pairs fused; the second 5 very large elongate plates, obtusely hexagonal, and about as high as the first and third combined; the third 5, pentagonal, converging to the tegmen. A prominent tubular fold runs from the column to each salient angle of the base, thence to the center of each plate of the second circlet and on to the upper corners, and from there less prominent up the middle of those of the third circlet. The wall of these folds is very thin, and they are frequently broken down, leaving open fissures; this is especially true in the third circlet, where with considerable uniformity large longitudinal openings are exposed. The tubular folds enclose diamond-shaped areas which are traversed by low vertical striations extending up the middle of the plates for their full height, with transverse bands of striae from center to center of the large plates of second circlet. The anal opening is outside the tegmen toward the apex of plates of third circlet.

¹ Icon. Crin. Suecc., 1878, p. 32, pl. 11, figs. 29-35; pl. 19, figs. 18a-c.

The tegmen is extremely small, in mature specimens about one fourth the greatest diameter of the cup; it contains a fourth circlet of five small plates enclosing a central space, from which shallow food-grooves extend outward. The apparently central plate shown by the drawing in fig. 22 proves to be a foreign substance.

This form is represented by four good specimens, two large and two small, ranging in height from 30 down to 10 mm. The former are typical for the species, while the latter, figs. 21, 22, represent a more globose species which may take the name *Lysocystites globosus*.

Horizon and locality. Laurel limestone, Niagaran; St. Paul, Indiana.

Caryocrinus bulbulus Miller and Gurley

Plate 33, figs. 23-27

Caryocrinus bulbulus Miller and Gurley, Bull. 5, Illinois St. Mus., 1894, p. 11, figs. 15-18.—Bassler, Bibliogr. Index, 1915, p. 186.

I have referred a series of five good specimens to this species, on account of the general contour and the strong protuberance on the middle of the second range of plates. In the description the surface is said to be otherwise smooth, but here it is marked by a variety of sculpture. They all have a juvenile aspect, and may be merely the young of *C. milliganae*.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

Caryocrinus persculptus new species

Plate 33, fig. 28

A strongly marked form of turbinate contour, in which the entire surface is covered with such an intense sculpture that the folds and pores are transformed into solid, rugose ridges, strongly elevated, not only on the lines of folds but between them. It is from a different horizon from all the described species, none of which agree with it in the characters stated. The one that nearest resembles it in surface markings is *Caryocrinus ellipticus* Miller and Gurley, from the Osgood.

Horizon and locality. Waldron shale, Niagaran; Newsom, Tennessee.

Caryocrinus milliganae Miller and Gurley

Plate 33, figs. 29-36

Caryocrinus milliganae M. & G., Bull. 9, Illinois St. Mus., 1896, p. 63, pl. 5, figs. 3, 4.—Wood, Bull. 64, U. S. Nat. Mus., 1909, p. 10.—Bassler, Bibliogr. Index, 1915, p. 186, for genus and species.

This is the most abundant cystid in the Tennessee area, and is subject to considerable variation in general form, from pear-shaped to ovoid, and from rotund to elongate. It attains a large size. The stereom folds radiating from the middle of the plates, and the lines of pores with their raised margins, characteristic of the genus, are abundantly shown in this material.

Troost proposed three species for different varieties, all of which are listed under the present species in Miss Wood's edition of his monograph. Among the earlier collectors this form was always referred to *C. ornatus* Say. I am figuring five rather well marked types, which will give a good idea of the range of variation shown by the collection, supplemented by an instructive figure of Say's well known genotype from Lockport.

Horizon and locality. Beech River formation, Niagaran; Decatur County, Tennessee.

Caryocrinus ornatus Say

Plate 33, fig. 37

Jour. Acad. Nat. Sci. Phil., 4, 1825, p. 290.—Bassler, Bibliogr. Index, 1915, p. 187.

For comparison with the forms from the Tennessee area, and to show the structure of the biserial arms.

Horizon and locality. Rochester shale, Niagaran; Lockport, New York.

PERSONAL RECORD ¹

FRANK SPRINGER

LAS VEGAS, NEW MEXICO

SKETCH ²

Frank Springer, son of Francis and Nancy R. Springer, was born at Wapello, Iowa, June 17, 1848. He received his education in the public schools and at the State University of Iowa. He was admitted to the bar in Iowa in 1869, and came to New Mexico in 1873, settling at Cimarron, where on October 10, 1876, he was married to Josephine M. Bishop. He immediately took foremost rank in his chosen profession, and has since been recognized as one of the leaders of the New Mexico bar. In 1890 he was elected president of the New Mexico Bar Association, in which capacity he delivered an address dealing entirely with the urgent necessity for the immediate settlement by the Congress of the United States, through a proper tribunal, of titles under Spanish and Mexican land grants. The bill, which finally became a law, was drafted principally by him. The bar association attached so much importance to this address that it was printed in full, and distributed among members of Congress, and filed with the Land Department at Washington. The subject matter of the address produced a profound impression at Washington with the President of the United States and members of Congress alike. A direct result was the passage of the act establishing the court of private land claims. As attorney for the Maxwell Land Grant Company, Mr. Springer achieved his greatest professional success. In the trial of many causes in the courts of New Mexico in which that company was a party, and in the final determination of the title of the company to the lands embraced within the limits of the grant, as confirmed by Congress, decided in favor of the company in *United States vs. The Maxwell Land Grant Company*, his preparation, argument, and management were masterly, eliciting commendation from members of the Supreme Court of the United States. He is the president of the board of trustees of the Maxwell Land Grant Company, whose business affairs under his management and direction have been phenomenally successful. He retired from the active practice of his profession in 1906, since which time he has given much attention to scientific study and research. He was twice a member of the legislative council of New Mexico, in which position he rendered the territory great service. As a paleontologist he takes first rank among scientists in America. He is a member of the Archaeological Institute of America, and a patron of the School of American Archaeology at Santa Fé, New Mexico. Through the generosity of Mr. Springer the restorations, in part, and many of the mural decorations in the Old Palace, have been accomplished under the direction of Dr. Edgar L. Hewett, director of the school.

(Further note by R. E. T.):

To the union of Frank and Josephine M. Springer seven children were born, four daughters and three sons, viz.:

Laura, wife of Mr. John J. K. Caskie, General Attorney of the Philadelphia Rapid Transit Company, residing in Philadelphia.

Helen, wife of Dr. John F. Fairbairn, of Buffalo, New York.

Ada, wife of Dr. Warren B. Davis, of Philadelphia.

¹ From *Twitchell's Leading Facts of New Mexican History*, Vol. II, p. 453.

² [EDITOR'S NOTE.—The biographical material given on pages 144-5 and 153 to 166 is taken from a memorial volume compiled by Ralph Emerson Twitchell, life long friend of Mr. Springer; it is here included at the express wish of Mr. Springer who feels that the present paper perhaps may be his last personal contribution to science, and desires to present the background for his scientific activities in its relation to affairs in his state.]

FRANK SPRINGER

These three daughters pursued different lines of study in American colleges, supplemented by several years of work in Europe. Ada received the degree of Master of Arts from Columbia University.

The husbands of the last two are surgeons eminent in their profession, each of whom, as well as the first, left a lucrative practice to serve the government as a volunteer during the Great War.

Eva Springer, an artist, whose paintings have been accepted at the exhibitions of the Paris Salon, the London Academy, the Philadelphia Academy of Fine Arts and other principal art exhibitions of this country. A collection of miniatures by her formed one of the attractions of the New Mexico Museum for several years.

Major Edward T. Springer, of Cimarron, New Mexico. Manager of the live stock business in which the family are interested.

Lieutenant L. Wallace Springer, of Cimarron, New Mexico. Manager of the irrigation enterprise which built the Eagle's Nest Dam.

The two last mentioned boys represented the family in the Great War, serving in the A. E. F. at the front during the American campaign in France. Edward was a captain in the 21st Field Artillery of the Fifth Division, being promoted to that rank during the Argonne offensive, and to that of Major in the Reserve Army after the war. Wallace entered the aviation service, and was made a first lieutenant in the Eleventh Squadron, taking part in the air battles on the Argonne front, in one of which he was wounded a few days before the Armistice.

Henry S. Springer, formerly of Cimarron, New Mexico. A well known live stock grower, who was selected by the family to remain in charge of necessary productive industry during the war. He was prominent in all the home activities relating to the war, and rendered valuable service on the Council of Defense. He died of pneumonia in 1920.

BIBLIOGRAPHY

A. SCIENTIFIC WORKS

1. IN COLLABORATION WITH CHARLES WACHSMUTH

1877. Revision of the Genus *Belemnocrinus*, and Description of Two New Species. (American Journal of Science (3), vol. xiv, 1877, pp. 253-266.)
1878. Transition Forms in Crinoids. (Proc. Acad. Nat. Sci. Phil., 1878, pp. 224-266; 1 plate.)
1879. Revision of the Palaeocrinoidea, part i. (Proc. Acad. Nat. Sci. Phil., 1879, pp. 226-379; plates 15-17.)
1881. Revision of the Palaeocrinoidea, part ii. (Proc. Acad. Nat. Sci. Phil., 1881, pp. 177-414; plates 17-19.)
1883. Remarks on *Glyptocrinus* and *Reticocrinus*, Two Genera of Lower Silurian Crinoids. (Am. Jour. Sci. (3), vol. xxv, 1883, pp. 255-268; 1 figure.)
1883. *Hybocrinus*, *Haplocrinus* and *Baerocrinus*. (Am. Jour. Sci. (3), vol. xxvi, 1883, pp. 365-367.)
1883. Descriptions of Fossil Invertebrates. (Illinois Geol. Surv., vol. vii, 1883, pp. 339-345.)
1885. Revision of the Palaeocrinoidea, part iii, section 1. (Proc. Acad. Nat. Sci. Phil., 1885, pp. 225-364; plates 4-9.)
1886. Revision of the Palaeocrinoidea, part iii, section 2. (Proc. Acad. Nat. Sci. Phil., 1886, pp. 64-226.)
1887. Summit Plates in Blastoids, Crinoids and Cystids, and Their Morphological Relations. (Proc. Acad. Nat. Sci. Phil., 1887, pp. 82-114; 1 plate.)
1888. Discovery of the Ventral Structure of *Taxocrinus* and *Haplocrinus*, and Consequent Modifications in the Classification of the Crinoids. (Proc. Acad. Nat. Sci. Phil., 1888, pp. 337-362; 2 plates.)
1888. *Crotalocrinus*, Its Structure and Zoological Position. (Proc. Acad. Nat. Sci. Phil., 1888, pp. 364-390; 2 plates.)
1890. Perisomic Plates of the Crinoids. (Proc. Acad. Nat. Sci. Phil., 1890, pp. 345-392; 1 plate.)
1890. New Species of Crinoids and Blastoids from the Kinderhook Group of Le Grand, Iowa. (Illinois Geol. Surv., vol. viii, 1890, pp. 157-251; 3 plates.)
1892. Description of Two New Genera and Eight Species of Camerate Crinoids from the Niagara Group. (American Geologist, vol. x, 1892, pp. 134-144.)
1897. Monograph of the North American Crinoidea Camerata. 4to. 3 vols., 800 pp. text, and atlas of 83 plates. (Memoirs Museum of Comparative Zoology, Harvard College, vols. xx and xxi, 1897. Reviewed under title "An Epoch in American Science" in *Annals of Iowa*, 3d series, vol. ii, 1896, pp. 345-364; under N. A. Fossil Crinoidea Camerata in Report of Iowa Geological Survey, vol. 22, 1913, pp. 69-88; and by F. A. Bather in Geol. Magazine London, Dec., iv, vol. v, 1898, in 5 numbers, Nov., etc.)

2. IN SOLE AUTHORSHIP

1884. On the Occurrence of the Lower Burlington Limestone in New Mexico. (American Journal of Science, vol. xxvii, 1884, pp. 97-103.)
1899. Notice of a New Discovery Concerning *Uintacrinus*. (American Geologist, vol. xxiv, 1899, p. 92.)
1899. Slab of *Uintacrinus* Presented to the Museum of Comparative Zoology, Harvard. (Annual Report Mus. Comp. Zool., 1899, p. 5, with plate. Burlington *Hawkeye*, Nov. 12, 1903.)

1900. On the Presence of Pores in the Ventral Sac of Fistulate Crinoids. (American Geologist, vol. xxvi, Sept., 1900, pp. 133-151; plate xvi.)
1901. Associate in Paleontology, Museum of Comparative Zoology, Harvard College.
1901. *Uintacrinus*, Its Structure and Relations, 4to. (Memoirs Museum of Comparative Zoology, Harvard College, vol. xxv, No. 1, August, 1901, pp. 1-89; plates 1-8. Cambridge, Mass. Noted in London Zoological Record for 1901, p. 1: Under Crinoidea: "The work of the year is Springer's magnificent Monograph on the Structure and Relations of *Uintacrinus*.")
1901. *Uintacrinus* Slab Presented to American Museum of Natural History, New York, and U. S. National Museum. (American Museum Journal, vol. ii, No. 2, February, 1902, with 2 plates; Burlington *Hawkeye*, Nov. 12, 1903.)
1901. Specimens of *Uintacrinus* Presented to Universities of Iowa and Chicago, British Museum, and Museum of Berlin. (Burlington *Hawkeye*, Nov. 12, 1903.)
1902. On the Crinoid Genera *Sagenocrinus*, *Forbesiocrinus*, and Allied Forms. (American Geologist, vol. xxx, August, 1902, pp. 88-97; 1 figure.)
1902. Notice of a New Comatula from the Florida Reefs. (American Geologist, vol. xxx, August, 1902, p. 98.)
1903. *Actinometra iowensis*, a New Unstalked Crinoid from the Florida Reefs. (Bulletin Laboratories of Natural History, State University of Iowa, vol. v, No. 3, June, 1903, pp. 217-221; plate 1.)
1905. *Cleiocrinus*. 4to. (Mem. Mus. Comp. Zool., Harvard College, vol. xxv, No. 2, Jan., 1905, pp. 93-113; plate 1. Cambridge, Mass.)
1905. *Hypsocrinus*, a New Genus of Crinoids from the Devonian. By Frank Springer and Arthur Ware Slocum. (Field Columbian Museum, Publication No. 114, Geological Series, vol. 2, No. 9, 1905, pp. 267-274; plate lxxxi, Chicago.)
1906. Discovery of the Disk of *Onychocrinus*, and Further Remarks on the Crinoidea Flexibilia. (Journal of Geology, vol. xiv, No. 6, Sept.-Oct., 1906, pp. 467-522; plates 4-7. Chicago.)
1906. Foreign Correspondent Geological Society of London.
1909. Some Noteworthy Accessions to the Division of Invertebrate Paleontology in the National Museum. By R. S. Bassler. (Smithsonian Miscellaneous Collections, vol. 52, part 2, No. 1823, Jan. 30, 1909, pp. 267-269; plates xvii, xviii.) Concerning specimens presented by F. S.
1909. A New American Jurassic Crinoid. (Proc. U. S. National Museum, vol. xxxvi, March, 1909, pp. 179-190; plate 4. Washington, D. C.)
1909. On the Name *Encrinus*. (Circular letter to Zoologists and Paleontologists generally, enclosing card for answers, May 1, 1909.)
1910. Collection of Fossil Crinoids Presented to the State University of Iowa. (Resolution by State Board of Education, at Cedar Rapids, March 2, 1910.)
1910. Field Session of School of American Archaeology. (*Science*, Nov. 4, 1910, pp. 622-624; Santa Fé *New Mexican*, Sept. 24, 1910.)
1911. The Crinoid Fauna of the Knobstone Formation. (Proc. U. S. National Museum, vol. xli, June, 1911, pp. 175-208. Washington, D. C.)
1911. New American Fossil Crinoids. 4to. (Memoirs Museum of Comparative Zoology, Harvard College, vol. xxv, no. 3, July, 1911, pp. 117-161; plates 1 to 6. Cambridge, Mass.)
1911. Farewell to Crinoids. Removal of collection and library to the U. S. National Museum. (Burlington *Hawkeye*, Nov. 12, 1911.)
1911. Great Library Comes Here. The Nicolaus Finck linguistic library presented to the School of American Archaeology. (Santa Fé *New Mexican*, May 25, 1911.)

1911. On a Trenton Echinoderm Fauna at Kirkfield, Ontario. (Mem. Geological Survey of Canada, No. 15-P, 1911, pp. 1-50; plates 1-5. Ottawa, Canada.)
1913. Notes on an Unusually Fine Slab of Fossil Crinoids. By R. S. Bassler. (Proc. U. S. National Museum, vol. 46, Nov. 29, 1913, pp. 57-59; plates 1-2.) Presented by F. S.
1913. Exploration and Field-Work of the Smithsonian Institution in 1913. (Smithsonian Miscellaneous Collections, vol. 63, no. 8, pp. 14-16, with plate.) Expedition supported by F. S.
1913. Chapters on Blastoids and Crinoids in Zittel-Eastman's Text-book of Paleontology, 2d edition, revised, 1913, pp. 161-243.
1914. Associate in Paleontology, United States National Museum.
1915. Review of Austin H. Clark's Monograph of the Existing Crinoids. (*Science* N. S., vol. 42, no. 1080, Sept. 10, 1915, pp. 342-345.)
1917. *Scyphocrinus* and Its Bulbous Root, *Camarocrinus*. 4to. (Smithsonian Institution, publication no. 2440, 1917, pp. 1-74, plates 1-9, text-figs. 1-19. Washington, D. C. Noted in *Santa Fé New Mexican*, Aug. 21, 1917.)
1918. Iowa Authors and Their Works. (State Historical Society of Iowa, 1918. Frank Springer, pp. 279-280; Wachsmuth and Springer, pp. 302-303.)
1918. On *Mysticocrinus*, a New Silurian Genus of Fossil Crinoids. (*American Journal of Science*, vol. xlvii, Nov., 1918, pp. 666-668, plate 2.)
1918. A New Species of Fossil *Pentacrinus* from the East Indies. (Nederlandische Timor-Expeditie, II, Jaarboek van het Mijnwesen, 45te Jaargang, 1916, pub. 1918, pp. 1-8; plate 1. Leiden, Holland.)
1919. A New Species of Crinoid, *Eucheirocrinus ontario*. (Geological Survey of Canada, Memoir III, No. 91, 1919, p. 127, figure 6.)
1920. Die Botryocriniden. E. Haarmann. Communication by F. S. (Jahrb. Preuss. geol. Land. Anst., Bd. xli, Teil 1, Berlin, 1920, 87 pp. Taf. 1-6.)
1920. Monograph of the Crinoidea Flexibilia, 4to, 2 vols. (Smithsonian Institution, Publication No. 2501, two vols., 486 pp., text, 51 text-figures, and 79 plates, 1920. Washington, D. C. Reviewed in *Geological Magazine*, London, vol. lix, April, 1922, pp. 177-178; *American Journal of Science*, April, 1921, pp. 369-370; *El Palacio*, Santa Fé, N. M., June, 1921; New York State Museum, 1921, pp. 1-43; Burlington *Hawkeye*, Oct. 23, 1921.)
1921. Degree of Doctor of Science by George Washington University. Washington, D. C., February 25, 1921.
1921. The Fossil Crinoid Genus *Dolatocrinus* and Its Allies. (United States National Museum, Bulletin 115, 1921, pp. 1-59; plates 1-16. Reviewed *American Journal of Science*, vol. 2, 1921, p. 57.)
1921. New Species of Devonian Crinoidea from Northern Canada. (Geological Survey of Canada, Bulletin No. 33, 1921, pp. 15-17; plate 1.)
1921. The Existing Crinoids. By Austin H. Clark. (Vol. 1, part 2, 4to. Bull. U. S. Nat. Mus. No. 82, 1921, p. 1.) Assistance to author.
1922. Crinoids from the Upper Cretaceous of Tamaulipas, Mexico. (Proc. U. S. National Museum, vol. 61, art. 5, 1922, pp. 1-4, plate 1.)
1923. The St. John Collection of Fossil Fishes Presented to the United States National Museum. (Report of the U. S. National Museum, June 30, 1923, pp. 84-85. Washington.)
1923. On the Fossil Crinoid Family Catillocrinidae. (Smithsonian Miscellaneous Collections, vol. 76, no. 3, Publication 2718, Aug. 3, 1923, pp. 1-41; plates 1-5. Washington, D. C.)

1923. A Flexible Crinoid from the Carboniferous Limestone of Fife, Scotland. By James Wright. Communication by F. S. (Geological Magazine, London, Nov., 1923, pp. 481-490.)
1924. Degree of Doctor of Philosophy Conferred by the Rheinische-Friedrich-Wilhelms University at Bonn, Germany, January 16, 1924, "upon Mr. Frank Springer, Las Vegas, New Mexico, as eminently successful investigator of Fossil Crinoidea; author of a reliable Treatise on the Palaeocrinoidea; disinterested promoter of Scientific Research; especially in recognition of the fact that all his scientific accomplishments have been gained outside of his main calling in life."
1924. A Tertiary Crinoid from the West Indies. (Proc. U. S. National Museum, vol. 65, art. 3, 1924, pp. 1-8; plate 1.)
1924. On the Genus *Holopus*, with description of a hitherto unrecorded specimen of *H. rangii* from Barbados. (State University of Iowa, Studies in Natural History, Aug., 1924; pp. 45-63; plates 1-3.)
1924. Die Permischen Krinoiden von Timor. By Johannes Wanner. (Teil II, 1894. The Hague, 348 pp., 22 plates. Introduction, p. 3.) Assistance to the author.
1924. A Remarkable Fossil Echinoderm Fauna in the East Indies. Review of Wanner's monograph on Timor Crinoids. (American Journal of Science, 8, Oct., 1924, pp. 325-335.)
1924. The Devonian Crinoids of New York, by Winifred Goldring. Review of monograph in Science, Oct. 24, 1924.
1925. The Genus *Pentacrinus* in Alaska. (Proc. U. S. National Museum, vol. 67, art. 5, 1924, pp. 1-7; plate 1.)
1925. Occurrence of the Crinoid Genus *Apiocrinus* in America. (Proc. U. S. National Museum, vol. 67, art. 18, 1925, pp. 1-5; plate 1.)
1926. Unusual Forms of Fossil Crinoids. (Proc. U. S. National Museum, vol. 67, art. 9, 1926, pp. 1-137; plates 1-26.)
1926. Upper Devonian Crinoids from Mackenzie River Valley. (Geological Survey of Canada, Bulletin 42, May, 1926, pp. 127-132, plate 24.)
1926. New Species of Crinoids from Anticosti. (Geological Survey of Canada. Awaiting publication.)
1926. American Silurian Crinoids. (Smithsonian Institution Publication no. 2871, 4to, pp. 1-239; plates 1-33.)

B. MISCELLANEOUS

1867. Science as a Civilizer. Graduating Oration, State University of Iowa, June 18, 1867.
1870. Oration at Second Commencement Meeting of the Alumni Association of the State University of Iowa, June, 1870. Noted by Hon. John P. Irish in the *Iowa State Press* of June 20, 1871: "The second meeting was last year, on which occasion Frank Springer, Esq., delivered one of the most remarkable speeches ever heard in Iowa. Remarkable for the reason that it sounded the slogan, and raised the standard of advance thought, and marked out the path which opinion will take when the alumni of our free western universities form a considerable part of the educated community."
1871. The Right of Rational Inquiry. Lecture for the benefit of the Burlington, Iowa, Public Library, at Burlington, March 9, 1872. (Published in *Die Iowa Tribune*, March 12, 1872.)
1880. Member of Territorial Senate, New Mexico. (Twitchell's Leading Facts of New Mexican History, vol. ii, p. 453.)
1884. Dedication of Monument to William R. Morley, at Las Vegas, New Mexico, November 9, 1884. (Published in the *Las Vegas Optic*, Nov. 10, 1884.)

1885. Quarter-Centennial of the State University of Iowa. Address as representative of the alumni, at Iowa City, June 24, 1885. (Published in *Iowa City Republican*, June 25, 1885.)
1887. Argument before the Supreme Court of the United States in the Maxwell Land Grant Case, at Washington, March 10, 1887. (Vol. 121, U. S. Supreme Court Reports, p. 377; 122 U. S., p. 375. Published in Twitchell's Spanish Archives of New Mexico, vol. i, pp. 53-60; Twitchell's Leading Facts of New Mexican History, vol. ii, p. 453; Geo. Wharton James, New Mexico, p. xix.)
1888. Trust Deed of Maxwell Land Grant Company, prepared and carried into execution, at Amsterdam, Holland, January 19, 1888, under which the property was administered for 38 years.
1889. Member of Constitutional Convention, New Mexico. (*Santa Fé New Mexican*, Oct. 10, 1889.)
1890. Land Titles in New Mexico. Address as President of New Mexico Bar Association, at Santa Fé, January 7, 1890. (Published in *Santa Fé daily New Mexican*, January 8, 1890; minutes of New Mexico Bar Association for 1890, and in pamphlet edition for distribution; Twitchell's Leading Facts of New Mexican History, vol. ii, pp. 453, 462.)
1890. President Board of Trustees Maxwell Land Grant Company. (Leading Facts of New Mexican History, vol. ii, p. 453.)
1890. Argument before District Court of New Mexico in *Clark vs. Maxwell Land Grant Company*, at Las Vegas, N. M., December, 1890. (Reported in *Las Vegas daily Optic*, June 2, 1891.)
1891. Argument before the United States Supreme Court in the Beales Grant case, March 11, 1891. (Vol. 139, U. S. Supreme Court Reports, p. 569; *Santa Fé daily New Mexican*, April 28, 1891.)
1896. Remarks at Funeral of Charles Wachsmuth, at Burlington, Iowa, February 9, 1896. (Published in *Burlington Hawkeye*, Feb. 10, 1896; *Annals of Iowa*, Des Moines, 3d series, vol. ii, 1896, pp. 354-357.)
1896. An Epoch in American Science. By C. R. Keyes. (*Annals of Iowa*, 3d series, vol. ii, 1896, pp. 345-364.)
1897. Argument before the United States Supreme Court in the Bent Suit, Washington, November 8, 1897. (Vol. 168, U. S. Supreme Court Reports, p. 451; *Las Vegas, New Mexico, daily Optic*, Nov. 9 and Dec. 7, 1897; see *daily New Mexican*, *Santa Fé*, Oct. 3, 1895.)
1899. The Flag and the Nation. Speech at the Rough Rider's banquet, at Santa Fé, New Mexico, February 10, 1899. (Published in the *daily New Mexican*, Feb. 11, 1899.)
1899. Dedication Address, New Mexico Normal University, at Las Vegas, March 4, 1899. (Published *Las Vegas daily Optic* and *Santa Fé daily New Mexican*, March 6, 1899; *Southwest* magazine, March, 1899, pp. 84-89.)
1899. Medal for Governor Roosevelt. Presentation on behalf of people of New Mexico at the First Rough Riders' reunion, at Las Vegas, June 25, 1899. (Published in *Las Vegas daily Optic*, June 26, 1899; *New York Sun*, June 26, 1899, and the daily press generally.)
1900. The Bar and the People. Speech at banquet of New Mexico Bar Association, at Santa Fé, January 3, 1900. (Published in the *daily New Mexican*, June 4, 1900.)
1900. University of New Mexico. Commencement address, at Albuquerque, June 7, 1900. (Published in *Albuquerque Morning Journal*, June 8, 1900.)

1901. Marshall Day Oration. On the centenary of the appointment of John Marshall as Chief Justice of the United States Supreme Court. By invitation of the Supreme Court of New Mexico; before the Supreme Court, the two Houses of the Legislature, and the New Mexico Bar Association, at Santa Fé, Feb. 4, 1901. (Published in the daily *New Mexican*, Feb. 5, 1901; and in the Minutes of the New Mexico Bar Association for 1901, pp. 15-25.)
1901. Member of Territorial Senate, New Mexico. (Twitchell's Leading Facts of New Mexican History, vol. ii, p. 453.)
1901. On the Olympic Games. Address on presentation of medal for athletics, New Mexico Normal University at Las Vegas, June 21, 1901. (Published in Las Vegas daily *Optic*, June 22, 1901.)
1901. Eulogy on William McKinley. Before the literary societies of the New Mexico Normal University, at Las Vegas, October 4, 1901. (Published in the Las Vegas daily *Optic*, October 5, 1901, and in pamphlet edition for distribution.)
1902. Colorado State Normal School. Commencement address, at Greeley, Colo., June 5, 1902. (See Las Vegas daily *Optic*, May 25, 1903.)
1902. A Prize for Oratory. Presentation of a medal for oratory, New Mexico Normal University, at Las Vegas, June 20, 1902. (Published in the daily *Optic*, June 21, 1902.)
1902. Portrait of Rev. William Salter. Address on presentation to State Historical Department of Iowa, at Des Moines, November 24, 1902; by invitation of citizens of Burlington. (Published in Burlington *Hawkeye*, Nov. 26, 1902, and in Des Moines *State Register and Capital*.)
1903. New Mexico Normal University. Commencement address, May 28, 1903. (Abstract in Las Vegas daily *Optic*, May 29, 1903; also May 25, 1903.)
1903. Portrait for State, Presented to Historical Department of Iowa on Request of Governor. (Robert J. Burdette in Burlington *Hawkeye*, November 12, and November 15, 1903; *Annals of Iowa*, vol. vi, 1903, p. 390.)
On complying F. S. wrote: "If I have done anything worth commemorating, it is to show that a man can, by husbanding his time, follow a reasonably successful business life, and at the same time give some thought to the wealth of nature that is around him."
1904. Burlington High School. Commencement address, June, 1904. (Burlington *Hawkeye*, June 8, 1904.)
1908. Biographical Sketch. Leading Facts of New Mexican History, Twitchell, p. 453.
1910. The Unknown Soldier. Remarks at the annual encampment of the New Mexico National Guard Regiment on Las Vegas Field, August, 1910. (Published in Las Vegas daily *Optic*, Nov. 18, 1920.)
1911. Opening of Republican Campaign for First State Election in New Mexico. Speech by request of Republican State Central Committee, at Las Vegas, Oct. 7, 1911. (Published in Santa Fé daily *New Mexican*, Las Vegas daily *Optic*, Albuquerque *Evening Herald*, and 20,000 copies in pamphlet distributed throughout the state.)
1914. A Man with a Vision. (Albuquerque *Morning Journal*, July 12, 1914.)
1915. A New Glory for New Mexico; New Museum at Santa Fé. (Magazine *Old Santa Fé*, vol. ii, no. 4, April, 1915, pp. 438-440.)
1916. Memorable Day in Annals of Santa Fé. (Daily *New Mexican*, April 12, 1916; *El Palacio*, April, 1916; Scribner's Magazine, March, 1923, p. 381.) Presentation of fund ensuring construction of new museum.
1916. Important Progress in Archaeology. Investigations of Frank Springer and party. (Albuquerque *Morning Journal*, Sept. 12, 1916.)
1916. The Secret of Keeping Young. (Albuquerque *Morning Journal*, Oct. 8, 1916.)

1917. When Dreams Come True. Dedication of new museum, at Santa Fé, November 25, 1917. (Published in daily *New Mexican*, Nov. 26, 1917; Las Vegas daily *Optic*, Nov. 26; *El Palacio*, Santa Fé, vol. iv, no. 4, for Nov., 1917, pp. 1-18; 90-97; *Art and Archaeology*, Washington, D. C., Jan., 1918, pp. 3-7; Geo. Wharton James, New Mexico, the Land of the Delight Makers, 1920, pp. 428-443, and pamphlet edition of 1,000 copies by the School of American Research.)
1918. A Scientist's Opinion. Dr. F. A. Bather on the New Mexico State Museum. (Museums Journal, London, March, 1918.)
1918. The St. Francis Murals; given by F. S. to New Mexico State Museum. (Art and Archaeology, Jan., 1918, p. 84.)
1918. The Paintings of Donald Beauregard; given by F. S. to New Mexico Museum. (Art and Archaeology, Jan., 1918, p. 79.)
1919. Theodore Roosevelt Memorial Service. Oration by invitation of the Legislature of the State of New Mexico, at Santa Fé, Feb. 9, 1919. (Published in daily *New Mexican*, Feb. 10, 1919; Las Vegas daily *Optic*, Feb. 10; Albuquerque *Morning Journal*, Feb. 10; two editions in pamphlet of 1,000 copies each for distribution in the State, and a third of 10,000 copies by order of the National Roosevelt Memorial Association, New York, for general distribution.)
1920. In Memoriam. Henry Strong Springer. Feb. 12, 1920.
1922. Bust by Scarpitta Presented to the State of New Mexico, at the State Museum, Santa Fé, Sept. 8, 1922. (Daily *New Mexican*, Sept. 9, 1922; *El Palacio*, Santa Fé, Oct. 2, 1922, pp. 75-88; *Pan American Geologist*, Des Moines, Iowa, Oct., 1922, pp. 177-186.)

HON. FRANK SPRINGER

(Bronze Bust by C. Scarpitta, presented to the State of New Mexico
by Friends of Mr. Springer)

PRESENTATION OF THE SCARPITTA BRONZE BUST
OF
HON. FRANK SPRINGER
TO THE
STATE OF NEW MEXICO³

Significant and memorable was the presentation of the bronze bust of Mr. Frank Springer to the state of New Mexico on the evening of September 8, 1922. The spirit of the celebration, its setting and its import, made it an historic event in the annals of a state overwhelmingly rich in traditions, thrilling events and picturesque episodes. To the hundreds of people who had gathered in the Saint Francis Auditorium to do honor to the man who had wrought so much for them and their commonwealth, and had achieved distinction and success in law, in business, in science, in statesmanship and in wise giving, it was an unforgettable occasion.

The glamour and the shouting of the 1922 Fiesta had filled the fore part of the week, and were followed by the earnestness and interest of the sessions of the Southwestern Division of the American Association for the Advancement of Science. Friday evening had been set apart by the Association for the exercises which proved to be the climax of a week crowded with stirring and notable incidents. It had been a typically sunny Santa Fé day, followed by a night rapturous with a brilliant harvest moon. The members of the Science Association, just returned from a field excursion to San Ildefonso, where the Tewas had delighted them with a tablita ceremony of wonderful color and grace—joined with the friends of Mr. Springer from far and near in filling comfortably the St. Francis Auditorium of the new museum. There were present former and present business associates from as far away as New York and St. Louis, from cities and towns in New Mexico and adjoining states; there were men and women in the audience who had won world-wide distinction in the fields of science, literature, art, music and the professions; there were represented the Indian, the Spanish American, the pioneers and the new comers to the Southwest—as cosmopolitan and fine an audience as ever gathered to do honor to a fellow man.

There were spontaneity and dignity about the program which transmitted themselves to the audience; a warmth of feeling and simplicity which characterize a neighborhood gathering rather than the formal state event; and yet, withal, the impressiveness of a solemn service in cathedral or abbey where all hearts are lifted to one purpose. Under the indirect, mellow glow of the high ceiling

³ Presented at the third annual session of the Southwestern Division of the American Association for the Advancement of Science at the St. Francis Auditorium of the New Museum, Santa Fé, N. M., Friday evening, September 8, 1922. Present account reprinted from *El Palacio*, Santa Fé, October 2, 1922.

lights of the nave the St. Francis murals took on new values, and the carved vigas with their primitive color designs stood out in massiveness from the shadows. The transepts and the chancel were flooded with light. To the fore on the altar platform, on an improvised pedestal, stood the veiled bronze bust, and to one side on a table the scientific volumes of which Mr. Springer is the author.

Dr. Edgar L. Hewett broke the hush of expectancy that fell upon the audience when the speakers of the evening took their places on the platform. It was obvious that the tribute he paid Mr. Springer was heartfelt. Like the speakers that followed, he commanded the closest attention of the audience. Closely associated with Mr. Springer for a quarter of a century in the upbuilding of such great institutions as the Normal University of New Mexico, the Museum of New Mexico, the School of American Research, and in the furthering of research work, the renaissance of native arts and handicrafts, the fostering of art and literature in the Southwest, it was quite fitting that Dr. Hewett emphasized Mr. Springer's contributions to science, education and art.

Col. Ralph E. Twitchell, historian, author, lawyer, eloquently and forcibly dwelt upon the high place Mr. Springer achieved as a member of the bar. He, too, had been close to Mr. Springer for decades, and prizes this association as the most precious in a life filled with activities for the advancement of his beloved state and city. He reviewed Mr. Springer's achievements with the sympathy of a friend, and yet with the preciseness of evaluation of the trained biographer and historian. Momentarily shaken with emotion as he paid an intimate tribute, he passed quickly to the orderly marshaling of dates, figures and historical facts, imposing in themselves and overwhelming in their aggregate.

In concluding his address Colonel Twitchell presented a handsomely bound volume entitled "Notes and Memoranda Concerning the Legal, Scientific and Literary Activities of Frank Springer as a Citizen of New Mexico," to serve as a permanent record in the library of the State Museum. It contains a sketch of Mr. Springer's career at the bar and his connection with public affairs, a history of the most notable cases in which he was engaged, together with comments by other lawyers and judges upon his arguments before the United States Supreme Court; a reprint of his oral argument in one of the cases; and contemporary newspaper comment upon these matters. Also reprints or original copies of many of his public addresses upon legal, educational and miscellaneous subjects. Also accounts and reviews of his scientific works and publications, taken from periodicals of high authority. This compilation exhibits from the original sources evidence of the wide range of Mr. Springer's intellectual activities. It is understood that a small number of copies of the work have been printed for reference in libraries, and for the use of his family and intimate friends.

Mr. James G. McNary, president of the First National Banks of El Paso, Texas, of Las Vegas and of Dawson, in New Mexico, who, too, had known Mr. Springer for a quarter of a century, spoke as a younger business man of a successful older man, emphasizing Mr. Springer's place as an empire builder, a man who touched the resources of northern New Mexico so that they yielded abundantly, creating irrigation works, developing mines, founding settlements, giving sustenance and bringing happiness to a great multitude of people.

Dr. D. T. MacDougal, general secretary of the American Association for the Advancement of Science and president of the Southwestern Division, who is in charge of the Desert Laboratory of the Carnegie Institution at Tucson, Arizona, spoke as one scientist of another, referring also with approval to the business man in science—in the case of Mr. Springer achieving doubly, by giving himself and giving of his means.

Then came the moment for unveiling. Dr. Hewett paid glowing tribute to his friend as he removed the cover from the bust which stood revealed as a splendid piece of art; he tendered it on the part of Mr. Springer's friends as a gift to the state. United States Senator A. A. Jones, delegated by Governor Merritt C. Mechem to receive the bronze on behalf of the state, was in a happy vein, his address of acceptance being felicitous in its warmth of feeling and eloquence.

At the conclusion of the addresses Homer Grunn, of Los Angeles, composer and pianist, rendered with exquisite feeling three of his themes, that stand among the most pleasing and beautiful modern compositions based on Pueblo melody and rhythm.

Following Mr. Grunn's numbers, Mr. Springer, who had not been present during the exercises, was escorted into the auditorium by Mrs. Laughlin and Mrs. Hewett, representing the Woman's Museum Board, and was given a warm and generous ovation, the audience rising and remaining standing while he responded briefly, first in a lighter vein, to mask his evident emotion, and then gravely, voicing his real feeling in a single touching sentence. He then presented to the audience Mr. Scarpitta, the sculptor, whose presence had not before been generally known, who with unaffected earnestness told of the deep feeling with which he had executed the portrait now before them. The remarks of the two gentlemen were as follows:

Mr. Springer: *My friends:* I did not come here to make a speech, but I am here in obedience to the command of my superior authorities, representatives of the women of Santa Fé, to whose example and encouragement is due a large part of what the men have been able to accomplish for the betterment of this community. When I was summoned to appear at this stage of the proceedings, I knew, of course, that the summons came from the hearts of the friends whose infinite kindness and courtesy have been so often manifested, the recollection of which makes this beautiful place seem even more beautiful. But I thought that it might also be intended to give to those in attendance here an opportunity to judge of the merit of the

work which they have been invited to view by a direct comparison with the original. On that point I think there can be but one opinion. Looking at such an example of the sculptor's art, one is inclined to marvel at his power to create out of inanimate materials so faithful a delineation of the human features; but come to think of it, it isn't so very wonderful after all; it is really quite simple; all the sculptor has to do is to procure a supply of the proper kind of clay, and then put it in the right place. This disclosure of the mystery is not original with me, but probably has done duty ever since studios were invented. I cribbed it from a painter, who told an inquisitive visitor seeking to learn the secret of his canvasses that it was easy, all that was necessary being two things: first, to select the proper colors; that was very important, he said; and then—put them in the right place. So it is with one of Fritz Kreisler's masterpieces: simple enough, just pressing the proper string, and drawing the bow the right way.

You know, it is against the rules for the subject who is being honored as I am to attempt to say anything serious; but I am wondering if in all this nonsense I may not inadvertently have stumbled upon a great truth, which is, that one of the chief problems of life is to learn how to select the proper things, and to use them in the right way.

Now, having broken the rule once, I am going to take another whack at it, and sum up what I really think and feel, by saying that the thing which is culminating tonight has touched me deeply; and that I would rather have it occur here, in this place, than anywhere else on earth.

One other privilege belongs to me, of which I avail myself with the most profound satisfaction; and that is, to present to this audience the man whose inspired art has created the work which you have come here to see—Scarpitta, the sculptor.

Mr. Scarpitta: *Mr. Springer, Dr. Hewett, Ladies and Gentlemen:* I would like to say a word about this work of mine. I am not a speaker, as you know, but I cannot go through this without first of all telling you how profoundly impressed I am that I was the one selected to make this portrait of Mr. Springer. The moment I came in contact with Mr. Springer I felt immediately the great force with which he has accomplished all that you have heard related by the Chairman and the other gentlemen. I went into this work with all my love, because I felt the sincerity of this man; and I have, I hope you will agree with me, put into this bronze something that will live on forever, and reveal the inner feelings of Mr. Springer, his intense love for humanity, and his power of convincing any one of his profound sincerity, which I have admired so, and of which I am so proud to know and see that you all agree with me. I thank you.

The program was then concluded by the gracious and beautiful Princess Tsianina in an exalted mood, who with the sympathetic accompaniment of Mrs. Doll, gave some of the finest Indian songs in her repertoire, among them the Zuni Invocation to the Sun-god, which, as she explained, she had substituted for another number because it better expressed her feeling after listening to the addresses which had preceded.

In the patio of the art museum the Woman's Museum Board had spread a table made colorful with magnificent dahlia blossoms. Moonlight flooded the scene, and garlands of colored lights swung across the sward. Here thronged the people after pouring past Mr. Springer, offering their words of appreciation for all he has done and all he has stood for, and lingered until midnight and after. The literary folk were present under the leadership of Mary Austin, who but recently returned from England; the anthropologists included Francis La Flesche and Miss Alice C. Fletcher, of Washington, D. C.; other branches

of science were represented by Dr. MacDougal, of the Desert Laboratory; Dr. Larkin, of the Mount Lowe Observatory; Dr. Douglass, of the Lowell Observatory, and many other members of the American Association for the Advancement of Science, most of them engaged in research work in the Southwest; the bar by Hon. Thomas B. Harlan, of St. Louis, and many others of the federal and state courts and bars; business by President Jan Van Houten, of the St. Louis, Rocky Mountain & Pacific Co.; President Levi Hughes, of the First National Bank of Santa Fé; President James G. McNary, of the First National Bank of El Paso; President George Ulrich, of the Exchange Bank of Carrizozo, and scores of others active and prominent in the upbuilding of the Southwest; statesmanship by United States Senator A. A. Jones and other federal and state officials; while the presence of members of the art colonies in Santa Fé and Taos, of leaders in music like Grunn and Tsianina, as well as those active in social circles, made the assembly a brilliant one indeed.

The art alcoves were thrown open. The Fiesta exhibit of paintings by Santa Fé and Taos artists made them unusually attractive. In one alcove the Springer bust was placed together with a Madonna group also by Scarpitta, a lovely piece of work. In the opinion of critics without exception the Springer bronze is a veritable masterpiece. It is more than a marvelous physical likeness, for it emphasizes the earnestness and power of concentration of the subject, at the same time giving an intimation of his spirituality and idealism. The bronze is a gift of friends of Mr. Springer, who contributed through a committee consisting of Paul A. F. Walter, chairman and treasurer, with Dr. Edgar L. Hewett in charge of contributions from artists, scientists and educators, Mr. Jan Van Houten of contributions from business associates of Mr. Springer, and Col. Ralph E. Twitchell, of contributions from members of the bar. The bust will have a permanent place in the art museum, whose creation is due primarily to the vision and generosity of Mr. Springer.

Herewith follows the text of the addresses delivered upon this memorable occasion, reflecting as they do the spirit of the celebration:

INTRODUCTORY

Dr. Hewett: *Friends:* We have met here tonight to celebrate the life and achievements of a man of science. The Southwestern Division of the American Association for the Advancement of Science joins on this occasion with the people of New Mexico in honoring a scientific man of the Southwest, Mr. Frank Springer. You, gentlemen of the bar and of the courts, may be surprised when I claim Mr. Springer as a man of science, for he has been one of your most distinguished associates for many years; and you men of business of the Southwest may also be somewhat startled, for he has long lived among you as an exceedingly busy man of affairs. The fact is he belongs equally with all of us.

To support my reference to Mr. Springer as a man of science, I ask you to inspect this shelf of works that would be a credit to one who had devoted his entire life to scientific pursuits. The majority of you have probably been ignorant of the fact that this silent, modest next door neighbor of yours has been steadily publishing scientific contributions that have made him known throughout the world.

So it is a many-sided man that we are honoring tonight. We are celebrating his achievements while he is still with us—achievements in science, in law, in business and in public affairs. It seems fitting that one of his old neighbors, who has known intimately his legal activities almost from the beginning, should speak of Mr. Springer as a lawyer and of his career at the bar. Col. R. E. Twitchell has been invited to perform that service. He can speak with the authority of the historian.

THE LAWYER AND CITIZEN

Colonel Twitchell: *Dr. Hewett, Ladies and Gentlemen:* I consider my having been selected for this great privilege a most notable distinction. To be associated in this way in these exercises is an honor which seldom comes to a member of our profession.

I first met Mr. Springer forty years ago, and there are only nine members of the bar today living who knew Mr. Springer during all that period.

Early in the '90s, I was twice honored with the presidency of the New Mexico Bar Association, and soon realized that the constructive power in the then "Territory of New Mexico," the history makers in the development of a great commonwealth, remained largely with the legal profession, and became certain that there was no one among the members of the bar in the entire territory who would feel like giving his time to the work of recording, biographically and otherwise, the efforts and achievements of that constructive element; and so I began a series of papers for the Bar Association, dealing with the history of the bench and bar during the preceding years of American occupation. That early effort, more than thirty years ago, resulted finally in an amplification and concrete compilation which was published a few years since.

In making a historical outline for these papers it became apparent that a certain classification was necessary for the reader to properly understand the position which, in certain decades, had been filled by the members of the bench and bar. That classification was as to time, and it may be repeated here: Those who came with the American Army of Occupation in 1846; those who came with Carleton's column from California during the Civil War period; and those who came shortly before and at the time of the coming of the railways into the southwest. Mr. Springer belongs in the last named classification, having come to New Mexico in 1873, anticipating the coming of the railways and the creation in a few years of a great state in the American Union.

Had it not been for the panic of 1873, and the failure of the great banking firm of Jay Cooke & Co., of Philadelphia, the geography and the history of the states of Colorado and New Mexico would not be such as we find it today. Had the efforts which were being made at that time by Mr. Springer and his eastern associates borne fruit, the city of Pueblo with all of its industries, all of its environment, in all probability would be located where the town of Cimarron, New Mexico, was then situated. The entire proposition of financing of the transcontinental railway into the southwest at that time had been consummated, but the failure of Jay Cooke & Co. in the panic of 1873 changed the entire status of affairs, and it was

not until several years later, due to the activities of certain people in Pueblo, Messrs. Thatcher and Reynolds and others, that a line of railway was built down the Arkansas to La Junta, where it joined the Atchison, Topeka & Santa Fé Railroad in its construction into the Southwest.

Mr. Springer was the retained counsel of the Santa Fé Railway during that period; and during all of the years of his active career at the bar, either as trial lawyer or as counsel, was consulted in every case of any consequence which was heard in the courts of New Mexico, as well as in many other jurisdictions where the interests of that great company were affected. These are matters of which I have personal cognizance.

My acquaintance with Mr. Springer began in the spring of 1883. In the summer of that year there was celebrated here in Santa Fé the Tertio-Millennial Anniversary of Spanish conquest and occupation of all the areas between the Pacific and the Mississippi.

When Mr. Springer came to New Mexico the bar was composed of many distinguished lawyers. The names of these eminent jurists are found in the published reports of our Supreme Court. Within two years after his coming Mr. Springer took the position and was recognized as one of the leading lawyers at the bar, and during his active career retained that position of leadership. And from my personal acquaintance with the volume and character of the litigation confided to his charge, the manner in which it was handled, in results obtained and in the satisfaction always accorded to his clients, he enjoyed a position pre-eminent among his fellows at the bar. This became an accepted fact when the final determination of the most important suit, involving title to great tracts of land, was had in the court of last resort in this country. I refer to the Maxwell Land Grant case. Lawyers of the greatest distinction in America had been employed in the litigation involving the title to that great property, but it remained for Mr. Springer to secure a final conclusion and determination favorable to the contentions of the grant owners.

The Supreme Court of the United States was right in following the argument and contentions of Mr. Springer. Few there were of the Anglo-American population who, with the advent of the transcontinental railways, could appreciate or who would attempt to understand how any one individual or company or organization could possibly have the right of ownership in a million or more acres of the public domain. The real question at that time was the one which a few years before had been dominant in California—the question of the squatter and the Don, or the Don's descendants and grantees. Few there were who understood or comprehended the policy of the Mexican government in making these grants; a protective policy, brought about largely owing to the experience which that Republic had passed through in the case of Texas, and the settlement on Mexican soil of Americans from Tennessee, Mississippi and Louisiana, the final divorcement of the province from the Mexican Republic and its final entrance into the American Union. As a policy of protection, the Mexican Government gave great grants of land on its northern frontier, which was the Arkansas, to Mexican citizens. Those granted were the Maxwell, the Montoya, the Tierra Amarilla, and other grants of lands, empires in area and extent; granted to Mexican citizens for the purpose of establishing upon the American frontier a zone of Mexican influence which would resist the advance of the American, brought about through the commerce of the prairies and the establishment of the Santa Fé trail. These are the facts which caused the creation of those great land grants, and it was the student of history, as well as jurisprudence, as exemplified in Mr. Springer, who was thus enabled to sustain his contentions in our court of last resort. The determination of that court in that case not only revealed him as the leader of the bar in New Mexico, but gave him an international reputation: a reputation of a character which any lawyer, no matter where the scene of his activities, would appreciate as the acme of professional success. His reputation must rest upon his argument in that great case. The information which he obtained in its preparation, the knowledge that was his, and exclusively his, relative to questions arising under the Treaty of Guadalupe Hidalgo in the matter of Spanish Land Grants and their rela-

tionship to American occupation, was so comprehensive that when it came to the enactment of a law by the American Congress whereby all of these titles, in so far as any claim made by the Government of the United States was concerned, could be settled, his influence as a lawyer and citizen of New Mexico before the Senate and Congress of the United States was so pronounced that he was in position to render the people of the southwest a great service. His testimony and his arguments before the various Congressional and Senate committees had more to do with the passage of the act providing for the establishment of the Court of Private Land Claims than of all other New Mexicans combined.

As I have said, when I undertook the writing of the biographies of the members of the bench and bar, dealing with those classified as having come with the American occupation period, I found very little recorded in New Mexico beyond the decisions of our Supreme Court, and there were only two volumes of reports at that time. And so it was that I concluded that future historians and searchers would not be put to that inconvenience if any of my time could be profitably used in recording and accumulating information with reference to all of the members of the bar of my own period. Naturally there was much of interest in securing all possible information from so prominent a man as Mr. Springer. While my acquaintance with him was not intimate at that time but was of a professional nature, I found him then, as he is today, one of the most modest and reticent of men. It was almost impossible to acquire any information in ordinary conversation if he happened to know its purpose, and while living in Las Vegas, I took occasion to invite him to my residence, when unknown to him I was able to draw out the story of all the facts relative to the early history of the Maxwell Land Grant and earlier events occurring in Colfax County, every thing which had to do with the various lawyers and judges who practised in the courts of Taos, Colfax, San Miguel, and Mora counties, and had it very carefully recorded by a stenographer who was concealed in an adjoining room, Mr. Springer being at the time unconscious of the fact until years later, when I published the *Leading Facts of New Mexican History*, and later, in some notes to my *Spanish Archives*. Recognizing that he had played a most prominent part in the making of New Mexico, I began preserving all of the printed records, briefs, arguments, newspaper clippings and accounts of his activities, and so it is that upon this occasion I am able, in a small way, to contribute a compilation of all of the printed accounts which have been thus preserved. This volume I am now tendering as a small contribution on my part to this really great New Mexican; a contribution not for today but as a source book which the future historian will appreciate, who will amplify the work of those of us who have sought to record the facts of history in the southwest. This compilation, I believe, will be appreciated by the present and future generations of Mr. Springer's family, as well as by the historical writers of the future.

His courteous treatment of his fellow members of the bar, his consideration for the shortcomings and limitations of the younger members of the profession, his great respect for judicial authority and personnel, and his mastery of many unsolved legal problems which confronted the early practitioners in the courts of New Mexico, are outstanding features of Mr. Springer's career.

Dr. Hewett: Thanks to Colonel Twitchell for this sketch of Mr. Springer's activities in those great days of the making of our state. From among the men who have more recently come into positions of influence and usefulness I wish to present one who can speak of Mr. Springer's place in business and public affairs. I introduce to you Mr. James G. McNary, president of the First National Banks of El Paso and Las Vegas.

THE MAN OF AFFAIRS

Mr. McNary: *Mr. President, Ladies and Gentlemen:* I esteem it a high honor and a great pleasure to be invited to participate with you in doing honor to the name of Frank Springer.

A quarter of a century ago through the instrumentality of Mr. Springer, president of the Board of Regents, and Mr. E. L. Hewett, president of the faculty of the Normal University of Las Vegas, the fates directed my youthful footsteps to New Mexico, for which I have always had reason to be deeply grateful.

From the day, just 24 years ago, when I first met Mr. Springer, I have held it a high honor to call him my friend, and who does not love to pay tribute to a friend of such illustrious achievements that moderation ceases to be a virtue?

It would be impossible to overestimate the value of scientific knowledge. Each stray bit of such truth from the farther shore of human thought and inquiry, has its value in applying the phenomena of the physical world to the needs of our complex social life.

Of Mr. Springer's unusual fund of scientific knowledge and of his generous use of it, you have heard and are to hear from others better qualified than I to speak, but I wish to trespass on their ground long enough to read a few lines dedicated to Mr. Springer and handed me by a mutual friend. Emerson said: "Next to the man who originates a beautiful thought is the man who quotes it."

"In solitude he played his flute and thought,
Till finally this miracle was wrought,
The ordered working of his cultured brain,
Gave power to his gaze and through the train
Of aeons of dead years his piercing eye
Sought out Earth's secrets where they underlie
The cold faced rocks. Then slowly page by page,
He read through Nature's book and age by age
He found a story there. Today the world
Is deeply in his debt, for he revealed
To man the mystery the Earth concealed."

I am, however, delegated to speak especially of Mr. Springer's services as a builder of the state. How, I beg of you, can I do justice to this theme in ten short minutes?

When Mr. Springer first came to New Mexico, over a half century ago, her population was a scant one hundred thousand, her developed resources scarcely amounted to the value of the property now within the limits of this city. Great resources of coal lay hidden in her mountains, which required money and brains to develop and railroads to haul to the consumer.

Her streams carried a wealth of water for the thirsty land, which it required the hand of man to conserve, impound and distribute.

Mr. Springer arrived in New Mexico in the year 1873, and engaged in the practice of law at Cimarron.

One of his first public services was the founding of a newspaper and though I have never seen a copy, I can judge of the high spirit which must have animated its pages.

Early in his legal career Mr. Springer became identified with the interests of the Maxwell Land Grant Company, and in due time became its president, which position he still occupies.

He exerted an important influence on the industrial life of the state, as he was one of the first to recognize the importance of coal resources of New Mexico, and took a prominent part in the organization of the St. Louis, Rocky Mountain and Pacific company, the most important fuel producing company in the west, is today one of its directors and has always exerted an important influence in its affairs.

The greatest enterprise ever undertaken in the state of New Mexico by private capital, the building of the so-called Eagle Nest Dam in the Cimarron Canyon, was carried through by Mr. Springer and his brother Charles.

While built primarily for the purpose of storing water for their own lands, it will also irrigate a large acreage of adjacent lands and result in the development of one of the most important agricultural areas of the state.

The builders of this enterprise, in investing their capital, have taken all the chances, while the public and the state receives a large portion of the benefits.

Mr. Springer's name has been frequently mentioned for the highest political honors within the gift of the state, but his inclination has led him away from the field of politics.

His wise counsel, his cool and deliberate judgment, have always been sought by the leaders of the state, and his influence and counsel has been a most important factor in guiding the state through her territorial period into full statehood and in framing her laws and constitution.

Mr. Springer was one of the first to see in New Mexico the meaning of her past, the value of her present, and the promise of her future.

In addition to his own constructive work, his assistance and support have been of immeasurable value to those other pioneers who have labored to establish and build up the educational and scientific institutions of the state.

Dr. Hewett: We have heard of Mr. Springer's activities in law and business. We may now hear something said of his achievements in science. I have the pleasure to introduce the distinguished President of the Southwestern Division of the American Association for the Advancement of Science, Dr. D. T. MacDougal, of the Carnegie Institute, head of the Desert Botanical Laboratories at Tucson and Carmel.

THE BUSINESS MAN IN SCIENCE

Dr. MacDougal: *Ladies and Gentlemen:* The honor which comes to me is in no sense personal. I feel that I am here representing the 12,000 members of the American Association for the Advancement of Science—an association made up of men whose biographies would be the history of science of the world for the last seventy-five years. I feel especially happy on this occasion because the subject of our felicitations represents the very ideal and essence of this association which has given so much to the world. I have no compromises to make. I can speak frankly, and it is not often that a speaker on these occasions can go without limits.

In order to put to you how a scientist would feel about this matter, I must go aside for the moment and consider the state of society in which we live. In a democratic form of government our educational, scientific and artistic work must take the form of generalized effort which mostly has no high peaks. To illustrate what I mean by that: an organization may employ a great artist to make pictures; may give him so much for the month or day; and in spite of the fact that they are in harness, our musicians, artists and poets do great things. But the great creative work of the world is done on individual initiative. Quite regardless of what may or may not happen in Russia or in any other experiment, rational or crazy, whose nature is always such that the group of people whom James Harvey Robinson has designated as "the wonderers"—and in that group he includes the scientists, the poets and the artists—the creative work of these people will be done not under direction but as a matter of personal effort—driven from within and not from without.

I am not here to decry democratic government, whatever its failings and failures may have been. But I do wish to emphasize some of the finer graces that may arise on the basis and on the substratum of sound democracy. The most precious thing that can come from such a substratum is a sense of service, a sense of indirect accomplishment. The artist or the scientist who gives himself to his own longings, who pleases his own curiosities, who does his work not because he is told to—these are the men who achieve.

Now it is quite obvious who have found the course of life during the first quarter of this century. I think this will be a notable quarter century in its accomplishments and its tragedies, but it will be most noticeable for the fact that we can discern a growing sense of service in men who accept our democratic limitations, who accept the firm knowledge that they live, and then seek by personal service to top it with something that neither democracy nor any other form of government can evolve. This personal service has taken two forms. I have seen the men who are eminent in great industrial enterprises devote the profits and fruits of their work to the promotion of the dreams of artists, scientists and poets and others of creative capacity. I have also seen those men who have shown great ability in industrial and other efforts give not only the money which they cannot use to the promotion of these things, but give themselves.

It seems to me that the subject of our felicitations this evening is a man who has become notable to you, to your state, to our country, and to the scientific world by giving service in both these ways; by giving substantial support for the promotion of the dreams of others, and by scientific effort in the way of research, the fruits of which would be ample for the achievement of a great scientist if he did nothing else during his life time.

THE FRIEND OF SCIENCE, ART AND EDUCATION

Dr. Hewett: I am glad that we have had this summing up of the activities of Mr. Springer, for few have ever known him in all these capacities. Some have known him in law, some in business, some in the fields of science, some in the development of our educational, scientific and art work here, but few have realized his eminence in all these lines.

But after all has been appraised, something greater still remains to be told. There are those, and the number is large, who desire to pay a proper tribute to him in another capacity, that of the staunch inspiring friend. No one has greater reason to speak of him in this vein than I.

Our old friend, Charles F. Lummis, on the occasion of one of our encampments in the Rito de los Frijoles, drew this portrait, than which I have seen nothing finer in color or bronze, and which you will all recognize as a perfect likeness:

"Grave and gentle and strong and still,
Sits the Chief in the Council Tent;
But when we come to a breakneck hill
His is the hand that is lent.
There's a Something we all can feel—
Power and poise of the Elder stamp;
Solomon must have made a deal
With Springer, Dean of the Rito camp."

No one will ever know how many have felt the support of his strong, kindly hand, always extended in such an unostentatious way that it was known only to the one receiving it. If we could assemble here the young artists, musicians, scientists, who have been helped to their higher opportunities by him, the number would astonish you. And a greater effect than the material aid afforded was always the determined effort inspired by his faith. It called out the best in you. He has always absolute faith in the visions of youth. He believed in you, trusted you implicitly; you simply had to make good.

When I was at the head of the Normal University of New Mexico and he was president of the Board of Regents, he gave me this admonition: "Mr. Hewett, see that no deserving student ever leaves this institution for lack of funds. Always let me know about such cases." I am venturing to tell some things about Mr. Springer that have not been told in public before. These were the things that gave him his greatest pleasure outside of his family life.

His home was a sacred precinct that some of us have been privileged to enter. Among the greatest of his successes, shared by a wise and beautiful wife and mother, has been the rearing of a remarkable family. Two things stand out in my mind as I recall what I have seen of his family life—the solicitous care for the opportunities of every son and daughter, and the veneration in which he always held his forebears. He is the son of a great father, whose memory he holds in profound reverence. He is one of those who believe that

"Whatever is Man in the sons of men,
Whatever is staunch and true,
We draw from our sires, and their sires again,
And mothers of mothers who mated when
The world and its heart were new."

We could speak long of the qualities that have endeared Mr. Springer to so many. He is all that has been claimed for him tonight, and something greater still. In the mind of every one of you who knows him well, he stands out as the incomparable gentleman. He has reached that highest of all distinctions through his interest in and practice of the finer things of life: The things in which he has taken deepest interest, science, art, music, education, public welfare, the good name of his state, the destiny of our country, which he has always staunchly held must be achieved on the lines laid down by the founders of our great republic, make for nobility. It is the highest possible ideal of the citizen—nobility in public and private life.

The friends of Mr. Springer have felt that at this time, when he has reached the fullness of life and is still with us, they would like to do something that would stand as an enduring contribution from them to this state in honor of its great citizen. In the olden times, men honored not only their soldiers and statesmen, but their scholars and all who contributed to the greatness of the state. It seemed to the friends of Mr. Springer that it would be a worthy thing to present some imperishable gift that would stand forever as a tribute to him, in this temple of art which he helped so much to create, in this institution which he has from its beginning enriched with his broad spirit and wise counsel, in this ancient capital which has felt in so many ways the beneficence of his presence, and in this great state of New Mexico, which he has helped to ennoble by his active, creative life, and which he has always cherished and acknowledged his obligations to as the place of his early opportunities.

On mentioning it to one after another, every one without exception said: "Let me have a part in that. I want to contribute to that purpose." So to a sculptor who had gained high eminence in our country, Cartaino Scarpitta, was entrusted this sacred task. He brought to it not only the skill of a master, but the spirit of a master—the discernment and zeal which produce results in art that live as master work. I am now going to show you the outcome of this plan that we have brought to a happy conclusion. When I told Mr. Springer what we were going to do here tonight, he promptly took to the woods; but I am in hopes that during the social hour that follows, a committee of our Woman's Board which provides these delightful social occasions for us, will see if they cannot find him and bring him into view. It gives me one of the greatest pleasures of my life to present to the state of New Mexico through you, Senator Jones, representing the Governor on this occasion, this bronze of our great citizen, Mr. Frank Springer.

ACCEPTANCE FOR THE STATE

Senator Jones: *Dr. Hewett, Ladies and Gentlemen:* This is to me a very gratifying occasion. Of course, I bring to you the regrets of our distinguished Governor that he is unable to be with us tonight, and in the name of the state of New Mexico I accept this very wonderful gift. I appreciate the fact that I am privileged to bear his commission and in his name perform this function. It is gratifying to me as a citizen of New Mexico and as a resident of Las Vegas, which has for so many years been the home of our distinguished citizen that we honor tonight.

It was my privilege to become acquainted with him back in the middle eighties and to have become intimately associated with him during subsequent years. During fourteen years, the latter part of his activity in law, he and I occupied the same offices—a close association which to me made an ever binding friendship. As a young lawyer at the bar I used to sit in council with him, and I can bear witness to the many splendid qualities and great ability of Mr. Springer which have been told to you in such an eloquent way by Colonel Twitchell, who has also been my friend during all that period of years. So, from a personal standpoint, it is more than gratifying to appear before you tonight and join in the tributes to my friend. I know his ability as a lawyer and as a business man, and I have known something of his work as a scientist and philanthropist and big hearted citizen.

On behalf of the state of New Mexico I am particularly pleased to speak on this occasion. To me it is unusual in history. We are doing tonight two things which have seldom been done in the past. First, we are honoring a private citizen. If you will observe the statues of men throughout the United States, you will find that they are dedicated generally to those who have become famous either in war or statesmanship—in some public capacity. Tonight we are met to honor a citizen who has become famous in other walks of life. It is my hope that this occasion may become an inspiration to other sections of our country and to other countries. Let us look into the lives of those with whom we associate; let us commend the great and good things they are doing. To be noted at the bar, in business, in science, is something which should appeal to the rising generations, and tonight when we honor a distinguished citizen for all these qualifications, we honor one whose career should be emulated by the youth of the land. So I am especially pleased in the name of the state of New Mexico to accept this gift to the state presented by the friends of the man who has become thus distinguished and thus famous in private life.

In another respect is this occasion unusual. The subject of our meeting this evening is still in life. Too often, almost universally, tributes of this character are brought after we have been laid under the sod. How much better it is that we should strew flowers along life's pathway and make it brighter and pleasanter for those with whom we associate. This is another act which I hope may be repeated in the future in New Mexico and in the nation. So, speaking for the chief executive of this state, and I know that I speak for the citizens of New Mexico, I say that I am proud and grateful to receive this monument, which will be preserved by the state for all time to come, to one who is worthy of the highest esteem of our great people.

INDEX

The names of genera, species and larger divisions considered to be valid are printed in bold-face type; those regarded as synonyms, in italics. The bold-face figures indicate the page at which the genus, species or term is defined.

- Abrachiocrinus**, 98.
Abracrinus, 42.
Actinocrinidae, defined by Jackel, 41.
Aethocrinus sculptus, 142.
 Agassiz, Louis, 1.
Allocrinus, 30.
 — *benedicti*, 31.
 — *longidactylus*, 30.
 — *ponderosus*, 30.
 — *typus*, 30.
 American Museum Nat. Hist., 7, 11.
Ampheristocrinus typus, 132.
 Anal structures, modifications of, 41.
 — tube, 96.
 Angelin, N. P., 9, 41.
Anisocrinus, 65.
 — *angelini*, 66.
 — *greenei*, 66.
 — *oswegoensis*, 66.
Anthemocrinus, 14.
Aorocrinus, 47.
 — *clarkensis*, 47.
 — *nodosus*, 47.
 Arms in Calceocrinidae, 89.
 — axil, 90, 94, 125.
 — branching of, 112, 125.
 — disappearance of, 89, 99.
 — lateral, 90.
 — median, 89, 90, 95, 125.
 — modifications of, 89, 94, 99.
Asaphocrinus, 66.
 — *bassleri*, 66.
 — *minor*, 67.
 Asymmetry of stem and arms, 89.
 Axil-arm system, 90, 92, 93.
 — development, diagram, 94.
 Bainbridge formation, 75, 79.
Barrandeocrinus, 43.
 Basals, atrophy of, 89.
 — fusion of, 90.
 Base, modifications of, 89, 90.
 — mobility of crown on, 96.
 Bassler, R. S., 2, 10, 11, 80, 117.
 — Bibliographic Index, 10.
 Bather, F. A., 9, 20, 82, 88, 96, 97, 100, 104, 126, 131, 136, 137.
Batocrinidae, 41.
 — rejected by Jackel, 41.
 Beachler, Charles S., 6, 9, 10.
 Beech River formation, 3, 6.
 Benedict, A. C., 6, 10, 76.
 Bibliography of author's works, 146-152.
 Bilateral symmetry, 89, 96, 99.
 Birmingham Museum, 8, 63.
Blastoidea and **Cystidea**, 141.
 Bob formation, 2.
Botryocrinus polyxo, 138.
 — *tenuidactylus*, 137.
 Brachials, 90.
Brahmacrinus, 53.
 — *elongatus*, 53.
 Braun, Frederick, 6, 7, 10, 11.
 British Museum, 6, 11.
 Brownsport group, 2.
Calceocrinidae, 71, 88, 101, 104.
 — diagram of Forms A, B, C, D, 91.
 — distribution, 100.
 — evolutionary series, 90, 91, 92, 99.
 — genera of, 104.
 — mobility of crown on stem, 98.
 — mode of life, 98.
 — nomenclature, 101.
 — orientation of calyx, 88.
 — series of, 91.
 — shifting of stem, 96.
 — tri-radiate form, 92.
Calceocrinus, 101, 102, 103, 104, 115, 118.
 — diagram, 91.
 — generic position of, 103.
 — *alleni*, 119.
 — *bassleri*, 100, 117.
 — *bidentatus*, 118.
 — *bifurcatus*, 113, 117.
 — *bradleyi*, 101.
 — *contractus*, 118.
 — *foersteri*, 94, 100, 116.
 — *furcillatus*, 110.
 — *gotlandicus*, 94, 100.
 — *halli*, 103, 118.
 — *interpres*, 94, 100.
 — *nitidus*, 100, 113, 119.
 — *pinnulatus*, 94, 95, 100, 113, 116.
 — *pugil*, 100, 117, 120.
 — *rugosus*, 110.
 — *stigmatus*, 101, 118.
 — *tenax*, 116.
 — *tucanus*, 100, 117.
 — *typus*, 101, 103, 119.
Callicrinus, 18, 39.
 — *beachleri*, 40.
 — spp., 40.
 Calvin, S., 127.
 Calycanthocrinus, 74.
Calyptocrinidae, 35.
Camerata, 12.
 Carpenter, P. Herbert, 104.

- Carpocrininae, 41.
 Carpocrinus, 41.
 — sculptus, 41.
 — simplex, 41.
 Caryocrinus bulbosus, 143.
 — milliganae, 143.
 — ornatus, 143.
 — persculptus, 143.
 Castocrinus, 101.
 — billingsianus, 110.
 Catillocrinidae, paper on, 96.
 Catillocrinus, 73, 82, 89.
 Chapman, K. M., 9, 11.
 Characters in classification, 41.
 — changes in, 125.
 Cheiocrinus, 101, 103.
 — clarus, 101, 102, 121.
 — dactylus, 101.
 — ventricosus, 123.
 Chicago area, 8.
 — University of, 6, 11.
 Classification, 41.
 Clidochirus, 69.
 — americanus, 69.
 Clonocrinus, 32, 34.
 — occidentalis, 34.
 Clusters of crinoids, 63.
 Coccoocrinus, 48.
 — zone, 4.
 — bacca, 3, 48, 49, 50.
 — rosaceus, 49, 50, 51.
 Coelocrinus, 47.
 Collections and localities:
 — American Mus. Nat. Hist., 7, 11.
 — Beachler, Chas. S., 6, 9, 10.
 — Braun, Frederick, 6, 7, 10, 11.
 — Chicago area, 8.
 — University of, 6, 11.
 — Decatur County, Tennessee, 4.
 — Dudley, England, 8.
 — Elrod, Moses, 6, 10.
 — Gotland, Sweden, 8.
 — Hall, James, 6, 7, 11.
 — Hammell, John F., 6, 10.
 — Illinois, University of, 11.
 — Jewett, Col., 7.
 — Lockport, New York, 7.
 — Milligan, Mrs. J. M., 2, 11.
 — Newsom, Tennessee, 10.
 — New York State Museum, 7, 11.
 — Pate, W. F., 3, 4, 11.
 — Ringueberg, E. N. S., 7, 10.
 — Safford, 1, 2.
 — Ste. Genevieve Co., Missouri, 75, 79.
 — St. Paul, Indiana, 6.
 — Troost, Gerard, 1, 11.
 — Ulrich, E. O., 11.
 — Wabash, Indiana, 6.
 — Wachsmuth, Charles, 3.
 — Waldron, Indiana, 6.
 — Walker, Sir Edmund, 11.
 — Worthen, A. H., 2.
 Corymbocrinus, 34.
 Cremacrinus, 101, 102, 103, 104, 105, 116.
 — diagram of, 91.
 — articulosus, 100, 108.
 — billingsianus, 110.
 — decatur, 107.
 — furcillatus, 110.
 — Kentuckiensis, 109.
 — punctatus, 101, 102, 110.
 — rugosus, 110.
 — simplex, 107.
 — tubuliferus, 94, 96, 106.
 — ulrichi, 103.
 Crotalocrinus, 128.
 — cora, 129.
 — pulcher, 130.
 — rugosus, 129.
 Crown, mobility on base, 96, 98.
 — recumbent position, 99, 124.
 Cryptodiscus, 40.
 Culicocrinus, 49, 50.
 — nodosus, 51.
 — spinosus, 51.
 Cupellaerinites, 55.
 Cyathocrinidae, 131.
 Cyathocrinus, 133.
 — decatur, 134.
 — polyxo, 138.
 — striolatus, 134.
 — wilsoni, 134.
 Cylicocrinus canaliculatus, 43.
 — spinosus, 44.
 Cypellaerinus, 55.
 Cyphocrinus, 14.
 — chicagoensis, 15.
 — gorbyi, 15.
 Cytocrinus laevis, 28.

 Decatur formation, 2, 5, 47.
 De Loriol, 104.
 Deltacrinus, 101, 102, 103.
 — clarus, 102, 121.
 Dendrocrinus nucleus, 138.
 Desmidocrinus, 43.
 — dubius, 43.
 — laurelianus, 43.
 Dimerocrinidae, 12.
 Dimerocrinus, 12.
 — inornatus, 13.
 — milliganae, 13.
 — nodobasis, 13.
 — planus, 13.
 Dolatocrinus, paper on, 23, 59.
 Dudley, England, 8.
 — collections at, 8.
 — Museum, 8, 11, 63.

 Ectenocrinus, 89.
 Elrod, Moses, 6, 10.
 Emperocrinus indianensis, 23.
 Eomylodactylus rotundus, 86.
 Eucalyptocrinus, 18, 35.
 — zone, 4, 37.
 — crassus, 16, 39.
 — depressus, 38.
 — elrodi, 36.
 — lindahli, 36.
 — magnus, 36.
 — milliganae, 37.
 — ovalis, 36.

- *pernodosus*, 38.
- *sculptilis*, 38.
- *splendidus*, 37.
- *ventricosus*, 37.
- spp., 38, 39.
- Eucheirocrinus*, 101, 103, 104, 111.
 - diagram of, 91.
 - *anglicus*, 111, 112, 113.
 - *chrysalis*, 111, 112.
 - *indianensis*, 114.
 - *minor*, 112.
 - *ontario*, 111, 112, 113.
- Euclinus*, 12, 14.
- Eudimerocrinus*, 14.
 - *multibrachiatus*, 14.
- Evolution in the act, 99.
- Evolutionary series, 90, 91.
- Faunas, American and European, 4, 8, 55, 72.
- Fistulata*, 85.
- Flexibilia*, 64.
- Florin, A., 8.
- Foerste, Aug. F., 2, 6, 7, 41, 117.
- Foreign Silurian, 8.
- Forms of Calceocrinidae, A, B, C, D, 90, 91.
- Fusion of basals, 90.
- Gasterocominae, 133.
- Gazacrinus*, 15.
 - *depressus*, 17.
 - *inornatus*, 16.
 - *magnus*, 17.
 - *milliganae*, 18.
 - *ramifer*, 18.
 - *stellatus*, 19.
 - *ventricosus*, 17.
- Genus and species indet., 21, 39, 58.
- Gissocrinus*, 135.
 - *approximatus*, 137.
 - *delicatus*, 136.
 - *lyoni*, 135.
 - *magnibrachiatus*, 136.
 - *quadratus*, 137.
- Glyptaster*, 12.
- Goldring, Winifred, 21, 27, 96.
- Gotland, Sweden, 8.
- Gnorimocrinus*, 70.
 - *cirrifer*, 70.
 - *varians*, 70.
- Gray, John, 8.
- Habrocrinus*, 42.
 - *ornatissimus*, 42.
 - *ornatus*, 42.
- Hall, James, 6, 7, 9, 101, 103.
- Halysiocrinus*, 101, 102, 104, 116, 121.
 - diagram of, 91.
 - *bradleyi*, 101.
 - *carinatus*, 122.
 - *dactylus*, 123.
 - *keyserensis*, 121.
 - *marylandensis*, 122.
 - *nodosus*, 124.
- *perplexus*, 97, 123.
- *secundus*, 96, 121.
- Hammell, John F., collection, 6, 10.
- Hapalocrinus*, 52.
 - *cirrifer*, 52.
 - *devonicus*, 53.
 - *gracilis*, 52.
 - *pinnulatus*, 52.
 - *retiaris*, 52.
 - *tennesseensis*, 53.
 - *tuberculatus*, 53.
- Haplocrinus*, 49, 50, 73, 81.
 - *mespiliformis*, 50, 81.
- Herpetocrinus*, 85.
- Heterocrinidae*, 85, 89.
- Heterocrinus*, 89, 91.
 - diagram of, 91.
 - *bellevillensis*, 92.
- Hinge, 96.
- Holcroft, Charles, 8, 63.
- Homocrinus*, 131.
- Hormocrinus*, 66.
 - *tennesseensis*, 66.
- Hypocrinus*, 133.
- Hyptiocrinus*, 15.
 - *typus*, 15.
- Ichthyocrinidae*, 68.
- Ichthyocrinus*, 68.
 - *subangularis*, 69.
- Idiocrinus*, 15, 16.
 - *elongatus*, 16.
- Illinois, University of, 11.
- Inadunata*, 71.
- Indiana area, 6.
- Insertae sedis, 54, 139.
- Iocrinus*, 85.
- Jaekel, O., 41, 49, 82, 99, 119.
- Jewett, Col., 7.
- Klinteberg, G., 8.
- Lampterocrinus*, 19.
 - *parvus*, 20.
 - *roemeri*, 21.
 - *sculptus*, 21.
 - *tennesseensis*, 20.
- Larviformia*, 71.
- Laurelocrinus*, 32.
 - *gibbosus*, 33.
 - *paulensis*, 33.
 - *spinoradialis*, 34.
 - *wilsoni*, 33.
- Lecanocrinidae*, 64.
- Lecanocrinus*, 65.
 - *caliculus*, 67.
 - *elongatus*, 65.
 - *meniscus*, 65.
 - *pisiformis*, 65.
 - *pusillus*, 65.
- Lecythiocrinus adamsi*, 133.
 - *problematicus*, 133.

- Lepadocrinus gebhardi*, 142.
Leptocrinus, 42.
 Liljevall, Georg, 88.
 Lobelville formation, 2.
 Lockport, New York, 7, 9, 10.
 Lyon, Sidney S., 2, 11, 50.
Lyonicrinus, 48.
 — *bacca*, 51.
Lyriocrinus dactylus, 24.
 — *melissa*, 24.
Lysocystites sculptus, 142.

Macrostylocrinus, 25.
 — *anglicus*, 27.
 — *fasciatus*, 25.
 — *granulosus*, 26.
 — *indianensis*, 27, 32.
 — *laevis*, 26.
 — *meeki*, 26.
 — *ornatus*, 27.
 — *pustulosus*, 26.
 — *striatus*, 27.
 Madely, William, 8.
 Main-axils, 90.
Mariacrinus, 28.
 — *aureatus*, 29.
 — *carleyi*, 29.
 — *rotundus*, 29.
 — *sp.*, 29.
Marsipocrininae, 55.
Marsipocrinus, 55.
 — *coelatus*, 52.
 — *concavus*, 60.
 — *excavatus*, 60.
 — *inflatus*, 58.
 — *magnificus*, 61.
 — *rosaeformis*, 57, 59.
 — *stellatus*, 60.
 — *striatissimus*, 61.
 — *striatus*, 58.
 — *tennesseensis*, 56, 57, 59.
 — *turbinatus*, 62.
 — *verneuili*, 59.
Marsupiocrinus, 55.
 Median arm, 89, 90, 95.
 Meek and Worthen, 101, 104.
Melocrinidae, 25.
 — genus and species indet., 32.
Melocrinus, 27.
 — *aequalis*, 28.
 — *obconicus*, 28.
 — *oblongus*, 28.
 — *roemeri*, 28.
 — *onondaga*, 28.
 — *spectabilis*, 27.
 — *tennesseensis*, 27.
 Merrill, Dr. Geo. P., 11.
 Miller, S. A., 6, 15, 16, 73, 76, 104.
 Miller and Gurley, 78, 131.
 Milligan, Mrs. J. M., 2, 11, 131.
 Mobility of crown, 98.
Monobrachiocrinus, 98.
 Moody, Margaret W., 11.

Mycocrinus, 73.
Myelodactylus, 7, 85, 98.
 — *ammonis*, 86.
 — *brachiatus*, 87.
 — *brevis*, 86.
 — *bridgeportensis*, 86.
 — *convolutus*, 86.
 — *extensus*, 87.
 — *fletcheri*, 86.
 — *gorbyi*, 86.
 — *keyserensis*, 87.
 — *nodosarius*, 87.
 — *schucherti*, 87.
Myrtillocrinus americanus, 49.
Mysticocrinus, 83.
 — *wilsoni*, 84.
 Münster, 74.
 Muscular articulation, 88.

 New Mexico, presentation to State, 153.
 Newsom, Tennessee, 10, 24, 36.
 New York area, 7.
 — State Museum, 7, 11.
 Niagaran, sections, 3, 5.
 Number of species, 8.
Nyctocrinus, 34.
 — *magnitubus*, 34.

 Oehlert, D., 81.
Ohioocrinus, 89.
 Oral plates, 73, 75, 82.
 Origin of this work, 9.

Palaeocrinus, 131.
Paragazacrinus, 22.
 — *rotundus*, 22.
Parastephanocrinus typus, 139.
Parisocrinus siluricus, 135.
 Pate, W. F., 2, 4, 9, 11.
Patelloocrinus, 31.
 — *laevis*, 32.
 — *ornatus*, 31.
 — *pinnulatus*, 31.
 — *rugosus*, 32.
 Patelloid plates, 137.
Paulocrinus, 22.
 — *biturbinatus*, 22.
Periechocrininae, 44.
Periechocrinus, 44.
 — *laevis*, 46.
 — *minor*, 46.
 — *moniliformis*, 44, 45.
 — *tennesseensis*, 45.
 — *whitei*, 44.
 — *sp.*, 45.
 Personal record, 144.
Petalocrinus, 126.
 — *inferior*, 127.
 — *longus*, 127, 128.
 — *mirabilis*, 126.
 — *visbyensis*, 127.
Phoenicocrinus, 42.
 Phylogenetic modifications, 95, 125.

- Pinnules, doubled, 20, 42.
 — openings for, 59.
Pionocrinus, 42.
Pisocrinus, 72, 89.
 — baccula, 75.
 — benedicti, 77.
 — campana, 76, 80.
 — gemmiformis, 74.
 — globosus, 79, 80.
 — gorbyi, 78.
 — granulosus, 75.
 — milliganae, 78.
 — ollula, 78, 80.
 — pilula, 76, 80.
 — pocillum, 76.
 — pyriformis, 80.
 — quinquelobus, 77, 78, 79.
 — sphericus, 75, 78, 79.
 — tennesseensis, 75, 79.
Platycrinidae, 82.
 — genus and species indet., No. 1, 51.
 — — No. 2, 51.
 — — No. 3, 52.
Platycrinus retarius, 52.
 — huntsvillae, 52, 53.
Primibrachs, unequal-faced, 94.
Proclivocrinus, 101.
 Progressive characters, 94, 95, 125.
Protaxocrinus, 69.
 — robustus, 70.
Pseudocrinus quadrifasciatus, 142.
Pycnosaccus, 67.
 — americanus, 67.
 — dubius, 68.
 — laurelianus, 67.
 — patci, 67.
 — welleri, 67.
 Racine dolomite, 8.
 Respiratory pores, 59.
 Resser, C. E., 11.
Rhodocrinidae, 22.
 — genus and species indet., 24.
 Riks Museum, Stockholm, 8.
 Ringueberg, E. N. S., 7, 9, 10, 88, 101, 103, 104.
 Rochester shale, 7, 10.
 Roemer, C. F., 1, 18.
 Rominger, Carl, 2.
 Rowley, R. R., 75, 79.
Saccocrinus, 44, 45.
 — benedicti, 46.
 — cuspidatus, 46.
 — speciosus, 45.
 — umbrosus, 46.
 Safford, J. M., 1, 11.
Sagenocrinidae, 68.
Sagenocrinoidea, 64.
Sagenocrinus, 68.
 — americanus, 68.
 — clarki, 68.
 St. Paul, Indiana, 6.
 Schuchert, C., 129.
 Schultze, L., 74, 81.
Scyphocrinus, paper on, 23, 57, 59.
 Segments of 1. ant. radial, 89, 94, 95.
 Series of Calceocrinidae, 91.
 Slocum, A. W., 8, 9, 129.
 Species, number of, 8.
 Springer, F., 23, 57, 59.
 — Paper on Catillocrinidae, 96.
 — — Dolatoocrinus, 21, 23, 59.
 — — Scyphocrinus, 23, 57, 59.
 — — Unusual Forms, 25, 85.
 — Zittel-Eastman Textb., 101.
Steganocrinus, 20.
 — sculptus, 20.
 Stem, asymmetric location, 89, 96.
 — bending of crown on, 88, 99.
 — indentation by, 110, 122.
 — sessile condition of, 99.
 — shifting of, 96.
Stephanocrinus, 139.
 — angulatus, 139.
 — gemmiformis, 140.
Stribalocystites gorbyi, 142.
 Subanal plate, 89.
 Suborals, 49.
 Supplementary plates, 97.
Symbathocrinus, 73.
 — tennesseensis, 79, 80.
Synchiocrinus anglicus, 119.
 T-piece, 89.
Taxocrinidae, 69.
Taxocrinoidea, 69.
 Tennessee area, 1.
Tetracystis fenestratus, 141.
 Text-figures, 91, 94, 112.
Thalamocrinus, 131.
 — cylindricus, 132.
 — elongatus, 132.
 — globosus, 132.
 — ovatus, 131.
 Thomas, A. O., 126.
Thysanocrinus, 12.
Tiarocrinus, 81, 82.
 — quadrifrons, 81.
 — soycei, 81.
Torynocrinus, 98.
Triacrinus, 74, 89.
 — altus, 74.
 — depressus, 74, 80.
 Troost, Gerard, 1, 11.
Troostocrinus, 141.
 — zone, 4.
 — reinwardti, 141.
 — sanctipaulensis, 141.
Uintacrinus, 137.
 Ulrich, E. O., 11, 97, 100, 101.
 United States National Museum, 8, 11.
 Unusual Forms of Fossil Crinoids, 25, 85.
 Vanderbilt University, 2, 11.
 Variation in characters, 124, 125.

- Wachsmuth, Charles, 2, 9, 11, 20.
Wachsmuth and Springer, 15, 16, 18, 21, 37, 49, 83,
96, 101, 129.
Walcott, Hon. Charles D., 11.
Waldron, Indiana, 6, 9, 64.
Walker, Sir Edmund, 6, 11.
Wanner, Joh., 133.
Washburn, C. C., 6, 10.
Weller, Stuart, 8, 9, 15, 80, 129.
Weller and Davidson, 126.
Wilson, Herrick E., 7, 10, 40.
- Wilsonicrinus**, 23.
—— **discoideus**, 23.
Wood, Elvira, 80.
Worthen, A. H., 2.

Zittel, K. A. von, 81.
Zittel-Eastman, Textb. Pal., 101.
Zophocrinus, 81, 82.
—— **globosus**, 83.
—— **howardi**, 82.
—— **pyriformis**, 83.

EXPLANATION OF PLATES

Enlargement, if any, of the drawings is indicated by the sign at the end of the paragraph. Unless so noted, the figure is of natural size. All the specimens figured, except as otherwise stated, are in the author's collection now in the United States National Museum.

PLATE 1

(All figures not otherwise noted natural size)

	PAGE
<i>Dimerocrinus planus</i> new species	13
FIG. 1. Complete crown of maximum size, with long stem; 10 heavy arms.	
2, 3. Lateral and posterior views of two other crowns.	
4. Crown with more delicate structure.	
5. Calyx showing anal side.	
6. Basal view of crushed calyx showing IBB.	
7, 7a. Large calyx with abnormal anal side, perhaps of this species; has IBB invisible from side view; posterior and lateral views.	
Beech River formation; Decatur County, Tennessee.	
<i>Dimerocrinus milliganae</i> (Miller and Gurley)	13
FIG. 8. Type specimen of " <i>Glyptaster</i> " <i>milliganae</i> , formerly in the collection of Mrs. J. M. Milligan.	
Horizon and locality, same as last.	
<i>Dimerocrinus nodobasis</i> new species	13
FIG. 9. Oblique basal view of large crown, showing nodes on BB, and minute IBB concealed by top columnal.	
9a. Another view of calyx, showing anal series.	
10. Basal view of smaller specimen.	
Horizon and locality, same as last.	
<i>Dimerocrinus inornatus</i> (Hall)	13
FIG. 11. Large calyx, from l. post. radius, with strongly projecting basals.	
11a. Basal view of same.	
12. Posterior view of smaller calyx.	
Waldron shale; Hartsville, Indiana.	
<i>Eudimerocrinus multibrachiatus</i> new species	14
FIG. 13. Nearly complete crown, with arms frequently branching, biserial above and below the bifurcations; basal and radial series surmounted by spinose cusps; IBB restricted to the column-facet. Anterior view.	
13a. Anal side of calyx, somewhat restored.	
13b. The base, with column removed, to show relation of IBB.	
14. Another complete crown, seen from l. post. radius.	
Beech River formation; Decatur County, Tennessee.	
<i>Cyphocrinus gorbyi</i> S. A. Miller	15
FIG. 15. Lateral view of calyx, showing drooping of arm-bases, pinnule openings between them, and spiniferous tegmen.	
16. Dorsal view of another specimen, showing arrangement of plates and distribution of arms.	
17, 18. Ventral views of two specimens showing composition of tegmen, position of anus, and spiniferous ambulacrals.	
Laurel formation; St. Paul, Indiana.	
— genus and species indet.	21
FIGS. 19, 20, 21, 22. Fragments of stem of unknown crinoid from the Laurel limestone, St. Paul, Indiana.

PLATE 2

(All figures natural size unless otherwise stated)

	PAGE
<i>Gazacrinus inornatus</i> S. A. Miller.....	
	16
FIG. 1. Post. view of calyx and rays to level of IBr. $\times 5/2$.	
2. Ant. view of crown with part of arms intact. $\times 2$.	
3. Dorsal view of calyx, showing excavate base with IBB concealed by a columnal; r. ant. side above. $\times 5/2$.	
4. Ventral view of cup, showing articulating faces of RR at level of IBr, the strong inter-radial processes for support of orals, which are wanting here. $\times 2$.	
5. The tegmen, with oral plates in position resting on the interradial processes, except the central part of post. oral, or a part corresponding to it, here broken away. $\times 2$.	
5a. Basal view of same, showing IBB at bottom of cavity. $\times 2$.	
6. A detached set of oral plates lacking central part. $\times 3/2$.	
7. Tegmen of another specimen with all oral plates in place, surmounted by ridges not present in fig. 5; fractured part restored in outline. $\times 2$.	
8, 9. Basal views of two other specimens. $\times 3/2$. The ventral structures vary in different specimens.	
Laurel limestone; St. Paul, Indiana.	
<i>Gazacrinus magnus</i> new species.....	
	17
FIG. 10. Very elongate central pyramid of tegmen, being the 5 oral plates divided by vertical sutures, one of which is seen at the right lower corner. $\times 2$.	
10a. Internal view of same, showing division of the plates; the outer lighter area corresponds to the lower shaded one in fig. 11. $\times 2$.	
11. An isolated plate, one of the 5 orals from a still larger specimen, showing the sharp sutural faces at either side; the lower shaded area is the surface of attachment to interradial processes, corresponding to the lighter area in fig. 10a. $\times 2$.	
12. Another ventral pyramid having only half as many plications as in fig. 10.	
Laurel limestone; St. Paul, Indiana.	
<i>Gazacrinus depressus</i> new species.....	
	17
FIG. 13. Specimen with depressed, rounded calyx and heavy arms; base crushed. $\times 3/2$.	
14. Detached base, perhaps of this species, somewhat resembling the Tennessee species. $\times 3/2$.	
Laurel limestone; St. Paul, Indiana.	
<i>Gazacrinus ventricosus</i> (Wachsmuth and Springer).....	
	17
FIGS. 15, 15a, 15b. Posterior, basal and ventral views of the type specimen, the latter having the low, radiating pyramid in place, somewhat imperfect in the median part. $\times 4$.	
16, 16a. Distal and proximal views of detached ventral pyramid from another specimen. $\times 2$.	
Laurel limestone; St. Paul, Indiana.	
<i>Gazacrinus ramifer</i> (Roemer)	
	18
FIG. 17. Dorsal view of calyx with anal side above, showing the broad basal cavity lodging IBB; with pentagonal projecting rim upon basals and less prominent triangles passing to the radial series.	
18. A nearly complete crown from 1. post. radius, with long and heavy biserial arms.	
19, 19a. Anterior and posterior views of crown with part of arms, the latter having the ventral pyramid clearly in view.	
20, 20a. A complete crown with the triangular sculpture on basals and radials sharply accentuated; ant. and r. post. iR views.	
21. Part of crown of similar well marked specimen from Wayne County, Tennessee.	
Beech River formation; all but the last from Decatur County, Tennessee.	
<i>Gazacrinus milliganae</i> (Miller and Gurley).....	
	18
FIG. 22. Basal view of type; plates raised into prominent nodes.	
23, 24. A detached ventral pyramid, and a larger specimen, perhaps of this species.	
Beech River formation; Decatur County, Tennessee.	

Gazacrinus stellatus new species.....

PAGE

19

FIGS. 25, 25a, b, c. Posterior, anterior, basal and ventral views of unique specimen, with highly sculptured surface, and pentagonal and triangular projections similar to those of *G. ramifer*, the latter forming a conspicuous star. In 25c there is a remnant of the ventral pyramid, much distorted.

Linden formation, Lower Devonian; Hardin County, Tennessee.

PLATE 3

(Figures natural size unless otherwise indicated)

	PAGE
<i>Lampterocrinus tennesseensis</i> Roemer	20
FIG. 1. A typical calyx from posterior side, showing the anal series with sharp median ridge leading to the tube.	
2. L. ant. view of another calyx, showing bulging of the anal side at the right, and strong ridges passing from basals to radial series.	
3, 4. Two other specimens differing in size and contour.	
5, 5a. Complete crown with tubular extensions single from each ray, bearing alternating pinuliferous side arms; with anal tube preserved; it is free from the matrix, and the two views show opposite sides.	
6. Fragment of another specimen showing details of arms, with two pinnules from each brachial. $\times 2$.	
Beech River formation; Decatur County, Tennessee.	
<i>Lampterocrinus sculptus</i> new species.....	21
FIGS. 7, 8. Two calices with more elaborate sculpturing and wider expanding contour—from same locality, and perhaps a variety of the preceding.	
<i>Lampterocrinus roemeri</i> new species	21
FIG. 9. Characteristic specimen, with bases of tubular extensions attached, and part of the pentagonal stem.	
10. Complete crown, with anal tube; radial extensions and side-arms well preserved.	
11, 12, 13. Three calices showing uniform character of the simplified ridges.	
14. Ventral view, showing base of anal tube.	
Beech River formation; Decatur County, Tennessee.	
————— genus and species indet.....	
FIG. 15. Fragment with six-sided stem, sharply angular, and only part of basals. $\times 2$.	
15a. Transverse view showing the large lumen, and angles of the hexagonal stem. $\times 2$.	
Beech River formation; Decatur County, Tennessee.	

PLATE 4

(Figures natural size unless otherwise stated)

	PAGE
Paragazacrinus rotundus new species.....	22
FIG. 1. Complete calyx, with tegmen surmounted by a central spine and divided by interambulacral and secondary ridges. $\times 5/3$.	
2. Dorsal view of calyx showing the 5 IBB at bottom of indented pit; RR separated all around by BB and iBr. $\times 5/2$.	
3. Another specimen, with smaller basal cavity. $\times 2$.	
3a. Ventral view of same with 5 large tegminal plates in place, but central spine lacking; shows openings for 10 arms. $\times 2$.	
4. Isolated pyramid with central spine attached. $\times 2$.
Laurel limestone; St. Paul, Indiana.	
Paulocrinus biturbinatus new genus and species.....	22
FIGS. 5, 5a. Post. and r. post. views of complete calyx with tegmen passing into a large anal tube; all plates strongly nodose or rugose; r. post. ray has only 1 IBr.	
5b. Basal view of same, showing irregular form of IBB, 2 large and 3 small, and 5 BB and 2 RR connecting with them.	
5c. Diagram showing arrangement of basal plates.
Laurel limestone; St. Paul, Indiana.	
Wilsonicrinus discoideus new genus and species.....	23
FIGS. 6, 6a. Dorsal and ventral views of the only specimen—a broad disk slightly convex below and passing into a strong anal tube above; 4 IBB; fixed pinnules from cuneate secundibrachs incorporated in calyx wall, leading to interrational pinnule openings at margin, conspicuous on ventral side, between the 10 arm-openings. $\times 3/2$.	
Emperocrinus indianensis Miller and Gurley.....	23
FIGS. 7, 7a. A typical specimen, of simpler structure than the preceding, and having 5 IBB; interrational pinnule openings shown; dorsal and ventral views.
Laurel limestone; St. Paul, Indiana.	
Rhodocrinidae , genus and species indet.....	24
FIGS. 8, 8a. Fragment of large stem standing at angle with calyx covering 4 large IBB as seen from the interior; RR separated all around; traces of injury by boring parasites. Outer and inner views.
Laurel limestone; St. Paul, Indiana.	
Macrostylocrinus fasciatus Hall.....	25
FIG. 9. Posterior view of highly elevated calyx, with anal series leading to opening at extreme margin of tegmen. $\times 2$.	
9a. Basal view of same. $\times 3/2$.	
9b. The tegmen, with ambulacra fully exposed flanked by numerous small plates; a small anal opening at extreme edge of tegmen. $\times 3/4$.
Waldron shale; Hartsville, Indiana.	
Macrostylocrinus granulatus Hall.....	26
FIG. 10. Post. view of calyx from Madison, Indiana, with anal series leading to opening; surface granulate. $\times 2$.	
10a. The tegmen, with anal opening covered by pyramid of small plates; a large posterior oral; and ambulacra radiating to arm-openings with numerous small plates between of uncertain arrangement. $\times 2$.	
11, 11a, 11b, 11c. R. ant., post., and ventral views of small specimen from Newsom, Tennessee, with tegmen well preserved. $\times 2$.	
12, 13. Specimens from Newsom and Decatur County, Tennessee, perhaps of this species. $\times 3/2$.
Waldron shale; Indiana and Tennessee.	

K. M. Chapman del.

	PAGE
<i>Macrostylocrinus laevis</i> new species	26
FIGS. 14, 15. Dorsal oblique views of two crowns, with strong biserial arms; calyx more or less crushed.	
16, 17, 18, 19. Lateral views of 4 calices, showing various contours; surface in all smooth; 16 and 17 flattened.	
Beech River formation; Decatur County, Tennessee.	
<i>Macrostylocrinus meeki</i> (Lyon)	26
FIGS. 20, 21, 22. Post., ant. and basal views of three specimens lacking the striate ornament; show the broadly rounded calyx, sharp ridges following the radial series, and the wide IBr; in 22 post. is at the right.	
23. A variant perhaps of different species, showing small size of arms.	
Beech River formation; Decatur County, Tennessee.	
<i>Macrostylocrinus ornatus</i> Hall	27
FIG. 24. The type species; a complete crown and stem with terminal root; biserial arms and pinules; from the Rochester shale at Lockport, New York, collected by Frederick Braun.	
(?) <i>Macrostylocrinus pustulosus</i> new	26
FIGS. 25, 25a. Post. and dorsal views of the only specimen, having anal side like <i>Macrostylocrinus</i> , but radial facets more like the <i>Platycrinidae</i> .	
Beech River formation; Decatur County, Tennessee.	

PLATE 5

(Figures natural size unless otherwise stated)

	PAGE
<i>Melocrinus tennesseensis</i> new species	27
FIG. 1. Crown with part of arms and long stem; the two main rami not fused; calyx plates displaced. Beech River formation; Decatur County, Tennessee.	
<i>Melocrinus</i> (?) <i>spectabilis</i> Angelin	27
FIG. 2. A very complete specimen with arms and stem from the Wenlockian at Dudley, England— for comparison with the Tennessee form. Note the numerous IBr plates.	
3. Dorsal view of another specimen from the same locality.	
<i>Melocrinus onondaga</i> new species	28
FIG. 4. A very small specimen from the Onondaga at Sylvania, Ohio. $\times 2$.	
<i>Melocrinus oblongus</i> Wachsmuth and Springer	28
FIGS. 5, 6. Lateral views of two calices. Louisville limestone; Jefferson County, Kentucky.	
<i>Cytocrinus laevis</i> Roemer.....	28
FIGS. 7, 7a. Lateral and ventral views of calyx.	
8. Post. view of smaller specimen, showing protuberant anal opening in tegmen. Beech River formation; Decatur County, Tennessee.	
<i>Mariacrinus carleyi</i> (Hall)	29
FIGS. 9, 10. Posterior and r. post. views of calyx. Laurel limestone; St. Paul, Indiana.	
<i>Mariacrinus aureatus</i> S. A. Miller	29
FIGS. 11, 11a. Lateral and dorsal views of calyx. Laurel limestone; St. Paul, Indiana.	
(?) <i>Mariacrinus rotundus</i> new species.....	29
FIGS. 12, 12a. L. post. and dorsal views of calyx. Decatur limestone; above Clifton on Tennessee River, Perry County, Tennessee.	
<i>Mariacrinus</i> sp.	29
FIG. 13. A well marked specimen with arms and stem, from the Wenlockian at Dudley, England.

K. M. Chapman del.

PLATE 6

(Figures natural size unless otherwise stated)

	PAGE
<i>Alloccrinus typus</i> Wachsmuth and Springer	30
FIGS. 1, 2. Two crowns, showing characteristic heavy arms, with calyx expanding upwards.	
3. Crown somewhat displaced, with complete stem.	
4. A smaller specimen.	
Beech River formation; Decatur County, Tennessee.	
<i>Alloccrinus ponderosus</i> new species	30
FIGS. 5, 6. Two crowns, showing the heavy arms, medially swollen, and very narrow calyx.	
Beech River formation; Decatur County, Tennessee.	
<i>Alloccrinus longidactylus</i> new species	30
FIG. 7. Complete crown with stem, showing extreme length of arms, and wide, nodose calyx	
8. Another specimen showing distribution of pinnules.	
9, 9a. Lateral and dorsal views of calyx.	
10. Var. <i>magniradialis</i> . Crown with nearly complete arms, and extremely large radials.	
Beech River formation; Decatur County, Tennessee.	
<i>Alloccrinus benedicti</i> S. A. Miller	31
FIGS. 11, 12. Two crowns with more rounded calyx plates, from Laurel limestone; St. Paul, Indiana.	
<i>Patelloccrinus ornatus</i> new species	31
FIGS. 13, 14. Lateral and basal views of calyx.	
15, 15a. Opposite lateral views of crown free from matrix, with tubercular surface, and 10 heavy, biserial arms. $\times 3/2$.	
Laurel limestone; St. Paul, Indiana.	
<i>Patelloccrinus rugosus</i> new species.....	32
FIGS. 16, 17. Two calices seen from l. ant. and r. post., with very large radials and rugose surface.	
18. Basal view of another specimen. $\times 2$.	
Laurel limestone; St. Paul, Indiana.	
<i>Patelloccrinus laevis</i> new species.....	32
FIGS. 19, 20. Lateral views of two specimens with smooth surface; IIBr in contact all around.	
$\times 3/2$.	
Laurel limestone; St. Paul, Indiana.	
— genus and species indet.....	32
FIG. 21. Complete crown, with 10 heavy biserial arms, 3 small BB, very large RR, followed by one IBr almost filling face of R; one small iBr.	
21a. Opposite, probably post., view of lower part with arms broken away, showing large iBr.	
$\times 2$.	
21b. The same part seen from a different angle. $\times 2$.	
Beech River formation; Decatur County, Tennessee.

PLATE 7

(Figures natural size unless otherwise stated)

	PAGE
<i>Laurelocrinus paulensis</i> new species	33
FIGS. 1, 2. Maximum calices to upper IIBr, elongate with broad base, sides rising nearly straight; first IBr about square; iBr elongate, 8- or 9-sided, followed by a second one much narrower; RR large, extending into basal cavity.	
3. Fragment with iBr.	
4, 5, 6. Bases of similar specimens with 3 unequal BB limited to shallow cavity, occupied largely by column-facet with very large lumen.	
7, 8. Variety with first IBr elongate and iBr acuminate, not followed by a second. Laurel limestone; St. Paul, Indiana.	
<i>Laurelocrinus wilsoni</i> new species	33
FIGS. 9, 10, 11. Much smaller form than last, with first IBr hexagonal, iBr 5-sided followed by one or two others; 3 calices. $\times 3/2$.	
12. Base of similar specimen, little depressed, with small column-facet and lumen. $\times 3/2$.	
13. Specimen showing general contour of calyx to higher level, upper plates displaced. $\times 3/2$.	
13a. Base of same, showing striate surface; one large basal bisected, making 4 BB. $\times 3/2$. Laurel limestone; St. Paul, Indiana.	
<i>Laurelocrinus gibbosus</i> new species	33
FIG. 14. Very small calyx, with extremely large RR and iBr, and small quadrangular IBr; all strongly gibbous. $\times 5/4$.	
14a. Base of same, very convex, with small column-facet and lumen. $\times 3/2$.	
15, 16. Base of another specimen, and isolated R and IBr of another. Laurel limestone; St. Paul, Indiana.	
<i>Laurelocrinus spinoradialis</i> new species	34
FIG. 17. Lateral view of calyx, with sharp spines projecting downward from RR; first IBr quadrangular; iBr two in succession, very elongate, 8- or 9-sided, and a similar iIIBr. $\times 3/2$.	
17a. Opposite view of lower part of same. $\times 3/2$.	
17b. Base of same. $\times 3/2$. Laurel limestone; St. Paul, Indiana.	
<i>Clonocrinus occidentalis</i> new species	34
FIGS. 18, 18a. Lateral and dorsal views of calyx, showing deeply sunken basal cavity, with 4 small BB at bottom. Decatur limestone; near Perryville, Tennessee.	
<i>Eucalyptocrinus crassus</i> Hall	39
FIGS. 19, 20. Lateral and ventral views showing the ventral pyramid, for comparison with <i>Gazacrinus</i> on plate 2. (See also N. A. Crin. Cam., pl. 83, figs. 12, 13, 14.) Waldron shale; Hartsville, Indiana.	
<i>Carpocrinus sculptus</i> new species	42
FIG. 21. Anterior view of the only specimen, showing the highly sculptured surface, and a few quadrangular ossicles of the arms. $\times 2$.	
21a. Posterior view, showing full series of anal plates leading to the opening; sculpture eroded on this side by weathering. $\times 2$.	
21b. Ventral view of same, showing the oral plates, ambulacra, and anal opening through tegmen. $\times 2$. Laurel limestone; St. Paul, Indiana.	

	PAGE
<i>Carpocrinus simplex</i> (Phillips).....	42
FIGS. 22, 22a. Complete crown, with stem, of specimen from Gotland, with pinnules well developed, two to each brachial, as shown in fig. 22a, enlarged. $\times 2$.	
23. Another specimen from Gotland, showing the tegmen with ambulacra exposed. Wenlockian; Gotland, Sweden.	
(?) <i>Desmidocrinus laurelianus</i> new species.....	43
FIG. 24. Crown with part of stem, r. ant. view; has 4 uniserial arms to the ray.	
24a. Posterior view of calyx, showing anal series. $\times 3/2$.	
24b. Base of same. $\times 3/2$. Laurel limestone; St. Paul, Indiana.	
(?) <i>Desmidocrinus dubius</i> new species.....	43
FIG. 25. Crown with uniserial arms, bifurcating once or twice; post. view. Same horizon and locality as last.	
<i>Cylicocrinus canaliculatus</i> S. A. Miller.....	43
FIGS. 26, 27. Posterior and anterior views of calyx.	
28, 28a. Lateral and basal views of calyx with 3 equal basal plates projecting in a wide rim.	
29. Basal view of small specimen with unequal basals and narrower rim. Laurel limestone; St. Paul, Indiana.	
<i>Cylicocrinus spinosus</i> new species.....	44
FIG. 30. Basal hexagon, with strong spinose projections on BB; dorsal view showing small cavity occupied by a columnal. $\times 3/2$.	
30a. Interior view of same, showing sutures connecting with lumen. $\times 3/2$. Same horizon and locality as last.	

PLATE 8

(Figures natural size unless otherwise stated)

	PAGE
<i>Eucalyptocrinus elrodi</i> S. A. Miller.....	36
FIG. 1. Dorsal view of unusually large calyx, with profuse tubercular ornamentation.	
2. Lateral view of another specimen with much fewer tubercles.	
Waldron shale; Newsom, Tennessee.	
<i>Eucalyptocrinus magnus</i> Worthen.....	36
FIG. 3. One of the largest known specimens belonging to this genus.	
Same horizon and locality.	
<i>Eucalyptocrinus lindahli</i> Wachsmuth and Springer.....	36
FIG. 4. Complete crown of typical specimen, showing the heavy, rounded arms, and partitions sunken in a deep groove.	
5. A smaller specimen, probably of this species.	
Beech River formation; Decatur County, Tennessee.	
<i>Eucalyptocrinus milliganae</i> Miller and Gurley.....	37
FIG. 6. A complete crown of maximum size, with stem; partitions widely projecting in upper part, and arms rapidly tapering upward.	
7, 8. Two similar crowns showing variation in shape and condition of partitions.	
Beech River formation; Decatur County, Tennessee.	
<i>Eucalyptocrinus ventricosus</i> Wachsmuth and Springer.....	37
FIGS. 9, 10, 11. Dorsal and two lateral views of typical specimens.	
12. Perfect crown, with nearly complete stem attached.	
Beech River formation; Decatur County, Tennessee.	
<i>Eucalyptocrinus</i> sp.	38
FIG. 13. Part of crown with broadly rounded calyx of very tumid plates.	
Beech River formation; Decatur County, Tennessee.

PLATE 9

(Figures natural size unless otherwise stated)

	PAGE
<i>Eucalyptocrinus pernodosus</i> new species.....	38
FIG. 1. Complete crown with part of stem attached, with extremely nodose plates, and heavy arms between the partitions. Nat. size.	
2, 2a. Lateral and dorsal views of a highly typical calyx, showing the very large radial plates; basals concealed in cavity.	
3, 3a. Similar views of a calyx with plates less prominently projecting.	
4. A smaller calyx with low convex plates.	
Decatur limestone; quarries above Perryville, Tennessee.	
<i>Eucalyptocrinus sculptilis</i> new species.....	38
FIG. 5. Complete crown with stem, showing its peculiar bands. Nat. size.	
6. Calyx of another specimen with very sharp sculpture.	
6a. Basal view of same, showing 4 BB at bottom of deep cavity. $\times 2$.	
Decatur limestone; near Perryville, Tennessee.	
<i>Eucalyptocrinus</i> sp.....	39
FIGS. 7, 7a. Lateral and basal views of perfectly smooth specimen, with narrow base.	
Decatur limestone; near Perryville, Tennessee.	
<i>Callicrinus beachleri</i> Wachsmuth and Springer.....	40
FIGS. 8, 9, 10. Lateral views of three calices showing the short partitions, base of smooth anal tube, and deep sculpturing of calyx plates.	
11, 12. Dorsal views of two specimens showing the 4 very small BB at bottom of cavity, large RR occupying about the entire base projecting in the middle, with sharp ridges following the radial series.	
13. A detached base, showing the 4 BB at bottom of cavity, and ornamentation of RR. $\times 2$.	
Laurel limestone; St. Paul, Indiana.	
<i>Callicrinus</i> sp.....	40
FIG. 14. One of the 4 fan-shaped summit plates surmounting the tube.	
14a. Opposite side of same, with some matrix attached, showing that it is not <i>Petalocrinus</i> .	
15. A similar smaller plate.	
16. An elongate, quadrant-shaped plate of same position.	
Laurel limestone; St. Paul, Indiana.	

K. M. Chapman del.

PLATE 10

(Figures natural size unless otherwise stated)

	PAGE
Periechocrinus tennesseensis (Hall)	45
FIG. 1. Lateral view of calyx of typical specimen with smooth surface; 3 and 4 arms to the ray.	
2. Posterior view of similar specimen.	
3, 4. Lateral and ventral views of smaller specimens, showing the numerous small plates of tegmen.	
Beech River formation; Decatur County, Tennessee, except fig. 4, which is from the Louisville limestone, Jefferson County, Kentucky.	
Periechocrinus sp.	45
FIG. 5. A large specimen with long arms and sharp ridges following the radial series; calyx partly broken away.	
5a. Opposite view of same with more of calyx shown.	
Beech River formation; Decatur County, Tennessee.	
Saccocrinus benedicti S. A. Miller.	46
FIGS. 6, 7, 8, 9. Four specimens with low convex plates; r. post., l. post. iR, l. ant. R, and post. views of calyx; all with 10 arms.	
10, 11, 12. Three calices with nodose plates; r. ant., l. ant., and basal views.	
13. Tegmen of specimen with strong stellate outline.	
14. Specimen with sharp stellate ornament, perhaps of a different species.	
Laurel limestone; St. Paul, Indiana.	
Saccocrinus cuspidatus new species.	46
FIG. 15. Calyx of very large specimen with plates surmounted by sharp cusps.	
Louisville limestone; Jefferson County, Kentucky.	
Aorocrinus nodosus new species.	47
FIGS. 16, 16a, 16b. Posterior, dorsal and ventral views of calyx, with strongly nodose plates, showing anal series leading to opening in tegmen.	
Decatur limestone; below Grandview, Perry County, Tennessee.	
Aorocrinus clarkensis new species.	47
FIGS. 17, 17a. Posterior and dorsal views of calyx, with strongly gibbous plates, 3 small BB, and very large RR.	
Hamilton, Middle Devonian; Clark County, Indiana.

PLATE 11

(Figures natural size unless otherwise stated)

	PAGE
<i>Coccocrinus rosaceus</i> Roemer.....	49, 51
FIGS. 1, 1a, 1b. Lateral, dorsal and ventral views of calyx, having the tegmen covered by 5 abutting orals, with narrow grooves for ambulacra, each underlaid by an interradial "suboral" plate.	
2. Oblique view of another specimen with larger orals and more open grooves than the preceding. (See Wachsmuth and Springer, N. A. Crin. Cam., pl. 3, fig. 14.)
Middle Devonian; Eifel, Germany.	
<i>Culicocrinus nodosus</i> Joh. Müller.....	51
FIG. 3. Lateral view of calyx from l. post. radius, showing long narrow iBr.	
3a. The tegmen entirely filled by orals closely interlocked, without marginal grooves, or iAmb.
After Müller.
Lower Devonian; Grauwacke, Coblenz, Germany.	
<i>Culicocrinus spinosus</i> new species.....	51
FIG. 4. Anterior lateral view of calyx showing iBr plates between the arm-bases.	
4a. Posterior view of same enlarged, showing anal opening with remnants of small plates covering it. $\times 2$.	
4b. Basal view of same.	
4c. Ventral view enlarged, showing 5 large orals abutting without radial grooves between; they follow the iBr in succession, and each is surmounted by a prominent spiniferous node. $\times 2$.
Beech River formation; Decatur County, Tennessee.	
<i>Myrtillocrinus americanus</i> Hall.....	49
FIG. 5. Lateral view of type. After Hall.	
5a. Tegmen, composed entirely of 5 large abutting orals. For comparison of tegmen with preceding.
Onondaga, Middle Devonian; Livingston County, New York.	
<i>Lyonicrinus bacca</i> (Roemer).....	51
FIGS. 6, 7. R. ant. and ant. views of calyx, showing two IBr, with interambulacra rising between them. $\times 3/2$.	
8. Dorsal view of calyx, with small column facet; small basal in l. post. position. $\times 3/2$.	
9. Dorsal view of another specimen, with base exceptionally divided into 5 plates. $\times 3/2$.	
10. Ventral view of specimen with tegmen partly covered by 4 large interambulacral plates not in contact, but separated by open ambulacral furrows leading from the arm-bases, and by a large open space in the middle, beyond which at the posterior side are two smaller interambulacral plates flanking the anal series leading to an opening through the tegmen. $\times 3/2$.	
11. Similar view of a specimen with some side pieces of ambulacra remaining. $\times 3/2$.	
12, 13, 14, 15, 16, 17, 18. A series of similar tegmental views showing the same structures in various positions. $\times 3/2$.	
19, 20. Lateral views from posterior side, showing succession of anal plates flanked by the two posterior interambulacra. $\times 3/2$.	
21, 21a. L. post. and opposite views of specimen with bases of arms and displaced remnants of tegmen plates; position of small basal at left posterior plainly shown. $\times 3/2$.	
22. Oblique view of calyx with uniserial arms attached, two to the ray; r. post. side up. $\times 3/2$.	
23. Part of a set of arms showing nearly their full length, with the pinnules, probably two to each brachial. $\times 3/2$.
Beech River formation; Decatur County, Tennessee.	
<i>Platycrinidae</i> gen. and sp. indet. No. 1.....	51
FIGS. 24, 24a, 24b. Dorsal cup only, with massive plates and large BB; 3 views. $\times 2$.
Beech River formation; Decatur County, Tennessee.	

	PAGE
Platycrinidae gen. and sp. indet. No. 2.....	51
FIG. 25. Calyx, with large interambulacral plates in position. $\times 3/2$. Laurel limestone; St. Paul, Indiana.	
Platycrinidae gen. and sp. indet. No. 3.....	52
FIGS. 26, 27, 28, 29. Lateral, dorsal and ventral views of small species from St. Paul, Indiana.	
Hapalocrinus gracilis new species.....	52
FIG. 30. Complete crown, with heavy cuneate arms and strong pinnules. $\times 2$.	
31. Another specimen with part of arms and stem, showing high interambulacral tegmen plates. $\times 2$.	
32. Small specimen with arms intact. $\times 3/2$. Beech River formation; Decatur County, Tennessee.	

PLATE 12

(Figures natural size unless otherwise stated)

	PAGE
Hapalocrinus cirrifer new species.....	52
FIGS. 1, 2, 3, 4. Four specimens with long and narrow arms, profuse pinnules, and long cirri. In fig. 4 the arms are extremely slender; the structure of the calyx is obscure, there being an abnormally enlarged anal opening. Figs. 1, 2, 4. $\times 3/2$; fig. 3, natural size.	
Hapalocrinus pinnulatus new species.....	52
FIG. 5. Crown with strong arms and pinnules, but much smaller and more conical calyx than the preceding. $\times 2$.	
Hapalocrinus tuberculatus new species.....	53
FIG. 6. Crown with arms and pinnules similar to the preceding, but calyx rotund and with tubercular surface. $\times 2$.	
Hapalocrinus tennesseensis new species.....	53
FIG. 7. Crown much displaced, with uniserial arms, quadrangular brachials, long stem and cirri. $\times 3/2$.
Figs. 1-7. Beech River formation; Decatur County, Tennessee.	
Hapalocrinus devonicus new species.....	53
FIG. 8. Crown with uniserial arms, 3 in some rays, quadrangular brachials, very strong stem and cirri, and tubercular surface.
Helderbergian; Keyser, West Virginia.	
Brahmacrinus elongatus new species.....	53
FIGS. 9, 10. Two calices showing brachial series of first two orders incorporated in calyx by secundibrachs in two ranges, and extremely large BB. $\times 2$.
11, 12. Two detached bases, from side view. $\times 2$.
13, 14, 15. Three bases from dorsal side, showing 3 equal basals, sharply excavate cavity, and minute axial opening. $\times 2$.
Laurel limestone; St. Paul, Indiana.	
Nyctocrinus magnitubus new genus and species.....	54
FIGS. 16, 16a. Complete crown with spiniferous anal tube rising above the branching, biserial arms; views from opposite sides; fig. 16 from post. interradius, the plate being somewhat larger than in other areas.
17. A similar specimen, from r. post. radius.
18, 18a. Opposite views of a larger crown, showing part of tube at the top.
19. A smaller specimen, with base of tube exposed.
20. Dorsal view of detached calyx, showing broad, shallow basal cavity, with 5 IBB at bottom.
21. Another detached base seen from the interior, with infrabasal cone plainly shown.
Beech River formation; Decatur County, Tennessee.	

PLATE 13

(Figures natural size unless otherwise stated)

PAGE
57**Marsipocrinus rosaeformis** (Troost)

- FIGS. 1, 1a, 1b, 1c. A typical hemispheric calyx, with surface characters finely preserved: 1. Lateral view, showing the great prominence of the arched tegmen over the flat dorsal cup; radial facets directed below the horizontal, iBr heptagonal, their distal margin projecting, with a strong lip, and corrugated. 1a. Dorsal side, with extremely sharp sculpturing in form of wrinkles radiating upon the plates and meeting in slightly raised ridges running from the angles, dividing the surface into triangles; very small axillary iBr, the much larger IIBr resting on distal face of RR; low rim around the lumen. 1b. Detail of these structures enlarged, showing more fully the succession of brachials leading to the biserial arm. $\times 2$. 1c. The tegmen, showing complete distribution of orals and other plates strongly nodose; iAmb connecting with the arched iBr and bearing numerous small and distinct tubercles; ambulacra defined but not conspicuous.
- 2, 2a, 2b, 2c. Similar views of a specimen with concavo-convex calyx; radial facets pointing strongly downward; surface ornament of much smaller wrinkles. 2b is enlarged. $\times 2$.
- 3, 3a, 3b, 3c. A smaller specimen with more granular ornamentation; similar views. 3b enlarged. $\times 2$.
- 4, 4a, 4b. Similar and larger specimen with highly sculptured tegmen and base finely granular.
- 5, 5a. A specimen of this type with more convex base than the preceding; iBr extremely prominent.
- 6, 6a, 6b. Specimen with tegmen strongly nodose.
- 7, 7a. Specimen with tegmen plates more angular than nodose.
- 8, 8a. A very mature specimen, with faint granulose-striate markings on dorsal cup, and tegmen plates profusely covered with small nodes; ambulacra inconspicuous.
Beech River formation; Decatur County, Tennessee.

K. M. Chapman del.

PLATE 14

(Figures natural size unless otherwise noted)

	PAGE
Marsipocrinus tennesseensis (Roemer).....	57
(See also pl. 15, figs. 9, 9a)	
FIGS. 1, 1a, 1b. Typical specimen with biconvex calyx, radial facets directed upward, granular surface on cup, tegmen plates rounded or angular; ambulacra well defined.	
2, 2a, 2b. Similar specimen with tegmen plates partly nodose. Original of W. and Spr. N. A. Crin. Cam., lxx, 16.	
3. Lateral view, for profile, of strongly convex calyx.	
4. Tegmen with nodose plates, intermediate from preceding species.	
5, 5a. Large specimen with massive convex tegmen plates and prominent ambulacra.	
6, 6a. Profile and tegmental view of maximum specimen with angular plates. Beech River formation; Decatur County, Tennessee.	
Marsipocrinus striatus Wachsmuth and Springer.....	58
FIGS. 7, 7a, 7b, 7c. Lateral, dorsal and ventral views of specimen with biconvex calyx, striate ornament on dorsal cup, tegmen plates low convex and smooth, ambulacra very distinct; 7c, detail of surface of cup enlarged. $\times 2$.	
8, 8a, 8b. Similar views of specimen with stronger striae.	
9, 9a. Lateral and ventral views of well marked specimen; 9b, oblique view of cup, showing very fine striation, enlarged $\times 2$.	
10, 10a. Lateral and dorsal view, enlarged $\times 2$, of specimen with coarser striae.	
11, 11a. Abnormal specimen, with 2 rays malformed.	
12, 13. Two fragments showing detail of striation on dorsal cup, enlarged $\times 2$. Beech River formation; Decatur County, Tennessee.

PLATE 15

(Figures natural size unless otherwise stated)

	PAGE
<i>Marsipocrinus inflatus</i> (Troost).....	58
- FIGS. 1, 1a, 1b. Lateral, dorsal and ventral views of characteristic calyx, with rather smooth tegmen, high conical and laterally constricted; dorsal surface elaborately sculptured; basal pentagon very large, slightly concave, and notched at the interbasal sutures; iBr not projecting, but bent over into the tegmen.
- 2, 2a, 2b. Similar views of specimen with tegmen plates tubercular and better defined.
3. Specimen with typical contour; tegmen profusely covered with small tubercles.
- 4, 4a, 5, 5a, 5b, 6, 6a. A series of dorsal cups, showing large size of basal pentagon.
- 7, 7a. A similar specimen, original of Wachsmuth and Springer, N. A. Crin. Cam., pl. 75, fig. 17.
8. Detail of tegmen of broken specimen, enlarged, showing course of ambulacra leading to the interbrachial pinnules. Reproduced from paper on *Scyphocrinus*, 1917, pl. 9, fig. 6, as *M. striatus*? $\times 2$.
From limestone beds top of Beech River formation; Decatur County, Tennessee.
<i>Marsipocrinus rosaeformis</i> (Troost).....	57, 59
- FIG. 9. Ventral side of an imbedded specimen, with bases of arms perfectly preserved, and pinnules springing both from the arms and interbrachial margins as continuations of tegmental ambulacra. Posterior is at upper right.
- 9a. Detail from posterior margin of same, enlarged $\times 3$, showing course of ambulacra leading to interbrachial pinnules. From *Scyphocrinus*, 1917, pl. 9, figs. 5a, b; *Dolatocrinus*, 1921, pl. 2, figs. 5, 6, as *M. tennesseensis*.
Beech River formation; Decatur County, Tennessee.

1a

1

1b

2a

2

2b

5a

4

5

5b

6a

4a

6

7

7a

9

8

9a

PLATE 16

(Figures natural size unless otherwise stated)

	PAGE
Marsipocrinus verneuili (Troost).....	59
Figs. 1, 1a, 1b. Typical specimen with striate wrinkles, with shallow depression almost full width of pentagon, bordered by sharp rim: lateral, dorsal and ventral views.	
1c. Oblique view of calyx, enlarged, showing surface ornament in greater detail. $\times 2$.	
2, 2a. Lateral and dorsal views of specimen with striate ornamentation.	
2b. Tegmen of same enlarged, with relatively few and large plates. $\times 2$.	
3. Crown of small specimen with strong rugose ornamentation, and broad, shallow, depressed base; form and proportions of arms well shown.	
3a. More direct view of calyx of same, enlarged, showing sculpturing in greater detail. $\times 2$.	
4. Larger specimen with arms and pinnules.	
5, 5a. Specimen with striate wrinkles, bases of arms coalesced, interrarial spaces depressed, producing stellate outline; basal depression narrow. Dorsal and ventral views.	
6. Similar specimen with arms partly preserved; basal depression narrow.	
Marsipocrinus excavatus new species.....	60
Figs. 7, 7a, 7b. Lateral, dorsal and ventral views of the only specimen; base abruptly depressed.	
Marsipocrinus concavus new species.....	60
FIG. 8. Dorsal cup only; small base at bottom of a broad and deep concavity into which the radials extend far down to narrow faces; strongly wrinkled ornament. By oversight posed with small B at upper right. $\times 2$.	
All Beech River formation. Decatur County, Tennessee.

1c

1b

2b

1a

1

4

2

2a

6

3a

3

5

8

5a

7a

7

7b

PLATE 17

(Figures natural size unless otherwise stated)

	PAGE
Marsipocrinus stellatus (Troost)	60
FIGS. 1, 1a, 1b. Lateral, dorsal and ventral views of specimen with dorsal surface perfectly smooth; numerous large plates in tegmen shown, but ambulacra much eroded; basal depression around facet small.	
2, 2a, 2b. Similar views of specimen with prominent ambulacra well defined; 2a and 2b enlarged $\times 2$.	
3, 3a. Lateral and ventral views of specimen with ambulacra still better defined; 3a enlarged $\times 2$.	
4, 5, 6, 7. Specimens with arms and stem more or less completely preserved.	
8, 9, 10. Tegmens in different conditions of preservation; fig. 10 may not belong here, dorsal side not being exposed. $\times 2$.	
Beech River formation; Decatur County, Tennessee.	
Marsipocrinus turbinatus new species	62
FIGS. 11, 12, 13. Fragmentary bases from St. Paul, Indiana.	
14. Lateral view of basals, showing extremely turbinate form.	
Laurel limestone; St. Paul, Indiana.

PLATE 18

(Figures natural size unless otherwise noted)

PAGE

Marsipocrinus striatissimus new species..... 61

- FIGS. 1, 1a. Dorsal and ventral views of a typical specimen, large and with very thin plates, more or less displaced by pressure; dorsal surface marked by extremely fine striae, and tegmen by numerous small tubercles.
2. Dorsal side of similar specimen not distorted, and with striae more distinct.
3. Specimen with arms and part of stem; striae indistinct.
- 4, 5. Two detached basal pentagons (as usually found) seen from the inner side, showing varying form of lumen.
- 6, 6a, 6b. Lateral, dorsal and ventral views of a well preserved dorsal cup of stronger construction than the preceding, and with striate marking remarkably distinct.
- 7, 7a. Lateral and ventral views of a very robust specimen having the surface covered with small tubercles. Not of this species, but probably a variant of *M. tennesseensis*. Beech River formation; Decatur County, Tennessee.

PLATE 19

(Figures natural size unless otherwise stated)

	PAGE
<i>Marsipocrinus magnificus</i> (Troost).....	61
FIGS. 1. 1a, 1b. Lateral, dorsal and ventral views of type in Troost collection, U. S. National Museum; dorsal cup highly sculptured with coarse rugae and prominent ridges; tegmen plates large and nodose; 4 arms to the ray. Beech River formation; Decatur County, Tennessee.	
<i>Marsipocrinus coelatus</i> (Phillips)	62
FIG. 2. Dorsal view of large specimen with outstretched arms, 4 to the ray.	
3, 3a. Specimen with arms nearly complete, and part of stem.	
4. Small specimen showing contour of calyx, and typical surface ornament; lateral view.	
5. Lateral view of similar specimen with part of arms attached.	
6. A very young specimen, with intense surface marking: gastropod seated over anus.	
6a. Basal view of same enlarged, showing sharp wrinkles vertical to the sutures. $\times 2$.	
7. A cluster of 9 stems springing from a common base on rock or coral, 4 with crown attached; reduced to 1/3. Wenlockian, Silurian; Dudley, England.

1a

1

1b

6

6a

7

3a

4

5

3

2

PLATE 20

(Figures natural size unless otherwise stated)

	PAGE
<i>Lecanocrinus pusillus</i> Hall.....	65
FIGS. 1, 1a. Posterior and basal views of calyx, showing radianal in oblique position, and projecting anal. $\times 2$.	
2. Dorsal view of specimen with abnormal base. $\times 2$.	
3. Ventral view showing distal faces of RR, and great thickness of plates. $\times 2$.	
4. Specimen with very acuminate anal, with part of rays in place. $\times 2$.	
5. Complete crown with part of stem. Coll. New York State Museum. Waldron shale; Waldron, Indiana.	
<i>Lecanocrinus pisiformis</i> (Roemer).....	65
FIGS. 6, 6a, 6b. Anterior, posterior and distal views of complete crown, showing oblique RA, and rapid taper of infolding arms. $\times 2$.	
7. Another complete crown. $\times 2$.	
8, 9. Dorsal views of two specimens, showing IBB, BB, and RA plates. $\times 2$.	
10, 11. Posterior views of two smaller specimens with smaller RA than usual. $\times 2$.	
12. Distal face of RR, showing comparative thinness of calyx wall. $\times 2$. Beech River formation; Decatur County, Tennessee.	
<i>Lecanocrinus elongatus</i> new species.....	65
FIG. 13. Anterior view of complete crown. $\times 2$.	
14. Posterior view of another specimen. $\times 2$. Laurel limestone; St. Paul, Indiana.	
<i>Lecanocrinus meniscus</i> Springer.....	65
FIGS. 15, 15a, 15b. Anterior, posterior and basal views of the type and only specimen. $\times 2$. Beech River formation; Decatur County, Tennessee.	
<i>Anisocrinus greenei</i> (Miller and Gurley).....	66
FIG. 16. A much distorted specimen. Same horizon and locality as last.	
17. The type, from Jefferson County, Kentucky.	
<i>Anisocrinus oswegoensis</i> (Miller and Gurley).....	66
FIG. 18. The type, from Oswego, Illinois.	
<i>Anisocrinus angelini</i> Wachsmuth and Springer.....	66
FIG. 19. The type, from Gotland, Sweden.	
<i>Asaphocrinus bassleri</i> Springer.....	66
FIGS. 20, 20a. Complete crown. Anterior and posterior views of principal type, showing arrangement of arms, anal plates, and massive tube with plates flanking it.	
21. Oblique view of specimen showing the base. Beech River formation; Decatur County, Tennessee.	
(?) <i>Asaphocrinus minor</i> new species.....	67
FIG. 22. Small specimen, with base lacking, perhaps of this genus. $\times 3/2$. Laurel limestone; St. Paul, Indiana. All figures on this plate, except 13, 14 and 22, are taken from the author's work on the Crinoidea Flexibilia, 1920.

PLATE 21

(Figures natural size unless otherwise noted)

	PAGE
<i>Pycnosaccus patei</i> Springer	67
FIG. 1. Complete crown with part of stem; r. ant. view.	
1a. Posterior view of same, showing anal and radianal plates. RA in oblique position.	
1b. An interradius, filled by an integument of small plates. $\times 2$.	
2. Another crown, showing arms more fully, and the interradianal integument.	
2a. The integument enlarged. $\times 2$.	
3, 3a. Anterior and dorsal views of specimen with very sharp ridges.	
4. Very large crown, with distal ends of arms curving down into the interradianal spaces. Beech River formation; Decatur County, Tennessee.	
<i>Pycnosaccus welleri</i> Springer	67
FIG. 5. Flattened crown with long stem; anterior view; has 3 and 4 IBr.	
5a, 5b. L. post. view of crown of same, and restoration in outline of natural contour.	
6. Crown with part of stem, ant. view.	
6a, 6b. Posterior and basal views of same.	
7. Larger crown, flattened, with stem; a small calyx and long arms, 2 and 3 IBB. Beech River formation; Decatur County, Tennessee.	
<i>Pycnosaccus laurelianus</i> new species	67
FIGS. 8, 8a. Anterior and posterior views of complete crown. $\times 2$. Laurel limestone; St. Paul, Indiana.	
(?) <i>Pycnosaccus dubius</i> Springer	68
FIG. 9. Crown with plates displaced, and complete stem tapering to very long and slender columnals.	
9a. Probable arrangement of calyx plates. Beech River formation; Decatur County, Tennessee.	
<i>Clidochirus americanus</i> Springer	69
FIGS. 10, 10a. Anterior and posterior views of crown, showing small RA at foot of r. post. ray.	
11. Another specimen with long stem; r. post. view, showing same small RA. Clinton group; Dayton, Ohio.	
All figures on this plate, except 8 and 8a, are taken from the author's work on the Crinoidea Flexibilia, 1920.

K. M. Chapman del.

PLATE 22

(Figures natural size unless otherwise stated)

	PAGE
Hormocrinus tennesseensis (Worthen).....	66
FIG. 1. Complete crown, with tapering stem formed of bead-like columnals, preserved for full length down to fine distal end.	
2. Crown of small specimen, showing full contour with infolding arms.	
2a. Dorsal view of same enlarged, showing base and arrangement of plates; no RA. $\times 2$.	
2b. Ventral view of same. $\times 2$.	
3, 3a. L. ant. and r. post. views of another complete crown.	
3b. L. ant. view of same enlarged, showing full details of arms with slender distal ends curving down into the interrarial spaces. $\times 2$.	
Beech River formation; Decatur County, Tennessee.	
Sagenocrinus clarki Springer.....	68
FIG. 4. The type. Flattened crown with stem as it lies in the rock; calyx plates much displaced.	
4a. Restoration from accurate measurements showing exact relation of plates; RA touching IBB.	
Beech River formation; Decatur County, Tennessee.	
Sagenocrinus americanus Springer.....	68
FIG. 5. Posterior view of type; RA not touching IBB.	
Waldron shale; Waldron, Indiana.	
Protaxocrinus robustus Springer.....	70
FIGS. 6, 7. Anterior and posterior views of types, showing small plates in interrarial areas RA in primitive position at base of r. post. ray, and large anal tube following.	
Beech River formation; Decatur County, Tennessee.	
Gnorimocrinus cirrifer Springer.....	70
FIG. 8. Complete crown, flattened, with long stem showing whorls of cirri; anterior view.	
9. Anterior view of another perfect crown, showing large anal tube, with RA in oblique position at its base.	
10. Dorsal view of smaller specimen, showing IBB and related plates. $\times 2$.	
Beech River formation; Decatur County, Tennessee.	
Gnorimocrinus varians Springer.....	70
FIGS. 11, 11a. Two views of specimen with strong anal tube.	
Beech River formation; Decatur County, Tennessee.	
Ichthyocrinus subangularis Hall.....	69
FIG. 12. R. post. interrarial view of crown, showing RA in primitive position at base of r. post. ray.	
13. Oblique view of another specimen, showing base with IBB, and small size of RA compared with RR in other rays.	
Waldron shale; Waldron, Indiana.	
All figures on this plate are taken from the author's work on the Crinoidea Flexibilia, 1920.	

PLATE 23

(All figures double size unless otherwise stated)

	PAGE
<i>Pisocrinus gemmiformis</i> S. A. Miller.....	74
FIG. 1. A characteristic specimen, with short arms complete; l. post. view.	
2, 3.	Calyx of mature specimens, radial and ant. views. Note the wide facets, low rectangular processes, convex base with small indented cavity for column facet.
4, 4a.	Basal and ventral views of similar specimen.
4b, 4c.	Details of radial facet of same, seen from side and above. $\times 8$.
5, 5a.	Radial and ventral views of average specimen.
6, 6a.	Similar views of smaller specimen.
7, 7a.	Basal and ventral views of minimum specimen.
8.	Ventral view of small specimen, with oral plates and first arm brachials in place; photograph unretouched.
8a.	The same further enlarged, with structures brought out plainer by the artist. $\times 4$.
All from Laurel formation; St. Paul, Indiana.	
<i>Pisocrinus baccula</i> S. A. Miller.....	75
FIG. 9. Mature specimen with arms complete, ant. view.	
10, 10a.	L. ant. and basal views of mature calyx.
11, 11a, 11b.	Radial, basal and ventral views of large specimen.
11c, 11d.	Details of radial facet and processes of same, seen from side and above. $\times 8$. Note smaller facets, large lance-head processes, broad, concave base and cavity.
12, 12a, 13.	Ant. and ventral views of similar mature specimens.
14, 14a, 14b.	Post., basal and ventral views of smaller specimen.
15, 15a, 15b.	L. ant., basal and ventral views of minimum specimen.
Laurel formation; St. Paul, Indiana.	
<i>Pisocrinus quinquelobus</i> Bather.....	77
FIG. 16. A typical specimen, with complete arms, and long stem; l. post. view.	
17, 18, 19.	L. ant., r. ant. and r. post. views of a series of nearly complete crowns, showing the general character of arms, facets and processes.
20.	One of the types of <i>P. milliganae</i> M. and G., r. post. view.
21, 22, 22a, 22b.	L. post., l. ant., dorsal and ventral views of calyx of mature specimens.
22c, 22d.	Details of radial facets and processes. $\times 4$. Note the relatively small facets, large processes, truncate base, minute basals hidden in deeply indented cavity—all with the strongly lobate form.
23.	Ant. view of mature specimen with unusually wide facets.
24, 24a, 24b.	R. ant., dorsal and ventral views of mature specimens.
25.	Dorsal view of smaller specimen.
26, 26a.	R. post. and ventral views of very small specimen.
27, 28.	Post. and ventral views of large specimen from Jack's Branch, Perry County, Tennessee, probably Lobelville formation.
29.	A calyx vertically fractured, showing in section the size of the minute basals.
All except Nos. 27, 28 from Beech River formation; Decatur County, Tennessee.	
<i>Pisocrinus sphericus</i> Rowley.....	79
FIG. 30. Complete crown, with very short arms. Decatur County.	
31, 31a, 31b.	Ant., dorsal, and ventral views of mature calyx from Rise Mill.
32, 33.	L. post. and ventral views of similar specimens, from Rise Mill.
34, 34a.	Similar specimen. R. ant. and ventral views, from Rise Mill.
35, 35a.	Radial and dorsal views of specimen from Flatwoods.
36.	Smaller specimen, but with perfectly round calyx. Flatwoods.
Top of Beech River; Rise Mill, Perry County; also Decatur and Wayne counties, Tennessee.	
<i>Pisocrinus granulatus</i> Rowley.....	75
FIGS. 37, 38. Ant. and ventral views of large calyx, with tubercular surface.	
39, 39a.	R. post. and basal views of smaller specimen. Note the good-sized basals combined with the rounded form and other characters like the preceding species.
In red shales, Dixon, below Beech River formation; Decatur County, Tennessee.	

	PAGE
<i>Pisocrinus gorbyi</i> S. A. Miller.....	78
FIGS. 40, 40a. Lobed and tubercular specimen, ant. and ventral views.	
41, 41a, 42. R. ant., ventral and basal views of similar specimens.	
Lobelville formation, above Beech River, Flatwoods, Tennessee.	
43, 44, 45. Ventral views of non-tubercular specimens from Rise Mill, Perry County, Tennessee.	
<i>Pisocrinus</i>	72
FIG. 46. Generic diagram.	

PLATE 24

(All figures double size unless otherwise stated)

	PAGE
<i>Pisocrinus tennesseensis</i> (Roemer).....	79
Figs. 1, 1a, 1b. Calyx of characteristic and very mature specimens; ant. and ventral views.	
2, 2a. R. ant. and ventral views of similar specimen.	
2b, 2c. Details of radial facet and processes of same. $\times 4$.	
3, 3a. Radial and ventral views of smaller specimen. Note the very wide facets and dove-tailed processes in all these; with basals limited to the column facet.	
4, 4a. L. post. and ventral views of smaller specimen, from Martin's Mill, Wayne County, Tennessee.	
5. Post. view of very mature specimen with unusual raised border of radials. Beech River formation; all except No. 4 from Decatur County, Tennessee.	
<i>Pisocrinus campana</i> S. A. Miller.....	76
Figs. 6, 7, 8. A series of crowns with very long arms, round calyx and high base, in contrast to <i>P. quinquelobus</i> . Fig. 8 shows ventral side of arms, with sockets of side-pieces of ambulacra.	
9. A set of long arms, with ventral furrow and traces of side-pieces, detached just above the calyx at the sutures, showing some of the brachial articulations.	
9a. Transverse view of proximal articular face of the arms, further enlarged; section of anal tube plainly seen in the middle. $\times 4$.	
10. Dorsal view of calyx from Rise Mill, Perry County, Tennessee, showing large base with abruptly indented column facet.	
11, 11a. R. ant. and ventral views of an average calyx with wide facets and nearly rectangular processes, tending to expand slightly inward; same locality as last.	
11b, 11c. Details of radial facets and processes of same, seen from side and above. $\times 6$.	
12, 12a, 12b. Ant., dorsal and ventral views of more elongate specimen; same locality.	
13, 14. L. ant. and ventral views of two other specimens from same locality.	
15. R. ant. view of similar specimen from Martin's Mill, Wayne County, Tennessee.	
16, 16a, 17. L. ant., ventral and ant. views of somewhat more globose specimens from Flatwoods, Tennessee.	
18. A much larger specimen, radial view, from Sinking Creek, Wayne County, Tennessee.	
19. Elongate specimen from Decatur County, Tennessee, art. view.	
20. Complete crown of elongate specimen, with very long arms and part of stem; l. ant. view.	
21, 21a. Extremely elongate form, r. ant. and ventral views; both the last from Decatur County, Tennessee.	
22, 22a, 23. Specimens from the typical area at Anderson, Marion Co., Indiana, about the usual size; r. ant. dorsal and post. views.	
24, 25, 26. Three abnormal specimens from the type locality at Wabash, Indiana; small RR connecting with BB; l. ant., r. post. and l. ant. views.	
27. The type specimen from Wabash, Indiana, formerly in the collection of A. C. Benedict; larger than average from typical area. All from Beech River formation, except Nos. 22 to 27, which are from the Niagaran of Northern Indiana.	
<i>Pisocrinus benedieti</i> S. A. Miller.....	77
Figs. 28, 29. Low and elongate forms from Anderson, Indiana, with angular processes; r. post. and ant. views.	
30, 30a. Large specimen from St. Paul, Indiana; radial and ventral views.	
31. Low specimen from Decatur County, Tennessee.	
32, 32a, 32b. Large oblate form from Perry County, Tennessee; radial, dorsal and ventral views.	
33. Similar form with long arms, incomplete, from same locality.	
34, 34a. Smaller low, rounded form from Decatur County, l. ant. and dorsal views.	
35, 36. L. ant. and ventral views of similar specimens from Flatwoods, Wayne County, Tennessee. Laurel and Racine formations, Indiana; Brownsport, Tennessee.

	PAGE
Pisocrinus pyriformis Ringueberg	80
- FIG. 37. The type, formerly in the collection of Dr. Ringueberg; l. ant. view. Inserted for comparison with *P. campana*.
Clinton formation; Lockport, New York.
Pisocrinus globosus Ringueberg	80
- FIG. 38. The type. Ant. view, for comparison with *P. benedicti*.
Same horizon and locality.

PLATE 25

(All figures twice natural size unless otherwise stated)

(Foreign species introduced for comparison)

PAGE

- Pisocrinus pilula* DeKoninck..... 80
- FIG. 1. Complete crown with stem, showing the extremely long arms. After Bather. Gotland.
 2, 2a, 2b. Conical specimen from Gotland; radial, basal and ventral views.
 3, 3a, 3b. A maximum high specimen from Dudley, England; ant., basal and ventral views.
 4, 4a, 4b. A smaller, more globose specimen from Dudley; l. ant., l. post. and ventral views; processes tending to lance-head form.
 5, 5a, 5b. A more depressed form from Dudley; r. post., basal and ventral views.
 6. A typical form with almost complete stem; first brachials in place enclosing the orals; ant. view. Dudley.
 6a. Ventral view of same showing the 5 orals, slightly fractured at the center. $\times 8$.
 6r. Ventral view of an average specimen from Gotland. After Bather.
 Wenlockian, Silurian; Gotland and England.
- Pisocrinus ollula* Angelin..... 80
- FIG. 7. L. ant. view of specimen from Gotland. After Bather.
 8. Specimen from Gotland, showing BB limited to column facet. After Bather.
 9. Ventral view of specimen showing wide facets, and processes widening inward. Gotland. After Bather from Angelin.
 10. Basal view of another specimen from Gotland, with extremely small BB at bottom of deep cavity and tendency to pentagonal outline; for comparison with *P. quinquelobus*.
- Pisocrinus pocillum* Angelin..... 80
- FIGS. 11, 11a. The type, from Gotland, post. and l. ant. views. After Bather from Angelin.
- Pisocrinus* sp. 80
- FIGS. 12, 12a. Abnormal specimen from Dudley, with processes bordered by a raised rim, and of a very different shape from those of other forms; lateral and ventral views.
- Pisocrinus* sp. 80
- FIGS. 13, 13a. A very small specimen from Gotland, having the form and structure of *P. gemmiformis*.
 13b, 13c. Details of radial facets and processes of same. $\times 8$.
 14. A larger specimen of similar form from Dudley, England.
 All from the Wenlockian formation, Silurian.
 All the specimens except those of figs. 1, 7, 8, 9, 11, are in the author's collection, U. S. National Museum.
- Triacrinus altus* (Müller)..... 74
- FIGS. 15, 15a, 15b. Characteristic specimen from the Middle Devonian, Eifel, Germany; r. post., dorsal and ventral views. Note the 3 BB, and facets and processes like *P. tennesseensis*.
- Triacrinus depressus* (Müller)..... 74
- FIGS. 16, 16a, 16b. L. ant. dorsal and ventral views of a typical specimen with 3 BB. Note the wholly different form of processes.
 17, 17a, 17b. Ant., dorsal and ventral views of specimen with 4 BB.
 18, 18a, 18b. Similar views of specimen with 5 BB.
 Middle Devonian; Eifel, Germany.

- 82
- Zophocrinus howardi* S. A. Miller.....
- FIGS. 19, 20. Posterior and anterior views of two specimens with the marginal ring of brachials and orals in place. $\times 4$.
21. Ventral view of another specimen in which the spade-shaped oral plates are very distinct, but somewhat frayed and displaced by pressure. $\times 4$.
- 21a. The same tegmen further enlarged, showing fossae on articulating face of some brachials. $\times 8$.
22. Complete view of tegmen, composed from two other specimens. Shows the 4 radii, and division of tegmen into 5 oral plates interlocking at the center, and extending to the periphery, with marginal plates between in 5 groups of 3 each. $\times 6$.
23. Base of another specimen, showing 3 unequal plates.
24. Calyx of a maximum specimen.
25. Specimen from Greenburg, Indiana.
26. Specimen from Flatwoods, Tennessee.
- Laurel formation, Niagaran; all from St. Paul, Indiana, except Nos. 25 and 26.
- 81
- Tiaracrinus quadrifrons* Schultze.....
- FIG. 27. Post. view of type, with the numerous marginal brachials bordering the tegmen in place, showing the same essential structures as *Zophocrinus*, but with a minute base.
- 27a. Basal view of same, showing the 4 unequal radii, and 3 minute BB.
- 27b. The same view, further enlarged. $\times 6$.
- 27c. The tegmen. Five interlocking oral plates meeting in the center, with narrower extensions to the outer margin, and some smaller plates inserted at the exterior angles; the posterior oral wedged in between the other 4. Surrounding these is the border of narrow marginal brachials in 5 groups of 11 or 12 each, instead of 3 as in *Zophocrinus*, separated by the extensions from the oral plates. $\times 4$.
- Middle Devonian; Eifel, Germany.
- 50, 81
- Haplocrinus mespiliformis* (Goldfuss)
- FIG. 28. Ventral view of specimen showing the interlocking orals and radial grooves for the reception of the arms, as usually seen. $\times 5$.
29. A specimen with arms preserved to 3 brachials fitting closely in the grooves between the orals. $\times 5$.
- 29a. Ventral view of same, showing the open grooves in 2 radii. $\times 5$. Figured here for comparison.
- Middle Devonian; Eifel, Germany.

PLATE 26

(Figures natural size unless otherwise stated)

	PAGE
<i>Mysticocrinus wilsoni</i> Springer.....	84
FIGS. 1a-f. Seven enlarged photographic views of the type specimen, with the very short arms preserved in 3 rays folding closely over the tegmen, showing the acuminate anal plate and the arrow-head processes between the arm-bases: 1, left anterior; 1a, left posterior; 1b, posterior; 1c, right posterior, showing the radianal; 1d, anterior; 1e, dorsal, showing the 3 BB; 1f, ventral, showing infolding of the arms. All $\times 4$	
1g. Diagram showing distribution of all the plates; missing brachials indicated by diagonal hachure. Laurel limestone; St. Paul, Indiana.	
<i>Thalamocrinus ovatus</i> Miller and Gurley.....	131
FIGS. 2, 2a, 2b. Posterior, dorsal and ventral views of large specimen; large IBB, RA in oblique position, anal x in line with RR, small column-facet, and ventral cavity.	
3, 3a, 4. Ant. iR and ventral views of similar specimen, and ant. R view of another more elongate.	
5. Specimen with long arms and extremely slender elongate brachials, displaced. Beech River formation; Decatur County, Tennessee.	
<i>Thalamocrinus cylindricus</i> Miller and Gurley.....	132
FIG. 6. Post. view of unusually large specimen.	
7, 7a, 7b. R. post., basal and ventral views of a typical specimen, with RR, anal and radianal plates in place.	
8, 8a. R. post. and ventral views of short specimen.	
9. The type; extremely elongate calyx, lacking the RR anal plates. Beech River formation; Decatur County, Tennessee.	
<i>Thalamocrinus globosus</i> new species.....	132
FIGS. 10, 10a, 10b. R. post., dorsal and ventral views of the only specimen. Beech River formation; Decatur County, Tennessee.	
<i>Thalamocrinus elongatus</i> new species.....	132
FIG. 11. Extremely large and elongate Devonian derivative, lacking the RR and anal plates; r. post. view; differs from all the others in the wide stem, covering the entire base. Linden beds, Helderbergian; near Camden, Benton County, Tennessee.	
<i>Petalocrinus mirabilis</i> Weller.....	126
FIG. 12. A set of separate arm-fans as usually found, artificially mounted; seen from the ventral side, showing 14 to 16 grooves.	
13. Similar set of arm-fans from dorsal side, showing complete fusion of arms in each radius into a rigid plate.	
14. A complete crown with calyx and arms in place, seen from the dorsal side, but the fused radii are broken away, exposing the inner floor of the ventral side, with the grooves indicated by ridges, in which covering plates are shown in some places. Coll. University of Iowa. $\times 2$. Hopkinton stage of Niagaran; Jones County, Iowa.	
<i>Petalocrinus inferior</i> Bather.....	127
FIG. 15. A nearly complete arm-fan, from ventral side, showing 28 grooves symmetrically divided in halves; angle of fan 70° . $\times 3/2$.	
15a. Dorsal side of same, fused to a rigid plate. $\times 3/2$.	
16. Another arm-fan with 30 grooves and angle of 65° ; small part missing at proximal end. Laurel formation; St. Paul, Indiana.

K. M. Chapman del.

- PAGE
- Crotalocrinus cora* (Hall)..... 129
- FIG. 17. Cast from an external mould showing surface markings, 3 ranges of calyx plates, and brachials within the radial facet.
Racine dolomite; Chicago, Illinois.
- Crotalocrinus rugosus* (J. S. Miller)..... 128, 129
- FIG. 18. Calyx of a small specimen from Dudley, England; from anal side, showing characteristic sculpture.
19. Detail of specimen from Gotland, showing arrangement of lower brachials at dorsal side.
× 2.
20. Detail of ventral side, showing covering plates of ambulacra, and the mode of connection of arm-branches by small lateral processes with larger spaces between, forming a net-work admitting flexibility of the leaf-like radii. × 3. (Figures after Wachsmuth and Springer.)
Wenlockian; England and Sweden.
- Crotalocrinus pulcher* (Hisinger)..... 128, 130
- FIG. 21. A nearly complete crown, with calyx injured by erosion, showing the radii in form of reticulate leaves which overlap each other. For comparison with the analogous rigid structures of *Petalocrinus*.
Wenlockian; Gotland, Sweden.

PLATE 27

(All figures natural size unless otherwise stated)

	PAGE
<i>Myelodactylus ammonis</i> Bather.....	86
FIG. 1. A close coil, with stem tapering to a point; cuneate interlocking columnals, with a cirrus from the broad end of each, converging to center completely enveloping the crown. $\times 3/2$.	
1a. The same seen from outer curve, showing the rapid taper of stem to narrow distal end; longitudinal sutures not visible. $\times 3/2$.	
2. A similar specimen natural size, with hour-glass shaped columnals bearing a cirrus at each end, and lenticular, non-cirriferous ossicles interposed.	
2a. The same from outer curve tapering to a point; longitudinal sutures strongly developed. $\times 3/2$.	
3. Similar specimen with alternate cuneate columnals. $\times 3/2$.	
3a. The tapering distal end. $\times 3/2$.	
3b. Form of the cuneate columnals in outline.	
4. A small specimen of similar type to last. $\times 3/2$.	
5. Fractured specimen having paired cirri on alternate columnals. $\times 3/2$.	
5a. Inner view of fragment showing form and arrangement of columnals for the paired cirri, with lenticular ossicles interposed. $\times 3/2$.	
Beech River formation; Decatur County, Tennessee.	
<i>Myelodactylus convolutus</i> Hall.....	86
FIG. 6. Specimen from St. Paul, Indiana, with close coil and beginning of broad curve of stem; cirri not preserved, were paired at each end of successive columnals, as shown by the facets.	
7. Specimen from the typical locality, Lockport, New York, showing the cirri in place, one to each successive columnal.	
8. Fragment from inner side of curve, showing form and arrangement of paired cirri at each end of successive columnals.	
Laurel and Rochester formations.	
<i>Myelodactylus brevis</i> Springer.....	86
FIG. 9. A nearly complete coil tapering to distal end, with paired cirri as in preceding species. $\times 3/2$.	
9a. Distal end of same, seen from outer curve. $\times 3/2$.	
Beech River formation; Decatur County, Tennessee.	
<i>Myelodactylus fletcheri</i> (Salter).....	86
FIG. 10. Specimen in author's collection with the crown in place, which is detachable for inspection at all sides, showing that it has only 4 radii. $\times 2$.	
10a, b. Posterior and anterior views of crown; anal tube in <i>a</i> resting upon left shoulder of r. post. Rs, indicating its Heterocrinid affinities. $\times 6$.	
Wenlockian; Dudley, England.	
<i>Myelodactylus extensus</i> Springer.....	87
FIG. 11. Large specimen with stem extended about 8 cm. beyond the close coil, nearly straight, not terminating in a point; cirri on alternating cuneate columnals.	
12. Specimen with about 5 cm. of stem beyond deviation from close coil, and proximal reverse curve in view; has paired cirri on hour-glass-shaped columnals.	
13. Another specimen with stem extension of about 6.5 cm., tapering near the end, and perhaps terminating in small roots; columnals as in preceding specimen.	
14. Large specimen with unusually small close coil, and loose coil partially restored, extended for about 12 cm. not yet terminated; cirri paired on hour-glass columnals.	
15. Smaller specimen with much of extended stem broken off; crown is seen following reversed curve, calyx indistinct; cirri alternating on cuneate columnals. $\times 3/2$.	
16. Large specimen with incomplete stem extension about 7 cm.; polished section obtained by grinding down to the axial canal, showing the proximal coil, neck, and crown in outline, with scattered cirri in some places; probably has less than 5 rays.	
17. Specimen with only the close coil remaining, having the cirri complete converging in the center.	
18. Specimen from the Wenlockian at Dudley, England, with paired cirri in place at the close coil, and stem extended far beyond that region not yet diminishing distalwards.	
All except fig. 18 from Beech River formation, Decatur County, Tennessee.	

- PAGE
87
- Myelodactylus brachiatus* Hall.....
- FIG. 19. Complete stem with crown preserved, with long, slender proximal neck following the reverse curve outside of principal coil; has branching cirri, limited to distal end. $\times 3/2$.
Rochester shale; Lockport, New York.
- Myelodactylus keyserensis* Springer..... 87
- FIG. 20. Nearly complete specimen, with crown fully exposed by removal of cirri; seen from anterior radius, with unbranched posterior radius to the left; cirri paired on successive columnals.
- 20a. Calyx and lower part of arms, from left anterior radius. $\times 2$.
Helderbergian, Keyser formation; Keyser, West Virginia.
- Myelodactylus schucherti* Springer..... 87
- FIG. 21. The type, with both distal and proximal portions broken off, leaving one coil of the crescentic region, followed by the reverse curve and very robust proximal neck which has alternate lenticular ossicles; cirri in crescentic region paired on successive columnals.
Helderbergian, Linden formation; Benton County, Tennessee.
- Myelodactylus nodosarius* (Hall)..... 87
- FIG. 22. One of Hall's types, showing great size of the ponderous cirri compared with smaller stem, and bulbous terminal at distal end. Coll. New York State Museum, Albany.
23. Distal end of stem, showing bulb, cirri, and relative size of small first cirrals. Same collection.
24. Inner side of same structures in another specimen. Coll. Yale University Museum.
Helderbergian, New Scotland formation; Schoharie County, New York.
- All the figures on this plate are taken from the author's work on "Unusual Forms of Crinoids," Proc. U. S. National Museum, vol. 67, 1926, plates 1 to 6.

PLATE 28

(Figures natural size unless otherwise stated)

	PAGE
<i>Cremacrinus ulrichi</i> new species.....	105
FIG. 1. Complete crown of maximum specimen, with nearly entire stem, in a partially erect position; anterior view, the arms somewhat displaced; r. ant. arm smaller than ant., the two rami of the latter occupying at least four fifths of the space on the exposed side.	
2. Another crown, ant. view, showing the dwarfed r. ant. ramus rather better than the preceding.	
Beech River formation; Decatur County, Tennessee.	
<i>Cremacrinus tubuliferus</i> new species.....	106
FIG. 3. Complete crown seen from the median arm transversely flattened, but free on both sides; showing quadrangular inferradial.	
3a. L. post. view of same, enlarged, showing median arm with 2 brachials beyond IBr; the large curved anal tube free from the arms lying to the left of stem; l. post. arm dividing into 2 symmetrical rami each with 8 bifurcations, and 9 long ramules. $\times 2$.	
3b. Anterior view of same, showing ant. arm with about 6 visible branches to each ramus; r. ant. arm dwarfed to about one fourth the size of ant., with 3 short remnants of ramules on one side. $\times 2$.	
4. Median view of another complete crown free on both sides.	
4a. L. post. view of same, showing arm with 8 and 9 long ramules to each ramus, and other features similar to 3a. $\times 2$.	
4b. Ant. view of same, showing by the diminutive branches lying against the tube all that is left of the dwarfed r. ant. arm—less than one fourth the size of ant. $\times 2$.	
5. Complete crown in recumbent position upon long stem, with tube bent like a bow; from ant. side, showing the two unequal arms. $\times 3/2$.	
5a. Anterior view of calyx, showing segments of l. ant. R. $\times 3/2$.	
6. Another crown similar to last, with peculiar hypertrophy of ramus of ant. arm. $\times 2$.	
Beech River formation; Decatur County, Tennessee.	
<i>Cremacrinus decatur</i> new species.....	107
FIG. 7. Anterior view of the only specimen, a nearly complete crown, showing the very stout median arm, and diminished r. ant. arm. $\times 3/2$.	
Decatur limestone; near Rise Mill, Perry County, Tennessee.	
<i>Cremacrinus simplex</i> new species.....	107
FIG. 8. The only specimen; l. post. view of crown, showing the ray dividing symmetrically, with further bifurcations giving 6 branches in all, more or less dichotomous; stem at right of anal tube. $\times 2$.	
8a. Anterior view, showing two arms, ant. somewhat dwarfed. $\times 2$.	
Beech River formation; Decatur County, Tennessee.	
<i>Cremacrinus articulatus</i> (Billings).....	108
FIG. 9. L. post. view of crown, showing the long median arm in profile, bifurcating near the end; the two rami of l. post. arm with their 7 or 8 zig-zag articulations, bearing long ramules.	
10. A complete crown in closed position on stem; ant. view showing long median arm in profile, the 2 rami of ant. arm fully developed about as long as median arm, and r. ant. arm reduced to less than half its size.	
11. A smaller crown, for comparison of arms and position of stem.	
12. Basal view of another calyx, showing hinge, and the 2 connected segments of l. ant. R.	
13. Large specimen from Woodford County, Kentucky, anterior view.	
Trenton group, Ordovician; all except 13 from Kirkfield, Ontario.	

- PAGE
- Cremacrinus kentuckiensis* (Miller and Gurley)..... 109
- FIG. 14. L. ant. view of crown, showing long median arm bifurcating on 14th brachial.
- 14a. Posterior view of same, turned to bring stem to middle, which is the plane of bilateral symmetry; tube projecting at distal end.
15. Post. view of another crown, showing anal tube in middle with stem to the right; one ramus of diminished r. ant. arm seen at right, to be compared with ramus of l. post. arm at left.
- Trenton group, Ordovician; Mercer County, Kentucky.
- Cremacrinus punctatus* Ulrich..... 110
- FIG. 16. The type specimen; l. post. view of crown.
- 16a. Ant. view of same; lower part of r. ant. arm broken away but some distal branches remaining.
17. R. post. view of a typical specimen with the hinged base closed, showing anal plates and eccentric position of stem to right of them; 4th (r. ant.) arm in place; superradials and inferradials of r. post. and r. ant. arms, with edges of the former deeply indented by pressure of stem; also the 4 unfused basals. $\times 2$.
18. Ant. view of similar specimen with the 4th arm (r. ant.) in view alongside of ant. arm. $\times 2$.
- 18a. R. post. view of same. $\times 3/2$.
19. Similar view of better preserved specimen, showing more fully the two lateral arms with their bifurcations, and the median arm in profile at the right. $\times 2$.
20. Calyx of another specimen seen from the hinge, with its widely gaping fissure lodging numerous "supplementary" plates, and its denticulate margin; also the 4 unfused basals, and the elongate inferradial.
- 20a. The same enlarged, showing the hinge in greater detail. $\times 2$.
- Black River (Decorah), Ordovician; Minneapolis, Minn.
- Calceocrinus bassleri* new species..... 117
- FIG. 21. A complete crown in partially erect position, with nearly entire stem attached; median and adjoining arms shown to full length, with large ramules continuous with the outer Betabrachs forming the most conspicuous part, while the main element of the axil-arms is visible in some places; hinge between radials and basals well shown.
- Beech River formation; Decatur County, Tennessee.

PLATE 29

(All figures natural size unless otherwise stated)

	PAGE
<i>Eucheirocrinus chrysalis</i> (Hall).....	112
FIGS. 1, 1a, 1b, 1c. Ant., l. ant., post. and basal views of the type specimen, showing the single lateral arm, narrow tapering inferradial, and large size of median arm.	
2. Basal view of calyx of another specimen, reversed; to show the 4 unfused basal plates. $\times 3/2$.	
3. Complete crown, showing arms to full length, recumbent on stem. Type of <i>Proclivocrinus radiculatus</i> Ring.
Rochester shale; Lockport, New York.	
<i>Eucheirocrinus indianensis</i> (S. A. Miller).....	114
FIGS. 4, 4a, 4b. L. post., ant. and basal views of type. $\times 2$.
Laurel limestone; St. Paul, Indiana.	
<i>Eucheirocrinus minor</i> new species.....	112
FIG. 5. Complete crown, recumbent, with stem to about its full length; l. post. view, showing great length of brachials in median arm, and dichotomous branching of lateral arm. $\times 5/3$.	
5a. Lower part of same from anterior side, showing base of arms. $\times 5/3$.	
5b. L. ant. view of calyx, showing connection of segments. $\times 5/3$.
Beech River formation; Decatur County, Tennessee.	
<i>Eucheirocrinus anglicus</i> new species.....	112, 113
FIG. 6. Complete crown from l. post. side, showing ponderous character of arms, and part of branchings of median arm. $\times 2$.	
6a. Shaded diagram from l. ant. view, showing calyx and complete bifurcation of median arm. $\times 3/2$. See also text-fig. 3c.
Wenlockian, Silurian; Dudley, England.	
<i>Calceocrinus foersteri</i> new species.....	116
FIGS. 7, 7a, 7b. L. post., l. ant., and post. views of crown, the two segments of l. ant. R. connected, inferradial elongate quadrilateral; l. ant. arm complete, unbranched; main branch of axil-arms entirely concealed by large ramules from outer Betabrachs; anal plates much foreshortened owing to curvature; 7b shows several plates of anal tube projecting beyond arms.	
8, 8a, 8b. Similar views of smaller specimen, also showing anal tube.	
9. Complete crown with stem, in partially erect position; tube projecting.	
10. A distorted crown, probably of this species.	
11. Set of arms of unusual size, with 7 axil-arms, probably of this species; to be compared with <i>C. tenax</i> Bather, of Gotland.
<i>Eucalyptocrinus</i> zone, Beech River formation; Decatur County, Tennessee.	
<i>Calceocrinus bassleri</i> new species.....	117
FIGS. 12, 12a, 12b. L. post., l. ant. and post. views of crown, showing the two triangular segments of l. ant. R. separated by the large RR meeting between them; l. ant. arm simple; main branch of axil-arms partly exposed at inner side of large ramule; anal tube projecting beyond the arms.	
12c. The hinge, showing structures when base is closed upon radials.	
13. A large crown with calyx plates slightly displaced, giving l. post. view of arms with l. ant. exposure of calyx and large median arm; main branch of axil-arms well exposed at inner side of large ramules, showing two more bifurcations.	
13a. Another view of hinge.	
14. L. post. view of still larger crown with part of stem, showing very short columnals; main branch of axil-arms nodose and further bifurcating. In this specimen the two segments of l. ant. R. almost connect by their points. (See also pl. 28, fig. 21.)
Horizon and locality, same as last.	

- PAGE
- 117
- Calceocrinus bifurcatus* new species.....
- FIG. 15. Crown in completely recumbent position, closed over the stem; main branch of axil-arms folded under, and only the large outer ramule visible.
- 15a. L. ant. view of same, showing rather slender median arm, bifurcating on 4th brachial, and 2 or 3 times beyond to at least 8 finials; segments of l. ant. R separated.
- 15b. Post. view, showing relation of stem to anal plates, with deep indentation; tube projecting. *Coccoocrinus* zone of Beech River formation; Decatur County, Tennessee.
- Calceocrinus stigmatus* Hall..... 118
- FIG. 16. L. ant. view of type. After Hall. Am. Mus. Nat. Hist.
Waldron shale, Niagaran; Waldron, Indiana.
- Calceocrinus typus* Ringueberg..... 118
- FIG. 17. Ringueberg's type specimen; ant. view showing large median arm in profile, and the axil-arm system well developed.
- 17b. Base of same.
- Calceocrinus* Hall 118
- FIG. 18. The original figure. After Hall.
- Calceocrinus halli* Ringueberg 119
- FIG. 19. Ringueberg's type; consolidated left basal.
Rochester shale; Lockport, New York.
- Calceocrinus nitidus* Bather..... 119
- FIG. 20. Complete crown recumbent on stem, l. post. view; showing extreme length of median arm; main branch of axil-arms hidden by outer ramules; only one primibrach in lateral arms.
Wenlockian, Silurian; Dudley, England.
- Calceocrinus pugil* Bather..... 120
- FIG. 21. Crown with short median arm, l. post. view; main branches of axil-arms unusually developed, larger than the outer ramules; axillaries strongly nodose.
Wenlockian, Silurian; Gotland, Sweden.
- Halysiocrinus keyserensis* new species..... 121
- FIG. 22. Ant. view of crown, showing median arm in profile, and axil-arms with main branches visible at inner side.
Helderbergian, Lower Devonian; Keyser, West Virginia.
- Halysiocrinus marylandensis* (Ohern)..... 122
- FIGS. 23, 23a, 23b. L. post., l. ant. and post. views of crown; showing large median arm, segments of R separated, inferradials of vanished arms meeting under anal x, main branches of axil-arms more or less visible. Coll. Geol. Surv. Maryland.
Oriskany sandstone, Lower Devonian; Cumberland, Md.
- Halysiocrinus carinatus* new species..... 122
- FIGS. 24, 24a. L. ant. and post. views of crown; showing small size of keeled median arm, and sharply angular anal tube deeply indented in lower part by pressure of stem. Alpena, Michigan.
25. Calyx of another specimen; l. ant. view showing hinge, and separated segments. New Buffalo, Iowa.
26. Another specimen probably of this species from near Louisville; l. ant. view. Clark County, Indiana.
Hamilton group, Middle Devonian; Michigan, Iowa and Indiana.

PLATE 30

(Figures natural size unless otherwise stated)

PAGE

Halysiocrinus dactylus (Hall)..... 123

- FIG. 1. The type specimen, anterior view. Mus. Comp. Zool., Harvard.
2. Ant. view of free crown in author's collection, with main ramus of axil-arms branching from nodose axillaries 2 and 3 times at intervals of 3 and 4 brachials; ant. R. extending far around towards anal tube.
- 2a. L. ant. view of same, showing median arm branching twice, proportions of lateral radials and separated segments of l. ant. R.
- 2b. Post. view of same; hinged base closed against RR; inferradials of vanished arms interposed between large RR, and meeting below anal α , forming part of base of tube.
- 3, 3a. A very young specimen probably of this species; crown with complete stem, 2 opposite views, enlarged; shows calyx higher than wide, narrowing downwards; l. post. view shows only one axil-arm and anal tube, and ant. view shows 2 arms and tube. $\times 2$. Upper Burlington limestone; Burlington, Iowa.

Halysiocrinus perplexus (Shumard)..... 123

- FIGS. 4, 5. L. ant. views of calyx, showing the hinge between radials and basals in position as they would be when fully extended; composed from weathered specimens with the two elements separate; markings of muscular attachment seen in the fissure, and different tubercular ornament upon the surface.
6. Another specimen similarly composed, seen from the interior, showing the same structures from the opposite side; marks of muscular attachment at inner surface of RR and BB; edges of opposing plates rounded; articular faces of median and lateral arms are shown, and the cavity which lodged the viscera.
7. Post. view of specimen with BB partly closed, their edges rounded, and showing arrangement of RR from the interior.
- 8, 9, 10. A series of radials to show the change in form with age from elongate with nearly parallel sides to relatively short and widely spreading downwards.
- 11, 12, 13. A series of bases seen from the interior. New Providence shale, basal Mississippian; Button-mould Knob, Kentucky.

Halysiocrinus nodosus (Hall)..... 124

- FIG. 14. Post. view of calyx, with basals closed against the lateral radials which have extended around to the anal series; inferradials of vanished arms now meeting between them and below anal α , and indented by stem.
15. Crown with the hinged basals widely opened until the stem is almost in line with the median arm; bifurcations of main branch of axil-arms well shown.
16. L. ant. view of crown, giving direct view of bifurcations of median arm to about 10 finials.
- 16a. Post. view of same, with hinged basals tightly closed against radials, inferradials of vanished arms meeting below anals and indented by stem; anal plates but little visible at base, but tube projects at distal end.
17. A typical complete crown in "humped" position recumbent upon the stem, with semi-elliptical outline; median arm bifurcates on 6th brachial, and axil-arms at intervals of 3 and 4.
18. A larger specimen in similar position, with stem underneath; showing the complete branching of the median arm on 7th brachial to a total of 10 finials.
19. A large crown closely recumbent upon the stem, with arms partly overlapping it. Main branch of axil-arms well exposed in the last five specimens.
- 20, 21. Ant. and l. post. views of two young specimens with elongate calyx, only 3 axil-arms, and median arm branching once and twice. Keokuk shale, Mississippian; Indian Creek, Indiana.

PLATE 31

(Figures natural size unless otherwise stated)

	PAGE
<i>Ampheristocrinus typus</i> Hall.....	132
FIG. 1. Complete crown with very delicate arms, sharply angular calyx and long stem.
Beech River formation; Decatur County, Tennessee.	
<i>Cyathocrinus decatur</i> new species.....	134
FIG. 2. Crown with long arms and stem, and smooth, elongate calyx.
Beech River formation; Decatur County, Tennessee.	
<i>Cyathocrinus wilsoni</i> new species.....	134
FIG. 3. Crown with very long brachials and conical calyx; r. ant. iR view. $\times 3/2$.	
3a. Posterior side of calyx, showing anal plate. $\times 3/2$.	
Laurel limestone; St. Paul, Indiana.	
<i>Cyathocrinus</i> cf. <i>striolatus</i> Angelin.....	134
FIGS. 4, 5. Post. and r. ant. views of two calices, with ridges passing from BB to adjoining plates.
$\times 3/2$.	
6, 6a. Post. and dorsal views of similar specimen with one IB smaller than the rest. $\times 3/2$.	
6b. R. ant. view of same, showing small IB. $\times 5/2$.	
7. L. ant. view of specimen having 6 IBB. $\times 3/2$.	
Laurel limestone; St. Paul, Indiana.	
<i>Parisocrinus siluricus</i> new species	135
FIG. 8. Crown with arms in place, and part of stem; ant. view. $\times 3/2$.	
8a. L. post. view of same, with massive anal tube. $\times 3/2$.	
Laurel limestone; St. Paul, Indiana.	
<i>Botryocrinus tenuidactylus</i> new species.....	137
FIG. 9. A set of arms with calyx wanting, having strong ramules bearing slender pinnules.
Beech River formation; Decatur County, Tennessee.	
<i>Botryocrinus polyxo</i> (Hall).....	138
FIG. 10. Calyx showing form and position of RA.
Waldron shale; Hartsville, Indiana.	
(?) <i>Leocythiocrinus problematicus</i> new species.....	133
FIGS. 11, 11a, 11b. Post., dorsal and ventral views of the only specimen, with anal opening below
level of RR, which close above it; 3 small, unequal IBB; very large RR, and tegmen
apparently composed of a few large plates. $\times 3$.	
Laurel limestone; St. Paul, Indiana.	
<i>Parastephanocrinus intermedius</i> new genus and species.....	139
FIGS. 12, 12a, 12b. Post., dorsal and ventral views of unique specimen, showing anal opening in line
with RR directly over post. B; 5 IBB at bottom of deep cavity; forked RR with sinus
for food-grooves passing in to central cavity; tegmen not preserved. $\times 2$.	
12c. Detail of RR and food grooves. $\times 4$.	
Louisville limestone; near Bob, Decatur County, Tennessee.	

	PAGE
<i>Stephanocrinus angulatus</i> Conrad.....	139
- FIG. 13. Post. iR view of specimen with about the usual elongate form, showing sculptured surface and vertical ridges. $\times 2$.
14. Specimen showing the extreme of elongation the species sometimes assumes; l. ant. iR view. $\times 2$.
15. Lateral view of specimen with cluster of biserial arms in place. $\times 2$.
- 15a. The same radius from a slightly different angle, enlarged. $\times 4$.
16. Tegminal view of specimen showing orals and ambulacral grooves; anus seen covered by pyramid of small plates. $\times 2$.
Rochester shale; Lockport, New York.
<i>Stephanocrinus gemmiformis</i> Hall.....	140
- FIG. 17. Lateral view of calyx, low and rotund, without ornament, showing fork of RR. $\times 2$.
18. Tegmen of another specimen, showing orals, ambulacra and anal opening. $\times 2$.
- 19, 19a. Lateral and ventral views of similar specimen from Flatwoods, Tennessee.
Waldron shale; Newsom, Tennessee, except 19, which may be from Beech River formation.

PLATE 32

(Figures natural size unless otherwise stated)

	PAGE
<i>Gissocrinus lyoni</i> new species.....	135
FIGS. 1, 1a. Nearly complete crown, showing arms, anal series, and 3 IBB; posterior and dorsal views.
Louisville limestone; Jefferson County, Kentucky.	
<i>Gissocrinus delicatus</i> new species.....	136
FIG. 2. Crown with extremely slender arms, and anal series passing into a tube.
Beech River formation; Decatur County, Tennessee.	
<i>Gissocrinus magnibrachiatus</i> new species.....	136
FIG. 3. Crown with parts of arms and stem; calyx small with plates displaced; shows dorsal aspect of the huge brachials, with the small connecting or "patelloid" articulating plates between them, usually two in succession.	
4. Fragment containing a few of the large brachials—one axillary with double articulation at distal end.	
5. Dorsal side of a better preserved set of arms, showing 2 and 3 "patelloid" plates, and several axillary brachials with double articulation.	
6. Another partial set of arms, exposing both dorsal and ventral sides, the latter with the food-grooves completely covered. $\times 2$.
Beech River formation; Decatur County, Tennessee.	
<i>Gissocrinus approximatus</i> new species.....	137
FIGS. 7, 8. Fragments of isolated axillary brachials somewhat similar to the preceding, but apparently lacking the "patelloid" plates; food groove open and covered. $\times 3/2$.	
9. Another axillary brachial, almost complete, extremely elongate; natural size.
Laurel limestone; St. Paul, Indiana.	
<i>Gissocrinus quadratus</i> new species.....	137
FIG. 10. An axillary plate and two following it, smaller and with rectangular brachials, also without "patelloid" plates; food-grooves open. $\times 3/2$.	
11. Part of an arm, with food-groove covered. $\times 3/2$.
Horizon and locality, same as last.	

1

3

1a

8

2

9

7

10

4

6

11

5

PLATE 33

(All figures natural size unless otherwise stated)

PAGE*

- Troostocrinus reinwardti** (Troost)..... 141
- FIGS. 1, 2. R. ant. and post. views of average specimens.
 3. R. post. view of similar specimen with part of stem.
 4, 5, 6. Lateral views of 3 small specimens.
 7. Ant. view of maximum specimen.
 8. Specimen with the brachioles in place. $\times 2$.
Troostocrinus zone of Beech River formation; Decatur and adjoining counties,
 Tennessee.
- Troostocrinus sanctipaulensis** Foerste 141
- FIG. 9. Imperfect specimen from the Laurel, St. Paul, Indiana.
- Tetracystis fenestratus** Schuchert 141
- FIG. 10. Post. view, showing pectinirhomb and one food-groove complete to base; also doubled
 hydropore.
 11. R. ant. view with rhomb, showing large size of stem.
 12. L. post. view, showing lower rhomb and one food groove.
 13. R. post. view, showing anus and l. p. food-groove, with some of the lower brachioles in
 place.
 14. Ventral aspect, showing the 4 food-grooves, and single hydropore.
 Beech River formation; Decatur County, Tennessee.
- Stribalocystites gorbyi** S. A. Miller..... 142
- FIGS. 15, 16. L. post. and ventral views of typical specimen. $\times 2$.
 17. Specimen with folds strongly elevated and pores protruding with raised margins; post.
 view. $\times 2$.
 18. Specimen with pores prominent, but folds ill defined. $\times 2$.
 Laurel limestone; St. Paul, Indiana.
- Lysocystites sculptus** (S. A. Miller)..... 142
- FIG. 19. R. post. view of large elongate specimen showing striation and tubular folds, the latter
 broken in several places.
 20. Ventral view of similar specimen, showing anal opening, and small area for tegmen.
 21. Small, rotund specimen, showing surface structure and proportions of the 3 ranges of
 plates; r. ant. view. $\times 2$.
 22. Ventral view of still smaller specimen showing anus, and tegmen covered by a few small
 plates. $\times 2$.
 Laurel limestone; St. Paul, Indiana.
- Caryocrinus bulbulus** Miller and Gurley..... 143
- FIGS. 23, 24, 25. R. post., basal and ventral views of typical smooth specimen.
 26, 27. L. post. and l. ant. views of specimens with median protuberance on BB, and protrud-
 ing pores.
 Beech River formation; Decatur County, Tennessee.
- Caryocrinus persculptus** new species 143
- FIG. 28. R. ant. view of specimen with pores coalesced into rugose ridges.
 Waldron formation; Newsom, Tennessee.

- PAGE
- Caryocrinus milliganae* Miller and Gurley..... 143
- FIG. 29. The type, formerly in the collection of Mrs. J. M. Milligan; post. view; of the pear-shaped contour as is most frequent; shows the stereom folds with pores running in lines to angles of plates.
- 30, 31, 32, 33. Lateral and ventral views of specimens of varying contour, with sharper folds, and pores outlined by raised margins.
34. Specimen with coarser folds, post. view.
- 35, 36. Ant. and l. post. views of two maximum elongate specimens, with folds and pores well defined.
- Beech River formation; Decatur County, Tennessee.
- Caryocrinus ornatus* Say..... 143
- FIG. 37. L. ant. view, showing biserial, pinnulate arms, with pores sharply defined.
- Rochester shale; Lockport, New York.

SMITHSONIAN INSTITUTION LIBRARIES

3 9088 00579 6651