589.2.

.M11

DEPARTMENT OF AGRICULTURE, VICTORIA.

SYSTEMATIC ARRANGEMENT

OF

AUSTRALIAN FUNGI,

TOGETHER WITH

HOST-INDEX AND LIST OF WORKS ON THE SUBJECT,

BY

D. McALPINE,

GOVERNMENT VEGETABLE PATHOLOGIST,

MEMBER OF THE INTERNATIONAL PHYTO-PATHOLOGIC COMMISSION;

MEMBER OF THE IMPERIAL LEOP. CAROL. GERMAN ACADEMY OF NATURALISTS;

HONOURMAN OF THE SCIENCE AND ART DEPARTMENT, SOUTH KENSINGTON, LONDON, ETC., ETC.

AUTHOR OF A "BOTANICAL ATLAS" IN 2 VOLS.; A "ZOOLOGICAL ATLAS" IN 2 VOLS.;

A "BIOLOGICAL ATLAS"; "SHORT NOTES FOR BIOLOGICAL STUDENTS";

"ATLAS OF ELEMENTARY PHYSIOLOGY AND PHYSIOLOGICAL ANATOMY";

"LIFE-HISTORIES OF PLANTS," ETC. ETC.

N. Y. Academy of Sciences.

Rec'd 28 Sept 1896

By Authority

ROBT. S. BRAIN, GOVERNMENT PRINTER, MELBOURNE.

1895.

LIBRARY
NEW YORK
BOTANICAL
GARDEN

P R E F A C E.

IN dealing with the diseases of plants due to Fungi, it is necessary to determine the name and nature of the Fungus causing the disease, in order to be able to cope with it and to take effectual measures for its prevention, palliation, or cure. Accordingly I considered it essential to have the various known Australian Fungi recorded for reference, just as the various higher forms of Australian vegetation are so ably set forth by the Government Botanist, Baron von Mueller, in his Systematic Census. The very useful *Handbook of Australian Fungi*, prepared by Dr. M. C. Cooke, the veteran mycologist, under the sanction and authority of the various colonial Governments, has been taken as a basis and prepared the way for the present publication. This *Systematic Arrangement of Australian Fungi* aims at giving in a compact and handy form a complete enumeration of all the known species up to date, systematically arranged so as to show their relationships, and briefly described, together with such additional information as may be of use in a future detailed and more directly useful account. The object being to bring together all the species recorded by the various workers in this field, to take stock, as it were, of what has been done, I had to consult the different works bearing on the subject previous to the addition of a large number of hitherto unrecorded Fungi to the list. I have accordingly prepared a "List of Works on Australian Fungi," the first of its kind. A complete list of Fungi having been compiled from the various publications, including several papers of my own read before the Royal Society of Victoria, together with the plants or parts of plants on which they occurred, in the case of parasitic forms, the material was supplied for a provisional Host-index; the term "host" being applied to the plant on which the Fungus lives or preys, the Fungus being an unwelcome guest as a rule. The necessity for a Host-index became apparent from the time I was appointed Vegetable Pathologist.

There are thus three connected and interdependent divisions in this publication, which may now be briefly glanced at and explained.

I.—SYSTEMATIC ARRANGEMENT OF AUSTRALIAN FUNGI.

The plan pursued is the following:—

1st. A consecutive number is given to each species, for convenience of reference, and all future additions will be numbered consecutively. Varieties are distinguished by having a letter added to the number of the species.

2nd. The number in Dr. Cooke's *Handbook of Australian Fungi* is next given for ready reference to the description of any species in that work. This serves a double purpose, and shows not only the species recorded in that work, but also species omitted.

3rd. The volume and number is next quoted for every Australian species given in Saccardo's *Sylloge Fungorum*, consisting at present of ten thick volumes, which are in the Melbourne Public Library. This is the standard work on Fungi, and is the most complete and exhaustive at the present time. The references to Cooke and Saccardo will leave no doubt as to the particular Fungus meant.

4th. The scientific name adopted for each species of Fungus follows next. It is absolutely necessary, for purposes of accuracy, to have the scientific names as well as the common names, for otherwise serious mistakes may arise. Thus, the name of "Peach Yellows" (the dreaded American disease) is often applied to a disease of the Peach in this colony, but, fortunately, it is a very different and much more harmless disease, being none other than the Peach-leaf Rust (*Puccinia Pruni*). It may be noted that the sub-genera of *Agaricus* are raised to the rank of genera; and, as the original generic name is thus set free, it is retained for the species to which the common Edible Mushroom belongs, and which were formerly included in the sub-genus *Pseudotsu*.

5th. The authority for the name is next stated. With so many different names often applied to the same Fungus, and even the same name often applied by different authors to entirely different plants, it is necessary to give the authority for the particular name, in order to indicate the precise Fungus meant. The name of the authority is usually given in a contracted form, and it will be noticed that it is sometimes printed in italics. The reason of this is that it is customary in works on Fungi often to give two authorities, the first to indicate the original describer of the Fungus, and the second where some one has classified it differently on good scientific grounds. I simply give one authority, the name of the original describer being printed in Roman characters; and, where the original name has been set aside, the correct classifier is given in italics. The year of publication is also stated.

As an illustration of the variety of naming, I may mention one kind of Rust of Wheat met with in the colony, and the following nine names have been given to it by the authors whose names are appended:—

Uredo rubigo-vera, De Candolle.

Uredo rubigo, Berkeley.

Caeoma rubigo, Link.

Trichobasis rubigo-vera, Leveille.

Trichobasis glumarum, Leveille.

Puccinia rubigo-vera, Winter,

Puccinia striiformis, Westendorp.

Puccinia straminis, Fuckel.

Eecidium asperifolii, Persoon.

By a recognised principle the name of *Puccinia rubigo-vera* is adopted, although Dr. Cooke in his Handbook uses the name of *Puccinia straminis*.

6th. The English name follows. This is merely an attempt to give an English rendering to the specific name, and something of the kind is necessary in naming Fungus diseases to the average farmer or fruit-grower: but, as these diseases become better known as to their cause, some characteristic feature of the disease may be used as a distinguishing name, such as Leaf-curl, Shot-hole, Bitter-rot, Club-root, &c.

7th. The "Habitat" is next given, the various colonies in which the species have been found being recorded. It has been thought advisable to add B. for British when it occurs there, as there may have been preventives or remedies applied in the old country which it would be profitable for us to know. I make no apology for dealing with Australian Fungi, including the five colonies of the Australian Continent and Tasmania, for Fungi do not respect our political boundaries and restrict themselves to artificial limits. There must be federation in the treatment of disease if it is to be thoroughly effectual, and this has been happily illustrated in dealing with the Rust in Wheat question, in which all the colonies are united for devising measures against a common enemy.

8th. The "Occurrence" follows, indicating on what plants or parts of plants the different kinds of Fungi may be looked for. This is afterwards collectively shown in the Host-index, each plant having all its known diseases due to Australian Fungi ranged under it.

9th. "General characters" conclude the whole, giving such superficial and easily-recognised characters as may serve as a guide in the rough discrimination of many species requiring immediate attention to check their spread.

From the very nature of this work and from our present very limited knowledge of the Fungi of Australia there will be constant additions made (in fact, I have quite a number of new species awaiting determination myself), and this will be met by the issue of supplements, when necessary, on the same lines. As Dr. Cooke truly says in his introduction to the Handbook—"It is quite probable that in the course of a few years, by working up the minute species, the total number contained in this volume would be more than doubled, even without the investigation of unexplored districts."

It ought also to be borne in mind that many of the more conspicuous Fungi—such as what are popularly called Mushrooms and Toadstools—work considerable mischief, although unseen and unnoticed. Thus, the Honey Agaric (*Armillaria mellea*), which is even considered edible, does a deal of damage, and by attacking the roots undermines the tree. It spreads from root to root in the soil by means of long purple-black cord-like strands, even in the absence of the tawny-yellow "Toadstools," which are simply the fructification of the Fungus, and I have seen orchard trees killed by this cause. In the soil and in the rotting roots or wood these strands are found, attacking the roots and bases of stems and often causing copious "gumming" there. The Vegetable Pathologist should therefore not only be more or less conversant with the Fungi of the different colonies, as they spread so readily by means of their spores, but he should be acquainted with Fungi as a whole, since even Mushrooms and Toadstools are not beyond his province.

II.—PROVISIONAL HOST-INDEX OF AUSTRALIAN FUNGI.

The list of Fungi, systematically arranged, enables us to classify them under their respective Host-plants. Strictly speaking, it is only those which are parasitic, or which prey upon living plants, that should be included; but it is so difficult with our present knowledge to distinguish between those which cause disease and those which attack decaying or decayed parts, that I have given all the Fungi found upon any particular plant. While special attention is paid to the Fungi occurring on the various vegetable products grown in the colony for commercial purposes, as given in the Government Statist's returns, the Fungi on so-called "weeds" are not neglected, because they may and often do pass over to the cultivated and therefore more delicate forms of vegetation. For example, the Fungus causing "Club-root" in Cabbages, Cauliflowers, Turnips, Radishes, Kale, &c., also infests two of our common weeds, viz., Shepherd's Purse (*Capsella Bursa-pastoris*, Moench) and Hedge Mustard (*Sisymbrium officinale*, Scop), and many similar instances could be given. This fact is strikingly put by Mr. Bailey, who says—"As we find in the animal kingdom the wild man preferring sheep to kangaroo, the flying fox peaches to quandong, the grasshopper the more succulent vegetation of our gardens to the dry herbage of the plains, so in like manner we shall doubtless find from time to time blight-fungi, at present unknown, will come from the indigenous plants to exotic ones which may be more congenial to their development."

There can be no doubt that many of the Fungi on our native vegetation will attack introduced plants, and it would be very desirable, both in the interests of science and of practical utility, to have a record of the Fungi preying upon our native plants. I have seen some of our richest soils with the decaying roots of Eucalypts and the mycelium of Fungi passing from them to the roots of orchard trees and causing their decay.

The Host-index should serve various useful purposes. First of all, it will enable the intelligent grower to determine with some degree of certainty the cause of the disease when it is due to a Fungus, and that is often the first step towards its eradication. Thus, if his Peach trees are affected with some Fungus disease on the leaves, he turns up the Index and finds two Fungi recorded there. He then turns to the General Characters in the "Systematic Arrangement" and can easily tell whether it is the "Peach-leaf Rust" or the "Leaf-curl." Or if his Cabbages and Cauliflowers begin to turn yellow and the roots become distorted, he finds from the Index that it is due to a Fungus, a knowledge of which enables him to battle with the disease. Having traced the disease to its source, he may find treatment already prescribed in some of the Government publications, or can apply to the Department for advice. If there is no record of the disease in the Index, then the grower knows it is a subject requiring investigation.

Further, the Host-index may be used in assisting growers to "spot" diseases due to Fungi before they have spread too far and become established. A great many Fungus diseases are overlooked for a number of years and allowed to spread freely before active measures are taken for their suppression, and thus what might have been easily nipped in the bud is now difficult to eradicate; so that another important use of this publication will be to enable Fungus diseases to be recognised at the earliest possible moment and action taken accordingly.

Onion Mould, Ergot in Rye and other Grasses, Powdery Mildew in Apple, and various other diseases, are not recorded in Cooke's Handbook, and, presumably, have been neglected.

A third use will be to assist in the carrying out of any legislation which may be passed for the suppression of Insect and Fungus pests. Many growers err in ignorance, because they are not aware of the disease being present until it has got a firm hold, but now a record of the various Fungus pests is available.

And there is a final purpose to be served which is not the least important. New diseases are continually cropping up, and the sooner they are recognised the better. If the disease is not recorded in the Index there is a strong probability of its being some new one, and then it can be traced to its source without delay.

The names of the Host-plants are given according to Baron von Mueller's *Second Systematic Census of Australian Plants* or Hooker and Jackson's *Index Kewensis*, as far as published. The Fungi belonging to Victoria are indicated by the letter V.

III.—LIST OF WORKS ON AUSTRALIAN FUNGI.

It was necessary, as already stated, to draw up a list of works in order to have the list of Fungi as complete as possible. I have only included those publications in which there is special reference to Australian forms, and no doubt several have been overlooked. To Dr. Alexr. Morrison I am much indebted for bringing under my notice some references to the subject in scattered publications. The "List of Works," the "Systematic Arrangement," and the "Host-index" should serve to focus our present knowledge and prepare the way for further additions to it.

In giving the general characters of the various Fungi, I have endeavoured to use as simple terms as possible, but it was difficult to avoid the employment of technical terms occasionally. For those who wish to enter into the subject more fully and to study in an elementary way the disease-causing Fungi, the following works among others may be mentioned:—

Diseases of Plants, by Professor Marshall Ward, and published by the Society for promoting Christian Knowledge (2s. 6d.). This is a readable little book, and treats in a popular manner such diseases as Rust in Wheat, Smut of Corn, Ergot of Rye, Hop disease, Potato disease, &c.

Diseases of Field and Garden Crops, by Worthington G. Smith, and published by MacMillan and Co. (4s. 6d.). This work is beautifully illustrated, and treats of Onion, Pea, Parsnip, Lettuce, Potato, and other diseases, in addition to those of Wheat and Oats.

Diseases of Crops and their Remedies, by Dr. A. B. Griffiths, and published in Bell's Agricultural Series (2s. 6d.). The diseases of leguminous, gramineous, root, and miscellaneous crops are considered, together with the Fungi or insects causing them, and the best methods of prevention.

Fungus Diseases of the Grape and other Plants, and their Treatment, by F. Lamson-Scribner, and published in America (5s.). This is a thoroughly practical work, and deals with the principal Fungus diseases of Fruit trees as well as of the Vine.

Fungi and Fungicides, by Dr. C. M. Weed, and published in New York (5s.). It is divided into five parts—Fungi affecting the larger fruits, the small fruits, shade trees, &c., vegetables, cereals, and forage crops; and practical remedies, as a rule, are given.

There is still a want of proper works dealing with the subject of Fungus disease from an Australian standpoint and suited to the wants of our orchardists and vignerons especially, but the strong necessity which exists for such information will probably soon lead to its being supplied.

The preparation of this work has entailed a vast amount of labour, done single handed and in my spare time, but it was absolutely necessary as a preliminary for the proper carrying out of my duties. To all those who have supplied me with information my best thanks are due and are hereby tendered. It is hardly necessary to mention special names, since the "List of Works" will afford the best evidence of work done. The Government Botanist, Baron von Mueller, has always aided me with the free use of his library and the benefit of his rare and critical knowledge in connexion with some of the Host-plants. Mr. F. M. Bailey, F.L.S., Colonial Botanist of Queensland, has given me every assistance in his power in connexion with Queensland Fungi, and Mrs. Flora Martin is well known for her indefatigable labours in extending our knowledge of Australian species. I am indebted to A. de Bary for a list of the Yeasts identified by him in Australia, and he adds that they will be largely increased from time to time. Wine Yeasts especially will yet play an important part in connexion with that industry, and there are kinds of Yeast causing decomposition and disease in Onions, &c. The officers in the neighbouring colonies have also willingly given me the benefit of their advice when asked. Amid such a mass of detail some important points may have been overlooked, and I shall be pleased to have any errors or omissions pointed out, such additions and corrections to be subsequently issued as a supplement.

It must not be imagined that because we have tabulated and briefly described a number of Fungi we therefore know all that is necessary about them. The most fascinating branch is the life-history—the story of their lives from year to year; and it is this knowledge as to their various and often disguised phases, how they spread, and where they winter, which will help us to cope with them successfully. There is room for plenty of workers, and it is hoped that some of our young and rising fruit-growers and farmers may be induced to attend to this subject, on account of its great interest and practical importance.

CONTENTS.

PREFACE		iii
I.—SYSTEMATIC ARRANGEMENT OF AUSTRALIAN FUNGI.		
Synopsis of Groups		1
General Classification of Hymenomycetes		3
Group I.—Hymenomycetes—	PAGE	
Order 1.—Agaricaceæ	4	
Additions to Agaricaceæ	30	
Order 2.—Polyporaceæ	34	
" 3.—Hydnaceæ	58	
" 4.—Thelephoraceæ	62	
" 5.—Clavariaceæ	72	
" 6.—Tremellaceæ	76	
Group II.—Gastromycetes—		
Order 7.—Pilaceraceæ	82	
" 8.—Phalloideaceæ	82	
" 9.—Nidulariaceæ	86	
" 10.—Lycoperdaceæ	88	
" 11.—Sclerodermaceæ	96	
" 12.—Hymenogastreae	98	
Group III.—Uredines—		
Order 13.—Uredinaceæ	102	
Group IV.—Pyrenomycetes—		
Order 14.—Hypocreaceæ	110	
" 15.—Xylariaceæ	114	
" 16.—Dothideaceæ	118	
" 17.—Melogrammaceæ	120	
" 18.—Diatrypaceæ	120	
" 19.—Valsaceæ	122	
" 20.—Entypaceæ	122	
" 21.—Cucurbitariaceæ	122	
" 22.—Superficiales	122	
" 23.—Pertusaceæ	124	
" 24.—Lophiostomaceæ	124	
" 25.—Ceratostomaceæ	124	
" 26.—Obtectaceæ	124	
" 27.—Caulicolaceæ	126	
" 28.—Folicolaceæ	126	
" 29.—Microthyriaceæ	128	
" 30.—Perisporiaceæ	128	
" 31.—Hysteriaceæ	132	
Group V.—Discomycetes—		
Order 32.—Cyttariaceæ	136	
" 33.—Helvellaceæ	136	
" 34.—Pezizaceæ	136	
" 35.—Ascobolaceæ	144	
" 36.—Dermataceæ	144	
Additions—(A.) New Australian Fungi		200
(B.) New Localities, Hosts, &c.		205
(C.) List of Australian Edible Fungi		207
Tables		208
II.—HOST-INDEX OF AUSTRALIAN FUNGI		213
III.—LIST OF WORKS ON AUSTRALIAN FUNGI		227

Group V.—Discomycetes— <i>continued</i> —		
Order 37.—Bulgariaceæ	141	
" 38.—Stictaceæ	146	
" 39.—Phacidiaceæ	146	
" 40.—Patellariaceæ	146	
" 41.—Gymnoascææ	146	
Group VI.—Tuberoides—		
Order 42.—Elaphomycetaceæ	150	
" 43.—Tuberaceæ	150	
" 44.—Eudogouaceæ	150	
Group VII.—Hyphomycetes—		
Order 45.—Mucedinaceæ	154	
" 46.—Dermatiaceæ	156	
" 47.—Stilbeaceæ	160	
" 48.—Tuberculariaceæ	162	
Group VIII.—Sphæropsides—		
Order 49.—Sphæroidaceæ	168	
" 50.—Nectrioidaceæ	174	
" 51.—Leptostromaceæ	174	
" 52.—Excipulaceæ	176	
" 53.—Melaucouiaceæ	176	
Group IX.—Saccharomycetes—		
Order 54.—Saccharomycetaceæ	180	
Group X.—Ustilagines—		
Order 55.—Ustilaginaceæ	184	
Group XI.—Phycomycetes—		
Order 56.—Mucoraceæ	190	
" 57.—Peronosporaceæ	190	
" 58.—Eutomophthoraceæ	190	
" 59.—Chytridiaceæ	190	
" 60.—Protomycetaceæ	190	
Group XII.—Myxomycetes—		
Order 61.—Tubulinaceæ	194	
" 62.—Cribrariaceæ	194	
" 63.—Stemonitaceæ	194	
" 64.—Lamprodermaceæ	194	
" 65.—Arcyriaceæ	194	
" 66.—Trichiaceæ	196	
" 67.—Didymaceæ	196	
" 68.—Physariaceæ	198	
" 69.—Plasmodiophoraceæ	198	

I.—SYSTEMATIC ARRANGEMENT OF AUSTRALIAN FUNGI.

SYNOPSIS OF GROUPS.

IN selecting a system of classification I have adopted that which best expresses the present state of our knowledge as regards the life-histories of the various forms, which after all constitute the ultimate court of appeal in settling affinity. But unfortunately there are numerous cases where the life-history has not been wrought out, and so certain groups have to be provisionally placed along with those to which they seem to be most nearly related. Saccardo's *Sylloge Fungorum* has been mainly followed, while Dr. Cooke's *Handbook of Australian Fungi*, G. Masec's *British Fungus-Flora*, De Bary's *Fungi*, and Brefeld's works have all been consulted.

GROUPS OF AUSTRALIAN FUNGI.

MYCOMYCETES ...	I.—HYMENOMYCETES	...	...	} Basidiomycetes.
	II.—GASTROMYCETES	...	...	
	III.—UREDINES	...	...	Æcidiomycetes.
	IV.—PYRENOMYCETES	...	...	} Ascomycetes.
	V.—DISCOMYCETES	...	...	
	VI.—TUBEROIDES	...	...	
	VII.—HYPHOMYCETES	...	...	} Imperfect forms of Ascomycetes ?
	VIII.—SPILEROPSIDES	...	...	
	IX.—SACCHAROMYCETES	...	...	
	X.—USTILAGINES	...	...	Transitional forms.
	XI.—PHYCOMYCETES	...	...	} Transitional to animals.
	XII.—MYXOMYCETES	...	...	

The systematic sequence of the groups is at present a matter of individual opinion, but they are arranged in the order in which they will be treated, and are reduced within the smallest limits consistent with clearness. The two main divisions are *Mycomycetes*, in which there are no sexually produced reproductive bodies, and *Phycomycetes*, or those approximating to sea-weeds, in which reproduction is sexual as well as asexual.

The *Mycomycetes* are divided into two chief classes—*Basidiomycetes* and *Ascomycetes*—the former producing naked spores at the ends of large terminal cells called *basidia*, and the latter producing spores in an *ascus* or bag. The *Ustilagines* are regarded as transition forms to the *Phycomycetes*. The *Mycomycetes* differ in important points from fungi, and are regarded as more nearly related to animals, but they are conveniently retained here for the present. The *Schizomycetes* or Bacteria also differ from fungi in chlorophyll being sometimes present, and the hyphæ or the threads of the ordinary fungus absent. To this group belong some of the organisms causing disease in plants, but as the greater part of the forms belong to medicine, I have finally decided to omit them. The *Uredines* are doubtful in their affinities, and they are placed in a class, *Æcidiomycetes*, between the other two until their position is properly settled.

The imperfect forms are those which are assumed to be genetically related to other fungi, probably *Ascomycetes*, in contradistinction to the perfect fungi, which have an independent life-history.

The characteristic features of each of the twelve groups are here shown, then the general classification of each group is given in its proper connexion.

BASIDIOMYCETES.—Naked spores borne on basidia. Receptacle distinct.

I.—HYMENOMYCETES.—Hymenium external.

II.—GASTROMYCETES.—Hymenium internal.

ÆCIDIOMYCETES.—Æcidium or cluster-cup forms a feature of the life-history.

III.—UREDINES.—Receptacle none or obsolete.

ASCOMYCETES.—Spores produced in asci or spore-sacs.

IV.—PYRENOMYCETES.—Receptacles (*Perithecia*) flask-shaped or spherical, opening at apex.

V.—DISCOMYCETES.—Receptacles (*Apothecia*) disc- or eup-shaped.

VI.—TUBEROIDES.—Subterranean, sub-globose, indehiscent.

IMPERFECT FORMS OF ASCOMYCETES?—

VII.—HYPHOMYCETES.—Perithecia absent.

VIII.—SPILÆROPSIDES.—Perithecia present.

IX.—SACCHAROMYCETES.—Multiplication by gemmation and ascospores.

TRANSITIONAL FORMS.—

X.—USTILAGINES.—Minute, parasitic, usually spores of one kind only.

ALGA-LIKE FORMS.—

XI.—PHYCOMYCETES.—Mycelium without septa. Sexual and asexual reproduction.

ANIMAL-LIKE FORMS.—

XII.—MYXOMYCETES.—Plasmodium or naked mass of motile protoplasm formed and hyphæ absent.

GENERAL CLASSIFICATION OF HYMENOMYCETES.

GROUP I.—HYMENOMYCETES, FRIES.

ARRANGEMENT OF ORDERS (6).

Hymenium or spore-bearing surface normally inferior—

1. AGARICACEÆ—Hymenium spread over gills.
2. POLYPORACEÆ—Hymenium spread over tubes or pores.
3. HYDNACEÆ—Hymenium spread over prickles.
4. THELEPHORACEÆ—Hymenium spread over an even surface.

Hymenium superior or encircling—

5. CLAVARIACEÆ—Plants club shaped or branched, rarely lobed.
6. TREMELLACEÆ—Plants lobed, convolute, or disc-like; gelatinous.

ORDER I.—AGARICACEÆ, FRIES.

ARRANGEMENT OF GENERA (55).

Section I.—Lencosporæ—Spores white, or nearly so.

Series 1. Haplophyllæ—Gills entire at edge.

Sub-section I. Molles—Plants fleshy, more or less firm, putrescent, not reviving when once dried.

Genera (16)—

1. Amanita, Pers.	5. Armillaria, Fries.	9. Mycena, Pers.	13. Hygrophorus, Fries.
2. Amanitopsis, Roze.	6. Tricholoma, Fries.	10. Hiatula, Fries.	14. Lactarius, D.C.
3. Lepiota, Pers.	7. Clitocybe, Fries.	11. Omphalia, Fries.	15. Russula, Pers.
4. Schulzeria, Bres.	8. Collybia, Fries.	12. Pleurotus, Fries.	16. Cantharellus, Pers.

Sub-section 2. Tenaces—Plants tough and leathery, or hard, reviving when moistened.

Genera (6)—

17. Marasmius, Fries.	19. Panus, Fries.	21. Trogia, Fries.	22. Lenzites, Fries.
18. Lentinus, Fries.	20. Xerotus, Fries.		

Series 2. Schizophyllæ—Gills split at edge.

Genus (1)—

23. Schizophyllum, Fries.

Section 2. Rhodosporeæ—Spores rosy or salmon pink.

Genera (10)—

24. Metrarhia, C. and M.	27. Pluteus, Fries.	30. Leptonia, Fries.	32. Eecilia, Fries.
25. Volvaria, Fries.	28. Entoloma, Fries.	31. Nolanea, Fries.	33. Cladopus, Smith.
26. Annularia, Schulz.	29. Clitopilus, Fries.		

Section 3. Ochrosporeæ—Spores ochrey brown or red brown.

Genera (11)—

34. Pholiota, Fries.	37. Hebeloma, Fries.	40. Galera, Fries.	43. Cortinarius, Pers.
35. Locellinia, Gill.	38. Flammula, Fries.	41. Tubaria, Smith.	44. Paxillus, Fries.
36. Inocybe, Fries.	39. Naucoria, Fries.	42. Crepidotus, Fries.	

Section 4. Melanosporæ—Spores blackish purple, purplish brown, black, or nearly black.

Genera (11)—

45. Agaricus, Linn.	48. Psilocybe, Fries.	51. Bolbitius, Fries.	54. Anellaria, Karst.
46. Stropharia, Fries.	49. Deconica, Smith.	52. Coprinus, Pers.	55. Psathyrella, Fries.
47. Hypholoma, Fries.	50. Psathyra, Fries.	53. Panaeolus, Fries.	

Total number of species = 552.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
---------	-----------------	--------------------	------------------	---------------------	---------------

GROUP I.—HYMENOMYCETES.—FRIES, SYST. MYC. I. 53 (1821).

1. AMANITA.—Pers. Syn. 246 (1801).

1	8	V. 36	<i>A. ananæceps</i> ...	Berk., Hook., Lond. Journ. VII. 572 (1848)	Pine-apple-headed amanita ...
2	16	" 23	<i>A. grossa</i> ...	Berk., Fl. Tasm. II. 242 (1860) ...	Large amanita ...
3	4	" 8	<i>A. mappa</i> ...	Fries, Epier. 6 (1838) ...	Napkin amanita ...
4	6	IX. 5	<i>A. murina</i> ...	Cooke and Mass., Grev. XVIII. 1 (1889)	Mouse-coloured amanita ...
5	7	V. 18	<i>A. muscaria</i> ...	Linn., in Fries S.M. I. 16 (1821)	Fly amanita ...
6	1	" 3	<i>A. ovoidea</i> ...	Fries, Hym. Eur. 18 (1874) ...	Ovoid amanita ...
7	2	" 4	<i>A. Preissii</i> ...	Fries, Pl. Preiss. II. 131 (1846)	Preiss's amanita ...
8	9	" 31	<i>A. spissa</i> ...	Fries, Epier. 9 (1838) ...	Clammy amanita ...
9	5	...	<i>A. strobilacea</i> ...	Cooke, Grev. XIX. 82 (1891) ...	Cone-like amanita ...
10	3	V. 7*	<i>A. verna</i> ...	Fries, Hym. Eur. 18 (1874) ...	Spring amanita ...

2. AMANITOPSIS.—Roze, in Karst.

11	14	IX. 9	<i>A. curta</i> ...	Cooke and Mass., Grev. XVI. 72 (1888) ...	Short-stalked amanitopsis ...
12	13	" 7	<i>A. farinacea</i> ...	Cooke and Mass., Grev. XVIII. 1 (1889)	Mealy amanitopsis ...
13	12	" 4	<i>A. illudens</i> ...	Cooke and Mass., Grev. XVI. 30 (1887)	Illusive amanitopsis ...
14	15	" 6	<i>A. pulchella</i> ...	Cooke and Mass., Grev. XVIII. 1 (1889)	Beautiful amanitopsis ...
15	11	V. 47	<i>A. vaginata</i> ...	Roze, Karst. Hattsv. I. 6 (1879)	Sheathed amanitopsis ...

3. LEPIOTA.—Pers.,

16	39	V. 150	<i>L. asprata</i> ...	Berk., Hook., Lond. Journ. VI. 481 (1847)	Warty lepiota ...
17	43	" 258	<i>L. australiana</i> ...	Fries, Pl. Preiss. II. 131 (1846) ...	Australian lepiota ...
18	25	" 185	<i>L. Beckleri</i> ...	Berk., Linn. Journ. XIII. 156 (1873)	Beckler's lepiota ...
19	32	" 243	<i>L. bubalina</i> ...	Berk., Linn. Journ. XIII. 156 (1873)	Gazelle lepiota ...
20	37	" 130	<i>L. cepæstipes</i> ...	Fries, Hym. Eur. 35 (1874) ...	Onion-stalked lepiota ...
20A	"	" "	<i>L. cepæstipes</i> , var. <i>cretacea</i>	Bull. Champ. 374 (1798) ...	Chalky lepiota ...
21	33	" 236	<i>L. cheimoniceps</i> ...	Berk. and Curt., Linn. Journ. X. 283 (1869)	Winter-capped lepiota ...
22	24	" 101	<i>L. clypeolaria</i> ...	Fries, S.M. I. 21 (1821)	Shield-like lepiota ...
23	29	" 111	<i>L. cristata</i> ...	Fries, S.M. I. 22 (1821)	Crested lepiota ...
24	20	" 86	<i>L. dolichaula</i> ...	Berk. and Br., Linn. Trans. XXVII. 150 (1869)	Long-tubed lepiota ...
25	18	" 83	<i>L. excoriata</i> ...	Fries, Hym. Eur. 30 (1874)	Flaky lepiota ...
26	28	IX. 29	<i>L. finetaria</i> ...	Cooke and Mass., Grev. XVIII. 1 (1889)	Dung lepiota ...
27	38	V. 145	<i>L. granulosa</i> ...	Fries, Hym. Eur. 36 (1874)	Granular lepiota ...
28	42	IX. 36	<i>L. lavendulae</i> ...	Cooke and Mass., Grev. XVI. 72 (1888)	Lavender lepiota ...
29	34	V. 234	<i>L. leontoderes</i> ...	Berk. and Br., Linn. Journ. XI. 499 (1871)	Tawny lepiota ...
30	21	" 170	<i>L. lepidophora</i> ...	Berk. and Br., Linn. Journ. XI. 498 (1871)	Scaly lepiota ...
31	36	" 132	<i>L. liemophora</i> ...	Berk. and Br., Linn. Journ. XI. 500 (1871)	Shovel-bearing lepiota ...
32	19	" 88	<i>L. mastoidea</i> ...	Fries, S.M. I. 20 (1821)	Bossed lepiota ...
33	...	...	<i>L. membranacea</i> ...	Cooke and Mass., Grev. XXI. 36 (1892)	Membranous lepiota ...
34	40	V. 165	<i>L. mesomorpha</i> ...	Fries, El. I. 2 (1828)	Intermediate lepiota ...
35	31	" 127	<i>L. naucina</i> ...	Fries, Epier. 16 (1838)	Short-stalked lepiota ...
35A	31	" 127	<i>L. naucina</i> , var. <i>sphaerospora</i>	Cooke and Mass., Grev. XVIII. 5 (1890)	Globose spored lepiota ...
36	35	IX. 37	<i>L. obclavata</i> ...	Cooke and Mass., Grev. XVI. 30 (1887)	Obclavate lepiota ...
37	23	IX. 13	<i>L. ochrophylla</i> ...	Cooke and Mass., Grev. XVIII. 2 (1889)	Ochre-gilled lepiota ...
38	16	V. 70	<i>L. procera</i> ...	Fries, S.M. I. 20 (1821)	Tall lepiota ...
39	17	" 74	<i>L. rhacodes</i> ...	Fries, Hym. Eur. 29 (1874)	Stripped lepiota ...

OF AUSTRALIAN FUNGI.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			

ORDER I.—AGARICACEÆ.—FRIES, PL. HOMON. 65 (1825).

Agaricus, Amanitopsis.

1	...	...	T.	...	...	Q.	...	Ground ...	Broad, smooth, shining, breaking into distinct angular spaces or areolae at the centre, each bearing a conical wart; pine-apple-like.
2	...	...	T.	...	...	...	...	Ground ...	White. Thick, fleshy, warted. Stem bulbous; ring obsolete.
3	...	...	...	V.	...	...	B	Ground ...	Delicate. Without separable enticle, dry, primrose, white, or buff.
4	...	...	...	V.	...	Q.	...	Sandy soil ...	Bell shaped, shining, mouse coloured. Stem thin, straight, whitish.
5	...	S.A.	...	V.	...	...	B	Woods ...	Large. Orange scarlet, clothed with scattered warts. Very common.
6	...	...	...	V.	...	...	...	Ground ...	White. Hemispherical, margin inflexed. <i>Edible</i> .
7	W.A.	...	...	...	...	...	...	Sandy soil, woods, &c.	Fleshy, viscid. Stem stuffed, mealy, rooting.
8	...	S.A.	...	...	...	...	B	Woods ...	Rough, with minute crowded mealy warts, amber grey.
9	...	...	...	V.	...	...	...	Ground ...	Hemispherical, covered with large persistent obtusely-conical warts, arranged after the manner of a fir-cone, ochrey yellow.
10	...	...	...	V.	...	Q.	B	Moist woods	Snow white, beautiful, viscid. Appearing in spring and summer.

Hattsv. I. 6 (1879).—Agaricus, Amanita.

11	...	...	...	V.	...	...	...	Ground ...	Ochrey white. Stem short, bulbous, brick red.
12	...	...	...	...	...	Q.	...	Ground ...	White and mealy. Fleshy, sprinkled with erect prominent warts, chiefly at disc.
13	...	...	...	V.	...	...	...	Ground ...	Ochrey yellow, clad with scattered broad unequal warts, which soon fall away.
14	...	...	...	V.	...	...	...	Ground ...	Vermilion, clad with irregular deciduous whitish warts. Stem hollow, white.
15	...	...	...	V.	N.S.W.	Q.	B	Woods and under trees	Size and colour very variable, grey, brown, &c. Thio, margin membranaceous, deeply furrowed. Common. <i>Edible</i> .

Tent. Disp. 68 (1797).—Agaricus.

16	...	...	...	...	N.S.W.	Q.	...	Ground, trunks ...	Hemispherical, pallid yellow, often deep orange, rough with warts.
17	W.A.	...	...	...	...	...	...	Sandy soil ...	Large. Slightly fleshy, viscid. Stem long, club shaped downwards.
18	...	...	...	...	N.S.W.	...	...	Ground, in scorched places	Spongy, rough about apex with little scales. Stem long, minutely warted.
19	W.A.	...	...	V	...	...	...	Cow-dung, &c. ...	Snow white. Cap ovate to hemispherical. Stem thickened downwards.
20	...	...	...	...	...	Q.	B	Ground ...	Sub-membranaceous, mealy and scaly, yellow. Stem hollow and bulbous.
20A	...	...	...	...	...	Q.	...	Ground ...	Chalky white, with darker scales.
21	...	...	...	...	...	Q.	...	Trunks ...	Snow white. Thin, powdery. Stem thickened downwards; mealy.
22	...	...	...	V.	...	Q.	B	Woods and hot-houses	Slender. Fleshy, variable in colour, yellowish or pinkish. Common.
23	...	...	T.	V.	...	...	B	Fields, lawns, &c. ...	Small and delicate. Slightly fleshy, whitish yellow, red scales.
24	...	...	...	...	...	Q.	...	Ground ...	Fleshy, centre smooth, otherwise with point-like scales, margin torn.
25	W.A.	...	...	V.	N.S.W.	Q.	B	Pastures ...	Small and delicate. Fleshy, pale-fawn colour. Enticle thin, breaking up into scattered papillae. <i>Edible</i> .
26	...	...	...	...	...	Q.	...	Dung ...	Fleshy, thin, pallid, ornamented with darker scales. Stem slender, scaly below.
27	...	...	...	V.	...	Q.	B	Woods and heaths ...	Small. Fleshy, tawny or dull reddish yellow, mealy and granular.
28	...	S.A.	...	V.	...	...	...	Ground ...	Rather fleshy, mealy, greyish-blue or dove colour. Stem cylindrical, whitish.
29	...	...	...	...	...	Q.	...	Ground ...	Tawny, with a few pallid warts, minutely tomentose or downy.
30	...	...	...	...	N.S.W.	...	...	Ground ...	Rather fleshy, white, sprinkled with minute reddish scales, yellow on drying.
31	...	...	...	V.	...	...	B	Ground ...	Membranaceous, lemon coloured, folded and furrowed, margin notched.
32	...	S.A.	...	V.	N.S.W.	Q.	B	Ground ...	Whitish brown. Rather fleshy, umbo or boss acute. scales papillate. <i>Edible</i> .
33	...	...	...	...	...	Q.	...	Chips of wood buried in ground	Membranaceous, thin, pale-cream colour. Stem slender, hollow.
34	...	...	...	V.	...	...	B	Ground ...	Slender. Rather fleshy, tawny. Stem even and smooth, as well as cap.
35	...	...	...	V.	...	...	B	Fields ...	Whitish. Fleshy, silky. Stem short, almost hollow, thickened at base.
35A	...	...	...	...	...	Q.	...	Ground ...	Spores globose.
36	...	S.A.	...	V.	...	...	...	Charred ground, under <i>Eucalyptus</i>	Rather fleshy, mealy, rufous with a tawny tinge. Stem slender, cylindrical
37	...	...	...	...	...	Q.	...	Sandy ground ...	Fleshy, pale ochre, variegated with darker concentric scales.
38	...	...	T.	V.	N.S.W.	Q.	B	Pastures ...	Large and tall, very shaggy, brownish. Fleshy, enticle thick and torn up into broad evanescent scales. <i>Edible</i> .
39	...	...	...	V.	...	...	B	Shady pastures ...	Very large. Fleshy, grey, enticle thin, broken into persistent brown scales. <i>Edible</i> .

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
3. LEPIDOTA.—Pers.,					
40	30	V. 119	<i>L. rhizobola</i> ...	Berk., Hook., Lond. Journ. IV. 42 (1845)	Bulbous-stalked lepiota ...
41	41	" 204	<i>L. rhyparophora</i> ...	Berk. and Br., Linn. Journ. XI. 500 (1871)	Spot-bearing lepiota ...
42	22	IX. 28	<i>L. rhytipelta</i> ...	F. v. M., Linn. Soc. N.S.W. 104 (1882)	Wrinkle-shield lepiota ...
43	26	" 16	<i>L. stenophylla</i> ...	Cooke and Mass., Grev. XV. 98 (1887)	Narrow-gilled lepiota ...
44	27	V. 237	<i>L. subclypeolaria</i> ...	Berk. and Curt., Linn. Journ. X. 283 (1869)	Sub-clypeolate lepiota ...
4. SCHULZERIA.—Bres,					
45	45	IX. 45	<i>S. revocans</i> ...	Cooke and Mass., Grev. XVIII. 2 (1889)	Recalling schulzeria ...
5. ARMILLARIA.—Fries,					
46	48	IX. 49	<i>A. fulgens</i> ...	Cooke and Mass., Grev. XVIII. 2 (1889)	Shining armillaria ...
47	47	V. 289	<i>A. mellea</i> ...	Fries, Hym. Eur. 44 (1874)	Honey-coloured armillaria ...
48	46	" 265	<i>A. robusta</i> ...	Fries, S.M. I. 26 (1821)	Robust armillaria ...
48A	"	" "	<i>A. robusta</i> , var. <i>subannulata</i>	Batsch, Consp. f. 17 (1783)	Smaller-ringed armillaria ...
6. TRICHOLOMA.—Fries,					
49	55	V. 415	<i>T. cerinum</i> ...	Fries, S.M. I. 89 (1821)	Wax-coloured tricholome ...
50	56	" 466	<i>T. civile</i> ...	Fries, Icon. t. 42, f. 1 (1867)	Civil tricholome ...
51	51	IX. 52	<i>T. coarctatum</i> ...	Cooke and Mass., Grev. XVIII. 2 (1889)	Compressed tricholome ...
52	53	V. 387	<i>T. cuneifolium</i> ...	Fries, S.M. I. 99 (1821)	Wedge-gill tricholome ...
53	59	" 485	<i>T. humile</i> ...	Fries, S.M. I. 51 (1821)	Humble tricholome ...
54	58	" 480	<i>T. melaleucum</i> ...	Fries, S.M. I. 114 (1821)	Black and white tricholome ...
55	50	" 329	<i>T. muculentum</i> ...	Berk., Hook., Lond. Journ. IV. 43 (1845)	Glutinous tricholome ...
56	57	" 470	<i>T. nudum</i> ...	Fries, Hym. Eur. 72 (1874)	Naked-margined tricholome ...
57	60	" 488	<i>T. persicinum</i> ...	Fries, S.M. I. 52 (1821)	Peach-coloured tricholome ...
58	61	" 501	<i>T. putidum</i> ...	Fries, Epier. 54 (1838)	Fetid tricholome ...
59	49	" 326	<i>T. resplendens</i> ...	Fries, Mon. I. 55 (1857)	Resplendent tricholome ...
60	52	" 344	<i>T. rutilans</i> ...	Fries, S.M. I. 41 (1821)	Red-haired tricholome ...
61	54	" 401	<i>T. sulphureum</i> ...	Fries, S.M. I. 110 (1821)	Sulphur-coloured tricholome ...
7. CLITOCYBE.—Fries,					
62	76	IX. 106	<i>C. canaliculata</i> ...	Cooke and Mass., Grev. XVIII. 2 (1889)	Channelled clitocybe ...
63	63	V. 553	<i>C. cerussata</i> ...	Fries, S.M. I. 92 (1821)	White clitocybe ...
64	62	" 517	<i>C. curtipes</i> ...	Fries, S.M. I. 88 (1821)	Short-stalked clitocybe ...
65	73	" 637	<i>C. expallens</i> ...	Fries, Mon. I. 129 (1857)	Bleaching clitocybe ...
66	71	" 621	<i>C. flaccida</i> ...	Fries, S.M. I. 81 (1821)	Flaccid clitocybe ...
67	64	" 580	<i>C. fumosa</i> ...	Fries, Hym. Eur. 91 (1874)	Smoky clitocybe ...
68	68	" 612	<i>C. gilva</i> ...	Fries, Hym. Eur. 95 (1874)	Yellowish-tan clitocybe ...
69	66	" 595	<i>C. infundibuliformis</i> ...	Fries, Hym. Eur. 93 (1874)	Funnel-shaped clitocybe ...
70	70	" 619	<i>C. inversa</i> ...	Fries, Hym. Eur. 96 (1874)	Inverted clitocybe ...
71	75	" 720	<i>C. laecata</i> ...	Fries, S.M. I. 106 (1821)	Scaling-wax clitocybe ...
72	67	IX. 75	<i>C. myriophylla</i> ...	Cooke and Mass., Grev. XVI. 113 (1888)	Myriad-gilled clitocybe ...
73	74	V. 643	<i>C. pruinosa</i> ...	Laseh, in Fries, Epier. 75 (1836)	Pruinose clitocybe ...
74	65	" 572	<i>C. schizophylla</i> ...	Berk., Fl. Tasm. II. 242 (1860)	Split-gilled clitocybe ...
75	69	IX. 96	<i>C. subsplendens</i> ...	Cooke and Mass., Grev. XVIII. 2 (1889)	Shining clitocybe ...
76	72	V. 632	<i>C. tuba</i> ...	Fries, Epier. 72 (1838)	Trumpet clitocybe ...
8. COLLYBIA.—Fries,					
77	83	V. 761	<i>C. butyracea</i> ...	Fries, Hym. Eur. 113 (1874)	Buttery collybia ...
78	92	" 865	<i>C. coagulata</i> ...	Berk. and Br., Linn. Trans. II. 53 (1883)	Coagulated collybia ...
79	93	" 871	<i>C. dryophila</i> ...	Fries, Hym. Eur. 122 (1874)	Wood-loving collybia ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Tent. Disp. 68 (1797).—Agaricus—continued.									
40	W.A.	...	...	...	...	...	...	Ground ...	Fleshy, shining, white, centre ornamented with pyramidal wart-like scales. Eaten largely by the smaller marsupials.
41	...	...	...	...	N.S.W.	...	...	Ground ...	Small, white, marked with brownish spots. Stem club shaped.
42	W.A.	...	...	V.	...	Q.	...	Ground ...	Fleshy, at first umber, then white. Stem rather bulbous at base.
43	...	...	...	V.	...	Q.	...	Ground ...	Fleshy, brownish, with depressed persistent scales. Stem long, bulbous.
44	...	...	...	V.	...	...	...	Ground, roots of trees, or dead wood	Thin, white, umbo or boss dusky. Stem smooth, white.
Trid. 7 (1881).—Agaricus.									
45	...	...	...	...	...	Q.	...	Gardens ...	Somewhat fleshy, pallid, spotted chiefly about disc, with darker scales.
S.M. I. 26 (1821).—Agaricus.									
46	...	...	...	...	...	Q.	...	Sandy soil	Bright golden yellow, smooth, shining. Stem erect, slender, hollow.
47	...	S.A.	...	V.	N.S.W.	Q.	B	Dead stumps	Fleshy, honey brown, scaly fibrous. In tufts on stumps. Very common. <i>Edible</i> .
48	...	...	...	V.	...	...	B	Woods, &c.	Robust. Fleshy, compact, brown. Stem solid, short, tapering downwards.
48A	...	...	...	V.	...	...	...	Woods, &c.	Smaller than typical form, with smaller ring.
S.M. I. 36 (1821).—Agaricus.									
49	...	...	...	V.	...	...	B	Lawns, &c.	Fleshy, brown or yellow. Stem stuffed, grooved, with fibrils. Rare.
50	...	...	...	V.	...	Q.	B	Pine woods	Fleshy, soft, moist, ash coloured, becoming pallid. Rare.
51	...	...	...	V.	...	...	...	Sandy soil	Pressed together and deformed. Fleshy, viscid, tan coloured, cracked when dry.
52	...	...	...	V.	...	...	B	Pastures ...	Small, very brittle. Rather fleshy, buff. Stem hollow, tapering downwards. Common.
53	...	...	...	V.	...	...	B	Ground, among grass	Fleshy, blackish brown to ash grey. Stem stuffed, powdery, and shaggy. Common.
54	...	S.A.	...	V.	...	...	B	Ground ...	Fleshy, moist, changing colour dingy black then livid brown. Gills white. Very common.
55	W.A.	...	...	...	...	...	...	Among moss	Rather fleshy, glutinous, whitish. Stem solid, viscid.
56	W.A.	...	T.	V.	...	...	B	Woods, &c., among dead leaves	Fleshy, rather thin, moist, changing colour. Rare. <i>Edible</i> .
57	...	...	...	V.	...	...	...	Grassy places	Fleshy, moist, thin at the naked margin. Stem stuffed, cartilaginous.
58	...	...	...	V.	...	...	B	Firwoods ...	Somewhat fleshy, olive grey, hoary when dry. Odour mealy, rancid.
59	...	...	...	...	...	Q.	B	Shady places	White. Fleshy, shining when dry. Odour agreeable.
60	W.A.	S.A.	...	...	...	...	B	Pine stumps	Fleshy, with red or purplish down. Odour strong. Common.
61	W.A.	...	...	...	...	...	B	Woods ...	Fleshy, more or less sulphur coloured. Odour disagreeable.
S.M. I. 78 (1821).—Agaricus, Laccaria.									
62	...	...	...	...	...	Q.	...	Under <i>Casuarina</i> (Sheoak) trees	Somewhat membranaceous, velvety, bright tawny, with radiating channels.
63	...	...	...	V.	...	Q.	B	Woods ...	White. Fleshy, moist. Stem spongy, elastic. Common. <i>Edible</i> .
64	...	...	T.	...	...	...	...	Grassy places	Rather fleshy, brown to livid. Stem solid, short, rigid.
65	...	...	...	V.	...	...	B	Ground ...	Fleshy to membranaceous, becoming tawny, when dry clay coloured. <i>Edible</i> .
66	...	S.A.	...	V.	...	...	B	Firwoods ...	Rather fleshy, flaccid, funnel shaped, bright brown, becoming pale.
67	W.A.	...	...	...	...	...	B	Woods, waste ground	Rigid. Fleshy, smoky, turning pale. <i>Edible</i> .
68	W.A.	...	...	...	...	...	B	Pine woods	Fleshy, moist, yellowish tan. Stem fleshy, solid, stout. <i>Edible</i> .
69	...	S.A.	...	V.	...	...	B	Fields and woods, among moss	Fleshy, downy, funnel shaped, flaccid, pale-tan colour or cinnamon. Common. <i>Edible</i> .
70	...	...	T.	V.	...	...	B	Woods ...	Fleshy, fragile, brownish red at first, then tan coloured. Margin inverted.
71	...	...	T.	V.	N.S.W.	Q.	B	Woods ...	Tall and slender. Membranaceous, red, brown, or amethyst, mealy. Very common. <i>Edible</i> .
72	...	...	...	V.	...	...	...	Grassy places	Fleshy, shining, tawny, grey or ochrey, white. Stem solid.
73	...	...	...	...	...	Q.	B	Pine woods and on trunks	Slender, rigid, inodorous. Fleshy to membranaceous, brownish or ash coloured, sprinkled with a greyish bloom. <i>Edible</i> .
74	...	...	T.	...	...	...	...	Rotten wood	In tufts. Gills splitting at the edge. Stem stringy.
75	...	...	...	...	...	Q.	...	Among grass in garden	Somewhat fleshy, shining, rufous or yellowish. Stem solid.
76	...	...	...	V.	...	...	B	Among leaves, chiefly of Pines	White. Fleshy, moist, shining with a whitish silky lustre. Stem soon hollow.
S.M. I. 129 (1821).—Agaricus, Amanita, Marasmius.									
77	...	...	...	V.	...	...	B	Woods ...	Small. Fleshy, changing colour, flesh becoming white. Very common.
78	...	...	...	...	...	Q.	...	Ground ...	Cream colour, yellow when dry. Stem slender, twisted, yellow.
79	...	...	...	...	...	Q.	B	Among leaves in woods	Somewhat fleshy, turning pale bay red, yellowish, clay coloured, white. Very common.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
8. COLLYBIA.—Fries,					
80	79	V. 734	<i>C. eradicata</i> ...	Kalch., Grev. VIII. 151 (1880) ...	Non-rooting collybia ...
81	91	" 840	<i>C. esculenta</i> ...	<i>Fries</i> , Hym. Eur. 121 (1874) ...	Esculent collybia ...
82	81	" 748	<i>C. fusipes</i> ...	<i>Fries</i> , Hym. Eur. 111 (1874) ...	Spindle-stalked collybia ...
83	77	" 807	<i>C. laceatina</i> ...	Berk., Linn. Journ. XVIII. 383 (1881) ...	Sealing-wax collybia ...
84	99	" 918	<i>C. lacerata</i> ...	Lasch, in <i>Fries</i> , Hym. Eur. 127 (1874) ...	Torn collybia ...
85	86	" 831	<i>C. lepidopoda</i> ...	<i>Fries</i> , in Pl. Preiss. II. 131 (1846) ...	Scaly-stalked collybia ...
86	80	" 735	<i>C. longipes</i> ...	<i>Fries</i> , Hym. Eur. 110 (1874) ...	Long-staked collybia ...
87	87	" 808	<i>C. morula</i> ...	Berk., Fl. Tasm. II. 243 (1860) ...	Mulberry-brown collybia ...
88	95	"	<i>C. nivosa</i> ...	Berk., Cuban Fungi III. ...	Snowy collybia ...
89	90	V. 839	<i>C. nummularia</i> ...	<i>Fries</i> , Epier. 91 (1838) ...	Coin-like collybia ...
90	82	IX. 116	<i>C. olivaceo-alba</i> ...	Cooke and Mass., Grev. XV. 93 (1887) ...	Olive-white collybia ...
91	97	V. 904	<i>C. ozes</i> ...	<i>Fries</i> , Epier. 95 (1838) ...	Smelling collybia ...
91A	"	IX. 139	<i>C. ozes</i> , var. <i>crassipes</i> ...	Cooke and Mass., Grev. XV. 93 (1887) ...	Thick-stalked collybia ...
92	98	V. 907	<i>C. plexipes</i> ...	<i>Fries</i> , S.M. I. 146 (1821) ...	Twisted-stalked collybia ...
93	78	" 728	<i>C. radicata</i> ...	<i>Fries</i> , Hym. Eur. 109 (1874) ...	Rooting collybia ...
93A	"	"	<i>C. radicata</i> , var. <i>superbicus</i> ...	Berk., Hook., Lond. Journ. IV. 43 (1845) ...	Superb collybia ...
94	96	" 899	<i>C. rancida</i> ...	<i>Fries</i> , S.M. I. 141 (1821) ...	Rancid collybia ...
95	430	" 784	<i>C. rheicolor</i> ...	<i>Sacc.</i> Syll. I. 214 (1887) ...	Rhubarb-coloured collybia ...
96	88	" 826	<i>C. tuberosa</i> ...	<i>Fries</i> , Hym. Eur. 119 (1874) ...	Tuberous collybia ...
97	100	" 929	<i>C. tylicolor</i> ...	<i>Fries</i> , S.M. I. 132 (1821) ...	Greyish collybia ...
98	94	IX. 115	<i>C. veluticeps</i> ...	Cooke and Mass., Grev. XVI. 30 (1887) ...	Velvet-head collybia ...
99	85	V. 773	<i>C. velutipes</i> ...	<i>Fries</i> , Hym. Eur. 115 (1874) ...	Velvet-stem collybia ...
100	89	" 836	<i>C. xanthopoda</i> ...	<i>Fries</i> , Hym. Eur. 120 (1874) ...	Yellow-stalked collybia ...
101	84	" 768	<i>C. xylophila</i> ...	Weinm., in Linn. X. 54 (1836) ...	Timber-loving collybia ...
9. MYCENA.—Pers.,					
102	109	V. 1050	<i>M. ætites</i> ...	<i>Fries</i> , Epier. 110 (1838) ...	Beaked mycena ...
103	107	" 1037	<i>M. atro-cyanea</i> ...	<i>Fries</i> , S.M. I. 147 (1821) ...	Dark-blue mycena ...
104	123	" 1152	<i>M. capillaris</i> ...	<i>Fries</i> , S.M. I. 160 (1821) ...	Thread-stalked mycena ...
105	432	" 991	<i>M. coharens</i> ...	<i>Fries</i> , Epier. 105 (1838) ...	Cohering mycena ...
106	120	" 147	<i>M. corticola</i> ...	<i>Fries</i> , S.M. I. 150 (1821) ...	Bark-growing mycena ...
107	113	" 1088	<i>M. crinalis</i> ...	Berk., Hook., Lond. Journ. IV. 44 (1845) ...	Hair-like mycena ...
108	111	" 1071	<i>M. debilis</i> ...	<i>Fries</i> , Epier. 112 (1838) ...	Tender mycena ...
109A	110	"	<i>M. filipes</i> , var. <i>acutata</i> ...	Kalch., Linn. Soc. N.S.W. 104 (1882) ...	Acutely conical mycena ...
110	104	IX. 45	<i>M. flavovirens</i> ...	Cooke and Mass., Grev. XIX. 45 (1890) ...	Yellowish-green mycena ...
111	105	V. 1002	<i>M. galericulata</i> ...	<i>Fries</i> , Hym. Eur. 138 (1874) ...	Little-cap mycena ...
112	115	" 1097	<i>M. hæmatopoda</i> ...	<i>Fries</i> , S.M. I. 149 (1821) ...	Dark-red juiced mycena ...
113	122	" 1148	<i>M. hiemalis</i> ...	Osbeck, in Retz. Supp. 19 (1805) ...	Winter mycena ...
114	119	" 1135	<i>M. interrupta</i> ...	Berk., Fl. Tasm. II. 243 (1860) ...	Interrupted mycena ...
115	124	" 1154	<i>M. juncicola</i> ...	<i>Fries</i> , Hym. Eur. 154 (1874) ...	Rush-growing mycena ...
116	108	" 1041	<i>M. leptcephala</i> ...	<i>Fries</i> , Hym. Eur. 141 (1874) ...	Delicate-head mycena ...
117	102	" 952	<i>M. pura</i> ...	<i>Fries</i> , Hym. Eur. 133 (1874) ...	Pure mauve-cap mycena ...
118	101	" 944	<i>M. rosella</i> ...	<i>Fries</i> , S.M. I. 151 (1821) ...	Rose-coloured mycena ...
119	116	" 1100	<i>M. sanguinolenta</i> ...	<i>Fries</i> , Hym. Eur. 148 (1874) ...	Light red juiced mycena ...
120	103	" 983	<i>M. Silenus</i> ...	Berk. and Br., Linn. Journ. XI. 524 (1871) ...	Bacchanalian mycena ...
121	112	" 1080	<i>M. speirea</i> ...	<i>Fries</i> , S.M. I. 159 (1821) ...	Twisted mycena ...
122	117	" 1124	<i>M. stylobates</i> ...	<i>Fries</i> , Hym. Eur. 150 (1874) ...	Pillar-shaped mycena ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
80	...	...	...	V.	N.S.W.	...	...	Ground ...	Resembling <i>C. radicata</i> , but not rooting.
81	...	...	...	V.	...	...	B	Pastures ...	Small, in clusters, buffish. Somewhat fleshy. <i>Edible</i> .
82	...	...	...	...	...	...	B	Stumps ...	Fleshy, dull vinous brown or chestnut. <i>Edible</i> .
83	...	...	...	...	...	Q.	...	Dead wood among leaves	Pale fleshy red, margin grooved. Stem paler, fibrillose.
84	...	...	...	V.	...	...	B	Pine woods	Fleshy to membranaceous, moist, streaked brown, lacerated when old.
85	W.A.	...	...	...	...	...	...	Ground ...	Rather fleshy, orange, stem rough with scales.
86	...	...	...	V.	...	Q.	B	Old stumps, &c.	Fleshy, dry, slightly velvety, tan brown. Stem stuffed, tall.
87	...	...	T.	V.	...	...	...	Dead wood	Purple red or dark mulberry brown. Fleshy. Stem horizontal, rough.
88	...	...	...	V.	...	...	...	Logs	Thin, whitish. Stem slender, smooth, solid, white, rooting copiously.
89	...	...	...	V.	...	Q.	B	Among leaves, on wood, &c.	Beautiful, white becoming pale, variegated with light yellow and red. Rather fleshy.
90	...	S.A.	...	V.	...	...	...	Ground under <i>Casuarina</i> (Sheoak)	Fleshy, shining, dark sooty olive. Stem smooth, whitish, black below.
91	...	...	...	V.	...	...	B	Pine leaves	Fleshy, ashy brown when moist, pale clayey brown when dry. Strong scented, odour of meal.
91A	...	S.A.	...	...	...	...	...	Low damp ground	Conical to bell shaped. Stem tapering upwards, umber.
92	...	...	...	V.	...	...	B	Trunks	Sub-membranaceous, grey. Stem hollow, silky fibrous. Inodorous.
93	W.A.	...	T.	V.	...	Q.	B	Ground around bases of stems	Fleshy, glutinous, with long, pale, slender, twisted, rooting stems.
93A	W.A.	...	...	...	...	...	...	Ground	Brown, and stem clad at base with velvety scurf.
94	...	...	...	V.	...	...	B	Under trees	Strong scented. Rather fleshy, grey, whitish, silky. Stem rooting, smooth.
95	...	...	...	...	...	Q.	...	Trunks	Rhubarb colour. Thin, and stem clothed with a velvety down.
96	...	...	...	...	...	Q.	B	Putrid Agarics, such as <i>Russula</i> , &c., and on ground	Slightly fleshy, white. Root springing from sclerotoid tuber. Common.
97	...	S.A.	...	V.	...	...	B	Woods	Rather fleshy, ash colour. Stem hollow, powdery.
98	...	...	...	V.	...	...	...	Fern-tree Gully	Fleshy, velvety, liver coloured. Stem short, pale upwards.
99	...	...	...	V.	...	...	B	Logs and trunks of trees—Willow, Beech, &c.	Fleshy, viscid, tawny yellow or fawn. Stem stuffed, velvety, dark bay. Common.
100	...	...	...	V.	...	...	B	About stumps of trees	Rather fleshy, becoming pale. Stem yellow, and rooting at base.
101	...	...	...	V.	...	...	B	Stumps	Rather fleshy, whitish or clay coloured, bell shaped. Stem hollow.

Tent. Disp. 69 (1797).—*Agaricus*, *Marasmius*.

102	...	S.A.	...	V.	...	...	B	Among mosses	Fragile. Membranaceous, brownish, growing pale, with broad obtuse prominent umbo.
103	...	...	T.	...	...	...	B	Ground	Fragile, inodorous. Membranaceous, brownish, then grey, becoming bluish.
104	...	...	T.	V.	...	...	B	Dead leaves in woods	Very delicate, white. Bell shaped, smooth. Stem thread-like, smooth.
105	...	...	T.	...	...	...	B	Ground	Rather fleshy, velvety, cinnamon brown, growing pale. Stem horny, rigid.
106	...	...	...	V.	N.S.W.	Q.	B	Mossy bark	Colour various—reddish brown, blue, or ash coloured. Stem incurved, scurfy.
107	W.A.	...	...	...	...	...	...	Decayed wood	Very delicate. Membranaceous, white. Stem thread-like, brown.
108	...	...	...	V.	N.S.W.	...	B	Woods	Tender. Membranaceous, brownish. Stem thread-like.
109A	...	...	...	V.	N.S.W.	...	...	Ground	Dark, ash coloured, and acutely conical. Stem thread-like, rooting.
110	...	...	...	V.	...	...	...	Tree ferns	Membranaceous, yellowish green. Stem slender, erect, hollow.
111	...	...	T.	V.	...	...	B	Trunks of trees	Sub-membranaceous, flesh coloured, drab, or various. Densely clustered. Common.
112	...	...	...	V.	...	...	B	Old dead trunks	Fleshy. Stem yielding a dark-red juice, rigid, powdery.
113	...	S.A.	...	V.	...	...	B	Trunks of trees	Thin, brighter coloured than <i>M. corticola</i> , hardly ashy brown.
114	...	...	T.	...	...	...	...	Bark	Rather thick, livid, gelatinously fleshy. Gills descending interruptedly into flesh of cap.
115	...	...	...	V.	...	...	B	Dead rushes in bogs.	Very delicate. Somewhat red. Stem thread-like, smooth, brownish.
116	...	...	...	V.	...	...	B	Trunks and ground	Fragile, with nitrous odour. Sub-membranaceous, furrowed, frosted.
117	...	...	...	V.	...	...	B	Ground in woods	Strong smelling, odour of radish. Rather fleshy, violet or roseate, becoming pallid and variously coloured.
118	...	...	...	V.	...	...	B	Among fir leaves	Rose coloured. Membranaceous, boss obtuse. Stem thin, juiceless.
119	...	S.A.	...	V.	...	...	B	Among leaves and damp moss	Delicate pale red, becoming brown, membranaceous. Stem yielding pale-red juice
120	...	...	...	...	...	Q.	...	Dead wood	Small. Fleshy, red to vinous brown. Stem short, hollow.
121	...	S.A.	...	V.	N.S.W.	...	B	Mossy trunks	Membranaceous, greyish brown, disc darker. Stem thread-like.
122	...	...	...	V.	...	Q.	B	Fern, twigs, &c.	Membranaceous, white, somewhat hairy. Stem thread-like.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number	Scientific Name.	Authority for Name.	English Name.
9. MYCENA.—Pers.,					
123	121	IX. 143	<i>M. subcorticalis</i> ...	Cooke and Mass., <i>Grev.</i> XV. 93 (1887) ...	Subcortical mycena ...
124	118	V. 1129	<i>M. tenerrima</i> ...	Fries, <i>Hym. Eur.</i> 151 (1874) ...	Very tender mycena ...
125	106	„ 1925	<i>M. trachycephala</i> ...	F. v. M. and Kalch., <i>Grev.</i> VIII. 151 (1880) ...	Rough-headed mycena ...
126	114	„ 1090	<i>M. tuberigena</i> ...	Berk., <i>Linn. Journ.</i> XIII. 156 (1873) ...	Tuber-bearing mycena ...
10. HIATULA.—Fries,					
127	359	V. 1168	<i>H. Wynnæ</i> ...	Berk. and Br., <i>Ann. Nat. Hist.</i> , III., 5. 206 (1879)...	Wynne's hiatula. Green-light fungus
11. OMPHALIA.—Fries.					
128	136	V. 1247	<i>O. carneo-rufula</i> ...	Berk., <i>Fl. Tasm.</i> II. 243 (1860) ...	Fleshy-red omphalia ...
129	126	„ 1181	<i>O. dumosa</i> ...	Fries, <i>Hym. Eur.</i> 155 (1874) ...	Thicket-loving omphalia ...
130	129	„ 1205	<i>O. epichysium</i> ...	Fries, S.M. I. 169 (1821) ...	Watery omphalia ...
131	140	„ 1283	<i>O. fibula</i> ...	Fries, <i>Hym. Eur.</i> 164 (1874) ...	Pin-like omphalia ...
132	137	„ 1265	<i>O. flavo-crocea</i> ...	Berk., <i>Fl. Tasm.</i> II. 244 (1860) ...	Bright-yellow omphalia ...
133	133	IX. 179	<i>O. glaucescens</i> ...	Kalch., <i>Linn. Soc. N.S.W.</i> 105 (1882) ...	Sage-green omphalia ...
134	141	V. 1286	<i>O. gomphomorpha</i> ...	Berk., <i>Linn. Journ.</i> XVIII. 383 (1881) ...	Club-shaped omphalia ...
135	142	„ 1289	<i>O. gracillima</i> ...	Weinm., <i>Ross</i> 121 (1836) ...	Slender omphalia ...
136	128	„ 1196	<i>O. holochlora</i> ...	Berk. and Br., <i>Linn. Journ.</i> XI. 525 (1871) ...	Green omphalia ...
137	125	„ 1179	<i>O. hydrogramma</i> ...	Fries, S.M. I. 169 (1821) ...	Water-line omphalia ...
138	143	„ 1313	<i>O. integrella</i> ...	Fries, <i>Hym. Eur.</i> 165 (1874) ...	Perfect omphalia ...
139	144	„ 1321	<i>O. Muelleriana</i> ...	Berk., in <i>Cooke's Handb. Aust. Fungi</i> 30 (1892) ...	Mueller's omphalia ...
140	134	„ 1239	<i>O. muralis</i> ...	Fries, <i>Hym. Eur.</i> 160 (1874) ...	Wall omphalia ...
141	130	„ 1208	<i>O. oniscus</i> ...	Fries, S.M. I. 172 (1821) ...	Grey omphalia ...
142	131	„ 1215	<i>O. pumilio</i> ...	Kalch., <i>Grev.</i> VIII. 151 (1880) ...	Dwarf omphalia ...
143	127	„ 1199	<i>O. pyxidata</i> ...	Fries, S.M. I. 164 (1821) ...	Box-like omphalia ...
144	132	„ 1216	<i>O. scyphiformis</i> ...	Fries, <i>Hym. Eur.</i> 159 (1874) ...	Goblet-shaped omphalia ...
145	139	„ 1282	<i>O. setipes</i> ...	Fries, <i>Hym. Eur.</i> 164 (1874) ...	Hairy-stalked omphalia ...
146	135	„ 1241	<i>O. umbellifera</i> ...	Fries, <i>Hym. Eur.</i> 160 (1874) ...	Umbrella-like omphalia ...
147	138	„ 1279	<i>O. umbratilis</i> ...	Fries, <i>Epier.</i> 127 (1838) ...	Shade-loving omphalia ...
12. PLEUROTUS.—Fries,					
148	147	V. 1346	<i>P. abbreviatus</i> ...	Kalch., <i>Grev.</i> VIII. 152 (1880) ...	Abbreviated pleurote ...
149	170	„ 1444	<i>P. affixus</i> ...	Berk., <i>Hook. Lond. Journ.</i> VII., 573 (1848) ...	Affixed pleurote ...
150	180	„ 1504	<i>P. applicatus</i> ...	Fries, <i>Hym. Eur.</i> 180 (1874) ...	Sessile pleurote ...
151	179	„ 1492	<i>P. atro-ceruleus</i> ...	Fries, S.M. I. 190 (1821) ...	Dark-blue pleurote ...
152	188	IX. 187	<i>P. australis</i> ...	Cooke and Mass., <i>Grev.</i> XV. 93 (1887) ...	Southern pleurote ...
153	177	V. 1487	<i>P. barsæiformis</i> ...	Berk., <i>Fl. Tasm.</i> II. 245 (1860) ...	Porch-like pleurote ...
154	156	„ 1400	<i>P. caudescens</i> ...	F. v. M., <i>Linn. Journ.</i> XIII. 157 (1873) ...	Glowing pleurote ...
155	169	„ 1442	<i>P. caryophyllus</i> ...	Berk., <i>Linn. Journ.</i> XIII. 157 (1873) ...	Clove-like pleurote ...
156	167	IX. 200	<i>P. chaetophyllus</i> ...	Sacc. <i>Hedw.</i> 126 (1889) ...	Hairy-gill pleurote ...
157	187	V. 1527	<i>P. chionus</i> ...	Pers. <i>M. Eur.</i> 3 (1828) ...	Snow-white pleurote ...
158	159	IX. 190	<i>P. clitocyboides</i> ...	Cooke and Mass., <i>Grev.</i> XV. 98 (1887) ...	Clitocybe-like pleurote ...
159	145	V. 1322	<i>P. corticatus</i> ...	Fries, S.M. I. 179 (1821) ...	Corticated pleurote ...
160	183	„ 1511	<i>P. diversipes</i> ...	Berk., <i>Fl. Tasm.</i> II. 244 (1860) ...	Variable-stalked pleurote ...
161	164	„ 1423	<i>P. Eucalyptorum</i> ...	Fries, <i>Pl. Preiss.</i> II. 131 (1846) ...	Eucalyptus pleurote ...
162	189	„ p. 387	<i>P. euphyllus</i> ...	Berk., in <i>Handb. N.Z. Flora</i> 755 (1867) ...	Broad-gilled pleurote ...
163	172	„ 1449	<i>P. flabellatus</i> ...	Berk. and Br., <i>Linn. Journ.</i> XI. 528 (1871) ...	Fan-shaped pleurote ...
164	149	„ 1369	<i>P. Gardneri</i> ...	Berk., <i>Hook., Journ.</i> II., 427 (1840) ...	Gardner's pleurote ...
165	161	„ 1409	<i>P. Guilfoylei</i> ...	Berk., <i>Linn. Journ.</i> XIII. 158 (1873) ...	Guilfoyle's pleurote ...
166	150	„ 1370	<i>P. illuminans</i> ...	F. v. M., <i>Linn. Journ.</i> XIII. 157 (1873) ...	Luminous pleurote ...
167	168	„ 1429	<i>P. imberbis</i> ...	Kalch., <i>Grev.</i> VIII. 152 (1880) ...	Beardless pleurote ...
168	148	„ 1347	<i>P. lacticolor</i> ...	Kalch., <i>Grev.</i> VIII. 151 (1880) ...	Bright-coloured pleurote ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Tent. Disp. 69 (1797).— <i>Agaricus</i> , <i>Marasmius</i> — <i>continued</i> .									
123	...	S.A.	...	...	...	...	...	Log of <i>Banksia</i> (Native Honeysuckle)	Thin, lilac, disc brick red. Stem ascending, thin, hollow.
124	...	...	...	...	...	Q.	B	Fir cones, sticks, &c.	White, very delicate. Cap frosted with senrfy granules. Stem hairy.
125	...	...	...	V.	...	...	...	Rotten trunks	Membranaceous, ashy brown, covered with papillæ. Stem thread-like.
126	...	...	...	V.	...	...	...	Ground	Tender and small, white. Stem thread-like, arising from sclerotium.
Nov. Symb. 27 (1851).									
127	...	...	...	V.	...	Q.	B	Ground	White. Tender, luminous, emitting a greenish light. Stem slender.
S.M. I. 162 (1821).— <i>Agaricus</i> .									
128	...	...	T.	...	...	...	...	Rotten wood	Rather fleshy, pale red. Stem flexuous, stuffed.
129	...	...	...	V.	...	...	...	Woods	Rather membranaceous, brick red. Stem hollow, smooth. Rare.
130	...	...	T.	...	...	...	...	Rotten wood	Soft. Membranaceous, sooty, ash coloured, silky or scaly.
131	W.A.	S.A.	...	V.	...	Q.	B	Among moss in moist places	Tiny. Membranaceous, nearly orange colour or orange fawn. Common.
132	...	...	T.	...	...	...	...	Branches	Yellow, and gills saffron yellow. Stem elongated, solid.
133	...	...	...	V.	N.S.W.	...	...	Ground	Small. Grey, sage green. Stem thread-like, greenish yellow.
134	...	...	...	...	...	Q.	...	Ground in tufts	Club-shaped, lurid. Stem thickened upwards, reddish brown.
135	...	...	...	V.	...	...	B	Marshy ground	Snow white. Membranaceous, furrowed. Stem thread-like.
136	...	...	...	...	...	Q.	...	Dead wood	Membranaceous, yellow brown, reddish brown when dry.
137	...	...	...	V.	N.S.W.	...	B	Dead leaves and moist places	Sub-membranaceous, livid; margin spreading, streaked. Stem hollow.
138	...	...	T.	...	...	...	B	Decayed wood	White, fragile. Membranaceous, pellucid. Stem very slender, short.
139	...	...	...	V.	...	...	...	Ground	Smooth, tawny. Stem elongated, smooth, or streaked lengthwise.
140	...	S.A.	...	V.	...	...	B	Ground, banks, and walls	Sub-membranaceous, reddish brown, radiately striate.
141	...	...	...	...	...	Q.	B	Swampy ground	Sub-membranaceous, dark ash coloured. Stem firm, partially hollow.
142	...	...	...	...	N.S.W.	Q.	...	Wood	Membranaceous, fawn coloured. Stem hollow, thin, curved.
143	...	S.A.	...	V.	...	...	B.	Among short grass, on lawns, &c.	Sub-membranaceous, brick red, funnel shaped. Stem stuffed at first. Common.
144	...	...	...	V.	...	Q.	...	Bare ground	White. Membranaceous. Stem rather hollow, short, thin.
145	...	...	...	V.	N.S.W.	...	...	Shady places	Membranaceous, brownish grey. Stem thread-like, downy at base.
146	W.A.	...	T.	V.	...	Q.	B	Swamps, exposed pastures, &c.	Fleshy to membranaceous, buff or variable in colour. Very common.
147	...	...	...	V.	...	...	B	Damp places	Sub-membranaceous, umber brown. Stem stuffed, tough.
S.M. I. 178 (1821).— <i>Agaricus</i> .									
148	...	...	...	...	N.S.W.	...	...	Wood	Entirely reddish brown. Stem shorter than diameter of cap.
149	...	...	T.	...	...	...	...	Bark of young <i>Eucalyptus amygdalina</i>	White. Cap at length attached by the side, cup shaped. Stem short.
150	W.A.	...	T.	V.	...	Q.	B	Dead fallen branches, and on <i>Eucalyptus viminalis</i>	Very small. Dark-ash colour. Sub-membranaceous, cup shaped.
151	W.A.	...	...	V.	...	...	B	Trunks	Fleshy, dark blue, rarely brown, downy.
152	...	S.A.	...	...	...	...	...	Roots of <i>Leptospermum</i> (Tea-tree)	Fleshy, umber. Stem rather lateral, thick, clad with white down.
153	...	...	T.	...	...	...	...	Rotting bark	Cap affixed behind, pouch-like, whitish. Stem short, almost smooth.
154	...	...	...	V.	...	...	...	Dead wood	Strongly phosphorescent. White, becoming dingy. Stem dilated above.
155	...	...	...	V.	N.S.W.	...	...	Wood	Pale-tawny colour. Fan shaped, and much lobed.
156	...	S.A.	...	...	...	...	...	Branches	Thin, white, tomentose, spoon shaped. Stem thick, wrinkled.
157	W.A.	...	...	V.	...	...	B	Wood or dung	Snow white. Very thin, woolly. Stem very short, bairy.
158	...	...	...	V.	...	...	...	Old fern stems	Thin, ochrey, becoming reddish.
159	...	...	...	...	...	Q.	B	Trunks, living and dead	Beautiful large species. Fleshy, greyish white. Stem firm.
160	...	...	T.	...	...	Q.	...	Rotten wood	Pellucid, covered with a gelatinous layer. Stem very variable.
161	W.A.	...	...	...	...	...	...	<i>Eucalyptus</i> bark	Fleshy, bay brown, clad with a rough wool, sessile, kidney shaped.
162	...	...	...	...	...	Q.	...	Wood	Pale chestnut. Stem none or obsolete. Gills broad.
163	...	...	...	...	...	Q.	...	Dead wood	Fan shaped, thin, white, becoming reddish. Adhering to wood by spongy base.
164	...	...	...	...	...	Q.	...	Petioles and half-rotten fronds of palms	Fleshy to leathery, yellow, funnel shaped. Stem short. Phosphorescent.
165	...	...	...	...	N.S.W.	Q.	...	Trunks	Whitish, very much wrinkled when dry, cap kidney shaped.
166	...	...	...	V.	N.S.W.	Q.	...	Dead wood	Phosphorescent. Tawny, smooth. Stem thick.
167	...	...	...	...	N.S.W.	...	...	Wood	Membranaceous, kidney shaped, sessile, horizontal.
168	...	...	...	...	N.S.W.	...	...	Wood (?)	Rather fleshy, golden yellow. Stem stuffed, naked.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
12. PLEUROTUS.—Fries,					
169	155	V. 1397	<i>P. lampas</i> ...	Berk., Hook., Lond. Journ. IV. 44 (1845) ...	Shining pleurote ...
170	181	" 1506	<i>P. lentienla</i> ...	Kalch., Grev. VIII. 151 (1880) ...	Freckled pleurote ...
171	166	" 1427	<i>P. limpidus</i> ...	Fries, Epier. 135 (1838) ...	Transparent pleurote ...
172	178	" 1488	<i>P. lividulus</i> ...	Berk. and Curt. Exp. No. 33 (1859) ...	Livid pleurote ...
173	151	" 1386	<i>P. luteo-aurantius</i> ...	Kalch., Grev. VIII. 151 (1880) ...	Orange-yellow pleurote ...
174	165	" 1425	<i>P. mitis</i> ...	Fries, S.M. I. 188 (1821) ...	Mild pleurote ...
175	154	" 1396	<i>P. nidiformis</i> ...	Berk., Hook., Lond. Journ. III. 185 (1844) ...	Nest-shaped pleurote ...
176	152	" 1390	<i>P. ostreatus</i> ...	Fries, Hym. Eur. 173 (1874) ...	Oyster-like pleurote ...
177	186	" 1523	<i>P. perpusillus</i> ...	Fries, S.M. I. 195 (1821) ...	Very small pleurote ...
178	162	" 1412	<i>P. petaloides</i> ...	Fries, S.M. I. 183 (1821) ...	Petal-like pleurote ...
179	157	" 1401	<i>P. phosphorus</i> ...	Berk., Hook., Lond. Journ. VII. 572 (1848) ...	Phosphorescent pleurote ...
180	153	IX. 196	<i>P. polyphemus</i> ...	Cooke and Mass., Grev. XVI. 72 (1888) ...	Variegated pleurote ...
181	163	V. 1416	<i>P. pulmonarius</i> ...	Fries, S.M. I. 187 (1821) ...	Lung-like pleurote ...
182	158	" 1405	<i>P. salignus</i> ...	Fries, Hym. Eur. 174 (1874) ...	Willow-sprout pleurote ...
183	176	" 1475	<i>P. scabriusculus</i> ...	Berk., Linn. Journ. XIII. 157 (1873) ...	Rough pleurote ...
184	174	" 1470	<i>P. semiliber</i> ...	Berk. and Br., Linn. Trans. II. 54 (1883) ...	Half-free pleurote ...
185	173	" 1469	<i>P. semisupinus</i> ...	Berk. and Br., Linn. Journ. XI. 529 (1871) ...	Semisupine pleurote ...
186	175	" 1473	<i>P. sordulentus</i> ...	Berk. and Br., Linn. Trans. II. 54 (1883) ...	Dirty-white pleurote ...
187	184	" 1518	<i>P. striatulus</i> ...	Fries, S.M. I. 193 (1821) ...	Striate pleurote ...
188	185	" 1522	<i>P. subbarbatus</i> ...	Berk. and Curt., Linn. Journ. X. 288 (1869) ...	Barbed pleurote ...
189	160	IX. 198	<i>P. sulceiceps</i> ...	Cooke and Mass., Grev. XVIII. 3 (1889) ...	Sulcate pleurote ...
190	182	V. 1510	<i>P. tasmanicus</i> ...	Berk., Fl. Tasm. II. 245 (1860) ...	Tasmanian pleurote ...
191	146	" 1343	<i>P. tephrophanus</i> ...	Berk., Fl. Tasm. II. 244 (1860) ...	Ash-growing pleurote ...
192	171	" 1445	<i>P. Thozetii</i> ...	Berk., Linn. Journ. XVIII. 383 (1881) ...	Thozet's pleurote ...
13. HYGROPHORUS.—Fries,					
193	379	IX. 216	<i>H. candidus</i> ...	Cooke and Mass., Grev. XVIII. 4 (1889) ...	White hygrophore ...
194	381	V. 1634	<i>H. ceraceus</i> ...	Fries, Epier. 330 (1838) ...	Waxy hygrophore ...
195	382	" 1637	<i>H. coccineus</i> ...	Fries, Epier. 330 (1838) ...	Scarlet hygrophore ...
196	384	" 1658	<i>H. conicus</i> ...	Fries, Epier. 331 (1838) ...	Conical hygrophore ...
197	376	" 1599	<i>H. flammans</i> ...	Berk., Linn. Journ. XIII. 160 (1873) ...	Flaming hygrophore ...
198	378	IX. 224	<i>H. gigasporus</i> ...	Cooke and Mass., Grev. XVI. 31 (1887) ...	Gigantic-spored hygrophore ...
199	375	" 228	<i>H. gilvus</i> ...	Kalch., Linn. Soc. N.S.W. 105 (1882) ...	Yellowish-tan hygrophore ...
200	873	V. 1570	<i>H. hypothejus</i> ...	Fries, Epier. 324 (1838) ...	Sulphur-yellow hygrophore ...
201	387	IX. 227	<i>H. Lewellinae</i> ...	Kalch., Linn. Soc. N.S.W. 105 (1882) ...	Lewellin's hygrophore ...
202	383	V. 1639	<i>H. micinatus</i> ...	Fries, Epier. 330 (1838) ...	Vermilion hygrophore ...
203	377	" 1600	<i>H. nigricans</i> ...	Berk., Linn. Journ. XIII. 160 (1873) ...	Blackening hygrophore ...
204	372	" 1555	<i>H. porphyrius</i> ...	Berk. and Br., Linn. Trans. II. 55 (1883) ...	Purple hygrophore ...
205	385	" 1677	<i>H. scarlatinus</i> ...	Kalch., Grev. VIII. 152 (1880) ...	Scarlet hygrophore ...
206	380	" 1628	<i>H. sciophanus</i> ...	Fries, Epier. 329 (1838) ...	Shadowy hygrophore ...
207	386	IX. 229	<i>H. subremotus</i> ...	Cooke and Mass., Grev. XVI. 113 (1888) ...	Sequestered hygrophore ...
208	374	V. 1590	<i>H. virgineus</i> ...	Fries, Epier. 327 (1838) ...	Virgin hygrophore ...
14. LACTARIUS.—D.C.,					
209	392	V. 1737	<i>L. pallidus</i> ...	Fries, Epier. 343 (1838) ...	Pale lactar ...
210	390	" 1727	<i>L. piperatus</i> ...	Fries, Epier. 340 (1838) ...	Peppery lactar ...
211	389	" 1720	<i>L. plumbeus</i> ...	Fries, Epier. 339 (1838) ...	Lead-colored lactar ...
212	388	" 1694	<i>L. stenophyllus</i> ...	Berk., Fl. Tasm. II. 248 (1860) ...	Narrow-gill lactar ...
213	391	" 1759	<i>L. subtomentosus</i> ...	Berk. and Rav., Ann. Nat. Hist. IV., 293 (1859) ...	Subtomentose lactar ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
S.M. I. 178 (1821).— <i>Agaricus</i> —continued.									
169	W.A.	...	T.	V.	...	...	...	Stems, languid, but not dead, of <i>Grevillea</i> (Silky oak)	Phosphorescent. Fleshy, tawny, turning black. Stem solid, sometimes splitting.
170	...	...	...	...	...	Q.	...	Trunks ...	Small. Olive brown, or powdered with white.
171	...	...	...	...	N.S.W.	...	B	Trunks ...	Rather fleshy, white, tapering behind into rudimentary stem.
172	...	...	...	...	N.S.W.	...	...	Dead branches ...	Becoming livid purple, clad with a powdery down, kidney or fan-shaped.
173	...	...	...	...	N.S.W.	...	...	Wood ...	Rather fleshy, orange yellow. Stem hollow, thin, short, curved upwards.
174	...	...	...	...	...	Q.	B	Dead wood—Pines, Firs, and Larches	Rather fleshy, growing pale, kidney shaped. Stem lateral, compressed, dilated upwards with white scales. Common.
175	W.A.	...	...	...	...	...	...	Ground ...	Very large. Fleshy, reddish brown, cup shaped. Stem central.
176	...	...	...	V.	N.S.W.	...	B	Trees ...	Soft, fleshy, shell-like, many overlapping, satiny, growing pale. Common. <i>Edible</i> .
177	W.A.	...	...	...	...	...	...	Trunks and branches	White, very delicate, tough, smooth.
178	...	...	...	V.	...	...	B	Ground ...	Fleshy, spoon shaped or fringed, disc woolly, brown to ashy buff. <i>Edible</i> .
179	...	...	T.	...	...	...	...	Roots of trees	Pale yellowish brown, funnel shaped. So phosphorescent that one was able to read books by its light, and even six days afterwards the light still served for reading.
180	S.A.	...	...	V.	...	...	...	Rotten wood	Fleshy, ochrey white, at length sulphur coloured, spotted with purple or sooty spots.
181	...	...	...	V.	N.S.W.	...	B	Trunks ...	Fleshy, greyish to tan colour, rather convex. Stem lateral, straight. <i>Edible</i> .
182	...	S.A.	...	V.	N.S.W.	...	B	Trunks, willow	Compact or spongy, shell shaped, white or grey. <i>Edible</i> .
183	...	...	...	V.	...	...	...	Rotten wood	White, sessile. Cap narrow or fan shaped, rough behind.
184	...	...	...	...	...	Q.	...	Wood ...	White. Cap half adherent, pale yellow. Stem lateral, short.
185	...	...	...	V.	...	Q.	...	Dead branches and leaves	Cap at first peziza-like, at length semisupine or half flattened out.
186	...	...	...	...	...	Q.	...	Wood ...	Dirty white, at first rather hairy, then becoming smooth. Stem obsolete.
187	...	...	...	...	...	Q.	B	Firwood twigs	Very delicate. Pale-ash colour, streaked, smooth.
188	...	...	...	V.	...	...	...	Rotten wood	Dark brown. Fan shaped, margin wavy. Stemless.
189	...	...	...	...	...	Q.	...	Rotten wood	Fleshy, sooty brown, disc darker, rather velvety. Stem thin, hollow.
190	...	...	T.	...	...	...	...	Rotten wood	Cap invested with gelatinous pellicle. Stem short, pure white, downy.
191	...	...	T.	...	...	...	...	Burnt wood	Excentric, funnel shaped, brown, powdery. Stem brown, hispid.
192	...	...	...	...	...	Q.	...	Dead leaves	Fan shaped and lobate, arising from rooting stem, whitish ochre.
Epicr. 320 (1838).— <i>Agaricus</i> .									
193	...	...	...	V.	...	...	...	Ground ...	White. Fleshy, tinged with brown. Stem rather flexuous, stuffed.
194	...	S.A.	...	V.	...	...	B	Pastures, lawns, &c.	Small, wax coloured, viscid, brittle, lustrous, translucent.
195	...	...	...	V.	...	...	B	Pastures ...	Fragile. Scarlet, shaded with orange and yellow, turning pale. <i>Edible</i> .
196	...	...	...	V.	...	Q.	B	Pastures and sandy land	Fragile, rarely red, commonly yellow, usually turning black where bruised. Sub-membranaceous, conical. Very common.
197	...	...	...	V.	...	...	...	Moist rocks	Small, dark red, funnel-shaped. Stem dilated upwards.
198	...	...	...	V.	...	...	...	Horse dung, and around it	Fleshy, sooty brown, shining, viscid. Stem straight, elongated.
199	...	...	...	V.	...	...	...	Ground ...	Yellowish orange, funnel shaped. Stem paler, thickened upwards.
200	...	...	...	V.	...	...	B	Pine woods, on sandy soil	Fleshy, covered with olive evanescent gluten and yellow beneath. Very common.
201	...	...	...	V.	...	...	...	Ground ...	Lilac, very elegant. Cap convex. Stem hollow, naked.
202	...	...	...	V.	...	Q.	B	Moist places	Fragile. Vermilion, changing colour, opaque. Stem scarlet.
203	...	...	...	V.	...	...	...	Ground ...	Small. Orange red, turning black. Stem thread-like.
204	...	...	...	...	...	Q.	...	Among grass	Fleshy, rather viscid, purple. Stem swollen in middle.
205	...	...	...	V.	...	Q.	...	Ground ...	Small. Rather fleshy, margin bright scarlet. Stem hollow, rosy white.
206	...	...	...	V.	...	Q.	B	Mossy places, wood	Somewhat brick red, rather fleshy, viscid, margin streaked.
207	...	...	...	V.	...	...	...	Among grass	Yellow, disc becoming reddish, viscid. Stem elongated, hollow.
208	...	S.A.	...	...	...	...	B	Downs and grassy places	Small. Satiny white, becoming tinted, fleshy. Stem stuffed, firm, short. Common. <i>Edible</i> .
Fl. Fr. II. 141 (1805).— <i>Agaricus</i> .									
209	...	...	...	V.	...	...	B	Woods ...	Fleshy, pallid, zoneless. Stem stuffed, then hollow. Milk mild, white. <i>Edible</i> .
210	...	...	...	...	...	Q.	B	Woods ...	White, turning black where bruised. Milk white, peppery. Common. <i>Edible</i> .
211	...	...	...	V.	...	...	B	Woods ...	Fleshy, yellowish to whitish, zoned. Gills very narrow, rather flesh coloured.
212	...	...	T.	...	...	...	...	Ground ...	Somewhat tomentose, uniber. Milk white, turning yellow, acrid.
213	...	...	...	V.	N.S.W.	...	...	Ground, in swamps	Dingy to blackish brown. Stem hollow, white at base. Milk acrid, white, unchangeable.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name	Authority for Name.	English Name.
15. RUSSULA.—Pers. Obs.,					
214	402	V. 1874	<i>R. alutacea</i> ...	Fries, <i>Epier.</i> 362 (1838) ...	Buff-gilled russule ...
215	394	IX. 249	<i>R. australiensis</i> ...	Cooke and Mass., <i>Grev.</i> XVI. 32 (1887) ...	Australian russule ...
216	399A	V. 1842	<i>R. Clusii</i> ...	Fries, <i>Hym. Eur.</i> 449 (1874) ...	Clusius' russule ...
217	399	„ 1841	<i>R. emetica</i> ...	Fries, <i>Epier.</i> 357 (1838) ...	Emetic russule ...
218	398	„ 1840	<i>R. expallens</i> ...	Gillet., <i>Tab.</i> 49 (1878) ...	Bleaching russule ...
219	400	„ 1852	<i>R. fragilis</i> ...	Fries, <i>Epier.</i> 359 (1838) ...	Fragile russule ...
220	397	„ 1818	<i>R. Linnei</i> ...	Fries, <i>Epier.</i> 356 (1838) ...	Linnaeus russule ...
221	395	„ 1805	<i>R. purpurea</i> ...	Gillet., <i>Tab.</i> 47 (1878) ...	Purple russule ...
222	396	„ 1817	<i>R. rubra</i> ...	Fries, <i>Epier.</i> 354 (1838) ...	Red russule ...
223	393	„ 1800	<i>R. sanguinea</i> ...	Fries, <i>Epier.</i> 351 (1838) ...	Blood-red russule ...
224	401	IX. 259	<i>R. subalbida</i> ...	Bres., <i>Pug., Myc. Austr.</i> 4 (1890) ...	Lurid white russule ...
16. CANTHARELLUS.					
225	404	V. 1886	<i>C. aurantiacus</i> ...	Fries, <i>S.M. I.</i> 318 (1821) ...	Orange-coloured chanterelle ...
226	405	IX. 266	<i>C. anaeolus</i> ...	Cooke and Mass., <i>Grev.</i> XVIII. 4 (1889) ...	Golden chanterelle ...
227	403	V. 1882	<i>C. cibarius</i> ...	Fries, <i>S.M. I.</i> 318 (1821) ...	Edible chanterelle ...
228	410	„ 1919	<i>C. cinereus</i> ...	Fries, <i>S.M. I.</i> 320 (1821) ...	Ash-grey chanterelle ...
229	412	„ 1934	<i>C. concinnus</i> ...	Berk., <i>Linn. Journ.</i> XVI. 38 (1878) ...	Elegant chanterelle ...
230	414	„ 1956	<i>C. foliolum</i> ...	Kalch., <i>Grev.</i> IX. 134 (1881) ...	Leaf-like chanterelle ...
231	411	„ 1920	<i>C. leucophaeus</i> ...	Noucl., <i>Mem. Lille</i> (1831) ...	White-looking chanterelle ...
232	413	„ 1952	<i>C. lobatus</i> ...	Fries, <i>S.M. I.</i> 323 (1821) ...	Lobed chanterelle ...
233	407	IX. 267	<i>C. politus</i> ...	Cooke and Mass., <i>Grev.</i> XVI. 32 (1887) ...	Polished chanterelle ...
234	409	V. 1914	<i>C. pusio</i> ...	Berk., <i>Hook., Journ.</i> VIII. 134 (1856) ...	Puny chanterelle ...
235	408	„ 1899	<i>C. strigipes</i> ...	Berk., <i>Fl. Tasm.</i> II. 248 (1860) ...	Hairy-stalked chanterelle ...
236	406	„ 1893	<i>C. viscosus</i> ...	Berk., <i>Hook., Lond. Journ.</i> IV. 49 (1845) ...	Viscid chanterelle ...
17. MARASMIUS.—Fries,					
237	444	V. 2223	<i>M. aciculaeformis</i> ...	Berk. and Curt., <i>Linn. Journ.</i> X. 297 (1869) ...	Needle-stalked marasmius ...
238	451	„ 2292	<i>M. affixus</i> ...	Berk., <i>Fl. Tasm.</i> II. 248 (1860) ...	Attached marasmius ...
239	442	„ 2218	<i>M. bicolor</i> ...	Sacc., and Cub., <i>Syll.</i> V. 555 (1887) ...	Two-coloured marasmius ...
240	438	„ 2187	<i>M. calobates</i> ...	Kalch., <i>Grev.</i> IV. 71 (1876) ...	Stilted marasmius ...
241	425	„ 2071	<i>M. calopus</i> ...	Fries, <i>Epier.</i> 379 (1838) ...	Beautiful-stemmed marasmius ...
242	431	„ 2122	<i>M. caudicinalis</i> ...	Fries, <i>Epier.</i> 383 (1838) ...	Craggy marasmius ...
243	418	„ 2046	<i>M. confertus</i> ...	Berk. and Br., <i>Linn. Journ.</i> XIV. 34 (1875) ...	Crowded marasmius ...
244	447	„ 2259	<i>M. de Toniatus</i> ...	Sacc. and Cub., <i>Syll.</i> V. 563 (1887) ...	De Toni's marasmius ...
245	450	„ 2286	<i>M. emergens</i> ...	Berk. in Cooke's <i>Handb., Aust. Fungi</i> , 88 (1892) ...	Emerging marasmius ...
246	445	„ 2239	<i>M. epiphyllus</i> ...	Fries, <i>Epier.</i> 386 (1838) ...	Leaf marasmius ...
247	441	„ 2203	<i>M. equicrinis</i> ...	F. v. M., <i>Grev.</i> VIII. 153 (1880) ...	Horse-hair marasmius ...
248	420	„ 2051	<i>M. erythropus</i> ...	Fries, <i>Epier.</i> 378 (1838) ...	Red-stalked marasmius ...
249	440	„ 2200	<i>M. Eucalypti</i> ...	Berk., <i>Fl. Tasm.</i> II. 249 (1860) ...	Eucalyptus marasmius ...
250	452	„ 2291	<i>M. Exocarpi</i> ...	Berk., <i>Linn. Journ.</i> XVIII. 384 (1881) ...	Native cherry marasmius ...
251	435	„ 2144	<i>M. ferrugineus</i> ...	Berk., <i>Hook. Lond. Journ.</i> II. 630 (1843) ...	Rust-red marasmius ...
252	423	„ 2063	<i>M. floriceps</i> ...	Berk. and Curt., <i>Linn. Journ.</i> X. 298 (1869) ...	Flower-capped marasmius ...
253	427	„ 2095	<i>M. foetidus</i> ...	Fries, <i>Epier.</i> 380 (1838) ...	Fetid marasmius ...
254	434	„ 2143	<i>M. haematoccephalus</i> ...	Mont., <i>Syll.</i> 109 (1856) ...	Blood-red capped marasmius ...
255	416	„ 2013	<i>M. hepaticus</i> ...	Berk., <i>Hook., Lond. Journ.</i> V. 1 (1846) ...	Liver-coloured marasmius ...
256	421	„ 2057	<i>M. impudicus</i> ...	Fries, <i>Epier.</i> 377 (1838) ...	Impure marasmius ...
257	422	IX. 278	<i>M. lanaripes</i> ...	Cooke and Mass., <i>Grev.</i> XVIII. 4 (1889) ...	Woolly-stalked marasmius ...
258	439	V. 2199	<i>M. liguoides</i> ...	Berk., <i>Linn. Journ.</i> XVIII. 384 (1881) ...	Smoky marasmius ...
259	436	„ 2147	<i>M. meloniformis</i> ...	Berk., <i>Fl. Tasm.</i> II. 249 (1860) ...	Melon-shaped marasmius ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
I. 100 (1796).—Agaricus.									
214	...	...	T.	...	...	...	B	Woods ...	Mild. Fleishy, dark lake, red, or purplish, margin flesh white. Gills bright buff. <i>Edible</i> .
215	...	...	...	V.	...	Q.	...	Ground ...	Acrid. Fleishy, red. Stem stuffed, then hollow, straw coloured.
216	...	...	...	V.	...	...	B	Woods ...	Blood red, flesh white to yellowish.
217	...	...	T.	V.	N.S.W.	...	B	Woods ...	Acrid. Fleishy, polished, shining, margin flesh white. Rose, varied with lilac or yellow.
218	...	...	...	V.	...	...	B	Under trees	Fleishy, viscid, bright, purple, becoming pale. Stem cylindrical, firm.
219	...	S.A.	...	V.	...	Q.	B	Woods ...	Very acrid, small. Fleishy, rose red, becoming pale, polished, slightly viscid.
220	...	...	...	...	...	Q.	B	Woods ...	Mild. Fleishy, polished, dry, white. Stem spongy, stout, red.
221	...	...	...	V.	...	...	B	Under trees	Rather mild. Fleishy, viscid, dark purple. Stem white at top, rosy middle.
222	...	...	...	V.	N.S.W.	Q.	B	Grassy places	Acrid. Fleishy, polished, dry, deep dark vermilion. Stem white or red.
223	...	...	...	...	...	Q.	B	Woods ...	Acrid. Fleishy, generally blood red, glistening. Stem white or red.
224	...	...	...	...	...	Q.	...	Ground ...	Rather fleshy, margin lurid white. Stem stuffed, then hollow.
Pers. Tent. Disp. 26 (1797).—Agaricus.									
225	...	...	...	...	...	Q.	B	Fir woods and heaths	Nearly orange colour or orange yellow. Fleishy, rather tomentose.
226	...	...	...	...	...	Q.	...	Ground ...	Golden. Thin, delicately downy. Stem slender, faintly streaked.
227	...	...	...	V.	N.S.W.	Q.	B	Woods ...	Apricot yellow and apricot scented. Fleishy. <i>Edible</i> .
228	...	...	T.	...	...	...	B	Woods ...	Sub-membranaceous, dingy black, hairy to scaly. Stem hollow.
229	...	...	...	...	N.S.W.	...	...	Ground ...	Small. Gills very narrow and forked. Stem thickened upwards.
230	...	...	...	...	...	Q.	...	Twigs, leaves, &c....	Membranaceous, whitish, pale ochre when dry or reddish brown. Colour and veins slightly prominent as in a dry leaf.
231	...	...	...	V.	...	...	B	Ground ...	Sub-membranaceous, umber. Stem stuffed, thin, of same colour.
232	...	...	...	V.	...	...	B	Mosses, in swamps	Membranaceous, gelatinous, sessile, dirty reddish brown.
233	...	S.A.	...	V.	...	...	...	Fern gully	Rather fleshy, viscid very shining, chestnut colour. Stem stuffed.
234	...	...	T.	V.	...	...	...	Ground ...	Becoming whitish. Funnel shaped, powdery, woolly. Stem brown when dry.
235	...	...	T.	...	...	...	...	Among ferns	Liver colour. Stem arising from tawny strigose hairs. Tapering upwards.
236	W.A.	...	...	...	...	...	...	Ground, among twigs	Beautifully yellow. Viscid, funnel shaped, somewhat wavy.
Epier. 372 (1838).—Agaricus.									
237	...	...	...	...	...	Q.	...	Rotten wood	Gregarious. Tawny. Stem hair-like, rigid, shining, brownish.
238	...	...	T.	...	...	Q.	...	Rotten wood	Whitish, mealy, tomentose, cup shaped, reflexed and attached by side.
239	...	...	...	...	...	Q.	...	Trunks ...	Small. White. Stem short, becoming red below, thread-like.
240	...	...	...	V.	...	Q.	...	Putrid leaves of <i>Bougainvillea</i>	Membranaceous, rust coloured, becoming brown. Stem horny, turning black.
241	...	...	...	V	...	Q.	B	Twigs, grass, roots, &c.	Inodorous. Rather fleshy. Stem shining, bay to red, hollow, not rooting.
242	...	...	...	V.	...	Q.	...	Ground, among leaves	Membranaceous, rust coloured, yellow, then ochre. Stem hollow, bay.
243	...	...	...	...	N.S.W.	Q.	...	Among dead vegetables	Sub-membranaceous, tawny. Stem of same colour, hollow.
244	...	...	...	...	N.S.W.	...	...	Branches ...	Cap scarcely the size of a mustard seed, brownish. Stem hair-like.
245	...	...	T.	...	...	...	...	Wood ...	Very minute, white, bursting through. Stem shortened or elongated.
246	...	...	...	V.	...	...	B	Fallen leaves, twigs, &c.	Minute. Membranaceous, creamy. Stem rather horny, finely velvety.
247	...	...	...	V.	N.S.W.	Q.	...	Branches ...	Whitish to tawny, small, membranaceous. Stem hair-like, rigid, black, shining, arising from black horsehair-like mycelium.
248	...	...	...	V.	...	Q.	B	Among leaves, near stumps	Inodorous. Rather fleshy. Stem dark red, hollow, streaked.
249	...	...	T.	...	...	...	...	Fruit and branches of <i>Eucalyptus</i>	Conical, brownish, silky. Stem hair-like, compressed, shining.
250	...	...	...	V.	...	Q.	...	Trunks of <i>Exocarpos latifolia</i> (Native Cherry)	White, wholly resupinate.
251	...	...	...	V.	...	...	...	Dead leaves, branches, &c.	Membranaceous, ferruginous yellow. Stem slender, twisted.
252	...	...	...	V.	...	...	...	Rotten wood	Conical, bright red brown. Stem twisted, hollow, shining.
253	...	...	...	...	...	Q.	B	Decayed twigs	Fætid. Sub-membranaceous, pellucid, tawny chestnut or somewhat red.
254	...	...	...	V.	...	Q.	...	Dead leaves	Membranaceous, blood red. Stem horny, hair-like, umber.
255	...	...	T.	...	...	...	...	Among ferns	Rather fleshy, liver coloured. Stem of stringy fibres, thickened below.
256	...	...	...	V.	...	...	B	About trunks	Fætid. Rather fleshy, chestnut red. Stem hollow, turning purple.
257	...	...	...	...	...	Q.	...	Rotten wood	Fleshy, lead colour or dirty dark blue. Stem red, densely velvety.
258	...	...	...	V.	...	...	...	Leaves of <i>Eucalyptus</i>	Furrowed. Stem black, shining, grooved.
259	...	...	T.	...	...	...	...	Leaves and branches of <i>Eucalyptus</i>	Minute. Bay brown, mealy. Stem thread-like, shining.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
17. MARASMIUS.—Fries,					
260	419	V. 2049	<i>M. Muelleri</i> ...	Berk., Linn. Journ. XVIII. 383 (1881) ...	Mueller's marasmius ...
261	429	„ 2106	<i>M. opacus</i> ...	Berk. and Curt., Hook. Journ. I. 99 (1849) ...	Opaque marasmius ...
262	417	„ 2014	<i>M. pilopus</i> ...	Kalch., Grev. VIII. 153 (1880) ...	Downy-stalked marasmius ...
263	446	„ 2257	<i>M. primulinus</i> ...	Berk., Linn. Journ. XVI. 38 (1878) ...	Pale-yellow marasmius ...
264	433	„ 2133	<i>M. putredinis</i> ...	Berk. and Curt., Linn., Journ. X. 295 (1869) ...	Putrid marasmius ...
265	428	„ 2103	<i>M. ramealis</i> ...	Fries, Epicr. 381 (1838) ...	Twig marasmius ...
266	449	„ 2261	<i>M. rhyticeps</i> ...	Kalch., Grev. IV. 71 (1876) ...	Wrinkle-capped marasmius ...
267	437	„ 2150	<i>M. rotula</i> ..	Fries, Epicr. 385 (1838) ...	Collared marasmius ...
268	443	„ 2219	<i>M. rufo-pallidus</i> ...	Kalch., Grev. IV. 71 (1876) ...	Pale-red marasmius ...
269	424	„ 2070	<i>M. scorodonius</i> ...	Fries, Epicr. 379 (1838) ...	Shallot marasmius ...
270	426	„ 2084	<i>M. stylobates</i> ...	Berk. and Curt., Linn. Journ. X. 296 (1869) ...	Pillar-shaped marasmius ...
271	...	„	<i>M. subroseus</i> ...	Cooke and Mass., Grev. XXI. 37 (1892) ...	Somewhat rosy marasmius ..
272	448	„ 2260	<i>M. subsupinus</i> ...	Berk., Fl. Tasm. II. 249 (1860) ...	Subsupine marasmius ...
273	415	„ 1976	<i>M. urens</i> ...	Fries, Epicr. 373 (1838) ...	Acrid marasmius ...
18. LENTINUS.—Fries,					
274	461	V. 2332	<i>L. blepharodes</i> ...	Berk. and Curt., Linn. Journ. X. 301 (1869) ...	Eyelashed lentine ...
275	463	„ 2348	<i>L. calvescens</i> ...	Berk., Hook., Journ. VIII. 141 (1856) ...	Bald lentine ...
276A	487	„ 2483	<i>L. castoreus</i> , var. <i>hirneoloides</i>	Berk. and Br., Linn. Journ. X. 302 (1869) ...	Hirneola-like lentine ...
277	479	„ 2456	<i>L. catervarius</i> ...	Berk. and Br., Linn. Trans. II. 55 (1883) ...	Crowded lentine ...
278	476	„ 2415	<i>L. cochleatus</i> ...	Fries, Hym. Eur. 484 (1874) ...	Cochleate lentine ...
279	470	„ 2376	<i>L. cretaceus</i> ...	Berk. and Br., Linn. Journ. XIV. 42 (1875) ...	Chalky lentine ...
280	475	„ 2398	<i>L. cyathus</i> ...	Berk. and Br., Linn. Trans. I. 399 (1879) ...	Goblet lentine ...
281	459	„ 2325	<i>L. dealbatus</i> ...	Fries, Pl. Preiss. II. 133 (1846) ...	White-washed lentine ...
282	465	„ 2350	<i>L. Dunalii</i> ...	Fries, Epicr. 390 (1838) ...	Dunal's lentine ...
283	467	„ 2358	<i>L. exasperatus</i> ...	Berk. and Br., Linn. Trans. II. 55 (1883) ...	Rough lentine ...
284	484	„ 2472	<i>L. exilis</i> ..	Fries, Epicr. 393 (1838) ...	Thin lentine ...
285	458	„ 2317	<i>L. fasciatus</i> ...	Berk., Hook., Journ. II. 146 (1840) ...	Clustered lentine ...
286	483	„ 2471	<i>L. fulvaster</i> ...	Berk. and Cooke, Linn. Journ. XV. 373 (1877) ...	Yellowish lentine ...
287	455	„ 2312	<i>L. fulvus</i> ...	Berk., Ann., Nat., Hist. X. 369 (1843) ...	Tawny lentine ...
288	456	„ 2315	<i>L. fusco-purpureus</i> ...	Kalch., Grev. VIII. 153 (1880) ...	Purple-brown lentine ...
289	486	IX. 322	<i>L. fusipes</i> ...	Cooke and Mass., Grev. XVI. 1 (1887) ...	Fusiform lentine ...
290	482	„ 301	<i>L. gracilentus</i> ...	Cooke and Mass., Grev. XVI. 73 (1888) ...	Slender lentine ...
291	480	V. 2458	<i>L. Guilfoylei</i> ...	Berk., Linn. Journ. XVIII. 384 (1881) ...	Guilfoyle's lentine ...
292	490	„ 2490	<i>L. hepatotrichus</i> ...	Berk., Fl. Tasm. II. 249 (1860) ...	Liver-coloured lentine ...
293	473	„ 2514	<i>L. holopogonius</i> ...	Berk., Grev. X. 63 (1881) ...	Bearded lentine ...
294	485	„ 2483	<i>L. hyracinus</i> ...	Kalch., Grev. VIII. 153 (1880) ...	Hyrax-coloured lentine ...
295	471	„ 2394	<i>L. Kurzianus</i> ...	Curr., Linn. Trans. I. 120 (1876) ...	Kurz's lentine ...
296	478	„ 2449	<i>L. kæviceps</i> ...	Kalch., Grev. VIII. 153 (1880) ...	Even-headed lentine ...
297	489	IX. 317	<i>L. lasiophyllus</i> ..	Cooke and Mass., Grev. XVI. 1 (1887) ...	Hairy-gilled lentine ...
298	481	V. 2459	<i>L. lateritius</i> ...	Berk., Linn. Journ. XVIII. 384 (1881) ...	Brick-red lentine ...
299	453	„ 2308	<i>L. Lecomtei</i> ...	Fries, Epicr. 368 (1838) ...	Lecomte's lentine ...
300	466	„ 2351	<i>L. lepidens</i> ...	Fries, Epicr. 390 (1838) ...	Scaly lentine ...
301	469	„ 2371	<i>L. manipularis</i> ...	Berk. and Br., Linn. Journ. XIV. 43 (1875) ...	Tufted lentine ...
302	493	„ 2499	<i>L. pelliculosus</i> ...	Fries, Epicr. 395 (1838) ...	Thin-skinned lentine ...
303	477	„ 2439	<i>L. pergamenus</i> ...	Lev., Champ., Mus. 117 (1846) ...	Parchment lentine ...
304	494	„ 2506	<i>L. pulvinulus</i> ...	Berk., Fl. Tasm. II. 250 (1860) ...	Pulvinate lentine ...
305	491	„ 2495	<i>L. punctaticeps</i> ...	Berk. and Br., Linn. Trans. II. 55 (1883) ...	Punctate-headed lentine ...
306	474	„ 2395	<i>L. radiceatus</i> ...	Cooke and Mass. Grev. XIV., 118 (1886) ...	Rooting lentine ...
307	472	„ 2512	<i>L. Schomburgkii</i> ...	Berk., Linn. Trans. XX. 111 (1851) ...	Schomburgk's lentine ...
308	462	„ 2333	<i>L. siparius</i> ...	Berk. and Curt., Linn. Journ. X. 301 (1869) ...	Curtain lentine ...
309	454	„ 2311	<i>L. strigosus</i> ...	Fries, Epicr. 388 (1838) ...	Strigose lentine ...
310	492	„ 2497	<i>L. subdulcis</i> ...	Berk., Hook., Journ. III. 46 (1851) ...	Sweet-scented lentine ...
311	468	„ 2361	<i>L. subnudus</i> ...	Berk., Hook., Lond. Journ. VI. 492 (1847) ...	Somewhat-naked lentine ...
312	460	„ 2330	<i>L. tener</i> ...	Klotsch in Fries, Epicr. 389 (1838) ...	Slender lentine ...
313	464	„ 2349	<i>L. tigrinus</i> ...	Fries, Epicr. 389 (1838) ...	Tiger-tuft lentine ...
314	457	„ 2316	<i>L. villosus</i> ...	Klotsch in Linn. 479 (1833) ...	Villous lentine ...
315	488	„ 2486	<i>L. vulpinus</i> ...	Fries, Mon. Hym. II. 238 (1857) ...	Fox-coloured lentine ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Epicr. 372 (1838).— <i>Agaricus</i> —continued.									
260	...	...	...	V.	...	Q.	...	Ground ...	Tawny, delicately tomentose. Stem thin, dilated at base.
261	...	...	...	...	N.S.W.	Q.	...	Leaves and twigs ...	Slender. Opaque, powdery, whitish. Stem mealy towards base.
262	...	...	...	...	N.S.W.	Q.	...	Wood ...	Leathery to membranaceous, yellowish tan. Stem with powdery ochrey down.
263	...	...	...	...	N.S.W.	...	...	...	Pale yellow, powdery. Stem short, slender, mealy.
264	...	...	...	V.	...	Q.	...	<i>Eucalyptus</i> ...	Thin, reddish yellow or grey. Stem of same colour, solid.
265	...	...	...	...	...	Q.	B	Dry dead branches	Inodorous. Somewhat fleshy, opaque, white, disc somewhat red. Very common.
266	...	...	...	...	...	Q.	...	Passion-flower twigs	Membranaceous, tawny to reddish brown. Stem thread-like, velvety.
267	...	...	...	V.	...	Q.	B	Fallen twigs, &c. ...	Minute. Membranaceous, whitish. Stem horny, shining, blackish, with collar from separating gills.
268	...	...	...	...	...	Q.	...	Ground, about trunks	Membranaceous, pale red. Stem thread-like, fixed at base by white mycelium.
269	...	...	...	V.	...	...	B	Dry ground	Small. Strong scented, oniony odour. Somewhat fleshy, red becoming white, or red buff. <i>Edible</i> .
270	...	...	...	...	...	Q.	...	Wood ...	White. Thin, smooth. Stem arising from circular base, smooth, hollow.
271	...	...	...	V.	...	...	...	Rotten wood	Membranaceous, pale-tan colour, tinged with pink. Stem horny, hollow.
272	...	...	T.	...	...	Q.	...	Rotten wood	Small. Mealy, adhering behind. Stem short, mealy.
273	...	...	...	...	...	Q.	B	Wood ...	Acrid, odourless. Fleshy to leathery. Usually ochrey tan. Stem fibrous.
Pl. Homon. 77 (1825).— <i>Agaricus</i> .									
274	...	...	...	...	...	Q.	...	Dead branches	Brown, hispid, margin ciliate. Stem velvety.
275	...	...	...	...	...	Q.	...	Rotten trunks	Pale, at first woolly then bald, margin lobed. Stem short, nearly naked.
276A	...	...	...	...	N.S.W.	...	...	Rotten logs, in woods	Pale, tawny. Gills paler. cap thin.
277	...	...	...	...	...	Q.	...	Trunks	Golden yellow. Cap convex, then flattened. Stem cylindrical.
278	W.A.	...	V.	...	...	Q.	B.	Trunks and ground	Tough, flaccid. Fleshy, reddish brown, somewhat lobed or contorted.
279	...	...	...	...	...	Q.	...	Ground	White, orbicular. Stem thin, at length furrowed.
280	...	...	...	...	...	Q.	...	Dense scrubs	Ochrey, with velvety brown lines radiating from centre, funnel shaped.
281	W.A.	...	...	...	...	...	...	Rotten trunks	Becoming white, leathery, woolly to hairy, zoneless. Stem short.
282	...	...	...	...	...	Q.	B.	Trunks — Willows and Poplars	Small, tufted. Fleshy to leathery, yellow white with brown scales.
283	...	...	...	...	...	Q.	...	Trunks	Rough, with rigid warts, ferruginous, powdery. Stem thickened downwards.
284	...	S.A.	V.	...	...	Q.	...	Rotten wood	Papery, pale tawny.
285	W.A.	...	T.	...	N.S.W.	Q.	...	Trunks	Thin, leathery, wine-glass shape, pale ochre. Stem velvety, tawny.
286	...	...	V.	...	...	...	...	Deadwood	Orbicular, white, becoming tawny when dry. Stem slender, smooth.
287	...	...	V.	...	N.S.W.	Q.	...	Rotten wood	Deeply funnel shaped, bay brown, somewhat zoned. Stem rough or downy.
288	...	...	...	...	N.S.W.	Q.	...	Wood	Leathery, funnel shaped, purple brown, hairy. Stem tall, bristly.
289	...	...	...	...	...	Q.	...	Rotten wood	Fleshy, white, downy. Stem lateral, spindle shaped, rooting.
290	...	...	V.	...	...	Q.	...	Rotten wood	Rather membranaceous, ochrey, funnel shaped. Stem slender, brown.
291	...	...	...	...	N.S.W.	Q.	...	Rotten wood	Umbel, smooth. Stem curved, of same colour.
292	...	...	T.	...	...	...	...	Bark of <i>Eucalyptus</i>	Hoof-like, liver coloured, becoming smooth in front, hispid behind.
293	...	...	...	...	...	Q.	...	Stumps	Dirty white, funnel shaped, densely hispid. Stem hispid.
294	...	...	...	...	N.S.W.	Q.	...	Wood	Orbicular or semi-orbicular, full red brown, downy, wrinkled behind.
295	...	...	...	...	...	Q.	...	Ground	Funnel shaped, mealy, tawny. Stem short, rusty brown.
296	...	...	...	...	...	...	...	...	Fleshy to leathery, yellowish white. Stem solid, scaly at base.
297	...	...	V.	...	N.S.W.	...	...	Stumps	Thin, ochrey, shining, lobed at margin. Stems discoid, downy.
298	...	...	...	...	...	...	...	Wood	Brick red, and quite smooth. Stem of same colour, rigid.
299	...	...	...	...	N.S.W.	Q.	...	Rotten wood	Leathery, funnel shaped, fawn colour. Stem hairy, of same colour.
300	...	...	...	...	...	Q.	B.	Stumps of firs, &c.	Fleshy, compact, pale ochrey, broken up into darker spot-like scales.
301	...	...	...	...	...	Q.	...	Dead wood	Tufted, orbicular, white, clad with mealy scales. Stem curved.
302	...	...	...	...	N.S.W.	...	...	Rotten trunks	Sessile, tough, very thin, tawny fawn colour, kidney shaped.
303	...	...	...	...	...	Q.	...	Ground	White, leathery to membranaceous, funnel shaped. Stem solid.
304	...	...	T.	...	...	...	...	Rotten wood	Cushion shaped, pale, margin furrowed, ochrey.
305	...	...	...	...	...	Q.	...	Trunks	Punctately hispid, presenting sponge-like appearance. Stem yellow.
306	...	...	...	...	...	Q.	...	Ground	Fleshy, funnel shaped, pale ochrey, shortly velvety. Stem velvety.
307	...	...	...	...	...	Q.	...	Wood	Leathery when dry, fawn coloured, broadly funnel shaped.
308	...	...	...	...	...	Q.	...	Rotten wood	Orange, brown when dry. Woolly with erect rigid hairs intermixed.
309	...	...	...	...	N.S.W.	Q.	...	Trunks	Reddish-fawn colour. Roughly hairy. Stem excentric and hairy.
310	...	...	...	...	...	Q.	...	Deadwood	White, sweet scented. Cap lobate, smooth. Stem obsolete or spurious.
311	...	S.A.	V.	...	...	Q.	...	Fallen trees	Rather funnel shaped, at first clad with mealy scales. Stem slender.
312	...	...	...	...	...	Q.	...	Rotten wood	Membranaceous, bay brown. Stem very slender, brownish.
313	...	...	...	...	...	Q.	B.	Trunks	Fleshy to leathery, whitish to yellow white, with tawny scales. <i>Edible</i> .
314	...	...	...	...	...	Q.	...	Rotten wood	Brown, leathery, with fibrous bristles. Stem solid, tawny.
315	...	...	...	...	N.S.W.	Q.	B.	Stumps	Sessile, imbricated. Fleshy, conchate, tan coloured, corrugated or woolly.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
19. PANUS.—Fries,					
316	511	...	<i>P. angustatus</i> ...	Berk. in Cooke's Handb. Aust. Fungi 98 (1892) ...	Narrow panus ...
317	503	V. 2567	<i>P. arenicola</i> ...	Berk., Linn., Journ. XVIII. 384 (1881) ...	Sand-growing panus ...
318	501	" 2552	<i>P. carbonarius</i> ...	Cooke and Mass., Grev. XV. 94 (1887) ...	Charcoal panus ...
319	505	" 2569	<i>P. cinnabarinus</i> ..	Fries, Pl. Preiss. II. 133 (1846) ...	Vermilion panus ...
320	495	" 2519	<i>P. conchatus</i> ...	Fries, Epicr. 398 (1838) ...	Conch panus ...
321	500	" 2547	<i>P. coriaceus</i> ...	Berk., Linn. Journ. XIII. 160 (1873) ...	Coriaceous panus ...
322	507	" 2573	<i>P. eugrammus</i> ...	Fries, Nov. Symb. 40 (1851) ...	Well-lined panus ...
323	498	" 2541	<i>P. incandescens</i> ...	Berk. and Br., Linn. Trans. II. 55 (1883) ...	Incandescent panus ...
324	506	IX. 329	<i>P. lateritius</i> ...	Sacc. Hedw. 125 (1889) ...	Brick-red panus ...
325	508	" 328	<i>P. olivaceo-flavidus</i> ...	Cooke and Mass., Grev. XVI. 1 (1887) ...	Yellowish-olive panus ...
326	510	V. 2588	<i>P. patellaris</i> ...	Fries, Epicr. 400 (1838) ...	Cup-shaped panus ...
327	497	" 2540	<i>P. rivulosus</i> ...	Berk., Linn. Journ. XVIII. 384 (1881) ...	Cracked panus ...
328	509	" 2578	<i>P. saccharinus</i> ...	Berk., Fl. Tasm. II. 250 (1860) ...	Saccharine panus ...
329	502	" 2557	<i>P. stypticus</i> ...	Fries, Epicr. 399 (1838) ...	Styptic panus ...
330	499	" 2542	<i>P. suborbicularis</i> ...	Berk. and Br., Linn. Trans. II. 56 (1883) ...	Sub-orbicular panus ...
331	496	" 2521	<i>P. torulosus</i> ...	Fries, Epicr. 397 (1838) ...	Twisted panus ...
332	504	" 2568	<i>P. viscidulus</i> ...	Berk. and Br., Linn. Trans. II. 56 (1883) ...	Gelatinous panus ...
20. XEROTUS.—Fries,					
333	516	V. 2609	<i>X. albidus</i> ...	Berk. and Br., Linn. Trans. II. 56 (1883) ...	Whitish xerote ...
334	520	" 2617	<i>X. Archeri</i> ...	Berk., Fl. Tasm. II. 250 (1860) ...	Archer's xerote ...
335	517	" 2611	<i>X. Berterii</i> ...	Mont. Chil. VII. 353 (1852) ...	Bertero's xerote ...
336	522	...	<i>X. Drummondii</i> ...	Berk. in Cooke's Handb. Aust. Fungi 100 (1892) ...	Drummond's xerote ...
337	521	...	<i>X. fulvus</i> ...	Berk. and Br. in Cooke's Handb. Aust. Fungi 100 (1892) ...	Tawny xerote ...
338	523	V. 2596	<i>X. griseus</i> ...	Berk., Hook., Lond. Journ. VI. 497 (1847) ...	Grey xerote ...
339	518	" 2613	<i>X. lateritius</i> ...	Berk. and Curt., Linn. Journ. X. 303 (1869) ...	Brick-red xerote ...
340	514	" 2606	<i>X. papuasius</i> ...	Kalch., Grev. VIII. 154 (1880) ...	Papuan xerote ...
341	519	" 2616	<i>X. papyraceus</i> ...	Berk., Fl. Tasm. II. 250 (1860) ...	Papery xerote ...
342	513	" 2601	<i>X. proximus</i> ...	Berk. and Br., Linn. Trans. II. 56 (1883) ...	Approximate xerote ...
343	515	" 2607	<i>X. rawakensis</i> ...	Pers. in Fries., Epicr. 401 (1838) ...	Rawak xerote ...
344	512	" 2599	<i>X. tener</i> ...	Berk. and Br., Linn. Journ. XIV. 45 (1875) ...	Tender xerote ...
21. TROGIA.—Fries,					
345	524	V. 2627	<i>T. crispa</i> ...	Fries, Mon. Hym. II. 244 (1857) ...	Crisped trogia ...
22. LENZITES.—Fries,					
346	529	V. 2638	<i>L. abietina</i> ...	Fries, Epicr. 407 (1838) ...	Larch lenzites ...
347	532	" 2654	<i>L. acuta</i> ...	Berk., Hook., Lond. Journ. I. 146 (1842) ...	Acutely-margined lenzites ...
348	534	" 2657	<i>L. applanata</i> ...	Fries, Epicr. 404 (1838) ...	Depressed lenzites ...
349	535	" 2658	<i>L. aspera</i> ...	Fries, Epicr. 405 (1838) ...	Rough lenzites ...
350	536	" 2664	<i>L. Beckleri</i> ...	Berk., Linn. Journ. XIII. 161 (1873) ...	Beckler's lenzites ...
351	530	" 2651	<i>L. Berkeleyi</i> ...	Lev., Ann. Sci. Nat. V. 122 (1846) ...	Berkeley's lenzites ...
352	526	" 2630	<i>L. betulina</i> ...	Fries, Epicr. 405 (1838) ...	Birch lenzites ...
352A	"	" 2651	<i>L. betulina</i> , var. <i>velutina</i> ...	Berk., Ann. Nat. Hist. III. 381 (1839) ...	Velvety lenzites ...
353	"	"	<i>L. bifasciatus</i> ...	Cooke and Mass., Grev. XXI. 37 (1892) ...	Bifasciate lenzites ...
354	533	V. 2656	<i>L. deplanata</i> ...	Fries, Epicr. I. 404 (1838) ...	Levelled lenzites ...
355	540	" 2685	<i>L. faventina</i> ...	Cald., Erb. Critt. Ital. No. 89 (1878, &c.) ...	Honeycombed lenzites ...
356	527	" 2631	<i>L. flaccida</i> ...	Fries, Epicr. 406 (1838) ...	Flaccid lenzites ...
357	539	" 2682	<i>L. Guilfoylei</i> ...	Berk. Grev. X. 64 (1881) ...	Guilfoyle's lenzites ...
358	538	" 2670	<i>L. nivea</i> ...	Cooke, Grev. XV. 94 (1887) ...	Snow-white lenzites ...
359	541	" 2687	<i>L. Palisotii</i> ...	Fries, Epicr. 404 (1838) ...	Palisot's lenzites ...
360	542	" 2688	<i>L. repanda</i> ...	Fries, Epicr. 404 (1838) ...	Repand lenzites ...
361	528	" 2636	<i>L. sæpiaria</i> ...	Fries, Epicr. 407 (1838) ...	Chocolate lenzites ...
362	531	" 2653	<i>L. striata</i> ...	Fries, Epicr. 406 (1838) ...	Striated lenzites ...
363	537	" 2665	<i>L. torrida</i> ...	Kalch., Grev. VIII. 154 (1880) ...	Torrid lenzites ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Epier. 396 (1838).— <i>Agaricus, Lentinus</i> .									
316	...	...	...	...	...	Q.	...	Logs ...	Spoon shaped, tawny, nearly sessile, with a few scattered hairs.
317	...	...	...	V.	...	...	...	Sandy soil ...	Brown, spoon shaped. Stem and cap covered with particles of sand.
318	...	S.A.	...	...	...	...	...	Among ferns where burnt	Fleshy, umber, fan or funnel shaped. Stem short, pale.
319	W.A.	...	...	...	...	Q.	...	Base of trunks ...	Leathery, vermilion, sessile, kidney shaped.
320	...	...	...	...	N.S.W.	...	B.	Trunks — Poplar, Beech, Birch	Largish. Fleshy, cinnamon, becoming pale, conchate. <i>Edible</i> .
321	...	...	...	V.	...	...	...	Bark ...	Very beautiful. Leathery, brown behind or black when young, sessile.
322	...	...	...	...	...	Q.	...	Bark ...	Sessile, imbricate. Leathery to membranaceous, pale, kidney shaped.
323	...	S.A.	...	V.	N.S.W.	Q.	...	Buried wood, but apparently on soil	Sometimes funnel shaped, smooth. Very luminous at night.
324	...	S.A.	...	...	...	...	...	Rotten wood of <i>Eucalyptus</i>	Membranaceous, tau coloured, sprinkled with brick-red point-like threads.
325	...	...	...	V.	N.S.W.	...	...	Burnt wood ...	Sooty brown, densely velvety, with yellowish-olive down, sessile.
326	...	...	...	...	...	Q.	B.	Branches — Beech, Cherry, &c.	Leathery, mealy to downy, flat cup shaped.
327	...	...	...	V.	...	...	...	Trunks ...	Ochrey, striately cracked. Stem excentric, similarly cracked.
328	...	...	T.	...	...	...	...	Rotten wood ...	Rather fleshy. Stem short, mealy or obsolete. Edge of gills as if dusted with sugar.
329	...	...	...	V.	...	...	B.	Stumps ...	Small. Leathery, kidney shaped, cinnamon turning pale. Astringent taste.
330	...	...	...	...	...	Q.	...	Old trunks ...	Sub-orbicular, white, delicately downy. Stem obsolete.
331	...	...	...	V.	...	Q.	B.	Trunks ...	Fleshy to leathery, flesh coloured to ochrey pink. <i>Edible</i> .
332	...	...	...	V.	N.S.W.	Q.	...	Rotten trunks, decaying bark ...	Upper layer gelatinous, rather viscid, dull slate coloured. Stem short, lateral, arising from spongy base.

Elench. I. 48 (1828).

333	...	...	...	...	...	Q.	...	Wood ...	Whitish, kidney shaped. Stem lateral, smooth or slightly velvety.
334	...	...	T.	V.	...	...	...	Sticks ...	Kidney shaped, reddish brown. Stem very short, lateral, powdery.
335	...	...	...	...	...	Q.	...	Fallen branches ...	Gregarious. Leathery to membranaceous, rust coloured, kidney shaped.
336	...	...	...	V.	...	...	...	Twigs ...	Gregarious. Kidney shaped, lobed or crispate, rust coloured.
337	...	...	...	...	...	Q.	...	Wood ...	Tawny ochrey. Membranaceous, kidney shaped. Stem lateral, thin.
338	...	...	...	V.	...	...	...	Old wood...	Funnel shaped, splitting, grey. Stem compressed, wedge shaped.
339	...	...	...	...	...	Q.	...	Dead bark ...	Sub-orbicular, brick red, furrowed.
340	...	...	...	...	N.S.W.	...	...	Bark ...	Ochrey tan colour. Membranaceous to leathery, radiately furrowed.
341	...	...	T.	V.	...	...	...	Rotten wood ...	Papery, pale. Stem very short or obsolete.
342	...	...	...	V.	...	Q.	...	Branches ...	White, then yellowish brown Sub-orbicular, delicately powdery.
343	...	...	...	...	...	Q.	...	Wood ...	Smooth, cinnamon. Leathery to membranaceous. Stem solid, short.
344	...	...	...	...	...	Q.	...	Dead wood ...	Kidney shaped, membranaceous, umber. Stem very short.

Epier. 402 (1838).—*Merulius, Cantharellus*.

345	...	...	...	V.	...	Q.	B.	Twigs—Birch, Beech, &c.	Tough, cup shaped, reddish yellow. Gills crisp, plaited, forked.
-----	-----	-----	-----	----	-----	----	----	-------------------------	--

Epier. 403 (1838).—*Agaricus, Dædalea*.

346	...	S.A.	...	...	...	Q.	B.	Wood ...	Leathery, clothed with umber down, at length smooth and whitish.
347	...	...	...	...	...	Q.	...	Wood ...	Kidney shaped, leathery, greyish umber. Stem distinct and disc shaped.
348	...	...	...	...	...	Q.	...	Wood ...	Kidney shaped, corky, whitish, zoneless, downy.
349	...	...	...	...	N.S.W.	Q.	...	Dead wood ...	Thick, spongy to corky, rough, pale, concentrically furrowed.
350	...	...	...	...	N.S.W.	Q.	...	Trunks ...	Woody, whitish, rather thick, margin ochrey.
351	...	...	...	...	...	Q.	...	Trunks ...	Leathery, flexible, hairy, brownish, sessile, somewhat kidney shaped.
352	...	...	...	V.	...	Q.	B.	Trunks ...	Corky to leathery, pale, concentrically grooved, downy.
352A	...	...	...	...	...	Q.	...	Trunks ...	Hard, sessile, lobed, deeply zoned, tawny, velvety.
353	...	...	...	V.	...	...	...	Bark ...	Kidney or shell shaped, leathery, greyish-fawn colour, silky.
354	...	...	...	...	N.S.W.	Q.	...	Trunks ...	Corky, kidney shaped, tan colour to whitish, downy, zoneless.
355	...	...	...	...	...	Q.	...	Trunks, chiefly Poplar	Corky, white, at length turning ash coloured, sessile, zoneless.
356	...	...	...	V.	N.S.W.	...	B.	Stumps of beech, &c.	Leathery, flaccid, hairy, pale, zoned.
357	...	...	...	...	N.S.W.	Q.	...	Trunks ...	Shell shaped, smoky behind, pale in front, dotted with tubercles.
358	...	...	...	V.	...	Q.	...	Trunks ...	Snow white, corky to leathery, pitted, rather discoid behind.
359	...	...	...	...	N.S.W.	Q.	...	Trunks ...	Corky, ochrey to white, hemispherical, zoned, margin lobed.
360	...	...	...	...	N.S.W.	Q.	...	Dead wood ...	Corky, white to pale, margin slightly waved.
361	...	...	...	V.	...	...	B.	Wood, pine ...	Leathery, bright brown, zoned, margin yellowish. Common.
362	...	S.A.	...	V.	...	Q.	...	Trunk ...	Leathery, soft, downy, rust coloured, obsoletely zoned.
363	...	...	...	...	N.S.W.	...	...	Wood ...	Entirely white. Compact, woody, sessile, concentrically furrowed.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
23. SCHIZOPHYLLUM.—Fries,					
364	525	V. 2705	<i>S. commune</i> ...	Fries, S.M. I. 333 (1821) ...	Common schizophyllum ...
365	"	" 2706	<i>S. multifidum</i> ...	Fries, S.M. I. 333 (1821) ...	Multifid schizophyllum ...
24. METRARIA.—Cooke and Mass.,					
366	190	IX. 348	<i>M. insignis</i> ...	Cooke and Mass., Grev. XIX. 105 (1891) ...	Remarkable metraria ...
25. VOLVARIA.—Fries,					
367	191	V. 2712	<i>V. bombycina</i> ...	Fries, S.M. I. 277 (1821) ...	Silky volvar ...
368	195	" 2740	<i>V. parvula</i> ...	Weinm., Ross 238 (1836) ...	Little volvar ...
369	194	" 2735	<i>V. speciosa</i> ...	Fries, S.M. I. 278 (1821) ...	Beautiful volvar ...
370	192	" 2717	<i>V. Taylora</i> ...	Berk., Outl. 140 (1860) ...	Taylor's volvar ...
371	193	" 2733	<i>V. xanthocephala</i> ...	Berk., Hook., Lond. Journ. IV. 45 (1845) ...	Yellow-headed volvar ...
26. ANNULARIA.—Schulz. Verh. Zool. Bot.					
372	196	IX. 350	<i>A. insignis</i> ..	Cooke and Mass., Grev. XVIII. 3 (1889) ...	Remarkable annularia ...
27. PLUTEUS.—Fries,					
373	197	V. 2747	<i>P. cervinus</i> ...	Fries, Epier. 140 (1838) ...	Fawn pluteus ..
374	198	" 2806	<i>P. Wehlianus</i> ...	F. v. M., Grev. XV. 93 (1887) ...	Wehl's pluteus ...
28. ENTOLOMA.—Fries,					
375	202	V. 2828	<i>E. Bloxami</i> ...	Berk. and Br., Outl. 143 (1860) ...	Bloxam's entolome ...
376	200	IX. 354	<i>E. galbineum</i> ...	Cooke and Mass., Grev. XVII. 7 (1888) ...	Yellowish entolome ...
377	201	" 359	<i>E. laeticolor</i> ...	Cooke and Mass., Grev. XVI. 31 (1887) ...	Bright-coloured entolome ...
378	199	" 360	<i>E. melaniceps</i> ...	Cooke and Mass., Grev. XVI. 31 (1887) ...	Black-headed entolome ...
379	203	V. 2863	<i>E. panniculus</i> ...	Berk., Fl. Tasm. II. 245 (1860) ...	Ragged entolome ...
29. CLITOPILUS.—Fries,					
380	204	V. 2900	<i>C. cancrinus</i> ...	Fries, Epier. 150 (1838) ...	Crab-like clitopile ...
381	...	...	<i>C. cyathoides</i> ...	Cooke and Mass., Grev. XXI. 36 (1892) ..	Goblet-like clitopile ...
30. LEPTONIA.—Fries,					
382	207	V. 2945	<i>L. aquila</i> ...	Fries, Epier. 154 (1838) ...	Eagle leptonia ...
383	205	" 2923	<i>L. lampropoda</i> ...	Fries, S.M. I. 203 (1821) ...	Brilliant-stalked leptonia ...
384	208	...	<i>L. melanura</i> ...	Cooke and Mass., Grev. XIX. 89 (1891) ...	Black-tailed leptonia ...
385	206	IX. 372	<i>L. quinquecolor</i> ...	Cooke and Mass., Grev. XVII. 7 (1888) ...	Five-coloured leptonia ...
31. NOLANEA.—Fries,					
386	210	V. 2967	<i>N. mammosa</i> ...	Fries, Mon. Hym. I. 293 (1837) ...	Papillate nolanea ...
387	209	" 2960	<i>N. pascua</i> ...	Fries, S.M. I. 205 (1821) ...	Pasture nolanea ...
388	211	" 2980	<i>N. rufo-carnea</i> ...	Berk. Outl. 148 (1860) ...	Reddish-brown nolanea ...
32. ECCILIA.—Fries,					
389	212	V. 3030	<i>E. rhodocylix</i> ...	Lasch. in Fries, Hym. Eur. 213 (1874) ...	Rose-cup eccilia ...
33. CLAUDOPUS.—Smith,					
390	213	V. 3037	<i>C. variabilis</i> ..	Fries, Hym. Eur. 213 (1874) ...	Variable claudopus ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.	
	W.A.	S.A.	T.	V.	N.S.W.	Q.				
Obs. I. 103 (1815).— <i>Agaricus</i> .										
364	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Dead wood	...	Dry, white or greyish, scarcely exceeding inch in diameter. Cosmopolitan.
365	...	...	...	V.	...	...	...	Wood	...	Deeply cut into numerous lobes, becoming pale yellow.
Grev. XIX. 104 (1891).										
366	...	...	...	V.	...	...	...	Woods	...	Margin cream colour, disc darker and reddish brown, viscid, shining when dry.
S.M. I. 277 (1821).— <i>Agaricus</i> , <i>Amanita</i> .										
367	...	...	...	V.	...	...	B.	Decayed wood	...	Large. Fleshy, silky, fawn to brown, globose and viscid at first. <i>Edible</i> .
368	...	...	...	...	...	...	Q.	Pastures, after stormy weather	...	White. Rather fleshy, downy, conical at first. Stem stuffed, silky.
369	...	...	...	V.	...	...	B.	Dung-hills, sides, &c.	road-	Large. Fleshy, smooth, viscid or polished, grey. Stem rather bulbous.
370	...	...	T.	V.	...	...	B.	Ground	...	Thin, livid. Stem pale, solid, smooth.
371	W.A.	...	...	...	...	...	...	Ground	...	Golden yellow, spotted with white from remains of volva. Stem bulbous.
Gesell. 49 (1868).— <i>Agaricus</i> , <i>Chamæota</i> .										
372	...	...	...	V.	...	...	...	Ground	...	Fleshy, pale, cuticle broken up into broad darker scales. Stem short, thick.
Epier. 140 (1838).— <i>Agaricus</i> .										
373	...	...	T.	V.	...	...	B.	Trunks of trees	...	Large. Fleshy, dull tawny, smooth, then clad with evanescent scales.
374	...	S.A.	...	V.	...	...	...	Rotten wood on ground	...	Fleshy, shining, ochrey to white, disc darker; stem 6 to 8 inches long, thick.
Epier. 143 (1838).— <i>Agaricus</i> .										
375	...	S.A.	...	V.	...	...	B.	Open exposed pastures	...	Fleshy, compact, blackish blue. Stem slightly tapering upwards.
376	...	...	...	V.	...	...	...	Ground	...	Sulphur colour. Rather fleshy, almost saffron colour. Stem hollow.
377	...	S.A.	...	V.	...	...	...	Sandy soil	...	Rather fleshy, shining, amethyst colour. Stem thin, nearly solid.
378	...	S.A.	...	V.	...	...	...	Ground	...	Fleshy, compact, dark sooty brown. Stem solid, short, pale.
379	...	...	T.	...	...	...	...	Among ferns	...	Thin, bell shaped, dark violet. Stem thickened at base, and downy.
Epier. 148 (1838).— <i>Agaricus</i> .										
380	...	S.A.	...	V.	...	Q.	B.	Grass fields	...	Small and beautiful. Fleshy to membranaceous, flesh colour to white.
381	...	...	...	V.	...	...	...	Under burnt logs	...	Rather thin, pale, finally funnel shaped. Stem hollow, white and woolly at base.
S.M. I. 201 (1821).— <i>Agaricus</i> .										
382	...	S.A.	...	...	...	...	...	Ground	...	Rather membranaceous, bay brown. Stem short, stuffed.
383	...	...	...	V.	...	...	B.	Pastures	...	Rather fleshy, mouse coloured, or steel grey, or sooty. Common.
384	...	...	...	V.	...	...	...	Ground	...	Bell shaped, shining black, silky. Stem cylindrical.
385	...	...	...	V.	...	...	...	Black loam	...	Membranaceous, disc brownish brick red, margin yellowish.
S.M. I. 204 (1821).— <i>Agaricus</i> .										
386	...	...	...	V.	...	...	B.	Meadows	...	Sub-membranaceous, papillate, tawny. Stem hollow, polished.
387	...	...	T.	...	...	...	B.	Pastures	...	Membranaceous, shining like silk when dry. Stem silky fibrous.
388	...	...	...	V.	...	...	B.	Heaths	...	Small. Sub-membranaceous, red brown, indistinctly scaly.
S.M. I. 207 (1821).— <i>Agaricus</i> .										
389	...	...	...	V.	...	...	B.	Rotten wood	...	Membranaceous, tawny, when dry flocculose, grey.
Seem. Journ.— <i>Agaricus</i> .										
390	...	...	...	V.	...	Q.	B.	Sticks, &c.	...	Very small. Sub-membranaceous, silky with white down. Common.

SYSTEMATIC ARRANGEMENT

Number	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
34. PHOLIOTA.—Fries,					
391	226	V. 3103	<i>P. allantopoda</i> ...	Berk., Hook., Lond. Journ. IV. 45 (1845)	Sausage-stalked pholiota ...
392	218	" 3053	<i>P. blattaria</i> ...	Fries, S.M. I. 246 (1821)	Cockroach-like pholiota ...
393	224	" 3094	<i>P. congesta</i> ...	Kalb., Grev. IX. 147 (1881)	Congested pholiota ...
394	220	...	<i>P. disrupta</i> ...	Cooke and Mass., Grev. XIX. 89 (1891)	Disrupted pholiota ...
395	223	V. 3104	<i>P. effusa</i> ...	Kalb., Grev. IX. 147 (1881)	Expanded pholiota ...
396	215	" 3050	<i>P. erebia</i> ...	Fries, S.M. I. 246 (1821)	Lurid pholiota ...
397	230	" 3128	<i>P. eriogena</i> ...	Fries, Pl. Preiss. II. 132 (1846)	Woolly pholiota ...
398	227	" 3109	<i>P. flammans</i> ...	Fries, S.M. I. 244 (1821)	Flame-coloured pholiota ...
399	229	" 3130	<i>P. marginata</i> ...	Fries, Hym. Eur. 225 (1874)	Margined pholiota ...
400	228	" 3129	<i>P. mutabilis</i> ...	Fries, S.M. I. 245 (1821)	Changeable pholiota ...
401	232	" 3137	<i>P. mycenoides</i> ...	Fries, S.M. I. 246 (1821)	Myccena-like pholiota ...
402	222	" 3071	<i>P. phylieigena</i> ...	Berk., Linn. Journ. XV. 52 (1877)	Phyllea-growing pholiota ...
403	219	" 3055	<i>P. praecox</i> ...	Fries, Hym. Eur. 217 (1874)	Precocious pholiota ...
404	221	" 3065	<i>P. pudica</i> ...	Fries, Hym. Eur. 218 (1874)	Modest pholiota ...
405	231	" 3135	<i>P. pumila</i> ...	Fries, El. 29 (1828)	Dwarfish pholiota ...
406	217	IX. 394	<i>P. recedens</i> ...	Cooke and Mass., Grev. XVIII. 25 (1889)	Receding pholiota ...
407	225	V. 3102	<i>P. spectabilis</i> ...	Fries, El. 28 (1828)	Notable pholiota ...
408	216	" 3052	<i>P. togularis</i> ...	Fries, Hym. Eur. 216 (1874)	Gowned pholiota ...
35. LOCELLINA.—Gill. Champ.					
409	214	V. 3141	<i>L. eyenopotamia</i> ...	Sacc. Syll., V. 762 (1887)	Swan river locellinia ...
36. INOCYBE.—Fries,					
410	234	V. 3149	<i>I. cincinnata</i> ...	Fries, S.M. I. 256 (1821)	Curly inocybe ...
411	236	" 3165	<i>I. flocculosa</i> ...	Fries, Hym. Eur. 229 (1874)	Flocculous inocybe ...
412	238	...	<i>I. gigaspora</i> ...	Cooke, Handb. Aust. Fungi 47 (1892)	Large-spored inocybe ...
413	237	V. 3235	<i>I. gomphodes</i> ...	Kalb., Grev. VIII. 152 (1880)	Pap-like inocybe ...
414	235	" 3155	<i>I. lanuginosa</i> ...	Fries, S.M. I. 257 (1821)	Woolly inocybe ...
415	233	" 3148	<i>I. plumosa</i> ...	Fries, Mon. Hym. (1857)	Downy inocybe ...
416	239	IX. 421	<i>I. victoriæ</i> ...	Cooke and Mass., Grev. XVI. 72 (1888)	Victorian inocybe ...
37. HEBELOMA.—Fries,					
417	245	IX. 426	<i>H. arenicolor</i> ...	Cooke and Mass., Grev. XVII. 7 (1888)	Sand-coloured hebelome ...
418	240	V. 3259	<i>H. fastibile</i> ...	Fries, Epier. 178 (1838)	Disagreeable hebelome ...
419	241	" 3260	<i>H. glutinosum</i> ...	Lindgr., Bot. Not. 199 (1845)	Glutinous hebelome ...
420	...	...	<i>H. griseum</i> ...	Cooke and Mass., Grev. XXI. 36 (1892)	Grey hebelome ...
421	242	V. 3268	<i>H. mesophæum</i> ...	Fries, Epier. 179 (1838)	Dusky-centred hebelome ...
421A	"	"	<i>H. mesophæum</i> , var. <i>holophæum</i> ...	Fries, Hym. Eur. 240 (1874)	Dusky hebelome ...
422	244	" 3291	<i>H. nudipes</i> ...	Fries, Epier. 181 (1838)	Naked-stalked hebelome ...
423	243	" 3275	<i>H. olidum</i> ...	Cooke and Mass., Grev. XV. 93 (1887)	Strong-smelling hebelome ...
424	246	" 3319	<i>H. petiginosum</i> ...	Fries, S.M. I. 259 (1821)	Scabby hebelome ...
38. FLAMMULA.—Fries,					
425	252	IX. 437	<i>F. avellanea</i> ...	Cooke and Mass., Grev. XVIII. 3 (1889)	Nut-brown flammula ...
426	253	V. 3344	<i>F. Baileyi</i> ...	Berk. and Br., Linn. Trans. II. 54 (1883)	Bailey's flammula ...
427	258	" 3369	<i>F. flavida</i> ...	Fries, S.M. I. 250 (1821)	Yellowish flammula ...
428	257	" 3363	<i>F. fusa</i> ...	Fries, Hym. Eur. 247 (1874)	Fusiform flammula ...
429	261	" 3382	<i>F. hybrida</i> ...	Fries, Mon. Hym. I. 360 (1857)	Hybrid flammula ...
430	251	IX. 438	<i>F. hyperion</i> ...	Cooke and Mass., Grev. XVI. 72 (1888)	Hyperion flammula ...
431	260	V. 3373	<i>F. inopoda</i> ...	Fries, S.M. I. 251 (1821)	Fibril-stalked flammula ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
S.M. I. 240 (1821).— <i>Agaricus</i> .									
391	W.A.	...	...	V.	...	...	...	Ground ...	Fleshy, golden yellow. Stem elongatedly bulbous at base.
392	...	...	...	V.	...	...	B.	Ground ...	Elegant, small. Rather fleshy, rust coloured, margin grooved.
393	...	...	...	V.	...	...	...	Trunks ...	Fleshy, the size of a pea, rather mealy, brownish.
394	...	...	...	V.	...	...	...	Ground ...	Fleshy, creamy white, at first smooth, then cracked deeply.
395	...	...	...	V.	...	...	...	Wood ...	Fleshy, white, breaking into polygonal wart-like spaces.
396	...	...	...	V.	...	...	B.	Grassy places ...	Fleshy, rather viscid, lurid. Stem hollow, fibrillose to scaly.
397	W.A.	...	...	...	...	...	...	Trunks ...	Fleshy, rust coloured. Stem with dense woolly mycelium at base.
398	...	...	...	...	...	...	Q.	B. Pinewoods ...	Fleshy, tawny, with sulphureous scales. Stem stuffed, thin, hollow.
399	...	...	...	V.	...	...	B.	Ground, among firs	Rather fleshy, cinnamon, margin striate. Stem hollow, not scaly.
400	...	...	T.	V.	...	...	B.	Trunks or ground ...	Fleshy, cinnamon, becoming pale. Stem rough with scales. <i>Edible</i> .
401	...	...	...	V.	...	...	B.	Ground, in damp places	Membranaceous, rust coloured, tawny or pale when dry.
402	...	...	...	...	...	...	Q.	Trunks of <i>Phytica</i> ...	Fleshy, tawny. Stem thick below, tapering upwards.
403	W.A.	...	...	...	...	...	B.	Gardens and pastures	Fleshy, soft, white to yellowish. Stem downy or mealy. <i>Edible</i> .
404	...	...	...	V.	...	...	B.	Trunks, &c. ...	Largish. Fleshy, dry, smooth, whitish buff, modestly coloured. <i>Edible</i> .
405	...	...	...	V.	N.S.W.	...	B.	Woods ...	Somewhat fleshy, hemispherical. Stem hollow, slender.
406	...	...	...	V.	...	...	...	Ground ...	Rather fleshy, golden tawny. Stem elongated, cylindrical
407	...	...	...	V.	N.S.W.	...	B.	Dead stumps—Oak, &c.	Large, compact. Golden orange or tawny buff, scales silky and broad. Odour bad.
408	...	...	...	V.	...	...	B.	Grassy places, &c....	Fleshy, pale ochre. Stem hollow, cracking. Ring hanging like toga.
Fr. 428 (1874).— <i>Agaricus</i> , <i>Acetabularia</i> .									
409	W.A.	...	...	...	...	...	...	Ground ...	Gills pale fawn colour, leaving a free space round stem. Very rare.
S.M. I. 254 (1821).— <i>Agaricus</i> , <i>Hebeloma</i> .									
410	...	...	...	V.	...	...	B.	Shady woods ...	Rather fleshy, scaly. Stem solid, thin, scaly.
411	...	...	...	V.	...	...	B.	Bare soil and among grass	Somewhat fleshy, tawny brown. Stem pale reddish.
412	...	...	...	V.	...	...	...	Ground ...	Rather fleshy, yellow brown. Stem abruptly rooting.
413	...	...	...	...	N.S.W.	...	...	Ground ...	Rather fleshy, with globose pap-like apex, tawny.
414	W.A.	...	...	...	...	...	B.	Ground ...	Rather fleshy, umber, becoming yellowish. Stem solid, thin, scaly.
415	...	...	...	V.	...	...	B.	Moist pine woods ...	Rather fleshy. Odour weak, not unpleasant. Stem slender.
416	...	...	...	V.	...	...	...	Grassy ground ...	Rather fleshy, viscid, whitish, shining. Stem stuffed, white.
S.M. I. 249 (1821).— <i>Agaricus</i> .									
417	...	...	...	V.	...	...	...	Ground ...	Fleshy, rather viscid, dingy ochre or sand colour. Stem cylindrical.
418	...	...	...	V.	...	...	B.	Woods ...	Compact, viscid, yellowish tan or tan colour. Stem solid, white.
419	...	...	...	V.	...	...	B.	Among dead leaves	Fleshy, viscous with a tenacious gluten, yellowish white. Stem stuffed.
420	...	...	...	V.	...	...	...	Ground in woods ...	Fleshy, mouse grey or pale silver grey, glutinous, smooth, shining when dry. Odour unpleasant.
421	...	...	...	V.	...	...	B.	Woods, &c. ...	Rather fleshy, viscid, yellowish tan, disc bay. Stem slender, white. Common.
421A	...	...	...	V.	...	...	...	Ground ...	Dark brown, veil resembling a ring. Stem becoming brown.
422	...	...	...	...	...	...	Q.	B. Ground ...	Fleshy, slightly viscid, tan coloured, thin, pale. Stem solid, white.
423	...	S.A.	...	...	...	...	...	Stony ground ...	Fleshy, viscid, full red brown. Stem hollow, smooth. Odour fetid.
424	...	...	...	V.	...	...	B.	Ground in shady places	Rather fleshy, dry, brown, circumference silky grey. Stem slender, powdery, brick red.
S.M. I. 250 (1821).— <i>Agaricus</i> , <i>Paxillus</i> .									
425	...	...	...	V.	...	...	Q.	Sandy ground ...	Fleshy, nut brown. Stem tapering upwards, grooved.
426	...	...	...	...	...	...	Q.	Rotten wood ...	Orange. Bell shaped to hemispherical, woolly, sprinkled with reddish-yellow mealy particles.
427	...	...	...	V.	N.S.W.	...	B.	Trunks—Pine, &c.	Fleshy, yellow, smooth, moist. Stem yellow, then rusty.
428	...	S.A.	...	V.	...	...	B.	Ground and fallen logs	Compact, rather viscid, flesh becoming yellow. Odour not unpleasant.
429	...	...	...	V.	...	...	B.	Fir stumps ...	Fleshy, moist, at first cinnamon brown then golden tawny. Veil forming ring.
430	...	...	...	V.	...	...	...	Stumps (?) ...	Fleshy, golden tawny, then darker. Stem tapering downwards, furrowed.
431	...	...	...	V.	...	...	B.	Pine trunks ...	Fleshy, moist, honey tan colour, becoming pale. Stem fibrillose.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
38. FLAMMULA.—Fries,					
432	259	V. 3379	<i>F. limonia</i> ...	Cooke and Mass., Grev. XV. 94 (1887) ...	Lemon-coloured flammula ...
433	369	" 3326	<i>F. paradoxa</i> ...	Kalch., Fung. Hung. t. 16, f. 1 (1873) ...	Paradoxical flammula ...
434	262	" 3381	<i>F. penetrans</i> ...	Fries, Hym. Eur. 250 (1874) ...	Penetrating flammula ...
434A	"	" "	<i>F. penetrans</i> , var. <i>australis</i> ...	F. v. M., Linn. Journ. XIII. 158 (1873) ...	Southern flammula ...
435	255	" 3346	<i>F. peregrina</i> ...	Fries, Elen. I. 31 (1828) ...	Foreign flammula ...
436	265	" 3389	<i>F. piecea</i> ...	Fries, Hym. Eur. 251 (1874) ...	Bitter flammula ...
437	254	IX. 446	<i>F. prasina</i> ...	Cooke and Mass., Grev. XVIII. 3 (1889) ...	Leek-green flammula ...
438	266	V. 3393	<i>F. purpureo-nitens</i> ...	Cooke and Mass., Grev. XV. 94 (1887) ...	Shining-purple flammula ...
439	249	IX. 445	<i>F. rubra</i> ...	Cooke and Mass., Grev. XIX. 46 (1890) ...	Red flammula ...
440	263	V. 3385	<i>F. sapinea</i> ...	Fries, Epicr. 189 (1838) ...	Pine-wood flammula ...
441	256	" 3358	<i>F. spumosa</i> ...	Fries, S.M. I. 252 (1821) ...	Frothy flammula ...
442	248	"	<i>F. veluticeps</i> ...	Cooke and Mass., Grev. XIX. 89 (1891) ...	Velvet-capped flammula ...
443	247	" 3323	<i>F. vinosa</i> ...	Fries, Hym. Eur. 244 (1874) ...	Wine-coloured flammula ...
444	250	"	<i>F. xanthophylla</i> ...	Cooke and Mass., Handb. Aust. Fungi 50 (1892) ...	Yellow-gilled flammula ...
39. NAUCORIA.—Fries,					
445	267	V. 3412	<i>N. anginea</i> ...	Fries, Epicr. 193 (1838) ...	Snake-like naucoria ...
446	281	" 3506	<i>N. Bowmani</i> ...	Berk., Linn. Journ. XIII. 158 (1873) ...	Bowman's naucoria ...
447	269	" 3435	<i>N. ceroides</i> ...	Fries, Epicr. 195 (1838) ...	Wax-like naucoria ...
448	278	" 3484	<i>N. Drummondii</i> ...	Berk., Hook., Lond. Journ. IV. 46 (1845) ...	Drummond's naucoria ...
449	282	" 3514	<i>N. escharoides</i> ...	Fries, S.M. I. 260 (1821) ...	Scabby naucoria ...
450	275	IX. 458	<i>N. fraterna</i> ...	Cooke and Mass., Grev. XVI. 31 (1887) ...	Fraternal naucoria ...
451	283	V. 3495	<i>N. frusticola</i> ...	Berk., Linn. Journ. XIII. 158 (1873) ...	Tufted naucoria ...
452	270	" 3437	<i>N. melinoides</i> ...	Fries, S.M. I. 266 (1821) ...	Honey-like naucoria ...
453	272	" 3426	<i>N. nasuta</i> ...	Kalch., Grev. VIII. 152 (1880) ...	Long-nosed naucoria ...
454	276	" 3469	<i>N. pediales</i> ...	Fries, S.M. I. 290 (1821) ...	Field naucoria ...
455	271	" 3440	<i>N. pusiola</i> ...	Fries, S.M. I. 264 (1821) ...	Little naucoria ...
456	268	" 3427	<i>N. russa</i> ...	Cooke and Mass., Grev. XV. 94 (1887) ...	Red naucoria ...
457	273	" 3450	<i>N. scolecina</i> ...	Fries, Epicr. 194 (1838) ...	Worm-eaten naucoria ...
458	277	" 3470	<i>N. semiorbicularis</i> ...	Fries, Mon. Hym. I. 376 (1857) ...	Hemispherical naucoria ...
459	280	" 3507	<i>N. stiparia</i> ...	Fries, S.M. I. 261 (1821) ...	Veiled naucoria ...
460	279	" 3486	<i>N. tenuilenta</i> ...	Fries, S.M. I. 268 (1821) ...	Dripping naucoria ...
461	274	" 3457	<i>N. triscopoda</i> ...	Fries, Mon. Hym. I. 375 (1857) ...	Hair-stalked naucoria ...
40. GALERA.—Fries,					
462	286	V. 3568	<i>G. hypnorum</i> ...	Fries, S.M. I. 267 (1821) ...	Moss galera ...
463	287	" 3574	<i>G. minuta</i> ...	Quel., Champ. Jura III. 10 (1873) ...	Minute galera ...
464	285	" 3549	<i>G. peroxydata</i> ...	Berk., Hook., Lond. Journ. II. 411 (1843) ...	Peroxide galera ...
465	284	" 3537	<i>G. tenera</i> ...	Fries, Hym. Eur. 267 (1874) ...	Delicate galera ...
41. TUBARIA.—Smith,					
466	288	V. 3584	<i>T. furfuracea</i> ...	Fries, Hym. Eur. 272 (1874) ...	Mealy tubaria ...
467	289	" 3597	<i>T. inquilina</i> ...	Fries, Hym. Eur. 274 (1874) ...	Little tubaria ...
467A	"	" "	<i>T. inquilina</i> , var. <i>ecbola</i> ...	Fries, Hym. Eur. 275 (1874) ...	Clay-coloured tubaria ...
468	"	"	<i>T. strigipes</i> ...	Cooke and Mass., Grev. XXI. 36 (1892) ...	Rough-stalked tubaria ...
42. CREPIDOTUS.—Fries,					
469	292	V. 3599	<i>C. alveolus</i> ...	Lasch., in Fries Epicr. 210 (1836) ...	Alveolate crepidotus ...
470	304	" 3665	<i>C. auricula</i> ...	Berk., Fl. Tasm. II. 246 (1860) ...	Eared crepidotus ...
471	300	" 3627	<i>C. cussiacolor</i> ...	Berk., Fl. Tasm. II. 246 (1860) ...	Cinnamon crepidotus ...
472	299	" 3601	<i>C. epigeus</i> ...	Berk. and Br., Ann. Nat. Hist. IX. 179 (1882) ...	Earth-borne crepidotus ...
473	294	" 3610	<i>C. globigera</i> ...	Berk., Linn. Journ. XIII. 158 (1873) ...	Globose-spored crepidotus ...
474	298	" 3620	<i>C. haustellaris</i> ...	Fries, S.M. I. 274 (1821) ...	Damp-loving crepidotus ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
S.M. I. 250 (1821).— <i>Agaricus Paxillus</i> —continued.									
432	...	S.A.	...	V.	N.S.W.	...	...	Rich soil ...	Fleshy, moist, sulphur coloured. Stem stuffed, yellowish white.
433	...	...	...	V.	...	Q.	B.	Ground ...	Fleshy, dry, downy, red umber. Stem solid, yellow or reddish.
434	...	S.A.	...	V.	N.S.W.	...	...	Wood ...	Fleshy, dry, yellow tawny or golden. Stem silky, with fleeting veil.
434A	...	...	...	V.	N.S.W.	...	...	Soil (probably covering pine chips)	Orange tawny, with stem and gills paler.
435	W.A.	...	...	V.	...	...	...	Trunks ...	Fleshy, rust coloured, corrugated. Stem solid, smooth.
436	...	...	...	V.	...	Q.	B.	Dead trunks of <i>Encyphalartos Denisonii</i>	Rather fleshy, moist, red to bay cinnamon, becoming pale. Stem thin, almost umber, tapering upwards, without veil.
437	...	...	...	V.	...	...	...	Ground ...	Fleshy, dry, silky, leek green. Stem straight, stuffed, lemon yellow.
438	W.A.	...	...	V.	...	Q.	...	Wood ...	Fleshy, shining, purple brown. Stem ascending, solid, paler.
439	...	...	...	V.	...	...	...	Ground ...	Fleshy, shining, red, with tinge of purple. Stem hollow, paler.
440	...	S.A.	...	V.	N.S.W.	Q.	B.	Fallen branches and chips	Compact, golden tawny, fluffy to scaly, then cracked. Strong odour.
441	...	S.A.	...	V.	...	...	B.	Woods ...	Small, stem tall. Fleshy, viscid, yellow. Very common.
442	...	...	...	V.	...	...	...	Among grass on hill-sides	Densely and shortly velvety, bay brown. Stem expanded upwards into cap, rather short, and of same colour.
443	...	S.A.	...	V.	...	...	B.	Ground ...	Fleshy, dry, rusty-fawn colour. Stem solid, delicately fluffy.
444	...	S.A.	...	V.	...	...	...	Wood ...	Ochrey yellow. Fleshy, compact, hard when dry. Stem lateral, short.
S.M. I. 260 (1821).— <i>Agaricus</i> .									
445	...	...	...	...	...	Q.	B.	Ground ...	Slightly fleshy, yellowish to tan colour. Stem with white fibrils.
446	...	...	...	...	...	Q.	...	Ground ...	Small. Rough with woolly tufts. Stem slender, fluffy.
447	...	...	...	V.	...	...	B.	Burnt soil ...	Rather fleshy, orbicular, ochrey. Stem naked, yellow, rusty at base.
448	W.A.	...	...	V.	...	...	...	Rotten wood ...	Viscid, when young very white. Stem mealy above, downy at base.
449	...	...	...	...	...	Q.	B.	Bare ground ...	Gregarious, fragile. Rather fleshy, whitish-tan colour, scabby scales.
450	...	...	...	V.	...	...	...	Logs ...	Tawny ferruginous. Stem elongated, thin, hollow, of same colour.
451	...	S.A.	...	...	N.S.W.	...	...	Roots of grass, &c.	Densely tufted. Tawny. Stem slender, mealy, thickened downwards.
452	W.A.	...	...	V.	...	Q.	B.	Among grass ...	Somewhat fleshy, tawny, ochrey when dry. Stem hollow, yellow.
453	...	...	...	...	N.S.W.	...	...	Ground ...	Rather fleshy, ochrey, with elongated teat-like umbo
454	...	S.A.	...	V.	...	...	B.	Pastures ...	Somewhat fleshy, yellow ochrey to tan colour. Very common.
455	...	...	...	V.	...	...	B.	Ground ...	Slightly fleshy, rather viscid, tawny yellow. Stem thread-like, shining.
456	...	...	...	V.	...	...	...	Ground ...	Thin, brick red. Stem nearly of same colour, whitish downy below.
457	...	S.A.	...	...	...	...	B.	Moist ground ...	Rather fleshy, ferruginous bay. Stem rusty, sprinkled with white meal.
458	...	...	...	V.	...	Q.	B.	Lawns and pastures	Rather fleshy, hemispherical, somewhat viscid, tawny or ochrey.
459	...	...	...	V.	...	...	B.	Soil, fern stems, &c.	Rather fleshy, with downy scales, red to rust colour.
460	...	...	...	V.	...	...	B.	Moist woods ...	Sub-membranaceous, rust colour, tan colour when dry. Stem polished.
461	...	...	...	V.	...	...	B.	Old wood ...	Rather fleshy, bay brown, ochrey when dry. Stem hair-like, rusty.
S.M. I. 264 (1821).— <i>Agaricus</i> .									
462	...	S.A.	...	V.	...	...	B.	Among moss ...	Minute. Membranaceous, bell shaped, sub-papillate, tawny. Common.
463	...	...	...	V.	...	...	B.	Decayed wood ...	Membranaceous, tawny, streaked. Stem almost hair-like.
464	...	...	...	...	...	Q.	...	Ground ...	Membranaceous, reddish brown, bell shaped. Stem very thin.
465	...	S.A.	T.	V.	...	...	B.	Grassy places, manure, &c.	Small, delicate. Sub-membranaceous, nearly conical, buff. Common.
Seem. Journ. (1870).— <i>Agaricus</i> .									
466	...	S.A.	T.	V.	...	Q.	B.	Chips, &c. ...	Small. Somewhat fleshy, at first clothed with silky evanescent scales, rich umber.
467	...	...	...	...	N.S.W.	...	B.	Chips ...	Minute. Sub-membranaceous, brown. Stem hollow, dark brown. Common.
467A	...	...	...	...	N.S.W.	...	B.	Grass roots ...	Clay coloured. Stem rooting; gills crowded, rusty.
468	...	...	...	V.	...	...	...	In tufts among grass	Hemispherical, tawny yellow, with conical spreading scales. Stem slender.
S.M. I. 272 (1821).— <i>Agaricus</i> .									
469	...	...	...	V.	...	...	B.	Old stumps ...	Fleshy, soft, ochrey brown, contracted, downy to shaggy behind.
470	...	...	T.	...	...	...	...	Dead wood ...	Sessile, shell shaped, cream colour. Flesh thick, brittle when dry.
471	...	S.A.	T.	...	...	...	...	Rotten bark ...	Mealy, cinnamon. Stem very short, slender, white, downy.
472	...	S.A.	...	V.	...	...	B.	Ground ...	Fragile, reddish grey, kidney shaped; base shaggy, whitish.
473	...	...	...	V.	...	...	...	Wood ...	Kidney shaped, tapering at base. About an inch long and wide.
474	...	S.A.	...	...	...	...	B.	Rotten trunks of <i>Eucalyptus viminalis</i>	Rather fleshy, flaccid, tan coloured. Stem tapering upwards, hairy.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
42. CREPIDOTUS.—Fries,					
475	295	V. 3602	<i>C. hepatochrous</i> ...	Berk., Hook., Lond. Journ. VII. 574 (1848)	Liver-coloured crepidotus ...
476	301	„ 3628	<i>C. insidiosus</i> ...	Berk., Hook., Lond. Journ. VII. 574 (1848)	Insidious crepidotus ...
477	296	„ 3603	<i>C. interceptus</i> ...	Berk., Fl. Tasm. II. 246 (1860)...	Interposed crepidotus ...
478	303	„ 3663	<i>C. leptomorphus</i> ...	Berk., Fl. Tasm. II. 246 (1860)...	Delicate crepidotus ...
479	302	„ 3641	<i>C. lepton</i> ...	Berk., Hook., Lond. Journ. IV. 46 (1845)	Thin crepidotus ...
480	293	„ 3600	<i>C. mollis</i> ...	Fries, S.M. I. 274 (1821)	Soft crepidotus ...
481	291	„ 3598	<i>C. palmatus</i> ...	Fries, Mon. Hym. I. 398 (1837)	Palmate crepidotus ...
482	290	IX. 481	<i>C. phaeton</i> ...	Cooke and Mass., Grev. XV. 99 (1887)	Brilliant crepidotus ...
483	297	V. 3655	<i>C. stromaticus</i> ...	Cooke and Mass., Grev. XV. 94 (1887)	Stromate crepidotus ...
484	305	„ 3664	<i>C. turbidulus</i> ...	Berk., in Cooke's Handb. Aust. Fungi 60 (1892)...	Turbid crepidotus ...
43. CORTINARIUS.—Pers.					
485	361	V. 3763	<i>C. Archeri</i> ...	Berk., Fl. Tasm. II. 247 (1860) ...	Archer's cortinar ...
486	366	„ 3903	<i>C. bovinus</i> ...	Fries, Epicr. 297 (1838)	Ox cortinar ...
487	365	„ 3848	<i>C. cinnabarinus</i> ...	Fries, Epicr. 288 (1838)	Vermilion cortinar ...
488	366	„ 3740	<i>C. decoloratus</i> ...	Fries, Epicr. 270 (1838)	Discoloured cortinar ...
489	362	...	<i>C. erythræus</i> ...	Berk., Hook., Lond. Journ. IV. 48 (1845)	Blood-red cortinar ...
490	364	V. 3849	<i>C. sanguineus</i> ...	Fries, Epicr. 288 (1838)	Dark-red cortinar ...
491	363	„ 3788	<i>C. violaceus</i> ...	Fries, Epicr. 279 (1838)	Violet-coloured cortinar ...
44. PAXILLUS.—Fries,					
492	370	V. 4020	<i>P. crassus</i> ...	Fries, Hym. Eur. 404 (1874)	Thick paxil ...
493	368	„ 4010	<i>P. Encalyptornis</i> ...	Berk., Hook., Lond. Journ. IV. 49 (1845)	Eucalypt paxil ...
494	367	„ 4008	<i>P. Muelléri</i> ...	Berk., Linn. Journ. XIII. 159 (1873)	Mueller's paxil ...
495	371	„ 4021	<i>P. pannoides</i> ...	Fries, Hym. Eur. 404 (1874)	Panus-like paxil ...
45. AGARICUS.—Linn. Sp. Pl.					
496	306	V. 4039	<i>A. arvensis</i> ...	Schaeff., Icon. t. 310. 311 (1762)...	Field agaric ...
497	307	„ 4053	<i>A. campestris</i> ...	Linn., Sp. Pl. 1173 (1753)	Pasture agaric ...
498	„	„ 4054	<i>A. silvicola</i> ...	Vitt. Mang. (1835)	Wood agaric ...
499	310	IX. 559	<i>A. elatior</i> ...	Cooke and Mass., Grev. XVIII. 3 (1889)	Tall agaric ...
500	308	V. 4061	<i>A. silvaticus</i> ...	Schaeff., Icon. t. 242 (1762)	Sylvan agaric ...
501	309	„ 4081	<i>A. versipes</i> ...	Berk. and Br., Linn. Trans. II. 54 (1883)	Twisted-stalked agaric ...
46. STROPHARIA.—Fries, Summ. Veg.					
502	311	V. 4120	<i>S. coronilla</i> ...	Fries, Hym. Eur. 285 (1874)	Crowned stropharia ...
503	314	„ 4144	<i>S. merdaria</i> ...	Fries, Hym. Eur. 286 (1874)	Dung-borne stropharia ...
504	313	„ 4151	<i>S. semiglobata</i> ...	Fries, Hym. Eur. 287 (1874)	Hemispherical stropharia ...
505	312	„ 4124	<i>S. squamosa</i> ...	Fries, Hym. Eur. 285 (1874)	Scaly stropharia ...
47. HYPHOLOMA.—Fries,					
506	317	IX. 565	<i>H. adustum</i> ...	Cooke and Mass., Grev. XVIII. 3 (1889)	Scorched hypholome ...
507	318	V. 4212	<i>H. Candolleianum</i> ...	Fries, S.M. I. 296 (1821)	De Candolle's hypholome
508	„	„	<i>H. discretum</i> ...	Cooke and Mass., Grev. XXI. 37 (1892)	Separated hypholome ...
509	316	V. 4182	<i>H. dispersum</i> ...	Fries, Epicr. 222 (1838)	Scattered hypholome ...
510	315	„ 4178	<i>H. fasciculare</i> ...	Fries, S.M. I. 288 (1821)	Tufted hypholome ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.	
	W.A.	S.A.	T.	V.	N.S.W.	Q.				
S.M. I. 272 (1821).— <i>Agaricus</i> — <i>continued</i> .										
475	...	...	T.	...	...	...	...	Bark ...	...	Rather fleshy, liver coloured. Cap globose at first, with short central stem.
476	...	...	T.	...	...	...	...	Bark ...	...	Membranaceous, margin downy, yellowish brown when dry. Stem short and slender.
477	...	...	T.	V.	...	...	...	Bark ...	...	Kidney shaped, ochrey white. Cap of three layers, the middle one, white, interposed between two darker ones.
478	...	...	T.	...	...	...	...	Dead wood	...	Sessile, whitish, downy, fixed at apex by a few white threads.
479	W.A.	...	...	...	...	...	...	Bark ...	...	Tawny ochre. Stem obsolete, extremely short if present.
480	W.A.	S.A.	...	V.	...	Q.	B.	Old stumps	...	Gelatinous to fleshy, flaccid, pale. Stem obsolete.
481	...	...	T.	...	...	...	B.	Trunks ...	...	Fleshy, compact, rust coloured. Stem excentric or lateral.
482	...	...	...	V.	...	...	...	Ground (?)	...	Sub-membranaceous, brick red. Stem lateral, elongated.
483	W.A.	...	...	...	...	...	...	Bark ...	...	Sessile, flaccid, tan colour, arising from white woolly stroma.
484	...	...	T.	...	...	...	...	Wood ...	...	Sessile, kidney shaped, ochrey, smooth.
Syn. 16 (1801).— <i>Agaricus</i> .										
485	...	...	T.	...	...	...	...	Ground ...	...	Fleshy, violet brown. Stem stout, viscid, violet.
486	...	...	...	V.	...	...	B.	Woods ...	...	Fleshy, watery cinnamon. Stem stout, spongy, bulbous, grey.
487	...	...	...	V.	...	Q.	B.	Under trees	...	Fleshy, silky, vermilion, shining. Stem stuffed, short, vermilion.
488	...	...	...	V.	...	...	B.	Woods ...	...	Fleshy, viscid, soon dry, floccose and discoloured. Stem tapering from base.
489	W.A.	...	...	...	...	...	...	Ground ...	...	Small, blood red. Cap clothed with thick gelatinous coat. Stem short, viscid.
490	...	...	...	V.	...	...	B.	Woods ...	...	Entirely dark red. Fleshy, silky, or scaly. Stem stuffed, thin, hollow.
491	...	...	...	V.	...	...	B.	Woods ...	...	Dark violet. Fleshy, woolly to scaly. Stem bulbous, spongy, shaggy. <i>Edible</i> .
Gen. Hym. 8 (1836).— <i>Agaricus</i> , <i>Merulius</i> .										
492	...	...	...	...	...	Q.	B.	Ground ...	...	Fleshy, rust coloured. Stem stuffed, excentric, very short.
493	W.A.	...	...	...	...	...	...	Under <i>Eucalyptus</i> trees	...	Thick and fleshy, compact, tawny yellow. Stem transversely scaly.
494	...	...	...	V.	N.S.W.	Q.	...	Meadows ...	...	Dark brown, convex. Stem tawny, frosted.
495	...	...	...	V.	...	...	B.	Cellars, on sawdust, &c.	...	Fleshy, shell-shaped, dirty yellow or whitish ochre, sessile.
(1753).— <i>Psalliota</i> , <i>Pratella</i> .										
496	W.A.	...	T.	V.	N.S.W.	...	B.	Meadows, &c.	...	Very large, expanding late. Fleshy, flesh turning slightly yellow where bruised. <i>Edible</i> .
497	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Rich pastures	...	Fleshy, silky floccose, or scaly. Stem stuffed, ring median. <i>Edible</i> .
498	...	...	...	V.	...	...	B.	Woods ...	...	Smooth, shining white. Stem stuffed, elongated, somewhat bulbous.
499	...	...	...	V.	...	...	...	Ground ...	...	Thinly fleshy, brown, scaly. Stem cylindrical, silky, whitish.
500	...	...	...	V.	...	...	B.	Woods ...	...	Fleshy, thin, bell shaped, fibrous or scaly. Stem hollow, whitish. <i>Edible</i> .
501	...	...	...	...	...	Q.	...	Roots of bamboos	...	White, smooth, like chamois leather. Stem loosely stuffed, tapering at base.
Scan. II. 295 (1849).— <i>Agaricus</i> .										
502	...	S.A.	...	V.	...	...	B.	By waysides	...	Fleshy, viscid, ochrey; margin whitish, fluffy. Stem white, stuffed.
503	...	...	...	V.	...	...	B.	Among grass	...	Moist, somewhat cinnamon colour, dry ochrey. Stem hollow, short.
504	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Dung ...	...	Small. Somewhat fleshy, hemispherical, mottled yellowish. Very common.
505	...	S.A.	...	V.	N.S.W.	...	B.	Woods ...	...	Fleshy, thin, somewhat viscid, yellowish tawny, sprinkled with superficial concentric scales.
S.M. I. 287 (1821).— <i>Agaricus</i> .										
506	...	...	...	...	...	Q.	...	Ground ...	...	Fleshy, dark brown, variegated with darker scales, yellowish within.
507	...	...	...	V.	N.S.W.	...	B.	Dead stumps	...	Somewhat fleshy, ochrey and whitish. Stem hollow, fragile, white.
508	...	...	...	V.	...	...	...	Ground ...	...	Bell shaped, tawny yellow. Stem slender, faintly streaked.
509	W.A.	S.A.	T.	V.	...	...	B.	Stumps and ground	...	Somewhat fleshy, tawny honey colour, margin silky. Stem thin.
510	...	S.A.	T.	V.	...	...	B.	Old stumps, &c.	...	Fleshy, yellowish, with greenish tinge. Stem hollow, flesh yellow.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
48. PSILOCYBE.—Fries,					
511	324	IX. 568	<i>P. ceres</i> ...	Cooke and Mass., Grev. XVI. 72 (1888) ...	Ceres psilocybe ...
512	322	V. 4269	<i>P. cernua</i> ...	Fries, S.M. I. 298 (1821) ...	Nodding psilocybe ...
513	320	„ 4259	<i>P. compta</i> ...	Fries, Hym. Eur. 301 (1874) ...	Ornamented psilocybe ...
514	319	„ 4235	<i>P. ericæa</i> ...	Fries, S.M. I. 291 (1821) ...	Heath-growing psilocybe ...
515	323	„ 4275	<i>P. fœnisecii</i> ...	Fries, S.M. I. 295 (1821) ...	Lawn psilocybe ...
516	321	„ 4267	<i>P. spadicea</i> ...	Fries, Epier. 225 (1838) ...	Date-brown psilocybe ...
49.—DECONICA.—Smith,					
517	325	V. 4293	<i>D. atro-rufa</i> ...	Sacc. Syll. V. 1059 (1887) ...	Dark-red deconica ...
518	326	„ 4294	<i>D. nunciseda</i> ...	Sacc. Syll. V. 1059 (1887) ...	Nutty deconica ...
50. PSATHYRA.—Fries.					
519	327	V. 4297	<i>P. conopileæ</i> ...	Fries, S.M. I. 504 (1821) ...	Cone-capped psathyra ...
520	330	„ 4344	<i>P. fatua</i> ...	Fries, S.M. I. 296 (1821) ...	Tasteless psathyra ...
521	331	„ 4349	<i>P. gossypina</i> ...	Fries, S.M. I. 310 (1821) ...	Cottony psathyra ...
522	329	„ 4320	<i>P. obtusata</i> ...	Fries, S.M. I. 293 (1821) ...	Obtuse psathyra ...
523	328	„ 4314	<i>P. Sonderiana</i> ...	Berk., Linn. Journ. XIII. 159 (1873) ...	Sonder's psathyra ...
51. BOLBITIUS.—Fries,					
524	...	...	<i>B. candidus</i> ...	Cooke and Mass., Grev. XXI. 37 (1892) ...	White bolbitius ...
525	358	V. 4357	<i>B. conocephalus</i> ...	Fries, Epier. 205 (1838) ...	Cone-headed bolbitius ...
526	356	„ 4355	<i>B. fragilis</i> ...	Fries, Epier. 254 (1838) ...	Fragile bolbitius ...
527	357	„ 4358	<i>B. titubans</i> ...	Fries, Epier. 254 (1838) ...	Tottering bolbitius ...
52. COPRINUS.—Pers.					
528	345	V. 4374	<i>C. comatus</i> ...	Fries, Epier. 242 (1838) ...	Maned coprin ...
529	351	„ 4429	<i>C. deliquescens</i> ...	Fries, Epier. 249 (1838) ...	Deliquescent coprin ...
530	354	„ 4480	<i>C. ephemerus</i> ...	Fries, Epier. 252 (1838) ...	Ephemeral coprin ...
531	347	„ 4404	<i>C. fimetarius</i> ...	Fries, Epier. 245 (1838) ...	Dung coprin ...
531A	347	„ „	<i>C. fimetarius, var. macrorrhizus</i> ...	(Fries), Hym. Eur. 324 (1874) ...	Large-rooting coprin ...
532	349	„ 4416	<i>C. micaceus</i> ...	Fries, Epier. 247 (1838) ...	Glistening coprin ...
533	353	„ 4477	<i>C. murinus</i> ...	Kalch., Grev. VIII. 152 (1880) ...	Mouse-coloured coprin ...
534	348bis.	„ 4407	<i>C. niveus</i> ...	Fries, Epier. 246 (1838) ...	Snowy coprin ...
535	346	„ 4394	<i>C. picaceus</i> ...	Fries, Epier. 244 (1838) ...	Variegated coprin ...
536	355	„ 4490	<i>C. plicatilis</i> ...	Fries, Epier. 252 (1838) ...	Plaited coprin ...
537	352	„ 4465	<i>C. stercorarius</i> ...	Fries, Epier. 251 (1838) ...	Dung-borne coprin ...
538	348	„ 4406	<i>C. tomentosus</i> ...	Fries, Epier. 246 (1838) ...	Tomentose coprin ...
539	350	„ 4420	<i>C. truncorum</i> ...	Fries, Epier. 248 (1838) ...	Trunk coprin ...
53. PANÆOLUS.—Fries,					
540	338	V. 4544	<i>P. campanulatus</i> ...	Fries, Hym. Eur. 311 (1874) ...	Bell panæolus ...
541	333	IX. 598	<i>P. eburneus</i> ...	Cooke and Mass., Grev. XVIII. 4 (1889) ...	Ivory-white panæolus ...
542	340	V. 4555	<i>P. fimicola</i> ...	Fries, Hym. Eur. 312 (1874) ...	Dung-borne panæolus ...
543	332	IX. 596	<i>P. ovatus</i> ...	Cooke and Mass., Grev. XVIII. 4 (1889) ...	Ovate panæolus ...
544	339	V. 4547	<i>P. papilionaceus</i> ...	Fries, Epier. 256 (1838) ...	Butterfly panæolus ...
545	335	„ 4536	<i>P. phalænarum</i> ...	Fries, Epier. 235 (1838) ...	Moth panæolus ...
546	336	„ 4539	<i>P. retrigis</i> ...	Fries, Epier. 235 (1838) ...	Wrinkled panæolus ...
547	337	IX. 597	<i>P. veluticeps</i> ...	Cooke and Mass., Grev. XVIII. 4 (1889) ...	Velvet-capped panæolus ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
S.M. I. 289 (1821).—Agaricus.									
511	...	...	...	V.	...	...	...	Ground ...	Thin, brick red. Stem elongated, ochrey, downy downwards.
512	...	...	...	V.	...	...	B.	Chips, decayed wood, &c.	Somewhat fleshy, wrinkled when dry, white. Stem hollow, white.
513	...	...	...	V.	...	...	B.	Woods ...	Pale ochrey, grooved, with scattered shining spots. Stem shining, silky.
514	W.A.	...	...	V.	...	...	B.	Exposed pastures after rain	Fleshy, rather viscid when moist, shining when dry, ferruginous tawny.
515	...	S.A.	...	...	...	...	B.	Among grass, lawn	Somewhat fleshy, dark brown, hemispherical or bell shaped. Stem pale red.
516	...	...	T.	V.	...	...	B.	Dead stumps, ground, &c., in woods	Fragile, rigid. Fleshy, bay to umber, moist. Stem hollow, tough, pale.
Seem. Journ. (1876).—Agaricus, Psilocybe.									
517	W.A.	...	...	...	...	...	...	Ground in woods ...	Dark red or purple brown. Rather fleshy, discoloured when dry.
518	...	...	...	...	...	Q.	...	Chips ...	Rather fleshy, yellowish, silky when dry.
S.M. I. 11 (1821).—Agaricus.									
519	...	...	...	V.	...	...	B.	Ground ...	Large, graceful. Sub-membranaceous, growing pale. Stem tall. Common.
520	...	...	...	...	...	...	B.	Gardens, &c. ...	Very fragile. Sub-membranaceous, clay coloured, rugged. Stem smooth.
521	...	...	...	...	...	Q.	B.	Ground ...	Ochrey to clayey. Sub-membranaceous, downy, becoming smooth.
522	W.A.	...	...	V.	...	...	B.	Oak trunks and ground	Sub-membranaceous, wrinkled, rather shining, obtuse. Stem rigid.
523	...	S.A.	...	...	...	...	...	Ground ...	Pale and dirty yellowish, acutely convex. Stem white, silky.
Epicr. 253 (1838).—Agaricus.									
524	...	...	...	V.	...	...	...	Stable refuse ...	Membranaceous, white, bell shaped. Stem long, hollow.
525	...	...	...	V.	N.S.W.	...	...	Moist ground ...	Fragile, graceful, from livid to clay white. Membranaceous, conical, rather viscid.
526	W.A.	...	...	V.	...	Q.	B.	Dung ...	Small, but rather tall. Sub-membranaceous, viscid, pellucid, yellow, becoming pale.
527	...	...	...	V.	...	...	B.	Among grass ...	Small, tall, very fragile, trembling and tottering. Membranaceous, yellow.
Tent. disp. 62 (1797).—Agaricus.									
528	...	...	...	V.	...	...	B.	Sides of roads, pastures	Large and tall, white. Rather fleshy, cylindrical. Cuticle broken and feathery. <i>Edible</i> .
529	...	...	...	...	...	Q.	B.	Old stumps ...	Large. Sub-membranaceous, livid, top papillate. Stem hollow.
530	...	...	...	...	...	Q.	B.	Dung-hills ...	Small. Very thin, splitting, somewhat mealy. Stem slender.
531	...	...	...	...	...	Q.	B.	Dung-heaps ...	Sub-membranaceous, soon torn, disc livid. Stem scaly.
531A	...	S.A.	...	...	...	...	B.	Dung-heaps ...	Leathery to scaly. Stem rooting, shaggy.
532	...	S.A.	...	V.	...	...	B.	About old stumps ...	Small. Sub-membranaceous, brown, covered with glittering particles.
533	...	...	...	V.	N.S.W.	...	...	Ground ...	Small. Sub-membranaceous, with prominent papilla at apex, grey.
534	...	...	...	V.	...	...	B.	Horse-dung ...	Small. Sub-membranaceous, clad with dense white down.
535	...	...	...	...	...	Q.	B.	Road-sides, &c. ...	Sub-membranaceous, deep black, variegated, with broad white superficial scales.
536	...	S.A.	...	V.	N.S.W.	Q.	B.	Pastures ...	Small, delicate. Very thin, splitting, furrowed and folded, grey.
537	...	...	T.	V.	N.S.W.	Q.	B.	Rich soil and dung	Very thin, ovate, covered with a dense white micaceous meal.
538	...	...	...	...	...	Q.	B.	Dung and rich pastures	Sub-membranaceous, cylindrical to conical, woolly to downy, whitish grey.
539	...	...	...	...	...	Q.	...	Wood ...	Membranaceous, deliquescent, ferruginous ochrey, at first densely micaceous.
Epicr. 234 (1838).—Agaricus.									
540	W.A.	...	...	V.	N.S.W.	...	B.	Rich soil ...	Fragile. Somewhat fleshy, bell shaped, shining, dry, red brown.
541	...	...	...	...	...	Q.	...	Dung ...	Rather fleshy, ivory white, shining. Stem fragile, elongated.
542	...	...	...	V.	...	Q.	B.	Dung, rich pastures &c.	Somewhat fleshy, marked near margin with a narrow brown zone. Stem fragile, elongated.
543	...	...	...	V.	...	...	...	Mannre ...	Rather fleshy, ovate, at length cracked, white. Stem erect, silky.
544	...	S.A.	...	V.	...	...	B.	Dung, rich pastures, &c.	Somewhat fleshy, pale tan, conico-convex, when dry cracked and scaly.
545	...	...	...	V.	...	...	B.	Dung ...	Rather fleshy, viscid, dirty clay colour. Veil fleeting.
546	...	...	...	V.	...	...	B.	Dung ...	Somewhat fleshy, reticulated with raised ribs, flesh to tan colour.
547	...	S.A.	...	...	...	Q.	...	In garden among grass	Velvety, grey, convex or bell shaped. Stem elongated, hollow, silvery grey.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
54. ANELLARIA.—Karst.					
48	334	V. 4561	<i>A. fimiputris</i> ...	Karst. Hattsv. I. 518 (1879, &c.) ...	Putrid dung anellaria ...
55. PSATHYRELLA.—Fries,					
49	343	V. 4595	<i>P. crenata</i> ...	Laseh., in Fries Hym. Eur. 315 (1874) ...	Crenate psathyrella ...
50	344	" 4597	<i>P. disseminata</i> ...	Fries, Hym. Eur. 316 (1836) ...	Scattered psathyrella ...
51	341	" 4572	<i>P. hiaseens</i> ...	Fries, Hym. Eur. 314 (1874) ...	Gaping psathyrella ...
52	342	" 4575	<i>P. trepida</i> ...	Fries, Epier. 238 (1838) ...	Trembling psathyrella ...
ADDITIONS TO					
53	...	...	<i>Amanita Forrestiae</i> ...	Kalch., Linn. Soc. N.S.W. VII. 638 (1882) ...	Forrest's amanita ...
54	...	V. 92	<i>Lepiota acute-squamosa</i> ...	Weinum., Syll. I. 70 (1836) ...	Acute-scaly lepiota ...
55	44	...	<i>L. megalotholes</i> ...	Kalch., Linn. Soc. N.S.W. VII. 563 (1882) ...	Large-nippled lepiota ...
56	...	...	<i>Tricholoma carneo-flavidum</i> ...	Kalch., Linn. Soc. N.S.W. VII. 639 (1882) ...	Fleshy-yellow tricholome ...
57	...	V. 474	<i>T. panæolum</i> ...	Fries, Epier. 49 (1838) ...	Variegated tricholome ...
58	...	...	<i>T. plagiolum</i> ...	Kalch., Linn. Soc. N.S.W. VII. 639 (1882) ...	Oblique tricholome ...
59	...	V. 496	<i>T. sordidum</i> ...	Fries, S.M. I. 51 (1821) ...	Sordid tricholome ...
60	...	...	<i>T. turbinipes</i> ...	Kalch., Linn. Soc. N.S.W. VII. 639 (1882) ...	Turbinate-stalked tricholome ...
61	...	V. 630	<i>Clitocybe catina</i> ...	Fries, Epier. 72 (1838) ...	Bowl-shaped clitocybe ...
62	...	" 673	<i>C. ditopoda</i> ...	Fries, S.M. I. 171 (1821) ...	Variable-stalked clitocybe ...
63	...	" 529	<i>C. ochro-purpurea</i> ...	Berk., Hook. Lond. Journ. Bot. IV. 299 (1845) ...	Ochrey-purple clitocybe ...
64	...	...	<i>Collybia museipula</i> ...	Cooke and Mass., Grev. XXII. 26 (1893) ...	Monsey collybia ...
65	...	V. 1109	<i>Mycena epipterygia</i> ...	Fries, S.M. I. 155 (1821) ...	Winged mycena ...
66	...	" 962	<i>M. luteo-alba</i> ...	Fries, S.M. (1821) ...	Yellowish-white mycena ...
67	...	" 1005	<i>M. polygramma</i> ...	Fries, S.M. I. 146 (1821) ...	Many-lined mycena ...
68	...	" 1137	<i>M. pterigena</i> ...	Fries, S.M. I. 160 (1821) ...	Pteris-borne mycena ...
69	...	" 992	<i>M. ræborrhiza</i> ...	Laseh., Linn. 539 (1829) ...	Crooked-root mycena ...
70	...	" 1407	<i>Pleurotus acerinus</i> ...	Fries, Epier. 134 (1838) ...	Maple pleurote ...
71	...	" 1502	<i>P. cyphellæformis</i> ...	Berk., Mag. Zool. and Bot. 511 (1837) ...	Cyphella-like pleurote ...
72	...	" 1339	<i>P. lignatilis</i> ...	Fries, Epier. 132 (1838) ...	Wood-growing pleurote ...
73	...	" 1561	<i>Hygrophorus discoideus</i> ...	Fries, Epier. 323 (1838) ...	Discoid hygrophore ...
74	...	" 1651	<i>H. puniceus</i> ...	Fries, Mon. Hym. II. 21 (1857) ...	Purple hygrophore ...
75	...	" 1738	<i>Lactarius quietus</i> ...	Fries, Epier. 343 (1838) ...	Mild lactar ...
76	...	" 1821	<i>Russula xerampelina</i> ...	Fries, Epier. 356 (1838) ...	Dark-red russule ...
77	...	" 1989	<i>Marasmius badius</i> ...	Berk. and Curt., Linn. Journ. X. 294 (1869) ...	Bay-brown marasmius ...
78	...	" 1996	<i>M. pellucidus</i> ...	Berk. and Br., Linn. Journ. XIV. 35 (1875) ...	Pellucid marasmius ...
79	...	" 1991	<i>M. rhyssophyllus</i> ...	Mont. in Berk and Curt., Linn. Journ. X. 294 (1869) ...	Wrinkle-gilled marasmius ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Hattsv. I. 25 (1879).— <i>Agaricus</i> , <i>Panaeolus</i> .									
548	...	...	...	V.	...	...	B.	Dung, &c.	Sub-membranaceous, viscid, dark grey. Stem slender, equal.
Epier. 237 (1838) — <i>Agaricus</i> .									
549	...	S.A.	...	...	...	...	B.	Grassy ground	Fragile. Membranaceous, ochrey to pale red, margin notched.
550	W.A.	...	T.	V.	...	Q.	B.	About trunks of trees and on ground	Densely clustered. Membranaceous, pearly white, scurfy. Stem lax, fragile.
551	...	...	...	...	...	Q.	B.	Ground	Membranaceous, fissured and furrowed, becoming yellow. Stem brittle
552	...	S.A.	...	V.	...	...	B.	Ground	Very fragile. Membranaceous, sooty, bell shaped, densely streaked. Stem nearly straight, transparent.
AGARICACEÆ.									
553	W.A.	...	...	...	...	...	...	...	Fleshy, convex to plane, smooth, naked, white to ashy grey or brown. Stem stout, stuffed, white, fibrous, not bulbous.
554	...	...	...	V.	...	...	B.	Grassy places	Fleshy, at first woolly hairy, then scaly, acute, dark tan. Stem somewhat stuffed, bulbous.
555	...	...	...	...	...	Q.	...	River-side	Fleshy, bell shaped, smooth, with brown adpressed scales. Stem somewhat hollow, dilated at base, naked, from white to brown.
556	W.A.	...	...	...	...	...	...	...	Fleshy, hemispherical, woolly scaly, fleshy yellow as if peach coloured. Stem solid, thick, of same colour
557	W.A.	...	...	...	...	...	B.	Grassy places	Small, spongy to compact, convex to plane, variegated with grey frosted spots. Stem solid, fibrous to striate.
558	W.A.	...	...	...	...	...	...	...	Fleshy, plane, depressed, smooth, pale reddish brown. Stem excentric, stuffed, cylindrical, base slightly thickened.
559	...	...	...	V.	...	...	B.	Meadows, dung-heaps, &c.	Somewhat fleshy, from bell shaped and convex to plane and depressed, finally squalid. Stem stuffed, base thickened.
560	W.A.	...	...	...	...	...	...	...	Large, pale reddish. Fleshy, irregularly convex. Stem top shaped, thick.
561	...	...	...	V.	...	...	B.	Among dead leaves	White, discoloured, odour pleasant. Fleshy, plane, then funnel shaped, dry. Stem stuffed, thick, elastic, white.
562	...	...	...	V.	...	...	B.	Woods	Strong smelling, like new meal. Rather fleshy, dingy, brownish grey, smooth. Stem hollow, almost smooth, of same colour.
563	...	...	...	...	...	...	...	Clayey soil in woods	Somewhat hemispherical, at length depressed, fleshy, compact, pale tan, becoming slightly purple. Stem thick, swollen in middle.
564	...	...	...	...	...	Q.	...	Ground	Fleshy, smooth, mouse grey or brown, wrinkled. Stem stuffed, tapering downwards and rooting, striate lengthwise.
565	...	...	...	...	...	Q.	B.	Among moss and leaves	Membranous, bell shaped, striate, very viscid and easily separable, usually grey. Stem elongated, tough, rooting, yellowish.
566	...	...	...	V.	...	...	B.	Among moss, &c.	Membranous, bell shaped, slightly grooved, pale, yellow. Stem thread-like, shining, smooth, becoming yellow.
567	...	...	...	V.	...	...	B.	Trunks	Rather membranous, conical to bell shaped, dry, grooved. Stem rigid, longitudinally furrowed and grooved, shining, rooting.
568	...	...	...	V.	...	...	B.	Dead fern stems, veins of leaves, &c.	Very elegant, delicate, rosy. Bell shaped, obtuse, and stem wavy, very thin, with disc at base.
569	...	...	...	V.	...	...	...	About trunks	Somewhat membranous, acutely bell shaped, dry, rather tawny or pale. Stem firm, thick, rooting.
570	...	...	...	...	...	Q.	B.	Trunks	White, firm. Fleshy, thin, unequal, silky hairy. Stem almost lateral, slender or nearly obsolete, downy.
571	...	...	...	...	...	Q.	B.	Moss and dead stems of herbaceous plants	Gregarious, small. Rather fleshy, sessile, cup shaped, grey, margin paler, delicately downy.
572	...	...	...	V.	...	...	B.	Trunks, rotten wood, &c.	Odour mealy. Fleshy, tough, convex to plane, dingy white. Stem stuffed, then hollow, slender base, rooting and downy.
573	...	...	...	V.	...	...	B.	Grassy places	Gregarious. Fleshy, smooth, very glutinous, yellowish tan, disc somewhat rusty. Stem stuffed, soft, viscid, pale white.
574	...	...	...	V.	...	...	B.	Mossy meadows, &c.	Very large, very showy, fragile. Bell shaped, viscid, scarlet to blood red. Stem hollow, thick, bulging, base white.
575	...	...	...	...	...	Q.	B.	Woods, &c.	Fleshy, viscid at first, slightly cinnamon, then dry and slightly silky, somewhat zoned. Stem spongy to stuffed, finally rusty red.
576	...	...	...	V.	...	...	B.	Woods	Mild. Fleshy, compact, dry, opaque, rose purple. Stem strong, firm, finally spongy to soft.
577	...	...	...	...	...	Q.	...	Wood	Reddish when fresh. Convex, striate, smooth, margin incurved. Stem frosted, becoming smooth.
578	...	...	...	...	...	Q.	...	Dead branches, &c.	Convex, pellucid. Stem equal, brown, delicately powdery.
579	...	...	...	...	...	Q.	...	Wood	Fibrous, smooth, pale yellow. Stem same colour, smooth, with rough-haired base.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name	Authority for Name.	English Name.
ADDITIONS TO					
580	...	V. 2379	<i>Lentinus descendens</i> ...	Fries, <i>Epier.</i> 290 (1838) ...	Descending lentinus ...
581	...	" 2924	<i>Leptonia æthiops</i> ...	Fries, <i>Epier.</i> 152 (1838) ...	Æthiopian leptonia ...
582	...	" 3406	<i>Nolanea sub-globosa</i> ...	Cooke, <i>Grev.</i> XVII. 38 (1888) ...	Sub-globose nolanea ...
583	...	" 3107	<i>Pholiota adiposa</i> ...	Fries, <i>S.M.</i> I. 242 (1821) ...	Glutinous pholiota ...
584	...	...	<i>P. bicincta</i> ...	Kalch., <i>Linn. Soc. N.S.W.</i> VII. 639 (1882) ...	Twice-girt pholiota ...
585	...	V. 3064	<i>P. radicata</i> ...	Fries, <i>S.M.</i> I. 242 (1821) ...	Rooting pholiota ...
586	...	...	<i>Hebeloma Kirtoni</i> ...	Kalch., <i>Linn. Soc. N.S.W.</i> VII. 564 (1882) ...	Kirton's hebeloma ...
587	...	V. 3359	<i>Flammula carbonaria</i> ...	Fries, <i>S.M.</i> I. 252 (1821) ...	Charcoal-loving flammula ...
588	...	" 3322	<i>F. gymnopodia</i> ...	Fries, <i>Hym. Eur.</i> 218 (1874) ...	Naked-stalked flammula ...
440A	...	" 3385	<i>F. sapinea</i> , var. <i>terrestris</i> ...	Fries, <i>S.M.</i> I. 239 (1821) ...	Terrestrial flammula ...
589	...	" 3433	<i>Naucoria abstrusa</i> ...	Fries, <i>Epier.</i> 194 (1838) ...	Concealed naucoria ...
590	...	" 3509	<i>N. conspersa</i> ...	Fries, <i>S.M.</i> I. 260 (1821) ...	Besprinkled naucoria ...
591	...	" 3499	<i>N. sobria</i> ...	Fries, <i>Epier.</i> 200 (1838) ...	Sober naucoria ...
592	...	" 3481	<i>N. tenax</i> ...	Fries, <i>Epier.</i> 198 (1838) ...	Firm naucoria ...
593	...	" 3605	<i>Crepidotus applanatus</i> ...	Fries, <i>Mon. Hym.</i> I. 399 (1857) ...	Depressed crepidotus ...
594	...	...	<i>Cortinarius Walkeri</i> ...	Cooke and Mass., <i>Grev.</i> XXII. 36 (1893) ...	Walker's cortinarius ...
595	...	IX. 553	<i>Paxillus hirtulus</i> ...	F.v.M., <i>Linn. Soc. N.S.W.</i> VIII. 175 (1883) ...	Hairy paxillus ...
596	...	...	<i>Hypoloma peltastes</i> ...	Kalch., <i>Linn. Soc. N.S.W.</i> VII. 564 (1882) ...	Shield-like hypoloma ...
597	...	V. 4261	<i>Psilocybe semilanceata</i> ...	Fries, <i>Obs.</i> II. 178 (1818) ...	Semi-pointed psilocybe ...
598	...	" 4560	<i>Anellaria separata</i> ...	Karst., <i>Hattsv.</i> I. 517 (1879) ...	Separate anellaria ...
599	...	" 4566	<i>Psathyrella impatiens</i> ...	Fries, <i>S.M.</i> I. 302 (1821) ...	Impatient psathyrella ...
600	...	V. 1960	<i>A. cupularis</i> ...	Fries, <i>S.V.</i> S. 312 (1849) ...	Cupular arrhenia ...

56 ARRHENIA.—Fries,

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
AGARICACEÆ—continued.									
580	...	...	...	...	...	Q.	...	Ground ...	Somewhat woody, funnel shaped, tan coloured, spotted with minute scales. Stem solid, very hard rooting.
581	...	...	...	V.	...	...	B.	Among grass, &c. ...	Fleshy, depressed, smooth, shining, sooty black. Stem slender, smooth, blackish brown, with black points towards top.
582	...	...	...	V.	...	...	B.	Ground ...	Fleshy, sub-globose, viscid, yellow. Stem thin, becoming hollow, longitudinally striate.
583	...	...	...	...	...	Q.	B.	Trunks ...	Tufted, very showy and large. Fleshy, glutinous, yellow, scaly. Stem stuffed, glutinous, base somewhat bulbous.
584	W.A.	...	...	...	...	...	...	...	Convex, obtuse, light umber, scaly. Stem solid, bulbous, fibrillose, about middle and above base doubly girt.
585	...	...	...	V.	...	...	B.	Woods, near to stumps	Large, beautiful, sweet odour. Fleshy, dry, smooth, spotted, almost clay colour. Stem solid, passing into tapering root.
586	...	...	...	...	N.S.W.	...	...	...	Compact, fleshy, convex to plane, smooth. Stem solid, fleshy, fibrillose, white to silky.
587	...	...	...	V.	...	...	B.	Burnt ground, charcoal, &c.	Gregarious. Fleshy, firm, viscid, tawny yellow, often depressed in centre. Stem rigid, scaly, pale, base usually darker.
588	...	...	...	...	...	Q.	B.	Ground ...	Often tufted, rusty brown. Fleshy, bell shaped to convex, scaly. Stem solid, almost smooth.
440A	...	...	...	...	...	Q.	B.	Ground ...	Tufted, stem elongated, rooting in a spindle-shaped manner.
589	W.A.	...	...	...	...	...	B.	Damp earth, leaf-soil, &c.	Rather fleshy, smooth, viscid, rusty tan. Stem cartilaginous, tough, hollow, polished, rusty.
590	...	...	...	V.	...	...	B.	Ground, among leaves, &c.	Gregarious, fragile. Fleshy, with scurfy scales, bay cinnamon when moist, ochrey when dry. Stem fibrillose, brownish cinnamon.
591	...	S.A.	...	...	...	...	B.	In moist woods or scrubs	Fleshy, slightly viscid, somewhat silky, honey colour when moist, not absorbent of moisture, hence the name. Stem thick, hollow, rusty brown below.
592	...	...	...	V.	...	...	B.	Woods, among grass	Somewhat fleshy, bell shaped then expanded, smooth, slightly viscid, cinnamon when moist, ochrey when dry. Stem stuffed then hollow, dusky yellow.
593	...	...	...	V.	...	...	B.	Rotten wood	Fleshy, soft, fragile, kidney to wedge shaped, whitish, at length depressed behind. Stem very short, whitish downy.
594	...	...	...	...	N.S.W.	...	...	Ground ...	Convex, then expanded, minutely silky, pale green then bluish green. Stem slightly thickened at base, stuffed, reddish.
595	...	...	...	...	...	Q.	...	Ground ...	Convex to depressed, becoming darkly lurid. Stem thickened downwards, base abruptly rooting, hairy.
596	...	...	...	...	N.S.W.	...	...	...	Fleshy, viscid, scutiform, becoming brown. Stem solid, thickened downwards, naked, white.
597	...	...	...	V.	...	...	B.	Among grass	Gregarious. Somewhat membranous, acutely conical, almost pointed, slightly viscid, pale yellow when dry. Stem tough, wavy, silky fibrous, shining.
598	...	...	...	...	...	...	B.	Dung ...	Fleshy, bell shaped, viscid, ochrey, then whitish and wrinkled when old, shining. Stem long, straight, shining, whitish, tapering upwards with persistent ring.
599	...	...	...	V.	...	...	...	Moist woods	Membranous, bell shaped, convex, smooth. Stem weak, smooth, white.

Summ. Veg. Scand. 312 (1849).—*Cantharellus*, *Merulius*.

600	...	...	...	...	...	Q.	...	Young pinnate leaf	Small, resupinate, soft, circular, shaggy, grey.
-----	-----	-----	-----	-----	-----	----	-----	--------------------	--

ORDER II.—POLYPORACEÆ, FRIES.

57. *Boletus*, Linn.
 58. *Strobilomyces*, Berk.
 59. *Fistulina*, Bull.
 60. *Polyporus*, Linn.

61. *Fomes*, Fries.
 62. *Polystictus*, Fries.
 63. *Poria*, Pers.
 64. *Trametes*, Fries.

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER II.—POLYPORACEÆ					
57. BOLETUS.—Linn., Sp. Pl. 1176 (1753).					
601	561	VI. 4749	<i>B. æreus</i>	Bull., Champ. 321 (1798)	Bronze bolet
602	562	„ 4755	<i>B. æstivalis</i>	Fries, Epier. 422 (1838)	Summer bolet
603	551	„ 4673	<i>B. alliciens</i>	Berk., Hook., Lond. Journ. IV. 50 (1845)	Attractive bolet
604	550	„ 4671	<i>B. arenarius</i>	Fries, Pl. Preiss. II. 134 (1846)... ..	Sand-loving bolet
605	552	„ 4674	<i>B. australis</i>	Cooke and Mass., Grev. XVI. 32 (1887)	Southern bolet
606	547	„ 4653	<i>B. badius</i>	Fries, Elench. 126 (1828)	Bay-brown bolet
607	557	...	<i>B. brunneus</i>	Cooke and Mass., Grev. XIX. 90 (1891)... ..	Brown bolet
608	565	VI. 4761	<i>B. caesareus</i>	Fries, Pl. Preiss. II. 134 (1846)... ..	Imperial bolet
609	558	„ 4726	<i>B. calopus</i>	Fries, S.M. I. 390 (1821)	Red-stalked bolet
610	553	„ 4680	<i>B. chrysenteron</i>	Fries, Epier. 415 (1838)	Red-crack bolet
611	560	„ 4748	<i>B. edulis</i>	Bull., Champ. 60 (1798)	Edible bolet
612	544	„ 4642	<i>B. elegans</i>	Schum., Saell. II. 374 (1801)	Elegant bolet
613	569	„ 4801	<i>B. felleus</i>	Bull., Champ. 379 (1798)	Bitter bolet
614	545	„ 4643	<i>B. flavus</i>	Wither., Fries, Epier. 410 (1838)	Yellow bolet
615	555	„ 4833 & IX. 633	<i>B. fruticicola</i>	Berk., Hook., Lond. Journ. VII. 574 (1848)	Shrub-growing bolet
616	546	VI. 4648	<i>B. granulatus</i>	Linn., Sp. Pl. 1177 (1753)	Granulated bolet
617	572	„ 4817	<i>B. badius</i>	Berk. and Br., Linn., Trans. II. 57 (1883)	Kid-like bolet
618	564	„ 4760	<i>B. infraetus</i>	Fries, Pl. Preiss. II. 134 (1846)	Fractured bolet
619	571	IX. 641	<i>B. lacunosus</i>	Cooke and Mass., Grev. XVIII. 5 (1889)	Pitted bolet
620	566	VI. 4768	<i>B. luridus</i>	Schaeff., Fung. 107 (1762)	Lurid bolet
621	543	„ 4641	<i>B. luteus</i>	Linn., Sp. Pl. 1177 (1753)	Yellow bolet
622	567	„ 4798	<i>B. marginatus</i>	Drum., Berk., Hook., Lond. Journ., Bot. IV. 50 (1845)	Margined bolet
623	570	„ 4803	<i>B. megalosporus</i>	Berk., Fl. Tasm. II. 251 (1860)	Large-spored bolet
624	573	„ 4830	<i>B. napipes</i>	F. v. M., Linn. Journ. XIII. 161 (1873)	Turnip-stalked bolet
625	559	„ 4728	<i>B. pachypus</i>	Fries, S.M. I. 390 (1821)	Thick-stalked bolet
626	563	„ 4756	<i>B. portentosus</i>	Berk. and Br., Linn. Journ. XIV. 46 (1875)	Monstrous bolet
627	568	„ 4800	<i>B. prunicolor</i>	Cooke and Mass., Grev. XVI. 32 (1887)	Plum-coloured bolet
628	548	„ 4656	<i>B. sanguineus</i>	With., Arr. IV. 319 (1796)	Blood-red bolet
629	...	„ 4792	<i>B. scaber</i>	Fries, S.M. I. 203 (1821)	Rough bolet

ARRANGEMENT OF GENERA (16).

65. Sclerodepsis, Cooke.
66. Hexagonia, Fries.
67. Dædalca, Pers.
68. Ceriomyces, Corda.

69. Favolus, Fries.
70. Laschia, Fries.
71. Campbellia, Cooke.
72. Merulius, Hall.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
601	...	...	...	...	...	Q.	B.	Woods ...	Cap smooth, olive brown, turning blackish. Stem stout and yellow. Pores sulphur yellow. Rare. <i>Edible</i> .
602	...	...	...	...	...	Q.	B.	Woods, and in pastures under trees	Largest of this genus. Cap smooth, whitish. Stem very thick and yellowish. Pores greenish yellow. <i>Edible</i> .
603	W.A.	...	...	...	...	...	...	Ground ...	Cap smooth, yellow, viscid. Stem downy. Pores yellow. <i>Edible</i> .
604	W.A.	...	...	...	...	...	...	Sandy soil ...	Cap flattened, viscid, excentric. Stem elongated, pale above, black below. Pores cinnamon.
605	...	S.A.	...	V.	...	...	...	Ground ...	Cap viscid, umber. Stem flesh colour. Pores hexagonal, sulphur colour.
606	...	...	...	...	...	Q.	B.	Woods ...	Cap soft, viscid, bay brown. Stem solid, with brownish bloom. Pores from yellowish white to greenish. <i>Edible</i> .
607	...	...	...	V.	...	...	...	Ground ...	Cap somewhat downy, reddish brown. Stem short, stout. Pores rather large, greenish grey.
608	W.A.	...	...	...	...	...	...	Sandy soil ...	Cap fleshy, blood red to purple. Stem stout, sulphur colour. Pores rounded, yellow, with oblique openings.
609	...	...	...	...	N.S.W.	...	B.	Woods ...	Cap globose, somewhat downy, olive. Stem firm and thick, scarlet. Pores yellow.
610	...	...	...	V.	...	Q.	B.	Woods, &c. ...	Cap, dull brown with red cracks. Stem rigid, crimson or yellow. Pores greenish yellow.
611	...	...	...	...	...	Q.	B.	Woods ...	Cap smooth, moist, brownish. Stem stout, pale brown. Pores lemon to yellowish-green. <i>Edible</i> .
612	...	...	...	...	...	Q.	B.	Woods ...	Golden yellow entirely. Cap viscid. Stem firm. Pores sulphur colour. <i>Edible</i> .
613	...	...	...	V.	...	Q.	B.	Woods ...	Cap soft, smooth, brown or reddish grey. Stem solid, stout. Pores angular, flesh pink.
614	...	...	...	V.	...	...	B.	Woods ...	Large, entirely yellow. Cap compact, viscid. Stem spotted with brown and with fugacious rings. Pores angular, yellow.
615	...	...	T.	...	...	...	...	Ground at roots of <i>Pleurocladus riparius</i>	Cap fleshy, smooth, red. Stem nearly smooth. Pores pale orange yellow.
616	...	...	...	V.	...	Q.	B.	Grassy places ...	Cap slimy. Stem covered with milky drops drying into brown granules. Pores granulated. <i>Edible</i> .
617	...	...	...	...	...	Q.	...	Ground ...	Cap thick, tan coloured. Stem similarly coloured. Pores pale.
618	W.A.	...	...	...	...	...	...	Ground ...	Cap smooth, purple, with margin much broken. Stem very short and tuberous. Pores sulphur colour.
619	...	...	...	...	...	Q.	...	Sandy ground ...	Cap soft, somewhat viscid, tawny. Stem deeply pitted. Pores angular whitish to flesh colour.
620	...	...	...	V.	...	Q.	B.	Ground ...	Large. Cap viscid, soft, olive brown or tawny. Stem stout, tall, vermillion. Pores orange, red, crimson. Common.
621	W.A.	...	...	V.	...	Q.	B.	Ground in Pine woods	Large. Cap viscid, soft, dingy yellow. Stem tall, firm, with broad dingy ring. Pores yellow. Common. <i>Edible</i> .
622	W.A.	...	...	V.	...	...	...	Ground ...	Cap compact, delicately velvety, margin thin and distinct from hymenium. Stem short, black. Pores internally palid.
623	...	...	T.	...	...	...	...	Ground in woods ...	Cap somewhat tan coloured. Stem warty. Pores flesh colour.
624	...	...	...	V.	...	...	...	Meadows ...	Cap reddish brown, at length blackish. Stem obconical. Pores lemon yellow.
625	...	...	...	...	N.S.W.	Q.	B.	Woods ...	Very large. Cap brownish, then pale tan colour. Stem thick, firm, yellow variegated with red, very bulbous. Pores round, yellow. Common.
626	...	...	...	V.	...	...	...	Ground ...	Very large. Cap depressed in centre. Stem thick and dilated at base. Pores lemon yellow.
627	...	...	...	V.	...	...	...	Ground ...	Cap soft, viscid, plum coloured. Stem club shaped, pale. Pores rounded, pale.
628	...	...	...	...	...	Q.	B.	Woods ...	Very small. Cap smooth, viscid, blood red. Stem yellow and red. Pores large, orange yellow. Rare.
629	...	...	...	V.	...	...	B.	Woods ...	Large. Dull brown, very rough. Cap cushion shaped, viscid. Stem solid, tall, scurfy. Very common. <i>Edible</i> .

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
57. BOLETUS.—Linn., Sp. Pl. 1176 (1753).					
630	549	VI. 4670	<i>B. subsimilis</i> ...	Fries, Pl. Preiss. II. 134 (1846) ...	Simulating bolet ...
631	554	„ 4682	<i>B. subtomentosus</i> ...	Linn., Sp. Pl. 1178 (1753) ...	Downy bolet... ..
632	556	„ 4703	<i>B. Thozetii</i> ...	Berk., Linn. Journ. XVIII. 384 (1881) ...	Thozet's bolet
58.—STROBILOMYCES.—Berk. in Hook.,					
633	581	VI. 4838	<i>S. ananæceps</i> ...	Sacc., Syll. VI. 50 (1888) ...	Pine-apple-headed strobilomyces
634	577	...	<i>S. fasciculatus</i> ...	Cooke, Grev. XX. 4 (1891) ...	Fasciculate strobilomyces ...
635	579	VI. 4835	<i>S. floccopus</i> ...	Vahl., Fl. Dan. t. 1252 (1764) ...	Woolly-stalked strobilomyces ...
636	578	...	<i>S. ligulatus</i> ...	Cooke, Grev. XX. 4 (1891) ...	Ligulate strobilomyces ...
637	574	VI. 4837	<i>S. nigricans</i> ...	Berk., Hook. Journ. 139 (1852) ...	Blackening strobilomyces ...
638	575	IX. 645	<i>S. pallescens</i> ...	Cooke and Mass., Grev. XVIII. 5 (1889) ...	Bleaching strobilomyces ...
639	576	„ 646	<i>S. rufescens</i> ...	Cooke and Mass., Grev. XVIII. 5 (1889) ...	Reddish-brown strobilomyces ...
640	580	„ 644	<i>S. velutipes</i> ...	Cooke and Mass., Grev. XVIII. 5 (1889) ...	Velvety-stalked strobilomyces ...
59. FISTULINA.—Bull.					
641	582	VI. 4849	<i>F. hepatica</i> ...	Fries, S.M. I. 396 (1821) ...	Liver-coloured <i>Fistulina</i> . Beef-steak fungus
60. POLYPORUS.—Adans, Fam. II. 10 (1763).—					
642	646	VI. 5146	<i>P. adustus</i> ...	Fries, S.M. I. 363 (1821) ...	Scorched polypore
643	594	„ 4913	<i>P. alveolarius</i> ...	Bose, Berl. Mag. IV. (1811) ...	Depressed polypore
644	666	„ 5253	<i>P. anebus</i> ...	Berk., Hook., Lond. Journ. VI. 504 (1847) ...	Beardless polypore
645	632	„ 5080	<i>P. angustus</i> ...	Berk., Fl. Tasm. II. 253 (1860) ...	Narrow-capped polypore
646	622	„ 5043	<i>P. anthracophilus</i> ...	Cooke, Grev. XII. 16 (1884) ...	Burnt-ground polypore
647	591	„ 4903	<i>P. arcularius</i> ...	Fries, S.M. I. 342 (1821) ...	Convex polypore
648	636	„ 5107	<i>P. argentatus</i> ...	Cooke, Grev. XV. 20 (1886) ...	Silvery polypore
649	667	„ 5255	<i>P. ascoboloides</i> ...	Berk., Linn. Journ. XIII. 162 (1873) ...	Ascobolus-like polypore
650	654	„ 5207	<i>P. betulinus</i> ...	Fries, S.M. I. 358 (1821) ...	Birch polypore
651	599	„ 4944	<i>P. biennis</i> ...	Fries, Epier. 433 (1838) ...	Biennial polypore
652	...	„ 5166	<i>P. biretum</i> ...	Kalch., Hedw. XV. 114 (1876) ...	Tawny polypore
653	652	„ 5187	<i>P. borealis</i> ...	Fries, S.M. I. 366 (1821) ...	Northern polypore
654	589	„ 4885	<i>P. brumalis</i> ...	Fries, S.M. I. 348 (1821) ...	Wintry polypore
655	637	„ 5106	<i>P. campylus</i> ...	Berk., Fl. Tasm. II. 252 (1860) ...	Curved polypore
656	659	„ 5232	<i>P. cartilagineus</i> ...	Berk. and Br., Linn. Journ. XIV. 49 (1875) ...	Cartilaginous polypore
657	635	„ 5093	<i>P. chioneus</i> ...	Fries, S.M. I. 359 (1821) ...	Snowy polypore

OF AUSTRALIAN FUNGI—*continued.*

Number.	Habitat.						R.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
630	W.A.	...	...	...	...	Q.	...	Ground ...	Cap fleshy, viscid, shining brown. Stem solid, sulphur yellow. Pores very short, sulphur yellow. Simulating <i>B. lividus</i> .
631	W.A.	S.A.	...	V.	N.S.W.	Q.	B.	Woods, &c.	Cap soft, dry, finely tomentose, bronze, with yellow cracks. Stem tall, yellow variegated with red. Pores yellow.
632	...	...	...	...	...	Q.	...	Barren soil	Yellow. Cap with granular warts. Stem slender and flexuous. Pores free.

*Tubiporus, Agaricus—continued.*Kew Misc. III. 78 (1851).—*Boletus.*

633	...	...	...	V.	N.S.W.	Q.	...	Ground ...	Cap broken up into flat, thick, broad, sealy warts.
634	...	...	...	V.	...	...	...	Ground ...	Cap reddish brown, sealy, with fascicles of strap-like scales. Stem paler. Pores angular, yellowish.
635	...	...	...	...	...	Q.	...	Ground ...	Ash coloured, becoming black. Cap soft, sealy, and veil silky. Stem stout and downy. Pores large, greyish white. Rare.
636	...	...	...	V.	...	...	...	Ground ...	Cap hemispherical, brown, with darker ligulate scales. Stem paler. Pores angular, yellowish, or tinted with red.
637	...	...	...	...	...	Q.	...	Woods ...	Small. Cap rough in centre, with hexagonal warts. Stem solid, with woolly scales, like margin of cap.
638	...	...	...	...	...	Q.	...	Base of trees	Cap rosy purple, turning pale, with thick obtuse conical warts overlapping. Stem thick, reddish. Pores large, angular, yellowish.
639	...	...	...	...	...	Q.	...	Base of trees	Entirely reddish brown. Cap hemispherical, thickly covered with overlapping conical warts. Stem solid, pale above, reddish brown below. Pores large, angular, tawny.
640	...	...	...	...	...	Q.	...	Ground ...	Blackening. Cap with thick irregular warts overlapping. Stem velvety. Pores large, angular.

Champ. I. 314 (1798).—*Boletus.*

641	W.A.	...	...	V.	...	...	B.	Trunks of living trees	Fleshy and red juiced. Cap flesh colour to blood red and liver colour, roundish, attached by broad base, internally streaked. <i>Edible.</i>
-----	------	-----	-----	----	-----	-----	----	------------------------	--

Polystictus, Fomes, Favolus, Trametes, Poria, Boletus, Dædalea.

642	...	...	...	V.	N.S.W.	Q.	B.	Trunks, stumps, &c.	Cap fleshy, tough, pale ash colour, with margin blackening. Pores minute, white to grey, becoming black.
643	...	...	...	...	...	Q.	...	Trunks ...	Cap fleshy, leathery, depressed, brown. Stem firm and thickened at base. Pores hexagonal, white.
644	...	...	...	...	...	Q.	...	Wood ...	Pale fawn colour. Cap thin, leathery, delicately velvety. Pores small, round, short.
645	...	...	T.	...	...	...	...	Rotten wood	Cap narrow, downy, brown. Pores small, angularly punctiform.
646	W.A.	...	V.	...	...	Q.	...	Burnt ground	Imbricate and much divided, very leathery and hardening. Caps growing together, overlapping, bay brown. Pores angular, white.
647	...	...	...	V.	N.S.W.	Q.	...	Trunks ...	Cap tough and leathery, without zones, brown to yellowish. Stem short greyish brown. Pores oblong rhomboid, large, whitish.
648	...	...	...	V.	...	Q.	...	Trunks ...	Cap fleshy to leathery, shell shaped, slightly silky, white in front, ashy brown behind. Pores white, rounded.
649	...	...	...	...	...	...	...	Trunks ...	Cap circular, thickish, white, downy. Pores hexagonal, small.
650	...	...	...	...	...	Q.	B.	Birch, &c.	Cap fleshy, then corky, hoof shaped, smooth. Pores minute, short, white, or brownish.
651	...	...	...	...	...	Q.	...	Ground, near trunks	Cap spongy, then corky to leathery, white to rusty colour. Stem short, thick, rust coloured, woolly. Pores torn and toothed, dull white to brownish.
652	...	...	...	...	N.S.W.	...	...	Trunks ...	Hemispherical, sessile, spongy to powdery, tawny, cinnamon or bay brown.
653	...	...	...	V.	...	...	B.	Trunks ...	White to yellowish. Cap spongy to corky, hairy. Pores unequal, torn, white.
654	...	...	...	...	...	Q.	B.	Trunks ...	Cap tough, fleshy to leathery, sooty brown. Stem thin, hairy, sealy. Pores angular, toothed, white.
655	...	...	T.	V.	...	...	...	Rotten wood	Cap palmate, lobed, white, smooth. Hymenium concave. Pores small, irregular.
656	...	...	...	...	...	Q.	...	Dead wood	Cap red brown to sooty brown, cuticle cartilaginous. Pores minute.
657	...	...	...	V.	N.S.W.	Q.	B.	Trunks and stumps	White. Cap fleshy, soft, smooth. Pores short, minute, rounded. Smell rather acid.

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
60. POLYPORUS.—Adans, Fam. II. 10 (1763).—					
658	620	VI. 5019	<i>P. confluens</i> ...	<i>Fries</i> , S.M. I. 355 (1821) ...	Confluent polypore ...
659	628	„ 5072	<i>P. corrivalis</i> ...	<i>Berk.</i> , Linn. Journ. XIII. 162 (1873) ...	Overgrowing polypore ...
660	663	„ 5241	<i>P. cubensis</i> ...	<i>Mont.</i> , Cuba 404 (1838) ...	Cuban polypore ...
661	590	„ 4902	<i>P. cupuliformis</i> ...	<i>Berk. and Curt.</i> , Grev. I. 38 (1872) ...	Cup-shaped polypore ...
662	644	„ 5140	<i>P. demissus</i> ...	<i>Berk.</i> , Hook., Lond. Journ. IV. 52 (1845) ...	Hood-shaped polypore ...
663	647	„ 5152	<i>P. dichrous</i> ...	<i>Fries</i> , S.M. I. 364 (1821) ...	Two-coloured polypore ...
664	613	„ 4982	<i>P. dictyopus</i> ...	<i>Mont.</i> , Fl. Fern. 14 (1835) ...	Net-stalked polypore ...
665	616	„ 5008	<i>P. doreadideus</i> ...	<i>Berk. and Br.</i> , Linn. Trans. II. 57 (1883) ...	Fawn-coloured polypore ...
666	609	„ 4971	<i>P. elegans</i> ...	<i>Fries</i> , Epicr. 440 (1838) ...	Elegant polypore ...
666A	„	„	<i>P. elegans</i> , var. <i>nummularius</i> ...	<i>Fries</i> , S.M. (1821) ...	Coin-like polypore ...
667	627	„ 5067	<i>P. epilencus</i> ...	<i>Fries</i> , Epicr. 452 (1838) ...	Whitish polypore ...
668	656	„ 5216	<i>P. Eucalyptorum</i> ...	<i>Fries</i> , Pl. Preiss. II. 135 (1846) ...	Eucalyptus polypore ...
669	639	„ 5123	<i>P. fædatus</i> ...	<i>Berk.</i> , Linn. Journ. XVI. 41 (1878) ...	Dirty polypore ...
670	633	„ 5082	<i>P. fragilis</i> ...	<i>Fries</i> , Elench. 86 (1828) ...	Fragile polypore ...
671	618	„ 5015	<i>P. frondosus</i> ...	<i>Fries</i> , S.M. I. 355 (1821) ...	Frondose polypore ...
672	649	„ 5169	<i>P. fruticum</i> ...	<i>Berk. and Curt.</i> , Linn. Journ. X. 310 (1869) ...	Shrub-growing polypore ...
673	617	„ 5009	<i>P. fusco-lineatus</i> ...	<i>Berk. and Br.</i> , Linn. Trans. I. 401 (1879) ...	Tawny-lined polypore ...
674	641	„ 5129	<i>P. gilvus</i> ...	<i>Schw.</i> , Carol. 897 (1822) ...	Yellowish-tan polypore ...
675	610	„ 4974	<i>P. glabratus</i> ...	<i>Kalch.</i> , in Hedw. XV. 114 (1876) ...	Smooth polypore ...
676	614	„ 5005	<i>P. grammocephalus</i> ...	<i>Berk.</i> , Hook., Lond. Journ. I. 148 (1842) ...	Line-headed polypore ...
676A	„	„	<i>P. grammocephalus</i> , var. <i>Emerici</i> ...	<i>Berk.</i> , Grev. X. 96 (1882) ...	Emericus polypore ...
676B	„	„	<i>P. grammocephalus</i> , var. <i>Muelleri</i> ...	<i>Kalch.</i> , Grev. X. 97 (1882) ...	Mueller's polypore ...
677	611	„ 4976	<i>P. Guilfoylei</i> ...	<i>Berk. and Br.</i> , Linn. Trans. II. 58 (1883) ...	Guilfoyle's polypore ...
678	631	„ 5079	<i>P. Gunnii</i> ...	<i>Berk.</i> , Fl. Tasm. II. 253 (1860) ...	Gunn's polypore ...
679	585	„ 4866	<i>P. Hartmanni</i> ...	<i>Cooke</i> , Grev. XII. 14 (1884) ...	Hartman's polypore ...
680	648	„ 5165	<i>P. hispidus</i> ...	<i>Fries</i> , S.M. I. 362 (1821) ...	Bristly polypore ...
681	„	IX. 689 } VI 5179 }	<i>P. hispidans</i> ...	<i>Berk.</i> , in <i>Fries</i> , Nov. Syn. 37 (1851) ...	Bristling polypore ...
682	658	VI 5227	<i>P. hypopolius</i> ...	<i>Kalch.</i> , Grev. X. 99 (1882) ...	Ihoary polypore ...
683	602	„ 4952	<i>P. hystriculus</i> ...	<i>Cooke</i> , Grev. XV. 16 (1886) ...	Porcupine-like polypore ...
684	607	„ 4965	<i>P. infernalis</i> ...	<i>Berk.</i> , Hook., Lond. Journ. II. 637 (1843) ...	Infernal polypore ...
685	619	„ 5017	<i>P. intybaceus</i> ...	<i>Fries</i> , Epicr. 446 (1838) ...	Endive polypore ...
686	623	„ 5047	<i>P. lætus</i> ...	<i>Cooke</i> , Grev. XII., 16 (1884) ...	Bright-coloured polypore ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			

Polystictus, Fomes, Favolus, Trametes, Poria, Boletus, Dædalea—continued.

658	...	...	...	...	N.S.W.	Q	...	Wood	...	Branching, firmly fleshy, fragile. Caps thick, overlapping, confluent, smooth, flesh colour to yellowish or dark. Stems very short, confluent. Pores short, minute, white.
659	...	S.A.	...	...	N.S.W.	Q.	...	Trunks	...	Imbricate. Caps shell shaped, whitish, downy. Pores small, angular.
660	...	...	...	...	...	Q.	...	Trunks	...	Pale white. Cap sessile, fleshy to corky, smooth. Pores round, small, white, then red brown.
661	...	...	...	V.	...	...	...	Trunks	...	Cap cup shaped, at first reddish brown, downy. Stem very short. Pores small, red brown.
662	W.A.	...	...	...	...	...	...	Rotten wood	...	Caps overlapping, hood shaped, corky, dependent, spongy, downy, pale yellow. Pores roundish, minute.
663	...	...	...	V.	...	...	...	Trunks	...	Cap fleshy, tough, silky, white. Pores short, minute, round, cinnamon brown. Pretty species.
664	...	...	...	V	...	Q.	...	Trunks	...	Cap fleshy to leathery, rigid, smooth, bay brown. Stem lateral, thick, reticulately wrinkled, bay to black. Pores minute, rounded, pale.
665	...	...	...	...	...	Q.	...	Trunks	...	Cap fan shaped, lobed, rich umber or fawn colour, with velvety bloom. Stem short, downy. Pores hexagonal.
666	...	...	T.	...	...	Q.	B.	Trunks	...	Cap fleshy, soon hardening and becoming woody, flat. Stem excentric or lateral. Pores minute, roundish, yellowish white.
666A	...	...	...	...	N.S.W.	...	B.	Trunks	...	Smaller, thinner, rather regular. Stem equal, excentric.
667	...	...	...	...	...	Q.	B.	Rotten trunks	...	Cap soft and cheesy, then firm, roughly hairy, whitish, semicircular. Pores minute, round, white.
668	W.A.	S.A.	...	V.	...	...	...	Eucalyptus trunks	...	Cap semicircular, hoof shaped, thick, very soft, white, invested with thin evanescent dark-brown crust. Pores short, small, falling away.
669	...	...	...	...	...	Q.	...	Trunks	...	Cap rather thin, kidney shaped, umber to sooty brown. Pores punctiform, pale cinnamon.
670	...	...	...	V.	...	...	B.	Wood	...	White, spotted with brown when touched. Cap fleshy, fragile, kidney shaped, rough. Pores very thin.
671	...	...	T.	...	...	...	B.	On trunks and at base	...	Elaborately branched, fibrously fleshy, tough. Caps very numerous, semicircular, rough, lobed, greyish to sooty brown. Stems growing together. Pores small, white.
672	...	...	...	...	...	Q.	...	Branches	...	Cap thin, soft, semicircular, rhubarb colour. Pores small, angular, toothed.
673	...	...	...	...	...	Q.	...	Trunks	...	Cap, thin, tough, ochrey, streaked with radiating brown, hispid lines. Stem ochrey, thicker above. Pores irregular, tawny brown.
674	W.A.	...	...	...	...	Q.	B.	Trunks	...	Cap fleshy, tough, yellowish tan. Pores minute, yellowish tan to rusty brown.
675	...	...	...	V.	...	...	...	Trunks	...	Cap excentric, fleshy, smooth, dark brown. Stem solid, tapering, becoming brown. Pores minute, round, white to yellowish.
676	...	...	...	...	N.S.W.	Q.	...	Trunks	...	Cap at first wedge shaped, then kidney shaped, flattened, pale umber. Stem lateral. Pores short, umber.
676A	...	...	...	...	...	Q.	...	Trunks	...	Whitish. Cap spoon shaped or kidney shaped. Pores angular.
676B	...	...	...	...	N.S.W.	...	...	Trunks	...	Whitish tan when dry. Cap thin, rigid. Pores short, angular, unequal.
677	...	...	...	...	...	Q	...	Trunks	...	Cap spoon shaped, lateral, fine, powdery. Stem black, cartilaginous. Pores punctiform.
678	...	...	T.	V.	...	...	...	Branches	...	Cap fan shaped, thin, whitish, downy, rough. Pores irregular, of medium size.
679	...	...	...	...	...	Q.	...	Ground	...	Cap fleshy, rather fragile, finely velvety, reddish brown. Stem swollen, thick, reddish. Pores small, round, pale. Elegant species.
680	...	...	...	...	...	Q.	B.	Trunks	...	Large, brown, juicy. Cap compact, spongy to fleshy, semicircular, thick set with bristly down. Pores minute, rounded, pale.
681	...	...	...	...	...	...	...	Trunks	...	Cap semicircular, fleshy to leathery, fawn or dusky, bristly. Pores angular.
682	...	...	...	...	...	Q.	...	Trunks	...	Cap leathery to woody, somewhat shell shaped, with rigid chestnut crust, rough with thick tubercles. Pores short, angular, white to hoary.
683	...	...	...	V.	...	...	...	About root	...	Cap tough, bristly, dark brown. Stem thick, shortened. Pores large, angular, torn or toothed.
684	...	...	...	V.	...	Q.	...	Trunks	...	Cap fan shaped, smooth, blackish, liver coloured. Stem short, lateral, black. Pores minute, round, very short.
685	...	...	...	V.	...	Q	B.	Trunks, and at foot of trees	...	Very much branched, fleshy, rather fragile. Caps very numerous, yellowish to tawny. Stems confluent into a very short trunk. Pores firm, white to tawny.
686	...	...	...	V.	...	...	...	Trunks	...	Imbricated and much divided, leathery, orange tawny. Caps grown together and converging behind into stem. Pores pale.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number	Scientific Name.	Authority for Name.	English Name.
60. POLYPORUS.—Adans, Fam. II. 10 (1763).—					
687	588	VI. 4884	<i>P. lentus</i> ...	Berk., Outl. 237 (1860) ...	Tough polypore ...
688	662	.. 5240	<i>P. lignosus</i> ...	Klotzsch., in Fries, Epicr. 471 (1838) ...	Woody polypore ...
689	604	.. 4958	<i>P. melanopus</i> ...	Fries, S.M. I. 347 (1821) ...	Black-stalked polypore ...
690	587	.. 4870	<i>P. myelodes</i> ...	Kalch., Grev. IV. 73 (1875) ...	Marrow-like polypore... ..
691	1331	...	<i>P. Mylittæ</i> ...	(Cooke and Mass., Grev. XXI. 37 (1892) ...	Mylitta polypore ...
692	638	VI. 5116	<i>P. nidulans</i> ...	(Sacc., Hedw. 56 (1893) ...	Native bread polypore ...
693	583	.. 4858	<i>P. ovinus</i> ...	Fries, S.M. I. 364 (1821) ...	Nest polypore ...
694	650	.. 4858	<i>P. ovinus</i> ...	Fries, S.M. I. 346 (1821) ...	Sheep polypore ...
695	...	IX. 673	<i>P. pelliculosus</i> ...	Berk., Hook., Lond. Journ. VII. 575 (1848) ...	Cuticular polypore ...
696	584	VI. 4862	<i>P. Pentzkei</i> ...	Kalch., Proe. Linn. Soc. N.S.W. VIII. 175 (1883) ...	Pentzke's polypore ...
697	613 bis	.. 4862	<i>P. pes-capræ</i> ...	Pers., Champ. Com. (1818) ...	Goat's-foot polypore ...
698	...	.. 4990	<i>P. petaloides</i> ...	Fries, Epicr. 444 (1838) ...	Petal-like polypore ...
699	605	.. 4999	<i>P. phlebophorus</i> ...	Berk., Fl. N.Z. II. 177 (1855) ...	Vein-bearing polypore ...
700	596	.. 4966	<i>P. picipes</i> ...	Fries, S.M. I. 353 (1821) ...	Pitch-stalked polypore ...
701	615	.. 4933	<i>P. pisiformis</i> ...	Kalch., Grev. X. 98 (1882) ...	Pea-shaped polypore ...
702	664	.. 5007	<i>P. platotis</i> ...	Berk. and Br., Linn. Trans I. 401 (1879) ...	Broad polypore ...
703	655	.. 5247	<i>P. plebeius</i> ...	Berk., Fl. N.Z. II. 179 (1855) ...	Plebeian polypore ...
704	601	.. 5212	<i>P. portentosus</i> ...	Berk., Hook., Journ. 188 (1844) ...	Monstrous polypore ...
705	625	.. 4951	<i>P. proteiporus</i> ...	Cooke, Grev. XII. 15 (1884) ...	Variable-pored polypore ...
706	645	.. 5034	<i>P. retiporus</i> ...	Cooke, Grev. XII. 15 (1884) ...	Net-pored polypore ...
707	640	.. 5141	<i>P. rhinocephalus</i> ...	Berk., Fl. Tasm. II. 253 (1860) ...	Rough-headed polypore ...
708	600	.. 5124	<i>P. rubidus</i> ...	Berk., Hook., Journ. 500 (1847) ...	Reddish polypore ...
709	621	.. 4946	<i>P. rufescens</i> ...	Fries, S.M. I. 351 (1821) ...	Reddening polypore ...
710	597	.. 5026	<i>P. scabriusculus</i> ...	Berk., Linn. Journ. XVIII. 384 (1881) ...	Roughish polypore ...
711	642	.. 4938	<i>P. Schweinitzii</i> ...	Fries, S.M. I. 351 (1821) ...	Schweinitz's polypore ...
711A	643	.. 5130	<i>P. scruposus</i> ...	Fries, Epicr. 473 (1838) ...	Rugged polypore ...
712	630	.. 5131	<i>P. scruposus, var. isidioides</i> ...	Cooke, Grev. XIII. 87 (1885) ...	Coral-like polypore ...
713	593	.. 5078	<i>P. semidigitaliformis</i> ...	Berk., Linn. Journ. XVI. 39 (1878) ...	Finger-like polypore ...
714	651	.. 4910	<i>P. similis</i> ...	Berk., Hook., Lond. Journ. II. 635 (1843) ...	Similar polypore ...
715	...	.. 5181	<i>P. spiculifer</i> ...	Cooke, Grev. XV. 20 (1886) ...	Spiculate polypore ...
716	603	.. 5186	<i>P. spumeus</i> ...	Fries, S.M. I. 358 (1821) ...	Frothy polypore ...
717	595	.. 4953	<i>P. squamosus</i> ...	Fries, S.M. I. 343 (1821) ...	Scaly polypore ...
718	606	IX. 667	<i>P. stipitarius</i> ...	Berk. and Curt., Linn. Journ. X. 304 (1869) ...	Stalked polypore ...
719	657	VI. 5220	<i>P. Strangeri</i> ...	F. v. M., Linn. Soc. N.S.W. 106 (1882) ...	Stranger's polypore ...
			<i>P. strumosus</i> ...	Fries, Epicr. 462 (1838) ...	Swollen polypore ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Polystictus, Fomes, Favolus, Trametes, Poria, Boletus, Daldalea—continued.									
687	...	...	...	V.	...	...	B.	Rotten stems, &c.	Cap fleshy to tough and leathery, pale ochrey. Stem short, rough haired and mealy. Pores irregular, white.
688	...	...	...	V.	...	...	...	Trunks	Cap fleshy to corky or woody, pale to yellowish. Pores long, small, very thin.
689	...	...	...	V.	...	Q.	B.	Ground and branches	Cap fleshy, tough, white to yellowish brown. Stem excentric, velvety at first, black. Pores rounded, small, white to yellowish
690	...	...	...	...	...	Q.	...	Ground, at base of trunks	Cap fleshy, fragile, tan or pale brown. Stem solid, obconical. Pores short, minute, whitish. <i>Edible</i> .
691	...	S.A.	T.	V.	N.S.W.	Q.	...	Sclerotium, known as <i>Mytilia australis</i>	Cap fleshy, tough, elastic, minutely velvety, white. Stem short, solid. Pores white, somewhat angular.
692	...	...	...	...	...	Q.	B.	Trunks	Cap fleshy, very soft, yellowish tan. Pores elongated, angular, tawny, brick red.
693	...	...	...	V.	...	...	...	Woods	Cap fleshy, fragile, whitish. Stem short, white. Pores minute, rounded, white to lemon yellow.
694	...	S.A.	T.	V.	...	Q.	...	Wood	Dark red, juicy, densely gregarious, shell shaped, at first with dense rough hairs. Pores rather angular.
695	...	...	...	...	...	Q.	...		Cap slender, leathery, base wedge shaped, smooth, zoned, ochrey, becoming dark. Stem cylindrical, pale.
696	...	...	...	V.	...	...	...	Pine woods	In tufts. Cap fleshy, fragile, bay brown to dark brown. Stem deformed, yellowish white. Pores broad, yellowish white.
697	...	...	...	V.	...	...	B.	Trunks	Cap membranaceous, spoon shaped, chestnut brown. Stem lateral, compressed, whitish. Pores very short, small, white.
698	...	...	...	...	...	Q.	...	Stems of <i>Eucalyptus hemiphloia</i>	Small, white. Cap fan shaped. Stem short, smooth, with undulating furrows.
699	...	...	...	V.	...	Q.	B.	Trunks	Cap fleshy to leathery, smooth, depressed. Stem excentric and lateral, firm, black. Pores rounded, small, white to yellowish.
700	...	...	...	V.	...	...	...	Wood	Entirely white, globose, sessile, size of pea or less. Pores minute, punctiform. Probably young stage of known species.
701	...	...	...	...	...	Q.	...	Wood	Cap club shaped to funnel shaped, fragile, ochrey. Stem elongated and thickened upwards. Pores angular, ochrey.
702	...	...	...	V.	...	Q.	...	Rotten wood	Pale, imbricate. Cap semicircular, delicately downy, corky. Pores minute, punctiform.
703	W.A.	S.A.	T.	V.	N.S.W.	Q.	...	Trunks	Sessile, very large. Cap fleshy, smooth, with thick tan cuticle. Pores small, externally brown, internally pale.
704	...	...	...	V.	...	Q.	...	Ground	Cap whitish, fleshy, tough, delicately downy and scurfy. Stem short, whitish. Pores angular, irregular, pale umber.
705	...	...	...	V.	...	Q.	...	Trunks	Tufted, forming clumps of juicy cheesy consistence, ochrey. Caps very broad. Pores angular, very short, with net-like partitions.
706	...	...	T.	...	...	...	...	Rotten wood	Cap semicircular, shell shaped, whitish, rough veined. Pores white to ashy.
707	...	...	...	...	N.S.W.	Q.	...	Wood	Reddish. Cap thin, leathery, kidney shaped, silky. Pores small, short, punctiform.
708	W.A.	...	...	V.	N.S.W.	Q.	B.	Grassy ground about old trunks	Flesh coloured. Cap spongy, hairy. Stem short, deformed. Pores large, sinuous, white to flesh colour.
709	...	...	...	...	...	Q.	...	Trunks	Caps spoon shaped, delicately downy, roughish. Stem cylindrical, repeatedly branched, central, sometimes nearly obsolete.
710	...	...	...	...	...	Q.	B.	Pine stumps	Very large. Cap thick, spongy to corky, bay brown. Stem thick, very short or obsolete, rusty brown. Pores large, sulphury to greenish.
711	W.A.	...	T	V.	N.S.W.	Q.	...	Dead wood	Cap corky, rough and rugged, umber. Pores minute, rounded, rusty brown.
711A	W.A.	...	...	...	...	...	...	Trunks and at foot of trees	Corky or woody, sessile, yellowish tan to rusty, rough, with thick tubercles.
712	...	...	...	...	N.S.W.	...	...	Trunks	Gregarious. Cap hoof shaped, whitish, rough, and downy. Pores large.
713	...	...	...	...	...	Q.	...	Trunks	Cap tough, leathery, smooth. Stem thickened downwards, velvety, becoming smooth. Pores small, angular, pale.
714	...	...	...	V.	...	...	...	Trunks	Cap fleshy, soft, and watery, sooty brown to black, beset with scattered obtuse spicules. Pores minute.
715	...	...	...	V.	...	...	B.	Old trunks of <i>Eucalypts</i>	White. Cap fleshy to spongy, compact, cushion shaped, rough haired, base stem-like.
716	...	...	...	V.	...	Q.	B.	Trunks	Cap fleshy to tough, fan shaped, ochrey, marked with brown scales. Stem stout, excentric, black at base. Very common.
717	...	...	...	...	...	Q.	...	Wood, &c.	White. Cap circular, thin, smooth. Stem slender,
718	...	...	...	...	N.S.W.	...	...	Trunks	Cap corky to leathery, kidney shaped, umber, turning blackish. Stem short, cylindrical, altogether black.
719	...	...	...	V.	...	...	...	Trunks	Fleshy, tough, afterwards very hard, sooty brown, margin acute, turning black.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
60. POLYPORUS.—Adans, Fam. II. 10 (1763).—					
720	634	VI. 5092	<i>P. stypticus</i> ...	<i>Fries</i> , S.M. I. 359 (1821) ...	Astringent polypore ...
721	653	„ 5194	<i>P. substuppeus</i> ...	<i>Berk. and Cooke</i> , Linn. Journ. XV. 380 (1877) ...	Tow-like polypore ...
722	661	IX. 696	<i>P. subzonalis</i> ...	<i>Cooke</i> , Grev. XIX. 44 (1890) ...	Slightly-zoned polypore ...
723	624	VI. 5050	<i>P. sulphureus</i> ...	<i>Fries</i> , S.M. I. 357 (1821) ...	Sulphur-coloured polypore ...
724	598	„ 4939	<i>P. tabuleformis</i> ...	<i>Berk., Hook.</i> , Lond. Journ. IV. 302 (1845) ...	Flat polypore ...
725	626	„ 5064	<i>P. tephronotus</i> ...	<i>Berk.</i> , Fl. Tasm. II. 252 (1860) ...	Ashy polypore ...
726	665	„ 5248	<i>P. testudo</i> ...	<i>Berk. and Broome</i> , Linn. Trans. II. 59 (1883) ...	Tortoise polypore ...
727	592	„ 4908	<i>P. tricholoma</i> ...	<i>Mont.</i> , Syll. I. 53 (1856) ...	Hairy polypore ...
728	586	IX. 650	<i>P. tumulosus</i> ...	<i>Cooke</i> , Grev. XVII. 55 (1889) ...	Buried polypore ...
729	...	VI. 4907	<i>P. umbilicatus</i> ...	<i>Berk., Hook.</i> , Journ. 79 (1851) ...	Umbilicate polypore ..
730	...	„ 5222	<i>P. unguatus</i> ...	<i>Cooke</i> , Grev. XIII. 116 (1885) ...	Hooked polypore ...
731	608	„ 4968	<i>P. varius</i> ...	<i>Fries</i> , S.M. I. 352 (1821) ...	Variable polypore ...
732	...	„ 5178	<i>P. Weinmanni</i> ...	<i>Fries</i> , Epier. 459 (1838) ...	Weinmann's polypore ...
733	660	„ 5237	<i>P. zonalis</i> ...	<i>Berk.</i> , Ann. Nat. Hist. X. 375 (1842) ...	Zoned polypore ...
61. FOMES.—Fries,					
734	672	VI. 5300	<i>F. amboinensis</i> ...	<i>Cooke</i> , Grev. XIII. 118 (1885) ...	Amboina fomes ...
734A	672	„ 5303	<i>F. amboinensis</i> , var. <i>gibbosus</i> ...	<i>Cooke</i> , Grev. XIII. 118 (1885) ...	Swollen fomes ...
735	708	„ 5487	<i>F. annosus</i> ...	<i>Cooke</i> , Grev. XIV. 20 (1885) ...	Aged fomes ...
736	683	„ 5397	<i>F. applanatus</i> ...	<i>Cooke</i> , Grev. XIV. 18 (1885) ...	Flattened fomes ...
737	681	„ 5394	<i>F. australis</i> ...	<i>Cooke</i> , Grev. XIV. 18 (1885) ...	Southern fomes ...
737A	681	IX. 723	<i>F. australis</i> , var. <i>arcuatus</i> ...	<i>Bres.</i> , Pug. Myc. Austr. (1890) ...	Bow-shaped fomes ...
738	723	VI. 5529	<i>F. bistratosus</i> ...	<i>Cooke</i> , Grev. XIV. 21 (1885) ...	Stratose fomes ...
739	717	„ 5509	<i>F. carneus</i> ...	<i>Cooke</i> , Grev. XIV. 21 (1885) ...	Flesh-coloured fomes ...
740	682	„ 5396	<i>F. chilensis</i> ...	<i>Cooke</i> , Grev. XIV. 18 (1885) ...	Chilian fomes ...
741	718	„ 5512	<i>F. cinereo-fusens</i> ...	<i>Cooke</i> , Grev. XIV. 21 (1885) ...	Ashy-brown fomes ...
742	709	„ 5491	<i>F. compressus</i> ...	<i>Cooke</i> , Grev. XV. 51 (1886) ...	Compressed fomes ...
743	679	IX. 705	<i>F. concavus</i> ...	<i>Cooke</i> , Grev. XIX. 44 (1890) ...	Concave fomes ...
744	680	VI. 5385	<i>F. conchatus</i> ...	<i>Cooke</i> , Grev. XIV. 18 (1885) ...	Shell-shaped fomes ...
745	710	„ 5485	<i>F. connatus</i> ...	<i>Cooke</i> , Grev. XIV. 20 (1885) ...	Connate fomes ...
746	694	„ 5451	<i>F. contrarius</i> ...	<i>Berk. and Curt.</i> , Grev. XV. 21 (1886) ...	Contrary fomes ...
747	...	„ 5524	<i>F. cryptarum</i> ...	<i>Fries</i> , S.M. I. 376 (1821) ...	Crypt fomes ...
748	705	„ 5481	<i>F. Curreyi</i> ...	<i>Berk.</i> , Grev. XV. 21 (1886) ...	Currey's fomes ...
749	678	„ 5359	<i>F. dochmius</i> ...	<i>Cooke</i> , Grev. XIV. 17 (1885) ...	Oblique fomes ...
750	704	„ 5478	<i>F. endapatus</i> ...	<i>Cooke</i> , Grev. XIV. 20 (1885) ...	Soft fomes ...
751	693	„ 5450	<i>F. exotephrus</i> ...	<i>Cooke</i> , Grev. XIV. 19 (1885) ...	Ashy fomes ...
752	716	„ 5499	<i>F. fasciatus</i> ...	<i>Cooke</i> , Grev. XIV. 21 (1885) ...	Banded fomes ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
720	W.A.	...	...	...	...	...	...	Trunks ...	Fleshy to corky, cushion shaped, fragile, whitish; margin obtuse, somewhat reddish.
721	...	...	...	...	...	Q.	...	Wood ...	Semicircular, decurrent behind, rough, short, invested with tow-like wool, pale ochrey when dry.
722	...	...	...	...	...	Q.	...	Wood ...	Corky, rather thin, sessile, entirely cream coloured, kidney shaped, faintly concentrically zoned.
723	...	...	T.	...	...	Q.	B.	Trunks ...	Large, tufted, and much divided; of juicy, cheesy, or doughy consistence. Caps very broad, overlapping, reddish or lemon yellow.
724	...	...	...	...	...	Q.	...	Trunks ...	Cap circular, thick in centre, thin at margin, somewhat lobed, slightly zoned, velvety, rusty bay. Stem short, central or lateral.
725	...	...	T.	...	N.S.W.	...	...	Rotten wood	Soft, downy, snow white. brown behind. Hymenium white, then turning slightly ashy.
726	...	...	...	...	...	Q.	...	Trunks ...	Caps overlapping, rigid, powdery, obscurely streaked and rough here and there.
727	...	...	...	...	...	Q.	...	Fallen branches	Cap leathery to membranous, rigid, convex, then funnel shaped, yellowish, with fringe of stiff brown hairs. Stem thin, yellowish brown.
728	...	...	...	...	...	Q.	...	Ground ...	Cap fleshy, firm, pale, with darker sunken scales. Stem short, thick, solid, ochrey.
729	...	...	T.	...	...	...	...	Wood ...	Cap fleshy to tough, umbilicate, then somewhat funnel shaped, snow white or cream coloured. Stem swelling above, and below minutely scaly.
730	...	S.A.	...	...	...	...	...	Trunks ...	Cap hard, whitish, shortly hooked behind, delicately downy, margin obtuse, wrinkled.
731	W.A.	...	T.	V.	...	Q.	B.	Trunks ...	Cap fleshy, tough, becoming woody, variable in form. Stem excentric and lateral, finally black.
732	...	...	...	V.	...	...	...	Trunks ...	White to red. Cap fleshy to spongy, firm, with reddish-brown hairs.
733	...	...	...	V.	...	Q.	...	Wood ...	Corky, thin, overlapping, sessile, semicircular, lobed, repeatedly zoned, covered with bloom, fawn colour.

Nov. Symb. 31 (1851).—*Agaricus*, *Polyporus*, *Boletus*, *Trametes*, *Ganoderma*.

734	...	...	...	...	...	Q.	...	Trunks ...	Cap corky to woody, somewhat ear-shaped, rough, pimpled. Stem lateral, very long, turning black.
734A	...	...	...	...	...	Q.	...	Trunks ...	Cap somewhat kidney shaped, and stem reddish brown.
735	...	...	...	...	...	Q.	B.	Trunks ...	Woody, rough, for the current year brown and silky; for previous season with blackened crust.
736	...	...	...	V.	...	Q.	B.	Trunks ...	Flattened, tuberculous, obsoletely zoned, powdery, cinnamon to hoary.
737	...	...	T.	V.	N.S.W.	Q.	...	Trunks ...	Very hard, semicircular, sessile, wavy, incrinated, somewhat bay brown.
737A	...	...	...	...	...	Q.	...	Trunks ...	Margin thicker, bow shaped.
738	...	...	...	...	...	Q.	...	Wood ...	Spread out, umber, with very little substance. Pores stratosse, punctiform.
739	...	S.A.	...	V.	...	Q.	B.	Trunks ...	Expanded and reflexed, woody, hard, thin, without zones, flesh coloured without and within.
740	...	...	...	...	...	Q.	...	Trunks ...	Corky, hoof shaped and dilated, with elevated ridges, brick red, turning pale.
741	...	...	...	...	...	Q.	...	Trunks ...	Semicircular, woody, very hard. Margin thin, ashy brown.
742	W.A.	...	...	V.	...	Q.	...	Wood ...	Small, obliquely compressed, hoof shaped, zoned, light brown at first, dark brown afterwards.
743	...	...	...	...	...	Q.	...	Trunks ...	Very hard, semicircular, turning nearly black, comparatively thin and covered with hard crust.
744	...	...	...	V.	...	Q.	B.	Trunks ...	Corky to tindery, thin, spread out, somewhat shell shaped, bay brown.
745	...	...	...	...	...	Q.	B.	Trunks ...	Corky to woody, spread out, overlapping and growing into each other, downy white or ashy.
746	...	...	...	...	...	Q.	...	Trunks ...	Somewhat zoned, brown at first, ochrey, thin, rigid, downy, finally smooth.
747	...	...	...	...	...	Q.	...	Logs and rotten wood	Corky to tow like, zoneless, silky, reddish to rusty brown, but variable.
748	...	...	...	...	...	Q.	...	Trunks ...	Rigid, corky to leathery, semicircular, kidney shaped, brown, with concentric elevated zones.
749	...	...	...	...	...	Q.	...	Wood ...	Semicircular, oblique, hard, zoned, banded.
750	...	...	...	...	N.S.W.	Q.	...	...	Overlapping each other, leathery, bay brown, longitudinally rough in lines, delicately downy, substance soft.
751	...	...	...	...	...	Q.	...	Trunks ...	Hard, zoned, delicately downy at first, then smooth. Margin furrowed, lobed.
752	...	...	...	...	...	Q.	...	Trunks ...	Woody, thin, flattened, rusty brown with black bands.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
61. FOMES.—Fries, Nov. Symb. 31 (1851).—					
753	713	VI. 5501	<i>F. ferrens</i>	<i>Cooke</i> , Grev. XIV. 21 (1885)	Hard fomes
754	686	„ 5409	<i>F. fomentarius</i>	<i>Cooke</i> , Grev. XIV. 18 (1885)	Tender fomes
755	689	„ 5417	<i>F. fulvus</i>	<i>Cooke</i> , Grev. XIV. 18 (1885)	Brown fomes
756	703	„ 5477	<i>F. Gourliei</i>	<i>Cooke</i> , Grev. XIV. 20 (1885)	Gourlie's fomes
757	690	„ 5424	<i>F. gryphæformis</i>	<i>Cooke</i> , Grev. XIV. 19 (1885)	Shell-shaped fomes
758	692	„ 5449	<i>F. hemileucus</i>	<i>Cooke</i> , Grev. XIV. 19 (1885)	Whitish fomes
759	711	„ 5497	<i>F. hemitephrus</i>	<i>Cooke</i> , Grev. XIV. 21 (1885)	Half-ashy fomes
760	719	„ 5519	<i>F. homalopilus</i>	<i>Cooke</i> , Grev. XIV. 21 (1885)	Smooth-piled fomes
761	687	„ 5412	<i>F. igniarius</i>	<i>Cooke</i> , Grev. XIV. 18 (1885)	Tinder fomes
762	720	„ 5523	<i>F. incrassatus</i>	<i>Cooke</i> , Grev. XIV. 21 (1885)	Thickened fomes
763	699	„ 5468	<i>F. inflexibilis</i>	<i>Cooke</i> , Grev. XIV. 20 (1885)	Inflexible fomes
764	697	„ 5465	<i>F. lineato-seaber</i>	<i>Cooke</i> , Grev. XV. 51 (1886)	Rough-lined fomes
765	700	„ 5470	<i>F. linteus</i>	<i>Cooke</i> , Grev. XIV. 20 (1885)	Linty fomes
766	673	„ 5305	<i>F. lucidus</i>	<i>Cooke</i> , Grev. XIII. 118 (1885)	Varnished fomes
767	722	„ 5528	<i>F. luridus</i>	<i>Cooke</i> , Grev. XIV. 21 (1885)	Lurid fomes
768	677	„ 5352	<i>F. marginatus</i>	<i>Cooke</i> , Grev. XIV. 17 (1885)	Margined fomes
769	668	„ 5272	<i>F. nigripes</i>	<i>Cooke</i> , Grev. XIII. 117 (1885)	Black-stalked fomes
770	685	„ 5401	<i>F. nigro-laccatus</i>	<i>Cooke</i> , Grev. XIV. 18 (1885)	Black-lacquered fomes
771	715	„ 5507	<i>F. oblinitus</i>	Berk., Grev. XV. 22 (1886)	Variegated fomes
772	721	„ 5527	<i>F. obliquus</i>	<i>Cooke</i> , Grev. XIV. 21 (1885)	Oblique fomes
773	684	„ 5400	<i>F. orbiformis</i>	<i>Cooke</i> , Grev. XIV. 18 (1885)	Orbicular fomes
774	712	„ 5520	<i>F. Palliseri</i>	<i>Cooke</i> , Grev. XIV. 21 (1885)	Palliser's fomes
775	701	„ 5469	<i>F. pectinatus</i>	<i>Cooke</i> , Grev. XIV. 20 (1885)	Comb-like fomes
776	707	„ 5484	<i>F. ponderosus</i>	<i>Cooke</i> , Grev. XIV. 20 (1885)	Weighty fomes
777	670	„ 5282	<i>F. pullatus</i>	<i>Cooke</i> , Grev. XIII. 117 (1885)	Mourning fomes
778	696	„ 5461	<i>F. pullus</i>	<i>Cooke</i> , Grev. XIV. 19 (1885)	Russet-brown fomes
779	688	„ 5415	<i>F. rimosus</i>	<i>Cooke</i> , Grev. XIV. 18 (1885)	Cracked fomes
780	702	„ 5473	<i>F. rubiginosus</i>	<i>Cooke</i> , Grev. XIV. 20 (1885)	Rusty fomes
781	669	„ 5281	<i>F. rudis</i>	<i>Cooke</i> , Grev. XIII. 117 (1885)	Rough fomes
782	671	„ 5283	<i>F. rugosus</i>	<i>Cooke</i> , Grev. XIII. 117 (1885)	Wrinkled fomes
783	691	„ 5429	<i>F. salicinus</i>	<i>Cooke</i> , Grev. XIV. 19 (1885)	Willow fomes
784	676	„ 5342	<i>F. scansilis</i>	<i>Cooke</i> , Grev. XIII. 119 (1885)	Climbing fomes
785	714	„ 5505	<i>F. scopulosus</i>	<i>Cooke</i> , Grev. XIV. 21 (1885)	Craggy fomes
786	675	„ 5335	<i>F. senex</i>	<i>Cooke</i> , Grev. XIII. 118 (1885)	Old fomes
787	698	„ 5466	<i>F. spadiceus</i>	<i>Cooke</i> , Grev. XIV. 20 (1885)	Bright-brown fomes
788	706	„ 5480	<i>F. strigatus</i>	<i>Cooke</i> , Grev. XIV. 20 (1885)	Stiff-haired fomes
789	674	„ 5315	<i>F. superpositus</i>	<i>Cooke</i> , Grev. XIII. 118 (1885)	Superposed fomes
790	695	„ 5452	<i>F. tasmanicus</i>	<i>Cooke</i> , Grev. XIV. 19 (1885)	Tasmanian fomes

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Agaricus, Polyporus, Boletus, Trametes, Ganoderma—continued.									
753	...	...	...	...	N.S.W.	Q	...	Wood ...	Hard like iron, corky. Caps of current year fawn coloured; velvety of previous year, banded with brown.
754	...	...	...	...	N.S.W.	...	B.	Stumps ...	Hoof shaped, cushion like, thick, sooty brown, becoming hoary, throwing off a snuff-like powder.
755	W.A.	S.A.	...	V.	N.S.W.	Q	B.	Trunks ...	Woody to corky, very hard at first; hairy, brown, then hoary.
756	...	...	T.	...	...	...	...	Bark ...	Semicircular, convex, sparingly zoned, velvety like tow, umber.
757	W.A.	...	...	...	...	...	...	Trunks ...	Very hard, hemispherical, shell shaped, cinnamon; margin rather thin, bay brown.
758	...	...	...	...	...	Q.	...	Stems ...	Corky, thick, rigid, semicircular, delicately downy, white.
759	...	...	...	V.	...	...	...	Trunks ...	Hoof shaped, concentrically furrowed, purple brown to ashy, very delicately downy.
760	...	...	...	...	...	Q.	...	Trunks ...	Semicircular, sessile, leathery to corky, rigid, thin, reddish brown.
761	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Trunks; very common on <i>Casuarinas</i>	Hoof shaped, with thin hoary cuticle, rusty brown, becoming blackish, substance zoned and very hard. Common.
762	...	...	...	...	...	Q	...	Trunks ...	Hard, kidney shaped, at first thin, without zones, at length thickened, repeatedly zoned, coffee colour.
763	...	...	...	...	...	Q	...	Trunks ...	Hoof shaped, brown, crested, furrowed, quite hard.
764	...	...	...	...	...	Q.	...	Trunks ...	Semicircular, rigid, brown; margin frequently zoned, lineately radiate, rough.
765	...	...	...	...	...	Q.	...	Bark ...	Hard, heavy, semicircular, furrowed, radiately cracked, brown, but rendered pale by lint-white down.
766	...	...	T.	...	...	Q.	B.	Base of stumps ...	Cap corky to woody, fan shaped, rough, shining as if lacquered, as well as lateral stem, yellowish red or brown. Very common.
767	...	...	...	...	N.S.W.	Q.	...	Branches ...	Spread out, closely adherent to the wood; pores white, then livid or lurid.
768	...	...	...	V.	N.S.W.	...	...	Trunks ...	Corky to woody, hoof shaped, somewhat flattened, incrustated, concentrically furrowed, covered with greyish bloom.
769	...	...	...	...	N.S.W.	...	...	Trunks ...	Corky to woody. Cap convex, zoneless, opaque, rusty brown. Stem rooting, shining as if lacquered, black.
770	...	...	...	...	...	Q.	...	Wood ...	Fan shaped, corky or woody, rough; margin wavy, chestnut brown to black, shining as if lacquered.
771	...	...	...	...	N.S.W.	...	...	Trunks ...	Corky to woody, convex to flattened, kidney shaped, variegated with faint concentric zones, red brown.
772	...	S.A.	...	V.	N.S.W.	Q.	...	Trunks, <i>Eucalyptus</i> , &c.	A magnificent fungus. Thick, casting off the bark, pale to bay brown, then blackish.
773	...	...	...	V.	...	...	...	Trunks ...	Very hard, convex, circular, concentrically furrowed, crustaceous.
774	...	...	...	V.	N.S.W.	Q.	...	Trunks ...	Fleshy to tough and leathery, shell shaped, slightly silky, white behind, ashy brown in front.
775	...	...	...	...	...	Q.	B.	Trunks ...	Corky to woody, hard, concentrically lamellately folded, scurfy, rusty brown.
776	...	...	...	...	...	Q.	...	Trunks ...	Remarkable for hardness and weight. Woody, semicircular, sessile, imbricate, shell shaped, lurid grey, at base brownish.
777	...	...	...	V.	...	Q.	...	Ground ...	Circular, rough, with furrowed zones, at first glaucous, delicately velvety, at length brown.
778	...	...	...	...	...	Q.	...	Branches of <i>Jasminum racemosum</i>	Small, somewhat imbricate, laterally confluent, hard, semicircular, shell shaped, bay brown.
779	W.A.	S.A.	T.	...	N.S.W.	...	...	Gum-tree trunks ...	Woody, very hard, hoof shaped, at length cracked, deeply furrowed, dark umber, and nearly black when old.
780	...	...	T.	...	...	...	...	Rotten wood ...	Horizontal, solid, sessile, thin, zoned, rough, minutely velvety when young, rusty, when old tinged with brown.
781	...	...	T.	...	...	...	...	Rotten wood ...	Cap circular, brown, rough, covered with a bloom or fine velvet. Stem nearly central, rooting in wood, brown, covered with tawny bloom, shiny.
782	...	...	...	V.	N.S.W.	Q.	...	Ground ...	Leathery, rigid, concentrically furrowed, bay brown, turning black.
783	...	...	...	...	...	Q.	B.	Trunks of Willows, &c.	Woody, quite hard, wavy, smooth, cinnamon brown or rusty; scent of aniseed.
784	...	...	...	...	...	Q.	...	Trunks ...	Cushion shaped, brown, repeatedly deeply furrowed and ribbed, coffee colour
785	...	...	...	...	...	Q.	...	Wood ...	Woody, hard, fan shaped, fixed by the vertex, whitish, zoned, rough.
786	...	...	...	V.	N.S.W.	Q.	...	Trunks ...	Large, nearly plane, corky, chestnut brown.
787	...	...	...	...	...	Q.	...	Trunks ...	Hard, leathery or corky, thin, bright brown, minutely velvety, closely zoned.
788	...	...	...	...	...	Q.	...	Trunks ...	Rigid, thin, semicircular, brown, zoned, with small scattered stiff hairs.
789	...	...	...	...	N.S.W.	...	...	Trunks ...	Cap shell shaped, imbricate, arising from a common lateral cylindrical stem, pale, covered with bloom.
790	...	...	T.	...	...	...	...	Rotten wood ...	Narrow, furrowed, pale brown, downy.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
62. POLYSTICTUS.—Fries, Nov. Symb. 54 (1831).—					
791	764	VI. 5702	<i>P. acutus</i> ...	<i>Cooke</i> , Grev. XIV. 82 (1886) ...	Acute polystictus ...
792	735	„ 5572	<i>P. adami</i> ...	<i>Cooke</i> , Grev. XIV. 78 (1886) ...	Adam's Peak polystictus ...
793	740	„ 5584	<i>P. affinis</i> ..	<i>Cooke</i> , Grev. XIV. 78 (1886) ...	Allied polystictus ...
794	799	„ 5869	<i>P. aratus</i> ...	<i>Cooke</i> , Grev. XIV. 86 (1886) ..	Furrowed polystictus ...
795	751	„ 5125	<i>P. Beckleri</i> ...	<i>Cooke</i> , Handb. Austr. Fungi 142 (1892)...	Beckler's polystictus ...
796	759	„ 5683	<i>P. biformis</i> ...	<i>Cooke</i> , Grev. XIV. 81 (1886) ...	Two-shaped polystictus ...
797	808	„ 5921	<i>P. bireflexus</i> ...	<i>Cooke</i> , Grev. XIV. 87 (1886) ...	Bireflexed polystictus ...
798	807	„ 5917	<i>P. Braunii</i> ...	<i>Cooke</i> , Grev. XIV. 87 (1886) ...	Braun's polystictus ...
799	806	„ 5909	<i>P. breviporus</i> ...	<i>Cooke</i> , Grev. XIV. 87 (1886) ...	Short-pored polystictus ...
800	810	„ 5931	<i>P. Broomei</i> ...	<i>Cooke</i> , Grev. XIV. 87 (1886) ...	Broome's polystictus ...
801	785	„ 5785	<i>P. brunneo-albus</i> ...	<i>Cooke</i> , Grev. XIV. 83 (1886) ...	Browish-white polystictus ...
802	743	„ 5616	<i>P. brunneolus</i> ...	<i>Cooke</i> , Grev. XIV. 79 (1886) ...	Brown polystictus ...
803	729	„ 5546	<i>P. bulbipes</i> ...	<i>Cooke</i> , Grev. XIV. 77 (1886) ...	Bulbous-stalked polystictus ...
804	796	„ 5846	<i>P. byrsinus</i> ...	<i>Cooke</i> , Grev. XIV. 85 (1886) ...	Leathery polystictus ...
805	805	„ 5887	<i>P. caperatus</i> ...	<i>Cooke</i> , Grev. XIV. 86 (1886) ...	Wrinkled polystictus ...
806	738	„ 5581	<i>P. carneo-niger</i> ...	<i>Cooke</i> , Grev. XIV. 78 (1886) ..	Fleshy black polystictus ...
807	803	„ 5878	<i>P. cichoraceus</i> ...	<i>Cooke</i> , Grev. XIV. 86 (1886) ..	Chicory-coloured polystictus ...
808	770	„ 5711	<i>P. cinabarius</i> ...	<i>Cooke</i> , Grev. XIV. 82 (1886) ...	Vermilion polystictus ...
809	727	„ 5542	<i>P. cinuamomeus</i> ...	<i>Sacc.</i> , Syll. VI. 210 (1888) ...	Cinnamon polystictus ...
810	798	„ 5866	<i>P. citreus</i> ...	<i>Cooke</i> , Grev. XIV. 85 (1886) ...	Lemon-yellow polystictus ...
811	763	„ 5700	<i>P. cristatus</i> ...	<i>Cooke</i> , Grev. XIV. 81 (1886) ...	Crested polystictus ...
812	792	„ 5836	<i>P. cupreo-roseus</i> ...	<i>Cooke</i> , Grev. XIV. 85 (1886) ...	Copper-rose polystictus ...
813	749	„ 5782	<i>P. dispar</i> ...	<i>Cooke</i> , Handb. Austr. Fungi 142 (1892)...	Unequal polystictus ...
814	750	„ 5640	<i>P. elongatus</i> ...	<i>Cooke</i> , Grev. XIV. 80 (1886) ...	Elongated polystictus ...
814A	750	„ 5640	<i>P. elongatus</i> , var. <i>Hodgkinsoniae</i>	<i>Kalch.</i> , Grev. X. 96 (1882) ...	Hodgkinson's polystictus ...
815	809	„ 5922	<i>P. eriophorus</i> ...	<i>Cooke</i> , Grev. XIV. 87 (1886) ...	Cottony polystictus ...
816	769	„ 5709	<i>P. Eucalypti</i> ...	<i>Cooke</i> , Grev. XIV. 82 (1886) ...	Eucalyptus polystictus ...
817	765	„ 5703	<i>P. extensus</i> ...	<i>Cooke</i> , Grev. XIV. 82 (1886) ...	Extended polystictus ...
818	768	„ 5706	<i>P. Feei</i> ...	<i>Cooke</i> , Grev. XIV. 82 (1886) ...	Fee's polystictus ...
819	733	„ 5569	<i>P. flabelliformis</i> ...	<i>Cooke</i> , Grev. XIV. 78 (1886) ...	Fan-shaped polystictus ...
820	773	„ 5733	<i>P. floridanus</i> ...	<i>Cooke</i> , Grev. XIV. 82 (1886) ...	Florida polystictus ...
821	754	„ 5646	<i>P. Friesii</i> ...	<i>Cooke</i> , Grev. XIV. 80 (1886) ...	Fries' polystictus ...
822	758	„ 5665	<i>P. funalis</i> ...	<i>Cooke</i> , Grev. XIV. 80 (1886) ...	Cord-like polystictus ...
823	756	„ 5656	<i>P. gallo-pavonis</i> ...	<i>Cooke</i> , Grev. XIV. 80 (1886) ...	Peacock polystictus ...
824	779	„ 5770	<i>P. gausapatus</i> ...	<i>Cooke</i> , Grev. XIV. 83 (1886) ...	Friezed polystictus ...
825	780	„ 5773	<i>P. glirinus</i> ...	<i>Cooke</i> , Grev. XIV. 83 (1886) ...	Dormouse polystictus ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.	
	W.A.	S.A.	T.	V.	N.S.W.	Q.				
Trametes, Polyporus, Boletus.										
791	...	...	...	...	N.S.W.	Q.	...	Branches	...	Spread out, reflexed, thin, pale ochrey, flexible, margin very acute.
792	...	...	...	...	N.S.W.	Q.	...	Rotten wood	...	Cap lateral, spoon shaped, thin, leathery, zoned, bay brown. Stem long, mealy, yellow.
793	...	...	...	...	N.S.W.	Q.	...	Branches	...	Cap papery, fan shaped, rigid, chestnut brown with darker zones. Stem lateral, thin, chestnut brown.
794	...	...	...	...	N.S.W.	Q.	...	Trunks	...	Yellowish olive, semicircular, flexible, concentrically furrowed, shining, smooth.
795	...	...	...	...	N.S.W.	...	...	Wood	...	Thin, semicircular, whitish or ochrey, roughened with very minute elevations.
796	...	...	...	V.	...	...	...	Trunks	...	Overlapping, cap reflexed, tow-like, leathery, soft, downy, white, not zoned.
797	...	...	...	...	...	Q.	...	Trunks	...	Spread out, attached; margin reflexed, incurved, pale, faintly linearly zoned, downy, then smooth.
798	...	...	...	V.	...	...	...	Trunks	...	Sessile, semicircular and overlapping in tufts or circular; soft when moist, hard when dry, chestnut brown.
799	...	...	...	...	...	Q.	...	Trunks	...	Rust coloured, membranous, rigid, radiately rough, shell shaped; margin crisped.
800	...	...	...	...	...	Q.	...	Old wood	...	Caps very numerous, minute, finally sessile, closely overlapping like scales, whitish yellow, brown when dry.
801	...	...	T.	...	...	...	...	Trunks	...	Overlapping, semicircular, somewhat zoned, thin, leathery, rusty to blackish brown, velvety.
802	...	...	...	...	...	Q.	...	Trunks	...	Corky to leathery, thin, kidney or fan shaped, pale fawn colour, thickly zoned, silky, and shining.
803	W.A.	...	T.	V.	...	...	...	Ground	...	Cinnamon. Cap leathery to membranous, fibrously radiate. Stem cylindrical, velvety, bulbous.
804	...	...	...	...	...	Q.	...	Bark	...	Broad, of one colour, leathery, soft, thin, elastic, reflexed, semicircular, downy, concentrically furrowed.
805	...	...	...	...	N.S.W.	Q.	...	Wood	...	Solid, woody to leathery, thin, zoned, variegated with brown, velvety at first, then naked, often shining.
806	...	...	...	...	...	Q.	...	Wood	...	Cap kidney shaped, thin, black, radiately rough, obscurely zoned. Stem of same colour, velvety, brown downwards.
807	...	...	...	V.	...	Q.	...	Trunks	...	Overlapping, thin, leathery, rigid, circular or kidney shaped, lobed, plaited, zoned, brown, silky.
808	W.A.	S.A.	T.	V.	N.S.W.	Q.	...	Trunks, &c.	...	Corky, somewhat zoned, rough, downy, becoming smooth, vermilion.
809	...	S.A.	...	...	...	...	B.	Woods, amongst moss	...	Bright cinnamon without and within. Cap zoned. Stem velvety.
810	...	...	...	...	...	...	...	Rotten wood	...	Narrow, spoon shaped, or split and fan shaped, lemon yellow, zoned a little.
811	...	...	...	...	...	Q.	...	Trunks	...	Widely extended with margins broadly reflexed, thin, bright ochre, flexible, fringed with nearly erect hairs.
812	...	...	...	...	...	...	...	Wood	...	Thin, leathery, copper coloured, silky, shining, radiately rough, thickly zoned.
813	...	...	...	V.	...	...	...	Trunks	...	Overlapping, confluent, base wedge shaped, sessile, fan shaped, lobed, slightly zoned, tan coloured.
814	...	...	...	V.	...	Q.	...	Dead leaves	...	Wedge shaped, rounded in front and lobed, tapering behind, thin, leathery, downy, pale ochre.
814A	...	...	...	...	N.S.W.	Q.	...	Wood	...	Cap rigid, spoon shaped, densely streaked, silky at first, then minutely warted, stem short, disc shaped.
815	...	...	...	...	N.S.W.	Q.	...	Branches	...	White, attached, cottony; margin slightly reflexed.
816	...	S.A.	...	V.	...	Q.	...	Trunks of <i>Eucalyptus</i>	...	Fleshy to corky (deformed), velvety, soft to the touch, zoneless, opaque, varying in colour from umber or bay to violet.
817	...	...	...	...	...	Q.	...	Trunks	...	Leathery, spread out, reflexed, radiately hispid, slightly concentrically zoned, thin, ochrey olive.
818	W.A.	...	T.	V.	N.S.W.	Q.	...	Trunks	...	Corky to leathery, with long hairs, zoned, brown, becoming hoary.
819	...	...	...	V.	N.S.W.	Q.	...	Wood	...	Cap leathery to membranous, zoned, covered with dingy evanescent down, somewhat bay brown. Stem very short, lateral.
820	...	...	...	...	...	Q.	...	Trunks	...	Pale bay, somewhat fan shaped, laterally growing together, thin, leathery, zoned, downy.
821	...	...	T.	...	...	...	...	Trunks	...	Leathery, thin, fan shaped, expanded from narrow base, densely concentrically streaked, silky, yellowish tan.
822	...	...	...	...	...	Q.	...	Trunks	...	Fibrously spongy, sessile, shell shaped, rusty, entirely resolved into rigid, much branched, cord-like fibres.
823	...	...	...	...	...	Q.	...	Trunks	...	Leathery, thin, rigid, flattened, shell shaped, slightly downy, greyish fawn colour, linearly zoned.
824	...	...	...	...	N.S.W.	...	...	Trunks	...	Fleshy to leathery, rigid, semicircular, sessile, shell shaped, zoned, hispid and rough to the touch, fawn to brownish.
825	...	...	...	...	...	Q.	...	Trunks	...	Semicircular or somewhat reniform, shell shaped, delicately downy, zoned, mouse or olive coloured.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
62. POLYSTICTUS.—Fries. Nov. Symb. 54 (1851).—					
826	801	VI. 5875	<i>P. Hasskarlii</i> ...	<i>Cooke, Grev. XIV. 86 (1886)</i> ...	Hasskarl's polystictus ...
827	777	„ 5760	<i>P. hirsutus</i> ...	<i>Cooke, Grev. XIV. 83 (1886)</i> ...	Hirsute polystictus ...
828	760	„ 5689	<i>P. hololeucus</i> ...	<i>Cooke, Grev. XIV. 81 (1886)</i> ...	Entirely white polystictus ...
829	784	„ 5784	<i>P. hypothejus</i> ...	<i>Cooke, Grev. XIV. 83 (1886)</i> ...	Under-yellow polystictus ...
830	775	„ 5755	<i>P. illotus</i> ...	<i>Cooke, Grev. XIV. 83 (1886)</i> ...	Dirty polystictus ...
831	742	„ 5591	<i>P. intonsus</i> ...	<i>Cooke, Grev. XIV. 79 (1886)</i> ...	Unshorn polystictus ...
832	748	„ 5639	<i>P. laceratus</i> ...	<i>Cooke, Grev. XIV. 80 (1886)</i> ...	Torn polystictus ...
833	811	„ 5933	<i>P. latus</i> ...	<i>Cooke, Grev. XIV. 87 (1886)</i> ...	Broad polystictus ...
834	757	„ 5663	<i>P. leonotis</i> ...	<i>Cooke, Grev. XIV. 80 (1886)</i> ...	Leonine polystictus ...
835	745	„ 5630	<i>P. libum</i> ...	<i>Cooke, Grev. XIV. 79 (1886)</i> ...	Cake polystictus ...
836	767	„ 5708	<i>P. lilacino-gilvus</i> ...	<i>Cooke, Grev. XIV. 82 (1886)</i> ...	Lilac-yellow polystictus ...
837	781	„ 5777	<i>P. limbatus</i> ...	<i>Cooke, Grev. XIV. 83 (1886)</i> ...	Bordered polystictus ...
838	725	„ 5538	<i>P. luteo-nitidus</i> ...	<i>Cooke, Grev. XIV. 77 (1886)</i> ...	Bright-yellow polystictus ...
839	800	„ 5870	<i>P. luteo-olivaceus</i> ...	<i>Cooke, Grev. XIV. 86 (1886)</i> ...	Olive-yellow polystictus ...
840	737	„ 5577	<i>P. luteus</i> ...	<i>Cooke, Grev. XIV. 78 (1886)</i> ...	Yellow polystictus ^a ...
841	753	„ 5644	<i>P. multilobus</i> ...	<i>Cooke, Grev. XIV. 80 (1886)</i> ...	Many-lobed polystictus ...
842	736	„ 5574	<i>P. mutabilis</i> ...	<i>Cooke, Grev. XIV. 78 (1886)</i> ...	Changeable polystictus ...
843	739	„ 5582	<i>P. nephridius</i> ...	<i>Cooke, Grev. XIV. 78 (1886)</i> ...	Kidney-shaped polystictus ...
844	728	„ 5545	<i>P. oblectans</i> ...	<i>Cooke, Grev. XIV. 77 (1886)</i> ...	Alluring polystictus ...
845	776	„ 5758	<i>P. obstinatus</i> ...	<i>Cooke, Grev. XIV. 83 (1886)</i> ...	Hard polystictus ...
846	794	„ 5843	<i>P. occidentalis</i> ...	<i>Cooke, Grev. XIV. 85 (1886)</i> ...	Western polystictus ...
847	752	„ 5642	<i>P. ornithorhyuchi</i> ...	<i>Cooke, Grev. XIV. 80 (1886)</i> ...	Ornithorhynchus polystictus ...
848	730	„ 5548	<i>P. parvulus</i> ...	<i>Cooke, Grev. XIV. 77 (1886)</i> ...	Very small polystictus ...
849	789	„ 5820	<i>P. peradeniæ</i> ...	<i>Cooke, Grev. XIV. 84 (1886)</i> ...	Peradenia polystictus ...
850	726	„ 5543	<i>P. perennis</i> ...	<i>Cooke, Grev. XIV. 77 (1886)</i> ...	Perennial polystictus ...
851	...	„ 5695	<i>P. pergamenus</i> ...	<i>Cooke, Grev. XIV. 81 (1886)</i> ...	Parchment polystictus ...
852	744	„ 5623	<i>P. peroxydatus</i> ...	<i>Cooke, Grev. XIV. 79 (1886)</i> ...	Peroxide polystictus ...
853	791	„ 5832	<i>P. Persoonii</i> ...	<i>Cooke, Grev. XIV. 85 (1886)</i> ...	Persoon's polystictus ...
854	783	„ 5781	<i>P. pinsitus</i> ...	<i>Cooke, Grev. XIV. 83 (1886)</i> ...	Crushed polystictus ...
855	734	„ 5571	<i>P. porphyrites</i> ...	<i>Cooke, Grev. XIV. 78 (1886)</i> ...	Purple-coloured polystictus ...
856	761	„ 5693	<i>P. proteiformis</i> ...	<i>Cooke, Grev. XIV. 81 (1886)</i> ...	Proteus-like polystictus ...
857	731	IX. 730	<i>P. quadrans</i> ...	<i>Cooke, Grev. XIV. 78 (1886)</i> ...	Quadrangle polystictus ...
858	771	VI. 5717	<i>P. radiatus</i> ...	<i>Cooke, Grev. XIV. 82 (1886)</i> ...	Radiate polystictus ...
859	755	„ 5648	<i>P. radiato-rugosus</i> ...	<i>Cooke, Grev. XIV. 80 (1886)</i> ...	Radiately-rough polystictus ...
860	747	„ 5634	<i>P. rasipes</i> ...	<i>Cooke, Grev. XIV. 79 (1886)</i> ...	Rough-stalked polystictus ...
861	795	„ 5844	<i>P. rigens</i> ...	<i>Sacc. and Cub., Syll. Fung. VI. (1888)</i> ...	Stiff polystictus ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Trametes, Polyporus, Boletus—continued.									
826	...	...	...	...	...	Q.	...	Trunks ...	Corky, overlapping, zoned, velvety, bay brown.
827	...	...	...	V.	N.S.W.	Q.	B.	Trunks ...	Corky to leathery, hirsute, with rigid hairs, zoned with concentric furrows, whitish to tawny. Common.
828	...	...	...	V.	...	...	...	Trunks ...	Wholly white, loosely corky, semicircular, sessile, concentrically furrowed, flesh tinder-like.
829	...	...	...	...	N.S.W.	...	...	Trunks ...	Thin, leathery, shell shaped, narrow at base, with silky hairs, white, zoned. Pores becoming bright yellow.
830	...	...	...	V.	...	...	...	Trunks ...	Leathery to membranous, softly velvety, concentrically furrowed, dingy grey, turning brownish.
831	...	...	T.	...	...	...	...	Rotten wood	Small. Cap fan shaped, thin, velvety, brown. Stem excentric, short.
832	...	...	...	...	N.S.W.	Q.	...	Branches	Spread out and reflexed, thin, zoned, rough, streaked, wood colour.
833	...	...	T.	...	...	...	...	Branches	Inverted, obscurely zoned, corky to leathery, dingy brown, with velvety bloom.
834	...	...	...	...	...	...	...	Trunks ...	Spongy to fleshy, overlapping and grown together, rough haired, zoneless, dark rusty brown.
835	...	...	...	...	N.S.W.	Q.	...	Wood ...	Leathery, polished, fixed behind by a disc, lobed, somewhat zoned, white; margin yellowish.
836	W.A.	S.A.	T.	V.	...	Q.	...	Rotten wood	Somewhat overlapping, corky to leathery, delicate, rough, more or less zoned, brown when old.
837	...	...	...	V.	...	...	...	Trunks ...	Leathery, thin, somewhat velvety, concentrically furrow-zoned, becoming rusty.
838	...	...	...	...	...	Q.	...	Ground ...	Cap rough, yellow, silky, shining, irregularly lobed, thickly zoned, rather velvety at first. Stem deformed, with spongy coating.
839	...	...	...	...	...	Q.	...	Trunks ...	Woody, rigid, sessile, thin, downy, deeply and concentrically zoned, warted; substance olive yellow.
840	...	...	...	...	N.S.W.	Q.	...	Wood ...	Cap thin, rigid, leathery, fan shaped, yellowish. Stem marginal, dilated at base, yellowish.
841	...	...	...	...	N.S.W.	...	...	Trunks ...	Leathery, thin, rigid, somewhat kidney shaped, narrowed into lateral and very short stem, white to tan; margin lobed.
842	...	...	...	...	...	Q.	...	Wood ...	Cap leathery, rigid, fan or spoon shaped, zoned, yellowish white, narrowed down into stem, which is lateral and variable in length.
843	...	...	...	...	...	Q.	...	Branches	Small, thin. Cap veined, smooth, kidney shaped, bay brown. Stem very short, black.
844	W.A.	S.A.	T.	V.	...	Q.	...	Sandy soil	Cap thin, leathery, cut, zoned about centre, shining, bright cinnamon. Stem velvety, reddish brown.
845	...	...	...	...	...	Q.	...	Trunks ...	Leathery to woody, hardening, thin, somewhat ash coloured, velvety, variegated with narrow zones.
846	...	S.A.	...	V.	N.S.W.	Q.	...	Trunks ...	Corky to leathery, spread out and reflexed, concentrically furrowed, yellowish tan, becoming pale.
847	...	...	...	...	N.S.W.	...	...	Trunks ...	Somewhat tufted, thin, leathery, wedge shaped, zoneless, hairy to downy, rusty umber, tapering into short or obsolete stem of same colour.
848	...	S.A.	...	V.	...	...	...	Ground ...	Cap leathery to membranous, obsoletely silky, zoned, bay brown. Stem thin, tuberous, velvety.
849	...	...	...	...	...	Q.	...	Wood ...	Semicircular, laterally running together, somewhat zoned, silky, membranous, olive.
850	...	...	...	...	...	Q.	B.	Ground ...	Cap leathery, funnel shaped, velvety, zoned, cinnamon to bay brown. Stem firm, thickened downwards, velvety.
851	...	...	...	...	...	Q.	...	Trunks ...	Leathery to membranous, rigid, downy, furrowed concentrically, white.
852	...	...	...	...	N.S.W.	...	...	Trunks ...	Thin, nearly circular, rusty, powdery. Cap slightly zoned, rough. Stem short, thick.
853	...	...	...	V.	N.S.W.	Q.	...	Wood ...	Leathery, flattened, obsoletely zoned, dark blood red, becoming pale.
854	...	...	...	...	...	Q.	...	Wood ...	Leathery to membranous, tough, hairy, concentrically furrowed, ash coloured.
855	...	...	...	...	...	Q.	...	Rotting branches ...	Thin, leathery. Cap fan shaped, with ochrey zones, shining brown, becoming purplish. Stem short, of same colour.
856	...	...	...	V.	...	...	...	Trunks ...	Spread out and reflexed, tow-like, leathery, white within, slightly concentrically furrowed. Very variable at different stages.
857	...	...	...	...	...	Q.	...	Wood ...	Cap rigid, smooth, thin, furrowed, zoned, dark ochre. Stem short, excentric, of same colour.
858	...	...	...	V.	...	Q.	B.	Trunks ...	Corky, leathery, rigid, radiately rough, velvety at first, brown, then rusty brown.
859	...	...	T.	...	...	...	...	Trunks ...	Thickly overlapping, thin, radiately rough, dingy white or grey.
860	...	...	...	...	...	Q.	...	Trunks ...	Cap fan and shell shaped, silky lineate, somewhat velvety, zoned, reddish brown when dry. Stem short, flattened, rather hispid.
861	...	...	...	...	N.S.W.	Q.	...	Trunks ...	Spread out, shortly reflexed, often run together, leathery, rigid, more or less concentrically furrowed, velvety, pale tawny to wood colour.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
62. POLYSTICTUS.—Fries, Nov. Symb. 54 (1851).—					
862	...	VI. 5831	<i>P. rigidus</i> ...	<i>Cooke, Grev. XIV. 85 (1886)</i> ...	Rigid polystictus ...
863	746	„ 5631	<i>P. sanguineus</i> ...	<i>Cooke, Grev. XIV. 79 (1886)</i> ...	Blood-red polystictus ...
864	788	„ 5811	<i>P. scorteus</i> ...	<i>Cooke, Grev. XIV. 84 (1886)</i> ...	Leathery polystictus ...
865	762	„ 5696	<i>P. seriatus</i> ...	<i>Cooke, Grev. XIV. 81 (1886)</i> ...	Seriate polystictus ...
866	741	„ 5585	<i>P. stereinus</i> ...	<i>Berk. and Curt., Linn. Journ. X. 308 (1869)</i> ...	Solid polystictus ...
867	787	„ 5808	<i>P. stereoides</i> ...	<i>Cooke, Grev. XIV. 78 (1886)</i> ...	Stereum-like polystictus ...
868	802	„ 5876	<i>P. tabacius</i> ...	<i>Cooke, Grev. XIV. 86 (1886)</i> ...	Tobacco-coloured polystictus ...
869	797	„ 5847	<i>P. tephroleucus</i> ...	<i>Cooke, Grev. XIV. 85 (1886)</i> ...	Pale-ashy polystictus ...
870	724	„ 5535	<i>P. tomentosus</i> ...	<i>Cooke, Grev. XIV. 77 (1886)</i> ...	Downy polystictus ...
871	786	„ 5787	<i>P. trizonatus</i> ...	<i>Cooke, Grev. XIV. 84 (1886)</i> ...	Three-zoned polystictus ...
872	782	„ 5779	<i>P. vellerens</i> ...	<i>Cooke, Grev. XIV. 83 (1886)</i> ...	Woolly polystictus ...
873	778	„ 5763	<i>P. velutinus</i> ...	<i>Cooke, Grev. XIV. 83 (1886)</i> ...	Velvety polystictus ...
874	772	„ 5732	<i>P. venustus</i> ...	<i>Cooke, Grev. XIV. 82 (1886)</i> ...	Graceful polystictus ...
875	790	„ 5827	<i>P. vernicifluus</i> ...	<i>Cooke, Grev. XIV. 84 (1886)</i> ...	Varnished polystictus ...
876	766	„ 5704	<i>P. versatilis</i> ...	<i>Cooke, Grev. XIV. 82 (1886)</i> ...	Changing polystictus ...
877	774	„ 5741	<i>P. versicolor</i> ...	<i>Cooke, Grev. XIV. 83 (1886)</i> ...	Variously-coloured polystictus ...
878	793	„ 5838	<i>P. vinosus</i> ...	<i>Cooke, Grev. XV. 51 (1886)</i> ...	Vinous polystictus ...
879	732	„ 5565	<i>P. xanthopus</i> ...	<i>Cooke, Grev. XIV. 78 (1886)</i> ...	Yellow-stalked polystictus ...
880	804	„ 5883	<i>P. xerampelinus</i> ...	<i>Cooke, Grev. XIV. 86 (1886)</i> ...	Purplish-umber polystictus ...
881	...	„ 5771	<i>P. zonatus</i> ...	<i>Cooke, Grev. XIV. 83 (1886)</i> ...	Zoned polystictus ...
63. PORIA.—Pers. Syn. 542 (1801).—					
882	831	VI. 6062	<i>P. aprica</i> ...	<i>Cooke, Grev. XIV. 112 (1886)</i> ...	Exposed poria ...
883	840	„ 6153	<i>P. Archeri</i> ...	<i>Cooke, Grev. XIV. 115 (1886)</i> ...	Archer's poria ...
884	826	„ 5995	<i>P. atro-vinosa</i> ...	<i>Cooke, Grev. XIV. 110 (1886)</i> ...	Dark-vinous poria ...
885	817	„ 5948	<i>P. calcea</i> ...	<i>Cooke, Grev. XIV. 109 (1886)</i> ...	Chalky-white poria ...
886	820	„ 5964	<i>P. callosa</i> ...	<i>Cooke, Grev. XIV. 110 (1886)</i> ...	Thick-skinned poria ...
887	836	„ 6126	<i>P. contigua</i> ...	<i>Cooke, Grev. XIV. 114 (1886)</i> ...	Contiguous poria ...
888	833	„ 6093	<i>P. corticola</i> ...	<i>Cooke, Grev. XIV. 113 (1886)</i> ...	Bark-growing poria ...
889	838	„ 6131	<i>P. dictyopora</i> ...	<i>Cooke, Grev. XII. 17 (1884)</i> ...	Net-pored poria ...
890	823	„ 5982	<i>P. epilintea</i> ...	<i>Cooke, Grev. XIV. 110 (1886)</i> ...	Fibrous poria ...
891	839	„ 6138	<i>P. faticens</i> ...	<i>Berk. and Rav., Grev. I. 65 (1872)</i> ...	Cracking poria ...
892	835	„ 6123	<i>P. ferruginosa</i> ...	<i>Cooke, Grev. XIV. 114 (1886)</i> ...	Rusty poria ...
893	814	„ 5998	<i>P. hyalina</i> ...	<i>Cooke, Grev. XIV. 109 (1886)</i> ...	Hyaline poria ...
894	824	„ 5984	<i>P. hyposclera</i> ...	<i>Cooke, Grev. XIV. 110 (1886)</i> ...	Hard poria ...
895	828	„ 6004	<i>P. livida</i> ...	<i>Cooke, Grev. X. 131 (1882)</i> ...	Livid poria ...
896	816	„ 5947	<i>P. medulla-panis</i> ...	<i>Cooke, Grev. XIV. 109 (1886)</i> ...	Pith-of-bread poria ...
897	830	„ 6057	<i>P. membranincta</i> ...	<i>Berk. Grev. XV. 26 (1886)</i> ...	Membrane-girt poria ...
898	832	„ 6068	<i>P. merulina</i> ...	<i>Cooke, Grev. XIV. 112 (1886)</i> ...	Merulius poria ...
899	813	„ 5936	<i>P. mollusca</i> ...	<i>Cooke, Grev. XIV. 109 (1886)</i> ...	Soft poria ...
900	818	„ 5949	<i>P. niphodes</i> ...	<i>Cooke, Grev. XIV. 109 (1886)</i> ...	Snow-like poria ...
901	837	„ 6130	<i>P. orbicularis</i> ...	<i>Cooke, Grev. XIV. 114 (1886)</i> ...	Circular poria ...
902	821	„ 5969	<i>P. parilis</i> ...	<i>Cooke, Grev. XIV. 110 (1886)</i> ...	Equal poria ...
903	825	„ 5994	<i>P. rufa</i> ...	<i>Cooke, Grev. XIV. 110 (1886)</i> ...	Red poria ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Trametes, Polyporus, Boletus—continued.									
862	...	...	...	...	...	Q.	...	Trunks	Leathery to corky, wedge shaped, light, lurid, brown zoned.
863	W.A.	S.A.	T.	V.	N.S.W.	Q.	...	Trunks	Vermilion. Cap leathery, thin, kidney shaped, shining, obsoletely concentrically furrowed. Stem lateral, short, circularly dilated at base.
864	...	...	...	...	N.S.W.	...	...	Trunks	Leathery, flattened on both sides, concentrically furrowed and zoned, shaggy haired, verdigris to grey.
865	...	...	...	V.	...	...	...	Trunks	Tow-like, semicircular, rigid, mostly run together in serial order, concentrically furrowed, velvety, brick red or yellowish tan.
866	...	...	...	...	N.S.W.	Q.	...	Trunks	Fan shaped, rigid when dry and inflexed. Cap thin, many zoned, chestnut red. Stem disc shaped.
867	...	...	...	V.	N.S.W.	Q.	...	Trunks	Leathery, thin, rigid, spread out and reflexed, downy, then smooth, grey brown.
868	...	...	...	...	N.S.W.	...	...	Trunks	Overlapping, rusty to bay, leathery, thin, rigid, spread out and reflexed, shell shaped, downy, concentrically zoned.
869	...	...	...	V.	...	...	...	Wood	Semicircular, rigid, leathery, white, zoned, velvety, with rough hairs.
870	...	...	...	V.	...	Q.	...	Ground	Corky, hard, deformed, zoneless, persistently downy, as well as the unequal stem, brown.
871	...	...	...	V.	...	Q.	...	Trunks	Leathery, thin, rigid, ochrey, flattened, silky, variegated with three elevated darker zones.
872	...	...	...	...	...	Q.	...	Trunks	Semicircular, leathery, soft, thin, white, deusely silky haired, zones darker, very narrow.
873	...	...	...	V.	...	Q.	B.	Trunks	Corky to leathery, velvety, soft, slightly zoned, white, at length yellowish.
874	W.A.	...	...	...	...	Q.	...	Trunks	Overlapping, forming elongated tufts, reflexed, leathery, zoned, whitish, downy, tufts in front.
875	...	...	T.	...	...	Q.	...	Rotten wood	Semicircular, somewhat fan shaped, thin, varnished and polished, reddish brown.
876	...	...	...	...	N.S.W.	Q.	...	Rotten wood	Very long, spread out, thin, whitish, flexible; margin broadly reflexed.
877	...	...	T.	V.	N.S.W.	Q.	B.	Trunks	Leathery, thin, rigid, flattened, velvety, shining, variegated with other coloured zones.
878	...	...	...	...	...	Q.	...	Rotten wood	Kidney shaped, thin, zoned, delicately velvety, turning smooth, dark wine colour.
879	...	...	...	V.	N.S.W.	Q.	...	Branches	Cap leathery, papery, funnel shaped, zoned, shining, bay brown. Stem short, shining, yellowish.
880	...	...	...	...	...	Q.	...	Trunks	Corky to leathery, densely overlapping, shell shaped, shaggy, becoming smooth, with concentric furrowed zones, purplish number.
881	...	...	...	...	...	Q.	...	Trunks	Corky to leathery, convex, tubercled and bulgiug behind, somewhat zoned, shaggy.
Polyporus, Boletus, Corticium.									
882	...	...	T.	...	...	...	...	Wood	Inverted, spread out, loosely adhering, pale fawn.
883	...	...	T.	...	...	...	...	Rotten wood	Inverted, spread out, leathery to membranous; margin downy.
884	...	...	...	V.	N.S.W.	...	...	Trunks	Spread out, undefined, dark vinous purple; margin powdery to downy.
885	...	...	...	...	...	Q.	...	Wood	Chalky white. Inverted, spread out; margin very thin and membranous.
886	...	...	...	...	...	Q.	B.	Wood	Widely expanded, tough, entire, separable like soft leather, white.
887	...	S.A.	...	V.	N.S.W.	Q.	B.	Old wood	Spread out, thick, firm, cinnamon when young, margin at first shaggy.
888	...	S.A.	...	V.	...	Q.	...	Rotting bark	Widely expanded, firm, white.
889	...	...	...	V.	...	Q.	...	Burnt wood	Spread out, undefined, thin, white, incrusting.
890	...	...	...	...	N.S.W.	...	...	Trunks	Inverted, spread out; margin and substance fibrous.
891	...	...	...	...	...	Q.	...	Wood	Inverted, very thin, powdery, white.
892	W.A.	...	...	V.	N.S.W.	Q.	...	Wood, on posts	Spread out, thick, firm, unequal, tawny, then rusty bay.
893	...	...	T.	...	...	...	...	Wood	Inverted, white, hyaline, thin, margin downy.
894	...	...	...	V.	...	Q.	...	Trunks	Spread out, rather thick, margin thin, and pale ochrey or tinged with flesh colour.
895	...	...	...	...	N.S.W.	...	...	Bark	Spread out, defined, crustaceous, livid to sooty brown.
896	...	...	...	...	...	Q.	B.	Old wood	Spread out, defined, somewhat wavy, firm, smooth, white.
897	...	...	T.	...	...	...	...	Wood	Spread out, thin, pale ochrey, seated on a paler membranous layer.
898	...	...	T.	...	...	...	...	Wood	Inverted, expanded, orange.
899	...	S.A.	...	V.	...	...	B.	Old wood. Dead bark of <i>Eucalyptus obliqua</i>	Expanded, thin, soft, white, circumference of finely filamentous texture.
900	...	...	...	...	N.S.W.	...	...	Wood	Inverted, snow white, margin very narrow.
901	...	...	T.	...	...	...	...	Living bark	Exactly circular, dark brown, margin membranous with rough down.
902	W.A.	...	...	...	...	...	...	Bark	Longitudinally expanded, closely attached, run together, dry, yellow, becoming pale.
903	...	...	...	V.	...	...	B.	Branches	Expanded, leathery, thin, attached, smooth, definite outline, blood red.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
63. PORIA.—Pers. Syn. 542 (1801).—					
904	828 bis	VI. 6029	<i>P. ruftineta</i> ...	Berk. and Curt., Grev. XV. 25 (1886) ...	Reddish poria ...
905	834	„ 6095	<i>P. sinuosa</i> ...	Cooke, Grev. XIV. 113 (1886) ...	Wavy poria ...
906	815	„ 5942	<i>P. subvineta</i> ...	Cooke, Grev. XIV. 109 (1886) ...	Somewhat-bound poria ...
907	819	„ 5952	<i>P. tarda</i> ...	Cooke, Grev. XIV. 109 (1886) ...	Slowly-developing poria ...
908	829	„ 6035	<i>P. vaporaria</i> ...	Cooke, Grev. XIV. 111 (1886) ...	Sweating poria ...
909	827	„ 6003	<i>P. victoriae</i> ...	Cooke, Grev. XIV. 111 (1886) ...	Victorian poria ...
910	822	„ 5983	<i>P. vineta</i> ...	Cooke, Grev. XIV. 110 (1886) ...	Bound poria ...
911	812	„ 5935	<i>P. vulgaris</i> ...	Fries, S.M. I. 381 (1832) ...	Common poria ...
64. TRAMETES.—Fries,					
912	859	VI. 6273	<i>T. Curreyi</i> ...	Cooke, Trans., Bot. Soc. Ed. 157 (1878) ...	Currey's trametes ...
913	848	„ 6197	<i>T. devexa</i> ...	Berk., Linn. Journ. XIII. 165 (1873) ...	Sloping trametes ...
914	854	„ 6240	<i>T. epitephra</i> ...	Berk., Linn. Journ. XIII. 165 (1873) ...	Ash-coloured trametes ...
915	852	„ 6220	<i>T. fibrosa</i> ...	Fries, Epicr. 490 (1838) ...	Fibrous trametes ...
916	861	„ ...	<i>T. gansapata</i> ...	Cooke, Grev. XV. 55 (1886) ...	Friezed trametes ...
917	846	„ 6188	<i>T. heteromalla</i> ...	Cooke, Grev. X. 132 (1882) ...	Shaggy trametes ...
918	853	„ 6235	<i>T. hispidula</i> ...	Berk. and Curt., Linn. Journ. X. 319 (1869) ...	Rough trametes ...
919	849	„ 6204	<i>T. lactinea</i> ...	Berk., Ann. Nat. Hist. X. 373 (1842) ...	Milk-white trametes ...
920	850	„ 6205	<i>T. laevis</i> ...	Berk., Hook., Lond. Journ. VI. 507 (1847) ...	Even trametes ...
921	858	„ 6264	<i>T. mollis</i> ...	Fries, Hym. Eur. 585 (1874) ...	Soft trametes ...
922	842	„ 6181	<i>T. Muelleri</i> ...	Berk., Linn. Journ. X. 320 (1869) ...	Mueller's trametes ...
923	856	„ 6247	<i>T. ochroflava</i> ...	Cooke, Grev. IX. 12 (1880) ...	Ochrey-yellow trametes ...
924	847	„ 5236	<i>T. ochroleuca</i> ...	Cooke, Grev. XIX. 99 (1891) ...	Ochrey-white trametes ...
925	841	„ 6170	<i>T. phellina</i> ...	Berk., Linn. Journ. XIII. 164 (1873) ...	Corky trametes ...
926	843	IX.p. 198	<i>T. picta</i> ...	Berk. and Br., Linn. Trans. II. 61 (1883) ...	Ornamented trametes ...
927	851	VI. 6213	<i>T. pini</i> ...	Fries, S.M. I. 336 (1821) ...	Pine trametes ...
928	855	„ 6241	<i>T. pyrochreas</i> ...	Berk., Linn. Journ. XIII. 164 (1873) ...	Fleshy trametes ...
929	857	„ 6249	<i>T. scrobiculata</i> ...	Berk., Grev. VI. 70 (1877) ...	Pitted trametes ...
930	862	„ 6267	<i>T. serpens</i> ...	Fries, Hym. Eur. 586 (1874) ...	Spreading trametes ...
931	845	„ 6185	<i>T. Sprucei</i> ...	Berk., Hook., Journ. 236 (1856) ...	Spruce's trametes ...
932	860	„ 5222	<i>T. ungulata</i> ...	Berk., Linn. Journ. XIII. 165 (1873) ...	Hoof-shaped trametes ...
933	844	„ 6183	<i>T. versiformis</i> ...	Berk., Linn. Journ. XIV. 56 (1875) ...	Variously-shaped trametes ...
65. SCLERODEPSIS.—Cooke,					
934	863	VI. 6237 IX.p. 194	<i>S. colliculosa</i> ...	Cooke, Grev. XIX. 49 (1890) ...	Hillocky sclerodepsis ...
66. HEXAGONIA.—Fries,					
935	883	VI. 6278	<i>H. crinigera</i> ...	Fries, Epicr. 496 (1838) ...	Hair-bearing hexagonia ...
936	894	„ 6333	<i>H. decipiens</i> ...	Berk., Linn. Journ. XIII. 166 (1873) ...	Deceptive hexagonia ...
937	890	„ 6320	<i>H. discolor</i> ...	Fries, Nov. Symb. 102 (1851) ...	Discoloured hexagonia ...
938	884	„ 6290	<i>H. durissima</i> ...	Berk. and Br., Linn. Journ. XIV. 57 (1875) ...	Very hard hexagonia ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Polyporus, Boletus, Corticium—continued.									
904	...	...	...	...	...	...	...	Wood ...	Expanded, rather thick, firm, tawny, rusty bay when old, margin woolly, golden brown.
905	...	...	...	...	N.S.W.	...	...	Wood and bark ...	Broadly expanded, attached, dry, springing from temporary mycelium, white to yellowish. Pores wavy.
906	...	...	T.	...	...	...	...	Wood ...	Rather thick, widely expanded, white, separable.
907	W.A.	...	...	V.	...	...	...	Wood ...	White, then ochrey, mycelium waxy, like <i>Corticium</i> ; margin narrow, downy. Pores slowly developed.
908	W.A.	...	T.	V.	...	Q.	B.	Creeping upon rotten wood, &c.	Expanded, mycelium creeping in the wood, woolly, brown or white. Very common.
909	...	...	...	V.	...	Q.	...	Trunks ...	Smoky colour, expanded, thin.
910	...	...	...	V.	...	Q.	...	Rotten wood ...	Inverted, thick in centre; margin thin, tinged above with red.
911	...	S.A.	...	V.	...	Q.	B.	Wood and branches	Broadly expanded, thin, closely attached and difficult to remove, dry, even, whitish. Very common.

Epier. 488 (1838).—Polyporus, Dædalea, Boletus.

912	...	...	...	...	N.S.W.	Q.	...	Trunks ...	Expanded, reflexed, lobed, membranous to leathery, rusty umber.
913	...	...	...	...	N.S.W.	Q.	...	Trunks ...	Woody, somewhat hoof-shaped, sloping behind, cap velvety, somewhat tawny.
914	...	S.A.	...	...	...	...	...	Trunks ...	Hoof-like, sloping behind, zoned, brown, somewhat rough; margin white.
915	...	S.A.	...	...	...	...	...	Trunks ...	Corky, thin, somewhat wavy, zoned, dark-brown, rough, with thickly grown branched fibres.
916	...	S.A.	...	...	...	Q.	...	Trunks ...	Spread out behind, reflexed, velvety, zoned, bright umber, leathery.
917	...	...	...	...	N.S.W.	...	...	Trunks ...	Corky, softish, shaggy, concentrically furrowed, whitish, then somewhat ashy.
918	...	S.A.	...	...	...	...	...	Wood ...	Small, hoof-shaped, rusty umber, rough behind, margin somewhat downy.
919	...	S.A.	...	...	N.S.W.	Q.	...	Wood ...	Sessile, irregular, thickish, hard, rigid, zoneless, corky, velvety, warty, milk white.
920	...	...	...	...	N.S.W.	Q.	...	Roots ...	Quite even, thick, hoof-shaped, pale-wood colour, delicately downy, somewhat zoned.
921	...	...	...	...	N.S.W.	...	B.	Branches ...	Inverted, distinct, somewhat membranous, pale wood-colour, finally brownish.
922	...	S.A.	...	V.	N.S.W.	Q.	...	Wood ...	Semicircular, corky, delicately downy, white, rough, margin lobed, concentrically furrowed.
923	...	...	...	...	...	Q.	...	Trunks ...	Entirely ochrey-yellow, often overlapping, corky, compact, tuberculose, concentrically zoned.
924	W.A.	...	T.	V.	N.S.W.	Q.	...	Bark ...	Hoof-shaped, corky, few-zoned, ochrey white, delicately downy at first; soon smooth and shining.
925	...	...	...	...	N.S.W.	...	...	Rotten wood ...	Corky, attached by circular disc, becoming whitish, rough, zoneless.
926	...	...	...	...	N.S.W.	Q.	...	Wood ...	Semicircular, corky, hard, smooth, pale, with darker concentric bands.
927	W.A.	...	...	...	...	...	B.	Trunks ...	Corky to woody, very hard, concentrically furrowed, cracked, rusty brown, turning blackish, odour slight.
928	...	...	...	...	N.S.W.	Q.	...	Trunks ...	Thickish, zoned, umber brown behind, at length velvety; substance compact but soft, tawny.
929	...	...	...	V.	...	...	...	Trunks ...	Ochrey, semicircular, slightly furrowed, pitted and dotted, substance corky.
930	...	...	...	...	...	Q.	B.	Bark ...	Dry, at first breaking through, tubercular, circular, white; margin distinct, downy.
931	...	...	...	V.	N.S.W.	Q.	...	Wood ...	Thickish, humped, obtuse, becoming white.
932	...	S.A.	...	...	...	...	...	Trunks ...	Hard, whitish, hoof-shaped, sloping behind, delicately downy.
933	...	...	...	...	...	Q.	...	Wood ...	White, reflexed, lobed, radiately rough, opaque, varying extremely in thickness.

Grev. XIX. 49 (1890).—Trametes.

934	...	...	...	...	N.S.W.	Q.	...	Wood ...	Sessile, semicircular, thin, hard, leathery, somewhat silky, zoned, ochrey.
-----	-----	-----	-----	-----	--------	----	-----	----------	---

Epier. 496 (1838).—Favolus, Polyporus.

935	...	...	...	...	...	Q.	...	Trunks ...	Corky to leathery, bristly, zoneless, brown, turning blackish.
936	W.A.	S.A.	...	...	V.	...	Q.	Trunks of <i>Casuarina</i>	Semicircular, zoned, umber, velvety to rough, furrowed. Species most distinct.
937	W.A.	...	...	...	...	...	...	Bark ...	Expanded, reflexed, unpolished, pale umber.
938	...	S.A.	...	V.	...	...	...	Wood and trunks	Hoof-shaped, rough, zoned with red and brown; substance rather fleshy.

SYSTEMATIC ARRANGEMENT

Number	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
66. HEXAGONIA.—Fries.					
939	887	VI. 6310	H. Gunnii ...	Berk., Hook., Lond. Journ. IV. 37 (1845) ...	Gunn's hexagonia ...
940	885	„ 6300	H. Muellerei ...	Berk., Linn. Journ. XIII. 166 (1873) ...	Mueller's hexagonia ...
941	892	„ 6325	H. polygramma ...	Cooke, Grev. XV. 58 (1886) ...	Many-lined hexagonia ...
942	888	„ 6315	H. rigida ...	Berk., Linn. Journ. XVI. 54 (1878) ...	Rigid hexagonia ...
943	886	„ 6306	H. sericea ...	Fries, Epier. 497 (1838) ...	Silky hexagonia ...
944	893	„ 6328	H. similis ...	Berk., Hook., Lond. Journ. V. 4 (1846) ...	Similar hexagonia ...
945	891	„ 6324	H. tenuis ...	Cooke, Grev. XV. 60 (1886) ...	Thin hexagonia ...
945A	891	...	H. tennis, var. sub-tenuis ...	Cooke, Grev. XIX. 103 (1891) ...	Thinnish hexagonia ...
946	889	VI. 6319	H. umbrinella ...	Fries, Fung. Nat. 17 (1848) ...	Umber hexagonia ...
947	882	„ 6274	H. Wightii ...	Cooke, Grev. XV. 60 (1886) ...	Wight's hexagonia ...
67. DÆDALEA.—Pers.					
948	866	VI. 6184	D. ambigua ...	Berk., Hook., Lond. Journ. IV. 305 (1845) ...	Ambiguous dædalea ...
949	874	„ 6374	D. aulacophylla ...	Berk., Linn. Journ. XIII. 166 (1873) ...	Furrow-leaved dædalea ...
950	880	„ 6409	D. Bowmanni ...	Berk., Linn. Journ. XIII. 166 (1873) ...	Bowman's dædalea ...
951	864	„ 6359	D. glabrescens ...	Berk., Linn. Journ. XVI. 39 (1878) ...	Smooth dædalea ...
952	876	„ 6382	D. Hobsoni ...	Berk., Linn. Journ. XIII. 165 (1873) ...	Hobson's dædalea ...
953	...	...	D. illudens ...	Cooke and Mass., Grev. XXI. 37 (1892) ...	Illusive dædalea ...
954	869	VI. 6363	D. incompta ...	Berk., Linn. Trans. II. 61 (1883) ...	Unadorned dædalea ...
955	865	„ 6361	D. intermedia ...	Berk., Linn. Journ. XVIII. 385 (1881) ...	Intermediate dædalea ...
956	877	„ 6404	D. latissima ...	Fries, S.M. I. 340 (1821) ...	Very broad dædalea ...
957	868	...	D. Muellerei ...	Berk., Grev. XIX. 93 (1891) ...	Mueller's dædalea ...
958	870	VI. 6364	D. scalaris ...	Berk. and Br., Linn. Trans. II. 61 (1883) ...	Ladder-like dædalea ...
959	872	„ 6368	D. Schomburgkii ...	Berk. in Cooke, Austr. Fun. 27 (1883) ...	Schomburgk's dædalea ...
960	878	„ 6406	D. sinuosa ...	Klotzsch, Linn. VIII. 482 (1833) ...	Flexuous dædalea ...
961	873	„ 6370	D. Sprucei ...	Berk., Hook., Journ. 236 (1856) ...	Spruce's dædalea ...
962	867	IX. 800	D. subcongener ...	Berk., Grev. XIX. 93 (1891) ...	Congeneric dædalea ...
963	879	VI. 6408	D. tasmanica ...	Sacc, Syll. VI. 384 (1888) ...	Tasmanian dædalea ...
964	871	„ 6367	D. tenuis ...	Berk., Hook., Lond. Journ. I. 151 (1842) ...	Thin dædalea ...
965	875	„ 6376	D. unicolor ...	Fries, S.M. I. 336 (1821) ...	One-coloured dædalea ...
68. CERIOMYCES.—Battarr.					
966	881	IX. 811	C. incomptus ...	Sacc, Bull. Soc. Myc. Fr. V. 115 (1880) ...	Unadorned ceriomyces ...
69. FAVOLUS.—Fries.					
967	896	VI. 6437	F. Boucheanus ...	Cooke, Grev. XV. 53 (1886) ...	Boucheanus favolus ...
968	900	„ 6465	F. hispidulus ...	Berk. and Curt., Linn. Journ. XIII. 167 (1873) ...	Hispid favolus ...
969	898	„ 6449	F. pusillus ...	Fries, Linn. V. 511 (1830) ...	Small favolus ...
970	899	„ 6460	F. Rhipidium ...	Cooke, Grev. XV. 54 (1886) ...	Fan-like favolus ...
971	897	„ 6439	F. scaber ...	Berk. and Br., Linn. Journ. XIV. 57 (1875) ...	Rough favolus ...
972	895	„ 6430	F. squamiger ...	Berk., Linn. Journ. XIII. 166 (1873) ...	Scale-bearing favolus ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						R.	Occurrence.	General Characters.	
	W.A.	S.A.	T.	V.	N.S.W.	Q.				
Epicr. 496 (1838).—Favolus, Polyporus—continued.										
939	W.A.	...	T.	V.	...	...	...	Bark, &c.	...	Sessile, somewhat hoof shaped, overlapping, wood coloured, slightly zoned towards the margin.
940	...	...	...	...	N.S.W.	Q.	...	Trunks of <i>Eucalyptus</i>	...	Thin, rigid, attached behind by circular disc, many zoned, lobed, ochrey, becoming brownish.
941	...	S.A.	...	...	...	Q.	...	Trunks	...	Leathery, thin, kidney shaped, shining, pale wood colour to brownish, zoned with concentric furrows.
942	...	...	...	...	N.S.W.	Q.	...	Trunks	...	Semicircular, sloping behind, rigid, amber, concentrically zoned and furrowed, radiately rough, finally smooth.
943	...	...	...	...	...	Q.	...	Trunks of <i>Hormoglyph cotinifolia</i>	...	Sessile, leathery to membranous, entire, somewhat bell shaped, silky hairy, pale.
944	...	S.A.	...	...	N.S.W.	Q.	...	Rotten wood	...	Corky to leathery, thin, pale wood colour to tawny, zoned, silky, shining.
945	...	S.A.	...	...	...	Q.	...	Trunks	...	Leathery, kidney shaped, rigid, becoming hoary, closely concentrically furrowed; margin thin, brown.
945A	...	S.A.	...	...	...	Q.	...	Branches	...	Only slightly different.
946	...	...	...	...	...	Q.	...	Trunks	...	Corky to leathery, kidney shaped, closely concentrically smooth, amber, opaque.
947	...	...	...	V.	...	Q.	...	Trunks	...	Corky to leathery, fibrous, bristly, zoneless, brown.

Syn. 499 (1801).—Trametes, Polyporus.

948	...	...	...	...	N.S.W.	...	...	Trunks	...	Corky, thick, convex, zoneless, becoming white.
949	...	S.A.	...	...	...	Q.	...	Trunks	...	Kidney shaped, sometimes with short stalk, whitish, downy, rather rough, slightly zoned.
950	...	...	...	...	...	Q.	...	Trunks	...	Narrowly reflexed, downy, pale.
951	...	...	...	...	N.S.W.	...	...	Trunks	...	Cushion shaped, thick, at first rough and downy, then smooth, zoned, pale.
952	...	S.A.	...	...	...	...	...	Trunks	...	Ochrey white, flabby, somewhat membranous.
953	...	...	...	V.	...	...	...	Trunks, &c.	...	Leathery, thin, running down behind, velvety, grey, with darker linear concentric zones.
954	...	...	...	...	...	Q.	...	Trunks	...	Overlapping, pale, variegated with dingy brown spots, split, hard, rough, zoned.
955	...	...	...	...	...	...	...	Trunks	...	Semicircular, pale, zoned in front, radiately rough.
956	...	...	...	...	...	Q.	B.	Old trunks	...	Expanded, often spreading for a foot or more in a continuous sheet, corky to woody, thick, wavy, pale wood colour, zoned within.
957	...	...	...	V.	...	...	...	Trunks	...	Corky, thickish, narrow behind, zoneless, rough, whitish.
958	...	...	...	...	...	Q.	...	Trunks	...	White, overlapping, thick, bleached above.
959	...	S.A.	...	...	...	...	...	Trunks	...	Pale ochrey, corky, thin, flattened, semicircular, zoned, roughish, shortly velvety.
960	...	...	...	...	...	Q.	...	Trunks	...	Expanded, corky to leathery, thin, pale wood colour. Pores flexuous.
961	...	...	...	...	...	Q.	...	Trunks	...	Corky, dirty amber, thin, roughish. Margin zoned.
962	...	...	...	...	...	Q.	...	Trunks	...	Corky, semicircular, flattened, velvety, pale wood colour, concentrically furrowed.
963	...	...	T.	...	...	...	...	Rotten wood	...	Inverted, somewhat circular, thin, brown.
964	...	S.A.	...	...	...	Q.	...	Stumps	...	Corky, amber to wood colour, semicircular, thin, zoned, rough, becoming almost smooth.
965	...	...	...	V.	...	Q.	B.	Trunks	...	Usually overlapping, leathery, velvety, grey, zones of same colour.

Hist. 62 (1759).

966	...	S.A.	...	...	...	...	...	Rotten wood	...	Somewhat globose, corky to woody, sessile, externally pale, internally sooty brown.
-----	-----	------	-----	-----	-----	-----	-----	-------------	-----	---

Syst. Myc. 342 (1821).—Hexagonia, Laschia.

967	...	...	...	V.	...	Q.	...	Trunks	...	Fleshy, tough, even, then scaly, yellowish tan.
968	...	S.A.	...	...	...	...	...	Stems	...	Cap thin, kidney shaped, hispid, netted, delicately downy. Stem short, cylindrical, rather rough.
969	...	...	T.	...	...	...	...	Trunks	...	Cap rather membranous, kidney shaped, smooth, tawny. Stem very short, blackening.
970	...	...	...	V.	N.S.W.	Q.	...	Branches, wood, &c.	...	Cap leathery, kidney shaped, concentrically furrowed, pale tan. Stem lateral, short, dilated above.
971	...	...	...	...	...	Q.	...	Wood	...	Cap white, then smoky, rough. Stem very short arising from orbicular disc.
972	...	...	...	...	N.S.W.	...	...	Trunks	...	Cap variegated with scales. Stem short, scaly, dilated upwards.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
70. LASCHIA.—Fries,					
973	901	VI. 6507	<i>L. caespitosa</i> ...	Sacc. Syll. VI. 407 (1888) ...	Tufted laschia ...
974	905	„ 6518	<i>L. micropus</i> ...	Berk., Linn. Journ. XIII. 170 (1873) ...	Small-stalked laschia ...
975	904	„ 6516	<i>L. pustulata</i> ...	Berk. and Br., Linn. Journ. XIV. 58 (1875) ...	Pustulate laschia ...
976	903	„ 6508	<i>L. Thwaitesii</i> ...	Berk. and Br., Linn. Journ. XIV. 58 (1875) ...	Thwaites' laschia ...
977	902	„ 6504	<i>L. tremellosa</i> ...	Fries, S.V. 325 (1849) ...	Gelatinous laschia ...
71. CAMPBELLIA.—Cooke and Mass.,					
978	906	VI. 6523 IX. p. 205	<i>C. infundibuliformis</i> ...	Cooke and Mass., Grev. XVIII. 87 (1890) ...	Funnel-shaped campbellia ...
72. MERULIUS.—Hall.					
979	910	VI. 6542	<i>M. aureus</i> ...	Fries, Elench. 62 (1828) ...	Golden merulius ...
980	909	„ 6538	<i>M. Baileyi</i> ...	Berk. and Br., Linn. Trans. II. 62 (1883) ...	Bailey's merulius ...
981	907	„ 6532	<i>M. corium</i> ...	Fries, Elench. 58 (1828) ...	Leathery merulius ...
982	914	„ 6563	<i>M. lacrymans</i> ...	Fries, S.M. I. 328 (1821) ...	Weeping merulius (Dry Rot) ...
983	913	„ 6559	<i>M. pallens</i> ...	Berk., Outl. 296 (1860) ...	Pale merulius ...
984	908	...	<i>M. pelliculosus</i> ...	Grev. XIX. 109 (1891) ...	Pellicle-like merulius ...
985	912	VI. 6553	<i>M. serpens</i> ...	Tode., Abh. Hall. I. 355 (1790) ..	Spreading merulius ...
986	911	„ 6550	<i>M. tenuissimus</i> ...	Berk. and Br., Linn. Trans. II. 62 (1883) ...	Very thin merulius ...
987	...	...	<i>Xylostroma giganteum</i> ...	Fries. ...	Gigantic xylostroma ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
in Linn. V. 533 (1830).									
973	...	...	...	...	N.S.W.	Q.	...	Branches ...	Densely tufted. Caps conical. Stems smooth, united at base.
974	...	...	...	V.	...	...	...	Trunks ...	Tawny yellow, minute, peziza-like, shortly stalked.
975	...	...	...	...	...	Q.	...	Rotten wood	Cap oehrey, then rusty, tuberculose, circular. Stem oblique, of same colour.
976	...	...	...	...	...	Q.	...	Wood ...	Tufted and gregarious. Cap tremelloid, often oblique, orange. Stem slender, white.
977	W.A.	...	...	V.	N.S.W.	Q.	...	Rotten wood	Bell shaped, attached behind, membranous, gelatinous, entirely dark red, rather thick, leathery when dry.
Grev. XVIII. 87 (1890).									
978	...	...	...	V.	...	...	...	Wood ...	Gelatinous, large, stalked. Cap deeply funnel shaped, thick. Stem short, thick, expanded disc at base.
Helv. 150 (1768).—Boletus.									
979	...	...	...	...	...	Q.	...	Wood ...	Expanded, thin, membranous, adherent, golden yellow.
980	...	...	...	...	...	Q.	...	Trunks ...	Fan shaped, viscid, smooth, almost orange coloured when fresh. Margin notched, rough, inflexed, flesh yellow.
981	W.A.	...	T.	V.	...	Q.	B.	Trunks and branches	Expanded, soft, rather papery, shaggy beneath, white.
982	W.A.	...	...	...	...	Q.	B.	Rotten wood chiefly of dwellings, in cellars	Large, spongy to fleshy, rusty yellow, web-like or velvety below.
983	...	...	T.	V.	...	...	B.	Trunks ...	Attached, fleshy, somewhat gelatinous, thin, slightly downy.
984	...	...	...	V.	...	...	...	Branches of <i>Acacia</i>	Broadly expanded, membranous, white, hyaline when dry, like a thin pellicle.
985	...	...	...	...	...	Q.	B.	Rotten wood	Crustaceous, attached, thin, pale, then reddish, spreading in wavy manner.
986	...	...	...	...	..	Q	...	Parasitic on <i>Hymenochaete</i>	Papery, forming very thin irregular yellowish-brown patches, nearly white at margin.
<hr/>									
987	...	...	...	V.	N.S.W.	Q.	...	Heart wood of various Eucalypts	This is the sterile state of a wood-destroying fungus, probably belonging to Polyporaceae, and consisting of thick dense leathery sheets like chamois leather.

ORDER III.—HYDNACEÆ, FRIES.

73. Hydnum, Linn.
74. Sistotrema, Pers.
75. Irpex, Fries.

76. Radulum, Fries.
77. Phlebia, Fries.

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER III.—HYDNACEÆ,					
73. HYDNUM.—Linn.,					
988	934	VI. 6761	<i>H. alutaceum</i> ...	Fries, S.M. I. 417 (1821) ...	Wash-leather hydnum ...
989	920	„ 6639	<i>H. ambustum</i> ...	Cooke and Mass., Grev. XVI. 32 (1887) ...	Scorched hydnum ...
990	...	...	<i>H. calcareum</i> ...	Cooke and Mass., Grev. XXI. 38 (1892) ...	Chalky-white hydnum ...
991	938	VI. 6809	<i>H. cervinum</i> ...	Berk., Fl. Tasm. II. 256 (1860) ...	Fawn hydnum ...
992	...	„ 6696	<i>H. cirrhatum</i> ...	Pers., Syn. 558 (1801) ...	Curled hydnum ...
993	925	„ 6677	<i>H. coralloides</i> ...	Scop. Carn. 472 (1772) ...	Coralline hydnum ...
994	921	IX. 840	<i>H. crocicens</i> ...	Cooke, Grev. XIX. 45 (1890) ...	Saffron-coloured hydnum ...
995	924	VI. 6664	<i>H. cyathiforme</i> ...	Schaeff., Fl. Dan. 1020 ...	Cup-shaped hydnum ...
996	930	„ 6731	<i>H. delicatulum</i> ...	Klotzsch., Ann. Nat. Hist. III. 395 (1839) ...	Delicate hydnum ...
997	933	„ 6737	<i>H. dispersum</i> ...	Berk., Hook., Lond. Journ. IV. 58 (1845) ...	Scattered hydnum ...
998	940	„ 6824	<i>H. filicicola</i> ...	Berk., Fl. Tasm. II. 256 (1860) ...	Fern-growing hydnum ...
999	927	„ 6722	<i>H. flavum</i> ...	Berk., Ann. Nat. Hist. X. 380 (1842) ...	Yellow hydnum ...
1000	931	„ 6733	<i>H. gilvum</i> ...	Berk., Hook., Journ. 168 (1851) ...	Yellowish-tan hydnum ...
1001	923	„ 6660	<i>H. graveolens</i> ...	Delast. in Fries, Hym. Eur. 605 (1874) ...	Strong-smelling hydnum ...
1002	936	„ 6779	<i>H. investiens</i> ...	Berk., Hook., Lond. Journ. IV. 57 (1845) ...	Lining hydnum ...
1003	941	„ 6836	<i>H. isidioides</i> ...	Berk., Hook., Lond. Journ. IV. 58 (1845) ...	Isidium-like hydnum ...
1004	918	„ 6624	<i>H. lævigatum</i> ...	Swartz., Vet. Akad. Handl. 243 (1810) ...	Smooth hydnum ...
1005	932	„ 6738	<i>H. membranaceum</i> ...	Bull., Champ. 481 (1798) ...	Membranous hydnum ...
1006	926	„ 6705	<i>H. merulioides</i> ...	Berk., Linn. Trans. II. 63 (1883) ...	Merulius-like hydnum ...
1007	939	„ 6812	<i>H. mucidum</i> ...	Pers., Syn. 561 (1801) ...	Hoary hydnum ...
1008	929	„ 6727	<i>H. Muelleri</i> ...	Berk., Linn. Journ. XIII. 167 (1873) ...	Mueller's hydnum ...
1009	922	„ 6657	<i>H. nigrum</i> ...	Fries, S.M. I. 404 (1821) ...	Black hydnum ...
1010	928	„ 6725	<i>H. ochraceum</i> ...	Pers., Syn. 559 (1801) ...	Ochrey hydnum ...
1011	919	„ 6633	<i>H. repandum</i> ...	Linn., Sp. Pl. 1178 (1753) ...	Repand hydnum ...
1012	937	„ 6795	<i>H. udum</i> ...	Fries, S.M. I. 422 (1821) ...	Moist hydnum ...
1013	935	„ 6778	<i>H. xanthum</i> ...	Berk. and Curt., Grev. I. 98 (1872) ...	Yellow hydnum ...
1013A	„	„	<i>H. xanthum</i> , var. <i>teretidens</i> ...	Cooke., Handb. Aust. Fung. 172 (1892) ...	Cylindrical hydnum ...
74. SISTOTREMA.—Pers.					
1014	943	VI. 6872	<i>S. irpicinum</i> ...	Berk. and Br., Linn. Trans. II. 62 (1883) ...	Irpex-like sistotrema ...
75. IRPEX.—Fries,					
1015	949	VI. 6925	<i>I. Archeri</i> ...	Berk., Fl. Tasm. II. 257 (1860) ...	Archer's irpex ...
1016	947	„ 6895	<i>I. flavus</i> ...	Klotzsch., Linn. VIII. 488 (1833) ...	Yellow irpex ...
1017	944	„ 6876	<i>I. hexagonoides</i> ...	Kalch., Grev. IX. 1 (1880) ...	Hexagonia-like irpex ...

ARRANGEMENT OF GENERA (9).

78. *Grandinia*, Fries.
79. *Porothelium*, Fries.

80. *Odontia*, Pers.
81. *Kueiffia*, Fries.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
988	...	...	...	V.	...	...	B.	Rotten wood	Spore-bearing surface expanded longitudinally, crustose, smooth, pale ochre. Spines acute.
989	...	...	...	V.	...	...	...	Sandy soil	Cap fleshy to membranous, brick red, turning black. Stem erect, slender, paler than cap.
990	...	...	...	V.	...	...	...	Bark	Chalky white, opaque, widely expanded. Spore-bearing surface crustose, smooth, mealy. Has a scorched appearance when dry.
991	...	...	T.	...	...	...	...	Rotten wood	Inverted, expanded, indistinct margin, pale vinous, at first delicately downy.
992	...	...	...	V.	...	...	B.	Trunks	Fleshy, expanded, colour variable, the upper surface with long curled abortive spines.
993	...	...	...	...	...	Q.	B.	Rotten wood, trunks	Much branched, creamy, like cauliflower at first, then a mass of coralline branches.
994	...	...	...	V.	...	...	...	Ground	Cap fleshy, thin, ochrey yellow. Stem central, slender, smooth when dry.
995	...	...	...	...	...	Q.	B.	Woods	Small, commonly grown together. Cap leathery, thin, funnel shaped, zoned, ashy pale. Stem smooth, slender, disc rather downy.
996	...	...	...	...	N.S.W.	...	...	Trunks	Cap expanded, reflexed, leathery, thin, margin yellowish.
997	W.A.	...	...	...	...	...	...	Rotten wood	In long patches. Spore-bearing surface, thin, waxy, at length disappearing.
998	...	...	T.	...	...	...	...	Fern stems	Expanded, indistinct margin, white, thin.
999	...	...	...	...	...	Q.	...	Branches	Sessile, nearly circular, thin, pale yellow, smooth.
1000	...	...	...	...	...	Q.	...	Rotten wood	Fan shaped, overlapping, thin, ochrey to yellowish tan, clothed with cartilaginous radiating hairs.
1001	...	...	...	V.	...	Q.	B.	Woods	Cap leathery, thin, soft, zoneless, rough, dark brown, margin whitish. Stem slender. Odour of melilot.
1002	W.A.	...	...	...	...	...	...	In cavities of trunks of <i>Xanthorrhoea</i>	Widely expanded, spore-bearing surface at first downy, then compact, smooth.
1003	W.A.	...	...	...	...	...	...	Hymenium of <i>Polyporus gryphaeformis</i>	Spore-bearing surface crustaceous, white, margin somewhat fringed, separating in chips.
1004	...	...	T.	...	...	Q.	B.	Solitary in wood	Cap fleshy, firm, smooth, amber. Stem thick, even.
1005	...	...	...	V.	...	Q.	B.	Rotten wood	Spore-bearing surface expanded, waxy, membranous, smooth, tawny to rusty.
1006	...	...	...	...	...	Q.	...	Wood	Semicircular, thick, smooth, pale, rough with prominent lines.
1007	...	...	...	...	...	Q.	...	Trunks	White. Spore-bearing surface very broad, membranous, soft, evanescent.
1008	...	...	...	...	N.S.W.	Q.	...	Wood	Inverted, then reflexed, sometimes growing together, velvety, somewhat zoned.
1009	...	...	...	...	N.S.W.	...	B.	Woods	Cap corky, rigid, downy, zoneless, bluish black. Stem stout, black.
1010	...	...	...	V.	...	Q.	B.	Trunks and sticks	Small. Expanded and reflexed, leathery, thin, zoned, ochrey.
1011	...	...	T.	...	...	...	B.	Woods	Gregarious. Cap fleshy, fragile, repand and wavy, deformed, pale, creamy buff. Stem rather short. Very common. <i>Edible</i> .
1012	...	...	T.	...	N.S.W.	Q.	B.	Rotten wood	Spore-bearing surface expanded, thin, somewhat gelatinous, flesh colour to watery yellow.
1013	...	...	...	...	N.S.W.	...	...	Trunks	Inverted, thin, margin delicately downy, then waxy. Spines compressed.
1013A	...	...	...	...	N.S.W.	...	...	Trunks	Spines not compressed and hanging from the brighter pale-orange cap.

Tent. Disp. 28 (1797).

1014	...	...	...	...	...	Q.	...	Dead branches	Somewhat cuticular, thick, delicately downy, pale, descending deeply behind.
------	-----	-----	-----	-----	-----	----	-----	---------------	--

Pl. Homon. 81 (1825).—*Polyporus*.

1015	...	...	T.	...	...	...	...	Rotten wood	White, inverted. Margin web-like, downy.
1016	...	S.A.	...	V.	N.S.W.	Q.	...	Trunks, &c.	Expanded, spongy soft, yellow, margin shortly reflexed, downy.
1017	...	...	...	...	N.S.W.	...	...	Trunks	Entirely white. Cap corky to leathery, running down behind, faintly zoned, teeth disposed in honeycomb manner.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
75. IRPEX.—Fries,					
1018	946	VI. 6891	<i>I. maximus</i>	Mont. Ann. Sci. Nat. 2 Ser. VIII. 364 (1837) ...	Largest irpex
1019	950	„ 6883	<i>I. sinuosus</i> , var. <i>cervicolor</i> ...	Berk. and Br., Linn. Journ. XIV. 60 (1875) ...	Fawn-coloured irpex
1020	948	„ 6902	<i>I. tabacinus</i>	Berk. and Curt., Grev. I. 102 (1872) ...	Tobacco-coloured irpex
1021	945	„ 6888	<i>I. zonatus</i>	Berk., Hook., Journ. 168 (1854) ...	Zoned irpex
76. RADULUM.—Fries,					
1022	951	VI. 6931	<i>R. molare</i>	Fries, Elench. 151 (1828)	Tubercled radulum
77. PHLEBIA.—Fries,					
1023	956	VI. 6965	<i>P. coriacea</i>	Berk., Linn. Journ. XVIII. 385 (1881) ...	Leathery phlebia
1024	954	„ 6960	<i>P. hispidula</i>	Berk. Linn., Journ. XIII. 167 (1873) ...	Rough phlebia
1025	952	„ 6950	<i>P. merismoides</i>	Fries, S.M. I. 427 (1821) ...	Merisma-like phlebia
1026	953	„ 6951	<i>P. radiata</i>	Fries, S.M. I. 427 (1821) ...	Radiate phlebia
1027	955	„ 6964	<i>P. reflexa</i>	Berk., Hook., Journ. 168 (1851) ...	Reflexed phlebia
78. GRANDINIA.—Fries,					
1028	960	VI. 6980	<i>G. australis</i>	Berk., Fl. Tasm. II. 257 (1860) ...	Southern grandinia
1029	959	„ 6976	<i>G. crustosa</i>	Fries, Epicr. 527 (1838) ...	Crustaceous grandinia
1030	961	IX. 865	<i>G. glauca</i>	Cooke, Grev. XVII. 55 (1889) ...	Glaucous grandinia
1031	957	VI. 6969	<i>G. granulosa</i>	Fries, Epicr. 527 (1838) ...	Granular grandinia
79. POROTHELIUM.—Fries,					
1032	915	VI. 6576	<i>P. subtile</i>	Fries, S.M. I. 506 (1821)	Delicate porothelium
80. ODONTIA.—Pers.					
1033	962	VI. 7018	<i>O. secernibilis</i>	Berk., Fl. Tasm. II. 257 (1860) ...	Separable odontia
81. KNEIFFIA.—Fries,					
1034	963	VI. 7022	<i>K. Muellieri</i>	Berk., Linn. Journ. XIII. 167 (1873) ...	Mueller's kneiffia
1035	963bis.	„ 7020	<i>K. setigera</i>	Fries, Epicr. 529 (1838) ...	Bristle-bearing kneiffia

OF AUSTRALIAN FUNGI—*continued*.

Number	Habitat.						B.	Occurrence.	General Characters.
	W. A.	S. A.	T.	V.	N. S. W.	Q.			
Pl. Homon. 81 (1825).—Polyporus—continued.									
1018	...	...	...	...	...	Q.	...	Trunks ...	Leathery, thin, kidney shaped, at first downy, then naked. concentrically furrowed.
1019	...	...	...	...	...	Q.	...	Wood ...	Expanded, shortly reflexed, thin, soft, zoneless, entirely fawn colour.
1020	...	...	...	...	...	Q.	...	Trunks ...	Slightly reflexed, somewhat zoned, bay brown, downy.
1021	...	...	...	V.	N. S. W.	Q.	...	Dead wood	Wood colour. Caps overlapping, somewhat fan shaped, leathery, zoned, delicately downy, becoming smooth.
Pl. Homon. 81 (1825).—Hydnum, Sistotrema.									
1022	...	...	...	...	...	Q.	...	Trunks, and on peach trees	Broadly expanded, crustaceous, smooth, somewhat yellow. Tubercles deformed, short.
S. M. I. 426 (1821).—Auricularia, Thelephora.									
1023	...	...	...	...	...	Q.	...	Ground ...	Spoon shaped, leathery, brown. Folds irregular.
1024	...	S. A.	...	...	...	...	...	Trunks ...	Inverted, reflexed, reddish, zoned, velvety and hispid.
1025	...	...	...	...	...	Q.	B.	Trunks, &c.	Expanded, even or incrusting, branched, flesh colour, white. Margin orange.
1026	...	...	...	...	...	Q.	B.	Bark ...	Roundish, smooth, red flesh colour. Margin radiately toothed.
1027	...	...	...	V.	N. S. W.	Q.	...	Wood ...	Inverted, semicircular, reflexed, cracked, clothed with spongy down, zoned, vinous brown.
Epier. 527 (1838).—Hydnum, Thelephora.									
1028	...	...	T.	V.	...	...	...	Rotten wood	Inverted, expanded, indistinct margin, pale, cracked, snow white within.
1029	...	...	...	V.	...	...	B.	Bark ...	Mealy, expanded irregularly, crustaceous, white.
1030	...	...	...	...	...	Q.	...	Wood ...	Waxy, broadly expanded, glaucous, margin distinct.
1031	...	...	T.	...	...	Q.	B.	Old wood...	Waxy, broadly expanded, tan coloured. Margin distinct. Granules hemispherical, crowded.
Obs. II. 272 (1818).—Boletus.									
1032	...	...	T.	...	...	...	...	Bark ...	Irregularly expanded, membranous, snow white, porous warts distinct.
Tent. Disp. 30 (1797).									
1033	...	...	T.	...	...	...	...	Rotten wood	Inverted, separable, membranous, white.
Epier. 529 (1838).									
1034	...	S. A.	...	...	...	...	...	Rotten wood	Resupinate, thin, mealy when young, cracking here and there.
1035	...	...	...	...	...	...	B.	Wood ...	Whitish, pale buff when dry, soft, forming a layer or sometimes fleshy. Bristles rigid, very minute, scattered.

ORDER IV.—THELEPHORACEÆ PERS.

82. Craterellus, Fries.
83. Thelephora, Ehrh.
84. Cladoderris, Pers

85. Stereum, Pers.
86. Hymenochaete, Lev.
87. Corticium, Fries.

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
---------	-----------------	--------------------	------------------	---------------------	---------------

ORDER IV.—THELEPHORACEÆ,

82. CRATERELLUS.—Fries, Epier. 531 (1838).—

1036	965	VI. 7046	<i>C. confluens</i> ...	Berk. and Curt., Linn. Journ. IX. 423 (1867) ...	Confluent craterellus ...
1037	964	„ 7042	<i>C. cornucopioides</i> ...	Pers. Myc., Eur. II. 5 (1822) ...	Trumpet-shaped craterellus ...
1038	967	IX. 882	<i>C. multiplex</i> ...	Cooke and Mass., Grev. XVIII. 25 (1889) ...	Multiplex craterellus ...
1039	966	...	<i>C. pnsio</i> ...	Berk., Fl. Tasm. II. 258 (1860) ...	Little craterellus ...

83. THELEPHORA.—Ehrh. in Roth.

1040	977	VI. 7116	<i>T. Archeri</i> ...	Berk., Fl. Tasm. II. 258 (1860) ...	Archer's thelephora ...
1041	...	„ 7146	<i>T. atra</i> ...	Weinm., Ross. 380 (1836) ...	Black thelephora ...
1042	975	„ 7098	<i>T. caryophyllea</i> ...	Pers., Syn. 565 (1801) ...	Clove thelephora ...
1043	974	„ 7087	<i>T. conerescens</i> ...	Fries, Pl. Preiss. II. 136 (1846) ...	Conerescent thelephora ...
1044	976	„ 7107	<i>T. congesta</i> ...	Berk., Linn. Journ. XVI. 168 (1878) ...	Congested thelephora ...
1045	984	„ 7159	<i>T. cristata</i> ...	Fries, S.M. I. 434 (1832) ...	Crested thelephora ...
1046	987	„ 7192	<i>T. exsculpta</i> ...	Berk., Linn. Journ. XIII. 168 (1873) ...	Carved thelephora ...
1047	980	„ 7144	<i>T. intybaea</i> ...	Pers., Syn. 567 (1801) ...	Endive thelephora ...
1048	982	„ 7147	<i>T. laciniata</i> ...	Pers., Syn. 567 (1801) ...	Cut thelephora ...
1049	979	„ 7142	<i>T. multipartita</i> ...	Schwein., in Fries, Epier. 536 (1838) ...	Multipartite thelephora ...
1050	978	„ 7129	<i>T. myriomera</i> ...	Fries, Pl. Preiss. II. 137 (1846) ...	Myriad-partite thelephora ...
1051	...	„ 7103	<i>T. palmata</i> ...	Fries, S.M. I. 432 (1832) ...	Palmate thelephora ...
1052	986	„ 7188	<i>T. pedicellata</i> ...	Schwein., Syn. Car. 108 (1822) ...	Elevated thelephora ...
1053	985	„ 7173	<i>T. riccioidea</i> ...	Berk., Fl. Tasm. II. 258 (1860) ...	Riccia-like thelephora ...
1054	...	„ 7088	<i>T. spongipes</i> ...	Berk., Linn. Trans. II. 63 (1883) ...	Spongy-stalked thelephora ...
1055	983	IX. 887	<i>T. stereoides</i> ...	Cooke and Mass., Grev. XVIII. 5 (1889) ...	Stereum-like thelephora ...
1056	981	VI. 7145	<i>T. terrestris</i> ...	Ehrh., Crypt. 179 (1788) ...	Ground thelephora ...

84. CLADODERRIS.—Pers. in Freye.

1057	989	VI. 7211	<i>C. australica</i> ...	Berk. in Grev. XI. 28 (1882) ...	Australian cladoderris ...
1058	988	„ 7210	<i>C. anstralis</i> ...	Kalch., in Thunb. Syn. Myc. Aust. II. (1878) ...	Southern cladoderris ...
1059	990	„ 7215	<i>C. dendritica</i> ...	Pers., Freye. Voy. (1826) ...	Tree-like cladoderris ...
1060	991	„ 7207	<i>C. spongiosa</i> ...	Fries, Fung. Nat. 20 (1848) ...	Spongy cladoderris ...

85. STEREUM.—Pers.

1061	1029	VI. 7418	<i>S. acerinum</i> ...	Fries, Hym. Eur. 645 (1874) ...	Maple sterem ...
1062	1027	„ 7375	<i>S. amœnum</i> ...	Mass., Mon. Thel. Linn. Journ. XXVII. 193 (1891) ...	Charming sterem ...
1063	992	„ 7074	<i>S. caperatum</i> ...	Mass., Mon. Thel. Linn. Journ. XXVII. 161 (1891) ...	Wrinkled sterem ...
1063A	„	„ 7075	<i>S. caperatum</i> , var. <i>lamellatum</i> ...	Cooke, Handb. Aust. Fung. 182 (1892) ...	Plaited sterem ...
1063B	„	„ 7088	<i>S. caperatum</i> , var. <i>spongipes</i> ...	Cooke, Handb. Aust. Fung. 182 (1892) ...	Spongy-stalked sterem ...
1064	1013	„ 7371	<i>S. complicatum</i> ...	Fries, Epier. 548 (1838) ...	Crisped sterem ...

ARRANGEMENT OF GENERA (12).

88. Peniophora, Cooke.
89. Coniophora, D. C.
90. Aleurodiscus, Rabb.

91. Cyphella, Fries.
92. Solenia, Hoffm.
93. Lachnocladium, Lev.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
1036	...	...	...	V.	...	Q.	...	Ground ...	Orange, deeply funnel shaped, margin incurved. Stem divided above into numerous caps.
1037	...	...	...	...	...	Q.	B.	Wood ...	Somewhat membranaceous, trumpet shaped, sooty brown, becoming black. Stem hollow, smooth, black. Common. <i>Edible</i> .
1038	...	...	T.	...	...	...	...	Ground ...	Kidney shaped, and attached at base to stem in series of five or six superimposed, ochrey. Stem slender, erect, wrinkled.
1039	...	...	T.	...	...	...	...	Among moss	Orange. Cap lateral, convex. Stem thickened upward.

PERS. MYC. EUR. I. 109 (1822).

Cantharellus, Merulius, Peziza, Helvella.

1040	...	...	T.	V.	...	Q.	...	Ground ...	Forked, branches compressed and dilated above, ochrey.
1041	...	...	...	V.	...	...	...	Ground ...	Growing in tufts. Expanded, soft, tufts arising from common tuber.
1042	W.A.	...	...	...	...	Q.	B.	Among grass	Purplish brown. Cap somewhat leathery, depressed, margin divided. Stem short and smooth. Inodorous.
1043	W.A.	...	...	...	...	...	...	Moist places	Tufted and growing together. Cap funnel shaped, brown as well as stem.
1044	...	...	...	V.	N.S.W.	Q.	...	Moist places	Small, gregarious, dark purple, sparingly branched in a forked manner.
1045	...	...	...	...	...	Q.	...	On moss, grass, &c.	Incrusting, tough, pale, spreading out into branches or fringes.
1046	...	...	...	V.	...	...	...	Bark ...	Circular, grey, tinged with pink, radiating with teeth towards the margin, powdery, dark purple beneath.
1047	...	...	...	...	N.S.W.	...	B.	Woods ...	Tufted, soft, whitish to rusty, then sooty brown.
1048	...	...	...	V.	...	...	B.	On trunks and ground	Incrusting, soft to leathery, rusty brown. Very common.
1049	...	...	...	...	...	Q.	...	Ground ...	Leathery, ashy brown. Cap smooth, much divided down to stem in a branching manner.
1050	W.A.	...	...	...	...	...	...	Ground ...	Tufted, papery, flattened and much branched, growing together.
1051	...	...	...	...	...	Q.	B.	Ground in woods	Fœtid, densely clustered. Brownish purple, soft, stem-like base with wedge-shaped branches.
1052	W.A.	...	...	V.	...	Q.	...	Branches	Spread out, finely filamentous, compact, cinnamon brown, rooting beneath with bundles of fibres.
1053	...	...	T.	...	...	...	...	Bare soil ...	Pale, closely attached and radiately branched.
1054	...	...	...	...	N.S.W.	Q.	...	Old wood	Funnel shaped, spongy and downy, radiately folded and wrinkled. Stem spongy.
1055	...	...	...	V.	...	...	...	Bark ...	Leathery, spread out, reflexed, downy, rust colour, crisped at the margin.
1056	...	...	...	V.	...	...	B.	Ground ...	Tufted, soft brown, turning black. Caps overlapping, flattened, running down into sub-lateral stem.

Voy. Bot. 176 (1826).—Thelephora.

1057	...	...	...	...	N.S.W.	...	...	Wood ...	Funnel shaped, oblique, fan shaped or semicircular. Stem excentric, short, of an umber colour like the cap.
1058	...	...	...	...	N.S.W.	...	...	Trunks ...	Cap somewhat excentric, funnel shaped, oblique, margin with fringed lobes. Stem woody and downy.
1059	...	...	...	V.	N.S.W.	Q.	...	Wood ...	Leathery to soft, yellowish tan. Cap kidney shaped, entire. Stem lateral, firm.
1060	...	...	...	V.	...	...	...	Trunks ...	Broadly funnel shaped, spongy, elastic, tan colour. Stem central, very short, woody.

Obs. Myc. I. 35 (1796).—Thelephora, Auricularia, Corticium, Elvelia.

1061	...	...	T.	V.	...	...	B.	Bark ...	Crustaceous, smooth, white, thin. Surface generally covered with minute particles of lime.
1062	...	S.A.	...	...	...	...	...	Fallen branches	Gregarious. Leathery, membranaceous, hairy, white.
1063	...	...	...	V.	N.S.W.	Q.	...	Trunks ...	Leathery to membranaceous, wrinkled, folded, ochrey, hairy in centre. Stem central, thick, downy.
1063A	...	...	...	...	...	Q.	...	Wood ...	Ochrey. Cap wrinkled, plaited. Stem elongated, velvety.
1063B	...	...	...	...	...	Q.	...	Wood ...	Spongy and downy, both in cap and stem.
1064	...	...	...	V.	...	Q.	...	Branches ...	Papery, furrowed, brownish or ochrey, much crisped and lobed.

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
85. STEREOUM.—Pers. Obs. Myc. I. 35 (1796).—					
1065	...	VI. 7299	<i>S. concolor</i> ...	Berk., Fl. Tasm. II. 259 (1860) ...	One-coloured sterium ...
1066	1000	" 7255	<i>S. crucebuliforme</i> ...	Mass., Mon. Thel. Linn. Journ. XXVII. 168 (1891)	Crucible-shaped sterium ...
1067	993	" 7223	<i>S. eyathiforme</i> ...	Fries, Epier. 245 (1838) ...	Cup-shaped sterium ...
1068	994	" 7233	<i>S. elegans</i> ...	Fries, Epier. 545 (1838) ...	Elegant sterium ...
1069	1011	" 7271	<i>S. fasciatum</i> ...	Fries, Epier. 546 (1838) ...	Banded sterium ...
1070	1012	" 7270	<i>S. gausapatum</i> ...	Fries, Hym. Eur. 638 (1874) ...	Rough-coated sterium ...
1071	1014	" 7288	<i>S. hirsutum</i> ...	Fries, Epier. 549 (1838) ...	Hairy sterium ...
1071A	...	...	<i>S. hirsutum</i> , var. <i>tenellum</i> ..	Sacc., Notes Myc. 5 (1890) ...	Tender sterium ...
1071B	...	...	<i>S. hirsutum</i> , var. <i>glaucellum</i> ..	Sacc., Notes Myc. 5 (1890) ...	Glaucous sterium ...
1072	1015	VI. 7329	<i>S. illudens</i> ...	Berk., Hook., Lond. Journ. Bot. IV. 59 (1845)	Deceptive sterium ...
1073	1009	" 7272	<i>S. involutum</i> ...	Klotzsch, Linnæa VII. 499 (1832) ...	Involute sterium ...
1074	...	" 7316	<i>S. Kalchbrenneri</i> ...	Sacc. Syll. VI. 568 (1888) ...	Kalchbrenner's sterium ...
1075	1007	" 7267	<i>S. Leichardtianum</i> ...	Sacc. Syll. VI. 559 (1888) ...	Leichardt's sterium ...
1076	1008	" 7311	<i>S. lobatum</i> ...	Fries, Epier. 547 (1838) ...	Lobed sterium ...
1077	1028	" 7360	<i>S. molle</i> ...	Sacc. Syll. VI. 577 (1888) ...	Soft sterium ...
1078	998	" 7247	<i>S. Moselei</i> ...	Berk., Linn. Journ. XVI. 48 (1878) ...	Moseley's sterium ...
1079	995	" 7229	<i>S. nitidulum</i> ...	Berk., Hook., Lond. Journ. Bot. II. 638 (1843)	Shining sterium ...
1080	1017	" 7283	<i>S. ochroleucum</i> ...	Fries, Hym. Eur. 639 (1874) ...	Yellowish-white sterium ...
1081	999	" 7253	<i>S. prolificus</i> ...	Berk., Linn. Journ. XVI. 41 (1878) ...	Prolific sterium ...
1082	1018	" 7284	<i>S. purpureum</i> ...	Pers. Obs. Myc. II. 92 (1796) ...	Purple sterium ...
1083	1005	" 7263	<i>S. pusillum</i> ...	Berk., Ann. Nat. Hist., X. 381 (1842) ...	Small sterium ...
1084	1001	" 7282	<i>S. radiatofissum</i> ...	Berk. and Broome, Linn. Trans. II. 63 (1883)	Radiately divided sterium ...
1085	1020	" 7526	<i>S. radicale</i> ...	Mass., Mon. Thel. Linn. Journ. XXVII. 187 (1891)	Root-growing sterium ...
1086	1019	" 7388	<i>S. retirugum</i> ...	Cooke, Proc., Roy. Soc. Ed. XI. 456 (1882)	Net-veined sterium ...
1087	1023	" 7336	<i>S. rugosum</i> ...	Fries, Epier. 552 (1838) ...	Wrinkled sterium ...
1088	1010	" 7278	<i>S. semilugens</i> ...	Kalch., Grev. IX. 1 (1880) ...	Gloomy sterium ...
1089	1021	" 7340	<i>S. simulans</i> ...	Berk. and Broome, Linn. Trans. II. 64 (1883)	Simulating sterium ...
1090	996	" 7070	<i>S. Sowerbei</i> ...	Mass., Mon. Thel. Linn. Journ. XXVII. 164 (1891)	Sowerby's sterium ...
1091	1022	" 7289	<i>S. spadiceum</i> ...	Fries, Epier. 549 (1838) ..	Bright-brown sterium ...
1092	1032	" 7387	<i>S. sparsum</i> ...	Berk., Linn. Journ. XIII. 169 (1873) ...	Scattered sterium ...
1093	1002	" 7257	<i>S. spathulatum</i> ...	Berk., Hook., Journ. VIII. 274 (1856) ...	Spoon-shaped sterium ...
1094	1006	" 7295	<i>S. striatum</i> ...	Fries, Hym. Eur. 641 (1874) ...	Streaked sterium ...
1095	1030 & 1031	" 7410	<i>S. strumosum</i> ...	Fries, Nov. Symb. Myc. 111 (1851) ...	Swollen sterium ...
1096	1024	" 7300	<i>S. sulphuratum</i> ...	Berk. and Rav., Linn. Journ. X. 331 (1869)	Sulphur-coloured sterium ...
1097	...	...	<i>S. theleporoides</i> ... (substituted for <i>S. pannosum</i>)	McAlpine, Syst. Arr. Austr. Fung. (1894) Cooke and Mass., Grev. XXI. 38 (1892)	Thelephora-like sterium ...
1098	997	VI. 7254	<i>S. Thozetii</i> ...	Berk., Linn. Journ. XVIII. 385 (1881) ...	Thozet's sterium ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Thelephora, Auricularia, Corticium, Elvella—continued.									
1065	...	...	T.	...	...	...	...	Dead branches	Dirty white, semicircular, decurrent behind, soft, downy, contracted when dry.
1066	...	...	...	V.	...	...	...	Wood	Hairy, margin inflexed. Resembling <i>Crucibulum vulgare</i> .
1067	...	S.A.	...	V.	...	...	...	Wood and ground	Leathery, whitish. Cap broad ($1\frac{1}{2}$ – $2\frac{1}{2}$ inches), wine-glass shaped, bristly. Stem about $\frac{1}{2}$ inch long, central, smooth.
1068	...	S.A.	...	V.	N.S.W.	Q.	...	Ground	Cartilaginous to leathery. Cap funnel shaped usually, chestnut, shining. Stem short and slender.
1069	...	S.A.	...	V.	...	Q.	...	Trunks of <i>Eucalypts</i>	Leathery, tufted, thin, ash grey, zoned with shining bay-brown bands, and tapering towards base.
1070	...	...	...	...	...	Q.	...	Trunks	Tufted, sessile. Caps soft, corky, shell shaped, rough haired, tawny, becoming pale, fringed.
1071	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Trunks and branches	Leathery, rigid. Caps overlapping, spread out, rough haired, somewhat zoned, dirty ochre, becoming pale. Common.
1071A	...	S.A.	...	...	...	...	...	Rotten wood of <i>Acacia</i>	Membranous. Caps reflexed, whitish, zoned.
1071B	...	S.A.	...	...	...	...	...	Trunks	Cap velvety, discoloured and zoned. Hymenium greyish, glaucous.
1072	W.A.	S.A.	T.	V.	N.S.W.	Q.	...	Branches, &c.	Leathery and rather rigid. Caps confluent, zoned, radiately plaited, bay brown.
1073	...	...	...	...	...	Q.	...	Trunks	Leathery, tufted. Caps overlapping, grown together, ear shaped, longitudinally furrowed, tapering at base into lateral black stem.
1074	...	...	...	...	...	...	...	...	Leathery to membranous, sessile, rough haired, whitish, zoned, whitish, finally tawny.
1075	...	S.A.	...	...	N.S.W.	Q.	...	Trunks	Leathery, flattened, tapering behind, zoned, velvety, tawny.
1076	...	...	T.	V.	N.S.W.	Q.	...	Trunks	Sessile, leathery. Cap 3–5 inches across, more or less lobed, downy, zoned, reddish cinnamon or brownish.
1077	...	...	...	...	...	Q.	...	Trunks	Sessile, leathery, flexible, tapering behind, zoned, soft and spongy, pale brown.
1078	...	...	...	V.	...	...	...	Sticks	Gregarious. Cap funnel to fan shaped, delicately velvety, somewhat zoned, fawn. Stem thin.
1079	...	...	...	V.	...	Q.	...	Among grass	Cap funnel shaped, thin, rather membranous, smooth, shining, zoned, brown. Stem central, thin.
1080	...	...	T.	...	...	Q.	B.	Bark and dead branches	Leathery, thickish, expanded, silky, zoned, hoary.
1081	...	...	...	V.	N.S.W.	Q.	...	Wood, &c.	Gregarious. Cap funnel shaped, zoned and furrowed, velvety, bay brown. Stem very short.
1082	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Trunks, branches, &c.	Tough, soft. Cap spread out, zoned, silky and downy. Hymenium more or less purple.
1083	...	...	T.	...	...	...	...	Wood	Cartilaginous to leathery. Cap fan shaped, shining, reddish umber, silky streaked. Stem short, $\frac{1}{2}$ inch high.
1084	...	...	...	...	...	Q.	...	Wood	Crowded, thin, fan shaped, silky, shining, bay brown, many zoned, usually split into numerous lobes.
1085	W.A.	...	...	...	...	...	...	Base of living shrubs	Cap thick, white within, spread out, rough haired, whitish tawny.
1086	...	...	...	...	...	...	...	Wood	Leathery to membranous, mouse coloured. Cap spread out, cup shaped, then flattened, margin fringed. Hymenium net veined.
1087	W.A.	...	...	...	N.S.W.	...	B.	Trunks	Corky, rigid, becoming red when bruised. Cap spread out and partly reflexed, bay brown.
1088	...	...	...	...	...	Q.	...	Trunks	Membranous, somewhat tufted, sessile, flattened, semicircular, zoned, rust coloured to umber.
1089	...	...	...	...	...	Q.	...	Branches	Cap circular, rigid, wrinkled, yellowish tan, zoned and slightly reflexed.
1090	...	S.A.	T.	V.	N.S.W.	Q.	B.	Ground	Snow white, funnel shaped, soon discoloured, rough, with radiating processes projecting from surface.
1091	...	S.A.	T.	V.	N.S.W.	Q.	B.	Trunks, &c.	Leathery, spread out or reflexed, hairy, rusty, blood red where scratched, with dark-coloured hymenium.
1092	...	...	...	V.	...	...	...	Bark	White or pale ochrey, in hard now and then confluent pustules. Evidently immature condition of some species.
1093	...	...	...	...	...	Q.	...	Wood	Cap spoon shaped or somewhat fan shaped, bristly behind, smooth in front. Stem yellowish, velvety, lateral.
1094	...	...	...	...	N.S.W.	...	...	Wood	Leathery, spread out, reflexed, wavy, roughly striated, rusty brown.
1095	...	...	...	...	...	Q.	...	Wood	Thick, firm, isolated patches, ochrey or lemon yellow. Including <i>S. sulphureum</i> , Fries.
1096	...	S.A.	...	V.	...	...	...	Dead wood and branches	Reflexed, lobed, crisped, sulphur coloured, rough haired, somewhat spongy.
1097	...	...	...	V.	...	...	...	Among moss and on ground	Tufted, growing together, sessile, soft, spongy, flexible, reflexed, pale umber, velvety, concentrically zoned. Species' name pre-occupied by Cooke himself.
1098	...	S.A.	...	V.	...	Q.	...	Trunks	Cap funnel shaped, downy, somewhat zoned, pale. Stem tapering downwards.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
85. STEREOUM.—Pers., Obs. Myc. I. 35 (1796).—					
1099	...	VI. 7316	<i>S. umbrinum</i> ...	Fries, Pl. Preiss. II. 137 (1846) ...	Umber sterium ...
1100	1004	„ 7367	<i>S. vellereum</i> ...	Berk., Fl. N. Zeal. 183 (1855) ...	Woolly sterium ...
1101	1003	„ 7276	<i>S. versicolor</i> ...	Fries, Epicr. 547 (1838) ...	Variously coloured sterium ...
1102	1026	„ 7376	<i>S. versiforme</i> ...	Berk. and Curt., Grev. I. 164 (1873) ...	Variously shaped sterium ...
1103	...	„ 7338	<i>S. vittæforme</i> ...	Fries, Pl. Preiss. II. 137 (1846) ...	Striped sterium ...
1104	1025	„ 7286	<i>S. vorticosum</i> ...	Fries, Obs. II. 275 (1818) ...	Obscurely zoned sterium ...
86. HYMENOCHÆTE.—Lev., Ann. Sci. Nat.					
1105	...	VI. 7476	<i>H. Archeri</i> ...	Cooke, Grev. VIII. 149 (1880) ...	Archer's hymenochæte ...
1106	1035	„ 7438	<i>H. cacao</i> ...	Berk., Linn. Trans. I. 403 (1879) ...	Chocolate-brown hymenochæte...
1107	1044	„ 7461	<i>H. crassa</i> ...	Berk., Grev. VIII. 148 (1885) ...	Thick-margined hymenochæte ...
1108	1040	„ 7473	<i>H. innata</i> ...	Cooke and Mass., Grev. XV. 99 (1887) ...	Innate hymenochæte ...
1109	1048	IX. 927	<i>H. Kalchbrenneri</i> ...	Mass., Mon. Thel. Linn. Journ. XXVII. 116 (1891)	Kalchbrenner's hymenochæte ...
1110	1041	VI. 7449	<i>H. Mougeotii</i> ...	Cooke, Grev. VIII. 1147 (1880) ...	Mougeot's hymenochæte ...
1111	1047	„ 7464	<i>H. olivacea</i> ...	Cooke, Grev. XIV. 11 (1885) ...	Dark-olive hymenochæte ...
1112	1034	„ 7441	<i>H. phœa</i> ...	Cooke, Grev. VIII. 146 (1880) ...	Dusky hymenochæte ...
1113	1046	„ 7462	<i>H. purpurea</i> ...	Mass., Mon. Thel. Linn. Journ. XXVII. 115 (1891)	Purple-gilled hymenochæte ...
1114	...	„ 7440	<i>H. rigidula</i> ...	Berk. and Curt., Linn. Journ. X. 334 (1869) ...	Rigid hymenochæte ...
1115	1043	„ 7467	<i>H. rhubarbarina</i> ...	Cooke, Grev. VIII. 148 (1880) ...	Rhubarb-gilled hymenochæte ...
1116	1033	„ 7427	<i>H. rubiginosa</i> ...	Lev., Ann. Sci. Nat. Ser. 3, V. (1846) ...	Rusty hymenochæte ...
1117	1045	„ 7312	<i>H. Schomburgkii</i> ...	Mass., Mon. Thel. Linn. Journ. XXVII. 115 (1891)	Schomburgk's hymenochæte ..
1118	1037	„ 7444	<i>H. spadicea</i> ...	Berk. and Br., Linn. Journ. XIV. 68 (1875) ...	Bay-brown hymenochæte ...
1119	1036	„ 7436	<i>H. strigosa</i> ...	Berk. and Br., Linn. Journ. XIV. 68 (1875) ...	Strigose hymenochæte ...
1120	1042	„ 7428	<i>H. tabacina</i> ...	Lev., Ann. Sci. Nat. Ser. 3, V. 151 (1846) ...	Dark-brown hymenochæte ...
1121	1039	IX. 923	<i>H. tasmanica</i> ...	Mass., Mon. Thel. Linn. Journ. XXVII. 105 (1891)	Tasmanian hymenochæte ...
1122	1038	VI. 7443	<i>H. tenuissima</i> ...	Berk., Linn. Journ. XIV. 67 (1875) ...	Very thin hymenochæte ...
87. CORTICIUM.—Fries, Epicr. 556 (1838).—Thelephora, Hypochnus,					
1123	1076	VI. 7786	<i>C. anthocroum</i> ...	Fries, Hym. Eur. 661 (1874) ...	Bright-coloured corticium ...
1124	1070	„ 7528	<i>C. arachnoideum</i> ...	Berk., Ann. Nat. Hist. 345 (1844) ...	Arachnoid corticium ...
1125	1069	„ 7668	<i>C. Archeri</i> ...	Berk., Fl. Tasm. II. 260 (1860) ...	Archer's corticium ...
1126	1079	„ 7540	<i>C. atrovirens</i> ...	Fries, Hym. Eur. 651 (1874) ...	Dark-green corticium ...
1127	1068	„ 7552	<i>C. Auberianum</i> ...	Mont., Crypt. Cnba 372 (1842) ...	Auber's corticium ...
1128	1064	„ 7596	<i>C. calceum</i> ...	Fries, Hym. Eur. 652 (1874) ...	Chalky corticium ...
1129	1074	IX. 946	<i>C. cinnabarinum</i> ...	Mass., Mon. Thel. Linn. Journ. XXVII. 140 (1891)	Vermilion-red corticium ...
1130	1078	VI. 7539	<i>C. cæruleum</i> ...	Fries, Hym. Eur. 651 (1874) ...	Blue corticium ...
1131	1080	„ 7616	<i>C. comedens</i> ...	Fries, Hym. Eur. 656 (1874) ...	Wasting corticium ...
1132	1067	„ 7527	<i>C. lacteum</i> ...	Fries, Hym. Eur. 649 (1874) ...	Milk-white corticium ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Thelephora, Auricularia, Corticium, Elvella—continued.									
1099	W.A.	...	...	...	...	...	...	Bark of <i>Banksia</i> ...	Sessile, hemispherical, leathery, rather flaccid, fan shaped, undulating radiations from base towards margin, umber.
1100	...	...	...	V.	...	...	...	Branches and dead twigs	Spreading, with broad lobed zoned margin, dirty white above, and clothed behind with coarse tow-like fibres.
1101	...	...	...	V.	...	...	...	Trunks ...	Spread out, reflexed, fan shaped, thin, rigid, with raised concentric zones, variously coloured, or whitish to brown.
1102	...	...	...	V.	...	...	...	Dead branches, &c.	Small, at first circular, often running together, thickish, bright brown. Margin thin, upraised.
1103	W.A.	...	...	...	...	...	...	Bark of <i>Acacia</i> ...	Entire, leathery, rigid, somewhat bell shaped, with elevated concentric lines, becoming mud colour.
1104	...	...	...	...	...	Q.	B.	Bark and wood ...	Leathery, spread out, reflexed, obscurely zoned, rough haired, pale.

Ser. 3, V. 151 (1846).—*Thelephora, Stereum, Corticium.*

1105	...	...	T.	...	...	...	...	Rotten wood	...	Spreading, without distinct margin, soon detached, wine colour to brown, bristly, umber within.
1106	...	...	...	...	...	Q.	...	Wood	...	Dense dark-brown circular patches closely overlapping, cap fan shaped, deeply lobed and folded, furrowed with a few zones, velvety.
1107	...	...	...	...	N.S.W.	...	B.	Trunks	...	Leathery, velvety, pale reddish brown, margin thick, at length free.
1108	...	...	...	...	...	Q.	...	Wood	...	Thin, innate, scarcely distinct from underlying matrix, externally fawn colour, internally brick red.
1109	...	...	...	V.	...	Q.	...	Dead trunks of <i>Eucalypts</i>	...	Brown, rather thickly membranous, broadly spread out, loosely adherent to matrix.
1110	...	...	T.	V.	N.S.W.	...	...	Trunks of <i>Pinus picea</i> , &c.	...	Broadly spread out, dry, attached, dark blood red.
1111	...	...	...	...	...	Q.	...	Branches...	...	Spread out, dark olive, rough, velvety. Margin thinner and paler.
1112	...	...	...	V.	N.S.W.	Q.	...	Bark and wood	...	Semicircular, sessile, thin, leathery, flexible, concentrically zoned, shortly hairy, somewhat velvety, bay brown.
1113	...	...	...	V.	N.S.W.	...	...	Wood	...	Broadly spread out, closely attached, texture soft and spongy.
1114	...	...	...	...	N.S.W.	...	...	Wood	...	Broadly spread out, reflexed, rigid, thickish, zoned, velvety, bay brown.
1115	...	...	...	...	...	Q.	...	Wood	...	Broadly spread out, closely attached. Hymenium velvety, rhubarb colour.
1116	W.A.	...	T.	...	...	Q.	B.	Hard wood, posts, &c.	...	Leathery, rigid, spread out, reflexed, velvety, reddish brown to bay brown.
1117	...	S.A.	...	...	...	Q.	...	Wood	...	Somewhat circular, then shell shaped, umber, somewhat zoned in front, velvety, and about 1 inch broad.
1118	...	...	...	...	N.S.W.	Q.	...	Wood	...	Thin, elastic when dry, semicircular or circular, attached behind, zoned, rusty to bay brown.
1119	...	...	...	...	...	...	...	Wood	...	Semicircular, thin, lobed, zoned, bay brown tinged with purple, strigose or rough haired, 1 to 3 inches across.
1120	...	...	...	V.	N.S.W.	...	B.	Trunks	...	Somewhat leathery, thin, flaccid, covering underside of fallen logs. Cap spread out, reflexed, silky, rusty. Margin golden yellow.
1121	W.A.	...	T.	V.	...	...	...	Wood	...	Broadly spread out, crustaceous, rather thick. Margin thinner and paler, and sometimes slightly curled.
1122	...	...	...	...	...	Q.	...	Wood and bark	...	Sessile, very thin and flexible, 1 inch or more long, rusty to tawny, zoned, clothed with coarse down.

Peniophora, Hymenochaete, Coniophora, Stereum, Auricularia.

1123	...	...	...	...	...	Q.	B.	Bark	...	Broadly spread out, membranous, brick red or rosy, turning pale.
1124	...	S.A.	T.	V.	...	Q.	B.	Wood, bark, &c.	...	Delicate, spread out, spider web-like patches, snow white, often remaining barren.
1125	...	...	T.	...	...	Q.	...	Charred wood	...	Pale red, white within, rather thick, cracking.
1126	...	...	...	...	...	Q.	B.	Rotten wood, sticks, &c.	...	Irregularly spread out, thin, blackish or verdigris green.
1127	...	...	...	V.	...	...	...	Branches...	...	Circular at first, thin, mealy, from white to pale ochre.
1128	...	...	T.	...	...	...	B.	Wood and branches	...	Broadly spread out, thin, waxy, smooth, white, hard, sometimes continuous, sometimes much cracked.
1129	...	...	...	...	N.S.W.	...	...	Wood	...	Spreading for several inches, rather thin, without distinct margin. Hymenium waxy, vermilion.
1130	...	...	...	...	N.S.W.	Q.	B.	Wood, branches, &c.	...	Irregularly spread out, attached, downy, bright blue. Said to be phosphorescent.
1131	W.A.	...	...	...	...	...	B.	Branches...	...	Spread out and developed beneath the bark, which eventually is ruptured, lilac then pale.
1132	...	...	...	V.	...	...	B.	Wood	...	Broadly spread out, somewhat membranous and usually broken up, whitish, ochrey or buff when dry.

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
87. CORTICIUM.—Fries, <i>Epicr.</i> 556 (1838).—Thelephora, Hypochnus,					
1133	1072	VI. 7530	<i>C. læve</i> ...	Pers., <i>Tent. Disp.</i> 30 (1797) ...	Even corticium ...
1134	...	„ 7597	<i>C. lividum</i> ...	Pers., <i>Obs.</i> I. 38 (1796) ...	Livid corticium ...
1135	...	„ 7643	<i>C. Marescalchiauum</i> ...	<i>Marc. and Sacc.</i> , <i>Syll.</i> VI. 633 (1888) ...	Marescalchi's corticium ...
1136	1075	„ 7676	<i>C. miniatum</i> ...	Cooke, <i>Grev.</i> IX. 2 (1880) ...	Vermilion corticium ...
1137	1073	„ 7609	<i>C. nudum</i> ...	Fries, <i>Epicr.</i> 564 (1838) ...	Naked corticium ...
1138	1071	...	<i>C. penetrans</i> ...	Cooke and Mass., <i>Grev.</i> XIX. 90 (1891) ...	Penetrating corticium ...
1139	1065	VI. 7162	<i>C. sebaceum</i> ...	<i>Mass.</i> , <i>Mon. Thel. Linn. Journ.</i> XXVII. 127 (1891) ...	Waxy corticium ...
1140	1066	„ 7593	<i>C. simulans</i> ...	Berk. and Broome, <i>Linn. Journ.</i> XIV. 72 (1875) ...	Simulating corticium ...
1141	1077	...	<i>C. sulphurellum</i> ...	Cooke and Mass., <i>Grev.</i> XX. 35 (1891) ...	Sulphur-yellow corticium ...
88. PENIOPHORA.—Cooke, <i>Grev.</i> VIII. 20 (1879).—					
1142	1053	VI. 7158	<i>P. albo-marginata</i> ...	<i>Mass.</i> , <i>Mon. Thel. Linn. Journ.</i> XXV. 144 (1890) ...	White-margined peniophora ...
1143	1058	„ 7712	<i>P. bambusicola</i> ...	<i>Sacc.</i> , <i>Syll.</i> VI. 647 (1888) ...	Bamboo-loving peniophora ...
1144	1059	„ 7697	<i>P. carnea</i> ...	Cooke, <i>Grev.</i> VIII. 21 (1879) ...	Flesh-coloured peniophora ...
1145	1051	„ 7694	<i>P. cinerea</i> ...	Cooke, <i>Grev.</i> VIII. 20 (1879) ...	Ash-coloured peniophora ...
1146	..	„ 7707	<i>P. deglubens</i> ...	Berk., <i>Linn. Journ.</i> XVIII. 385 (1881) ...	Peeling peniophora ...
1147	1057	„ 7605	<i>P. incarvata</i> ...	<i>Mass.</i> , <i>Mon. Thel. Linn. Journ.</i> XXV. 147 (1890) ...	Bright peniophora ...
1148	1049	„ 7688	<i>P. papyrina</i> ...	Cooke, <i>Grev.</i> VIII. 20 (1879) ...	Papery peniophora ...
1149	1050	IX. 969	<i>P. puberula</i> ...	<i>Sacc.</i> , <i>Syll.</i> IX. 288 (1891) ...	Downy peniophora ...
1150	1056	VI. 7531	<i>P. rosea</i> ...	<i>Mass.</i> , <i>Mon. Thel. Linn. Journ.</i> XXV. 146 (1890) ...	Rosy peniophora ...
1151	1060	„ 7702	<i>P. sparsa</i> ...	Cooke, <i>Grev.</i> VIII. 21 (1879) ...	Scattered peniophora ...
1152	1052	„ 7695	<i>P. tephra</i> ...	Cooke, <i>Grev.</i> VIII. 20 (1879) ...	Ashy peniophora ...
1153	1055	„ 7477	<i>P. vinosa</i> ...	<i>Mass.</i> , <i>Mon. Thel. Linn. Journ.</i> XXV. 145 (1890) ...	Vinous peniophora ...
89. CONIOPHORA.—De Candolle, <i>Fl. Fr.</i> V. 34 (1815).—					
1154	1084	VI. 7719	<i>C. luteo-cincta</i> ...	Cooke, <i>Grev.</i> VIII. 89 (1880) ...	Yellow-margined conioophora ...
1155	1083	„ 7724	<i>C. membranacea</i> ...	<i>D. C.</i> , <i>Fl. Fr.</i> V. 634 (1815) ...	Membranous conioophora ...
1156	1086	„ 7683	<i>C. murina</i> ...	<i>Mass.</i> , <i>Mon. Thel. Linn. Journ.</i> XXV. 135 (1890) ...	Mouse-coloured conioophora ...
1157	1082	„ 7723	<i>C. olivacea</i> ...	Cooke, <i>Grev.</i> VIII. 89 (1880) ...	Olive-coloured conioophora ...
1158	1085	„ 7535	<i>C. sulphurea</i> ...	<i>Mass.</i> , <i>Mon. Thel. Linn. Journ.</i> XXV. 132 (1890) ...	Sulphur-coloured conioophora ...
1159	1081	„ 7203	<i>C. viridis</i> ...	Cooke, <i>Grev.</i> VIII. 89 (1879) ...	Green conioophora ...
90. ALEURODISCUS.—Rabh.—					
1160	1062	IX. 930	<i>A. albidus</i> ...	<i>Mass.</i> , <i>Grev.</i> XVII. 55 (1889) ...	White aleurodiscus ...
1161	1061	VI. 7506	<i>A. amorphus</i> ...	<i>Rabh.</i> , <i>Fung. Eur.</i> No. 1824 ...	Shapeless aleurodiscus ...
1162	1063	„ 7510	<i>A. tabacinus</i> ...	Cooke, <i>Grev.</i> XIV. 11 (1885) ...	Dark-brown aleurodiscus ...
91. CYPHELLA.—Fries, <i>Syst. Myc.</i> II. 201 (1823).—					
1163	1087	VI. 7817	<i>C. albo-violascens</i> ...	<i>Karst.</i> , <i>Fung. Fenn. Exs.</i> No. 715 ...	Pale-violet cyphella ...
1164	1091	...	<i>C. australiensis</i> ...	Cooke, <i>Grev.</i> XX. 9 (1891) ...	Australian cyphella ...

OF AUSTRALIAN FUNGI—*continued.*

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Peniophora, Hymenochaete, Coniophora, Stereum, Auricularia— <i>continued.</i>									
1133	...	...	T.	V.	...	Q.	B.	Wood, bark, &c. ...	Spread out over fibrils, membranous, separating from matrix, pinkish or pale.
1134	...	...	...	...	...	Q.	B.	Rotten wood ...	Expanded, soft, waxy, thin, irregular, smooth, bluish grey tinged with purple.
1135	...	...	...	...	...	Q.	...	Wood ...	Hymenium powdery, rhubarb colour, broken into little elevations.
1136	...	...	...	...	N.S.W.	Q.	...	Bark ...	Spread out, vermilion, margin fringed, whitish. When dry hymenium resembles patches of dried blood (Cooke).
1137	...	...	...	...	...	Q.	B.	Bark of <i>Citrus</i> ...	Waxy, cracking, flesh colour to pale. Margin decided, smooth.
1138	...	...	...	V.	...	...	...	Rotten wood, &c. ...	White, spreading, encrusting, thick, soft, with profuse penetrating mycelium.
1139	...	...	T.	...	...	...	B.	On ground and running up stems of plants	Spread out, fleshy or waxy, turning hard, incrusting and variable in form, whitish.
1140	...	...	...	V.	...	...	...	Running over mosses and twigs	Soft, tawny, arising from a white membranous woolly mycelium.
1141	...	...	...	V.	...	...	...	Dead branches ...	Broadly spread out, usually forming a thin, powdery, bright sulphur-yellowish stratum.

Corticium, Stereum, Thelephora, Hymenochaete.

1142	...	...	...	V.	N.S.W.	...	...	Bark and wood ...	Very broadly spread and confluent, umber, velvety in centre, and margin white and downy.
1143	...	...	...	...	...	Q.	...	Rotting bamboo ...	Roundish patches, yellowish tan, tough, thin, fringed, cracked in drying.
1144	...	...	...	...	...	...	...	On <i>Pinus contorta</i> ...	Broadly spread, ochrey to flesh colour. Margin white and with loose fibres.
1145	...	...	...	V.	...	...	B.	Branches... ...	Waxy, cracking, confluent, ash coloured or brownish.
1146	...	...	...	...	...	Q.	...	Trunks ...	Pale, spreading, thick, peeling, downy.
1147	W.A.	...	...	...	...	Q.	B.	Thin layer on wood and bark	Broadly spread, thin, waxy, radiating at margin, red or orange with pink bloom.
1148	...	...	...	V.	...	Q.	...	Bark ...	Very thin, leathery to papery, very broadly spread out and reflexed, rough haired, ash coloured, and margin tawny.
1149	...	...	...	...	...	Q.	...	Wood ...	Membranous, leathery, broadly spread out, ochrey yellow, downy. Margin reflexed.
1150	...	...	T.	...	...	...	B.	Wood and bark ...	Spread out, clear rose pink, turning white, rather fleshy, fringed. Beautiful species, sometimes in scattered patches.
1151	...	...	...	V.	...	Q.	...	Bark ...	Minute, snowy white, rather circular scattered patches.
1152	...	S.A.	...	...	...	...	...	Bark ...	Spread out, margin reddish brown, notched, free, downy.
1153	W.A.	...	...	...	...	...	...	Wood and bark ...	Isolated round patches becoming confluent and widely extending, irregularly lobed, wine colour or dark brown.

Thelephora, Coniophora, Auricularia, Merulius, Corticium, Hypoclinus.

1154	...	...	...	V.	...	...	...	On ground and bark	Spreading. Hymenium brown, powdery, yellow at circumference.
1155	...	...	...	V.	...	...	B.	Walls, &c. ...	In thin patches, foot or more in diameter, fragile, yellowish.
1156	...	...	...	V.	...	...	...	Branches, &c. ...	Widely spreading, at length breaking up, mouse coloured.
1157	...	...	...	V.	...	Q.	B.	Decayed pine ...	Widely spreading, membranous. Margin whitish, fringed. Hymenium dark olive, powdery.
1158	...	...	T.	...	...	...	B.	Bark, wood, leaves, &c.	Spreading, often spongy, passing into radiating cord-like branching sulphur-coloured threads.
1159	...	...	T.	...	...	...	...	Rotten wood ...	Developed beneath bark, spreading, downy, greenish.

Corticium, Cyphella, Peziza.

1160	...	...	...	...	...	Q.	...	Branches... ...	Pure white, at first scattered, becoming confluent and forming irregular patches, cup shaped at first, then expanded and flattened.
1161	...	...	...	...	...	Q.	...	Fir trunks and branches	Waxy, tough, rather leathery, cup shaped, then expanded, confluent, white, downy.
1162	...	...	...	V.	N.S.W.	...	...	Wood ...	Gregarious, cup shaped, cap somewhat elliptical, wrinkled, crisped, brown, downy.

Peziza, Thelephora, Cantharellus.

1163	...	S.A.	...	V.	...	...	B.	Bark and wood, branches of Vine	Somewhat corky, sessile or nearly so, spherical to hemispherical, white. Hymenium pale violet.
1164	...	...	...	V.	...	...	...	Bark ...	Gregarious, cup shaped, sessile, pale, with closely pressed silky hairs.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
91. CYPHELLA.—Fries, Syst. Myc. II. 201 (1823).—					
1165	1088	VI. 7856	<i>C. capula</i> ...	<i>Fries</i> , Epicr. 568 (1838) ...	Cup-like cyphella ...
1166	...	...	<i>C. filicola</i> ...	<i>Cooke</i> , Grev. XIV. 129 (1886) ...	Fern-growing cyphella ...
1167	...	...	<i>C. longipes</i> ...	<i>Cooke</i> and <i>Mass.</i> , Grev. XXI. 38 (1892) ...	Long-stalked cyphella ...
1168	1092	VI. 7889	<i>C. muscigena</i> ...	<i>Fries</i> , Epicr. 567 (1838) ...	Moss-growing cyphella ...
1169	...	7906	<i>C. parasitica</i> ...	<i>Berk.</i> and <i>Br.</i> , Linn. Journ. XIV. 74 (1875) ...	Parasitic cyphella ...
1170	1090	IX. 1006	<i>C. polycephala</i> ...	<i>Sacc.</i> , Hedw. 126 (1889) ...	Many-headed cyphella ...
1171	...	1010	<i>C. Schneideri</i> ...	<i>Berk.</i> and <i>Br.</i> , Linn. Trans. II. 220 (1887) ...	Schneider's cyphella ...
1172	1089	VI. 7868	<i>C. villosa</i> ...	<i>Karst.</i> , Myc. Fenn. III. 325 (1871) ...	Villous cyphella ...
92. SOLENIA.—Hoffm.,					
1173	916	VI. 6589	<i>S. candida</i> ...	<i>Pers.</i> , Tent. Disp. 36 (1797) ...	White solenia ...
1174A	917	„ 6594	<i>S. anomala</i> , <i>Fries</i> , var. <i>ochracea</i> (<i>Mass.</i>)	<i>Brit.</i> , Fung. Fl. I. 144 (1892) ...	Ochrey solenia ...
1175	917*	„ 6596	<i>S. sulphurea</i> ...	<i>Sacc.</i> and <i>Ellis.</i> , Mich. II. 564 (1882) ...	Sulphur solenia ...
93. LACHNOCLADIUM.—Lev., Orb. Dict. VIII. 487 (1849).—					
1176	969	VI. 8177	<i>L. Brasiliense</i> ...	<i>Sacc.</i> , Syll. VI. 738 (1888) ...	Brazilian lachnocladium ...
1177	973	„ 8018	<i>L. flagelliforme</i> ...	<i>Cooke</i> , Handb. Aust. Fung. 179 (1892) ...	Whip-shaped lachnocladium ...
1178	968	„ 8175	<i>L. furcellatum</i> ...	<i>Sacc.</i> , Syll. VI. 738 (1888) ...	Forked lachnocladium ...
1179	972	„ 8183	<i>L. rameale</i> ...	<i>Berk.</i> and <i>Broome</i> , Linn. Journ. XIV. 67 (1875) ...	Branch-growing lachnocladium ...
1180	970	„ 8180	<i>L. semivestitum</i> ...	<i>Berk.</i> and <i>Cnrt.</i> , Grev. I. 161 (1873) ...	Half-clothed lachnocladium ...
1181	971	„ 8188	<i>L. setulosum</i> ...	<i>Sacc.</i> , Syll. VI. 740 (1888) ...	Bristly lachnocladium ...
1182	...	IX. 1043	<i>L. simulans</i> ...	<i>Berk.</i> and <i>Broome</i> , Linn. Trans. II. 219 (1887) ...	Simulating lachnocladium ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Peziza, Thelephora, Cantharellus—continued.									
1165	...	...	T.	...	...	Q.	B.	Herb stems ...	Membranous, obliquely bell shaped, running down into oblique stem, whitish.
1166	...	...	..	V.	...	...	...	Fronds of <i>Adiantum</i> ...	Whitish, somewhat discoid, concave, smooth. Margin entire or two cusped.
1167	...	...	...	...	...	Q.	...	Logs and stems of trees in wet scrubs	Gregarious, membranous, white, cap narrow, funnel shaped, tapering into long thin curved stem.
1168	...	...	T.	V.	...	...	B.	On larger mosses ...	Membranous, soft, flattened nearly semicircular, white.
1169	...	...	...	...	...	...	...	Parasitic on some <i>sphæria</i>	Minute, cup shaped, snow white, externally hairy. Margin inflexed.
1170	...	S.A.	...	...	N.S.W.	...	...	Herb stems, <i>Senecio hypoleucus</i>	Gregarious. Caps closely joined in common base, whitish brown, urn shaped.
1171	...	...	...	...	...	Q.	...	Wood ...	Gregarious, tubular, membranous, pale yellow.
1172	...	...	...	V.	...	Q.	B.	Herbstems and rotting stems of Castor Oil plant	Sessile, dry, spherical, white, covered with snow-white persistent villous down.

Bot. Tasch. 68 (1795).

1173	...	...	...	...	...	Q.	...	Rotten wood	...	Scattered, cylindrical, smooth, white.
1174a	...	...	T.	...	...	...	B.	Rotten wood	...	Scattered, club shaped to cylindrical, downy, ochrey.
1175	...	...	...	V.	...	...	...	Branches	...	Thickly crowded, minute, cup-like, shortly stalked, sulphur coloured, with rough hairs.

Clavaria, Eriocladus.

1176	...	...	...	...	...	Q.	...	Trunks ...	...	Very shortly stalked, much branched. Branches tapering, forked, ochrey white.
1177	...	...	...	...	...	Q.	...	Bare ground	...	Very much branched, divided to the base. Branches tufted, cylindrical, elongated, forked.
1178	...	...	...	...	...	Q.	...	Rotten wood	...	Ascending, somewhat rusty. Branches solid, repeatedly forked, distaut, tough, velvety.
1179	...	...	...	...	...	Q.	...	Branches, &c.	...	Dark purple, thread-like, forked, encrusting fresh branches, leaf stalks and leaves.
1180	...	...	...	...	...	Q.	...	Ground ...	...	Delicate, repeatedly forked. Branches downy.
1181	W.A.	...	...	...	...	...	...	Ground ...	...	Small, ochrey. Stem short, irregularly divided. Branches compressed, forked, downy.
1182	...	...	...	...	...	Q.	...	Ground ...	...	Dark brown when dry, downy. Stem simple below, repeatedly branched above. Branches slender, tips forked.

ORDER V.—CLAVARIACEÆ, CORDA.

94. Sparassis, Fries.

Number.	Corda's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
---------	-----------------	--------------------	------------------	---------------------	---------------

ORDER V.—CLAVARIACEÆ,

94. SPARASSIS.—Fries,

1183	1094	VI. 7923	<i>S. crispa</i>	Fries, S.M. I. 465 (1821)	Crisped sparassis
------	------	----------	-------------------------	----------------------------------	--------------------------

95. CLAVARIA.—Linn.,

1184	1110	VI. 7970	<i>C. abietina</i>	Pers., Comm. 46 (1797)	Fir clavaria
1185	1125	" 8077	<i>C. Archeri</i>	Berk., Fl. Tasm. II. 261 (1860)	Archer's clavaria
1186	1124	" 8072	<i>C. argillacea</i>	Fries, S.M. I. 482 (1821)	Clay-coloured clavaria
1187	1131	" 8108	<i>C. aurantia</i>	Cooke and Mass., Grev. XVI. 33 (1887)	Orange clavaria
1188	1108	" 7963	<i>C. aurea</i>	Schaeff. 287 (1762)	Golden clavaria
1189	1096	" 7931	<i>C. botrytes</i>	Pers., Syn. 587 (1801)	Clustered clavaria
1190	1100	" 7942	<i>C. cinerea</i>	Bull, Champ. 354 (1791)	Ash-coloured clavaria
1191	1119	" 8039	<i>C. Colensoi</i>	Berk., Fl. N. Zeal. 186 (1855)	Colenso's clavaria
1192	1099	" 7941	<i>C. coralloides</i>	Linn., Sp. Pl. 1182 (1753)	Coral-like clavaria
1193	1115	" 7991	<i>C. crispula</i>	Fries, S.M. I. 470 (1821)	Curled clavaria
1194	1102	" 7944	<i>C. cristata</i>	Pers., Syn. 591 (1801)	Crested clavaria
1195	1112	" 7973	<i>C. crucea</i>	Pers., Ic. and Desc. 36 (1798)	Saffron clavaria
1196	1097	" 7937	<i>C. fastigiata</i>	Linn., Sp. Pl. 1183 (1753)	Fastigate clavaria
1197	1095	" 7929	<i>C. flava</i>	Schaeff. 175 (1762)	Yellow clavaria
1198	1109	" 7967	<i>C. formosa</i>	Pers., Ic. and Desc. 11 (1798)	(Pollard fungus) Elegant clavaria
1199	1127	" 8080	<i>C. fragilis</i>	Holmsk. I. 7 (1818)	Brittle clavaria
1200	1122	" 8067	<i>C. fusiformis</i>	Sow., Fung. 224 (1797)	Spindle-shaped clavaria
1201	1113	" 7979	<i>C. grisea</i>	Pers., Comm. 44 (1797)	Dingy-grey clavaria
1202	1123	" 8069	<i>C. inæqualis</i>	Muell., Fl. Dan. 873	Unequal clavaria
1203	1129	" 8093	<i>C. juncea</i>	Fries, S.M. I. 472 (1821)	Rush-like clavaria
1204	1111	IX. 1024	<i>C. Kalchbrenneri</i>	F. v. M., Linn. Soc. N.S.W. 105 (1882)	Kalchbrenner's clavaria
1205	1104	VI. 7949	<i>C. Krombolzii</i>	Fries, Epier. 572 (1838)	Krombolz's clavaria
1206	1105	" 7951	<i>C. Kunzei</i>	Fries, S.M. I. 474 (1821)	Kunze's clavaria
1207	...	" 8064	<i>C. lactea</i>	Berk. and Br., Linn. Journ. XIV. 76 (1875)	Bright clavaria
1208	1117	" 8008	<i>C. lactissima</i>	Pers., in Linn. Journ. XVIII. 386 (1881)	Very bright clavaria
1209	1118	" 8029	<i>C. lorithamnus</i>	Berk., Linn. Journ. XIII. 169 (1873)	Crooked bush clavaria
1210	1101	IX. 1028	<i>C. lurida</i>	Kalch, Linn. Soc. N.S.W. 103 (1882)	Lurid clavaria
1211	1133	VI. 8116	<i>C. militina</i>	Berk., Hook., Journ. 140 (1852)	Crimson clavaria
1212	1134	" 8125	<i>C. mucida</i>	Pers., Comm. 2 (1797)	Musty clavaria

ARRANGEMENT OF GENERA (2).

95. *Clavaria*, Linn.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
CORDA, IC. FUNG. II. 35 (1838).									
S.M. I. 464 (1821).—Clavaria.									
1183	...	...	...	...	N.S.W.	...	B.	Wood ...	Very much branched, whitish. Branches intricate, zoneless, serrate. <i>Edible.</i>
Sp. Pl. 1182 (1753).—Typhula.									
1184	...	...	...	V.	...	...	B.	Firwood ...	Ochrey, very much branched, green where bruised. Trunk whitish, downy, thick. Branches crowded.
1185	...	...	T.	...	...	Q.	...	Ground ...	Tufted, short, orange, fan to club shaped, rather rough.
1186	...	...	...	V.	N.S.W.	Q.	B.	Ground ...	Tufted, fragile, clay coloured, pale. Clubs simple, variable. Stem shining, yellow.
1187	...	...	...	V.	...	...	...	Ground ...	Orange, quite simple, straight, thickened upwards into club, tapering downwards into stem.
1188	...	...	...	V.	N.S.W.	Q.	B.	Woods ...	Yellow. Trunk thick, divided into stout straight forked much-divided tapering branches.
1189	W.A.	...	T.	V.	N.S.W.	Q.	B.	Ground ...	Fragile, white. Trunk very thick, much branched. Branches swollen, and tips red.
1190	...	...	...	V.	...	...	B.	Woods ...	Fragile, stuffed, grey. Trunk rather thick, short, much branched, wrinkled.
1191	...	...	...	...	...	Q.	...	Bare ground and decayed wood	Small, branched from compressed base, branches erect, forked, brown when dry.
1192	...	...	...	V.	...	...	B.	Moist woods	Rather fragile, usually tufted, white, hollow inside. Trunk thick, repeatedly and irregularly branched.
1193	W.A.	...	...	...	...	...	B.	At base of trunks	Very much branched, tan to ochrey. Trunk thin, shaggy, and rooting. Branches flexuous, spreading.
1194	...	...	T.	V.	...	Q.	B.	Woods ...	Tough, stuffed, dingy white. Branches dilated above, and fringed.
1195	...	...	...	V.	...	...	B.	Woods ...	Minute, thin, saffron yellow. Stem naked, pale. Branches and branchlets somewhat forked.
1196	...	...	...	V.	N.S.W.	Q.	B.	Pastures ...	Yellow tufted, tough, much branched. Branches equal, short, spreading, branchlets twiggy.
1197	...	S.A.	T.	V.	N.S.W.	Q.	B.	Gravelly ground	Fragile. Trunk thick, fleshy, white, very much branched. Branches tapering, twiggy, yellow.
1198	...	...	...	V.	N.S.W.	Q.	B.	Woods ...	Trunk thick, whitish, very much branched. Branches rosy orange, branchlets yellowish.
1199	...	...	...	V.	...	...	B.	Ground ...	In bundles, very delicate, yellow above, white below, sometimes entirely white. Clubs hollow, variable.
1200	...	...	...	...	...	Q.	B.	Among grass	Tufted and run together, yellow, soon hollow. Clubs somewhat spindle shaped, simple and toothed.
1201	...	...	...	V.	...	...	B.	Ground in woods	Grey, firm, fragrant. Trunk thick, whitish. Branches tapering and dingy grey, as well as branchlets.
1202	...	...	T.	V.	N.S.W.	...	B.	Among sand	Tufted, yellow, fragile. Clubs various, simple or forked. Apex jagged.
1203	...	...	T.	V.	...	...	B.	Among dead leaves	Gregarious. Thin, thread-like, flabby, hollow, pale to reddish brown, with creeping base of fibrils.
1204	...	...	...	V.	...	...	...	Ground ...	Thin, pale, orange yellow, somewhat tufted. Trunk, thin, naked. Branches short, forked, or tufted.
1205	...	...	...	V.	...	...	B.	Ground ...	Fragile, tufted, white, sparingly branched. Branches rather compressed, obtuse.
1206	...	...	...	...	...	Q.	B.	Woods ...	Rather fragile, very much branched from the thin base, white. Branches elongated, crowded, repeatedly forked.
1207	...	...	...	V.	...	...	...	Red soil ...	Simple, tufted, acute, shining red, without evident stem.
1208	...	...	...	...	...	Q.	...	Ground ...	Tufted, orange, repeatedly forked, compressed. Apices dilated, sub-divided, tawny.
1209	...	...	...	V.	...	...	...	Ground ...	Pale amber. Branches straight. Apices shortly bifid, and rather acute.
1210	...	...	...	V.	...	...	...	Ground ...	Tufted, very much branched, dirty white. Trunks thin. Branches and branchlets crowded, elongated, tawny when dry.
1211	...	...	...	V.	...	Q.	...	Rotten wood	Gregarious, fragile. Clubs simple, acute, crimson.
1212	...	...	...	...	N.S.W.	Q.	...	Ground and musty wood	Gregarious, minute, simple or very sparingly branched, white, becoming yellowish, sometimes rosy.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
95. CLAVARIA.— <i>Linn.</i> ,					
1213	1136	...	<i>C. Muelleri</i> ...	Berk., Grev. XX. 10 (1891) ...	Mueller's clavaria ...
1214	1098	VI. 7938	<i>C. muscoides</i> ...	Linn., Sp. Pl. 1183 (1753) ...	Moss-like clavaria ...
1215	1130	„ 8096	<i>C. paludicola</i> ...	Libert., in Fries Hym. Eur. 678 (1874) ...	Marsh-growing clavaria ...
1216	1128	„ 8085	<i>C. pistillaris</i> ...	Linn., Sp. Pl. 1182 (1753) ...	Pestle-shaped clavaria ...
1217	1106	„ 7955	<i>C. plebeja</i> ...	Fries, Pl. Preiss. II. 137 (1846) ...	Plebeian clavaria ...
1218	1116	„ 7996	<i>C. portentosa</i> ...	Berk. and Br., Linn. Trans. II. 65 (1883) ...	Monstrous clavaria ...
1219	1107	„ 7957	<i>C. pyxidata</i> ...	Pers., Comm. (1797) ...	Box-like clavaria ...
1220	1132	„ 8112	<i>C. rhizomorpha</i> ...	Berk., Fl. Tasm. II. 261 (1860) ...	Root-shaped clavaria ...
1221	1121	„ 8063	<i>C. rosea</i> ...	Fries, S.M. I. 482 (1821) ...	Rosy clavaria ...
1221a	1121	...	<i>C. rosea</i> , var. <i>attenuata</i> ...	Fries, Obs. 2 (1818) ...	Attenuated clavaria ...
1222	1120	VI. 8062	<i>C. rufa</i> ...	Muell., Fl. Dan. 755 ...	Reddish clavaria ...
1223	1103	„ 7947	<i>C. rugosa</i> ...	Bull, Champ. t. 448 (1798) ...	Wrinkled clavaria ...
1224	1114	„ 7958	<i>C. stricta</i> ...	Pers., Comm. 45 (1797) ...	Straight clavaria ...
1225	...	„ 7953	<i>C. subtilis</i> ...	Pers., Comm. (1797) ...	Slender clavaria ...
1226	1135	...	<i>C. tasmanica</i> ...	Berk., in Herb., Grev. XX. 10 (1891) ...	Tasmanian clavaria ...
1227	1126	VI. 8079	<i>C. vermicularis</i> ...	Scop., in Fries, S.M. I. 484 (1821) ...	Worm-like clavaria ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Sp. Pl. 1182 (1753).—Typhula—continued.									
1213	...	...	...	V.	...	Q.	...	Ground ...	Simple, club shaped, white, slender, tapering below into thin cylindrical stem.
1214	...	...	...	V.	N.S.W.	...	B.	Pastures among moss	Tough, slender, yellow, two to three times forked. Stem thin. Branches curved, long, graceful.
1215	...	S.A.	...	...	...	...	...	Moist places among ferns	Small, simple, slightly compressed, rough, yellow, orange when dry. Clubs short, obtuse.
1216	...	...	...	V.	...	...	B.	Among grass	Simple, tall, fleshy, stuffed, yellow to reddish. Club large.
1217	W.A.	...	...	...	...	...	...	Sandy soil	Tough, white, becoming yellow. Trunk thickish. Branches and branchlets very much divided and crested at top.
1218	...	...	...	...	...	Q.	...	Among leaves	Whitish. Stem somewhat cylindrical, rough, repeatedly much branched. Apices elongated.
1219	...	...	...	V.	N.S.W.	...	B.	Rotten wood	Pale tan to reddish. Trunk thin, branched. Branches and branchlets all excavated at the tips into little cups.
1220	...	...	T.	...	...	...	...	Dead bark	Eruptent, confluent, chestnut red, nearly simple.
1221	...	...	...	V.	...	...	B.	Ground among moss	In bundles, fragile, rosy. Clubs stuffed, at length yellowish at tips, tapering downwards, and whitish.
1221A	...	...	...	V.	...	...	...	Ground among moss	Clubs tapering at apex.
1222	...	...	...	V.	...	Q.	B.	Among grass	Tufted, rufous. Clubs stuffed, thickened, sometimes bifid, acute.
1223	...	S.A.	...	V.	...	Q.	B.	Moist places	Tough, simple or sparingly branched, thickened upwards and wrinkled, white. Branches deformed.
1224	...	...	...	...	N.S.W.	Q.	B.	Trunks	Very much branched, pale yellow, turning brown when bruised. Trunk thickish. Branches and branchlets straight.
1225	...	...	...	V.	...	...	...	Ground in woods	Scattered, slender, somewhat tough, whitish, becoming pale, smooth at base.
1226	...	...	T.	...	...	...	...	Tree ferns, wood, &c.	Clubs simple, single or two or three together, sooty brown, base expanded in a white woolly web.
1227	...	...	...	V.	...	...	B.	Among grass	Tufted, fragile, white. Clubs tufted, simple, cylindrical, often flexuous or incurved.

ORDER VI.—TREMELLACEÆ, FRIES.

SUB-ORDER I.—Auriculariæ, Bref.—Basidia or spore-bearing

Genera (2)—

96. Auricularia.

SUB-ORDER II.—Tremelleæ, Bref.—Basidia globose or ovoid,

Genera (5)—

98. Exidia, Fries.

99. Ulocolla, Bref.

100. Tremella, Linn.

SUB-ORDER III.—Dacryomycetæ, Bref.—Basidia

Genera (3)—

103. Dacryomyces, Nees.

104. Guepinia, Fries.

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER VI.—TREMELLACEÆ,					
96. AURICULARIA.—Bull,					
1228	1145	VI. 8302	<i>A. albicans</i> ...	Berk., Linn. Journ. XIII. 170 (1873) ...	Whitish auricularia ...
1229	1144	„ 8293	<i>A. lobata</i> ...	Somm., Mag. Nat. Vid. (1827) ...	Lobate auricularia ...
1230	1143	„ 8294	<i>A. mesenterica</i> ..	Fries, Epier. 555 (1838) ...	Intestine-like auricularia ...
1231	1146	„ 8303	<i>A. minuta</i> ..	Berk., Hook., Lond. Journ. IV. 59 (1845) ...	Minute auricularia ...
1232	1147	„ 8305	<i>A. pusio</i> ..	Berk., Linn. Journ. XVIII. 386 (1881) ...	Small auricularia ...
97. HIRNEOLA.—Fries,					
1233	1150	VI. 8312	<i>H. auricula-judæ</i> ...	Berk., Ontl. 289 (1860) ...	Jew's ear hirneola ...
1234	1148	„ 8309	<i>H. auriformis</i> ...	Fries, Fung. Nat. 26 (1848) ...	Ear-shaped hirneola ...
1235	1151	„ 8319	<i>H. fusco-succinea</i> ...	Mont., Cuba 364 (1842) ...	Amber-brown hirneola ...
1236	1153	„ 8323	<i>H. hispidula</i> ...	Sacc. Syll. VI. 769 (1888) ...	Hispid hirneola ..
1237	1149	„ 8311	<i>H. polytricha</i> ...	Mont., in Bel. Voy. 154. ...	Many-haired hirneola ...
1238	1152	„ 8320	<i>H. rufa</i> ...	Fries, Fung. Nat. 27 (1848) ...	Reddish-brown hirneola ...
1239	1154	„ 8328	<i>H. vitellina</i> ...	Fries, Fung. Nat. 27 (1848) ...	Egg-yellow hirneola ...
1239A	„	„	<i>H. vitellina</i> , var. <i>tasmanica</i>	Berk., Fl. Tasm. II. 262 (1860) ...	Tasmanian hirneola ...
98. EXIDIA.—Fries,					
1240	1156	VI. 8352	<i>E. albida</i> ..	Bref., Unters. VII. 94 (1888) ...	Whitish exidia ...
1241	1155	„ 8347	<i>E. glandulosa</i> ...	Fries, S.M. II. 224 (1821) ...	Glandulous exidia ...
99. ULOCOLLA.—Bref.,					
1242	1157	VI. 8367	<i>U. foliacea</i> ...	Bref., Unters. VII. 98 (1888) ...	Leafy ulocolla ...
100. TREMELLA.—Linn.,					
1243	1158	VI. 8375	<i>T. frondosa</i> ...	Fries, S.M. II. 212 (1821) ...	Frondose tremella ...
1244	1160	„ 8384	<i>T. fuciformis</i> ...	Berk., Hook., Journ. 277 (1856) ...	Seaweed-like tremella ...
1245	1159	„ 8377	<i>T. lutescens</i> ...	Pers., Syn. 622 (1801) ...	Yellowish tremella ...

ARRANGEMENT OF GENERA (10).

bodies, elongated or fusoid, transversely divided.

97. Hirneola.

four partite in a cruciate manner when mature.

101. Seismosarca, Cooke.

102. Tremellodon, Pers.

cylindrical or club shaped, forked upwards.

105. Calocera, Fries.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			

FRIES, SYST. MYC. II. 207 (1823).

Herb. Fr. I. 36 (1787).—*Helvella*, *Thelephora*.

1228	...	...	...	...	...	Q.	...	Trunks	...	Circular, whitish, delicately downy beneath.
1229	...	...	...	...	N.S.W.	Q.	B.	Bark	...	Expanded, reflexed, lobed, variegated with hispid zones, or velvety, or smooth, dusky to whitish.
1230	W.A.	...	...	...	...	Q.	B.	Trunks	...	Reflexed, entire, shaggy, zoned, brownish to ashy. Hymenium ribbed and folded.
1231	W.A.	...	T.	...	...	...	...	Dead branches	...	Gregarious, expanded behind. Caps minute, lobed, hispid, zoned, tawny umber.
1232	...	...	...	...	...	Q.	...	Trunks	...	Cap attached behind, reflexed, white, downy, wrinkled. Margin lobed. Small but distinct species.

Pl. Homon. 93 (1825).—*Tremella*, *Peziza*, *Auricularia*, *Exidia*.

1233	...	...	T.	V.	N.S.W.	Q.	B.	Trunks	...	Hollow ear-like cups, flexuous, thin, blackish, with vein-like folds on both sides, downy beneath.
1234	...	...	...	...	...	Q.	...	Trunks	...	Tufted, stalked, glaucous brown. Caps semicircular, veined beneath. Stem short, twisted, lateral.
1235	W.A.	...	...	...	N.S.W.	Q.	...	Bark	...	Broad, sessile, shell shaped, then flattened. Margin wavy, internally netted with veins, amber brown.
1236	...	...	...	V.	...	Q.	...	Wood	...	Globose to bell shaped, oblique, sessile, internally dark brown, externally with short fawn woolly hairs.
1237	...	S.A.	...	V.	N.S.W.	Q.	B. (introduced)	Trunks	...	Cups hemispherical, expanded, ear shaped, shaggy, grey, produced into very short oblique stem.
1238	...	...	...	...	...	Q.	...	Trunks	...	Cap shaped, somewhat lateral, sessile, beset with tufted short reddish-brown bristles.
1239	...	...	T.	...	...	...	...	Trunks	...	Cap shaped, sessile, excavated. Hymenium egg yellow.
1239A	...	...	T.	...	...	...	...	Wood	...	Pale, circular, wavy, small. Stem short, compressed.

S.M. II. 220 (1823).—*Tremella*.

1240	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Branches and bark of dead logs	...	Ascending, tough, expanded, wavy, whitish, tawny when dry.
1241	W.A.	...	T.	...	...	...	B.	Trunks and woods	...	Expanded, rather flattened, thick, turning black, with conical pimples, ashy beneath and somewhat downy.

Unters. VII. 95 (1888).—*Tremella*.

1242	W.A.	...	T.	V.	...	Q.	B.	Old trunks	...	Tufted, wavy, cinnamon to flesh colour, folded at the base.
------	------	-----	----	----	-----	----	----	------------	-----	---

Sp. Pl. 1157 (1753).—*Elvella*, *Thelephora*.

1243	...	...	T.	...	...	...	B.	Old trunks	...	Tufted, large, yellow to pale, folded at base. Lobes folded and waved.
1244	...	...	...	...	...	Q.	...	Trunks	...	White, tufted, repeatedly lobed or forked. Lobes dilated in fan-like manner.
1245	...	S.A.	T.	V.	N.S.W.	Q.	B.	Fallen branches	...	Tufted, small, very soft, wavy, and folded, yellowish. Lobes entire, naked.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
100. TREMELLA.—Linn.,					
1246	1161	VI. 8387	<i>T. mesenterica</i> ...	Retz., Vet. Ak. Handl. 249 (1769) ...	Contorted tremella ...
1247	...	IX. 1071	<i>T. microscopica</i> ...	Berk. and Br., Linn. Trans. II. 220 (1887) ...	Microscopic tremella ...
1248	1162	VI. 8397	<i>T. olens</i> ...	Berk., Fl. Tasm. II. 262 (1860) ...	Scented tremella ...
1249	...	„ 8444	<i>T. sarcoides</i> ...	Fries, S.M. I. 215 (1821) ...	Flesh-coloured tremella ...
1250	1163	„ 8402	<i>T. viscosa</i> ...	Berk., Outl. 288 (1860) ...	Sticky tremella ...
101. SEISMOSARCA.—Cooke,					
1251	1164	IX. 1082	<i>S. hydrophora</i> ..	Cooke, Grev. XVIII. 25 (1889) ...	Watery seismosarca ...
102. TREMELLODON.—Pers.,					
1252	942	VI. 6862	<i>T. gelatinosum</i> ...	Pers., Myc. Eur. II. 172 (1822) ...	Gelatinous tremellodon ...
103. DACRYOMYCES.—Nees,					
1253	1167	VI. 8472	<i>D. deliquescent</i> ...	Duby, Bot. Gall. 729 (1822) ...	Deliquescent dacryomyces ..
1254	1165	„ 8469	<i>D. militinus</i> ...	Berk., Fl. Tasm. II. 263 (1860) ...	Vermilion dacryomyces ...
1255	1166	„ 8471	<i>D. rubrofusus</i> ...	Berk., Hook., Lond. Journ. IV. 61 (1845) ...	Reddish-brown dacryomyces ...
1256	1169	„ 8483	<i>D. sacchari</i> ...	Berk. and Br., Linn. Trans. II. 65 (1883) ...	Sugar-cane dacryomyces ...
1257	1171	„ 8502	<i>D. sclerotoides</i> ...	Berk., Fl. Tasm. II. 263 (1860) ...	Sclerotium-like dacryomyces ...
1258	1170	„ 8488	<i>D. seriatus</i> ...	Berk., Fl. Tasm. II. 263 (1860) ...	Seriate dacryomyces ...
1259	1168	„ 8473	<i>D. stillatus</i> ...	Nees, Syst. 89 (1816) ...	Dripping dacryomyces ...
104. GUEPINIA.—Fries,					
1260	1172	VI. 8514	<i>G. merulina</i> ...	Quelet Quelq. Esp. II. 11 (1878) ...	Merulius-like guepinia ...
1261	1173	„ 8518	<i>G. pezizæformis</i> ...	Berk., Hook., Lond. Journ. IV. 60 (1845) ...	Cup-shaped guepinia ...
1262	1174	„ 8520	<i>G. spathularia</i> ...	Fries, Elench. II. 32 (1828) ...	Spoon-shaped guepinia ...
105. CALOCERA.—Fries,					
1263	1139	VI. 8158	<i>C. cornea</i> ...	Fries, S.M. I. 485 (1821) ...	Horny calocera ...
1264	1138	IX. 1042	<i>C. digitata</i> ...	Cooke and Mass., Grev. XVII. 7 (1888) ...	Digitate calocera ...
1265	1142	VI. 8165	<i>C. glossoides</i> ...	Fries, S.M. I. 487 (1821) ...	Tongue-like calocera ...
1266	1137	„ 8154	<i>C. guepiniioides</i> ...	Berk., Hook., Lond. Journ. IV. 61 (1845) ...	Guepinia-like calocera ...
1267	1141	IX. 1041	<i>C. nutans</i> ...	Sacc. Hedw. 154 (1890) ...	Nodding calocera ...
1268	1140	VI. 8163	<i>C. stricta</i> ...	Fries, Epicr. 581 (1838) ...	Erect calocera ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Sp. Pl. 1157 (1753).— <i>Elvella</i> , <i>Thelephora</i> — <i>continued</i> .									
1246	W.A.	S.A.	...	V.	N.S.W.	Q.	B.	Dead branches	... Toughish, variable in form, folded and wavy, bright orange.
1247	...	...	...	V.	...	Q.	...	Leaves	... Minute, hemispherical, point-like, dark-green scattered spots on upper surface of leaf.
1248	...	...	T.	...	...	...	...	Rotten wood	... Irregular, gelatinous, pale, scented.
1249	...	...	...	V.	...	...	B.	Trunks	... Tufted, soft, viscid, pale-flesh colour, at first club shaped, then compressed. Lobed and folded. Conidial stage of <i>Ombrophila sarcoides</i> .
1250	...	...	T.	...	...	...	B.	Old wood	... Expanded, flattened, wavy, rather viscid, white to hyaline, brown when dry.
Grev. XVIII. 25 (1889).									
1251	...	...	...	...	N.S.W.	...	...	Wood	... Inflated, gelatinous, lobate, pale sooty brown, very soft and watery, covered with scattered coloured hairs.
Myc. Eur. II. 172 (1825).— <i>Hydnium</i> .									
1252	...	...	...	...	...	Q.	B.	Ground and trunks	... Gelatinous, tremulous, semicircular, somewhat stalked, greyish green to brownish. Only tremelloid fungus with true spines.
Syst. 89 (1816).— <i>Tremella</i> .									
1253	...	...	T.	V.	...	...	B.	Rotten wood	... Nearly round, rooting, yellowish, contorted at length, hyaline.
1254	...	S.A.	T.	...	N.S.W.	...	...	Dry wood	... Small, vermilion, lobate and folded.
1255	W.A.	...	...	...	...	...	...	Rotten branches	... Small, reddish brown, black when dry, cracked and folded.
1256	...	...	...	...	...	Q.	...	Stems of <i>Saccharum</i>	... Irregular, thin, gelatinous, orange red, seated on whitish layer, spreading over charred stems of sugar cane.
1257	...	...	T.	...	...	...	...	Bark	... White, circular, depressed in centre, cup shaped.
1258	...	...	T.	...	...	...	...	Bark	... Bursting through, arranged in a row, whitish, then yellowish.
1259	...	S.A.	...	V.	N.S.W.	...	B.	Rotten wood	... Nearly round, folded, yellow to orange, colour persistent.
Pl. Homon. 92 (1825).— <i>Dacryomyces</i> , <i>Merulius</i> , <i>Tremella</i> .									
1260	...	S.A.	...	...	...	...	...	Rotten wood of <i>Melaleuca</i>	... Jelly-like to tough, orange yellow, solitary or in tufts. Stem at first clubbed, then expanding into cup shape.
1261	W.A.	...	T.	V.	...	Q.	...	Wood	... Minute, velvety, red. Stem short, velvety. Hymenium obliquely cup shaped.
1262	...	...	...	V.	N.S.W.	Q.	...	Wood and fences	... Tufted, somewhat erect, rooting. Cap semicircular, spoon shaped. Stem downy, glaucous.
Syst. Myc. I. 485 (1821).— <i>Clavaria</i> .									
1263	...	...	...	V.	...	Q.	B.	Rotten wood	... Tufted, rooting, viscid, orange yellow. Clubs short, grown together at base.
1264	...	...	...	V.	...	...	...	Damp logs	... Branched, tough, pale. Trunk thin, twice or thrice forked. Branches expanded at apex like a spoon each bearing from three to five finger-like processes.
1265	...	...	T.	V.	...	...	B.	Trunks	... Simple, solitary, jelly-like, yellow. Clubs thickened, compressed. Stem tapering.
1266	W.A.	S.A.	T.	V.	...	Q.	...	Rotten wood	... Small, bursting through, reddish brown. Stem compressed, palmate above. Branches few.
1267	...	S.A.	...	V.	...	...	...	Trunks	... Scattered, tapering, compressed, honey yellow, curved. Stem very short, but distinct.
1268	...	...	...	V.	...	...	B.	Wood and dead fir leaves	... Simple, solitary, elongated, linear, yellow. Clubs short, grown together at base.

GENERAL CLASSIFICATION OF GASTROMYCETES.

GROUP II.—GASTROMYCETES, WILLD.

ARRANGEMENT OF ORDERS (6).

Above ground—

7. PILACREACEÆ—Minute. Peridium eventually disintegrating. Intermediate between Hymenomyces and Gastromycetes.
8. PHALLOIDEACEÆ—Fleshy to gelatinous. Receptacle and spore-bearing surface enclosed in universal volva
9. NIDULARIACEÆ—Leathery. Spores never powdery.
10. LYCOPERDACEÆ—Membranous to leathery. Spores forming powdery mass when mature.
11. SCLERODERMACEÆ—Leathery. Peridium thick, sessile or stalked, opening at apex.

Subterranean—

12. HYMENOGASTRACEÆ—Fleshy to firm. Peridium indehiscant.

ORDER VII.—PILACREACEÆ, BREF.

Genus (1)—

106. Pilacre, Fries.

ORDER VIII.—PHALLOIDEACEÆ, FRIES.

Genera (8)—

107. Dictyophora, Desv.	109. Mutinus, Fries.	111. Colus, Cav. and Sec.	113. Anthurus, Kalch.
108. Ithyphallus, Fries.	110. Clathrus, Linn.	112. Lysurus, Fries.	114. Aseroë, La Bill.

ORDER IX.—NIDULARIACEÆ, FRIES.

Genera (3)—

115. Cyathus, Hall.	116. Crucibulum, Tul.	117. Sphærobelus, Tode.

ORDER X.—LYCOPERDACEÆ, EHR.

Genera (18)—

118. Secotium, Kunze.	123. Gymnoglossum, Mass.	127. Calostoma, Desv.	132. Areolaria, Forq.
119. Chainoderma, Mass.	124. Protoglossum, Mass.	128. Geaster, Scop.	133. Castoreum, Cooke and Mass.
120. Cycloderma, Klot.	125. Tulostoma, Pers.	129. Diploderma, Link.	134. Xylopodium, Mont.
121. Mesophellia, Berk.	126. Battarreia, Pers.	130. Bovista, Scop.	135. Favillea, Fries.
122. Podaxou, Fries.		131. Lycoperdon, Linn.	

ORDER XI.—SCLERODERMACEÆ, FRIES.

Genera (4)—

136. Scleroderma, Pers.	137. Polysaceum, D.C.	138. Arachnion, Schw.	139. Paurocotylis, Berk.

ORDER XII.—HYMENOGASTRACEÆ, VITT.

Genera (5)—

140. Octaviania, Vitt.	142. Hymenogaster, Vitt.	143. Hydngangium, Wallr.	144. Gautieria, Vitt.
141. Rhizopogon, Fries.			

GROUP II.—GASTROMYCETES.—

ORDER VII.—PILACREACEÆ,

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
106. PILACRE.—Fries,					
1269	1989	IV. 2750	<i>P. divisa</i>	Berk., Fl. N.Z. II. 197 (1855)	Divided pilacre
1270	1990	„ 2752	<i>P. Petersii</i>	Berk. and Curt., Ann. Nat. Hist. III., 3rd Ser. 362 (1859)	Peters' pilacre

ORDER VIII.—PHALLOIDEACEÆ,

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
107. DICTYOPHORA.—Desv., Journ.					
1271	1179	VII. 13	<i>D. merulina</i>	Berk., Linn. Journ. XIII. 172 (1873)	Merulius-like dictyophora
1272	1178	„ 11	<i>D. multicolor</i>	Berk. and Br., Linn. Trans. II. 65 (1883)	Many-coloured dictyophora
1273	1175	„ 2	<i>D. phalloidea</i>	Desv., Journ. Bot. II. 88 (1809)	Phallus-like dictyophora
1274	1177	„ 6	<i>D. speciosa</i>	Meyen., Nov. Act. XIX. 239 (1843)	Handsome dictyophora
1275	1176	„ 3	<i>D. tahitensis</i>	Fisch., Sacc. Syll. VII. 4 (1888)	Tahitian dictyophora

108. ITHYPHALLUS.—Fries, S.M. II. 283 (1823).—

1276	1183	VII. 22	<i>I. aurantiacus</i>	Fisch., Sacc. Syll. VII. 9 (1888)	Orange-coloured ithyphallus
1277	1182	„ 21	<i>I. calyptratus</i>	Fisch., Sacc. Syll. VII. 9 (1888)	Capped ithyphallus
1278	1180	„ 18	<i>I. impudicus</i>	Fisch., Sacc. Syll. VII. 8 (1888)	Impure ithyphallus (Stinkhorn)
1279	1184	„ 23	<i>I. novæ hollandiæ</i>	Fisch., Sacc. Syll. VII. 10 (1888)	New Holland ithyphallus
1280	1181	„ 20	<i>I. quadricolor</i>	Fisch., Sacc. Syll. VII. 9 (1888)	Four-coloured ithyphallus
1281	1185	„ 27	<i>I. retusus</i>	Fisch., Sacc. Syll. VII. 11 (1888)	Blunt ithyphallus
1282	1186	„ 29	<i>I. rubicundus</i>	Fisch., Sacc. Syll. VII. 11 (1888)	Rubicund ithyphallus

WILLD., BEMERK. FARR. (1802).

BREF. UNT. VII. (1888).

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
1269	...	...	T.	...	...	...	...	Bark ...	Head globose, clay coloured. Stem divided, brownish.
1270	...	...	...	...	N.S.W.	...	...	Trunks ...	Gregarious, often covering half-dead trunks for a considerable distance. Head relatively large. Stem short, whitish.

Pl. Homon. 364 (1825).

FRIES, S.M. II. 281 (1823).

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
1271	...	...	...	...	...	Q.	...	Ground ...	Gregarious. Cap bell shaped, ochrey, covered with fœtid brown spore-bearing mass. Stem distinct, white.
1272	...	...	...	...	N.S.W.	Q.	...	Ground ...	About 7 inches high. Cap conical, orange, netted. Stem cream coloured, tapering to base.
1273	...	...	...	...	...	Q.	...	Sandy soil	Cap thickened at apex, bell shaped, white, netted. Stem white and pitted.
1274	...	...	...	...	...	Q.	...	Ground ...	Stem tapering upwards, white. Cap joined to stem at apex with short collar, bell shaped.
1275	...	...	...	...	N.S.W.	Q.	...	Ground ...	Stem cylindrical, walls pitted. Cap at apex of stem without collar ovate, roughly netted.

Bot. II. 92 (1809).—Phallus.

1276	...	...	...	...	...	Q.	...	Ground ...	Stem 6 to 8 inches long, $\frac{3}{4}$ inch thick, orange. Cap without collar or ring, thimble shaped, orange.
1277	...	...	...	...	...	Q.	...	Among grass	Scarcely 2 inches high. Stem slightly tapering upwards. Cap somewhat hemispherical, orange.
1278	...	...	...	...	...	Q.	B.	Ground ...	Up to 10 inches high. Stem tapering above and below, white. Cap conical, netted; gelatinous mass of spores, dark olive.
1279	...	...	...	V.	N.S.W.	Q.	...	Ground ...	Stem white, slender, tapering upwards. Cap narrow bell shaped, netted.
1280	...	...	...	...	...	Q.	...	Ground ...	About $4\frac{1}{2}$ inches high. Stem cylindrical, lemon colour, veil white; mycelium purple. Cap orange coloured.
1281	...	...	...	...	N.S.W.	...	...	Ground ...	About 6 inches high. Stem 1 inch or more thick, cylindrical. Cap ovate, blunt above apex of stem, white.
1282	...	...	...	V.	...	...	...	Ground ...	Stem between 5 and 6 inches high, spindle shaped, red. Cap conical to bell shaped, clad with brownish gluten.

Phallus, Omphalo-phallus, Cynophallus.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
109. MUTINUS.—Fries, Summ. Veg.					
1283	...	VII. 30	<i>M. caninus</i> ...	<i>Fries</i> , S.V.S. II. (1849) ...	Dog mutinus ...
1284	1188	„ 38	<i>M. curtus</i> ...	<i>Fisch.</i> , Sacc. Syll. VII. 13 (1888) ...	Short mutinus ...
1285	1190	„ 40	<i>M. discolor</i> ...	<i>Fisch.</i> , Sacc. Syll. VII. 14 (1888) ...	Discoloured mutinus ...
1286	1189	„ 39	<i>M. papuasius</i> ...	Kalch., Grev. IV. 74 (1875) ...	Papuan mutinus ...
1287	1187	„ 37	<i>M. Watsoni</i> ...	<i>Fisch.</i> , Sacc. Syll. VII. 13 (1888) ...	Watson's mutinus ...
110. CLATHRUS.— <i>Linn.</i> ,					
1288	1194	VII. 61	<i>C. albidus</i> ...	Becker, Schles. Gesell. 81 (1874) ...	Whitish clathrus ...
1289	1195	„ 59	<i>C. cibarius</i> ...	<i>Fisch.</i> , Sacc. Syll. VII. 20 (1888) ...	Edible clathrus ...
1290	1196	„ 60	<i>C. crispus</i> ...	<i>Turp.</i> , Dict. Sci. Nat. (1822) ...	Curled clathrus ...
1291	1193	„ 58	<i>C. gracilis</i> ...	<i>Schl.</i> , Linn. 166 (1861)... ...	Graceful clathrus ...
1292	1192	„ 55	<i>C. pusillus</i> ...	Berk., Hook., Lond. Journ. IV. 67 (1845) ...	Small clathrus ...
1293	1191	„ 51	<i>C. triscapus</i> ...	<i>Fries</i> , S.M. II. 287 (1823) ...	Three-branched clathrus ...
111. COLUS.—Cav. and Sec.,					
1294	1197	VII. 62	<i>C. hirudinosus</i> ...	Cav. and Sec., Ann. Sci. Nat. 253 (1835) ...	Leech colus ...
1295	...	IX. 1093	<i>C. Muelleri</i> ...	<i>Fisch.</i> , Unters. Phall. 61 (1890) ...	Mueller's colus ...
112. LYSURUS.—Fries,					
1296	1198	IX. 1095	<i>L. australiensis</i> ...	Cooke and Mass., Grev. XVIII. 6 (1889) ...	Australian lysurus ...
113. ANTHURUS.—Kalch.,					
1297	1200	VII. 71	<i>A. Archeri</i> ...	<i>Fisch.</i> , Sacc. Syll. VII. 24 (1888) ...	Archer's anthurus ...
1298	1199	„ 69	<i>A. Muellerianus</i> ...	Kalch., Grev. IX. 2 (1880) ...	Mueller's anthurus ...
114. ASEROË.—La Bill.,					
1299	1202	VII. 78	<i>A. lysuroides</i> ...	<i>Fisch.</i> , Jahrb. Bot. Gart. IV. (1886) ...	Lysurus-like aseroë ...
1300	1201	„ 76	<i>A. rubra</i> ...	La Bill. Voy. 44 (1798) ...	Red aseroë ...
1300A	„	„ „	<i>A. rubra</i> , var. <i>pentactina</i> ...	(<i>Sacc.</i>) Syll. VII. 26 (1888) ...	Five-rayed aseroë ..

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Scan. II. 434 (1849).—Phallus, Cynophallus.									
1283	...	...	...	...	...	...	B.	Woods ...	Receptacle somewhat spindle shaped, white or rosy, spore-bearing portion short, red. Stem 3 to 4 inches high, scentless.
1284	W.A.	...	...	...	...	...	...	Ground ...	About 1 inch; receptacle broadly truncate at apex. Stem yellow; very foetid.
1285	...	...	...	...	...	Q.	...	Ground ...	Stem cylindrical, orange; spore-bearing part one-sixth of receptacle; apex yellowish grey or turning black.
1286	...	...	...	...	...	Q.	...	Ground ...	About 3 to 4 inches high; receptacle thin and slender; spore-bearing part pear shaped, black.
1287	...	...	...	...	...	Q.	...	Ground ...	About 2½ inches high; spore-bearing part conical, minutely veined, red.
Sp. Pl. II. 1179 (1753).—Neodictyon.									
1288	...	...	...	V.	...	...	...	Ground ...	Branches of receptacle with broad channel, white, then yellowish.
1289	...	...	...	V.	...	Q.	...	Ground ...	Receptacle spherical or ovoid, white, interstices broad; gregarious and common. <i>Edible</i> .
1290	...	...	...	...	N.S.W.	Q.	...	Sandy soil	Receptacle spherical or obovate, vermilion or salmon colour; interstices rounded or oval.
1291	W.A.	S.A.	T.	V.	N.S.W.	Q.	...	Ground ...	Veil globose, splitting into about four lobes; receptacle ovoid, white; interstices hexagonal.
1292	W.A.	...	...	...	...	Q.	...	Ground ...	Small; veil nearly cylindrical; receptacle bright ruby red.
1293	...	...	...	...	...	Q.	...	Ground ...	Receptacle of three vertical branches, slender, thin, straight, white below, vermilion about apex.
Ann. Sci. Nat. 2 Ser. III. 253 (1835).—Clathrus.									
1294	W.A.	...	...	...	...	...	...	Ground ...	Receptacle spindle shaped, white, reddening at apex. Stem inversely conical, with meshes at top.
1295	...	...	...	V.	...	...	...	Ground ...	Receptacle unequally perforated, the superior meshes of equal diameter, inferior greatly elongated. Stem short.
S.M. II. 285 (1823).—Mutinus.									
1296	...	...	...	...	...	Q.	...	Ground ...	Receptacle tawny, mostly five lobed. Stem cylindrical, hollow, whitish. Veil globose, torn in lobes.
Grev. IX. 2 (1880).—Lysurus.									
1297	...	...	T.	...	...	...	...	Ground ...	Receptacle 3½ inches high, rosy. Stem very short, divided into five long erect lobes.
1298	...	...	...	V.	N.S.W.	...	...	Ground ...	Receptacle yellow to reddish, cup shaped or funnel shaped above, dilated.
Voy. 44 (1798).—Lysurus.									
1299	...	...	T.	...	...	...	...	Ground ...	Stem white, slender, long, disc carmine rose above, externally distinct from stem.
1300	...	S.A.	...	V.	N.S.W.	Q.	B. (introduced)	Ground ...	Stem becoming red, margin divided into five to eight teeth, vermilion above, forked.
1300A	...	...	...	...	N.S.W.	...	...	Ground ...	With five bifid rays.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER IX.—NIDULARIACEÆ,					
115. CYATHUS.—Hall,					
1301	...	VII. 101	<i>C. ambiguus</i> ...	Tul., Ann. Sci. Nat. 75 (1844) ...	Ambiguous cyathus ...
1302	...	...	<i>C. Baileyi</i> ...	Mass., Grev. XXI. 3 (1892) ...	Bailey's cyathus ...
1303	1214	VII. 127	<i>C. Colensoi</i> ...	Berk., Fl. N. Zeal. II. 192 (1855) ...	Colenso's cyathus ...
1304	1212	" 113	<i>C. dasypus</i> ...	Nees, Phys. Ber. 41 (1820) ...	Hairy-stalked cyathus ...
1305	1206	" 104	<i>C. desertorum</i> ...	F. v. M., Linn. Journ. XVIII. 387 (1881) ...	Desert cyathus ...
1306	...	...	<i>C. dimorphus</i> ...	Cobb., Ag. Gaz. N.S.W. III. Pt. 12 (1892) ...	Dimorphic cyathus ...
1307	1213	VII. 121	<i>C. fimetarius</i> ...	D. C., Fl. Fr. V. 104 (1815) ...	Dung cyathus ..
1308	1209	" 116	<i>C. fimicola</i> ...	Berk., Linn. Journ. XVIII. 387 (1881) ...	Dung-borne cyathus ...
1309	1205	" 98	<i>C. intermedius</i> ...	Tul., Mon. Nid. Ann. Sci. Nat. 72 (1844)...	Intermediate cyathus ...
1310	1207	" 109	<i>C. Lesueurii</i> ...	Tul., Mon. Nid. Ann. Sci. Nat. 79 (1844)...	Lesueur's cyathus ...
1311	1204	" 96	<i>C. Montagnei</i> ...	Tul., Mon. Nid. Ann. Sci. Nat. 70 (1844)...	Montagne's cyathus ...
1312	1203	" 95	<i>C. novæ-zealandiæ</i> ...	Tul., Mon. Nid. Ann. Sci. Nat. 66 (1844)...	New Zealand cyathus ...
1313	1210	" 117	<i>C. pezizoides</i> ...	Berk., Linn. Journ. XVIII. 387 (1881) ...	Peziza-like cyathus ...
1314	1211	" 118	<i>C. pusio</i> ...	Berk., Linn. Journ. XVIII. 387 (1881) ...	Small cyathus ...
1315	1208	" 110	<i>C. varicosus</i> ...	D. C., Flor. Fr. II. 270 (1805) ...	Varnished cyathus ...
116. CRUCIBULUM.—Tul., Mon. Nid. Ann. Sci.					
1316	1216	...	<i>C. simile</i> ...	Mass., Grev. XIX. 94 (1891) ...	Similar crucibulum ...
1317	1215	VII. 128	<i>C. vulgare</i> ...	Tul., Mon. Nid. Ann. Sci. Nat. 90 (1844)...	Common crucibulum ...
117. SPHEROBOLUS.—Tode,					
1318	1217	VII. 136	<i>S. stellatus</i> ...	Tode, Meck. 43 (1790) ...	Stellate sphaerobolus ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
1301	...	...	...	...	...	Q.	...	Garden soil	... Oboconical to cup shaped, cartilaginous to membranous, tawny, rusty, or tawny cinnamon, beautiful silky gloss.
1302	...	...	...	...	...	Q.	...	Dung	... More or less gregarious, inversely conical or bell shaped, thin and cartilaginous, minutely downy, cinnamon.
1303	...	S.A.	T.	...	N.S.W.	...	...	Ground	... Densely crowded, cup shaped, thin, flexible, dirty umber, downy.
1304	...	...	...	V.	...	...	...	Bare ground	... Bell shaped, somewhat cylindrical, pale ochre, and minutely downy.
1305	...	S.A.	...	V.	N.S.W.	...	...	Sandy soil	... Pale, downy, smooth within and even.
1306	...	...	...	...	N.S.W.	...	...	Ground	... Gregarious. Peridium ash colour, bell shaped, with obscure circle of small markings half way up.
1307	...	...	...	...	...	Q.	...	Dung	... Hemispherical, brown to tawny, velvety.
1308	...	...	...	V.	...	Q.	...	Dung	... Cup shaped, umber, becoming pale, minutely velvety.
1309	...	...	...	...	...	Q.	...	Rubbish	... Cup shaped, inversely conical, rusty, becoming yellowish, hairy, slightly streaked.
1310	...	...	...	V.	N.S.W.	Q.	...	Rotten wood	... Membranous, thin, grey, clad with somewhat star-shaped hairs or naked.
1311	W.A.	...	...	...	...	Q.	...	Chips	... Crucible shaped, rusty, with a few woolly hairs, internally smooth, streaked and ciliate above.
1312	...	...	...	...	N.S.W.	...	...	Rotten wood	... Elongated, narrow, brown, woolly outside, streaked and furrowed inside.
1313	...	...	...	...	...	Q.	...	Rotting herbs	... Cup shaped, densely downy outside with flexuous hairs, umber, very small, smooth inside.
1314	...	...	...	...	...	Q.	...	Trunks of <i>Eucalyptus</i>	... Wine-glass shaped, whitish, clad outside with a fine down, smooth inside.
1315	W.A.	...	...	V.	...	Q.	B.	Ground and twigs	... Clustered, bell shaped, nearly sessile, pale ochre to ash colour, then dusky, downy, lead colour or brown within.

Nat. 3 Ser. I. 89 (1844).—Cyathus, Nidularia, Peziza.

1316	...	...	...	...	...	...	...	Bark and wood	... Crowded or scattered, somewhat cylindrical, thin, flexible, externally densely covered with ochrey-brown shaggy down.
1317	...	...	...	V.	...	Q.	B.	Twigs, wood, &c.	... Gregarious, cylindrical to bell shaped, ochrey, then rusty, downy when young, then smooth, inside shining yellow.

Meck. I. 43 (1790).—Lycoperdon.

1318	...	...	T.	V.	...	...	B.	Wood, chips, &c.	... Nearly spherical, fleshy, dehiscing with five to eight acute teeth, yellow, interior whitish.
------	-----	-----	----	----	-----	-----	----	------------------	---

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER X.—LYCOPERDACEÆ,					
118. SECOTIUM.—Kunze					
1319	1218	VII. 146	<i>S. acuminatum</i> ...	Mont., Fl. Alg. I. 371 (1846) ...	Acuminate secotium ...
1320	1219	„ 150	<i>S. coarctatum</i> ...	Berk., Hook., Lond. Journ. IV. 63 (1845) ...	Compact secotium ...
1321	1221	„ 152	<i>S. erythrocephalum</i> ...	Tul., Ann. Sci. Nat. 115 (1844) ...	Red-headed secotium ...
1322	1222	„ 156	<i>S. Gunnii</i>	Berk., in Cooke's Handb. Austr. Fung. 221 (1892) ...	Gunn's secotium ...
1323	1220	„ 151	<i>S. melanosporum</i> ...	Berk., Hook., Lond. Journ. IV. 62 (1845) ...	Dark-spored secotium ..
1324	1223	...	<i>S. scabrosum</i> ...	Cooke and Mass., Grev. XX. 35 (1891) ...	Scabrous secotium ...
119. CHAINODERMA.—Massee,					
1325	1224	...	<i>C. Drummondii</i> ...	Mass., Grev. XIX. 46 (1890) ...	Drummond's chainoderma ...
120. CYCLODERMA.—Klotzsch,					
1326	1225	VII. 1584	<i>C. platyspora</i> ...	Cooke and Mass., Grev. XVI. 73 (1888) ...	Broad-spored cycloiderma ...
121. MESOPHELLIA.—Berk.,					
1327	1226	VII. 162	<i>M. arenaria</i> ...	Berk., Linn. Trans. XXII. 131 (1857) ...	Sandy mesophellia ...
1328	1227	„ 163	<i>M. ingratisima</i> ...	De Toni, Sacc. Syll. VII. 57 (1888) ...	Strong-smelling mesophellia ...
122. PODAXON.—Fries,					
1329	1231	VII. 170	<i>P. axata</i> ...	Mass., Mon. Pol. Journ. Bot. (1890) ...	Axate podaxon ...
1330	1229	„ 168	<i>P. carcinomalis</i> ...	Fries, S.M. III. 62 (1829) ...	Cancerous podaxon ...
1331	1230	„ 171	<i>P. indica</i> ...	Spreng., Syst. Veg. V. 518 (1828) ...	Indian podaxon ...
123. GYMNOGLOSSUM.—Massee,					
1332	1232	...	<i>G. stipitatum</i> ..	Mass., Grev. XIX. 97 (1891) ...	Stalked gymnoglossum ...
124. PROTOGLOSSUM.—Mass.,					
1333	1233	...	<i>P. luteum</i> ...	Mass., Grev. XIX. 97 (1891) ...	Yellow protoglossum ...
125. TULOSTOMA.—Pers.,					
1334	1240	...	<i>T. album</i> ...	Mass., Grev. XIX. 95 (1891) ...	White tulostoma ...
1335	1238	VII. 185	<i>T. fimbriatum</i> ...	Fries, S.M. III. 43 (1829) ...	Fringed tulostoma ...
1336	1239	„ 193	<i>T. granulosum</i> ...	Lev., Demid. Voy. IV. 120 (1842) ...	Granular tulostoma ...
1337	1235	„ 177	<i>T. leprosum</i> ...	Kalch., Grev. IV. 72 (1875) ...	Leprous tulostoma ...
1338	1234	„ 175	<i>T. mammosum</i> ...	Fries, S.M. III. 42 (1829) ...	Teat-like tulostoma ...
1339	1237	„ 184	<i>T. maximum</i> ...	Cooke and Mass., Grev. XV. 94 (1887) ...	Maximum tulostoma ...
1340	1241	IX. 1113	<i>T. pulchellum</i> ...	Sacc., Bull. Soc. Myc. V. 118 (1889) ...	Beautiful tulostoma ...
1341	1236	VII. 182	<i>T. Wightii</i> ...	Berk., Hook., Lond. Journ. I. 157 (1842) ...	Wight's tulostoma ..

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
EHR. SYLV. BER. 14 (1818).									
Fl. 321 (1840).									
1319	W.A.	...	...	...	...	...	...	Ground ...	Solitary, ovoid, tan or ochrey, acuminate at apex, cuticle breaking up into scales. Stem short, twisted.
1320	W.A.	S.A.	...	...	...	...	...	Ground ...	Minute, strong scented, inversely ovate; margin bent inwards. Stem slender, cylindrical.
1321	...	...	T.	...	...	...	...	Ground ...	Gregarious, rather long stemmed. Stem erect, naked, white. Peridium smooth, carmine red.
1322	...	...	T.	...	...	...	...	Ground ...	Small. Stem slender, solid, pale brown. Peridium somewhat globose, pale brown, smooth.
1323	W.A.	...	...	...	...	...	...	Ground ...	Tufted. Cap somewhat globose, mealy. Margin obtuse and rounded. Stem solid. Spores dark chocolate brown.
1324	...	...	...	V.	...	...	...	Ground ...	Hemispherical, dingy olive or greyish, rugged.
Grev. XIX. 46 (1890).—Secotium.									
1325	W.A.	...	...	...	...	...	...	Ground ...	Club to spindle shaped, dingy brown, smooth. Mass of spores dingy brown.
Linn. VII. 203 (1832).									
1326	...	...	...	V.	...	...	...	Ground ...	Ovate, external thick, flexible, ochrey, internal thin, shining.
Fl. Tasm. II. 266 (1860).—Inoderma.									
1327	...	...	T.	V.	...	...	...	Ground ...	Thick, elliptical, externally clad with whitish woolly threads.
1328	...	...	...	V.	...	...	...	Ground ...	Strong scented, crustaceous, very fragile, somewhat globose.
S.M. III. 62 (1829).—Lycoperdon, Mitremyces.									
1329	...	...	...	...	N.S.W.	...	...	Dry sandy places ...	Tuberous rooting, oblong. Stem hollow, substance woody. Peridium ovate.
1330	...	S.A.	...	...	...	Q.	...	Sandy places ...	Peridium ovate, oblong, whitish. Stem cylindrical, curved.
1331	...	...	...	V.	...	Q.	...	Ground ...	Stem corded, often twisted lengthwise. Peridium club shaped, invested with saffron-yellow membrane.
Grev. XIX. 97 (1891).									
1332	...	...	...	...	N.S.W.	...	...	Ground ...	Obtusely conical, pitted, pale brown. Stem solid, pale brown.
Grev. XIX. 97 (1891).									
1333	...	...	...	V.	...	Q.	...	In rich black mould	Cylindrical, growing vertically, with extreme apex above, orange yellow, underground portion yellowish. No stem.
Tent. Disp. 6 (1797).—Lycoperdon, Schizostoma.									
1334	W.A.	...	...	...	...	...	...	Ground ...	Stem ochrey, wrinkled lengthwise. Peridium globose, pure white shining.
1335	W.A.	...	...	V.	...	...	...	Sandy soil ...	Peridium almost naked, scales falling away, becoming tawny. Stem tawny ochre, mouth torn, fringed.
1336	W.A.	...	...	V.	...	...	...	Ground ...	Peridium globose, depressed, brown; mouth teat-like; margin torn and toothed. Stem thickish.
1337	...	...	...	...	...	Q.	...	Ground ...	Peridium clad with a lurid umber mealy scurf, at length falling away.
1338	...	...	...	V.	...	Q.	B.	Ground ...	Stem hollow, covered more or less with falling scales. Peridium globose, with minute prominent teat-like mouth.
1339	W.A.	...	...	...	...	...	...	Ground ...	Peridium smooth, ochrey, with rounded mouth. Stem elongated, of same colour.
1340	...	S.A.	...	V.	...	...	...	Branches ...	Minute, shortly stalked. Stem cylindrical, smooth, whitish. Peridium membranous, sub-globose.
1341	...	...	...	...	...	Q.	...	Ground ...	Peridium papery, egg shaped to globular. Stem somewhat scaly.

SYSTEMATIC ARRANGEMENT

Number	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
126. BATTARREA.—Pers.,					
1342	1244	VII. 199	<i>B. Muelleri</i> ...	Kalch., Grev. IX. 3 (1880) ...	Mueller's battarrea ...
1343	1242	„ 195	<i>B. pballoides</i> ...	<i>Pers.</i> , Syn. 129 (1801) ...	Phallus-like battarrea ...
1344	1243	„ 196	<i>B. Steveni</i> ...	<i>Fries</i> , S.M. III. 7 (1829) ...	Steven's battarrea ...
1345	1245	IX. 1115	<i>B. Tepperiana</i> ...	Ludw., Bot. Centr. 337 (1889) ...	Tepper's battarrea ...
127. CALOSTOMA.—Desv.,					
1346	1249	...	<i>C. æruginosa</i> ...	Mass., Grev. XIX. 96 (1891) ...	Verdigris-green calostoma ...
1347	1247	VII. 206	<i>C. fusca</i> ...	<i>Mass.</i> , Ann. Bot. II. 43 (1888) ...	Brown calostoma ...
1348	1246	„ 205	<i>C. lurida</i> ...	<i>Mass.</i> , Ann. Bot. II. 43 (1888) ...	Lurid calostoma ...
1349	1248	„ 207	<i>C. viridis</i> ...	<i>Mass.</i> , Ann. Bot. II. 40 (1888) ...	Green calostoma ...
128. GEASTER.—Scop.,					
1350	1270	VII. 1590	<i>G. Archeri</i> ...	Berk., Fl. Tasm. II. 264 (1860) ...	Archer's geaster ...
1351	1271	IX. 1123	<i>G. argenteus</i> ...	Cooke, Grev. XVII. 75 (1889) ...	Silvery geaster ...
1352	1259	VII. 1592	<i>G. australis</i> ...	Berk., Fl. Tasm. II. 265 (1860) ...	Southern geaster ...
1353	...	IX. 1119	<i>G. Berkeleyi</i> ...	Mass., Ann. Bot. 79 (1889) ...	Berkeley's geaster ...
1354	1253	VII. 226	<i>G. Drummondii</i> ...	Berk., Hook., Lond. Journ. IV. 63 (1845) ...	Drummond's geaster ...
1355	1269	„ 261	<i>G. dubius</i> ...	Berk., Linn. Journ. XVI. 40 (1878) ...	Doubtful geaster ...
1356	1264	„ 248	<i>G. floriformis</i> ...	Vitt., Mon. Lycop. 167 (1842) ...	Flower-shaped geaster ...
1357	1258	„ 238	<i>G. fimbriatus</i> ...	Fries, S.M. III. 16 (1829) ...	Fringed geaster ...
1358	1268	„ 257	<i>G. hygrometricus</i> ...	<i>Pers.</i> , Syn. 135 (1801) ...	Hygrometric geaster ...
1359	1262	„ 246	<i>G. lageniformis</i> ...	Vitt., Mon. Lycop. 160 (1842) ...	Flask-shaped geaster ...
1360	1256	„ 229	<i>G. lignicola</i> ...	Berk., Linn. Journ. XVIII. 386 (1881) ...	Wood-growing geaster ...
1361	1267	„ 255	<i>G. lugubris</i> ...	Kalch., Gast. 10 (1883) ...	Mourning geaster ...
1362	1257	„ 232	<i>G. minimus</i> ...	Schwein., Syn. Car. 327 (1822) ...	Least geaster ...
1363	1265	„ 249	<i>G. pusillus</i> ...	Fries, Pl. Preiss. II. 139 (1846) ...	Small geaster...
1364	1254	„ 1591	<i>G. Readeri</i> ...	Cooke and Mass., Grev. XVI. 73 (1888) ...	Reader's geaster ...
1365	1266	„ 251	<i>G. rufescens</i> ...	<i>Pers.</i> , Syn. 134 (1801) ...	Reddish geaster ...
1366	1261	„ 245	<i>G. saccatus</i> ...	Fries, S.M. III. 16 (1829) ...	Saccate geaster ...
1367	1263	„ 247	<i>G. Spegazzinianus</i> ...	<i>De Ton.</i> , Rev. Geast. in Rev. Myc. 19 (1887) ...	Spegazzini's geaster ...
1368	1252	„ 224	<i>G. striatulus</i> ...	Kalch., Grev. IX. 3 (1880) ...	Furrowed geaster ...
1369	1251	„ 222	<i>G. striatus</i> ...	D. C., Fl. Fr. II. 267 (1805) ...	Streaked geaster ...
1370	1255	„ 228	<i>G. subiculosus</i> ...	Cooke and Mass., Grev. XV. 97 (1887) ...	Subiculous geaster ...
1371	1250	„ 218	<i>G. tenuipes</i> ...	Berk., Hook., Lond. Journ. VII. 576 (1848) ...	Slender-stalked geaster ...
1372	1260	„ 242	<i>G. vittatus</i> ...	Kalch., Grev. IX. 3 (1880) ...	Vittate geaster ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.	
	W.A.	S.A.	T.	V.	N.S.W.	Q.				
Syn. 129 (1801).—Lycoperdon, Dendromyces.										
1342	...	S.A.	...	...	...	...	...	Ground	...	Entirely white, at length rusty from the scattered spores. Peridium bell to mitre shaped, seated on solid very long stem.
1343	W.A.	...	...	V.	...	...	B.	Sandy soil	...	Veil ovate, whitish, with mucus. Stem cylindrical, tapering a little to each end. Peridium bell shaped, smooth below, powdery brown above.
1344	W.A.	...	...	...	...	...	...	Sandy soil	...	Stem bellied, covered with scales, hollow. Peridium somewhat plane leathery, thin.
1345	...	S.A.	...	V.	...	...	...	Sandy soil	...	Stem very long, thickened upwards, woody, hollow, in upper part torn into large membranous scales.
Journ. Bot. II. 94 (1809).—Mitremyces.										
1346	...	...	...	V.	...	...	...	Ground	...	Outer peridium becoming broken up into small verdigris-green scales; inner peridium sub-globose, dingy green.
1347	W.A.	...	T.	V.	...	...	...	Ground	...	Simple or tufted, outer peridium dark brown, dingy red within; inner peridium pale brown, sub-globose, mouth vermilion, teeth erect.
1348	W.A.	...	...	...	...	...	...	Sandy soil	...	Outer peridium breaking up early into small blackish granules, adhering to ochrey inner peridium.
1349	...	...	...	V.	...	Q.	...	Ground and timber	dead	Outer peridium in form of dingy-green irregular scales, adhering to pale-green inner peridium, stem-like, base stout, greenish.
Carn. II. 489 (1772).—Lycoperdon.										
1350	...	...	T.	...	...	Q.	...	Ground	...	Outer peridium cut to the middle into six to seven lobes; inner peridium globose, purplish umber.
1351	...	...	...	V.	...	...	...	Ground	...	Outer peridium cut into eight to ten teeth, whitish and shining, internally dingy umber; inner peridium globose.
1352	W.A.	...	T.	...	...	...	...	Ground	...	Outer peridium leathery, rigid, cut to the middle in eight to ten lobes; inner peridium sub-globose, pale umber.
1353	...	...	...	...	...	...	B.	Ground	...	Outer peridium thinnish, split to centre into a number of segments; inner peridium short stalked, thick, pale brown.
1354	W.A.	S.A.	...	V.	...	...	...	Ground	...	Outer peridium simple, rigid, flattened, many lobed; inner peridium globose, delicately rough.
1355	...	...	...	...	N.S.W.	Q.	...	Ground	...	Outer peridium thick, globose, delicately powdery, fawn colour, seated on stem-like mycelium.
1356	...	S.A.	...	V.	...	Q.	...	Ground	...	Outer peridium cut into five to eight lobes; inner peridium ovate-oblong, papery, greyish white, shining.
1357	...	S.A.	T.	V.	...	Q.	B.	Grassy spots	...	Outer peridium simple, five to fifteen lobed, flattened, tawny brown; inner peridium sub-globose, whitish, yellow or umber.
1358	W.A.	...	...	...	...	Q.	B.	Ground	...	Outer peridium cut to the base into seven to twenty lobes, rarely six; inner peridium compressed, brown or grey.
1359	...	...	...	...	...	Q.	B.	Ground	...	Outer peridium cut nearly to middle in six to nine lobes; inner peridium nearly spherical, soft, membranous.
1360	...	...	...	...	...	Q.	...	Trunks	...	Outer peridium downy to granular, pale, irregularly ruptured; inner peridium brown.
1361	W.A.	...	...	...	...	...	...	Ground	...	Outer peridium cut into seven to eight narrow lance-shaped teeth, with thin continuous black layer; inner peridium clay colour to brownish.
1362	W.A.	S.A.	...	V.	N.S.W.	Q.	...	Moist clay soil	...	Outer peridium for most part seven to nine lobed; inner peridium shortly but distinctly stalked, size of pea, white.
1363	W.A.	...	...	...	...	...	...	Sandy soil	...	Outer peridium splitting into eight lobes; inner peridium globose, becoming whitish.
1364	...	...	...	V.	N.S.W.	...	...	Ground	...	Outer peridium thin, cut into seven to nine lobes, umber within; inner peridium somewhat stalked, globose, ochrey umber.
1365	W.A.	...	...	...	N.S.W.	Q.	B.	Ground	...	Outer peridium rigid, cut into about six lobes, reddish; inner peridium somewhat ovate, pale.
1366	W.A.	...	T.	...	N.S.W.	Q.	B.	Ground	...	Outer peridium cut into six to nine lobes, thin, soft; inner peridium globose, collapsed.
1367	...	S.A.	...	V.	...	Q.	...	Ground	...	Outer peridium split into eight to sixteen stellate fringes; inner peridium globose, tough, yellowish tan.
1368	...	S.A.	...	...	...	Q.	...	Ground	...	Small. Outer peridium with few lobes, mealy outside, smooth inside, umber or tawny; inner peridium conically globose.
1369	W.A.	...	...	V.	...	Q.	B.	Ground	...	Outer peridium often multifid beyond the middle, brown within; inner peridium globose, umber.
1370	...	...	...	...	...	Q.	...	Rotten wood	...	Gregarious, springing from expanded, white subiculum, or filamentous mass. Outer peridium mealy, wood colour; inner peridium darker, globose.
1371	...	...	T.	...	...	...	...	Ground	...	Outer peridium soft, papery, pale umber, about seven lobed; inner peridium on long stalk, globose, dark brown.
1372	...	...	...	...	...	...	...	Ground	...	Outer peridium membranous to leathery, cut into about eight fringes, tan colour, cracked lengthwise as if channelled; inner peridium globose, tawny.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
129. DIPLODERMA.—Link,					
1373	1275	IX. 1125	<i>D. album</i> ...	... Cooke and Mass., Grev. XVI. 2 (1887) ...	... White diploderma ...
1374	1276	„ 1126	<i>D. fumosum</i> ...	... Cooke and Mass., Grev. XVI. 2 (1887) ...	... Smoke-coloured spored diploderma
1375	1272	VII. 269	<i>D. glaucum</i> ...	... Cooke and Mass., Grev. XV. 99 (1887) ...	... Glaucons diploderma ...
1376	1277	...	<i>D. melaspermium</i> ...	... Cooke and Mass., Grev. XX. 35 (1891) ...	... Dark-spored diploderma ...
1377	1274	IX. 1127	<i>D. pachythrix</i> ...	... Cooke and Mass., Grev. XVIII. 50 (1890) ...	... Thick-fibred diploderma ...
1378	...	...	<i>D. sabulosum</i> ...	... Cooke and Mass., Grev. XXI. 38 (1892) ...	... Sandy diploderma ...
1379	1273	VII. 270	<i>D. suberosum</i> ...	... Cooke and Mass., Grev. XV. 100 (1887) ...	... Corky diploderma ...
130. BOVISTA.—Scop.,					
1380	1282	IX. 1130	<i>B. anomala</i> ...	... Cooke and Mass., Grev. XVIII. 6 (1889) ...	... Anomalous bovista ...
1381	...	VII. 283	<i>B. brasiliensis</i> ...	... <i>De Ton.</i> , Sacc. Syll. VII. 100 (1888) ...	... Brazilian bovista ...
1382	1278	„ 283	<i>B. brunnea</i> ...	... Berk., Fl. N. Zeal. II. 189 (1855) ...	... Brown bovista ...
1383	1284	„ 291	<i>B. cervina</i> ...	... Berk., Ann. Nat. Hist. IX. 447 (1842) ...	... Fawn-coloured bovista ...
1384	1303	„ 296	<i>B. dermoxantha</i> ...	... <i>De Ton.</i> , Sacc. Syll. VII. 100 (1888) ...	... Yellow-skinned bovista ...
1385	1280	„ 1605	<i>B. hyalothrix</i> ...	... Cooke and Mass., Grev. XVI. 73 (1888) ...	... Colourless-threaded bovista ...
1386	1281	...	<i>B. hypogæa</i> ...	... Cooke and Mass., Grev. XX. 35 (1891) ...	... Subterranean bovista ...
1387	1279	VII. 293	<i>B. Mnelleri</i> ...	... Berk., Linn. Journ. XIII. 171 (1873) ...	... Mneller's bovista ...
1388	1307	„ 286	<i>B. mundula</i> ...	... <i>De Ton.</i> , Sacc. Syll. VII. 98 (1888) ...	... Neat bovista ...
1389	1283	„ 1600	<i>B. olivacea</i> ...	... Cooke and Mass., Grev. XVI. 77 (1888) ...	... Olive bovista ...
1390	1306	„ 325	<i>B. pusilla</i> ...	... <i>De Ton.</i> , Rev. Geast (1887) ...	... Little bovista ...
131. LYCOPERDON.—Linn.,					
1391	1298	VII. 387	<i>L. australe</i> ...	... Berk., Fl. Tasm. II. 266 (1860) ...	... Southern lycoperdon (or puff-ball)
1392	1293	„ 324	<i>L. Bovista</i> ...	... Linn., Sp. Pl. 1653 (1753) ...	... Bovista lycoperdon ...
1393	1287	IX. 1133	<i>L. bovistoides</i> ...	... Sacc., Bull. Soc. Myc. Fr. V. 118 (1889) ..	... Bovista-like lycoperdon ...
1394	1295	VII. 352	<i>L. cælatum</i> ...	... Bull., Champ. 430 (1812) ...	... Embossed lycoperdon ...
1395	1296	„ 1615	<i>L. Cookei</i> ...	... Mass., Mon. Lyc. Trans. R.M.S. 714 (1887) ...	... Cooke's lycoperdon ...
1396	1301	...	<i>L. coprophilum</i> ...	... Cooke and Mass., Grev. ...	... Dung-loving lycoperdon ..
1397	1289	VII. 320	<i>L. gemmatum</i> ...	... Batsch., Elen. 147 (1783) ...	... Warty lycoperdon ... (Root fungus)
1398	1292	„ 386	<i>L. glabrescens</i> ...	... Berk., Fl. Tasm. II. 226 (1860) ...	... Smooth lycoperdon ...
1399	1309	„ 341	<i>L. Gunnii</i> ...	... Berk., Fl. Tasm. II. 265 (1860) .	... Gunn's lycoperdon ...
1400	1285	„ 403	<i>L. lilacinum</i> ...	... <i>Spæg.</i> , Fuog. Arg. 110 (1882) ...	... Lilac lycoperdon ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
1373	...	...	...	...	N.S.W.	...	...	Ground	Somewhat globose. Outer peridium thin, persistent; inner peridium whitish, cartilaginous.
1374	...	...	...	V.	N.S.W.	...	...	Ground	Globose, depressed, white. Outer peridium fibrous; inner peridium pale, fragile.
1375	...	...	T.	...	...	...	...	Ground, amongst sand	Somewhat globose, greyish green. Outer peridium fragile, soon falling away; inner peridium thin, yellowish.
1376	...	...	...	V.	...	...	...	Ground	Somewhat globose. Outer peridium thin, persistent, densely velvety, grey; inner peridium cinnamon.
1377	...	...	...	V.	...	...	...	Underground	Somewhat globose, about 1 inch in diameter. Outer peridium thin, ashy; inner peridium pale. Sterile threads thick, fibrous.
1378	...	S.A.	...	...	...	...	...	Sandy soil	Nearly globose, pale. Outer peridium thick, somewhat gelatinous, collecting grains of sand; inner peridium membranous.
1379	...	S.A.	...	...	...	Q.	...	Ground	Somewhat globose. Outer peridium corky, ochrey; inner peridium cartilaginous, turning black.

Berl. Mag. VII. 44 (1816).

Carn. II. 487 (1772).

1380	...	...	...	V.	...	...	...	Ground	Somewhat globose, depressed, whitish. Peridium thick, leathery, delicately velvety, dingy ochre.
1381	...	...	...	...	...	Q.	...	Trunks	Tufted. Peridium membranous, persistent, globose, rough pointed, rooting.
1382	...	...	...	...	N.S.W.	...	...	Ground	Globose, about 1 inch across, with minute apiculate rooting base. Peridium brownish umber, smooth, shining.
1383	...	...	...	V.	...	Q.	...	Ground	Small, globose. Peridium membranous, pale, fawn-coloured, cortex rather rigid.
1384	...	...	...	V.	...	...	...	Grassy places	Peridium very thin, sessile, irregularly globose, root rather long, slender, bright yellow becoming brownish.
1385	...	...	...	V.	...	...	...	Ground	Somewhat globose. Cortex very thick and fibrous, forming persistent base like acorn cup. Peridium minutely rugged.
1386	...	...	...	V.	...	...	...	Subterranean or partly exposed	Globose and depressed. Outer cortex persistent, thin, white, silky; inner layer thin, whitish, flexible.
1387	...	...	...	...	...	Q.	...	Ground	Somewhat globose, with short stout rooting base. Cortex soon broken up into minute warts. Peridium firm, brown.
1388	...	...	...	V.	...	...	...	Ground	Peridium fluffy, becoming smooth, white, size of hazel-nut.
1389	...	...	...	V.	...	...	B.	Ground	Globose, with short stout rooting base. Cortex very thin and evanescent. Peridium thick, soft, and pliant like leather, pale ochrey.
1390	W.A.	...	...	V.	N.S.W.	Q.	B.	Ground	Among the smallest of puff-balls. Somewhat globose, slightly tapering below, with minute scurfy scales, becoming smooth, pale-olive ochre.

Sp. Pl. II. 1183 (1753).—Bovista.

1391	W.A.	S.A.	T.	V.	N.S.W.	Q.	...	Sand turfy meadows	Sessile, globose, and depressed, densely covered with small pointed warts. Root long tapering.
1392	...	...	...	...	...	Q.	B.	Ground	Peridium somewhat spherical, sessile, fragile above, greyish white to yellowish white, then olive grey. <i>Edible</i> .
1393	...	S.A.	...	V.	...	...	...	Ground	Peridium nearly sessile, attached by a broad base, globose, then depressed, membranous, yellowish.
1394	...	...	T.	V.	N.S.W.	Q.	B.	Among grass	Solitary, sessile or stalked, large, somewhat globose or depressed. Cortex or bark pale-creamy ochre, broken into large angular patches.
1395	...	...	...	...	N.S.W.	...	B.	Ground	Hemispherical or globose, abruptly contracted into short thick stem-like base, smoky brown above, white below.
1396	...	...	...	...	...	Q.	...	Dung	Globose, sessile, whitish, covered with delicate persistent spines.
1397	W.A.	...	T.	...	N.S.W.	Q.	B.	Ground	Stalked, somewhat globose, depressed above, with prominent spiny warts, eventually falling off. Stem stout.
1398	...	...	T.	V.	...	...	...	Ground	Nearly hemispherical, at first covered with slender spines becoming smooth. Stem short, stout.
1399	...	...	T.	V.	...	Q.	...	Pastures	Sessile, somewhat globose, with very minute stellate warts.
1400	W.A.	...	T.	V.	N.S.W.	Q.	...	Ground	Broadly obovate, contracted below into stout stem-like base. Peridium thin; cortex white, polished.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
131. LYCOPERDON.— <i>Linn.</i> ,					
1401	1288	VII. 1610	<i>L. natalense</i> ...	Cooke and Mass., in Mon. Lye. Trans. R.M.S. 709 (1887)	Natal lycoperdon ...
1402	1291	„ 359	<i>L. pyriforme</i> ...	Schaeff., Leon 185 (1762) ...	Pear-shaped lycoperdon
1403	1304	.. 330	<i>L. reticulatum</i> ...	Berk., Fl. N. Zeal. II. 190 (1855) ...	Reticulated lycoperdon ...
1404	1299	„ 333	<i>L. stellatum</i> ...	Cooke and Mass., Grev. XV. 97 (1887) ...	Stellate lycoperdon ...
1405	1300	„ 1621	<i>L. substellatum</i> ...	Berk. and Curt., in Mass. Mon. Lye. Trans. R.M.S. 720 (1887)	Sub-stellate lycoperdon ...
1406	1305	„ 437	<i>L. tephrum</i> ...	Berk., in Mass. Mon. Lye. Trans. R.M.S. 723 (1887)	Ash-coloured lycoperdon ...
1407	1286	„ 1607	<i>L. violascens</i> ...	Cooke and Mass., Mon. Lye. Trans. R.M.S. 706 (1887)	Violet lycoperdon ...
132. AREOLARIA.— <i>Forq.</i> , Champ.					
1408	1318	VII. 481	<i>A. strobilina</i> ...	<i>Forq.</i> , Champ. Exot. 155 (1886) ...	Cone-like areolaria ...
133. CASTOREUM.—Cooke and Mass.,					
1409	1322	VII. 476	<i>C. radicans</i> ...	Cooke and Mass., Grev. XV. 100 (1887) ...	Rooting castoreum ...
134. XYLOPODIUM.— <i>Mont.</i> ,					
1410	1323	VII. 479	<i>X. australe</i> ...	Berk., Linn. Journ. XIII. 171 (1873) ...	Southern xylopodium ...
1411	1324	„ 478	<i>X. ochroleucum</i> ...	Cooke and Mass., Grev. XV. 95 (1887) ...	Whitish-ochre xylopodium ...
135. FAVILLEA.— <i>Fries</i> ,					
1412	1325	VII. 487	<i>F. argillacea</i> ...	Fries, Fung. Nat. 32 (1848) ...	Clay-coloured favillea ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Sp. Pl. II. 1183 (1753).— <i>Bovista</i> — <i>continued</i> .									
1401	...	...	...	V.	...	...	...	Ground ...	Globose, sessile, passing abruptly into short tapering root. Peridium thick, minutely warted, becoming smooth.
1402	...	...	T.	V.	N.S.W.	Q.	B.	On stumps or soil, attached to branches, &c.	Densely tufted, pear shaped, membranous, covered with minute-pointed warts, brownish, rooting.
1403	...	...	...	V.	...	...	...	Ground ...	Globose, tapering downward, with slightly-raised reticulations, eventually disappearing and leaving polished surface.
1404	W.A.	...	...	...	...	...	...	Ground ...	Sessile, nearly globose. Peridium thin, covered at first with stout stellate spiny warts, falling away in patches and leaving smooth surface.
1405	..	...	...	...	...	Q.	...	Rotten wood, &c. ...	Globose, sessile, whitish, covered with delicate spines, which become smaller downwards.
1406	...	...	...	...	...	Q.	...		Globose, sessile, thick and rigid, brown, minutely velvety.
1407	...	...	...	V.	...	...	...	Ground ...	Globose, sessile, terminating in short slender root. Peridium papery, covered at first with minute warts, becoming smooth and shining.

Exot. 155 (1886).—*Scleroderma*, *Phellorina*.

1408	...	...	...	...	...	Q.	...	Ground ...	Globose and depressed, with stout angular scales above. Stem solid, rather woody.
------	-----	-----	-----	-----	-----	----	-----	------------	---

Grev. XV. 100 (1887).

1409	...	...	T.	...	...	...	...	Ground ...	Tufted, nearly globose, confluent below in tough rooting stem. Outer peridium tawny, leathery; inner peridium at length horny.
------	-----	-----	----	-----	-----	-----	-----	------------	--

Ann. Sci. Nat. 3 Ser. IV. 364 (1845).

1410	...	S.A.	...	V.	N.S.W.	...	...	Ground, trunk of <i>Eucalyptus hemiphloia</i>	Peridium volvate when young. Stem rooting, broken into scales.
1411	...	...	...	...	...	Q.	...	Ground ...	Stalked. Peridium globose, with large warts. Stem erect, thick, solid, with overlapping scales.

Fung. Nat. 32 (1848).

1412	...	...	...	...	...	...	...	Ground ...	Peridium club shaped, simple, without special cortex, membranous above.
------	-----	-----	-----	-----	-----	-----	-----	------------	---

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER XI.—SCLERODERMACEÆ,					
136. SCLERODERMA.—Pers., Syn.					
1413	1316	IX. 1142	<i>S. australe</i> ...	Mass., Grev. XVIII. 26 (1889) ...	Southern scleroderma ...
1414	1311	VII. 446	<i>S. Bovista</i> ...	Fries, S.M. III. 40 (1829) ...	Bovista scleroderma ...
1415	1319	„ 474	<i>S. corium</i> ...	<i>Grav.</i> , in Duby. Bot. Gall. II. 892 (1830) ...	Leathery scleroderma ...
1416	1310	„ 459	<i>S. Geaster</i> ...	Fries, S.M. III. 46 (1829) ...	Earth-star scleroderma ...
1417	1321	„ 1629	<i>S. olivaceum</i> ...	<i>De Ton.</i> , Sacc. Syll. VII. 489 (1888) ...	Olive scleroderma ...
1418	1314	„ 454	<i>S. Pandanaceum</i> ...	F. v. M., Linn. Journ. XIII. 171 (1873) ...	Pandanus scleroderma...
1419	1320	„ 461	<i>S. phæotrichum</i> ...	<i>De Ton.</i> , Sacc. Syll. VII. 139 (1888) ...	Dusky-haired scleroderma ...
1420	...	...	<i>S. pileolatum</i> ...	Kalch., Linn. Soc. N.S.W. VII. 565 (1882) ...	Capped scleroderma ...
1421	1317	IX. 1146	<i>S. umbrinum</i> ...	Cooke and Mass., Grev. XIX. 45 (1890) ...	Umber scleroderma ...
1422	1313	VII. 447	<i>S. verrucosum</i> ...	<i>Pers.</i> , Syn. 154 (1801) ...	Warted scleroderma ...
1423	1312	„ 445	<i>S. vulgare</i> ...	Fries, S.M. III. 46 (1829) ...	Common scleroderma ...
137. POLYSACCUM.—D. C., Fl. Fr. V. 103 (1815).—					
1424	1334	...	<i>P. album</i> ...	Cooke and Mass., Grev. XX. 36 (1891) ...	White polysaccum ...
1425	1333	VII. 490	<i>P. australe</i> ...	Lev., Ann. Sci. Nat. Ser. 3, IX. 136 (1848) ...	Southern polysaccum ...
1426	1332	„ 1632	<i>P. confusum</i> ...	Cooke, Grev. XVI. 76 (1888) ...	Confused polysaccum ...
1427	1328	„ 491	<i>P. crassipes</i> ...	D. C., Fl. Fr. V. 103 (1815) ...	Thick-stalked polysaccum ...
1428	1335	„ 500	<i>P. degenerans</i> ...	Fries, Pl. Preiss. 139 (1846) ...	Degenerating polysaccum ...
1429	1331	„ 499	<i>P. marmoratum</i> ...	Berk., Linn. Journ. XIII. 171 (1873) ...	Marbled polysaccum ...
1430	1327	„ 1633	<i>P. microcarpum</i> ...	Cooke and Mass., Grev. XVI. 28 (1887) ...	Small-fruited polysaccum ...
1431	1326	„ 494	<i>P. pisocarpium</i> ...	Fries, S.M. III. 54 (1829) ...	Pea-fruited polysaccum ...
1431a	„	„	<i>P. pisocarpium</i> , var. <i>acaule</i> ...	D. C., Fl. Fr. V. 103 (1815) ...	Stemless polysaccum ...
1432	1330	„ 501	<i>P. tuberosum</i> ...	Fries, Linn. V. 694 (1830) ...	Tuberous polysaccum ...
1433	1329	„ 489	<i>P. turgidum</i> ...	Fries, S.M. III. 53 (1829) ...	Turgid polysaccum ...
138. ARACHNION.—Schwein.,					
1434	1336	VII. 507	<i>A. Drummondi</i> ...	Berk., Linn. Journ. XVIII. 389 (1887) ...	Drummond's arachnion ...
139. PAUROCOTYLIS.—Berk.,					
1435	1338	VII. 512	<i>P. echinosperma</i> ...	Cooke, Grev. VIII. 59 (1879) ...	Spiny-spored paurocotylis ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.	
	W.A.	S.A.	T.	V.	N.S.W.	Q.				
FRIES, S.M. III. 5 (1829).										
Fung. 150 (1801).—Lycoperdon, Mycenastrum.										
1413	...	...	...	...	...	Q.	...	Soil	...	Nearly globose, sessile. Peridium thick, externally minutely felty, dirty ochre, with rooting base.
1414	...	...	...	V.	N.S.W.	Q.	B.	Sandy ground	...	Obovoid to spherical, somewhat stalked or possibly with rooting base, warty, turning yellow.
1415	...	...	...	V.	...	Q.	...	Sandy ground	...	Nearly globose, whitish, then greyish brown, leathery, splitting above in stellate manner.
1416	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Sandy ground	...	Sessile, nearly globose, opening at top in stellate fringes.
1417	...	...	...	...	N.S.W.	Q.	...	Ground	...	Nearly globose, sessile, lead colour above, whitish below; sterile threads and mass of spores olive.
1418	...	...	...	...	...	Q.	...	Ground	...	Globose, sessile, yellow, tessellate above.
1419	W.A.	...	...	...	N.S.W.	Q.	...	Ground	...	Nearly globose; sterile threads and mass of spores dark purple brown.
1420	...	...	...	...	N.S.W.	...	...	...	...	Peridium globose, opaque, umber beneath, concave, size of hazel-nut, with slender stem.
1421	...	...	...	...	...	Q.	...	Ground	...	Peridium globose, coarsely wrinkled below, dirty pale ochre. Stem coarsely furrowed, dark brown, passing into dense bulbous mass.
1422	...	...	...	...	...	Q.	B.	Sandy ground	...	Peridium rounded, at first hard, then fragile, covered with rather warted cortex, dingy yellowish.
1423	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Ground under trees, &c.	...	Almost sessile, deformed. Peridium corky, citron then reddish brown, growing pale with age.

Lycoperdon, Lycoperdoides.

1424	...	...	...	...	...	Q.	...	Ground	...	Peridium globose, white, polished and shining, tapering below into very short stout irregular stem-like base.
1425	W.A.	...	...	...	...	Q.	...	Ground	...	Stem rooting, almost cylindrical, shining, blackish brown, dilated into like-coloured nearly globose peridium.
1426	...	S.A.	...	V.	...	...	...	Ground	...	Peridium nearly globose, slightly narrowed below into short thick stem, or pear shaped, olivaceous umber with polygonal cavities.
1427	W.A.	S.A.	...	V.	...	...	...	Immersed in sand	...	Peridium from spherical to clavate, pale ochre becoming darker; stem-like base stout, minute cavities.
1428	W.A.	...	...	...	...	...	...	River banks	...	Club shaped with rooting stem. Peridium simple, ochrey-tan colour.
1429	W.A.	S.A.	...	...	N.S.W.	...	...	Ground	...	Peridium somewhat globose, tapering to stem-like base, dirty ochre, marbled with darker patches.
1430	...	...	...	V.	...	Q.	...	Ground	...	Peridium nearly globose, coarsely tubercled, ochrey brown, stout, with stem-like base, bright citrine.
1431	W.A.	...	...	V.	...	Q.	B.	Sandy ground	...	Peridium nearly globose, passing down into short stem-like base, reddish brown, tinged olive.
1431A	...	...	...	...	...	Q.	...	Sandy ground	...	Without distinct stem.
1432	...	...	...	...	N.S.W.	Q.	...	Ground	...	Peridium nearly globose or deformed, with short stem-like base, ochrey; cavities large, angular.
1433	W.A.	...	...	...	...	...	...	Sandy soil	...	Peridium cylindric to club shaped, covered at first with fine spider-like web, dark umber, passing into stem-like base.

Syn. Car. 14 (1822).

1434	W.A.	...	...	...	...	...	...	Attached to <i>Locellinia cychnotamia</i>	Globose, depressed a little, pale.
------	------	-----	-----	-----	-----	-----	-----	---	------------------------------------

Fl. N. Zeal. II. 188 (1855).

1435	...	...	...	V.	...	...	...	Trunks	...	Globose, depressed, tawny flesh colour, spores spiny.
------	-----	-----	-----	----	-----	-----	-----	--------	-----	---

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER XII.—HYMENOGASTRACEÆ,					
140. OCTAVIANIA.—Vitt., Mon.					
1436	1340	IX. 1150	<i>O. alveolata</i> ...	Cooke and Mass., Grev. XVI. 2 (1887) ...	Alveolate octaviana ...
1437	1341	VII. 529	<i>O. Archeri</i> ...	Berk., Fl. Tasm. II. 263 (1860) ...	Archer's octaviana ...
1438	1339	, 578	<i>O. australiense</i> ...	Berk., in Cooke's Handb. Austr. Fung. 246 (1892)	Australian octaviana ...
141. RHIZOPOGON.—Fries,					
1439	1342	VII. 534	<i>R. luteolus</i> ...	Fries, Symb. Gast. 5 (1818) ...	Yellowish rhizopogon ...
142. HYMENOGASTER.—Vitt., Mon. Tub. 30 (1831).—					
1440	1343	VII. 560	<i>H. Klotzschii</i> ...	Tul., Fung. Hyp. 64 (1851) ...	Klotzsch's hymenogaster ...
1441	1344	, 563	<i>H. lycoperdineus</i> ...	Vitt. Mon. Tub. 22 (1831) ...	Puff-ball hymenogaster ...
1442	1345	, 564 bis.	<i>H. Moselei</i> ...	<i>De Ton.</i> , Sacc., Syll. VII. 172 (1888) ...	Moseley's hymenogaster ...
143. HYDNANGIUM.—Wallr., in Corda					
1443	1346	...	<i>H. brisbanensis</i> ...	Berk. and Br., in Cooke's Handb. Austr. Fung. 247 (1892)	Brisbane hydnangium ...
1444	1347	...	<i>H. tasmanicum</i> ...	Kalch., in Grev. XIX. 95 (1891) ...	Tasmanian hydnangium ...
144. GAUTIERIA.—Vitt.,					
1445	1348	...	<i>G. Drummondi</i> ...	Berk., in Cooke's Handb. Austr. Fung. 247 (1892)	Drummond's gautieria ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
VITT. MON. TUB. 11 (1831).									
Tub. 15 (1831).—Hyduangium.									
1436	...	...	...	...	N.S.W.	...	...	In the ground	... Somewhat globose or irregular, whitish, then ochrey. Spores globose, alveolate.
1437	...	...	T.	...	...	...	...	Ground	... Inversely egg shaped, small, with large sterile base, no fibrils.
1438	...	...	...	V.	...	...	Q.	Under tea tree	... Somewhat globose, irregular. Peridium thin, wrinkled, ochrey.
Symb. Gast. 5 (1818).—Hysterangium, Tuber.									
1439	...	...	...	V.	...	...	B.	In sandy soil	... Deformed, usually spherical, kidney or egg shaped; fibrils rooting becoming yellowish.
Rhizopogon, Splachnomyces, Hymenangium.									
1440	W.A.	...	...	...	...	...	B.	In ground	... Obovate, fibrillose at base. Peridium membranous, whitish, somewhat, downy.
1441	...	...	...	...	...	Q.	...	Ground	... Gregarious, with strong odour of garlic. Peridium rounded, deformed, white, then brownish.
1442	...	...	...	...	N.S.W.	...	...	Soil	... Almost globose, tapering at base, citrine yellow, smooth.
Ic. Fung. V. 28 (1842).—Octaviania.									
1443	...	...	...	V.	...	Q.	...	In soil	... Almost globose. Peridium thick, dry, wrinkled, reddish brown.
1444	...	...	T.	...	...	...	...	Ground	... Nearly globose. Peridium thick, dark brown, angular when dry.
Mon. Tub. 25 (1831).									
1445	W.A.	...	...	...	...	...	...	In soil	... Nearly globose, small; cells wavy. Spores ellipsoid.

GENERAL CLASSIFICATION OF UREDINES.

GROUP III.—UREDINES, BRONGN.

ORDER XIII.—UREDINACEÆ—Parasitic. Spores usually of more than one kind and not all unicellular.

ARRANGEMENT OF GENERA (9).

Section 1. Amerosporæ, Sacc. and De Toni—Teleutospores continuous, one-celled.

Genera (3)—

145. Uromyces, Link. | 146. Melampsora, Cast. | 147. Cronartium, Fries.

Section 2. Didymosporæ, Sacc. and De Toni—Teleutospores bilocular.

Genus (1)—

148. Puccinia, Pers.

Section 3. Phragmosporæ, Sacc. and De Toni—Teleutospores 3, or many-celled.

Genera (2)—

149. Phragmidium, Link. | 150. Hamaspora, Kœrn.

Imperfect Forms—

Genera (3)—

151. Æcidium, Pers. | 152. Roestelia, Reb. | 153. Uredo, Pers.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
GROUP III.—UREDINES.—					
ORDER XIII.—UREDINACEÆ,					
145 UROMYCES.—Link, Berl. Mag. VII. 28 (1816).—					
1446	1732	VII. 1928	U. Betæ ...	... <i>Kuehn</i> , Bot. Zeit. 540 (1869) ...	Beet uredomyces ...
1447	1738 bis.	„ 2033	U. Bulbinis ...	... <i>Thuem.</i> Fl. 410 (1877) ...	Balbine uredomyces ...
1448	1737	„ 1982	U. digitatus ...	... <i>Wint.</i> Rev. Myc. 209 (1886) ...	Digitate uredomyces ...
1449	1738	IX. 1212	U. Diploglottidis ...	... <i>Cooke and Mass.</i> , Grev. XVII. 55 (1889) ...	Diploglottis uredomyces ...
1450	1735	VII. 1980	U. fusisporus ...	... <i>Cooke and Mass.</i> , Grev. XVI. 2 (1887) ...	Spindle-spored uredomyces ...
1451	...	„ 1940	U. Junci ...	... <i>Tul.</i> , Ann. Sci. Nat. 146 (1854) ...	Rush uredomyces ...
1452	...	...	U. Kuehnii ...	... <i>Krueger</i> , Bericht. Versuchs. Java I. (1891) ...	Kuehn's uredomyces (Cane rust) ...
1453	...	VII. 1209	U. Limosellæ ...	... <i>Ludw.</i> , Diet. Hedw. 182 (1888) ..	Limosella uredomyces ...
1454	1740	„ 2066	U. Microtidis ...	... <i>Cooke</i> , Grev. XIV. 12 (1885) ...	Microtis uredomyces ...
1455	1741	„ 2071	U. orchidearum ...	... <i>Cooke and Mass.</i> , Grev. XVI. 74 (1888) ...	Orchid uredomyces ...
1456	1736	IX. 1203	U. phyllodii ...	... <i>Cooke and Mass.</i> , Grev. XVII. 70 (1889)...	Phyllode uredomyces ...
1457	1742	VII 2100	U. puccinioides ..	... <i>Berk.</i> and <i>F. v. M.</i> , Linn. Journ. XIII. 173 (1873)	Puccinia-like uredomyces ..
1458	1739	IX. 1204	U. Tepperianus ...	... <i>Sacc.</i> , Hedw. 126 (1889) ...	Tepper's uredomyces ...
1459	1731	VII. 1925	U. Trifolii ...	... <i>Lev.</i> in <i>Winter's Die Pilze</i> I. 159 (1884) ...	Clover uredomyces ...
1460	1733	„ 1955	U. vesiculosa ...	... <i>Wint.</i> , Hedw. 2 (1885) ...	Vesicular uredomyces ...
146. MELAMPSORA.—Cast.,					
1461	1743	VII. 2107	M. Lini ...	... <i>Tul.</i> , Ann. Sci. Nat. 93 (1854) ...	Flax melampsora ...
1462	1741	„ 2124	M. Nesodaphnes ...	... <i>Berk.</i> and <i>Br.</i> , Linn. Trans. II. 67 (1883)	Nesodaphnes melampsora ...
1463	1745	„ 2123	M. phyllodiorum ...	... <i>Berk.</i> and <i>Br.</i> , Linn. Trans. II. 67 (1883)	Phyllode melampsora ...
147. CRONARTIUM.—Fries,					
1464	1746	VII. 2137	C. Asclepiadeum ...	... <i>Fries</i> , Obs. Myc. I. 220 (1815) ...	Aselepiad Cronartium ...
148. PUCCINIA.—Pers.,					
1465	1760	VII. 2218	P. Acetosæ ...	... <i>Karn.</i> , Hedw. 184 (1876) ...	Sorrel puccinia ..
1466	1751	„ 2174	P. ægra ...	... <i>Grove</i> , Journ. Bot. 274 (1883) ...	Sick puccinia ...
1467	1768	„ 2494	P. Alyxiæ ...	... <i>Cooke and Mass.</i> , Grev. XVI. 2 (1887) ...	Alyxia puccinia ...
1468	1757	„ 2211	P. Apii ...	... <i>Corda</i> , Icon. VI. 30 (1854) ...	Celery puccinia ...
1469	1765	„ 2337	P. aucta ...	... <i>Berk.</i> and <i>F. v. M.</i> , Linn. Journ. XIII. 173 (1873)	Abundant puccinia ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
1446	...	S.A.	...	V.	...	...	B.	Leaves of <i>Beta</i> ...	Cluster cups on circular or oblong yellow spots. Uredospore pustules cinnamon. Teleutospore pustules dark brown.
1447	...	...	...	V.	N.S.W.	...	...	Leaves of <i>Bulbine bulbosa</i>	Pustules on both surfaces, small, densely crowded, covered by cuticle, brown.
1448	...	S.A.	...	...	...	...	...	Leaves of <i>Acacia notabilis</i>	Pustules in centre of circular spots, bounded by narrow brown line, black. Teleutospores with finger-like processes.
1449	...	...	...	...	...	Q.	...	Fading leaves of <i>Diploglottis</i>	Pustules scattered, convex, minute, at length splitting, pale brown, on circular greenish spots.
1450	...	...	...	V.	...	...	...	Phyllodes of <i>Acacia salicina</i>	On both surfaces. Pustules disc-like, bursting through, black, surrounded by ruptured cuticle.
1451	...	...	...	V.	...	...	B.	<i>Juncus maritima</i> ...	Cluster cups cup shaped with whitish torn edges. Uredospore pustules on brownish spots. Teleutospore pustules round or elongated.
1452	...	...	...	...	N.S.W.	...	...	Leaves of Sugar cane	Mostly on under surface, spots finally orange. Pustules in streaks, elongated, narrow, brownish or blackish, bursting through, often, run together.
1453	...	S.A.	...	...	...	...	...	Leaves of <i>Limosella</i>	Cluster cups on both sides, scattered or gregarious. Teleutospore pustules mixed with cluster cups.
1454	...	...	...	...	N.S.W.	...	...	Leaves of <i>Microtis porrifolia</i>	On both surfaces. Pustules gregarious, dark brown, surrounded by epidermis.
1455	...	...	...	...	N.S.W.	...	...	Leaves of <i>Chiloglottis diphylla</i>	Pustules blistered, at length bursting, brown.
1456	...	...	...	...	...	Q.	...	Phyllodes of <i>Acacia</i>	Pustules minute, circular, compact, brown, crowded on blistered spots, at length naked.
1457	...	S.A.	T.	V.	...	Q.	...	Leaves and flower stalks of <i>Goodenia</i> and <i>Selliera</i>	Cluster cups on brown circular spots (<i>Æcidium Goodeniacearum</i> , Berk.). Pustules blistered.
1458	...	S.A.	...	V.	...	...	...	Living branches of <i>Acacia salicina</i> , <i>A. myrtifolia</i> , and <i>A. hakeoides</i>	Long and broad, flattened, growing beneath cuticle and casting off bark, bright cinnamon.
1459	...	...	...	V.	...	...	B.	Clover ...	Cluster cups circularly arranged, pale orange. Uredospore clusters chestnut brown. Teleutospore clusters smaller.
1460	...	S.A.	...	...	...	...	...	Leaves and stems of <i>Zygophyllum ammodendron</i>	Pustules scattered or gregarious, often run together, covered by ash- vesicular epidermis.

Obs. II. 18 (1843).—Uredo, Cæoma.

1461	...	S.A.	...	V.	N.S.W.	...	B.	Leaves of <i>Linum marginale</i> and <i>L. usitatissimum</i>	Uredospore pustules scattered, rounded, orange, minute. Teleutospore pustules at first red brown, then almost black.
1462	...	...	...	...	...	Q.	...	Fruit of <i>Nesolaphnes obtusifolia</i>	Spore masses powdery, shaggy, ochrey.
1463	...	...	...	...	...	Q.	...	Phyllodes of <i>Acacia</i>	Pustules in tubercles; spores arising from delicate filaments, granulated.

Obs. Myc. I. 220 (1815).—Erineum, Uredo, Cæoma.

1464	...	...	...	...	...	Q.	...	Leaves of <i>Jacksonia scoparia</i>	Uredospore pustules on under surface, scattered or clustered, brown. Teleutospore pustules yellowish.
------	-----	-----	-----	-----	-----	----	-----	-------------------------------------	---

Tent. Disp. Meth. 38 (1797).—Uredo, Æcidium, Cæoma, Trichobasis, Uromyces.

1465	...	...	...	...	...	Q.	...	Leaves, stems, &c., of <i>Rumex</i>	Pustules of both forms scattered, minute on leaves and irregularly rounded, oblong on stems and leaf stalks.
1466	...	...	...	V.	...	...	B.	Leaves of <i>Viola hederacea</i>	Cluster cups scattered, white, spores orange yellow. Uredospore pustules on yellow spots. Teleutospore pustules similar.
1467	...	...	...	V.	...	...	...	Leaves of <i>Alyxia birifolia</i>	On under surface. Pustules disc-like, compact, dark brown, girt by ruptured epidermis.
1468	...	...	...	V.	...	...	B.	Celery ...	Cluster cups causing long yellow swellings on stem. Uredospore pustules large, cinnamon brown. Teleutospore pustules blackish brown.
1469	...	S.A.	...	V.	N.S.W.	...	...	Leaves of <i>Lobelia anceps</i> , <i>L. pedunculata</i> , <i>L. platycalyx</i>	Cluster cups occupying entire surface of leaves or leaf stalks, ochrey. Teleutospore pustules blistered.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name
148. PUCCINIA.—Pers.,					
1470	1769	VII. 2506	<i>P. Berkeleyana</i> ...	<i>De Toni</i> , in Sacc. Syll. VII. 717 (1888) ...	Berkeley's puccinia ...
1471	...	...	<i>P. Burchardiae</i> ...	Ludw., Zeit. f. Pilzkr. III. 137 (1893) ...	Burchardia puccinia ...
1472	...	...	<i>P. Carissae</i> ...	Cooke and Mass. XXII. 37 (1893) ...	Carissa puccinia ...
1473	...	VII. 2162	<i>P. Epilobii</i> ...	D. C., Fl. Fr. VI. 61 (1805) ...	Epilobium puccinia ...
1474	...	„ 2409	<i>P. Geranii</i> ...	Corda, Icon. IV. 12 (1854) ...	Geranium puccinia ...
1475	1753	„ 2191	<i>P. graminis</i> ...	Pers., Disp. Fung. 39 (1797) ...	Grass puccinia (Rust in wheat) ...
1476	1747	„ 2150	<i>P. Helianthi</i> ...	Schw., Syn. Car. 73 (1822) ...	Sunflower puccinia ...
1477	1767	„ 2403	<i>P. heterospora</i> ...	Berk. and Curt., Linn. Journ. X. 356 (1869) ...	Heterosporous puccinia ...
1478	1749	„ 2210	<i>P. Hieracii</i> ...	<i>Mart.</i> , Fl. Mosq. 226 (1812) ...	Hawk-weed puccinia ...
1479	...	...	<i>P. Junciphila</i> ...	Cooke and Mass. XXII. 37 (1893) ...	Rush-loving puccinia ...
1480	1761	VII. 2240	<i>P. Kalchbrenneri</i> ...	<i>De Toni</i> , Sacc. Syll. VII. 645 (1888) ...	Kalchbrenner's puccinia ...
1481	...	...	<i>P. Kochiae</i> ...	Mass., Grev. XXII. 17 (1893) ...	Kochia puccinia ...
1482	1752	VII. 2169	<i>P. Lagenophorae</i> ...	Cooke, Grev. XIII. 6 (1884) ...	Lagenophora puccinia ...
1483	1759	...	<i>P. Ludwighi</i> ...	Tepper, Bot. Centr. Blatt. 6 (1890) ...	Ludwig's puccinia ...
1484	...	VII. 2205	<i>P. Magnusiana</i> ...	Körn., Hedw. 179 (1876) ...	Magnus' puccinia ...
1485	1766	„ 2368	<i>P. Malvacearum</i> ...	Mont., in Gay's Hist. Chil. VIII. 43 (1845) ...	Mallow puccinia (Hollyhock fungus) ...
1486	...	...	<i>P. munita</i> ...	Ludw., Zeit. f. Pilzkr. II. 133 (1892) ...	Protected puccinia ...
1487	1787	VII. 2202	<i>P. obscura</i> ...	Schrot., Nuov. Giorn. Bot. Ital. IX. 256 (1875) ...	Obscure puccinia ...
1488	1756	„ 2204	<i>P. Phragmitis</i> ...	Körn., Hedw. 179 (1876) ...	Phragmitis puccinia ...
1489	1755	„ 2195	<i>P. Poarum</i> ...	Niels., Bot. Tids. II. 26 (1877) ...	Poa puccinia ...
1490	1748	„ 2157	<i>P. Prenanthis</i> ...	Fekl., Sym. Myc. 45 (1875) ...	Prenanthes puccinia ...
1491	1734	„ 2252	<i>P. Pruni</i> ...	Pers., Syn. Fung. 226 (1808) ...	Plum puccinia (Peach and Plum leaf rust) ...
1492	1770	...	<i>P. rimosa</i> ...	Link, Winter, Hedw. 28 (1880) ...	Cracking puccinia ...
1493	1754	VII. 2194	<i>P. Rubigo-vera</i> ...	Wint., Die Pilze. I. 217 (1884) ...	True-rust puccinia ...
1494	1758	„ 2214	<i>P. Rumicis-scutati</i> ...	Wint., Die Pilze. I. 187 (1884) ...	Rumex puccinia ...
1494A	„	IX. 1236	<i>P. Rumicis-scutati</i> , var. <i>Muehlenbeckia</i> ...	Cooke, Grev. XIX. 47 (1890) ...	Muehlenbeckia puccinia ...
1495	1764	„ 1280	<i>P. Saccardoi</i> ...	Ludw., Hedw. 362 (1889) ...	Saccardo's puccinia ...
1496	1762	VII. 2289	<i>P. Sorghi</i> ...	Schw., N. Am. Fung. 295 (1831) ...	Sorghum puccinia ...
1497	...	...	<i>P. Tepperi</i> ...	Ludw., Zeit. f. Pilzkr. II. 130 (1892) ...	Tepper's puccinia ...
1498	1750	VII. 2163	<i>P. Viole</i> ...	Wint., Die Pilze. I. 215 (1884) ...	Violet puccinia ...
1499	1763	„ 2304	<i>P. Wurnbseae</i> ...	Cooke and Mass., Grev. XVI. 71 (1888) ...	Wurnbseae puccinia ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Tent. Disp. Meth. 38 (1797).—Uredo, <i>Æcidium</i> , <i>Cæoma</i> , <i>Trichobasis</i> , <i>Uromyces</i> —continued.									
1470	...	...	...	V.	...	...	...	Leaves of <i>Dichondra repens</i>	Pustules minute on lower, rarely on upper surface, scattered, resembling a cluster cup.
1471	...	S.A.	...	V.	...	...	...	Leaves of <i>Burchardia umbellata</i>	Pustules bursting through, circular or elliptical, black. Uredospores and teleutospores.
1472	...	...	...	...	...	Q.	...	Living leaves of <i>Carissa ovata</i>	On under surface. Only teleutospores seen. Pustules small, gregarious on circular spots, dark brown.
1473	...	...	...	V.	...	...	B.	Leaves of <i>Epilobium glabellum</i>	Pustules small, roundish, rather crowded, soon naked, surrounded by torn epidermis, dark brown.
1474	...	...	...	V.	...	...	...	Leaves of <i>Pelargonium australe</i>	Pustules small, brown, powdery, on under surface. Teleutospore pustules black.
1475	...	S.A.	T	V.	N.S.W.	Q.	B.	<i>Avena</i> and <i>Triticum</i>	Cluster cups forming circular, reddish-yellow spots. Uredospore pustules linear (<i>Uredo linearis</i>). Teleutospore pustules elongated, black.
1476	...	...	...	V.	N.S.W.	Q.	...	Sunflower leaves	Cluster cups crowded or circular, spores orange red. Uredospore pustules minute, chestnut brown. Teleutospore pustules dark brown or black.
1477	...	...	...	...	N.S.W.	Q.	...	Leaves of <i>Abutilon crispum</i> and <i>A. arvense</i>	Spots purplish or yellow. Pustules on under surface, minute, soon naked, brown.
1478	...	...	...	...	...	Q.	B.	<i>Hypochaeris glabra</i>	Pustules oblong, mostly solitary, reddish brown, girt by ruptured epidermis.
1479	...	...	...	V.	...	...	...	<i>Juncus</i> ...	Pustules rusty, powdery, elliptical or confluent. Uredospores and teleutospores intermixed.
1480	...	...	...	V.	...	...	...	Leaves of <i>Helichrysium</i>	Pustules on both surfaces, scattered or gregarious, covered at first, then free, ochrey.
1481	...	...	...	V.	...	...	...	Leaves of <i>Kochia sedifolia</i>	On both surfaces of leaf. Pustules discoid, blackish brown, girt by torn epidermis.
1482	...	...	...	V.	...	...	...	Living leaves of <i>Lagenophora Biltardieri</i>	Cluster cups on upper surface. Uredospore pustules small, brown. Teleutospore pustules dark brown.
1483	...	...	...	V.	...	...	...	Leaves of <i>Rumex Brownii</i>	Uredospores pale yellow brown. Teleutospore pustules minute, circular girt by ruptured epidermis.
1484	...	S.A.	...	...	...	...	...	<i>Arundo phragmites</i>	Cluster cups mostly on under surface of leaves. Uredospore pustules orange brown. Teleutospore pustules black.
1485	...	S.A.	...	V.	N.S.W.	Q.	B.	Leaves and stems of <i>Althæa rosea</i> and <i>Malva rotundifolia</i>	Pustules greyish brown, compact, round, elongated on stems, scattered, pale reddish brown.
1486	...	S.A.	...	...	...	...	...	Leaves of <i>Hydrocotyle hirta</i>	On under surface. Pustules cinnamon brown. Teleutospores protected by white pseudo-peridium.
1487	...	...	...	V.	...	...	B.	<i>Bellis perennis</i> ...	Cluster cups on large roundish spots, mostly on upper surface (<i>Æcidium Bellidis</i>).
1488	...	...	...	V.	...	...	B.	<i>Arundo phragmites</i> . &c.	Cluster cups on circular red spots. Uredospore pustules large, dark brown. Teleutospore pustules sooty black.
1489	...	...	...	V.	N.S.W.	...	B.	<i>Poa</i> species ...	Cluster-cup spots yellow. Uredospore pustules small, orange. Teleutospore pustules black.
1490	...	S.A.	...	V.	N.S.W.	...	B.	<i>Lactuca</i> , &c. ...	Cluster cups in circular or elongated patches. Uredospore pustules reddish brown. Teleutospore pustules blackish.
1491	...	S.A.	T.	V.	N.S.W.	Q.	B.	Peach, Plum, and Almond leaves, also on fruit of Peach	Uredospore pustules light brown, small, round, crowded. Teleutospore pustules almost black.
1492	...	...	...	V.	...	...	...	<i>Isolepis nodosa</i> ...	Producing narrow cracks, often encircling stem and causing brown spots.
1493	...	S.A.	T.	V.	N.S.W.	Q.	B.	Cereals, <i>Poa annua</i> , &c.	Cluster cups on large circular spots. Uredospore pustules rust colour (<i>Uredo rubigo-vera</i>). Teleutospore pustules black, covered a long time.
1494	...	...	...	...	...	Q.	...	Leaves, leaf stalks, and stems of <i>Rumex</i>	Pustules scattered or arranged in circle, rounded or elongated, girt by torn epidermis, brown.
1494A	...	...	...	V.	...	...	...	Leaves of <i>Muehlenbeckia adpressa</i>	Pustules on upper surface, scattered, blistered at first.
1495	...	S.A.	...	V.	...	...	...	Leaves of <i>Goodenia geniculata</i>	Cluster cups in groups, on brownish spots. Teleutospore pustules rounded or elongated.
1496	...	...	...	...	N.S.W.	Q.	...	Maize ...	Uredospore pustules on both surfaces, reddish brown (<i>Uredo Maydis</i>). Teleutospores not noted.
1497	...	S.A.	...	...	...	...	...	<i>Arundo phragmites</i>	Generally resembling <i>P. phragmites</i> , but the teleutospores differ in size and shape, and are yellowish to pale yellowish brown.
1498	...	...	...	V.	...	...	B.	Different species of <i>Viola</i>	Cluster cups on all green parts (<i>Æcidium Violæ</i>). Uredospore pustules brown. Teleutospore pustules black.
1499	...	...	...	...	...	Q.	...	Leaves of <i>Wurmbea dioica</i>	Pustules elongated, blistered, dark brown. Uredospores brown. Teleutospores darker.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
149. PHRAGMIDIUM.—Link, Sp. II. 84 (1824).—					
1500	1771	VII. 2621	<i>P. Barnardi</i> ...	Plow. and Wint., Rev. Myc. 208 (1886) ...	Barnard's phragmidium ...
1501	1773	„ 2616	<i>P. Potentillæ</i> ...	Karst., Fung. Fenn. 94 (1887) ...	Potentilla phragmidium ...
1502	1772	„ 2622	<i>P. subcorticium</i> ...	Winter, Die Pilze. I. 228 (1884) ...	Sub-cortical phragmidium ...
150. HAMASPORA.—Körn.					
1503	1774	VII. 2630	<i>H. longissima</i> ...	Körn., Hedw. XVI. 23 (1877) ...	Very long hamaspora ...
151. ÆCIDIUM.—Pers., in Gmel.					
1504	1780	VII. 2857	<i>A. Apocyni</i> ...	Schwein, Syn. Car. 68 (1822) ...	Apocynum æcidium ...
1505	1776	„ 2719	<i>A. Barbareæ</i> ...	D. C., Fl. Fr. II. 244 (1815) ...	Barbarea æcidium ...
1506	1779	„ 2815	<i>A. Compositarum</i> ...	Mart., Erl. 314 (1817) ...	Composite æcidium ...
1507	1785	„ 2970	<i>A. cystoseiroides</i> ...	Berk., Fl. Tasm. 270 (1860) ...	Cystoseira-like æcidium ...
1508	1786	„ 2100	<i>A. Goodeniacearum</i> ...	Berk., Linn. Journ. XIII. 173 (1873) ...	Goodenia æcidium ...
1509	1781	„ 2864	<i>A. Nymphoidis</i> ...	D. C., Fl. Fr. II. 597 (1815) ...	Nymphoides æcidium ...
1510	1782	„ 2879	<i>A. Plantaginis</i> ...	Ces., Erb. Critt. Ital. 247 (1878) ...	Plantain æcidium ...
1511	1775	„ 2707	<i>A. Ranunculacearum</i> ...	D. C., Fl. Fr. V. 97 (1805) ...	Buttercup æcidium ...
1512	1790	„ 2313	<i>A. Senecionis</i> ...	Desm., Ann. Sci. Nat. 243 (1836) ...	Groundsel æcidium ...
1513	1778	„ 2770	<i>A. soleniforme</i> ...	Berk., Fl. Tasm. II. 270 (1860) ...	Solenia-like æcidium ...
1514	1789	„ 2196	<i>A. Urticæ</i> ...	Schum., Fl. Saccl. II. 223 (1801) ...	Nettle æcidium ...
1515	1783	„ 2887	<i>A. Veronicæ</i> ...	Berk., Grev. XI. 97 (1883) ...	Speedwell æcidium ...
152. RÆSTELIA.—Rebent.,					
1516	1791	VII. 2974	<i>R. polita</i> ...	Berk., Linn. Journ. XIII. 174 (1873) ...	Polished ræstelia ...
153. UREDO.—Pers.,					
1517	1792	VII. 2999	<i>U. angiosperma</i> ...	Thuem., Myc. Austr. IV. 95 (1880) ...	Angiospermons uredo ...
1518	1801	„	<i>U. armillata</i> ...	Ludw., Bot. Centr. 6 (1890) ...	Collared uredo ...
1519	1799	VII. 2210	<i>U. Cichoracearum</i> ...	D. C., Fl. Fr. II. 229 (1815) ...	Chicory uredo... ...
1520	1800	„ 3139	<i>U. Clematidis</i> ...	Berk., Hook., Journ. VI. 205 (1854) ...	Clematis uredo ...
1521	1798	„	<i>U. leguminum</i> ...	Desm., Ann. Sci. Nat. X. 310 (1838) ...	Legume uredo ...
1522	1793	„	<i>U. notabilis</i> ...	Ludw., Bot. Centr. 5 (1890) ...	Notable uredo ...
1523	...	„	<i>U. pallidula</i> ...	Cooke and Mass., Grev. XXII. 37 (1893) ...	Pallid uredo... ...
1524	1795	VII. 3101	<i>U. Rhagodiae</i> ...	Cooke and Mass., Grev. XV. 99 (1887) ...	Rhagodia uredo ...
1525	1796	„ 3111	<i>U. Spyridii</i> ...	Cooke and Mass., Grev. XV. 99 (1887) ...	Spyridium uredo ...
1526	1794	„ 3000	<i>U. Wurmbææ=U. Anguillaræ</i> ...	Cooke, Grev. XIV. 11 (1885) ...	Wurmbæa nredo ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Uredo, Cæoma, Puccinia, Lycoperdon, Hamaspora.									
1500	...	S.A.	...	V.	...	...	...	Leaves of <i>Rubus parvifolius</i>	Uredospore pustules scattered on under surface. Teleutospore pustules scattered or gregarious.
1501	...	...	...	V.	...	...	B.	Leaves of <i>Aciena Sanguisorba</i>	Spots roundish, orange yellow. Uredospores pustules orange red. Teleutospore pustules black.
1502	...	...	...	V.	...	...	B.	Rose leaves	Uredo-spore pustules on under surface, scattered or crowded, yellow. Teleutospore pustules black.

Hedw. 22 (1877).—Phragmidium, Uredo.

1503	...	...	...	...	...	Q.	...	Leaves of <i>Rubus Moluccanus</i>	Uredospores pustules on under surface, scattered or gregarious, bright orange. Teleutospore pustules gregarious, pale ochre.
------	-----	-----	-----	-----	-----	----	-----	-----------------------------------	--

Syst. 1472 (1791).—Cæoma, Uredo, Puccinia, Trichobasis.

1504	...	...	...	...	...	Q.	...	Leaves of <i>Tabernaemontana orientalis</i>	Spots thin, circular, large, orange, pale beneath.
1505	...	...	...	...	N.S.W.	...	B.	Crucifers...	Cluster cups on both surfaces on ruddy spots in irregular clusters, large, distorting leaf.
1506	...	...	...	V.	N.S.W.	Q.	B.	Leaves of <i>Senecio Vellegioides</i> and other <i>Compositæ</i>	Spots purplish, nearly round and run together. Cluster cups crowded on the spots in circular patches. Probably stage of <i>Puccinia Hieracii</i> and others.
1507	...	...	T.	...	...	...	...	<i>Opereularia</i> ...	Pustulate, deforming the leaves. Cluster cups immersed.
1508	...	S.A.	...	V.	N.S.W.	Q.	...	Leaves of <i>Selliera</i> , <i>Goodenia</i> , and <i>Scævola</i>	Spots circular, brown beneath. Cluster cups scattered. May belong to <i>Puccinia Saccardi</i> or <i>Uromyces puccinioides</i> .
1509	...	...	...	...	...	Q.	...	Leaves of <i>Limnanthemum indicum</i>	Cluster cups gregarious, disposed without order on rounded spots or in concentric zones.
1510	...	...	...	V.	N.S.W.	...	...	Leaves of <i>Plantago</i> ...	Spots mostly small, sometimes broad. Cluster cups loosely scattered on both surfaces.
1511	...	...	T.	V.	...	...	B.	Leaves of <i>Ranunculus rivularis</i> and <i>R. inundatus</i>	Cluster cups on under surface, in circular or elongated clusters, cup shaped.
1512	...	...	...	V.	N.S.W.	...	...	<i>Senecio</i> ...	Cluster cups on brown spots often bordered with black, arranged in clusters without definite order.
1513	...	...	T.	...	...	...	...	<i>Goodia latifolia</i> ...	Spots circular, brown. Spores orange.
1514	...	...	...	V.	...	...	B.	Nettles, &c.	Referred to <i>Puccinia caricis</i> . Cluster cups arranged in single or double series on yellow or red spots, on leaves, or stems.
1515	...	...	...	V.	...	...	...	<i>Veronica</i> ...	Scattered half-immersed cluster cups with margins fringed.

Fl. Neom. 330 (1804).

1516	...	...	...	V.	...	Q.	...	Branches of <i>Muehlenbeckia Cunninghamii</i> and <i>Jacksonia scoparia</i>	Ochrey, cylindrical, polished, delicately downy.
------	-----	-----	-----	----	-----	----	-----	---	--

in Usteri. n. Ann. IX. 16 (1795).

1517	W.A.	...	...	...	...	...	...	Leaves of <i>Hakea</i> ...	Pustules on both surfaces, large, commonly disposed about a circle, powdery, brown.
1518	...	...	...	V.	...	...	...	<i>Juncus pallidus</i> ...	Pustules red brown, run together, surrounded by ruptured epidermis.
1519	...	...	...	V.	...	Q.	B.	<i>Bidens pilosa</i> ...	Referred to <i>Puccinia Hieracii</i> . Spots very minute. Pustules on both surfaces, scattered, small, circular.
1520	...	...	...	V.	...	Q.	...	Leaves of <i>Clematis aristata</i> and <i>C. microphylla</i>	Pustules on under surface, solitary or gregarious, pale yellow, more or less rounded, flattened.
1521	...	...	...	...	...	Q.	...	Pods of <i>Acacia</i> ...	Pustules rounded, solitary, rather large, girt by ruptured epidermis.
1522	...	S.A.	...	V.	...	...	...	Phyllodes of <i>Acacia notabilis</i>	Pustules large, red brown, seated on distorted inflated tubercle.
1523	...	...	...	...	...	Q.	...	Leaves, twigs, and legumes of <i>Cassia</i>	Pustules on both surfaces, pallid, convex, gregarious, splitting irregularly, and then girt by ruptured epidermis.
1524	...	...	...	V.	...	...	...	Leaves of <i>Rhagodia Billardieri</i>	Pustules on under surface, scattered, covered a long time, at length torn, brown.
1525	...	...	...	V.	...	...	...	Leaves of <i>Spyridium parvifolium</i>	Pustules on under surface, scattered, yellowish, powdery.
1526	...	...	...	...	N.S.W.	...	...	Leaves of <i>Wurmbsea dioica</i>	Pustules on both surfaces, gregarious, blistered, long covered by epidermis.

GENERAL CLASSIFICATION OF PYRENOMYCETES.

GROUP IV.—PYRENOMYCETES, FRIES.

ARRANGEMENT OF ORDERS (18).

14. HYPOCREACEÆ—Simple or compound. Receptacles rather fleshy or waxy, bright coloured, never carbonaceous.
15. XYLARIACEÆ—Stroma erect, compound. Receptacles carbonaceous.
16. DOTHIIDEACEÆ—Composite, leathery or carbonaceous, blackish.
17. MELOGRAMMACEÆ—Receptacles formed from the stroma, or confluent with it.
18. DIATRYPACEÆ—Receptacles immersed in a heterogeneous stroma.
19. VALSACEÆ—Receptacles distinct, circinating, or in a single row.
20. EUTYPACEÆ—Receptacles immersed in stroma, densely gregarious for the most part.
21. CUCURBITARIACEÆ—Receptacles tufted or gregarious; eruptent, carbonaceous.
22. SUPERFICIALES—Receptacles distinct from each other, superficial or nearly so.
23. PERTUSACEÆ—Receptacles emergent, smooth, flattened at base.
24. LOPHIOSTOMACEÆ—Receptacles nearly superficial, opening compressed.
25. CERATOSTOMACEÆ—Receptacles for the most part immersed or sometimes nearly superficial.
26. OBTECTACEÆ—Receptacles innate in bark, and covered by cuticle.
27. CAULICOLACEÆ—Immersed, innate, observed mostly on dead stems of herbaceous plants.
28. FOLICOLACEÆ—Receptacles minute, membranous, innate, growing mostly on leaves.
29. MICROTHYRIACEÆ—Simple, receptacles nearly superficial, membranous or carbonaceous.
30. PERISPORIACEÆ—Receptacles membranous, leathery or somewhat carbonaceous, wholly closed.
31. HYSTERIACEÆ—Receptacles more or less elongated, leathery or somewhat carbonaceous.

ORDER XIV.—HYPOCREACEÆ, DE NOT.

ARRANGEMENT OF GENERA (15).

Sub-order 1. Hypocreoidæ, Cooke—Composite forms.

Genera (6)—

154. Claviceps, Tul.	156. Epichloë, Fries.	158. Hypocrella, Sacc.
155. Cordyceps, Fries.	157. Hypocrea, Fries.	159. Polystigma, Pers.

Sub-order 2. Nectriæ, Cooke—Simple or tufted forms.

Genera (9)—

160. Sphaerostilbe, Tul.	163. Hypomyces, Fries.	166. Gibberella, Sacc.
161. Nectria, Fries.	164. Dialonectria, Sacc.	167. Lisiella, Cooke.
162. Calonectria, De Not.	165. Ophionectria, Sacc.	168. Melanospora, Corda.

ORDER XV.—XYLARIACEÆ, COOKE.

Genera (7)—

169. Xylaria, Hill.	172. Ustilina, Tul.	174. Daldinia, De Not. and Ces.
170. Poronia, Willd.	173. Nummularia, Tul.	175. Hypoxylon, Bull.
171. Kretzschmaria, Fries.		

ORDER XVI.—DOTHIIDEACEÆ, NITS. AND FCKL.

ARRANGEMENT OF GENERA (7).

Sub-order 1. Dothideoidæ.

Genera (5)—

176. Phyllachora, Nits.	178. Montagnella, Speg.	180. Darwiniella, Speg.
177. Dothidella, Speg.	179. Bagnisiella, Speg.	

Sub-order 2. Rhytismoidæ.

Genus (1)—

181. Rhytisma, Fries.

Sub-order 3. Stigmatoidæ.

Genus (1)—

182. Trabutia, Sacc. and Roum.

ORDER XVII.—MELOGRAMMACEÆ, NITS.

Genera (4)—

183. Sarcosydon, Cooke.	185. Botryosphaeria, De Not.	186. Melogramma, Tul.
184. Gibellia, Sacc.		

ORDER XVIII.—DIATRYPACEÆ, FRIES.

Genera (2)—

187. *Diatrype*, Fries.| 188. *Cœlosphæria*, Sacc.

ORDER XIX.—VALSACEÆ, FRIES.

Genera (2)—

189. *Valsa*, Fries.| 190. *Eutypella*, Nits.

ORDER XX.—EUTYPACEÆ, COOKE.

Genera (3)—

191. *Cryptovalsa*, Ces. and De Not.| 192. *Cryptosphærella*, Sacc.| 193. *Eutypa*, Tul.

ORDER XXI.—CUCURBITARIACEÆ, COOKE.

Genus (1)—

194. *Gibberidea*, Fekl.

ORDER XXII.—SUPERFICIALES, FRIES.

Genera (7)—

195. *Byssosphæria*, Cooke.196. *Lasiosphæria*, Ces. and De Not.| 197. *Pleosphæria*, Speg.| 198. *Venturia*, De Not. and Ces.| 199. *Chaetomium*, Kunze.200. *Rosellinia*, De Not.201. *Trematosphæria*, Fekl.

ORDER XXIII.—PERTUSACEÆ, FRIES.

Genus (1)—

202. *Conisphæria*, Cooke.

ORDER XXIV.—LOPHIOSTOMACEÆ, SACC.

Genus (1)—

203. *Lophiostoma*, Ces. and De Not.

ORDER XXV.—CERATOSTOMACEÆ, FRIES.

Genus (1)—

204. *Rhamphoria*, Niessl.

ORDER XXVI.—OBTECTACEÆ, FRIES.

Genera (2)—

205. *Massariella*, Speg.| 206. *Didymosphæria*, Fekl.

ORDER XXVII.—CAULICOLACEÆ, FRIES.

Genera (4)—

207. *Physalospora*, Niessl.208. *Didymella*, Sacc.| 209. *Anthostomella*, Sacc.| 210. *Pleospora*, Rabh.

ORDER XXVIII.—FOLLICOLACEÆ, FRIES.

Genera (3)—

211. *Laetadiala*, Auersw.| 212. *Sphærella*, Ces. and De Not.| 213. *Sphaerulina*, Sacc.

ORDER XXIX.—MICROTHYRIACEÆ, SACC.

Genera (2)—

214. *Microthyrium*, Desm.| 215. *Micropeltis*, Mont.

ORDER XXX.—PERISPORIACEÆ, FRIES.

Genera (13)—

216. *Podosphæra*, Kunze.217. *Sphaerothera*, Lev.218. *Erysiphe*, Hedw.219. *Eurotium*, Link.220. *Asterina*, Lev.| 221. *Asterella*, Sacc.| 222. *Dimerosporium*, Fekl.| 223. *Parodiella*, Speg.| 224. *Meliola*, Fries.225. *Zukalia*, Sacc.226. *Asteridium*, Sacc.227. *Capnodium*, Mont.228. *Antennaria*, Link.

ORDER XXXI.—HYSTERIACEÆ, CORDA.

Genera (8)—

229. *Aulographum*, Lib.230. *Glonium*, Muhl.231. *Lembosia*, Lev.| 232. *Hysterium*, Tode.| 233. *Trybliidiella*, Sacc.| 234. *Rhytidhysterium*, Speg.235. *Platycheilus*, Cooke.236. *Hysterographium*, Corda.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
GROUP IV.—PYRENOMYCETES.—					
ORDER XIV.—HYPOCREACEÆ,					
154. CLAVICEPS.—Tul.,					
1527	...	II. 5005	<i>C. purpurea</i>	<i>Tul.</i> , Ann. Sci. Nat. XX. (1853)	Purple claviceps (Ergot)
155. CORDYCEPS.—Fries,					
1528	1488	II. 5012	<i>C. entomorrhiza</i>	<i>Fries</i> , S.V.S. 381 (1849)	Insect-root cordyceps
1528A	1488	...	<i>C. entomorrhiza</i> , var. <i>Menesteridis</i>	<i>Cooke</i> , Handb. Aust. Fung. 277 (1892)	Menesteridis cordyceps
1529	1486	II. 5030	<i>C. Gunnii</i>	<i>Berk.</i> , Hook., Lond. Journ. VII. 577 (1848)	Gunn's cordyceps
1530	1487	IX. 4013	<i>C. Hawkesii</i>	<i>Gray</i> , Not. Insects (1858)	Hawkes' cordyceps
1531	...	II. 5038	<i>C. ophioglossoides</i>	<i>Link</i> , Handb. III. 347 (1833)	Ophioglossum-like cordyceps
1532	...	„ 5041	<i>C. Taylori</i>	<i>Sacc.</i> , Mich. I. 320 (1878)	Taylor's cordyceps
156. EPICHLÆ.—Fries,					
1533	1489	II. 5059	<i>E. cinerea</i>	<i>Berk. and Br.</i> , Linn. Journ. XIV. 111 (1875)	Ashy epichlæ
157. HYPOCREA.—Fries,					
1534	1490	II. 4897	<i>H. cerebriformis</i>	<i>Berk.</i> , Linn. Journ. XIII. 179 (1873)	Brain-like hypocreæ
1535	1492	„ 4875	<i>H. citrinea</i>	<i>Fries</i> , S.V.S. 185 (1849)	Lemon-yellow hypocreæ
1536	1493	„ 4834	<i>H. rufa</i>	<i>Fries</i> , S.V.S. 383 (1849)	Reddish hypocreæ
1537	1491	„ 4898	<i>H. semiorbis</i>	<i>Berk.</i> , Fl. Tasm. II. 278 (1860)	Semicircular hypocreæ
158. HYPOCRELLA.—Sacc.,					
1538	1495	...	<i>H. axillaris</i>	<i>Cooke</i> , Grev. XX. 4 (1891)	Axillary hypocrella
1539	1494	II. 5061	<i>H. discoidea</i>	<i>Sacc.</i> , Mich. I. 322 (1878)	Discoid hypocrella
159. POLYSTIGMA.—D. C.,					
1540	1496	IX. 3802	<i>P. australiens</i>	<i>Sacc.</i> , Bull., Soc. Myc. Fr. V. 119 (1889)	Australian polystigma
160. SPHEROSTILBE.—Tul.,					
1541	1497	II. 4817	<i>S. cinnabarina</i>	<i>Tul.</i> , Carp. III. 103 (1865)	Cinnabar sphaerostilbe
1542	1500	„ 4825	<i>S. dubia</i>	<i>Berk.</i> , Linn. Journ. XVIII. 389 (1881)	Doubtful sphaerostilbe
1543	1498	„ 4820	<i>S. hypocreoides</i>	<i>Kalch. and Cooke</i> , Grev. IX. 26 (1880)	Hypocreæ-like sphaerostilbe
1544	1499	IX. 3912	<i>S. microspora</i>	<i>Cooke and Mass.</i> , Grev. XVI. 4 (1887)	Small-spored sphaerostilbe

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
FRIES, S.M. I. 51 (1821).									
DE NOT. OSS. PIR. GIORN. BOT. I. (1844).									
Ann. Sci. Nat. Ser. III., XX. 43 (1853).—Sphaeria.									
1527	...	...	...	V.	...	...	B.	On <i>Lolium perenne</i> , <i>L. temulentum</i> , <i>Triticum sativum</i> , &c.	Stroma stalked, erect, arising from a sclerotium, club headed.
S.M. II. 324 (1821).—Sphaeria.									
1528	...	...	...	V.	...	...	B.	Insect larvæ (<i>Tinea</i> , &c.)	Fleshy. Club somewhat globose, brown. Stem thin, very long.
1528A	...	...	...	V.	...	...	...	Larvæ of <i>Menes-terdis</i> , &c.	Club elliptic, reddish, at first powdery. Stem thickened upwards.
1529	...	...	T.	V.	N.S.W.	...	...	Larvæ of some <i>Cossus</i> or <i>Hepialus</i>	Fleshy. Club cylindrical, yellow, blackening above. Stem elongated, white.
1530	...	...	T	...	...	...	...	Larvæ of insects	Cylindrical, tapering and truncate at apex. Stem flexuous, sometimes forked with three or four clubs.
1531	...	...	...	...	...	...	B.	...	Solitary, rarely tufted, simple, rarely branched, fleshy, yellow within. Club oblong. Stem olive, then blackening.
1532	...	...	...	V.	N.S.W.	...	...	Larvæ of insects	Stems tufted, running together in net-like manner. Clubs reddish yellow, delicately velvety.
S.V.S. 381 (1849).									
1533	...	...	...	...	...	Q.	...	Various grasses, preferably <i>Sporobolus</i>	Encircling stems of grasses, dark ashy, and dotted with the darker openings of immersed receptacles.
S.V.S. 383 (1849).									
1534	...	S.A.	...	...	...	...	...	Trunks	Cushion shaped, wrinkled, lobed, fawn, substance thick.
1535	...	...	T.	...	...	...	B.	Soil, mosses, rotting leaves, &c.	Fleshy, spread out, lemon yellow, prominent openings of receptacles brownish.
1536	...	...	T.	...	...	...	B.	Wood or bark	Gregarious, hemispherical when moist, collapsing when dry, wrinkled, reddish, soft and fleshy.
1537	...	...	T.	...	...	...	...	Bark and wood	Hemispherical, rather fleshy, ochrey, darker than the bark. Receptacles immersed, openings minute.
Mich. I. 322 (1878).—Hypocrea.									
1538	...	...	...	...	...	Q.	...	Grasses (<i>Eragrostis stricta</i>)	Inversely club shaped, seated in the upper axils, black, opaque, minutely granular with the openings of the receptacles.
1539	...	...	...	...	...	Q.	...	Leaves	Circular, separating from matrix, scarlet. Receptacles rather prominent.
Fl. Fr. V. 164 (1815).									
1540	...	...	...	V.	...	...	...	Leaves, rarely stems, of <i>Leguminosæ</i>	Immersed, rather swollen, covering half or entire leaf, dull rosy, rather fleshy.
Carp. III. 103 (1865).									
1541	...	...	...	...	N.S.W	Q.	...	Bark	Receptacles at base of conidia-bearing layers, small, sessile, globose, smooth, orange red.
1542	...	...	...	...	...	Q.	...	Bark of <i>Ægiceras</i>	Only Stilbum form known, bearing conidia.
1543	...	...	...	...	...	Q.	...	Bark	Pale rose, convex. Receptacles as in <i>Hypocrea</i> , associated with club-shaped conidia bearers.
1544	...	...	...	V.	...	...	...	Bark	Receptacles associated with conidia bearers, minute, scattered, ovate, orange.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
161. NECTRIA.—Fries,					
1545	1502	II. 4670	<i>N. coccinea</i> ...	<i>Fries</i> , S.V.S. 368 (1849) ...	Scarlet nectria ...
1546	..	IX. 3858	<i>N. ferruginea</i> ...	<i>Cooke</i> , Grev. XIII. 8 (1884) ...	Rusty nectria... ..
1547	1501	II. 4561	<i>N. fusarioides</i> ...	<i>Berk.</i> , Fl. Tasm. II. 279 (1860) ...	Fusarium-like nectria ...
1548	1503	„ 4705	<i>N. tasmanica</i> ...	<i>Berk.</i> , Fl. Tasm. II. 279 (1860) ..	Tasmanian nectria ...
1549	1504	„ 4678	<i>N. zealandica</i> ...	<i>Cooke</i> , Grev. VIII. 65 (1879) ...	New Zealand nectria ...
162. CALONECTRIA.—De Not.,					
1550	...	II. 6174	<i>C. otagensis</i> ...	<i>Sacc.</i> Syll. II. LXVIII. (1883) ...	Otago calonectria ...
163. HYPOMYCES.—Fries, Pl. Homon. 105 (1825).—					
1551	1508	II. 4622	<i>H. aurantius</i> ...	<i>Fekl.</i> , Symb. Myc. 183 (1875) ...	Golden hypomyces ...
1552	1505	„ 4614	<i>H. chrysospermus</i> ...	<i>Tul.</i> , Sel. Fung. Carp. III. 51 (1865) ...	Golden-seeded hypomyces ...
1553	1509	...	<i>H. membranaceus</i> ...	<i>Cooke</i> , Handb. Austr. Fung. 281 (1892) ...	Membranous hypomyces ...
1554	1506	II. 4617	<i>H. rosellus</i> ...	<i>Tul.</i> , Sel. Fung. Carp. III. 45 (1865) ...	Rosy-red hypomyces ...
1555	1507	„ 4643	<i>H. tomentosus</i> ...	<i>Fries</i> , in Grev. IV. 15 (1875) ...	Downy hypomyces ...
164. DIALONECTRIA.—Sacc.					
1556	1511	II. 4733	<i>D. quisquilaris</i> ...	<i>Cooke</i> , Handb. Austr. Fung. 282 (1882) ...	Rubbish-loving dialonectria ...
1557	1510	„ 4721	<i>D. sanguinea</i> ...	<i>Fries</i> , S.V.S. 388 (1849) ...	Blood-red dialonectria ...
1558	1512	„ 4742	<i>D. tephrothele</i> ...	<i>Berk.</i> , Fl. Tasm. II. 278 (1860) ...	Dark-nippled dialonectria ...
165. OPHIONECTRIA.—Sacc.,					
1559	1513	II. 5001	<i>O. agaricieola</i> ...	<i>Sacc.</i> Syll. II. 563 (1883) ...	Agaricus-growing ophiouectria ...
166. GIBBERELLA.—Sacc., Mich. I. 43 (1878).—					
1560	1514	II. 4977	<i>G. Saubinetii</i> ...	<i>Sacc.</i> , Mich. I. 513 (1878) ...	Saubinet's gibberella ...
167. LISIELLA.—Cooke,					
1561	1515	IX. 3804	<i>L. Passiflorae</i> ...	<i>Cooke</i> , Grev. XVI. 5 (1887) ...	Passion-flower lisiella ...
168. MELANOSPORA.—Corda,					
1562	1516	II. 4599	<i>M. capriua</i> ...	<i>Sacc.</i> Syll. II. 462 (1883) ...	Shaggy melanospora ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
S.V.S. 387 (1849).—Sphaeria.									
1545	...	...	T.	V.	N.S.W.	Q.	B.	Bark and dead branches	Receptacles in tufts, on a convex yellowish layer, ovoid, bright red, sometimes ochrey red.
1546	...	...	...	V.	...	...	...	Living leaves, bracts, &c. of <i>Styphelia</i>	Bursting through, tufted. Receptacles waxy, almost globose, dark brown, at length naked.
1547	...	...	T.	...	...	...	...	Dead bark	Pale crimson. Receptacles ovate to pap-like, with a bloom half immersed in umber-coloured layer.
1548	...	...	T.	...	...	...	...	Dead bark	Tufted, red. Receptacles ovate, with pap-like openings, often arising from circular disc.
1549	...	...	...	V.	...	...	...	Bark	Tufted, brick red, bursting through, pustules convex. Receptacles almost globose, pap-like, soon concave.
Comm. Critt. II. 477 (1867).—Nectria.									
1550	...	...	...	...	...	Q.	...	Twigs of <i>Capparis Mitchellii</i>	Receptacles densely tufted, pale-apricot colour, with openings a little deeper in colour, papillate.
Sphaeria, Hypoerea, Nectria.									
1551	...	...	...	...	...	Q.	B.	On <i>Polyporus</i> , &c.	Conidia-bearing mycelium white, then orange. Receptacles seated on fluffy ochrey base, often white at margin, golden yellow or orange.
1552	W.A.	...	...	V.	...	Q.	B.	On <i>Boleti</i> chiefly and <i>Polyporus</i>	Conidia-bearing mycelium penetrating matrix. Receptacles closely packed in a rough layer, pale yellow brown.
1553	...	...	...	...	...	Q.	...	On <i>Polyporus</i>	Forming at first fine filaments, gradually becoming membranous, tan coloured.
1554	W.A.	...	...	...	...	...	B.	On <i>Polyporus</i> , &c.	Receptacles gregarious, at first white, then bright-red mycelium.
1555	...	...	T.	...	...	...	...	On <i>Agarics</i>	Stratum white, delicate downy.
Syll. II. 490 (1883).—Nectria, Sphaeria.									
1556	...	...	...	V.	...	...	...	Bark, chips, &c.	Scattered, umber coloured or somewhat orange. Receptacles crowded here and there.
1557	W.A.	...	...	...	...	...	B.	Wood and bark	Receptacles scattered, egg shaped, openings pap-like, blood red.
1558	...	...	T.	...	...	...	...	<i>Hypoxylon</i>	Receptacles scattered, crimson, egg shaped, openings darker and pap-like.
Mich. I. 323 (1878).—Nectria.									
1559	...	...	T.	V.	...	...	...	Putrid <i>Agarics</i>	Vermilion. Receptacles egg shaped, with fibrous swollen texture.
Gibbera, Botryosphaeria.									
1560	...	...	T.	V.	...	...	B.	Herbaceous stems	Receptacles gregarious, growing together in tufts, somewhat membranous, warted, folded, blue.
Grev. XVI. 5 (1887).—Gibberella.									
1561	...	...	...	...	...	Q.	...	Stems of <i>Passiflora</i>	Receptacles bursting through, collected in small clusters, globose, substance bright blue.
Icon. Fung. I. 24 (1837).—Sphaeria.									
1562	...	...	T.	...	...	...	B.	Wood and chips	Receptacles globose, shaggy, white, openings turning blackish.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER XV.—XYLARIACEÆ,					
169. XYLARIA.—Hill. Hist. Plant. 62 (1773).—					
1563	1543	IX. 2175	<i>X. agariciformis</i> ...	Cooke and Mass., Grev. XVII. 81 (1889) ...	Agaricoid xylaria ...
1564	1532	I. 1178	<i>X. allantoidea</i> ...	Berk., Linn. Journ. X. 380 (1869) ...	Allantoid xylaria ...
1565	1535	„ 1219	<i>X. anisopleura</i> ...	Mont., Syll. 688 (1856) ...	Unequal-ribbed xylaria ...
1566	1545	„ 1241	<i>X. aphrodisiaca</i> ...	Welw. and Curr., Fung. Angol. 280 (1867) ...	Aphrodisiac xylaria ...
1567	1517	IX. 2137	<i>X. australis</i> ...	Cooke, Grev. XI. 84 (1883) ...	Southern xylaria ...
1568	1531	I. 1246	<i>X. castorea</i> ...	Berk., Fl. N. Zeal. II. 204 (1855) ...	Beaver xylaria ...
1569	1528	IX. 2157	<i>X. cerebriiformis</i> ...	Cooke, Grev. XI. 86 (1883) ...	Brain-like xylaria ...
1570	1541	„ 2173	<i>X. cinnabarina</i> ...	Cooke and Mass., XV. 101 (1887) ...	Vermilion xylaria ...
1571	1544	I. 1239	<i>X. corniformis</i> ...	Fries, S.V.S. 381 (1849) ...	Horn-shaped xylaria ...
1572	1538	II. 5963	<i>X. cretacea</i> ...	Berk. and Br., Linn. Trans. 405 (1879) ...	Chalky xylaria ...
1573	1529	IX. 2160	<i>X. cynoglossa</i> ...	Cooke, Grev. XII. 1 (1883) ...	Dog-tongue xylaria ...
1574	1534	I. 1223	<i>X. dealbata</i> ...	Berk. and Curt., Exot. Fung. 284 (1853) ...	Bleached xylaria ...
1575	1546	„ 1283	<i>X. digitata</i> ...	Grev., Fl. Ed. 356 (1824) ...	Digitate xylaria ...
1576	1540	IX. 2172	<i>X. elastica</i> ...	Cooke, Grev. XVI. 4 (1887) ...	Elastic xylaria ...
1577	1536	„ 2169	<i>X. ellipsozona</i> ...	Cooke and Mass., Grev. XVI. 33 (1887) ...	Elliptical-spored xylaria ...
1578	1533	I. 1185	<i>X. fistulosa</i> ...	Fries, Nov. Symb. 126 (1851) ...	Hollow xylaria ...
1579	1539	...	<i>X. gigas</i> ...	Cooke, Handb. Austr. Fung. 287 (1892) ...	Gigantic xylaria ...
1580	1526	I. 1188	<i>X. gracilis</i> ...	Sacc. Syll. I. 317 (1882) ...	Graceful xylaria ...
1581	1519	„ 1189	<i>X. grammica</i> ...	Mont., Syll. 680 (1856) ...	Lined xylaria ...
1582	...	„ 1303	<i>X. hippotrichoides</i> ...	Sacc. Syll. I. 344 (1882) ...	Horsehair-like xylaria ...
1583	1547	„ 1260	<i>X. hypoxylon</i> ...	Grev., Fl. Ed. 355 (1824) ...	Hypoxylon xylaria ... (Candle-snuff fungus)
1584	1518	„ 1228	<i>X. involuta</i> ...	Klotzsch, Linn. VII. (1832) ...	Involute xylaria ...
1585	1530	IX. 2171	<i>X. lobata</i> ...	Cooke, Grev. XI. 86 (1883) ...	Lobed xylaria ...
1586	1525	„ 2154	<i>X. ovispora</i> ...	Cooke and Mass., Grev. XV. 101 (1887) ...	Egg-spored xylaria ...
1587	1537	I. 1167	<i>X. phosphorea</i> ...	Berk., Linn. Journ. XIII. 177 (1873) ...	Phosphorous xylaria ...
1588	1527	„ 1150	<i>X. polymorpha</i> ...	Grev., Fl. Ed. 35 (1824) ...	Polymorphous xylaria ...
1588A	...	...	<i>X. polymorpha</i> , var. <i>pachystroma</i> ...	Sacc., Myc. Austr. 13 (1890) ...	Thick stroma xylaria ...
1589	...	...	<i>X. Readeri</i> ...	F. v. M., Grev. XXII. 17 (1893) ...	Reader's xylaria ...
1590	...	...	<i>X. rhizophila</i> ...	Cooke and Mass., XXII. 37 (1893) ...	Root-loving xylaria ...
1591	1520	I. 1234	<i>X. rhopaloides</i> ...	Mont., Ann. Sci. Nat. III. 99 (1855) ...	Club-like xylaria ...
1592	1522	„ 1200	<i>X. rhytidophloea</i> ...	Mont., Syll. 687 (1856) ...	Wrinkled xylaria ...
1593	1521	„ 1222	<i>X. Schweinitzii</i> ...	Berk. and Curt., Exot. Fung. 284 (1853) ...	Schweinitz's xylaria ...
1594	1524	„ 1288	<i>X. scopiformis</i> ...	Mont., Ann. Sci. Nat. XIII. 319 (1840) ...	Stalk-like xylaria ...
1595	1542	„ 1225	<i>X. tuberiformis</i> ...	Berk., Fl. N. Zeal. II. 204 (1855) ...	Tuber-like xylaria ...
1596	1523	„ 1209	<i>X. zealandica</i> ...	Cooke, Grev. VIII. 66 (1879) ...	New Zealand xylaria ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
COOKE, GREV. XIII. 9 (1884).									
Sphaeria, Clavaria, Hypoxylon, Rhizomorpha.									
1563	...	S.A.	...	..	...	...	...	Stumps ...	Half globose, glaucous, dotted with black openings. Stem equal, with barren black ring round it.
1564	W.A.	...	...	V.	...	Q	...	Stumps ...	Club shaped, encrusted with black, leathery. Stem very short. Receptacles minute, globose.
1565	...	...	...	...	...	Q.	...	Dead wood	Solitary, woody, very hard. Club inversely egg shaped as well as stem, scaly.
1566	...	...	...	...	...	Q	...	Rotten trunks	Tufted, smooth. Clubs somewhat cylindrical, brown to ashy, substance white.
1567	...	...	...	...	...	Q.	...	Wood ...	Club shaped, thickened upwards, brown. Stem elongated, blackening.
1568	...	...	T.	..	...	Q.	...	Rotten wood	Stem short, spongy and velvety at first, at length naked, black. Clubs obtuse, ovate, compressed.
1569	...	...	...	...	...	Q.	...	Wood ...	Large, corky, stalked, sooty brown. Stem woody, smooth, furrowed. Clubs somewhat elliptic, brain-like.
1570	...	...	...	...	...	Q.	...	Wood ...	Corky, somewhat globose, brown, vermilion under thin skin. Stem obsolete.
1571	...	...	T.	...	...	...	B.	Prostrate trunks ...	Cylindrical, thick, brown, then black, horn shaped.
1572	...	...	...	...	...	Q.	...	Trunks ...	Nearly globose, stalked, white, rather wrinkled, netted with thin brown lines.
1573	...	...	...	...	...	Q.	...	Wood ...	Umber, tongue shaped, shortly stalked, flesh white.
1574	...	...	...	...	...	Q.	...	Rotten trunks	Club elliptical, tapering downwards into short cylindrical stem, covered with whitened crust.
1575	...	...	...	V.	N.S.W.	...	B	Rotten wood	Erect, thick, brown, velvety, tapering towards apex, sometimes divided into forked branches; short stem.
1576	...	...	...	V.	...	Q.	...	Rotten wood	Corky to elastic, nearly globose, or hemispherical, sessile, becoming black.
1577	...	...	T.	...	...	...	...	Rotten wood	Club shaped, obtuse, black. Stem shortened, smooth. Receptacles immersed, not prominent.
1578	...	...	...	...	...	Q.	..	Trunks ...	Corky, simple, club shaped, with varnished crust, black, hollow, confluent with short stem.
1579	...	...	...	...	N.S.W.	Q.	...	Stumps, &c.	Large, ochrey, then brown. Clubs oval or irregular, wrinkled. Stem stout and irregular.
1580	...	...	...	...	...	Q.	...	Wood of <i>Acacia harpophylla</i>	Leathery. Stem smooth, forked, tufted. Clubs cylindrical, narrow, wrinkled, black.
1581	...	...	...	V.	...	Q.	...	Trunks ...	Large, corky, club shaped, rigid, sooty black, becoming whitish. Clubs with longitudinal lines running into one another.
1582	...	...	...	...	...	...	B.	...	Somewhat tufted, thread-like, ascending, black, branched.
1583	...	...	...	...	N.S.W.	Q.	B.	Stumps ...	Erect, compressed, and dilated, black, shaggy about base. Stem usually shorter than club.
1584	...	...	...	...	N.S.W.	Q.	...	Woods ...	Leathery, club shaped, ochrey to brownish yellow or fawn, tapering below into long slender stem.
1585	...	...	...	...	...	Q.	...	Wood ...	Large, corky, shell shaped, sessile, circumference lobed, brown, lobes rounded.
1586	...	...	...	...	...	Q.	...	Stumps ...	Leathery, black, stalked, erect, forked above, tapering downwards into smooth stem.
1587	...	...	...	V.	...	...	...	Trunks ...	Reddish brown, small. Stem short, cylindrical, streaked, expanding upwards into short club.
1588	...	...	...	...	N.S.W.	Q.	B.	Wood ...	Clubs in clusters, rarely solitary erect, thick, smooth, brown, then black, variously shaped.
1588A	...	...	...	...	...	Q.	...	Trunks ...	Stroma or receptacle-bearing layer very hard and thick above middle branching into finger-like processes.
1589	...	...	...	V.	...	...	...	Sandy desert	Black, globose or broadly elliptical, crowned by short spine, mealy, with white conidia. Stem erect.
1590	...	...	...	...	...	Q.	...	Roots of herbs and grasses	Stroma club shaped, divided nearly to base into two to six clubs. Clubs spoon shaped, flattened, bright brown.
1591	...	...	...	...	...	Q.	...	Putrid wood	Clubs cylindrical, obtuse, tapering downwards into short smooth stem.
1592	...	...	...	V.	...	Q.	...	Wood ...	Horny, compressed, obtuse or horn shaped, black. Stem very short and wrinkled in a netted manner.
1593	...	...	...	V.	N.S.W.	...	...	Rotten wood	Club elliptic, obtuse, corky, compact. Stem elongated, smooth, slightly cracked.
1594	...	...	...	...	...	Q.	...	Decaying fruit of <i>Flindersia australis</i>	Simple, slender. Clubs cylindrical, acute at apex, black. Stem as long as club, compressed.
1595	..	...	...	...	...	Q.	...	Rotten wood	Corky, almost globose, wrinkled, cap like, thick. Stem short or obsolete.
1596	...	...	...	...	...	Q.	...	Rotten wood	Simple, slender, stalked, black. Clubs cylindrical, wrinkled. Stem smooth, channelled.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
170. PORONIA.—Willd.,					
1597	1549	I. 1322	<i>P. edipus</i> ...	Mont., Syll. 209 (1856) ...	Swollen-stalked poronia ...
1598	1550	„ 1323	<i>P. pileiformis</i> ...	<i>Fries</i> , Nov. Symb. 129 (1851) ...	Cap-shaped poronia ...
1599	1548	„ 1321	<i>P. punctata</i> ...	<i>Fries</i> , S.V.S. 382 (1849) ...	Punctate poronia ...
171. KRETZSCHMARIA.—Fries,					
1600	1552	I. 1519	<i>K. angolensis</i> ...	<i>Sacc.</i> Syll. IX. 565 (1891) ...	Angola kretzschmaria ...
1601	1551	„ 1489	<i>K. cetrarioides</i> ...	<i>Sacc.</i> Syll. IX. 567 (1891) ...	Cetrarium-like kretzschmaria ...
172. USTULINA.—Tul., Sel. Fung.					
1602	1553	I. 1328	<i>U. vulgaris</i> ...	Tul., Sel. Fung. Carp. II. 23 (1863) ...	Common ustulina ...
173. NUMMULARIA.—Tul., Sel. Fung.					
1603	1558	IX. 2295	<i>N. australis</i> ...	Cooke, Grev. XII. 6 (1883) ...	Southern nummularia ..
1604	1554	„ 2292	<i>N. Baileyi</i> ...	Cooke, Grev. XII. 6 (1883) ...	Bailey's nummularia ...
1605	1556	I. 1524	<i>N. Bulliardii</i> ...	Tul., Sel. Fung. Carp. II. 43 (1863) ...	Bulliard's nummularia ...
1606	1557	„ 1105	<i>N. exutans</i> ...	Cooke, Handb. Austr. Fung. 291 (1892) ...	Shedding nummularia ...
1607	1555	„ 1528	<i>N. lutea</i> ...	<i>Nits.</i> Pyr. Germ. 59 (1867) ...	Yellow nummularia ...
1608	1559	„ 1112	<i>N. microplaea</i> ...	Cooke, Grev. XIII. 13 (1884) ...	Cake-like nummularia ...
1609	1560	IX. 2300	<i>N. pusilla</i> ...	Sacc., Hedw. (1889) ...	Small nummularia ...
174. DALDINIA.—De Not. and Ces.,					
1610	1561	I. 1515	<i>D. concentrica</i> ...	<i>Ces. and De Not.</i> , Schema Sfer. Ital. 198 (1870) ...	Concentric daldinia ...
1611	1562	„ 1516	<i>D. vernicosa</i> ...	<i>Ces. and De Not.</i> , Schema Sfer. Ital. 198 (1870) ...	Varnished daldinia ...
175. HYPOXYLON.—Bull.,					
1612	1573	I. 1384	<i>H. annulatum</i> ...	Mont., Syll. 213 (1856) ...	Annulate hypoxylon ...
1613	1580	„ 1449	<i>H. Archeri</i> ...	Berk., Fl. Tasm. II. 280 (1860) ...	Archer's hypoxylon ...
1614	1567	„ 1337	<i>H. argillaceum</i> ...	Berk., Outl. 387 (1860) ...	Clay-coloured hypoxylon ...
1615	...	„ 1433	<i>H. atro-purpureum</i> ...	Fries, S.V.S. 384 (1849) ...	Dark-purple hypoxylon ...
1616	1576	„ 1113	<i>H. capnodes</i> ...	Cooke, Grev. XI. 147 (1883) ...	Capnodium-like hypoxylon ...
1617	1566	„ 1333	<i>H. coccineum</i> ...	Bull., Champ. I. 174 (1798) ...	Brick-red hypoxylon ...
1618	1571	„ 1370	<i>H. coherens</i> ...	Fries, S.V.S. 384 (1849) ...	Cohering hypoxylon ...
1619	1581	IX. 2264	<i>H. ellipticum</i> ...	Cooke and Mass., Grev. XVII. 70 (1889) ...	Elliptic hypoxylon ...
1620	...	„ 2230	<i>H. flavo-fusceum</i> ...	Berk. and Br., Linn. Traus. II. 222 (1887) ...	Yellow-brown hypoxylon ...
1621	1569	I. 1368	<i>H. fuscum</i> ...	Fries, S.V.S. 384 (1849) ...	Brown hypoxylon ...
1622	1575	IX. 2254	<i>H. hæmatites</i> ...	Lev., Grev. XI. 133 (1883) ...	Orange-red hypoxylon ...
1623	1572	I. 1435	<i>H. hæmato-stroma</i> ...	Mont., Syll. 737 (1856) ...	Blood-red stroma hypoxylon ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Fl. Berol. 400 (1787).—Sphæria, Peziza.									
1597	...	...	...	V.	N.S.W.	Q.	...	Dung ...	Erect, externally blackish, simple or branched. Stem long, clubbed at base, expanded into cup at apex.
1598	...	...	...	...	...	Q.	...	Rotten wood ...	Branched, nearly even, branches ending in pileiform cups, blackish.
1599	W.A.	S.A.	T.	V.	...	...	B.	Dung ...	Erect, simple, at first club shaped, soon cup shaped, tapering into long black downy stem, disc white, punctate with black openings.
S.V.S. 409 (1849).—Rhopalopsis, Daldinia, Hypoxylon.									
1600	...	...	...	...	...	Q.	...	Rotting bark ...	Club shaped, black, bright, and shining. Receptacles thickly crowded, openings minute.
1601	...	...	...	...	...	Q.	...	Trunks ...	Resembling in habit and mode of growth the Lichen <i>Cetraria tristis</i> , fringed, fringes passing into receptacles at apex.
Carp. II. 23 (1861).—Sphæria, Hypoxylon.									
1602	...	...	...	...	...	Q.	B.	Branches and logs	Spread out, large, thick, wavy, at length quite black, carbonaceous, hollow within.
Carp. II. 42 (1863).—Hypoxylon, Sphæria, Anthostoma, Diatrype.									
1603	...	...	...	...	N.S.W.	...	...	Branches ...	Developed within bark, then bursting through, unpolished, black. Receptacles nearly globose, small.
1604	...	...	...	...	...	Q.	...	Wood ...	Bursting through, circular, cup shaped, disc rough with prominent openings of receptacles, which are immersed in centre.
1605	...	...	T.	...	...	...	B.	Branches ...	Bursting through, circular or oval, broadly expanded, black without and within.
1606	...	...	...	V.	...	...	...	Branches ...	Broadly expanded, marginate, black, bursting through, and throwing off epidermis.
1607	...	...	...	...	...	Q.	B.	Wood ...	Superficial on decorticated wood, circular, thick, cup shaped, surface brown, then blackish.
1608	...	...	...	...	...	Q.	...	Bark ...	Thin, circular, sparingly dotted with minute pap-like openings of receptacles, black.
1609	...	S.A.	...	...	...	...	...	Branches of <i>Bursaria spinosa</i>	Small, flattened, becoming black, shining. Receptacles crowded, oblong, openings point-like.
Schema Sfer. Ital. I. 197 (1870).—Hypoxylon, Sphæria.									
1610	W.A.	...	T.	V.	N.S.W.	Q.	B.	Trunks ...	Spherical or hemispherical, zoned internally with concentric layers, black or brown.
1611	...	...	...	V.	...	...	...	Trunks ...	Large, tapering below into thick stem, surface black, varnished. Receptacles in many rows, black.
Champ. I. 168 (1798).—Sphæria, Diatrype, Anthostoma, Nummularia.									
1612	...	...	T.	...	...	...	...	Bark and wood ...	Hemispherical, confluent, blackening. Receptacles nearly globose, opening in centre of disc, with rather prominent annulate margin.
1613	...	...	T.	...	...	...	...	Rotten wood ...	Quite black, spread out. Receptacles nearly globose, wrinkled, opening in centre of depressed disc.
1614	...	...	T.	...	...	...	B.	Trunks ...	Bursting through, somewhat globose, clay coloured, turning black within. Receptacles small, ovate, crowded.
1615	...	...	...	...	...	...	B.	Wood ...	Stroma in wood turning black and widely spread out, circumference variously and often interrupted.
1616	W.A.	...	...	...	...	Q.	...	Branches ...	Spread out, greyish black, dotted with the prominent openings of the receptacles.
1617	...	...	T.	V.	...	...	B.	Branches ...	Bursting through, nearly globose, violet brown or fawn, then brick red. Receptacles minute, ovate, crowded.
1618	...	...	T.	...	...	...	B.	Branches ...	Bursting through, nearly globose or flattened, thick, dirty brown, then black. Receptacles large, globose.
1619	...	...	...	...	...	Q.	...	Decorticated wood	Parallel, elliptic, black, openings of receptacles minute, crowded, dotted.
1620	...	...	...	...	...	Q.	...	Roots and stumps of grasses	Convex, irregularly lobed, yellow brown, mealy; mouths black, prominently punctate.
1621	...	...	...	...	...	Q.	B.	Bark ...	Bursting through, spot-like, hemispherical, purple brown then black. Receptacles globose, crowded.
1622	...	...	...	...	...	Q.	...	Wood ...	Expanded, crustaceous, wrinkled, bright orange red, at length rusty. Receptacles thickly crowded.
1623	...	...	...	...	...	Q.	...	Bark ...	Irregularly spread out, confluent, purplish black. Receptacles immersed rather prominent, layer blood red.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
175. HYPOXYLON.—Bull,					
1623A	...	IX. 2232	II. hæmato-stroma, var. hæmatozonum	Sacc., Pug. Austr. 13 (1890) ...	Blood-red zoned hypoxylon ...
1624	...	„ 2227	II. hians	Berk. and Cooke, Grev. XI. 129 (1883) ...	Gaping hypoxylon ...
1625	1574	I. 1414	II. marginatum	Berk., Linn. Journ. X. 385 (1869) ...	Marginate hypoxylon ...
1626	1570	„ 1376	II. multiforme	Fries, S.V.S. 384 (1849) ...	Multiform hypoxylon ..
1627	1577	„ 1456	II. oodes	Berk. and Br., Linn. Journ. 122 (1875) ...	Egg-like hypoxylon ...
1628	1564	„ 1535	II. placentaforme	Berk. and Curt., Linn. Journ. 383 (1869) ...	Placenta-shaped hypoxylon
1629	1578	„ 1534	II. punctulatum	Berk. and Rav., Grev. IV. 94 (1875) ...	Punctulate hypoxylon ...
1630	1568	„ 1344	II. rutilum	Tul., Sel. Fung. Carp. II. 38 (1863) ...	Reddish hypoxylon ...
1631	1563	„ 1341	II. sclerophæum	Berk. and Curt., Linn. Journ. XIII. 177 (1873) ...	Hard-dusky hypoxylon ...
1632	1579	„ 1448	II. serpens	Fries, S.V.S. 384 (1849) ...	Creeping hypoxylon ...
1633	1565	IX. 2212	II. stratosum	Sacc., Pug. Austr. 13 (1890) ...	Stratose hypoxylon ...

ORDER XVI.—DOTHIDEACEÆ.

176. PHYLLACHORA.—Nits,					
1634	1585	IX. 4083	P. Alpinie	Sacc. and Berl., Misc. Myc. II. 5 (1884) ...	Alpinia phyllachora ..
1635	1586	„ 4091	P. anceps	Sacc., Hedw. 156 (1890) ...	Two-sided phyllachora ...
1636	1588	„ 4089	P. Fimbristylis	Sacc. Syll. IX. 1025 (1891) ...	Fimbristylis phyllachora ...
1637	1582	II. 5132	P. granulinis	Fekl., Symb. Myc. 216 (1869) ...	Grass phyllachora ...
1638	1589	„ 5144	P. junci...	Fekl., Symb. Myc. 216 (1869) ...	Rush phyllachora ...
1639	1587	...	P. maculata	Cooke, Grev. XX. 4 (1891) ...	Spotted phyllachora ...
1640	...	IX. 4088	P. nervisequia	Winter, Hedw. 9 (1885) ...	Nerve-following phyllachora ...
177. DOTHIDELLA.—Speg.,					
1641	1583	II. 5093	P. rhytismoides	Sacc. Syll. II. 594 (1883) ...	Rhytisma-like phyllachora ...
1642	1584	„ 5184	P. Trifolii	Fekl., Symb. Myc. 218 (1869) ...	Clover phyllachora ...
178. MONTAGNELLA.—Speg.,					
1643	1590	IX. 4124	D. apiculata	Sacc. and Berl., F. Austr. 4 (1885) ...	Apiculate-spored dothidella ...
1644	1592	...	D. inæqualis	Cooke, Grev. XX. 5 (1891) ...	Unequal dothidella ...
1645	1591	II. 5260	D. tephrosia	Sacc. Syll. II. 630 (1883) ...	Ash-coloured dothidella ...
178. MONTAGNELLA.—Speg.,					
1646	1593	IX. 4165	M. Eucalypti	Cooke and Mass., Grev. XVI. 5 (1887) ...	Eucalypt montagnella ...
1647	1594	...	M. rugulosa	Cooke, Grev. XX. 5 (1891) ...	Wrinkled montagnella ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Clamp. I. 168 (1798).— <i>Sphaeria</i> , <i>Diatrype</i> , <i>Anthostoma</i> , <i>Nummularia</i> — <i>continued</i> .									
1623A	...	...	...	...	...	Q.	...	Wood ...	Rather thick, externally, minutely and densely papillate, clay to red internally.
1624	...	...	T.	...	...	...	...	Wood ...	Hemispherical or nearly globose, superficial, black, shining dusky black within. Cup-shaped depression around opening of receptacle.
1625	...	...	...	...	...	Q.	B.	Bark and wood ...	Hemispherical, confluent, finally black, openings of receptacles singly in distinct marginate disc.
1626	W.A.	...	T.	V.	...	...	B.	Bark or wood ...	Bursting through, hemispherical, thick, variously shaped, often deformed. Receptacles rather large, globose.
1627	...	...	...	...	...	Q.	...	Rotten wood ...	Broadly expanded, coffee coloured, blackening. Receptacles globose, sparingly confluent.
1628	...	...	...	...	...	Q.	...	Old trunks ...	Large, margin inflexed, substance black, surface rusty to black. Receptacles oblong, immersed.
1629	...	...	...	...	...	Q.	...	Rotten branches ...	Very broadly expanded, black, girt by ruptured epidermis. Receptacles small, ovoid, crowded.
1630	...	..	...	V.	...	Q.	...	Bark and wood ...	Bursting through, cushion shaped, irregular form when young, clay colour to bright red, old red brown or dark red.
1631	...	S.A.	..	...	...	...	...	Trunks ...	Expanded, cushion shaped, thick, surface rust coloured, substance black. Receptacles oblong.
1632	...	...	...	...	...	Q.	B.	Rotten wood ...	On wood, rarely on bark, spread out in narrow thin crust, dark brown or black. Receptacles large, thickly crowded.
1633	...	...	...	..	...	Q.	...	Bark of dead trees	Hemispherical, large, corky to woody, sooty brown. Receptacles in outer layer black, oblong.

NITS AND FCKL, SYMB. 214 (1869).

Fekl, Smyb. Myc. 216 (1869).—*Dothidea*, *Sphaeria*.

1634	...	...	...	...	...	Q.	...	Fading leaves of <i>Alpinia carulea</i>	Spots brown, then pitch black, elongated, running together here and there.
1635	...	S.A.	...	...	...	...	...	Stems of <i>Scirpus nodosus</i>	Elongated, immersed, making matrix brownish. Receptacles parallel, globose, sporidia unequal-sided.
1636	...	...	...	...	...	Q.	...	<i>Fimbristylis</i> ...	Black, covered by epidermis, openings granulate.
1637	...	...	...	V.	...	...	B.	Leaves of grass, dying or dead	Distinct or run together in parenchyma of leaf, covered by shining blackened epidermis.
1638	...	...	...	V.	...	...	B.	<i>Juncus</i> ...	Internal, brown, epidermis ultimately brownish and blackish, cracked lengthwise.
1639	...	...	...	V.	...	...	...	Leaves of Eucalypts	Gregarious, on blistered tawny spots of living leaves, black, half immersed.
1640	...	...	...	...	...	Q.	...	Living and dry leaves of <i>Cordyline terminalis</i> , var. <i>Candollei</i>	Elongated, lance shaped, running parallel with veins, shining black, in brown spots.
1641	...	...	...	...	...	Q.	...	Phyllodes of <i>Acacia penninervis</i> and leaves of Figs	Immersed, black, shining, pustules black, minute, warty, warts with openings.
1642	...	...	...	V.	...	...	B.	Leaves of clover ...	Internal, forming brownish spots, at first producing conidia.

Fung. Arg. Pug. IV. 186 (1882).—*Phyllachora*, *Sphaeria*.

1643	...	...	...	...	...	Q.	...	Fading leaves of <i>Litsea dealbata</i>	On ochrey-brown spots, loosely gregarious, at first covered with epidermis, shining black.
1644	...	...	...	V.	...	...	...	Dead leaves of Eucalypts	Bursting through on both surfaces, nearly circular, black, shining.
1645	W.A.	...	...	...	...	...	...	Leaves ...	Internal, circular, plane then convex, ash coloured, openings point-like, black.

Fung. Arg. Pug. IV. 188 (1882).

1646	...	...	...	V.	...	...	...	Dead leaves of Eucalypts	Circular, convex, shining, black.
1647	...	...	...	V	...	...	...	Leaves of Eucalypts	On upper or under service, thin, somewhat circular, black, wrinkled.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
179. BAGNISIELLA.—Speg., F. Arg.					
1648	1597	...	<i>B. catervaria</i> ...	<i>Cooke, Handb. Austr. Fung.</i> 299 (1892) ...	Crowded bagnisiella ...
1649	1596	IX. 4026	<i>B. eudopyria</i> ...	<i>Sacc., Hedw.</i> 155 (1890) ...	Fiery bagnisiella ...
1650	1595	„ 4025	<i>B. rugulosa</i> ...	<i>Cooke, Grev.</i> XIX. 45 (1890) ...	Wrinkled bagnisiella ...
180. DARWINIELLA.—Speg.,					
1651	1598	II. 4167	<i>D. globulosa</i> ...	<i>Sacc., Syll.</i> IX. 1049 (1891) ...	Globose darwiniella ...
181. RHYTISMA.—Fries,					
1652	1600	VIII. 3027	<i>R. filicinum</i> ...	<i>Berk. and Br., Linn. Journ.</i> XIV. 130 (1875) ...	Fern rhytisma ...
1653	1599	X. 4655	<i>R. hypoxanthum</i> ...	<i>Berk. and Br., Fung. Brisb.</i> II. 71 (1883) ...	Buff-coloured rhytisma ...
182. TRABUTIA.—Sacc. and					
1654	1603	IX. 2439	<i>T. Eucalypti</i> ...	<i>Cooke and Mass., Grev.</i> XVII. 43 (1888) ...	Eucalypt trabutia ...
1655	1601	...	<i>T. parvicapsa</i> ...	<i>Cooke, Grev.</i> XX. 5 (1891) ...	Small-receptacled trabutia ...
1656	1602	...	<i>T. phylloidiæ</i> ...	<i>Cooke and Mass., Grev.</i> XIX. 60 (1891) ...	Phyllode trabutia ...
ORDER XVII.—					
183. SARCOXYLON.—Cooke,					
1657	1605	I. 1231	<i>S. compunctum</i> ...	<i>Cooke, Grev.</i> XIII. 107 (1885) ...	Punctate sarcoxylon ...
184. GIBELLIA.—Sacc.,					
1658	1606	IX. 2470	<i>G. dothideoides</i> ...	<i>Sacc. and Berl., Misc. Myc.</i> II. 23 (1885) ...	Dothidea-like gibellia...
185. BOTRYOSPHERIA.—Ces. and De Not.,					
1659	...	...	<i>B. hypoxyloidea</i> ...	<i>Cooke, Grev.</i> XIII. 102 (1885) ...	Hypoxylon-like botryosphaeria...
186. MELOGRAMMA.—Tul., Carp. II. 81 (1863).—					
1660	1607	I. 2814	<i>M. rubricosa</i> ...	<i>Cooke, Handb. Austr. Fung.</i> 301 (1892) ...	Red melogramma ...
ORDER XVIII.—					
187. DIATRYPE.—Fries, p.p. Nits					
1661	1610	I. 715	<i>D. chlorosarca</i> ...	<i>Berk. and Br., Linn. Journ.</i> 123 (1875) ...	Green-fleshed diatrype ...
1662	1608	„ 740	<i>D. glomeraria</i> ...	<i>Berk., Fl. N. Zeal.</i> II. 205 (1855) ...	Ball-like diatrype ...
1663	1609	„ 705	<i>D. stigma</i> ...	<i>Fries, S.V.S.</i> 385 (1849) ...	Stigma diatrype ...
188. CÆLOSOPHERIA.—Sacc.,					
1664	...	II. 5890 IX. 1836	<i>C. leptosporoides</i> ...	<i>Wint., Hedw.</i> 2 (1883) ...	Leptospora-like cælosphaeria ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Pug. III. 22 (1882).—Phyllachora, Dothidea.									
1648	...	...	...	...	...	Q.	...	Leaves of <i>Ficus</i> ...	Pustules minute, crowded in orbicular or irregular spots, openings pap-like.
1649	...	...	..	V.	...	...	...	Leaves of <i>Myoporum platycarpum</i>	Minute, disc shaped, black, crowded here and there, surface wrinkled, leathery, substance bright fiery orange.
1650	...	...	...	V.	...	...	...	Leaves of Eucalypts	On upper or under surface, gregarious, globose, black, wrinkled.
F. Fueg. 279 (1887).—Dothidea.									
1651	...	...	T.	...	...	...	...	Leaves of <i>Tasmania aromatica</i>	On both surfaces, globose, wrinkled, black, opaque.
S.M. II. 569 (1822).—Marchalia.									
1652	...	...	...	...	...	Q.	...	Fronds of <i>Alsophila</i>	Spots rather circular, thin. Cells elongated, wavy, thin.
1653	...	...	...	...	...	Q.	...	Leaves of <i>Cudrania javanensis</i>	Spots irregular, thickened, buff coloured, layer shining black, margin distinct.
Roum., Rev. Myc. 27 (1881).									
1654	...	...	T.	V.	...	...	...	Leaves of Eucalypts	Leathery, somewhat circular, convex, wrinkled, black, shining.
1655	...	...	...	V.	...	...	...	Phyllodes of <i>Acacia</i>	Internal. Receptacles on brown elliptical spots, crowded, small, black, shining.
1656	...	...	...	V.	...	...	...	Phyllodes of <i>Acacia longifolia</i>	Receptacles convex, brown, four to ten, seated on circular spots, with pore at apex.

MELOGRAMMACEÆ, NITS.

Grev. XIII. 107 (1885).

1657	...	...	...	...	...	Q.	...	Prostrate trunks ...	Globose, deformed, constricted at base, smooth, pale tan, punctate with black openings.
------	-----	-----	-----	-----	-----	----	-----	----------------------	---

Misc. Myc. II. 12 (1885).

1658	...	...	...	...	...	Q.	...	Bark ...	Cushion shaped, loosely gregarious, black, paler within, openings of receptacles point-like.
------	-----	-----	-----	-----	-----	----	-----	----------	--

Schem. Sfer. Comm. I. 211 (1861).

1659	...	...	...	...	...	...	...	Branches ...	Soon superficial, with habit and appearance of <i>Hypoxydon</i> , but not carbonaceous. Receptacles very small.
------	-----	-----	-----	-----	-----	-----	-----	--------------	---

Venturia, Hypoxydon, Sphaeria, Valsaria.

1660	W.A.	...	...	...	...	Q.	B.	Bark, dead wood ...	Deformed, tubercled, wrinkled and cracked, reddish. Receptacles immersed, black, shining.
------	------	-----	-----	-----	-----	----	----	---------------------	---

DIATRYPACEÆ, FRIES.

Pyr. Germ. 64 (1867).—Sphaeria.

1661	...	...	...	...	...	Q.	...	Branches and bark of trees	Small, pustulate, circular, green inside.
1662	...	...	...	V.	...	...	...	Branches of <i>Rhipigonum parviflorum</i>	Bursting through, angular or run together, and elongated. Receptacles ovate.
1663	...	...	...	...	...	Q.	B.	Branches ...	Receptacle-bearing layer, bursting through, long and broad, at length black. Receptacles ovoid.

Myc. Ven. Spec. 115 (1873)

1664	...	...	...	...	...	...	...	Branches ...	Receptacles gregarious, at first moist and globose, at length dry and cup shaped, dark brown.
------	-----	-----	-----	-----	-----	-----	-----	--------------	---

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER XIX.—					
189. VALSA.—Fries,					
1665	...	I. 476	V. decorticans ...	Fries, S.V.S. 412 (1849) ...	Decorticating valsa ...
1666	1612	„ 498	V. echidua ..	Cooke, Grev. IX. 4 (1880) ...	Hedgehog valsa ...
190. EUTYPELLA.—Nits,					
1667	1611	I. 571	E. stellulata ...	Sacc. Syll. I. 149 (1882) ...	Stellate eutypella ...
ORDER XX.—EUTYPACEÆ,					
191. CRYPTOVALSA.—Ces. and De Not.,					
1668	1613	I. 702	C. elevata ...	Sacc. Syll. I. 191 (1882) ...	Elevated cryptovalsa ...
192. CRYPTOSPHÆRELLA.—Sacc.,					
1669	...	IX. 1940	C. Macrozamia ...	Sacc. Syll. IX. 471 (1891) ...	Macrozamia cryptosphaerella ...
193. EUTYPA.—Tul., Sel. Fung.					
1670	1614	I. 637	E. lata ...	Tul., Sel. Fung. Carp. II. 56 (1861) ...	Broad eutypa ...
1671	1616	„ 632	E. ludibunda ...	Sacc., Mich. I. 15 (1878) ...	Sportive eutypa ...
1672	1615	IX. 1926	E. polyscia ...	Berl. and Vogl., in Sacc. Syll. IX. 467 (1891) ...	Shadowy eutypa ...
ORDER XXI.—CUCURBITARIACEÆ,					
191. GIBBERIDEA.—Fekl.,					
1673	1617	II. 3637	G. Archeri ...	Cooke, Handb. Aust. Fung. 304 (1892) ...	Archer's gibberidea ...
1674	1618	IX. 3340	G. plagia ...	Sacc. Syll. IX. 820 (1891) ...	Striped gibberidea ...
ORDER XXII.—					
195. BYSSOSPHERIA.—Cooke,					
1675	1620	IX. 2450	B. acanthostroma ...	Cooke, Handb. Aust. Fung. 305 (1892) ...	Thorny byssospheria ...
1676	1619	I. 916	B. aquila ...	Cooke, Grev. XV. 122 (1887) ...	Eagle byssospheria ...
196. LASIOSPHERIA.—Ces. and De Not.,					
1677	1622	IX. 3469	L. larvaspora ...	Cooke and Mass, Grev. XIX. 83 (1891) ...	Larva-spored lasiosphaeria ...
1678	1621	II. 3568	L. ovina ...	Ces. and De Not., Schema Sfer. 229 (1870) ...	Woolly lasiosphaeria ...
197. PLEOSPHERIA.—Speg.,					
1679	1623	II. 3927	P. pulvinula ...	Sacc. Syll. II. (1883) ...	Cushion-shaped pleospheria ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
VALSACEÆ, FRIES.									
S.V.S., 410 (1849).—Sphæria.									
1665	...	...	...	...	...	Q.	...	Dry branches	Stroma somewhat circular or often oval, abruptly tapering into a disc of openings, becoming dusky, at length black.
1666	...	...	...	...	N.S.W.	Q.	...	Bark	Bursting through. Receptacles nestling in a white powdery layer.
Pyr. Germ. 163 (1867).—Valsa, Sphæria.									
1667	...	...	...	V.	...	...	B.	Branches...	Somewhat round, immersed; receptacle-bearing layer, white or dirty white; openings radiately stellate.
COOKE, GREV. XIV. 93 (1886).									
Schema Sfer. It. 29 (1870).—Diatrype, Sphæria, Eutypa.									
1668	W.A.	...	T.	...	...	...	B.	Dead branches	Elongated, emergent, black or grey. Receptacles gregarious, globose, immersed in wood.
Syll. I. 186 (1882).—Sphæria.									
1669	...	...	...	...	...	Q.	...	Fruit of <i>Macrozamia Hopei</i>	Receptacles under epidermis, at first scattered, then crowded, black, springing from dark-brown filaments.
Carp. II. 52 (1861).—Sphæria, Valsa.									
1670	...	...	T.	...	...	...	B.	Wood and bark	Receptacle-bearing layer long and broad, innate in wood or bark, brown or ashy. Receptacles immersed, sphaeroid.
1671	...	...	...	V.	...	...	...	Branches...	Receptacle-bearing layer spreading. Receptacles globose, externally black, at first mealy white.
1672	...	...	...	...	...	Q.	...	Epicarp of <i>Cucurbita lagenaria</i>	Receptacle-bearing layer black, point-like. Receptacles immersed.
COOKE, GREV. XV. 83 (1887).									
Symb. Myc. 168 (1869).—Zignoella, Sphæria, Melanomma.									
1673	...	...	T.	...	...	...	...	Rotten wood	Crowded. Receptacles wrinkled, at length collapsing and cup shaped.
1674	...	...	...	V.	...	...	...	Living twigs of <i>Cassinia aculeata</i>	Receptacles densely crowded, at length run together in large patches, globose, black, shining, smooth.
SUPERFICIALES, FRIES.									
Grev. VII. 84 (1879).—Scortechinia, Rosellinia, Sphæria.									
1675	...	...	...	...	...	Q.	...	Wood and bark	Finely filamentous. Receptacles very small, globose, crowded, black, not pap-like.
1676	...	...	T.	...	...	...	B.	Wood and bark	Receptacles gregarious or densely crowded, globose, dark brown, pap-like.
Schema Sfer. 55 (1870).—Sphæria, Leptospora.									
1677	...	...	...	V.	...	...	...	Bark	Receptacles loosely gregarious, globose, covered with mealy fluffy sulphur-coloured coat. Sporidia long, spindle shaped, with fifteen to nineteen partitions.
1678	...	...	...	V.	...	...	B.	Rotten wood	Receptacles gregarious, almost globose, covered with mealy fluffy lemon-coloured coat.
Fung. Arg. IV. 65 (1882).—Sphæria, Coniochaeta, Lasiosphæria.									
1679	W.A.	...	...	...	...	...	...	Rotten wood	Scattered, somewhat globose, collapsed and depressed at length, hairy, black.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
198. VENTURIA.—De Not. and Ces.,					
1680	1624	I. 2311	V. circinans	Sacc., Mich. I. 499 (1868)	Circinate venturia
199. CHÆTOMIUM.—Kunze,					
1681	1626	IX. 1997	C. cymatotrichum	Cooke, Grev. XII. 21 (1883)	Wavy-haired chætomium
1682	1625	I. 793	C. comatum	Fries, S.M. III. 253 (1832)	Hairy chætomium
200. ROSELLINIA.—De Not.,					
1683	1627	I. 970	R. inspersa	Sacc. Syll. I. 265 (1882)	Scattered rosellinia
1684	1628	IX. 2055	R. tremellicola	Cooke and Mass., Grev. XVIII. 6 (1889)	Tremella rosellinia
201. TREMATOSPHERIA.—Fekl.,					
1685	1629	IX. 3307	T. congesta	Berl. and Vogl., in Sacc. Syll. IX. 811 (1891)	Congested trematosphæria
ORDER XXIII.—					
202. CONISPHERIA.—Cooke,					
1686	1632	IX. 3539	C. australica	Cooke and Mass., Handb. Austr. Fung. 307 (1892)	Australian conisphæria
1687	1631	...	C. erumpens	Cooke, Handb. Austr. Fung. 307 (1892)	Erumpent conisphæria
1688	1630	IX. 3527	C. subcorticalis	Cooke, Handb. Austr. Fung. 307 (1892)	Underbark conisphæria
ORDER XXIV.—LOPHIOSTOMACEÆ,					
203. LOPHIOSTOMA.—Ces. and De Not.,					
1689	1633	II. 5405	L. Schomburgkii	Cooke, Handb. Austr. Fung. 307 (1892)	Schomburgk's lophiostoma
ORDER XX.—					
204. RHAMPHORIA.—Niesl.,					
1690	1634	IX. 3684	R. tenella	Sacc., Hedw. 155 (1890)	Delicate rhamphoria
ORDER XXVI.—					
205. MASSARIELLA.—Speg.,					
1691	1635	I. 2707	M. australis	Sacc. Syll. I. 716 (1882)	Southern massariella
206. DIDYMOSPHERIA.—Fekl.,					
1692	1638	IX. 3000	D. Banksiæ	Cooke, Grev. XIX. 90 (1891)	Banksia didymosphæria
1693	1637	„ 2979	D. conoidella	Sacc. and Berl., Misc. Myc. II. 26 (1885)	Conical didymosphæria

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Schema Sf. I. 225 (1870).—Perisporium, Stigmatea.									
1680	...	...	...	V.	...	...	B.	Leaves of <i>Geranium</i>	Receptacles clustered in patches or spots, hardly circinating, hairs thickened at base.
Myc. Heft. I. 15 (1817).—Sphaeria.									
1681	...	...	...	...	...	Q.	...	Leaves of <i>Solanum Dallachyi</i>	Gregarious. Receptacles depressed globose, woolly, sooty olive.
1682	...	S.A.	...	V.	...	...	B.	Rotting grass, &c. ...	Receptacles gregarious, nearly ovoid, thin black hairs radiating from base.
Giorn. Bot. Ital. II. 334 (1847).—Sphaeria.									
1683	W.A.	...	...	...	...	...	...	Rotten wood	Crowded or scattered, black. Receptacles nearly globose, wrinkled
1684	...	...	...	...	...	Q.	...	<i>Tremella fuciformis</i>	Receptacles scattered, globose, superficial, black, pap-like, smooth.
Symb. Myc. 161 (1869) emended.—Psilosphaeria, Melanomma.									
1685	...	...	...	...	N.S.W.	...	...	Bark	Gregarious, crowded, black. Receptacles convex, smooth, pierced with a pore.

PERTUSACEÆ, FRIES.

Grev. XVI. 87 (1888).—Zignoëlla.

1686	...	...	...	V.	...	...	...	Naked branches	Receptacles scattered, half immersed, rather conical, base buried in wood.
1687	...	...	...	V.	...	...	...	Twigs	Scattered or collected together, erumpent. Receptacles globose, smooth, black.
1688	...	...	...	...	...	...	...	Inside dead bark of trees	Scattered. Receptacles half immersed, pierced, black.

SACC, MICH. I. 333 (1879).

Schem. Sfer. 45 (1870).—Schizostoma, Sphaeria.

1689	...	...	...	...	...	Q.	...	Wood	Receptacles large, free, black, openings linear.
------	-----	-----	-----	-----	-----	----	-----	------	--

CERATOSTOMACEÆ, FRIES.

Notiz. 44 (1876).

1690	...	S.A.	...	...	...	...	...	Rotten wood of <i>Eucalyptus viminalis</i>	Receptacles almost superficial, or base buried in wood, small, globose, black, thinly carbonaceous.
------	-----	------	-----	-----	-----	-----	-----	--	---

OBTECTACEÆ, FRIES., S.V.S. (1849).

Fung. Arg. Pug. 2 (1880).—Massaria.

1691	...	...	...	V.	...	...	...	Bark	Scattered, covered, inconspicuous. Receptacles depressed.
------	-----	-----	-----	----	-----	-----	-----	------	---

Symb. Myc. 140 (1869) emended.

1692	...	...	...	V.	...	...	...	Living leaves of <i>Banksia</i>	Spots circular, pale with indistinct brown margin. Receptacles few, black, bursting through.
1693	...	...	...	...	...	Q.	...	Dead branches of <i>Copparis</i>	Receptacles loosely gregarious, globose, then conical, black.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER XXVII.—CAULICOLACEÆ,					
207. PHYSALOSPORA.—Niessl., Notiz.					
1694	1639	I. 1660	<i>P. gregaria</i>	Sacc. Syll. I. (1882)	Gregarious physalospora ..
1695	...	„ 1723	<i>P. labeaula</i>	Sacc. Syll. I. 447 (1882)	Stain physalospora
1696	...	IX. 2427	<i>P. Sacchari</i>	Sacc. Syll. IX. 599 (1891)	Sugar-cane physalospora ...
208. DIDYMELLA.—Sacc., Mich. I. 377 (1878).—					
1697	1640	I. 2174	<i>D. Bryonie</i>	Rehm., Ascom. III. 99 (1881)	Bryonia didymella
1698	1636	„ 2126	<i>D. cladophila</i>	Sacc. Syll. I. 545 (1882)	Brauch-loving didymella ...
209. ANTHOSTOMELLA.—Sacc., Consp.					
1699	1641	...	<i>A. Lepidospermæ</i>	Cooke, Grev. XX. 5 (1891)	Lepidosperma anthostomella ...
210. PLEOSPORA.—Rabh., Herb. Myc.					
1700	1643	II. 3776	<i>P. Aucubæ</i>	Lamb., Fl. Myc. Belg. II. 268 (1889)	Aucuba pleospora
1701	1642	„ 3730	<i>P. herbarum</i>	Rabh., Herb. Myc. Ed. II. 547 (1858)	Herb pleospora
ORDER XXVIII.—FOLIICOLACEÆ,					
211. LÆSTADIA.—Auersw., in Hedw. 177					
1702	..	IX. 2368	<i>L. Dammaræ</i>	Sacc. Syll. IX. 586 (1891)	Dammara læstadia
1703	1644	„ 2378	<i>L. destructiva</i>	Berl. and Vogl., in Sacc. Syll. IX. 588 (1891)	Destructive læstadia
1704	„ bis.	...	<i>L. Litscæ</i>	Cooke, Grev. XX. 65 (1892)	Litscæ læstadia
1705	...	IX. 2346	<i>L. Melaleucæ</i>	Sacc. Syll. IX. 581 (1891)	Melaleuca læstadia
1706	1645	„ 2400	<i>L. phyllodiæ</i>	Cooke, Handb. Austr. Fung. 310 (1892)	Phyllode læstadia
212. SPHERELLA.—Ces. and De Not.,					
1707	1651	IX. 2556	<i>S. Alyxiæ</i>	Cooke and Mass., Grev. XVI. 5 (1887)	Alyxia sphærella
1708	...	I. 1921	<i>S. atra</i>	Sacc. Syll. I. 498 (1882)	Black sphærella
1709	1650	IX. 2578	<i>S. Banksiæ</i>	Cooke and Mass., Grev. XVI. 114 (1888)	Banksia sphærella
1710	1647	...	<i>S. cryptica</i>	Cooke, Grev. XX. 5 (1891)	Hidden sphærella
1711	1649	I. 1906	<i>S. Euonymi</i>	Auersw., Myc. Eur. 10	Euonymus sphærella
1712	...	„ 1951	<i>S. Fragariæ</i>	Sacc. Syll. I. 505 (1882)	Strawberry sphærella... ..
1713	...	...	<i>S. Goodiæfolia</i>	Cooke, Grev. XXI. 38 (1892)	Goodiæ-leaf sphærella ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
FRIES, S.M. II. 503 (1823).									
Kr. Pyr. 10 (1876).—Sphaeria.									
1694	...	...	...	...	...	Q.	...	Stems of <i>Ricinus communis</i>	Receptacles densely gregarious, covered by epidermis, globose, black white within.
1695	...	...	...	...	...	...	...	Leaves of <i>Acacia verticillata</i>	Innate, upper or under surface spot-like, black, shining. Receptacles nearly globose, black within.
1696	...	...	...	...	...	Q.	...	Leaves of Sugar cane	Receptacles scattered or gregarious, minute, black, bursting through, filaments dark brown.
Sphaeria, Sphaerella, Didymosphaeria.									
1697	...	...	...	...	...	Q.	B.	Twigs of cucurbitaceous plant	Receptacles beneath epidermis, at length almost free, gregarious, very minute, black.
1698	...	S.A.	...	...	...	...	...	Branches of Grape vine	Receptacles loosely gregarious, covered with bleached cuticle, hemispherical, black, leathery.
Gen. Pyr. Ital. 8 (1875).									
1699	...	...	...	V.	...	...	...	<i>Lepidosperma</i> ...	Receptacles on bleached elongated spots, with dark-brown border, globular.
Ed. II. 347 (1858).—Sphaeria, Phoma.									
1700	...	...	...	V.	...	...	...	Leaves of <i>Aucuba japonica</i>	Receptacles spherical, immersed, black, scattered upon somewhat circular brown spots.
1701	...	...	...	V.	...	...	B.	Herbaceous stems	Receptacles somewhat gregarious, at first covered, then almost naked, spherical, depressed.
FRIES, S.M. II. 513 (1823).									
(1869).—Sphaerella, Physalospora.									
1702	...	...	...	...	...	Q.	...	Leaves of <i>Dammara robusta</i> = <i>Agathis</i>	Spots pale brown or yellow, margin amber, then dark brown. Receptacles black, internal.
1703	...	...	...	V.	N.S.W.	Q.	...	Leaves of Lucerne	Receptacles minute, seated on brown spots; raised above general surface.
1704	...	...	...	...	...	Q.	...	Leaves of <i>Litsea</i> ...	Spots irregular or somewhat circular on upper surface, becoming pale, with broad brown border.
1705	...	...	...	...	N.S.W.	...	...	Leaves of <i>Melaleuca</i>	Spots circular, minute, brown, convex. Receptacles black, in crowded spots.
1706	...	...	...	V.	...	...	...	Phyllodes of <i>Acacia suaveolens</i>	Receptacles scattered, very thin, immersed, black, covered by blackened cuticle.
Schema Sfer. Ital. 62 (1870).—Sphaeria.									
1707	...	...	...	V.	...	...	...	Dead leaves of <i>Alyxia buxifolia</i>	On both sides. Receptacles gregarious, arising from within, nearly globose, covered by blackened cuticle.
1708	...	...	...	...	...	...	...	Leaves of <i>Grevillea</i>	Gregarious or confluent. Receptacles globose, white within, covered by a black expanded layer.
1709	...	...	...	V.	...	...	...	Fading leaves of <i>Banksia integrifolia</i>	On upper surface, spots none. Receptacles gregarious, black, pierced with a pore.
1710	...	...	...	V.	...	...	...	Fading leaves of Eucalypts	On both sides, spots reddish brown, large. Receptacles immersed in substance of leaf and hidden.
1711	...	...	...	V.	...	...	...	Dead leaves of <i>Euonymus</i>	On under surface. Receptacles black, beneath epidermis, globose, on greyish spots.
1712	...	S.A.	...	V.	N.S.W.	...	...	Leaves of Strawberry	Spots becoming purple, then pale towards the centre. Receptacles very minute, globular, black.
1713	...	...	...	V.	...	...	...	Leaves of <i>Goodia latifolia</i>	Spots circular, brown, surrounded by darker line. Receptacles gregarious, minute.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
212. SPHERELLA.—Ces. and De Not.,					
1714	1653	I. 2048	<i>S. graminicola</i> ...	Fekl, Symb. Myc. 101 (1875) ...	Grass-growing sphaerella ...
1715	1646	...	<i>S. nubilosa</i> ...	Cooke, Grev. XIX. 61 (1891) ..	Cloudy sphaerella ...
1716	1648	IX. 2598	<i>S. rubiginosa</i> ...	Cooke, Grev. XIV. 90 (1886) ...	Rubiginous sphaerella ...
1717	1652	I. 2028	<i>S. smilacicola</i> ...	Cooke, Grev. VI. 146 (1878) ...	Smilax sphaerella ...
213. SPHERULINA.—Sacc.,					
1718	1654	IX. 3189	<i>S. Camelliae</i> ...	Cooke, Handb. Austr. Fung. 312 (1892) ...	Camellia sphaerulina ...
ORDER XXIX.—MICROTHYRIACEÆ,					
214. MICROTHYRIUM.—Desm., Ann.					
1719	1655	IX. 4196	<i>M. amygdalinum</i> ...	Cooke and Mass., Grev. XIX. 90 (1891) ...	Amygdalina microthyrium ...
215. MICROPELTIS.—Mont., Ann.					
1720	1656	II. 5390	<i>M. applanata</i> ...	Mont., Ann. Sci. Nat. XVII. 122 (1842) ...	Flattened micropeltis ...
1720A	...	IX. 4236	<i>M. applanata</i> , var. <i>depau- perata</i>	Sacc. and Berl., Rev. Myc. (1885) ...	Impoverishing micropeltis ...
ORDER XXX.—PERISPORIACEÆ,					
216. PODOSPHERA.—Kunze,					
1721	...	I. 3	<i>P. tridactyla</i> ...	De Bary, Syst. Erys. in Hedw. 68 (1871) ...	Three-fingered podosphaera ... (Powdery mildew of apple)
217. SPHEROTHECA.—Lev., Ann. Sci.					
1722	1658	I. 6	<i>S. pannosa</i> ...	Lev., Ann. Sci. Nat. XV. 138 (1851) ...	Cloth-like sphaerotheca ... (Rose blight)
218. ERYSIPIHE.—Hedw., Lev.					
1723	...	I. 70	<i>E. communis</i> ...	Fries, S.V.S. 406 (1849) ...	Common erysiphe ...
1724	...	„ 74	<i>E. graminis</i> ...	D. C., Fl. Fr. VI. 106 (1815) ...	Grass erysiphe (Grass mildew)...
1725	1657	IX. 1571	<i>E. vitigera</i> ...	Cooke and Mass., Grev. XV. 98 (1887) ...	Vine-growing erysiphe ...
219. EUROTIIUM.—Link, Berl.					
1726	1659	I. 101	<i>E. herbariorum</i> ...	Link, Sp. Pl. I. 79 (1824) ...	Herbarium eurotium ...
1727	1660	„ 104	<i>E. lateritium</i> ...	Mont., Syll. 918 (1856) ...	Brick-red eurotium ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Schema Sfer. Ital. 62 (1870).— <i>Sphaeria</i> — <i>continued</i> .									
1714	...	...	...	V.	...	...	...	Leaves of grass	On upper surface. Receptacles growing from within, prominent, small, black, occupying entire surface.
1715	...	...	...	V.	...	...	...	Living leaves of Eucalypts	On under surface, spots circular or irregular, greyish brown, soon falling away. Receptacles very minute.
1716	...	...	...	...	...	Q.	...	Leaves of <i>Pittosporum rubiginosum</i>	On upper surface. Receptacles scattered, minute, point-like, prominent, black.
1717	...	...	...	...	...	Q.	...	Leaves of <i>Dioscorea</i> and <i>Smilax</i>	Spots reddish brown with black margin. Receptacles point-like, conical, black.

Mich. l. 399 (1878).

1718	...	...	...	V.	...	...	...	Living leaves of Camellia	Spots on upper surface, turning brownish, indeterminate. Receptacles scattered.
------	-----	-----	-----	----	-----	-----	-----	---------------------------	---

SACC. SYLL. II. 658 (1883).

Sci. Nat. XV. 137 (1841).

1719	...	S.A.	...	...	...	...	...	Living leaves of <i>Eucalyptus amygdalina</i>	Receptacles gregarious or scattered, on both surfaces, membranous, very dark brown.
------	-----	------	-----	-----	-----	-----	-----	---	---

Sci. Nat. XVII. 122 (1842).

1720	...	...	...	...	...	...	...	Languid leaves of <i>Eucalyptus tereticornis</i>	Receptacles on both surfaces, convex to flattened, black, finally opening in centre.
1720a	...	...	...	...	...	Q.	...		

FRIES, S.V.S. 375 (1849).

Mye. Heft. II. 111 (1817).—*Alphitomorpha*.

1721	...	...	...	V.	N.S.W.	...	...	Young leaves and shoots of Apple	On both surfaces of leaf, forming a white felt, and the spores so numerous as to make it powdery; conidia barrel shaped.
------	-----	-----	-----	----	--------	-----	-----	----------------------------------	--

Nat. XV. 138 (1851).—*Alphitomorpha*.

1722	...	...	...	V.	N.S.W.	Q.	B.	Rose leaves	Mycelium woolly, then cloth-like, white, persistent. <i>Oidium leucoconium</i> is the conidial stage.
------	-----	-----	-----	----	--------	----	----	-------------	---

Ann. Sci. Nat. XV. 161 (1851).

1723	...	...	...	V.	...	...	B.	Leaves of various plants	Mycelium spreading, arachnoid, evanescent or persistent. Receptacles scattered or gregarious, minute.
1724	...	...	...	V.	N.S.W.	...	B.	Leaves and stems of various grasses	Mycelium spreading, fluffly to woolly, ochrey white, jointed, persistent. Early or conidial stage is <i>Oidium monilioides</i> , Link.
1725	...	...	...	V.	...	...	...	Vine leaves	On both sides of leaf. Mycelium fluffly, persistent. Receptacles gregarious, very minute, spherical.

Mag. III. 31 (1809).—*Mucor*.

1726	...	...	...	...	...	Q.	B.	Plants in herbaria, decaying organic matter, &c.	Mycelium creeping, fluffly, branched, uncoloured. Conidial stage is <i>Aspergillus glaucus</i> , Link, with glaucous conidia.
1727	W.A.	...	...	...	...	Q.	...	Leaves of <i>Piperomia</i>	Mycelium dense, woolly, orange yellow. Receptacles membranous, yellow, then ochre, immersed.

SYSTEMATIC ARRANGEMENT

Number	Cooke's Number	Saccardo's Number	Scientific Name	Authority for Name	English Name
220. ASTERINA.—Lev., Ann.					
1728	1661	IX. 1609	<i>A. Baileyi</i>	Berk. and Br., Linn. Trans. II. 71 (1883)	Bailey's asterina
1729	1662	„ 1622	<i>A. correicola</i>	Cooke and Mass., Grev. XVI. 5 (1887) ...	Correa asterina
1730	...	...	<i>A. hoveaefolia</i>	Cooke and Mass., Grev. XXII. 36 (1893) ...	Hovea-leaved asterina
1731	1665	IX. 1607	<i>A. microthyrioides</i>	Winter, Hedw. 3 (1885)	Microthyrium-like asterina
1732	1663	I. 260	<i>A. pelliculosa</i>	Berk., Autare. Crypt. II. 453 (1847) ...	Pellicle asterina
1733	1664	IX. 1621	<i>A. platystoma</i>	Cooke and Mass., Grev. XVIII. 6 (1889)	Broad-mouthed asterina
1734	1668	I. 198	<i>A. reptans</i>	Berk. and Curt., Linn. Trans. X. 373 (1869)	Creeping asterina
221. ASTERELLA.—Sacc.					
1735	1666	IX. 1690	<i>A. Alsophila</i>	Cooke and Mass., Grev. XVIII. 81 (1889)	Alsophila asterella
1736	1667	„ 1679	<i>A. subcuticulosa</i>	Cooke, Grev. XVII. 81 (1889)	Subcuticular asterella
222. DIMEROSPORIUM.—Fekl.					
1737	1670	IX. 1710	<i>D. Ludwigianum</i>	Sacc., Hedw. 127 (1889)	Ludwig's dimerosporium
1738	1671	...	<i>D. parvulum</i>	Cooke, Grev. XX. 5 (1891)	Small dimerosporium
1739	...	...	<i>D. secedens</i>	Sacc., Hedw. 57 (1893)	Seceding dimerosporium
1740	...	...	<i>D. strigosum</i>	Sacc., Grev. XXI. 68 (1893)	Hispid dimerosporium
223. PARODIELLA.—Speg.					
1741	...	IX. 1723	<i>P. Banksiae</i>	Sacc. and Bizz., Syll. IX. 410 (1891) ...	Banksia parodiella
1742	1604	I. 2711	<i>P. Perisporioides</i>	Speg., Fung. Arg. Png. I. 178 (1880) ...	Perisporia-like parodiella
224. MELIOLA.—Fries,					
1743	1673	I. 287	<i>M. amphitricha</i>	Fries, Elench. II. 109 (1828)	Amphitrichous meliola
1744	...	„ 284	<i>M. elatricha</i>	Lev., Ann. Sci. Nat. V. 266 (1846) ...	Branch-haired meliola
1745	1672	„ 279	<i>M. corallina</i>	Mont., Syll. 910 (1856)	Coralline meliola
1746	1676	IX. 1758	<i>M. densa</i>	Cooke, Grev. XII. 85 (1881)	Dense meliola
1747	1679	I. 237	<i>M. mollis</i>	Berk. and Br., Linn. Trans. XIV. 136 (1875)	Soft meliola
1748	1674	„ 291	<i>M. Musae</i>	Mont., Syll. 905 (1856)	Musa meliola
1749	1678	IX. 1752	<i>M. octospora</i>	Cooke, Grev. XI. 38 (1882)	Eight-spored meliola
1750	1675	I. 294	<i>M. orbicularis</i>	Berk. and Curt., Linn. Trans. X. 392 (1869)	Orbicular meliola
1751	...	„ 301	<i>M. polytricha</i>	Kalch. and Cooke, Grev. VIII. 72 (1879)	Many-haired meliola
1752	1680	„ 310	<i>M. Tetracera</i>	Thuem., Symb. Myc. Aust. II. 92 (1878)	Tetracera meliola

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Sci. Nat. III. 59 (1845).									
1728	...	...	...	...	...	Q.	...	Leaves of <i>Hakea lorea</i> and other shrubs	Mycelial threads brown, knotted, branching, forming reddish-brown patches. Receptacles minute, wrinkled, brown.
1729	...	...	...	V.	...	...	...	Living leaves of <i>Correa Lawrenceana</i>	Circular black spots on leaves. Receptacles convex to flat, black, crowded on spots.
1730	...	...	...	...	...	Q.	...	Leaves of <i>Hovea longifolia</i>	Spots black, or with brown centre, nearly circular. Receptacles usually arranged in a ring, black.
1731	...	...	...	V.	...	...	...	Leaves of <i>Eucalyptus pilularis</i>	On under surface. Receptacles scattered or loosely gregarious, very minute, black, wrinkled, margin brown.
1732	...	...	...	...	...	Q.	...	Leaves of <i>Trema aspera</i>	Mycelium forming a pellicle, spot-like, black. Receptacles globose, depressed, black.
1733	...	...	...	...	...	Q.	...	Living leaves of <i>Castanospermum australe</i>	Mycelium thin, more or less circular, tree-like, black. Receptacles convex to flattened, black.
1734	...	...	...	...	...	Q.	...	Leaves of <i>Eugenia</i>	Mycelium thin, rather netted. Receptacles minute, formed from the radiating cells.
Syll. I. 42 (1882).									
1735	...	...	...	...	...	Q.	...	Fronds of <i>Alsophila Rebecca</i>	Receptacles membranous, disc-like, nearly circular, mostly running together in oblong or irregular pitchy-black patches.
1736	...	...	...	V.	N.S.W.	...	...	Fading and dead leaves of <i>Aster argophyllus</i>	Receptacles thin, flattened, without mycelium, black or brown.
Symb. Myc. 89 (1869).									
1737	...	S.A.	...	V.	...	...	...	Fading leaves of <i>Lagenophora Billardieri</i>	Mycelium forming pale sooty spots on both surfaces. Receptacles crowded here and there, globular, superficial, dark sooty brown.
1738	...	...	...	...	...	Q.	...	Living leaves of <i>Trema aspera</i>	Mycelium brown, sparse, radiating, on irregular black spots. Receptacles minute, nearly globose.
1739	...	...	...	...	...	Q.	...	Leaves of living plants	Broadly expanded, pitch black, easily separating, capnodium-like. Receptacles thickly clustered, globose, shining black.
1740	...	...	...	...	...	...	...	...	(Belongs to genus <i>Dicmasporium</i> .)
Fung. Arg. Pug. I. 178 (1880).—Dothidea.									
1741	...	...	...	...	N.S.W.	...	...	Lower surface of languid leaves of <i>Banksia marginata</i>	Receptacles globose, black, superficial, like dots, densely clustered. Mycelium almost none.
1742	...	...	...	V.	N.S.W.	...	...	Living leaves of <i>Leguminosæ</i>	Receptacles globose, black, superficial, thickly clustered, often covering entire surface, sooty olive.
Elench. II. 100 (1828).—Sphaeria, Dothidea.									
1743	...	...	...	V.	...	Q.	...	Leaves of <i>Cupania Eucalyptus</i> , <i>Flindersia</i> , <i>Acacia</i> , &c.	Lower surface. Mycelium spot-like, radiating from centre, continuous, black. Receptacles globose, surrounded by black rigid erect appendages.
1744	...	...	...	V.	...	...	...	Leaves of <i>Eugenia</i>	Mycelium spread out, with radiating branched circumference. Receptacles globose, with erect bifid appendages.
1745	...	...	...	...	...	Q.	...	Leaves ...	Both surfaces. Mycelium spot-like, spots circular, black. Receptacles large, globose, surrounded by rigid shiny black appendages.
1746	...	...	...	...	...	Q.	...	Leaves of <i>Eucalyptus</i>	Under rarely upper surface, forming circular very black velvety spots. Receptacles globose, black, surrounded by crowded erect appendages.
1747	...	...	...	...	...	Q.	...	Leaves ...	Mycelium of soft black threads. Receptacles globose, appendages erect, brown.
1748	...	...	...	...	...	Q.	...	<i>Muscæ</i> ...	Spot like, large, black tufts. Receptacles very minute and inconspicuous; appendages erect, simple.
1749	...	...	...	...	...	Q.	...	Leaves of <i>Tristania conferta</i>	Spots circular, minute, velvety. Receptacles medium sized; appendages erect.
1750	...	...	...	...	...	Q.	...	Branches and leaves	Spots thick, orbicular. Receptacles globose; appendages flexuous, curved.
1751	...	...	...	...	...	Q.	...	Leaves of <i>Callistemon</i>	On under or both surfaces, black, spot-like. Mycelium spread out, radiating, conidia bearing. Receptacles globose, with erect acute wavy appendages.
1752	...	...	...	...	...	Q.	...	Leaves of <i>Tretacera Wuthiana</i>	Both surfaces, spots more or less circular, black, fading. Receptacles carbonaceous, globose.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
225. ZUKALIA.—Sacc.					
1753	1677	IX. 1792	<i>Z. loganiensis</i> ...	<i>Sacc. and Berl., Syll. IX. 431 (1891)</i> ...	Logan zukalia ...
226. ASTERIDIUM.—Sacc.					
1754	1669	IX. 1808	<i>A. Eucalypti</i> ...	<i>Cooke and Mass., Grev. XVI. 74 (1888)</i> ...	Eucalyptus asteridium ...
227. CAPNODIUM.—Mont., Ann.					
1755	1683	I. 324	<i>C. australe</i> ...	<i>Mont., Syll. 916 (1856)</i> ...	Southern capnodium ...
1756	1681	„ 346	<i>C. citri</i> ...	<i>Berk. and Desm., Journ. Hort. Lond. IV. 11 (1849)</i> ...	Citrus capnodium ...
1757	1682	„ 329	<i>C. elongatum</i> ...	<i>Berk. and Desm., Journ. Hort. Lond. IV. 251 (1849)</i> ...	Elongated capnodium ...
1758	1684	„ 323	<i>C. salicinum</i> ...	<i>Mont., Syll. 915 (1856)</i> ...	Willow capnodium ...
1759	...	...	<i>C. Walteri</i> ...	<i>Sacc., Hedw. 58 (1893)</i> ...	Walter's capnodium ...
228. ANTENNARIA.—Link,					
1760	1687	I. 362	<i>A. Robinsoni</i> ...	<i>Berk. and Mont., in Mont. Syll. 1066 (1856)</i> ...	Robinson's antennaria ...
1761	1686	„ 364	<i>A. scoriadea</i> ...	<i>Berk., Hook., Journ. 70 (1845)...</i> ...	Drossy antennaria ...
1762	1688	„ 366	<i>A. semiovata</i> ...	<i>Berk. and Br., Ann. Nat. Hist. XIII., 2 Ser. 468 (1854)</i> ...	Semi-ovate antennaria ...
ORDER XXXI.—HYSTERIACEÆ,					
229. AULOGRAPHUM.—Lib., Crypt.					
1763	1475	IX. 4344	<i>A. Eucalypti</i> ...	<i>Cooke and Mass., Grev. XVIII. 6 (1889)</i> ...	Eucalypt aulographum ...
1764	1474	„ 4343	<i>A. melioides</i> ...	<i>Cooke and Mass., Grev. XVIII. 6 (1889)</i> ...	Meliola-like aulographum ...
230. GLONIUM.—Muhl., Cat.					
1765	1476	II. 5586	<i>G. stellatum</i> ...	<i>Muhl., Cat. Am. 101 (1813)</i> ...	Stellate glonium ...
1766	1477	„ 5607	<i>G. tardum</i> ...	<i>Sacc., Syll. II. 737 (1883)</i> ...	Slow-opening glonium ...
231. LEMBOSIA.—Lev., Ann.					
1767	1478	IX. 4353	<i>L. graphioides</i> ...	<i>Sacc. and Berl., Misc. Myc. II. 6 (1885)</i> ...	Graphium-like lembosia ...
1768	...	„ 4354	<i>L. orbicularis</i> ...	<i>Winter, Hedw. 29 (1885)</i> ...	Orbicular lembosia ...
232. HYSTERIUM.—Tode, Fung.					
1769	1479	II. 5634	<i>H. pulicæ</i> ...	<i>Pers., Syn. 98 (1801)</i> ...	Flea-like hystereum ...
233. TRYBLIDIELLA.—Sacc.					
1770	1480	II. 5694	<i>T. rufula</i> ...	<i>Sacc., Syll. II. 757 (1883)</i> ...	Reddish tryblidiella ...
234. RHYTHIDHYSTERIUM.—Speg.,					
1771	1481	IX. 4378	<i>R. Scortechini</i> ...	<i>Sacc. and Berl., Misc. Myc. II. 7 (1885)</i> ...	Scortechini's rhytidhystereum ...
235. PLATYCHEILUS.—Cooke,					
1772	1482	...	<i>P. cæspitosus</i> ...	<i>Cooke and Mass., Handb. Austr. Fung. 409 (1892)</i> ...	Tufted platycheilus ...
236. HYSTEROGRAPHIUM.—Corda,					
1773	1483	II. 5759	<i>H. elongatum</i> ...	<i>Corda, Icon. V. 77 (1842)</i> ...	Elongated hystereographium ...
1774	1484	„ 5771	<i>H. hiæcens</i> ...	<i>Rehm., Ascom. 314 (1881)</i> ...	Gaping hystereographium ...
1775	1485	„ 5768	<i>H. Roussellii</i> ...	<i>Sacc., Syll. II. 779 (1883)</i> ...	Roussel's hystereographium ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Syll. IX. 431 (1891).— <i>Meliola</i> .									
1753	...	...	...	...	...	Q.	...	Leaves of <i>Smilax</i> ...	Upper surface. Mycelium thin, widely spreading. Receptacles globose, black, sparingly beset with sooty-brown bristles.
Syll. I. 25 (1882).									
1754	...	...	...	V.	...	...	...	Dead leaves of <i>Eucalyptus amygdalina</i>	Very thickly gregarious. Receptacles minute, disc-like, flattened, black
Sci. Nat. 3 XI. 233 (1848).— <i>Fumago</i> .									
1755	W.A.	...	...	...	...	Q.	...	Branches of Conifers, <i>Cycas</i> , &c.	Involved, velvety. Mycelium of branched necklace-like fibres. Receptacles somewhat forked, obtuse.
1756	...	...	...	V.	...	...	...	Leaves of <i>Citrus</i> ...	Scattered, bristle-like. Mycelium branched, necklace-like and netted. Receptacles elongated.
1757	...	S.A.	...	...	...	Q.	...	Leaves, &c. ...	Bristle-like. Receptacles elongated, pointed, and apex fringed.
1758	...	...	...	...	...	Q.	B.	<i>Xanthoxylon</i> ...	Mass of dark-brown branched threads, bearing receptacles, fleshy, club to horn shaped.
1759	...	...	...	V.	...	...	...	Branches and living leaves of <i>Bursaria spinosa</i>	Black, broadly expanded, separating as a pellicle; threads creeping, branched, sooty brown.
Schröd. Journ. III. 16 (1810).									
1760	...	...	...	V.	...	Q.	...	Ferns, &c. ...	Mycelium expanded, cloth-like, fibres very thin, elongated. Receptacles oblong.
1761	W.A.	...	...	V.	N.S.W.	...	...	Branches, &c. ...	Spongy, black, woolly tufts, filaments necklace-like or even. Receptacles elliptic.
1762	...	...	...	...	...	Q.	B.	Ferns, &c. ...	Mycelium dense, black, forming cloth-like coating on leaves. Receptacles curved.
CORDA ANL. 142 (1842).									
Ard. 272 (1837).— <i>Schizothyrium</i> .									
1763	...	...	...	V.	...	...	...	Dead leaves of <i>Eucalyptus</i>	On both surfaces, spots circular, reddish brown. Receptacles gregarious, minute, linear or run together, black.
1764	...	...	...	...	...	Q.	...	Leathery leaves, living and languid	Spots black, circular or run together. Receptacles gregarious, elongated, linear, flexuous, black.
Am. 101 (1813).— <i>Solenarium, Hysterium</i> .									
1765	...	...	T.	...	...	...	...	Rotten wood ...	Receptacle-bearing layer spread out irregularly, dark brown. Receptacles diverging from centre in a radiate manner for an inch or two.
1766	...	...	T.	...	...	...	...	Leaves of <i>Cyathodes straminea</i>	Receptacles elliptical, obtuse, opening slowly.
Sci. Nat. III. 58 (1845).									
1767	...	...	...	...	...	Q.	...	Leaves of <i>Olea paniculata</i>	Receptacles gregarious, linear oblong or forked, black, receptacle-bearing layer obsolete.
1768	...	...	...	...	...	...	...	Leaves of <i>Eucalyptus pilularis</i>	Spots somewhat circular, black, distinctly defined, and for the most part on upper surface. Receptacles elongated, shining black.
Meckl. II. 4 (1790).— <i>Hysterographium</i> .									
1769	...	...	...	...	N.S.W.	...	B.	Bark ...	Receptacles scattered or gregarious, superficial, variable in form, mostly oblong, striate lengthwise, black.
Syll. II. 757 (1883).— <i>Tryblidium, Hysterium</i> .									
1770	...	...	...	...	...	Q.	...	Bark of trees ...	Bursting through, wavy or triangular, black, swollen lips, disc turning red.
Fung. Arg. Pug. IV. 191 (1882).									
1771	...	...	...	...	...	Q.	...	Bark of trees ...	Receptacles scattered, superficial, oblong to elongated, somewhat leathery, disc reddish to brown.
Handb. Austr. Fung. 409 (1892).— <i>Triblidiopsis, Tryblidium</i> .									
1772	...	...	...	V.	...	...	...	Bark ...	Tufts scattered, bursting through, black, hemispherical, leathery.
Icon. V. 34 (1842).— <i>Hysterium</i> .									
1773	W.A.	...	...	...	...	...	B.	Decorticated wood...	Receptacles superficial, on black spot-like crust, oblong, straight, black.
1774	...	...	...	V.	...	Q.	...	Rotten wood ...	Receptacles elongated, linear, straight, parallel, black, with narrow fissure, lips stout.
1775	...	...	...	V.	...	...	B.	Wood ...	Receptacles bursting through, at length superficial, oblong to linear, in parallel lines, black.

GENERAL CLASSIFICATION OF DISCOMYCETES.

GROUP V.—DISCOMYCETES, FRIES.

ARRANGEMENT OF ORDERS (10).

32. CYTTARIACEÆ—Receptacle sub-globose.
 33. HELVELLACEÆ—Receptacle vertical, stalked; cap club or mitre shaped.
 34. PEZIZACEÆ—Receptacle cup shaped or disc shaped, sessile or stalked, fleshy or waxy.
 35. ASCOBOLACEÆ—Receptacle plane or convex, sessile or sub-sessile, fleshy.
 36. DERMATACEÆ—Receptacle concave or plane, sessile or somewhat stalked, corky, leathery, or horny.
 37. BULGARIACEÆ—Receptacle top, cup, or disc shaped, gelatinous, becoming cartilaginous or horny.
 38. STICTACEÆ—Receptacle immersed, usually bright coloured, waxy.
 39. PHACIDIACEÆ—Receptacle immersed, usually blackish, waxy.
 40. PATELLARIACEÆ—Receptacle superficial, often blackish, leathery to horny.
 41. GYMNOASCEÆ—No proper receptacle.

Genus (1)—
 237. *Cyttaria*, Berk.

ORDER XXXII.—CYTTARIACEÆ, LEV.

Genera (5)—
 238. *Morehella*, Linn.
 239. *Helvella*, Linn.

ORDER XXXIII.—HELVELLACEÆ, SCHW.

240. <i>Mitrula</i> , Fries.	242. <i>Geoglossum</i> , Pers.
241. <i>Leotia</i> , Fries.	

Genera (23)—
 243. *Rhizina*, Fries.
 244. *Geopyxis*, Pers.
 245. *Peziza*, Linn.
 246. *Otidea*, Pers.
 247. *Discina*, Fries.
 248. *Pyronema*, Carus.
 249. *Humaria*, Fries.
 250. *Phyllipsia*, Berk.

ORDER XXXIV.—PEZIZACEÆ, FRIES.

251. <i>Sarcoscypha</i> , Fries.	259. <i>Mollisia</i> , Fries.
252. <i>Trichoscypha</i> , Cooke.	260. <i>Tapesia</i> , Pers.
253. <i>Lachnea</i> , Fries.	261. <i>Trichopeziza</i> , Fekl.
254. <i>Ciboria</i> , Fekl.	262. <i>Dasyscypha</i> , Fries.
255. <i>Helotium</i> , Fries.	263. <i>Phaeopeziza</i> , Sacc.
256. <i>Phialea</i> , Fries.	264. <i>Belonidium</i> , Mont. and Dur.
257. <i>Pseudohelotium</i> , Fekl.	265. <i>Erinella</i> , Sacc.
258. <i>Chlorosplenium</i> , Fries.	

Genus (1)—
 266. *Ascobolus*, Pers.

ORDER XXXV.—ASCOBOLACEÆ, BOND.

Genera (3)—
 267. *Urnula*, Fries.

ORDER XXXVI.—DERMATACEÆ, FRIES.

268. <i>Cenangium</i> , Fries.	269. <i>Tympanis</i> , Tode.
--------------------------------	------------------------------

Genera (3)—
 270. *Ombrophila*, Fries.

ORDER XXXVII.—BULGARIACEÆ, FRIES.

271. <i>Orbilia</i> , Fries.	272. <i>Coryne</i> , Tul.
------------------------------	---------------------------

Genus (1)—
 273. *Stictis*, Pers.

ORDER XXXVIII.—STICTACEÆ, FRIES.

Genera (3)—
 274. *Pseudopeziza*, Fekl.

ORDER XXXIX.—PHACIDIACEÆ, FRIES.

275. <i>Fabreia</i> , Sacc.	276. <i>Coccomyces</i> , De Not.
-----------------------------	----------------------------------

Genera (2)—
 277. *Patinella*, Sacc.

ORDER XL.—PATELLARIACEÆ, FRIES.

278. <i>Karschia</i> , Kærnb.

Genus (1)—
 279. *Exoascus*, Fekl.

ORDER XLI.—GYMNOASCEÆ, BAR.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
GROUP V.—DISCOMYCETES.—FRIES,					
ORDER XXXII.—CYTTARIACEÆ,					
237. CYTTARIA.—Berk.,					
1776	1352	VIII. 1	<i>C. Gunnii</i>	Berk., Hook., Lond. Journ. 576 (1848)	Gunn's cyttaria

ORDER XXXIII.—HELVELLACEÆ,

238. MORCHELLA.—Link,					
1777	1354	VIII. 10	<i>M. conica</i> ...	Pers., Champ. Com. 257 (1818)	Conical morel
1778	1355	" 13	<i>M. deliciosa</i>	Fries, S.M. II. 8 (1821)	Delicious morel
1779	1353	" 8	<i>M. esculenta</i>	Pers., Syn. 618 (1801)	Esculent morel
1780	1356	" 23	<i>M. semilibera</i>	D. C., Fl. Fr. II. 212 (1815)	Half-free morel
239. HELVELLA.—Linn.,					
1781	1357	VIII. 62	<i>H. monachella</i>	Fries, S.M. II. 18 (1821)	Monkish helvel
240. MITRULA.—Fries,					
1782	1358	VIII. 118	<i>M. vinosa</i>	Berk., Fl. Tasm. II. 273 (1860)	Wine-coloured mitrula
241. LEOTIA.—Fries,					
1783	1359	VIII. 2510	<i>L. lubrica</i>	Pers., Syn. 613 (1801)	Slimy leotia
242. GEOGLOSSUM.—Pers.,					
1784	1364	VIII. 141	<i>G. australe</i>	Berk., in Cooke Myco. 6 (1879)	Southern geoglossum
1785	1363	" 141	<i>G. glabrum</i>	Pers., Syn. 608 (1801)	Smooth geoglossum
1786	1360	" 150	<i>G. hirsutum</i>	Pers., Syn. 608 (1801)	Hairy geoglossum
1787	1362	" 138	<i>G. Muelleri</i>	Cooke, Myco. 4 (1879)	Mueller's geoglossum
1788	1365	" 145	<i>G. nigrinum</i>	Cooke, Myco. 205 (1879)	Black geoglossum
1789	1366	" 147	<i>G. Peckianum</i>	Cooke, Myco. 5 (1879)	Peck's geoglossum
1790	1361	" 149	<i>G. Walteri</i>	Berk., in Cooke Myco. 4 (1879)	Walter's geoglossum

ORDER XXXIV.—PEZIZACEÆ, FRIES,

243. RHIZINA.—Fries,					
1791	1367	VIII. 182	<i>R. ferruginea</i>	Phil., Grev. XVI. 74 (1888)	Rusty rhizina
244. GEOPYXIS.—Pers.,					
1792	1368	VIII. 210	<i>G. aluticolor</i>	Berk., Linn. Journ. XIII. 176 (1873)	Tan-coloured geopyxis

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
EPICR. I. (1836).									
LEV. CONSID. MYC. 117 (1846).									
Linn. Trans. XIX. 37 (1841).									
1776	...	...	T.	V.	...	...	...	Living branches of <i>Fagus Cunninghamii</i>	Gregarious, globose or pear shaped, at length hollow, tapering below, without distinct stem.
SCHWARTZ, SUMM. VEG. SCAND. (1814).									
Berl. Mag. III. 41 (1809).—Phallus, Helvella.									
1777	...	S.A.	T.	V.	N.S.W.	...	B.	Shady places	... Oblong conic, attached at base, brown, bay to black. Stem whitish, cylindrical. <i>Edible</i> .
1778	...	...	...	V.	...	...	B.	Grassy places	... Rather cylindrical, acute, livid yellow. Stem short. <i>Edible</i> .
1779	...	...	...	...	...	...	B.	Moist places	... Round or ovate, attached at base to stem, dingy yellow. Stem white, inflated, faint odour. <i>Edible</i> .
1780	...	...	...	V.	N.S.W.	...	B.	Grassy places	... Conical, free to middle, yellowish to dirty tawny when dry. Stem whitish. <i>Edible</i> .
Sp. Pl., 1649 (1763).—Phallus, Boletus.									
1781	...	...	T.	...	...	...	B.	Woods, on sandy ground	Bent downwards, lobed, attached at base, somewhat bay brown. Stem hollow, white.
S.M. I. 491 (1822).									
1782	...	...	T.	...	...	...	...	Rotten wood	... Vinous purple, slender, linear to club shaped. Stem thread-like, straight.
S.M. II. 29 (1821).—Elvela.									
1783	...	...	T.	V.	...	...	B.	Moist ground	... Gregarious, jelly-like, swollen, greenish yellow. Stem hollow, yellow.
Obs. I. 11 (1795).—Clavaria.									
1784	...	...	T.	V.	...	...	...	Among moss	... Smooth, dry, brown to black, clubs compressed, almost distinct from scaly stem.
1785	...	...	...	V.	...	Q.	B.	Grassy places	... Somewhat gregarious, smooth, dry, blackish. Stem slender, crooked, scaly.
1786	...	...	...	V.	...	...	B.	Among grass	... Hairy, black. Club often elongated, compressed. Stem erect, cylindrical.
1787	...	...	...	V.	...	...	...	Grassy places	... Smooth, rather viscid, blackening. Club compressed, equal in length to stem.
1788	...	...	...	V.	...	...	...	Grassy places	... Tufted, fragile, black, hollow. Clubs rather compressed, equal in length to slender stem.
1789	...	...	...	...	...	...	...	Moist ground	... Smooth, somewhat viscid, blackening.
1790	...	...	...	V.	...	...	...	Stems of <i>Dicksonia</i>	... Hairy, dark brown, blackening. Clubs spoon shaped, compressed. Stem slender.
S.M. II. 38 (1821).									
Obs. Myc. I. 161 (1815).									
1791	...	...	...	V.	...	...	...	Rotten wood	... Circular, sessile, concave, rigid when dry, tough and gelatinous when moist.
M.E. I. 42 (1822).									
1792	...	...	...	...	N.S.W.	Q.	...	Wood	... Tan coloured, stalked, funnel shaped, folded at base. Stem dilated at base.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
245. PEZIZA.— <i>Linn.</i> , Sp. Pl. (1753).— <i>Ciboria</i> . <i>Dasycephala</i> . <i>Discina</i> . <i>Geopyxis</i> . <i>Humaria</i> ,					
1793	...	VIII. 343	<i>P. applanata</i> ...	<i>Fries</i> , S.M. II. 64 (1821) ...	Flattened peziza ...
1794	1369	„ 253	<i>P. aurantia</i> ...	<i>Pers.</i> , Obs. II. 76 (1796) ...	Orange-coloured peziza ...
1795	1371	„ 293	<i>P. badia</i> ...	<i>Pers.</i> , Obs. II. 78 (1796) ...	Chestnut-brown peziza ...
1796	1375	„ 341	<i>P. brunneo-atra</i> ...	<i>Desm.</i> , Des. Esp. Nouv. 9 (1836) ...	Dark-brown peziza ...
1797	1372	„ 307	<i>P. cochleata</i> ...	<i>Linn.</i> , Sp. Pl. 1625 (1753) ...	Cochleate peziza ...
1798	1370	„ 279	<i>P. Drummondii</i> ...	<i>Berk.</i> , Hook., Journ. 71 (1845) ...	Drummond's peziza ...
1799	...	„ 306	<i>P. funerata</i> ...	<i>Cooke</i> , Myco. (1879) ...	Funereal peziza ...
1800	1374	„ 335	<i>P. Saccardiana</i> ...	<i>Cooke</i> , Myco. 174 (1879) ...	Saccardo's peziza ...
1801	1373	„ 297	<i>P. vesiculosa</i> ...	<i>Bull.</i> , Champ. 457 (1812) ...	Swollen peziza ...
246. OTIDEA.— <i>Pers.</i> ,					
1802	...	VIII. 354	<i>O. apophysata</i> ...	<i>Cooke and Phil.</i> , Grev. V. 60 (1876) ...	Branching otidea ...
1803	1377	X. 4473	<i>O. darjeelensis</i> ...	<i>Berk.</i> , Hook., Journ. 202 (1851) ...	Darjeeling otidea ...
1804	1378	VIII. 358	<i>O. hirneoloides</i> ...	<i>Sacc.</i> , Syll. VIII. 96 (1889) ...	Hirneola-like otidea ...
1805	1379	„ 362	<i>O. phlebophora</i> ...	<i>Sacc.</i> , Syll. VIII. 97 (1889) ...	Veined otidea ...
247. DISCINA.— <i>Fries</i> ,					
1806	1383	...	<i>D. australica</i> ...	<i>Cooke</i> , Handb. Austr. Fung. 255 (1892) ...	Australian discina ...
1807	1381	VIII. 377	<i>D. lumbricalis</i> ...	<i>Sacc.</i> , Syll. VIII. 101 (1889) ...	Worm-like discina ...
1808	1380	„ 373	<i>D. repanda</i> ...	<i>Sacc.</i> , Syll. VIII. 100 (1889) ...	Repand discina ...
1809	1382	„ 391	<i>D. venosa</i> ...	<i>Sacc.</i> , Syll. VIII. 104 (1889) ...	Veined discina ...
248. PYRONEMA.— <i>Carus</i> ,					
1810	1384	VIII. 401	<i>P. melaloma</i> ...	<i>Fekl.</i> , Symb. Myc. 319 (1869) ...	Black-bordered pyronema ...
1811	1385	„ 400	<i>P. omphalodes</i> ...	<i>Fekl.</i> , Symb. Myc. 319 (1869) ...	Navel-like pyronema ...
249. HUMARIA.— <i>Fries</i> ,					
1812	1393	VIII. 506	<i>H. carbonigena</i> ...	<i>Sacc.</i> , Syll. VIII. 130 (1889) ...	Charcoal-growing humaria ...
1813	1395	„ 520	<i>H. fusispora</i> ...	<i>Sacc.</i> , Syll. VIII. 133 (1889) ...	Spindle-spored humaria ...
1814	1388	„ 431	<i>H. globifera</i> ...	<i>Cooke</i> , Myco. (1879) ...	Globose-spored humaria ...
1814	...	„ 503	<i>H. granulata</i> ...	<i>Bull.</i> , Champ. 258 (1791) ...	Granulated humaria ...
bis					
1815	1392	„ 481	<i>H. Hartmanni</i> ...	<i>Phil.</i> , Grev. XVI. 5 (1887) ...	Hartmann's humaria ...
1816	1387	„ 424	<i>H. militina</i> ...	<i>Cooke</i> , Handb. Austr. Fung. 256 (1892) ...	Crimson humaria ...
1817	1386	„ 416	<i>H. miniata</i> ...	<i>Cooke</i> , Handb. Austr. Fung. 256 (1892) ...	Scarlet humaria ...
1818	1391	„ 455	<i>H. Muellerei</i> ...	<i>Berk.</i> , Linn. Journ. XIII. 176 (1873) ...	Mueller's humaria ...
1819	1390	„ 443	<i>H. recurva</i> ...	<i>Cooke</i> , Handb. Austr. Fung. 257 (1892) ...	Recurved humaria ...
1820	1394	„ 518	<i>H. rutilans</i> ...	<i>Sacc.</i> , Syll. VIII. 133 (1889) ...	Reddish humaria ...
1821	...	„ 588	<i>H. scatigena</i> ...	<i>Sacc.</i> , Syll. VIII. 147 (1889) ...	Springing humaria ...
1822	1396	„ 577	<i>H. tenacella</i> ...	<i>Sacc.</i> , Syll. VIII. 145 (1889) ...	Toughish humaria ...
1823	1397	„ 569	<i>H. Thozetii</i> ...	<i>Sacc.</i> , Syll. VIII. 144 (1889) ...	Thozet's humaria ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Lachnea, Mollisia, Otidea, Patinella, Phialea, Pseudohelotium, Pyronema, Sarcoscypha, Urnula.									
1793	...	...	...	V.	...	...	...	Moist places	Sessile, depressed, reddish; disc at length somewhat wrinkled, delicately frosted, fleshy.
1794	...	...	T.	V.	N.S.W.	...	B.	Ground	Gregarious, almost sessile, irregular, oblique, orange; base prolonged into short stem.
1795	W.A.	S.A.	...	...	...	...	B.	Moist ground	Nearly sessile, entire, dark brown, fleshy base often passing into short stem.
1796	...	...	...	V.	...	...	B.	Ground	Sessile, solitary, largish, entire, fleshy, fragile, smooth, brownish black or dark brown.
1797	...	S.A.	T.	V.	...	Q.	B.	Ground	Often densely tufted and much twisted, sessile, large, fleshy, umber.
1798	W.A.	...	...	V.	...	...	...	Ground	Cup shaped, sessile, medium sized, bay brown.
1799	...	S.A.	...	...	...	...	...	Immersed in sand	Bell shaped, brown, margin reflexed, somewhat lobed, thin, fragile.
1800	...	...	...	V.	...	...	...	Moist ground	Sessile, fleshy, fragile, concave, flesh red, margin often torn.
1801	...	S.A.	...	V.	...	...	B.	Ground, manure heaps, &c.	Large, entire, sessile, at first globose or top shaped, then bell shaped; base of cup very fleshy.
M.E. I. 229 (1822).—Peziza.									
1802	...	...	...	...	...	Q.	B.	Moist places	Tufted or gregarious, sessile, lobed, margin indented, umber brown; paraphyses peculiarly branched.
1803	...	...	...	V.	...	...	...	Ground	Expanded, somewhat cochleate, usually elongated on one side, umber.
1804	...	...	...	V.	...	Q.	...	Rotten wood	Sessile or very shortly stalked, red, white beneath.
1805	...	...	...	V.	...	...	B.	Ground	Cup shaped, oblique, ochrey yellow, finely powdery, with short stem-like base veined and ribbed.
S.V. 348 (1849).—Peziza.									
1806	...	...	...	V.	...	...	...	Ground	Large, cup shaped, then expanded, smooth, ochrey, tapering downwards into short thick rooting stem.
1807	...	...	...	V.	...	...	...	Ground	Large, cup shaped, at length expanded, internally pale brown, externally nearly smooth or mealy.
1808	...	...	...	V.	N.S.W.	Q.	B.	Rotten trunks and ground	Solitary or tufted, large, ineised, repand or bent backwards, internally brown, externally whitish.
1809	...	...	...	V.	...	...	B.	Ground	Sessile or somewhat stalked, umber brown, externally whitish, rough with ribbed veins, strong nitrous odour.
Nov. Aet. Cur. XVII. 370 (1835).									
1810	W.A.	...	...	V.	...	...	B.	Burnt ground	Sessile, crowded, dingy orange, margin with delicate black hairs or prominent cells.
1811	...	...	...	V.	...	...	B.	Burnt ground and cinder heaps	Sessile, crowded, often running together, minute, orange red or orange yellow.
S.M. II. 42 (1822).—Peziza.									
1812	...	...	T.	V.	...	...	B.	Charred ground	Gregarious, orange yellow, sessile, flexuous, slightly granular, margin wavy.
1813	...	...	T.	...	...	...	B.	Charred and heathy ground	Gregarious, rather crowded, sessile, hemispherical, yellow, downy; spores spindle shaped.
1814	...	...	...	V.	...	...	...	Sandy soil	Yellow, saucer shaped, margin turned in, sometimes lobed.
1814 bis	...	...	...	...	...	Q.	B.	Cow dung	Sessile, scattered or crowded, margin thick externally, tawny brown, and coarsely granular.
1815	...	...	...	...	...	Q.	...	Decayed branches	Gregarious, sessile or somewhat stalked, concave, margin splitting, disc pale crimson.
1816	...	...	...	V.	...	...	...	Sandy ground	Sessile, scattered, crimson, flattened, margin paler beneath, free.
1817	...	...	...	V.	...	...	B.	Among moss	Fleshy, firm, pitcher to plate like, scarlet.
1818	...	...	T.	V.	...	...	...	Ground	Scattered, sessile, cups irregular, marginate delicately downy, externally, disc crimson.
1819	...	...	T.	...	...	...	...	Ground	Nearly sessile, wavy, convex, recurved, bay brown, smooth.
1820	W.A.	S.A.	...	...	...	...	B.	Ground among moss	Gregarious, nearly sessile, bell or beaker shaped; disc orange yellow, externally paler, and slightly downy.
1821	...	...	...	...	...	Q.	...	Dung	Hemispherical, dark-wine colour, somewhat green when fresh, externally mealy white.
1822	...	...	...	V.	...	...	...	Ground	Sessile, slightly concave, smooth, umber brown, margin entire, flesh firm.
1823	...	...	...	...	N.S.W.	Q.	...	On <i>Nepenthes</i>	Dish shaped, fleshy, brown.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
250. PHILLIPSIA.—Berk.,					
1824	1399	VIII. 608	<i>P. polyporoides</i> ...	Berk., Linn. Journ. XVIII. 388 (1881) ...	Polyporous-like phillipsia ...
1825	1398	„ 607	<i>P. subpurpurea</i> ...	Berk. and Br., Linn. Soc. N.S.W. 88 (1880) ...	Purplish phillipsia ...
251. SARCOZYPHA.—Fries,					
1826	1403	VIII. 638	<i>S. bulbosa</i> ...	Cooke, Handb. Austr. Fung. 259 (1892) ...	Bulbous sarcosypha ...
1827	1400	„ 618	<i>S. coccinea</i> ...	Sacc. Syll. VIII. 154 (1889) ...	Scarlet sarcosypha ...
1828	1401	„ 620	<i>S. lepida</i> ...	Sacc. Syll. VIII. 154 (1889) ...	Handsome sarcosypha ...
1829	1404	„ 657	<i>S. melastoma</i> ...	Cooke, Handb. Austr. Fung. 259 (1892) ...	Black-mouthed sarcosypha ...
1830	1402	„ 630	<i>S. rhenana</i> ...	Sacc. Syll. VIII. 157 (1889) ...	Woolly sarcosypha ...
252. TRICHOSCYPHA.—Cooke,					
1831	1405	VIII. 652	<i>T. lindsi</i> ...	Sacc. Syll. VIII. 161 (1889) ...	Lind's trichoscypha ...
1832	1406	„ 647	<i>T. tricholoma</i> ...	Sacc. Syll. VIII. 160 (1889) ...	Hairy-edged trichoscypha ...
253. LACHNEA.—Fries,					
1833	1415	VIII. 733	<i>L. alpina</i> ...	Sacc. Syll. VIII. 180 (1889) ...	Alpine lachnea ...
1834	1410	„ 699	<i>L. badio-berbis</i> ...	Sacc. Syll. VIII. 173 (1889) ...	Bay-bearded lachnea ...
1835	1408	„ 772	<i>L. confusa</i> ...	Cooke, Handb. Austr. Fung. 260 (1892) ...	Confused lachnea ...
1836	1419	„ 735	<i>L. coprogena</i> ...	Sacc. Syll. VIII. 181 (1892) ...	Dung-borne lachnea ...
1837	1414	„ 730	<i>L. dalmeniensis</i> ...	Phil., Disc. 227 (1887) ...	Dalmeny lachnea ...
1838	1420	„ 741	<i>L. Erinaceus</i> ...	Sacc. Syll. VIII. 182 (1889) ...	Hedgehog lachnea ...
1839	1413	„ 705	<i>L. hirta</i> ...	Gill., Champ. 75 (1879) ...	Hairy lachnea ...
1840	1416	„ 722	<i>L. lusatie</i> ...	Sacc. Syll. VIII. 178 (1889) ...	Lusatian lachnea ...
1841	1411	„ 700	<i>L. margaritacea</i> ...	Sacc. Syll. VIII. 173 (1889) ...	Pearly lachnea ...
1842	1417	„ 725	<i>L. scubalouta</i> ...	Sacc. Syll. VIII. 179 (1889) ...	Refuse lachnea ...
1843	1409	„ 698	<i>L. scutellata</i> ...	Gill., Champ. 75 (1879) ...	Saucer-shaped lachnea ...
1844	1421	„ 744	<i>L. stercorea</i> ...	Gill., Champ. 76 (1879) ...	Dung lachnea ...
1845	1418	„ 728	<i>L. theleboloides</i> ...	Gill., Champ. 74 (1879) ...	Thelebolus-like lachnea ...
1846	1412	„ 701	<i>L. umbrata</i> ...	Phil., Disc. 222 (1887) ...	Shaded lachnea ...
1846	„	„	<i>L. umbrata</i> , var. <i>pallida</i> ...	Rehm., Asco. No. 456 (1873) ...	Pale lachnea ...
1847	1407	„ 687	<i>L. vinoso-brunnea</i> ...	Sacc. Syll. VIII. 171 (1889) ...	Vinous-brown lachnea ...
254. CIBORIA.—Fekl.,					
1848	1423	VIII. 829	<i>C. firma</i> ...	Fekl., Sym. Myc. 312 (1869) ...	Firm eiboria ...
255. HELOTIUM.—Fries,					
1849	1425	VIII. 910	<i>H. citrinum</i> ...	Fries, S.V. 355 (1849) ...	Lemon-yellow helotium ...
1850	1426	„ 914	<i>H. claro-flavum</i> ...	Berk., Outl. 372 (1860) ...	Light-yellow helotium ...
1851	1429	„ 925	<i>H. epiphyllum</i> ...	Fries, S.V. 356 (1849) ...	Leaf-growing helotium ...
1852	1427	„ 918	<i>H. gratum</i> ...	Cooke, Austr. Fung. 51 (1883) ...	Agreeable helotium ...
1853	1424	„ 876	<i>H. nigripes</i> ...	Fries, S.V. 356 (1849) ...	Black-stalked helotium ...
1854	1428	„ 1028	<i>H. pateræforme</i> ...	Cooke, Austr. Fung. 51 (1883) ...	Saucer-shaped helotium ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Linn. Journ. XVIII. 388 (1881).—Peziza.									
1824	...	...	...	...	...	Q.	...	Dead stems of <i>Vitis</i>	Expanded, attached, thick, flesh colour.
1825	...	...	...	...	...	Q.	...	Wood ...	Plane, margin lobed, fixed at centre; disc purplish, brown when dry.
S.M. II. 78 (1822).—Peziza, Lachnea, Macropodia.									
1826	...	...	...	V.	...	...	B.	Ground in sandy soil	Hemispherical, turning ashy, minutely scaly, disc brown. Stem firm, tuberos at base.
1827	...	...	T.	...	...	...	B.	Rotten branches ...	Funnel shaped, externally whitish, downy, as well as stem, disc carmine, most handsome.
1828	..	S.A.	...	...	...	...	...	Ground ...	Funnel shaped, medium sized, with gradually tapering stem, disc crimson.
1829	..	...	...	...	...	Q.	B.	Old branches, &c. ...	Fleshy, almost globose, externally brick red, woolly, disc black. Stem short, with rooting black hairs.
1830	W.A.	...	...	...	...	...	...	Bare ground ...	Tufted, united in thick stem, whitish downy, nearly globose, margin turned in, disc orange.
Myc. 252 (1879).									
1831	...	...	...	...	...	Q.	...	Rotten wood ...	Bright red, cup shaped, externally with delicate bloom, tawny yellow. Stem tapering downwards.
1832	...	...	...	...	...	Q.	...	Rotten wood ...	Stalked, fleshy, hemispherical, top shaped, yellow, hairy edged. Stem smooth.
S.M. II. 77 (1822).—Peziza, Humaria, Sphaerospora.									
1833	...	...	...	V.	...	...	...	Cow dung ...	Gregarious, closed at first, then flattened, circular, margin orange yellow with jointed hairs.
1834	...	S.A.	...	...	N.S.W.	...	...	Rotten wood ...	Concave, disc vermilion, margin clad with very long hairs.
1835	...	...	...	...	...	Q.	...	Charcoal ...	Gregarious or crowded, sessile, almost spherical, brown, clad with short hairs.
1836	...	...	...	...	...	Q.	...	Dung ...	Nearly orange, invested with pale-bay obtuse hairs.
1837	...	...	...	V.	...	...	B.	Ground in shady woods	Sessile, fleshy, hemispherical, becoming expanded, bright yellow, fringed with long erect yellow hairs.
1838	...	...	...	...	...	Q.	...	Rotten wood ...	Gregarious, circular, depressed, ochrey white, externally beset with long bay-brown hairs.
1839	...	...	...	V.	...	...	B.	Ground among moss	Sessile, scattered, somewhat hemispherical, externally clad with brown hairs, disc scarlet.
1840	...	...	...	V.	...	...	...	Rotten wood ...	Gregarious, sessile, cup shaped, flattened at length, orange red, with erect brown hairs externally.
1841	...	...	...	V.	...	...	...	Rotten wood ...	Sessile, hemispherical, at length expanded, vermilion, rough externally, with short brown hairs.
1842	...	...	...	V.	...	...	...	Dung ...	Scattered, sessile, fleshy, hemispherical, thickly clad externally, with septate brown hairs.
1843	W.A.	...	T.	V.	N.S.W.	Q.	B.	Rotten wood ...	Gregarious, sessile, flattened, vermilion red, rough towards margin, with long straight black hairs.
1844	...	...	T.	V.	...	...	B.	Dung ...	Gregarious, sessile, concave, dingy red or tawny, beset with brown septate hairs.
1845	...	...	...	V.	...	...	B.	Earth, &c. ...	Gregarious, spherical, then open, externally whitish, clad with pale hairs, disc pale yellow.
1846	...	...	...	...	...	...	B.	...	...
1846a	...	...	...	V.	...	...	B.	Charred wood, &c. ...	Brownish flesh colour, with scattered obtuse hairs.
1847	...	...	...	...	...	Q.	...	Burnt ground ...	Sessile, hemispherical, flattened, vinous brown, rough, with short obtuse brown scattered hairs.
Sym. Myc. 311 (1869).—Peziza, Hymenocypha.									
1848	...	...	T.	...	...	...	B.	Rotting branches ...	Funnel shaped, then expanded, firm, pale brown. Stem long, tapering downwards, becoming blackish.
S.V. 354 (1849).—Peziza, Phialea.									
1849	...	...	T.	...	...	Q.	B.	Dead stumps and branches	Gregarious or crowded, shortly stalked or sessile, flattened, concave, lemon yellow, waxy.
1850	...	...	...	V.	...	...	B.	Decayed wood and fallen branches	Very minute, shortly stalked or sessile, clear yellow, smooth, margin somewhat lobed.
1851	...	...	...	...	...	Q.	B.	Dead leaves ...	Almost sessile, convex to plane, smooth, marginate, firm, pale ochrey.
1852	...	...	T.	...	...	...	...	Dead wood ...	Plane, transparent, marginate, shortly stalked, nearly orange. Stem paler, cylindrical.
1853	...	...	T.	...	...	...	...	Trunks and rotting leaves	Flattened, concave, pale, smooth, marginate. Stem longish, blackening.
1854	...	...	T.	...	...	...	...	Rotten wood ...	Ochrey, sessile, somewhat lobed, concave, somewhat wrinkled and delicately downy beneath.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
256. PHIALEA.—Fries,					
1855	1432	VIII. 1048	<i>P. Berggrenii</i> ...	<i>Sacc.</i> Syll. VIII. 254 (1889) ...	Berggren's phialea ...
1856	1434	„ 1104	<i>P. byssogena</i> ...	<i>Sacc.</i> Syll. VIII. 267 (1889) ...	Thread-borne phialca ...
1857	1433	„ 1102	<i>P. ccratina</i> ...	<i>Sacc.</i> Syll. VIII. 267 (1889) ...	Horny phialea ...
257. PSEUDOHELOTIUM.—Fekl., Symb. Myc. 298 (1869).—					
1858	1435	VIII. 1215	<i>P. hyalinum</i> ...	Fekl., Symb. Myc. 298 (1869) ...	Hyaline pseudohelotium ...
1859	1436	„ 1267	<i>P. ilicinolum</i> ...	<i>Sacc.</i> Syll. VIII. 304 (1889) ...	Holly-growing pseudohelotium ...
258. CHLOROSPLENIUM.—Fries,					
1860	1430	VIII. 1311	<i>C. æruginosum</i> ...	<i>De Not.</i> , Disc. 22 (1864) ...	Verdigris chlorosplenium ...
1861	1431	„ 1313	<i>C. omnivirens</i> ...	<i>Cooke</i> , Austr. Fung. 51 (1883) ...	All-green chlorosplenium ...
259. MOLLISIA.—Fries,					
1862	1437	VIII. 1393	<i>M. cinerea</i> ...	<i>Karst.</i> , M.F. I. 189 (1871) ...	Ash-coloured mollisia ...
260. TAPESIA.—Pers.,					
1863	1438	VIII. 1573	<i>T. epithepha</i> ...	<i>Sacc.</i> Syll. VIII. 381 (1889) ...	Woolly-base tapesia ...
261. TRICHOPEZIZA.—Fekl.,					
1864	1439	X. 4540	<i>T. Sphærula</i> ...	<i>Sacc.</i> , Hedw. 155 (1890) ...	Sphærula trichopeziza...
262. DASYSCEPHA.—Fries, S.M. II. 89 (1822).—					
1865	1443	VIII. 1924	<i>D. Eucalypti</i> ...	<i>Sacc.</i> Syll. VIII. 462 (1889) ...	Eucalypt dasyscypha ...
1866	1442	„ 1876	<i>D. glabrescens</i> ...	<i>Sacc.</i> Syll. VIII. 451 (1889) ...	Smooth dasyscypha ...
1867	1441	„ 1804	<i>D. lachnoderma</i> ...	<i>Rehm.</i> , Asco. No. 303 (1873) ...	Downy dasyscypha ...
1868	1444	„ 1938	<i>D. lanariceps</i> ...	<i>Sacc.</i> Syll. VIII. 465 (1889) ...	Woolly-capped dasyscypha ...
1869	1445	„ 1947	<i>D. terrestris</i> ...	<i>Sacc.</i> Syll. VIII. (1889) ...	Terrestrial dasyscypha ...
1870	1440	„ 1801	<i>D. virginea</i> ...	Fekl., Symb. Myc. 305 (1869) ...	Virgin dasyscypha ...
263. PILEOPEZIZA.—Sacc.,					
1871	1376	VIII. 1966	<i>P. apiculata</i> ...	<i>Sacc.</i> , Mich. I. 71 (1877) ...	Apiculate phæopeziza ...
264. BELONIDIUM.—Mont.					
1872	1446	VIII. 2064	<i>B. araucosum</i> ...	<i>Sacc.</i> Syll. VIII. 500 (1889) ...	Web-like belonidium ...
265. ERINELLA.—Sacc.					
1873	1447	...	<i>E. lutca</i> ...	<i>Phil.</i> , Grev. XIX. 61 (1891) ...	Yellow crinella ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Obs. II. 305 (1818).									
1855	...	...	...	V.	...	...	...	Rotting leaves	Pale, stalked, scattered, wine-glass shaped. Stem slender.
1856	...	...	T.	...	...	...	...	Wool	Ochrey, concave. Stem elongated, cylindrical, arising from interwoven radiating threads.
1857	...	...	T.	...	...	...	...	Leaves of Eucalypts	Top shaped, stalked, smooth, pale, horny brown, minute.
Peziza, Lachnea, Helotium, Lachnella, Mollisia.									
1858	...	...	T.	...	...	...	B.	Rotten trunks and inside bark	Minute, gregarious, sessile, globose, then expanded, transparent when moist, downy externally.
1859	...	...	...	V.	...	...	B.	Holly branches and lichen, <i>Myriangium</i> growing on Holly	Tufted, hemispherical, then expanded, externally dirty white, disc brown, purple, or rosy grey.
S.V. 356 (1849).—Peziza, Helotium.									
1860	...	...	...	V.	...	Q.	B.	Fallen wood	Shortly stalked or sessile, verdigris green, disc becoming whitish. Stem short, stout. Wood stained employed as green oak in manufacture of Tunbridge ware.
1861	...	...	T.	...	...	...	...	Rotten wood	Verdigris green, shortly stalked, rather top shaped.
S.M. II. 137 (1822).—Peziza.									
1862	...	...	T.	V.	...	...	B.	Decaying wood	Gregarious or scattered, sessile, soft, minute, saucer-like, ashy, with entire whitish margin.
M.E. I. 220 (1828).—Peziza, Lachnella.									
1863	...	...	T.	...	...	...	...	Leaves	Minute, white, hemispherical or almost globose, concave, arising from crisp interwoven threads.
Symb. Myc. 295 (1869).									
1864	...	S.A.	...	...	...	...	...	Dead bark of <i>Casuarina</i>	Scattered, minute, sessile, globose, bright sulphur yellow, sprinkled with rough hairs.
Peziza, Hymenoseypha, Lachnella, Helotium.									
1865	...	...	T.	V.	...	...	...	Leaves of <i>Eucalyptus</i> and <i>Casuarina</i>	Pale olive, plane, margin fringed with rigid dark-purple hairs. Stem cylindrical.
1866	...	...	...	V.	...	...	...	On <i>Rhipogonum</i>	Scattered, stalked, white, wine-glass shaped, at first shaggy, then naked, smooth.
1867	...	...	T.	...	...	Q.	...	Dead bark	Almost hemispherical, shortly stalked, externally snowy white and downy, vermillion within.
1868	...	...	...	V.	...	...	...	On <i>Rhipogonum</i>	Scattered, stalked, ochrey brown, top shaped, at length open, shaggy, sprinkled with purple granules.
1869	...	...	...	...	...	Q.	...	Bare ground	Small, stalked, horn colour, lurid, externally shaggy.
1870	...	...	T.	...	...	...	B.	Wood, bark, branches, &c.	Gregarious, stalked, white, hemispherical, with crowded spreading hair. Stem short.
Mich. I. 71 (1877).									
1871	...	...	...	V.	...	...	...	Bark	Sessile, saucer shaped, fleshy, rather tough, black, smooth, disc with margin entire.
and Dur., Fl. Alg. (1846).									
1872	...	...	T.	...	...	...	...	Wood	At first nearly globose, then hemispherical, externally web-like, arising from creeping threads.
Syll. VIII. (1889).									
1873	...	...	...	V.	...	...	...	Crevices of bark	Gregarious or scattered, shortly stalked, cup shaped, with short whitish hairs, becoming yellow, then yellowish brown.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
ORDER XXXV.—ASCOBOLACEÆ,					
266. ASCOBOLUS.—Pers.,					
1874	1452	VIII. 2161	<i>A. Archeri</i> ...	Berk., Fl. Tasm. II. 276 (1860) ...	Archer's ascobolus ...
1875	1449	„ 2149	<i>A. australis</i> ...	Berk., Linn. Journ. XVIII. 398 (1881) ...	Southern ascobolus ...
1876	1451	„ 2150	<i>A. Baileyi</i> ...	Berk. and Br., Linn. Trans. II. 69 (1883) ...	Bailey's ascobolus ...
1877	1448	„ 2143	<i>A. furfuraceus</i> ...	Pers., Tent. Disp. Meth. 25 (1797) ...	Scurfy ascobolus ...
1878	1450	...	<i>A. Phillipsii</i> ...	Berk., in Cooke's Handb. Austr. Fung. 268 (1892)	Phillips' ascobolus ...

ORDER XXXVI.—DERMATACEÆ,

267. URNULA.—Fries,					
1879	1453	VIII. 640 and 218	<i>U. campylospora</i> ...	Cooke, Handb. Austr. Fung. 268 (1892) ...	Curve-spored urnula ...
1880	1454	VIII. 331	<i>U. rhytidea</i> ...	Cooke, Austr. Fung. 52 (1883) ...	Wrinkled urnula ...
268. CENANGIUM.—Fries,					
1881	1455	VIII. 2323	<i>C. liehenoides</i> ...	Berk. and Br., Linn. Trans. I. 404 (1879) ...	Lichenoid cenangium ...
269. TYMPANIS.—Tode,					
1882	...	X. 4603	<i>T. Toomausis</i> ...	Berk. and Br., Linn. Trans. II. 222 (1887) ...	Tooma tympanis ...

ORDER XXXVII.—BULGARIACEÆ,

270. OMBROPHILA.—Fries,					
1883	1459	X. 4612	<i>O. bulgarioides</i> ...	Sacc., Myc. Austr. 14 (1890) ...	Bulgaria-like ombrophila ...
1884	1457	VIII. 2532	<i>O. radicata</i> ...	Phil., Grev. XVI. 33 (1887) ...	Rooting ombrophila ...
1885	1458	„ 2553	<i>O. terrestris</i> ...	Phil., Grev. XVI. 75 (1888) ...	Terrestrial ombrophila ...
1886	1460	...	<i>O. trachycarpa</i> ...	Phil., Grev. XIX. 61 (1891) ...	Rough-spored ombrophila ...
1887	1456	VIII. 2526	<i>O. violacea</i> ...	Fries, S.V.S. 357 (1849) ...	Violet ombrophila ...
1887a	1456A	„ „	<i>O. violacea</i> , var. <i>australis</i> ...	Cooke, Grev. VIII. 64 (1879) ...	Southern ombrophila ...
271. ORBILIA.—Fries,					
1888	1461	VIII. 2572	<i>O. chrysocoma</i> ...	Sacc., Syll. VIII. 621 (1889) ...	Golden-yellow orbilia ...
1889	1462	„ 2568	<i>O. decipiens</i> ...	Sacc., Syll. VIII. 623 (1889) ...	Deceptive orbilia ...
272. CORYNE.—Tul., Carp. III. 190 (1865).—					
1890	1463	VIII. 2647	<i>C. sareoides</i> ...	Tul., Carp. III. 190 (1865) ...	Flesh-like coryne ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.	
	W.A.	S.A.	T.	V.	N.S.W.	Q.				
BOND., MEM. ASCOB. 20 (1869).										
in Gmel. Syst. 1461 (1791).—Peziza.										
1874	...	...	T.	...	...	...	...	Charcoal	...	Wavy, sessile, vinous brown.
1875	...	...	..	...	...	Q.	...	Dung	...	Brown, cup shaped.
1876	...	...	...	...	...	Q.	...	Dung	...	Concave, at first ochrey, then vinous brown and flattened, slightly granulate externally.
1877	W.A.	...	...	...	...	..	B.	Old cow dung, &c....	...	Sessile, globose, then expanded; externally greenish yellow, mealy; disc slightly concave at first, yellowish green, turning blackish brown when old.
1878	...	...	...	...	...	Q.	...	Cow dung	...	Concave with elevated margin, externally wax colour, then tawny, disc ash coloured.

FRIES, S.V. 345 (1849).S.V.S. II. 364 (1849).—*Peziza*, *Macropodia*.

1879	...	...	...	...	...	Q.	...	Rotten wood	Funnel shaped, sooty or ashy, stalked, deeply wrinkled, margin incurved. Stem similarly coloured.
1880	...	...	...	...	N.S.W.	...	...	Ground	Sooty brown, nearly sessile, hemispherical, cut, undulately wrinkled, flesh olive.

S.M. II. 177 (1822).

1881	...	...	...	...	...	Q.	...	Trunks	Tufted, ashy, top shaped, stalked, invested with irregular ashy warts, disc red brown.
------	-----	-----	-----	-----	-----	----	-----	--------	--

Meekl. I. 23 (1790).

1882	...	...	...	...	...	Q.	...	Fruit of <i>Eauskia</i>	Gregarious, at first mealy, sphaeria shaped.
------	-----	-----	-----	-----	-----	----	-----	-------------------------	--

FRIES, S.V. 345 (1849).S.V.S. 357 (1849).—*Peziza*.

1883	...	...	...	...	...	Q.	...	Rotten wood	Clustered, gelatinous, then hard, at first sessile, then shortly stalked, externally bright ochrey yellow, disc reddish brown.
1884	...	...	...	V.	...	...	...	Swampy places	Solitary or tufted, stalked, rather gelatinous. Stem elongated, rooting; disc depressed, wrinkled, liver colour.
1885	...	...	...	V.	...	...	...	Ground	Circular, sessile, gelatinous, concave or flattened, amber brown, a little paler externally.
1886	...	...	...	V.	...	...	...	Sandy ground among mosses	Somewhat gregarious, sessile, concave, externally wrinkled horizontally, firm, dark red brown.
1887	...	...	...	V.	...	...	...	Trunks	Gregarious or scattered, finally distinctly stalked, violet. Stem obconic, short.
1887A	...	...	...	V.	...	...	...	Branches. &c., in swampy places	Stem longer, flexuous, more ash coloured than type.

S.V. 357 (1849).—*Peziza*, *Calloria*.

1888	...	...	...	V.	...	...	...	Wood	Gregarious, almost globose at first, soon flattened, and rather jelly-like, golden yellow, horny when dry.
1889	...	...	...	...	...	Q.	...	Old rope	Gregarious or scattered, sessile; disc pale-flesh colour, orange red, or pale brown; externally granulose.

Peziza, *Ombrophila*, *Lichen*, *Bulgaria*, *Helvella*, *Tremella*.

1890	...	...	T.	V.	...	...	B.	Trunks and branches of trees	Tufted, sessile or somewhat stalked, firm, fleshy red, veined below, disc hollowed out.
------	-----	-----	----	----	-----	-----	----	------------------------------	---

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
---------	-----------------	--------------------	------------------	---------------------	---------------

ORDER XXXVIII.—STICTACEÆ,

273. STICTIS.—Pers., Obs. II. 73 (1796).—

1891	1465	X. 4635	<i>S. emarginata</i>	Cooke and Mass., Grev. XVIII. 7 (1889)...	Emarginate stictis
1892	1464	VIII. 2795	<i>S. radiata</i>	Pers., Obs. II. 73 (1796)	Radiating stictis
1892a	...	" ..	<i>S. radiata</i> , var. <i>brachyspora</i>	Sacc. and Berl., Rev. Myc. (1885)	Short-spored stictis

ORDER XXXIX.—PHACIDIACEÆ,

274. PSEUDOPEZIZA.—Fekl., Symb.

1893	...	VIII. 2976	<i>P. Cerastiorum</i>	<i>Fekl.</i> , Symb. Myc. 291 (1869)	Chickweed pseudopeziza
1894	1467	" 2971	<i>P. Medicaginis</i>	<i>Sacc.</i> , Fung. Ard. No. 90 (1888)	Medicago pseudopeziza
1895	1466	" 2970	<i>P. Trifolii</i>	<i>Fekl.</i> , Symb. Myc. 291 (1869)	Trifolium pseudopeziza

275. FABRÆA.—Sacc.,

1896	1468	X. 4651	<i>F. rhytismoides</i>	<i>Sacc.</i> , Syll. X. 50 (1892)	Rhytisma-like fabræa
------	------	---------	-------------------------------	--	-----------------------------

276. COCCOMYCES.—De Not.,

1897	1469	...	<i>C. delta</i>	<i>Cooke</i> , Handb. Austr. Fung. 272 (1892)	Deltoid coccomyces
------	------	-----	------------------------	--	---------------------------

ORDER XL.—PATELLARIACEÆ,

277. PATINELLA.—Sacc.,

1898	1470	VIII. 3162	<i>P. tasmanica</i>	<i>Sacc.</i> , Syll. VIII. 770 (1889)	Tasmanian patinella
1899	1471	" 3178	<i>P. Adamsoni</i>	<i>Sacc.</i> , Syll. VIII. 772 (1889)	Adamson's patinella

278. KARSCHIA.—Kærh.,

1900	1472	VIII. 3200	<i>K. lignyata</i>	<i>Sacc.</i> , Syll. VIII. 779 (1889)	Wood-growing karschia
------	------	------------	---------------------------	--	------------------------------

ORDER XLI.—GYMNOASCEÆ,

279. EXOASCUS.—Fekl., Enum. Fung.

1901	1473	VIII. 3341	<i>E. deformans</i>	<i>Fekl.</i> , Symb. Myc. 252 (1869)	Deforming exoascus (Peach-leaf curl)
------	------	------------	----------------------------	---	--

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
FRIES, PL. HOM. 86 (1825).									
Peziza, Lycoperdon, Sphaerobolus, Schmitzonia.									
1891	...	...	...	V.	...	...	...	Leaves of Eucalypts	Very minute, gregarious, immersed, bursting through, pierced at apex. Gregarious or scattered, flesh coloured or yellowish, deeply immersed; margin four to six rayed, white, scurfy. Margin narrower than type, and disc ashy grey to violet.
1892	...	...	T.	...	...	Q.	B.	Wood and branches	
1892a	...	...	...	...	...	Q.	...	Rotten stem	

FRIES, S.M. II. 317 (1822).

Myc. 290 (1869).—Phacidium, Ascobolus, Peziza, Mollisia.

1893	...	...	...	...	N.S.W.	...	B.	Leaves and more rarely calyx of <i>Cerastium vulgatum</i>	Gregarious, sessile, minute, round, smooth, at first white with reddish-grey rim, at last buff with dark-brown rim.
1894	...	...	...	V.	...	...	...	Leaves of <i>Medicago</i>	Scattered, minute, soon flattened, ochrey brown, originating on yellowish spots, then girt by three to four toothed skin.
1895	...	...	...	V.	...	...	B.	Living but languishing clover leaves	Gregarious, sessile, minute, circular, plane, smooth, smoky yellow; margin thin, torn.

Mich. II. 331 (1881).—Phacidium, Pseudopeziza.

1896	...	...	...	V.	...	...	...	Living leaves of <i>Cotula</i>	Cups clustered together, usually six to eight, minute, externally dark brown; disc closing in drying, blackening, and then resembling <i>Rhytisma</i> .
------	-----	-----	-----	----	-----	-----	-----	--------------------------------	---

Mem. II. 38 (1847).—Phacidium.

1897	...	...	...	V.	...	...	...	Leaves of Eucalypts	Innate, three-angled, with three elevated joints, opening in three valves, disc brown.
------	-----	-----	-----	----	-----	-----	-----	---------------------	--

FRIES, S.V. 345 (1849).

Grev. IV. 22 (1875).—Patellaria, Peziza.

1898	...	...	T.	...	...	...	...	Dead wood	Small, sessile, cups concave, then plane; disc reddish brown, then black.
1899	...	...	...	V.	...	...	...	Branches of Eucalypts	Circular, plane, cups with distinct margin, quite black.

Parerg. 459 (1865).—Patellaria.

1900	...	...	...	V.	...	...	B.	Rotten wood	Scattered or slightly gregarious, sessile, saucer shaped, horny when dry, externally dark red; disc concave, quite black.
------	-----	-----	-----	----	-----	-----	----	-------------	---

BAR. BOT. ZEIT. 158 (1872).

Nass. 29 (1860).—Taphrina, Aseomyces.

1901	...	S.A.	...	V.	N.S.W.	Q.	B.	Peach, &c., leaves.	On under surface of leaves causing blisters, and covered with a whitish bloom.
------	-----	------	-----	----	--------	----	----	---------------------	--

GENERAL CLASSIFICATION OF TUBEROIDES.

GROUP VI.—TUBEROIDES, VITT.

ARRANGEMENT OF ORDERS (3).

- 42. ELAPHOMYCETACEÆ—Gleba or spore-bearing tissue traversed by silky filaments.
 - 43. TUBERACEÆ—Gleba traversed by branched filaments, or with cavities.
 - 44. ENDOGONACEÆ—Gleba destitute of internal cavities, continuous.
-

ORDER XLII.—ELAPHOMYCETACEÆ, TUL.

Genus (1)—

- 280. Elaphomyces, Nees.

ORDER XLIII.—TUBERACEÆ, FRIES.

Genus (1)—

- 281. Stephensia, Tul.

ORDER XLIV.—ENDOGONACEÆ, FRIES

Genus (1)—

- 282. Endogone, Link.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
---------	-----------------	--------------------	------------------	---------------------	---------------

GROUP VI.—TUBEROIDES.—

ORDER XLII.—ELAPHOMYCETACEÆ,

280. ELAPHOMYCES.—Nees,

1902	1349	VIII. 3481	E. Leveillei	Tul., Ann. Sci. Nat. 2 Ser. XVI. 21 (1841) ...	Leveille's elaphomyces ...
------	------	------------	---------------------	--	----------------------------

ORDER XLIII.—TUBERACEÆ,

281. STEPHENSIA.—Tul, Compt.

1903	...	...	S. arenivaga	Cooke and Mass., Grev. XXI. 38 (1892) ...	Desert stephensia
------	-----	-----	---------------------	---	--------------------------

ORDER XLIV.—ENDOGONACEÆ,

282. ENDOGONE.—Link,

1904	1350	VIII. 3597	E. australis	Berk., Fl. Tasm. II. 282 (1860) ...	Southern endogone
------	------	------------	---------------------	-------------------------------------	--------------------------

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			

VITT. MON. TUBER. 12 (1831).

TUL. FUNG. HYP. 100 (1851).

Syn. Pl Myc. (1820).

1902	...	...	...	...	...	Q.	...	Under trees	...	Rounded or depressed, hollowed out on both sides, arising from green crustaceous mycelium.
------	-----	-----	-----	-----	-----	----	-----	-------------	-----	--

FRIES, S.V.S. 437 (1849).

Rend. XXI. 1433 (1845).—Genoa.

1903	...	S.A.	...	...	...	...	...	Sandy soil	...	Nearly globose, irregular, pale, soft, becoming hard, gathering particles of sand which cohere.
------	-----	------	-----	-----	-----	-----	-----	------------	-----	---

FRIES, S.V.S. 438 (1849).

Obs. Pl. III. 33 (1809).

1904	...	...	T	...	...	...	...	Ground	...	Hemispherical, white.
------	-----	-----	---	-----	-----	-----	-----	--------	-----	-----------------------

GENERAL CLASSIFICATION OF HYPHOMYCETES.

GROUP VII.—HYPHOMYCETES, MARTIUS.

ARRANGEMENT OF ORDERS (4).

45. MUCEDINACEÆ—Finely filamentous, pale or bright coloured (rarely brownish).
46. DEMATIACEÆ—Finely filamentous, brown or black, rather rigid.
47. STILBEACEÆ—Finely filamentous, pale or brown: fertile threads collected in fascicles (stroma).
48. TUBERCULARIACEÆ—Compact, wart-like, globose, disc-like, superficial or erumpent, waxy or somewhat gelatinous

ORDER XLV.—MUCEDINACEÆ, LINK.

ARRANGEMENT OF GENERA (13).

Section 1. Amerosporæ, Sacc.—Conidia spherical or shortly cylindrical, continuous, transparent or brightly coloured.

Sub-section 1. Micronemæ, Sacc.—Threads very short, or scarcely distinct from conidia.

Genera (3)—

283. Oospora, Wallr.		284. Monilia, Pers.		285. Oidium, Link.
----------------------	--	---------------------	--	--------------------

Sub-section 2. Macronemæ, Sacc.—Threads elongated, distinct from conidia.

Genera (9)—

286. Trichoderma, Pers.		289. Rhinotrichum, Corda.		292. Sepedonium, Link.
287. Aspergillus, Adans.		290. Sporotrichum, Link.		293. Verticillium, Nees.
288. Penicillium, Link.		291. Botrytis, Adans.		294. Nematogonium, Desm.

Section 2. Didymosporæ, Sacc.—Conidia ovoid, oblong or shortly fusoid, one septate, hyaline or brightly coloured.

Genus (1)—

295. Tricothecium, Link.

ORDER XLVI.—DEMATIACEÆ, FRIES.

ARRANGEMENT OF GENERA (19).

Section 1. Amerosporæ, Sacc.—Conidia continuous, globose, ovoid or oblong.

Sub-section 1. Micronemæ, Sacc.—Threads very short, or scarcely distinct from conidia.

Genera (4)—

296. Coniosporium, Link.		298. Hormiscium, Kunze.		299. Heterobotrys, Sacc.
297. Torula, Pers.				

Sub-section 2. Macronemæ, Sacc.—Threads evident and distinct from conidia.

Genera (3)—

300. Periconia, Tode.		301. Monotropa, Corda.		302. Botryotrichum, Sacc. and March.
-----------------------	--	------------------------	--	--------------------------------------

Section 2. Didymosporæ, Sacc.—Conidia ovoid or oblong, typically one septate.

Sub-section 1. Micronemæ, Sacc.—Threads very short or scarcely distinct.

Genus (1)—

303. Bipora, Corda.

Sub-section 2. Macronemæ, Sacc.—Threads evident and distinct from conidia.

Genera (3)—

304. Fusieladium, Bön.		305. Scolecotrichum, Kunze.		306. Cladosporium, Link.
------------------------	--	-----------------------------	--	--------------------------

Section 3. Phragmosporæ, Sacc.—Conidia ovoid, oblong, cylindrical or worm-shaped, two or more septate.

Sub-section 2. Macronemæ, Sacc.—Threads evident, and distinct from conidia.

Genera (4)—

307. Helminthosporium, Link.		309. Cercospora, Fres.		310. Heterosporium, Klotzsch.
308. Brachysporium, Sacc.				

Section 4. Dictyosporæ, Sacc.—Conidia globose, transversely and longitudinally septate, brown.

Sub-section 1. Micronemeæ, Sacc.—Threads very short, or scarcely distinct.

Genus (1)—

311. Sporodesmium, Link.

Sub-section 2. Macronemeæ, Sacc.—Threads evident and distinct from conidia.

Genera (3)—

312. Stemphylium, Wallr. | 313. Macrosporium, Fries. | 314. Fumago, Pers.

ORDER XLVII.—STILBEACEÆ, FRIES.

ARRANGEMENT OF GENERA (6).

Series 1. Hyalostilbeæ, Sacc.—Threads and conidia pallid.

Section 1. Amerosporæ, Sacc.—Conidia globular, ellipsoid or oblong, continuous, transparent or pallid.

Genera (3)—

315. Stilbum, Tode. | 316. Isaria, Pers. | 317. Ceratium, Alb. and Schw

Series 2. Phæostilbeæ, Sacc.—Threads and conidia brown, rigid.

Section 1. Amerosporæ, Sacc.—Conidia globose, oblong or elongated, continuous.

Genus (1)—

318. Harpographium, Sacc.

Section 4. Phragmosporæ, Sacc.—Conidia oblong or cylindrical, two to more septate.

Genera (2)—

319. Podosporium, Schw. | 320. Isariopsis, Fries.

ORDER XLVIII.—TUBERCULARIACEÆ, EHRL.

ARRANGEMENT OF GENERA (13).

Series 1. Tuberculariæ mucedineæ, Sacc.—Threads and conidia white or bright coloured.

Section 1. Amerosporæ, Sacc.—Conidia continuous, ovoid, sigmoid, shortly cylindrical or fusoid.

Genera (6)—

321. Tubercularia, Tode. | 323. Ægerita, Pers. | 325. Hymenula, Fries.
322. Illosporium, Mart. | 324. Fusicolla, Bon. | 326. Thozetia, Berk. and F. v. M.

Section 3. Phragmosporæ, Sacc.—Conidia elongated, fusoid or sickle shaped, typically two or more septate.

Genera (3)—

327. Bactridium, Kunze. | 328. Fusarium, Link. | 329. Microcera, Desm.

Series 2. Tuberculariæ dematiæ, Sacc.—Threads olive or sooty black, conidia same colour or hyaline.

Section 1. Amerosporæ, Sacc.—Conidia continuous, globose, ovoid or elongated.

Genera (4)—

330. Epicoccum, Link. | 332. Myrothecium, Tode. | 333. Actinomma, Sacc.
331. Strumella, Sacc. |

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
GROUP VII.—HYPHOMYCETES.—MARTIUS,					
ORDER XLV.—MUCEDINACEÆ,					
283. OOSPORA.—Wallr. Fl.					
1905	1916	X. 7062	<i>O. Aphidis</i> ...	Cooke and Mass., Grev. XVI. 76 (1888) ...	Aphis oospora ...
1906	...	...	<i>O. rutilans</i> ...	Cooke and Mass., Grev. XXI. 39 (1892) ...	Red oospora ...
1907	...	IV. 69	<i>O. vinosella</i> ...	Sacc., Fung. Ital. 874 (1886) ...	Vinous oospora ...
284. MONILIA.—Pers., Syn. 691					
1908	...	IV. 157	<i>M. fructigena</i> ...	Pers., Syn. 693 (1801) ...	Fruit-growing monilia (Brown rot)
285. OIDIUM.—Link, Berl.					
1909	...	IV. 199	<i>O. Chrysanthemi</i> ...	Rabh., Hedw. I. 19 (1852) ...	Chrysanthemum oidium ...
1910	1917	„ 189	<i>O. erysipoides</i> ...	Fries, S.M. III. 432 (1829) ...	Erysiphe-like oidium ...
1911	1918	„ 190	<i>O. leucoconium</i> ...	Desm., Ann. Sci. Nat. XIII. 102 (1829) ...	White-dust oidium ...
1912	1920	X. 7093	<i>O. Lycopersium</i> ...	Cooke and Mass., Grev. XVI. 114 (1888) ...	Tomato oidium ...
1913	...	IV. 219	<i>O. monilioides</i> ...	Link, Sp. Pl. 122 (1824) ...	Necklace oidium ...
1914	1919	„ 191	<i>O. Tuckeri</i> ...	Berk., Gard. Chron. 779 (1847) ...	Tucker's oidium (Powdery mildew)
286. TRICHODERMA.—Pers..					
1915	1921	IV. 284	<i>T. viride</i> ...	Pers., Syn. 230 (1801) ...	Green trichoderma ...
287. ASPERGILLUS.—Adans..					
1916	1926	IV. 342	<i>A. Cookei</i> ...	Sacc. Syll. IV. 71 (1886) ...	Cook's aspergillus ...
1917	1922	„ 304	<i>A. glaucus</i> ...	Link, Sp. Pl. Fung. I. 67 (1824) ...	Glaucous aspergillus ...
1918	1924	„ 319	<i>A. Muelleri</i> ...	Berk., Linn. Journ. XIII. 175 (1873) ...	Mueller's aspergillus ...
1919	1925	„ 326	<i>A. roseus</i> ...	Link, Sp. Pl. Fung. I. 68 (1824) ...	Rose-coloured aspergillus ...
288. PENICILLIUM.—Link, Berl. Mag. III. 16					
1920	...	IV. 381	<i>P. candidum</i> ...	Link, Obs. Myc. I. 15 (1809) ...	White penicillium ...
1921	1927	„ 373	<i>P. glaucum</i> ...	Link, Obs. Myc. I. 15 (1809) ...	Glaucous penicillium (Common blue mould)
289. RHINOTRICHUM.—Corda,					
1922	1929	IV. 448	<i>R. Carteri</i> ...	Cooke, Fung. Aust. 60 (1883) ...	Carter's rhinotrichum ...
1923	1928	„ 447	<i>R. microsporum</i> ...	Berk., Fl. Tasm. II. 272 (1860) ...	Small-spored rhinotrichum ...
1924	1930	„ 460	<i>R. pulchrum</i> ...	Berk., Linn. Journ. XIII. 175 (1873) ...	Beautiful rhinotrichum ...
1925	1931	„ 469	<i>R. ramosissimum</i> ...	Berk. and Curt., Grev. III. (1875) ...	Much-branched rhinotrichum ...
290. SPOROTRICHUM.—Link,					
1926	1932	IV. 507	<i>S. densum</i> ...	Link, Obs. Myc. I. 11 (1809) ...	Dense sporotrichum ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
FL. CRYPT. ERLANG. 334 (1817).									
LINK, BERL. MAG. III. 10 (1809).									
Crypt. II. 182 (1833).—Torula.									
1905	...	...	...	...	...	Q	...	Aphides upon Pump-kin leaves	Threads short, continuous, somewhat tufted, transparent. Conidia in little chains.
1906	...	...	...	V.	...	...	...	Dung ...	Expanded, crustaceous, red or orange red. Conidia in chains.
1907	...	...	...	V.	...	...	...	With moist <i>Daldinia concentrica</i>	Tufted, cushion shaped, dirty yellow to wine colour, minute. Conidia in more or less elongated chains.
(1801).—Oidium, Torula.									
1908	...	...	...	V.	...	...	B.	Apples, Pears, &c.	Compact tufts, cushion shaped, usually running together, downy, whitish then fleshy ochre. Common.
Mag. III. 18 (1809).									
1909	...	...	...	V.	...	...	B.	Leaves of Chrysanthemum	Expanded, white. Threads creeping, continuous, transparent. Conidia in long chains.
1910	...	...	...	V.	N.S.W.	Q.	B.	Living leaves of various plants	Broadly expanded, indeterminate, white. Tufts conspicuous, rosy white, threads erect, very slender.
1911	...	...	...	V.	N.S.W.	Q.	B.	Rose leaves, &c. ...	Tufts expanded, white. Threads creeping, with fertile branches short and erect. Conidial stage of <i>Spherotheca pannosa</i> .
1912	...	...	...	V.	...	...	...	Stems and leaves of Tomato (<i>Solanum lycopersicum</i>)	Tufts expanded, indeterminate, white, spiderweb-like. Threads short, branching, erect.
1913	...	...	...	V.	N.S.W.	...	B.	Living leaves, &c. ...	Tufts widely spread, oebrey white. Conidia forming chains like a necklace, dirty white. Conidial stage of <i>Erysiphe graminis</i> .
1914	...	S.A.	T.	V.	N.S.W.	Q.	B.	Viue leaves and grapes	Tufts densely clustered, often running together and forming whitish web-like layer. Conidia barrel shaped.
Obs. Myc. I. 99 (1796).									
1915	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Bark, wood, leaves, and branches	Tufts nearly circular, cushion shaped, compact, then expanded, first white, then bluish green, afterwards yellowish. Conidial stage of <i>Hypocrea rufa</i> .
Fam. II. 2 (1763).—Mucor, Monilia.									
1916	W.A.	S.A.	...	V.	...	Q.	...	Dead plants and leaves	Gregarious, white, intricately interwoven. Fertile threads erect, transparent, crowned with large globose vesicle.
1917	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Vegetable substances	Creeping threads fluffy, branched, nocoloured. Fertile threads erect, simple, transparent or glaucous, swelling into spherical vesicle. Conidial stage of <i>Eurotium herbariorum</i> .
1918	W.A.	...	...	V.	...	...	...	<i>Lepiota bubalina</i> , &c.	Snowy white, creeping. Fertile threads erect, rather flexuous.
1919	...	...	...	V.	...	...	B.	Soil, damp paper, linen, &c.	Thin, creeping. Fertile threads simple. Conidia globular, rose colour.
(1809).—Aspergillus, Botrytis, Mucor.									
1920	...	...	...	V.	...	...	B.	Leaves and decaying substances	Tufts running together, white. Sterile threads creeping, interwoven white, fertile threads ascending or erect.
1921	...	...	...	V.	N.S.W.	Q.	B.	Decaying vegetables	Expanded, creeping, white. Sterile threads creeping, intricate. Fertile threads erect, branched at top in a pencil-like manner. Branches erect, once or twice forked. Conidia verdigris green.
Ic. Fung. I. 17 (1837).									
1922	...	...	...	...	N.S.W.	...	...	Wood ...	White, peziza shaped, rather compact. Threads branched, club shaped, somewhat knotted.
1923	...	...	T.	...	...	...	...	Ground ...	Threads stuck together, forming nearly cylindrical clubs, with spicules.
1924	...	...	...	V.	N.S.W.	...	...	Rotten wood	Forming thin saffron-coloured layer. Threads globosely clavate at top.
1925	...	S.A.	...	...	N.S.W.	...	B.	Rotten wood	Pale fawn or tan colour. Threads very much branched, ultimate joints elongated and toothed.
Berl. Mag. III. 12 (1809).									
1926	...	...	...	...	N.S.W.	Q.	...	Dead insects, &c. ...	Threads sparingly branched, transparent, white, densely crowded in a rather thick layer. Allied to <i>Botrytis Bassiana</i> , which causes the disease known as "Muscardine" in silkworms.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
291. BOTRYTIS.— <i>Adans.</i> , Fam. II. 3 (1763).—					
1927	1933	IV. 664	<i>B. vulgaris</i> ...	Fries, S.M. III. 398 (1829) ...	Common botrytis ...
292. SEPEDONIUM.—Link,					
1928	1935	X. 7206	<i>S. aureo-fulvum</i> ..	Cooke and Mass., Grev. XVI. 76 (1888) ...	Golden-yellow sepedonium ...
1929	1934	IV. 754	<i>S. chrysospermum</i> ...	Fries, S.M. III. 438 (1829) ...	Golden-spored sepedonium ...
293. VERTICILLIUM.—Nees,					
1930	1936	IV. 792	<i>V. eximium</i> ...	Berk., Linn. Journ. XIII. 175 (1873) ...	Excellent verticillium ...
1931	1938	„ 808	<i>V. lateritium</i> ...	Berk., in Cooke's Handb. Brit. Fung. 635 (1871) ...	Brick-red verticillium ..
1932	1937	„ 797	<i>V. nivium</i> ...	Berk., Fl. Tasm. II. 271 (1860) ...	Snow-white verticillium ...
294. NEMATOGONIUM.—Desm., Ann.					
1933	1939	IV. 867	<i>N. aurantiacum</i> ...	Desm., Ann. Sci. Nat. II. 70 (1834) ...	Orange nematogonium ...
1934	1940	„ 868	<i>N. aureum</i> ...	Sacc. Syll. IV. 8 (1886) ...	Golden nematogonium ...
295. TRICHOHECIUM.—Link,					
1935	1941	IV. 881	<i>T. roseum</i> ...	Link, Obs. Myc. I. 16 (1809) ...	Rosy trichothecium ...

ORDER XLVI.—DEMATIACEÆ, FRIES,

296. CONIOSPORIUM.—Link,					
1936	1942	IV. 1152	<i>C. inquinans</i> ...	Dur. and Mont., Fl. Alg. I. 327 (1846) ...	Black coniosporium ...
1937	1943	X. 7334	<i>C. pterospermum</i> ...	Cooke and Mass., Grev. XIX. 90 (1891) ...	Wing-spored coniosporium ...
297. TORULA.—Pers.,					
1938	1944	IV. 1230	<i>T. herbarum</i> ...	Link, Sp. Pl. Fung. I. 128 (1824) ...	Herb-growing torula ...
193	1945	X. 7363	<i>T. mycetophila</i> ...	Cooke and Mass., Grev. XVI. 3 (1887) ...	Fungus-loving torula ...
298. HORMISCIMUM.—Kunze, Myk.					
1940	1947	IV. 1286	<i>H. pithyophilum</i> ...	Sacc. Syll. IV. 265 (1886) ...	Pine-loving hormiscium ...
1941	1946	„ 1283	<i>H. stilbosporum</i> ...	Sacc. Syll. IV. 264 (1886) ...	Stilbum-spored hormiscium ...
299. HETEROBOTRYS.—Sacc.,					
1942	1948	IV. 1296	<i>H. paradoxa</i> ...	Sacc., Mich. II. 124 (1880) ...	Paradoxical heterobotrys ...
300. PERICONIA.—Tode,					
1943	1949	IV. 1328	<i>P. nigrella</i> ...	Sacc. Syll. IV. 274 (1886) ...	Black periconia ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Polyactus, Monilia, Peziza.									
1927	...	S.A.	T.	V.	N.S.W.	Q.	B.	Herbs, fruit, flowers, leaves, and branches in decay	Tufts olive grey. Threads fluffy, erect, olive, branched above. Branches shortened, spreading, and branchlets bearing the conglomerate conidia. Mould-like <i>Botrytis</i> is only the conidial form of <i>Peziza</i> .
Berl. Mag. III. 18 (1809).—Mucor, Uredo.									
1928	...	...	...	V.	...	...	...	<i>Polyporus</i>	Threads creeping, branched. Conidia profuse, globose, forming a golden-tawny powder within decaying <i>Polypori</i> .
1929	W.A.	S.A.	...	V.	N.S.W.	Q.	B.	<i>Boletus</i> , &c.	Threads spread out, then interwoven, rather thick, almost transparent, variously forked, conidia yellow or golden yellow.
Syst. Pilz. 56 (1816).									
1930	...	...	...	...	N.S.W.	...	...	<i>Clavaria</i> ...	Threads branched, branches short, thickened at ends, with radiating acute conidia-bearing processes. A beautiful species.
1931	...	...	...	V.	...	...	B.	Maize ...	Threads elegantly and many times branched in whorls, collected in brick-red velvety or woolly tufts.
1932	...	...	T.	...	...	...	...	Dead Agarics	White, branched. Branches rather short, thickened at base.
Sci. Nat. II. 69 (1834).—Aspergillus.									
1933	...	...	...	...	...	...	B.	Bark and wood	Tufts velvety, orange tawny, expanded. Sterile threads creeping, thin. Fertile threads erect, swollen at each end.
1934	...	...	...	...	...	Q.	B.	Bark ...	Fertile threads erect, short, club shaped, with about four joints. Conidia golden yellow.
Berl. Mag. III. 18 (1809).—Trichoderma, Puecinia.									
1935	...	...	...	V.	...	...	B.	Rotting fruit, branches, leaves, paper, cheese, &c.	Tufts cushion shaped, velvety, rather large; at first white, then rosy.
SYST. MYC. III. 335 (1832).									
Berl. Mag III. 8 (1809).									
1936	W.A.	...	...	...	...	...	...	Stems of <i>Arundo</i> ...	Spread out, very black. Tufts rounded or oblong, run together and irregular.
1937	...	...	...	V.	...	...	...	<i>Lepidosperma</i> ...	Pustles gregarious, small, bursting through, blackish. Spore body globose, with membranous expansion.
Ust. Ann. IX. 25 (1795).—Monilia.									
1938	...	...	...	...	...	Q.	B.	Rotting herb stems	Tufts expanded, olive to ochrey, then becoming black, somewhat velvety. Sterile threads ereeping, sooty. Fertile threads erect, olive, then black.
1939	...	...	...	V.	...	...	...	Pileus or cap of <i>Polyporus cinnabarinus</i>	Tufts minute, very thin, scattered, black. Threads sparingly branched, nearly straight.
Heft. I. 12 (1817).—Antennaria, Torula.									
1940	...	S.A.	...	...	N.S.W.	...	B.	Branches and leaves of <i>Conifera</i> and <i>Eucalyptus</i>	Expanded, thick, superficial, quite black. Chains of conidia indistinctly branched, branches tapering towards apex and slightly curved.
1941	...	...	...	...	...	Q.	B.	Branches ...	Tufts bursting through, powdery, run together, quite black. Chains unequal, branched, or simple.
Mich. II. 21 (1880).									
1942	...	S.A.	...	...	...	...	...	Upper surface of leaves of <i>Bertya rotundifolia</i>	Tufted, gregarious, black. Threads straggling, ereeping, pale, sooty. Conidia brown.
Fung. Meek. I. 2 (1790).—Sporocybe.									
1943	...	...	...	...	...	Q.	B.	Leaves of <i>Andropogon</i>	Very minute, black. Fertile threads simple, thin. Head globose or ellipsoid.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
301. MONOTOSPORA.—Corda,					
1944	...	...	<i>M. fasciculata</i> ...	Cooke and Mass., Grev. XXI. 39 (1892)	Fasciculate monotospora ...
302. BOTRYOTRICHUM.—Sacc. and					
1945	...	...	<i>B. Lachnella</i> ...	Sacc., Hedw. 58 (1893)	Lachnella botryotrichum ...
303. BISPORA.—Corda,					
1946	1950	IV. 1632	<i>B. monilioides</i> ...	Corda, Ic. Fung. I. 9 (1837)	Necklace bispora ...
304. FUSICLADIUM.—Bon. Handb. 80					
1947	1951	IV. 1642	<i>F. dendriticum</i> ..	Fekl., S.M. 357 (1875) ...	Tree-like fusicladium ... (Black spot of Apple)
1948	1952	„ 1643	<i>F. pyrinum</i> ...	Fekl., S.M. 357 (1875) ...	Pear fusicladium ...
305. SCOLECOTRICHUM.—Kunze and					
1949	1953	X. 7478	<i>S. atriellum</i> ...	Cooke and Mass., Grev. XVI. 3 (1887)	Black scolecotrichum ...
1950	...	IV. 1656	<i>S. graminis</i> , var. <i>Avenae</i> ...	Erikss., Zeit. Pflkrk. I. 23 (1891)	Oat scolecotrichum ...
306. CLADOSPORIUM.—Link,					
1951	1957	IV. 1698	<i>C. Asteroma</i> ...	Fekl., S.M. 355 (1875) ...	Asteroma cladosporium ...
1951A	„	...	<i>C. Asteroma</i> , var. <i>minor</i> ...	Cooke, Handb. Austr. Fung. 376 (1892)	Lesser cladosporium ...
1952	1959	IV. 1718	<i>C. epiphyllum</i> ...	Mart., Erl. 351 (1817) ...	Epiphyllous cladosporium ...
1953	1954	IV. 1665	<i>C. herbarum</i> ...	Link, Obs. Myc. II. 37 (1809)	Herb-growing cladosporium ...
1953A	„	...	<i>C. herbarum</i> , var. <i>epixylinum</i>	Corda ...	Wood-growing cladosporium ...
1954	1958	IV. 1714	<i>C. hypophyllum</i> ...	Fekl., S.M. 356 (1875) ...	Hypophyllous cladosporium ...
1955	1955	„ 1669	<i>C. oligocarpum</i> ...	Corda, Ic. Fung. I. 14 (1837)	Few-spored cladosporium ...
1956	1961	„ 1774	<i>C. papyricolum</i> ...	Berk. and Br., Linn. Trans. II. 68 (1883)	Paper-growing cladosporium ...
1957	1956	„ 1670	<i>C. stenosporum</i> ...	Berk. and Curt., Grev. III. (1875)	Slender-spored cladosporium ...
1958	1960	„ 1750	<i>C. Typharum</i> ...	Desm., Exs. 304	Typha cladosporium ...
307. HELMINTHOSPORIUM.—Link,					
1959	1966	IV. 1969	<i>H. inconspicuum</i> ...	Cooke and Ell., Grev. VI. 88 (1878)	Inconspicuous helminthosporium
1960	1963	„ 1973	<i>H. macrocarpum</i> ...	Grev., Scot. III. 148 (1825)	Large-spored helminthosporium
1961	1965	„ 1966	<i>H. puccinioides</i> ...	Sacc. and Berl., Rev. Myc. (1885)	Puccinia-like helminthosporium
1962	1962	„ 1971	<i>H. Ravenelii</i> ...	Curt., in Sill. Journ. 352 (1848)	Raveuel's helminthosporium ...
1963	1964	„ 2010	<i>H. rhabdiferum</i> ...	Berk. and Br., Ann. Nat. Hist. XV. 402 (1865)	Rod-bearing helminthosporium
308. BRACHYSPORIUM.—Sacc.,					
1964	1967	IV. 2039	<i>B. oligocarpum</i> ...	Sacc. Syll. IV. 424 (1886)	Few-spored brachysporium ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Ic. Fung. I. 11 (1837).									
1944	...	...	...	V.	...	...	...	Bark ...	Tufts bursting through, gregarious, black. Fertile threads erect, densely fasciculate in awl-shaped tufts.
March., Champ. Copr. Belg. 34 (1885).									
1945	...	...	...	V.	...	...	...	Branches and spines of <i>Bursaria spinosa</i> , not yet dead	Gregarious, number to dusky. Tufts of threads in circular bundles, like <i>Lachnella</i> . Sterile threads erect. Conidia spherical, rather transparent, granular.
Ic. Fung. I. 9 (1837).									
1946	...	...	...	...	N.S.W.	Q.	B.	Wood ...	Expanded, dark brown, powdery. Threads short, rather conical. Conidia sooty brown. Conidial stage of <i>Bisporella monilifera</i> .
(1851).—Cladosporium, Helminthosporium.									
1947	...	S.A.	T.	V.	N.S.W.	Q.	B.	Leaves and fruit of Apple and Pear	Expanded, velvety, olive, often tree-like on leaves. Threads filiform, erect, in bundles. Conidia olive.
1948	...	S.A.	T.	V.	N.S.W.	Q.	B.	Leaves and fruit of Pear	Expanded, velvety, olive. Threads short, tapering, toothed at apex. Conidia olive.
Schm., Myk. Heft. I. 10 (1817).									
1949	...	...	...	...	...	Q.	...	Twigs of <i>Passiflora</i>	Tufts spread out, run together, black. Threads erect, simple, brown. Conidia dark brown.
1950	...	...	...	V.	...	...	...	Oats ...	Spots on leaves elongated, ochrey, drying up. Threads in bundles, point-like, densely clustered. Conidia olive to brown.
Berl. Mag. VII. 37 (1816).—Dematium.									
1951	...	...	...	...	...	...	...	...	...
1951A	...	...	...	...	...	Q.	...	Foliage of Grape Vine	Tufts in centre of brown spot, disposed in a tree like manner, minute, yellow, becoming greenish.
1952	...	...	...	V.	...	...	B.	Leaves of Oak, &c.	Tufts arranged in a circle, olive, then blackish, large, thick. Threads at first erect, then declining.
1953	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Stems, leaves, &c. ...	Tufts densely clustered, run together, forming a velvety yellow-olive then dark-olive layer.
1953A	...	...	...	...	...	Q.	...	Wood.	...
1954	...	...	...	...	...	Q.	...	Lower surface of leaves of <i>Serjania</i>	Tufts spread out, then greyish green. Threads branched, flexuous, yellow.
1955	...	...	...	...	...	Q.	...	Wood ...	Tufts minute, solitary, black. Threads tufted, erect, long, slender, slightly branched, olive.
1956	...	...	...	...	...	Q.	...	On paper forming dark-grey layer	Threads irregularly branched, transparent above, brown below. Conidia pale brown.
1957	...	...	...	...	...	Q.	...	Leaves of Pear, &c.	Threads simple, thin, divided, arising from a creeping mycelium, greyish brown below.
1958	...	...	...	V.	...	...	...	Leaves of <i>Typha</i> (Bulrush)	Tufts elongated or oblong, scattered, turning black, seated at first on distinct greyish spot.
Berl. Mag. III. 10 (1809).—Macrosporium.									
1959	...	...	...	...	...	Q.	B.	Fading leaves of Maize	Thin cloud-like stain. Threads elongated, septate knotted, pale brown.
1960	...	...	...	...	...	Q.	B.	Trunks and branches	Expanded, velvety, dark olive or sooty brown. Threads clustered, lax, simple or sparingly branched.
1961	...	...	...	...	...	Q.	...	Fading or dead leaves of <i>Tristania laurina</i>	Tufts on both surfaces, very black, loosely gregarious, disc shaped, compact, resembling <i>Puccinia</i> . Conidia pale sooty brown.
1962	...	...	...	...	...	Q.	...	Inflorescence of grasses (<i>Sporobolus indicus</i>)	Spongy. Threads flaccid, flexuous, knotted, branched. Conidia brown.
1963	...	S.A.	...	...	...	...	B.	Ripe peaches ...	Expanded, internally black. Threads erect, sparingly branched. Conidia straight, at first oblong, then elongated and somewhat linear, dark brown.
Mich II. 28 (1880).—Helminthosporium.									
1964	...	...	...	...	...	Q.	...	Wood ...	Tufts minute, linear, nearly parallel. Threads flexuous, simple, in bundles, dark brown. Conidia yellow brown.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name
309. CERCOSPORA.—Fres., Beitr. 90					
1965	1971	X. 7685	<i>C. Daviesiæ</i> ...	Cooke and Mass., Grev. XVIII. 7 (1889) ...	Daviesia cercospora ...
1966	1968	„ 7696	<i>C. epicoccoides</i> ...	Cooke and Mass., Grev. XIX. 91 (1891) ...	Epicoccum-like cercospora ...
1967	1973	„ 7697	<i>C. Eucalypti</i> ...	Cooke and Mass., Grev. XVIII. 7 (1889) ...	Eucalyptus cercospora ...
1968	...	...	<i>C. Glycines</i> ...	Cooke and Mass., Grev. XXI. 39 (1892) ...	Glycine cercospora ...
1969	1969	X. 7678	<i>C. Kennedyæ</i> ...	Cooke and Mass., Grev. XIX. 90 (1891) ...	Kennedyia cercospora ...
1970	1972	IV. 2161	<i>C. Solanacea</i> ...	Sacc. and Berl., Rev. Myc. (1885) ...	Solanum cercospora ...
1971	1970	„ 2200	<i>C. viticola</i> ...	Sacc., Syll. IV. 458 (1886) ...	Vine cercospora (Tufted leaf blight) ...
310. HETEROSPORIUM.—Klotzsch,					
1972	1974	X. 7769	<i>H. epimyces</i> ...	Cooke and Mass., Grev. XVI. 80 (1888) ...	Fungus heterosporium ...
311. SPORODESMIUM.—Link, Berl					
1973	1975	IV. 2391	<i>S. atrofusum</i> ...	Cooke, Grev. XII. 12 (1883) ...	Blackish-brown sporodesmium ...
1974	1976	„ 2356	<i>S. melanopodium</i> ...	Berk. and Br., Ann. Nat. Hist. V. 159 (1850) ...	Black-stalked sporodesmium ...
312. STEMPHYLIUM.—Wallr., Fl.					
1975	1977	IV. 2487	<i>S. pulchrum</i> ...	Sacc., Syll. IV. 521 (1886) ...	Beautiful stemphylium ...
313. MACROSPORIUM.—Fries,					
1976	1980	X. 7837	<i>M. Camelliæ</i> ...	Cooke and Mass., Grev. XVII. 42 (1888) ...	Camellia macrosporium ...
1977	1978	IV. 2501	<i>M. cladosporioides</i> ...	Desm., Pl. Crypt. 3 (1857) ...	Cladosporium-like macrosporium ...
1978	1979	„ 2499	<i>M. commune</i> ...	Rabh., Fung. Eur. Exs. 1360 ...	Common macrosporium ...
1979	...	X. 7853	<i>M. graminum</i> ...	Cooke, Grev. XVII. 66 (1889) ...	Grass macrosporium ...
1980	1982	IV. 2549	<i>M. peponicolum</i> ...	Rabh., in Sitz. 101 (1867) ...	Gourd-growing macrosporium ...
1981	1981	X. 7841	<i>M. Readeri</i> ...	Winter, Rev. Myc. 212 (1886) ...	Reader's macrosporium ...
1982	1980 bis.	IV. 2552	<i>M. Tomato</i> ...	Cooke, Grev. XII. 32 (1883) ...	Tomato macrosporium ...
314. FUMAGO.—Pers., Myc.					
1983	1983	IV. 2618	<i>F. vagans</i> ...	Pers., Myc. Eur. I. 9 (1822) ...	Creeping fumago ...
ORDER XLVII.—STILBEACEÆ,					
315. STILBUM.—Tode,					
1981	...	IV. 2714	<i>S. aurantiacum</i> ...	Bab., Linn. Trans. (1839) ...	Orange stilbum ...
1985	1986	X. 7894	<i>S. caninum</i> ...	Cooke and Mass., Grev. XX. 36 (1891) ...	Dog's dung stilbum ...
1986	1988	IV. 2705	<i>S. cinnabarium</i> ...	Mont., Fl. Cub. 308 (1812) ...	Vermilion stilbum ...
1987	1987	X. 7879	<i>S. corallinum</i> ...	Cooke and Mass., Grev. XIX. 91 (1891) ...	Coralline stilbum ...
1988	1984	IV. 2680	<i>S. erythrocephalum</i> ...	Ditm., Sturm. Fl. III. (1817) ...	Red-headed stilbum ...
1989	1985	X. 7893	<i>S. Formicarum</i> ...	Cooke and Mass., Grev. XVIII. 8 (1889) ...	Ant stilbum ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
(1850).—Cladosporium, Helminthosporium.									
1965	...	...	...	V.	...	...	...	Fading leaves of <i>Daviesia latifolia</i>	Spots brown, irregular, angular. Threads in bundles, shortened. Conidia bent like a bow, pale brown.
1966	...	...	...	V.	...	...	...	<i>Eucalyptus</i> leaves ...	Spots small or run together, purple. Tufts gregarious, rather compact, black. Conidia pale olive.
1967	...	...	...	V.	...	...	...	Fading <i>Eucalyptus</i> leaves	Spots rather circular, or run together, pale, with rosy margin. Threads shortened. Conidia curved, pale.
1968	...	...	...	V.	...	...	...	Living leaves of <i>Glycine clandestina</i>	On both surfaces, but chiefly upper, spots definite, irregular, angular, umber. Tufts gregarious on the spots, point-like, black. Conidia almost transparent.
1969	...	...	...	V.	...	...	...	Leaves of <i>Kennedya prostrata</i>	Spots cinnamon brown, irregular and run together. Tufts scattered, black, point-like. Conidia clear olive.
1970	...	...	...	...	...	Q.	...	Leaves of <i>Solanum verbascifolium</i>	Spots nearly circular, brown. Tufts point-like, sooty olive. Conidia rod-like, curved, almost hyaline.
1971	...	...	...	...	N.S.W.	Q.	...	Vine leaves ...	Spots on both surfaces, somewhat circular or irregular, when dry ochrey. Threads often on under surface, here and there densely tufted. Conidia ochrey olive.
Herb. Myc. I. 67 (1832).									
1972	...	...	...	V.	...	...	B.	Decayed Agarics ...	Tufts olive, spread out in more or less dense velvety patches. Threads sparingly forked, pale brown. Conidia minutely warted, pale olive.
Mag. III. 41 (1809).—Spiloma.									
1973	...	...	...	V.	...	...	...	Wood ...	Expanded, velvety black. Conidia elongated, club shaped, divided in all directions into quadrate cells, dark brown.
1974	...	...	...	V.	...	...	B.	Bark ...	Tufts broad, black. Conidia nearly globose, opaque, seated on base of variable size.
Crypt. II. 300 (1833).—Mystrosporium.									
1975	W.A.	...	...	...	...	...	...	Rotten wood ...	Expanded, olive. Threads white, wrinkled, forked or trifid. Conidia a little rough, blackish.
S.M. III. 373 (1832).									
1976	...	...	...	V.	...	...	B.	Living leaves of <i>Camellia</i>	Spots circular or run together, pale, with broad brown margin. Threads tufted, pale olive.
1977	...	...	...	V.	...	...	B.	Leaves and stems of herbs	Spots large, tawny, irregular. Tufts numerous, minute, rounded, velvety, dark olive.
1978	...	...	...	...	...	Q.	B.	Stems, leaves, &c. ...	Tufts densely clustered, numerous, brown. Threads in bundles, ascending, brown. Conidia olive. Considered to be conidial condition of <i>Pleospora herbarum</i> .
1979	...	...	...	...	N.S.W.	...	...	Wheat, Sugar cane, &c.	Expanded, very thin. *Threads creeping, at length with erect branches, greyish brown. Conidia same colour.
1980	...	...	...	...	...	Q.	...	Papaw fruit ...	Spots large, circular, black. Sterile threads, slender, creeping. Fertile threads, short, erect. Conidia amber brown.
1981	...	...	...	V.	...	...	...	Dry stems of Artichoke	Tufts forming expanded black layer. Threads in minute bundles, erect, brown. Conidia brown.
1982	...	...	...	V.	...	Q.	E.	Ripe Tomatoes ...	Circular, black. Threads short, robust, flexuose. Conidia brown.
Eur. I. 9 (1822).—Cladosporium, Torula.									
1983	...	...	...	...	...	Q.	B.	Living leaves of Vine, &c.	Threads creeping, branched in a straggling manner, olive or sooty brown, forming a thin membranous black layer. Conidia in short chains.
FRIES, MICH. II. 31 (1880).									
Meckl. I. 10 (1790).—Sphaerostilbe.									
1984	...	...	...	...	...	Q.	B.	Dead branches ...	Somewhat fasciculate, orange coloured. Stem even, darker downwards. Head somewhat club shaped.
1985	...	...	...	V.	...	...	...	Dog's dung ...	Gregarious, flesh coloured. Head darker, continuous with smooth stem, which is sometimes forked. Conidia transparent.
1986	...	...	...	...	...	Q.	...	Bark ...	Gregarious, flesh coloured. Head convex to hemispherical. Stem short, mealy.
1987	...	...	...	V.	...	...	...	Bark ...	Tufted, flesh coloured. Stems tapering upwards, shortly branched, mealy. Head orange red.
1988	W.A.	...	...	V.	...	...	B.	Dung ...	Gregarious or somewhat scattered. Stem rather thick, terminated by rosy or red, globose, mealy head.
1989	...	...	...	V.	...	...	...	Dead ants (<i>Formica</i>)	Stems elongated, slender, black, flexuous, slightly thickened below, Head inversely egg shaped, rosy.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
316. ISARIA.—Pers., Tent.					
1990	1997	IV. 2851	<i>I. arbuscula</i> ...	<i>Bres. and Roum.</i> , Rev. Myc. 38 (1890) ...	Dendritic isaria ...
1991	1994	„ 2841	<i>I. Cicadæ</i> ...	<i>Miq.</i> , Ann. Sci. Nat. X. 378 (1838) ...	Cicada isaria ... (Brazilian Cicada clubs)
1992	1993	„ 2842 and 2839	<i>I. graminiperda</i> (including <i>I. fuciformis</i> , Berk.)	<i>Berk. and F. v. M.</i> , Gard. Chron. 596 (1873)	Grass-destroying isaria ...
1993	1992	IV. 2828	<i>I. radians</i> ...	<i>Berk.</i> , Fl. Tasm. II. 271 (1860) ...	Radiating isaria ...
1994	1995	X. 7921	<i>I. suffruticosa</i> ...	<i>Cooke and Mass.</i> , Grev. XVIII. 45 (1890)	Shrubby isaria ...
1995	1991	IV. 2807	<i>I. umbrina</i> ...	<i>Pers.</i> , Syn. 689 (1801) ...	Umber isaria ...
317. CERATIUM.—Alb. and Schw., Consp.					
1996	1996	IV. 2845	<i>C. hydroides</i> ...	<i>Alb. and Schw.</i> , Consp. Fung. Lus. 358 (1805) ...	Hydium-like ceratium ...
318. HARPOGRAPHIUM.—					
1997	1998	X. 7949	<i>H. corynelioides</i> ...	<i>Cooke and Mass.</i> , Grev. XVI. 76 (1888) ...	Corynelia-like harpoglyphium ...
1998	1999	„ 7948	<i>H. quaternarium</i> ...	<i>Cooke and Mass.</i> , Grev. XVI. 3 (1887) ...	Quaternate harpoglyphium ...
319. PODOSPORIUM.—Schw.,					
1999	2000	IV. 2982	<i>P. grande</i> ...	<i>Cooke</i> , Grev. XII. 11 (1883) ...	Large podosporium ...
320. ISARIOPSIS.—Fries, in Sacc.					
2000	2001	IV. 2998	<i>I. clavispora</i> ...	<i>Sacc.</i> , Syll. IV. 631 (1886) ...	Clavate-spored isariopsis ...

ORDER XLVIII.—TUBERCULARIACEÆ,

321. TUBERCULARIA.—Tode, Fung.					
2001	2002	X. 7990	<i>T. leguminum</i> ...	<i>Cooke and Mass.</i> , Grev. XVI. 33 (1887) ...	Legume-growing tubercularia ...
2002	...	IV. 3002	<i>T. vulgaris</i> ...	<i>Tode</i> , Fung. Meck. I. 18 (1790) ...	Common tubercularia ...
322. ILLOSPORIUM.—Mart.,					
2003	2004	...	<i>I. flvellum</i> ...	<i>Berk. and Br.</i> , Linn. Trans. II. 68 (1883)	Yellow illosporium ...
2004	2006	IV. 3106	<i>I. flaveolum</i> ...	<i>Sacc.</i> , Mich. II. 297 (1880) ...	Yellowish illosporium ...
2005	2005	X. 8019	<i>I. obscurum</i> ...	<i>Cooke and Mass.</i> , Grev. XVI. 113 (1888)	Obscure illosporium ...
323. ÆGERITA.—Pers., Tent.					
2006	2007	IV. 3124	<i>A. candida</i> ...	<i>Pers.</i> , Syn. 684 (1801) ...	White agerita ...
324. FUSICOLLA.—Bon.,					
2007	2008	X. 8024	<i>F. incarnata</i> ...	<i>Cooke and Mass.</i> , Grev. XVII. 8 (1888) ...	Flesh-coloured fusicolla ...
325. HYMENULA.—Fries.					
2008	...	...	<i>H. Eucalypti</i> ...	<i>Cooke and Mass.</i> , Grev. XXI. 39 (1892) ...	Eucalypt hymenula ...
326. THOZETIA.—Berk. and F. v. M.,					
2009	2009	IV. 3213	<i>T. nivea</i> ...	<i>Berk.</i> , Linn. Journ. XVIII. 388 (1881) ...	Snow-white thozetia ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						E.	Occurrence.	General Characters.	
	W.A.	S.A.	T.	V.	N.S.W.	Q.				
Disp. 41 (1797).—Ceratium.										
1990	...	...	...	...	...	Q.	...	Rotten wood	...	Snowy white. Conidia-bearing layer tree-like, branched in tufts from simple base.
1991	...	...	...	V.	...	...	...	Head of Cicada	...	Within and growing through joints of dead <i>Cicada</i> . Conidia-bearing layer hard and compact, with shortened stem.
1992	...	S.A.	...	V.	N.S.W.	Q.	B.	Grasses (<i>Lolium</i>) and germinating cereals	...	Bright orange, gelatinous, slender, sparingly branched. Conidia minute, globose.
1993	...	...	T.	...	...	...	...	Bark	...	Greyish fawn, circular, branched, covered with whorled or forked woolly tufts.
1994	...	...	...	...	N.S.W.	...	...	Hairy caterpillar	...	Tufted, white. Stem smooth or slightly mealy, with slender branches interwoven with lateral branchlets. Conidia minute, ellipsoid.
1995	...	...	...	V.	...	...	B.	Wood, and <i>Hypoxylon coccineum</i> , of which it is conidial form	...	Clubs without stem, fawn colour, in radiating tufts. Conidia inversely egg shaped, umber.
Fung. Lus. 358 (1805).—Isaria, Tremella.										
1996	...	...	...	...	N.S.W.	Q.	B.	Rotten wood	...	Conidia-bearing layer tapering, simple or sparingly branched, white or yellowish. Conidia ovoid or globose, transparent.
Sacc., Mich. II. 33 (1880).										
1997	...	...	...	V.	...	...	...	Branches of <i>Leptospermum scoparium</i>	...	Tufted, bursting through, black. Stems composite, radiating, club shaped above, simple or forked. Conidia, curved, transparent.
1998	...	...	...	...	...	Q.	...	Dead twigs of <i>Passiflora</i>	...	Tufts black, minute. Stems composite, club shaped above. Conidia spindle shaped, transparent, for the most part quarternate.
Trans. Amer. Phil. Soc. IV. (1832).										
1999	...	...	...	V.	...	...	...	Stems of <i>Aster argophyllus</i>	...	Large, black, woolly, forming dense tufts. Threads erect, crowded together, dark brown.
Mich. II. 33 (1880).—Graphium.										
2000	...	...	...	...	...	Q.	...	Vine leaves	...	Minute, olive, arising from circular brown spots. Threads relaxed above and flexuous.
EHRB. SYLV. MYC. 12 (1818).										
Meck. I. 18 (1719).—Tremella, Sphæria.										
2001	...	...	...	...	...	Q.	...	Legumes of <i>Cassia</i>	...	Minute, bursting through, flesh colour. Conidia bearers short, straight.
2002	...	...	...	...	...	...	B.	Branches...	...	Gregarious, bursting through, vermilion coloured, globular to depressed, more or less shortly stalked.
Fl. Crypt. Erl. 325 (1817).										
2003	...	...	...	...	...	Q.	...	Lichens...	...	Stalked, yellow. Stem short. Conidia globose.
2004	...	...	...	...	...	Q.	...	Rotten wood	...	Very minute, gregarious, yellow. Threads branched in a forked manner, twisted.
2005	...	...	...	V.	...	...	...	Leaves of <i>Eucalyptus globulus</i>	...	Somewhat gregarious, circular, bursting through, minute, sooty brown. Threads branched in a forked manner.
Disp. 40 (1797).—Tubercularia, Sclerotium.										
2006	...	...	...	V.	...	...	B.	Wood and bark	...	Crowded, granule-like, globose to hemispherical, size of poppy or turnip seed, white but yellowish when dry.
Handb. 150 (1851).										
2007	...	...	...	...	...	Q.	...	Dead eoriaceous leaves	...	Pustules small, gregarious, seated on paler spots, rosy flesh colour, somewhat gelatinous or scattered over leaf, stalks, and midribs.
Pl. Homon 94 (1825).										
2008	...	...	...	V.	...	...	...	<i>Eucalyptus</i> leaves...	...	On both surfaces; pustules bursting through, disc-like, brownish. Conidia bearers simple, rather thick.
Linn. Journ. XVIII. 388 (1881).										
2009	...	...	...	...	...	Q.	...	Rotten wood	...	White. Conidia oblong, transparent, acute at each extremity, and terminated by long bristle.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
327. BACTRIDIDIUM.—Kunze,					
2010	2010	IV. 3268	<i>B. flavum</i>	Kunze and Schw., Myk. Heft. I. 5 (1817)	Orange bactridium
2011	2011	„ 3273	<i>B. magnum</i>	Cooke, Grev. VIII. 60 (1879)	Great bactridium
328. FUSARIUM.—Link,					
2012	2016	X. 8105	<i>F. hypocreoideum</i>	Cooke and Mass., Grev. XVI. 76 (1888)	Hypocrea-like fusarium
2013	2012	IV. 3283	<i>F. lateritium</i>	Nees, Syst. 31 (1816)	Brick-red fusarium
2014	2015	X. 8074	<i>F. longisporum</i>	Cooke and Mass., Grev. XVI. 4 (1887)	Long-spored fusarium
2015	2013	...	<i>F. rubicolor</i>	Berk. and Br., Linn. Trans. 11. 68 (1883)	Ruby-coloured fusarium
329. MICROCERA.—Desm.,					
2016	2017	IV. 3473	<i>M. coccophila</i>	Desm., Ann. Sci. Nat. X. 359 (1848)	Coccus-loving microcera
2017	2018	X. 8119	<i>M. rectispora</i>	Cooke and Mass., Grev. XVI. 4 (1887)	Straight-spored microcera
330. EPICOCCUM.—Link,					
2018	2019	IV. 3491	<i>E. scabrum</i>	Corda, Ic. Fung. I. 5 (1837)	Rough epicoccum (False potato disease)
331. STRUMELLA.—Sacc.,					
2019	2020	X. 8127	<i>S. hysteroidea</i>	Cooke and Mass., Grev. XVII. 69 (1889)	Hysterium-like strumella
2020	2022	„ 8128	<i>S. patelloidea</i>	Cooke and Mass., Grev. XX. 7 (1891)	Patelloid strumella
2021	2021	„ 8130	<i>S. sacchari</i>	Cooke, Grev. XIX. 45 (1890)	Sugar-cane strumella (Cane spume)
332. MYROTHECIUM.—Tode,					
2022	2025	IV. 3552	<i>M. inundatum</i>	Tode, Meck. I. 25 (1790)	Inundated myrothecium
2023	2024	„ 3550	<i>M. roridum</i>	Tode, Meck. I. 25 (1790)	Bedewed myrothecium
333. ACTINOMMA.—Sacc.,					
2024	2023	IV. 3564	<i>A. Gastonis</i>	Sacc., Misc. Myc. I. 28 (1884)	Gaston's actinomma

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Myk. Heft. I. 5 (1817).—Tremella.									
2010	...	...	...	...	...	Q.	B.	Rotten wood	... Tubercles globose to hemispherical, beautiful orange, rather large. Conidia honey coloured.
2011	...	...	...	V.	...	...	...	Bare wood	... Cushion shaped, somewhat hemispherical or irregular, pale. Conidia club shaped, transparent, large, long.
Berl. Mag. III. 10 (1809).—Fusicporium, Selenosporium.									
2012	...	...	...	...	...	Q.	...	Fading leaves of <i>Ficus aspera</i>	Convex, cushion shaped, resembling <i>Hypocrea</i> , somewhat disc shaped, orange.
2013	W.A.	...	...	...	...	...	B.	Branches	... Pustules various, bursting through, brick red. Conidia bow shaped, tapering to each end.
2014	...	...	...	...	...	Q.	...	Twigs of <i>Passiflora</i>	Tufts bursting through, convex, rosy, then whitish. Threads repeatedly forked, transparent.
2015	...	...	...	...	...	Q.	...	<i>Eucalyptus</i> leaves, spreading over galls	Expanded, greyish flesh colour. Conidia elongated. They spread over the leaves and colour the veins with a tint like that of raspberry cream.
Ann. Sci. Nat. X. 359 (1848).									
2016	...	...	...	...	...	Q.	B.	Cocci attached to branches	Small, rather tufted, horn-like to conical, rosy, girt at base by thin whitish membrane. Conidial stage of <i>Spharostilbe</i> .
2017	...	...	...	...	...	Q.	...	Coccus of Orange—scale insect (<i>Chionaspis citri</i>)	Tufts rather spherical, almost sessile, reddish at first, then pale. Conidia elongated, spindle shape.
Berl. Mag. VII. 32 (1816).									
2018	...	...	...	...	...	Q.	...	Leaves and stems of Potato	Gregarious, no spots. Conidia-bearing layer somewhat globose, fleshy, brownish. Conidia brown, rough.
Mich. II. 36 (1880).									
2019	...	...	...	...	...	Q.	...	Decorticated branches	Pustules gregarious, bursting through, rather prominent, resembling <i>Hysterium</i> , black. Conidia olive.
2020	...	...	T.	...	...	...	...	Naked wood	... Conidia-bearing layer circular, plate-like, scattered, superficial, black. Conidia dark olive.
2021	...	...	...	...	N.S.W.	Q.	...	Sugar cane, stalk and leaf	Pustules gregarious, bursting through, black, with short stem-like base. Conidia continuous, dusky.
Meck. I. 25 (1790).									
2022	...	...	...	...	...	Q.	B.	Putrid Agarics, &c.	Pustules disc shaped or variable, dark olive with a white margin. Conidia olive.
2023	...	...	...	...	...	Q.	B.	Old twine	... Pustules disc shaped, then run together and distorted, black with a white margin. Conidia pale olive.
Misc. Myc. I. 28 (1884).									
2024	...	...	...	...	...	Q.	...	Phyllodes of fading <i>Acacia</i>	Gregarious, superficial, black, flattened, contracted when dry, star shaped. Conidia pale brown.

GENERAL CLASSIFICATION OF SPHÆROPSIDES.

GROUP VIII.—SPHÆROPSIDES, LEV.

ARRANGEMENT OF ORDERS (5).

- 49. SPHERIOIDACEÆ—Receptacles black, never fleshy nor brightly coloured, entire.
- 50. NECTRIOIDACEÆ—Receptacles brightly coloured, fleshy or waxy.
- 51. LEPTOSTROMACEÆ—Receptacles more or less distinctly semicircular.
- 52. EXCIPULACEÆ—Receptacles cup shaped, sancer shaped, or Hysterium-like.
- 53. MELANCONIACEÆ—Receptacles absent.

ORDER XLIX.—SPHERIOIDACEÆ, SACC.

ARRANGEMENT OF GENERA (24).

Section 1. Hyalosporæ, Sacc.—Spores hyaline.

Genera (7)—

334. Phyllosticta, Pers.	337. Asteromella, Pass. and Thuem.	339. Dothiorella, Sacc.
335. Phoma, Fries.	338. Chætophoma, Cooke.	340. Cytospora, Ehr.
336. Aposphæria, Berk.		

Section 2. Phæosporæ, Sacc.—Spores olive or sooty brown.

Genera (4)—

341. Sphæropsis, Lev.	343. Capnodiastrum, Speg.	344. Chætomella, Fekl.
342. Coniothyrium, Corda.		

Section 3. Phæodidymæ, Sacc.—Spores uni-septate, brown.

Genus (1)—

- 345. Diplodia, Fries.

Section 4. Hyalodidymæ, Sacc.—Spores uni-septate, hyaline or green.

Genera (5)—

346. Ascochyta, Lib.	348. Actinonema, Fries.	350. Diplodina, West.
347. Robillarda, Sacc.	349. Darluca, Cast.	

Section 5. Phragmosporæ, Sacc.—Spores two or many septate, brown.

Genera (2)—

351. Hendersonia, Berk.	352. Stagonospora, Sacc.
-------------------------	--------------------------

Section 6. Dictyosporæ, Sacc.—Spores two or many septate, wall-like, coloured.

Genus (1)—

- 353. Camarosporium, Schulz.

Section 7. Scolecosporæ, Sacc.—Spores rod shaped, thread-like or elongated, spindle shape, continuous or septate, hyaline or green.

Genera (4)—

354. Septoria, Fries.	356. Phlyctæna, Mont. and Desm.	357. Gamospora, Sacc.
355. Phleospora, Wallr.		

ORDER L.—NECTRIOIDACEÆ, SACC.

ARRANGEMENT OF GENERA (3).

Sub-division 1. Zythiæ, Sacc.—Receptacles nearly globose, Sphæria-like.

Section 1. Hyalosporæ, Sacc.—Spores globose, ovoid or oblong, continuous, hyaline.

Genera (2)—

358. Sphæronamella, Karst. | 359. Aschersonia, Mont.

Section 2. Scolecosporæ, Sacc.—Spores thread-like or rod shaped, continuous or many septate, hyaline.

Genus (1)—

360. Martinella, Cooke and Mass.

ORDER LI.—LEPTOSTROMACEÆ, REHB.

ARRANGEMENT OF GENERA (7).

Section 1. Hyalosporæ, Sacc.—Spores globose, ellipsoid or oblong, continuous, hyaline.

Genera (5)—

361. Leptothyrium, Kunze. | 363. Melasmia, Lev. | 365. Saeidinum, Nees.
362. Piggotia, Berk. and Br. | 364. Actinothecium, Ces. |

Section 2. Scolecosporæ, Sacc.—Spores thread-like or rod shaped, continuous or septate, hyaline.

Genera (2)—

366. Melophia, Sacc. | 367. Leptostromella, Sacc.

ORDER LII.—EXCIPULACEÆ, CORDA.

ARRANGEMENT OF GENERA (2).

Section 1. Hyalosporæ, Sacc.—Spores globose, ellipsoid or oblong.

Genus (1)—

368. Dinemasporium, Lev.

Section 2. Scolecosporæ, Sacc.—Spores filiform, elongated.

Genus (1)—

369. Protostegia, Cooke.

ORDER LIII.—MELANCONIACEÆ, CORDA.

ARRANGEMENT OF GENERA (7).

Section 1. Hyalosporæ, Sacc.—Conidia globose, ovoid or oblong, continuous.

Genera (2)—

370. Glæosporium, Desm. and Mont. | 371. Pestalozziella, Sacc. and Ell.

Section 2. Didymosporæ, Sacc.—Conidia ovoid or oblong, uniseptate.

Genus (1)—

372. Marsonia, Fisch.

Section 3. Phragmosporæ, Sacc.—Conidia oblong or cylindrical, two to many septate.

Genera (4)—

373. Stilbospora, Pers. | 375. Hyaloceras, Dur. and Mont. | 376. Pestalozzia, De Not.
374. Coryneum, Nees. |

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
GROUP VIII.—SPHÆROPSIDES.—					
ORDER XLIX.—SPHÆRIOIDACEÆ,					
334. PHYLLOSTICTA.—Pers.,					
2025	1802	III. 15	<i>P. circumscissa</i>	Cooke, Grev. XI. 150 (1882)	Circular phyllosticta ... (Shot-hole fungus)
2026	1813	X. 5072	<i>P. Cordyliues</i>	Sacc. and Berl. Misc. Myc. II. 36 (1885) ...	Cordylina phyllosticta ...
2027	1805	III. 33	<i>P. Eucalypti</i>	Thuem., Lusit. 374 (1878)	Eucalyptus phyllosticta ...
2028	1812	.. 219	<i>P. fragaricola</i>	Desm., Pl. Crypt. III. 686	Strawberry phyllosticta ..
2029	1807	X. 4881	<i>P. Hardenbergiæ</i>	Cooke and Mass., Grev. XVI. 3 (1887) ...	Hardenbergia phyllosticta ...
2030	1810	.. 4903	<i>P. neurospileæ</i>	Sacc. and Berl. Misc. Myc. II. 37 (1885) ...	Vein-spot phyllosticta ...
2031	...	.. 5066	<i>P. palmicola</i>	Cooke, Grev. XIV. 89 (1886)	Palm-growing phyllosticta ...
2032	1806	.. 4886	<i>P. phyllodiorum</i>	Sacc., Hedw. 156 (1890)	Phyllode phyllosticta... ..
2033	1809	...	<i>P. Platylobii</i>	Cooke and Mass., Grev. XIX. 61 (1891) ...	Platylobium phyllosticta ...
2034	...	...	<i>P. Prostantheræ</i>	Cooke, Grev. XXI. 39 (1892)	Prostanthera phyllosticta ...
2035	1803	III. 31	<i>P. Rosæ</i>	Desm., Exs. 687	Rose phyllosticta
2036	1804	.. 30	<i>P. Ruborum</i>	Sacc., Mich. II. 342 (1882)	Bramble phyllosticta
2037	1808	X. 4979	<i>P. soriformis</i>	Cooke and Mass., Grev., XIX. 47 (1890) ...	Sorus-shaped phyllosticta ...
335. PHOMA.—Fries,					
2038	1828	III. 940	<i>P. allii</i>	Sacc. and Roum., Reliq. Lib. Ser. IV. 79 (1884) ...	Allium phoma
2039	1814	.. 467	<i>P. ampelina</i> = <i>Sphaceloma ampelinum</i>	B. and C., Grev. II. (1873)	Vine phoma (Black spot)
2040	1820	...	<i>P. australis</i>	Cooke, Grev. XV. 17 (1886)	Southern phoma
2041	1829	III. 965	<i>P. Cordylines</i>	Sacc. Syll. III. 162 (1884)	Lily palm phoma
2042	1822	X. 5084	<i>P. Daviesiæ</i>	Cooke and Mass., Grev. XVIII. 7 (1889) ...	Daviesia phoma
2043	1823	.. 5201	<i>P. Diploglottidis</i>	Cooke and Mass., Grev. XVII. 56 (1889) ...	Diploglottis phoma
2044	1818	III. 649	<i>P. eucalyptidæ</i>	Thuem., Lus. 563 (1878)	Eucalyptus phoma
2045	...	.. 923	<i>P. folliculorum</i>	Sacc. Syll. III. 155 (1884)	Follicle phoma
2046	1825	X. 5310	<i>P. Goodeniarum</i>	Cooke and Mass., Grev. XVI. 2 (1887) ...	Goodenia phoma
2047	1830	III. 998	<i>P. graminis</i>	West. in Kieckx. Fl. Fland. I. 441 (1867)...	Grass phoma
2048	1826	.. 793	<i>P. herbarum</i>	West, Exs. 965	Herb phoma
2049	1824	X. 5337	<i>P. Lythri</i>	Cooke and Mass., Grev. XVI. 75 (1888) ...	Lythrum phoma
2050	1817	III. 650	<i>P. Mollerianæ</i>	Sacc. Syll. III. 110 (1884)	Moller's phoma
2051	1831	.. 997	<i>P. nitida</i>	Rob., in Desm. Exs.	Shining phoma
2052	1816	.. 556	<i>P. notla</i>	Berk., Ann. Nat. Hist. 2 Ser. V. 369 (1850) ...	Spurious phoma

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
2025	...	S.A.	T.	V.	N.S.W.	Q.	...	Leaves and fruit of <i>Prunus Armeniaca</i> and <i>P. Cerasus</i> , &c.	Both surfaces, spots circular, reddish brown, finally falling out and leaving leaf as if riddled with shot.
2026	...	...	...	...	...	Q.	...	Leaves of <i>Cordyline terminalis</i>	Spots indistinct, becoming pale. Receptacles on upper surface, point-like, pierced.
2027	...	...	...	V.	...	...	...	Leaves of <i>Eucalyptus globulus</i>	Spots large, irregular, at first dingy brown then whitish, with narrow purple border.
2028	...	S.A.	...	...	...	Q.	...	Strawberry leaves	Spots straggling, becoming bleached, with red margin.
2029	...	...	...	V.	...	...	...	Leaves of <i>Hardenbergia</i>	Spots on both surfaces, various, tawny.
2030	...	...	...	...	...	Q.	...	Leaves of <i>Vitis antarctica</i>	Spots on upper surface, limited by the veins, hence angular, reddish ochrey.
2031	...	...	...	...	...	Q.	...	Palm leaves	Spots on both surfaces, irregular, whitish to ashy. Margin somewhat elevated, brown.
2032	...	...	...	V.	...	...	...	Phyllodes of <i>Acacia</i>	Spots on both surfaces, somewhat circular, whitish, with brown margin.
2033	...	...	...	V.	...	...	...	Leaves of <i>Platylobium</i>	Spots on both surfaces, irregular, pale with narrow brown margin. Receptacles minute, gregarious, black.
2034	...	...	...	V.	...	...	...	Leaves of <i>Prostanthera lasiantha</i>	Spots somewhat circular, pale amber, with raised dark marginal line.
2035	...	...	...	V.	...	...	B.	Rose leaves	Spots rather circular, greenish, then brownish or greyish, with purple border.
2036	...	...	...	V.	...	...	...	Fading leaves of <i>Rubus fruticosus</i>	Spots minute, whitish, often near the veins.
2037	...	...	...	V.	...	...	...	Leaves of some <i>Proteaceæ</i>	Spots on both surfaces, brown, circular, with darker margin.

Novit. Fl. Suec. V. (1819).—Sphærospis.

2038	...	...	...	V.	...	...	...	Scapes of <i>Allium</i>	Receptacles gregarious, spherical, black, very small, obtuse.
2039	...	S.A.	T.	V.	N.S.W.	Q.	...	Vine twigs	Sub-cuticular, Hysterium-like, swollen.
2040	...	...	...	V.	...	...	...	Leaves of <i>Eucalyptus</i>	Spots brownish, elliptical, surrounded by brown line. Receptacles black, point-like, half immersed.
2041	...	...	...	...	...	Q.	...	Old leaves of <i>Crinum pedunculatum</i>	Receptacles numerous, thickly clustered, on under surface, pustular, quite black.
2042	...	...	...	V.	...	...	...	Dead leaves of <i>Daviesia latifolia</i>	Chiefly on under surface. Receptacles very minute, covered, black, forming nebulous spots.
2043	...	...	...	...	...	Q.	...	Fading leaves of <i>Diploglottis Cunninghamii</i>	On under surface, gregarious. Receptacles half immersed, minute, black, pap-like.
2044	...	...	...	V.	...	...	...	Living or fading leaves of <i>Eucalyptus globulus</i>	Receptacles on under surface, scattered, conically elevated, black, minute.
2045	...	...	...	...	...	Q.	...	Follicles of a <i>Marsdenia</i>	Bursting through. Receptacles gregarious, immersed, black, conical, girt by whitish spots.
2046	...	...	...	V.	...	...	...	Fading leaves of <i>Goodenia ovata</i>	Receptacles scattered, dot-like, minute, black, membranous.
2047	...	...	...	...	...	Q.	...	Grass stems (<i>Poa</i>)	Receptacles globose or angular, black, arranged in series, and forming elongated pustules, wrinkled, dark grey.
2048	...	...	...	V.	...	...	B.	Herbaceous stems...	Receptacles gregarious, depressed globose, pap-like, black, everywhere.
2049	...	...	...	V.	...	...	...	Fading leaves of <i>Lathrum hyssopifolia</i>	On upper surface. Receptacles scattered or gregarious, globose, covered, at length bursting through.
2050	...	...	...	V.	...	...	...	Fallen leaves of <i>Eucalyptus globulus</i>	Receptacles on both surfaces, large, thickly clustered, turgid, shining, dark chestnut.
2051	...	...	...	V.	...	...	...	Grass	Scattered, minute, shining. Receptacles hemispherical, white within, covered by epidermis, which splits lengthwise.
2052	...	...	...	...	...	Q.	B.	Dead branches of <i>Platanus</i>	Receptacles spurious, circular, elevated here and there.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
335. PHOMA.—Fries,					
2053	1827	X. 5349	<i>P. plagia</i> ...	Cooke and Mass., Grev. XVII. 55 (1889)	Defined phoma ...
2054	1832	„ 5390	<i>P. portentosa</i> ...	Cooke and Mass., Grev. XVI. 2 (1887)	Monstrous phoma ...
2055	1821	„ 5250	<i>P. purpurea</i> ...	Cooke and Mass., Grev. XV. 97 (1887)	Purple phoma ...
2056	1815	III. 451	<i>P. Rosarum</i> ...	Dur. and Mont., Alg. 604 (1849)	Rose phoma ...
2057	...	„ 994	<i>P. Sacchari</i> ...	Sacc. Syll. III. 166 (1884)	Sugar-caue phoma ...
2058	...	„ 887	<i>P. uvicola</i> ... = <i>Laestadia Bidwellii</i>	B. and C., Grev. II. (1873) (not Arcang.)	Grape-growing phoma (Black rot) ...
2059	1819	X. 5115	<i>P. viminalis</i> ...	Cooke and Mass., Grev. XVI. 75 (1888)	Viminalis phoma ...
336. APOSPHÆRIA.—Berk.,					
2060	1833	X. 5466	<i>A. Leptospermi</i> ...	Cooke, Grev. XIX. 91 (1891)	<i>Leptospermum aposphæria</i> ...
337. ASTEROMELLA.—Pass.					
2061	1834	X. 5489	<i>A. acaciæ</i> ...	Cooke, Grev. XIX. 5 (1890)	<i>Acacia asteromella</i> ...
2062	1835	...	<i>A. epitrema</i> ...	Cooke, Grev. XX. 6 (1891)	<i>Trema asteromella</i> ...
2063	1836	...	<i>A. Homalanthi</i> ...	Cooke and Mass., Grev. XX. 65 (1892)	<i>Homalanthus asteromella</i> ...
338. CHÆTOPHOMA.—Cooke,					
2064	1837	X. 5510	<i>C. eutricha</i> ...	Sacc. and Berl., Misc. Myc. II. 8 (1885)	Well-haired chætophoma ...
339. DOTHIORELLA.—Sacc.,					
2065	1838	X. 5578	<i>D. Amygdali</i> ...	Cooke and Mass., Grev. XIX. 91 (1891)	Almond dothiorella ...
2066	...	„ 5579	<i>D. Eucalypti</i> ...	Sacc. Syll. X. 229 (1892)	<i>Eucalyptus dothiorella</i> ...
2067	1839	„ 5599	<i>D. pericarpica</i> ...	Sacc., Pug. Austr. 15 (1890)	<i>Pericarp dothiorella</i> ...
340. CYTOSPORA.—Ehr.,					
2068	1841	X. 5677	<i>C. verrucula</i> ...	Sacc. and Berl., Misc. Myc. II. 8 (1885)	Warty cytospora ...
2069	1840	III. 1531	<i>C. xanthosperma</i> ...	Fries, S.M. II. 543 (1823)	Yellow-spored cytospora
341. SPHEROPSIS.—Lev.,					
2070	1844	...	<i>S. numerosa</i> ...	Cooke and Mass., Grev. XX. 65 (1892)	Numerous sphæropsis ...
2071	1845	X. 5711	<i>S. phomatoidea</i> ...	Cooke and Mass., Grev. XVIII. 49 (1890)	Phoma-like sphæropsis ...
2072	1843	III. 1649	<i>S. Rosarum</i> ...	Cooke and Ellis, Grev. VI. 2 (1877)	Rose sphæropsis ...
2073	1842	„ 1720	<i>S. Tricorynes</i> ...	Berk. and Br., Linn. Trans. II. 68 (1883)	<i>Tricoryne sphæropsis</i> ...
2074	1846	X. 5734	<i>S. Tritici</i> ...	Cooke and Mass., Grev. XVI. 75 (1888)	Wheat sphæropsis ...
342. CONIOTHYRIUM.—Corda,					
2075	1847	X. 5723	<i>C. olivaceum</i> ...	Bon. in Fekl., Sym. 377 (1875)	Olive coniothyrium ...
2076	1848	„ 5752	<i>C. septorioides</i> ...	Cooke and Mass., Grev. XX. 36 (1891)	Septoria-like coniothyrium ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Novit. Fl. Suec. V. (1819).— <i>Sphæropsis</i> —continued.									
2053	...	...	...	...	...	Q.	...	Palm leaves ..	Spots distinctly defined, glaucous, elliptic or confluent. Receptacles very minute, black.
2054	...	...	...	V.	...	...	...	Cap of <i>Polyporus portentosus</i>	Scattered. Receptacles innate, covered by blackened cuticle, pap-like, black, shining.
2055	...	...	...	...	...	Q.	...	Foliage of Eucalypts and Tristanias	On both surfaces. Spots circular, purple. Receptacles gregarious, half immersed, black, shining.
2056	...	...	...	...	...	Q.	...	Rose twigs	Receptacles rather minute, covered by the unbroken or stellately-split epidermis, dark brown, white within.
2057	...	...	...	...	N.S.W.	...	...	Leaves and stems of Sugar cane	Receptacles nearly spherical, black, bursting through, scattered or gregarious.
2058	...	...	...	V.	...	...	...	Grapes ..	Irregular, bursting through, and then surrounded by the narrow cuticle.
2059	...	...	...	V.	...	...	...	Leaves of <i>Eucalyptus viminalis</i>	On upper surface. Receptacles immersed, bursting through, black, somewhat globose.
Outl. 315 (1860).									
2060	...	...	...	V.	...	...	...	Bark of <i>Leptospermum</i>	Receptacles scattered, bursting through, then superficial, minute, black, pap-like, white within.
and Thuem., in M.U. 1689 (1877).									
2061	...	...	...	V.	...	...	...	Phyllodes of <i>Acacia</i>	Receptacles very numerous, densely crowded, and forming blackish spots, minute.
2062	...	...	...	...	...	Q.	...	Living leaves of <i>Trema aspera</i>	Spots on upper surface, black, somewhat circular. Receptacles minute, rather globose, seated on brown mycelium.
2063	...	...	...	...	...	Q.	...	Leaves of <i>Homalanthus populifolius</i>	Spots somewhat circular, on both surfaces, sooty brown, dotted with minute black receptacles.
Grev. III. 25 (1874).									
2064	...	...	...	...	...	Q.	...	Languid leaves of <i>Custanosperrum australe</i>	Spots black, often running together. Threads of mycelium, sooty brown. Receptacles dot-like, black.
Mich. II. 5 (1882).									
2065	...	...	...	V.	...	...	...	Bark of Peach and Almond	Receptacles inuate, clustered, transversely bursting through, black, opaque, somewhat gelatinous when moist.
2066	...	...	...	V.	...	...	...	Leaves of Eucalypts	Receptacles globose, seated on a sclerotoid body, black, shining. Sclerotia loosely spongy, pale brown within.
2067	...	...	...	...	...	Q.	...	Pericarp of <i>Macrozamia Denisonii</i>	Receptacles in clusters bursting through, black, cushion shaped, tuberculose, globose or angular.
Syl. Berol. 28 (1820).									
2068	...	...	...	...	...	Q.	...	Branches	Receptacles few, immersed, seated on layer soon bursting through, globose or depressed, black.
2069	...	...	...	V.	...	...	...	Branches of <i>Salix Babylonica</i>	Receptacles none. Spores issuing in golden tendrils.
Ann. Sci. Nat. III. 62 (1846).									
2070	...	...	...	V.	...	...	...	Dead bark	Receptacles gregarious, half immersed, globose, black, becoming flattened.
2071	...	...	...	V.	...	...	...	<i>Eucalyptus</i> leaves	On under surface. Receptacles scattered over irregular brown spots, at first covered, black.
2072	...	...	...	...	...	Q.	...	Rose branches	Gregarious or scattered. Receptacles covered splitting the epidermis.
2073	...	...	...	...	...	Q.	...	Leaves of <i>Tricoryne anceps</i>	Receptacles minute, black, immersed in substance of leaf.
2074	...	...	...	V.	...	...	...	Dead leaves and sheaths of Wheat	Receptacles very minute, thickly clustered, at first covered, point-like, black.
Icon. IV. 38 (1854).									
2075	...	...	...	V.	...	...	...	Involucres of <i>Leptospermum lavigatum</i>	Receptacles scattered, at first covered, then bursting through, rather large, pap-like.
2076	...	...	...	V.	...	...	...	Leaves of <i>Prostanthura lasiantha</i>	Spots circular, tawny, with broad purple margin. Receptacles mostly in circles upon spots, black.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
2077	1849	X. 5809	<i>C. orbiculatum</i> ...	Cooke and Mass., Grev. XVIII. 49 (1890)	343. CAPNODIASTRUM.—Speg., Orbicular capnodiastrum ...
2078	1850	III. 1807	<i>C. brachyspora</i> ...	Sacc. and Speg., Mich. I. 260 (1879) ...	344. CHÆTOMELLA.—Fekl., Short-spored chætomella ...
2079	1853	X. 5873	<i>D. canthifolia</i> ...	Cooke and Mass., Grev. XX. 36 (1891) ...	345. DIPLODIA.—Fries, Canthium-leaved diplodia ...
2080	1851	„ 5829	<i>D. lichenopsis</i> ...	Cooke and Mass., Grev. XVI. 2 (1887) ...	Lichen-like diplodia ...
2081	...	...	<i>D. Marsdeniæ</i> ...	Cooke and Mass., Grev. XXI. 75 (1893) ...	Marsdenia diplodia ...
2082	1852	III. 1990	<i>D. phyllodiorum</i> ...	Penz. and Sacc., Fung. Mort. ...	Phyllode diplodia ...
2083	1855	X. 5957	<i>A. apiospora</i> ...	Cooke and Mass., Grev. XV. 98 (1887) ...	346. ASCOCHYTA.— Pear-shaped spored ascochyta ...
2084	1854	„ 5964	<i>A. brunnea</i> ...	Cooke and Mass., Grev. XV. 98 (1887) ...	Brown ascochyta ...
2085	1856	III. 2253	<i>R. sessilis</i> ...	Sacc., Mich. II. 8 (1880) ...	347. ROBILLARDA.—Sacc., Sessile robillarda ...
2086	1857	III. 2257	<i>A. Rosæ</i> ...	Fries, S.V.S. 424 (1849) ...	348. ACTINONEMA.—Fries, Rose actinonema ...
2087	...	III. 2263	<i>D. filum</i> ...	Cast., Cat. Mars. Supp. 53 (1851) ...	349. DARLUCA.—Cast., Cat. Mars. Thread darluca ...
2088	1858	X. 6054	<i>D. Dendrobii</i> ...	Cooke and Mass., Grev. XVI. 3 (1887) ...	350. DIPLODINA.—West, Dendrobium diplodina ...
2089	1859	III. 2320	<i>II. Eucalypti</i> ...	Cooke and Hark., Grev. IX. 128 (1881) ...	351. HENDERSONIA.—Berk., Eucalyptus hendersonia ...
2090	1860	X. 6140	<i>S. orbicularis</i> ...	Cooke, Grev. XX. 6 (1891) ...	352. STAGONOSPORA.—Sacc., Orbicular stagonospora ...
2091	1861	X. 6191	<i>C. Eucalypti</i> ...	Wint., Rev. Myc. 212 (1886) ...	353. CAMAROSPORIUM.—Schulz, Eucalyptus camarosporium ...
2092	1870	III. 3051	<i>S. Bromi</i> ...	Sacc., Mich. I. 194 (1879) ...	354. SEPTORIA.—Fries, Brome septoria ...
2093	1868	X. 6241	<i>S. epiphyllodea</i> ... substituted for <i>S. phyllodi-</i> <i>orum</i>	Cooke, Handb. Aust. Fung. 356 (1892) ... Sacc., Hedw. 156 (1890).	Epiphyllode septoria ...
2094	1866	„ 6242	<i>S. Hardenbergiæ</i> ...	Sacc., Hedw. 156 (1890) ...	Hardenbergia septoria ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Guar. I. 145 (1883).									
2077	...	...	...	...	...	Q.	...	Leathery leaves	Spots circular, of black interwoven mycelium. Receptacles minute, globose, rather membranous.
Sym. 402 (1875).									
2078	...	S.A.	...	V.	...	...	...	Bark, branches of Grape vine	Receptacles scattered, rather superficial, globose, then depressed, black, clad with stiff brown bristles.
S.V.S. 416 (1849).									
2079	...	...	...	V.	...	...	...	Leaves of <i>Canthium latifolium</i>	Receptacles scattered, immersed, membranous, dark brown piercing cuticle.
2080	...	...	...	...	...	Q.	...	Phyllodes of <i>Acacia complanata</i>	Spots brick red, determinate, pale at length, or girt with red zone. Receptacles half-internal, point-like, black.
2081	...	...	...	...	...	Q.	...	Follicles of a <i>Marsdenia</i>	Receptacles gregarious, black, bursting through. In company with <i>Phoma folliculorum</i> , Sacc.
2082	...	...	...	V.	...	...	...	Living or fading phyllodes of <i>Acacia</i>	Receptacles gregarious, minute, under cuticle then bursting through, black, seated on whitish spots.
Lib. Exs. (1837).									
2083	...	...	...	...	...	Q.	...	Leaves of <i>Myrtus</i> and <i>Buckhousia</i>	Spots on upper surface, circular or irregular, tawny, girt by purple zone. Receptacles minute, innate.
2084	...	...	...	...	...	Q.	...	Leaves of Brisbane Box (<i>Tristania conferta</i>)	Spots on both surfaces, circular or irregular, pale brown or ochrey, girt by darker elevated line. Receptacles minute, point-like, black.
Mich. II. 8 (1880).—Pestalozzia.									
2085	...	...	...	...	...	Q.	...	Fading leaves of Vine	Spots small, angular, turning whitish, encircled with red. Receptacles on upper surface.
S.V.S. 424 (1849).—Asteroma.									
2086	...	...	...	V.	...	Q.	B.	Rose leaves	On upper surface. Spots purplish. Receptacle-like tubercles scattered and collapsing, blackish.
Supp. 53 (1851).—Sphaeria, Diplodia.									
2087	...	...	...	...	...	Q.	B.	Leaves of <i>Sorghum</i> and <i>Muehlenbeckia</i> infested with Uredines	Gregarious, very minute. Receptacles globose, black, shining, pierced.
5 Not. 19 (1866).									
2088	...	...	...	...	...	Q.	...	Leaves of <i>Dendrobium speciosum</i>	Receptacles gregarious, innate, black, convex, at length splitting cuticle, shining.
Ann. Nat. Hist. VI. 430 (1841).									
2089	...	...	...	V.	...	...	...	Dead branches and leaves of <i>Eucalyptus</i>	Receptacles in circular spots, immersed.
Mich. II. 8 (1880).									
2090	...	...	...	V.	...	...	...	Dead leaves of <i>Eucalyptus</i>	Spots on both surfaces, small, circular, pale, surrounded by brown line. Receptacles few, in centre of spots, black.
Myk. Beitr. 649 (1870).									
2091	...	...	...	V.	...	...	...	Leaves of <i>Eucalyptus</i>	Receptacles on irregular spots, which are pale brown or grey, limited by darker line.
S.M. III. 480 (1832).									
2092	...	S.A.	...	V.	...	...	...	Leaves of <i>Bromus</i> , &c.	Spots obsolete, becoming pale, elongated. Receptacles plentiful, globose to flattened, pierced.
2093	...	S.A.	...	V.	...	...	...	Phyllodes of <i>Acacia</i>	Spots on both surfaces, circular, whitish, encircled by brown. Receptacles crowded, point-like, becoming black.
2094	...	S.A.	...	...	...	...	...	Leaves of <i>Hardenbergia monophylla</i>	Spots on both surfaces, broad, pale, brown at margin. Receptacles point-like, ochrey.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
354. SEPTORIA.—Fries,					
2095	1869	X. 6429	<i>S. Lepidospermi</i> ...	Cooke and Mass., Grev. XIX. 91 (1891) ...	<i>Lepidosperma septoria</i> ...
2096	1863	„ 6353	<i>S. Martinii</i> ...	Cooke, Grev. XIX. 5 (1890) ...	Martin's septoria ...
2097	1865	„ 6264	<i>S. Myoporii</i> ...	Cooke and Mass., Grev. XVI. 113 (1888)...	<i>Myoporum septoria</i> ...
2098	1864	III. 2683	<i>S. oleandrina</i> ...	Sacc., Fung. Ven. V. 205 (1873-82) ...	<i>Oleander septoria</i> ...
2099	1867	X. 6245	<i>S. phyllodiorum</i> ...	Cooke and Mass., Grev. XIX. 47 (1890) ..	<i>Phyllode septoria</i> ...
2100	...	III. 3042	= <i>S. Martiniana</i> ... <i>S. Tritici</i> ...	Sacc., Syll. X. 351 (1892). Desm. IX., Not. 17 (1842) ...	Wheat septoria ...
2101	1862	„ 2811	<i>S. Violæ</i> ...	West, Exs. Fasc. 2, 91 ...	<i>Violet septoria</i> ...
355. PHLEOSPORA.—Wallr., Fl. Crypt. 7 (1833).—					
2102	...	III. 3136	<i>P. Mori</i> ...	Sacc. Syll. III. 577 (1884) ...	<i>Mulberry phleospora</i> ... (Leaf-spot of Mulberry)
356. PHLYCTÆNA.—Mont. and Desm.					
2103	1871	X. 6518	<i>P. Passifloræ</i> ...	Cooke and Mass., Grev. XVI. 3 (1887) ...	<i>Passion-flower phlyctæna</i> ...
357. GAMOSPORA.—Sacc.,					
2104	1872	X. 6529	<i>G. eriosporoides</i> ...	Sacc. and Berl., Rev. Myc. (1885) ...	<i>Eriospora-like gamospora</i> ...
ORDER L.—					
358. SPILERONÆMELLA.—Karst.,					
2105	1873	III. 3308	<i>S. rufa</i> ...	Sacc. Syll. III. 618 (1884) ...	<i>Red sphæronæmella</i> ...
359. ASCHERSONIA.—Mont.,					
2106	1874	III. 3313	<i>A. tahitensis</i> ...	Mont., Ann. Sci. Nat. 122 (1848) ...	<i>Tahitian aschersonia</i> ...
360. MARTINELLA.—Cooke and Mass.,					
2107	1875	X. 6555	<i>M. Eucalypti</i> ...	Cooke and Mass., Grev. XVIII. 7 (1889) ...	<i>Eucalyptus martinella</i> ...
ORDER LI.—					
361. LEPTOTHYRIUM.—Kunze and					
2108	1877	X. 6567	<i>L. aristatum</i> ...	Cooke, Grev. XX. 6 (1891) ...	<i>Bristly leptothyrium</i> ...
2109	1876	„ 6568	<i>L. Eucalyptorum</i> ...	Cooke and Mass., Grev. XVIII. 7 (1889) ...	<i>Eucalypt leptothyrium</i> ...
362. PIGGOTIA.—Berk. and Br.,					
2110	1878	X. 6596	<i>P. substellata</i> ...	Cooke, Grev. XX. 6 (1891) ...	<i>Stellate piggotia</i> ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
S.M. III. 480 (1832)—continued.									
2095	...	...	...	V.	...	...	...	Leaves of <i>Lepidosperma</i>	Spots on both surfaces, greyish, then white, oblong, with broad brown margin. Receptacles small, black.
2096	...	...	...	V.	...	...	...	Leaves of <i>Senecio Bedfordii</i>	Spots on upper surface, grey, run together, surrounded by black line. Receptacles point-like, globose.
2097	...	...	...	V.	...	...	...	Leaves of <i>Myoporum insulare</i>	Spots on upper surface, circular, whitish, girt by brown line. Receptacles half immersed, point-like, black.
2098	...	...	...	...	...	Q.	...	Leaves of <i>Nerium Oleander</i>	Spots on upper surface, rounded or angular, and run together, turning whitish. Receptacles somewhat large, globose.
2099	...	...	...	V.	..	...	...	Phyllodes of <i>Acacia longifolia</i>	Receptacles closely crowded on both surfaces, without definite spots, often occupying entire surface, immersed, black.
2100	...	...	...	V.	N.S.W.	...	B.	Fading leaves of Wheat, &c., also stem and ear	On both surfaces. Spots linear lengthwise, whitish with dark-purple margin. Receptacles innate, very minute, black.
2101	...	...	...	V.	...	...	B.	Fading Violet leaves	Receptacles minute, numerous, brownish yellow, seated on pale zoned circular spots, girt by reddish-brown ring.

Septoria, Sphærella, Fusarium, Fusisporium.

2102	...	...	...	V.	...	...	B.	Leaves of Mulberry	Spots whitish or ochrey, surrounded by brown. Receptacles innate, globose, gregarious.
------	-----	-----	-----	----	-----	-----	----	--------------------	--

Ann. Sci. Nat. 16 (1847).

2103	...	...	...	...	...	Q.	...	Twigs of Passiflora growing on stem	Receptacles very thickly clustered, minute, innate, at length bursting through.
------	-----	-----	-----	-----	-----	----	-----	-------------------------------------	---

Rev. Myc. (1885).

2104	...	...	...	...	...	Q.	...	Languid leathery leaves	Receptacles on upper surface, interspersed on thin spot-like brown mycelium, point-like, globose to depressed.
------	-----	-----	-----	-----	-----	----	-----	-------------------------	--

NECTRIOIDACEÆ, SACC. SYLL. III. 613 (1884).

Hedw. 17 (1884).—Sphæronæma.

2105	...	...	T.	...	...	...	...	Pine chips	Receptacles awl shaped, acute, reddish brown, paler downwards.
------	-----	-----	----	-----	-----	-----	-----	------------	--

Ann. Sci. Nat. 3 Ser. X. 121 (1848).

2106	...	...	...	...	...	Q.	...	Leaves of climber...	Receptacles minute, seated on hemispherical obtuse yellow layer.
------	-----	-----	-----	-----	-----	----	-----	----------------------	--

Grev. XVIII. 7 (1889).

2107	...	...	...	V.	...	...	...	Leaves of <i>Eucalyptus</i>	On upper surface. Receptacles very minute, immersed, cracked at mouth, seated on rather circular fleshy reddish-brown layer.
------	-----	-----	-----	----	-----	-----	-----	-----------------------------	--

LEPTOSTROMACEÆ, RCHB. NOM. GEN. 6 (1841).

Schm., Myk. Heft. II. 79 (1823).

2108	...	...	...	V.	...	...	...	Dead leaves of <i>Eucalyptus</i>	Receptacles scattered, superficial, circular, dark brown. Spores with oblique bristle at one end.
2109	...	...	...	V.	...	...	...	Fallen leaves of <i>Eucalyptus</i>	Receptacles scattered over bleached spots, shield shaped, flattened, black, dehiscing in middle with star-like fissure.

Ann. Nat. Hist. VII. 2 Ser. 95 (1851).

2110	...	...	...	V.	...	...	...	Leaves of <i>Eucalyptus</i>	On under surface, forming small somewhat circular stellate black patches, composed of flattened receptacles run together.
------	-----	-----	-----	----	-----	-----	-----	-----------------------------	---

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
363. MELASMIA.—Lev.,					
2111	1879	X. 6602	M. Eucalypti ...	Cooke and Mass., Grev. XVI. 75 (1888) ...	Eucalyptus melasmia ...
2112	...	...	M. Tecomatis ...	Cooke and Mass., Grev. XXII. 37 (1893) ...	Tecoma melasmia ...
364. ACTINOTHECIUM.—Ces.,					
2113	1879 bis	III. 3398	A. Scortechinii ...	Sacc. and Berl., in Sacc. Syll. III. 639 (1884) ...	Scortechini's actinothecium ...
365. SACIDIUM.—Nees, in Kunze and Schm.,					
2114	1881	X. 6615	S. Camelliae ...	Cooke and Mass., Grev. XVI. 3 (1887) ...	Camellia sacidium ...
2115	1880	„ 6616	S. Eucalypti ...	Cooke and Mass., Grev. XVI. 75 (1888) ...	Eucalyptus sacidium ...
366. MELOPHIA.—Sacc.					
2116	1883	X. 6643	M. Leptospermi ... = M. Victoriae ...	Cooke, Grev. XX. 65 (1892) ... Sacc. Syll. X. 428 (1892).	Leptospermum melophia ...
2117	1882	III. 3512	M. Woodsiana ...	Sacc. and Berl., in Sacc. Syll. III. 659 (1884) ...	Woodsio's melophia ...
367. LEPTOSTROMELLA.—Sacc.,					
2118	1884	X. 6652	L. Eucalypti ...	Cooke and Mass., Grev. XIX. 91 (1891) ...	Eucalypt leptostromella ...
ORDER LII.—					
368. DINEMASPORIUM.—Lev.,					
2119	1885	III. 3619	D. hispidulum ...	Sacc., Mich. II. 281 (1882) ...	Hispid dinemasporium...
369. PROTOSTEGIA.—					
2120	1886	X. 6715	P. Eucalypti ...	Cooke and Mass., Grev. XVI. 75 (1888) ...	Eucalypt protostegia ...
ORDER LIII.—					
370. GLÆOSPORIUM.—Desm. and Mont.,					
2121	...	...	G. Alphitoniæ ..	Cooke and Mass., Grev. XXII. 37 (1893) ...	Alphitonia glæosporium ..
2122	1900	III. 3755	G. ampelophagum ...	Sacc., Mich. I. 217 (1879) ...	Grape-destroying glæosporium ..
2123	1887	X. 6737	G. Citri ...	Cooke and Mass., Grev. XIX. 92 (1891) ...	Citrus glæosporium ...
2124	1889	„ 6739	G. citricolum ...	Cooke and Mass., Grev. XVI. 3 (1887) ...	Citrus-growing glæosporium ...
2125	1895	„ 6786	G. Denisonii ..	Sacc. and Berl., Misc. Myc. II. 10 (1885)...	Denison's glæosporium ...
2126	1894	„ 6819	G. epiciadæi ...	Cooke and Mass., Grev. XIX. 92 (1891) ...	Cladium glæosporium ...
2127	1898	III. 3751	G. fructigenum ...	Berk., Gard. Chron. 245 (1856) ...	Fruit glæosporium ... (Ripe rot)
2128	1891	X. 6787	G. glaucum ...	Cooke and Mass., Grev. XVI. 75 (1888)...	Glaucous glæosporium ...
2129	1892	„ 6726	G. Hedycaryæ ...	Cooke and Mass., Grev. XVIII. 7 (1889) ...	Hedycarya glæosporium ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Ann. Sci. Nat. 276 (1846).									
2111	...	...	...	V.	...	Q	...	Leaves of <i>Eucalyptus</i>	Spots circular or confluent, black. Receptacles few, somewhat gregarious, elliptical, opening by a fissure.
2112	...	...	...	...	...	Q.	...	Leaves of <i>Tecoma jasminoides</i>	Receptacles on both surfaces, superficial, circular, wrinkled, black, disc brownish.
Hedw. I. (1852).									
2113	...	...	...	...	...	Q.	...	Leaves of <i>Smilax</i> ...	Receptacles linear, straight or curved, somewhat superficial, forked or variously branched, furrowed lengthwise.
Myc. Heft. II. 64 (1823).									
2114	..	...	..	V.	...	...	...	Fading leaves of <i>Camellia</i>	Receptacles scattered, superficial, hemispherical, black, opaque, mostly minute.
2115	...	...	...	V.	...	...	...	Dead leaves of <i>Eucalyptus globulus</i>	On both surfaces. Receptacles gregarious, bursting through, small, flattened, black.
Syll. III. 658 (1884).									
2116	...	...	...	V.	...	...	...	Leaves of <i>Leptospermum laevigatum</i>	Receptacles scattered on both surfaces, convex, flattened at base, black, white within.
2117	...	...	...	...	...	Q.	...	Phyllodes of <i>Acacia harpophylla</i>	Receptacles distantly scattered, inserted in a thin white filamentous spot-like mass.
Mich. III. 632 (1882).									
2118	...	...	...	V.	...	...	...	Fading leaves of <i>Eucalyptus</i> of	Spots somewhat circular, on both surfaces, reddish brown, then sooty brown. Receptacles scattered over spots, black.

EXCIPULACEÆ, CORDA, IC. FUNG. V. 35 (1842).

Ann. Sci. Nat. 274 (1846).—Peziza, Polynema, Excipula.

2119	W.A.	...	...	...	...	...	...	Wood ...	Receptacles gregarious or scattered, rather large, cup shaped, black long rigid straight hairs. (No. 1740 wrongly entered as this species.)
Cooke, Grev. IX. 19 (1880).									
2120	...	...	...	V.	...	...	...	Dead leaves of <i>Eucalyptus incrassatus</i>	Receptacles immersed, cup shaped, gelatinous, orange coloured, covered by epidermis, at length split.

MELANCONIACEÆ, CORDA, IC. FUNG. V. 33 (1842).

Ann. Sci. Nat. XII. 295 (1849).—Ramularia, Fusarium.

2121	...	...	...	...	...	...	Q.	Leaves of <i>Alphitonia excelsa</i>	Spots irregular or confluent, pale. Pustules bursting through, small, gregarious on spots.
2122	...	...	...	V.	...	...	...	Grapes, rarely vine leaves or branches	Spots rather circular, often run together.
2123	...	...	...	V.	...	...	...	Branches of Lemon	Gregarious, bursting through, pale sooty brown. Pustules rather small, often run together.
2124	...	...	...	...	...	...	Q.	Orange leaves	Spots dark brown, small, rather disc-like, often run together. Pustules immersed.
2125	...	...	...	...	...	...	Q.	Leaves of <i>Encephalartos Denisonii</i>	Pustules gregarious, minutely pustulate, covered by epidermis hardly broken, yellowish within.
2126	...	...	...	V.	...	...	...	<i>Gahnia tetraquetra</i> ...	Pustules gregarious in centre of irregular spots, caused by blackened cuticle.
2127	...	...	...	...	...	...	Q. B.	Pears	Pustules concentric, dull rose colour bursting through, with single pore or fringed mouth.
2128	...	...	...	...	...	...	Q.	Living leaves	Spots rather circular on one or both surfaces, becoming glaucous, rather mealy.
2129	...	...	...	V.	...	...	...	Fading leaves of <i>Hedyocarya Cunninghamii</i>	On upper surface. Spots circular, turning black. Pustules solitary or gregarious.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
370. GLOEOSPORIUM.—Desm. and Mont.,					
2130	1888	III. 3675	<i>G. intermedium</i>	Sacc., Mich. II. 118 (1882)	Intermediate gloeosporium
2131	1902	„ 3757	<i>G. lagenarium</i>	Sacc. and Roum., Rev. Myc. 201 (1880)	Lagenaria gloeosporium
2131a	1903	„ 3758	<i>G. lagenarium</i> , var. <i>Cucurbitarum</i>	Cooke, Handb. Austr. Fung. 363 (1892)	Gourd gloeosporium
2132	1897	„ 3748	<i>G. leguminis</i>	Cooke and Hark., Grev. IX. 7 (1889)	Legume gloeosporium
2133	1896	„ 3747	<i>G. Lindemuthianum</i> = <i>Colletotrichum</i>	Sacc. and Magn., Mich. I. 129 (1878)	Lindemuth's gloeosporium (Bean spot)
2134	1899	X. 6811	<i>G. Musarum</i>	Cooke and Mass., Grev. XVI. 3 (1887)	Musa gloeosporium
2135	1893	„ 6748	<i>G. nigricans</i>	Cooke and Mass., Grev., XIX. 91 (1891)	Blackening gloeosporium
2136	1901	„ 6733	<i>G. pestiferum</i>	Cooke and Mass., Grev. XIX. 61 (1891)	Pestiferous gloeosporium
2137	1890	„ 6801	<i>G. subglobosum</i>	Cooke and Mass., Grev. XV. 3 (1887)	Sub-globose gloeosporium
2138	...	III. 3752	<i>G. versicolor</i>	Berk. and Curt., Grev. III. (1874)	Colour-changing gloeosporium ... (Bitter rot)
371. PESTALOZZIELLA.—Sacc. and Ellis,					
2139	1904	X. 6858	<i>P. circularis</i>	Cooke and Mass., Grev. XVIII. 80 (1890)	Circular pestalozziella
372. MARSONIA.—Fisch.,					
2140	1905	X. 6884	<i>M. Acaciæ</i>	Cooke and Mass., Grev. XIX. 47 (1890)	Acacia marsonia
2141	1906	...	<i>M. deformans</i>	Cooke and Mass., Grev. XIX. 62 (1891)	Deforming marsonia
373. STILBOSPORA.—Pers.,					
2142	1907	X. 6904	<i>S. foliorum</i>	Cooke, Grev. XX. 6 (1891)	Leaf stilbospora
374. CORYNEUM, Nees, Syst.					
2143	1908	X. 6911	<i>C. viminalis</i>	Cooke and Mass., Grev. XX. 36 (1891)	Viminalis coryneum
375. HYALOCERAS.—Dur. and					
2144	1909	X. 6925	<i>H. diflophosporum</i>	Cooke, Grev. XIX. 5 (1890)	Triseptate-spored hyaloceras
376. PESTALOZZIA.—De Not.,					
2145	1911	III. 4110	<i>P. Acaciæ</i>	Thuem., Lusit. 576 (1878)	Acacia pestalozzia
2146	1914	X. 6951	<i>P. Casuarinæ</i>	Cooke and Mass., Grev. XVIII. 114 (1888)	Sheoak pestalozzia
2147	1913	III. 4135	<i>P. funerea</i>	Desm., Ann. Sci. Nat. XIX. 235 (1843)	Gloomy pestalozzia
2148	...	„ 4146	<i>P. Guepini</i>	Desm., Ann. Sci. Nat. XIII. 182 (1840)	Camellia-leaf fungus
2149	1915	„ 4161	<i>P. monochaeta</i>	Desm., Ann. Sci. Nat. 3 Ser. X. 355 (1848)	One-haired pestalozzia
2150	1910	„ 4128	<i>P. uvicola</i>	Speg., in Thuem. Pilz. Min. 13 (1878)	Grape pestalozzia
2151	1912	„ 4134	<i>P. versicolor</i>	Speg., in Sacc. Mich. I. 479 (1879)	Parti-coloured pestalozzia

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T	V.	N.S.W.	Q.			
Ann. Sci. Nat. XII. 295 (1849).— <i>Ramularia</i> , <i>Fusarium</i> — <i>continued</i> .									
2130	...	...	...	...	...	Q.	...	Leaves of <i>Hoya australis</i>	Pustules gregarious, point-like, black, then bursting through.
2131	..	...	...	...	...	Q.	...	Epicarp of Melons, Mango fruit, &c.	Pustules beneath cuticle, bursting through, minute, cushion shaped, somewhat rosy.
2131A	...	...	...	...	...	Q.	...	On Gourds, Bananas, and Melons	Spots bright orange, depressed. Conidia club-shaped, shortly stalked.
2132	...	...	...	V.	...	...	...	Legumes of <i>Acacia melanoxylon</i>	Scattered, covered by cuticle. Conidia oval, transparent.
2133	...	...	...	V.	N.S.W.	Q.	B	Legumes of Bean, Pea, &c.	Spots on fruit, rarely on stem or leaves, roundish, bleached, at first with reddish-brown margin.
2134	...	...	...	...	...	Q.	...	Ripe Bananas, rendering them dry and insipid	Pustules innate, bursting through, gregarious, rather rosy.
2135	...	...	...	V.	...	...	...	Leaves of <i>Eucalyptus pauciflora</i>	Without distinct spots, on both surfaces. Pustules densely clustered, becoming black, convex.
2136	...	...	...	V.	...	Q.	...	Twigs, flower stalks, and fruit of Vine	Pustules gregarious, small, discoid, convex, rosy.
2137	...	...	...	V.	...	...	...	Fading leaves of <i>Goodenia ovata</i>	Pustules scattered, pale, inconspicuous. Conidia sub-globose.
2138	...	...	...	V.	N.S.W.	...	...	Rotting Apples	Spots brown, small, circular, running together. Pustules bursting through, arranged in rings.
Mich. II. 575 (1882).									
2139	...	...	...	V.	...	...	...	Dead leaves of <i>Eucalyptus pauciflora</i>	On both surfaces. False receptacles usually arranged in circles, at first brown, then nearly black and shining.
in Rab. Fl. Eur. No. 1857.									
2140	...	...	..	V.	...	...	...	Phyllodes of <i>Acacia</i>	Spots irregular or run together, pale or whitish, with brown margin. Pustules gregarious on the spots.
2141	..	S.A.	..	V.	...	...	...	Cultivated Peas, chiefly on leaves, stipules, leaf stalks, &c.	Pustules gregarious, often run together, brown, distorting the foliage, sometimes on large discoloured spots.
Syn. Fung. 96 (1801).									
2142	...	...	...	V.	...	...	...	Dead leaves of <i>Eucalyptus</i>	Pustules in circular paler spots, splitting the cuticle with three or four openings.
Pilz. 34 (1816).									
2143	...	...	...	V.	...	..	...	Leaves of <i>Eucalyptus viminalis</i>	Pustules point-like, flattened, scattered, black, not seated on definite spots.
Mont. Fl. Alg. 587 (1849).									
2144	...	...	...	V.	...	...	...	Leaves of <i>Leptospermum scoparium</i>	Pustules gregarious, minute, brown, bursting through, splitting irregularly in centre.
Mier. Ital. II. (1842).— <i>Coryneum</i> .									
2145	...	...	...	V.	...	...	...	Living <i>Acacia</i> leaves	On under surface. Pustules gregarious or solitary, hemispherical, seated on irregular dirty ochre spots, with broad rusty margin.
2146	...	...	...	V.	...	...	...	Branches of <i>Casuarina</i>	Pustules gregarious, minute, elliptic, encircled by ruptured epidermis.
2147	...	...	...	...	...	Q.	B.	Leaves of <i>Eleodendron</i> and <i>Myrtus</i>	Pustules scattered, point-like, black, covered by epidermis, then bursting through.
2148	...	...	...	...	...	Q.	B	Foliage of <i>Alphitonia excelsa</i> (Red Ash)	Pustules minute, point-like, convex, black, covered, then bursting through.
2149	...	...	...	...	...	Q.	...	Leaves of <i>Eucalyptus</i>	Pustules scattered or gregarious, often on under surface. Spots variable, becoming stained with black.
2150	...	...	...	...	N.S.W.	Q.	...	Vine leaves, Grapes, and Mangos	Pustules globose, then lens shaped, black, beneath cuticle, bursting through.
2151	...	...	..	...	...	Q.	...	Leaves of <i>Cupania anacardioides</i>	Pustules somewhat lens shaped, covered, then bursting through, causing surrounding parts to blacken.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
---------	-----------------	--------------------	------------------	---------------------	---------------

GENERAL CLASSIFICATION

GROUP IX.—

ORDER LIV.—SACCHAROMYCETACEÆ—Unicellular, multiplying by budding and by ascospores.

GROUP IX.—SACCHAROMYCETES,

ORDER LIV.—SACCHAROMYCETACEÆ,

377. SACCHAROMYCES.—Meyen, in Wieg.

2152	2028	VIII. 3632	S. apiculatus	...	...	Reess, Bot. Unt. 84 (1870)	...	...	...	Apiculate yeast	...	...
2153	2026	„ 3620	S. Cerevisiæ	...	...	Meyen, in Wieg. Archiv. IV. 109 (1883)...	...	...	...	Beer yeast	...	...
2154	2027	„ 3621	S. ellipsoideus	...	...	Reess, Bot. Unt. 82 (1870)	...	...	...	Elliptic yeast	...	...
2155	2029	„ 3625	S. Mycoderma	...	...	Reess, Bot. Unt. 83 (1870)	...	...	...	Scum yeast	...	...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			

OF SACCHAROMYCETES.

SACCHAROMYCETES, REESS.

Genus (1)—

377. *Saccharomyces*, Meyen.

REESS, BOT. UNT. (1870).

REESS, BOT. UNT. (1870).

Arch. IV. 2 (1838).—*Mycoderma*, *Torula*.

2152	...	...	...	...	...	...	B.	In fermentation of wine	Cells lemon shaped, shortly apiculate at each end, rarely united in small scarcely-branched colonies.
2153	...	...	...	...	...	...	B.	In beer ...	Cells mostly round or oval, solitary or united in small colonies.
2154	...	...	...	...	...	...	B.	Producing spontaneous fermentation in must	Cells elliptical, solitary or united in little branched colonies.
2155	...	...	...	...	...	...	B.	On fermented fluids, &c.	Cells oval, elliptical or cylindrical, united in very much branched colonies.

GENERAL CLASSIFICATION OF USTILAGINES.

GROUP X.—USTILAGINES, TUL.

ORDER LV.—USTILAGINACEÆ—Parasitic. Mycelium soon disappearing. Spores virtually all unicellular.

ARRANGEMENT OF GENERA (11).

Section 1. *Americosporæ*, Sacc. and De Toni—Spores continuous, sub-solitary.

Genera (4)—

378. <i>Ustilago</i> , Pers.		380. <i>Entyloma</i> , De Bary.		381. <i>Sphacelotheca</i> , De Bary.
379. <i>Tilletia</i> , Tul.				

Section 2. *Dietyosporæ*, Sacc. and De Toni—Spores agglomerated.

Genera (4)—

382. <i>Doassansia</i> , Cornu.		384. <i>Sorosporium</i> , Rud.		385. <i>Urocystis</i> , Rabh.
383. <i>Thecaphora</i> , Fing.				

Exceptional—Genera (3)—

386. <i>Graphiola</i> , Poit.		387. <i>Cerebella</i> , Ces.		388. <i>Schinzia</i> , Næg.
-------------------------------	--	------------------------------	--	-----------------------------

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
<p style="text-align: right;">GROUP X.—USTILAGINES, ORDER LV.—USTILAGINACEÆ, 378. USTILAGO.—Pers.,</p>					
2156	1701	VII. 1657	<i>U. australis</i> ...	Cooke, Grev. VIII. 34 (1879) ...	Southern ustilago ...
2157	1705	IX. 1172	<i>U. axieola</i> ...	Berk., Ann. Nat. Hist. 2 Ser. IX. 200 (1852) ...	Axis-growing ustilago ...
2158	1710	VII. 1677	<i>U. bromivora</i> ...	Waldh., Ustil. 215 (1877) ...	Brome-destroying ustilago ...
2159	1713	„ 1704	<i>U. bullata</i> ...	Berk., Fl. N. Zeal. II. 196 (1855) ...	Blistered ustilago ...
2160	...	„ 1726	<i>U. bursa</i> ...	Berk., Hook., Journ. 206 (1854) ...	Purse ustilago ...
2161	...	...	<i>U. catenata</i> ...	Ludw., Zeitsch. f. Pflkrk. III. 139 (1893) ...	Chain ustilago ...
2162	1716	VII. 1728	<i>U. Cesatii</i> ...	Waldh., Ustil. 25 (1877) ...	Cesati's ustilago ...
2163	...	...	<i>U. comburens</i> ...	Ludw., Zeitsch. f. Pflkrk. III. 139 (1893) ...	Burning ustilago ...
2164	1702	...	<i>U. confusa</i> ...	Mass., Grev. XX. 65 (1892) ...	Confused ustilago ...
2165	1703	VII. 1645	<i>U. destruens</i> ...	Schlecht., Berol. 130 (1823) ...	Destructive ustilago ...
2166	...	„ 1644	<i>U. Digitariæ</i> ...	Rabh., Fung. Eur. 1199 ...	Digitaria ustilago ...
2167	1714	„ 1712	<i>U. emodensis</i> ...	Berk., Hook., Journ. III. 202 (1851) ...	Dark-lilac ustilago ...
2168	1707	„ 1671	<i>U. leucoderma</i> ...	Berk., Ann. Nat. Hist., 2 Ser. IX. 200 (1852) ...	White-skinned ustilago ...
2169	1708	„ 1675	<i>U. marmorata</i> ...	Berk., Linn. Journ. XIII. 174 (1873) ...	Marbling ustilago ...
2170	...	„ 1723	<i>U. maydis</i> ...	Corda, Icon. V. 3 (1854) ...	Maize ustilago ...
2171	1704	„ 1664	<i>U. Muelleriana</i> ...	Thuem., Myc. Univ. 623 (1879) ...	Mueller's ustilago ...
2172	1706	„ 1665	<i>U. pilulæformis</i> ...	Tul., Ann. Sci. Nat. 93 (1847) ...	Pill-shaped ustilago ...
2173	1709	„ 1676	<i>U. segetum</i> ...	Ditm., in Sturm's Deutsch. Fl. (1817-51) ...	Corn ustilago ...
2173A	...	...	<i>U. segetum</i> , var. <i>tritici</i> ...	Jensen, Journ. Roy. Ag. Soc. Eng. 407 (1888) ...	Wheat ustilago ...
2173B	...	...	<i>U. segetum</i> , var. <i>avenæ</i> ...	Jensen, Journ. Roy. Ag. Soc. Eng. 407 (1888) ...	Oat ustilago ...
2173C	...	...	<i>U. segetum</i> , var. <i>nuda hordei</i> ...	Jensen, Journ. Roy. Ag. Soc. Eng. 407 (1888) ...	Naked Barley ustilago ...
2174	...	...	<i>U. Spinificis</i> ...	Ludw., Zeitsch. f. Pflkrk. III. 138 (1893) ...	Spinifex ustilago ...
2175	1712	...	<i>U. Tepperi</i> ...	Ludw., Bot. Centr. 341 (1889) ...	Tepper's ustilago ...
2176	1717	VII. 1737	<i>U. utriculosa</i> ...	Tul., Mem. Ust. 102 (1847) ...	Swelling ustilago ...
<p style="text-align: right;">379. TILLETIA.—Tul., Ann.</p>					
2177	1719	VII. 1783	<i>T. epiphylla</i> ...	Berk. and Br., Linn. Trans. II. 67 (1883) ...	Epiphyllous tilletia ...
2178	1718	„ 1760	<i>T. tritici</i> ...	Winter, Die Pilze 110 (1884) ...	Wheat tilletia ...
<p style="text-align: right;">380. ENTYLOMA.—De Bary.</p>					
2179	1720	...	<i>E. Eugéniarum</i> ...	Cooke and Mass., Grev. XIX. 92 (1891) ...	Eugenia entyloma ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
TUL., ANN. SCI. NAT. BOT. 14 (1847).									
TUL., ANN. SCI. NAT. BOT. 14 (1847).									
Syn. 224 (1808).—Uredo, Tilletia, Caeoma, Cintractia.									
2156	...	...	...	V.	...	...	...	Spikelets of <i>Eriachne</i>	Produced within the ovaries. Spores black, somewhat globose or deformed.
2157	...	...	...	V.	...	Q.	...	Fruits and panicles of <i>Cyperus</i> and <i>Fimbristylis</i>	Little dusty irregular balls in axis of lower spikelets. Spores rather pellucid.
2158	...	S.A.	...	V.	N.S.W.	Q.	B.	<i>Bromus mollis</i> and <i>arenarius</i> , <i>Anthistiria ciliata</i>	Produced in inflorescence. Pustules dark brown, soon powdery.
2159	...	S.A.	...	V.	N.S.W.	...	...	Inflorescence of <i>Triticum</i>	Pustules black. Spores very pale olive brown.
2160	...	...	...	...	...	Q.	...	Grain of <i>Anthistiria frondosa</i>	Pustules greenish. Spores brownish black.
2161	...	S.A.	...	...	...	...	...	Spikes of <i>Cyperus lucidus</i>	Pustules crumb-like, ashy-black spores joined in a chain.
2162	...	...	...	V.	...	Q.	...	<i>Paspalum scrobiculatum</i>	Pustules black. Spores dark brown.
2163	...	S.A.	...	...	...	...	...	Species of <i>Stipa</i> ...	Pustules black, powdery, in stems and panicles which are almost totally destroyed.
2164	...	...	...	V.	...	...	...	<i>Panicum paradoxum</i>	Pustules produced in ovary, soon naked. Mass of spores powdery, violet black.
2165	...	...	...	V.	...	...	...	<i>Danthonia</i> ...	Pustules black, powdery, blackening flowers and panicles, and destroying ovaries.
2166	...	...	...	V.	...	...	...	<i>Panicum</i> ...	Pustules black. Spores brown to orange.
2167	...	...	...	...	...	Q.	...	Stems, &c., of <i>Polygonum</i>	Pustules lobate. Spores dark lilac.
2168	...	...	...	V.	...	Q.	...	Sheaths of <i>Carex</i> , <i>Danthonia</i> , &c.	Pustules black, seated on large spots, covered by whitish crust.
2169	...	S.A.	...	V.	...	...	...	Leaves of <i>Scirpus prolifer</i>	Compact. Marbling the yet unbroken epidermis.
2170	...	...	...	...	N.S.W.	...	B.	Indian Corn (<i>Zea Mays</i>)	Brown in mass with tinge of olive. Spores pale brown, warty.
2171	...	S.A.	...	V.	...	...	...	Seeds of <i>Juncus planifolius</i>	Spores at length clustered together, brown.
2172	...	...	...	V.	...	...	...	Ovaries of <i>Juncus</i>	Compact, black. Spores black.
2173	...	S.A.	...	V.	N.S.W.	Q.	B.	<i>Aristida</i> , <i>Danthonia</i>	Pustules black to olive brown, powdery, covered by soon ruptured epidermis.
2173A	...	S.A.	...	V.	N.S.W.	Q.	B.	Wheat ...	} Spores of one variety do not germinate on the host-plant of another variety.
2173B	...	S.A.	...	V.	N.S.W.	Q.	B.	Oats ...	
2173C	...	S.A.	...	V.	N.S.W.	Q.	B.	Barley ...	
2174	...	S.A.	...	...	...	...	...	Flowers and spikes of <i>Spinifex hirsutus</i>	Pustules olive, destroying the ovaries. Spores grey olive.
2175	...	S.A.	...	...	...	...	...	<i>Amphipogon strictus</i> , <i>Neurachnes alopecuroides</i> , and <i>Danthonia penicillata</i>	Spores powdery, black, destroying flowers and upper portion of stems.
2176	...	S.A.	...	V.	...	...	B.	Ovaries and stems of <i>Polygonum minus</i> and <i>P. gracile</i>	Pustules dark violet, turning violet brown, powdery, causing blossoms to swell.

Sci. Nat. 112 (1847).—Uredo, Ustilago, Lycoperdon, Caeoma.

2177	...	...	...	...	...	Q.	...	Leaves of Maize ...	Pustules short. Spores brown.
2178	...	S.A.	T.	V.	N.S.W.	Q.	B.	Grains of Wheat ...	Pustules olive black, odour of stinking fish, always covered by epidermis, soon powdery.

Bot. Zeit. 101 (1874).

2179	...	...	...	...	...	Q.	...	Leaves of <i>Eugenia</i>	Pustules irregular, dark brown, flattened, rounded, or confluent, in large patches.
------	-----	-----	-----	-----	-----	----	-----	--------------------------	---

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
381. SPHACELOTHECA.—De Bary,					
2180	1721	VII. 1834	<i>S. hydropiperis</i> ...	<i>De Bary</i> , <i>Vergl. Morph.</i> 187 (1884) ...	Hydropiper sphacelotheca ...
2180A	"	...	<i>S. hydropiperis</i> , var. <i>columellifera</i>	<i>Berk.</i> , in <i>Cooke Handb. Anstr. Fung.</i> 327 (1892)...	Columella bearing sphacelotheca
382. DOASSANSIA.—Cornu.,					
2181	1722	VII. 1847	<i>D. punctiformis</i> ...	<i>Winter</i> , <i>Rev. Myc.</i> 207 (1886) ...	Point-like doassausia ...
383. THECAPHORA.—Fing.,					
2182	1723	VII. 1861	<i>T. inquinans</i> ..	<i>Berk. and Br.</i> , <i>Linn. Journ.</i> XIV. 94 (1875) ...	Defiling thecaphora ...
2183	1724	" 1868	<i>T. Leptocarpi</i> ...	<i>Berk.</i> , <i>Linn. Journ.</i> XVIII. 388 (1881) ...	Leptocarpus thecaphora ...
384. SOROSPORIUM.—Rud.,					
2184	1725	VII. 1885	<i>S. Eriachnes</i> ...	<i>Thuem.</i> , <i>Symb. Austr.</i> II. 4 (1878) ...	Eriachoe sorosporium ...
2185	1726	" 1884	<i>S. Muellerianum</i> ...	<i>Thuem.</i> , <i>Symb. Austr.</i> II. 5 (1878) ...	Mueller's sorosporium ...
385. UROCYSTIS.—Rabh., Klotzsch.					
2186	...	VII. 1891	<i>U. occulta</i> ...	<i>Rabh.</i> , <i>Klotzsch, Herb. Myc.</i> II. 393 (1860) ...	Hidden urocystis ...
2187	1727	" 1910	<i>U. solida</i> ...	<i>Waldh.</i> , <i>Ustil.</i> 38 (1877) ...	Compact urocystis ...
386. GRAPHIOLA.—Poit., Ann.					
2188	1728	VII. 1915	<i>G. Phœnicis</i> ...	<i>Poit.</i> , <i>Ann. Sci. Nat.</i> 473 (1824) ...	Date graphiola ...
387. CEREBELLA.—Ces.,					
2189	1730	VII. 1919	<i>C. Andropogonis</i> ...	<i>Ces.</i> , <i>Klotzsch, Herb.</i> 1587 (1851) ...	Andropogon cerebella ...
2190	1729	" 1920	<i>C. Paspali</i> ...	<i>Cooke and Mass.</i> , <i>Grev.</i> XVI. 20 (1887) ...	Paspalum cerebella ...
388. SCHINZIA.—Nægeli,					
2191	...	...	<i>S. Leguminosarum</i> ...	<i>Frank.</i> , <i>Krauk. Pfl.</i> 652 (1881) ...	Leguminous schinzia ...

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Vergl. Morph. Pilze 187 (1884).—Ustilago, Uredo.									
2180	...	...	...	...	...	Q.	B.	Ovaries of <i>Polygonum</i>	Spore masses black, elongated, projecting from flower, opening to allow escape of spores.
2180A	...	...	...	...	...	...	...	Ovaries of <i>Polygonum</i>	Differs only in more distinct columella.
Ann. Sci. Nat. 285 (1883).									
2181	...	S.A.	...	V.	...	...	...	Leaves of <i>Lythrum hyssopifolium</i>	Pustules on both sides, globose, point-like, scattered or gregarious, brownish.
Linn. X. 230 (1835).									
2182	...	...	...	...	N.S.W.	Q.	...	Inflorescence of Rice grass (<i>Leersia hexandra</i>)	Pustules almost globose, nestling in pales. Spores pale brown.
2183	...	...	...	V.	...	...	...	Ovaries of <i>Leptocarpus tenax</i>	Spore balls composed of about ten globose spores, ultimately falling away into black powder.
Linn. IV. 116 (1829).									
2184	...	...	...	...	N.S.W.	Q.	...	Spikes of <i>Eriachne</i>	Mature fruit changed into black powdery mass. Spores brown.
2185	...	S.A.	...	V.	...	...	...	Panicles of <i>Gahnia filum</i>	Infesting inflorescence, but hardly visible to naked eye. Spores up to 100, in dark-brown balls.
Herb. Myc. II. 393 (1860).—Polycystis, Ustilago.									
2186	...	S.A.	...	V.	N.S.W.	...	B.	Wheat stems, leaves, glumes	Pustules forming long black streaks. Spores dark brown, one to three celled, surrounded by bladder-like sterile cells.
2187	...	...	T.	V.	N.S.W.	...	...	<i>Schanus imberbis</i> ...	Pustules black, globose, compact. Spore balls of three to eight.
Sci. Nat. III. 473 (1824).—Phacidium.									
2188	...	...	...	...	...	Q.	B.	Date palms ...	Conceptacles bursting through, opening above, outer layer black and horny. Spores yellow in mass.
Klotzsch, Herb. 1587 (1851).									
2189	...	...	...	...	...	Q.	...	<i>Heteropogon contortus</i>	Olive brown, at first covered with spores of same colour. Spores stuck together.
2190	...	...	...	...	...	Q.	...	Glumes of <i>Paspalum scrobiculatum</i>	Convex, hemispherical, twisted and folded, dark olive. Spores olive.
Linn. XVI. 278 (1842).									
2191	...	...	...	V.	...	...	B.	Roots of leguminous plants	Tubercles varying in size and form, coloured like root, containing hyphæ and innumerable minute corpuscles.

GENERAL CLASSIFICATION OF PHYCOMYCETES.

GROUP XI.—PHYCOMYCETES, DE BARY.

ARRANGEMENT OF ORDERS (5).

Hyphæ well developed—

56. MUCORACEÆ—Threads producing spore sacs.
57. PERONOSPORACEÆ—Threads often branched, bearing active or passive conidia.
58. ENTOMOPHTHORACEÆ—Threads bearing conidia mostly on insects.

Hyphæ obsolete—

59. CHYTRIDIACEÆ—Spore sacs alone, without threads.
60. PROTOMYCETACEÆ—Spore sacs thick walled, slender threads soon disappearing.
-

ORDER LVI.—MUCORACEÆ, DE BARY.

Genera (5)—

389. Pilobolus, Tode.		391. Phycomyces, Kunze.		393. Circinella, V. Tiegh.
390. Mucor, Mich.		392. Spinellus, V. Tiegh.		

ORDER LVII.—PERONOSPORACEÆ, DE BARY.

Genera (4)—

394. Cystopus, Lev.		396. Plasmopara, Schr.		397. Peronospora, Corda.
395. Sclerospora, Schr.				

ORDER LVIII.—ENTOMOPHTHORACEÆ, NOWAK.

Genus (1)—

398. Empusa, Cohn.

ORDER LIX.—CHYTRIDIACEÆ, DE BARY.

Genus (1)—

399. Synchytrium, De Bary.

ORDER LX.—PROTOMYCETACEÆ, DE BARY.

Genus (1)—

400. Protomyces, Unger.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
GROUP XI.—PHYCOMYCETES, DE BARY,					
ORDER LVI.—MUCORACEÆ,					
389. PILOBOLUS.—Tode,					
2192	1689	VII. 592	<i>P. crystallinus</i> ...	<i>Tode, Meek. 41 (1790) ..</i>	Crystalline pilobolus ...
390. MUCOR.—Linn.,					
2193	1691	IX. 1412	<i>M. cervinolenus</i> ...	<i>Berk., Fl. Tasm. II. 282 (1860) ...</i>	Fawn-white mucor ...
2194	1690	VII. 615	<i>M. mucedo</i> ...	<i>Linn., Sp. Pl. II. 1655 (1753) ...</i>	Mould mucor... ..
391. PHYCOMYCES.—Kunze,					
2195	1692	VII. 696	<i>P. nitens</i> ...	<i>Kunze, Myk. II. 113 (1823) ...</i>	Shining phycomyces ...
392. SPINELLUS.—Van Tiegh,					
2196	1693	IX. 1414	<i>S. gigasporus</i> ...	<i>Cooke and Mass., Grev. XVIII. 26 (1889) ...</i>	Large-spored spinellus... ..
393. CIRCINELLA.—Van Tiegh and Mon.,					
2197	1694	VII. 732	<i>C. umbellata</i> ...	<i>Van Tiegh and Mon., Ann. Sci. Nat. 300 (1853) ...</i>	Umbellate circinella ...
ORDER LVII.—PERONOSPORACEÆ,					
394. CYSTOPUS.—Lev., Ann. Sci.					
2198	1695	VII. 792	<i>C. candidus</i> ...	<i>Lev., Ann. Sci. Nat. 371 (1847) ...</i>	White cystopus ...
395. SCLEROSPORA.—Schr., in Cohn's					
2199	1696	IX. 1434	<i>S. macrospora</i> ...	<i>Sacc., Hedw. 155 (1890) ...</i>	Large-spored sclerospora ...
396. PLASMOPARA.—Schr., in Cohn's					
2200	...	VII. 806	<i>P. viticola</i> ...	<i>Berl. and De Ton., Sacc. Syll. VII. 239 (1888) ...</i>	Vine-growing plasmopara (Brown rot or downy mildew) ...
397. PERONOSPORA.—Corda,					
2201	1697	VII. 877	<i>P. hyoscyami</i> ...	<i>De Bary, Ann. Sci. Nat. 123 (1863) ...</i>	Henbane peronospora ...
2202	...	„ 857	<i>P. Schleideni</i> ...	<i>Unger, Bot. Ztg. 315 (1847) ...</i>	Schleiden's peronospora (Onion mildew) ...
ORDER LVIII.—ENTOMOPHTHORACEÆ,					
398. EMPUSA.—Cohn,					
2203	...	VII. 968	<i>E. Muscæ</i> ...	<i>Cohn, Nov. Act. Acad. XXV. 317 (1855)..</i>	House-fly empusa * ...
ORDER LIX.—CHYTRIDIACEÆ,					
399. SYNCHYTRIUM.—De Bary and Wor.,					
2204	1699	VII. 1002	<i>S. Succisæ</i> ...	<i>De Bary and Wor., Chytr. (1863) ...</i>	Succisa synchytrium ...
2205	1698	„ 999	<i>S. Taraxaci</i> ...	<i>De Bary and Wor., Chytr. (1863) ...</i>	Dandelion synchytrium ...
ORDER LX.—PROTOMYCETEEÆ,					
400. PROTOMYCES.—Unger,					
2206	1700	VII. 1120	<i>P. macrosporus</i> ...	<i>Unger, Exanth. 344 (1833) ...</i>	Large-spored protomyces ...

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
IN FCKL. SYMB. MYC. 66 (1875).									
DE BARY.									
Meek. 41 (1790).—Mucor.									
2192	...	...	...	...	...	Q.	B.	Dung ...	Gregarious, threads slender, pellucid, weeping, yellowish, club shaped at apex.
Sp. II. 1185 (1753).									
2193	...	...	T.	...	...	...	...	Dung ...	Threads simple, erect, white below, ochrey above.
2194	W.A.	S.A.	T.	V.	N.S.W.	Q.	B.	Putrid organic substances	Spore-bearing threads simple, erect, brownish. Spore sacs spherical, dark brown when dry.
Myk. II. 113 (1823).—Ulva, Mucor.									
2195	...	...	...	V.	N.S.W.	...	B.	Fatty substances ...	Spore-bearing threads bending, shining, brass colour, continuous. Spore sacs globose, turning black.
Ann. Sci. Nat. 66 (1875).—Mucor.									
2196	...	...	...	V.	...	...	...	Decaying Agarics...	Spore-bearing threads simple, bending, shining, olive. Spore sacs somewhat globose.
Ann. Sci. Nat. 300 (1873).—Mucor, Helicostylum.									
2197	...	...	...	...	...	Q.	...	Putrid substances ...	Spore-bearing threads erect, simple or branched, brown. Spore sacs spherical, becoming bluish.
DE BARY.									
Nat. 371 (1847).—Uredo, Æcidium.									
2198	...	S.A.	...	V.	N.S.W.	...	B.	Leaves, stems, &c., of <i>Crucifera</i>	Pustules bursting through, white, variable. Conidia uniform, globose, colourless. Very common.
Krypt. Fl. Schl. 236 (1888).									
2199	...	...	...	V.	...	...	...	Leaves of <i>Alopecurus</i> along with <i>Puccinia rubigo-vera</i>	Conidial stage unknown. Reproductive organs covered by epidermis, becoming brownish.
Krypt. Fl. Schl. 236 (1888).—Peronospora, Botrytis.									
2200	...	...	.	V.	...	...	...	Leaves of Grape vine	Threads thick, frequently constricted and swollen, with minute suckers.
Icon. I. 20 (1854).—Botrytis.									
2201	...	...	...	V.	N.S.W.	Q.	B.	Tobacco leaves ...	Conidia-bearing threads thick, tall, forking. Branches spreading, tapering.
2202	...	...	...	V.	N.S.W.	...	B.	Leaves of Onion, Garlic, &c.	Forming whitish-grey or greyish-lilac tufts, sometimes covering leaves. Conidia-bearing threads large, erect, forked.
NOWAK.									
Hedw. 57 (1885).—Sporendonema, Entomophthora.									
2203	...	...	...	V.	...	...	B.	Bodies of house flies and dipterous insects	White mould-like growth. Conidia bearers simple, crowded, club shaped.
DE BARY AND WORON.									
Ber. Nat. Ges. III. 22 (1863).									
2204	...	S.A.	...	V.	...	...	...	Leaves and leaf stalks of Goodeniaceous plants	Cells containing spore sacs orange red. Galls wart-like, solitary or forming brown crust.
2205	...	...	...	V.	...	...	B.	Leaves of <i>Compositae</i>	Spots crust-like, running together, orange red. Galls small, flattened, scarcely projecting.
DE BARY.									
Exanth. 341 (1833).—Physoderma.									
2206	...	...	...	...	...	Q.	B.	<i>Hydrocotyle asiatica</i> ...	Spores usually collected in scattered bulging spots, which are at first translucent, then brown.

GENERAL CLASSIFICATION OF MYXOMYCETES.

GROUP XII.—MYXOMYCETES, WALLR.

ARRANGEMENT OF ORDERS (9).

A.—Wall of spore sac not encrusted with lime—

Section 1. Peritrichæ—Capillitium absent or formed from wall of spore sac.

- 61. TUBULINACEÆ—Wall of spore sac not perforated.
- 62. CRIBRARIACEÆ—Wall of spore sac perforated.

Section 2. Columelliferæ—Capillitium originating from central columella.

- 63. STEMONITACEÆ—Arising from every part of elongated columella.
- 64. LAMPRODERMACEÆ—Arising from apical portion of columella.

Section 3. Calotricheæ—Capillitium not springing from columella.

- 65. ARCYRIACEÆ—Threads attached.
- 66. TRICHIACEÆ—Threads free.

B.—Wall of spore sac with external deposit of lime—

Section 4. Lithodermeæ—Capillitium present.

- 67. DIDYMIACEÆ—Threads without lime.
- 68. PHYSARACEÆ—Threads with lime.

C.—Without special spore sac.

- 69. PLASMIDIOPHORACEÆ—Plasmodia, or naked motile masses of protoplasm, formed.

ORDER LXI.—TUBULINACEÆ, MASS.

Genus (1)—

- 401. Tubulina, Pers.

ORDER LXII.—CRIBRARIACEÆ, MASS.

Genera (2)—

- 402. Enteridium, Ehrh.
- 403. Clathroptychium, Rost.

ORDER LXIII.—STEMONITACEÆ, ROST.

Genus (1)—

- 404. Stemonitis, Gled.

ORDER LXIV.—LAMPRODERMACEÆ, MASS.

Genus (1)—

- 405. Lamproderma, Rost.

ORDER LXV.—ARCYRIACEÆ, ROST.

Genera (4)—

- 406. Perichæna, Fries.
- 407. Lycopala, Pers.
- 408. Prototrichia, Rost.
- 409. Arcyria, Hill.

3868.

ORDER LXVI.—TRICHIACEÆ, ROST.

Genus (1)—

- 410. Trichia, Hall.

ORDER LXVII.—DIDYMIACEÆ, ROST.

Genera (4)—

- 411. Chondrioderma, Rost.
- 412. Didymium, Schrad.
- 413. Spumaria, Pers.
- 414. Diachæa, Fries.

ORDER LXVIII.—PHYSARACEÆ, ROST.

Genera (6)—

- 415. Craterium, Trent.
- 416. Physarum, Pers.
- 417. Badhamia, Berk.
- 418. Tilmadoche, Fries.
- 419. Leocarpus, Link.
- 420. Fuligo, Hall.

ORDER LXIX.—PLASMIDIOPHORACEÆ, ZOPF.

Genus (1)—

- 421. Plasmodiophora, Zopf.

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
GROUP XII.—MYXOMYCETES,					
ORDER LXI.—TUBULINACEÆ,					
401. TUBULINA.—Pers., Syn. 197 (1808).—					
2207	2032	VII. 1391	<i>T. cylindrica</i>	<i>D. C.</i> , Fl. Fr. II. 249 (1815)	Cylindrical tubulina
2207A	„	„ 1394	<i>T. cylindrica</i> , var. <i>nitidissima</i>	Cooke, Handb. Austr. Fung. 392 (1892)	Shining tubulina
2208	2033	X. 4818	<i>T. spumarioidea</i>	Cooke and Mass., Mon. Myx. 42 (1892)	Spumaria-like tubulina
ORDER LXII.—CRIBRARIACEÆ,					
402. ENTERIDIUM.—Ehrh. in Link Jahrb.,					
2209	2034	VII. 1399	<i>E. olivaceum</i>	Ehrh., Sylv. Ber. II. 54 (1818)	Olive enteridium
403. CLATHROPTYCHUM.—Rost.,					
2210	2035	VII. 1396	<i>C. rugulosum</i>	<i>Rost.</i> , Mon. 225 (1875)	Wrinkled clathroptychium
ORDER LXIII.—STEMONITACEÆ,					
404. STEMONITIS.—Gled., Meth. 140 (1753).—					
2211	2038	VII. 1365	<i>S. ferruginea</i>	Ehr., Sylv. Berl. 20 (1818)	Ferruginous stemonitis
2212	2037	„ 1356	<i>S. Friesiana</i>	De Bary, Rabh., Fung. Eur. 568 (1861-81)	Fries' stemonitis
2213	2036	„ 1362	<i>S. fusca</i>	Roth., Fl. Germ. I. 548 (1802)	Brown stemonitis
ORDER LXIV.—LAMPRODERMACEÆ,					
405. LAMPRODERMA.—Rost., Vers.					
2214	2039	VII. 1344	<i>L. echinulatum</i>	<i>Rost.</i> , Mon. App. 25 (1875)	Echinulate lamproderma
2215	2040	„	<i>L. Listeri</i>	Mass., Mon. Myx. 97 (1892)	Lister's lamproderma
ORDER LXV.—ARCYRIACEÆ,					
406. PERICHÆNA.—Fries, Symb.					
2216	2042	VII. 1515	<i>P. applanata</i>	Cooke and Mass., Mon. Myx. 116 (1892)	Flattened perichæna
2217	2041	„ 1435	<i>P. corticalis</i>	<i>Rost.</i> , Mon. 293 (1875)	Cortical perichæna
407. LYCOGALA.—Pers., Tent. Disp. 7 (1797).—					
2218	2043	VII. 1484	<i>L. epidendrum</i>	<i>Rost.</i> , Mon. 285 (1875)	Tree-growing lycogala
408. PROTOTRICHIA.—Rost.,					
2219	2044	VII. 1492	<i>P. metallica</i>	<i>Mass.</i> , Mon. Myx. 127 (1892)	Metallic prototrichia

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.	
	W.A.	S.A.	T.	V.	N.S.W.	Q.				
WALLR., FL. CRYPT. II. 333 (1833).										
MASS. MON. MYX. (1892).										
Tubulifera, Mucor, Licea, &c.										
2207	...	...	T.	...	...	Q.	B.	Rotten wood ...	Spore sacs cylindrical, rounded at apex, gregarious, mostly crowded. Mass of spores chestnut.	
2207A	...	...	...	...	...	Q.	...	<i>Eucalyptus microtheca</i>	Spore sacs shining, golden yellow.	
2208	...	...	...	V.	...	...	...	Running over twigs and on ground	Æthelium irregular, ashy. Cortex membranous, netted with branched veins.	
MASS. MON. MYX. (1892).										
Grev. II. 51 (1873).—Lycoperdon, Licea.										
2209	W.A.	...	...	...	...	...	B.	Wood ...	Æthelium very variable in form, flattened or cushion shaped, olive.	
Mon. 225 (1875).—Fuligo, Licea.										
2210	W.A.	...	...	...	...	Q.	B.	Dead twigs, &c. ...	Colour of æthelium variable, red brown or ochrey. Spore sacs bell shaped at apex.	
ROST. MON. (1875).										
Clathrus, Comatricha, Trichia.										
2211	...	...	...	...	...	Q.	B.	Rotten wood ...	Spore sacs cylindrical, gregarious, on violet-black expansion. Spores rusty cinnamon.	
2212	...	...	T.	...	...	Q.	B.	Rotten wood ...	Spore sacs globose, ovate, erect. Stem black, shining.	
2213	W.A.	...	T.	V.	...	Q.	B.	Rotten wood, &c. ...	Spore sacs cylindrical, obtuse, on strongly-developed expansion, which is violet black.	
MASS. MON. MYX. (1892).										
Syst. Myc. 7 (1873).—Stemonitis.										
2214	...	...	T.	...	...	...	...	Among moss ...	Spore sacs stalked, dark steel blue or blackish, iridescent. Stem short. Spores spiny.	
2215	...	...	T.	...	...	...	B.	Moss, wood, &c. ...	Spore sacs stalked, globose, blackish purple, iridescent. Stem elongated, blackish brown.	
ROST. MON. (1875).										
Gast. 9 (1818).—Hemiareyria, Lycoperdon.										
2216	...	...	...	...	...	Q.	...	Rotting <i>Cycas</i> ...	Sessile, much depressed, circular or irregular in outline. Spores in mass clear orange yellow.	
2217	W.A.	...	...	...	...	...	B.	Bark and wood ...	Spore sacs spherical to depressed, crowded. Spores in mass pale yellow.	
Fungus, Lycoperdon, Bovista.										
2218	W.A.	...	...	V.	...	Q.	B.	Stumps ...	Æthelium gregarious, rounded, size of pea, shining, distinctly warted, rose colour.	
Mon. Appl. 38 (1875).										
2219	...	...	T.	...	...	...	...	Wood ...	Spore sacs scattered, stalked or sessile, copper colour with metallic tints. Stem very short.	

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
409. ARCYRIA.—Hall, Hist. 47 (1768).—Clathrus,					
2220	2049	VII. 1459	<i>A. cinerea</i> ...	Schum., Saell. 1480 (1801) ...	Grey arcyria ...
2221	2046	„ 1470	<i>A. ferruginea</i> ...	Rost., Mon. 280 (1875) ...	Ferruginous arcyria ...
2222	2052	X. 4857	<i>A. fuliginea</i> ...	Mass., Mon. Myx. 169 (1892) ...	Sooty-brown arcyria ...
2223	2047	VII. 1461	<i>A. incarnata</i> ...	Pers., Obs. (1796) ...	Fleshy arcyria ...
2224	2048	„ 1464	<i>A. nutans</i> ...	Grev., Fl. Ed. 435 (1824) ...	Drooping arcyria ...
2225	2045	„ 1457	<i>A. punicea</i> ...	Pers., Disp. 10 (1797) ...	Reddish arcyria ...
2226	2050	„ 1512	<i>A. rubiformis</i> ...	Mass., Mon. Myx. 158 (1892) ...	Lustrous arcyria ...
2227	2051	„ 1514	<i>A. serpula</i> ...	Mass., Mon. Myx. 164 (1892) ...	Creeping arcyria ...

ORDER LXVI.—TRICHIACEÆ,

410. TRICHIA.—Hall, Helv. III. 114 (1768).—

2228	2057	VII. 1499	<i>T. affinis</i> ...	De Bary, in Rost. Mon. 257 (1875) ...	Allied trichia ...
2229	2055	„ 1503	<i>T. contorta</i> ...	Rost., Mon. 259 (1875) ...	Contorted trichia ...
2230	2053	„ 1494	<i>T. fragilis</i> ...	Rost., Mon. 246 (1875) ...	Fragile trichia ...
2231	...	X. 4848	<i>T. Kalbreyeri</i> ...	Mass., Mon. Myx. 191 (1892) ...	Kalbreyer's trichia ...
2232	2054	VII. 1497	<i>T. varia</i> ...	Pers., Tent. Disp. 10 (1797) ...	Variable trichia ...
2233	2056	X. 4847	<i>T. verrucosa</i> ...	Berk., Fl. Tasm. II. 269 (1860) ...	Warted trichia ...

ORDER LXVII.—DIDYMIACEÆ,

411. CHONDRIODERMA.—Rost.,

2234	2058	VII. 1282	<i>C. difforme</i> ...	Rost., Mon. 177 (1875) ...	Deformed chondrioderma ...
2235	...	„ 1257	<i>C. Muelleri</i> ...	Rost., Mon. 15 (1875) ...	Mueller's chondrioderma ...

412. DIDYMIUM.—Schröd., Nov. Pl. Gen. I. 22 (1797).—

2236	2064	X. 4803	<i>D. australe</i> ...	Mass., Grev. XVII. 7 (1888) ...	Southern didymium ...
2237	2059	VII. 1309	<i>D. farinaceum</i> ...	Schröd., Nov. Pl. Gen. I. (1797) ...	Mealy didymium ...
2238	2065	„ 1193	<i>D. flavicomum</i> ...	Mass., Mon. Myx. 242 (1892) ...	Yellow-haired didymium ...
2239	2063	„ 1256	<i>D. pezizoideum</i> ...	Mass., Mon. Myx. 239 (1892) ...	Peziza-like didymium ...
2240	2062	„ 1297	<i>D. serpula</i> ...	Fries, S.M. III. 126 (1832) ...	Creeping didymium ...
2241	2061	„ 1269	<i>D. spumarioides</i> ...	Fries, Symb. Gast. 20 (1818) ...	Spumaria-like didymium ...
2242	2060	„ 1301	<i>D. squamulosum</i> ...	Fries, S.M. III. 118 (1832) ...	Scaly didymium ...

413. SPUMARIA.—Pers.,

2243	2066	VII. 1338	<i>S. alba</i> ...	D.C., Fl. Fr. II. 261 (1815) ...	White spumaria ...
------	------	-----------	--------------------	----------------------------------	--------------------

OF AUSTRALIAN FUNGI—continued.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Trichia, Stemonitis, Hemiareyria, Clathroides, Mucor.									
2220	...	...	...	...	...	Q.	B.	Stumps ...	Gregarious, stalked, ovoid or elongated ovoid. Stem erect, straight, long. Spores usually bright grey.
2221	...	...	...	...	...	Q.	B.	Rotten wood, &c. ...	Spore sacs ovate. Stem usually short. Spores in mass usually brick red, now and then rusty.
2222	...	...	...	...	N.S.W.	...	...	Leaves of <i>Atherospermum</i> ...	Threads forming a net-work, spiny. Spores globose, smooth, in mass sooty brown.
2223	W.A.	...	...	...	...	Q.	B.	Rotten wood ...	Spore sacs egg shaped, with evanescent short erect stem. Spores in mass usually flesh colour.
2224	W.A.	...	...	V.	...	Q.	B.	Rotten wood ...	Spore sacs cylindrical, with short evanescent stem. Capillitium drooping.
2225	...	...	...	...	...	Q.	B.	Rotten stumps ...	Spore sacs more or less egg shaped, of beautiful lustre, usually with erect stem.
2226	...	...	T.	...	...	...	B.	Dead wood ...	Spore sacs usually tufted, collected in short stem, often of beautiful metallic lustre.
2227	W.A.	...	...	...	...	...	B.	Rotten wood ...	Vein-like, creeping, forming a net, or somewhat globose, and sessile on broad base, yellow.

ROST. MON. (1875).**Lycogala, Licea, Lycoperdon.**

2228	...	...	T.	...	...	Q.	B.	Rotten wood, &c. ...	Spore sacs clustered, circular or elliptical, sessile on broad base, clear yellow.
2229	...	...	T.	...	...	Q.	B.	Rotten wood ...	Variable in form, sometimes elongated and twisted, sometimes veined, creeping, bay brown.
2230	W.A.	...	T.	...	...	...	B.	Rotten wood, &c. ...	Spore sacs stalked, or tufted on common stem; wall smooth, blackish or yellowish.
2231	...	...	T.	...	...	...	...	Fragments of rotting plants ...	Spore sacs crowded, sessile, globose, yellow, nearly the same colour within.
2232	...	...	T.	...	...	Q.	B.	Stumps ...	Spore sacs variously developed, either stalked or sessile.
2233	...	...	T.	...	...	...	...	Wood ...	Spore sacs brown or chestnut, shining, passing down into long slender stem.

ROST. MON. (1875.)**Mon. 167 (1875).—Physarum.**

2234	...	S.A.	...	V.	...	Q.	B.	Bark, leaves, twigs, grass, &c. ...	Spore sacs sessile, roundish, deformed; outer wall crustaceous, chalky white.
2235	...	...	...	...	...	Q.	...		Spore sacs discoid, curved upwards, snow white, stalked. Stem straight, rigid, with rusty-brown furrows.

Mucor, Trichia, Physarum, Chondrioderma, Spumaria.

2236	...	...	...	...	...	Q.	...	Old Auricularia ...	Spore sacs globose or slightly compressed, covered with dense white layer of lime.
2237	...	...	...	V.	...	Q.	B.	Dead leaves, twigs, decaying fruit, &c. ...	Spore sacs hemispherical or a little flattened, greyish white with lime, or black and shining.
2238	W.A.	...	...	...	...	...	...	Rotten wood ...	Fructification hemispherical, violet or lilac. Stem elongated, slender, yellowish tan or copper colour.
2239	...	...	...	...	...	Q.	...	Dead wood of <i>Erythrina vespertilio</i> ...	Fructification somewhat nodding, ashy white, arising from mealy crusty cracking membrane.
2240	...	...	...	...	...	Q.	B.	Fallen leaves and rotten wood ...	Fructification either cushion-like, flattened, or vein-like, creeping.
2241	...	...	T.	...	...	Q.	B.	Leaves, moss, clover, &c. ...	Spore sacs irregular in shape, snow white or greyish, always in clusters.
2242	...	...	T.	...	...	...	B.	Wood, dead leaves, &c. ...	Spore sacs either hemispherical and flattened or globose. Stem snow white.

Syn. 162 (1808).—Reticularia.

2243	...	...	...	V.	N.S.W.	Q.	B.	Grass ...	Æthelium complex, branching, whitish to grey, spongy.
------	-----	-----	-----	----	--------	----	----	-----------	---

SYSTEMATIC ARRANGEMENT

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
414. DIACHÆA.—Fries, Syst. Orb.					
2244	2067	VII. 1335	<i>D. leucopoda</i>	Rost., Mon. 191 (1875)...	White-stalked diachæa ...
ORDER LXVIII.—PHYSARACEÆ,					
415. CRATERIUM.—Trent.,					
2245	2068	VII. 1233	<i>C. confusum</i>	Mass., Mon. Myx. 263 (1892)	Confused craterium
416. PHYSARUM.—Pers., Obs. Myc. 5 (1799).—					
2246	2073	VII. 1189	<i>P. cinereum</i>	Pers., Syn. 170 (1808)...	Ashy physarum
2247	...	„ 1171	<i>P. didermoides</i>	Rost., Mon. 97 (1875)	Two-membraued physarum ...
2248	2072	„ 1192	<i>P. leucophænum</i>	Fries, Symb. Gast. 24 (1818)	Grey physarum
2249	2071	„ 1188	<i>P. leucopus</i>	Rost., Mon. 101 (1875)...	White-stalked physarum ...
2250	2070	...	<i>P. Readeri</i>	Mass., Mon. Myx. 282 (1892)	Reader's physarum
2251	2069	VII. 1251	<i>P. rufibasis</i>	Berk. and Br., Linn. Journ. XIV. 85 (1875)	Red-based physarum
2252	2074	„ 1189	<i>P. scrobiculatum</i>	Mass., Mon. Myx. 300 (1892)	Pitted physarum
417. BADHAMIA.—Berk.,					
2253	2075	VII. 1150	<i>B. varia</i>	Mass., Mon. Myx. 319 (1892)	Variable badhamia
418. TILMADOCHÆ.—Fries,					
2254	2077	VII. 1247	<i>T. mutabilis</i>	Rost., Mon. 130 (1875)...	Changeable tilmadochæ ...
2255	2076	„ 1244	<i>T. nutans</i>	Rost., Mon. 127 (1875)	Nodding tilmadochæ
419. LEOCARPUS.—Link,					
2256	2078	VII. 1242	<i>L. fragilis</i>	Rost., Mon. 132 (1875)...	Fragile leocarpus
420. FULIGO.—Hall, Hist.					
2257	2079	VII. 1228	<i>F. varians</i>	Sommf., Fl. Lapp. 231 (1826)	Variable fuligo
ORDER LXIX.—PLASMIDIOPHORACEÆ,					
421. PLASMIDIOPHORA.—Woron.					
2258	...	VII. 1568	<i>P. Brassicæ</i>	Woron. Pringsh. Jahrb. XI. 548 (1878)	Turnip Plasmodiophora (Club-Root) (Fingers and Toes)

OF AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
Veg. I. 143 (1825).— <i>Trichia</i> , <i>Stemonitis</i> .									
2244	...	...	...	V.	...	...	B.	Leaves, &c. ...	Spore sacs cylindrical, stalked. Stem short, thickened at base, snow white.
ROST. MON. (1875).									
in Roth. Cat. II. 224 (1806).									
2245	W.A.	...	T.	...	...	...	B.	Leaves, &c. ...	Spore sacs variable in form, stalked or rarely somewhat sessile, bright, brown, ochrey, or white.
Lycoperdon, <i>Trichia</i> , <i>Didymium</i> , <i>Tilmadocho</i> .									
2246	W.A.	...	...	V.	...	...	B.	Bark, wood, leaves, &c.	Spore sacs globose or hemispherical, sessile or gregarious.
2247	...	...	...	...	...	...	Q.	Scales of Onions, and bracts of Maize, Grass, &c.	Spore sacs ovoid, ash coloured, with white mealy covering, and separate membranous outer coat.
2248	..	...	...	...	...	...	B.	Leaves, &c. ...	Spore sacs somewhat globose, stalked or sessile, wall thin with white lime patches.
2249	...	...	...	...	...	...	B.	Wood, &c. ...	Spore sacs globose, stalked or sessile, wall covered with snow-white coat of lime. Stem white.
2250	...	...	...	V.	...	...	...	Wood ...	Spore sacs stalked, greyish, covered with flakes of lime. Stem very thick, brown.
2251	...	...	...	...	...	...	Q.	Moss ...	Scattered or gregarious, stalked. Stem elongated, slender, expanding into circular bright-brown base. Spore sacs globose, dull yellow or tawny.
2252	W.A.	...	...	...	...	...	B.	Charred wood ...	Spore sacs sessile, on broad or narrowed base, seated on thick spreading expansion.
Outl. 308 (1860).— <i>Physarum</i> .									
2253	...	...	T.	V.	N.S.W.	...	B.	Rotten wood, &c. ...	Spore sacs more or less clustered, sessile or stalked, globose, grey or opaque.
S.V.S. 454 (1849).— <i>Stemonitis</i> .									
2254	W.A.	...	...	...	...	...	Q.	Decayed wood ...	Spore sacs globose or flattened, usually cracked, yellow or rusty orange, stalked, nodding.
2255	W.A.	...	T.	...	...	...	Q.	Rotten wood, &c. ...	Spore sacs lens shaped, usually cracked, greyish white, stalked, nodding.
Sp. Pl. I. 25 (1824).— <i>Lycoperdon</i> .									
2256	...	...	T.	...	...	...	B.	Grass, twigs, moss, &c.	Spore sacs somewhat roundish, sessile, or with thin thread-like coloured stem.
Helv. III. 110 (1768).— <i>Mucor</i> .									
2257	W.A.	S.A.	T.	V.	...	Q.	B.	Wood, tan, soil, &c.	Spore sacs more or less closely interwoven, bark not always developed; walls of spore sac usually coloured.
ZOPF. PILZTH. 129 (1885).									
Pringsh. Jahrb. XI., 548 (1878).									
2258	...	...	...	V.	N.S.W.	...	B.	Roots of Crucifers— <i>Brassica</i> , &c.	Producing the malformation of the roots of cabbages, cauliflowers, &c., which gives them a clubbed appearance, or several misshapen roots like "fingers and toes."

ADDITIONS.—A.—NEW

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
<p style="text-align: right;">GROUP I.— ORDER II.— Genus 422.—<i>Laccocephalum</i>, McAlp., Stone-like-base laccocephalum ...</p>					
2259	...	...	<i>L. basilapiloides</i> ...	McAlp. and Tepp., Proc. Roy. Soc. Vic. VII. N.S. 166 Pl. X. (1894)	Stone-like-base laccocephalum ...
<p style="text-align: right;">GROUP III.— ORDER XIII.—</p>					
2260	...	VII. 1926	<i>Uromyces Phaseoli</i> ...	Winter, Die Pilze 157 (1834) ...	Bean uromyces (Bean rust) ...
2261	...	...	<i>Puccinia Correæ</i> ...	McAlp., Proc. Roy. Soc. Vic. VII. N.S. 215 (1894)	Correa puccinia ...
2262	...	...	<i>P. Erechthitis</i> ...	McAlp., Proc. Roy. Soc. Vic. VII. N.S. 216 (1894)	Erechthites puccinia ...
2263	...	...	<i>P. Hypochæris</i> ...	McAlp., Proc. Roy. Soc. Vic. VII. N.S. 217 (1894)	Hypochæris puccinia ...
2264	...	...	<i>P. Plagianthi</i> ...	McAlp., Proc. Roy. Soc. Vic. VII. N.S. 218 (1894)	Plagianthus puccinia ...
2265	...	...	<i>Æcidium eburneum</i> ...	McAlp., Proc. Roy. Soc. Vic. VII. N.S. 218 (1894)	Ivory æcidium ...
2266	...	IX. 1318	<i>A. monocystis</i> ...	Berk., Flor. N.Z. II. 196 (1855)...	Walled æcidium ...
2267	...	...	<i>Puccinia Coprosomatis</i> ...	Morrison, Vict. Nat. XI. No. 6, 90 (1894)	Coprosma puccinia ...
2267A	...	...	<i>P. Coprosomatis</i> , var. <i>Operculariæ</i>	Morrison, Vict. Nat. XI. No. 8, 119 (1894)	Opercularia puccinia ...
2268	...	VII. 2457	<i>P. investita</i> ...	Schwein, Syn. N. Am. Fungi (1831) ...	Invested puccinia ...
<p style="text-align: right;">GROUP IV.— ORDER XV.—</p>					
2269	...	I. 1166	<i>Xylaria fulvella</i> ...	Berk. and Curt., Linn. Journ. X. 380 (1869)	Tawny xylaria ...
2270	...	I. 1282	<i>X. ianthino-velutina</i> ...	Mont., Syll. Crypt. (1856) ...	Violet-haired xylaria ...
2271	...	IX. 2282	<i>Kretzschmaria confusa</i> ...	Sacc., Syll. IX. 566 (1891) ..	Confused kretzschmaria ...
2272	...	...	<i>Hypoxylon atosphaericum</i> ...	Cooke and Mass., Grev. XXII. 68 (1894)	Black-sphered hypoxylon ...
<p style="text-align: right;">ORDER XVI.—</p>					
2273	...	II. 5107	<i>Phyllachora Grevilleæ</i> ..	Sacc., Syll. II. 597 (1883) ...	Grevillea phyllachora ...
<p style="text-align: right;">GROUP V.— ORDER XXXIV.—</p>					
2274	...	...	<i>Peziza Lyonsiæ</i> ...	Cobb., Ag. Gaz. N.S.W. V. 6, 390 (1894)	Lyonsia peziza ...
2275	...	...	<i>Belonidium parasiticum</i> ..	Cooke and Mass., Grev. XXII. 68 (1894)	Parasitic belonidium ...

AUSTRALIAN FUNGI.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
HYMENOMYCETES, FRIES.									
POLYPORACEÆ, FRIES.									
Proc. Roy. Soc. Vic. VII. N.S. 166 (1894).									
2259	...	S.A.	...	...	...	...	...	Sandy soil in Mallee scrub	Solitary. Cap woody, brownish fawn, surface pitted. Stem compressed oval, dirty fawn, hardened like cap. Pores moderately large, crowded, nearly oval.
UREDINES, BRONGN.									
UREDINACEÆ, BRONGN.									
2260	...	...	...	...	N.S.W.	...	B.	All parts of Bean plant (<i>Phaseolus vulgaris</i>), more especially on leaves	Pustules brown, scattered, bursting through, surrounded by ruptured cuticle.
2261	...	...	T.	...	...	...	...	Under surface of leaves of <i>Correa Lawrenciana</i>	Pustules cushion shaped, circular or interruptedly so, dirty brown, scattered, soon naked.
2262	...	...	...	V.	...	...	...	Stem and leaves of <i>Erechtites quadridentata</i>	Cluster cups pale yellow to orange yellow, causing distortion and swelling. Pustules black, crowded together, and forming swelling.
2263	...	...	...	V.	...	...	...	Leaves of <i>Hypochaeris radicata</i>	Cluster cups on greenish-yellow or brownish circular patches. Pustules intermixed with cluster cups, black, and on both surfaces of leaf.
2264	...	...	T.	...	...	...	...	Leaves and flowers of <i>Plagianthus sidioides</i>	Pustules reddish brown, naked, blistered, scattered. Very common.
2265	...	...	T.	V.	...	...	...	Stem, leaves, flowers, and legumes of <i>Bossia cinerea</i>	Cluster cups ivory colour to brownish, clustered together without definite order.
2266	...	...	T.	...	...	...	...	Leaves of <i>Abrotanella forsterioides</i>	Cluster cups near tips of leaves, large, solitary, surrounded by strong wall arising from matrix.
2267	...	...	...	V.	...	...	...	Leaves of <i>Coprosma Billardieri</i>	Pustules on under surface, seldom on upper, prominent, deep brown, coalescing. Teleutospores compact, brown.
2267A	...	...	...	V.	...	...	...	Leaves and petioles of young plants of <i>Opercularia varia</i>	Pustules on under surface, deforming leaf and forming concavity on opposite side, reddish-brown. Teleutospores pale yellowish brown.
2268	...	...	...	V.	...	...	...	Leaves and stems of <i>Gnaphalium purpureum</i>	Pustules on both surfaces of leaf, bursting through epidermis, and bordered by ruptured cuticle. Uredospores and teleutospores generally mixed.
PYRENOMYCETES, FRIES.									
XYLARIACEÆ, COOKE.									
2269	...	...	...	...	...	Q.	...	At base of dead stump	Club shaped, rust coloured, papillate. Stem cylindrical, pale tawny. Receptacles with black openings.
2270	...	...	...	...	...	Q.	...	Old fruit of a <i>Flindersia</i>	Simple or branched, cylindrical and tapering, apex compressed, long violet-brown hair all over.
2271	...	...	...	...	...	Q.	...	Bark of dead log	Gregarious, stalked, simple. Heads depressed, globose, glaucous, at length black.
2272	...	...	...	...	...	Q.	...	Bark	Gregarious, sub-globose, black. Receptacles around the circumference ovate, teat-like.
DOTHIDEACEÆ, NITS.									
2273	W.A.	...	...	...	...	...	...	Leaves of <i>Grevillea</i>	On both surfaces, scattered, circular, shining black. Receptacles globose, immersed.
DISCOMYCETES, FRIES.									
PEZIZACEÆ, FRIES.									
2274	...	...	...	...	N.S.W.	...	...	Leaves of <i>Lyonsia reticulata</i>	Cups somewhat gregarious on both sides, on ashy-grey roundish spots, flat, sessile, round. Leaves appear at a little distance as if attacked by scale insect.
2275	...	...	...	...	...	Q.	...	Parasitic on <i>Asterina</i> , growing on leaflets of <i>Tarrietia trifoliolata</i>	White Cups very minute, hairless, concave, attached by central papilla, hardly visible to naked eye.

ADDITIONS.—(A.)—NEW

Number.	Cooke's Number.	Saccardo's Number.	Scientific Name.	Authority for Name.	English Name.
GROUP VII.— ORDER XLV.—					
2276	...	...	Oidium Oxalidis ...	McAlp., Proc. Roy. Soc. Vic. VII. N.S. 219 (1894)	Wood-sorrel oidium ...
ORDER XLVI.— Genus 423.—Stachybotrys,					
2277	...	IV. 1304	Stachybotrys lobulata ...	Berk., Ontl. 343 (1860) ...	Lobed stachybotrys ...
2278	...	„ 1675	Cladosporium carpophilum ...	Thuem., Fung. Pom. 13 (1878) ...	Fruit-loving cladosporium (Peach freckle)
ORDER XLVII.—					
2279	...	...	Isaria Oncopterae ...	McAlp., Proc. Roy. Soc. Vic. VII. N.S. 159 (1894)	Oncoptera isaria ...
GROUP VIII.— ORDER XLIX.—					
2280	...	III. 2796	Septoria Dianthi ..	Desm., 17 Not. 6, p. 20... ..	Carnation septoria ...
GROUP IX.—					
2281	...	VIII. 3622	Saccharomyces conglomeratus	Reess, Bot. Unt. 82 (1870) ..	Conglomerate yeast ...
2282	...	„ 3623	S. exiguus ...	Reess, Bot. Unt. 82 (1870) ...	Small yeast ...
2283	...	„ 3629	S. Marxianus ...	Hansen in Ann. de Microg. (1888) ...	Marx yeast ...
2284	...	„ 3630	S. membranifaciens ...	Hansen Bot. Zeil. 772 (1888) ...	Membrane-forming yeast ...
2285	...	„ 3635	S. minor ...	Engel, Ferm. (1872) ...	Lesser yeast ...
2286	...	„ 3624	S. Pasteurianus ...	Reess, Bot. Unt. 83 (1870) ...	Pasteur's yeast ...
GROUP X.— ORDER LV.—					
2287	...	...	Ustilago Allii ...	McAlp., Proc. Roy. Soc. Vic. VII. N.S. 220 (1894)	Onion ustilago ...
2288	...	...	U. Poarum ...	McAlp., Proc. Roy. Soc. Vic. VII. N.S. 220 (1894)	Poa ustilago ...
Genus 424.—Tolyposporium, Woron.					
2289	...	...	Tolyposporium Anthistiriae	Cobb, Ag. Gaz. N.S.W. III., pt. 12 (1892) ...	Kaogaroo-grass tolyposporium ...
GROUP XI.— ORDER LVII.—					
2290A	...	...	Peronospora parasitica, var. Lepidii	McAlp., Proc. Roy. Soc. Vic. VII. N.S. 221 (1894)	Lepidium peronospora ...

AUSTRALIAN FUNGI—*continued*.

Number.	Habitat.						B.	Occurrence.	General Characters.
	W.A.	S.A.	T.	V.	N.S.W.	Q.			
HYPHOMYCETES, MARTIUS.									
MUCEDINACEÆ, LINK.									
2276	...	...	...	V.	...	...	...	Leaves, leaf stalks, stem, and fruit of <i>Oxalis corniculata</i>	Mostly on upper surface of leaves, sometimes on lower, spread out, greyish, powdery.
DEMATIACEÆ, FRIES.									
Corda, Anleit. 57 (1842).									
2277	...	...	...	...	...	Q.	B.	Damp wall-paper ...	Black, sterile hyphæ creeping, fertile branches ascending, simple or branched, pale upwards.
2278	...	...	...	...	N.S.W.	...	...	Peaches ...	Spots circular, small at first, often confluent, dark green, finally causing cracking of peach.
STILBEACEÆ, FRIES.									
2279	...	...	...	V.	...	...	...	Dead larvæ of <i>Oncop-tera intricata</i> , Walk.	Dirty-brown root colour. Stem branched, velvety, slender, tips of branches fertile.
SPHÆROPSIDES, LEV.									
SPHÆRIOIDACEÆ, SACC.									
2280	...	...	...	V.	...	...	B.	Leaves of Carnations	Spots yellowish, oblong, round, or irregular. Receptacles globose to depressed, black to brown.
SACCHAROMYCETES, REESS.									
2281	...	...	...	...	...	...	B.	In fermentation and putrefaction of wine	Cells spheroidal, and forming a conglomeration instead of chains or flakes.
2282	...	...	...	...	...	...	B.	In juices of fermented fruits and fermentation of beer and wine	Cells spherical or top-shaped, united into a few branched colonies, and very small.
2283	...	...	...	...	...	...	...	Grape berry ...	Cells somewhat resembling <i>C. ellipsoideus</i> .
2284	...	...	...	...	...	...	...	Saccharine liquids...	The asci are many-spored.
2285	...	...	...	...	...	...	B.	In fermentation of bread	Cells spherical, isolated, double or sometimes in threes; like ordinary yeast but smaller.
2286	...	...	...	...	...	...	B.	In fermentation of wine and self-fermenting beer	Cells oval, oblong to club-shaped, varying in size; in branched colonies or flakes.
USTILAGINES, TUL.									
USTILAGINACEÆ, TUL.									
2287	...	...	...	V.	...	...	...	Scale leaves of stored Onion bulbs	Pustules minute, dark coloured, in parallel lines along veins of leaves, at first covered by epidermis; then powdery black.
2288	...	...	...	V.	...	...	...	<i>Poa annua</i> , especially foliage	Distorting, discolouring, and stunting plants, and forming black powdery masses.
in Schroet. Pilzfl. Schles. 276 (1882).									
2289	...	...	...	...	N.S.W.	...	...	Inflorescence of Kangaroo-grass (<i>Anthistiria ciliata</i>)	Fructification black. Spores compound, consisting of from a few dozen to several hundred thick-walled brown spores.
PHYCOMYCETES, DE BARY.									
PERONOSPORACEÆ, DE BARY.									
2290A	...	...	...	V.	...	...	...	Leaves, stem, and fruit of <i>Lepidium rudemale</i>	Lower surface of leaf attacked first, causing it to curl. Dense white mould forming a felt.

B.—NEW LOCALITIES, HOSTS, ETC.

GROUP I.—HYMENOMYCETES, FRIES.

ORDER IV.—THELEPHORACEÆ, PERS.

Genus 425.—*Soppittiella*, Mass. Brit. Fung. Fl. 106 (1892).

1045. *THELEPHORA CRISTATA* = *Soppittiella cristata*, Mass. B.

1080. *STEREUM OCHROLEUCUM*—V.

GROUP II.—GASTROMYCETES, WILLD.

ORDER X.—LYCOPERDACEÆ, EHRLH.

1343. *BATTARREA PHALLOIDES*—Q.

GROUP III.—UREDINES, BRONGN.

ORDER XIII.—UREDINACEÆ, BRONGN.

1446. *UROMYCES BETÆ*—N.S.W.

1455. *UROMYCES ORCHIDEARUM*—V. Uredospores intermixed with teleutospores.

1461. *MELAMPSORA LINI*—T.

1477. *PUCCINIA HETEROSPORA*—On leaves of a native *Hibiscus* in Queensland.

1478. *PUCCINIA HIERACHII*—V. On leaves of flowering stems of *Hypochaeris radicata*.

1511. *ÆCIDIUM RANUNCULACEARUM*—On *Ranunculus parviflorus* in Victoria.

GROUP IV.—PYRENOAMYCETES, FRIES.

ORDER XIV.—HYPOCREACEÆ, DE NOT.

1527. *CLAVICEPS PURPUREA*—T., N.S.W.

ORDER XV.—XYLARIACEÆ, COOKE.

1619. *HYPOXYLON ELLIPTICUM*—V. Rotten wood.

ORDER XXVIII.—FOLIICOLACEÆ, FRIES.

1712. *SPHÆRELLA FRAGRARIÆ*—W.A., T.

ORDER XXX.—PERISPORIACEÆ, FRIES.

1756. *CAPNODIUM CITRI*—W.A., Q. Leaves of oranges and lemons.

GROUP V.—DISCOMYCETES, FRIES.

ORDER XXXIII.—HELVELLACEÆ, LINK.

1778. *MORCHELLA DELICIOSA*—Q. Amongst rotten bark, near stem of gum-tree.

ORDER XXXIV.—PEZIZACEÆ, FRIES.

1869. *DASYSCYPHA TERRESTRIS*—This form should be restored to its original genus, *Helotium terrestre*, Berk. and Broome, Linn. Trans. II., 69 (1883). Cooke remarks in Grev. XX, 36 (1891), that this species was originally described in error as being externally villous, whereas it is externally smooth and naked; hence it was wrongly transferred by Saccardo to *Dasyscypa*.

ORDER XXXIX.—PHACIDIACEÆ, FRIES.

1894. *PSEUDOPEZIZA MEDICAGINIS*—On both surfaces of leaflets of *Medicago sativa* in Victoria.

ORDER XLI.—GYMNOASCEÆ, BAR.

1901. *EXOASCUS DEFORMANS*—T., W.A. On leaves of peaches and nectarines in West Australia.

GROUP VII.—HYPHOMYCETES, MARTIUS.

ORDER XLV.—MUCEDINACEÆ, LINK.

1911. *OIDIUM LEUCOCONIUM*—T.
 1914. *OIDIUM TUCKERI*—W.A.
 1921. *PENICILLIUM GLAUCUM*—T.

ORDER XLVI.—DEMATIACEÆ, FRIES.

1947. *FUSICLADIUM DENDRITICUM*—W.A.
 1948. *FUSICLADIUM PYRINUM*—W.A.
 1962. *HELMINTHOSPORIUM RAVENELII*—N.S.W.
 1982. *MACROSPORIUM TOMATO*—N.S.W.

GROUP VIII.—SPHÆROPSIDES, LEV.

ORDER XLIX.—SPHÆRIOIDACEÆ, SACC.

2025. *PHYLLOSTICTA CIRCUMSCISSA*—W.A. On apricots.
 2039. *PHOMA AMPELINA*—W.A. On vines.
 2058. *PHOMA UVICOLA*—N.S.W.
 2102. *PHLEOSPORA MORI*—T.
-

C.—LIST OF AUSTRALIAN EDIBLE FUNGI.

In the body of the work a number of species are marked Edible, but it is thought desirable to show them together, and give as complete a list as possible. Edible Fungi refers to those which may be eaten with impunity, not necessarily to those which may be eaten with relish. Our native species have still to be tested in most cases, but I have mainly given those Australian species which have been found wholesome in Britain or America. The number is 84.

No.	Name.	No.	Name.
6.	<i>Amanita ovoidea</i> .	602.	<i>Boletus æstivalis</i> .
15.	<i>Amanitopsis vaginata</i> .	603.	<i>Boletus alliciens</i> .
20.	<i>Lepiota cepæstipes</i> .	606.	<i>Boletus badius</i> .
25.	<i>Lepiota excoziata</i> .	611.	<i>Boletus edulis</i> .
32.	<i>Lepiota mastoidea</i> .	612.	<i>Boletus elegans</i> .
35.	<i>Lepiota naucina</i> .	616.	<i>Boletus granulatus</i> .
38.	<i>Lepiota procera</i> .	621.	<i>Boletus luteus</i> .
39.	<i>Lepiota rhacodes</i> .	629.	<i>Boletus scaber</i> .
47.	<i>Armillaria mellea</i> .	641.	<i>Fistulina hepatica</i> .
56.	<i>Tricholoma nudum</i> .	685.	<i>Polyporus intybaceus</i> .
63.	<i>Clitocybe cerussata</i> .	690.	<i>Polyporus myelodes</i> .
65.	<i>Clitocybe expallens</i> .	691.	<i>Polyporus Mylittæ</i> (<i>Sclerotium</i> , known as "Native Bread").
67.	<i>Clitocybe fumosa</i> .	699.	<i>Polyporus picipes</i> .
69.	<i>Clitocybe infundibuliformis</i> .	723.	<i>Polyporus sulphureus</i> .
71.	<i>Clitocybe laccata</i> .	728.	<i>Polyporus tumulosus</i> (eaten by Aborigines).
73.	<i>Clitocybe pruinosa</i> .	993.	<i>Hydnum coralloides</i> .
81.	<i>Collybia esculenta</i> .	1004.	<i>Hydnum lævigatum</i> .
82.	<i>Collybia fusipes</i> .	1011.	<i>Hydnum repandum</i> .
93.	<i>Collybia radicata</i> .	1037.	<i>Craterellus cornucopioides</i> .
176.	<i>Pleurotus ostreatus</i> .	1183.	<i>Sparassis crispa</i> .
178.	<i>Pleurotus petaloides</i> .	1188.	<i>Clavaria aurea</i> .
181.	<i>Pleurotus pulmonarius</i> .	1189.	<i>Clavaria botrytes</i> .
182.	<i>Pleurotus salignus</i> .	1194.	<i>Clavaria cristata</i> .
195.	<i>Hygrophorus coccineus</i> .	1196.	<i>Clavaria fastigiata</i> .
202.	<i>Hygrophorus miniatus</i> .	1197.	<i>Clavaria flava</i> .
208.	<i>Hygrophorus virgineus</i> .	1198.	<i>Clavaria formosa</i> .
209.	<i>Lactarius pallidus</i> .	1223.	<i>Clavaria rugosa</i> .
210.	<i>Lactarius piperatus</i> .	1233.	<i>Hirneola anricula-judæ</i> .
214.	<i>Russula alutacea</i> .	1237.	<i>Hirneola polytricha</i> .
227.	<i>Cantharellus cibarius</i> .	1245.	<i>Tremella lutescens</i> .
269.	<i>Marasmius scorodonius</i> .	1289.	<i>Clathrus cibarius</i> .
313.	<i>Lentinus tigrinus</i> .	1392.	<i>Lycoperdon Bovista</i> .
320.	<i>Panus conchatus</i> .	1397.	<i>Lycoperdon gemmatum</i> .
331.	<i>Panus torulosus</i> .	1400.	<i>Lycoperdon lilacinum</i> .
367.	<i>Volvaria bombycina</i> .	1776.	<i>Cyttaria Gunnii</i> .
400.	<i>Pholiota mutabilis</i> .	1777.	<i>Morchella conica</i> .
403.	<i>Pholiota præcox</i> .	1778.	<i>Morchella deliciosa</i> .
404.	<i>Pholiota pudica</i> .	1779.	<i>Morchella esculenta</i> .
496.	<i>Agaricus arvensis</i> .	1780.	<i>Morchella semilibera</i> .
497.	<i>Agaricus campestris</i> .	1783.	<i>Leotia lubrica</i> .
500.	<i>Agaricus silvaticus</i> .	1797.	<i>Peziza cochleata</i> .
528.	<i>Coprinus comatus</i> .		
601.	<i>Boletus æreus</i> .		

TABLE I.

Number of Orders, Genera, Species, and Varieties in the different Groups and different Colonies, together with those common to Britain.

Groups.	No. of Orders.	No. of Genera.	No. of Species.	No. of Varieties.	Australian.				W.A.				S.A.				T.				V.				N.S.W.				Q.				B.			
					Orders.	Genera.	Species.	Varieties.	Orders.	Genera.	Species.	Varieties.	Orders.	Genera.	Species.	Varieties.	Orders.	Genera.	Species.	Varieties.	Orders.	Genera.	Species.	Varieties.	Orders.	Genera.	Species.	Varieties.	Orders.	Genera.	Species.	Varieties.				
1. Hymenomyces ...	6	106	1,266	28	4	14	19	...	6	57	137	2	6	61	151	5	6	64	186	2	6	89	600	5	6	62	247	8	6	86	618	11	6	81	455	5
2. Gastromycetes ...	6	39	177	2	3	4	5	...	5	18	44	...	4	12	25	...	6	16	28	...	5	28	72	...	6	18	42	1	5	24	88	1	6	16	30	...
3. Uredines ...	1	9	90	2	...	...	...	...	1	1	1	...	1	6	21	...	1	4	12	...	1	7	58	1	1	5	23	...	1	8	28	...	1	6	25	...
4. Pyrenomycetes ...	18	83	253	4	8	9	9	...	9	16	19	...	9	13	13	...	8	22	36	...	14	48	89	1	7	19	27	...	15	49	126	3	11	33	53	...
5. Discomycetes ...	10	43	128	3	2	3	3	...	3	7	8	...	3	7	10	...	8	24	35	...	8	30	68	2	5	9	12	...	7	22	37	1	8	23	49	1
6. Tuberoïdes ...	3	3	3	...	...	...	...	...	...	...	...	...	1	1	1	...	1	1	1	...	...	...	...	...	...	...	...	...	1	1	1	...	...	...	...	...
7. Hyphomycetes ...	4	52	123	2	2	2	2	...	4	10	13	...	3	12	14	...	4	11	13	...	4	30	59	...	4	19	28	...	4	38	60	2	4	32	50	...
8. Sphaeropsides ...	5	43	128	...	...	...	...	...	2	3	3	...	2	5	8	...	2	4	4	...	5	31	77	...	2	5	8	...	4	23	53	1	2	8	13	...
9. Saccharomycetes ...	1	1	10	...	1	1	10	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	1	1	8	...
10. Ustilagines ...	1	12	39	4	1	1	...	1	...	...	...	...	1	5	14	3	1	2	2	...	1	7	23	3	1	5	10	3	1	8	16	3	1	6	9	3
11. Phycomycetes ...	5	12	15	1	...	...	...	...	1	1	1	...	3	3	3	...	1	1	2	...	4	9	11	1	2	4	5	...	3	5	5	...	5	8	9	...
12. Myxomycetes ...	9	21	52	1	1	1	2	...	6	12	16	...	2	2	2	...	7	12	20	...	6	12	13	...	4	4	4	...	7	13	28	1	9	20	38	...
Totals ...	69	424	2,284	47	22	35	50	1	37	125	242	2	35	127	262	8	45	161	339	2	54	291	1,070	13	38	150	406	12	54	277	1,060	23	54	234	739	9

TABLE II.

Number of Australian Species compared with British and total known Species.

Groups.				No. of Australian Species (1894).	No. of British Species (1892).	Total known Species (1892), (Approximate).
1. Hymenomycetes ...	...	...	...	1,266	1,902	10,163
2. Gastromycetes ...	...	...	...	177	78	718
3. Uredines ...	...	...	...	90	53	1,428
4. Pyrenomycetes ...	...	...	...	253	1,275	9,247
5. Discomycetes ...	...	...	...	128		3,944
6. Tuberoïdes ...	...	...	...	3		145
7. Hyphomycetes ...	...	...	...	123	580	5,004
8. Sphærospides ...	...	...	...	128	685	6,745
9. Saccharomycetes ...	...	...	...	10	8	30
10. Ustilagines ...	...	...	...	39	177	329
11. Phycomycetes ...	...	...	...	15	145	686
12. Myxomycetes ...	...	...	...	52	137	510
Other Groups ...	...	...	...	...	...	714
Totals ...	...	...	...	2,284	5,040	39,663

TABLE III.

Number of Species of Fungi found in Victoria on the following Plants of economic importance.

Name of Plant.	No. of Species of Fungi.	Name of Plant.	No. of Species of Fungi.
Acacia species	16	Lucerne	2
Almond	3	Maize	1
Apple	5	Mulberry	1
Apricot	2	Oats	4
Bean	1	Onion..	2
Beet	1	Orange	1
Cabbage	1	Pea	2
Cauliflower	1	Peach	4
Celery	1	Pear	3
Cherry	1	Plum	3
Clover	3	Rye-grass	3
Eucalyptus species	54	Strawberry	1
Flax	1	Tomato	2
Garlic	1	Vine	8
Lemon	1	Wheat	10
Lettuce	1		

Total number of Orders	69
" " Genera	424
" " Species	2,284
Species common to Australia and Britain	739
Species in West Australia	242
" South Australia... ..	262
" Tasmania	339
" Victoria	1,070
" New South Wales	406
" Queensland	1,060
Species not referred to their respective Colonies	50

Proportion of Species of Fungi found in the different Colonies:—

West Australia	10·6 per cent.
South Australia	11·5 "
Tasmania	14·8 "
Victoria	46·8 "
New South Wales... ..	17·8 "
Queensland	46·4 "

NOTE.—It is not to be inferred from the relatively high percentage of Fungi in Victoria and Queensland, for instance, as compared with New South Wales, that they are absent from the latter colony, but rather that they still await investigation and determination there.

HOST-INDEX OF AUSTRALIAN FUNGI.

II.—HOST-INDEX OF AUSTRALIAN FUNGI.

- Abrotanella forsterioides**, J. Hook.
2266. *Aecidium monocystis*.—Leaves.
- Abutilon Avicennæ**, Ger.
1477. *Puccinia heterospora*.—Leaves.
- Abutilon crispum**, Don.
1477. *Puccinia heterospora*.—Leaves.
- Acacia** sp.
984. *Merulius pelliculosus*. V.—Branches.
1071A. *Stereum hirsutum*, var. *tinellum*.—Rotten wood.
1103. *Stereum vittaeforme*.—Bark.
1456. *Uromyces phyllodii*.—Phyllodes.
1463. *Melampsora phyllodiorum*.—Phyllodes.
1521. *Uredo leguminum*.—Pods.
1655. *Trabutia parvicapsa*. V.—Phyllodes.
1743. *Meliola amphitricha*. V.—Leaves.
2024. *Actinomma Gastonis*.—Phyllodes.
2032. *Phyllosticta phyllodiorum*. V.—Phyllodes.
2061. *Asteromella Acaciæ*. V.—Phyllodes.
2082. *Diplodia phyllodiorum*. V.—Phyllodes.
2093. *Septoria epiphyllodea*. V.—Phyllodes.
2140. *Marsonia Acaciæ*. V.—Phyllodes.
2145. *Pestalozzia Acaciæ*. V.—Leaves.
- Acacia complanata**, Cunn.
2080. *Diplodia lichenopsis*.—Phyllodes.
- Acacia hakeoides**, Cunn.
1458. *Uromyces Tepperianus*. V.—Branches.
- Acacia harpophylla**, F. v. M.
1580. *Xylaria gracilis*.—Wood.
2117. *Melophia Woodsiana*.—Phyllodes.
- Acacia longifolia**, Willd.
1656. *Trabutia phyllodii*. V.—Phyllodes.
2099. *Septoria phyllodiorum*. V.—Phyllodes.
- Acacia melanoxylon**, R. Br. (Blackwood Tree).
2132. *Gleosporium leguminis*. V.—Legumes.
- Acacia myrtifolia**, Willd.
1458. *Uromyces Tepperianus*. V.—Branches.
- Acacia notabilis**, F. v. M.
1448. *Uromyces digitatus*.—Phyllodes.
1522. *Uredo notabilis*. V.—Phyllodes.
- Acacia penninervis**, Sieb.
1641. *Phyllachora rhytismoides*.—Phyllodes.
- Acacia salicina**, Lind.
1450. *Uromyces fusisporus*. V.—Phyllodes.
1458. *Uromyces Tepperianus*. V.—Branches.
- Acacia suaveolens**, Willd. (Sweet-scented Acacia.)
1706. *Læstidia phyllodii*. V.—Phyllodes.
- Acacia verticillata**, Willd.
1695. *Physalospora labecula*.—Leaves.
- Acæna sanguisorbæ**, Vahl. (Sheep's-burr.)
1501. *Phragmidium Potentillæ*. V.—Leaves.
- Adiantum** sp. (Maiden-hair Fern.)
1166. *Cyphella filicola*. V.—Fronds.
- Ægiceras** sp.
1542. *Sphærostilbe dubia*.—Bark.
- Agathis robusta**, Masters = Dammar. (Queensland Kauri.)
1702. *Læstidia Dammaræ*.—Leaves.
- Agarics** (Fungi).
1555. *Hypomyces tomentosus*.
1559. *Ophionectria agaricicola*. V.—Putrid Agarics.
1932. *Verticillium niveum*.—Dead Agarics.
1972. *Heterosporium epinyces*. V.—Decayed Agarics.
2022. *Myrothecium inundatum*.—Putrid Agarics.
2196. *Spinellus gigasporus*. V.—Decaying Agarics.
- Allium** sp.
2038. *Phoma alliicola*. V.—Scapes.
- Allium cepa**, L. (Onion.)
2202. *Peronospora Schleideni*. V.—Leaves.
2247. *Physarum didermoides*.—Scales.
2281. *Ustilago Allii*. V.—Scale leaves of stored Onion bulbs.
- Allium sativum**, Bauh. (Garlic.)
2202. *Peronospora Schleideni*. V.—Leaves.
- Alopecurus** sp. (Fox-tail Grass.)
2199. *Sclerospora macrospora*. V.
1493. *Puccinia Rubigo-vera*. V.—Leaves.
- Alphitonia excelsa**, Reiss. (Red Ash.)
2121. *Gleosporium Alphitoninæ*.—Leaves.
2148. *Pestalozzia Guepini*.—Leaves.
- Alpinia cœrulea**, Benth.
1634. *Phyllachora Alpininæ*.—Leaves.
- Alsophila** sp.
1652. *Rhytisma filicinum*.—Fronds.
- Alsophila Rebeccæ**, F. v. M.
1735. *Asterella Alsophilæ*.—Fronds.
- Althæa rosea**, Cav. (Hollyhock.)
1485. *Puccinia Malvacearum*. V.—Leaves and stems.
- Alyxia buxifolia**, R. Br.
1467. *Puccinia Alyxiæ*. V.—Leaves.
1707. *Sphærella Alyxiæ*. V.—Leaves.
- Amphipogon strictus**, R. Br. (Bearded-heads.)
2175. *Ustilago Tepperi*.
- Andropogon** sp.
1943. *Periconia nigrella*.—Leaves.

Andropogon contortus, L. = *Heteropogon contortus*.
(Bunch Spear-grass.)
2189. *Cerebella Andropogonis*.

Ant-Formica sp.

Anthistiria ciliata, L. fil. (Kangaroo-grass.)
2158. *Ustilago bromivora*. V.—Inflorescence.
2283. *Tolyposporium Anthistiriae*.—Inflorescence.

Anthistiria frondosa, R. Br.
2160. *Ustilago bursa*.—Grain.

Aphides. (On Pumpkin Leaves.)
1905. *Oospora Aphides*.

Apium graveolens, L. (Celery.)
1468. *Puccinia Apii*. V.

Apple—*Pyrus Malus*, L.

Apricot—*Prunus Armeniaca*, L.

Aristida sp.
2173. *Ustilago segetum*. V.

Artichoke—*Cynara Scolymus*, L.

Arundo sp. (Reeds.)
1936. *Coniosporium iniquians*.—Stems.

Arundo Phragmites, L. (Thatch-reed.)
1484. *Puccinia Magnusiana*.
1488. *Puccinia Phragmitis*. V.
1497. *Puccinia Tepperi*.

Aster argophyllus, Labill. = *Olearia*. (Musk Tree.)
1736. *Asterella subcuticulosa*. V.—Fading and dead leaves.
1999. *Podosporium grande*. V.—Stems.

Asterina sp. Growing on leaflets of *Tarrietia trifoliolata*.
F. v. M.
2275. *Belonidium parasiticum*.

Atherosperma moschatum, Labill. (Native Sassafras.)
2221. *Arcyria fuliginea*.—Leaves.

Aucuba japonica, L.
1700. *Pleospora Aucubæ*. V.—Leaves.

Auricularia sp. (Fungus.)
2236. *Didymium australe*.

Australian Beech—*Fagus Cunninghamii*, Hook.

Avena fatua, L. (Wild Oats.)
1475. *Puccinia graminis*. V.

Avena sativa, L. (Oats.)
1475. *Puccinia graminis*. V.
1493. *Puccinia rubigo-vera*. V.
1950. *Scolecotrichum graminis*, var. *Avenæ*. V.
2173B. *Ustilago segetum*, var. *Avenæ*. V.—Ear.

Backhousia sp.
2083. *Aschyta apiospora*.—Leaves.

Bambusa sp. (Bamboo.)
501. *Agaricus versipes*.—Roots.
1143. *Peniophora bambusicola*.—Rotting bamboo.

Banana—*Musa Cavendishii*, Lamb.

Banksia sp. (Native Honeysuckle.)
123. *Mycena subcorticalis*.—Log.
1699. *Stereum umbrinum*.—Bark.
1692. *Didymosphaeria Banksiæ*. V.—Leaves.
1882. *Tympanis Toomansii*.—Fruit.

Banksia integrifolia, L. (Beefwood.)
1709. *Spharella Banksiæ*. V.—Fading leaves.

Banksia marginata, Cav.
1741. *Parodiella Banksiæ*.—Languid leaves.

Barley—*Hordeum distichon*, Bauh.

Bean—*Phaseolus vulgaris*, L.

Bean Caper—*Zygophyllum ammophilum*, F. v. M.

Bearded-heads—*Amphipogon strictus*, R. Br.

Beefwood—*Banksia integrifolia*, L.

Beet—*Beta vulgaris*, L.

Beilschmiedia obtusifolia, Benth. = *Nesodaphne*.
1462. *Melampsora Nesodaphnes*.—Fruit.

Bellis perennis, L. (Daisy.)
1487. *Puccinia obscura*, Schrot. V.

Bertya rotundifolia, F. v. M.
1942. *Heterobotrys paradoxa*.—Leaves.

Beta vulgaris, L. (Beet.)
1446. *Uromyces Betæ*. V.—Leaves.

Betula alba, L. (Birch.)
650. *Polyporus betulinus*.

Bidens pilosus, L.
1519. *Uredo Cichoracearum*. V.

Birch—*Betula alba*, L.

Bitter Almond—*Prunus Amygdalus* var. *Amara*.

Bitter Bark—*Tabernæmontana orientalis*, R. Br.

Black Ash—*Litsea dealbata*, Nees.

Blackbutt Tree—*Eucalyptus pilularis*, Sm.

Blackwood Tree—*Acacia Melanoxylon*, R. Br.

Bloodwood Tree—*Eucalyptus corymbosa*, Sm.

Bluegum Tree—*Encalyptus globulus*, Labill.

Boletus sp. (Fungus.)
1552. *Hypomyces chrysospermus*. V.
1929. *Sepedonium chrysospermum*. V.

Bossia cinerea, R. Br.
2265. *Ecidium eburneum*. V.—Stems, leaves, flowers, and fruit.

Bottle Gourd—*Lagenaria vulgaris*, Ser.

Bottle Thistle—*Lagenophora Billardieri*, Cass.

Bougainvillea sp.
240. *Marasmius calobates*. V.—Putrid leaves.

Box Eucalypt—*Eucalyptus hemiphloia*, F. v. M.
,, *Eucalyptus largiflorens*, F. v. M.

Box Thorn—*Bursaria spinosa*, Cav.

Bramble—*Rubus fruticosus*, L.

Brassica oleracea, L. (Cabbage and Cauliflower.)
2258. *Plasmodiophora Brassicæ*. V.—Roots.

- Brisbane Box**—*Tristania conferta*, R. Br.
- Bromus** sp.
2092. *Septoria Bromi*. V.—Leaves.
- Bromus arenarius**, Labill.
2092. *Septoria Bromi*. V.—Leaves.
2158. *Ustilago bromivora*. V.—Inflorescence.
- Bromus mollis**, L.
2158. *Ustilago bromivora*. V.—Inflorescence.
- Bromus sterilis**, L.
1475. *Puccinia graminis*. V.
- Bulbine bulbosa**, Haw.
1447. *Uromyces Bulbinis*. V.—Leaves.
- Bulrush**—*Typha* sp.
- Bunch Spear-grass**—*Andropogon contortus*, L.
- Burchardia umbellata**, R. Br.
1471. *Puccinia Burchardiae*. V.—Leaves.
- Bursaria spinosa**, Cav. (Boxthorn)
1609. *Nummularia pusilla*.—Branches.
1759. *Capnodium Walteri*. V.—Branches and living leaves.
1945. *Botryotrichum Lachnella*. V.—Branches and spines.
- Cabbage**—*Brassica oleracea*, L.
- Callistemon** sp.
1751. *Meliola polytricha*.—Leaves.
- Camellia japonica**, L. (Camellia.)
1718. *Sphaerulina Camelliae*. V.—Leaves.
1976. *Macrosporium Camelliae*. V.—Leaves.
2114. *Sacidium Camelliae*. V.—Leaves.
- Canthium latifolium**, F. v. M.
2079. *Diplodia canthifolia*. V.—Leaves.
- Capparis** sp. (Caper.)
1693. *Didymospheria conoidella*.—Dead branches.
- Capparis Mitchelli**, Lind.
1550. *Calonectria otagensis*.—Twigs.
- Carex** sp. (Sedge.)
1943. *Periconia nigrella*.—Leaves.
2168. *Ustilago leucoderma*. V.—Sheaths.
- Carica Papaya**, L. (Papaw.)
1980. *Macrosporium peponicolum*.—Fruit.
- Carissa ovata**, R. Br.
1472. *Puccinia Carissæ*.—Leaves.
- Carnation**—*Dianthus Caryophyllus*, L.
- Cassia** sp.
1523. *Uredo pallidula*.—Leaves, twigs, and legumes.
2001. *Tubercularia leguminum*.—Legumes.
- Cassinia aculeata**, R. Br.
1674. *Gibberidea plagia*. V.—Twigs.
- Castanospermum australe**, Cunn. and Fraser. (Moreton Bay Chestnut.)
1733. *Asterina platystoma*.—Leaves.
2064. *Chaetophoma entricha*.—Leaves.
- Castor-oil Plant**—*Ricinus communis*, L.
- Casuarina** sp. (Sheoak.)
761. *Fomes igniarius*. V.—Trunks.
936. *Hexagonia decipiens*. V.—Trunks.
1864. *Trichopeziza Sphaerula*.—Dead bark.
1865. *Dasycephala Eucalypti*. V.—Leaves.
2146. *Pestalozzia Casuarinae*. V.—Branches.
- Cat's-ear**—*Hypochaeris glabra*, L.
- Cauliflower**—*Brassica oleracea*, L.
- Celery**—*Apium graveolens*, L.
- Cerastium glomeratum**, Thuill.
1893. *Pseudopeziza Cerastiorum*.—Leaves and calyx.
- Cherry**—*Prunus Cerasus*, L.
- Chiloglottis diphylla**, R. Br.
1455. *Uromyces orchidearum*. V.—Leaves.
- Chionaspis Citri**. (Coccus of Orange.)
2017. *Microcera rectispora*.
- Chrysanthemum** sp.
1909. *Oidium Chrysanthemi*. V.—Leaves.
- Cicada** sp.
1991. *Isaria Cicadae*. V.
- Citrus** sp.
1136. *Corticium nudum*.—Bark.
1756. *Capnodium citri*. V.—Leaves.
- Citrus Aurantium**, L. (Orange.)
2124. *Gloeosporium citricolum*.—Leaves.
1756. *Capnodium citri*. V.—Leaves.
- Citrus Limonium**, Risso (Lemon.)
2123. *Gloeosporium Citri*. V.—Branches.
- Cladium** = *Gahnia*.
- Clavaria** sp. (Fungus.)
1930. *Verticillium eximium*.
- Clematis aristata**, R. Br. (Native Supple-jack.)
1520. *Uredo Clematidis*. V.—Leaves.
- Clematis microphylla**, D. C.
1520. *Uredo Clematidis*. V.—Leaves.
- Clover**—*Trifolium* sp.
- Club Rush**—*Scirpus nodosus*, Rott.
- Coccus** sp.
2016. *Microcera coccophila*.
- Compositæ**.
1506. *Æcidium Compositarum*. V.
2205. *Synchytrium Taraxaci*. V.—Leaves.
- Coniferæ**.
47. *Armillaria mellea*. V.—Stumps.
724. *Polyporus tabulæformis*.—Trunks.
735. *Fomes annosus*.—Trunks.
1755. *Capnodium australe*.—Branches.
1940. *Hormiscium pithyophilum*.—Branches and leaves.
- Coprosma Billardieri**, J. Hook.
2267. *Puccinia Coprosmatis*. V.—Leaves.
- Cordyline australis**, Hook.
2041. *Phoma Cordylines*.
- Cordyline terminalis**, Kunth.
2026. *Phyllosticta Cordylines*.—Leaves.

Cordyline terminalis, var. *Cannæfolia*.
1640. *Phyllachora nervisequia*.—Living and dry leaves.

Correa Lawrenciana, Hook.
1729. *Asterina correicola*. V.—Leaves.
2261. *Puccinia Correæ*.—Leaves.

Cotula sp.
1896. *Fabraea rhytismoides*. V.—Leaves.

Crinum pedunculatum, R. Br.
2041. *Phoma Cordylines*.—Old leaves.

Cruciferae.
1505. *Accidium Barbareæ*.
2198. *Cystopus candidus*. V.—Leaves, stems, &c.
2258. *Plasmodiophora Brassicæ*. V.—Roots.

Cryptandra parvifolia, Turcz. = *Spyridium*.
1525. *Uredo Spyridii*. V.—Leaves.

Cucumis Melo, L. (Melon.)
2131. *Glœosporium lagenarium*—Epicarp.
2131A. *Glœosporium lagenarium*, var. *Cucurbitarum*.

Cucurbita sp.
2131A. *Glœosporium lagenarium*, var. *Cucurbitarum*.—Fruit.

Cucurbita lagenaria = *Lagenaria vulgaris*.

Cucurbitaceæ.
1697. *Didymella Bryoniæ*.—Twigs.

Cudrania javanensis, Tréc.
1653. *Rhytisma hypoxanthum*.—Leaves.

Cupania sp.
1743. *Meliola amphitricha*. V.—Leaves.

Cupania anacardioides, Rich.
2151. *Pestalozzia versicolor*.—Leaves.

Cycas sp.
1755. *Capnodium australe*.—Branches.
2216. *Perichæna appianata*.—Rotting Cycads.

Cynara Scolymus, L. (Artichoke.)
1981. *Macrosporium Readeri*. V.—Dry stems.

Cyperus sp.
2157. *Ustilago axicola*. V.—Fruits and panicles.

Cyperus Incidus, R. Br.
2161. *Ustilago catenata*.—Spikes.

Daisy—*Bellis perennis*, L.

Danthonia sp. (Oat grass.)
2165. *Ustilago destruens*. V.—Inflorescence.
2168. *Ustilago leucoderma*. V.—Sheaths.
2173. *Ustilago segetum*. V.

Danthonia penicillata, F. v. M.
2165. *Ustilago destruens*. V.
2175. *Ustilago Tepperi*.—Flowers and upper portions of stem.

Darnel—*Lolium temulentum*, L.

Date Palm—*Phoenix dactylifera*, L.

Daviesia latifolia, R. Br.
1965. *Cercospora Daviesiæ*. V.—Fading leaves.
2042. *Phoma Daviesiæ*. V.—Dead leaves.

Dendrobium speciosum, Smith.
2088. *Diplodina Dendrobii*.—Leaves.

Dianthus caryophyllus, L. (Carnation.)
2280. *Septoria Dianthi*. V.—Leaves.

Dichondra repens, R. and G. Forst.
1470. *Puccinia Berkeleyana*. V.—Leaves.

Dicksonia sp.
1790. *Geoglossum Walteri*. V.—Stems.

Dioscorea sp. (Native Yam.)
1717. *Sphærella smilacicola*.—Leaves.

Diploglottis Cunninghamii, J. Hook.
1449. *Uromyces Diploglottidis*.—Fading leaves.
2043. *Phoma Diploglottidis*.—Fading leaves.

Ditch Millet—*Paspalum serobiculatum*, L.

Dock—*Rumex* sp.

Drake—*Lolium temulentum*, L.

Drimys aromatica, F. v. M. = *Tasmannia aromatica*.
(Native Pepper Tree.)
1651. *Darwiniella globulosa*.—Leaves.

Dwarf Mallow—*Malva rotundifolia*, L.

Elæodendron sp.
2147. *Pestalozzia funerea*.—Leaves.

Encephalartos Denisonii, F. v. M. = *E. Hopei*. = *Macrozamia*.
436. *Flammula picrea*. V.—Dead trunks.
1669. *Cryptosphærella Macrozamia*.—Fruit.
2067. *Dothiorella pericarpica*.—Pericarp.
2125. *Glœosporium Denisonii*.—Leaves.

Epilobium glabellum, Forst.
1473. *Puccinia Epilobii*. V.—Leaves.

Eragrostis Brownei, Nees.
1538. *Hypocrella axillaris*.

Erechtites quadridentata, D. C.
2262. *Puccinia Erechtitis*. V.

Eriachne sp.
2156. *Ustilago australis*. V.—Spikelets.
2184. *Sorosporium Eriachnes*.—Spikelets.

Erythrina vespertilio, Benth.
2239. *Didymium pezizoidem*.—Dead wood.

Eucalyptus sp.
264. *Marasmius putredinis*. V.
987. *Xylostroma giganteum*. V.—Heart-wound

A. ON LEAVES.

258. *Marasmius lignyodes*. V.
259. *Marasmius meloniformis*.—Leaves and branches
1639. *Phyllachora maculata*. V.
1644. *Dothidella inæqualis*. V.
1646. *Montagnella Eucalypti*. V.
1647. *Montagnella rugulosa*. V.
1650. *Bagnisiella rugulosa*. V.
1654. *Trabutia Eucalypti*. V.
1710. *Sphærella cryptica*. V.
1715. *Sphærella nubilosa*. V.
1743. *Meliola amphitricha*. V.
1746. *Meliola densa*.
1763. *Anlographum Eucalypti*. V.
1857. *Phialea ceratina*.
1865. *Dasycephala Eucalypti*. V.
1891. *Stictis emarginata*. V.
1897. *Coccomyces delta*. V.

1940. *Hormiscium pithyophilum*.—Leaves and branches.
 1966. *Cercospora epicoccoides*. V.
 1967. *Cercospora Eucalypti*. V.
 2008. *Hymenium Eucalypti*. V.
 2015. *Fusarium rubicolor*.—Spreading over galls on leaves.
 2040. *Phoma australis*. V.
 2055. *Phoma purpurea*.
 2066. *Dothiorella Eucalypti*. V.
 2071. *Sphaeropsis phomatoides*. V.
 2090. *Stagonospora obicularis*. V.
 2091. *Camarosporium Eucalypti*. V.
 2107. *Martinella Eucalypti*. V.
 2108. *Leptothyrium aristatum*. V.
 2109. *Leptothyrium Eucalyptorum*. V.—Fallen leaves.
 2110. *Piggotia substellata*. V.
 2111. *Melasma Eucalypti*. V.
 2118. *Leptostromella Eucalypti*. V.
 2142. *Stilbospora foliorum*. V.
 2149. *Pestalozzia monochaeta*.

B. ON BRANCHES.

249. *Marasmius Eucalypti*.—And fruit.
 259. *Marasmius meloniformis*.—And leaves.
 1899. *Patinella Adamsoni*. V.
 1940. *Hormiscium pithyophilum*.—And leaves.
 2089. *Hendersonia Eucalypti*. V.—And leaves.

C. ON BARK.

161. *Pleurotus Eucalyptorum*.
 292. *Lentius hepatoirichus*.

D. ON TRUNKS.

668. *Polyporus Eucalyptorum*. V.
 715. *Polyporus spumeus*. V.
 772. *Fomes obliquus*. V.
 816. *Polystictus Eucalypti*. V.
 940. *Hexagonia Muelleri*.
 1069. *Stereum fasciatum*. V.
 1109. *Hymenochaete Kalkbrenneri*. V.
 1314. *Cyathus pusio*.

E. ON ROTTEN WOOD.

324. *Panus lateritius*.

Eucalyptus amygdalina, Labill. (Peppermint Gumtree.)

149. *Pleurotus affixus*.—Bark of young trees.
 987. *Xylostroma giganteum*. V.—Wood.
 1719. *Microthyrium amygdalinum*.—Living leaves.
 1754. *Asteridium Eucalypti*.—Dead leaves.

Eucalyptus corymbosa, Sm. (Bloodwood Tree.)

987. *Xylostroma giganteum*. V.—Wood.

Eucalyptus Globulus, Labill. (Bluegum Tree.)

2005. *Illosporium obscurum*. V.—Leaves.
 2027. *Phyllosticta Eucalypti*. V.—Leaves.
 2044. *Phoma eucalyptidea*. V.—Living or fading leaves.
 2050. *Phoma Molleriana*. V.—Fallen leaves.
 2115. *Sacidium Eucalypti*. V.—Dead leaves.

Eucalyptus hemiphloia, F. v. M. (Box Eucalypt.)

698. *Polyporus phlebochrous*.—Stems.
 1410. *Xyloporium australe*. V.—Trunk.

Eucalyptus incrassata, Labill. (Mallee.)

2120. *Protostegia Eucalypti*. V.—Dead leaves.

Eucalyptus largiflorens, F. v. M. = *E. bicolor*. (Box Eucalypt.)

2139. *Pestalozziella circulare*. V.—Dead leaves.

Eucalyptus maculata, Hook. (Spotted Gum.)

987. *Xylostroma giganteum*.—Wood.

Eucalyptus microtheca, F. v. M.

- 2207A. *Tubulina cylindrica*, var. *Nitidissima*.

Eucalyptus obliqua, L'Her. (Messmate = Stringybark Tree.)

899. *Poria mollusca*. V.—Dead bark.

Eucalyptus pauciflora, Sieber. (Whitegum Tree.)

2135. *Glæosporium nigricans*. V.—Leaves.
 2139. *Pestalozziella circulare*. V.—Dead leaves.

Eucalyptus pilularis, Smith. (Blackbutt Tree.)

1731. *Asterina microthyrioides*. V.—Leaves.
 1768. *Lembosia orbicularis*.—Leaves.

Eucalyptus siderophloia, Benth. (Ironbark Tree.)

987. *Xylostroma giganteum*.—Wood.

Eucalyptus tereticornis, Smith. (Flooded Gumtree.)

474. *Crepidotus haustellaris*.—Rotten trunks.
 1720A. *Micropeltis applanata*, var. *depauperata*.—Languid leaves. *

Eucalyptus viminalis, Labill. (Manna Gumtree.)

150. *Pleurotus applicatus*.—V.
 1690. *Rhamphoria tenella*.—Rotten wood.
 2059. *Phoma viminalis*. V.—Leaves.
 2143. *Coryneum viminale*. V.—Leaves.

Eucalyptus virgata, Sieb. (Mountain Ash.)

987. *Xylostroma giganteum*.—Wood.

Eugenia sp.

1734. *Asterina reptans*.—Leaves.
 1744. *Meliola cladotricha*. V.—Leaves.
 2179. *Entyloma Eugeniæ*.—Leaves.

Euonymus sp. (Spindle Tree.)

1711. *Sphaerella Euonymi*. V.—Dead leaves.

Everlastings—*Helichrysum* sp.

Exocarpos latifolia, R. Br.

250. *Marasmius Exocarpi*. V.—Trunks.

Fagus Cunninghamii, Hook. (Native Beech.)

1776. *Cyttaria Gunnii*. V.—Branches.

Fennel—*Feniculum vulgare*, Mill.

Ferns.

110. *Mycena flavovirens*. V.—Tree ferns.
 122. *Mycena stylobates*. V.
 158. *Pleurotus elitocyboides*. V.—Stems.
 459. *Naucoria siparia*. V.—Stems.
 568. *Mycena pterigena*. V.—Stems.
 998. *Hydnum filicola*. Stems.
 1226. *Clavaria tasmanica*.—Tree ferns.
 1760. *Antennaria Robinsoni*.—V.
 1762. *Antennaria semiovata*.

Ficus sp. (Fig.)

1648. *Bagnisiella catervaria*.—Leaves.

Ficus aspera, Forst.

2012. *Fusarium hypocreoides*.—Fading leaves.

Ficus Carica, L. (Fig.)

1641. *Phyllachora rhytismoides*.—Leaves.

Fimbristylis sp.

1636. *Phyllachora Fimbristylis*.
 2157. *Ustilago axicola*. V.—Fruits and panicles.

Fir.

1161. *Aleurodiscus amorphus*.—Trunks and branches.
1268. *Calocera stricta*. V.—Dead leaves.

Flat-leaved Rush—*Juncus planifolius*, R. Br.

Flat-weed—*Hypochaeris radicata*, L.

Flax—*Linum usitatissimum*, L.

Flindersia sp.

1743. *Meliola amphitricha*. V.—Leaves.
2270. *Xylaria ianthino-velutina*.—Old fruit.

Flindersia australis, R. Br.

1594. *Xylaria scopiformis*.—Decaying fruit.

Flooded Gum-tree—*Eucalyptus tereticornis*, Sm.

Fœniculum vulgare, Mill. (Fennel.)

1953. *Cladosporium herbarum*. V.—Stems.

Fomes gryphæformis, Cooke = *Polyporus* (Fungus).

1003. *Hydnum isidioides*.—Hymenium.

Formica sp. (Ant.)

1989. *Stilbum Formicarum*. V.—Dead ants.

Fox-tail Grass—*Alopecurus* sp.

Fragaria vesca, L. (Strawberry.)

1712. *Sphaerella Fragariae*. V.—Leaves.
2028. *Phyllosticta fragaricola*.—Leaves.

Gahnia filum, F. v. M. = *Cladium filum*.

2185. *Sorosporium Muellerianum*. V.—Panicles.

Gahnia tetraquetra, F. v. M.

2126. *Glebosporium epicaladii*.—V.

Garlic—*Allium sativum*, Bauh.

Geranium sp.

1680. *Venturia circinans*. V.—Leaves.

Glycine clandestina, Wendl.

1968. *Cercospora Glycines*. V.—Living leaves.

Gnaphalium purpureum, L.

2268. *Puccinia investita*, Schwein. V.—Leaves and stems.

Goodeniaceæ.

2204. *Synchytrium Succisæ*. V.—Leaves and leaf-stalks.

Goodenia sp.

1457. *Uromyces puccinioides*. V.—Leaves and flower-stalks.
1508. *Æcidium Goodeniacearum*. V.—Leaves.

Goodenia geniculata, R. Br.

1495. *Puccinia Saccardoi*. V.—Leaves.

Goodenia pinnatifida, Schlecht.

1504. *Æcidium Goodeniacearum*. V.—Leaves.

Goodenia ovata, Smith.

2046. *Phoma Goodeniarum*. V.—Fading leaves.
2137. *Glebosporium subglobosum*. V.—Fading leaves.

Goodia lotifolia, Sal.

1513. *Æcidium soleniforme*.
1713. *Sphaerella Goodiaefolia*. V.—Leaves.

Goose-grass—*Poa annua*, L.

Grape-vine—*Vitis vinifera*, Bauh.

Grass.

241. *Marasmius calopus*. V.—Roots.
451. *Naucoria frusticola*. Roots.
467A. *Tubaria inquilina*, var. *Echola*.—Roots.
1045. *Thelephora cristata*.—Running over grass.
1533. *Epichloe cinerea*.
1637. *Phyllachora graminis*. V.—Leaves.
1682. *Chaetomium comatum*. V.—Rotting grass.
1714. *Sphaerella graminicola*. V.—Leaves.
1724. *Erysiphe graminis*. V.—Leaves and stems.
1992. *Isaria graminiperda*.—V.
2051. *Phoma nitida*.—V.
2243. *Spumaria alba*.—V.
2256. *Leocarpus fragilis*.

Grass-tree—*Xanthorrhoea* sp.

Grevillea sp. (Silky Oak.)

169. *Pleurotus lampas*. V.—Languid stems.
1708. *Sphaerella atra*.—Leaves.
2273. *Phyllachora Grevilleae*.—Leaves.

Groundsel—*Senecio* sp.

Hakea sp.

1517. *Uredo angiosperma*.—Leaves.

Hakea lorea, R. Br.

1728. *Asterina Baileyi*.—Leaves.

Hardenbergia = *Kennedya*.

Hedycarya Cunninghami, Tul.

2129. *Glebosporium Hedycaryæ*. V.—Fading leaves.

Helianthus annuus, L. (Sun Flower.)

1476. *Puccinia Helianthi*. V.—Leaves.

Helichrysum sp. (Everlastings.)

1480. *Puccinia Kalkbrenneri*. V.—Leaves.

Hemarthria compressa, R. Br. (Sugar Grass.)

1493. *Puccinia Rubigo-vera*.

Heteropogon = *Andropogon*.

Hibiscus sp.

1477. *Puccinia heterospora*.—Leaves.

Holly—*Ilex Aquifolium*, L.

Hollyhock—*Althæa rosea*, Cav.

Hordeum distichon, Bauh. (Barley.)

1475. *Puccinia graminis*. V.
1493. *Puccinia Rubigo-vera*. V.
2173c. *Ustilago segetum*, var. *Nuda Hordei*. V.—Ear.

Hormogyne cotinifolia, D. C.

943. *Hexagonia sericea*.—Trunks.

House Fly—*Musca domestica*.

Hovea longifolia, R. Br.

1730. *Asterina hoveafolia*.—Leaves.

Hoya australis, R. Br.

2130. *Glebosporium intermedium*.

Hydrocotyle asiatica, L.

2206. *Protomyces macrosporus*.

Hydrocotyle hirta, R. Br.

1486. *Puccinia munita*.—Leaves.

Hymenochaete sp. (Fungus.)

986. *Merulius tenuissimus*.

Hypochaeris glabra, L. (Cat's Ear.)1478. *Puccinia Hieracii*.**Hypochaeris radicata**, L. (Flatweed.)1478. *Puccinia Hieracii*. V.—Leaves of flowering stems.
2263. *Puccinia Hypochaeris*. V.—Leaves.**Hypoxydon** sp. (Fungus.)1558. *Dialonectria tephrothela*.**Ilex Aquifolium**, L. (Holly.)1859. *Pseudohelotium ilicinolum*. V.—Branches.**Insects.**1528. *Cordyceps entomorrhiza*. V.—Larvæ.
1528A. *Cordyceps entomorrhiza*, var. *Menesteridis*. V.—Larvæ.
1529. *Cordyceps Gunnii*. V.—Larvæ.
1530. *Cordyceps Hawkesii*.—Larvæ.
1531. *Cordyceps ophioglossoides*.
1532. *Cordyceps Tylori*. V.—Larvæ.
1926. *Sporotrichum densum*.—Dead insects.
1994. *Isaria suffruticosa*.—Hairy caterpillar.**Ironbark-tree**—*Eucalyptus siderophloia*, Hook.**Isolepis** = *Scirpus*.**Jacksonia scoparia**, R. Br.1464. *Cronartium Asepiadenm*.—Leaves.
1516. *Ræstelia polita*. V.—Branches.**Jasminum racemosum**, F. v. M.778. *Fomes pnllus*.—Branches.**Juncus** sp. (Rush.)115. *Mycena juncicola*. V.—Dead rushes.
1479. *Puccinia Junciphila*. V.
1638. *Phyllachora Junci*. V.
2172. *Ustilago piluleformis*. V.—Ovaries.**Juncus maritimus**, Lam.1451. *Uromyces Junci*. V.**Juncus pallidus**, R. Br. (Sheathed Rush.)1518. *Uredo armillata*. V.**Juncus planifolius**, R. Br. (Flat-leaved Rush.)2171. *Ustilago Muelleriana*. V.—Seeds.**Kangaroo Grape-vine**—*Vitis antarctica*, Benth.**Kangaroo Grass**—*Anthistria ciliata*, L.**Kennedya** sp. = *Hardenbergia*.2029. *Phyllosticta Hardenbergiae*. V.—Leaves.**Kennedya monophylla**, Vent.2094. *Septoria Hardenbergiae*.—Leaves.**Kennedya prostrata**, R. Br. (Native Scarlet-runner.)1969. *Cercospora Kennedya*. V.—Leaves.**Kochia sedifolia**, F. v. M.1481. *Puccinia Kochiae*. V.—Leaves.**Lactuca** sp. (Lettuce.)1490. *Puccinia Prenanthis*. V.**Lagenaria vulgaris**, Sér. = *Cucurbita lagenaria*. (Bottle Gourd.)1672. *Eutypa polyscia*—Epicarp.**Lagenophora Billardieri**, Cass. (Bottle Thistle.)1482. *Puccinia Lagenophoræ*. V.—Leaves.
1737. *Dimerosporium Ludwigianum*. V.—Fading leaves.**Leersia hexandra**, Swartz. (Native Rice-grass.)2182. *Thecaphora inquinans*.—Inflorescence.**Leguminosæ.**1540. *Polystigma australiense*. V.—Leaves, rarely stems.1742. *Parodiella Perisporioides*. V.—Leaves.2191. *Schinzia Leguminosarum*. V.—Roots.**Lemon**—*Citrus Limonium*, Risso.**Lepidium ruderales**, L.2284A. *Peronospora parasitica*, var. *Lepidii*. V.—Leaves, stem, and fruit.**Lepidosperma** sp. (Sword-rush.)1699. *Anthostomella Lepidospermæ*.—V.1937. *Coniosporium pterospermum*. V.2095. *Septoria Lepidospermi*. V.—Leaves.**Lepiota bubalina**, Berk. (Fungus.)1918. *Aspergillus Muelleri*. V.**Leptocarpus tenax**, R. Br.2183. *Thecaphora Leptocarpi*. V.—Ovaries.**Leptospermum** sp. (Tea-tree.)152. *Plenrotas australis*.—Roots.2060. *Aposphaeria Leptospermi*. V.—Bark.**Leptospermum lævigatum**, F. v. M. (Sandstay.)2075. *Coniothyrium olivaceum*. V.—Involucres.2116. *Melophia Leptospermi*. V.—Leaves.**Leptospermum scoparium**, Forst.1997. *Harpographium Corynelioides*. V.—Branches.2144. *Hyaloceras dilophosporum*. V.—Leaves.**Lettuce**—*Lactuca* sp.**Lichens.**2003. *Hilosporium flavellum*.**Limnanthemum indicum**, Thwaites.1509. *Æcidium nymphoidis*.—Leaves.**Limosella** sp.1453. *Uromyces Limosellæ*.—Leaves.**Linum marginale**, Cuon. (Native Flax.)1461. *Melampsora Lini*. V.—Leaves.**Linum usitatissimum**, L. (Flax.)1461. *Melampsora Lini*. V.—Leaves.**Litsea** sp.1704. *Læstadia Litseæ*.—Leaves.**Litsea dealbata**, Nees. (Black Ash.)1643. *Dothidella apiculata*.—Fading leaves.**Lobelia anceps**, Thun.1469. *Puccinia aucta*. V.—Leaves.**Lobelia pedunculata**, R. Br.1469. *Puccinia aucta*. V.—Leaves.**Lobelia platycalyx**, F. v. M.1469. *Puccinia aucta*. V.—Leaves.**Locellinia cynopotamia**, Sacc. (Fungus.)1434. *Arachnium Drummondii*.—Attached to above fungus.

Lolium sp. (Rye-grass.)1992. *Isaria graminiperda*. V.**Lolium perenne**, L. (Rye-grass.)1493. *Puccinia rubigo-vera*. V.1527. *Claviceps purpurea*. V.—Inflorescence.1992. *Isaria graminiperda*. V.**Lolium temulentum**, L. (Darnel, Drake.)1527. *Claviceps purpurea*. V.—Inflorescence.**Loosestrife**—*Lythrum hyssopifolia*, L.**Lucerne**—*Medicago sativa*, L.**Lyonsia reticulata**, F. v. M.2274. *Peziza Lyonsiae*.—Leaves.**Lythrum hyssopifolia**, L. (Loosestrife.)2049. *Phoma Lythri*. V.—Fading leaves.2181. *Doassansia punctiformis*. V.—Leaves.**Macrozamia** = *Encephalartos*.**Maidenhair Fern**—*Adiantum* sp.**Maize**—*Zea Mays*, L.**Mallee**—*Eucalyptus incrassata*, Labill.**Malva rotundifolia**, L. (Dwarf Mallow.)1485. *Puccinia Malvacearum*. V.—Leaves and stems.**Mangifera indica**, L. (Mango.)2131. *Gloeosporium lagenarium*.—Fruit.2150. *Pestalozzia uvicola*.—Fruit.**Manna Gum-tree**—*Eucalyptus viminalis*, Labill.**Marsdenia** sp.2045. *Phoma folliculorum*.—Follicles.2081. *Diplodia Marsdeniae*.—Follicles.**Meadow Grass**—*Poa* sp.**Medicago sativa**, L. (Lucerne.)1703. *Læstadia destructiva*. V.—Leaves.1894. *Pseudopeziza Medicaginis*. V.—Leaves.**Melaleuca** sp. (Tea-tree.)1260. *Guepinia merulina*.—Rotten wood.1705. *Læstadia Melaleucæ*.—Leaves.**Melon**—*Cucumis Melo*, L.**Messmate**—*Stringybark Tree*—*Eucalyptus obliqua*, L'Hér.**Microtis porrifolia**, R. Br.1454. *Uromyces Microtidis*.—Leaves.**Mint Tree**—*Prostanthera lasiantha*, Labill.**Moluccas Bramble**—*Rubus Moluccanus*, L.**Moreton Bay Chestnut**—*Castanospermum australe*, Cunn.**Morus** sp. (Mulberry.)2102. *Phleospora Mori*. V.—Leaves.**Moss.**232. *Cantharellus lobatus*. V.571. *Pleurotus cyphella-formis*.1045. *Thelephora cristata*.—*Soppitticella cristata*.1140. *Corticium simulans*. V.1168. *Cyphella muscigena*. V.2215. *Lamproderma Listeri*.2241. *Didymium spumarioides*.2251. *Physarum rubbasis*.2256. *Leocarpus fragilis*.**Mountain Ash**—*Eucalyptus virgata*, Sieb.**Muehlenbeckia** sp.2087. *Darluca filum*.**Muehlenbeckia adpressa**, Meiss.1494A. *Puccinia runcicis-scutati*, var. *Muehlenbeckia*. V.—Leaves.**Muehlenbeckia Cunninghamii**, F. v. M.1516. *Ræstelia polita*. V.—Branches.**Mulberry**—*Morus* sp.**Musa** sp.1748. *Meliola Musæ*.**Musa Cavendishii**, Lamb. (Banana.)2131A. *Gloeosporium lagenarium*, var. *Cucurbitarum*.2134. *Gloeosporium Musarum*.—Fruit.**Musca domestica**, L. (House-fly.)2203. *Empusa Muscæ*. V.**Musk-Tree**—*Aster argophyllus*, Labill.**Myoporum insulare**, R. Br.2097. *Septoria Myoporii*. V.—Leaves.**Myoporum platycarpum**, R. Br.1649. *Bagnisiella endopyria*. V.—Leaves**Myriangium** sp. (Lichen.)1859. *Pseudohelotium ilicinolum*.—V.**Myrtus** sp. (Myrtle.)2083. *Ascochyta apiospora*.—Leaves.2147. *Pestalozzia funerea*.—Leaves.**Native Beech**—*Fagus Cunninghamii*, Hook.**Native Flax**—*Linum marginale*, Cunn.**Native Honeysuckle**—*Banksia* sp.**Native Pepper-tree**—*Drimys aromatica*, F. v. M.**Native Raspberry**—*Rubus parvifolius*, L.**Native Rice-grass**—*Leersia hexandra*, Swartz.**Native Sassafras**—*Atherosperma moschatum*, Labill.**Native Scarlet-runner**—*Kennedyia prostrata*, R. Br.**Native Supple-jack**—*Clematis aristata*, R. Br.**Native Yam**—*Dioscorea* sp.**Nepenthes** sp. (Pitcher plant.)1823. *Hunnaria Thozetii*.**Nerium Oleander**, L. (Oleander.)2098. *Septoria oleandrina*.—Leaves.**Nesodaphne** = *Beilschmiedia*.**Nettle**—*Urtica* sp.**Neurachne alopecuroides**, R. Br.2175. *Ustilago Tepperi*.**Nicotiana Tabacum**, L. (Tobacco.)2201. *Peronospora Hyoscyami*. V.—Leaves.**Norway-Spruce**—*Pinus picea*, Du Roi.**Oak**—*Quercus Robur*, L.**Oat-grass**—*Danthonia* sp.

- Oats**—*Avena sativa*, L.
- Olea paniculata**, R. Br.
1767. *Lembosia graphioides*.—Leaves.
- Oleander**—*Nerium Oleander*, L.
- Omalthus populifolius**, Grah.
2063. *Asteromella Omalthi*.—Leaves.
- Oncoptera intricata**, Walk. (Insect.)
2279. *Isaria Oncopterae*. V.—Dead Larvæ.
- Onion**—*Allium cepa*, L.
- Opercularia varia**, J. Hook.
1507. *Æcidium cystoseiroides*.
2267A. *Puccinia Coprosmatidis* var. *Opercularia*. V.—Leaves and leaf-stalks.
- Orange**—*Citrus Aurantium*, L.
- Oxalis corniculata**, L.
2276. *Oidium Oxalidis*. V.—Leaves, stem, and fruit.
- Palm**.
164. *Pleurotus Gardneri*.—Petioles and half-rotten fronds.
2031. *Phyllosticta palmicola*.—Leaves.
2053. *Phoma plagia*.—Leaves.
- Panicum** sp.
2166. *Ustilago Digitariae*. V.
- Panicum paradoxum**, R. Br.
2164. *Ustilago confusa*. V.
- Papaw**—*Carica Papaya*, L.
- Paspalum scrobiculatum** L. (Ditch Millet.)
2162. *Ustilago Cعاتii*. V.
2190. *Cerebella Paspali*.—Glumes.
- Passiflora** sp. (Passion-flower.)
266. *Marasmius rhyticeps*.—Twigs.
1561. *Lisiella Passifloræ*.—Stems.
1949. *Scoleotrichum atriellum*.—Twigs.
1998. *Harpographium quaternarium*.—Dead twigs.
2014. *Fusarium longisporum*.—Twigs.
2103. *Phlyctæna Passifloræ*.—Twigs.
- Pea**—*Pisum sativum*, L.
- Peach**—*Prunus Persica*, J. Hook.
- Pear**—*Pyrus communis*, L.
- Pelargonium australe**, Willd.
1474. *Puccinia Geranii*. V.—Leaves.
- Peppermint Gumtree**—*Eucalyptus amygdalina*, Labill.
- Phalaris minor**, Retz.
1475. *Puccinia graminis*. V.
- Phaseolus vulgaris**, L. (Bean.)
2133. *Glœosporium Lindemuthianum*. V.—Legume.
2260. *Uromyces Phaseoli*.—All parts, more especially leaves.
- Phoenix dactylifera**, L. (Date Palm.)
2188. *Graphiola Phœnicis*.
- Phyllis** sp.
402. *Pholiota phylligena*.—Trunks.
- Pinus** sp.
1157. *Coniophora olivacea*.—V.—Decayed wood.
1940. *Hormiscium pithyophilum*.—Branches.
2105. *Sphærœmella pinifolia*.—Pine chips.
- Pinus contorta**, Dougl.
1144. *Peniophora carnea*.
- Pinus picea**, Du Roi. (Norway Spruce.)
1110. *Hymenochaete Mougeotii*. V.—Trunks.
- Piperomia** sp.
1727. *Eurotium lateritium*.—Leaves.
- Pisum sativum**, L. (Pea.)
2133. *Glœosporium Lindemuthianum*. V.—Legume.
2141. *Marsonia deformans*. V.—Leaves, stipules, &c.
- Pitcher-plant**.—*Nepenthes* sp.
- Pittosporum rubiginosum**, Cunn.
1716. *Sphærella rubiginosa*.—Leaves.
- Plagianthus sidoides**, Hook. *
2264. *Puccinia Plagianthi*.—Leaves and flowers.
- Plane-tree**.—*Platanus* sp.
- Plantago** sp.
1510. *Æcidium Plantaginis*. V.—Leaves.
- Plantain**.—*Plantago* sp.
- Platanus** sp. (Plane-tree.)
2052. *Phoma notha*.—Dead branches.
- Platylobium** sp.
2033. *Phyllosticta Platylobii*. V.—Leaves.
- Plum**.—*Prunus domestica*, L.
- Poa** sp. (Meadow-grass.)
1489. *Puccinia Poarum*. V.
2047. *Phoma graminis*.—Stems.
- Poa annua**, L.—(Goose-grass.)
1489. *Puccinia Poarum*. V.
1493. *Puccinia rubigo-vera*. V.
2282. *Ustilago Poarum*. V.—Foliage especially.
- Polygonum** sp.
2167. *Ustilago emodensis*.—Stems, &c.
2180. *Sphacelotheca hydropiperis*.—Ovaries.
2180. *Sphacelotheca hydropiperis*, var. *Columellifera*.—Ovaries.
- Polygonum Hydropiper**, L. (Water-pepper.)
2176. *Ustilago utriculosa*. V.—Stems and ovaries.
- Polygonum minus**, Huds.
2176. *Ustilago utriculosa*. V.—Stems and ovaries.
- Polyporus**, sp. (Fungus.)
1551. *Hypomyces aurantius*.
1552. *Hypomyces chrysospermus*. V.
1553. *Hypomyces membranaceus*.
1554. *Hypomyces rosellus*.
1928. *Sepedonium aureo-fulvum*. V.
- Polyporus gryphæformis** = *Fomes gryphæformis*.
- Polyporus portentosus**, Berk. (Fungus.)
2054. *Phoma portentosa*. V.—Cap.
- Polystictus cinnabarinus**, Cooke = *Polyporus*. (Fungus.)
1939. *Torula mycetophila*. V.—Cap.
- Potato**.—*Solanum tuberosum*, L.
- Prostanthera lasiantha**, Labill (Mint-tree.)
2034. *Phyllosticta Prostantheræ*. V.—Leaves.
2076. *Coniothyrium septorioides*. V.—Leaves.

Proteaceæ.

2037. *Phyllosticta soriformis*. V.—Leaves.

Prunus Amygdalus, J. Hook. (Almond.)

1491. *Puccinia Pruni*. V.—Leaves.
1901. *Exoascus deformans*. V.—Leaves.
2065. *Dothiorella Amygdali*. V.—Bark.

Prunus Amygdalus, var. *amara*. (Bitter Almond.)

1935. *Trichothecium roseum*. V.—Fruit.

Prunus Armeniaca, L. (Apricot.)

1908. *Monilia fructigena*. V.—Fruit.
2025. *Phyllosticta circumscissa*. V.—Leaves and fruit.

Prunus Cerasus, L. (Cherry.)

2025. *Phyllosticta circumscissa*. V.—Leaves and fruit.

Prunus domestica, L. (Plum.)

1491. *Puccinia Pruni*. V.—Leaf.
1721. *Podosphaera tridactyla*. V.—Leaf.
1908. *Monilia fructigena*. V.—Fruit.

Prunus Persica, J. Hook. (Peach.)

1491. *Puccinia Pruni*. V.—Leaves and fruit.
1901. *Exoascus deformans*. V.—Leaves.
1908. *Monilia fructigena*. V.—Fruit.
1963. *Helminthosporium rhabdiferum*.—Ripe fruit.
2065. *Dothiorella Amygdali*. V.—Bark.
2278. *Cladosporium carpophilum*.—Fruit.

Pyrus communis, L. (Pear.)

1908. *Monilia fructigena*. V.—Fruit.
1947. *Fusicladium dendriticum*. V.—Leaves and fruit.
1948. *Fusicladium pyrinum*. V.—Leaves and fruit.
1957. *Cladosporium stenosporum*.—Leaves.
2127. *Glœosporium fructigenum*.—Fruit.

Pyrus Malus, L. (Apple.)

1721. *Podosphaera tridactyla*. V.—Young leaves and shoots.
1908. *Monilia fructigena*. V.—Fruit.
1935. *Trichothecium roseum*. V.—Fruit.
1947. *Fusicladium dendriticum*. V.—Leaves.
2138. *Glœosporium versicolor*. V.—Rotting fruit.

Queensland-Kauri.—*Agathis robusta*, *Masters*.**Quercus Robur**, L. (Oak.)

1952. *Cladosporium epiphyllum*. V.—Leaves.

Ranunculus parviflorus, L.

1511. *Æcidium Ranunculacearum*. V.—Leaves.

Ranunculus rivularis, Banks and Sol. (Water Crowfoot.)

1511. *Æcidium Ranunculacearum*. V.—Leaves.

Red Ash.—*Alphitonia excelsa*, Reiss.**Reeds**.—*Arundo* sp.**Rhagodia Billardieri**, R. Br.

1524. *Uredo Rhagodiae*. V.—Leaves.

Rhipogonum sp.

1866. *Dasyscypha glabrescens*. V.
1868. *Dasyscypha lanariceps*. V.

Rhipogonum parviflorum, R. Br.

1662. *Diatrype glomeraria*. V.—Branches.

Ricinus communis L. (Castor Oil plant.)

1172. *Cyphella villosa*. V.—Rotting stems
1694. *Physalospora gregaria*.—Stems.

Rosa sp. (Rose.)

1502. *Phragmidium subcorticium*. V.—Leaves.
1722. *Sphærotheca pannosa*. V.—Leaves.
1911. *Oidium leucoconium*. V.—Leaves.
2035. *Phyllosticta Rosæ*. V.—Leaves.
2056. *Phoma Rosarum*.—Twigs.
2072. *Sphæropsis Rosarum*.—Branches.
2086. *Actinonema Rosæ*. V.—Leaves.

Rubus fruticosus, L. (Bramble.)

2036. *Pyhllosticta Ruborum*. V.—Fading leaves.

Rubus Moluccanus, L. (Moluccas Bramble.)

1503. *Hamaspora longissima*.—Leaves.

Rubus parvifolius, L. (Native Raspberry.)

1500. *Phragmidium Barnardi*. V.—Leaves.

Rumex sp. (Dock.)

1465. *Puccinia Acetosæ*.—Leaves and stems.
1494. *Puccinia Rumicis-scutati*.—Leaves, leaf-stalks, and stems.

Rumex Brownii, Campd.

1483. *Puccinia Ludwigii*. V.—Leaves.

Rush (*Juncus* sp.).**Russula** sp. (Fungus.)

96. *Collybia tuberosa*.—Putrid specimens.

Rye-grass—*Lolium perenne*, L.**Saccharum officinarum**, L. (Sugar-cane.)

1256. *Daeryomyces Sacchari*.—Stems.
1452. *Uromyces Kuehnii*.—Leaves.
1696. *Physalospora Sacchari*.—Leaves.
1979. *Macrosporium graninum*.—Leaves.
2021. *Strumella Sacchari*.—Stalk and leaf.
2057. *Phoma Sacchari*.—Leaves and stems.

Salix sp. (Willow.)

182. *Pleurotus salignus*. V.
783. *Fomes salicinus*.—Trunks.

Salix Babylonica, Tourn. (Weeping willow.)

2069. *Cytospora xanthosperma*. V.—Branches.

Sandstay—*Leptospermum levigatum*, F. v. M.**Sarsaparilla**—*Smilax* sp.**Scævola** sp.

1508. *Æcidium Goodeniacearum*. V.—Leaves.

Schœnus imberbis, R. Br.

2187. *Uroystis solida*. V.

Scirpus nodosus, Rott. = *Isolepis*. (Club Rush.)

1492. *Puccinia rimosa*. V.
1635. *Phyllachora anceps*.—Stems.

Scirpus prolifer, Rott.

2169. *Ustilago marmorata*. V.—Leaves.

Sedge.—*Carex* sp.**Selliera** sp.

1457. *Uromyces puccinioides*. V.—Leaves and flower-stalks.
1508. *Æcidium Goodeniacearum*. V.—Leaves.

Senecio sp. (Groundsel.)

1512. *Æcidium Senecionis*. V.

Senecio Bedfordii, F. v. M.

2096. *Septoria Martinii*. V.—Leaves.

Senecio odoratus, Horn.1170. *Cyphella polycephala*.**Senecio velleioides**, Cunn.1506. *Æcidium Compositarum*. V.—Leaves.**Serjania**.1954. *Cladosporium hypophyllum*.—Leaves.**Sheathed Rush**—*Juncus pallidus*, R. Br.**Sheep's Burr**—*Acæna sanguisorbæ*, Vahl.**She Oak**—*Casuarina* sp.**Silky Oak**—*Grevillea* sp.**Smilax** sp. (*Sarsaparilla*.)1717. *Sphaerella smilacicola*.—Leaves.1753. *Zukalia loganiensis*.—Leaves.2113. *Actinothercium Scortechinii*.—Leaves.**Solanum Dallachyi**, Benth.1681. *Chaetomium cymatotrichum*.—Leaves.**Solanum Lycopersicum**, L. (Tomato.)1912. *Oidium Lycopersicum*. V.—Stem and leaves.1982. *Macrosporium Tomato*. V.—Ripe tomatoes.**Solanum tuberosum**, L. (Potato.)2018. *Epicoccum scabrum*.—Leaves and stems.**Solanum verbascifolium**, L.1970. *Cercospora Solanacea*.—Leaves.**Sorghum** sp.2087. *Darlucæ filum*.**Speedwell**—*Veronica* sp.**Sphæria** sp. (Fungus.)1169. *Cyphella parasitica*.**Spindle-tree**—*Enonymus* sp.**Spinifex hirsutus**, Labill.2174. *Ustilago Spinificæ*.—Flowers and spikes.**Sporobolus** sp.1533. *Epichloæ cinerea*.**Sporobolus indicus**, R. Br. (Tussock Grass.)1962. *Helminthosporium Ravenelii*.—Inflorescence.**Spotted Gum**—*Eucalyptus maculata*, Hook.**Spyridium**=*Cryptandra*.**Stipa** sp. (Spear Grass.)2163. *Ustilago combrens*.**Strawberry**—*Fragaria vesca*, L.**Styphelia** sp.1546. *Nectria ferruginea*. V.—Leaves, bracts, &c.**Styphelia straminea**, Spreng.=*Cyathodes*.1766. *Glonium tardum*.—Leaves.**Sugar Cane**—*Saccharum officinarum*, L.**Sugar Grass**—*Hemarthria compressa*, R. Br.**Sunflower**—*Helianthus annuus*, L.**Sweet-scented Acacia**—*Acacia suaveolens*, Willd.**Sword Rush**—*Lepidosperma* sp.**Tabernæmontana orientalis**, R. Br. (Bitter Bark.)1504. *Æcidium Apocyni*.—Leaves.**Tasmannia**=*Drimys*.**Tea-tree**—*Leptospermum* sp., and *Melaleuca* sp.**Tecoma jasminoides**, Lind.2112. *Melasmia Tecomatis*.—Leaves.**Tetracera Wuthiana**, F. v. M.1752. *Meliola Tetracera*.—Leaves.**Thatch-reed**—*Arundo Phragmites*, L.**Tinea**. (Insect.)1528. *Cordyceps entomorrhiza*. V.—Larva.**Tobacco**—*Nicotiana Tabacum*, L.**Tomato**—*Solanum Lycopersicum*, L.**Trema aspera**, Blume.1732. *Asterina pelliculosa*.—Leaves.1738. *Dimerosporium parvulum*.—Living leaves.2062. *Asteromella epitrema*.—Living leaves.**Tremella fuciformis**, Berk. (Fungus.)1684. *Rosellinia tremellicola*.**Tricoryne anceps**, R. Br.2073. *Sphaeropsis Tricorynes*.—Leaves.**Trifolium** sp. (Clover.)1459. *Uromyces Trifolii*.—V.1642. *Phyllachora Trifolii*. V.—Leaves.1895. *Pseudopeziza Trifolii*. V.—Languishing leaves.2241. *Didymium spumarioides*.**Tristania** sp.2055. *Phoma purpurea*.—Leaves.**Tristania conferta**, R. Br. (Brisbane Box.)1749. *Meliola octospora*.—Leaves.2084. *Ascochyta brunnea*.—Leaves.**Tristania laurina**, R. Br.1961. *Helminthosporium puccinioides*.—Fading or dead leaves.**Triticum vulgare**, Vill. (Wheat.)1475. *Puccinia graminis*. V.—Leaves, stem, and ear.1493. *Puccinia rubigo-vera*. V.—Leaves, stem, and ear.1527. *Claviceps purpurea*. V.—Inflorescence.1953. *Cladosporium herbarum*. V.—Leaves, stem, and ear.1979. *Macrosporium graminum*.—Leaves.2074. *Sphaeropsis Tritici*. V.—Dead leaves and sheaths.2100. *Septoria Tritici*. V.—Stem, leaves, and ear.2159. *Ustilago bullata*. V.—Ears.2173A. *Ustilago segetum*, var. *Tritici*. V.—Ear.2178. *Tilletia Tritici*. V.—Grains.2186. *Urocystis oenantha*. V.—Stem, leaves, glumes.**Tussock Grass**—*Sporobolus indicus*, R. Br.**Typha** sp. (Bulrush).1958. *Cladosporium Typharum*. V.—Leaves.**Urtica** sp. (Nettle).1514. *Æcidium Urticæ*. V.**Veronica** sp. (Speedwell).1515. *Æcidium Veronicæ*.—V.**Vine**—*Vitis vinifera*, Bauh.**Viola** sp.1498. *Puccinia Violæ*.—V.2101. *Septoria Violæ*. V.—Fading leaves.

Viola hederacea, Labill.1466. *Puccinia agra*. V.—Leaves.**Vitis** sp.1824. *Phillipsia polyporoides*.—Dead stems.**Vitis antarctica**, Benth. (Kangaroo Grape-vine).2030. *Phyllosticta neurospile*a.—Leaves.**Vitis vinifera**, Bauh. (Grape Vine.)1163. *Cyphella albo-violascens*. V.—Branches.1698. *Didymella cladophila*.—Branches.1725. *Erysiphe vitigera*. V.—Leaves.1914. *Oidium Tuckeri*. V.—Leaves and grapes.1951A. *Cladosporium Asteroma*, var. *minor*.—Foliage.1971. *Cercospora viticola*.—Leaves.1983. *Fumago vagans*.—Living leaves.2000. *Isariopsis clavisporea*.—Leaves.2039. *Phoma ampelina*. V.—Twigs.2058. *Phoma uvicola*. V.—Grapes.2078. *Chatomella brachyspora*. V.—Bark and branches.2085. *Robillarda sessilis*.—Fading leaves.2122. *Gleospodium ampelophagum*. V.—Grapes, rarely leaves or branches.2136. *Gleospodium pestiferum*. V.—Twigs, flower-stalks, and grapes.2150. *Pestalozzia uvicola*.—Leaves and grapes.2200. *Plasmopara viticola*.—Leaves.**Water Crowfoot**—*Ranunculus rivularis*, Banks and Sol.**Water-pepper**—*Polygonum Hydropiper*, L.**Weeping Willow**—*Salix Babylonica*, Tournef.**Wheat**—*Triticum vulgare*, Vill.**White Gumtree**—*Eucalyptus pauciflora*, Sieb.**Wild Oats**—*Avena fatua*, L.**Willow**—*Salix* sp.**Wurmbsea dioica**, F. v. M.1499. *Puccinia Wurmbseae*.—Leaves.1526. *Uredo Wurmbseae*.—Leaves.(This is probably the *Uredo*-stage of the same fungus.)**Xanthorrhoea** sp. (Grass tree.)1002. *Hydnum investiens*.—In cavities of trunk.**Xanthoxylum** sp.1758. *Capnodium salicinum*.**Zea Mays**, L. (Maize.)1496. *Puccinia Sorghi*.1931. *Verticillium lateritium*. V.1959. *Helminthosporium inconspicuum*.—Fading leaves.2170. *Ustilago Maydis*.2177. *Tilletia epiphylla*.—Leaves.2247. *Physarum didermoides*.—Bracts.**Zygophyllum ammophilum**, F. v. M. (Bean Caper.)1460. *Uromyces vesiculosa*.—Leaves and stems.

LIST OF WORKS ON AUSTRALIAN FUNGI.

III.—LIST OF WORKS ON AUSTRALIAN FUNGI.

1. ABBOTT (F.)—"Smut in Wheat." Proc. Roy. Soc., Tasm., 1889.
The nature of smut stated, and means for prevention given.
2. AGRICULTURE DEPARTMENT.—New South Wales, Queensland, South Australia, Tasmania, and Victoria.
Reports and Bulletins issued up to date contain various papers on parasitic fungi, the more important of which are noted under authors' names.
3. Agricultural Gazette of New South Wales. Department of Agriculture, Vols. I.—V.—(continued.) Sydney, 1890-94.
Contains numerous articles on fungus diseases, the more important of which are given under authors' names.
4. ANDERSON (H. C. L.)—"Rust in Wheat: Experiments, and their Objects." Ag. Gaz., N.S.W., I, Pt. I., 1890.
A variety of measures suggested for trial in order to minimize the effects of rust.
5. BACKHOUSE (J.)—"A Narrative of a Visit to the Australian Colonies." 8vo. London, 1843.
Reference at p. 119 to Punk and an edible fungus near Emu Bay in Tasmania; also in Appendix D, p. xl, to the common Mushroom and *Mylitta australis*.
6. BAILEY (F. M.)—"A General Account of the Flora of Tropical Queensland." Proc. Linu. Soc., N.S.W., II, 1878.
Some of the more important fungi referred to.
7. ————"Medicinal Plants of Queensland." Ibid. V., 1880.
Hirneola auricula-jude, or Jew's Ear, referred to.
8. ————"A Synopsis of the Queensland Flora." 8vo. Brisbane, 1883.
9. ————"A Classified Index of the Indigenous and Naturalized Plants of Queensland." 8vo. Brisbane, 1883.
10. ————"Contributions to the Queensland Flora." Proc. Roy. Soc., Q., I., Pt. I., 1884.
Eighty-two species of fungi recorded.
11. ————"Contributions to the Queensland Flora." Pt. II., ibid., 1884.
Eleven species of fungi recorded.
12. ————"Contributions to the Queensland Flora." Part III., ibid., 1884.
Seven species of fungi recorded.
13. ————"A Synopsis of the Queensland Flora." First Supplement. 8vo. Brisbane, 1886.
14. ————"Classified Index of the First Supplement to the Indigenous and Naturalized Plants of Queensland." 8vo. Brisbane, 1886.
15. ————"A Synopsis of the Queensland Flora." Second Supplement. 8vo. Brisbane, 1888.
16. BAILEY (F. M.)—"Classified Index of the Second Supplement to the Indigenous and Naturalized Plants of Queensland." 8vo. Brisbane, 1888.
17. ————"Supplement to the Report of the Botany of the Bellenden-Ker Expedition—Fungi collected or observed about the Bellenden-Ker Range." Ann. Rep. Dept. Ag., Q., 1890.
Fifty-seven species of fungi are recorded, fourteen of which are new to Australia.
18. ————"A Synopsis of the Queensland Flora." Third Supplement. 8vo. Brisbane, 1890.
Includes first addendum to third supplement.
19. ————"Catalogue of the Plants of Queensland." 8vo. Brisbane, 1890.
Includes second addendum to third supplement.
20. ————"Contributions to the Queensland Flora." Bull. 4, or Bot. Bulletin I. Dept. Ag., Q., 1890.
Strumella sacchari (Cooke) and *Peronospora hyoscyami* (De Bary) recorded.
21. ————"Final Supplement to the Report of the Botany of the Bellenden-Ker Expedition." Ann. Rep. Dept. Ag., Q., 1891.
Aschersonia tahitensis (Mont.), *Entyloma Eugenia-rum* (Cooke and Mass.), *Asterina reptans*, (Berk. and Cooke) added.
22. ————"Additional Fungus Blights observed to have injured Plants during the Year." Ibid., 1891.
Five species are mentioned altogether, occurring on Vines, Hollyhocks, Tobacco plant, and native plants.
23. ————"Contributions to the Queensland Flora." Bull. 7, or Bot. Bull. II. Dept. Ag., Q., 1891.
24. ————"Contributions to the Queensland Flora." Bull. 9, or Bot. Bull. III. Dept. Ag., Q., 1891.
Gleospodium pestiferum (Cooke and Mass.) briefly defined.
25. ————"Contributions to the Queensland Flora." Bull. 13, or Bot. Bull. IV. Dept. Ag., Q., 1891.
Fourteen species of fungi described.
26. ————"Additional Fungus Blights." Report of Colonial Botanist in Ann. Rep. Dept. Ag., Q., 1892.
Twelve species of fungi are recorded.
27. ————"Contributions to the Queensland Flora." Bull. 18, or Bot. Bull. V. Dept. Ag., Q., 1892.
Ten species of fungi are recorded, and most of them described.
28. ————"A Review of the Fungus Blights which have been observed to injure Living Vegetation in the Colony of Queensland." Report Aust. Assoc. Adv. Sci., Hobart, IV., 388, 1892.
Blights are classed under epiphytes and parasites, twenty-five species under the former and one hundred and fifteen under the latter, with hosts.

29. BAILEY (F. M.)—"Contributions to the Queensland Flora." Botany Bull. VIII. Dept. Ag., Q., 1893.
Forty-five species of fungi are recorded.
30. ——— "Additional Fungus Blights." Report of Colonial Botanist in Ann. Rep. Dept. Ag., Q., 1893.
Six fungi are noted and two new.
31. ——— "Companion for the Queensland Student of Plant Life."
32. ——— "Botany abridged." Dept. Ag., Q., 1894.
Edible fungi in Queensland recorded.
33. ——— "Contributions to the Queensland Flora." Botany Bull. IX. Dept. Ag., Q., 1894.
Thirty-six species described and two recorded without description.
34. BAILEY (F. M.) and GORDON (P. R.)—"Plants reputed Poisonous and Injurious to Stock." 8vo. Brisbane, 1887.
A few fungi are added, injurious to fodder plants.
Nine altogether, with an illustration.
- BAILEY (F. M.)—[See "Tension-Woods (J. E.)"]
35. BANCROFT (J.)—"Experiments with Indian Wheats in Queensland." Proc. Roy. Soc., Queensland, I., Pt. 4, 1884.
Indian Wheats of the tall dark-bearded kinds found to be rust-resisting.
36. BANCROFT (T. L.)—"Notes on Bacterial Diseases of the Roots of Leguminosæ." Proc. Linn. Soc., N.S.W., Vol. VIII., Pt. I., 1893.
Five leguminous plants affected—Mimosa, Sesbania, Desmodium, Medicago, and Crotalaria.
37. BARWICK (J.)—"Smut in Wheat." Proc. Roy. Soc., Tasm., 1889.
Considers that it is grain damaged in threshing which is smutty, and self-sown grain is never smutty.
38. BELL (R.)—"Some Account of Red Rust and its Remedy." Pp. 10. Ballarat, 1893.
The remedy given is to apply a solution of common salt to the growing wheat plant—1 lb. of salt to 1 gallon of water.
39. BENNETT (G.)—"Gatherings of a Naturalist in Australasia." 8vo. Lond., 1860.
Reference to a luminous agaric.
40. BENSON (A. H.)—"Principal Insect and Fungus Pests in New South Wales, and their remedies." Ag. Gaz., N.S.W., III., Pt. 8, 1892.
Notices injurious fungi, with their remedies, on Citrus, Apple, Pear, Apricot, Plum, and Peach trees, and Vines.
41. ——— "Apple Culture." Ibid. V., Pt. 6., 1894.
Fungus diseases of Apple described and illustrated.
42. BERKELEY (M. J.)—"Contributions towards the Flora of Van Diemen's Land." Fungi. Ann. Nat. Hist. III., 1839.
Twenty-seven species given, twelve of which are common European fungi.
43. ——— "Description of Two New Fungi in the Collection of Sir W. J. Hooker." Hook., Journ. Bot. II., Pl. 1, 1840.
Lentinus fasciatus is described from Tasmania.
44. BERKELEY (M. J.)—"On some Entomogenous Sphæriæ." Hook., Lond. Journ. Bot. II., Pl. 1, 1843.
Seven described altogether, and one (*Sphæria Taylora*) described and figured from N.S.W.
45. ——— "Decades of Fungi." Decade I. Ibid. III., Pl. 2, 1844.
Three new species are described from Australia—*Agaricus nidiformis*, *Polyporus portentosus*, and *Asera rubra*.
46. ——— "Decades of Fungi." Decades III.–VII. Ibid. IV., Pl. 2, 1845.
Forty-nine new species described, and some figured.
47. ——— "Decades of Fungi." Decades VIII.–X. Ibid. IV., Pl. 2, 1845.
Three new species described—*Sphæria elevata*, *S. pulvinulus*, and *S. inspersa*.
48. ——— "Decades of Fungi." Decade XI. Ibid. V., 1846.
Four new species described for Australia—*Marasmius hepaticus*, *Hexagonia similis*, *Polyporus brunneo-leucus*, and *Peziza fusispora*.
49. ——— "On Cordyceps Gunnii." Hook., Lond. Journ. Bot. VII., 577, Pl. 22, 1848.
First described and figured.
50. ——— "On some Entomogenous Sphæriæ." Linn. Journ. I., Pl. 1, 1856.
Entomogenous species of *Cordyceps* mentioned for Australia—*C. Gunnii* and *C. Taylora*.
51. ——— "Introduction to Cryptogamic Botany." 8vo. London, 1857.
Refers to various Australian fungi, and gives drawings of some such as *Cyttaria Gunnii*.
52. ——— "Flora of Tasmania." Fungi. Hooker's Botany of the Antarctic Voyage. 4to. Pt. III., Vol. II., Pl. 4. London, 1860.
Two hundred and seventy-five species are described, only about eight of which are peculiarly Australian.
53. ——— "Outlines of British Fungology." 8vo. Lond., 1860.
Tasmanian fungi referred to at pp. 34 and 35.
54. ——— "On a Collection of Fungi from Cuba." Journ. Linn. Soc. X., 1868.
Habitats given for species occurring also in Australia.
55. ——— "Australian Fungi, received principally from Baron F. von Mueller and Dr. R. Schomburgk." Ibid. XIII., 1873.
Fungi characterized, and where new described; received during a period of nearly twenty years.
56. ——— "Enumeration of the Fungi collected during the Expedition of H.M.S. *Challenger*." 1874–5. (Third notice.) Ibid. XVI., 1877.
Forty-nine species are recorded altogether from the neighbourhood of Sydney, N.S.W., and new species are described.
57. ——— "Gardener's Chronicle." 791, Fig. 130. 1878.
Description of *Cordyceps Menesteridis*—the same as, or a variety of, *C. entomorrhiza*.
58. ——— "Australian Fungi." Part II. Received principally from Baron F. von Mueller. Ibid. XVIII., 1880.
A number of new species described.

59. BERKELEY (M. J.) and BROOME (C. E.)—"On some species of the genus *Agaricus* from Ceylon." Trans. Linn. Soc. XXVII., 149-152, Pl. 33-34, 1868.
60. ——— "The Fungi of Ceylon." Linn. Journ. Bot., Vols. XI., XIV., and XV., 1870-75.
Contain descriptions of fungi common to Ceylon and Australia, between which countries remarkable coincidences occur in the distribution of some species.
61. ——— "List of Fungi from Brisbane, Queensland, with Descriptions of New Species." Ibid. I., 2nd Ser., Bot. I., Pl. 2, 1878.
About one hundred and twenty species recorded.
62. ——— Ibid. II., 2nd Ser., Pl. 6, 1882.
Fifty-three species recorded, for the most part common European ones.
63. ——— "List of Fungi from Queensland and other Parts of Australia, with Descriptions of New Species." Ibid. II., 2nd Ser., Pt. 3, Pl. 1, 1886.
Supplementary to previous lists.
64. BERKELEY (M. J.) and CURTIS (M. A.)—"Fungi Cubenses (Hymenomycetes)." Journ. Linn. Soc. X., 1867.
Habitats given for species occurring also in Australia.
65. BICHENO (J. E.)—"On the Potato as an Article of National Diet, and the Potato Disease in connexion with Distress in Ireland." Proc. Roy. Soc., Van Diemen's Land, I., Pt. III., 1851.
Sceptical as to the disease originating with the Aphis; that more likely the insect is the effect than the cause.
66. BLIGHT IN WHEAT.—Dept. Ag., Vict., Bull. No. 1. June, 1888.
Disease similar to "Take-all."
67. BRESADOLA (J.) and SACCARDO (P. A.)—"Pugillus mycetum Australiensium." Malpighia, Genoa, 1890.
Eighty-three species are recorded, with plate, five of which are new.
68. BROWN (R.)—"Miscellaneous Botanical Works (Ray Society): General Remarks—Geographical and Systematical—on the Botany of Terra Australis." I., 1866. (Reprinted from the voyage to Terra Australis, by M. Flinders, London, 1814.)
Ten species of fungi are noted.
69. CAMPBELL (F. M.) [now Mrs. Martin].—"Victorian Fungi hitherto unrecorded." Vict. Nat. II., No. 11, 127, 1886.
Twenty-one species are given.
70. ——— "Thirty Species of Fungi hitherto unrecorded for Victoria." Vict. Nat. IV., No. 6, 95, 1887.
71. ——— "Vegetable Pathology." Vict. Nat. IV., No. 8, 124, 1887.
Reference is made to the importance of the subject, and the great damage done by fungus pests to our forest trees and cultivated plants.
72. ——— "Fungus Pests." Vict. Roy. Com. Veg. Prod., 5th Progress Report, 1888.
Various fungus diseases pointed out, and a large and interesting collection exhibited.
73. CLARSON (W.)—"Blights and their Teachings." Bull. No. 5, Dept. Ag., Vict., Sept., 1889.
General reference to various fungus pests of the orchard.
74. CLARSON (W.)—"The Fruit Garden." Pts. I. and II. Melb., 1889.
Reference is made to fungus diseases under the different fruits.
75. COBB (N. A.)—"Peach Rust in our Orchards (*Uromyces amygdali*)." Ag. Gaz., N.S.W., I., Pt. 1, 1890.
Description, with figure, and treatment recommended.
76. ——— "Report on Pumpkin Mould." Ibid. 1890.
Description and drawing of mould belonging to the *Erysipheæ*.
77. ——— "Contributions to an Economic Knowledge of the Australian Rusts (Uredineæ)." Ag. Gaz., N.S.W., I., Pt. 3, 1890.
Methods of investigation given, and rust occurring around wheat paddocks recorded.
78. ——— "Notes on Diseases of Plants." Ibid. II., Pt. 1, 1891.
Anthracnose on Vines, *Fusicladium pyrinum*, *Sphaerella fragariae*, and *Puccinia malvacearum* noted.
79. ——— "Pathological Notes." Ibid. II., Pt. 2, 1891.
Sphaerella destructiva (B. and Br.) on Lucerne, and red incrustation on fence-rails noted.
80. ——— "Notes on Diseases of Plants." Ibid. II., Pt. 3, 1891.
Bitter Rot of Apple (*Glucosporium versicolor*), Flax Rust (*Maclospora lini*), and Peach Rust (*Puccinia pruni*) noted.
81. ——— "Pathological Notes." Ibid. II., Pt. 4, 1891.
Maize Rust (*Puccinia maydis*) and Apple Scab (*Fusicladium dendriticum*) noted.
82. ——— "Notes on Diseases of Plants." Ibid. II., Pt. 5, 1881.
Cystopus candidus or White Rust, *Ustilago maydis* or Maize Smut, *Puccinia maydis* or Maize Rust, *Sphaerella destructiva* on Lucerne, and Water Core in Apples noted.
83. ——— "Notes on Diseases of Plants." Ibid. II., Pt. 6, 1891.
Mouldy Core in Apples and *Glucosporium pestiferum* of the Vine noted.
84. ——— "Notes on Diseases of Plants." Ibid. II., Pt. 8, 1891.
Apple Scab (*Fusicladium dendriticum*) and Strawberry Leaf Blight (*Sphaerella fragariae*) again noted.
85. ——— "Notes on Diseases of Plants." Ibid. II., Pt. 10, 1891.
Onion Mildew (*Peronospora Schleideniana*), Tobacco Mildew (*Peronospora hyoscyami*), Banana Disease, and Bread Mould on Oranges noted. Potato Blight described, but not found in Australia.
86. ——— "Smut." Ibid. II., Pt. 11, 1891.
Oat Smut (*Ustilago avenae*), Wheat Smut (*Ustilago tritici* and *Urocystis occulta*), Maize Smut (*Ustilago maydis*), and Stinking Smut of Wheat (*Tilletia foetens*) described.
87. ——— "Dialogue concerning the manner in which a Poisonous Spray does its work in Preventing or Checking Blight." Ibid. II., Pt. 12, 1891.
With various drawings showing how a fine spray acts upon the spores of a fungus.

88. COBB (N. A.)—"Contributions to an Economic Knowledge of the Australian Rusts (Uredineæ)." Ibid. III., Pt. 1, 1892.

What has been found out in this and other countries concerning Wheat Rust together with the examination of a number of varieties of wheat, and the kind of rust determined.

89. ————"Contributions to an Economic Knowledge of the Australian Rusts (Uredineæ)." Ibid. III., Pt. 3, 1892.

Stiff flag, tough cuticle, and glaucousness found to be characteristic of rust-resistant Wheats. Various illustrative drawings and tables, among which the rust-devouring *Diplosis* is given.

90. ————"Economic Notes on Plant Diseases." Ibid. III., Pt. 4, 1892.

Apple Scab (*Fusicladium dendriticum*), Powdery Mildew of Apple (*Podosphora Kunzei*), Bitter Rot (*Glaeosporium versicolor*), Mouldy Core and Water Core in Apples. Pear Scab (*Fusicladium pyrinum*), Shot Hole (*Phyllosticta circumscissa*), Anthracnose (*Glaeosporium ampelinum*), Tufted-leaf Blight (*Cercospora viticola*), Strawberry-leaf Blight (*Sphaerella fragariae*), White Rust (*Cystopus candidus*), Pumpkin-leaf Oidium (*Oidium erysiphoides*), and Powdery Mildew of Rose (*Sphaerotheca pannosa*) illustrated and described.

91. ————"Plant Diseases and how to Prevent them." Ibid. III., Pt. 6, 1892.

Pourridie or Mouldy Root of the Vine, Tufted-leaf Blight of Bean, and Apple Canker described, and remedies prescribed.

92. ————"Plant Diseases and how to Prevent them." Ibid. III., Pt. 12, 1892.

"Take-all" and Dry Blight of Wheat and Oats and Leaf Curl of Peach described and illustrated. Also two new species of fungi—*Cyathus dimorphus* and *Tolyposporium anthistiria*.

93. ————"Contributions to an Economic Knowledge of Australian Rusts (Uredineæ)." Ibid. IV., Pt. 6, 1893.

Seventy-one varieties of Wheat described and illustrated.

94. ————"Contributions to an Economic Knowledge of Australian Rusts (Uredineæ)." Ibid. IV., Pt. 7, 1893.

Artificial crossing of Wheat and improving Wheats by selection fully described and illustrated.

95. ————"Plant Diseases and their Remedies—Diseases of the Sugar Cane." Ibid. IV., Pt. 10, 1893.

The gumming of Sugar Cane is due to *Bacillus vascularum*, and the following five species of fungi are described and illustrated:—*Uromyces Kuhnii*, *Strumella sacchari*, *Macrosporium graminum*, *Phoma sacchari*, and an undetermined species causing "Red Rot."

96. ————"Host and Habitat Index of the Australian Fungi." Ag. Dept., N.S.W. (Miscellaneous Publication, No. 16), p. 44, 1893.

Contains all the fungi recorded in Dr. Cooke's "Handbook of Australian Fungi."

97. ————"Contributions to an Economic Knowledge of Australian Rusts (Uredineæ)." Ibid. V., Pt. 4, 1894.

Improving Wheat by selection.

98. ————"Notes on Diseases of Plants." Ibid. V., Pt. 6, 1894.

Bean Anthracnose (*Colletotrichum Lindemuthianum*), Bean Rust (*Uromyces Phaseoli*), Peach Freckle (*Cladosporium carpophilum*), Black Rot of Tomato (*Macrosporium Tomato*), a Mango Blight (*Pestalozzia uricola*, Speg.), Disease of Grass (*Helminthosporium Ravenelii*) illustrated and described.

99. COBB (N. A.)—"A New Australian Fungus." Ibid. V., Pt. 6, 1894.

Peziza Lyonsiae described and illustrated.

100. COBB (N. A.) and OLLIFF (A. S.)—"Insect Larva (*Cecidomyia*, sp.) Eating Rust on Wheat and Flax." Ag. Gaz., N.S.W., II., Pt. 2, 1891. Also Ann. Nat. Hist. VII., 6th Ser. 1891.

Larva observed under the microscope unmistakably feeding on rust spores. The larva not only devours the spores but spreads the rust.

101. COOKE (M. C.)—"The Beech Morels of the Southern Hemisphere." Pharm. Journ. (3), I., 264, 1870.

Cyttaria Gunnii (B.), found on living branches of *Fagus Cunninghamii* and *F. Gunnii* in Tasmania, is figured and described.

102. ————"Jew's Ear (*Hirneola auricula-judae*)." Ibid. 681, 1871.

Figured and popularly described.

103. ————"Fungi: Their Nature, Influence, and Uses." 8vo. London, 1875.

Various Australian species referred to.

104. ————"Australian Fungi." Grev. VI. 70, 1877.

Twelve species collected in neighbourhood of Melbourne by Mr. Le Fevre, one of which was undescribed, viz., *Trametes serobiculata* (Berk.).

105. ————"New Zealand Fungi." Grev. VIII. 54, 1879.

Thirty-six Australian species recorded.

106. ————"Australian Fungi." Grev. IX. 142, 1881.

107. ————"Australian Fungi." Grev. X. 60, 93, and 131, 1882.

108. ————"Australian Fungi." Grev. XI. 28 and 57, 1882.

109. ————"On Xylaria and its Allies." Grev. XI. 81, 1883.

Three new Australian species described.

110. ————"Australian Fungi." Grev. XI. 97 and 145, 1883.

111. ————"Hypoxylon and its Allies." Grev. XI. 121, 1883.

Three Australian species described.

112. ————"Fungi." Trans. Roy. Soc., S.A., Vol. XVI., Pt. 2, 1883.

Eleven species recorded and two described.

113. ————"Fungi Australiani." 8vo., Pl. 4, pp. 72, London and Melbourne, 1883.

Hymenomyces, 783 species; *Gastromyces*, 111; *Myxomyces*, 33; *Ecidomyces*, 47; *Discomyces*, 84; *Pyrenomyces*, 94; *Hypomyces*, 47; *Phycomyces* 4; total 1,203 species.

114. ————"Some Exotic Fungi." Grev. XII. 85, 1884.

Meliola densa (Cooke) described.

115. ————"Some Exotic Fungi—Australasia." Grev. XIV. 11, 1885.

Five species described.

116. ————"Some Exotic Fungi." Grev. XIV. 89, 1886.

Phyllosticta palmicola (Cooke) described.

117. ————"Exotic Fungi." Grev. XV. 16, 1886.

Six species described.

118. ————"Some Australian Fungi." Grev. XV. 93, 1887.

Fourteen species described.

119. COOKE (M. C.)—"Some Australian Fungi." Grev. XVI. 97, 1887.
Nineteen species described.
120. ——— "New Australian Fungi." Grev. XVI. 1, 1887.
Thirty-five species described.
121. ——— "Two Remarkable Fungi." Grev. XVI. 20, 1887.
Cerebella paspali and *Hemiarcyria applanata*.
122. ——— "Australasian Fungi." Grev. XVI. 30, 1887.
Nineteen species described.
123. ——— "Australian Fungi." Grev. XVI. 72, 1888.
Twenty-nine species described or referred to.
124. ——— "Australasian Fungi." Grev. XVI. 113, 1888.
Eight species described.
125. ——— "Australasian Fungi." Grev. XVII. 7, 1888.
Seven species described.
126. ——— "Australian Fungi." Grev. XVII. 55, 1889.
Seven species described.
127. ——— "Some Brisbane Fungi." Grev. XVII. 69, 1889.
Four species described.
128. ——— "New Australian Fungi." Grev. XVIII. 1, 1889.
Forty-two species described.
129. ——— "New Australian Fungi." Grev. XVIII. 25, 1889.
Six species described, and one new genus—*Seismosarca*.
130. ——— "Australian Fungi." Grev. XVIII. 1, 1890.
Thirty-six species mostly described.
131. ——— "Australian Fungi." Grev. XVIII. 49, 1890.
Two species described—*Sphaeropsis phomatoidea* and *Capnodiastrum orbiculatum*.
132. ——— "Australian Fungi." Grev. XVIII. 80, 1890.
Three species described.
133. ——— "On Campbellia—New Genus." Grev. XVIII. 87, 1890.
134. ——— "Australian Fungi." Grev. XIX. 5, 1890.
Five species described.
135. ——— "Australian Fungi." Grev. XIX. 44, 1890.
Fifteen species and new genus (*Chainoderma*) described.
136. ——— "Australian Fungi." Grev. XIX. 89, 1891.
Twenty-one species described.
137. ——— "Additions to Dædalea." Grev. XIX. 93, 1891.
Dædalea Muellerei described.
138. ——— "Trametes and its Allies." Grev. XIX. 98, 1891.
Sclerodopsis, *Trametes*, and *Hexagonia* noticed.
139. ——— "New Sub-genus of Agaricus." Grev. XIX. 104, 1891.
Metrararia insignis described.
140. ——— "Additions to Merulius." Grev. XIX. 108, 1891.
Merulius pelliculosus described.
141. COOKE (M. C.)—"Australian Fungi." Grev. XX. 4, 1891.
Seventeen species described.
142. ——— "Species of Cyphella." Grev. XX. 9, 1891.
Cyphella australiensis described.
143. ——— "Notes on Clavariel." Grev. XX. 19, 1891.
Clavaria Muellerei (Berk.) and *C. tasmanica* (Berk.) are described.
144. ——— "Notes on Thelephoræ." Grev. XX. 11, 1891.
Species of *Hymenochaete* and *Corticium* referred to.
145. ——— "Apple Scab (*Fusicladium dendriticum*)." Grev. XX. 27, 1891.
Noticing profusion of examples from Australia and methods of checking quoted.
146. ——— "Ceylon in Australia." Grev. XX. 29, 1891.
Ceylon species of fungi found also in Australia.
147. ——— "Australian Fungi." Grev. XX. 65, 1892.
Four species described.
148. ——— "A Mystery Solved." *Gardener's Chronicle*, 20th Oct., 1892.
Sclerotium of *Mylitta* with *Polyporus ovinus*.
149. ——— "Vegetable Wasps and Plant Worms—A Popular History of Entomogenous Fungi or Fungi Parasitic upon Insects." S.P.C.K., Lond., 1892.
Seven Australian forms are noted and described.
150. ——— "Handbook of Australian Fungi." 8vo., pp. 458. Pl. 36. London, 1892.
Gives a technical description of over two thousand species, and is intended to include all Australian fungi known to date.
151. ——— "Australian Fungi—A Supplement to Handbook." Grev. XXI. 35, 1892.
Twenty-four species recorded as additions and corrections to the Handbook.
152. ——— "Fungi." Trans. Roy. Soc., S.A., XVI, Pt. II., 1893.
Eleven species recorded, and two described—*Stephensia arenivaga* (Cooke and Mass.) and *Diploderma Sabulosum* (Cooke and Mass.).
153. ——— "Exotic Fungi." Grev. XXI. 75, 1893.
A new species (*Diplodia Marsdeniæ*, C. and M.) described.
154. ——— "Australian Fungi." Grev. XXII. 36, 1893.
Nine new species described, and seven old ones recorded.
155. ——— "Australian Fungi." Grev. XXII. 68, 1893.
Two new species described—*Hypoxyylon atrophoricum* (Cooke and Mass.) and *Belonidium parasiticum* (Cooke and Mass.).
COOKE (M. C.)—[See "Kaltbrenner (K.)"]
156. COOKE (M. C.) and MASSEE (G.)—"Glæosporium pestiferum." *Gardener's Chronicle*, London, 1891.
157. CORDA (A. C. J.)—"Icones Fungorum hucusque cognitorum." 6 vols. Folio. 1837-54.
158. COUESLAND (F.)—"Disease of the Vine (or Oidium Tuckeri), and its Remedy." Pamphlet. Melb., 1876.
159. CRAWFORD (F. S.)—"The Apricot Disease." Proc. Roy. Soc., S.A., 1884.
Phyllosticta circumscissa and *Helminthosporium rhabdiferum*.

160. CRAWFORD (F. S.)—"Report on the Fusieladiums, the Codlin Moth, and certain other Fungus and Insect Pests attacking Apple and Pear Trees in South Australia." Pp. 70. Pl. 7. Adelaide, 1886.
Fusielodium dendriticum and *F. pyrinum* fully described and illustrated, and their history given, together with several remedies suggested.
161. ——"Insect and Fungus Pests." Viet. Roy. Com. Veg. Prod., 4th Progress Report and App. No. 5, 1887.
 Refers to Fusieladiums and other fungus pests, and suggests a central department in Victoria for Vegetable Pathology.
162. ——"Insect and Fungus Pests." Proc. of First Congress of Ag. Bureau of S.A. Pl. 6. Adelaide, 1890.
 Describes and illustrates rust in Wheat.
163. CRICHTON (D. A.)—"The Australasian Fruit Culturist." 8vo. Melbourne, 1893-4.
 Under the heading of the various fruits gives the principal fungus diseases.
164. DESPEISSIS (J. A.)—"Anthracosis, or Black Spot of the Grape." Ag. Gaz., N.S.W., II., Pt. 7, 1891.
 Drawings of *Sphaceloma ampelinum* and methods of treatment and remedies.
165. ——"Oidium in Grapes." Ibid. V., Pt. 10, 1894.
 Experiments with different preparations in its treatment.
166. DRUMMOND (J.)—"Fungi of Swan River, West Australia." Hook., Lord. Journ. Bot. I., 1842.
 A general account of some luminous fungi, in letter dated 1841.
167. ELLIS (Smith).—"Preventive for Rust in Wheat." Pp. 15, Melbourne, 1890.
 Recommends that the seed-wheat be reaped when perfectly ripe, then to be kept perfectly dry, and finally sown in a wet seed bed, as the "Rust Smoke" perishes in water.
168. ENDLICHER (S.)—"Iconographia Generum Plantarum." Folio Vindob. 1838.
Ascrea pentactina figured = *A. rubra*, var. *pentactina*.
169. FISCHER (Ed.)—"Versuch einer Syst. Uebersicht über die bisher bekannten Phalloideen." 8vo. Berlin, 1886.
170. ——"Unters. z. vergl. Entwicklungsgeschichte und Systematik der Phalloideen, in Neue Denkschr. d. Allgem.-Schweiz. gesellsch. f. d. ges. Naturwiss." Bd. XXXII., p. 1, Zürich., 1889.
171. ——"Beiträge zur Kenntniss exotischer Pilze." Hedwigia, 2, 1891.
 Refers to *Mylitta australis* (Berk.) and *Lentinus Cyathus* (Berk. and Br.).
172. FRIES (E. M.)—"Plantae Preissianae in Australasia Coll. Fungi." 8vo. Hamburg, 1846-7.
 Forty-one species enumerated.
173. FULTON (T. W.)—"The Dispersion of the Spores of Fungi by the Agency of Insects, with Special Reference to the Phalloidei." Ann. Bot. III., No. X., 1889.
 Ten Australian species referred to.
174. GALLOWAY (T. B.)—"Rust of Flax." Journ. Myc. V., No. 4, 1889.
Melampsora lini (D. C.) Tul., received from S.A.
 GORDON (P. R.)—[See "Bailey (F. M.)"]
175. GRAY (G. R.)—"Notices of Insects that are known to form the bases of Fungoid Parasites." London, 1858 (privately printed).
Cordyceps Hawkesii (Gray) described and figured.
176. "Grevillea: A Monthly, now Quarterly, Record of Cryptogamic Botany and its Literature." Edited by Dr. M. C. Cooke, and now by G. Massee. 8vo. Illustrated. Vols. I.-XXII. London. 1872-94—(continued).
177. HAMLET (W. M.)—"Anthrax in Australia, with some Account of Pasteur's Method of Vaccination." Trans. Int. Med. Cong. Austr., 522, 1889.
178. HAVILAND (E.)—"On a Microscopic Fungus (Oidium monilioides) Parasitic on Cueurbitaceae." Proc. Linn. Soc., N.S.W., I., 2nd Ser., 1886.
179. HEDWIGIA—"Ein Notizblatt für Cryptogamische Studien." 8vo. Illustrated. Vols. I.-XXXII.—(continued). Dresden, 1852-94.
- HILL (W. H. F.)—[See "McAlpine (D.)"]
180. JENSEN (F. L.)—"The Strawberry (Fragaria): Its History and Cultivation." Ag. Gaz., N.S.W., III., Pt. 7, 1892.
 Strawberry-leaf Blight referred to.
181. "Journal of the Bureau of Agriculture, South Australia." Monthly. Edited by A. Molinieux, F.L.S. Vols. I.-VII. Adelaide, 1889-94—(continued).
 Contains numerous useful notes on disease-causing fungi.
182. KALCHRENNER (K.)—"Phalloidei novi vel minus cogniti." 8vo. Buda-Pest, 1880.
183. ——"Fungi in reg. div. Australiæ et Asiæ a Jul. Remy collecti." 1863-6. 8vo. Toulouse, 1880.
Pholiota prominens described and illustrated (will be given in Supplement).
184. ——"Fungi of Australia—Basidiomycetes." Grev. VIII. 151, 1880.
 Twenty species described.
185. ——"Definition of some new Australian Fungi." Proc. Linn. Soc., N.S.W., VII., 1882.
 Eight species described.
186. ——"Fungi aliquot Australiæ orientalis." Proc. Linn. Soc., N.S.W., VII., 1882.
 Five species described.
187. ——"New Species of Agaricus discovered in Western Australia." Proc. Linn. Soc., N.S.W., VII., 1882.
 Five new species described, and seven species recorded.
188. ——"Description of Two New Fungi." Proc. Linn. Soc., N.S.W., VIII., 1883.
Polyporus Pentzkei (Kalch.) and *Paxillus hirtulus* (F. v. M.) described.
189. ——"Gastromycetes novi vel minus cogniti." 8vo. Buda-Pest, 1883.
 Australian species described, with coloured illustrations.
190. KALCHRENNER (K.) and COOKE (M. C.)—"Australian Fungi." Grev. IX. 1, 1880.
 Sixteen species recorded, fifteen of which are described, and a new genus (*Anthurus*) constituted.
191. KYNGDON (F. B.)—"Rust in Wheat." Address at Conference. Ag. Gaz., N.S.W., I., Pt. 1, 1890.
 Suggests experiments for dealing with it.

192. LABILLARDIÈRE (J. J.)—"Relation du Voyage à la Recherche de la Pérouse." 2 vols., 4to, with atlas in folio, Paris, An. VIII., 1799-1800; or English translation, in 1 vol., 4to., or 2 vols., 8vo., London, 1800.
193. ————"Novæ Hollandiæ Plantarum Specimen." 2 vols., 4to. Paris 1801-1806.
Genus *Aseroë* founded, and *A. rubra* described and figured, from a specimen found in Tasmania.
194. LAMB (S.)—"Tobacco: Its Cultivation in Northern Queensland." Bull. 6, Dept. Ag., Q., 1890.
Tobacco Blight referred to, and remedy recommended.
195. ————"Tobacco: Its Cultivation in Southern Queensland." Bull. 15, Dept. Ag., Q., 1892.
Blue Mould (*Peronospora hyoscyami*) stated to be very destructive in North Queensland, but not as common in South Queensland.
196. LÉVEILLÉ (J. H.)—"Champignons Exotiques," in Ann. Sci. Nat., 3rd Ser., Vols. II.-III., 8vo. Paris, 1844.
Sphæria (*Conferia*) *atra*, N. sp. [*Spharella*], and *S. (Conf.) labeecula*, N. sp. [*Physalospora*] from New Holland, described.
197. ————"Descr. des Champ. de l'Herbier du Mus. de Paris," in Ann. Sci. Nat. (3) V. III., 249, 1846.
Thelephora (*Stereum*) *Leichardtiana*, Lev., from Moreton Bay, and *Sphaeria* (*Conferia*) *tephrosia*, Lev., with *Dothidea Greville*, Lev., from Swan River, described.
198. LIVERSIDGE (A.)—"Disease in the Sugar Cane, Queensland." Pamphlet, pp. 34 (no date).
Disease known as "Rust" and fungus considered to be the consequence of and not cause of disease.
199. LUDWIG (F.)—"Ueber einige neue Pilze aus Australien" (On some New Fungi from Australia). Bot. Cent., Pt. XLIII., 1890, and Cassel 1889.
200. ————"Contributions to the Fungal Flora of Australia." Translated and communicated by J. G. O. Tepper. Roy. Soc., S. A., XIV. 55, 1891.
Contains lists of the Australian *Uredineæ*, *Ustilagineæ*, and the parasitic enemies of Eucalypts and Acacias.
201. ————"Ueber neue Australische Rostkrankheiten" (On New Australian Rust Diseases). Zeitschrift f. Pflanzenkrankheiten. Vol. II., Pt. 3, 130-4, 1892.
Two new species are described—*Puccinia Tepperi* (Ludw.) on *Arundo Phragmites*, and *Puccinia munita* on the underside of the leaves of *Hydrocotyle hirta*. Also *Puccinia Magnusiana* (Körn) on *Arundo Phragmites*.
202. ————"Ueber einige Rost- und Brand-pilze, Australiens" (On some Rust and Smut Australian Fungi). Zeitschrift f. Pflanzenkrankheiten. Vol. III., Pt. 3, 137-9, 1893.
Five species described, four of which are new—viz., *Puccinia Burchardie*, *Ustilago Spinificis*, *U. eombarens*, and *U. catenata*.
203. MAIDEN (J. H.)—"Australian Indigenous Plants providing Human Food and Food Adjuncts." Proc. Linn. Soc., N.S.W., 1888.
Agaricus (*Psalliota*) *campestris* and *Mylitta australis* mentioned.
204. ————"The Useful Native Plants of Australia (including Tasmania)." 8vo. Sydney and London, 1889.
Agaricus campestris and *Mylitta australis* noted with reference to their edible qualities.
205. MAIDEN (J. H.)—"A Bibliography of Australian Economic Botany." Tech. Ed. Series, No. 10. Sydney, 1892.
References given to economic fungi.
206. ————"Native Bread or Native Truffle (Polyporus Mylittæ, C. and M., syn. *Mylitta australis*, Berk.)" Ag. Gaz., N.S.W., IV., Pt. 12, 1893.
Description and drawing given and preliminary chemical examination made.
207. MASSEE (G.)—"Monograph of the Genus *Lycoperdon*." Journ. Roy. Micr. Soc., Pl. II, 1887.
One hundred and twenty-nine species recorded, eleven of which are found in Australia.
208. ————"Monograph of the Genus *Calostoma*. Desv. (*Mitremyces*, Nees)." Ann. Bot. Fl., No. V., 1888.
Calostoma fuscum and *C. lurida* described.
209. ————"A Revision of the *Trichiaceæ*." Journ. Roy. Micr. Soc., Pl. 4, 1889.
Fifty-one species recorded, seven of which are found in Australia.
210. ————"Monograph of the British Gastromycetes." Ann. Bot. IV., No. XIII., 1889.
The habitats are given for Australia.
211. ————"Monograph of the Genus *Podaxis*." Journ. Bot., Feb., 1890.
212. ————"Monograph of the *Thelephoreæ*." Part I.—Journ. Linn. Soc., Pl. 3, 1890. Part II.—Journ. Linn. Soc., Pl. 3, 1891.
A number recorded for Australia.
213. ————"Monograph of *Myxogastres*." 8vo. Coloured plates. Lond., 1892.
Forty-five Australian species are described.
214. ————"Notes on Exotic Fungi in the Royal Herbarium, Kew." Grev. XXI. 1, 1892.
Cyathus Baileyi (Mass.) from Brisbane described.
215. ————"Australian Fungi." Grev. XXII. 17, 1893.
Two new species described—*Puccinia Kochii* (Mass.) and *Xylaria Readeri* (F. v. M.); and *Phoma wiccola* recorded.
- MASSEE (G.)—[See "Cooke (M. C.)"]
216. McALPINE (D.)—"Report of the Vegetable Pathologist." Dept. Ag., Vict., Bull. 12, 1891.
Objects of section of vegetable pathology dealing with fungus pests stated, and reference to reports upon *Uromyces betæ* and *Plasmiodiophora Brassicæ*.
217. ————"The Life-history of the Rust of Wheat." Ibid., Bull. 14, Pl. I.-II., 1891.
A popular account of the different phases of rust, (*Puccinia graminis*).
218. ————"Rust of Wheat." Ibid., Bull. 14, 1891.
Uredospores of *Puccinia graminis* found during the winter season as well as in summer, and *Puccinia poarum* also observed all the year round.
219. ————"Report of Wheat Blight." Ibid., Bull. 14, 1891.
Gives an account of *Septoria tritici*.
220. ————"Report on Club Root of Cauliflowers, Cabbages, Turnips, and other Cruciferous Plants." Ibid., Bull. 14, Pl. III., 1891.
Description of *Plasmiodiophora Brassicæ*, with preventive measures.

221. McALPINE (D.)—"Beet-leaf Rust or Blighted Mangel Leaves." *Ibid.*, Bull. 14, 1891.
Description of *Uromyces beta*, with preventive measures.
222. ——— "Root Fungus of Raspberry (Raspberry-root Disease)." *Ibid.*, Bull. 14, 1891.
Mycelium of a fungus found on the roots.
223. ——— "Report on Peach and Plum Leaf Rust (*Puccinia pruni*)." *Ibid.*, Bull. 14, Pl. V.-VI., 1891.
Notes the appearance of this disease not only on the leaves but also on the fruit of the Peach.
A full account of it is given, together with preventive and remedial measures.
224. ——— "Report on Rust in Wheat as Victorian Delegate to New South Wales." *Proc. Rust in Wheat Conf.*, Sydney, 1891.
225. ——— "Ueber die Verwendung geschrumpfter Körner von rostigem Weizen als Saatgut" (On the use of Shrivelled Grains of Rusty Wheat for Seed). *Zeitschrift f. Pflanzenkrankheiten*, III. Pl. 4, 1892.
Gives the results of experiments with rust-shrivelled and plump grain, showing that 87 per cent. of the former germinated, as against 67 per cent. of the latter.
226. ——— "Report on Rust in Wheat as Victorian Delegate to South Australia." *Proc. Rust in Wheat Conf.*, Adelaide, 1892.
227. ——— "Native Bread (*Polyporus Mylittæ* C. and M.)." *Aust. Jour. Pharm.*, VIII. 291, 1893.
Fresh specimen of sclerotium described, along with its edible qualities.
228. ——— "The Undescribed Uredospores of *Puccinia Burchardiae*, Ludwig." *Vict. Nat.* X. 192, 1894. (Read Nov. 1893.)
Uredospores on stem and leaves described.
229. ——— "Report on Rust in Wheat Experiments, 1892-3." Pp. 66. Illustrated with maps and drawings. Govt. Printer, Melbourne, 1894.
Results obtained from 315 experimental plots.
230. ——— "Report on Rust in Wheat as Victorian Delegate to Brisbane." *Proc. Rust in Wheat Conf.*, Brisbane, 1894.
Results obtained from 285 plots at School of Horticulture, Burnley; also from farmers' wheat-testing experiments.
231. ——— "Botanical Nomenclature, with special reference to the Fungi." *Report Aust. Assoc. Adv. Sci.*, Adelaide, V., 414, 1893.
232. ——— "Australian Fungi." *Proc. Roy. Soc., Vict.*, VII., N.S., 1895.
Twenty-eight species recorded, eight of which are new to science, in addition to one new variety.
233. ——— "Systematic Arrangement of Australian Fungi, together with Host-index and List of Works on the subject" *Dept. of Agriculture. Govt. Printing Office*, Melbourne, 1894.
Gives a list of all known Australian fungi up to date, numbering 2278 species, with habitats, occurrence, general characters, &c. Also the host-plants with their accompanying fungi, and a numbered list of all works relating to the subject.
234. McALPINE (D.) and HILL (W. H. F.)—"The Entomogenous Fungi of Victoria." *Proc. Roy. Soc., Vict.* VII., N.S., 159, 1895.
Isaria Oncopterae (McAlp.) described on dead larvæ of *Oncoptera intricata*.
235. McALPINE (D.) and TEPPER (J. G. O.)—"A New Australian Stone-making Fungus" (*Laccocephalum basila piloides*, McAlp. and Tepp.) *Proc. Roy. Soc., Vict.* VII., N.S. 166. Pl. X., 3 Figs., 1895.
A new genus (*Laccocephalum*, McAlp.) constituted and the stone-forming species fully described.
236. "Michelia, Commentarium Mycologicæ Italianæ." Edited by P. A. Saccardo. 8vo.—(continued.) Padua, 1877-94.
Published at irregular intervals, and devoted exclusively to fungi.
237. MORRISON (A.)—"Notices of Victorian Fungi: New or imperfectly described *Uredineæ*." *Vict. Nat.* XI., No. 6, 90, 1894.
Puccinia Coprosomatis new and uredospores of *Uromyces Orchidearum* (Cooke and Mass.) first described.
238. ——— "Notices of Victorian Fungi: New or imperfectly described *Uredineæ*—(continued.)" *Vict. Nat.* XI., No. 8, 1894.
Puccinia Coprosomatis, var. *Operculariæ*, new, and *Puccinia investita* (Schwein) new to Australia.
239. MÜCKE (C.)—"The 'Take-All' " (*Xenodochus cerealium*, F. v. M.), with five plates. Prize Essay, pp. 19. Melb., 1870.
The cause of this disease supposed to be due to a fungus.
240. MUELLER (F. v.)—"Fragmenta Phytographiæ Australiæ—Supplement." The fungi determined by Berkeley, von Thunem, Kalkbrenner, and Cooke. 8vo., Vol. XI. Melb. 1880.
One thousand and sixty-nine species of fungi are recorded.
241. ——— "Census of the Genera of Plants, hitherto known as Indigenous to Australia." *Proc. Roy. Soc., N.S.W.*, XV., 1881.
The genera of Australian fungi are recorded, with authority for names and year of publication.
242. ——— "Additions to Census of the Genera of Plants." *Proc. Roy. Soc., N.S.W.*, XVII., 1883.
Several genera of fungi are given.
243. ——— "Notes on Victorian Fungs." *Vict. Nat.* II., No. 6, 76, 1885.
Gives list of fungi obtained by the Botanic Department of Melbourne from 1882 to 1884 as new for Victoria. Compiled from successive records furnished by Dr. M. C. Cooke, M.A. One hundred and one species are enumerated.
244. ——— "Further Additions to Census of the Genera of Plants" *Proc. Roy. Soc., N.S.W.*, XX., 1886.
Several genera of fungi recorded.
245. ——— "Notes on Rare Victorian Fungi." *Vict. Nat.* III., No. 10, 140, 1887.
Cyrtaria Gunnii (Berk.) and *Cordyceps Taylori* are given.
246. ——— "Notes on some New and Rare Plants." *Proc. Roy. Soc., Tasm.*, 1887.
Two new fungi for Tasmania are given—*Diploderma glaucum* (Cooke and Mass.) and *Castoreum radicum* (Cooke and Mass.).
247. ——— "List of Fungi named by Dr. M. C. Cooke—collected near Lake Bonney by Miss Wehl." *Proc. Roy. Soc., S.A.*, XI., 1888.
Thirty-five species are recorded.

248. MUELLER (F. v.)—"Select Extra-tropical Plants, readily Eligible for Industrial Culture or Naturalization." Melb. Successive editions and translations up to 1891.
References to useful and edible fungi.
249. MUSSON (C. T.)—"Notes on Insect and Fungous Pests." Ag. Gaz., N.S.W., V., Pt. 8, 1894.
Notices fungus diseases of Apple, Pear, Peach, Apricot, and Vine, and suggests remedies.
250. ————"Notes on Insect and Fungous Pests." Ibid. V., Pt. 9, 1894.
Suggests measures for dealing with such pests in the spring.
251. ————"Notes on Insect and Fungous Pests." Ibid. V., Pt. 10, 1894.
Suggestions for carrying out quarantine against such pests.
252. "Native Bread (*Mylitta australis*)." Ag. Gaz., N.S.W., III., Pt. 1, 1892.
Referred to in general notes.
253. "New Victorian Fungs." Vict. Nat. III., No. 6, 80, 1886.
Fifteen species named by Dr. M. C. Cooke, forwarded by Baron von Mueller.
254. PLOWRIGHT (C. B.)—"British Uredineæ and Ustilagineæ." 8vo. London, 1889.
Occasional references to Australian species.
255. RALPH (T. S.)—"On Dry Rot." Proc. Roy. Soc., Vict., VI., 1861.
Remarks chiefly in reference to Pines.
256. ————"On the *Æcidium* affecting the *Senecio vulgaris*, or Groundsel." Vict. Nat. VII., No. 2, 18, 1890.
257. READER (F.)—"Notes on some hitherto Unrecorded Victorian Fungi." Vict. Nat. II., 66, 1885.
258. ————"Note on *Hirneola polytricha*." Vict. Nat. IV., 174, 1887.
259. REICHARDT (H. W.)—"Fungi, in Reise der Oesterreichischen regatte Novara um die Erde in den Jahren 1857, 1858, 1859." Botanischer Theil. 4, Wien., 1870.
Two species—*Lycogala leiosporum* (Rehdt.) and *Hydnum griseo-fuscescens* (Rehdt.)—described, from Sydney.
- REMY (J.)—[See "Kalchbrenner (K.)."]
260. "Revue Mycologique, recueil trimestriel illustré consacré à l'Étude des Champignons. Edited by C. Roumeguère. 8vo., Vols. I.—XV.—(continued)." Toulouse, 1879-94.
261. "Rust in Wheat—Minutes of Proceedings at a Conference of Delegates from Victoria, South Australia, New South Wales, and Queensland." Melb., March, 1890.
A series of experiments, and the issuing of questions to farmers suggested. Appendix to report on "The Nature of Rust in Wheat," by A. N. Pearson.
262. "Rust in Wheat—Report of the Conference at Sydney." Sydney, 1891.
Delegates from the four colonies as above, and results of experiments, together with replies from farmers given.
263. "Rust in Wheat—Report of the Proceedings of the Conference at Adelaide (Third Session)." Adelaide, 1892.
Delegate from Tasmania in addition, and reports on replies from farmers and on experiments given.
264. "Rust in Wheat." Ag. Gaz., N.S.W., III., Pt. 7, 1892.
Details of mode of conducting experiments.
265. "Rust in Wheat—Report of the Proceedings of the Rust in Wheat Conference at Brisbane (Fourth Session)." Brisbane, 1894.
Reports from the various delegates and descriptions of prominent varieties of Wheat with illustrations.
266. SACCARDO (P. A.)—"Sylloge Fungorum omnium hucusque cognitorum." 8vo., Vols. I—X., Padua, 1882-92.
Describes all the known species of fungi up to June, 1892 = 39,663.
267. ————"Notes Mycologiques—Mycetes aliquot australienses." Series I., Hedwigia, 125, 1889. Series II., Bull. Soc. Myc., France, V., 116, 1890. Series III., Hedwigia, 1890. Series IV., Hedwigia, 1893.
Series II. consists of twenty-two species, four of which are new. Series III. consists of eighteen species, nine of which are new. Series IV. consists of twelve species, four of which are new.
268. SACCARDO (P. A.) and BERLESE (A. N.)—"Fungi australienses." Rev. Myc. No. 26, 1885, or Atti del. R. Instit. Veneto di Scienze, &c., Venetia, 1885.
Fifty-one species recorded with figures. Three new genera are constituted—*Scortechinia*, *Gibellia*, and *Gamospora*; and eighteen new species described.
- SACCARDO (P. A.)—[See "Bresadola (J.)."]
269. SCHLECTENDAL (D. F. L. V.)—"De Aseroës genere Dissertatio." Berlin, 1847.
270. ————"Eine neue Phalloidee, nebst Bemerkungen ü. d. ganze Familie derselben." Linnæa. Bd. XXXI., 101, 1861.
Contains general remarks on the *Phalloideæ*, which are perhaps better represented in Australia than in any of the other great divisions of the globe.
271. SKUSE (F. A. A.)—"The New Zealand Vegetable Caterpillar." Vict. Nat. VIII., Nos. 2 and 3, 47, 1891.
Criticises paper by T. Steel on the subject, and expresses doubts as to the *Spharria Robertsii* being associated with the caterpillar of *Hepialus virescens*.
272. SOUTHALL (W.)—"Note on a Specimen of *Mylitta australis* (with figure)." Pharm. Journ. (3) XV., 210, 1884.
273. SPICER (W. W.)—"Ergot." Proc. Roy. Soc., Tasm., 1877.
On *Lolium perenne*.
274. STEELE (T.)—"The New Zealand Vegetable Caterpillar." Vict. Nat. VIII., No. 8, 110, 1890.
Refers to Australian specimens of entomogenous fungi.
275. STEPHEN (T.)—"Smut in Wheat." Proc. Roy. Soc., Tasm., 1889.
Steps for prevention given.
276. SUTHERLAND (G.)—"The South Australian Vine-growers' Manual." Adelaide, Gov. Printer, 1892.
Oidium, Black Rot, *Peronospora viticola*, Black Spot or Anthracnose, Pox or *Gloeosporium ampelophagum* referred to at pp. 101-104.
277. TATE (R.)—"A List of the Charas, Mosses, Liverworts, Lichens, Fungs, and Algae of Extra-tropical S.A." Proc. Roy. Soc., S.A., Vol. IV., 1881.
Eighty-six species of fungi are recorded.

278. TAYLOR (R.)—"Description of the Bulrush Caterpillar (*Sphaeria Robertsi*)." Tasm. Journ. I., Pl. 1, 1842.
A drawing of *Sphaeria inominata* is given from N.S.W.
279. TENISON-WOODS (J. E.)—"Botanical Notes on Queensland." Proc. Linn. Soc., N.S.W., VII., 1882.
Hexagonia crinigera (Fr.) mentioned.
280. TENISON-WOODS (J. E.) and BAILEY (F. M.)—"On some Fungi of New South Wales and Queensland." Proc. Linn. Soc., N.S.W., 1880.
Notes on the genera and more remarkable species, followed by a classified list.
281. TEPPER (J. G. O.)—"Red Rust: Its Nature, Approximate Cause, and Probable Cure." Proc. Roy. Soc., S.A., III., 1879.
282. ————"Botanical Notes Relating to S.A." Proc. Roy. Soc., S.A., VI., 1883.
Additions to the list of Australian fungi—eighteen species (eight unrecorded for Australia and ten new to S.A.).
283. ————"Fungi collected near Clarendon 1882-3." Proc. Roy. Soc., S.A., VIII., 1885.
Eleven species are recorded identified by Dr. Cooke.
284. ————"Additional Lichens and Fungi of S.A., collected from 1880-85." Proc. Roy. Soc., S.A., IX., 1887.
Five species of fungi recorded.
285. ————"Notes on South Australian Fungi." Proc. Roy. Soc., S.A., XII., 1889.
List of fungi new or rare for S.A.—fourteen species—together with Australian *Ustilaginae*.
286. ————"Additional Species of Australian Fungi." Proc. Roy. Soc., S.A., XIII., 1890.
Records twenty-four species collected by himself in S.A., and described by Winter, Saccardo, and Ludwig in various journals.
287. ————"Take-all and its Remedies." Ag. Gaz. N.S.W., III., Pt. 1, 1892.
Fungus not considered to be the cause of the disease, but simply starvation of the crop.
- TEPPER (J. G. O.)—[See "Ludwig (F. M.)" and also "McAlpine (D.)"].
288. THOMPSON (E. H.)—"A Handbook to the Insect Pests of Farm and Orchard." Bull. No. 1. Dept. Ag., Tasmania.
Treats also of fungus pests, particularly *Puccinia pruni* and *Fusicladium dendriticum*.
289. THUMEN (F. von.)—"Symbolae ad Floram Mycologicam Australiæ I." Grev. f., 1875."
290. ———— Ibid. II., Flora, 1878.
291. ————"Mycotheca Universalis." Bayreuth 1879.
Ustilago Muelleriana recorded as a new species.
292. TISDALL (H. T.)—"Fungi of Country East of Mount Baw Baw." Vict. Nat. I., No. 15, 169, 1885.
Seven species of *Agaricus* in its extended sense are recorded, determined by Dr. Cooke.
293. ————"Fungi of North Gippsland." Vict. Nat. II., No. 9, 106, 1886.
Eight species described, and found a *Polyporus* developed from *Mylitta australis*.
294. ————"Notes on Fungi in Mines." Proc. Roy. Soc., Vict., XXIV., Parts I.-II., 1887.
295. ————"Victorian Agarics." Vict. Nat. IV., No. 12, 203, 1888.
Forty-three species briefly described, and localities given.
296. TISDALL (H. T.)—"Fungi of the Season." Vict. Nat. VI., No. 7, 107, 1889.
Fungi found in or near Melbourne.
297. ————"A Curious Fungus." Vict. Nat. VI., No. 7, 119, 1889.
Species of *Cordyceps* growing from an ant *Formica corisobrina*, and found by Mr. C. French, Government Entomologist.
298. ————"Victorian Fungs new to Science." Vict. Nat. VII., No. 7, 96, 1890.
Seven new species recorded, and six of these described.
299. ————"On a Species of Isaria." Vict. Nat. X., No. 6, 90, 1893.
Found on a cocoon, supposed to be that of the moth *Darula ocellata*.
300. "Tobacco Industry in the Adelong and Tumut Districts." Ag. Gaz., N.S.W., II., Pt. I, 1891.
Tobacco Blight (*Peronospora*) referred to and remedies suggested.
301. TRYON (H.)—"Report on Insect and Fungus Pests." Dept. Ag., Queensland. 8vo., pp. 238. Brisbane, 1889.
Records fungus diseases in Apple, Pear, Peach, Almond, Orange, Vine, Pumpkin, Potato, Maize, and Wheat.
302. TURNER (F.)—"Xylostroma giganteum, Fr. (a peculiar fungus)." Ag. Gaz., N.S.W., III., Pt. 6, 1892.
Obtained from the heart-wood of several Eucalypts.
303. TULASNE (L. R.)—"Fungi Hypogaei, p. 199, folio, 1851." Refers to *Mylitta australis*, &c.
304. "Victorian Royal Commission on Vegetable Products, 1885-94." 8vo. Melbourne.
Ten Progress Reports issued, and fungus pests occasionally referred to.
305. WALLACE (R.)—"The Rural Economy and Agriculture of Australia and New Zealand." 8vo. London, 1891.
References to rust in Wheat, Anthracnose of the Vine, Oidium, Ergot, and *Peronospora* in Australia.
306. WALLIS (A. R.)—"The Vine Disease, Oidium Tuckeri." Two plates. Ann. Rep. Dept. Ag., Vict., 1873.
307. ————"A New Disease among Rye Grass." Two plates. Ibid., 1873.
Isaria graminiperda (Berk. and F. v. M.) as the cause of it, described in *Gardener's Chronicle*, 596, 1873.
308. WEHL (Miss.)—"List of Species of Agaricus and Panus, discovered near Lake Bonney." Proc. Roy. Soc., S.A., X., 1887.
Seven species of *Agaricus* recorded, and one of *Panus* (*P. carbonarius*).
309. WINTER (G.)—"Exotische Pilze II., Hedwigia, 1885." Four new species described.—*Uromyces vesiculosa*, *Asterina microthyrioides*, *Phyllachora nervisequia*, and *Lembosia orbicularis*. *Meliola cladotricha* (Lev.) also further described.
310. ————"Fungi Australienses." Revue Mycologique, Toulouse, 1886, and Rev. Myc. 1888.
311. WOOLLS (W.)—"A Contribution to the Flora of Australia." 8vo., pp. 255. Sydney, 1867.
Occasional references are made to fungi in the neighbourhood of Sydney.

CORRECTIONS.

P. 13.—No. 180.—S.A. in wrong column.

P. 124.—Order XX. should be XXV.

P. 156.—No. 193 should be 1939.

P. 178.—No. 2133.—*Colletotrichum* should be in brackets as a synonym.

P. 180.—No. 2152.—Add *Carpozyma* as a synonym, since Apiculate yeast is considered by some to belong to this genus.

3 5185 00143 2002