Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

62.71

INDEX

Price List of Fruit and Nut Trees

Season 1923-24

PEARCY BROS. NURSERY

237 State St., Salem, Oregon

These prices are subject to change without notice.

Terms: 10 % cash with order, balance C .O .D. No trees reserved unless covered by a 10 % deposit. Packing for shipment at cost.

100 rates apply to 50 trees or more

We predict a shortage in Walnuts, Royal Ann cherries, Mazzard cherry seedlings, filberts and Bosc Pears before the season is over.

Cover your needs in these lines early.

GRAFTED FRANQUETTE WALNUTS.

	Each	Pre 10	Per 100
6-8 ft.	\$1.85	\$1.75	\$1.60
4-6 ft.	1.60	1.50	1.35
3-4 ft.	1.25	1.20	1.10
Selects	2.00		

FILBERTS

We furnish Barcelona variety with enough DuChilly, Daviana and White Aveline for pollenators. The bulk of our stock is layered stock and one year old nursery grown stock. We have a small amount of two and three year trees. No pollenizers sold except where we furnish Barcelonas.

LAYERED OR NURSERY TREES

No.1	.60	.50	.50
No.2	.45	.40	.35
2 or 3 yr. trees	.70	.65	.60
Selects	.75		

ITALIAN PRUNES

6-8 ft.	.30		.25
4-6 ft.	.25		.20
3-4 ft.	.20		.15

SWEET CHERRIES

ROYAL ANN, BING, LAMBERT, WATERHOUSE
BLACK REPUBLICAN

4-6 ft.	.75	.70	.65
3-4 ft.		.60	.55
2-3 ft.		.45	.40

Bing and Lambert 5¢ under the above
and Black Republican 5¢ higher.

Sour Cherries—Large Montmorency and
E. Richmond (Kentish).

Prices same as Royal Ann except that best
grade is 3½-5 ft. instead of 4-6 ft.

MAZZARD CHERRY SEEDLINGS

One year 4-6 ft.	.25	.20
One year 3-4 ft.	.20	.15
One year 2-3 ft.	.15	.10

Two year trees with heavier root systems
add 5¢ to above price.

PEACHES

Charlotte, Early Crawford, Late Crawford,
Muir, Elberta, Champion, J.H. Hale, Hales
Early, Triumph.

	Each	Per 10	Per 100
3½-5 ft.	.45	.40	.35
3-4 ft.			.30
2-3 ft			.25

ALMONDS

Texas Prolific and Drake

3½-5 ft.	.50	.45
----------	-----	-----

APRICOT

TILTON	.50	.45
--------	-----	-----

APPLES

Delicious, Spitzenburg, winter Banana,
King, Yellow Newtown, Yellow Transparent,
Grimes Golden, Jonathan, Northern Spy ,
Rome, Gravenstein, Ortely, Baldwin, Whitney
Crab.

4-6 ft.	.45	.40	.30
---------	-----	-----	-----

PEARS

4-6 (except Bosc).	.45	.40	.35
--------------------	-----	-----	-----

Bosc 5¢ higher.

PLUMS

Hungarian, Shiro, Blue Damson, Peach

4-6 ft.	.45	.40	.35
---------	-----	-----	-----

QUINCE

Orange

GRAPES

Concord, Campbells Early, Worden, Niagara,
Moores Early.

Two year	.30	.25	.20
----------	-----	-----	-----

Campbells Early and Moores Early add 5¢
per plant to above.

GOOSEBERRIES—OREGON CHAMPION

	Each	Per 100	Per 1000
No. 1	.15	\$5.50	\$50.00

CURRANTS PERFECTION

.15	5.50
-----	------

RASPBERRIES CUTHBERT SUCKERS

5¢	1.50	7.50
----	------	------

BLACKCAP RASPBERRIES

Munger One year old transplants

Each	Per100	Per1000
\$.15	\$3.50	\$30.00

BLACKBERRIES

Himalaya 1 yr. transplants

.15	6.00
-----	------

Lawton Suckers.15 5 .0 0

¶ Pollenizers--The Barcelona filbert will not bear commercial crops unless certain other varieties are planted close to it. Certain varieties have a greater effect upon it than have others. We advise planting of about 83% Barcelona, 9% DuChilly and 4% each of Daviana and White Aveline.

The sweet cherries are also self-sterile. They do not set commercial crops unless pollenizers are planted with them. Royal Ann, Bing and Lambert are all in this class. Neither do these varieties pollinize each other. Black Republican, Waterhouse or Governor Wood pollinize these varieties. About one pollenizer in nine is needed.

Mazzard cherries. Our sweet cherries are subject to a gumming disease called Gummosis. This disease often girdles and kills or badly injures the tree. It cannot be controlled by spraying. The best method of avoiding this disease is to plant Mazzard seedling cherry trees in the orchard and top work them to the desired varieties after two or three years. The gummosis seldom attacks the Mazzard hence in this method of building an orchard the disease is kept out of the trunk and main crotches of the tree.

Fall planting is preferable provided well ripened trees can be had from the nursery before the ground becomes too muddy for planting.