Dec 105

1873

W. C. C. C.

Paid laundry at home

5.06

376

due me 1.26

doc. 108

3338

Charles Schuchert
Yale University
New Haven, Conn.
1914.

39057

Eastern Canada with
McClellan

Prof. E. Hageroll
Acadia College
Wolfville, Nova Scotia.

Met H. A. Bell at Windsor
May 26, 1914.

Took him to Tor Brook, Nictaux
and Dictyonema websteri locality

Hageroll.

Took him Helder. and Crick. fossils.
See about slides of life restored for
each period.

New Haven, May 16-1914.

Left New Haven with McLearn at 5.50 P.M. for Boston. Got there at 9.40. Put up at the Adams House and then walked about the city for an hour.

Boston, May 17-1914. Sunday.

After breakfast saw the Capitol, Bunker Hill Monument. Then to Harvard where we spent most of the afternoon seeing the Museum of Comparative Zoology. Then the Boston Society of Natural History and the Boston Library!

Had dinner at the Adams House and then took the 7.30 at North Station for Boston. Had an afternoon rest.

St. John, N. B. May 18. Sunday
Got here at 12 noon eastern time.
Stopping at the Hotel.

In the afternoon collecting eastward
at the Suspension Bridge. To the west is
a mass of much white limestone
of the Silurian. This is Frégère's age. Then
a fault zone with the Permian or Upper
Carbonian strata again, the limestone but
with a less steep dip. The thickness did
not make out but was 200 ft. or more.
This is a series of light green thin bedded
sandstones and sandy shales with thin
single layer bedding for most of the thickness.

Then another fault zone followed by
the Tetragraptus shales with a thickness of
100 ft. or more than 30 ft. There are also
standing at a right angle and much crumpled
and tilted so that the beds are not
in place. At the very base one got one stop
[D6085]

[D6086] —→

Stalen v. v.

We often stop at some small place, and
 make a short stay. Usually the road leads us
 to the top of a high mountain, and we
 often find a large number of small
 lakes and streams.

Each of the order of name is to be
the same. The due

The Etcheminian on the Lower Peninsula
and the whole of the region. 'Dend' is a very distinct.
For base see legend.

gastrolites and coryphids. About at the top of the exposure we get the large lot of gastrolites. These two zones may be 20 feet apart but hardly more.

The brachiopods (orthids) come from str 10 feet below the upper gastrolite bed. They do not occur in calcareous nodules, but are always changed to iron pyrite.

Friday, Tuesday May 19 - 1914

At the Cambrian locality on July 2nd.

One of the localities are abrupt cliffs of a greenish shale (trachyte) of green material. All along the side of the exposure are cliffs and to the south the land descends into the Cambrian shale and the upper Cambrian sandstone of the same color.

Above the Trachyte (contact with green shale) 30 to 40 feet of fine grained micaceous shale without fossils; but put in the others. This is the Silurian. (See page to left)

This ^{sandstone} zone is seen of wide distribution in
north of John.

These shales and others with thin sandstone
conform to the next strata above and
then are soft dark green shales to the bottom
of the valley in which lies a stream. On the
other side of the stream the land rises into
the Upper Cambrian.

If there is no repetition the Cambrian must
be several thousand feet thick. This can give
any idea of the thickness.

Then comes in a white coarse leamy bedded
sandstone about 30 feet thick. In the middle
of the section it is fine grained at the top and more even bedded. The upper
surface shows fine ripple marks.

This seems to be ^{the} structural base of the Acadian Cambrian.

On the latter reference, can say
that sandstone is fine grained, and
is sandy shales more sandy than shales.
The middle Cambrian is fine grained

can
towards the top.
here. As they are in the red
of the top of the white quartzite
All ~~that~~ become more and more fine grained to
near the top, where the fossils abound in great numbers.

About 1/4 mile farther east we are in the
public park where the contact between the trachyte
and the St. John's granite can be clearly seen.

The basal 8 feet of the St. John's granite is an almost
dark red in color, very coarse in crystalline with zones

May the Elohemian be of continental character, or at least of shore deposition?

See specimens)

which are
of pebbles up to 3 inches across. These pebbles are not
well rounded and are of hard material usually a
red granite. Then there are pieces of red gneiss iron ore,
some red shale pieces, quartzites and schists. The
variety is considerable but rarely are any of the
more rare ones. The whole weathered down rather
easily into a coarse sand.

A little farther south the white quartzite of
early stage is seen to on a mountain 15 feet off the
trachyte cutting out near the Italian American.

Porter gneiss. The dolomite is practically
recrystallized.

Below it is a thick zone of quartzite
also much recrystallized.

Both are cut by granites and other igneous
rocks.

Left at 12:40 for Tucson where we arrived
at 8:00. Stopped at the Learmont Hotel.

Thurso, May 20 - Wednesday

Before breakfast visited Victoria Park.

At 9.40 on way to Gairloch, where we arrived at 1 P.M.

Had dinner at the Royal George and then by carriage to Arisaig. The team and man cost £5⁰⁰. Started at 2.30 and got to Mrs John A. McDonald's at 5.30. Drove 21 miles.

Arisaig, May 21. Thursday

Spent the day with Mac ^{neil} along the sea cliffs from Arisaig Brook to the end of Strachan's cliffs. Much water has fallen down making it more easy to get the fossils and the horizon exposed. Fossils however are not abundant.

Beetey Hill shale,

sandy shale (sand variegated)

Apophyllite

Beetey Hill shale with local small
lenses of banded sandstone.

Friday May 22 - Friday.

Started our car to see the basal Silurian beds.

On the McDonalds road we saw the Beech Hill formation resting on the Apophyllite. Here it is a fine grained sandstone with several zones of arenaceous limestone and all of them have fossils in abundance but of few species. Much of the formation goes over into shales and finally even into the Ross Island shale. This fossiliferous zone is about 15 feet thick and is toward the top of the Beech Hill formation. The reason why it is sandy here is because it is adjacent to an island of apophyllite that is very fine and its sand goes into the sedimentary muds. These beds stand out well.

We then went further east to Beech Hill Cove. Here on the northern side of the cove the Beech Hill formation again stands out well due to an increase in the rock distance. The formation is a blue sandy, arenaceous shales with very local

Masses of bedded sandstone. There are many of these
in the lower 100 feet. Very rare one of them is
long when it occurs, piece of a single species
of Corbicularia - Corbicularia can be seen
thomasioides. Salterella intermedia. seen in
about 75 feet above the base (base however not
seen).

The lower Beechey Hill shales are marked
by a delicate Psithyrion; are pressed flat
and of no value as fossils.

On the eastern side of Beechey Hill one
sees considerable of the Beechey Hill formation
but no contact with the apophyllite is seen.
Here again the upper part is more shaly. In all
of these places the contact with the Ross Brook
shale is clear and one can make out the
contact easily. Saw but few fossils here.

We then went to Doctor Brook where the
Beechey Hill formation stands on and as always
in contact with the apophyllite. In the lower

reach the bridge over the brook we see the
contact with the apophyllite. The basal water
appears to be down a bed down apophyllite clay
with quartz pebbles as it is the weathered soil
and then a fine grained conglomerate. In any
event the apophyllite was weathered and was
land on which the Lilliman sea rippled.

At the base of the brook the Lilliman
formation is hard and sandy and makes cliffs.
The bridge stands on the uppermost beds of it
and to the north the land rises down to the
the base of the Lilliman shale.
Bed runs from the base of the Lilliman shale as 200 feet.

May 23, Saturday.
The Devonian contact. The Devonian is the
highest land here and is the base of the
Devonian series.

We then went down Mc Alas brook and where
in the Devonian. It is ^{in the lowest part} a series of red to maroon
shales interbedded with thin zones of greenish sandstones.
The shales nearly always have small nodular con-
cretions some of which stand out like the reddish.
These I interpret to be caliche deposits, the sweating
out of the water under a dry climate due to
capillarity. Therefore concretionary deposits, probably on
a delta. ^{than} the top of the land, within a 1000 feet
the road left in the red shale a fish bone and
three bivalves. If these are fresh-water shells it is
quite a find. This horizon is no down in the
Devonian and ^{series and probably several hundred feet lower than}
the fossil locality beside the road.

Towards the lower part. Mc Alas brook
we came upon the ^{even yesterday} trip ^{above} the

5/11/20

Estimated
in thickness

Niagara Falls
on the 1st of Nov.
see them on the water.

over the 1700000000
the Western

The Area of the Area

Sept 5. 1820 about
the distance from the
village to the river

Andean Concomitant
No. 1 W.

My dear
John

in this block
is an section. Beneath the diabase lies the
Hendon conglomerate, showing here at the stream
level about 4 feet. The conglomerate is a dark
maroon with flat pebbles ^{of the Lewisian series and Devonian}, all dimensions up to
3 inches here (not elsewhere up to 6 inches), fairly
well rounded and lying all on their flat sides and
flat with the bedding. It is not a stream deposit
but a beach shingle.

On this conglomerate lies the diabase which to
me seems to be an intrusive for it lies regularly
upon the bedding of the conglomerate, has its dip and
the basal beds are vesicular and quite even. The
vesicles are small from $\frac{1}{4}$ to $\frac{3}{8}$ inch on the
average. About 20 feet higher is another more decided
vesicular zone.

Further examination shows that it is un-
mistakably a flow and not an intrusive.

Beyond is ^{a second} flow and like the previous one. On
the sea side one sees the upper part. Here it is
a breccia full of vesicles filled with ^{and calcite} quartz.
Then comes a considerable thickness of ^{basalt} on

trend
 into
 position
 as seen
 from base
 of cliff.
 correct.

This
 is a
 fault
 as seen
 from top
 of cliff.
 correct

Old raised beach surface
 sea level

Did the shale have any
 the distance of the top?
 Or is there a natural
 exposure here?
 Mac seen with and why?

The dip is 10 degrees
 The strike is 10 degrees
 The same as the other

sea level

conglomerate. The ^{little} beds just above the second trap
is made up of the diabase worn down into natural
fall size. Finally to the west we see a third
flow. Here again the base is highly vesicular
and lies fairly level upon the Devonian conglom-
erate.

All of this evidence goes to show that this
country was depressed into mountains during the
Devonian, that the topography was rough in
Ordovician ^{time}, and that active volcanoes were abundant
during the ^{early} Devonian time.

All previous writers assumed that the
red beds on the sea shore beneath the diabase
was the end of the Silurian section. Making out
the intrusive nature of the diabase and that
the same flow lies upon the Silurian, Devonian
and Ordovician I concluded it to be worth while
to see if the Silurian indeed as supposed. To
my great surprise I saw the Silurian dipping

into the land, ^{with west} and therefore that there was no
more Silurian. In going up the little Stonehouse
road I soon saw that above the red Silurian
lay considerable more massive rocks. These are
flaggy ^{light green} sandstones and shales of greenish ^(light blue) and
sandy greenish shales. Fossils are scarce
but brachiozoa turn up frequently and occasional
chonetes or nautilus. ^{These are the} ^{trilobite} rocks in which I got so many brachiozoa
when collecting here. In fact, I think we
could not go far from this place for the trilobite
seen ^{still} here and in connection with the
lower beds. We have collected some hundreds
of them. It was here where the dip was
like the one we saw. This seems to show that
the Arising Silurian are entirely unbroken into
the Devonian.

I am quite certain that many hundreds of
years ago and the shales must be added
to the Silurian section as determined by Town-

is in line between the red Stonehouse and the
Lafel. Probably ^{red Devonian} as much as 1000 feet. If
this observation is correct then the Silurian
marine series has a thickness of not less than
4000 feet followed by about another ¹⁰⁰⁰ ^{or more} feet of
red Devonian beds.

Over the Silurian and Devonian un-
conformably lie the Lower Devonian series. It
has been deposited since its deposition.

The question will then be asked as to what
is the age of the "Red Devonian". If it is synchronous
with the Silurian it is either a Silurian, or
to judge of the Gorge sequence it may go up
into the Devonian quite high. At Arvian con-
ditional conditions known in late Stonehouse time
(may mean early Lelina time), while at Gorge
it did not start until Middle Devonian time.

August 22 Sunday.

In morning a nice courteous young Mr. MacDonald was vol. to take me to Antigonish. He started up and was off at 8.30 A.M. Got to Antigonish at 11.15 A.M.

Mac Learn made out my expense for the trip at 61 dollars which is paid by the Canadian Government. This does not include meals and hotel en route. I am allowed 33 dollars to get me to my haven. The transportation charges to Antigonish is \$23.80 including my Pullman fare.

Therefore Mac Learn will owe me in the fall when he gets back the \$33. I paid out on the way back.

Stopping at King George Hotel.

After dinner strolled around Antigonish

and saw the ~~conglomerate~~ conglomerate. Here
it is like that seen ^{west} of Strehouse Brook, a dark
brown pebble conglomerate, each piece with desert
like patina. This patina is on all the pieces and
is of the color of a fine sandy soil. The pebbles
are full - in size up to 8 inches or even one foot across.
In other places the conglomerate takes on the
color of a brown sandstone a light red and
then the pebbles are in the matrix of the sandstone.

I first saw the conglomerate where the Arising
road crosses the Dickinson road. Probably
1000 feet or more lower down geologically I came
across a gypsum bed, 50 feet or more thick.
It was covered by either a red clay or a fine
red conglomerate. I saw it across a small stream
beside the railway but could not get near to it.

As the Dickinson is conglomeratic in so many
places and is always thick, the evidence in-
dicates that the country was mountainous, filling
in the valleys below. The climate was arid

and fairly cool. The water must have been massive
on account of the great amount of gypsum de-
posited.

Antigonish May 25. Sunday.

Forded around the village all morning
and at 1.04 P.M. off for Truro. Arrived
here at 5 P.M. Stopping again at the
Seamark Hotel.

Truro May 26. Tuesday.

Left on Dominion Atlantic on the 6.40 A.M.
Got to Windsor at 8.20 A.M. Met ^{H.A.} Bell at the
station and we are off at 9 A.M. for Horton
Bluff. Returned to Windsor at 5.10 P.M.

These very coarse granites have not only broken
through the Silurian but also cut the
Victory Gneiss. They then belong to
the great Permian disturbance. These granites are
in great exposures all through eastern Nova
Scotia and especially in the southern and
portion of Nova Scotia.

the arkoses rounded pebbles of green shales
pieces of some mud hard torn up and rolled
into the arkoses. All in all it gives one the
idea of stream action rather ^{than} sea wave as-
sisting. Rarely there are small accumulations of quartz pebbles ^{in sizes from 1 to 2 1/2 inches.}

The red slays are Windsor like in their coloring,
have much muscovite and sand and have a
lacy fracture. In places one may irregularly deposits
by glacial observation one also saw here and there
rootlets more horizontal than vertical in position.

The black shales ^{are more local and} are coarser and are associated
with the arkoses as if they formed in local pools
during the intervals between the freshets. The black color
is due to plant material and muscovite.

The arkoses have at times great masses of red
shale inclusions, masses torn up by the stream
and ^{moved or} cut into leaning beds 5 to 75 feet
long standing out off in each end by the arkoses.

One sees no evidence of logs in the arkoses
but occasionally in the dark shales there are
pieces of ^{small} *Leptodendron*.

It may be seen from the statement that the
andesitic section becomes more flat bearing as
one rises in this series.

The entire section may be 1000 feet thick.

One possible idea is that the western section is younger
than the eastern. However there is no definite proof.
and the section consists of andesitic section of small size
the western section. Perhaps the andesitic section is from 500 ft.

In a blue fine grained shale between
 and one layer we saw considerable *Leptidodonta*
 and not like *Leptidodonta* ^{with the plants} associated in certain layers
~~about the same time~~ we got *Estheria*. Saw no
ostrea but did see some cases. This locality
 is due to the east of the fault. It is here probably
 that the *Leptidodonta* were collected.

A few hundred feet farther west and the
 strata stand on end.

Carbonaceous and dark shale

Exposed zone
 has dark shale
 500 feet across.
 End of section

Regulation Horton.

It looks to me as if the masses should be at the base
 of the Horton but here says the evidence seems to point
 that they are at the top.

The outcrop Horton is probably 200 or more feet
 thick and consists of dark blue to almost black shales
 with more or less thick zones of fine grained sandstone.
 All are banded due to the great deformation.

The rest of the Horton or more than one-half

of the section consists in the main of dark blue to almost black shales with at first no sandstone zones, then here and there a thin one, with more and more of sandstones towards the top. These however are never dominant in the section and at times, are also in tabular layers of fine grain. Most of the sandstones are rippled, and sun-cracked.

One conventional limestone layer, more or less with septaria was seen. The layer at the top may be said to be decidedly lumpy. The thickness ranged from about 2 to 5 inches. It formed while the beds were still soft.

Ostracodes are very abundant along with goniatid oolites in the shales near the base of the western half ^{or lower part} of the section. Otherwise ostracodes are absent. Plants are scarce throughout but at the top are more common. In one zone of sandstone about 6 inches thick and in the adjacent shales saw many vertical *Lycopods*. In size they ranged from 3 to 10 inches ^{in diameter} and there is one to two square feet

of ground. A forest over-shelved by the sand-
stone with the top rotted away in the muds into
which some projected.

There can be no doubt that the arkosic Horton
is of continental origin. Not a marine fossil is
seen here and the *Estheria* and *Leania* must be
of fresh water origin.

The great shale Horton series is probably
also of continental origin although the even bedding
of the shales looks more like marine deposits. The
Ostracoda here occur near the base and as
Beyrichia occur here maybe of brackish water
origin. However higher up one sees only land
plants, garnet scales and bones, rippling of very
shallow water and often of very irregular pattern,
sun-cracking and root cut prints.

The dark color of these shales is probably due
plant material, broken up fragments, spores and
spore cases (the latter are common). At first one is
disposed to regard them as marine because of
their even bedding.

Windsor - Wolfville May 27, Wednesday.

Left Windsor at 9 A.M. for Wolfville to see Prof. E. Hageroll and the Dictyonema web-
steri locality at Beech Hill.

This is Commencement day at Acadia here at Wolfville and Prof. Hageroll was at the train to see one up to the exercises. Had to go and then upon the stage. It lasted from 10.30 to 1 P.M. About 75 women and men took their degrees and 3 honorary degrees were conferred upon their own graduates.

In the afternoon with an automobile we covered the country here about. First we went to Hageroll's fruit farm and then to Beech Hill.

The Horton lies only in the lower ground going into the Sackville and Cornwallis valleys. It is the impression that it runs west over the higher hills to the east but may

have gone over the Silurian hill to the west
of the Gasperan valley.

The present general topography was established
since the Triassic for these deposits ^{are} in
the valleys but probably did not outcrop the
Gasperan ridge. It is the general belief that
there is no Triassic in the ^{lower} Gasperan valley
for it is to be seen only in the lower part of
this valley. However Hays Creek states that
in coal digging Triassic was encountered in
the upper Gasperan valley. Is this Triassic
deposit that has preserved the topography of
the Permian times.

At White Rock in the upper Gasperan
valley we came upon schistose quartzites and
green slates. The dip of them is hard to
make out. Everywhere where we saw the
slates schistosity is developed, in other words
it is regional schistosity. Finally we found the
locality for Clity mena oresteri at Buck Hill

Deer tells me it is the opinion of Fletcher and others that the Silurian is conformable upon the Gold Slate series and that all were folded together. This hardly seems possible when one learns that the latter has a thickness of 36,000 feet, one-half of which is quartzite the other slate.

although we did not find this fossil. It is
always a light colored more green to dark
blue slate with glossy surfaces. Occasionally
there are zones of a coarse grained quartzite.

This series of slates and quartzites may
be the Archaean series. As corals have
been found about Cannan with probably
Chonetes norasectica the fossil evidence is in
favor of Upper Archaean.

Prof. Hageroll showed me a large amount
of Oriskany material he collected in a pasture
but a short distance to the east of Oriskany
Falls. The fossils while not good show
an Oriskany ^{fauna} not in keeping with the Sarco-
nian or Fall aspect. There are several species
of Spurifer, 2 or 3 of Strophomena, Ectoclema
near E. multilobis of the New Scotland, at least
one species of Rhipidomella, ? Hippoclema, Proteroclema,
Leptocoelia, Strophomena (large) and possibly a
Megaclema (small). There are L. arenosus.

Nictaux, Thursday May 28.

Left Wolfville at 9.30 and got to Middleton at 10.45. Stopping at the American House.

After dinner took the train east to Nictaux 4 miles and then walked north-east to Tor Brook. The iron mines are around Tor Brook but they are all shut down now. On the dumps we saw considerable Oridany fossils in the "shell rock". *Spirifer* are very common and the largest among them is *S. arenosus*. The variety is also not large.

The ore is a very fine grained flax-sand as lies that of the Clinton. The iron runs to 50%. The bed is said to range from 3 feet 11 1/4' to 6". It lies in an olive green mud-squeezed slate in which near the ore one sees a few poor *Spirifers*. These slates are like those of the *Siphonaria*.

All of the brachiopods are single valves and ^{some} show ^{some} marking together. Saw one species of a *Trochotoma*, *pleurac* of *Stromatolites*.

The strata lie in a syncline. Tor Brook is on the eastern limb and Nictaux on the western. The syncline may have a width of one mile.

For report on the Ojibwa-Torbrook area
see Fletcher, Summary Report C. E. S.
for 1904 (295-6), 293-318. Has also a
map.

If Helderberg are present it must lie more
eastward than the Oriskany.

Quincy Brook.

Friday, May 29-1914.

Drove from Middleton this morning to Hersenger Brook which is east of Phinney Brook. Both are east of Tor Brook Mills, and the latter is about 2 miles west of Hersenger Brook. Tor Brook Mills is two miles east of Tor Brook where we saw the iron mines yesterday.

Menomun Brook is a small brook flowing north down the hills of Paleozoic strata passing northward and the Triassic valley. The northern margin of this wide valley is made by North Mountain of Triassic sandstone and loam.

At the extreme southern margin of the
Triassic one soon comes in (Mazonia Brook
upon critical standing ^{existing} state and then a thick ^{series of}
shaly limestones and shale ^{is also} more ^{is} true. True

At the top of the fossiliferous Gristany Bell
got a very large head of *Præps* over 2 inches
across from side to side.

In the Lower Gristany shale Bell got
several large *Rhipidomella* *plata*.

are also some sandstones and arenaceous limestones.
The entire series is at least 300 feet thick, not counting
the ^{higher} ~~limestones~~ ^{slates}. Fossils occur in it almost throughout
but more commonly at the top beneath the dark
slates. The specimens are all poor due to the
great movement all of these beds have under-
gone. Initially the commonest fossil is Turbid-
leptus carinatus (see if near the European
Algonian form) and the rarest is Leptodus arvensis.
I saw no Rensselaeria and I don't think I saw
one Hippodamia parvius. Spizifer are common.
The Strophodontes are present in at least two
forms but saw no Leptodus lambdoides.
A very large Strophodontes is often seen in frag-
ments. I saw no Leptodus flabellites. Stroph-
odontes are not rare.

The Bedfordian is a series of thin
sandstones, siltstone like sandstones, sandy shales
and ^{in lower part of} arenaceous slates. In the main the bed.
is sandy. The thickness may be as much
as 800 feet. The bed is marked at the top

Overanga Bank

Location
Terning Overanga Farm

Triassic

from middle of the
all those for the
= Upper Devonian

Triassic
about 300'

from the
from the

from the

from the

from the

about 300'

about 500'

with occasional
has a cell or date of diatase, and the sandy
nature of the Held. distinguishes it at once
from the green slates of the Silurian below. Of
the Silurian slates at least 400 are exposed
at the top of the Hill where the road has its
origin. (50-75 feet)

About ^{here} the diatase still occurs a
lign sandstone abounding in *Spinifers* and other
fossils none of which are familiar. Above
the ^{lign} exposure seems to be *Cecroft*. Partially
300 feet ^{low} more fossils are common though
none of 50 to 100 feet. Here the *Spinifers*
are small associated with small *Stropho-*
donta of *Saplauna* and ^{in abundance of} *Phipdomella*. The
fauna is directly connected with the *Cecroft*
but the horizon marked in the upper New
Scotland though I saw nothing to make
this correlation probable. Below this fauna
the sandy ^{and shaly} nature of the Held continues to
the base but the fossils if present are not

seen.

The olive green Silurian ^{slates} & shales are like those seen at Beech Hill and elsewhere to the north. Bell found one fossil that maybe a *Chonetes* or an orthid.

The strata throughout practically stand on end or at least at 80 degrees. If there is a change in dip I did not notice it.

The section is a descending one from Wp. Crickany through to Beech Hill at least and probably into the island. Then probably a great time break (the contact seems to be a disconformable one) with the Silurian slates at the base of the section. Fletcher measured the section as upwards of 1500 feet thick. Of this 1100 feet is Lower Devonian (top not seen), and over 400 feet is Silurian (base nowhere seen).

Middleton - Yarmouth. May 30

Saw Bull eastward at 9. A.M.

I left Middleton at 11.24 for Yarmouth.

Got here at 4.45 P.M.

On board steamer Prince George. Left Yarmouth in a rain with some wind at 6. P.M. Have room 74 all to myself,

Got to Boston on a fine Sunday morning at 9 A.M. The steamer was ahead of schedule one-half hour. Left for N. Haven at 10. A.M.

May 24.	McClelland for car	London	£ 3.00
" 25	Antigonish to Toronto	25¢	25¢
" 25	" Royal Gorge Hotel		2.25
" 26	Learmark at Toronto		1.00
" 26	Toronto to Windsor		1.80
" 30	Middleton to Boston		8.00
" 30	" " Jarmouth Chain		.60
" 30	Meal on steamer (dinner)		.75
" 30	Berth " "		3.60
" 31	Breakfast " "		.50
" 31	Boston to N. H.		3.25
" 31	Baggage transfer 35¢ 35¢		.70
" 31	Dinner on cars		1.00
			<hr/>
			7 29.40

2.60
2.17
2.17
1.60
8.40
6.00
7.00
3.60
50
2.25
70
00
50

19.60

lyn
ae